

Computer algebra independent integration tests

Summer 2022 edition

5-Inverse-trig-functions/5.6-Inverse-cosecant/159-5.6.2-Inverse-cosecant-functions

Nasser M. Abbasi

September 27, 2022 Compiled on September 27, 2022 at 7:27am

Contents

1	Introduction	3
2	detailed summary tables of results	19
3	Listing of integrals	37
4	Appendix	263

Chapter 1

Introduction

Local contents

1.1	Listing of CAS systems tested	4
1.2	Results	5
1.3	Time and leaf size Performance	9
1.4	list of integrals that has no closed form antiderivative	11
1.5	List of integrals solved by CAS but has no known antiderivative	12
1.6	list of integrals solved by CAS but failed verification	13
1.7	Timing	13
1.8	Verification	14
1.9	Important notes about some of the results	14
1.10	Design of the test system	17

This report gives the result of running the computer algebra independent integration test. The download section in the appendix contains links to download the problems in plain text format used for all CAS systems.

The number of integrals in this report is [49]. This is test number [159].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 13.1 (June 29, 2022) on windows 10.
2. Rubi 4.16.1 (Dec 19, 2018) on Mathematica 13.0.1 on windows 10.
3. Maple 2022.1 (June 1, 2022) on windows 10.
4. Maxima 5.46 (April 13, 2022) using Lisp SBCL 2.1.11.debian on Linux via sagemath 9.6.
5. Fricas 1.3.8 (June 21, 2022) based on sbcl 2.1.11.debian on Linux via sagemath 9.6.
6. Giac/Xcas 1.9.0-13 (July 3, 2022) on Linux via sagemath 9.6.
7. Sympy 1.10.1 (March 20, 2022) Using Python 3.10.4 on Linux.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was called directly from Python.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed.

If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (49)	0.00 (0)
Mathematica	100.00 (49)	0.00 (0)
Maple	73.47 (36)	26.53 (13)
Fricas	55.10 (27)	44.90 (22)
Giac	51.02 (25)	48.98 (24)
Maxima	30.61 (15)	69.39 (34)
Mupad	24.49 (12)	75.51 (37)
Sympy	24.49 (12)	75.51 (37)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	100.00	0.00	0.00	0.00
Mathematica	77.55	12.24	10.20	0.00
Maple	61.22	12.24	0.00	26.53
Fricas	44.90	10.20	0.00	44.90
Maxima	28.57	2.04	0.00	69.39
Giac	24.49	26.53	0.00	48.98
Sympy	8.16	0.00	16.33	75.51
Mupad	N/A	24.49	0.00	75.51

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the CAS systems for each grade level.



The following table shows the distribution of the different types of failure for each CAS. There are 3 types of reasons why it can fail. The first is when CAS returns back the input within the time limit, which means it could not solve it. This is the typical normal failure **F**.

The second is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned **F(-1)**.

The third is due to an exception generated. Assigned **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and

Giac) or it could be an indication of an internal error in CAS. This type of error requires more investigations to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00 %	0.00 %	0.00 %
Mathematica	0	0.00 %	0.00 %	0.00 %
Maple	13	100.00 %	0.00 %	0.00 %
Fricas	22	90.91 %	0.00 %	9.09 %
Giac	24	95.83 %	0.00 %	4.17 %
Maxima	34	100.00 %	0.00 %	0.00 %
Sympy	37	94.59 %	2.70 %	2.70 %
Mupad	37	100.00 %	0.00 %	0.00 %

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.


Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)	Mean size	Normalized mean	Median size	Normalized median
Rubi	0.10	124.59	1.00	79.00	1.00
Mathematica	0.44	166.90	1.42	99.00	1.11
Maple	1.87	278.14	1.99	138.50	1.61
Maxima	0.34	50.80	1.20	52.00	1.14
Fricas	0.38	121.11	1.34	51.00	0.83
Sympy	12.80	67.42	1.55	54.50	1.63
Giac	0.47	155.00	1.80	81.00	1.75
Mupad	0.77	41.50	0.91	40.00	0.92

Table 1.5: Time and leaf size performance for each CAS

The following are bar charts for the normalized leafsize and time used from the above table.



1.4 list of integrals that has no closed form antiderivative

{}

1.5 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Mupad {}

1.6 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not mean necessarily that the anti-derivative is wrong, as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it easier to do further investigation to determine why it was not possible to verify the result produced.

Rubi {}

Mathematica {27, 28, 33}

Maple Verification phase not implemented yet.

Maxima Verification phase not implemented yet.

Fricas Verification phase not implemented yet.

Sympy Verification phase not implemented yet.

Giac Verification phase not implemented yet.

Mupad Verification phase not implemented yet.

1.7 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each `integrate` call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int', int(expr,x)), output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the `integrate` command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.8 Verification

A verification phase was applied on the result of integration for Rubi and Mathematica.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.9 Important notes about some of the results

1.9.1 Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima `integrate` was run using SageMath with the following settings set by default

```
'besselexpand : true'
'display2d : false'
'domain : complex'
'keepfloat : true'
'load(to_poly_solve)'
'load(simplify_sum)'
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib
maxima_lib.set('extra_definite_integration_methods', '[]')
```

```
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

1.9.2 Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError.`

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

1.9.3 Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    """
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount = 1
```

1.9.4 Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^{2/2}$

1.10 Design of the test system

The following diagram gives a high level view of the current test build system.



1. integer, the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "F"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified.

The following fields are present only in Rubi Table file

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Chapter 2

detailed summary tables of results

Local contents

2.1	List of integrals sorted by grade for each CAS	20
2.2	Detailed conclusion table per each integral for all CAS systems	23
2.3	Detailed conclusion table specific for Rubi results	34

2.1 List of integrals sorted by grade for each CAS

Local contents

2.1.1	Rubi	21
2.1.2	Mathematica	21
2.1.3	Maple	21
2.1.4	Maxima	21
2.1.5	FriCAS	22
2.1.6	Sympy	22
2.1.7	Giac	22
2.1.8	Mupad	22

2.1.1 Rubi

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49 }

B grade: { }

C grade: { }

F grade: { }

2.1.2 Mathematica

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 14, 15, 17, 18, 19, 20, 22, 27, 28, 29, 30, 31, 33, 34, 35, 36, 37, 41, 42, 43, 44, 45, 46, 47, 48, 49 }

B grade: { 13, 21, 32, 38, 39, 40 }

C grade: { 16, 23, 24, 25, 26 }

F grade: { }

2.1.3 Maple

A grade: { 2, 3, 4, 5, 6, 7, 8, 9, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 23, 27, 28, 29, 30, 32, 33, 34, 35, 38, 40, 49 }

B grade: { 10, 11, 22, 24, 25, 26 }

C grade: { }

F grade: { 1, 31, 36, 37, 39, 41, 42, 43, 44, 45, 46, 47, 48 }

2.1.4 Maxima

A grade: { 2, 3, 4, 5, 7, 9, 10, 11, 13, 14, 15, 21, 38, 39 }

B grade: { 8 }

C grade: { }

F grade: { 1, 6, 12, 16, 17, 18, 19, 20, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49 }

2.1.5 FriCAS

A grade: { 2, 3, 4, 5, 7, 8, 9, 10, 11, 14, 15, 17, 18, 19, 20, 24, 25, 26, 45, 46, 47, 48 }

B grade: { 13, 21, 23, 38, 39 }

C grade: { }

F grade: { 1, 6, 12, 16, 22, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 40, 41, 42, 43, 44, 49 }

2.1.6 Sympy

A grade: { 9, 10, 11, 15 }

B grade: { }

C grade: { 2, 3, 4, 5, 7, 8, 13, 14 }

F grade: { 1, 6, 12, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49 }

2.1.7 Giac

A grade: { 7, 8, 9, 10, 11, 14, 15, 20, 23, 39, 45, 49 }

B grade: { 2, 3, 4, 5, 13, 17, 18, 19, 21, 24, 25, 26, 38 }

C grade: { }

F grade: { 1, 6, 12, 16, 22, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 40, 41, 42, 43, 44, 46, 47, 48 }

2.1.8 Mupad

A grade: { }

B grade: { 1, 5, 6, 7, 10, 11, 12, 13, 21, 38, 39, 40 }

C grade: { }

F grade: { 2, 3, 4, 8, 9, 14, 15, 16, 17, 18, 19, 20, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 41, 42, 43, 44, 45, 46, 47, 48, 49 }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by table below. The elapsed time is in seconds. For failed result it is given as F(-1) if the failure was due to timeout. It is given as F(-2) if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given just an F.

In this table, the column N.S. in the table below, which stands for **normalized size** is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To help make the table fit, **Mathematica** was abbreviated to MMA.

	Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
MMA	grade	A	A	A	F	F	F	F	F	B
	verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
	size	62	62	56	0	0	0	0	0	58
	N.S.	1	1.00	0.90	0.00	0.00	0.00	0.00	0.00	0.94
	time (sec)	N/A	0.076	0.031	0.095	0.000	0.000	0.000	0.000	0.827

	Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
MMA	grade	A	A	A	A	A	A	C	B	F
	verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
	size	58	58	40	43	66	32	119	152	-1
	N.S.	1	1.00	0.69	0.74	1.14	0.55	2.05	2.62	-0.02
	time (sec)	N/A	0.015	0.021	0.208	0.255	0.356	61.545	0.435	0.000

	Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
MMA	grade	A	A	A	A	A	A	C	B	F
	verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
	size	47	47	35	38	52	27	90	116	-1
	N.S.	1	1.00	0.74	0.81	1.11	0.57	1.91	2.47	-0.02
	time (sec)	N/A	0.016	0.016	0.206	0.259	0.350	20.002	0.453	0.000

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	C	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	36	27	31	38	20	58	80	-1
N.S.	1	1.00	0.75	0.86	1.06	0.56	1.61	2.22	-0.03
time (sec)	N/A	0.009	0.017	0.215	0.259	0.338	6.266	0.452	0.000

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	C	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	16	24	20	12	29	41	20
N.S.	1	1.00	1.00	1.50	1.25	0.75	1.81	2.56	1.25
time (sec)	N/A	0.003	0.004	0.204	0.255	0.341	2.602	0.438	1.057

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	56	56	54	105	0	0	0	0	42
N.S.	1	1.00	0.96	1.88	0.00	0.00	0.00	0.00	0.75
time (sec)	N/A	0.056	0.026	0.595	0.000	0.000	0.000	0.000	0.847

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	C	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	32	46	51	21	76	32	28
N.S.	1	1.00	0.84	1.21	1.34	0.55	2.00	0.84	0.74
time (sec)	N/A	0.012	0.018	0.209	0.487	0.367	14.026	0.427	0.689

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	C	A	F	
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	54	54	55	57	80	30	146	59	-1
N.S.	1	1.00	1.02	1.06	1.48	0.56	2.70	1.09	-0.02
time (sec)	N/A	0.015	0.036	0.204	0.492	0.359	45.170	0.438	0.000

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	56	56	42	66	54	39	51	68	-1
N.S.	1	1.00	0.75	1.18	0.96	0.70	0.91	1.21	-0.02
time (sec)	N/A	0.031	0.023	0.456	0.480	0.339	0.127	0.455	0.000

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	47	47	44	90	46	38	41	48	38
N.S.	1	1.00	0.94	1.91	0.98	0.81	0.87	1.02	0.81
time (sec)	N/A	0.015	0.019	0.493	0.455	0.335	0.090	0.412	0.612

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	25	52	23	26	22	26	23
N.S.	1	1.00	1.00	2.08	0.92	1.04	0.88	1.04	0.92
time (sec)	N/A	0.007	0.008	0.486	0.260	0.340	0.061	0.427	0.098

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	59	59	54	125	0	0	0	0	49
N.S.	1	1.00	0.92	2.12	0.00	0.00	0.00	0.00	0.83
time (sec)	N/A	0.044	0.026	0.845	0.000	0.000	0.000	0.000	0.608

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	A	A	B	C	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	32	32	93	42	52	65	27	61	30
N.S.	1	1.00	2.91	1.31	1.62	2.03	0.84	1.91	0.94
time (sec)	N/A	0.022	0.100	0.073	0.262	0.382	1.391	0.449	0.590

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	C	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	35	54	32	38	51	61	-1
N.S.	1	1.00	0.92	1.42	0.84	1.00	1.34	1.61	-0.03
time (sec)	N/A	0.016	0.021	0.507	0.462	0.376	0.730	0.458	0.000

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	60	60	69	91	64	92	99	80	-1
N.S.	1	1.00	1.15	1.52	1.07	1.53	1.65	1.33	-0.02
time (sec)	N/A	0.031	0.037	0.509	0.451	0.364	1.606	0.448	0.000

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F(-2)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	63	155	0	0	0	0	-1
N.S.	1	1.00	0.91	2.25	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.071	0.063	1.320	0.000	0.000	0.000	0.000	0.000

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	197	197	174	329	0	151	0	408	-1
N.S.	1	1.00	0.88	1.67	0.00	0.77	0.00	2.07	-0.01
time (sec)	N/A	0.172	0.147	0.301	0.000	0.397	0.000	0.553	0.000

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	155	155	149	249	0	129	0	300	-1
N.S.	1	1.00	0.96	1.61	0.00	0.83	0.00	1.94	-0.01
time (sec)	N/A	0.107	0.185	0.308	0.000	0.374	0.000	0.473	0.000

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	116	116	129	190	0	117	0	203	-1
N.S.	1	1.00	1.11	1.64	0.00	1.01	0.00	1.75	-0.01
time (sec)	N/A	0.064	0.125	0.301	0.000	0.369	0.000	0.468	0.000

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	79	79	110	108	0	102	0	134	-1
N.S.	1	1.00	1.39	1.37	0.00	1.29	0.00	1.70	-0.01
time (sec)	N/A	0.039	0.081	0.301	0.000	0.395	0.000	0.466	0.000

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	A	A	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	36	336	43	55	75	0	81	33
N.S.	1	1.00	9.33	1.19	1.53	2.08	0.00	2.25	0.92
time (sec)	N/A	0.018	0.184	0.074	0.256	0.368	0.000	0.455	0.884

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	210	210	375	1144	0	0	0	0	-1
N.S.	1	1.00	1.79	5.45	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.215	0.288	6.025	0.000	0.000	0.000	0.000	0.000

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	B	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	115	127	0	278	0	96	-1
N.S.	1	1.00	1.67	1.84	0.00	4.03	0.00	1.39	-0.01
time (sec)	N/A	0.068	0.275	0.634	0.000	0.397	0.000	0.469	0.000

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	A	F	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	123	123	199	456	0	428	0	217	-1
N.S.	1	1.00	1.62	3.71	0.00	3.48	0.00	1.76	-0.01
time (sec)	N/A	0.150	0.568	0.644	0.000	0.424	0.000	0.484	0.000

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	A	F	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	180	180	241	970	0	548	0	450	-1
N.S.	1	1.00	1.34	5.39	0.00	3.04	0.00	2.50	-0.01
time (sec)	N/A	0.220	0.344	0.650	0.000	0.416	0.000	0.498	0.000

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	A	F	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	239	239	307	1702	0	673	0	841	-1
N.S.	1	1.00	1.28	7.12	0.00	2.82	0.00	3.52	-0.00
time (sec)	N/A	0.323	0.354	0.668	0.000	0.443	0.000	0.490	0.000

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F(-2)	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	366	366	420	703	0	0	0	0	-1
N.S.	1	1.00	1.15	1.92	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.202	3.483	12.517	0.000	0.000	0.000	0.000	0.000

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	272	272	314	500	0	0	0	0	-1
N.S.	1	1.00	1.15	1.84	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.157	3.137	10.441	0.000	0.000	0.000	0.000	0.000

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	145	145	213	214	0	0	0	0	-1
N.S.	1	1.00	1.47	1.48	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.094	0.623	1.051	0.000	0.000	0.000	0.000	0.000

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	86	86	99	150	0	0	0	0	-1
N.S.	1	1.00	1.15	1.74	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.044	0.096	0.563	0.000	0.000	0.000	0.000	0.000

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	324	324	408	0	0	0	0	0	-1
N.S.	1	1.00	1.26	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.340	0.238	2.384	0.000	0.000	0.000	0.000	0.000

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	254	254	802	302	0	0	0	0	-1
N.S.	1	1.00	3.16	1.19	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.285	2.125	6.843	0.000	0.000	0.000	0.000	0.000

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	464	464	656	749	0	0	0	0	-1
N.S.	1	1.00	1.41	1.61	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.280	5.986	9.851	0.000	0.000	0.000	0.000	0.000

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	264	264	314	425	0	0	0	0	-1
N.S.	1	1.00	1.19	1.61	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.185	0.589	6.128	0.000	0.000	0.000	0.000	0.000

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	140	140	162	224	0	0	0	0	-1
N.S.	1	1.00	1.16	1.60	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.076	0.076	1.148	0.000	0.000	0.000	0.000	0.000

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	448	448	554	0	0	0	0	0	-1
N.S.	1	1.00	1.24	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.386	0.224	2.533	0.000	0.000	0.000	0.000	0.000

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	378	378	289	0	0	0	0	0	-1
N.S.	1	1.00	0.76	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.415	0.341	2.861	0.000	0.000	0.000	0.000	0.000

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	A	A	B	F(-1)	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	48	48	127	54	63	88	0	91	44
N.S.	1	1.00	2.65	1.12	1.31	1.83	0.00	1.90	0.92
time (sec)	N/A	0.045	0.168	0.079	0.258	0.372	0.000	0.496	1.034

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	F	A	B	F(-2)	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	48	48	130	0	66	94	0	74	42
N.S.	1	1.00	2.71	0.00	1.38	1.96	0.00	1.54	0.88
time (sec)	N/A	0.052	0.191	0.042	0.265	0.390	0.000	0.497	0.726

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	A	F	F(-2)	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	85	85	280	188	0	0	0	0	91
N.S.	1	1.00	3.29	2.21	0.00	0.00	0.00	0.00	1.07
time (sec)	N/A	0.063	0.461	1.206	0.000	0.000	0.000	0.000	1.247

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	95	95	79	0	0	0	0	0	-1
N.S.	1	1.00	0.83	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.083	0.338	0.040	0.000	0.000	0.000	0.000	0.000

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	87	87	101	0	0	0	0	0	-1
N.S.	1	1.00	1.16	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.077	0.187	0.037	0.000	0.000	0.000	0.000	0.000

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	87	87	54	0	0	0	0	0	-1
N.S.	1	1.00	0.62	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.066	0.070	0.038	0.000	0.000	0.000	0.000	0.000

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	75	0	0	0	0	0	-1
N.S.	1	1.00	1.74	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.039	0.040	0.037	0.000	0.000	0.000	0.000	0.000

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	A	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	39	39	33	0	0	23	0	42	-1
N.S.	1	1.00	0.85	0.00	0.00	0.59	0.00	1.08	-0.03
time (sec)	N/A	0.021	0.030	0.037	0.000	0.400	0.000	0.454	0.000

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	A	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	30	0	0	32	0	0	-1
N.S.	1	1.00	0.73	0.00	0.00	0.78	0.00	0.00	-0.02
time (sec)	N/A	0.034	0.036	0.040	0.000	0.413	0.000	0.000	0.000

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	A	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	84	84	54	0	0	41	0	0	-1
N.S.	1	1.00	0.64	0.00	0.00	0.49	0.00	0.00	-0.01
time (sec)	N/A	0.046	0.100	0.037	0.000	0.418	0.000	0.000	0.000

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	A	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	81	81	50	0	0	51	0	0	-1
N.S.	1	1.00	0.62	0.00	0.00	0.63	0.00	0.00	-0.01
time (sec)	N/A	0.046	0.095	0.037	0.000	0.402	0.000	0.000	0.000

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	59	167	0	0	0	114	-1
N.S.	1	1.00	0.86	2.42	0.00	0.00	0.00	1.65	-0.01
time (sec)	N/A	0.066	0.041	0.878	0.000	0.000	0.000	0.660	0.000

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is given. The larger this ratio is, the harder the integral was to solve. In this test, problem number [49] had the largest ratio of [19]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	7	6	1.00	10	0.600
2	A	4	3	1.00	10	0.300
3	A	4	3	1.00	10	0.300
4	A	4	3	1.00	8	0.375
5	A	3	3	1.00	6	0.500
6	A	7	6	1.00	10	0.600
7	A	5	5	1.00	10	0.500
8	A	6	5	1.00	10	0.500
9	A	5	4	1.00	10	0.400
10	A	4	4	1.00	8	0.500
11	A	3	3	1.00	6	0.500
12	A	6	6	1.00	10	0.600
13	A	5	5	1.00	10	0.500
14	A	3	3	1.00	10	0.300
15	A	6	6	1.00	10	0.600
16	A	7	6	1.00	10	0.600
17	A	9	8	1.00	10	0.800
18	A	8	7	1.00	10	0.700
19	A	7	6	1.00	10	0.600
20	A	6	6	1.00	8	0.750
21	A	5	5	1.00	6	0.833
22	A	14	8	1.00	10	0.800
23	A	6	6	1.00	10	0.600
24	A	8	8	1.00	10	0.800
25	A	9	9	1.00	10	0.900

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	<u>number of rules</u> <u>integrand leaf size</u>
26	A	10	9	1.00	10	0.900
27	A	20	9	1.00	12	0.750
28	A	17	9	1.00	12	0.750
29	A	11	8	1.00	10	0.800
30	A	8	6	1.00	8	0.750
31	A	17	9	1.00	12	0.750
32	A	12	8	1.00	12	0.667
33	A	25	14	1.00	12	1.167
34	A	16	12	1.00	10	1.200
35	A	10	7	1.00	8	0.875
36	A	20	10	1.00	12	0.833
37	A	14	9	1.00	12	0.750
38	A	6	6	1.00	12	0.500
39	A	6	6	1.00	14	0.429
40	A	7	7	1.00	10	0.700
41	A	6	4	1.00	10	0.400
42	A	6	4	1.00	8	0.500
43	A	5	3	1.00	6	0.500
44	A	6	5	1.00	10	0.500
45	A	3	3	1.00	10	0.300
46	A	5	4	1.00	10	0.400
47	A	6	4	1.00	10	0.400
48	A	6	4	1.00	10	0.400
49	A	8	8	1.00	19	0.421

Chapter 3

Listing of integrals

Local contents

3.1	$\int \frac{\csc^{-1}(ax^5)}{x} dx$	38
3.2	$\int x^3 \csc^{-1}(\sqrt{x}) dx$	42
3.3	$\int x^2 \csc^{-1}(\sqrt{x}) dx$	46
3.4	$\int x \csc^{-1}(\sqrt{x}) dx$	50
3.5	$\int \csc^{-1}(\sqrt{x}) dx$	54
3.6	$\int \frac{\csc^{-1}(\sqrt{x})}{x} dx$	57
3.7	$\int \frac{\csc^{-1}(\sqrt{x})}{x^2} dx$	61
3.8	$\int \frac{\csc^{-1}(\sqrt{x})}{x^3} dx$	65
3.9	$\int x^2 \csc^{-1}\left(\frac{a}{x}\right) dx$	69
3.10	$\int x \csc^{-1}\left(\frac{a}{x}\right) dx$	73
3.11	$\int \csc^{-1}\left(\frac{a}{x}\right) dx$	77
3.12	$\int \frac{\csc^{-1}\left(\frac{a}{x}\right)}{x} dx$	80
3.13	$\int \frac{\csc^{-1}\left(\frac{a}{x}\right)}{x^2} dx$	84
3.14	$\int \frac{\csc^{-1}\left(\frac{a}{x}\right)}{x^3} dx$	88
3.15	$\int \frac{\csc^{-1}\left(\frac{a}{x}\right)}{x^4} dx$	92
3.16	$\int \frac{\csc^{-1}(ax^n)}{x} dx$	97
3.17	$\int x^4 \csc^{-1}(a+bx) dx$	101
3.18	$\int x^3 \csc^{-1}(a+bx) dx$	107
3.19	$\int x^2 \csc^{-1}(a+bx) dx$	112
3.20	$\int x \csc^{-1}(a+bx) dx$	117
3.21	$\int \csc^{-1}(a+bx) dx$	121
3.22	$\int \frac{\csc^{-1}(a+bx)}{x} dx$	125
3.23	$\int \frac{\csc^{-1}(a+bx)}{x^2} dx$	130
3.24	$\int \frac{\csc^{-1}(a+bx)}{x^3} dx$	135

3.25	$\int \frac{\csc^{-1}(a+bx)}{x^4} dx$	141
3.26	$\int \frac{\csc^{-1}(a+bx)}{x^5} dx$	147
3.27	$\int x^3 \csc^{-1}(a+bx)^2 dx$	155
3.28	$\int x^2 \csc^{-1}(a+bx)^2 dx$	161
3.29	$\int x \csc^{-1}(a+bx)^2 dx$	166
3.30	$\int \csc^{-1}(a+bx)^2 dx$	171
3.31	$\int \frac{\csc^{-1}(a+bx)^2}{x} dx$	175
3.32	$\int \frac{\csc^{-1}(a+bx)^2}{x^2} dx$	180
3.33	$\int x^2 \csc^{-1}(a+bx)^3 dx$	186
3.34	$\int x \csc^{-1}(a+bx)^3 dx$	193
3.35	$\int \csc^{-1}(a+bx)^3 dx$	200
3.36	$\int \frac{\csc^{-1}(a+bx)^3}{x} dx$	205
3.37	$\int \frac{\csc^{-1}(a+bx)^3}{x^2} dx$	211
3.38	$\int x^3 \csc^{-1}(a+bx^4) dx$	217
3.39	$\int x^{-1+n} \csc^{-1}(a+bx^n) dx$	222
3.40	$\int \csc^{-1}(ce^{a+bx}) dx$	226
3.41	$\int e^{\csc^{-1}(ax)} x^2 dx$	231
3.42	$\int e^{\csc^{-1}(ax)} x dx$	234
3.43	$\int e^{\csc^{-1}(ax)} dx$	238
3.44	$\int \frac{e^{\csc^{-1}(ax)}}{x} dx$	241
3.45	$\int \frac{e^{\csc^{-1}(ax)}}{x^2} dx$	245
3.46	$\int \frac{e^{\csc^{-1}(ax)}}{x^3} dx$	248
3.47	$\int \frac{e^{\csc^{-1}(ax)}}{x^4} dx$	251
3.48	$\int \frac{e^{\csc^{-1}(ax)}}{x^5} dx$	254
3.49	$\int \frac{\csc^{-1}(a+bx)}{\frac{ad}{b}+dx} dx$	257

3.1 $\int \frac{\csc^{-1}(ax^5)}{x} dx$

Optimal. Leaf size=62

$$\frac{1}{10}i \csc^{-1}(ax^5)^2 - \frac{1}{5} \csc^{-1}(ax^5) \log\left(1 - e^{2i \csc^{-1}(ax^5)}\right) + \frac{1}{10}i \text{PolyLog}\left(2, e^{2i \csc^{-1}(ax^5)}\right)$$

[Out] $\frac{1}{10}I*\text{arccsc}(a*x^5)^2 - \frac{1}{5}\text{arccsc}(a*x^5)*\ln(1 - (I/a/x^5 + (1 - 1/a^2/x^10)^{(1/2)})^2) + \frac{1}{10}I*\text{polylog}(2, (I/a/x^5 + (1 - 1/a^2/x^10)^{(1/2)})^2)$

Rubi [A]

time = 0.08, antiderivative size = 62, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.600, Rules used = {5327, 4721, 3798, 2221, 2317, 2438}

$$\frac{1}{10}i \text{Li}_2\left(e^{2i \csc^{-1}(ax^5)}\right) + \frac{1}{10}i \csc^{-1}(ax^5)^2 - \frac{1}{5} \csc^{-1}(ax^5) \log\left(1 - e^{2i \csc^{-1}(ax^5)}\right)$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{ArcCsc}[a*x^5]/x, x]$

[Out] $(I/10)*\text{ArcCsc}[a*x^5]^2 - (\text{ArcCsc}[a*x^5]*\text{Log}[1 - E^{((2*I)*\text{ArcCsc}[a*x^5])}])/5 + (I/10)*\text{PolyLog}[2, E^{((2*I)*\text{ArcCsc}[a*x^5])}]$

Rule 2221

```
Int[((F_)^((g_.)*(e_.) + (f_.)*(x_)))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/((a_) + (b_.)*((F_)^((g_.)*(e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] :> Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x]; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2317

```
Int[Log[(a_) + (b_.)*((F_)^((e_.)*(c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] :> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x]; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2438

```
Int[Log[(c_.)*(d_ + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x]; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 3798

```
Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + Pi*(k_.) + (f_.)*(x_)], x_Symbol] :> Simp[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Dist[2*I, Int[(c + d*x)^m
```

```
*E^(2*I*k*Pi)*(E^(2*I*(e + f*x))/(1 + E^(2*I*k*Pi)*E^(2*I*(e + f*x)))) , x] ,
x] /; FreeQ[{c, d, e, f}, x] && IntegerQ[4*k] && IGtQ[m, 0]
```

Rule 4721

```
Int[((a_.) + ArcSin[(c_)*(x_)]*(b_.))^(n_.)/(x_), x_Symbol] :> Subst[Int[(a + b*x)^n*Cot[x], x], x, ArcSin[c*x]] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0]
```

Rule 5327

```
Int[((a_.) + ArcCsc[(c_)*(x_)]*(b_.))/(x_), x_Symbol] :> -Subst[Int[(a + b*ArcSin[x/c])/x, x, 1/x]] /; FreeQ[{a, b, c}, x]
```

Rubi steps

$$\begin{aligned}
\int \frac{\csc^{-1}(ax^5)}{x} dx &= \frac{1}{5} \text{Subst}\left(\int \frac{\csc^{-1}(ax)}{x} dx, x, x^5\right) \\
&= -\left(\frac{1}{5} \text{Subst}\left(\int \frac{\sin^{-1}\left(\frac{x}{a}\right)}{x} dx, x, \frac{1}{x^5}\right)\right) \\
&= -\left(\frac{1}{5} \text{Subst}\left(\int x \cot(x) dx, x, \csc^{-1}(ax^5)\right)\right) \\
&= \frac{1}{10} i \csc^{-1}(ax^5)^2 + \frac{2}{5} i \text{Subst}\left(\int \frac{e^{2ix}x}{1-e^{2ix}} dx, x, \csc^{-1}(ax^5)\right) \\
&= \frac{1}{10} i \csc^{-1}(ax^5)^2 - \frac{1}{5} \csc^{-1}(ax^5) \log\left(1 - e^{2i \csc^{-1}(ax^5)}\right) + \frac{1}{5} \text{Subst}\left(\int \log(1 - e^{2ix}) dx, x, \csc^{-1}(ax^5)\right) \\
&= \frac{1}{10} i \csc^{-1}(ax^5)^2 - \frac{1}{5} \csc^{-1}(ax^5) \log\left(1 - e^{2i \csc^{-1}(ax^5)}\right) - \frac{1}{10} i \text{Subst}\left(\int \frac{\log(1-x)}{x} dx, x, \csc^{-1}(ax^5)\right) \\
&= \frac{1}{10} i \csc^{-1}(ax^5)^2 - \frac{1}{5} \csc^{-1}(ax^5) \log\left(1 - e^{2i \csc^{-1}(ax^5)}\right) + \frac{1}{10} i \text{Li}_2\left(e^{2i \csc^{-1}(ax^5)}\right)
\end{aligned}$$

Mathematica [A]

time = 0.03, size = 56, normalized size = 0.90

$$\frac{1}{10} i \left(\csc^{-1}(ax^5) \left(\csc^{-1}(ax^5) + 2i \log\left(1 - e^{2i \csc^{-1}(ax^5)}\right) \right) + \text{PolyLog}\left(2, e^{2i \csc^{-1}(ax^5)}\right) \right)$$

Antiderivative was successfully verified.

[In] `Integrate[ArcCsc[a*x^5]/x, x]`

[Out] `(I/10)*(ArcCsc[a*x^5]*(ArcCsc[a*x^5] + (2*I)*Log[1 - E^((2*I)*ArcCsc[a*x^5])]) + PolyLog[2, E^((2*I)*ArcCsc[a*x^5])])`

Maple [F]

time = 0.10, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{arccsc}(ax^5)}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(arccsc(a*x^5)/x,x)`

[Out] `int(arccsc(a*x^5)/x,x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(a*x^5)/x,x, algorithm="maxima")`

[Out] `5*a^2*integrate(sqrt(a*x^5 + 1)*sqrt(a*x^5 - 1)*log(x)/(a^4*x^11 - a^2*x), x) - 5*I*a^2*integrate(log(x)/(a^4*x^11 - a^2*x), x) + (arctan2(1, sqrt(a*x^5 + 1)*sqrt(a*x^5 - 1)) + I*log(a))*log(x) - 1/2*I*log(a^2*x^10)*log(x) + 1/2*I*log(a*x^5 + 1)*log(x) + 1/2*I*log(-a*x^5 + 1)*log(x) + 5/2*I*log(x)^2 + 1/10*I*dilog(a*x^5) + 1/10*I*dilog(-a*x^5)`

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(a*x^5)/x,x, algorithm="fricas")`

[Out] `integral(arccsc(a*x^5)/x, x)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{acsc}(ax^5)}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(acsc(a*x**5)/x,x)`

[Out] `Integral(acsc(a*x**5)/x, x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(arccsc(a*x^5)/x,x, algorithm="giac")

[Out] integrate(arccsc(a*x^5)/x, x)

Mupad [B]

time = 0.83, size = 58, normalized size = 0.94

$$-\frac{\ln \left(1 - e^{\operatorname{asin}\left(\frac{1}{ax^5}\right) 2i}\right) \operatorname{asin}\left(\frac{1}{ax^5}\right)}{5} + \frac{\operatorname{polylog}\left(2, e^{\operatorname{asin}\left(\frac{1}{ax^5}\right) 2i}\right) 1i}{10} + \frac{\operatorname{asin}\left(\frac{1}{ax^5}\right)^2 1i}{10}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(asin(1/(a*x^5))/x,x)

[Out]
$$\frac{(\operatorname{polylog}(2, \exp(\operatorname{asin}(1/(a*x^5))*2i))*1i)/10 - (\log(1 - \exp(\operatorname{asin}(1/(a*x^5))*2i))*\operatorname{asin}(1/(a*x^5)))/5 + (\operatorname{asin}(1/(a*x^5))^2*1i)/10}{}$$

3.2 $\int x^3 \csc^{-1}(\sqrt{x}) dx$

Optimal. Leaf size=58

$$\frac{\sqrt{-1+x}}{4} + \frac{1}{4}(-1+x)^{3/2} + \frac{3}{20}(-1+x)^{5/2} + \frac{1}{28}(-1+x)^{7/2} + \frac{1}{4}x^4 \csc^{-1}(\sqrt{x})$$

[Out] $1/4*(-1+x)^{(3/2)}+3/20*(-1+x)^{(5/2)}+1/28*(-1+x)^{(7/2)}+1/4*x^4*\text{arccsc}(x^{(1/2)})+1/4*(-1+x)^{(1/2)}$

Rubi [A]

time = 0.01, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.300, Rules used = {5379, 12, 45}

$$\frac{1}{4}x^4 \csc^{-1}(\sqrt{x}) + \frac{1}{28}(x-1)^{7/2} + \frac{3}{20}(x-1)^{5/2} + \frac{1}{4}(x-1)^{3/2} + \frac{\sqrt{x-1}}{4}$$

Antiderivative was successfully verified.

[In] $\text{Int}[x^3 \text{ArcCsc}[\text{Sqrt}[x]], x]$

[Out] $\text{Sqrt}[-1+x]/4 + (-1+x)^{(3/2)}/4 + (3*(-1+x)^{(5/2)})/20 + (-1+x)^{(7/2)}/28 + (x^4 \text{ArcCsc}[\text{Sqrt}[x]])/4$

Rule 12

```
Int[(a_)*(u_), x_Symbol] :> Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]
```

Rule 45

```
Int[((a_.) + (b_.)*(x_.))^(m_.)*((c_.) + (d_.)*(x_.))^(n_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])
```

Rule 5379

```
Int[((a_.) + ArcCsc[u_]*(b_.))*((c_.) + (d_.)*(x_.))^(m_.), x_Symbol] :> Simp[((c + d*x)^(m + 1)*((a + b*ArcCsc[u])/(d*(m + 1))), x] + Dist[b*(u/(d*(m + 1)*Sqrt[u^2])), Int[SimplifyIntegrand[((c + d*x)^(m + 1)*(D[u, x]/(u*Sqrt[u^2 - 1])), x], x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1] && InverseFunctionFreeQ[u, x] && !FunctionOfQ[(c + d*x)^(m + 1), u, x] && !FunctionOfExponentialQ[u, x]
```

Rubi steps

$$\begin{aligned}
\int x^3 \csc^{-1}(\sqrt{x}) dx &= \frac{1}{4}x^4 \csc^{-1}(\sqrt{x}) + \frac{1}{4} \int \frac{x^3}{2\sqrt{-1+x}} dx \\
&= \frac{1}{4}x^4 \csc^{-1}(\sqrt{x}) + \frac{1}{8} \int \frac{x^3}{\sqrt{-1+x}} dx \\
&= \frac{1}{4}x^4 \csc^{-1}(\sqrt{x}) + \frac{1}{8} \int \left(\frac{1}{\sqrt{-1+x}} + 3\sqrt{-1+x} + 3(-1+x)^{3/2} + (-1+x)^{5/2} \right) \\
&= \frac{\sqrt{-1+x}}{4} + \frac{1}{4}(-1+x)^{3/2} + \frac{3}{20}(-1+x)^{5/2} + \frac{1}{28}(-1+x)^{7/2} + \frac{1}{4}x^4 \csc^{-1}(\sqrt{x})
\end{aligned}$$

Mathematica [A]

time = 0.02, size = 40, normalized size = 0.69

$$\frac{1}{140}\sqrt{-1+x}(16+8x+6x^2+5x^3) + \frac{1}{4}x^4 \csc^{-1}(\sqrt{x})$$

Antiderivative was successfully verified.

[In] Integrate[x^3*ArcCsc[Sqrt[x]], x]

[Out] (Sqrt[-1 + x]*(16 + 8*x + 6*x^2 + 5*x^3))/140 + (x^4*ArcCsc[Sqrt[x]])/4

Maple [A]

time = 0.21, size = 43, normalized size = 0.74

method	result	size
derivative divides	$\frac{x^4 \operatorname{arccsc}(\sqrt{x})}{4} + \frac{(x-1)(5x^3+6x^2+8x+16)}{140 \sqrt{\frac{x-1}{x}}} \sqrt{x}$	43
default	$\frac{x^4 \operatorname{arccsc}(\sqrt{x})}{4} + \frac{(x-1)(5x^3+6x^2+8x+16)}{140 \sqrt{\frac{x-1}{x}}} \sqrt{x}$	43

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^3*arccsc(x^(1/2)), x, method=_RETURNVERBOSE)

[Out] 1/4*x^4*arccsc(x^(1/2))+1/140*(x-1)*(5*x^3+6*x^2+8*x+16)/((x-1)/x)^(1/2)/x^(1/2)

Maxima [A]

time = 0.25, size = 66, normalized size = 1.14

$$\frac{1}{28}x^{\frac{7}{2}}\left(-\frac{1}{x}+1\right)^{\frac{7}{2}} + \frac{3}{20}x^{\frac{5}{2}}\left(-\frac{1}{x}+1\right)^{\frac{5}{2}} + \frac{1}{4}x^4 \operatorname{arccsc}(\sqrt{x}) + \frac{1}{4}x^{\frac{3}{2}}\left(-\frac{1}{x}+1\right)^{\frac{3}{2}} + \frac{1}{4}\sqrt{x} \sqrt{-\frac{1}{x}+1}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x^3*arccsc(x^(1/2)),x, algorithm="maxima")
[Out] 1/28*x^(7/2)*(-1/x + 1)^(7/2) + 3/20*x^(5/2)*(-1/x + 1)^(5/2) + 1/4*x^4*arc
csc(sqrt(x)) + 1/4*x^(3/2)*(-1/x + 1)^(3/2) + 1/4*sqrt(x)*sqrt(-1/x + 1)
```

Fricas [A]

time = 0.36, size = 32, normalized size = 0.55

$$\frac{1}{4} x^4 \operatorname{arccsc}(\sqrt{x}) + \frac{1}{140} (5x^3 + 6x^2 + 8x + 16)\sqrt{x-1}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x^3*arccsc(x^(1/2)),x, algorithm="fricas")
[Out] 1/4*x^4*arccsc(sqrt(x)) + 1/140*(5*x^3 + 6*x^2 + 8*x + 16)*sqrt(x - 1)
```

Sympy [C] Result contains complex when optimal does not.

time = 61.55, size = 119, normalized size = 2.05

$$\frac{x^4 \operatorname{acsc}(\sqrt{x})}{4} + \begin{cases} \frac{2x^3\sqrt{x-1}}{7} + \frac{12x^2\sqrt{x-1}}{35} + \frac{16x\sqrt{x-1}}{35} + \frac{32\sqrt{x-1}}{35} & \text{for } |x| > 1 \\ \frac{2ix^3\sqrt{1-x}}{7} + \frac{12ix^2\sqrt{1-x}}{35} + \frac{16ix\sqrt{1-x}}{35} + \frac{32i\sqrt{1-x}}{35} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x**3*acsc(x**(1/2)),x)
[Out] x**4*acsc(sqrt(x))/4 + Piecewise((2*x**3*sqrt(x - 1)/7 + 12*x**2*sqrt(x - 1)/35 + 16*x*sqrt(x - 1)/35 + 32*sqrt(x - 1)/35, Abs(x) > 1), (2*I*x**3*sqrt(1 - x)/7 + 12*I*x**2*sqrt(1 - x)/35 + 16*I*x*sqrt(1 - x)/35 + 32*I*sqrt(1 - x)/35, True))/8
```

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 152 vs. 2(38) = 76.

time = 0.43, size = 152, normalized size = 2.62

$$\frac{1}{3584} x^{\frac{7}{2}} \left(\sqrt{-\frac{1}{x} + 1} - 1 \right)^7 + \frac{7}{2560} x^{\frac{5}{2}} \left(\sqrt{-\frac{1}{x} + 1} - 1 \right)^5 + \frac{1}{4} x^4 \arcsin \left(\frac{1}{\sqrt{x}} \right) + \frac{7}{512} x^{\frac{3}{2}} \left(\sqrt{-\frac{1}{x} + 1} - 1 \right)^3 + \frac{35}{512} \sqrt{x} \left(\sqrt{-\frac{1}{x} + 1} - 1 \right) - \frac{1225 x^3 \left(\sqrt{-\frac{1}{x} + 1} - 1 \right)^6 + 245 x^2 \left(\sqrt{-\frac{1}{x} + 1} - 1 \right)^4 + 49 x \left(\sqrt{-\frac{1}{x} + 1} - 1 \right)^2 + 5}{17920 x^{\frac{1}{2}} \left(\sqrt{-\frac{1}{x} + 1} - 1 \right)^7}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x^3*arccsc(x^(1/2)),x, algorithm="giac")
[Out] 1/3584*x^(7/2)*(sqrt(-1/x + 1) - 1)^7 + 7/2560*x^(5/2)*(sqrt(-1/x + 1) - 1)^5 + 1/4*x^4*arcsin(1/sqrt(x)) + 7/512*x^(3/2)*(sqrt(-1/x + 1) - 1)^3 + 35/512*sqrt(x)*(sqrt(-1/x + 1) - 1) - 1/17920*(1225*x^3*(sqrt(-1/x + 1) - 1)^6
```

$$+ 245*x^2*(sqrt(-1/x + 1) - 1)^4 + 49*x*(sqrt(-1/x + 1) - 1)^2 + 5)/(x^{(7/2)}*(sqrt(-1/x + 1) - 1)^7)$$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int x^3 \arcsin\left(\frac{1}{\sqrt{x}}\right) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^3*asin(1/x^(1/2)),x)

[Out] int(x^3*asin(1/x^(1/2)), x)

3.3 $\int x^2 \csc^{-1}(\sqrt{x}) dx$

Optimal. Leaf size=47

$$\frac{\sqrt{-1+x}}{3} + \frac{2}{9}(-1+x)^{3/2} + \frac{1}{15}(-1+x)^{5/2} + \frac{1}{3}x^3 \csc^{-1}(\sqrt{x})$$

[Out] $2/9*(-1+x)^{(3/2)}+1/15*(-1+x)^{(5/2)}+1/3*x^3*\text{arccsc}(x^{(1/2)})+1/3*(-1+x)^{(1/2)}$

Rubi [A]

time = 0.02, antiderivative size = 47, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.300, Rules used = {5379, 12, 45}

$$\frac{1}{3}x^3 \csc^{-1}(\sqrt{x}) + \frac{1}{15}(x-1)^{5/2} + \frac{2}{9}(x-1)^{3/2} + \frac{\sqrt{x-1}}{3}$$

Antiderivative was successfully verified.

[In] Int[x^2*ArcCsc[Sqrt[x]], x]

[Out] $\text{Sqrt}[-1+x]/3 + (2*(-1+x)^{(3/2)})/9 + (-1+x)^{(5/2)}/15 + (x^3*\text{ArcCsc}[\text{Sqr}t[x]])/3$

Rule 12

Int[(a_)*(u_), x_Symbol] :> Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]

Rule 45

Int[((a_.) + (b_.)*(x_.))^(m_.)*((c_.) + (d_.)*(x_.))^n_, x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])

Rule 5379

Int[((a_.) + ArcCsc[u_]*(b_.))*((c_.) + (d_.)*(x_.))^m_, x_Symbol] :> Simplify[((c + d*x)^(m + 1)*((a + b*ArcCsc[u])/(d*(m + 1))), x] + Dist[b*(u/(d*(m + 1)*Sqrt[u^2 - 1])), Int[SimplifyIntegrand[((c + d*x)^(m + 1)*(D[u, x]/(u*Sqrt[u^2 - 1])), x], x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1] && InverseFunctionFreeQ[u, x] && !FunctionOfQ[(c + d*x)^(m + 1), u, x] && !FunctionOfExponentialQ[u, x]

Rubi steps

$$\begin{aligned}
\int x^2 \csc^{-1}(\sqrt{x}) dx &= \frac{1}{3}x^3 \csc^{-1}(\sqrt{x}) + \frac{1}{3} \int \frac{x^2}{2\sqrt{-1+x}} dx \\
&= \frac{1}{3}x^3 \csc^{-1}(\sqrt{x}) + \frac{1}{6} \int \frac{x^2}{\sqrt{-1+x}} dx \\
&= \frac{1}{3}x^3 \csc^{-1}(\sqrt{x}) + \frac{1}{6} \int \left(\frac{1}{\sqrt{-1+x}} + 2\sqrt{-1+x} + (-1+x)^{3/2} \right) dx \\
&= \frac{\sqrt{-1+x}}{3} + \frac{2}{9}(-1+x)^{3/2} + \frac{1}{15}(-1+x)^{5/2} + \frac{1}{3}x^3 \csc^{-1}(\sqrt{x})
\end{aligned}$$

Mathematica [A]

time = 0.02, size = 35, normalized size = 0.74

$$\frac{1}{45}\sqrt{-1+x}(8+4x+3x^2) + \frac{1}{3}x^3 \csc^{-1}(\sqrt{x})$$

Antiderivative was successfully verified.

[In] Integrate[x^2*ArcCsc[Sqrt[x]], x]

[Out] (Sqrt[-1 + x]*(8 + 4*x + 3*x^2))/45 + (x^3*ArcCsc[Sqrt[x]]))/3

Maple [A]

time = 0.21, size = 38, normalized size = 0.81

method	result	size
derivativedivides	$\frac{x^3 \operatorname{arccsc}(\sqrt{x})}{3} + \frac{(x-1)(3x^2+4x+8)}{45 \sqrt{\frac{x-1}{x}} \sqrt{x}}$	38
default	$\frac{x^3 \operatorname{arccsc}(\sqrt{x})}{3} + \frac{(x-1)(3x^2+4x+8)}{45 \sqrt{\frac{x-1}{x}} \sqrt{x}}$	38

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^2*arccsc(x^(1/2)), x, method=_RETURNVERBOSE)

[Out] 1/3*x^3*arccsc(x^(1/2))+1/45*(x-1)*(3*x^2+4*x+8)/((x-1)/x)^(1/2)/x^(1/2)

Maxima [A]

time = 0.26, size = 52, normalized size = 1.11

$$\frac{1}{15}x^{\frac{5}{2}}\left(-\frac{1}{x}+1\right)^{\frac{5}{2}} + \frac{1}{3}x^3 \operatorname{arccsc}(\sqrt{x}) + \frac{2}{9}x^{\frac{3}{2}}\left(-\frac{1}{x}+1\right)^{\frac{3}{2}} + \frac{1}{3}\sqrt{x} \sqrt{-\frac{1}{x}+1}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x^2*arccsc(x^(1/2)),x, algorithm="maxima")
[Out] 1/15*x^(5/2)*(-1/x + 1)^(5/2) + 1/3*x^3*arccsc(sqrt(x)) + 2/9*x^(3/2)*(-1/x + 1)^(3/2) + 1/3*sqrt(x)*sqrt(-1/x + 1)
```

Fricas [A]

time = 0.35, size = 27, normalized size = 0.57

$$\frac{1}{3} x^3 \operatorname{arccsc}(\sqrt{x}) + \frac{1}{45} (3x^2 + 4x + 8)\sqrt{x-1}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x^2*arccsc(x^(1/2)),x, algorithm="fricas")
[Out] 1/3*x^3*arccsc(sqrt(x)) + 1/45*(3*x^2 + 4*x + 8)*sqrt(x - 1)
```

Sympy [C] Result contains complex when optimal does not.

time = 20.00, size = 90, normalized size = 1.91

$$\frac{x^3 \operatorname{acsc}(\sqrt{x})}{3} + \frac{\begin{cases} \frac{2x^2\sqrt{x-1}}{5} + \frac{8x\sqrt{x-1}}{15} + \frac{16\sqrt{x-1}}{15} & \text{for } |x| > 1 \\ \frac{2ix^2\sqrt{1-x}}{5} + \frac{8ix\sqrt{1-x}}{15} + \frac{16i\sqrt{1-x}}{15} & \text{otherwise} \end{cases}}{6}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x**2*acsc(x**(1/2)),x)
[Out] x**3*acsc(sqrt(x))/3 + Piecewise((2*x**2*sqrt(x - 1)/5 + 8*x*sqrt(x - 1)/15 + 16*sqrt(x - 1)/15, Abs(x) > 1), (2*I*x**2*sqrt(1 - x)/5 + 8*I*x*sqrt(1 - x)/15 + 16*I*sqrt(1 - x)/15, True))/6
```

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 116 vs. 2(31) = 62.

time = 0.45, size = 116, normalized size = 2.47

$$\frac{1}{480} x^{\frac{5}{2}} \left(\sqrt{-\frac{1}{x} + 1} - 1 \right)^5 + \frac{5}{288} x^{\frac{3}{2}} \left(\sqrt{-\frac{1}{x} + 1} - 1 \right)^3 + \frac{1}{3} x^3 \arcsin \left(\frac{1}{\sqrt{x}} \right) + \frac{5}{48} \sqrt{x} \left(\sqrt{-\frac{1}{x} + 1} - 1 \right) - \frac{150 x^2 \left(\sqrt{-\frac{1}{x} + 1} - 1 \right)^4 + 25 x \left(\sqrt{-\frac{1}{x} + 1} - 1 \right)^2 + 3}{1440 x^{\frac{5}{2}} \left(\sqrt{-\frac{1}{x} + 1} - 1 \right)^5}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x^2*arccsc(x^(1/2)),x, algorithm="giac")
[Out] 1/480*x^(5/2)*(sqrt(-1/x + 1) - 1)^5 + 5/288*x^(3/2)*(sqrt(-1/x + 1) - 1)^3 + 1/3*x^3*arcsin(1/sqrt(x)) + 5/48*sqrt(x)*(sqrt(-1/x + 1) - 1) - 1/1440*(
```

$150*x^2*(\sqrt{-1/x + 1} - 1)^4 + 25*x*(\sqrt{-1/x + 1} - 1)^2 + 3)/(x^{(5/2)} * (\sqrt{-1/x + 1} - 1)^5)$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int x^2 \arcsin\left(\frac{1}{\sqrt{x}}\right) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{int}(x^2 * \arcsin(1/x^{(1/2)}), x)$

[Out] $\text{int}(x^2 * \arcsin(1/x^{(1/2)}), x)$

$$\mathbf{3.4} \quad \int x \csc^{-1}(\sqrt{x}) \, dx$$

Optimal. Leaf size=36

$$\frac{\sqrt{-1+x}}{2} + \frac{1}{6}(-1+x)^{3/2} + \frac{1}{2}x^2 \csc^{-1}(\sqrt{x})$$

[Out] $1/6*(-1+x)^{(3/2)}+1/2*x^2*\text{arccsc}(x^{(1/2)})+1/2*(-1+x)^{(1/2)}$

Rubi [A]

time = 0.01, antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {5379, 12, 45}

$$\frac{1}{2}x^2 \csc^{-1}(\sqrt{x}) + \frac{1}{6}(x-1)^{3/2} + \frac{\sqrt{x-1}}{2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[x*\text{ArcCsc}[\text{Sqrt}[x]], x]$

[Out] $\text{Sqrt}[-1+x]/2 + (-1+x)^{(3/2)}/6 + (x^2*\text{ArcCsc}[\text{Sqrt}[x]])/2$

Rule 12

```
Int[(a_)*(u_), x_Symbol] :> Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]
```

Rule 45

```
Int[((a_.) + (b_.)*(x_.))^(m_.)*((c_.) + (d_.)*(x_.))^n_, x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])
```

Rule 5379

```
Int[((a_.) + ArcCsc[u_]*(b_.))*((c_.) + (d_.)*(x_.))^m_, x_Symbol] :> Simplify[((c + d*x)^(m + 1)*((a + b*ArcCsc[u])/(d*(m + 1))), x] + Dist[b*(u/(d*(m + 1)*Sqrt[u^2])), Int[SimplifyIntegrand[(c + d*x)^(m + 1)*(D[u, x]/(u*Sqrt[u^2 - 1])), x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1] && InverseFunctionFreeQ[u, x] && !FunctionOfQ[(c + d*x)^(m + 1), u, x] && !FunctionOfExponentialQ[u, x]]
```

Rubi steps

$$\begin{aligned}
\int x \csc^{-1}(\sqrt{x}) dx &= \frac{1}{2}x^2 \csc^{-1}(\sqrt{x}) + \frac{1}{2} \int \frac{x}{2\sqrt{-1+x}} dx \\
&= \frac{1}{2}x^2 \csc^{-1}(\sqrt{x}) + \frac{1}{4} \int \frac{x}{\sqrt{-1+x}} dx \\
&= \frac{1}{2}x^2 \csc^{-1}(\sqrt{x}) + \frac{1}{4} \int \left(\frac{1}{\sqrt{-1+x}} + \sqrt{-1+x} \right) dx \\
&= \frac{\sqrt{-1+x}}{2} + \frac{1}{6}(-1+x)^{3/2} + \frac{1}{2}x^2 \csc^{-1}(\sqrt{x})
\end{aligned}$$

Mathematica [A]

time = 0.02, size = 27, normalized size = 0.75

$$\frac{1}{6}(\sqrt{-1+x}(2+x) + 3x^2 \csc^{-1}(\sqrt{x}))$$

Antiderivative was successfully verified.

[In] `Integrate[x*ArcCsc[Sqrt[x]], x]`[Out] `(Sqrt[-1 + x]*(2 + x) + 3*x^2*ArcCsc[Sqrt[x]])/6`**Maple [A]**

time = 0.22, size = 31, normalized size = 0.86

method	result	size
derivativedivides	$\frac{x^2 \operatorname{arccsc}(\sqrt{x})}{2} + \frac{(x-1)(x+2)}{6 \sqrt{\frac{x-1}{x}} \sqrt{x}}$	31
default	$\frac{x^2 \operatorname{arccsc}(\sqrt{x})}{2} + \frac{(x-1)(x+2)}{6 \sqrt{\frac{x-1}{x}} \sqrt{x}}$	31

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x*arccsc(x^(1/2)), x, method=_RETURNVERBOSE)`[Out] `1/2*x^2*arccsc(x^(1/2))+1/6*(x-1)*(x+2)/((x-1)/x)^(1/2)/x^(1/2)`**Maxima [A]**

time = 0.26, size = 38, normalized size = 1.06

$$\frac{1}{6}x^{\frac{3}{2}}\left(-\frac{1}{x} + 1\right)^{\frac{3}{2}} + \frac{1}{2}x^2 \operatorname{arccsc}(\sqrt{x}) + \frac{1}{2}\sqrt{x} \sqrt{-\frac{1}{x} + 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x*arccsc(x^(1/2)),x, algorithm="maxima")`
[Out] $1/6x^{(3/2)}(-1/x + 1)^{(3/2)} + 1/2x^2\text{arccsc}(\sqrt{x}) + 1/2\sqrt{x}\sqrt{-1/x + 1}$

Fricas [A]

time = 0.34, size = 20, normalized size = 0.56

$$\frac{1}{2}x^2 \text{arccsc}(\sqrt{x}) + \frac{1}{6}(x+2)\sqrt{x-1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x*arccsc(x^(1/2)),x, algorithm="fricas")`
[Out] $1/2x^2\text{arccsc}(\sqrt{x}) + 1/6(x+2)\sqrt{x-1}$

Sympy [C] Result contains complex when optimal does not.

time = 6.27, size = 58, normalized size = 1.61

$$\frac{x^2 \text{acsc}(\sqrt{x})}{2} + \frac{\begin{cases} \frac{x\sqrt{x-1}}{3} + \frac{2\sqrt{x-1}}{3} & \text{for } |x| > 1 \\ \frac{ix\sqrt{1-x}}{3} + \frac{2i\sqrt{1-x}}{3} & \text{otherwise} \end{cases}}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x*acsc(x**(1/2)),x)`
[Out] $x^{(3/2)}\text{acsc}(\sqrt{x})/2 + \text{Piecewise}((x\sqrt{x-1})/3 + 2\sqrt{x-1}/3, \text{Abs}(x) > 1, (I*x\sqrt{1-x})/3 + 2*I\sqrt{1-x}/3, \text{True})/2$

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 80 vs. $2(24) = 48$.
time = 0.45, size = 80, normalized size = 2.22

$$\frac{1}{48}x^{\frac{3}{2}}\left(\sqrt{-\frac{1}{x}+1}-1\right)^3 + \frac{1}{2}x^2\arcsin\left(\frac{1}{\sqrt{x}}\right) + \frac{3}{16}\sqrt{x}\left(\sqrt{-\frac{1}{x}+1}-1\right) - \frac{9x\left(\sqrt{-\frac{1}{x}+1}-1\right)^2+1}{48x^{\frac{3}{2}}\left(\sqrt{-\frac{1}{x}+1}-1\right)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x*arccsc(x^(1/2)),x, algorithm="giac")`
[Out] $1/48x^{(3/2)}(\sqrt{-1/x + 1} - 1)^3 + 1/2x^2\arcsin(1/\sqrt{x}) + 3/16\sqrt{x}(\sqrt{-1/x + 1} - 1) - 1/48*(9x*(\sqrt{-1/x + 1} - 1)^2 + 1)/(x^{(3/2)}(\sqrt{-1/x + 1} - 1)^3)$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.03

$$\int x \arcsin\left(\frac{1}{\sqrt{x}}\right) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x*asin(1/x^(1/2)),x)`

[Out] `int(x*asin(1/x^(1/2)), x)`

$$3.5 \quad \int \csc^{-1}(\sqrt{x}) \, dx$$

Optimal. Leaf size=16

$$\sqrt{-1+x} + x \csc^{-1}(\sqrt{x})$$

[Out] $x \operatorname{arccsc}(x^{1/2}) + (-1+x)^{1/2}$

Rubi [A]

time = 0.00, antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 6, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.500, Rules used = {5377, 12, 32}

$$\sqrt{x-1} + x \csc^{-1}(\sqrt{x})$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[\operatorname{ArcCsc}[\operatorname{Sqrt}[x]], x]$

[Out] $\operatorname{Sqrt}[-1+x] + x \operatorname{ArcCsc}[\operatorname{Sqrt}[x]]$

Rule 12

$\operatorname{Int}[(a_*)*(u_), x_{\text{Symbol}}] \Rightarrow \operatorname{Dist}[a, \operatorname{Int}[u, x], x] /; \operatorname{FreeQ}[a, x] \& \& \operatorname{!MatchQ}[u, (b_*)*(v_*) /; \operatorname{FreeQ}[b, x]]$

Rule 32

$\operatorname{Int}[((a_.) + (b_.)*(x_.))^{(m_.)}, x_{\text{Symbol}}] \Rightarrow \operatorname{Simp}[(a + b*x)^{(m + 1)} / (b*(m + 1)), x] /; \operatorname{FreeQ}[\{a, b, m\}, x] \& \& \operatorname{NeQ}[m, -1]$

Rule 5377

$\operatorname{Int}[\operatorname{ArcCsc}[u_], x_{\text{Symbol}}] \Rightarrow \operatorname{Simp}[x \operatorname{ArcCsc}[u], x] + \operatorname{Dist}[u / \operatorname{Sqrt}[u^2], \operatorname{Int}[\operatorname{SimplifyIntegrand}[x * (\operatorname{D}[u, x] / (u * \operatorname{Sqrt}[u^2 - 1])), x], x] /; \operatorname{InverseFunctionFreeQ}[u, x] \& \& \operatorname{!FunctionOfExponentialQ}[u, x]$

Rubi steps

$$\begin{aligned} \int \csc^{-1}(\sqrt{x}) \, dx &= x \csc^{-1}(\sqrt{x}) + \int \frac{1}{2\sqrt{-1+x}} \, dx \\ &= x \csc^{-1}(\sqrt{x}) + \frac{1}{2} \int \frac{1}{\sqrt{-1+x}} \, dx \\ &= \sqrt{-1+x} + x \csc^{-1}(\sqrt{x}) \end{aligned}$$

Mathematica [A]

time = 0.00, size = 16, normalized size = 1.00

$$\sqrt{-1+x} + x \csc^{-1}(\sqrt{x})$$

Antiderivative was successfully verified.

[In] `Integrate[ArcCsc[Sqrt[x]],x]`[Out] `Sqrt[-1 + x] + x*ArcCsc[Sqrt[x]]`**Maple [A]**

time = 0.20, size = 24, normalized size = 1.50

method	result	size
derivativedivides	$x \operatorname{arccsc}(\sqrt{x}) + \frac{\frac{x-1}{\sqrt{\frac{x-1}{x}}}}{\sqrt{x}}$	24
default	$x \operatorname{arccsc}(\sqrt{x}) + \frac{\frac{x-1}{\sqrt{\frac{x-1}{x}}}}{\sqrt{x}}$	24

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(arccsc(x^(1/2)),x,method=_RETURNVERBOSE)`[Out] `x*arccsc(x^(1/2))+1/((x-1)/x)^(1/2)/x^(1/2)*(x-1)`**Maxima [A]**

time = 0.25, size = 20, normalized size = 1.25

$$x \operatorname{arccsc}(\sqrt{x}) + \sqrt{x} \sqrt{-\frac{1}{x} + 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(x^(1/2)),x, algorithm="maxima")`[Out] `x*arccsc(sqrt(x)) + sqrt(x)*sqrt(-1/x + 1)`**Fricas [A]**

time = 0.34, size = 12, normalized size = 0.75

$$x \operatorname{arccsc}(\sqrt{x}) + \sqrt{x - 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(x^(1/2)),x, algorithm="fricas")`[Out] `x*arccsc(sqrt(x)) + sqrt(x - 1)`

Sympy [C] Result contains complex when optimal does not.
time = 2.60, size = 29, normalized size = 1.81

$$x \operatorname{acsc}(\sqrt{x}) + \frac{\begin{cases} 2\sqrt{x-1} & \text{for } |x| > 1 \\ 2i\sqrt{1-x} & \text{otherwise} \end{cases}}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(acsc(x**(1/2)),x)`

[Out] `x*acsc(sqrt(x)) + Piecewise((2*sqrt(x - 1), Abs(x) > 1), (2*I*sqrt(1 - x), True))/2`

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 41 vs. $2(12) = 24$.
time = 0.44, size = 41, normalized size = 2.56

$$x \arcsin\left(\frac{1}{\sqrt{x}}\right) + \frac{1}{2} \sqrt{x} \left(\sqrt{-\frac{1}{x} + 1} - 1 \right) - \frac{1}{2 \sqrt{x} \left(\sqrt{-\frac{1}{x} + 1} - 1 \right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(x^(1/2)),x, algorithm="giac")`

[Out] `x*arcsin(1/sqrt(x)) + 1/2*sqrt(x)*(sqrt(-1/x + 1) - 1) - 1/2/(sqrt(x)*(sqrt(-1/x + 1) - 1))`

Mupad [B]

time = 1.06, size = 20, normalized size = 1.25

$$x \arcsin\left(\frac{1}{\sqrt{x}}\right) + \sqrt{x} \sqrt{1 - \frac{1}{x}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(asin(1/x^(1/2)),x)`

[Out] `x*asin(1/x^(1/2)) + x^(1/2)*(1 - 1/x)^(1/2)`

3.6 $\int \frac{\csc^{-1}(\sqrt{x})}{x} dx$

Optimal. Leaf size=56

$$i \csc^{-1}(\sqrt{x})^2 - 2 \csc^{-1}(\sqrt{x}) \log\left(1 - e^{2i \csc^{-1}(\sqrt{x})}\right) + i \text{PolyLog}\left(2, e^{2i \csc^{-1}(\sqrt{x})}\right)$$

[Out] $I * \text{arccsc}(x^{1/2})^2 - 2 \text{arccsc}(x^{1/2}) * \ln(1 - (I/x^{1/2} + (1-1/x)^{1/2})^2) + I * \text{polylog}(2, (I/x^{1/2} + (1-1/x)^{1/2})^2)$

Rubi [A]

time = 0.06, antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.600, Rules used = {5327, 4721, 3798, 2221, 2317, 2438}

$$i \text{Li}_2\left(e^{2i \csc^{-1}(\sqrt{x})}\right) + i \csc^{-1}(\sqrt{x})^2 - 2 \csc^{-1}(\sqrt{x}) \log\left(1 - e^{2i \csc^{-1}(\sqrt{x})}\right)$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{ArcCsc}[\text{Sqrt}[x]]/x, x]$

[Out] $I * \text{ArcCsc}[\text{Sqrt}[x]]^2 - 2 * \text{ArcCsc}[\text{Sqrt}[x]] * \text{Log}[1 - E^{((2*I)*\text{ArcCsc}[\text{Sqrt}[x]])}] + I * \text{PolyLog}[2, E^{((2*I)*\text{ArcCsc}[\text{Sqrt}[x]])}]$

Rule 2221

```
Int[((F_)^((g_.)*(e_.) + (f_.)*(x_)))^(n_.)*(c_.) + (d_.)*(x_))^(m_.))/((a_.) + (b_.)*(F_)^((g_.)*(e_.) + (f_.)*(x_)))^(n_.)), x_Symbol] :> Simplify[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x]; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2317

```
Int[Log[(a_.) + (b_.)*(F_)^((e_.)*(c_.) + (d_.)*(x_)))^(n_.)], x_Symbol] :> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x]; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2438

```
Int[Log[(c_.)*(d_.) + (e_.)*(x_)^(n_.)]/(x_), x_Symbol] :> Simplify[-PolyLog[2, (-c)*e*x^n]/n, x]; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 3798

```
Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + Pi*(k_.) + (f_.)*(x_)], x_Symbol] :> Simplify[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Dist[2*I, Int[(c + d*x)^m
```

```
*E^(2*I*k*Pi)*(E^(2*I*(e + f*x))/(1 + E^(2*I*k*Pi)*E^(2*I*(e + f*x)))), x],  
x] /; FreeQ[{c, d, e, f}, x] && IntegerQ[4*k] && IGtQ[m, 0]
```

Rule 4721

```
Int[((a_.) + ArcSin[(c_.)*(x_)]*(b_._))^(n_.)/(x_), x_Symbol] :> Subst[Int[(  
a + b*x)^n*Cot[x], x, ArcSin[c*x]] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0]
```

Rule 5327

```
Int[((a_.) + ArcCsc[(c_.*(x_))*b_._]/(x_), x_Symbol] :> -Subst[Int[(a + b  
*ArcSin[x/c])/x, x, 1/x] /; FreeQ[{a, b, c}, x]
```

Rubi steps

$$\begin{aligned} \int \frac{\csc^{-1}(\sqrt{x})}{x} dx &= 2\text{Subst}\left(\int \frac{\csc^{-1}(x)}{x} dx, x, \sqrt{x}\right) \\ &= -\left(2\text{Subst}\left(\int \frac{\sin^{-1}(x)}{x} dx, x, \frac{1}{\sqrt{x}}\right)\right) \\ &= -\left(2\text{Subst}\left(\int x \cot(x) dx, x, \sin^{-1}\left(\frac{1}{\sqrt{x}}\right)\right)\right) \\ &= i \sin^{-1}\left(\frac{1}{\sqrt{x}}\right)^2 + 4i\text{Subst}\left(\int \frac{e^{2ix}x}{1-e^{2ix}} dx, x, \sin^{-1}\left(\frac{1}{\sqrt{x}}\right)\right) \\ &= i \sin^{-1}\left(\frac{1}{\sqrt{x}}\right)^2 - 2 \sin^{-1}\left(\frac{1}{\sqrt{x}}\right) \log\left(1 - e^{2i \sin^{-1}\left(\frac{1}{\sqrt{x}}\right)}\right) + 2\text{Subst}\left(\int \log(1 - e^{2ix}) dx, x, \sin^{-1}\left(\frac{1}{\sqrt{x}}\right)\right) \\ &= i \sin^{-1}\left(\frac{1}{\sqrt{x}}\right)^2 - 2 \sin^{-1}\left(\frac{1}{\sqrt{x}}\right) \log\left(1 - e^{2i \sin^{-1}\left(\frac{1}{\sqrt{x}}\right)}\right) - i\text{Subst}\left(\int \frac{\log(1-x)}{x} dx, x, \sin^{-1}\left(\frac{1}{\sqrt{x}}\right)\right) \\ &= i \sin^{-1}\left(\frac{1}{\sqrt{x}}\right)^2 - 2 \sin^{-1}\left(\frac{1}{\sqrt{x}}\right) \log\left(1 - e^{2i \sin^{-1}\left(\frac{1}{\sqrt{x}}\right)}\right) + i\text{Li}_2\left(e^{2i \sin^{-1}\left(\frac{1}{\sqrt{x}}\right)}\right) \end{aligned}$$

Mathematica [A]

time = 0.03, size = 54, normalized size = 0.96

$$i \left(\csc^{-1}(\sqrt{x}) \left(\csc^{-1}(\sqrt{x}) + 2i \log\left(1 - e^{2i \csc^{-1}(\sqrt{x})}\right) \right) + \text{PolyLog}\left(2, e^{2i \csc^{-1}(\sqrt{x})}\right) \right)$$

Antiderivative was successfully verified.

[In] Integrate[ArcCsc[Sqrt[x]]/x, x]

[Out] $I * (\text{ArcCsc}[\text{Sqrt}[x]] * (\text{ArcCsc}[\text{Sqrt}[x]] + (2*I) * \text{Log}[1 - E^{((2*I) * \text{ArcCsc}[\text{Sqrt}[x]])}) + \text{PolyLog}[2, E^{((2*I) * \text{ArcCsc}[\text{Sqrt}[x])}])$

Maple [A]

time = 0.60, size = 105, normalized size = 1.88

method	result
derivativedivides	$i \text{arccsc}(\sqrt{x})^2 - 2 \text{arccsc}(\sqrt{x}) \ln\left(1 - \frac{i}{\sqrt{x}} - \sqrt{1 - \frac{1}{x}}\right) - 2 \text{arccsc}(\sqrt{x}) \ln\left(1 + \frac{i}{\sqrt{x}} - \sqrt{1 - \frac{1}{x}}\right)$
default	$i \text{arccsc}(\sqrt{x})^2 - 2 \text{arccsc}(\sqrt{x}) \ln\left(1 - \frac{i}{\sqrt{x}} - \sqrt{1 - \frac{1}{x}}\right) - 2 \text{arccsc}(\sqrt{x}) \ln\left(1 + \frac{i}{\sqrt{x}} - \sqrt{1 - \frac{1}{x}}\right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(arccsc(x^(1/2))/x,x,method=_RETURNVERBOSE)`

[Out] $I * \text{arccsc}(x^{(1/2)})^2 - 2 * \text{arccsc}(x^{(1/2)}) * \ln(1 - I/x^{(1/2)} - (1 - 1/x)^{(1/2)}) - 2 * \text{arccsc}(x^{(1/2)}) * \ln(1 + I/x^{(1/2)} + (1 - 1/x)^{(1/2)}) + 2 * I * \text{polylog}(2, -I/x^{(1/2)} - (1 - 1/x)^{(1/2)}) + 2 * I * \text{polylog}(2, I/x^{(1/2)} + (1 - 1/x)^{(1/2)})$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(x^(1/2))/x,x, algorithm="maxima")`

[Out] `integrate(arccsc(sqrt(x))/x, x)`

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(x^(1/2))/x,x, algorithm="fricas")`

[Out] `integral(arccsc(sqrt(x))/x, x)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\text{acsc}(\sqrt{x})}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(acsc(x**(1/2))/x,x)`

[Out] `Integral(acsc(sqrt(x))/x, x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(x^(1/2))/x,x, algorithm="giac")`

[Out] `integrate(arccsc(sqrt(x))/x, x)`

Mupad [B]

time = 0.85, size = 42, normalized size = 0.75

$$\operatorname{polylog}\left(2, e^{\operatorname{asin}\left(\frac{1}{\sqrt{x}}\right) 2i}\right) 1i + \operatorname{asin}\left(\frac{1}{\sqrt{x}}\right)^2 1i - 2 \ln\left(1 - e^{\operatorname{asin}\left(\frac{1}{\sqrt{x}}\right) 2i}\right) \operatorname{asin}\left(\frac{1}{\sqrt{x}}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(asin(1/x^(1/2))/x,x)`

[Out] `polylog(2, exp(asin(1/x^(1/2))*2i))*1i + asin(1/x^(1/2))^2*1i - 2*log(1 - e^xp(asin(1/x^(1/2))*2i))*asin(1/x^(1/2))`

3.7 $\int \frac{\csc^{-1}(\sqrt{x})}{x^2} dx$

Optimal. Leaf size=38

$$-\frac{\sqrt{-1+x}}{2x} - \frac{\csc^{-1}(\sqrt{x})}{x} - \frac{1}{2}\text{ArcTan}(\sqrt{-1+x})$$

[Out] $-\text{arccsc}(x^{1/2})/x - 1/2\text{arctan}((-1+x)^{1/2}) - 1/2*(-1+x)^{1/2}/x$

Rubi [A]

time = 0.01, antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.500, Rules used = {5379, 12, 44, 65, 209}

$$-\frac{1}{2}\text{ArcTan}(\sqrt{x-1}) - \frac{\sqrt{x-1}}{2x} - \frac{\csc^{-1}(\sqrt{x})}{x}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{ArcCsc}[\text{Sqrt}[x]]/x^2, x]$

[Out] $-1/2*\text{Sqrt}[-1+x]/x - \text{ArcCsc}[\text{Sqrt}[x]]/x - \text{ArcTan}[\text{Sqrt}[-1+x]]/2$

Rule 12

```
Int[(a_)*(u_), x_Symbol] :> Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]
```

Rule 44

```
Int[((a_.) + (b_.)*(x_.))^(m_)*((c_.) + (d_.)*(x_.))^n_, x_Symbol] :> Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Dist[d*((m + n + 2)/((b*c - a*d)*(m + 1))), Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && ILtQ[m, -1] && !IntegerQ[n] && LtQ[n, 0]
```

Rule 65

```
Int[((a_.) + (b_.)*(x_.))^(m_)*((c_.) + (d_.)*(x_.))^n_, x_Symbol] :> With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^(p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 209

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
```

```
, 0] || GtQ[b, 0])
```

Rule 5379

```
Int[((a_.) + ArcCsc[u_]*(b_.))*((c_.) + (d_.)*(x_.))^(m_.), x_Symbol] :> Simp[((c + d*x)^(m + 1)*((a + b*ArcCsc[u])/(d*(m + 1))), x] + Dist[b*(u/(d*(m + 1)*Sqrt[u^2])), Int[SimplifyIntegrand[(c + d*x)^(m + 1)*(D[u, x]/(u*Sqrt[u^2 - 1])), x], x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1] && InverseFunctionFreeQ[u, x] && !FunctionOfQ[(c + d*x)^(m + 1), u, x] && !FunctionOfExponentialQ[u, x]
```

Rubi steps

$$\begin{aligned}
\int \frac{\csc^{-1}(\sqrt{x})}{x^2} dx &= -\frac{\csc^{-1}(\sqrt{x})}{x} - \int \frac{1}{2\sqrt{-1+x} x^2} dx \\
&= -\frac{\csc^{-1}(\sqrt{x})}{x} - \frac{1}{2} \int \frac{1}{\sqrt{-1+x} x^2} dx \\
&= -\frac{\sqrt{-1+x}}{2x} - \frac{\csc^{-1}(\sqrt{x})}{x} - \frac{1}{4} \int \frac{1}{\sqrt{-1+x} x} dx \\
&= -\frac{\sqrt{-1+x}}{2x} - \frac{\csc^{-1}(\sqrt{x})}{x} - \frac{1}{2} \text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \sqrt{-1+x}\right) \\
&= -\frac{\sqrt{-1+x}}{2x} - \frac{\csc^{-1}(\sqrt{x})}{x} - \frac{1}{2} \tan^{-1}(\sqrt{-1+x})
\end{aligned}$$

Mathematica [A]

time = 0.02, size = 32, normalized size = 0.84

$$-\frac{\sqrt{-1+x} + 2 \csc^{-1}(\sqrt{x}) - x \text{ArcSin}\left(\frac{1}{\sqrt{x}}\right)}{2x}$$

Antiderivative was successfully verified.

```
[In] Integrate[ArcCsc[Sqrt[x]]/x^2, x]
[Out] -1/2*(Sqrt[-1 + x] + 2*ArcCsc[Sqrt[x]] - x*ArcSin[1/Sqrt[x]])/x
```

Maple [A]

time = 0.21, size = 46, normalized size = 1.21

method	result	size
--------	--------	------

derivativedivides	$-\frac{\operatorname{arccsc}(\sqrt{x})}{x} + \frac{\sqrt{x-1} \left(\arctan\left(\frac{1}{\sqrt{x-1}}\right) x - \sqrt{x-1} \right)}{2 \sqrt{\frac{x-1}{x}} x^{\frac{3}{2}}} \quad 46$
default	$-\frac{\operatorname{arccsc}(\sqrt{x})}{x} + \frac{\sqrt{x-1} \left(\arctan\left(\frac{1}{\sqrt{x-1}}\right) x - \sqrt{x-1} \right)}{2 \sqrt{\frac{x-1}{x}} x^{\frac{3}{2}}} \quad 46$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(arccsc(x^(1/2))/x^2, x, method=_RETURNVERBOSE)`

[Out] $-\operatorname{arccsc}(x^{(1/2)})/x + 1/2*(x-1)^{(1/2)}*(\arctan(1/(x-1)^{(1/2})*x - (x-1)^{(1/2})/((x-1)/x)^{(1/2})/x^{(3/2)}$

Maxima [A]

time = 0.49, size = 51, normalized size = 1.34

$$\frac{\sqrt{x} \sqrt{-\frac{1}{x} + 1}}{2 (x (\frac{1}{x} - 1) - 1)} - \frac{\operatorname{arccsc}(\sqrt{x})}{x} - \frac{1}{2} \arctan\left(\sqrt{x} \sqrt{-\frac{1}{x} + 1}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(x^(1/2))/x^2, x, algorithm="maxima")`

[Out] $1/2*\sqrt{x}*\sqrt{-1/x + 1}/(x*(1/x - 1) - 1) - \operatorname{arccsc}(\sqrt{x})/x - 1/2*\arctan(\sqrt{x}*\sqrt{-1/x + 1})$

Fricas [A]

time = 0.37, size = 21, normalized size = 0.55

$$\frac{(x-2) \operatorname{arccsc}(\sqrt{x}) - \sqrt{x-1}}{2 x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(x^(1/2))/x^2, x, algorithm="fricas")`

[Out] $1/2*((x - 2)*\operatorname{arccsc}(\sqrt{x}) - \sqrt{x - 1})/x$

Sympy [C]

Result contains complex when optimal does not.

time = 14.03, size = 76, normalized size = 2.00

$$-\frac{\begin{cases} i \operatorname{acosh}\left(\frac{1}{\sqrt{x}}\right) - \frac{i}{\sqrt{x} \sqrt{-1 + \frac{1}{x}}} + \frac{i}{x^{\frac{3}{2}} \sqrt{-1 + \frac{1}{x}}} & \text{for } \frac{1}{|x|} > 1 \\ -\operatorname{asin}\left(\frac{1}{\sqrt{x}}\right) + \frac{\sqrt{1 - \frac{1}{x}}}{\sqrt{x}} & \text{otherwise} \end{cases}}{2} - \frac{\operatorname{acsc}(\sqrt{x})}{x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(acsc(x**(1/2))/x**2,x)`

[Out] `-Piecewise((I*acosh(1/sqrt(x)) - I/(sqrt(x)*sqrt(-1 + 1/x)) + I/(x**3/2)*sqrt(-1 + 1/x)), 1/Abs(x) > 1), (-asin(1/sqrt(x)) + sqrt(1 - 1/x)/sqrt(x), True))/2 - acsc(sqrt(x))/x`

Giac [A]

time = 0.43, size = 32, normalized size = 0.84

$$-\left(\frac{1}{x} - 1\right) \arcsin\left(\frac{1}{\sqrt{x}}\right) - \frac{\sqrt{-\frac{1}{x} + 1}}{2\sqrt{x}} - \frac{1}{2} \arcsin\left(\frac{1}{\sqrt{x}}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(x^(1/2))/x^2,x, algorithm="giac")`

[Out] `-(1/x - 1)*arcsin(1/sqrt(x)) - 1/2*sqrt(-1/x + 1)/sqrt(x) - 1/2*arcsin(1/sqrt(x))`

Mupad [B]

time = 0.69, size = 28, normalized size = 0.74

$$-\frac{\sqrt{1 - \frac{1}{x}}}{2\sqrt{x}} - \frac{\arcsin\left(\frac{1}{\sqrt{x}}\right) (\frac{2}{x} - 1)}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(asin(1/x^(1/2))/x^2,x)`

[Out] `- (1 - 1/x)^(1/2)/(2*x^(1/2)) - (asin(1/x^(1/2))*(2/x - 1))/2`

3.8 $\int \frac{\csc^{-1}(\sqrt{x})}{x^3} dx$

Optimal. Leaf size=54

$$-\frac{\sqrt{-1+x}}{8x^2} - \frac{3\sqrt{-1+x}}{16x} - \frac{\csc^{-1}(\sqrt{x})}{2x^2} - \frac{3}{16}\text{ArcTan}(\sqrt{-1+x})$$

[Out] $-1/2*\text{arccsc}(x^{(1/2)})/x^2 - 3/16*\text{arctan}((-1+x)^{(1/2)}) - 1/8*(-1+x)^{(1/2)}/x^2 - 3/16*(-1+x)^{(1/2)}/x$

Rubi [A]

time = 0.02, antiderivative size = 54, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.500, Rules used = {5379, 12, 44, 65, 209}

$$-\frac{3}{16}\text{ArcTan}(\sqrt{x-1}) - \frac{\sqrt{x-1}}{8x^2} - \frac{\csc^{-1}(\sqrt{x})}{2x^2} - \frac{3\sqrt{x-1}}{16x}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{ArcCsc}[\text{Sqrt}[x]]/x^3, x]$

[Out] $-1/8*\text{Sqrt}[-1+x]/x^2 - (3*\text{Sqrt}[-1+x])/(16*x) - \text{ArcCsc}[\text{Sqrt}[x]]/(2*x^2) - (3*\text{ArcTan}[\text{Sqrt}[-1+x]])/16$

Rule 12

```
Int[(a_)*(u_), x_Symbol] :> Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]
```

Rule 44

```
Int[((a_.) + (b_.)*(x_.))^(m_)*((c_.) + (d_.)*(x_.))^(n_), x_Symbol] :> Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Dist[d*((m + n + 2)/((b*c - a*d)*(m + 1))), Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && ILtQ[m, -1] && !IntegerQ[n] && LtQ[n, 0]
```

Rule 65

```
Int[((a_.) + (b_.)*(x_.))^(m_)*((c_.) + (d_.)*(x_.))^(n_), x_Symbol] :> With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 209

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

Rule 5379

```
Int[((a_) + ArcCsc[u_]*(b_))*(c_*) + (d_*)*(x_)^(m_), x_Symbol] :> Simplify[Int[(c + d*x)^(m + 1)*(a + b*ArcCsc[u])/(d*(m + 1)), x] + Dist[b*(u/(d*(m + 1)*Sqrt[u^2])), Int[Simplify[Integrand[(c + d*x)^(m + 1)*(D[u, x]/(u*Sqrt[u^2 - 1])), x], x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1] && InverseFunctionFreeQ[u, x] && !FunctionOfQ[(c + d*x)^(m + 1), u, x] && !FunctionOfExponentialQ[u, x]]
```

Rubi steps

$$\begin{aligned}
\int \frac{\csc^{-1}(\sqrt{x})}{x^3} dx &= -\frac{\csc^{-1}(\sqrt{x})}{2x^2} - \frac{1}{2} \int \frac{1}{2\sqrt{-1+x} x^3} dx \\
&= -\frac{\csc^{-1}(\sqrt{x})}{2x^2} - \frac{1}{4} \int \frac{1}{\sqrt{-1+x} x^3} dx \\
&= -\frac{\sqrt{-1+x}}{8x^2} - \frac{\csc^{-1}(\sqrt{x})}{2x^2} - \frac{3}{16} \int \frac{1}{\sqrt{-1+x} x^2} dx \\
&= -\frac{\sqrt{-1+x}}{8x^2} - \frac{3\sqrt{-1+x}}{16x} - \frac{\csc^{-1}(\sqrt{x})}{2x^2} - \frac{3}{32} \int \frac{1}{\sqrt{-1+x} x} dx \\
&= -\frac{\sqrt{-1+x}}{8x^2} - \frac{3\sqrt{-1+x}}{16x} - \frac{\csc^{-1}(\sqrt{x})}{2x^2} - \frac{3}{16} \text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \sqrt{-1+x}\right) \\
&= -\frac{\sqrt{-1+x}}{8x^2} - \frac{3\sqrt{-1+x}}{16x} - \frac{\csc^{-1}(\sqrt{x})}{2x^2} - \frac{3}{16} \tan^{-1}(\sqrt{-1+x})
\end{aligned}$$

Mathematica [A]

time = 0.04, size = 55, normalized size = 1.02

$$\left(-\frac{1}{8x^{3/2}} - \frac{3}{16\sqrt{x}}\right) \sqrt{\frac{-1+x}{x}} - \frac{\csc^{-1}(\sqrt{x})}{2x^2} + \frac{3}{16} \text{ArcSin}\left(\frac{1}{\sqrt{x}}\right)$$

Antiderivative was successfully verified.

[In] `Integrate[ArcCsc[Sqrt[x]]/x^3,x]`

[Out] $(-1/8*1/x^{(3/2)} - 3/(16*Sqrt[x]))*Sqrt[(-1 + x)/x] - \text{ArcCsc}[Sqrt[x]]/(2*x^2) + (3*\text{ArcSin}[1/Sqrt[x]])/16$

Maple [A]

time = 0.20, size = 57, normalized size = 1.06

method	result	size
derivativedivides	$-\frac{\operatorname{arccsc}(\sqrt{x})}{2x^2} + \frac{\sqrt{x-1} \left(3 \arctan\left(\frac{1}{\sqrt{x-1}}\right) x^2 - 3 \sqrt{x-1} x - 2 \sqrt{x-1}\right)}{16 \sqrt{\frac{x-1}{x}} x^{\frac{5}{2}}}$	57
default	$-\frac{\operatorname{arccsc}(\sqrt{x})}{2x^2} + \frac{\sqrt{x-1} \left(3 \arctan\left(\frac{1}{\sqrt{x-1}}\right) x^2 - 3 \sqrt{x-1} x - 2 \sqrt{x-1}\right)}{16 \sqrt{\frac{x-1}{x}} x^{\frac{5}{2}}}$	57

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(arccsc(x^(1/2))/x^3,x,method=_RETURNVERBOSE)`

[Out]
$$-1/2*\operatorname{arccsc}(x^{(1/2)})/x^2 + 1/16*(x-1)^{(1/2)}*(3*\arctan(1/(x-1)^{(1/2})*x^2 - 3*(x-1)^{(1/2})*x - 2*(x-1)^{(1/2}))/((x-1)/x)^{(1/2})*x^{(5/2)}$$

Maxima [B] Leaf count of result is larger than twice the leaf count of optimal. 80 vs. $2(38) = 76$.

time = 0.49, size = 80, normalized size = 1.48

$$-\frac{3 x^{\frac{3}{2}} \left(-\frac{1}{x} + 1\right)^{\frac{3}{2}} + 5 \sqrt{x} \sqrt{-\frac{1}{x} + 1}}{16 \left(x^2 \left(\frac{1}{x} - 1\right)^2 - 2 x \left(\frac{1}{x} - 1\right) + 1\right)} - \frac{\operatorname{arccsc}(\sqrt{x})}{2 x^2} - \frac{3}{16} \arctan\left(\sqrt{x} \sqrt{-\frac{1}{x} + 1}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(x^(1/2))/x^3,x, algorithm="maxima")`

[Out]
$$-1/16*(3*x^{(3/2)}*(-1/x + 1)^{(3/2}) + 5*sqrt(x)*sqrt(-1/x + 1))/(x^2*(1/x - 1)^2 - 2*x*(1/x - 1) + 1) - 1/2*\operatorname{arccsc}(\sqrt{x})/x^2 - 3/16*\arctan(\sqrt{x})*sqrt(-1/x + 1))$$

Fricas [A]

time = 0.36, size = 30, normalized size = 0.56

$$\frac{(3 x^2 - 8) \operatorname{arccsc}(\sqrt{x}) - (3 x + 2) \sqrt{x-1}}{16 x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(x^(1/2))/x^3,x, algorithm="fricas")`

[Out]
$$1/16*((3*x^2 - 8)*\operatorname{arccsc}(\sqrt{x}) - (3*x + 2)*sqrt(x - 1))/x^2$$

Sympy [C] Result contains complex when optimal does not.

time = 45.17, size = 146, normalized size = 2.70

$$\begin{aligned}
 & -\frac{\begin{cases} \frac{3i \operatorname{acosh}\left(\frac{1}{\sqrt{x}}\right)}{4} - \frac{3i}{4\sqrt{x}\sqrt{-1+\frac{1}{x}}} + \frac{i}{4x^{\frac{3}{2}}\sqrt{-1+\frac{1}{x}}} + \frac{i}{2x^{\frac{5}{2}}\sqrt{-1+\frac{1}{x}}} & \text{for } \frac{1}{|x|} > 1 \\ -\frac{3\operatorname{asin}\left(\frac{1}{\sqrt{x}}\right)}{4} + \frac{3}{4\sqrt{x}\sqrt{1-\frac{1}{x}}} - \frac{1}{4x^{\frac{3}{2}}\sqrt{1-\frac{1}{x}}} - \frac{1}{2x^{\frac{5}{2}}\sqrt{1-\frac{1}{x}}} & \text{otherwise} \end{cases}}{4} - \frac{\operatorname{acsc}\left(\sqrt{x}\right)}{2x^2}
 \end{aligned}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(acsc(x**(1/2))/x**3,x)`

[Out] `-Piecewise((3*I*acosh(1/sqrt(x))/4 - 3*I/(4*sqrt(x)*sqrt(-1 + 1/x)) + I/(4*x**3/2)*sqrt(-1 + 1/x)) + I/(2*x**5/2)*sqrt(-1 + 1/x)), 1/Abs(x) > 1), (-3*asin(1/sqrt(x))/4 + 3/(4*sqrt(x)*sqrt(1 - 1/x)) - 1/(4*x**3/2)*sqrt(1 - 1/x)) - 1/(2*x**5/2)*sqrt(1 - 1/x)), True))/4 - acsc(sqrt(x))/(2*x**2)`

Giac [A]

time = 0.44, size = 59, normalized size = 1.09

$$-\frac{1}{2} \left(\frac{1}{x} - 1\right)^2 \arcsin\left(\frac{1}{\sqrt{x}}\right) - \left(\frac{1}{x} - 1\right) \arcsin\left(\frac{1}{\sqrt{x}}\right) + \frac{\left(-\frac{1}{x} + 1\right)^{\frac{3}{2}}}{8\sqrt{x}} - \frac{5\sqrt{-\frac{1}{x} + 1}}{16\sqrt{x}} - \frac{5}{16} \arcsin\left(\frac{1}{\sqrt{x}}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(x^(1/2))/x^3,x, algorithm="giac")`

[Out] `-1/2*(1/x - 1)^2*arcsin(1/sqrt(x)) - (1/x - 1)*arcsin(1/sqrt(x)) + 1/8*(-1/x + 1)^(3/2)/sqrt(x) - 5/16*sqrt(-1/x + 1)/sqrt(x) - 5/16*arcsin(1/sqrt(x))`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\operatorname{asin}\left(\frac{1}{\sqrt{x}}\right)}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(asin(1/x^(1/2))/x^3,x)`

[Out] `int(asin(1/x^(1/2))/x^3, x)`

$$3.9 \quad \int x^2 \csc^{-1} \left(\frac{a}{x} \right) dx$$

Optimal. Leaf size=56

$$\frac{1}{3}a^3\sqrt{1 - \frac{x^2}{a^2}} - \frac{1}{9}a^3\left(1 - \frac{x^2}{a^2}\right)^{3/2} + \frac{1}{3}x^3\text{ArcSin}\left(\frac{x}{a}\right)$$

[Out] $-1/9*a^3*(1-x^2/a^2)^(3/2)+1/3*x^3*\text{arcsin}(x/a)+1/3*a^3*(1-x^2/a^2)^(1/2)$

Rubi [A]

time = 0.03, antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.400, Rules used = {5373, 4723, 272, 45}

$$-\frac{1}{9}a^3\left(1 - \frac{x^2}{a^2}\right)^{3/2} + \frac{1}{3}a^3\sqrt{1 - \frac{x^2}{a^2}} + \frac{1}{3}x^3\text{ArcSin}\left(\frac{x}{a}\right)$$

Antiderivative was successfully verified.

[In] Int[x^2*ArcCsc[a/x], x]

[Out] $(a^3*\text{Sqrt}[1 - x^2/a^2])/3 - (a^3*(1 - x^2/a^2)^(3/2))/9 + (x^3*\text{ArcSin}[x/a])/3$

Rule 45

```
Int[((a_.) + (b_ .)*(x_ ))^(m_ .)*((c_ .) + (d_ .)*(x_ ))^(n_ .), x_Symbol] :> Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && Le
Q[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])
```

Rule 272

```
Int[(x_)^(m_ .)*((a_) + (b_ .)*(x_)^(n_ ))^(p_ ), x_Symbol] :> Dist[1/n, Subst[
Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 4723

```
Int[((a_.) + ArcSin[(c_ .)*(x_ )]*(b_ .))^(n_ .)*((d_ .)*(x_ ))^(m_ .), x_Symbol]
:> Simp[(d*x)^(m + 1)*((a + b*ArcSin[c*x])^n/(d*(m + 1))), x] - Dist[b*c*(n/(d*(m + 1))), Int[(d*x)^(m + 1)*((a + b*ArcSin[c*x])^(n - 1)/Sqrt[1 - c^2*x^2]), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]
```

Rule 5373

```
Int[ArcCsc[(c_ .)/((a_ .) + (b_ .)*(x_)^(n_ .))]^(m_ .)*(u_ .), x_Symbol] :> Int[
u*ArcSin[a/c + b*(x^n/c)]^m, x] /; FreeQ[{a, b, c, n, m}, x]
```

Rubi steps

$$\begin{aligned}
\int x^2 \csc^{-1} \left(\frac{a}{x} \right) dx &= \int x^2 \sin^{-1} \left(\frac{x}{a} \right) dx \\
&= \frac{1}{3} x^3 \sin^{-1} \left(\frac{x}{a} \right) - \frac{\int \frac{x^3}{\sqrt{1 - \frac{x^2}{a^2}}} dx}{3a} \\
&= \frac{1}{3} x^3 \sin^{-1} \left(\frac{x}{a} \right) - \frac{\text{Subst} \left(\int \frac{x}{\sqrt{1 - \frac{x}{a^2}}} dx, x, x^2 \right)}{6a} \\
&= \frac{1}{3} x^3 \sin^{-1} \left(\frac{x}{a} \right) - \frac{\text{Subst} \left(\int \left(\frac{a^2}{\sqrt{1 - \frac{x}{a^2}}} - a^2 \sqrt{1 - \frac{x}{a^2}} \right) dx, x, x^2 \right)}{6a} \\
&= \frac{1}{3} x^3 \sin^{-1} \left(\frac{x}{a} \right) - \frac{1}{3} a^3 \sqrt{1 - \frac{x^2}{a^2}} - \frac{1}{9} a^3 \left(1 - \frac{x^2}{a^2} \right)^{3/2} + \frac{1}{3} x^3 \sin^{-1} \left(\frac{x}{a} \right)
\end{aligned}$$

Mathematica [A]

time = 0.02, size = 42, normalized size = 0.75

$$\frac{1}{9} a (2a^2 + x^2) \sqrt{1 - \frac{x^2}{a^2}} + \frac{1}{3} x^3 \csc^{-1} \left(\frac{a}{x} \right)$$

Antiderivative was successfully verified.

[In] `Integrate[x^2*ArcCsc[a/x], x]`[Out] `(a*(2*a^2 + x^2)*Sqrt[1 - x^2/a^2])/9 + (x^3*ArcCsc[a/x])/3`Maple [A]

time = 0.46, size = 66, normalized size = 1.18

method	result	size
derivativedivides	$-a^3 \left(-\frac{x^3 \text{arccsc}(\frac{a}{x})}{3a^3} - \frac{\left(\frac{a^2}{x^2}-1\right)\left(\frac{2a^2}{x^2}+1\right)x^4}{9\sqrt{\frac{\left(\frac{a^2}{x^2}-1\right)x^2}{a^2}} a^4} \right)$	66

default	$-a^3 \left(-\frac{x^3 \operatorname{arccsc}(\frac{a}{x})}{3a^3} - \frac{\left(\frac{a^2}{x^2}-1\right)\left(\frac{2a^2}{x^2}+1\right)x^4}{9\sqrt{\frac{\left(\frac{a^2}{x^2}-1\right)x^2}{a^2}} a^4} \right)$	66
---------	---	----

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x^2*arccsc(a/x),x,method=_RETURNVERBOSE)`

[Out] $-a^3(-1/3x^3/a^3\operatorname{arccsc}(a/x)-1/9(a^2/x^2-1)(2a^2/x^2+1)/((a^2/x^2-1)*x^2/a^2)^{(1/2)}*x^4/a^4)$

Maxima [A]

time = 0.48, size = 54, normalized size = 0.96

$$\frac{1}{3} x^3 \operatorname{arccsc}\left(\frac{a}{x}\right) + \frac{2 a^4 \sqrt{-\frac{x^2}{a^2} + 1} + a^2 x^2 \sqrt{-\frac{x^2}{a^2} + 1}}{9 a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^2*arccsc(a/x),x, algorithm="maxima")`

[Out] $\frac{1}{3}x^3\operatorname{arccsc}\left(\frac{a}{x}\right) + \frac{1}{9}(2a^4\sqrt{-x^2/a^2 + 1} + a^2x^2\sqrt{-x^2/a^2 + 1})/a$

Fricas [A]

time = 0.34, size = 39, normalized size = 0.70

$$\frac{1}{3} x^3 \operatorname{arccsc}\left(\frac{a}{x}\right) + \frac{1}{9} (2 a^2 x + x^3) \sqrt{\frac{a^2 - x^2}{x^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^2*arccsc(a/x),x, algorithm="fricas")`

[Out] $\frac{1}{3}x^3\operatorname{arccsc}\left(\frac{a}{x}\right) + \frac{1}{9}(2a^2x + x^3)\sqrt{(a^2 - x^2)/x^2}$

Sympy [A]

time = 0.13, size = 51, normalized size = 0.91

$$\begin{cases} \frac{2a^3\sqrt{1-\frac{x^2}{a^2}}}{9} + \frac{ax^2\sqrt{1-\frac{x^2}{a^2}}}{9} + \frac{x^3\operatorname{acsc}(\frac{a}{x})}{3} & \text{for } a \neq 0 \\ \infty x^3 & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x**2*acsc(a/x),x)`

[Out] Piecewise((2*a**3*sqrt(1 - x**2/a**2)/9 + a*x**2*sqrt(1 - x**2/a**2)/9 + x**3*acsc(a/x)/3, Ne(a, 0)), (zoo*x**3, True))

Giac [A]

time = 0.45, size = 68, normalized size = 1.21

$$\frac{1}{3} a^2 x \left(\frac{x^2}{a^2} - 1 \right) \arcsin \left(\frac{x}{a} \right) - \frac{1}{9} a^3 \left(-\frac{x^2}{a^2} + 1 \right)^{\frac{3}{2}} + \frac{1}{3} a^2 x \arcsin \left(\frac{x}{a} \right) + \frac{1}{3} a^3 \sqrt{-\frac{x^2}{a^2} + 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2*arccsc(a/x),x, algorithm="giac")

[Out] $\frac{1}{3} a^2 x^2 \arccsc \left(\frac{x}{a} \right) - \frac{1}{9} a^3 (-x^2/a^2 + 1)^{(3/2)} + \frac{1}{3} a^3 x^2 \arcsin \left(\frac{x}{a} \right) + \frac{1}{3} a^3 \sqrt{-x^2/a^2 + 1}$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\begin{cases} \frac{x^3 \arcsin(\frac{x}{a})}{3} + \frac{\sqrt{a^2 - x^2} (2a^2 + x^2)}{9} & \text{if } 0 < a \\ \int x^2 \arcsin \left(\frac{x}{a} \right) dx & \text{if } -0 < a \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^2*asin(x/a),x)

[Out] piecewise(0 < a, (x^3*asin(x/a))/3 + ((a^2 - x^2)^(1/2)*(2*a^2 + x^2))/9, ~ 0 < a, int(x^2*asin(x/a), x))

3.10 $\int x \csc^{-1} \left(\frac{a}{x} \right) dx$

Optimal. Leaf size=47

$$\frac{1}{4}ax\sqrt{1 - \frac{x^2}{a^2}} - \frac{1}{4}a^2\text{ArcSin}\left(\frac{x}{a}\right) + \frac{1}{2}x^2\text{ArcSin}\left(\frac{x}{a}\right)$$

[Out] $-1/4*a^2*\arcsin(x/a)+1/2*x^2*\arcsin(x/a)+1/4*a*x*(1-x^2/a^2)^(1/2)$

Rubi [A]

time = 0.01, antiderivative size = 47, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.500, Rules used = {5373, 4723, 327, 222}

$$-\frac{1}{4}a^2\text{ArcSin}\left(\frac{x}{a}\right) + \frac{1}{4}ax\sqrt{1 - \frac{x^2}{a^2}} + \frac{1}{2}x^2\text{ArcSin}\left(\frac{x}{a}\right)$$

Antiderivative was successfully verified.

[In] $\text{Int}[x*\text{ArcCsc}[a/x], x]$

[Out] $(a*x*\text{Sqrt}[1 - x^2/a^2])/4 - (a^2*\text{ArcSin}[x/a])/4 + (x^2*\text{ArcSin}[x/a])/2$

Rule 222

```
Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] :> Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]
```

Rule 327

```
Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] :> Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 4723

```
Int[((a_) + ArcSin[(c_)*(x_)]*(b_))^(n_)*((d_)*(x_))^(m_), x_Symbol] :> Simp[(d*x)^(m + 1)*((a + b*ArcSin[c*x])^n/(d*(m + 1))), x] - Dist[b*c*(n/(d*(m + 1))), Int[(d*x)^(m + 1)*((a + b*ArcSin[c*x])^(n - 1)/Sqrt[1 - c^2*x^2]), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]
```

Rule 5373

```
Int[ArcCsc[(c_)/((a_) + (b_)*(x_)^(n_))]^(m_)*(u_), x_Symbol] :> Int[u*ArcSin[a/c + b*(x^n/c)]^m, x] /; FreeQ[{a, b, c, n, m}, x]
```

Rubi steps

$$\begin{aligned}
\int x \csc^{-1} \left(\frac{a}{x} \right) dx &= \int x \sin^{-1} \left(\frac{x}{a} \right) dx \\
&= \frac{1}{2} x^2 \sin^{-1} \left(\frac{x}{a} \right) - \frac{\int \frac{x^2}{\sqrt{1 - \frac{x^2}{a^2}}} dx}{2a} \\
&= \frac{1}{4} ax \sqrt{1 - \frac{x^2}{a^2}} + \frac{1}{2} x^2 \sin^{-1} \left(\frac{x}{a} \right) - \frac{1}{4} a \int \frac{1}{\sqrt{1 - \frac{x^2}{a^2}}} dx \\
&= \frac{1}{4} ax \sqrt{1 - \frac{x^2}{a^2}} - \frac{1}{4} a^2 \sin^{-1} \left(\frac{x}{a} \right) + \frac{1}{2} x^2 \sin^{-1} \left(\frac{x}{a} \right)
\end{aligned}$$

Mathematica [A]

time = 0.02, size = 44, normalized size = 0.94

$$\frac{1}{4} \left(ax \sqrt{1 - \frac{x^2}{a^2}} + 2x^2 \csc^{-1} \left(\frac{a}{x} \right) - a^2 \text{ArcSin} \left(\frac{x}{a} \right) \right)$$

Antiderivative was successfully verified.

[In] `Integrate[x*ArcCsc[a/x], x]`[Out] `(a*x*.Sqrt[1 - x^2/a^2] + 2*x^2*ArcCsc[a/x] - a^2*ArcSin[x/a])/4`Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 89 vs. 2(39) = 78.

time = 0.49, size = 90, normalized size = 1.91

method	result	size
--------	--------	------

	$-a^2 \left(-\frac{x^2 \operatorname{arccsc}(\frac{a}{x})}{2a^2} - \frac{\sqrt{\frac{a^2}{x^2} - 1} \left(-\frac{\arctan\left(\frac{1}{\sqrt{\frac{a^2}{x^2} - 1}}\right) a^2}{4 \sqrt{\frac{(a^2 - 1)x^2}{a^2}}} + \sqrt{\frac{a^2}{x^2} - 1} \right) x^3}{4 \sqrt{\frac{(a^2 - 1)x^2}{a^2}} a^3} \right)$	90
derivative divides	$-a^2 \left(-\frac{x^2 \operatorname{arccsc}(\frac{a}{x})}{2a^2} - \frac{\sqrt{\frac{a^2}{x^2} - 1} \left(-\frac{\arctan\left(\frac{1}{\sqrt{\frac{a^2}{x^2} - 1}}\right) a^2}{4 \sqrt{\frac{(a^2 - 1)x^2}{a^2}}} + \sqrt{\frac{a^2}{x^2} - 1} \right) x^3}{4 \sqrt{\frac{(a^2 - 1)x^2}{a^2}} a^3} \right)$	90
default	$-a^2 \left(-\frac{x^2 \operatorname{arccsc}(\frac{a}{x})}{2a^2} - \frac{\sqrt{\frac{a^2}{x^2} - 1} \left(-\frac{\arctan\left(\frac{1}{\sqrt{\frac{a^2}{x^2} - 1}}\right) a^2}{4 \sqrt{\frac{(a^2 - 1)x^2}{a^2}}} + \sqrt{\frac{a^2}{x^2} - 1} \right) x^3}{4 \sqrt{\frac{(a^2 - 1)x^2}{a^2}} a^3} \right)$	90

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x*arccsc(a/x),x,method=_RETURNVERBOSE)`

[Out] `-a^2*(-1/2*x^2/a^2*arccsc(a/x)-1/4*(a^2/x^2-1)^(1/2)*(-arctan(1/(a^2/x^2-1)^(1/2))/x^2*a^2+(a^2/x^2-1)^(1/2))/((a^2/x^2-1)*x^2/a^2)^(1/2)*x^3/a^3)`

Maxima [A]

time = 0.46, size = 46, normalized size = 0.98

$$\frac{1}{2} x^2 \operatorname{arccsc}\left(\frac{a}{x}\right) - \frac{a^3 \arcsin\left(\frac{x}{a}\right) - a^2 x \sqrt{-\frac{x^2}{a^2} + 1}}{4 a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x*arccsc(a/x),x, algorithm="maxima")`

[Out] `1/2*x^2*arccsc(a/x) - 1/4*(a^3*arcsin(x/a) - a^2*x*sqrt(-x^2/a^2 + 1))/a`

Fricas [A]

time = 0.34, size = 38, normalized size = 0.81

$$\frac{1}{4} x^2 \sqrt{\frac{a^2 - x^2}{x^2}} - \frac{1}{4} (a^2 - 2x^2) \operatorname{arccsc}\left(\frac{a}{x}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x*arccsc(a/x),x, algorithm="fricas")`[Out] `1/4*x^2*sqrt((a^2 - x^2)/x^2) - 1/4*(a^2 - 2*x^2)*arccsc(a/x)`**Sympy [A]**

time = 0.09, size = 41, normalized size = 0.87

$$\begin{cases} -\frac{a^2 \operatorname{acsc}\left(\frac{a}{x}\right)}{4} + \frac{ax \sqrt{1 - \frac{x^2}{a^2}}}{4} + \frac{x^2 \operatorname{acsc}\left(\frac{a}{x}\right)}{2} & \text{for } a \neq 0 \\ \infty x^2 & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x*acsc(a/x),x)`[Out] `Piecewise((-a**2*acsc(a/x)/4 + a*x*sqrt(1 - x**2/a**2)/4 + x**2*acsc(a/x)/2, Ne(a, 0)), (zoo*x**2, True))`**Giac [A]**

time = 0.41, size = 48, normalized size = 1.02

$$\frac{1}{2} a^2 \left(\frac{x^2}{a^2} - 1 \right) \arcsin\left(\frac{x}{a}\right) + \frac{1}{4} a^2 \arcsin\left(\frac{x}{a}\right) + \frac{1}{4} ax \sqrt{-\frac{x^2}{a^2} + 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x*arccsc(a/x),x, algorithm="giac")`[Out] `1/2*a^2*(x^2/a^2 - 1)*arcsin(x/a) + 1/4*a^2*arcsin(x/a) + 1/4*a*x*sqrt(-x^2/a^2 + 1)`**Mupad [B]**

time = 0.61, size = 38, normalized size = 0.81

$$\frac{a^2 \sin\left(\frac{x}{a}\right) \left(\frac{2x^2}{a^2} - 1\right)}{4} + \frac{ax \sqrt{1 - \frac{x^2}{a^2}}}{4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x*asin(x/a),x)`[Out] `(a^2*asin(x/a)*((2*x^2)/a^2 - 1))/4 + (a*x*(1 - x^2/a^2)^(1/2))/4`

3.11 $\int \csc^{-1} \left(\frac{a}{x} \right) dx$

Optimal. Leaf size=25

$$a \sqrt{1 - \frac{x^2}{a^2}} + x \operatorname{ArcSin} \left(\frac{x}{a} \right)$$

[Out] $x \operatorname{arcsin}(x/a) + a * (1 - x^2/a^2)^{(1/2)}$

Rubi [A]

time = 0.01, antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 6, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.500, Rules used = {5373, 4715, 267}

$$a \sqrt{1 - \frac{x^2}{a^2}} + x \operatorname{ArcSin} \left(\frac{x}{a} \right)$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[\operatorname{ArcCsc}[a/x], x]$

[Out] $a * \operatorname{Sqrt}[1 - x^2/a^2] + x * \operatorname{ArcSin}[x/a]$

Rule 267

```
Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] :> Simp[(a + b*x^n)^(p + 1)/(b*n*(p + 1)), x] /; FreeQ[{a, b, m, n, p}, x] && EqQ[m, n - 1] && NeQ[p, -1]
```

Rule 4715

```
Int[((a_) + ArcSin[(c_)*(x_)]*(b_))^(n_), x_Symbol] :> Simp[x*(a + b*ArcSin[c*x])^n, x] - Dist[b*c*n, Int[x*((a + b*ArcSin[c*x])^(n - 1))/Sqrt[1 - c^2*x^2]], x, x] /; FreeQ[{a, b, c}, x] && GtQ[n, 0]
```

Rule 5373

```
Int[ArcCsc[(c_.)/((a_.) + (b_)*(x_)^(n_.))]^(m_)*(u_), x_Symbol] :> Int[u*ArcSin[a/c + b*(x^n/c)]^m, x] /; FreeQ[{a, b, c, n, m}, x]
```

Rubi steps

$$\begin{aligned}
\int \csc^{-1} \left(\frac{a}{x} \right) dx &= \int \sin^{-1} \left(\frac{x}{a} \right) dx \\
&= x \sin^{-1} \left(\frac{x}{a} \right) - \frac{\int \frac{x}{\sqrt{1 - \frac{x^2}{a^2}}} dx}{a} \\
&= a \sqrt{1 - \frac{x^2}{a^2}} + x \sin^{-1} \left(\frac{x}{a} \right)
\end{aligned}$$

Mathematica [A]

time = 0.01, size = 25, normalized size = 1.00

$$a \sqrt{1 - \frac{x^2}{a^2}} + x \csc^{-1} \left(\frac{a}{x} \right)$$

Antiderivative was successfully verified.

[In] Integrate[ArcCsc[a/x], x]

[Out] a*.Sqrt[1 - x^2/a^2] + x*ArcCsc[a/x]

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 51 vs. 2(23) = 46.

time = 0.49, size = 52, normalized size = 2.08

method	result	size
derivativedivides	$-a \left(-\frac{x \operatorname{arccsc} \left(\frac{a}{x} \right)}{a} - \frac{x^2 \left(\frac{a^2}{x^2} - 1 \right)}{\sqrt{\frac{\left(\frac{a^2}{x^2} - 1 \right) x^2}{a^2}} a^2} \right)$	52
default	$-a \left(-\frac{x \operatorname{arccsc} \left(\frac{a}{x} \right)}{a} - \frac{x^2 \left(\frac{a^2}{x^2} - 1 \right)}{\sqrt{\frac{\left(\frac{a^2}{x^2} - 1 \right) x^2}{a^2}} a^2} \right)$	52

Verification of antiderivative is not currently implemented for this CAS.

[In] int(arccsc(a/x), x, method=_RETURNVERBOSE)

[Out] -a*(-x/a*arccsc(a/x)-1/((a^2/x^2-1)*x^2/a^2)^(1/2)*x^2/a^2*(a^2/x^2-1))

Maxima [A]

time = 0.26, size = 23, normalized size = 0.92

$$x \operatorname{arccsc} \left(\frac{a}{x} \right) + a \sqrt{-\frac{x^2}{a^2} + 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(a/x),x, algorithm="maxima")`

[Out] $x \operatorname{arccsc}(a/x) + a \sqrt{-x^2/a^2 + 1}$

Fricas [A]

time = 0.34, size = 26, normalized size = 1.04

$$x \operatorname{arccsc} \left(\frac{a}{x} \right) + x \sqrt{\frac{a^2 - x^2}{x^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(a/x),x, algorithm="fricas")`

[Out] $x \operatorname{arccsc}(a/x) + x \sqrt{(a^2 - x^2)/x^2}$

Sympy [A]

time = 0.06, size = 22, normalized size = 0.88

$$\begin{cases} a \sqrt{1 - \frac{x^2}{a^2}} + x \operatorname{acsc} \left(\frac{a}{x} \right) & \text{for } a \neq 0 \\ \infty x & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(acsc(a/x),x)`

[Out] `Piecewise((a*sqrt(1 - x**2/a**2) + x*acsc(a/x), Ne(a, 0)), (zoo*x, True))`

Giac [A]

time = 0.43, size = 26, normalized size = 1.04

$$a \left(\frac{x \arcsin \left(\frac{x}{a} \right)}{a} + \sqrt{-\frac{x^2}{a^2} + 1} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(a/x),x, algorithm="giac")`

[Out] $a * (x * \operatorname{arcsin}(x/a) / a + \sqrt{-x^2/a^2 + 1})$

Mupad [B]

time = 0.10, size = 23, normalized size = 0.92

$$a \sqrt{1 - \frac{x^2}{a^2}} + x \operatorname{asin} \left(\frac{x}{a} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(asin(x/a),x)`

[Out] $a * (1 - x^2/a^2)^{(1/2)} + x * \operatorname{asin}(x/a)$

3.12 $\int \frac{\csc^{-1}\left(\frac{a}{x}\right)}{x} dx$

Optimal. Leaf size=59

$$-\frac{1}{2}i\text{ArcSin}\left(\frac{x}{a}\right)^2 + \text{ArcSin}\left(\frac{x}{a}\right) \log\left(1 - e^{2i\text{ArcSin}\left(\frac{x}{a}\right)}\right) - \frac{1}{2}i\text{PolyLog}\left(2, e^{2i\text{ArcSin}\left(\frac{x}{a}\right)}\right)$$

[Out] $-1/2*I*\arcsin(x/a)^2 + \arcsin(x/a)*\ln(1 - (I*x/a + (1-x^2/a^2)^(1/2))^2) - 1/2*I*\text{polylog}(2, (I*x/a + (1-x^2/a^2)^(1/2))^2)$

Rubi [A]

time = 0.04, antiderivative size = 59, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.600, Rules used = {5373, 4721, 3798, 2221, 2317, 2438}

$$-\frac{1}{2}i\text{Li}_2\left(e^{2i\text{ArcSin}\left(\frac{x}{a}\right)}\right) - \frac{1}{2}i\text{ArcSin}\left(\frac{x}{a}\right)^2 + \text{ArcSin}\left(\frac{x}{a}\right) \log\left(1 - e^{2i\text{ArcSin}\left(\frac{x}{a}\right)}\right)$$

Antiderivative was successfully verified.

[In] Int[ArcCsc[a/x]/x, x]

[Out] $(-1/2*I)*\text{ArcSin}[x/a]^2 + \text{ArcSin}[x/a]*\text{Log}[1 - E^{((2*I)*\text{ArcSin}[x/a])}] - (I/2)*\text{PolyLog}[2, E^{((2*I)*\text{ArcSin}[x/a])}]$

Rule 2221

```
Int[((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] :> Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]]
```

Rule 2317

```
Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)]], x_Symbol] :> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]]
```

Rule 2438

```
Int[Log[(c_.)*(d_ + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]]
```

Rule 3798

```
Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + Pi*(k_.) + (f_.)*(x_)], x_Symbol] :> Simp[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Dist[2*I, Int[(c + d*x)^m
```

```
*E^(2*I*k*Pi)*(E^(2*I*(e + f*x))/(1 + E^(2*I*k*Pi)*E^(2*I*(e + f*x)))), x],  
x] /; FreeQ[{c, d, e, f}, x] && IntegerQ[4*k] && IGtQ[m, 0]
```

Rule 4721

```
Int[((a_.) + ArcSin[(c_)*(x_)]*(b_.))^(n_.)/(x_), x_Symbol] :> Subst[Int[(  
a + b*x)^n*Cot[x], x], x, ArcSin[c*x]] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0]
```

Rule 5373

```
Int[ArcCsc[(c_.)/((a_.) + (b_.)*(x_)^(n_.))]^(m_.)*(u_.), x_Symbol] :> Int[  
u*ArcSin[a/c + b*(x^n/c)]^m, x] /; FreeQ[{a, b, c, n, m}, x]
```

Rubi steps

$$\begin{aligned} \int \frac{\csc^{-1}\left(\frac{a}{x}\right)}{x} dx &= \int \frac{\sin^{-1}\left(\frac{x}{a}\right)}{x} dx \\ &= \text{Subst}\left(\int x \cot(x) dx, x, \sin^{-1}\left(\frac{x}{a}\right)\right) \\ &= -\frac{1}{2}i \sin^{-1}\left(\frac{x}{a}\right)^2 - 2i \text{Subst}\left(\int \frac{e^{2ix}x}{1-e^{2ix}} dx, x, \sin^{-1}\left(\frac{x}{a}\right)\right) \\ &= -\frac{1}{2}i \sin^{-1}\left(\frac{x}{a}\right)^2 + \sin^{-1}\left(\frac{x}{a}\right) \log\left(1 - e^{2i \sin^{-1}\left(\frac{x}{a}\right)}\right) - \text{Subst}\left(\int \log\left(1 - e^{2ix}\right) dx, x, \sin^{-1}\left(\frac{x}{a}\right)\right) \\ &= -\frac{1}{2}i \sin^{-1}\left(\frac{x}{a}\right)^2 + \sin^{-1}\left(\frac{x}{a}\right) \log\left(1 - e^{2i \sin^{-1}\left(\frac{x}{a}\right)}\right) + \frac{1}{2}i \text{Subst}\left(\int \frac{\log(1-x)}{x} dx, x, e^{2i \sin^{-1}\left(\frac{x}{a}\right)}\right) \\ &= -\frac{1}{2}i \sin^{-1}\left(\frac{x}{a}\right)^2 + \sin^{-1}\left(\frac{x}{a}\right) \log\left(1 - e^{2i \sin^{-1}\left(\frac{x}{a}\right)}\right) - \frac{1}{2}i \text{Li}_2\left(e^{2i \sin^{-1}\left(\frac{x}{a}\right)}\right) \end{aligned}$$

Mathematica [A]

time = 0.03, size = 54, normalized size = 0.92

$$\csc^{-1}\left(\frac{a}{x}\right) \log\left(1 - e^{2i \csc^{-1}\left(\frac{a}{x}\right)}\right) - \frac{1}{2}i \left(\csc^{-1}\left(\frac{a}{x}\right)^2 + \text{PolyLog}\left(2, e^{2i \csc^{-1}\left(\frac{a}{x}\right)}\right) \right)$$

Antiderivative was successfully verified.

[In] `Integrate[ArcCsc[a/x]/x, x]`

[Out] `ArcCsc[a/x]*Log[1 - E^((2*I)*ArcCsc[a/x])] - (I/2)*(ArcCsc[a/x]^2 + PolyLog[2, E^((2*I)*ArcCsc[a/x])])`

Maple [A]

time = 0.84, size = 125, normalized size = 2.12

method	result
derivativedivides	$-\frac{i \operatorname{arccsc}\left(\frac{a}{x}\right)^2}{2} + \operatorname{arccsc}\left(\frac{a}{x}\right) \ln \left(1 - \frac{ix}{a} - \sqrt{1 - \frac{x^2}{a^2}}\right) + \operatorname{arccsc}\left(\frac{a}{x}\right) \ln \left(1 + \frac{ix}{a} + \sqrt{1 - \frac{x^2}{a^2}}\right)$
default	$-\frac{i \operatorname{arccsc}\left(\frac{a}{x}\right)^2}{2} + \operatorname{arccsc}\left(\frac{a}{x}\right) \ln \left(1 - \frac{ix}{a} - \sqrt{1 - \frac{x^2}{a^2}}\right) + \operatorname{arccsc}\left(\frac{a}{x}\right) \ln \left(1 + \frac{ix}{a} + \sqrt{1 - \frac{x^2}{a^2}}\right)$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(arccsc(a/x)/x,x,method=_RETURNVERBOSE)
[Out] -1/2*I*arccsc(a/x)^2+arccsc(a/x)*ln(1-I*x/a-(1-x^2/a^2)^(1/2))+arccsc(a/x)*
ln(1+I*x/a+(1-x^2/a^2)^(1/2))-I*polylog(2,I*x/a+(1-x^2/a^2)^(1/2))-I*polylo
g(2,-I*x/a-(1-x^2/a^2)^(1/2))
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(arccsc(a/x)/x,x, algorithm="maxima")
[Out] integrate(arccsc(a/x)/x, x)
```

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(arccsc(a/x)/x,x, algorithm="fricas")
[Out] integral(arccsc(a/x)/x, x)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{acsc}\left(\frac{a}{x}\right)}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(acsc(a/x)/x,x)
[Out] Integral(acsc(a/x)/x, x)
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(arccsc(a/x)/x,x, algorithm="giac")

[Out] integrate(arccsc(a/x)/x, x)

Mupad [B]

time = 0.61, size = 49, normalized size = 0.83

$$-\frac{\text{polylog}\left(2, e^{\text{asin}\left(\frac{x}{a}\right) 2 i}\right) 1 i}{2} + \ln \left(1 - e^{\text{asin}\left(\frac{x}{a}\right) 2 i}\right) \text{asin}\left(\frac{x}{a}\right) - \frac{\text{asin}\left(\frac{x}{a}\right)^2 1 i}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(asin(x/a)/x,x)

[Out] $\log(1 - \exp(\text{asin}(x/a)*2i)) * \text{asin}(x/a) - (\text{polylog}(2, \exp(\text{asin}(x/a)*2i)) * 1i)/2 - (\text{asin}(x/a)^2 * 1i)/2$

3.13 $\int \frac{\csc^{-1}\left(\frac{a}{x}\right)}{x^2} dx$

Optimal. Leaf size=32

$$-\frac{\text{ArcSin}\left(\frac{x}{a}\right)}{x} - \frac{\tanh^{-1}\left(\sqrt{1 - \frac{x^2}{a^2}}\right)}{a}$$

[Out] $-\arcsin(x/a)/x - \operatorname{arctanh}((1-x^2/a^2)^{(1/2)})/a$

Rubi [A]

time = 0.02, antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.500, Rules used = {5373, 4723, 272, 65, 214}

$$-\frac{\tanh^{-1}\left(\sqrt{1 - \frac{x^2}{a^2}}\right)}{a} - \frac{\text{ArcSin}\left(\frac{x}{a}\right)}{x}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[\operatorname{ArcCsc}[a/x]/x^2, x]$

[Out] $-(\text{ArcSin}[x/a]/x) - \operatorname{ArcTanh}[\operatorname{Sqrt}[1 - x^2/a^2]]/a$

Rule 65

```
Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] :> With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 214

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] :> Simplify[(Rt[-a/b, 2]/a)*ArcTanh[x/
Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 272

```
Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] :> Dist[1/n, Subst[
Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b,
m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 4723

```

Int[((a_.) + ArcSin[(c_.)*(x_)]*(b_.) )^(n_.)*((d_.)*(x_))^(m_.), x_Symbol]
:> Simp[(d*x)^(m+1)*((a + b*ArcSin[c*x])^n/(d*(m+1))), x] - Dist[b*c*(n/(d*(m+1))), Int[(d*x)^(m+1)*((a + b*ArcSin[c*x])^(n-1)/Sqrt[1 - c^2*x^2]), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]

```

Rule 5373

```

Int[ArcCsc[(c_.)/((a_.) + (b_.)*(x_)^(n_.))]^(m_.)*(u_.), x_Symbol] :> Int[
u*ArcSin[a/c + b*(x^n/c)]^m, x] /; FreeQ[{a, b, c, n, m}, x]

```

Rubi steps

$$\begin{aligned}
\int \frac{\csc^{-1}\left(\frac{a}{x}\right)}{x^2} dx &= \int \frac{\sin^{-1}\left(\frac{x}{a}\right)}{x^2} dx \\
&= -\frac{\sin^{-1}\left(\frac{x}{a}\right)}{x} + \frac{\int \frac{1}{x \sqrt{1 - \frac{x^2}{a^2}}} dx}{a} \\
&= -\frac{\sin^{-1}\left(\frac{x}{a}\right)}{x} + \frac{\text{Subst}\left(\int \frac{1}{x \sqrt{1 - \frac{x^2}{a^2}}} dx, x, x^2\right)}{2a} \\
&= -\frac{\sin^{-1}\left(\frac{x}{a}\right)}{x} - a \text{Subst}\left(\int \frac{1}{a^2 - a^2 x^2} dx, x, \sqrt{1 - \frac{x^2}{a^2}}\right) \\
&= -\frac{\sin^{-1}\left(\frac{x}{a}\right)}{x} - \frac{\tanh^{-1}\left(\sqrt{1 - \frac{x^2}{a^2}}\right)}{a}
\end{aligned}$$

Mathematica [B] Leaf count is larger than twice the leaf count of optimal. 93 vs. 2(32) = 64.

time = 0.10, size = 93, normalized size = 2.91

$$-\frac{\csc^{-1}\left(\frac{a}{x}\right)}{x} - \frac{\sqrt{-1 + \frac{a^2}{x^2}} x \left(-\log\left(1 - \frac{a}{\sqrt{-1 + \frac{a^2}{x^2}} x}\right) + \log\left(1 + \frac{a}{\sqrt{-1 + \frac{a^2}{x^2}} x}\right)\right)}{2a^2 \sqrt{1 - \frac{x^2}{a^2}}}$$

Antiderivative was successfully verified.

[In] Integrate[ArcCsc[a/x]/x^2,x]

[Out] $-(\text{ArcCsc}[a/x]/x) - (\text{Sqrt}[-1 + a^2/x^2]*x*(-\text{Log}[1 - a/(\text{Sqrt}[-1 + a^2/x^2]*x)] + \text{Log}[1 + a/(\text{Sqrt}[-1 + a^2/x^2]*x)]))/(2*a^2*\text{Sqrt}[1 - x^2/a^2])$

Maple [A]

time = 0.07, size = 42, normalized size = 1.31

method	result	size
derivativedivides	$\frac{\frac{\text{arccsc}\left(\frac{a}{x}\right)a}{x} + \ln\left(\frac{\frac{a}{x} + \sqrt{1 - \frac{x^2}{a^2}}}{x}\right)}{a}$	42
default	$\frac{\frac{\text{arccsc}\left(\frac{a}{x}\right)a}{x} + \ln\left(\frac{\frac{a}{x} + \sqrt{1 - \frac{x^2}{a^2}}}{x}\right)}{a}$	42

Verification of antiderivative is not currently implemented for this CAS.

[In] int(arccsc(a/x)/x^2,x,method=_RETURNVERBOSE)

[Out] $-1/a*(\text{arccsc}(a/x)*a/x + \ln(a/x+a/x*(1-x^2/a^2)^(1/2)))$

Maxima [A]

time = 0.26, size = 52, normalized size = 1.62

$$-\frac{\frac{2 a \text{arccsc}\left(\frac{a}{x}\right)}{x} + \log\left(\sqrt{-\frac{x^2}{a^2} + 1} + 1\right) - \log\left(-\sqrt{-\frac{x^2}{a^2} + 1} + 1\right)}{2 a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(arccsc(a/x)/x^2,x, algorithm="maxima")

[Out] $-1/2*(2*a*\text{arccsc}(a/x)/x + \log(\sqrt{-x^2/a^2 + 1} + 1) - \log(-\sqrt{-x^2/a^2 + 1} + 1))/a$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 65 vs. $2(30) = 60$.

time = 0.38, size = 65, normalized size = 2.03

$$-\frac{2 a \text{arccsc}\left(\frac{a}{x}\right) + x \log\left(x \sqrt{\frac{a^2 - x^2}{x^2}} + a\right) - x \log\left(x \sqrt{\frac{a^2 - x^2}{x^2}} - a\right)}{2 a x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(arccsc(a/x)/x^2,x, algorithm="fricas")

[Out] $-1/2*(2*a*arccsc(a/x) + x*log(x*sqrt((a^2 - x^2)/x^2) + a) - x*log(x*sqrt((a^2 - x^2)/x^2) - a))/(a*x)$

Sympy [C] Result contains complex when optimal does not.
time = 1.39, size = 27, normalized size = 0.84

$$-\frac{\operatorname{acsc}\left(\frac{a}{x}\right)}{x} + \frac{\begin{cases} -\operatorname{acosh}\left(\frac{a}{x}\right) & \text{for } \left|\frac{a^2}{x^2}\right| > 1 \\ i \operatorname{asin}\left(\frac{a}{x}\right) & \text{otherwise} \end{cases}}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(acsc(a/x)/x**2,x)`

[Out] $-\operatorname{acsc}(a/x)/x + \operatorname{Piecewise}((- \operatorname{acosh}(a/x), \operatorname{Abs}(a^2/x^2) > 1), (I * \operatorname{asin}(a/x), \text{True}))/a$

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 61 vs. $2(30) = 60$.
time = 0.45, size = 61, normalized size = 1.91

$$-\frac{a \left(\frac{\log(|a + \sqrt{a^2 - x^2}|)}{a} - \frac{\log(|-a + \sqrt{a^2 - x^2}|)}{a} \right)}{2|a|} - \frac{\arcsin\left(\frac{x}{a}\right)}{x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(a/x)/x^2,x, algorithm="giac")`

[Out] $-1/2*a*(\log(\operatorname{abs}(a + \sqrt{a^2 - x^2}))/a - \log(\operatorname{abs}(-a + \sqrt{a^2 - x^2}))/a)/\operatorname{abs}(a) - \arcsin(x/a)/x$

Mupad [B]

time = 0.59, size = 30, normalized size = 0.94

$$-\frac{\operatorname{asin}\left(\frac{x}{a}\right)}{x} - \frac{\operatorname{atanh}\left(\frac{1}{\sqrt{1 - \frac{x^2}{a^2}}}\right)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(asin(x/a)/x^2,x)`

[Out] $-\operatorname{asin}(x/a)/x - \operatorname{atanh}(1/(1 - x^2/a^2)^{(1/2)})/a$

3.14 $\int \frac{\csc^{-1}\left(\frac{a}{x}\right)}{x^3} dx$

Optimal. Leaf size=38

$$-\frac{\sqrt{1 - \frac{x^2}{a^2}}}{2ax} - \frac{\text{ArcSin}\left(\frac{x}{a}\right)}{2x^2}$$

[Out] $-1/2*\arcsin(x/a)/x^2 - 1/2*(1-x^2/a^2)^(1/2)/a/x$

Rubi [A]

time = 0.02, antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.300, Rules used = {5373, 4723, 270}

$$-\frac{\sqrt{1 - \frac{x^2}{a^2}}}{2ax} - \frac{\text{ArcSin}\left(\frac{x}{a}\right)}{2x^2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{ArcCsc}[a/x]/x^3, x]$

[Out] $-1/2*\text{Sqrt}[1 - x^2/a^2]/(a*x) - \text{ArcSin}[x/a]/(2*x^2)$

Rule 270

```
Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] :> Simp[(c*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*c*(m + 1))), x] /; FreeQ[{a, b, c, m, n, p}, x] && EqQ[(m + 1)/n + p + 1, 0] && NeQ[m, -1]
```

Rule 4723

```
Int[((a_) + ArcSin[(c_)*(x_)]*(b_))^(n_)*((d_)*(x_))^(m_), x_Symbol]
:> Simp[(d*x)^(m + 1)*((a + b*ArcSin[c*x])^n/(d*(m + 1))), x] - Dist[b*c*(n/(d*(m + 1))), Int[(d*x)^(m + 1)*((a + b*ArcSin[c*x])^(n - 1)/Sqrt[1 - c^2*x^2]), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]
```

Rule 5373

```
Int[ArcCsc[(c_)/((a_) + (b_)*(x_)^(n_))]^(m_)*(u_), x_Symbol] :> Int[u*ArcSin[a/c + b*(x^n/c)]^m, x] /; FreeQ[{a, b, c, n, m}, x]
```

Rubi steps

$$\begin{aligned}
\int \frac{\csc^{-1}\left(\frac{a}{x}\right)}{x^3} dx &= \int \frac{\sin^{-1}\left(\frac{x}{a}\right)}{x^3} dx \\
&= -\frac{\sin^{-1}\left(\frac{x}{a}\right)}{2x^2} + \frac{\int \frac{1}{\sqrt{1 - \frac{x^2}{a^2}}} dx}{2a} \\
&= -\frac{\sqrt{1 - \frac{x^2}{a^2}}}{2ax} - \frac{\sin^{-1}\left(\frac{x}{a}\right)}{2x^2}
\end{aligned}$$

Mathematica [A]

time = 0.02, size = 35, normalized size = 0.92

$$-\frac{x \sqrt{1 - \frac{x^2}{a^2}} + a \csc^{-1}\left(\frac{a}{x}\right)}{2ax^2}$$

Antiderivative was successfully verified.

[In] Integrate[ArcCsc[a/x]/x^3, x]

[Out] $-1/2*(x* \text{Sqrt}[1 - x^2/a^2] + a*\text{ArcCsc}[a/x])/(a*x^2)$ **Maple [A]**

time = 0.51, size = 54, normalized size = 1.42

method	result	size
derivativedivides	$ -\frac{\frac{a^2 \text{arccsc}\left(\frac{a}{x}\right)}{2x^2} + \frac{x \left(\frac{a^2}{x^2} - 1\right)}{2 \sqrt{\frac{\left(\frac{a^2}{x^2} - 1\right) x^2}{a^2}}}}{a^2} $	54
default	$ -\frac{\frac{a^2 \text{arccsc}\left(\frac{a}{x}\right)}{2x^2} + \frac{x \left(\frac{a^2}{x^2} - 1\right)}{2 \sqrt{\frac{\left(\frac{a^2}{x^2} - 1\right) x^2}{a^2}}}}{a^2} $	54

Verification of antiderivative is not currently implemented for this CAS.

[In] int(arccsc(a/x)/x^3, x, method=_RETURNVERBOSE)

[Out] $-1/a^2*(1/2/x^2*a^2*arccsc(a/x)+1/2/((a^2/x^2-1)*x^2/a^2)^(1/2)*x/a*(a^2/x^2-1))$

Maxima [A]

time = 0.46, size = 32, normalized size = 0.84

$$-\frac{\operatorname{arccsc}\left(\frac{a}{x}\right)}{2x^2} - \frac{\sqrt{-\frac{x^2}{a^2} + 1}}{2ax}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(a/x)/x^3,x, algorithm="maxima")`[Out] $-1/2*\operatorname{arccsc}(a/x)/x^2 - 1/2*\sqrt{(-x^2/a^2 + 1)/(a*x)}$ **Fricas [A]**

time = 0.38, size = 38, normalized size = 1.00

$$-\frac{a^2 \operatorname{arccsc}\left(\frac{a}{x}\right) + x^2 \sqrt{\frac{a^2 - x^2}{x^2}}}{2a^2 x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(a/x)/x^3,x, algorithm="fricas")`[Out] $-1/2*(a^2*\operatorname{arccsc}(a/x) + x^2*\sqrt{((a^2 - x^2)/x^2)})/(a^2*x^2)$ **Sympy [C]** Result contains complex when optimal does not.

time = 0.73, size = 51, normalized size = 1.34

$$-\frac{\operatorname{acsc}\left(\frac{a}{x}\right)}{2x^2} + \begin{cases} -\frac{\sqrt{\frac{a^2}{x^2} - 1}}{a} & \text{for } \left|\frac{a^2}{x^2}\right| > 1 \\ -\frac{i\sqrt{-\frac{a^2}{x^2} + 1}}{a} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(acsc(a/x)/x**3,x)`[Out] $-\operatorname{acsc}(a/x)/(2*x^2) + \operatorname{Piecewise}((-sqrt(a^2/x^2 - 1)/a, \operatorname{Abs}(a^2/x^2) > 1), (-I*sqrt(-a^2/x^2 + 1)/a, \operatorname{True}))/ (2*a)$ **Giac [A]**

time = 0.46, size = 61, normalized size = 1.61

$$-\frac{a \left(\frac{a+\sqrt{a^2-x^2}}{a^2 x} - \frac{x}{(a+\sqrt{a^2-x^2}) a^2} \right)}{4|a|} - \frac{\arcsin\left(\frac{x}{a}\right)}{2x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(a/x)/x^3,x, algorithm="giac")`
[Out]
$$\frac{-1/4*a*((a + \sqrt{a^2 - x^2})*(a^2*x) - x*((a + \sqrt{a^2 - x^2})*a^2))/\text{abs}(a) - 1/2*\arcsin(x/a)/x^2}{x^3}$$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{\arcsin\left(\frac{x}{a}\right)}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(asin(x/a)/x^3,x)`
[Out] `int(asin(x/a)/x^3, x)`

3.15 $\int \frac{\csc^{-1}\left(\frac{a}{x}\right)}{x^4} dx$

Optimal. Leaf size=60

$$-\frac{\sqrt{1 - \frac{x^2}{a^2}}}{6ax^2} - \frac{\text{ArcSin}\left(\frac{x}{a}\right)}{3x^3} - \frac{\tanh^{-1}\left(\sqrt{1 - \frac{x^2}{a^2}}\right)}{6a^3}$$

[Out] $-1/3*\arcsin(x/a)/x^3 - 1/6*\operatorname{arctanh}((1-x^2/a^2)^{(1/2)})/a^3 - 1/6*(1-x^2/a^2)^{(1/2)}/a/x^2$

Rubi [A]

time = 0.03, antiderivative size = 60, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.600, Rules used = {5373, 4723, 272, 44, 65, 214}

$$-\frac{\sqrt{1 - \frac{x^2}{a^2}}}{6ax^2} - \frac{\tanh^{-1}\left(\sqrt{1 - \frac{x^2}{a^2}}\right)}{6a^3} - \frac{\text{ArcSin}\left(\frac{x}{a}\right)}{3x^3}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{ArcCsc}[a/x]/x^4, x]$

[Out] $-1/6*\text{Sqrt}[1 - x^2/a^2]/(a*x^2) - \text{ArcSin}[x/a]/(3*x^3) - \text{ArcTanh}[\text{Sqrt}[1 - x^2/a^2]]/(6*a^3)$

Rule 44

```
Int[((a_.) + (b_.*(x_))^(m_)*((c_.) + (d_.*(x_))^(n_)), x_Symbol] :> Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Dist[d*((m + n + 2)/((b*c - a*d)*(m + 1))), Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x]
/; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && ILtQ[m, -1] && !IntegerQ[n] && LtQ[n, 0]
```

Rule 65

```
Int[((a_.) + (b_.*(x_))^(m_)*((c_.) + (d_.*(x_))^(n_)), x_Symbol] :> With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 214

```
Int[((a_) + (b_.*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 272

```
Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] :> Dist[1/n, Subst[
Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 4723

```
Int[((a_) + ArcSin[(c_)*(x_)]*(b_))^(n_)*((d_)*(x_))^(m_), x_Symbol]
:> Simp[(d*x)^(m + 1)*((a + b*ArcSin[c*x])^n/(d*(m + 1))), x] - Dist[b*c*(n/(d*(m + 1))), Int[(d*x)^(m + 1)*((a + b*ArcSin[c*x])^(n - 1)/Sqrt[1 - c^2*x^2]), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]
```

Rule 5373

```
Int[ArcCsc[(c_)/((a_) + (b_)*(x_)^(n_))]^(m_)*(u_), x_Symbol] :> Int[
u*ArcSin[a/c + b*(x^n/c)]^m, x] /; FreeQ[{a, b, c, n, m}, x]
```

Rubi steps

$$\begin{aligned}
\int \frac{\csc^{-1}\left(\frac{a}{x}\right)}{x^4} dx &= \int \frac{\sin^{-1}\left(\frac{x}{a}\right)}{x^4} dx \\
&= -\frac{\sin^{-1}\left(\frac{x}{a}\right)}{3x^3} + \frac{\int \frac{1}{x^3 \sqrt{1 - \frac{x^2}{a^2}}} dx}{3a} \\
&= -\frac{\sin^{-1}\left(\frac{x}{a}\right)}{3x^3} + \frac{\text{Subst}\left(\int \frac{1}{x^2 \sqrt{1 - \frac{x}{a^2}}} dx, x, x^2\right)}{6a} \\
&= -\frac{\sqrt{1 - \frac{x^2}{a^2}}}{6ax^2} - \frac{\sin^{-1}\left(\frac{x}{a}\right)}{3x^3} + \frac{\text{Subst}\left(\int \frac{1}{x \sqrt{1 - \frac{x}{a^2}}} dx, x, x^2\right)}{12a^3} \\
&= -\frac{\sqrt{1 - \frac{x^2}{a^2}}}{6ax^2} - \frac{\sin^{-1}\left(\frac{x}{a}\right)}{3x^3} - \frac{\text{Subst}\left(\int \frac{1}{a^2 - a^2x^2} dx, x, \sqrt{1 - \frac{x^2}{a^2}}\right)}{6a} \\
&= -\frac{\sqrt{1 - \frac{x^2}{a^2}}}{6ax^2} - \frac{\sin^{-1}\left(\frac{x}{a}\right)}{3x^3} - \frac{\tanh^{-1}\left(\sqrt{1 - \frac{x^2}{a^2}}\right)}{6a^3}
\end{aligned}$$

Mathematica [A]

time = 0.04, size = 69, normalized size = 1.15

$$\frac{a^2 x \sqrt{1 - \frac{x^2}{a^2}} + 2a^3 \csc^{-1}(\frac{a}{x}) - x^3 \log(x) + x^3 \log\left(1 + \sqrt{1 - \frac{x^2}{a^2}}\right)}{6a^3 x^3}$$

Antiderivative was successfully verified.

[In] Integrate[ArcCsc[a/x]/x^4, x]

[Out] $\frac{-1/6*(a^2*x*SQRT[1 - x^2/a^2] + 2*a^3*ArcCsc[a/x] - x^3*Log[x] + x^3*Log[1 + SQRT[1 - x^2/a^2]]/(a^3*x^3))}{a^3*x^3}$ **Maple [A]**

time = 0.51, size = 91, normalized size = 1.52

method	result	size
derivativedivides	$\frac{\frac{a^3 \text{arccsc}(\frac{a}{x})}{3x^3} + \frac{\sqrt{\frac{a^2}{x^2} - 1} \left(\frac{a \sqrt{\frac{a^2}{x^2} - 1}}{x} + \ln\left(\frac{a}{x} + \sqrt{\frac{a^2}{x^2} - 1}\right) \right)x}{6 \sqrt{\frac{(\frac{a^2}{x^2} - 1)x^2}{a^2}}}}{a^3}$	91
default	$\frac{\frac{a^3 \text{arccsc}(\frac{a}{x})}{3x^3} + \frac{\sqrt{\frac{a^2}{x^2} - 1} \left(\frac{a \sqrt{\frac{a^2}{x^2} - 1}}{x} + \ln\left(\frac{a}{x} + \sqrt{\frac{a^2}{x^2} - 1}\right) \right)x}{6 \sqrt{\frac{(\frac{a^2}{x^2} - 1)x^2}{a^2}}}}{a^3}$	91

Verification of antiderivative is not currently implemented for this CAS.

[In] int(arccsc(a/x)/x^4, x, method=_RETURNVERBOSE)

[Out] $\frac{-1/a^3*(1/3*x^3*a^3*arccsc(a/x)+1/6*(a^2/x^2-1)^(1/2)*(1/x*a*(a^2/x^2-1)^(1/2)+ln(a/x+(a^2/x^2-1)^(1/2)))/((a^2/x^2-1)*x^2/a^2)^(1/2)*x/a)}{a^3*x^3}$ **Maxima [A]**

time = 0.45, size = 64, normalized size = 1.07

$$\frac{\log\left(\frac{2 \sqrt{-\frac{x^2}{a^2} + 1}}{|x|} + \frac{2}{|x|}\right)}{6 a} + \frac{\sqrt{-\frac{x^2}{a^2} + 1}}{a^2} - \frac{\text{arccsc}(\frac{a}{x})}{3 x^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(a/x)/x^4,x, algorithm="maxima")`

[Out]
$$\frac{-1/6 * (\log(2*\sqrt(-x^2/a^2 + 1)/\text{abs}(x) + 2/\text{abs}(x))/a^2 + \sqrt(-x^2/a^2 + 1)/x^2)/a - 1/3 * \text{arccsc}(a/x)/x^3}{x^2}$$

Fricas [A]

time = 0.36, size = 92, normalized size = 1.53

$$\frac{4 a^3 \text{arccsc}\left(\frac{a}{x}\right) + x^3 \log\left(x \sqrt{\frac{a^2 - x^2}{x^2}} + a\right) - x^3 \log\left(x \sqrt{\frac{a^2 - x^2}{x^2}} - a\right) + 2 a x^2 \sqrt{\frac{a^2 - x^2}{x^2}}}{12 a^3 x^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(a/x)/x^4,x, algorithm="fricas")`

[Out]
$$\frac{-1/12 * (4*a^3 * \text{arccsc}(a/x) + x^3 * \log(x * \sqrt((a^2 - x^2)/x^2) + a) - x^3 * \log(x * \sqrt((a^2 - x^2)/x^2) - a) + 2*a*x^2 * \sqrt((a^2 - x^2)/x^2))/(a^3 * x^3)}$$

Sympy [A]

time = 1.61, size = 99, normalized size = 1.65

$$\frac{-\frac{\sqrt{\frac{a^2}{x^2} - 1}}{2ax} - \frac{\text{acosh}(\frac{a}{x})}{2a^2}}{3x^3} + \begin{cases} \frac{\frac{ia}{2x^3} \sqrt{-\frac{a^2}{x^2} + 1} - \frac{i}{2ax} \sqrt{-\frac{a^2}{x^2} + 1} + \frac{i \text{asin}(\frac{a}{x})}{2a^2}}{3a} & \text{for } \left|\frac{a^2}{x^2}\right| > 1 \\ \frac{\frac{ia}{2x^3} \sqrt{-\frac{a^2}{x^2} + 1} - \frac{i}{2ax} \sqrt{-\frac{a^2}{x^2} + 1} + \frac{i \text{asin}(\frac{a}{x})}{2a^2}}{3a} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(acsc(a/x)/x**4,x)`

[Out]
$$\frac{-\text{acsc}(a/x)/(3*x**3) + \text{Piecewise}((- \sqrt(a**2/x**2 - 1)/(2*a*x) - \text{acosh}(a/x)/(2*a**2), \text{Abs}(a**2/x**2) > 1), (\text{I}*a/(2*x**3 * \sqrt(-a**2/x**2 + 1)) - \text{I}/(2*a*x * \sqrt(-a**2/x**2 + 1)) + \text{I} * \text{asin}(a/x)/(2*a**2), \text{True})/(3*a))}{3x^3}$$

Giac [A]

time = 0.45, size = 80, normalized size = 1.33

$$\frac{a \left(\frac{\log\left(|a + \sqrt{a^2 - x^2}|\right)}{a^3} - \frac{\log\left(|-a + \sqrt{a^2 - x^2}|\right)}{a^3} + \frac{2 \sqrt{a^2 - x^2}}{a^2 x^2} \right)}{12 |a|} - \frac{\text{arcsin}(\frac{x}{a})}{3 x^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(a/x)/x^4,x, algorithm="giac")`

[Out]
$$\frac{-1/12*a*(\log(\text{abs}(a + \sqrt{a^2 - x^2}))/a^3 - \log(\text{abs}(-a + \sqrt{a^2 - x^2}))/a^3 + 2*\sqrt{a^2 - x^2}/(a^2*x^2))/\text{abs}(a) - 1/3*\arcsin(x/a)/x^3}{x^4}$$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\arcsin\left(\frac{x}{a}\right)}{x^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(asin(x/a)/x^4,x)`

[Out] `int(asin(x/a)/x^4, x)`

3.16 $\int \frac{\csc^{-1}(ax^n)}{x} dx$

Optimal. Leaf size=69

$$\frac{i \csc^{-1}(ax^n)^2}{2n} - \frac{\csc^{-1}(ax^n) \log\left(1 - e^{2i \csc^{-1}(ax^n)}\right)}{n} + \frac{i \text{PolyLog}\left(2, e^{2i \csc^{-1}(ax^n)}\right)}{2n}$$

[Out] $\frac{1}{2} I \arccsc(a x^n)^2/n - \arccsc(a x^n) \ln(1 - (I/a/(x^n) + (1 - 1/a^2/(x^n)^2)^{(1/2)})^2)/n + 1/2 I \text{polylog}(2, (I/a/(x^n) + (1 - 1/a^2/(x^n)^2)^{(1/2)})^2)/n$

Rubi [A]

time = 0.07, antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.600,

Rules used = {5327, 4721, 3798, 2221, 2317, 2438}

$$\frac{i \text{Li}_2\left(e^{2i \csc^{-1}(ax^n)}\right)}{2n} + \frac{i \csc^{-1}(ax^n)^2}{2n} - \frac{\csc^{-1}(ax^n) \log\left(1 - e^{2i \csc^{-1}(ax^n)}\right)}{n}$$

Antiderivative was successfully verified.

[In] Int[ArcCsc[a*x^n]/x, x]

[Out] $((I/2)*\text{ArcCsc}[a*x^n]^2)/n - (\text{ArcCsc}[a*x^n]*\text{Log}[1 - E^{((2*I)*\text{ArcCsc}[a*x^n])}])/n + ((I/2)*\text{PolyLog}[2, E^{((2*I)*\text{ArcCsc}[a*x^n])}])/n$

Rule 2221

```
Int[((((F_)^((g_.)*(e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/((a_) + (b_.)*((F_)^((g_.)*(e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] :> Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x]; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2317

```
Int[Log[(a_) + (b_.)*((F_)^((e_.)*(c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] :> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x]; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2438

```
Int[Log[(c_.)*(d_) + (e_.)*(x_)^(n_.)]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x]; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 3798

```
Int[((c_) + (d_)*(x_))^(m_)*tan[(e_) + Pi*(k_) + (f_)*(x_)], x_Symbol]
] :> Simp[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Dist[2*I, Int[(c + d*x)^m
*E^(2*I*k*Pi)*(E^(2*I*(e + f*x))/(1 + E^(2*I*k*Pi)*E^(2*I*(e + f*x)))), x],
x] /; FreeQ[{c, d, e, f}, x] && IntegerQ[4*k] && IGtQ[m, 0]
```

Rule 4721

```
Int[((a_) + ArcSin[(c_)*(x_)]*(b_))/x_, x_Symbol] :> Subst[Int[(a +
b*x)^n*Cot[x], x, ArcSin[c*x]] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0]
```

Rule 5327

```
Int[((a_) + ArcCsc[(c_)*(x_)]*(b_))/x_, x_Symbol] :> -Subst[Int[(a + b
*ArcSin[x/c])/x, x, 1/x] /; FreeQ[{a, b, c}, x]
```

Rubi steps

$$\begin{aligned}
\int \frac{\csc^{-1}(ax^n)}{x} dx &= \frac{\text{Subst}\left(\int \frac{\csc^{-1}(ax)}{x} dx, x, x^n\right)}{n} \\
&= -\frac{\text{Subst}\left(\int \frac{\sin^{-1}(\frac{x}{a})}{x} dx, x, x^{-n}\right)}{n} \\
&= -\frac{\text{Subst}\left(\int x \cot(x) dx, x, \sin^{-1}\left(\frac{x^{-n}}{a}\right)\right)}{n} \\
&= \frac{i \sin^{-1}\left(\frac{x^{-n}}{a}\right)^2}{2n} + \frac{(2i)\text{Subst}\left(\int \frac{e^{2ix}x}{1-e^{2ix}} dx, x, \sin^{-1}\left(\frac{x^{-n}}{a}\right)\right)}{n} \\
&= \frac{i \sin^{-1}\left(\frac{x^{-n}}{a}\right)^2}{2n} - \frac{\sin^{-1}\left(\frac{x^{-n}}{a}\right) \log\left(1 - e^{2i \sin^{-1}\left(\frac{x^{-n}}{a}\right)}\right)}{n} + \frac{\text{Subst}\left(\int \log(1 - e^{2ix}) dx, x, \sin^{-1}\left(\frac{x^{-n}}{a}\right)\right)}{n} \\
&= \frac{i \sin^{-1}\left(\frac{x^{-n}}{a}\right)^2}{2n} - \frac{\sin^{-1}\left(\frac{x^{-n}}{a}\right) \log\left(1 - e^{2i \sin^{-1}\left(\frac{x^{-n}}{a}\right)}\right)}{n} - \frac{i \text{Subst}\left(\int \frac{\log(1-x)}{x} dx, x, e^{2i \sin^{-1}\left(\frac{x^{-n}}{a}\right)}\right)}{2n} \\
&= \frac{i \sin^{-1}\left(\frac{x^{-n}}{a}\right)^2}{2n} - \frac{\sin^{-1}\left(\frac{x^{-n}}{a}\right) \log\left(1 - e^{2i \sin^{-1}\left(\frac{x^{-n}}{a}\right)}\right)}{n} + \frac{i \text{Li}_2\left(e^{2i \sin^{-1}\left(\frac{x^{-n}}{a}\right)}\right)}{2n}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.06, size = 63, normalized size = 0.91

$$-\frac{x^{-n} {}_3F_2\left(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}; \frac{3}{2}, \frac{3}{2}; \frac{x^{-2n}}{a^2}\right)}{an} + \left(\csc^{-1}(ax^n) - \text{ArcSin}\left(\frac{x^{-n}}{a}\right)\right) \log(x)$$

Antiderivative was successfully verified.

[In] `Integrate[ArcCsc[a*x^n]/x,x]`

[Out] $-\text{HypergeometricPFQ}[\{1/2, 1/2, 1/2\}, \{3/2, 3/2\}, 1/(a^2 x^{(2n)})]/(a^n x^n) + (\text{ArcCsc}[a x^n] - \text{ArcSin}[1/(a x^n)]) \text{Log}[x]$

Maple [A]

time = 1.32, size = 155, normalized size = 2.25

method	result
derivativedivides	$\frac{i \arccsc(a x^n)^2}{2} - \arccsc(a x^n) \ln\left(1 - \frac{ix^{-n}}{a} - \sqrt{1 - \frac{x^{-2n}}{a^2}}\right) - \arccsc(a x^n) \ln\left(1 + \frac{ix^{-n}}{a} + \sqrt{1 - \frac{x^{-2n}}{a^2}}\right) + i \text{polylog}\left(2, \frac{ix^{-n}}{a} + \sqrt{1 - \frac{x^{-2n}}{a^2}}\right)$
default	$\frac{i \arccsc(a x^n)^2}{2} - \arccsc(a x^n) \ln\left(1 - \frac{ix^{-n}}{a} - \sqrt{1 - \frac{x^{-2n}}{a^2}}\right) - \arccsc(a x^n) \ln\left(1 + \frac{ix^{-n}}{a} + \sqrt{1 - \frac{x^{-2n}}{a^2}}\right) + i \text{polylog}\left(2, \frac{ix^{-n}}{a} + \sqrt{1 - \frac{x^{-2n}}{a^2}}\right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(arccsc(a*x^n)/x,x,method=_RETURNVERBOSE)`

[Out] $1/n*(1/2*I*\arccsc(a*x^n)^2 - \arccsc(a*x^n)*\ln(1-I/a/(x^n) - (1-1/a^2/(x^n)^2)^(1/2)) - \arccsc(a*x^n)*\ln(1+I/a/(x^n) + (1-1/a^2/(x^n)^2)^(1/2)) + I*\text{polylog}(2,I/a/(x^n) + (1-1/a^2/(x^n)^2)^(1/2)) + I*\text{polylog}(2,-I/a/(x^n) - (1-1/a^2/(x^n)^2)^(1/2)))$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(a*x^n)/x,x, algorithm="maxima")`

[Out] $a^{2n} \int \sqrt{a x^n + 1} \sqrt{a x^n - 1} \log(x) / (a^{4n} x^{2n} - a^2 x), x) + \text{arctan2}(1, \sqrt{a x^n + 1} \sqrt{a x^n - 1}) \log(x)$

Fricas [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(a*x^n)/x,x, algorithm="fricas")`

[Out] Exception raised: TypeError >> Error detected within library code: integ rate: implementation incomplete (constant residues)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{acsc}(ax^n)}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(acsc(a*x**n)/x,x)`

[Out] `Integral(acsc(a*x**n)/x, x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(a*x^n)/x,x, algorithm="giac")`

[Out] `integrate(arccsc(a*x^n)/x, x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\operatorname{asin}\left(\frac{1}{ax^n}\right)}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(asin(1/(a*x^n))/x,x)`

[Out] `int(asin(1/(a*x^n))/x, x)`

$$\mathbf{3.17} \quad \int x^4 \csc^{-1}(a + bx) dx$$

Optimal. Leaf size=197

$$\frac{a(20 + 53a^2)(a + bx)\sqrt{1 - \frac{1}{(a + bx)^2}}}{30b^5} - \frac{11ax^2(a + bx)\sqrt{1 - \frac{1}{(a + bx)^2}}}{60b^3} + \frac{x^3(a + bx)\sqrt{1 - \frac{1}{(a + bx)^2}}}{20b^2} + \dots$$

[Out] $\frac{1}{5}a^5 \operatorname{arccsc}(bx+a)/b^5 + \frac{1}{5}x^5 \operatorname{arccsc}(bx+a) + \frac{1}{40}(40a^4 + 40a^2 + 3) \operatorname{arctanh}\left(\frac{1}{\sqrt{1 - \frac{1}{(a+bx)^2}}}\right) + \frac{1}{30}a^5(53a^2 + 20)(bx+a) \operatorname{arccsc}\left(\frac{1}{\sqrt{1 - \frac{1}{(a+bx)^2}}}\right) + \frac{1}{60}a^5x^2(bx+a) \operatorname{arccsc}\left(\frac{1}{\sqrt{1 - \frac{1}{(a+bx)^2}}}\right) + \frac{1}{120}a^5x^4(bx+a) \operatorname{arccsc}\left(\frac{1}{\sqrt{1 - \frac{1}{(a+bx)^2}}}\right)$

Rubi [A]

time = 0.17, antiderivative size = 197, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.800$, Rules used = {5367, 4512, 3867, 4141, 4133, 3855, 3852, 8}

$$\frac{a^5 \csc^{-1}(a + bx)}{5b^5} - \frac{(53a^2 + 20)a(a + bx)\sqrt{1 - \frac{1}{(a + bx)^2}}}{30b^5} + \frac{(58a^2 + 9)(a + bx)^2\sqrt{1 - \frac{1}{(a + bx)^2}}}{120b^5} + \frac{(40a^4 + 40a^2 + 3)\tanh^{-1}\left(\sqrt{1 - \frac{1}{(a + bx)^2}}\right)}{40b^5} - \frac{11ax^2(a + bx)\sqrt{1 - \frac{1}{(a + bx)^2}}}{60b^3} + \frac{x^3(a + bx)\sqrt{1 - \frac{1}{(a + bx)^2}}}{20b^2} + \frac{1}{5}x^5 \csc^{-1}(a + bx)$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[x^4 \operatorname{ArcCsc}[a + b*x], x]$

[Out] $\frac{-1}{30}(a(20 + 53a^2)(a + b*x)\operatorname{Sqrt}[1 - (a + b*x)^{-2}])/b^5 - \frac{(11a^2x^2(a + b*x)\operatorname{Sqrt}[1 - (a + b*x)^{-2}])/(60b^3) + (x^3(a + b*x)\operatorname{Sqrt}[1 - (a + b*x)^{-2}])/(20b^2) + ((9 + 58a^2)(a + b*x)^2\operatorname{Sqrt}[1 - (a + b*x)^{-2}])/(120b^5) + (a^5\operatorname{ArcCsc}[a + b*x])/(5b^5) + (x^5\operatorname{ArcCsc}[a + b*x])/5 + ((3 + 40a^2 + 40a^4)\operatorname{ArcTanh}[\operatorname{Sqrt}[1 - (a + b*x)^{-2}]])/(40b^5)}$

Rule 8

$\operatorname{Int}[a_, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[a*x, x] /; \operatorname{FreeQ}[a, x]$

Rule 3852

$\operatorname{Int}[\csc[(c_.) + (d_.)*(x_.)]^{n_}, x_{\text{Symbol}}] \rightarrow \operatorname{Dist}[-d^{(-1)}, \operatorname{Subst}[\operatorname{Int}[\operatorname{ExpandIntegrand}[(1 + x^2)^{(n/2 - 1)}, x], x, \operatorname{Cot}[c + d*x]], x] /; \operatorname{FreeQ}[\{c, d\}, x] \&& \operatorname{IGtQ}[n/2, 0]$

Rule 3855

$\operatorname{Int}[\csc[(c_.) + (d_.)*(x_.)], x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[-\operatorname{ArcTanh}[\operatorname{Cos}[c + d*x]]/d, x] /; \operatorname{FreeQ}[\{c, d\}, x]$

Rule 3867

```

Int[(csc[(c_.) + (d_ .)*(x_)]*(b_ .) + (a_ ))^(n_), x_Symbol] :> Simp[(-b^2)*C
ot[c + d*x]*((a + b*Csc[c + d*x])^(n - 2)/(d*(n - 1))), x] + Dist[1/(n - 1)
, Int[(a + b*Csc[c + d*x])^(n - 3)*Simp[a^3*(n - 1) + (b*(b^2*(n - 2) + 3*a
~2*(n - 1)))*Csc[c + d*x] + (a*b^2*(3*n - 4))*Csc[c + d*x]^2, x], x], x] /;
FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[n, 2] && IntegerQ[2*n]

```

Rule 4133

```

Int[((A_.) + csc[(e_.) + (f_ .)*(x_)]*(B_ .) + csc[(e_.) + (f_ .)*(x_)]^2*(C_ .
))* (csc[(e_.) + (f_ .)*(x_)]*(b_ .) + (a_ )), x_Symbol] :> Simp[(-b)*C*Csc[e +
f*x]*(Cot[e + f*x]/(2*f)), x] + Dist[1/2, Int[Simp[2*A*a + (2*B*a + b*(2*A
+ C))*Csc[e + f*x] + 2*(a*C + B*b)*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, e, f, A, B, C}, x]

```

Rule 4141

```

Int[((A_.) + csc[(e_.) + (f_ .)*(x_)]*(B_ .) + csc[(e_.) + (f_ .)*(x_)]^2*(C_ .
))* (csc[(e_.) + (f_ .)*(x_)]*(b_ .) + (a_ ))^(m_.), x_Symbol] :> Simp[(-C)*Cot
[e + f*x]*((a + b*Csc[e + f*x])^m/(f*(m + 1))), x] + Dist[1/(m + 1), Int[(a
+ b*Csc[e + f*x])^(m - 1)*Simp[a*A*(m + 1) + ((A*b + a*B)*(m + 1) + b*C*m
)*Csc[e + f*x] + (b*B*(m + 1) + a*C*m)*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, e, f, A, B, C}, x] && NeQ[a^2 - b^2, 0] && IGtQ[2*m, 0]

```

Rule 4512

```

Int[Cot[(c_.) + (d_ .)*(x_)]*Csc[(c_.) + (d_ .)*(x_)]*(Csc[(c_.) + (d_ .)*(x_ )
]*(b_ .) + (a_ ))^(n_ .)*(e_.) + (f_ .)*(x_ ))^(m_.), x_Symbol] :> Simp[(-(e +
f*x)^m)*((a + b*Csc[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[f*(m/(b*d*(n
+ 1))), Int[(e + f*x)^(m - 1)*(a + b*Csc[c + d*x])^(n + 1), x], x] /; Fre
eQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]

```

Rule 5367

```

Int[((a_.) + ArcCsc[(c_) + (d_ .)*(x_)]*(b_ .))^(p_ .)*((e_.) + (f_ .)*(x_ ))^(m
_.), x_Symbol] :> Dist[-(d^(m + 1))^(-1), Subst[Int[(a + b*x)^p*Csc[x]*Cot[
x]*(d*e - c*f + f*Csc[x])^m, x], x, ArcCsc[c + d*x]], x] /; FreeQ[{a, b, c,
d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]

```

Rubi steps

$$\begin{aligned}
\int x^4 \csc^{-1}(a + bx) dx &= -\frac{\text{Subst}(\int x \cot(x) \csc(x)(-a + \csc(x))^4 dx, x, \csc^{-1}(a + bx))}{b^5} \\
&= \frac{1}{5}x^5 \csc^{-1}(a + bx) - \frac{\text{Subst}(\int (-a + \csc(x))^5 dx, x, \csc^{-1}(a + bx))}{5b^5} \\
&= \frac{x^3(a + bx)\sqrt{1 - \frac{1}{(a + bx)^2}}}{20b^2} + \frac{1}{5}x^5 \csc^{-1}(a + bx) - \frac{\text{Subst}(\int (-a + \csc(x))^2(-4a^3 - \\
&\quad 11ax^2(a + bx)\sqrt{1 - \frac{1}{(a + bx)^2}} + \frac{x^3(a + bx)\sqrt{1 - \frac{1}{(a + bx)^2}}}{20b^2} + \frac{1}{5}x^5 \csc^{-1}(a + bx) \\
&\quad - \frac{11ax^2(a + bx)\sqrt{1 - \frac{1}{(a + bx)^2}}}{60b^3} + \frac{x^3(a + bx)\sqrt{1 - \frac{1}{(a + bx)^2}}}{20b^2} + \frac{(9 + 58a^2)(a + bx)}{60b^3} \\
&\quad - \frac{11ax^2(a + bx)\sqrt{1 - \frac{1}{(a + bx)^2}}}{60b^3} + \frac{x^3(a + bx)\sqrt{1 - \frac{1}{(a + bx)^2}}}{20b^2} + \frac{(9 + 58a^2)(a + bx)}{60b^3} \\
&\quad - \frac{11ax^2(a + bx)\sqrt{1 - \frac{1}{(a + bx)^2}}}{60b^3} + \frac{x^3(a + bx)\sqrt{1 - \frac{1}{(a + bx)^2}}}{20b^2} + \frac{(9 + 58a^2)(a + bx)}{60b^3} \\
&\quad - \frac{a(20 + 53a^2)(a + bx)\sqrt{1 - \frac{1}{(a + bx)^2}}}{30b^5} - \frac{11ax^2(a + bx)\sqrt{1 - \frac{1}{(a + bx)^2}}}{60b^3} + \frac{x^3(a + bx)\sqrt{1 - \frac{1}{(a + bx)^2}}}{20b^2} + \frac{(9 + 58a^2)(a + bx)}{60b^3})}{120b^5}
\end{aligned}$$

Mathematica [A]

time = 0.15, size = 174, normalized size = 0.88

$$-\frac{\sqrt{\frac{-1 + a^2 + 2abx + b^2x^2}{(a + bx)^2}} (a^2(71 + 154a^2) + 2a(31 + 48a^2)bx - 9(1 + 4a^2)b^2x^2 + 16ab^3x^3 - 6b^4x^4) + 24b^5x^5 \csc^{-1}(a + bx) + 24a^5 \text{ArcSin}\left(\frac{1}{a + bx}\right) + 3(3 + 40a^2 + 40a^4) \log\left((a + bx) \left(1 + \sqrt{\frac{-1 + a^2 + 2abx + b^2x^2}{(a + bx)^2}}\right)\right)}{120b^5}$$

Antiderivative was successfully verified.

[In] `Integrate[x^4*ArcCsc[a + b*x], x]`

[Out] $\begin{aligned}
&(-(Sqrt[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2]*(a^2*(71 + 154*a^2) + 2 \\
&*a*(31 + 48*a^2)*b*x - 9*(1 + 4*a^2)*b^2*x^2 + 16*a*b^3*x^3 - 6*b^4*x^4)) + \\
&24*b^5*x^5*ArcCsc[a + b*x] + 24*a^5*ArcSin[(a + b*x)^{-1}] + 3*(3 + 40*a^2 \\
&+ 40*a^4)*Log[(a + b*x)*(1 + Sqrt[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2])])/(120*b^5)
\end{aligned}$

Maple [A]

time = 0.30, size = 329, normalized size = 1.67

method	result
derivativedivides	$-\frac{\operatorname{arccsc}(bx+a)a^5}{5} + \operatorname{arccsc}(bx+a)a^4(bx+a) - 2\operatorname{arccsc}(bx+a)a^3(bx+a)^2 + 2\operatorname{arccsc}(bx+a)a^2(bx+a)^3 - \operatorname{arccsc}(bx+a)a(bx+a)^4$
default	$-\frac{\operatorname{arccsc}(bx+a)a^5}{5} + \operatorname{arccsc}(bx+a)a^4(bx+a) - 2\operatorname{arccsc}(bx+a)a^3(bx+a)^2 + 2\operatorname{arccsc}(bx+a)a^2(bx+a)^3 - \operatorname{arccsc}(bx+a)a(bx+a)^4$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x^4*arccsc(b*x+a), x, method=_RETURNVERBOSE)`[Out]
$$\begin{aligned} & 1/b^5 * (-1/5 * \operatorname{arccsc}(b*x+a)*a^5 + \operatorname{arccsc}(b*x+a)*a^4*(b*x+a) - 2*\operatorname{arccsc}(b*x+a)*a^3 \\ & *(b*x+a)^2 + 2*\operatorname{arccsc}(b*x+a)*a^2*(b*x+a)^3 - \operatorname{arccsc}(b*x+a)*a*(b*x+a)^4 + 1/5*\operatorname{arccsc}(b*x+a)*(b*x+a)^5 - 1/120*((b*x+a)^2-1)^{(1/2)} * (-24*a^5*\operatorname{arctan}(1/((b*x+a)^2-1)^{(1/2)}) - 120*a^4*\ln(b*x+a+((b*x+a)^2-1)^{(1/2)}) + 240*a^3*((b*x+a)^2-1)^{(1/2)} - 120*a^2*((b*x+a)^2-1)^{(1/2)} + 40*a*(b*x+a)^2*((b*x+a)^2-1)^{(1/2)} - 6*((b*x+a)^3*((b*x+a)^2-1)^{(1/2)} - 120*a^2*\ln(b*x+a+((b*x+a)^2-1)^{(1/2)}) + 80*a*((b*x+a)^2-1)^{(1/2)} - 9*(b*x+a)*((b*x+a)^2-1)^{(1/2)} - 9*\ln(b*x+a+((b*x+a)^2-1)^{(1/2)})) / (((b*x+a)^2-1)/(b*x+a)^2)^{(1/2)} / (b*x+a) \end{aligned}$$
Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^4*arccsc(b*x+a), x, algorithm="maxima")`[Out]
$$\begin{aligned} & 1/5*x^5*\operatorname{arctan2}(1, \sqrt{b*x + a + 1})*\sqrt{b*x + a - 1}) + \operatorname{integrate}(1/5*(b^2*x^6 + a*b*x^5)*e^{(1/2*\log(b*x + a + 1) + 1/2*\log(b*x + a - 1)) / (b^2*x^2 + 2*a*b*x + a^2 + (b^2*x^2 + 2*a*b*x + a^2 - 1)*e^{(\log(b*x + a + 1) + \log(b*x + a - 1)) - 1}), x) \end{aligned}$$
Fricas [A]

time = 0.40, size = 151, normalized size = 0.77

$$\frac{24 b^5 x^5 \operatorname{arccsc}(bx+a) - 48 a^5 \arctan(-bx - a + \sqrt{b^2 x^2 + 2 abx + a^2 - 1}) - 3(40 a^4 + 40 a^2 + 3) \log(-bx - a + \sqrt{b^2 x^2 + 2 abx + a^2 - 1}) + (6 b^3 x^3 - 22 ab^2 x^2 - 154 a^3 + (58 a^2 + 9)bx - 71 a)\sqrt{b^2 x^2 + 2 abx + a^2 - 1}}{120 b^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^4*arccsc(b*x+a),x, algorithm="fricas")`
[Out] $\frac{1}{120} \left(24b^5x^5 \operatorname{arccsc}(bx + a) - 48a^5 \operatorname{arctan}(-bx - a + \sqrt{b^2x^2 + 2ax + a^2 - 1}) - 3(40a^4 + 40a^2 + 3) \log(-bx - a + \sqrt{b^2x^2 + 2ax + a^2 - 1}) + (6b^3x^3 - 22a^2b^2x^2 - 154a^3 + (58a^2 + 9)a^2 - 71a) \sqrt{b^2x^2 + 2ax + a^2 - 1} \right) / b^5$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x^4 \operatorname{acsc}(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x**4*acsc(b*x+a),x)`
[Out] `Integral(x**4*acsc(a + b*x), x)`

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 408 vs. $2(173) = 346$.

time = 0.55, size = 408, normalized size = 2.07

$$\frac{1}{320} \left(\frac{320 \operatorname{atan}\left(\frac{\sqrt{-\frac{1}{(bx+a)^2}}}{\sqrt{\frac{1}{(bx+a)^2}+1}}\right) + 48 \operatorname{atan}\left(\frac{\sqrt{-\frac{1}{(bx+a)^2}}}{\sqrt{\frac{1}{(bx+a)^2}+1}}\right)^3 + 960 \operatorname{atan}\left(\frac{\sqrt{-\frac{1}{(bx+a)^2}}}{\sqrt{\frac{1}{(bx+a)^2}+1}}\right)^5 + 3840 \operatorname{atan}\left(\frac{\sqrt{-\frac{1}{(bx+a)^2}}}{\sqrt{\frac{1}{(bx+a)^2}+1}}\right)^7 + 24(3a+b)x^7 \operatorname{atan}\left(\frac{\sqrt{-\frac{1}{(bx+a)^2}}}{\sqrt{\frac{1}{(bx+a)^2}+1}}\right) + 24(18a^2+40a^2+21) \operatorname{atan}\left(\frac{\sqrt{-\frac{1}{(bx+a)^2}}}{\sqrt{\frac{1}{(bx+a)^2}+1}}\right)^3 + 24(18a^2+40a^2+21) \operatorname{atan}\left(\frac{\sqrt{-\frac{1}{(bx+a)^2}}}{\sqrt{\frac{1}{(bx+a)^2}+1}}\right)^5 + 24(18a^2+40a^2+21) \operatorname{atan}\left(\frac{\sqrt{-\frac{1}{(bx+a)^2}}}{\sqrt{\frac{1}{(bx+a)^2}+1}}\right)^7 + 360 \operatorname{atan}\left(\frac{\sqrt{-\frac{1}{(bx+a)^2}}}{\sqrt{\frac{1}{(bx+a)^2}+1}}\right)^9 + 360 \operatorname{atan}\left(\frac{\sqrt{-\frac{1}{(bx+a)^2}}}{\sqrt{\frac{1}{(bx+a)^2}+1}}\right)^{11} + 360 \operatorname{atan}\left(\frac{\sqrt{-\frac{1}{(bx+a)^2}}}{\sqrt{\frac{1}{(bx+a)^2}+1}}\right)^{13} + 360 \operatorname{atan}\left(\frac{\sqrt{-\frac{1}{(bx+a)^2}}}{\sqrt{\frac{1}{(bx+a)^2}+1}}\right)^{15} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^4*arccsc(b*x+a),x, algorithm="giac")`
[Out] $\begin{aligned} & -\frac{1}{960} b^6 (192(bx + a)^5 (5a/(bx + a) - 10a^2/(bx + a)^2 + 10a^3/(bx + a)^3 - 5a^4/(bx + a)^4 - 1) \arcsin(-1/((bx + a)(a/(bx + a) - 1) - a)) + (3(bx + a)^4 (\sqrt{-1/(bx + a)^2 + 1} - 1)^4 + 40(bx + a)^3 a (\sqrt{-1/(bx + a)^2 + 1} - 1)^3 + 240(bx + a)^2 a^2 (\sqrt{-1/(bx + a)^2 + 1} - 1)^2 + 960(bx + a) a^3 (\sqrt{-1/(bx + a)^2 + 1} - 1) + 24(bx + a)^2 (\sqrt{-1/(bx + a)^2 + 1} - 1)^2 + 360(bx + a) a^4 (\sqrt{-1/(bx + a)^2 + 1} - 1) + 24(40a^4 + 40a^2 + 3) \log(-(\sqrt{-1/(bx + a)^2 + 1} - 1) * \operatorname{abs}(bx + a)) - (120(8a^3 + 3a) * (bx + a)^3 * (\sqrt{-1/(bx + a)^2 + 1} - 1)^3 + 24(10a^2 + 1) * (bx + a)^2 * (\sqrt{-1/(bx + a)^2 + 1} - 1)^2 + 40(bx + a) * a * (\sqrt{-1/(bx + a)^2 + 1} - 1)^3 + 3) / ((bx + a)^4 * (\sqrt{-1/(bx + a)^2 + 1} - 1)^4)) / b^6 \end{aligned}$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int x^4 \operatorname{asin}\left(\frac{1}{a + bx}\right) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x^4*asin(1/(a + b*x)),x)`
[Out] `int(x^4*asin(1/(a + b*x)), x)`

3.18 $\int x^3 \csc^{-1}(a + bx) dx$

Optimal. Leaf size=155

$$\frac{(2 + 17a^2)(a + bx)\sqrt{1 - \frac{1}{(a + bx)^2}}}{12b^4} + \frac{x^2(a + bx)\sqrt{1 - \frac{1}{(a + bx)^2}}}{12b^2} - \frac{a(a + bx)^2\sqrt{1 - \frac{1}{(a + bx)^2}}}{3b^4} - \frac{a^4 \csc^{-1}(a + bx)}{4b^4}$$

[Out] $-1/4*a^4*\text{arccsc}(b*x+a)/b^4+1/4*x^4*\text{arccsc}(b*x+a)-1/2*a*(2*a^2+1)*\text{arctanh}((1-1/(b*x+a)^2)^(1/2))/b^4+1/12*(17*a^2+2)*(b*x+a)*(1-1/(b*x+a)^2)^(1/2)/b^4+1/12*x^2*(b*x+a)*(1-1/(b*x+a)^2)^(1/2)/b^2-1/3*a*(b*x+a)^2*(1-1/(b*x+a)^2)^(1/2)/b^4$

Rubi [A]

time = 0.11, antiderivative size = 155, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.700, Rules used = {5367, 4512, 3867, 4133, 3855, 3852, 8}

$$-\frac{a^4 \csc^{-1}(a + bx)}{4b^4} + \frac{(17a^2 + 2)(a + bx)\sqrt{1 - \frac{1}{(a + bx)^2}}}{12b^4} - \frac{(2a^2 + 1)a \tanh^{-1}\left(\sqrt{1 - \frac{1}{(a + bx)^2}}\right)}{2b^4} - \frac{a(a + bx)^2\sqrt{1 - \frac{1}{(a + bx)^2}}}{3b^4} + \frac{x^2(a + bx)\sqrt{1 - \frac{1}{(a + bx)^2}}}{12b^2} + \frac{1}{4}x^4 \csc^{-1}(a + bx)$$

Antiderivative was successfully verified.

[In] $\text{Int}[x^3 \text{ArcCsc}[a + b*x], x]$

[Out] $((2 + 17*a^2)*(a + b*x)*\text{Sqrt}[1 - (a + b*x)^{-2}])/(12*b^4) + (x^2*(a + b*x)*\text{Sqrt}[1 - (a + b*x)^{-2}])/(12*b^2) - (a*(a + b*x)^2*\text{Sqrt}[1 - (a + b*x)^{-2}])/(3*b^4) - (a^4*\text{ArcCsc}[a + b*x])/(4*b^4) + (x^4*\text{ArcCsc}[a + b*x])/4 - (a*(1 + 2*a^2)*\text{ArcTanh}[\text{Sqrt}[1 - (a + b*x)^{-2}]])/(2*b^4)$

Rule 8

$\text{Int}[a_, x_Symbol] :> \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

Rule 3852

$\text{Int}[\csc[(c_.) + (d_.)*(x_.)]^{(n_)}, x_Symbol] :> \text{Dist}[-d^{(-1)}, \text{Subst}[\text{Int}[\text{ExpandIntegrand}[(1 + x^2)^{(n/2 - 1)}, x], x, \text{Cot}[c + d*x]], x] /; \text{FreeQ}[\{c, d\}, x] \&& \text{IGtQ}[n/2, 0]$

Rule 3855

$\text{Int}[\csc[(c_.) + (d_.)*(x_.)], x_Symbol] :> \text{Simp}[-\text{ArcTanh}[\text{Cos}[c + d*x]]/d, x] /; \text{FreeQ}[\{c, d\}, x]$

Rule 3867

```

Int[(csc[(c_.) + (d_ .)*(x_)]*(b_ .) + (a_ ))^(n_), x_Symbol] :> Simp[(-b^2)*C
ot[c + d*x]*((a + b*Csc[c + d*x])^(n - 2)/(d*(n - 1))), x] + Dist[1/(n - 1)
, Int[(a + b*Csc[c + d*x])^(n - 3)*Simp[a^3*(n - 1) + (b*(b^2*(n - 2) + 3*a
^2*(n - 1)))*Csc[c + d*x] + (a*b^2*(3*n - 4))*Csc[c + d*x]^2, x], x], x] /;
FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[n, 2] && IntegerQ[2*n]

```

Rule 4133

```

Int[((A_.) + csc[(e_ .) + (f_ .)*(x_)]*(B_ .) + csc[(e_ .) + (f_ .)*(x_)]^2*(C_ .
))*csc[(e_ .) + (f_ .)*(x_)]*(b_ .) + (a_ ), x_Symbol] :> Simp[(-b)*C*Csc[e +
f*x]*(Cot[e + f*x]/(2*f)), x] + Dist[1/2, Int[Simp[2*A*a + (2*B*a + b*(2*A
+ C))*Csc[e + f*x] + 2*(a*C + B*b)*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, e, f, A, B, C}, x]

```

Rule 4512

```

Int[Cot[(c_ .) + (d_ .)*(x_)]*Csc[(c_ .) + (d_ .)*(x_)]*(Csc[(c_ .) + (d_ .)*(x_ )
]*(b_ .) + (a_ ))^(n_ .)*(e_ .) + (f_ .)*(x_ ))^(m_ .), x_Symbol] :> Simp[(-(e +
f*x)^m)*((a + b*Csc[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[f*(m/(b*d*(n
+ 1))), Int[(e + f*x)^(m - 1)*(a + b*Csc[c + d*x])^(n + 1), x], x] /; Fre
eQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]

```

Rule 5367

```

Int[((a_ .) + ArcCsc[(c_ .) + (d_ .)*(x_)]*(b_ .))^(p_ .)*((e_ .) + (f_ .)*(x_ ))^(m
_.), x_Symbol] :> Dist[-(d^(m + 1))^{(-1)}, Subst[Int[(a + b*x)^p*Csc[x]*Cot[
x]*(d*e - c*f + f*Csc[x])^m, x], x, ArcCsc[c + d*x]], x] /; FreeQ[{a, b, c,
d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]

```

Rubi steps

$$\begin{aligned}
\int x^3 \csc^{-1}(a + bx) dx &= -\frac{\text{Subst}(\int x \cot(x) \csc(x)(-a + \csc(x))^3 dx, x, \csc^{-1}(a + bx))}{b^4} \\
&= \frac{1}{4}x^4 \csc^{-1}(a + bx) - \frac{\text{Subst}(\int (-a + \csc(x))^4 dx, x, \csc^{-1}(a + bx))}{4b^4} \\
&= \frac{x^2(a + bx)\sqrt{1 - \frac{1}{(a + bx)^2}}}{12b^2} + \frac{1}{4}x^4 \csc^{-1}(a + bx) - \frac{\text{Subst}(\int (-a + \csc(x))(-3a^3 + 17a^2b^2)x^3 dx, x, \csc^{-1}(a + bx))}{4b^4} \\
&= \frac{x^2(a + bx)\sqrt{1 - \frac{1}{(a + bx)^2}}}{12b^2} - \frac{a(a + bx)^2\sqrt{1 - \frac{1}{(a + bx)^2}}}{3b^4} + \frac{1}{4}x^4 \csc^{-1}(a + bx) - \frac{\text{Subst}(\int (-a + \csc(x))(-3a^3 + 17a^2b^2)x^3 dx, x, \csc^{-1}(a + bx))}{4b^4} \\
&= \frac{x^2(a + bx)\sqrt{1 - \frac{1}{(a + bx)^2}}}{12b^2} - \frac{a(a + bx)^2\sqrt{1 - \frac{1}{(a + bx)^2}}}{3b^4} - \frac{a^4 \csc^{-1}(a + bx)}{4b^4} + \frac{1}{4} \\
&= \frac{x^2(a + bx)\sqrt{1 - \frac{1}{(a + bx)^2}}}{12b^2} - \frac{a(a + bx)^2\sqrt{1 - \frac{1}{(a + bx)^2}}}{3b^4} - \frac{a^4 \csc^{-1}(a + bx)}{4b^4} + \frac{1}{4} \\
&= \frac{(2 + 17a^2)(a + bx)\sqrt{1 - \frac{1}{(a + bx)^2}}}{12b^4} + \frac{x^2(a + bx)\sqrt{1 - \frac{1}{(a + bx)^2}}}{12b^2} - \frac{a(a + bx)^2}{12b^2}
\end{aligned}$$

Mathematica [A]

time = 0.19, size = 149, normalized size = 0.96

$$\frac{\sqrt{\frac{-1 + a^2 + 2abx + b^2x^2}{(a + bx)^2}} (2a + 13a^3 + 2bx + 9a^2bx - 3ab^2x^2 + b^3x^3) + 3b^4x^4 \csc^{-1}(a + bx) - 3a^4 \text{ArcSin}\left(\frac{1}{a+bx}\right) - 6a(1 + 2a^2) \log\left((a + bx) \left(1 + \sqrt{\frac{-1 + a^2 + 2abx + b^2x^2}{(a + bx)^2}}\right)\right)}{12b^4}$$

Antiderivative was successfully verified.

[In] `Integrate[x^3*ArcCsc[a + b*x], x]`

[Out] `(Sqrt[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2]*(2*a + 13*a^3 + 2*b*x + 9*a^2*b*x - 3*a*b^2*x^2 + b^3*x^3) + 3*b^4*x^4*ArcCsc[a + b*x] - 3*a^4*ArcSin[(a + b*x)^(-1)] - 6*a*(1 + 2*a^2)*Log[(a + b*x)*(1 + Sqrt[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2])])/(12*b^4)`

Maple [A]

time = 0.31, size = 249, normalized size = 1.61

method	result
--------	--------

	$\frac{\text{arccsc}(bx+a)a^4}{4} - \text{arccsc}(bx+a)a^3(bx+a) + \frac{3\text{arccsc}(bx+a)a^2(bx+a)^2}{2} - \text{arccsc}(bx+a)a(bx+a)^3 + \frac{\text{arccsc}(bx+a)(bx+a)^4}{4} - \frac{3\text{arccsc}(bx+a)a^3(bx+a)^3}{2} + \frac{9\text{arccsc}(bx+a)a^2(bx+a)^4}{4}$
derivativedivides	$\sqrt{(bx+a)^2}$
default	$\sqrt{(bx+a)^2}$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(x^3*arccsc(b*x+a),x,method=_RETURNVERBOSE)
[Out] 1/b^4*(1/4*arccsc(b*x+a)*a^4-arccsc(b*x+a)*a^3*(b*x+a)+3/2*arccsc(b*x+a)*a^2*(b*x+a)^2-arccsc(b*x+a)*a*(b*x+a)^3+1/4*arccsc(b*x+a)*(b*x+a)^4-1/12*((b*x+a)^2-1)^(1/2)*(3*a^4*arctan(1/((b*x+a)^2-1)^(1/2))+12*a^3*ln(b*x+a+((b*x+a)^2-1)^(1/2))-18*a^2*((b*x+a)^2-1)^(1/2)+6*a*(b*x+a)*((b*x+a)^2-1)^(1/2)-(b*x+a)^2*((b*x+a)^2-1)^(1/2)+6*a*ln(b*x+a+((b*x+a)^2-1)^(1/2))-2*((b*x+a)^2-1)^(1/2))/(((b*x+a)^2-1)/(b*x+a)^2)^(1/2)/(b*x+a))
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x^3*arccsc(b*x+a),x, algorithm="maxima")
[Out] 1/4*x^4*arctan2(1, sqrt(b*x + a + 1)*sqrt(b*x + a - 1)) + integrate(1/4*(b^2*x^5 + a*b*x^4)*e^(1/2*log(b*x + a + 1) + 1/2*log(b*x + a - 1))/(b^2*x^2 + 2*a*b*x + a^2 + (b^2*x^2 + 2*a*b*x + a^2 - 1)*e^(log(b*x + a + 1) + log(b*x + a - 1)) - 1), x)
```

Fricas [A]

time = 0.37, size = 129, normalized size = 0.83

$$\frac{3b^4x^4 \text{arccsc}(bx+a) + 6a^4 \arctan\left(-bx - a + \sqrt{b^2x^2 + 2abx + a^2 - 1}\right) + 6(2a^3 + a) \log\left(-bx - a + \sqrt{b^2x^2 + 2abx + a^2 - 1}\right) + \sqrt{b^2x^2 + 2abx + a^2 - 1} (b^2x^2 - 4abx + 13a^2 + 2)}{12b^4}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x^3*arccsc(b*x+a),x, algorithm="fricas")
[Out] 1/12*(3*b^4*x^4*arccsc(b*x + a) + 6*a^4*arctan(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)) + 6*(2*a^3 + a)*log(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)))
```

$$\hat{2} - 1)) + \sqrt{b^2 x^2 + 2 a b x + a^2 - 1} * (b^2 x^2 - 4 a b x + 13 a^2 + 2) / b^4$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x^3 \operatorname{acsc}(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x**3*acsc(b*x+a),x)

[Out] Integral(x**3*acsc(a + b*x), x)

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 300 vs. 2(135) = 270.

time = 0.47, size = 300, normalized size = 1.94

$$-\frac{1}{30} \left| \frac{24 (bx + a)^2 \left(\frac{dx}{ax + b} + \frac{4x^2}{(bx + a)^2} + \frac{4x^3}{(bx + a)^3} - 1\right) \arcsin\left(\frac{1}{\sqrt{(bx + a)^2 - a^2}}\right)}{b^5} - \frac{\left((bx + a)^2 \left(\sqrt{-\frac{1}{(bx + a)^2} + 1}\right)^2 + 12 (bx + a)^2 \sqrt{\left(\sqrt{-\frac{1}{(bx + a)^2} + 1}\right)^2 + 72 (bx + a) a^2 \left(\sqrt{-\frac{1}{(bx + a)^2} + 1}\right) + 9 (bx + a) \left(\sqrt{-\frac{1}{(bx + a)^2} + 1}\right) - 1\right) + 48 (2 a^2 + a) \log\left(-\left(\sqrt{-\frac{1}{(bx + a)^2} + 1}\right) - 1\right) - \frac{9 (8 a^2 + 1) (bx + a)^2 \sqrt{\left(\sqrt{-\frac{1}{(bx + a)^2} + 1}\right)^2 + 12 (bx + a) a^2 \left(\sqrt{-\frac{1}{(bx + a)^2} + 1}\right) + 1}}{(bx + a)^2 \left(\sqrt{-\frac{1}{(bx + a)^2} + 1}\right)^2}}}{b^5} \right|$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^3*arccsc(b*x+a),x, algorithm="giac")

[Out] $-1/96*b*(24*(b*x + a)^4*(4*a/(b*x + a) - 6*a^2/(b*x + a)^2 + 4*a^3/(b*x + a)^3 - 1)*\arcsin(-1/((b*x + a)*(a/(b*x + a) - 1) - a))/b^5 - ((b*x + a)^3*(sqrt(-1/(b*x + a)^2 + 1) - 1)^3 + 12*(b*x + a)^2*a*(sqrt(-1/(b*x + a)^2 + 1) - 1)^2 + 72*(b*x + a)*a^2*(sqrt(-1/(b*x + a)^2 + 1) - 1) + 9*(b*x + a)*(sqrt(-1/(b*x + a)^2 + 1) - 1) + 48*(2*a^3 + a)*log(-(sqrt(-1/(b*x + a)^2 + 1) - 1)*abs(b*x + a)) - (9*(8*a^2 + 1)*(b*x + a)^2*(sqrt(-1/(b*x + a)^2 + 1) - 1)^2 + 12*(b*x + a)*a*(sqrt(-1/(b*x + a)^2 + 1) - 1) + 1)/((b*x + a)^3*(sqrt(-1/(b*x + a)^2 + 1) - 1)^3))/b^5)$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int x^3 \sin\left(\frac{1}{a + bx}\right) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^3*asin(1/(a + b*x)),x)

[Out] int(x^3*asin(1/(a + b*x)), x)

$$\int x^2 \csc^{-1}(a + bx) dx$$

Optimal. Leaf size=116

$$-\frac{5a(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{6b^3} + \frac{x(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{6b^2} + \frac{a^3 \csc^{-1}(a+bx)}{3b^3} + \frac{1}{3}x^3 \csc^{-1}(a+bx) + \dots$$

[Out] $\frac{1}{3}a^3 \operatorname{arccsc}(bx+a)/b^3 + \frac{1}{3}x^3 \operatorname{arccsc}(bx+a) + \frac{1}{6}*(6*a^2+1)*\operatorname{arctanh}((1-1/(bx+a)^2)^{(1/2)})/b^3 - \frac{5}{6}a*(bx+a)*(1-1/(bx+a)^2)^{(1/2)}/b^3 + \frac{1}{6}x*(bx+a)*(1-1/(bx+a)^2)^{(1/2)}/b^2$

Rubi [A]

time = 0.06, antiderivative size = 116, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.600,

Rules used = {5367, 4512, 3867, 3855, 3852, 8}

$$\frac{a^3 \csc^{-1}(a+bx)}{3b^3} + \frac{(6a^2+1) \tanh^{-1}\left(\sqrt{1-\frac{1}{(a+bx)^2}}\right)}{6b^3} - \frac{5a(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{6b^3} + \frac{x(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{6b^2} + \frac{1}{3}x^3 \csc^{-1}(a+bx)$$

Antiderivative was successfully verified.

[In] Int[x^2*ArcCsc[a + b*x], x]

[Out] $\frac{(-5*a*(a+b*x)*\operatorname{Sqrt}[1-(a+b*x)^{-2}])/(6*b^3)}{(6*b^2)} + \frac{(x*(a+b*x)*\operatorname{Sqrt}[1-(a+b*x)^{-2}])/(6*b^2)}{(6*b^3)} + \frac{(a^3*\operatorname{ArcCsc}[a+b*x])/(3*b^3)}{(6*b^3)} + \frac{(x^3*\operatorname{ArcCsc}[a+b*x])/(3)}{(6*b^3)} + \frac{((1+6*a^2)*\operatorname{ArcTanh}[\operatorname{Sqrt}[1-(a+b*x)^{-2}]])/(6*b^3)}{(6*b^3)}$

Rule 8

Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]

Rule 3852

Int[csc[(c_.) + (d_.)*(x_.)]^(n_), x_Symbol] :> Dist[-d^(-1), Subst[Int[ExpandoIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]

Rule 3855

Int[csc[(c_.) + (d_.)*(x_.)], x_Symbol] :> Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rule 3867

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.) + (a_.))^(n_), x_Symbol] :> Simp[(-b^2)*Cot[c + d*x]*((a + b*Csc[c + d*x])^(n - 2)/(d*(n - 1))), x] + Dist[1/(n - 1)

```
, Int[(a + b*Csc[c + d*x])^(n - 3)*Simp[a^3*(n - 1) + (b*(b^2*(n - 2) + 3*a^2*(n - 1)))*Csc[c + d*x] + (a*b^2*(3*n - 4))*Csc[c + d*x]^2, x], x] /;
FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[n, 2] && IntegerQ[2*n]
```

Rule 4512

```
Int[Cot[(c_.) + (d_.)*(x_.)]*Csc[(c_.) + (d_.)*(x_.)]*(Csc[(c_.) + (d_.)*(x_.)]*(b_.) + (a_.)^(n_.)*(e_.) + (f_.)*(x_.))^(m_.), x_Symbol] :> Simplify[(-(e + f*x)^m)*((a + b*Csc[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[f*(m/(b*d*(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Csc[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]
```

Rule 5367

```
Int[((a_.) + ArcCsc[(c_.) + (d_.)*(x_.)]*(b_.))^p*((e_.) + (f_.)*(x_.))^(m_.), x_Symbol] :> Dist[-(d^(m + 1))^(-1), Subst[Int[(a + b*x)^p*Csc[x]*Cot[x]*(d*e - c*f + f*Csc[x])^m, x], x, ArcCsc[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Rubi steps

$$\begin{aligned}
\int x^2 \csc^{-1}(a + bx) dx &= -\frac{\text{Subst}\left(\int x \cot(x) \csc(x) (-a + \csc(x))^2 dx, x, \csc^{-1}(a + bx)\right)}{b^3} \\
&= \frac{1}{3} x^3 \csc^{-1}(a + bx) - \frac{\text{Subst}\left(\int (-a + \csc(x))^3 dx, x, \csc^{-1}(a + bx)\right)}{3b^3} \\
&= \frac{x(a + bx) \sqrt{1 - \frac{1}{(a + bx)^2}}}{6b^2} + \frac{1}{3} x^3 \csc^{-1}(a + bx) - \frac{\text{Subst}\left(\int (-2a^3 + (1 + 6a^2) \csc(x))^2 dx, x, \csc^{-1}(a + bx)\right)}{6b^3} \\
&= \frac{x(a + bx) \sqrt{1 - \frac{1}{(a + bx)^2}}}{6b^2} + \frac{a^3 \csc^{-1}(a + bx)}{3b^3} + \frac{1}{3} x^3 \csc^{-1}(a + bx) + \frac{(5a)\text{Subst}\left(\int (-2a^3 + (1 + 6a^2) \csc(x))^2 dx, x, \csc^{-1}(a + bx)\right)}{18b^5} \\
&= \frac{x(a + bx) \sqrt{1 - \frac{1}{(a + bx)^2}}}{6b^2} + \frac{a^3 \csc^{-1}(a + bx)}{3b^3} + \frac{1}{3} x^3 \csc^{-1}(a + bx) + \frac{(1 + 6a^2) \tan^{-1}\left(\frac{a + bx}{\sqrt{1 - \frac{1}{(a + bx)^2}}}\right)}{6b^2} \\
&= -\frac{5a(a + bx) \sqrt{1 - \frac{1}{(a + bx)^2}}}{6b^3} + \frac{x(a + bx) \sqrt{1 - \frac{1}{(a + bx)^2}}}{6b^2} + \frac{a^3 \csc^{-1}(a + bx)}{3b^3} +
\end{aligned}$$

Mathematica [A]

time = 0.13, size = 129, normalized size = 1.11

$$\frac{(-5a^2 - 4abx + b^2x^2) \sqrt{\frac{-1 + a^2 + 2abx + b^2x^2}{(a + bx)^2}} + 2b^3x^3 \csc^{-1}(a + bx) + 2a^3 \text{ArcSin}\left(\frac{1}{a+bx}\right) + (1 + 6a^2) \log\left((a + bx) \left(1 + \sqrt{\frac{-1 + a^2 + 2abx + b^2x^2}{(a + bx)^2}}\right)\right)}{6b^3}$$

Antiderivative was successfully verified.

[In] `Integrate[x^2*ArcCsc[a + b*x], x]`

[Out] $\frac{((-5a^2 - 4a*b*x + b^2*x^2)*\text{Sqrt}[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2] + 2*b^3*x^3*\text{ArcCsc}[a + b*x] + 2*a^3*\text{ArcSin}[(a + b*x)^{-1}] + (1 + 6*a^2)*\text{Log}[(a + b*x)*(1 + \text{Sqrt}[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2])])/(6*b^3)}$

Maple [A]

time = 0.30, size = 190, normalized size = 1.64

method	result
derivativedivides	$\frac{-\frac{\text{arccsc}(bx+a)a^3}{3} + \text{arccsc}(bx+a)a^2(bx+a) - \text{arccsc}(bx+a)a(bx+a)^2 + \frac{\text{arccsc}(bx+a)(bx+a)^3}{3} + \dots}{\sqrt{(bx+a)^2 - 1}}$
default	$\frac{-\frac{\text{arccsc}(bx+a)a^3}{3} + \text{arccsc}(bx+a)a^2(bx+a) - \text{arccsc}(bx+a)a(bx+a)^2 + \frac{\text{arccsc}(bx+a)(bx+a)^3}{3} + \dots}{\sqrt{(bx+a)^2 - 1}}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x^2*arccsc(b*x+a), x, method=_RETURNVERBOSE)`

[Out] $\frac{1}{b^3} \cdot (-\frac{1}{3} \text{arccsc}(b*x+a) * a^3 + \text{arccsc}(b*x+a) * a^2 * (b*x+a) - \text{arccsc}(b*x+a) * a * (b*x+a)^2 + \frac{1}{3} \text{arccsc}(b*x+a) * (b*x+a)^3 + \frac{1}{6} ((b*x+a)^2 - 1)^{(1/2)} * (2*a^3 * \text{arctan}(1/((b*x+a)^2 - 1)^{(1/2)}) + 6*a^2 * \ln(b*x+a + ((b*x+a)^2 - 1)^{(1/2)}) - 6*a * ((b*x+a)^2 - 1)^{(1/2)} * (b*x+a) * ((b*x+a)^2 - 1)^{(1/2)} + \ln(b*x+a + ((b*x+a)^2 - 1)^{(1/2)})) / (((b*x+a)^2 - 1) / (b*x+a)^2)^{(1/2)} / (b*x+a))$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^2*arccsc(b*x+a), x, algorithm="maxima")`

[Out] $\frac{1}{3}x^3 \operatorname{arctan}(1, \sqrt{bx+a+1})\sqrt{bx+a-1} + \operatorname{integrate}(1/3*(b^2x^4 + a*b*x^3)*e^{(1/2*\log(b*x+a+1) + 1/2*\log(b*x+a-1))/(b^2*x^2 + 2*a*b*x + a^2 + (b^2*x^2 + 2*a*b*x + a^2 - 1)*e^{(\log(b*x+a+1) + \log(b*x+a-1)) - 1}), x)$

Fricas [A]

time = 0.37, size = 117, normalized size = 1.01

$$\frac{2b^3x^3 \operatorname{arccsc}(bx+a) - 4a^3 \arctan(-bx-a+\sqrt{b^2x^2+2abx+a^2-1}) - (6a^2+1)\log(-bx-a+\sqrt{b^2x^2+2abx+a^2-1}) + \sqrt{b^2x^2+2abx+a^2-1}(bx-5a)}{6b^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^2*arccsc(b*x+a), x, algorithm="fricas")`

[Out] $\frac{1}{6}*(2*b^3*x^3*arccsc(b*x+a) - 4*a^3*arctan(-b*x-a+\sqrt{b^2*x^2+2*a*b*x+a^2-1}) - (6*a^2+1)*log(-b*x-a+\sqrt{b^2*x^2+2*a*b*x+a^2-1}) + \sqrt{b^2*x^2+2*a*b*x+a^2-1}*(b*x-5*a))/b^3$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x^2 \operatorname{acsc}(a+bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x**2*acsc(b*x+a), x)`

[Out] `Integral(x**2*acsc(a+b*x), x)`

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 203 vs. 2(100) = 200.

time = 0.47, size = 203, normalized size = 1.75

$$-\frac{1}{24}b \left(\frac{\frac{8(bx+a)^3 \left(\frac{3a}{bx+a} - \frac{3a^2}{(bx+a)^2} - 1\right) \arcsin\left(\frac{1}{(bx+a)(\frac{3a}{bx+a}-1)-a}\right)}{b^4} + \frac{(bx+a)^2 \left(\sqrt{-\frac{1}{(bx+a)^2}+1} - 1\right)^2 + 12(bx+a)a \left(\sqrt{-\frac{1}{(bx+a)^2}+1} - 1\right) + 4(6a^2+1)\log\left(-\left(\sqrt{-\frac{1}{(bx+a)^2}+1} - 1\right)|bx+a|\right) - \frac{12(bx+a)a \left(\sqrt{-\frac{1}{(bx+a)^2}+1} - 1\right)}{\left(\frac{1}{(bx+a)^2} \left(\sqrt{-\frac{1}{(bx+a)^2}+1} - 1\right)\right)^2}}{b^4} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^2*arccsc(b*x+a), x, algorithm="giac")`

[Out] $\frac{-1/24*b*(8*(b*x+a)^3*(3*a/(b*x+a) - 3*a^2/(b*x+a)^2 - 1)*\arcsin(-1/((b*x+a)*(a/(b*x+a) - 1) - a))/b^4 + ((b*x+a)^2*(\sqrt{-1/(b*x+a)^2 + 1} - 1)^2 + 12*(b*x+a)*a*(\sqrt{-1/(b*x+a)^2 + 1} - 1) + 4*(6*a^2 + 1)*\log(-(\sqrt{-1/(b*x+a)^2 + 1} - 1)*\abs(b*x+a)) - (12*(b*x+a)*a*(\sqrt{-1/(b*x+a)^2 + 1} - 1) + 1)/((b*x+a)^2*(\sqrt{-1/(b*x+a)^2 + 1} - 1)^2))/b^4}$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int x^2 \arcsin\left(\frac{1}{a + b x}\right) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x^2*asin(1/(a + b*x)),x)`[Out] `int(x^2*asin(1/(a + b*x)), x)`

3.20 $\int x \csc^{-1}(a + bx) dx$

Optimal. Leaf size=79

$$\frac{(a + bx)\sqrt{1 - \frac{1}{(a + bx)^2}}}{2b^2} - \frac{a^2 \csc^{-1}(a + bx)}{2b^2} + \frac{1}{2}x^2 \csc^{-1}(a + bx) - \frac{a \tanh^{-1}\left(\sqrt{1 - \frac{1}{(a + bx)^2}}\right)}{b^2}$$

[Out] $-1/2*a^2*\text{arccsc}(b*x+a)/b^2+1/2*x^2*\text{arccsc}(b*x+a)-a*\text{arctanh}((1-1/(b*x+a)^2)^(1/2))/b^2+1/2*(b*x+a)*(1-1/(b*x+a)^2)^(1/2)/b^2$

Rubi [A]

time = 0.04, antiderivative size = 79, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.750, Rules used = {5367, 4512, 3858, 3855, 3852, 8}

$$-\frac{a^2 \csc^{-1}(a + bx)}{2b^2} + \frac{(a + bx)\sqrt{1 - \frac{1}{(a + bx)^2}}}{2b^2} - \frac{a \tanh^{-1}\left(\sqrt{1 - \frac{1}{(a + bx)^2}}\right)}{b^2} + \frac{1}{2}x^2 \csc^{-1}(a + bx)$$

Antiderivative was successfully verified.

[In] Int[x*ArcCsc[a + b*x], x]

[Out] $((a + b*x)*\text{Sqrt}[1 - (a + b*x)^{-2}])/(2*b^2) - (a^2*\text{ArcCsc}[a + b*x])/(2*b^2) + (x^2*\text{ArcCsc}[a + b*x])/2 - (a*\text{ArcTanh}[\text{Sqrt}[1 - (a + b*x)^{-2}]])/b^2$

Rule 8

Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]

Rule 3852

Int[csc[(c_.) + (d_.)*(x_.)]^(n_), x_Symbol] :> Dist[-d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]]

Rule 3855

Int[csc[(c_.) + (d_.)*(x_.)], x_Symbol] :> Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rule 3858

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.) + (a_.))^2, x_Symbol] :> Simp[a^2*x, x] + (Dist[2*a*b, Int[Csc[c + d*x], x], x] + Dist[b^2, Int[Csc[c + d*x]^2, x], x]) /; FreeQ[{a, b, c, d}, x]

Rule 4512

```
Int[Cot[(c_.) + (d_.)*(x_)]*Csc[(c_.) + (d_.)*(x_)]*(Csc[(c_.) + (d_.)*(x_)]
]*((b_.) + (a_.))^(n_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] :> Simp[(-(e +
f*x)^m)*((a + b*Csc[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[f*(m/(b*d*(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Csc[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]
```

Rule 5367

```
Int[((a_.) + ArcCsc[(c_.) + (d_.)*(x_)]*(b_.))^p*((e_.) + (f_.)*(x_))^(m - .), x_Symbol] :> Dist[-(d^(m + 1))^{(-1)}, Subst[Int[(a + b*x)^p*Csc[x]*Cot[x]*(d*e - c*f + f*Csc[x])^m, x], x, ArcCsc[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Rubi steps

$$\begin{aligned} \int x \csc^{-1}(a + bx) dx &= -\frac{\text{Subst}\left(\int x \cot(x) \csc(x)(-a + \csc(x)) dx, x, \csc^{-1}(a + bx)\right)}{b^2} \\ &= \frac{1}{2} x^2 \csc^{-1}(a + bx) - \frac{\text{Subst}\left(\int (-a + \csc(x))^2 dx, x, \csc^{-1}(a + bx)\right)}{2b^2} \\ &= -\frac{a^2 \csc^{-1}(a + bx)}{2b^2} + \frac{1}{2} x^2 \csc^{-1}(a + bx) - \frac{\text{Subst}\left(\int \csc^2(x) dx, x, \csc^{-1}(a + bx)\right)}{2b^2} + \\ &= -\frac{a^2 \csc^{-1}(a + bx)}{2b^2} + \frac{1}{2} x^2 \csc^{-1}(a + bx) - \frac{a \tanh^{-1}\left(\sqrt{1 - \frac{1}{(a + bx)^2}}\right)}{b^2} + \text{Subst}\left(\right. \\ &\quad \left. -\frac{(a + bx) \sqrt{1 - \frac{1}{(a + bx)^2}}}{2b^2} - \frac{a^2 \csc^{-1}(a + bx)}{2b^2} + \frac{1}{2} x^2 \csc^{-1}(a + bx) - \frac{a \tanh^{-1}\left(\sqrt{1 - \frac{1}{(a + bx)^2}}\right)}{b^2}\right) \end{aligned}$$

Mathematica [A]

time = 0.08, size = 110, normalized size = 1.39

$$\frac{(a + bx) \sqrt{\frac{-1 + a^2 + 2abx + b^2x^2}{(a + bx)^2}} + b^2x^2 \csc^{-1}(a + bx) - a^2 \text{ArcSin}\left(\frac{1}{a + bx}\right) - 2a \log\left((a + bx) \left(1 + \sqrt{\frac{-1 + a^2 + 2abx + b^2x^2}{(a + bx)^2}}\right)\right)}{2b^2}$$

Antiderivative was successfully verified.

[In] `Integrate[x*ArcCsc[a + b*x], x]`

[Out] $((a + b*x)*\text{Sqrt}[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2] + b^2*x^2*\text{ArcCs}\text{c}[a + b*x] - a^2*\text{ArcSin}[(a + b*x)^{(-1)}] - 2*a*\text{Log}[(a + b*x)*(1 + \text{Sqrt}[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2])])/(2*b^2)$

Maple [A]

time = 0.30, size = 108, normalized size = 1.37

method	result
derivativedivides	$\frac{\frac{\arccsc(bx+a)(bx+a)^2}{2} - \arccsc(bx+a)a(bx+a) - \frac{\sqrt{(bx+a)^2 - 1} \left(-\sqrt{(bx+a)^2 - 1} + 2a \ln(bx+a) \sqrt{(bx+a)^2 - 1} \right)}{2(bx+a) \sqrt{\frac{(bx+a)^2 - 1}{(bx+a)^2}}}{b^2}$
default	$\frac{\frac{\arccsc(bx+a)(bx+a)^2}{2} - \arccsc(bx+a)a(bx+a) - \frac{\sqrt{(bx+a)^2 - 1} \left(-\sqrt{(bx+a)^2 - 1} + 2a \ln(bx+a) \sqrt{(bx+a)^2 - 1} \right)}{2(bx+a) \sqrt{\frac{(bx+a)^2 - 1}{(bx+a)^2}}}{b^2}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x*arccsc(b*x+a),x,method=_RETURNVERBOSE)`
[Out]
$$\frac{1}{b^2} \cdot \frac{1}{2} \cdot \frac{(1/2 \cdot \arccsc(b*x+a) \cdot (b*x+a)^2 - \arccsc(b*x+a) \cdot a \cdot (b*x+a) - 1/2 \cdot ((b*x+a)^2 - 1)^{(1/2)} \cdot (-((b*x+a)^2 - 1)^{(1/2)} + 2*a \cdot \ln(b*x+a + ((b*x+a)^2 - 1)^{(1/2)}))}{(b*x+a) \cdot ((b*x+a)^2 - 1)^{(1/2)}}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x*arccsc(b*x+a),x, algorithm="maxima")`
[Out]
$$\frac{1}{2} \cdot x^2 \cdot \arctan2(1, \sqrt{b*x + a + 1}) \cdot \sqrt{b*x + a - 1} + \text{integrate}(1/2 \cdot (b^2 \cdot x^3 + a \cdot b \cdot x^2) \cdot e^{-(1/2 \cdot \log(b*x + a + 1) + 1/2 \cdot \log(b*x + a - 1)) / (b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + (b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 - 1) \cdot e^{(\log(b*x + a + 1) + \log(b*x + a - 1)) - 1}), x)$$

Fricas [A]

time = 0.39, size = 102, normalized size = 1.29

$$\frac{b^2 x^2 \arccsc(bx + a) + 2 a^2 \arctan\left(-bx - a + \sqrt{b^2 x^2 + 2 abx + a^2 - 1}\right) + 2 a \log\left(-bx - a + \sqrt{b^2 x^2 + 2 abx + a^2 - 1}\right) + \sqrt{b^2 x^2 + 2 abx + a^2 - 1}}{2 b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x*arccsc(b*x+a),x, algorithm="fricas")`
[Out]
$$\frac{1}{2} \cdot (b^2 \cdot x^2 \cdot \arccsc(b*x + a) + 2 \cdot a^2 \cdot \arctan(-b*x - a + \sqrt{b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 - 1}) + 2 \cdot a \cdot \log(-b*x - a + \sqrt{b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 - 1})) + \sqrt{b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 - 1} / b^2$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x \operatorname{acsc}(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x*acsc(b*x+a),x)`[Out] `Integral(x*acsc(a + b*x), x)`**Giac [A]**

time = 0.47, size = 134, normalized size = 1.70

$$-\frac{1}{4} b \left(\frac{2 (bx + a)^2 \left(\frac{2 a}{bx + a} - 1\right) \arcsin \left(-\frac{1}{(bx + a) \left(\frac{2 a}{bx + a} - 1\right) - a}\right)}{b^3} - \frac{(bx + a) \left(\sqrt{-\frac{1}{(bx + a)^2} + 1} - 1\right) + 4 a \log \left(-\left(\sqrt{-\frac{1}{(bx + a)^2} + 1} - 1\right) |bx + a|\right)}{b^3} - \frac{(bx + a) \left(\sqrt{-\frac{1}{(bx + a)^2} + 1} - 1\right)}{b^3} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x*arccsc(b*x+a),x, algorithm="giac")`[Out] $-1/4*b*(2*(b*x + a)^2*(2*a/(b*x + a) - 1)*\arcsin(-1/((b*x + a)*(a/(b*x + a) - 1) - a))/b^3 - ((b*x + a)*(\sqrt(-1/(b*x + a)^2 + 1) - 1) + 4*a*\log(-(\sqrt(-1/(b*x + a)^2 + 1) - 1)*abs(b*x + a)) - 1/((b*x + a)*(\sqrt(-1/(b*x + a)^2 + 1) - 1)))/b^3)$ **Mupad [F]**

time = 0.00, size = -1, normalized size = -0.01

$$\int x \operatorname{asin}\left(\frac{1}{a + bx}\right) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x*asin(1/(a + b*x)),x)`[Out] `int(x*asin(1/(a + b*x)), x)`

$$\mathbf{3.21} \quad \int \csc^{-1}(a + bx) dx$$

Optimal. Leaf size=36

$$\frac{(a + bx) \csc^{-1}(a + bx)}{b} + \frac{\tanh^{-1}\left(\sqrt{1 - \frac{1}{(a + bx)^2}}\right)}{b}$$

[Out] $(b*x+a)*\text{arccsc}(b*x+a)/b+\text{arctanh}((1-1/(b*x+a)^2)^{(1/2)})/b$

Rubi [A]

time = 0.02, antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 6, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.833, Rules used = {5359, 379, 272, 65, 212}

$$\frac{(a + bx) \csc^{-1}(a + bx)}{b} + \frac{\tanh^{-1}\left(\sqrt{1 - \frac{1}{(a + bx)^2}}\right)}{b}$$

Antiderivative was successfully verified.

[In] Int[ArcCsc[a + b*x], x]

[Out] $((a + b*x)*\text{ArcCsc}[a + b*x])/b + \text{ArcTanh}[\text{Sqrt}[1 - (a + b*x)^{-2}]]/b$

Rule 65

```
Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] :> With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ[
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 212

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

Rule 272

```
Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] :> Dist[1/n, Subst[
Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b,
m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 379

```
Int[(u_)^(m_)*((a_) + (b_)*(v_)^(n_))^(p_), x_Symbol] :> Dist[u^m/(Coeff
icient[v, x, 1]*v^m), Subst[Int[x^m*(a + b*x^n)^p, x], x, v], x] /; FreeQ[{a, b, m, n, p}, x] && LinearPairQ[u, v, x]
```

Rule 5359

```
Int[ArcCsc[(c_) + (d_)*(x_)], x_Symbol] :> Simp[(c + d*x)*(ArcCsc[c + d*x]
/d), x] + Int[1/((c + d*x)*Sqrt[1 - 1/(c + d*x)^2]), x] /; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned}
\int \csc^{-1}(a + bx) dx &= \frac{(a + bx) \csc^{-1}(a + bx)}{b} + \int \frac{1}{(a + bx) \sqrt{1 - \frac{1}{(a + bx)^2}}} dx \\
&= \frac{(a + bx) \csc^{-1}(a + bx)}{b} + \frac{\text{Subst}\left(\int \frac{1}{\sqrt{1 - \frac{1}{x^2}}} dx, x, a + bx\right)}{b} \\
&= \frac{(a + bx) \csc^{-1}(a + bx)}{b} - \frac{\text{Subst}\left(\int \frac{1}{\sqrt{1 - x}} dx, x, \frac{1}{(a+bx)^2}\right)}{2b} \\
&= \frac{(a + bx) \csc^{-1}(a + bx)}{b} + \frac{\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \sqrt{1 - \frac{1}{(a+bx)^2}}\right)}{b} \\
&= \frac{(a + bx) \csc^{-1}(a + bx)}{b} + \frac{\tanh^{-1}\left(\sqrt{1 - \frac{1}{(a+bx)^2}}\right)}{b}
\end{aligned}$$

Mathematica [B] Leaf count is larger than twice the leaf count of optimal. 336 vs. 2(36) = 72.

time = 0.18, size = 336, normalized size = 9.33

$$x \csc^{-1}(a + bx) + \frac{(a + bx) \sqrt{\frac{-1 + a^2 + 2ax + b^2x^2}{(a + bx)^2}} \left(2a\left(b + \sqrt{b^2}x - \sqrt{-1 + a^2 + 2ax + b^2x^2}\right) \operatorname{ArcTan}\left(a + \sqrt{b^2}x + \sqrt{-1 + a^2 + 2ax + b^2x^2}\right) + 2a\left(-b + \sqrt{b^2}\right) \operatorname{ArcTan}\left(a - \sqrt{b^2}x + \sqrt{-1 + a^2 + 2ax + b^2x^2}\right) - \sqrt{b^2} \log\left(-a + \sqrt{b^2}x + \sqrt{-1 + a^2 + 2ax + b^2x^2}\right) + b \log\left(a - \sqrt{b^2}x + \sqrt{-1 + a^2 + 2ax + b^2x^2}\right) - \sqrt{b^2} \log\left(a - \sqrt{b^2}x + \sqrt{-1 + a^2 + 2ax + b^2x^2}\right) - b \log\left(ab^2 + (b^2)^{3/2}x - b^2 \sqrt{-1 + a^2 + 2ax + b^2x^2}\right)\right)}{2b \sqrt{-1 + a^2 + 2ax + b^2x^2}}$$

Antiderivative was successfully verified.

[In] `Integrate[ArcCsc[a + b*x], x]`

[Out] $x \operatorname{ArcCsc}[a + b*x] + ((a + b*x)*\operatorname{Sqrt}[-1 + a^2 + 2*a*b*x + b^2*x^2]/(a + b*x)^2)*(2*a*(b + \operatorname{Sqrt}[b^2])* \operatorname{ArcTan}[a + \operatorname{Sqrt}[b^2]*x] - \operatorname{Sqrt}[-1 + a^2 + 2*a*b*x$

$$+ b^{2*x^2}] + 2*a*(-b + \sqrt{b^2})*\text{ArcTan}[a - \sqrt{b^2}*x + \sqrt{-1 + a^2 + 2*a*b*x + b^{2*x^2}}] - \sqrt{b^2}*\text{Log}[-a - \sqrt{b^2}*x + \sqrt{-1 + a^2 + 2*a*b*x + b^{2*x^2}}] + b*\text{Log}[a - \sqrt{b^2}*x + \sqrt{-1 + a^2 + 2*a*b*x + b^{2*x^2}}] - \sqrt{b^2}*\text{Log}[a - \sqrt{b^2}*x + \sqrt{-1 + a^2 + 2*a*b*x + b^{2*x^2}}] - b*\text{Log}[a*b^2 + (b^2)^{(3/2)}*x - b^{2*}\sqrt{-1 + a^2 + 2*a*b*x + b^{2*x^2}}])/(2*b^{2*}\sqrt{-1 + a^2 + 2*a*b*x + b^{2*x^2}})$$

Maple [A]

time = 0.07, size = 43, normalized size = 1.19

method	result	size
derivativedivides	$\frac{\text{arccsc}(bx+a)(bx+a)+\ln\left(bx+a+(bx+a)\sqrt{1-\frac{1}{(bx+a)^2}}\right)}{b}$	43
default	$\frac{\text{arccsc}(bx+a)(bx+a)+\ln\left(bx+a+(bx+a)\sqrt{1-\frac{1}{(bx+a)^2}}\right)}{b}$	43

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(arccsc(b*x+a),x,method=_RETURNVERBOSE)`[Out] $1/b*(\text{arccsc}(b*x+a)*(b*x+a)+\ln(b*x+a+(b*x+a)*(1-1/(b*x+a)^2)^(1/2)))$ **Maxima [A]**

time = 0.26, size = 55, normalized size = 1.53

$$\frac{2(bx+a)\text{arccsc}(bx+a)+\log\left(\sqrt{-\frac{1}{(bx+a)^2}+1}+1\right)-\log\left(-\sqrt{-\frac{1}{(bx+a)^2}+1}+1\right)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(b*x+a),x, algorithm="maxima")`[Out] $1/2*(2*(b*x+a)*\text{arccsc}(b*x+a)+\log(\sqrt{-1/(b*x+a)^2+1}+1)-\log(-\sqrt{-1/(b*x+a)^2+1}+1))/b$ **Fricas [B]** Leaf count of result is larger than twice the leaf count of optimal. 75 vs. $2(34) = 68$.

time = 0.37, size = 75, normalized size = 2.08

$$\frac{bx\text{arccsc}(bx+a)-2a\arctan\left(-bx-a+\sqrt{b^2x^2+2abx+a^2-1}\right)-\log\left(-bx-a+\sqrt{b^2x^2+2abx+a^2-1}\right)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(b*x+a),x, algorithm="fricas")`

[Out] $(b*x*\operatorname{arccsc}(b*x + a) - 2*a*\operatorname{arctan}(-b*x - a + \sqrt{b^2*x^2 + 2*a*b*x + a^2 - 1}) - \log(-b*x - a + \sqrt{b^2*x^2 + 2*a*b*x + a^2 - 1}))/b$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \operatorname{acsc}(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(acsc(b*x+a),x)`

[Out] `Integral(acsc(a + b*x), x)`

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 81 vs. $2(34) = 68$.
time = 0.46, size = 81, normalized size = 2.25

$$\frac{1}{2} b \left(\frac{2(bx + a) \arcsin\left(\frac{1}{(bx+a)\left(\frac{a}{bx+a}-1\right)-a}\right)}{b^2} + \frac{\log\left(\sqrt{-\frac{1}{(bx+a)^2}+1} + 1\right) - \log\left(-\sqrt{-\frac{1}{(bx+a)^2}+1} + 1\right)}{b^2} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(b*x+a),x, algorithm="giac")`

[Out] $\frac{1}{2}b*(2*(b*x + a)*\arcsin(-1/((b*x + a)*(a/(b*x + a) - 1) - a))/b^2 + (\log(\sqrt{-1/(b*x + a)^2 + 1} + 1) - \log(-\sqrt{-1/(b*x + a)^2 + 1} + 1))/b^2)$

Mupad [B]

time = 0.88, size = 33, normalized size = 0.92

$$\frac{\operatorname{atanh}\left(\frac{1}{\sqrt{1 - \frac{1}{(a + b x)^2}}}\right) + \operatorname{asin}\left(\frac{1}{a + b x}\right)(a + b x)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(asin(1/(a + b*x)),x)`

[Out] `(\operatorname{atanh}(1/(1 - 1/(a + b*x)^2)^{(1/2)}) + \operatorname{asin}(1/(a + b*x))*(a + b*x))/b`

$$\mathbf{3.22} \quad \int \frac{\csc^{-1}(a+bx)}{x} dx$$

Optimal. Leaf size=210

$$\csc^{-1}(a+bx) \log \left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{1 - \sqrt{1 - a^2}} \right) + \csc^{-1}(a+bx) \log \left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{1 + \sqrt{1 - a^2}} \right) - \csc^{-1}(a+bx) \log \left(1 - e^{2i \csc^{-1}(a+bx)} \right)$$

[Out] $-\text{arccsc}(b*x+a)*\ln(1-(I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))^2)+\text{arccsc}(b*x+a)*\ln(1+I*a*(I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))/(1-(-a^2+1)^(1/2)))+\text{arccsc}(b*x+a)*\ln(1+I*a*(I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))/(1+(-a^2+1)^(1/2)))+1/2*I*\text{polylog}(2,(I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))^2)-I*\text{polylog}(2,-I*a*(I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))/(1-(-a^2+1)^(1/2))-I*\text{polylog}(2,-I*a*(I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))/(1+(-a^2+1)^(1/2)))$

Rubi [A]

time = 0.21, antiderivative size = 210, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 8, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.800$, Rules used = {5367, 4648, 4625, 3798, 2221, 2317, 2438, 4615}

$$-i\text{Li}_2\left(-\frac{iae^{i \csc^{-1}(a+bx)}}{1 - \sqrt{1 - a^2}}\right) - i\text{Li}_2\left(-\frac{iae^{i \csc^{-1}(a+bx)}}{\sqrt{1 - a^2} + 1}\right) + \csc^{-1}(a + bx) \log \left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{1 - \sqrt{1 - a^2}} \right) + \csc^{-1}(a + bx) \log \left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{\sqrt{1 - a^2} + 1} \right) + \frac{1}{2}i\text{Li}_2\left(e^{2i \csc^{-1}(a+bx)}\right) - \csc^{-1}(a + bx) \log \left(1 - e^{2i \csc^{-1}(a+bx)} \right)$$

Antiderivative was successfully verified.

[In] Int[ArcCsc[a + b*x]/x, x]

[Out] $\text{ArcCsc}[a + b*x]*\text{Log}[1 + (I*a*E^(\text{ArcCsc}[a + b*x]))/(1 - \text{Sqrt}[1 - a^2])] + \text{ArcCsc}[a + b*x]*\text{Log}[1 + (I*a*E^(\text{ArcCsc}[a + b*x]))/(1 + \text{Sqrt}[1 - a^2])] - \text{ArcCsc}[a + b*x]*\text{Log}[1 - E^((2*I)*\text{ArcCsc}[a + b*x])] - I*\text{PolyLog}[2, ((-I)*a*E^(\text{ArcCsc}[a + b*x]))/(1 - \text{Sqrt}[1 - a^2])] - I*\text{PolyLog}[2, ((-I)*a*E^(\text{ArcCsc}[a + b*x]))/(1 + \text{Sqrt}[1 - a^2])] + (I/2)*\text{PolyLog}[2, E^((2*I)*\text{ArcCsc}[a + b*x])]$

Rule 2221

```
Int[((F_)^((g_.)*(e_.) + (f_.)*(x_.)))^(n_.)*((c_.) + (d_.)*(x_.))^(m_.))/((a_) + (b_.)*(F_)^((g_.)*(e_.) + (f_.)*(x_.)))^(n_.)), x_Symbol] :> Simplify[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2317

```
Int[Log[(a_) + (b_.)*(F_)^((e_.)*(c_.) + (d_.)*(x_.)))^(n_.)], x_Symbol] :> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2438

```
Int[Log[(c_)*(d_) + (e_)*(x_)^(n_)]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 3798

```
Int[((c_) + (d_)*(x_))^(m_)*tan[(e_) + Pi*(k_*) + (f_*)*(x_)], x_Symbol] :> Simp[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Dist[2*I, Int[(c + d*x)^m *E^(2*I*k*Pi)*(E^(2*I*(e + f*x))/(1 + E^(2*I*k*Pi)*E^(2*I*(e + f*x)))), x], x] /; FreeQ[{c, d, e, f}, x] && IntegerQ[4*k] && IGtQ[m, 0]
```

Rule 4615

```
Int[(Cos[(c_) + (d_)*(x_)]*((e_) + (f_*)*(x_))^(m_))/((a_) + (b_))*Sin[(c_) + (d_)*(x_)], x_Symbol] :> Simp[(-I)*((e + f*x)^(m + 1)/(b*f*(m + 1))), x] + (Int[(e + f*x)^m*(E^(I*(c + d*x))/(a - Rt[a^2 - b^2, 2] - I*b*E^(I*(c + d*x)))), x] + Int[(e + f*x)^m*(E^(I*(c + d*x))/(a + Rt[a^2 - b^2, 2] - I*b*E^(I*(c + d*x)))), x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && PosQ[a^2 - b^2]
```

Rule 4625

```
Int[(Cot[(c_) + (d_)*(x_)]^(n_)*((e_) + (f_*)*(x_))^(m_))/((a_) + (b_))*Sin[(c_) + (d_)*(x_)], x_Symbol] :> Dist[1/a, Int[(e + f*x)^m*Cot[c + d*x]^n, x], x] - Dist[b/a, Int[(e + f*x)^m*Cos[c + d*x]*(Cot[c + d*x]^(n - 1)/(a + b*Sin[c + d*x])), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rule 4648

```
Int[((e_) + (f_*)*(x_))^(m_)*(F_)[(c_) + (d_)*(x_)]^(n_)*(G_)[(c_) + (d_)*(x_)]^(p_)/((Csc[(c_) + (d_)*(x_)]*(b_) + (a_)), x_Symbol) :> Int[(e + f*x)^m*Sin[c + d*x]*F[c + d*x]^n*(G[c + d*x]^p/(b + a*Sin[c + d*x])), x] /; FreeQ[{a, b, c, d, e, f}, x] && TrigQ[F] && TrigQ[G] && IntegersQ[m, n, p]
```

Rule 5367

```
Int[((a_) + ArcCsc[(c_) + (d_)*(x_)]*(b_))^(p_)*((e_) + (f_*)*(x_))^(m_), x_Symbol] :> Dist[-(d^(m + 1))^(-1), Subst[Int[(a + b*x)^p*Csc[x]*Cot[x]*(d*e - c*f + f*Csc[x])^m, x], x, ArcCsc[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Rubi steps

$$\begin{aligned}
\int \frac{\csc^{-1}(a+bx)}{x} dx &= -\text{Subst}\left(\int \frac{x \cot(x) \csc(x)}{-a + \csc(x)} dx, x, \csc^{-1}(a+bx)\right) \\
&= -\text{Subst}\left(\int \frac{x \cot(x)}{1 - a \sin(x)} dx, x, \csc^{-1}(a+bx)\right) \\
&= -\left(a \text{Subst}\left(\int \frac{x \cos(x)}{1 - a \sin(x)} dx, x, \csc^{-1}(a+bx)\right)\right) - \text{Subst}\left(\int x \cot(x) dx, x, \csc^{-1}(a+bx)\right) \\
&= 2i \text{Subst}\left(\int \frac{e^{2ix} x}{1 - e^{2ix}} dx, x, \csc^{-1}(a+bx)\right) - a \text{Subst}\left(\int \frac{e^{ix} x}{1 - \sqrt{1 - a^2} + iae^{ix}} dx, x, \csc^{-1}(a+bx)\right) \\
&= \csc^{-1}(a+bx) \log\left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{1 - \sqrt{1 - a^2}}\right) + \csc^{-1}(a+bx) \log\left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{1 + \sqrt{1 - a^2}}\right) - c \\
&= \csc^{-1}(a+bx) \log\left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{1 - \sqrt{1 - a^2}}\right) + \csc^{-1}(a+bx) \log\left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{1 + \sqrt{1 - a^2}}\right) - c \\
&= \csc^{-1}(a+bx) \log\left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{1 - \sqrt{1 - a^2}}\right) + \csc^{-1}(a+bx) \log\left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{1 + \sqrt{1 - a^2}}\right) - c
\end{aligned}$$

Mathematica [A]

time = 0.29, size = 375, normalized size = 1.79

$$\frac{1}{2} \left((x - 2 \operatorname{acsc}^2(a + bx))^2 - 2i \operatorname{AcBis} \left(\frac{\sqrt{1 - a^2}}{\sqrt{1 - b^2}}, \operatorname{ArcTan} \left(\frac{2 + a \operatorname{cot}(2x) + 2 \operatorname{csc}(2x) + b \operatorname{cot}(2x)}{\sqrt{1 - b^2}} \right) \right) - \left(x - 2 \operatorname{acsc}^2(a + bx) + 4 \operatorname{AcBis} \left(\frac{\sqrt{1 - a^2}}{\sqrt{1 - b^2}}, \operatorname{ArcTan} \left(\frac{i(-1 + \sqrt{1 - a^2}) e^{i \operatorname{csc}^{-1}(a + bx)}}{a} \right) \right) \right) \operatorname{Log} \left(1 + \frac{i(-1 + \sqrt{1 - a^2}) e^{i \operatorname{csc}^{-1}(a + bx)}}{a} \right) - \left(x - 2 \operatorname{acsc}^2(a + bx) - 4 \operatorname{AcBis} \left(\frac{\sqrt{1 - a^2}}{\sqrt{1 - b^2}}, \operatorname{ArcTan} \left(\frac{i(-1 + \sqrt{1 - a^2}) e^{i \operatorname{csc}^{-1}(a + bx)}}{a} \right) \right) \right) \operatorname{Log} \left(1 - \frac{i(-1 + \sqrt{1 - a^2}) e^{i \operatorname{csc}^{-1}(a + bx)}}{a} \right) - \operatorname{Im} \left(\operatorname{acsc}^2(a + bx) \operatorname{Log} \left(\frac{1}{a} \right) + \operatorname{acsc}^2(a + bx) \operatorname{Log} \left(\frac{1}{a} \right) + \operatorname{PolyLog} \left(2, \frac{i(-1 + \sqrt{1 - a^2}) e^{i \operatorname{csc}^{-1}(a + bx)}}{a} \right) + \operatorname{PolyLog} \left(2, \frac{i(-1 + \sqrt{1 - a^2}) e^{i \operatorname{csc}^{-1}(a + bx)}}{a} \right) \right) + \operatorname{Re} \left(\operatorname{acsc}^2(a + bx) \operatorname{Log} \left(\frac{1}{a} \right) + \operatorname{acsc}^2(a + bx) \operatorname{Log} \left(\frac{1}{a} \right) + \operatorname{PolyLog} \left(2, \frac{i(-1 + \sqrt{1 - a^2}) e^{i \operatorname{csc}^{-1}(a + bx)}}{a} \right) + \operatorname{PolyLog} \left(2, \frac{i(-1 + \sqrt{1 - a^2}) e^{i \operatorname{csc}^{-1}(a + bx)}}{a} \right) \right) + \operatorname{Re} \left(\operatorname{acsc}^2(a + bx) \operatorname{Log} \left(\frac{1}{a} \right) + \operatorname{acsc}^2(a + bx) \operatorname{Log} \left(\frac{1}{a} \right) + \operatorname{PolyLog} \left(2, \frac{i(-1 + \sqrt{1 - a^2}) e^{i \operatorname{csc}^{-1}(a + bx)}}{a} \right) + \operatorname{PolyLog} \left(2, \frac{i(-1 + \sqrt{1 - a^2}) e^{i \operatorname{csc}^{-1}(a + bx)}}{a} \right) \right)$$

Antiderivative was successfully verified.

[In] Integrate[ArcCsc[a + b*x]/x, x]

[Out] $(I*(\Pi - 2*\operatorname{ArcCsc}[a + b*x])^2 - (32*I)*\operatorname{ArcSin}[\operatorname{Sqrt}[(-1 + a)/a]/\operatorname{Sqrt}[2]]*\operatorname{ArcTan}[(1 + a)*\operatorname{Cot}[(\Pi + 2*\operatorname{ArcCsc}[a + b*x])/4]]/\operatorname{Sqrt}[1 - a^2] - 4*(\Pi - 2*\operatorname{ArcCsc}[a + b*x] + 4*\operatorname{ArcSin}[\operatorname{Sqrt}[(-1 + a)/a]/\operatorname{Sqrt}[2]])*\operatorname{Log}[1 + (I*(-1 + \operatorname{Sqrt}[1 - a^2]))/(a*E^(I*\operatorname{ArcCsc}[a + b*x]))] - 4*(\Pi - 2*\operatorname{ArcCsc}[a + b*x] - 4*\operatorname{ArcSin}[\operatorname{Sqrt}[(-1 + a)/a]/\operatorname{Sqrt}[2]])*\operatorname{Log}[1 - (I*(1 + \operatorname{Sqrt}[1 - a^2]))/(a*E^(I*\operatorname{ArcCsc}[a + b*x]))] - 8*\operatorname{ArcCsc}[a + b*x]*\operatorname{Log}[1 - E^((2*I)*\operatorname{ArcCsc}[a + b*x])] + 4*(\Pi - 2*\operatorname{ArcCsc}[a + b*x])* \operatorname{Log}[(b*x)/(a + b*x)] + 8*\operatorname{ArcCsc}[a + b*x]*\operatorname{Log}[(b*x)/(a + b*x)] + (8*I)*(\operatorname{PolyLog}[2, ((-I)*(-1 + \operatorname{Sqrt}[1 - a^2]))/(a*E^(I*\operatorname{ArcCsc}[a + b*x]))] + \operatorname{PolyLog}[2, (I*(1 + \operatorname{Sqrt}[1 - a^2]))/(a*E^(I*\operatorname{ArcCsc}[a + b*x]))] + (4*I)*(\operatorname{ArcCsc}[a + b*x]^2 + \operatorname{PolyLog}[2, E^((2*I)*\operatorname{ArcCsc}[a + b*x]))])/8$

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1143 vs. $2(269) = 538$.

time = 6.02, size = 1144, normalized size = 5.45 Too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(arccsc(b*x+a)/x,x,method=_RETURNVERBOSE)
[Out] -1/2*I*arccsc(b*x+a)/(a^2-1)^(1/2)*ln((-I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))*a+I-(a^2-1)^(1/2))/(I-(a^2-1)^(1/2))-I/(a^2-1)*dilog((-I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))*a+(a^2-1)^(1/2)+I)/(I+(a^2-1)^(1/2))*a^2+1/2*a^2*arccsc(b*x+a)/(a^2-1)*ln((-I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))*a+I-(a^2-1)^(1/2))/(I-(a^2-1)^(1/2))-1/2*arccsc(b*x+a)/(a^2-1)*ln(((I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))*a+(a^2-1)^(1/2)-I)/(I+(a^2-1)^(1/2)))+1/2*I/(a^2-1)*dilog((-I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))*a+I-(a^2-1)^(1/2))/(I-(a^2-1)^(1/2))+1/2*arccsc(b*x+a)/(a^2-1)*ln(((I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))*a+(a^2-1)^(1/2)-I)/(I+(a^2-1)^(1/2)))*a^2-1/2*I/(a^2-1)*dilog(((I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))*a+(a^2-1)^(1/2)-I)/(I+(a^2-1)^(1/2)))*a^2+I/(a^2-1)*dilog((-I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))*a+(a^2-1)^(1/2)+I)/(I+(a^2-1)^(1/2))+arccsc(b*x+a)/(a^2-1)*ln((-I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))*a+(a^2-1)^(1/2)+I)/(I+(a^2-1)^(1/2)))+1/2*I*arccsc(b*x+a)/(a^2-1)*ln((-I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))*a+(a^2-1)^(1/2)-I)/(I+(a^2-1)^(1/2))+1/2*I*arccsc(b*x+a)/(a^2-1)*ln((-I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))*a+(a^2-1)^(1/2)-I)/(I+(a^2-1)^(1/2))+1/2/(a^2-1)^(1/2)*dilog(((I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))*a+(a^2-1)^(1/2)-I)/(I+(a^2-1)^(1/2)))-1/2*I*a^2/(a^2-1)*dilog((-I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))*a+I-(a^2-1)^(1/2))/(I-(a^2-1)^(1/2))-I*dilog(I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))*a+(a^2-1)^(1/2)+I/(I+(a^2-1)^(1/2))+1/2*I/(a^2-1)*dilog(((I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))*a+(a^2-1)^(1/2)-I)/(I+(a^2-1)^(1/2)))-arccsc(b*x+a)/(a^2-1)*ln((-I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))*a+(a^2-1)^(1/2)+I)/(I+(a^2-1)^(1/2))-ln(1+I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))*arccsc(b*x+a)-1/2*arccsc(b*x+a)/(a^2-1)*ln((-I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))*a+I-(a^2-1)^(1/2))/(I-(a^2-1)^(1/2))-1/2/(a^2-1)^(1/2)*dilog((-I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))*a+I-(a^2-1)^(1/2))/(I-(a^2-1)^(1/2)))
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(arccsc(b*x+a)/x,x, algorithm="maxima")
[Out] integrate(arccsc(b*x + a)/x, x)
```

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(arccsc(b*x+a)/x,x, algorithm="fricas")
```

[Out] integral(arccsc(b*x + a)/x, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{acsc}(a + bx)}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(acsc(b*x+a)/x,x)

[Out] Integral(acsc(a + b*x)/x, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(arccsc(b*x+a)/x,x, algorithm="giac")

[Out] integrate(arccsc(b*x + a)/x, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\operatorname{asin}\left(\frac{1}{a+bx}\right)}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(asin(1/(a + b*x))/x,x)

[Out] int(asin(1/(a + b*x))/x, x)

3.23 $\int \frac{\csc^{-1}(a+bx)}{x^2} dx$

Optimal. Leaf size=69

$$-\frac{b \csc^{-1}(a + bx)}{a} - \frac{\csc^{-1}(a + bx)}{x} - \frac{2 b \text{ArcTan}\left(\frac{a - \tan\left(\frac{1}{2} \csc^{-1}(a + bx)\right)}{\sqrt{1 - a^2}}\right)}{a \sqrt{1 - a^2}}$$

[Out] $-\frac{b \text{arccsc}(b x + a)}{a} - \frac{\text{arccsc}(b x + a)}{x} - \frac{2 b \text{arctan}\left(\frac{a - \tan\left(\frac{1}{2} \text{csc}^{-1}(a + b x)\right)}{\sqrt{1 - a^2}}\right)}{a \sqrt{1 - a^2}}$

Rubi [A]

time = 0.07, antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {5367, 4512, 3868, 2739, 632, 210}

$$-\frac{2 b \text{ArcTan}\left(\frac{a - \tan\left(\frac{1}{2} \csc^{-1}(a + bx)\right)}{\sqrt{1 - a^2}}\right)}{a \sqrt{1 - a^2}} - \frac{b \csc^{-1}(a + bx)}{a} - \frac{\csc^{-1}(a + bx)}{x}$$

Antiderivative was successfully verified.

[In] Int[ArcCsc[a + b*x]/x^2, x]

[Out] $-\frac{((b \text{ArcCsc}[a + b x])/a) - \text{ArcCsc}[a + b x]/x - (2 b \text{ArcTan}[(a - \text{Tan}[\text{ArcCsc}[a + b x]/2])/\text{Sqrt}[1 - a^2]])/(a \text{Sqrt}[1 - a^2])}{a}$

Rule 210

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] :> Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] & & (LtQ[a, 0] || LtQ[b, 0])

Rule 632

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] :> Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 2739

Int[((a_) + (b_)*sin[(c_*) + (d_)*(x_)])^(-1), x_Symbol] :> With[{e = FreFactors[Tan[(c + d*x)/2], x]}, Dist[2*(e/d), Subst[Int[1/(a + 2*b*e*x + a*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]

Rule 3868

```
Int[(csc[(c_.) + (d_ .)*(x_ )]*(b_ .) + (a_ ))^(-1), x_Symbol] :> Simp[x/a, x]
- Dist[1/a, Int[1/(1 + (a/b)*Sin[c + d*x]), x], x] /; FreeQ[{a, b, c, d}, x]
] && NeQ[a^2 - b^2, 0]
```

Rule 4512

```
Int[Cot[(c_.) + (d_ .)*(x_ )]*Csc[(c_.) + (d_ .)*(x_ )]*(Csc[(c_.) + (d_ .)*(x_ )
]*(b_ .) + (a_ ))^(n_ .)*(e_ . + (f_ .)*(x_ ))^(m_ .), x_Symbol] :> Simp[(-(e +
f*x)^m)*(a + b*Csc[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[f*(m/(b*d*(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Csc[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]
```

Rule 5367

```
Int[((a_.) + ArcCsc[(c_.) + (d_ .)*(x_ )]*(b_ .))^p*((e_.) + (f_ .)*(x_ ))^(m_ .), x_Symbol] :> Dist[-(d^(m + 1))(-1), Subst[Int[(a + b*x)^p*Csc[x]*Cot[x]*(d*e - c*f + f*Csc[x])^m, x], x, ArcCsc[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Rubi steps

$$\begin{aligned}
\int \frac{\csc^{-1}(a+bx)}{x^2} dx &= - \left(b \text{Subst} \left(\int \frac{x \cot(x) \csc(x)}{(-a + \csc(x))^2} dx, x, \csc^{-1}(a+bx) \right) \right) \\
&= - \frac{\csc^{-1}(a+bx)}{x} + b \text{Subst} \left(\int \frac{1}{-a + \csc(x)} dx, x, \csc^{-1}(a+bx) \right) \\
&= - \frac{b \csc^{-1}(a+bx)}{a} - \frac{\csc^{-1}(a+bx)}{x} + \frac{b \text{Subst} \left(\int \frac{1}{1-a \sin(x)} dx, x, \csc^{-1}(a+bx) \right)}{a} \\
&= - \frac{b \csc^{-1}(a+bx)}{a} - \frac{\csc^{-1}(a+bx)}{x} + \frac{(2b) \text{Subst} \left(\int \frac{1}{1-2ax+x^2} dx, x, \tan(\frac{1}{2} \csc^{-1}(a+bx)) \right)}{a} \\
&= - \frac{b \csc^{-1}(a+bx)}{a} - \frac{\csc^{-1}(a+bx)}{x} - \frac{(4b) \text{Subst} \left(\int \frac{1}{-4(1-a^2)-x^2} dx, x, -2a + 2 \tan(\frac{1}{2} \csc^{-1}(a+bx)) \right)}{a} \\
&= - \frac{b \csc^{-1}(a+bx)}{a} - \frac{\csc^{-1}(a+bx)}{x} - \frac{2b \tan^{-1} \left(\frac{a - \tan(\frac{1}{2} \csc^{-1}(a+bx))}{\sqrt{1-a^2}} \right)}{a \sqrt{1-a^2}}
\end{aligned}$$

Mathematica [C] Result contains complex when optimal does not.

time = 0.27, size = 115, normalized size = 1.67

$$\begin{aligned} & b \left(-\text{ArcSin}\left(\frac{1}{a+bx}\right) + \frac{i \log \left(\frac{2 \left(-\frac{ia(-1+a^2+abx)}{\sqrt{1-a^2}} - a(a+bx) \right) \sqrt{\frac{-1+a^2+2abx+b^2x^2}{(a+bx)^2}}}{bx} \right)}{\sqrt{1-a^2}} \right) \\ & - \frac{\csc^{-1}(a+bx)}{x} + \frac{a}{a} \end{aligned}$$

Antiderivative was successfully verified.

[In] Integrate[ArcCsc[a + b*x]/x^2,x]

[Out] $-(\text{ArcCsc}[a + b*x]/x) + (b*(-\text{ArcSin}[(a + b*x)^{-1}] + (I*\text{Log}[(2*((-I)*a*(-1 + a^2 + a*b*x))/\text{Sqrt}[1 - a^2] - a*(a + b*x)*\text{Sqrt}[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2]))/\text{Sqrt}[1 - a^2]))/a$

Maple [A]

time = 0.63, size = 127, normalized size = 1.84

method	result
derivative divides	$b \left(-\frac{\text{arccsc}(bx+a)}{bx} - \frac{\sqrt{(bx+a)^2 - 1} \left(\arctan \left(\frac{1}{\sqrt{(bx+a)^2 - 1}} \right) \sqrt{a^2 - 1} - \ln \left(\frac{2\sqrt{a^2 - 1} \sqrt{(bx+a)^2 - 1}}{(bx+a)a\sqrt{a^2 - 1}} \right) \right)}{\sqrt{\frac{(bx+a)^2 - 1}{(bx+a)^2}} (bx+a)a\sqrt{a^2 - 1}} \right)$
default	$b \left(-\frac{\text{arccsc}(bx+a)}{bx} - \frac{\sqrt{(bx+a)^2 - 1} \left(\arctan \left(\frac{1}{\sqrt{(bx+a)^2 - 1}} \right) \sqrt{a^2 - 1} - \ln \left(\frac{2\sqrt{a^2 - 1} \sqrt{(bx+a)^2 - 1}}{(bx+a)a\sqrt{a^2 - 1}} \right) \right)}{\sqrt{\frac{(bx+a)^2 - 1}{(bx+a)^2}} (bx+a)a\sqrt{a^2 - 1}} \right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(arccsc(b*x+a)/x^2,x,method=_RETURNVERBOSE)

[Out] $b*(-1/b/x*\text{arccsc}(b*x+a) - ((b*x+a)^2 - 1)^{1/2} * (\arctan(1/((b*x+a)^2 - 1)^{1/2}) * (a^2 - 1)^{1/2} - \ln(2*((a^2 - 1)^{1/2} * ((b*x+a)^2 - 1)^{1/2} + a*(b*x+a) - 1)/b/x)) / ((b*x+a)^2 - 1) / (b*x+a)^2)^{1/2} / (b*x+a) / a / (a^2 - 1)^{1/2})$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(b*x+a)/x^2,x, algorithm="maxima")`
[Out]
$$\frac{-(x \cdot \text{integrate}((b^2 \cdot x + a \cdot b) \cdot e^{(1/2 \cdot \log(b \cdot x + a + 1) + 1/2 \cdot \log(b \cdot x + a - 1))}, x) + \arctan(1, \sqrt{b \cdot x + a + 1}) \cdot \sqrt{b \cdot x + a - 1}))}{(b^2 \cdot x^3 + 2 \cdot a \cdot b \cdot x^2 + (a^2 - 1) \cdot x + (b^2 \cdot x^3 + 2 \cdot a \cdot b \cdot x^2 + (a^2 - 1) \cdot x) \cdot e^{(\log(b \cdot x + a + 1) + \log(b \cdot x + a - 1))}})$$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 132 vs. 2(63) = 126.

time = 0.40, size = 278, normalized size = 4.03

$$\left[\frac{2(a^2-1)bx \arctan\left(-bx-a+\sqrt{b^2x^2+2abx+a^2-1}\right)+\sqrt{a^2-1}bx \log\left(\frac{(a^2+bx^2+\sqrt{b^2x^2+2abx+a^2-1})(a^2+bx^2-\sqrt{a^2-1}-a)}{x}\right)-(a^2-a)\arccos(bx+a)}{(a^2-a)x}, \frac{2(a^2-1)bx \arctan\left(-bx-a+\sqrt{b^2x^2+2abx+a^2-1}\right)-2\sqrt{-a^2+1}bx \arctan\left(\frac{-\sqrt{-a^2+1}bx-\sqrt{b^2x^2+2abx+a^2-1}\sqrt{-a^2+1}}{x}\right)-(a^2-a)\arccos(bx+a)}{(a^2-a)x} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(b*x+a)/x^2,x, algorithm="fricas")`
[Out]
$$[(2*(a^2-1)*b*x*\arctan(-b*x-a+\sqrt{b^2*x^2+2*a*b*x+a^2-1}))+\sqrt{a^2-1}*b*x*\log((a^2*b*x+a^3+\sqrt{b^2*x^2+2*a*b*x+a^2-1})*(a^2-1)*(a^2+sqrt(a^2-1)*a-1)+(a*b*x+a^2-1)*sqrt(a^2-1))/x-(a^3-a)*\arccsc(b*x+a))/((a^3-a)*x), (2*(a^2-1)*b*x*\arctan(-b*x-a+\sqrt{b^2*x^2+2*a*b*x+a^2-1}))-2*sqrt(-a^2+1)*b*x*\arctan(-(sqrt(-a^2+1)*b*x-sqrt(b^2*x^2+2*a*b*x+a^2-1)*sqrt(-a^2+1))/(a^2-1))-(a^3-a)*\arccsc(b*x+a))/((a^3-a)*x)]$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{acsc}(a+bx)}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(acsc(b*x+a)/x**2,x)`
[Out] `Integral(acsc(a+b*x)/x**2, x)`

Giac [A]

time = 0.47, size = 96, normalized size = 1.39

$$-b \left(\frac{2 \arctan \left(\frac{(bx+a) \left(\sqrt{-\frac{1}{(bx+a)^2} + 1} - 1 \right) + a}{\sqrt{-a^2+1}} \right)}{\sqrt{-a^2+1} a} - \frac{\arcsin \left(-\frac{1}{(bx+a) \left(\frac{a}{bx+a} - 1 \right) - a} \right)}{a \left(\frac{a}{bx+a} - 1 \right)} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(b*x+a)/x^2,x, algorithm="giac")`

[Out]
$$\frac{-b*(2*\arctan((b*x + a)*(\sqrt{-1/(b*x + a)^2 + 1} - 1) + a)/\sqrt{-a^2 + 1})}{(\sqrt{-a^2 + 1}*a) - \arcsin(-1/((b*x + a)*(a/(b*x + a) - 1) - a))/(a*(a/(b*x + a) - 1)))}$$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sin \left(\frac{1}{a+b x} \right)}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(asin(1/(a + b*x))/x^2,x)`

[Out] `int(asin(1/(a + b*x))/x^2, x)`

3.24 $\int \frac{\csc^{-1}(a+bx)}{x^3} dx$

Optimal. Leaf size=123

$$\frac{b(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{2a(1-a^2)x} + \frac{b^2\csc^{-1}(a+bx)}{2a^2} - \frac{\csc^{-1}(a+bx)}{2x^2} + \frac{(1-2a^2)b^2\text{ArcTan}\left(\frac{a-\tan(\frac{1}{2}\csc^{-1}(a+bx))}{\sqrt{1-a^2}}\right)}{a^2(1-a^2)^{3/2}}$$

[Out] $1/2*b^2*\text{arccsc}(b*x+a)/a^2 - 1/2*\text{arccsc}(b*x+a)/x^2 + (-2*a^2+1)*b^2*\text{arctan}((a-\tan(1/2*\text{arccsc}(b*x+a)))/(-a^2+1)^(1/2))/a^2/(-a^2+1)^(3/2) - 1/2*b*(b*x+a)*(1-1/(b*x+a)^2)^(1/2)/a/(-a^2+1)/x$

Rubi [A]

time = 0.15, antiderivative size = 123, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.800$, Rules used = {5367, 4512, 3870, 4004, 3916, 2739, 632, 210}

$$\frac{(1-2a^2)b^2\text{ArcTan}\left(\frac{a-\tan(\frac{1}{2}\csc^{-1}(a+bx))}{\sqrt{1-a^2}}\right)}{a^2(1-a^2)^{3/2}} + \frac{b^2\csc^{-1}(a+bx)}{2a^2} - \frac{b(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{2a(1-a^2)x} - \frac{\csc^{-1}(a+bx)}{2x^2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{ArcCsc}[a+b*x]/x^3, x]$

[Out] $-1/2*(b*(a+b*x)*\text{Sqrt}[1-(a+b*x)^{-2}])/(a*(1-a^2)*x) + (b^2*\text{ArcCsc}[a+b*x])/(2*a^2) - \text{ArcCsc}[a+b*x]/(2*x^2) + ((1-2*a^2)*b^2*\text{ArcTan}[(a-\tan[\text{ArcCsc}[a+b*x]/2])/Sqrt[1-a^2]])/(a^2*(1-a^2)^{3/2})$

Rule 210

$\text{Int}[(a_.) + (b_.)*(x_.)^2^{-1}, x_{\text{Symbol}}] \Rightarrow \text{Simp}[-(\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{(-1)}*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \& \text{PosQ}[a/b] \& (\text{LtQ}[a, 0] \text{ || } \text{LtQ}[b, 0])$

Rule 632

$\text{Int}[(a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2^{-1}, x_{\text{Symbol}}] \Rightarrow \text{Dist}[-2, \text{Subst}[\text{Int}[1/\text{Simp}[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; \text{FreeQ}[\{a, b, c\}, x] \& \text{NeQ}[b^2 - 4*a*c, 0]$

Rule 2739

$\text{Int}[(a_.) + (b_.)*\sin[(c_.) + (d_.)*(x_.)]^{-1}, x_{\text{Symbol}}] \Rightarrow \text{With}[\{e = \text{FreeFactors}[\text{Tan}[(c+d*x)/2], x]\}, \text{Dist}[2*(e/d), \text{Subst}[\text{Int}[1/(a + 2*b*e*x + a*x^2), x], x, \text{Tan}[(c+d*x)/2]/e], x]] /; \text{FreeQ}[\{a, b, c, d\}, x] \& \text{NeQ}[$

$a^2 - b^2, 0]$

Rule 3870

```
Int[(csc[(c_.) + (d_ .)*(x_)]*(b_ .) + (a_ ))^(n_), x_Symbol] :> Simp[b^2*Cot[c + d*x]*((a + b*Csc[c + d*x])^(n + 1)/(a*d*(n + 1)*(a^2 - b^2))), x] + Dist[1/(a*(n + 1)*(a^2 - b^2)), Int[(a + b*Csc[c + d*x])^(n + 1)*Simp[(a^2 - b^2)*(n + 1) - a*b*(n + 1)*Csc[c + d*x] + b^2*(n + 2)*Csc[c + d*x]^2, x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]
```

Rule 3916

```
Int[csc[(e_.) + (f_ .)*(x_)]/(csc[(e_.) + (f_ .)*(x_)]*(b_ .) + (a_ )), x_Symbol] :> Dist[1/b, Int[1/(1 + (a/b)*Sin[e + f*x]), x], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a^2 - b^2, 0]
```

Rule 4004

```
Int[(csc[(e_.) + (f_ .)*(x_)]*(d_ .) + (c_ ))/(csc[(e_.) + (f_ .)*(x_)]*(b_ .) + (a_ )), x_Symbol] :> Simp[c*(x/a), x] - Dist[(b*c - a*d)/a, Int[Csc[e + f*x]/(a + b*Csc[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]
```

Rule 4512

```
Int[Cot[(c_.) + (d_ .)*(x_)]*Csc[(c_.) + (d_ .)*(x_)]*(Csc[(c_.) + (d_ .)*(x_)]*(b_ .) + (a_ ))^(n_ .)*((e_.) + (f_ .)*(x_))^(m_.), x_Symbol] :> Simp[((-e + f*x)^m)*((a + b*Csc[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[f*(m/(b*d*(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Csc[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]
```

Rule 5367

```
Int[((a_.) + ArcCsc[(c_) + (d_ .)*(x_)]*(b_ .))^(p_ .)*((e_.) + (f_ .)*(x_))^(m_.), x_Symbol] :> Dist[-(d^(m + 1))^(-1), Subst[Int[(a + b*x)^p*Csc[x]*Cot[x]*(d*e - c*f + f*Csc[x])^m, x], x, ArcCsc[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Rubi steps

$$\begin{aligned}
\int \frac{\csc^{-1}(a+bx)}{x^3} dx &= - \left(b^2 \text{Subst} \left(\int \frac{x \cot(x) \csc(x)}{(-a + \csc(x))^3} dx, x, \csc^{-1}(a+bx) \right) \right) \\
&= - \frac{\csc^{-1}(a+bx)}{2x^2} + \frac{1}{2} b^2 \text{Subst} \left(\int \frac{1}{(-a + \csc(x))^2} dx, x, \csc^{-1}(a+bx) \right) \\
&= - \frac{b(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{2a(1-a^2)x} - \frac{\csc^{-1}(a+bx)}{2x^2} - \frac{b^2 \text{Subst} \left(\int \frac{1-a^2-a \csc(x)}{-a+\csc(x)} dx, x, \csc^{-1}(a+bx) \right)}{2a(1-a^2)} \\
&= - \frac{b(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{2a(1-a^2)x} + \frac{b^2 \csc^{-1}(a+bx)}{2a^2} - \frac{\csc^{-1}(a+bx)}{2x^2} - \frac{((1-2a^2)b^2) \text{Subst} \left(\int \frac{1-a^2-a \csc(x)}{-a+\csc(x)} dx, x, \csc^{-1}(a+bx) \right)}{2a(1-a^2)} \\
&= - \frac{b(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{2a(1-a^2)x} + \frac{b^2 \csc^{-1}(a+bx)}{2a^2} - \frac{\csc^{-1}(a+bx)}{2x^2} - \frac{((1-2a^2)b^2) \text{Subst} \left(\int \frac{1-a^2-a \csc(x)}{-a+\csc(x)} dx, x, \csc^{-1}(a+bx) \right)}{2a(1-a^2)} \\
&= - \frac{b(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{2a(1-a^2)x} + \frac{b^2 \csc^{-1}(a+bx)}{2a^2} - \frac{\csc^{-1}(a+bx)}{2x^2} - \frac{((1-2a^2)b^2) \text{Subst} \left(\int \frac{1-a^2-a \csc(x)}{-a+\csc(x)} dx, x, \csc^{-1}(a+bx) \right)}{2a(1-a^2)} \\
&= - \frac{b(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{2a(1-a^2)x} + \frac{b^2 \csc^{-1}(a+bx)}{2a^2} - \frac{\csc^{-1}(a+bx)}{2x^2} + \frac{(2(1-2a^2)b^2) \text{Subst} \left(\int \frac{1-a^2-a \csc(x)}{-a+\csc(x)} dx, x, \csc^{-1}(a+bx) \right)}{2a(1-a^2)} \\
&= - \frac{b(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{2a(1-a^2)x} + \frac{b^2 \csc^{-1}(a+bx)}{2a^2} - \frac{\csc^{-1}(a+bx)}{2x^2} + \frac{(1-2a^2)b^2 \tan^{-1} \frac{a+bx}{\sqrt{1-a^2}}}{a^2(1-a^2)^{3/2}}
\end{aligned}$$

Mathematica [C] Result contains complex when optimal does not.

time = 0.57, size = 199, normalized size = 1.62

$$\frac{bx(a+bx)\sqrt{\frac{-1+a^2+2abx+b^2x^2}{(a+bx)^2}}}{a(-1+a^2)} - \csc^{-1}(a+bx) + \frac{\frac{b^2x^2 \text{ArcSin}(\frac{1}{a+bx})}{a^2} + \frac{i(-1+2a^2)b^2x^2 \log \left(\frac{\sqrt{\frac{-1+a^2+2abx+b^2x^2}{(a+bx)^2}}}{\sqrt{1-a^2}} \right)}{(-1+2a^2)b^2x}}{2x^2}$$

Antiderivative was successfully verified.

[In] Integrate[ArcCsc[a + b*x]/x^3, x]

[Out] $((b*x*(a + b*x)*\text{Sqrt}[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2])/(a*(-1 + a^2)) - \text{ArcCsc}[a + b*x] + (b^2*x^2*\text{ArcSin}[(a + b*x)^{-1}])/a^2 + (I*(-1 + 2$

$$*a^2)*b^2*x^2*Log[(4*(-1 + a)*a^2*(1 + a)*((I*(-1 + a^2 + a*b*x))/Sqrt[1 - a^2] + (a + b*x)*Sqrt[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2]))/((-1 + 2*a^2)*b^2*x)])/(a^2*(1 - a^2)^(3/2)))/(2*x^2)$$

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 455 vs. $2(109) = 218$.

time = 0.64, size = 456, normalized size = 3.71

method	result
derivativedivides	$b^2 \left(-\frac{\operatorname{arccsc}(bx+a)}{2b^2x^2} + \frac{\sqrt{(bx+a)^2 - 1} \left(-\arctan\left(\frac{1}{\sqrt{(bx+a)^2 - 1}}\right) (a^2-1)^{\frac{3}{2}} a^3 + \arctan\left(\frac{1}{\sqrt{(bx+a)^2 - 1}}\right) (a^2-1)^{\frac{3}{2}} a^3\right)}{(bx+a)^2}\right)$
default	$b^2 \left(-\frac{\operatorname{arccsc}(bx+a)}{2b^2x^2} + \frac{\sqrt{(bx+a)^2 - 1} \left(-\arctan\left(\frac{1}{\sqrt{(bx+a)^2 - 1}}\right) (a^2-1)^{\frac{3}{2}} a^3 + \arctan\left(\frac{1}{\sqrt{(bx+a)^2 - 1}}\right) (a^2-1)^{\frac{3}{2}} a^3\right)}{(bx+a)^2}\right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(arccsc(b*x+a)/x^3,x,method=_RETURNVERBOSE)`

[Out] $b^2*(-1/2/b^2/x^2*arccsc(b*x+a)+1/2*((b*x+a)^2-1)^(1/2)*(-arctan(1/((b*x+a)^2-1)^(1/2))*(a^2-1)^(3/2)*a^2*(b*x+a)+2*ln(2*((a^2-1)^(1/2)*((b*x+a)^2-1)^(1/2)+a*(b*x+a)-1)/b/x)*a^5-2*ln(2*((a^2-1)^(1/2)*((b*x+a)^2-1)^(1/2)+a*(b*x+a)-1)/b/x)*a^4*(b*x+a)+((b*x+a)^2-1)^(1/2)*(a^2-1)^(3/2)*a+arctan(1/((b*x+a)^2-1)^(1/2))*(a^2-1)^(3/2)*a-arctan(1/((b*x+a)^2-1)^(1/2))*(a^2-1)^(3/2)*(b*x+a)-3*ln(2*((a^2-1)^(1/2)*((b*x+a)^2-1)^(1/2)+a*(b*x+a)-1)/b/x)*a^3+3*ln(2*((a^2-1)^(1/2)*((b*x+a)^2-1)^(1/2)+a*(b*x+a)-1)/b/x)*a^2+ln(2*((a^2-1)^(1/2)*((b*x+a)^2-1)^(1/2)+a*(b*x+a)-1)/b/x)-ln(2*((a^2-1)^(1/2)*((b*x+a)^2-1)^(1/2)+a*(b*x+a)-1)/b/x)*(b*x+a))/(((b*x+a)^2-1)/(b*x+a)^(1/2)/(b*x+a)/a^2/(a^2-1)^(5/2)/b/x)$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(b*x+a)/x^3,x, algorithm="maxima")`

[Out] $-1/2*(2*x^2*integrate(1/2*(b^2*x + a*b)*e^(1/2*log(b*x + a + 1) + 1/2*log(b*x + a - 1))/(b^2*x^4 + 2*a*b*x^3 + (a^2 - 1)*x^2 + (b^2*x^4 + 2*a*b*x^3 + (a^2 - 1)*x^2)*e^(log(b*x + a + 1) + log(b*x + a - 1))), x) + arctan2(1, sqrt(b*x + a + 1)*sqrt(b*x + a - 1)))/x^2$

Fricas [A]

time = 0.42, size = 428, normalized size = 3.48

$$\frac{(2(x^2 - 1)\sqrt{a^2 - 1})^{1/2}x^2 \operatorname{sgn}\left(\frac{\sinh(x\sqrt{b^2 x^2 + 2 a b x + a^2 - 1}) (e^{i x \sqrt{b^2 x^2 + 2 a b x + a^2 - 1}})}{2(x^2 - 2 a^2 + 1)^{1/2} x^2 \operatorname{arctan}\left(-4 x - a + \sqrt{b^2 x^2 + 2 a b x + a^2 - 1}\right)} + (a^2 - a)x^2 + \sqrt{b^2 x^2 + 2 a b x + a^2 - 1}) (x^2 - a^2 x - (a^2 - 2 a^2 + a^2)x \operatorname{arccos}(bx + a) + 2(2(x^2 - 1))^{1/2} x^2 \operatorname{arctan}\left(-\frac{\sqrt{b^2 x^2 + 2 a b x + a^2 - 1} \operatorname{sgn}(bx + a)}{2(x^2 - 2 a^2 + a^2)x^2}\right) - 2(x^2 - 2 a^2 + 1)^{1/2} x^2 \operatorname{arctan}\left(-\frac{\sqrt{b^2 x^2 + 2 a b x + a^2 - 1} \operatorname{sgn}(bx + a)}{2(x^2 - 2 a^2 + a^2)x^2}\right) + (a^2 - a)x^2 + \sqrt{b^2 x^2 + 2 a b x + a^2 - 1})^2 - a)x \operatorname{arccos}(bx + a))}{2(x^2 - 2 a^2 + a^2)x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(b*x+a)/x^3,x, algorithm="fricas")`

[Out] $[1/2*((2*a^2 - 1)*sqrt(a^2 - 1)*b^2*x^2*log((a^2*b*x + a^3 + sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)*(a^2 - sqrt(a^2 - 1)*a - 1) - (a*b*x + a^2 - 1)*sqrt(a^2 - 1) - a)/x) - 2*(a^4 - 2*a^2 + 1)*b^2*x^2*arctan(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)) + (a^3 - a)*b^2*x^2 + sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)*(a^3 - a)*b*x - (a^6 - 2*a^4 + a^2)*arccsc(b*x + a))/((a^6 - 2*a^4 + a^2)*x^2), 1/2*(2*(2*a^2 - 1)*sqrt(-a^2 + 1)*b^2*x^2*arctan(-(sqrt(-a^2 + 1)*b*x - sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)*sqrt(-a^2 + 1))/(a^2 - 1)) - 2*(a^4 - 2*a^2 + 1)*b^2*x^2*arctan(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)) + (a^3 - a)*b^2*x^2 + sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)*(a^3 - a)*b*x - (a^6 - 2*a^4 + a^2)*arccsc(b*x + a))/((a^6 - 2*a^4 + a^2)*x^2)]$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{acsc}(a + bx)}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(acsc(b*x+a)/x**3,x)`

[Out] `Integral(acsc(a + b*x)/x**3, x)`

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 217 vs. 2(108) = 216.

time = 0.48, size = 217, normalized size = 1.76

$$\frac{1}{2} b \left(\frac{2 (2 a^2 b - b) \operatorname{arctan}\left(\frac{\sqrt{\frac{1}{(bx+a)^2} + 1} - 1}{\sqrt{-a^2 + 1}}\right) + a}{(a^4 - a^2) \sqrt{-a^2 + 1}} + \frac{2 \left((bx + a) ab \left(\sqrt{\frac{1}{(bx+a)^2} + 1} - 1\right) + b\right)}{\left((bx + a)^2 \left(\sqrt{\frac{1}{(bx+a)^2} + 1} - 1\right)^2 + 2 (bx + a) a \left(\sqrt{\frac{1}{(bx+a)^2} + 1} - 1\right) + 1\right) (a^3 - a)} - \frac{(\frac{2 a b}{bx+a} - b) \arcsin\left(-\frac{1}{\frac{a^2}{bx+a} - 1}\right)}{a^2 \left(\frac{a}{bx+a} - 1\right)^2} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(b*x+a)/x^3,x, algorithm="giac")`

[Out] $\frac{1}{2} b \left(2 \left(2 a^2 b - b\right) \arctan \left(\left(b x + a\right) \sqrt{-\frac{1}{(b x + a)^2} + 1}\right) - a\right) \sqrt{-a^2 + 1} / \left(a^4 - a^2\right) \sqrt{-a^2 + 1} + 2 \left(\left(b x + a\right) a b \sqrt{-\frac{1}{(b x + a)^2} + 1} - 1\right) \sqrt{-a^2 + 1} + 2 \left(\left(b x + a\right)^2 \left(\sqrt{-\frac{1}{(b x + a)^2} + 1} - 1\right) + 1\right) \left(a^3 - a\right) - \left(2 a b \left(b x + a\right) - b\right) \arcsin \left(-\frac{1}{\left(b x + a\right)} \left(a / \left(b x + a\right) - 1\right) - a\right) / \left(a^2 \left(a / \left(b x + a\right) - 1\right)^2\right)$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\arcsin\left(\frac{1}{a+b x}\right)}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(asin(1/(a + b*x))/x^3,x)`

[Out] `int(asin(1/(a + b*x))/x^3, x)`

3.25 $\int \frac{\csc^{-1}(a+bx)}{x^4} dx$

Optimal. Leaf size=180

$$\frac{b(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{6a(1-a^2)x^2} + \frac{(2-5a^2)b^2(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{6a^2(1-a^2)^2x} - \frac{b^3\csc^{-1}(a+bx)}{3a^3} - \frac{\csc^{-1}(a+bx)}{3x^3} -$$

[Out] $-1/3*b^3*\arccsc(b*x+a)/a^3-1/3*\arccsc(b*x+a)/x^3-1/3*(6*a^4-5*a^2+2)*b^3*\arctan((a-\tan(1/2*\arccsc(b*x+a)))/(-a^2+1)^(1/2))/a^3/(-a^2+1)^(5/2)-1/6*b*(b*x+a)*(1-1/(b*x+a)^2)^(1/2)/a/(-a^2+1)/x^2+1/6*(-5*a^2+2)*b^2*(b*x+a)*(1-1/(b*x+a)^2)^(1/2)/a^2/(-a^2+1)^2/x$

Rubi [A]

time = 0.22, antiderivative size = 180, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 9, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.900, Rules used = {5367, 4512, 3870, 4145, 4004, 3916, 2739, 632, 210}

$$-\frac{b^3\csc^{-1}(a+bx)}{3a^3} + \frac{(2-5a^2)b^2(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{6a^2(1-a^2)^2x} - \frac{b(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{6a(1-a^2)x^2} - \frac{(6a^4-5a^2+2)b^3\text{ArcTan}\left(\frac{a-\tan(\frac{1}{2}\csc^{-1}(a+bx))}{\sqrt{1-a^2}}\right)}{3a^3(1-a^2)^{5/2}} - \frac{\csc^{-1}(a+bx)}{3x^3}$$

Antiderivative was successfully verified.

[In] Int[ArcCsc[a + b*x]/x^4, x]

[Out] $-1/6*(b*(a+b*x)*Sqrt[1-(a+b*x)^{-2}]}/{(a*(1-a^2)*x^2)} + ((2-5*a^2)*b^2*(a+b*x)*Sqrt[1-(a+b*x)^{-2}])/(6*a^2*(1-a^2)^2*x) - (b^3*ArcCsc[a+b*x])/(3*a^3) - ArcCsc[a+b*x]/(3*x^3) - ((2-5*a^2+6*a^4)*b^3*ArcTan[(a-Tan[ArcCsc[a+b*x]/2])/Sqrt[1-a^2]])/(3*a^3*(1-a^2)^(5/2))$

Rule 210

Int[((a_) + (b_)*(x_)^2)^{-1}, x_Symbol] :> Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 632

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^{-1}, x_Symbol] :> Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 2739

Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^{-1}, x_Symbol] :> With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Dist[2*(e/d), Subst[Int[1/(a + 2*b*e*x + a*

```
e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]
```

Rule 3870

```
Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.) + (a_.))^n_, x_Symbol] :> Simp[b^2*Cot[c + d*x]*((a + b*Csc[c + d*x])^(n + 1)/(a*d*(n + 1)*(a^2 - b^2))), x] + Dist[1/(a*(n + 1)*(a^2 - b^2)), Int[(a + b*Csc[c + d*x])^(n + 1)*Simp[(a^2 - b^2)*(n + 1) - a*b*(n + 1)*Csc[c + d*x] + b^2*(n + 2)*Csc[c + d*x]^2, x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]
```

Rule 3916

```
Int[csc[(e_.) + (f_.)*(x_.)]/(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)), x_Symbol] :> Dist[1/b, Int[1/(1 + (a/b)*Sin[e + f*x]), x], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a^2 - b^2, 0]
```

Rule 4004

```
Int[(csc[(e_.) + (f_.)*(x_.)]*(d_.) + (c_.))/(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.)), x_Symbol] :> Simp[c*(x/a), x] - Dist[(b*c - a*d)/a, Int[Csc[e + f*x]/(a + b*Csc[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]
```

Rule 4145

```
Int[((A_.) + csc[(e_.) + (f_.)*(x_.)]*(B_.) + csc[(e_.) + (f_.)*(x_.)]^2*(C_.))*csc[(e_.) + (f_.)*(x_.)]*(b_.)^(m_), x_Symbol] :> Simp[(A*b^2 - a*b*B + a^2*C)*Cot[e + f*x]*((a + b*Csc[e + f*x])^(m + 1)/(a*f*(m + 1)*(a^2 - b^2))), x] + Dist[1/(a*(m + 1)*(a^2 - b^2)), Int[(a + b*Csc[e + f*x])^(m + 1)*Simp[A*(a^2 - b^2)*(m + 1) - a*(A*b - a*B + b*C)*(m + 1)*Csc[e + f*x] + (A*b^2 - a*b*B + a^2*C)*(m + 2)*Csc[e + f*x]^2, x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && NeQ[a^2 - b^2, 0] && LtQ[m, -1]
```

Rule 4512

```
Int[Cot[(c_.) + (d_.)*(x_.)]*Csc[(c_.) + (d_.)*(x_.)]*(Csc[(c_.) + (d_.)*(x_.)]*(b_.)^(n_)*(e_.)^(m_), x_Symbol] :> Simp[(-(e + f*x)^m)*((a + b*Csc[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[f*(m/(b*d*(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Csc[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]
```

Rule 5367

```
Int[((a_.) + ArcCsc[(c_.) + (d_.)*(x_.)]*(b_.))^(p_)*(e_.)^(m_ - ), x_Symbol] :> Dist[-(d^(m + 1))^{(-1)}, Subst[Int[(a + b*x)^p*Csc[x]*Cot[
```

```
x]*(d*e - c*f + f*Csc[x])^m, x], x, ArcCsc[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Rubi steps

$$\begin{aligned}
\int \frac{\csc^{-1}(a+bx)}{x^4} dx &= - \left(b^3 \text{Subst} \left(\int \frac{x \cot(x) \csc(x)}{(-a+\csc(x))^4} dx, x, \csc^{-1}(a+bx) \right) \right) \\
&= - \frac{\csc^{-1}(a+bx)}{3x^3} + \frac{1}{3} b^3 \text{Subst} \left(\int \frac{1}{(-a+\csc(x))^3} dx, x, \csc^{-1}(a+bx) \right) \\
&= - \frac{b(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{6a(1-a^2)x^2} - \frac{\csc^{-1}(a+bx)}{3x^3} - \frac{b^3 \text{Subst} \left(\int \frac{2(1-a^2)-2a \csc(x)-\csc^2(x)}{(-a+\csc(x))^2} dx, x, \csc^{-1}(a+bx) \right)}{6a(1-a^2)} \\
&= - \frac{b(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{6a(1-a^2)x^2} + \frac{(2-5a^2)b^2(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{6a^2(1-a^2)^2 x} - \frac{\csc^{-1}(a+bx)}{3x^3} \\
&= - \frac{b(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{6a(1-a^2)x^2} + \frac{(2-5a^2)b^2(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{6a^2(1-a^2)^2 x} - \frac{b^3 \csc^{-1}(a+bx)}{3a^3} \\
&= - \frac{b(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{6a(1-a^2)x^2} + \frac{(2-5a^2)b^2(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{6a^2(1-a^2)^2 x} - \frac{b^3 \csc^{-1}(a+bx)}{3a^3} \\
&= - \frac{b(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{6a(1-a^2)x^2} + \frac{(2-5a^2)b^2(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{6a^2(1-a^2)^2 x} - \frac{b^3 \csc^{-1}(a+bx)}{3a^3} \\
&= - \frac{b(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{6a(1-a^2)x^2} + \frac{(2-5a^2)b^2(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{6a^2(1-a^2)^2 x} - \frac{b^3 \csc^{-1}(a+bx)}{3a^3} \\
&= - \frac{b(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{6a(1-a^2)x^2} + \frac{(2-5a^2)b^2(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{6a^2(1-a^2)^2 x} - \frac{b^3 \csc^{-1}(a+bx)}{3a^3}
\end{aligned}$$

Mathematica [C] Result contains complex when optimal does not.

time = 0.34, size = 241, normalized size = 1.34

$$\frac{1}{6} \left(\frac{b \sqrt{\frac{-1 + a^2 + 2abx + b^2x^2}{(a+bx)^2}} (a^4 + abx - 4a^3bx + 2b^2x^2 - a^2(1 + 5b^2x^2))}{a^2(-1 + a^2)^2 x^2} - \frac{2 \csc^{-1}(a+bx)}{x^3} - \frac{2b^3 \text{ArcSin}\left(\frac{1}{a+bx}\right)}{a^3} + \frac{i(2 - 5a^2 + 6a^4) b^3 \log\left(\frac{12a^3(-1+a^2)^2 \left(-\frac{i(-1+a^2+abx)}{\sqrt{1-a^2}}-(a+bx)\right) \sqrt{\frac{-1+a^2+2abx+b^2x^2}{(a+bx)^2}}}{(2-5a^2+6a^4)b^3x}\right)}{a^3(1-a^2)^{5/2}} \right)$$

Antiderivative was successfully verified.

[In] Integrate[ArcCsc[a + b*x]/x^4,x]

[Out] $\frac{((b*\text{Sqrt}[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2]*(a^4 + a*b*x - 4*a^3*b*x + 2*b^2*x^2 - a^2*(1 + 5*b^2*x^2)))/(a^2*(-1 + a^2)^2*x^2) - (2*\text{ArcCsc}[a + b*x])/x^3 - (2*b^3*\text{ArcSin}[(a + b*x)^{-1}])/a^3 + (I*(2 - 5*a^2 + 6*a^4)*b^3*\text{Log}[(12*a^3*(-1 + a^2)^2*((-I)*(-1 + a^2 + a*b*x))/\text{Sqrt}[1 - a^2] - (a + b*x)*\text{Sqrt}[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2]))/((2 - 5*a^2 + 6*a^4)*b^3*x)]/(a^3*(1 - a^2)^(5/2)))/6$

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 969 vs. 2(160) = 320.

time = 0.65, size = 970, normalized size = 5.39 Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int(arccsc(b*x+a)/x^4,x,method=_RETURNVERBOSE)

[Out] $b^3*(-1/3/b^3/x^3*arccsc(b*x+a)+1/6*((b*x+a)^2-1)^(1/2)*(-8*arctan(1/((b*x+a)^2-1)^(1/2))*(a^2-1)^(3/2)*a^3*(b*x+a)+4*arctan(1/((b*x+a)^2-1)^(1/2))*(a^2-1)^(3/2)*a^2*(b*x+a)^2+4*arctan(1/((b*x+a)^2-1)^(1/2))*(a^2-1)^(3/2)*a^5*(b*x+a)-2*arctan(1/((b*x+a)^2-1)^(1/2))*(a^2-1)^(3/2)*a^4*(b*x+a)^2-5*(b*x+a)^2-1)^(1/2)*(a^2-1)^(3/2)*a^3*(b*x+a)-2*ln(2*((a^2-1)^(1/2)*((b*x+a)^2-1)^(1/2)+a*(b*x+a)-1)/b/x)*a^2+4*arctan(1/((b*x+a)^2-1)^(1/2))*(a^2-1)^(3/2)*a^5*(b*x+a)-2*ln(2*((a^2-1)^(1/2)*((b*x+a)^2-1)^(1/2)+a*(b*x+a)-1)/b/x)*a^5*(b*x+a)+11*ln(2*((a^2-1)^(1/2)*((b*x+a)^2-1)^(1/2)+a*(b*x+a)-1)/b/x)*a^4*(b*x+a)^2-3*((b*x+a)^2-1)^(1/2)*(a^2-1)^(3/2)*a^2-2*arctan(1/((b*x+a)^2-1)^(1/2))*(a^2-1)^(3/2)*(b*x+a)^2-12*ln(2*((a^2-1)^(1/2)*((b*x+a)^2-1)^(1/2)+a*(b*x+a)-1)/b/x)*a^7*(b*x+a)+6*ln(2*((a^2-1)^(1/2)*((b*x+a)^2-1)^(1/2)+a*(b*x+a)-1)/b/x)*a^6*(b*x+a)^2+6*((b*x+a)^2-1)^(1/2)*(a^2-1)^(3/2)*a^4-2*arctan(1/((b*x+a)^2-1)^(1/2))*(a^2-1)^(3/2)*a^6+4*ln(2*((a^2-1)^(1/2)*((b*x+a)^2-1)^(1/2)+a*(b*x+a)-1)/b/x)*a^7*(b*x+a)-14*ln(2*((a^2-1)^(1/2)*((b*x+a)^2-1)^(1/2)+a*(b*x+a)-1)/b/x)*a^3*(b*x+a)+7*ln(2*((a^2-1)^(1/2)*((b*x+a)^2-1)^(1/2)+a*(b*x+a)-1)/b/x)*a^2*(b*x+a)^2-2*ln(2*((a^2-1)^(1/2)*((b*x+a)^2-1)^(1/2)+a*(b*x+a)-1)/b/x)*(b*x+a)^2+6*ln(2*((a^2-1)^(1/2)*((b*x+a)^2-1)^(1/2)+a*(b*x+a)-1)/b/x)*a^8-11*ln(2*((a^2-1)^(1/2)*((b*x+a)^2-1)^(1/2)+a*(b*x+a)-1)/b/x)*a^6+7*ln(2*((a^2-1)^(1/2)*((b*x+a)^2-1)^(1/2)+a*(b*x+a)-1)/b/x)*a^4+2*((b*x+a)^2-1)^(1/2)*(a^2-1)^(3/2))$

$2*a*(b*x+a)+4*arctan(1/((b*x+a)^2-1)^(1/2))*(a^2-1)^(3/2)*a*(b*x+a))/(((b*x+a)^2-1)/(b*x+a)^2)^(1/2)/(b*x+a)/a^3/(a^2-1)^(7/2)/b^2/x^2)$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(b*x+a)/x^4,x, algorithm="maxima")`

[Out] $-1/3*(3*x^3*integrate(1/3*(b^2*x + a*b)*e^(1/2*log(b*x + a + 1) + 1/2*log(b*x + a - 1))/(b^2*x^5 + 2*a*b*x^4 + (a^2 - 1)*x^3 + (b^2*x^5 + 2*a*b*x^4 + (a^2 - 1)*x^3)*e^(log(b*x + a + 1) + log(b*x + a - 1))), x) + arctan2(1, sqrt(b*x + a + 1)*sqrt(b*x + a - 1)))/x^3$

Fricas [A]

time = 0.42, size = 548, normalized size = 3.04



Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(b*x+a)/x^4,x, algorithm="fricas")`

[Out] $[1/6*((6*a^4 - 5*a^2 + 2)*sqrt(a^2 - 1)*b^3*x^3*log((a^2*b*x + a^3 + sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)*(a^2 + sqrt(a^2 - 1)*a - 1) + (a*b*x + a^2 - 1)*sqrt(a^2 - 1) - a)/x) + 4*(a^6 - 3*a^4 + 3*a^2 - 1)*b^3*x^3*arctan(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)) - (5*a^5 - 7*a^3 + 2*a)*b^3*x^3 - 2*(a^9 - 3*a^7 + 3*a^5 - a^3)*arccsc(b*x + a) - ((5*a^5 - 7*a^3 + 2*a)*b^2*x^2 - (a^6 - 2*a^4 + a^2)*b*x)*sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1))/((a^9 - 3*a^7 + 3*a^5 - a^3)*x^3), -1/6*(2*(6*a^4 - 5*a^2 + 2)*sqrt(-a^2 + 1)*b^3*x^3*arctan(-(sqrt(-a^2 + 1)*b*x - sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)*sqrt(-a^2 + 1))/(a^2 - 1)) - 4*(a^6 - 3*a^4 + 3*a^2 - 1)*b^3*x^3*arctan(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)) + (5*a^5 - 7*a^3 + 2*a)*b^3*x^3 + 2*(a^9 - 3*a^7 + 3*a^5 - a^3)*arccsc(b*x + a) + ((5*a^5 - 7*a^3 + 2*a)*b^2*x^2 - (a^6 - 2*a^4 + a^2)*b*x)*sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1))/((a^9 - 3*a^7 + 3*a^5 - a^3)*x^3)]$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{acsc}(a + bx)}{x^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(acsc(b*x+a)/x**4,x)`

[Out] $\text{Integral}(\text{acsc}(a + b*x)/x^{**4}, x)$

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 450 vs. $2(156) = 312$.

time = 0.50, size = 450, normalized size = 2.50

$$\frac{\left(\frac{(ba)^2 - 5a^2b^2 + 2b^2}{(a^2 - 2a^2 + a^2)x^2 + a^2}\right) \operatorname{atan}\left(\sqrt{\frac{1}{(ba + a^2)^2} + 1} - 1\right) + 4(ba + a^2)a^2b^2 \left(\sqrt{\frac{1}{(ba + a^2)^2} + 1} - 1\right)^3 + 10(ba + a^2)a^2b^2 \left(\sqrt{\frac{1}{(ba + a^2)^2} + 1} - 1\right)^5 - (ba + a^2)a^2b^2 \left(\sqrt{\frac{1}{(ba + a^2)^2} + 1} - 1\right)^7 + 16(ba + a^2)a^2b^2 \left(\sqrt{\frac{1}{(ba + a^2)^2} + 1} - 1\right)^9 - 2(ba + a^2)a^2b^2 \left(\sqrt{\frac{1}{(ba + a^2)^2} + 1} - 1\right)^{11} - 7(ba + a^2)a^2b^2 \left(\sqrt{\frac{1}{(ba + a^2)^2} + 1} - 1\right)^{13} + 5a^2b^2 - 2b^2}{(a^2 - 2a^2 + a^2)x^2 + a^2} + \frac{\left(\frac{4ba^2 - 2ab^2 - 4a^2b^2}{(a^2 - 2a^2 + a^2)x^2 + a^2}\right) \operatorname{arcsin}\left(\frac{1}{\sqrt{a^2 + (ba + a^2)^2}}\right)}{a^2\left(\frac{1}{a^2} - 1\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{integrate}(\text{arccsc}(b*x+a)/x^4, x, \text{algorithm}=\text{"giac"})$

[Out] $-1/3*b*((6*a^4*b^2 - 5*a^2*b^2 + 2*b^2)*\operatorname{arctan}((b*x + a)*(\sqrt{-1/(b*x + a)^2 + 1} - 1) + a)/\sqrt{-a^2 + 1})/((a^7 - 2*a^5 + a^3)*\sqrt{-a^2 + 1}) + (4*(b*x + a)^3*a^3*b^2*(\sqrt{-1/(b*x + a)^2 + 1} - 1)^3 + 10*(b*x + a)^2*a^4*b^2*(\sqrt{-1/(b*x + a)^2 + 1} - 1)^2 - (b*x + a)^3*a*b^2*(\sqrt{-1/(b*x + a)^2 + 1} - 1)^3 + (b*x + a)^2*a^2*b^2*(\sqrt{-1/(b*x + a)^2 + 1} - 1)^2 + 16*(b*x + a)^3*b^2*(\sqrt{-1/(b*x + a)^2 + 1} - 1)^2 - 2*(b*x + a)^2*b^2*(\sqrt{-1/(b*x + a)^2 + 1} - 1)^3 - 7*(b*x + a)*a*b^2*(\sqrt{-1/(b*x + a)^2 + 1} - 1)^2 + 5*a^2*b^2 - 2*b^2)/((a^6 - 2*a^4 + a^2)*((b*x + a)^2*(\sqrt{-1/(b*x + a)^2 + 1} - 1)^2 + 2*(b*x + a)*(\sqrt{-1/(b*x + a)^2 + 1} - 1)^2) + (3*a*b^2/(b*x + a) - 3*a^2*b^2/(b*x + a)^2 - b^2)*\operatorname{arcsin}(-1/((b*x + a)*(a/(b*x + a) - 1) - a))/(a^3*(a/(b*x + a) - 1)^3))$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sin\left(\frac{1}{a+b x}\right)}{x^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{int}(\text{asin}(1/(a + b*x))/x^4, x)$

[Out] $\text{int}(\text{asin}(1/(a + b*x))/x^4, x)$

3.26 $\int \frac{\csc^{-1}(a+bx)}{x^5} dx$

Optimal. Leaf size=239

$$\frac{b(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{12a(1-a^2)x^3} + \frac{(3-8a^2)b^2(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{24a^2(1-a^2)^2x^2} - \frac{(6-17a^2+26a^4)b^3(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{24a^3(1-a^2)^3x}$$

[Out] $1/4*b^4*arccsc(b*x+a)/a^4 - 1/4*arccsc(b*x+a)/x^4 + 1/4*(-8*a^6+8*a^4-7*a^2+2)*b^4*arctan((a-tan(1/2*arccsc(b*x+a)))/(-a^2+1)^(1/2))/a^4/(-a^2+1)^(7/2) - 1/12*b*(b*x+a)*(1-1/(b*x+a)^2)^(1/2)/a/(-a^2+1)/x^3 + 1/24*(-8*a^2+3)*b^2*(b*x+a)*(1-1/(b*x+a)^2)^(1/2)/a^2/(-a^2+1)^2/x^2 - 1/24*(26*a^4-17*a^2+6)*b^3*(b*x+a)*(1-1/(b*x+a)^2)^(1/2)/a^3/(-a^2+1)^3/x$

Rubi [A]

time = 0.32, antiderivative size = 239, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 9, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.900$, Rules used = {5367, 4512, 3870, 4145, 4004, 3916, 2739, 632, 210}

$$\frac{b^4 \csc^{-1}(a+bx)}{4a^4} + \frac{(3-8a^2)b^2(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{24a^2(1-a^2)^2x^2} - \frac{b(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{12a(1-a^2)x^3} + \frac{(-8a^6+8a^4-7a^2+2)b^4 \operatorname{ArcTan}\left(\frac{a-\tan(\frac{1}{2}\csc^{-1}(a+bx))}{\sqrt{1-a^2}}\right)}{4a^4(1-a^2)^{7/2}} - \frac{(26a^4-17a^2+6)b^3(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{24a^3(1-a^2)^3x} - \frac{\csc^{-1}(a+bx)}{4x^4}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{ArcCsc}[a+b*x]/x^5, x]$

[Out] $-1/12*(b*(a+b*x)*\text{Sqrt}[1-(a+b*x)^{-2}])/(a*(1-a^2)*x^3) + ((3-8*a^2)*b^2*(a+b*x)*\text{Sqrt}[1-(a+b*x)^{-2}])/(24*a^2*(1-a^2)^2*x^2) - ((6-17*a^2+26*a^4)*b^3*(a+b*x)*\text{Sqrt}[1-(a+b*x)^{-2}])/(24*a^3*(1-a^2)^3*x) + (b^4*\text{ArcCsc}[a+b*x])/(4*a^4) - \text{ArcCsc}[a+b*x]/(4*x^4) + ((2-7*a^2+8*a^4-8*a^6)*b^4*\text{ArcTan}[(a-\text{Tan}[\text{ArcCsc}[a+b*x]/2])/\text{Sqrt}[1-a^2]])/(4*a^4*(1-a^2)^{(7/2)})$

Rule 210

```
Int[((a_) + (b_)*(x_)^2)^{-1}, x_Symbol] :=> Simp[(-(Rt[-a, 2]*Rt[-b, 2])^{(-1)})*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])
```

Rule 632

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^{-1}, x_Symbol] :=> Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 2739

```
Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-1), x_Symbol] :> With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Dist[2*(e/d), Subst[Int[1/(a + 2*b*e*x + a*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]
```

Rule 3870

```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.)^(n_), x_Symbol] :> Simp[b^2*Cot[c + d*x]*((a + b*Csc[c + d*x])^(n + 1)/(a*d*(n + 1)*(a^2 - b^2))), x] + Dist[1/(a*(n + 1)*(a^2 - b^2)), Int[(a + b*Csc[c + d*x])^(n + 1)*Simp[(a^2 - b^2)*(n + 1) - a*b*(n + 1)*Csc[c + d*x] + b^2*(n + 2)*Csc[c + d*x]^2, x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]
```

Rule 3916

```
Int[csc[(e_.) + (f_.)*(x_)]/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol] :> Dist[1/b, Int[1/(1 + (a/b)*Sin[e + f*x]), x], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a^2 - b^2, 0]
```

Rule 4004

```
Int[(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_))/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol] :> Simp[c*(x/a), x] - Dist[(b*c - a*d)/a, Int[Csc[e + f*x]/(a + b*Csc[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]
```

Rule 4145

```
Int[((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.))*((csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_), x_Symbol] :> Simp[(A*b^2 - a*b*B + a^2*C)*Cot[e + f*x]*((a + b*Csc[e + f*x])^(m + 1)/(a*f*(m + 1)*(a^2 - b^2))), x] + Dist[1/(a*(m + 1)*(a^2 - b^2)), Int[(a + b*Csc[e + f*x])^(m + 1)*Simp[A*(a^2 - b^2)*(m + 1) - a*(A*b - a*B + b*C)*(m + 1)*Csc[e + f*x] + (A*b^2 - a*b*B + a^2*C)*(m + 2)*Csc[e + f*x]^2, x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && NeQ[a^2 - b^2, 0] && LtQ[m, -1]
```

Rule 4512

```
Int[Cot[(c_.) + (d_.)*(x_)]*Csc[(c_.) + (d_.)*(x_)]*(Csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_))^(n_)*((e_.) + (f_.)*(x_))^(m_), x_Symbol] :> Simp[(-(e + f*x)^m)*((a + b*Csc[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[f*(m/(b*d*(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Csc[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IntegerQ[m, 0] && NeQ[n, -1]
```

Rule 5367

```
Int[((a_.) + ArcCsc[(c_) + (d_)*(x_)]*(b_.))^(p_.)*((e_.) + (f_)*(x_))^(m_),
x_Symbol] :> Dist[-(d^(m + 1))^{(-1)}, Subst[Int[(a + b*x)^p*Csc[x]*Cot[x]*(d*e - c*f + f*Csc[x])^m, x], x, ArcCsc[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Rubi steps

$$\begin{aligned}
\int \frac{\csc^{-1}(a+bx)}{x^5} dx &= - \left(b^4 \text{Subst} \left(\int \frac{x \cot(x) \csc(x)}{(-a + \csc(x))^5} dx, x, \csc^{-1}(a+bx) \right) \right) \\
&= - \frac{\csc^{-1}(a+bx)}{4x^4} + \frac{1}{4} b^4 \text{Subst} \left(\int \frac{1}{(-a + \csc(x))^4} dx, x, \csc^{-1}(a+bx) \right) \\
&= - \frac{b(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{12a(1-a^2)x^3} - \frac{\csc^{-1}(a+bx)}{4x^4} - \frac{b^4 \text{Subst} \left(\int \frac{3(1-a^2)-3a \csc(x)-2 \csc^2(x)}{(-a+\csc(x))^3} dx, x, \csc^{-1}(a+bx) \right)}{12a(1-a^2)} \\
&= - \frac{b(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{12a(1-a^2)x^3} + \frac{(3-8a^2)b^2(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{24a^2(1-a^2)^2x^2} - \frac{\csc^{-1}(a+bx)}{4x^4} \\
&= - \frac{b(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{12a(1-a^2)x^3} + \frac{(3-8a^2)b^2(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{24a^2(1-a^2)^2x^2} - \frac{(6-17a^2+bx^2)\sqrt{1-\frac{1}{(a+bx)^2}}}{24a^2(1-a^2)^2x^2} \\
&= - \frac{b(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{12a(1-a^2)x^3} + \frac{(3-8a^2)b^2(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{24a^2(1-a^2)^2x^2} - \frac{(6-17a^2+bx^2)\sqrt{1-\frac{1}{(a+bx)^2}}}{24a^2(1-a^2)^2x^2} \\
&= - \frac{b(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{12a(1-a^2)x^3} + \frac{(3-8a^2)b^2(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{24a^2(1-a^2)^2x^2} - \frac{(6-17a^2+bx^2)\sqrt{1-\frac{1}{(a+bx)^2}}}{24a^2(1-a^2)^2x^2} \\
&= - \frac{b(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{12a(1-a^2)x^3} + \frac{(3-8a^2)b^2(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{24a^2(1-a^2)^2x^2} - \frac{(6-17a^2+bx^2)\sqrt{1-\frac{1}{(a+bx)^2}}}{24a^2(1-a^2)^2x^2} \\
&= - \frac{b(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{12a(1-a^2)x^3} + \frac{(3-8a^2)b^2(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{24a^2(1-a^2)^2x^2} - \frac{(6-17a^2+bx^2)\sqrt{1-\frac{1}{(a+bx)^2}}}{24a^2(1-a^2)^2x^2} \\
&= - \frac{b(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{12a(1-a^2)x^3} + \frac{(3-8a^2)b^2(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{24a^2(1-a^2)^2x^2} - \frac{(6-17a^2+bx^2)\sqrt{1-\frac{1}{(a+bx)^2}}}{24a^2(1-a^2)^2x^2}
\end{aligned}$$

Mathematica [C] Result contains complex when optimal does not.

time = 0.35, size = 307, normalized size = 1.28

$$\frac{1}{8} \left(\frac{\frac{b \sqrt{\frac{-1 + a^2 + 2 a b x + b^2 x^2}{(a + b x)^3}} (2 a^7 - 6 a^6 b x + 3 a^5 b^2 x^2 + 6 a^5 b^3 x^3 + a^3 (2 - 6 b^2 x^2) + 2 a^5 (-2 + 9 b^2 x^2) + a^4 b x (7 + 26 b^2 x^2) - a^2 (b x + 17 b^3 x^3))}{3 a^3 (-1 + a^2)^3 x^3} - \frac{2 \csc^{-1}(a + b x)}{x^4} + \frac{2 b^4 \text{ArcSin}\left(\frac{1}{\sqrt{a^2 + b^2 x^2}}\right)}{a^4} + \frac{i (-2 + 7 a^2 - 8 a^4 + 8 a^6) b^4 \log \left(\frac{16 a^4 (-1 + a^2)^3 \left(\frac{(i (-1 + a^2 + 8 a^4) \sqrt{1 - a^2} + (a + b x) \sqrt{-1 + a^2 + 2 a b x + b^2 x^2})}{(-2 + 7 a^2 - 8 a^4 + 8 a^6) b^2 x} \right)}{a^4 (1 - a^2)^{7/2}} \right)}{a^4 (1 - a^2)^{7/2}} \right)$$

Antiderivative was successfully verified.

[In] Integrate[ArcCsc[a + b*x]/x^5, x]

[Out] $\frac{((b * \text{Sqrt}[(-1 + a^2 + 2 * a * b * x + b^2 * x^2) / (a + b * x)^2] * (2 * a^7 - 6 * a^6 * b * x + 3 * a * b^2 * x^2 + 6 * b^3 * x^3 + a^3 * (2 - 6 * b^2 * x^2) + 2 * a^5 * (-2 + 9 * b^2 * x^2) + a^4 * b * x * (7 + 26 * b^2 * x^2) - a^2 * (b * x + 17 * b^3 * x^3))) / (3 * a^3 * (-1 + a^2)^3 * x^3) - (2 * \text{ArcCsc}[a + b * x]) / x^4 + (2 * b^4 * \text{ArcSin}[(a + b * x)^{-1}] / a^4 + (I * (-2 + 7 * a^2 - 8 * a^4 + 8 * a^6) * b^4 * \text{Log}[(16 * a^4 * (-1 + a^2)^3 * ((I * (-1 + a^2 + a * b * x)) / \text{Sqrt}[1 - a^2] + (a + b * x) * \text{Sqrt}[(-1 + a^2 + 2 * a * b * x + b^2 * x^2) / (a + b * x)^2])) / ((-2 + 7 * a^2 - 8 * a^4 + 8 * a^6) * b^4 * x)) / (a^4 * (1 - a^2)^{7/2})) / 8}{2(215) = 430}$

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 1701 vs. $2(215) = 430$.

time = 0.67, size = 1702, normalized size = 7.12

method	result	size
derivativeDivides	Expression too large to display	1702
default	Expression too large to display	1702

Verification of antiderivative is not currently implemented for this CAS.

[In] int(arccsc(b*x+a)/x^5, x, method=_RETURNVERBOSE)

[Out] $b^4 * (-1/4/b^4/x^4 * \text{arccsc}(b*x+a) - 1/24 * ((b*x+a)^2 - 1)^{1/2} * (18 * \text{arctan}(1 / ((b*x+a)^2 - 1)^{1/2})) * (a^2 - 1)^{3/2} * a^7 * (b*x+a)^2 + 27 * \ln(2 * ((a^2 - 1)^{1/2} * ((b*x+a)^2 - 1)^{1/2} + a * (b*x+a) - 1) / b*x) * a^5 - 6 * \text{arctan}(1 / ((b*x+a)^2 - 1)^{1/2}) * (a^2 - 1)^{3/2} * a^6 * (b*x+a)^3 + 54 * \text{arctan}(1 / ((b*x+a)^2 - 1)^{1/2}) * (a^2 - 1)^{3/2} * a^6 * (b*x+a)^2 - 54 * \text{arctan}(1 / ((b*x+a)^2 - 1)^{1/2}) * (a^2 - 1)^{3/2} * a^5 * (b*x+a)^2 + 18 * \text{arctan}(1 / ((b*x+a)^2 - 1)^{1/2}) * (a^2 - 1)^{3/2} * a^4 * (b*x+a)^3 + 60 * (a^2 - 1)^{3/2} * ((b*x+a)^2 - 1)^{1/2} * a^5 * (b*x+a)^2 - 26 * (a^2 - 1)^{3/2} * ((b*x+a)^2 - 1)^{1/2} * a^5 * (b*x+a)^2 - 54 * \text{arctan}(1 / ((b*x+a)^2 - 1)^{1/2}) * (a^2 - 1)^{3/2} * a^4 * (b*x+a) + 54 * \text{arctan}(1 / ((b*x+a)^2 - 1)^{1/2}) * (a^2 - 1)^{3/2} * a^3 * (b*x+a)^2 - 18 * \text{arctan}(1 / ((b*x+a)^2 - 1)^{1/2}) * (a^2 - 1)^{3/2} * a^4 * (b*x+a) + 17 * (a^2 - 1)^{3/2} * ((b*x+a)^2 - 1)^{1/2} * a^3 * (b*x+a)^2 - 18 * \text{arctan}(1 / ((b*x+a)^2 - 1)^{1/2}) * (a^2 - 1)^{3/2} * a^2 * (b*x+a)^2 + 15 * (a^2 - 1)^{3/2} * ((b*x+a)^2 - 1)^{1/2} * a^2 * (b*x+a)^2 - 6 * (a^2 - 1)^{3/2} * ((b*x+a)^2 - 1)^{1/2} * a * (b*x+a)^2 - 18 * \text{arctan}(1 / ((b*x+a)^2 - 1)^{1/2}) * (a^2 - 1)^{3/2} * a^8 * (b*x+a)^2 - 6 * \ln(2 * ((a^2 - 1)^{1/2} * ((b*x+a)^2 - 1)^{1/2} + a * (b*x+a) - 1) / b*x) * a^3 + 6 * \text{arctan}(1 / ((b*x+a)^2 - 1)^{1/2}) * (a^2 - 1)^{3/2} * (b*x+a)^3 - 11 * (a^2 - 1)^{3/2} * ((b*x+a)^2 - 1)^{1/2} * a^3 + 81 * \ln(2 * ((a^2 - 1)^{1/2} * ((b*x+a)^2 - 1)^{1/2} + a * (b*x+a) - 1) / b*x)$

$$\begin{aligned}
& \frac{1}{2} * ((b*x+a)^{2-1})^{(1/2)} + a * (b*x+a) - 1 / b/x * a^3 * (b*x+a)^{2-27} * \ln(2 * ((a^{2-1})^{(1/2)} * ((b*x+a)^{2-1})^{(1/2)} + a * (b*x+a) - 1 / b/x * a^2 * (b*x+a)^{3+72} * \ln(2 * ((a^{2-1})^{(1/2)} * ((b*x+a)^{2-1})^{(1/2)} + a * (b*x+a) - 1 / b/x * a^{10} * (b*x+a) - 72 * \ln(2 * ((a^{2-1})^{(1/2)} * ((b*x+a)^{2-1})^{(1/2)} + a * (b*x+a) - 1 / b/x * a^9 * (b*x+a)^{2+24} * \ln(2 * ((a^{2-1})^{(1/2)} * ((b*x+a)^{2-1})^{(1/2)} + a * (b*x+a) - 1 / b/x * a^8 * (b*x+a)^{3-18} * \arctan(1 / ((b*x+a)^{2-1})^{(1/2)} * (a^{2-1})^{(3/2)} * a^{7-36} * (a^{2-1})^{(3/2)} * ((b*x+a)^{2-1})^{(1/2)} * a^{7-14} * \ln(2 * ((a^{2-1})^{(1/2)} * ((b*x+a)^{2-1})^{(1/2)} + a * (b*x+a) - 1 / b/x * a^8 * (b*x+a) + 144 * \ln(2 * ((a^{2-1})^{(1/2)} * ((b*x+a)^{2-1})^{(1/2)} + a * (b*x+a) - 1 / b/x * a^7 * (b*x+a)^{2-48} * \ln(2 * ((a^{2-1})^{(1/2)} * ((b*x+a)^{2-1})^{(1/2)} + a * (b*x+a) - 1 / b/x * a^6 * (b*x+a)^{3+18} * \arctan(1 / ((b*x+a)^{2-1})^{(1/2)} * (a^{2-1})^{(3/2)} * a^{5+32} * (a^{2-1})^{(3/2)} * ((b*x+a)^{2-1})^{(1/2)} * a^{5+135} * \ln(2 * ((a^{2-1})^{(1/2)} * ((b*x+a)^{2-1})^{(1/2)} + a * (b*x+a) - 1 / b/x * a^6 * (b*x+a) - 135 * \ln(2 * ((a^{2-1})^{(1/2)} * ((b*x+a)^{2-1})^{(1/2)} + a * (b*x+a) - 1 / b/x * a^5 * (b*x+a)^{2+6} * \arctan(1 / ((b*x+a)^{2-1})^{(1/2)} * (a^{2-1})^{(3/2)} * a^{9-18} * \ln(2 * ((a^{2-1})^{(1/2)} * ((b*x+a)^{2-1})^{(1/2)} + a * (b*x+a) - 1 / b/x * a^4 * (b*x+a)^{2+45} * \ln(2 * ((a^{2-1})^{(1/2)} * ((b*x+a)^{2-1})^{(1/2)} + a * (b*x+a) - 1 / b/x * a^4 * (b*x+a)^{3-6} * \arctan(1 / ((b*x+a)^{2-1})^{(1/2)} * (a^{2-1})^{(3/2)} * a^{3-24} * \ln(2 * ((a^{2-1})^{(1/2)} * ((b*x+a)^{2-1})^{(1/2)} + a * (b*x+a) - 1 / b/x * a^{11+48} * \ln(2 * ((a^{2-1})^{(1/2)} * ((b*x+a)^{2-1})^{(1/2)} + a * (b*x+a) - 1 / b/x * a^9-45 * \ln(2 * ((a^{2-1})^{(1/2)} * ((b*x+a)^{2-1})^{(1/2)} + a * (b*x+a) - 1 / b/x * a^7+6 * \ln(2 * ((a^{2-1})^{(1/2)} * ((b*x+a)^{2-1})^{(1/2)} + a * (b*x+a) - 1 / b/x * (b*x+a)^{3+18} * \ln(2 * ((a^{2-1})^{(1/2)} * ((b*x+a)^{2-1})^{(1/2)} + a * (b*x+a) - 1 / b/x * a^2 * (b*x+a) - 81 * \ln(2 * ((a^{2-1})^{(1/2)} * ((b*x+a)^{2-1})^{(1/2)} + a * (b*x+a) - 1 / b/x * a^4 * (b*x+a) + 18 * \arctan(1 / ((b*x+a)^{2-1})^{(1/2)} * (a^{2-1})^{(3/2)} * a^{2 * (b*x+a)}) / (((b*x+a)^{2-1}) / (b*x+a)^2)^{(1/2)} / (b*x+a) / a^4 / (a^{2-1})^{(9/2)} / b^3 / x^3)
\end{aligned}$$
Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(arccsc(b*x+a)/x^5,x, algorithm="maxima")
[Out] -1/4*(4*x^4*integrate(1/4*(b^2*x + a*b)*e^(1/2*log(b*x + a + 1) + 1/2*log(b*x + a - 1)) / (b^2*x^6 + 2*a*b*x^5 + (a^2 - 1)*x^4 + (b^2*x^6 + 2*a*b*x^5 + (a^2 - 1)*x^4)*e^(log(b*x + a + 1) + log(b*x + a - 1))), x) + arctan2(1, sqrt(b*x + a + 1)*sqrt(b*x + a - 1))) / x^4
```

Fricas [A]

time = 0.44, size = 673, normalized size = 2.82

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(arccsc(b*x+a)/x^5,x, algorithm="fricas")
```

[Out] $\frac{1}{24} \cdot (3 \cdot (8 \cdot a^6 - 8 \cdot a^4 + 7 \cdot a^2 - 2) \cdot \sqrt{a^2 - 1} \cdot b^4 \cdot x^4 \cdot \log((a^2 \cdot b \cdot x + a^3 + \sqrt{b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 - 1}) \cdot (a^2 - \sqrt{a^2 - 1} \cdot a - 1) - (a \cdot b \cdot x + a^2 - 1) \cdot \sqrt{a^2 - 1} - a) / x) - 12 \cdot (a^8 - 4 \cdot a^6 + 6 \cdot a^4 - 4 \cdot a^2 + 1) \cdot b^4 \cdot x^4 \cdot \arctan(-b \cdot x - a + \sqrt{b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 - 1}) + (26 \cdot a^7 - 43 \cdot a^5 + 23 \cdot a^3 - 6 \cdot a) \cdot b^4 \cdot x^4 - 6 \cdot (a^{12} - 4 \cdot a^{10} + 6 \cdot a^8 - 4 \cdot a^6 + a^4) \cdot \text{arccsc}(b \cdot x + a) + ((26 \cdot a^7 - 43 \cdot a^5 + 23 \cdot a^3 - 6 \cdot a) \cdot b^3 \cdot x^3 - (8 \cdot a^8 - 19 \cdot a^6 + 14 \cdot a^4 - 3 \cdot a^2) \cdot b^2 \cdot x^2 + 2 \cdot (a^9 - 3 \cdot a^7 + 3 \cdot a^5 - a^3) \cdot b \cdot x) \cdot \sqrt{b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 - 1}) / ((a^{12} - 4 \cdot a^{10} + 6 \cdot a^8 - 4 \cdot a^6 + a^4) \cdot x^4), \frac{1}{24} \cdot (6 \cdot (8 \cdot a^6 - 8 \cdot a^4 + 7 \cdot a^2 - 2) \cdot \sqrt{-a^2 + 1} \cdot b^4 \cdot x^4 \cdot \arctan(-\sqrt{-a^2 + 1} \cdot b \cdot x - \sqrt{b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 - 1}) \cdot \sqrt{-a^2 + 1}) / (a^2 - 1) - 12 \cdot (a^8 - 4 \cdot a^6 + 6 \cdot a^4 - 4 \cdot a^2 + 1) \cdot b^4 \cdot x^4 \cdot \arctan(-b \cdot x - a + \sqrt{b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 - 1}) + (26 \cdot a^7 - 43 \cdot a^5 + 23 \cdot a^3 - 6 \cdot a) \cdot b^4 \cdot x^4 - 6 \cdot (a^{12} - 4 \cdot a^{10} + 6 \cdot a^8 - 4 \cdot a^6 + a^4) \cdot \text{arccsc}(b \cdot x + a) + ((26 \cdot a^7 - 43 \cdot a^5 + 23 \cdot a^3 - 6 \cdot a) \cdot b^3 \cdot x^3 - (8 \cdot a^8 - 19 \cdot a^6 + 14 \cdot a^4 - 3 \cdot a^2) \cdot b^2 \cdot x^2 + 2 \cdot (a^9 - 3 \cdot a^7 + 3 \cdot a^5 - a^3) \cdot b \cdot x) \cdot \sqrt{b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 - 1}) / ((a^{12} - 4 \cdot a^{10} + 6 \cdot a^8 - 4 \cdot a^6 + a^4) \cdot x^4)]$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{acsc}(a + bx)}{x^5} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(acsc(b*x+a)/x**5,x)

[Out] Integral(acsc(a + b*x)/x**5, x)

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 841 vs. 2(209) = 418.

time = 0.49, size = 841, normalized size = 3.52

$$\frac{1}{b^5} \left(\frac{1}{x^5} \operatorname{acsc}\left(\frac{a+b x}{b}\right) + \frac{5}{b^4} \left(\frac{1}{x^4} \operatorname{acsc}\left(\frac{a+b x}{b}\right) - \frac{1}{x^3} \operatorname{acsc}\left(\frac{a+b x}{b}\right) + \frac{1}{x^2} \operatorname{acsc}\left(\frac{a+b x}{b}\right) - \frac{1}{x} \operatorname{acsc}\left(\frac{a+b x}{b}\right) + \operatorname{acsc}\left(\frac{a+b x}{b}\right) \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(arccsc(b*x+a)/x**5,x, algorithm="giac")

[Out] $\frac{1}{12} \cdot b \cdot (3 \cdot (8 \cdot a^6 \cdot b^3 - 8 \cdot a^4 \cdot b^3 + 7 \cdot a^2 \cdot b^3 - 2 \cdot b^3) \cdot \arctan(((b \cdot x + a) \cdot (\sqrt{-1/(b \cdot x + a)^2 + 1} - 1) + a) / \sqrt{-a^2 + 1})) / ((a^{10} - 3 \cdot a^8 + 3 \cdot a^6 - a^4) \cdot \sqrt{-a^2 + 1}) + (18 \cdot (b \cdot x + a)^5 \cdot a^5 \cdot b^3 \cdot (\sqrt{-1/(b \cdot x + a)^2 + 1} - 1)^5 + 84 \cdot (b \cdot x + a)^4 \cdot a^6 \cdot b^3 \cdot (\sqrt{-1/(b \cdot x + a)^2 + 1} - 1)^4 + 104 \cdot (b \cdot x + a)^3 \cdot a^7 \cdot b^3 \cdot (\sqrt{-1/(b \cdot x + a)^2 + 1} - 1)^3 - 6 \cdot (b \cdot x + a)^5 \cdot a^3 \cdot b^3 \cdot (\sqrt{-1/(b \cdot x + a)^2 + 1} - 1)^5 - 12 \cdot (b \cdot x + a)^4 \cdot a^4 \cdot b^3 \cdot (\sqrt{-1/(b \cdot x + a)^2 + 1} - 1)^4 + 88 \cdot (b \cdot x + a)^3 \cdot a^5 \cdot b^3 \cdot (\sqrt{-1/(b \cdot x + a)^2 + 1} - 1)^3 + 3 \cdot (b \cdot x + a)^5 \cdot a^2 \cdot b^3 \cdot (\sqrt{-1/(b \cdot x + a)^2 + 1} - 1)^5 + 228 \cdot (b \cdot x + a)^2 \cdot a^6 \cdot b^3 \cdot (\sqrt{-1/(b \cdot x + a)^2 + 1} - 1)^2 - 3 \cdot (b \cdot x + a)^4 \cdot a^2 \cdot b^3 \cdot (\sqrt{-1/(b \cdot x + a)^2 + 1} - 1))$

$$\begin{aligned} & \sim 2 + 1) - 1)^4 - 78*(b*x + a)^3*a^3*b^3*(\sqrt{-1/(b*x + a)^2 + 1}) - 1)^3 - \\ & 114*(b*x + a)^2*a^4*b^3*(\sqrt{-1/(b*x + a)^2 + 1} - 1)^2 + 6*(b*x + a)^4*b^3*(\sqrt{-1/(b*x + a)^2 + 1} - 1)^4 + 138*(b*x + a)*a^5*b^3*(\sqrt{-1/(b*x + a)^2 + 1} - 1) + 36*(b*x + a)^3*a*b^3*(\sqrt{-1/(b*x + a)^2 + 1} - 1)^3 + 24*(b*x + a)^2*a^2*b^3*(\sqrt{-1/(b*x + a)^2 + 1} - 1)^2 - 96*(b*x + a)*a^3*b^3*(\sqrt{-1/(b*x + a)^2 + 1} - 1) + 26*a^4*b^3 + 12*(b*x + a)^2*b^3*(\sqrt{-1/(b*x + a)^2 + 1} - 1)^2 + 33*(b*x + a)*a*b^3*(\sqrt{-1/(b*x + a)^2 + 1} - 1) - 17*a^2*b^3 + 6*b^3)/((a^9 - 3*a^7 + 3*a^5 - a^3)*((b*x + a)^2*(\sqrt{-1/(b*x + a)^2 + 1} - 1)^2 + 2*(b*x + a)*a*(\sqrt{-1/(b*x + a)^2 + 1} - 1) + 1)^3) - 3*(4*a*b^3/(b*x + a) - 6*a^2*b^3/(b*x + a)^2 + 4*a^3*b^3/(b*x + a)^3 - b^3)*\arcsin(-1/((b*x + a)*(a/(b*x + a) - 1) - a))/(a^4*(a/(b*x + a) - 1)^4)) \end{aligned}$$
Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\arcsin\left(\frac{1}{a+bx}\right)}{x^5} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(asin(1/(a + b*x))/x^5, x)

[Out] int(asin(1/(a + b*x))/x^5, x)

$$3.27 \quad \int x^3 \csc^{-1}(a + bx)^2 dx$$

Optimal. Leaf size=366

$$-\frac{ax}{b^3} + \frac{(a+bx)^2}{12b^4} + \frac{(a+bx)\sqrt{1 - \frac{1}{(a+bx)^2}} \csc^{-1}(a+bx)}{3b^4} + \frac{3a^2(a+bx)\sqrt{1 - \frac{1}{(a+bx)^2}} \csc^{-1}(a+bx)}{b^4} - \frac{a(a+bx)^3}{b^5}$$

[Out] $-a*x/b^3 + 1/12*(b*x+a)^2/b^4 - 1/4*a^4*arccsc(b*x+a)^2/b^4 + 1/4*x^4*arccsc(b*x+a)^2 - 2*a*arccsc(b*x+a)*arctanh(I/(b*x+a) + (1-1/(b*x+a)^2)^(1/2))/b^4 - 4*a^3*a*rccsc(b*x+a)*arctanh(I/(b*x+a) + (1-1/(b*x+a)^2)^(1/2))/b^4 + 1/3*ln(b*x+a)/b^4 + 3*a^2*ln(b*x+a)/b^4 + 2*I*a^3*polylog(2, -I/(b*x+a) - (1-1/(b*x+a)^2)^(1/2))/b^4 - 2*I*a^3*polylog(2, I/(b*x+a) + (1-1/(b*x+a)^2)^(1/2))/b^4 - I*a*polylog(2, I/(b*x+a) + (1-1/(b*x+a)^2)^(1/2))/b^4 + I*a*polylog(2, -I/(b*x+a) - (1-1/(b*x+a)^2)^(1/2))/b^4 + 1/4*I*a*polylog(2, -I/(b*x+a) - (1-1/(b*x+a)^2)^(1/2))/b^4 + 1/3*(b*x+a)*arccsc(b*x+a)*(1-1/(b*x+a)^2)^(1/2)/b^4 + 3*a^2*(b*x+a)*arccsc(b*x+a)*(1-1/(b*x+a)^2)^(1/2)/b^4 - a*(b*x+a)^2*(b*x+a)*(1-1/(b*x+a)^2)^(1/2)/b^4 + 1/6*(b*x+a)^3*arccsc(b*x+a)*(1-1/(b*x+a)^2)^(1/2)/b^4$

Rubi [A]

time = 0.20, antiderivative size = 366, normalized size of antiderivative = 1.00, number of steps used = 20, number of rules used = 9, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {5367, 4512, 4275, 4268, 2317, 2438, 4269, 3556, 4270}

$$\frac{x^3(a+bx)^2}{4b^5} + \frac{2ia^2\text{Li}_2\left(\frac{-e^{i\pi(-1+bx)}}{b}\right)}{b^5} - \frac{2ia^2\text{Li}_2\left(\frac{e^{i\pi(-1+bx)}}{b}\right)}{b^5} - \frac{4a^3\csc^{-1}(a+bx)\tanh^{-1}\left(\frac{e^{i\pi(-1+bx)}}{b}\right)}{b^5} + \frac{3a^2(a+bx)\sqrt{1-(a+bx)^2}\csc^{-1}(a+bx)}{b^4} + \frac{i\text{Li}_2\left(\frac{-e^{i\pi(-1+bx)}}{b}\right)}{b^5} - \frac{i\text{Li}_2\left(\frac{e^{i\pi(-1+bx)}}{b}\right)}{b^5} + \frac{(a+bx)^2}{12b^4} + \frac{\log(a+bx)}{3b^2} - \frac{a(a+bx)^2}{6b^3} - \frac{(a+bx)^2\sqrt{1-(a+bx)^2}\csc^{-1}(a+bx)}{6b^2} + \frac{(a+bx)^2}{3b^2} - \frac{(a+bx)\sqrt{1-(a+bx)^2}\csc^{-1}(a+bx)}{3b^2} + \frac{2a\csc^{-1}(a+bx)\tanh^{-1}\left(\frac{e^{i\pi(-1+bx)}}{b}\right)}{b^5} - \frac{a\pi}{b^5} + \frac{1}{4}x^2\csc^{-1}(a+bx)^2$$

Antiderivative was successfully verified.

[In] $\text{Int}[x^3 \text{ArcCsc}[a + b*x]^2, x]$

[Out] $-((a*x)/b^3) + (a + b*x)^2/(12*b^4) + ((a + b*x)*\text{Sqrt}[1 - (a + b*x)^{-2}]*\text{ArcCsc}[a + b*x])/(3*b^4) + (3*a^2*(a + b*x)*\text{Sqrt}[1 - (a + b*x)^{-2}]*\text{ArcCsc}[a + b*x])/b^4 - (a*(a + b*x)^2*\text{Sqrt}[1 - (a + b*x)^{-2}]*\text{ArcCsc}[a + b*x])/b^4 + ((a + b*x)^3*\text{Sqrt}[1 - (a + b*x)^{-2}]*\text{ArcCsc}[a + b*x])/(6*b^4) - (a^4*\text{ArcCsc}[a + b*x]^2)/(4*b^4) + (x^4*\text{ArcCsc}[a + b*x]^2)/4 - (2*a*\text{ArcCsc}[a + b*x]*\text{ArcTanh}[E^{(I*\text{ArcCsc}[a + b*x])}])/b^4 - (4*a^3*\text{ArcCsc}[a + b*x]*\text{ArcTanh}[E^{(I*\text{ArcCsc}[a + b*x])}])/b^4 + \text{Log}[a + b*x]/(3*b^4) + (3*a^2*\text{Log}[a + b*x])/b^4 + (I*a*\text{PolyLog}[2, -E^{(I*\text{ArcCsc}[a + b*x])}])/b^4 + ((2*I)*a^3*\text{PolyLog}[2, -E^{(I*\text{ArcCsc}[a + b*x])}])/b^4 - (I*a*\text{PolyLog}[2, E^{(I*\text{ArcCsc}[a + b*x])}])/b^4 - ((2*I)*a^3*\text{PolyLog}[2, E^{(I*\text{ArcCsc}[a + b*x])}])/b^4$

Rule 2317

```
Int[Log[(a_) + (b_)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol]
:= Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2438

```
Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 3556

```
Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

Rule 4268

```
Int[csc[(e_.) + (f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_), x_Symbol] :> Simp[-2*(c + d*x)^m*(ArcTanh[E^(I*(e + f*x))/f]), x] + (-Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Log[1 - E^(I*(e + f*x))], x], x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Log[1 + E^(I*(e + f*x))], x], x]) /; FreeQ[{c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 4269

```
Int[csc[(e_.) + (f_.)*(x_)]^2*((c_.) + (d_.)*(x_))^(m_), x_Symbol] :> Simp[((-c + d*x)^m)*(Cot[e + f*x]/f), x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Cot[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

Rule 4270

```
Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^n*((c_.) + (d_.)*(x_)), x_Symbol] :> Simp[(-b^2)*(c + d*x)*Cot[e + f*x]*((b*Csc[e + f*x])^(n - 2)/(f*(n - 1))), x] + (Dist[b^2*((n - 2)/(n - 1)), Int[(c + d*x)*(b*Csc[e + f*x])^(n - 2), x], x] - Simp[b^2*d*((b*Csc[e + f*x])^(n - 2)/(f^2*(n - 1)*(n - 2))), x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && NeQ[n, 2]
```

Rule 4275

```
Int[(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^n*((c_.) + (d_.)*(x_))^(m_), x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (a + b*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rule 4512

```
Int[Cot[(c_.) + (d_.)*(x_)]*Csc[(c_.) + (d_.)*(x_)]*(Csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_.))^n*((e_.) + (f_.)*(x_))^(m_), x_Symbol] :> Simp[((-e + f*x)^m)*((a + b*Csc[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[f*(m/(b*d*(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Csc[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]
```

Rule 5367

```
Int[((a_) + ArcCsc[(c_) + (d_)*(x_)]*(b_.))^(p_.)*((e_.) + (f_)*(x_))^m_, x_Symbol] :> Dist[-(d^(m + 1))^{-1}, Subst[Int[(a + b*x)^p*Csc[x]*Cot[x]*(d*e - c*f + f*Csc[x])^m, x], x, ArcCsc[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Rubi steps

$$\begin{aligned}
\int x^3 \csc^{-1}(a + bx)^2 dx &= -\frac{\text{Subst}\left(\int x^2 \cot(x) \csc(x)(-a + \csc(x))^3 dx, x, \csc^{-1}(a + bx)\right)}{b^4} \\
&= \frac{1}{4} x^4 \csc^{-1}(a + bx)^2 - \frac{\text{Subst}\left(\int x(-a + \csc(x))^4 dx, x, \csc^{-1}(a + bx)\right)}{2b^4} \\
&= \frac{1}{4} x^4 \csc^{-1}(a + bx)^2 - \frac{\text{Subst}\left(\int (a^4 x - 4a^3 x \csc(x) + 6a^2 x \csc^2(x) - 4ax \csc^3(x) + \dots) dx, x, \csc^{-1}(a + bx)\right)}{2b^4} \\
&= -\frac{a^4 \csc^{-1}(a + bx)^2}{4b^4} + \frac{1}{4} x^4 \csc^{-1}(a + bx)^2 - \frac{\text{Subst}\left(\int x \csc^4(x) dx, x, \csc^{-1}(a + bx)\right)}{2b^4} \\
&= -\frac{ax}{b^3} + \frac{(a + bx)^2}{12b^4} + \frac{3a^2(a + bx)\sqrt{1 - \frac{1}{(a + bx)^2}} \csc^{-1}(a + bx)}{b^4} - \frac{a(a + bx)^2\sqrt{1 - \frac{1}{(a + bx)^2}}}{b^4} \\
&= -\frac{ax}{b^3} + \frac{(a + bx)^2}{12b^4} + \frac{(a + bx)\sqrt{1 - \frac{1}{(a + bx)^2}} \csc^{-1}(a + bx)}{3b^4} + \frac{3a^2(a + bx)\sqrt{1 - \frac{1}{(a + bx)^2}}}{b^4} \\
&= -\frac{ax}{b^3} + \frac{(a + bx)^2}{12b^4} + \frac{(a + bx)\sqrt{1 - \frac{1}{(a + bx)^2}} \csc^{-1}(a + bx)}{3b^4} + \frac{3a^2(a + bx)\sqrt{1 - \frac{1}{(a + bx)^2}}}{b^4} \\
&= -\frac{ax}{b^3} + \frac{(a + bx)^2}{12b^4} + \frac{(a + bx)\sqrt{1 - \frac{1}{(a + bx)^2}} \csc^{-1}(a + bx)}{3b^4} + \frac{3a^2(a + bx)\sqrt{1 - \frac{1}{(a + bx)^2}}}{b^4}
\end{aligned}$$

Mathematica [A]

time = 3.48, size = 420, normalized size = 1.15

```
-----
```

Warning: Unable to verify antiderivative.

[In] `Integrate[x^3*ArcCsc[a + b*x]^2, x]`

[Out] `(-16*(6*a - 2*(1 + 9*a^2)*ArcCsc[a + b*x] + 3*(a + 2*a^3)*ArcCsc[a + b*x]^2)*Cot[ArcCsc[a + b*x]/2] + 2*(2 - 24*a*ArcCsc[a + b*x] + (3 + 36*a^2)*ArcCsc[a + b*x]^2)*Csc[ArcCsc[a + b*x]/2]^2 + 3*ArcCsc[a + b*x]^2*Csc[ArcCsc[a + b*x]/2] + 3*ArcCsc[a + b*x]^2*Csc[ArcCsc[a + b*x]/2]^2)`

$$\begin{aligned} & b*x]/2]^4 - (2*ArcCsc[a + b*x]*(-1 + 6*a*ArcCsc[a + b*x])*Csc[ArcCsc[a + b*x]/2]^4)/(a + b*x) - 64*(1 + 9*a^2)*Log[(a + b*x)^{-1}] + 192*(a + 2*a^3)*(ArcCsc[a + b*x]*(Log[1 - E^(I*ArcCsc[a + b*x])] - Log[1 + E^(I*ArcCsc[a + b*x])]) + I*(PolyLog[2, -E^(I*ArcCsc[a + b*x])] - PolyLog[2, E^(I*ArcCsc[a + b*x])])) + 2*(2 + 24*a*ArcCsc[a + b*x] + (3 + 36*a^2)*ArcCsc[a + b*x]^2)*Sec[ArcCsc[a + b*x]/2]^2 + 3*ArcCsc[a + b*x]^2*Sec[ArcCsc[a + b*x]/2]^4 - 32*(a + b*x)^3*ArcCsc[a + b*x]*(1 + 6*a*ArcCsc[a + b*x])*Sin[ArcCsc[a + b*x]/2]^4 - 16*(6*a + 2*(1 + 9*a^2)*ArcCsc[a + b*x] + 3*(a + 2*a^3)*ArcCsc[a + b*x]^2)*Tan[ArcCsc[a + b*x]/2])/(192*b^4) \end{aligned}$$

Maple [A]

time = 12.52, size = 703, normalized size = 1.92

method	result
derivative divides	$\frac{(bx+a)^2}{12} - \frac{\ln\left(1+\frac{i}{bx+a}+\sqrt{1-\frac{1}{(bx+a)^2}}\right)}{3} + \frac{2\ln\left(\frac{i}{bx+a}+\sqrt{1-\frac{1}{(bx+a)^2}}\right)}{3} - \frac{\ln\left(\frac{i}{bx+a}+\sqrt{1-\frac{1}{(bx+a)^2}-1}\right)}{3} - a$
default	$\frac{(bx+a)^2}{12} - \frac{\ln\left(1+\frac{i}{bx+a}+\sqrt{1-\frac{1}{(bx+a)^2}}\right)}{3} + \frac{2\ln\left(\frac{i}{bx+a}+\sqrt{1-\frac{1}{(bx+a)^2}}\right)}{3} - \frac{\ln\left(\frac{i}{bx+a}+\sqrt{1-\frac{1}{(bx+a)^2}-1}\right)}{3} - a$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^3*arccsc(b*x+a)^2,x,method=_RETURNVERBOSE)

[Out]

$$\begin{aligned} & 1/b^4*(-1/3*\ln(1+I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))+1/12*(b*x+a)^2+2/3*\ln(I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))-1/3*\ln(I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2)-1)-a*(b*x+a)-a*arccsc(b*x+a)*\ln(1+I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))+I*a*polylog(2,-I/(b*x+a)-(1-1/(b*x+a)^2)^(1/2))+a*arccsc(b*x+a)*\ln(1-I/(b*x+a)-(1-1/(b*x+a)^2)^(1/2))-I*a*polylog(2,I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))-2*a^3*arccsc(b*x+a)*\ln(1+I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))+2*I*a^3*polylog(2,-I/(b*x+a)-(1-1/(b*x+a)^2)^(1/2))+2*a^3*arccsc(b*x+a)*\ln(1-I/(b*x+a)-(1-1/(b*x+a)^2)^(1/2))-2*I*a^3*polylog(2,I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))-arccsc(b*x+a)^2*a^3*(b*x+a)+3/2*arccsc(b*x+a)^2*a^2*(b*x+a)^2-arccsc(b*x+a)^2*a^2*(b*x+a)^3+1/6*arccsc(b*x+a)*((b*x+a)^2-1)/(b*x+a)^2)^(1/2)*(b*x+a)^3-3*I*a^2*arccsc(b*x+a)+1/3*arccsc(b*x+a)*((b*x+a)^2-1)/(b*x+a)^2)^(1/2)*(b*x+a)^3+3*arccsc(b*x+a)*((b*x+a)^2-1)/(b*x+a)^2)^(1/2)*a^2*(b*x+a)-arccsc(b*x+a)*((b*x+a)^2-1)/(b*x+a)^2)^(1/2)*a*(b*x+a)^2-3*a^2*ln(I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2)-1)-3*a^2*ln(1+I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))+6*a^2*ln(I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))+1/4*arccsc(b*x+a)^2*(b*x+a)^4-1/3*I*arccsc(b*x+a)) \end{aligned}$$
Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^3*arccsc(b*x+a)^2,x, algorithm="maxima")`

[Out]
$$\frac{1}{4}x^4 \operatorname{arctan2}\left(\sqrt{bx + a + 1}, \sqrt{bx + a - 1}\right)^2 - \frac{1}{16}x^4 \log(b^2x^2 + 2ax + a^2)^2 + \operatorname{integrate}\left(\frac{1}{4}(2\sqrt{bx + a + 1})^2 \sqrt{bx + a - 1} \cdot \frac{1}{4}x^4 \operatorname{arctan2}\left(\sqrt{bx + a + 1}, \sqrt{bx + a - 1}\right) - 4(b^3x^6 + 3a^2b^2x^5 + (3a^2 - 1)b^2x^4 + (a^3 - a)x^3) \log(bx + a)^2 + (b^3x^6 + 2a^2b^2x^5 + (a^2 - 1)b^2x^4 + 4(b^3x^6 + 3a^2b^2x^5 + (3a^2 - 1)b^2x^4 + (a^3 - a)x^3) \log(bx + a)) \log(b^2x^2 + 2ax + a^2) / (b^3x^3 + 3a^2b^2x^2 + a^3 + (3a^2 - 1)b^2x - a), x\right)$$

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^3*arccsc(b*x+a)^2,x, algorithm="fricas")`

[Out] `integral(x^3*arccsc(b*x + a)^2, x)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x^3 \operatorname{acsc}^2(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x**3*acsc(b*x+a)**2,x)`

[Out] `Integral(x**3*acsc(a + b*x)**2, x)`

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: RuntimeError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^3*arccsc(b*x+a)^2,x, algorithm="giac")`

[Out] Exception raised: RuntimeError >> An error occurred running a Giac command:
 INPUT:sage2OUTPUT:Warning, integration of abs or sign assumes constant sign
 by intervals (correct if the argument is real):Check [abs(sageVARa+sageVAR
 b*sageVAR

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int x^3 \operatorname{asin}\left(\frac{1}{a + bx}\right)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x^3*asin(1/(a + b*x))^2,x)`

[Out] `int(x^3*asin(1/(a + b*x))^2, x)`

$$3.28 \quad \int x^2 \csc^{-1}(a + bx)^2 dx$$

Optimal. Leaf size=272

$$\frac{x}{3b^2} - \frac{2a(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}\csc^{-1}(a+bx)}{b^3} + \frac{(a+bx)^2\sqrt{1-\frac{1}{(a+bx)^2}}\csc^{-1}(a+bx)}{3b^3} + \frac{a^3\csc^{-1}(a+bx)}{3b^3}$$

[Out] $\frac{1}{3}x/b^2 + \frac{1}{3}a^3 \operatorname{arccsc}(b*x+a)^2/b^3 + \frac{1}{3}x^3 \operatorname{arccsc}(b*x+a)^2 + \frac{2}{3} \operatorname{arccsc}(b*x+a) \operatorname{arctanh}\left(\frac{I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2)}{b^3}\right) + \frac{4*a^2 \operatorname{arccsc}(b*x+a) \operatorname{arctanh}\left(\frac{I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2)}{b^3}\right)}{3} - \frac{2*a^3 \operatorname{arccsc}(b*x+a) \operatorname{ln}(b*x+a)}{b^3} - \frac{1}{3}I \operatorname{polylog}(2, -I/(b*x+a) - (1-1/(b*x+a)^2)^(1/2)) + \frac{2*I \operatorname{polylog}(2, I/(b*x+a) + (1-1/(b*x+a)^2)^(1/2))}{b^3} + \frac{2*I^2 \operatorname{polylog}(2, I/(b*x+a) + (1-1/(b*x+a)^2)^(1/2))}{b^3} - \frac{2*a^2 \operatorname{polylog}(2, I/(b*x+a) + (1-1/(b*x+a)^2)^(1/2))}{b^3} + \frac{2*a^3 \operatorname{arccsc}(b*x+a) \operatorname{arccsc}(b*x+a) \operatorname{arctanh}\left(\frac{I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2)}{b^3}\right)}{3} + \frac{1}{3}x^3 \operatorname{arccsc}(b*x+a) \operatorname{arccsc}(b*x+a)$

Rubi [A]

time = 0.16, antiderivative size = 272, normalized size of antiderivative = 1.00, number of steps used = 17, number of rules used = 9, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {5367, 4512, 4275, 4268, 2317, 2438, 4269, 3556, 4270}

$$\frac{a^3 \csc^{-1}(a+bx)^2}{3b^3} - \frac{2a^2 \operatorname{Li}_2\left(-e^{i \csc^{-1}(a+bx)}\right)}{b^3} + \frac{2ia^2 \operatorname{Li}_2\left(e^{i \csc^{-1}(a+bx)}\right)}{b^3} + \frac{4a^2 \csc^{-1}(a+bx) \tanh^{-1}\left(e^{i \csc^{-1}(a+bx)}\right)}{b^3} - \frac{i \operatorname{Li}_2\left(-e^{i \csc^{-1}(a+bx)}\right)}{3b^3} + \frac{i \operatorname{Li}_2\left(e^{i \csc^{-1}(a+bx)}\right)}{3b^3} - \frac{2a \log(a+bx)}{b^3} - \frac{2a(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}\csc^{-1}(a+bx)}{b^3} + \frac{(a+bx)^2\sqrt{1-\frac{1}{(a+bx)^2}}\csc^{-1}(a+bx)}{3b^3} + \frac{2 \csc^{-1}(a+bx) \tanh^{-1}\left(e^{i \csc^{-1}(a+bx)}\right)}{3b^3} + \frac{1}{3}x^3 \csc^{-1}(a+bx)^2 + \frac{x}{3b^2}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[x^2 \operatorname{ArcCsc}[a + b*x]^2, x]$

[Out] $x/(3*b^2) - (2*a*(a + b*x)*\operatorname{Sqrt}[1 - (a + b*x)^{-2}]*\operatorname{ArcCsc}[a + b*x])/b^3 + ((a + b*x)^2*\operatorname{Sqrt}[1 - (a + b*x)^{-2}]*\operatorname{ArcCsc}[a + b*x])/(3*b^3) + (a^3*\operatorname{ArcCsc}[a + b*x]^2)/(3*b^3) + (x^3*\operatorname{ArcCsc}[a + b*x]^2)/3 + (2*\operatorname{ArcCsc}[a + b*x]*\operatorname{ArcTanh}[E^(\operatorname{I}*\operatorname{ArcCsc}[a + b*x]))]/(3*b^3) + (4*a^2*\operatorname{ArcCsc}[a + b*x]*\operatorname{ArcTanh}[E^(\operatorname{I}*\operatorname{ArcCsc}[a + b*x]))]/b^3 - (2*a*\operatorname{Log}[a + b*x])/b^3 - ((\operatorname{I}/3)*\operatorname{PolyLog}[2, -E^(\operatorname{I}*\operatorname{ArcCsc}[a + b*x]))]/b^3 - ((2*\operatorname{I})*a^2*\operatorname{PolyLog}[2, -E^(\operatorname{I}*\operatorname{ArcCsc}[a + b*x]))]/b^3 + ((\operatorname{I}/3)*\operatorname{PolyLog}[2, E^(\operatorname{I}*\operatorname{ArcCsc}[a + b*x))]/b^3 + ((2*\operatorname{I})*a^2*\operatorname{PolyLog}[2, E^(\operatorname{I}*\operatorname{ArcCsc}[a + b*x))]/b^3$

Rule 2317

```
Int[Log[(a_) + (b_.)*((F_)^((e_.)*(c_.) + (d_.)*(x_)))^(n_.)], x_Symbol]
 :> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]]
```

Rule 2438

```
Int[Log[(c_.)*(d_.) + (e_.)*(x_)^(n_.)]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]]
```

Rule 3556

```
Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-Log[RemoveContent[Cos[c + d *x], x]]/d, x] /; FreeQ[{c, d}, x]
```

Rule 4268

```
Int[csc[(e_.) + (f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] :> Simp[-2*(c + d*x)^m*(ArcTanh[E^(I*(e + f*x))/f], x] + (-Dist[d*(m/f), Int[(c + d *x)^(m - 1)*Log[1 - E^(I*(e + f*x))], x], x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Log[1 + E^(I*(e + f*x))], x], x]) /; FreeQ[{c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 4269

```
Int[csc[(e_.) + (f_.)*(x_)]^2*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] :> Simp[[-(c + d*x)^m)*(Cot[e + f*x]/f), x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Cot[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

Rule 4270

```
Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(n_)*((c_.) + (d_.)*(x_)), x_Symbol] :> Simp[(-b^2)*(c + d*x)*Cot[e + f*x]*((b*Csc[e + f*x])^(n - 2)/(f*(n - 1))), x] + (Dist[b^2*((n - 2)/(n - 1)), Int[(c + d*x)*(b*Csc[e + f*x])^(n - 2), x], x] - Simp[b^2*d*((b*Csc[e + f*x])^(n - 2)/(f^2*(n - 1)*(n - 2))), x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && NeQ[n, 2]
```

Rule 4275

```
Int[(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(n_)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (a + b*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rule 4512

```
Int[Cot[(c_.) + (d_.)*(x_)]*Csc[(c_.) + (d_.)*(x_)]*(Csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_.))^(n_)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] :> Simp[(-(e + f*x)^m)*((a + b*Csc[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[f*(m/(b*d*(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Csc[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]
```

Rule 5367

```
Int[((a_.) + ArcCsc[(c_.) + (d_.)*(x_)]*(b_.))^(p_)*((e_.) + (f_.)*(x_))^(m _.), x_Symbol] :> Dist[-(d^(m + 1))^(-1), Subst[Int[(a + b*x)^p*Csc[x]*Cot[x]*(d*e - c*f + f*Csc[x])^m, x], x, ArcCsc[c + d*x]], x] /; FreeQ[{a, b, c,
```

$d, e, f}, x] \&& \text{IGtQ}[p, 0] \&& \text{IntegerQ}[m]$

Rubi steps

$$\begin{aligned}
\int x^2 \csc^{-1}(a+bx)^2 dx &= -\frac{\text{Subst}(\int x^2 \cot(x) \csc(x)(-a+\csc(x))^2 dx, x, \csc^{-1}(a+bx))}{b^3} \\
&= \frac{1}{3}x^3 \csc^{-1}(a+bx)^2 - \frac{2\text{Subst}(\int x(-a+\csc(x))^3 dx, x, \csc^{-1}(a+bx))}{3b^3} \\
&= \frac{1}{3}x^3 \csc^{-1}(a+bx)^2 - \frac{2\text{Subst}(\int (-a^3x + 3a^2x \csc(x) - 3ax \csc^2(x) + x \csc^3(x)) dx, x, \csc^{-1}(a+bx))}{3b^3} \\
&= \frac{a^3 \csc^{-1}(a+bx)^2}{3b^3} + \frac{1}{3}x^3 \csc^{-1}(a+bx)^2 - \frac{2\text{Subst}(\int x \csc^3(x) dx, x, \csc^{-1}(a+bx))}{3b^3} \\
&= \frac{x}{3b^2} - \frac{2a(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}} \csc^{-1}(a+bx)}{b^3} + \frac{(a+bx)^2\sqrt{1-\frac{1}{(a+bx)^2}} \csc^{-1}(a+bx)}{3b^3} \\
&= \frac{x}{3b^2} - \frac{2a(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}} \csc^{-1}(a+bx)}{b^3} + \frac{(a+bx)^2\sqrt{1-\frac{1}{(a+bx)^2}} \csc^{-1}(a+bx)}{3b^3} \\
&= \frac{x}{3b^2} - \frac{2a(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}} \csc^{-1}(a+bx)}{b^3} + \frac{(a+bx)^2\sqrt{1-\frac{1}{(a+bx)^2}} \csc^{-1}(a+bx)}{3b^3} \\
&= \frac{x}{3b^2} - \frac{2a(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}} \csc^{-1}(a+bx)}{b^3} + \frac{(a+bx)^2\sqrt{1-\frac{1}{(a+bx)^2}} \csc^{-1}(a+bx)}{3b^3}
\end{aligned}$$

Mathematica [A]

time = 3.14, size = 314, normalized size = 1.15

$$\begin{aligned}
&-22 - 12\pi \cos^{-1}(a+bx) + (1 + 6a^2)\cos^{-1}(a+bx)^2 \cos(2\pi \cos^{-1}(a+bx)) + 2\cos^{-1}(a+bx)(-1 + 3a\cos^{-1}(a+bx))\cos^2(\frac{1}{2}\cos^{-1}(a+bx)) - \frac{-27\cos^{-1}(a+bx)(-1 + 3a\cos^{-1}(a+bx))}{288} - 36a\log(\frac{1}{2}\pi b^2) + 6\pi^2 \left(\cos^{-1}(a+bx) \log\left(1 - e^{i\pi \cos^{-1}(a+bx)}\right) + i \left(\text{PolyLog}_2\left(2, -e^{i\pi \cos^{-1}(a+bx)}\right) - \text{PolyLog}_2\left(2, e^{i\pi \cos^{-1}(a+bx)}\right)\right)\right) + 2\cos^{-1}(a+bx)(1 + 3a\cos^{-1}(a+bx))\cos^2(\frac{1}{2}\cos^{-1}(a+bx)) - 6(a+bx)^2\cos^2(\frac{1}{2}\cos^{-1}(a+bx)) - 2(2 + 13\cos^{-1}(a+bx) + (1 + 6a^2)\cos^{-1}(a+bx)^2) \tan(\frac{1}{2}\cos^{-1}(a+bx))
\end{aligned}$$

Warning: Unable to verify antiderivative.

[In] `Integrate[x^2*ArcCsc[a + b*x]^2, x]`

[Out]
$$\begin{aligned}
&-1/24*(-2*(2 - 12*a*ArcCsc[a + b*x] + (1 + 6*a^2)*ArcCsc[a + b*x]^2)*Cot[Ar\\
&cCsc[a + b*x]/2] + 2*ArcCsc[a + b*x]*(-1 + 3*a*ArcCsc[a + b*x])*Csc[ArcCsc[\\
&a + b*x]/2]^2 - (ArcCsc[a + b*x]^2*Csc[ArcCsc[a + b*x]/2]^4)/(2*(a + b*x))\\
&- 48*a*Log[(a + b*x)^{-1}] + 8*(1 + 6*a^2)*(ArcCsc[a + b*x]*(Log[1 - E^{(I*Ar\\
&cCsc[a + b*x])}] - Log[1 + E^{(I*ArcCsc[a + b*x])}]) + I*(PolyLog[2, -E^{(I*Ar\\
&cCsc[a + b*x])}] - PolyLog[2, E^{(I*ArcCsc[a + b*x])}])) + 2*ArcCsc[a + b*x]*(
\end{aligned}$$

$$1 + 3*a*ArcCsc[a + b*x])*Sec[ArcCsc[a + b*x]/2]^2 - 8*(a + b*x)^3*ArcCsc[a + b*x]^2*Sin[ArcCsc[a + b*x]/2]^4 - 2*(2 + 12*a*ArcCsc[a + b*x] + (1 + 6*a^2)*ArcCsc[a + b*x]^2)*Tan[ArcCsc[a + b*x]/2])/b^3$$

Maple [A]

time = 10.44, size = 500, normalized size = 1.84

method	result
derivativedivides	$\arccsc(bx+a)^2 a^2 (bx+a) - \arccsc(bx+a)^2 a (bx+a)^2 + \frac{\arccsc(bx+a)^2 (bx+a)^3}{3} - 2 \arccsc(bx+a) \sqrt{\frac{(bx+a)^2 - 1}{(bx+a)^2}} a (bx+a) + \dots$
default	$\arccsc(bx+a)^2 a^2 (bx+a) - \arccsc(bx+a)^2 a (bx+a)^2 + \frac{\arccsc(bx+a)^2 (bx+a)^3}{3} - 2 \arccsc(bx+a) \sqrt{\frac{(bx+a)^2 - 1}{(bx+a)^2}} a (bx+a) + \dots$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x^2*arccsc(b*x+a)^2,x,method=_RETURNVERBOSE)`

[Out]

$$\begin{aligned} & 1/b^3 * (\arccsc(b*x+a)^2 * a^2 * (b*x+a) - \arccsc(b*x+a)^2 * a * (b*x+a)^2 + 1/3 * \arccsc(b*x+a)^2 * a^3 * (b*x+a)^3) \\ & + 1/3 * \arccsc(b*x+a) * ((b*x+a)^2 - 1) / (b*x+a)^2)^{(1/2)} * a * (b*x+a) \\ & + 1/3 * \arccsc(b*x+a) * ((b*x+a)^2 - 1) / (b*x+a)^2)^{(1/2)} * (b*x+a)^2 + 1/3 * I * \text{polylog}(2, I / (b*x+a) + (1 - 1 / (b*x+a)^2)^{(1/2)}) + 1/3 * b*x + 1/3 * \arccsc(b*x+a) * \ln(1 + I / (b*x+a) + (1 - 1 / (b*x+a)^2)^{(1/2)}) + 2 * I * \arccsc(b*x+a) * a - 1/3 * \arccsc(b*x+a) * \ln(1 - I / (b*x+a) - (1 - 1 / (b*x+a)^2)^{(1/2)}) - 2 * I * a^2 * \text{polylog}(2, -I / (b*x+a) - (1 - 1 / (b*x+a)^2)^{(1/2)}) + 2 * a * \ln(1 + I / (b*x+a) + (1 - 1 / (b*x+a)^2)^{(1/2)}) + 2 * a * \ln(I / (b*x+a) + (1 - 1 / (b*x+a)^2)^{(1/2)} - 1) - 4 * a * \ln(I / (b*x+a) + (1 - 1 / (b*x+a)^2)^{(1/2)}) + 2 * a^2 * \arccsc(b*x+a) * \ln(1 + I / (b*x+a) + (1 - 1 / (b*x+a)^2)^{(1/2)}) - 1/3 * I * \text{polylog}(2, -I / (b*x+a) - (1 - 1 / (b*x+a)^2)^{(1/2)}) - 2 * a^2 * \arccsc(b*x+a) * \ln(1 - I / (b*x+a) - (1 - 1 / (b*x+a)^2)^{(1/2)}) + 2 * I * a^2 * \text{polylog}(2, I / (b*x+a) + (1 - 1 / (b*x+a)^2)^{(1/2)})) \end{aligned}$$
Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^2*arccsc(b*x+a)^2,x, algorithm="maxima")`

[Out]

$$\begin{aligned} & 1/3 * x^3 * \text{arctan2}(1, \sqrt{b*x + a + 1}) * \sqrt{b*x + a - 1})^2 - 1/12 * x^3 * \log(b^2 * x^2 + 2 * a * b * x + a^2)^2 + \text{integrate}(1/3 * (2 * \sqrt{b*x + a + 1}) * \sqrt{b*x + a - 1}) * b*x^3 * \text{arctan2}(1, \sqrt{b*x + a + 1}) * \sqrt{b*x + a - 1}) - 3 * (b^3 * x^5 + 3 * a * b^2 * x^4 + (3 * a^2 - 1) * b * x^3 + (a^3 - a) * x^2) * \log(b*x + a)^2 + (b^3 * x^5 + 2 * a * b^2 * x^4 + (a^2 - 1) * b * x^3 + 3 * (b^3 * x^5 + 3 * a * b^2 * x^4 + (3 * a^2 - 1) * b * x^3 + (a^3 - a) * x^2) * \log(b*x + a)) * \log(b^2 * x^2 + 2 * a * b * x + a^2)) / (b^3 * x^3 + 3 * a * b^2 * x^2 + a^3 + (3 * a^2 - 1) * b * x - a), x) \end{aligned}$$

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^2*arccsc(b*x+a)^2,x, algorithm="fricas")`[Out] `integral(x^2*arccsc(b*x + a)^2, x)`**Sympy [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int x^2 \operatorname{acsc}^2(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x**2*acsc(b*x+a)**2,x)`[Out] `Integral(x**2*acsc(a + b*x)**2, x)`**Giac [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^2*arccsc(b*x+a)^2,x, algorithm="giac")`[Out] `integrate(x^2*arccsc(b*x + a)^2, x)`**Mupad [F]**

time = 0.00, size = -1, normalized size = -0.00

$$\int x^2 \sin\left(\frac{1}{a + bx}\right)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x^2*asin(1/(a + b*x))^2,x)`[Out] `int(x^2*asin(1/(a + b*x))^2, x)`

$$3.29 \quad \int x \csc^{-1}(a + bx)^2 dx$$

Optimal. Leaf size=145

$$\frac{(a + bx) \sqrt{1 - \frac{1}{(a + bx)^2}} \csc^{-1}(a + bx)}{b^2} - \frac{a^2 \csc^{-1}(a + bx)^2}{2b^2} + \frac{1}{2} x^2 \csc^{-1}(a + bx)^2 - \frac{4a \csc^{-1}(a + bx) \tanh^{-1}\left(\frac{e^{i \csc^{-1}(a + bx)}}{b}\right)}{b^2}$$

[Out] $-1/2*a^2*arccsc(b*x+a)^2/b^2+1/2*x^2*arccsc(b*x+a)^2-4*a*arccsc(b*x+a)*\text{arctanh}\left(I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2)\right)/b^2+\ln(b*x+a)/b^2+2*I*a*\text{polylog}(2,-I/(b*x+a)-(1-1/(b*x+a)^2)^(1/2))/b^2-2*I*a*\text{polylog}(2,I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))/b^2+(b*x+a)*arccsc(b*x+a)*(1-1/(b*x+a)^2)^(1/2)/b^2$

Rubi [A]

time = 0.09, antiderivative size = 145, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 8, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.800$, Rules used = {5367, 4512, 4275, 4268, 2317, 2438, 4269, 3556}

$$-\frac{a^2 \csc^{-1}(a + bx)^2}{2b^2} + \frac{2ia \text{Li}_2\left(-e^{i \csc^{-1}(a + bx)}\right)}{b^2} - \frac{2ia \text{Li}_2\left(e^{i \csc^{-1}(a + bx)}\right)}{b^2} + \frac{\log(a + bx)}{b^2} + \frac{(a + bx) \sqrt{1 - \frac{1}{(a + bx)^2}} \csc^{-1}(a + bx)}{b^2} - \frac{4a \csc^{-1}(a + bx) \tanh^{-1}\left(e^{i \csc^{-1}(a + bx)}\right)}{b^2} + \frac{1}{2} x^2 \csc^{-1}(a + bx)^2$$

Antiderivative was successfully verified.

[In] Int[x*ArcCsc[a + b*x]^2, x]

[Out] $((a + b*x)*\text{Sqrt}[1 - (a + b*x)^{-2}]*\text{ArcCsc}[a + b*x])/b^2 - (a^2*\text{ArcCsc}[a + b*x]^2)/(2*b^2) + (x^2*\text{ArcCsc}[a + b*x]^2)/2 - (4*a*\text{ArcCsc}[a + b*x]*\text{ArcTanh}[E^{(I*\text{ArcCsc}[a + b*x])}])/b^2 + \text{Log}[a + b*x]/b^2 + ((2*I)*a*\text{PolyLog}[2, -E^{(I*\text{ArcCsc}[a + b*x])}])/b^2 - ((2*I)*a*\text{PolyLog}[2, E^{(I*\text{ArcCsc}[a + b*x])}])/b^2$

Rule 2317

```
Int[Log[(a_) + (b_)*((F_)^((e_.)*(c_.) + (d_.)*(x_))))^(n_.)], x_Symbol]
:> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2438

```
Int[Log[(c_.)*(d_) + (e_.)*(x_)^(n_.)]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 3556

```
Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

Rule 4268

```
Int[csc[(e_.) + (f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] :> Simp[-2*(c + d*x)^m*(ArcTanh[E^(I*(e + f*x))/f], x] + (-Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Log[1 - E^(I*(e + f*x))], x], x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Log[1 + E^(I*(e + f*x))], x], x]) /; FreeQ[{c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 4269

```
Int[csc[(e_.) + (f_.)*(x_)]^2*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] :> Simp[((-c + d*x)^m)*(Cot[e + f*x]/f), x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Cot[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

Rule 4275

```
Int[(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^n*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (a + b*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rule 4512

```
Int[Cot[(c_.) + (d_.)*(x_)]*Csc[(c_.) + (d_.)*(x_)]*(Csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_.))^n*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] :> Simp[((-e + f*x)^m)*((a + b*Csc[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[f*(m/(b*d*(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Csc[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]
```

Rule 5367

```
Int[((a_.) + ArcCsc[(c_.) + (d_.)*(x_)]*(b_.))^p*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] :> Dist[-(d^(m + 1))^{(-1)}, Subst[Int[(a + b*x)^p*Csc[x]*Cot[x]*(d*e - c*f + f*Csc[x])^m, x], x, ArcCsc[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Rubi steps

$$\begin{aligned}
\int x \csc^{-1}(a+bx)^2 dx &= -\frac{\text{Subst}(\int x^2 \cot(x) \csc(x)(-a+\csc(x)) dx, x, \csc^{-1}(a+bx))}{b^2} \\
&= \frac{1}{2}x^2 \csc^{-1}(a+bx)^2 - \frac{\text{Subst}(\int x(-a+\csc(x))^2 dx, x, \csc^{-1}(a+bx))}{b^2} \\
&= \frac{1}{2}x^2 \csc^{-1}(a+bx)^2 - \frac{\text{Subst}(\int (a^2x - 2ax \csc(x) + x \csc^2(x)) dx, x, \csc^{-1}(a+bx))}{b^2} \\
&= -\frac{a^2 \csc^{-1}(a+bx)^2}{2b^2} + \frac{1}{2}x^2 \csc^{-1}(a+bx)^2 - \frac{\text{Subst}(\int x \csc^2(x) dx, x, \csc^{-1}(a+bx))}{b^2} \\
&= \frac{(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}} \csc^{-1}(a+bx)}{b^2} - \frac{a^2 \csc^{-1}(a+bx)^2}{2b^2} + \frac{1}{2}x^2 \csc^{-1}(a+bx)^2 \\
&= \frac{(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}} \csc^{-1}(a+bx)}{b^2} - \frac{a^2 \csc^{-1}(a+bx)^2}{2b^2} + \frac{1}{2}x^2 \csc^{-1}(a+bx)^2 \\
&= \frac{(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}} \csc^{-1}(a+bx)}{b^2} - \frac{a^2 \csc^{-1}(a+bx)^2}{2b^2} + \frac{1}{2}x^2 \csc^{-1}(a+bx)^2
\end{aligned}$$

Mathematica [A]

time = 0.62, size = 213, normalized size = 1.47

$$\frac{2a\sqrt{\frac{-1+a^2+2abx+b^2x^2}{(a+bx)^2}}\csc^{-1}(a+bx)+2bx\sqrt{\frac{-1+a^2+2abx+b^2x^2}{(a+bx)^2}}\csc^{-1}(a+bx)-a^2\csc^{-1}(a+bx)^2+b^2x^2\csc^{-1}(a+bx)^2+4a\csc^{-1}(a+bx)\log\left(1-e^{i\csc^{-1}(a+bx)}\right)-4a\csc^{-1}(a+bx)\log\left(1+e^{i\csc^{-1}(a+bx)}\right)-2\log\left(\frac{1}{a+bx}\right)+4ia\text{PolyLog}\left(2,-e^{i\csc^{-1}(a+bx)}\right)-4ia\text{PolyLog}\left(2,e^{i\csc^{-1}(a+bx)}\right)}{2b^2}$$

Antiderivative was successfully verified.

[In] `Integrate[x*ArcCsc[a + b*x]^2,x]`

[Out] $(2*a*\text{Sqrt}[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2]*\text{ArcCsc}[a + b*x] + 2*b*x*\text{Sqrt}[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2]*\text{ArcCsc}[a + b*x] - a^2*ArcCsc[a + b*x]^2 + b^2*x^2*ArcCsc[a + b*x]^2 + 4*a*ArcCsc[a + b*x]*\text{Log}[1 - E^(I*ArcCsc[a + b*x])] - 4*a*ArcCsc[a + b*x]*\text{Log}[1 + E^(I*ArcCsc[a + b*x])] - 2*\text{Log}[(a + b*x)^{(-1)}] + (4*I)*a*\text{PolyLog}[2, -E^(I*ArcCsc[a + b*x])] - (4*I)*a*\text{PolyLog}[2, E^(I*ArcCsc[a + b*x])])/(2*b^2)$

Maple [A]

time = 1.05, size = 214, normalized size = 1.48

method	result
derivative divides	$-a\text{arccsc}(bx+a)^2(bx+a)+2a\text{arccsc}(bx+a)\ln\left(1-\frac{i}{bx+a}-\sqrt{1-\frac{1}{(bx+a)^2}}\right)-2a\text{arccsc}(bx+a)\ln\left(1+\frac{i}{bx+a}+\sqrt{1-\frac{1}{(bx+a)^2}}\right)$

default	$\frac{-a \operatorname{arccsc}(bx+a)^2(bx+a) + 2a \operatorname{arccsc}(bx+a) \ln\left(1 - \frac{i}{bx+a} - \sqrt{1 - \frac{1}{(bx+a)^2}}\right) - 2a \operatorname{arccsc}(bx+a) \ln\left(1 + \frac{i}{bx+a} + \sqrt{1 - \frac{1}{(bx+a)^2}}\right)}{...}$
---------	--

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x*arccsc(b*x+a)^2,x,method=_RETURNVERBOSE)`

[Out]
$$\frac{1}{b^2}(-a \operatorname{arccsc}(bx+a)^2(bx+a) + 2a \operatorname{arccsc}(bx+a) \ln(1 - I/(bx+a) - (1 - 1/(bx+a)^2)^{(1/2)}) - 2a \operatorname{arccsc}(bx+a) \ln(1 + I/(bx+a) + (1 - 1/(bx+a)^2)^{(1/2)}) + 2I \operatorname{adilog}(1 + I/(bx+a) + (1 - 1/(bx+a)^2)^{(1/2)}) - 2I \operatorname{ailog}(1 - I/(bx+a) - (1 - 1/(bx+a)^2)^{(1/2)}) + 1/2 \operatorname{arccsc}(bx+a)^2 + \operatorname{arccsc}(bx+a) (((bx+a)^2 - 1)/(bx+a)^2)^{(1/2)} * (bx+a) - \ln(1/(bx+a)))$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x*arccsc(b*x+a)^2,x, algorithm="maxima")`

[Out]
$$\frac{1}{2}x^2 \operatorname{arctan2}(1, \sqrt{bx + a + 1}) \sqrt{bx + a - 1})^2 - \frac{1}{8}x^2 \log(b^2 x^2 + 2a b x + a^2)^2 + \operatorname{integrate}\left(\frac{1}{2} \cdot 2 \sqrt{bx + a + 1} \sqrt{bx + a - 1} \cdot bx^2 \operatorname{arctan2}(1, \sqrt{bx + a + 1}) \sqrt{bx + a - 1}) - 2(b^3 x^4 + 3a b^2 x^3 + (3a^2 - 1) b x^2 + (a^3 - a) x) \log(bx + a)^2 + (b^3 x^4 + 2a b^2 x^3 + (a^2 - 1) b x^2 + 2(b^3 x^4 + 3a b^2 x^3 + (3a^2 - 1) b x^2 + (a^3 - a) x) \log(bx + a)) \log(b^2 x^2 + 2a b x + a^2)\right) / (b^3 x^3 + 3a b^2 x^2 + a^3 + (3a^2 - 1) b x - a), x)$$

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x*arccsc(b*x+a)^2,x, algorithm="fricas")`

[Out] `integral(x*arccsc(b*x + a)^2, x)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x \operatorname{acsc}^2(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x*acsc(b*x+a)**2,x)`
[Out] `Integral(x*acsc(a + b*x)**2, x)`

Giac [F]
time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x*arccsc(b*x+a)^2,x, algorithm="giac")`
[Out] `integrate(x*arccsc(b*x + a)^2, x)`

Mupad [F]
time = 0.00, size = -1, normalized size = -0.01

$$\int x \arcsin\left(\frac{1}{a + bx}\right)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x*asin(1/(a + b*x))^2,x)`
[Out] `int(x*asin(1/(a + b*x))^2, x)`

$$\text{3.30} \quad \int \csc^{-1}(a + bx)^2 dx$$

Optimal. Leaf size=86

$$\frac{(a + bx) \csc^{-1}(a + bx)^2}{b} + \frac{4 \csc^{-1}(a + bx) \tanh^{-1}\left(e^{i \csc^{-1}(a + bx)}\right)}{b} - \frac{2 i \text{PolyLog}\left(2, -e^{i \csc^{-1}(a + bx)}\right)}{b} + \frac{2 i \text{PolyLog}\left(2, e^{i \csc^{-1}(a + bx)}\right)}{b}$$

[Out] $(b*x+a)*\text{arccsc}(b*x+a)^2/b + 4*\text{arccsc}(b*x+a)*\text{arctanh}(I/(b*x+a)+(1-1/(b*x+a)^2)^{(1/2)})/b - 2*I*\text{polylog}(2, -I/(b*x+a)-(1-1/(b*x+a)^2)^{(1/2)})/b + 2*I*\text{polylog}(2, I/(b*x+a)+(1-1/(b*x+a)^2)^{(1/2)})/b$

Rubi [A]

time = 0.04, antiderivative size = 86, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 6, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.750, Rules used = {5361, 5325, 3843, 4268, 2317, 2438}

$$-\frac{2 i \text{Li}_2\left(-e^{i \csc^{-1}(a + bx)}\right)}{b} + \frac{2 i \text{Li}_2\left(e^{i \csc^{-1}(a + bx)}\right)}{b} + \frac{(a + bx) \csc^{-1}(a + bx)^2}{b} + \frac{4 \csc^{-1}(a + bx) \tanh^{-1}\left(e^{i \csc^{-1}(a + bx)}\right)}{b}$$

Antiderivative was successfully verified.

[In] Int[ArcCsc[a + b*x]^2, x]

[Out] $((a + b*x)*\text{ArcCsc}[a + b*x]^2)/b + (4*\text{ArcCsc}[a + b*x]*\text{ArcTanh}[E^(\text{I}*\text{ArcCsc}[a + b*x])])/b - ((2*\text{I})*\text{PolyLog}[2, -E^(\text{I}*\text{ArcCsc}[a + b*x])])/b + ((2*\text{I})*\text{PolyLog}[2, E^(\text{I}*\text{ArcCsc}[a + b*x])])/b$

Rule 2317

```
Int[Log[(a_) + (b_.)*((F_)^((e_.)*(c_.) + (d_.)*(x_))))^(n_.)], x_Symbol]
: > Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2438

```
Int[Log[(c_.)*(d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] : > Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 3843

```
Int[Cot[(a_.) + (b_.)*(x_)^(n_.)]^(q_.)*Csc[(a_.) + (b_.)*(x_)^(n_.)]^(p_.)*(x_)^(m_), x_Symbol] : > Simp[(-x^(m - n + 1))*(Csc[a + b*x^n]^p/(b*n*p)), x] + Dist[(m - n + 1)/(b*n*p), Int[x^(m - n)*Csc[a + b*x^n]^p, x], x] /; FreeQ[{a, b, p}, x] && IntegerQ[n] && GeQ[m, n] && EqQ[q, 1]
```

Rule 4268

```
Int[csc[(e_.) + (f_ .)*(x_)]*((c_.) + (d_ .)*(x_))^(m_.), x_Symbol] :> Simp[-2*(c + d*x)^m*(ArcTanh[E^(I*(e + f*x))/f], x] + (-Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Log[1 - E^(I*(e + f*x))], x], x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Log[1 + E^(I*(e + f*x))], x], x]) /; FreeQ[{c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 5325

```
Int[((a_.) + ArcCsc[(c_.)*(x_)]*(b_ .))^n, x_Symbol] :> Dist[-c^(-1), Subst[Int[(a + b*x)^n*Csc[x]*Cot[x], x], x, ArcCsc[c*x]], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[n, 0]
```

Rule 5361

```
Int[((a_.) + ArcCsc[(c_.) + (d_ .)*(x_)]*(b_ .))^p, x_Symbol] :> Dist[1/d, Subst[Int[(a + b*ArcCsc[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[p, 0]
```

Rubi steps

$$\begin{aligned} \int \csc^{-1}(a + bx)^2 dx &= \frac{\text{Subst}\left(\int \csc^{-1}(x)^2 dx, x, a + bx\right)}{b} \\ &= -\frac{\text{Subst}\left(\int x^2 \cot(x) \csc(x) dx, x, \csc^{-1}(a + bx)\right)}{b} \\ &= \frac{(a + bx) \csc^{-1}(a + bx)^2}{b} - \frac{2\text{Subst}\left(\int x \csc(x) dx, x, \csc^{-1}(a + bx)\right)}{b} \\ &= \frac{(a + bx) \csc^{-1}(a + bx)^2}{b} + \frac{4 \csc^{-1}(a + bx) \tanh^{-1}\left(e^{i \csc^{-1}(a+bx)}\right)}{b} + \frac{2\text{Subst}\left(\int \log(1\right.}{b} \\ &= \frac{(a + bx) \csc^{-1}(a + bx)^2}{b} + \frac{4 \csc^{-1}(a + bx) \tanh^{-1}\left(e^{i \csc^{-1}(a+bx)}\right)}{b} - \frac{(2i)\text{Subst}\left(\int \log(1\right.}{b} \\ &= \frac{(a + bx) \csc^{-1}(a + bx)^2}{b} + \frac{4 \csc^{-1}(a + bx) \tanh^{-1}\left(e^{i \csc^{-1}(a+bx)}\right)}{b} - \frac{2i \text{Li}_2\left(-e^{i \csc^{-1}(a+bx)}\right)}{b} \end{aligned}$$

Mathematica [A]

time = 0.10, size = 99, normalized size = 1.15

$$\frac{\csc^{-1}(a + bx) \left((a + bx) \csc^{-1}(a + bx) - 2 \log \left(1 - e^{i \csc^{-1}(a+bx)}\right) + 2 \log \left(1 + e^{i \csc^{-1}(a+bx)}\right)\right) - 2i \text{PolyLog}\left(2, -e^{i \csc^{-1}(a+bx)}\right) + 2i \text{PolyLog}\left(2, e^{i \csc^{-1}(a+bx)}\right)}{b}$$

Antiderivative was successfully verified.

[In] `Integrate[ArcCsc[a + b*x]^2, x]`

[Out] $(\text{ArcCsc}[a + b*x]*((a + b*x)*\text{ArcCsc}[a + b*x] - 2*\text{Log}[1 - E^{\text{I}}*\text{ArcCsc}[a + b*x]]) + 2*\text{Log}[1 + E^{\text{I}}*\text{ArcCsc}[a + b*x]]) - (2*\text{I})*\text{PolyLog}[2, -E^{\text{I}}*\text{ArcCsc}[a + b*x]] + (2*\text{I})*\text{PolyLog}[2, E^{\text{I}}*\text{ArcCsc}[a + b*x]])/b$

Maple [A]

time = 0.56, size = 150, normalized size = 1.74

method	result
derivativedivides	$\frac{\text{arccsc}(bx+a)^2(bx+a) - 2 \text{arccsc}(bx+a) \ln\left(1 - \frac{i}{bx+a} - \sqrt{1 - \frac{1}{(bx+a)^2}}\right) + 2 \text{arccsc}(bx+a) \ln\left(1 + \frac{i}{bx+a} + \sqrt{1 - \frac{1}{(bx+a)^2}}\right)}{b}$
default	$\frac{\text{arccsc}(bx+a)^2(bx+a) - 2 \text{arccsc}(bx+a) \ln\left(1 - \frac{i}{bx+a} - \sqrt{1 - \frac{1}{(bx+a)^2}}\right) + 2 \text{arccsc}(bx+a) \ln\left(1 + \frac{i}{bx+a} + \sqrt{1 - \frac{1}{(bx+a)^2}}\right)}{b}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(arccsc(b*x+a)^2,x,method=_RETURNVERBOSE)`

[Out] $1/b*(\text{arccsc}(b*x+a)^2*(b*x+a) - 2*\text{arccsc}(b*x+a)*\ln(1-\text{I}/(b*x+a)-(1-1/(b*x+a)^2)^{(1/2)}) + 2*\text{arccsc}(b*x+a)*\ln(1+\text{I}/(b*x+a)+(1-1/(b*x+a)^2)^{(1/2)}) - 2*\text{I}*\text{dilog}(1+\text{I}/(b*x+a)+(1-1/(b*x+a)^2)^{(1/2)}) + 2*\text{I}*\text{dilog}(1-\text{I}/(b*x+a)-(1-1/(b*x+a)^2)^{(1/2)}))$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(b*x+a)^2,x, algorithm="maxima")`

[Out] $x*\text{arctan2}(1, \sqrt{b*x + a + 1})*\sqrt{b*x + a - 1})^2 - 1/4*x*\log(b^2*x^2 + 2*a*b*x + a^2)^2 + \text{integrate}((2*\sqrt{b*x + a + 1})*\sqrt{b*x + a - 1})*b*x*\text{arctan2}(1, \sqrt{b*x + a + 1})*\sqrt{b*x + a - 1}) - (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2 - 1)*b*x - a)*\log(b*x + a)^2 + (b^3*x^3 + 2*a*b^2*x^2 + (a^2 - 1)*b*x + (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2 - 1)*b*x - a)*\log(b*x + a))*\log(b^2*x^2 + 2*a*b*x + a^2))/(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2 - 1)*b*x - a), x)$

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(b*x+a)^2,x, algorithm="fricas")`

[Out] integral(arccsc(b*x + a)^2, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \operatorname{acsc}^2(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(acsc(b*x+a)**2,x)

[Out] Integral(acsc(a + b*x)**2, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(arccsc(b*x+a)^2,x, algorithm="giac")

[Out] integrate(arccsc(b*x + a)^2, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \operatorname{asin}\left(\frac{1}{a + b x}\right)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(asin(1/(a + b*x))^2,x)

[Out] int(asin(1/(a + b*x))^2, x)

3.31 $\int \frac{\csc^{-1}(a+bx)^2}{x} dx$

Optimal. Leaf size=324

$$\csc^{-1}(a+bx)^2 \log \left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{1 - \sqrt{1 - a^2}} \right) + \csc^{-1}(a+bx)^2 \log \left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{1 + \sqrt{1 - a^2}} \right) - \csc^{-1}(a+bx)^2 \log \left(1 - e^{2i \csc^{-1}(a+bx)} \right)$$

[Out] $-\text{arccsc}(b*x+a)^2 \ln(1-(I/(b*x+a)+(1-1/(b*x+a)^2)^{(1/2)})^2)+\text{arccsc}(b*x+a)^2 \ln(1+I*a*(I/(b*x+a)+(1-1/(b*x+a)^2)^{(1/2)})/(1-(-a^2+1)^{(1/2)}))+\text{arccsc}(b*x+a)^2 \ln(1+I*a*(I/(b*x+a)+(1-1/(b*x+a)^2)^{(1/2)})/(1+(-a^2+1)^{(1/2)}))+I*\text{arccsc}(b*x+a)*\text{polylog}(2,(I/(b*x+a)+(1-1/(b*x+a)^2)^{(1/2)})^2)-2*I*\text{arccsc}(b*x+a)*\text{polylog}(2,-I*a*(I/(b*x+a)+(1-1/(b*x+a)^2)^{(1/2)})/(1-(-a^2+1)^{(1/2)}))-2*I*\text{arccsc}(b*x+a)*\text{polylog}(2,-I*a*(I/(b*x+a)+(1-1/(b*x+a)^2)^{(1/2)})/(1+(-a^2+1)^{(1/2)}))-1/2*\text{polylog}(3,(I/(b*x+a)+(1-1/(b*x+a)^2)^{(1/2)})^2)+2*\text{polylog}(3,-I*a*(I/(b*x+a)+(1-1/(b*x+a)^2)^{(1/2)})/(1-(-a^2+1)^{(1/2)}))+2*\text{polylog}(3,-I*a*(I/(b*x+a)+(1-1/(b*x+a)^2)^{(1/2)})/(1+(-a^2+1)^{(1/2)}))$

Rubi [A]

time = 0.34, antiderivative size = 324, normalized size of antiderivative = 1.00, number of steps used = 17, number of rules used = 9, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {5367, 4648, 4625, 3798, 2221, 2611, 2320, 6724, 4615}

$$-2i \csc^{-1}(a+bx) \text{Li}_2\left(\frac{i a e^{i \csc^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right) - 2i \csc^{-1}(a+bx) \text{Li}_2\left(\frac{i a e^{i \csc^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right) + 2 \text{Li}_3\left(\frac{i a e^{i \csc^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right) + 2 \text{Li}_3\left(\frac{i a e^{i \csc^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right) + \csc^{-1}(a+bx)^2 \log\left(1+\frac{i a e^{i \csc^{-1}(a+bx)}}{\sqrt{1-a^2}}\right) + \csc^{-1}(a+bx)^2 \log\left(1+\frac{i a e^{i \csc^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right) + i \csc^{-1}(a+bx) \text{Li}_2\left(e^{2i \csc^{-1}(a+bx)}\right) - \frac{1}{2} \text{Li}_2\left(e^{2i \csc^{-1}(a+bx)}\right) - \csc^{-1}(a+bx)^2 \log\left(1-e^{2i \csc^{-1}(a+bx)}\right)$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{ArcCsc}[a+b*x]^2/x, x]$

[Out] $\text{ArcCsc}[a+b*x]^2 \text{Log}[1+(I*a*E^(I*ArcCsc[a+b*x]))/(1-\text{Sqrt}[1-a^2])]+\text{ArcCsc}[a+b*x]^2 \text{Log}[1+(I*a*E^(I*ArcCsc[a+b*x]))/(1+\text{Sqrt}[1-a^2])]-\text{ArcCsc}[a+b*x]^2 \text{Log}[1-E^((2*I)*\text{ArcCsc}[a+b*x])]-(2*I)*\text{ArcCsc}[a+b*x]*\text{PolyLog}[2, ((-I)*a*E^(I*ArcCsc[a+b*x]))/(1-\text{Sqrt}[1-a^2])]-(2*I)*\text{ArcCsc}[a+b*x]*\text{PolyLog}[2, ((-I)*a*E^(I*ArcCsc[a+b*x]))/(1+\text{Sqrt}[1-a^2])]+\text{I}*\text{ArcCsc}[a+b*x]*\text{PolyLog}[2, E^((2*I)*\text{ArcCsc}[a+b*x])]+2*\text{PolyLog}[3, ((-I)*a*E^(I*ArcCsc[a+b*x]))/(1-\text{Sqrt}[1-a^2])]+2*\text{PolyLog}[3, ((-I)*a*E^(I*ArcCsc[a+b*x]))/(1+\text{Sqrt}[1-a^2])]-\text{PolyLog}[3, E^((2*I)*\text{ArcCsc}[a+b*x])]/2$

Rule 2221

```
Int[((F_)^((g_.)*(e_.)+(f_)*(x_)))^(n_.)*((c_.)+(d_)*(x_))^(m_.))/((a_.)+(b_)*(F_)^((g_.)*(e_.)+(f_)*(x_)))^(n_.)), x_Symbol] :> Simplify[((c+d*x)^m/(b*f*g*n*Log[F]))*Log[1+b*((F^(g*(e+f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c+d*x)^(m-1)*Log[1+b*((F^(g*(e+f*x)))^n/a)], x], x]; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2320

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*(a_.)*(v_)^(n_.)^(m_.)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*(a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2611

```
Int[Log[1 + (e_)*(F_)^((c_)*(a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] :> Simp[((-f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]
```

Rule 3798

```
Int[((c_.) + (d_.)*(x_))^m*tan[(e_.) + Pi*(k_.) + (f_.)*(x_)], x_Symbol] :> Simp[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Dist[2*I, Int[(c + d*x)^m *E^(2*I*k*Pi)*(E^(2*I*(e + f*x))/(1 + E^(2*I*k*Pi)*E^(2*I*(e + f*x)))), x], x] /; FreeQ[{c, d, e, f}, x] && IntegerQ[4*k] && IGtQ[m, 0]
```

Rule 4615

```
Int[(Cos[(c_.) + (d_.)*(x_)]*((e_.) + (f_.)*(x_))^(m_.))/((a_) + (b_.)*Sin[(c_.) + (d_.)*(x_)]), x_Symbol] :> Simp[(-I)*((e + f*x)^(m + 1)/(b*f*(m + 1))), x] + (Int[(e + f*x)^m*(E^(I*(c + d*x))/(a - Rt[a^2 - b^2, 2] - I*b*E^(I*(c + d*x)))), x] + Int[(e + f*x)^m*(E^(I*(c + d*x))/(a + Rt[a^2 - b^2, 2] - I*b*E^(I*(c + d*x)))), x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && PosQ[a^2 - b^2]
```

Rule 4625

```
Int[(Cot[(c_.) + (d_.)*(x_)]^n*((e_.) + (f_.)*(x_))^(m_.))/((a_) + (b_.)*Sin[(c_.) + (d_.)*(x_)]), x_Symbol] :> Dist[1/a, Int[(e + f*x)^m*Cot[c + d*x]^n, x], x] - Dist[b/a, Int[(e + f*x)^m*Cos[c + d*x]*(Cot[c + d*x]^(n - 1)/(a + b*Sin[c + d*x])), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rule 4648

```
Int[((e_.) + (f_.)*(x_))^(m_.)*(F_)[(c_.) + (d_.)*(x_)]^(n_.)*(G_)[(c_.) + (d_.)*(x_)]^(p_.)]/(Csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_)), x_Symbol] :> Int[((e + f*x)^m*Sin[c + d*x]*F[c + d*x]^n*(G[c + d*x]^p/(b + a*Sin[c + d*x])), x] /; FreeQ[{a, b, c, d, e, f}, x] && TrigQ[F] && TrigQ[G] && IntegersQ[m]
```

, n, p]

Rule 5367

```
Int[((a_) + ArcCsc[(c_) + (d_)*(x_)]*(b_.))^(p_)*(e_.) + (f_)*(x_))^(m_),
x_Symbol] :> Dist[-(d^(m + 1))^{(-1)}, Subst[Int[(a + b*x)^p*Csc[x]*Cot[x]*(d*e - c*f + f*Csc[x])^m, x], x, ArcCsc[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Rule 6724

```
Int[PolyLog[n_, (c_)*(a_.) + (b_)*(x_))^(p_.)]/((d_.) + (e_)*(x_)), x_Symbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

Rubi steps

$$\begin{aligned} \int \frac{\csc^{-1}(a+bx)^2}{x} dx &= -\text{Subst}\left(\int \frac{x^2 \cot(x) \csc(x)}{-a+\csc(x)} dx, x, \csc^{-1}(a+bx)\right) \\ &= -\text{Subst}\left(\int \frac{x^2 \cot(x)}{1-a \sin(x)} dx, x, \csc^{-1}(a+bx)\right) \\ &= -\left(a \text{Subst}\left(\int \frac{x^2 \cos(x)}{1-a \sin(x)} dx, x, \csc^{-1}(a+bx)\right)\right) - \text{Subst}\left(\int x^2 \cot(x) dx, x, \csc^{-1}(a+bx)\right) \\ &= 2i \text{Subst}\left(\int \frac{e^{2ix} x^2}{1-e^{2ix}} dx, x, \csc^{-1}(a+bx)\right) - a \text{Subst}\left(\int \frac{e^{ix} x^2}{1-\sqrt{1-a^2}+iae^{ix}} dx, x, \csc^{-1}(a+bx)\right) \\ &= \csc^{-1}(a+bx)^2 \log\left(1+\frac{iae^{i \csc^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right) + \csc^{-1}(a+bx)^2 \log\left(1+\frac{iae^{i \csc^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right) \\ &= \csc^{-1}(a+bx)^2 \log\left(1+\frac{iae^{i \csc^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right) + \csc^{-1}(a+bx)^2 \log\left(1+\frac{iae^{i \csc^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right) \\ &= \csc^{-1}(a+bx)^2 \log\left(1+\frac{iae^{i \csc^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right) + \csc^{-1}(a+bx)^2 \log\left(1+\frac{iae^{i \csc^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right) \\ &= \csc^{-1}(a+bx)^2 \log\left(1+\frac{iae^{i \csc^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right) + \csc^{-1}(a+bx)^2 \log\left(1+\frac{iae^{i \csc^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right) \\ &= \csc^{-1}(a+bx)^2 \log\left(1+\frac{iae^{i \csc^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right) + \csc^{-1}(a+bx)^2 \log\left(1+\frac{iae^{i \csc^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right) \end{aligned}$$

Mathematica [A]

time = 0.24, size = 408, normalized size = 1.26

$$\frac{1}{2} \csc^{-1}(a+bx)^2 - \csc^{-1}(a+bx)^2 \log\left(1-e^{-i \csc^{-1}(a+bx)}\right) - \csc^{-1}(a+bx)^2 \log\left(1+e^{i \csc^{-1}(a+bx)}\right) + \csc^{-1}(a+bx)^2 \log\left(1-\frac{i a e^{i \csc^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right) + \csc^{-1}(a+bx)^2 \log\left(1+\frac{i a e^{i \csc^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right) - 2 i \csc^{-1}(a+bx) \text{PolyLog}\left[\frac{1}{2}, e^{-i \csc^{-1}(a+bx)}\right] + 2 i \csc^{-1}(a+bx) \text{PolyLog}\left[\frac{1}{2}, e^{i \csc^{-1}(a+bx)}\right] - 2 i \csc^{-1}(a+bx) \text{PolyLog}\left[\frac{1}{2}, \frac{i a e^{i \csc^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right] - 2 i \csc^{-1}(a+bx) \text{PolyLog}\left[\frac{1}{2}, -\frac{i a e^{i \csc^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right] - 2 i \csc^{-1}(a+bx) \text{PolyLog}\left[\frac{1}{2}, -e^{-i \csc^{-1}(a+bx)}\right] + 2 i \csc^{-1}(a+bx) \text{PolyLog}\left[\frac{1}{2}, e^{-i \csc^{-1}(a+bx)}\right] + 2 i \csc^{-1}(a+bx) \text{PolyLog}\left[\frac{1}{2}, -e^{i \csc^{-1}(a+bx)}\right] - 2 i \csc^{-1}(a+bx) \text{PolyLog}\left[\frac{1}{2}, -\frac{i a e^{i \csc^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right] + 2 i \csc^{-1}(a+bx) \text{PolyLog}\left[\frac{1}{2}, \frac{i a e^{i \csc^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right] + 2 i \csc^{-1}(a+bx) \text{PolyLog}\left[\frac{1}{2}, \frac{i a e^{i \csc^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right]$$

Antiderivative was successfully verified.

```
[In] Integrate[ArcCsc[a + b*x]^2/x,x]
[Out] (I/6)*Pi^3 - (I/3)*ArcCsc[a + b*x]^3 - ArcCsc[a + b*x]^2*Log[1 - E^((-I)*ArcCsc[a + b*x])] - ArcCsc[a + b*x]^2*Log[1 + E^(I*ArcCsc[a + b*x])] + ArcCsc[a + b*x]^2*Log[1 - (I*a*E^(I*ArcCsc[a + b*x]))/(-1 + Sqrt[1 - a^2])] + ArcCsc[a + b*x]^2*Log[1 + (I*a*E^(I*ArcCsc[a + b*x]))/(1 + Sqrt[1 - a^2])] - (2*I)*ArcCsc[a + b*x]*PolyLog[2, E^((-I)*ArcCsc[a + b*x])] + (2*I)*ArcCsc[a + b*x]*PolyLog[2, -E^(I*ArcCsc[a + b*x])] - (2*I)*ArcCsc[a + b*x]*PolyLog[2, (I*a*E^(I*ArcCsc[a + b*x]))/(-1 + Sqrt[1 - a^2])] - (2*I)*ArcCsc[a + b*x]*PolyLog[2, ((-I)*a*E^(I*ArcCsc[a + b*x]))/(1 + Sqrt[1 - a^2])] - 2*PolyLog[3, E^((-I)*ArcCsc[a + b*x])] - 2*PolyLog[3, -E^(I*ArcCsc[a + b*x])] + 2*PolyLog[3, (I*a*E^(I*ArcCsc[a + b*x]))/(-1 + Sqrt[1 - a^2])] + 2*PolyLog[3, ((-I)*a*E^(I*ArcCsc[a + b*x]))/(1 + Sqrt[1 - a^2])]
```

Maple [F]

time = 2.38, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{arccsc}(bx+a)^2}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(arccsc(b*x+a)^2/x,x)
```

```
[Out] int(arccsc(b*x+a)^2/x,x)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(arccsc(b*x+a)^2/x,x, algorithm="maxima")
```

```
[Out] integrate(arccsc(b*x + a)^2/x, x)
```

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(arccsc(b*x+a)^2/x,x, algorithm="fricas")
```

```
[Out] integral(arccsc(b*x + a)^2/x, x)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{acsc}^2(a + bx)}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(acsc(b*x+a)**2/x,x)`

[Out] `Integral(acsc(a + b*x)**2/x, x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(b*x+a)^2/x,x, algorithm="giac")`

[Out] `integrate(arccsc(b*x + a)^2/x, x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\arcsin\left(\frac{1}{a+bx}\right)^2}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(asin(1/(a + b*x))^2/x,x)`

[Out] `int(asin(1/(a + b*x))^2/x, x)`

$$3.32 \quad \int \frac{\csc^{-1}(a+bx)^2}{x^2} dx$$

Optimal. Leaf size=254

$$\frac{b \csc^{-1}(a+bx)^2}{a} - \frac{\csc^{-1}(a+bx)^2}{x} - \frac{2ib \csc^{-1}(a+bx) \log\left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} + \frac{2ib \csc^{-1}(a+bx) \log\left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{\sqrt{1-a^2+1}}\right)}{a\sqrt{1-a^2}}$$

[Out] $-b*\text{arccsc}(b*x+a)^2/a - \text{arccsc}(b*x+a)^2/x - 2*I*b*\text{arccsc}(b*x+a)*\ln(1+I*a*(I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))/(1-(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2) + 2*I*b*\text{arccsc}(b*x+a)*\ln(1+I*a*(I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))/(1-(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2) - 2*b*\text{polylog}(2, -I*a*(I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))/(1-(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2) + 2*b*\text{polylog}(2, -I*a*(I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))/(1-(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)$

Rubi [A]

time = 0.28, antiderivative size = 254, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 8, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {5367, 4512, 4276, 3404, 2296, 2221, 2317, 2438}

$$-\frac{2b\text{Li}_2\left(\frac{-iae^{i \csc^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} + \frac{2b\text{Li}_2\left(\frac{-iae^{i \csc^{-1}(a+bx)}}{\sqrt{1-a^2+1}}\right)}{a\sqrt{1-a^2}} - \frac{2ib \csc^{-1}(a+bx) \log\left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} + \frac{2ib \csc^{-1}(a+bx) \log\left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{\sqrt{1-a^2+1}}\right)}{a\sqrt{1-a^2}} - \frac{b \csc^{-1}(a+bx)^2}{a} - \frac{\csc^{-1}(a+bx)^2}{x}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{ArcCsc}[a+b*x]^2/x^2, x]$

[Out] $-(\text{ArcCsc}[a+b*x]^2/a) - \text{ArcCsc}[a+b*x]^2/x - ((2*I)*b*\text{ArcCsc}[a+b*x]*\text{Log}[1+(I*a*E^{(I*\text{ArcCsc}[a+b*x])}/(1-Sqrt[1-a^2]))]/(a*Sqrt[1-a^2]) + ((2*I)*b*\text{ArcCsc}[a+b*x]*\text{Log}[1+(I*a*E^{(I*\text{ArcCsc}[a+b*x])}/(1+Sqrt[1-a^2]))]/(a*Sqrt[1-a^2]) - (2*b*\text{PolyLog}[2, ((-I)*a*E^{(I*\text{ArcCsc}[a+b*x])}/(1-Sqrt[1-a^2]))]/(a*Sqrt[1-a^2]) + (2*b*\text{PolyLog}[2, ((-I)*a*E^{(I*\text{ArcCsc}[a+b*x])}/(1+Sqrt[1-a^2]))]/(a*Sqrt[1-a^2]))/(a*Sqrt[1-a^2])$

Rule 2221

$\text{Int}[(((F_*)^((g_*)*((e_*)+(f_*)*(x_))))^{(n_*)}*((c_*)+(d_*)*(x_))^{(m_*)})/((a_*)+(b_*)*((F_*)^((g_*)*((e_*)+(f_*)*(x_))))^{(n_*)}), x]\text{Symbol} \Rightarrow \text{Simp}[((c+d*x)^m/(b*f*g*n*\text{Log}[F]))*\text{Log}[1+b*((F^(g*(e+f*x)))^n/a)], x] - \text{Dist}[d*(m/(b*f*g*n*\text{Log}[F])), \text{Int}[(c+d*x)^(m-1)*\text{Log}[1+b*((F^(g*(e+f*x)))^n/a)], x], x]; \text{FreeQ}\{F, a, b, c, d, e, f, g, n\}, x] \&& \text{IGtQ}[m, 0]$

Rule 2296

$\text{Int}[((F_*)^((u_*)*((f_*)+(g_*)*(x_)))^{(m_*)})/((a_*)+(b_*)*(F_*)^((u_*)+(c_*)*(F_*)^((v_)))), x)\text{Symbol} \Rightarrow \text{With}[\{q=\text{Rt}[b^2-4*a*c, 2]\}, \text{Dist}[2*(c/q), \text{Int}[(f+g*x)^m*(F^u/(b-q+2*c*F^u)), x], x] - \text{Dist}[2*(c/q), \text{Int}[(f+g*x)^m$

```

$$*(F^u/(b + q + 2*c*F^u)), x], x]] /; FreeQ[{F, a, b, c, f, g}, x] && EqQ[v, 2*u] && LinearQ[u, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[m, 0]$$

```

Rule 2317

```
Int[Log[(a_) + (b_)*((F_)^((e_)*(c_) + (d_)*(x_))))^(n_)], x_Symbol] :> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2438

```
Int[Log[(c_)*(d_) + (e_)*(x_)^(n_)]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 3404

```
Int[((c_) + (d_)*(x_))^(m_)/((a_) + (b_)*sin[(e_) + (f_)*(x_)]), x_Symbol] :> Dist[2, Int[(c + d*x)^m*(E^(I*(e + f*x))/(I*b + 2*a*E^(I*(e + f*x))) - I*b*E^(2*I*(e + f*x))), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[a^2 - b^2, 0] && IGtQ[m, 0]
```

Rule 4276

```
Int[(csc[(e_) + (f_)*(x_)]*(b_) + (a_))^(n_)*((c_) + (d_)*(x_))^(m_), x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, 1/(Sin[e + f*x]^n/(b + a*Sin[e + f*x])^n)], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && ILtQ[n, 0] && IGtQ[m, 0]
```

Rule 4512

```
Int[Cot[(c_) + (d_)*(x_)]*Csc[(c_) + (d_)*(x_)]*(Csc[(c_) + (d_)*(x_)]*(b_) + (a_))^(n_)*((e_) + (f_)*(x_))^(m_), x_Symbol] :> Simp[(-(e + f*x)^m)*((a + b*Csc[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[f*(m/(b*d*(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Csc[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]
```

Rule 5367

```
Int[((a_) + ArcCsc[(c_) + (d_)*(x_)]*(b_))^(p_)*((e_) + (f_)*(x_))^(m_), x_Symbol] :> Dist[-(d^(m + 1))^(-1), Subst[Int[(a + b*x)^p*Csc[x]*Cot[x]*(d*e - c*f + f*Csc[x])^m, x], x, ArcCsc[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Rubi steps

$$\begin{aligned}
\int \frac{\csc^{-1}(a+bx)^2}{x^2} dx &= - \left(b \text{Subst} \left(\int \frac{x^2 \cot(x) \csc(x)}{(-a + \csc(x))^2} dx, x, \csc^{-1}(a+bx) \right) \right) \\
&= - \frac{\csc^{-1}(a+bx)^2}{x} + (2b) \text{Subst} \left(\int \frac{x}{-a + \csc(x)} dx, x, \csc^{-1}(a+bx) \right) \\
&= - \frac{\csc^{-1}(a+bx)^2}{x} + (2b) \text{Subst} \left(\int \left(-\frac{x}{a} + \frac{x}{a(1-a \sin(x))} \right) dx, x, \csc^{-1}(a+bx) \right) \\
&= - \frac{b \csc^{-1}(a+bx)^2}{a} - \frac{\csc^{-1}(a+bx)^2}{x} + \frac{(2b) \text{Subst} \left(\int \frac{x}{1-a \sin(x)} dx, x, \csc^{-1}(a+bx) \right)}{a} \\
&= - \frac{b \csc^{-1}(a+bx)^2}{a} - \frac{\csc^{-1}(a+bx)^2}{x} + \frac{(4b) \text{Subst} \left(\int \frac{e^{ix} x}{-ia+2e^{ix}+iae^{2ix}} dx, x, \csc^{-1}(a+bx) \right)}{a} \\
&= - \frac{b \csc^{-1}(a+bx)^2}{a} - \frac{\csc^{-1}(a+bx)^2}{x} + \frac{(4ib) \text{Subst} \left(\int \frac{e^{ix} x}{2-2\sqrt{1-a^2}+2iae^{ix}} dx, x, \csc^{-1}(a+bx) \right)}{\sqrt{1-a^2}} \\
&= - \frac{b \csc^{-1}(a+bx)^2}{a} - \frac{\csc^{-1}(a+bx)^2}{x} - \frac{2ib \csc^{-1}(a+bx) \log \left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right)}{a \sqrt{1-a^2}} + \\
&= - \frac{b \csc^{-1}(a+bx)^2}{a} - \frac{\csc^{-1}(a+bx)^2}{x} - \frac{2ib \csc^{-1}(a+bx) \log \left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right)}{a \sqrt{1-a^2}} + \\
&= - \frac{b \csc^{-1}(a+bx)^2}{a} - \frac{\csc^{-1}(a+bx)^2}{x} - \frac{2ib \csc^{-1}(a+bx) \log \left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right)}{a \sqrt{1-a^2}}
\end{aligned}$$

Mathematica [B] Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 802 vs. $2(254) = 508$.

time = 2.13, size = 802, normalized size = 3.16

$$\left[\dots \right]$$

Antiderivative was successfully verified.

[In] `Integrate[ArcCsc[a + b*x]^2/x^2,x]`

[Out]
$$\begin{aligned}
&-((b*((a+b*x)*ArcCsc[a+b*x]^2)/(b*x)) + (2*\pi*ArcTan[(a - Tan[ArcCsc[a+b*x]/2])/Sqrt[1 - a^2]])/Sqrt[1 - a^2] + (2*(-2*ArcCos[a^(-1)]*ArcTanh[((1 + a)*Cot[(Pi + 2*ArcCsc[a + b*x])/4])/Sqrt[-1 + a^2]]) + (Pi - 2*ArcCsc[a + b*x])*ArcTanh[((-1 + a)*Tan[(Pi + 2*ArcCsc[a + b*x])/4])/Sqrt[-1 + a^2]] + (ArcCos[a^(-1)] + (2*I)*(-ArcTanh[((1 + a)*Cot[(Pi + 2*ArcCsc[a + b*x])/4])
\end{aligned}$$

$$\begin{aligned}
& \text{]}/\text{Sqrt}[-1 + a^2] + \text{ArcTanh}[((-1 + a)*\text{Tan}[(\text{Pi} + 2*\text{ArcCsc}[a + b*x])/4])/\text{Sqr}\\
& \text{t}[-1 + a^2]]]*\text{Log}[((1/2 + I/2)*\text{Sqrt}[-1 + a^2])/(\text{Sqrt}[a]*E^{((I/2)*\text{ArcCsc}[a}\\
& + b*x])* \text{Sqrt}[-((b*x)/(a + b*x))])] + (\text{ArcCos}[a^{(-1)}] + (2*I)*\text{ArcTanh}[((1 +\\
& a)*\text{Cot}[(\text{Pi} + 2*\text{ArcCsc}[a + b*x])/4])/\text{Sqrt}[-1 + a^2]] - (2*I)*\text{ArcTanh}[(((-1 +\\
& a)*\text{Tan}[(\text{Pi} + 2*\text{ArcCsc}[a + b*x])/4])/\text{Sqrt}[-1 + a^2]])*\text{Log}[((1/2 - I/2)*\text{Sqr}\\
& \text{t}[-1 + a^2]*E^{((I/2)*\text{ArcCsc}[a + b*x])}/(\text{Sqrt}[a]*\text{Sqrt}[-((b*x)/(a + b*x))])] -\\
& (\text{ArcCos}[a^{(-1)}] - (2*I)*\text{ArcTanh}[((1 + a)*\text{Cot}[(\text{Pi} + 2*\text{ArcCsc}[a + b*x])/4])/\text{S}\\
& \text{qr}[-1 + a^2]]*\text{Log}[((-1 + a)*(I + I*a + \text{Sqrt}[-1 + a^2]))*(-I + \text{Cot}[(\text{Pi} + 2*\text{A}\\
& \text{rcCsc}[a + b*x])/4])/(a*(-1 + a + \text{Sqrt}[-1 + a^2]*\text{Cot}[(\text{Pi} + 2*\text{ArcCsc}[a + b*x])/4])] -\\
& (\text{ArcCos}[a^{(-1)}] + (2*I)*\text{ArcTanh}[((1 + a)*\text{Cot}[(\text{Pi} + 2*\text{ArcCsc}[a + b*x])/4])/\text{S}\\
& \text{qr}[-1 + a^2]]*\text{Log}[((-1 + a)*(-I - I*a + \text{Sqrt}[-1 + a^2]))*(I + \\
& \text{Cot}[(\text{Pi} + 2*\text{ArcCsc}[a + b*x])/4])/(a*(-1 + a + \text{Sqrt}[-1 + a^2]*\text{Cot}[(\text{Pi} + 2*\text{A}\\
& \text{rcCsc}[a + b*x])/4])] + I*(-\text{PolyLog}[2, ((1 - I*\text{Sqrt}[-1 + a^2])*(1 - a + \text{Sqr}\\
& \text{t}[-1 + a^2]*\text{Cot}[(\text{Pi} + 2*\text{ArcCsc}[a + b*x])/4]))/(a*(-1 + a + \text{Sqrt}[-1 + a^2]*\text{C}\\
& \text{ot}[(\text{Pi} + 2*\text{ArcCsc}[a + b*x])/4])) + \text{PolyLog}[2, ((1 + I*\text{Sqrt}[-1 + a^2])*(1 - a + \text{Sqr}\\
& \text{t}[-1 + a^2]*\text{Cot}[(\text{Pi} + 2*\text{ArcCsc}[a + b*x])/4]))/(a*(-1 + a + \text{Sqrt}[-1 + a^2]*\text{C}\\
& \text{ot}[(\text{Pi} + 2*\text{ArcCsc}[a + b*x])/4]))]))/\text{Sqrt}[-1 + a^2]))/a)
\end{aligned}$$

Maple [A]

time = 6.84, size = 302, normalized size = 1.19

method	result
derivativedivides	$ b \left(-\frac{(bx+a)\text{arccsc}(bx+a)^2}{bxa} - \frac{2 \text{arccsc}(bx+a) \ln \left(\frac{-\left(\frac{i}{bx+a} + \sqrt{1 - \frac{1}{(bx+a)^2}}\right)a + \sqrt{a^2 - 1} + i}{i + \sqrt{a^2 - 1}} \right)}{a\sqrt{a^2 - 1}} + \right. \\ \left. \frac{2 \text{arccsc}(bx+a) \ln \left(\frac{-\left(\frac{i}{bx+a} + \sqrt{1 - \frac{1}{(bx+a)^2}}\right)a + \sqrt{a^2 - 1} + i}{i + \sqrt{a^2 - 1}} \right)}{a\sqrt{a^2 - 1}} \right) $
default	$ b \left(-\frac{(bx+a)\text{arccsc}(bx+a)^2}{bxa} - \frac{2 \text{arccsc}(bx+a) \ln \left(\frac{-\left(\frac{i}{bx+a} + \sqrt{1 - \frac{1}{(bx+a)^2}}\right)a + \sqrt{a^2 - 1} + i}{i + \sqrt{a^2 - 1}} \right)}{a\sqrt{a^2 - 1}} + \right. \\ \left. \frac{2 \text{arccsc}(bx+a) \ln \left(\frac{-\left(\frac{i}{bx+a} + \sqrt{1 - \frac{1}{(bx+a)^2}}\right)a + \sqrt{a^2 - 1} + i}{i + \sqrt{a^2 - 1}} \right)}{a\sqrt{a^2 - 1}} \right) $

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(arccsc(b*x+a)^2/x^2,x,method=_RETURNVERBOSE)

[Out] b*(-(b*x+a)*arccsc(b*x+a)^2/b/x/a-2/a*arccsc(b*x+a)/(a^2-1)^(1/2)*ln((-I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))*a+(a^2-1)^(1/2)+I)/(I+(a^2-1)^(1/2))+2/a*arc
csc(b*x+a)/(a^2-1)^(1/2)*ln((-I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))*a+I-(a^2-1)^(1/2))/(I-(a^2-1)^(1/2))+2*I/a/(a^2-1)^(1/2)*dilog((-I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))*a+(a^2-1)^(1/2)+I)/(I+(a^2-1)^(1/2))-2*I/a/(a^2-1)^(1/2)*dil
og((-I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))*a+I-(a^2-1)^(1/2))/(I-(a^2-1)^(1/2)))
)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(arccsc(b*x+a)^2/x^2,x, algorithm="maxima")
[Out] -1/4*(4*arctan2(1, sqrt(b*x + a + 1)*sqrt(b*x + a - 1))^2 + 4*x*integrate((2*sqrt(b*x + a + 1)*sqrt(b*x + a - 1)*b*x*arctan2(1, sqrt(b*x + a + 1)*sqrt(b*x + a - 1)) + (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2 - 1)*b*x - a)*log(b*x + a)^2 + (b^3*x^3 + 2*a*b^2*x^2 + (a^2 - 1)*b*x - (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2 - 1)*b*x - a)*log(b*x + a))*log(b^2*x^2 + 2*a*b*x + a^2))/(b^3*x^5 + 3*a*b^2*x^4 + (3*a^2 - 1)*b*x^3 + (a^3 - a)*x^2), x) - log(b^2*x^2 + 2*a*b*x + a^2)^2)/x
```

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(arccsc(b*x+a)^2/x^2,x, algorithm="fricas")
[Out] integral(arccsc(b*x + a)^2/x^2, x)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{acsc}^2(a + bx)}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(acsc(b*x+a)**2/x**2,x)
[Out] Integral(acsc(a + b*x)**2/x**2, x)
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(arccsc(b*x+a)^2/x^2,x, algorithm="giac")
[Out] integrate(arccsc(b*x + a)^2/x^2, x)
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\arcsin\left(\frac{1}{a+bx}\right)^2}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(asin(1/(a + b*x))^2/x^2,x)

[Out] int(asin(1/(a + b*x))^2/x^2, x)

$$\int x^2 \csc^{-1}(a + bx)^3 dx$$

Optimal. Leaf size=464

$$\frac{(a + bx) \csc^{-1}(a + bx)}{b^3} - \frac{3ia \csc^{-1}(a + bx)^2}{b^3} - \frac{3a(a + bx) \sqrt{1 - \frac{1}{(a + bx)^2}} \csc^{-1}(a + bx)^2}{b^3} + \frac{(a + bx)^2 \sqrt{1 - \frac{1}{(a + bx)^2}}}{b^3}$$

[Out] $(b*x+a)*\text{arccsc}(b*x+a)/b^3 + I*\text{arccsc}(b*x+a)*\text{polylog}(2, I/(b*x+a) + (1-1/(b*x+a)^2)^(1/2))/b^3 + 1/3*a^3*\text{arccsc}(b*x+a)^3/b^3 + 1/3*x^3*\text{arccsc}(b*x+a)^3 + \text{arccsc}(b*x+a)^2*\text{arctanh}(I/(b*x+a) + (1-1/(b*x+a)^2)^(1/2))/b^3 + 6*a^2*\text{arccsc}(b*x+a)^2*a*\text{rctanh}(I/(b*x+a) + (1-1/(b*x+a)^2)^(1/2))/b^3 + \text{arctanh}((1-1/(b*x+a)^2)^(1/2))/b^3 + 3+6*a^2*\text{arccsc}(b*x+a)^2*a*\text{rctanh}(I/(b*x+a) + (1-1/(b*x+a)^2)^(1/2))/b^3 + 6*a^2*\text{arccsc}(b*x+a)*\text{ln}(1-(I/(b*x+a) + (1-1/(b*x+a)^2)^(1/2))^2)/b^3 - 3*I*a*polylog(2, (I/(b*x+a) + (1-1/(b*x+a)^2)^(1/2))^2)/b^3 - 3*I*a*\text{arccsc}(b*x+a)^2/b^3 - I*\text{arccsc}(b*x+a)*\text{polylog}(2, -I/(b*x+a) - (1-1/(b*x+a)^2)^(1/2))/b^3 + 6*I*a^2*\text{arccsc}(b*x+a)*\text{polylog}(2, I/(b*x+a) + (1-1/(b*x+a)^2)^(1/2))/b^3 - 6*I*a^2*\text{arccsc}(b*x+a)*\text{polylog}(2, -I/(b*x+a) - (1-1/(b*x+a)^2)^(1/2))/b^3 + \text{polylog}(3, -I/(b*x+a) - (1-1/(b*x+a)^2)^(1/2))/b^3 + 6*a^2*\text{polylog}(3, -I/(b*x+a) - (1-1/(b*x+a)^2)^(1/2))/b^3 - 3*\text{polylog}(3, I/(b*x+a) + (1-1/(b*x+a)^2)^(1/2))/b^3 - 6*a^2*\text{polylog}(3, I/(b*x+a) + (1-1/(b*x+a)^2)^(1/2))/b^3 - 3*a*(b*x+a)*\text{arccsc}(b*x+a)^2*(1-1/(b*x+a)^2)^(1/2)/b^3 + 1/2*(b*x+a)^2*\text{arccsc}(b*x+a)^2*(1-1/(b*x+a)^2)^(1/2)/b^3$

Rubi [A]

time = 0.28, antiderivative size = 464, normalized size of antiderivative = 1.00, number of steps used = 25, number of rules used = 14, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}}$ = 1.167, Rules used = {5367, 4512, 4275, 4268, 2611, 2320, 6724, 4269, 3798, 2221, 2317, 2438, 4271, 3855}

$$\frac{x^2 \csc ^{-1}(a+b x)^3}{b^3}+\frac{3 i a \csc ^{-1}(a+b x)^2}{b^3}-\frac{3 a (a+b x) \sqrt{1-\frac{1}{(a+b x)^2}} \csc ^{-1}(a+b x)^2}{b^3}+\frac{(a+b x)^2 \sqrt{1-\frac{1}{(a+b x)^2}}}{b^3}$$

Antiderivative was successfully verified.

[In] $\text{Int}[x^2 \text{ArcCsc}[a + b*x]^3, x]$

[Out] $((a + b*x)*\text{ArcCsc}[a + b*x])/b^3 - ((3*I)*a*\text{ArcCsc}[a + b*x]^2)/b^3 - (3*a*(a + b*x)*\text{Sqrt}[1 - (a + b*x)^{-2}]*\text{ArcCsc}[a + b*x]^2)/b^3 + ((a + b*x)^2*\text{Sqrt}[1 - (a + b*x)^{-2}]*\text{ArcCsc}[a + b*x]^2)/(2*b^3) + (a^3*\text{ArcCsc}[a + b*x]^3)/(3*b^3) + (x^3*\text{ArcCsc}[a + b*x]^3)/3 + (\text{ArcCsc}[a + b*x]^2*\text{ArcTanh}[E^(I*\text{ArcCsc}[a + b*x])])/b^3 + (6*a^2*\text{ArcCsc}[a + b*x]^2*\text{ArcTanh}[E^(I*\text{ArcCsc}[a + b*x])])/b^3 + \text{ArcTanh}[\text{Sqrt}[1 - (a + b*x)^{-2}]])/b^3 + (6*a*\text{ArcCsc}[a + b*x]*\text{Log}[1 - E^(-(2*I)*\text{ArcCsc}[a + b*x])])/b^3 - (I*\text{ArcCsc}[a + b*x]*\text{PolyLog}[2, -E^(I*\text{ArcCsc}[a + b*x])])/b^3 - ((6*I)*a^2*\text{ArcCsc}[a + b*x]*\text{PolyLog}[2, -E^(I*\text{ArcCsc}[a + b*x])])/b^3 + (I*\text{ArcCsc}[a + b*x]*\text{PolyLog}[2, E^(I*\text{ArcCsc}[a + b*x])])/b^3 + ((6*I)*a^2*\text{ArcCsc}[a + b*x]*\text{PolyLog}[2, E^(I*\text{ArcCsc}[a + b*x])])/b^3 - ((3*I)*a*\text{PolyLog}[2, E^(-(2*I)*\text{ArcCsc}[a + b*x])])/b^3 + \text{PolyLog}[3, -E^(I*\text{ArcCsc}[a + b*x])] + \frac{1}{2} x^2 \csc ^{-1}(a+b x)^3$

$$b*x)]/b^3 + (6*a^2*PolyLog[3, -E^(I*ArcCsc[a + b*x])])/b^3 - PolyLog[3, E^(I*ArcCsc[a + b*x])]/b^3 - (6*a^2*PolyLog[3, E^(I*ArcCsc[a + b*x])])/b^3$$
Rule 2221

```
Int[((F_)^((g_.)*(e_.) + (f_.)*(x_)))*((c_.) + (d_.)*(x_))^((m_.))/((a_) + (b_.)*((F_)^((g_.)*(e_.) + (f_.)*(x_))))^((n_.))), x_Symbol] :> Simplify[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x]; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2317

```
Int[Log[(a_) + (b_.)*((F_)^((e_.)*(c_.) + (d_.)*(x_))))^((n_.))], x_Symbol] :> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x]; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2320

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]]; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^((n_.)))^((m_.))]; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_]]; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2438

```
Int[Log[(c_.)*(d_.) + (e_.)*(x_)^((n_.))]/(x_), x_Symbol] :> Simplify[-PolyLog[2, (-c)*e*x^n]/n, x]; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 2611

```
Int[Log[1 + (e_.)*((F_)^((c_.)*(a_.) + (b_.)*(x_)))]*((f_.) + (g_.)*(x_)^((m_.)), x_Symbol] :> Simplify[((-f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x]; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]
```

Rule 3798

```
Int[((c_.) + (d_.)*(x_)^((m_.))*tan[(e_.) + Pi*(k_.) + (f_.)*(x_)], x_Symbol] :> Simplify[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Dist[2*I, Int[(c + d*x)^m * E^(2*I*k*Pi)*(E^(2*I*(e + f*x))/(1 + E^(2*I*k*Pi)*E^(2*I*(e + f*x)))), x], x]; FreeQ[{c, d, e, f}, x] && IntegerQ[4*k] && IGtQ[m, 0]
```

Rule 3855

```
Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

Rule 4268

```
Int[csc[(e_.) + (f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_), x_Symbol] :> Simp[-2*(c + d*x)^m*(ArcTanh[E^(I*(e + f*x))/f], x] + (-Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Log[1 - E^(I*(e + f*x))], x], x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Log[1 + E^(I*(e + f*x))], x], x]) /; FreeQ[{c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 4269

```
Int[csc[(e_.) + (f_.)*(x_)]^2*((c_.) + (d_.)*(x_))^(m_), x_Symbol] :> Simp[[-(c + d*x)^m)*(Cot[e + f*x]/f), x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Cot[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

Rule 4271

```
Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(n_)*((c_.) + (d_.)*(x_))^(m_), x_Symbol] :> Simp[(-b^2)*(c + d*x)^m*Cot[e + f*x]*((b*Csc[e + f*x])^(n - 2)/(f^(n - 1))), x] + (Dist[b^2*d^2*m*((m - 1)/(f^2*(n - 1)*(n - 2))), Int[(c + d*x)^(m - 2)*(b*Csc[e + f*x])^(n - 2), x], x] + Dist[b^2*((n - 2)/(n - 1)), Int[(c + d*x)^m*(b*Csc[e + f*x])^(n - 2), x], x] - Simp[b^2*d*m*(c + d*x)^(m - 1)*((b*Csc[e + f*x])^(n - 2)/(f^2*(n - 1)*(n - 2))), x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && NeQ[n, 2] && GtQ[m, 1]
```

Rule 4275

```
Int[(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(n_)*((c_.) + (d_.)*(x_))^(m_), x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (a + b*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rule 4512

```
Int[Cot[(c_.) + (d_.)*(x_)]*Csc[(c_.) + (d_.)*(x_)]*(Csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_.))^(n_)*((e_.) + (f_.)*(x_))^(m_), x_Symbol] :> Simp[(-(e + f*x)^m)*((a + b*Csc[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[f*(m/(b*d*(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Csc[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]
```

Rule 5367

```
Int[((a_.) + ArcCsc[(c_.) + (d_.)*(x_)]*(b_.))^(p_)*((e_.) + (f_.)*(x_))^(m_), x_Symbol] :> Dist[-(d^(m + 1))^(-1), Subst[Int[(a + b*x)^p*Csc[x]*Cot[x]*(d*e - c*f + f*Csc[x])^m, x], x, ArcCsc[c + d*x]], x] /; FreeQ[{a, b, c,
```

```
d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Rule 6724

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^p]/((d_.) + (e_.)*(x_)), x_Symbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

Rubi steps

$$\begin{aligned}
\int x^2 \csc^{-1}(a + bx)^3 dx &= -\frac{\text{Subst}\left(\int x^3 \cot(x) \csc(x)(-a + \csc(x))^2 dx, x, \csc^{-1}(a + bx)\right)}{b^3} \\
&= \frac{1}{3}x^3 \csc^{-1}(a + bx)^3 - \frac{\text{Subst}\left(\int x^2(-a + \csc(x))^3 dx, x, \csc^{-1}(a + bx)\right)}{b^3} \\
&= \frac{1}{3}x^3 \csc^{-1}(a + bx)^3 - \frac{\text{Subst}\left(\int (-a^3x^2 + 3a^2x^2 \csc(x) - 3ax^2 \csc^2(x) + x^2 \csc^3(x)) dx, x, \csc^{-1}(a + bx)\right)}{b^3} \\
&= \frac{a^3 \csc^{-1}(a + bx)^3}{3b^3} + \frac{1}{3}x^3 \csc^{-1}(a + bx)^3 - \frac{\text{Subst}\left(\int x^2 \csc^3(x) dx, x, \csc^{-1}(a + bx)\right)}{b^3} \\
&= \frac{(a + bx) \csc^{-1}(a + bx)}{b^3} - \frac{3a(a + bx) \sqrt{1 - \frac{1}{(a + bx)^2}} \csc^{-1}(a + bx)^2}{b^3} + \frac{(a + bx)^2 \csc^{-1}(a + bx)^2}{b^3} \\
&= \frac{(a + bx) \csc^{-1}(a + bx)}{b^3} - \frac{3ia \csc^{-1}(a + bx)^2}{b^3} - \frac{3a(a + bx) \sqrt{1 - \frac{1}{(a + bx)^2}} \csc^{-1}(a + bx)^2}{b^3} \\
&= \frac{(a + bx) \csc^{-1}(a + bx)}{b^3} - \frac{3ia \csc^{-1}(a + bx)^2}{b^3} - \frac{3a(a + bx) \sqrt{1 - \frac{1}{(a + bx)^2}} \csc^{-1}(a + bx)^2}{b^3} \\
&= \frac{(a + bx) \csc^{-1}(a + bx)}{b^3} - \frac{3ia \csc^{-1}(a + bx)^2}{b^3} - \frac{3a(a + bx) \sqrt{1 - \frac{1}{(a + bx)^2}} \csc^{-1}(a + bx)^2}{b^3} \\
&= \frac{(a + bx) \csc^{-1}(a + bx)}{b^3} - \frac{3ia \csc^{-1}(a + bx)^2}{b^3} - \frac{3a(a + bx) \sqrt{1 - \frac{1}{(a + bx)^2}} \csc^{-1}(a + bx)^2}{b^3}
\end{aligned}$$

Mathematica [A]

time = 5.99, size = 656, normalized size = 1.41

Warning: Unable to verify antiderivative.

```
[In] Integrate[x^2*ArcCsc[a + b*x]^3,x]
[Out] -1/24*((72*I)*a*ArcCsc[a + b*x]^2 - 12*ArcCsc[a + b*x]*Cot[ArcCsc[a + b*x]/2] + 36*a*ArcCsc[a + b*x]^2*Cot[ArcCsc[a + b*x]/2] - 2*ArcCsc[a + b*x]^3*Cos[ArcCsc[a + b*x]/2] - 12*a^2*ArcCsc[a + b*x]^3*Cot[ArcCsc[a + b*x]/2] - 3*ArcCsc[a + b*x]^2*Csc[ArcCsc[a + b*x]/2]^2 + 6*a*ArcCsc[a + b*x]^3*Csc[ArcCsc[a + b*x]/2]^2 - (ArcCsc[a + b*x]^3*Csc[ArcCsc[a + b*x]/2]^4)/(2*(a + b*x)) + 12*ArcCsc[a + b*x]^2*Log[1 - E^(I*ArcCsc[a + b*x])] + 72*a^2*ArcCsc[a + b*x]^2*Log[1 - E^(I*ArcCsc[a + b*x])] - 12*ArcCsc[a + b*x]^2*Log[1 + E^(I*ArcCsc[a + b*x])] - 72*a^2*ArcCsc[a + b*x]^2*Log[1 + E^(I*ArcCsc[a + b*x])] - 144*a*ArcCsc[a + b*x]*Log[1 - E^((2*I)*ArcCsc[a + b*x])] + 24*Log[Tan[ArcCsc[a + b*x]/2]] + (24*I)*(1 + 6*a^2)*ArcCsc[a + b*x]*PolyLog[2, -E^(I*ArcCsc[a + b*x])] - (24*I)*(1 + 6*a^2)*ArcCsc[a + b*x]*PolyLog[2, E^(I*ArcCsc[a + b*x])] + (72*I)*a*PolyLog[2, E^((2*I)*ArcCsc[a + b*x])] - 24*PolyLog[3, -E^(I*ArcCsc[a + b*x])] - 144*a^2*PolyLog[3, -E^(I*ArcCsc[a + b*x])] + 24*PolyLog[3, E^(I*ArcCsc[a + b*x])] + 144*a^2*PolyLog[3, E^(I*ArcCsc[a + b*x])] + 3*ArcCsc[a + b*x]^2*Sec[ArcCsc[a + b*x]/2]^2 + 6*a*ArcCsc[a + b*x]^3*Sec[ArcCsc[a + b*x]/2]^2 - 8*(a + b*x)^3*ArcCsc[a + b*x]^3*Sin[ArcCsc[a + b*x]/2]^4 - 12*ArcCsc[a + b*x]*Tan[ArcCsc[a + b*x]/2] - 36*a*ArcCsc[a + b*x]^2*Tan[ArcCsc[a + b*x]/2] - 2*ArcCsc[a + b*x]^3*Tan[ArcCsc[a + b*x]/2] - 12*a^2*ArcCsc[a + b*x]^3*Tan[ArcCsc[a + b*x]/2])/b^3
```

Maple [A]

time = 9.85, size = 749, normalized size = 1.61 Too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(x^2*arccsc(b*x+a)^3,x,method=_RETURNVERBOSE)
[Out] 1/b^3*(1/6*arccsc(b*x+a)*(6*arccsc(b*x+a)^2*a^2*(b*x+a) - 6*arccsc(b*x+a)^2*a*(b*x+a)^2 + 2*arccsc(b*x+a)^2*(b*x+a)^3 - 18*arccsc(b*x+a)*((b*x+a)^2 - 1)/(b*x+a)^2)^(1/2)*a*(b*x+a) + 3*arccsc(b*x+a)*(((b*x+a)^2 - 1)/(b*x+a)^2)^(1/2)*(b*x+a)^2 + 18*I*arccsc(b*x+a)*a + 6*b*x + 6*a - 1/2*arccsc(b*x+a)^2*ln(1 - I/(b*x+a) - (1 - 1/(b*x+a)^2)^(1/2)) - 6*I*a*polylog(2, -I/(b*x+a) - (1 - 1/(b*x+a)^2)^(1/2)) - polylog(3, I/(b*x+a) + (1 - 1/(b*x+a)^2)^(1/2)) + 1/2*arccsc(b*x+a)^2*ln(1 + I/(b*x+a) + (1 - 1/(b*x+a)^2)^(1/2)) - I*arccsc(b*x+a)*polylog(2, -I/(b*x+a) - (1 - 1/(b*x+a)^2)^(1/2)) + polylog(3, -I/(b*x+a) - (1 - 1/(b*x+a)^2)^(1/2)) + 2*arctanh(I/(b*x+a) + (1 - 1/(b*x+a)^2)^(1/2)) - 6*I*a*polylog(2, I/(b*x+a) + (1 - 1/(b*x+a)^2)^(1/2)) + 6*a*arccsc(b*x+a)*ln(1 - I/(b*x+a) - (1 - 1/(b*x+a)^2)^(1/2)) + 6*I*a^2*arccsc(b*x+a)*polylog(2, I/(b*x+a) + (1 - 1/(b*x+a)^2)^(1/2)) + 6*a*arccsc(b*x+a)*ln(1 + I/(b*x+a) + (1 - 1/(b*x+a)^2)^(1/2)) + I*arccsc(b*x+a)*polylog(2, I/(b*x+a) + (1 - 1/(b*x+a)^2)^(1/2)) - 3*a^2*arccsc(b*x+a)^2*ln(1 - I/(b*x+a) - (1 - 1/(b*x+a)^2)^(1/2)) - 6*I*arccsc(b*x+a)^2*a - 6*a^2*polylog(3, I/(b*x+a) + (1 - 1/(b*x+a)^2)^(1/2)) + 3*a^2*arccsc(b*x+a)^2*ln(1 + I/(b*x+a) + (1 - 1/(b*x+a)^2)^(1/2)) - 6*I*a^2*arccsc(b*x+a)*polylog(2, -I/(b*x+a) - (1 - 1/(b*x+a)^2)^(1/2)) + 6*a^2*polylog(3, -I/(b*x+a) - (1 - 1/(b*x+a)^2)^(1/2)))
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^2*arccsc(b*x+a)^3,x, algorithm="maxima")`

[Out]
$$\begin{aligned} & \frac{1}{3}x^3 \operatorname{arctan2}(1, \sqrt{bx + a + 1}) \sqrt{bx + a - 1})^3 - \frac{1}{4}x^3 \operatorname{arctan2}(1, \sqrt{bx + a + 1}) \sqrt{bx + a - 1}) \log(b^2x^2 + 2axb + a^2)^2 - i \\ & \text{ntegrate}\left(\frac{1}{4} \cdot (12 \cdot (b^3x^5 \operatorname{arctan2}(1, \sqrt{bx + a + 1}) \sqrt{bx + a - 1}) + 3a^2b^2x^4 \operatorname{arctan2}(1, \sqrt{bx + a + 1}) \sqrt{bx + a - 1}) + (3a^2 \operatorname{arcta} \\ & n2(1, \sqrt{bx + a + 1}) \sqrt{bx + a - 1}) - \operatorname{arctan2}(1, \sqrt{bx + a + 1}) \sqrt{bx + a - 1})) \cdot b^2x^3 + (a^3 \operatorname{arctan2}(1, \sqrt{bx + a + 1}) \sqrt{bx + a - 1}) - a \operatorname{arctan2}(1, \sqrt{bx + a + 1}) \sqrt{bx + a - 1}) \cdot x^2) \cdot \log(bx + a) \\ & ^2 - (4b^2x^3 \operatorname{arctan2}(1, \sqrt{bx + a + 1}) \sqrt{bx + a - 1})^2 - b^2x^3 \log(b^2x^2 + 2axb + a^2)^2) \cdot \sqrt{bx + a + 1} \cdot \sqrt{bx + a - 1} - 4(b^3x^5 \operatorname{arctan2}(1, \sqrt{bx + a + 1}) \sqrt{bx + a - 1}) + 2a^2b^2x^4 \operatorname{arctan2}(1, \sqrt{bx + a + 1}) \sqrt{bx + a - 1}) + (a^2 \operatorname{arctan2}(1, \sqrt{bx + a + 1}) \sqrt{bx + a - 1}) - \operatorname{arctan2}(1, \sqrt{bx + a + 1}) \sqrt{bx + a - 1}) \cdot b^2x^3 \\ & + 3(b^3x^5 \operatorname{arctan2}(1, \sqrt{bx + a + 1}) \sqrt{bx + a - 1}) + 3a^2b^2x^4 \operatorname{arctan2}(1, \sqrt{bx + a + 1}) \sqrt{bx + a - 1}) + (3a^2 \operatorname{arctan2}(1, \sqrt{bx + a + 1}) \sqrt{bx + a - 1}) - \operatorname{arctan2}(1, \sqrt{bx + a + 1}) \sqrt{bx + a - 1}) \cdot b^2x^3 + (a^3 \operatorname{arctan2}(1, \sqrt{bx + a + 1}) \sqrt{bx + a - 1}) - a \operatorname{arct} \\ & an2(1, \sqrt{bx + a + 1}) \sqrt{bx + a - 1}) \cdot x^2) \cdot \log(bx + a)) \cdot \log(b^2x^2 + 2axb + a^2)) / (b^3x^3 + 3a^2b^2x^2 + a^3 + (3a^2 - 1)b^2x - a), x) \end{aligned}$$

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^2*arccsc(b*x+a)^3,x, algorithm="fricas")`[Out] `integral(x^2*arccsc(b*x + a)^3, x)`**Sympy [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int x^2 \operatorname{acsc}^3(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x**2*acsc(b*x+a)**3,x)`

[Out] `Integral(x**2*acsc(a + b*x)**3, x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^2*arccsc(b*x+a)^3,x, algorithm="giac")`

[Out] `integrate(x^2*arccsc(b*x + a)^3, x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int x^2 \arcsin\left(\frac{1}{a + bx}\right)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x^2*asin(1/(a + b*x))^3,x)`

[Out] `int(x^2*asin(1/(a + b*x))^3, x)`

$$\text{3.34} \quad \int x \csc^{-1}(a + bx)^3 dx$$

Optimal. Leaf size=264

$$\frac{3i \csc^{-1}(a + bx)^2}{2b^2} + \frac{3(a + bx) \sqrt{1 - \frac{1}{(a + bx)^2}} \csc^{-1}(a + bx)^2}{2b^2} - \frac{a^2 \csc^{-1}(a + bx)^3}{2b^2} + \frac{1}{2} x^2 \csc^{-1}(a + bx)^3 - \frac{6a \csc^{-1}(a + bx)^2}{b^2}$$

[Out] $\frac{3}{2} \cdot \frac{I \cdot \operatorname{arccsc}(bx+a)^2}{b^2} - \frac{1}{2} \cdot \frac{3(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}} \csc^{-1}(a+bx)^2}{b^2} - \frac{a^2 \csc^{-1}(a+bx)^3}{2b^2} + \frac{1}{2} x^2 \csc^{-1}(a+bx)^3 - \frac{6a \csc^{-1}(a+bx)^2}{b^2}$

$\sim 3 - 6a \operatorname{arccsc}(bx+a)^2 \operatorname{arctanh}\left(\frac{I}{(bx+a)} + \frac{(1-1/(bx+a)^2)^{(1/2)}}{b^2}\right) - 3 \operatorname{arccsc}(bx+a) \ln\left(1 - \left(\frac{I}{(bx+a)} + \frac{(1-1/(bx+a)^2)^{(1/2)}}{b^2}\right)^2\right) + 6 \cdot I \cdot a \cdot \operatorname{arccsc}(bx+a) \cdot \operatorname{polylog}(2, -\frac{I}{(bx+a)} - \frac{(1-1/(bx+a)^2)^{(1/2)}}{b^2}) - 6 \cdot I \cdot a \cdot \operatorname{arccsc}(bx+a) \cdot \operatorname{polylog}(2, \frac{I}{(bx+a)} + \frac{(1-1/(bx+a)^2)^{(1/2)}}{b^2}) + 3 \cdot 2 \cdot I \cdot \operatorname{polylog}(2, \frac{I}{(bx+a)} + \frac{(1-1/(bx+a)^2)^{(1/2)}}{b^2}) - 2 \cdot 6 \cdot a \cdot \operatorname{polylog}(3, -\frac{I}{(bx+a)} - \frac{(1-1/(bx+a)^2)^{(1/2)}}{b^2}) - 2 \cdot 6 \cdot a \cdot \operatorname{polylog}(3, \frac{I}{(bx+a)} + \frac{(1-1/(bx+a)^2)^{(1/2)}}{b^2}) + 3 \cdot 2 \cdot (bx+a) \cdot \operatorname{arccsc}(bx+a) \cdot 2 \cdot (1-1/(bx+a)^2)^{(1/2)} / b^2$

Rubi [A]

time = 0.19, antiderivative size = 264, normalized size of antiderivative = 1.00, number of steps used = 16, number of rules used = 12, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 1.200$, Rules used = {5367, 4512, 4275, 4268, 2611, 2320, 6724, 4269, 3798, 2221, 2317, 2438}

$$-\frac{a^2 \csc^{-1}(a + bx)^2}{2b^2} + \frac{6i a \csc^{-1}(a + bx) \operatorname{Li}_2\left(-e^{i \csc^{-1}(a + bx)}\right)}{b^2} - \frac{6i a \csc^{-1}(a + bx) \operatorname{Li}_2\left(e^{i \csc^{-1}(a + bx)}\right)}{b^2} + \frac{3i \operatorname{Li}_2\left(e^{2i \csc^{-1}(a + bx)}\right)}{2b^2} + \frac{6i \operatorname{Li}_2\left(e^{i \csc^{-1}(a + bx)}\right)}{b^2} + \frac{3(a + bx) \sqrt{1 - \frac{1}{(a + bx)^2}} \csc^{-1}(a + bx)^2}{2b^2} + \frac{3i \csc^{-1}(a + bx)^2}{2b^2} - \frac{3 \csc^{-1}(a + bx) \log\left(1 - e^{2i \csc^{-1}(a + bx)}\right)}{b^2} - \frac{6i \csc^{-1}(a + bx)^2 \tanh^{-1}\left(e^{i \csc^{-1}(a + bx)}\right)}{b^2} + \frac{1}{2} x^2 \csc^{-1}(a + bx)^3$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[x \cdot \operatorname{ArcCsc}[a + b x]^3, x]$

[Out] $\frac{((3I)/2) \cdot \operatorname{ArcCsc}[a + b x]^2}{b^2} + \frac{3 \cdot (a + b x) \cdot \operatorname{Sqrt}[1 - (a + b x)^{-2}] \cdot \operatorname{ArcCsc}[a + b x]^2}{(2 b^2)} - \frac{(a^2 \cdot \operatorname{ArcCsc}[a + b x]^3)}{(2 b^2)} + \frac{(x^2 \cdot \operatorname{ArcCsc}[a + b x]^3)}{2} - \frac{(6 \cdot a \cdot \operatorname{ArcCsc}[a + b x]^2 \cdot \operatorname{ArcTanh}[E^{(I \cdot \operatorname{ArcCsc}[a + b x])}])}{b^2} - \frac{(3 \cdot \operatorname{ArcCsc}[a + b x] \cdot \operatorname{Log}[1 - E^{((2I) \cdot \operatorname{ArcCsc}[a + b x])}])}{b^2} + \frac{((6I) \cdot a \cdot \operatorname{ArcCsc}[a + b x] \cdot \operatorname{PolyLog}[2, -E^{(I \cdot \operatorname{ArcCsc}[a + b x])}])}{b^2} - \frac{((6I) \cdot a \cdot \operatorname{ArcCsc}[a + b x] \cdot \operatorname{PolyLog}[2, E^{(I \cdot \operatorname{ArcCsc}[a + b x])}])}{b^2} + \frac{((3I)/2) \cdot \operatorname{PolyLog}[2, E^{((2I) \cdot \operatorname{ArcCsc}[a + b x])}])}{b^2} - \frac{(6 \cdot a \cdot \operatorname{PolyLog}[3, -E^{(I \cdot \operatorname{ArcCsc}[a + b x])}])}{b^2} + \frac{(6 \cdot a \cdot \operatorname{PolyLog}[3, E^{(I \cdot \operatorname{ArcCsc}[a + b x])}])}{b^2}$

Rule 2221

```
Int[((F_)^((g_.)*(e_.)+(f_.)*(x_.))^(n_.)*((c_.)+(d_.)*(x_.))^(m_.))/((a_.)+(b_.)*((F_)^((g_.)*(e_.)+(f_.)*(x_.))))^(n_.)), x_Symbol] := Simp[((c+d*x)^m/(b*f*g*n*Log[F]))*Log[1+b*((F^(g*(e+f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c+d*x)^(m-1)*Log[1+b*((F^(g*(e+f*x)))^n/a)], x], x]; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2317

```

Int[Log[(a_) + (b_.)*((F_)^((e_)*(c_.) + (d_.)*(x_))))^(n_.)], x_Symbol]
:> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]

```

Rule 2320

```

Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x]
, Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*(a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]

```

Rule 2438

```

Int[Log[((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]

```

Rule 2611

```

Int[Log[1 + (e_.)*((F_)^((c_.)*(a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)
*(x_)^m, x_Symbol] :> Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^
(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e,
f, g, n}, x] && GtQ[m, 0]

```

Rule 3798

```

Int[((c_.) + (d_.)*(x_)^m)*tan[(e_.) + Pi*(k_.) + (f_.)*(x_)], x_Symbol]
:> Simp[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Dist[2*I, Int[(c + d*x)^m
*E^(2*I*k*Pi)*(E^(2*I*(e + f*x))/(1 + E^(2*I*k*Pi)*E^(2*I*(e + f*x)))), x],
x] /; FreeQ[{c, d, e, f}, x] && IntegerQ[4*k] && IGtQ[m, 0]

```

Rule 4268

```

Int[csc[(e_.) + (f_.)*(x_)]*((c_.) + (d_.)*(x_)^m, x_Symbol] :> Simp[-2*(c + d*x)^m*(ArcTanh[E^(I*(e + f*x))/f]), x] + (-Dist[d*(m/f), Int[(c + d
*x)^(m - 1)*Log[1 - E^(I*(e + f*x))], x], x] + Dist[d*(m/f), Int[(c + d*x)^
(m - 1)*Log[1 + E^(I*(e + f*x))], x], x]) /; FreeQ[{c, d, e, f}, x] && IGtQ
[m, 0]

```

Rule 4269

```

Int[csc[(e_.) + (f_.)*(x_)]^2*((c_.) + (d_.)*(x_)^m, x_Symbol] :> Simp
[(-(c + d*x)^m)*(Cot[e + f*x]/f), x] + Dist[d*(m/f), Int[(c + d*x)^
(m - 1)*Cot[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]

```

Rule 4275

```
Int[(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(n_.)*((c_.) + (d_.)*(x_.))^(m_.),
x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (a + b*Csc[e + f*x])^n, x],
x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rule 4512

```
Int[Cot[(c_.) + (d_.)*(x_.)]*Csc[(c_.) + (d_.)*(x_.)]*(Csc[(c_.) + (d_.)*(x_.)
]* (b_.) + (a_.))^(n_.)*((e_.) + (f_.)*(x_.))^(m_.), x_Symbol] :> Simp[(-(e +
f*x)^m)*(a + b*Csc[c + d*x])^(n + 1)/(b*d*(n + 1)), x] + Dist[f*(m/(b*d*(n + 1))),
Int[(e + f*x)^(m - 1)*(a + b*Csc[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]
```

Rule 5367

```
Int[((a_.) + ArcCsc[(c_.) + (d_.)*(x_.)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_.))^(m
_.), x_Symbol] :> Dist[-(d^(m + 1))^{(-1)}, Subst[Int[(a + b*x)^p*Csc[x]*Cot[
x]*(d*e - c*f + f*Csc[x])^m, x], x, ArcCsc[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Rule 6724

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_.))^p]/((d_.) + (e_.)*(x_.)), x_Symbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

Rubi steps

$$\begin{aligned}
\int x \csc^{-1}(a + bx)^3 dx &= -\frac{\text{Subst}(\int x^3 \cot(x) \csc(x)(-a + \csc(x)) dx, x, \csc^{-1}(a + bx))}{b^2} \\
&= \frac{1}{2} x^2 \csc^{-1}(a + bx)^3 - \frac{3\text{Subst}(\int x^2(-a + \csc(x))^2 dx, x, \csc^{-1}(a + bx))}{2b^2} \\
&= \frac{1}{2} x^2 \csc^{-1}(a + bx)^3 - \frac{3\text{Subst}(\int (a^2x^2 - 2ax^2 \csc(x) + x^2 \csc^2(x)) dx, x, \csc^{-1}(a + bx))}{2b^2} \\
&= -\frac{a^2 \csc^{-1}(a + bx)^3}{2b^2} + \frac{1}{2} x^2 \csc^{-1}(a + bx)^3 - \frac{3\text{Subst}(\int x^2 \csc^2(x) dx, x, \csc^{-1}(a + bx))}{2b^2} \\
&= \frac{3(a + bx) \sqrt{1 - \frac{1}{(a + bx)^2}} \csc^{-1}(a + bx)^2}{2b^2} - \frac{a^2 \csc^{-1}(a + bx)^3}{2b^2} + \frac{1}{2} x^2 \csc^{-1}(a + bx)^3 \\
&= \frac{3i \csc^{-1}(a + bx)^2}{2b^2} + \frac{3(a + bx) \sqrt{1 - \frac{1}{(a + bx)^2}} \csc^{-1}(a + bx)^2}{2b^2} - \frac{a^2 \csc^{-1}(a + bx)}{2b^2} \\
&= \frac{3i \csc^{-1}(a + bx)^2}{2b^2} + \frac{3(a + bx) \sqrt{1 - \frac{1}{(a + bx)^2}} \csc^{-1}(a + bx)^2}{2b^2} - \frac{a^2 \csc^{-1}(a + bx)}{2b^2} \\
&= \frac{3i \csc^{-1}(a + bx)^2}{2b^2} + \frac{3(a + bx) \sqrt{1 - \frac{1}{(a + bx)^2}} \csc^{-1}(a + bx)^2}{2b^2} - \frac{a^2 \csc^{-1}(a + bx)}{2b^2} \\
&= \frac{3i \csc^{-1}(a + bx)^2}{2b^2} + \frac{3(a + bx) \sqrt{1 - \frac{1}{(a + bx)^2}} \csc^{-1}(a + bx)^2}{2b^2} - \frac{a^2 \csc^{-1}(a + bx)}{2b^2}
\end{aligned}$$

Mathematica [A]

time = 0.59, size = 314, normalized size = 1.19

$$\frac{2i \csc^{-1}(a + bx)^2 + 2a \sqrt{\frac{-1 + a^2 + 2abx + b^2x^2}{(a + bx)^2}} \csc^{-1}(a + bx)^2 + 2ba \sqrt{\frac{-1 + a^2 + 2abx + b^2x^2}{(a + bx)^2}} \csc^{-1}(a + bx)^2 - a^2 \csc^{-1}(a + bx)^2 + b^2x^2 \csc^{-1}(a + bx)^2 + 6a \csc^{-1}(a + bx) \log(1 - e^{i \pi c \operatorname{atan}(a/bx)}) - 6a \csc^{-1}(a + bx) \log(1 + e^{i \pi c \operatorname{atan}(a/bx)}) - 6i \csc^{-1}(a + bx) \log(1 - e^{i \pi c \operatorname{atan}(a/bx)}) + 12ia \csc^{-1}(a + bx) \operatorname{PolyLog}(2, e^{i \pi c \operatorname{atan}(a/bx)}) - 12ia \csc^{-1}(a + bx) \operatorname{PolyLog}(2, e^{-i \pi c \operatorname{atan}(a/bx)}) + 3i \operatorname{PolyLog}(2, e^{i \pi c \operatorname{atan}(a/bx)}) - 12i \operatorname{PolyLog}(2, e^{-i \pi c \operatorname{atan}(a/bx)}) + 12i \operatorname{PolyLog}(3, e^{i \pi c \operatorname{atan}(a/bx)}) + 12i \operatorname{PolyLog}(3, e^{-i \pi c \operatorname{atan}(a/bx)})}{2b^2}$$

Antiderivative was successfully verified.

[In] `Integrate[x*ArcCsc[a + b*x]^3, x]`

[Out]
$$\begin{aligned}
&((3*I)*\operatorname{ArcCsc}[a + b*x]^2 + 3*a*\operatorname{Sqrt}[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)]^2)*\operatorname{ArcCsc}[a + b*x]^2 + 3*b*x*\operatorname{Sqrt}[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)]^2*\operatorname{ArcCsc}[a + b*x]^2 - a^2*\operatorname{ArcCsc}[a + b*x]^3 + b^2*x^2*\operatorname{ArcCsc}[a + b*x]^3 + 6*a*\operatorname{ArcCsc}[a + b*x]^2*\operatorname{Log}[1 - E^{(I*\operatorname{ArcCsc}[a + b*x])}] - 6*a*\operatorname{ArcCsc}[a + b*x]^2*\operatorname{Log}[1 + E^{(I*\operatorname{ArcCsc}[a + b*x])}] - 6*\operatorname{ArcCsc}[a + b*x]*\operatorname{Log}[1 - E^{((2*I)*\operatorname{ArcCsc}[a + b*x])}] + (12*I)*a*\operatorname{ArcCsc}[a + b*x]*\operatorname{PolyLog}[2, -E^{(I*\operatorname{ArcCsc}[a + b*x])}]
\end{aligned}$$

$$] - (12*I)*a*ArcCsc[a + b*x]*PolyLog[2, E^(I*ArcCsc[a + b*x])] + (3*I)*PolyLog[2, E^((2*I)*ArcCsc[a + b*x])] - 12*a*PolyLog[3, -E^(I*ArcCsc[a + b*x])] + 12*a*PolyLog[3, E^(I*ArcCsc[a + b*x])])/(2*b^2)$$

Maple [A]

time = 6.13, size = 425, normalized size = 1.61

method	result
derivativedivides	$\frac{\arccsc(bx+a)^2 \left(2 \arccsc(bx+a) a (bx+a) - \arccsc(bx+a) (bx+a)^2 - 3 \sqrt{\frac{(bx+a)^2 - 1}{(bx+a)^2}} (bx+a) + 3i \right)}{2} - 3a \arccsc(bx+a)^2 \ln\left(1 + \frac{i}{bx+a}\right)$
default	$\frac{\arccsc(bx+a)^2 \left(2 \arccsc(bx+a) a (bx+a) - \arccsc(bx+a) (bx+a)^2 - 3 \sqrt{\frac{(bx+a)^2 - 1}{(bx+a)^2}} (bx+a) + 3i \right)}{2} - 3a \arccsc(bx+a)^2 \ln\left(1 + \frac{i}{bx+a}\right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x*arccsc(b*x+a)^3,x,method=_RETURNVERBOSE)

[Out]
$$\begin{aligned} & 1/b^2*(-1/2*arccsc(b*x+a)^2*(2*arccsc(b*x+a)*a*(b*x+a)-arccsc(b*x+a)*(b*x+a)^2-3*((b*x+a)^2-1)/(b*x+a)^2)^{(1/2)}*(b*x+a)+3*I)-3*a*arccsc(b*x+a)^2*ln(1+I/(b*x+a)+(1-1/(b*x+a)^2)^{(1/2)})+3*a*arccsc(b*x+a)^2*ln(1-I/(b*x+a)-(1-1/(b*x+a)^2)^{(1/2)})-3*arccsc(b*x+a)*ln(1+I/(b*x+a)+(1-1/(b*x+a)^2)^{(1/2)})+6*I*a*arccsc(b*x+a)*polylog(2,-I/(b*x+a)-(1-1/(b*x+a)^2)^{(1/2)})-6*a*polylog(3,-I/(b*x+a)-(1-1/(b*x+a)^2)^{(1/2)})-3*arccsc(b*x+a)*ln(1-I/(b*x+a)-(1-1/(b*x+a)^2)^{(1/2)})-6*I*a*arccsc(b*x+a)*polylog(2,I/(b*x+a)+(1-1/(b*x+a)^2)^{(1/2)})+6*a*polylog(3,I/(b*x+a)+(1-1/(b*x+a)^2)^{(1/2)})+3*I*arccsc(b*x+a)^2+3*I*polylog(2,-I/(b*x+a)-(1-1/(b*x+a)^2)^{(1/2)})+3*I*polylog(2,I/(b*x+a)+(1-1/(b*x+a)^2)^{(1/2)})) \end{aligned}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x*arccsc(b*x+a)^3,x, algorithm="maxima")

[Out]
$$\begin{aligned} & 1/2*x^2*arctan2(1, sqrt(b*x + a + 1)*sqrt(b*x + a - 1))^3 - 3/8*x^2*arctan2(1, sqrt(b*x + a + 1)*sqrt(b*x + a - 1))*log(b^2*x^2 + 2*a*b*x + a^2)^2 - i*integrate(3/8*(8*(b^3*x^4*arctan2(1, sqrt(b*x + a + 1)*sqrt(b*x + a - 1))) + 3*a*b^2*x^3*arctan2(1, sqrt(b*x + a + 1)*sqrt(b*x + a - 1))) + (3*a^2*arctan2(1, sqrt(b*x + a + 1)*sqrt(b*x + a - 1))) - arctan2(1, sqrt(b*x + a + 1)*sqrt(b*x + a - 1)))*b*x^2 + (a^3*arctan2(1, sqrt(b*x + a + 1)*sqrt(b*x + a - 1))) - a*arctan2(1, sqrt(b*x + a + 1)*sqrt(b*x + a - 1)))*x)*log(b*x + a)^2 \end{aligned}$$

```

- (4*b*x^2*arctan2(1, sqrt(b*x + a + 1)*sqrt(b*x + a - 1))^2 - b*x^2*log(b^
2*x^2 + 2*a*b*x + a^2)^2)*sqrt(b*x + a + 1)*sqrt(b*x + a - 1) - 4*(b^3*x^4*
arctan2(1, sqrt(b*x + a + 1)*sqrt(b*x + a - 1)) + 2*a*b^2*x^3*arctan2(1, sq
rt(b*x + a + 1)*sqrt(b*x + a - 1)) + (a^2*arctan2(1, sqrt(b*x + a + 1)*sqrt(
b*x + a - 1)) - arctan2(1, sqrt(b*x + a + 1)*sqrt(b*x + a - 1)))*b*x^2 + 2*
(b^3*x^4*arctan2(1, sqrt(b*x + a + 1)*sqrt(b*x + a - 1)) + 3*a*b^2*x^3*arc
tan2(1, sqrt(b*x + a + 1)*sqrt(b*x + a - 1)) + (3*a^2*arctan2(1, sqrt(b*x +
a + 1)*sqrt(b*x + a - 1)) - arctan2(1, sqrt(b*x + a + 1)*sqrt(b*x + a - 1))
)*b*x^2 + (a^3*arctan2(1, sqrt(b*x + a + 1)*sqrt(b*x + a - 1)) - a*arctan2(
1, sqrt(b*x + a + 1)*sqrt(b*x + a - 1)))*x)*log(b*x + a))*log(b^2*x^2 + 2*
a*b*x + a^2))/(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2 - 1)*b*x - a), x)

```

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x*arccsc(b*x+a)^3,x, algorithm="fricas")

[Out] integral(x*arccsc(b*x + a)^3, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x \operatorname{acsc}^3(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x*acsc(b*x+a)**3,x)

[Out] Integral(x*acsc(a + b*x)**3, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x*arccsc(b*x+a)^3,x, algorithm="giac")

[Out] integrate(x*arccsc(b*x + a)^3, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int x \operatorname{asin}\left(\frac{1}{a + bx}\right)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\int x \cdot \text{asin}(1/(a + b \cdot x))^3 dx$
[Out] $\int x \cdot \text{asin}(1/(a + b \cdot x))^3 dx$, x

$$3.35 \quad \int \csc^{-1}(a + bx)^3 dx$$

Optimal. Leaf size=140

$$\frac{(a + bx) \csc^{-1}(a + bx)^3}{b} + \frac{6 \csc^{-1}(a + bx)^2 \tanh^{-1}\left(e^{i \csc^{-1}(a + bx)}\right)}{b} - \frac{6 i \csc^{-1}(a + bx) \text{PolyLog}\left(2, -e^{i \csc^{-1}(a + bx)}\right)}{b}$$

[Out] $(b*x+a)*\text{arccsc}(b*x+a)^3/b + 6*\text{arccsc}(b*x+a)^2*\text{arctanh}(I/(b*x+a)+(1-1/(b*x+a)^2)^{(1/2)})/b - 6*I*\text{arccsc}(b*x+a)*\text{polylog}(2, -I/(b*x+a)-(1-1/(b*x+a)^2)^{(1/2)})/b + 6*I*\text{arccsc}(b*x+a)*\text{polylog}(2, I/(b*x+a)+(1-1/(b*x+a)^2)^{(1/2)})/b + 6*\text{polylog}(3, -I/(b*x+a)-(1-1/(b*x+a)^2)^{(1/2)})/b - 6*\text{polylog}(3, I/(b*x+a)+(1-1/(b*x+a)^2)^{(1/2)})/b$

Rubi [A]

time = 0.08, antiderivative size = 140, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 7, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.875$, Rules used = {5361, 5325, 3843, 4268, 2611, 2320, 6724}

$$-\frac{6 i \csc^{-1}(a + bx) \text{Li}_2\left(-e^{i \csc^{-1}(a + bx)}\right)}{b} + \frac{6 i \csc^{-1}(a + bx) \text{Li}_2\left(e^{i \csc^{-1}(a + bx)}\right)}{b} + \frac{6 \text{Li}_3\left(-e^{i \csc^{-1}(a + bx)}\right)}{b} - \frac{6 \text{Li}_3\left(e^{i \csc^{-1}(a + bx)}\right)}{b} + \frac{(a + bx) \csc^{-1}(a + bx)^3}{b} + \frac{6 \csc^{-1}(a + bx)^2 \tanh^{-1}\left(e^{i \csc^{-1}(a + bx)}\right)}{b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{ArcCsc}[a + b*x]^3, x]$

[Out] $((a + b*x)*\text{ArcCsc}[a + b*x]^3)/b + (6*\text{ArcCsc}[a + b*x]^2*\text{ArcTanh}[E^(\text{I}*\text{ArcCsc}[a + b*x])])/b - ((6*\text{I})*\text{ArcCsc}[a + b*x]*\text{PolyLog}[2, -E^(\text{I}*\text{ArcCsc}[a + b*x])])/b + ((6*\text{I})*\text{ArcCsc}[a + b*x]*\text{PolyLog}[2, E^(\text{I}*\text{ArcCsc}[a + b*x])])/b + (6*\text{PolyLo}g[3, -E^(\text{I}*\text{ArcCsc}[a + b*x])])/b - (6*\text{PolyLog}[3, E^(\text{I}*\text{ArcCsc}[a + b*x])])/b$

Rule 2320

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*(a_.) + (b_.)*x))*(F_)[v_]] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2611

```
Int[Log[1 + (e_)*(F_)^((c_)*(a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g_.)*(x_.))^(m_.), x_Symbol] :> Simp[((-f + g*x)^m)*(PolyLog[2, (-e)*(F^((c*(a + b*x))))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^((c*(a + b*x))))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]
```

Rule 3843

```
Int[Cot[(a_.) + (b_ .)*(x_)^(n_.)]^(q_.)*Csc[(a_.) + (b_ .)*(x_)^(n_.)]^(p_.)
*(x_)^(m_.), x_Symbol] :> Simp[(-x^(m - n + 1))*(Csc[a + b*x^n]^p/(b*n*p)),
x] + Dist[(m - n + 1)/(b*n*p), Int[x^(m - n)*Csc[a + b*x^n]^p, x], x]; FreeQ[{a, b, p}, x] && IntegerQ[n] && GeQ[m, n] && EqQ[q, 1]
```

Rule 4268

```
Int[csc[(e_.) + (f_ .)*(x_)]*((c_.) + (d_ .)*(x_))^(m_.), x_Symbol] :> Simp[-2*(c + d*x)^m*(ArcTanh[E^(I*(e + f*x))/f], x] + (-Dist[d*(m/f), Int[(c + d
*x)^(m - 1)*Log[1 - E^(I*(e + f*x))], x], x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Log[1 + E^(I*(e + f*x))], x], x]) /; FreeQ[{c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 5325

```
Int[((a_.) + ArcCsc[(c_.)*(x_)]*(b_ .))^n, x_Symbol] :> Dist[-c^(-1), Subst[Int[(a + b*x)^n*Csc[x]*Cot[x], x], x, ArcCsc[c*x]], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[n, 0]
```

Rule 5361

```
Int[((a_.) + ArcCsc[(c_.) + (d_ .)*(x_)]*(b_ .))^p, x_Symbol] :> Dist[1/d, Subst[Int[(a + b*ArcCsc[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[p, 0]
```

Rule 6724

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_ .)*(x_))^(p_.)]/((d_.) + (e_ .)*(x_)), x_Symbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

Rubi steps

$$\begin{aligned}
\int \csc^{-1}(a+bx)^3 dx &= \frac{\text{Subst}(\int \csc^{-1}(x)^3 dx, x, a+bx)}{b} \\
&= -\frac{\text{Subst}(\int x^3 \cot(x) \csc(x) dx, x, \csc^{-1}(a+bx))}{b} \\
&= \frac{(a+bx) \csc^{-1}(a+bx)^3}{b} - \frac{3\text{Subst}(\int x^2 \csc(x) dx, x, \csc^{-1}(a+bx))}{b} \\
&= \frac{(a+bx) \csc^{-1}(a+bx)^3}{b} + \frac{6 \csc^{-1}(a+bx)^2 \tanh^{-1}(e^{i \csc^{-1}(a+bx)})}{b} + \frac{6\text{Subst}(\int x \log} \\
&= \frac{(a+bx) \csc^{-1}(a+bx)^3}{b} + \frac{6 \csc^{-1}(a+bx)^2 \tanh^{-1}(e^{i \csc^{-1}(a+bx)})}{b} - \frac{6i \csc^{-1}(a+bx)}{b} \\
&= \frac{(a+bx) \csc^{-1}(a+bx)^3}{b} + \frac{6 \csc^{-1}(a+bx)^2 \tanh^{-1}(e^{i \csc^{-1}(a+bx)})}{b} - \frac{6i \csc^{-1}(a+bx)}{b} \\
&= \frac{(a+bx) \csc^{-1}(a+bx)^3}{b} + \frac{6 \csc^{-1}(a+bx)^2 \tanh^{-1}(e^{i \csc^{-1}(a+bx)})}{b} - \frac{6i \csc^{-1}(a+bx)}{b}
\end{aligned}$$

Mathematica [A]

time = 0.08, size = 162, normalized size = 1.16

$$\frac{a \csc^{-1}(a+bx)^3 + bx \csc^{-1}(a+bx)^3 - 3 \csc^{-1}(a+bx)^2 \log(1 - e^{i \csc^{-1}(a+bx)}) + 3 \csc^{-1}(a+bx)^2 \log(1 + e^{i \csc^{-1}(a+bx)}) - 6i \csc^{-1}(a+bx) \text{PolyLog}(2, -e^{i \csc^{-1}(a+bx)}) + 6i \csc^{-1}(a+bx) \text{PolyLog}(2, e^{i \csc^{-1}(a+bx)}) + 6 \text{PolyLog}(3, -e^{i \csc^{-1}(a+bx)}) - 6 \text{PolyLog}(3, e^{i \csc^{-1}(a+bx)})}{b}$$

Antiderivative was successfully verified.

[In] Integrate[ArcCsc[a + b*x]^3, x]

[Out]
$$\begin{aligned}
&(a \text{ArcCsc}[a+b x]^3 + b x \text{ArcCsc}[a+b x]^3 - 3 \text{ArcCsc}[a+b x]^2 \text{Log}[1 - E^{\text{ArcCsc}[a+b x]}]) + 3 \text{ArcCsc}[a+b x]^2 \text{Log}[1 + E^{\text{ArcCsc}[a+b x]}] \\
&- (6 I) \text{ArcCsc}[a+b x] \text{PolyLog}[2, -E^{\text{ArcCsc}[a+b x]}] + (6 I) \text{ArcCsc}[a+b x] \text{PolyLog}[2, E^{\text{ArcCsc}[a+b x]}] + 6 \text{PolyLog}[3, -E^{\text{ArcCsc}[a+b x]}] - 6 \text{PolyLog}[3, E^{\text{ArcCsc}[a+b x]}])/b
\end{aligned}$$

Maple [A]

time = 1.15, size = 224, normalized size = 1.60

method	result
derivative divides	$ \text{arccsc}(bx+a)^3(bx+a) - 3\text{arccsc}(bx+a)^2 \ln\left(1 - \frac{i}{bx+a} - \sqrt{1 - \frac{1}{(bx+a)^2}}\right) + 6i \text{arccsc}(bx+a) \text{polylog}\left(2, \frac{i}{bx+a} + \sqrt{1 - \frac{1}{(bx+a)^2}}\right) $
default	$ \text{arccsc}(bx+a)^3(bx+a) - 3\text{arccsc}(bx+a)^2 \ln\left(1 - \frac{i}{bx+a} - \sqrt{1 - \frac{1}{(bx+a)^2}}\right) + 6i \text{arccsc}(bx+a) \text{polylog}\left(2, \frac{i}{bx+a} + \sqrt{1 - \frac{1}{(bx+a)^2}}\right) $

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(arccsc(b*x+a)^3,x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{b} \left(\text{arccsc}(b*x+a)^3 (b*x+a) - 3 \text{arccsc}(b*x+a)^2 \ln\left(\frac{1-I/(b*x+a)-(1-1/(b*x+a)^2)^{(1/2)}}{2}\right) + 6*I \text{arccsc}(b*x+a) \text{polylog}(2, I/(b*x+a) + (1-1/(b*x+a)^2)^{(1/2)}) - 6 \text{polylog}(3, I/(b*x+a) + (1-1/(b*x+a)^2)^{(1/2)}) + 3 \text{arccsc}(b*x+a)^2 \ln\left(\frac{1+I/(b*x+a)+(1-1/(b*x+a)^2)^{(1/2)}}{2}\right) - 6*I \text{arccsc}(b*x+a) \text{polylog}(2, -I/(b*x+a) - (1-1/(b*x+a)^2)^{(1/2)}) + 6 \text{polylog}(3, -I/(b*x+a) - (1-1/(b*x+a)^2)^{(1/2)}) \right)$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(b*x+a)^3,x, algorithm="maxima")`

[Out] $x \arctan^2(1, \sqrt{b*x + a + 1}) \sqrt{b*x + a - 1})^3 - \frac{3}{4} x \arctan^2(1, \sqrt{b*x + a + 1}) \sqrt{b*x + a - 1}) \log(b^2*x^2 + 2*a*b*x + a^2)^2 - \text{integrate}\left(\frac{3}{4} (4*(b^3*x^3 \arctan^2(1, \sqrt{b*x + a + 1}) \sqrt{b*x + a - 1}) + 3*a*b^2*x^2 \arctan^2(1, \sqrt{b*x + a + 1}) \sqrt{b*x + a - 1}) + a^3 \arctan^2(1, \sqrt{b*x + a + 1}) \sqrt{b*x + a - 1}) + (3*a^2 \arctan^2(1, \sqrt{b*x + a + 1}) \sqrt{b*x + a - 1}) - \arctan^2(1, \sqrt{b*x + a + 1}) \sqrt{b*x + a - 1}) * b*x - a \arctan^2(1, \sqrt{b*x + a + 1}) \sqrt{b*x + a - 1}) * \log(b*x + a)^2 - (4*b*x \arctan^2(1, \sqrt{b*x + a + 1}) \sqrt{b*x + a - 1})^2 - b*x \log(b^2*x^2 + 2*a*b*x + a^2)^2 * \sqrt{b*x + a + 1} \sqrt{b*x + a - 1} - 4*(b^3*x^3 \arctan^2(1, \sqrt{b*x + a + 1}) \sqrt{b*x + a - 1}) + 2*a*b^2*x^2 \arctan^2(1, \sqrt{b*x + a + 1}) \sqrt{b*x + a - 1}) + (a^2 \arctan^2(1, \sqrt{b*x + a + 1}) \sqrt{b*x + a - 1}) - a \arctan^2(1, \sqrt{b*x + a + 1}) \sqrt{b*x + a - 1}) * b*x + (b^3*x^3 \arctan^2(1, \sqrt{b*x + a + 1}) \sqrt{b*x + a - 1}) + 3*a*b^2*x^2 \arctan^2(1, \sqrt{b*x + a + 1}) \sqrt{b*x + a - 1}) + a^3 \arctan^2(1, \sqrt{b*x + a + 1}) \sqrt{b*x + a - 1}) + (3*a^2 \arctan^2(1, \sqrt{b*x + a + 1}) \sqrt{b*x + a - 1}) - \arctan^2(1, \sqrt{b*x + a + 1}) \sqrt{b*x + a - 1}) * b*x - a \arctan^2(1, \sqrt{b*x + a + 1}) \sqrt{b*x + a - 1}) * \log(b*x + a) * \log(b^2*x^2 + 2*a*b*x + a^2)) / (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2 - 1)*b*x - a), x\right)$

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(b*x+a)^3,x, algorithm="fricas")`

[Out] integral(arccsc(b*x + a)^3, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \operatorname{acsc}^3(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(acsc(b*x+a)**3,x)

[Out] Integral(acsc(a + b*x)**3, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(arccsc(b*x+a)^3,x, algorithm="giac")

[Out] integrate(arccsc(b*x + a)^3, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \operatorname{asin}\left(\frac{1}{a + b x}\right)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(asin(1/(a + b*x))^3,x)

[Out] int(asin(1/(a + b*x))^3, x)

$$\text{3.36} \quad \int \frac{\csc^{-1}(a+bx)^3}{x} dx$$

Optimal. Leaf size=448

$$\csc^{-1}(a+bx)^3 \log \left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{1 - \sqrt{1 - a^2}} \right) + \csc^{-1}(a+bx)^3 \log \left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{1 + \sqrt{1 - a^2}} \right) - \csc^{-1}(a+bx)^3 \log \left(1 - e^{2i \csc^{-1}(a+bx)} \right)$$

[Out] $-\operatorname{arccsc}(b*x+a)^3 \ln(1-(I/(b*x+a)+(1-1/(b*x+a)^2)^{(1/2)})^2)+\operatorname{arccsc}(b*x+a)^3 \ln(1+I*a*(I/(b*x+a)+(1-1/(b*x+a)^2)^{(1/2)})/(1-(-a^2+1)^{(1/2)}))+\operatorname{arccsc}(b*x+a)^3 \ln(1+I*a*(I/(b*x+a)+(1-1/(b*x+a)^2)^{(1/2)})/(1+(-a^2+1)^{(1/2)}))+3/2*I*\operatorname{arccsc}(b*x+a)^2*\operatorname{polylog}(2,(I/(b*x+a)+(1-1/(b*x+a)^2)^{(1/2)})^2)-3*I*\operatorname{arccsc}(b*x+a)^2*\operatorname{polylog}(2,-I*a*(I/(b*x+a)+(1-1/(b*x+a)^2)^{(1/2)})/(1-(-a^2+1)^{(1/2)}))-3*I*\operatorname{arccsc}(b*x+a)^2*\operatorname{polylog}(2,-I*a*(I/(b*x+a)+(1-1/(b*x+a)^2)^{(1/2)})/(1+(-a^2+1)^{(1/2)}))-3/2*\operatorname{arccsc}(b*x+a)*\operatorname{polylog}(3,(I/(b*x+a)+(1-1/(b*x+a)^2)^{(1/2)})^2)+6*I*\operatorname{arccsc}(b*x+a)*\operatorname{polylog}(3,-I*a*(I/(b*x+a)+(1-1/(b*x+a)^2)^{(1/2)})/(1-(-a^2+1)^{(1/2)}))+6*\operatorname{arccsc}(b*x+a)*\operatorname{polylog}(3,-I*a*(I/(b*x+a)+(1-1/(b*x+a)^2)^{(1/2)})/(1+(-a^2+1)^{(1/2)}))-3/4*I*\operatorname{polylog}(4,(I/(b*x+a)+(1-1/(b*x+a)^2)^{(1/2)})^2)+6*I*\operatorname{polylog}(4,-I*a*(I/(b*x+a)+(1-1/(b*x+a)^2)^{(1/2)})/(1-(-a^2+1)^{(1/2)}))+6*I*\operatorname{polylog}(4,-I*a*(I/(b*x+a)+(1-1/(b*x+a)^2)^{(1/2)})/(1+(-a^2+1)^{(1/2)}))$

Rubi [A]

time = 0.39, antiderivative size = 448, normalized size of antiderivative = 1.00, number of steps used = 20, number of rules used = 10, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {5367, 4648, 4625, 3798, 2221, 2611, 6744, 2320, 6724, 4615}

$$-3 \operatorname{cn}^{-1}(a+bx)^2 I_2 \left(\frac{\operatorname{izn}(\operatorname{cn}^{-1}(a+bx))}{1-\sqrt{1-a^2}} \right) - 3 i \operatorname{cn}^{-1}(a+bx)^2 I_3 \left(\frac{\operatorname{izn}(\operatorname{cn}^{-1}(a+bx))}{\sqrt{1-a^2}} \right) + 6 \operatorname{cn}^{-1}(a+bx) I_4 \left(\frac{\operatorname{izn}(\operatorname{cn}^{-1}(a+bx))}{1-\sqrt{1-a^2}} \right) + 6 i \operatorname{cn}^{-1}(a+bx) I_5 \left(\frac{\operatorname{izn}(\operatorname{cn}^{-1}(a+bx))}{\sqrt{1-a^2}} \right) + 6 i \operatorname{dn} I_1 \left(\frac{\operatorname{izn}(\operatorname{cn}^{-1}(a+bx))}{\sqrt{1-a^2}} \right) + \operatorname{cn}^{-1}(a+bx)^2 \log \left(1+\frac{\operatorname{izn}(\operatorname{cn}^{-1}(a+bx))}{\sqrt{1-a^2}} \right) + \operatorname{cn}^{-1}(a+bx)^2 \log \left(1+\frac{\operatorname{izn}(\operatorname{cn}^{-1}(a+bx))}{\sqrt{1-a^2}} \right) + \frac{3}{2} \operatorname{cn}^{-1}(a+bx)^2 I_2 \left(\operatorname{cn}(\operatorname{cn}^{-1}(a+bx)) \right) - \frac{3}{2} \operatorname{cn}^{-1}(a+bx)^2 \log \left(1-\frac{\operatorname{izn}(\operatorname{cn}^{-1}(a+bx))}{\sqrt{1-a^2}} \right) - \operatorname{cn}^{-1}(a+bx)^2 \log \left(1-\frac{\operatorname{izn}(\operatorname{cn}^{-1}(a+bx))}{\sqrt{1-a^2}} \right)$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[\operatorname{ArcCsc}[a+b*x]^3/x, x]$

[Out] $\operatorname{ArcCsc}[a+b*x]^3 \operatorname{Log}[1+(I*a*E^(\operatorname{ArcCsc}[a+b*x]))/(1-\operatorname{Sqrt}[1-a^2])]+\operatorname{ArcCsc}[a+b*x]^3 \operatorname{Log}[1+(I*a*E^(\operatorname{ArcCsc}[a+b*x]))/(1+\operatorname{Sqrt}[1-a^2])]-\operatorname{ArcCsc}[a+b*x]^3 \operatorname{Log}[1-E^((2*I)*\operatorname{ArcCsc}[a+b*x])]-(3*I)*\operatorname{ArcCsc}[a+b*x]^2 \operatorname{PolyLog}[2, ((-I)*a*E^(\operatorname{ArcCsc}[a+b*x]))/(1-\operatorname{Sqrt}[1-a^2])]- (3*I)*\operatorname{ArcCsc}[a+b*x]^2 \operatorname{PolyLog}[2, ((-I)*a*E^(\operatorname{ArcCsc}[a+b*x]))/(1+\operatorname{Sqrt}[1-a^2])]+((3*I)/2)*\operatorname{ArcCsc}[a+b*x]^2 \operatorname{PolyLog}[2, E^((2*I)*\operatorname{ArcCsc}[a+b*x])]+6*\operatorname{ArcCsc}[a+b*x] \operatorname{PolyLog}[3, ((-I)*a*E^(\operatorname{ArcCsc}[a+b*x]))/(1-\operatorname{Sqrt}[1-a^2])]+6*\operatorname{ArcCsc}[a+b*x] \operatorname{PolyLog}[3, ((-I)*a*E^(\operatorname{ArcCsc}[a+b*x]))/(1+\operatorname{Sqrt}[1-a^2])]- (3*\operatorname{ArcCsc}[a+b*x] \operatorname{PolyLog}[3, E^((2*I)*\operatorname{ArcCsc}[a+b*x]))]/2+(6*I)*\operatorname{PolyLog}[4, ((-I)*a*E^(\operatorname{ArcCsc}[a+b*x]))/(1-\operatorname{Sqrt}[1-a^2])]+(6*I)*\operatorname{PolyLog}[4, ((-I)*a*E^(\operatorname{ArcCsc}[a+b*x]))/(1+\operatorname{Sqrt}[1-a^2])]-((3*I)/4)*\operatorname{PolyLog}[4, E^((2*I)*\operatorname{ArcCsc}[a+b*x])]$

Rule 2221

```

Int[((F_)^((g_.)*(e_.) + (f_.)*(x_.)))^(n_.)*((c_.) + (d_.)*(x_.))^((m_.))/((a_.) + (b_.)*(F_)^((g_.)*(e_.) + (f_.)*(x_.)))^(n_.)), x_Symbol] :> Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

```

Rule 2320

```

Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_.))^(m_.)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_[v_]) /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]

```

Rule 2611

```

Int[Log[1 + (e_.)*((F_)^((c_.)*(a_.) + (b_.)*(x_.)))^(n_.))*((f_.) + (g_.)*(x_.))^((m_.)), x_Symbol] :> Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]

```

Rule 3798

```

Int[((c_.) + (d_.)*(x_.))^((m_.))*tan[(e_.) + Pi*(k_.) + (f_.)*(x_.)], x_Symbol] :> Simp[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Dist[2*I, Int[(c + d*x)^m *E^(2*I*k*Pi)*(E^(2*I*(e + f*x))/(1 + E^(2*I*k*Pi)*E^(2*I*(e + f*x)))), x], x] /; FreeQ[{c, d, e, f}, x] && IntegerQ[4*k] && IGtQ[m, 0]

```

Rule 4615

```

Int[(Cos[(c_.) + (d_.)*(x_.)]*((e_.) + (f_.)*(x_.))^((m_.)))/((a_.) + (b_.)*Sin[(c_.) + (d_.)*(x_.)]), x_Symbol] :> Simp[(-I)*((e + f*x)^(m + 1)/(b*f*(m + 1))), x] + (Int[(e + f*x)^m*(E^(I*(c + d*x))/(a - Rt[a^2 - b^2, 2] - I*b*E^(I*(c + d*x)))), x] + Int[(e + f*x)^m*(E^(I*(c + d*x))/(a + Rt[a^2 - b^2, 2] - I*b*E^(I*(c + d*x)))), x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && PosQ[a^2 - b^2]

```

Rule 4625

```

Int[(Cot[(c_.) + (d_.)*(x_.)]^((n_.))*((e_.) + (f_.)*(x_.))^((m_.)))/((a_.) + (b_.)*Sin[(c_.) + (d_.)*(x_.)]), x_Symbol] :> Dist[1/a, Int[(e + f*x)^m*Cot[c + d*x]^n, x], x] - Dist[b/a, Int[(e + f*x)^m*Cos[c + d*x]*(Cot[c + d*x]^(n - 1)/(a + b*Sin[c + d*x])), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && IGtQ[n, 0]

```

Rule 4648

```
Int[((e_.) + (f_)*(x_))^(m_)*(F_)[(c_.) + (d_)*(x_)]^(n_)*(G_)[(c_.) +
(d_)*(x_)]^(p_))/(Csc[(c_.) + (d_)*(x_)]*(b_) + (a_)), x_Symbol] :> In
t[(e + f*x)^m*Sin[c + d*x]*F[c + d*x]^n*(G[c + d*x]^p/(b + a*Sin[c + d*x]))),
x] /; FreeQ[{a, b, c, d, e, f}, x] && TrigQ[F] && TrigQ[G] && IntegersQ[m,
n, p]
```

Rule 5367

```
Int[((a_.) + ArcCsc[(c_) + (d_)*(x_)]*(b_))^(p_)*((e_.) + (f_)*(x_))^m,
x_Symbol] :> Dist[-(d^(m + 1))^{(-1)}, Subst[Int[(a + b*x)^p*Csc[x]*Cot[
x]*(d*e - c*f + f*Csc[x])]^m, x], x, ArcCsc[c + d*x]], x] /; FreeQ[{a, b, c,
d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Rule 6724

```
Int[PolyLog[n_, (c_)*((a_.) + (b_)*(x_))^p]/((d_.) + (e_)*(x_)), x_Symbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d,
e, n, p}, x] && EqQ[b*d, a*e]
```

Rule 6744

```
Int[((e_.) + (f_)*(x_))^(m_)*PolyLog[n_, (d_)*((F_)^((c_)*((a_.) + (b_.
)*(x_))))^p], x_Symbol] :> Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Dist[f*(m/(b*c*p*Log[F])), Int[(e + f*x)^
(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c,
d, e, f, n, p}, x] && GtQ[m, 0]
```

Rubi steps

$$\begin{aligned}
\int \frac{\csc^{-1}(a+bx)^3}{x} dx &= -\text{Subst}\left(\int \frac{x^3 \cot(x) \csc(x)}{-a + \csc(x)} dx, x, \csc^{-1}(a+bx)\right) \\
&= -\text{Subst}\left(\int \frac{x^3 \cot(x)}{1 - a \sin(x)} dx, x, \csc^{-1}(a+bx)\right) \\
&= -\left(a \text{Subst}\left(\int \frac{x^3 \cos(x)}{1 - a \sin(x)} dx, x, \csc^{-1}(a+bx)\right)\right) - \text{Subst}\left(\int x^3 \cot(x) dx, x, \csc^{-1}(a+bx)\right) \\
&= 2i \text{Subst}\left(\int \frac{e^{2ix} x^3}{1 - e^{2ix}} dx, x, \csc^{-1}(a+bx)\right) - a \text{Subst}\left(\int \frac{e^{ix} x^3}{1 - \sqrt{1 - a^2} + iae^{ix}} dx, x, \csc^{-1}(a+bx)\right) \\
&= \csc^{-1}(a+bx)^3 \log\left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{1 - \sqrt{1 - a^2}}\right) + \csc^{-1}(a+bx)^3 \log\left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{1 + \sqrt{1 - a^2}}\right) \\
&= \csc^{-1}(a+bx)^3 \log\left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{1 - \sqrt{1 - a^2}}\right) + \csc^{-1}(a+bx)^3 \log\left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{1 + \sqrt{1 - a^2}}\right) \\
&= \csc^{-1}(a+bx)^3 \log\left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{1 - \sqrt{1 - a^2}}\right) + \csc^{-1}(a+bx)^3 \log\left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{1 + \sqrt{1 - a^2}}\right) \\
&= \csc^{-1}(a+bx)^3 \log\left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{1 - \sqrt{1 - a^2}}\right) + \csc^{-1}(a+bx)^3 \log\left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{1 + \sqrt{1 - a^2}}\right) \\
&= \csc^{-1}(a+bx)^3 \log\left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{1 - \sqrt{1 - a^2}}\right) + \csc^{-1}(a+bx)^3 \log\left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{1 + \sqrt{1 - a^2}}\right)
\end{aligned}$$

Mathematica [A]

time = 0.22, size = 554, normalized size = 1.24

```
<--> (1 - E^(((-I)*ArcCsc[a + b*x])^4))^(1/4) + ArcCsc[a + b*x]^3*Log[1 - E^(((-I)*ArcCsc[a + b*x]))^4] - ArcCsc[a + b*x]^3*Log[1 + E^((I*ArcCsc[a + b*x]))^4] + ArcCsc[a + b*x]^3*Log[1 - (I*a*E^((I*ArcCsc[a + b*x])))/(-1 + Sqrt[1 - a^2])] + ArcCsc[a + b*x]^3*Log[1 + (I*a*E^((I*ArcCsc[a + b*x])))/(1 + Sqrt[1 - a^2])] - (3*I)*ArcCsc[a + b*x]^2*PolyLog[2, E^((-I)*ArcCsc[a + b*x])] + (3*I)*ArcCsc[a + b*x]^2*PolyLog[2, -E^((I*ArcCsc[a + b*x]))] - (3*I)*ArcCsc[a + b*x]^2*PolyLog[2, (I*a*E^((I*ArcCsc[a + b*x])))/(-1 + Sqrt[1 - a^2])] - (3*I)*ArcCsc[a + b*x]^2*PolyLog[2, ((-I)*a*E^((I*ArcCsc[a + b*x])))/(1 + Sqrt[1 - a^2])] - 6
```

Antiderivative was successfully verified.

[In] Integrate[ArcCsc[a + b*x]^3/x, x]

[Out] $(I/8)*Pi^4 - (I/4)*ArcCsc[a + b*x]^4 - ArcCsc[a + b*x]^3*Log[1 - E^((-I)*ArcCsc[a + b*x])] - ArcCsc[a + b*x]^3*Log[1 + E^((I*ArcCsc[a + b*x]))] + ArcCsc[a + b*x]^3*Log[1 - (I*a*E^((I*ArcCsc[a + b*x])))/(-1 + Sqrt[1 - a^2])] + ArcCsc[a + b*x]^3*Log[1 + (I*a*E^((I*ArcCsc[a + b*x])))/(1 + Sqrt[1 - a^2])] - (3*I)*ArcCsc[a + b*x]^2*PolyLog[2, E^((-I)*ArcCsc[a + b*x])] + (3*I)*ArcCsc[a + b*x]^2*PolyLog[2, -E^((I*ArcCsc[a + b*x]))] - (3*I)*ArcCsc[a + b*x]^2*PolyLog[2, (I*a*E^((I*ArcCsc[a + b*x])))/(-1 + Sqrt[1 - a^2])] - (3*I)*ArcCsc[a + b*x]^2*PolyLog[2, ((-I)*a*E^((I*ArcCsc[a + b*x])))/(1 + Sqrt[1 - a^2])] - 6$

```
*ArcCsc[a + b*x]*PolyLog[3, E^((-I)*ArcCsc[a + b*x])] - 6*ArcCsc[a + b*x]*PolyLog[3, -E^(I*ArcCsc[a + b*x])] + 6*ArcCsc[a + b*x]*PolyLog[3, (I*a*E^(I*ArcCsc[a + b*x]))/(-1 + Sqrt[1 - a^2])] + 6*ArcCsc[a + b*x]*PolyLog[3, ((-I)*a*E^(I*ArcCsc[a + b*x]))/(1 + Sqrt[1 - a^2])] + (6*I)*PolyLog[4, E^((-I)*ArcCsc[a + b*x])] - (6*I)*PolyLog[4, -E^(I*ArcCsc[a + b*x])] + (6*I)*PolyLog[4, (I*a*E^(I*ArcCsc[a + b*x]))/(-1 + Sqrt[1 - a^2])] + (6*I)*PolyLog[4, ((-I)*a*E^(I*ArcCsc[a + b*x]))/(1 + Sqrt[1 - a^2])]
```

Maple [F]

time = 2.53, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{arccsc}(bx + a)^3}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(arccsc(b*x+a)^3/x,x)

[Out] int(arccsc(b*x+a)^3/x,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(arccsc(b*x+a)^3/x,x, algorithm="maxima")

[Out] integrate(arccsc(b*x + a)^3/x, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(arccsc(b*x+a)^3/x,x, algorithm="fricas")

[Out] integral(arccsc(b*x + a)^3/x, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{acsc}^3(a + bx)}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(acsc(b*x+a)**3/x, x)`
[Out] `Integral(acsc(a + b*x)**3/x, x)`

Giac [F]
time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(b*x+a)^3/x, x, algorithm="giac")`
[Out] `integrate(arccsc(b*x + a)^3/x, x)`

Mupad [F]
time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\arcsin\left(\frac{1}{a+bx}\right)^3}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(asin(1/(a + b*x))^3/x, x)`
[Out] `int(asin(1/(a + b*x))^3/x, x)`

$$3.37 \quad \int \frac{\csc^{-1}(a+bx)^3}{x^2} dx$$

Optimal. Leaf size=378

$$\frac{b \csc^{-1}(a+bx)^3}{a} - \frac{\csc^{-1}(a+bx)^3}{x} - \frac{3ib \csc^{-1}(a+bx)^2 \log \left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} + \frac{3ib \csc^{-1}(a+bx)^2 \log \left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}}$$

[Out] $-b*\arccsc(b*x+a)^3/a - \arccsc(b*x+a)^3/x - 3*I*b*\arccsc(b*x+a)^2*\ln(1+I*a*(I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))/(1-(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2) + 3*I*b*\arccsc(b*x+a)^2*\ln(1+I*a*(I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))/(1+(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2) - 6*b*\arccsc(b*x+a)*\text{polylog}(2, -I*a*(I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))/(1-(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2) + 6*b*\arccsc(b*x+a)*\text{polylog}(2, -I*a*(I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))/(1+(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2) - 6*I*b*\text{polylog}(3, -I*a*(I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))/(1-(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2) + 6*I*b*\text{polylog}(3, -I*a*(I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))/(1+(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2) + 6*I*b*\text{polylog}(3, -I*a*(I/(b*x+a)+(1-1/(b*x+a)^2)^(1/2))/(1+(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)$

Rubi [A]

time = 0.42, antiderivative size = 378, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 9, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {5367, 4512, 4276, 3404, 2296, 2221, 2611, 2320, 6724}

$$-\frac{6b \csc^{-1}(a+bx) \text{Li}_2\left(-\frac{iae^{i \csc^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} + \frac{6b \csc^{-1}(a+bx) \text{Li}_2\left(-\frac{iae^{i \csc^{-1}(a+bx)}}{1-\sqrt{1-a^2+1}}\right)}{a\sqrt{1-a^2}} - \frac{6ib \text{Li}_3\left(-\frac{iae^{i \csc^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} + \frac{6ib \text{Li}_3\left(-\frac{iae^{i \csc^{-1}(a+bx)}}{1-\sqrt{1-a^2+1}}\right)}{a\sqrt{1-a^2}} - \frac{3ib \csc^{-1}(a+bx)^2 \log \left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} + \frac{3ib \csc^{-1}(a+bx)^2 \log \left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{1-\sqrt{1-a^2+1}}\right)}{a\sqrt{1-a^2}} - \frac{b \csc^{-1}(a+bx)^3}{a} - \frac{\csc^{-1}(a+bx)^3}{x}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{ArcCsc}[a+b*x]^3/x^2, x]$

[Out] $-((b*\text{ArcCsc}[a+b*x]^3)/a) - \text{ArcCsc}[a+b*x]^3/x - ((3*I)*b*\text{ArcCsc}[a+b*x]^2*\text{Log}[1+(I*a*E^(\text{ArcCsc}[a+b*x]))/(1-Sqrt[1-a^2])])/(a*Sqrt[1-a^2]) + ((3*I)*b*\text{ArcCsc}[a+b*x]^2*\text{Log}[1+(I*a*E^(\text{ArcCsc}[a+b*x]))/(1+Sqrt[1-a^2])])/(a*Sqrt[1-a^2]) - (6*b*\text{ArcCsc}[a+b*x]*\text{PolyLog}[2, ((-I)*a*E^(\text{ArcCsc}[a+b*x]))/(1-Sqrt[1-a^2])])/(a*Sqrt[1-a^2]) + (6*b*\text{ArcCsc}[a+b*x]*\text{PolyLog}[2, ((-I)*a*E^(\text{ArcCsc}[a+b*x]))/(1+Sqrt[1-a^2])])/(a*Sqrt[1-a^2]) - ((6*I)*b*\text{PolyLog}[3, ((-I)*a*E^(\text{ArcCsc}[a+b*x]))/(1-Sqrt[1-a^2])])/(a*Sqrt[1-a^2]) + ((6*I)*b*\text{PolyLog}[3, ((-I)*a*E^(\text{ArcCsc}[a+b*x]))/(1+Sqrt[1-a^2])])/(a*Sqrt[1-a^2])$

Rule 2221

```
Int[((F_)^((g_.)*(e_.) + (f_.)*(x_.)))^(n_.)*((c_.) + (d_.)*(x_.))^(m_.))/((a_) + (b_.)*(F_)^((g_.)*(e_.) + (f_.)*(x_.)))^(n_.)), x_Symbol] :> Simplify[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x]; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2296

```
Int[((F_)^(u_)*((f_.) + (g_.)*(x_.))^(m_.))/((a_.) + (b_.)*(F_)^(u_) + (c_.)
*(F_)^(v_)), x_Symbol] :> With[{q = Rt[b^2 - 4*a*c, 2]}, Dist[2*(c/q), Int[
(f + g*x)^m*(F^u/(b - q + 2*c*F^u)), x], x] - Dist[2*(c/q), Int[(f + g*x)^m
*(F^u/(b + q + 2*c*F^u)), x], x]] /; FreeQ[{F, a, b, c, f, g}, x] && EqQ[v,
2*u] && LinearQ[u, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[m, 0]
```

Rule 2320

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x]
, Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_.))^(m_)] /; FreeQ[
{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(
F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2611

```
Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)
*(x_.))^(m_.), x_Symbol] :> Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^m
- 1]*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e,
f, g, n}, x] && GtQ[m, 0]
```

Rule 3404

```
Int[((c_.) + (d_.)*(x_.))^(m_.)/((a_) + (b_.)*sin[(e_.) + (f_.)*(x_.)]), x_Sy
mbol] :> Dist[2, Int[(c + d*x)^m*(E^(I*(e + f*x))/(I*b + 2*a*E^(I*(e + f*x)
) - I*b*E^(2*I*(e + f*x)))), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[
a^2 - b^2, 0] && IGtQ[m, 0]
```

Rule 4276

```
Int[(csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(n_.)*((c_.) + (d_.)*(x_.))^(m_.)
, x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, 1/(Sin[e + f*x]^n/(b + a*Si
n[e + f*x])^n), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && ILtQ[n, 0] && IGt
Q[m, 0]
```

Rule 4512

```
Int[Cot[(c_.) + (d_.)*(x_.)]*Csc[(c_.) + (d_.)*(x_.)]*(Csc[(c_.) + (d_.)*(x_)]
]*(b_.) + (a_.))^(n_.)*((e_.) + (f_.)*(x_.))^(m_.), x_Symbol] :> Simp[(-(e +
f*x)^m)*((a + b*Csc[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[f*(m/(b*d*(n
+ 1))), Int[(e + f*x)^(m - 1)*(a + b*Csc[c + d*x])^(n + 1), x], x] /; Fre
eQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]
```

Rule 5367

```
Int[((a_.) + ArcCsc[(c_) + (d_)*(x_)]*(b_.))^(p_.)*((e_.) + (f_)*(x_))^m_, x_Symbol] :> Dist[-(d^(m + 1))^{-1}, Subst[Int[(a + b*x)^p*Csc[x]*Cot[x]*(d*e - c*f + f*Csc[x])^m, x], x, ArcCsc[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Rule 6724

```
Int[PolyLog[n_, (c_)*(a_.) + (b_)*(x_)]^(p_.)]/((d_.) + (e_)*(x_)), x_Symbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

Rubi steps

$$\begin{aligned}
\int \frac{\csc^{-1}(a+bx)^3}{x^2} dx &= - \left(b \text{Subst} \left(\int \frac{x^3 \cot(x) \csc(x)}{(-a + \csc(x))^2} dx, x, \csc^{-1}(a+bx) \right) \right) \\
&= - \frac{\csc^{-1}(a+bx)^3}{x} + (3b) \text{Subst} \left(\int \frac{x^2}{-a + \csc(x)} dx, x, \csc^{-1}(a+bx) \right) \\
&= - \frac{\csc^{-1}(a+bx)^3}{x} + (3b) \text{Subst} \left(\int \left(-\frac{x^2}{a} + \frac{x^2}{a(1-a \sin(x))} \right) dx, x, \csc^{-1}(a+bx) \right) \\
&= - \frac{b \csc^{-1}(a+bx)^3}{a} - \frac{\csc^{-1}(a+bx)^3}{x} + \frac{(3b) \text{Subst} \left(\int \frac{x^2}{1-a \sin(x)} dx, x, \csc^{-1}(a+bx) \right)}{a} \\
&= - \frac{b \csc^{-1}(a+bx)^3}{a} - \frac{\csc^{-1}(a+bx)^3}{x} + \frac{(6b) \text{Subst} \left(\int \frac{e^{ix} x^2}{-ia+2e^{ix}+iae^{2ix}} dx, x, \csc^{-1}(a+bx) \right)}{a} \\
&= - \frac{b \csc^{-1}(a+bx)^3}{a} - \frac{\csc^{-1}(a+bx)^3}{x} + \frac{(6ib) \text{Subst} \left(\int \frac{e^{ix} x^2}{2-2\sqrt{1-a^2}+2iae^{ix}} dx, x, \csc^{-1}(a+bx) \right)}{\sqrt{1-a^2}} \\
&= - \frac{b \csc^{-1}(a+bx)^3}{a} - \frac{\csc^{-1}(a+bx)^3}{x} - \frac{3ib \csc^{-1}(a+bx)^2 \log \left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right)}{a \sqrt{1-a^2}} + \\
&= - \frac{b \csc^{-1}(a+bx)^3}{a} - \frac{\csc^{-1}(a+bx)^3}{x} - \frac{3ib \csc^{-1}(a+bx)^2 \log \left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right)}{a \sqrt{1-a^2}} + \\
&= - \frac{b \csc^{-1}(a+bx)^3}{a} - \frac{\csc^{-1}(a+bx)^3}{x} - \frac{3ib \csc^{-1}(a+bx)^2 \log \left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right)}{a \sqrt{1-a^2}} + \\
&= - \frac{b \csc^{-1}(a+bx)^3}{a} - \frac{\csc^{-1}(a+bx)^3}{x} - \frac{3ib \csc^{-1}(a+bx)^2 \log \left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right)}{a \sqrt{1-a^2}} +
\end{aligned}$$

Mathematica [A]

time = 0.34, size = 289, normalized size = 0.76

$$\begin{aligned}
&\frac{(a+bx) \csc^{-1}(a+bx)^3}{x} + \frac{3ib \left(\csc^{-1}(a+bx)^2 \log \left(1 - \frac{iae^{i \csc^{-1}(a+bx)}}{-1+\sqrt{1-a^2}} \right) - \csc^{-1}(a+bx)^2 \log \left(1 + \frac{iae^{i \csc^{-1}(a+bx)}}{1+\sqrt{1-a^2}} \right) - 2i \csc^{-1}(a+bx) \text{PolyLog} \left(2, \frac{iae^{i \csc^{-1}(a+bx)}}{-1+\sqrt{1-a^2}} \right) + 2i \csc^{-1}(a+bx) \text{PolyLog} \left(2, \frac{iae^{i \csc^{-1}(a+bx)}}{1+\sqrt{1-a^2}} \right) + 2 \text{PolyLog} \left(3, \frac{iae^{i \csc^{-1}(a+bx)}}{-1+\sqrt{1-a^2}} \right) - 2 \text{PolyLog} \left(3, \frac{iae^{i \csc^{-1}(a+bx)}}{1+\sqrt{1-a^2}} \right) \right)}{\sqrt{1-a^2}}
\end{aligned}$$

Antiderivative was successfully verified.

[In] `Integrate[ArcCsc[a + b*x]^3/x^2, x]`

[Out] $-(((a+b*x)*ArcCsc[a+b*x]^3)/x + ((3*I)*b*(ArcCsc[a+b*x]^2*Log[1 - (I*a*E^(I*ArcCsc[a+b*x]))/(-1 + Sqrt[1 - a^2])]) - ArcCsc[a+b*x]^2*Log[1 +$

$$\begin{aligned} & \left(I * a * E^{\text{ArcCsc}[a + b*x]} \right) / (1 + \text{Sqrt}[1 - a^2]) - (2*I) * \text{ArcCsc}[a + b*x] * \text{PolyLog}[2, \left(I * a * E^{\text{ArcCsc}[a + b*x]} \right) / (-1 + \text{Sqrt}[1 - a^2])] + (2*I) * \text{ArcCsc}[a + b*x] * \text{PolyLog}[2, ((-I) * a * E^{\text{ArcCsc}[a + b*x]}) / (1 + \text{Sqrt}[1 - a^2])] + 2 * \text{PolyLog}[3, \left(I * a * E^{\text{ArcCsc}[a + b*x]} \right) / (-1 + \text{Sqrt}[1 - a^2])] - 2 * \text{PolyLog}[3, ((-I) * a * E^{\text{ArcCsc}[a + b*x]}) / (1 + \text{Sqrt}[1 - a^2])]) / \text{Sqrt}[1 - a^2] / a \end{aligned}$$
Maple [F]

time = 2.86, size = 0, normalized size = 0.00

$$\int \frac{\text{arccsc}(bx + a)^3}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(arccsc(b*x+a)^3/x^2,x)`
[Out] `int(arccsc(b*x+a)^3/x^2,x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(b*x+a)^3/x^2,x, algorithm="maxima")`
[Out]
$$\begin{aligned} & -1/4 * (4 * \text{arctan2}(1, \sqrt{b*x + a + 1}) * \sqrt{b*x + a - 1})^3 - 3 * \text{arctan2}(1, \sqrt{b*x + a + 1}) * \sqrt{b*x + a - 1} * \log(b^2*x^2 + 2*a*b*x + a^2)^2 - 4*x * \text{integrate}(-3/4 * (4 * (\sqrt{b^3*x^3} * \text{arctan2}(1, \sqrt{b*x + a + 1}) * \sqrt{b*x + a - 1})) + 3 * a * b^2 * x^2 * \text{arctan2}(1, \sqrt{b*x + a + 1}) * \sqrt{b*x + a - 1}) + a^3 * \text{arctan2}(1, \sqrt{b*x + a + 1}) * \sqrt{b*x + a - 1}) + (3*a^2 * \text{arctan2}(1, \sqrt{b*x + a + 1}) * \sqrt{b*x + a - 1}) - \text{arctan2}(1, \sqrt{b*x + a + 1}) * \sqrt{b*x + a - 1}) * b*x - a * \text{arctan2}(1, \sqrt{b*x + a + 1}) * \sqrt{b*x + a - 1}) * \log(b*x + a)^2 + (4*b*x * \text{arctan2}(1, \sqrt{b*x + a + 1}) * \sqrt{b*x + a - 1})^2 - b*x * \log(b^2*x^2 + 2*a*b*x + a^2)^2 * \sqrt{b*x + a + 1} * \sqrt{b*x + a - 1} + 4 * (\sqrt{b^3*x^3} * \text{arctan2}(1, \sqrt{b*x + a + 1}) * \sqrt{b*x + a - 1}) + 2 * a * b^2 * x^2 * \text{arctan2}(1, \sqrt{b*x + a + 1}) * \sqrt{b*x + a - 1}) + (a^2 * \text{arctan2}(1, \sqrt{b*x + a + 1}) * \sqrt{b*x + a - 1}) - \text{arctan2}(1, \sqrt{b*x + a + 1}) * \sqrt{b*x + a - 1}) * b*x - (b^3*x^3 * \text{arctan2}(1, \sqrt{b*x + a + 1}) * \sqrt{b*x + a - 1}) + 3 * a * b^2 * x^2 * \text{arctan2}(1, \sqrt{b*x + a + 1}) * \sqrt{b*x + a - 1}) + a^3 * \text{arctan2}(1, \sqrt{b*x + a + 1}) * \sqrt{b*x + a - 1}) + a^2 * \text{arctan2}(1, \sqrt{b*x + a + 1}) * \sqrt{b*x + a - 1}) - \text{arctan2}(1, \sqrt{b*x + a + 1}) * \sqrt{b*x + a - 1}) * b*x - a * \text{arctan2}(1, \sqrt{b*x + a + 1}) * \sqrt{b*x + a - 1}) * \log(b*x + a)) * \log(b^2*x^2 + 2*a*b*x + a^2)) / (b^3*x^5 + 3*a*b^2*x^4 + (3*a^2 - 1)*b*x^3 + (a^3 - a)*x^2), x) / x \end{aligned}$$

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(b*x+a)^3/x^2,x, algorithm="fricas")`
[Out] `integral(arccsc(b*x + a)^3/x^2, x)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{acsc}^3(a + bx)}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(acsc(b*x+a)**3/x**2,x)`
[Out] `Integral(acsc(a + b*x)**3/x**2, x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(b*x+a)^3/x^2,x, algorithm="giac")`
[Out] `integrate(arccsc(b*x + a)^3/x^2, x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\operatorname{asin}\left(\frac{1}{a+bx}\right)^3}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(asin(1/(a + b*x))^3/x^2,x)`
[Out] `int(asin(1/(a + b*x))^3/x^2, x)`

$$\mathbf{3.38} \quad \int x^3 \csc^{-1}(a + bx^4) dx$$

Optimal. Leaf size=48

$$\frac{(a + bx^4) \csc^{-1}(a + bx^4)}{4b} + \frac{\tanh^{-1}\left(\sqrt{1 - \frac{1}{(a + bx^4)^2}}\right)}{4b}$$

[Out] $\frac{1}{4}*(b*x^4+a)*\text{arccsc}(b*x^4+a)/b+1/4*\text{arctanh}((1-1/(b*x^4+a)^2)^(1/2))/b$

Rubi [A]

time = 0.05, antiderivative size = 48, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.500, Rules used = {6847, 5359, 379, 272, 65, 212}

$$\frac{(a + bx^4) \csc^{-1}(a + bx^4)}{4b} + \frac{\tanh^{-1}\left(\sqrt{1 - \frac{1}{(a + bx^4)^2}}\right)}{4b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[x^3 \text{ArcCsc}[a + b*x^4], x]$

[Out] $((a + b*x^4)*\text{ArcCsc}[a + b*x^4])/(4*b) + \text{ArcTanh}[\text{Sqrt}[1 - (a + b*x^4)^{-2}]]/(4*b)$

Rule 65

```
Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] :> With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 212

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] :> Simplify[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])
```

Rule 272

```
Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] :> Dist[1/n, Subst[
Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b,
m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 379

```
Int[(u_)^m_*((a_) + (b_)*(v_)^n_)^p_, x_Symbol] :> Dist[u^m/(Coeff
icient[v, x, 1]*v^m), Subst[Int[x^m*(a + b*x^n)^p, x], x, v], x] /; FreeQ[{a, b, m, n, p}, x] && LinearPairQ[u, v, x]
```

Rule 5359

```
Int[ArcCsc[(c_) + (d_)*(x_)], x_Symbol] :> Simp[(c + d*x)*(ArcCsc[c + d*x]
/d), x] + Int[1/((c + d*x)*Sqrt[1 - 1/(c + d*x)^2]), x] /; FreeQ[{c, d}, x]
```

Rule 6847

```
Int[(u_)*(x_)^m_, x_Symbol] :> Dist[1/(m + 1), Subst[Int[SubstFor[x^(m +
1), u, x], x], x, x^(m + 1)], x] /; FreeQ[m, x] && NeQ[m, -1] && FunctionQ
fQ[x^(m + 1), u, x]
```

Rubi steps

$$\begin{aligned}
\int x^3 \csc^{-1}(a + bx^4) dx &= \frac{1}{4} \text{Subst}\left(\int \csc^{-1}(a + bx) dx, x, x^4\right) \\
&= \frac{(a + bx^4) \csc^{-1}(a + bx^4)}{4b} + \frac{1}{4} \text{Subst}\left(\int \frac{1}{(a + bx)\sqrt{1 - \frac{1}{(a + bx)^2}}} dx, x, x^4\right) \\
&= \frac{(a + bx^4) \csc^{-1}(a + bx^4)}{4b} + \frac{\text{Subst}\left(\int \frac{1}{\sqrt{1 - \frac{1}{x^2}}} dx, x, a + bx^4\right)}{4b} \\
&= \frac{(a + bx^4) \csc^{-1}(a + bx^4)}{4b} - \frac{\text{Subst}\left(\int \frac{1}{\sqrt{1 - x}} dx, x, \frac{1}{(a+bx^4)^2}\right)}{8b} \\
&= \frac{(a + bx^4) \csc^{-1}(a + bx^4)}{4b} + \frac{\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \sqrt{1 - \frac{1}{(a+bx^4)^2}}\right)}{4b} \\
&= \frac{(a + bx^4) \csc^{-1}(a + bx^4)}{4b} + \frac{\tanh^{-1}\left(\sqrt{1 - \frac{1}{(a+bx^4)^2}}\right)}{4b}
\end{aligned}$$

Mathematica [B] Leaf count is larger than twice the leaf count of optimal. 127 vs. 2(48) = 96.

time = 0.17, size = 127, normalized size = 2.65

$$\frac{(a + bx^4) \csc^{-1}(a + bx^4)}{4b} + \frac{\sqrt{-1 + (a + bx^4)^2} \left(-\log \left(1 - \frac{a + bx^4}{\sqrt{-1 + (a + bx^4)^2}} \right) + \log \left(1 + \frac{a + bx^4}{\sqrt{-1 + (a + bx^4)^2}} \right) \right)}{8b(a + bx^4) \sqrt{1 - \frac{1}{(a + bx^4)^2}}}$$

Antiderivative was successfully verified.

[In] `Integrate[x^3*ArcCsc[a + b*x^4], x]`

[Out] $\frac{((a + b*x^4)*\text{ArcCsc}[a + b*x^4])/(4*b) + (\text{Sqrt}[-1 + (a + b*x^4)^2]*(-\text{Log}[1 - (a + b*x^4)/\text{Sqrt}[-1 + (a + b*x^4)^2]] + \text{Log}[1 + (a + b*x^4)/\text{Sqrt}[-1 + (a + b*x^4)^2]]))}{8*b*(a + b*x^4)*\text{Sqrt}[1 - (a + b*x^4)^{-2}]}$

Maple [A]

time = 0.08, size = 54, normalized size = 1.12

method	result	size
derivativedivides	$\frac{\text{arccsc}(bx^4+a)(bx^4+a)+\ln(bx^4+a+(bx^4+a)\sqrt{1-\frac{1}{(bx^4+a)^2}})}{4b}$	54
default	$\frac{\text{arccsc}(bx^4+a)(bx^4+a)+\ln(bx^4+a+(bx^4+a)\sqrt{1-\frac{1}{(bx^4+a)^2}})}{4b}$	54

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x^3*arccsc(b*x^4+a), x, method=_RETURNVERBOSE)`

[Out] $\frac{1/4/b*(\text{arccsc}(b*x^4+a)*(b*x^4+a)+\ln(b*x^4+a+(b*x^4+a)*(1-1/(b*x^4+a)^2)^(1/2)))}{8b}$

Maxima [A]

time = 0.26, size = 63, normalized size = 1.31

$$\frac{2(bx^4 + a) \text{arccsc}(bx^4 + a) + \log \left(\sqrt{-\frac{1}{(bx^4 + a)^2} + 1} + 1 \right) - \log \left(-\sqrt{-\frac{1}{(bx^4 + a)^2} + 1} + 1 \right)}{8b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^3*arccsc(b*x^4+a), x, algorithm="maxima")`

[Out] $\frac{1/8*(2*(b*x^4 + a)*\text{arccsc}(b*x^4 + a) + \log(\text{sqrt}(-1/(b*x^4 + a)^2 + 1) + 1) - \log(-\text{sqrt}(-1/(b*x^4 + a)^2 + 1) + 1))/b}{8b}$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 88 vs. $2(42) = 84$.

time = 0.37, size = 88, normalized size = 1.83

$$\frac{bx^4 \operatorname{arccsc}(bx^4 + a) - 2a \arctan\left(-bx^4 - a + \sqrt{b^2x^8 + 2abx^4 + a^2 - 1}\right) - \log\left(-bx^4 - a + \sqrt{b^2x^8 + 2abx^4 + a^2 - 1}\right)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^3*arccsc(b*x^4+a),x, algorithm="fricas")`

[Out] $\frac{1}{4}*(b*x^4*\operatorname{arccsc}(b*x^4 + a) - 2*a*\arctan(-b*x^4 - a + \sqrt{b^2*x^8 + 2*a*b*x^4 + a^2 - 1})) - \log(-b*x^4 - a + \sqrt{b^2*x^8 + 2*a*b*x^4 + a^2 - 1})/b$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x**3*acsc(b*x**4+a),x)`

[Out] Timed out

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 91 vs. $2(42) = 84$.
time = 0.50, size = 91, normalized size = 1.90

$$\frac{1}{8} b \left(\frac{2(bx^4 + a) \arcsin\left(\frac{1}{(bx^4 + a)(\frac{a}{bx^4 + a} - 1) - a}\right)}{b^2} + \frac{\log\left(\sqrt{-\frac{1}{(bx^4 + a)^2} + 1} + 1\right) - \log\left(-\sqrt{-\frac{1}{(bx^4 + a)^2} + 1} + 1\right)}{b^2} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^3*arccsc(b*x^4+a),x, algorithm="giac")`

[Out] $\frac{1}{8}b*(2*(b*x^4 + a)*\arcsin(-1/((b*x^4 + a)*(a/(b*x^4 + a) - 1) - a))/b^2 + (\log(\sqrt{-1/(b*x^4 + a)^2 + 1} + 1) - \log(-\sqrt{-1/(b*x^4 + a)^2 + 1} + 1))/b^2)$

Mupad [B]

time = 1.03, size = 44, normalized size = 0.92

$$\frac{\operatorname{atanh}\left(\frac{1}{\sqrt{1 - \frac{1}{(bx^4 + a)^2}}}\right)}{4b} + \frac{\operatorname{asin}\left(\frac{1}{bx^4 + a}\right) (bx^4 + a)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\int x^3 \operatorname{asin}(1/(a + b*x^4)) dx$

[Out] $\operatorname{atanh}(1/(1 - 1/(a + b*x^4)^2)^{1/2})/(4*b) + (\operatorname{asin}(1/(a + b*x^4))*(a + b*x^4))/(4*b)$

$$\text{3.39} \quad \int x^{-1+n} \csc^{-1}(a + bx^n) dx$$

Optimal. Leaf size=48

$$\frac{(a + bx^n) \csc^{-1}(a + bx^n)}{bn} + \frac{\tanh^{-1}\left(\sqrt{1 - \frac{1}{(a + bx^n)^2}}\right)}{bn}$$

[Out] $(a+b*x^n)*\text{arccsc}(a+b*x^n)/b/n + \text{arctanh}((1-1/(a+b*x^n)^2)^{(1/2)})/b/n$

Rubi [A]

time = 0.05, antiderivative size = 48, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.429, Rules used = {6847, 5359, 379, 272, 65, 212}

$$\frac{(a + bx^n) \csc^{-1}(a + bx^n)}{bn} + \frac{\tanh^{-1}\left(\sqrt{1 - \frac{1}{(a + bx^n)^2}}\right)}{bn}$$

Antiderivative was successfully verified.

[In] $\text{Int}[x^{(-1 + n)} * \text{ArcCsc}[a + b*x^n], x]$

[Out] $((a + b*x^n)*\text{ArcCsc}[a + b*x^n])/(b*n) + \text{ArcTanh}[\text{Sqrt}[1 - (a + b*x^n)^{(-2)}]]/(b*n)$

Rule 65

```
Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] :> With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ[
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 212

```
Int[((a_) + (b_)*(x_)^2)^{(-1)}, x_Symbol] :> Simplify[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])
```

Rule 272

```
Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] :> Dist[1/n, Subst[
Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 379

```
Int[(u_)^m_*((a_) + (b_)*(v_)^n_)^p_, x_Symbol] :> Dist[u^m/(Coeff
icient[v, x, 1]*v^m), Subst[Int[x^m*(a + b*x^n)^p, x], x, v], x] /; FreeQ[{a, b, m, n, p}, x] && LinearPairQ[u, v, x]
```

Rule 5359

```
Int[ArcCsc[(c_) + (d_)*(x_)], x_Symbol] :> Simp[(c + d*x)*(ArcCsc[c + d*x]
/d), x] + Int[1/((c + d*x)*Sqrt[1 - 1/(c + d*x)^2]), x] /; FreeQ[{c, d}, x]
```

Rule 6847

```
Int[(u_)*(x_)^m_, x_Symbol] :> Dist[1/(m + 1), Subst[Int[SubstFor[x^(m +
1), u, x], x, x^(m + 1)], x] /; FreeQ[m, x] && NeQ[m, -1] && FunctionO
fQ[x^(m + 1), u, x]
```

Rubi steps

$$\begin{aligned}
\int x^{-1+n} \csc^{-1}(a + bx^n) dx &= \frac{\text{Subst}\left(\int \csc^{-1}(a + bx) dx, x, x^n\right)}{n} \\
&= \frac{(a + bx^n) \csc^{-1}(a + bx^n)}{bn} + \frac{\text{Subst}\left(\int \frac{1}{(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}} dx, x, x^n\right)}{n} \\
&= \frac{(a + bx^n) \csc^{-1}(a + bx^n)}{bn} + \frac{\text{Subst}\left(\int \frac{1}{\sqrt{1-\frac{1}{x^2}}} dx, x, a + bx^n\right)}{bn} \\
&= \frac{(a + bx^n) \csc^{-1}(a + bx^n)}{bn} - \frac{\text{Subst}\left(\int \frac{1}{\sqrt{1-x^2}} dx, x, \frac{1}{(a+bx^n)^2}\right)}{2bn} \\
&= \frac{(a + bx^n) \csc^{-1}(a + bx^n)}{bn} + \frac{\text{Subst}\left(\int \frac{1}{\sqrt{1-\frac{1}{(a+bx^n)^2}}} dx, x, \sqrt{1-\frac{1}{(a+bx^n)^2}}\right)}{bn} \\
&= \frac{(a + bx^n) \csc^{-1}(a + bx^n)}{bn} + \frac{\tanh^{-1}\left(\sqrt{1-\frac{1}{(a+bx^n)^2}}\right)}{bn}
\end{aligned}$$

Mathematica [B] Leaf count is larger than twice the leaf count of optimal. 130 vs. 2(48) = 96.

time = 0.19, size = 130, normalized size = 2.71

$$\frac{(a + bx^n) \csc^{-1}(a + bx^n)}{bn} + \frac{\sqrt{-1 + (a + bx^n)^2} \left(-\log \left(1 - \frac{a + bx^n}{\sqrt{-1 + (a + bx^n)^2}} \right) + \log \left(1 + \frac{a + bx^n}{\sqrt{-1 + (a + bx^n)^2}} \right) \right)}{2bn(a + bx^n) \sqrt{1 - \frac{1}{(a + bx^n)^2}}}$$

Antiderivative was successfully verified.

[In] Integrate[x^(-1 + n)*ArcCsc[a + b*x^n], x]

[Out] $\frac{((a + b x^n) \operatorname{ArcCsc}[a + b x^n])/(b n) + (\operatorname{Sqrt}[-1 + (a + b x^n)^2] * (-\operatorname{Log}[1 - (a + b x^n)/\operatorname{Sqrt}[-1 + (a + b x^n)^2]]) + \operatorname{Log}[1 + (a + b x^n)/\operatorname{Sqrt}[-1 + (a + b x^n)^2]])) / (2 b n (a + b x^n) \operatorname{Sqrt}[1 - (a + b x^n)^{-2}])}{2 b n}$

Maple [F]

time = 0.04, size = 0, normalized size = 0.00

$$\int x^{-1+n} \operatorname{arccsc}(a + b x^n) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^(-1+n)*arccsc(a+b*x^n), x)

[Out] int(x^(-1+n)*arccsc(a+b*x^n), x)

Maxima [A]

time = 0.27, size = 66, normalized size = 1.38

$$\frac{2 (bx^n + a) \operatorname{arccsc}(bx^n + a) + \log \left(\sqrt{-\frac{1}{(bx^n + a)^2} + 1} + 1 \right) - \log \left(-\sqrt{-\frac{1}{(bx^n + a)^2} + 1} + 1 \right)}{2 bn}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^(-1+n)*arccsc(a+b*x^n), x, algorithm="maxima")

[Out] $\frac{1/2 * (2 * (b x^n + a) * \operatorname{arccsc}(b x^n + a) + \log(\sqrt{-1/(b x^n + a)^2 + 1} + 1) - \log(-\sqrt{-1/(b x^n + a)^2 + 1} + 1)) / (b n)}$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 94 vs. 2(46) = 92.

time = 0.39, size = 94, normalized size = 1.96

$$\frac{bx^n \operatorname{arccsc}(bx^n + a) - 2 a \operatorname{arctan} \left(-bx^n - a + \sqrt{b^2 x^{2n} + 2 abx^n + a^2 - 1} \right) - \log \left(-bx^n - a + \sqrt{b^2 x^{2n} + 2 abx^n + a^2 - 1} \right)}{bn}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^(-1+n)*arccsc(a+b*x^n),x, algorithm="fricas")`

[Out]
$$\frac{(b*x^n * \text{arccsc}(b*x^n + a) - 2*a*\text{arctan}(-b*x^n - a + \sqrt{b^2*x^{(2*n)} + 2*a*b*x^n + a^2 - 1}) - \log(-b*x^n - a + \sqrt{b^2*x^{(2*n)} + 2*a*b*x^n + a^2 - 1})) / (b*n)}$$

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x**(-1+n)*acsc(a+b*x**n),x)`

[Out] Exception raised: SystemError >> excessive stack use: stack is 5008 deep

Giac [A]

time = 0.50, size = 74, normalized size = 1.54

$$\frac{b \left(\frac{\frac{2(bx^n+a)\arcsin(\frac{1}{bx^n+a})}{b^2} + \frac{\log\left(\sqrt{-\frac{1}{(bx^n+a)^2}+1}\right)+1}{b^2} - \log\left(-\sqrt{-\frac{1}{(bx^n+a)^2}+1}\right)+1 \right)}{2n}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^(-1+n)*arccsc(a+b*x^n),x, algorithm="giac")`

[Out]
$$\frac{1/2*b*(2*(b*x^n + a)*\arcsin(1/(b*x^n + a))/b^2 + (\log(\sqrt{-1/(b*x^n + a)^2 + 1}) + 1) - \log(-\sqrt{-1/(b*x^n + a)^2 + 1} + 1))/b^2)/n}{n}$$

Mupad [B]

time = 0.73, size = 42, normalized size = 0.88

$$\frac{\operatorname{atanh}\left(\frac{1}{\sqrt{1 - \frac{1}{(a + b x^n)^2}}}\right) + \operatorname{asin}\left(\frac{1}{a + b x^n}\right) (a + b x^n)}{b n}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x^(n - 1)*asin(1/(a + b*x^n)),x)`

[Out]
$$\frac{(\operatorname{atanh}(1/(1 - 1/(a + b*x^n)^2)^{(1/2)}) + \operatorname{asin}(1/(a + b*x^n))*(a + b*x^n))/(b*n)}$$

3.40 $\int \csc^{-1}(ce^{a+bx}) dx$

Optimal. Leaf size=85

$$\frac{i \csc^{-1}(ce^{a+bx})^2}{2b} - \frac{\csc^{-1}(ce^{a+bx}) \log(1 - e^{2i \csc^{-1}(ce^{a+bx})})}{b} + \frac{i \text{PolyLog}\left(2, e^{2i \csc^{-1}(ce^{a+bx})}\right)}{2b}$$

[Out] $1/2*I*\text{arccsc}(c*\exp(b*x+a))^2/b - \text{arccsc}(c*\exp(b*x+a))*\ln(1 - (I/c/\exp(b*x+a) + (1 - 1/c^2/\exp(b*x+a)^2)^{(1/2)})^2)/b + 1/2*I*\text{polylog}(2, (I/c/\exp(b*x+a) + (1 - 1/c^2/e^{xp(b*x+a)^2})^{(1/2)})^2)/b$

Rubi [A]

time = 0.06, antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.700, Rules used = {2320, 5327, 4721, 3798, 2221, 2317, 2438}

$$\frac{i \text{Li}_2\left(e^{2i \csc^{-1}(ce^{a+bx})}\right)}{2b} + \frac{i \csc^{-1}(ce^{a+bx})^2}{2b} - \frac{\csc^{-1}(ce^{a+bx}) \log(1 - e^{2i \csc^{-1}(ce^{a+bx})})}{b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{ArcCsc}[c*E^(a + b*x)], x]$

[Out] $((I/2)*\text{ArcCsc}[c*E^(a + b*x)]^2)/b - (\text{ArcCsc}[c*E^(a + b*x)]*\text{Log}[1 - E^{((2*I)*\text{ArcCsc}[c*E^(a + b*x)])}])/b + ((I/2)*\text{PolyLog}[2, E^{((2*I)*\text{ArcCsc}[c*E^(a + b*x)])}])/b$

Rule 2221

```
Int[((((F_)^((g_.)*(e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/((a_) + (b_.)*((F_)^((g_.)*(e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] :> Simplify[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x]; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2317

```
Int[Log[(a_) + (b_.)*((F_)^((e_.)*(c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] :> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x]; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2320

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_.))^(m_.)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))]*
```

$(F_{_})[v_{_}] /; \text{FreeQ}[\{a, b, c\}, x] \&& \text{InverseFunctionQ}[F[x]]]$

Rule 2438

```
Int[Log[(c_.)*(d_.) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 3798

```
Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + Pi*(k_.) + (f_.)*(x_)], x_Symbol] :> Simp[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Dist[2*I, Int[(c + d*x)^m *E^(2*I*k*Pi)*(E^(2*I*(e + f*x))/(1 + E^(2*I*k*Pi)*E^(2*I*(e + f*x)))), x], x] /; FreeQ[{c, d, e, f}, x] && IntegerQ[4*k] && IGtQ[m, 0]
```

Rule 4721

```
Int[((a_.) + ArcSin[(c_.)*(x_)]*(b_.))^(n_.)/(x_), x_Symbol] :> Subst[Int[(a + b*x)^n*Cot[x], x], x, ArcSin[c*x]] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0]
```

Rule 5327

```
Int[((a_.) + ArcCsc[(c_.)*(x_)]*(b_.))/x, x_Symbol] :> -Subst[Int[(a + b*ArcSin[x/c])/x, x], x, 1/x] /; FreeQ[{a, b, c}, x]
```

Rubi steps

$$\begin{aligned}
\int \csc^{-1}(ce^{a+bx}) dx &= \frac{\text{Subst}\left(\int \frac{\csc^{-1}(cx)}{x} dx, x, e^{a+bx}\right)}{b} \\
&= -\frac{\text{Subst}\left(\int \frac{\sin^{-1}(\frac{x}{c})}{x} dx, x, e^{-a-bx}\right)}{b} \\
&= -\frac{\text{Subst}\left(\int x \cot(x) dx, x, \sin^{-1}\left(\frac{e^{-a-bx}}{c}\right)\right)}{b} \\
&= \frac{i \sin^{-1}\left(\frac{e^{-a-bx}}{c}\right)^2}{2b} + \frac{(2i)\text{Subst}\left(\int \frac{e^{2ix}x}{1-e^{2ix}} dx, x, \sin^{-1}\left(\frac{e^{-a-bx}}{c}\right)\right)}{b} \\
&= \frac{i \sin^{-1}\left(\frac{e^{-a-bx}}{c}\right)^2}{2b} - \frac{\sin^{-1}\left(\frac{e^{-a-bx}}{c}\right) \log\left(1 - e^{2i \sin^{-1}\left(\frac{e^{-a-bx}}{c}\right)}\right)}{b} + \frac{\text{Subst}\left(\int \log(1 - e^{2i \sin^{-1}\left(\frac{e^{-a-bx}}{c}\right)}) dx, x, \sin^{-1}\left(\frac{e^{-a-bx}}{c}\right)\right)}{b} \\
&= \frac{i \sin^{-1}\left(\frac{e^{-a-bx}}{c}\right)^2}{2b} - \frac{\sin^{-1}\left(\frac{e^{-a-bx}}{c}\right) \log\left(1 - e^{2i \sin^{-1}\left(\frac{e^{-a-bx}}{c}\right)}\right)}{b} - \frac{i \text{Subst}\left(\int \frac{\log(1-x)}{x} dx, x, \sin^{-1}\left(\frac{e^{-a-bx}}{c}\right)\right)}{b} \\
&= \frac{i \sin^{-1}\left(\frac{e^{-a-bx}}{c}\right)^2}{2b} - \frac{\sin^{-1}\left(\frac{e^{-a-bx}}{c}\right) \log\left(1 - e^{2i \sin^{-1}\left(\frac{e^{-a-bx}}{c}\right)}\right)}{b} + \frac{i \text{Li}_2\left(e^{2i \sin^{-1}\left(\frac{e^{-a-bx}}{c}\right)}\right)}{2b}
\end{aligned}$$

Mathematica [B] Leaf count is larger than twice the leaf count of optimal. 280 vs. 2(85) = 170.

time = 0.46, size = 280, normalized size = 3.29

$$x \csc^{-1}(ce^{a+bx}) + \frac{e^{-a-bx} \left(4 \sqrt{-1 + c^2 e^{2(a+b x)}} \operatorname{ArcTan}\left(\sqrt{-1 + c^2 e^{2(a+b x)}}\right) (2 b x - \log(c^2 e^{2(a+b x)})) + \sqrt{1 - c^2 e^{2(a+b x)}} \left(\log^2(c^2 e^{2(a+b x)}) - 4 \log(c^2 e^{2(a+b x)}) \log\left(\frac{1}{2} \left(1 + \sqrt{1 - c^2 e^{2(a+b x)}}\right)\right) + 2 \log^2\left(\frac{1}{2} \left(1 + \sqrt{1 - c^2 e^{2(a+b x)}}\right)\right)\right) - 4 \sqrt{1 - c^2 e^{2(a+b x)}} \operatorname{PolyLog}\left(2, \frac{1}{2} \left(1 - \sqrt{1 - c^2 e^{2(a+b x)}}\right)\right)\right)}{8 b c \sqrt{1 - \frac{e^{-2(a+b x)}}{c^2}}}$$

Antiderivative was successfully verified.

[In] `Integrate[ArcCsc[c*E^(a + b*x)], x]`

[Out] $x \operatorname{ArcCsc}[c \operatorname{E}^{(a+b x)}] + (\operatorname{E}^{(-a-b x)} (4 \operatorname{Sqrt}[-1+c^2 \operatorname{E}^{(2(a+b x))}] * \operatorname{rcTan}[\operatorname{Sqrt}[-1+c^2 \operatorname{E}^{(2(a+b x))}]]) * (2 b x-\operatorname{Log}[c^2 \operatorname{E}^{(2(a+b x))}]) + \operatorname{Sqrt}[1-c^2 \operatorname{E}^{(2(a+b x))}] * (\operatorname{Log}[c^2 \operatorname{E}^{(2(a+b x))}]^2-4 \operatorname{Log}[c^2 \operatorname{E}^{(2(a+b x))}] * \operatorname{Log}[(1+\operatorname{Sqrt}[1-c^2 \operatorname{E}^{(2(a+b x))}])] / 2)+2 \operatorname{Log}[(1+\operatorname{Sqrt}[1-c^2 \operatorname{E}^{(2(a+b x))}])] / 2^2-4 \operatorname{Sqrt}[1-c^2 \operatorname{E}^{(2(a+b x))}] * \operatorname{PolyLog}[2,(1-\operatorname{Sqrt}[1-c^2 \operatorname{E}^{(2(a+b x))}])] / 2])) / (8 b c \operatorname{Sqrt}[1-1/(c^2 \operatorname{E}^{(2(a+b x))})])$

Maple [A]

time = 1.21, size = 188, normalized size = 2.21

method	result
derivativedivides	$\frac{i \operatorname{arccsc}\left(e^{bx+a} c\right)^2}{2}-\operatorname{arccsc}\left(e^{bx+a} c\right) \ln \left(1-\frac{i e^{-bx-a}}{c}-\sqrt{1-\frac{e^{-2bx-2a}}{c^2}}\right)-\operatorname{arccsc}\left(e^{bx+a} c\right) \ln \left(1+\frac{i e^{-bx-a}}{c}+\sqrt{1-\frac{e^{-2bx-2a}}{c^2}}\right)$
default	$\frac{i \operatorname{arccsc}\left(e^{bx+a} c\right)^2}{2}-\operatorname{arccsc}\left(e^{bx+a} c\right) \ln \left(1-\frac{i e^{-bx-a}}{c}-\sqrt{1-\frac{e^{-2bx-2a}}{c^2}}\right)-\operatorname{arccsc}\left(e^{bx+a} c\right) \ln \left(1+\frac{i e^{-bx-a}}{c}+\sqrt{1-\frac{e^{-2bx-2a}}{c^2}}\right)$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(arccsc(exp(b*x+a)*c),x,method=_RETURNVERBOSE)
[Out] 1/b*(1/2*I*arccsc(exp(b*x+a)*c)^2-arccsc(exp(b*x+a)*c)*ln(1-I/exp(b*x+a)/c-
(1-1/exp(b*x+a)^2/c^2)^(1/2))-arccsc(exp(b*x+a)*c)*ln(1+I/exp(b*x+a)/c+(1-1/
exp(b*x+a)^2/c^2)^(1/2))+I*polylog(2,I/exp(b*x+a)/c+(1-1/exp(b*x+a)^2/c^2)^(1/2))+I*polylog(2,-I/exp(b*x+a)/c-(1-1/exp(b*x+a)^2/c^2)^(1/2)))
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(arccsc(c*exp(b*x+a)),x, algorithm="maxima")
[Out] 1/2*(2*b^2*c^2*integrate(x*e^(2*b*x + 2*a + 1/2*log(c*e^(b*x + a) + 1) + 1/2*log(c*e^(b*x + a) - 1))/(c^2*e^(2*b*x + 2*a) + (c^2*e^(2*b*x + 2*a) - 1)*e^(log(c*e^(b*x + a) + 1) + log(c*e^(b*x + a) - 1)) - 1), x) - 2*I*b^2*c^2*integrate(x*e^(2*b*x + 2*a)/(c^2*e^(2*b*x + 2*a) + (c^2*e^(2*b*x + 2*a) - 1)*e^(log(c*e^(b*x + a) + 1) + log(c*e^(b*x + a) - 1)) - 1), x) + I*b^2*x^2 - I*b*x*log(c^2*e^(2*b*x + 2*a)) + I*b*x*log(c*e^(b*x + a) + 1) + I*b*x*log(-c*e^(b*x + a) + 1) - 2*((-I*a - arctan2(1, sqrt(c*e^(b*x + a) + 1))*sqrt(c*e^(b*x + a) - 1)))*b - I*b*log(c)*x + I*dilog(c*e^(b*x + a)) + I*dilog(-c*e^(b*x + a)))/b
```

Fricas [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(arccsc(c*exp(b*x+a)),x, algorithm="fricas")
[Out] Exception raised: TypeError >> Error detected within library code:   integ-
rate: implementation incomplete (constant residues)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \operatorname{acsc}(ce^{a+bx}) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(acsc(c*exp(b*x+a)),x)

[Out] Integral(acsc(c*exp(a + b*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(arccsc(c*exp(b*x+a)),x, algorithm="giac")

[Out] integrate(arccsc(c*e^(b*x + a)), x)

Mupad [B]

time = 1.25, size = 91, normalized size = 1.07

$$\frac{\text{polylog}\left(2, e^{\operatorname{asin}\left(\frac{e^{-a-b x}}{c}\right) 2 i}\right) \operatorname{li}{1i}}{2 b} + \frac{\operatorname{asin}\left(\frac{e^{-a-b x}}{c}\right)^2 \operatorname{li}{1i}}{2 b} - \frac{\ln \left(1 - e^{\operatorname{asin}\left(\frac{e^{-a-b x}}{c}\right) 2 i}\right) \operatorname{asin}\left(\frac{e^{-a-b x}}{c}\right)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(asin(exp(- a - b*x)/c),x)

[Out] $\frac{\text{polylog}(2, \exp(\operatorname{asin}(\exp(- a - b*x)/c)*2i))*1i}{(2*b)} + \frac{(\operatorname{asin}(\exp(- a - b*x)/c)^2*1i)}{(2*b)} - \frac{(\log(1 - \exp(\operatorname{asin}(\exp(- a - b*x)/c)*2i))*\operatorname{asin}(\exp(- a - b*x)/c))}{b}$

3.41 $\int e^{\csc^{-1}(ax)} x^2 dx$

Optimal. Leaf size=95

$$\frac{\left(\frac{4}{5} - \frac{12i}{5}\right) e^{(1+3i)\csc^{-1}(ax)} {}_2F_1\left(\frac{3}{2} - \frac{i}{2}, 3; \frac{5}{2} - \frac{i}{2}; e^{2i\csc^{-1}(ax)}\right)}{a^3} - \frac{\left(\frac{8}{5} - \frac{24i}{5}\right) e^{(1+3i)\csc^{-1}(ax)} {}_2F_1\left(\frac{3}{2} - \frac{i}{2}, 4; \frac{5}{2} - \frac{i}{2}; e^{2i\csc^{-1}(ax)}\right)}{a^3}$$

[Out] $(4/5 - 12/5*I)*\exp((1+3*I)*\text{arccsc}(a*x))*\text{hypergeom}([3, 3/2 - 1/2*I], [5/2 - 1/2*I], (I/a/x + (1 - 1/a^2/x^2)^(1/2))^2)/a^3 + (-8/5 + 24/5*I)*\exp((1+3*I)*\text{arccsc}(a*x))*\text{hypergeom}([4, 3/2 - 1/2*I], [5/2 - 1/2*I], (I/a/x + (1 - 1/a^2/x^2)^(1/2))^2)/a^3$

Rubi [A]

time = 0.08, antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.400, Rules used = {5375, 12, 4559, 2283}

$$\frac{\left(\frac{4}{5} - \frac{12i}{5}\right) e^{(1+3i)\csc^{-1}(ax)} {}_2F_1\left(\frac{3}{2} - \frac{i}{2}, 3; \frac{5}{2} - \frac{i}{2}; e^{2i\csc^{-1}(ax)}\right)}{a^3} - \frac{\left(\frac{8}{5} - \frac{24i}{5}\right) e^{(1+3i)\csc^{-1}(ax)} {}_2F_1\left(\frac{3}{2} - \frac{i}{2}, 4; \frac{5}{2} - \frac{i}{2}; e^{2i\csc^{-1}(ax)}\right)}{a^3}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{\wedge} \text{ArcCsc}[a*x]*x^2, x]$

[Out] $((4/5 - (12*I)/5)*E^((1 + 3*I)*\text{ArcCsc}[a*x]))*\text{Hypergeometric2F1}[3/2 - I/2, 3, 5/2 - I/2, E^((2*I)*\text{ArcCsc}[a*x]))]/a^3 - ((8/5 - (24*I)/5)*E^((1 + 3*I)*\text{ArcCsc}[a*x]))*\text{Hypergeometric2F1}[3/2 - I/2, 4, 5/2 - I/2, E^((2*I)*\text{ArcCsc}[a*x))]/a^3$

Rule 12

$\text{Int}[(a_)*(u_), x_Symbol] :> \text{Dist}[a, \text{Int}[u, x], x] /; \text{FreeQ}[a, x] \&& \text{!MatchQ}[u, (b_)*(v_) /; \text{FreeQ}[b, x]]$

Rule 2283

$\text{Int}[(a_*) + (b_*)*(F_*)^((e_*)*((c_*) + (d_*)*(x_))))^((p_*)*(G_*)^((h_*)*((f_*) + (g_*)*(x_)))), x_Symbol] :> \text{Simp}[a^p*(G^{(h*(f + g*x))}/(g*h*\text{Log}[G]))*\text{Hypergeometric2F1}[-p, g*h*(\text{Log}[G]/(d*e*\text{Log}[F])), g*h*(\text{Log}[G]/(d*e*\text{Log}[F])) + 1, \text{Simplify}[(-b/a)*F^((e*(c + d*x)))]], x] /; \text{FreeQ}[\{F, G, a, b, c, d, e, f, g, h, p\}, x] \&& (\text{ILtQ}[p, 0] \text{ || } \text{GtQ}[a, 0])$

Rule 4559

$\text{Int}[(F_*)^((c_*)*((a_*) + (b_*)*(x_)))*(G_*[(d_*) + (e_*)*(x_)]^((m_*)*(H_*[(d_*) + (e_*)*(x_)]^((n_*), x_Symbol) :> \text{Int}[\text{ExpandTrigToExp}[F^((c*(a + b*x)), G[d + e*x]^m*H[d + e*x]^n, x], x] /; \text{FreeQ}[\{F, a, b, c, d, e\}, x] \&& \text{IGtQ}[m, 0] \&& \text{IGtQ}[n, 0] \&& \text{TrigQ}[G] \&& \text{TrigQ}[H]$

Rule 5375

```
Int[(u_)*(f_)^(ArcCsc[(a_.) + (b_.)*(x_.)]^(n_.)*(c_.)), x_Symbol] :> Dist[-b^(-1), Subst[Int[(u /. x -> -a/b + Csc[x]/b)*f^(c*x^n)*Csc[x]*Cot[x], x], x, ArcCsc[a + b*x]], x] /; FreeQ[{a, b, c, f}, x] && IGtQ[n, 0]
```

Rubi steps

$$\begin{aligned} \int e^{\csc^{-1}(ax)} x^2 dx &= -\frac{\text{Subst}\left(\int \frac{e^x \cot(x) \csc^3(x)}{a^2} dx, x, \csc^{-1}(ax)\right)}{a} \\ &= -\frac{\text{Subst}\left(\int e^x \cot(x) \csc^3(x) dx, x, \csc^{-1}(ax)\right)}{a^3} \\ &= -\frac{\text{Subst}\left(\int \left(\frac{16e^{(1+3i)x}}{(-1+e^{2ix})^4} + \frac{8e^{(1+3i)x}}{(-1+e^{2ix})^3}\right) dx, x, \csc^{-1}(ax)\right)}{a^3} \\ &= -\frac{8\text{Subst}\left(\int \frac{e^{(1+3i)x}}{(-1+e^{2ix})^3} dx, x, \csc^{-1}(ax)\right)}{a^3} - \frac{16\text{Subst}\left(\int \frac{e^{(1+3i)x}}{(-1+e^{2ix})^4} dx, x, \csc^{-1}(ax)\right)}{a^3} \\ &= \frac{\left(\frac{4}{5} - \frac{12i}{5}\right) e^{(1+3i) \csc^{-1}(ax)} {}_2F_1\left(\frac{3}{2} - \frac{i}{2}, 3; \frac{5}{2} - \frac{i}{2}; e^{2i \csc^{-1}(ax)}\right)}{a^3} - \frac{\left(\frac{8}{5} - \frac{24i}{5}\right) e^{(1+3i) \csc^{-1}(ax)} {}_2F_1\left(\frac{1}{2} - \frac{i}{2}, 2; \frac{3}{2} - \frac{i}{2}; e^{2i \csc^{-1}(ax)}\right)}{a^3} \end{aligned}$$

Mathematica [A]

time = 0.34, size = 79, normalized size = 0.83

$$\frac{e^{\csc^{-1}(ax)} \left((4 + 4i)e^{i \csc^{-1}(ax)} {}_2F_1\left(\frac{1}{2} - \frac{i}{2}, 1; \frac{3}{2} - \frac{i}{2}; e^{2i \csc^{-1}(ax)}\right) + a^3 x^3 (5 - \cos(2 \csc^{-1}(ax)) + \sin(2 \csc^{-1}(ax)))\right)}{12a^3}$$

Antiderivative was successfully verified.

[In] Integrate[E^ArcCsc[a*x]*x^2,x]

[Out] (E^ArcCsc[a*x]*((4 + 4*I)*E^(I*ArcCsc[a*x])*Hypergeometric2F1[1/2 - I/2, 1, 3/2 - I/2, E^((2*I)*ArcCsc[a*x])] + a^3*x^3*(5 - Cos[2*ArcCsc[a*x]] + Sin[2*ArcCsc[a*x]])))/(12*a^3)

Maple [F]

time = 0.04, size = 0, normalized size = 0.00

$$\int e^{\arccsc(ax)} x^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(arccsc(a*x))*x^2,x)

[Out] $\int \exp(\operatorname{arccsc}(ax))x^2 dx$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] $\operatorname{integrate}(\exp(\operatorname{arccsc}(ax))x^2, x, \text{algorithm}=\text{"maxima"})$

[Out] $\operatorname{integrate}(x^2 e^{\operatorname{arccsc}(ax)}, x)$

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] $\operatorname{integrate}(\exp(\operatorname{arccsc}(ax))x^2, x, \text{algorithm}=\text{"fricas"})$

[Out] $\operatorname{integral}(x^2 e^{\operatorname{arccsc}(ax)}, x)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x^2 e^{\operatorname{acsc}(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\operatorname{integrate}(\exp(\operatorname{acsc}(ax))x^{**2}, x)$

[Out] $\operatorname{Integral}(x^{**2} \exp(\operatorname{acsc}(ax)), x)$

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] $\operatorname{integrate}(\exp(\operatorname{arccsc}(ax))x^2, x, \text{algorithm}=\text{"giac"})$

[Out] $\operatorname{integrate}(x^2 e^{\operatorname{arccsc}(ax)}, x)$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int x^2 e^{\operatorname{asin}(\frac{1}{ax})} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\operatorname{int}(x^2 \exp(\operatorname{asin}(1/(ax))), x)$

[Out] $\operatorname{int}(x^2 \exp(\operatorname{asin}(1/(ax))), x)$

3.42 $\int e^{\csc^{-1}(ax)} x \, dx$

Optimal. Leaf size=87

$$\frac{\left(\frac{8}{5} + \frac{4i}{5}\right) e^{(1+2i)\csc^{-1}(ax)} {}_2F_1\left(1 - \frac{i}{2}, 2; 2 - \frac{i}{2}; e^{2i\csc^{-1}(ax)}\right)}{a^2} - \frac{\left(\frac{16}{5} + \frac{8i}{5}\right) e^{(1+2i)\csc^{-1}(ax)} {}_2F_1\left(1 - \frac{i}{2}, 3; 2 - \frac{i}{2}; e^{2i\csc^{-1}(ax)}\right)}{a^2}$$

[Out] $(8/5 + 4/5*I)*\exp((1+2*I)*\text{arccsc}(a*x))*\text{hypergeom}([2, 1-1/2*I], [2-1/2*I], (I/a/x+(1-1/a^2/x^2)^(1/2))^2)/a^2 - (16/5 + 8/5*I)*\exp((1+2*I)*\text{arccsc}(a*x))*\text{hypergeom}([3, 1-1/2*I], [2-1/2*I], (I/a/x+(1-1/a^2/x^2)^(1/2))^2)/a^2$

Rubi [A]

time = 0.08, antiderivative size = 87, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5375, 12, 4559, 2283}

$$\frac{\left(\frac{8}{5} + \frac{4i}{5}\right) e^{(1+2i)\csc^{-1}(ax)} {}_2F_1\left(1 - \frac{i}{2}, 2; 2 - \frac{i}{2}; e^{2i\csc^{-1}(ax)}\right)}{a^2} - \frac{\left(\frac{16}{5} + \frac{8i}{5}\right) e^{(1+2i)\csc^{-1}(ax)} {}_2F_1\left(1 - \frac{i}{2}, 3; 2 - \frac{i}{2}; e^{2i\csc^{-1}(ax)}\right)}{a^2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{\wedge} \text{ArcCsc}[a*x]*x, x]$

[Out] $((8/5 + (4*I)/5)*E^((1 + 2*I)*\text{ArcCsc}[a*x]))*\text{Hypergeometric2F1}[1 - I/2, 2, 2 - I/2, E^((2*I)*\text{ArcCsc}[a*x]))/a^2 - ((16/5 + (8*I)/5)*E^((1 + 2*I)*\text{ArcCsc}[a*x]))*\text{Hypergeometric2F1}[1 - I/2, 3, 2 - I/2, E^((2*I)*\text{ArcCsc}[a*x]))/a^2$

Rule 12

$\text{Int}[(a_*)*(u_), x_Symbol] :> \text{Dist}[a, \text{Int}[u, x], x] /; \text{FreeQ}[a, x] \&& \text{!MatchQ}[u, (b_*)*(v_) /; \text{FreeQ}[b, x]]$

Rule 2283

$\text{Int}[((a_) + (b_*)*(F_))^{((e_*)*((c_.) + (d_*)*(x_)))})^{(p_*)*(G_)}^{((h_.)*(f_.) + (g_*)*(x_))), x_Symbol] :> \text{Simp}[a^p*(G^{(h*(f + g*x))/(g*h*\text{Log}[G])})*\text{Hypergeometric2F1}[-p, g*h*(\text{Log}[G]/(d*e*\text{Log}[F])), g*h*(\text{Log}[G]/(d*e*\text{Log}[F])) + 1, \text{Simplify}[(-b/a)*F^((e*(c + d*x)))]], x] /; \text{FreeQ}[\{F, G, a, b, c, d, e, f, g, h, p\}, x] \&& (\text{ILtQ}[p, 0] \text{ || } \text{GtQ}[a, 0])$

Rule 4559

$\text{Int}[(F_)^{((c_.)*(a_.) + (b_*)*(x_)))*(G_)[(d_.) + (e_*)*(x_)]^{(m_.)*(H_.)}[(d_.) + (e_*)*(x_)]^{(n_.)}, x_Symbol] :> \text{Int}[\text{ExpandTrigToExp}[F^((c*(a + b*x)), G[d + e*x]^m*H[d + e*x]^n, x], x] /; \text{FreeQ}[\{F, a, b, c, d, e\}, x] \&& \text{IGtQ}[m, 0] \&& \text{IGtQ}[n, 0] \&& \text{TrigQ}[G] \&& \text{TrigQ}[H]$

Rule 5375

```
Int[(u_)*(f_)^(ArcCsc[(a_.) + (b_.)*(x_)]^(n_.)*(c_.)), x_Symbol] :> Dist[-b^(-1), Subst[Int[(u /. x -> -a/b + Csc[x]/b)*f^(c*x^n)*Csc[x]*Cot[x], x], x, ArcCsc[a + b*x]], x] /; FreeQ[{a, b, c, f}, x] && IGtQ[n, 0]
```

Rubi steps

$$\begin{aligned} \int e^{\csc^{-1}(ax)} x dx &= -\frac{\text{Subst}\left(\int \frac{e^x \cot(x) \csc^2(x)}{a} dx, x, \csc^{-1}(ax)\right)}{a} \\ &= -\frac{\text{Subst}\left(\int e^x \cot(x) \csc^2(x) dx, x, \csc^{-1}(ax)\right)}{a^2} \\ &= -\frac{\text{Subst}\left(\int \left(-\frac{8ie^{(1+2i)x}}{(-1+e^{2ix})^3} - \frac{4ie^{(1+2i)x}}{(-1+e^{2ix})^2}\right) dx, x, \csc^{-1}(ax)\right)}{a^2} \\ &= \frac{(4i)\text{Subst}\left(\int \frac{e^{(1+2i)x}}{(-1+e^{2ix})^2} dx, x, \csc^{-1}(ax)\right)}{a^2} + \frac{(8i)\text{Subst}\left(\int \frac{e^{(1+2i)x}}{(-1+e^{2ix})^3} dx, x, \csc^{-1}(ax)\right)}{a^2} \\ &= \frac{\left(\frac{8}{5} + \frac{4i}{5}\right) e^{(1+2i)\csc^{-1}(ax)} {}_2F_1\left(1 - \frac{i}{2}, 2; 2 - \frac{i}{2}; e^{2i\csc^{-1}(ax)}\right)}{a^2} - \frac{\left(\frac{16}{5} + \frac{8i}{5}\right) e^{(1+2i)\csc^{-1}(ax)} {}_2F_1\left(1 - \frac{i}{2}, 2 - \frac{i}{2}; 2 - \frac{i}{2}; e^{2i\csc^{-1}(ax)}\right)}{a^2} \end{aligned}$$

Mathematica [A]

time = 0.19, size = 101, normalized size = 1.16

$$\frac{\left(\frac{1}{5} + \frac{i}{10}\right) e^{\csc^{-1}(ax)} \left((2-i)ax \left(\sqrt{1 - \frac{1}{a^2 x^2}} + ax\right) + (1+2i) {}_2F_1\left(-\frac{i}{2}, 1; 1 - \frac{i}{2}; e^{2i\csc^{-1}(ax)}\right) + e^{2i\csc^{-1}(ax)} {}_2F_1\left(1, 1 - \frac{i}{2}; 2 - \frac{i}{2}; e^{2i\csc^{-1}(ax)}\right)\right)}{a^2}$$

Antiderivative was successfully verified.

[In] `Integrate[E^ArcCsc[a*x]*x,x]`

[Out] $((1/5 + I/10)*E^{\text{ArcCsc}[a*x]}*((2 - I)*a*x*(\text{Sqrt}[1 - 1/(a^2*x^2)] + a*x) + (1 + 2*I)*\text{Hypergeometric2F1}[-1/2*I, 1, 1 - I/2, E^{((2*I)*\text{ArcCsc}[a*x])}] + E^{((2*I)*\text{ArcCsc}[a*x])}*\text{Hypergeometric2F1}[1, 1 - I/2, 2 - I/2, E^{((2*I)*\text{ArcCsc}[a*x])}]))/a^2$

Maple [F]

time = 0.04, size = 0, normalized size = 0.00

$$\int e^{\text{arccsc}(ax)} x dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\int \exp(\operatorname{arccsc}(ax)) \cdot x \, dx$

[Out] $\int \exp(\operatorname{arccsc}(ax)) \cdot x \, dx$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] $\operatorname{integrate}(\exp(\operatorname{arccsc}(ax)) \cdot x, x, \text{algorithm}=\text{"maxima"})$

[Out] $\operatorname{integrate}(x \cdot e^{\operatorname{arccsc}(ax)}, x)$

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] $\operatorname{integrate}(\exp(\operatorname{arccsc}(ax)) \cdot x, x, \text{algorithm}=\text{"fricas"})$

[Out] $\operatorname{integral}(x \cdot e^{\operatorname{arccsc}(ax)}, x)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x e^{\operatorname{acsc}(ax)} \, dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\operatorname{integrate}(\exp(\operatorname{acsc}(ax)) \cdot x, x)$

[Out] $\operatorname{Integral}(x \cdot \exp(\operatorname{acsc}(ax)), x)$

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] $\operatorname{integrate}(\exp(\operatorname{arccsc}(ax)) \cdot x, x, \text{algorithm}=\text{"giac"})$

[Out] $\operatorname{integrate}(x \cdot e^{\operatorname{arccsc}(ax)}, x)$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int x e^{\operatorname{asin}\left(\frac{1}{ax}\right)} \, dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x*exp(asin(1/(a*x))),x)`
[Out] `int(x*exp(asin(1/(a*x))), x)`

3.43 $\int e^{\csc^{-1}(ax)} dx$

Optimal. Leaf size=87

$$\frac{(1-i)e^{(1+i)\csc^{-1}(ax)} {}_2F_1\left(\frac{1}{2}-\frac{i}{2}, 1; \frac{3}{2}-\frac{i}{2}; e^{2i\csc^{-1}(ax)}\right)}{a} + \frac{(2-2i)e^{(1+i)\csc^{-1}(ax)} {}_2F_1\left(\frac{1}{2}-\frac{i}{2}, 2; \frac{3}{2}-\frac{i}{2}; e^{2i\csc^{-1}(ax)}\right)}{a}$$

[Out] $(-1+I)*\exp((1+I)*\text{arccsc}(a*x))*\text{hypergeom}([1, 1/2-1/2*I], [3/2-1/2*I], (I/a/x+(1-1/a^2/x^2)^(1/2))^2)/a + (2-2*I)*\exp((1+I)*\text{arccsc}(a*x))*\text{hypergeom}([2, 1/2-1/2*I], [3/2-1/2*I], (I/a/x+(1-1/a^2/x^2)^(1/2))^2)/a$

Rubi [A]

time = 0.07, antiderivative size = 87, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, integrand size = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5375, 4559, 2283}

$$\frac{(2-2i)e^{(1+i)\csc^{-1}(ax)} {}_2F_1\left(\frac{1}{2}-\frac{i}{2}, 2; \frac{3}{2}-\frac{i}{2}; e^{2i\csc^{-1}(ax)}\right)}{a} - \frac{(1-i)e^{(1+i)\csc^{-1}(ax)} {}_2F_1\left(\frac{1}{2}-\frac{i}{2}, 1; \frac{3}{2}-\frac{i}{2}; e^{2i\csc^{-1}(ax)}\right)}{a}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{\wedge} \text{ArcCsc}[a*x], x]$

[Out] $((-1 + I)*E^{\wedge}((1 + I)*\text{ArcCsc}[a*x])* \text{Hypergeometric2F1}[1/2 - I/2, 1, 3/2 - I/2, E^{\wedge}((2*I)*\text{ArcCsc}[a*x]))/a + ((2 - 2*I)*E^{\wedge}((1 + I)*\text{ArcCsc}[a*x])* \text{Hypergeometric2F1}[1/2 - I/2, 2, 3/2 - I/2, E^{\wedge}((2*I)*\text{ArcCsc}[a*x]))/a$

Rule 2283

$\text{Int}[((a_) + (b_*)*(F_*)^((e_*)*((c_*) + (d_*)*(x_))))^((p_)*(G_*)^((h_*)*((f_*) + (g_*)*(x_)))), x_Symbol] :> \text{Simp}[a^p*(G^h*(h*(f + g*x))/(g*h*\text{Log}[G]))*\text{Hypergeometric2F1}[-p, g*h*(\text{Log}[G]/(d*e*\text{Log}[F])), g*h*(\text{Log}[G]/(d*e*\text{Log}[F])) + 1, \text{Simplify}[(-b/a)*F^e*(e*(c + d*x))]], x] /; \text{FreeQ}[\{F, G, a, b, c, d, e, f, g, h, p\}, x] \&& (\text{ILtQ}[p, 0] \text{ || } \text{GtQ}[a, 0])$

Rule 4559

$\text{Int}[(F_*)^((c_*)*((a_*) + (b_*)*(x_)))*(G_*)[(d_*) + (e_*)*(x_)]^((m_*)*(H_*)[((d_*) + (e_*)*(x_))^(n_*)], x_Symbol] :> \text{Int}[\text{ExpandTrigToExp}[F^c*(a + b*x), G[d + e*x]^m*H[d + e*x]^n, x], x] /; \text{FreeQ}[\{F, a, b, c, d, e\}, x] \&& \text{IGtQ}[m, 0] \&& \text{IGtQ}[n, 0] \&& \text{TrigQ}[G] \&& \text{TrigQ}[H]$

Rule 5375

$\text{Int}[(u_*)*(f_*)^(\text{ArcCsc}[(a_*) + (b_*)*(x_)]^((n_*)*(c_))), x_Symbol] :> \text{Dist}[-b^(-1), \text{Subst}[\text{Int}[(u /. x \rightarrow -a/b + \text{Csc}[x]/b)*f^c*(c*x^n)*\text{Csc}[x]*\text{Cot}[x], x], x, \text{ArcCsc}[a + b*x]], x] /; \text{FreeQ}[\{a, b, c, f\}, x] \&& \text{IGtQ}[n, 0]$

Rubi steps

$$\begin{aligned}
 \int e^{\csc^{-1}(ax)} dx &= -\frac{\text{Subst}\left(\int e^x \cot(x) \csc(x) dx, x, \csc^{-1}(ax)\right)}{a} \\
 &= -\frac{\text{Subst}\left(\int \left(\frac{2e^{(1+i)x}}{1-e^{2ix}} - \frac{4e^{(1+i)x}}{(-1+e^{2ix})^2}\right) dx, x, \csc^{-1}(ax)\right)}{a} \\
 &= -\frac{2\text{Subst}\left(\int \frac{e^{(1+i)x}}{1-e^{2ix}} dx, x, \csc^{-1}(ax)\right)}{a} + \frac{4\text{Subst}\left(\int \frac{e^{(1+i)x}}{(-1+e^{2ix})^2} dx, x, \csc^{-1}(ax)\right)}{a} \\
 &= -\frac{(1-i)e^{(1+i)\csc^{-1}(ax)} {}_2F_1\left(\frac{1}{2} - \frac{i}{2}, 1; \frac{3}{2} - \frac{i}{2}; e^{2i\csc^{-1}(ax)}\right)}{a} + \frac{(2-2i)e^{(1+i)\csc^{-1}(ax)} {}_2F_1\left(\frac{1}{2} - \frac{i}{2}, 1; \frac{3}{2} - \frac{i}{2}; e^{2i\csc^{-1}(ax)}\right)}{a}
 \end{aligned}$$

Mathematica [A]

time = 0.07, size = 54, normalized size = 0.62

$$\frac{e^{\csc^{-1}(ax)} \left(ax + (1+i)e^{i\csc^{-1}(ax)} {}_2F_1\left(\frac{1}{2} - \frac{i}{2}, 1; \frac{3}{2} - \frac{i}{2}; e^{2i\csc^{-1}(ax)}\right)\right)}{a}$$

Antiderivative was successfully verified.

[In] `Integrate[E^ArcCsc[a*x], x]`

[Out] `(E^ArcCsc[a*x]*(a*x + (1 + I)*E^(I*ArcCsc[a*x])*Hypergeometric2F1[1/2 - I/2, 1, 3/2 - I/2, E^((2*I)*ArcCsc[a*x])]))/a`

Maple [F]

time = 0.04, size = 0, normalized size = 0.00

$$\int e^{\arccsc(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(arccsc(a*x)), x)`

[Out] `int(exp(arccsc(a*x)), x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(arccsc(a*x)), x, algorithm="maxima")`

[Out] integrate($e^{\text{arccsc}(ax)}$), x

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(arccsc(ax)),x, algorithm="fricas")

[Out] integral($e^{\text{arccsc}(ax)}$), x

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int e^{\text{acsc}(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(acsc(ax)),x)

[Out] Integral(exp(acsc(ax)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(arccsc(ax)),x, algorithm="giac")

[Out] integrate($e^{\text{arccsc}(ax)}$), x

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int e^{\text{asin}(\frac{1}{ax})} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(asin(1/(ax))),x)

[Out] int(exp(asin(1/(ax))), x)

3.44 $\int \frac{e^{\csc^{-1}(ax)}}{x} dx$

Optimal. Leaf size=43

$$-ie^{\csc^{-1}(ax)} + 2ie^{\csc^{-1}(ax)} {}_2F_1\left(-\frac{i}{2}, 1; 1 - \frac{i}{2}; e^{2i\csc^{-1}(ax)}\right)$$

[Out] $-I*\exp(\operatorname{arccsc}(a*x))+2*I*\exp(\operatorname{arccsc}(a*x))*\text{hypergeom}([1, -1/2*I], [1-1/2*I], (I/a/x+(1-1/a^2/x^2)^(1/2))^2)$

Rubi [A]

time = 0.04, antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5375, 12, 4528, 2225, 2283}

$$2ie^{\csc^{-1}(ax)} {}_2F_1\left(-\frac{i}{2}, 1; 1 - \frac{i}{2}; e^{2i\csc^{-1}(ax)}\right) - ie^{\csc^{-1}(ax)}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{\operatorname{ArcCsc}[a*x]}/x, x]$

[Out] $(-I)*E^{\operatorname{ArcCsc}[a*x]} + (2*I)*E^{\operatorname{ArcCsc}[a*x]}*\text{Hypergeometric2F1}[-1/2*I, 1, 1 - I/2, E^{((2*I)*\operatorname{ArcCsc}[a*x])}]$

Rule 12

$\operatorname{Int}[(a_*)(u_), x_Symbol] := \operatorname{Dist}[a, \operatorname{Int}[u, x], x] /; \operatorname{FreeQ}[a, x] \&& \operatorname{!MatchQ}[u, (b_*)(v_) /; \operatorname{FreeQ}[b, x]]$

Rule 2225

$\operatorname{Int}[((F_)^((c_.)*(a_.) + (b_.)*(x_.)))^((n_.)), x_Symbol] := \operatorname{Simp}[(F^(c*(a + b*x)))^n/(b*c*n*\operatorname{Log}[F]), x] /; \operatorname{FreeQ}[\{F, a, b, c, n\}, x]$

Rule 2283

$\operatorname{Int}[((a_) + (b_*)(F_)^((e_.)*(c_.) + (d_.)*(x_.))))^((p_.)*(G_)^((h_.)*(f_.) + (g_.)*(x_.))), x_Symbol] := \operatorname{Simp}[a^p*(G^(h*(f + g*x))/(g*h*\operatorname{Log}[G]))*\text{Hypergeometric2F1}[-p, g*h*(\operatorname{Log}[G]/(d*e*\operatorname{Log}[F])), g*h*(\operatorname{Log}[G]/(d*e*\operatorname{Log}[F])) + 1, \operatorname{Simplify}[(-b/a)*F^(e*(c + d*x))]], x] /; \operatorname{FreeQ}[\{F, G, a, b, c, d, e, f, g, h, p\}, x] \&& (\operatorname{ILtQ}[p, 0] \|\operatorname{GtQ}[a, 0])$

Rule 4528

$\operatorname{Int}[\operatorname{Cot}[(d_.) + (e_.)*(x_.)]^((n_.)*(F_)^((c_.)*(a_.) + (b_.)*(x_.))), x_Symbol] := \operatorname{Dist}[(-I)^n, \operatorname{Int}[\operatorname{ExpandIntegrand}[F^(c*(a + b*x))]*((1 + E^(2*I*(d + e*x)))/x), x]] /; \operatorname{FreeQ}[\{F, G, a, b, c, d, e, f, g, h, p\}, x] \&& (\operatorname{ILtQ}[p, 0] \|\operatorname{GtQ}[a, 0])$

```
*x)))^n/(1 - E^(2*I*(d + e*x)))^n), x], x], x] /; FreeQ[{F, a, b, c, d, e}, x] && IntegerQ[n]
```

Rule 5375

```
Int[(u_)*(f_)^(ArcCsc[(a_.) + (b_.)*(x_)]^(n_.)*(c_.)), x_Symbol] :> Dist[-b^(-1), Subst[Int[(u /. x -> -a/b + Csc[x]/b)*f^(c*x^n)*Csc[x]*Cot[x], x], ArcCsc[a + b*x]], x] /; FreeQ[{a, b, c, f}, x] && IGtQ[n, 0]
```

Rubi steps

$$\begin{aligned} \int \frac{e^{\csc^{-1}(ax)}}{x} dx &= -\frac{\text{Subst}\left(\int ae^x \cot(x) dx, x, \csc^{-1}(ax)\right)}{a} \\ &= -\text{Subst}\left(\int e^x \cot(x) dx, x, \csc^{-1}(ax)\right) \\ &= i\text{Subst}\left(\int \left(-e^x - \frac{2e^x}{-1+e^{2ix}}\right) dx, x, \csc^{-1}(ax)\right) \\ &= -\left(i\text{Subst}\left(\int e^x dx, x, \csc^{-1}(ax)\right)\right) - 2i\text{Subst}\left(\int \frac{e^x}{-1+e^{2ix}} dx, x, \csc^{-1}(ax)\right) \\ &= -ie^{\csc^{-1}(ax)} + 2ie^{\csc^{-1}(ax)} {}_2F_1\left(-\frac{i}{2}, 1; 1 - \frac{i}{2}; e^{2i\csc^{-1}(ax)}\right) \end{aligned}$$

Mathematica [A]

time = 0.04, size = 75, normalized size = 1.74

$$-i\left(-e^{\csc^{-1}(ax)} {}_2F_1\left(-\frac{i}{2}, 1; 1 - \frac{i}{2}; e^{2i\csc^{-1}(ax)}\right) - \left(\frac{1}{5} - \frac{2i}{5}\right) e^{(1+2i)\csc^{-1}(ax)} {}_2F_1\left(1, 1 - \frac{i}{2}; 2 - \frac{i}{2}; e^{2i\csc^{-1}(ax)}\right)\right)$$

Antiderivative was successfully verified.

[In] `Integrate[E^ArcCsc[a*x]/x, x]`

[Out] `(-I)*(-(E^ArcCsc[a*x]*Hypergeometric2F1[-1/2*I, 1, 1 - I/2, E^((2*I)*ArcCsc[a*x])]) - (1/5 - (2*I)/5)*E^((1 + 2*I)*ArcCsc[a*x])*Hypergeometric2F1[1, 1 - I/2, 2 - I/2, E^((2*I)*ArcCsc[a*x])])`

Maple [F]

time = 0.04, size = 0, normalized size = 0.00

$$\int \frac{e^{\arccsc(ax)}}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(arccsc(a*x))/x,x)`

[Out] `int(exp(arccsc(a*x))/x,x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(arccsc(a*x))/x,x, algorithm="maxima")`

[Out] `integrate(e^(arccsc(a*x))/x, x)`

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(arccsc(a*x))/x,x, algorithm="fricas")`

[Out] `integral(e^(arccsc(a*x))/x, x)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{\operatorname{acsc}(ax)}}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(acsc(a*x))/x,x)`

[Out] `Integral(exp(acsc(a*x))/x, x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(arccsc(a*x))/x,x, algorithm="giac")`

[Out] `integrate(e^(arccsc(a*x))/x, x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{e^{\operatorname{asin}\left(\frac{1}{ax}\right)}}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(asin(1/(a*x)))/x,x)`
[Out] `int(exp(asin(1/(a*x)))/x, x)`

3.45 $\int \frac{e^{\csc^{-1}(ax)}}{x^2} dx$

Optimal. Leaf size=39

$$-\frac{1}{2}ae^{\csc^{-1}(ax)}\sqrt{1 - \frac{1}{a^2x^2}} - \frac{e^{\csc^{-1}(ax)}}{2x}$$

[Out] $-1/2*\exp(\operatorname{arccsc}(a*x))/x - 1/2*a*\exp(\operatorname{arccsc}(a*x))*(1 - 1/a^2/x^2)^{(1/2)}$

Rubi [A]

time = 0.02, antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.300, Rules used = {5375, 12, 4518}

$$-\frac{1}{2}a\sqrt{1 - \frac{1}{a^2x^2}} e^{\csc^{-1}(ax)} - \frac{e^{\csc^{-1}(ax)}}{2x}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{\wedge} \operatorname{ArcCsc}[a*x]/x^2, x]$

[Out] $-1/2*(a*E^{\wedge} \operatorname{ArcCsc}[a*x]*\operatorname{Sqrt}[1 - 1/(a^2*x^2)]) - E^{\wedge} \operatorname{ArcCsc}[a*x]/(2*x)$

Rule 12

```
Int[(a_)*(u_), x_Symbol] :> Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]
```

Rule 4518

```
Int[Cos[d_.] + (e_.)*(x_.)]*(F_)^((c_.)*((a_.) + (b_.)*(x_))), x_Symbol] :>
Simp[b*c*Log[F]*F^(c*(a + b*x))*(Cos[d + e*x]/(e^2 + b^2*c^2*Log[F]^2)), x] +
Simp[e*F^(c*(a + b*x))*(Sin[d + e*x]/(e^2 + b^2*c^2*Log[F]^2)), x] /; FreeQ[{F, a, b, c, d, e}, x] && NeQ[e^2 + b^2*c^2*Log[F]^2, 0]
```

Rule 5375

```
Int[(u_.)*(f_)^(ArcCsc[(a_.) + (b_.)*(x_.)]^(n_.)*(c_.)), x_Symbol] :> Dist[-b^(-1), Subst[Int[(u /. x -> -a/b + Csc[x]/b)*f^(c*x^n)*Csc[x]*Cot[x], x], x, ArcCsc[a + b*x]], x] /; FreeQ[{a, b, c, f}, x] && IGtQ[n, 0]
```

Rubi steps

$$\begin{aligned} \int \frac{e^{\csc^{-1}(ax)}}{x^2} dx &= -\frac{\text{Subst}\left(\int a^2 e^x \cos(x) dx, x, \csc^{-1}(ax)\right)}{a} \\ &= -\left(a \text{Subst}\left(\int e^x \cos(x) dx, x, \csc^{-1}(ax)\right)\right) \\ &= -\frac{1}{2} a e^{\csc^{-1}(ax)} \sqrt{1 - \frac{1}{a^2 x^2}} - \frac{e^{\csc^{-1}(ax)}}{2x} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 33, normalized size = 0.85

$$-\frac{1}{2} a e^{\csc^{-1}(ax)} \left(\sqrt{1 - \frac{1}{a^2 x^2}} + \frac{1}{ax} \right)$$

Antiderivative was successfully verified.

[In] `Integrate[E^ArcCsc[a*x]/x^2, x]`[Out] `-1/2*(a*E^ArcCsc[a*x]*(Sqrt[1 - 1/(a^2*x^2)] + 1/(a*x)))`**Maple [F]**

time = 0.04, size = 0, normalized size = 0.00

$$\int \frac{e^{\arccsc(ax)}}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(arccsc(a*x))/x^2, x)`[Out] `int(exp(arccsc(a*x))/x^2, x)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(arccsc(a*x))/x^2, x, algorithm="maxima")`[Out] `integrate(e^(arccsc(a*x))/x^2, x)`**Fricas [A]**

time = 0.40, size = 23, normalized size = 0.59

$$-\frac{\left(\sqrt{a^2 x^2 - 1} + 1\right) e^{(\arccsc(ax))}}{2x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(arccsc(a*x))/x^2,x, algorithm="fricas")`
[Out] $-1/2 * (\sqrt{a^2 * x^2 - 1} + 1) * e^{arccsc(a*x)} / x$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{\operatorname{acsc}(ax)}}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(acsc(a*x))/x**2,x)`
[Out] `Integral(exp(acsc(a*x))/x**2, x)`

Giac [A]

time = 0.45, size = 42, normalized size = 1.08

$$-\frac{1}{2} \left(\sqrt{-\frac{1}{a^2 x^2} + 1} e^{(\arcsin(\frac{1}{ax}))} + \frac{e^{(\arcsin(\frac{1}{ax}))}}{ax} \right) a$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(arccsc(a*x))/x^2,x, algorithm="giac")`
[Out] $-1/2 * (\sqrt{-1/(a^2 * x^2) + 1} * e^{(\arcsin(1/(a*x)))} + e^{(\arcsin(1/(a*x)))} / (a * x)) * a$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{e^{\operatorname{asin}(\frac{1}{ax})}}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(asin(1/(a*x)))/x^2,x)`
[Out] `int(exp(asin(1/(a*x)))/x^2, x)`

3.46 $\int \frac{e^{\csc^{-1}(ax)}}{x^3} dx$

Optimal. Leaf size=41

$$\frac{1}{5}a^2 e^{\csc^{-1}(ax)} \cos(2 \csc^{-1}(ax)) - \frac{1}{10}a^2 e^{\csc^{-1}(ax)} \sin(2 \csc^{-1}(ax))$$

[Out] $1/5*a^2*\exp(\operatorname{arccsc}(a*x))*\cos(2*\operatorname{arccsc}(a*x))-1/10*a^2*\exp(\operatorname{arccsc}(a*x))*\sin(2*\operatorname{arccsc}(a*x))$

Rubi [A]

time = 0.03, antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.400, Rules used = {5375, 12, 4557, 4517}

$$\frac{1}{5}a^2 e^{\csc^{-1}(ax)} \cos(2 \csc^{-1}(ax)) - \frac{1}{10}a^2 e^{\csc^{-1}(ax)} \sin(2 \csc^{-1}(ax))$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{\operatorname{ArcCsc}[a*x]}/x^3, x]$

[Out] $(a^2 E^{\operatorname{ArcCsc}[a*x]} \cos[2 \operatorname{ArcCsc}[a*x]])/5 - (a^2 E^{\operatorname{ArcCsc}[a*x]} \sin[2 \operatorname{ArcCsc}[a*x]])/10$

Rule 12

$\operatorname{Int}[(a_*)*(u_), x_{\text{Symbol}}] := \operatorname{Dist}[a, \operatorname{Int}[u, x], x] /; \operatorname{FreeQ}[a, x] \&& \operatorname{!MatchQ}[u, (b_*)*(v_) /; \operatorname{FreeQ}[b, x]]$

Rule 4517

$\operatorname{Int}[(F_*)^{((c_*)*((a_*) + (b_*)*(x_))) * \sin((d_*) + (e_*)*(x_)), x_{\text{Symbol}}] := \operatorname{Simp}[b*c*\operatorname{Log}[F]*F^{(c*(a + b*x))} * (\sin[d + e*x]/(e^2 + b^2*c^2*\operatorname{Log}[F]^2)), x] - \operatorname{Simp}[e*F^{(c*(a + b*x))} * (\cos[d + e*x]/(e^2 + b^2*c^2*\operatorname{Log}[F]^2)), x] /; \operatorname{FreeQ}[\{F, a, b, c, d, e\}, x] \&& \operatorname{NeQ}[e^2 + b^2*c^2*\operatorname{Log}[F]^2, 0]$

Rule 4557

$\operatorname{Int}[\cos((f_*) + (g_*)*(x_))^{(n_*)} * (F_*)^{((c_*)*((a_*) + (b_*)*(x_))) * \sin((d_*) + (e_*)*(x_))^{(m_*)}, x_{\text{Symbol}}] := \operatorname{Int}[\operatorname{ExpandTrigReduce}[F^{(c*(a + b*x))}, \sin[d + e*x]^m * \cos[f + g*x]^n, x], x] /; \operatorname{FreeQ}[\{F, a, b, c, d, e, f, g\}, x] \&& \operatorname{IGtQ}[m, 0] \&& \operatorname{IGtQ}[n, 0]$

Rule 5375

$\operatorname{Int}[(u_*)*(f_*)^{(\operatorname{ArcCsc}[(a_*) + (b_*)*(x_)]^{(n_*)} * (c_*)}), x_{\text{Symbol}}] := \operatorname{Dist}[-b^{(-1)}, \operatorname{Subst}[\operatorname{Int}[(u /. x \rightarrow -a/b + \operatorname{Csc}[x]/b) * f^{(c*x^n)} * \operatorname{Csc}[x] * \operatorname{Cot}[x], x],$

$x, \text{ArcCsc}[a + b*x], x] /; \text{FreeQ}[\{a, b, c, f\}, x] \&& \text{IGtQ}[n, 0]$

Rubi steps

$$\begin{aligned} \int \frac{e^{\csc^{-1}(ax)}}{x^3} dx &= -\frac{\text{Subst}\left(\int a^3 e^x \cos(x) \sin(x) dx, x, \csc^{-1}(ax)\right)}{a} \\ &= -\left(a^2 \text{Subst}\left(\int e^x \cos(x) \sin(x) dx, x, \csc^{-1}(ax)\right)\right) \\ &= -\left(a^2 \text{Subst}\left(\int \frac{1}{2} e^x \sin(2x) dx, x, \csc^{-1}(ax)\right)\right) \\ &= -\left(\frac{1}{2} a^2 \text{Subst}\left(\int e^x \sin(2x) dx, x, \csc^{-1}(ax)\right)\right) \\ &= \frac{1}{5} a^2 e^{\csc^{-1}(ax)} \cos(2 \csc^{-1}(ax)) - \frac{1}{10} a^2 e^{\csc^{-1}(ax)} \sin(2 \csc^{-1}(ax)) \end{aligned}$$

Mathematica [A]

time = 0.04, size = 30, normalized size = 0.73

$$-\frac{1}{10} a^2 e^{\csc^{-1}(ax)} (-2 \cos(2 \csc^{-1}(ax)) + \sin(2 \csc^{-1}(ax)))$$

Antiderivative was successfully verified.

[In] `Integrate[E^ArcCsc[a*x]/x^3,x]`

[Out] `-1/10*(a^2 E^ArcCsc[a*x]*(-2*Cos[2*ArcCsc[a*x]] + Sin[2*ArcCsc[a*x]]))`

Maple [F]

time = 0.04, size = 0, normalized size = 0.00

$$\int \frac{e^{\arccsc(ax)}}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(arccsc(a*x))/x^3,x)`

[Out] `int(exp(arccsc(a*x))/x^3,x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(arccsc(a*x))/x^3,x, algorithm="maxima")`

[Out] `integrate(e^(arccsc(a*x))/x^3, x)`

Fricas [A]

time = 0.41, size = 32, normalized size = 0.78

$$\frac{\left(a^2 x^2 - \sqrt{a^2 x^2 - 1} - 2\right) e^{(\operatorname{arccsc}(ax))}}{5 x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(arccsc(a*x))/x^3,x, algorithm="fricas")`

[Out] `1/5*(a^2*x^2 - sqrt(a^2*x^2 - 1) - 2)*e^(arccsc(a*x))/x^2`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{\operatorname{acsc}(ax)}}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(acsc(a*x))/x**3,x)`

[Out] `Integral(exp(acsc(a*x))/x**3, x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(arccsc(a*x))/x^3,x, algorithm="giac")`

[Out] `integrate(e^(arccsc(a*x))/x^3, x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{e^{\operatorname{asin}\left(\frac{1}{ax}\right)}}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(asin(1/(a*x)))/x^3,x)`

[Out] `int(exp(asin(1/(a*x)))/x^3, x)`

$$\mathbf{3.47} \quad \int \frac{e^{\csc^{-1}(ax)}}{x^4} dx$$

Optimal. Leaf size=84

$$-\frac{1}{8}a^3e^{\csc^{-1}(ax)}\sqrt{1-\frac{1}{a^2x^2}}-\frac{a^2e^{\csc^{-1}(ax)}}{8x}+\frac{1}{40}a^3e^{\csc^{-1}(ax)}\cos(3\csc^{-1}(ax))+\frac{3}{40}a^3e^{\csc^{-1}(ax)}\sin(3\csc^{-1}(ax))$$

[Out] $-1/8*a^2*\exp(\operatorname{arccsc}(a*x))/x+1/40*a^3*\exp(\operatorname{arccsc}(a*x))*\cos(3*\operatorname{arccsc}(a*x))+3/40*a^3*\exp(\operatorname{arccsc}(a*x))*\sin(3*\operatorname{arccsc}(a*x))-1/8*a^3*\exp(\operatorname{arccsc}(a*x))*(1-1/a^2/x^2)^{(1/2)}$

Rubi [A]

time = 0.05, antiderivative size = 84, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.400, Rules used = {5375, 12, 4557, 4518}

$$\frac{1}{40}a^3e^{\csc^{-1}(ax)}\cos(3\csc^{-1}(ax))+\frac{3}{40}a^3e^{\csc^{-1}(ax)}\sin(3\csc^{-1}(ax))-\frac{a^2e^{\csc^{-1}(ax)}}{8x}-\frac{1}{8}a^3\sqrt{1-\frac{1}{a^2x^2}}e^{\csc^{-1}(ax)}$$

Antiderivative was successfully verified.

[In] Int[E^ArcCsc[a*x]/x^4, x]

[Out] $-1/8*(a^3E^{\operatorname{ArcCsc}[a*x]}*\operatorname{Sqrt}[1 - 1/(a^2*x^2)]) - (a^2E^{\operatorname{ArcCsc}[a*x]})/(8*x) + (a^3E^{\operatorname{ArcCsc}[a*x]}*\operatorname{Cos}[3*\operatorname{ArcCsc}[a*x]])/40 + (3*a^3E^{\operatorname{ArcCsc}[a*x]}*\operatorname{Sin}[3*\operatorname{ArcCsc}[a*x]])/40$

Rule 12

```
Int[(a_)*(u_), x_Symbol] :> Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]
```

Rule 4518

```
Int[Cos[(d_.) + (e_.)*(x_)]*(F_)^((c_.)*((a_.) + (b_.)*(x_))), x_Symbol] :>
Simp[b*c*Log[F]*F^(c*(a + b*x))*(Cos[d + e*x]/(e^2 + b^2*c^2*Log[F]^2)), x] +
Simp[e*F^(c*(a + b*x))*(Sin[d + e*x]/(e^2 + b^2*c^2*Log[F]^2)), x] /; FreeQ[{F, a, b, c, d, e}, x] && NeQ[e^2 + b^2*c^2*Log[F]^2, 0]
```

Rule 4557

```
Int[Cos[(f_.) + (g_.)*(x_)]^(n_.)*(F_)^((c_.)*((a_.) + (b_.)*(x_)))*Sin[(d_.) + (e_.)*(x_)]^(m_.), x_Symbol] :> Int[ExpandTrigReduce[F^(c*(a + b*x)), Sin[d + e*x]^m*Cos[f + g*x]^n, x], x] /; FreeQ[{F, a, b, c, d, e, f, g}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rule 5375

```
Int[(u_)*(f_)^(ArcCsc[(a_.) + (b_.)*(x_.)]^(n_.)*(c_.)), x_Symbol] :> Dist[
-b^(-1), Subst[Int[(u /. x -> -a/b + Csc[x]/b)*f^(c*x^n)*Csc[x]*Cot[x], x],
x, ArcCsc[a + b*x]], x] /; FreeQ[{a, b, c, f}, x] && IGtQ[n, 0]
```

Rubi steps

$$\begin{aligned} \int \frac{e^{\csc^{-1}(ax)}}{x^4} dx &= -\frac{\text{Subst}\left(\int a^4 e^x \cos(x) \sin^2(x) dx, x, \csc^{-1}(ax)\right)}{a} \\ &= -\left(a^3 \text{Subst}\left(\int e^x \cos(x) \sin^2(x) dx, x, \csc^{-1}(ax)\right)\right) \\ &= -\left(a^3 \text{Subst}\left(\int \left(\frac{1}{4}e^x \cos(x) - \frac{1}{4}e^x \cos(3x)\right) dx, x, \csc^{-1}(ax)\right)\right) \\ &= -\left(\frac{1}{4}a^3 \text{Subst}\left(\int e^x \cos(x) dx, x, \csc^{-1}(ax)\right)\right) + \frac{1}{4}a^3 \text{Subst}\left(\int e^x \cos(3x) dx, x, \csc^{-1}(ax)\right) \\ &= -\frac{1}{8}a^3 e^{\csc^{-1}(ax)} \sqrt{1 - \frac{1}{a^2 x^2}} - \frac{a^2 e^{\csc^{-1}(ax)}}{8x} + \frac{1}{40}a^3 e^{\csc^{-1}(ax)} \cos(3 \csc^{-1}(ax)) + \frac{3}{40}a^3 e^{\csc^{-1}(ax)} \sin(3 \csc^{-1}(ax)) \end{aligned}$$

Mathematica [A]

time = 0.10, size = 54, normalized size = 0.64

$$\frac{1}{40}a^3 e^{\csc^{-1}(ax)} \left(-5\sqrt{1 - \frac{1}{a^2 x^2}} - \frac{5}{ax} + \cos(3 \csc^{-1}(ax)) + 3 \sin(3 \csc^{-1}(ax)) \right)$$

Antiderivative was successfully verified.

[In] `Integrate[E^ArcCsc[a*x]/x^4, x]`

[Out] `(a^3 E^ArcCsc[a*x]*(-5*Sqrt[1 - 1/(a^2*x^2)] - 5/(a*x) + Cos[3*ArcCsc[a*x]] + 3*Sin[3*ArcCsc[a*x]]))/40`

Maple [F]

time = 0.04, size = 0, normalized size = 0.00

$$\int \frac{e^{\arccsc(ax)}}{x^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(arccsc(a*x))/x^4, x)`

[Out] `int(exp(arccsc(a*x))/x^4, x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(arccsc(a*x))/x^4,x, algorithm="maxima")`
[Out] `integrate(e^(arccsc(a*x))/x^4, x)`

Fricas [A]

time = 0.42, size = 41, normalized size = 0.49

$$\frac{\left(a^2 x^2 - (a^2 x^2 + 1) \sqrt{a^2 x^2 - 1} - 3\right) e^{(\operatorname{arccsc}(ax))}}{10 x^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(arccsc(a*x))/x^4,x, algorithm="fricas")`
[Out] `1/10*(a^2*x^2 - (a^2*x^2 + 1)*sqrt(a^2*x^2 - 1) - 3)*e^(arccsc(a*x))/x^3`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{\operatorname{acsc}(ax)}}{x^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(acsc(a*x))/x**4,x)`
[Out] `Integral(exp(acsc(a*x))/x**4, x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(arccsc(a*x))/x^4,x, algorithm="giac")`
[Out] `integrate(e^(arccsc(a*x))/x^4, x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{e^{\operatorname{asin}\left(\frac{1}{ax}\right)}}{x^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(asin(1/(a*x)))/x^4,x)`
[Out] `int(exp(asin(1/(a*x)))/x^4, x)`

3.48 $\int \frac{e^{\csc^{-1}(ax)}}{x^5} dx$

Optimal. Leaf size=81

$$\frac{1}{10}a^4 e^{\csc^{-1}(ax)} \cos(2 \csc^{-1}(ax)) - \frac{1}{34}a^4 e^{\csc^{-1}(ax)} \cos(4 \csc^{-1}(ax)) - \frac{1}{20}a^4 e^{\csc^{-1}(ax)} \sin(2 \csc^{-1}(ax)) + \frac{1}{136}a^4 e^{\csc^{-1}(ax)} \sin(4 \csc^{-1}(ax))$$

[Out] $1/10*a^4*\exp(\text{arccsc}(a*x))*\cos(2*\text{arccsc}(a*x))-1/34*a^4*\exp(\text{arccsc}(a*x))*\cos(4*\text{arccsc}(a*x))-1/20*a^4*\exp(\text{arccsc}(a*x))*\sin(2*\text{arccsc}(a*x))+1/136*a^4*\exp(a*\text{rccsc}(a*x))*\sin(4*\text{arccsc}(a*x))$

Rubi [A]

time = 0.05, antiderivative size = 81, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.400, Rules used = {5375, 12, 4557, 4517}

$$\frac{1}{10}a^4 e^{\csc^{-1}(ax)} \cos(2 \csc^{-1}(ax)) - \frac{1}{34}a^4 e^{\csc^{-1}(ax)} \cos(4 \csc^{-1}(ax)) - \frac{1}{20}a^4 e^{\csc^{-1}(ax)} \sin(2 \csc^{-1}(ax)) + \frac{1}{136}a^4 e^{\csc^{-1}(ax)} \sin(4 \csc^{-1}(ax))$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{\wedge} \text{ArcCsc}[a*x]/x^5, x]$

[Out] $(a^4 E^{\wedge} \text{ArcCsc}[a*x] * \text{Cos}[2 \text{ArcCsc}[a*x]])/10 - (a^4 E^{\wedge} \text{ArcCsc}[a*x] * \text{Cos}[4 \text{ArcCsc}[a*x]])/34 - (a^4 E^{\wedge} \text{ArcCsc}[a*x] * \text{Sin}[2 \text{ArcCsc}[a*x]])/20 + (a^4 E^{\wedge} \text{ArcCsc}[a*x] * \text{Sin}[4 \text{ArcCsc}[a*x]])/136$

Rule 12

$\text{Int}[(a_*)*(u_), x_Symbol] :> \text{Dist}[a, \text{Int}[u, x], x] /; \text{FreeQ}[a, x] \&& \text{!MatchQ}[u, (b_*)*(v_) /; \text{FreeQ}[b, x]]$

Rule 4517

$\text{Int}[(F_*)^{((c_*)*((a_*) + (b_*)*(x_*)))*\text{Sin}[(d_*) + (e_*)*(x_*)], x_Symbol] :> \text{Simp}[b*c*\text{Log}[F]*F^{(c*(a + b*x))*(\text{Sin}[d + e*x]/(e^2 + b^2*c^2*\text{Log}[F]^2))}, x] - \text{Simp}[e*F^{(c*(a + b*x))*(\text{Cos}[d + e*x]/(e^2 + b^2*c^2*\text{Log}[F]^2))}, x] /; \text{FreeQ}[\{F, a, b, c, d, e\}, x] \&& \text{NeQ}[e^2 + b^2*c^2*\text{Log}[F]^2, 0]$

Rule 4557

$\text{Int}[\text{Cos}[(f_*) + (g_*)*(x_*)]^{(n_*)}*(F_*)^{((c_*)*((a_*) + (b_*)*(x_*)))*\text{Sin}[(d_*) + (e_*)*(x_*)]^{(m_*)}, x_Symbol] :> \text{Int}[\text{ExpandTrigReduce}[F^{(c*(a + b*x)), \text{Sin}[d + e*x]^m*\text{Cos}[f + g*x]^n, x], x] /; \text{FreeQ}[\{F, a, b, c, d, e, f, g\}, x] \&& \text{IGtQ}[m, 0] \&& \text{IGtQ}[n, 0]$

Rule 5375

```
Int[(u_)*(f_)^(ArcCsc[(a_.) + (b_.)*(x_)]^(n_.)*(c_.)), x_Symbol] :> Dist[
-b^(-1), Subst[Int[(u /. x -> -a/b + Csc[x]/b)*f^(c*x^n)*Csc[x]*Cot[x], x],
x, ArcCsc[a + b*x]], x] /; FreeQ[{a, b, c, f}, x] && IGtQ[n, 0]
```

Rubi steps

$$\begin{aligned} \int \frac{e^{\csc^{-1}(ax)}}{x^5} dx &= -\frac{\text{Subst}\left(\int a^5 e^x \cos(x) \sin^3(x) dx, x, \csc^{-1}(ax)\right)}{a} \\ &= -\left(a^4 \text{Subst}\left(\int e^x \cos(x) \sin^3(x) dx, x, \csc^{-1}(ax)\right)\right) \\ &= -\left(a^4 \text{Subst}\left(\int \left(\frac{1}{4}e^x \sin(2x) - \frac{1}{8}e^x \sin(4x)\right) dx, x, \csc^{-1}(ax)\right)\right) \\ &= \frac{1}{8}a^4 \text{Subst}\left(\int e^x \sin(4x) dx, x, \csc^{-1}(ax)\right) - \frac{1}{4}a^4 \text{Subst}\left(\int e^x \sin(2x) dx, x, \csc^{-1}(ax)\right) \\ &= \frac{1}{10}a^4 e^{\csc^{-1}(ax)} \cos(2 \csc^{-1}(ax)) - \frac{1}{34}a^4 e^{\csc^{-1}(ax)} \cos(4 \csc^{-1}(ax)) - \frac{1}{20}a^4 e^{\csc^{-1}(ax)} \sin(2 \csc^{-1}(ax)) \end{aligned}$$

Mathematica [A]

time = 0.09, size = 50, normalized size = 0.62

$$-\frac{1}{680}a^4 e^{\csc^{-1}(ax)} (-68 \cos(2 \csc^{-1}(ax)) + 20 \cos(4 \csc^{-1}(ax)) + 34 \sin(2 \csc^{-1}(ax)) - 5 \sin(4 \csc^{-1}(ax)))$$

Antiderivative was successfully verified.

[In] `Integrate[E^ArcCsc[a*x]/x^5,x]`

[Out] `-1/680*(a^4*E^ArcCsc[a*x]*(-68*Cos[2*ArcCsc[a*x]] + 20*Cos[4*ArcCsc[a*x]] + 34*Sin[2*ArcCsc[a*x]] - 5*Sin[4*ArcCsc[a*x]]))`

Maple [F]

time = 0.04, size = 0, normalized size = 0.00

$$\int \frac{e^{\arccsc(ax)}}{x^5} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(arccsc(a*x))/x^5,x)`

[Out] `int(exp(arccsc(a*x))/x^5,x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(arccsc(a*x))/x^5,x, algorithm="maxima")`

[Out] `integrate(e^(arccsc(a*x))/x^5, x)`

Fricas [A]

time = 0.40, size = 51, normalized size = 0.63

$$\frac{\left(6 a^4 x^4 + 3 a^2 x^2 - (6 a^2 x^2 + 5) \sqrt{a^2 x^2 - 1} - 20\right) e^{(\text{arccsc}(ax))}}{85 x^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(arccsc(a*x))/x^5,x, algorithm="fricas")`

[Out] `1/85*(6*a^4*x^4 + 3*a^2*x^2 - (6*a^2*x^2 + 5)*sqrt(a^2*x^2 - 1) - 20)*e^(arccsc(a*x))/x^4`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{\text{acsc}(ax)}}{x^5} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(acsc(a*x))/x**5,x)`

[Out] `Integral(exp(acsc(a*x))/x**5, x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(arccsc(a*x))/x^5,x, algorithm="giac")`

[Out] `integrate(e^(arccsc(a*x))/x^5, x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{e^{\text{asin}(\frac{1}{ax})}}{x^5} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(asin(1/(a*x)))/x^5,x)`

[Out] `int(exp(asin(1/(a*x)))/x^5, x)`

3.49 $\int \frac{\csc^{-1}(a+bx)}{\frac{ad}{b}+dx} dx$

Optimal. Leaf size=69

$$\frac{i \csc^{-1}(a+bx)^2}{2d} - \frac{\csc^{-1}(a+bx) \log\left(1 - e^{2i \csc^{-1}(a+bx)}\right)}{d} + \frac{i \text{PolyLog}\left(2, e^{2i \csc^{-1}(a+bx)}\right)}{2d}$$

[Out] $\frac{1}{2} \text{I} \cdot \text{arccsc}(b*x+a)^2/d - \text{arccsc}(b*x+a) \cdot \ln(1 - (I/(b*x+a) + (1 - 1/(b*x+a)^2)^{(1/2)})^2)/d + \frac{1}{2} \text{I} \cdot \text{polylog}(2, (I/(b*x+a) + (1 - 1/(b*x+a)^2)^{(1/2)})^2)/d$

Rubi [A]

time = 0.07, antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.421, Rules used = {5365, 12, 5327, 4721, 3798, 2221, 2317, 2438}

$$\frac{i \text{Li}_2\left(e^{2i \csc^{-1}(a+bx)}\right)}{2d} + \frac{i \csc^{-1}(a+bx)^2}{2d} - \frac{\csc^{-1}(a+bx) \log\left(1 - e^{2i \csc^{-1}(a+bx)}\right)}{d}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{ArcCsc}[a+b*x]/((a*d)/b+d*x), x]$

[Out] $\frac{((I/2)*\text{ArcCsc}[a+b*x]^2)/d - (\text{ArcCsc}[a+b*x]*\text{Log}[1 - E^{((2*I)*\text{ArcCsc}[a+b*x])}])/d + ((I/2)*\text{PolyLog}[2, E^{((2*I)*\text{ArcCsc}[a+b*x])}])/d}{d}$

Rule 12

$\text{Int}[(a_)*(u_), x_Symbol] \rightarrow \text{Dist}[a, \text{Int}[u, x], x] /; \text{FreeQ}[a, x] \&& \text{!MatchQ}[u, (b_)*(v_) /; \text{FreeQ}[b, x]]$

Rule 2221

$\text{Int}[(((F_)^((g_)*(e_)+(f_)*(x_))))^{(n_)}*((c_)+(d_)*(x_))^{(m_)})/((a_)+(b_)*(F_)^((g_)*(e_)+(f_)*(x_)))^{(n_)}], x_Symbol] \rightarrow \text{Simp}[((c+d*x)^m/(b*f*g*n*\text{Log}[F]))*\text{Log}[1 + b*((F^(g*(e+f*x)))^n/a)], x] - \text{Dist}[d*(m/(b*f*g*n*\text{Log}[F])), \text{Int}[(c+d*x)^(m-1)*\text{Log}[1 + b*((F^(g*(e+f*x)))^n/a)], x], x] /; \text{FreeQ}[\{F, a, b, c, d, e, f, g, n\}, x] \&& \text{IGtQ}[m, 0]$

Rule 2317

$\text{Int}[\text{Log}[(a_)+(b_)*(F_)^((e_)*(c_)+(d_)*(x_)))^{(n_)}], x_Symbol] \rightarrow \text{Dist}[1/(d*e*n*\text{Log}[F]), \text{Subst}[\text{Int}[\text{Log}[a+b*x]/x, x], x, (F^(e*(c+d*x)))^n], x] /; \text{FreeQ}[\{F, a, b, c, d, e, n\}, x] \&& \text{GtQ}[a, 0]$

Rule 2438

```
Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 3798

```
Int[((c_.) + (d_.)*(x_)^(m_.))*tan[(e_.) + Pi*(k_.) + (f_.)*(x_)], x_Symbol] :> Simp[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Dist[2*I, Int[(c + d*x)^m *E^(2*I*k*Pi)*(E^(2*I*(e + f*x))/(1 + E^(2*I*k*Pi)*E^(2*I*(e + f*x)))), x], x] /; FreeQ[{c, d, e, f}, x] && IntegerQ[4*k] && IGtQ[m, 0]
```

Rule 4721

```
Int[((a_.) + ArcSin[(c_.)*(x_)]*(b_.)^(n_.)/(x_), x_Symbol] :> Subst[Int[(a + b*x)^n*Cot[x], x], x, ArcSin[c*x]] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0]
```

Rule 5327

```
Int[((a_.) + ArcCsc[(c_.)*(x_)]*(b_.)/(x_), x_Symbol] :> -Subst[Int[(a + b *ArcSin[x/c])/x, x], x, 1/x] /; FreeQ[{a, b, c}, x]
```

Rule 5365

```
Int[((a_.) + ArcCsc[(c_) + (d_.)*(x_)]*(b_.)^(p_.)*((e_.) + (f_.)*(x_))^(m _.), x_Symbol] :> Dist[1/d, Subst[Int[(f*(x/d))^m*(a + b*ArcCsc[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[d*e - c*f, 0] && IGtQ[p, 0]
```

Rubi steps

$$\begin{aligned}
\int \frac{\csc^{-1}(a+bx)}{\frac{ad}{b} + dx} dx &= \frac{\text{Subst}\left(\int \frac{b \csc^{-1}(x)}{dx} dx, x, a+bx\right)}{b} \\
&= \frac{\text{Subst}\left(\int \frac{\csc^{-1}(x)}{x} dx, x, a+bx\right)}{d} \\
&= -\frac{\text{Subst}\left(\int \frac{\sin^{-1}(x)}{x} dx, x, \frac{1}{a+bx}\right)}{d} \\
&= -\frac{\text{Subst}\left(\int x \cot(x) dx, x, \sin^{-1}\left(\frac{1}{a+bx}\right)\right)}{d} \\
&= \frac{i \sin^{-1}\left(\frac{1}{a+bx}\right)^2}{2d} + \frac{(2i)\text{Subst}\left(\int \frac{e^{2ix}}{1-e^{2ix}} dx, x, \sin^{-1}\left(\frac{1}{a+bx}\right)\right)}{d} \\
&= \frac{i \sin^{-1}\left(\frac{1}{a+bx}\right)^2}{2d} - \frac{\sin^{-1}\left(\frac{1}{a+bx}\right) \log\left(1 - e^{2i \sin^{-1}\left(\frac{1}{a+bx}\right)}\right)}{d} + \frac{\text{Subst}\left(\int \log(1 - e^{2ix}) dx, x\right)}{d} \\
&= \frac{i \sin^{-1}\left(\frac{1}{a+bx}\right)^2}{2d} - \frac{\sin^{-1}\left(\frac{1}{a+bx}\right) \log\left(1 - e^{2i \sin^{-1}\left(\frac{1}{a+bx}\right)}\right)}{d} - \frac{i \text{Subst}\left(\int \frac{\log(1-x)}{x} dx, x, e^{2i \sin^{-1}\left(\frac{1}{a+bx}\right)}\right)}{2d} \\
&= \frac{i \sin^{-1}\left(\frac{1}{a+bx}\right)^2}{2d} - \frac{\sin^{-1}\left(\frac{1}{a+bx}\right) \log\left(1 - e^{2i \sin^{-1}\left(\frac{1}{a+bx}\right)}\right)}{d} + \frac{i \text{Li}_2\left(e^{2i \sin^{-1}\left(\frac{1}{a+bx}\right)}\right)}{2d}
\end{aligned}$$

Mathematica [A]

time = 0.04, size = 59, normalized size = 0.86

$$\frac{-\csc^{-1}(a+bx) \log\left(1 - e^{2i \csc^{-1}(a+bx)}\right) + \frac{1}{2}i \left(\csc^{-1}(a+bx)^2 + \text{PolyLog}\left(2, e^{2i \csc^{-1}(a+bx)}\right)\right)}{d}$$

Antiderivative was successfully verified.

[In] `Integrate[ArcCsc[a + b*x]/((a*d)/b + d*x), x]`[Out] `(-(ArcCsc[a + b*x]*Log[1 - E^((2*I)*ArcCsc[a + b*x])]) + (I/2)*(ArcCsc[a + b*x]^2 + PolyLog[2, E^((2*I)*ArcCsc[a + b*x])]))/d`**Maple [A]**

time = 0.88, size = 167, normalized size = 2.42

method	result
--------	--------

derivative divides	$\frac{\frac{b \operatorname{arccsc}(bx+a) \ln \left(1 - \frac{i}{bx+a} - \sqrt{1 - \frac{1}{(bx+a)^2}}\right)}{d} + \frac{i b \operatorname{polylog} \left(2, \frac{i}{bx+a} + \sqrt{1 - \frac{1}{(bx+a)^2}}\right)}{d}}{b}$
default	$\frac{\frac{b \operatorname{arccsc}(bx+a) \ln \left(1 - \frac{i}{bx+a} - \sqrt{1 - \frac{1}{(bx+a)^2}}\right)}{d} + \frac{i b \operatorname{polylog} \left(2, \frac{i}{bx+a} + \sqrt{1 - \frac{1}{(bx+a)^2}}\right)}{d}}{b}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(arccsc(b*x+a)/(a*d/b+d*x),x,method=_RETURNVERBOSE)`
[Out]
$$\frac{1}{b} \left(\frac{1}{2} I b/d \operatorname{arccsc}(b*x+a)^2 - b/d \operatorname{arccsc}(b*x+a) \ln \left(1 - \frac{I}{b*x+a} - (1 - 1/(b*x+a)^2)^{(1/2)} \right) + I b/d \operatorname{polylog} \left(2, \frac{I}{b*x+a} + (1 - 1/(b*x+a)^2)^{(1/2)} \right) - b/d \operatorname{arccsc}(b*x+a) \ln \left(1 + \frac{I}{b*x+a} + (1 - 1/(b*x+a)^2)^{(1/2)} \right) + I b/d \operatorname{polylog} \left(2, -\frac{I}{b*x+a} - (1 - 1/(b*x+a)^2)^{(1/2)} \right) \right)$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(b*x+a)/(a*d/b+d*x),x, algorithm="maxima")`
[Out]
$$\frac{1}{2} \left(2 b^2 d \operatorname{integrate}(\sqrt{b*x + a + 1} * \sqrt{b*x + a - 1} * \log(b*x + a) / (b^3 * d * x^3 + 3*a*b^2*d*x^2 + (3*a^2 - 1)*b*d*x + (a^3 - a)*d), x) - 2*I*b*d \operatorname{integrate}(\log(b*x + a) / (b^3*d*x^3 + 3*a*b^2*d*x^2 + (3*a^2 - 1)*b*d*x + (a^3 - a)*d), x) + (2*\operatorname{arctan2}(1, \sqrt{b*x + a + 1}) * \sqrt{b*x + a - 1}) + I * \log(-b*x - a + 1) * \log(b*x + a) - I * \log(b^2*x^2 + 2*a*b*x + a^2) * \log(b*x + a) + I * \log(b*x + a + 1) * \log(b*x + a) + I * \log(b*x + a)^2 + I * \operatorname{dilog}(b*x + a) + I * \operatorname{dilog}(-b*x - a) \right) / d$$

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(b*x+a)/(a*d/b+d*x),x, algorithm="fricas")`
[Out] `integral(b*arccsc(b*x + a)/(b*d*x + a*d), x)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\frac{b \int \frac{\operatorname{acsc}(a+bx)}{a+bx} dx}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(acsc(b*x+a)/(a*d/b+d*x),x)`
[Out] `b*Integral(acsc(a + b*x)/(a + b*x), x)/d`

Giac [A]

time = 0.66, size = 114, normalized size = 1.65

$$-\frac{1}{4} b^2 \left(\frac{\frac{2 (bx + a)^2 \arcsin\left(\frac{1}{((bx+a)\left(\frac{a}{bx+a}-1)-a)\left(\frac{a}{bx+a}-1)+a}\right)}{b^3 d} + \frac{(bx + a) \left(\sqrt{-\frac{1}{(bx + a)^2} + 1} - 1\right)}{b^3 d} - \frac{(bx + a) \left(\sqrt{-\frac{1}{(bx + a)^2} + 1} - 1\right)}{b^3 d}}{b^3 d} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(arccsc(b*x+a)/(a*d/b+d*x),x, algorithm="giac")`
[Out] `-1/4*b^2*(2*(b*x + a)^2*arcsin(1/(((b*x + a)*(a/(b*x + a) - 1) - a)*(a/(b*x + a) - 1) + a))/(b^3*d) + ((b*x + a)*(sqrt(-1/(b*x + a)^2 + 1) - 1) - 1/((b*x + a)*(sqrt(-1/(b*x + a)^2 + 1) - 1)))/(b^3*d))`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\operatorname{asin}\left(\frac{1}{a+bx}\right)}{dx + \frac{ad}{b}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(asin(1/(a + b*x))/(d*x + (a*d)/b),x)`
[Out] `int(asin(1/(a + b*x))/(d*x + (a*d)/b), x)`

Chapter 4

Appendix

Local contents

4.1	Download section	264
4.2	Listing of Grading functions	264

4.1 Download section

The following zip files contain the raw integrals used in this test.

Mathematica format Mathematica_syntax.zip

Maple and Mupad format Maple_syntax.zip

Sympy format SYMPY_syntax.zip

Sage math format SAGE_syntax.zip

4.2 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

4.2.1 Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(* Small rewrite of logic in main function to make it*)
(* match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)
```

```

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafC
expnResult = ExpnType[result];
expnOptimal = ExpnType[optimal];
leafCountResult = LeafCount[result];
leafCountOptimal = LeafCount[optimal];

(*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
If[expnResult<=expnOptimal,
  If[Not[FreeQ[result,Complex]], (*result contains complex*)
    If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
      If[leafCountResult<=2*leafCountOptimal,
        finalresult={"A","none"}
        ,(*ELSE*)
        finalresult={"B","Both result and optimal contain complex but leaf count
      ]
      ,(*ELSE*)
      finalresult={"C","Result contains complex when optimal does not."}
    ]
    ,(*ELSE*)(*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A","none"}
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal. $"
    ]
    ]
  ,(*ELSE*)(*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order "<
    ,
    finalresult={"F","Contains unresolved integral."}
  ]
];
finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hypergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)

```

```
(*9 = unknown function*)
```

```
ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
  If[ListQ[expn],
    Max[Map[ExpnType, expn]],
  If[Head[expn] === Power,
    If[IntegerQ[expn[[2]]],
      ExpnType[expn[[1]]],
    If[Head[expn[[2]]] === Rational,
      If[IntegerQ[expn[[1]]] || Head[expn[[1]]] === Rational,
        1,
      Max[ExpnType[expn[[1]]], 2]],
      Max[ExpnType[expn[[1]]], ExpnType[expn[[2]], 3]]],
    If[Head[expn] === Plus || Head[expn] === Times,
      Max[ExpnType[First[expn]], ExpnType[Rest[expn]]],
    If[ElementaryFunctionQ[Head[expn]],
      Max[3, ExpnType[expn[[1]]]],
    If[SpecialFunctionQ[Head[expn]],
      Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 4]],
    If[HypergeometricFunctionQ[Head[expn]],
      Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 5]],
    If[AppellFunctionQ[Head[expn]],
      Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 6]],
    If[Head[expn] === RootSum,
      Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 7]],
    If[Head[expn] === Integrate || Head[expn] === Int,
      Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 8]],
    9]]]]]]]]]]]
```

```
ElementaryFunctionQ[func_] :=
  MemberQ[{  

    Exp, Log,  

    Sin, Cos, Tan, Cot, Sec, Csc,  

    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,  

    Sinh, Cosh, Tanh, Coth, Sech, Csch,  

    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch  

  }, func]
```

```
SpecialFunctionQ[func_] :=
  MemberQ[{  

    Erf, Erfc, Erfi,  

    FresnelS, FresnelC,
```

```

ExpIntegralE, ExpIntegralEi, LogIntegral,
SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
Gamma, LogGamma, PolyGamma,
Zeta, PolyLog, ProductLog,
EllipticF, EllipticE, EllipticPi
},func]

HypergeometricFunctionQ[func_] :=
MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

AppellFunctionQ[func_] :=
MemberQ[{AppellF1}, func]

```

4.2.2 Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#           if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#           see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
      #do NOT call ExpnType() if leaf size is too large. Recursion problem
      if leaf_count_result > 500000 then
          return "B","result has leaf size over 500,000. Avoiding possible recursion issues
      fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

```

```

if debug then
    print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A","");
            else
                return "B",cat("Both result and optimal contain complex but leaf count of r
                                convert(leaf_count_result,string)," vs. $2 (
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf_c
                end if
            else #result contains complex but optimal is not
                if debug then
                    print("result contains complex but optimal is not");
                fi;
                return "C","Result contains complex when optimal does not.";
            fi;
        else # result do not contain complex
    fi;
fi;

```

```

        # this assumes optimal do not as well. No check is needed here.
        if debug then
            print("result do not contain complex, this assumes optimal do not as well")
        fi;
        if leaf_count_result<=2*leaf_count_optimal then
            if debug then
                print("leaf_count_result<=2*leaf_count_optimal");
            fi;
            return "A","");
        else
            if debug then
                print("leaf_count_result>2*leaf_count_optimal");
            fi;
            return "B",cat("Leaf count of result is larger than twice the leaf count of optimal",
                           convert(leaf_count_result,string)," vs. $2(",
                           convert(leaf_count_optimal,string),")=",convert(2*leaf_count_optimal,string));
        fi;
    fi;
else #ExpnType(result) > ExpnType(optimal)
    if debug then
        print("ExpnType(result) > ExpnType(optimal)");
    fi;
    return "C",cat("Result contains higher order function than in optimal. Order ",
                   convert(ExpnType_result,string)," vs. order ",
                   convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hypergeometric function
# 6 = appell function
# 7 = rootsum function

```

```

# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'`^`) then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'`+`) or type(expn,'`*`) then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
  9
  end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func,[
    exp,log,ln,
    sin,cos,tan,cot,sec,csc,

```

```

arcsin,arccos,arctan,arccot,arcsec,arccsc,
sinh,cosh,tanh,coth,sech,csch,
arcsinh,arccosh,arctanh,arccoth,arcsech,arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func,[
    erf,erfc,erfi,
    FresnelS,FresnelC,
    Ei,Ei,Li,Si,Ci,Shi,Chi,
    GAMMA,lnGAMMA,Psi,Zeta,polylog,dilog,LambertW,
    EllipticF,EllipticE,EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func,[Hypergeometric1F1,hypergeom,HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func,[AppellF1])
end proc:

# u is a sum or product.  rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u),op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc:

```

4.2.3 Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#          Port of original Maple grading function by
#          Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#          added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):
    if isinstance(expr,Pow):
        if expr.args[1] == Rational(1,2):
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
                   asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
                   asinh,acosh,atanh,acoth,asech,acsch
                   ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
                    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
                    gamma,loggamma,digamma,zeta,polylog,LambertW,
                    elliptic_f,elliptic_e,elliptic_pi,exp_polar
                    ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False
    except:
        return False
```

```

except AttributeError as error:
    return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn))  #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow):  #type(expn,'`^`)
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnT
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'`+') or type(expn,'`*`)
        m1 = expnType(expn.args[0])
        m2 = expnType(list(expn.args[1:]))
        return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
    elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
        return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
    elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
    elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif is_appell_function(expn.func):
        m1 = max(map(expnType, list(expn.args)))
        return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif isinstance(expn,RootSum):
        m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
        return max(7,m1)
    elif str(expn).find("Integral") != -1:

```

```

m1 = max(map(expnType, list(expn.args)))
    return max(8,m1)  #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

#print ("Enter grade_antiderivative for sageMath")
#print("Enter grade_antiderivative, result=",result, " optimal=",optimal)

leaf_count_result = leaf_count(result)
leaf_count_optimal = leaf_count(optimal)

#print("leaf_count_result=",leaf_count_result)
#print("leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if str(result).find("Integral") != -1:
    grade = "F"
    grade_annotation = ""
else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation ="Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal."
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation ="Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = ""
            else:
                grade = "B"
                grade_annotation ="Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count(result)-2*leaf_count(optimal))
    else:
        grade = "C"
        grade_annotation ="Result contains higher order function than in optimal. Order "+str(ExpnType_result-ExpnType_optimal)

```

```
#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation
```

4.2.4 SageMath grading function

```
#Dec 24, 2019. Nasser: Ported original Maple grading function by
#           Albert Rich to use with Sagemath. This is used to
#           grade Fricas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#           'arctan2','floor','abs','log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#           issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    """
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow:  #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False
```

```

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arccoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:
            print ("func ", func , " is elementary_function")
        else:
            print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']  #[appellf1] can't find this in sage

```

```

def is_atom(expn):

    #debug=False
    if debug:
        print ("Enter is_atom, expn=",expn)

    if not hasattr(expn, 'parent'):
        return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(),"base_ring") and hasattr(expn.parent(),"gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__)
    return False


def expnType(expn):

    if debug:
        print (">>>>Enter expnType, expn=", expn)
        print (">>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],Rational)):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #isinstance(expn,Pow)
        if type(expn.operands()[1])==Integer: #isinstance(expn.args[1],Integer)
            return expnType(expn.operands()[0]) #expnType(expn.args[0])
        elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
            if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)

```

```

        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    else:
        return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn.op
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isinstan
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    if debug:
        print ("Enter grade_antiderivative for sagemath")
        print("Enter grade_antiderivative, result=",result)
        print("Enter grade_antiderivative, optimal=",optimal)
        print("type(anti)=",type(result))
        print("type(optimal)=",type(optimal))

        leaf_count_result = tree_size(result) #leaf_count(result)
        leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

        #if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

```

```

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger than optimal"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_result)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```