

Computer algebra independent integration tests

Summer 2022 edition

6-Hyperbolic-functions/6.3-Hyperbolic-tangent/172-6.3.2-
Hyperbolic-tangent-functions

Nasser M. Abbasi

September 27, 2022

Compiled on September 27, 2022 at 5:45am

Contents

1	Introduction	3
2	detailed summary tables of results	19
3	Listing of integrals	83
4	Appendix	1095

Chapter 1

Introduction

Local contents

1.1	Listing of CAS systems tested	4
1.2	Results	5
1.3	Time and leaf size Performance	9
1.4	list of integrals that has no closed form antiderivative	11
1.5	List of integrals solved by CAS but has no known antiderivative	12
1.6	list of integrals solved by CAS but failed verification	13
1.7	Timing	13
1.8	Verification	14
1.9	Important notes about some of the results	14
1.10	Design of the test system	17

This report gives the result of running the computer algebra independent integration test. The download section in the appendix contains links to download the problems in plain text format used for all CAS systems.

The number of integrals in this report is [247]. This is test number [172].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 13.1 (June 29, 2022) on windows 10.
2. Rubi 4.16.1 (Dec 19, 2018) on Mathematica 13.0.1 on windows 10.
3. Maple 2022.1 (June 1, 2022) on windows 10.
4. Maxima 5.46 (April 13, 2022) using Lisp SBCL 2.1.11.debian on Linux via sagemath 9.6.
5. Fricas 1.3.8 (June 21, 2022) based on sbcl 2.1.11.debian on Linux via sagemath 9.6.
6. Giac/Xcas 1.9.0-13 (July 3, 2022) on Linux via sagemath 9.6.
7. Sympy 1.10.1 (March 20, 2022) Using Python 3.10.4 on Linux.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was called directly from Python.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or `Hypergeometric2F1` functions. `RootSum` and `RootOf` are not allowed.

If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (247)	0.00 (0)
Mathematica	100.00 (247)	0.00 (0)
Fricas	84.62 (209)	15.38 (38)
Maple	83.81 (207)	16.19 (40)
Giac	75.71 (187)	24.29 (60)
Mupad	70.85 (175)	29.15 (72)
Maxima	61.13 (151)	38.87 (96)
Sympy	27.53 (68)	72.47 (179)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	100.00	0.00	0.00	0.00
Mathematica	81.38	6.48	12.15	0.00
Giac	56.68	16.60	2.43	24.29
Maple	56.28	15.38	12.15	16.19
Maxima	42.51	16.19	2.43	38.87
Fricas	15.38	66.80	2.43	15.38
Sympy	13.77	13.77	0.00	72.47
Mupad	N/A	70.04	0.00	29.15

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the CAS systems for each grade level.



The following table shows the distribution of the different types of failure for each CAS. There are 3 types of reasons why it can fail. The first is when CAS returns back the input within the time limit, which means it could not solve it. This is the typical normal failure **F**.

The second is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned **F(-1)**.

The third is due to an exception generated. Assigned **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and

Giac) or it could be an indication of an internal error in CAS. This type of error requires more investigations to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00 %	0.00 %	0.00 %
Mathematica	0	0.00 %	0.00 %	0.00 %
Maple	40	100.00 %	0.00 %	0.00 %
Fricas	38	100.00 %	0.00 %	0.00 %
Giac	60	81.67 %	15.00 %	3.33 %
Maxima	96	84.38 %	0.00 %	15.62 %
Sympy	179	96.65 %	2.23 %	1.12 %
Mupad	72	100.00 %	0.00 %	0.00 %

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.



Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)	Mean size	Normalized mean	Median size	Normalized median
Rubi	0.08	78.32	0.99	60.00	1.00
Mathematica	0.84	81.64	1.12	56.00	1.00
Maple	1.05	89.85	1.34	57.00	1.10
Maxima	0.37	88.25	1.57	65.00	1.14
Fricas	0.42	727.07	8.16	230.00	4.33
Sympy	2.55	417.85	4.96	75.00	1.76
Giac	0.42	91.60	1.41	63.00	1.24
Mupad	1.15	85.99	1.35	48.00	0.92

Table 1.5: Time and leaf size performance for each CAS

The following are bar charts for the normalized leafsize and time used from the above table.



1.4 list of integrals that has no closed form antiderivative

{243, 247}

1.5 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Mupad {}

1.6 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not mean necessarily that the anti-derivative is wrong, as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it easier to do further investigation to determine why it was not possible to verify the result produced.

Rubi {}

Mathematica {163, 164, 167, 168, 169, 170, 171, 192, 193, 238}

Maple Verification phase not implemented yet.

Maxima Verification phase not implemented yet.

Fricas Verification phase not implemented yet.

Sympy Verification phase not implemented yet.

Giac Verification phase not implemented yet.

Mupad Verification phase not implemented yet.

1.7 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.8 Verification

A verification phase was applied on the result of integration for `Rubi` and `Mathematica`.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.9 Important notes about some of the results

1.9.1 Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima `integrate` was run using SageMath with the following settings set by default

```
'besselexpand : true'
'display2d : false'
'domain : complex'
'keepfloat : true'
'load(to_poly_solve)'
'load(simplify_sum)'
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib
maxima_lib.set('extra_definite_integration_methods', '[]')
```

```
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

1.9.2 Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

1.9.3 Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount =1
```

1.9.4 Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand,the_variable)
```

Which gives $\sin(x)^2/2$

1.10 Design of the test system

The following diagram gives a high level view of the current test build system.



One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "F"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified.

The following fields are present only in Rubi Table file

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,..}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Chapter 2

detailed summary tables of results

Local contents

2.1	List of integrals sorted by grade for each CAS	20
2.2	Detailed conclusion table per each integral for all CAS systems	25
2.3	Detailed conclusion table specific for Rubi results	75

2.1 List of integrals sorted by grade for each CAS

Local contents

2.1.1	Rubi	21
2.1.2	Mathematica	21
2.1.3	Maple	21
2.1.4	Maxima	22
2.1.5	FriCAS	22
2.1.6	Sympy	23
2.1.7	Giac	23
2.1.8	Mupad	23

2.1.1 Rubi

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247 }

B grade: { }

C grade: { }

F grade: { }

2.1.2 Mathematica

A grade: { 1, 2, 3, 4, 5, 6, 7, 9, 11, 13, 14, 15, 16, 17, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 37, 38, 39, 40, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 148, 150, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 164, 165, 166, 176, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 193, 194, 195, 196, 197, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 213, 215, 216, 226, 228, 229, 232, 233, 234, 235, 236, 237, 239, 240, 241, 242, 243, 244, 245, 246, 247 }

B grade: { 41, 42, 146, 163, 169, 170, 171, 172, 173, 174, 175, 177, 178, 192, 230, 231 }

C grade: { 8, 10, 12, 18, 19, 20, 21, 36, 145, 147, 149, 151, 167, 168, 198, 199, 211, 212, 214, 217, 218, 219, 220, 221, 222, 223, 224, 225, 227, 238 }

F grade: { }

2.1.3 Maple

A grade: { 1, 2, 3, 4, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 73, 80, 81, 82, 83, 84, 85, 86, 87, 88, 90, 95, 96, 105, 106, 107, 108, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 146, 153, 176, 186, 187, 188, 194, 195, 196, 197, 198, 199, 200, 201, 202, 208, 209, 210, 211, 212, 213, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247 }

B grade: { 5, 6, 7, 8, 69, 70, 71, 72, 74, 75, 76, 77, 78, 79, 89, 91, 92, 93, 94, 97, 98, 99, 100, 101, 102, 103, 104, 109, 121, 122, 123, 124, 144, 145, 150, 157, 183, 237 }

C grade: { 147, 148, 149, 151, 152, 154, 155, 156, 158, 159, 205, 206, 207, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 234, 235, 236 }

F grade: { 22, 23, 40, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 177, 178, 179, 180, 181, 182, 184, 185, 189, 190, 191, 192, 193, 203, 204, 228, 229, 230, 231, 232, 233 }

2.1.4 Maxima

A grade: { 5, 6, 7, 8, 24, 25, 37, 38, 39, 44, 45, 46, 47, 48, 49, 50, 56, 59, 60, 61, 62, 65, 66, 69, 70, 71, 72, 73, 74, 80, 82, 90, 91, 92, 93, 94, 95, 96, 105, 106, 107, 108, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 176, 183, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 224, 225, 234, 235, 236, 237, 238, 239, 243, 247 }

B grade: { 1, 2, 3, 4, 9, 10, 11, 12, 26, 41, 42, 43, 51, 52, 53, 54, 55, 57, 58, 63, 64, 75, 76, 77, 78, 79, 85, 87, 89, 97, 98, 99, 100, 101, 102, 103, 104, 186, 187, 188 }

C grade: { 27, 28, 29, 30, 31, 32 }

F grade: { 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 33, 34, 35, 36, 40, 67, 68, 81, 83, 84, 86, 88, 109, 110, 111, 112, 113, 114, 125, 126, 127, 128, 129, 130, 131, 132, 144, 145, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 177, 178, 179, 180, 181, 182, 184, 185, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 222, 223, 226, 227, 228, 229, 230, 231, 232, 233, 240, 241, 242, 244, 245, 246 }

2.1.5 FriCAS

A grade: { 61, 65, 66, 71, 72, 84, 90, 93, 94, 95, 106, 112, 136, 137, 138, 139, 146, 147, 148, 149, 150, 152, 153, 154, 155, 159, 209, 215, 218, 219, 220, 224, 236, 237, 241, 243, 245, 247 }

B grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 24, 25, 26, 33, 34, 35, 36, 37, 38, 39, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 62, 63, 64, 67, 68, 69, 70, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 85, 86, 87, 88, 89, 91, 92, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 107, 108, 109, 110, 111, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 140, 141, 142, 143, 144, 145, 151, 156, 157, 158, 176, 183, 186, 187, 188, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 210, 211, 212, 213, 214, 216, 217, 221, 222, 223, 225, 226, 227, 234, 235, 238, 239, 240, 242, 244, 246 }

C grade: { 27, 28, 29, 30, 31, 32 }

F grade: { 22, 23, 40, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 177, 178, 179, 180, 181, 182, 184, 185, 189, 190, 191, 192, 193, 228, 229, 230, 231, 232, 233 }

2.1.6 Sympy

A grade: { 1, 2, 3, 4, 5, 21, 41, 42, 43, 44, 57, 58, 59, 60, 65, 66, 95, 125, 126, 127, 128, 129, 130, 131, 132, 150, 157, 176, 187, 188, 196, 197, 243, 247 }

B grade: { 6, 7, 8, 9, 10, 11, 12, 45, 46, 47, 48, 49, 61, 62, 63, 70, 72, 92, 94, 106, 115, 116, 117, 118, 119, 120, 133, 134, 135, 136, 137, 138, 183, 186 }

C grade: { }

F grade: { 13, 14, 15, 16, 17, 18, 19, 20, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 50, 51, 52, 53, 54, 55, 56, 64, 67, 68, 69, 71, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 93, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 107, 108, 109, 110, 111, 112, 113, 114, 121, 122, 123, 124, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 151, 152, 153, 154, 155, 156, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 177, 178, 179, 180, 181, 182, 184, 185, 189, 190, 191, 192, 193, 194, 195, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 244, 245, 246 }

2.1.7 Giac

A grade: { 1, 2, 3, 4, 5, 8, 9, 10, 11, 12, 16, 17, 24, 26, 33, 37, 38, 39, 41, 42, 43, 44, 45, 46, 47, 48, 49, 52, 53, 54, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 69, 70, 71, 72, 73, 74, 76, 77, 78, 79, 80, 81, 82, 83, 84, 86, 88, 90, 91, 92, 93, 94, 95, 98, 99, 100, 101, 106, 107, 108, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 131, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 183, 187, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 234, 235, 236, 237, 238, 239, 243, 247 }

B grade: { 6, 7, 13, 14, 15, 18, 19, 20, 21, 25, 34, 35, 36, 50, 51, 55, 56, 75, 85, 87, 89, 96, 97, 102, 103, 104, 105, 109, 125, 126, 127, 128, 129, 130, 132, 143, 176, 186, 188, 226, 227 }

C grade: { 27, 28, 29, 30, 31, 32 }

F grade: { 22, 23, 40, 67, 68, 144, 145, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 177, 178, 179, 180, 181, 182, 184, 185, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 228, 229, 230, 231, 232, 233, 240, 241, 242, 244, 245, 246 }

2.1.8 Mupad

A grade: { 243, 247 }

B grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 26, 30, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 176, 183, 186, 187, 188, 194, 195, 196, 197, 198, 199, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227 }

C grade: { }

F grade: { 22, 23, 24, 25, 27, 28, 29, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 144, 145, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 177, 178, 179, 180, 181, 182, 184, 185, 189, 190, 191, 192, 193, 200, 201, 202, 203, 204, 205, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 244, 245, 246 }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by table below. The elapsed time is in seconds. For failed result it is given as F(-1) if the failure was due to timeout. It is given as F(-2) if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given just an F.

In this table, the column N.S. in the table below, which stands for **normalized size** is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To help make the table fit, **Mathematica** was abbrev-

	Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
viated to MMA.	grade	A	A	A	A	B	B	A	A	B
	verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
	size	43	43	53	56	115	254	39	74	34
	N.S.	1	1.00	1.23	1.30	2.67	5.91	0.91	1.72	0.79
	time (sec)	N/A	0.017	0.015	0.265	0.280	0.407	0.146	0.404	0.109

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	B	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	42	42	37	48	102	968	42	67	37
N.S.	1	1.00	0.88	1.14	2.43	23.05	1.00	1.60	0.88
time (sec)	N/A	0.023	0.073	0.260	0.474	0.375	0.116	0.424	0.117

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	B	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	28	28	38	46	71	119	27	52	24
N.S.	1	1.00	1.36	1.64	2.54	4.25	0.96	1.86	0.86
time (sec)	N/A	0.013	0.009	0.255	0.260	0.522	0.091	0.406	1.010

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	B	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	27	38	61	339	31	48	27
N.S.	1	1.00	1.00	1.41	2.26	12.56	1.15	1.78	1.00
time (sec)	N/A	0.013	0.012	0.259	0.465	0.382	0.077	0.426	1.013

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	13	13	23	36	25	33	15	24	13
N.S.	1	1.00	1.77	2.77	1.92	2.54	1.15	1.85	1.00
time (sec)	N/A	0.006	0.007	0.250	0.257	0.496	0.061	0.413	0.064

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	B	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	11	11	11	28	11	37	17	24	16
N.S.	1	1.00	1.00	2.55	1.00	3.36	1.55	2.18	1.45
time (sec)	N/A	0.005	0.005	0.247	0.263	0.658	0.060	0.401	1.019

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	B	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	11	11	19	28	11	37	27	25	21
N.S.	1	1.00	1.73	2.55	1.00	3.36	2.45	2.27	1.91
time (sec)	N/A	0.005	0.009	0.255	0.263	0.487	0.218	0.402	0.043

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	A	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	13	13	27	36	25	33	87	24	13
N.S.	1	1.00	2.08	2.77	1.92	2.54	6.69	1.85	1.00
time (sec)	N/A	0.007	0.008	0.249	0.267	0.531	0.628	0.403	1.019

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	34	38	79	346	112	49	68
N.S.	1	1.00	1.26	1.41	2.93	12.81	4.15	1.81	2.52
time (sec)	N/A	0.013	0.055	0.254	0.256	0.385	0.868	0.401	0.050

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	B	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	28	28	31	46	71	108	100	52	24
N.S.	1	1.00	1.11	1.64	2.54	3.86	3.57	1.86	0.86
time (sec)	N/A	0.013	0.009	0.264	0.260	0.362	1.380	0.436	0.058

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	42	42	44	48	122	978	126	70	159
N.S.	1	1.00	1.05	1.14	2.90	23.29	3.00	1.67	3.79
time (sec)	N/A	0.026	0.115	0.263	0.262	0.420	2.407	0.409	0.988

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	B	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	31	56	115	239	114	74	34
N.S.	1	1.00	0.72	1.30	2.67	5.56	2.65	1.72	0.79
time (sec)	N/A	0.018	0.008	0.262	0.266	0.376	3.852	0.438	0.066

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	97	97	83	74	0	1556	0	293	83
N.S.	1	1.00	0.86	0.76	0.00	16.04	0.00	3.02	0.86
time (sec)	N/A	0.058	0.173	1.948	0.000	0.387	0.000	0.529	1.515

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	78	78	68	60	0	980	0	173	62
N.S.	1	1.00	0.87	0.77	0.00	12.56	0.00	2.22	0.79
time (sec)	N/A	0.037	0.154	1.866	0.000	0.373	0.000	0.512	1.254

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	75	75	61	58	0	638	0	131	61
N.S.	1	1.00	0.81	0.77	0.00	8.51	0.00	1.75	0.81
time (sec)	N/A	0.036	0.064	1.745	0.000	0.506	0.000	0.448	1.164

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	51	48	0	593	0	88	41
N.S.	1	1.00	0.88	0.83	0.00	10.22	0.00	1.52	0.71
time (sec)	N/A	0.024	0.029	2.091	0.000	0.405	0.000	0.437	1.083

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	57	57	49	48	0	599	0	89	38
N.S.	1	1.00	0.86	0.84	0.00	10.51	0.00	1.56	0.67
time (sec)	N/A	0.022	0.025	2.089	0.000	0.373	0.000	0.475	1.123

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	78	78	36	62	0	924	0	133	64
N.S.	1	1.00	0.46	0.79	0.00	11.85	0.00	1.71	0.82
time (sec)	N/A	0.039	0.060	1.915	0.000	0.443	0.000	0.464	1.201

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	79	79	38	63	0	1436	0	173	63
N.S.	1	1.00	0.48	0.80	0.00	18.18	0.00	2.19	0.80
time (sec)	N/A	0.037	0.051	2.008	0.000	0.425	0.000	0.571	1.356

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	100	38	77	0	2144	0	293	80
N.S.	1	1.00	0.38	0.77	0.00	21.44	0.00	2.93	0.80
time (sec)	N/A	0.053	0.077	2.069	0.000	0.524	0.000	0.631	1.506

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	B	A	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	26	102	0	179	63	110	71
N.S.	1	1.00	0.38	1.48	0.00	2.59	0.91	1.59	1.03
time (sec)	N/A	0.043	0.020	0.497	0.000	0.369	1.951	0.419	1.318

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	45	0	0	0	0	0	-1
N.S.	1	1.00	1.05	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.017	0.048	1.850	0.000	0.000	0.000	0.000	0.000

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	48	48	51	0	0	0	0	0	-1
N.S.	1	1.00	1.06	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.021	0.034	1.778	0.000	0.000	0.000	0.000	0.000

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	28	30	42	467	0	52	-1
N.S.	1	1.00	0.80	0.86	1.20	13.34	0.00	1.49	-0.03
time (sec)	N/A	0.015	0.020	0.746	0.472	0.395	0.000	0.426	0.000

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	16	26	19	72	0	31	-1
N.S.	1	1.00	1.00	1.62	1.19	4.50	0.00	1.94	-0.06
time (sec)	N/A	0.011	0.006	0.762	0.485	0.407	0.000	0.414	0.000

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	16	29	31	76	0	1	14
N.S.	1	1.00	1.00	1.81	1.94	4.75	0.00	0.06	0.88
time (sec)	N/A	0.010	0.007	0.788	0.486	0.404	0.000	0.417	1.200

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	C	C	F	C	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	88	56	67	113	180	0	142	-1
N.S.	1	1.00	0.64	0.76	1.28	2.05	0.00	1.61	-0.01
time (sec)	N/A	0.039	0.211	1.461	0.499	0.383	0.000	0.446	0.000

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	C	C	F	C	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	60	60	46	53	66	100	0	92	-1
N.S.	1	1.00	0.77	0.88	1.10	1.67	0.00	1.53	-0.02
time (sec)	N/A	0.024	0.072	1.334	0.508	0.386	0.000	0.419	0.000

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	C	C	F	C	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	31	45	28	23	0	54	-1
N.S.	1	1.00	1.00	1.45	0.90	0.74	0.00	1.74	-0.03
time (sec)	N/A	0.013	0.025	1.449	0.497	0.365	0.000	0.423	0.000

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	C	C	F	C	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	39	56	45	23	0	63	24
N.S.	1	1.00	1.26	1.81	1.45	0.74	0.00	2.03	0.77
time (sec)	N/A	0.013	0.057	1.635	0.478	0.420	0.000	0.414	1.224

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	C	C	F	C	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	60	60	51	81	85	100	0	100	-1
N.S.	1	1.00	0.85	1.35	1.42	1.67	0.00	1.67	-0.02
time (sec)	N/A	0.024	0.105	1.667	0.492	0.424	0.000	0.422	0.000

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	C	C	F	C	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	88	63	91	132	180	0	126	-1
N.S.	1	1.00	0.72	1.03	1.50	2.05	0.00	1.43	-0.01
time (sec)	N/A	0.035	0.175	1.737	0.475	0.416	0.000	0.425	0.000

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	B	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	57	57	38	43	0	106	0	55	-1
N.S.	1	1.00	0.67	0.75	0.00	1.86	0.00	0.96	-0.02
time (sec)	N/A	0.028	0.028	0.868	0.000	0.435	0.000	0.414	0.000

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	B	F	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	86	86	55	76	0	1269	0	342	-1
N.S.	1	1.00	0.64	0.88	0.00	14.76	0.00	3.98	-0.01
time (sec)	N/A	0.029	0.046	1.121	0.000	0.411	0.000	0.472	0.000

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	B	F	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	40	59	0	376	0	115	-1
N.S.	1	1.00	0.63	0.94	0.00	5.97	0.00	1.83	-0.02
time (sec)	N/A	0.021	0.024	1.150	0.000	0.397	0.000	0.431	0.000

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	B	F	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	64	64	26	65	0	516	0	123	-1
N.S.	1	1.00	0.41	1.02	0.00	8.06	0.00	1.92	-0.02
time (sec)	N/A	0.022	0.014	1.140	0.000	0.454	0.000	0.434	0.000

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	56	46	82	2114	0	45	-1
N.S.	1	1.00	0.81	0.67	1.19	30.64	0.00	0.65	-0.01
time (sec)	N/A	0.019	0.111	0.726	0.489	0.442	0.000	0.421	0.000

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	19	32	19	213	0	16	-1
N.S.	1	1.00	0.61	1.03	0.61	6.87	0.00	0.52	-0.03
time (sec)	N/A	0.010	0.013	0.724	0.485	0.368	0.000	0.423	0.000

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	19	32	23	238	0	19	-1
N.S.	1	1.00	0.61	1.03	0.74	7.68	0.00	0.61	-0.03
time (sec)	N/A	0.011	0.025	0.739	0.484	0.363	0.000	0.408	0.000

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	57	57	55	0	0	0	0	0	-1
N.S.	1	1.00	0.96	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.030	0.038	180.000	0.000	0.000	0.000	0.000	0.000

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	A	B	B	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	100	202	58	302	907	95	85	65
N.S.	1	1.00	2.02	0.58	3.02	9.07	0.95	0.85	0.65
time (sec)	N/A	0.056	0.802	0.264	0.484	0.353	0.135	0.401	0.160

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	A	B	B	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	77	77	178	48	196	562	76	71	53
N.S.	1	1.00	2.31	0.62	2.55	7.30	0.99	0.92	0.69
time (sec)	N/A	0.041	0.574	0.263	0.478	0.392	0.110	0.420	0.108

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	B	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	56	56	103	38	116	299	61	57	43
N.S.	1	1.00	1.84	0.68	2.07	5.34	1.09	1.02	0.77
time (sec)	N/A	0.026	0.491	0.256	0.470	0.346	0.101	0.407	0.094

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	36	58	28	50	117	44	39	33
N.S.	1	1.00	1.61	0.78	1.39	3.25	1.22	1.08	0.92
time (sec)	N/A	0.017	0.241	0.260	0.264	0.390	0.075	0.428	1.067

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	28	28	39	43	31	50	73	30	25
N.S.	1	1.00	1.39	1.54	1.11	1.79	2.61	1.07	0.89
time (sec)	N/A	0.010	0.084	0.524	0.260	0.392	0.307	0.411	1.071

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	51	51	60	55	43	101	223	42	41
N.S.	1	1.00	1.18	1.08	0.84	1.98	4.37	0.82	0.80
time (sec)	N/A	0.023	0.132	0.504	0.267	0.356	0.543	0.423	1.088

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	73	73	83	67	56	160	430	53	58
N.S.	1	1.00	1.14	0.92	0.77	2.19	5.89	0.73	0.79
time (sec)	N/A	0.034	0.184	0.660	0.272	0.437	0.766	0.420	1.099

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	96	96	88	79	67	220	694	64	75
N.S.	1	1.00	0.92	0.82	0.70	2.29	7.23	0.67	0.78
time (sec)	N/A	0.048	0.185	0.640	0.290	0.374	1.083	0.421	1.138

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	121	121	109	91	78	287	1018	75	92
N.S.	1	1.00	0.90	0.75	0.64	2.37	8.41	0.62	0.76
time (sec)	N/A	0.065	0.211	0.641	0.291	0.401	1.557	0.421	1.160

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	57	57	65	43	83	434	0	140	44
N.S.	1	1.00	1.14	0.75	1.46	7.61	0.00	2.46	0.77
time (sec)	N/A	0.033	0.147	0.582	0.479	0.382	0.000	0.418	0.192

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	45	45	39	35	70	252	0	96	54
N.S.	1	1.00	0.87	0.78	1.56	5.60	0.00	2.13	1.20
time (sec)	N/A	0.024	0.122	0.546	0.478	0.488	0.000	0.418	0.113

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	33	33	33	27	57	129	0	52	26
N.S.	1	1.00	1.00	0.82	1.73	3.91	0.00	1.58	0.79
time (sec)	N/A	0.016	0.050	0.556	0.469	0.354	0.000	0.394	1.119

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	21	21	21	17	43	50	0	27	16
N.S.	1	1.00	1.00	0.81	2.05	2.38	0.00	1.29	0.76
time (sec)	N/A	0.009	0.035	0.661	0.479	0.381	0.000	0.434	0.132

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	32	32	32	27	57	85	0	50	26
N.S.	1	1.00	1.00	0.84	1.78	2.66	0.00	1.56	0.81
time (sec)	N/A	0.016	0.056	0.634	0.465	0.579	0.000	0.415	0.128

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	49	53	35	69	166	0	95	32
N.S.	1	1.00	1.08	0.71	1.41	3.39	0.00	1.94	0.65
time (sec)	N/A	0.025	0.105	0.599	0.480	0.366	0.000	0.409	0.115

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	61	61	62	43	79	266	0	139	40
N.S.	1	1.00	1.02	0.70	1.30	4.36	0.00	2.28	0.66
time (sec)	N/A	0.032	0.168	0.588	0.474	0.466	0.000	0.421	0.121

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	B	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	142	142	114	182	310	2739	211	224	153
N.S.	1	1.00	0.80	1.28	2.18	19.29	1.49	1.58	1.08
time (sec)	N/A	0.153	0.460	0.273	0.481	0.647	0.164	0.426	1.164

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	B	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	101	101	91	134	201	1389	144	152	113
N.S.	1	1.00	0.90	1.33	1.99	13.75	1.43	1.50	1.12
time (sec)	N/A	0.088	0.242	0.269	0.470	0.365	0.135	0.429	1.086

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	67	93	118	646	100	97	77
N.S.	1	1.00	0.97	1.35	1.71	9.36	1.45	1.41	1.12
time (sec)	N/A	0.047	0.235	0.263	0.477	0.408	0.108	0.439	0.112

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	54	61	49	201	54	56	44
N.S.	1	1.00	1.42	1.61	1.29	5.29	1.42	1.47	1.16
time (sec)	N/A	0.017	0.075	0.257	0.273	0.379	0.075	0.409	1.044

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	50	50	64	71	56	62	224	62	60
N.S.	1	1.00	1.28	1.42	1.12	1.24	4.48	1.24	1.20
time (sec)	N/A	0.038	0.062	0.852	0.267	0.411	0.723	0.431	1.143

Problem 62	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	85	85	94	93	127	422	3597	132	127
N.S.	1	1.00	1.11	1.09	1.49	4.96	42.32	1.55	1.49
time (sec)	N/A	0.070	0.829	0.759	0.272	0.606	14.102	0.426	1.429

Problem 63	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	129	129	122	130	325	1427	16657	205	304
N.S.	1	1.00	0.95	1.01	2.52	11.06	129.12	1.59	2.36
time (sec)	N/A	0.128	1.689	0.635	0.299	0.387	82.404	0.420	2.112

Problem 64	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	B	F(-1)	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	169	169	160	163	525	3693	0	305	452
N.S.	1	1.00	0.95	0.96	3.11	21.85	0.00	1.80	2.67
time (sec)	N/A	0.189	2.314	0.796	0.328	0.452	0.000	0.437	2.989

Problem 65	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	55	42	28	49	42	30	34
N.S.	1	1.00	1.77	1.35	0.90	1.58	1.35	0.97	1.10
time (sec)	N/A	0.031	0.028	0.631	0.264	0.438	0.231	0.406	0.132

Problem 66	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	55	42	29	48	42	25	33
N.S.	1	1.00	1.77	1.35	0.94	1.55	1.35	0.81	1.06
time (sec)	N/A	0.031	0.027	0.638	0.279	0.355	0.233	0.410	0.120

Problem 67	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F(-2)	B	F	F(-2)	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	74	74	74	70	0	2203	0	0	151
N.S.	1	1.00	1.00	0.95	0.00	29.77	0.00	0.00	2.04
time (sec)	N/A	0.053	0.063	1.747	0.000	0.651	0.000	0.000	1.337

Problem 68	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F(-2)	B	F	F(-2)	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	74	74	74	70	0	2279	0	0	240
N.S.	1	1.00	1.00	0.95	0.00	30.80	0.00	0.00	3.24
time (sec)	N/A	0.044	0.047	1.565	0.000	0.568	0.000	0.000	1.393

Problem 69	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	60	60	42	98	36	92	0	42	34
N.S.	1	1.00	0.70	1.63	0.60	1.53	0.00	0.70	0.57
time (sec)	N/A	0.047	0.064	0.521	0.250	0.413	0.000	0.412	1.312

Problem 70	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	34	72	27	56	134	25	23
N.S.	1	1.00	1.36	2.88	1.08	2.24	5.36	1.00	0.92
time (sec)	N/A	0.119	0.049	0.486	0.261	0.439	0.370	0.394	1.179

Problem 71	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	24	68	22	51	0	30	22
N.S.	1	1.00	0.63	1.79	0.58	1.34	0.00	0.79	0.58
time (sec)	N/A	0.042	0.029	0.372	0.279	0.334	0.000	0.407	1.145

Problem 72	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	17	19	42	11	23	48	11	11
N.S.	1	1.00	1.12	2.47	0.65	1.35	2.82	0.65	0.65
time (sec)	N/A	0.076	0.012	0.325	0.264	0.411	0.194	0.398	1.119

Problem 73	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	14	17	21	38	0	18	21
N.S.	1	1.00	1.17	1.42	1.75	3.17	0.00	1.50	1.75
time (sec)	N/A	0.078	0.016	0.562	0.261	0.391	0.000	0.417	0.059

Problem 74	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	11	32	29	77	0	29	23
N.S.	1	1.00	0.73	2.13	1.93	5.13	0.00	1.93	1.53
time (sec)	N/A	0.027	0.024	0.500	0.262	0.375	0.000	0.407	1.079

Problem 75	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	20	39	48	209	0	34	48
N.S.	1	1.00	1.11	2.17	2.67	11.61	0.00	1.89	2.67
time (sec)	N/A	0.112	0.058	0.612	0.263	0.356	0.000	0.404	1.086

Problem 76	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	17	20	48	75	84	0	18	18
N.S.	1	1.00	1.18	2.82	4.41	4.94	0.00	1.06	1.06
time (sec)	N/A	0.031	0.035	0.611	0.257	0.383	0.000	0.398	1.063

Problem 77	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	34	34	49	55	74	640	0	49	117
N.S.	1	1.00	1.44	1.62	2.18	18.82	0.00	1.44	3.44
time (sec)	N/A	0.135	0.090	0.629	0.268	0.436	0.000	0.416	1.093

Problem 78	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	33	33	27	80	149	185	0	24	24
N.S.	1	1.00	0.82	2.42	4.52	5.61	0.00	0.73	0.73
time (sec)	N/A	0.037	0.037	0.616	0.261	0.329	0.000	0.406	0.096

Problem 79	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	44	44	68	103	98	1260	0	61	207
N.S.	1	1.00	1.55	2.34	2.23	28.64	0.00	1.39	4.70
time (sec)	N/A	0.146	0.194	0.631	0.274	0.359	0.000	0.400	1.166

Problem 80	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	147	147	144	246	163	1226	0	214	135
N.S.	1	1.00	0.98	1.67	1.11	8.34	0.00	1.46	0.92
time (sec)	N/A	0.261	0.425	0.703	0.282	0.360	0.000	0.419	1.671

Problem 81	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F(-2)	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	137	137	180	166	0	1861	0	163	261
N.S.	1	1.00	1.31	1.21	0.00	13.58	0.00	1.19	1.91
time (sec)	N/A	0.204	0.838	0.686	0.000	0.362	0.000	0.422	1.456

Problem 82	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	84	84	73	145	83	334	0	101	81
N.S.	1	1.00	0.87	1.73	0.99	3.98	0.00	1.20	0.96
time (sec)	N/A	0.114	0.176	0.677	0.262	0.399	0.000	0.411	1.434

Problem 83	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F(-2)	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	79	92	0	427	0	60	157
N.S.	1	1.00	1.10	1.28	0.00	5.93	0.00	0.83	2.18
time (sec)	N/A	0.085	0.163	0.656	0.000	0.349	0.000	0.394	1.363

Problem 84	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F(-2)	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	52	52	59	53	0	237	0	60	177
N.S.	1	1.00	1.13	1.02	0.00	4.56	0.00	1.15	3.40
time (sec)	N/A	0.100	0.072	0.738	0.000	0.392	0.000	0.406	1.397

Problem 85	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	29	29	28	56	65	122	0	78	323
N.S.	1	1.00	0.97	1.93	2.24	4.21	0.00	2.69	11.14
time (sec)	N/A	0.040	0.070	0.738	0.252	0.361	0.000	0.421	1.509

Problem 86	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F(-2)	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	82	82	123	110	0	1165	0	125	506
N.S.	1	1.00	1.50	1.34	0.00	14.21	0.00	1.52	6.17
time (sec)	N/A	0.208	0.257	0.878	0.000	0.383	0.000	0.417	1.511

Problem 87	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	78	78	70	146	161	912	0	202	123
N.S.	1	1.00	0.90	1.87	2.06	11.69	0.00	2.59	1.58
time (sec)	N/A	0.074	0.211	0.749	0.289	0.363	0.000	0.418	1.330

Problem 88	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F(-2)	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	255	255	296	227	0	5347	0	273	753
N.S.	1	1.00	1.16	0.89	0.00	20.97	0.00	1.07	2.95
time (sec)	N/A	0.402	0.609	1.015	0.000	0.453	0.000	0.421	3.307

Problem 89	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	130	130	119	277	308	2972	0	412	237
N.S.	1	1.00	0.92	2.13	2.37	22.86	0.00	3.17	1.82
time (sec)	N/A	0.111	0.448	0.754	0.296	0.397	0.000	0.402	1.326

Problem 90	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	33	33	46	29	34	43	0	28	61
N.S.	1	1.00	1.39	0.88	1.03	1.30	0.00	0.85	1.85
time (sec)	N/A	0.071	0.051	0.933	0.526	0.380	0.000	0.402	0.458

Problem 91	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	60	60	42	116	36	95	0	42	34
N.S.	1	1.00	0.70	1.93	0.60	1.58	0.00	0.70	0.57
time (sec)	N/A	0.042	0.034	0.491	0.268	0.364	0.000	0.405	1.286

Problem 92	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	29	29	36	80	33	60	134	31	29
N.S.	1	1.00	1.24	2.76	1.14	2.07	4.62	1.07	1.00
time (sec)	N/A	0.029	0.038	0.465	0.262	0.419	0.347	0.400	1.187

Problem 93	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	30	76	22	52	0	30	22
N.S.	1	1.00	0.79	2.00	0.58	1.37	0.00	0.79	0.58
time (sec)	N/A	0.035	0.023	0.500	0.263	0.377	0.000	0.410	0.126

Problem 94	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	23	40	17	25	48	19	17
N.S.	1	1.00	1.21	2.11	0.89	1.32	2.53	1.00	0.89
time (sec)	N/A	0.022	0.022	0.486	0.260	0.415	0.180	0.400	1.051

Problem 95	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	10	10	7	11	6	9	8	6	6
N.S.	1	1.00	0.70	1.10	0.60	0.90	0.80	0.60	0.60
time (sec)	N/A	0.015	0.005	0.257	0.266	0.366	0.157	0.413	0.038

Problem 96	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	5	5	7	6	5	20	0	13	13
N.S.	1	1.00	1.40	1.20	1.00	4.00	0.00	2.60	2.60
time (sec)	N/A	0.022	0.005	0.509	0.267	0.341	0.000	0.408	1.047

Problem 97	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	6	6	12	21	22	48	0	18	18
N.S.	1	1.00	2.00	3.50	3.67	8.00	0.00	3.00	3.00
time (sec)	N/A	0.023	0.020	0.496	0.482	0.354	0.000	0.415	1.041

Problem 98	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	11	11	11	34	37	53	0	10	16
N.S.	1	1.00	1.00	3.09	3.36	4.82	0.00	0.91	1.45
time (sec)	N/A	0.023	0.019	0.546	0.261	0.372	0.000	0.408	1.042

Problem 99	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	24	24	24	41	49	288	0	31	61
N.S.	1	1.00	1.00	1.71	2.04	12.00	0.00	1.29	2.54
time (sec)	N/A	0.030	0.024	0.566	0.465	0.425	0.000	0.406	0.091

Problem 100	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	20	56	93	140	0	18	18
N.S.	1	1.00	0.80	2.24	3.72	5.60	0.00	0.72	0.72
time (sec)	N/A	0.027	0.023	0.552	0.262	0.327	0.000	0.397	1.038

Problem 101	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	34	34	34	67	73	670	0	45	137
N.S.	1	1.00	1.00	1.97	2.15	19.71	0.00	1.32	4.03
time (sec)	N/A	0.034	0.026	0.574	0.463	0.363	0.000	0.403	0.085

Problem 102	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	140	140	137	413	386	5275	0	593	301
N.S.	1	1.00	0.98	2.95	2.76	37.68	0.00	4.24	2.15
time (sec)	N/A	0.117	0.441	0.759	0.492	0.408	0.000	0.411	1.435

Problem 103	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	83	83	92	222	204	1827	0	316	169
N.S.	1	1.00	1.11	2.67	2.46	22.01	0.00	3.81	2.04
time (sec)	N/A	0.081	0.276	0.705	0.486	0.379	0.000	0.419	1.290

Problem 104	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	40	40	49	99	89	430	0	104	88
N.S.	1	1.00	1.22	2.48	2.22	10.75	0.00	2.60	2.20
time (sec)	N/A	0.048	0.132	0.711	0.469	0.428	0.000	0.413	1.266

Problem 105	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	11	11	20	12	11	42	0	45	50
N.S.	1	1.00	1.82	1.09	1.00	3.82	0.00	4.09	4.55
time (sec)	N/A	0.028	0.036	0.517	0.252	0.376	0.000	0.411	0.207

Problem 106	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	39	39	29	55	41	42	146	43	35
N.S.	1	1.00	0.74	1.41	1.05	1.08	3.74	1.10	0.90
time (sec)	N/A	0.032	0.041	0.238	0.272	0.397	0.264	0.406	1.114

Problem 107	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	91	91	75	153	86	331	0	111	84
N.S.	1	1.00	0.82	1.68	0.95	3.64	0.00	1.22	0.92
time (sec)	N/A	0.106	0.092	0.713	0.286	0.404	0.000	0.417	1.284

Problem 108	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	155	155	156	265	165	1281	0	227	143
N.S.	1	1.00	1.01	1.71	1.06	8.26	0.00	1.46	0.92
time (sec)	N/A	0.171	0.152	0.714	0.272	0.493	0.000	0.412	1.511

Problem 109	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F(-2)	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	157	187	166	323	0	6509	0	326	447
N.S.	1	1.19	1.06	2.06	0.00	41.46	0.00	2.08	2.85
time (sec)	N/A	0.196	0.356	1.071	0.000	0.512	0.000	0.415	7.293

Problem 110	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F(-2)	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	102	109	116	164	0	2043	0	152	265
N.S.	1	1.07	1.14	1.61	0.00	20.03	0.00	1.49	2.60
time (sec)	N/A	0.118	0.186	0.987	0.000	0.472	0.000	0.416	5.224

Problem 111	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F(-2)	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	56	56	65	77	0	309	0	63	119
N.S.	1	1.00	1.16	1.38	0.00	5.52	0.00	1.12	2.12
time (sec)	N/A	0.063	0.064	0.846	0.000	0.364	0.000	0.409	3.811

Problem 112	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F(-2)	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	37	46	39	0	148	0	35	35
N.S.	1	1.00	1.24	1.05	0.00	4.00	0.00	0.95	0.95
time (sec)	N/A	0.024	0.024	0.448	0.000	0.443	0.000	0.405	0.128

Problem 113	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F(-2)	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	73	73	80	93	0	435	0	61	157
N.S.	1	1.00	1.10	1.27	0.00	5.96	0.00	0.84	2.15
time (sec)	N/A	0.066	0.161	0.699	0.000	0.567	0.000	0.412	1.358

Problem 114	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F(-2)	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	132	132	258	176	0	1871	0	162	221
N.S.	1	1.00	1.95	1.33	0.00	14.17	0.00	1.23	1.67
time (sec)	N/A	0.124	0.320	0.706	0.000	0.405	0.000	0.410	2.254

Problem 115	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	40	40	55	571	104	47	35
N.S.	1	1.00	0.93	0.93	1.28	13.28	2.42	1.09	0.81
time (sec)	N/A	0.069	0.072	0.376	0.458	0.454	0.222	0.430	0.099

Problem 116	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	37	33	32	43	354	85	39	29
N.S.	1	1.00	0.89	0.86	1.16	9.57	2.30	1.05	0.78
time (sec)	N/A	0.049	0.041	0.363	0.468	0.560	0.199	0.407	1.045

Problem 117	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	27	28	29	186	75	35	21
N.S.	1	1.00	0.87	0.90	0.94	6.00	2.42	1.13	0.68
time (sec)	N/A	0.038	0.037	0.293	0.474	0.415	0.189	0.420	0.075

Problem 118	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	23	24	17	73	61	17	21
N.S.	1	1.00	1.21	1.26	0.89	3.84	3.21	0.89	1.11
time (sec)	N/A	0.027	0.023	0.293	0.474	0.332	0.175	0.425	1.065

Problem 119	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	18	24	10	26	27	10	10
N.S.	1	1.00	1.12	1.50	0.62	1.62	1.69	0.62	0.62
time (sec)	N/A	0.014	0.021	0.240	0.256	0.375	0.164	0.408	0.066

Problem 120	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	18	24	10	26	27	10	10
N.S.	1	1.00	1.12	1.50	0.62	1.62	1.69	0.62	0.62
time (sec)	N/A	0.007	0.013	0.230	0.250	0.345	0.170	0.401	1.033

Problem 121	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	23	43	24	73	0	18	17
N.S.	1	1.00	1.21	2.26	1.26	3.84	0.00	0.95	0.89
time (sec)	N/A	0.031	0.023	0.423	0.266	0.387	0.000	0.417	0.065

Problem 122	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	29	29	27	59	38	196	0	36	29
N.S.	1	1.00	0.93	2.03	1.31	6.76	0.00	1.24	1.00
time (sec)	N/A	0.052	0.035	0.426	0.257	0.598	0.000	0.402	1.033

Problem 123	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	37	33	75	54	357	0	40	35
N.S.	1	1.00	0.89	2.03	1.46	9.65	0.00	1.08	0.95
time (sec)	N/A	0.069	0.046	0.493	0.261	0.412	0.000	0.430	0.079

Problem 124	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	42	91	64	582	0	48	69
N.S.	1	1.00	0.98	2.12	1.49	13.53	0.00	1.12	1.60
time (sec)	N/A	0.081	0.099	0.494	0.275	0.425	0.000	0.419	1.097

Problem 125	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	B	A	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	45	45	39	35	0	252	78	96	34
N.S.	1	1.00	0.87	0.78	0.00	5.60	1.73	2.13	0.76
time (sec)	N/A	0.037	0.139	0.503	0.000	0.605	6.719	0.440	0.146

Problem 126	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	B	A	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	32	32	32	26	0	129	66	53	25
N.S.	1	1.00	1.00	0.81	0.00	4.03	2.06	1.66	0.78
time (sec)	N/A	0.028	0.036	0.572	0.000	0.457	1.286	0.419	1.061

Problem 127	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	B	A	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	30	30	30	25	0	85	63	52	24
N.S.	1	1.00	1.00	0.83	0.00	2.83	2.10	1.73	0.80
time (sec)	N/A	0.027	0.040	0.670	0.000	0.338	1.562	0.414	0.129

Problem 128	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	B	A	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	49	51	35	0	168	78	73	32
N.S.	1	1.00	1.04	0.71	0.00	3.43	1.59	1.49	0.65
time (sec)	N/A	0.036	0.086	0.591	0.000	0.411	6.784	0.428	1.088

Problem 129	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	B	A	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	45	45	53	35	0	429	78	140	34
N.S.	1	1.00	1.18	0.78	0.00	9.53	1.73	3.11	0.76
time (sec)	N/A	0.044	0.145	0.627	0.000	0.357	10.124	0.428	1.111

Problem 130	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	B	A	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	34	34	34	26	0	237	68	96	25
N.S.	1	1.00	1.00	0.76	0.00	6.97	2.00	2.82	0.74
time (sec)	N/A	0.037	0.052	0.654	0.000	0.352	1.853	0.434	0.109

Problem 131	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	B	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	42	42	37	35	0	182	75	54	36
N.S.	1	1.00	0.88	0.83	0.00	4.33	1.79	1.29	0.86
time (sec)	N/A	0.040	0.066	0.737	0.000	0.331	2.139	0.419	0.132

Problem 132	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	B	A	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	49	53	35	0	168	78	95	31
N.S.	1	1.00	1.08	0.71	0.00	3.43	1.59	1.94	0.63
time (sec)	N/A	0.057	0.073	0.681	0.000	0.362	7.672	0.423	1.075

Problem 133	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	94	94	105	95	150	1296	546	142	85
N.S.	1	1.00	1.12	1.01	1.60	13.79	5.81	1.51	0.90
time (sec)	N/A	0.264	0.410	0.315	0.490	0.413	0.566	0.407	0.229

Problem 134	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	76	88	76	100	644	442	98	68
N.S.	1	1.00	1.16	1.00	1.32	8.47	5.82	1.29	0.89
time (sec)	N/A	0.157	0.217	0.323	0.482	0.370	0.481	0.401	0.166

Problem 135	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	64	64	65	67	71	264	330	75	59
N.S.	1	1.00	1.02	1.05	1.11	4.12	5.16	1.17	0.92
time (sec)	N/A	0.097	0.103	0.359	0.490	0.433	0.378	0.413	1.089

Problem 136	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	49	60	56	76	243	58	46
N.S.	1	1.00	0.78	0.95	0.89	1.21	3.86	0.92	0.73
time (sec)	N/A	0.067	0.070	0.255	0.479	0.420	0.323	0.407	0.121

Problem 137	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	39	39	29	55	40	43	141	43	36
N.S.	1	1.00	0.74	1.41	1.03	1.10	3.62	1.10	0.92
time (sec)	N/A	0.047	0.046	0.250	0.263	0.338	0.267	0.423	1.059

Problem 138	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	39	39	29	55	41	42	146	43	35
N.S.	1	1.00	0.74	1.41	1.05	1.08	3.74	1.10	0.90
time (sec)	N/A	0.036	0.037	0.141	0.266	0.349	0.282	0.411	0.002

Problem 139	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	51	51	46	78	65	73	0	58	58
N.S.	1	1.00	0.90	1.53	1.27	1.43	0.00	1.14	1.14
time (sec)	N/A	0.057	0.064	0.622	0.268	0.376	0.000	0.416	0.412

Problem 140	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	60	60	64	100	86	271	0	75	73
N.S.	1	1.00	1.07	1.67	1.43	4.52	0.00	1.25	1.22
time (sec)	N/A	0.128	0.099	0.664	0.270	0.416	0.000	0.414	1.348

Problem 141	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	76	91	135	121	641	0	97	111
N.S.	1	1.00	1.20	1.78	1.59	8.43	0.00	1.28	1.46
time (sec)	N/A	0.215	0.142	0.641	0.285	0.428	0.000	0.412	1.416

Problem 142	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	97	97	108	177	173	1299	0	142	163
N.S.	1	1.00	1.11	1.82	1.78	13.39	0.00	1.46	1.68
time (sec)	N/A	0.336	0.232	0.685	0.283	0.417	0.000	0.419	1.502

Problem 143	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	55	55	49	73	68	182	0	174	69
N.S.	1	1.00	0.89	1.33	1.24	3.31	0.00	3.16	1.25
time (sec)	N/A	0.061	0.141	1.213	0.410	0.371	0.000	0.421	1.154

Problem 144	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	B	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	231	231	223	953	0	1516	0	0	-1
N.S.	1	1.00	0.97	4.13	0.00	6.56	0.00	0.00	-0.00
time (sec)	N/A	0.400	1.646	2.932	0.000	0.416	0.000	0.000	0.000

Problem 145	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	B	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	351	351	316	1186	0	2110	0	0	-1
N.S.	1	1.00	0.90	3.38	0.00	6.01	0.00	0.00	-0.00
time (sec)	N/A	0.634	1.654	2.862	0.000	0.403	0.000	0.000	0.000

Problem 146	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	29	29	64	24	23	28	0	23	21
N.S.	1	1.00	2.21	0.83	0.79	0.97	0.00	0.79	0.72
time (sec)	N/A	0.028	0.020	0.494	0.276	0.339	0.000	0.401	1.083

Problem 147	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	151	151	64	37	128	158	0	123	47
N.S.	1	1.00	0.42	0.25	0.85	1.05	0.00	0.81	0.31
time (sec)	N/A	0.088	8.661	0.615	0.489	0.350	0.000	0.399	1.096

Problem 148	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	35	41	19	22	0	19	25
N.S.	1	1.00	1.52	1.78	0.83	0.96	0.00	0.83	1.09
time (sec)	N/A	0.017	0.154	0.625	0.471	0.339	0.000	0.406	1.072

Problem 149	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	145	145	58	33	124	154	0	119	44
N.S.	1	1.00	0.40	0.23	0.86	1.06	0.00	0.82	0.30
time (sec)	N/A	0.067	0.133	0.615	0.498	0.367	0.000	0.412	1.068

Problem 150	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	12	26	10	18	15	20	15
N.S.	1	1.00	1.00	2.17	0.83	1.50	1.25	1.67	1.25
time (sec)	N/A	0.010	0.019	0.675	0.262	0.362	0.090	0.412	1.123

Problem 151	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	A	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	147	147	59	42	125	204	0	121	45
N.S.	1	1.00	0.40	0.29	0.85	1.39	0.00	0.82	0.31
time (sec)	N/A	0.075	0.125	0.612	0.483	0.377	0.000	0.427	1.091

Problem 152	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	40	44	19	21	0	16	24
N.S.	1	1.00	2.00	2.20	0.95	1.05	0.00	0.80	1.20
time (sec)	N/A	0.019	0.111	0.622	0.472	0.347	0.000	0.423	1.045

Problem 153	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	47	47	86	42	40	58	0	39	39
N.S.	1	1.00	1.83	0.89	0.85	1.23	0.00	0.83	0.83
time (sec)	N/A	0.042	0.085	0.380	0.268	0.355	0.000	0.414	1.122

Problem 154	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	173	173	174	53	144	231	0	139	67
N.S.	1	1.00	1.01	0.31	0.83	1.34	0.00	0.80	0.39
time (sec)	N/A	0.107	0.528	0.504	0.473	0.346	0.000	0.403	1.123

Problem 155	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	40	40	41	57	35	50	0	35	41
N.S.	1	1.00	1.02	1.42	0.88	1.25	0.00	0.88	1.02
time (sec)	N/A	0.033	0.271	0.507	0.474	0.371	0.000	0.416	1.050

Problem 156	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	A	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	165	165	146	47	138	228	0	133	61
N.S.	1	1.00	0.88	0.28	0.84	1.38	0.00	0.81	0.37
time (sec)	N/A	0.090	0.389	0.533	0.505	0.358	0.000	0.400	1.091

Problem 157	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	14	14	24	35	21	28	12	19	28
N.S.	1	1.00	1.71	2.50	1.50	2.00	0.86	1.36	2.00
time (sec)	N/A	0.016	0.027	0.710	0.267	0.334	0.120	0.408	1.065

Problem 158	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	A	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	190	190	181	64	146	274	0	143	68
N.S.	1	1.00	0.95	0.34	0.77	1.44	0.00	0.75	0.36
time (sec)	N/A	0.093	0.548	0.487	0.474	0.400	0.000	0.401	1.103

Problem 159	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	59	59	40	66	37	51	0	39	47
N.S.	1	1.00	0.68	1.12	0.63	0.86	0.00	0.66	0.80
time (sec)	N/A	0.036	0.268	0.497	0.477	0.370	0.000	0.418	1.063

Problem 160	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	60	60	47	0	0	0	0	0	-1
N.S.	1	1.00	0.78	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.025	0.068	0.405	0.000	0.000	0.000	0.000	0.000

Problem 161	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	79	79	79	0	0	0	0	0	-1
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.055	0.121	0.295	0.000	0.000	0.000	0.000	0.000

Problem 162	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	176	176	111	0	0	0	0	0	-1
N.S.	1	1.00	0.63	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.124	0.162	0.296	0.000	0.000	0.000	0.000	0.000

Problem 163	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	F	F	F	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	79	79	259	0	0	0	0	0	-1
N.S.	1	1.00	3.28	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.029	1.462	0.615	0.000	0.000	0.000	0.000	0.000

Problem 164	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	99	99	126	0	0	0	0	0	-1
N.S.	1	1.00	1.27	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.061	2.206	0.403	0.000	0.000	0.000	0.000	0.000

Problem 165	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	51	51	76	0	0	0	0	0	-1
N.S.	1	1.00	1.49	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.018	2.083	0.611	0.000	0.000	0.000	0.000	0.000

Problem 166	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	106	106	121	0	0	0	0	0	-1
N.S.	1	1.00	1.14	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.058	1.995	0.625	0.000	0.000	0.000	0.000	0.000

Problem 167	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	158	158	177	0	0	0	0	0	-1
N.S.	1	1.00	1.12	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.088	2.819	0.589	0.000	0.000	0.000	0.000	0.000

Problem 168	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	190	190	177	0	0	0	0	0	-1
N.S.	1	1.00	0.93	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.102	2.242	0.628	0.000	0.000	0.000	0.000	0.000

Problem 169	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	F	F	F	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	61	61	171	0	0	0	0	0	-1
N.S.	1	1.00	2.80	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.023	1.167	0.636	0.000	0.000	0.000	0.000	0.000

Problem 170	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	F	F	F	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	61	61	171	0	0	0	0	0	-1
N.S.	1	1.00	2.80	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.023	1.318	0.625	0.000	0.000	0.000	0.000	0.000

Problem 171	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	F	F	F	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	61	61	171	0	0	0	0	0	-1
N.S.	1	1.00	2.80	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.025	1.360	0.597	0.000	0.000	0.000	0.000	0.000

Problem 172	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	59	59	127	0	0	0	0	0	-1
N.S.	1	1.00	2.15	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.046	5.754	1.113	0.000	0.000	0.000	0.000	0.000

Problem 173	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	136	0	0	0	0	0	-1
N.S.	1	1.00	2.16	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.043	5.504	0.824	0.000	0.000	0.000	0.000	0.000

Problem 174	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	55	55	122	0	0	0	0	0	-1
N.S.	1	1.00	2.22	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.037	5.342	0.820	0.000	0.000	0.000	0.000	0.000

Problem 175	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	126	0	0	0	0	0	-1
N.S.	1	1.00	2.38	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.036	6.278	0.481	0.000	0.000	0.000	0.000	0.000

Problem 176	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	24	48	24	76	36	74	34
N.S.	1	1.00	0.96	1.92	0.96	3.04	1.44	2.96	1.36
time (sec)	N/A	0.015	0.040	2.108	0.273	0.474	1.882	0.497	1.061

Problem 177	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	59	59	126	0	0	0	0	0	-1
N.S.	1	1.00	2.14	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.042	2.288	0.482	0.000	0.000	0.000	0.000	0.000

Problem 178	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	56	56	120	0	0	0	0	0	-1
N.S.	1	1.00	2.14	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.043	2.182	0.477	0.000	0.000	0.000	0.000	0.000

Problem 179	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	133	133	159	0	0	0	0	0	-1
N.S.	1	1.00	1.20	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.124	5.706	0.709	0.000	0.000	0.000	0.000	0.000

Problem 180	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	137	137	169	0	0	0	0	0	-1
N.S.	1	1.00	1.23	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.116	5.538	0.469	0.000	0.000	0.000	0.000	0.000

Problem 181	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	131	131	155	0	0	0	0	0	-1
N.S.	1	1.00	1.18	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.111	5.539	0.472	0.000	0.000	0.000	0.000	0.000

Problem 182	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	127	127	163	0	0	0	0	0	-1
N.S.	1	1.00	1.28	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.107	6.453	0.471	0.000	0.000	0.000	0.000	0.000

Problem 183	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	28	28	51	63	36	72	70	37	34
N.S.	1	1.00	1.82	2.25	1.29	2.57	2.50	1.32	1.21
time (sec)	N/A	0.020	0.068	2.081	0.317	0.514	4.123	0.493	1.063

Problem 184	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	135	135	162	0	0	0	0	0	-1
N.S.	1	1.00	1.20	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.121	2.431	0.369	0.000	0.000	0.000	0.000	0.000

Problem 185	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	136	136	159	0	0	0	0	0	-1
N.S.	1	1.00	1.17	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.115	2.337	0.363	0.000	0.000	0.000	0.000	0.000

Problem 186	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	B	B	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	43	56	304	566	73	127	94
N.S.	1	1.00	1.00	1.30	7.07	13.16	1.70	2.95	2.19
time (sec)	N/A	0.028	0.111	2.069	0.314	0.375	0.773	0.470	1.067

Problem 187	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	B	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	45	45	62	69	494	194	70	67	162
N.S.	1	1.00	1.38	1.53	10.98	4.31	1.56	1.49	3.60
time (sec)	N/A	0.027	0.079	2.023	0.347	0.347	1.464	0.493	1.094

Problem 188	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	66	66	55	71	829	1568	92	161	227
N.S.	1	1.00	0.83	1.08	12.56	23.76	1.39	2.44	3.44
time (sec)	N/A	0.043	0.152	2.283	0.375	0.379	2.912	0.494	1.058

Problem 189	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	88	160	0	0	0	0	0	-1
N.S.	1	1.00	1.82	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.052	9.628	0.496	0.000	0.000	0.000	0.000	0.000

Problem 190	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	169	169	317	0	0	0	0	0	-1
N.S.	1	1.00	1.88	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.150	12.976	0.493	0.000	0.000	0.000	0.000	0.000

Problem 191	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	307	307	606	0	0	0	0	0	-1
N.S.	1	1.00	1.97	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.319	13.713	0.609	0.000	0.000	0.000	0.000	0.000

Problem 192	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	F	F	F	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	115	115	387	0	0	0	0	0	-1
N.S.	1	1.00	3.37	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.083	2.653	1.309	0.000	0.000	0.000	0.000	0.000

Problem 193	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	135	135	174	0	0	0	0	0	-1
N.S.	1	1.00	1.29	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.105	3.431	0.723	0.000	0.000	0.000	0.000	0.000

Problem 194	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	B	F(-2)	F(-1)	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	73	73	64	76	0	625	0	0	65
N.S.	1	1.00	0.88	1.04	0.00	8.56	0.00	0.00	0.89
time (sec)	N/A	0.039	0.239	4.853	0.000	0.391	0.000	0.000	2.247

Problem 195	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	B	F	F(-1)	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	70	57	74	0	334	0	0	51
N.S.	1	1.00	0.81	1.06	0.00	4.77	0.00	0.00	0.73
time (sec)	N/A	0.036	0.129	4.176	0.000	0.384	0.000	0.000	1.804

Problem 196	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	B	A	F(-1)	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	48	48	48	61	0	303	66	0	39
N.S.	1	1.00	1.00	1.27	0.00	6.31	1.38	0.00	0.81
time (sec)	N/A	0.028	0.060	4.148	0.000	0.362	1.562	0.000	1.452

Problem 197	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	B	A	F(-1)	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	47	47	47	37	0	305	66	0	36
N.S.	1	1.00	1.00	0.79	0.00	6.49	1.40	0.00	0.77
time (sec)	N/A	0.028	0.085	4.159	0.000	0.391	2.468	0.000	1.546

Problem 198	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	B	F	F(-1)	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	71	71	44	76	0	625	0	0	65
N.S.	1	1.00	0.62	1.07	0.00	8.80	0.00	0.00	0.92
time (sec)	N/A	0.037	0.121	4.120	0.000	0.355	0.000	0.000	1.718

Problem 199	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	B	F	F(-1)	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	46	74	0	1110	0	0	64
N.S.	1	1.00	0.64	1.03	0.00	15.42	0.00	0.00	0.89
time (sec)	N/A	0.036	0.153	4.203	0.000	0.368	0.000	0.000	2.476

Problem 200	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	B	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	135	135	136	149	0	8891	0	0	-1
N.S.	1	1.00	1.01	1.10	0.00	65.86	0.00	0.00	-0.01
time (sec)	N/A	0.251	0.964	1.220	0.000	1.202	0.000	0.000	0.000

Problem 201	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	B	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	105	105	105	90	0	6663	0	0	-1
N.S.	1	1.00	1.00	0.86	0.00	63.46	0.00	0.00	-0.01
time (sec)	N/A	0.142	0.052	0.991	0.000	1.005	0.000	0.000	0.000

Problem 202	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	B	F	F(-1)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	58	52	0	1748	0	0	-1
N.S.	1	1.00	1.00	0.90	0.00	30.14	0.00	0.00	-0.02
time (sec)	N/A	0.080	0.025	1.264	0.000	0.766	0.000	0.000	0.000

Problem 203	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	B	F	F(-1)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	106	106	109	0	0	6705	0	0	-1
N.S.	1	1.00	1.03	0.00	0.00	63.25	0.00	0.00	-0.01
time (sec)	N/A	0.164	0.262	1.970	0.000	0.949	0.000	0.000	0.000

Problem 204	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	B	F	F(-1)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	183	183	142	0	0	9168	0	0	-1
N.S.	1	1.00	0.78	0.00	0.00	50.10	0.00	0.00	-0.01
time (sec)	N/A	0.237	0.435	1.821	0.000	1.176	0.000	0.000	0.000

Problem 205	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	B	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	132	132	131	850	0	7896	0	0	-1
N.S.	1	1.00	0.99	6.44	0.00	59.82	0.00	0.00	-0.01
time (sec)	N/A	0.161	0.256	0.968	0.000	1.423	0.000	0.000	0.000

Problem 206	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	A	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	107	107	76	92	94	604	0	68	155
N.S.	1	1.00	0.71	0.86	0.88	5.64	0.00	0.64	1.45
time (sec)	N/A	0.048	0.125	1.451	0.468	0.368	0.000	0.419	0.101

Problem 207	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	A	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	77	77	60	80	69	339	0	52	93
N.S.	1	1.00	0.78	1.04	0.90	4.40	0.00	0.68	1.21
time (sec)	N/A	0.035	0.088	1.401	0.475	0.353	0.000	0.438	0.080

Problem 208	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	51	51	40	48	47	147	0	41	58
N.S.	1	1.00	0.78	0.94	0.92	2.88	0.00	0.80	1.14
time (sec)	N/A	0.024	0.073	0.454	0.462	0.334	0.000	0.401	1.067

Problem 209	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	22	27	23	38	0	23	34
N.S.	1	1.00	0.88	1.08	0.92	1.52	0.00	0.92	1.36
time (sec)	N/A	0.012	0.013	0.704	0.463	0.324	0.000	0.404	0.055

Problem 210	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	22	27	38	49	0	32	38
N.S.	1	1.00	0.88	1.08	1.52	1.96	0.00	1.28	1.52
time (sec)	N/A	0.012	0.016	0.714	0.254	0.344	0.000	0.441	0.083

Problem 211	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	A	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	179	48	62	198	0	56	62
N.S.	1	1.00	3.38	0.91	1.17	3.74	0.00	1.06	1.17
time (sec)	N/A	0.028	1.365	0.456	0.255	0.368	0.000	0.447	0.072

Problem 212	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	A	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	81	81	286	77	88	459	0	72	97
N.S.	1	1.00	3.53	0.95	1.09	5.67	0.00	0.89	1.20
time (sec)	N/A	0.036	1.800	1.612	0.259	0.361	0.000	0.440	1.110

Problem 213	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	113	113	115	88	110	796	0	83	160
N.S.	1	1.00	1.02	0.78	0.97	7.04	0.00	0.73	1.42
time (sec)	N/A	0.051	10.079	1.276	0.273	0.353	0.000	0.437	0.100

Problem 214	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	A	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	113	113	48	35	89	168	0	89	86
N.S.	1	1.00	0.42	0.31	0.79	1.49	0.00	0.79	0.76
time (sec)	N/A	0.061	0.035	0.810	0.469	0.364	0.000	0.407	0.246

Problem 215	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	95	95	95	24	78	113	0	78	81
N.S.	1	1.00	1.00	0.25	0.82	1.19	0.00	0.82	0.85
time (sec)	N/A	0.045	0.024	0.802	0.489	0.393	0.000	0.410	1.256

Problem 216	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	A	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	16	36	22	31	0	23	26
N.S.	1	1.00	1.00	2.25	1.38	1.94	0.00	1.44	1.62
time (sec)	N/A	0.010	0.011	0.849	0.476	0.329	0.000	0.408	0.167

Problem 217	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	A	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	113	48	34	230	0	35	38
N.S.	1	1.00	3.23	1.37	0.97	6.57	0.00	1.00	1.09
time (sec)	N/A	0.020	1.156	0.859	0.466	0.359	0.000	0.414	1.179

Problem 218	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	113	113	97	59	81	160	0	81	86
N.S.	1	1.00	0.86	0.52	0.72	1.42	0.00	0.72	0.76
time (sec)	N/A	0.156	0.062	0.954	0.484	0.358	0.000	0.415	0.323

Problem 219	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	97	97	24	47	69	107	0	69	70
N.S.	1	1.00	0.25	0.48	0.71	1.10	0.00	0.71	0.72
time (sec)	N/A	0.142	0.011	0.913	0.469	0.368	0.000	0.423	0.264

Problem 220	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	85	85	22	138	75	113	0	76	81
N.S.	1	1.00	0.26	1.62	0.88	1.33	0.00	0.89	0.95
time (sec)	N/A	0.089	0.013	0.793	0.482	0.334	0.000	0.414	0.259

Problem 221	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	A	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	108	108	113	150	87	628	0	88	93
N.S.	1	1.00	1.05	1.39	0.81	5.81	0.00	0.81	0.86
time (sec)	N/A	0.104	1.368	0.824	0.476	0.350	0.000	0.427	1.378

Problem 222	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	382	382	51	36	0	1373	0	263	474
N.S.	1	1.00	0.13	0.09	0.00	3.59	0.00	0.69	1.24
time (sec)	N/A	0.307	0.040	0.885	0.000	0.386	0.000	0.423	3.985

Problem 223	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	366	366	24	24	0	1089	0	251	457
N.S.	1	1.00	0.07	0.07	0.00	2.98	0.00	0.69	1.25
time (sec)	N/A	0.174	0.011	0.845	0.000	0.382	0.000	0.419	3.819

Problem 224	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	116	116	22	56	97	134	0	98	104
N.S.	1	1.00	0.19	0.48	0.84	1.16	0.00	0.84	0.90
time (sec)	N/A	0.052	0.015	0.985	0.499	0.353	0.000	0.409	1.333

Problem 225	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	A	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	134	134	113	68	109	213	0	110	122
N.S.	1	1.00	0.84	0.51	0.81	1.59	0.00	0.82	0.91
time (sec)	N/A	0.073	1.430	1.021	0.470	0.371	0.000	0.410	1.395

Problem 226	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F(-2)	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	107	107	81	87	0	440	0	328	163
N.S.	1	1.00	0.76	0.81	0.00	4.11	0.00	3.07	1.52
time (sec)	N/A	0.087	0.063	1.270	0.000	0.377	0.000	0.391	2.448

Problem 227	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F(-2)	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	152	152	107	191	0	1163	0	456	280
N.S.	1	1.00	0.70	1.26	0.00	7.65	0.00	3.00	1.84
time (sec)	N/A	0.127	0.121	1.373	0.000	0.397	0.000	0.391	23.355

Problem 228	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	167	167	205	0	0	0	0	0	-1
N.S.	1	1.00	1.23	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.132	2.934	1.444	0.000	0.000	0.000	0.000	0.000

Problem 229	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	117	117	169	0	0	0	0	0	-1
N.S.	1	1.00	1.44	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.092	2.315	1.301	0.000	0.000	0.000	0.000	0.000

Problem 230	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	67	67	141	0	0	0	0	0	-1
N.S.	1	1.00	2.10	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.048	1.125	1.026	0.000	0.000	0.000	0.000	0.000

Problem 231	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	65	65	134	0	0	0	0	0	-1
N.S.	1	1.00	2.06	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.047	1.433	1.173	0.000	0.000	0.000	0.000	0.000

Problem 232	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	113	113	145	0	0	0	0	0	-1
N.S.	1	1.00	1.28	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.086	2.163	1.183	0.000	0.000	0.000	0.000	0.000

Problem 233	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	161	161	185	0	0	0	0	0	-1
N.S.	1	1.00	1.15	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.122	2.154	1.295	0.000	0.000	0.000	0.000	0.000

Problem 234	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	A	B	F(-2)	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	311	311	133	324	145	1226	0	184	-1
N.S.	1	1.00	0.43	1.04	0.47	3.94	0.00	0.59	-0.00
time (sec)	N/A	0.628	0.167	4.819	0.481	0.358	0.000	0.451	0.000

Problem 235	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	A	B	F(-1)	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	193	193	104	301	90	458	0	129	-1
N.S.	1	1.00	0.54	1.56	0.47	2.37	0.00	0.67	-0.01
time (sec)	N/A	0.199	0.122	4.723	0.481	0.349	0.000	0.429	0.000

Problem 236	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	A	A	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	83	83	51	218	35	53	0	60	-1
N.S.	1	1.00	0.61	2.63	0.42	0.64	0.00	0.72	-0.01
time (sec)	N/A	0.105	0.040	4.914	0.483	0.328	0.000	0.409	0.000

Problem 237	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	83	83	51	213	56	70	0	88	-1
N.S.	1	1.00	0.61	2.57	0.67	0.84	0.00	1.06	-0.01
time (sec)	N/A	0.142	0.096	7.083	0.483	0.344	0.000	0.433	0.000

Problem 238	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	A	B	F	A	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	197	197	334	298	112	613	0	161	-1
N.S.	1	1.00	1.70	1.51	0.57	3.11	0.00	0.82	-0.01
time (sec)	N/A	0.599	7.256	6.768	0.485	0.357	0.000	0.487	0.000

Problem 239	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F(-1)	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	319	319	164	320	167	1617	0	215	-1
N.S.	1	1.00	0.51	1.00	0.52	5.07	0.00	0.67	-0.00
time (sec)	N/A	1.224	10.478	6.552	0.512	0.370	0.000	0.524	0.000

Problem 240	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	B	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	157	157	124	118	0	296	0	0	-1
N.S.	1	1.00	0.79	0.75	0.00	1.89	0.00	0.00	-0.01
time (sec)	N/A	0.278	0.175	1.288	0.000	0.383	0.000	0.000	0.000

Problem 241	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	115	115	88	88	0	155	0	0	-1
N.S.	1	1.00	0.77	0.77	0.00	1.35	0.00	0.00	-0.01
time (sec)	N/A	0.192	0.117	1.026	0.000	0.366	0.000	0.000	0.000

Problem 242	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	B	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	77	77	59	58	0	149	0	0	-1
N.S.	1	1.00	0.77	0.75	0.00	1.94	0.00	0.00	-0.01
time (sec)	N/A	0.108	0.100	0.966	0.000	0.348	0.000	0.000	0.000

Problem 243	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	A
verified	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	67	0	0	0	0	0	0	0	-1
N.S.	1	0.00	0.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.058	1.999	1.517	0.000	0.000	0.000	0.000	0.000

Problem 244	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	B	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	157	157	124	118	0	298	0	0	-1
N.S.	1	1.00	0.79	0.75	0.00	1.90	0.00	0.00	-0.01
time (sec)	N/A	0.276	0.219	1.288	0.000	0.395	0.000	0.000	0.000

Problem 245	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	115	115	88	88	0	155	0	0	-1
N.S.	1	1.00	0.77	0.77	0.00	1.35	0.00	0.00	-0.01
time (sec)	N/A	0.190	0.126	1.052	0.000	0.388	0.000	0.000	0.000

Problem 246	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	B	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	77	77	62	58	0	151	0	0	-1
N.S.	1	1.00	0.81	0.75	0.00	1.96	0.00	0.00	-0.01
time (sec)	N/A	0.106	0.082	0.992	0.000	0.369	0.000	0.000	0.000

Problem 247	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	A
verified	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	67	0	0	0	0	0	0	0	-1
N.S.	1	0.00	0.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.056	3.563	1.164	0.000	0.000	0.000	0.000	0.000

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is given. The larger this ratio is, the harder the integral was to solve. In this test, problem number [145] had the largest ratio of [26]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	4	2	1.00	8	0.250
2	A	3	2	1.00	8	0.250
3	A	3	2	1.00	8	0.250
4	A	2	2	1.00	8	0.250
5	A	2	2	1.00	8	0.250
6	A	1	1	1.00	6	0.167
7	A	1	1	1.00	6	0.167
8	A	2	2	1.00	8	0.250
9	A	2	2	1.00	8	0.250
10	A	3	2	1.00	8	0.250
11	A	3	2	1.00	8	0.250
12	A	4	2	1.00	8	0.250
13	A	7	6	1.00	12	0.500
14	A	6	6	1.00	12	0.500
15	A	6	6	1.00	12	0.500
16	A	5	5	1.00	12	0.417
17	A	5	5	1.00	12	0.417
18	A	6	6	1.00	12	0.500
19	A	6	6	1.00	12	0.500
20	A	7	6	1.00	12	0.500
21	A	9	9	1.00	8	1.125
22	A	2	2	1.00	8	0.250
23	A	2	2	1.00	10	0.200
24	A	3	3	1.00	10	0.300
25	A	2	2	1.00	10	0.200

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
26	A	2	2	1.00	10	0.200
27	A	4	3	1.00	14	0.214
28	A	3	3	1.00	14	0.214
29	A	2	2	1.00	14	0.143
30	A	2	2	1.00	14	0.143
31	A	3	3	1.00	14	0.214
32	A	4	3	1.00	14	0.214
33	A	7	7	1.00	8	0.875
34	A	8	7	1.00	10	0.700
35	A	7	7	1.00	10	0.700
36	A	7	7	1.00	10	0.700
37	A	5	3	1.00	10	0.300
38	A	3	3	1.00	10	0.300
39	A	3	3	1.00	10	0.300
40	A	3	3	1.00	12	0.250
41	A	5	3	1.00	12	0.250
42	A	4	3	1.00	12	0.250
43	A	3	3	1.00	12	0.250
44	A	2	2	1.00	12	0.167
45	A	2	2	1.00	12	0.167
46	A	3	2	1.00	12	0.167
47	A	4	2	1.00	12	0.167
48	A	5	2	1.00	12	0.167
49	A	6	2	1.00	12	0.167
50	A	5	3	1.00	8	0.375
51	A	4	3	1.00	8	0.375
52	A	3	3	1.00	8	0.375
53	A	2	2	1.00	8	0.250
54	A	3	3	1.00	8	0.375
55	A	4	3	1.00	8	0.375
56	A	5	3	1.00	8	0.375
57	A	5	4	1.00	12	0.333
58	A	4	4	1.00	12	0.333
59	A	3	3	1.00	12	0.250
60	A	2	2	1.00	12	0.167

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
61	A	2	2	1.00	12	0.167
62	A	3	3	1.00	12	0.250
63	A	4	4	1.00	12	0.333
64	A	5	4	1.00	12	0.333
65	A	2	2	1.00	12	0.167
66	A	2	2	1.00	12	0.167
67	A	5	4	1.00	14	0.286
68	A	5	4	1.00	14	0.286
69	A	5	4	1.00	11	0.364
70	A	9	7	1.00	11	0.636
71	A	5	4	1.00	11	0.364
72	A	8	6	1.00	9	0.667
73	A	8	7	1.00	9	0.778
74	A	3	2	1.00	11	0.182
75	A	8	7	1.00	11	0.636
76	A	4	3	1.00	11	0.273
77	A	9	8	1.00	11	0.727
78	A	4	3	1.00	11	0.273
79	A	10	8	1.00	11	0.727
80	A	5	3	1.00	13	0.231
81	A	10	9	1.00	13	0.692
82	A	4	3	1.00	13	0.231
83	A	6	6	1.00	11	0.546
84	A	6	5	1.00	11	0.454
85	A	3	2	1.00	13	0.154
86	A	15	11	1.00	13	0.846
87	A	3	2	1.00	13	0.154
88	A	29	13	1.00	13	1.000
89	A	3	2	1.00	13	0.154
90	A	6	5	1.00	11	0.454
91	A	4	3	1.00	11	0.273
92	A	3	2	1.00	11	0.182
93	A	4	3	1.00	11	0.273
94	A	2	2	1.00	9	0.222
95	A	1	1	1.00	9	0.111

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
96	A	2	2	1.00	11	0.182
97	A	2	2	1.00	11	0.182
98	A	2	1	1.00	11	0.091
99	A	3	3	1.00	11	0.273
100	A	3	2	1.00	11	0.182
101	A	4	3	1.00	11	0.273
102	A	3	2	1.00	13	0.154
103	A	3	2	1.00	13	0.154
104	A	3	2	1.00	13	0.154
105	A	2	2	1.00	13	0.154
106	A	2	2	1.00	8	0.250
107	A	4	3	1.00	13	0.231
108	A	5	4	1.00	13	0.308
109	A	14	6	1.19	13	0.462
110	A	9	6	1.07	13	0.462
111	A	5	5	1.00	13	0.385
112	A	2	2	1.00	11	0.182
113	A	5	5	1.00	11	0.454
114	A	9	6	1.00	13	0.462
115	A	5	4	1.00	11	0.364
116	A	4	4	1.00	11	0.364
117	A	3	3	1.00	11	0.273
118	A	3	2	1.00	11	0.182
119	A	2	2	1.00	9	0.222
120	A	2	2	1.00	6	0.333
121	A	4	4	1.00	9	0.444
122	A	4	4	1.00	11	0.364
123	A	5	4	1.00	11	0.364
124	A	6	4	1.00	11	0.364
125	A	4	4	1.00	11	0.364
126	A	3	3	1.00	11	0.273
127	A	3	3	1.00	11	0.273
128	A	4	4	1.00	11	0.364
129	A	4	4	1.00	13	0.308
130	A	3	3	1.00	13	0.231

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
131	A	4	4	1.00	13	0.308
132	A	4	4	1.00	13	0.308
133	A	7	7	1.00	13	0.538
134	A	6	6	1.00	13	0.462
135	A	5	5	1.00	13	0.385
136	A	4	4	1.00	13	0.308
137	A	2	2	1.00	11	0.182
138	A	2	2	1.00	8	0.250
139	A	3	3	1.00	11	0.273
140	A	4	4	1.00	13	0.308
141	A	5	5	1.00	13	0.385
142	A	6	6	1.00	13	0.462
143	A	3	3	1.00	14	0.214
144	A	9	6	1.00	24	0.250
145	A	11	7	1.00	26	0.269
146	A	4	3	1.00	11	0.273
147	A	11	8	1.00	11	0.727
148	A	4	4	1.00	9	0.444
149	A	11	8	1.00	7	1.143
150	A	2	1	1.00	11	0.091
151	A	11	8	1.00	11	0.727
152	A	4	4	1.00	11	0.364
153	A	4	3	1.00	13	0.231
154	A	12	9	1.00	13	0.692
155	A	5	5	1.00	11	0.454
156	A	13	9	1.00	9	1.000
157	A	3	2	1.00	13	0.154
158	A	12	9	1.00	13	0.692
159	A	5	5	1.00	13	0.385
160	A	3	3	1.00	13	0.231
161	A	4	4	1.00	15	0.267
162	A	5	5	1.00	15	0.333
163	A	3	3	1.00	9	0.333
164	A	3	3	1.00	15	0.200
165	A	2	2	1.00	11	0.182

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
166	A	4	4	1.00	11	0.364
167	A	5	5	1.00	11	0.454
168	A	5	5	1.00	11	0.454
169	A	3	3	1.00	7	0.429
170	A	3	3	1.00	9	0.333
171	A	3	3	1.00	9	0.333
172	A	4	4	1.00	17	0.235
173	A	4	4	1.00	17	0.235
174	A	4	4	1.00	15	0.267
175	A	4	4	1.00	13	0.308
176	A	2	1	1.00	17	0.059
177	A	4	4	1.00	17	0.235
178	A	4	4	1.00	17	0.235
179	A	5	5	1.00	19	0.263
180	A	5	5	1.00	19	0.263
181	A	5	5	1.00	17	0.294
182	A	5	5	1.00	15	0.333
183	A	3	2	1.00	19	0.105
184	A	5	5	1.00	19	0.263
185	A	5	5	1.00	19	0.263
186	A	3	2	1.00	17	0.118
187	A	4	2	1.00	17	0.118
188	A	4	2	1.00	17	0.118
189	A	4	4	1.00	19	0.210
190	A	5	5	1.00	21	0.238
191	A	6	6	1.00	21	0.286
192	A	4	4	1.00	15	0.267
193	A	4	4	1.00	21	0.190
194	A	7	6	1.00	19	0.316
195	A	7	6	1.00	19	0.316
196	A	6	5	1.00	19	0.263
197	A	6	5	1.00	19	0.263
198	A	7	6	1.00	19	0.316
199	A	7	6	1.00	19	0.316
200	A	8	7	1.00	23	0.304

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
201	A	7	6	1.00	23	0.261
202	A	4	4	1.00	21	0.190
203	A	8	5	1.00	21	0.238
204	A	11	6	1.00	23	0.261
205	A	8	7	1.00	21	0.333
206	A	7	6	1.00	16	0.375
207	A	7	6	1.00	16	0.375
208	A	5	4	1.00	16	0.250
209	A	3	3	1.00	14	0.214
210	A	3	3	1.00	14	0.214
211	A	5	4	1.00	16	0.250
212	A	7	6	1.00	16	0.375
213	A	7	6	1.00	16	0.375
214	A	13	9	1.00	10	0.900
215	A	11	8	1.00	8	1.000
216	A	5	5	1.00	8	0.625
217	A	7	6	1.00	10	0.600
218	A	14	9	1.00	10	0.900
219	A	12	8	1.00	8	1.000
220	A	12	8	1.00	8	1.000
221	A	14	9	1.00	10	0.900
222	A	23	9	1.00	10	0.900
223	A	21	8	1.00	8	1.000
224	A	15	12	1.00	8	1.500
225	A	17	13	1.00	10	1.300
226	A	5	5	1.00	14	0.357
227	A	7	6	1.00	14	0.429
228	A	6	3	1.00	18	0.167
229	A	5	3	1.00	18	0.167
230	A	4	3	1.00	16	0.188
231	A	4	3	1.00	16	0.188
232	A	5	3	1.00	18	0.167
233	A	6	3	1.00	18	0.167
234	A	9	7	1.00	25	0.280
235	A	8	7	1.00	25	0.280

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
236	A	4	4	1.00	25	0.160
237	A	4	4	1.00	25	0.160
238	A	8	7	1.00	25	0.280
239	A	9	7	1.00	25	0.280
240	A	19	5	1.00	9	0.556
241	A	13	5	1.00	9	0.556
242	A	9	4	1.00	7	0.571
243	A	0	0	0.00	0	0.000
244	A	19	5	1.00	9	0.556
245	A	13	5	1.00	9	0.556
246	A	9	4	1.00	7	0.571
247	A	0	0	0.00	0	0.000

Chapter 3

Listing of integrals

Local contents

3.1	$\int \tanh^6(a + bx) dx$	84
3.2	$\int \tanh^5(a + bx) dx$	87
3.3	$\int \tanh^4(a + bx) dx$	91
3.4	$\int \tanh^3(a + bx) dx$	94
3.5	$\int \tanh^2(a + bx) dx$	97
3.6	$\int \tanh(a + bx) dx$	100
3.7	$\int \coth(a + bx) dx$	103
3.8	$\int \coth^2(a + bx) dx$	106
3.9	$\int \coth^3(a + bx) dx$	109
3.10	$\int \coth^4(a + bx) dx$	113
3.11	$\int \coth^5(a + bx) dx$	116
3.12	$\int \coth^6(a + bx) dx$	120
3.13	$\int (b \tanh(c + dx))^{7/2} dx$	124
3.14	$\int (b \tanh(c + dx))^{5/2} dx$	129
3.15	$\int (b \tanh(c + dx))^{3/2} dx$	134
3.16	$\int \sqrt{b \tanh(c + dx)} dx$	139
3.17	$\int \frac{1}{\sqrt{b \tanh(c + dx)}} dx$	143
3.18	$\int \frac{1}{(b \tanh(c + dx))^{3/2}} dx$	147
3.19	$\int \frac{1}{(b \tanh(c + dx))^{5/2}} dx$	152
3.20	$\int \frac{1}{(b \tanh(c + dx))^{7/2}} dx$	157
3.21	$\int \sqrt[3]{\tanh(8x)} dx$	163
3.22	$\int \tanh^n(a + bx) dx$	168
3.23	$\int (b \tanh(c + dx))^n dx$	171
3.24	$\int (a \tanh^2(x))^{3/2} dx$	174
3.25	$\int \sqrt{a \tanh^2(x)} dx$	178

3.26	$\int \frac{1}{\sqrt{a \tanh^2(x)}} dx$	181
3.27	$\int (-\tanh^2(c+dx))^{5/2} dx$	184
3.28	$\int (-\tanh^2(c+dx))^{3/2} dx$	188
3.29	$\int \sqrt{-\tanh^2(c+dx)} dx$	192
3.30	$\int \frac{1}{\sqrt{-\tanh^2(c+dx)}} dx$	195
3.31	$\int \frac{1}{(-\tanh^2(c+dx))^{3/2}} dx$	198
3.32	$\int \frac{1}{(-\tanh^2(c+dx))^{5/2}} dx$	202
3.33	$\int \sqrt{\tanh^3(x)} dx$	206
3.34	$\int (a \tanh^3(x))^{3/2} dx$	211
3.35	$\int \sqrt{a \tanh^3(x)} dx$	217
3.36	$\int \frac{1}{\sqrt{a \tanh^3(x)}} dx$	222
3.37	$\int (a \tanh^4(x))^{3/2} dx$	227
3.38	$\int \sqrt{a \tanh^4(x)} dx$	232
3.39	$\int \frac{1}{\sqrt{a \tanh^4(x)}} dx$	236
3.40	$\int (b \tanh^m(c+dx))^n dx$	240
3.41	$\int (a + a \tanh(c+dx))^5 dx$	243
3.42	$\int (a + a \tanh(c+dx))^4 dx$	247
3.43	$\int (a + a \tanh(c+dx))^3 dx$	251
3.44	$\int (a + a \tanh(c+dx))^2 dx$	255
3.45	$\int \frac{1}{a+a \tanh(c+dx)} dx$	258
3.46	$\int \frac{1}{(a+a \tanh(c+dx))^2} dx$	261
3.47	$\int \frac{1}{(a+a \tanh(c+dx))^3} dx$	264
3.48	$\int \frac{1}{(a+a \tanh(c+dx))^4} dx$	268
3.49	$\int \frac{1}{(a+a \tanh(c+dx))^5} dx$	272
3.50	$\int (1 + \tanh(x))^{7/2} dx$	277
3.51	$\int (1 + \tanh(x))^{5/2} dx$	281
3.52	$\int (1 + \tanh(x))^{3/2} dx$	285
3.53	$\int \sqrt{1 + \tanh(x)} dx$	289
3.54	$\int \frac{1}{\sqrt{1 + \tanh(x)}} dx$	292
3.55	$\int \frac{1}{(1+\tanh(x))^{3/2}} dx$	296
3.56	$\int \frac{1}{(1+\tanh(x))^{5/2}} dx$	300
3.57	$\int (a + b \tanh(c+dx))^5 dx$	304
3.58	$\int (a + b \tanh(c+dx))^4 dx$	310
3.59	$\int (a + b \tanh(c+dx))^3 dx$	315

3.60	$\int (a + b \tanh(c + dx))^2 dx$	319
3.61	$\int \frac{1}{a + b \tanh(c + dx)} dx$	322
3.62	$\int \frac{1}{(a + b \tanh(c + dx))^2} dx$	326
3.63	$\int \frac{1}{(a + b \tanh(c + dx))^3} dx$	331
3.64	$\int \frac{1}{(a + b \tanh(c + dx))^4} dx$	337
3.65	$\int \frac{1}{4 + 6 \tanh(c + dx)} dx$	343
3.66	$\int \frac{1}{4 - 6 \tanh(c + dx)} dx$	346
3.67	$\int \sqrt{a + b \tanh(c + dx)} dx$	349
3.68	$\int \frac{1}{\sqrt{a + b \tanh(c + dx)}} dx$	354
3.69	$\int \frac{\sinh^4(x)}{1 + \tanh(x)} dx$	359
3.70	$\int \frac{\sinh^3(x)}{1 + \tanh(x)} dx$	363
3.71	$\int \frac{\sinh^2(x)}{1 + \tanh(x)} dx$	367
3.72	$\int \frac{\sinh(x)}{1 + \tanh(x)} dx$	371
3.73	$\int \frac{\operatorname{csch}(x)}{1 + \tanh(x)} dx$	375
3.74	$\int \frac{\operatorname{csch}^2(x)}{1 + \tanh(x)} dx$	379
3.75	$\int \frac{\operatorname{csch}^3(x)}{1 + \tanh(x)} dx$	382
3.76	$\int \frac{\operatorname{csch}^4(x)}{1 + \tanh(x)} dx$	386
3.77	$\int \frac{\operatorname{csch}^5(x)}{1 + \tanh(x)} dx$	390
3.78	$\int \frac{\operatorname{csch}^6(x)}{1 + \tanh(x)} dx$	395
3.79	$\int \frac{\operatorname{csch}^7(x)}{1 + \tanh(x)} dx$	399
3.80	$\int \frac{\sinh^4(x)}{a + b \tanh(x)} dx$	405
3.81	$\int \frac{\sinh^3(x)}{a + b \tanh(x)} dx$	410
3.82	$\int \frac{\sinh^2(x)}{a + b \tanh(x)} dx$	416
3.83	$\int \frac{\sinh(x)}{a + b \tanh(x)} dx$	420
3.84	$\int \frac{\operatorname{csch}(x)}{a + b \tanh(x)} dx$	424
3.85	$\int \frac{\operatorname{csch}^2(x)}{a + b \tanh(x)} dx$	428
3.86	$\int \frac{\operatorname{csch}^3(x)}{a + b \tanh(x)} dx$	431
3.87	$\int \frac{\operatorname{csch}^4(x)}{a + b \tanh(x)} dx$	437
3.88	$\int \frac{\operatorname{csch}^5(x)}{a + b \tanh(x)} dx$	441
3.89	$\int \frac{\operatorname{csch}^6(x)}{a + b \tanh(x)} dx$	449
3.90	$\int \frac{\operatorname{csch}(x)}{i + \tanh(x)} dx$	455
3.91	$\int \frac{\cosh^4(x)}{1 + \tanh(x)} dx$	459

3.92	$\int \frac{\cosh^3(x)}{1+\tanh(x)} dx$	463
3.93	$\int \frac{\cosh^2(x)}{1+\tanh(x)} dx$	466
3.94	$\int \frac{\cosh(x)}{1+\tanh(x)} dx$	469
3.95	$\int \frac{\operatorname{sech}(x)}{1+\tanh(x)} dx$	472
3.96	$\int \frac{\operatorname{sech}^2(x)}{1+\tanh(x)} dx$	475
3.97	$\int \frac{\operatorname{sech}^3(x)}{1+\tanh(x)} dx$	478
3.98	$\int \frac{\operatorname{sech}^4(x)}{1+\tanh(x)} dx$	481
3.99	$\int \frac{\operatorname{sech}^5(x)}{1+\tanh(x)} dx$	484
3.100	$\int \frac{\operatorname{sech}^6(x)}{1+\tanh(x)} dx$	488
3.101	$\int \frac{\operatorname{sech}^7(x)}{1+\tanh(x)} dx$	491
3.102	$\int \frac{\operatorname{sech}^8(x)}{a+b\tanh(x)} dx$	495
3.103	$\int \frac{\operatorname{sech}^6(x)}{a+b\tanh(x)} dx$	501
3.104	$\int \frac{\operatorname{sech}^4(x)}{a+b\tanh(x)} dx$	506
3.105	$\int \frac{\operatorname{sech}^2(x)}{a+b\tanh(x)} dx$	510
3.106	$\int \frac{1}{a+b\tanh(x)} dx$	513
3.107	$\int \frac{\cosh^2(x)}{a+b\tanh(x)} dx$	516
3.108	$\int \frac{\cosh^4(x)}{a+b\tanh(x)} dx$	520
3.109	$\int \frac{\operatorname{sech}^7(x)}{a+b\tanh(x)} dx$	525
3.110	$\int \frac{\operatorname{sech}^5(x)}{a+b\tanh(x)} dx$	531
3.111	$\int \frac{\operatorname{sech}^3(x)}{a+b\tanh(x)} dx$	536
3.112	$\int \frac{\operatorname{sech}(x)}{a+b\tanh(x)} dx$	540
3.113	$\int \frac{\cosh(x)}{a+b\tanh(x)} dx$	543
3.114	$\int \frac{\cosh^3(x)}{a+b\tanh(x)} dx$	547
3.115	$\int \frac{\tanh^5(x)}{1+\tanh(x)} dx$	552
3.116	$\int \frac{\tanh^4(x)}{1+\tanh(x)} dx$	556
3.117	$\int \frac{\tanh^3(x)}{1+\tanh(x)} dx$	560
3.118	$\int \frac{\tanh^2(x)}{1+\tanh(x)} dx$	563
3.119	$\int \frac{\tanh(x)}{1+\tanh(x)} dx$	566
3.120	$\int \frac{1}{1+\tanh(x)} dx$	569
3.121	$\int \frac{\operatorname{coth}(x)}{1+\tanh(x)} dx$	572
3.122	$\int \frac{\operatorname{coth}^2(x)}{1+\tanh(x)} dx$	575

3.123	$\int \frac{\coth^3(x)}{1+\tanh(x)} dx$	579
3.124	$\int \frac{\coth^4(x)}{1+\tanh(x)} dx$	583
3.125	$\int \tanh(x)(1+\tanh(x))^{3/2} dx$	587
3.126	$\int \tanh(x)\sqrt{1+\tanh(x)} dx$	591
3.127	$\int \frac{\tanh(x)}{\sqrt{1+\tanh(x)}} dx$	595
3.128	$\int \frac{\tanh(x)}{(1+\tanh(x))^{3/2}} dx$	599
3.129	$\int \tanh^2(x)(1+\tanh(x))^{3/2} dx$	603
3.130	$\int \tanh^2(x)\sqrt{1+\tanh(x)} dx$	607
3.131	$\int \frac{\tanh^2(x)}{\sqrt{1+\tanh(x)}} dx$	611
3.132	$\int \frac{\tanh^2(x)}{(1+\tanh(x))^{3/2}} dx$	615
3.133	$\int \frac{\tanh^5(x)}{a+b\tanh(x)} dx$	619
3.134	$\int \frac{\tanh^4(x)}{a+b\tanh(x)} dx$	625
3.135	$\int \frac{\tanh^3(x)}{a+b\tanh(x)} dx$	630
3.136	$\int \frac{\tanh^2(x)}{a+b\tanh(x)} dx$	634
3.137	$\int \frac{\tanh(x)}{a+b\tanh(x)} dx$	638
3.138	$\int \frac{1}{a+b\tanh(x)} dx$	641
3.139	$\int \frac{\coth(x)}{a+b\tanh(x)} dx$	644
3.140	$\int \frac{\coth^2(x)}{a+b\tanh(x)} dx$	647
3.141	$\int \frac{\coth^3(x)}{a+b\tanh(x)} dx$	651
3.142	$\int \frac{\coth^4(x)}{a+b\tanh(x)} dx$	656
3.143	$\int \frac{x\operatorname{sech}^2(x)}{(a+b\tanh(x))^2} dx$	661
3.144	$\int \frac{x\operatorname{sech}^2(c+dx)}{a+b\tanh^2(c+dx)} dx$	665
3.145	$\int \frac{x^2\operatorname{sech}^2(c+dx)}{a+b\tanh^2(c+dx)} dx$	671
3.146	$\int x^3 \tanh(a+2\log(x)) dx$	678
3.147	$\int x^2 \tanh(a+2\log(x)) dx$	681
3.148	$\int x \tanh(a+2\log(x)) dx$	685
3.149	$\int \tanh(a+2\log(x)) dx$	688
3.150	$\int \frac{\tanh(a+2\log(x))}{x} dx$	692
3.151	$\int \frac{\tanh(a+2\log(x))}{x^2} dx$	695
3.152	$\int \frac{\tanh(a+2\log(x))}{x^3} dx$	699
3.153	$\int x^3 \tanh^2(a+2\log(x)) dx$	702
3.154	$\int x^2 \tanh^2(a+2\log(x)) dx$	705
3.155	$\int x \tanh^2(a+2\log(x)) dx$	710
3.156	$\int \tanh^2(a+2\log(x)) dx$	713
3.157	$\int \frac{\tanh^2(a+2\log(x))}{x} dx$	718

3.158	$\int \frac{\tanh^2(a+2\log(x))}{x^2} dx$	721
3.159	$\int \frac{\tanh^2(a+2\log(x))}{x^3} dx$	726
3.160	$\int (ex)^m \tanh(a + 2\log(x)) dx$	730
3.161	$\int (ex)^m \tanh^2(a + 2\log(x)) dx$	733
3.162	$\int (ex)^m \tanh^3(a + 2\log(x)) dx$	736
3.163	$\int \tanh^p(a + b\log(x)) dx$	740
3.164	$\int (ex)^m \tanh^p(a + b\log(x)) dx$	743
3.165	$\int \tanh^p\left(a + \frac{\log(x)}{2}\right) dx$	746
3.166	$\int \tanh^p\left(a + \frac{\log(x)}{4}\right) dx$	749
3.167	$\int \tanh^p\left(a + \frac{\log(x)}{6}\right) dx$	752
3.168	$\int \tanh^p\left(a + \frac{\log(x)}{8}\right) dx$	756
3.169	$\int \tanh^p(a + \log(x)) dx$	760
3.170	$\int \tanh^p(a + 2\log(x)) dx$	763
3.171	$\int \tanh^p(a + 3\log(x)) dx$	766
3.172	$\int x^3 \tanh(d(a + b\log(cx^n))) dx$	769
3.173	$\int x^2 \tanh(d(a + b\log(cx^n))) dx$	772
3.174	$\int x \tanh(d(a + b\log(cx^n))) dx$	775
3.175	$\int \tanh(d(a + b\log(cx^n))) dx$	778
3.176	$\int \frac{\tanh(d(a+b\log(cx^n)))}{x} dx$	781
3.177	$\int \frac{\tanh(d(a+b\log(cx^n)))}{x^2} dx$	784
3.178	$\int \frac{\tanh(d(a+b\log(cx^n)))}{x^3} dx$	787
3.179	$\int x^3 \tanh^2(d(a + b\log(cx^n))) dx$	790
3.180	$\int x^2 \tanh^2(d(a + b\log(cx^n))) dx$	794
3.181	$\int x \tanh^2(d(a + b\log(cx^n))) dx$	798
3.182	$\int \tanh^2(d(a + b\log(cx^n))) dx$	802
3.183	$\int \frac{\tanh^2(d(a+b\log(cx^n)))}{x} dx$	806
3.184	$\int \frac{\tanh^2(d(a+b\log(cx^n)))}{x^2} dx$	809
3.185	$\int \frac{\tanh^2(d(a+b\log(cx^n)))}{x^3} dx$	813
3.186	$\int \frac{\tanh^3(a+b\log(cx^n))}{x} dx$	817
3.187	$\int \frac{\tanh^4(a+b\log(cx^n))}{x} dx$	821
3.188	$\int \frac{\tanh^5(a+b\log(cx^n))}{x} dx$	825
3.189	$\int (ex)^m \tanh(d(a + b\log(cx^n))) dx$	830
3.190	$\int (ex)^m \tanh^2(d(a + b\log(cx^n))) dx$	833
3.191	$\int (ex)^m \tanh^3(d(a + b\log(cx^n))) dx$	837
3.192	$\int \tanh^p(d(a + b\log(cx^n))) dx$	841
3.193	$\int (ex)^m \tanh^p(d(a + b\log(cx^n))) dx$	844
3.194	$\int \frac{\tanh^{\frac{5}{2}}(a+b\log(cx^n))}{x} dx$	847
3.195	$\int \frac{\tanh^{\frac{3}{2}}(a+b\log(cx^n))}{x} dx$	851
3.196	$\int \frac{\sqrt{\tanh(a + b\log(cx^n))}}{x} dx$	855

3.197	$\int \frac{1}{x \sqrt{\tanh(a + b \log(cx^n))}} dx$	859
3.198	$\int \frac{1}{x \tanh^{\frac{3}{2}}(a + b \log(cx^n))} dx$	863
3.199	$\int \frac{1}{x \tanh^{\frac{5}{2}}(a + b \log(cx^n))} dx$	868
3.200	$\int \frac{\tanh^5(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx$	873
3.201	$\int \frac{\tanh^3(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx$	880
3.202	$\int \frac{\tanh(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx$	886
3.203	$\int \frac{\coth(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx$	891
3.204	$\int \frac{\coth^3(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx$	897
3.205	$\int \tanh(x) \sqrt{a + b \tanh^2(x) + c \tanh^4(x)} dx$	903
3.206	$\int e^{a+bx} \tanh^4(a + bx) dx$	909
3.207	$\int e^{a+bx} \tanh^3(a + bx) dx$	914
3.208	$\int e^{a+bx} \tanh^2(a + bx) dx$	918
3.209	$\int e^{a+bx} \tanh(a + bx) dx$	922
3.210	$\int e^{a+bx} \coth(a + bx) dx$	925
3.211	$\int e^{a+bx} \coth^2(a + bx) dx$	928
3.212	$\int e^{a+bx} \coth^3(a + bx) dx$	932
3.213	$\int e^{a+bx} \coth^4(a + bx) dx$	937
3.214	$\int e^x \tanh^2(2x) dx$	942
3.215	$\int e^x \tanh(2x) dx$	947
3.216	$\int e^x \coth(2x) dx$	952
3.217	$\int e^x \coth^2(2x) dx$	955
3.218	$\int e^x \tanh^2(3x) dx$	959
3.219	$\int e^x \tanh(3x) dx$	964
3.220	$\int e^x \coth(3x) dx$	969
3.221	$\int e^x \coth^2(3x) dx$	974
3.222	$\int e^x \tanh^2(4x) dx$	979
3.223	$\int e^x \tanh(4x) dx$	986
3.224	$\int e^x \coth(4x) dx$	992
3.225	$\int e^x \coth^2(4x) dx$	997
3.226	$\int \frac{e^x}{a - \tanh(2x)} dx$	1003
3.227	$\int \frac{e^x}{(a - \tanh(2x))^2} dx$	1007
3.228	$\int e^{c(a+bx)} \tanh^3(d + ex) dx$	1013
3.229	$\int e^{c(a+bx)} \tanh^2(d + ex) dx$	1017
3.230	$\int e^{c(a+bx)} \tanh(d + ex) dx$	1020
3.231	$\int e^{c(a+bx)} \coth(d + ex) dx$	1023
3.232	$\int e^{c(a+bx)} \coth^2(d + ex) dx$	1026

3.233	$\int e^{c(a+bx)} \coth^3(d+ex) dx$	1029
3.234	$\int e^{c(a+bx)} \tanh^2(ac+bcx)^{5/2} dx$	1033
3.235	$\int e^{c(a+bx)} \tanh^2(ac+bcx)^{3/2} dx$	1039
3.236	$\int e^{c(a+bx)} \sqrt{\tanh^2(ac+bcx)} dx$	1044
3.237	$\int \frac{e^{c(a+bx)}}{\sqrt{\tanh^2(ac+bcx)}} dx$	1048
3.238	$\int \frac{e^{c(a+bx)}}{\tanh^2(ac+bcx)^{3/2}} dx$	1052
3.239	$\int \frac{e^{c(a+bx)}}{\tanh^2(ac+bcx)^{5/2}} dx$	1058
3.240	$\int \sin^3(\tanh(a+bx)) dx$	1064
3.241	$\int \sin^2(\tanh(a+bx)) dx$	1068
3.242	$\int \sin(\tanh(a+bx)) dx$	1072
3.243	$\int \csc(\tanh(a+bx)) dx$	1076
3.244	$\int \cos^3(\tanh(a+bx)) dx$	1079
3.245	$\int \cos^2(\tanh(a+bx)) dx$	1083
3.246	$\int \cos(\tanh(a+bx)) dx$	1087
3.247	$\int \sec(\tanh(a+bx)) dx$	1091

3.1 $\int \tanh^6(a + bx) dx$

Optimal. Leaf size=43

$$x - \frac{\tanh(a + bx)}{b} - \frac{\tanh^3(a + bx)}{3b} - \frac{\tanh^5(a + bx)}{5b}$$

[Out] $x - \tanh(b*x+a)/b - 1/3*\tanh(b*x+a)^3/b - 1/5*\tanh(b*x+a)^5/b$

Rubi [A]

time = 0.02, antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3554, 8}

$$-\frac{\tanh^5(a + bx)}{5b} - \frac{\tanh^3(a + bx)}{3b} - \frac{\tanh(a + bx)}{b} + x$$

Antiderivative was successfully verified.

[In] Int[Tanh[a + b*x]^6, x]

[Out] $x - \text{Tanh}[a + b*x]/b - \text{Tanh}[a + b*x]^3/(3*b) - \text{Tanh}[a + b*x]^5/(5*b)$

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 3554

Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Dist[b^2, Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]

Rubi steps

$$\begin{aligned} \int \tanh^6(a + bx) dx &= -\frac{\tanh^5(a + bx)}{5b} + \int \tanh^4(a + bx) dx \\ &= -\frac{\tanh^3(a + bx)}{3b} - \frac{\tanh^5(a + bx)}{5b} + \int \tanh^2(a + bx) dx \\ &= -\frac{\tanh(a + bx)}{b} - \frac{\tanh^3(a + bx)}{3b} - \frac{\tanh^5(a + bx)}{5b} + \int 1 dx \\ &= x - \frac{\tanh(a + bx)}{b} - \frac{\tanh^3(a + bx)}{3b} - \frac{\tanh^5(a + bx)}{5b} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 53, normalized size = 1.23

$$\frac{\tanh^{-1}(\tanh(a + bx))}{b} - \frac{\tanh(a + bx)}{b} - \frac{\tanh^3(a + bx)}{3b} - \frac{\tanh^5(a + bx)}{5b}$$

Antiderivative was successfully verified.

`[In] Integrate[Tanh[a + b*x]^6, x]``[Out] ArcTanh[Tanh[a + b*x]]/b - Tanh[a + b*x]/b - Tanh[a + b*x]^3/(3*b) - Tanh[a + b*x]^5/(5*b)`**Maple [A]**

time = 0.26, size = 56, normalized size = 1.30

method	result	size
derivativedivides	$\frac{-\frac{(\tanh^5(bx+a))}{5} - \frac{(\tanh^3(bx+a))}{3} - \tanh(bx+a) - \frac{\ln(-1+\tanh(bx+a))}{2} + \frac{\ln(\tanh(bx+a)+1)}{2}}{b}$	56
default	$\frac{-\frac{(\tanh^5(bx+a))}{5} - \frac{(\tanh^3(bx+a))}{3} - \tanh(bx+a) - \frac{\ln(-1+\tanh(bx+a))}{2} + \frac{\ln(\tanh(bx+a)+1)}{2}}{b}$	56
risch	$x + \frac{6e^{8bx+8a} + 12e^{6bx+6a} + \frac{56e^{4bx+4a}}{3} + \frac{28e^{2bx+2a}}{3} + \frac{46}{15}}{b(e^{2bx+2a}+1)^5}$	67

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(tanh(b*x+a)^6, x, method=_RETURNVERBOSE)``[Out] 1/b*(-1/5*tanh(b*x+a)^5-1/3*tanh(b*x+a)^3-tanh(b*x+a)-1/2*ln(-1+tanh(b*x+a))+1/2*ln(tanh(b*x+a)+1))`**Maxima [B]** Leaf count of result is larger than twice the leaf count of optimal. 115 vs. 2(39) = 78.

time = 0.28, size = 115, normalized size = 2.67

$$x + \frac{a}{b} - \frac{2(70e^{(-2bx-2a)} + 140e^{(-4bx-4a)} + 90e^{(-6bx-6a)} + 45e^{(-8bx-8a)} + 23)}{15b(5e^{(-2bx-2a)} + 10e^{(-4bx-4a)} + 10e^{(-6bx-6a)} + 5e^{(-8bx-8a)} + e^{(-10bx-10a)} + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(tanh(b*x+a)^6, x, algorithm="maxima")``[Out] x + a/b - 2/15*(70*e^(-2*b*x - 2*a) + 140*e^(-4*b*x - 4*a) + 90*e^(-6*b*x - 6*a) + 45*e^(-8*b*x - 8*a) + 23)/(b*(5*e^(-2*b*x - 2*a) + 10*e^(-4*b*x - 4*a) + 10*e^(-6*b*x - 6*a) + 5*e^(-8*b*x - 8*a) + e^(-10*b*x - 10*a) + 1))`**Fricas [B]** Leaf count of result is larger than twice the leaf count of optimal. 254 vs. 2(39) = 78.

time = 0.41, size = 254, normalized size = 5.91

(15bx + 23) cosh(bx + a)² + 5(15bx + 23) cosh(bx + a) sinh(bx + a) - 23 sinh(bx + a)² + 5(15bx + 23) cosh(bx + a)² - 5(46 cosh(bx + a)² + 5) sinh(bx + a)² + 5(2(15bx + 23) cosh(bx + a)² + 3(15bx + 23) cosh(bx + a) sinh(bx + a)² + 10(15bx + 23) cosh(bx + a) - 5(23 cosh(bx + a)⁴ + 15 cosh(bx + a)² + 10) sinh(bx + a)² + 15(b cosh(bx + a)⁵ + 5b cosh(bx + a) sinh(bx + a)³ + 5b cosh(bx + a)³ + 5(2b cosh(bx + a)² + 3b cosh(bx + a) sinh(bx + a)² + 10b cosh(bx + a))

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(b*x+a)^6,x, algorithm="fricas")

[Out] $\frac{1}{15}((15bx + 23)\cosh(bx + a)^5 + 5(15bx + 23)\cosh(bx + a)\sinh(bx + a)^4 - 23\sinh(bx + a)^5 + 5(15bx + 23)\cosh(bx + a)^3 - 5(46\cosh(bx + a)^2 + 5)\sinh(bx + a)^3 + 5(2(15bx + 23)\cosh(bx + a)^3 + 3(15bx + 23)\cosh(bx + a))\sinh(bx + a)^2 + 10(15bx + 23)\cosh(bx + a) - 5(23\cosh(bx + a)^4 + 15\cosh(bx + a)^2 + 10)\sinh(bx + a))/(b\cosh(bx + a)^5 + 5b\cosh(bx + a)\sinh(bx + a)^4 + 5b\cosh(bx + a)^3 + 5(2b\cosh(bx + a)^3 + 3b\cosh(bx + a))\sinh(bx + a)^2 + 10b\cosh(bx + a))$

Sympy [A]

time = 0.15, size = 39, normalized size = 0.91

$$\begin{cases} x - \frac{\tanh^5(a+bx)}{5b} - \frac{\tanh^3(a+bx)}{3b} - \frac{\tanh(a+bx)}{b} & \text{for } b \neq 0 \\ x \tanh^6(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(b*x+a)**6,x)

[Out] Piecewise((x - tanh(a + b*x)**5/(5*b) - tanh(a + b*x)**3/(3*b) - tanh(a + b*x)/b, Ne(b, 0)), (x*tanh(a)**6, True))

Giac [A]

time = 0.40, size = 74, normalized size = 1.72

$$\frac{15bx + 15a + \frac{2(45e^{(8bx+8a)} + 90e^{(6bx+6a)} + 140e^{(4bx+4a)} + 70e^{(2bx+2a)} + 23)}{(e^{(2bx+2a)} + 1)^5}}{15b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(b*x+a)^6,x, algorithm="giac")

[Out] $\frac{1}{15}(15bx + 15a + 2(45e^{(8bx+8a)} + 90e^{(6bx+6a)} + 140e^{(4bx+4a)} + 70e^{(2bx+2a)} + 23))/(e^{(2bx+2a)} + 1)^5/b$

Mupad [B]

time = 0.11, size = 34, normalized size = 0.79

$$x - \frac{\frac{\tanh(a+bx)^5}{5} + \frac{\tanh(a+bx)^3}{3} + \tanh(a+bx)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(a + b*x)^6,x)

[Out] $x - (\tanh(a + b*x) + \tanh(a + b*x)^3/3 + \tanh(a + b*x)^5/5)/b$

3.2 $\int \tanh^5(a + bx) dx$

Optimal. Leaf size=42

$$\frac{\log(\cosh(a + bx))}{b} - \frac{\tanh^2(a + bx)}{2b} - \frac{\tanh^4(a + bx)}{4b}$$

[Out] $\ln(\cosh(b*x+a))/b-1/2*\tanh(b*x+a)^2/b-1/4*\tanh(b*x+a)^4/b$

Rubi [A]

time = 0.02, antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3554, 3556}

$$-\frac{\tanh^4(a + bx)}{4b} - \frac{\tanh^2(a + bx)}{2b} + \frac{\log(\cosh(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Tanh}[a + b*x]^5, x]$

[Out] $\text{Log}[\text{Cosh}[a + b*x]]/b - \text{Tanh}[a + b*x]^2/(2*b) - \text{Tanh}[a + b*x]^4/(4*b)$

Rule 3554

$\text{Int}[(b*\tan[(c_.) + (d_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[b*((b*\text{Tan}[c + d*x])^{(n-1)})/(d*(n-1)), x] - \text{Dist}[b^2, \text{Int}[(b*\text{Tan}[c + d*x])^{(n-2)}, x], x] /;$ $\text{FreeQ}\{b, c, d, x\} \ \&\& \ \text{GtQ}[n, 1]$

Rule 3556

$\text{Int}[\tan[(c_.) + (d_.)*(x_.)], x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] /;$ $\text{FreeQ}\{c, d, x\}$

Rubi steps

$$\begin{aligned} \int \tanh^5(a + bx) dx &= -\frac{\tanh^4(a + bx)}{4b} + \int \tanh^3(a + bx) dx \\ &= -\frac{\tanh^2(a + bx)}{2b} - \frac{\tanh^4(a + bx)}{4b} + \int \tanh(a + bx) dx \\ &= \frac{\log(\cosh(a + bx))}{b} - \frac{\tanh^2(a + bx)}{2b} - \frac{\tanh^4(a + bx)}{4b} \end{aligned}$$

Mathematica [A]

time = 0.07, size = 37, normalized size = 0.88

$$\frac{4 \log(\cosh(a + bx)) - 2 \tanh^2(a + bx) - \tanh^4(a + bx)}{4b}$$

Antiderivative was successfully verified.

[In] Integrate[Tanh[a + b*x]^5,x]

[Out] (4*Log[Cosh[a + b*x]] - 2*Tanh[a + b*x]^2 - Tanh[a + b*x]^4)/(4*b)

Maple [A]

time = 0.26, size = 48, normalized size = 1.14

method	result	size
derivativedivides	$\frac{-\frac{(\tanh^4(bx+a))}{4} - \frac{(\tanh^2(bx+a))}{2} - \frac{\ln(-1+\tanh(bx+a))}{b} - \frac{\ln(\tanh(bx+a)+1)}{2}}{b}$	48
default	$\frac{-\frac{(\tanh^4(bx+a))}{4} - \frac{(\tanh^2(bx+a))}{2} - \frac{\ln(-1+\tanh(bx+a))}{b} - \frac{\ln(\tanh(bx+a)+1)}{2}}{b}$	48
risch	$-x - \frac{2a}{b} + \frac{4e^{2bx+2a}(e^{4bx+4a}+e^{2bx+2a}+1)}{b(e^{2bx+2a}+1)^4} + \frac{\ln(e^{2bx+2a}+1)}{b}$	74

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(b*x+a)^5,x,method=_RETURNVERBOSE)

[Out] 1/b*(-1/4*tanh(b*x+a)^4-1/2*tanh(b*x+a)^2-1/2*ln(-1+tanh(b*x+a))-1/2*ln(tanh(b*x+a)+1))

Maxima [B] Leaf count of result is larger than twice the leaf count of optimal. 102 vs. 2(38) = 76.

time = 0.47, size = 102, normalized size = 2.43

$$x + \frac{a}{b} + \frac{\log(e^{(-2bx-2a)} + 1)}{b} + \frac{4(e^{(-2bx-2a)} + e^{(-4bx-4a)} + e^{(-6bx-6a)})}{b(4e^{(-2bx-2a)} + 6e^{(-4bx-4a)} + 4e^{(-6bx-6a)} + e^{(-8bx-8a)} + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(b*x+a)^5,x, algorithm="maxima")

[Out] x + a/b + log(e^(-2*b*x - 2*a) + 1)/b + 4*(e^(-2*b*x - 2*a) + e^(-4*b*x - 4*a) + e^(-6*b*x - 6*a))/(b*(4*e^(-2*b*x - 2*a) + 6*e^(-4*b*x - 4*a) + 4*e^(-6*b*x - 6*a) + e^(-8*b*x - 8*a) + 1))

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 968 vs. 2(38) = 76.

time = 0.37, size = 968, normalized size = 23.05

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(b*x+a)^5,x, algorithm="fricas")

```
[Out] -(b*x*cosh(b*x + a)^8 + 8*b*x*cosh(b*x + a)*sinh(b*x + a)^7 + b*x*sinh(b*x + a)^8 + 4*(b*x - 1)*cosh(b*x + a)^6 + 4*(7*b*x*cosh(b*x + a)^2 + b*x - 1)*sinh(b*x + a)^6 + 8*(7*b*x*cosh(b*x + a)^3 + 3*(b*x - 1)*cosh(b*x + a))*sinh(b*x + a)^5 + 2*(3*b*x - 2)*cosh(b*x + a)^4 + 2*(35*b*x*cosh(b*x + a)^4 + 30*(b*x - 1)*cosh(b*x + a)^2 + 3*b*x - 2)*sinh(b*x + a)^4 + 8*(7*b*x*cosh(b*x + a)^5 + 10*(b*x - 1)*cosh(b*x + a)^3 + (3*b*x - 2)*cosh(b*x + a))*sinh(b*x + a)^3 + 4*(b*x - 1)*cosh(b*x + a)^2 + 4*(7*b*x*cosh(b*x + a)^6 + 15*(b*x - 1)*cosh(b*x + a)^4 + 3*(3*b*x - 2)*cosh(b*x + a)^2 + b*x - 1)*sinh(b*x + a)^2 + b*x - (cosh(b*x + a)^8 + 8*cosh(b*x + a)*sinh(b*x + a)^7 + sinh(b*x + a)^8 + 4*(7*cosh(b*x + a)^2 + 1)*sinh(b*x + a)^6 + 4*cosh(b*x + a)^6 + 8*(7*cosh(b*x + a)^3 + 3*cosh(b*x + a))*sinh(b*x + a)^5 + 2*(35*cosh(b*x + a)^4 + 30*cosh(b*x + a)^2 + 3)*sinh(b*x + a)^4 + 6*cosh(b*x + a)^4 + 8*(7*cosh(b*x + a)^5 + 10*cosh(b*x + a)^3 + 3*cosh(b*x + a))*sinh(b*x + a)^3 + 4*(7*cosh(b*x + a)^6 + 15*cosh(b*x + a)^4 + 9*cosh(b*x + a)^2 + 1)*sinh(b*x + a)^2 + 4*cosh(b*x + a)^2 + 8*(cosh(b*x + a)^7 + 3*cosh(b*x + a)^5 + 3*cosh(b*x + a)^3 + cosh(b*x + a))*sinh(b*x + a) + 1)*log(2*cosh(b*x + a)/(cosh(b*x + a) - sinh(b*x + a))) + 8*(b*x*cosh(b*x + a)^7 + 3*(b*x - 1)*cosh(b*x + a)^5 + (3*b*x - 2)*cosh(b*x + a)^3 + (b*x - 1)*cosh(b*x + a))*sinh(b*x + a))/(b*cosh(b*x + a)^8 + 8*b*cosh(b*x + a)*sinh(b*x + a)^7 + b*sinh(b*x + a)^8 + 4*b*cosh(b*x + a)^6 + 4*(7*b*cosh(b*x + a)^2 + b)*sinh(b*x + a)^6 + 8*(7*b*cosh(b*x + a)^3 + 3*b*cosh(b*x + a))*sinh(b*x + a)^5 + 6*b*cosh(b*x + a)^4 + 2*(35*b*cosh(b*x + a)^4 + 30*b*cosh(b*x + a)^2 + 3*b)*sinh(b*x + a)^4 + 8*(7*b*cosh(b*x + a)^5 + 10*b*cosh(b*x + a)^3 + 3*b*cosh(b*x + a))*sinh(b*x + a)^3 + 4*b*cosh(b*x + a)^2 + 4*(7*b*cosh(b*x + a)^6 + 15*b*cosh(b*x + a)^4 + 9*b*cosh(b*x + a)^2 + b)*sinh(b*x + a)^2 + 8*(b*cosh(b*x + a)^7 + 3*b*cosh(b*x + a)^5 + 3*b*cosh(b*x + a)^3 + b*cosh(b*x + a))*sinh(b*x + a) + b)
```

Sympy [A]

time = 0.12, size = 42, normalized size = 1.00

$$\begin{cases} x - \frac{\log(\tanh(\frac{a+bx}{b})+1)}{b} - \frac{\tanh^4(\frac{a+bx}{b})}{4b} - \frac{\tanh^2(\frac{a+bx}{b})}{2b} & \text{for } b \neq 0 \\ x \tanh^5(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(b*x+a)**5,x)

[Out] Piecewise((x - log(tanh(a + b*x) + 1)/b - tanh(a + b*x)**4/(4*b) - tanh(a + b*x)**2/(2*b), Ne(b, 0)), (x*tanh(a)**5, True))

Giac [A]

time = 0.42, size = 67, normalized size = 1.60

$$\frac{bx + a - \frac{4(e^{(6bx+6a)} + e^{(4bx+4a)} + e^{(2bx+2a)})}{(e^{(2bx+2a)} + 1)^4} - \log(e^{(2bx+2a)} + 1)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(b*x+a)^5,x, algorithm="giac")

[Out] $-(b*x + a - 4*(e^{6*b*x + 6*a}) + e^{(4*b*x + 4*a)} + e^{(2*b*x + 2*a)})/(e^{(2*b*x + 2*a)} + 1)^4 - \log(e^{(2*b*x + 2*a)} + 1))/b$

Mupad [B]

time = 0.12, size = 37, normalized size = 0.88

$$x - \frac{\ln(\tanh(a + bx) + 1) + \frac{\tanh(a+bx)^2}{2} + \frac{\tanh(a+bx)^4}{4}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(a + b*x)^5,x)

[Out] $x - (\log(\tanh(a + b*x) + 1) + \tanh(a + b*x)^2/2 + \tanh(a + b*x)^4/4)/b$

3.3 $\int \tanh^4(a + bx) dx$

Optimal. Leaf size=28

$$x - \frac{\tanh(a + bx)}{b} - \frac{\tanh^3(a + bx)}{3b}$$

[Out] x-tanh(b*x+a)/b-1/3*tanh(b*x+a)^3/b

Rubi [A]

time = 0.01, antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3554, 8}

$$-\frac{\tanh^3(a + bx)}{3b} - \frac{\tanh(a + bx)}{b} + x$$

Antiderivative was successfully verified.

[In] Int[Tanh[a + b*x]^4,x]

[Out] x - Tanh[a + b*x]/b - Tanh[a + b*x]^3/(3*b)

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 3554

Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Dist[b^2, Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]

Rubi steps

$$\begin{aligned} \int \tanh^4(a + bx) dx &= -\frac{\tanh^3(a + bx)}{3b} + \int \tanh^2(a + bx) dx \\ &= -\frac{\tanh(a + bx)}{b} - \frac{\tanh^3(a + bx)}{3b} + \int 1 dx \\ &= x - \frac{\tanh(a + bx)}{b} - \frac{\tanh^3(a + bx)}{3b} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 38, normalized size = 1.36

$$\frac{\tanh^{-1}(\tanh(a + bx))}{b} - \frac{\tanh(a + bx)}{b} - \frac{\tanh^3(a + bx)}{3b}$$

Antiderivative was successfully verified.

[In] Integrate[Tanh[a + b*x]^4, x]

[Out] ArcTanh[Tanh[a + b*x]]/b - Tanh[a + b*x]/b - Tanh[a + b*x]^3/(3*b)

Maple [A]

time = 0.26, size = 46, normalized size = 1.64

method	result	size
risch	$x + \frac{4e^{4bx+4a} + 4e^{2bx+2a} + \frac{8}{3}}{b(e^{2bx+2a} + 1)^3}$	45
derivativedivides	$\frac{-\frac{(\tanh^3(bx+a))}{3} - \tanh(bx+a) - \frac{\ln(-1+\tanh(bx+a))}{2} + \frac{\ln(\tanh(bx+a)+1)}{2}}{b}$	46
default	$\frac{-\frac{(\tanh^3(bx+a))}{3} - \tanh(bx+a) - \frac{\ln(-1+\tanh(bx+a))}{2} + \frac{\ln(\tanh(bx+a)+1)}{2}}{b}$	46

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(b*x+a)^4, x, method=_RETURNVERBOSE)

[Out] 1/b*(-1/3*tanh(b*x+a)^3-tanh(b*x+a)-1/2*ln(-1+tanh(b*x+a))+1/2*ln(tanh(b*x+a)+1))

Maxima [B] Leaf count of result is larger than twice the leaf count of optimal. 71 vs. 2(26) = 52.

time = 0.26, size = 71, normalized size = 2.54

$$x + \frac{a}{b} - \frac{4(3e^{(-2bx-2a)} + 3e^{(-4bx-4a)} + 2)}{3b(3e^{(-2bx-2a)} + 3e^{(-4bx-4a)} + e^{(-6bx-6a)} + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(b*x+a)^4, x, algorithm="maxima")

[Out] x + a/b - 4/3*(3*e^(-2*b*x - 2*a) + 3*e^(-4*b*x - 4*a) + 2)/(b*(3*e^(-2*b*x - 2*a) + 3*e^(-4*b*x - 4*a) + e^(-6*b*x - 6*a) + 1))

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 119 vs. 2(26) = 52.

time = 0.52, size = 119, normalized size = 4.25

$$\frac{(3bx+4)\cosh(bx+a)^3 + 3(3bx+4)\cosh(bx+a)\sinh(bx+a)^2 - 12\cosh(bx+a)^2\sinh(bx+a) - 4\sinh(bx+a)^3 + 3(3bx+4)\cosh(bx+a)}{3(b\cosh(bx+a)^3 + 3b\cosh(bx+a)\sinh(bx+a)^2 + 3b\cosh(bx+a))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(b*x+a)^4, x, algorithm="fricas")

[Out] 1/3*((3*b*x + 4)*cosh(b*x + a)^3 + 3*(3*b*x + 4)*cosh(b*x + a)*sinh(b*x + a)^2 - 12*cosh(b*x + a)^2*sinh(b*x + a) - 4*sinh(b*x + a)^3 + 3*(3*b*x + 4)*

$\cosh(b*x + a)/(b*\cosh(b*x + a)^3 + 3*b*\cosh(b*x + a)*\sinh(b*x + a)^2 + 3*b*\cosh(b*x + a))$

Sympy [A]

time = 0.09, size = 27, normalized size = 0.96

$$\begin{cases} x - \frac{\tanh^3(a+bx)}{3b} - \frac{\tanh(a+bx)}{b} & \text{for } b \neq 0 \\ x \tanh^4(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(b*x+a)**4,x)

[Out] Piecewise((x - tanh(a + b*x)**3/(3*b) - tanh(a + b*x)/b, Ne(b, 0)), (x*tanh(a)**4, True))

Giac [A]

time = 0.41, size = 52, normalized size = 1.86

$$\frac{3bx + 3a + \frac{4(3e^{(4bx+4a)} + 3e^{(2bx+2a)} + 2)}{(e^{(2bx+2a)} + 1)^3}}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(b*x+a)^4,x, algorithm="giac")

[Out] 1/3*(3*b*x + 3*a + 4*(3*e^(4*b*x + 4*a) + 3*e^(2*b*x + 2*a) + 2)/(e^(2*b*x + 2*a) + 1)^3)/b

Mupad [B]

time = 1.01, size = 24, normalized size = 0.86

$$x - \frac{\frac{\tanh(a+bx)^3}{3} + \tanh(a + bx)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(a + b*x)^4,x)

[Out] x - (tanh(a + b*x) + tanh(a + b*x)^3/3)/b

3.4 $\int \tanh^3(a + bx) dx$

Optimal. Leaf size=27

$$\frac{\log(\cosh(a + bx))}{b} - \frac{\tanh^2(a + bx)}{2b}$$

[Out] $\ln(\cosh(b*x+a))/b-1/2*\tanh(b*x+a)^2/b$

Rubi [A]

time = 0.01, antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3554, 3556}

$$\frac{\log(\cosh(a + bx))}{b} - \frac{\tanh^2(a + bx)}{2b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Tanh}[a + b*x]^3, x]$

[Out] $\text{Log}[\text{Cosh}[a + b*x]]/b - \text{Tanh}[a + b*x]^2/(2*b)$

Rule 3554

$\text{Int}[(b*.)*\tan[(c_.) + (d_.)*(x_.)]^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[b*((b*\text{Tan}[c + d*x])^{(n - 1)/(d*(n - 1))}), x] - \text{Dist}[b^2, \text{Int}[(b*\text{Tan}[c + d*x])^{(n - 2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x] \&\& \text{GtQ}[n, 1]$

Rule 3556

$\text{Int}[\tan[(c_.) + (d_.)*(x_.)], x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] /; \text{FreeQ}\{c, d\}, x]$

Rubi steps

$$\begin{aligned} \int \tanh^3(a + bx) dx &= -\frac{\tanh^2(a + bx)}{2b} + \int \tanh(a + bx) dx \\ &= \frac{\log(\cosh(a + bx))}{b} - \frac{\tanh^2(a + bx)}{2b} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 27, normalized size = 1.00

$$\frac{\log(\cosh(a + bx))}{b} - \frac{\tanh^2(a + bx)}{2b}$$

Antiderivative was successfully verified.

[In] Integrate[Tanh[a + b*x]^3,x]

[Out] Log[Cosh[a + b*x]]/b - Tanh[a + b*x]^2/(2*b)

Maple [A]

time = 0.26, size = 38, normalized size = 1.41

method	result	size
derivativedivides	$-\frac{\frac{\tanh^2(bx+a)}{2}}{b} - \frac{\ln(-1+\tanh(bx+a))}{b} - \frac{\ln(\tanh(bx+a)+1)}{2}$	38
default	$-\frac{\frac{\tanh^2(bx+a)}{2}}{b} - \frac{\ln(-1+\tanh(bx+a))}{b} - \frac{\ln(\tanh(bx+a)+1)}{2}$	38
risch	$-x - \frac{2a}{b} + \frac{2e^{2bx+2a}}{b(e^{2bx+2a}+1)^2} + \frac{\ln(e^{2bx+2a}+1)}{b}$	54

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(b*x+a)^3,x,method=_RETURNVERBOSE)

[Out] 1/b*(-1/2*tanh(b*x+a)^2-1/2*ln(-1+tanh(b*x+a))-1/2*ln(tanh(b*x+a)+1))

Maxima [B] Leaf count of result is larger than twice the leaf count of optimal. 61 vs. 2(25) = 50.

time = 0.46, size = 61, normalized size = 2.26

$$x + \frac{a}{b} + \frac{\log(e^{(-2bx-2a)} + 1)}{b} + \frac{2e^{(-2bx-2a)}}{b(2e^{(-2bx-2a)} + e^{(-4bx-4a)} + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(b*x+a)^3,x, algorithm="maxima")

[Out] x + a/b + log(e^(-2*b*x - 2*a) + 1)/b + 2*e^(-2*b*x - 2*a)/(b*(2*e^(-2*b*x - 2*a) + e^(-4*b*x - 4*a) + 1))

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 339 vs. 2(25) = 50.

time = 0.38, size = 339, normalized size = 12.56

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(b*x+a)^3,x, algorithm="fricas")

[Out] -(b*x*cosh(b*x + a))^4 + 4*b*x*cosh(b*x + a)*sinh(b*x + a)^3 + b*x*sinh(b*x + a)^4 + 2*(b*x - 1)*cosh(b*x + a)^2 + 2*(3*b*x*cosh(b*x + a)^2 + b*x - 1)*

$$\sinh(b*x + a)^2 + b*x - (\cosh(b*x + a)^4 + 4*\cosh(b*x + a)*\sinh(b*x + a)^3 + \sinh(b*x + a)^4 + 2*(3*\cosh(b*x + a)^2 + 1)*\sinh(b*x + a)^2 + 2*\cosh(b*x + a)^2 + 4*(\cosh(b*x + a)^3 + \cosh(b*x + a))*\sinh(b*x + a) + 1)*\log(2*\cosh(b*x + a)/(\cosh(b*x + a) - \sinh(b*x + a))) + 4*(b*x*\cosh(b*x + a)^3 + (b*x - 1)*\cosh(b*x + a))*\sinh(b*x + a)/(b*\cosh(b*x + a)^4 + 4*b*\cosh(b*x + a)*\sinh(b*x + a)^3 + b*\sinh(b*x + a)^4 + 2*b*\cosh(b*x + a)^2 + 2*(3*b*\cosh(b*x + a)^2 + b)*\sinh(b*x + a)^2 + 4*(b*\cosh(b*x + a)^3 + b*\cosh(b*x + a))*\sinh(b*x + a) + b)$$

Sympy [A]

time = 0.08, size = 31, normalized size = 1.15

$$\begin{cases} x - \frac{\log(\tanh(a+bx)+1)}{b} - \frac{\tanh^2(a+bx)}{2b} & \text{for } b \neq 0 \\ x \tanh^3(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(b*x+a)**3,x)

[Out] Piecewise((x - log(tanh(a + b*x) + 1)/b - tanh(a + b*x)**2/(2*b), Ne(b, 0)), (x*tanh(a)**3, True))

Giac [A]

time = 0.43, size = 48, normalized size = 1.78

$$-\frac{bx + a - \frac{2e^{(2bx+2a)}}{(e^{(2bx+2a)}+1)^2} - \log(e^{(2bx+2a)} + 1)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(b*x+a)^3,x, algorithm="giac")

[Out] -(b*x + a - 2*e^(2*b*x + 2*a)/(e^(2*b*x + 2*a) + 1)^2 - log(e^(2*b*x + 2*a) + 1))/b

Mupad [B]

time = 1.01, size = 27, normalized size = 1.00

$$x - \frac{\ln(\tanh(a + bx) + 1) + \frac{\tanh(a+bx)^2}{2}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(a + b*x)^3,x)

[Out] x - (log(tanh(a + b*x) + 1) + tanh(a + b*x)^2/2)/b

3.5 $\int \tanh^2(a + bx) dx$

Optimal. Leaf size=13

$$x - \frac{\tanh(a + bx)}{b}$$

[Out] x-tanh(b*x+a)/b

Rubi [A]

time = 0.01, antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3554, 8}

$$x - \frac{\tanh(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] Int[Tanh[a + b*x]^2,x]

[Out] x - Tanh[a + b*x]/b

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 3554

Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Dist[b^2, Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]

Rubi steps

$$\begin{aligned} \int \tanh^2(a + bx) dx &= -\frac{\tanh(a + bx)}{b} + \int 1 dx \\ &= x - \frac{\tanh(a + bx)}{b} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 23, normalized size = 1.77

$$\frac{\tanh^{-1}(\tanh(a + bx))}{b} - \frac{\tanh(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[Tanh[a + b*x]^2,x]

[Out] ArcTanh[Tanh[a + b*x]]/b - Tanh[a + b*x]/b

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 35 vs. 2(13) = 26.

time = 0.25, size = 36, normalized size = 2.77

method	result	size
risch	$x + \frac{2}{b(e^{2bx+2a}+1)}$	21
derivativedivides	$\frac{-\tanh(bx+a) - \frac{\ln(-1+\tanh(bx+a))}{2} + \frac{\ln(\tanh(bx+a)+1)}{2}}{b}$	36
default	$\frac{-\tanh(bx+a) - \frac{\ln(-1+\tanh(bx+a))}{2} + \frac{\ln(\tanh(bx+a)+1)}{2}}{b}$	36

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(b*x+a)^2,x,method=_RETURNVERBOSE)

[Out] 1/b*(-tanh(b*x+a)-1/2*ln(-1+tanh(b*x+a))+1/2*ln(tanh(b*x+a)+1))

Maxima [A]

time = 0.26, size = 25, normalized size = 1.92

$$x + \frac{a}{b} - \frac{2}{b(e^{-2bx-2a} + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(b*x+a)^2,x, algorithm="maxima")

[Out] x + a/b - 2/(b*(e^(-2*b*x - 2*a) + 1))

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 33 vs. 2(13) = 26.

time = 0.50, size = 33, normalized size = 2.54

$$\frac{(bx + 1) \cosh(bx + a) - \sinh(bx + a)}{b \cosh(bx + a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(b*x+a)^2,x, algorithm="fricas")

[Out] ((b*x + 1)*cosh(b*x + a) - sinh(b*x + a))/(b*cosh(b*x + a))

Sympy [A]

time = 0.06, size = 15, normalized size = 1.15

$$\begin{cases} x - \frac{\tanh(a+bx)}{b} & \text{for } b \neq 0 \\ x \tanh^2(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(b*x+a)**2,x)

[Out] Piecewise((x - tanh(a + b*x)/b, Ne(b, 0)), (x*tanh(a)**2, True))

Giac [A]

time = 0.41, size = 24, normalized size = 1.85

$$\frac{bx + a + \frac{2}{e^{(2bx+2a)+1}}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(b*x+a)^2,x, algorithm="giac")

[Out] (b*x + a + 2/(e^(2*b*x + 2*a) + 1))/b

Mupad [B]

time = 0.06, size = 13, normalized size = 1.00

$$x - \frac{\tanh(a + bx)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(a + b*x)^2,x)

[Out] x - tanh(a + b*x)/b

3.6 $\int \tanh(a + bx) dx$

Optimal. Leaf size=11

$$\frac{\log(\cosh(a + bx))}{b}$$

[Out] $\ln(\cosh(b*x+a))/b$

Rubi [A]

time = 0.00, antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3556}

$$\frac{\log(\cosh(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] `Int[Tanh[a + b*x], x]`

[Out] `Log[Cosh[a + b*x]]/b`

Rule 3556

`Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

Rubi steps

$$\int \tanh(a + bx) dx = \frac{\log(\cosh(a + bx))}{b}$$

Mathematica [A]

time = 0.01, size = 11, normalized size = 1.00

$$\frac{\log(\cosh(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] `Integrate[Tanh[a + b*x], x]`

[Out] `Log[Cosh[a + b*x]]/b`

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 27 vs. $2(11) = 22$.

time = 0.25, size = 28, normalized size = 2.55

method	result	size
risch	$-x - \frac{2a}{b} + \frac{\ln(e^{2bx+2a}+1)}{b}$	27
derivativdivides	$\frac{-\frac{\ln(-1+\tanh(bx+a))}{2} - \frac{\ln(\tanh(bx+a)+1)}{2}}{b}$	28
default	$\frac{-\frac{\ln(-1+\tanh(bx+a))}{2} - \frac{\ln(\tanh(bx+a)+1)}{2}}{b}$	28

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tanh(b*x+a),x,method=_RETURNVERBOSE)`

[Out] `1/b*(-1/2*ln(-1+tanh(b*x+a))-1/2*ln(tanh(b*x+a)+1))`

Maxima [A]

time = 0.26, size = 11, normalized size = 1.00

$$\frac{\log(\cosh(bx+a))}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(b*x+a),x,algorithm="maxima")`

[Out] `log(cosh(b*x + a))/b`

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 37 vs. 2(11) = 22.

time = 0.66, size = 37, normalized size = 3.36

$$-\frac{bx - \log\left(\frac{2 \cosh(bx+a)}{\cosh(bx+a) - \sinh(bx+a)}\right)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(b*x+a),x,algorithm="fricas")`

[Out] `-(b*x - log(2*cosh(b*x + a)/(cosh(b*x + a) - sinh(b*x + a))))/b`

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 17 vs. 2(8) = 16.

time = 0.06, size = 17, normalized size = 1.55

$$\begin{cases} x - \frac{\log(\tanh(a+bx)+1)}{b} & \text{for } b \neq 0 \\ x \tanh(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(b*x+a),x)

[Out] Piecewise((x - log(tanh(a + b*x) + 1)/b, Ne(b, 0)), (x*tanh(a), True))

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 24 vs. 2(11) = 22.
time = 0.40, size = 24, normalized size = 2.18

$$-\frac{bx + a - \log(e^{(2bx+2a)} + 1)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(b*x+a),x, algorithm="giac")

[Out] -(b*x + a - log(e^(2*b*x + 2*a) + 1))/b

Mupad [B]

time = 1.02, size = 16, normalized size = 1.45

$$x - \frac{\ln(\tanh(a + bx) + 1)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(a + b*x),x)

[Out] x - log(tanh(a + b*x) + 1)/b

3.7 $\int \coth(a + bx) dx$

Optimal. Leaf size=11

$$\frac{\log(\sinh(a + bx))}{b}$$

[Out] $\ln(\sinh(b*x+a))/b$

Rubi [A]

time = 0.00, antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3556}

$$\frac{\log(\sinh(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Coth}[a + b*x], x]$

[Out] $\text{Log}[\text{Sinh}[a + b*x]]/b$

Rule 3556

$\text{Int}[\tan[(c_.) + (d_.)*(x_.)], x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] /; \text{FreeQ}\{c, d\}, x]$

Rubi steps

$$\int \coth(a + bx) dx = \frac{\log(\sinh(a + bx))}{b}$$

Mathematica [A]

time = 0.01, size = 19, normalized size = 1.73

$$\frac{\log(\cosh(a + bx)) + \log(\tanh(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] $\text{Integrate}[\text{Coth}[a + b*x], x]$

[Out] $(\text{Log}[\text{Cosh}[a + b*x]] + \text{Log}[\text{Tanh}[a + b*x]])/b$

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 27 vs. $2(11) = 22$.

time = 0.26, size = 28, normalized size = 2.55

method	result	size
risch	$-x - \frac{2a}{b} + \frac{\ln(e^{2bx+2a}-1)}{b}$	27
derivativdivides	$\frac{-\frac{\ln(\coth(bx+a)-1)}{2} - \frac{\ln(\coth(bx+a)+1)}{2}}{b}$	28
default	$\frac{-\frac{\ln(\coth(bx+a)-1)}{2} - \frac{\ln(\coth(bx+a)+1)}{2}}{b}$	28

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(coth(b*x+a),x,method=_RETURNVERBOSE)`

[Out] $1/b*(-1/2*\ln(\coth(b*x+a)-1)-1/2*\ln(\coth(b*x+a)+1))$

Maxima [A]

time = 0.26, size = 11, normalized size = 1.00

$$\frac{\log(\sinh(bx+a))}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(coth(b*x+a),x, algorithm="maxima")`

[Out] $\log(\sinh(b*x + a))/b$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 37 vs. $2(11) = 22$.

time = 0.49, size = 37, normalized size = 3.36

$$-\frac{bx - \log\left(\frac{2 \sinh(bx+a)}{\cosh(bx+a) - \sinh(bx+a)}\right)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(coth(b*x+a),x, algorithm="fricas")`

[Out] $-(b*x - \log(2*\sinh(b*x + a)/(cosh(b*x + a) - sinh(b*x + a))))/b$

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 27 vs. $2(8) = 16$.

time = 0.22, size = 27, normalized size = 2.45

$$\begin{cases} x - \frac{\log(\tanh(a+bx)+1)}{b} + \frac{\log(\tanh(a+bx))}{b} & \text{for } b \neq 0 \\ x \coth(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(b*x+a),x)

[Out] Piecewise((x - log(tanh(a + b*x) + 1)/b + log(tanh(a + b*x))/b, Ne(b, 0)),
(x*coth(a), True))

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 25 vs. 2(11) = 22.
time = 0.40, size = 25, normalized size = 2.27

$$-\frac{bx + a - \log(|e^{(2bx+2a)} - 1|)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(b*x+a),x, algorithm="giac")

[Out] -(b*x + a - log(abs(e^(2*b*x + 2*a) - 1)))/b

Mupad [B]

time = 0.04, size = 21, normalized size = 1.91

$$\frac{\ln(e^{2a} e^{2bx} - 1)}{b} - x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(coth(a + b*x),x)

[Out] log(exp(2*a)*exp(2*b*x) - 1)/b - x

3.8 $\int \coth^2(a + bx) dx$

Optimal. Leaf size=13

$$x - \frac{\coth(a + bx)}{b}$$

[Out] x-coth(b*x+a)/b

Rubi [A]

time = 0.01, antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3554, 8}

$$x - \frac{\coth(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] Int[Coth[a + b*x]^2,x]

[Out] x - Coth[a + b*x]/b

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 3554

Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Dist[b^2, Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]

Rubi steps

$$\begin{aligned} \int \coth^2(a + bx) dx &= -\frac{\coth(a + bx)}{b} + \int 1 dx \\ &= x - \frac{\coth(a + bx)}{b} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 27, normalized size = 2.08

$$-\frac{\coth(a + bx) {}_2F_1\left(-\frac{1}{2}, 1; \frac{1}{2}; \tanh^2(a + bx)\right)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[Coth[a + b*x]^2,x]

[Out] -((Coth[a + b*x]*Hypergeometric2F1[-1/2, 1, 1/2, Tanh[a + b*x]^2])/b)

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 35 vs. $2(13) = 26$.

time = 0.25, size = 36, normalized size = 2.77

method	result	size
risch	$x - \frac{2}{b(e^{2bx+2a}-1)}$	21
derivativedivides	$-\frac{\coth(bx+a) - \frac{\ln(\coth(bx+a)-1)}{2} + \frac{\ln(\coth(bx+a)+1)}{2}}{b}$	36
default	$-\frac{\coth(bx+a) - \frac{\ln(\coth(bx+a)-1)}{2} + \frac{\ln(\coth(bx+a)+1)}{2}}{b}$	36

Verification of antiderivative is not currently implemented for this CAS.

[In] int(coth(b*x+a)^2,x,method=_RETURNVERBOSE)

[Out] 1/b*(-coth(b*x+a)-1/2*ln(coth(b*x+a)-1)+1/2*ln(coth(b*x+a)+1))

Maxima [A]

time = 0.27, size = 25, normalized size = 1.92

$$x + \frac{a}{b} + \frac{2}{b(e^{(-2bx-2a)} - 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(b*x+a)^2,x, algorithm="maxima")

[Out] x + a/b + 2/(b*(e^(-2*b*x - 2*a) - 1))

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 33 vs. $2(13) = 26$.

time = 0.53, size = 33, normalized size = 2.54

$$\frac{(bx + 1) \sinh(bx + a) - \cosh(bx + a)}{b \sinh(bx + a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(b*x+a)^2,x, algorithm="fricas")

[Out] ((b*x + 1)*sinh(b*x + a) - cosh(b*x + a))/(b*sinh(b*x + a))

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 87 vs. $2(8) = 16$.

time = 0.63, size = 87, normalized size = 6.69

$$\begin{cases} -\frac{\log(-e^{-bx}) \coth^2(bx + \log(-e^{-bx}))}{b} & \text{for } a = \log(-e^{-bx}) \\ -\frac{\log(e^{-bx}) \coth^2(bx + \log(e^{-bx}))}{b} & \text{for } a = \log(e^{-bx}) \\ x \coth^2(a) & \text{for } b = 0 \\ x - \frac{1}{b \tanh(a + bx)} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(b*x+a)**2,x)

[Out] Piecewise((-log(-exp(-b*x))*coth(b*x + log(-exp(-b*x))))**2/b, Eq(a, log(-exp(-b*x)))), (-log(exp(-b*x))*coth(b*x + log(exp(-b*x))))**2/b, Eq(a, log(exp(-b*x)))), (x*coth(a)**2, Eq(b, 0)), (x - 1/(b*tanh(a + b*x)), True))

Giac [A]

time = 0.40, size = 24, normalized size = 1.85

$$\frac{bx + a - \frac{2}{e^{(2bx+2a)} - 1}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(b*x+a)^2,x, algorithm="giac")

[Out] (b*x + a - 2/(e^(2*b*x + 2*a) - 1))/b

Mupad [B]

time = 1.02, size = 13, normalized size = 1.00

$$x - \frac{\coth(a + bx)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(coth(a + b*x)^2,x)

[Out] x - coth(a + b*x)/b

3.9 $\int \coth^3(a + bx) dx$

Optimal. Leaf size=27

$$-\frac{\coth^2(a + bx)}{2b} + \frac{\log(\sinh(a + bx))}{b}$$

[Out] $-1/2*\coth(b*x+a)^2/b+\ln(\sinh(b*x+a))/b$

Rubi [A]

time = 0.01, antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3554, 3556}

$$\frac{\log(\sinh(a + bx))}{b} - \frac{\coth^2(a + bx)}{2b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Coth}[a + b*x]^3, x]$

[Out] $-1/2*\text{Coth}[a + b*x]^2/b + \text{Log}[\text{Sinh}[a + b*x]]/b$

Rule 3554

$\text{Int}[(b \cdot \tan[(c \cdot x) + (d \cdot x)])^n, x_Symbol] \rightarrow \text{Simp}[b \cdot ((b \cdot \tan[c + d \cdot x])^{n-1}) / (d \cdot (n-1)), x] - \text{Dist}[b^2, \text{Int}[(b \cdot \tan[c + d \cdot x])^{n-2}, x], x] /;$ FreeQ[{b, c, d}, x] && GtQ[n, 1]

Rule 3556

$\text{Int}[\tan[(c \cdot x) + (d \cdot x)], x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d \cdot x], x]]/d, x] /;$ FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \coth^3(a + bx) dx &= -\frac{\coth^2(a + bx)}{2b} + \int \coth(a + bx) dx \\ &= -\frac{\coth^2(a + bx)}{2b} + \frac{\log(\sinh(a + bx))}{b} \end{aligned}$$

Mathematica [A]

time = 0.05, size = 34, normalized size = 1.26

$$\frac{\coth^2(a + bx) - 2 \log(\cosh(a + bx)) - 2 \log(\tanh(a + bx))}{2b}$$

$\sinh(b*x + a)^2 + b*x - (\cosh(b*x + a)^4 + 4*\cosh(b*x + a)*\sinh(b*x + a)^3 + \sinh(b*x + a)^4 + 2*(3*\cosh(b*x + a)^2 - 1)*\sinh(b*x + a)^2 - 2*\cosh(b*x + a)^2 + 4*(\cosh(b*x + a)^3 - \cosh(b*x + a))*\sinh(b*x + a) + 1)*\log(2*\sinh(b*x + a)/(\cosh(b*x + a) - \sinh(b*x + a))) + 4*(b*x*\cosh(b*x + a)^3 - (b*x - 1)*\cosh(b*x + a))*\sinh(b*x + a)/(b*\cosh(b*x + a)^4 + 4*b*\cosh(b*x + a)*\sinh(b*x + a)^3 + b*\sinh(b*x + a)^4 - 2*b*\cosh(b*x + a)^2 + 2*(3*b*\cosh(b*x + a)^2 - b)*\sinh(b*x + a)^2 + 4*(b*\cosh(b*x + a)^3 - b*\cosh(b*x + a))*\sinh(b*x + a) + b)$

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 112 vs. $2(20) = 40$.

time = 0.87, size = 112, normalized size = 4.15

$$\begin{cases} -\frac{\log(-e^{-bx}) \coth^3(bx + \log(-e^{-bx}))}{b} & \text{for } a = \log(-e^{-bx}) \\ -\frac{\log(e^{-bx}) \coth^3(bx + \log(e^{-bx}))}{b} & \text{for } a = \log(e^{-bx}) \\ x \coth^3(a) & \text{for } b = 0 \\ x - \frac{\log(\tanh(a+bx)+1)}{b} + \frac{\log(\tanh(a+bx))}{b} - \frac{1}{2b \tanh^2(a+bx)} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(b*x+a)**3,x)

[Out] Piecewise((-log(-exp(-b*x))*coth(b*x + log(-exp(-b*x))))**3/b, Eq(a, log(-exp(-b*x)))), (-log(exp(-b*x))*coth(b*x + log(exp(-b*x))))**3/b, Eq(a, log(exp(-b*x)))), (x*coth(a)**3, Eq(b, 0)), (x - log(tanh(a + b*x) + 1)/b + log(tanh(a + b*x))/b - 1/(2*b*tanh(a + b*x)**2), True))

Giac [A]

time = 0.40, size = 49, normalized size = 1.81

$$\frac{bx + a + \frac{2e^{(2bx+2a)}}{(e^{(2bx+2a)}-1)^2} - \log(|e^{(2bx+2a)} - 1|)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(b*x+a)^3,x, algorithm="giac")

[Out] $-(b*x + a + 2*e^{(2*b*x + 2*a)}/(e^{(2*b*x + 2*a)} - 1)^2 - \log(\text{abs}(e^{(2*b*x + 2*a)} - 1)))/b$

Mupad [B]

time = 0.05, size = 68, normalized size = 2.52

$$\frac{\ln(e^{2a} e^{2bx} - 1)}{b} - x - \frac{2}{b(e^{2a+2bx} - 1)} - \frac{2}{b(e^{4a+4bx} - 2e^{2a+2bx} + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(coth(a + b*x)^3,x)
```

```
[Out] log(exp(2*a)*exp(2*b*x) - 1)/b - x - 2/(b*(exp(2*a + 2*b*x) - 1)) - 2/(b*(exp(4*a + 4*b*x) - 2*exp(2*a + 2*b*x) + 1))
```

3.10 $\int \coth^4(a + bx) dx$

Optimal. Leaf size=28

$$x - \frac{\coth(a + bx)}{b} - \frac{\coth^3(a + bx)}{3b}$$

[Out] x-coth(b*x+a)/b-1/3*coth(b*x+a)^3/b

Rubi [A]

time = 0.01, antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3554, 8}

$$-\frac{\coth^3(a + bx)}{3b} - \frac{\coth(a + bx)}{b} + x$$

Antiderivative was successfully verified.

[In] Int[Coth[a + b*x]^4,x]

[Out] x - Coth[a + b*x]/b - Coth[a + b*x]^3/(3*b)

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 3554

Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Dist[b^2, Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]

Rubi steps

$$\begin{aligned} \int \coth^4(a + bx) dx &= -\frac{\coth^3(a + bx)}{3b} + \int \coth^2(a + bx) dx \\ &= -\frac{\coth(a + bx)}{b} - \frac{\coth^3(a + bx)}{3b} + \int 1 dx \\ &= x - \frac{\coth(a + bx)}{b} - \frac{\coth^3(a + bx)}{3b} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 31, normalized size = 1.11

$$-\frac{\coth^3(a + bx) {}_2F_1\left(-\frac{3}{2}, 1; -\frac{1}{2}; \tanh^2(a + bx)\right)}{3b}$$

Antiderivative was successfully verified.

[In] Integrate[Coth[a + b*x]^4,x]

[Out] $-1/3*(\text{Coth}[a + b*x]^3*\text{Hypergeometric2F1}[-3/2, 1, -1/2, \text{Tanh}[a + b*x]^2])/b$

Maple [A]

time = 0.26, size = 46, normalized size = 1.64

method	result	size
risch	$x - \frac{4(3e^{4bx+4a} - 3e^{2bx+2a} + 2)}{3b(e^{2bx+2a} - 1)^3}$	45
derivativedivides	$-\frac{(\coth^3(bx+a))}{3} - \coth(bx+a) - \frac{\ln(\coth(bx+a)-1)}{2} + \frac{\ln(\coth(bx+a)+1)}{2}$ b	46
default	$-\frac{(\coth^3(bx+a))}{3} - \coth(bx+a) - \frac{\ln(\coth(bx+a)-1)}{2} + \frac{\ln(\coth(bx+a)+1)}{2}$ b	46

Verification of antiderivative is not currently implemented for this CAS.

[In] int(coth(b*x+a)^4,x,method=_RETURNVERBOSE)

[Out] $1/b*(-1/3*\coth(b*x+a)^3 - \coth(b*x+a) - 1/2*\ln(\coth(b*x+a)-1) + 1/2*\ln(\coth(b*x+a)+1))$

Maxima [B] Leaf count of result is larger than twice the leaf count of optimal. 71 vs. $2(26) = 52$.

time = 0.26, size = 71, normalized size = 2.54

$$x + \frac{a}{b} - \frac{4(3e^{(-2bx-2a)} - 3e^{(-4bx-4a)} - 2)}{3b(3e^{(-2bx-2a)} - 3e^{(-4bx-4a)} + e^{(-6bx-6a)} - 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(b*x+a)^4,x, algorithm="maxima")

[Out] $x + a/b - 4/3*(3*e^{(-2*b*x - 2*a)} - 3*e^{(-4*b*x - 4*a)} - 2)/(b*(3*e^{(-2*b*x - 2*a)} - 3*e^{(-4*b*x - 4*a)} + e^{(-6*b*x - 6*a)} - 1))$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 108 vs. $2(26) = 52$.

time = 0.36, size = 108, normalized size = 3.86

$$\frac{(3bx+4)\sinh(bx+a)^3 - 4\cosh(bx+a)^3 - 12\cosh(bx+a)\sinh(bx+a)^2 + 3((3bx+4)\cosh(bx+a)^2 - 3bx-4)\sinh(bx+a)}{3(b\sinh(bx+a))^3 + 3(b\cosh(bx+a))^2 - b\sinh(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(b*x+a)^4,x, algorithm="fricas")

[Out] $\frac{1}{3}((3bx + 4)\sinh(bx + a)^3 - 4\cosh(bx + a)^3 - 12\cosh(bx + a)\sinh(bx + a)^2 + 3((3bx + 4)\cosh(bx + a)^2 - 3bx - 4)\sinh(bx + a))/(b\sinh(bx + a)^3 + 3(b\cosh(bx + a)^2 - b)\sinh(bx + a))$

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 100 vs. $2(20) = 40$.

time = 1.38, size = 100, normalized size = 3.57

$$\begin{cases} -\frac{\log(-e^{-bx}) \coth^4(bx + \log(-e^{-bx}))}{b} & \text{for } a = \log(-e^{-bx}) \\ -\frac{\log(e^{-bx}) \coth^4(bx + \log(e^{-bx}))}{b} & \text{for } a = \log(e^{-bx}) \\ x \coth^4(a) & \text{for } b = 0 \\ x - \frac{1}{b \tanh(a+bx)} - \frac{1}{3b \tanh^3(a+bx)} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(coth(b*x+a)**4,x)`

[Out] `Piecewise((-log(-exp(-b*x))*coth(b*x + log(-exp(-b*x))))**4/b, Eq(a, log(-exp(-b*x)))), (-log(exp(-b*x))*coth(b*x + log(exp(-b*x))))**4/b, Eq(a, log(exp(-b*x)))), (x*coth(a)**4, Eq(b, 0)), (x - 1/(b*tanh(a + b*x)) - 1/(3*b*tanh(a + b*x)**3), True))`

Giac [A]

time = 0.44, size = 52, normalized size = 1.86

$$\frac{3bx + 3a - \frac{4(3e^{(4bx+4a)} - 3e^{(2bx+2a)} + 2)}{(e^{(2bx+2a)} - 1)^3}}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(coth(b*x+a)^4,x, algorithm="giac")`

[Out] $\frac{1}{3}(3bx + 3a - 4(3e^{(4bx+4a)} - 3e^{(2bx+2a)} + 2)/(e^{(2bx+2a)} - 1)^3)/b$

Mupad [B]

time = 0.06, size = 24, normalized size = 0.86

$$x - \frac{\frac{\coth(a+bx)^3}{3} + \coth(a+bx)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(coth(a + b*x)^4,x)`

[Out] $x - (\coth(a + b*x) + \coth(a + b*x)^3/3)/b$

3.11 $\int \coth^5(a + bx) dx$

Optimal. Leaf size=42

$$-\frac{\coth^2(a + bx)}{2b} - \frac{\coth^4(a + bx)}{4b} + \frac{\log(\sinh(a + bx))}{b}$$

[Out] $-1/2*\coth(b*x+a)^2/b-1/4*\coth(b*x+a)^4/b+\ln(\sinh(b*x+a))/b$

Rubi [A]

time = 0.03, antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3554, 3556}

$$-\frac{\coth^4(a + bx)}{4b} - \frac{\coth^2(a + bx)}{2b} + \frac{\log(\sinh(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] Int[Coth[a + b*x]^5,x]

[Out] $-1/2*\text{Coth}[a + b*x]^2/b - \text{Coth}[a + b*x]^4/(4*b) + \text{Log}[\text{Sinh}[a + b*x]]/b$

Rule 3554

Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Dist[b^2, Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]

Rule 3556

Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \coth^5(a + bx) dx &= -\frac{\coth^4(a + bx)}{4b} + \int \coth^3(a + bx) dx \\ &= -\frac{\coth^2(a + bx)}{2b} - \frac{\coth^4(a + bx)}{4b} + \int \coth(a + bx) dx \\ &= -\frac{\coth^2(a + bx)}{2b} - \frac{\coth^4(a + bx)}{4b} + \frac{\log(\sinh(a + bx))}{b} \end{aligned}$$

Mathematica [A]

time = 0.11, size = 44, normalized size = 1.05

$$\frac{2 \coth^2(a + bx) + \coth^4(a + bx) - 4 \log(\cosh(a + bx)) - 4 \log(\tanh(a + bx))}{4b}$$

Antiderivative was successfully verified.

[In] Integrate[Coth[a + b*x]^5,x]

[Out] $-1/4*(2*\text{Coth}[a + b*x]^2 + \text{Coth}[a + b*x]^4 - 4*\text{Log}[\text{Cosh}[a + b*x]] - 4*\text{Log}[\text{Tanh}[a + b*x]])/b$

Maple [A]

time = 0.26, size = 48, normalized size = 1.14

method	result	size
derivativedivides	$\frac{-\frac{(\coth^4(bx+a))}{4} - \frac{(\coth^2(bx+a))}{2} - \frac{\ln(\coth(bx+a)-1)}{2} - \frac{\ln(\coth(bx+a)+1)}{2}}{b}$	48
default	$\frac{-\frac{(\coth^4(bx+a))}{4} - \frac{(\coth^2(bx+a))}{2} - \frac{\ln(\coth(bx+a)-1)}{2} - \frac{\ln(\coth(bx+a)+1)}{2}}{b}$	48
risch	$-x - \frac{2a}{b} - \frac{4e^{2bx+2a}(e^{4bx+4a} - e^{2bx+2a} + 1)}{b(e^{2bx+2a} - 1)^4} + \frac{\ln(e^{2bx+2a} - 1)}{b}$	76

Verification of antiderivative is not currently implemented for this CAS.

[In] int(coth(b*x+a)^5,x,method=_RETURNVERBOSE)

[Out] $1/b*(-1/4*\text{coth}(b*x+a)^4 - 1/2*\text{coth}(b*x+a)^2 - 1/2*\ln(\text{coth}(b*x+a)-1) - 1/2*\ln(\text{coth}(b*x+a)+1))$

Maxima [B] Leaf count of result is larger than twice the leaf count of optimal. 122 vs. 2(38) = 76.

time = 0.26, size = 122, normalized size = 2.90

$$x + \frac{a}{b} + \frac{\log(e^{(-bx-a)} + 1)}{b} + \frac{\log(e^{(-bx-a)} - 1)}{b} + \frac{4(e^{(-2bx-2a)} - e^{(-4bx-4a)} + e^{(-6bx-6a)})}{b(4e^{(-2bx-2a)} - 6e^{(-4bx-4a)} + 4e^{(-6bx-6a)} - e^{(-8bx-8a)} - 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(b*x+a)^5,x, algorithm="maxima")

[Out] $x + a/b + \log(e^{(-b*x - a)} + 1)/b + \log(e^{(-b*x - a)} - 1)/b + 4*(e^{(-2*b*x - 2*a)} - e^{(-4*b*x - 4*a)} + e^{(-6*b*x - 6*a)})/(b*(4*e^{(-2*b*x - 2*a)} - 6*e^{(-4*b*x - 4*a)} + 4*e^{(-6*b*x - 6*a)} - e^{(-8*b*x - 8*a)} - 1))$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 978 vs. 2(38) = 76.

time = 0.42, size = 978, normalized size = 23.29

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(b*x+a)^5,x, algorithm="fricas")

```
[Out] -(b*x*cosh(b*x + a)^8 + 8*b*x*cosh(b*x + a)*sinh(b*x + a)^7 + b*x*sinh(b*x + a)^8 - 4*(b*x - 1)*cosh(b*x + a)^6 + 4*(7*b*x*cosh(b*x + a)^2 - b*x + 1)*sinh(b*x + a)^6 + 8*(7*b*x*cosh(b*x + a)^3 - 3*(b*x - 1)*cosh(b*x + a))*sinh(b*x + a)^5 + 2*(3*b*x - 2)*cosh(b*x + a)^4 + 2*(35*b*x*cosh(b*x + a)^4 - 30*(b*x - 1)*cosh(b*x + a)^2 + 3*b*x - 2)*sinh(b*x + a)^4 + 8*(7*b*x*cosh(b*x + a)^5 - 10*(b*x - 1)*cosh(b*x + a)^3 + (3*b*x - 2)*cosh(b*x + a))*sinh(b*x + a)^3 - 4*(b*x - 1)*cosh(b*x + a)^2 + 4*(7*b*x*cosh(b*x + a)^6 - 15*(b*x - 1)*cosh(b*x + a)^4 + 3*(3*b*x - 2)*cosh(b*x + a)^2 - b*x + 1)*sinh(b*x + a)^2 + b*x - (cosh(b*x + a)^8 + 8*cosh(b*x + a)*sinh(b*x + a)^7 + sinh(b*x + a)^8 + 4*(7*cosh(b*x + a)^2 - 1)*sinh(b*x + a)^6 - 4*cosh(b*x + a)^6 + 8*(7*cosh(b*x + a)^3 - 3*cosh(b*x + a))*sinh(b*x + a)^5 + 2*(35*cosh(b*x + a)^4 - 30*cosh(b*x + a)^2 + 3)*sinh(b*x + a)^4 + 6*cosh(b*x + a)^4 + 8*(7*cosh(b*x + a)^5 - 10*cosh(b*x + a)^3 + 3*cosh(b*x + a))*sinh(b*x + a)^3 + 4*(7*cosh(b*x + a)^6 - 15*cosh(b*x + a)^4 + 9*cosh(b*x + a)^2 - 1)*sinh(b*x + a)^2 - 4*cosh(b*x + a)^2 + 8*(cosh(b*x + a)^7 - 3*cosh(b*x + a)^5 + 3*cosh(b*x + a)^3 - cosh(b*x + a))*sinh(b*x + a) + 1)*log(2*sinh(b*x + a)/(cosh(b*x + a) - sinh(b*x + a))) + 8*(b*x*cosh(b*x + a)^7 - 3*(b*x - 1)*cosh(b*x + a)^5 + (3*b*x - 2)*cosh(b*x + a)^3 - (b*x - 1)*cosh(b*x + a))*sinh(b*x + a))/(b*cosh(b*x + a)^8 + 8*b*cosh(b*x + a)*sinh(b*x + a)^7 + b*sinh(b*x + a)^8 - 4*b*cosh(b*x + a)^6 + 4*(7*b*cosh(b*x + a)^2 - b)*sinh(b*x + a)^6 + 8*(7*b*cosh(b*x + a)^3 - 3*b*cosh(b*x + a))*sinh(b*x + a)^5 + 6*b*cosh(b*x + a)^4 + 2*(35*b*cosh(b*x + a)^4 - 30*b*cosh(b*x + a)^2 + 3*b)*sinh(b*x + a)^4 + 8*(7*b*cosh(b*x + a)^5 - 10*b*cosh(b*x + a)^3 + 3*b*cosh(b*x + a))*sinh(b*x + a)^3 - 4*b*cosh(b*x + a)^2 + 4*(7*b*cosh(b*x + a)^6 - 15*b*cosh(b*x + a)^4 + 9*b*cosh(b*x + a)^2 - b)*sinh(b*x + a)^2 + 8*(b*cosh(b*x + a)^7 - 3*b*cosh(b*x + a)^5 + 3*b*cosh(b*x + a)^3 - b*cosh(b*x + a))*sinh(b*x + a) + b)
```

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 126 vs. $2(32) = 64$.

time = 2.41, size = 126, normalized size = 3.00

$$\begin{cases} -\frac{\log(-e^{-bx}) \coth^5(bx + \log(-e^{-bx}))}{b} & \text{for } a = \log(-e^{-bx}) \\ -\frac{\log(e^{-bx}) \coth^5(bx + \log(e^{-bx}))}{b} & \text{for } a = \log(e^{-bx}) \\ x \coth^5(a) & \text{for } b = 0 \\ x - \frac{\log(\tanh(a+bx)+1)}{b} + \frac{\log(\tanh(a+bx))}{b} - \frac{1}{2b \tanh^2(a+bx)} - \frac{1}{4b \tanh^4(a+bx)} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(b*x+a)**5,x)

```
[Out] Piecewise((-log(-exp(-b*x))*coth(b*x + log(-exp(-b*x))))**5/b, Eq(a, log(-exp(-b*x))), (-log(exp(-b*x))*coth(b*x + log(exp(-b*x))))**5/b, Eq(a, log(exp(-b*x))), (x*coth(a)**5, Eq(b, 0)), (x - log(tanh(a + b*x) + 1)/b + log(ta
```

```
nh(a + b*x))/b - 1/(2*b*tanh(a + b*x)**2) - 1/(4*b*tanh(a + b*x)**4), True)
)
```

Giac [A]

time = 0.41, size = 70, normalized size = 1.67

$$\frac{bx + a + \frac{4(e^{(6bx+6a)} - e^{(4bx+4a)} + e^{(2bx+2a)})}{(e^{(2bx+2a)} - 1)^4} - \log(|e^{(2bx+2a)} - 1|)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(coth(b*x+a)^5,x, algorithm="giac")
```

```
[Out] -(b*x + a + 4*(e^(6*b*x + 6*a) - e^(4*b*x + 4*a) + e^(2*b*x + 2*a))/(e^(2*b*x + 2*a) - 1)^4 - log(abs(e^(2*b*x + 2*a) - 1)))/b
```

Mupad [B]

time = 0.99, size = 159, normalized size = 3.79

$$\frac{\ln(e^{2a}e^{2bx} - 1)}{b} - x - \frac{4}{b(e^{2a+2bx} - 1)} - \frac{8}{b(e^{4a+4bx} - 2e^{2a+2bx} + 1)} - \frac{8}{b(3e^{2a+2bx} - 3e^{4a+4bx} + e^{6a+6bx} - 1)} - \frac{4}{b(6e^{4a+4bx} - 4e^{2a+2bx} - 4e^{6a+6bx} + e^{8a+8bx} + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(coth(a + b*x)^5,x)
```

```
[Out] log(exp(2*a)*exp(2*b*x) - 1)/b - x - 4/(b*(exp(2*a + 2*b*x) - 1)) - 8/(b*(exp(4*a + 4*b*x) - 2*exp(2*a + 2*b*x) + 1)) - 8/(b*(3*exp(2*a + 2*b*x) - 3*exp(4*a + 4*b*x) + exp(6*a + 6*b*x) - 1)) - 4/(b*(6*exp(4*a + 4*b*x) - 4*exp(2*a + 2*b*x) - 4*exp(6*a + 6*b*x) + exp(8*a + 8*b*x) + 1))
```

3.12 $\int \coth^6(a + bx) dx$

Optimal. Leaf size=43

$$x - \frac{\coth(a + bx)}{b} - \frac{\coth^3(a + bx)}{3b} - \frac{\coth^5(a + bx)}{5b}$$

[Out] $x - \coth(b*x+a)/b - 1/3*\coth(b*x+a)^3/b - 1/5*\coth(b*x+a)^5/b$

Rubi [A]

time = 0.02, antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3554, 8}

$$-\frac{\coth^5(a + bx)}{5b} - \frac{\coth^3(a + bx)}{3b} - \frac{\coth(a + bx)}{b} + x$$

Antiderivative was successfully verified.

[In] Int[Coth[a + b*x]^6, x]

[Out] $x - \text{Coth}[a + b*x]/b - \text{Coth}[a + b*x]^3/(3*b) - \text{Coth}[a + b*x]^5/(5*b)$

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 3554

Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Dist[b^2, Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]

Rubi steps

$$\begin{aligned} \int \coth^6(a + bx) dx &= -\frac{\coth^5(a + bx)}{5b} + \int \coth^4(a + bx) dx \\ &= -\frac{\coth^3(a + bx)}{3b} - \frac{\coth^5(a + bx)}{5b} + \int \coth^2(a + bx) dx \\ &= -\frac{\coth(a + bx)}{b} - \frac{\coth^3(a + bx)}{3b} - \frac{\coth^5(a + bx)}{5b} + \int 1 dx \\ &= x - \frac{\coth(a + bx)}{b} - \frac{\coth^3(a + bx)}{3b} - \frac{\coth^5(a + bx)}{5b} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 31, normalized size = 0.72

$$\frac{\coth^5(a + bx) {}_2F_1\left(-\frac{5}{2}, 1; -\frac{3}{2}; \tanh^2(a + bx)\right)}{5b}$$

Antiderivative was successfully verified.

[In] Integrate[Coth[a + b*x]^6, x]

[Out] -1/5*(Coth[a + b*x]^5*Hypergeometric2F1[-5/2, 1, -3/2, Tanh[a + b*x]^2])/b

Maple [A]

time = 0.26, size = 56, normalized size = 1.30

method	result	size
derivativedivides	$\frac{-\frac{\coth^5(bx+a)}{5} - \frac{\coth^3(bx+a)}{3} - \coth(bx+a) - \frac{\ln(\coth(bx+a)-1)}{2} + \frac{\ln(\coth(bx+a)+1)}{2}}{b}$	56
default	$\frac{-\frac{\coth^5(bx+a)}{5} - \frac{\coth^3(bx+a)}{3} - \coth(bx+a) - \frac{\ln(\coth(bx+a)-1)}{2} + \frac{\ln(\coth(bx+a)+1)}{2}}{b}$	56
risch	$x - \frac{2(45e^{8bx+8a} - 90e^{6bx+6a} + 140e^{4bx+4a} - 70e^{2bx+2a} + 23)}{15b(e^{2bx+2a}-1)^5}$	67

Verification of antiderivative is not currently implemented for this CAS.

[In] int(coth(b*x+a)^6, x, method=_RETURNVERBOSE)

[Out] 1/b*(-1/5*coth(b*x+a)^5-1/3*coth(b*x+a)^3-coth(b*x+a)-1/2*ln(coth(b*x+a)-1)+1/2*ln(coth(b*x+a)+1))

Maxima [B] Leaf count of result is larger than twice the leaf count of optimal. 115 vs. 2(39) = 78.

time = 0.27, size = 115, normalized size = 2.67

$$x + \frac{a}{b} - \frac{2(70e^{(-2bx-2a)} - 140e^{(-4bx-4a)} + 90e^{(-6bx-6a)} - 45e^{(-8bx-8a)} - 23)}{15b(5e^{(-2bx-2a)} - 10e^{(-4bx-4a)} + 10e^{(-6bx-6a)} - 5e^{(-8bx-8a)} + e^{(-10bx-10a)} - 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(b*x+a)^6, x, algorithm="maxima")

[Out] x + a/b - 2/15*(70*e^(-2*b*x - 2*a) - 140*e^(-4*b*x - 4*a) + 90*e^(-6*b*x - 6*a) - 45*e^(-8*b*x - 8*a) - 23)/(b*(5*e^(-2*b*x - 2*a) - 10*e^(-4*b*x - 4*a) + 10*e^(-6*b*x - 6*a) - 5*e^(-8*b*x - 8*a) + e^(-10*b*x - 10*a) - 1))

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 239 vs. 2(39) = 78.

time = 0.38, size = 239, normalized size = 5.56

(15bx + 23)sinh(bx + a)^2 - 23cosh(bx + a)^2 - 115cosh(bx + a)sinh(bx + a) + 5(2(15bx + 23)cosh(bx + a)^2 - 15bx - 23)sinh(bx + a)^2 + 25cosh(bx + a)^2 - 5(46cosh(bx + a) - 15cosh(bx + a)sinh(bx + a) + 5(15bx + 23)cosh(bx + a)^2 - 3(15bx + 23)cosh(bx + a)^2 + 30bx + 46)sinh(bx + a) - 50cosh(bx + a)sinh(bx + a)^2 + 5(2kosh(bx + a)^2 - 8)sinh(bx + a)^2 + 5(kosh(bx + a)^2 - 3kosh(bx + a)^2 + 24)sinh(bx + a)

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(b*x+a)^6,x, algorithm="fricas")

[Out] $\frac{1}{15}((15bx + 23)\sinh(bx + a)^5 - 23\cosh(bx + a)^5 - 115\cosh(bx + a)\sinh(bx + a)^4 + 5(2(15bx + 23)\cosh(bx + a)^2 - 15bx - 23)\sinh(bx + a)^3 + 25\cosh(bx + a)^3 - 5(46\cosh(bx + a)^3 - 15\cosh(bx + a))\sinh(bx + a)^2 + 5((15bx + 23)\cosh(bx + a)^4 - 3(15bx + 23)\cosh(bx + a)^2 + 30bx + 46)\sinh(bx + a) - 50\cosh(bx + a))/(b\sinh(bx + a)^5 + 5(2b\cosh(bx + a)^2 - b)\sinh(bx + a)^3 + 5(b\cosh(bx + a)^4 - 3b\cosh(bx + a)^2 + 2b)\sinh(bx + a))$

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 114 vs. $2(32) = 64$.

time = 3.85, size = 114, normalized size = 2.65

$$\begin{cases} -\frac{\log(-e^{-bx}) \coth^6(bx + \log(-e^{-bx}))}{b} & \text{for } a = \log(-e^{-bx}) \\ -\frac{\log(e^{-bx}) \coth^6(bx + \log(e^{-bx}))}{b} & \text{for } a = \log(e^{-bx}) \\ x \coth^6(a) & \text{for } b = 0 \\ x - \frac{1}{b \tanh(a+bx)} - \frac{1}{3b \tanh^3(a+bx)} - \frac{1}{5b \tanh^5(a+bx)} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(b*x+a)**6,x)

[Out] Piecewise((-log(-exp(-b*x))*coth(b*x + log(-exp(-b*x))))**6/b, Eq(a, log(-exp(-b*x)))), (-log(exp(-b*x))*coth(b*x + log(exp(-b*x))))**6/b, Eq(a, log(exp(-b*x)))), (x*coth(a)**6, Eq(b, 0)), (x - 1/(b*tanh(a + b*x)) - 1/(3*b*tanh(a + b*x)**3) - 1/(5*b*tanh(a + b*x)**5), True))

Giac [A]

time = 0.44, size = 74, normalized size = 1.72

$$\frac{15bx + 15a - \frac{2(45e^{(8bx+8a)} - 90e^{(6bx+6a)} + 140e^{(4bx+4a)} - 70e^{(2bx+2a)} + 23)}{(e^{(2bx+2a)} - 1)^5}}{15b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(b*x+a)^6,x, algorithm="giac")

[Out] $\frac{1}{15}(15bx + 15a - 2(45e^{(8bx+8a)} - 90e^{(6bx+6a)} + 140e^{(4bx+4a)} - 70e^{(2bx+2a)} + 23))/(e^{(2bx+2a)} - 1)^5/b$

Mupad [B]

time = 0.07, size = 34, normalized size = 0.79

$$x - \frac{\frac{\coth(a+bx)^5}{5} + \frac{\coth(a+bx)^3}{3} + \coth(a+bx)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(coth(a + b*x)^6,x)`

[Out] $x - (\coth(a + b*x) + \coth(a + b*x)^3/3 + \coth(a + b*x)^5/5)/b$

3.13 $\int (b \tanh(c + dx))^{7/2} dx$

Optimal. Leaf size=97

$$\frac{b^{7/2} \operatorname{ArcTan}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{d} + \frac{b^{7/2} \tanh^{-1}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{d} - \frac{2b^3 \sqrt{b \tanh(c + dx)}}{d} - \frac{2b(b \tanh(c + dx))^{5/2}}{5d}$$

[Out] $b^{(7/2)*\arctan((b*\tanh(d*x+c))^{(1/2)}/b^{(1/2)})/d+b^{(7/2)*\operatorname{arctanh}((b*\tanh(d*x+c))^{(1/2)}/b^{(1/2)})/d-2*b^3*(b*\tanh(d*x+c))^{(1/2)}/d-2/5*b*(b*\tanh(d*x+c))^{(5/2)}/d}$

Rubi [A]

time = 0.06, antiderivative size = 97, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3554, 3557, 335, 218, 212, 209}

$$\frac{b^{7/2} \operatorname{ArcTan}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{d} + \frac{b^{7/2} \tanh^{-1}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{d} - \frac{2b^3 \sqrt{b \tanh(c + dx)}}{d} - \frac{2b(b \tanh(c + dx))^{5/2}}{5d}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(b*\operatorname{Tanh}[c + d*x])^{(7/2)}, x]$

[Out] $(b^{(7/2)*\operatorname{ArcTan}[\operatorname{Sqrt}[b*\operatorname{Tanh}[c + d*x]]/\operatorname{Sqrt}[b]])/d + (b^{(7/2)*\operatorname{ArcTanh}[\operatorname{Sqrt}[b*\operatorname{Tanh}[c + d*x]]/\operatorname{Sqrt}[b]])/d - (2*b^3*\operatorname{Sqrt}[b*\operatorname{Tanh}[c + d*x]])/d - (2*b*(b*\operatorname{Tanh}[c + d*x])^{(5/2)})/(5*d)$

Rule 209

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[b, 2]))*\operatorname{rcTan}[\operatorname{Rt}[b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{PosQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{GtQ}[b, 0])$

Rule 212

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$

Rule 218

$\operatorname{Int}[(a_ + (b_)*(x_)^4)^{-1}, x_Symbol] \rightarrow \operatorname{With}\{r = \operatorname{Numerator}[\operatorname{Rt}[-a/b, 2]], s = \operatorname{Denominator}[\operatorname{Rt}[-a/b, 2]]\}, \operatorname{Dist}[r/(2*a), \operatorname{Int}[1/(r - s*x^2), x], x] + \operatorname{Dist}[r/(2*a), \operatorname{Int}[1/(r + s*x^2), x], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \ !\operatorname{GtQ}[a/b, 0]$

Rule 335

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
  Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
  ))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
  ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 3554

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d
  *x])^(n - 1)/(d*(n - 1))), x] - Dist[b^2, Int[(b*Tan[c + d*x])^(n - 2), x],
  x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]
```

Rule 3557

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[b/d, Subst[Int[
  x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !
  IntegerQ[n]
```

Rubi steps

$$\begin{aligned}
 \int (b \tanh(c + dx))^{7/2} dx &= -\frac{2b(b \tanh(c + dx))^{5/2}}{5d} + b^2 \int (b \tanh(c + dx))^{3/2} dx \\
 &= -\frac{2b^3 \sqrt{b \tanh(c + dx)}}{d} - \frac{2b(b \tanh(c + dx))^{5/2}}{5d} + b^4 \int \frac{1}{\sqrt{b \tanh(c + dx)}} dx \\
 &= -\frac{2b^3 \sqrt{b \tanh(c + dx)}}{d} - \frac{2b(b \tanh(c + dx))^{5/2}}{5d} - \frac{b^5 \text{Subst}\left(\int \frac{1}{\sqrt{x}(-b^2+x^2)} dx, x, \sqrt{b \tanh(c + dx)}\right)}{d} \\
 &= -\frac{2b^3 \sqrt{b \tanh(c + dx)}}{d} - \frac{2b(b \tanh(c + dx))^{5/2}}{5d} - \frac{(2b^5) \text{Subst}\left(\int \frac{1}{-b^2+x^4} dx, x, \sqrt{b \tanh(c + dx)}\right)}{d} \\
 &= -\frac{2b^3 \sqrt{b \tanh(c + dx)}}{d} - \frac{2b(b \tanh(c + dx))^{5/2}}{5d} + \frac{b^4 \text{Subst}\left(\int \frac{1}{b-x^2} dx, x, \sqrt{b \tanh(c + dx)}\right)}{d} \\
 &= \frac{b^{7/2} \tan^{-1}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{d} + \frac{b^{7/2} \tanh^{-1}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{d} - \frac{2b^3 \sqrt{b \tanh(c + dx)}}{d}
 \end{aligned}$$

Mathematica [A]

time = 0.17, size = 83, normalized size = 0.86

$$\frac{b^3 \sqrt{b \tanh(c + dx)} \left(5 \text{ArcTan}\left(\sqrt{\tanh(c + dx)}\right) + 5 \tanh^{-1}\left(\sqrt{\tanh(c + dx)}\right) - 10 \sqrt{\tanh(c + dx)} - 2 \tanh^{\frac{5}{2}}(c + dx)\right)}{5d \sqrt{\tanh(c + dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Tanh[c + d*x])^(7/2),x]

[Out] (b^3*Sqrt[b*Tanh[c + d*x]]*(5*ArcTan[Sqrt[Tanh[c + d*x]]] + 5*ArcTanh[Sqrt[Tanh[c + d*x]]] - 10*Sqrt[Tanh[c + d*x]] - 2*Tanh[c + d*x]^(5/2)))/(5*d*Sqrt[Tanh[c + d*x]])

Maple [A]

time = 1.95, size = 74, normalized size = 0.76

method	result
derivativedivides	$2b \left(\frac{(b \tanh(dx+c))^{\frac{5}{2}}}{5} + b^2 \sqrt{b \tanh(dx+c)} - \frac{b^{\frac{5}{2}} \arctan\left(\frac{\sqrt{b \tanh(dx+c)}}{\sqrt{b}}\right)}{2} - \frac{b^{\frac{5}{2}} \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(dx+c)}}{\sqrt{b}}\right)}{2} \right) \frac{1}{d}$
default	$2b \left(\frac{(b \tanh(dx+c))^{\frac{5}{2}}}{5} + b^2 \sqrt{b \tanh(dx+c)} - \frac{b^{\frac{5}{2}} \arctan\left(\frac{\sqrt{b \tanh(dx+c)}}{\sqrt{b}}\right)}{2} - \frac{b^{\frac{5}{2}} \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(dx+c)}}{\sqrt{b}}\right)}{2} \right) \frac{1}{d}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*tanh(d*x+c))^(7/2),x,method=_RETURNVERBOSE)

[Out] -2/d*b*(1/5*(b*tanh(d*x+c))^(5/2)+b^2*(b*tanh(d*x+c))^(1/2)-1/2*b^(5/2)*arctan((b*tanh(d*x+c))^(1/2)/b^(1/2))-1/2*b^(5/2)*arctanh((b*tanh(d*x+c))^(1/2)/b^(1/2)))/d

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*tanh(d*x+c))^(7/2),x, algorithm="maxima")

[Out] integrate((b*tanh(d*x + c))^(7/2), x)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 751 vs. 2(79) = 158.

time = 0.39, size = 1556, normalized size = 16.04

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*tanh(d*x+c))^(7/2),x, algorithm="fricas")

[Out]
$$\begin{aligned} & [-1/20*(10*(b^3*\cosh(d*x + c)^4 + 4*b^3*\cosh(d*x + c)*\sinh(d*x + c)^3 + b^3 \\ & * \sinh(d*x + c)^4 + 2*b^3*\cosh(d*x + c)^2 + b^3 + 2*(3*b^3*\cosh(d*x + c)^2 + \\ & b^3)*\sinh(d*x + c)^2 + 4*(b^3*\cosh(d*x + c)^3 + b^3*\cosh(d*x + c))*\sinh(d* \\ & x + c))*\sqrt{-b}*\arctan((\cosh(d*x + c)^2 + 2*\cosh(d*x + c)*\sinh(d*x + c) + \\ & \sinh(d*x + c)^2)*\sqrt{-b}*\sqrt{b*\sinh(d*x + c)/\cosh(d*x + c)})/(b*\cosh(d*x + \\ & c)^2 + 2*b*\cosh(d*x + c)*\sinh(d*x + c) + b*\sinh(d*x + c)^2 - b)) - 5*(b^3* \\ & \cosh(d*x + c)^4 + 4*b^3*\cosh(d*x + c)*\sinh(d*x + c)^3 + b^3*\sinh(d*x + c)^4 \\ & + 2*b^3*\cosh(d*x + c)^2 + b^3 + 2*(3*b^3*\cosh(d*x + c)^2 + b^3)*\sinh(d*x + \\ & c)^2 + 4*(b^3*\cosh(d*x + c)^3 + b^3*\cosh(d*x + c))*\sinh(d*x + c))*\sqrt{-b} \\ & * \log(-(b*\cosh(d*x + c)^4 + 4*b*\cosh(d*x + c)^3*\sinh(d*x + c) + 6*b*\cosh(d*x \\ & + c)^2*\sinh(d*x + c)^2 + 4*b*\cosh(d*x + c)*\sinh(d*x + c)^3 + b*\sinh(d*x + \\ & c)^4 + 2*(\cosh(d*x + c)^2 + 2*\cosh(d*x + c)*\sinh(d*x + c) + \sinh(d*x + c)^2 \\ & + 1)*\sqrt{-b}*\sqrt{b*\sinh(d*x + c)/\cosh(d*x + c)} - 2*b)/(\cosh(d*x + c)^4 \\ & + 4*\cosh(d*x + c)^3*\sinh(d*x + c) + 6*\cosh(d*x + c)^2*\sinh(d*x + c)^2 + 4*c \\ & \cosh(d*x + c)*\sinh(d*x + c)^3 + \sinh(d*x + c)^4)) + 16*(3*b^3*\cosh(d*x + c)^4 \\ & + 12*b^3*\cosh(d*x + c)*\sinh(d*x + c)^3 + 3*b^3*\sinh(d*x + c)^4 + 4*b^3*co \\ & sh(d*x + c)^2 + 3*b^3 + 2*(9*b^3*\cosh(d*x + c)^2 + 2*b^3)*\sinh(d*x + c)^2 + \\ & 4*(3*b^3*\cosh(d*x + c)^3 + 2*b^3*\cosh(d*x + c))*\sinh(d*x + c))*\sqrt{b*\sinh \\ & (d*x + c)/\cosh(d*x + c)})/(d*\cosh(d*x + c)^4 + 4*d*\cosh(d*x + c)*\sinh(d*x + \\ & c)^3 + d*\sinh(d*x + c)^4 + 2*d*\cosh(d*x + c)^2 + 2*(3*d*\cosh(d*x + c)^2 + \\ & d)*\sinh(d*x + c)^2 + 4*(d*\cosh(d*x + c)^3 + d*\cosh(d*x + c))*\sinh(d*x + c) \\ & + d), -1/20*(10*(b^3*\cosh(d*x + c)^4 + 4*b^3*\cosh(d*x + c)*\sinh(d*x + c)^3 \\ & + b^3*\sinh(d*x + c)^4 + 2*b^3*\cosh(d*x + c)^2 + b^3 + 2*(3*b^3*\cosh(d*x + c) \\ &)^2 + b^3)*\sinh(d*x + c)^2 + 4*(b^3*\cosh(d*x + c)^3 + b^3*\cosh(d*x + c))*\si \\ & nh(d*x + c))*\sqrt{b}*\arctan(\sqrt{b}*\sqrt{b*\sinh(d*x + c)/\cosh(d*x + c)})/(b* \\ & \cosh(d*x + c)^2 + 2*b*\cosh(d*x + c)*\sinh(d*x + c) + b*\sinh(d*x + c)^2 - b)) \\ & - 5*(b^3*\cosh(d*x + c)^4 + 4*b^3*\cosh(d*x + c)*\sinh(d*x + c)^3 + b^3*\sinh(\\ & d*x + c)^4 + 2*b^3*\cosh(d*x + c)^2 + b^3 + 2*(3*b^3*\cosh(d*x + c)^2 + b^3)* \\ & \sinh(d*x + c)^2 + 4*(b^3*\cosh(d*x + c)^3 + b^3*\cosh(d*x + c))*\sinh(d*x + c) \\ &)*\sqrt{b}*\log(2*b*\cosh(d*x + c)^4 + 8*b*\cosh(d*x + c)^3*\sinh(d*x + c) + 12* \\ & b*\cosh(d*x + c)^2*\sinh(d*x + c)^2 + 8*b*\cosh(d*x + c)*\sinh(d*x + c)^3 + 2*b \\ & *\sinh(d*x + c)^4 + 2*(\cosh(d*x + c)^4 + 4*\cosh(d*x + c)*\sinh(d*x + c)^3 + s \\ & inh(d*x + c)^4 + (6*\cosh(d*x + c)^2 + 1)*\sinh(d*x + c)^2 + \cosh(d*x + c)^2 \\ & + 2*(2*\cosh(d*x + c)^3 + \cosh(d*x + c))*\sinh(d*x + c))*\sqrt{b}*\sqrt{b*\sinh(\\ & d*x + c)/\cosh(d*x + c)} - b) + 16*(3*b^3*\cosh(d*x + c)^4 + 12*b^3*\cosh(d*x \\ & + c)*\sinh(d*x + c)^3 + 3*b^3*\sinh(d*x + c)^4 + 4*b^3*\cosh(d*x + c)^2 + 3*b^ \\ & 3 + 2*(9*b^3*\cosh(d*x + c)^2 + 2*b^3)*\sinh(d*x + c)^2 + 4*(3*b^3*\cosh(d*x + \\ & c)^3 + 2*b^3*\cosh(d*x + c))*\sinh(d*x + c))*\sqrt{b*\sinh(d*x + c)/\cosh(d*x + \\ & c)})/(d*\cosh(d*x + c)^4 + 4*d*\cosh(d*x + c)*\sinh(d*x + c)^3 + d*\sinh(d*x + \\ & c)^4 + 2*d*\cosh(d*x + c)^2 + 2*(3*d*\cosh(d*x + c)^2 + d)*\sinh(d*x + c)^2 + \\ & 4*(d*\cosh(d*x + c)^3 + d*\cosh(d*x + c))*\sinh(d*x + c) + d)] \end{aligned}$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (b \tanh(c + dx))^{\frac{7}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*tanh(d*x+c))**(7/2),x)

[Out] Integral((b*tanh(c + d*x))**(7/2), x)

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 293 vs. 2(79) = 158.

time = 0.53, size = 293, normalized size = 3.02

$$\frac{10b^{\frac{7}{2}} \arctan\left(\frac{-\sqrt{b} e^{2dx+2c} - \sqrt{b(4dx+4c)-b}}{\sqrt{b}}\right) - 5b^{\frac{7}{2}} \log\left(\frac{-\sqrt{b} e^{2dx+2c} + \sqrt{b(4dx+4c)-b}}{\sqrt{b}}\right) - \frac{16\left(\left(\sqrt{b} e^{2dx+2c} - \sqrt{b(4dx+4c)-b}\right)^{10} + 10\left(\sqrt{b} e^{2dx+2c} - \sqrt{b(4dx+4c)-b}\right)^8 + 20\left(\sqrt{b} e^{2dx+2c} - \sqrt{b(4dx+4c)-b}\right)^6 + 10\left(\sqrt{b} e^{2dx+2c} - \sqrt{b(4dx+4c)-b}\right)^4 + 3b^{\frac{7}{2}}\right)}{10d \left(\sqrt{b} e^{2dx+2c} - \sqrt{b(4dx+4c)-b} + \sqrt{b}\right)^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*tanh(d*x+c))^(7/2),x, algorithm="giac")

[Out] 1/10*(10*b^(7/2)*arctan(-sqrt(b)*e^(2*d*x + 2*c) - sqrt(b*e^(4*d*x + 4*c) - b))/sqrt(b) - 5*b^(7/2)*log(abs(-sqrt(b)*e^(2*d*x + 2*c) + sqrt(b*e^(4*d*x + 4*c) - b))) - 16*(5*(sqrt(b)*e^(2*d*x + 2*c) - sqrt(b*e^(4*d*x + 4*c) - b))^4*b^4 + 10*(sqrt(b)*e^(2*d*x + 2*c) - sqrt(b*e^(4*d*x + 4*c) - b))^3*b^(9/2) + 20*(sqrt(b)*e^(2*d*x + 2*c) - sqrt(b*e^(4*d*x + 4*c) - b))^2*b^5 + 10*(sqrt(b)*e^(2*d*x + 2*c) - sqrt(b*e^(4*d*x + 4*c) - b))*b^(11/2) + 3*b^6)/(sqrt(b)*e^(2*d*x + 2*c) - sqrt(b*e^(4*d*x + 4*c) - b) + sqrt(b))^5/d

Mupad [B]

time = 1.51, size = 83, normalized size = 0.86

$$\frac{b^{7/2} \operatorname{atan}\left(\frac{\sqrt{b} \tanh(c + dx)}{\sqrt{b}}\right)}{d} - \frac{2b^3 \sqrt{b} \tanh(c + dx)}{d} - \frac{2b(b \tanh(c + dx))^{5/2}}{5d} - \frac{b^{7/2} \operatorname{atan}\left(\frac{\sqrt{b} \tanh(c + dx)}{\sqrt{b}}\right) \operatorname{li}}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*tanh(c + d*x))^(7/2),x)

[Out] (b^(7/2)*atan((b*tanh(c + d*x))^(1/2)/b^(1/2)))/d - (2*b^3*(b*tanh(c + d*x))^(1/2))/d - (2*b*(b*tanh(c + d*x))^(5/2))/(5*d) - (b^(7/2)*atan((b*tanh(c + d*x))^(1/2)*1i)/b^(1/2))*1i)/d

3.14 $\int (b \tanh(c + dx))^{5/2} dx$

Optimal. Leaf size=78

$$\frac{b^{5/2} \operatorname{ArcTan}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{d} + \frac{b^{5/2} \tanh^{-1}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{d} - \frac{2b(b \tanh(c + dx))^{3/2}}{3d}$$

[Out] $-b^{(5/2)*\arctan((b*\tanh(d*x+c))^{(1/2)}/b^{(1/2)})/d+b^{(5/2)*\operatorname{arctanh}((b*\tanh(d*x+c))^{(1/2)}/b^{(1/2)})/d-2/3*b*(b*\tanh(d*x+c))^{(3/2)}/d}$

Rubi [A]

time = 0.04, antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3554, 3557, 335, 304, 209, 212}

$$\frac{b^{5/2} \operatorname{ArcTan}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{d} + \frac{b^{5/2} \tanh^{-1}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{d} - \frac{2b(b \tanh(c + dx))^{3/2}}{3d}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(b*\operatorname{Tanh}[c + d*x])^{(5/2)}, x]$

[Out] $-((b^{(5/2)*\operatorname{ArcTan}[\operatorname{Sqrt}[b*\operatorname{Tanh}[c + d*x]]/\operatorname{Sqrt}[b]])/d) + (b^{(5/2)*\operatorname{ArcTanh}[\operatorname{Sqrt}[b*\operatorname{Tanh}[c + d*x]]/\operatorname{Sqrt}[b]])/d - (2*b*(b*\operatorname{Tanh}[c + d*x])^{(3/2)})/(3*d)$

Rule 209

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[b, 2]))*\operatorname{ArcTan}[\operatorname{Rt}[b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{PosQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{GtQ}[b, 0])$

Rule 212

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$

Rule 304

$\operatorname{Int}[x^2/((a_ + (b_)*(x_)^4), x_Symbol] \rightarrow \operatorname{With}[\{r = \operatorname{Numerator}[\operatorname{Rt}[-a/b, 2]], s = \operatorname{Denominator}[\operatorname{Rt}[-a/b, 2]]\}, \operatorname{Dist}[s/(2*b), \operatorname{Int}[1/(r + s*x^2), x], x] - \operatorname{Dist}[s/(2*b), \operatorname{Int}[1/(r - s*x^2), x], x]] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \ !\operatorname{GtQ}[a/b, 0]$

Rule 335


```
Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
  Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
  )]^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 3554

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d
*x])^(n - 1)/(d*(n - 1))), x] - Dist[b^2, Int[(b*Tan[c + d*x])^(n - 2), x],
x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]
```

Rule 3557

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[b/d, Subst[Int[
x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !
IntegerQ[n]
```

Rubi steps

$$\begin{aligned}
\int (b \tanh(c + dx))^{5/2} dx &= -\frac{2b(b \tanh(c + dx))^{3/2}}{3d} + b^2 \int \sqrt{b \tanh(c + dx)} dx \\
&= -\frac{2b(b \tanh(c + dx))^{3/2}}{3d} - \frac{b^3 \text{Subst}\left(\int \frac{\sqrt{x}}{-b^2+x^2} dx, x, b \tanh(c + dx)\right)}{d} \\
&= -\frac{2b(b \tanh(c + dx))^{3/2}}{3d} - \frac{(2b^3) \text{Subst}\left(\int \frac{x^2}{-b^2+x^4} dx, x, \sqrt{b \tanh(c + dx)}\right)}{d} \\
&= -\frac{2b(b \tanh(c + dx))^{3/2}}{3d} + \frac{b^3 \text{Subst}\left(\int \frac{1}{b-x^2} dx, x, \sqrt{b \tanh(c + dx)}\right)}{d} - \frac{b^3 \text{Subst}\left(\int \frac{1}{b+x^2} dx, x, \sqrt{b \tanh(c + dx)}\right)}{d} \\
&= -\frac{b^{5/2} \tan^{-1}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{d} + \frac{b^{5/2} \tanh^{-1}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{d} - \frac{2b(b \tanh(c + dx))^{3/2}}{3d}
\end{aligned}$$

Mathematica [A]

time = 0.15, size = 68, normalized size = 0.87

$$\frac{(b \tanh(c + dx))^{5/2} \left(3 \text{ArcTan}\left(\sqrt{\tanh(c + dx)}\right) - 3 \tanh^{-1}\left(\sqrt{\tanh(c + dx)}\right) + 2 \tanh^{3/2}(c + dx) \right)}{3d \tanh^{5/2}(c + dx)}$$

Antiderivative was successfully verified.

```
[In] Integrate[(b*Tanh[c + d*x])^(5/2),x]
```

[Out] $-1/3*((b*\text{Tanh}[c + d*x])^{(5/2)}*(3*\text{ArcTan}[\text{Sqrt}[\text{Tanh}[c + d*x]]] - 3*\text{ArcTanh}[\text{Sqrt}[\text{Tanh}[c + d*x]]] + 2*\text{Tanh}[c + d*x]^{(3/2)}))/(d*\text{Tanh}[c + d*x]^{(5/2)})$

Maple [A]

time = 1.87, size = 60, normalized size = 0.77

method	result	size
derivativedivides	$2b \frac{\left(\frac{(b \tanh(dx+c))^{3/2}}{3} - \frac{b^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(dx+c)}}{\sqrt{b}}\right)}{2} + \frac{b^{3/2} \operatorname{arctan}\left(\frac{\sqrt{b \tanh(dx+c)}}{\sqrt{b}}\right)}{2} \right)}{d}$	60
default	$2b \frac{\left(\frac{(b \tanh(dx+c))^{3/2}}{3} - \frac{b^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(dx+c)}}{\sqrt{b}}\right)}{2} + \frac{b^{3/2} \operatorname{arctan}\left(\frac{\sqrt{b \tanh(dx+c)}}{\sqrt{b}}\right)}{2} \right)}{d}$	60

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*tanh(d*x+c))^(5/2),x,method=_RETURNVERBOSE)`

[Out] $-2/d*b*(1/3*(b*\text{tanh}(d*x+c))^{(3/2)}-1/2*b^{(3/2)}*\text{arctanh}((b*\text{tanh}(d*x+c))^{(1/2)}/b^{(1/2)})+1/2*b^{(3/2)}*\text{arctan}((b*\text{tanh}(d*x+c))^{(1/2)}/b^{(1/2)}))$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*tanh(d*x+c))^(5/2),x, algorithm="maxima")`

[Out] `integrate((b*tanh(d*x + c))^(5/2), x)`

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 463 vs. $2(62) = 124$.

time = 0.37, size = 980, normalized size = 12.56

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*tanh(d*x+c))^(5/2),x, algorithm="fricas")`

[Out] $[-1/12*(6*(b^2*\cosh(d*x + c))^2 + 2*b^2*\cosh(d*x + c)*\sinh(d*x + c) + b^2*\sinh(d*x + c)^2 + b^2)*\text{sqrt}(-b)*\text{arctan}((\cosh(d*x + c))^2 + 2*\cosh(d*x + c)*\sinh(d*x + c) + \sinh(d*x + c)^2)*\text{sqrt}(-b)*\text{sqrt}(b*\sinh(d*x + c)/\cosh(d*x + c))]/$

```
(b*cosh(d*x + c)^2 + 2*b*cosh(d*x + c)*sinh(d*x + c) + b*sinh(d*x + c)^2 -
b)) - 3*(b^2*cosh(d*x + c)^2 + 2*b^2*cosh(d*x + c)*sinh(d*x + c) + b^2*sinh
(d*x + c)^2 + b^2)*sqrt(-b)*log(-(b*cosh(d*x + c)^4 + 4*b*cosh(d*x + c)^3*s
inh(d*x + c) + 6*b*cosh(d*x + c)^2*sinh(d*x + c)^2 + 4*b*cosh(d*x + c)*sinh
(d*x + c)^3 + b*sinh(d*x + c)^4 - 2*(cosh(d*x + c)^2 + 2*cosh(d*x + c)*sinh
(d*x + c) + sinh(d*x + c)^2 + 1)*sqrt(-b)*sqrt(b*sinh(d*x + c)/cosh(d*x + c
)) - 2*b)/(cosh(d*x + c)^4 + 4*cosh(d*x + c)^3*sinh(d*x + c) + 6*cosh(d*x +
c)^2*sinh(d*x + c)^2 + 4*cosh(d*x + c)*sinh(d*x + c)^3 + sinh(d*x + c)^4))
+ 8*(b^2*cosh(d*x + c)^2 + 2*b^2*cosh(d*x + c)*sinh(d*x + c) + b^2*sinh(d*
x + c)^2 - b^2)*sqrt(b*sinh(d*x + c)/cosh(d*x + c)))/(d*cosh(d*x + c)^2 + 2
*d*cosh(d*x + c)*sinh(d*x + c) + d*sinh(d*x + c)^2 + d), 1/12*(6*(b^2*cosh(
d*x + c)^2 + 2*b^2*cosh(d*x + c)*sinh(d*x + c) + b^2*sinh(d*x + c)^2 + b^2)
*sqrt(b)*arctan(sqrt(b)*sqrt(b*sinh(d*x + c)/cosh(d*x + c)))/(b*cosh(d*x + c
)^2 + 2*b*cosh(d*x + c)*sinh(d*x + c) + b*sinh(d*x + c)^2 - b)) + 3*(b^2*co
sh(d*x + c)^2 + 2*b^2*cosh(d*x + c)*sinh(d*x + c) + b^2*sinh(d*x + c)^2 + b
^2)*sqrt(b)*log(2*b*cosh(d*x + c)^4 + 8*b*cosh(d*x + c)^3*sinh(d*x + c) + 1
2*b*cosh(d*x + c)^2*sinh(d*x + c)^2 + 8*b*cosh(d*x + c)*sinh(d*x + c)^3 + 2
*b*sinh(d*x + c)^4 + 2*(cosh(d*x + c)^4 + 4*cosh(d*x + c)*sinh(d*x + c)^3 +
sinh(d*x + c)^4 + (6*cosh(d*x + c)^2 + 1)*sinh(d*x + c)^2 + cosh(d*x + c)^
2 + 2*(2*cosh(d*x + c)^3 + cosh(d*x + c))*sinh(d*x + c))*sqrt(b)*sqrt(b*sin
h(d*x + c)/cosh(d*x + c)) - b) - 8*(b^2*cosh(d*x + c)^2 + 2*b^2*cosh(d*x +
c)*sinh(d*x + c) + b^2*sinh(d*x + c)^2 - b^2)*sqrt(b*sinh(d*x + c)/cosh(d*x
+ c)))/(d*cosh(d*x + c)^2 + 2*d*cosh(d*x + c)*sinh(d*x + c) + d*sinh(d*x +
c)^2 + d)]
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (b \tanh(c + dx))^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*tanh(d*x+c))**(5/2),x)

[Out] Integral((b*tanh(c + d*x))**(5/2), x)

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 173 vs. 2(62) = 124.

time = 0.51, size = 173, normalized size = 2.22

$$6b^{\frac{5}{2}} \arctan\left(-\frac{\sqrt{b} e^{(2dx+2c)} - \sqrt{be^{(4dx+4c)} - b}}{\sqrt{b}}\right) + 3b^{\frac{5}{2}} \log\left(\left|-\sqrt{b} e^{(2dx+2c)} + \sqrt{be^{(4dx+4c)} - b}\right|\right) + \frac{8\left(3\left(\sqrt{b} e^{(2dx+2c)} - \sqrt{be^{(4dx+4c)} - b}\right)^2 b^3 + b^4\right)}{\left(\sqrt{b} e^{(2dx+2c)} - \sqrt{be^{(4dx+4c)} - b} + \sqrt{b}\right)^3}$$

6d

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*tanh(d*x+c))^(5/2),x, algorithm="giac")

[Out]
$$\frac{-1/6*(6*b^{5/2}*\arctan(-(\sqrt{b})*e^{(2*d*x + 2*c)} - \sqrt{b*e^{(4*d*x + 4*c)} - b))/\sqrt{b}) + 3*b^{5/2}*\log(\text{abs}(-\sqrt{b})*e^{(2*d*x + 2*c)} + \sqrt{b*e^{(4*d*x + 4*c)} - b})) + 8*(3*(\sqrt{b})*e^{(2*d*x + 2*c)} - \sqrt{b*e^{(4*d*x + 4*c)} - b})^2*b^3 + b^4)/(\sqrt{b})*e^{(2*d*x + 2*c)} - \sqrt{b*e^{(4*d*x + 4*c)} - b} + \sqrt{b})^3/d$$

Mupad [B]

time = 1.25, size = 62, normalized size = 0.79

$$\frac{b^{5/2} \operatorname{atanh}\left(\frac{\sqrt{b} \tanh(c + dx)}{\sqrt{b}}\right)}{d} - \frac{b^{5/2} \operatorname{atan}\left(\frac{\sqrt{b} \tanh(c + dx)}{\sqrt{b}}\right)}{d} - \frac{2b(b \tanh(c + dx))^{3/2}}{3d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\operatorname{int}((b*\tanh(c + d*x))^{5/2}, x)$

[Out] $(b^{5/2}*\operatorname{atanh}((b*\tanh(c + d*x))^{1/2}/b^{1/2}))/d - (b^{5/2}*\operatorname{atan}((b*\tanh(c + d*x))^{1/2}/b^{1/2}))/d - (2*b*(b*\tanh(c + d*x))^{3/2})/(3*d)$

3.15 $\int (b \tanh(c + dx))^{3/2} dx$

Optimal. Leaf size=75

$$\frac{b^{3/2} \operatorname{ArcTan}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{d} + \frac{b^{3/2} \tanh^{-1}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{d} - \frac{2b\sqrt{b \tanh(c + dx)}}{d}$$

[Out] $b^{(3/2)*\arctan((b*\tanh(d*x+c))^{(1/2)}/b^{(1/2)})/d+b^{(3/2)*\operatorname{arctanh}((b*\tanh(d*x+c))^{(1/2)}/b^{(1/2)})/d-2*b*(b*\tanh(d*x+c))^{(1/2)}/d}$

Rubi [A]

time = 0.04, antiderivative size = 75, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3554, 3557, 335, 218, 212, 209}

$$\frac{b^{3/2} \operatorname{ArcTan}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{d} + \frac{b^{3/2} \tanh^{-1}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{d} - \frac{2b\sqrt{b \tanh(c + dx)}}{d}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(b*\operatorname{Tanh}[c + d*x])^{(3/2)}, x]$

[Out] $(b^{(3/2)*\operatorname{ArcTan}[\operatorname{Sqrt}[b*\operatorname{Tanh}[c + d*x]]/\operatorname{Sqrt}[b]])/d + (b^{(3/2)*\operatorname{ArcTanh}[\operatorname{Sqrt}[b*\operatorname{Tanh}[c + d*x]]/\operatorname{Sqrt}[b]])/d - (2*b*\operatorname{Sqrt}[b*\operatorname{Tanh}[c + d*x]])/d$

Rule 209

$\operatorname{Int}[(a_+ + (b_+)*(x_+)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[b, 2]))*\operatorname{ArcTan}[\operatorname{Rt}[b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{PosQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{GtQ}[b, 0])$

Rule 212

$\operatorname{Int}[(a_+ + (b_+)*(x_+)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$

Rule 218

$\operatorname{Int}[(a_+ + (b_+)*(x_+)^4)^{-1}, x_Symbol] \rightarrow \operatorname{With}\{r = \operatorname{Numerator}[\operatorname{Rt}[-a/b, 2]], s = \operatorname{Denominator}[\operatorname{Rt}[-a/b, 2]]\}, \operatorname{Dist}[r/(2*a), \operatorname{Int}[1/(r - s*x^2), x], x] + \operatorname{Dist}[r/(2*a), \operatorname{Int}[1/(r + s*x^2), x], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \ !\operatorname{GtQ}[a/b, 0]$

Rule 335

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
  Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 3554

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d
*x])^(n - 1)/(d*(n - 1))), x] - Dist[b^2, Int[(b*Tan[c + d*x])^(n - 2), x],
x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]
```

Rule 3557

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[b/d, Subst[Int[
x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !
IntegerQ[n]
```

Rubi steps

$$\begin{aligned}
\int (b \tanh(c + dx))^{3/2} dx &= -\frac{2b\sqrt{b \tanh(c + dx)}}{d} + b^2 \int \frac{1}{\sqrt{b \tanh(c + dx)}} dx \\
&= -\frac{2b\sqrt{b \tanh(c + dx)}}{d} - \frac{b^3 \text{Subst}\left(\int \frac{1}{\sqrt{x}(-b^2+x^2)} dx, x, b \tanh(c + dx)\right)}{d} \\
&= -\frac{2b\sqrt{b \tanh(c + dx)}}{d} - \frac{(2b^3) \text{Subst}\left(\int \frac{1}{-b^2+x^4} dx, x, \sqrt{b \tanh(c + dx)}\right)}{d} \\
&= -\frac{2b\sqrt{b \tanh(c + dx)}}{d} + \frac{b^2 \text{Subst}\left(\int \frac{1}{b-x^2} dx, x, \sqrt{b \tanh(c + dx)}\right)}{d} + \frac{b^2 \text{Subst}\left(\int \frac{1}{b+x^2} dx, x, \sqrt{b \tanh(c + dx)}\right)}{d} \\
&= \frac{b^{3/2} \tan^{-1}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{d} + \frac{b^{3/2} \tanh^{-1}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{d} - \frac{2b\sqrt{b \tanh(c + dx)}}{d}
\end{aligned}$$

Mathematica [A]

time = 0.06, size = 61, normalized size = 0.81

$$\frac{\left(\text{ArcTan}\left(\sqrt{\tanh(c + dx)}\right) + \tanh^{-1}\left(\sqrt{\tanh(c + dx)}\right) - 2\sqrt{\tanh(c + dx)}\right) (b \tanh(c + dx))^{3/2}}{d \tanh^{3/2}(c + dx)}$$

Antiderivative was successfully verified.

```
[In] Integrate[(b*Tanh[c + d*x])^(3/2), x]
```

[Out] ((ArcTan[Sqrt[Tanh[c + d*x]]] + ArcTanh[Sqrt[Tanh[c + d*x]]] - 2*Sqrt[Tanh[c + d*x]])*(b*Tanh[c + d*x])^(3/2))/(d*Tanh[c + d*x]^(3/2))

Maple [A]

time = 1.74, size = 58, normalized size = 0.77

method	result
derivativedivides	$\frac{2b \left(\sqrt{b \tanh(dx + c)} - \frac{\sqrt{b} \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(dx + c)}}{\sqrt{b}}\right)}{2} - \frac{\sqrt{b} \operatorname{arctan}\left(\frac{\sqrt{b \tanh(dx + c)}}{\sqrt{b}}\right)}{2} \right)}{d}$
default	$\frac{2b \left(\sqrt{b \tanh(dx + c)} - \frac{\sqrt{b} \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(dx + c)}}{\sqrt{b}}\right)}{2} - \frac{\sqrt{b} \operatorname{arctan}\left(\frac{\sqrt{b \tanh(dx + c)}}{\sqrt{b}}\right)}{2} \right)}{d}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*tanh(d*x+c))^(3/2),x,method=_RETURNVERBOSE)

[Out] -2/d*b*((b*tanh(d*x+c))^(1/2)-1/2*b^(1/2)*arctanh((b*tanh(d*x+c))^(1/2)/b^(1/2))-1/2*b^(1/2)*arctan((b*tanh(d*x+c))^(1/2)/b^(1/2)))

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate


Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*tanh(d*x+c))^(3/2),x, algorithm="maxima")

[Out] integrate((b*tanh(d*x + c))^(3/2), x)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 291 vs. 2(61) = 122.

time = 0.51, size = 638, normalized size = 8.51



Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*tanh(d*x+c))^(3/2),x, algorithm="fricas")

[Out] [-1/4*(2*sqrt(-b)*b*arctan((cosh(d*x + c))^2 + 2*cosh(d*x + c)*sinh(d*x + c) + sinh(d*x + c)^2)*sqrt(-b)*sqrt(b*sinh(d*x + c)/cosh(d*x + c)))/(b*cosh(d*

$x + c)^2 + 2*b*cosh(d*x + c)*sinh(d*x + c) + b*sinh(d*x + c)^2 - b)) - sqrt$
 $(-b)*b*log(-(b*cosh(d*x + c)^4 + 4*b*cosh(d*x + c)^3*sinh(d*x + c) + 6*b*cosh$
 $sh(d*x + c)^2*sinh(d*x + c)^2 + 4*b*cosh(d*x + c)*sinh(d*x + c)^3 + b*sinh$
 $d*x + c)^4 + 2*(cosh(d*x + c)^2 + 2*cosh(d*x + c)*sinh(d*x + c) + sinh(d*x$
 $+ c)^2 + 1)*sqrt(-b)*sqrt(b*sinh(d*x + c)/cosh(d*x + c)) - 2*b)/(cosh(d*x +$
 $c)^4 + 4*cosh(d*x + c)^3*sinh(d*x + c) + 6*cosh(d*x + c)^2*sinh(d*x + c)^2$
 $+ 4*cosh(d*x + c)*sinh(d*x + c)^3 + sinh(d*x + c)^4)) + 8*b*sqrt(b*sinh(d*$
 $x + c)/cosh(d*x + c))/d, -1/4*(2*b^(3/2)*arctan(sqrt(b)*sqrt(b*sinh(d*x +$
 $c)/cosh(d*x + c))/(b*cosh(d*x + c)^2 + 2*b*cosh(d*x + c)*sinh(d*x + c) + b*$
 $sinh(d*x + c)^2 - b)) - b^(3/2)*log(2*b*cosh(d*x + c)^4 + 8*b*cosh(d*x + c)$
 $^3*sinh(d*x + c) + 12*b*cosh(d*x + c)^2*sinh(d*x + c)^2 + 8*b*cosh(d*x + c)$
 $*sinh(d*x + c)^3 + 2*b*sinh(d*x + c)^4 + 2*(cosh(d*x + c)^4 + 4*cosh(d*x +$
 $c)*sinh(d*x + c)^3 + sinh(d*x + c)^4 + (6*cosh(d*x + c)^2 + 1)*sinh(d*x +$
 $c)^2 + cosh(d*x + c)^2 + 2*(2*cosh(d*x + c)^3 + cosh(d*x + c))*sinh(d*x + c)$
 $)*sqrt(b)*sqrt(b*sinh(d*x + c)/cosh(d*x + c)) - b) + 8*b*sqrt(b*sinh(d*x +$
 $c)/cosh(d*x + c))/d]$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (b \tanh(c + dx))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*tanh(d*x+c))**(3/2),x)

[Out] Integral((b*tanh(c + d*x))**(3/2), x)

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 131 vs. 2(61) = 122.

time = 0.45, size = 131, normalized size = 1.75

$$\frac{\left(2\sqrt{b} \arctan\left(-\frac{\sqrt{b} e^{(2dx+2c)} - \sqrt{be^{(4dx+4c)} - b}}{\sqrt{b}}\right) - \sqrt{b} \log\left(\left|-\sqrt{b} e^{(2dx+2c)} + \sqrt{be^{(4dx+4c)} - b}\right|\right) - \frac{8b}{\sqrt{b} e^{(2dx+2c)} - \sqrt{be^{(4dx+4c)} - b} + \sqrt{b}}\right)b}{2d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*tanh(d*x+c))^(3/2),x, algorithm="giac")

[Out] 1/2*(2*sqrt(b)*arctan(-sqrt(b)*e^(2*d*x + 2*c) - sqrt(b*e^(4*d*x + 4*c) - b))/sqrt(b) - sqrt(b)*log(abs(-sqrt(b)*e^(2*d*x + 2*c) + sqrt(b*e^(4*d*x + 4*c) - b))) - 8*b/(sqrt(b)*e^(2*d*x + 2*c) - sqrt(b*e^(4*d*x + 4*c) - b) + sqrt(b))*b/d

Mupad [B]

time = 1.16, size = 61, normalized size = 0.81

$$\frac{b^{3/2} \operatorname{atan}\left(\frac{\sqrt{b} \tanh(c + dx)}{\sqrt{b}}\right)}{d} - \frac{2b \sqrt{b} \tanh(c + dx)}{d} + \frac{b^{3/2} \operatorname{atanh}\left(\frac{\sqrt{b} \tanh(c + dx)}{\sqrt{b}}\right)}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*tanh(c + d*x))^(3/2),x)`

[Out] $(b^{3/2}*\operatorname{atan}((b*\tanh(c + d*x))^{1/2}/b^{1/2}))/d - (2*b*(b*\tanh(c + d*x))^{1/2})/d + (b^{3/2}*\operatorname{atanh}((b*\tanh(c + d*x))^{1/2}/b^{1/2}))/d$

3.16 $\int \sqrt{b \tanh(c + dx)} dx$

Optimal. Leaf size=58

$$-\frac{\sqrt{b} \operatorname{ArcTan}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{d} + \frac{\sqrt{b} \tanh^{-1}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{d}$$

[Out] $-\arctan((b \cdot \tanh(d \cdot x + c))^{1/2} / b^{1/2}) \cdot b^{1/2} / d + \operatorname{arctanh}((b \cdot \tanh(d \cdot x + c))^{1/2} / b^{1/2}) \cdot b^{1/2} / d$

Rubi [A]

time = 0.02, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {3557, 335, 304, 209, 212}

$$\frac{\sqrt{b} \tanh^{-1}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{d} - \frac{\sqrt{b} \operatorname{ArcTan}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{d}$$

Antiderivative was successfully verified.

[In] `Int[Sqrt[b*Tanh[c + d*x]],x]`

[Out] $-\left(\frac{\operatorname{Sqrt}[b] \operatorname{ArcTan}\left[\frac{\operatorname{Sqrt}[b \operatorname{Tanh}[c + d \cdot x]]}{\operatorname{Sqrt}[b]}\right]}{d}\right) + \left(\frac{\operatorname{Sqrt}[b] \operatorname{ArcTanh}\left[\frac{\operatorname{Sqrt}[b \operatorname{Tanh}[c + d \cdot x]]}{\operatorname{Sqrt}[b]}\right]}{d}\right)$

Rule 209

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

Rule 212

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rule 304

`Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

Rule 335

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
  Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
  ))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 3557

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[b/d, Subst[Int[
x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !
IntegerQ[n]
```

Rubi steps

$$\begin{aligned}
 \int \sqrt{b \tanh(c + dx)} \, dx &= -\frac{b \operatorname{Subst}\left(\int \frac{\sqrt{x}}{-b^2+x^2} \, dx, x, b \tanh(c + dx)\right)}{d} \\
 &= -\frac{(2b) \operatorname{Subst}\left(\int \frac{x^2}{-b^2+x^4} \, dx, x, \sqrt{b \tanh(c + dx)}\right)}{d} \\
 &= \frac{b \operatorname{Subst}\left(\int \frac{1}{b-x^2} \, dx, x, \sqrt{b \tanh(c + dx)}\right)}{d} - \frac{b \operatorname{Subst}\left(\int \frac{1}{b+x^2} \, dx, x, \sqrt{b \tanh(c + dx)}\right)}{d} \\
 &= -\frac{\sqrt{b} \tan^{-1}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{d} + \frac{\sqrt{b} \tanh^{-1}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{d}
 \end{aligned}$$

Mathematica [A]

time = 0.03, size = 51, normalized size = 0.88

$$\frac{\left(-\operatorname{ArcTan}\left(\sqrt{\tanh(c + dx)}\right) + \tanh^{-1}\left(\sqrt{\tanh(c + dx)}\right)\right) \sqrt{b \tanh(c + dx)}}{d \sqrt{\tanh(c + dx)}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sqrt[b*Tanh[c + d*x]], x]
```

```
[Out] ((-ArcTan[Sqrt[Tanh[c + d*x]]] + ArcTanh[Sqrt[Tanh[c + d*x]]])*Sqrt[b*Tanh[
c + d*x]])/(d*Sqrt[Tanh[c + d*x]])
```

Maple [A]

time = 2.09, size = 48, normalized size = 0.83

method	result	size
--------	--------	------

derivativedivides	$2b \frac{\left(\frac{\arctan\left(\frac{\sqrt{b} \tanh(dx+c)}{\sqrt{b}}\right)}{2\sqrt{b}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{b} \tanh(dx+c)}{\sqrt{b}}\right)}{2\sqrt{b}} \right)}{d}$	48
default	$2b \frac{\left(\frac{\arctan\left(\frac{\sqrt{b} \tanh(dx+c)}{\sqrt{b}}\right)}{2\sqrt{b}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{b} \tanh(dx+c)}{\sqrt{b}}\right)}{2\sqrt{b}} \right)}{d}$	48

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((b*tanh(d*x+c))^(1/2),x,method=_RETURNVERBOSE)
```

```
[Out] -2/d*b*(1/2/b^(1/2)*arctan((b*tanh(d*x+c))^(1/2)/b^(1/2))-1/2/b^(1/2)*arctanh((b*tanh(d*x+c))^(1/2)/b^(1/2)))
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*tanh(d*x+c))^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(sqrt(b*tanh(d*x + c)), x)
```

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 269 vs. 2(46) = 92.

time = 0.41, size = 593, normalized size = 10.22

$$\frac{\sqrt{-b} \operatorname{arctan}\left(\frac{\sqrt{b} \tanh(dx+c)}{\sqrt{b}}\right) - \sqrt{-b} \operatorname{arctanh}\left(\frac{\sqrt{b} \tanh(dx+c)}{\sqrt{b}}\right)}{2\sqrt{b}} - \frac{\sqrt{-b} \operatorname{arctan}\left(\frac{\sqrt{b} \tanh(dx+c)}{\sqrt{b}}\right) - \sqrt{-b} \operatorname{arctanh}\left(\frac{\sqrt{b} \tanh(dx+c)}{\sqrt{b}}\right)}{2\sqrt{b}}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*tanh(d*x+c))^(1/2),x, algorithm="fricas")
```

```
[Out] [-1/4*(2*sqrt(-b)*arctan((cosh(d*x + c))^2 + 2*cosh(d*x + c)*sinh(d*x + c) + sinh(d*x + c)^2)*sqrt(-b)*sqrt(b*sinh(d*x + c)/cosh(d*x + c)))/(b*cosh(d*x + c)^2 + 2*b*cosh(d*x + c)*sinh(d*x + c) + b*sinh(d*x + c)^2 - b) - sqrt(-b)*log(-(b*cosh(d*x + c)^4 + 4*b*cosh(d*x + c)^3*sinh(d*x + c) + 6*b*cosh(d*x + c)^2*sinh(d*x + c)^2 + 4*b*cosh(d*x + c)*sinh(d*x + c)^3 + b*sinh(d*x + c)^4 - 2*(cosh(d*x + c)^2 + 2*cosh(d*x + c)*sinh(d*x + c) + sinh(d*x + c)^2 + 1)*sqrt(-b)*sqrt(b*sinh(d*x + c)/cosh(d*x + c)) - 2*b)/(cosh(d*x + c)^4 + 4*cosh(d*x + c)^3*sinh(d*x + c) + 6*cosh(d*x + c)^2*sinh(d*x + c)^2 + 4
```

*cosh(d*x + c)*sinh(d*x + c)^3 + sinh(d*x + c)^4))/d, 1/4*(2*sqrt(b)*arctan(sqrt(b)*sqrt(b*sinh(d*x + c)/cosh(d*x + c))/(b*cosh(d*x + c)^2 + 2*b*cosh(d*x + c)*sinh(d*x + c) + b*sinh(d*x + c)^2 - b)) + sqrt(b)*log(2*b*cosh(d*x + c)^4 + 8*b*cosh(d*x + c)^3*sinh(d*x + c) + 12*b*cosh(d*x + c)^2*sinh(d*x + c)^2 + 8*b*cosh(d*x + c)*sinh(d*x + c)^3 + 2*b*sinh(d*x + c)^4 + 2*(cosh(d*x + c)^4 + 4*cosh(d*x + c)*sinh(d*x + c)^3 + sinh(d*x + c)^4 + (6*cosh(d*x + c)^2 + 1)*sinh(d*x + c)^2 + cosh(d*x + c)^2 + 2*(2*cosh(d*x + c)^3 + cosh(d*x + c))*sinh(d*x + c))*sqrt(b)*sqrt(b*sinh(d*x + c)/cosh(d*x + c)) - b))/d]

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \tanh(c + dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*tanh(d*x+c))**(1/2),x)

[Out] Integral(sqrt(b*tanh(c + d*x)), x)

Giac [A]

time = 0.44, size = 88, normalized size = 1.52

$$\frac{2\sqrt{b} \arctan\left(\frac{-\sqrt{b}e^{(2dx+2c)} - \sqrt{be^{(4dx+4c)} - b}}{\sqrt{b}}\right) + \sqrt{b} \log\left(\left|-\sqrt{b}e^{(2dx+2c)} + \sqrt{be^{(4dx+4c)} - b}\right|\right)}{2d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*tanh(d*x+c))^(1/2),x, algorithm="giac")

[Out] -1/2*(2*sqrt(b)*arctan(-(sqrt(b)*e^(2*d*x + 2*c) - sqrt(b*e^(4*d*x + 4*c) - b))/sqrt(b)) + sqrt(b)*log(abs(-sqrt(b)*e^(2*d*x + 2*c) + sqrt(b*e^(4*d*x + 4*c) - b))))/d

Mupad [B]

time = 1.08, size = 41, normalized size = 0.71

$$\frac{\sqrt{b} \left(\operatorname{atan}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right) - \operatorname{atanh}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right) \right)}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*tanh(c + d*x))^(1/2),x)

[Out] -(b^(1/2)*(atan((b*tanh(c + d*x))^(1/2)/b^(1/2)) - atanh((b*tanh(c + d*x))^(1/2)/b^(1/2))))/d

$$3.17 \quad \int \frac{1}{\sqrt{b \tanh(c + dx)}} dx$$

Optimal. Leaf size=57

$$\frac{\text{ArcTan}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{\sqrt{b} d} + \frac{\tanh^{-1}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{\sqrt{b} d}$$

[Out] arctan((b*tanh(d*x+c))^(1/2)/b^(1/2))/d/b^(1/2)+arctanh((b*tanh(d*x+c))^(1/2)/b^(1/2))/d/b^(1/2)

Rubi [A]

time = 0.02, antiderivative size = 57, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {3557, 335, 218, 212, 209}

$$\frac{\text{ArcTan}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{\sqrt{b} d} + \frac{\tanh^{-1}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{\sqrt{b} d}$$

Antiderivative was successfully verified.

[In] Int[1/Sqrt[b*Tanh[c + d*x]],x]

[Out] ArcTan[Sqrt[b*Tanh[c + d*x]]/Sqrt[b]]/(Sqrt[b]*d) + ArcTanh[Sqrt[b*Tanh[c + d*x]]/Sqrt[b]]/(Sqrt[b]*d)

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 218

Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 335

Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && FractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 3557

Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[b/d, Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && IntegerQ[n]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{b \tanh(c + dx)}} dx &= -\frac{b \operatorname{Subst}\left(\int \frac{1}{\sqrt{x(-b^2+x^2)}} dx, x, b \tanh(c + dx)\right)}{d} \\ &= -\frac{(2b) \operatorname{Subst}\left(\int \frac{1}{-b^2+x^4} dx, x, \sqrt{b \tanh(c + dx)}\right)}{d} \\ &= \frac{\operatorname{Subst}\left(\int \frac{1}{b-x^2} dx, x, \sqrt{b \tanh(c + dx)}\right)}{d} + \frac{\operatorname{Subst}\left(\int \frac{1}{b+x^2} dx, x, \sqrt{b \tanh(c + dx)}\right)}{d} \\ &= \frac{\tan^{-1}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{\sqrt{b} d} + \frac{\tanh^{-1}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{\sqrt{b} d} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 49, normalized size = 0.86

$$\frac{\left(\operatorname{ArcTan}\left(\sqrt{\tanh(c + dx)}\right) + \tanh^{-1}\left(\sqrt{\tanh(c + dx)}\right)\right) \sqrt{\tanh(c + dx)}}{d \sqrt{b \tanh(c + dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[1/Sqrt[b*Tanh[c + d*x]], x]

[Out] ((ArcTan[Sqrt[Tanh[c + d*x]]] + ArcTanh[Sqrt[Tanh[c + d*x]]])*Sqrt[Tanh[c + d*x]])/(d*Sqrt[b*Tanh[c + d*x]])

Maple [A]

time = 2.09, size = 48, normalized size = 0.84

$4 + 4*\cosh(d*x + c)^3*\sinh(d*x + c) + 6*\cosh(d*x + c)^2*\sinh(d*x + c)^2 + 4$
 $*\cosh(d*x + c)*\sinh(d*x + c)^3 + \sinh(d*x + c)^4)))/(b*d), -1/4*(2*\sqrt{b}*$
 $\arctan(\sqrt{b}*\sqrt{b*\sinh(d*x + c)/\cosh(d*x + c)})/(b*\cosh(d*x + c)^2 + 2*b$
 $*\cosh(d*x + c)*\sinh(d*x + c) + b*\sinh(d*x + c)^2 - b)) - \sqrt{b}*\log(2*b*\co$
 $sh(d*x + c)^4 + 8*b*\cosh(d*x + c)^3*\sinh(d*x + c) + 12*b*\cosh(d*x + c)^2*si$
 $nh(d*x + c)^2 + 8*b*\cosh(d*x + c)*\sinh(d*x + c)^3 + 2*b*\sinh(d*x + c)^4 + 2$
 $*(\cosh(d*x + c)^4 + 4*\cosh(d*x + c)*\sinh(d*x + c)^3 + \sinh(d*x + c)^4 + (6*$
 $\cosh(d*x + c)^2 + 1)*\sinh(d*x + c)^2 + \cosh(d*x + c)^2 + 2*(2*\cosh(d*x + c)$
 $^3 + \cosh(d*x + c))*\sinh(d*x + c))*\sqrt{b}*\sqrt{b*\sinh(d*x + c)/\cosh(d*x +$
 $c)) - b))/(b*d)]$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{b \tanh(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*tanh(d*x+c))**(1/2),x)

[Out] Integral(1/sqrt(b*tanh(c + d*x)), x)

Giac [A]

time = 0.47, size = 89, normalized size = 1.56

$$\frac{2 \arctan\left(\frac{-\sqrt{b} e^{(2dx+2c)} - \sqrt{be^{(4dx+4c)} - b}}{\sqrt{b}}\right)}{\sqrt{b}} - \frac{\log\left(\left|-\sqrt{b} e^{(2dx+2c)} + \sqrt{be^{(4dx+4c)} - b}\right|\right)}{\sqrt{b}}}{2d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*tanh(d*x+c))^(1/2),x, algorithm="giac")

[Out] 1/2*(2*arctan(-sqrt(b)*e^(2*d*x + 2*c) - sqrt(b*e^(4*d*x + 4*c) - b))/sqrt(b))/sqrt(b) - log(abs(-sqrt(b)*e^(2*d*x + 2*c) + sqrt(b*e^(4*d*x + 4*c) - b)))/sqrt(b))/d

Mupad [B]

time = 1.12, size = 38, normalized size = 0.67

$$\frac{\operatorname{atan}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right) + \operatorname{atanh}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{\sqrt{b} d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*tanh(c + d*x))^(1/2),x)

[Out] (atan((b*tanh(c + d*x))^(1/2)/b^(1/2)) + atanh((b*tanh(c + d*x))^(1/2)/b^(1/2)))/(b^(1/2)*d)

$$3.18 \quad \int \frac{1}{(b \tanh(c+dx))^{3/2}} dx$$

Optimal. Leaf size=78

$$-\frac{\operatorname{ArcTan}\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right)}{b^{3/2}d} + \frac{\tanh^{-1}\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right)}{b^{3/2}d} - \frac{2}{bd\sqrt{b \tanh(c+dx)}}$$

[Out] $-\arctan((b*\tanh(d*x+c))^{(1/2)}/b^{(1/2)})/b^{(3/2)}/d+\operatorname{arctanh}((b*\tanh(d*x+c))^{(1/2)}/b^{(1/2)})/b^{(3/2)}/d-2/b/d/(b*\tanh(d*x+c))^{(1/2)}$

Rubi [A]

time = 0.04, antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3555, 3557, 335, 304, 209, 212}

$$-\frac{\operatorname{ArcTan}\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right)}{b^{3/2}d} + \frac{\tanh^{-1}\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right)}{b^{3/2}d} - \frac{2}{bd\sqrt{b \tanh(c+dx)}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(b*\operatorname{Tanh}[c + d*x])^{(-3/2)}, x]$

[Out] $-(\operatorname{ArcTan}[\operatorname{Sqrt}[b*\operatorname{Tanh}[c + d*x]]/\operatorname{Sqrt}[b]]/(b^{(3/2)*d})) + \operatorname{ArcTanh}[\operatorname{Sqrt}[b*\operatorname{Tanh}[c + d*x]]/\operatorname{Sqrt}[b]]/(b^{(3/2)*d}) - 2/(b*d*\operatorname{Sqrt}[b*\operatorname{Tanh}[c + d*x]])$

Rule 209

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[b, 2]))*\operatorname{ArcTan}[\operatorname{Rt}[b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{PosQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{GtQ}[b, 0])$

Rule 212

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$

Rule 304

$\operatorname{Int}(x_)^2/((a_ + (b_)*(x_)^4), x_Symbol] \rightarrow \operatorname{With}\{r = \operatorname{Numerator}[\operatorname{Rt}[-a/b, 2]], s = \operatorname{Denominator}[\operatorname{Rt}[-a/b, 2]]\}, \operatorname{Dist}[s/(2*b), \operatorname{Int}[1/(r + s*x^2), x], x] - \operatorname{Dist}[s/(2*b), \operatorname{Int}[1/(r - s*x^2), x], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \ !\operatorname{GtQ}[a/b, 0]$

Rule 335

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
  Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
  )^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 3555

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(b*Tan[c + d*x]
)^n + 1)/(b*d*(n + 1)), x] - Dist[1/b^2, Int[(b*Tan[c + d*x])^(n + 2), x],
x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1]
```

Rule 3557

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[b/d, Subst[Int[
x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !
IntegerQ[n]
```

Rubi steps

$$\begin{aligned}
\int \frac{1}{(b \tanh(c + dx))^{3/2}} dx &= -\frac{2}{bd \sqrt{b \tanh(c + dx)}} + \frac{\int \sqrt{b \tanh(c + dx)} dx}{b^2} \\
&= -\frac{2}{bd \sqrt{b \tanh(c + dx)}} - \frac{\text{Subst}\left(\int \frac{\sqrt{x}}{-b^2 + x^2} dx, x, b \tanh(c + dx)\right)}{bd} \\
&= -\frac{2}{bd \sqrt{b \tanh(c + dx)}} - \frac{2 \text{Subst}\left(\int \frac{x^2}{-b^2 + x^4} dx, x, \sqrt{b \tanh(c + dx)}\right)}{bd} \\
&= -\frac{2}{bd \sqrt{b \tanh(c + dx)}} + \frac{\text{Subst}\left(\int \frac{1}{b-x^2} dx, x, \sqrt{b \tanh(c + dx)}\right)}{bd} - \frac{\text{Subst}\left(\int \frac{1}{b+x^2} dx, x, \sqrt{b \tanh(c + dx)}\right)}{bd} \\
&= -\frac{\tan^{-1}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{b^{3/2}d} + \frac{\tanh^{-1}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{b^{3/2}d} - \frac{2}{bd \sqrt{b \tanh(c + dx)}}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.06, size = 36, normalized size = 0.46

$$-\frac{{}_2F_1\left(-\frac{1}{4}, 1; \frac{3}{4}; \tanh^2(c + dx)\right)}{bd \sqrt{b \tanh(c + dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Tanh[c + d*x])^(-3/2),x]

[Out] (-2*Hypergeometric2F1[-1/4, 1, 3/4, Tanh[c + d*x]^2])/(b*d*Sqrt[b*Tanh[c + d*x]])

Maple [A]

time = 1.92, size = 62, normalized size = 0.79

method	result	size
derivativedivides	$2b \left(\frac{\operatorname{arctanh}\left(\frac{\sqrt{b \tanh(dx+c)}}{\sqrt{b}}\right)}{2b^{\frac{5}{2}}} + \frac{\operatorname{arctan}\left(\frac{\sqrt{b \tanh(dx+c)}}{\sqrt{b}}\right)}{2b^{\frac{5}{2}}} + \frac{1}{b^2 \sqrt{b \tanh(dx+c)}} \right)$	62
default	$2b \left(\frac{\operatorname{arctanh}\left(\frac{\sqrt{b \tanh(dx+c)}}{\sqrt{b}}\right)}{2b^{\frac{5}{2}}} + \frac{\operatorname{arctan}\left(\frac{\sqrt{b \tanh(dx+c)}}{\sqrt{b}}\right)}{2b^{\frac{5}{2}}} + \frac{1}{b^2 \sqrt{b \tanh(dx+c)}} \right)$	62

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*tanh(d*x+c))^(3/2),x,method=_RETURNVERBOSE)

[Out] -2/d*b*(-1/2/b^(5/2)*arctanh((b*tanh(d*x+c))^(1/2)/b^(1/2))+1/2/b^(5/2)*arctan((b*tanh(d*x+c))^(1/2)/b^(1/2))+1/b^2/(b*tanh(d*x+c))^(1/2))

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*tanh(d*x+c))^(3/2),x, algorithm="maxima")

[Out] integrate((b*tanh(d*x + c))^(3/2), x)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 435 vs. 2(64) = 128.

time = 0.44, size = 924, normalized size = 11.85

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*tanh(d*x+c))^(3/2),x, algorithm="fricas")

```
[Out] [-1/4*(2*(cosh(d*x + c)^2 + 2*cosh(d*x + c)*sinh(d*x + c) + sinh(d*x + c)^2 - 1)*sqrt(-b)*arctan((cosh(d*x + c)^2 + 2*cosh(d*x + c)*sinh(d*x + c) + sinh(d*x + c)^2)*sqrt(-b)*sqrt(b*sinh(d*x + c)/cosh(d*x + c)))/(b*cosh(d*x + c)^2 + 2*b*cosh(d*x + c)*sinh(d*x + c) + b*sinh(d*x + c)^2 - b)) + (cosh(d*x + c)^2 + 2*cosh(d*x + c)*sinh(d*x + c) + sinh(d*x + c)^2 - 1)*sqrt(-b)*log(-(b*cosh(d*x + c)^4 + 4*b*cosh(d*x + c)^3*sinh(d*x + c) + 6*b*cosh(d*x + c)^2*sinh(d*x + c)^2 + 4*b*cosh(d*x + c)*sinh(d*x + c)^3 + b*sinh(d*x + c)^4 + 2*(cosh(d*x + c)^2 + 2*cosh(d*x + c)*sinh(d*x + c) + sinh(d*x + c)^2 + 1)*sqrt(-b)*sqrt(b*sinh(d*x + c)/cosh(d*x + c)) - 2*b)/(cosh(d*x + c)^4 + 4*cosh(d*x + c)^3*sinh(d*x + c) + 6*cosh(d*x + c)^2*sinh(d*x + c)^2 + 4*cosh(d*x + c)*sinh(d*x + c)^3 + sinh(d*x + c)^4)) + 8*(cosh(d*x + c)^2 + 2*cosh(d*x + c)*sinh(d*x + c) + sinh(d*x + c)^2 + 1)*sqrt(b*sinh(d*x + c)/cosh(d*x + c)))/(b^2*d*cosh(d*x + c)^2 + 2*b^2*d*cosh(d*x + c)*sinh(d*x + c) + b^2*d*sinh(d*x + c)^2 - b^2*d), 1/4*(2*(cosh(d*x + c)^2 + 2*cosh(d*x + c)*sinh(d*x + c) + sinh(d*x + c)^2 - 1)*sqrt(b)*arctan(sqrt(b)*sqrt(b*sinh(d*x + c)/cosh(d*x + c)))/(b*cosh(d*x + c)^2 + 2*b*cosh(d*x + c)*sinh(d*x + c) + b*sinh(d*x + c)^2 - b)) + (cosh(d*x + c)^2 + 2*cosh(d*x + c)*sinh(d*x + c) + sinh(d*x + c)^2 - 1)*sqrt(b)*log(2*b*cosh(d*x + c)^4 + 8*b*cosh(d*x + c)^3*sinh(d*x + c) + 12*b*cosh(d*x + c)^2*sinh(d*x + c)^2 + 8*b*cosh(d*x + c)*sinh(d*x + c)^3 + 2*b*sinh(d*x + c)^4 + 2*(cosh(d*x + c)^4 + 4*cosh(d*x + c)*sinh(d*x + c)^3 + sinh(d*x + c)^4 + (6*cosh(d*x + c)^2 + 1)*sinh(d*x + c)^2 + cosh(d*x + c)^2 + 2*(2*cosh(d*x + c)^3 + cosh(d*x + c))*sinh(d*x + c))*sqrt(b)*sqrt(b*sinh(d*x + c)/cosh(d*x + c)) - b) - 8*(cosh(d*x + c)^2 + 2*cosh(d*x + c)*sinh(d*x + c) + sinh(d*x + c)^2 + 1)*sqrt(b*sinh(d*x + c)/cosh(d*x + c)))/(b^2*d*cosh(d*x + c)^2 + 2*b^2*d*cosh(d*x + c)*sinh(d*x + c) + b^2*d*sinh(d*x + c)^2 - b^2*d)]
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \tanh(c + dx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(b*tanh(d*x+c))**(3/2), x)
```

```
[Out] Integral((b*tanh(c + d*x))**(-3/2), x)
```

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 133 vs. 2(64) = 128.

time = 0.46, size = 133, normalized size = 1.71

$$\frac{2 \arctan\left(\frac{-\sqrt{b} e^{(2 dx+2 c)} - \sqrt{b e^{(4 dx+4 c)} - b}}{\sqrt{b}}\right)}{\sqrt{b}} + \frac{\log\left(\left|-\sqrt{b} e^{(2 dx+2 c)} + \sqrt{b e^{(4 dx+4 c)} - b}\right|\right)}{\sqrt{b}} - \frac{8}{\sqrt{b} e^{(2 dx+2 c)} - \sqrt{b e^{(4 dx+4 c)} - b} - \sqrt{b}}$$

2 b d

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*tanh(d*x+c))^(3/2),x, algorithm="giac")

[Out] $-1/2*(2*\arctan(-(\sqrt{b}*e^{(2*d*x + 2*c)} - \sqrt{b*e^{(4*d*x + 4*c)} - b}))/\sqrt{t(b)}/\sqrt{b} + \log(\text{abs}(-\sqrt{b}*e^{(2*d*x + 2*c)} + \sqrt{b*e^{(4*d*x + 4*c)} - b}))/\sqrt{b} - 8/(\sqrt{b}*e^{(2*d*x + 2*c)} - \sqrt{b*e^{(4*d*x + 4*c)} - b} - \sqrt{b}))/b*d$

Mupad [B]

time = 1.20, size = 64, normalized size = 0.82

$$\frac{\operatorname{atanh}\left(\frac{\sqrt{b} \tanh(c + dx)}{\sqrt{b}}\right)}{b^{3/2} d} - \frac{\operatorname{atan}\left(\frac{\sqrt{b} \tanh(c + dx)}{\sqrt{b}}\right)}{b^{3/2} d} - \frac{2}{b d \sqrt{b \tanh(c + dx)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*tanh(c + d*x))^(3/2),x)

[Out] $\operatorname{atanh}((b*\tanh(c + d*x))^{(1/2)}/b^{(1/2)})/(b^{(3/2)}*d) - \operatorname{atan}((b*\tanh(c + d*x))^{(1/2)}/b^{(1/2)})/(b^{(3/2)}*d) - 2/(b*d*(b*\tanh(c + d*x))^{(1/2)})$

$$3.19 \quad \int \frac{1}{(b \tanh(c+dx))^{5/2}} dx$$

Optimal. Leaf size=79

$$\frac{\text{ArcTan}\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right)}{b^{5/2}d} + \frac{\tanh^{-1}\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right)}{b^{5/2}d} - \frac{2}{3bd(b \tanh(c+dx))^{3/2}}$$

[Out] arctan((b*tanh(d*x+c))^(1/2)/b^(1/2))/b^(5/2)/d+arctanh((b*tanh(d*x+c))^(1/2)/b^(1/2))/b^(5/2)/d-2/3/b/d/(b*tanh(d*x+c))^(3/2)

Rubi [A]

time = 0.04, antiderivative size = 79, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3555, 3557, 335, 218, 212, 209}

$$\frac{\text{ArcTan}\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right)}{b^{5/2}d} + \frac{\tanh^{-1}\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right)}{b^{5/2}d} - \frac{2}{3bd(b \tanh(c+dx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[(b*Tanh[c + d*x])^(-5/2),x]

[Out] ArcTan[Sqrt[b*Tanh[c + d*x]]/Sqrt[b]]/(b^(5/2)*d) + ArcTanh[Sqrt[b*Tanh[c + d*x]]/Sqrt[b]]/(b^(5/2)*d) - 2/(3*b*d*(b*Tanh[c + d*x])^(3/2))

Rule 209

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 218

Int[((a_) + (b_)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 335

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
  Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
  ))^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
  ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 3555

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(b*Tan[c + d*x]
  )^(n + 1)/(b*d*(n + 1)), x] - Dist[1/b^2, Int[(b*Tan[c + d*x])^(n + 2), x],
  x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1]
```

Rule 3557

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[b/d, Subst[Int[
  x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !
  IntegerQ[n]
```

Rubi steps

$$\begin{aligned}
 \int \frac{1}{(b \tanh(c + dx))^{5/2}} dx &= -\frac{2}{3bd(b \tanh(c + dx))^{3/2}} + \frac{\int \frac{1}{\sqrt{b \tanh(c + dx)}} dx}{b^2} \\
 &= -\frac{2}{3bd(b \tanh(c + dx))^{3/2}} - \frac{\text{Subst}\left(\int \frac{1}{\sqrt{x(-b^2+x^2)}} dx, x, b \tanh(c + dx)\right)}{bd} \\
 &= -\frac{2}{3bd(b \tanh(c + dx))^{3/2}} - \frac{2\text{Subst}\left(\int \frac{1}{-b^2+x^4} dx, x, \sqrt{b \tanh(c + dx)}\right)}{bd} \\
 &= -\frac{2}{3bd(b \tanh(c + dx))^{3/2}} + \frac{\text{Subst}\left(\int \frac{1}{b-x^2} dx, x, \sqrt{b \tanh(c + dx)}\right)}{b^2d} + \frac{\text{Subst}\left(\int \frac{1}{b+x^2} dx, x, \sqrt{b \tanh(c + dx)}\right)}{b^2d} \\
 &= \frac{\tan^{-1}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{b^{5/2}d} + \frac{\tan^{-1}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{b^{5/2}d} - \frac{2}{3bd(b \tanh(c + dx))^{3/2}}
 \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.05, size = 38, normalized size = 0.48

$$-\frac{{}_2F_1\left(-\frac{3}{4}, 1; \frac{1}{4}; \tanh^2(c + dx)\right)}{3bd(b \tanh(c + dx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Tanh[c + d*x])^(-5/2),x]

[Out] $(-2*\text{Hypergeometric2F1}[-3/4, 1, 1/4, \text{Tanh}[c + d*x]^2])/(3*b*d*(b*\text{Tanh}[c + d*x])^{(3/2)})$

Maple [A]

time = 2.01, size = 63, normalized size = 0.80

method	result	size
derivativedivides	$2b \left(\frac{\arctan\left(\frac{\sqrt{b \tanh(dx+c)}}{\sqrt{b}}\right)}{2b^{7/2}} + \frac{1}{3b^2(b \tanh(dx+c))^{3/2}} - \frac{\text{arctanh}\left(\frac{\sqrt{b \tanh(dx+c)}}{\sqrt{b}}\right)}{2b^{7/2}} \right) \frac{1}{d}$	63
default	$2b \left(\frac{\arctan\left(\frac{\sqrt{b \tanh(dx+c)}}{\sqrt{b}}\right)}{2b^{7/2}} + \frac{1}{3b^2(b \tanh(dx+c))^{3/2}} - \frac{\text{arctanh}\left(\frac{\sqrt{b \tanh(dx+c)}}{\sqrt{b}}\right)}{2b^{7/2}} \right) \frac{1}{d}$	63

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*tanh(d*x+c))^(5/2),x,method=_RETURNVERBOSE)

[Out] $-2/d*b*(-1/2/b^{(7/2)}*\arctan((b*\tanh(d*x+c))^{(1/2)}/b^{(1/2)})+1/3/b^{(2)}/(b*\tanh(d*x+c))^{(3/2)}-1/2/b^{(7/2)}*\text{arctanh}((b*\tanh(d*x+c))^{(1/2)}/b^{(1/2)}))$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*tanh(d*x+c))^(5/2),x, algorithm="maxima")

[Out] integrate((b*tanh(d*x + c))^(5/2), x)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 691 vs. 2(63) = 126.

time = 0.43, size = 1436, normalized size = 18.18

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*tanh(d*x+c))^(5/2),x, algorithm="fricas")

[Out] $[-1/12*(6*(\cosh(d*x + c))^4 + 4*\cosh(d*x + c)*\sinh(d*x + c)^3 + \sinh(d*x + c)^4 + 2*(3*\cosh(d*x + c)^2 - 1)*\sinh(d*x + c)^2 - 2*\cosh(d*x + c)^2 + 4*(\co$

```

sh(d*x + c)^3 - cosh(d*x + c))*sinh(d*x + c) + 1)*sqrt(-b)*arctan((cosh(d*x
+ c)^2 + 2*cosh(d*x + c)*sinh(d*x + c) + sinh(d*x + c)^2)*sqrt(-b)*sqrt(b*
sinh(d*x + c)/cosh(d*x + c)))/(b*cosh(d*x + c)^2 + 2*b*cosh(d*x + c)*sinh(d*
x + c) + b*sinh(d*x + c)^2 - b)) + 3*(cosh(d*x + c)^4 + 4*cosh(d*x + c)*sin
h(d*x + c)^3 + sinh(d*x + c)^4 + 2*(3*cosh(d*x + c)^2 - 1)*sinh(d*x + c)^2
- 2*cosh(d*x + c)^2 + 4*(cosh(d*x + c)^3 - cosh(d*x + c))*sinh(d*x + c) + 1
)*sqrt(-b)*log(-(b*cosh(d*x + c)^4 + 4*b*cosh(d*x + c)^3*sinh(d*x + c) + 6*
b*cosh(d*x + c)^2*sinh(d*x + c)^2 + 4*b*cosh(d*x + c)*sinh(d*x + c)^3 + b*s
inh(d*x + c)^4 - 2*(cosh(d*x + c)^2 + 2*cosh(d*x + c)*sinh(d*x + c) + sinh(
d*x + c)^2 + 1)*sqrt(-b)*sqrt(b*sinh(d*x + c)/cosh(d*x + c)) - 2*b)/(cosh(d
*x + c)^4 + 4*cosh(d*x + c)^3*sinh(d*x + c) + 6*cosh(d*x + c)^2*sinh(d*x +
c)^2 + 4*cosh(d*x + c)*sinh(d*x + c)^3 + sinh(d*x + c)^4)) + 8*(cosh(d*x +
c)^4 + 4*cosh(d*x + c)*sinh(d*x + c)^3 + sinh(d*x + c)^4 + 2*(3*cosh(d*x +
c)^2 + 1)*sinh(d*x + c)^2 + 2*cosh(d*x + c)^2 + 4*(cosh(d*x + c)^3 + cosh(d
*x + c))*sinh(d*x + c) + 1)*sqrt(b*sinh(d*x + c)/cosh(d*x + c)))/(b^3*d*cos
h(d*x + c)^4 + 4*b^3*d*cosh(d*x + c)*sinh(d*x + c)^3 + b^3*d*sinh(d*x + c)^
4 - 2*b^3*d*cosh(d*x + c)^2 + b^3*d + 2*(3*b^3*d*cosh(d*x + c)^2 - b^3*d)*s
inh(d*x + c)^2 + 4*(b^3*d*cosh(d*x + c)^3 - b^3*d*cosh(d*x + c))*sinh(d*x +
c)), -1/12*(6*(cosh(d*x + c)^4 + 4*cosh(d*x + c)*sinh(d*x + c)^3 + sinh(d*x
+ c)^4 + 2*(3*cosh(d*x + c)^2 - 1)*sinh(d*x + c)^2 - 2*cosh(d*x + c)^2 +
4*(cosh(d*x + c)^3 - cosh(d*x + c))*sinh(d*x + c) + 1)*sqrt(b)*arctan(sqrt(
b)*sqrt(b*sinh(d*x + c)/cosh(d*x + c)))/(b*cosh(d*x + c)^2 + 2*b*cosh(d*x +
c)*sinh(d*x + c) + b*sinh(d*x + c)^2 - b)) - 3*(cosh(d*x + c)^4 + 4*cosh(d*x
+ c)*sinh(d*x + c)^3 + sinh(d*x + c)^4 + 2*(3*cosh(d*x + c)^2 - 1)*sinh(d*x
+ c)^2 - 2*cosh(d*x + c)^2 + 4*(cosh(d*x + c)^3 - cosh(d*x + c))*sinh(d*x
+ c) + 1)*sqrt(b)*log(2*b*cosh(d*x + c)^4 + 8*b*cosh(d*x + c)^3*sinh(d*x
+ c) + 12*b*cosh(d*x + c)^2*sinh(d*x + c)^2 + 8*b*cosh(d*x + c)*sinh(d*x +
c)^3 + 2*b*sinh(d*x + c)^4 + 2*(cosh(d*x + c)^4 + 4*cosh(d*x + c)*sinh(d*x
+ c)^3 + sinh(d*x + c)^4 + (6*cosh(d*x + c)^2 + 1)*sinh(d*x + c)^2 + cosh(d
*x + c)^2 + 2*(2*cosh(d*x + c)^3 + cosh(d*x + c))*sinh(d*x + c))*sqrt(b)*sq
rt(b*sinh(d*x + c)/cosh(d*x + c)) - b) + 8*(cosh(d*x + c)^4 + 4*cosh(d*x +
c)*sinh(d*x + c)^3 + sinh(d*x + c)^4 + 2*(3*cosh(d*x + c)^2 + 1)*sinh(d*x
+ c)^2 + 2*cosh(d*x + c)^2 + 4*(cosh(d*x + c)^3 + cosh(d*x + c))*sinh(d*x +
c) + 1)*sqrt(b*sinh(d*x + c)/cosh(d*x + c)))/(b^3*d*cosh(d*x + c)^4 + 4*b^3
*d*cosh(d*x + c)*sinh(d*x + c)^3 + b^3*d*sinh(d*x + c)^4 - 2*b^3*d*cosh(d*x
+ c)^2 + b^3*d + 2*(3*b^3*d*cosh(d*x + c)^2 - b^3*d)*sinh(d*x + c)^2 + 4*(
b^3*d*cosh(d*x + c)^3 - b^3*d*cosh(d*x + c))*sinh(d*x + c))]

```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \tanh(c + dx))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*tanh(d*x+c))**(5/2),x)

[Out] Integral((b*tanh(c + d*x))**(-5/2), x)

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 173 vs. 2(63) = 126.

time = 0.57, size = 173, normalized size = 2.19

$$\frac{6 \arctan\left(\frac{-\sqrt{b} e^{(2dx+2c)} - \sqrt{be^{(4dx+4c)} - b}}{\sqrt{b}}\right) - 3 \log\left(\left|-\sqrt{b} e^{(2dx+2c)} + \sqrt{be^{(4dx+4c)} - b}\right|\right) + \frac{8 \left(3 \left(\sqrt{b} e^{(2dx+2c)} - \sqrt{be^{(4dx+4c)} - b}\right)^2 + b\right)}{\left(\sqrt{b} e^{(2dx+2c)} - \sqrt{be^{(4dx+4c)} - b} - \sqrt{b}\right)^3}}{6d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*tanh(d*x+c))^(5/2),x, algorithm="giac")

[Out] 1/6*(6*arctan(-sqrt(b)*e^(2*d*x + 2*c) - sqrt(b*e^(4*d*x + 4*c) - b))/sqrt(b))/b^(5/2) - 3*log(abs(-sqrt(b)*e^(2*d*x + 2*c) + sqrt(b*e^(4*d*x + 4*c) - b)))/b^(5/2) + 8*(3*(sqrt(b)*e^(2*d*x + 2*c) - sqrt(b*e^(4*d*x + 4*c) - b))^2 + b)/((sqrt(b)*e^(2*d*x + 2*c) - sqrt(b*e^(4*d*x + 4*c) - b) - sqrt(b))^3*b^2)/d

Mupad [B]

time = 1.36, size = 63, normalized size = 0.80

$$\frac{\operatorname{atan}\left(\frac{\sqrt{b} \tanh(c + dx)}{\sqrt{b}}\right)}{b^{5/2} d} - \frac{2}{3 b d (b \tanh(c + dx))^{3/2}} + \frac{\operatorname{atanh}\left(\frac{\sqrt{b} \tanh(c + dx)}{\sqrt{b}}\right)}{b^{5/2} d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*tanh(c + d*x))^(5/2),x)

[Out] atan((b*tanh(c + d*x))^(1/2)/b^(1/2))/(b^(5/2)*d) - 2/(3*b*d*(b*tanh(c + d*x))^(3/2)) + atanh((b*tanh(c + d*x))^(1/2)/b^(1/2))/(b^(5/2)*d)

$$3.20 \quad \int \frac{1}{(b \tanh(c+dx))^{7/2}} dx$$

Optimal. Leaf size=100

$$-\frac{\operatorname{ArcTan}\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right)}{b^{7/2}d} + \frac{\tanh^{-1}\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right)}{b^{7/2}d} - \frac{2}{5bd(b \tanh(c+dx))^{5/2}} - \frac{2}{b^3d\sqrt{b \tanh(c+dx)}}$$

[Out] $-\arctan((b*\tanh(d*x+c))^{(1/2)}/b^{(1/2)})/b^{(7/2)}/d+\operatorname{arctanh}((b*\tanh(d*x+c))^{(1/2)}/b^{(1/2)})/b^{(7/2)}/d-2/b^3/d/(b*\tanh(d*x+c))^{(1/2)}-2/5/b/d/(b*\tanh(d*x+c))^{(5/2)}$

Rubi [A]

time = 0.05, antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3555, 3557, 335, 304, 209, 212}

$$-\frac{\operatorname{ArcTan}\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right)}{b^{7/2}d} + \frac{\tanh^{-1}\left(\frac{\sqrt{b \tanh(c+dx)}}{\sqrt{b}}\right)}{b^{7/2}d} - \frac{2}{b^3d\sqrt{b \tanh(c+dx)}} - \frac{2}{5bd(b \tanh(c+dx))^{5/2}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(b*\operatorname{Tanh}[c + d*x])^{(-7/2)}, x]$

[Out] $-(\operatorname{ArcTan}[\operatorname{Sqrt}[b*\operatorname{Tanh}[c + d*x]]/\operatorname{Sqrt}[b]]/(b^{(7/2)*d})) + \operatorname{ArcTanh}[\operatorname{Sqrt}[b*\operatorname{Tanh}[c + d*x]]/\operatorname{Sqrt}[b]]/(b^{(7/2)*d}) - 2/(5*b*d*(b*\operatorname{Tanh}[c + d*x])^{(5/2)}) - 2/(b^3*d*\operatorname{Sqrt}[b*\operatorname{Tanh}[c + d*x]])$

Rule 209

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[b, 2]))*\operatorname{ArcTan}[\operatorname{Rt}[b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{PosQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{GtQ}[b, 0])$

Rule 212

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$

Rule 304

$\operatorname{Int}[(x_)^2/((a_ + (b_)*(x_)^4), x_Symbol] \rightarrow \operatorname{With}\{r = \operatorname{Numerator}[\operatorname{Rt}[-a/b, 2]], s = \operatorname{Denominator}[\operatorname{Rt}[-a/b, 2]]\}, \operatorname{Dist}[s/(2*b), \operatorname{Int}[1/(r + s*x^2), x], x] - \operatorname{Dist}[s/(2*b), \operatorname{Int}[1/(r - s*x^2), x], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \ !\operatorname{GtQ}[a/b, 0]$

Rule 335

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
  Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
  ))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
  ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 3555

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(b*Tan[c + d*x]
  )^(n + 1)/(b*d*(n + 1)), x] - Dist[1/b^2, Int[(b*Tan[c + d*x])^(n + 2), x],
  x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1]
```

Rule 3557

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[b/d, Subst[Int[
  x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !
  IntegerQ[n]
```

Rubi steps

$$\begin{aligned}
 \int \frac{1}{(b \tanh(c + dx))^{7/2}} dx &= -\frac{2}{5bd(b \tanh(c + dx))^{5/2}} + \frac{\int \frac{1}{(b \tanh(c + dx))^{3/2}} dx}{b^2} \\
 &= -\frac{2}{5bd(b \tanh(c + dx))^{5/2}} - \frac{2}{b^3 d \sqrt{b \tanh(c + dx)}} + \frac{\int \sqrt{b \tanh(c + dx)} dx}{b^4} \\
 &= -\frac{2}{5bd(b \tanh(c + dx))^{5/2}} - \frac{2}{b^3 d \sqrt{b \tanh(c + dx)}} - \frac{\text{Subst}\left(\int \frac{\sqrt{x}}{-b^2 + x^2} dx, x, b \tanh(c + dx)\right)}{b^3 d} \\
 &= -\frac{2}{5bd(b \tanh(c + dx))^{5/2}} - \frac{2}{b^3 d \sqrt{b \tanh(c + dx)}} - \frac{2 \text{Subst}\left(\int \frac{x^2}{-b^2 + x^4} dx, x, \sqrt{b \tanh(c + dx)}\right)}{b^3 d} \\
 &= -\frac{2}{5bd(b \tanh(c + dx))^{5/2}} - \frac{2}{b^3 d \sqrt{b \tanh(c + dx)}} + \frac{\text{Subst}\left(\int \frac{1}{b - x^2} dx, x, \sqrt{b \tanh(c + dx)}\right)}{b^3 d} \\
 &= -\frac{\tan^{-1}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{b^{7/2} d} + \frac{\tanh^{-1}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{b^{7/2} d} - \frac{2}{5bd(b \tanh(c + dx))^{5/2}}
 \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.08, size = 38, normalized size = 0.38

$$-\frac{{}_2F_1\left(-\frac{5}{4}, 1; -\frac{1}{4}; \tanh^2(c + dx)\right)}{5bd(b \tanh(c + dx))^{5/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Tanh[c + d*x])^(-7/2),x]

[Out] (-2*Hypergeometric2F1[-5/4, 1, -1/4, Tanh[c + d*x]^2])/(5*b*d*(b*Tanh[c + d*x])^(5/2))

Maple [A]

time = 2.07, size = 77, normalized size = 0.77

method	result
derivativedivides	$2b \left(\frac{1}{b^4 \sqrt{b \tanh(dx + c)}} + \frac{1}{5b^2 (b \tanh(dx + c))^{5/2}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{b \tanh(dx + c)}}{\sqrt{b}}\right)}{2b^{9/2}} + \frac{\operatorname{arctan}\left(\frac{\sqrt{b \tanh(dx + c)}}{\sqrt{b}}\right)}{2b^{9/2}} \right)$
default	$2b \left(\frac{1}{b^4 \sqrt{b \tanh(dx + c)}} + \frac{1}{5b^2 (b \tanh(dx + c))^{5/2}} - \frac{d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(dx + c)}}{\sqrt{b}}\right)}{2b^{9/2}} + \frac{d \operatorname{arctan}\left(\frac{\sqrt{b \tanh(dx + c)}}{\sqrt{b}}\right)}{2b^{9/2}} \right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*tanh(d*x+c))^(7/2),x,method=_RETURNVERBOSE)

[Out] -2/d*b*(1/b^4/(b*tanh(d*x+c))^(1/2)+1/5/b^2/(b*tanh(d*x+c))^(5/2)-1/2/b^(9/2)*arctanh((b*tanh(d*x+c))^(1/2)/b^(1/2))+1/2/b^(9/2)*arctan((b*tanh(d*x+c))^(1/2)/b^(1/2)))

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*tanh(d*x+c))^(7/2),x, algorithm="maxima")

[Out] integrate((b*tanh(d*x + c))^(7/2), x)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 1045 vs. 2(82) = 164.

time = 0.52, size = 2144, normalized size = 21.44

Too large to display

+ c)^4 + 8*b*cosh(d*x + c)^3*sinh(d*x + c) + 12*b*cosh(d*x + c)^2*sinh(d*x + c)^2 + 8*b*cosh(d*x + c)*sinh(d*x + c)^3 + 2*b*sinh(d*x + c)^4 + 2*(cosh(d*x + c)^4 + 4*cosh(d*x + c)*sinh(d*x + c)^3 + sinh(d*x + c)^4 + (6*cosh(d*x + c)^2 + 1)*sinh(d*x + c)^2 + cosh(d*x + c)^2 + 2*(2*cosh(d*x + c)^3 + cosh(d*x + c))*sinh(d*x + c))*sqrt(b)*sqrt(b*sinh(d*x + c)/cosh(d*x + c)) - b) - 16*(3*cosh(d*x + c)^6 + 18*cosh(d*x + c)*sinh(d*x + c)^5 + 3*sinh(d*x + c)^6 + (45*cosh(d*x + c)^2 - 1)*sinh(d*x + c)^4 - cosh(d*x + c)^4 + 4*(15*cosh(d*x + c)^3 - cosh(d*x + c))*sinh(d*x + c)^3 + (45*cosh(d*x + c)^4 - 6*cosh(d*x + c)^2 - 1)*sinh(d*x + c)^2 - cosh(d*x + c)^2 + 2*(9*cosh(d*x + c)^5 - 2*cosh(d*x + c)^3 - cosh(d*x + c))*sinh(d*x + c) + 3)*sqrt(b*sinh(d*x + c)/cosh(d*x + c)))/(b^4*d*cosh(d*x + c)^6 + 6*b^4*d*cosh(d*x + c)*sinh(d*x + c)^5 + b^4*d*sinh(d*x + c)^6 - 3*b^4*d*cosh(d*x + c)^4 + 3*b^4*d*cosh(d*x + c)^2 - b^4*d + 3*(5*b^4*d*cosh(d*x + c)^2 - b^4*d)*sinh(d*x + c)^4 + 4*(5*b^4*d*cosh(d*x + c)^3 - 3*b^4*d*cosh(d*x + c))*sinh(d*x + c)^3 + 3*(5*b^4*d*cosh(d*x + c)^4 - 6*b^4*d*cosh(d*x + c)^2 + b^4*d)*sinh(d*x + c)^2 + 6*(b^4*d*cosh(d*x + c)^5 - 2*b^4*d*cosh(d*x + c)^3 + b^4*d*cosh(d*x + c))*sinh(d*x + c))]

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \tanh(c + dx))^{\frac{7}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*tanh(d*x+c))**(7/2),x)

[Out] Integral((b*tanh(c + d*x))**(-7/2), x)

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 293 vs. 2(82) = 164.

time = 0.63, size = 293, normalized size = 2.93

$$\frac{10 \arctan\left(\frac{-\sqrt{b} e^{2dx+2c} - \sqrt{be^{4dx+4c} - b}}{\sqrt{b}}\right) + 5 \log\left(\frac{-\sqrt{b} e^{2dx+2c} + \sqrt{be^{4dx+4c} - b}}{\sqrt{b}}\right) - 16 \left(\sqrt{b} e^{2dx+2c} - \sqrt{be^{4dx+4c} - b}\right)^{-1} - 10 \left(\sqrt{b} e^{2dx+2c} - \sqrt{be^{4dx+4c} - b}\right)^3 \sqrt{b} + 20 \left(\sqrt{b} e^{2dx+2c} - \sqrt{be^{4dx+4c} - b}\right)^5 - 10 \left(\sqrt{b} e^{2dx+2c} - \sqrt{be^{4dx+4c} - b}\right)^3 b^{3/2}}{\left(\sqrt{b} e^{2dx+2c} - \sqrt{be^{4dx+4c} - b} - \sqrt{b}\right)^5} \frac{1}{10d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*tanh(d*x+c))^(7/2),x, algorithm="giac")

[Out] -1/10*(10*arctan(-sqrt(b)*e^(2*d*x + 2*c) - sqrt(b*e^(4*d*x + 4*c) - b))/sqrt(b))/b^(7/2) + 5*log(abs(-sqrt(b)*e^(2*d*x + 2*c) + sqrt(b*e^(4*d*x + 4*c) - b)))/b^(7/2) - 16*(5*(sqrt(b)*e^(2*d*x + 2*c) - sqrt(b*e^(4*d*x + 4*c) - b))^4 - 10*(sqrt(b)*e^(2*d*x + 2*c) - sqrt(b*e^(4*d*x + 4*c) - b))^3*sqrt(b) + 20*(sqrt(b)*e^(2*d*x + 2*c) - sqrt(b*e^(4*d*x + 4*c) - b))^2*b - 10*(sqrt(b)*e^(2*d*x + 2*c) - sqrt(b*e^(4*d*x + 4*c) - b))*b^(3/2) + 3*b^2)/((sqrt(b)*e^(2*d*x + 2*c) - sqrt(b*e^(4*d*x + 4*c) - b) - sqrt(b))^5*b^3)/d

Mupad [B]

time = 1.51, size = 80, normalized size = 0.80

$$\frac{\operatorname{atanh}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{b^{7/2} d} - \frac{\operatorname{atan}\left(\frac{\sqrt{b \tanh(c + dx)}}{\sqrt{b}}\right)}{b^{7/2} d} - \frac{\frac{2}{5b} + \frac{2 \tanh(c + dx)^2}{b}}{d (b \tanh(c + dx))^{5/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(b*tanh(c + d*x))^(7/2),x)`

[Out] `atanh((b*tanh(c + d*x))^(1/2)/b^(1/2))/(b^(7/2)*d) - atan((b*tanh(c + d*x))^(1/2)/b^(1/2))/(b^(7/2)*d) - (2/(5*b) + (2*tanh(c + d*x)^2)/b)/(d*(b*tanh(c + d*x))^(5/2))`

3.21 $\int \sqrt[3]{\tanh(8x)} dx$

Optimal. Leaf size=69

$$-\frac{1}{16}\sqrt{3} \operatorname{ArcTan}\left(\frac{1+2\tanh^{\frac{2}{3}}(8x)}{\sqrt{3}}\right) - \frac{1}{16} \log\left(1 - \tanh^{\frac{2}{3}}(8x)\right) + \frac{1}{32} \log\left(1 + \tanh^{\frac{2}{3}}(8x) + \tanh^{\frac{4}{3}}(8x)\right)$$

[Out] -1/16*ln(1-tanh(8*x)^(2/3))+1/32*ln(1+tanh(8*x)^(2/3)+tanh(8*x)^(4/3))-1/16*arctan(1/3*(1+2*tanh(8*x)^(2/3))*3^(1/2))*3^(1/2)

Rubi [A]

time = 0.04, antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 9, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 1.125$, Rules used = {3557, 335, 281, 298, 31, 648, 632, 210, 642}

$$-\frac{1}{16}\sqrt{3} \operatorname{ArcTan}\left(\frac{2\tanh^{\frac{2}{3}}(8x)+1}{\sqrt{3}}\right) - \frac{1}{16} \log\left(1 - \tanh^{\frac{2}{3}}(8x)\right) + \frac{1}{32} \log\left(\tanh^{\frac{4}{3}}(8x) + \tanh^{\frac{2}{3}}(8x) + 1\right)$$

Antiderivative was successfully verified.

[In] Int[Tanh[8*x]^(1/3),x]

[Out] -1/16*(Sqrt[3]*ArcTan[(1 + 2*Tanh[8*x]^(2/3))/Sqrt[3]]) - Log[1 - Tanh[8*x]^(2/3)]/16 + Log[1 + Tanh[8*x]^(2/3) + Tanh[8*x]^(4/3)]/32

Rule 31

Int[((a_) + (b_.)*(x_))^(n_), x_Symbol] := Simp[Log[RemoveContent[a + b*x, x]]/b, x] /; FreeQ[{a, b}, x]

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(n_), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(n_+1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 281

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Dist[1/k, Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1 /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]

Rule 298

Int[(x_)/((a_) + (b_.)*(x_)^3), x_Symbol] := Dist[-(3*Rt[a, 3]*Rt[b, 3])^(n_+1), Int[1/(Rt[a, 3] + Rt[b, 3]*x), x], x] + Dist[1/(3*Rt[a, 3]*Rt[b, 3]), Int[(Rt[a, 3] + Rt[b, 3]*x)/(Rt[a, 3]^2 - Rt[a, 3]*Rt[b, 3]*x + Rt[b, 3]^2*x

$^2), x], x] /; \text{FreeQ}\{a, b\}, x]$

Rule 335

$\text{Int}[(c_.)*(x_.)^{(m_)}*((a_.) + (b_.)*(x_.)^{(n_)})^{(p_)}, x_Symbol] \rightarrow \text{With}\{k = \text{Denominator}[m]\}, \text{Dist}[k/c, \text{Subst}[\text{Int}[x^{(k*(m+1)-1)}*(a + b*(x^{(k*n)}/c^n)]^{(p)}, x], x, (c*x)^{(1/k)}], x] /; \text{FreeQ}\{a, b, c, p\}, x] \&\& \text{IGtQ}[n, 0] \&\& \text{FractionQ}[m] \&\& \text{IntBinomialQ}[a, b, c, n, m, p, x]$

Rule 632

$\text{Int}[(a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2]^{(-1)}, x_Symbol] \rightarrow \text{Dist}[-2, \text{Subst}[\text{Int}[1/\text{Simp}[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; \text{FreeQ}\{a, b, c\}, x] \&\& \text{NeQ}[b^2 - 4*a*c, 0]$

Rule 642

$\text{Int}[(d_.) + (e_.)*(x_.)]/((a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2), x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}\{a, b, c, d, e\}, x] \&\& \text{EqQ}[2*c*d - b*e, 0]$

Rule 648

$\text{Int}[(d_.) + (e_.)*(x_.)]/((a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2), x_Symbol] \rightarrow \text{Dist}[(2*c*d - b*e)/(2*c), \text{Int}[1/(a + b*x + c*x^2), x], x] + \text{Dist}[e/(2*c), \text{Int}[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; \text{FreeQ}\{a, b, c, d, e\}, x] \&\& \text{NeQ}[2*c*d - b*e, 0] \&\& \text{NeQ}[b^2 - 4*a*c, 0] \&\& \text{!NiceSqrtQ}[b^2 - 4*a*c]$

Rule 3557

$\text{Int}[(b_.)*\tan[(c_.) + (d_.)*(x_.)])^{(n_)}, x_Symbol] \rightarrow \text{Dist}[b/d, \text{Subst}[\text{Int}[x^n/(b^2 + x^2), x], x, b*\text{Tan}[c + d*x]], x] /; \text{FreeQ}\{b, c, d, n\}, x] \&\& \text{!IntegerQ}[n]$

Rubi steps

$$\begin{aligned}
\int \sqrt[3]{\tanh(8x)} dx &= -\left(\frac{1}{8}\text{Subst}\left(\int \frac{\sqrt[3]{x}}{-1+x^2} dx, x, \tanh(8x)\right)\right) \\
&= -\left(\frac{3}{8}\text{Subst}\left(\int \frac{x^3}{-1+x^6} dx, x, \sqrt[3]{\tanh(8x)}\right)\right) \\
&= -\left(\frac{3}{16}\text{Subst}\left(\int \frac{x}{-1+x^3} dx, x, \tanh^{\frac{2}{3}}(8x)\right)\right) \\
&= -\left(\frac{1}{16}\text{Subst}\left(\int \frac{1}{-1+x} dx, x, \tanh^{\frac{2}{3}}(8x)\right)\right) + \frac{1}{16}\text{Subst}\left(\int \frac{-1+x}{1+x+x^2} dx, x, \tanh^{\frac{2}{3}}(8x)\right) \\
&= -\frac{1}{16}\log\left(1 - \tanh^{\frac{2}{3}}(8x)\right) + \frac{1}{32}\text{Subst}\left(\int \frac{1+2x}{1+x+x^2} dx, x, \tanh^{\frac{2}{3}}(8x)\right) - \frac{3}{32}\text{Subst}\left(\int \frac{1}{-3-x} dx, x, \tanh^{\frac{2}{3}}(8x)\right) \\
&= -\frac{1}{16}\log\left(1 - \tanh^{\frac{2}{3}}(8x)\right) + \frac{1}{32}\log\left(1 + \tanh^{\frac{2}{3}}(8x) + \tanh^{\frac{4}{3}}(8x)\right) + \frac{3}{16}\text{Subst}\left(\int \frac{1}{-3-x} dx, x, \tanh^{\frac{2}{3}}(8x)\right) \\
&= -\frac{1}{16}\sqrt{3}\tan^{-1}\left(\frac{1+2\tanh^{\frac{2}{3}}(8x)}{\sqrt{3}}\right) - \frac{1}{16}\log\left(1 - \tanh^{\frac{2}{3}}(8x)\right) + \frac{1}{32}\log\left(1 + \tanh^{\frac{2}{3}}(8x) + \tanh^{\frac{4}{3}}(8x)\right)
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 26, normalized size = 0.38

$$\frac{3}{32} {}_2F_1\left(\frac{2}{3}, 1; \frac{5}{3}; \tanh^2(8x)\right) \tanh^{\frac{4}{3}}(8x)$$

Antiderivative was successfully verified.

[In] Integrate[Tanh[8*x]^(1/3), x]

[Out] (3*Hypergeometric2F1[2/3, 1, 5/3, Tanh[8*x]^2]*Tanh[8*x]^(4/3))/32

Maple [A]

time = 0.50, size = 102, normalized size = 1.48

method	result
derivativedivides	$-\frac{\ln(\tanh^{\frac{1}{3}}(8x)-1)}{16} + \frac{\ln(\tanh^{\frac{2}{3}}(8x)+\tanh^{\frac{1}{3}}(8x)+1)}{32} + \frac{\sqrt{3}\arctan\left(\frac{(2(\tanh^{\frac{1}{3}}(8x)+1)\sqrt{3})}{3}\right)}{16} + \frac{\ln(\tanh^{\frac{2}{3}}(8x))}{32}$
default	$-\frac{\ln(\tanh^{\frac{1}{3}}(8x)-1)}{16} + \frac{\ln(\tanh^{\frac{2}{3}}(8x)+\tanh^{\frac{1}{3}}(8x)+1)}{32} + \frac{\sqrt{3}\arctan\left(\frac{(2(\tanh^{\frac{1}{3}}(8x)+1)\sqrt{3})}{3}\right)}{16} + \frac{\ln(\tanh^{\frac{2}{3}}(8x))}{32}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tanh(8*x)^(1/3),x,method=_RETURNVERBOSE)`

[Out] $-1/16*\ln(\tanh(8*x)^{(1/3)}-1)+1/32*\ln(\tanh(8*x)^{(2/3)}+\tanh(8*x)^{(1/3)}+1)+1/16*3^{(1/2)}*\arctan(1/3*(2*\tanh(8*x)^{(1/3)}+1)*3^{(1/2)})+1/32*\ln(\tanh(8*x)^{(2/3)}-\tanh(8*x)^{(1/3)}+1)-1/16*3^{(1/2)}*\arctan(1/3*(2*\tanh(8*x)^{(1/3)}-1)*3^{(1/2)})-1/16*\ln(\tanh(8*x)^{(1/3)}+1)$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(8*x)^(1/3),x, algorithm="maxima")`

[Out] `integrate(tanh(8*x)^(1/3), x)`

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 179 vs. 2(52) = 104.

time = 0.37, size = 179, normalized size = 2.59

$$-\frac{1}{16}\sqrt{3}\arctan\left(\frac{2}{3}\sqrt{3}\frac{\sinh(8x)}{\cosh(8x)}+\frac{1}{3}\sqrt{3}\right)-\frac{1}{16}\log\left(\frac{\sinh(8x)}{\cosh(8x)}-1\right)+\frac{1}{32}\log\left(\frac{\cosh(8x)^2+2\cosh(8x)\sinh(8x)+\sinh(8x)^2+(\cosh(8x)^2+2\cosh(8x)\sinh(8x)+\sinh(8x)^2+1)\left(\frac{\sinh(8x)}{\cosh(8x)}\right)^{\frac{2}{3}}+(\cosh(8x)^2+2\cosh(8x)\sinh(8x)+\sinh(8x)^2-1)\left(\frac{\sinh(8x)}{\cosh(8x)}\right)^{\frac{1}{3}}+1}{\cosh(8x)^2+2\cosh(8x)\sinh(8x)+\sinh(8x)^2+1}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(8*x)^(1/3),x, algorithm="fricas")`

[Out] $-1/16*\sqrt{3}*\arctan(2/3*\sqrt{3}*(\sinh(8*x)/\cosh(8*x))^{(2/3)}+1/3*\sqrt{3})-1/16*\log((\sinh(8*x)/\cosh(8*x))^{(2/3)}-1)+1/32*\log((\cosh(8*x)^2+2*\cosh(8*x)*\sinh(8*x)+\sinh(8*x)^2+(\cosh(8*x)^2+2*\cosh(8*x)*\sinh(8*x)+\sinh(8*x)^2+1)*(\sinh(8*x)/\cosh(8*x))^{(2/3)}+(\cosh(8*x)^2+2*\cosh(8*x)*\sinh(8*x)+\sinh(8*x)^2-1)*(\sinh(8*x)/\cosh(8*x))^{(1/3)}+1)/(\cosh(8*x)^2+2*\cosh(8*x)*\sinh(8*x)+\sinh(8*x)^2+1))$

Sympy [A]

time = 1.95, size = 63, normalized size = 0.91

$$-\frac{\log\left(\tanh^{\frac{2}{3}}(8x)-1\right)}{16}+\frac{\log\left(\tanh^{\frac{4}{3}}(8x)+\tanh^{\frac{2}{3}}(8x)+1\right)}{32}-\frac{\sqrt{3}\operatorname{atan}\left(\frac{2\sqrt{3}\left(\tanh^{\frac{2}{3}}(8x)+\frac{1}{2}\right)}{3}\right)}{16}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(8*x)**(1/3),x)`

[Out] $-\log(\tanh(8*x)**(2/3)-1)/16+\log(\tanh(8*x)**(4/3)+\tanh(8*x)**(2/3)+1)/32-\sqrt{3}*\operatorname{atan}(2*\sqrt{3}*(\tanh(8*x)**(2/3)+1/2)/3)/16$

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 110 vs. 2(52) = 104.

time = 0.42, size = 110, normalized size = 1.59

$$-\frac{1}{16}\sqrt{3}\arctan\left(\frac{1}{3}\sqrt{3}\left(2\left(\frac{e^{16x}-1}{e^{16x}+1}\right)^{\frac{2}{3}}+1\right)\right)+\frac{1}{32}\log\left(\left(\frac{e^{16x}-1}{e^{16x}+1}\right)^{\frac{2}{3}}+\frac{\left(\frac{e^{16x}-1}{e^{16x}+1}\right)^{\frac{1}{3}}(e^{16x}-1)}{e^{16x}+1}+1\right)-\frac{1}{16}\log\left(\left|\left(\frac{e^{16x}-1}{e^{16x}+1}\right)^{\frac{2}{3}}-1\right|\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(8*x)^(1/3),x, algorithm="giac")

[Out] -1/16*sqrt(3)*arctan(1/3*sqrt(3)*(2*((e^(16*x) - 1)/(e^(16*x) + 1))^(2/3) + 1)) + 1/32*log(((e^(16*x) - 1)/(e^(16*x) + 1))^(2/3) + ((e^(16*x) - 1)/(e^(16*x) + 1))^(1/3)*(e^(16*x) - 1)/(e^(16*x) + 1) + 1) - 1/16*log(abs(((e^(16*x) - 1)/(e^(16*x) + 1))^(2/3) - 1))

Mupad [B]

time = 1.32, size = 71, normalized size = 1.03

$$-\frac{\ln(81\tanh(8x)^{2/3}-81)}{16}-\ln\left(162\tanh(8x)^{2/3}\left(-\frac{1}{4}+\frac{\sqrt{3}i}{4}\right)-81\right)\left(-\frac{1}{32}+\frac{\sqrt{3}i}{32}\right)+\ln\left(-162\tanh(8x)^{2/3}\left(\frac{1}{4}+\frac{\sqrt{3}i}{4}\right)-81\right)\left(\frac{1}{32}+\frac{\sqrt{3}i}{32}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(8*x)^(1/3),x)

[Out] log(-162*tanh(8*x)^(2/3)*((3^(1/2)*i)/4 + 1/4) - 81)*((3^(1/2)*i)/32 + 1/32) - log(162*tanh(8*x)^(2/3)*((3^(1/2)*i)/4 - 1/4) - 81)*((3^(1/2)*i)/32 - 1/32) - log(81*tanh(8*x)^(2/3) - 81)/16

3.22 $\int \tanh^n(a + bx) dx$

Optimal. Leaf size=43

$$\frac{{}_2F_1\left(1, \frac{1+n}{2}; \frac{3+n}{2}; \tanh^2(a + bx)\right) \tanh^{1+n}(a + bx)}{b(1+n)}$$

[Out] hypergeom([1, 1/2+1/2*n], [3/2+1/2*n], tanh(b*x+a)^2)*tanh(b*x+a)^(1+n)/b/(1+n)

Rubi [A]

time = 0.02, antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3557, 371}

$$\frac{\tanh^{n+1}(a + bx) {}_2F_1\left(1, \frac{n+1}{2}; \frac{n+3}{2}; \tanh^2(a + bx)\right)}{b(n+1)}$$

Antiderivative was successfully verified.

[In] Int[Tanh[a + b*x]^n, x]

[Out] (Hypergeometric2F1[1, (1 + n)/2, (3 + n)/2, Tanh[a + b*x]^2]*Tanh[a + b*x]^(1 + n))/(b*(1 + n))

Rule 371

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p * ((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])

Rule 3557

Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[b/d, Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\begin{aligned} \int \tanh^n(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{x^n}{-1+x^2} dx, x, \tanh(a + bx)\right)}{b} \\ &= \frac{{}_2F_1\left(1, \frac{1+n}{2}; \frac{3+n}{2}; \tanh^2(a + bx)\right) \tanh^{1+n}(a + bx)}{b(1+n)} \end{aligned}$$

Mathematica [A]

time = 0.05, size = 45, normalized size = 1.05

$$\frac{{}_2F_1\left(1, \frac{1+n}{2}; 1 + \frac{1+n}{2}; \tanh^2(a + bx)\right) \tanh^{1+n}(a + bx)}{b(1 + n)}$$

Antiderivative was successfully verified.

`[In] Integrate[Tanh[a + b*x]^n, x]``[Out] (Hypergeometric2F1[1, (1 + n)/2, 1 + (1 + n)/2, Tanh[a + b*x]^2]*Tanh[a + b*x]^(1 + n))/(b*(1 + n))`**Maple [F]**

time = 1.85, size = 0, normalized size = 0.00

$$\int \tanh^n(bx + a) dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(tanh(b*x+a)^n, x)``[Out] int(tanh(b*x+a)^n, x)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(tanh(b*x+a)^n, x, algorithm="maxima")``[Out] integrate(tanh(b*x + a)^n, x)`**Fricas [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(tanh(b*x+a)^n, x, algorithm="fricas")``[Out] integral(tanh(b*x + a)^n, x)`**Sympy [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int \tanh^n(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(b*x+a)**n,x)`

[Out] `Integral(tanh(a + b*x)**n, x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(b*x+a)^n,x, algorithm="giac")`

[Out] `integrate(tanh(b*x + a)^n, x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \tanh(a + b x)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tanh(a + b*x)^n,x)`

[Out] `int(tanh(a + b*x)^n, x)`

3.23 $\int (b \tanh(c + dx))^n dx$

Optimal. Leaf size=48

$$\frac{{}_2F_1\left(1, \frac{1+n}{2}; \frac{3+n}{2}; \tanh^2(c + dx)\right) (b \tanh(c + dx))^{1+n}}{bd(1+n)}$$

[Out] hypergeom([1, 1/2+1/2*n], [3/2+1/2*n], tanh(d*x+c)^2)*(b*tanh(d*x+c))^(1+n)/b/d/(1+n)

Rubi [A]

time = 0.02, antiderivative size = 48, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3557, 371}

$$\frac{(b \tanh(c + dx))^{n+1} {}_2F_1\left(1, \frac{n+1}{2}; \frac{n+3}{2}; \tanh^2(c + dx)\right)}{bd(n+1)}$$

Antiderivative was successfully verified.

[In] Int[(b*Tanh[c + d*x])^n,x]

[Out] (Hypergeometric2F1[1, (1 + n)/2, (3 + n)/2, Tanh[c + d*x]^2]*(b*Tanh[c + d*x])^(1 + n))/(b*d*(1 + n))

Rule 371

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])

Rule 3557

Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[b/d, Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\begin{aligned} \int (b \tanh(c + dx))^n dx &= -\frac{b \text{Subst}\left(\int \frac{x^n}{-b^2+x^2} dx, x, b \tanh(c + dx)\right)}{d} \\ &= \frac{{}_2F_1\left(1, \frac{1+n}{2}; \frac{3+n}{2}; \tanh^2(c + dx)\right) (b \tanh(c + dx))^{1+n}}{bd(1+n)} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 51, normalized size = 1.06

$$\frac{{}_2F_1\left(1, \frac{1+n}{2}; 1 + \frac{1+n}{2}; \tanh^2(c + dx)\right) \tanh(c + dx) (b \tanh(c + dx))^n}{d(1 + n)}$$

Antiderivative was successfully verified.

`[In] Integrate[(b*Tanh[c + d*x])^n,x]``[Out] (Hypergeometric2F1[1, (1 + n)/2, 1 + (1 + n)/2, Tanh[c + d*x]^2]*Tanh[c + d*x]*(b*Tanh[c + d*x])^n)/(d*(1 + n))`**Maple [F]**

time = 1.78, size = 0, normalized size = 0.00

$$\int (b \tanh(dx + c))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((b*tanh(d*x+c))^n,x)``[Out] int((b*tanh(d*x+c))^n,x)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((b*tanh(d*x+c))^n,x, algorithm="maxima")``[Out] integrate((b*tanh(d*x + c))^n, x)`**Fricas [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((b*tanh(d*x+c))^n,x, algorithm="fricas")``[Out] integral((b*tanh(d*x + c))^n, x)`**Sympy [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int (b \tanh(c + dx))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*tanh(d*x+c))**n,x)

[Out] Integral((b*tanh(c + d*x))**n, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*tanh(d*x+c))^n,x, algorithm="giac")

[Out] integrate((b*tanh(d*x + c))^n, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int (b \tanh(c + dx))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*tanh(c + d*x))^n,x)

[Out] int((b*tanh(c + d*x))^n, x)

3.24 $\int (a \tanh^2(x))^{3/2} dx$

Optimal. Leaf size=35

$$a \coth(x) \log(\cosh(x)) \sqrt{a \tanh^2(x)} - \frac{1}{2} a \tanh(x) \sqrt{a \tanh^2(x)}$$

[Out] a*coth(x)*ln(cosh(x))*(a*tanh(x)^2)^(1/2)-1/2*a*(a*tanh(x)^2)^(1/2)*tanh(x)

Rubi [A]

time = 0.02, antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3739, 3554, 3556}

$$a \coth(x) \sqrt{a \tanh^2(x)} \log(\cosh(x)) - \frac{1}{2} a \tanh(x) \sqrt{a \tanh^2(x)}$$

Antiderivative was successfully verified.

[In] Int[(a*Tanh[x]^2)^(3/2),x]

[Out] a*Coth[x]*Log[Cosh[x]]*Sqrt[a*Tanh[x]^2] - (a*Tanh[x]*Sqrt[a*Tanh[x]^2])/2

Rule 3554

Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Dist[b^2, Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]

Rule 3556

Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]

Rule 3739

Int[(u_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_)^(p_), x_Symbol] :> With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[(b*ff^n)^IntPart[p]*((b*Tan[e + f*x])^n)^FracPart[p]/(Tan[e + f*x]/ff)^(n*FracPart[p])], Int[ActivateTrig[u]*(Tan[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]

Rubi steps

$$\begin{aligned}
\int (a \tanh^2(x))^{3/2} dx &= \left(a \coth(x) \sqrt{a \tanh^2(x)} \right) \int \tanh^3(x) dx \\
&= -\frac{1}{2} a \tanh(x) \sqrt{a \tanh^2(x)} + \left(a \coth(x) \sqrt{a \tanh^2(x)} \right) \int \tanh(x) dx \\
&= a \coth(x) \log(\cosh(x)) \sqrt{a \tanh^2(x)} - \frac{1}{2} a \tanh(x) \sqrt{a \tanh^2(x)}
\end{aligned}$$

Mathematica [A]

time = 0.02, size = 28, normalized size = 0.80

$$\frac{1}{2} a (2 \coth(x) \log(\cosh(x)) + \operatorname{csch}(x) \operatorname{sech}(x)) \sqrt{a \tanh^2(x)}$$

Antiderivative was successfully verified.

`[In] Integrate[(a*Tanh[x]^2)^(3/2), x]``[Out] (a*(2*Coth[x]*Log[Cosh[x]] + Csch[x]*Sech[x])*Sqrt[a*Tanh[x]^2])/2`**Maple [A]**

time = 0.75, size = 30, normalized size = 0.86

method	result	size
derivativedivides	$-\frac{(a(\tanh^2(x)))^{\frac{3}{2}}(\tanh^2(x)+\ln(\tanh(x)-1)+\ln(1+\tanh(x)))}{2\tanh(x)^3}$	30
default	$-\frac{(a(\tanh^2(x)))^{\frac{3}{2}}(\tanh^2(x)+\ln(\tanh(x)-1)+\ln(1+\tanh(x)))}{2\tanh(x)^3}$	30
risch	$-\frac{a(1+e^{2x})\sqrt{\frac{a(e^{2x}-1)^2}{(1+e^{2x})^2}}}{e^{2x}-1} + \frac{2a\sqrt{\frac{a(e^{2x}-1)^2}{(1+e^{2x})^2}}e^{2x}}{(e^{2x}-1)(1+e^{2x})} + \frac{a(1+e^{2x})\sqrt{\frac{a(e^{2x}-1)^2}{(1+e^{2x})^2}}\ln(1+e^{2x})}{e^{2x}-1}$	126

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((a*tanh(x)^2)^(3/2), x, method=_RETURNVERBOSE)``[Out] -1/2*(a*tanh(x)^2)^(3/2)*(tanh(x)^2+ln(tanh(x)-1)+ln(1+tanh(x)))/tanh(x)^3`**Maxima [A]**

time = 0.47, size = 42, normalized size = 1.20

$$-a^{\frac{3}{2}}x - a^{\frac{3}{2}} \log(e^{(-2x)} + 1) - \frac{2a^{\frac{3}{2}}e^{(-2x)}}{2e^{(-2x)} + e^{(-4x)} + 1}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((a*tanh(x)^2)^(3/2), x, algorithm="maxima")`

[Out] $-a^{3/2}x - a^{3/2}\log(e^{-2x} + 1) - 2a^{3/2}e^{-2x}/(2e^{-2x} + e^{-4x} + 1)$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 467 vs. 2(29) = 58.

time = 0.39, size = 467, normalized size = 13.34

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a*tanh(x)^2)^(3/2),x, algorithm="fricas")`

[Out] $-(a*x*\cosh(x)^4 + (a*x*e^{2x} + a*x)*\sinh(x)^4 + 4*(a*x*\cosh(x)*e^{2x} + a*x*\cosh(x))*\sinh(x)^3 + 2*(a*x - a)*\cosh(x)^2 + 2*(3*a*x*\cosh(x)^2 + a*x + (3*a*x*\cosh(x)^2 + a*x - a)*e^{2x} - a)*\sinh(x)^2 + a*x + (a*x*\cosh(x)^4 + 2*(a*x - a)*\cosh(x)^2 + a*x)*e^{2x} - (a*\cosh(x)^4 + (a*e^{2x} + a)*\sinh(x)^4 + 4*(a*\cosh(x)*e^{2x} + a*\cosh(x))*\sinh(x)^3 + 2*a*\cosh(x)^2 + 2*(3*a*\cosh(x)^2 + (3*a*\cosh(x)^2 + a)*e^{2x} + a)*\sinh(x)^2 + (a*\cosh(x)^4 + 2*a*\cosh(x)^2 + a)*e^{2x} + 4*(a*\cosh(x)^3 + a*\cosh(x) + (a*\cosh(x)^3 + a*\cosh(x))*e^{2x}))*\sinh(x) + a)*\log(2*\cosh(x)/(cosh(x) - sinh(x))) + 4*(a*x*\cosh(x)^3 + (a*x - a)*\cosh(x) + (a*x*\cosh(x)^3 + (a*x - a)*\cosh(x))*e^{2x}))*\sinh(x))*\sqrt{(a*e^{4x} - 2*a*e^{2x} + a)/(e^{4x} + 2*e^{2x} + 1))/((e^{2x} - 1)*\sinh(x)^4 - \cosh(x)^4 + 4*(\cosh(x)*e^{2x} - \cosh(x))*\sinh(x)^3 - 2*(3*\cosh(x)^2 - (3*\cosh(x)^2 + 1)*e^{2x} + 1)*\sinh(x)^2 - 2*\cosh(x)^2 + (\cosh(x)^4 + 2*\cosh(x)^2 + 1)*e^{2x} - 4*(\cosh(x)^3 - (\cosh(x)^3 + \cosh(x))*e^{2x} + \cosh(x))*\sinh(x) - 1)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (a \tanh^2(x))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a*tanh(x)**2)**(3/2),x)`

[Out] `Integral((a*tanh(x)**2)**(3/2), x)`

Giac [A]

time = 0.43, size = 52, normalized size = 1.49

$$-\left(x \operatorname{sgn}(e^{4x} - 1) - \log(e^{2x} + 1) \operatorname{sgn}(e^{4x} - 1) - \frac{2e^{2x} \operatorname{sgn}(e^{4x} - 1)}{(e^{2x} + 1)^2}\right) a^{\frac{3}{2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a*tanh(x)^2)^(3/2),x, algorithm="giac")`

[Out] $-(x*\text{sgn}(e^{4*x} - 1) - \log(e^{2*x} + 1)*\text{sgn}(e^{4*x} - 1) - 2*e^{2*x}*\text{sgn}(e^{4*x} - 1))/(e^{2*x} + 1)^2*a^{3/2}$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.03

$$\int (a \tanh(x)^2)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a*tanh(x)^2)^(3/2),x)`

[Out] `int((a*tanh(x)^2)^(3/2), x)`

3.25 $\int \sqrt{a \tanh^2(x)} dx$

Optimal. Leaf size=16

$$\coth(x) \log(\cosh(x)) \sqrt{a \tanh^2(x)}$$

[Out] $\coth(x) \ln(\cosh(x)) (a \tanh(x)^2)^{1/2}$

Rubi [A]

time = 0.01, antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3739, 3556}

$$\coth(x) \sqrt{a \tanh^2(x)} \log(\cosh(x))$$

Antiderivative was successfully verified.

[In] Int[Sqrt[a*Tanh[x]^2], x]

[Out] Coth[x]*Log[Cosh[x]]*Sqrt[a*Tanh[x]^2]

Rule 3556

Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]

Rule 3739

Int[(u_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_)^(p_), x_Symbol] :> With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[(b*ff^n)^IntPart[p]*((b*Tan[e + f*x]^n)^FracPart[p]/(Tan[e + f*x]/ff)^(n*FracPart[p])), Int[ActivateTrig[u]*(Tan[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]

Rubi steps

$$\begin{aligned} \int \sqrt{a \tanh^2(x)} dx &= \left(\coth(x) \sqrt{a \tanh^2(x)} \right) \int \tanh(x) dx \\ &= \coth(x) \log(\cosh(x)) \sqrt{a \tanh^2(x)} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 16, normalized size = 1.00

$$\coth(x) \log(\cosh(x)) \sqrt{a \tanh^2(x)}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[a*Tanh[x]^2], x]

[Out] Coth[x]*Log[Cosh[x]]*Sqrt[a*Tanh[x]^2]

Maple [A]

time = 0.76, size = 26, normalized size = 1.62

method	result	size
derivativedivides	$-\frac{\sqrt{a(\tanh^2(x))} (\ln(\tanh(x)-1)+\ln(1+\tanh(x)))}{2 \tanh(x)}$	26
default	$-\frac{\sqrt{a(\tanh^2(x))} (\ln(\tanh(x)-1)+\ln(1+\tanh(x)))}{2 \tanh(x)}$	26
risch	$-\frac{\sqrt{\frac{a(e^{2x}-1)^2}{(1+e^{2x})^2}} (1+e^{2x})x}{e^{2x}-1} + \frac{\sqrt{\frac{a(e^{2x}-1)^2}{(1+e^{2x})^2}} (1+e^{2x}) \ln(1+e^{2x})}{e^{2x}-1}$	81

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*tanh(x)^2)^(1/2), x, method=_RETURNVERBOSE)

[Out] -1/2*(a*tanh(x)^2)^(1/2)*(ln(tanh(x)-1)+ln(1+tanh(x)))/tanh(x)

Maxima [A]

time = 0.48, size = 19, normalized size = 1.19

$$-\sqrt{a} x - \sqrt{a} \log(e^{-2x} + 1)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*tanh(x)^2)^(1/2), x, algorithm="maxima")

[Out] -sqrt(a)*x - sqrt(a)*log(e^(-2*x) + 1)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 72 vs. 2(14) = 28.

time = 0.41, size = 72, normalized size = 4.50

$$-\frac{\left(xe^{2x} - (e^{2x} + 1) \log\left(\frac{2 \cosh(x)}{\cosh(x) - \sinh(x)}\right) + x\right) \sqrt{\frac{ae^{4x} - 2ae^{2x} + a}{e^{4x} + 2e^{2x} + 1}}}{e^{2x} - 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*tanh(x)^2)^(1/2), x, algorithm="fricas")

[Out] -(x*e^(2*x) - (e^(2*x) + 1)*log(2*cosh(x)/(cosh(x) - sinh(x))) + x)*sqrt((a*e^(4*x) - 2*a*e^(2*x) + a)/(e^(4*x) + 2*e^(2*x) + 1))/(e^(2*x) - 1)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{a \tanh^2(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*tanh(x)**2)**(1/2),x)**[Out]** Integral(sqrt(a*tanh(x)**2), x)**Giac [B]** Leaf count of result is larger than twice the leaf count of optimal. 31 vs. 2(14) = 28.
time = 0.41, size = 31, normalized size = 1.94

$$-(x \operatorname{sgn}(e^{4x} - 1) - \log(e^{2x} + 1) \operatorname{sgn}(e^{4x} - 1)) \sqrt{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*tanh(x)^2)^(1/2),x, algorithm="giac")**[Out]** -(x*sgn(e^(4*x) - 1) - log(e^(2*x) + 1)*sgn(e^(4*x) - 1))*sqrt(a)**Mupad [F]**

time = 0.00, size = -1, normalized size = -0.06

$$\int \sqrt{a \tanh(x)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*tanh(x)^2)^(1/2),x)**[Out]** int((a*tanh(x)^2)^(1/2), x)

$$3.26 \quad \int \frac{1}{\sqrt{a \tanh^2(x)}} dx$$

Optimal. Leaf size=16

$$\frac{\log(\sinh(x)) \tanh(x)}{\sqrt{a \tanh^2(x)}}$$

[Out] $\ln(\sinh(x)) \cdot \tanh(x) / (a \cdot \tanh(x)^2)^{(1/2)}$

Rubi [A]

time = 0.01, antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3739, 3556}

$$\frac{\tanh(x) \log(\sinh(x))}{\sqrt{a \tanh^2(x)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[1/\text{Sqrt}[a \cdot \text{Tanh}[x]^2], x]$

[Out] $(\text{Log}[\text{Sinh}[x]] \cdot \text{Tanh}[x]) / \text{Sqrt}[a \cdot \text{Tanh}[x]^2]$

Rule 3556

$\text{Int}[\tan[(c_.) + (d_.)(x_.)], x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d \cdot x], x]]/d, x] /;$ $\text{FreeQ}\{c, d\}, x]$

Rule 3739

$\text{Int}[(u_.) \cdot ((b_.) \cdot \tan[(e_.) + (f_.)(x_.)]^{(n_.)})^{(p_.)}, x_Symbol] \rightarrow \text{With}\{\{ff = \text{FreeFactors}[\text{Tan}[e + f \cdot x], x]\}, \text{Dist}[(b \cdot ff^n)^{\text{IntPart}[p]} \cdot ((b \cdot \text{Tan}[e + f \cdot x]^n)^{\text{FracPart}[p]} / (\text{Tan}[e + f \cdot x] / ff)^{(n \cdot \text{FracPart}[p])}), \text{Int}[\text{ActivateTrig}[u] \cdot (\text{Tan}[e + f \cdot x] / ff)^{(n \cdot p)}, x], x]\} /;$ $\text{FreeQ}\{b, e, f, n, p\}, x \ \&\& \ !\text{IntegerQ}[p] \ \&\& \ \text{IntegerQ}[n] \ \&\& \ (\text{EqQ}[u, 1] \ || \ \text{MatchQ}[u, ((d_.) \cdot (\text{trig}_)[e + f \cdot x])^{(m_.)}]) /;$ $\text{FreeQ}\{d, m\}, x \ \&\& \ \text{MemberQ}\{\{\sin, \cos, \tan, \cot, \sec, \csc\}, \text{trig}\}]$

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{a \tanh^2(x)}} dx &= \frac{\tanh(x) \int \coth(x) dx}{\sqrt{a \tanh^2(x)}} \\ &= \frac{\log(\sinh(x)) \tanh(x)}{\sqrt{a \tanh^2(x)}} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 16, normalized size = 1.00

$$\frac{\log(\sinh(x)) \tanh(x)}{\sqrt{a \tanh^2(x)}}$$

Antiderivative was successfully verified.

`[In] Integrate[1/Sqrt[a*Tanh[x]^2], x]``[Out] (Log[Sinh[x]]*Tanh[x])/Sqrt[a*Tanh[x]^2]`**Maple [A]**

time = 0.79, size = 29, normalized size = 1.81

method	result	size
derivativedivides	$-\frac{\tanh(x)(\ln(1+\tanh(x))+\ln(\tanh(x)-1)-2\ln(\tanh(x)))}{2\sqrt{a(\tanh^2(x))}}$	29
default	$-\frac{\tanh(x)(\ln(1+\tanh(x))+\ln(\tanh(x)-1)-2\ln(\tanh(x)))}{2\sqrt{a(\tanh^2(x))}}$	29
risch	$-\frac{(e^{2x}-1)x}{\sqrt{\frac{a(e^{2x}-1)^2}{(1+e^{2x})^2}}(1+e^{2x})} + \frac{(e^{2x}-1)\ln(e^{2x}-1)}{\sqrt{\frac{a(e^{2x}-1)^2}{(1+e^{2x})^2}}(1+e^{2x})}$	81

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/(a*tanh(x)^2)^(1/2), x, method=_RETURNVERBOSE)``[Out] -1/2*tanh(x)*(ln(1+tanh(x))+ln(tanh(x)-1)-2*ln(tanh(x)))/(a*tanh(x)^2)^(1/2)`**Maxima [B]** Leaf count of result is larger than twice the leaf count of optimal. 31 vs. 2(14) = 28.

time = 0.49, size = 31, normalized size = 1.94

$$-\frac{x}{\sqrt{a}} - \frac{\log(e^{-x} + 1)}{\sqrt{a}} - \frac{\log(e^{-x} - 1)}{\sqrt{a}}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/(a*tanh(x)^2)^(1/2), x, algorithm="maxima")``[Out] -x/sqrt(a) - log(e^(-x) + 1)/sqrt(a) - log(e^(-x) - 1)/sqrt(a)`**Fricas [B]** Leaf count of result is larger than twice the leaf count of optimal. 76 vs. 2(14) = 28.

time = 0.40, size = 76, normalized size = 4.75

$$\frac{\left(xe^{(2x)} - (e^{(2x)} + 1) \log\left(\frac{2 \sinh(x)}{\cosh(x) - \sinh(x)}\right) + x\right) \sqrt{\frac{ae^{(4x)} - 2ae^{(2x)} + a}{e^{(4x)} + 2e^{(2x)} + 1}}}{ae^{(2x)} - a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*tanh(x)^2)^(1/2),x, algorithm="fricas")

[Out] -(x*e^(2*x) - (e^(2*x) + 1)*log(2*sinh(x)/(cosh(x) - sinh(x))) + x)*sqrt((a*e^(4*x) - 2*a*e^(2*x) + a)/(e^(4*x) + 2*e^(2*x) + 1))/(a*e^(2*x) - a)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{a \tanh^2(x)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*tanh(x)**2)**(1/2),x)

[Out] Integral(1/sqrt(a*tanh(x)**2), x)

Giac [A]

time = 0.42, size = 1, normalized size = 0.06

0

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*tanh(x)^2)^(1/2),x, algorithm="giac")

[Out] 0

Mupad [B]

time = 1.20, size = 14, normalized size = 0.88

$$\frac{\operatorname{atanh}\left(\frac{\tanh(x)}{\sqrt{\tanh(x)^2}}\right)}{\sqrt{a}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a*tanh(x)^2)^(1/2),x)

[Out] atanh(tanh(x)/(tanh(x)^2)^(1/2))/a^(1/2)

3.27 $\int (-\tanh^2(c + dx))^{5/2} dx$

Optimal. Leaf size=88

$$\frac{\coth(c + dx) \log(\cosh(c + dx)) \sqrt{-\tanh^2(c + dx)}}{d} - \frac{\tanh(c + dx) \sqrt{-\tanh^2(c + dx)}}{2d} - \frac{\tanh^3(c + dx) \sqrt{-\tanh^2(c + dx)}}{4d}$$

[Out] $\coth(d*x+c)*\ln(\cosh(d*x+c))*(-\tanh(d*x+c)^2)^{(1/2)}/d-1/2*(-\tanh(d*x+c)^2)^{(1/2)}*\tanh(d*x+c)/d-1/4*(-\tanh(d*x+c)^2)^{(1/2)}*\tanh(d*x+c)^3/d$

Rubi [A]

time = 0.04, antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {3739, 3554, 3556}

$$-\frac{\sqrt{-\tanh^2(c + dx)} \tanh(c + dx)}{2d} - \frac{\sqrt{-\tanh^2(c + dx)} \tanh^3(c + dx)}{4d} + \frac{\sqrt{-\tanh^2(c + dx)} \coth(c + dx) \log(\cosh(c + dx))}{d}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(-\text{Tanh}[c + d*x]^2)^{(5/2)}, x]$

[Out] $(\text{Coth}[c + d*x]*\text{Log}[\text{Cosh}[c + d*x]]*\text{Sqrt}[-\text{Tanh}[c + d*x]^2])/d - (\text{Tanh}[c + d*x]*\text{Sqrt}[-\text{Tanh}[c + d*x]^2])/(2*d) - (\text{Tanh}[c + d*x]^3*\text{Sqrt}[-\text{Tanh}[c + d*x]^2])/(4*d)$

Rule 3554

$\text{Int}[(b*.)*\tan[(c*.) + (d*.)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Simp}[b*((b*\text{Tan}[c + d*x])^{(n - 1)}/(d*(n - 1))), x] - \text{Dist}[b^2, \text{Int}[(b*\text{Tan}[c + d*x])^{(n - 2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x \ \&\& \ \text{GtQ}[n, 1]$

Rule 3556

$\text{Int}[\tan[(c*.) + (d*.)*(x_)], x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] /; \text{FreeQ}\{c, d\}, x]$

Rule 3739

$\text{Int}[(u*.)*((b*.)*\tan[(e*.) + (f*.)*(x_)]^{(n_)})^{(p_)}, x_Symbol] \rightarrow \text{With}\{\{ff = \text{FreeFactors}[\text{Tan}[e + f*x], x]\}, \text{Dist}[(b*ff^n)^{\text{IntPart}[p]}*((b*\text{Tan}[e + f*x])^n)^{\text{FracPart}[p]}/(\text{Tan}[e + f*x]/ff)^{(n*\text{FracPart}[p])}], \text{Int}[\text{ActivateTrig}[u]*(\text{Tan}[e + f*x]/ff)^{(n*p)}, x], x] /; \text{FreeQ}\{b, e, f, n, p\}, x \ \&\& \ !\text{IntegerQ}[p] \ \&\& \ \text{IntegerQ}[n] \ \&\& \ (\text{EqQ}[u, 1] \ || \ \text{MatchQ}[u, ((d*.)*(\text{trig}_)[e + f*x])^{(m_)}]) /; \text{FreeQ}\{d, m\}, x \ \&\& \ \text{MemberQ}\{\{\sin, \cos, \tan, \cot, \sec, \csc\}, \text{trig}\}]$

Rubi steps

$$\begin{aligned}
\int (-\tanh^2(c+dx))^{5/2} dx &= \left(\coth(c+dx) \sqrt{-\tanh^2(c+dx)} \right) \int \tanh^5(c+dx) dx \\
&= -\frac{\tanh^3(c+dx) \sqrt{-\tanh^2(c+dx)}}{4d} + \left(\coth(c+dx) \sqrt{-\tanh^2(c+dx)} \right) \int \tanh^4(c+dx) dx \\
&= -\frac{\tanh(c+dx) \sqrt{-\tanh^2(c+dx)}}{2d} - \frac{\tanh^3(c+dx) \sqrt{-\tanh^2(c+dx)}}{4d} + \left(\coth(c+dx) \sqrt{-\tanh^2(c+dx)} \right) \int \tanh^3(c+dx) dx \\
&= \frac{\coth(c+dx) \log(\cosh(c+dx)) \sqrt{-\tanh^2(c+dx)}}{d} - \frac{\tanh(c+dx) \sqrt{-\tanh^2(c+dx)}}{2d}
\end{aligned}$$

Mathematica [A]

time = 0.21, size = 56, normalized size = 0.64

$$\frac{\coth(c+dx) (-1 - 2 \coth^2(c+dx) + 4 \coth^4(c+dx) \log(\cosh(c+dx))) (-\tanh^2(c+dx))^{5/2}}{4d}$$

Antiderivative was successfully verified.

`[In] Integrate[(-Tanh[c + d*x]^2)^(5/2), x]``[Out] (Coth[c + d*x]*(-1 - 2*Coth[c + d*x]^2 + 4*Coth[c + d*x]^4*Log[Cosh[c + d*x]])*(-Tanh[c + d*x]^2)^(5/2))/(4*d)`**Maple [A]**

time = 1.46, size = 67, normalized size = 0.76

method	result
derivativdivides	$-\frac{(-(\tanh^2(dx+c)))^{5/2} (\tanh^4(dx+c)+2(\tanh^2(dx+c))+2 \ln(\tanh(dx+c)-1)+2 \ln(\tanh(dx+c)+1))}{4d \tanh(dx+c)^5}$
default	$-\frac{(-(\tanh^2(dx+c)))^{5/2} (\tanh^4(dx+c)+2(\tanh^2(dx+c))+2 \ln(\tanh(dx+c)-1)+2 \ln(\tanh(dx+c)+1))}{4d \tanh(dx+c)^5}$
risch	$\frac{(1+e^{2dx+2c}) \sqrt{-\frac{(e^{2dx+2c}-1)^2}{(1+e^{2dx+2c})^2}}}{e^{2dx+2c}-1} x - \frac{2(1+e^{2dx+2c}) \sqrt{-\frac{(e^{2dx+2c}-1)^2}{(1+e^{2dx+2c})^2}} (dx+c)}{(e^{2dx+2c}-1)d} + \frac{4 \sqrt{-\frac{(e^{2dx+2c}-1)^2}{(1+e^{2dx+2c})^2}}}{(e^{2dx+2c}-1)(1+e^{2dx+2c})}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((-tanh(d*x+c)^2)^(5/2), x, method=_RETURNVERBOSE)``[Out] -1/4/d*(-tanh(d*x+c)^2)^(5/2)*(tanh(d*x+c)^4+2*tanh(d*x+c)^2+2*ln(tanh(d*x+c)-1)+2*ln(tanh(d*x+c)+1))/tanh(d*x+c)^5`

Maxima [C] Result contains complex when optimal does not.

time = 0.50, size = 113, normalized size = 1.28

$$\frac{i(dx+c)}{d} - \frac{i \log(e^{(-2dx-2c)} + 1)}{d} + \frac{4(-ie^{(-2dx-2c)} - ie^{(-4dx-4c)} - ie^{(-6dx-6c)})}{d(4e^{(-2dx-2c)} + 6e^{(-4dx-4c)} + 4e^{(-6dx-6c)} + e^{(-8dx-8c)} + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-tanh(d*x+c)^2)^(5/2),x, algorithm="maxima")

[Out] -I*(d*x + c)/d - I*log(e^(-2*d*x - 2*c) + 1)/d + 4*(-I*e^(-2*d*x - 2*c) - I*e^(-4*d*x - 4*c) - I*e^(-6*d*x - 6*c))/(d*(4*e^(-2*d*x - 2*c) + 6*e^(-4*d*x - 4*c) + 4*e^(-6*d*x - 6*c) + e^(-8*d*x - 8*c) + 1))

Fricas [C] Result contains complex when optimal does not.

time = 0.38, size = 180, normalized size = 2.05

$$\frac{-idxe^{(8dx+8c)} - idx - 4(dx-i)e^{(6dx+6c)} - 2(3idx-2i)e^{(4dx+4c)} - 4(dx-i)e^{(2dx+2c)} + (ie^{(8dx+8c)} + 4ie^{(6dx+6c)} + 6ie^{(4dx+4c)} + 4ie^{(2dx+2c)} + i) \log(e^{(2dx+2c)} + 1)}{de^{(8dx+8c)} + 4de^{(6dx+6c)} + 6de^{(4dx+4c)} + 4de^{(2dx+2c)} + d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-tanh(d*x+c)^2)^(5/2),x, algorithm="fricas")

[Out] (-I*d*x*e^(8*d*x + 8*c) - I*d*x - 4*(I*d*x - I)*e^(6*d*x + 6*c) - 2*(3*I*d*x - 2*I)*e^(4*d*x + 4*c) - 4*(I*d*x - I)*e^(2*d*x + 2*c) + (I*e^(8*d*x + 8*c) + 4*I*e^(6*d*x + 6*c) + 6*I*e^(4*d*x + 4*c) + 4*I*e^(2*d*x + 2*c) + I)*1*log(e^(2*d*x + 2*c) + 1))/(d*e^(8*d*x + 8*c) + 4*d*e^(6*d*x + 6*c) + 6*d*e^(4*d*x + 4*c) + 4*d*e^(2*d*x + 2*c) + d)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (-\tanh^2(c + dx))^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-tanh(d*x+c)**2)**(5/2),x)

[Out] Integral((-tanh(c + d*x)**2)**(5/2), x)

Giac [C] Result contains complex when optimal does not.

time = 0.45, size = 142, normalized size = 1.61

$$\frac{i(dx+c)\operatorname{sgn}(-e^{(4dx+4c)} + 1) - i \log(e^{(2dx+2c)} + 1) \operatorname{sgn}(-e^{(4dx+4c)} + 1) - \frac{4i(e^{(6dx+6c)} \operatorname{sgn}(-e^{(4dx+4c)} + 1) + e^{(4dx+4c)} \operatorname{sgn}(-e^{(4dx+4c)} + 1) + e^{(2dx+2c)} \operatorname{sgn}(-e^{(4dx+4c)} + 1))}{(e^{(2dx+2c)} + 1)^4}}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-tanh(d*x+c)^2)^(5/2),x, algorithm="giac")

```
[Out] (I*(d*x + c)*sgn(-e^(4*d*x + 4*c) + 1) - I*log(e^(2*d*x + 2*c) + 1)*sgn(-e^(4*d*x + 4*c) + 1) - 4*I*(e^(6*d*x + 6*c)*sgn(-e^(4*d*x + 4*c) + 1) + e^(4*d*x + 4*c)*sgn(-e^(4*d*x + 4*c) + 1) + e^(2*d*x + 2*c)*sgn(-e^(4*d*x + 4*c) + 1))/(e^(2*d*x + 2*c) + 1)^4/d
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int (-\tanh(c + dx)^2)^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((-tanh(c + d*x)^2)^(5/2), x)
```

```
[Out] int((-tanh(c + d*x)^2)^(5/2), x)
```

3.28 $\int (-\tanh^2(c + dx))^{3/2} dx$

Optimal. Leaf size=60

$$-\frac{\coth(c + dx) \log(\cosh(c + dx)) \sqrt{-\tanh^2(c + dx)}}{d} + \frac{\tanh(c + dx) \sqrt{-\tanh^2(c + dx)}}{2d}$$

[Out] $-\coth(d*x+c)*\ln(\cosh(d*x+c))*(-\tanh(d*x+c)^2)^{(1/2)}/d+1/2*(-\tanh(d*x+c)^2)^{(1/2)}*\tanh(d*x+c)/d$

Rubi [A]

time = 0.02, antiderivative size = 60, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {3739, 3554, 3556}

$$\frac{\tanh(c + dx) \sqrt{-\tanh^2(c + dx)}}{2d} - \frac{\sqrt{-\tanh^2(c + dx)} \coth(c + dx) \log(\cosh(c + dx))}{d}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(-\text{Tanh}[c + d*x]^2)^{(3/2)}, x]$

[Out] $-\left(\frac{\text{Coth}[c + d*x] \text{Log}[\text{Cosh}[c + d*x]] \text{Sqrt}[-\text{Tanh}[c + d*x]^2]}{d}\right) + \frac{\text{Tanh}[c + d*x] \text{Sqrt}[-\text{Tanh}[c + d*x]^2]}{(2*d)}$

Rule 3554

$\text{Int}[(b \cdot \tan(c) + d \cdot x)^n, x_Symbol] \rightarrow \text{Simp}[b \cdot (b \cdot \tan(c + d \cdot x))^n, x] - \text{Dist}[b^2, \text{Int}[(b \cdot \tan(c + d \cdot x))^{n-2}, x], x] /;$ FreeQ[{b, c, d}, x] && GtQ[n, 1]

Rule 3556

$\text{Int}[\tan(c) + d \cdot x, x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d \cdot x], x]]/d, x] /;$ FreeQ[{c, d}, x]

Rule 3739

$\text{Int}[u \cdot (b \cdot \tan(e) + f \cdot x)^n, x_Symbol] \rightarrow \text{With}[\{ff = \text{FreeFactors}[\tan[e + f \cdot x], x]\}, \text{Dist}[(b \cdot ff^n)^{\text{IntPart}[p]} \cdot (b \cdot \tan[e + f \cdot x])^{\text{FracPart}[p]} / (\tan[e + f \cdot x] / ff)^{n \cdot \text{FracPart}[p]}], \text{Int}[\text{ActivateTrig}[u] \cdot (\tan[e + f \cdot x] / ff)^{n \cdot p}, x], x] /;$ FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d \cdot x) \cdot (trig_)[e + f \cdot x])^m] /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])

Rubi steps

$$\begin{aligned}
\int (-\tanh^2(c+dx))^{3/2} dx &= -\left(\left(\coth(c+dx) \sqrt{-\tanh^2(c+dx)} \right) \int \tanh^3(c+dx) dx \right) \\
&= \frac{\tanh(c+dx) \sqrt{-\tanh^2(c+dx)}}{2d} - \left(\coth(c+dx) \sqrt{-\tanh^2(c+dx)} \right) \int \tanh^2(c+dx) dx \\
&= -\frac{\coth(c+dx) \log(\cosh(c+dx)) \sqrt{-\tanh^2(c+dx)}}{d} + \frac{\tanh(c+dx) \sqrt{-\tanh^2(c+dx)}}{2d}
\end{aligned}$$

Mathematica [A]

time = 0.07, size = 46, normalized size = 0.77

$$\frac{\coth(c+dx) (-1 + 2 \coth^2(c+dx) \log(\cosh(c+dx))) (-\tanh^2(c+dx))^{3/2}}{2d}$$

Antiderivative was successfully verified.

`[In] Integrate[(-Tanh[c + d*x]^2)^(3/2), x]``[Out] (Coth[c + d*x]*(-1 + 2*Coth[c + d*x]^2*Log[Cosh[c + d*x]])*(-Tanh[c + d*x]^2)^(3/2))/(2*d)`**Maple [A]**

time = 1.33, size = 53, normalized size = 0.88

method	result
derivativedivides	$-\frac{(-\tanh^2(dx+c))^{3/2} (\tanh^2(dx+c) + \ln(\tanh(dx+c)-1) + \ln(\tanh(dx+c)+1))}{2d \tanh(dx+c)^3}$
default	$-\frac{(-\tanh^2(dx+c))^{3/2} (\tanh^2(dx+c) + \ln(\tanh(dx+c)-1) + \ln(\tanh(dx+c)+1))}{2d \tanh(dx+c)^3}$
risch	$-\frac{(1+e^{2dx+2c}) \sqrt{-\frac{(e^{2dx+2c}-1)^2}{(1+e^{2dx+2c})^2}}}{e^{2dx+2c}-1} x + \frac{2(1+e^{2dx+2c}) \sqrt{-\frac{(e^{2dx+2c}-1)^2}{(1+e^{2dx+2c})^2}}}{(e^{2dx+2c}-1)d} (dx+c) - \frac{2 \sqrt{-\frac{(e^{2dx+2c}-1)^2}{(1+e^{2dx+2c})^2}}}{(e^{2dx+2c}-1)(1+e^{2dx+2c})} e^{2dx}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((-tanh(d*x+c)^2)^(3/2), x, method=_RETURNVERBOSE)``[Out] -1/2/d*(-tanh(d*x+c)^2)^(3/2)*(tanh(d*x+c)^2+ln(tanh(d*x+c)-1)+ln(tanh(d*x+c)+1))/tanh(d*x+c)^3`**Maxima [C]** Result contains complex when optimal does not.

time = 0.51, size = 66, normalized size = 1.10

$$\frac{i(dx+c)}{d} + \frac{i \log(e^{(-2dx-2c)} + 1)}{d} + \frac{2i e^{(-2dx-2c)}}{d(2e^{(-2dx-2c)} + e^{(-4dx-4c)} + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-tanh(d*x+c)^2)^(3/2),x, algorithm="maxima")

[Out] $I*(d*x + c)/d + I*\log(e^{(-2*d*x - 2*c)} + 1)/d + 2*I*e^{(-2*d*x - 2*c)}/(d*(2*e^{(-2*d*x - 2*c)} + e^{(-4*d*x - 4*c)} + 1))$

Fricas [C] Result contains complex when optimal does not.

time = 0.39, size = 100, normalized size = 1.67

$$\frac{i dx e^{(4 dx+4 c)} + i dx - 2(-i dx + i) e^{(2 dx+2 c)} + (-i e^{(4 dx+4 c)} - 2i e^{(2 dx+2 c)} - i) \log(e^{(2 dx+2 c)} + 1)}{d e^{(4 dx+4 c)} + 2 d e^{(2 dx+2 c)} + d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-tanh(d*x+c)^2)^(3/2),x, algorithm="fricas")

[Out] $(I*d*x*e^{(4*d*x + 4*c)} + I*d*x - 2*(-I*d*x + I)*e^{(2*d*x + 2*c)} + (-I*e^{(4*d*x + 4*c)} - 2*I*e^{(2*d*x + 2*c)} - I)*\log(e^{(2*d*x + 2*c)} + 1))/(d*e^{(4*d*x + 4*c)} + 2*d*e^{(2*d*x + 2*c)} + d)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (-\tanh^2(c + dx))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-tanh(d*x+c)**2)**(3/2),x)

[Out] Integral((-tanh(c + d*x)**2)**(3/2), x)

Giac [C] Result contains complex when optimal does not.

time = 0.42, size = 92, normalized size = 1.53

$$\frac{-i(dx + c)\operatorname{sgn}(-e^{(4 dx+4 c)} + 1) + i \log(e^{(2 dx+2 c)} + 1) \operatorname{sgn}(-e^{(4 dx+4 c)} + 1) + \frac{2i e^{(2 dx+2 c)} \operatorname{sgn}(-e^{(4 dx+4 c)} + 1)}{(e^{(2 dx+2 c)} + 1)^2}}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-tanh(d*x+c)^2)^(3/2),x, algorithm="giac")

[Out] $(-I*(d*x + c)*\operatorname{sgn}(-e^{(4*d*x + 4*c)} + 1) + I*\log(e^{(2*d*x + 2*c)} + 1)*\operatorname{sgn}(-e^{(4*d*x + 4*c)} + 1) + 2*I*e^{(2*d*x + 2*c)}*\operatorname{sgn}(-e^{(4*d*x + 4*c)} + 1))/(e^{(2*d*x + 2*c)} + 1)^2/d$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int (-\tanh(c + dx)^2)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((-tanh(c + d*x)^2)^(3/2),x)
```

```
[Out] int((-tanh(c + d*x)^2)^(3/2), x)
```

3.29 $\int \sqrt{-\tanh^2(c+dx)} dx$

Optimal. Leaf size=31

$$\frac{\coth(c+dx) \log(\cosh(c+dx)) \sqrt{-\tanh^2(c+dx)}}{d}$$

[Out] $\coth(d*x+c)*\ln(\cosh(d*x+c))*(-\tanh(d*x+c)^2)^{(1/2)}/d$

Rubi [A]

time = 0.01, antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3739, 3556}

$$\frac{\sqrt{-\tanh^2(c+dx)} \coth(c+dx) \log(\cosh(c+dx))}{d}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sqrt}[-\text{Tanh}[c + d*x]^2], x]$

[Out] $(\text{Coth}[c + d*x]*\text{Log}[\text{Cosh}[c + d*x]]*\text{Sqrt}[-\text{Tanh}[c + d*x]^2])/d$

Rule 3556

$\text{Int}[\tan[(c_.) + (d_.)*(x_.)], x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] /; \text{FreeQ}\{c, d\}, x]$

Rule 3739

$\text{Int}[(u_.)*((b_.)*\tan[(e_.) + (f_.)*(x_.)]^{(n_.)})^{(p_.)}, x_Symbol] \rightarrow \text{With}\{\{ff = \text{FreeFactors}[\text{Tan}[e + f*x], x]\}, \text{Dist}[(b*ff^n)^{\text{IntPart}[p]}*(b*\text{Tan}[e + f*x]^n)^{\text{FracPart}[p]} / (\text{Tan}[e + f*x]/ff)^{(n*\text{FracPart}[p])}], \text{Int}[\text{ActivateTrig}[u]*(\text{Tan}[e + f*x]/ff)^{(n*p)}, x], x]\} /; \text{FreeQ}\{b, e, f, n, p\}, x \ \&\& \ !\text{IntegerQ}[p] \ \&\& \ \text{IntegerQ}[n] \ \&\& \ (\text{EqQ}[u, 1] \ || \ \text{MatchQ}[u, ((d_.)*(trig_)[e + f*x])^{(m_.)}] /; \text{FreeQ}\{d, m\}, x] \ \&\& \ \text{MemberQ}\{\{\sin, \cos, \tan, \cot, \sec, \csc\}, \text{trig}\})$

Rubi steps

$$\begin{aligned} \int \sqrt{-\tanh^2(c+dx)} dx &= \left(\coth(c+dx) \sqrt{-\tanh^2(c+dx)} \right) \int \tanh(c+dx) dx \\ &= \frac{\coth(c+dx) \log(\cosh(c+dx)) \sqrt{-\tanh^2(c+dx)}}{d} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 31, normalized size = 1.00

$$\frac{\coth(c + dx) \log(\cosh(c + dx)) \sqrt{-\tanh^2(c + dx)}}{d}$$

Antiderivative was successfully verified.

`[In] Integrate[Sqrt[-Tanh[c + d*x]^2], x]``[Out] (Coth[c + d*x]*Log[Cosh[c + d*x]]*Sqrt[-Tanh[c + d*x]^2])/d`**Maple [A]**

time = 1.45, size = 45, normalized size = 1.45

method	result
derivativedivides	$-\frac{\sqrt{-(\tanh^2(dx+c))} (\ln(\tanh(dx+c)-1)+\ln(\tanh(dx+c)+1))}{2d \tanh(dx+c)}$
default	$-\frac{\sqrt{-(\tanh^2(dx+c))} (\ln(\tanh(dx+c)-1)+\ln(\tanh(dx+c)+1))}{2d \tanh(dx+c)}$
risch	$\frac{(1+e^{2dx+2c}) \sqrt{-\frac{(e^{2dx+2c}-1)^2}{(1+e^{2dx+2c})^2}}}{e^{2dx+2c}-1} x - \frac{2(1+e^{2dx+2c}) \sqrt{-\frac{(e^{2dx+2c}-1)^2}{(1+e^{2dx+2c})^2}}}{(e^{2dx+2c}-1)d} (dx+c) + \frac{(1+e^{2dx+2c}) \sqrt{-\frac{(e^{2dx+2c}-1)^2}{(1+e^{2dx+2c})^2}}}{(e^{2dx+2c}-1)}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((-tanh(d*x+c)^2)^(1/2), x, method=_RETURNVERBOSE)``[Out] -1/2/d*(-tanh(d*x+c)^2)^(1/2)*(ln(tanh(d*x+c)-1)+ln(tanh(d*x+c)+1))/tanh(d*x+c)`**Maxima [C]** Result contains complex when optimal does not.

time = 0.50, size = 28, normalized size = 0.90

$$-\frac{i(dx+c)}{d} - \frac{i \log(e^{-2dx-2c} + 1)}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((-tanh(d*x+c)^2)^(1/2), x, algorithm="maxima")``[Out] -I*(d*x + c)/d - I*log(e^(-2*d*x - 2*c) + 1)/d`**Fricas [C]** Result contains complex when optimal does not.

time = 0.36, size = 23, normalized size = 0.74

$$\frac{-i dx + i \log(e^{(2dx+2c)} + 1)}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-tanh(d*x+c)^2)^(1/2),x, algorithm="fricas")

[Out] $(-I*d*x + I*\log(e^{(2*d*x + 2*c)} + 1))/d$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{-\tanh^2(c + dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-tanh(d*x+c)**2)**(1/2),x)

[Out] Integral(sqrt(-tanh(c + d*x)**2), x)

Giac [C] Result contains complex when optimal does not.

time = 0.42, size = 54, normalized size = 1.74

$$\frac{i(dx + c)\operatorname{sgn}(-e^{(4dx+4c)} + 1) - i\log(e^{(2dx+2c)} + 1)\operatorname{sgn}(-e^{(4dx+4c)} + 1)}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-tanh(d*x+c)^2)^(1/2),x, algorithm="giac")

[Out] $(I*(d*x + c)*\operatorname{sgn}(-e^{(4*d*x + 4*c)} + 1) - I*\log(e^{(2*d*x + 2*c)} + 1)*\operatorname{sgn}(-e^{(4*d*x + 4*c)} + 1))/d$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.03

$$\int \sqrt{-\tanh(c + dx)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((-tanh(c + d*x)^2)^(1/2),x)

[Out] int((-tanh(c + d*x)^2)^(1/2), x)

$$3.30 \quad \int \frac{1}{\sqrt{-\tanh^2(c+dx)}} dx$$

Optimal. Leaf size=31

$$\frac{\log(\sinh(c+dx)) \tanh(c+dx)}{d\sqrt{-\tanh^2(c+dx)}}$$

[Out] $\ln(\sinh(d*x+c))*\tanh(d*x+c)/d/(-\tanh(d*x+c)^2)^{(1/2)}$

Rubi [A]

time = 0.01, antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3739, 3556}

$$\frac{\tanh(c+dx) \log(\sinh(c+dx))}{d\sqrt{-\tanh^2(c+dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[1/\text{Sqrt}[-\text{Tanh}[c + d*x]^2], x]$

[Out] $(\text{Log}[\text{Sinh}[c + d*x]]*\text{Tanh}[c + d*x])/(d*\text{Sqrt}[-\text{Tanh}[c + d*x]^2])$

Rule 3556

$\text{Int}[\tan[(c_.) + (d_.)*(x_.)], x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] /; \text{FreeQ}\{c, d\}, x]$

Rule 3739

$\text{Int}[(u_.)*((b_.)*\tan[(e_.) + (f_.)*(x_.)]^{(n_.)})^{(p_.)}, x_Symbol] \rightarrow \text{With}\{\{ff = \text{FreeFactors}[\text{Tan}[e + f*x], x]\}, \text{Dist}[(b*ff^n)^{\text{IntPart}[p]}*((b*\text{Tan}[e + f*x]^n)^{\text{FracPart}[p]})/(\text{Tan}[e + f*x]/ff)^{(n*\text{FracPart}[p])}], \text{Int}[\text{ActivateTrig}[u]*(\text{Tan}[e + f*x]/ff)^{(n*p)}, x], x]\} /; \text{FreeQ}\{b, e, f, n, p\}, x \ \&\& \ !\text{IntegerQ}[p] \ \&\& \ \text{IntegerQ}[n] \ \&\& \ (\text{EqQ}[u, 1] \ || \ \text{MatchQ}[u, ((d_.)*(trig_)[e + f*x])^{(m_.)}] /; \text{FreeQ}\{d, m\}, x \ \&\& \ \text{MemberQ}\{\{\sin, \cos, \tan, \cot, \sec, \csc\}, \text{trig}\})$

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{-\tanh^2(c+dx)}} dx &= \frac{\tanh(c+dx) \int \coth(c+dx) dx}{\sqrt{-\tanh^2(c+dx)}} \\ &= \frac{\log(\sinh(c+dx)) \tanh(c+dx)}{d\sqrt{-\tanh^2(c+dx)}} \end{aligned}$$

Mathematica [A]

time = 0.06, size = 39, normalized size = 1.26

$$\frac{(\log(\cosh(c + dx)) + \log(\tanh(c + dx))) \tanh(c + dx)}{d\sqrt{-\tanh^2(c + dx)}}$$

Antiderivative was successfully verified.

`[In] Integrate[1/Sqrt[-Tanh[c + d*x]^2], x]``[Out] ((Log[Cosh[c + d*x]] + Log[Tanh[c + d*x]])*Tanh[c + d*x])/(d*Sqrt[-Tanh[c + d*x]^2])`**Maple [A]**

time = 1.64, size = 56, normalized size = 1.81

method	result
derivativedivides	$\frac{\tanh(dx+c)(2\ln(\tanh(dx+c))-\ln(\tanh(dx+c)-1)-\ln(\tanh(dx+c)+1))}{2d\sqrt{-(\tanh^2(dx+c))}}$
default	$\frac{\tanh(dx+c)(2\ln(\tanh(dx+c))-\ln(\tanh(dx+c)-1)-\ln(\tanh(dx+c)+1))}{2d\sqrt{-(\tanh^2(dx+c))}}$
risch	$\frac{(e^{2dx+2c}-1)x}{\sqrt{-\frac{(e^{2dx+2c}-1)^2}{(1+e^{2dx+2c})^2}}(1+e^{2dx+2c})} - \frac{2(e^{2dx+2c}-1)(dx+c)}{\sqrt{-\frac{(e^{2dx+2c}-1)^2}{(1+e^{2dx+2c})^2}}(1+e^{2dx+2c})d} + \frac{(e^{2dx+2c}-1)\ln(e^{2dx+2c}-1)}{\sqrt{-\frac{(e^{2dx+2c}-1)^2}{(1+e^{2dx+2c})^2}}(1+e^{2dx+2c})d}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/(-tanh(d*x+c)^2)^(1/2), x, method=_RETURNVERBOSE)``[Out] 1/2/d*tanh(d*x+c)*(2*ln(tanh(d*x+c))-ln(tanh(d*x+c)-1)-ln(tanh(d*x+c)+1))/(-tanh(d*x+c)^2)^(1/2)`**Maxima [C]** Result contains complex when optimal does not.

time = 0.48, size = 45, normalized size = 1.45

$$\frac{i(dx+c)}{d} + \frac{i \log(e^{-dx-c} + 1)}{d} + \frac{i \log(e^{-dx-c} - 1)}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/(-tanh(d*x+c)^2)^(1/2), x, algorithm="maxima")``[Out] I*(d*x + c)/d + I*log(e^(-d*x - c) + 1)/d + I*log(e^(-d*x - c) - 1)/d`**Fricas [C]** Result contains complex when optimal does not.

time = 0.42, size = 23, normalized size = 0.74

$$\frac{i dx - i \log(e^{2dx+2c} - 1)}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-tanh(d*x+c)^2)^(1/2),x, algorithm="fricas")

[Out] (I*d*x - I*log(e^(2*d*x + 2*c) - 1))/d

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{-\tanh^2(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-tanh(d*x+c)**2)**(1/2),x)

[Out] Integral(1/sqrt(-tanh(c + d*x)**2), x)

Giac [C] Result contains complex when optimal does not.

time = 0.41, size = 63, normalized size = 2.03

$$\frac{\frac{i dx + i c}{\operatorname{sgn}(-e^{(4 dx + 4 c)})} - \frac{i \log(-i e^{(2 dx + 2 c)} + i)}{\operatorname{sgn}(-e^{(4 dx + 4 c)})}}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-tanh(d*x+c)^2)^(1/2),x, algorithm="giac")

[Out] -((I*d*x + I*c)/sgn(-e^(4*d*x + 4*c) + 1) - I*log(-I*e^(2*d*x + 2*c) + I)/sgn(-e^(4*d*x + 4*c) + 1))/d

Mupad [B]

time = 1.22, size = 24, normalized size = 0.77

$$\frac{\operatorname{atan}\left(\frac{\tanh(c+dx)}{\sqrt{-\tanh(c+dx)^2}}\right)}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-tanh(c + d*x)^2)^(1/2),x)

[Out] atan(tanh(c + d*x)/(-tanh(c + d*x)^2)^(1/2))/d

$$3.31 \quad \int \frac{1}{(-\tanh^2(c+dx))^{3/2}} dx$$

Optimal. Leaf size=60

$$\frac{\coth(c+dx)}{2d\sqrt{-\tanh^2(c+dx)}} - \frac{\log(\sinh(c+dx))\tanh(c+dx)}{d\sqrt{-\tanh^2(c+dx)}}$$

[Out] 1/2*coth(d*x+c)/d/(-tanh(d*x+c)^2)^(1/2)-ln(sinh(d*x+c))*tanh(d*x+c)/d/(-tanh(d*x+c)^2)^(1/2)

Rubi [A]

time = 0.02, antiderivative size = 60, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {3739, 3554, 3556}

$$\frac{\coth(c+dx)}{2d\sqrt{-\tanh^2(c+dx)}} - \frac{\tanh(c+dx)\log(\sinh(c+dx))}{d\sqrt{-\tanh^2(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[(-Tanh[c + d*x]^2)^(-3/2), x]

[Out] Coth[c + d*x]/(2*d*Sqrt[-Tanh[c + d*x]^2]) - (Log[Sinh[c + d*x]]*Tanh[c + d*x])/(d*Sqrt[-Tanh[c + d*x]^2])

Rule 3554

Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Dist[b^2, Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]

Rule 3556

Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]

Rule 3739

Int[(u_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[(b*ff^n)^IntPart[p]*((b*Tan[e + f*x])^n)^FracPart[p]/(Tan[e + f*x]/ff)^(n*FracPart[p])], Int[ActivateTrig[u]*(Tan[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig]])

Rubi steps

$$\begin{aligned}
 \int \frac{1}{(-\tanh^2(c+dx))^{3/2}} dx &= -\frac{\tanh(c+dx) \int \coth^3(c+dx) dx}{\sqrt{-\tanh^2(c+dx)}} \\
 &= \frac{\coth(c+dx)}{2d\sqrt{-\tanh^2(c+dx)}} - \frac{\tanh(c+dx) \int \coth(c+dx) dx}{\sqrt{-\tanh^2(c+dx)}} \\
 &= \frac{\coth(c+dx)}{2d\sqrt{-\tanh^2(c+dx)}} - \frac{\log(\sinh(c+dx)) \tanh(c+dx)}{d\sqrt{-\tanh^2(c+dx)}}
 \end{aligned}$$

Mathematica [A]

time = 0.10, size = 51, normalized size = 0.85

$$\frac{\coth(c+dx) - 2(\log(\cosh(c+dx)) + \log(\tanh(c+dx))) \tanh(c+dx)}{2d\sqrt{-\tanh^2(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(-Tanh[c + d*x]^2)^(-3/2), x]

[Out] (Coth[c + d*x] - 2*(Log[Cosh[c + d*x]] + Log[Tanh[c + d*x]])*Tanh[c + d*x]) / (2*d*Sqrt[-Tanh[c + d*x]^2])

Maple [A]

time = 1.67, size = 81, normalized size = 1.35

method	result
derivativedivides	$\frac{\tanh(dx+c)(2\ln(\tanh(dx+c))(\tanh^2(dx+c)) - \ln(\tanh(dx+c)-1)(\tanh^2(dx+c)) - \ln(\tanh(dx+c)+1)(\tanh^2(dx+c)) - 1)}{2d(-\tanh^2(dx+c))^{\frac{3}{2}}}$
default	$\frac{\tanh(dx+c)(2\ln(\tanh(dx+c))(\tanh^2(dx+c)) - \ln(\tanh(dx+c)-1)(\tanh^2(dx+c)) - \ln(\tanh(dx+c)+1)(\tanh^2(dx+c)) - 1)}{2d(-\tanh^2(dx+c))^{\frac{3}{2}}}$
risch	$-\frac{(e^{2dx+2c}-1)x}{\sqrt{-\frac{(e^{2dx+2c}-1)^2}{(1+e^{2dx+2c})^2}}(1+e^{2dx+2c})} + \frac{2(e^{2dx+2c}-1)(dx+c)}{\sqrt{-\frac{(e^{2dx+2c}-1)^2}{(1+e^{2dx+2c})^2}}(1+e^{2dx+2c})d} + \frac{2e^{2dx+2c}}{(e^{2dx+2c}-1)(1+e^{2dx+2c})\sqrt{-\frac{(e^{2dx+2c}-1)^2}{(1+e^{2dx+2c})^2}}}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-tanh(d*x+c)^2)^(3/2), x, method=_RETURNVERBOSE)

[Out] 1/2/d*tanh(d*x+c)*(2*ln(tanh(d*x+c))*tanh(d*x+c)^2-ln(tanh(d*x+c)-1)*tanh(d*x+c)^2-ln(tanh(d*x+c)+1)*tanh(d*x+c)^2-1)/(-tanh(d*x+c)^2)^(3/2)

Maxima [C] Result contains complex when optimal does not.

time = 0.49, size = 85, normalized size = 1.42

$$\frac{i(dx+c)}{d} - \frac{i \log(e^{(-dx-c)} + 1)}{d} - \frac{i \log(e^{(-dx-c)} - 1)}{d} - \frac{2i e^{(-2dx-2c)}}{d(2e^{(-2dx-2c)} - e^{(-4dx-4c)} - 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-tanh(d*x+c)^2)^(3/2),x, algorithm="maxima")

[Out] -I*(d*x + c)/d - I*log(e^(-d*x - c) + 1)/d - I*log(e^(-d*x - c) - 1)/d - 2*I*e^(-2*d*x - 2*c)/(d*(2*e^(-2*d*x - 2*c) - e^(-4*d*x - 4*c) - 1))

Fricas [C] Result contains complex when optimal does not.

time = 0.42, size = 100, normalized size = 1.67

$$\frac{-i dx e^{(4dx+4c)} - i dx - 2(-i dx + i)e^{(2dx+2c)} + (i e^{(4dx+4c)} - 2i e^{(2dx+2c)} + i) \log(e^{(2dx+2c)} - 1)}{d e^{(4dx+4c)} - 2d e^{(2dx+2c)} + d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-tanh(d*x+c)^2)^(3/2),x, algorithm="fricas")

[Out] (-I*d*x*e^(4*d*x + 4*c) - I*d*x - 2*(-I*d*x + I)*e^(2*d*x + 2*c) + (I*e^(4*d*x + 4*c) - 2*I*e^(2*d*x + 2*c) + I)*log(e^(2*d*x + 2*c) - 1))/(d*e^(4*d*x + 4*c) - 2*d*e^(2*d*x + 2*c) + d)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(-\tanh^2(c + dx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-tanh(d*x+c)**2)**(3/2),x)

[Out] Integral((-tanh(c + d*x)**2)**(-3/2), x)

Giac [C] Result contains complex when optimal does not.

time = 0.42, size = 100, normalized size = 1.67

$$\frac{\frac{i dx + i c}{\operatorname{sgn}(-e^{(4 dx + 4 c)} + 1)} - \frac{i \log(e^{(2 dx + 2 c)} - 1)}{\operatorname{sgn}(-e^{(4 dx + 4 c)} + 1)} + \frac{2i e^{(2 dx + 2 c)}}{(e^{(2 dx + 2 c)} - 1)^2 \operatorname{sgn}(-e^{(4 dx + 4 c)} + 1)}}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-tanh(d*x+c)^2)^(3/2),x, algorithm="giac")

[Out] $((I*d*x + I*c)/\text{sgn}(-e^{(4*d*x + 4*c)} + 1) - I*\log(e^{(2*d*x + 2*c)} - 1)/\text{sgn}(-e^{(4*d*x + 4*c)} + 1) + 2*I*e^{(2*d*x + 2*c)}/((e^{(2*d*x + 2*c)} - 1)^2*\text{sgn}(-e^{(4*d*x + 4*c)} + 1)))/d$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{(-\tanh(c + dx)^2)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(-tanh(c + d*x)^2)^(3/2),x)`

[Out] `int(1/(-tanh(c + d*x)^2)^(3/2), x)`

3.32 $\int \frac{1}{(-\tanh^2(c+dx))^{5/2}} dx$

Optimal. Leaf size=88

$$-\frac{\coth(c+dx)}{2d\sqrt{-\tanh^2(c+dx)}} - \frac{\coth^3(c+dx)}{4d\sqrt{-\tanh^2(c+dx)}} + \frac{\log(\sinh(c+dx))\tanh(c+dx)}{d\sqrt{-\tanh^2(c+dx)}}$$

[Out] $-1/2*\coth(d*x+c)/d/(-\tanh(d*x+c)^2)^{(1/2)}-1/4*\coth(d*x+c)^3/d/(-\tanh(d*x+c)^2)^{(1/2)}+\ln(\sinh(d*x+c))*\tanh(d*x+c)/d/(-\tanh(d*x+c)^2)^{(1/2)}$

Rubi [A]

time = 0.03, antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {3739, 3554, 3556}

$$-\frac{\coth^3(c+dx)}{4d\sqrt{-\tanh^2(c+dx)}} - \frac{\coth(c+dx)}{2d\sqrt{-\tanh^2(c+dx)}} + \frac{\tanh(c+dx)\log(\sinh(c+dx))}{d\sqrt{-\tanh^2(c+dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(-\text{Tanh}[c + d*x]^2)^{-5/2}, x]$

[Out] $-1/2*\text{Coth}[c + d*x]/(d*\text{Sqrt}[-\text{Tanh}[c + d*x]^2]) - \text{Coth}[c + d*x]^3/(4*d*\text{Sqrt}[-\text{Tanh}[c + d*x]^2]) + (\text{Log}[\text{Sinh}[c + d*x]]*\text{Tanh}[c + d*x])/(d*\text{Sqrt}[-\text{Tanh}[c + d*x]^2])$

Rule 3554

$\text{Int}[(b_*)*\tan[(c_*) + (d_*)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Simp}[b*((b*\tan[c + d*x])^{(n-1)})/(d*(n-1))], x] - \text{Dist}[b^2, \text{Int}[(b*\tan[c + d*x])^{(n-2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x] \ \&\& \ \text{GtQ}[n, 1]$

Rule 3556

$\text{Int}[\tan[(c_*) + (d_*)*(x_)], x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] /; \text{FreeQ}\{c, d\}, x]$

Rule 3739

$\text{Int}[(u_)*((b_*)*\tan[(e_*) + (f_*)*(x_)]^{(n_)})^{(p_)}, x_Symbol] \rightarrow \text{With}\{\{ff = \text{FreeFactors}[\tan[e + f*x], x]\}, \text{Dist}[(b*ff^n)^{\text{IntPart}[p]}*((b*\tan[e + f*x])^{\text{FracPart}[p]}/(\tan[e + f*x]/ff)^{(n*\text{FracPart}[p])})], \text{Int}[\text{ActivateTrig}[u]*(\tan[e + f*x]/ff)^{(n*p)}, x], x]\} /; \text{FreeQ}\{b, e, f, n, p\}, x] \ \&\& \ !\text{IntegerQ}[p] \ \&\& \ \text{IntegerQ}[n] \ \&\& \ (\text{EqQ}[u, 1] \ || \ \text{MatchQ}[u, ((d_*)*(\text{trig}_)[e + f*x])^{(m_)}] /; \text{FreeQ}\{d, m\}, x] \ \&\& \ \text{MemberQ}\{\{\sin, \cos, \tan, \cot, \sec, \csc\}, \text{trig}\})$

Rubi steps

$$\begin{aligned}
\int \frac{1}{(-\tanh^2(c+dx))^{5/2}} dx &= \frac{\tanh(c+dx) \int \coth^5(c+dx) dx}{\sqrt{-\tanh^2(c+dx)}} \\
&= -\frac{\coth^3(c+dx)}{4d\sqrt{-\tanh^2(c+dx)}} + \frac{\tanh(c+dx) \int \coth^3(c+dx) dx}{\sqrt{-\tanh^2(c+dx)}} \\
&= -\frac{\coth(c+dx)}{2d\sqrt{-\tanh^2(c+dx)}} - \frac{\coth^3(c+dx)}{4d\sqrt{-\tanh^2(c+dx)}} + \frac{\tanh(c+dx) \int \coth(c+dx) dx}{\sqrt{-\tanh^2(c+dx)}} \\
&= -\frac{\coth(c+dx)}{2d\sqrt{-\tanh^2(c+dx)}} - \frac{\coth^3(c+dx)}{4d\sqrt{-\tanh^2(c+dx)}} + \frac{\log(\sinh(c+dx)) \tanh(c+dx)}{d\sqrt{-\tanh^2(c+dx)}}
\end{aligned}$$

Mathematica [A]

time = 0.18, size = 63, normalized size = 0.72

$$\frac{-2 \coth(c+dx) - \coth^3(c+dx) + 4(\log(\cosh(c+dx)) + \log(\tanh(c+dx))) \tanh(c+dx)}{4d\sqrt{-\tanh^2(c+dx)}}$$

Antiderivative was successfully verified.

`[In] Integrate[(-Tanh[c + d*x]^2)^(-5/2), x]`

```
[Out] (-2*Coth[c + d*x] - Coth[c + d*x]^3 + 4*(Log[Cosh[c + d*x]] + Log[Tanh[c + d*x]])*Tanh[c + d*x])/(4*d*Sqrt[-Tanh[c + d*x]^2])
```

Maple [A]

time = 1.74, size = 91, normalized size = 1.03

method	result
derivativedivides	$\frac{\tanh(dx+c)(4 \ln(\tanh(dx+c))(\tanh^4(dx+c)) - 2 \ln(\tanh(dx+c)+1)(\tanh^4(dx+c)) - 2 \ln(\tanh(dx+c)-1)(\tanh^4(dx+c)))}{4d(-\tanh^2(dx+c))^{5/2}}$
default	$\frac{\tanh(dx+c)(4 \ln(\tanh(dx+c))(\tanh^4(dx+c)) - 2 \ln(\tanh(dx+c)+1)(\tanh^4(dx+c)) - 2 \ln(\tanh(dx+c)-1)(\tanh^4(dx+c)))}{4d(-\tanh^2(dx+c))^{5/2}}$
risch	$\frac{(e^{2dx+2c}-1)x}{\sqrt{-\frac{(e^{2dx+2c}-1)^2}{(1+e^{2dx+2c})^2}}(1+e^{2dx+2c})} - \frac{2(e^{2dx+2c}-1)(dx+c)}{\sqrt{-\frac{(e^{2dx+2c}-1)^2}{(1+e^{2dx+2c})^2}}(1+e^{2dx+2c})d} - \frac{4e^{2dx+2c}(e^{4dx+4c}-e^{2dx+2c})}{(e^{2dx+2c}-1)^3(1+e^{2dx+2c})\sqrt{-\frac{(e^{2dx+2c}-1)^2}{(1+e^{2dx+2c})^2}}}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/(-tanh(d*x+c)^2)^(5/2), x, method=_RETURNVERBOSE)`

[Out] $1/4/d*\tanh(d*x+c)*(4*\ln(\tanh(d*x+c))*\tanh(d*x+c)^4-2*\ln(\tanh(d*x+c)+1)*\tanh(d*x+c)^4-2*\ln(\tanh(d*x+c)-1)*\tanh(d*x+c)^4-2*\tanh(d*x+c)^2-1)/(-\tanh(d*x+c))^2)^{(5/2)}$

Maxima [C] Result contains complex when optimal does not.

time = 0.48, size = 132, normalized size = 1.50

$$\frac{i(dx+c)}{d} + \frac{i \log(e^{-dx-c} + 1)}{d} + \frac{i \log(e^{-dx-c} - 1)}{d} - \frac{4(-i e^{-2dx-2c} + i e^{-4dx-4c} - i e^{-6dx-6c})}{d(4e^{-2dx-2c} - 6e^{-4dx-4c} + 4e^{-6dx-6c} - e^{-8dx-8c} - 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(-tanh(d*x+c))^2)^(5/2),x, algorithm="maxima")`

[Out] $I*(d*x + c)/d + I*\log(e^{-d*x - c} + 1)/d + I*\log(e^{-d*x - c} - 1)/d - 4*(-I*e^{-2*d*x - 2*c} + I*e^{-4*d*x - 4*c} - I*e^{-6*d*x - 6*c})/(d*(4*e^{-2*d*x - 2*c} - 6*e^{-4*d*x - 4*c} + 4*e^{-6*d*x - 6*c} - e^{-8*d*x - 8*c} - 1))$

Fricas [C] Result contains complex when optimal does not.

time = 0.42, size = 180, normalized size = 2.05

$$\frac{i dx e^{(8 dx+8 c)} + i dx - 4(i dx - i) e^{(6 dx+6 c)} - 2(-3i dx + 2i) e^{(4 dx+4 c)} - 4(i dx - i) e^{(2 dx+2 c)} + (-i e^{(8 dx+8 c)} + 4i e^{(6 dx+6 c)} - 6i e^{(4 dx+4 c)} + 4i e^{(2 dx+2 c)} - i) \log(e^{(2 dx+2 c)} - 1)}{d e^{(8 dx+8 c)} - 4 d e^{(6 dx+6 c)} + 6 d e^{(4 dx+4 c)} - 4 d e^{(2 dx+2 c)} + d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(-tanh(d*x+c))^2)^(5/2),x, algorithm="fricas")`

[Out] $(I*d*x*e^{(8*d*x + 8*c)} + I*d*x - 4*(I*d*x - I)*e^{(6*d*x + 6*c)} - 2*(-3*I*d*x + 2*I)*e^{(4*d*x + 4*c)} - 4*(I*d*x - I)*e^{(2*d*x + 2*c)} + (-I*e^{(8*d*x + 8*c)} + 4*I*e^{(6*d*x + 6*c)} - 6*I*e^{(4*d*x + 4*c)} + 4*I*e^{(2*d*x + 2*c)} - I)*\log(e^{(2*d*x + 2*c)} - 1))/(d*e^{(8*d*x + 8*c)} - 4*d*e^{(6*d*x + 6*c)} + 6*d*e^{(4*d*x + 4*c)} - 4*d*e^{(2*d*x + 2*c)} + d)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(-\tanh^2(c + dx))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(-tanh(d*x+c)**2)**(5/2),x)`

[Out] `Integral((-tanh(c + d*x)**2)**(-5/2), x)`

Giac [C] Result contains complex when optimal does not.

time = 0.42, size = 126, normalized size = 1.43

$$-\frac{\frac{i dx+i c}{\operatorname{sgn}(-e^{(4 dx+4 c)}+1)} - \frac{i \log(e^{(2 dx+2 c)}-1)}{\operatorname{sgn}(-e^{(4 dx+4 c)}+1)} + \frac{4(i e^{(6 dx+6 c)}-i e^{(4 dx+4 c)}+i e^{(2 dx+2 c)})}{(e^{(2 dx+2 c)}-1)^4 \operatorname{sgn}(-e^{(4 dx+4 c)}+1)}}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(-tanh(d*x+c)^2)^(5/2),x, algorithm="giac")

[Out] -((I*d*x + I*c)/sgn(-e^(4*d*x + 4*c) + 1) - I*log(e^(2*d*x + 2*c) - 1)/sgn(-e^(4*d*x + 4*c) + 1) + 4*(I*e^(6*d*x + 6*c) - I*e^(4*d*x + 4*c) + I*e^(2*d*x + 2*c))/((e^(2*d*x + 2*c) - 1)^4*sgn(-e^(4*d*x + 4*c) + 1))/d

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(-\tanh(c + dx)^2)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(-tanh(c + d*x)^2)^(5/2),x)

[Out] int(1/(-tanh(c + d*x)^2)^(5/2), x)

3.33 $\int \sqrt{\tanh^3(x)} dx$

Optimal. Leaf size=57

$$-2 \coth(x) \sqrt{\tanh^3(x)} + \frac{\operatorname{ArcTan}\left(\sqrt{\tanh(x)}\right) \sqrt{\tanh^3(x)}}{\tanh^{\frac{3}{2}}(x)} + \frac{\tanh^{-1}\left(\sqrt{\tanh(x)}\right) \sqrt{\tanh^3(x)}}{\tanh^{\frac{3}{2}}(x)}$$

[Out] $-2*\coth(x)*(\tanh(x)^3)^{(1/2)}+\arctan(\tanh(x)^{(1/2)})*(\tanh(x)^3)^{(1/2)}/\tanh(x)^{(3/2)}+\operatorname{arctanh}(\tanh(x)^{(1/2)})*(\tanh(x)^3)^{(1/2)}/\tanh(x)^{(3/2)}$

Rubi [A]

time = 0.03, antiderivative size = 57, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.875$, Rules used = {3739, 3554, 3557, 335, 218, 212, 209}

$$\frac{\sqrt{\tanh^3(x)} \operatorname{ArcTan}\left(\sqrt{\tanh(x)}\right)}{\tanh^{\frac{3}{2}}(x)} + \frac{\tanh^{-1}\left(\sqrt{\tanh(x)}\right) \sqrt{\tanh^3(x)}}{\tanh^{\frac{3}{2}}(x)} - 2\sqrt{\tanh^3(x)} \coth(x)$$

Antiderivative was successfully verified.

[In] `Int[Sqrt[Tanh[x]^3], x]`

[Out] $-2*\operatorname{Coth}[x]*\operatorname{Sqrt}[\operatorname{Tanh}[x]^3] + (\operatorname{ArcTan}[\operatorname{Sqrt}[\operatorname{Tanh}[x]]]*\operatorname{Sqrt}[\operatorname{Tanh}[x]^3])/\operatorname{Tanh}[x]^{(3/2)} + (\operatorname{ArcTanh}[\operatorname{Sqrt}[\operatorname{Tanh}[x]]]*\operatorname{Sqrt}[\operatorname{Tanh}[x]^3])/\operatorname{Tanh}[x]^{(3/2)}$

Rule 209

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

Rule 212

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rule 218

`Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

Rule 335

```
Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
  Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
  ))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 3554

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d
*x])^(n - 1)/(d*(n - 1))), x] - Dist[b^2, Int[(b*Tan[c + d*x])^(n - 2), x],
x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]
```

Rule 3557

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[b/d, Subst[Int[
x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !
IntegerQ[n]
```

Rule 3739

```
Int[(u_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff
= FreeFactors[Tan[e + f*x], x]}, Dist[(b*ff^n)^IntPart[p]*((b*Tan[e + f*x]^
n)^FracPart[p]/(Tan[e + f*x]/ff)^(n*FracPart[p])), Int[ActivateTrig[u]*(Tan
[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p]
&& IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /;
FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]
```

Rubi steps

$$\begin{aligned}
\int \sqrt{\tanh^3(x)} dx &= \frac{\sqrt{\tanh^3(x)} \int \tanh^{\frac{3}{2}}(x) dx}{\tanh^{\frac{3}{2}}(x)} \\
&= -2 \coth(x) \sqrt{\tanh^3(x)} + \frac{\sqrt{\tanh^3(x)} \int \frac{1}{\sqrt{\tanh(x)}} dx}{\tanh^{\frac{3}{2}}(x)} \\
&= -2 \coth(x) \sqrt{\tanh^3(x)} - \frac{\sqrt{\tanh^3(x)} \operatorname{Subst}\left(\int \frac{1}{\sqrt{x}(-1+x^2)} dx, x, \tanh(x)\right)}{\tanh^{\frac{3}{2}}(x)} \\
&= -2 \coth(x) \sqrt{\tanh^3(x)} - \frac{\left(2\sqrt{\tanh^3(x)}\right) \operatorname{Subst}\left(\int \frac{1}{-1+x^4} dx, x, \sqrt{\tanh(x)}\right)}{\tanh^{\frac{3}{2}}(x)} \\
&= -2 \coth(x) \sqrt{\tanh^3(x)} + \frac{\sqrt{\tanh^3(x)} \operatorname{Subst}\left(\int \frac{1}{1-x^2} dx, x, \sqrt{\tanh(x)}\right)}{\tanh^{\frac{3}{2}}(x)} + \frac{\sqrt{\tanh^3(x)}}{\tanh^{\frac{3}{2}}(x)} \\
&= -2 \coth(x) \sqrt{\tanh^3(x)} + \frac{\tan^{-1}\left(\sqrt{\tanh(x)}\right) \sqrt{\tanh^3(x)}}{\tanh^{\frac{3}{2}}(x)} + \frac{\tanh^{-1}\left(\sqrt{\tanh(x)}\right) \sqrt{\tanh^3(x)}}{\tanh^{\frac{3}{2}}(x)}
\end{aligned}$$

Mathematica [A]

time = 0.03, size = 38, normalized size = 0.67

$$\frac{\left(\operatorname{ArcTan}\left(\sqrt{\tanh(x)}\right) + \tanh^{-1}\left(\sqrt{\tanh(x)}\right) - 2\sqrt{\tanh(x)}\right) \sqrt{\tanh^3(x)}}{\tanh^{\frac{3}{2}}(x)}$$

Antiderivative was successfully verified.

`[In] Integrate[Sqrt[Tanh[x]^3], x]``[Out] ((ArcTan[Sqrt[Tanh[x]]] + ArcTanh[Sqrt[Tanh[x]]] - 2*Sqrt[Tanh[x]])*Sqrt[Tanh[x]^3])/Tanh[x]^(3/2)`**Maple [A]**

time = 0.87, size = 43, normalized size = 0.75

method	result
derivativedivides	$ -\frac{\sqrt{\tanh^3(x)} \left(4\left(\sqrt{\tanh(x)}\right)_{+\ln\left(\sqrt{\tanh(x)}-1\right)-\ln\left(\sqrt{\tanh(x)}+1\right)-2\arctan\left(\sqrt{\tanh(x)}\right)}\right)}{2 \tanh(x)^{\frac{3}{2}}} $

default	$-\frac{\sqrt{\tanh^3(x)} \left(4 \left(\sqrt{\tanh(x)} \right) + \ln \left(\sqrt{\tanh(x)} - 1 \right) - \ln \left(\sqrt{\tanh(x)} + 1 \right) - 2 \arctan \left(\sqrt{\tanh(x)} \right) \right)}{2 \tanh(x)^{\frac{3}{2}}}$
---------	--

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((tanh(x)^3)^(1/2),x,method=_RETURNVERBOSE)`

[Out] `-1/2*(tanh(x)^3)^(1/2)*(4*tanh(x)^(1/2)+ln(tanh(x)^(1/2)-1)-ln(tanh(x)^(1/2)+1)-2*arctan(tanh(x)^(1/2)))/tanh(x)^(3/2)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((tanh(x)^3)^(1/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(tanh(x)^3), x)`

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 106 vs. 2(43) = 86.

time = 0.44, size = 106, normalized size = 1.86

$$-2 \sqrt{\frac{\sinh(x)}{\cosh(x)}} + \arctan \left(-\cosh(x)^2 - 2 \cosh(x) \sinh(x) - \sinh(x)^2 + (\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 + 1) \sqrt{\frac{\sinh(x)}{\cosh(x)}} \right) - \frac{1}{2} \log \left(-\cosh(x)^2 - 2 \cosh(x) \sinh(x) - \sinh(x)^2 + (\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 + 1) \sqrt{\frac{\sinh(x)}{\cosh(x)}} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((tanh(x)^3)^(1/2),x, algorithm="fricas")`

[Out] `-2*sqrt(sinh(x)/cosh(x)) + arctan(-cosh(x)^2 - 2*cosh(x)*sinh(x) - sinh(x)^2 + (cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)*sqrt(sinh(x)/cosh(x))) - 1/2*log(-cosh(x)^2 - 2*cosh(x)*sinh(x) - sinh(x)^2 + (cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)*sqrt(sinh(x)/cosh(x)))`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{\tanh^3(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((tanh(x)**3)**(1/2),x)`

[Out] `Integral(sqrt(tanh(x)**3), x)`

Giac [A]

time = 0.41, size = 55, normalized size = 0.96

$$\frac{4}{\sqrt{e^{(4x)} - 1} - e^{(2x)} - 1} + \arctan\left(\sqrt{e^{(4x)} - 1} - e^{(2x)}\right) - \frac{1}{2} \log\left(-\sqrt{e^{(4x)} - 1} + e^{(2x)}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((tanh(x)^3)^(1/2),x, algorithm="giac")**[Out]** 4/(sqrt(e^(4*x) - 1) - e^(2*x) - 1) + arctan(sqrt(e^(4*x) - 1) - e^(2*x)) - 1/2*log(-sqrt(e^(4*x) - 1) + e^(2*x))**Mupad [F]**

time = 0.00, size = -1, normalized size = -0.02

$$\int \sqrt{\tanh(x)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((tanh(x)^3)^(1/2),x)**[Out]** int((tanh(x)^3)^(1/2), x)

3.34 $\int (a \tanh^3(x))^{3/2} dx$

Optimal. Leaf size=86

$$-\frac{2}{3}a\sqrt{a \tanh^3(x)} - \frac{a \operatorname{ArcTan}\left(\sqrt{\tanh(x)}\right) \sqrt{a \tanh^3(x)}}{\tanh^{\frac{3}{2}}(x)} + \frac{a \tanh^{-1}\left(\sqrt{\tanh(x)}\right) \sqrt{a \tanh^3(x)}}{\tanh^{\frac{3}{2}}(x)} - \frac{2}{7}a \tanh^2(x)$$

[Out] $-2/3*a*(a*\tanh(x)^3)^{(1/2)}-a*\arctan(\tanh(x)^{(1/2)})*(a*\tanh(x)^3)^{(1/2)}/\tanh(x)^{(3/2)}+a*\operatorname{arctanh}(\tanh(x)^{(1/2)})*(a*\tanh(x)^3)^{(1/2)}/\tanh(x)^{(3/2)}-2/7*a*(a*\tanh(x)^3)^{(1/2)}*\tanh(x)^2$

Rubi [A]

time = 0.03, antiderivative size = 86, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.700$, Rules used = {3739, 3554, 3557, 335, 304, 209, 212}

$$-\frac{a\sqrt{a \tanh^3(x)} \operatorname{ArcTan}\left(\sqrt{\tanh(x)}\right)}{\tanh^{\frac{3}{2}}(x)} - \frac{2}{3}a\sqrt{a \tanh^3(x)} - \frac{2}{7}a \tanh^2(x) \sqrt{a \tanh^3(x)} + \frac{a \tanh^{-1}\left(\sqrt{\tanh(x)}\right) \sqrt{a \tanh^3(x)}}{\tanh^{\frac{3}{2}}(x)}$$

Antiderivative was successfully verified.

[In] `Int[(a*Tanh[x]^3)^(3/2),x]`

[Out] $(-2*a*\operatorname{Sqrt}[a*\operatorname{Tanh}[x]^3])/3 - (a*\operatorname{ArcTan}[\operatorname{Sqrt}[\operatorname{Tanh}[x]]]*\operatorname{Sqrt}[a*\operatorname{Tanh}[x]^3])/ \operatorname{Tanh}[x]^{(3/2)} + (a*\operatorname{ArcTanh}[\operatorname{Sqrt}[\operatorname{Tanh}[x]]]*\operatorname{Sqrt}[a*\operatorname{Tanh}[x]^3])/ \operatorname{Tanh}[x]^{(3/2)} - (2*a*\operatorname{Tanh}[x]^2*\operatorname{Sqrt}[a*\operatorname{Tanh}[x]^3])/7$

Rule 209

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

Rule 212

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rule 304

`Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

Rule 335

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
  Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
  )^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 3554

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d
*x])^(n - 1)/(d*(n - 1))), x] - Dist[b^2, Int[(b*Tan[c + d*x])^(n - 2), x],
x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]
```

Rule 3557

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[b/d, Subst[Int[
x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !
IntegerQ[n]
```

Rule 3739

```
Int[(u_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff
= FreeFactors[Tan[e + f*x], x]}, Dist[(b*ff^n)^IntPart[p]*(b*Tan[e + f*x]^
n)^FracPart[p]/(Tan[e + f*x]/ff)^(n*FracPart[p])], Int[ActivateTrig[u]*(Tan
[e + f*x]/ff)^(n*p), x], x]] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p]
&& IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /;
FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]
```

Rubi steps

$$\begin{aligned}
\int (a \tanh^3(x))^{3/2} dx &= \frac{\left(a \sqrt{a \tanh^3(x)}\right) \int \tanh^{\frac{9}{2}}(x) dx}{\tanh^{\frac{3}{2}}(x)} \\
&= -\frac{2}{7} a \tanh^2(x) \sqrt{a \tanh^3(x)} + \frac{\left(a \sqrt{a \tanh^3(x)}\right) \int \tanh^{\frac{5}{2}}(x) dx}{\tanh^{\frac{3}{2}}(x)} \\
&= -\frac{2}{3} a \sqrt{a \tanh^3(x)} - \frac{2}{7} a \tanh^2(x) \sqrt{a \tanh^3(x)} + \frac{\left(a \sqrt{a \tanh^3(x)}\right) \int \sqrt{\tanh(x)} dx}{\tanh^{\frac{3}{2}}(x)} \\
&= -\frac{2}{3} a \sqrt{a \tanh^3(x)} - \frac{2}{7} a \tanh^2(x) \sqrt{a \tanh^3(x)} - \frac{\left(a \sqrt{a \tanh^3(x)}\right) \text{Subst}\left(\int \frac{\sqrt{x}}{-1+x} dx\right)}{\tanh^{\frac{3}{2}}(x)} \\
&= -\frac{2}{3} a \sqrt{a \tanh^3(x)} - \frac{2}{7} a \tanh^2(x) \sqrt{a \tanh^3(x)} - \frac{\left(2a \sqrt{a \tanh^3(x)}\right) \text{Subst}\left(\int \frac{x^2}{-1+x} dx\right)}{\tanh^{\frac{3}{2}}(x)} \\
&= -\frac{2}{3} a \sqrt{a \tanh^3(x)} - \frac{2}{7} a \tanh^2(x) \sqrt{a \tanh^3(x)} + \frac{\left(a \sqrt{a \tanh^3(x)}\right) \text{Subst}\left(\int \frac{1}{1-x^2} dx\right)}{\tanh^{\frac{3}{2}}(x)} \\
&= -\frac{2}{3} a \sqrt{a \tanh^3(x)} - \frac{a \tan^{-1}\left(\sqrt{\tanh(x)}\right) \sqrt{a \tanh^3(x)}}{\tanh^{\frac{3}{2}}(x)} + \frac{a \tanh^{-1}\left(\sqrt{\tanh(x)}\right)}{\tanh^{\frac{3}{2}}(x)}
\end{aligned}$$

Mathematica [A]

time = 0.05, size = 55, normalized size = 0.64

$$\frac{(a \tanh^3(x))^{3/2} \left(21 \text{ArcTan}\left(\sqrt{\tanh(x)}\right) - 21 \tanh^{-1}\left(\sqrt{\tanh(x)}\right) + 14 \tanh^{\frac{3}{2}}(x) + 6 \tanh^{\frac{7}{2}}(x)\right)}{21 \tanh^{\frac{9}{2}}(x)}$$

Antiderivative was successfully verified.

`[In] Integrate[(a*Tanh[x]^3)^(3/2), x]`

```
[Out] -1/21*((a*Tanh[x]^3)^(3/2)*(21*ArcTan[Sqrt[Tanh[x]]] - 21*ArcTanh[Sqrt[Tanh[x]]] + 14*Tanh[x]^(3/2) + 6*Tanh[x]^(7/2)))/Tanh[x]^(9/2)
```

Maple [A]

time = 1.12, size = 76, normalized size = 0.88

method	result
--------	--------

derivativedivides	$\frac{(a(\tanh^3(x)))^{\frac{3}{2}} \left(21a^{\frac{7}{2}} \arctan\left(\frac{\sqrt{a \tanh(x)}}{\sqrt{a}}\right) - 21a^{\frac{7}{2}} \operatorname{arctanh}\left(\frac{\sqrt{a \tanh(x)}}{\sqrt{a}}\right) + 6(a \tanh(x))^{\frac{7}{2}} + 14a^2 \right)}{21 \tanh(x)^3 (a \tanh(x))^{\frac{3}{2}} a^2}$
default	$\frac{(a(\tanh^3(x)))^{\frac{3}{2}} \left(21a^{\frac{7}{2}} \arctan\left(\frac{\sqrt{a \tanh(x)}}{\sqrt{a}}\right) - 21a^{\frac{7}{2}} \operatorname{arctanh}\left(\frac{\sqrt{a \tanh(x)}}{\sqrt{a}}\right) + 6(a \tanh(x))^{\frac{7}{2}} + 14a^2 \right)}{21 \tanh(x)^3 (a \tanh(x))^{\frac{3}{2}} a^2}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a*tanh(x)^3)^(3/2),x,method=_RETURNVERBOSE)`

[Out]
$$-1/21*(a*\tanh(x)^3)^{(3/2)}*(21*a^{(7/2)}*\arctan((a*\tanh(x))^{(1/2)}/a^{(1/2)})-21*a^{(7/2)}*\operatorname{arctanh}((a*\tanh(x))^{(1/2)}/a^{(1/2)})+6*(a*\tanh(x))^{(7/2)}+14*a^2*(a*\tanh(x))^{(3/2)})/\tanh(x)^3/(a*\tanh(x))^{(3/2)}/a^2$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a*tanh(x)^3)^(3/2),x, algorithm="maxima")`

[Out] `integrate((a*tanh(x)^3)^(3/2), x)`

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 615 vs. 2(66) = 132.

time = 0.41, size = 1269, normalized size = 14.76

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a*tanh(x)^3)^(3/2),x, algorithm="fricas")`

[Out]
$$\begin{aligned} & [-1/84*(42*(a*\cosh(x))^6 + 6*a*\cosh(x)*\sinh(x)^5 + a*\sinh(x)^6 + 3*a*\cosh(x)^4 \\ & + 3*(5*a*\cosh(x)^2 + a)*\sinh(x)^4 + 4*(5*a*\cosh(x)^3 + 3*a*\cosh(x))*\sinh(x)^3 \\ & + 3*a*\cosh(x)^2 + 3*(5*a*\cosh(x)^4 + 6*a*\cosh(x)^2 + a)*\sinh(x)^2 + 6 \\ & *(a*\cosh(x)^5 + 2*a*\cosh(x)^3 + a*\cosh(x))*\sinh(x) + a)*\sqrt{-a}*\arctan((\cosh(x)^2 \\ & + 2*\cosh(x)*\sinh(x) + \sinh(x)^2)*\sqrt{-a}*\sqrt{a*\sinh(x)/\cosh(x)})/(a*\cosh(x)^2 \\ & + 2*a*\cosh(x)*\sinh(x) + a*\sinh(x)^2 - a) - 21*(a*\cosh(x))^6 + 6 \\ & *a*\cosh(x)*\sinh(x)^5 + a*\sinh(x)^6 + 3*a*\cosh(x)^4 + 3*(5*a*\cosh(x)^2 + a)* \\ & \sinh(x)^4 + 4*(5*a*\cosh(x)^3 + 3*a*\cosh(x))*\sinh(x)^3 + 3*a*\cosh(x)^2 + 3*(\\ & 5*a*\cosh(x)^4 + 6*a*\cosh(x)^2 + a)*\sinh(x)^2 + 6*(a*\cosh(x))^5 + 2*a*\cosh(x)^3 \\ & + a*\cosh(x))*\sinh(x) + a)*\sqrt{-a}*\log(-(a*\cosh(x))^4 + 4*a*\cosh(x)^3*\sinh(x) \\ & + 3*a*\cosh(x)^2 + a)*\sinh(x) + a) \end{aligned}$$

$h(x) + 6*a*cosh(x)^2*sinh(x)^2 + 4*a*cosh(x)*sinh(x)^3 + a*sinh(x)^4 - 2*(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)*sqrt(-a)*sqrt(a*sinh(x)/cosh(x)) - 2*a)/(cosh(x)^4 + 4*cosh(x)^3*sinh(x) + 6*cosh(x)^2*sinh(x)^2 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4) + 16*(5*a*cosh(x)^6 + 30*a*cosh(x)*sinh(x)^5 + 5*a*sinh(x)^6 - a*cosh(x)^4 + (75*a*cosh(x)^2 - a)*sinh(x)^4 + 4*(25*a*cosh(x)^3 - a*cosh(x))*sinh(x)^3 + a*cosh(x)^2 + (75*a*cosh(x)^4 - 6*a*cosh(x)^2 + a)*sinh(x)^2 + 2*(15*a*cosh(x)^5 - 2*a*cosh(x)^3 + a*cosh(x))*sinh(x) - 5*a)*sqrt(a*sinh(x)/cosh(x)))/(cosh(x)^6 + 6*cosh(x)*sinh(x)^5 + sinh(x)^6 + 3*(5*cosh(x)^2 + 1)*sinh(x)^4 + 3*cosh(x)^4 + 4*(5*cosh(x)^3 + 3*cosh(x))*sinh(x)^3 + 3*(5*cosh(x)^4 + 6*cosh(x)^2 + 1)*sinh(x)^2 + 3*cosh(x)^2 + 6*(cosh(x)^5 + 2*cosh(x)^3 + cosh(x))*sinh(x) + 1), -1/84*(42*(a*cosh(x)^6 + 6*a*cosh(x)*sinh(x)^5 + a*sinh(x)^6 + 3*a*cosh(x)^4 + 3*(5*a*cosh(x)^2 + a)*sinh(x)^4 + 4*(5*a*cosh(x)^3 + 3*a*cosh(x))*sinh(x)^3 + 3*a*cosh(x)^2 + 3*(5*a*cosh(x)^4 + 6*a*cosh(x)^2 + a)*sinh(x)^2 + 6*(a*cosh(x)^5 + 2*a*cosh(x)^3 + a*cosh(x))*sinh(x) + a)*sqrt(a)*arctan((cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)*sqrt(a*sinh(x)/cosh(x))/sqrt(a)) - 21*(a*cosh(x)^6 + 6*a*cosh(x)*sinh(x)^5 + a*sinh(x)^6 + 3*a*cosh(x)^4 + 3*(5*a*cosh(x)^2 + a)*sinh(x)^4 + 4*(5*a*cosh(x)^3 + 3*a*cosh(x))*sinh(x)^3 + 3*a*cosh(x)^2 + 3*(5*a*cosh(x)^4 + 6*a*cosh(x)^2 + a)*sinh(x)^2 + 6*(a*cosh(x)^5 + 2*a*cosh(x)^3 + a*cosh(x))*sinh(x) + a)*sqrt(a)*log(2*a*cosh(x)^4 + 8*a*cosh(x)^3*sinh(x) + 12*a*cosh(x)^2*sinh(x)^2 + 8*a*cosh(x)*sinh(x)^3 + 2*a*sinh(x)^4 + 2*(cosh(x)^4 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4 + (6*cosh(x)^2 + 1)*sinh(x)^2 + cosh(x)^2 + 2*(2*cosh(x)^3 + cosh(x))*sinh(x))*sqrt(a)*sqrt(a*sinh(x)/cosh(x)) - a) + 16*(5*a*cosh(x)^6 + 30*a*cosh(x)*sinh(x)^5 + 5*a*sinh(x)^6 - a*cosh(x)^4 + (75*a*cosh(x)^2 - a)*sinh(x)^4 + 4*(25*a*cosh(x)^3 - a*cosh(x))*sinh(x)^3 + a*cosh(x)^2 + (75*a*cosh(x)^4 - 6*a*cosh(x)^2 + a)*sinh(x)^2 + 2*(15*a*cosh(x)^5 - 2*a*cosh(x)^3 + a*cosh(x))*sinh(x) - 5*a)*sqrt(a*sinh(x)/cosh(x)))/(cosh(x)^6 + 6*cosh(x)*sinh(x)^5 + sinh(x)^6 + 3*(5*cosh(x)^2 + 1)*sinh(x)^4 + 3*cosh(x)^4 + 4*(5*cosh(x)^3 + 3*cosh(x))*sinh(x)^3 + 3*(5*cosh(x)^4 + 6*cosh(x)^2 + 1)*sinh(x)^2 + 3*cosh(x)^2 + 6*(cosh(x)^5 + 2*cosh(x)^3 + cosh(x))*sinh(x) + 1)]$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (a \tanh^3(x))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*tanh(x)**3)**(3/2), x)

[Out] Integral((a*tanh(x)**3)**(3/2), x)

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 342 vs. 2(66) = 132.

time = 0.47, size = 342, normalized size = 3.98

$$\frac{1}{8} \left(4\sqrt{a} \operatorname{arctan} \left(\frac{\sqrt{2a} \operatorname{tanh}(x) - \sqrt{2a} \operatorname{sech}(x)}{2} \right) \operatorname{sech}(x)^{-1} + 2\sqrt{2} \log \left(\frac{-\sqrt{2a} \operatorname{tanh}(x) + \sqrt{2a} \operatorname{sech}(x)}{\operatorname{sech}(x)^{-1}} \right) - \frac{38 \left(2 \left(\sqrt{2a} \operatorname{tanh}(x) - \sqrt{2a} \operatorname{sech}(x) \right) \operatorname{sech}(x)^{-1} - 1 \right) + 42 \left(\sqrt{2a} \operatorname{tanh}(x) - \sqrt{2a} \operatorname{sech}(x) \right) \operatorname{sech}(x)^{-1} - 110 \left(\sqrt{2a} \operatorname{tanh}(x) - \sqrt{2a} \operatorname{sech}(x) \right) \operatorname{sech}(x)^{-1} + 38 \left(\sqrt{2a} \operatorname{tanh}(x) - \sqrt{2a} \operatorname{sech}(x) \right) \operatorname{sech}(x)^{-1} + 42 \left(\sqrt{2a} \operatorname{tanh}(x) - \sqrt{2a} \operatorname{sech}(x) \right) \operatorname{sech}(x)^{-1} - 110 \left(\sqrt{2a} \operatorname{tanh}(x) - \sqrt{2a} \operatorname{sech}(x) \right) \operatorname{sech}(x)^{-1} + 38 \left(\sqrt{2a} \operatorname{tanh}(x) - \sqrt{2a} \operatorname{sech}(x) \right) \operatorname{sech}(x)^{-1} \right)}{\left(\sqrt{2a} \operatorname{tanh}(x) - \sqrt{2a} \operatorname{sech}(x) \right)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*tanh(x)^3)^(3/2),x, algorithm="giac")

[Out]
$$\begin{aligned} & -1/42*(42*\sqrt{a}*\arctan(-(\sqrt{a}*e^{2x}) - \sqrt{a*e^{4x} - a})/\sqrt{a})* \\ & \operatorname{sgn}(e^{4x} - 1) + 21*\sqrt{a}*\log(\operatorname{abs}(-\sqrt{a}*e^{2x}) + \sqrt{a*e^{4x} - a} \\ &))*\operatorname{sgn}(e^{4x} - 1) + 16*(21*(\sqrt{a}*e^{2x}) - \sqrt{a*e^{4x} - a})^6*a*s \\ & \operatorname{gn}(e^{4x} - 1) + 42*(\sqrt{a}*e^{2x}) - \sqrt{a*e^{4x} - a})^5*a^{3/2}* \operatorname{sgn}(\\ & e^{4x} - 1) + 119*(\sqrt{a}*e^{2x}) - \sqrt{a*e^{4x} - a})^4*a^2*\operatorname{sgn}(e^{4x} \\ &) - 1) + 56*(\sqrt{a}*e^{2x}) - \sqrt{a*e^{4x} - a})^3*a^{5/2}* \operatorname{sgn}(e^{4x} - \\ & 1) + 63*(\sqrt{a}*e^{2x}) - \sqrt{a*e^{4x} - a})^2*a^3*\operatorname{sgn}(e^{4x} - 1) + 1 \\ & 4*(\sqrt{a}*e^{2x}) - \sqrt{a*e^{4x} - a})*a^{7/2}* \operatorname{sgn}(e^{4x} - 1) + 5*a^4* \\ & \operatorname{sgn}(e^{4x} - 1))/(\sqrt{a}*e^{2x}) - \sqrt{a*e^{4x} - a} + \sqrt{a})^7)*a \end{aligned}$$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int (a \tanh(x)^3)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*tanh(x)^3)^(3/2),x)

[Out] int((a*tanh(x)^3)^(3/2), x)

3.35 $\int \sqrt{a \tanh^3(x)} dx$

Optimal. Leaf size=63

$$-2 \coth(x) \sqrt{a \tanh^3(x)} + \frac{\operatorname{ArcTan}\left(\sqrt{\tanh(x)}\right) \sqrt{a \tanh^3(x)}}{\tanh^{\frac{3}{2}}(x)} + \frac{\tanh^{-1}\left(\sqrt{\tanh(x)}\right) \sqrt{a \tanh^3(x)}}{\tanh^{\frac{3}{2}}(x)}$$

[Out] $-2*\coth(x)*(a*\tanh(x)^3)^{(1/2)}+\arctan(\tanh(x)^{(1/2)})*(a*\tanh(x)^3)^{(1/2)}/\tanh(x)^{(3/2)}+\operatorname{arctanh}(\tanh(x)^{(1/2)})*(a*\tanh(x)^3)^{(1/2)}/\tanh(x)^{(3/2)}$

Rubi [A]

time = 0.02, antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.700$, Rules used = {3739, 3554, 3557, 335, 218, 212, 209}

$$\frac{\sqrt{a \tanh^3(x)} \operatorname{ArcTan}\left(\sqrt{\tanh(x)}\right)}{\tanh^{\frac{3}{2}}(x)} + \frac{\tanh^{-1}\left(\sqrt{\tanh(x)}\right) \sqrt{a \tanh^3(x)}}{\tanh^{\frac{3}{2}}(x)} - 2 \coth(x) \sqrt{a \tanh^3(x)}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[a*Tanh[x]^3],x]

[Out] $-2*\operatorname{Coth}[x]*\operatorname{Sqrt}[a*\operatorname{Tanh}[x]^3] + (\operatorname{ArcTan}[\operatorname{Sqrt}[\operatorname{Tanh}[x]]]*\operatorname{Sqrt}[a*\operatorname{Tanh}[x]^3])/\operatorname{Tanh}[x]^{(3/2)} + (\operatorname{ArcTanh}[\operatorname{Sqrt}[\operatorname{Tanh}[x]]]*\operatorname{Sqrt}[a*\operatorname{Tanh}[x]^3])/\operatorname{Tanh}[x]^{(3/2)}$

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 218

Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] :> With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 335


```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
  Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
  ))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 3554

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d
*x])^(n - 1)/(d*(n - 1))), x] - Dist[b^2, Int[(b*Tan[c + d*x])^(n - 2), x],
x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]
```

Rule 3557

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[b/d, Subst[Int[
x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !
IntegerQ[n]
```

Rule 3739

```
Int[(u_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_)^(p_), x_Symbol] := With[{ff
= FreeFactors[Tan[e + f*x], x]}, Dist[(b*ff^n)^IntPart[p]*((b*Tan[e + f*x]^
n)^FracPart[p]/(Tan[e + f*x]/ff)^(n*FracPart[p])), Int[ActivateTrig[u]*(Tan
[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p]
&& IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /;
FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]
```

Rubi steps

$$\begin{aligned}
\int \sqrt{a \tanh^3(x)} dx &= \frac{\sqrt{a \tanh^3(x)} \int \tanh^{\frac{3}{2}}(x) dx}{\tanh^{\frac{3}{2}}(x)} \\
&= -2 \coth(x) \sqrt{a \tanh^3(x)} + \frac{\sqrt{a \tanh^3(x)} \int \frac{1}{\sqrt{\tanh(x)}} dx}{\tanh^{\frac{3}{2}}(x)} \\
&= -2 \coth(x) \sqrt{a \tanh^3(x)} - \frac{\sqrt{a \tanh^3(x)} \operatorname{Subst}\left(\int \frac{1}{\sqrt{x}(-1+x^2)} dx, x, \tanh(x)\right)}{\tanh^{\frac{3}{2}}(x)} \\
&= -2 \coth(x) \sqrt{a \tanh^3(x)} - \frac{\left(2\sqrt{a \tanh^3(x)}\right) \operatorname{Subst}\left(\int \frac{1}{-1+x^4} dx, x, \sqrt{\tanh(x)}\right)}{\tanh^{\frac{3}{2}}(x)} \\
&= -2 \coth(x) \sqrt{a \tanh^3(x)} + \frac{\sqrt{a \tanh^3(x)} \operatorname{Subst}\left(\int \frac{1}{1-x^2} dx, x, \sqrt{\tanh(x)}\right)}{\tanh^{\frac{3}{2}}(x)} + \frac{\sqrt{a \tanh^3(x)}}{\tanh^{\frac{3}{2}}(x)} \\
&= -2 \coth(x) \sqrt{a \tanh^3(x)} + \frac{\tan^{-1}\left(\sqrt{\tanh(x)}\right) \sqrt{a \tanh^3(x)}}{\tanh^{\frac{3}{2}}(x)} + \frac{\tanh^{-1}\left(\sqrt{\tanh(x)}\right)}{\tanh^{\frac{3}{2}}(x)}
\end{aligned}$$

Mathematica [A]

time = 0.02, size = 40, normalized size = 0.63

$$\frac{\left(\operatorname{ArcTan}\left(\sqrt{\tanh(x)}\right) + \tanh^{-1}\left(\sqrt{\tanh(x)}\right) - 2\sqrt{\tanh(x)}\right) \sqrt{a \tanh^3(x)}}{\tanh^{\frac{3}{2}}(x)}$$

Antiderivative was successfully verified.

`[In] Integrate[Sqrt[a*Tanh[x]^3], x]`

```
[Out] ((ArcTan[Sqrt[Tanh[x]]] + ArcTanh[Sqrt[Tanh[x]]] - 2*Sqrt[Tanh[x]])*Sqrt[a*
Tanh[x]^3])/Tanh[x]^(3/2)
```

Maple [A]

time = 1.15, size = 59, normalized size = 0.94

method	result
derivativedivides	$ \frac{\sqrt{a(\tanh^3(x))} \left(-2\sqrt{a \tanh(x)} + \sqrt{a} \arctan\left(\frac{\sqrt{a \tanh(x)}}{\sqrt{a}}\right) + \sqrt{a} \operatorname{arctanh}\left(\frac{\sqrt{a \tanh(x)}}{\sqrt{a}}\right)\right)}{\tanh(x) \sqrt{a \tanh(x)}} $

default	$\frac{\sqrt{a (\tanh^3(x))} \left(-2\sqrt{a \tanh(x)} + \sqrt{a} \arctan\left(\frac{\sqrt{a \tanh(x)}}{\sqrt{a}}\right) + \sqrt{a} \operatorname{arctanh}\left(\frac{\sqrt{a \tanh(x)}}{\sqrt{a}}\right) \right)}{\tanh(x) \sqrt{a \tanh(x)}}$
---------	--

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((a*tanh(x)^3)^(1/2),x,method=_RETURNVERBOSE)
```

```
[Out] (a*tanh(x)^3)^(1/2)/tanh(x)/(a*tanh(x))^(1/2)*(-2*(a*tanh(x))^(1/2)+a^(1/2)
*arctan((a*tanh(x))^(1/2)/a^(1/2))+a^(1/2)*arctanh((a*tanh(x))^(1/2)/a^(1/2)
))
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a*tanh(x)^3)^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(sqrt(a*tanh(x)^3), x)
```

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 173 vs. 2(49) = 98.

time = 0.40, size = 376, normalized size = 5.97

$$\frac{1}{4} \sqrt{-a} \log\left(\frac{\cosh(x)^4 + 4a \cosh(x)^3 \sinh(x) + 6a^2 \cosh(x)^2 \sinh(x)^2 + 4a^3 \cosh(x) \sinh(x)^3 + a^4 \sinh(x)^4}{\cosh(x)^4 + 4a \cosh(x)^3 \sinh(x) + 6a^2 \cosh(x)^2 \sinh(x)^2 + 4a^3 \cosh(x) \sinh(x)^3 + a^4 \sinh(x)^4}\right) - \frac{1}{4} \sqrt{-a} \log\left(\frac{\cosh(x)^4 + 4a \cosh(x)^3 \sinh(x) + 6a^2 \cosh(x)^2 \sinh(x)^2 + 4a^3 \cosh(x) \sinh(x)^3 + a^4 \sinh(x)^4}{\cosh(x)^4 + 4a \cosh(x)^3 \sinh(x) + 6a^2 \cosh(x)^2 \sinh(x)^2 + 4a^3 \cosh(x) \sinh(x)^3 + a^4 \sinh(x)^4}\right) - \frac{1}{4} \sqrt{-a} \log\left(\frac{\cosh(x)^4 + 4a \cosh(x)^3 \sinh(x) + 6a^2 \cosh(x)^2 \sinh(x)^2 + 4a^3 \cosh(x) \sinh(x)^3 + a^4 \sinh(x)^4}{\cosh(x)^4 + 4a \cosh(x)^3 \sinh(x) + 6a^2 \cosh(x)^2 \sinh(x)^2 + 4a^3 \cosh(x) \sinh(x)^3 + a^4 \sinh(x)^4}\right) - \frac{1}{4} \sqrt{-a} \log\left(\frac{\cosh(x)^4 + 4a \cosh(x)^3 \sinh(x) + 6a^2 \cosh(x)^2 \sinh(x)^2 + 4a^3 \cosh(x) \sinh(x)^3 + a^4 \sinh(x)^4}{\cosh(x)^4 + 4a \cosh(x)^3 \sinh(x) + 6a^2 \cosh(x)^2 \sinh(x)^2 + 4a^3 \cosh(x) \sinh(x)^3 + a^4 \sinh(x)^4}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a*tanh(x)^3)^(1/2),x, algorithm="fricas")
```

```
[Out] [-1/2*sqrt(-a)*arctan((cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2)*sqrt(-a)*
sqrt(a*sinh(x)/cosh(x))/(a*cosh(x)^2 + 2*a*cosh(x)*sinh(x) + a*sinh(x)^2 -
a)) + 1/4*sqrt(-a)*log(-(a*cosh(x)^4 + 4*a*cosh(x)^3*sinh(x) + 6*a*cosh(x)^
2*sinh(x)^2 + 4*a*cosh(x)*sinh(x)^3 + a*sinh(x)^4 + 2*(cosh(x)^2 + 2*cosh(x)
)*sinh(x) + sinh(x)^2 + 1)*sqrt(-a)*sqrt(a*sinh(x)/cosh(x)) - 2*a)/(cosh(x)
^4 + 4*cosh(x)^3*sinh(x) + 6*cosh(x)^2*sinh(x)^2 + 4*cosh(x)*sinh(x)^3 + si
nh(x)^4) - 2*sqrt(a*sinh(x)/cosh(x)), -1/2*sqrt(a)*arctan(sqrt(a)*sqrt(a*s
inh(x)/cosh(x))/(a*cosh(x)^2 + 2*a*cosh(x)*sinh(x) + a*sinh(x)^2 - a)) + 1/
4*sqrt(a)*log(2*a*cosh(x)^4 + 8*a*cosh(x)^3*sinh(x) + 12*a*cosh(x)^2*sinh(x)
)^2 + 8*a*cosh(x)*sinh(x)^3 + 2*a*sinh(x)^4 + 2*(cosh(x)^4 + 4*cosh(x)*sinh
(x)^3 + sinh(x)^4 + (6*cosh(x)^2 + 1)*sinh(x)^2 + cosh(x)^2 + 2*(2*cosh(x)^
3 + cosh(x))*sinh(x))*sqrt(a)*sqrt(a*sinh(x)/cosh(x)) - a) - 2*sqrt(a*sinh(
x)/cosh(x))]
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{a \tanh^3(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*tanh(x)**3)**(1/2),x)**[Out]** Integral(sqrt(a*tanh(x)**3), x)**Giac [B]** Leaf count of result is larger than twice the leaf count of optimal. 115 vs. 2(49) = 98.

time = 0.43, size = 115, normalized size = 1.83

$$\sqrt{a} \arctan\left(-\frac{\sqrt{a} e^{2x} - \sqrt{ae^{4x} - a}}{\sqrt{a}}\right) \operatorname{sgn}(e^{4x} - 1) - \frac{1}{2} \sqrt{a} \log\left(|-\sqrt{a} e^{2x} + \sqrt{ae^{4x} - a}|\right) \operatorname{sgn}(e^{4x} - 1) - \frac{4 a \operatorname{sgn}(e^{4x} - 1)}{\sqrt{a} e^{2x} - \sqrt{ae^{4x} - a} + \sqrt{a}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*tanh(x)^3)^(1/2),x, algorithm="giac")

[Out] sqrt(a)*arctan(-(sqrt(a)*e^(2*x) - sqrt(a*e^(4*x) - a))/sqrt(a))*sgn(e^(4*x) - 1) - 1/2*sqrt(a)*log(abs(-sqrt(a)*e^(2*x) + sqrt(a*e^(4*x) - a)))*sgn(e^(4*x) - 1) - 4*a*sgn(e^(4*x) - 1)/(sqrt(a)*e^(2*x) - sqrt(a*e^(4*x) - a) + sqrt(a))

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \sqrt{a \tanh^3(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*tanh(x)^3)^(1/2),x)**[Out]** int((a*tanh(x)^3)^(1/2), x)

$$3.36 \quad \int \frac{1}{\sqrt{a \tanh^3(x)}} dx$$

Optimal. Leaf size=64

$$-\frac{2 \tanh(x)}{\sqrt{a \tanh^3(x)}} - \frac{\text{ArcTan}\left(\sqrt{\tanh(x)}\right) \tanh^{\frac{3}{2}}(x)}{\sqrt{a \tanh^3(x)}} + \frac{\tanh^{-1}\left(\sqrt{\tanh(x)}\right) \tanh^{\frac{3}{2}}(x)}{\sqrt{a \tanh^3(x)}}$$

[Out] $-2*\tanh(x)/(a*\tanh(x)^3)^{(1/2)}-\arctan(\tanh(x)^{(1/2)})*\tanh(x)^{(3/2)}/(a*\tanh(x)^3)^{(1/2)}+\text{arctanh}(\tanh(x)^{(1/2)})*\tanh(x)^{(3/2)}/(a*\tanh(x)^3)^{(1/2)}$

Rubi [A]

time = 0.02, antiderivative size = 64, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.700$, Rules used = {3739, 3555, 3557, 335, 304, 209, 212}

$$-\frac{\tanh^{\frac{3}{2}}(x)\text{ArcTan}\left(\sqrt{\tanh(x)}\right)}{\sqrt{a \tanh^3(x)}} - \frac{2 \tanh(x)}{\sqrt{a \tanh^3(x)}} + \frac{\tanh^{-1}\left(\sqrt{\tanh(x)}\right) \tanh^{\frac{3}{2}}(x)}{\sqrt{a \tanh^3(x)}}$$

Antiderivative was successfully verified.

[In] Int[1/Sqrt[a*Tanh[x]^3], x]

[Out] $(-2*\text{Tanh}[x])/ \text{Sqrt}[a*\text{Tanh}[x]^3] - (\text{ArcTan}[\text{Sqrt}[\text{Tanh}[x]]]*\text{Tanh}[x]^{(3/2)})/ \text{Sqrt}[a*\text{Tanh}[x]^3] + (\text{ArcTanh}[\text{Sqrt}[\text{Tanh}[x]]]*\text{Tanh}[x]^{(3/2)})/ \text{Sqrt}[a*\text{Tanh}[x]^3]$

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 304

Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 335

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
  Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
  ))^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 3555

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(b*Tan[c + d*x]
)^n/(b*d*(n + 1)), x] - Dist[1/b^2, Int[(b*Tan[c + d*x])^(n + 2), x],
x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1]
```

Rule 3557

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[b/d, Subst[Int[
x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !
IntegerQ[n]
```

Rule 3739

```
Int[(u_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff
= FreeFactors[Tan[e + f*x], x]}, Dist[(b*ff^n)^IntPart[p]*((b*Tan[e + f*x]^
n)^FracPart[p]/(Tan[e + f*x]/ff)^(n*FracPart[p])), Int[ActivateTrig[u]*(Tan
[e + f*x]/ff)^(n*p), x], x]] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p]
&& IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /;
FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]
```

Rubi steps

$$\begin{aligned}
\int \frac{1}{\sqrt{a \tanh^3(x)}} dx &= \frac{\tanh^{\frac{3}{2}}(x) \int \frac{1}{\tanh^{\frac{3}{2}}(x)} dx}{\sqrt{a \tanh^3(x)}} \\
&= -\frac{2 \tanh(x)}{\sqrt{a \tanh^3(x)}} + \frac{\tanh^{\frac{3}{2}}(x) \int \sqrt{\tanh(x)} dx}{\sqrt{a \tanh^3(x)}} \\
&= -\frac{2 \tanh(x)}{\sqrt{a \tanh^3(x)}} - \frac{\tanh^{\frac{3}{2}}(x) \text{Subst}\left(\int \frac{\sqrt{x}}{-1+x^2} dx, x, \tanh(x)\right)}{\sqrt{a \tanh^3(x)}} \\
&= -\frac{2 \tanh(x)}{\sqrt{a \tanh^3(x)}} - \frac{\left(2 \tanh^{\frac{3}{2}}(x)\right) \text{Subst}\left(\int \frac{x^2}{-1+x^4} dx, x, \sqrt{\tanh(x)}\right)}{\sqrt{a \tanh^3(x)}} \\
&= -\frac{2 \tanh(x)}{\sqrt{a \tanh^3(x)}} + \frac{\tanh^{\frac{3}{2}}(x) \text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \sqrt{\tanh(x)}\right)}{\sqrt{a \tanh^3(x)}} - \frac{\tanh^{\frac{3}{2}}(x) \text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \sqrt{\tanh(x)}\right)}{\sqrt{a \tanh^3(x)}} \\
&= -\frac{2 \tanh(x)}{\sqrt{a \tanh^3(x)}} - \frac{\tan^{-1}\left(\sqrt{\tanh(x)}\right) \tanh^{\frac{3}{2}}(x)}{\sqrt{a \tanh^3(x)}} + \frac{\tanh^{-1}\left(\sqrt{\tanh(x)}\right) \tanh^{\frac{3}{2}}(x)}{\sqrt{a \tanh^3(x)}}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 26, normalized size = 0.41

$$-\frac{{}_2F_1\left(-\frac{1}{4}, 1; \frac{3}{4}; \tanh^2(x)\right) \tanh(x)}{\sqrt{a \tanh^3(x)}}$$

Antiderivative was successfully verified.

[In] Integrate[1/Sqrt[a*Tanh[x]^3], x]

[Out] (-2*Hypergeometric2F1[-1/4, 1, 3/4, Tanh[x]^2]*Tanh[x])/Sqrt[a*Tanh[x]^3]

Maple [A]

time = 1.14, size = 65, normalized size = 1.02

method	result
--------	--------

derivativedivides	$\frac{\tanh(x) \left(2a^{\frac{5}{2}} + \arctan\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a}}\right) a^2 \sqrt{a \tanh(x)} - \operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a}}\right) a^2 \sqrt{a \tanh(x)} \right)}{\sqrt{a} (\tanh^3(x)) a^{\frac{5}{2}}}$
default	$\frac{\tanh(x) \left(2a^{\frac{5}{2}} + \arctan\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a}}\right) a^2 \sqrt{a \tanh(x)} - \operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a}}\right) a^2 \sqrt{a \tanh(x)} \right)}{\sqrt{a} (\tanh^3(x)) a^{\frac{5}{2}}}$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(a*tanh(x)^3)^(1/2),x,method=_RETURNVERBOSE)
```

```
[Out] -tanh(x)*(2*a^(5/2)+arctan((a*tanh(x))^(1/2)/a^(1/2))*a^2*(a*tanh(x))^(1/2)
-arctanh((a*tanh(x))^(1/2)/a^(1/2))*a^2*(a*tanh(x))^(1/2))/(a*tanh(x)^3)^(1
/2)/a^(5/2)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(a*tanh(x)^3)^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(1/sqrt(a*tanh(x)^3), x)
```

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 239 vs. 2(50) = 100.

time = 0.45, size = 516, normalized size = 8.06



Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(a*tanh(x)^3)^(1/2),x, algorithm="fricas")
```

```
[Out] [-1/4*(2*(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - 1)*sqrt(-a)*arctan((c
osh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2)*sqrt(-a)*sqrt(a*sinh(x)/cosh(x)))/
(a*cosh(x)^2 + 2*a*cosh(x)*sinh(x) + a*sinh(x)^2 - a) + (cosh(x)^2 + 2*cos
h(x)*sinh(x) + sinh(x)^2 - 1)*sqrt(-a)*log(-(a*cosh(x)^4 + 4*a*cosh(x)^3*si
nh(x) + 6*a*cosh(x)^2*sinh(x)^2 + 4*a*cosh(x)*sinh(x)^3 + a*sinh(x)^4 + 2*(
cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)*sqrt(-a)*sqrt(a*sinh(x)/cosh
(x)) - 2*a)/(cosh(x)^4 + 4*cosh(x)^3*sinh(x) + 6*cosh(x)^2*sinh(x)^2 + 4*co
sh(x)*sinh(x)^3 + sinh(x)^4)) + 8*(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^
2 + 1)*sqrt(a*sinh(x)/cosh(x)))/(a*cosh(x)^2 + 2*a*cosh(x)*sinh(x) + a*sinh
```


$(x)^2 - a)$, $-1/4*(2*(\cosh(x)^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2 - 1)*\sqrt{a}*\arctan((\cosh(x)^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2 + 1)*\sqrt{a*\sinh(x)/\cosh(x)})/\sqrt{a}) - (\cosh(x)^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2 - 1)*\sqrt{a}*\log(2*a*\cosh(x)^4 + 8*a*\cosh(x)^3*\sinh(x) + 12*a*\cosh(x)^2*\sinh(x)^2 + 8*a*\cosh(x)*\sinh(x)^3 + 2*a*\sinh(x)^4 + 2*(\cosh(x)^4 + 4*\cosh(x)*\sinh(x)^3 + \sinh(x)^4 + (6*\cosh(x)^2 + 1)*\sinh(x)^2 + \cosh(x)^2 + 2*(2*\cosh(x)^3 + \cosh(x))*\sinh(x))*\sqrt{a}*\sqrt{a*\sinh(x)/\cosh(x)} - a) + 8*(\cosh(x)^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2 + 1)*\sqrt{a*\sinh(x)/\cosh(x)})/(a*\cosh(x)^2 + 2*a*\cosh(x)*\sinh(x) + a*\sinh(x)^2 - a)]$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{a \tanh^3(x)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*tanh(x)**3)**(1/2),x)

[Out] Integral(1/sqrt(a*tanh(x)**3), x)

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 123 vs. 2(50) = 100.

time = 0.43, size = 123, normalized size = 1.92

$$-\frac{\arctan\left(\frac{-\sqrt{a}e^{2x}-\sqrt{ae^{4x}-a}}{\sqrt{a}}\right)}{\sqrt{a}\operatorname{sgn}(e^{4x}-1)} - \frac{\log\left(\left|-\sqrt{a}e^{2x}+\sqrt{ae^{4x}-a}\right|\right)}{2\sqrt{a}\operatorname{sgn}(e^{4x}-1)} + \frac{4}{\left(\sqrt{a}e^{2x}-\sqrt{ae^{4x}-a}-\sqrt{a}\right)\operatorname{sgn}(e^{4x}-1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*tanh(x)^3)^(1/2),x, algorithm="giac")

[Out] $-\arctan(-(\sqrt{a}*e^{2*x} - \sqrt{a*e^{4*x} - a})/\sqrt{a})/(\sqrt{a}*\operatorname{sgn}(e^{4*x} - 1)) - 1/2*\log(\operatorname{abs}(-\sqrt{a}*e^{2*x} + \sqrt{a*e^{4*x} - a}))/(\sqrt{a}*\operatorname{sgn}(e^{4*x} - 1)) + 4/((\sqrt{a}*e^{2*x} - \sqrt{a*e^{4*x} - a} - \sqrt{a})*\operatorname{sgn}(e^{4*x} - 1))$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\sqrt{a \tanh^3(x)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a*tanh(x)^3)^(1/2),x)

[Out] int(1/(a*tanh(x)^3)^(1/2), x)

3.37 $\int (a \tanh^4(x))^{3/2} dx$

Optimal. Leaf size=69

$$-a \coth(x) \sqrt{a \tanh^4(x)} + ax \coth^2(x) \sqrt{a \tanh^4(x)} - \frac{1}{3} a \tanh(x) \sqrt{a \tanh^4(x)} - \frac{1}{5} a \tanh^3(x) \sqrt{a \tanh^4(x)}$$

[Out] $-a \coth(x) * (a * \tanh(x)^4)^{(1/2)} + a * x * \coth(x)^2 * (a * \tanh(x)^4)^{(1/2)} - 1/3 * a * (a * \tanh(x)^4)^{(1/2)} * \tanh(x) - 1/5 * a * (a * \tanh(x)^4)^{(1/2)} * \tanh(x)^3$

Rubi [A]

time = 0.02, antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3739, 3554, 8}

$$-\frac{1}{3} a \tanh(x) \sqrt{a \tanh^4(x)} - \frac{1}{5} a \tanh^3(x) \sqrt{a \tanh^4(x)} + ax \coth^2(x) \sqrt{a \tanh^4(x)} - a \coth(x) \sqrt{a \tanh^4(x)}$$

Antiderivative was successfully verified.

[In] Int[(a*Tanh[x]^4)^(3/2), x]

[Out] $-(a * \text{Coth}[x] * \text{Sqrt}[a * \text{Tanh}[x]^4]) + a * x * \text{Coth}[x]^2 * \text{Sqrt}[a * \text{Tanh}[x]^4] - (a * \text{Tanh}[x] * \text{Sqrt}[a * \text{Tanh}[x]^4]) / 3 - (a * \text{Tanh}[x]^3 * \text{Sqrt}[a * \text{Tanh}[x]^4]) / 5$

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 3554

Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Dist[b^2, Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]

Rule 3739

Int[(u_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[(b*ff^n)^IntPart[p]*((b*Tan[e + f*x]^n)^FracPart[p]/(Tan[e + f*x]/ff)^(n*FracPart[p])), Int[ActivateTrig[u]*(Tan[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]

Rubi steps

$$\begin{aligned}
\int (a \tanh^4(x))^{3/2} dx &= \left(a \coth^2(x) \sqrt{a \tanh^4(x)} \right) \int \tanh^6(x) dx \\
&= -\frac{1}{5} a \tanh^3(x) \sqrt{a \tanh^4(x)} + \left(a \coth^2(x) \sqrt{a \tanh^4(x)} \right) \int \tanh^4(x) dx \\
&= -\frac{1}{3} a \tanh(x) \sqrt{a \tanh^4(x)} - \frac{1}{5} a \tanh^3(x) \sqrt{a \tanh^4(x)} + \left(a \coth^2(x) \sqrt{a \tanh^4(x)} \right) \int \tanh^2(x) dx \\
&= -a \coth(x) \sqrt{a \tanh^4(x)} - \frac{1}{3} a \tanh(x) \sqrt{a \tanh^4(x)} - \frac{1}{5} a \tanh^3(x) \sqrt{a \tanh^4(x)} \\
&= -a \coth(x) \sqrt{a \tanh^4(x)} + ax \coth^2(x) \sqrt{a \tanh^4(x)} - \frac{1}{3} a \tanh(x) \sqrt{a \tanh^4(x)}
\end{aligned}$$

Mathematica [A]

time = 0.11, size = 56, normalized size = 0.81

$$\frac{1}{240} \coth(x) \operatorname{csch}^5(x) (150x \cosh(x) + 75x \cosh(3x) + 15x \cosh(5x) - 50 \sinh(x) - 25 \sinh(3x) - 23 \sinh(5x)) (a \tanh^4(x))^{3/2}$$

Antiderivative was successfully verified.

[In] Integrate[(a*Tanh[x]^4)^(3/2),x]**[Out]** (Coth[x]*Csch[x]^5*(150*x*Cosh[x] + 75*x*Cosh[3*x] + 15*x*Cosh[5*x] - 50*Sinh[x] - 25*Sinh[3*x] - 23*Sinh[5*x])*(a*Tanh[x]^4)^(3/2))/240**Maple [A]**

time = 0.73, size = 46, normalized size = 0.67

method	result	size
derivativedivides	$-\frac{(a(\tanh^4(x)))^{\frac{3}{2}}(6(\tanh^5(x))+10(\tanh^3(x))+15\ln(\tanh(x)-1)-15\ln(1+\tanh(x))+30\tanh(x))}{30\tanh(x)^6}$	46
default	$-\frac{(a(\tanh^4(x)))^{\frac{3}{2}}(6(\tanh^5(x))+10(\tanh^3(x))+15\ln(\tanh(x)-1)-15\ln(1+\tanh(x))+30\tanh(x))}{30\tanh(x)^6}$	46
risch	$\frac{a(1+e^{2x})^2 \sqrt{\frac{a(e^{2x}-1)^4}{(1+e^{2x})^4}} x}{(e^{2x}-1)^2} + \frac{2a \sqrt{\frac{a(e^{2x}-1)^4}{(1+e^{2x})^4}} (45e^{8x}+90e^{6x}+140e^{4x}+70e^{2x}+23)}{15(e^{2x}-1)^2(1+e^{2x})^3}$	106

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*tanh(x)^4)^(3/2),x,method=_RETURNVERBOSE)**[Out]** -1/30*(a*tanh(x)^4)^(3/2)*(6*tanh(x)^5+10*tanh(x)^3+15*ln(tanh(x)-1)-15*ln(1+tanh(x))+30*tanh(x))/tanh(x)^6

Maxima [A]

time = 0.49, size = 82, normalized size = 1.19

$$a^{\frac{3}{2}}x - \frac{2 \left(70 a^{\frac{3}{2}} e^{-2x} + 140 a^{\frac{3}{2}} e^{-4x} + 90 a^{\frac{3}{2}} e^{-6x} + 45 a^{\frac{3}{2}} e^{-8x} + 23 a^{\frac{3}{2}} \right)}{15 \left(5 e^{-2x} + 10 e^{-4x} + 10 e^{-6x} + 5 e^{-8x} + e^{-10x} + 1 \right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*tanh(x)^4)^(3/2),x, algorithm="maxima")

[Out] $a^{3/2}x - 2/15*(70*a^{3/2}*e^{-2*x} + 140*a^{3/2}*e^{-4*x} + 90*a^{3/2}*e^{-6*x} + 45*a^{3/2}*e^{-8*x} + 23*a^{3/2})/(5*e^{-2*x} + 10*e^{-4*x} + 10*e^{-6*x} + 5*e^{-8*x} + e^{-10*x} + 1)$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 2114 vs. 2(57) = 114.

time = 0.44, size = 2114, normalized size = 30.64

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*tanh(x)^4)^(3/2),x, algorithm="fricas")

[Out] $1/15*(15*a*x*\cosh(x)^{10} + 15*(a*x*e^{4*x} + 2*a*x*e^{2*x} + a*x)*\sinh(x)^{10} + 150*(a*x*\cosh(x)*e^{4*x} + 2*a*x*\cosh(x)*e^{2*x} + a*x*\cosh(x))*\sinh(x)^9 + 15*(5*a*x + 6*a)*\cosh(x)^8 + 15*(45*a*x*\cosh(x)^2 + 5*a*x + (45*a*x*\cosh(x)^2 + 5*a*x + 6*a)*e^{4*x} + 2*(45*a*x*\cosh(x)^2 + 5*a*x + 6*a)*e^{2*x} + 6*a)*\sinh(x)^8 + 120*(15*a*x*\cosh(x)^3 + (5*a*x + 6*a)*\cosh(x) + (15*a*x*\cosh(x)^3 + (5*a*x + 6*a)*\cosh(x))*e^{4*x} + 2*(15*a*x*\cosh(x)^3 + (5*a*x + 6*a)*\cosh(x))*e^{2*x})*\sinh(x)^7 + 30*(5*a*x + 6*a)*\cosh(x)^6 + 30*(105*a*x*\cosh(x)^4 + 14*(5*a*x + 6*a)*\cosh(x)^2 + 5*a*x + (105*a*x*\cosh(x)^4 + 14*(5*a*x + 6*a)*\cosh(x)^2 + 5*a*x + 6*a)*e^{4*x} + 2*(105*a*x*\cosh(x)^4 + 14*(5*a*x + 6*a)*\cosh(x)^2 + 5*a*x + 6*a)*e^{2*x} + 6*a)*\sinh(x)^6 + 60*(63*a*x*\cosh(x)^5 + 14*(5*a*x + 6*a)*\cosh(x)^3 + 3*(5*a*x + 6*a)*\cosh(x) + (63*a*x*\cosh(x)^5 + 14*(5*a*x + 6*a)*\cosh(x)^3 + 3*(5*a*x + 6*a)*\cosh(x))*e^{4*x} + 2*(63*a*x*\cosh(x)^5 + 14*(5*a*x + 6*a)*\cosh(x)^3 + 3*(5*a*x + 6*a)*\cosh(x))*e^{2*x})*\sinh(x)^5 + 10*(15*a*x + 28*a)*\cosh(x)^4 + 10*(315*a*x*\cosh(x)^6 + 105*(5*a*x + 6*a)*\cosh(x)^4 + 45*(5*a*x + 6*a)*\cosh(x)^2 + 15*a*x + (315*a*x*\cosh(x)^6 + 105*(5*a*x + 6*a)*\cosh(x)^4 + 45*(5*a*x + 6*a)*\cosh(x)^2 + 15*a*x + 28*a)*e^{4*x} + 2*(315*a*x*\cosh(x)^6 + 105*(5*a*x + 6*a)*\cosh(x)^4 + 45*(5*a*x + 6*a)*\cosh(x)^2 + 15*a*x + 28*a)*e^{2*x} + 28*a)*\sinh(x)^4 + 40*(45*a*x*\cosh(x)^7 + 21*(5*a*x + 6*a)*\cosh(x)^5 + 15*(5*a*x + 6*a)*\cosh(x)^3 + (15*a*x + 28*a)*\cosh(x) + (45*a*x*\cosh(x)^7 + 21*(5*a*x + 6*a)*\cosh(x)^5 + 15*(5*a*x + 6*a)*\cosh(x)^3 + (15*a*x + 28*a)*\cosh(x))*e^{4*x} + 2*(45*a*x*\cosh(x)^7 + 21*(5*a*x + 6*a)*\cosh(x)^5 + 15*(5*a*x + 6*a)*\cosh(x)^3 + (15*a*x + 28*a)*\cosh(x))*e^{2*x})*\sinh(x)^3 + 5*(15*a*x + 28*a)*\cosh(x)^$

$2 + 5*(135*a*x*cosh(x)^8 + 84*(5*a*x + 6*a)*cosh(x)^6 + 90*(5*a*x + 6*a)*cosh(x)^4 + 12*(15*a*x + 28*a)*cosh(x)^2 + 15*a*x + (135*a*x*cosh(x)^8 + 84*(5*a*x + 6*a)*cosh(x)^6 + 90*(5*a*x + 6*a)*cosh(x)^4 + 12*(15*a*x + 28*a)*cosh(x)^2 + 15*a*x + 28*a)*e^{(4*x)} + 2*(135*a*x*cosh(x)^8 + 84*(5*a*x + 6*a)*cosh(x)^6 + 90*(5*a*x + 6*a)*cosh(x)^4 + 12*(15*a*x + 28*a)*cosh(x)^2 + 15*a*x + 28*a)*e^{(2*x)} + 28*a)*sinh(x)^2 + 15*a*x + (15*a*x*cosh(x)^10 + 15*(5*a*x + 6*a)*cosh(x)^8 + 30*(5*a*x + 6*a)*cosh(x)^6 + 10*(15*a*x + 28*a)*cosh(x)^4 + 5*(15*a*x + 28*a)*cosh(x)^2 + 15*a*x + 46*a)*e^{(4*x)} + 2*(15*a*x*cosh(x)^10 + 15*(5*a*x + 6*a)*cosh(x)^8 + 30*(5*a*x + 6*a)*cosh(x)^6 + 10*(15*a*x + 28*a)*cosh(x)^4 + 5*(15*a*x + 28*a)*cosh(x)^2 + 15*a*x + 46*a)*e^{(2*x)} + 10*(15*a*x*cosh(x)^9 + 12*(5*a*x + 6*a)*cosh(x)^7 + 18*(5*a*x + 6*a)*cosh(x)^5 + 4*(15*a*x + 28*a)*cosh(x)^3 + (15*a*x + 28*a)*cosh(x))*e^{(4*x)} + 2*(15*a*x*cosh(x)^9 + 12*(5*a*x + 6*a)*cosh(x)^7 + 18*(5*a*x + 6*a)*cosh(x)^5 + 4*(15*a*x + 28*a)*cosh(x)^3 + (15*a*x + 28*a)*cosh(x))*e^{(2*x)})*sinh(x) + 46*a)*sqrt((a*e^{(8*x)} - 4*a*e^{(6*x)} + 6*a*e^{(4*x)} - 4*a*e^{(2*x)} + a)/(e^{(8*x)} + 4*e^{(6*x)} + 6*e^{(4*x)} + 4*e^{(2*x)} + 1))/((e^{(4*x)} - 2*e^{(2*x)} + 1)*sinh(x)^10 + cosh(x)^10 + 10*(cosh(x)*e^{(4*x)} - 2*cosh(x)*e^{(2*x)} + cosh(x))*sinh(x)^9 + 5*(9*cosh(x)^2 + (9*cosh(x)^2 + 1)*e^{(4*x)} - 2*(9*cosh(x)^2 + 1)*e^{(2*x)} + 1)*sinh(x)^8 + 5*cosh(x)^8 + 40*(3*cosh(x)^3 + (3*cosh(x)^3 + cosh(x))*e^{(4*x)} - 2*(3*cosh(x)^3 + cosh(x))*e^{(2*x)} + cosh(x))*sinh(x)^7 + 10*(21*cosh(x)^4 + 14*cosh(x)^2 + (21*cosh(x)^4 + 14*cosh(x)^2 + 1)*e^{(4*x)} - 2*(21*cosh(x)^4 + 14*cosh(x)^2 + 1)*e^{(2*x)} + 1)*sinh(x)^6 + 10*cosh(x)^6 + 4*(63*cosh(x)^5 + 70*cosh(x)^3 + (63*cosh(x)^5 + 70*cosh(x)^3 + 15*cosh(x))*e^{(4*x)} - 2*(63*cosh(x)^5 + 70*cosh(x)^3 + 15*cosh(x))*e^{(2*x)} + 15*cosh(x))*sinh(x)^5 + 10*(21*cosh(x)^6 + 35*cosh(x)^4 + 15*cosh(x)^2 + (21*cosh(x)^6 + 35*cosh(x)^4 + 15*cosh(x)^2 + 1)*e^{(4*x)} - 2*(21*cosh(x)^6 + 35*cosh(x)^4 + 15*cosh(x)^2 + 1)*e^{(2*x)} + 1)*sinh(x)^4 + 10*cosh(x)^4 + 40*(3*cosh(x)^7 + 7*cosh(x)^5 + 5*cosh(x)^3 + (3*cosh(x)^7 + 7*cosh(x)^5 + 5*cosh(x)^3 + cosh(x))*e^{(4*x)} - 2*(3*cosh(x)^7 + 7*cosh(x)^5 + 5*cosh(x)^3 + cosh(x))*e^{(2*x)} + cosh(x))*sinh(x)^3 + 5*(9*cosh(x)^8 + 28*cosh(x)^6 + 30*cosh(x)^4 + 12*cosh(x)^2 + (9*cosh(x)^8 + 28*cosh(x)^6 + 30*cosh(x)^4 + 12*cosh(x)^2 + 1)*e^{(4*x)} - 2*(9*cosh(x)^8 + 28*cosh(x)^6 + 30*cosh(x)^4 + 12*cosh(x)^2 + 1)*e^{(2*x)} + 1)*sinh(x)^2 + 5*cosh(x)^2 + (cosh(x)^10 + 5*cosh(x)^8 + 10*cosh(x)^6 + 10*cosh(x)^4 + 5*cosh(x)^2 + 1)*e^{(4*x)} - 2*(cosh(x)^10 + 5*cosh(x)^8 + 10*cosh(x)^6 + 10*cosh(x)^4 + 5*cosh(x)^2 + 1)*e^{(2*x)} + 10*(cosh(x)^9 + 4*cosh(x)^7 + 6*cosh(x)^5 + 4*cosh(x)^3 + cosh(x))*e^{(4*x)} - 2*(cosh(x)^9 + 4*cosh(x)^7 + 6*cosh(x)^5 + 4*cosh(x)^3 + cosh(x))*e^{(2*x)} + cosh(x))*sinh(x) + 1)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (a \tanh^4(x))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*tanh(x)**4)**(3/2),x)

[Out] Integral((a*tanh(x)**4)**(3/2), x)

Giac [A]

time = 0.42, size = 45, normalized size = 0.65

$$\frac{1}{15} a^{\frac{3}{2}} \left(15x + \frac{2(45e^{8x} + 90e^{6x} + 140e^{4x} + 70e^{2x} + 23)}{(e^{2x} + 1)^5} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*tanh(x)^4)^(3/2),x, algorithm="giac")

[Out] 1/15*a^(3/2)*(15*x + 2*(45*e^(8*x) + 90*e^(6*x) + 140*e^(4*x) + 70*e^(2*x) + 23)/(e^(2*x) + 1)^5)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int (a \tanh(x)^4)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*tanh(x)^4)^(3/2),x)

[Out] int((a*tanh(x)^4)^(3/2), x)

3.38 $\int \sqrt{a \tanh^4(x)} dx$

Optimal. Leaf size=31

$$-\coth(x)\sqrt{a \tanh^4(x)} + x \coth^2(x)\sqrt{a \tanh^4(x)}$$

[Out] $-\coth(x)*(a*\tanh(x)^4)^{(1/2)}+x*\coth(x)^2*(a*\tanh(x)^4)^{(1/2)}$

Rubi [A]

time = 0.01, antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3739, 3554, 8}

$$x \coth^2(x)\sqrt{a \tanh^4(x)} - \coth(x)\sqrt{a \tanh^4(x)}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sqrt}[a*\text{Tanh}[x]^4], x]$

[Out] $-(\text{Coth}[x]*\text{Sqrt}[a*\text{Tanh}[x]^4]) + x*\text{Coth}[x]^2*\text{Sqrt}[a*\text{Tanh}[x]^4]$

Rule 8

$\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

Rule 3554

$\text{Int}[(b_*)*\tan[(c_*) + (d_*)(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Simp}[b*((b*\tan[c + d*x])^{(n-1)})/(d*(n-1)), x] - \text{Dist}[b^2, \text{Int}[(b*\tan[c + d*x])^{(n-2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x] \ \&\& \ \text{GtQ}[n, 1]$

Rule 3739

$\text{Int}[(u_)*((b_*)*\tan[(e_*) + (f_*)(x_)]^{(n_)})^{(p_)}, x_Symbol] \rightarrow \text{With}\{\text{ff} = \text{FreeFactors}[\tan[e + f*x], x]\}, \text{Dist}[(b*\text{ff}^n)^{\text{IntPart}[p]}*(b*\tan[e + f*x]^n)^{\text{FracPart}[p]} / (\tan[e + f*x]/\text{ff})^{(n*\text{FracPart}[p])}], \text{Int}[\text{ActivateTrig}[u]*(\tan[e + f*x]/\text{ff})^{(n*p)}, x], x] /; \text{FreeQ}\{b, e, f, n, p\}, x] \ \&\& \ !\text{IntegerQ}[p] \ \&\& \ \text{IntegerQ}[n] \ \&\& \ (\text{EqQ}[u, 1] \ || \ \text{MatchQ}[u, ((d_*)(\text{trig}_)[e + f*x])^{(m_)}]) /; \text{FreeQ}\{d, m\}, x] \ \&\& \ \text{MemberQ}\{\sin, \cos, \tan, \cot, \sec, \csc\}, \text{trig}])$

Rubi steps

$$\begin{aligned}
\int \sqrt{a \tanh^4(x)} dx &= \left(\coth^2(x) \sqrt{a \tanh^4(x)} \right) \int \tanh^2(x) dx \\
&= -\coth(x) \sqrt{a \tanh^4(x)} + \left(\coth^2(x) \sqrt{a \tanh^4(x)} \right) \int 1 dx \\
&= -\coth(x) \sqrt{a \tanh^4(x)} + x \coth^2(x) \sqrt{a \tanh^4(x)}
\end{aligned}$$

Mathematica [A]

time = 0.01, size = 19, normalized size = 0.61

$$\coth(x)(-1 + x \coth(x)) \sqrt{a \tanh^4(x)}$$

Antiderivative was successfully verified.

`[In] Integrate[Sqrt[a*Tanh[x]^4], x]``[Out] Coth[x]*(-1 + x*Coth[x])*Sqrt[a*Tanh[x]^4]`**Maple [A]**

time = 0.72, size = 32, normalized size = 1.03

method	result	size
derivativedivides	$-\frac{\sqrt{a(\tanh^4(x))} (2 \tanh(x) + \ln(\tanh(x)-1) - \ln(1+\tanh(x)))}{2 \tanh(x)^2}$	32
default	$-\frac{\sqrt{a(\tanh^4(x))} (2 \tanh(x) + \ln(\tanh(x)-1) - \ln(1+\tanh(x)))}{2 \tanh(x)^2}$	32
risch	$\frac{\sqrt{\frac{a(e^{2x}-1)^4}{(1+e^{2x})^4}} (1+e^{2x})^2 x}{(e^{2x}-1)^2} + \frac{2 \sqrt{\frac{a(e^{2x}-1)^4}{(1+e^{2x})^4}} (1+e^{2x})}{(e^{2x}-1)^2}$	76

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((a*tanh(x)^4)^(1/2), x, method=_RETURNVERBOSE)``[Out] -1/2*(a*tanh(x)^4)^(1/2)*(2*tanh(x)+ln(tanh(x)-1)-ln(1+tanh(x)))/tanh(x)^2`**Maxima [A]**

time = 0.48, size = 19, normalized size = 0.61

$$\sqrt{a} x - \frac{2 \sqrt{a}}{e^{(-2x)} + 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*tanh(x)^4)^(1/2),x, algorithm="maxima")

[Out] sqrt(a)*x - 2*sqrt(a)/(e^(-2*x) + 1)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 213 vs. 2(27) = 54.

time = 0.37, size = 213, normalized size = 6.87

$$\frac{(x \cosh(x)^2 + (x e^{4x} + 2x e^{2x} + x) \sinh(x)^2 + (x \cosh(x)^2 + x + 2) e^{4x} + 2(x \cosh(x)^2 + x + 2) e^{2x} + 2(x \cosh(x) e^{4x} + 2x \cosh(x) e^{2x} + x \cosh(x) \sinh(x) + x + 2) \sqrt{\frac{a e^{8x} - 4a e^{6x} + 6a e^{4x} - 4a e^{2x} + a}{e^{8x} + 4e^{6x} + 6e^{4x} + 4e^{2x} + 1}}}{(e^{4x} - 2e^{2x} + 1) \sinh(x)^2 + \cosh(x)^2 + (\cosh(x)^2 + 1) e^{4x} - 2(\cosh(x)^2 + 1) e^{2x} + 2(\cosh(x) e^{4x} - 2 \cosh(x) e^{2x} + \cosh(x) \sinh(x) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*tanh(x)^4)^(1/2),x, algorithm="fricas")

[Out] (x*cosh(x)^2 + (x*e^(4*x) + 2*x*e^(2*x) + x)*sinh(x)^2 + (x*cosh(x)^2 + x + 2)*e^(4*x) + 2*(x*cosh(x)^2 + x + 2)*e^(2*x) + 2*(x*cosh(x)*e^(4*x) + 2*x*cosh(x)*e^(2*x) + x*cosh(x))*sinh(x) + x + 2)*sqrt((a*e^(8*x) - 4*a*e^(6*x) + 6*a*e^(4*x) - 4*a*e^(2*x) + a)/(e^(8*x) + 4*e^(6*x) + 6*e^(4*x) + 4*e^(2*x) + 1))/((e^(4*x) - 2*e^(2*x) + 1)*sinh(x)^2 + cosh(x)^2 + (cosh(x)^2 + 1)*e^(4*x) - 2*(cosh(x)^2 + 1)*e^(2*x) + 2*(cosh(x)*e^(4*x) - 2*cosh(x)*e^(2*x) + cosh(x))*sinh(x) + 1)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{a \tanh^4(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*tanh(x)**4)**(1/2),x)

[Out] Integral(sqrt(a*tanh(x)**4), x)

Giac [A]

time = 0.42, size = 16, normalized size = 0.52

$$\sqrt{a} \left(x + \frac{2}{e^{2x} + 1} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*tanh(x)^4)^(1/2),x, algorithm="giac")

[Out] sqrt(a)*(x + 2/(e^(2*x) + 1))

Mupad [F]

time = 0.00, size = -1, normalized size = -0.03

$$\int \sqrt{a \tanh(x)^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((a*tanh(x)^4)^(1/2),x)
```

```
[Out] int((a*tanh(x)^4)^(1/2), x)
```

$$3.39 \quad \int \frac{1}{\sqrt{a \tanh^4(x)}} dx$$

Optimal. Leaf size=31

$$-\frac{\tanh(x)}{\sqrt{a \tanh^4(x)}} + \frac{x \tanh^2(x)}{\sqrt{a \tanh^4(x)}}$$

[Out] $-\tanh(x)/(a*\tanh(x)^4)^{(1/2)}+x*\tanh(x)^2/(a*\tanh(x)^4)^{(1/2)}$

Rubi [A]

time = 0.01, antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3739, 3554, 8}

$$\frac{x \tanh^2(x)}{\sqrt{a \tanh^4(x)}} - \frac{\tanh(x)}{\sqrt{a \tanh^4(x)}}$$

Antiderivative was successfully verified.

[In] Int[1/Sqrt[a*Tanh[x]^4],x]

[Out] $-(\text{Tanh}[x]/\text{Sqrt}[a*\text{Tanh}[x]^4]) + (x*\text{Tanh}[x]^2)/\text{Sqrt}[a*\text{Tanh}[x]^4]$

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 3554

Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Dist[b^2, Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]

Rule 3739

Int[(u_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[(b*ff^n)^IntPart[p]*((b*Tan[e + f*x])^n)^FracPart[p]/(Tan[e + f*x]/ff)^(n*FracPart[p])], Int[ActivateTrig[u]*(Tan[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]

Rubi steps

$$\begin{aligned}
\int \frac{1}{\sqrt{a \tanh^4(x)}} dx &= \frac{\tanh^2(x) \int \coth^2(x) dx}{\sqrt{a \tanh^4(x)}} \\
&= -\frac{\tanh(x)}{\sqrt{a \tanh^4(x)}} + \frac{\tanh^2(x) \int 1 dx}{\sqrt{a \tanh^4(x)}} \\
&= -\frac{\tanh(x)}{\sqrt{a \tanh^4(x)}} + \frac{x \tanh^2(x)}{\sqrt{a \tanh^4(x)}}
\end{aligned}$$

Mathematica [A]

time = 0.03, size = 19, normalized size = 0.61

$$\frac{\tanh(x)(-1 + x \tanh(x))}{\sqrt{a \tanh^4(x)}}$$

Antiderivative was successfully verified.

`[In] Integrate[1/Sqrt[a*Tanh[x]^4], x]``[Out] (Tanh[x]*(-1 + x*Tanh[x]))/Sqrt[a*Tanh[x]^4]`**Maple [A]**

time = 0.74, size = 32, normalized size = 1.03

method	result	size
derivativedivides	$\frac{\tanh(x)(\ln(1+\tanh(x)) \tanh(x) - \ln(\tanh(x)-1) \tanh(x)-2)}{2\sqrt{a(\tanh^4(x))}}$	32
default	$\frac{\tanh(x)(\ln(1+\tanh(x)) \tanh(x) - \ln(\tanh(x)-1) \tanh(x)-2)}{2\sqrt{a(\tanh^4(x))}}$	32
risch	$\frac{(e^{2x}-1)^2 x}{\sqrt{\frac{a(e^{2x}-1)^4}{(1+e^{2x})^4}} (1+e^{2x})^2} - \frac{2(e^{2x}-1)}{\sqrt{\frac{a(e^{2x}-1)^4}{(1+e^{2x})^4}} (1+e^{2x})^2}}$	76

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/(a*tanh(x)^4)^(1/2), x, method=_RETURNVERBOSE)``[Out] 1/2*tanh(x)*(ln(1+tanh(x))*tanh(x)-ln(tanh(x)-1)*tanh(x)-2)/(a*tanh(x)^4)^(1/2)`**Maxima [A]**

time = 0.48, size = 23, normalized size = 0.74

$$\frac{x}{\sqrt{a}} + \frac{2\sqrt{a}}{ae^{(-2x)} - a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*tanh(x)^4)^(1/2),x, algorithm="maxima")

[Out] x/sqrt(a) + 2*sqrt(a)/(a*e^(-2*x) - a)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 238 vs. 2(27) = 54.

time = 0.36, size = 238, normalized size = 7.68

$$\frac{(x \cosh(x)^2 + (x e^{4x} + 2x e^{2x} + x) \sinh(x)^2 + (x \cosh(x)^2 - x - 2) e^{4x} + 2(x \cosh(x)^2 - x - 2) e^{2x} + 2(x \cosh(x) e^{4x} + 2x \cosh(x) e^{2x} + x \cosh(x) \sinh(x) - x - 2) \sqrt{\frac{a e^{8x} - 4a e^{6x} + 6a e^{4x} - 4a e^{2x} + a}{e^{8x} + 4e^{6x} + 6e^{4x} + 4e^{2x} + 1}}}{a \cosh(x)^2 + (a e^{4x} - 2a e^{2x} + a) \sinh(x)^2 + (a \cosh(x)^2 - a) e^{4x} - 2(a \cosh(x)^2 - a) e^{2x} + 2(a \cosh(x) e^{4x} - 2a \cosh(x) e^{2x} + a \cosh(x) \sinh(x) - a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*tanh(x)^4)^(1/2),x, algorithm="fricas")

[Out] (x*cosh(x)^2 + (x*e^(4*x) + 2*x*e^(2*x) + x)*sinh(x)^2 + (x*cosh(x)^2 - x - 2)*e^(4*x) + 2*(x*cosh(x)^2 - x - 2)*e^(2*x) + 2*(x*cosh(x)*e^(4*x) + 2*x*cosh(x)*e^(2*x) + x*cosh(x))*sinh(x) - x - 2)*sqrt((a*e^(8*x) - 4*a*e^(6*x) + 6*a*e^(4*x) - 4*a*e^(2*x) + a)/(e^(8*x) + 4*e^(6*x) + 6*e^(4*x) + 4*e^(2*x) + 1))/(a*cosh(x)^2 + (a*e^(4*x) - 2*a*e^(2*x) + a)*sinh(x)^2 + (a*cosh(x)^2 - a)*e^(4*x) - 2*(a*cosh(x)^2 - a)*e^(2*x) + 2*(a*cosh(x)*e^(4*x) - 2*a*cosh(x)*e^(2*x) + a*cosh(x))*sinh(x) - a)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{a \tanh^4(x)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*tanh(x)**4)**(1/2),x)

[Out] Integral(1/sqrt(a*tanh(x)**4), x)

Giac [A]

time = 0.41, size = 19, normalized size = 0.61

$$\frac{x}{\sqrt{a}} - \frac{2}{\sqrt{a} (e^{2x} - 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*tanh(x)^4)^(1/2),x, algorithm="giac")

[Out] x/sqrt(a) - 2/(sqrt(a)*(e^(2*x) - 1))

Mupad [F]

time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{1}{\sqrt{a \tanh(x)^4}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a*tanh(x)^4)^(1/2),x)

[Out] int(1/(a*tanh(x)^4)^(1/2), x)

3.40 $\int (b \tanh^m(c + dx))^n dx$

Optimal. Leaf size=57

$$\frac{{}_2F_1\left(1, \frac{1}{2}(1 + mn); \frac{1}{2}(3 + mn); \tanh^2(c + dx)\right) \tanh(c + dx) (b \tanh^m(c + dx))^n}{d(1 + mn)}$$

[Out] hypergeom([1, 1/2*m*n+1/2], [1/2*m*n+3/2], tanh(d*x+c)^2)*tanh(d*x+c)*(b*tanh(d*x+c)^m)^n/d/(m*n+1)

Rubi [A]

time = 0.03, antiderivative size = 57, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3740, 3557, 371}

$$\frac{\tanh(c + dx) (b \tanh^m(c + dx))^n {}_2F_1\left(1, \frac{1}{2}(mn + 1); \frac{1}{2}(mn + 3); \tanh^2(c + dx)\right)}{d(mn + 1)}$$

Antiderivative was successfully verified.

[In] Int[(b*Tanh[c + d*x]^m)^n,x]

[Out] (Hypergeometric2F1[1, (1 + m*n)/2, (3 + m*n)/2, Tanh[c + d*x]^2]*Tanh[c + d*x]*(b*Tanh[c + d*x]^m)^n)/(d*(1 + m*n))

Rule 371

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[a^p *((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])

Rule 3557

Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Dist[b/d, Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rule 3740

Int[(u_.)*((b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] :> Dist[b^IntPart[p]*((b*(c*Tan[e + f*x])^n)^FracPart[p]/(c*Tan[e + f*x])^(n*FracPart[p])), Int[ActivateTrig[u]*(c*Tan[e + f*x])^(n*p), x], x] /; FreeQ[{b, c, e, f, n, p}, x] && !IntegerQ[p] && !IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig]])

Rubi steps

$$\begin{aligned}
\int (b \tanh^m(c + dx))^n dx &= (\tanh^{-mn}(c + dx) (b \tanh^m(c + dx))^n) \int \tanh^{mn}(c + dx) dx \\
&= -\frac{(\tanh^{-mn}(c + dx) (b \tanh^m(c + dx))^n) \operatorname{Subst}\left(\int \frac{x^{mn}}{-1+x^2} dx, x, \tanh(c + dx)\right)}{d} \\
&= \frac{{}_2F_1\left(1, \frac{1}{2}(1 + mn); \frac{1}{2}(3 + mn); \tanh^2(c + dx)\right) \tanh(c + dx) (b \tanh^m(c + dx))^n}{d(1 + mn)}
\end{aligned}$$

Mathematica [A]

time = 0.04, size = 55, normalized size = 0.96

$$\frac{{}_2F_1\left(1, \frac{1}{2}(1 + mn); \frac{1}{2}(3 + mn); \tanh^2(c + dx)\right) \tanh(c + dx) (b \tanh^m(c + dx))^n}{d + dm n}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Tanh[c + d*x]^m)^n,x]

[Out] (Hypergeometric2F1[1, (1 + m*n)/2, (3 + m*n)/2, Tanh[c + d*x]^2]*Tanh[c + d*x]*(b*Tanh[c + d*x]^m)^n)/(d + d*m*n)

Maple [F]

time = 180.00, size = 0, normalized size = 0.00

$$\int (b(\tanh^m(dx + c)))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*tanh(d*x+c)^m)^n,x)

[Out] int((b*tanh(d*x+c)^m)^n,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*tanh(d*x+c)^m)^n,x, algorithm="maxima")

[Out] integrate((b*tanh(d*x + c)^m)^n, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*tanh(d*x+c)^m)^n,x, algorithm="fricas")

[Out] integral((b*tanh(d*x + c)^m)^n, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (b \tanh^m(c + dx))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*tanh(d*x+c)**m)**n,x)

[Out] Integral((b*tanh(c + d*x)**m)**n, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*tanh(d*x+c)^m)^n,x, algorithm="giac")

[Out] integrate((b*tanh(d*x + c)^m)^n, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int (b \tanh(c + dx)^m)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*tanh(c + d*x)^m)^n,x)

[Out] int((b*tanh(c + d*x)^m)^n, x)

3.41 $\int (a + a \tanh(c + dx))^5 dx$

Optimal. Leaf size=100

$$16a^5x + \frac{16a^5 \log(\cosh(c + dx))}{d} - \frac{8a^5 \tanh(c + dx)}{d} - \frac{2a^2(a + a \tanh(c + dx))^3}{3d} - \frac{a(a + a \tanh(c + dx))^4}{4d} - \frac{2a(a^2 + a^2 \tanh(c + dx))^2}{d}$$

[Out] 16*a^5*x+16*a^5*ln(cosh(d*x+c))/d-8*a^5*tanh(d*x+c)/d-2/3*a^2*(a+a*tanh(d*x+c))^3/d-1/4*a*(a+a*tanh(d*x+c))^4/d-2*a*(a^2+a^2*tanh(d*x+c))^2/d

Rubi [A]

time = 0.06, antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3559, 3558, 3556}

$$-\frac{8a^5 \tanh(c + dx)}{d} + \frac{16a^5 \log(\cosh(c + dx))}{d} + 16a^5x - \frac{2a^2(a \tanh(c + dx) + a)^3}{3d} - \frac{2a(a^2 \tanh(c + dx) + a^2)^2}{d} - \frac{a(a \tanh(c + dx) + a)^4}{4d}$$

Antiderivative was successfully verified.

[In] Int[(a + a*Tanh[c + d*x])^5, x]

[Out] 16*a^5*x + (16*a^5*Log[Cosh[c + d*x]])/d - (8*a^5*Tanh[c + d*x])/d - (2*a^2*(a + a*Tanh[c + d*x])^3)/(3*d) - (a*(a + a*Tanh[c + d*x])^4)/(4*d) - (2*a*(a^2 + a^2*Tanh[c + d*x])^2)/d

Rule 3556

Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]

Rule 3558

Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_.)])^2, x_Symbol] := Simp[(a^2 - b^2)*x, x] + (Dist[2*a*b, Int[Tan[c + d*x], x], x] + Simp[b^2*(Tan[c + d*x]/d), x]) /; FreeQ[{a, b, c, d}, x]

Rule 3559

Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_.)])^(n_), x_Symbol] := Simp[b*((a + b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] + Dist[2*a, Int[(a + b*Tan[c + d*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && GtQ[n, 1]

Rubi steps

$$\begin{aligned}
\int (a + a \tanh(c + dx))^5 dx &= -\frac{a(a + a \tanh(c + dx))^4}{4d} + (2a) \int (a + a \tanh(c + dx))^4 dx \\
&= -\frac{2a^2(a + a \tanh(c + dx))^3}{3d} - \frac{a(a + a \tanh(c + dx))^4}{4d} + (4a^2) \int (a + a \tanh(c + dx))^3 dx \\
&= -\frac{2a^3(a + a \tanh(c + dx))^2}{d} - \frac{2a^2(a + a \tanh(c + dx))^3}{3d} - \frac{a(a + a \tanh(c + dx))^4}{4d} \\
&= 16a^5x - \frac{8a^5 \tanh(c + dx)}{d} - \frac{2a^3(a + a \tanh(c + dx))^2}{d} - \frac{2a^2(a + a \tanh(c + dx))^3}{3d} \\
&= 16a^5x + \frac{16a^5 \log(\cosh(c + dx))}{d} - \frac{8a^5 \tanh(c + dx)}{d} - \frac{2a^3(a + a \tanh(c + dx))^2}{d}
\end{aligned}$$

Mathematica [B] Leaf count is larger than twice the leaf count of optimal. 202 vs. 2(100) = 200.

time = 0.80, size = 202, normalized size = 2.02

$a^5 \text{sech}(c) \cosh^5(c + dx) (18 \cosh(3c + 2dx) + 48d \cosh(3c + 2dx) + 12d \cosh(3c + 4dx) + 12d \cosh(3c + 4dx) + 48 \cosh(3c + 2dx) \log(\cosh(c + dx)) + 12 \cosh(3c + 4dx) \log(\cosh(c + dx)) + 12 \cosh(3c + 4dx) \log(\cosh(c + dx)) + 6 \cosh(c) (33 + 72dx + 72 \log(\cosh(c + dx))) + 75 \sinh(c) - 70 \sinh(c + 2dx) + 30 \sinh(3c + 2dx) - 25 \sinh(3c + 4dx)) / (12d)$

Antiderivative was successfully verified.

[In] Integrate[(a + a*Tanh[c + d*x])^5, x]

[Out] (a^5*Sech[c]*Sech[c + d*x]^4*(18*Cosh[3*c + 2*d*x] + 48*d*x*Cosh[3*c + 2*d*x] + 12*d*x*Cosh[3*c + 4*d*x] + 12*d*x*Cosh[5*c + 4*d*x] + 48*Cosh[3*c + 2*d*x]*Log[Cosh[c + d*x]] + 12*Cosh[3*c + 4*d*x]*Log[Cosh[c + d*x]] + 12*Cosh[5*c + 4*d*x]*Log[Cosh[c + d*x]] + 6*Cosh[c + 2*d*x]*(3 + 8*d*x + 8*Log[Cosh[c + d*x]]) + Cosh[c]*(33 + 72*d*x + 72*Log[Cosh[c + d*x]]) + 75*Sinh[c] - 70*Sinh[c + 2*d*x] + 30*Sinh[3*c + 2*d*x] - 25*Sinh[3*c + 4*d*x]))/(12*d)

Maple [A]

time = 0.26, size = 58, normalized size = 0.58

method	result	size
derivativedivides	$a^5 \left(\frac{-\frac{\tanh^4(dx+c)}{4} - \frac{5(\tanh^3(dx+c))}{3} - \frac{11(\tanh^2(dx+c))}{2} - 15 \tanh(dx+c) - 16 \ln(\tanh(dx+c)-1)}{d} \right)$	58
default	$a^5 \left(\frac{-\frac{\tanh^4(dx+c)}{4} - \frac{5(\tanh^3(dx+c))}{3} - \frac{11(\tanh^2(dx+c))}{2} - 15 \tanh(dx+c) - 16 \ln(\tanh(dx+c)-1)}{d} \right)$	58
risch	$-\frac{32a^5c}{d} + \frac{4a^5(48e^{6dx+6c} + 108e^{4dx+4c} + 88e^{2dx+2c} + 25)}{3d(1+e^{2dx+2c})^4} + \frac{16a^5 \ln(1+e^{2dx+2c})}{d}$	87

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a+a*tanh(d*x+c))^5,x,method=_RETURNVERBOSE)

[Out] $1/d*a^5*(-1/4*\tanh(d*x+c)^4-5/3*\tanh(d*x+c)^3-11/2*\tanh(d*x+c)^2-15*\tanh(d*x+c)-16*\ln(\tanh(d*x+c)-1))$

Maxima [B] Leaf count of result is larger than twice the leaf count of optimal. 302 vs. 2(96) = 192.

time = 0.48, size = 302, normalized size = 3.02

$$\frac{5}{3}a^5\left(3x + \frac{3c}{d} - \frac{4(3e^{(-2dx-2)} + 3e^{(-4dx-4)} + 2)}{d(3e^{(-2dx-2)} + 3e^{(-4dx-4)} + e^{(-6dx-6)} + 1)}\right) + a^5\left(x + \frac{c}{d} + \frac{\log(e^{(-2dx-2)} + 1)}{d} + \frac{4(e^{(-2dx-2)} + e^{(-4dx-4)} + e^{(-6dx-6)})}{d(4e^{(-2dx-2)} + 6e^{(-4dx-4)} + 4e^{(-6dx-6)} + e^{(-8dx-8)} + 1)}\right) + 10a^5\left(x + \frac{c}{d} + \frac{\log(e^{(-2dx-2)} + 1)}{d} + \frac{2e^{(-2dx-2)}}{d(2e^{(-2dx-2)} + e^{(-4dx-4)} + 1)}\right) + 10a^5\left(x + \frac{c}{d} - \frac{2}{d(e^{(-2dx-2)} + 1)}\right) + a^5x + \frac{5a^5 \log(\cosh(dx+c))}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+a*tanh(d*x+c))^5,x, algorithm="maxima")`

[Out] $5/3*a^5*(3*x + 3*c/d - 4*(3*e^{(-2*d*x - 2*c)} + 3*e^{(-4*d*x - 4*c)} + 2)/(d*(3*e^{(-2*d*x - 2*c)} + 3*e^{(-4*d*x - 4*c)} + e^{(-6*d*x - 6*c)} + 1))) + a^5*(x + c/d + \log(e^{(-2*d*x - 2*c)} + 1)/d + 4*(e^{(-2*d*x - 2*c)} + e^{(-4*d*x - 4*c)} + e^{(-6*d*x - 6*c)})/(d*(4*e^{(-2*d*x - 2*c)} + 6*e^{(-4*d*x - 4*c)} + 4*e^{(-6*d*x - 6*c)} + e^{(-8*d*x - 8*c)} + 1))) + 10*a^5*(x + c/d + \log(e^{(-2*d*x - 2*c)} + 1)/d + 2*e^{(-2*d*x - 2*c)}/(d*(2*e^{(-2*d*x - 2*c)} + e^{(-4*d*x - 4*c)} + 1))) + 10*a^5*(x + c/d - 2/(d*(e^{(-2*d*x - 2*c)} + 1))) + a^5*x + 5*a^5*\log(\cosh(d*x + c))/d$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 907 vs. 2(96) = 192.

time = 0.35, size = 907, normalized size = 9.07

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+a*tanh(d*x+c))^5,x, algorithm="fricas")`

[Out] $4/3*(48*a^5*\cosh(d*x + c)^6 + 288*a^5*\cosh(d*x + c)*\sinh(d*x + c)^5 + 48*a^5*\sinh(d*x + c)^6 + 108*a^5*\cosh(d*x + c)^4 + 88*a^5*\cosh(d*x + c)^2 + 25*a^5 + 36*(20*a^5*\cosh(d*x + c)^2 + 3*a^5)*\sinh(d*x + c)^4 + 48*(20*a^5*\cosh(d*x + c)^3 + 9*a^5*\cosh(d*x + c))*\sinh(d*x + c)^3 + 8*(90*a^5*\cosh(d*x + c)^4 + 81*a^5*\cosh(d*x + c)^2 + 11*a^5)*\sinh(d*x + c)^2 + 12*(a^5*\cosh(d*x + c)^8 + 8*a^5*\cosh(d*x + c)*\sinh(d*x + c)^7 + a^5*\sinh(d*x + c)^8 + 4*a^5*\cosh(d*x + c)^6 + 6*a^5*\cosh(d*x + c)^4 + 4*a^5*\cosh(d*x + c)^2 + 4*(7*a^5*\cosh(d*x + c)^2 + a^5)*\sinh(d*x + c)^6 + 8*(7*a^5*\cosh(d*x + c)^3 + 3*a^5*\cosh(d*x + c))*\sinh(d*x + c)^5 + a^5 + 2*(35*a^5*\cosh(d*x + c)^4 + 30*a^5*\cosh(d*x + c)^2 + 3*a^5)*\sinh(d*x + c)^4 + 8*(7*a^5*\cosh(d*x + c)^5 + 10*a^5*\cosh(d*x + c)^3 + 3*a^5*\cosh(d*x + c))*\sinh(d*x + c)^3 + 4*(7*a^5*\cosh(d*x + c)^6 + 15*a^5*\cosh(d*x + c)^4 + 9*a^5*\cosh(d*x + c)^2 + a^5)*\sinh(d*x + c)^2 + 8*(a^5*\cosh(d*x + c)^7 + 3*a^5*\cosh(d*x + c)^5 + 3*a^5*\cosh(d*x + c)^3 + a^5*\cosh(d*x + c))*\sinh(d*x + c))*\log(2*\cosh(d*x + c)/(cosh(d*x + c) - sinh(d*x + c))) + 16*(18*a^5*\cosh(d*x + c)^5 + 27*a^5*\cosh(d*x + c)^3 + 11*a^5*\cosh(d*x + c))*\sinh(d*x + c))/(d*\cosh(d*x + c)^8 + 8*d*\cosh(d*x + c)*\sinh$

$$(d*x + c)^7 + d*\sinh(d*x + c)^8 + 4*d*\cosh(d*x + c)^6 + 4*(7*d*\cosh(d*x + c)^2 + d)*\sinh(d*x + c)^6 + 8*(7*d*\cosh(d*x + c)^3 + 3*d*\cosh(d*x + c))*\sinh(d*x + c)^5 + 6*d*\cosh(d*x + c)^4 + 2*(35*d*\cosh(d*x + c)^4 + 30*d*\cosh(d*x + c)^2 + 3*d)*\sinh(d*x + c)^4 + 8*(7*d*\cosh(d*x + c)^5 + 10*d*\cosh(d*x + c)^3 + 3*d*\cosh(d*x + c))*\sinh(d*x + c)^3 + 4*d*\cosh(d*x + c)^2 + 4*(7*d*\cosh(d*x + c)^6 + 15*d*\cosh(d*x + c)^4 + 9*d*\cosh(d*x + c)^2 + d)*\sinh(d*x + c)^2 + 8*(d*\cosh(d*x + c)^7 + 3*d*\cosh(d*x + c)^5 + 3*d*\cosh(d*x + c)^3 + d*\cosh(d*x + c))*\sinh(d*x + c) + d$$

Sympy [A]

time = 0.14, size = 95, normalized size = 0.95

$$\begin{cases} 32a^5x - \frac{16a^5 \log(\tanh(c+dx)+1)}{d} - \frac{a^5 \tanh^4(c+dx)}{4d} - \frac{5a^5 \tanh^3(c+dx)}{3d} - \frac{11a^5 \tanh^2(c+dx)}{2d} - \frac{15a^5 \tanh(c+dx)}{d} & \text{for } d \neq 0 \\ x(a \tanh(c) + a)^5 & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+a*tanh(d*x+c))**5,x)

[Out] Piecewise((32*a**5*x - 16*a**5*log(tanh(c + d*x) + 1)/d - a**5*tanh(c + d*x)**4/(4*d) - 5*a**5*tanh(c + d*x)**3/(3*d) - 11*a**5*tanh(c + d*x)**2/(2*d) - 15*a**5*tanh(c + d*x)/d, Ne(d, 0)), (x*(a*tanh(c) + a)**5, True))

Giac [A]

time = 0.40, size = 85, normalized size = 0.85

$$\frac{4 \left(12 a^5 \log(e^{(2 dx+2 c)} + 1) + \frac{48 a^5 e^{(6 dx+6 c)} + 108 a^5 e^{(4 dx+4 c)} + 88 a^5 e^{(2 dx+2 c)} + 25 a^5}{(e^{(2 dx+2 c)} + 1)^4} \right)}{3 d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+a*tanh(d*x+c))^5,x, algorithm="giac")

[Out] 4/3*(12*a^5*log(e^(2*d*x + 2*c) + 1) + (48*a^5*e^(6*d*x + 6*c) + 108*a^5*e^(4*d*x + 4*c) + 88*a^5*e^(2*d*x + 2*c) + 25*a^5)/(e^(2*d*x + 2*c) + 1)^4)/d

Mupad [B]

time = 0.16, size = 65, normalized size = 0.65

$$32 a^5 x - \frac{a^5 (192 \ln(\tanh(c + dx) + 1) + 180 \tanh(c + dx) + 66 \tanh(c + dx)^2 + 20 \tanh(c + dx)^3 + 3 \tanh(c + dx)^4)}{12 d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + a*tanh(c + d*x))^5,x)

[Out] 32*a^5*x - (a^5*(192*log(tanh(c + d*x) + 1) + 180*tanh(c + d*x) + 66*tanh(c + d*x)^2 + 20*tanh(c + d*x)^3 + 3*tanh(c + d*x)^4))/(12*d)

3.42 $\int (a + a \tanh(c + dx))^4 dx$

Optimal. Leaf size=77

$$8a^4x + \frac{8a^4 \log(\cosh(c + dx))}{d} - \frac{4a^4 \tanh(c + dx)}{d} - \frac{a(a + a \tanh(c + dx))^3}{3d} - \frac{(a^2 + a^2 \tanh(c + dx))^2}{d}$$

[Out] $8*a^4*x + 8*a^4*\ln(\cosh(d*x+c))/d - 4*a^4*\tanh(d*x+c)/d - 1/3*a*(a+a*\tanh(d*x+c))^3/d - (a^2+a^2*\tanh(d*x+c))^2/d$

Rubi [A]

time = 0.04, antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3559, 3558, 3556}

$$-\frac{4a^4 \tanh(c + dx)}{d} + \frac{8a^4 \log(\cosh(c + dx))}{d} + 8a^4x - \frac{(a^2 \tanh(c + dx) + a^2)^2}{d} - \frac{a(a \tanh(c + dx) + a)^3}{3d}$$

Antiderivative was successfully verified.

[In] `Int[(a + a*Tanh[c + d*x])^4, x]`

[Out] $8*a^4*x + (8*a^4*\text{Log}[\text{Cosh}[c + d*x]])/d - (4*a^4*\text{Tanh}[c + d*x])/d - (a*(a + a*\text{Tanh}[c + d*x])^3)/(3*d) - (a^2 + a^2*\text{Tanh}[c + d*x])^2/d$

Rule 3556

`Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

Rule 3558

`Int[((a_.) + (b_.)*tan[(c_.) + (d_.)*(x_)])^2, x_Symbol] := Simp[(a^2 - b^2)*x, x] + (Dist[2*a*b, Int[Tan[c + d*x], x], x] + Simp[b^2*(Tan[c + d*x]/d), x]) /; FreeQ[{a, b, c, d}, x]`

Rule 3559

`Int[((a_.) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((a + b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] + Dist[2*a, Int[(a + b*Tan[c + d*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && GtQ[n, 1]`

Rubi steps

$$\begin{aligned}
\int (a + a \tanh(c + dx))^4 dx &= -\frac{a(a + a \tanh(c + dx))^3}{3d} + (2a) \int (a + a \tanh(c + dx))^3 dx \\
&= -\frac{a(a + a \tanh(c + dx))^3}{3d} - \frac{(a^2 + a^2 \tanh(c + dx))^2}{d} + (4a^2) \int (a + a \tanh(c + dx))^2 dx \\
&= 8a^4 x - \frac{4a^4 \tanh(c + dx)}{d} - \frac{a(a + a \tanh(c + dx))^3}{3d} - \frac{(a^2 + a^2 \tanh(c + dx))^2}{d} \\
&= 8a^4 x + \frac{8a^4 \log(\cosh(c + dx))}{d} - \frac{4a^4 \tanh(c + dx)}{d} - \frac{a(a + a \tanh(c + dx))^3}{3d}
\end{aligned}$$

Mathematica [B] Leaf count is larger than twice the leaf count of optimal. 178 vs. $2(77) = 154$.

time = 0.57, size = 178, normalized size = 2.31

$$\frac{a^4 \operatorname{sech}(c) \operatorname{sech}^2(c + dx) (\cosh(4dx) + \sinh(4dx)) (6dx \cosh(2c + 3dx) + 6dx \cosh(4c + 3dx) + 6 \cosh(2c + 3dx) \log(\cosh(c + dx)) + 6 \cosh(4c + 3dx) \log(\cosh(c + dx)) + 6 \cosh(dx) (1 + 3dx + 3 \log(\cosh(c + dx))) + 6 \cosh(2c + dx) (1 + 3dx + 3 \log(\cosh(c + dx))) - 21 \sinh(dx) + 12 \sinh(2c + dx) - 11 \sinh(2c + 3dx))}{6d(\cosh(dx) + \sinh(dx))^4}$$

Antiderivative was successfully verified.

[In] Integrate[(a + a*Tanh[c + d*x])^4, x]

[Out] (a^4*Sech[c]*Sech[c + d*x]^3*(Cosh[4*d*x] + Sinh[4*d*x])*(6*d*x*Cosh[2*c + 3*d*x] + 6*d*x*Cosh[4*c + 3*d*x] + 6*Cosh[2*c + 3*d*x]*Log[Cosh[c + d*x]] + 6*Cosh[4*c + 3*d*x]*Log[Cosh[c + d*x]] + 6*Cosh[d*x]*(1 + 3*d*x + 3*Log[Cosh[c + d*x]])) + 6*Cosh[2*c + d*x]*(1 + 3*d*x + 3*Log[Cosh[c + d*x]]) - 21*Sinh[d*x] + 12*Sinh[2*c + d*x] - 11*Sinh[2*c + 3*d*x]))/(6*d*(Cosh[d*x] + Sinh[d*x])^4)

Maple [A]

time = 0.26, size = 48, normalized size = 0.62

method	result	size
derivativedivides	$a^4 \left(\frac{-\frac{\tanh^3(dx+c)}{3} - 2(\tanh^2(dx+c)) - 7 \tanh(dx+c) - 8 \ln(\tanh(dx+c)-1)}{d} \right)$	48
default	$a^4 \left(\frac{-\frac{\tanh^3(dx+c)}{3} - 2(\tanh^2(dx+c)) - 7 \tanh(dx+c) - 8 \ln(\tanh(dx+c)-1)}{d} \right)$	48
risch	$-\frac{16a^4c}{d} + \frac{4a^4(18e^{4dx+4c} + 27e^{2dx+2c} + 11)}{3d(1+e^{2dx+2c})^3} + \frac{8a^4 \ln(1+e^{2dx+2c})}{d}$	76

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a+a*tanh(d*x+c))^4,x,method=_RETURNVERBOSE)

[Out] 1/d*a^4*(-1/3*tanh(d*x+c)^3-2*tanh(d*x+c)^2-7*tanh(d*x+c)-8*ln(tanh(d*x+c)-1))

Maxima [B] Leaf count of result is larger than twice the leaf count of optimal. 196 vs. $2(75) = 150$.
time = 0.48, size = 196, normalized size = 2.55

$$\frac{1}{3}a^4 \left(3x + \frac{3c}{d} - \frac{4(3e^{(-2dx-2c)} + 3e^{(-4dx-4c)} + 2)}{d(3e^{(-2dx-2c)} + 3e^{(-4dx-4c)} + e^{(-6dx-6c)} + 1)} \right) + 4a^4 \left(x + \frac{c}{d} + \frac{\log(e^{(-2dx-2c)} + 1)}{d} + \frac{2e^{(-2dx-2c)}}{d(2e^{(-2dx-2c)} + e^{(-4dx-4c)} + 1)} \right) + 6a^4 \left(x + \frac{c}{d} - \frac{2}{d(e^{(-2dx-2c)} + 1)} \right) + a^4x + \frac{4a^4 \log(\cosh(dx + c))}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+a*tanh(d*x+c))^4,x, algorithm="maxima")

[Out] $\frac{1}{3}a^4(3x + 3c/d - 4(3e^{(-2dx-2c)} + 3e^{(-4dx-4c)} + 2)/(d(3e^{(-2dx-2c)} + 3e^{(-4dx-4c)} + e^{(-6dx-6c)} + 1))) + 4a^4(x + c/d + \log(e^{(-2dx-2c)} + 1)/d + 2e^{(-2dx-2c)}/(d(2e^{(-2dx-2c)} + e^{(-4dx-4c)} + 1))) + 6a^4(x + c/d - 2/(d(e^{(-2dx-2c)} + 1))) + a^4x + 4a^4 \log(\cosh(dx + c))/d$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 562 vs. $2(75) = 150$.
time = 0.39, size = 562, normalized size = 7.30

[[[The number of nodes in the result of the antiderivative is larger than twice the number of nodes in the original expression. This is a warning that the result is not optimal.]]]]

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+a*tanh(d*x+c))^4,x, algorithm="fricas")

[Out] $\frac{4}{3}(18a^4 \cosh(dx + c)^4 + 72a^4 \cosh(dx + c) \sinh(dx + c)^3 + 18a^4 \sinh(dx + c)^4 + 27a^4 \cosh(dx + c)^2 + 11a^4 + 27(4a^4 \cosh(dx + c)^2 + a^4) \sinh(dx + c)^2 + 6(a^4 \cosh(dx + c)^6 + 6a^4 \cosh(dx + c) \sinh(dx + c)^5 + a^4 \sinh(dx + c)^6 + 3a^4 \cosh(dx + c)^4 + 3a^4 \cosh(dx + c)^2 + 3(5a^4 \cosh(dx + c)^2 + a^4) \sinh(dx + c)^4 + a^4 + 4(5a^4 \cosh(dx + c)^3 + 3a^4 \cosh(dx + c) \sinh(dx + c)^3 + 3(5a^4 \cosh(dx + c)^4 + 6a^4 \cosh(dx + c)^2 + a^4) \sinh(dx + c)^2 + 6(a^4 \cosh(dx + c)^5 + 2a^4 \cosh(dx + c)^3 + a^4 \cosh(dx + c) \sinh(dx + c)) \log(2 \cosh(dx + c) / (\cosh(dx + c) - \sinh(dx + c))) + 18(4a^4 \cosh(dx + c)^3 + 3a^4 \cosh(dx + c) \sinh(dx + c)) / (d \cosh(dx + c)^6 + 6d \cosh(dx + c) \sinh(dx + c)^5 + d \sinh(dx + c)^6 + 3d \cosh(dx + c)^4 + 3(5d \cosh(dx + c)^2 + d) \sinh(dx + c)^4 + 4(5d \cosh(dx + c)^3 + 3d \cosh(dx + c) \sinh(dx + c)^3 + 3d \cosh(dx + c)^2 + 3(5d \cosh(dx + c)^4 + 6d \cosh(dx + c)^2 + d) \sinh(dx + c)^2 + 6(d \cosh(dx + c)^5 + 2d \cosh(dx + c)^3 + d \cosh(dx + c)) \sinh(dx + c) + d)$

Sympy [A]

time = 0.11, size = 76, normalized size = 0.99

$$\begin{cases} 16a^4x - \frac{8a^4 \log(\tanh(c+dx)+1)}{d} - \frac{a^4 \tanh^3(c+dx)}{3d} - \frac{2a^4 \tanh^2(c+dx)}{d} - \frac{7a^4 \tanh(c+dx)}{d} & \text{for } d \neq 0 \\ x(a \tanh(c) + a)^4 & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+a*tanh(d*x+c))**4,x)

[Out] Piecewise((16*a**4*x - 8*a**4*log(tanh(c + d*x) + 1)/d - a**4*tanh(c + d*x)**3/(3*d) - 2*a**4*tanh(c + d*x)**2/d - 7*a**4*tanh(c + d*x)/d, Ne(d, 0)), (x*(a*tanh(c) + a)**4, True))

Giac [A]

time = 0.42, size = 71, normalized size = 0.92

$$\frac{4 \left(6 a^4 \log \left(e^{(2 dx+2 c)} + 1 \right) + \frac{18 a^4 e^{(4 dx+4 c)} + 27 a^4 e^{(2 dx+2 c)} + 11 a^4}{\left(e^{(2 dx+2 c)} + 1 \right)^3} \right)}{3 d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+a*tanh(d*x+c))^4,x, algorithm="giac")

[Out] 4/3*(6*a^4*log(e^(2*d*x + 2*c) + 1) + (18*a^4*e^(4*d*x + 4*c) + 27*a^4*e^(2*d*x + 2*c) + 11*a^4)/(e^(2*d*x + 2*c) + 1)^3)/d

Mupad [B]

time = 0.11, size = 53, normalized size = 0.69

$$16 a^4 x - \frac{a^4 \left(24 \ln \left(\tanh(c + dx) + 1 \right) + 21 \tanh(c + dx) + 6 \tanh(c + dx)^2 + \tanh(c + dx)^3 \right)}{3 d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + a*tanh(c + d*x))^4,x)

[Out] 16*a^4*x - (a^4*(24*log(tanh(c + d*x) + 1) + 21*tanh(c + d*x) + 6*tanh(c + d*x)^2 + tanh(c + d*x)^3))/(3*d)

3.43 $\int (a + a \tanh(c + dx))^3 dx$

Optimal. Leaf size=56

$$4a^3x + \frac{4a^3 \log(\cosh(c + dx))}{d} - \frac{2a^3 \tanh(c + dx)}{d} - \frac{a(a + a \tanh(c + dx))^2}{2d}$$

[Out] $4*a^3*x + 4*a^3*\ln(\cosh(d*x+c))/d - 2*a^3*\tanh(d*x+c)/d - 1/2*a*(a+a*\tanh(d*x+c))^2/d$

Rubi [A]

time = 0.03, antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3559, 3558, 3556}

$$-\frac{2a^3 \tanh(c + dx)}{d} + \frac{4a^3 \log(\cosh(c + dx))}{d} + 4a^3x - \frac{a(a \tanh(c + dx) + a)^2}{2d}$$

Antiderivative was successfully verified.

[In] Int[(a + a*Tanh[c + d*x])^3, x]

[Out] $4*a^3*x + (4*a^3*\text{Log}[\text{Cosh}[c + d*x]])/d - (2*a^3*\text{Tanh}[c + d*x])/d - (a*(a + a*\text{Tanh}[c + d*x])^2)/(2*d)$

Rule 3556

Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]

Rule 3558

Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^2, x_Symbol] := Simp[(a^2 - b^2)*x, x] + (Dist[2*a*b, Int[Tan[c + d*x], x], x] + Simp[b^2*(Tan[c + d*x]/d), x]) /; FreeQ[{a, b, c, d}, x]

Rule 3559

Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((a + b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] + Dist[2*a, Int[(a + b*Tan[c + d*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && GtQ[n, 1]

Rubi steps

$$\begin{aligned}
\int (a + a \tanh(c + dx))^3 dx &= -\frac{a(a + a \tanh(c + dx))^2}{2d} + (2a) \int (a + a \tanh(c + dx))^2 dx \\
&= 4a^3 x - \frac{2a^3 \tanh(c + dx)}{d} - \frac{a(a + a \tanh(c + dx))^2}{2d} + (4a^3) \int \tanh(c + dx) dx \\
&= 4a^3 x + \frac{4a^3 \log(\cosh(c + dx))}{d} - \frac{2a^3 \tanh(c + dx)}{d} - \frac{a(a + a \tanh(c + dx))^2}{2d}
\end{aligned}$$

Mathematica [A]

time = 0.49, size = 103, normalized size = 1.84

$$\frac{a^3 \operatorname{sech}(c) \operatorname{sech}^2(c + dx) (2dx \cosh(3c + 2dx) + 2 \cosh(3c + 2dx) \log(\cosh(c + dx)) + 2 \cosh(c + 2dx) (dx + \log(\cosh(c + dx))) + \cosh(c) (1 + 4dx + 4 \log(\cosh(c + dx))) + 3 \sinh(c) - 3 \sinh(c + 2dx))}{2d}$$

Antiderivative was successfully verified.

`[In] Integrate[(a + a*Tanh[c + d*x])^3, x]`

```
[Out] (a^3*Sech[c]*Sech[c + d*x]^2*(2*d*x*Cosh[3*c + 2*d*x] + 2*Cosh[3*c + 2*d*x]
*Log[Cosh[c + d*x]] + 2*Cosh[c + 2*d*x]*(d*x + Log[Cosh[c + d*x]]) + Cosh[c
]*(1 + 4*d*x + 4*Log[Cosh[c + d*x])) + 3*Sinh[c] - 3*Sinh[c + 2*d*x]))/(2*d
)
```

Maple [A]

time = 0.26, size = 38, normalized size = 0.68

method	result	size
derivativeldivides	$\frac{a^3 \left(-\frac{\tanh^2(dx+c)}{2} - 3 \tanh(dx+c) - 4 \ln(\tanh(dx+c)-1) \right)}{d}$	38
default	$\frac{a^3 \left(-\frac{\tanh^2(dx+c)}{2} - 3 \tanh(dx+c) - 4 \ln(\tanh(dx+c)-1) \right)}{d}$	38
risch	$-\frac{8a^3 c}{d} + \frac{2a^3 (4e^{2dx+2c}+3)}{d(1+e^{2dx+2c})^2} + \frac{4a^3 \ln(1+e^{2dx+2c})}{d}$	65

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((a+a*tanh(d*x+c))^3,x,method=_RETURNVERBOSE)`

```
[Out] 1/d*a^3*(-1/2*tanh(d*x+c)^2-3*tanh(d*x+c)-4*ln(tanh(d*x+c)-1))
```

Maxima [B] Leaf count of result is larger than twice the leaf count of optimal. 116 vs. 2(54) = 108.

time = 0.47, size = 116, normalized size = 2.07

$$a^3 \left(x + \frac{c}{d} + \frac{\log(e^{(-2dx-2c)} + 1)}{d} + \frac{2e^{(-2dx-2c)}}{d(2e^{(-2dx-2c)} + e^{(-4dx-4c)} + 1)} \right) + 3a^3 \left(x + \frac{c}{d} - \frac{2}{d(e^{(-2dx-2c)} + 1)} \right) + a^3 x + \frac{3a^3 \log(\cosh(dx+c))}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+a*tanh(d*x+c))^3,x, algorithm="maxima")

[Out] $a^3*(x + c/d + \log(e^{(-2*d*x - 2*c)} + 1)/d + 2*e^{(-2*d*x - 2*c)}/(d*(2*e^{(-2*d*x - 2*c)} + e^{(-4*d*x - 4*c)} + 1))) + 3*a^3*(x + c/d - 2/(d*(e^{(-2*d*x - 2*c)} + 1))) + a^3*x + 3*a^3*\log(\cosh(d*x + c))/d$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 299 vs. $2(54) = 108$.

time = 0.35, size = 299, normalized size = 5.34

$$\frac{2(4a^3 \cosh(dx+c)^2 + 8a^3 \cosh(dx+c) \sinh(dx+c) + 4a^3 \sinh(dx+c)^2 + 3a^3 + 2(e^{(-2dx-2c)} + e^{(-4dx-4c)} + 1)/d + 2e^{(-2dx-2c)}/(d(2e^{(-2dx-2c)} + e^{(-4dx-4c)} + 1))) + 3a^3(x + c/d - 2/(d(e^{(-2dx-2c)} + 1))) + a^3x + 3a^3 \log(\cosh(dx+c))}{d \cosh(dx+c)^4 + 4d \cosh(dx+c) \sinh(dx+c) + d \sinh(dx+c)^2 + 2d \cosh(dx+c)^2 + 2(3d \cosh(dx+c)^2 + d) \sinh(dx+c)^2 + 4(d \cosh(dx+c) + d \sinh(dx+c)) \sinh(dx+c) + d \log\left(\frac{2 \cosh(dx+c)}{\cosh(dx+c) - \sinh(dx+c)}\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+a*tanh(d*x+c))^3,x, algorithm="fricas")

[Out] $2*(4*a^3*\cosh(d*x + c)^2 + 8*a^3*\cosh(d*x + c)*\sinh(d*x + c) + 4*a^3*\sinh(d*x + c)^2 + 3*a^3 + 2*(a^3*\cosh(d*x + c)^4 + 4*a^3*\cosh(d*x + c)*\sinh(d*x + c)^3 + a^3*\sinh(d*x + c)^4 + 2*a^3*\cosh(d*x + c)^2 + a^3 + 2*(3*a^3*\cosh(d*x + c)^2 + a^3)*\sinh(d*x + c)^2 + 4*(a^3*\cosh(d*x + c)^3 + a^3*\cosh(d*x + c))*\sinh(d*x + c))*\log(2*\cosh(d*x + c)/(\cosh(d*x + c) - \sinh(d*x + c)))/(d*\cosh(d*x + c)^4 + 4*d*\cosh(d*x + c)*\sinh(d*x + c)^3 + d*\sinh(d*x + c)^4 + 2*d*\cosh(d*x + c)^2 + 2*(3*d*\cosh(d*x + c)^2 + d)*\sinh(d*x + c)^2 + 4*(d*\cosh(d*x + c)^3 + d*\cosh(d*x + c))*\sinh(d*x + c) + d)$

Sympy [A]

time = 0.10, size = 61, normalized size = 1.09

$$\begin{cases} 8a^3x - \frac{4a^3 \log(\tanh(c+dx)+1)}{d} - \frac{a^3 \tanh^2(c+dx)}{2d} - \frac{3a^3 \tanh(c+dx)}{d} & \text{for } d \neq 0 \\ x(a \tanh(c) + a)^3 & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+a*tanh(d*x+c))**3,x)

[Out] Piecewise((8*a**3*x - 4*a**3*log(tanh(c + d*x) + 1)/d - a**3*tanh(c + d*x)*2/(2*d) - 3*a**3*tanh(c + d*x)/d, Ne(d, 0)), (x*(a*tanh(c) + a)**3, True))

Giac [A]

time = 0.41, size = 57, normalized size = 1.02

$$\frac{2 \left(2 a^3 \log \left(e^{(2 dx+2 c)} + 1 \right) + \frac{4 a^3 e^{(2 dx+2 c)} + 3 a^3}{\left(e^{(2 dx+2 c)} + 1 \right)^2} \right)}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+a*tanh(d*x+c))^3,x, algorithm="giac")

[Out] $2*(2*a^3*\log(e^{(2*d*x + 2*c)} + 1) + (4*a^3*e^{(2*d*x + 2*c)} + 3*a^3)/(e^{(2*d*x + 2*c)} + 1)^2)/d$

Mupad [B]

time = 0.09, size = 43, normalized size = 0.77

$$8a^3x - \frac{a^3(8\ln(\tanh(cx + d)) + 1) + 6\tanh(cx + d) + \tanh(cx + d)^2}{2d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + a*tanh(c + d*x))^3,x)

[Out] $8*a^3*x - (a^3*(8*\log(\tanh(c + d*x) + 1) + 6*\tanh(c + d*x) + \tanh(c + d*x)^2))/(2*d)$

3.44 $\int (a + a \tanh(c + dx))^2 dx$

Optimal. Leaf size=36

$$2a^2x + \frac{2a^2 \log(\cosh(c + dx))}{d} - \frac{a^2 \tanh(c + dx)}{d}$$

[Out] $2*a^2*x+2*a^2*\ln(\cosh(d*x+c))/d-a^2*\tanh(d*x+c)/d$

Rubi [A]

time = 0.02, antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3558, 3556}

$$-\frac{a^2 \tanh(c + dx)}{d} + \frac{2a^2 \log(\cosh(c + dx))}{d} + 2a^2x$$

Antiderivative was successfully verified.

[In] $\text{Int}[(a + a*\text{Tanh}[c + d*x])^2, x]$

[Out] $2*a^2*x + (2*a^2*\text{Log}[\text{Cosh}[c + d*x]])/d - (a^2*\text{Tanh}[c + d*x])/d$

Rule 3556

$\text{Int}[\tan[(c_.) + (d_.)*(x_)], x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] /; \text{FreeQ}[\{c, d\}, x]$

Rule 3558

$\text{Int}[((a_.) + (b_.)*\tan[(c_.) + (d_.)*(x_)])^2, x_Symbol] \rightarrow \text{Simp}[(a^2 - b^2)*x, x] + (\text{Dist}[2*a*b, \text{Int}[\text{Tan}[c + d*x], x], x] + \text{Simp}[b^2*(\text{Tan}[c + d*x]/d), x]) /; \text{FreeQ}[\{a, b, c, d\}, x]$

Rubi steps

$$\begin{aligned} \int (a + a \tanh(c + dx))^2 dx &= 2a^2x - \frac{a^2 \tanh(c + dx)}{d} + (2a^2) \int \tanh(c + dx) dx \\ &= 2a^2x + \frac{2a^2 \log(\cosh(c + dx))}{d} - \frac{a^2 \tanh(c + dx)}{d} \end{aligned}$$

Mathematica [A]

time = 0.24, size = 58, normalized size = 1.61

$\frac{a^2 \text{sech}(c) \text{sech}(c + dx) (\cosh(dx) (dx + \log(\cosh(c + dx))) + \cosh(2c + dx) (dx + \log(\cosh(c + dx))) - \sinh(dx))}{d}$

Antiderivative was successfully verified.

[In] Integrate[(a + a*Tanh[c + d*x])^2,x]

[Out] (a^2*Sech[c]*Sech[c + d*x]*(Cosh[d*x]*(d*x + Log[Cosh[c + d*x]]) + Cosh[2*c + d*x]*(d*x + Log[Cosh[c + d*x]]) - Sinh[d*x]))/d

Maple [A]

time = 0.26, size = 28, normalized size = 0.78

method	result	size
derivativedivides	$\frac{a^2(-\tanh(dx+c)-2\ln(\tanh(dx+c)-1))}{d}$	28
default	$\frac{a^2(-\tanh(dx+c)-2\ln(\tanh(dx+c)-1))}{d}$	28
risch	$-\frac{4a^2c}{d} + \frac{2a^2}{d(1+e^{2dx+2c})} + \frac{2a^2\ln(1+e^{2dx+2c})}{d}$	52

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a+a*tanh(d*x+c))^2,x,method=_RETURNVERBOSE)

[Out] 1/d*a^2*(-tanh(d*x+c)-2*ln(tanh(d*x+c)-1))

Maxima [A]

time = 0.26, size = 50, normalized size = 1.39

$$a^2 \left(x + \frac{c}{d} - \frac{2}{d(e^{(-2dx-2c)} + 1)} \right) + a^2x + \frac{2a^2 \log(\cosh(dx+c))}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+a*tanh(d*x+c))^2,x, algorithm="maxima")

[Out] a^2*(x + c/d - 2/(d*(e^(-2*d*x - 2*c) + 1))) + a^2*x + 2*a^2*log(cosh(d*x + c))/d

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 117 vs. 2(36) = 72.

time = 0.39, size = 117, normalized size = 3.25

$$\frac{2 \left(a^2 + (a^2 \cosh(dx+c))^2 + 2a^2 \cosh(dx+c) \sinh(dx+c) + a^2 \sinh(dx+c)^2 + a^2 \log \left(\frac{2 \cosh(dx+c)}{\cosh(dx+c) - \sinh(dx+c)} \right) \right)}{d \cosh(dx+c)^2 + 2d \cosh(dx+c) \sinh(dx+c) + d \sinh(dx+c)^2 + d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+a*tanh(d*x+c))^2,x, algorithm="fricas")

[Out] 2*(a^2 + (a^2*cosh(d*x + c))^2 + 2*a^2*cosh(d*x + c)*sinh(d*x + c) + a^2*sinh(d*x + c)^2 + a^2)*log(2*cosh(d*x + c)/(cosh(d*x + c) - sinh(d*x + c)))/d

$d \cdot \cosh(dx + c)^2 + 2 \cdot d \cdot \cosh(dx + c) \cdot \sinh(dx + c) + d \cdot \sinh(dx + c)^2 + d$
)

Sympy [A]

time = 0.08, size = 44, normalized size = 1.22

$$\begin{cases} 4a^2x - \frac{2a^2 \log(\tanh(c+dx)+1)}{d} - \frac{a^2 \tanh(c+dx)}{d} & \text{for } d \neq 0 \\ x(a \tanh(c) + a)^2 & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+a*tanh(d*x+c))**2,x)

[Out] Piecewise((4*a**2*x - 2*a**2*log(tanh(c + d*x) + 1)/d - a**2*tanh(c + d*x)/d, Ne(d, 0)), (x*(a*tanh(c) + a)**2, True))

Giac [A]

time = 0.43, size = 39, normalized size = 1.08

$$\frac{2 \left(a^2 \log(e^{(2dx+2c)} + 1) + \frac{a^2}{e^{(2dx+2c)} + 1} \right)}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+a*tanh(d*x+c))^2,x, algorithm="giac")

[Out] 2*(a^2*log(e^(2*d*x + 2*c) + 1) + a^2/(e^(2*d*x + 2*c) + 1))/d

Mupad [B]

time = 1.07, size = 33, normalized size = 0.92

$$4a^2x - \frac{a^2(2 \ln(\tanh(c + dx) + 1) + \tanh(c + dx))}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + a*tanh(c + d*x))^2,x)

[Out] 4*a^2*x - (a^2*(2*log(tanh(c + d*x) + 1) + tanh(c + d*x)))/d

$$3.45 \quad \int \frac{1}{a+a \tanh(c+dx)} dx$$

Optimal. Leaf size=28

$$\frac{x}{2a} - \frac{1}{2d(a + a \tanh(c + dx))}$$

[Out] 1/2*x/a-1/2/d/(a+a*tanh(d*x+c))

Rubi [A]

time = 0.01, antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3560, 8}

$$\frac{x}{2a} - \frac{1}{2d(a \tanh(c + dx) + a)}$$

Antiderivative was successfully verified.

[In] Int[(a + a*Tanh[c + d*x])^(-1),x]

[Out] x/(2*a) - 1/(2*d*(a + a*Tanh[c + d*x]))

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 3560

Int[((a_) + (b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n)), x] + Dist[1/(2*a), Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{a+a \tanh(c+dx)} dx &= -\frac{1}{2d(a + a \tanh(c + dx))} + \frac{\int 1 dx}{2a} \\ &= \frac{x}{2a} - \frac{1}{2d(a + a \tanh(c + dx))} \end{aligned}$$

Mathematica [A]

time = 0.08, size = 39, normalized size = 1.39

$$\frac{-1 + 2dx + (1 + 2dx) \tanh(c + dx)}{4ad(1 + \tanh(c + dx))}$$

Antiderivative was successfully verified.

[In] Integrate[(a + a*Tanh[c + d*x])^(-1),x]

[Out] (-1 + 2*d*x + (1 + 2*d*x)*Tanh[c + d*x])/(4*a*d*(1 + Tanh[c + d*x]))

Maple [A]

time = 0.52, size = 43, normalized size = 1.54

method	result	size
risch	$\frac{x}{2a} - \frac{e^{-2dx-2c}}{4ad}$	25
derivativedivides	$-\frac{1}{2(\tanh(dx+c)+1)} + \frac{\ln(\tanh(dx+c)+1) - \ln(\tanh(dx+c)-1)}{4da}$	43
default	$-\frac{1}{2(\tanh(dx+c)+1)} + \frac{\ln(\tanh(dx+c)+1) - \ln(\tanh(dx+c)-1)}{4da}$	43

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a+a*tanh(d*x+c)),x,method=_RETURNVERBOSE)

[Out] 1/d/a*(-1/2/(tanh(d*x+c)+1)+1/4*ln(tanh(d*x+c)+1)-1/4*ln(tanh(d*x+c)-1))

Maxima [A]

time = 0.26, size = 31, normalized size = 1.11

$$\frac{dx + c}{2ad} - \frac{e^{(-2dx-2c)}}{4ad}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+a*tanh(d*x+c)),x, algorithm="maxima")

[Out] 1/2*(d*x + c)/(a*d) - 1/4*e^(-2*d*x - 2*c)/(a*d)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 50 vs. 2(24) = 48.

time = 0.39, size = 50, normalized size = 1.79

$$\frac{(2dx - 1) \cosh(dx + c) + (2dx + 1) \sinh(dx + c)}{4(ad \cosh(dx + c) + ad \sinh(dx + c))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+a*tanh(d*x+c)),x, algorithm="fricas")

[Out] 1/4*((2*d*x - 1)*cosh(d*x + c) + (2*d*x + 1)*sinh(d*x + c))/(a*d*cosh(d*x + c) + a*d*sinh(d*x + c))

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 73 vs. $2(19) = 38$.

time = 0.31, size = 73, normalized size = 2.61

$$\begin{cases} \frac{dx \tanh(c+dx)}{2ad \tanh(c+dx)+2ad} + \frac{dx}{2ad \tanh(c+dx)+2ad} - \frac{1}{2ad \tanh(c+dx)+2ad} & \text{for } d \neq 0 \\ \frac{x}{a \tanh(c)+a} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(a+a*tanh(d*x+c)),x)`

[Out] `Piecewise((d*x*tanh(c + d*x)/(2*a*d*tanh(c + d*x) + 2*a*d) + d*x/(2*a*d*tanh(c + d*x) + 2*a*d) - 1/(2*a*d*tanh(c + d*x) + 2*a*d), Ne(d, 0)), (x/(a*tanh(c) + a), True))`

Giac [A]

time = 0.41, size = 30, normalized size = 1.07

$$\frac{\frac{2(dx+c)}{a} - \frac{e^{(-2dx-2c)}}{a}}{4d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(a+a*tanh(d*x+c)),x, algorithm="giac")`

[Out] `1/4*(2*(d*x + c)/a - e^(-2*d*x - 2*c)/a)/d`

Mupad [B]

time = 1.07, size = 25, normalized size = 0.89

$$\frac{x}{2a} - \frac{1}{2ad(\tanh(c+dx)+1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(a + a*tanh(c + d*x)),x)`

[Out] `x/(2*a) - 1/(2*a*d*(tanh(c + d*x) + 1))`

$$3.46 \quad \int \frac{1}{(a+a \tanh(c+dx))^2} dx$$

Optimal. Leaf size=51

$$\frac{x}{4a^2} - \frac{1}{4d(a+a \tanh(c+dx))^2} - \frac{1}{4d(a^2+a^2 \tanh(c+dx))}$$

[Out] 1/4*x/a^2-1/4/d/(a+a*tanh(d*x+c))^2-1/4/d/(a^2+a^2*tanh(d*x+c))

Rubi [A]

time = 0.02, antiderivative size = 51, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3560, 8}

$$-\frac{1}{4d(a^2 \tanh(c+dx)+a^2)} + \frac{x}{4a^2} - \frac{1}{4d(a \tanh(c+dx)+a)^2}$$

Antiderivative was successfully verified.

[In] Int[(a + a*Tanh[c + d*x])^(-2),x]

[Out] x/(4*a^2) - 1/(4*d*(a + a*Tanh[c + d*x])^2) - 1/(4*d*(a^2 + a^2*Tanh[c + d*x]))

Rule 8

Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]

Rule 3560

Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n)), x] + Dist[1/(2*a), Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{(a+a \tanh(c+dx))^2} dx &= -\frac{1}{4d(a+a \tanh(c+dx))^2} + \frac{\int \frac{1}{a+a \tanh(c+dx)} dx}{2a} \\ &= -\frac{1}{4d(a+a \tanh(c+dx))^2} - \frac{1}{4d(a^2+a^2 \tanh(c+dx))} + \frac{\int 1 dx}{4a^2} \\ &= \frac{x}{4a^2} - \frac{1}{4d(a+a \tanh(c+dx))^2} - \frac{1}{4d(a^2+a^2 \tanh(c+dx))} \end{aligned}$$

Mathematica [A]

time = 0.13, size = 60, normalized size = 1.18

$$\frac{\operatorname{sech}^2(c + dx)(-4 + (-1 + 4dx) \cosh(2(c + dx)) + (1 + 4dx) \sinh(2(c + dx)))}{16a^2d(1 + \tanh(c + dx))^2}$$

Antiderivative was successfully verified.

`[In] Integrate[(a + a*Tanh[c + d*x])^(-2), x]`

```
[Out] (Sech[c + d*x]^2*(-4 + (-1 + 4*d*x)*Cosh[2*(c + d*x)] + (1 + 4*d*x)*Sinh[2*(c + d*x)]))/(16*a^2*d*(1 + Tanh[c + d*x])^2)
```

Maple [A]

time = 0.50, size = 55, normalized size = 1.08

method	result	size
risch	$\frac{x}{4a^2} - \frac{e^{-2dx-2c}}{4a^2d} - \frac{e^{-4dx-4c}}{16a^2d}$	42
derivativedivides	$\frac{-\frac{1}{4(\tanh(dx+c)+1)^2} - \frac{1}{4(\tanh(dx+c)+1)} + \frac{\ln(\tanh(dx+c)+1)}{8} - \frac{\ln(\tanh(dx+c)-1)}{8}}{da^2}$	55
default	$\frac{-\frac{1}{4(\tanh(dx+c)+1)^2} - \frac{1}{4(\tanh(dx+c)+1)} + \frac{\ln(\tanh(dx+c)+1)}{8} - \frac{\ln(\tanh(dx+c)-1)}{8}}{da^2}$	55

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/(a+a*tanh(d*x+c))^2,x,method=_RETURNVERBOSE)`

```
[Out] 1/d/a^2*(-1/4/(tanh(d*x+c)+1)^2-1/4/(tanh(d*x+c)+1)+1/8*ln(tanh(d*x+c)+1)-1/8*ln(tanh(d*x+c)-1))
```

Maxima [A]

time = 0.27, size = 43, normalized size = 0.84

$$\frac{dx + c}{4a^2d} - \frac{4e^{(-2dx-2c)} + e^{(-4dx-4c)}}{16a^2d}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/(a+a*tanh(d*x+c))^2,x, algorithm="maxima")`

```
[Out] 1/4*(d*x + c)/(a^2*d) - 1/16*(4*e^(-2*d*x - 2*c) + e^(-4*d*x - 4*c))/(a^2*d)
```

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 101 vs. 2(45) = 90.

time = 0.36, size = 101, normalized size = 1.98

$$\frac{(4dx - 1) \cosh(dx + c)^2 + 2(4dx + 1) \cosh(dx + c) \sinh(dx + c) + (4dx - 1) \sinh(dx + c)^2 - 4}{16(a^2d \cosh(dx + c)^2 + 2a^2d \cosh(dx + c) \sinh(dx + c) + a^2d \sinh(dx + c)^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+a*tanh(d*x+c))^2,x, algorithm="fricas")

[Out] $1/16*((4*d*x - 1)*\cosh(d*x + c)^2 + 2*(4*d*x + 1)*\cosh(d*x + c)*\sinh(d*x + c) + (4*d*x - 1)*\sinh(d*x + c)^2 - 4)/(a^2*d*\cosh(d*x + c)^2 + 2*a^2*d*\cosh(d*x + c)*\sinh(d*x + c) + a^2*d*\sinh(d*x + c)^2)$

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 223 vs. $2(41) = 82$.

time = 0.54, size = 223, normalized size = 4.37

$$\begin{cases} \frac{\frac{dx \tanh^2(c+dx)}{4a^2d \tanh^2(c+dx)+8a^2d \tanh(c+dx)+4a^2d} + \frac{2dx \tanh(c+dx)}{4a^2d \tanh^2(c+dx)+8a^2d \tanh(c+dx)+4a^2d} + \frac{dx}{4a^2d \tanh^2(c+dx)+8a^2d \tanh(c+dx)+4a^2d} - \frac{\tanh(c+dx)}{4a^2d \tanh^2(c+dx)+8a^2d \tanh(c+dx)+4a^2d} - \frac{2}{4a^2d \tanh^2(c+dx)+8a^2d \tanh(c+dx)+4a^2d}}{(a \tanh(c)+a)^2} & \text{for } d \neq 0 \\ \frac{x}{(a \tanh(c)+a)^2} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+a*tanh(d*x+c))^2,x)

[Out] Piecewise((d*x*tanh(c + d*x)**2/(4*a**2*d*tanh(c + d*x)**2 + 8*a**2*d*tanh(c + d*x) + 4*a**2*d) + 2*d*x*tanh(c + d*x)/(4*a**2*d*tanh(c + d*x)**2 + 8*a**2*d*tanh(c + d*x) + 4*a**2*d) + d*x/(4*a**2*d*tanh(c + d*x)**2 + 8*a**2*d*tanh(c + d*x) + 4*a**2*d) - tanh(c + d*x)/(4*a**2*d*tanh(c + d*x)**2 + 8*a**2*d*tanh(c + d*x) + 4*a**2*d) - 2/(4*a**2*d*tanh(c + d*x)**2 + 8*a**2*d*tanh(c + d*x) + 4*a**2*d), Ne(d, 0)), (x/(a*tanh(c) + a)**2, True))

Giac [A]

time = 0.42, size = 42, normalized size = 0.82

$$-\frac{(4e^{(2dx+2c)}+1)e^{(-4dx-4c)}}{a^2} - \frac{4(dx+c)}{a^2}$$

$16d$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+a*tanh(d*x+c))^2,x, algorithm="giac")

[Out] $-1/16*((4*e^{(2*d*x + 2*c)} + 1)*e^{(-4*d*x - 4*c)}/a^2 - 4*(d*x + c)/a^2)/d$

Mupad [B]

time = 1.09, size = 41, normalized size = 0.80

$$\frac{x}{4a^2} - \frac{e^{-2c-2dx}}{4a^2d} - \frac{e^{-4c-4dx}}{16a^2d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a + a*tanh(c + d*x))^2,x)

[Out] $x/(4*a^2) - \exp(-2*c - 2*d*x)/(4*a^2*d) - \exp(-4*c - 4*d*x)/(16*a^2*d)$

$$3.47 \quad \int \frac{1}{(a+a \tanh(c+dx))^3} dx$$

Optimal. Leaf size=73

$$\frac{x}{8a^3} - \frac{1}{6d(a+a \tanh(c+dx))^3} - \frac{1}{8ad(a+a \tanh(c+dx))^2} - \frac{1}{8d(a^3+a^3 \tanh(c+dx))}$$

[Out] 1/8*x/a^3-1/6/d/(a+a*tanh(d*x+c))^3-1/8/a/d/(a+a*tanh(d*x+c))^2-1/8/d/(a^3+a^3*tanh(d*x+c))

Rubi [A]

time = 0.03, antiderivative size = 73, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3560, 8}

$$-\frac{1}{8d(a^3 \tanh(c+dx)+a^3)} + \frac{x}{8a^3} - \frac{1}{8ad(a \tanh(c+dx)+a)^2} - \frac{1}{6d(a \tanh(c+dx)+a)^3}$$

Antiderivative was successfully verified.

[In] Int[(a + a*Tanh[c + d*x])^(-3), x]

[Out] x/(8*a^3) - 1/(6*d*(a + a*Tanh[c + d*x])^3) - 1/(8*a*d*(a + a*Tanh[c + d*x])^2) - 1/(8*d*(a^3 + a^3*Tanh[c + d*x]))

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 3560

Int[((a_) + (b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n)), x] + Dist[1/(2*a), Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{(a+a \tanh(c+dx))^3} dx &= -\frac{1}{6d(a+a \tanh(c+dx))^3} + \frac{\int \frac{1}{(a+a \tanh(c+dx))^2} dx}{2a} \\ &= -\frac{1}{6d(a+a \tanh(c+dx))^3} - \frac{1}{8ad(a+a \tanh(c+dx))^2} + \frac{\int \frac{1}{a+a \tanh(c+dx)} dx}{4a^2} \\ &= -\frac{1}{6d(a+a \tanh(c+dx))^3} - \frac{1}{8ad(a+a \tanh(c+dx))^2} - \frac{1}{8d(a^3+a^3 \tanh(c+dx))} \\ &= \frac{x}{8a^3} - \frac{1}{6d(a+a \tanh(c+dx))^3} - \frac{1}{8ad(a+a \tanh(c+dx))^2} - \frac{1}{8d(a^3+a^3 \tanh(c+dx))} \end{aligned}$$

Mathematica [A]

time = 0.18, size = 83, normalized size = 1.14

$$\frac{\operatorname{sech}^3(c+dx)(-27 \cosh(c+dx) + 2(-1+6dx) \cosh(3(c+dx)) - 9 \sinh(c+dx) + 2 \sinh(3(c+dx)) + 12dx \sinh(3(c+dx)))}{96a^3d(1+\tanh(c+dx))^3}$$

Antiderivative was successfully verified.

`[In] Integrate[(a + a*Tanh[c + d*x])^(-3), x]`

`[Out] (Sech[c + d*x]^3*(-27*Cosh[c + d*x] + 2*(-1 + 6*d*x)*Cosh[3*(c + d*x)] - 9*Sinh[c + d*x] + 2*Sinh[3*(c + d*x)] + 12*d*x*Sinh[3*(c + d*x)]))/(96*a^3*d*(1 + Tanh[c + d*x])^3)`

Maple [A]

time = 0.66, size = 67, normalized size = 0.92

method	result	size
risch	$\frac{x}{8a^3} - \frac{3e^{-2dx-2c}}{16a^3d} - \frac{3e^{-4dx-4c}}{32a^3d} - \frac{e^{-6dx-6c}}{48a^3d}$	59
derivativedivides	$\frac{-\frac{1}{6(\tanh(dx+c)+1)^3} - \frac{1}{8(\tanh(dx+c)+1)^2} - \frac{1}{8(\tanh(dx+c)+1)} + \frac{\ln(\tanh(dx+c)+1)}{16} - \frac{\ln(\tanh(dx+c)-1)}{16}}{da^3}$	67
default	$\frac{-\frac{1}{6(\tanh(dx+c)+1)^3} - \frac{1}{8(\tanh(dx+c)+1)^2} - \frac{1}{8(\tanh(dx+c)+1)} + \frac{\ln(\tanh(dx+c)+1)}{16} - \frac{\ln(\tanh(dx+c)-1)}{16}}{da^3}$	67

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/(a+a*tanh(d*x+c))^3,x,method=_RETURNVERBOSE)`

`[Out] 1/d/a^3*(-1/6/(tanh(d*x+c)+1)^3-1/8/(tanh(d*x+c)+1)^2-1/8/(tanh(d*x+c)+1)+1/16*ln(tanh(d*x+c)+1)-1/16*ln(tanh(d*x+c)-1))`

Maxima [A]

time = 0.27, size = 56, normalized size = 0.77

$$\frac{dx+c}{8a^3d} - \frac{18e^{(-2dx-2c)} + 9e^{(-4dx-4c)} + 2e^{(-6dx-6c)}}{96a^3d}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/(a+a*tanh(d*x+c))^3,x, algorithm="maxima")`

`[Out] 1/8*(d*x + c)/(a^3*d) - 1/96*(18*e^(-2*d*x - 2*c) + 9*e^(-4*d*x - 4*c) + 2*e^(-6*d*x - 6*c))/(a^3*d)`

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 160 vs. 2(65) = 130.

time = 0.44, size = 160, normalized size = 2.19

$$\frac{2(6dx-1)\cosh(dx+c)^3 + 6(6dx-1)\cosh(dx+c)\sinh(dx+c)^2 + 2(6dx+1)\sinh(dx+c)^3 + 3(2(6dx+1)\cosh(dx+c)^2 - 3)\sinh(dx+c) - 27\cosh(dx+c)}{96(a^3d\cosh(dx+c)^3 + 3a^3d\cosh(dx+c)^2\sinh(dx+c) + 3a^3d\cosh(dx+c)\sinh(dx+c)^2 + a^3d\sinh(dx+c)^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+a*tanh(d*x+c))^3,x, algorithm="fricas")

[Out] $\frac{1}{96} \cdot (2 \cdot (6 \cdot d \cdot x - 1) \cdot \cosh(d \cdot x + c)^3 + 6 \cdot (6 \cdot d \cdot x - 1) \cdot \cosh(d \cdot x + c) \cdot \sinh(d \cdot x + c)^2 + 2 \cdot (6 \cdot d \cdot x + 1) \cdot \sinh(d \cdot x + c)^3 + 3 \cdot (2 \cdot (6 \cdot d \cdot x + 1) \cdot \cosh(d \cdot x + c)^2 - 3) \cdot \sinh(d \cdot x + c) - 27 \cdot \cosh(d \cdot x + c)) / (a^3 \cdot d \cdot \cosh(d \cdot x + c)^3 + 3 \cdot a^3 \cdot d \cdot \cosh(d \cdot x + c)^2 \cdot \sinh(d \cdot x + c) + 3 \cdot a^3 \cdot d \cdot \cosh(d \cdot x + c) \cdot \sinh(d \cdot x + c)^2 + a^3 \cdot d \cdot \sinh(d \cdot x + c)^3)$

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 430 vs. 2(60) = 120.

time = 0.77, size = 430, normalized size = 5.89

fricas

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+a*tanh(d*x+c))**3,x)

[Out] Piecewise((3*d*x*tanh(c + d*x)**3/(24*a**3*d*tanh(c + d*x)**3 + 72*a**3*d*tanh(c + d*x)**2 + 72*a**3*d*tanh(c + d*x) + 24*a**3*d) + 9*d*x*tanh(c + d*x)**2/(24*a**3*d*tanh(c + d*x)**3 + 72*a**3*d*tanh(c + d*x)**2 + 72*a**3*d*tanh(c + d*x) + 24*a**3*d) + 9*d*x*tanh(c + d*x)/(24*a**3*d*tanh(c + d*x)**3 + 72*a**3*d*tanh(c + d*x)**2 + 72*a**3*d*tanh(c + d*x) + 24*a**3*d) + 3*d*x/(24*a**3*d*tanh(c + d*x)**3 + 72*a**3*d*tanh(c + d*x)**2 + 72*a**3*d*tanh(c + d*x) + 24*a**3*d) - 3*tanh(c + d*x)**2/(24*a**3*d*tanh(c + d*x)**3 + 72*a**3*d*tanh(c + d*x)**2 + 72*a**3*d*tanh(c + d*x) + 24*a**3*d) - 9*tanh(c + d*x)/(24*a**3*d*tanh(c + d*x)**3 + 72*a**3*d*tanh(c + d*x)**2 + 72*a**3*d*tanh(c + d*x) + 24*a**3*d) - 10/(24*a**3*d*tanh(c + d*x)**3 + 72*a**3*d*tanh(c + d*x)**2 + 72*a**3*d*tanh(c + d*x) + 24*a**3*d), Ne(d, 0)), (x/(a*tanh(c) + a)**3, True))

Giac [A]

time = 0.42, size = 53, normalized size = 0.73

$$-\frac{(18e^{(4dx+4c)}+9e^{(2dx+2c)}+2)e^{(-6dx-6c)} - \frac{12(dx+c)}{a^3}}{96d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+a*tanh(d*x+c))^3,x, algorithm="giac")

[Out] $-1/96 \cdot ((18 \cdot e^{(4 \cdot d \cdot x + 4 \cdot c)} + 9 \cdot e^{(2 \cdot d \cdot x + 2 \cdot c)} + 2) \cdot e^{(-6 \cdot d \cdot x - 6 \cdot c)} / a^3 - 12 \cdot (d \cdot x + c) / a^3) / d$

Mupad [B]

time = 1.10, size = 58, normalized size = 0.79

$$\frac{x}{8a^3} - \frac{3e^{-2c-2dx}}{16a^3d} - \frac{3e^{-4c-4dx}}{32a^3d} - \frac{e^{-6c-6dx}}{48a^3d}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(a + a*tanh(c + d*x))^3,x)
```

```
[Out] x/(8*a^3) - (3*exp(- 2*c - 2*d*x))/(16*a^3*d) - (3*exp(- 4*c - 4*d*x))/(32*  
a^3*d) - exp(- 6*c - 6*d*x)/(48*a^3*d)
```

$$3.48 \quad \int \frac{1}{(a+a \tanh(c+dx))^4} dx$$

Optimal. Leaf size=96

$$\frac{x}{16a^4} - \frac{1}{8d(a+a \tanh(c+dx))^4} - \frac{1}{12ad(a+a \tanh(c+dx))^3} - \frac{1}{16d(a^2+a^2 \tanh(c+dx))^2} - \frac{1}{16d(a^4+a^4 \tanh(c+dx))}$$

[Out] 1/16*x/a^4-1/8/d/(a+a*tanh(d*x+c))^4-1/12/a/d/(a+a*tanh(d*x+c))^3-1/16/d/(a^2+a^2*tanh(d*x+c))^2-1/16/d/(a^4+a^4*tanh(d*x+c))

Rubi [A]

time = 0.05, antiderivative size = 96, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3560, 8}

$$-\frac{1}{16d(a^4 \tanh(c+dx) + a^4)} + \frac{x}{16a^4} - \frac{1}{16d(a^2 \tanh(c+dx) + a^2)^2} - \frac{1}{12ad(a \tanh(c+dx) + a)^3} - \frac{1}{8d(a \tanh(c+dx) + a)^4}$$

Antiderivative was successfully verified.

[In] Int[(a + a*Tanh[c + d*x])^(-4),x]

[Out] x/(16*a^4) - 1/(8*d*(a + a*Tanh[c + d*x])^4) - 1/(12*a*d*(a + a*Tanh[c + d*x])^3) - 1/(16*d*(a^2 + a^2*Tanh[c + d*x])^2) - 1/(16*d*(a^4 + a^4*Tanh[c + d*x]))

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 3560

Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n)), x] + Dist[1/(2*a), Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]

Rubi steps

$$\begin{aligned}
\int \frac{1}{(a + a \tanh(c + dx))^4} dx &= -\frac{1}{8d(a + a \tanh(c + dx))^4} + \frac{\int \frac{1}{(a + a \tanh(c + dx))^3} dx}{2a} \\
&= -\frac{1}{8d(a + a \tanh(c + dx))^4} - \frac{1}{12ad(a + a \tanh(c + dx))^3} + \frac{\int \frac{1}{(a + a \tanh(c + dx))^2} dx}{4a^2} \\
&= -\frac{1}{8d(a + a \tanh(c + dx))^4} - \frac{1}{12ad(a + a \tanh(c + dx))^3} - \frac{1}{16d(a^2 + a^2 \tanh(c + dx))^2} \\
&= -\frac{1}{8d(a + a \tanh(c + dx))^4} - \frac{1}{12ad(a + a \tanh(c + dx))^3} - \frac{1}{16d(a^2 + a^2 \tanh(c + dx))^2} \\
&= \frac{x}{16a^4} - \frac{1}{8d(a + a \tanh(c + dx))^4} - \frac{1}{12ad(a + a \tanh(c + dx))^3} - \frac{1}{16d(a^2 + a^2 \tanh(c + dx))^2}
\end{aligned}$$

Mathematica [A]

time = 0.18, size = 88, normalized size = 0.92

$$\frac{\operatorname{sech}^4(c + dx)(-36 - 64 \cosh(2(c + dx)) + 3(-1 + 8dx) \cosh(4(c + dx)) - 32 \sinh(2(c + dx)) + 3 \sinh(4(c + dx)) + 24dx \sinh(4(c + dx)))}{384a^4d(1 + \tanh(c + dx))^4}$$

Antiderivative was successfully verified.

`[In] Integrate[(a + a*Tanh[c + d*x])^(-4), x]`

```
[Out] (Sech[c + d*x]^4*(-36 - 64*Cosh[2*(c + d*x)] + 3*(-1 + 8*d*x)*Cosh[4*(c + d*x)] - 32*Sinh[2*(c + d*x)] + 3*Sinh[4*(c + d*x)] + 24*d*x*Sinh[4*(c + d*x)]))/(384*a^4*d*(1 + Tanh[c + d*x])^4)
```

Maple [A]

time = 0.64, size = 79, normalized size = 0.82

method	result	S
risch	$\frac{x}{16a^4} - \frac{e^{-2dx-2c}}{8a^4d} - \frac{3e^{-4dx-4c}}{32a^4d} - \frac{e^{-6dx-6c}}{24a^4d} - \frac{e^{-8dx-8c}}{128a^4d}$	7
derivativedivides	$\frac{-\frac{1}{8(\tanh(dx+c)+1)^4} - \frac{1}{12(\tanh(dx+c)+1)^3} - \frac{1}{16(\tanh(dx+c)+1)^2} - \frac{1}{16(\tanh(dx+c)+1)} + \frac{\ln(\tanh(dx+c)+1)}{32} - \frac{\ln(\tanh(dx+c)-1)}{32}}{da^4}$	7
default	$\frac{-\frac{1}{8(\tanh(dx+c)+1)^4} - \frac{1}{12(\tanh(dx+c)+1)^3} - \frac{1}{16(\tanh(dx+c)+1)^2} - \frac{1}{16(\tanh(dx+c)+1)} + \frac{\ln(\tanh(dx+c)+1)}{32} - \frac{\ln(\tanh(dx+c)-1)}{32}}{da^4}$	7

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/(a+a*tanh(d*x+c))^4,x,method=_RETURNVERBOSE)`

```
[Out] 1/d/a^4*(-1/8/(tanh(d*x+c)+1)^4-1/12/(tanh(d*x+c)+1)^3-1/16/(tanh(d*x+c)+1)^2-1/16/(tanh(d*x+c)+1)+1/32*ln(tanh(d*x+c)+1)-1/32*ln(tanh(d*x+c)-1))
```

Maxima [A]

time = 0.29, size = 67, normalized size = 0.70

$$\frac{dx + c}{16 a^4 d} - \frac{48 e^{(-2 dx - 2 c)} + 36 e^{(-4 dx - 4 c)} + 16 e^{(-6 dx - 6 c)} + 3 e^{(-8 dx - 8 c)}}{384 a^4 d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+a*tanh(d*x+c))^4,x, algorithm="maxima")

[Out] 1/16*(d*x + c)/(a^4*d) - 1/384*(48*e^(-2*d*x - 2*c) + 36*e^(-4*d*x - 4*c) + 16*e^(-6*d*x - 6*c) + 3*e^(-8*d*x - 8*c))/(a^4*d)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 220 vs. 2(86) = 172.

time = 0.37, size = 220, normalized size = 2.29

$$\frac{3(8 dx - 1) \cosh(dx + c)^4 + 12(8 dx + 1) \cosh(dx + c) \sinh(dx + c)^3 + 3(8 dx - 1) \sinh(dx + c)^4 + 2(9(8 dx - 1) \cosh(dx + c)^2 - 32) \sinh(dx + c)^2 - 64 \cosh(dx + c)^2 + 4(3(8 dx + 1) \cosh(dx + c)^3 - 16 \cosh(dx + c) \sinh(dx + c) - 36)}{384(a^4 d \cosh(dx + c)^4 + 4 a^4 d \cosh(dx + c)^3 \sinh(dx + c) + 6 a^4 d \cosh(dx + c)^2 \sinh(dx + c)^2 + 4 a^4 d \cosh(dx + c) \sinh(dx + c)^3 + a^4 d \sinh(dx + c)^4)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+a*tanh(d*x+c))^4,x, algorithm="fricas")

[Out] 1/384*(3*(8*d*x - 1)*cosh(d*x + c)^4 + 12*(8*d*x + 1)*cosh(d*x + c)*sinh(d*x + c)^3 + 3*(8*d*x - 1)*sinh(d*x + c)^4 + 2*(9*(8*d*x - 1)*cosh(d*x + c)^2 - 32)*sinh(d*x + c)^2 - 64*cosh(d*x + c)^2 + 4*(3*(8*d*x + 1)*cosh(d*x + c)^3 - 16*cosh(d*x + c)*sinh(d*x + c) - 36)/(a^4*d*cosh(d*x + c)^4 + 4*a^4*d*cosh(d*x + c)^3*sinh(d*x + c) + 6*a^4*d*cosh(d*x + c)^2*sinh(d*x + c)^2 + 4*a^4*d*cosh(d*x + c)*sinh(d*x + c)^3 + a^4*d*sinh(d*x + c)^4)

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 694 vs. 2(80) = 160.

time = 1.08, size = 694, normalized size = 7.23

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+a*tanh(d*x+c))**4,x)

[Out] Piecewise((3*d*x*tanh(c + d*x)**4/(48*a**4*d*tanh(c + d*x)**4 + 192*a**4*d*tanh(c + d*x)**3 + 288*a**4*d*tanh(c + d*x)**2 + 192*a**4*d*tanh(c + d*x) + 48*a**4*d) + 12*d*x*tanh(c + d*x)**3/(48*a**4*d*tanh(c + d*x)**4 + 192*a**4*d*tanh(c + d*x)**3 + 288*a**4*d*tanh(c + d*x)**2 + 192*a**4*d*tanh(c + d*x) + 48*a**4*d) + 18*d*x*tanh(c + d*x)**2/(48*a**4*d*tanh(c + d*x)**4 + 192*a**4*d*tanh(c + d*x)**3 + 288*a**4*d*tanh(c + d*x)**2 + 192*a**4*d*tanh(c + d*x) + 48*a**4*d) + 12*d*x*tanh(c + d*x)/(48*a**4*d*tanh(c + d*x)**4 + 192*a**4*d*tanh(c + d*x)**3 + 288*a**4*d*tanh(c + d*x)**2 + 192*a**4*d*tanh(c + d*x) + 48*a**4*d) + 3*d*x/(48*a**4*d*tanh(c + d*x)**4 + 192*a**4*d*tanh(c + d*x)**3 + 288*a**4*d*tanh(c + d*x)**2 + 192*a**4*d*tanh(c + d*x) + 48*a**4*d)), (0, False))

$c + d*x)^{**3} + 288*a^{**4}*d*\tanh(c + d*x)^{**2} + 192*a^{**4}*d*\tanh(c + d*x) + 48*a^{**4}*d) - 3*\tanh(c + d*x)^{**3}/(48*a^{**4}*d*\tanh(c + d*x)^{**4} + 192*a^{**4}*d*\tanh(c + d*x)^{**3} + 288*a^{**4}*d*\tanh(c + d*x)^{**2} + 192*a^{**4}*d*\tanh(c + d*x) + 48*a^{**4}*d) - 12*\tanh(c + d*x)^{**2}/(48*a^{**4}*d*\tanh(c + d*x)^{**4} + 192*a^{**4}*d*\tanh(c + d*x)^{**3} + 288*a^{**4}*d*\tanh(c + d*x)^{**2} + 192*a^{**4}*d*\tanh(c + d*x) + 48*a^{**4}*d) - 19*\tanh(c + d*x)/(48*a^{**4}*d*\tanh(c + d*x)^{**4} + 192*a^{**4}*d*\tanh(c + d*x)^{**3} + 288*a^{**4}*d*\tanh(c + d*x)^{**2} + 192*a^{**4}*d*\tanh(c + d*x) + 48*a^{**4}*d) - 16/(48*a^{**4}*d*\tanh(c + d*x)^{**4} + 192*a^{**4}*d*\tanh(c + d*x)^{**3} + 288*a^{**4}*d*\tanh(c + d*x)^{**2} + 192*a^{**4}*d*\tanh(c + d*x) + 48*a^{**4}*d), Ne(d, 0)), (x/(a*tanh(c) + a)^{**4}, True))$

Giac [A]

time = 0.42, size = 64, normalized size = 0.67

$$-\frac{(48e^{(6dx+6c)}+36e^{(4dx+4c)}+16e^{(2dx+2c)}+3)e^{(-8dx-8c)}}{a^4} - \frac{24(dx+c)}{a^4}$$

384 d

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+a*tanh(d*x+c))^4,x, algorithm="giac")

[Out] -1/384*((48*e^(6*d*x + 6*c) + 36*e^(4*d*x + 4*c) + 16*e^(2*d*x + 2*c) + 3)*e^(-8*d*x - 8*c)/a^4 - 24*(d*x + c)/a^4)/d

Mupad [B]

time = 1.14, size = 75, normalized size = 0.78

$$\frac{x}{16a^4} - \frac{e^{-2c-2dx}}{8a^4d} - \frac{3e^{-4c-4dx}}{32a^4d} - \frac{e^{-6c-6dx}}{24a^4d} - \frac{e^{-8c-8dx}}{128a^4d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a + a*tanh(c + d*x))^4,x)

[Out] x/(16*a^4) - exp(- 2*c - 2*d*x)/(8*a^4*d) - (3*exp(- 4*c - 4*d*x))/(32*a^4*d) - exp(- 6*c - 6*d*x)/(24*a^4*d) - exp(- 8*c - 8*d*x)/(128*a^4*d)

$$3.49 \quad \int \frac{1}{(a+a \tanh(c+dx))^5} dx$$

Optimal. Leaf size=121

$$\frac{x}{32a^5} - \frac{1}{10d(a+a \tanh(c+dx))^5} - \frac{1}{16ad(a+a \tanh(c+dx))^4} - \frac{1}{24a^2d(a+a \tanh(c+dx))^3} - \frac{1}{32ad(a^2+a^2 \tanh(c+dx))^2}$$

[Out] 1/32*x/a^5-1/10/d/(a+a*tanh(d*x+c))^5-1/16/a/d/(a+a*tanh(d*x+c))^4-1/24/a^2/d/(a+a*tanh(d*x+c))^3-1/32/a/d/(a^2+a^2*tanh(d*x+c))^2-1/32/d/(a^5+a^5*tanh(d*x+c))

Rubi [A]

time = 0.06, antiderivative size = 121, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3560, 8}

$$-\frac{1}{32d(a^5 \tanh(c+dx) + a^5)} + \frac{x}{32a^5} - \frac{1}{32ad(a^2 \tanh(c+dx) + a^2)^2} - \frac{1}{24a^2d(a \tanh(c+dx) + a)^3} - \frac{1}{16ad(a \tanh(c+dx) + a)^4} - \frac{1}{10d(a \tanh(c+dx) + a)^5}$$

Antiderivative was successfully verified.

[In] Int[(a + a*Tanh[c + d*x])^(-5),x]

[Out] x/(32*a^5) - 1/(10*d*(a + a*Tanh[c + d*x])^5) - 1/(16*a*d*(a + a*Tanh[c + d*x])^4) - 1/(24*a^2*d*(a + a*Tanh[c + d*x])^3) - 1/(32*a*d*(a^2 + a^2*Tanh[c + d*x])^2) - 1/(32*d*(a^5 + a^5*Tanh[c + d*x]))

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 3560

Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n), x] + Dist[1/(2*a), Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]

Rubi steps

$$\begin{aligned}
\int \frac{1}{(a + a \tanh(c + dx))^5} dx &= -\frac{1}{10d(a + a \tanh(c + dx))^5} + \frac{\int \frac{1}{(a+a \tanh(c+dx))^4} dx}{2a} \\
&= -\frac{1}{10d(a + a \tanh(c + dx))^5} - \frac{1}{16ad(a + a \tanh(c + dx))^4} + \frac{\int \frac{1}{(a+a \tanh(c+dx))^3} dx}{4a^2} \\
&= -\frac{1}{10d(a + a \tanh(c + dx))^5} - \frac{1}{16ad(a + a \tanh(c + dx))^4} - \frac{1}{24a^2d(a + a \tanh(c + dx))^3} \\
&= -\frac{1}{10d(a + a \tanh(c + dx))^5} - \frac{1}{16ad(a + a \tanh(c + dx))^4} - \frac{1}{24a^2d(a + a \tanh(c + dx))^3} \\
&= -\frac{1}{10d(a + a \tanh(c + dx))^5} - \frac{1}{16ad(a + a \tanh(c + dx))^4} - \frac{1}{24a^2d(a + a \tanh(c + dx))^3} \\
&= \frac{x}{32a^5} - \frac{1}{10d(a + a \tanh(c + dx))^5} - \frac{1}{16ad(a + a \tanh(c + dx))^4} - \frac{1}{24a^2d(a + a \tanh(c + dx))^3}
\end{aligned}$$

Mathematica [A]

time = 0.21, size = 109, normalized size = 0.90

$$\frac{\operatorname{sech}^5(c + dx)(-500 \cosh(c + dx) - 375 \cosh(3(c + dx)) - 12 \cosh(5(c + dx)) + 120dx \cosh(5(c + dx)) - 100 \sinh(c + dx) - 225 \sinh(3(c + dx)) + 12 \sinh(5(c + dx)) + 120dx \sinh(5(c + dx)))}{3840a^5d(1 + \tanh(c + dx))^5}$$

Antiderivative was successfully verified.

`[In] Integrate[(a + a*Tanh[c + d*x])^(-5),x]`

```
[Out] (Sech[c + d*x]^5*(-500*Cosh[c + d*x] - 375*Cosh[3*(c + d*x)] - 12*Cosh[5*(c + d*x)] + 120*d*x*Cosh[5*(c + d*x)] - 100*Sinh[c + d*x] - 225*Sinh[3*(c + d*x)] + 12*Sinh[5*(c + d*x)] + 120*d*x*Sinh[5*(c + d*x)]))/(3840*a^5*d*(1 + Tanh[c + d*x])^5)
```

Maple [A]

time = 0.64, size = 91, normalized size = 0.75

method	result
derivativdivides	$\frac{-\frac{\ln(\tanh(dx+c)-1)}{64} - \frac{1}{10(\tanh(dx+c)+1)^5} - \frac{1}{16(\tanh(dx+c)+1)^4} - \frac{1}{24(\tanh(dx+c)+1)^3} - \frac{1}{32(\tanh(dx+c)+1)^2} - \frac{1}{32(\tanh(dx+c)+1)} + \frac{1}{da^5}}$
default	$\frac{-\frac{\ln(\tanh(dx+c)-1)}{64} - \frac{1}{10(\tanh(dx+c)+1)^5} - \frac{1}{16(\tanh(dx+c)+1)^4} - \frac{1}{24(\tanh(dx+c)+1)^3} - \frac{1}{32(\tanh(dx+c)+1)^2} - \frac{1}{32(\tanh(dx+c)+1)} + \frac{1}{da^5}}$
risch	$\frac{x}{32a^5} - \frac{5e^{-2dx-2c}}{64a^5d} - \frac{5e^{-4dx-4c}}{64a^5d} - \frac{5e^{-6dx-6c}}{96a^5d} - \frac{5e^{-8dx-8c}}{256a^5d} - \frac{e^{-10dx-10c}}{320a^5d}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/(a+a*tanh(d*x+c))^5,x,method=_RETURNVERBOSE)`

[Out] $1/d/a^5*(-1/64*\ln(\tanh(d*x+c)-1)-1/10/(\tanh(d*x+c)+1)^5-1/16/(\tanh(d*x+c)+1)^4-1/24/(\tanh(d*x+c)+1)^3-1/32/(\tanh(d*x+c)+1)^2-1/32/(\tanh(d*x+c)+1)+1/64*\ln(\tanh(d*x+c)+1))$

Maxima [A]

time = 0.29, size = 78, normalized size = 0.64

$$\frac{dx + c}{32 a^5 d} - \frac{300 e^{(-2 dx - 2 c)} + 300 e^{(-4 dx - 4 c)} + 200 e^{(-6 dx - 6 c)} + 75 e^{(-8 dx - 8 c)} + 12 e^{(-10 dx - 10 c)}}{3840 a^5 d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(a+a*tanh(d*x+c))^5,x, algorithm="maxima")`

[Out] $1/32*(d*x + c)/(a^5*d) - 1/3840*(300*e^{(-2*d*x - 2*c)} + 300*e^{(-4*d*x - 4*c)} + 200*e^{(-6*d*x - 6*c)} + 75*e^{(-8*d*x - 8*c)} + 12*e^{(-10*d*x - 10*c)})/(a^5*d)$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 287 vs. 2(109) = 218.

time = 0.40, size = 287, normalized size = 2.37

$$\frac{12(10 dx - 1) \cosh(dx + c)^4 + 60(10 dx - 1) \cosh(dx + c) \sinh(dx + c)^4 + 12(10 dx + 1) \sinh(dx + c)^5 + 15(8(10 dx + 1) \cosh(dx + c)^2 - 15) \sinh(dx + c)^3 - 375 \cosh(dx + c)^3 + 15(8(10 dx - 1) \cosh(dx + c)^2 - 75 \cosh(dx + c)) \sinh(dx + c)^2 + 5(12(10 dx + 1) \cosh(dx + c)^4 - 135 \cosh(dx + c)^2 - 20) \sinh(dx + c) - 500 \cosh(dx + c)}{3840 a^5 d \cosh(dx + c)^5 + 5 a^5 d \cosh(dx + c)^4 \sinh(dx + c) + 10 a^5 d \cosh(dx + c)^3 \sinh(dx + c)^2 + 10 a^5 d \cosh(dx + c)^2 \sinh(dx + c)^3 + 5 a^5 d \cosh(dx + c) \sinh(dx + c)^4 + a^5 d \sinh(dx + c)^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(a+a*tanh(d*x+c))^5,x, algorithm="fricas")`

[Out] $1/3840*(12*(10*d*x - 1)*\cosh(d*x + c)^5 + 60*(10*d*x - 1)*\cosh(d*x + c)*\sinh(d*x + c)^4 + 12*(10*d*x + 1)*\sinh(d*x + c)^5 + 15*(8*(10*d*x + 1)*\cosh(d*x + c)^2 - 15)*\sinh(d*x + c)^3 - 375*\cosh(d*x + c)^3 + 15*(8*(10*d*x - 1)*\cosh(d*x + c)^2 - 75*\cosh(d*x + c))*\sinh(d*x + c)^2 + 5*(12*(10*d*x + 1)*\cosh(d*x + c)^4 - 135*\cosh(d*x + c)^2 - 20)*\sinh(d*x + c) - 500*\cosh(d*x + c))/(a^5*d*\cosh(d*x + c)^5 + 5*a^5*d*\cosh(d*x + c)^4*\sinh(d*x + c) + 10*a^5*d*\cosh(d*x + c)^3*\sinh(d*x + c)^2 + 10*a^5*d*\cosh(d*x + c)^2*\sinh(d*x + c)^3 + 5*a^5*d*\cosh(d*x + c)*\sinh(d*x + c)^4 + a^5*d*\sinh(d*x + c)^5)$

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 1018 vs. 2(102) = 204.

time = 1.56, size = 1018, normalized size = 8.41

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(a+a*tanh(d*x+c))**5,x)`

[Out] $\text{Piecewise}((15*d*x*tanh(c + d*x))**5/(480*a**5*d*tanh(c + d*x)**5 + 2400*a**5*d*tanh(c + d*x)**4 + 4800*a**5*d*tanh(c + d*x)**3 + 4800*a**5*d*tanh(c + d$

```

*x)**2 + 2400*a**5*d*tanh(c + d*x) + 480*a**5*d) + 75*d*x*tanh(c + d*x)**4/
(480*a**5*d*tanh(c + d*x)**5 + 2400*a**5*d*tanh(c + d*x)**4 + 4800*a**5*d*t
anh(c + d*x)**3 + 4800*a**5*d*tanh(c + d*x)**2 + 2400*a**5*d*tanh(c + d*x)
+ 480*a**5*d) + 150*d*x*tanh(c + d*x)**3/(480*a**5*d*tanh(c + d*x)**5 + 240
0*a**5*d*tanh(c + d*x)**4 + 4800*a**5*d*tanh(c + d*x)**3 + 4800*a**5*d*tanh
(c + d*x)**2 + 2400*a**5*d*tanh(c + d*x) + 480*a**5*d) + 150*d*x*tanh(c + d
*x)**2/(480*a**5*d*tanh(c + d*x)**5 + 2400*a**5*d*tanh(c + d*x)**4 + 4800*a
**5*d*tanh(c + d*x)**3 + 4800*a**5*d*tanh(c + d*x)**2 + 2400*a**5*d*tanh(c
+ d*x) + 480*a**5*d) + 75*d*x*tanh(c + d*x)/(480*a**5*d*tanh(c + d*x)**5 +
2400*a**5*d*tanh(c + d*x)**4 + 4800*a**5*d*tanh(c + d*x)**3 + 4800*a**5*d*t
anh(c + d*x)**2 + 2400*a**5*d*tanh(c + d*x) + 480*a**5*d) + 15*d*x/(480*a**
5*d*tanh(c + d*x)**5 + 2400*a**5*d*tanh(c + d*x)**4 + 4800*a**5*d*tanh(c +
d*x)**3 + 4800*a**5*d*tanh(c + d*x)**2 + 2400*a**5*d*tanh(c + d*x) + 480*a**
5*d) - 15*tanh(c + d*x)**4/(480*a**5*d*tanh(c + d*x)**5 + 2400*a**5*d*tanh
(c + d*x)**4 + 4800*a**5*d*tanh(c + d*x)**3 + 4800*a**5*d*tanh(c + d*x)**2
+ 2400*a**5*d*tanh(c + d*x) + 480*a**5*d) - 75*tanh(c + d*x)**3/(480*a**5*d
*tanh(c + d*x)**5 + 2400*a**5*d*tanh(c + d*x)**4 + 4800*a**5*d*tanh(c + d*x
)**3 + 4800*a**5*d*tanh(c + d*x)**2 + 2400*a**5*d*tanh(c + d*x) + 480*a**5*
d) - 155*tanh(c + d*x)**2/(480*a**5*d*tanh(c + d*x)**5 + 2400*a**5*d*tanh(c
+ d*x)**4 + 4800*a**5*d*tanh(c + d*x)**3 + 4800*a**5*d*tanh(c + d*x)**2 +
2400*a**5*d*tanh(c + d*x) + 480*a**5*d) - 175*tanh(c + d*x)/(480*a**5*d*tan
h(c + d*x)**5 + 2400*a**5*d*tanh(c + d*x)**4 + 4800*a**5*d*tanh(c + d*x)**3
+ 4800*a**5*d*tanh(c + d*x)**2 + 2400*a**5*d*tanh(c + d*x) + 480*a**5*d) -
128/(480*a**5*d*tanh(c + d*x)**5 + 2400*a**5*d*tanh(c + d*x)**4 + 4800*a**
5*d*tanh(c + d*x)**3 + 4800*a**5*d*tanh(c + d*x)**2 + 2400*a**5*d*tanh(c +
d*x) + 480*a**5*d), Ne(d, 0)), (x/(a*tanh(c) + a)**5, True))

```

Giac [A]

time = 0.42, size = 75, normalized size = 0.62

$$\frac{(300e^{(8dx+8c)}+300e^{(6dx+6c)}+200e^{(4dx+4c)}+75e^{(2dx+2c)}+12)e^{(-10dx-10c)}}{a^5} - \frac{120(dx+c)}{a^5}$$

3840 d

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+a*tanh(d*x+c))^5,x, algorithm="giac")

[Out] -1/3840*((300*e^(8*d*x + 8*c) + 300*e^(6*d*x + 6*c) + 200*e^(4*d*x + 4*c) + 75*e^(2*d*x + 2*c) + 12)*e^(-10*d*x - 10*c)/a^5 - 120*(d*x + c)/a^5)/d

Mupad [B]

time = 1.16, size = 92, normalized size = 0.76

$$\frac{x}{32a^5} - \frac{5e^{-2c-2dx}}{64a^5d} - \frac{5e^{-4c-4dx}}{64a^5d} - \frac{5e^{-6c-6dx}}{96a^5d} - \frac{5e^{-8c-8dx}}{256a^5d} - \frac{e^{-10c-10dx}}{320a^5d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a + a*tanh(c + d*x))^5,x)

[Out] $x/(32*a^5) - (5*\exp(-2*c - 2*d*x))/(64*a^5*d) - (5*\exp(-4*c - 4*d*x))/(64*a^5*d) - (5*\exp(-6*c - 6*d*x))/(96*a^5*d) - (5*\exp(-8*c - 8*d*x))/(256*a^5*d) - \exp(-10*c - 10*d*x)/(320*a^5*d)$

3.50 $\int (1 + \tanh(x))^{7/2} dx$

Optimal. Leaf size=57

$$8\sqrt{2} \tanh^{-1} \left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}} \right) - 8\sqrt{1 + \tanh(x)} - \frac{4}{3}(1 + \tanh(x))^{3/2} - \frac{2}{5}(1 + \tanh(x))^{5/2}$$

[Out] 8*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))*2^(1/2)-8*(1+tanh(x))^(1/2)-4/3*(1+tanh(x))^(3/2)-2/5*(1+tanh(x))^(5/2)

Rubi [A]

time = 0.03, antiderivative size = 57, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {3559, 3561, 212}

$$8\sqrt{2} \tanh^{-1} \left(\frac{\sqrt{\tanh(x) + 1}}{\sqrt{2}} \right) - \frac{2}{5}(\tanh(x) + 1)^{5/2} - \frac{4}{3}(\tanh(x) + 1)^{3/2} - 8\sqrt{\tanh(x) + 1}$$

Antiderivative was successfully verified.

[In] Int[(1 + Tanh[x])^(7/2), x]

[Out] 8*Sqrt[2]*ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]] - 8*Sqrt[1 + Tanh[x]] - (4*(1 + Tanh[x])^(3/2))/3 - (2*(1 + Tanh[x])^(5/2))/5

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 3559

Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((a + b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] + Dist[2*a, Int[(a + b*Tan[c + d*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && GtQ[n, 1]

Rule 3561

Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Dist[-2*(b/d), Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]

Rubi steps

$$\begin{aligned}
\int (1 + \tanh(x))^{7/2} dx &= -\frac{2}{5}(1 + \tanh(x))^{5/2} + 2 \int (1 + \tanh(x))^{5/2} dx \\
&= -\frac{4}{3}(1 + \tanh(x))^{3/2} - \frac{2}{5}(1 + \tanh(x))^{5/2} + 4 \int (1 + \tanh(x))^{3/2} dx \\
&= -8\sqrt{1 + \tanh(x)} - \frac{4}{3}(1 + \tanh(x))^{3/2} - \frac{2}{5}(1 + \tanh(x))^{5/2} + 8 \int \sqrt{1 + \tanh(x)} dx \\
&= -8\sqrt{1 + \tanh(x)} - \frac{4}{3}(1 + \tanh(x))^{3/2} - \frac{2}{5}(1 + \tanh(x))^{5/2} + 16 \operatorname{Subst}\left(\int \frac{1}{2 - x^2} dx\right) \\
&= 8\sqrt{2} \tanh^{-1}\left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}}\right) - 8\sqrt{1 + \tanh(x)} - \frac{4}{3}(1 + \tanh(x))^{3/2} - \frac{2}{5}(1 + \tanh(x))^{5/2}
\end{aligned}$$

Mathematica [A]

time = 0.15, size = 65, normalized size = 1.14

$$\frac{\cosh^3(x) \left(8\sqrt{2} \tanh^{-1}\left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}}\right) (1 + \tanh(x))^3 - \frac{2}{15}(1 + \tanh(x))^{7/2} (76 - 3\operatorname{sech}^2(x) + 16 \tanh(x)) \right)}{(\cosh(x) + \sinh(x))^3}$$

Antiderivative was successfully verified.

`[In] Integrate[(1 + Tanh[x])^(7/2), x]`

```
[Out] (Cosh[x]^3*(8*Sqrt[2]*ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]]*(1 + Tanh[x])^3 -
(2*(1 + Tanh[x])^(7/2)*(76 - 3*Sech[x]^2 + 16*Tanh[x]))/15))/(Cosh[x] + Sin
h[x])^3
```

Maple [A]

time = 0.58, size = 43, normalized size = 0.75

method	result
derivativedivides	$8 \operatorname{arctanh}\left(\frac{\sqrt{1 + \tanh(x)} \sqrt{2}}{2}\right) \sqrt{2} - 8\sqrt{1 + \tanh(x)} - \frac{4(1 + \tanh(x))^{3/2}}{3} - \frac{2(1 + \tanh(x))^{5/2}}{5}$
default	$8 \operatorname{arctanh}\left(\frac{\sqrt{1 + \tanh(x)} \sqrt{2}}{2}\right) \sqrt{2} - 8\sqrt{1 + \tanh(x)} - \frac{4(1 + \tanh(x))^{3/2}}{3} - \frac{2(1 + \tanh(x))^{5/2}}{5}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((1+tanh(x))^(7/2), x, method=_RETURNVERBOSE)`

```
[Out] 8*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))*2^(1/2)-8*(1+tanh(x))^(1/2)-4/3*(1
+tanh(x))^(3/2)-2/5*(1+tanh(x))^(5/2)
```

Maxima [A]

time = 0.48, size = 83, normalized size = 1.46

$$-4\sqrt{2} \log\left(-\frac{\sqrt{2} - \frac{\sqrt{2}}{\sqrt{e^{(-2x)} + 1}}}{\sqrt{2} + \frac{\sqrt{2}}{\sqrt{e^{(-2x)} + 1}}}\right) - \frac{8\sqrt{2}}{\sqrt{e^{(-2x)} + 1}} - \frac{8\sqrt{2}}{3(e^{(-2x)} + 1)^{\frac{3}{2}}} - \frac{8\sqrt{2}}{5(e^{(-2x)} + 1)^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+tanh(x))^(7/2),x, algorithm="maxima")

[Out] $-4*\sqrt{2}*\log(-(\sqrt{2} - \sqrt{2}/\sqrt{e^{(-2*x)} + 1)})/(\sqrt{2} + \sqrt{2}/\sqrt{e^{(-2*x)} + 1})) - 8*\sqrt{2}/\sqrt{e^{(-2*x)} + 1} - 8/3*\sqrt{2}/(e^{(-2*x)} + 1)^{(3/2)} - 8/5*\sqrt{2}/(e^{(-2*x)} + 1)^{(5/2)}$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 434 vs. $2(42) = 84$.

time = 0.38, size = 434, normalized size = 7.61

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+tanh(x))^(7/2),x, algorithm="fricas")

[Out] $-4/15*(2*\sqrt{2}*(23*\sqrt{2}*\cosh(x)^5 + 115*\sqrt{2}*\cosh(x)*\sinh(x)^4 + 23*\sqrt{2}*\sinh(x)^5 + 5*(46*\sqrt{2}*\cosh(x)^2 + 7*\sqrt{2})*\sinh(x)^3 + 35*\sqrt{2}*\cosh(x)^3 + 5*(46*\sqrt{2}*\cosh(x)^3 + 21*\sqrt{2}*\cosh(x))*\sinh(x)^2 + 5*(23*\sqrt{2}*\cosh(x)^4 + 21*\sqrt{2}*\cosh(x)^2 + 3*\sqrt{2})*\sinh(x) + 15*\sqrt{2}*\cosh(x))*\sqrt{\cosh(x)/(\cosh(x) - \sinh(x))} - 15*(\sqrt{2}*\cosh(x)^6 + 6*\sqrt{2}*\cosh(x)*\sinh(x)^5 + \sqrt{2}*\sinh(x)^6 + 3*(5*\sqrt{2}*\cosh(x)^2 + \sqrt{2})*\sinh(x)^4 + 3*\sqrt{2}*\cosh(x)^4 + 4*(5*\sqrt{2}*\cosh(x)^3 + 3*\sqrt{2}*\cosh(x))*\sinh(x)^3 + 3*(5*\sqrt{2}*\cosh(x)^4 + 6*\sqrt{2}*\cosh(x)^2 + \sqrt{2})*\sinh(x)^2 + 3*\sqrt{2}*\cosh(x)^2 + 6*(\sqrt{2}*\cosh(x)^5 + 2*\sqrt{2}*\cosh(x)^3 + \sqrt{2}*\cosh(x))*\sinh(x) + \sqrt{2})*\log(-2*\sqrt{2}*\sqrt{\cosh(x)/(\cosh(x) - \sinh(x))}*(\cosh(x) + \sinh(x)) - 2*\cosh(x)^2 - 4*\cosh(x)*\sinh(x) - 2*\sinh(x)^2 - 1))/(\cosh(x)^6 + 6*\cosh(x)*\sinh(x)^5 + \sinh(x)^6 + 3*(5*\cosh(x)^2 + 1)*\sinh(x)^4 + 3*\cosh(x)^4 + 4*(5*\cosh(x)^3 + 3*\cosh(x))*\sinh(x)^3 + 3*(5*\cosh(x)^4 + 6*\cosh(x)^2 + 1)*\sinh(x)^2 + 3*\cosh(x)^2 + 6*(\cosh(x)^5 + 2*\cosh(x)^3 + \cosh(x))*\sinh(x) + 1)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (\tanh(x) + 1)^{\frac{7}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+tanh(x))**(7/2),x)

[Out] Integral((tanh(x) + 1)**(7/2), x)

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 140 vs. 2(42) = 84.

time = 0.42, size = 140, normalized size = 2.46

$$\frac{4}{15} \sqrt{2} \left(\frac{2 \left(45 \left(\sqrt{e^{4x} + e^{2x}} - e^{2x} \right)^4 - 135 \left(\sqrt{e^{4x} + e^{2x}} - e^{2x} \right)^3 + 170 \left(\sqrt{e^{4x} + e^{2x}} - e^{2x} \right)^2 - 100 \sqrt{e^{4x} + e^{2x}} + 100 e^{2x} + 23 \right)}{\left(\sqrt{e^{4x} + e^{2x}} - e^{2x} - 1 \right)^5} - 15 \log \left(-2 \sqrt{e^{4x} + e^{2x}} + 2 e^{2x} + 1 \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+tanh(x))^(7/2),x, algorithm="giac")

[Out] 4/15*sqrt(2)*(2*(45*(sqrt(e^(4*x) + e^(2*x)) - e^(2*x)))^4 - 135*(sqrt(e^(4*x) + e^(2*x)) - e^(2*x))^3 + 170*(sqrt(e^(4*x) + e^(2*x)) - e^(2*x))^2 - 100*sqrt(e^(4*x) + e^(2*x)) + 100*e^(2*x) + 23)/(sqrt(e^(4*x) + e^(2*x)) - e^(2*x) - 1)^5 - 15*log(-2*sqrt(e^(4*x) + e^(2*x)) + 2*e^(2*x) + 1))

Mupad [B]

time = 0.19, size = 44, normalized size = 0.77

$$-8 \sqrt{\tanh(x) + 1} - \frac{4(\tanh(x) + 1)^{3/2}}{3} - \frac{2(\tanh(x) + 1)^{5/2}}{5} - \sqrt{2} \operatorname{atan} \left(\frac{\sqrt{2} \sqrt{\tanh(x) + 1} \operatorname{li}}{2} \right) 8i$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((tanh(x) + 1)^(7/2),x)

[Out] - 2^(1/2)*atan((2^(1/2)*(tanh(x) + 1)^(1/2)*1i)/2)*8i - 8*(tanh(x) + 1)^(1/2) - (4*(tanh(x) + 1)^(3/2))/3 - (2*(tanh(x) + 1)^(5/2))/5

3.51 $\int (1 + \tanh(x))^{5/2} dx$

Optimal. Leaf size=45

$$4\sqrt{2} \tanh^{-1} \left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}} \right) - 4\sqrt{1 + \tanh(x)} - \frac{2}{3}(1 + \tanh(x))^{3/2}$$

[Out] 4*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))*2^(1/2)-4*(1+tanh(x))^(1/2)-2/3*(1+tanh(x))^(3/2)

Rubi [A]

time = 0.02, antiderivative size = 45, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {3559, 3561, 212}

$$4\sqrt{2} \tanh^{-1} \left(\frac{\sqrt{\tanh(x) + 1}}{\sqrt{2}} \right) - \frac{2}{3}(\tanh(x) + 1)^{3/2} - 4\sqrt{\tanh(x) + 1}$$

Antiderivative was successfully verified.

[In] Int[(1 + Tanh[x])^(5/2), x]

[Out] 4*Sqrt[2]*ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]] - 4*Sqrt[1 + Tanh[x]] - (2*(1 + Tanh[x])^(3/2))/3

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 3559

Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((a + b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] + Dist[2*a, Int[(a + b*Tan[c + d*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && GtQ[n, 1]

Rule 3561

Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Dist[-2*(b/d), Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]

Rubi steps

$$\begin{aligned}
\int (1 + \tanh(x))^{5/2} dx &= -\frac{2}{3}(1 + \tanh(x))^{3/2} + 2 \int (1 + \tanh(x))^{3/2} dx \\
&= -4\sqrt{1 + \tanh(x)} - \frac{2}{3}(1 + \tanh(x))^{3/2} + 4 \int \sqrt{1 + \tanh(x)} dx \\
&= -4\sqrt{1 + \tanh(x)} - \frac{2}{3}(1 + \tanh(x))^{3/2} + 8 \operatorname{Subst}\left(\int \frac{1}{2 - x^2} dx, x, \sqrt{1 + \tanh(x)}\right) \\
&= 4\sqrt{2} \tanh^{-1}\left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}}\right) - 4\sqrt{1 + \tanh(x)} - \frac{2}{3}(1 + \tanh(x))^{3/2}
\end{aligned}$$

Mathematica [A]

time = 0.12, size = 39, normalized size = 0.87

$$4\sqrt{2} \tanh^{-1}\left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}}\right) - \frac{2}{3}\sqrt{1 + \tanh(x)}(7 + \tanh(x))$$

Antiderivative was successfully verified.

`[In] Integrate[(1 + Tanh[x])^(5/2), x]``[Out] 4*Sqrt[2]*ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]] - (2*Sqrt[1 + Tanh[x]]*(7 + Tanh[x]))/3`**Maple [A]**

time = 0.55, size = 35, normalized size = 0.78

method	result	size
derivativedivides	$4 \operatorname{arctanh}\left(\frac{\sqrt{1 + \tanh(x)} \sqrt{2}}{2}\right) \sqrt{2} - 4\sqrt{1 + \tanh(x)} - \frac{2(1 + \tanh(x))^{3/2}}{3}$	35
default	$4 \operatorname{arctanh}\left(\frac{\sqrt{1 + \tanh(x)} \sqrt{2}}{2}\right) \sqrt{2} - 4\sqrt{1 + \tanh(x)} - \frac{2(1 + \tanh(x))^{3/2}}{3}$	35

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((1+tanh(x))^(5/2), x, method=_RETURNVERBOSE)``[Out] 4*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))*2^(1/2)-4*(1+tanh(x))^(1/2)-2/3*(1+tanh(x))^(3/2)`**Maxima [B]** Leaf count of result is larger than twice the leaf count of optimal. 70 vs. $2(34) = 68$.

time = 0.48, size = 70, normalized size = 1.56

$$-2\sqrt{2} \log\left(\frac{\sqrt{2} - \frac{\sqrt{2}}{\sqrt{e^{(-2x)} + 1}}}{\sqrt{2} + \frac{\sqrt{2}}{\sqrt{e^{(-2x)} + 1}}}\right) - \frac{4\sqrt{2}}{\sqrt{e^{(-2x)} + 1}} - \frac{4\sqrt{2}}{3(e^{(-2x)} + 1)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+tanh(x))^(5/2),x, algorithm="maxima")

[Out] -2*sqrt(2)*log(-sqrt(2) - sqrt(2)/sqrt(e^(-2*x) + 1))/(sqrt(2) + sqrt(2)/sqrt(e^(-2*x) + 1)) - 4*sqrt(2)/sqrt(e^(-2*x) + 1) - 4/3*sqrt(2)/(e^(-2*x) + 1)^(3/2)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 252 vs. 2(34) = 68.

time = 0.49, size = 252, normalized size = 5.60

$$\frac{2\left(2\sqrt{2}\left(4\sqrt{2}\cosh(x)^2 + 12\sqrt{2}\cosh(x)\sinh(x) + 4\sqrt{2}\sinh(x)^2 + 3\left(4\sqrt{2}\cosh(x)^2 + \sqrt{2}\sinh(x) + 3\sqrt{2}\cosh(x)\right)\frac{\cosh(x)}{\cosh(x) - \sinh(x)}\right) - 3\left(\sqrt{2}\cosh(x)^2 + 4\sqrt{2}\cosh(x)\sinh(x) + \sqrt{2}\sinh(x)^2 + 2\left(3\sqrt{2}\cosh(x)^2 + \sqrt{2}\sinh(x) + 2\sqrt{2}\cosh(x)\right) + 4\left(\sqrt{2}\cosh(x)^2 + \sqrt{2}\sinh(x) + \sqrt{2}\right)\frac{\cosh(x)}{\cosh(x) - \sinh(x)}\right)\log\left(\frac{-2\sqrt{2}\frac{\cosh(x)}{\cosh(x) - \sinh(x)}(\cosh(x) + \sinh(x)) - 2\cosh(x) - 4\cosh(x)\sinh(x) - 2\sinh(x) - 1}{3(\cosh(x)^2 + 4\cosh(x)\sinh(x) + \sinh(x)^2 + 2(3\cosh(x)^2 + \sqrt{2}\sinh(x) + 2\sqrt{2}\cosh(x)) + 4(\cosh(x)^2 + \sinh(x) + \sqrt{2}))}\right)}{3(\cosh(x)^2 + 4\cosh(x)\sinh(x) + \sinh(x)^2 + 2(3\cosh(x)^2 + \sqrt{2}\sinh(x) + 2\sqrt{2}\cosh(x)) + 4(\cosh(x)^2 + \sinh(x) + \sqrt{2}))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+tanh(x))^(5/2),x, algorithm="fricas")

[Out] -2/3*(2*sqrt(2)*(4*sqrt(2)*cosh(x)^3 + 12*sqrt(2)*cosh(x)*sinh(x)^2 + 4*sqrt(2)*sinh(x)^3 + 3*(4*sqrt(2)*cosh(x)^2 + sqrt(2))*sinh(x) + 3*sqrt(2)*cosh(x))*sqrt(cosh(x)/(cosh(x) - sinh(x))) - 3*(sqrt(2)*cosh(x)^4 + 4*sqrt(2)*cosh(x)*sinh(x)^3 + sqrt(2)*sinh(x)^4 + 2*(3*sqrt(2)*cosh(x)^2 + sqrt(2))*sinh(x)^2 + 2*sqrt(2)*cosh(x)^2 + 4*(sqrt(2)*cosh(x)^3 + sqrt(2)*cosh(x))*sinh(x) + sqrt(2))*log(-2*sqrt(2)*sqrt(cosh(x)/(cosh(x) - sinh(x)))*(cosh(x) + sinh(x)) - 2*cosh(x)^2 - 4*cosh(x)*sinh(x) - 2*sinh(x)^2 - 1)/(cosh(x)^4 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4 + 2*(3*cosh(x)^2 + 1)*sinh(x)^2 + 2*cosh(x)^2 + 4*(cosh(x)^3 + cosh(x))*sinh(x) + 1)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (\tanh(x) + 1)^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+tanh(x))**(5/2),x)

[Out] Integral((tanh(x) + 1)**(5/2), x)

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 96 vs. $2(34) = 68$.
time = 0.42, size = 96, normalized size = 2.13

$$\frac{2}{3} \sqrt{2} \left(\frac{2 \left(6 \left(\sqrt{e^{(4x)} + e^{(2x)}} - e^{(2x)} \right)^2 - 9 \sqrt{e^{(4x)} + e^{(2x)}} + 9 e^{(2x)} + 4 \right)}{\left(\sqrt{e^{(4x)} + e^{(2x)}} - e^{(2x)} - 1 \right)^3} - 3 \log \left(-2 \sqrt{e^{(4x)} + e^{(2x)}} + 2 e^{(2x)} + 1 \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+tanh(x))^(5/2),x, algorithm="giac")

[Out] $\frac{2}{3} \sqrt{2} * (2 * (6 * (\sqrt{e^{(4*x)} + e^{(2*x)}} - e^{(2*x)})^2 - 9 * \sqrt{e^{(4*x)} + e^{(2*x)}} + 9 * e^{(2*x)} + 4) / (\sqrt{e^{(4*x)} + e^{(2*x)}} - e^{(2*x)} - 1)^3 - 3 * \log(-2 * \sqrt{e^{(4*x)} + e^{(2*x)}} + 2 * e^{(2*x)} + 1))$

Mupad [B]

time = 0.11, size = 54, normalized size = 1.20

$$\sqrt{8} \ln \left(-2 \sqrt{8} \sqrt{\tanh(x) + 1} - 8 \right) - \frac{2 (\tanh(x) + 1)^{3/2}}{3} - 2 \sqrt{2} \ln \left(4 \sqrt{2} \sqrt{\tanh(x) + 1} - 8 \right) - 4 \sqrt{\tanh(x) + 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((tanh(x) + 1)^(5/2),x)

[Out] $8^{(1/2)} * \log(-2 * 8^{(1/2)} * (\tanh(x) + 1)^{(1/2)} - 8) - (2 * (\tanh(x) + 1)^{(3/2)}) / 3 - 2 * 2^{(1/2)} * \log(4 * 2^{(1/2)} * (\tanh(x) + 1)^{(1/2)} - 8) - 4 * (\tanh(x) + 1)^{(1/2)}$

3.52 $\int (1 + \tanh(x))^{3/2} dx$

Optimal. Leaf size=33

$$2\sqrt{2} \tanh^{-1} \left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}} \right) - 2\sqrt{1 + \tanh(x)}$$

[Out] 2*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))*2^(1/2)-2*(1+tanh(x))^(1/2)

Rubi [A]

time = 0.02, antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {3559, 3561, 212}

$$2\sqrt{2} \tanh^{-1} \left(\frac{\sqrt{\tanh(x) + 1}}{\sqrt{2}} \right) - 2\sqrt{\tanh(x) + 1}$$

Antiderivative was successfully verified.

[In] Int[(1 + Tanh[x])^(3/2), x]

[Out] 2*Sqrt[2]*ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]] - 2*Sqrt[1 + Tanh[x]]

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 3559

Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((a + b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] + Dist[2*a, Int[(a + b*Tan[c + d*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && GtQ[n, 1]

Rule 3561

Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Dist[-2*(b/d), Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]

Rubi steps

$$\begin{aligned}
\int (1 + \tanh(x))^{3/2} dx &= -2\sqrt{1 + \tanh(x)} + 2 \int \sqrt{1 + \tanh(x)} dx \\
&= -2\sqrt{1 + \tanh(x)} + 4 \operatorname{Subst} \left(\int \frac{1}{2 - x^2} dx, x, \sqrt{1 + \tanh(x)} \right) \\
&= 2\sqrt{2} \tanh^{-1} \left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}} \right) - 2\sqrt{1 + \tanh(x)}
\end{aligned}$$

Mathematica [A]

time = 0.05, size = 33, normalized size = 1.00

$$2\sqrt{2} \tanh^{-1} \left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}} \right) - 2\sqrt{1 + \tanh(x)}$$

Antiderivative was successfully verified.

`[In] Integrate[(1 + Tanh[x])^(3/2), x]``[Out] 2*Sqrt[2]*ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]] - 2*Sqrt[1 + Tanh[x]]`**Maple [A]**

time = 0.56, size = 27, normalized size = 0.82

method	result	size
derivativedivides	$2 \operatorname{arctanh} \left(\frac{\sqrt{1 + \tanh(x)} \sqrt{2}}{2} \right) \sqrt{2} - 2\sqrt{1 + \tanh(x)}$	27
default	$2 \operatorname{arctanh} \left(\frac{\sqrt{1 + \tanh(x)} \sqrt{2}}{2} \right) \sqrt{2} - 2\sqrt{1 + \tanh(x)}$	27

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((1+tanh(x))^(3/2), x, method=_RETURNVERBOSE)``[Out] 2*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))*2^(1/2)-2*(1+tanh(x))^(1/2)`**Maxima [B]** Leaf count of result is larger than twice the leaf count of optimal. 57 vs. $2(26) = 52.$

time = 0.47, size = 57, normalized size = 1.73

$$-\sqrt{2} \log \left(\frac{\sqrt{2} - \frac{\sqrt{2}}{\sqrt{e^{(-2x)} + 1}}}{\sqrt{2} + \frac{\sqrt{2}}{\sqrt{e^{(-2x)} + 1}}} \right) - \frac{2\sqrt{2}}{\sqrt{e^{(-2x)} + 1}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+tanh(x))^(3/2),x, algorithm="maxima")

[Out] -sqrt(2)*log(-(sqrt(2) - sqrt(2)/sqrt(e^(-2*x) + 1))/(sqrt(2) + sqrt(2)/sqrt(e^(-2*x) + 1))) - 2*sqrt(2)/sqrt(e^(-2*x) + 1)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 129 vs. 2(26) = 52.

time = 0.35, size = 129, normalized size = 3.91

$$\frac{2\sqrt{2}(\sqrt{2}\cosh(x) + \sqrt{2}\sinh(x))\sqrt{\frac{\cosh(x)}{\cosh(x) - \sinh(x)}} - (\sqrt{2}\cosh(x)^2 + 2\sqrt{2}\cosh(x)\sinh(x) + \sqrt{2}\sinh(x)^2 + \sqrt{2})\log\left(-2\sqrt{2}\sqrt{\frac{\cosh(x)}{\cosh(x) - \sinh(x)}}(\cosh(x) + \sinh(x)) - 2\cosh(x)^2 - 4\cosh(x)\sinh(x) - 2\sinh(x)^2 - 1\right)}{\cosh(x)^2 + 2\cosh(x)\sinh(x) + \sinh(x)^2 + 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+tanh(x))^(3/2),x, algorithm="fricas")

[Out] -(2*sqrt(2)*(sqrt(2)*cosh(x) + sqrt(2)*sinh(x))*sqrt(cosh(x)/(cosh(x) - sinh(x))) - (sqrt(2)*cosh(x)^2 + 2*sqrt(2)*cosh(x)*sinh(x) + sqrt(2)*sinh(x)^2 + sqrt(2))*log(-2*sqrt(2)*sqrt(cosh(x)/(cosh(x) - sinh(x)))*(cosh(x) + sinh(x)) - 2*cosh(x)^2 - 4*cosh(x)*sinh(x) - 2*sinh(x)^2 - 1))/(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (\tanh(x) + 1)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+tanh(x))**(3/2),x)

[Out] Integral((tanh(x) + 1)**(3/2), x)

Giac [A]

time = 0.39, size = 52, normalized size = 1.58

$$\sqrt{2} \left(\frac{2}{\sqrt{e^{(4x)} + e^{(2x)}} - e^{(2x)} - 1} - \log\left(-2\sqrt{e^{(4x)} + e^{(2x)}} + 2e^{(2x)} + 1\right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+tanh(x))^(3/2),x, algorithm="giac")

[Out] sqrt(2)*(2/(sqrt(e^(4*x) + e^(2*x)) - e^(2*x) - 1) - log(-2*sqrt(e^(4*x) + e^(2*x)) + 2*e^(2*x) + 1))

Mupad [B]

time = 1.12, size = 26, normalized size = 0.79

$$2\sqrt{2} \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{\tanh(x) + 1}}{2}\right) - 2\sqrt{\tanh(x) + 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((tanh(x) + 1)^(3/2), x)`

[Out] `2*2^(1/2)*atanh((2^(1/2)*(tanh(x) + 1)^(1/2))/2) - 2*(tanh(x) + 1)^(1/2)`

3.53 $\int \sqrt{1 + \tanh(x)} dx$

Optimal. Leaf size=21

$$\sqrt{2} \tanh^{-1} \left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}} \right)$$

[Out] arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))*2^(1/2)

Rubi [A]

time = 0.01, antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3561, 212}

$$\sqrt{2} \tanh^{-1} \left(\frac{\sqrt{\tanh(x) + 1}}{\sqrt{2}} \right)$$

Antiderivative was successfully verified.

[In] Int[Sqrt[1 + Tanh[x]],x]

[Out] Sqrt[2]*ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]]

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 3561

Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] :> Dist[-2*(b/d), Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]

Rubi steps

$$\begin{aligned} \int \sqrt{1 + \tanh(x)} dx &= 2 \text{Subst} \left(\int \frac{1}{2 - x^2} dx, x, \sqrt{1 + \tanh(x)} \right) \\ &= \sqrt{2} \tanh^{-1} \left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}} \right) \end{aligned}$$

Mathematica [A]

time = 0.04, size = 21, normalized size = 1.00

$$\sqrt{2} \tanh^{-1} \left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}} \right)$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[1 + Tanh[x]],x]

[Out] Sqrt[2]*ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]]

Maple [A]

time = 0.66, size = 17, normalized size = 0.81

method	result	size
derivativedivides	$\operatorname{arctanh}\left(\frac{\sqrt{1 + \tanh(x)} \sqrt{2}}{2}\right) \sqrt{2}$	17
default	$\operatorname{arctanh}\left(\frac{\sqrt{1 + \tanh(x)} \sqrt{2}}{2}\right) \sqrt{2}$	17

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+tanh(x))^(1/2),x,method=_RETURNVERBOSE)

[Out] arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))*2^(1/2)

Maxima [B] Leaf count of result is larger than twice the leaf count of optimal. 43 vs. 2(16) = 32.

time = 0.48, size = 43, normalized size = 2.05

$$-\frac{1}{2} \sqrt{2} \log \left(-\frac{\sqrt{2} - \frac{\sqrt{2}}{\sqrt{e^{(-2x)} + 1}}}{\sqrt{2} + \frac{\sqrt{2}}{\sqrt{e^{(-2x)} + 1}}} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+tanh(x))^(1/2),x, algorithm="maxima")

[Out] -1/2*sqrt(2)*log(-(sqrt(2) - sqrt(2)/sqrt(e^(-2*x) + 1))/(sqrt(2) + sqrt(2)/sqrt(e^(-2*x) + 1)))

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 50 vs. 2(16) = 32.

time = 0.38, size = 50, normalized size = 2.38

$$\frac{1}{2} \sqrt{2} \log \left(-2 \sqrt{2} \sqrt{\frac{\cosh(x)}{\cosh(x) - \sinh(x)}} (\cosh(x) + \sinh(x)) - 2 \cosh(x)^2 - 4 \cosh(x) \sinh(x) - 2 \sinh(x)^2 - 1 \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+tanh(x))^(1/2),x, algorithm="fricas")

[Out] $\frac{1}{2}\sqrt{2}\log(-2\sqrt{2}\sqrt{\cosh(x)/(\cosh(x) - \sinh(x))}*(\cosh(x) + \sinh(x)) - 2\cosh(x)^2 - 4\cosh(x)\sinh(x) - 2\sinh(x)^2 - 1)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{\tanh(x) + 1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+tanh(x))**(1/2),x)`

[Out] `Integral(sqrt(tanh(x) + 1), x)`

Giac [A]

time = 0.43, size = 27, normalized size = 1.29

$$-\frac{1}{2}\sqrt{2}\log\left(-2\sqrt{e^{4x} + e^{2x}} + 2e^{2x} + 1\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+tanh(x))^(1/2),x, algorithm="giac")`

[Out] `-1/2*sqrt(2)*log(-2*sqrt(e^(4*x) + e^(2*x)) + 2*e^(2*x) + 1)`

Mupad [B]

time = 0.13, size = 16, normalized size = 0.76

$$\sqrt{2}\operatorname{atanh}\left(\frac{\sqrt{2}\sqrt{\tanh(x) + 1}}{2}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((tanh(x) + 1)^(1/2),x)`

[Out] `2^(1/2)*atanh((2^(1/2)*(tanh(x) + 1)^(1/2))/2)`

$$3.54 \quad \int \frac{1}{\sqrt{1 + \tanh(x)}} dx$$

Optimal. Leaf size=32

$$\frac{\tanh^{-1}\left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{1}{\sqrt{1 + \tanh(x)}}$$

[Out] 1/2*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))*2^(1/2)-1/(1+tanh(x))^(1/2)

Rubi [A]

time = 0.02, antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {3560, 3561, 212}

$$\frac{\tanh^{-1}\left(\frac{\sqrt{\tanh(x) + 1}}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{1}{\sqrt{\tanh(x) + 1}}$$

Antiderivative was successfully verified.

[In] Int[1/Sqrt[1 + Tanh[x]],x]

[Out] ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]]/Sqrt[2] - 1/Sqrt[1 + Tanh[x]]

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 3560

Int[((a_) + (b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n), x] + Dist[1/(2*a), Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]

Rule 3561

Int[Sqrt[(a_) + (b_)*tan[(c_) + (d_)*(x_)]], x_Symbol] := Dist[-2*(b/d), Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]

Rubi steps

$$\begin{aligned}
\int \frac{1}{\sqrt{1 + \tanh(x)}} dx &= -\frac{1}{\sqrt{1 + \tanh(x)}} + \frac{1}{2} \int \sqrt{1 + \tanh(x)} dx \\
&= -\frac{1}{\sqrt{1 + \tanh(x)}} + \text{Subst}\left(\int \frac{1}{2 - x^2} dx, x, \sqrt{1 + \tanh(x)}\right) \\
&= \frac{\tanh^{-1}\left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{1}{\sqrt{1 + \tanh(x)}}
\end{aligned}$$

Mathematica [A]

time = 0.06, size = 32, normalized size = 1.00

$$\frac{\tanh^{-1}\left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{1}{\sqrt{1 + \tanh(x)}}$$

Antiderivative was successfully verified.

`[In] Integrate[1/Sqrt[1 + Tanh[x]], x]``[Out] ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]]/Sqrt[2] - 1/Sqrt[1 + Tanh[x]]`**Maple [A]**

time = 0.63, size = 27, normalized size = 0.84

method	result	size
derivativedivides	$\frac{\operatorname{arctanh}\left(\frac{\sqrt{1 + \tanh(x)} \sqrt{2}}{2}\right) \sqrt{2}}{2} - \frac{1}{\sqrt{1 + \tanh(x)}}$	27
default	$\frac{\operatorname{arctanh}\left(\frac{\sqrt{1 + \tanh(x)} \sqrt{2}}{2}\right) \sqrt{2}}{2} - \frac{1}{\sqrt{1 + \tanh(x)}}$	27

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/(1+tanh(x))^(1/2), x, method=_RETURNVERBOSE)``[Out] 1/2*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))*2^(1/2)-1/(1+tanh(x))^(1/2)`**Maxima [B]** Leaf count of result is larger than twice the leaf count of optimal. 57 vs. $2(26) = 52$.

time = 0.47, size = 57, normalized size = 1.78

$$-\frac{1}{4}\sqrt{2}\log\left(-\frac{\sqrt{2}-\frac{\sqrt{2}}{\sqrt{e^{(-2x)}+1}}}{\sqrt{2}+\frac{\sqrt{2}}{\sqrt{e^{(-2x)}+1}}}\right)-\frac{1}{2}\sqrt{2}\sqrt{e^{(-2x)}+1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+tanh(x))^(1/2),x, algorithm="maxima")

[Out] -1/4*sqrt(2)*log(-(sqrt(2) - sqrt(2)/sqrt(e^(-2*x) + 1))/(sqrt(2) + sqrt(2)/sqrt(e^(-2*x) + 1))) - 1/2*sqrt(2)*sqrt(e^(-2*x) + 1)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 85 vs. 2(26) = 52.

time = 0.58, size = 85, normalized size = 2.66

$$\frac{(\sqrt{2}\cosh(x) + \sqrt{2}\sinh(x))\log\left(-2\sqrt{2}\sqrt{\frac{\cosh(x)}{\cosh(x) - \sinh(x)}}(\cosh(x) + \sinh(x)) - 2\cosh(x)^2 - 4\cosh(x)\sinh(x) - 2\sinh(x)^2 - 1\right) - 4\sqrt{\frac{\cosh(x)}{\cosh(x) - \sinh(x)}}}{4(\cosh(x) + \sinh(x))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+tanh(x))^(1/2),x, algorithm="fricas")

[Out] 1/4*((sqrt(2)*cosh(x) + sqrt(2)*sinh(x))*log(-2*sqrt(2)*sqrt(cosh(x)/(cosh(x) - sinh(x)))*(cosh(x) + sinh(x)) - 2*cosh(x)^2 - 4*cosh(x)*sinh(x) - 2*sinh(x)^2 - 1) - 4*sqrt(cosh(x)/(cosh(x) - sinh(x))))/(cosh(x) + sinh(x))

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{\tanh(x) + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+tanh(x))**(1/2),x)

[Out] Integral(1/sqrt(tanh(x) + 1), x)

Giac [A]

time = 0.42, size = 50, normalized size = 1.56

$$-\frac{1}{4}\sqrt{2}\left(\frac{2}{\sqrt{e^{(4x)}+e^{(2x)}}-e^{(2x)}}+\log\left(-2\sqrt{e^{(4x)}+e^{(2x)}}+2e^{(2x)}+1\right)\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+tanh(x))^(1/2),x, algorithm="giac")

[Out] $-1/4*\sqrt{2}*(2/(\sqrt{e^{4*x} + e^{2*x}}) - e^{2*x}) + \log(-2*\sqrt{e^{4*x} + e^{2*x}}) + 2*e^{2*x} + 1)$

Mupad [B]

time = 0.13, size = 26, normalized size = 0.81

$$\frac{\sqrt{2} \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{\tanh(x) + 1}}{2}\right)}{2} - \frac{1}{\sqrt{\tanh(x) + 1}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(tanh(x) + 1)^(1/2),x)

[Out] $(2^{(1/2)}*\operatorname{atanh}((2^{(1/2)}*(\tanh(x) + 1)^{(1/2)})/2))/2 - 1/(\tanh(x) + 1)^{(1/2)}$

$$3.55 \quad \int \frac{1}{(1+\tanh(x))^{3/2}} dx$$

Optimal. Leaf size=49

$$\frac{\tanh^{-1}\left(\frac{\sqrt{1+\tanh(x)}}{\sqrt{2}}\right)}{2\sqrt{2}} - \frac{1}{3(1+\tanh(x))^{3/2}} - \frac{1}{2\sqrt{1+\tanh(x)}}$$

[Out] 1/4*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))*2^(1/2)-1/2/(1+tanh(x))^(1/2)-1/3/(1+tanh(x))^(3/2)

Rubi [A]

time = 0.02, antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {3560, 3561, 212}

$$\frac{\tanh^{-1}\left(\frac{\sqrt{\tanh(x)+1}}{\sqrt{2}}\right)}{2\sqrt{2}} - \frac{1}{2\sqrt{\tanh(x)+1}} - \frac{1}{3(\tanh(x)+1)^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[(1 + Tanh[x])^(-3/2), x]

[Out] ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]]/(2*Sqrt[2]) - 1/(3*(1 + Tanh[x])^(3/2)) - 1/(2*Sqrt[1 + Tanh[x]])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 3560

Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n)), x] + Dist[1/(2*a), Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]

Rule 3561

Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Dist[-2*(b/d), Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]

Rubi steps

$$\begin{aligned}
\int \frac{1}{(1 + \tanh(x))^{3/2}} dx &= -\frac{1}{3(1 + \tanh(x))^{3/2}} + \frac{1}{2} \int \frac{1}{\sqrt{1 + \tanh(x)}} dx \\
&= -\frac{1}{3(1 + \tanh(x))^{3/2}} - \frac{1}{2\sqrt{1 + \tanh(x)}} + \frac{1}{4} \int \sqrt{1 + \tanh(x)} dx \\
&= -\frac{1}{3(1 + \tanh(x))^{3/2}} - \frac{1}{2\sqrt{1 + \tanh(x)}} + \frac{1}{2} \text{Subst} \left(\int \frac{1}{2 - x^2} dx, x, \sqrt{1 + \tanh(x)} \right) \\
&= \frac{\tanh^{-1} \left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}} \right)}{2\sqrt{2}} - \frac{1}{3(1 + \tanh(x))^{3/2}} - \frac{1}{2\sqrt{1 + \tanh(x)}}
\end{aligned}$$

Mathematica [A]

time = 0.10, size = 53, normalized size = 1.08

$$\frac{1}{12} \left(3\sqrt{2} \tanh^{-1} \left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}} \right) - \frac{2(\cosh(x) - \sinh(x))(5 \cosh(x) + 3 \sinh(x))}{\sqrt{1 + \tanh(x)}} \right)$$

Antiderivative was successfully verified.

`[In] Integrate[(1 + Tanh[x])^(-3/2), x]`

```
[Out] (3*Sqrt[2]*ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]] - (2*(Cosh[x] - Sinh[x])*(5*Cosh[x] + 3*Sinh[x]))/Sqrt[1 + Tanh[x]])/12
```

Maple [A]

time = 0.60, size = 35, normalized size = 0.71

method	result	size
derivativedivides	$ \frac{\operatorname{arctanh} \left(\frac{\sqrt{1 + \tanh(x)} \sqrt{2}}{2} \right) \sqrt{2}}{4} - \frac{1}{2\sqrt{1 + \tanh(x)}} - \frac{1}{3(1 + \tanh(x))^{3/2}} $	35
default	$ \frac{\operatorname{arctanh} \left(\frac{\sqrt{1 + \tanh(x)} \sqrt{2}}{2} \right) \sqrt{2}}{4} - \frac{1}{2\sqrt{1 + \tanh(x)}} - \frac{1}{3(1 + \tanh(x))^{3/2}} $	35

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/(1+tanh(x))^(3/2), x, method=_RETURNVERBOSE)`

```
[Out] 1/4*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))*2^(1/2)-1/2/(1+tanh(x))^(1/2)-1/3/(1+tanh(x))^(3/2)
```


Maxima [B] Leaf count of result is larger than twice the leaf count of optimal. 69 vs. $2(34) = 68$.
time = 0.48, size = 69, normalized size = 1.41

$$-\frac{1}{12} \sqrt{2} \left(\frac{3}{e^{(-2x)} + 1} + 1 \right) (e^{(-2x)} + 1)^{\frac{3}{2}} - \frac{1}{8} \sqrt{2} \log \left(-\frac{\sqrt{2} - \frac{\sqrt{2}}{\sqrt{e^{(-2x)} + 1}}}{\sqrt{2} + \frac{\sqrt{2}}{\sqrt{e^{(-2x)} + 1}}} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+tanh(x))^(3/2),x, algorithm="maxima")

[Out] -1/12*sqrt(2)*(3/(e^(-2*x) + 1) + 1)*(e^(-2*x) + 1)^(3/2) - 1/8*sqrt(2)*log(-(sqrt(2) - sqrt(2)/sqrt(e^(-2*x) + 1))/(sqrt(2) + sqrt(2)/sqrt(e^(-2*x) + 1)))

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 166 vs. $2(34) = 68$.
time = 0.37, size = 166, normalized size = 3.39

$$\frac{2\sqrt{2}(4\sqrt{2}\cosh(x)^2 + 8\sqrt{2}\cosh(x)\sinh(x) + 4\sqrt{2}\sinh(x)^2 + \sqrt{2})\sqrt{\frac{\cosh(x)}{\cosh(x) - \sinh(x)}} - 3(\sqrt{2}\cosh(x)^3 + 3\sqrt{2}\cosh(x)^2\sinh(x) + 3\sqrt{2}\cosh(x)\sinh(x)^2 + \sqrt{2}\sinh(x)^3)\log\left(-2\sqrt{2}\sqrt{\frac{\cosh(x)}{\cosh(x) - \sinh(x)}}(\cosh(x) + \sinh(x)) - 2\cosh(x)^2 - 4\cosh(x)\sinh(x) - 2\sinh(x)^2 - 1\right)}{24(\cosh(x)^3 + 3\cosh(x)^2\sinh(x) + 3\cosh(x)\sinh(x)^2 + \sinh(x)^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+tanh(x))^(3/2),x, algorithm="fricas")

[Out] -1/24*(2*sqrt(2)*(4*sqrt(2)*cosh(x)^2 + 8*sqrt(2)*cosh(x)*sinh(x) + 4*sqrt(2)*sinh(x)^2 + sqrt(2))*sqrt(cosh(x)/(cosh(x) - sinh(x))) - 3*(sqrt(2)*cosh(x)^3 + 3*sqrt(2)*cosh(x)^2*sinh(x) + 3*sqrt(2)*cosh(x)*sinh(x)^2 + sqrt(2)*sinh(x)^3)*log(-2*sqrt(2)*sqrt(cosh(x)/(cosh(x) - sinh(x)))*(cosh(x) + sinh(x)) - 2*cosh(x)^2 - 4*cosh(x)*sinh(x) - 2*sinh(x)^2 - 1))/(cosh(x)^3 + 3*cosh(x)^2*sinh(x) + 3*cosh(x)*sinh(x)^2 + sinh(x)^3)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(\tanh(x) + 1)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+tanh(x))**(3/2),x)

[Out] Integral((tanh(x) + 1)**(-3/2), x)

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 95 vs. $2(34) = 68$.
time = 0.41, size = 95, normalized size = 1.94

$$-\frac{1}{24} \sqrt{2} \left(\frac{2 \left(6 \left(\sqrt{e^{(4x)} + e^{(2x)}} - e^{(2x)} \right)^2 - 3 \sqrt{e^{(4x)} + e^{(2x)}} + 3e^{(2x)} + 1 \right)}{\left(\sqrt{e^{(4x)} + e^{(2x)}} - e^{(2x)} \right)^3} + 3 \log \left(-2 \sqrt{e^{(4x)} + e^{(2x)}} + 2e^{(2x)} + 1 \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+tanh(x))^(3/2),x, algorithm="giac")

[Out] -1/24*sqrt(2)*(2*(6*(sqrt(e^(4*x) + e^(2*x)) - e^(2*x))^2 - 3*sqrt(e^(4*x) + e^(2*x)) + 3*e^(2*x) + 1)/(sqrt(e^(4*x) + e^(2*x)) - e^(2*x))^3 + 3*log(-2*sqrt(e^(4*x) + e^(2*x)) + 2*e^(2*x) + 1))

Mupad [B]

time = 0.12, size = 32, normalized size = 0.65

$$\frac{\sqrt{2} \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{\tanh(x) + 1}}{2}\right)}{4} - \frac{\frac{\tanh(x)}{2} + \frac{5}{6}}{(\tanh(x) + 1)^{3/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(tanh(x) + 1)^(3/2),x)

[Out] (2^(1/2)*atanh((2^(1/2)*(tanh(x) + 1)^(1/2))/2))/4 - (tanh(x)/2 + 5/6)/(tanh(x) + 1)^(3/2)

$$3.56 \quad \int \frac{1}{(1+\tanh(x))^{5/2}} dx$$

Optimal. Leaf size=61

$$\frac{\tanh^{-1}\left(\frac{\sqrt{1+\tanh(x)}}{\sqrt{2}}\right)}{4\sqrt{2}} - \frac{1}{5(1+\tanh(x))^{5/2}} - \frac{1}{6(1+\tanh(x))^{3/2}} - \frac{1}{4\sqrt{1+\tanh(x)}}$$

[Out] 1/8*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))*2^(1/2)-1/4/(1+tanh(x))^(1/2)-1/5/(1+tanh(x))^(5/2)-1/6/(1+tanh(x))^(3/2)

Rubi [A]

time = 0.03, antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {3560, 3561, 212}

$$\frac{\tanh^{-1}\left(\frac{\sqrt{\tanh(x)+1}}{\sqrt{2}}\right)}{4\sqrt{2}} - \frac{1}{4\sqrt{\tanh(x)+1}} - \frac{1}{6(\tanh(x)+1)^{3/2}} - \frac{1}{5(\tanh(x)+1)^{5/2}}$$

Antiderivative was successfully verified.

[In] Int[(1 + Tanh[x])^(-5/2), x]

[Out] ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]]/(4*Sqrt[2]) - 1/(5*(1 + Tanh[x])^(5/2)) - 1/(6*(1 + Tanh[x])^(3/2)) - 1/(4*Sqrt[1 + Tanh[x]])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 3560

Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n)), x] + Dist[1/(2*a), Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]

Rule 3561

Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Dist[-2*(b/d), Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]

Rubi steps

$$\begin{aligned}
\int \frac{1}{(1 + \tanh(x))^{5/2}} dx &= -\frac{1}{5(1 + \tanh(x))^{5/2}} + \frac{1}{2} \int \frac{1}{(1 + \tanh(x))^{3/2}} dx \\
&= -\frac{1}{5(1 + \tanh(x))^{5/2}} - \frac{1}{6(1 + \tanh(x))^{3/2}} + \frac{1}{4} \int \frac{1}{\sqrt{1 + \tanh(x)}} dx \\
&= -\frac{1}{5(1 + \tanh(x))^{5/2}} - \frac{1}{6(1 + \tanh(x))^{3/2}} - \frac{1}{4\sqrt{1 + \tanh(x)}} + \frac{1}{8} \int \sqrt{1 + \tanh(x)} dx \\
&= -\frac{1}{5(1 + \tanh(x))^{5/2}} - \frac{1}{6(1 + \tanh(x))^{3/2}} - \frac{1}{4\sqrt{1 + \tanh(x)}} + \frac{1}{4} \text{Subst} \left(\int \frac{1}{2 - x^2} \right. \\
&\quad \left. \tanh^{-1} \left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}} \right) \right) \\
&= \frac{\tanh^{-1} \left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}} \right)}{4\sqrt{2}} - \frac{1}{5(1 + \tanh(x))^{5/2}} - \frac{1}{6(1 + \tanh(x))^{3/2}} - \frac{1}{4\sqrt{1 + \tanh(x)}}
\end{aligned}$$

Mathematica [A]

time = 0.17, size = 62, normalized size = 1.02

$$\frac{\tanh^{-1} \left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}} \right)}{4\sqrt{2}} + \frac{(-\cosh(2x) + \sinh(2x))(11 + 26 \cosh(2x) + 20 \sinh(2x))}{60\sqrt{1 + \tanh(x)}}$$

Antiderivative was successfully verified.

`[In] Integrate[(1 + Tanh[x])^(-5/2), x]`

```
[Out] ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]]/(4*Sqrt[2]) + ((-Cosh[2*x] + Sinh[2*x])*(11 + 26*Cosh[2*x] + 20*Sinh[2*x]))/(60*Sqrt[1 + Tanh[x]])
```

Maple [A]

time = 0.59, size = 43, normalized size = 0.70

method	result	size
derivativedivides	$\frac{\operatorname{arctanh} \left(\frac{\sqrt{1 + \tanh(x)} \sqrt{2}}{2} \right) \sqrt{2}}{8} - \frac{1}{4\sqrt{1 + \tanh(x)}} - \frac{1}{5(1 + \tanh(x))^{5/2}} - \frac{1}{6(1 + \tanh(x))^{3/2}}$	43
default	$\frac{\operatorname{arctanh} \left(\frac{\sqrt{1 + \tanh(x)} \sqrt{2}}{2} \right) \sqrt{2}}{8} - \frac{1}{4\sqrt{1 + \tanh(x)}} - \frac{1}{5(1 + \tanh(x))^{5/2}} - \frac{1}{6(1 + \tanh(x))^{3/2}}$	43

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/(1+tanh(x))^(5/2), x, method=_RETURNVERBOSE)`

[Out] $\frac{1}{8} \operatorname{arctanh}\left(\frac{1}{2} \cdot (1 + \tanh(x))^{1/2}\right) \cdot 2^{1/2} - \frac{1}{4} (1 + \tanh(x))^{1/2} - \frac{1}{5} (1 + \tanh(x))^{5/2} - \frac{1}{6} (1 + \tanh(x))^{3/2}$

Maxima [A]

time = 0.47, size = 79, normalized size = 1.30

$$-\frac{1}{120} \sqrt{2} \left(\frac{5}{e^{(-2x)} + 1} + \frac{15}{(e^{(-2x)} + 1)^2} + 3 \right) (e^{(-2x)} + 1)^{\frac{5}{2}} - \frac{1}{16} \sqrt{2} \log \left(-\frac{\sqrt{2} - \frac{\sqrt{2}}{\sqrt{e^{(-2x)} + 1}}}{\sqrt{2} + \frac{\sqrt{2}}{\sqrt{e^{(-2x)} + 1}}} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+tanh(x))^(5/2),x, algorithm="maxima")`

[Out] $-1/120 \sqrt{2} (5/(e^{(-2x)} + 1) + 15/(e^{(-2x)} + 1)^2 + 3) (e^{(-2x)} + 1)^{5/2} - 1/16 \sqrt{2} \log(-(\sqrt{2} - \sqrt{2}/\sqrt{e^{(-2x)} + 1})/(\sqrt{2} + \sqrt{2}/\sqrt{e^{(-2x)} + 1}))$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 266 vs. 2(42) = 84.

time = 0.47, size = 266, normalized size = 4.36

$$\frac{2 \sqrt{2} (23 \sqrt{2} \cosh(x)^2 + 22 \sqrt{2} \cosh(x) \sinh(x) + 23 \sqrt{2} \sinh(x)^2 + (138 \sqrt{2} \cosh(x)^2 + 11 \sqrt{2}) \sinh(x) + 11 \sqrt{2} \cosh(x)^2 + 2 (46 \sqrt{2} \cosh(x) + 11 \sqrt{2} \sinh(x)) \sinh(x) + 3 \sqrt{2})}{340 \cosh(x)^2 + 5 \cosh(x) \sinh(x) + 10 \sinh(x)^2 + 10 \cosh(x)^2 \sinh(x) + 5 \sinh(x) \cosh(x) + \sinh(x)^2} \log \left(-\frac{\sqrt{2} \cosh(x) - 1}{\sqrt{2} \cosh(x) + 1} \right) \frac{\cosh(x) + \sinh(x)}{\cosh(x) - \sinh(x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+tanh(x))^(5/2),x, algorithm="fricas")`

[Out] $-1/240 (2 \sqrt{2} (23 \sqrt{2} \cosh(x)^2 + 22 \sqrt{2} \cosh(x) \sinh(x) + 23 \sqrt{2} \sinh(x)^2 + (138 \sqrt{2} \cosh(x)^2 + 11 \sqrt{2}) \sinh(x) + 11 \sqrt{2} \cosh(x)^2 + 2 (46 \sqrt{2} \cosh(x) + 11 \sqrt{2} \sinh(x)) \sinh(x) + 3 \sqrt{2}) \sqrt{\cosh(x) / (\cosh(x) - \sinh(x))} - 15 (\sqrt{2} \cosh(x)^5 + 5 \sqrt{2} \cosh(x)^4 \sinh(x) + 10 \sqrt{2} \cosh(x)^3 \sinh(x)^2 + 10 \sqrt{2} \cosh(x)^2 \sinh(x)^3 + 5 \sqrt{2} \cosh(x) \sinh(x)^4 + \sqrt{2} \sinh(x)^5) \log(-2 \sqrt{2} \sqrt{\cosh(x) / (\cosh(x) - \sinh(x))} (\cosh(x) + \sinh(x)) - 2 \cosh(x)^2 - 4 \cosh(x) \sinh(x) - 2 \sinh(x)^2 - 1) / (\cosh(x)^5 + 5 \cosh(x)^4 \sinh(x) + 10 \cosh(x)^3 \sinh(x)^2 + 10 \cosh(x)^2 \sinh(x)^3 + 5 \cosh(x) \sinh(x)^4 + \sinh(x)^5)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(\tanh(x) + 1)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+tanh(x))**(5/2),x)`

[Out] Integral((tanh(x) + 1)**(-5/2), x)

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 139 vs. 2(42) = 84.

time = 0.42, size = 139, normalized size = 2.28

$$-\frac{1}{240} \sqrt{2} \left(\frac{2 \left(45 \left(\sqrt{e^{4x} + e^{2x}} - e^{2x} \right)^4 - 45 \left(\sqrt{e^{4x} + e^{2x}} - e^{2x} \right)^3 + 35 \left(\sqrt{e^{4x} + e^{2x}} - e^{2x} \right)^2 - 15 \sqrt{e^{4x} + e^{2x}} + 15 e^{2x} + 3 \right)}{\left(\sqrt{e^{4x} + e^{2x}} - e^{2x} \right)^5} + 15 \log \left(-2 \sqrt{e^{4x} + e^{2x}} + 2 e^{2x} + 1 \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+tanh(x))^(5/2),x, algorithm="giac")

[Out] -1/240*sqrt(2)*(2*(45*(sqrt(e^(4*x) + e^(2*x)) - e^(2*x))^4 - 45*(sqrt(e^(4*x) + e^(2*x)) - e^(2*x))^3 + 35*(sqrt(e^(4*x) + e^(2*x)) - e^(2*x))^2 - 15*sqrt(e^(4*x) + e^(2*x)) + 15*e^(2*x) + 3)/(sqrt(e^(4*x) + e^(2*x)) - e^(2*x))^5 + 15*log(-2*sqrt(e^(4*x) + e^(2*x)) + 2*e^(2*x) + 1))

Mupad [B]

time = 0.12, size = 40, normalized size = 0.66

$$\frac{\sqrt{2} \operatorname{atanh} \left(\frac{\sqrt{2} \sqrt{\tanh(x) + 1}}{2} \right)}{8} - \frac{\frac{\tanh(x)}{6} + \frac{(\tanh(x)+1)^2}{4} + \frac{11}{30}}{(\tanh(x) + 1)^{5/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(tanh(x) + 1)^(5/2),x)

[Out] (2^(1/2)*atanh((2^(1/2)*(tanh(x) + 1)^(1/2))/2))/8 - (tanh(x)/6 + (tanh(x) + 1)^2/4 + 11/30)/(tanh(x) + 1)^(5/2)

3.57 $\int (a + b \tanh(c + dx))^5 dx$

Optimal. Leaf size=142

$$a(a^4 + 10a^2b^2 + 5b^4)x + \frac{b(5a^4 + 10a^2b^2 + b^4) \log(\cosh(c + dx))}{d} - \frac{4ab^2(a^2 + b^2) \tanh(c + dx)}{d} - \frac{b(3a^2 + b^2)}{d}$$

[Out] a*(a^4+10*a^2*b^2+5*b^4)*x+b*(5*a^4+10*a^2*b^2+b^4)*ln(cosh(d*x+c))/d-4*a*b^2*(a^2+b^2)*tanh(d*x+c)/d-1/2*b*(3*a^2+b^2)*(a+b*tanh(d*x+c))^2/d-2/3*a*b*(a+b*tanh(d*x+c))^3/d-1/4*b*(a+b*tanh(d*x+c))^4/d

Rubi [A]

time = 0.15, antiderivative size = 142, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3563, 3609, 3606, 3556}

$$-\frac{b(3a^2 + b^2)(a + b \tanh(c + dx))^2}{2d} - \frac{4ab^2(a^2 + b^2) \tanh(c + dx)}{d} + \frac{b(5a^4 + 10a^2b^2 + b^4) \log(\cosh(c + dx))}{d} + ax(a^4 + 10a^2b^2 + 5b^4) - \frac{b(a + b \tanh(c + dx))^4}{4d} - \frac{2ab(a + b \tanh(c + dx))^3}{3d}$$

Antiderivative was successfully verified.

[In] Int[(a + b*Tanh[c + d*x])^5,x]

[Out] a*(a^4 + 10*a^2*b^2 + 5*b^4)*x + (b*(5*a^4 + 10*a^2*b^2 + b^4)*Log[Cosh[c + d*x]])/d - (4*a*b^2*(a^2 + b^2)*Tanh[c + d*x])/d - (b*(3*a^2 + b^2)*(a + b*Tanh[c + d*x])^2)/(2*d) - (2*a*b*(a + b*Tanh[c + d*x])^3)/(3*d) - (b*(a + b*Tanh[c + d*x])^4)/(4*d)

Rule 3556

Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]

Rule 3563

Int[((a_.) + (b_.)*tan[(c_.) + (d_.)*(x_.)])^(n_), x_Symbol] := Simp[b*((a + b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] + Int[(a^2 - b^2 + 2*a*b*Tan[c + d*x])*(a + b*Tan[c + d*x])^(n - 2), x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0] && GtQ[n, 1]

Rule 3606

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Dist[b*c + a*d, Int[Tan[e + f*x], x], x] + Simp[b*d*(Tan[e + f*x]/f), x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]

Rule 3609

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])], x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int
[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x]
, x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2,
0] && GtQ[m, 0]
```

Rubi steps

$$\begin{aligned}
\int (a + b \tanh(c + dx))^5 dx &= -\frac{b(a + b \tanh(c + dx))^4}{4d} + \int (a + b \tanh(c + dx))^3 (a^2 + b^2 + 2ab \tanh(c + dx)) dx \\
&= -\frac{2ab(a + b \tanh(c + dx))^3}{3d} - \frac{b(a + b \tanh(c + dx))^4}{4d} + \int (a + b \tanh(c + dx)) dx \\
&= -\frac{b(3a^2 + b^2)(a + b \tanh(c + dx))^2}{2d} - \frac{2ab(a + b \tanh(c + dx))^3}{3d} - \frac{b(a + b \tanh(c + dx))^4}{4d} \\
&= a(a^4 + 10a^2b^2 + 5b^4)x - \frac{4ab^2(a^2 + b^2) \tanh(c + dx)}{d} - \frac{b(3a^2 + b^2)(a + b \tanh(c + dx))^4}{2d} \\
&= a(a^4 + 10a^2b^2 + 5b^4)x + \frac{b(5a^4 + 10a^2b^2 + b^4) \log(\cosh(c + dx))}{d} - \frac{4ab^2(a^2 + b^2) \tanh(c + dx)}{d}
\end{aligned}$$

Mathematica [A]

time = 0.46, size = 114, normalized size = 0.80

$$\frac{6(a+b)^5 \log(1 - \tanh(c+dx)) - 6(a-b)^5 \log(1 + \tanh(c+dx)) + 60ab^2(2a^2 + b^2) \tanh(c+dx) + 6b^3(10a^2 + b^2) \tanh^2(c+dx) + 20ab^4 \tanh^3(c+dx) + 3b^5 \tanh^4(c+dx)}{12d}$$

Antiderivative was successfully verified.

```
[In] Integrate[(a + b*Tanh[c + d*x])^5, x]
```

```
[Out] -1/12*(6*(a + b)^5*Log[1 - Tanh[c + d*x]] - 6*(a - b)^5*Log[1 + Tanh[c + d*x]] + 60*a*b^2*(2*a^2 + b^2)*Tanh[c + d*x] + 6*b^3*(10*a^2 + b^2)*Tanh[c + d*x]^2 + 20*a*b^4*Tanh[c + d*x]^3 + 3*b^5*Tanh[c + d*x]^4)/d
```

Maple [A]

time = 0.27, size = 182, normalized size = 1.28

method	result
derivativedivides	$-\frac{b^5 \tanh^4(dx+c)}{4} - 5a^2 b^3 \tanh^2(dx+c) - \frac{5a b^4 \tanh^3(dx+c)}{3} - 5a b^4 \tanh(dx+c) - 10a^3 b^2 \tanh(dx+c) - \frac{b^5 \tanh^2(dx+c)}{2}$
default	$-\frac{b^5 \tanh^4(dx+c)}{4} - 5a^2 b^3 \tanh^2(dx+c) - \frac{5a b^4 \tanh^3(dx+c)}{3} - 5a b^4 \tanh(dx+c) - 10a^3 b^2 \tanh(dx+c) - \frac{b^5 \tanh^2(dx+c)}{2}$
risch	$a^5 x - 5a^4 b x + 10a^3 b^2 x - 10a^2 b^3 x + 5a b^4 x - b^5 x - \frac{10b a^4 c}{d} - \frac{20b^3 a^2 c}{d} - \frac{2b^5 c}{d} + \frac{4b^2 (15a^3 e^{6dx+c} + \dots)}{d}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a+b*tanh(d*x+c))^5,x,method=_RETURNVERBOSE)`

[Out] $1/d*(-1/4*b^5*tanh(d*x+c)^4-5*a^2*b^3*tanh(d*x+c)^2-5/3*a*b^4*tanh(d*x+c)^3-5*a*b^4*tanh(d*x+c)-10*a^3*b^2*tanh(d*x+c)-1/2*b^5*tanh(d*x+c)^2-1/2*(a^5+5*a^4*b+10*a^3*b^2+10*a^2*b^3+5*a*b^4+b^5)*\ln(\tanh(d*x+c)-1)+1/2*(a^5-5*a^4*b+10*a^3*b^2-10*a^2*b^3+5*a*b^4-b^5)*\ln(\tanh(d*x+c)+1))$

Maxima [B] Leaf count of result is larger than twice the leaf count of optimal. 310 vs. $2(136) = 272$.

time = 0.48, size = 310, normalized size = 2.18

$$\frac{5}{3}ab^4\left(3x + \frac{3c}{d} - \frac{4(3e^{d(-2dx-2c)} + 3e^{d(-4dx-4c)} + 2)}{d(3e^{d(-2dx-2c)} + 3e^{d(-4dx-4c)} + 1)}\right) + b^5\left(x + \frac{c}{d} + \frac{\log(e^{d(-2dx-2c)} + 1)}{d} + \frac{4(e^{d(-2dx-2c)} + e^{d(-4dx-4c)} + e^{d(-6dx-6c)})}{d(4e^{d(-2dx-2c)} + 6e^{d(-4dx-4c)} + 4e^{d(-6dx-6c)} + d^{d(-8dx-8c)} + 1)}\right) + 10a^2b^3\left(x + \frac{c}{d} + \frac{\log(e^{d(-2dx-2c)} + 1)}{d} + \frac{2e^{d(-2dx-2c)}}{d(2e^{d(-2dx-2c)} + e^{d(-4dx-4c)} + 1)}\right) + 10a^2b^3\left(x + \frac{c}{d} - \frac{2}{d(2e^{d(-2dx-2c)} + e^{d(-4dx-4c)} + 1)}\right) + a^5x + \frac{5a^4b \log(\cosh(dx+c))}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+b*tanh(d*x+c))^5,x, algorithm="maxima")`

[Out] $5/3*a*b^4*(3*x + 3*c/d - 4*(3*e^{(-2*d*x - 2*c)} + 3*e^{(-4*d*x - 4*c)} + 2)/(d*(3*e^{(-2*d*x - 2*c)} + 3*e^{(-4*d*x - 4*c)} + e^{(-6*d*x - 6*c)} + 1))) + b^5*(x + c/d + \log(e^{(-2*d*x - 2*c)} + 1)/d + 4*(e^{(-2*d*x - 2*c)} + e^{(-4*d*x - 4*c)} + e^{(-6*d*x - 6*c)})/(d*(4*e^{(-2*d*x - 2*c)} + 6*e^{(-4*d*x - 4*c)} + 4*e^{(-6*d*x - 6*c)} + e^{(-8*d*x - 8*c)} + 1))) + 10*a^2*b^3*(x + c/d + \log(e^{(-2*d*x - 2*c)} + 1)/d + 2*e^{(-2*d*x - 2*c)}/(d*(2*e^{(-2*d*x - 2*c)} + e^{(-4*d*x - 4*c)} + 1))) + 10*a^2*b^3*(x + c/d - 2/(d*(e^{(-2*d*x - 2*c)} + 1))) + a^5*x + 5*a^4*b*\log(\cosh(dx+c))/d$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 2739 vs. $2(136) = 272$.

time = 0.65, size = 2739, normalized size = 19.29

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+b*tanh(d*x+c))^5,x, algorithm="fricas")`

[Out] $1/3*(3*(a^5 - 5*a^4*b + 10*a^3*b^2 - 10*a^2*b^3 + 5*a*b^4 - b^5)*d*x*\cosh(dx+c)^8 + 24*(a^5 - 5*a^4*b + 10*a^3*b^2 - 10*a^2*b^3 + 5*a*b^4 - b^5)*d*x*\cosh(dx+c)*\sinh(dx+c)^7 + 3*(a^5 - 5*a^4*b + 10*a^3*b^2 - 10*a^2*b^3 + 5*a*b^4 - b^5)*d*x*\sinh(dx+c)^8 + 12*(5*a^3*b^2 + 5*a^2*b^3 + 5*a*b^4 + b^5 + (a^5 - 5*a^4*b + 10*a^3*b^2 - 10*a^2*b^3 + 5*a*b^4 - b^5)*d*x)*\cosh(dx+c)^6 + 12*(5*a^3*b^2 + 5*a^2*b^3 + 5*a*b^4 + b^5 + 7*(a^5 - 5*a^4*b + 10*a^3*b^2 - 10*a^2*b^3 + 5*a*b^4 - b^5)*d*x*\cosh(dx+c)^2 + (a^5 - 5*a^4*b + 10*a^3*b^2 - 10*a^2*b^3 + 5*a*b^4 - b^5)*d*x)*\sinh(dx+c)^6 + 24$

$$\begin{aligned}
&*(7*(a^5 - 5*a^4*b + 10*a^3*b^2 - 10*a^2*b^3 + 5*a*b^4 - b^5)*d*x*cosh(d*x \\
&+ c)^3 + 3*(5*a^3*b^2 + 5*a^2*b^3 + 5*a*b^4 + b^5 + (a^5 - 5*a^4*b + 10*a^3 \\
&*b^2 - 10*a^2*b^3 + 5*a*b^4 - b^5)*d*x)*cosh(d*x + c))*sinh(d*x + c)^5 + 60 \\
&*a^3*b^2 + 40*a*b^4 + 6*(30*a^3*b^2 + 20*a^2*b^3 + 20*a*b^4 + 2*b^5 + 3*(a^5 \\
&- 5*a^4*b + 10*a^3*b^2 - 10*a^2*b^3 + 5*a*b^4 - b^5)*d*x)*cosh(d*x + c)^4 \\
&+ 6*(35*(a^5 - 5*a^4*b + 10*a^3*b^2 - 10*a^2*b^3 + 5*a*b^4 - b^5)*d*x*cosh \\
&(d*x + c)^4 + 30*a^3*b^2 + 20*a^2*b^3 + 20*a*b^4 + 2*b^5 + 3*(a^5 - 5*a^4*b \\
&+ 10*a^3*b^2 - 10*a^2*b^3 + 5*a*b^4 - b^5)*d*x + 30*(5*a^3*b^2 + 5*a^2*b^3 \\
&+ 5*a*b^4 + b^5 + (a^5 - 5*a^4*b + 10*a^3*b^2 - 10*a^2*b^3 + 5*a*b^4 - b^5 \\
&)*d*x)*cosh(d*x + c)^2)*sinh(d*x + c)^4 + 24*(7*(a^5 - 5*a^4*b + 10*a^3*b^2 \\
&- 10*a^2*b^3 + 5*a*b^4 - b^5)*d*x*cosh(d*x + c)^5 + 10*(5*a^3*b^2 + 5*a^2* \\
&b^3 + 5*a*b^4 + b^5 + (a^5 - 5*a^4*b + 10*a^3*b^2 - 10*a^2*b^3 + 5*a*b^4 - \\
&b^5)*d*x)*cosh(d*x + c)^3 + (30*a^3*b^2 + 20*a^2*b^3 + 20*a*b^4 + 2*b^5 + 3 \\
&*(a^5 - 5*a^4*b + 10*a^3*b^2 - 10*a^2*b^3 + 5*a*b^4 - b^5)*d*x)*cosh(d*x + \\
&c))*sinh(d*x + c)^3 + 3*(a^5 - 5*a^4*b + 10*a^3*b^2 - 10*a^2*b^3 + 5*a*b^4 \\
&- b^5)*d*x + 4*(45*a^3*b^2 + 15*a^2*b^3 + 25*a*b^4 + 3*b^5 + 3*(a^5 - 5*a^4 \\
&*b + 10*a^3*b^2 - 10*a^2*b^3 + 5*a*b^4 - b^5)*d*x)*cosh(d*x + c)^2 + 4*(21* \\
&(a^5 - 5*a^4*b + 10*a^3*b^2 - 10*a^2*b^3 + 5*a*b^4 - b^5)*d*x*cosh(d*x + c) \\
&^6 + 45*a^3*b^2 + 15*a^2*b^3 + 25*a*b^4 + 3*b^5 + 45*(5*a^3*b^2 + 5*a^2*b^3 \\
&+ 5*a*b^4 + b^5 + (a^5 - 5*a^4*b + 10*a^3*b^2 - 10*a^2*b^3 + 5*a*b^4 - b^5 \\
&)*d*x)*cosh(d*x + c)^4 + 3*(a^5 - 5*a^4*b + 10*a^3*b^2 - 10*a^2*b^3 + 5*a*b \\
&^4 - b^5)*d*x + 9*(30*a^3*b^2 + 20*a^2*b^3 + 20*a*b^4 + 2*b^5 + 3*(a^5 - 5* \\
&a^4*b + 10*a^3*b^2 - 10*a^2*b^3 + 5*a*b^4 - b^5)*d*x)*cosh(d*x + c)^2)*sinh \\
&(d*x + c)^2 + 3*((5*a^4*b + 10*a^2*b^3 + b^5)*cosh(d*x + c)^8 + 8*(5*a^4*b \\
&+ 10*a^2*b^3 + b^5)*cosh(d*x + c)*sinh(d*x + c)^7 + (5*a^4*b + 10*a^2*b^3 + \\
&b^5)*sinh(d*x + c)^8 + 4*(5*a^4*b + 10*a^2*b^3 + b^5)*cosh(d*x + c)^6 + 4* \\
&(5*a^4*b + 10*a^2*b^3 + b^5 + 7*(5*a^4*b + 10*a^2*b^3 + b^5)*cosh(d*x + c)^ \\
&2)*sinh(d*x + c)^6 + 8*(7*(5*a^4*b + 10*a^2*b^3 + b^5)*cosh(d*x + c)^3 + 3* \\
&(5*a^4*b + 10*a^2*b^3 + b^5)*cosh(d*x + c))*sinh(d*x + c)^5 + 5*a^4*b + 10* \\
&a^2*b^3 + b^5 + 6*(5*a^4*b + 10*a^2*b^3 + b^5)*cosh(d*x + c)^4 + 2*(15*a^4*b \\
&b + 30*a^2*b^3 + 3*b^5 + 35*(5*a^4*b + 10*a^2*b^3 + b^5)*cosh(d*x + c)^4 + \\
&30*(5*a^4*b + 10*a^2*b^3 + b^5)*cosh(d*x + c)^2)*sinh(d*x + c)^4 + 8*(7*(5* \\
&a^4*b + 10*a^2*b^3 + b^5)*cosh(d*x + c)^5 + 10*(5*a^4*b + 10*a^2*b^3 + b^5) \\
&)*cosh(d*x + c)^3 + 3*(5*a^4*b + 10*a^2*b^3 + b^5)*cosh(d*x + c))*sinh(d*x + \\
&c)^3 + 4*(5*a^4*b + 10*a^2*b^3 + b^5)*cosh(d*x + c)^2 + 4*(7*(5*a^4*b + 10 \\
&a^2*b^3 + b^5)*cosh(d*x + c)^6 + 5*a^4*b + 10*a^2*b^3 + b^5 + 15*(5*a^4*b \\
&+ 10*a^2*b^3 + b^5)*cosh(d*x + c)^4 + 9*(5*a^4*b + 10*a^2*b^3 + b^5)*cosh(d \\
&*x + c)^2)*sinh(d*x + c)^2 + 8*((5*a^4*b + 10*a^2*b^3 + b^5)*cosh(d*x + c)^ \\
&7 + 3*(5*a^4*b + 10*a^2*b^3 + b^5)*cosh(d*x + c)^5 + 3*(5*a^4*b + 10*a^2*b^ \\
&3 + b^5)*cosh(d*x + c)^3 + (5*a^4*b + 10*a^2*b^3 + b^5)*cosh(d*x + c))*sinh \\
&(d*x + c))*log(2*cosh(d*x + c)/(cosh(d*x + c) - sinh(d*x + c))) + 8*(3*(a^5 \\
&- 5*a^4*b + 10*a^3*b^2 - 10*a^2*b^3 + 5*a*b^4 - b^5)*d*x*cosh(d*x + c)^7 + \\
&9*(5*a^3*b^2 + 5*a^2*b^3 + 5*a*b^4 + b^5 + (a^5 - 5*a^4*b + 10*a^3*b^2 - 1 \\
&0*a^2*b^3 + 5*a*b^4 - b^5)*d*x)*cosh(d*x + c)^5 + 3*(30*a^3*b^2 + 20*a^2*b^ \\
&3 + 20*a*b^4 + 2*b^5 + 3*(a^5 - 5*a^4*b + 10*a^3*b^2 - 10*a^2*b^3 + 5*a*b^4
\end{aligned}$$

$$- b^5 * d * x) * \cosh(d * x + c)^3 + (45 * a^3 * b^2 + 15 * a^2 * b^3 + 25 * a * b^4 + 3 * b^5 + 3 * (a^5 - 5 * a^4 * b + 10 * a^3 * b^2 - 10 * a^2 * b^3 + 5 * a * b^4 - b^5) * d * x) * \cosh(d * x + c) * \sinh(d * x + c) / (d * \cosh(d * x + c)^8 + 8 * d * \cosh(d * x + c) * \sinh(d * x + c)^7 + d * \sinh(d * x + c)^8 + 4 * d * \cosh(d * x + c)^6 + 4 * (7 * d * \cosh(d * x + c)^2 + d) * \sinh(d * x + c)^6 + 8 * (7 * d * \cosh(d * x + c)^3 + 3 * d * \cosh(d * x + c)) * \sinh(d * x + c)^5 + 6 * d * \cosh(d * x + c)^4 + 2 * (35 * d * \cosh(d * x + c)^4 + 30 * d * \cosh(d * x + c)^2 + 3 * d) * \sinh(d * x + c)^4 + 8 * (7 * d * \cosh(d * x + c)^5 + 10 * d * \cosh(d * x + c)^3 + 3 * d * \cosh(d * x + c)) * \sinh(d * x + c)^3 + 4 * d * \cosh(d * x + c)^2 + 4 * (7 * d * \cosh(d * x + c)^6 + 15 * d * \cosh(d * x + c)^4 + 9 * d * \cosh(d * x + c)^2 + d) * \sinh(d * x + c)^2 + 8 * (d * \cosh(d * x + c)^7 + 3 * d * \cosh(d * x + c)^5 + 3 * d * \cosh(d * x + c)^3 + d * \cosh(d * x + c)) * \sinh(d * x + c) + d)$$

Sympy [A]

time = 0.16, size = 211, normalized size = 1.49

$$\left\{ \begin{array}{l} \frac{a^5 x + 5a^4 b x - \frac{5a^3 b \log(\tanh(\frac{c+dx}{d})+1)}{d} + 10a^2 b^2 x - \frac{10a^2 b^2 \tanh(\frac{c+dx}{d})}{d} + 10a^2 b^2 x - \frac{10a^2 b^2 \log(\tanh(\frac{c+dx}{d})+1)}{d} - \frac{5a^2 b^2 \tanh^2(\frac{c+dx}{d})}{d} + 5ab^4 x - \frac{5ab^4 \tanh^3(\frac{c+dx}{d})}{3d} - \frac{5ab^4 \tanh(\frac{c+dx}{d})}{d} + b^5 x - \frac{b^5 \log(\tanh(\frac{c+dx}{d})+1)}{d} - \frac{b^5 \tanh^4(\frac{c+dx}{d})}{4d} - \frac{b^5 \tanh^2(\frac{c+dx}{d})}{2d} \end{array} \right. \text{for } d \neq 0$$

otherwise

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+b*tanh(d*x+c))**5,x)

[Out] Piecewise((a**5*x + 5*a**4*b*x - 5*a**4*b*log(tanh(c + d*x) + 1)/d + 10*a**3*b**2*x - 10*a**3*b**2*tanh(c + d*x)/d + 10*a**2*b**3*x - 10*a**2*b**3*log(tanh(c + d*x) + 1)/d - 5*a**2*b**3*tanh(c + d*x)**2/d + 5*a*b**4*x - 5*a*b**4*tanh(c + d*x)**3/(3*d) - 5*a*b**4*tanh(c + d*x)/d + b**5*x - b**5*log(tanh(c + d*x) + 1)/d - b**5*tanh(c + d*x)**4/(4*d) - b**5*tanh(c + d*x)**2/(2*d), Ne(d, 0)), (x*(a + b*tanh(c))**5, True))

Giac [A]

time = 0.43, size = 224, normalized size = 1.58

$$\frac{3(a^5 - 5a^4b + 10a^3b^2 - 10a^2b^3 + 5ab^4 - b^5)(dx + c) + 3(5a^4b + 10a^2b^3 + b^5) \log(e^{(2dx+2c)} + 1) + \frac{4(15a^3b^2 + 10ab^4 + 3(5a^3b^2 + 5a^2b^3 + 5ab^4 + b^5)e^{(6dx+6c)} + 3(15a^3b^2 + 10a^2b^3 + 10ab^4 + b^5)e^{(4dx+4c)} + (45a^3b^2 + 15a^2b^3 + 25ab^4 + 3b^5)e^{(2dx+2c)})}{(e^{(2dx+2c)} + 1)^4}}{3d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+b*tanh(d*x+c))^5,x, algorithm="giac")

[Out] 1/3*(3*(a^5 - 5*a^4*b + 10*a^3*b^2 - 10*a^2*b^3 + 5*a*b^4 - b^5)*(d*x + c) + 3*(5*a^4*b + 10*a^2*b^3 + b^5)*log(e^(2*d*x + 2*c) + 1) + 4*(15*a^3*b^2 + 10*a*b^4 + 3*(5*a^3*b^2 + 5*a^2*b^3 + 5*a*b^4 + b^5)*e^(6*d*x + 6*c) + 3*(15*a^3*b^2 + 10*a^2*b^3 + 10*a*b^4 + b^5)*e^(4*d*x + 4*c) + (45*a^3*b^2 + 15*a^2*b^3 + 25*a*b^4 + 3*b^5)*e^(2*d*x + 2*c))/(e^(2*d*x + 2*c) + 1)^4/d

Mupad [B]

time = 1.16, size = 153, normalized size = 1.08

$$x(a^5 + 5a^4b + 10a^3b^2 + 10a^2b^3 + 5ab^4 + b^5) - \frac{5 \tanh(c + dx)(2a^3b^2 + ab^4)}{d} - \frac{b^5 \tanh(c + dx)^2}{4d} - \frac{\ln(\tanh(c + dx) + 1)(5a^4b + 10a^2b^3 + b^5)}{d} - \frac{\tanh(c + dx)^2(10a^2b^3 + b^5)}{2d} - \frac{5ab^4 \tanh(c + dx)^3}{3d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*tanh(c + d*x))^5,x)

[Out] $x*(5*a*b^4 + 5*a^4*b + a^5 + b^5 + 10*a^2*b^3 + 10*a^3*b^2) - (5*\tanh(c + d*x)*(a*b^4 + 2*a^3*b^2))/d - (b^5*\tanh(c + d*x)^4)/(4*d) - (\log(\tanh(c + d*x) + 1)*(5*a^4*b + b^5 + 10*a^2*b^3))/d - (\tanh(c + d*x)^2*(b^5 + 10*a^2*b^3))/(2*d) - (5*a*b^4*\tanh(c + d*x)^3)/(3*d)$

3.58 $\int (a + b \tanh(c + dx))^4 dx$

Optimal. Leaf size=101

$$(a^4 + 6a^2b^2 + b^4)x + \frac{4ab(a^2 + b^2) \log(\cosh(c + dx))}{d} - \frac{b^2(3a^2 + b^2) \tanh(c + dx)}{d} - \frac{ab(a + b \tanh(c + dx))^2}{d}$$

[Out] $(a^4 + 6a^2b^2 + b^4)x + 4a*b*(a^2 + b^2)*\ln(\cosh(d*x + c))/d - b^2*(3a^2 + b^2)*\tanh(d*x + c)/d - a*b*(a + b*\tanh(d*x + c))^2/d - 1/3*b*(a + b*\tanh(d*x + c))^3/d$

Rubi [A]

time = 0.09, antiderivative size = 101, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3563, 3609, 3606, 3556}

$$-\frac{b^2(3a^2 + b^2) \tanh(c + dx)}{d} + \frac{4ab(a^2 + b^2) \log(\cosh(c + dx))}{d} + x(a^4 + 6a^2b^2 + b^4) - \frac{b(a + b \tanh(c + dx))^3}{3d} - \frac{ab(a + b \tanh(c + dx))^2}{d}$$

Antiderivative was successfully verified.

[In] Int[(a + b*Tanh[c + d*x])^4, x]

[Out] $(a^4 + 6a^2b^2 + b^4)x + (4a*b*(a^2 + b^2)*\text{Log}[\text{Cosh}[c + d*x]])/d - (b^2*(3a^2 + b^2)*\text{Tanh}[c + d*x])/d - (a*b*(a + b*\text{Tanh}[c + d*x])^2)/d - (b*(a + b*\text{Tanh}[c + d*x])^3)/(3*d)$

Rule 3556

Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]

Rule 3563

Int[((a_.) + (b_.)*tan[(c_.) + (d_.)*(x_.)])^(n_), x_Symbol] := Simp[b*((a + b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] + Int[(a^2 - b^2 + 2*a*b*Tan[c + d*x])*(a + b*Tan[c + d*x])^(n - 2), x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0] && GtQ[n, 1]

Rule 3606

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Dist[b*c + a*d, Int[Tan[e + f*x], x], x] + Simp[b*d*(Tan[e + f*x]/f), x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]

Rule 3609

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int

```
[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x]
, x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2,
0] && GtQ[m, 0]
```

Rubi steps

$$\begin{aligned} \int (a + b \tanh(c + dx))^4 dx &= -\frac{b(a + b \tanh(c + dx))^3}{3d} + \int (a + b \tanh(c + dx))^2 (a^2 + b^2 + 2ab \tanh(c + dx)) dx \\ &= -\frac{ab(a + b \tanh(c + dx))^2}{d} - \frac{b(a + b \tanh(c + dx))^3}{3d} + \int (a + b \tanh(c + dx)) (a^2 + b^2 + 2ab \tanh(c + dx)) dx \\ &= (a^4 + 6a^2b^2 + b^4)x - \frac{b^2(3a^2 + b^2) \tanh(c + dx)}{d} - \frac{ab(a + b \tanh(c + dx))^2}{d} - \frac{b^3 \tanh^3(c + dx)}{3d} \\ &= (a^4 + 6a^2b^2 + b^4)x + \frac{4ab(a^2 + b^2) \log(\cosh(c + dx))}{d} - \frac{b^2(3a^2 + b^2) \tanh(c + dx)}{d} \end{aligned}$$

Mathematica [A]

time = 0.24, size = 91, normalized size = 0.90

$$\frac{3(a+b)^4 \log(1 - \tanh(c+dx)) - 3(a-b)^4 \log(1 + \tanh(c+dx)) + 6b^2(6a^2 + b^2) \tanh(c+dx) + 12ab^3 \tanh^2(c+dx) + 2b^4 \tanh^3(c+dx)}{6d}$$

Antiderivative was successfully verified.

```
[In] Integrate[(a + b*Tanh[c + d*x])^4, x]
```

```
[Out] -1/6*(3*(a + b)^4*Log[1 - Tanh[c + d*x]] - 3*(a - b)^4*Log[1 + Tanh[c + d*x]
]) + 6*b^2*(6*a^2 + b^2)*Tanh[c + d*x] + 12*a*b^3*Tanh[c + d*x]^2 + 2*b^4*T
anh[c + d*x]^3)/d
```

Maple [A]

time = 0.27, size = 134, normalized size = 1.33

method	result
derivativedivides	$-\frac{b^4(\tanh^3(dx+c))}{3} - 2ab^3(\tanh^2(dx+c)) - 6a^2b^2 \tanh(dx+c) - b^4 \tanh(dx+c) - \frac{(a^4 + 4a^3b + 6a^2b^2 + 4ab^3 + b^4) \ln(\tanh(dx+c))}{2d}$
default	$-\frac{b^4(\tanh^3(dx+c))}{3} - 2ab^3(\tanh^2(dx+c)) - 6a^2b^2 \tanh(dx+c) - b^4 \tanh(dx+c) - \frac{(a^4 + 4a^3b + 6a^2b^2 + 4ab^3 + b^4) \ln(\tanh(dx+c))}{2d}$
risch	$xa^4 - 4a^3bx + 6a^2b^2x - 4b^3ax + b^4x - \frac{8ba^3c}{d} - \frac{8b^3ac}{d} + \frac{4b^2(9a^2e^{4dx+4c} + 6abe^{4dx+4c} + 3b^2e^{4dx+4c} + 3b^3e^{4dx+4c})}{3d(1 - \tanh^2(dx+c))}$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((a+b*tanh(d*x+c))^4, x, method=_RETURNVERBOSE)
```

[Out] $1/d*(-1/3*b^4*\tanh(d*x+c)^3-2*a*b^3*\tanh(d*x+c)^2-6*a^2*b^2*\tanh(d*x+c)-b^4*\tanh(d*x+c)-1/2*(a^4+4*a^3*b+6*a^2*b^2+4*a*b^3+b^4)*\ln(\tanh(d*x+c)-1)+1/2*(a^4-4*a^3*b+6*a^2*b^2-4*a*b^3+b^4)*\ln(\tanh(d*x+c)+1))$

Maxima [B] Leaf count of result is larger than twice the leaf count of optimal. 201 vs. 2(99) = 198.

time = 0.47, size = 201, normalized size = 1.99

$$\frac{1}{3}b^4\left(3x + \frac{3c}{d} - \frac{4(3e^{(-2dx-2c)} + 3e^{(-4dx-4c)} + 2)}{d(3e^{(-2dx-2c)} + 3e^{(-4dx-4c)} + e^{(-6dx-6c)} + 1)}\right) + 4ab^3\left(x + \frac{c}{d} + \frac{\log(e^{(-2dx-2c)} + 1)}{d} + \frac{2e^{(-2dx-2c)}}{d(2e^{(-2dx-2c)} + e^{(-4dx-4c)} + 1)}\right) + 6a^2b^2\left(x + \frac{c}{d} - \frac{2}{d(e^{(-2dx-2c)} + 1)}\right) + a^4x + \frac{4a^3b \log(\cosh(dx + c))}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+b*tanh(d*x+c))^4,x, algorithm="maxima")`

[Out] $1/3*b^4*(3*x + 3*c/d - 4*(3*e^{(-2*d*x - 2*c)} + 3*e^{(-4*d*x - 4*c)} + 2)/(d*(3*e^{(-2*d*x - 2*c)} + 3*e^{(-4*d*x - 4*c)} + e^{(-6*d*x - 6*c)} + 1))) + 4*a*b^3*(x + c/d + \log(e^{(-2*d*x - 2*c)} + 1)/d + 2*e^{(-2*d*x - 2*c)}/(d*(2*e^{(-2*d*x - 2*c)} + e^{(-4*d*x - 4*c)} + 1))) + 6*a^2*b^2*(x + c/d - 2/(d*(e^{(-2*d*x - 2*c)} + 1))) + a^4*x + 4*a^3*b*\log(\cosh(d*x + c))/d$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 1389 vs. 2(99) = 198.

time = 0.37, size = 1389, normalized size = 13.75

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+b*tanh(d*x+c))^4,x, algorithm="fricas")`

[Out] $1/3*(3*(a^4 - 4*a^3*b + 6*a^2*b^2 - 4*a*b^3 + b^4)*d*x*\cosh(d*x + c)^6 + 18*(a^4 - 4*a^3*b + 6*a^2*b^2 - 4*a*b^3 + b^4)*d*x*\cosh(d*x + c)*\sinh(d*x + c)^5 + 3*(a^4 - 4*a^3*b + 6*a^2*b^2 - 4*a*b^3 + b^4)*d*x*\sinh(d*x + c)^6 + 3*(12*a^2*b^2 + 8*a*b^3 + 4*b^4 + 3*(a^4 - 4*a^3*b + 6*a^2*b^2 - 4*a*b^3 + b^4)*d*x)*\cosh(d*x + c)^4 + 3*(15*(a^4 - 4*a^3*b + 6*a^2*b^2 - 4*a*b^3 + b^4)*d*x*\cosh(d*x + c)^2 + 12*a^2*b^2 + 8*a*b^3 + 4*b^4 + 3*(a^4 - 4*a^3*b + 6*a^2*b^2 - 4*a*b^3 + b^4)*d*x)*\sinh(d*x + c)^4 + 36*a^2*b^2 + 8*b^4 + 12*(5*(a^4 - 4*a^3*b + 6*a^2*b^2 - 4*a*b^3 + b^4)*d*x*\cosh(d*x + c)^3 + (12*a^2*b^2 + 8*a*b^3 + 4*b^4 + 3*(a^4 - 4*a^3*b + 6*a^2*b^2 - 4*a*b^3 + b^4)*d*x)*\cosh(d*x + c))*\sinh(d*x + c)^3 + 3*(a^4 - 4*a^3*b + 6*a^2*b^2 - 4*a*b^3 + b^4)*d*x + 3*(24*a^2*b^2 + 8*a*b^3 + 4*b^4 + 3*(a^4 - 4*a^3*b + 6*a^2*b^2 - 4*a*b^3 + b^4)*d*x)*\cosh(d*x + c)^2 + 3*(15*(a^4 - 4*a^3*b + 6*a^2*b^2 - 4*a*b^3 + b^4)*d*x*\cosh(d*x + c)^4 + 24*a^2*b^2 + 8*a*b^3 + 4*b^4 + 3*(a^4 - 4*a^3*b + 6*a^2*b^2 - 4*a*b^3 + b^4)*d*x + 6*(12*a^2*b^2 + 8*a*b^3 + 4*b^4 + 3*(a^4 - 4*a^3*b + 6*a^2*b^2 - 4*a*b^3 + b^4)*d*x)*\cosh(d*x + c)^2)*\sinh(d*x + c)^2 + 12*((a^3*b + a*b^3)*\cosh(d*x + c)^6 + 6*(a^3*b + a*b^3)*\cosh(d*x + c)*\sinh(d*x + c)^5 + (a^3*b + a*b^3)*\sinh(d*x + c)^6 + 3*(a^3*b + a*b^3)*\cosh(d*x + c)^4 + 3*(a^3*b + a*b^3 + 5*(a^3*b + a*b^3)*\cosh(d*x + c)^2)*$

$$\begin{aligned} & \sinh(dx + c)^4 + a^3b + ab^3 + 4*(5*(a^3b + ab^3)*\cosh(dx + c)^3 + 3* \\ & (a^3b + ab^3)*\cosh(dx + c))*\sinh(dx + c)^3 + 3*(a^3b + ab^3)*\cosh(dx \\ & + c)^2 + 3*(5*(a^3b + ab^3)*\cosh(dx + c)^4 + a^3b + ab^3 + 6*(a^3b + \\ & ab^3)*\cosh(dx + c)^2)*\sinh(dx + c)^2 + 6*((a^3b + ab^3)*\cosh(dx + c) \\ & ^5 + 2*(a^3b + ab^3)*\cosh(dx + c)^3 + (a^3b + ab^3)*\cosh(dx + c))*\sin \\ & h(dx + c))*\log(2*\cosh(dx + c)/(\cosh(dx + c) - \sinh(dx + c))) + 6*(3*(a^ \\ & 4 - 4*a^3b + 6*a^2b^2 - 4*ab^3 + b^4)*d*x*\cosh(dx + c)^5 + 2*(12*a^2b^ \\ & 2 + 8*ab^3 + 4*b^4 + 3*(a^4 - 4*a^3b + 6*a^2b^2 - 4*ab^3 + b^4)*d*x)*co \\ & sh(dx + c)^3 + (24*a^2b^2 + 8*ab^3 + 4*b^4 + 3*(a^4 - 4*a^3b + 6*a^2b^2 - \\ & 4*ab^3 + b^4)*d*x)*\cosh(dx + c))*\sinh(dx + c))/(d*\cosh(dx + c)^6 + \\ & 6*d*\cosh(dx + c)*\sinh(dx + c)^5 + d*\sinh(dx + c)^6 + 3*d*\cosh(dx + c)^4 \\ & + 3*(5*d*\cosh(dx + c)^2 + d)*\sinh(dx + c)^4 + 4*(5*d*\cosh(dx + c)^3 + 3 \\ & *d*\cosh(dx + c))*\sinh(dx + c)^3 + 3*d*\cosh(dx + c)^2 + 3*(5*d*\cosh(dx + \\ & c)^4 + 6*d*\cosh(dx + c)^2 + d)*\sinh(dx + c)^2 + 6*(d*\cosh(dx + c)^5 + 2 \\ & *d*\cosh(dx + c)^3 + d*\cosh(dx + c))*\sinh(dx + c) + d) \end{aligned}$$

Sympy [A]

time = 0.13, size = 144, normalized size = 1.43

$$\begin{cases} a^4x + 4a^3bx - \frac{4a^3b \log(\tanh(\frac{c+dx}{d})+1)}{d} + 6a^2b^2x - \frac{6a^2b^2 \tanh(\frac{c+dx}{d})}{d} + 4ab^3x - \frac{4ab^3 \log(\tanh(\frac{c+dx}{d})+1)}{d} - \frac{2ab^3 \tanh^2(\frac{c+dx}{d})}{d} + b^4x - \frac{b^4 \tanh^3(\frac{c+dx}{d})}{3d} - \frac{b^4 \tanh(\frac{c+dx}{d})}{d} & \text{for } d \neq 0 \\ x(a + b \tanh(c))^4 & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+b*tanh(dx+c))**4,x)

[Out] Piecewise((a**4*x + 4*a**3*b*x - 4*a**3*b*log(tanh(c + dx) + 1)/d + 6*a**2*b**2*x - 6*a**2*b**2*tanh(c + dx)/d + 4*a*b**3*x - 4*a*b**3*log(tanh(c + dx) + 1)/d - 2*a*b**3*tanh(c + dx)**2/d + b**4*x - b**4*tanh(c + dx)**3/(3*d) - b**4*tanh(c + dx)/d, Ne(d, 0)), (x*(a + b*tanh(c))**4, True))

Giac [A]

time = 0.43, size = 152, normalized size = 1.50

$$\frac{3(a^4 - 4a^3b + 6a^2b^2 - 4ab^3 + b^4)(dx + c) + 12(a^3b + ab^3) \log(e^{(2dx+2c)} + 1) + \frac{4(9a^2b^2 + 2b^4 + 3(3a^2b^2 + 2ab^3 + b^4)e^{(4dx+4c)} + 3(6a^2b^2 + 2ab^3 + b^4)e^{(2dx+2c)})}{(e^{(2dx+2c)} + 1)^3}}{3d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+b*tanh(dx+c))⁴,x, algorithm="giac")

[Out] 1/3*(3*(a⁴ - 4*a³b + 6*a²b² - 4*ab³ + b⁴)*(dx + c) + 12*(a³b + ab³)*log(e^(2*dx + 2*c) + 1) + 4*(9*a²b² + 2*b⁴ + 3*(3*a²b² + 2*a*b³ + b⁴)*e^(4*d*x + 4*c) + 3*(6*a²b² + 2*a*b³ + b⁴)*e^(2*d*x + 2*c))/(e^(2*d*x + 2*c) + 1)³/d

Mupad [B]

time = 1.09, size = 113, normalized size = 1.12

$$x(a^4 + 4a^3b + 6a^2b^2 + 4ab^3 + b^4) - \frac{b^4 \tanh(c + dx)^3}{3d} - \frac{\ln(\tanh(c + dx) + 1)(4a^3b + 4ab^3)}{d} - \frac{2ab^3 \tanh(c + dx)^2}{d} - \frac{b^2 \tanh(c + dx)(6a^2 + b^2)}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*tanh(c + d*x))^4,x)`

[Out] $x*(4*a*b^3 + 4*a^3*b + a^4 + b^4 + 6*a^2*b^2) - (b^4*tanh(c + d*x)^3)/(3*d)$
 $- (\log(tanh(c + d*x) + 1)*(4*a*b^3 + 4*a^3*b))/d - (2*a*b^3*tanh(c + d*x)^2)/d - (b^2*tanh(c + d*x)*(6*a^2 + b^2))/d$

3.59 $\int (a + b \tanh(c + dx))^3 dx$

Optimal. Leaf size=69

$$a(a^2 + 3b^2)x + \frac{b(3a^2 + b^2) \log(\cosh(c + dx))}{d} - \frac{2ab^2 \tanh(c + dx)}{d} - \frac{b(a + b \tanh(c + dx))^2}{2d}$$

[Out] a*(a^2+3*b^2)*x+b*(3*a^2+b^2)*ln(cosh(d*x+c))/d-2*a*b^2*tanh(d*x+c)/d-1/2*b*(a+b*tanh(d*x+c))^2/d

Rubi [A]

time = 0.05, antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3563, 3606, 3556}

$$\frac{b(3a^2 + b^2) \log(\cosh(c + dx))}{d} + ax(a^2 + 3b^2) - \frac{2ab^2 \tanh(c + dx)}{d} - \frac{b(a + b \tanh(c + dx))^2}{2d}$$

Antiderivative was successfully verified.

[In] Int[(a + b*Tanh[c + d*x])^3,x]

[Out] a*(a^2 + 3*b^2)*x + (b*(3*a^2 + b^2)*Log[Cosh[c + d*x]])/d - (2*a*b^2*Tanh[c + d*x])/d - (b*(a + b*Tanh[c + d*x])^2)/(2*d)

Rule 3556

Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]

Rule 3563

Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((a + b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] + Int[(a^2 - b^2 + 2*a*b*Tan[c + d*x])*(a + b*Tan[c + d*x])^(n - 2), x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0] && GtQ[n, 1]

Rule 3606

Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Dist[b*c + a*d, Int[Tan[e + f*x], x], x] + Simp[b*d*(Tan[e + f*x]/f), x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]

Rubi steps

$$\begin{aligned} \int (a + b \tanh(c + dx))^3 dx &= -\frac{b(a + b \tanh(c + dx))^2}{2d} + \int (a + b \tanh(c + dx)) (a^2 + b^2 + 2ab \tanh(c + dx)) dx \\ &= a(a^2 + 3b^2) x - \frac{2ab^2 \tanh(c + dx)}{d} - \frac{b(a + b \tanh(c + dx))^2}{2d} + (b(3a^2 + b^2)) \\ &= a(a^2 + 3b^2) x + \frac{b(3a^2 + b^2) \log(\cosh(c + dx))}{d} - \frac{2ab^2 \tanh(c + dx)}{d} - \frac{b(a + b \tanh(c + dx))^2}{2d} \end{aligned}$$

Mathematica [A]

time = 0.24, size = 67, normalized size = 0.97

$$\frac{(a + b)^3 \log(1 - \tanh(c + dx)) - (a - b)^3 \log(1 + \tanh(c + dx)) + 6ab^2 \tanh(c + dx) + b^3 \tanh^2(c + dx)}{2d}$$

Antiderivative was successfully verified.

`[In] Integrate[(a + b*Tanh[c + d*x])^3,x]`

```
[Out] -1/2*((a + b)^3*Log[1 - Tanh[c + d*x]] - (a - b)^3*Log[1 + Tanh[c + d*x]] +
6*a*b^2*Tanh[c + d*x] + b^3*Tanh[c + d*x]^2)/d
```

Maple [A]

time = 0.26, size = 93, normalized size = 1.35

method	result
derivativedivides	$\frac{-\frac{b^3 (\tanh^2(dx+c))}{2} - 3ab^2 \tanh(dx+c) - \frac{(a^3 + 3a^2b + 3ab^2 + b^3) \ln(\tanh(dx+c)-1)}{2} + \frac{(a^3 - 3a^2b + 3ab^2 - b^3) \ln(\tanh(dx+c)+1)}{2}}{d}$
default	$\frac{-\frac{b^3 (\tanh^2(dx+c))}{2} - 3ab^2 \tanh(dx+c) - \frac{(a^3 + 3a^2b + 3ab^2 + b^3) \ln(\tanh(dx+c)-1)}{2} + \frac{(a^3 - 3a^2b + 3ab^2 - b^3) \ln(\tanh(dx+c)+1)}{2}}{d}$
risch	$a^3x - 3a^2bx + 3ab^2x - b^3x - \frac{6a^2bc}{d} - \frac{2b^3c}{d} + \frac{2b^2(3ae^{2dx+2c} + be^{2dx+2c} + 3a)}{d(1+e^{2dx+2c})^2} + \frac{3b \ln(1+e^{2dx+2c})a^2}{d}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((a+b*tanh(d*x+c))^3,x,method=_RETURNVERBOSE)`

```
[Out] 1/d*(-1/2*b^3*tanh(d*x+c)^2-3*a*b^2*tanh(d*x+c)-1/2*(a^3+3*a^2*b+3*a*b^2+b^3)*ln(tanh(d*x+c)-1)+1/2*(a^3-3*a^2*b+3*a*b^2-b^3)*ln(tanh(d*x+c)+1))
```

Maxima [A]

time = 0.48, size = 118, normalized size = 1.71

$$b^3 \left(x + \frac{c}{d} + \frac{\log(e^{-2dx-2c} + 1)}{d} + \frac{2e^{-2dx-2c}}{d(2e^{-2dx-2c} + e^{-4dx-4c} + 1)} \right) + 3ab^2 \left(x + \frac{c}{d} - \frac{2}{d(e^{-2dx-2c} + 1)} \right) + a^3x + \frac{3a^2b \log(\cosh(dx+c))}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+b*tanh(d*x+c))^3,x, algorithm="maxima")

[Out] $b^3*(x + c/d + \log(e^{-2*d*x - 2*c} + 1)/d + 2*e^{-2*d*x - 2*c}/(d*(2*e^{-2*d*x - 2*c} + e^{-4*d*x - 4*c} + 1))) + 3*a*b^2*(x + c/d - 2/(d*(e^{-2*d*x - 2*c} + 1))) + a^3*x + 3*a^2*b*\log(\cosh(d*x + c))/d$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 646 vs. $2(67) = 134$.

time = 0.41, size = 646, normalized size = 9.36

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+b*tanh(d*x+c))^3,x, algorithm="fricas")

[Out] $((a^3 - 3*a^2*b + 3*a*b^2 - b^3)*d*x*\cosh(d*x + c)^4 + 4*(a^3 - 3*a^2*b + 3*a*b^2 - b^3)*d*x*\cosh(d*x + c)*\sinh(d*x + c)^3 + (a^3 - 3*a^2*b + 3*a*b^2 - b^3)*d*x*\sinh(d*x + c)^4 + 6*a*b^2 + (a^3 - 3*a^2*b + 3*a*b^2 - b^3)*d*x + 2*(3*a*b^2 + b^3 + (a^3 - 3*a^2*b + 3*a*b^2 - b^3)*d*x)*\cosh(d*x + c)^2 + 2*(3*(a^3 - 3*a^2*b + 3*a*b^2 - b^3)*d*x*\cosh(d*x + c)^2 + 3*a*b^2 + b^3 + (a^3 - 3*a^2*b + 3*a*b^2 - b^3)*d*x)*\sinh(d*x + c)^2 + ((3*a^2*b + b^3)*\cosh(d*x + c)^4 + 4*(3*a^2*b + b^3)*\cosh(d*x + c)*\sinh(d*x + c)^3 + (3*a^2*b + b^3)*\sinh(d*x + c)^4 + 3*a^2*b + b^3 + 2*(3*a^2*b + b^3)*\cosh(d*x + c)^2 + 2*(3*a^2*b + b^3 + 3*(3*a^2*b + b^3)*\cosh(d*x + c)^2)*\sinh(d*x + c)^2 + 4*((3*a^2*b + b^3)*\cosh(d*x + c)^3 + (3*a^2*b + b^3)*\cosh(d*x + c))*\sinh(d*x + c))*\log(2*\cosh(d*x + c)/(\cosh(d*x + c) - \sinh(d*x + c))) + 4*((a^3 - 3*a^2*b + 3*a*b^2 - b^3)*d*x*\cosh(d*x + c)^3 + (3*a*b^2 + b^3 + (a^3 - 3*a^2*b + 3*a*b^2 - b^3)*d*x)*\cosh(d*x + c))*\sinh(d*x + c))/(d*\cosh(d*x + c)^4 + 4*d*\cosh(d*x + c)*\sinh(d*x + c)^3 + d*\sinh(d*x + c)^4 + 2*d*\cosh(d*x + c)^2 + 2*(3*d*\cosh(d*x + c)^2 + d)*\sinh(d*x + c)^2 + 4*(d*\cosh(d*x + c)^3 + d*\cosh(d*x + c))*\sinh(d*x + c) + d)$

Sympy [A]

time = 0.11, size = 100, normalized size = 1.45

$$\begin{cases} a^3x + 3a^2bx - \frac{3a^2b \log(\tanh(c+dx)+1)}{d} + 3ab^2x - \frac{3ab^2 \tanh(c+dx)}{d} + b^3x - \frac{b^3 \log(\tanh(c+dx)+1)}{d} - \frac{b^3 \tanh^2(c+dx)}{2d} & \text{for } d \neq 0 \\ x(a + b \tanh(c))^3 & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+b*tanh(d*x+c))**3,x)

[Out] $\text{Piecewise}((a**3*x + 3*a**2*b*x - 3*a**2*b*\log(\tanh(c + d*x) + 1)/d + 3*a*b**2*x - 3*a*b**2*\tanh(c + d*x)/d + b**3*x - b**3*\log(\tanh(c + d*x) + 1)/d - b**3*\tanh(c + d*x)**2/(2*d), \text{Ne}(d, 0)), (x*(a + b*\tanh(c))**3, \text{True}))$

Giac [A]

time = 0.44, size = 97, normalized size = 1.41

$$\frac{(a^3 - 3a^2b + 3ab^2 - b^3)(dx + c) + (3a^2b + b^3) \log(e^{(2dx+2c)} + 1) + \frac{2(3ab^2 + (3ab^2 + b^3)e^{(2dx+2c)})}{(e^{(2dx+2c)} + 1)^2}}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+b*tanh(d*x+c))^3,x, algorithm="giac")

[Out] ((a^3 - 3*a^2*b + 3*a*b^2 - b^3)*(d*x + c) + (3*a^2*b + b^3)*log(e^(2*d*x + 2*c) + 1) + 2*(3*a*b^2 + (3*a*b^2 + b^3)*e^(2*d*x + 2*c)))/(e^(2*d*x + 2*c) + 1)^2/d

Mupad [B]

time = 0.11, size = 77, normalized size = 1.12

$$x(a^3 + 3a^2b + 3ab^2 + b^3) - \frac{\ln(\tanh(c + dx) + 1)(3a^2b + b^3)}{d} - \frac{b^3 \tanh(c + dx)^2}{2d} - \frac{3ab^2 \tanh(c + dx)}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*tanh(c + d*x))^3,x)

[Out] x*(3*a*b^2 + 3*a^2*b + a^3 + b^3) - (log(tanh(c + d*x) + 1)*(3*a^2*b + b^3))/d - (b^3*tanh(c + d*x)^2)/(2*d) - (3*a*b^2*tanh(c + d*x))/d

3.60 $\int (a + b \tanh(c + dx))^2 dx$

Optimal. Leaf size=38

$$(a^2 + b^2)x + \frac{2ab \log(\cosh(c + dx))}{d} - \frac{b^2 \tanh(c + dx)}{d}$$

[Out] $(a^2+b^2)*x+2*a*b*\ln(\cosh(d*x+c))/d-b^2*\tanh(d*x+c)/d$

Rubi [A]

time = 0.02, antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3558, 3556}

$$x(a^2 + b^2) + \frac{2ab \log(\cosh(c + dx))}{d} - \frac{b^2 \tanh(c + dx)}{d}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(a + b*\text{Tanh}[c + d*x])^2, x]$

[Out] $(a^2 + b^2)*x + (2*a*b*\text{Log}[\text{Cosh}[c + d*x]])/d - (b^2*\text{Tanh}[c + d*x])/d$

Rule 3556

$\text{Int}[\tan[(c_.) + (d_.)*(x_.)], x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] /; \text{FreeQ}\{c, d\}, x]$

Rule 3558

$\text{Int}[((a_.) + (b_.)*\tan[(c_.) + (d_.)*(x_.)])^2, x_Symbol] \rightarrow \text{Simp}[(a^2 - b^2)*x, x] + (\text{Dist}[2*a*b, \text{Int}[\text{Tan}[c + d*x], x], x] + \text{Simp}[b^2*(\text{Tan}[c + d*x]/d), x]) /; \text{FreeQ}\{a, b, c, d\}, x]$

Rubi steps

$$\begin{aligned} \int (a + b \tanh(c + dx))^2 dx &= (a^2 + b^2)x - \frac{b^2 \tanh(c + dx)}{d} + (2ab) \int \tanh(c + dx) dx \\ &= (a^2 + b^2)x + \frac{2ab \log(\cosh(c + dx))}{d} - \frac{b^2 \tanh(c + dx)}{d} \end{aligned}$$

Mathematica [A]

time = 0.08, size = 54, normalized size = 1.42

$$\frac{-(a + b)^2 \log(1 - \tanh(c + dx)) + (a - b)^2 \log(1 + \tanh(c + dx)) - 2b^2 \tanh(c + dx)}{2d}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*Tanh[c + d*x])^2,x]

[Out] $(-((a + b)^2 \cdot \text{Log}[1 - \text{Tanh}[c + d \cdot x]]) + (a - b)^2 \cdot \text{Log}[1 + \text{Tanh}[c + d \cdot x]] - 2 \cdot b^2 \cdot \text{Tanh}[c + d \cdot x]) / (2 \cdot d)$

Maple [A]

time = 0.26, size = 61, normalized size = 1.61

method	result	size
derivativedivides	$\frac{-b^2 \tanh(dx+c) - \frac{(a^2+2ab+b^2) \ln(\tanh(dx+c)-1)}{2} + \frac{(a^2-2ab+b^2) \ln(\tanh(dx+c)+1)}{2}}{d}$	61
default	$\frac{-b^2 \tanh(dx+c) - \frac{(a^2+2ab+b^2) \ln(\tanh(dx+c)-1)}{2} + \frac{(a^2-2ab+b^2) \ln(\tanh(dx+c)+1)}{2}}{d}$	61
risch	$a^2x - 2abx + b^2x - \frac{4abc}{d} + \frac{2b^2}{d(1+e^{2dx+2c})} + \frac{2ab \ln(1+e^{2dx+2c})}{d}$	65

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a+b*tanh(d*x+c))^2,x,method=_RETURNVERBOSE)

[Out] $1/d * (-b^2 * \tanh(d*x+c) - 1/2 * (a^2 + 2*a*b + b^2) * \ln(\tanh(d*x+c) - 1) + 1/2 * (a^2 - 2*a*b + b^2) * \ln(\tanh(d*x+c) + 1))$

Maxima [A]

time = 0.27, size = 49, normalized size = 1.29

$$b^2 \left(x + \frac{c}{d} - \frac{2}{d(e^{(-2dx-2c)} + 1)} \right) + a^2x + \frac{2ab \log(\cosh(dx+c))}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+b*tanh(d*x+c))^2,x, algorithm="maxima")

[Out] $b^2 * (x + c/d - 2/(d * (e^{(-2*d*x - 2*c)} + 1))) + a^2 * x + 2 * a * b * \log(\cosh(d * x + c)) / d$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 201 vs. 2(38) = 76.

time = 0.38, size = 201, normalized size = 5.29

$$\frac{(a^2 - 2ab + b^2) dx \cosh(dx+c)^2 + 2(a^2 - 2ab + b^2) dx \cosh(dx+c) \sinh(dx+c) + (a^2 - 2ab + b^2) dx \sinh(dx+c)^2 + (a^2 - 2ab + b^2) dx + 2b^2 + 2(ab \cosh(dx+c)^2 + 2ab \cosh(dx+c) \sinh(dx+c) + ab \sinh(dx+c)^2 + ab) \log\left(\frac{2 \cosh(dx+c)}{\cosh(dx+c) - \sinh(dx+c)}\right)}{d \cosh(dx+c)^2 + 2d \cosh(dx+c) \sinh(dx+c) + d \sinh(dx+c)^2 + d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+b*tanh(d*x+c))^2,x, algorithm="fricas")

[Out] $((a^2 - 2*a*b + b^2) * d * x * \cosh(d * x + c)^2 + 2 * (a^2 - 2*a*b + b^2) * d * x * \cosh(d * x + c) * \sinh(d * x + c) + (a^2 - 2*a*b + b^2) * d * x * \sinh(d * x + c)^2 + (a^2 - 2 *$

$a*b + b^2)*d*x + 2*b^2 + 2*(a*b*cosh(d*x + c)^2 + 2*a*b*cosh(d*x + c)*sinh(d*x + c) + a*b*sinh(d*x + c)^2 + a*b)*log(2*cosh(d*x + c)/(cosh(d*x + c) - sinh(d*x + c)))/(d*cosh(d*x + c)^2 + 2*d*cosh(d*x + c)*sinh(d*x + c) + d*sinh(d*x + c)^2 + d)$

Sympy [A]

time = 0.07, size = 54, normalized size = 1.42

$$\begin{cases} a^2x + 2abx - \frac{2ab \log(\tanh(c+dx)+1)}{d} + b^2x - \frac{b^2 \tanh(c+dx)}{d} & \text{for } d \neq 0 \\ x(a + b \tanh(c))^2 & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+b*tanh(d*x+c))**2,x)

[Out] Piecewise((a**2*x + 2*a*b*x - 2*a*b*log(tanh(c + d*x) + 1)/d + b**2*x - b**2*tanh(c + d*x)/d, Ne(d, 0)), (x*(a + b*tanh(c))**2, True))

Giac [A]

time = 0.41, size = 56, normalized size = 1.47

$$\frac{2ab \log(e^{(2dx+2c)} + 1) + (a^2 - 2ab + b^2)(dx + c) + \frac{2b^2}{e^{(2dx+2c)} + 1}}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+b*tanh(d*x+c))^2,x, algorithm="giac")

[Out] (2*a*b*log(e^(2*d*x + 2*c) + 1) + (a^2 - 2*a*b + b^2)*(d*x + c) + 2*b^2/(e^(2*d*x + 2*c) + 1))/d

Mupad [B]

time = 1.04, size = 44, normalized size = 1.16

$$x(a^2 + 2ab + b^2) - \frac{b^2 \tanh(c + dx)}{d} - \frac{2ab \ln(\tanh(c + dx) + 1)}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*tanh(c + d*x))^2,x)

[Out] x*(2*a*b + a^2 + b^2) - (b^2*tanh(c + d*x))/d - (2*a*b*log(tanh(c + d*x) + 1))/d

3.61 $\int \frac{1}{a+b \tanh(c+dx)} dx$

Optimal. Leaf size=50

$$\frac{ax}{a^2 - b^2} - \frac{b \log(a \cosh(c + dx) + b \sinh(c + dx))}{(a^2 - b^2) d}$$

[Out] $a*x/(a^2-b^2)-b*\ln(a*\cosh(d*x+c)+b*\sinh(d*x+c))/(a^2-b^2)/d$

Rubi [A]

time = 0.04, antiderivative size = 50, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3565, 3611}

$$\frac{ax}{a^2 - b^2} - \frac{b \log(a \cosh(c + dx) + b \sinh(c + dx))}{d(a^2 - b^2)}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(a + b*\text{Tanh}[c + d*x])^{-1}, x]$

[Out] $(a*x)/(a^2 - b^2) - (b*\text{Log}[a*\text{Cosh}[c + d*x] + b*\text{Sinh}[c + d*x]])/((a^2 - b^2)*d)$

Rule 3565

$\text{Int}[(a + (b_*)*\text{tan}[(c_*) + (d_*)*(x_)])^{-1}, x_Symbol] \rightarrow \text{Simp}[a*(x/(a^2 + b^2)), x] + \text{Dist}[b/(a^2 + b^2), \text{Int}[(b - a*\text{Tan}[c + d*x])/(a + b*\text{Tan}[c + d*x]), x], x] /;$ $\text{FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{NeQ}[a^2 + b^2, 0]$

Rule 3611

$\text{Int}[(c + (d_*)*\text{tan}[(e_*) + (f_*)*(x_)])/(a + (b_*)*\text{tan}[(e_*) + (f_*)*(x_)]), x_Symbol] \rightarrow \text{Simp}[(c/(b*f))*\text{Log}[\text{RemoveContent}[a*\text{Cos}[e + f*x] + b*\text{Sin}[e + f*x], x]], x] /;$ $\text{FreeQ}\{a, b, c, d, e, f\}, x \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{EqQ}[a*c + b*d, 0]$

Rubi steps

$$\begin{aligned} \int \frac{1}{a + b \tanh(c + dx)} dx &= \frac{ax}{a^2 - b^2} - \frac{(ib) \int \frac{-ib - ia \tanh(c+dx)}{a + b \tanh(c+dx)} dx}{a^2 - b^2} \\ &= \frac{ax}{a^2 - b^2} - \frac{b \log(a \cosh(c + dx) + b \sinh(c + dx))}{(a^2 - b^2) d} \end{aligned}$$

Mathematica [A]

time = 0.06, size = 64, normalized size = 1.28

$$\frac{(-a + b) \log(1 - \tanh(c + dx)) + (a + b) \log(1 + \tanh(c + dx)) - 2b \log(a + b \tanh(c + dx))}{2(a - b)(a + b)d}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*Tanh[c + d*x])^(-1),x]

[Out] ((-a + b)*Log[1 - Tanh[c + d*x]] + (a + b)*Log[1 + Tanh[c + d*x]] - 2*b*Log[a + b*Tanh[c + d*x]])/(2*(a - b)*(a + b)*d)

Maple [A]

time = 0.85, size = 71, normalized size = 1.42

method	result	size
derivativedivides	$\frac{\frac{\ln(\tanh(dx+c)+1)}{2a-2b} - \frac{\ln(\tanh(dx+c)-1)}{2b+2a} - \frac{b \ln(a+b \tanh(dx+c))}{(a-b)(a+b)}}{d}$	71
default	$\frac{\frac{\ln(\tanh(dx+c)+1)}{2a-2b} - \frac{\ln(\tanh(dx+c)-1)}{2b+2a} - \frac{b \ln(a+b \tanh(dx+c))}{(a-b)(a+b)}}{d}$	71
risch	$\frac{x}{a+b} + \frac{2bx}{a^2-b^2} + \frac{2bc}{d(a^2-b^2)} - \frac{b \ln\left(e^{2dx+2c} + \frac{a-b}{a+b}\right)}{d(a^2-b^2)}$	81

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a+b*tanh(d*x+c)),x,method=_RETURNVERBOSE)

[Out] 1/d*(1/(2*a-2*b)*ln(tanh(d*x+c)+1)-1/(2*b+2*a)*ln(tanh(d*x+c)-1)-b/(a-b)/(a+b)*ln(a+b*tanh(d*x+c)))

Maxima [A]

time = 0.27, size = 56, normalized size = 1.12

$$-\frac{b \log\left(-\frac{(a-b)e^{(-2dx-2c)} - a - b}{(a^2 - b^2)d}\right) + \frac{dx + c}{(a + b)d}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+b*tanh(d*x+c)),x, algorithm="maxima")

[Out] -b*log(-(a - b)*e^(-2*d*x - 2*c) - a - b)/((a^2 - b^2)*d) + (d*x + c)/((a + b)*d)

Fricas [A]

time = 0.41, size = 62, normalized size = 1.24

$$\frac{(a + b)dx - b \log\left(\frac{2(a \cosh(dx+c) + b \sinh(dx+c))}{\cosh(dx+c) - \sinh(dx+c)}\right)}{(a^2 - b^2)d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(a+b*tanh(d*x+c)),x, algorithm="fricas")`

[Out] $((a + b)*d*x - b*\log(2*(a*\cosh(d*x + c) + b*\sinh(d*x + c))/(\cosh(d*x + c) - \sinh(d*x + c))))/((a^2 - b^2)*d)$

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 224 vs. 2(37) = 74.

time = 0.72, size = 224, normalized size = 4.48

$$\left\{ \begin{array}{ll} \frac{\infty x}{\tanh(c)} & \text{for } a = 0 \wedge b = 0 \wedge d = 0 \\ -\frac{dx \tanh(c+dx)}{2bd \tanh(c+dx)-2bd} + \frac{dx}{2bd \tanh(c+dx)-2bd} + \frac{1}{2bd \tanh(c+dx)-2bd} & \text{for } a = -b \\ \frac{dx \tanh(c+dx)}{2bd \tanh(c+dx)+2bd} + \frac{dx}{2bd \tanh(c+dx)+2bd} - \frac{1}{2bd \tanh(c+dx)+2bd} & \text{for } a = b \\ \frac{x}{a+b \tanh(c)} & \text{for } d = 0 \\ \frac{x}{a} & \text{for } b = 0 \\ \frac{adx}{a^2d-b^2d} - \frac{bdx}{a^2d-b^2d} - \frac{b \log\left(\frac{a}{b} + \tanh(c+dx)\right)}{a^2d-b^2d} + \frac{b \log(\tanh(c+dx)+1)}{a^2d-b^2d} & \text{otherwise} \end{array} \right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(a+b*tanh(d*x+c)),x)`

[Out] `Piecewise((zoo*x/tanh(c), Eq(a, 0) & Eq(b, 0) & Eq(d, 0)), (-d*x*tanh(c + d*x)/(2*b*d*tanh(c + d*x) - 2*b*d) + d*x/(2*b*d*tanh(c + d*x) - 2*b*d) + 1/(2*b*d*tanh(c + d*x) - 2*b*d), Eq(a, -b)), (d*x*tanh(c + d*x)/(2*b*d*tanh(c + d*x) + 2*b*d) + d*x/(2*b*d*tanh(c + d*x) + 2*b*d) - 1/(2*b*d*tanh(c + d*x) + 2*b*d), Eq(a, b)), (x/(a + b*tanh(c)), Eq(d, 0)), (x/a, Eq(b, 0)), (a*d*x/(a**2*d - b**2*d) - b*d*x/(a**2*d - b**2*d) - b*log(a/b + tanh(c + d*x))/(a**2*d - b**2*d) + b*log(tanh(c + d*x) + 1)/(a**2*d - b**2*d), True))`

Giac [A]

time = 0.43, size = 62, normalized size = 1.24

$$-\frac{\frac{b \log(|ae^{(2dx+2c)}+be^{(2dx+2c)}+a-b|)}{a^2-b^2} - \frac{dx+c}{a-b}}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(a+b*tanh(d*x+c)),x, algorithm="giac")`

[Out] $-(b*\log(\text{abs}(a*e^{(2*d*x + 2*c)} + b*e^{(2*d*x + 2*c)} + a - b)))/(a^2 - b^2) - (d*x + c)/(a - b)/d$

Mupad [B]

time = 1.14, size = 60, normalized size = 1.20

$$\frac{ax - bx}{a^2 - b^2} + \frac{b(\ln(\tanh(c + dx) + 1) - \ln(a + b \tanh(c + dx)))}{d(a^2 - b^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(a + b*tanh(c + d*x)),x)
```

```
[Out] (a*x - b*x)/(a^2 - b^2) + (b*(log(tanh(c + d*x) + 1) - log(a + b*tanh(c + d*x))))/(d*(a^2 - b^2))
```

$$3.62 \quad \int \frac{1}{(a+b \tanh(c+dx))^2} dx$$

Optimal. Leaf size=85

$$\frac{(a^2 + b^2)x}{(a^2 - b^2)^2} - \frac{2ab \log(a \cosh(c + dx) + b \sinh(c + dx))}{(a^2 - b^2)^2 d} + \frac{b}{(a^2 - b^2)d(a + b \tanh(c + dx))}$$

[Out] (a^2+b^2)*x/(a^2-b^2)^2-2*a*b*ln(a*cosh(d*x+c)+b*sinh(d*x+c))/(a^2-b^2)^2/d+b/(a^2-b^2)/d/(a+b*tanh(d*x+c))

Rubi [A]

time = 0.07, antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$,

Rules used = {3564, 3612, 3611}

$$\frac{b}{d(a^2 - b^2)(a + b \tanh(c + dx))} - \frac{2ab \log(a \cosh(c + dx) + b \sinh(c + dx))}{d(a^2 - b^2)^2} + \frac{x(a^2 + b^2)}{(a^2 - b^2)^2}$$

Antiderivative was successfully verified.

[In] Int[(a + b*Tanh[c + d*x])^(-2),x]

[Out] ((a^2 + b^2)*x)/(a^2 - b^2)^2 - (2*a*b*Log[a*Cosh[c + d*x] + b*Sinh[c + d*x]])/((a^2 - b^2)^2*d) + b/((a^2 - b^2)*d*(a + b*Tanh[c + d*x]))

Rule 3564

Int[((a_) + (b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[b*((a + b*Tan[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 + b^2))), x] + Dist[1/(a^2 + b^2), Int[(a - b*Tan[c + d*x])*(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0] && LtQ[n, -1]

Rule 3611

Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] :> Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]

Rule 3612

Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] :> Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Dist[(b*c - a*d)/(a^2 + b^2), Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{(a + b \tanh(c + dx))^2} dx &= \frac{b}{(a^2 - b^2) d(a + b \tanh(c + dx))} + \frac{\int \frac{a - b \tanh(c + dx)}{a + b \tanh(c + dx)} dx}{a^2 - b^2} \\ &= \frac{(a^2 + b^2) x}{(a^2 - b^2)^2} + \frac{b}{(a^2 - b^2) d(a + b \tanh(c + dx))} - \frac{(2iab) \int \frac{-ib - ia \tanh(c + dx)}{a + b \tanh(c + dx)} dx}{(a^2 - b^2)^2} \\ &= \frac{(a^2 + b^2) x}{(a^2 - b^2)^2} - \frac{2ab \log(a \cosh(c + dx) + b \sinh(c + dx))}{(a^2 - b^2)^2 d} + \frac{b}{(a^2 - b^2) d(a + b \tanh(c + dx))} \end{aligned}$$

Mathematica [A]

time = 0.83, size = 94, normalized size = 1.11

$$\frac{-\frac{\log(1 - \tanh(c + dx))}{(a + b)^2} + \frac{\log(1 + \tanh(c + dx))}{(a - b)^2} + \frac{2b(-2a \log(a + b \tanh(c + dx)) + \frac{a^2 - b^2}{a + b \tanh(c + dx)})}{(a^2 - b^2)^2}}{2d}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*Tanh[c + d*x])^(-2), x]

[Out] (-Log[1 - Tanh[c + d*x]]/(a + b)^2) + Log[1 + Tanh[c + d*x]]/(a - b)^2 + (2*b*(-2*a*Log[a + b*Tanh[c + d*x]] + (a^2 - b^2)/(a + b*Tanh[c + d*x])))/(a^2 - b^2)^2/(2*d)

Maple [A]

time = 0.76, size = 93, normalized size = 1.09

method	result
derivativedivides	$\frac{\frac{\ln(\tanh(dx+c)+1)}{2(a-b)^2} - \frac{\ln(\tanh(dx+c)-1)}{2(a+b)^2} + \frac{b}{(a-b)(a+b)(a+b \tanh(dx+c))} - \frac{2ab \ln(a+b \tanh(dx+c))}{(a+b)^2(a-b)^2}}{d}$
default	$\frac{\frac{\ln(\tanh(dx+c)+1)}{2(a-b)^2} - \frac{\ln(\tanh(dx+c)-1)}{2(a+b)^2} + \frac{b}{(a-b)(a+b)(a+b \tanh(dx+c))} - \frac{2ab \ln(a+b \tanh(dx+c))}{(a+b)^2(a-b)^2}}{d}$
risch	$\frac{x}{a^2 + 2ab + b^2} + \frac{4abx}{a^4 - 2a^2b^2 + b^4} + \frac{4abc}{d(a^4 - 2a^2b^2 + b^4)} + \frac{2b^2}{(a-b)d(a^2 + 2ab + b^2)(ae^{2dx+2c} + be^{2dx+2c} + a - b)} - \frac{2ab \ln(e^{2a}}{d(a^4 - 2a^2b^2 + b^4)}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a+b*tanh(d*x+c))^2,x,method=_RETURNVERBOSE)

[Out] 1/d*(1/2/(a-b)^2*ln(tanh(d*x+c)+1)-1/2/(a+b)^2*ln(tanh(d*x+c)-1)+b/(a-b)/(a+b)/(a+b*tanh(d*x+c))-2*a*b/(a+b)^2/(a-b)^2*ln(a+b*tanh(d*x+c)))

Maxima [A]

time = 0.27, size = 127, normalized size = 1.49

$$-\frac{2ab \log(-(a-b)e^{-2dx-2c}) - a - b}{(a^4 - 2a^2b^2 + b^4)d} - \frac{2b^2}{(a^4 - 2a^2b^2 + b^4 + (a^4 - 2a^3b + 2ab^3 - b^4)e^{-2dx-2c})d} + \frac{dx + c}{(a^2 + 2ab + b^2)d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+b*tanh(d*x+c))^2,x, algorithm="maxima")

[Out] $-2*a*b*\log(-(a - b)*e^{(-2*d*x - 2*c)} - a - b)/((a^4 - 2*a^2*b^2 + b^4)*d) - 2*b^2/((a^4 - 2*a^2*b^2 + b^4 + (a^4 - 2*a^3*b + 2*a*b^3 - b^4)*e^{(-2*d*x - 2*c)})*d) + (d*x + c)/((a^2 + 2*a*b + b^2)*d)$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 422 vs. 2(85) = 170.

time = 0.61, size = 422, normalized size = 4.96

$\frac{(a^3 + 3a^2b + 3ab^2 + b^3)dx \cosh(dx + c)^2 + 2(a^3 + 3a^2b + 3ab^2 + b^3)dx \sinh(dx + c) + (a^3 + 3a^2b + 3ab^2 + b^3)dx \sinh(dx + c)^2 + 2a^2b^2 + (a^2 + a^2b - ab^2 - b^2)dx - 2(a^2b - ab^2 + (a^2b + ab^2) \cosh(dx + c)^2 + 2(a^2b + ab^2) \cosh(dx + c) \sinh(dx + c) + (a^2b + ab^2) \sinh(dx + c)^2) \log\left(\frac{2(\cosh(dx + c) - \sinh(dx + c))}{\cosh(dx + c) + \sinh(dx + c)}\right)}{(a^2 + a^2b - 2a^2b^2 - 2a^2b^2 + ab^2 + b^2)dx \cosh(dx + c)^2 + 2(a^2 + a^2b - 2a^2b^2 - 2a^2b^2 + ab^2 + b^2)dx \sinh(dx + c) + (a^2 + a^2b - 2a^2b^2 - 2a^2b^2 + ab^2 + b^2)dx \sinh(dx + c)^2 + (a^2 - a^2b - 2a^2b^2 + 2a^2b^2 + ab^2 - b^2)d}$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+b*tanh(d*x+c))^2,x, algorithm="fricas")

[Out] $((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*d*x*cosh(d*x + c)^2 + 2*(a^3 + 3*a^2*b + 3*a*b^2 + b^3)*d*x*cosh(d*x + c)*sinh(d*x + c) + (a^3 + 3*a^2*b + 3*a*b^2 + b^3)*d*x*sinh(d*x + c)^2 + 2*a*b^2 - 2*b^3 + (a^3 + a^2*b - a*b^2 - b^3)*d*x - 2*(a^2*b - a*b^2 + (a^2*b + a*b^2)*cosh(d*x + c)^2 + 2*(a^2*b + a*b^2)*cosh(d*x + c)*sinh(d*x + c) + (a^2*b + a*b^2)*sinh(d*x + c)^2)*log(2*(a*cosh(d*x + c) + b*sinh(d*x + c))/(cosh(d*x + c) - sinh(d*x + c)))/((a^5 + a^4*b - 2*a^3*b^2 - 2*a^2*b^3 + a*b^4 + b^5)*d*cosh(d*x + c)^2 + 2*(a^5 + a^4*b - 2*a^3*b^2 - 2*a^2*b^3 + a*b^4 + b^5)*d*cosh(d*x + c)*sinh(d*x + c) + (a^5 + a^4*b - 2*a^3*b^2 - 2*a^2*b^3 + a*b^4 + b^5)*d*sinh(d*x + c)^2 + (a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*d)$

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 3597 vs. 2(70) = 140.

time = 14.10, size = 3597, normalized size = 42.32

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+b*tanh(d*x+c))**2,x)

[Out] $\text{Piecewise}((zoo*x/\tanh(c)**2, \text{Eq}(a, 0) \& \text{Eq}(b, 0) \& \text{Eq}(d, 0)), (d*x*tanh(c + d*x)**2/(4*b**2*d*tanh(c + d*x)**2 - 8*b**2*d*tanh(c + d*x) + 4*b**2*d) - 2*d*x*tanh(c + d*x)/(4*b**2*d*tanh(c + d*x)**2 - 8*b**2*d*tanh(c + d*x) + 4*b**2*d) + d*x/(4*b**2*d*tanh(c + d*x)**2 - 8*b**2*d*tanh(c + d*x) + 4*b**2*d) - tanh(c + d*x)/(4*b**2*d*tanh(c + d*x)**2 - 8*b**2*d*tanh(c + d*x) + 4*b**2*d) + 2/(4*b**2*d*tanh(c + d*x)**2 - 8*b**2*d*tanh(c + d*x) + 4*b**2*d)), \text{Eq}(a, -b)), (d*x*tanh(c + d*x)**2/(4*b**2*d*tanh(c + d*x)**2 + 8*b**2*d*tanh(c + d*x) + 4*b**2*d) + 2*d*x*tanh(c + d*x)/(4*b**2*d*tanh(c + d*x)**2 + 8*b**2*d*tanh(c + d*x) + 4*b**2*d) + d*x/(4*b**2*d*tanh(c + d*x)**2 + 8*b$

$$\begin{aligned}
& **2*d*\tanh(c + d*x) + 4*b**2*d) - \tanh(c + d*x)/(4*b**2*d*\tanh(c + d*x)**2 \\
& + 8*b**2*d*\tanh(c + d*x) + 4*b**2*d) - 2/(4*b**2*d*\tanh(c + d*x)**2 + 8*b** \\
& 2*d*\tanh(c + d*x) + 4*b**2*d), \text{Eq}(a, b)), (d*x*\exp(4*c)*\exp(4*d*x)*\tanh(c + \\
& d*x)**2/(4*b**2*d*\exp(4*c)*\exp(4*d*x)*\tanh(c + d*x)**2 - 8*b**2*d*\exp(4*c) \\
& *\exp(4*d*x)*\tanh(c + d*x) + 4*b**2*d*\exp(4*c)*\exp(4*d*x) + 8*b**2*d*\exp(2*c) \\
&)*\exp(2*d*x)*\tanh(c + d*x)**2 - 8*b**2*d*\exp(2*c)*\exp(2*d*x) + 4*b**2*d*\tan \\
& h(c + d*x)**2 + 8*b**2*d*\tanh(c + d*x) + 4*b**2*d) - 2*d*x*\exp(4*c)*\exp(4*d \\
& *x)*\tanh(c + d*x)/(4*b**2*d*\exp(4*c)*\exp(4*d*x)*\tanh(c + d*x)**2 - 8*b**2*d \\
& *\exp(4*c)*\exp(4*d*x)*\tanh(c + d*x) + 4*b**2*d*\exp(4*c)*\exp(4*d*x) + 8*b**2*d \\
& *\exp(2*c)*\exp(2*d*x)*\tanh(c + d*x)**2 - 8*b**2*d*\exp(2*c)*\exp(2*d*x) + 4*b \\
& **2*d*\tanh(c + d*x)**2 + 8*b**2*d*\tanh(c + d*x) + 4*b**2*d) + d*x*\exp(4*c)* \\
& \exp(4*d*x)/(4*b**2*d*\exp(4*c)*\exp(4*d*x)*\tanh(c + d*x)**2 - 8*b**2*d*\exp(4*c) \\
& *\exp(4*d*x)*\tanh(c + d*x) + 4*b**2*d*\exp(4*c)*\exp(4*d*x) + 8*b**2*d*\exp(2 \\
& *c)*\exp(2*d*x)*\tanh(c + d*x)**2 - 8*b**2*d*\exp(2*c)*\exp(2*d*x) + 4*b**2*d*\t \\
& anh(c + d*x)**2 + 8*b**2*d*\tanh(c + d*x) + 4*b**2*d) - 4*d*x*\exp(2*c)*\exp(2 \\
& *d*x)*\tanh(c + d*x)**2/(4*b**2*d*\exp(4*c)*\exp(4*d*x)*\tanh(c + d*x)**2 - 8*b \\
& **2*d*\exp(4*c)*\exp(4*d*x)*\tanh(c + d*x) + 4*b**2*d*\exp(4*c)*\exp(4*d*x) + 8 \\
& *b**2*d*\exp(2*c)*\exp(2*d*x)*\tanh(c + d*x)**2 - 8*b**2*d*\exp(2*c)*\exp(2*d*x) \\
& + 4*b**2*d*\tanh(c + d*x)**2 + 8*b**2*d*\tanh(c + d*x) + 4*b**2*d) + 4*d*x*\exp \\
& p(2*c)*\exp(2*d*x)/(4*b**2*d*\exp(4*c)*\exp(4*d*x)*\tanh(c + d*x)**2 - 8*b**2*d \\
& *\exp(4*c)*\exp(4*d*x)*\tanh(c + d*x) + 4*b**2*d*\exp(4*c)*\exp(4*d*x) + 8*b**2*d \\
& *\exp(2*c)*\exp(2*d*x)*\tanh(c + d*x)**2 - 8*b**2*d*\exp(2*c)*\exp(2*d*x) + 4*b \\
& **2*d*\tanh(c + d*x)**2 + 8*b**2*d*\tanh(c + d*x) + 4*b**2*d) + d*x*\tanh(c + \\
& d*x)**2/(4*b**2*d*\exp(4*c)*\exp(4*d*x)*\tanh(c + d*x)**2 - 8*b**2*d*\exp(4*c)* \\
& \exp(4*d*x)*\tanh(c + d*x) + 4*b**2*d*\exp(4*c)*\exp(4*d*x) + 8*b**2*d*\exp(2*c) \\
& *\exp(2*d*x)*\tanh(c + d*x)**2 - 8*b**2*d*\exp(2*c)*\exp(2*d*x) + 4*b**2*d*\tanh \\
& (c + d*x)**2 + 8*b**2*d*\tanh(c + d*x) + 4*b**2*d) + 2*d*x*\tanh(c + d*x)/(4* \\
& b**2*d*\exp(4*c)*\exp(4*d*x)*\tanh(c + d*x)**2 - 8*b**2*d*\exp(4*c)*\exp(4*d*x)* \\
& \tanh(c + d*x) + 4*b**2*d*\exp(4*c)*\exp(4*d*x) + 8*b**2*d*\exp(2*c)*\exp(2*d*x) \\
& *\tanh(c + d*x)**2 - 8*b**2*d*\exp(2*c)*\exp(2*d*x) + 4*b**2*d*\tanh(c + d*x)** \\
& 2 + 8*b**2*d*\tanh(c + d*x) + 4*b**2*d) + d*x/(4*b**2*d*\exp(4*c)*\exp(4*d*x)* \\
& \tanh(c + d*x)**2 - 8*b**2*d*\exp(4*c)*\exp(4*d*x)*\tanh(c + d*x) + 4*b**2*d*\exp \\
& p(4*c)*\exp(4*d*x) + 8*b**2*d*\exp(2*c)*\exp(2*d*x)*\tanh(c + d*x)**2 - 8*b**2*d \\
& *\exp(2*c)*\exp(2*d*x) + 4*b**2*d*\tanh(c + d*x)**2 + 8*b**2*d*\tanh(c + d*x) \\
& + 4*b**2*d) - \exp(4*c)*\exp(4*d*x)*\tanh(c + d*x)/(4*b**2*d*\exp(4*c)*\exp(4*d* \\
& x)*\tanh(c + d*x)**2 - 8*b**2*d*\exp(4*c)*\exp(4*d*x)*\tanh(c + d*x) + 4*b**2*d \\
& *\exp(4*c)*\exp(4*d*x) + 8*b**2*d*\exp(2*c)*\exp(2*d*x)*\tanh(c + d*x)**2 - 8*b* \\
& **2*d*\exp(2*c)*\exp(2*d*x) + 4*b**2*d*\tanh(c + d*x)**2 + 8*b**2*d*\tanh(c + d* \\
& x) + 4*b**2*d) + 2*\exp(4*c)*\exp(4*d*x)/(4*b**2*d*\exp(4*c)*\exp(4*d*x)*\tanh(c \\
& + d*x)**2 - 8*b**2*d*\exp(4*c)*\exp(4*d*x)*\tanh(c + d*x) + 4*b**2*d*\exp(4*c) \\
& *\exp(4*d*x) + 8*b**2*d*\exp(2*c)*\exp(2*d*x)*\tanh(c + d*x)**2 - 8*b**2*d*\exp(\\
& 2*c)*\exp(2*d*x) + 4*b**2*d*\tanh(c + d*x)**2 + 8*b**2*d*\tanh(c + d*x) + 4*b* \\
& **2*d) + 4*\exp(2*c)*\exp(2*d*x)*\tanh(c + d*x)/(4*b**2*d*\exp(4*c)*\exp(4*d*x)*\t \\
& anh(c + d*x)**2 - 8*b**2*d*\exp(4*c)*\exp(4*d*x)*\tanh(c + d*x) + 4*b**2*d*\exp \\
& (4*c)*\exp(4*d*x) + 8*b**2*d*\exp(2*c)*\exp(2*d*x)*\tanh(c + d*x)**2 - 8*b**2*d
\end{aligned}$$

*exp(2*c)*exp(2*d*x) + 4*b**2*d*tanh(c + d*x)**2 + 8*b**2*d*tanh(c + d*x) + 4*b**2*d) - tanh(c + d*x)/(4*b**2*d*exp(4*c)*exp(4*d*x)*tanh(c + d*x)**2 - 8*b**2*d*exp(4*c)*exp(4*d*x)*tanh(c + d*x) + 4*b**2*d*exp(4*c)*exp(4*d*x) + 8*b**2*d*exp(2*c)*exp(2*d*x)*tanh(c + d*x)**2 - 8*b**2*d*exp(2*c)*exp(2*d*x) + 4*b**2*d*tanh(c + d*x)**2 + 8*b**2*d*tanh(c + d*x) + 4*b**2*d) - 2/(4*b**2*d*exp(4*c)*exp(4*d*x)*tanh(c + d*x)**2 - 8*b**2*d*exp(4*c)*exp(4*d*x)*tanh(c + d*x) + 4*b**2*d*exp(4*c)*exp(4*d*x) + 8*b**2*d*exp(2*c)*exp(2*d*x)*tanh(c + d*x)**2 - 8*b**2*d*exp(2*c)*exp(2*d*x) + 4*b**2*d*tanh(c + d*x)**2 + 8*b**2*d*tanh(c + d*x) + 4*b**2*d), Eq(a, (-b*exp(2*c)*exp(2*d*x) + b)/(exp(2*c)*exp(2*d*x) + 1)), (x/(a + b*tanh(c))**2, Eq(d, 0)), (x/a**2, Eq(b, 0)), (a**3*d*x/(a**5*d + a**4*b*d*tanh(c + d*x) - 2*a**3*b**2*d - 2*a**2*b**3*d*tanh(c + d*x) + a*b**4*d + b**5*d*tanh(c + d*x)) + a**2*b*d*x*tanh(c + d*x)/(a**5*d + a**4*b*d*tanh(c + d*x) - 2*...

Giac [A]

time = 0.43, size = 132, normalized size = 1.55

$$\frac{2ab \log\left(\frac{-ae^{(2dx+2c)} - be^{(2dx+2c)} - a + b}{a^4 - 2a^2b^2 + b^4}\right) - \frac{dx+c}{a^2 - 2ab + b^2} - \frac{2(ab^2 - b^3)}{(ae^{(2dx+2c)} + be^{(2dx+2c)} + a - b)(a+b)^2(a-b)^2}}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+b*tanh(d*x+c))^2,x, algorithm="giac")

[Out] -(2*a*b*log(abs(-a*e^(2*d*x + 2*c) - b*e^(2*d*x + 2*c) - a + b))/(a^4 - 2*a^2*b^2 + b^4) - (d*x + c)/(a^2 - 2*a*b + b^2) - 2*(a*b^2 - b^3)/((a*e^(2*d*x + 2*c) + b*e^(2*d*x + 2*c) + a - b)*(a + b)^2*(a - b)^2))/d

Mupad [B]

time = 1.43, size = 127, normalized size = 1.49

$$\frac{\frac{ax}{(a+b)^2} + \frac{bx \tanh(c+dx)}{(a+b)^2} - \frac{b^2 \tanh(c+dx)}{ad(a^2-b^2)}}{a + b \tanh(c + dx)} - \frac{2ab \ln(a + b \tanh(c + dx))}{d(a^4 - 2a^2b^2 + b^4)} + \frac{2ab \ln(\tanh(c + dx) + 1)}{d(a^2 - b^2)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a + b*tanh(c + d*x))^2,x)

[Out] ((a*x)/(a + b)^2 + (b*x*tanh(c + d*x))/(a + b)^2 - (b^2*tanh(c + d*x))/(a*d*(a^2 - b^2)))/(a + b*tanh(c + d*x)) - (2*a*b*log(a + b*tanh(c + d*x)))/(d*(a^4 + b^4 - 2*a^2*b^2)) + (2*a*b*log(tanh(c + d*x) + 1))/(d*(a^2 - b^2)^2)

3.63 $\int \frac{1}{(a+b \tanh(c+dx))^3} dx$

Optimal. Leaf size=129

$$\frac{a(a^2 + 3b^2)x}{(a^2 - b^2)^3} - \frac{b(3a^2 + b^2) \log(a \cosh(c + dx) + b \sinh(c + dx))}{(a^2 - b^2)^3 d} + \frac{b}{2(a^2 - b^2)d(a + b \tanh(c + dx))^2} + \frac{1}{(a^2 - b^2)}$$

[Out] $a*(a^2+3*b^2)*x/(a^2-b^2)^3-b*(3*a^2+b^2)*\ln(a*\cosh(d*x+c)+b*\sinh(d*x+c))/(a^2-b^2)^3/d+1/2*b/(a^2-b^2)/d/(a+b*\tanh(d*x+c))^2+2*a*b/(a^2-b^2)^2/d/(a+b*\tanh(d*x+c))$

Rubi [A]

time = 0.13, antiderivative size = 129, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3564, 3610, 3612, 3611}

$$\frac{2ab}{d(a^2 - b^2)^2(a + b \tanh(c + dx))} + \frac{b}{2d(a^2 - b^2)(a + b \tanh(c + dx))^2} - \frac{b(3a^2 + b^2) \log(a \cosh(c + dx) + b \sinh(c + dx))}{d(a^2 - b^2)^3} + \frac{ax(a^2 + 3b^2)}{(a^2 - b^2)^3}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(a + b*\text{Tanh}[c + d*x])^{-3}, x]$

[Out] $(a*(a^2 + 3*b^2)*x)/(a^2 - b^2)^3 - (b*(3*a^2 + b^2)*\text{Log}[a*\text{Cosh}[c + d*x] + b*\text{Sinh}[c + d*x]])/((a^2 - b^2)^3*d) + b/(2*(a^2 - b^2)*d*(a + b*\text{Tanh}[c + d*x])^2) + (2*a*b)/((a^2 - b^2)^2*d*(a + b*\text{Tanh}[c + d*x]))$

Rule 3564

$\text{Int}[(a + b*\text{Tan}[c + d*x])^n, x] \text{ :> } \text{Simp}[b*(a + b*\text{Tan}[c + d*x])^{n+1}/(d*(n+1)*(a^2 + b^2)), x] + \text{Dist}[1/(a^2 + b^2), \text{Int}[(a - b*\text{Tan}[c + d*x])*(a + b*\text{Tan}[c + d*x])^{n+1}, x], x] /;$ FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0] && LtQ[n, -1]

Rule 3610

$\text{Int}[(a + b*\text{Tan}[e + f*x])^m, x] \text{ :> } \text{Simp}[(b*c - a*d)*(a + b*\text{Tan}[e + f*x])^{m+1}/(f*(m+1)*(a^2 + b^2)), x] + \text{Dist}[1/(a^2 + b^2), \text{Int}[(a + b*\text{Tan}[e + f*x])^{m+1}*\text{Simp}[a*c + b*d - (b*c - a*d)*\text{Tan}[e + f*x], x], x] /;$ FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]

Rule 3611

$\text{Int}[(c + d*\text{Tan}[e + f*x])/(a + b*\text{Tan}[e + f*x]), x] \text{ :> } \text{Simp}[(c/(b*f))*\text{Log}[\text{RemoveContent}[a*\text{Cos}[e + f*x] + b*\text{Sin}[e + f*x], x]], x] /;$ FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] &&

NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]

Rule 3612

Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Dist[(b*c - a*d)/(a^2 + b^2), Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{(a + b \tanh(c + dx))^3} dx &= \frac{b}{2(a^2 - b^2) d(a + b \tanh(c + dx))^2} + \frac{\int \frac{a - b \tanh(c + dx)}{(a + b \tanh(c + dx))^2} dx}{a^2 - b^2} \\ &= \frac{b}{2(a^2 - b^2) d(a + b \tanh(c + dx))^2} + \frac{2ab}{(a^2 - b^2)^2 d(a + b \tanh(c + dx))} + \frac{\int \frac{a^2}{(a + b \tanh(c + dx))^2} dx}{2(a^2 - b^2)^2 d} \\ &= \frac{a(a^2 + 3b^2)x}{(a^2 - b^2)^3} + \frac{b}{2(a^2 - b^2) d(a + b \tanh(c + dx))^2} + \frac{2ab}{(a^2 - b^2)^2 d(a + b \tanh(c + dx))} \\ &= \frac{a(a^2 + 3b^2)x}{(a^2 - b^2)^3} - \frac{b(3a^2 + b^2) \log(a \cosh(c + dx) + b \sinh(c + dx))}{(a^2 - b^2)^3 d} + \frac{2ab}{2(a^2 - b^2)^2 d} \end{aligned}$$

Mathematica [A]

time = 1.69, size = 122, normalized size = 0.95

$$\frac{-\frac{\log(1 - \tanh(c + dx))}{(a + b)^3} + \frac{\log(1 + \tanh(c + dx))}{(a - b)^3} + \frac{b \left(-2(3a^2 + b^2) \log(a + b \tanh(c + dx)) + \frac{(a^2 - b^2)(5a^2 - b^2 + 4ab \tanh(c + dx))}{(a + b \tanh(c + dx))^2} \right)}{(a^2 - b^2)^3}}{2d}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*Tanh[c + d*x])^(-3), x]

[Out] (-(Log[1 - Tanh[c + d*x]]/(a + b)^3) + Log[1 + Tanh[c + d*x]]/(a - b)^3 + (b*(-2*(3*a^2 + b^2)*Log[a + b*Tanh[c + d*x]] + ((a^2 - b^2)*(5*a^2 - b^2 + 4*a*b*Tanh[c + d*x]))/(a + b*Tanh[c + d*x])^2))/(a^2 - b^2)^3)/(2*d)

Maple [A]

time = 0.64, size = 130, normalized size = 1.01

method	result
--------	--------

derivativedivides	$\frac{\frac{\ln(\tanh(dx+c)+1)}{2(a-b)^3} + \frac{b}{2(a-b)(a+b)(a+b \tanh(dx+c))^2} + \frac{2ab}{(a+b)^2(a-b)^2(a+b \tanh(dx+c))} - \frac{b(3a^2+b^2) \ln(a+b \tanh(dx+c))}{(a+b)^3(a-b)^3} - \frac{\ln(\tanh(dx+c))}{2(a+b)}}{d}$
default	$\frac{\frac{\ln(\tanh(dx+c)+1)}{2(a-b)^3} + \frac{b}{2(a-b)(a+b)(a+b \tanh(dx+c))^2} + \frac{2ab}{(a+b)^2(a-b)^2(a+b \tanh(dx+c))} - \frac{b(3a^2+b^2) \ln(a+b \tanh(dx+c))}{(a+b)^3(a-b)^3} - \frac{\ln(\tanh(dx+c))}{2(a+b)}}{d}$
risch	$\frac{x}{a^3+3a^2b+3ab^2+b^3} + \frac{6ba^2x}{a^6-3a^4b^2+3a^2b^4-b^6} + \frac{2b^3x}{a^6-3a^4b^2+3a^2b^4-b^6} + \frac{6ba^2c}{d(a^6-3a^4b^2+3a^2b^4-b^6)} + \frac{2b^3c}{d(a^6-3a^4b^2+3a^2b^4-b^6)}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(a+b*tanh(d*x+c))^3,x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{d} * \left(\frac{1}{2} * \frac{1}{(a-b)^3} * \ln(\tanh(d*x+c)+1) + \frac{1}{2} * \frac{b}{(a-b)} * \frac{1}{(a+b)} * \frac{1}{(a+b \tanh(d*x+c))} \right)^{2+2} * \frac{a*b}{(a+b)^2} * \frac{1}{(a-b)^2} * \frac{1}{(a+b \tanh(d*x+c))} - b * \frac{(3a^2+b^2)}{(a+b)^3} * \frac{1}{(a-b)^3} * \ln(a+b \tanh(d*x+c)) - \frac{1}{2} * \frac{1}{(a+b)^3} * \ln(\tanh(d*x+c)-1) \right)$

Maxima [B] Leaf count of result is larger than twice the leaf count of optimal. 325 vs. 2(127) = 254.

time = 0.30, size = 325, normalized size = 2.52

$$\frac{(3a^2b+b^3) \log(-(a-b)e^{-2dx-2c}-a-b)}{(a^6-3a^4b^2+3a^2b^4-b^6)d} - \frac{2(3a^2b^2+3ab^3+(3a^2b-2ab^2-b^3)e^{-2dx-2c})}{(a^6-3a^4b^2+3a^2b^4-b^6)d} + \frac{dx+c}{(a^6-3a^4b^2+3a^2b^4-b^6)d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(a+b*tanh(d*x+c))^3,x, algorithm="maxima")`

[Out] $-(3a^2b + b^3) * \log(-(a - b) * e^{-2*d*x - 2*c} - a - b) / ((a^6 - 3a^4b^2 + 3a^2b^4 - b^6) * d) - 2 * (3a^2b^2 + 3a*b^3 + (3a^2b^2 - 2a*b^3 - b^4) * e^{-2*d*x - 2*c}) / ((a^7 + a^6b - 3a^5b^2 - 3a^4b^3 + 3a^3b^4 + 3a^2b^5 - a*b^6 - b^7 + 2*(a^7 - a^6b - 3a^5b^2 + 3a^4b^3 + 3a^3b^4 - 3a^2b^5 - a*b^6 + b^7) * e^{-2*d*x - 2*c} + (a^7 - 3a^6b + a^5b^2 + 5a^4b^3 - 5a^3b^4 - a^2b^5 + 3a*b^6 - b^7) * e^{-4*d*x - 4*c}) * d) + (d*x + c) / ((a^3 + 3a^2b + 3a*b^2 + b^3) * d)$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 1427 vs. 2(127) = 254.

time = 0.39, size = 1427, normalized size = 11.06

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(a+b*tanh(d*x+c))^3,x, algorithm="fricas")`

[Out] $((a^5 + 5a^4b + 10a^3b^2 + 10a^2b^3 + 5a*b^4 + b^5) * d*x * \cosh(d*x + c))^4 + 4 * (a^5 + 5a^4b + 10a^3b^2 + 10a^2b^3 + 5a*b^4 + b^5) * d*x * \cosh(d*x + c) * \sinh(d*x + c)^3 + (a^5 + 5a^4b + 10a^3b^2 + 10a^2b^3 + 5a*b^4 + b^5) * d*x * \cosh(d*x + c) * \sinh(d*x + c)^2 + (a^5 + 5a^4b + 10a^3b^2 + 10a^2b^3 + 5a*b^4 + b^5) * d*x * \cosh(d*x + c) * \sinh(d*x + c) + (a^5 + 5a^4b + 10a^3b^2 + 10a^2b^3 + 5a*b^4 + b^5) * d*x * \cosh(d*x + c) * \sinh(d*x + c)$

$$\begin{aligned}
&^4 + b^5) * d * x * \sinh(d * x + c)^4 + 6 * a^3 * b^2 - 12 * a^2 * b^3 + 6 * a * b^4 + (a^5 + a \\
&^4 * b - 2 * a^3 * b^2 - 2 * a^2 * b^3 + a * b^4 + b^5) * d * x + 2 * (3 * a^3 * b^2 - a^2 * b^3 - \\
&3 * a * b^4 + b^5 + (a^5 + 3 * a^4 * b + 2 * a^3 * b^2 - 2 * a^2 * b^3 - 3 * a * b^4 - b^5) * d * x \\
&)* \cosh(d * x + c)^2 + 2 * (3 * a^3 * b^2 - a^2 * b^3 - 3 * a * b^4 + b^5 + 3 * (a^5 + 5 * a^4 \\
&* b + 10 * a^3 * b^2 + 10 * a^2 * b^3 + 5 * a * b^4 + b^5) * d * x * \cosh(d * x + c)^2 + (a^5 + \\
&3 * a^4 * b + 2 * a^3 * b^2 - 2 * a^2 * b^3 - 3 * a * b^4 - b^5) * d * x) * \sinh(d * x + c)^2 - (3 * \\
&a^4 * b - 6 * a^3 * b^2 + 4 * a^2 * b^3 - 2 * a * b^4 + b^5 + (3 * a^4 * b + 6 * a^3 * b^2 + 4 * a^ \\
&2 * b^3 + 2 * a * b^4 + b^5) * \cosh(d * x + c)^4 + 4 * (3 * a^4 * b + 6 * a^3 * b^2 + 4 * a^2 * b^3 \\
&+ 2 * a * b^4 + b^5) * \cosh(d * x + c) * \sinh(d * x + c)^3 + (3 * a^4 * b + 6 * a^3 * b^2 + 4 * \\
&a^2 * b^3 + 2 * a * b^4 + b^5) * \sinh(d * x + c)^4 + 2 * (3 * a^4 * b - 2 * a^2 * b^3 - b^5) * \co \\
&sh(d * x + c)^2 + 2 * (3 * a^4 * b - 2 * a^2 * b^3 - b^5 + 3 * (3 * a^4 * b + 6 * a^3 * b^2 + 4 * a \\
&^2 * b^3 + 2 * a * b^4 + b^5) * \cosh(d * x + c)^2) * \sinh(d * x + c)^2 + 4 * ((3 * a^4 * b + 6 * \\
&a^3 * b^2 + 4 * a^2 * b^3 + 2 * a * b^4 + b^5) * \cosh(d * x + c)^3 + (3 * a^4 * b - 2 * a^2 * b^3 \\
&- b^5) * \cosh(d * x + c)) * \sinh(d * x + c)) * \log(2 * (a * \cosh(d * x + c) + b * \sinh(d * x + \\
&c)) / (\cosh(d * x + c) - \sinh(d * x + c))) + 4 * ((a^5 + 5 * a^4 * b + 10 * a^3 * b^2 + 10 \\
&* a^2 * b^3 + 5 * a * b^4 + b^5) * d * x * \cosh(d * x + c)^3 + (3 * a^3 * b^2 - a^2 * b^3 - 3 * a * \\
&b^4 + b^5 + (a^5 + 3 * a^4 * b + 2 * a^3 * b^2 - 2 * a^2 * b^3 - 3 * a * b^4 - b^5) * d * x) * \co \\
&sh(d * x + c)) * \sinh(d * x + c)) / ((a^8 + 2 * a^7 * b - 2 * a^6 * b^2 - 6 * a^5 * b^3 + 6 * a^3 \\
&* b^5 + 2 * a^2 * b^6 - 2 * a * b^7 - b^8) * d * \cosh(d * x + c)^4 + 4 * (a^8 + 2 * a^7 * b - 2 * \\
&a^6 * b^2 - 6 * a^5 * b^3 + 6 * a^3 * b^5 + 2 * a^2 * b^6 - 2 * a * b^7 - b^8) * d * \cosh(d * x + c \\
&)* \sinh(d * x + c)^3 + (a^8 + 2 * a^7 * b - 2 * a^6 * b^2 - 6 * a^5 * b^3 + 6 * a^3 * b^5 + 2 * \\
&a^2 * b^6 - 2 * a * b^7 - b^8) * d * \sinh(d * x + c)^4 + 2 * (a^8 - 4 * a^6 * b^2 + 6 * a^4 * b^4 \\
&- 4 * a^2 * b^6 + b^8) * d * \cosh(d * x + c)^2 + 2 * (3 * (a^8 + 2 * a^7 * b - 2 * a^6 * b^2 - 6 \\
&* a^5 * b^3 + 6 * a^3 * b^5 + 2 * a^2 * b^6 - 2 * a * b^7 - b^8) * d * \cosh(d * x + c)^2 + (a^8 \\
&- 4 * a^6 * b^2 + 6 * a^4 * b^4 - 4 * a^2 * b^6 + b^8) * d) * \sinh(d * x + c)^2 + (a^8 - 2 * a^ \\
&7 * b - 2 * a^6 * b^2 + 6 * a^5 * b^3 - 6 * a^3 * b^5 + 2 * a^2 * b^6 + 2 * a * b^7 - b^8) * d + 4 * \\
&((a^8 + 2 * a^7 * b - 2 * a^6 * b^2 - 6 * a^5 * b^3 + 6 * a^3 * b^5 + 2 * a^2 * b^6 - 2 * a * b^7 - \\
&b^8) * d * \cosh(d * x + c)^3 + (a^8 - 4 * a^6 * b^2 + 6 * a^4 * b^4 - 4 * a^2 * b^6 + b^8) * d \\
&* \cosh(d * x + c)) * \sinh(d * x + c))
\end{aligned}$$

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 16657 vs. $2(107) = 214$.

time = 82.40, size = 16657, normalized size = 129.12

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+b*tanh(d*x+c))**3,x)

[Out] Piecewise((zoo*x/tanh(c)**3, Eq(a, 0) & Eq(b, 0) & Eq(d, 0)), (x/a**3, Eq(b, 0)), (-3*d*x*tanh(c + d*x)**3/(24*b**3*d*tanh(c + d*x)**3 - 72*b**3*d*tanh(c + d*x)**2 + 72*b**3*d*tanh(c + d*x) - 24*b**3*d) + 9*d*x*tanh(c + d*x)**2/(24*b**3*d*tanh(c + d*x)**3 - 72*b**3*d*tanh(c + d*x)**2 + 72*b**3*d*tanh(c + d*x) - 24*b**3*d) - 9*d*x*tanh(c + d*x)/(24*b**3*d*tanh(c + d*x)**3 - 72*b**3*d*tanh(c + d*x)**2 + 72*b**3*d*tanh(c + d*x) - 24*b**3*d) + 3*d*x/

$$\begin{aligned}
& (24*b**3*d*tanh(c + d*x)**3 - 72*b**3*d*tanh(c + d*x)**2 + 72*b**3*d*tanh(c \\
& + d*x) - 24*b**3*d) + 3*tanh(c + d*x)**2/(24*b**3*d*tanh(c + d*x)**3 - 72* \\
& b**3*d*tanh(c + d*x)**2 + 72*b**3*d*tanh(c + d*x) - 24*b**3*d) - 9*tanh(c + \\
& d*x)/(24*b**3*d*tanh(c + d*x)**3 - 72*b**3*d*tanh(c + d*x)**2 + 72*b**3*d* \\
& tanh(c + d*x) - 24*b**3*d) + 10/(24*b**3*d*tanh(c + d*x)**3 - 72*b**3*d*tan \\
& h(c + d*x)**2 + 72*b**3*d*tanh(c + d*x) - 24*b**3*d), Eq(a, -b)), (3*d*x*tan \\
& h(c + d*x)**3/(24*b**3*d*tanh(c + d*x)**3 + 72*b**3*d*tanh(c + d*x)**2 + 7 \\
& 2*b**3*d*tanh(c + d*x) + 24*b**3*d) + 9*d*x*tanh(c + d*x)**2/(24*b**3*d*tan \\
& h(c + d*x)**3 + 72*b**3*d*tanh(c + d*x)**2 + 72*b**3*d*tanh(c + d*x) + 24*b \\
& **3*d) + 9*d*x*tanh(c + d*x)/(24*b**3*d*tanh(c + d*x)**3 + 72*b**3*d*tanh(c \\
& + d*x)**2 + 72*b**3*d*tanh(c + d*x) + 24*b**3*d) + 3*d*x/(24*b**3*d*tanh(c \\
& + d*x)**3 + 72*b**3*d*tanh(c + d*x)**2 + 72*b**3*d*tanh(c + d*x) + 24*b**3 \\
& *d) - 3*tanh(c + d*x)**2/(24*b**3*d*tanh(c + d*x)**3 + 72*b**3*d*tanh(c + d \\
& *x)**2 + 72*b**3*d*tanh(c + d*x) + 24*b**3*d) - 9*tanh(c + d*x)/(24*b**3*d* \\
& tanh(c + d*x)**3 + 72*b**3*d*tanh(c + d*x)**2 + 72*b**3*d*tanh(c + d*x) + 2 \\
& 4*b**3*d) - 10/(24*b**3*d*tanh(c + d*x)**3 + 72*b**3*d*tanh(c + d*x)**2 + 7 \\
& 2*b**3*d*tanh(c + d*x) + 24*b**3*d), Eq(a, b)), (-3*d*x*exp(6*c)*exp(6*d*x) \\
& *tanh(c + d*x)**3/(24*b**3*d*exp(6*c)*exp(6*d*x)*tanh(c + d*x)**3 - 72*b**3 \\
& *d*exp(6*c)*exp(6*d*x)*tanh(c + d*x)**2 + 72*b**3*d*exp(6*c)*exp(6*d*x)*tan \\
& h(c + d*x) - 24*b**3*d*exp(6*c)*exp(6*d*x) + 72*b**3*d*exp(4*c)*exp(4*d*x)* \\
& tanh(c + d*x)**3 - 72*b**3*d*exp(4*c)*exp(4*d*x)*tanh(c + d*x)**2 - 72*b**3 \\
& *d*exp(4*c)*exp(4*d*x)*tanh(c + d*x) + 72*b**3*d*exp(4*c)*exp(4*d*x) + 72*b \\
& **3*d*exp(2*c)*exp(2*d*x)*tanh(c + d*x)**3 + 72*b**3*d*exp(2*c)*exp(2*d*x)* \\
& tanh(c + d*x)**2 - 72*b**3*d*exp(2*c)*exp(2*d*x)*tanh(c + d*x) - 72*b**3*d* \\
& exp(2*c)*exp(2*d*x) + 24*b**3*d*tanh(c + d*x)**3 + 72*b**3*d*tanh(c + d*x)* \\
& *2 + 72*b**3*d*tanh(c + d*x) + 24*b**3*d) + 9*d*x*exp(6*c)*exp(6*d*x)*tanh(\\
& c + d*x)**2/(24*b**3*d*exp(6*c)*exp(6*d*x)*tanh(c + d*x)**3 - 72*b**3*d*exp \\
& (6*c)*exp(6*d*x)*tanh(c + d*x)**2 + 72*b**3*d*exp(6*c)*exp(6*d*x)*tanh(c + \\
& d*x) - 24*b**3*d*exp(6*c)*exp(6*d*x) + 72*b**3*d*exp(4*c)*exp(4*d*x)*tanh(c \\
& + d*x)**3 - 72*b**3*d*exp(4*c)*exp(4*d*x)*tanh(c + d*x)**2 - 72*b**3*d*exp \\
& (4*c)*exp(4*d*x)*tanh(c + d*x) + 72*b**3*d*exp(4*c)*exp(4*d*x) + 72*b**3*d* \\
& exp(2*c)*exp(2*d*x)*tanh(c + d*x)**3 + 72*b**3*d*exp(2*c)*exp(2*d*x)*tanh(c \\
& + d*x)**2 - 72*b**3*d*exp(2*c)*exp(2*d*x)*tanh(c + d*x) - 72*b**3*d*exp(2* \\
& c)*exp(2*d*x) + 24*b**3*d*tanh(c + d*x)**3 + 72*b**3*d*tanh(c + d*x)**2 + 7 \\
& 2*b**3*d*tanh(c + d*x) + 24*b**3*d) - 9*d*x*exp(6*c)*exp(6*d*x)*tanh(c + d* \\
& x)/(24*b**3*d*exp(6*c)*exp(6*d*x)*tanh(c + d*x)**3 - 72*b**3*d*exp(6*c)*exp \\
& (6*d*x)*tanh(c + d*x)**2 + 72*b**3*d*exp(6*c)*exp(6*d*x)*tanh(c + d*x) - 24 \\
& *b**3*d*exp(6*c)*exp(6*d*x) + 72*b**3*d*exp(4*c)*exp(4*d*x)*tanh(c + d*x)** \\
& 3 - 72*b**3*d*exp(4*c)*exp(4*d*x)*tanh(c + d*x)**2 - 72*b**3*d*exp(4*c)*exp \\
& (4*d*x)*tanh(c + d*x) + 72*b**3*d*exp(4*c)*exp(4*d*x) + 72*b**3*d*exp(2*c)* \\
& exp(2*d*x)*tanh(c + d*x)**3 + 72*b**3*d*exp(2*c)*exp(2*d*x)*tanh(c + d*x)** \\
& 2 - 72*b**3*d*exp(2*c)*exp(2*d*x)*tanh(c + d*x) - 72*b**3*d*exp(2*c)*exp(2* \\
& d*x) + 24*b**3*d*tanh(c + d*x)**3 + 72*b**3*d*tanh(c + d*x)**2 + 72*b**3*d* \\
& tanh(c + d*x) + 24*b**3*d) + 3*d*x*exp(6*c)*exp(6*d*x)/(24*b**3*d*exp(6*c)* \\
& exp(6*d*x)*tanh(c + d*x)**3 - 72*b**3*d*exp(6*c)*exp(6*d*x)*tanh(c + d*x)**
\end{aligned}$$

$2 + 72*b**3*d*exp(6*c)*exp(6*d*x)*tanh(c + d*x) - 24*b**3*d*exp(6*c)*exp(6*d*x) + 72*b**3*d*exp(4*c)*exp(4*d*x)*tanh(c + d*x)**3 - 72*b**3*d*exp(4*c)*exp(4*d*x)*tanh(c + d*x)**2 - 72*b**3*d*exp(4*c)*exp(4*d*x)*tanh(c + d*x) + 72*b**3*d*exp(4*c)*exp(4*d*x) + 72*b**3*d*exp(2*c)*exp(2*d*x)*tanh(c + d*x)**3 + 72*b**3*d*exp(2*c)*exp(2*d*x)*tanh(c + d*x)**2 - 72*b**3*d*exp(2*c)*exp(2*d*x)*tanh(c + d*x) - 72*b**3*d*exp(2*c)*exp(2*d*x) + 24*b**3*d*tanh(c + d*x)**3 + 72*b**3*d*tanh(c + d*x)**2 + 72*b**3*d*tanh(c + d*x) + 24*b**3*d) + 27*d*x*exp(4*c)*exp(4*d*x)*tanh(c + d*x)**3/(24*b**3*d*exp(6*c)*exp(6*d*x)*tanh(c + d*x)**3 - 72*b**3*d*exp(6*c)*exp(6*d*x)*tanh(c + d*x)**2 + 72*b**3*d*exp(6*c)*exp(6*d*x)*tanh(c + d*x) - 24*b**3*d*exp(6*c)*exp(6*d*x) + 72*b**3*d*exp(4*c)*exp(4*d*x)*tanh(c + d*x)**3 - 72*b**3*d*exp(4*c)*exp(4*d*x)*tanh(c + d*x)**2 - 72*b**3*d*exp(4*c)*exp(4*d*x)*tanh(c + d*x) + 72*b**3*d*exp(4*c)*exp(4*d*x) + 72*b**3*d*exp(2*c)*exp(2*d*x)*tanh(c + d*x)**3 + 72*b**3*d*exp(2*c)*exp(2*d*x)*tanh(c + d*x)**2 - 72*b**3*d*exp(2*c)*exp(2*d*x)*tanh(c + d*x) - 72*b**3*d*exp(2*c)*exp(2*d*x) + 24*b**3*d*tanh(c + d*x)**3 + 72*b**3*d*tanh(c + d*x)**2 + 72*b**3*d*...$

Giac [A]

time = 0.42, size = 205, normalized size = 1.59

$$\frac{(3a^2b+b^3)\log\left(\frac{-ae^{(2dx+2c)}-be^{(2dx+2c)}-a+b}{a^6-3a^4b^2+3a^2b^4-b^6}\right) - \frac{dx+c}{a^3-3a^2b+3ab^2-b^3} - \frac{2\left((3a^2b^2-4ab^3+b^4)e^{(2dx+2c)} + \frac{3(a^3b^2-2a^2b^3+ab^4)}{a+b}\right)}{(ae^{(2dx+2c)}+be^{(2dx+2c)}+a-b)^2(a+b)^2(a-b)^3}}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+b*tanh(d*x+c))^3,x, algorithm="giac")

[Out] $-\left(\frac{(3a^2b + b^3)\log(\text{abs}(-ae^{(2dx+2c)} - be^{(2dx+2c)} - a + b))}{(a^6 - 3a^4b^2 + 3a^2b^4 - b^6)} - \frac{(dx + c)}{(a^3 - 3a^2b + 3ab^2 - b^3)} - 2\left(\frac{(3a^2b^2 - 4ab^3 + b^4)e^{(2dx+2c)} + 3(a^3b^2 - 2a^2b^3 + ab^4)}{(a+b)^2(a-b)^2(a-b)^3}\right)\right)/d$

Mupad [B]

time = 2.11, size = 304, normalized size = 2.36

$$\frac{\tanh(c+dx)\left(\frac{1}{ad} - \frac{a^2+a^2b^2}{ad(a^2-2a^2b^2+b^4)}\right) + \frac{a^2x}{(a+b)(a^2+2ab+b^2)} + \frac{b^2x\tanh(c+dx)^2}{a^3+3a^2b+3ab^2+b^3} + \frac{\tanh(c+dx)^2\left(\frac{b^2}{a^2} - \frac{5a^2b^2}{a^2}\right)}{a^2d(a^2-2a^2b^2+b^4)} + \frac{2abx\tanh(c+dx)}{a^4+3a^2b+3ab^2+b^3} - \frac{\ln(a+b\tanh(c+dx))(3a^2b+b^3)}{d(a^6-3a^4b^2+3a^2b^4-b^6)} + \frac{\ln(\tanh(c+dx)+1)(3a^2b+b^3)}{d(a^2-b^2)^3}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a + b*tanh(c + d*x))^3,x)

[Out] $(\tanh(c + d*x)*(1/(a*d) - (a^4 + a^2*b^2)/(a*d*(a^4 + b^4 - 2*a^2*b^2))) + (a^2*x)/((a + b)*(2*a*b + a^2 + b^2)) + (b^2*x*tanh(c + d*x)^2)/(3*a*b^2 + 3*a^2*b + a^3 + b^3) + (\tanh(c + d*x)^2*(b^5/2 - (5*a^2*b^3)/2))/(a^2*d*(a^4 + b^4 - 2*a^2*b^2)) + (2*a*b*x*tanh(c + d*x))/(3*a*b^2 + 3*a^2*b + a^3 + b^3))/(a^2 + b^2*tanh(c + d*x)^2 + 2*a*b*tanh(c + d*x)) - (\log(a + b*tanh(c + d*x))*(3*a^2*b + b^3))/(d*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)) + (\log(\tanh(c + d*x) + 1)*(3*a^2*b + b^3))/(d*(a^2 - b^2)^3)$

3.64 $\int \frac{1}{(a+b \tanh(c+dx))^4} dx$

Optimal. Leaf size=169

$$\frac{(a^4 + 6a^2b^2 + b^4)x}{(a^2 - b^2)^4} - \frac{4ab(a^2 + b^2) \log(a \cosh(c + dx) + b \sinh(c + dx))}{(a^2 - b^2)^4 d} + \frac{b}{3(a^2 - b^2)d(a + b \tanh(c + dx))^3} + \dots$$

[Out] $(a^4+6*a^2*b^2+b^4)*x/(a^2-b^2)^4-4*a*b*(a^2+b^2)*\ln(a*\cosh(d*x+c)+b*\sinh(d*x+c))/(a^2-b^2)^4/d+1/3*b/(a^2-b^2)/d/(a+b*\tanh(d*x+c))^3+a*b/(a^2-b^2)^2/d/(a+b*\tanh(d*x+c))^2+b*(3*a^2+b^2)/(a^2-b^2)^3/d/(a+b*\tanh(d*x+c))$

Rubi [A]

time = 0.19, antiderivative size = 169, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3564, 3610, 3612, 3611}

$$\frac{b(3a^2 + b^2)}{d(a^2 - b^2)^3(a + b \tanh(c + dx))} + \frac{ab}{d(a^2 - b^2)^2(a + b \tanh(c + dx))^2} + \frac{b}{3d(a^2 - b^2)(a + b \tanh(c + dx))^3} - \frac{4ab(a^2 + b^2) \log(a \cosh(c + dx) + b \sinh(c + dx))}{d(a^2 - b^2)^4} + \frac{x(a^4 + 6a^2b^2 + b^4)}{(a^2 - b^2)^4}$$

Antiderivative was successfully verified.

[In] Int[(a + b*Tanh[c + d*x])^(-4), x]

[Out] $((a^4 + 6*a^2*b^2 + b^4)*x)/(a^2 - b^2)^4 - (4*a*b*(a^2 + b^2)*\text{Log}[a*\text{Cosh}[c + d*x] + b*\text{Sinh}[c + d*x]])/((a^2 - b^2)^4*d) + b/(3*(a^2 - b^2)*d*(a + b*\text{Tanh}[c + d*x])^3) + (a*b)/((a^2 - b^2)^2*d*(a + b*\text{Tanh}[c + d*x])^2) + (b*(3*a^2 + b^2))/((a^2 - b^2)^3*d*(a + b*\text{Tanh}[c + d*x]))$

Rule 3564

Int[((a_) + (b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[b*((a + b*Tan[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 + b^2))), x] + Dist[1/(a^2 + b^2), Int[(a - b*Tan[c + d*x])*(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0] && LtQ[n, -1]

Rule 3610

Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] :> Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Dist[1/(a^2 + b^2), Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]

Rule 3611

Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/(a_ + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] :> Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] &&

NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]

Rule 3612

Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] :> Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Dist[(b*c - a*d)/(a^2 + b^2), Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]

Rubi steps

$$\begin{aligned}
 \int \frac{1}{(a + b \tanh(c + dx))^4} dx &= \frac{b}{3(a^2 - b^2) d(a + b \tanh(c + dx))^3} + \frac{\int \frac{a - b \tanh(c + dx)}{(a + b \tanh(c + dx))^3} dx}{a^2 - b^2} \\
 &= \frac{b}{3(a^2 - b^2) d(a + b \tanh(c + dx))^3} + \frac{ab}{(a^2 - b^2)^2 d(a + b \tanh(c + dx))^2} + \frac{\int \frac{a}{(a + b \tanh(c + dx))^3} dx}{a^2 - b^2} \\
 &= \frac{b}{3(a^2 - b^2) d(a + b \tanh(c + dx))^3} + \frac{ab}{(a^2 - b^2)^2 d(a + b \tanh(c + dx))^2} + \frac{\int \frac{a}{(a + b \tanh(c + dx))^3} dx}{a^2 - b^2} \\
 &= \frac{(a^4 + 6a^2b^2 + b^4)x}{(a^2 - b^2)^4} + \frac{b}{3(a^2 - b^2) d(a + b \tanh(c + dx))^3} + \frac{ab}{(a^2 - b^2)^2 d(a + b \tanh(c + dx))^2} + \frac{\int \frac{a}{(a + b \tanh(c + dx))^3} dx}{a^2 - b^2} \\
 &= \frac{(a^4 + 6a^2b^2 + b^4)x}{(a^2 - b^2)^4} - \frac{4ab(a^2 + b^2) \log(a \cosh(c + dx) + b \sinh(c + dx))}{(a^2 - b^2)^4 d} + \frac{\int \frac{a}{(a + b \tanh(c + dx))^3} dx}{a^2 - b^2}
 \end{aligned}$$

Mathematica [A]

time = 2.31, size = 160, normalized size = 0.95

$$\frac{-\frac{3 \log(1 - \tanh(c + dx))}{(a + b)^4} + \frac{3 \log(1 + \tanh(c + dx))}{(a - b)^4} + \frac{2b \left(-12a(a^2 + b^2) \log(a + b \tanh(c + dx)) + \frac{(a^2 - b^2)(13a^4 - 2a^2b^2 + b^4 + 3ab(7a^2 + b^2) \tanh(c + dx) + 3b^2(3a^2 + b^2) \tanh^2(c + dx))}{(a + b \tanh(c + dx))^3} \right)}{(a^2 - b^2)^4}}{6d}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*Tanh[c + d*x])^(-4), x]

[Out] ((-3*Log[1 - Tanh[c + d*x]])/(a + b)^4 + (3*Log[1 + Tanh[c + d*x]])/(a - b)^4 + (2*b*(-12*a*(a^2 + b^2)*Log[a + b*Tanh[c + d*x]] + ((a^2 - b^2)*(13*a^4 - 2*a^2*b^2 + b^4 + 3*a*b*(7*a^2 + b^2)*Tanh[c + d*x] + 3*b^2*(3*a^2 + b^2)*Tanh[c + d*x]^2))/(a + b*Tanh[c + d*x]^3))/(a^2 - b^2)^4)/(6*d)

Maple [A]

time = 0.80, size = 163, normalized size = 0.96

method	result
derivativedivides	$\frac{-\frac{\ln(\tanh(dx+c)-1)}{2(a+b)^4} + \frac{\ln(\tanh(dx+c)+1)}{2(a-b)^4} + \frac{b}{3(a-b)(a+b)(a+b \tanh(dx+c))^3} + \frac{ab}{(a+b)^2(a-b)^2(a+b \tanh(dx+c))^2} + \frac{b(3a^2+b^2)}{(a+b)^3(a-b)^3(a+b \tanh(dx+c))}}{d}$
default	$\frac{-\frac{\ln(\tanh(dx+c)-1)}{2(a+b)^4} + \frac{\ln(\tanh(dx+c)+1)}{2(a-b)^4} + \frac{b}{3(a-b)(a+b)(a+b \tanh(dx+c))^3} + \frac{ab}{(a+b)^2(a-b)^2(a+b \tanh(dx+c))^2} + \frac{b(3a^2+b^2)}{(a+b)^3(a-b)^3(a+b \tanh(dx+c))}}{d}$
risch	$\frac{x}{a^4+4a^3b+6a^2b^2+4ab^3+b^4} + \frac{8ba^3x}{a^8-4a^6b^2+6b^4a^4-4a^2b^6+b^8} + \frac{8b^3ax}{a^8-4a^6b^2+6b^4a^4-4a^2b^6+b^8} + \frac{8ba^3c}{d(a^8-4a^6b^2+6b^4a^4-4a^2b^6+b^8)}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(a+b*tanh(d*x+c))^4,x,method=_RETURNVERBOSE)`

[Out] $1/d*(-1/2/(a+b)^4*\ln(\tanh(d*x+c)-1)+1/2/(a-b)^4*\ln(\tanh(d*x+c)+1)+1/3*b/(a-b)/(a+b)/(a+b*tanh(d*x+c))^3+a*b/(a+b)^2/(a-b)^2/(a+b*tanh(d*x+c))^2+b*(3*a^2+b^2)/(a+b)^3/(a-b)^3/(a+b*tanh(d*x+c))-4*b*a*(a^2+b^2)/(a+b)^4/(a-b)^4*\ln(a+b*tanh(d*x+c)))$

Maxima [B] Leaf count of result is larger than twice the leaf count of optimal. 525 vs. $2(167) = 334$.

time = 0.33, size = 525, normalized size = 3.11

$$\frac{4(a^9 + a^7) \log\left(\frac{-1 - b \tanh(dx+c)}{a-b}\right) - a - b}{(a^4 + 4a^3b + 6a^2b^2 + 4ab^3 + b^4)(a^8 - 4a^6b^2 + 6b^4a^4 - 4a^2b^6 + b^8)} + \frac{4(9a^9 + 18a^7b + 11a^5b^2 + 4a^3b^3 + 3b^4)}{3(a-b)(a+b)(a+b \tanh(dx+c))^3} + \frac{4a^2b}{(a+b)^2(a-b)^2(a+b \tanh(dx+c))^2} + \frac{4a^2b^2}{(a+b)^3(a-b)^3(a+b \tanh(dx+c))} + \frac{dx+c}{d(a^8 - 4a^6b^2 + 6b^4a^4 - 4a^2b^6 + b^8)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(a+b*tanh(d*x+c))^4,x, algorithm="maxima")`

[Out] $-4*(a^3*b + a*b^3)*\log\left(\frac{-(a-b)*e^{-2*d*x-2*c}-a-b}{(a^8-4*a^6*b^2+6*a^4*b^4-4*a^2*b^6+b^8)*d}\right) - \frac{4/3*(9*a^4*b^2+18*a^3*b^3+11*a^2*b^4+4*a*b^5+2*b^6+3*(6*a^4*b^2+2*a^3*b^3-5*a^2*b^4-2*a*b^5-b^6))*e^{-2*d*x-2*c} + 3*(3*a^4*b^2-4*a^3*b^3+b^6)*e^{-4*d*x-4*c}}{(a^{10}+2*a^9*b-3*a^8*b^2-8*a^7*b^3+2*a^6*b^4+12*a^5*b^5+2*a^4*b^6-8*a^3*b^7-3*a^2*b^8+2*a*b^9+b^{10}+3*(a^{10}-5*a^8*b^2+10*a^6*b^4-10*a^4*b^6+5*a^2*b^8-b^{10}))*e^{-2*d*x-2*c} + 3*(a^{10}-2*a^9*b-3*a^8*b^2+8*a^7*b^3+2*a^6*b^4-12*a^5*b^5+2*a^4*b^6+8*a^3*b^7-3*a^2*b^8-2*a*b^9+b^{10}))*e^{-4*d*x-4*c} + (a^{10}-4*a^9*b+3*a^8*b^2+8*a^7*b^3-14*a^6*b^4+14*a^4*b^6-8*a^3*b^7-3*a^2*b^8+4*a*b^9-b^{10})*e^{-6*d*x-6*c}}*d) + (d*x+c)/((a^4+4*a^3*b+6*a^2*b^2+4*a*b^3+b^4)*d)$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 3693 vs. $2(167) = 334$.

time = 0.45, size = 3693, normalized size = 21.85

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+b*tanh(d*x+c))^4,x, algorithm="fricas")

[Out] $\frac{1}{3} \cdot (3 \cdot (a^7 + 7 \cdot a^6 \cdot b + 21 \cdot a^5 \cdot b^2 + 35 \cdot a^4 \cdot b^3 + 35 \cdot a^3 \cdot b^4 + 21 \cdot a^2 \cdot b^5 + 7 \cdot a \cdot b^6 + b^7) \cdot d \cdot x \cdot \cosh(d \cdot x + c)^6 + 18 \cdot (a^7 + 7 \cdot a^6 \cdot b + 21 \cdot a^5 \cdot b^2 + 35 \cdot a^4 \cdot b^3 + 35 \cdot a^3 \cdot b^4 + 21 \cdot a^2 \cdot b^5 + 7 \cdot a \cdot b^6 + b^7) \cdot d \cdot x \cdot \cosh(d \cdot x + c) \cdot \sinh(d \cdot x + c)^5 + 3 \cdot (a^7 + 7 \cdot a^6 \cdot b + 21 \cdot a^5 \cdot b^2 + 35 \cdot a^4 \cdot b^3 + 35 \cdot a^3 \cdot b^4 + 21 \cdot a^2 \cdot b^5 + 7 \cdot a \cdot b^6 + b^7) \cdot d \cdot x \cdot \sinh(d \cdot x + c)^6 + 36 \cdot a^5 \cdot b^2 - 108 \cdot a^4 \cdot b^3 + 116 \cdot a^3 \cdot b^4 - 60 \cdot a^2 \cdot b^5 + 24 \cdot a \cdot b^6 - 8 \cdot b^7 + 3 \cdot (12 \cdot a^5 \cdot b^2 + 4 \cdot a^4 \cdot b^3 - 16 \cdot a^3 \cdot b^4 + 4 \cdot a \cdot b^6 - 4 \cdot b^7 + 3 \cdot (a^7 + 5 \cdot a^6 \cdot b + 9 \cdot a^5 \cdot b^2 + 5 \cdot a^4 \cdot b^3 - 5 \cdot a^3 \cdot b^4 - 9 \cdot a^2 \cdot b^5 - 5 \cdot a \cdot b^6 - b^7) \cdot d \cdot x) \cdot \cosh(d \cdot x + c)^4 + 3 \cdot (12 \cdot a^5 \cdot b^2 + 4 \cdot a^4 \cdot b^3 - 16 \cdot a^3 \cdot b^4 + 4 \cdot a \cdot b^6 - 4 \cdot b^7 + 15 \cdot (a^7 + 7 \cdot a^6 \cdot b + 21 \cdot a^5 \cdot b^2 + 35 \cdot a^4 \cdot b^3 + 35 \cdot a^3 \cdot b^4 + 21 \cdot a^2 \cdot b^5 + 7 \cdot a \cdot b^6 + b^7) \cdot d \cdot x \cdot \cosh(d \cdot x + c))^2 + 3 \cdot (a^7 + 5 \cdot a^6 \cdot b + 9 \cdot a^5 \cdot b^2 + 5 \cdot a^4 \cdot b^3 - 5 \cdot a^3 \cdot b^4 - 9 \cdot a^2 \cdot b^5 - 5 \cdot a \cdot b^6 - b^7) \cdot d \cdot x) \cdot \sinh(d \cdot x + c)^4 + 12 \cdot (5 \cdot (a^7 + 7 \cdot a^6 \cdot b + 21 \cdot a^5 \cdot b^2 + 35 \cdot a^4 \cdot b^3 + 35 \cdot a^3 \cdot b^4 + 21 \cdot a^2 \cdot b^5 + 7 \cdot a \cdot b^6 + b^7) \cdot d \cdot x \cdot \cosh(d \cdot x + c))^3 + (12 \cdot a^5 \cdot b^2 + 4 \cdot a^4 \cdot b^3 - 16 \cdot a^3 \cdot b^4 + 4 \cdot a \cdot b^6 - 4 \cdot b^7 + 3 \cdot (a^7 + 5 \cdot a^6 \cdot b + 9 \cdot a^5 \cdot b^2 + 5 \cdot a^4 \cdot b^3 - 5 \cdot a^3 \cdot b^4 - 9 \cdot a^2 \cdot b^5 - 5 \cdot a \cdot b^6 - b^7) \cdot d \cdot x) \cdot \cosh(d \cdot x + c)) \cdot \sinh(d \cdot x + c)^3 + 3 \cdot (a^7 + a^6 \cdot b - 3 \cdot a^5 \cdot b^2 - 3 \cdot a^4 \cdot b^3 + 3 \cdot a^3 \cdot b^4 + 3 \cdot a^2 \cdot b^5 - a \cdot b^6 - b^7) \cdot d \cdot x + 3 \cdot (24 \cdot a^5 \cdot b^2 - 32 \cdot a^4 \cdot b^3 - 12 \cdot a^3 \cdot b^4 + 28 \cdot a^2 \cdot b^5 - 12 \cdot a \cdot b^6 + 4 \cdot b^7 + 3 \cdot (a^7 + 3 \cdot a^6 \cdot b + a^5 \cdot b^2 - 5 \cdot a^4 \cdot b^3 - 5 \cdot a^3 \cdot b^4 + a^2 \cdot b^5 + 3 \cdot a \cdot b^6 + b^7) \cdot d \cdot x) \cdot \cosh(d \cdot x + c)^2 + 3 \cdot (24 \cdot a^5 \cdot b^2 - 32 \cdot a^4 \cdot b^3 - 12 \cdot a^3 \cdot b^4 + 28 \cdot a^2 \cdot b^5 - 12 \cdot a \cdot b^6 + 4 \cdot b^7 + 15 \cdot (a^7 + 7 \cdot a^6 \cdot b + 21 \cdot a^5 \cdot b^2 + 35 \cdot a^4 \cdot b^3 + 35 \cdot a^3 \cdot b^4 + 21 \cdot a^2 \cdot b^5 + 7 \cdot a \cdot b^6 + b^7) \cdot d \cdot x \cdot \cosh(d \cdot x + c))^4 + 3 \cdot (a^7 + 3 \cdot a^6 \cdot b + a^5 \cdot b^2 - 5 \cdot a^4 \cdot b^3 - 5 \cdot a^3 \cdot b^4 + a^2 \cdot b^5 + 3 \cdot a \cdot b^6 + b^7) \cdot d \cdot x + 6 \cdot (12 \cdot a^5 \cdot b^2 + 4 \cdot a^4 \cdot b^3 - 16 \cdot a^3 \cdot b^4 + 4 \cdot a \cdot b^6 - 4 \cdot b^7 + 3 \cdot (a^7 + 5 \cdot a^6 \cdot b + 9 \cdot a^5 \cdot b^2 + 5 \cdot a^4 \cdot b^3 - 5 \cdot a^3 \cdot b^4 - 9 \cdot a^2 \cdot b^5 - 5 \cdot a \cdot b^6 - b^7) \cdot d \cdot x) \cdot \cosh(d \cdot x + c))^2 \cdot \sinh(d \cdot x + c))^2 - 12 \cdot (a^6 \cdot b - 3 \cdot a^5 \cdot b^2 + 4 \cdot a^4 \cdot b^3 - 4 \cdot a^3 \cdot b^4 + 3 \cdot a^2 \cdot b^5 - a \cdot b^6 + (a^6 \cdot b + 3 \cdot a^5 \cdot b^2 + 4 \cdot a^4 \cdot b^3 + 4 \cdot a^3 \cdot b^4 + 3 \cdot a^2 \cdot b^5 + a \cdot b^6) \cdot \cosh(d \cdot x + c))^6 + 6 \cdot (a^6 \cdot b + 3 \cdot a^5 \cdot b^2 + 4 \cdot a^4 \cdot b^3 + 4 \cdot a^3 \cdot b^4 + 3 \cdot a^2 \cdot b^5 + a \cdot b^6) \cdot \cosh(d \cdot x + c) \cdot \sinh(d \cdot x + c))^5 + (a^6 \cdot b + 3 \cdot a^5 \cdot b^2 + 4 \cdot a^4 \cdot b^3 + 4 \cdot a^3 \cdot b^4 + 3 \cdot a^2 \cdot b^5 + a \cdot b^6) \cdot \sinh(d \cdot x + c))^6 + 3 \cdot (a^6 \cdot b + a^5 \cdot b^2 - a^2 \cdot b^5 - a \cdot b^6) \cdot \cosh(d \cdot x + c))^4 + 3 \cdot (a^6 \cdot b + a^5 \cdot b^2 - a^2 \cdot b^5 - a \cdot b^6 + 5 \cdot (a^6 \cdot b + 3 \cdot a^5 \cdot b^2 + 4 \cdot a^4 \cdot b^3 + 4 \cdot a^3 \cdot b^4 + 3 \cdot a^2 \cdot b^5 + a \cdot b^6) \cdot \cosh(d \cdot x + c))^2 \cdot \sinh(d \cdot x + c))^4 + 4 \cdot (5 \cdot (a^6 \cdot b + 3 \cdot a^5 \cdot b^2 + 4 \cdot a^4 \cdot b^3 + 4 \cdot a^3 \cdot b^4 + 3 \cdot a^2 \cdot b^5 + a \cdot b^6) \cdot \cosh(d \cdot x + c))^3 + 3 \cdot (a^6 \cdot b + a^5 \cdot b^2 - a^2 \cdot b^5 - a \cdot b^6) \cdot \cosh(d \cdot x + c)) \cdot \sinh(d \cdot x + c))^3 + 3 \cdot (a^6 \cdot b - a^5 \cdot b^2 - a^2 \cdot b^5 + a \cdot b^6 + 5 \cdot (a^6 \cdot b + 3 \cdot a^5 \cdot b^2 + 4 \cdot a^4 \cdot b^3 + 4 \cdot a^3 \cdot b^4 + 3 \cdot a^2 \cdot b^5 + a \cdot b^6) \cdot \cosh(d \cdot x + c))^4 + 6 \cdot (a^6 \cdot b + a^5 \cdot b^2 - a^2 \cdot b^5 - a \cdot b^6) \cdot \cosh(d \cdot x + c))^2 \cdot \sinh(d \cdot x + c))^2 + 6 \cdot ((a^6 \cdot b + 3 \cdot a^5 \cdot b^2 + 4 \cdot a^4 \cdot b^3 + 4 \cdot a^3 \cdot b^4 + 3 \cdot a^2 \cdot b^5 + a \cdot b^6) \cdot \cosh(d \cdot x + c))^5 + 2 \cdot (a^6 \cdot b + a^5 \cdot b^2 - a^2 \cdot b^5 - a \cdot b^6) \cdot \cosh(d \cdot x + c))^3 + (a^6 \cdot b - a^5 \cdot b^2 - a^2 \cdot b^5 + a \cdot b^6) \cdot \cosh(d \cdot x + c)) \cdot \sinh(d \cdot x + c)) \cdot \log(2 \cdot (a \cdot \cosh(d \cdot x + c) + b \cdot \sinh(d \cdot x + c)) / (\cosh(d \cdot x + c) - \sinh(d \cdot x + c))) + 6 \cdot (3 \cdot$

[Out]
$$-1/3*(12*(a^3*b + a*b^3)*\log(\text{abs}(-a*e^{(2*d*x + 2*c)} - b*e^{(2*d*x + 2*c)} - a + b))/(a^8 - 4*a^6*b^2 + 6*a^4*b^4 - 4*a^2*b^6 + b^8) - 3*(d*x + c)/(a^4 - 4*a^3*b + 6*a^2*b^2 - 4*a*b^3 + b^4) - 4*(3*(3*a^4*b^2 - 2*a^3*b^3 - 2*a^2*b^4 + 2*a*b^5 - b^6)*e^{(4*d*x + 4*c)} + 3*(6*a^4*b^2 - 14*a^3*b^3 + 11*a^2*b^4 - 4*a*b^5 + b^6)*e^{(2*d*x + 2*c)} + (9*a^5*b^2 - 27*a^4*b^3 + 29*a^3*b^4 - 15*a^2*b^5 + 6*a*b^6 - 2*b^7)/(a + b))/((a*e^{(2*d*x + 2*c)} + b*e^{(2*d*x + 2*c)} + a - b)^3*(a + b)^3*(a - b)^4)/d$$

Mupad [B]

time = 2.99, size = 452, normalized size = 2.67

$$\frac{\ln(\tanh(c+dx)+1)}{2da^4-8da^3b+12da^2b^2-8da^2b^2+2db^4} - \frac{\frac{\tanh(c+dx)(6a^4b^2-3a^3b^3+3b^4)}{d(a^6-3a^4b^2+3a^2b^4-b^6)} + \frac{\tanh(c+dx)^2(10a^4b^2-3a^3b^3+3b^4)}{a^3+3a^2b\tanh(c+dx)+3ab^2\tanh(c+dx)^2+b^3\tanh(c+dx)^3}}{a^3+3a^2b\tanh(c+dx)+3ab^2\tanh(c+dx)^2+b^3\tanh(c+dx)^3} - \frac{\ln(1-\tanh(c+dx))}{2da^4+8da^3b+12da^2b^2+8da^2b^2+2db^4} - \frac{4\ln(a+b\tanh(c+dx))(a^3b+ab^3)}{d(a^4-4a^3b+6a^2b^2-4ab^3+b^4)(a^4+4a^3b+6a^2b^2+4ab^3+b^4)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{int}(1/(a + b*\tanh(c + d*x))^4, x)$

[Out]
$$\log(\tanh(c + d*x) + 1)/(2*a^4*d + 2*b^4*d + 12*a^2*b^2*d - 8*a*b^3*d - 8*a^3*b*d) - ((\tanh(c + d*x)*(b^6 - 3*a^2*b^4 + 6*a^4*b^2))/(a*d*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)) + (\tanh(c + d*x)^2*(b^7 - 3*a^2*b^5 + 10*a^4*b^3))/(a^2*d*(3*a*b^2 - 3*a^2*b + a^3 - b^3)*(3*a*b^2 + 3*a^2*b + a^3 + b^3)) + (\tanh(c + d*x)^3*(b^8/3 - (2*a^2*b^6)/3 + (13*a^4*b^4)/3))/(a^3*d*(3*a*b^2 - 3*a^2*b + a^3 - b^3)*(3*a*b^2 + 3*a^2*b + a^3 + b^3)))/(a^3 + b^3*\tanh(c + d*x)^3 + 3*a*b^2*\tanh(c + d*x)^2 + 3*a^2*b*\tanh(c + d*x)) - \log(1 - \tanh(c + d*x))/(2*a^4*d + 2*b^4*d + 12*a^2*b^2*d + 8*a*b^3*d + 8*a^3*b*d) - (4*\log(a + b*\tanh(c + d*x))*(a*b^3 + a^3*b))/(d*(a^4 - 4*a^3*b - 4*a*b^3 + b^4 + 6*a^2*b^2)*(4*a*b^3 + 4*a^3*b + a^4 + b^4 + 6*a^2*b^2))$$

$$3.65 \quad \int \frac{1}{4+6 \tanh(c+dx)} dx$$

Optimal. Leaf size=31

$$-\frac{x}{5} + \frac{3 \log(2 \cosh(c+dx) + 3 \sinh(c+dx))}{10d}$$

[Out] -1/5*x+3/10*ln(2*cosh(d*x+c)+3*sinh(d*x+c))/d

Rubi [A]

time = 0.03, antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3565, 3611}

$$\frac{3 \log(3 \sinh(c+dx) + 2 \cosh(c+dx))}{10d} - \frac{x}{5}$$

Antiderivative was successfully verified.

[In] Int[(4 + 6*Tanh[c + d*x])^(-1), x]

[Out] -1/5*x + (3*Log[2*Cosh[c + d*x] + 3*Sinh[c + d*x]])/(10*d)

Rule 3565

Int[((a_) + (b_.)*tan[(c_) + (d_.)*(x_)])^(-1), x_Symbol] := Simp[a*(x/(a^2 + b^2)), x] + Dist[b/(a^2 + b^2), Int[(b - a*Tan[c + d*x])/(a + b*Tan[c + d*x]), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0]

Rule 3611

Int[((c_) + (d_.)*tan[(e_) + (f_.)*(x_)])/((a_) + (b_.)*tan[(e_) + (f_.)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{4+6 \tanh(c+dx)} dx &= -\frac{x}{5} + \frac{3}{10} i \int \frac{-6i - 4i \tanh(c+dx)}{4+6 \tanh(c+dx)} dx \\ &= -\frac{x}{5} + \frac{3 \log(2 \cosh(c+dx) + 3 \sinh(c+dx))}{10d} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 55, normalized size = 1.77

$$-\frac{\log(6 - 6 \tanh(c+dx))}{20d} + \frac{3 \log(4 + 6 \tanh(c+dx))}{10d} - \frac{\log(6 + 6 \tanh(c+dx))}{4d}$$

Antiderivative was successfully verified.

```
[In] Integrate[(4 + 6*Tanh[c + d*x])^(-1),x]
```

```
[Out] -1/20*Log[6 - 6*Tanh[c + d*x]]/d + (3*Log[4 + 6*Tanh[c + d*x]])/(10*d) - Log[6 + 6*Tanh[c + d*x]]/(4*d)
```

Maple [A]

time = 0.63, size = 42, normalized size = 1.35

method	result	size
risch	$-\frac{x}{2} - \frac{3c}{5d} + \frac{3 \ln(e^{2dx+2c} - \frac{1}{5})}{10d}$	28
derivativdivides	$\frac{\frac{3 \ln(2+3 \tanh(dx+c))}{5} - \frac{\ln(\tanh(dx+c)-1)}{10} - \frac{\ln(\tanh(dx+c)+1)}{2}}{2d}$	42
default	$\frac{\frac{3 \ln(2+3 \tanh(dx+c))}{5} - \frac{\ln(\tanh(dx+c)-1)}{10} - \frac{\ln(\tanh(dx+c)+1)}{2}}{2d}$	42

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(4+6*tanh(d*x+c)),x,method=_RETURNVERBOSE)
```

```
[Out] 1/2/d*(3/5*ln(2+3*tanh(d*x+c))-1/10*ln(tanh(d*x+c)-1)-1/2*ln(tanh(d*x+c)+1))
```

Maxima [A]

time = 0.26, size = 28, normalized size = 0.90

$$\frac{dx + c}{10d} + \frac{3 \log(e^{(-2dx-2c)} - 5)}{10d}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(4+6*tanh(d*x+c)),x, algorithm="maxima")
```

```
[Out] 1/10*(d*x + c)/d + 3/10*log(e^(-2*d*x - 2*c) - 5)/d
```

Fricas [A]

time = 0.44, size = 49, normalized size = 1.58

$$\frac{5dx - 3 \log\left(\frac{2(2 \cosh(dx+c) + 3 \sinh(dx+c))}{\cosh(dx+c) - \sinh(dx+c)}\right)}{10d}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(4+6*tanh(d*x+c)),x, algorithm="fricas")
```

```
[Out] -1/10*(5*d*x - 3*log(2*(2*cosh(d*x + c) + 3*sinh(d*x + c))/(cosh(d*x + c) - sinh(d*x + c))))/d
```

Sympy [A]

time = 0.23, size = 42, normalized size = 1.35

$$\begin{cases} \frac{x}{10} - \frac{3 \log(\tanh(c+dx)+1)}{10d} + \frac{3 \log(3 \tanh(c+dx)+2)}{10d} & \text{for } d \neq 0 \\ \frac{x}{6 \tanh(c)+4} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(4+6*tanh(d*x+c)),x)

[Out] Piecewise((x/10 - 3*log(tanh(c + d*x) + 1)/(10*d) + 3*log(3*tanh(c + d*x) + 2)/(10*d), Ne(d, 0)), (x/(6*tanh(c) + 4), True))

Giac [A]

time = 0.41, size = 30, normalized size = 0.97

$$-\frac{5 dx + 5 c - 3 \log(|5 e^{(2 dx + 2 c)} - 1|)}{10 d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(4+6*tanh(d*x+c)),x, algorithm="giac")

[Out] -1/10*(5*d*x + 5*c - 3*log(abs(5*e^(2*d*x + 2*c) - 1)))/d

Mupad [B]

time = 0.13, size = 34, normalized size = 1.10

$$\frac{x}{10} - \frac{\frac{3 \ln(\tanh(c+dx)+1)}{10} - \frac{3 \ln(3 \tanh(c+dx)+2)}{10}}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(6*tanh(c + d*x) + 4),x)

[Out] x/10 - ((3*log(tanh(c + d*x) + 1))/10 - (3*log(3*tanh(c + d*x) + 2))/10)/d

$$3.66 \quad \int \frac{1}{4-6 \tanh(c+dx)} dx$$

Optimal. Leaf size=31

$$-\frac{x}{5} - \frac{3 \log(2 \cosh(c+dx) - 3 \sinh(c+dx))}{10d}$$

[Out] $-1/5*x-3/10*\ln(2*\cosh(d*x+c)-3*\sinh(d*x+c))/d$

Rubi [A]

time = 0.03, antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3565, 3611}

$$-\frac{3 \log(2 \cosh(c+dx) - 3 \sinh(c+dx))}{10d} - \frac{x}{5}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(4 - 6*\text{Tanh}[c + d*x])^{-1}, x]$

[Out] $-1/5*x - (3*\text{Log}[2*\text{Cosh}[c + d*x] - 3*\text{Sinh}[c + d*x]])/(10*d)$

Rule 3565

$\text{Int}[(a_ + (b_)*\text{tan}[(c_ + (d_)*(x_)]))^{-1}, x_Symbol] := \text{Simp}[a*(x/(a^2 + b^2)), x] + \text{Dist}[b/(a^2 + b^2), \text{Int}[(b - a*\text{Tan}[c + d*x])/(a + b*\text{Tan}[c + d*x]), x], x] /; \text{FreeQ}\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[a^2 + b^2, 0]$

Rule 3611

$\text{Int}[(c_ + (d_)*\text{tan}[(e_ + (f_)*(x_)]))/(a_ + (b_)*\text{tan}[(e_ + (f_)*(x_)]), x_Symbol] := \text{Simp}[(c/(b*f))*\text{Log}[\text{RemoveContent}[a*\text{Cos}[e + f*x] + b*\text{Sin}[e + f*x], x]], x] /; \text{FreeQ}\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{EqQ}[a*c + b*d, 0]$

Rubi steps

$$\begin{aligned} \int \frac{1}{4-6 \tanh(c+dx)} dx &= -\frac{x}{5} - \frac{3}{10} i \int \frac{6i - 4i \tanh(c+dx)}{4-6 \tanh(c+dx)} dx \\ &= -\frac{x}{5} - \frac{3 \log(2 \cosh(c+dx) - 3 \sinh(c+dx))}{10d} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 55, normalized size = 1.77

$$-\frac{3 \log(4 - 6 \tanh(c+dx))}{10d} + \frac{\log(6 - 6 \tanh(c+dx))}{4d} + \frac{\log(6 + 6 \tanh(c+dx))}{20d}$$

Antiderivative was successfully verified.

[In] Integrate[(4 - 6*Tanh[c + d*x])^(-1),x]

[Out] (-3*Log[4 - 6*Tanh[c + d*x]])/(10*d) + Log[6 - 6*Tanh[c + d*x]]/(4*d) + Log[6 + 6*Tanh[c + d*x]]/(20*d)

Maple [A]

time = 0.64, size = 42, normalized size = 1.35

method	result	size
risch	$\frac{x}{10} + \frac{3c}{5d} - \frac{3 \ln(e^{2dx+2c}-5)}{10d}$	28
derivativedivides	$\frac{\frac{\ln(\tanh(dx+c)+1)}{10} - \frac{3 \ln(-2+3 \tanh(dx+c))}{5} + \frac{\ln(\tanh(dx+c)-1)}{2}}{2d}$	42
default	$\frac{\frac{\ln(\tanh(dx+c)+1)}{10} - \frac{3 \ln(-2+3 \tanh(dx+c))}{5} + \frac{\ln(\tanh(dx+c)-1)}{2}}{2d}$	42

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(4-6*tanh(d*x+c)),x,method=_RETURNVERBOSE)

[Out] 1/2/d*(1/10*ln(tanh(d*x+c)+1)-3/5*ln(-2+3*tanh(d*x+c))+1/2*ln(tanh(d*x+c)-1))

Maxima [A]

time = 0.28, size = 29, normalized size = 0.94

$$-\frac{1}{2}x - \frac{c}{2d} - \frac{3 \log(5e^{(-2dx-2c)} - 1)}{10d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(4-6*tanh(d*x+c)),x, algorithm="maxima")

[Out] -1/2*x - 1/2*c/d - 3/10*log(5*e^(-2*d*x - 2*c) - 1)/d

Fricas [A]

time = 0.35, size = 48, normalized size = 1.55

$$\frac{dx - 3 \log\left(-\frac{2(2 \cosh(dx+c)-3 \sinh(dx+c))}{\cosh(dx+c)-\sinh(dx+c)}\right)}{10d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(4-6*tanh(d*x+c)),x, algorithm="fricas")

[Out] 1/10*(d*x - 3*log(-2*(2*cosh(d*x + c) - 3*sinh(d*x + c))/(cosh(d*x + c) - sinh(d*x + c))))/d

Sympy [A]

time = 0.23, size = 42, normalized size = 1.35

$$\begin{cases} -\frac{x}{2} + \frac{3 \log(\tanh(c+dx)+1)}{10d} - \frac{3 \log(3 \tanh(c+dx)-2)}{10d} & \text{for } d \neq 0 \\ \frac{x}{4-6 \tanh(c)} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/(4-6*tanh(d*x+c)),x)``[Out] Piecewise((-x/2 + 3*log(tanh(c + d*x) + 1)/(10*d) - 3*log(3*tanh(c + d*x) - 2)/(10*d), Ne(d, 0)), (x/(4 - 6*tanh(c))), True))`**Giac [A]**

time = 0.41, size = 25, normalized size = 0.81

$$\frac{dx + c - 3 \log(|e^{(2dx+2c)} - 5|)}{10d}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/(4-6*tanh(d*x+c)),x, algorithm="giac")``[Out] 1/10*(d*x + c - 3*log(abs(e^(2*d*x + 2*c) - 5)))/d`**Mupad [B]**

time = 0.12, size = 33, normalized size = 1.06

$$\frac{\frac{3 \ln(\tanh(c+dx)+1)}{10} - \frac{3 \ln(3 \tanh(c+dx)-2)}{10}}{d} - \frac{x}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(-1/(6*tanh(c + d*x) - 4),x)``[Out] ((3*log(tanh(c + d*x) + 1))/10 - (3*log(3*tanh(c + d*x) - 2))/10)/d - x/2`

3.67 $\int \sqrt{a + b \tanh(c + dx)} dx$

Optimal. Leaf size=74

$$-\frac{\sqrt{a-b} \tanh^{-1}\left(\frac{\sqrt{a+b \tanh(c+dx)}}{\sqrt{a-b}}\right)}{d} + \frac{\sqrt{a+b} \tanh^{-1}\left(\frac{\sqrt{a+b \tanh(c+dx)}}{\sqrt{a+b}}\right)}{d}$$

[Out] $-\operatorname{arctanh}\left(\frac{(a+b \tanh(d*x+c))^{1/2}}{(a-b)^{1/2}}\right) \cdot (a-b)^{1/2}/d + \operatorname{arctanh}\left(\frac{(a+b \tanh(d*x+c))^{1/2}}{(a+b)^{1/2}}\right) \cdot (a+b)^{1/2}/d$

Rubi [A]

time = 0.05, antiderivative size = 74, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3566, 714, 1144, 213}

$$\frac{\sqrt{a+b} \tanh^{-1}\left(\frac{\sqrt{a+b \tanh(c+dx)}}{\sqrt{a+b}}\right)}{d} - \frac{\sqrt{a-b} \tanh^{-1}\left(\frac{\sqrt{a+b \tanh(c+dx)}}{\sqrt{a-b}}\right)}{d}$$

Antiderivative was successfully verified.

[In] `Int[Sqrt[a + b*Tanh[c + d*x]],x]`

[Out] $-\left(\frac{\operatorname{Sqrt}[a-b] \operatorname{ArcTanh}\left[\frac{\operatorname{Sqrt}[a+b \operatorname{Tanh}[c+d*x]]}{\operatorname{Sqrt}[a-b]}\right]}{d}\right) + \left(\frac{\operatorname{Sqrt}[a+b] \operatorname{ArcTanh}\left[\frac{\operatorname{Sqrt}[a+b \operatorname{Tanh}[c+d*x]]}{\operatorname{Sqrt}[a+b]}\right]}{d}\right)$

Rule 213

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`

Rule 714

`Int[Sqrt[(d_) + (e_.)*(x_)]/((a_) + (c_.)*(x_)^2), x_Symbol] := Dist[2*e, Subst[Int[x^2/(c*d^2 + a*e^2 - 2*c*d*x^2 + c*x^4), x], x, Sqrt[d + e*x]], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0]`

Rule 1144

`Int[((d_.)*(x_))^(m_)/((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Dist[(d^2/2)*(b/q + 1), Int[(d*x)^(m-2)/(b/2 + q/2 + c*x^2), x], x] - Dist[(d^2/2)*(b/q - 1), Int[(d*x)^(m-2)/(b/2 - q/2 + c*x^2), x], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b^2 - 4*a*c, 0] && GeQ[m, 2]`

Rule 3566

$\text{Int}[(a + (b \cdot \tan(c + d \cdot x)))^n, x_Symbol] \rightarrow \text{Dist}[b/d, \text{Subst}[\text{Int}[(a + x)^n/(b^2 + x^2), x], x, b \cdot \text{Tan}[c + d \cdot x]], x] /;$ FreeQ[{a, b, c, d, n}, x] && NeQ[a^2 + b^2, 0]

Rubi steps

$$\begin{aligned} \int \sqrt{a + b \tanh(c + dx)} dx &= -\frac{b \text{Subst}\left(\int \frac{\sqrt{a+x}}{-b^2+x^2} dx, x, b \tanh(c+dx)\right)}{d} \\ &= -\frac{(2b) \text{Subst}\left(\int \frac{x^2}{a^2-b^2-2ax^2+x^4} dx, x, \sqrt{a+b \tanh(c+dx)}\right)}{d} \\ &= \frac{(a-b) \text{Subst}\left(\int \frac{1}{-a+b+x^2} dx, x, \sqrt{a+b \tanh(c+dx)}\right)}{d} - \frac{(a+b) \text{Subst}\left(\int \frac{1}{a+b+x^2} dx, x, \sqrt{a+b \tanh(c+dx)}\right)}{d} \\ &= -\frac{\sqrt{a-b} \tanh^{-1}\left(\frac{\sqrt{a+b \tanh(c+dx)}}{\sqrt{a-b}}\right)}{d} + \frac{\sqrt{a+b} \tanh^{-1}\left(\frac{\sqrt{a+b \tanh(c+dx)}}{\sqrt{a+b}}\right)}{d} \end{aligned}$$

Mathematica [A]

time = 0.06, size = 74, normalized size = 1.00

$$-\frac{\sqrt{a-b} \tanh^{-1}\left(\frac{\sqrt{a+b \tanh(c+dx)}}{\sqrt{a-b}}\right)}{d} + \frac{\sqrt{a+b} \tanh^{-1}\left(\frac{\sqrt{a+b \tanh(c+dx)}}{\sqrt{a+b}}\right)}{d}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[a + b*Tanh[c + d*x]],x]

[Out] -((Sqrt[a - b]*ArcTanh[Sqrt[a + b*Tanh[c + d*x]]/Sqrt[a - b]])/d) + (Sqrt[a + b]*ArcTanh[Sqrt[a + b*Tanh[c + d*x]]/Sqrt[a + b]])/d

Maple [A]

time = 1.75, size = 70, normalized size = 0.95

method	result
derivativedivides	$-\frac{\sqrt{a+b} \arctan\left(\frac{\sqrt{a+b \tanh(dx+c)}}{\sqrt{a+b}}\right) + \sqrt{-a+b} \arctan\left(\frac{\sqrt{a+b \tanh(dx+c)}}{\sqrt{-a+b}}\right)}{d}$

default	$-\frac{\sqrt{a+b} \operatorname{arctanh}\left(\frac{\sqrt{a+b \tanh(dx+c)}}{\sqrt{a+b}}\right) + \sqrt{-a+b} \operatorname{arctan}\left(\frac{\sqrt{a+b \tanh(dx+c)}}{\sqrt{-a+b}}\right)}{d}$
---------	---

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((a+b*tanh(d*x+c))^(1/2),x,method=_RETURNVERBOSE)
```

```
[Out] -2/d*b*(-1/2*(a+b)^(1/2)/b*arctanh((a+b*tanh(d*x+c))^(1/2)/(a+b)^(1/2))+1/2
*(-a+b)^(1/2)/b*arctan((a+b*tanh(d*x+c))^(1/2)/(-a+b)^(1/2)))
```

Maxima [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a+b*tanh(d*x+c))^(1/2),x, algorithm="maxima")
```

```
[Out] Exception raised: ValueError >> Computation failed since Maxima requested a
dditional constraints; using the 'assume' command before evaluation *may* h
elp (example of legal syntax is 'assume(4*a-4*b>0)', see 'assume?' for more
detail
```

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 332 vs. 2(62) = 124.

time = 0.65, size = 2203, normalized size = 29.77

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a+b*tanh(d*x+c))^(1/2),x, algorithm="fricas")
```

```
[Out] [1/4*(sqrt(a + b)*log(2*(a^2 + 2*a*b + b^2)*cosh(d*x + c)^4 + 8*(a^2 + 2*a*
b + b^2)*cosh(d*x + c)*sinh(d*x + c)^3 + 2*(a^2 + 2*a*b + b^2)*sinh(d*x + c
)^4 + 4*(a^2 + a*b)*cosh(d*x + c)^2 + 4*(3*(a^2 + 2*a*b + b^2)*cosh(d*x + c
)^2 + a^2 + a*b)*sinh(d*x + c)^2 + 2*a^2 - b^2 + 2*((a + b)*cosh(d*x + c)^4
+ 4*(a + b)*cosh(d*x + c)*sinh(d*x + c)^3 + (a + b)*sinh(d*x + c)^4 + (2*a
+ b)*cosh(d*x + c)^2 + (6*(a + b)*cosh(d*x + c)^2 + 2*a + b)*sinh(d*x + c)
^2 + 2*(2*(a + b)*cosh(d*x + c)^3 + (2*a + b)*cosh(d*x + c))*sinh(d*x + c)
+ a)*sqrt(a + b)*sqrt((a*cosh(d*x + c) + b*sinh(d*x + c))/cosh(d*x + c)) +
8*((a^2 + 2*a*b + b^2)*cosh(d*x + c)^3 + (a^2 + a*b)*cosh(d*x + c))*sinh(d*
x + c)) + sqrt(a - b)*log(((2*a^2 - b^2)*cosh(d*x + c)^4 + 4*(2*a^2 - b^2)*
cosh(d*x + c)*sinh(d*x + c)^3 + (2*a^2 - b^2)*sinh(d*x + c)^4 + 4*(a^2 - a*
```

$b) \cosh(dx + c)^2 + 2*(3*(2*a^2 - b^2)*\cosh(dx + c)^2 + 2*a^2 - 2*a*b)*\sinh(dx + c)^2 + 2*a^2 - 4*a*b + 2*b^2 - 2*(a*\cosh(dx + c)^4 + 4*a*\cosh(dx + c)*\sinh(dx + c)^3 + a*\sinh(dx + c)^4 + (2*a - b)*\cosh(dx + c)^2 + (6*a*\cosh(dx + c)^2 + 2*a - b)*\sinh(dx + c)^2 + 2*(2*a*\cosh(dx + c)^3 + (2*a - b)*\cosh(dx + c))*\sinh(dx + c) + a - b)*\sqrt{a - b}*\sqrt{(a*\cosh(dx + c) + b*\sinh(dx + c))/\cosh(dx + c)} + 4*((2*a^2 - b^2)*\cosh(dx + c)^3 + 2*(a^2 - a*b)*\cosh(dx + c))*\sinh(dx + c))/(\cosh(dx + c)^4 + 4*\cosh(dx + c)^3*\sinh(dx + c) + 6*\cosh(dx + c)^2*\sinh(dx + c)^2 + 4*\cosh(dx + c)*\sinh(dx + c)^3 + \sinh(dx + c)^4))/d, -1/4*(2*\sqrt{-a - b}*\arctan(((a + b)*\cosh(dx + c)^2 + 2*(a + b)*\cosh(dx + c)*\sinh(dx + c) + (a + b)*\sinh(dx + c)^2 + a)*\sqrt{-a - b}*\sqrt{(a*\cosh(dx + c) + b*\sinh(dx + c))/\cosh(dx + c)}))/((a^2 + 2*a*b + b^2)*\cosh(dx + c)^2 + 2*(a^2 + 2*a*b + b^2)*\cosh(dx + c)*\sinh(dx + c) + (a^2 + 2*a*b + b^2)*\sinh(dx + c)^2 + a^2 - b^2)) - \sqrt{a - b}*\log(((2*a^2 - b^2)*\cosh(dx + c)^4 + 4*(2*a^2 - b^2)*\cosh(dx + c)*\sinh(dx + c)^3 + (2*a^2 - b^2)*\sinh(dx + c)^4 + 4*(a^2 - a*b)*\cosh(dx + c)^2 + 2*(3*(2*a^2 - b^2)*\cosh(dx + c)^2 + 2*a^2 - 2*a*b)*\sinh(dx + c)^2 + 2*a^2 - 4*a*b + 2*b^2 - 2*(a*\cosh(dx + c)^4 + 4*a*\cosh(dx + c)*\sinh(dx + c)^3 + a*\sinh(dx + c)^4 + (2*a - b)*\cosh(dx + c)^2 + (6*a*\cosh(dx + c)^2 + 2*a - b)*\sinh(dx + c)^2 + 2*(2*a*\cosh(dx + c)^3 + (2*a - b)*\cosh(dx + c))*\sinh(dx + c) + a - b)*\sqrt{a - b}*\sqrt{(a*\cosh(dx + c) + b*\sinh(dx + c))/\cosh(dx + c)} + 4*((2*a^2 - b^2)*\cosh(dx + c)^3 + 2*(a^2 - a*b)*\cosh(dx + c))*\sinh(dx + c))/(\cosh(dx + c)^4 + 4*\cosh(dx + c)^3*\sinh(dx + c) + 6*\cosh(dx + c)^2*\sinh(dx + c)^2 + 4*\cosh(dx + c)*\sinh(dx + c)^3 + \sinh(dx + c)^4))/d, -1/4*(2*\sqrt{-a + b}*\arctan(-(a*\cosh(dx + c)^2 + 2*a*\cosh(dx + c)*\sinh(dx + c) + a*\sinh(dx + c)^2 + a - b)*\sqrt{-a + b}*\sqrt{(a*\cosh(dx + c) + b*\sinh(dx + c))/\cosh(dx + c)}))/((a^2 - b^2)*\cosh(dx + c)^2 + 2*(a^2 - b^2)*\cosh(dx + c)*\sinh(dx + c) + (a^2 - b^2)*\sinh(dx + c)^2 + a^2 - 2*a*b + b^2)) - \sqrt{a + b}*\log(2*(a^2 + 2*a*b + b^2)*\cosh(dx + c)^4 + 8*(a^2 + 2*a*b + b^2)*\cosh(dx + c)*\sinh(dx + c)^3 + 2*(a^2 + 2*a*b + b^2)*\sinh(dx + c)^4 + 4*(a^2 + a*b)*\cosh(dx + c)^2 + 4*(3*(a^2 + 2*a*b + b^2)*\cosh(dx + c)^2 + a^2 + a*b)*\sinh(dx + c)^2 + 2*a^2 - b^2 + 2*((a + b)*\cosh(dx + c)^4 + 4*(a + b)*\cosh(dx + c)*\sinh(dx + c)^3 + (a + b)*\sinh(dx + c)^4 + (2*a + b)*\cosh(dx + c)^2 + (6*(a + b)*\cosh(dx + c)^2 + 2*a + b)*\sinh(dx + c)^2 + 2*(2*(a + b)*\cosh(dx + c)^3 + (2*a + b)*\cosh(dx + c))*\sinh(dx + c) + a)*\sqrt{a + b}*\sqrt{(a*\cosh(dx + c) + b*\sinh(dx + c))/\cosh(dx + c)} + 8*((a^2 + 2*a*b + b^2)*\cosh(dx + c)^3 + (a^2 + a*b)*\cosh(dx + c))*\sinh(dx + c))/d, -1/2*(\sqrt{-a + b}*\arctan(-(a*\cosh(dx + c)^2 + 2*a*\cosh(dx + c)*\sinh(dx + c) + a*\sinh(dx + c)^2 + a - b)*\sqrt{-a + b}*\sqrt{(a*\cosh(dx + c) + b*\sinh(dx + c))/\cosh(dx + c)}))/((a^2 - b^2)*\cosh(dx + c)^2 + 2*(a^2 - b^2)*\cosh(dx + c)*\sinh(dx + c) + (a^2 - b^2)*\sinh(dx + c)^2 + a^2 - 2*a*b + b^2)) + \sqrt{-a - b}*\arctan(((a + b)*\cosh(dx + c)^2 + 2*(a + b)*\cosh(dx + c)*\sinh(dx + c) + (a + b)*\sinh(dx + c)^2 + a)*\sqrt{-a - b}*\sqrt{(a*\cosh(dx + c) + b*\sinh(dx + c))/\cosh(dx + c)}))/((a^2 + 2*a*b + b^2)*\cosh(dx + c)^2 + 2*(a^2 + 2*a*b + b^2)*\cosh(dx + c)*\sinh(dx + c) + (a^2 + 2*a*b + b^2)*\sinh(dx + c)^2 + a^2 - b^2))/d$

]

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{a + b \tanh(c + dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+b*tanh(d*x+c))**(1/2),x)

[Out] Integral(sqrt(a + b*tanh(c + d*x)), x)

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+b*tanh(d*x+c))^(1/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
 UT:sage2:=int(sage0,sageVARx)::OUTPUT:index.cc index_m operator + Error: Ba
 d Argument Value

Mupad [B]

time = 1.34, size = 151, normalized size = 2.04

$$\frac{\operatorname{atan}\left(\frac{\sqrt{a+b}\sqrt{a+b\tanh(c+dx)} - \sqrt{a-b}\sqrt{a+b\tanh(c+dx)}}{a^2-b^2}\right)\sqrt{a+b} + \operatorname{atan}\left(\frac{\sqrt{a-b}\sqrt{a+b\tanh(c+dx)} - \sqrt{a+b}\sqrt{a+b\tanh(c+dx)}}{a^2-b^2}\right)\sqrt{a-b}}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*tanh(c + d*x))^(1/2),x)

[Out] (atan((b^2*(a + b)^(1/2)*(a + b*tanh(c + d*x))^(1/2)*1i - a*b*(a + b)^(1/2)
 *(a + b*tanh(c + d*x))^(1/2)*1i)/(a^2*b - b^3))*(a + b)^(1/2)*1i)/d + (atan
 ((b^2*(a - b)^(1/2)*(a + b*tanh(c + d*x))^(1/2)*1i + a*b*(a - b)^(1/2)*(a +
 b*tanh(c + d*x))^(1/2)*1i)/(a^2*b - b^3))*(a - b)^(1/2)*1i)/d

$$3.68 \quad \int \frac{1}{\sqrt{a + b \tanh(c + dx)}} dx$$

Optimal. Leaf size=74

$$-\frac{\tanh^{-1}\left(\frac{\sqrt{a + b \tanh(c + dx)}}{\sqrt{a - b}}\right)}{\sqrt{a - b} d} + \frac{\tanh^{-1}\left(\frac{\sqrt{a + b \tanh(c + dx)}}{\sqrt{a + b}}\right)}{\sqrt{a + b} d}$$

[Out] $-\operatorname{arctanh}\left(\frac{(a+b*\tanh(d*x+c))^{(1/2)}}{(a-b)^{(1/2)}}\right)/d/(a-b)^{(1/2)}+\operatorname{arctanh}\left(\frac{(a+b*\tanh(d*x+c))^{(1/2)}}{(a+b)^{(1/2)}}\right)/d/(a+b)^{(1/2)}$

Rubi [A]

time = 0.04, antiderivative size = 74, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3566, 722, 1107, 213}

$$\frac{\tanh^{-1}\left(\frac{\sqrt{a + b \tanh(c + dx)}}{\sqrt{a + b}}\right)}{d\sqrt{a + b}} - \frac{\tanh^{-1}\left(\frac{\sqrt{a + b \tanh(c + dx)}}{\sqrt{a - b}}\right)}{d\sqrt{a - b}}$$

Antiderivative was successfully verified.

[In] Int[1/Sqrt[a + b*Tanh[c + d*x]],x]

[Out] $-(\operatorname{ArcTanh}[\operatorname{Sqrt}[a + b*\operatorname{Tanh}[c + d*x]]/\operatorname{Sqrt}[a - b]]/(\operatorname{Sqrt}[a - b]*d)) + \operatorname{ArcTanh}[\operatorname{Sqrt}[a + b*\operatorname{Tanh}[c + d*x]]/\operatorname{Sqrt}[a + b]]/(\operatorname{Sqrt}[a + b]*d)$

Rule 213

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])

Rule 722

Int[1/(Sqrt[(d_) + (e_)*(x_)]*((a_) + (c_)*(x_)^2)), x_Symbol] := Dist[2*e, Subst[Int[1/(c*d^2 + a*e^2 - 2*c*d*x^2 + c*x^4), x], x, Sqrt[d + e*x]], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0]

Rule 1107

Int[((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(-1), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Dist[c/q, Int[1/(b/2 - q/2 + c*x^2), x], x] - Dist[c/q, Int[1/(b/2 + q/2 + c*x^2), x], x]] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[b^2 - 4*a*c]

Rule 3566

```
Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[b/d, Sub
st[Int[(a + x)^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{a, b, c,
d, n}, x] && NeQ[a^2 + b^2, 0]
```

Rubi steps

$$\begin{aligned}
\int \frac{1}{\sqrt{a + b \tanh(c + dx)}} dx &= -\frac{b \operatorname{Subst}\left(\int \frac{1}{\sqrt{a + x}(-b^2 + x^2)} dx, x, b \tanh(c + dx)\right)}{d} \\
&= -\frac{(2b) \operatorname{Subst}\left(\int \frac{1}{a^2 - b^2 - 2ax^2 + x^4} dx, x, \sqrt{a + b \tanh(c + dx)}\right)}{d} \\
&= -\frac{\operatorname{Subst}\left(\int \frac{1}{-a - b + x^2} dx, x, \sqrt{a + b \tanh(c + dx)}\right)}{d} + \frac{\operatorname{Subst}\left(\int \frac{1}{-a + b + x^2} dx, x, \sqrt{a + b \tanh(c + dx)}\right)}{d} \\
&= -\frac{\tanh^{-1}\left(\frac{\sqrt{a + b \tanh(c + dx)}}{\sqrt{a - b}}\right)}{\sqrt{a - b} d} + \frac{\tanh^{-1}\left(\frac{\sqrt{a + b \tanh(c + dx)}}{\sqrt{a + b}}\right)}{\sqrt{a + b} d}
\end{aligned}$$

Mathematica [A]

time = 0.05, size = 74, normalized size = 1.00

$$-\frac{\tanh^{-1}\left(\frac{\sqrt{a + b \tanh(c + dx)}}{\sqrt{a - b}}\right)}{\sqrt{a - b} d} + \frac{\tanh^{-1}\left(\frac{\sqrt{a + b \tanh(c + dx)}}{\sqrt{a + b}}\right)}{\sqrt{a + b} d}$$

Antiderivative was successfully verified.

```
[In] Integrate[1/Sqrt[a + b*Tanh[c + d*x]],x]
```

```
[Out] -(ArcTanh[Sqrt[a + b*Tanh[c + d*x]]/Sqrt[a - b]]/(Sqrt[a - b]*d)) + ArcTanh
[Sqrt[a + b*Tanh[c + d*x]]/Sqrt[a + b]]/(Sqrt[a + b]*d)
```

Maple [A]

time = 1.56, size = 70, normalized size = 0.95

method	result	size
derivativedivides	$ -\frac{2b \left(\frac{\operatorname{arctanh}\left(\frac{\sqrt{a + b \tanh(dx + c)}}{\sqrt{-a + b}}\right)}{2b\sqrt{-a + b}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{a + b \tanh(dx + c)}}{\sqrt{a + b}}\right)}{2b\sqrt{a + b}} \right)}{d} $	70

default	$\frac{2b \left(\frac{\arctan\left(\frac{\sqrt{a+b \tanh(dx+c)}}{\sqrt{-a+b}}\right)}{2b\sqrt{-a+b}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \tanh(dx+c)}}{\sqrt{a+b}}\right)}{2b\sqrt{a+b}} \right)}{d}$	70
---------	--	----

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(a+b*tanh(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

[Out]
$$-2/d*b*(-1/2/b/(-a+b)^{(1/2)}*\arctan((a+b*\tanh(d*x+c))^{(1/2)/(-a+b)^{(1/2)})}-1/2/b/(a+b)^{(1/2)}*\operatorname{arctanh}((a+b*\tanh(d*x+c))^{(1/2)/(a+b)^{(1/2)}))$$

Maxima [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(a+b*tanh(d*x+c))^(1/2),x, algorithm="maxima")`

[Out] Exception raised: ValueError >> Computation failed since Maxima requested a dditional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a-4*b>0)', see 'assume?' for more detail

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 351 vs. 2(62) = 124.

time = 0.57, size = 2279, normalized size = 30.80

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(a+b*tanh(d*x+c))^(1/2),x, algorithm="fricas")`

[Out]
$$\begin{aligned} & [1/4*(\sqrt{a+b}*(a-b)*\log(2*(a^2+2*a*b+b^2)*\cosh(dx+c)^4+8*(a^2+2*a*b+b^2)*\cosh(dx+c)*\sinh(dx+c)^3+2*(a^2+2*a*b+b^2)*\sinh(dx+c)^4+4*(a^2+a*b)*\cosh(dx+c)^2+4*(3*(a^2+2*a*b+b^2)*\cosh(dx+c)^2+a^2+a*b)*\sinh(dx+c)^2+2*a^2-b^2+2*((a+b)*\cosh(dx+c)^4+4*(a+b)*\cosh(dx+c)*\sinh(dx+c)^3+(a+b)*\sinh(dx+c)^4+(2*a+b)*\cosh(dx+c)^2+(6*(a+b)*\cosh(dx+c)^2+2*a+b)*\sinh(dx+c)^2+2*(2*(a+b)*\cosh(dx+c)^3+(2*a+b)*\cosh(dx+c))*\sinh(dx+c)+a)*\sqrt{a+b}*\sqrt{(a*\cosh(dx+c)+b*\sinh(dx+c))/\cosh(dx+c)})+8*((a^2+2*a*b+b^2)*\cosh(dx+c)^3+(a^2+a*b)*\cosh(dx+c))*\sinh(dx+c)+(a+b)*\sqrt{a-b}*\log(((2*a^2-b^2)*\cosh(dx+c)^4+4*(2*a^2-b^2)*\cosh(dx+c)*\sinh(dx+c)^3+(2*a^2-b^2)*\sinh(dx+c) \end{aligned}$$

$$\begin{aligned}
&^4 + 4*(a^2 - a*b)*\cosh(d*x + c)^2 + 2*(3*(2*a^2 - b^2)*\cosh(d*x + c)^2 + 2 \\
&*a^2 - 2*a*b)*\sinh(d*x + c)^2 + 2*a^2 - 4*a*b + 2*b^2 - 2*(a*\cosh(d*x + c)^ \\
&4 + 4*a*\cosh(d*x + c)*\sinh(d*x + c)^3 + a*\sinh(d*x + c)^4 + (2*a - b)*\cosh(\\
&d*x + c)^2 + (6*a*\cosh(d*x + c)^2 + 2*a - b)*\sinh(d*x + c)^2 + 2*(2*a*\cosh(\\
&d*x + c)^3 + (2*a - b)*\cosh(d*x + c))*\sinh(d*x + c) + a - b)*\sqrt{a - b)*\sqrt{ \\
&rt((a*\cosh(d*x + c) + b*\sinh(d*x + c))/\cosh(d*x + c))} + 4*((2*a^2 - b^2)*\co \\
&sh(d*x + c)^3 + 2*(a^2 - a*b)*\cosh(d*x + c))*\sinh(d*x + c))/(\cosh(d*x + c)^ \\
&4 + 4*\cosh(d*x + c)^3*\sinh(d*x + c) + 6*\cosh(d*x + c)^2*\sinh(d*x + c)^2 + 4 \\
&*cosh(d*x + c)*\sinh(d*x + c)^3 + \sinh(d*x + c)^4))/((a^2 - b^2)*d), -1/4*(\\
&2*(a - b)*\sqrt{-a - b)*\arctan(((a + b)*\cosh(d*x + c)^2 + 2*(a + b)*\cosh(d*x \\
&+ c)*\sinh(d*x + c) + (a + b)*\sinh(d*x + c)^2 + a)*\sqrt{-a - b)*\sqrt{(a*\cos \\
&h(d*x + c) + b*\sinh(d*x + c))/\cosh(d*x + c)))/((a^2 + 2*a*b + b^2)*\cosh(d*x \\
&+ c)^2 + 2*(a^2 + 2*a*b + b^2)*\cosh(d*x + c)*\sinh(d*x + c) + (a^2 + 2*a*b + \\
&b^2)*\sinh(d*x + c)^2 + a^2 - b^2)) - (a + b)*\sqrt{a - b)*\log(((2*a^2 - b^2) \\
&)*\cosh(d*x + c)^4 + 4*(2*a^2 - b^2)*\cosh(d*x + c)*\sinh(d*x + c)^3 + (2*a^2 \\
&- b^2)*\sinh(d*x + c)^4 + 4*(a^2 - a*b)*\cosh(d*x + c)^2 + 2*(3*(2*a^2 - b^2) \\
&)*\cosh(d*x + c)^2 + 2*a^2 - 2*a*b)*\sinh(d*x + c)^2 + 2*a^2 - 4*a*b + 2*b^2 - \\
&2*(a*\cosh(d*x + c)^4 + 4*a*\cosh(d*x + c)*\sinh(d*x + c)^3 + a*\sinh(d*x + c) \\
&^4 + (2*a - b)*\cosh(d*x + c)^2 + (6*a*\cosh(d*x + c)^2 + 2*a - b)*\sinh(d*x + \\
&c)^2 + 2*(2*a*\cosh(d*x + c)^3 + (2*a - b)*\cosh(d*x + c))*\sinh(d*x + c) + a \\
&- b)*\sqrt{a - b)*\sqrt{(a*\cosh(d*x + c) + b*\sinh(d*x + c))/\cosh(d*x + c)} + \\
&4*((2*a^2 - b^2)*\cosh(d*x + c)^3 + 2*(a^2 - a*b)*\cosh(d*x + c))*\sinh(d*x + \\
&c))/(\cosh(d*x + c)^4 + 4*\cosh(d*x + c)^3*\sinh(d*x + c) + 6*\cosh(d*x + c)^2 \\
&)*\sinh(d*x + c)^2 + 4*\cosh(d*x + c)*\sinh(d*x + c)^3 + \sinh(d*x + c)^4))/((a \\
&^2 - b^2)*d), -1/4*(2*(a + b)*\sqrt{-a + b)*\arctan(-(a*\cosh(d*x + c)^2 + 2*a \\
&)*\cosh(d*x + c)*\sinh(d*x + c) + a*\sinh(d*x + c)^2 + a - b)*\sqrt{-a + b)*\sqrt{ \\
&((a*\cosh(d*x + c) + b*\sinh(d*x + c))/\cosh(d*x + c)))/((a^2 - b^2)*\cosh(d*x + \\
&c)^2 + 2*(a^2 - b^2)*\cosh(d*x + c)*\sinh(d*x + c) + (a^2 - b^2)*\sinh(d*x + \\
&c)^2 + a^2 - 2*a*b + b^2)) - \sqrt{a + b)*(a - b)*\log(2*(a^2 + 2*a*b + b^2)* \\
&\cosh(d*x + c)^4 + 8*(a^2 + 2*a*b + b^2)*\cosh(d*x + c)*\sinh(d*x + c)^3 + 2*(\\
&a^2 + 2*a*b + b^2)*\sinh(d*x + c)^4 + 4*(a^2 + a*b)*\cosh(d*x + c)^2 + 4*(3*(\\
&a^2 + 2*a*b + b^2)*\cosh(d*x + c)^2 + a^2 + a*b)*\sinh(d*x + c)^2 + 2*a^2 - b \\
&^2 + 2*((a + b)*\cosh(d*x + c)^4 + 4*(a + b)*\cosh(d*x + c)*\sinh(d*x + c)^3 + \\
&(a + b)*\sinh(d*x + c)^4 + (2*a + b)*\cosh(d*x + c)^2 + (6*(a + b)*\cosh(d*x \\
&+ c)^2 + 2*a + b)*\sinh(d*x + c)^2 + 2*(2*(a + b)*\cosh(d*x + c)^3 + (2*a + b) \\
&)*\cosh(d*x + c))*\sinh(d*x + c) + a)*\sqrt{a + b)*\sqrt{(a*\cosh(d*x + c) + b*\sinh \\
&(d*x + c))/\cosh(d*x + c)} + 8*((a^2 + 2*a*b + b^2)*\cosh(d*x + c)^3 + (a^2 \\
&+ a*b)*\cosh(d*x + c))*\sinh(d*x + c))/((a^2 - b^2)*d), -1/2*((a + b)*\sqrt{ \\
&-a + b)*\arctan(-(a*\cosh(d*x + c)^2 + 2*a*\cosh(d*x + c)*\sinh(d*x + c) + a*s \\
&\sinh(d*x + c)^2 + a - b)*\sqrt{-a + b)*\sqrt{(a*\cosh(d*x + c) + b*\sinh(d*x + c) \\
&))/\cosh(d*x + c)))/((a^2 - b^2)*\cosh(d*x + c)^2 + 2*(a^2 - b^2)*\cosh(d*x + c) \\
&)*\sinh(d*x + c) + (a^2 - b^2)*\sinh(d*x + c)^2 + a^2 - 2*a*b + b^2)) + (a - \\
&b)*\sqrt{-a - b)*\arctan(((a + b)*\cosh(d*x + c)^2 + 2*(a + b)*\cosh(d*x + c)*\sinh \\
&(d*x + c) + (a + b)*\sinh(d*x + c)^2 + a)*\sqrt{-a - b)*\sqrt{(a*\cosh(d*x + c) \\
&+ b*\sinh(d*x + c))/\cosh(d*x + c)))/((a^2 + 2*a*b + b^2)*\cosh(d*x + c)^2
\end{aligned}$$

+ 2*(a^2 + 2*a*b + b^2)*cosh(d*x + c)*sinh(d*x + c) + (a^2 + 2*a*b + b^2)*sinh(d*x + c)^2 + a^2 - b^2))/((a^2 - b^2)*d]

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{a + b \tanh(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+b*tanh(d*x+c))**(1/2),x)

[Out] Integral(1/sqrt(a + b*tanh(c + d*x)), x)

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+b*tanh(d*x+c))^(1/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP UT:sage2:=int(sage0,sageVARx):;OUTPUT:index.cc index_m operator + Error: Bad Argument Value

Mupad [B]

time = 1.39, size = 240, normalized size = 3.24

$$\frac{\operatorname{atanh}\left(\frac{(a d^3 + b d^3) \sqrt{a + b \tanh(c + dx)}}{b d^3 \sqrt{a + b}} - \frac{16 a b^2 \sqrt{a + b \tanh(c + dx)}}{\left(\frac{16 b^4 d^3}{a d^3 + b d^3} + \frac{16 a b^3 d^3}{a d^3 + b d^3}\right) \sqrt{a + b}}\right)}{d \sqrt{a + b}} + \frac{\operatorname{atanh}\left(\frac{16 a b^2 \sqrt{a + b \tanh(c + dx)}}{\left(\frac{16 b^4 d^3}{a d^3 - b d^3} - \frac{16 a b^3 d^3}{a d^3 - b d^3}\right) \sqrt{a - b}} + \frac{(a d^3 - b d^3) \sqrt{a + b \tanh(c + dx)}}{b d^3 \sqrt{a - b}}\right)}{d \sqrt{a - b}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a + b*tanh(c + d*x))^(1/2),x)

[Out] atanh(((a*d^3 + b*d^3)*(a + b*tanh(c + d*x))^(1/2))/(b*d^3*(a + b)^(1/2)) - (16*a*b^2*(a + b*tanh(c + d*x))^(1/2))/(((16*b^4*d^3)/(a*d^3 + b*d^3) + (16*a*b^3*d^3)/(a*d^3 + b*d^3))*(a + b)^(1/2)))/(d*(a + b)^(1/2)) + atanh((16*a*b^2*(a + b*tanh(c + d*x))^(1/2))/(((16*b^4*d^3)/(a*d^3 - b*d^3) - (16*a*b^3*d^3)/(a*d^3 - b*d^3))*(a - b)^(1/2)) + ((a*d^3 - b*d^3)*(a + b*tanh(c + d*x))^(1/2))/(b*d^3*(a - b)^(1/2)))/(d*(a - b)^(1/2))

$$3.69 \quad \int \frac{\sinh^4(x)}{1+\tanh(x)} dx$$

Optimal. Leaf size=60

$$\frac{x}{16} + \frac{1}{32(1-\tanh(x))^2} - \frac{1}{8(1-\tanh(x))} - \frac{1}{24(1+\tanh(x))^3} + \frac{5}{32(1+\tanh(x))^2} - \frac{3}{16(1+\tanh(x))}$$

[Out] 1/16*x+1/32/(1-tanh(x))^2-1/8/(1-tanh(x))-1/24/(1+tanh(x))^3+5/32/(1+tanh(x))^2-3/16/(1+tanh(x))

Rubi [A]

time = 0.05, antiderivative size = 60, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {3597, 862, 90, 213}

$$\frac{x}{16} - \frac{1}{8(1-\tanh(x))} - \frac{3}{16(\tanh(x)+1)} + \frac{1}{32(1-\tanh(x))^2} + \frac{5}{32(\tanh(x)+1)^2} - \frac{1}{24(\tanh(x)+1)^3}$$

Antiderivative was successfully verified.

[In] Int[Sinh[x]^4/(1 + Tanh[x]), x]

[Out] x/16 + 1/(32*(1 - Tanh[x])^2) - 1/(8*(1 - Tanh[x])) - 1/(24*(1 + Tanh[x])^3) + 5/(32*(1 + Tanh[x])^2) - 3/(16*(1 + Tanh[x]))

Rule 90

Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && IntegersQ[m, n] && (IntegerQ[p] || (GtQ[m, 0] && GeQ[n, -1]))

Rule 213

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])

Rule 862

Int[((d_) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))^(n_)*((a_) + (c_.)*(x_)^2)^(p_.), x_Symbol] :> Int[(d + e*x)^(m + p)*(f + g*x)^n*(a/d + (c/e)*x)^p, x] /; FreeQ[{a, c, d, e, f, g, m, n}, x] && NeQ[e*f - d*g, 0] && EqQ[c*d^2 + a*e^2, 0] && (IntegerQ[p] || (GtQ[a, 0] && GtQ[d, 0] && EqQ[m + p, 0]))

Rule 3597

```
Int[sin[(e_.) + (f_.)*(x_)]^(m_)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_
), x_Symbol] :> Dist[b/f, Subst[Int[x^m*((a + x)^n/(b^2 + x^2)^(m/2 + 1)),
x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, n}, x] && IntegerQ[m/2]
```

Rubi steps

$$\begin{aligned} \int \frac{\sinh^4(x)}{1 + \tanh(x)} dx &= -\text{Subst}\left(\int \frac{x^4}{(1+x)(-1+x^2)^3} dx, x, \tanh(x)\right) \\ &= -\text{Subst}\left(\int \frac{x^4}{(-1+x)^3(1+x)^4} dx, x, \tanh(x)\right) \\ &= -\text{Subst}\left(\int \left(\frac{1}{16(-1+x)^3} + \frac{1}{8(-1+x)^2} - \frac{1}{8(1+x)^4} + \frac{5}{16(1+x)^3} - \frac{3}{16(1+x)^2} + \frac{1}{16}\right) dx, x, \tanh(x)\right) \\ &= \frac{1}{32(1 - \tanh(x))^2} - \frac{1}{8(1 - \tanh(x))} - \frac{1}{24(1 + \tanh(x))^3} + \frac{5}{32(1 + \tanh(x))^2} - \frac{1}{16(1 + \tanh(x))} \\ &= \frac{x}{16} + \frac{1}{32(1 - \tanh(x))^2} - \frac{1}{8(1 - \tanh(x))} - \frac{1}{24(1 + \tanh(x))^3} + \frac{5}{32(1 + \tanh(x))^2} - \frac{1}{16(1 + \tanh(x))} \end{aligned}$$

Mathematica [A]

time = 0.06, size = 42, normalized size = 0.70

$$\frac{1}{192}(12x - 15 \cosh(2x) + 6 \cosh(4x) - \cosh(6x) - 3 \sinh(2x) - 3 \sinh(4x) + \sinh(6x))$$

Antiderivative was successfully verified.

```
[In] Integrate[Sinh[x]^4/(1 + Tanh[x]),x]
```

```
[Out] (12*x - 15*Cosh[2*x] + 6*Cosh[4*x] - Cosh[6*x] - 3*Sinh[2*x] - 3*Sinh[4*x]
+ Sinh[6*x])/192
```

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 97 vs. 2(48) = 96.

time = 0.52, size = 98, normalized size = 1.63

method	result
risch	$\frac{x}{16} + \frac{e^{4x}}{128} - \frac{3e^{2x}}{64} - \frac{e^{-2x}}{32} + \frac{3e^{-4x}}{128} - \frac{e^{-6x}}{192}$
default	$-\frac{1}{3(\tanh(\frac{x}{2})+1)^6} + \frac{1}{(\tanh(\frac{x}{2})+1)^5} - \frac{7}{8(\tanh(\frac{x}{2})+1)^4} + \frac{1}{12(\tanh(\frac{x}{2})+1)^3} + \frac{1}{8(\tanh(\frac{x}{2})+1)^2} + \frac{\ln(\tanh(\frac{x}{2})+1)}{16} + \frac{1}{8(\tanh(\frac{x}{2})+1)}$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sinh(x)^4/(1+tanh(x)),x,method=_RETURNVERBOSE)
```

[Out] $-1/3/(\tanh(1/2*x)+1)^6+1/(\tanh(1/2*x)+1)^5-7/8/(\tanh(1/2*x)+1)^4+1/12/(\tanh(1/2*x)+1)^3+1/8/(\tanh(1/2*x)+1)^2+1/16*\ln(\tanh(1/2*x)+1)+1/8/(\tanh(1/2*x)-1)^4+1/4/(\tanh(1/2*x)-1)^3-1/8/(\tanh(1/2*x)-1)-1/16*\ln(\tanh(1/2*x)-1)$

Maxima [A]

time = 0.25, size = 36, normalized size = 0.60

$$-\frac{1}{128} (6e^{(-2x)} - 1)e^{(4x)} + \frac{1}{16} x - \frac{1}{32} e^{(-2x)} + \frac{3}{128} e^{(-4x)} - \frac{1}{192} e^{(-6x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sinh(x)^4/(1+tanh(x)),x, algorithm="maxima")`

[Out] $-1/128*(6*e^{(-2*x)} - 1)*e^{(4*x)} + 1/16*x - 1/32*e^{(-2*x)} + 3/128*e^{(-4*x)} - 1/192*e^{(-6*x)}$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 92 vs. $2(44) = 88$.

time = 0.41, size = 92, normalized size = 1.53

$$\frac{\cosh(x)^5 + 5 \cosh(x) \sinh(x)^4 + 5 \sinh(x)^5 + (50 \cosh(x)^2 - 27) \sinh(x)^3 - 9 \cosh(x)^3 + (10 \cosh(x)^3 - 27 \cosh(x)) \sinh(x)^2 + 12(2x - 1) \cosh(x) + (25 \cosh(x)^4 - 81 \cosh(x)^2 + 24x + 12) \sinh(x)}{384(\cosh(x) + \sinh(x))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sinh(x)^4/(1+tanh(x)),x, algorithm="fricas")`

[Out] $1/384*(\cosh(x)^5 + 5*\cosh(x)*\sinh(x)^4 + 5*\sinh(x)^5 + (50*\cosh(x)^2 - 27)*\sinh(x)^3 - 9*\cosh(x)^3 + (10*\cosh(x)^3 - 27*\cosh(x))*\sinh(x)^2 + 12*(2*x - 1)*\cosh(x) + (25*\cosh(x)^4 - 81*\cosh(x)^2 + 24*x + 12)*\sinh(x))/(\cosh(x) + \sinh(x))$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sinh^4(x)}{\tanh(x) + 1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sinh(x)**4/(1+tanh(x)),x)`

[Out] `Integral(sinh(x)**4/(tanh(x) + 1), x)`

Giac [A]

time = 0.41, size = 42, normalized size = 0.70

$$-\frac{1}{384} (22e^{(6x)} + 12e^{(4x)} - 9e^{(2x)} + 2)e^{(-6x)} + \frac{1}{16} x + \frac{1}{128} e^{(4x)} - \frac{3}{64} e^{(2x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sinh(x)^4/(1+tanh(x)),x, algorithm="giac")

[Out] $-1/384*(22*e^{6*x} + 12*e^{4*x} - 9*e^{2*x} + 2)*e^{-6*x} + 1/16*x + 1/128*e^{4*x} - 3/64*e^{2*x}$

Mupad [B]

time = 1.31, size = 34, normalized size = 0.57

$$\frac{x}{16} - \frac{e^{-2x}}{32} - \frac{3e^{2x}}{64} + \frac{3e^{-4x}}{128} + \frac{e^{4x}}{128} - \frac{e^{-6x}}{192}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sinh(x)^4/(tanh(x) + 1),x)

[Out] $x/16 - \exp(-2*x)/32 - (3*\exp(2*x))/64 + (3*\exp(-4*x))/128 + \exp(4*x)/128 - \exp(-6*x)/192$

3.70 $\int \frac{\sinh^3(x)}{1+\tanh(x)} dx$

Optimal. Leaf size=25

$$-\frac{1}{3} \cosh^3(x) + \frac{\cosh^5(x)}{5} - \frac{\sinh^5(x)}{5}$$

[Out] $-1/3*\cosh(x)^3+1/5*\cosh(x)^5-1/5*\sinh(x)^5$

Rubi [A]

time = 0.12, antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 7, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.636$, Rules used = {3599, 3187, 3186, 2645, 14, 2644, 30}

$$-\frac{\sinh^5(x)}{5} + \frac{\cosh^5(x)}{5} - \frac{\cosh^3(x)}{3}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sinh}[x]^3/(1 + \text{Tanh}[x]), x]$

[Out] $-1/3*\text{Cosh}[x]^3 + \text{Cosh}[x]^5/5 - \text{Sinh}[x]^5/5$

Rule 14

$\text{Int}[(u_*)((c_)*(x_))^{(m_)}], x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c*x)^m*u, x], x] /;$ FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_)*(v_)] /; FreeQ[{a, b}, x] && InverseFunctionQ[v]

Rule 30

$\text{Int}[(x_)^{(m_)}], x_Symbol] \rightarrow \text{Simp}[x^{(m+1)}/(m+1), x] /;$ FreeQ[m, x] && NeQ[m, -1]

Rule 2644

$\text{Int}[\cos[(e_)+(f_)*(x_)]^{(n_)}*((a_)*\sin[(e_)+(f_)*(x_)]^{(m_)}], x_Symbol] \rightarrow \text{Dist}[1/(a*f), \text{Subst}[\text{Int}[x^m*(1-x^2/a^2)^{((n-1)/2)}, x], x, a*\text{Sin}[e+f*x]], x] /;$ FreeQ[{a, e, f, m}, x] && IntegerQ[(n-1)/2] && !(IntegerQ[(m-1)/2] && LtQ[0, m, n])

Rule 2645

$\text{Int}[(\cos[(e_)+(f_)*(x_)]*(a_))^{(m_)}*\sin[(e_)+(f_)*(x_)]^{(n_)}], x_Symbol] \rightarrow \text{Dist}[-(a*f)^{-1}, \text{Subst}[\text{Int}[x^m*(1-x^2/a^2)^{((n-1)/2)}, x], x, a*\text{Cos}[e+f*x]], x] /;$ FreeQ[{a, e, f, m}, x] && IntegerQ[(n-1)/2] && !(IntegerQ[(m-1)/2] && GtQ[m, 0] && LeQ[m, n])

Rule 3186

```
Int[cos[(c_.) + (d_.)*(x_)]^(m_.)*sin[(c_.) + (d_.)*(x_)]^(n_.)*(cos[(c_.)
+ (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)]^(p_.), x_Symbol] := In
t[ExpandTrig[cos[c + d*x]^m*sin[c + d*x]^n*(a*cos[c + d*x] + b*sin[c + d*x]
)^p, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && IGtQ[p, 0]
```

Rule 3187

```
Int[cos[(c_.) + (d_.)*(x_)]^(m_.)*sin[(c_.) + (d_.)*(x_)]^(n_.)*(cos[(c_.)
+ (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)]^(p_.), x_Symbol] := Dis
t[a^p*b^p, Int[(Cos[c + d*x]^m*Sin[c + d*x]^n)/(b*Cos[c + d*x] + a*Sin[c +
d*x])^p, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[a^2 + b^2, 0] && ILt
Q[p, 0]
```

Rule 3599

```
Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]^(n
_.), x_Symbol] := Int[Sin[e + f*x]^m*((a*Cos[e + f*x] + b*Sin[e + f*x])^n/C
os[e + f*x]^n), x] /; FreeQ[{a, b, e, f}, x] && IntegerQ[(m - 1)/2] && ILtQ
[n, 0] && ((LtQ[m, 5] && GtQ[n, -4]) || (EqQ[m, 5] && EqQ[n, -1]))
```

Rubi steps

$$\begin{aligned}
\int \frac{\sinh^3(x)}{1 + \tanh(x)} dx &= \int \frac{\cosh(x) \sinh^3(x)}{\cosh(x) + \sinh(x)} dx \\
&= i \int \cosh(x) (-i \cosh(x) + i \sinh(x)) \sinh^3(x) dx \\
&= - \int (-\cosh^2(x) \sinh^3(x) + \cosh(x) \sinh^4(x)) dx \\
&= \int \cosh^2(x) \sinh^3(x) dx - \int \cosh(x) \sinh^4(x) dx \\
&= i \text{Subst} \left(\int x^4 dx, x, i \sinh(x) \right) - \text{Subst} \left(\int x^2 (1 - x^2) dx, x, \cosh(x) \right) \\
&= -\frac{1}{5} \sinh^5(x) - \text{Subst} \left(\int (x^2 - x^4) dx, x, \cosh(x) \right) \\
&= -\frac{1}{3} \cosh^3(x) + \frac{\cosh^5(x)}{5} - \frac{\sinh^5(x)}{5}
\end{aligned}$$

Mathematica [A]

time = 0.05, size = 34, normalized size = 1.36

$$\frac{1}{120} (\cosh(x) - \sinh(x)) (-20 \cosh(2x) + 4 \cosh(4x) - 10 \sinh(2x) + \sinh(4x))$$

Antiderivative was successfully verified.

[In] Integrate[Sinh[x]^3/(1 + Tanh[x]),x]

[Out] ((Cosh[x] - Sinh[x])*(-20*Cosh[2*x] + 4*Cosh[4*x] - 10*Sinh[2*x] + Sinh[4*x]))/120

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 71 vs. 2(19) = 38.

time = 0.49, size = 72, normalized size = 2.88

method	result
risch	$\frac{e^{3x}}{48} - \frac{e^x}{8} - \frac{e^{-3x}}{24} + \frac{e^{-5x}}{80}$
default	$-\frac{1}{6(\tanh(\frac{x}{2})-1)^3} - \frac{1}{4(\tanh(\frac{x}{2})-1)^2} + \frac{1}{8\tanh(\frac{x}{2})-8} + \frac{2}{5(\tanh(\frac{x}{2})+1)^5} - \frac{1}{(\tanh(\frac{x}{2})+1)^4} + \frac{2}{3(\tanh(\frac{x}{2})+1)^3} - \frac{1}{8(\tanh(\frac{x}{2})+1)}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sinh(x)^3/(1+tanh(x)),x,method=_RETURNVERBOSE)

[Out] -1/6/(tanh(1/2*x)-1)^3-1/4/(tanh(1/2*x)-1)^2+1/8/(tanh(1/2*x)-1)+2/5/(tanh(1/2*x)+1)^5-1/(tanh(1/2*x)+1)^4+2/3/(tanh(1/2*x)+1)^3-1/8/(tanh(1/2*x)+1)

Maxima [A]

time = 0.26, size = 27, normalized size = 1.08

$$-\frac{1}{48} (6e^{(-2x)} - 1)e^{(3x)} - \frac{1}{24} e^{(-3x)} + \frac{1}{80} e^{(-5x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sinh(x)^3/(1+tanh(x)),x, algorithm="maxima")

[Out] -1/48*(6*e^(-2*x) - 1)*e^(3*x) - 1/24*e^(-3*x) + 1/80*e^(-5*x)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 56 vs. 2(19) = 38.

time = 0.44, size = 56, normalized size = 2.24

$$\frac{\cosh(x)^4 + \cosh(x)\sinh(x)^3 + \sinh(x)^4 + (6\cosh(x)^2 - 5)\sinh(x)^2 - 5\cosh(x)^2 + (\cosh(x)^3 - 5\cosh(x))\sinh(x)}{30(\cosh(x) + \sinh(x))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sinh(x)^3/(1+tanh(x)),x, algorithm="fricas")

[Out] 1/30*(cosh(x)^4 + cosh(x)*sinh(x)^3 + sinh(x)^4 + (6*cosh(x)^2 - 5)*sinh(x)^2 - 5*cosh(x)^2 + (cosh(x)^3 - 5*cosh(x))*sinh(x))/(cosh(x) + sinh(x))

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 134 vs. $2(19) = 38$.

time = 0.37, size = 134, normalized size = 5.36

$$\frac{3 \sinh^3(x) \tanh(x)}{15 \tanh(x) + 15} - \frac{3 \sinh^3(x)}{15 \tanh(x) + 15} + \frac{6 \sinh^2(x) \cosh(x) \tanh(x)}{15 \tanh(x) + 15} + \frac{9 \sinh^2(x) \cosh(x)}{15 \tanh(x) + 15} - \frac{6 \sinh(x) \cosh^2(x) \tanh(x)}{15 \tanh(x) + 15} + \frac{6 \sinh(x) \cosh^2(x)}{15 \tanh(x) + 15} - \frac{8 \cosh^3(x) \tanh(x)}{15 \tanh(x) + 15} - \frac{2 \cosh^3(x)}{15 \tanh(x) + 15}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sinh(x)**3/(1+tanh(x)),x)

[Out] $3*\sinh(x)**3*\tanh(x)/(15*\tanh(x) + 15) - 3*\sinh(x)**3/(15*\tanh(x) + 15) + 6*\sinh(x)**2*\cosh(x)*\tanh(x)/(15*\tanh(x) + 15) + 9*\sinh(x)**2*\cosh(x)/(15*\tanh(x) + 15) - 6*\sinh(x)*\cosh(x)**2*\tanh(x)/(15*\tanh(x) + 15) + 6*\sinh(x)*\cosh(x)**2/(15*\tanh(x) + 15) - 8*\cosh(x)**3*\tanh(x)/(15*\tanh(x) + 15) - 2*\cosh(x)**3/(15*\tanh(x) + 15)$

Giac [A]

time = 0.39, size = 25, normalized size = 1.00

$$-\frac{1}{240} (10 e^{2x} - 3) e^{(-5x)} + \frac{1}{48} e^{(3x)} - \frac{1}{8} e^x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sinh(x)^3/(1+tanh(x)),x, algorithm="giac")

[Out] $-1/240*(10*e^{2*x} - 3)*e^{(-5*x)} + 1/48*e^{(3*x)} - 1/8*e^x$

Mupad [B]

time = 1.18, size = 23, normalized size = 0.92

$$\frac{e^{3x}}{48} - \frac{e^{-3x}}{24} + \frac{e^{-5x}}{80} - \frac{e^x}{8}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sinh(x)^3/(tanh(x) + 1),x)

[Out] $\exp(3*x)/48 - \exp(-3*x)/24 + \exp(-5*x)/80 - \exp(x)/8$

3.71 $\int \frac{\sinh^2(x)}{1+\tanh(x)} dx$

Optimal. Leaf size=38

$$-\frac{x}{8} + \frac{1}{8(1-\tanh(x))} - \frac{1}{8(1+\tanh(x))^2} + \frac{1}{4(1+\tanh(x))}$$

[Out] $-1/8*x+1/8/(1-\tanh(x))-1/8/(1+\tanh(x))^2+1/4/(1+\tanh(x))$

Rubi [A]

time = 0.04, antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {3597, 862, 90, 213}

$$-\frac{x}{8} + \frac{1}{8(1-\tanh(x))} + \frac{1}{4(\tanh(x)+1)} - \frac{1}{8(\tanh(x)+1)^2}$$

Antiderivative was successfully verified.

[In] Int[Sinh[x]^2/(1 + Tanh[x]),x]

[Out] $-1/8*x + 1/(8*(1 - Tanh[x])) - 1/(8*(1 + Tanh[x])^2) + 1/(4*(1 + Tanh[x]))$

Rule 90

Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && IntegersQ[m, n] && (IntegerQ[p] || (GtQ[m, 0] && GeQ[n, -1]))

Rule 213

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1)*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])

Rule 862

Int[((d_) + (e_.)*(x_))^(m_.)*((f_.) + (g_.)*(x_))^(n_.)*((a_) + (c_.)*(x_)^2)^(p_.), x_Symbol] :> Int[(d + e*x)^(m + p)*(f + g*x)^n*(a/d + (c/e)*x)^p, x] /; FreeQ[{a, c, d, e, f, g, m, n}, x] && NeQ[e*f - d*g, 0] && EqQ[c*d^2 + a*e^2, 0] && (IntegerQ[p] || (GtQ[a, 0] && GtQ[d, 0] && EqQ[m + p, 0]))

Rule 3597

Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] :> Dist[b/f, Subst[Int[x^m*((a + x)^n/(b^2 + x^2)^(m/2 + 1)),

$x], x, b*\text{Tan}[e + f*x]], x] /; \text{FreeQ}[\{a, b, e, f, n\}, x] \ \&\& \ \text{IntegerQ}[m/2]$

Rubi steps

$$\begin{aligned} \int \frac{\sinh^2(x)}{1 + \tanh(x)} dx &= \text{Subst} \left(\int \frac{x^2}{(1+x)(-1+x^2)^2} dx, x, \tanh(x) \right) \\ &= \text{Subst} \left(\int \frac{x^2}{(-1+x)^2(1+x)^3} dx, x, \tanh(x) \right) \\ &= \text{Subst} \left(\int \left(\frac{1}{8(-1+x)^2} + \frac{1}{4(1+x)^3} - \frac{1}{4(1+x)^2} + \frac{1}{8(-1+x^2)} \right) dx, x, \tanh(x) \right) \\ &= \frac{1}{8(1 - \tanh(x))} - \frac{1}{8(1 + \tanh(x))^2} + \frac{1}{4(1 + \tanh(x))} + \frac{1}{8} \text{Subst} \left(\int \frac{1}{-1+x^2} dx, x, \tanh(x) \right) \\ &= -\frac{x}{8} + \frac{1}{8(1 - \tanh(x))} - \frac{1}{8(1 + \tanh(x))^2} + \frac{1}{4(1 + \tanh(x))} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 24, normalized size = 0.63

$$\frac{1}{32}(-4x + 4 \cosh(2x) - \cosh(4x) + \sinh(4x))$$

Antiderivative was successfully verified.

[In] Integrate[Sinh[x]^2/(1 + Tanh[x]),x]

[Out] (-4*x + 4*Cosh[2*x] - Cosh[4*x] + Sinh[4*x])/32

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 67 vs. 2(30) = 60.

time = 0.37, size = 68, normalized size = 1.79

method	result
risch	$-\frac{x}{8} + \frac{e^{2x}}{16} + \frac{e^{-2x}}{16} - \frac{e^{-4x}}{32}$
default	$\frac{1}{4(\tanh(\frac{x}{2})-1)^2} + \frac{1}{4 \tanh(\frac{x}{2})-4} + \frac{\ln(\tanh(\frac{x}{2})-1)}{8} - \frac{1}{2(\tanh(\frac{x}{2})+1)^4} + \frac{1}{(\tanh(\frac{x}{2})+1)^3} - \frac{1}{2(\tanh(\frac{x}{2})+1)^2} - \frac{\ln(\tanh(\frac{x}{2}))}{8}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sinh(x)^2/(1+tanh(x)),x,method=_RETURNVERBOSE)

[Out] 1/4/(tanh(1/2*x)-1)^2+1/4/(tanh(1/2*x)-1)+1/8*ln(tanh(1/2*x)-1)-1/2/(tanh(1/2*x)+1)^4+1/(tanh(1/2*x)+1)^3-1/2/(tanh(1/2*x)+1)^2-1/8*ln(tanh(1/2*x)+1)

Maxima [A]

time = 0.28, size = 22, normalized size = 0.58

$$-\frac{1}{8}x + \frac{1}{16}e^{(2x)} + \frac{1}{16}e^{(-2x)} - \frac{1}{32}e^{(-4x)}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sinh(x)^2/(1+tanh(x)),x, algorithm="maxima")``[Out] -1/8*x + 1/16*e^(2*x) + 1/16*e^(-2*x) - 1/32*e^(-4*x)`**Fricas [A]**

time = 0.33, size = 51, normalized size = 1.34

$$\frac{\cosh(x)^3 + 3 \cosh(x) \sinh(x)^2 + 3 \sinh(x)^3 - 2(2x - 1) \cosh(x) + (9 \cosh(x)^2 - 4x - 2) \sinh(x)}{32(\cosh(x) + \sinh(x))}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sinh(x)^2/(1+tanh(x)),x, algorithm="fricas")``[Out] 1/32*(cosh(x)^3 + 3*cosh(x)*sinh(x)^2 + 3*sinh(x)^3 - 2*(2*x - 1)*cosh(x) + (9*cosh(x)^2 - 4*x - 2)*sinh(x))/(cosh(x) + sinh(x))`**Sympy [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sinh^2(x)}{\tanh(x) + 1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sinh(x)**2/(1+tanh(x)),x)``[Out] Integral(sinh(x)**2/(tanh(x) + 1), x)`**Giac [A]**

time = 0.41, size = 30, normalized size = 0.79

$$\frac{1}{32} (3e^{(4x)} + 2e^{(2x)} - 1)e^{(-4x)} - \frac{1}{8}x + \frac{1}{16}e^{(2x)}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sinh(x)^2/(1+tanh(x)),x, algorithm="giac")``[Out] 1/32*(3*e^(4*x) + 2*e^(2*x) - 1)*e^(-4*x) - 1/8*x + 1/16*e^(2*x)`**Mupad [B]**

time = 1.14, size = 22, normalized size = 0.58

$$\frac{e^{-2x}}{16} - \frac{x}{8} + \frac{e^{2x}}{16} - \frac{e^{-4x}}{32}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sinh(x)^2/(tanh(x) + 1),x)
```

```
[Out] exp(-2*x)/16 - x/8 + exp(2*x)/16 - exp(-4*x)/32
```

3.72 $\int \frac{\sinh(x)}{1+\tanh(x)} dx$

Optimal. Leaf size=17

$$\frac{\cosh^3(x)}{3} - \frac{\sinh^3(x)}{3}$$

[Out] 1/3*cosh(x)^3-1/3*sinh(x)^3

Rubi [A]

time = 0.08, antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 6, integrand size = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {3599, 3187, 3186, 2645, 30, 2644}

$$\frac{\cosh^3(x)}{3} - \frac{\sinh^3(x)}{3}$$

Antiderivative was successfully verified.

[In] Int[Sinh[x]/(1 + Tanh[x]),x]

[Out] Cosh[x]^3/3 - Sinh[x]^3/3

Rule 30

Int[(x_)^(m_), x_Symbol] :> Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2644

Int[cos[(e_) + (f_)*(x_)]^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] :> Dist[1/(a*f), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && LtQ[0, m, n])

Rule 2645

Int[(cos[(e_) + (f_)*(x_)]*(a_))^(m_)*sin[(e_) + (f_)*(x_)]^(n_), x_Symbol] :> Dist[-(a*f)^(-1), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])

Rule 3186

Int[cos[(c_) + (d_)*(x_)]^(m_)*sin[(c_) + (d_)*(x_)]^(n_)*(cos[(c_) + (d_)*(x_)]*(a_) + (b_)*sin[(c_) + (d_)*(x_)])^(p_), x_Symbol] :> Int[ExpandTrig[cos[c + d*x]^m*sin[c + d*x]^n*(a*cos[c + d*x] + b*sin[c + d*x])^p, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && IGtQ[p, 0]

Rule 3187

```
Int[cos[(c_.) + (d_.)*(x_.)]^(m_.)*sin[(c_.) + (d_.)*(x_.)]^(n_.)*(cos[(c_.)
+ (d_.)*(x_.)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_.)])^(p_), x_Symbol] := Dis
t[a^p*b^p, Int[(Cos[c + d*x]^m*Sin[c + d*x]^n)/(b*Cos[c + d*x] + a*Sin[c +
d*x])^p, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[a^2 + b^2, 0] && ILt
Q[p, 0]
```

Rule 3599

```
Int[sin[(e_.) + (f_.)*(x_.)]^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(n
_.), x_Symbol] := Int[Sin[e + f*x]^m*((a*Cos[e + f*x] + b*Sin[e + f*x])^n/C
os[e + f*x]^n), x] /; FreeQ[{a, b, e, f}, x] && IntegerQ[(m - 1)/2] && ILtQ
[n, 0] && ((LtQ[m, 5] && GtQ[n, -4]) || (EqQ[m, 5] && EqQ[n, -1]))
```

Rubi steps

$$\begin{aligned}
\int \frac{\sinh(x)}{1 + \tanh(x)} dx &= \int \frac{\cosh(x) \sinh(x)}{\cosh(x) + \sinh(x)} dx \\
&= i \int \cosh(x) (-i \cosh(x) + i \sinh(x)) \sinh(x) dx \\
&= \int (\cosh^2(x) \sinh(x) - \cosh(x) \sinh^2(x)) dx \\
&= \int \cosh^2(x) \sinh(x) dx - \int \cosh(x) \sinh^2(x) dx \\
&= -\left(i \text{Subst} \left(\int x^2 dx, x, i \sinh(x) \right) \right) + \text{Subst} \left(\int x^2 dx, x, \cosh(x) \right) \\
&= \frac{\cosh^3(x)}{3} - \frac{\sinh^3(x)}{3}
\end{aligned}$$

Mathematica [A]

time = 0.01, size = 19, normalized size = 1.12

$$\frac{1}{12} (3 \cosh(x) + \cosh(3x) - 4 \sinh^3(x))$$

Antiderivative was successfully verified.

[In] Integrate[Sinh[x]/(1 + Tanh[x]),x]

[Out] (3*Cosh[x] + Cosh[3*x] - 4*Sinh[x]^3)/12

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 41 vs. 2(13) = 26.

time = 0.32, size = 42, normalized size = 2.47

method	result	size
risch	$\frac{e^x}{4} + \frac{e^{-3x}}{12}$	12
default	$\frac{2}{3(\tanh(\frac{x}{2})+1)^3} - \frac{1}{(\tanh(\frac{x}{2})+1)^2} + \frac{1}{2\tanh(\frac{x}{2})+2} - \frac{1}{2(\tanh(\frac{x}{2})-1)}$	42

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sinh(x)/(1+tanh(x)),x,method=_RETURNVERBOSE)
```

```
[Out] 2/3/(tanh(1/2*x)+1)^3-1/(tanh(1/2*x)+1)^2+1/2/(tanh(1/2*x)+1)-1/2/(tanh(1/2*x)-1)
```

Maxima [A]

time = 0.26, size = 11, normalized size = 0.65

$$\frac{1}{12} e^{(-3x)} + \frac{1}{4} e^x$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sinh(x)/(1+tanh(x)),x, algorithm="maxima")
```

```
[Out] 1/12*e^(-3*x) + 1/4*e^x
```

Fricas [A]

time = 0.41, size = 23, normalized size = 1.35

$$\frac{\cosh(x)^2 + \cosh(x) \sinh(x) + \sinh(x)^2}{3(\cosh(x) + \sinh(x))}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sinh(x)/(1+tanh(x)),x, algorithm="fricas")
```

```
[Out] 1/3*(cosh(x)^2 + cosh(x)*sinh(x) + sinh(x)^2)/(cosh(x) + sinh(x))
```

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 48 vs. $2(12) = 24$.

time = 0.19, size = 48, normalized size = 2.82

$$\frac{\sinh(x) \tanh(x)}{3 \tanh(x) + 3} - \frac{\sinh(x)}{3 \tanh(x) + 3} + \frac{2 \cosh(x) \tanh(x)}{3 \tanh(x) + 3} + \frac{\cosh(x)}{3 \tanh(x) + 3}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sinh(x)/(1+tanh(x)),x)
```

```
[Out] sinh(x)*tanh(x)/(3*tanh(x) + 3) - sinh(x)/(3*tanh(x) + 3) + 2*cosh(x)*tanh(x)/(3*tanh(x) + 3) + cosh(x)/(3*tanh(x) + 3)
```

Giac [A]

time = 0.40, size = 11, normalized size = 0.65

$$\frac{1}{12} e^{(-3x)} + \frac{1}{4} e^x$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sinh(x)/(1+tanh(x)),x, algorithm="giac")``[Out] 1/12*e^(-3*x) + 1/4*e^x`**Mupad [B]**

time = 1.12, size = 11, normalized size = 0.65

$$\frac{e^{-3x}}{12} + \frac{e^x}{4}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sinh(x)/(tanh(x) + 1),x)``[Out] exp(-3*x)/12 + exp(x)/4`

3.73 $\int \frac{\operatorname{csch}(x)}{1+\tanh(x)} dx$

Optimal. Leaf size=12

$$-\tanh^{-1}(\cosh(x)) + \cosh(x) - \sinh(x)$$

[Out] -arctanh(cosh(x))+cosh(x)-sinh(x)

Rubi [A]

time = 0.08, antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, integrand size = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.778$, Rules used = {3599, 3187, 3186, 2717, 2672, 327, 212}

$$-\sinh(x) + \cosh(x) - \tanh^{-1}(\cosh(x))$$

Antiderivative was successfully verified.

[In] Int[Csch[x]/(1 + Tanh[x]),x]

[Out] -ArcTanh[Cosh[x]] + Cosh[x] - Sinh[x]

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 327

Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 2672

Int[((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[ff/f, Subst[Int[(ff*x)^(m + n)/(a^2 - ff^2*x^2)^((n + 1)/2), x], x, a*(Sin[e + f*x]/ff)], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2]

Rule 2717

Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]

Rule 3186

```
Int[cos[(c_.) + (d_.)*(x_)]^(m_.)*sin[(c_.) + (d_.)*(x_)]^(n_.)*(cos[(c_.)
+ (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)]^(p_.), x_Symbol] := In
t[ExpandTrig[cos[c + d*x]^m*sin[c + d*x]^n*(a*cos[c + d*x] + b*sin[c + d*x]
)^p, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && IGtQ[p, 0]
```

Rule 3187

```
Int[cos[(c_.) + (d_.)*(x_)]^(m_.)*sin[(c_.) + (d_.)*(x_)]^(n_.)*(cos[(c_.)
+ (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)]^(p_.), x_Symbol] := Dis
t[a^p*b^p, Int[(Cos[c + d*x]^m*Sin[c + d*x]^n)/(b*Cos[c + d*x] + a*Sin[c +
d*x])^p, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[a^2 + b^2, 0] && ILtQ
[p, 0]
```

Rule 3599

```
Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]^(n
_.), x_Symbol] := Int[Sin[e + f*x]^m*((a*Cos[e + f*x] + b*Sin[e + f*x])^n/C
os[e + f*x]^n), x] /; FreeQ[{a, b, e, f}, x] && IntegerQ[(m - 1)/2] && ILtQ
[n, 0] && ((LtQ[m, 5] && GtQ[n, -4]) || (EqQ[m, 5] && EqQ[n, -1]))
```

Rubi steps

$$\begin{aligned}
\int \frac{\operatorname{csch}(x)}{1 + \tanh(x)} dx &= \int \frac{\operatorname{coth}(x)}{\cosh(x) + \sinh(x)} dx \\
&= i \int \operatorname{coth}(x)(-i \cosh(x) + i \sinh(x)) dx \\
&= - \int (\cosh(x) - \cosh(x) \operatorname{coth}(x)) dx \\
&= - \int \cosh(x) dx + \int \cosh(x) \operatorname{coth}(x) dx \\
&= -\sinh(x) - \operatorname{Subst}\left(\int \frac{x^2}{1-x^2} dx, x, \cosh(x)\right) \\
&= \cosh(x) - \sinh(x) - \operatorname{Subst}\left(\int \frac{1}{1-x^2} dx, x, \cosh(x)\right) \\
&= -\tanh^{-1}(\cosh(x)) + \cosh(x) - \sinh(x)
\end{aligned}$$

Mathematica [A]

time = 0.02, size = 14, normalized size = 1.17

$$\cosh(x) + \log\left(\tanh\left(\frac{x}{2}\right)\right) - \sinh(x)$$

Antiderivative was successfully verified.

[In] Integrate[Csch[x]/(1 + Tanh[x]),x]

[Out] Cosh[x] + Log[Tanh[x/2]] - Sinh[x]

Maple [A]

time = 0.56, size = 17, normalized size = 1.42

method	result	size
default	$\frac{2}{\tanh(\frac{x}{2})+1} + \ln\left(\tanh\left(\frac{x}{2}\right)\right)$	17
risch	$e^{-x} - \ln(e^x + 1) + \ln(e^x - 1)$	18

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csch(x)/(1+tanh(x)),x,method=_RETURNVERBOSE)

[Out] 2/(tanh(1/2*x)+1)+ln(tanh(1/2*x))

Maxima [A]

time = 0.26, size = 21, normalized size = 1.75

$$e^{(-x)} - \log(e^{(-x)} + 1) + \log(e^{(-x)} - 1)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csch(x)/(1+tanh(x)),x, algorithm="maxima")

[Out] e^(-x) - log(e^(-x) + 1) + log(e^(-x) - 1)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 38 vs. 2(12) = 24.

time = 0.39, size = 38, normalized size = 3.17

$$\frac{(\cosh(x) + \sinh(x)) \log(\cosh(x) + \sinh(x) + 1) - (\cosh(x) + \sinh(x)) \log(\cosh(x) + \sinh(x) - 1) - 1}{\cosh(x) + \sinh(x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csch(x)/(1+tanh(x)),x, algorithm="fricas")

[Out] -((cosh(x) + sinh(x))*log(cosh(x) + sinh(x) + 1) - (cosh(x) + sinh(x))*log(cosh(x) + sinh(x) - 1) - 1)/(cosh(x) + sinh(x))

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{csch}(x)}{\tanh(x) + 1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csch(x)/(1+tanh(x)),x)`

[Out] `Integral(csch(x)/(tanh(x) + 1), x)`

Giac [A]

time = 0.42, size = 18, normalized size = 1.50

$$e^{(-x)} - \log(e^x + 1) + \log(|e^x - 1|)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csch(x)/(1+tanh(x)),x, algorithm="giac")`

[Out] `e^(-x) - log(e^x + 1) + log(abs(e^x - 1))`

Mupad [B]

time = 0.06, size = 21, normalized size = 1.75

$$\ln(2 - 2e^x) - \ln(-2e^x - 2) + e^{-x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(sinh(x)*(tanh(x) + 1)),x)`

[Out] `log(2 - 2*exp(x)) - log(- 2*exp(x) - 2) + exp(-x)`

3.74 $\int \frac{\operatorname{csch}^2(x)}{1+\tanh(x)} dx$

Optimal. Leaf size=15

$$-\coth(x) - \log(\tanh(x)) + \log(1 + \tanh(x))$$

[Out] -coth(x)-ln(tanh(x))+ln(1+tanh(x))

Rubi [A]

time = 0.03, antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {3597, 46}

$$-\coth(x) - \log(\tanh(x)) + \log(\tanh(x) + 1)$$

Antiderivative was successfully verified.

[In] Int[Csch[x]^2/(1 + Tanh[x]), x]

[Out] -Coth[x] - Log[Tanh[x]] + Log[1 + Tanh[x]]

Rule 46

Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && ILtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m + n + 2, 0])

Rule 3597

Int[sin[(e_.) + (f_.)*(x_)]^(m_)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> Dist[b/f, Subst[Int[x^m*((a + x)^n/(b^2 + x^2)^(m/2 + 1)), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, n}, x] && IntegerQ[m/2]

Rubi steps

$$\begin{aligned} \int \frac{\operatorname{csch}^2(x)}{1+\tanh(x)} dx &= \operatorname{Subst}\left(\int \frac{1}{x^2(1+x)} dx, x, \tanh(x)\right) \\ &= \operatorname{Subst}\left(\int \left(\frac{1}{x^2} - \frac{1}{x} + \frac{1}{1+x}\right) dx, x, \tanh(x)\right) \\ &= -\coth(x) - \log(\tanh(x)) + \log(1 + \tanh(x)) \end{aligned}$$

Mathematica [A]

time = 0.02, size = 11, normalized size = 0.73

$$x - \coth(x) - \log(\sinh(x))$$

Antiderivative was successfully verified.

[In] Integrate[Csch[x]^2/(1 + Tanh[x]),x]

[Out] x - Coth[x] - Log[Sinh[x]]

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 31 vs. 2(15) = 30.

time = 0.50, size = 32, normalized size = 2.13

method	result	size
risch	$2x - \frac{2}{e^{2x}-1} - \ln(e^{2x} - 1)$	24
default	$-\frac{\tanh(\frac{x}{2})}{2} - \frac{1}{2\tanh(\frac{x}{2})} - \ln(\tanh(\frac{x}{2})) + 2\ln(\tanh(\frac{x}{2}) + 1)$	32

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csch(x)^2/(1+tanh(x)),x,method=_RETURNVERBOSE)

[Out] -1/2*tanh(1/2*x)-1/2/tanh(1/2*x)-ln(tanh(1/2*x))+2*ln(tanh(1/2*x)+1)

Maxima [A]

time = 0.26, size = 29, normalized size = 1.93

$$\frac{2}{e^{(-2x)} - 1} - \log(e^{(-x)} + 1) - \log(e^{(-x)} - 1)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csch(x)^2/(1+tanh(x)),x, algorithm="maxima")

[Out] 2/(e^(-2*x) - 1) - log(e^(-x) + 1) - log(e^(-x) - 1)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 77 vs. 2(15) = 30.

time = 0.37, size = 77, normalized size = 5.13

$$\frac{2x \cosh(x)^2 + 4x \cosh(x) \sinh(x) + 2x \sinh(x)^2 - (\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 - 1) \log\left(\frac{2 \sinh(x)}{\cosh(x) - \sinh(x)}\right) - 2x - 2}{\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 - 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csch(x)^2/(1+tanh(x)),x, algorithm="fricas")

[Out] (2*x*cosh(x)^2 + 4*x*cosh(x)*sinh(x) + 2*x*sinh(x)^2 - (cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - 1)*log(2*sinh(x)/(cosh(x) - sinh(x))) - 2*x - 2)/(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - 1)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{csch}^2(x)}{\tanh(x) + 1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csch(x)**2/(1+tanh(x)),x)

[Out] Integral(csch(x)**2/(tanh(x) + 1), x)

Giac [A]

time = 0.41, size = 29, normalized size = 1.93

$$2x + \frac{e^{(2x)} - 3}{e^{(2x)} - 1} - \log(|e^{(2x)} - 1|)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csch(x)^2/(1+tanh(x)),x, algorithm="giac")

[Out] 2*x + (e^(2*x) - 3)/(e^(2*x) - 1) - log(abs(e^(2*x) - 1))

Mupad [B]

time = 1.08, size = 23, normalized size = 1.53

$$2x - \ln(e^{2x} - 1) - \frac{2}{e^{2x} - 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(sinh(x)^2*(tanh(x) + 1)),x)

[Out] 2*x - log(exp(2*x) - 1) - 2/(exp(2*x) - 1)

$$3.75 \quad \int \frac{\operatorname{csch}^3(x)}{1+\tanh(x)} dx$$

Optimal. Leaf size=18

$$-\frac{1}{2} \tanh^{-1}(\cosh(x)) + \operatorname{csch}(x) - \frac{1}{2} \coth(x) \operatorname{csch}(x)$$

[Out] -1/2*arctanh(cosh(x))+csch(x)-1/2*coth(x)*csch(x)

Rubi [A]

time = 0.11, antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.636$, Rules used = {3599, 3187, 3186, 2686, 8, 2691, 3855}

$$\operatorname{csch}(x) - \frac{1}{2} \tanh^{-1}(\cosh(x)) - \frac{1}{2} \coth(x) \operatorname{csch}(x)$$

Antiderivative was successfully verified.

[In] Int[Csch[x]^3/(1 + Tanh[x]), x]

[Out] -1/2*ArcTanh[Cosh[x]] + Csch[x] - (Coth[x]*Csch[x])/2

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 2686

Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Dist[a/f, Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])

Rule 2691

Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[b*(a*Sec[e + f*x])^m*((b*Tan[e + f*x])^(n - 1)/(f*(m + n - 1))), x] - Dist[b^2*((n - 1)/(m + n - 1)), Int[(a*Sec[e + f*x])^m*(b*Tan[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] && NeQ[m + n - 1, 0] && IntegerQ[2*m, 2*n]

Rule 3186

Int[cos[(c_.) + (d_.)*(x_)]^(m_.)*sin[(c_.) + (d_.)*(x_)]^(n_.)*(cos[(c_.) + (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)]^(p_.), x_Symbol] := Int[ExpandTrig[cos[c + d*x]^m*sin[c + d*x]^n*(a*cos[c + d*x] + b*sin[c + d*x]

)^p, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && IGtQ[p, 0]

Rule 3187

Int[cos[(c_.) + (d_.)*(x_)]^(m_.)*sin[(c_.) + (d_.)*(x_)]^(n_.)*(cos[(c_.) + (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)]^(p_)), x_Symbol] := Dist[a^p*b^p, Int[(Cos[c + d*x]^m*Sin[c + d*x]^n)/(b*Cos[c + d*x] + a*Sin[c + d*x])^p, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[a^2 + b^2, 0] && ILtQ[p, 0]

Rule 3599

Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]^(n_.)), x_Symbol] := Int[Sin[e + f*x]^m*((a*Cos[e + f*x] + b*Sin[e + f*x])^n/Cos[e + f*x]^n), x] /; FreeQ[{a, b, e, f}, x] && IntegerQ[(m - 1)/2] && ILtQ[n, 0] && ((LtQ[m, 5] && GtQ[n, -4]) || (EqQ[m, 5] && EqQ[n, -1]))

Rule 3855

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned}
 \int \frac{\operatorname{csch}^3(x)}{1 + \tanh(x)} dx &= \int \frac{\operatorname{coth}(x)\operatorname{csch}^2(x)}{\cosh(x) + \sinh(x)} dx \\
 &= i \int \operatorname{coth}(x)\operatorname{csch}^2(x)(-i \cosh(x) + i \sinh(x)) dx \\
 &= \int (-\operatorname{coth}(x)\operatorname{csch}(x) + \operatorname{coth}^2(x)\operatorname{csch}(x)) dx \\
 &= -\int \operatorname{coth}(x)\operatorname{csch}(x) dx + \int \operatorname{coth}^2(x)\operatorname{csch}(x) dx \\
 &= -\frac{1}{2} \operatorname{coth}(x)\operatorname{csch}(x) + i \operatorname{Subst}\left(\int 1 dx, x, -i \operatorname{csch}(x)\right) + \frac{1}{2} \int \operatorname{csch}(x) dx \\
 &= -\frac{1}{2} \tanh^{-1}(\cosh(x)) + \operatorname{csch}(x) - \frac{1}{2} \operatorname{coth}(x)\operatorname{csch}(x)
 \end{aligned}$$

Mathematica [A]

time = 0.06, size = 20, normalized size = 1.11

$$\frac{1}{2} \left(-((-2 + \operatorname{coth}(x))\operatorname{csch}(x)) + \log\left(\tanh\left(\frac{x}{2}\right)\right) \right)$$

Antiderivative was successfully verified.

[In] Integrate[Csch[x]^3/(1 + Tanh[x]),x]

[Out] $(-((-2 + \text{Coth}[x])\text{Csch}[x]) + \text{Log}[\text{Tanh}[x/2]])/2$

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 38 vs. $2(14) = 28$.

time = 0.61, size = 39, normalized size = 2.17

method	result	size
risch	$\frac{e^x(e^{2x}-3)}{(e^{2x}-1)^2} + \frac{\ln(e^x-1)}{2} - \frac{\ln(e^x+1)}{2}$	33
default	$\frac{(\tanh^2(\frac{x}{2}))}{8} - \frac{\tanh(\frac{x}{2})}{2} - \frac{1}{8 \tanh(\frac{x}{2})^2} + \frac{\ln(\tanh(\frac{x}{2}))}{2} + \frac{1}{2 \tanh(\frac{x}{2})}$	39

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csch(x)^3/(1+tanh(x)),x,method=_RETURNVERBOSE)

[Out] $1/8*\tanh(1/2*x)^2-1/2*\tanh(1/2*x)-1/8/\tanh(1/2*x)^2+1/2*\ln(\tanh(1/2*x))+1/2/\tanh(1/2*x)$

Maxima [B] Leaf count of result is larger than twice the leaf count of optimal. 48 vs. $2(14) = 28$.

time = 0.26, size = 48, normalized size = 2.67

$$-\frac{e^{-x} - 3e^{-3x}}{2e^{-2x} - e^{-4x} - 1} - \frac{1}{2} \log(e^{-x} + 1) + \frac{1}{2} \log(e^{-x} - 1)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csch(x)^3/(1+tanh(x)),x, algorithm="maxima")

[Out] $-(e^{-x} - 3e^{-3x})/(2e^{-2x} - e^{-4x} - 1) - 1/2*\log(e^{-x} + 1) + 1/2*\log(e^{-x} - 1)$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 209 vs. $2(14) = 28$.

time = 0.36, size = 209, normalized size = 11.61

$$\frac{2 \cosh(x)^2 + 6 \cosh(x) \sinh(x)^2 + 2 \sinh(x)^2 - (\cosh(x) + 4 \cosh(x) \sinh(x)^2 + \sinh(x)^2 + 2(3 \cosh(x)^2 - 1) \sinh(x)^2 - 2 \cosh(x)^2 + 4(\cosh(x)^2 - \cosh(x)) \sinh(x) + 1) \log(\cosh(x) + \sinh(x) + 1) + (\cosh(x)^2 + 4 \cosh(x) \sinh(x)^2 + \sinh(x)^2 + 2(3 \cosh(x)^2 - 1) \sinh(x)^2 - 2 \cosh(x)^2 + 4(\cosh(x)^2 - \cosh(x)) \sinh(x) + 1) \log(\cosh(x) - \sinh(x) - 1) + 6(\cosh(x)^2 - 1) \sinh(x) - 6 \cosh(x)}{2(\cosh(x)^2 + 4 \cosh(x) \sinh(x)^2 + \sinh(x)^2 + 2(3 \cosh(x)^2 - 1) \sinh(x)^2 - 2 \cosh(x)^2 + 4(\cosh(x)^2 - \cosh(x)) \sinh(x) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csch(x)^3/(1+tanh(x)),x, algorithm="fricas")

[Out] $1/2*(2*\cosh(x)^3 + 6*\cosh(x)*\sinh(x)^2 + 2*\sinh(x)^3 - (\cosh(x)^4 + 4*\cosh(x)*\sinh(x)^3 + \sinh(x)^4 + 2*(3*\cosh(x)^2 - 1)*\sinh(x)^2 - 2*\cosh(x)^2 + 4*(\cosh(x)^3 - \cosh(x))*\sinh(x) + 1)*\log(\cosh(x) + \sinh(x) + 1) + (\cosh(x)^4$

+ 4*cosh(x)*sinh(x)^3 + sinh(x)^4 + 2*(3*cosh(x)^2 - 1)*sinh(x)^2 - 2*cosh(x)^2 + 4*(cosh(x)^3 - cosh(x))*sinh(x) + 1)*log(cosh(x) + sinh(x) - 1) + 6*(cosh(x)^2 - 1)*sinh(x) - 6*cosh(x))/(cosh(x)^4 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4 + 2*(3*cosh(x)^2 - 1)*sinh(x)^2 - 2*cosh(x)^2 + 4*(cosh(x)^3 - cosh(x))*sinh(x) + 1)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{csch}^3(x)}{\tanh(x) + 1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csch(x)**3/(1+tanh(x)),x)

[Out] Integral(csch(x)**3/(tanh(x) + 1), x)

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 34 vs. 2(14) = 28.

time = 0.40, size = 34, normalized size = 1.89

$$\frac{e^{(3x)} - 3e^x}{(e^{(2x)} - 1)^2} - \frac{1}{2} \log(e^x + 1) + \frac{1}{2} \log(|e^x - 1|)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csch(x)^3/(1+tanh(x)),x, algorithm="giac")

[Out] (e^(3*x) - 3*e^x)/(e^(2*x) - 1)^2 - 1/2*log(e^x + 1) + 1/2*log(abs(e^x - 1))

Mupad [B]

time = 1.09, size = 48, normalized size = 2.67

$$\frac{\ln(1 - e^x)}{2} - \frac{\ln(-e^x - 1)}{2} + \frac{e^x}{e^{2x} - 1} - \frac{2e^x}{e^{4x} - 2e^{2x} + 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(sinh(x)^3*(tanh(x) + 1)),x)

[Out] log(1 - exp(x))/2 - log(- exp(x) - 1)/2 + exp(x)/(exp(2*x) - 1) - (2*exp(x))/(exp(4*x) - 2*exp(2*x) + 1)

$$3.76 \quad \int \frac{\operatorname{csch}^4(x)}{1+\tanh(x)} dx$$

Optimal. Leaf size=17

$$\frac{\operatorname{coth}^2(x)}{2} - \frac{\operatorname{coth}^3(x)}{3}$$

[Out] 1/2*coth(x)^2-1/3*coth(x)^3

Rubi [A]

time = 0.03, antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3597, 862, 45}

$$\frac{\operatorname{coth}^2(x)}{2} - \frac{\operatorname{coth}^3(x)}{3}$$

Antiderivative was successfully verified.

[In] Int[Csch[x]^4/(1 + Tanh[x]),x]

[Out] Coth[x]^2/2 - Coth[x]^3/3

Rule 45

Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])

Rule 862

Int[((d_.) + (e_.)*(x_))^(m_.)*((f_.) + (g_.)*(x_))^(n_.)*((a_.) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Int[(d + e*x)^(m + p)*(f + g*x)^n*(a/d + (c/e)*x)^p, x] /; FreeQ[{a, c, d, e, f, g, m, n}, x] && NeQ[e*f - d*g, 0] && EqQ[c*d^2 + a*e^2, 0] && (IntegerQ[p] || (GtQ[a, 0] && GtQ[d, 0] && EqQ[m + p, 0]))

Rule 3597

Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Dist[b/f, Subst[Int[x^m*((a + x)^n/(b^2 + x^2)^(m/2 + 1)), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, n}, x] && IntegerQ[m/2]

Rubi steps

$$\begin{aligned}
\int \frac{\operatorname{csch}^4(x)}{1 + \tanh(x)} dx &= -\operatorname{Subst}\left(\int \frac{-1 + x^2}{x^4(1 + x)} dx, x, \tanh(x)\right) \\
&= -\operatorname{Subst}\left(\int \frac{-1 + x}{x^4} dx, x, \tanh(x)\right) \\
&= -\operatorname{Subst}\left(\int \left(-\frac{1}{x^4} + \frac{1}{x^3}\right) dx, x, \tanh(x)\right) \\
&= \frac{\operatorname{coth}^2(x)}{2} - \frac{\operatorname{coth}^3(x)}{3}
\end{aligned}$$

Mathematica [A]

time = 0.04, size = 20, normalized size = 1.18

$$-\frac{1}{6} \operatorname{csch}(x)(2 \cosh(x) + (-3 + 2 \operatorname{coth}(x)) \operatorname{csch}(x))$$

Antiderivative was successfully verified.

[In] Integrate[Csch[x]^4/(1 + Tanh[x]), x]

[Out] -1/6*(Csch[x]*(2*Cosh[x] + (-3 + 2*Coth[x])*Csch[x]))

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 47 vs. 2(13) = 26.

time = 0.61, size = 48, normalized size = 2.82

method	result	size
risch	$-\frac{2(3e^{2x}+1)}{3(e^{2x}-1)^3}$	19
default	$-\frac{(\tanh^3(\frac{x}{2}))}{24} + \frac{(\tanh^2(\frac{x}{2}))}{8} - \frac{\tanh(\frac{x}{2})}{8} + \frac{1}{8 \tanh(\frac{x}{2})^2} - \frac{1}{8 \tanh(\frac{x}{2})} - \frac{1}{24 \tanh(\frac{x}{2})^3}$	48

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csch(x)^4/(1+tanh(x)), x, method=_RETURNVERBOSE)

[Out] -1/24*tanh(1/2*x)^3+1/8*tanh(1/2*x)^2-1/8*tanh(1/2*x)+1/8/tanh(1/2*x)^2-1/8/tanh(1/2*x)-1/24/tanh(1/2*x)^3

Maxima [B] Leaf count of result is larger than twice the leaf count of optimal. 75 vs. 2(13) = 26.

time = 0.26, size = 75, normalized size = 4.41

$$-\frac{2e^{(-2x)}}{3e^{(-2x)} - 3e^{(-4x)} + e^{(-6x)} - 1} + \frac{4e^{(-4x)}}{3e^{(-2x)} - 3e^{(-4x)} + e^{(-6x)} - 1} + \frac{2}{3(3e^{(-2x)} - 3e^{(-4x)} + e^{(-6x)} - 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csch(x)^4/(1+tanh(x)),x, algorithm="maxima")

[Out] $-2e^{-2x}/(3e^{-2x} - 3e^{-4x} + e^{-6x} - 1) + 4e^{-4x}/(3e^{-2x} - 3e^{-4x} + e^{-6x} - 1) + 2/3/(3e^{-2x} - 3e^{-4x} + e^{-6x} - 1)$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 84 vs. 2(13) = 26.

time = 0.38, size = 84, normalized size = 4.94

$$\frac{4(2 \cosh(x) + \sinh(x))}{3(\cosh(x)^5 + 5 \cosh(x) \sinh(x)^4 + \sinh(x)^5 + (10 \cosh(x)^2 - 3) \sinh(x)^3 - 3 \cosh(x)^3 + (10 \cosh(x)^3 - 9 \cosh(x) \sinh(x)^2 + (5 \cosh(x)^4 - 9 \cosh(x)^2 + 4) \sinh(x) + 2 \cosh(x))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csch(x)^4/(1+tanh(x)),x, algorithm="fricas")

[Out] $-4/3*(2*\cosh(x) + \sinh(x))/(\cosh(x)^5 + 5*\cosh(x)*\sinh(x)^4 + \sinh(x)^5 + (10*\cosh(x)^2 - 3)*\sinh(x)^3 - 3*\cosh(x)^3 + (10*\cosh(x)^3 - 9*\cosh(x))*\sinh(x)^2 + (5*\cosh(x)^4 - 9*\cosh(x)^2 + 4)*\sinh(x) + 2*\cosh(x))$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{csch}^4(x)}{\tanh(x) + 1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csch(x)**4/(1+tanh(x)),x)

[Out] Integral(csch(x)**4/(tanh(x) + 1), x)

Giac [A]

time = 0.40, size = 18, normalized size = 1.06

$$-\frac{2(3e^{2x} + 1)}{3(e^{2x} - 1)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csch(x)^4/(1+tanh(x)),x, algorithm="giac")

[Out] $-2/3*(3e^{2x} + 1)/(e^{2x} - 1)^3$

Mupad [B]

time = 1.06, size = 18, normalized size = 1.06

$$-\frac{2(3e^{2x} + 1)}{3(e^{2x} - 1)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(sinh(x)^4*(tanh(x) + 1)),x)
```

```
[Out] -(2*(3*exp(2*x) + 1))/(3*(exp(2*x) - 1)^3)
```

3.77 $\int \frac{\operatorname{csch}^5(x)}{1+\tanh(x)} dx$

Optimal. Leaf size=34

$$\frac{1}{8} \tanh^{-1}(\cosh(x)) - \frac{1}{8} \coth(x) \operatorname{csch}(x) + \frac{\operatorname{csch}^3(x)}{3} - \frac{1}{4} \coth(x) \operatorname{csch}^3(x)$$

[Out] 1/8*arctanh(cosh(x))-1/8*coth(x)*csch(x)+1/3*csch(x)^3-1/4*coth(x)*csch(x)^3

Rubi [A]

time = 0.14, antiderivative size = 34, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.727$, Rules used = {3599, 3187, 3186, 2686, 30, 2691, 3853, 3855}

$$\frac{\operatorname{csch}^3(x)}{3} + \frac{1}{8} \tanh^{-1}(\cosh(x)) - \frac{1}{4} \coth(x) \operatorname{csch}^3(x) - \frac{1}{8} \coth(x) \operatorname{csch}(x)$$

Antiderivative was successfully verified.

[In] Int[Csch[x]^5/(1 + Tanh[x]), x]

[Out] ArcTanh[Cosh[x]]/8 - (Coth[x]*Csch[x])/8 + Csch[x]^3/3 - (Coth[x]*Csch[x]^3)/4

Rule 30

Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2686

Int[((a_)*sec[(e_) + (f_)*(x_)])^(m_)*((b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Dist[a/f, Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])

Rule 2691

Int[((a_)*sec[(e_) + (f_)*(x_)])^(m_)*((b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(a*Sec[e + f*x])^m*((b*Tan[e + f*x])^(n - 1)/(f*(m + n - 1))), x] - Dist[b^2*((n - 1)/(m + n - 1)), Int[(a*Sec[e + f*x])^m*(b*Tan[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] && NeQ[m + n - 1, 0] && IntegerQ[2*m, 2*n]

Rule 3186

```
Int[cos[(c_.) + (d_.)*(x_)]^(m_.)*sin[(c_.) + (d_.)*(x_)]^(n_.)*(cos[(c_.)
+ (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)]^(p_.), x_Symbol] := In
t[ExpandTrig[cos[c + d*x]^m*sin[c + d*x]^n*(a*cos[c + d*x] + b*sin[c + d*x]
)^p, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && IGtQ[p, 0]
```

Rule 3187

```
Int[cos[(c_.) + (d_.)*(x_)]^(m_.)*sin[(c_.) + (d_.)*(x_)]^(n_.)*(cos[(c_.)
+ (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)]^(p_.), x_Symbol] := Dis
t[a^p*b^p, Int[(Cos[c + d*x]^m*Sin[c + d*x]^n)/(b*Cos[c + d*x] + a*Sin[c +
d*x])^p, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[a^2 + b^2, 0] && ILtQ
[p, 0]
```

Rule 3599

```
Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]^(n
_.), x_Symbol] := Int[Sin[e + f*x]^m*((a*Cos[e + f*x] + b*Sin[e + f*x])^n/C
os[e + f*x]^n), x] /; FreeQ[{a, b, e, f}, x] && IntegerQ[(m - 1)/2] && ILtQ
[n, 0] && ((LtQ[m, 5] && GtQ[n, -4]) || (EqQ[m, 5] && EqQ[n, -1]))
```

Rule 3853

```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*
x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Dist[b^2*((n - 2)/(n - 1)),
Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] &
& IntegerQ[2*n]
```

Rule 3855

```
Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned}
\int \frac{\operatorname{csch}^5(x)}{1 + \tanh(x)} dx &= \int \frac{\operatorname{coth}(x)\operatorname{csch}^4(x)}{\cosh(x) + \sinh(x)} dx \\
&= i \int \operatorname{coth}(x)\operatorname{csch}^4(x)(-i \cosh(x) + i \sinh(x)) dx \\
&= - \int (\operatorname{coth}(x)\operatorname{csch}^3(x) - \operatorname{coth}^2(x)\operatorname{csch}^3(x)) dx \\
&= - \int \operatorname{coth}(x)\operatorname{csch}^3(x) dx + \int \operatorname{coth}^2(x)\operatorname{csch}^3(x) dx \\
&= -\frac{1}{4} \operatorname{coth}(x)\operatorname{csch}^3(x) - i \operatorname{Subst}\left(\int x^2 dx, x, -i\operatorname{csch}(x)\right) + \frac{1}{4} \int \operatorname{csch}^3(x) dx \\
&= -\frac{1}{8} \operatorname{coth}(x)\operatorname{csch}(x) + \frac{\operatorname{csch}^3(x)}{3} - \frac{1}{4} \operatorname{coth}(x)\operatorname{csch}^3(x) - \frac{1}{8} \int \operatorname{csch}(x) dx \\
&= \frac{1}{8} \tanh^{-1}(\cosh(x)) - \frac{1}{8} \operatorname{coth}(x)\operatorname{csch}(x) + \frac{\operatorname{csch}^3(x)}{3} - \frac{1}{4} \operatorname{coth}(x)\operatorname{csch}^3(x)
\end{aligned}$$

Mathematica [A]

time = 0.09, size = 49, normalized size = 1.44

$$-\frac{1}{192} \operatorname{csch}^4(x) \left(42 \cosh(x) + 6 \cosh(3x) + 2 \sinh(x) \left(-32 - 9 \log\left(\tanh\left(\frac{x}{2}\right)\right) \sinh(x) + 3 \log\left(\tanh\left(\frac{x}{2}\right)\right) \sinh(3x) \right) \right)$$

Antiderivative was successfully verified.

`[In] Integrate[Csch[x]^5/(1 + Tanh[x]), x]`

```
[Out] -1/192*(Csch[x]^4*(42*Cosh[x] + 6*Cosh[3*x] + 2*Sinh[x]*(-32 - 9*Log[Tanh[x/2]]*Sinh[x] + 3*Log[Tanh[x/2]]*Sinh[3*x])))
```

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 54 vs. 2(26) = 52.

time = 0.63, size = 55, normalized size = 1.62

method	result	size
risch	$-\frac{e^x(3e^{6x} - 11e^{4x} + 53e^{2x} + 3)}{12(e^{2x} - 1)^4} + \frac{\ln(e^x + 1)}{8} - \frac{\ln(e^x - 1)}{8}$	48
default	$\frac{(\tanh^4(\frac{x}{2}))}{64} - \frac{(\tanh^3(\frac{x}{2}))}{24} + \frac{\tanh(\frac{x}{2})}{8} - \frac{1}{64 \tanh(\frac{x}{2})^4} - \frac{1}{8 \tanh(\frac{x}{2})} - \frac{\ln(\tanh(\frac{x}{2}))}{8} + \frac{1}{24 \tanh(\frac{x}{2})^3}$	55

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(csch(x)^5/(1+tanh(x)), x, method=_RETURNVERBOSE)`

```
[Out] 1/64*tanh(1/2*x)^4-1/24*tanh(1/2*x)^3+1/8*tanh(1/2*x)-1/64/tanh(1/2*x)^4-1/8/tanh(1/2*x)-1/8*ln(tanh(1/2*x))+1/24/tanh(1/2*x)^3
```

Maxima [B] Leaf count of result is larger than twice the leaf count of optimal. 74 vs. 2(26) = 52.

time = 0.27, size = 74, normalized size = 2.18

$$\frac{3e^{(-x)} - 11e^{(-3x)} + 53e^{(-5x)} + 3e^{(-7x)}}{12(4e^{(-2x)} - 6e^{(-4x)} + 4e^{(-6x)} - e^{(-8x)} - 1)} + \frac{1}{8} \log(e^{(-x)} + 1) - \frac{1}{8} \log(e^{(-x)} - 1)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csch(x)^5/(1+tanh(x)),x, algorithm="maxima")

[Out] 1/12*(3*e^(-x) - 11*e^(-3*x) + 53*e^(-5*x) + 3*e^(-7*x))/(4*e^(-2*x) - 6*e^(-4*x) + 4*e^(-6*x) - e^(-8*x) - 1) + 1/8*log(e^(-x) + 1) - 1/8*log(e^(-x) - 1)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 640 vs. 2(26) = 52.

time = 0.44, size = 640, normalized size = 18.82

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csch(x)^5/(1+tanh(x)),x, algorithm="fricas")

[Out] -1/24*(6*cosh(x)^7 + 42*cosh(x)*sinh(x)^6 + 6*sinh(x)^7 + 2*(63*cosh(x)^2 - 11)*sinh(x)^5 - 22*cosh(x)^5 + 10*(21*cosh(x)^3 - 11*cosh(x))*sinh(x)^4 + 2*(105*cosh(x)^4 - 110*cosh(x)^2 + 53)*sinh(x)^3 + 106*cosh(x)^3 + 2*(63*cosh(x)^5 - 110*cosh(x)^3 + 159*cosh(x))*sinh(x)^2 - 3*(cosh(x)^8 + 8*cosh(x)*sinh(x)^7 + sinh(x)^8 + 4*(7*cosh(x)^2 - 1)*sinh(x)^6 - 4*cosh(x)^6 + 8*(7*cosh(x)^3 - 3*cosh(x))*sinh(x)^5 + 2*(35*cosh(x)^4 - 30*cosh(x)^2 + 3)*sinh(x)^4 + 6*cosh(x)^4 + 8*(7*cosh(x)^5 - 10*cosh(x)^3 + 3*cosh(x))*sinh(x)^3 + 4*(7*cosh(x)^6 - 15*cosh(x)^4 + 9*cosh(x)^2 - 1)*sinh(x)^2 - 4*cosh(x)^2 + 8*(cosh(x)^7 - 3*cosh(x)^5 + 3*cosh(x)^3 - cosh(x))*sinh(x) + 1)*log(cosh(x) + sinh(x) + 1) + 3*(cosh(x)^8 + 8*cosh(x)*sinh(x)^7 + sinh(x)^8 + 4*(7*cosh(x)^2 - 1)*sinh(x)^6 - 4*cosh(x)^6 + 8*(7*cosh(x)^3 - 3*cosh(x))*sinh(x)^5 + 2*(35*cosh(x)^4 - 30*cosh(x)^2 + 3)*sinh(x)^4 + 6*cosh(x)^4 + 8*(7*cosh(x)^5 - 10*cosh(x)^3 + 3*cosh(x))*sinh(x)^3 + 4*(7*cosh(x)^6 - 15*cosh(x)^4 + 9*cosh(x)^2 - 1)*sinh(x)^2 - 4*cosh(x)^2 + 8*(cosh(x)^7 - 3*cosh(x)^5 + 3*cosh(x)^3 - cosh(x))*sinh(x) + 1)*log(cosh(x) + sinh(x) - 1) + 2*(21*cosh(x)^6 - 55*cosh(x)^4 + 159*cosh(x)^2 + 3)*sinh(x) + 6*cosh(x))/(cosh(x)^8 + 8*cosh(x)*sinh(x)^7 + sinh(x)^8 + 4*(7*cosh(x)^2 - 1)*sinh(x)^6 - 4*cosh(x)^6 + 8*(7*cosh(x)^3 - 3*cosh(x))*sinh(x)^5 + 2*(35*cosh(x)^4 - 30*cosh(x)^2 + 3)*sinh(x)^4 + 6*cosh(x)^4 + 8*(7*cosh(x)^5 - 10*cosh(x)^3 + 3*cosh(x))*sinh(x)^3 + 4*(7*cosh(x)^6 - 15*cosh(x)^4 + 9*cosh(x)^2 - 1)*sinh(x)^2 - 4*cosh(x)^2 + 8*(cosh(x)^7 - 3*cosh(x)^5 + 3*cosh(x)^3 - cosh(x))*sinh(x) + 1)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{csch}^5(x)}{\tanh(x) + 1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(csch(x)**5/(1+tanh(x)),x)``[Out] Integral(csch(x)**5/(tanh(x) + 1), x)`**Giac [A]**

time = 0.42, size = 49, normalized size = 1.44

$$-\frac{3e^{(7x)} - 11e^{(5x)} + 53e^{(3x)} + 3e^x}{12(e^{(2x)} - 1)^4} + \frac{1}{8} \log(e^x + 1) - \frac{1}{8} \log(|e^x - 1|)$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(csch(x)^5/(1+tanh(x)),x, algorithm="giac")``[Out] -1/12*(3*e^(7*x) - 11*e^(5*x) + 53*e^(3*x) + 3*e^x)/(e^(2*x) - 1)^4 + 1/8*log(e^x + 1) - 1/8*log(abs(e^x - 1))`**Mupad [B]**

time = 1.09, size = 117, normalized size = 3.44

$$\frac{\ln\left(\frac{e^x}{4} + \frac{1}{4}\right)}{8} - \frac{\ln\left(\frac{e^x}{4} - \frac{1}{4}\right)}{8} - \frac{e^x}{4(e^{2x} - 1)} - \frac{2e^{3x} + 2e^x}{6e^{4x} - 4e^{2x} - 4e^{6x} + e^{8x} + 1} - \frac{4e^x}{3(3e^{2x} - 3e^{4x} + e^{6x} - 1)} + \frac{e^x}{6(e^{4x} - 2e^{2x} + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/(sinh(x)^5*(tanh(x) + 1)),x)``[Out] log(exp(x)/4 + 1/4)/8 - log(exp(x)/4 - 1/4)/8 - exp(x)/(4*(exp(2*x) - 1)) - (2*exp(3*x) + 2*exp(x))/(6*exp(4*x) - 4*exp(2*x) - 4*exp(6*x) + exp(8*x) + 1) - (4*exp(x))/(3*(3*exp(2*x) - 3*exp(4*x) + exp(6*x) - 1)) + exp(x)/(6*(exp(4*x) - 2*exp(2*x) + 1))`

3.78 $\int \frac{\operatorname{csch}^6(x)}{1+\tanh(x)} dx$

Optimal. Leaf size=33

$$-\frac{1}{2} \operatorname{coth}^2(x) + \frac{\operatorname{coth}^3(x)}{3} + \frac{\operatorname{coth}^4(x)}{4} - \frac{\operatorname{coth}^5(x)}{5}$$

[Out] $-1/2*\operatorname{coth}(x)^2+1/3*\operatorname{coth}(x)^3+1/4*\operatorname{coth}(x)^4-1/5*\operatorname{coth}(x)^5$

Rubi [A]

time = 0.04, antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3597, 862, 76}

$$-\frac{1}{5} \operatorname{coth}^5(x) + \frac{\operatorname{coth}^4(x)}{4} + \frac{\operatorname{coth}^3(x)}{3} - \frac{\operatorname{coth}^2(x)}{2}$$

Antiderivative was successfully verified.

[In] `Int[Csch[x]^6/(1 + Tanh[x]),x]`

[Out] $-1/2*\operatorname{Coth}[x]^2 + \operatorname{Coth}[x]^3/3 + \operatorname{Coth}[x]^4/4 - \operatorname{Coth}[x]^5/5$

Rule 76

`Int[((d_.)*(x_))^(n_.)*((a_) + (b_.)*(x_))*((e_) + (f_.)*(x_))^(p_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)*(d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, d, e, f, n}, x] && IGtQ[p, 0] && EqQ[b*e + a*f, 0] && !(ILTQ[n + p + 2, 0] && GtQ[n + 2*p, 0])`

Rule 862

`Int[((d_) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))^(n_)*((a_) + (c_.)*(x_)^2)^(p_.), x_Symbol] :> Int[(d + e*x)^(m + p)*(f + g*x)^n*(a/d + (c/e)*x)^p, x] /; FreeQ[{a, c, d, e, f, g, m, n}, x] && NeQ[e*f - d*g, 0] && EqQ[c*d^2 + a*e^2, 0] && (IntegerQ[p] || (GtQ[a, 0] && GtQ[d, 0] && EqQ[m + p, 0]))`

Rule 3597

`Int[sin[(e_.) + (f_.)*(x_)]^(m_)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]^(n_)), x_Symbol] :> Dist[b/f, Subst[Int[x^m*((a + x)^n/(b^2 + x^2)^(m/2 + 1)), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, n}, x] && IntegerQ[m/2]`

Rubi steps

$$\begin{aligned}
\int \frac{\operatorname{csch}^6(x)}{1 + \tanh(x)} dx &= \operatorname{Subst} \left(\int \frac{(-1 + x^2)^2}{x^6(1 + x)} dx, x, \tanh(x) \right) \\
&= \operatorname{Subst} \left(\int \frac{(-1 + x)^2(1 + x)}{x^6} dx, x, \tanh(x) \right) \\
&= \operatorname{Subst} \left(\int \left(\frac{1}{x^6} - \frac{1}{x^5} - \frac{1}{x^4} + \frac{1}{x^3} \right) dx, x, \tanh(x) \right) \\
&= -\frac{1}{2} \operatorname{coth}^2(x) + \frac{\operatorname{coth}^3(x)}{3} + \frac{\operatorname{coth}^4(x)}{4} - \frac{\operatorname{coth}^5(x)}{5}
\end{aligned}$$

Mathematica [A]

time = 0.04, size = 27, normalized size = 0.82

$$\frac{1}{120} \operatorname{csch}^5(x) (-20 \cosh(x) - 5 \cosh(3x) + \cosh(5x) + 30 \sinh(x))$$

Antiderivative was successfully verified.

`[In] Integrate[Csch[x]^6/(1 + Tanh[x]), x]``[Out] (Csch[x]^5*(-20*Cosh[x] - 5*Cosh[3*x] + Cosh[5*x] + 30*Sinh[x]))/120`**Maple [B]** Leaf count of result is larger than twice the leaf count of optimal. 79 vs. 2(25) = 50.

time = 0.62, size = 80, normalized size = 2.42

method	result
risch	$-\frac{4(20e^{4x} + 5e^{2x} - 1)}{15(e^{2x} - 1)^5}$
default	$-\frac{(\tanh^5(\frac{x}{2}))}{160} + \frac{(\tanh^4(\frac{x}{2}))}{64} + \frac{(\tanh^3(\frac{x}{2}))}{96} - \frac{(\tanh^2(\frac{x}{2}))}{16} + \frac{\tanh(\frac{x}{2})}{16} - \frac{1}{160 \tanh(\frac{x}{2})^5} + \frac{1}{64 \tanh(\frac{x}{2})^4} + \frac{1}{96 \tanh(\frac{x}{2})^3}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(csch(x)^6/(1+tanh(x)), x, method=_RETURNVERBOSE)`
`[Out] -1/160*tanh(1/2*x)^5+1/64*tanh(1/2*x)^4+1/96*tanh(1/2*x)^3-1/16*tanh(1/2*x)^2+1/16*tanh(1/2*x)-1/160/tanh(1/2*x)^5+1/64/tanh(1/2*x)^4+1/96/tanh(1/2*x)^3+1/16/tanh(1/2*x)-1/16/tanh(1/2*x)^2`
Maxima [B] Leaf count of result is larger than twice the leaf count of optimal. 149 vs. 2(25) = 50.

time = 0.26, size = 149, normalized size = 4.52

$$\frac{4e^{-2x}}{3(5e^{-2x} - 10e^{-4x} + 10e^{-6x} - 5e^{-8x} + e^{-10x}) - 1} - \frac{8e^{-4x}}{3(5e^{-2x} - 10e^{-4x} + 10e^{-6x} - 5e^{-8x} + e^{-10x}) - 1} + \frac{8e^{-6x}}{5e^{-2x} - 10e^{-4x} + 10e^{-6x} - 5e^{-8x} + e^{-10x}) - 1} - \frac{4}{15(5e^{-2x} - 10e^{-4x} + 10e^{-6x} - 5e^{-8x} + e^{-10x}) - 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csch(x)^6/(1+tanh(x)),x, algorithm="maxima")

[Out] $\frac{4}{3}e^{-2x}/(5e^{-2x} - 10e^{-4x} + 10e^{-6x} - 5e^{-8x} + e^{-10x} - 1) - \frac{8}{3}e^{-4x}/(5e^{-2x} - 10e^{-4x} + 10e^{-6x} - 5e^{-8x} + e^{-10x} - 1) + \frac{8}{5}e^{-6x}/(5e^{-2x} - 10e^{-4x} + 10e^{-6x} - 5e^{-8x} + e^{-10x} - 1) - \frac{4}{15}e^{-8x}/(5e^{-2x} - 10e^{-4x} + 10e^{-6x} - 5e^{-8x} + e^{-10x} - 1)$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 185 vs. 2(25) = 50.

time = 0.33, size = 185, normalized size = 5.61

$\frac{4(19 \cosh(x)^2 + 42 \cosh(x) \sinh(x) + 19 \sinh(x)^2 + 5)}{15(\cosh(x)^8 + 8 \cosh(x) \sinh(x)^7 + \sinh(x)^8 + (28 \cosh(x)^2 - 5) \sinh(x)^6 - 5 \cosh(x)^6 + 2(28 \cosh(x)^3 - 15 \cosh(x)) \sinh(x)^5 + 5(14 \cosh(x)^4 - 15 \cosh(x)^2 + 2) \sinh(x)^4 + 10 \cosh(x)^4 + 4(14 \cosh(x)^5 - 25 \cosh(x)^3 + 10 \cosh(x)) \sinh(x)^3 + (28 \cosh(x)^6 - 75 \cosh(x)^4 + 60 \cosh(x)^2 - 11) \sinh(x)^2 - 11 \cosh(x)^2 + 2(4 \cosh(x)^7 - 15 \cosh(x)^5 + 20 \cosh(x)^3 - 9 \cosh(x)) \sinh(x) + 5)}$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csch(x)^6/(1+tanh(x)),x, algorithm="fricas")

[Out] $\frac{-4}{15} \frac{(19 \cosh(x)^2 + 42 \cosh(x) \sinh(x) + 19 \sinh(x)^2 + 5)}{(\cosh(x)^8 + 8 \cosh(x) \sinh(x)^7 + \sinh(x)^8 + (28 \cosh(x)^2 - 5) \sinh(x)^6 - 5 \cosh(x)^6 + 2(28 \cosh(x)^3 - 15 \cosh(x)) \sinh(x)^5 + 5(14 \cosh(x)^4 - 15 \cosh(x)^2 + 2) \sinh(x)^4 + 10 \cosh(x)^4 + 4(14 \cosh(x)^5 - 25 \cosh(x)^3 + 10 \cosh(x)) \sinh(x)^3 + (28 \cosh(x)^6 - 75 \cosh(x)^4 + 60 \cosh(x)^2 - 11) \sinh(x)^2 - 11 \cosh(x)^2 + 2(4 \cosh(x)^7 - 15 \cosh(x)^5 + 20 \cosh(x)^3 - 9 \cosh(x)) \sinh(x) + 5)}$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{csch}^6(x)}{\tanh(x) + 1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csch(x)**6/(1+tanh(x)),x)

[Out] Integral(csch(x)**6/(tanh(x) + 1), x)

Giac [A]

time = 0.41, size = 24, normalized size = 0.73

$$\frac{4(20e^{4x} + 5e^{2x} - 1)}{15(e^{2x} - 1)^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csch(x)^6/(1+tanh(x)),x, algorithm="giac")

[Out] $-4/15*(20*e^{(4*x)} + 5*e^{(2*x)} - 1)/(e^{(2*x)} - 1)^5$

Mupad [B]

time = 0.10, size = 24, normalized size = 0.73

$$-\frac{4(5e^{2x} + 20e^{4x} - 1)}{15(e^{2x} - 1)^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(sinh(x)^6*(tanh(x) + 1)),x)`

[Out] $-(4*(5*\exp(2*x) + 20*\exp(4*x) - 1))/(15*(\exp(2*x) - 1)^5)$

3.79 $\int \frac{\operatorname{csch}^7(x)}{1+\tanh(x)} dx$

Optimal. Leaf size=44

$$-\frac{1}{16} \tanh^{-1}(\cosh(x)) + \frac{1}{16} \coth(x) \operatorname{csch}(x) - \frac{1}{24} \coth(x) \operatorname{csch}^3(x) + \frac{\operatorname{csch}^5(x)}{5} - \frac{1}{6} \coth(x) \operatorname{csch}^5(x)$$

[Out] $-1/16*\operatorname{arctanh}(\cosh(x))+1/16*\coth(x)*\operatorname{csch}(x)-1/24*\coth(x)*\operatorname{csch}(x)^3+1/5*\operatorname{csch}(x)^5-1/6*\coth(x)*\operatorname{csch}(x)^5$

Rubi [A]

time = 0.15, antiderivative size = 44, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 8, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.727$, Rules used = {3599, 3187, 3186, 2686, 30, 2691, 3853, 3855}

$$\frac{\operatorname{csch}^5(x)}{5} - \frac{1}{16} \tanh^{-1}(\cosh(x)) - \frac{1}{6} \coth(x) \operatorname{csch}^5(x) - \frac{1}{24} \coth(x) \operatorname{csch}^3(x) + \frac{1}{16} \coth(x) \operatorname{csch}(x)$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[\operatorname{Csch}[x]^7/(1 + \operatorname{Tanh}[x]), x]$

[Out] $-1/16*\operatorname{ArcTanh}[\operatorname{Cosh}[x]] + (\operatorname{Coth}[x]*\operatorname{Csch}[x])/16 - (\operatorname{Coth}[x]*\operatorname{Csch}[x]^3)/24 + \operatorname{Csch}[x]^5/5 - (\operatorname{Coth}[x]*\operatorname{Csch}[x]^5)/6$

Rule 30

$\operatorname{Int}[(x_)^{(m_.)}, x_Symbol] \rightarrow \operatorname{Simp}[x^{(m+1)}/(m+1), x] /; \operatorname{FreeQ}[m, x] \ \&\& \operatorname{NeQ}[m, -1]$

Rule 2686

$\operatorname{Int}[(a_.)*\operatorname{sec}[(e_.) + (f_.)*(x_)]^{(m_.)}*((b_.)*\operatorname{tan}[(e_.) + (f_.)*(x_)]^{(n_.)}, x_Symbol] \rightarrow \operatorname{Dist}[a/f, \operatorname{Subst}[\operatorname{Int}[(a*x)^{(m-1)}*(-1+x^2)^{((n-1)/2)}, x], x, \operatorname{Sec}[e+f*x]], x] /; \operatorname{FreeQ}[\{a, e, f, m\}, x] \ \&\& \operatorname{IntegerQ}[(n-1)/2] \ \&\& \operatorname{IntegerQ}[m/2] \ \&\& \operatorname{LtQ}[0, m, n+1]$

Rule 2691

$\operatorname{Int}[(a_.)*\operatorname{sec}[(e_.) + (f_.)*(x_)]^{(m_.)}*((b_.)*\operatorname{tan}[(e_.) + (f_.)*(x_)]^{(n_.)}, x_Symbol] \rightarrow \operatorname{Simp}[b*(a*\operatorname{Sec}[e+f*x])^{(m)}*((b*\operatorname{Tan}[e+f*x])^{(n-1)})/(f*(m+n-1)), x] - \operatorname{Dist}[b^2*((n-1)/(m+n-1)), \operatorname{Int}[(a*\operatorname{Sec}[e+f*x])^{(m)}*(b*\operatorname{Tan}[e+f*x])^{(n-2)}, x], x] /; \operatorname{FreeQ}[\{a, b, e, f, m\}, x] \ \&\& \operatorname{GtQ}[n, 1] \ \&\& \operatorname{NeQ}[m+n-1, 0] \ \&\& \operatorname{IntegersQ}[2*m, 2*n]$

Rule 3186

```
Int[cos[(c_.) + (d_.)*(x_)]^(m_.)*sin[(c_.) + (d_.)*(x_)]^(n_.)*(cos[(c_.)
+ (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)]^(p_.), x_Symbol] := Int
t[ExpandTrig[cos[c + d*x]^m*sin[c + d*x]^n*(a*cos[c + d*x] + b*sin[c + d*x]
)^p, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && IGtQ[p, 0]
```

Rule 3187

```
Int[cos[(c_.) + (d_.)*(x_)]^(m_.)*sin[(c_.) + (d_.)*(x_)]^(n_.)*(cos[(c_.)
+ (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)]^(p_.), x_Symbol] := Dis
t[a^p*b^p, Int[(Cos[c + d*x]^m*Sin[c + d*x]^n)/(b*Cos[c + d*x] + a*Sin[c +
d*x])^p, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[a^2 + b^2, 0] && ILt
Q[p, 0]
```

Rule 3599

```
Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]^(n
_.), x_Symbol] := Int[Sin[e + f*x]^m*((a*cos[e + f*x] + b*sin[e + f*x])^n/C
os[e + f*x]^n), x] /; FreeQ[{a, b, e, f}, x] && IntegerQ[(m - 1)/2] && ILtQ
[n, 0] && ((LtQ[m, 5] && GtQ[n, -4]) || (EqQ[m, 5] && EqQ[n, -1]))
```

Rule 3853

```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*
x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Dist[b^2*((n - 2)/(n - 1)),
Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] &
& IntegerQ[2*n]
```

Rule 3855

```
Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned}
\int \frac{\operatorname{csch}^7(x)}{1 + \tanh(x)} dx &= \int \frac{\operatorname{coth}(x)\operatorname{csch}^6(x)}{\cosh(x) + \sinh(x)} dx \\
&= i \int \operatorname{coth}(x)\operatorname{csch}^6(x)(-i \cosh(x) + i \sinh(x)) dx \\
&= \int (-\operatorname{coth}(x)\operatorname{csch}^5(x) + \operatorname{coth}^2(x)\operatorname{csch}^5(x)) dx \\
&= -\int \operatorname{coth}(x)\operatorname{csch}^5(x) dx + \int \operatorname{coth}^2(x)\operatorname{csch}^5(x) dx \\
&= -\frac{1}{6} \operatorname{coth}(x)\operatorname{csch}^5(x) + i \operatorname{Subst}\left(\int x^4 dx, x, -i\operatorname{csch}(x)\right) + \frac{1}{6} \int \operatorname{csch}^5(x) dx \\
&= -\frac{1}{24} \operatorname{coth}(x)\operatorname{csch}^3(x) + \frac{\operatorname{csch}^5(x)}{5} - \frac{1}{6} \operatorname{coth}(x)\operatorname{csch}^5(x) - \frac{1}{8} \int \operatorname{csch}^3(x) dx \\
&= \frac{1}{16} \operatorname{coth}(x)\operatorname{csch}(x) - \frac{1}{24} \operatorname{coth}(x)\operatorname{csch}^3(x) + \frac{\operatorname{csch}^5(x)}{5} - \frac{1}{6} \operatorname{coth}(x)\operatorname{csch}^5(x) + \frac{1}{16} \int \operatorname{csch}(x) dx \\
&= -\frac{1}{16} \tanh^{-1}(\cosh(x)) + \frac{1}{16} \operatorname{coth}(x)\operatorname{csch}(x) - \frac{1}{24} \operatorname{coth}(x)\operatorname{csch}^3(x) + \frac{\operatorname{csch}^5(x)}{5} - \frac{1}{6} \operatorname{coth}(x)\operatorname{csch}^5(x)
\end{aligned}$$

Mathematica [A]

time = 0.19, size = 68, normalized size = 1.55

$$\frac{\operatorname{csch}^6(x) (-1140 \cosh(x) - 170 \cosh(3x) + 30 \cosh(5x) + 6 \sinh(x) (256 + 50 \log(\tanh(\frac{x}{2})) \sinh(x) - 25 \log(\tanh(\frac{x}{2})) \sinh(3x) + 5 \log(\tanh(\frac{x}{2})) \sinh(5x))}{7680}$$

Antiderivative was successfully verified.

`[In] Integrate[Csch[x]^7/(1 + Tanh[x]), x]`

```
[Out] (Csch[x]^6*(-1140*Cosh[x] - 170*Cosh[3*x] + 30*Cosh[5*x] + 6*Sinh[x]*(256 + 50*Log[Tanh[x/2]]*Sinh[x] - 25*Log[Tanh[x/2]]*Sinh[3*x] + 5*Log[Tanh[x/2]]*Sinh[5*x]))/7680
```

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 102 vs. 2(34) = 68.

time = 0.63, size = 103, normalized size = 2.34

method	result
risch	$\frac{e^x (15 e^{10x} - 85 e^{8x} + 198 e^{6x} - 1338 e^{4x} - 85 e^{2x} + 15)}{120(e^{2x} - 1)^6} + \frac{\ln(e^x - 1)}{16} - \frac{\ln(e^x + 1)}{16}$
default	$\frac{(\tanh^6(\frac{x}{2}))}{384} - \frac{(\tanh^5(\frac{x}{2}))}{160} - \frac{(\tanh^4(\frac{x}{2}))}{128} + \frac{(\tanh^3(\frac{x}{2}))}{32} - \frac{(\tanh^2(\frac{x}{2}))}{128} - \frac{\tanh(\frac{x}{2})}{16} + \frac{1}{128 \tanh(\frac{x}{2})^4} - \frac{1}{32 \tanh(\frac{x}{2})^3} + \dots$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(csch(x)^7/(1+tanh(x)), x, method=_RETURNVERBOSE)`

[Out] $1/384*\tanh(1/2*x)^6-1/160*\tanh(1/2*x)^5-1/128*\tanh(1/2*x)^4+1/32*\tanh(1/2*x)^3-1/128*\tanh(1/2*x)^2-1/16*\tanh(1/2*x)+1/128/\tanh(1/2*x)^4-1/32/\tanh(1/2*x)^3+1/16*\ln(\tanh(1/2*x))+1/128/\tanh(1/2*x)^2-1/384/\tanh(1/2*x)^6+1/16/\tanh(1/2*x)+1/160/\tanh(1/2*x)^5$

Maxima [B] Leaf count of result is larger than twice the leaf count of optimal. 98 vs. $2(34) = 68$.

time = 0.27, size = 98, normalized size = 2.23

$$\frac{15e^{-x} - 85e^{-3x} + 198e^{-5x} - 1338e^{-7x} - 85e^{-9x} + 15e^{-11x}}{120(6e^{-2x} - 15e^{-4x} + 20e^{-6x} - 15e^{-8x} + 6e^{-10x} - e^{-12x} - 1)} - \frac{1}{16} \log(e^{-x} + 1) + \frac{1}{16} \log(e^{-x} - 1)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csch(x)^7/(1+tanh(x)),x, algorithm="maxima")`

[Out] $-1/120*(15*e^{-x} - 85*e^{-3*x} + 198*e^{-5*x} - 1338*e^{-7*x} - 85*e^{-9*x} + 15*e^{-11*x})/(6*e^{-2*x} - 15*e^{-4*x} + 20*e^{-6*x} - 15*e^{-8*x} + 6*e^{-10*x} - e^{-12*x} - 1) - 1/16*\log(e^{-x} + 1) + 1/16*\log(e^{-x} - 1)$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 1260 vs. $2(34) = 68$.

time = 0.36, size = 1260, normalized size = 28.64

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csch(x)^7/(1+tanh(x)),x, algorithm="fricas")`

[Out] $1/240*(30*\cosh(x)^{11} + 330*\cosh(x)*\sinh(x)^{10} + 30*\sinh(x)^{11} + 10*(165*\cosh(x)^2 - 17)*\sinh(x)^9 - 170*\cosh(x)^9 + 90*(55*\cosh(x)^3 - 17*\cosh(x))*\sinh(x)^8 + 36*(275*\cosh(x)^4 - 170*\cosh(x)^2 + 11)*\sinh(x)^7 + 396*\cosh(x)^7 + 84*(165*\cosh(x)^5 - 170*\cosh(x)^3 + 33*\cosh(x))*\sinh(x)^6 + 12*(1155*\cosh(x)^6 - 1785*\cosh(x)^4 + 693*\cosh(x)^2 - 223)*\sinh(x)^5 - 2676*\cosh(x)^5 + 60*(165*\cosh(x)^7 - 357*\cosh(x)^5 + 231*\cosh(x)^3 - 223*\cosh(x))*\sinh(x)^4 + 10*(495*\cosh(x)^8 - 1428*\cosh(x)^6 + 1386*\cosh(x)^4 - 2676*\cosh(x)^2 - 17)*\sinh(x)^3 - 170*\cosh(x)^3 + 6*(275*\cosh(x)^9 - 1020*\cosh(x)^7 + 1386*\cosh(x)^5 - 4460*\cosh(x)^3 - 85*\cosh(x))*\sinh(x)^2 - 15*(\cosh(x)^{12} + 12*\cosh(x))*\sinh(x)^{11} + \sinh(x)^{12} + 6*(11*\cosh(x)^2 - 1)*\sinh(x)^{10} - 6*\cosh(x)^{10} + 20*(11*\cosh(x)^3 - 3*\cosh(x))*\sinh(x)^9 + 15*(33*\cosh(x)^4 - 18*\cosh(x)^2 + 1)*\sinh(x)^8 + 15*\cosh(x)^8 + 24*(33*\cosh(x)^5 - 30*\cosh(x)^3 + 5*\cosh(x))*\sinh(x)^7 + 4*(231*\cosh(x)^6 - 315*\cosh(x)^4 + 105*\cosh(x)^2 - 5)*\sinh(x)^6 - 20*\cosh(x)^6 + 24*(33*\cosh(x)^7 - 63*\cosh(x)^5 + 35*\cosh(x)^3 - 5*\cosh(x))*\sinh(x)^5 + 15*(33*\cosh(x)^8 - 84*\cosh(x)^6 + 70*\cosh(x)^4 - 20*\cosh(x)^2 + 1)*\sinh(x)^4 + 15*\cosh(x)^4 + 20*(11*\cosh(x)^9 - 36*\cosh(x)^7 + 42*\cosh(x)^5 - 20*\cosh(x)^3 + 3*\cosh(x))*\sinh(x)^3 + 6*(11*\cosh(x)^{10} - 45*\cosh(x)^8 + 70*\cosh(x)^6 - 50*\cosh(x)^4 + 15*\cosh(x)^2 - 1)*\sinh(x)^2 - 6*\cosh(x)$

$x)^2 + 12*(\cosh(x)^{11} - 5*\cosh(x)^9 + 10*\cosh(x)^7 - 10*\cosh(x)^5 + 5*\cosh(x)^3 - \cosh(x))*\sinh(x) + 1)*\log(\cosh(x) + \sinh(x) + 1) + 15*(\cosh(x)^{12} + 12*\cosh(x)*\sinh(x)^{11} + \sinh(x)^{12} + 6*(11*\cosh(x)^2 - 1)*\sinh(x)^{10} - 6*\cosh(x)^{10} + 20*(11*\cosh(x)^3 - 3*\cosh(x))*\sinh(x)^9 + 15*(33*\cosh(x)^4 - 18*\cosh(x)^2 + 1)*\sinh(x)^8 + 15*\cosh(x)^8 + 24*(33*\cosh(x)^5 - 30*\cosh(x)^3 + 5*\cosh(x))*\sinh(x)^7 + 4*(231*\cosh(x)^6 - 315*\cosh(x)^4 + 105*\cosh(x)^2 - 5)*\sinh(x)^6 - 20*\cosh(x)^6 + 24*(33*\cosh(x)^7 - 63*\cosh(x)^5 + 35*\cosh(x)^3 - 5*\cosh(x))*\sinh(x)^5 + 15*(33*\cosh(x)^8 - 84*\cosh(x)^6 + 70*\cosh(x)^4 - 20*\cosh(x)^2 + 1)*\sinh(x)^4 + 15*\cosh(x)^4 + 20*(11*\cosh(x)^9 - 36*\cosh(x)^7 + 42*\cosh(x)^5 - 20*\cosh(x)^3 + 3*\cosh(x))*\sinh(x)^3 + 6*(11*\cosh(x)^{10} - 45*\cosh(x)^8 + 70*\cosh(x)^6 - 50*\cosh(x)^4 + 15*\cosh(x)^2 - 1)*\sinh(x)^2 - 6*\cosh(x)^2 + 12*(\cosh(x)^{11} - 5*\cosh(x)^9 + 10*\cosh(x)^7 - 10*\cosh(x)^5 + 5*\cosh(x)^3 - \cosh(x))*\sinh(x) + 1)*\log(\cosh(x) + \sinh(x) - 1) + 6*(55*\cosh(x)^{10} - 255*\cosh(x)^8 + 462*\cosh(x)^6 - 2230*\cosh(x)^4 - 85*\cosh(x)^2 + 5)*\sinh(x) + 30*\cosh(x))/(\cosh(x)^{12} + 12*\cosh(x)*\sinh(x)^{11} + \sinh(x)^{12} + 6*(11*\cosh(x)^2 - 1)*\sinh(x)^{10} - 6*\cosh(x)^{10} + 20*(11*\cosh(x)^3 - 3*\cosh(x))*\sinh(x)^9 + 15*(33*\cosh(x)^4 - 18*\cosh(x)^2 + 1)*\sinh(x)^8 + 15*\cosh(x)^8 + 24*(33*\cosh(x)^5 - 30*\cosh(x)^3 + 5*\cosh(x))*\sinh(x)^7 + 4*(231*\cosh(x)^6 - 315*\cosh(x)^4 + 105*\cosh(x)^2 - 5)*\sinh(x)^6 - 20*\cosh(x)^6 + 24*(33*\cosh(x)^7 - 63*\cosh(x)^5 + 35*\cosh(x)^3 - 5*\cosh(x))*\sinh(x)^5 + 15*(33*\cosh(x)^8 - 84*\cosh(x)^6 + 70*\cosh(x)^4 - 20*\cosh(x)^2 + 1)*\sinh(x)^4 + 15*\cosh(x)^4 + 20*(11*\cosh(x)^9 - 36*\cosh(x)^7 + 42*\cosh(x)^5 - 20*\cosh(x)^3 + 3*\cosh(x))*\sinh(x)^3 + 6*(11*\cosh(x)^{10} - 45*\cosh(x)^8 + 70*\cosh(x)^6 - 50*\cosh(x)^4 + 15*\cosh(x)^2 - 1)*\sinh(x)^2 - 6*\cosh(x)^2 + 12*(\cosh(x)^{11} - 5*\cosh(x)^9 + 10*\cosh(x)^7 - 10*\cosh(x)^5 + 5*\cosh(x)^3 - \cosh(x))*\sinh(x) + 1)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{csch}^7(x)}{\tanh(x) + 1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csch(x)**7/(1+tanh(x)),x)

[Out] Integral(csch(x)**7/(tanh(x) + 1), x)

Giac [A]

time = 0.40, size = 61, normalized size = 1.39

$$\frac{15 e^{(11x)} - 85 e^{(9x)} + 198 e^{(7x)} - 1338 e^{(5x)} - 85 e^{(3x)} + 15 e^x}{120 (e^{(2x)} - 1)^6} - \frac{1}{16} \log(e^x + 1) + \frac{1}{16} \log(|e^x - 1|)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csch(x)^7/(1+tanh(x)),x, algorithm="giac")

[Out] $1/120*(15*e^{(11*x)} - 85*e^{(9*x)} + 198*e^{(7*x)} - 1338*e^{(5*x)} - 85*e^{(3*x)} + 15*e^x)/(e^{(2*x)} - 1)^6 - 1/16*\log(e^x + 1) + 1/16*\log(\text{abs}(e^x - 1))$

Mupad [B]

time = 1.17, size = 207, normalized size = 4.70

$$\frac{\ln\left(\frac{1}{8} - \frac{e^x}{8}\right)}{16} - \frac{\ln\left(-\frac{e^x}{8} - \frac{1}{8}\right)}{16} - \frac{\frac{16e^{3x}}{3} + \frac{16e^{5x}}{3}}{15e^{4x} - 6e^{2x} - 20e^{6x} + 15e^{8x} - 6e^{10x} + e^{12x} + 1} - \frac{\frac{8e^{3x}}{3} + \frac{8e^x}{3}}{5e^{2x} - 10e^{4x} + 10e^{6x} - 5e^{8x} + e^{10x} - 1} - \frac{6e^x}{5(6e^{4x} - 4e^{2x} - 4e^{6x} + e^{8x} + 1)} + \frac{e^x}{8(e^{2x} - 1)} + \frac{e^x}{15(3e^{2x} - 3e^{4x} + e^{6x} - 1)} - \frac{e^x}{12(e^{4x} - 2e^{2x} + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(sinh(x)^7*(tanh(x) + 1)),x)`

[Out] $\log(1/8 - \exp(x)/8)/16 - \log(-\exp(x)/8 - 1/8)/16 - ((16*\exp(3*x))/3 + (16*\exp(5*x))/3)/(15*\exp(4*x) - 6*\exp(2*x) - 20*\exp(6*x) + 15*\exp(8*x) - 6*\exp(10*x) + \exp(12*x) + 1) - ((8*\exp(3*x))/3 + (8*\exp(x))/5)/(5*\exp(2*x) - 10*\exp(4*x) + 10*\exp(6*x) - 5*\exp(8*x) + \exp(10*x) - 1) - (6*\exp(x))/(5*(6*\exp(4*x) - 4*\exp(2*x) - 4*\exp(6*x) + \exp(8*x) + 1)) + \exp(x)/(8*(\exp(2*x) - 1)) + \exp(x)/(15*(3*\exp(2*x) - 3*\exp(4*x) + \exp(6*x) - 1)) - \exp(x)/(12*(\exp(4*x) - 2*\exp(2*x) + 1))$

3.80 $\int \frac{\sinh^4(x)}{a+b \tanh(x)} dx$

Optimal. Leaf size=147

$$-\frac{a(3a+b)\log(1-\tanh(x))}{16(a+b)^3} + \frac{a(3a-b)\log(1+\tanh(x))}{16(a-b)^3} - \frac{a^4 b \log(a+b \tanh(x))}{(a^2-b^2)^3} - \frac{\cosh^4(x)(b-a \tanh(x))}{4(a^2-b^2)}$$

[Out] $-1/16*a*(3*a+b)*\ln(1-\tanh(x))/(a+b)^3+1/16*a*(3*a-b)*\ln(1+\tanh(x))/(a-b)^3-a^4*b*\ln(a+b*\tanh(x))/(a^2-b^2)^3-1/4*\cosh(x)^4*(b-a*\tanh(x))/(a^2-b^2)+1/8*\cosh(x)^2*(4*b*(2*a^2-b^2)-a*(5*a^2-b^2)*\tanh(x))/(a^2-b^2)^2$

Rubi [A]

time = 0.26, antiderivative size = 147, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$,

Rules used = {3597, 1661, 815}

$$-\frac{\cosh^4(x)(b-a \tanh(x))}{4(a^2-b^2)} + \frac{\cosh^2(x)(4b(2a^2-b^2)-a(5a^2-b^2)\tanh(x))}{8(a^2-b^2)^2} - \frac{a^4 b \log(a+b \tanh(x))}{(a^2-b^2)^3} - \frac{a(3a+b)\log(1-\tanh(x))}{16(a+b)^3} + \frac{a(3a-b)\log(\tanh(x)+1)}{16(a-b)^3}$$

Antiderivative was successfully verified.

[In] `Int[Sinh[x]^4/(a + b*Tanh[x]), x]`

[Out] $-1/16*(a*(3*a + b)*\text{Log}[1 - \text{Tanh}[x]]/(a + b)^3 + (a*(3*a - b)*\text{Log}[1 + \text{Tanh}[x]])/(16*(a - b)^3) - (a^4*b*\text{Log}[a + b*\text{Tanh}[x]]/(a^2 - b^2)^3 - (\text{Cosh}[x]^4*(b - a*\text{Tanh}[x]))/(4*(a^2 - b^2)) + (\text{Cosh}[x]^2*(4*b*(2*a^2 - b^2) - a*(5*a^2 - b^2)*\text{Tanh}[x]))/(8*(a^2 - b^2)^2)$

Rule 815

`Int[(((d_.) + (e_.)*(x_))^(m_.)*((f_.) + (g_.)*(x_)))/((a_.) + (c_.)*(x_)^2), x_Symbol] :> Int[ExpandIntegrand[(d + e*x)^m*((f + g*x)/(a + c*x^2)), x], x] /; FreeQ[{a, c, d, e, f, g}, x] && NeQ[c*d^2 + a*e^2, 0] && IntegerQ[m]`

Rule 1661

`Int[(Pq_)*((d_) + (e_.)*(x_))^(m_.)*((a_) + (c_.)*(x_)^2)^(p_), x_Symbol] :> With[{Q = PolynomialQuotient[(d + e*x)^m*Pq, a + c*x^2, x], f = Coeff[PolynomialRemainder[(d + e*x)^m*Pq, a + c*x^2, x], x, 0], g = Coeff[PolynomialRemainder[(d + e*x)^m*Pq, a + c*x^2, x], x, 1]}, Simp[(a*g - c*f*x)*((a + c*x^2)^(p + 1)/(2*a*c*(p + 1))), x] + Dist[1/(2*a*c*(p + 1)), Int[(d + e*x)^m*(a + c*x^2)^(p + 1)*ExpandToSum[(2*a*c*(p + 1)*Q]/(d + e*x)^m + (c*f*(2*p + 3))/(d + e*x)^m, x], x] /; FreeQ[{a, c, d, e}, x] && PolyQ[Pq, x] && NeQ[c*d^2 + a*e^2, 0] && LtQ[p, -1] && ILtQ[m, 0]`

Rule 3597

```
Int[sin[(e_.) + (f_.)*(x_)]^(m_)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_
), x_Symbol] :> Dist[b/f, Subst[Int[x^m*((a + x)^n/(b^2 + x^2)^(m/2 + 1)),
x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, n}, x] && IntegerQ[m/2]
```

Rubi steps

$$\begin{aligned} \int \frac{\sinh^4(x)}{a + b \tanh(x)} dx &= - \left(b \text{Subst} \left(\int \frac{x^4}{(a+x)(-b^2+x^2)^3} dx, x, b \tanh(x) \right) \right) \\ &= - \frac{\cosh^4(x)(b - a \tanh(x))}{4(a^2 - b^2)} - \frac{\text{Subst} \left(\int \frac{\frac{a^2 b^4}{a^2 - b^2} - \frac{3ab^4 x}{a^2 - b^2} + 4b^2 x^2}{(a+x)(-b^2+x^2)^2} dx, x, b \tanh(x) \right)}{4b} \\ &= - \frac{\cosh^4(x)(b - a \tanh(x))}{4(a^2 - b^2)} + \frac{\cosh^2(x)(4b(2a^2 - b^2) - a(5a^2 - b^2) \tanh(x))}{8(a^2 - b^2)^2} - \frac{\text{Subst} \left(\int \frac{a^2 b^4}{(a+x)(-b^2+x^2)^2} dx, x, b \tanh(x) \right)}{4b} \\ &= - \frac{\cosh^4(x)(b - a \tanh(x))}{4(a^2 - b^2)} + \frac{\cosh^2(x)(4b(2a^2 - b^2) - a(5a^2 - b^2) \tanh(x))}{8(a^2 - b^2)^2} - \frac{\text{Subst} \left(\int \frac{a^2 b^4}{(a+x)(-b^2+x^2)^2} dx, x, b \tanh(x) \right)}{4b} \\ &= - \frac{a(3a + b) \log(1 - \tanh(x))}{16(a + b)^3} + \frac{a(3a - b) \log(1 + \tanh(x))}{16(a - b)^3} - \frac{a^4 b \log(a + b \tanh(x))}{(a^2 - b^2)^3} \end{aligned}$$

Mathematica [A]

time = 0.42, size = 144, normalized size = 0.98

$$\frac{12a^5x + 24a^3b^2x - 4ab^4x + 4b(3a^4 - 4a^2b^2 + b^4) \cosh(2x) - b(a^2 - b^2)^2 \cosh(4x) - 32a^4b \log(a \cosh(x) + b \sinh(x)) - 8a^3(a^2 - b^2) \sinh(2x) + a^5 \sinh(4x) - 2a^2b^2 \sinh(4x) + ab^4 \sinh(4x)}{32(a-b)^3(a+b)^3}$$

Antiderivative was successfully verified.

[In] Integrate[Sinh[x]^4/(a + b*Tanh[x]), x]

[Out] (12*a^5*x + 24*a^3*b^2*x - 4*a*b^4*x + 4*b*(3*a^4 - 4*a^2*b^2 + b^4)*Cosh[2*x] - b*(a^2 - b^2)^2*Cosh[4*x] - 32*a^4*b*Log[a*Cosh[x] + b*Sinh[x]] - 8*a^3*(a^2 - b^2)*Sinh[2*x] + a^5*Sinh[4*x] - 2*a^3*b^2*Sinh[4*x] + a*b^4*Sinh[4*x])/(32*(a - b)^3*(a + b)^3)

Maple [A]

time = 0.70, size = 246, normalized size = 1.67

method	result
risch	$\frac{3a^2x}{8(a+b)^3} + \frac{axb}{8(a+b)^3} + \frac{e^{4x}}{64a+64b} - \frac{e^{2x}a}{8(a+b)^2} - \frac{e^{2x}b}{16(a+b)^2} - \frac{e^{-2x}b}{16(a-b)^2} + \frac{e^{-2x}a}{8(a-b)^2} - \frac{e^{-4x}}{64(a-b)} + \frac{2a^4bx}{a^6-3a^4b^2+3a^2b^4-b^6} -$

default	$-\frac{8}{(32a-32b)(\tanh(\frac{x}{2})+1)^4} + \frac{32}{(64a-64b)(\tanh(\frac{x}{2})+1)^3} - \frac{-a-b}{8(a-b)^2(\tanh(\frac{x}{2})+1)^2} - \frac{3a-b}{8(a-b)^2(\tanh(\frac{x}{2})+1)} + \frac{a(3a-b)\ln(\tanh(\frac{x}{2}))}{8(a-b)^2}$
---------	---

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sinh(x)^4/(a+b*tanh(x)),x,method=_RETURNVERBOSE)`

[Out]
$$-8/(32*a-32*b)/(\tanh(1/2*x)+1)^4+32/(64*a-64*b)/(\tanh(1/2*x)+1)^3-1/8*(-a-b)/(a-b)^2/(\tanh(1/2*x)+1)^2-1/8*(3*a-b)/(a-b)^2/(\tanh(1/2*x)+1)+1/8*a*(3*a-b)/(a-b)^3*\ln(\tanh(1/2*x)+1)+8/(32*a+32*b)/(\tanh(1/2*x)-1)^4+32/(64*a+64*b)/(\tanh(1/2*x)-1)^3-1/8*(a-b)/(a+b)^2/(\tanh(1/2*x)-1)^2-1/8*(3*a+b)/(a+b)^2/(\tanh(1/2*x)-1)-1/8*a*(3*a+b)/(a+b)^3*\ln(\tanh(1/2*x)-1)-a^4*b/(a-b)^3/(a+b)^3*\ln(a*tanh(1/2*x)^2+2*b*tanh(1/2*x)+a)$$

Maxima [A]

time = 0.28, size = 163, normalized size = 1.11

$$-\frac{a^4 b \log(-(a-b)e^{-2x}-a-b)}{a^6-3a^4b^2+3a^2b^4-b^6} + \frac{(3a^2+ab)x}{8(a^3+3a^2b+3ab^2+b^3)} - \frac{(4(2a+b)e^{-2x}-a-b)e^{4x}}{64(a^2+2ab+b^2)} + \frac{4(2a-b)e^{-2x}-(a-b)e^{-4x}}{64(a^2-2ab+b^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sinh(x)^4/(a+b*tanh(x)),x, algorithm="maxima")`

[Out]
$$-a^4*b*\log(-(a-b)*e^{-2*x}-a-b)/(a^6-3*a^4*b^2+3*a^2*b^4-b^6)+1/8*(3*a^2+3*a*b)*x/(a^3+3*a^2*b+3*a*b^2+b^3)-1/64*(4*(2*a+b)*e^{-2*x}-a-b)*e^{4*x}/(a^2+2*a*b+b^2)+1/64*(4*(2*a-b)*e^{-2*x}-(a-b)*e^{-4*x})/(a^2-2*a*b+b^2)$$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 1226 vs. 2(139) = 278.

time = 0.36, size = 1226, normalized size = 8.34

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sinh(x)^4/(a+b*tanh(x)),x, algorithm="fricas")`

[Out]
$$1/64*((a^5-a^4*b-2*a^3*b^2+2*a^2*b^3+a*b^4-b^5)*\cosh(x)^8+8*(a^5-a^4*b-2*a^3*b^2+2*a^2*b^3+a*b^4-b^5)*\cosh(x)*\sinh(x)^7+(a^5-a^4*b-2*a^3*b^2+2*a^2*b^3+a*b^4-b^5)*\sinh(x)^8-4*(2*a^5-3*a^4*b-b-2*a^3*b^2+4*a^2*b^3-b^5)*\cosh(x)^6-4*(2*a^5-3*a^4*b-2*a^3*b^2+4*a^2*b^3-b^5-7*(a^5-a^4*b-2*a^3*b^2+2*a^2*b^3+a*b^4-b^5))*\cosh(x)^2*\sinh(x)^6+8*(3*a^5+8*a^4*b+6*a^3*b^2-a*b^4)*x*\cosh(x)^4+8*(7*(a^5-a^4*b-2*a^3*b^2+2*a^2*b^3+a*b^4-b^5)*\cosh(x)^3-3*(2*a^5-3*a^4*b-2*a^3*b^2+4*a^2*b^3-b^5)*\cosh(x))*\sinh(x)^5-a^5-a^4*b+2*a^3*b^2+2*a^2*b^3-a*b^4-b^5+2*(35*(a^5-a^4*b-2*a^3*b^2+2*a^2*b^3+a*b^4-b^5)*\cosh(x)^4-30*(2*a^5-3*a^4*b-2*a^3*b^2+4*$$

$$\begin{aligned}
& a^2 b^3 - b^5) \cosh(x)^2 + 4(3a^5 + 8a^4 b + 6a^3 b^2 - a b^4) x \sinh(x)^4 + 8(7(a^5 - a^4 b - 2a^3 b^2 + 2a^2 b^3 + a b^4 - b^5) \cosh(x)^5 - \\
& 10(2a^5 - 3a^4 b - 2a^3 b^2 + 4a^2 b^3 - b^5) \cosh(x)^3 + 4(3a^5 + 8a^4 b + 6a^3 b^2 - a b^4) x \cosh(x) \sinh(x)^3 + 4(2a^5 + 3a^4 b - 2a^3 b^2 - 4a^2 b^3 + \\
& b^5 - 15(2a^5 - 3a^4 b - 2a^3 b^2 + 4a^2 b^3 - b^5) \cosh(x)^4 + 12(3a^5 + 8a^4 b + 6a^3 b^2 - a b^4) x \cosh(x)^2 \sinh(x)^2 - 64(a^4 b \cosh(x)^4 + 4a^4 b \cosh(x)^3 \sinh(x) + 6a^4 b \cosh(x)^2 \sinh(x)^2 + 4a^4 b \cosh(x) \sinh(x)^3 + a^4 b \sinh(x)^4) \log(2(a \cosh(x) + b \sinh(x)) / (\cosh(x) - \sinh(x))) + 8((a^5 - a^4 b - 2a^3 b^2 + 2a^2 b^3 + a b^4 - b^5) \cosh(x)^7 - 3(2a^5 - 3a^4 b - 2a^3 b^2 + 4a^2 b^3 - b^5) \cosh(x)^5 + 4(3a^5 + 8a^4 b + 6a^3 b^2 - a b^4) x \cosh(x)^3 + (2a^5 + 3a^4 b - 2a^3 b^2 - 4a^2 b^3 + b^5) \cosh(x) \sinh(x)) / ((a^6 - 3a^4 b^2 + 3a^2 b^4 - b^6) \cosh(x)^4 + 4(a^6 - 3a^4 b^2 + 3a^2 b^4 - b^6) \cosh(x)^3 \sinh(x) + 6(a^6 - 3a^4 b^2 + 3a^2 b^4 - b^6) \cosh(x)^2 \sinh(x)^2 + 4(a^6 - 3a^4 b^2 + 3a^2 b^4 - b^6) \cosh(x) \sinh(x)^3 + (a^6 - 3a^4 b^2 + 3a^2 b^4 - b^6) \sinh(x)^4)
\end{aligned}$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sinh^4(x)}{a + b \tanh(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sinh(x)**4/(a+b*tanh(x)),x)

[Out] Integral(sinh(x)**4/(a + b*tanh(x)), x)

Giac [A]

time = 0.42, size = 214, normalized size = 1.46

$$-\frac{a^4 b \log((a e^{2x}) + b e^{4x} + a - b)}{a^6 - 3a^4 b^2 + 3a^2 b^4 - b^6} + \frac{(3a^2 - ab)x}{8(a^3 - 3a^2 b + 3ab^2 - b^3)} - \frac{(18a^2 e^{4x}) - 6abe^{4x} - 8a^2 e^{2x} + 12abe^{2x} - 4b^2 e^{2x} + a^2 - 2ab + b^2) e^{-4x}}{64(a^3 - 3a^2 b + 3ab^2 - b^3)} + \frac{ae^{4x} + be^{4x} - 8ae^{2x} - 4be^{2x}}{64(a^2 + 2ab + b^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sinh(x)^4/(a+b*tanh(x)),x, algorithm="giac")

[Out] $-a^4 b \log(\text{abs}(a e^{2x} + b e^{4x} + a - b)) / (a^6 - 3a^4 b^2 + 3a^2 b^4 - b^6) + 1/8(3a^2 - a b) x / (a^3 - 3a^2 b + 3a b^2 - b^3) - 1/64(18a^2 e^{4x} - 6a b e^{4x} - 8a^2 e^{2x} + 12a b e^{2x} - 4b^2 e^{2x}) + a^2 - 2a b + b^2) e^{-4x} / (a^3 - 3a^2 b + 3a b^2 - b^3) + 1/64(a e^{4x} + b e^{4x} - 8a e^{2x} - 4b e^{2x}) / (a^2 + 2a b + b^2)$

Mupad [B]

time = 1.67, size = 135, normalized size = 0.92

$$\frac{e^{4x}}{64a + 64b} - \frac{e^{-4x}}{64a - 64b} + \frac{e^{-2x}(2a - b)}{16(a - b)^2} - \frac{e^{2x}(2a + b)}{16(a + b)^2} - \frac{a^4 b \ln(a - b + a e^{2x} + b e^{2x})}{a^6 - 3a^4 b^2 + 3a^2 b^4 - b^6} + \frac{a x (3a - b)}{8(a - b)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sinh(x)^4/(a + b*tanh(x)),x)
```

```
[Out] exp(4*x)/(64*a + 64*b) - exp(-4*x)/(64*a - 64*b) + (exp(-2*x)*(2*a - b))/(16*(a - b)^2) - (exp(2*x)*(2*a + b))/(16*(a + b)^2) - (a^4*b*log(a - b + a*exp(2*x) + b*exp(2*x)))/(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2) + (a*x*(3*a - b))/(8*(a - b)^3)
```


3.81 $\int \frac{\sinh^3(x)}{a+b \tanh(x)} dx$

Optimal. Leaf size=137

$$-\frac{a^3 b \operatorname{ArcTan}\left(\frac{b \cosh(x) + a \sinh(x)}{\sqrt{a^2 - b^2}}\right)}{(a^2 - b^2)^{5/2}} - \frac{ab^2 \cosh(x)}{(a^2 - b^2)^2} - \frac{a \cosh(x)}{a^2 - b^2} + \frac{a \cosh^3(x)}{3(a^2 - b^2)} + \frac{a^2 b \sinh(x)}{(a^2 - b^2)^2} - \frac{b \sinh^3(x)}{3(a^2 - b^2)}$$

[Out] $-a^3 b \arctan((b \cosh(x) + a \sinh(x)) / (a^2 - b^2)^{1/2}) / (a^2 - b^2)^{5/2} - a b^2 \cosh(x) / (a^2 - b^2)^2 - a \cosh(x) / (a^2 - b^2) + 1/3 a \cosh(x)^3 / (a^2 - b^2) + a^2 b \sinh(x) / (a^2 - b^2)^2 - 1/3 b \sinh(x)^3 / (a^2 - b^2)$

Rubi [A]

time = 0.20, antiderivative size = 137, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 9, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.692$, Rules used = {3599, 3188, 2644, 30, 2713, 3178, 3153, 212, 2718}

$$-\frac{b \sinh^3(x)}{3(a^2 - b^2)} + \frac{a^2 b \sinh(x)}{(a^2 - b^2)^2} + \frac{a \cosh^3(x)}{3(a^2 - b^2)} - \frac{a \cosh(x)}{a^2 - b^2} - \frac{ab^2 \cosh(x)}{(a^2 - b^2)^2} - \frac{a^3 b \operatorname{ArcTan}\left(\frac{a \sinh(x) + b \cosh(x)}{\sqrt{a^2 - b^2}}\right)}{(a^2 - b^2)^{5/2}}$$

Antiderivative was successfully verified.

[In] `Int[Sinh[x]^3/(a + b*Tanh[x]),x]`

[Out] $-\left(\frac{a^3 b \operatorname{ArcTan}\left[\frac{b \operatorname{Cosh}[x] + a \operatorname{Sinh}[x]}{\sqrt{a^2 - b^2}}\right]}{(a^2 - b^2)^{5/2}}\right) - \frac{a b^2 \operatorname{Cosh}[x]}{(a^2 - b^2)^2} - \frac{a \operatorname{Cosh}[x]}{a^2 - b^2} + \frac{a \operatorname{Cosh}[x]^3}{3(a^2 - b^2)} + \frac{a^2 b \operatorname{Sinh}[x]}{(a^2 - b^2)^2} - \frac{b \operatorname{Sinh}[x]^3}{3(a^2 - b^2)}$

Rule 30

`Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]`

Rule 212

`Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rule 2644

`Int[cos[(e_) + (f_)*(x_)]^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] := Dist[1/(a*f), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(In`

tegerQ[(m - 1)/2] && LtQ[0, m, n])

Rule 2713

Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Dist[-d^(-1), Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]

Rule 2718

Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]

Rule 3153

Int[(cos[(c_.) + (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-1), x_Symbol] := Dist[-d^(-1), Subst[Int[1/(a^2 + b^2 - x^2), x], x, b*Cos[c + d*x] - a*Sin[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0]

Rule 3178

Int[sin[(c_.) + (d_.)*(x_)]^(m_)/(cos[(c_.) + (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)]), x_Symbol] := Simp[(-a)*(Sin[c + d*x]^(m - 1)/(d*(a^2 + b^2)*(m - 1))), x] + (Dist[a^2/(a^2 + b^2), Int[Sin[c + d*x]^(m - 2)/(a*Cos[c + d*x] + b*Sin[c + d*x]), x], x] + Dist[b/(a^2 + b^2), Int[Sin[c + d*x]^(m - 1), x], x]) /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0] && GtQ[m, 1]

Rule 3188

Int[(cos[(c_.) + (d_.)*(x_)]^(m_)*sin[(c_.) + (d_.)*(x_)]^(n_))/(cos[(c_.) + (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)]), x_Symbol] := Dist[b/(a^2 + b^2), Int[Cos[c + d*x]^m*Sin[c + d*x]^(n - 1), x], x] + (Dist[a/(a^2 + b^2), Int[Cos[c + d*x]^(m - 1)*Sin[c + d*x]^n, x], x] - Dist[a*(b/(a^2 + b^2)), Int[Cos[c + d*x]^(m - 1)*(Sin[c + d*x]^(n - 1)/(a*Cos[c + d*x] + b*Sin[c + d*x])), x], x]) /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0] && IGtQ[m, 0] && IGtQ[n, 0]

Rule 3599

Int[sin[(e_.) + (f_.)*(x_)]^(m_)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Int[Sin[e + f*x]^m*((a*Cos[e + f*x] + b*Sin[e + f*x])^n/Cos[e + f*x]^n), x] /; FreeQ[{a, b, e, f}, x] && IntegerQ[(m - 1)/2] && ILtQ[n, 0] && ((LtQ[m, 5] && GtQ[n, -4]) || (EqQ[m, 5] && EqQ[n, -1]))

Rubi steps

$$\begin{aligned}
\int \frac{\sinh^3(x)}{a + b \tanh(x)} dx &= \int \frac{\cosh(x) \sinh^3(x)}{a \cosh(x) + b \sinh(x)} dx \\
&= \frac{a \int \sinh^3(x) dx}{a^2 - b^2} - \frac{b \int \cosh(x) \sinh^2(x) dx}{a^2 - b^2} + \frac{(ab) \int \frac{\sinh^2(x)}{a \cosh(x) + b \sinh(x)} dx}{a^2 - b^2} \\
&= \frac{a^2 b \sinh(x)}{(a^2 - b^2)^2} - \frac{(a^3 b) \int \frac{1}{a \cosh(x) + b \sinh(x)} dx}{(a^2 - b^2)^2} - \frac{(ab^2) \int \sinh(x) dx}{(a^2 - b^2)^2} - \frac{a \text{Subst}(\int (1 - x^2) dx)}{a^2 - b^2} \\
&= -\frac{ab^2 \cosh(x)}{(a^2 - b^2)^2} - \frac{a \cosh(x)}{a^2 - b^2} + \frac{a \cosh^3(x)}{3(a^2 - b^2)} + \frac{a^2 b \sinh(x)}{(a^2 - b^2)^2} - \frac{b \sinh^3(x)}{3(a^2 - b^2)} - \frac{(ia^3 b) \text{Subst}(\int (1 - x^2) dx)}{a^2 - b^2} \\
&= -\frac{a^3 b \tan^{-1}\left(\frac{b \cosh(x) + a \sinh(x)}{\sqrt{a^2 - b^2}}\right)}{(a^2 - b^2)^{5/2}} - \frac{ab^2 \cosh(x)}{(a^2 - b^2)^2} - \frac{a \cosh(x)}{a^2 - b^2} + \frac{a \cosh^3(x)}{3(a^2 - b^2)} + \frac{a^2 b \sinh(x)}{(a^2 - b^2)^2}
\end{aligned}$$

Mathematica [A]

time = 0.84, size = 180, normalized size = 1.31

$$\frac{-3a\sqrt{a-b}\sqrt{a+b}(3a^2+b^2)\cosh(x) + a\sqrt{a-b}\sqrt{a+b}(a^2-b^2)\cosh(3x) + b\left(-24a^3\text{ArcTan}\left(\frac{b+a\tanh\left(\frac{x}{2}\right)}{\sqrt{a-b}\sqrt{a+b}}\right) + 3\sqrt{a-b}\sqrt{a+b}(5a^2-b^2)\sinh(x) - \sqrt{a-b}\sqrt{a+b}(a^2-b^2)\sinh(3x)\right)}{12(a-b)^{5/2}(a+b)^{5/2}}$$

Antiderivative was successfully verified.

`[In] Integrate[Sinh[x]^3/(a + b*Tanh[x]), x]`

```
[Out] (-3*a*Sqrt[a - b]*Sqrt[a + b]*(3*a^2 + b^2)*Cosh[x] + a*Sqrt[a - b]*Sqrt[a + b]*(a^2 - b^2)*Cosh[3*x] + b*(-24*a^3*ArcTan[(b + a*Tanh[x/2])]/(Sqrt[a - b]*Sqrt[a + b])) + 3*Sqrt[a - b]*Sqrt[a + b]*(5*a^2 - b^2)*Sinh[x] - Sqrt[a - b]*Sqrt[a + b]*(a^2 - b^2)*Sinh[3*x])/(12*(a - b)^(5/2)*(a + b)^(5/2))
```

Maple [A]

time = 0.69, size = 166, normalized size = 1.21

method	result
default	$ -\frac{16}{3(\tanh(\frac{x}{2})-1)^3(16a+16b)} - \frac{8}{(16a+16b)(\tanh(\frac{x}{2})-1)^2} + \frac{a}{2(a+b)^2(\tanh(\frac{x}{2})-1)} - \frac{8}{(16a-16b)(\tanh(\frac{x}{2})+1)^2} + \frac{1}{3(\tanh(\frac{x}{2})+1)} $
risch	$ \frac{e^{3x}}{24a+24b} - \frac{3e^x a}{8(a+b)^2} - \frac{e^x b}{8(a+b)^2} - \frac{3e^{-x} a}{8(a-b)^2} + \frac{e^{-x} b}{8(a-b)^2} + \frac{e^{-3x}}{24a-24b} - \frac{b a^3 \ln\left(e^x + \frac{a-b}{\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2}(a+b)^2(a-b)^2} + \frac{b a^3 \ln\left(e^x - \frac{a-b}{\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2}(a+b)^2(a-b)^2} $

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sinh(x)^3/(a+b*tanh(x)), x, method=_RETURNVERBOSE)`

[Out]
$$-16/3/(\tanh(1/2*x)-1)^3/(16*a+16*b)-8/(16*a+16*b)/(\tanh(1/2*x)-1)^2+1/2*a/(a+b)^2/(\tanh(1/2*x)-1)-8/(16*a-16*b)/(\tanh(1/2*x)+1)^2+16/3/(\tanh(1/2*x)+1)^3/(16*a-16*b)-1/2*a/(a-b)^2/(\tanh(1/2*x)+1)-2*a^3*b/(a-b)^2/(a+b)^2/(a^2-b^2)^{1/2}*\arctan(1/2*(2*a*\tanh(1/2*x)+2*b)/(a^2-b^2)^{1/2})$$

Maxima [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sinh(x)^3/(a+b*tanh(x)),x, algorithm="maxima")`

[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see 'assume?' for more details)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 903 vs. 2(129) = 258.

time = 0.36, size = 1861, normalized size = 13.58

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sinh(x)^3/(a+b*tanh(x)),x, algorithm="fricas")`

[Out]
$$\begin{aligned} & [1/24*((a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*\cosh(x)^6 + 6*(a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*\cosh(x)*\sinh(x)^5 + (a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*\sinh(x)^6 + a^5 + a^4*b - 2*a^3*b^2 - 2*a^2*b^3 + a*b^4 + b^5 - 3*(3*a^5 - 5*a^4*b - 2*a^3*b^2 + 6*a^2*b^3 - a*b^4 - b^5)*\cosh(x)^4 - 3*(3*a^5 - 5*a^4*b - 2*a^3*b^2 + 6*a^2*b^3 - a*b^4 - b^5 - 5*(a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*\cosh(x)^2)*\sinh(x)^4 + 4*(5*(a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*\cosh(x)^3 - 3*(3*a^5 - 5*a^4*b - 2*a^3*b^2 + 6*a^2*b^3 - a*b^4 - b^5)*\cosh(x))*\sinh(x)^3 - 3*(3*a^5 + 5*a^4*b - 2*a^3*b^2 - 6*a^2*b^3 - a*b^4 + b^5)*\cosh(x)^2 - 3*(3*a^5 + 5*a^4*b - 2*a^3*b^2 - 6*a^2*b^3 + a*b^4 - b^5)*\cosh(x)^4 + 6*(3*a^5 - 5*a^4*b - 2*a^3*b^2 + 6*a^2*b^3 - a*b^4 - b^5)*\cosh(x)^2)*\sinh(x)^2 - 24*(a^3*b*\cosh(x)^3 + 3*a^3*b*\cosh(x)^2*\sinh(x) + 3*a^3*b*\cosh(x)*\sinh(x)^2 + a^3*b*\sinh(x)^3)*\sqrt{-a^2 + b^2}*\log(((a + b)*\cosh(x)^2 + 2*(a + b)*\cosh(x)*\sinh(x) + (a + b)*\sinh(x)^2 + 2*\sqrt{-a^2 + b^2}*(\cosh(x) + \sinh(x)) - a + b)/((a + b)*\cosh(x)^2 + 2*(a + b)*\cosh(x)*\sinh(x) + (a + b)*\sinh(x)^2 + a - b)) + 6*((a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*\cosh(x)^5 - 2*(3*a^5 - 5*a^4*b - 2*a^3*b^2 + 6*a^2*b^3 - a*b^4 - b^5)*\cosh(x)^3 - (3*a^5 + 5*a^4*b - 2*a^3*b^2 - 6*a^2*b^3 - a*b^4 + b^5)*\cosh(x))*\sinh(x)]/((a^6 - 3*a \end{aligned}$$

$$\begin{aligned} &^4*b^2 + 3*a^2*b^4 - b^6)*\cosh(x)^3 + 3*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6) \\ &*\cosh(x)^2*\sinh(x) + 3*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\cosh(x)*\sinh(x)^2 \\ &+ (a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\sinh(x)^3), 1/24*((a^5 - a^4*b - 2* \\ &a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*\cosh(x)^6 + 6*(a^5 - a^4*b - 2*a^3*b^2 + \\ &2*a^2*b^3 + a*b^4 - b^5)*\cosh(x)*\sinh(x)^5 + (a^5 - a^4*b - 2*a^3*b^2 + 2* \\ &a^2*b^3 + a*b^4 - b^5)*\sinh(x)^6 + a^5 + a^4*b - 2*a^3*b^2 - 2*a^2*b^3 + a* \\ &b^4 + b^5 - 3*(3*a^5 - 5*a^4*b - 2*a^3*b^2 + 6*a^2*b^3 - a*b^4 - b^5)*\cosh(\\ &x)^4 - 3*(3*a^5 - 5*a^4*b - 2*a^3*b^2 + 6*a^2*b^3 - a*b^4 - b^5 - 5*(a^5 - \\ &a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*\cosh(x)^2)*\sinh(x)^4 + 4*(5*(a \\ &^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*\cosh(x)^3 - 3*(3*a^5 - 5* \\ &a^4*b - 2*a^3*b^2 + 6*a^2*b^3 - a*b^4 - b^5)*\cosh(x))*\sinh(x)^3 - 3*(3*a^5 \\ &+ 5*a^4*b - 2*a^3*b^2 - 6*a^2*b^3 - a*b^4 + b^5)*\cosh(x)^2 - 3*(3*a^5 + 5*a \\ &^4*b - 2*a^3*b^2 - 6*a^2*b^3 - a*b^4 + b^5 - 5*(a^5 - a^4*b - 2*a^3*b^2 + 2* \\ &a^2*b^3 + a*b^4 - b^5)*\cosh(x)^4 + 6*(3*a^5 - 5*a^4*b - 2*a^3*b^2 + 6*a^2* \\ &b^3 - a*b^4 - b^5)*\cosh(x)^2)*\sinh(x)^2 + 48*(a^3*b*\cosh(x)^3 + 3*a^3*b*\cos \\ &h(x)^2*\sinh(x) + 3*a^3*b*\cosh(x)*\sinh(x)^2 + a^3*b*\sinh(x)^3)*\sqrt{a^2 - b^2} \\ &)*\arctan(\sqrt{a^2 - b^2}/((a + b)*\cosh(x) + (a + b)*\sinh(x))) + 6*((a^5 - \\ &a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*\cosh(x)^5 - 2*(3*a^5 - 5*a^4*b \\ &- 2*a^3*b^2 + 6*a^2*b^3 - a*b^4 - b^5)*\cosh(x)^3 - (3*a^5 + 5*a^4*b - 2*a^3* \\ &b^2 - 6*a^2*b^3 - a*b^4 + b^5)*\cosh(x))*\sinh(x))/((a^6 - 3*a^4*b^2 + 3*a^2* \\ &b^4 - b^6)*\cosh(x)^3 + 3*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\cosh(x)^2*\sinh(x) \\ &+ 3*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\cosh(x)*\sinh(x)^2 + (a^6 - 3* \\ &a^4*b^2 + 3*a^2*b^4 - b^6)*\sinh(x)^3)] \end{aligned}$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sinh^3(x)}{a + b \tanh(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sinh(x)**3/(a+b*tanh(x)),x)

[Out] Integral(sinh(x)**3/(a + b*tanh(x)), x)

Giac [A]

time = 0.42, size = 163, normalized size = 1.19

$$\frac{2a^3b \arctan\left(\frac{ae^x + be^x}{\sqrt{a^2 - b^2}}\right)}{(a^4 - 2a^2b^2 + b^4)\sqrt{a^2 - b^2}} - \frac{(9ae^{2x} - 3be^{2x} - a + b)e^{-3x}}{24(a^2 - 2ab + b^2)} + \frac{a^2e^{3x} + 2abe^{3x} + b^2e^{3x} - 9a^2e^x - 12abe^x - 3b^2e^x}{24(a^3 + 3a^2b + 3ab^2 + b^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sinh(x)^3/(a+b*tanh(x)),x, algorithm="giac")

[Out] -2*a^3*b*arctan((a*e^x + b*e^x)/sqrt(a^2 - b^2))/((a^4 - 2*a^2*b^2 + b^4)*sqrt(a^2 - b^2)) - 1/24*(9*a*e^(2*x) - 3*b*e^(2*x) - a + b)*e^(-3*x)/(a^2 -

$$2*a*b + b^2) + 1/24*(a^2*e^{(3*x)} + 2*a*b*e^{(3*x)} + b^2*e^{(3*x)} - 9*a^2*e^x - 12*a*b*e^x - 3*b^2*e^x)/(a^3 + 3*a^2*b + 3*a*b^2 + b^3)$$

Mupad [B]

time = 1.46, size = 261, normalized size = 1.91

$$\frac{e^{-3x}}{24a - 24b} + \frac{e^{3x}}{24a + 24b} - \frac{e^x(3a + b)}{8(a + b)^2} - \frac{e^{-x}(3a - b)}{8(a - b)^2} - \frac{2 \operatorname{atan}\left(\frac{a^3 b e^x \sqrt{a^{10} - 5a^8 b^2 + 10a^6 b^4 - 10a^4 b^6 + 5a^2 b^8 - b^{10}}}{a^5 \sqrt{a^6 b^2} - b^5 \sqrt{a^6 b^2} + 2a^2 b^3 \sqrt{a^6 b^2} - 2a^3 b^2 \sqrt{a^6 b^2} + a b^4 \sqrt{a^6 b^2} - a^4 b \sqrt{a^6 b^2}}\right) \sqrt{a^6 b^2}}{\sqrt{a^{10} - 5a^8 b^2 + 10a^6 b^4 - 10a^4 b^6 + 5a^2 b^8 - b^{10}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sinh(x)^3/(a + b*tanh(x)),x)`

[Out] `exp(-3*x)/(24*a - 24*b) + exp(3*x)/(24*a + 24*b) - (exp(x)*(3*a + b))/(8*(a + b)^2) - (exp(-x)*(3*a - b))/(8*(a - b)^2) - (2*atan((a^3*b*exp(x)*(a^10 - b^10 + 5*a^2*b^8 - 10*a^4*b^6 + 10*a^6*b^4 - 5*a^8*b^2)^(1/2))/(a^5*(a^6*b^2)^(1/2) - b^5*(a^6*b^2)^(1/2) + 2*a^2*b^3*(a^6*b^2)^(1/2) - 2*a^3*b^2*(a^6*b^2)^(1/2) + a*b^4*(a^6*b^2)^(1/2) - a^4*b*(a^6*b^2)^(1/2)))/(a^6*b^2)^(1/2))/(a^10 - b^10 + 5*a^2*b^8 - 10*a^4*b^6 + 10*a^6*b^4 - 5*a^8*b^2)^(1/2)`

3.82 $\int \frac{\sinh^2(x)}{a+b \tanh(x)} dx$

Optimal. Leaf size=84

$$\frac{a \log(1 - \tanh(x))}{4(a+b)^2} - \frac{a \log(1 + \tanh(x))}{4(a-b)^2} + \frac{a^2 b \log(a + b \tanh(x))}{(a^2 - b^2)^2} - \frac{\cosh^2(x)(b - a \tanh(x))}{2(a^2 - b^2)}$$

[Out] $1/4*a*\ln(1-\tanh(x))/(a+b)^2-1/4*a*\ln(1+\tanh(x))/(a-b)^2+a^2*b*\ln(a+b*\tanh(x))/(a^2-b^2)^2-1/2*\cosh(x)^2*(b-a*\tanh(x))/(a^2-b^2)$

Rubi [A]

time = 0.11, antiderivative size = 84, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {3597, 1661, 815}

$$\frac{a^2 b \log(a + b \tanh(x))}{(a^2 - b^2)^2} - \frac{\cosh^2(x)(b - a \tanh(x))}{2(a^2 - b^2)} + \frac{a \log(1 - \tanh(x))}{4(a+b)^2} - \frac{a \log(\tanh(x) + 1)}{4(a-b)^2}$$

Antiderivative was successfully verified.

[In] Int[Sinh[x]^2/(a + b*Tanh[x]),x]

[Out] $(a*\text{Log}[1 - \text{Tanh}[x]])/(4*(a + b)^2) - (a*\text{Log}[1 + \text{Tanh}[x]])/(4*(a - b)^2) + (a^2*b*\text{Log}[a + b*\text{Tanh}[x]])/(a^2 - b^2)^2 - (\text{Cosh}[x]^2*(b - a*\text{Tanh}[x]))/(2*(a^2 - b^2))$

Rule 815

Int[(((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_)))/((a_) + (c_)*(x_)^2), x_Symbol] :> Int[ExpandIntegrand[(d + e*x)^m*((f + g*x)/(a + c*x^2)), x], x] /; FreeQ[{a, c, d, e, f, g}, x] && NeQ[c*d^2 + a*e^2, 0] && IntegerQ[m]

Rule 1661

Int[(Pq_)*((d_) + (e_)*(x_))^(m_)*((a_) + (c_)*(x_)^2)^(p_), x_Symbol] :> With[{Q = PolynomialQuotient[(d + e*x)^m*Pq, a + c*x^2, x], f = Coeff[PolynomialRemainder[(d + e*x)^m*Pq, a + c*x^2, x], x, 0], g = Coeff[PolynomialRemainder[(d + e*x)^m*Pq, a + c*x^2, x], x, 1]}, Simp[(a*g - c*f*x)*((a + c*x^2)^(p + 1)/(2*a*c*(p + 1))), x] + Dist[1/(2*a*c*(p + 1)), Int[(d + e*x)^m*(a + c*x^2)^(p + 1)*ExpandToSum[(2*a*c*(p + 1)*Q)/(d + e*x)^m + (c*f*(2*p + 3))/(d + e*x)^m, x], x] /; FreeQ[{a, c, d, e}, x] && PolyQ[Pq, x] && NeQ[c*d^2 + a*e^2, 0] && LtQ[p, -1] && ILtQ[m, 0]

Rule 3597

Int[sin[(e_) + (f_)*(x_)]^(m_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] :> Dist[b/f, Subst[Int[x^m*((a + x)^n/(b^2 + x^2)^(m/2 + 1)),

`x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, n}, x] && IntegerQ[m/2]`

Rubi steps

$$\begin{aligned} \int \frac{\sinh^2(x)}{a + b \tanh(x)} dx &= b \text{Subst} \left(\int \frac{x^2}{(a+x)(-b^2+x^2)^2} dx, x, b \tanh(x) \right) \\ &= -\frac{\cosh^2(x)(b-a \tanh(x))}{2(a^2-b^2)} + \frac{\text{Subst} \left(\int \frac{\frac{a^2 b^2}{a^2-b^2} - \frac{a b^2 x}{a^2-b^2}}{(a+x)(-b^2+x^2)} dx, x, b \tanh(x) \right)}{2b} \\ &= -\frac{\cosh^2(x)(b-a \tanh(x))}{2(a^2-b^2)} + \frac{\text{Subst} \left(\int \left(-\frac{ab}{2(a+b)^2(b-x)} + \frac{2a^2 b^2}{(a-b)^2(a+b)^2(a+x)} - \frac{ab}{2(a-b)^2(b+x)} \right) dx, x, b \tanh(x) \right)}{2b} \\ &= \frac{a \log(1 - \tanh(x))}{4(a+b)^2} - \frac{a \log(1 + \tanh(x))}{4(a-b)^2} + \frac{a^2 b \log(a + b \tanh(x))}{(a^2 - b^2)^2} - \frac{\cosh^2(x)(b-a \tanh(x))}{2(a^2 - b^2)} \end{aligned}$$

Mathematica [A]

time = 0.18, size = 73, normalized size = 0.87

$$\frac{(-a^2 b + b^3) \cosh(2x) + a(-2(a^2 + b^2)x + 4ab \log(a \cosh(x) + b \sinh(x)) + (a^2 - b^2) \sinh(2x))}{4(a-b)^2(a+b)^2}$$

Antiderivative was successfully verified.

`[In] Integrate[Sinh[x]^2/(a + b*Tanh[x]), x]`

`[Out] ((-(a^2*b) + b^3)*Cosh[2*x] + a*(-2*(a^2 + b^2)*x + 4*a*b*Log[a*Cosh[x] + b*Sinh[x]] + (a^2 - b^2)*Sinh[2*x]))/(4*(a - b)^2*(a + b)^2)`

Maple [A]

time = 0.68, size = 145, normalized size = 1.73

method	result
risch	$-\frac{ax}{2(a+b)^2} + \frac{e^{2x}}{8a+8b} - \frac{e^{-2x}}{8(a-b)} - \frac{2a^2bx}{a^4-2a^2b^2+b^4} + \frac{a^2b \ln\left(e^{2x} + \frac{a-b}{a+b}\right)}{a^4-2a^2b^2+b^4}$
default	$\frac{a^2b \ln\left(a \left(\tanh^2\left(\frac{x}{2}\right)\right) + 2b \tanh\left(\frac{x}{2}\right) + a\right)}{(a-b)^2(a+b)^2} + \frac{4}{(8a+8b)\left(\tanh\left(\frac{x}{2}\right)-1\right)^2} + \frac{8}{(16a+16b)\left(\tanh\left(\frac{x}{2}\right)-1\right)} + \frac{a \ln\left(\tanh\left(\frac{x}{2}\right)-1\right)}{2(a+b)^2} - \frac{4}{(8a-8b)\left(\tanh\left(\frac{x}{2}\right)+1\right)^2}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sinh(x)^2/(a+b*tanh(x)), x, method=_RETURNVERBOSE)`

`[Out] a^2*b/(a-b)^2/(a+b)^2*ln(a*tanh(1/2*x)^2+2*b*tanh(1/2*x)+a)+4/(8*a+8*b)/(tanh(1/2*x)-1)^2+8/(16*a+16*b)/(tanh(1/2*x)-1)+1/2*a/(a+b)^2*ln(tanh(1/2*x)-1)`

)-4/(8*a-8*b)/(tanh(1/2*x)+1)^2+8/(16*a-16*b)/(tanh(1/2*x)+1)-1/2*a/(a-b)^2
*ln(tanh(1/2*x)+1)

Maxima [A]

time = 0.26, size = 83, normalized size = 0.99

$$\frac{a^2 b \log\left(-\frac{(a-b)e^{-2x}}{a-b}\right)}{a^4 - 2a^2 b^2 + b^4} - \frac{ax}{2(a^2 + 2ab + b^2)} + \frac{e^{(2x)}}{8(a+b)} - \frac{e^{(-2x)}}{8(a-b)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sinh(x)^2/(a+b*tanh(x)),x, algorithm="maxima")

[Out] a^2*b*log(-(a - b)*e^(-2*x) - a - b)/(a^4 - 2*a^2*b^2 + b^4) - 1/2*a*x/(a^2 + 2*a*b + b^2) + 1/8*e^(2*x)/(a + b) - 1/8*e^(-2*x)/(a - b)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 334 vs. 2(79) = 158.

time = 0.40, size = 334, normalized size = 3.98

$$\frac{(a^2 - a^2b - ab^2 + b^3) \cosh(x)^4 + 4(a^3 - a^2b - ab^2 + b^3) \cosh(x)^3 \sinh(x) + (a^3 - a^2b - ab^2 + b^3) \sinh(x)^4 - 4(a^3 + 2a^2b + ab^2 + b^3) \cosh(x)^2 - a^3 - a^2b + ab^2 + b^3 + 2(3(a^3 - a^2b - ab^2 + b^3) \cosh(x)^2 - 2(a^3 + 2a^2b + ab^2 + b^3) \sinh(x)^2 + 8(a^2b \cosh(x) \sinh(x) + a^2b \sinh(x)^2) \log\left(\frac{2(a \cosh(x) + b \sinh(x))}{\cosh(x) - \sinh(x)}\right) + 4((a^2 - a^2b - ab^2 + b^3) \cosh(x)^2 - 2(a^2 + 2ab + b^2) x \cosh(x)) \sinh(x)}{8((a^2 - 2a^2b + b^3) \cosh(x)^2 + 2(a^2 - 2a^2b + b^3) \cosh(x) \sinh(x) + (a^2 - 2a^2b + b^3) \sinh(x)^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sinh(x)^2/(a+b*tanh(x)),x, algorithm="fricas")

[Out] 1/8*((a^3 - a^2*b - a*b^2 + b^3)*cosh(x)^4 + 4*(a^3 - a^2*b - a*b^2 + b^3)*cosh(x)*sinh(x)^3 + (a^3 - a^2*b - a*b^2 + b^3)*sinh(x)^4 - 4*(a^3 + 2*a^2*b + a*b^2)*x*cosh(x)^2 - a^3 - a^2*b + a*b^2 + b^3 + 2*(3*(a^3 - a^2*b - a*b^2 + b^3)*cosh(x)^2 - 2*(a^3 + 2*a^2*b + a*b^2)*x)*sinh(x)^2 + 8*(a^2*b*cosh(x)^2 + 2*a^2*b*cosh(x)*sinh(x) + a^2*b*sinh(x)^2)*log(2*(a*cosh(x) + b*sinh(x))/(cosh(x) - sinh(x))) + 4*((a^3 - a^2*b - a*b^2 + b^3)*cosh(x)^3 - 2*(a^3 + 2*a^2*b + a*b^2)*x*cosh(x))*sinh(x))/((a^4 - 2*a^2*b^2 + b^4)*cosh(x)^2 + 2*(a^4 - 2*a^2*b^2 + b^4)*cosh(x)*sinh(x) + (a^4 - 2*a^2*b^2 + b^4)*sinh(x)^2)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sinh^2(x)}{a + b \tanh(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sinh(x)**2/(a+b*tanh(x)),x)

[Out] Integral(sinh(x)**2/(a + b*tanh(x)), x)

Giac [A]

time = 0.41, size = 101, normalized size = 1.20

$$\frac{a^2 b \log(|ae^{(2x)} + be^{(2x)} + a - b|)}{a^4 - 2a^2 b^2 + b^4} - \frac{ax}{2(a^2 - 2ab + b^2)} + \frac{(2ae^{(2x)} - a + b)e^{(-2x)}}{8(a^2 - 2ab + b^2)} + \frac{e^{(2x)}}{8(a + b)}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sinh(x)^2/(a+b*tanh(x)),x, algorithm="giac")`

```
[Out] a^2*b*log(abs(a*e^(2*x) + b*e^(2*x) + a - b))/(a^4 - 2*a^2*b^2 + b^4) - 1/2
*a*x/(a^2 - 2*a*b + b^2) + 1/8*(2*a*e^(2*x) - a + b)*e^(-2*x)/(a^2 - 2*a*b
+ b^2) + 1/8*e^(2*x)/(a + b)
```

Mupad [B]

time = 1.43, size = 81, normalized size = 0.96

$$\frac{e^{2x}}{8a + 8b} - \frac{e^{-2x}}{8a - 8b} - \frac{ax}{2(a - b)^2} + \frac{a^2 b \ln(a - b + ae^{2x} + be^{2x})}{a^4 - 2a^2 b^2 + b^4}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sinh(x)^2/(a + b*tanh(x)),x)`

```
[Out] exp(2*x)/(8*a + 8*b) - exp(-2*x)/(8*a - 8*b) - (a*x)/(2*(a - b)^2) + (a^2*b
*log(a - b + a*exp(2*x) + b*exp(2*x)))/(a^4 + b^4 - 2*a^2*b^2)
```

$$3.83 \quad \int \frac{\sinh(x)}{a+b \tanh(x)} dx$$

Optimal. Leaf size=72

$$\frac{ab \operatorname{ArcTan}\left(\frac{b \cosh(x)+a \sinh(x)}{\sqrt{a^2-b^2}}\right)}{(a^2-b^2)^{3/2}} + \frac{a \cosh(x)}{a^2-b^2} - \frac{b \sinh(x)}{a^2-b^2}$$

[Out] $a*b*\arctan((b*\cosh(x)+a*\sinh(x))/(a^2-b^2)^{(1/2)))/(a^2-b^2)^{(3/2)}+a*\cosh(x)/(a^2-b^2)-b*\sinh(x)/(a^2-b^2)$

Rubi [A]

time = 0.09, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.546$, Rules used = {3599, 3188, 2717, 2718, 3153, 212}

$$\frac{ab \operatorname{ArcTan}\left(\frac{a \sinh(x)+b \cosh(x)}{\sqrt{a^2-b^2}}\right)}{(a^2-b^2)^{3/2}} - \frac{b \sinh(x)}{a^2-b^2} + \frac{a \cosh(x)}{a^2-b^2}$$

Antiderivative was successfully verified.

[In] `Int[Sinh[x]/(a + b*Tanh[x]),x]`

[Out] $(a*b*\operatorname{ArcTan}[(b*\operatorname{Cosh}[x] + a*\operatorname{Sinh}[x])/ \operatorname{Sqrt}[a^2 - b^2]])/(a^2 - b^2)^{(3/2)} + (a*\operatorname{Cosh}[x])/(a^2 - b^2) - (b*\operatorname{Sinh}[x])/(a^2 - b^2)$

Rule 212

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rule 2717

`Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

Rule 2718

`Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

Rule 3153

`Int[(cos[(c_.) + (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-1), x_Symbol] := Dist[-d^(-1), Subst[Int[1/(a^2 + b^2 - x^2), x], x, b*Cos[c + d`

`*x] - a*Sin[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0]`

Rule 3188

`Int[(cos[(c_.) + (d_.)*(x_.)]^(m_.)*sin[(c_.) + (d_.)*(x_.)]^(n_.))/(cos[(c_.) + (d_.)*(x_.)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_.)]), x_Symbol] := Dist[b/(a^2 + b^2), Int[Cos[c + d*x]^m*Sin[c + d*x]^(n - 1), x], x] + (Dist[a/(a^2 + b^2), Int[Cos[c + d*x]^(m - 1)*Sin[c + d*x]^n, x], x] - Dist[a*(b/(a^2 + b^2)), Int[Cos[c + d*x]^(m - 1)*(Sin[c + d*x]^(n - 1)/(a*Cos[c + d*x] + b*Sin[c + d*x]), x], x]) /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0] && IGtQ[m, 0] && IGtQ[n, 0]`

Rule 3599

`Int[sin[(e_.) + (f_.)*(x_.)]^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Int[Sin[e + f*x]^m*((a*Cos[e + f*x] + b*Sin[e + f*x])^n/Cos[e + f*x]^n), x] /; FreeQ[{a, b, e, f}, x] && IntegerQ[(m - 1)/2] && ILtQ[n, 0] && ((LtQ[m, 5] && GtQ[n, -4]) || (EqQ[m, 5] && EqQ[n, -1]))`

Rubi steps

$$\begin{aligned} \int \frac{\sinh(x)}{a + b \tanh(x)} dx &= \int \frac{\cosh(x) \sinh(x)}{a \cosh(x) + b \sinh(x)} dx \\ &= \frac{a \int \sinh(x) dx}{a^2 - b^2} - \frac{b \int \cosh(x) dx}{a^2 - b^2} + \frac{(ab) \int \frac{1}{a \cosh(x) + b \sinh(x)} dx}{a^2 - b^2} \\ &= \frac{a \cosh(x)}{a^2 - b^2} - \frac{b \sinh(x)}{a^2 - b^2} + \frac{(iab) \text{Subst}\left(\int \frac{1}{a^2 - b^2 - x^2} dx, x, -ib \cosh(x) - ia \sinh(x)\right)}{a^2 - b^2} \\ &= \frac{ab \tan^{-1}\left(\frac{b \cosh(x) + a \sinh(x)}{\sqrt{a^2 - b^2}}\right)}{(a^2 - b^2)^{3/2}} + \frac{a \cosh(x)}{a^2 - b^2} - \frac{b \sinh(x)}{a^2 - b^2} \end{aligned}$$

Mathematica [A]

time = 0.16, size = 79, normalized size = 1.10

$$\frac{2ab \text{ArcTan}\left(\frac{b + a \tanh\left(\frac{x}{2}\right)}{\sqrt{a - b} \sqrt{a + b}}\right)}{(a - b)^{3/2} (a + b)^{3/2}} + \frac{a \cosh(x)}{a^2 - b^2} + \frac{b \sinh(x)}{-a^2 + b^2}$$

Antiderivative was successfully verified.

`[In] Integrate[Sinh[x]/(a + b*Tanh[x]), x]`

$$\frac{h(x)^2 + 2(a+b)\cosh(x)\sinh(x) + (a+b)\sinh(x)^2 + 2\sqrt{-a^2 + b^2}(\cosh(x) + \sinh(x) - a + b)/((a+b)\cosh(x)^2 + 2(a+b)\cosh(x)\sinh(x) + (a+b)\sinh(x)^2 + a - b))}{((a^4 - 2a^2b^2 + b^4)\cosh(x) + (a^4 - 2a^2b^2 + b^4)\sinh(x))} + \frac{1/2(a^3 + a^2b - ab^2 - b^3 + (a^3 - a^2b - ab^2 + b^3)\cosh(x)^2 + 2(a^3 - a^2b - ab^2 + b^3)\cosh(x)\sinh(x) + (a^3 - a^2b - ab^2 + b^3)\sinh(x)^2 - 4(ab\cosh(x) + ab\sinh(x))\sqrt{a^2 - b^2})\arctan(\sqrt{a^2 - b^2}/((a+b)\cosh(x) + (a+b)\sinh(x)))}{((a^4 - 2a^2b^2 + b^4)\cosh(x) + (a^4 - 2a^2b^2 + b^4)\sinh(x))}$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sinh(x)}{a + b \tanh(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sinh(x)/(a+b*tanh(x)),x)

[Out] Integral(sinh(x)/(a + b*tanh(x)), x)

Giac [A]

time = 0.39, size = 60, normalized size = 0.83

$$\frac{2ab \arctan\left(\frac{ae^x + be^x}{\sqrt{a^2 - b^2}}\right)}{(a^2 - b^2)^{\frac{3}{2}}} + \frac{e^{-x}}{2(a-b)} + \frac{e^x}{2(a+b)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sinh(x)/(a+b*tanh(x)),x, algorithm="giac")

[Out] $2*a*b*\arctan((a*e^x + b*e^x)/\sqrt{a^2 - b^2})/(a^2 - b^2)^{(3/2)} + 1/2*e^{(-x)}/(a - b) + 1/2*e^x/(a + b)$

Mupad [B]

time = 1.36, size = 157, normalized size = 2.18

$$\frac{e^x}{2a + 2b} + \frac{e^{-x}}{2a - 2b} + \frac{2 \operatorname{atan}\left(\frac{ab e^x \sqrt{a^6 - 3a^4 b^2 + 3a^2 b^4 - b^6}}{a^3 \sqrt{a^2 b^2} + b^3 \sqrt{a^2 b^2} - a b^2 \sqrt{a^2 b^2} - a^2 b \sqrt{a^2 b^2}}\right) \sqrt{a^2 b^2}}{\sqrt{a^6 - 3a^4 b^2 + 3a^2 b^4 - b^6}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sinh(x)/(a + b*tanh(x)),x)

[Out] $\exp(x)/(2*a + 2*b) + \exp(-x)/(2*a - 2*b) + (2*\operatorname{atan}((a*b*\exp(x))*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{(1/2)})/(a^3*(a^2*b^2)^{(1/2)} + b^3*(a^2*b^2)^{(1/2)} - a*b^2*(a^2*b^2)^{(1/2)} - a^2*b*(a^2*b^2)^{(1/2)}))*(a^2*b^2)^{(1/2)})/(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{(1/2)}$

3.84 $\int \frac{\operatorname{csch}(x)}{a+b \tanh(x)} dx$

Optimal. Leaf size=52

$$-\frac{b \operatorname{ArcTan}\left(\frac{b \cosh(x)+a \sinh(x)}{\sqrt{a^2-b^2}}\right)}{a \sqrt{a^2-b^2}} - \frac{\tanh^{-1}(\cosh(x))}{a}$$

[Out] $-\operatorname{arctanh}(\cosh(x))/a - b \operatorname{arctan}((b \cosh(x)+a \sinh(x))/(a^2-b^2)^{(1/2)})/a/(a^2-b^2)^{(1/2)}$

Rubi [A]

time = 0.10, antiderivative size = 52, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.454$, Rules used = {3599, 3189, 3855, 3153, 212}

$$-\frac{b \operatorname{ArcTan}\left(\frac{a \sinh(x)+b \cosh(x)}{\sqrt{a^2-b^2}}\right)}{a \sqrt{a^2-b^2}} - \frac{\tanh^{-1}(\cosh(x))}{a}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[\operatorname{Csch}[x]/(a + b \operatorname{Tanh}[x]), x]$

[Out] $-\left(\frac{b \operatorname{ArcTan}[b \operatorname{Cosh}[x] + a \operatorname{Sinh}[x]]/\operatorname{Sqrt}[a^2 - b^2]}{a \operatorname{Sqrt}[a^2 - b^2]}\right) - \operatorname{ArcTanh}[\operatorname{Cosh}[x]]/a$

Rule 212

$\operatorname{Int}[(a_.) + (b_.) \cdot (x_.)^2]^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2] \operatorname{Rt}[-b, 2])) \cdot \operatorname{ArcTanh}[\operatorname{Rt}[-b, 2] \cdot (x/\operatorname{Rt}[a, 2])], x] /;$ $\operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{Gt} Q[a, 0] \ || \ \operatorname{Lt} Q[b, 0])$

Rule 3153

$\operatorname{Int}[(\cos[(c_.) + (d_.) \cdot (x_.)]) \cdot (a_.) + (b_.) \cdot \sin[(c_.) + (d_.) \cdot (x_.)])^{-1}, x_Symbol] \rightarrow \operatorname{Dist}[-d^{-1}, \operatorname{Subst}[\operatorname{Int}[1/(a^2 + b^2 - x^2), x], x, b \cos[c + d \cdot x] - a \sin[c + d \cdot x]], x] /;$ $\operatorname{FreeQ}\{a, b, c, d\}, x \ \&\& \operatorname{NeQ}[a^2 + b^2, 0]$

Rule 3189

$\operatorname{Int}[(\cos[(c_.) + (d_.) \cdot (x_.)])^{(m_.)} \cdot \sin[(c_.) + (d_.) \cdot (x_.)]^{(n_.)})/(\cos[(c_.) + (d_.) \cdot (x_.)] \cdot (a_.) + (b_.) \cdot \sin[(c_.) + (d_.) \cdot (x_.)]), x_Symbol] \rightarrow \operatorname{Int}[\operatorname{ExpandTrig}[\cos[c + d \cdot x]^m \cdot \sin[c + d \cdot x]^n / (a \cos[c + d \cdot x] + b \sin[c + d \cdot x])], x] /;$ $\operatorname{FreeQ}\{a, b, c, d, m, n\}, x \ \&\& \operatorname{Integers} Q[m, n]$

Rule 3599

```
Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Int[Sin[e + f*x]^m*((a*Cos[e + f*x] + b*Sin[e + f*x])^n/Cos[e + f*x]^n), x] /; FreeQ[{a, b, e, f}, x] && IntegerQ[(m - 1)/2] && ILtQ[n, 0] && ((LtQ[m, 5] && GtQ[n, -4]) || (EqQ[m, 5] && EqQ[n, -1]))
```

Rule 3855

```
Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned}
 \int \frac{\operatorname{csch}(x)}{a + b \tanh(x)} dx &= \int \frac{\operatorname{coth}(x)}{a \cosh(x) + b \sinh(x)} dx \\
 &= i \int \left(-\frac{i \operatorname{csch}(x)}{a} + \frac{ib}{a(a \cosh(x) + b \sinh(x))} \right) dx \\
 &= \frac{\int \operatorname{csch}(x) dx}{a} - \frac{b \int \frac{1}{a \cosh(x) + b \sinh(x)} dx}{a} \\
 &= -\frac{\tanh^{-1}(\cosh(x))}{a} - \frac{(ib) \operatorname{Subst}\left(\int \frac{1}{a^2 - b^2 - x^2} dx, x, -ib \cosh(x) - ia \sinh(x)\right)}{a} \\
 &= -\frac{b \tan^{-1}\left(\frac{b \cosh(x) + a \sinh(x)}{\sqrt{a^2 - b^2}}\right)}{a \sqrt{a^2 - b^2}} - \frac{\tanh^{-1}(\cosh(x))}{a}
 \end{aligned}$$

Mathematica [A]

time = 0.07, size = 59, normalized size = 1.13

$$\frac{-2b \operatorname{ArcTan}\left(\frac{b + a \tanh\left(\frac{x}{2}\right)}{\sqrt{a-b} \sqrt{a+b}}\right)}{\sqrt{a-b} \sqrt{a+b}} + \log\left(\tanh\left(\frac{x}{2}\right)\right)$$

Antiderivative was successfully verified.

```
[In] Integrate[Csch[x]/(a + b*Tanh[x]), x]
```

```
[Out] ((-2*b*ArcTan[(b + a*Tanh[x/2])/(Sqrt[a - b]*Sqrt[a + b])])/(Sqrt[a - b]*Sqrt[a + b]) + Log[Tanh[x/2]])/a
```

Maple [A]

time = 0.74, size = 53, normalized size = 1.02

method	result	size
default	$-\frac{2b \arctan\left(\frac{2a \tanh\left(\frac{x}{2}\right) + 2b}{2\sqrt{a^2 - b^2}}\right)}{a\sqrt{a^2 - b^2}} + \frac{\ln(\tanh(\frac{x}{2}))}{a}$	53
risch	$-\frac{\ln(e^x + 1)}{a} + \frac{\ln(e^x - 1)}{a} - \frac{b \ln\left(e^x + \frac{a-b}{\sqrt{-a^2 + b^2}}\right)}{\sqrt{-a^2 + b^2} a} + \frac{b \ln\left(e^x - \frac{a-b}{\sqrt{-a^2 + b^2}}\right)}{\sqrt{-a^2 + b^2} a}$	97

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(csch(x)/(a+b*tanh(x)),x,method=_RETURNVERBOSE)
```

```
[Out] -2/a*b/(a^2-b^2)^(1/2)*arctan(1/2*(2*a*tanh(1/2*x)+2*b)/(a^2-b^2)^(1/2))+1/a*ln(tanh(1/2*x))
```

Maxima [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csch(x)/(a+b*tanh(x)),x, algorithm="maxima")
```

```
[Out] Exception raised: ValueError >> Computation failed since Maxima requested a
dditional constraints; using the 'assume' command before evaluation *may* h
elp (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see 'assume?' for
more de
```

Fricas [A]

time = 0.39, size = 237, normalized size = 4.56

$$\left[\frac{\sqrt{-a^2 + b^2} b \log\left(\frac{(a+b)\cosh(x)^2 + 2(a+b)\cosh(x)\sinh(x) + (a-b)\sinh(x)^2 + \sqrt{-a^2 + b^2}\cosh(x)\sinh(x) - a^2}{(a+b)\cosh(x)^2 + 2(a+b)\cosh(x)\sinh(x) + (a-b)\sinh(x)^2 + a - b}\right) + (a^2 - b^2) \log(\cosh(x) + \sinh(x) + 1) - (a^2 - b^2) \log(\cosh(x) + \sinh(x) - 1)}{a^3 - ab^2}, \frac{2\sqrt{a^2 - b^2} b \arctan\left(\frac{\sqrt{a^2 - b^2}}{(a+b)\cosh(x) + (a-b)\sinh(x)}\right) - (a^2 - b^2) \log(\cosh(x) + \sinh(x) + 1) + (a^2 - b^2) \log(\cosh(x) + \sinh(x) - 1)}{a^3 - ab^2} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csch(x)/(a+b*tanh(x)),x, algorithm="fricas")
```

```
[Out] [-(sqrt(-a^2 + b^2)*b*log(((a + b)*cosh(x)^2 + 2*(a + b)*cosh(x)*sinh(x) +
(a + b)*sinh(x)^2 + 2*sqrt(-a^2 + b^2)*(cosh(x) + sinh(x)) - a + b)/((a + b
)*cosh(x)^2 + 2*(a + b)*cosh(x)*sinh(x) + (a + b)*sinh(x)^2 + a - b)) + (a^
2 - b^2)*log(cosh(x) + sinh(x) + 1) - (a^2 - b^2)*log(cosh(x) + sinh(x) - 1
))/(a^3 - a*b^2), (2*sqrt(a^2 - b^2)*b*arctan(sqrt(a^2 - b^2)/((a + b)*cosh
(x) + (a + b)*sinh(x))) - (a^2 - b^2)*log(cosh(x) + sinh(x) + 1) + (a^2 - b
^2)*log(cosh(x) + sinh(x) - 1))/(a^3 - a*b^2)]
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{csch}(x)}{a + b \tanh(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(csch(x)/(a+b*tanh(x)),x)``[Out] Integral(csch(x)/(a + b*tanh(x)), x)`**Giac [A]**

time = 0.41, size = 60, normalized size = 1.15

$$-\frac{2b \arctan\left(\frac{ae^x + be^x}{\sqrt{a^2 - b^2}}\right)}{\sqrt{a^2 - b^2} a} - \frac{\log(e^x + 1)}{a} + \frac{\log(|e^x - 1|)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(csch(x)/(a+b*tanh(x)),x, algorithm="giac")``[Out] -2*b*arctan((a*e^x + b*e^x)/sqrt(a^2 - b^2))/(sqrt(a^2 - b^2)*a) - log(e^x + 1)/a + log(abs(e^x - 1))/a`**Mupad [B]**

time = 1.40, size = 177, normalized size = 3.40

$$\frac{\ln(32ab - 32a^2 + 32a^2e^x - 32abe^x)}{a} - \frac{\ln(32ab - 32a^2 - 32a^2e^x + 32abe^x)}{a} - \frac{b \ln\left(\frac{32ab^2e^x + 32a^2be^x - 32ab\sqrt{b^2 - a^2}}{a^2 - a^3}\right) \sqrt{b^2 - a^2}}{a^2 - a^3} + \frac{b \ln\left(\frac{32ab^2e^x + 32a^2be^x + 32ab\sqrt{b^2 - a^2}}{a^2 - a^3}\right) \sqrt{b^2 - a^2}}{a^2 - a^3}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/(sinh(x)*(a + b*tanh(x))),x)`

```
[Out] log(32*a*b - 32*a^2 + 32*a^2*exp(x) - 32*a*b*exp(x))/a - log(32*a*b - 32*a^2 - 32*a^2*exp(x) + 32*a*b*exp(x))/a - (b*log(32*a*b^2*exp(x) + 32*a^2*b*exp(x) - 32*a*b*(b^2 - a^2)^(1/2))*(b^2 - a^2)^(1/2))/(a*b^2 - a^3) + (b*log(32*a*b^2*exp(x) + 32*a^2*b*exp(x) + 32*a*b*(b^2 - a^2)^(1/2))*(b^2 - a^2)^(1/2))/(a*b^2 - a^3)
```

3.85 $\int \frac{\operatorname{csch}^2(x)}{a+b \tanh(x)} dx$

Optimal. Leaf size=29

$$-\frac{\operatorname{coth}(x)}{a} - \frac{b \log(\tanh(x))}{a^2} + \frac{b \log(a + b \tanh(x))}{a^2}$$

[Out] $-\operatorname{coth}(x)/a - b \cdot \ln(\tanh(x))/a^2 + b \cdot \ln(a + b \cdot \tanh(x))/a^2$

Rubi [A]

time = 0.04, antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {3597, 46}

$$-\frac{b \log(\tanh(x))}{a^2} + \frac{b \log(a + b \tanh(x))}{a^2} - \frac{\operatorname{coth}(x)}{a}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Csch}[x]^2/(a + b \cdot \text{Tanh}[x]), x]$

[Out] $-(\text{Coth}[x]/a) - (b \cdot \text{Log}[\text{Tanh}[x]])/a^2 + (b \cdot \text{Log}[a + b \cdot \text{Tanh}[x]])/a^2$

Rule 46

$\text{Int}[(a_+ + (b_+)(x_+))^{m_+}((c_+ + (d_+)(x_+))^{n_+}), x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b \cdot x)^m(c + d \cdot x)^n, x], x] /; \text{FreeQ}\{a, b, c, d, x\} \ \&\& \ \text{NeQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \text{ILtQ}[m, 0] \ \&\& \ \text{IntegerQ}[n] \ \&\& \ !(\text{IGtQ}[n, 0] \ \&\& \ \text{LtQ}[m + n + 2, 0])$

Rule 3597

$\text{Int}[\sin[(e_+ + (f_+)(x_+))^{m_+}((a_+ + (b_+)\tan[(e_+ + (f_+)(x_+))^{n_+})]), x_Symbol] \rightarrow \text{Dist}[b/f, \text{Subst}[\text{Int}[x^m((a + x)^n/(b^2 + x^2)^{(m/2 + 1)}), x], x, b \cdot \text{Tan}[e + f \cdot x]], x] /; \text{FreeQ}\{a, b, e, f, n\}, x\} \ \&\& \ \text{IntegerQ}[m/2]$

Rubi steps

$$\begin{aligned} \int \frac{\operatorname{csch}^2(x)}{a+b \tanh(x)} dx &= b \operatorname{Subst} \left(\int \frac{1}{x^2(a+x)} dx, x, b \tanh(x) \right) \\ &= b \operatorname{Subst} \left(\int \left(\frac{1}{ax^2} - \frac{1}{a^2x} + \frac{1}{a^2(a+x)} \right) dx, x, b \tanh(x) \right) \\ &= -\frac{\operatorname{coth}(x)}{a} - \frac{b \log(\tanh(x))}{a^2} + \frac{b \log(a + b \tanh(x))}{a^2} \end{aligned}$$

Mathematica [A]

time = 0.07, size = 28, normalized size = 0.97

$$\frac{a \coth(x) + b \log(\sinh(x)) - b \log(a \cosh(x) + b \sinh(x))}{a^2}$$

Antiderivative was successfully verified.

`[In] Integrate[Csch[x]^2/(a + b*Tanh[x]), x]``[Out] -((a*Coth[x] + b*Log[Sinh[x]] - b*Log[a*Cosh[x] + b*Sinh[x]])/a^2)`**Maple [A]**

time = 0.74, size = 56, normalized size = 1.93

method	result	size
risch	$-\frac{2}{a(e^{2x}-1)} + \frac{b \ln(e^{2x} + \frac{a-b}{a+b})}{a^2} - \frac{b \ln(e^{2x}-1)}{a^2}$	50
default	$-\frac{\tanh(\frac{x}{2})}{2a} + \frac{b \ln(a(\tanh^2(\frac{x}{2})) + 2b \tanh(\frac{x}{2}) + a)}{a^2} - \frac{1}{2a \tanh(\frac{x}{2})} - \frac{b \ln(\tanh(\frac{x}{2}))}{a^2}$	56

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(csch(x)^2/(a+b*tanh(x)), x, method=_RETURNVERBOSE)``[Out] -1/2/a*tanh(1/2*x)+b/a^2*ln(a*tanh(1/2*x)^2+2*b*tanh(1/2*x)+a)-1/2/a/tanh(1/2*x)-b/a^2*ln(tanh(1/2*x))`**Maxima [B]** Leaf count of result is larger than twice the leaf count of optimal. 65 vs. 2(29) = 58.

time = 0.25, size = 65, normalized size = 2.24

$$\frac{b \log(-(a-b)e^{(-2x)} - a - b)}{a^2} - \frac{b \log(e^{(-x)} + 1)}{a^2} - \frac{b \log(e^{(-x)} - 1)}{a^2} + \frac{2}{ae^{(-2x)} - a}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(csch(x)^2/(a+b*tanh(x)), x, algorithm="maxima")``[Out] b*log(-(a - b)*e^(-2*x) - a - b)/a^2 - b*log(e^(-x) + 1)/a^2 - b*log(e^(-x) - 1)/a^2 + 2/(a*e^(-2*x) - a)`**Fricas [B]** Leaf count of result is larger than twice the leaf count of optimal. 122 vs. 2(29) = 58.

time = 0.36, size = 122, normalized size = 4.21

$$\frac{(b \cosh(x)^2 + 2b \cosh(x) \sinh(x) + b \sinh(x)^2 - b) \log\left(\frac{2(a \cosh(x) + b \sinh(x))}{\cosh(x) - \sinh(x)}\right) - (b \cosh(x)^2 + 2b \cosh(x) \sinh(x) + b \sinh(x)^2 - b) \log\left(\frac{2 \sinh(x)}{\cosh(x) - \sinh(x)}\right) - 2a}{a^2 \cosh(x)^2 + 2a^2 \cosh(x) \sinh(x) + a^2 \sinh(x)^2 - a^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csch(x)^2/(a+b*tanh(x)),x, algorithm="fricas")

[Out] ((b*cosh(x)^2 + 2*b*cosh(x)*sinh(x) + b*sinh(x)^2 - b)*log(2*(a*cosh(x) + b*sinh(x))/(cosh(x) - sinh(x))) - (b*cosh(x)^2 + 2*b*cosh(x)*sinh(x) + b*sinh(x)^2 - b)*log(2*sinh(x)/(cosh(x) - sinh(x)))) - 2*a/(a^2*cosh(x)^2 + 2*a^2*cosh(x)*sinh(x) + a^2*sinh(x)^2 - a^2)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{csch}^2(x)}{a + b \tanh(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csch(x)**2/(a+b*tanh(x)),x)

[Out] Integral(csch(x)**2/(a + b*tanh(x)), x)

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 78 vs. 2(29) = 58.

time = 0.42, size = 78, normalized size = 2.69

$$\frac{(ab + b^2) \log(|ae^{(2x)} + be^{(2x)} + a - b|)}{a^3 + a^2b} - \frac{b \log(|e^{(2x)} - 1|)}{a^2} + \frac{be^{(2x)} - 2a - b}{a^2(e^{(2x)} - 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csch(x)^2/(a+b*tanh(x)),x, algorithm="giac")

[Out] (a*b + b^2)*log(abs(a*e^(2*x) + b*e^(2*x) + a - b))/(a^3 + a^2*b) - b*log(abs(e^(2*x) - 1))/a^2 + (b*e^(2*x) - 2*a - b)/(a^2*(e^(2*x) - 1))

Mupad [B]

time = 1.51, size = 323, normalized size = 11.14

$$\frac{2 \operatorname{atan}\left(\frac{(a^2 (b^2)^{3/2} - a^2 \sqrt{b^2}) (a b^2 \sqrt{-a^2 - b^2} \sqrt{-a^2 + a^2 b^2} \sqrt{-a^2 - a^2 b^2} \sqrt{-a^2 + a^2 b^2} \sqrt{-a^2 - 2 a^2 b^2 + \sqrt{-a^2 + a^2 b^2} \sqrt{-a^2}}{-3 a^2 b^2 + 3 a^2 b^2 - 3 a^2 b^2 + a^2 b^2}) (a b^2 \sqrt{-a^2 - b^2} \sqrt{-a^2 + a^2 b^2} \sqrt{-a^2 - a^2 b^2} \sqrt{-a^2 + a^2 b^2} \sqrt{-a^2 - 2 a^2 b^2 + \sqrt{-a^2 + a^2 b^2} \sqrt{-a^2}})}{\sqrt{-a^2}}\right) \sqrt{b^2}}{a (e^{2x} - 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(sinh(x)^2*(a + b*tanh(x))),x)

[Out] (2*atan((b*(a^4*(b^2)^(3/2) - a^6*(b^2)^(1/2))*(a*b^5*(-a^4)^(1/2) - b^6*(-a^4)^(1/2) + a^2*b^4*(-a^4)^(1/2) - a^3*b^3*(-a^4)^(1/2) + b^6*exp(2*x)*(-a^4)^(1/2) - 2*a^2*b^4*exp(2*x)*(-a^4)^(1/2) + a^4*b^2*exp(2*x)*(-a^4)^(1/2)) + b^2*(a^3*(b^2)^(3/2) - a^5*(b^2)^(1/2))*(a*b^5*(-a^4)^(1/2) - b^6*(-a^4)^(1/2) + a^2*b^4*(-a^4)^(1/2) - a^3*b^3*(-a^4)^(1/2) + b^6*exp(2*x)*(-a^4)^(1/2) - 2*a^2*b^4*exp(2*x)*(-a^4)^(1/2) + a^4*b^2*exp(2*x)*(-a^4)^(1/2)))/(a^6*b^10 - 3*a^8*b^8 + 3*a^10*b^6 - a^12*b^4)*(b^2)^(1/2)/(-a^4)^(1/2) - 2/(a*(exp(2*x) - 1))

3.86 $\int \frac{\operatorname{csch}^3(x)}{a+b \tanh(x)} dx$

Optimal. Leaf size=82

$$\frac{b\sqrt{a^2-b^2} \operatorname{ArcTan}\left(\frac{b \cosh(x)+a \sinh(x)}{\sqrt{a^2-b^2}}\right)}{a^3} + \frac{\tanh^{-1}(\cosh(x))}{2a} - \frac{b^2 \tanh^{-1}(\cosh(x))}{a^3} + \frac{b \operatorname{csch}(x)}{a^2} - \frac{\coth(x) \operatorname{csch}(x)}{2a}$$

[Out] 1/2*arctanh(cosh(x))/a-b^2*arctanh(cosh(x))/a^3+b*csch(x)/a^2-1/2*coth(x)*csch(x)/a+b*arctan((b*cosh(x)+a*sinh(x))/(a^2-b^2)^(1/2))*(a^2-b^2)^(1/2)/a^3

Rubi [A]

time = 0.21, antiderivative size = 82, normalized size of antiderivative = 1.00, number of steps used = 15, number of rules used = 11, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.846$, Rules used = {3599, 3189, 3853, 3855, 2701, 327, 213, 2702, 3183, 3153, 212}

$$-\frac{b^2 \tanh^{-1}(\cosh(x))}{a^3} + \frac{b \operatorname{csch}(x)}{a^2} + \frac{b\sqrt{a^2-b^2} \operatorname{ArcTan}\left(\frac{a \sinh(x)+b \cosh(x)}{\sqrt{a^2-b^2}}\right)}{a^3} + \frac{\tanh^{-1}(\cosh(x))}{2a} - \frac{\coth(x) \operatorname{csch}(x)}{2a}$$

Antiderivative was successfully verified.

[In] Int[Csch[x]^3/(a + b*Tanh[x]),x]

[Out] (b*Sqrt[a^2 - b^2]*ArcTan[(b*Cosh[x] + a*Sinh[x])/Sqrt[a^2 - b^2]])/a^3 + ArcTanh[Cosh[x]]/(2*a) - (b^2*ArcTanh[Cosh[x]])/a^3 + (b*Csch[x])/a^2 - (Coth[x]*Csch[x])/(2*a)

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 213

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1)*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])

Rule 327

Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n-1)*(c*x)^(m-n+1)*((a + b*x^n)^(p+1)/(b*(m+n*p+1))), x] - Dist[a*c^n*((m-n+1)/(b*(m+n*p+1))), Int[(c*x)^(m-n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n-1] && NeQ[m+n*p

+ 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 2701

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*sec[(e_.) + (f_.)*(x_.)]^(n_), x_Symbol] := Dist[-(f*a^n)^(-1), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2), x], x, a*Csc[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rule 2702

Int[csc[(e_.) + (f_.)*(x_.)]^(n_)*((a_.)*sec[(e_.) + (f_.)*(x_.)])^(m_), x_Symbol] := Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2), x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rule 3153

Int[(cos[(c_.) + (d_.)*(x_.)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_.)])^(-1), x_Symbol] := Dist[-d^(-1), Subst[Int[1/(a^2 + b^2 - x^2), x], x, b*Cos[c + d*x] - a*Sin[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0]

Rule 3183

Int[cos[(c_.) + (d_.)*(x_.)]^(m_)/(cos[(c_.) + (d_.)*(x_.)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_.)]), x_Symbol] := Simp[-Cos[c + d*x]^(m + 1)/(b*d*(m + 1)), x] + (-Dist[a/b^2, Int[Cos[c + d*x]^(m + 1), x], x] + Dist[(a^2 + b^2)/b^2, Int[Cos[c + d*x]^(m + 2)/(a*cos[c + d*x] + b*sin[c + d*x]), x], x]) /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]

Rule 3189

Int[(cos[(c_.) + (d_.)*(x_.)]^(m_)*sin[(c_.) + (d_.)*(x_.)]^(n_))/(cos[(c_.) + (d_.)*(x_.)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_.)]), x_Symbol] := Int[ExpandTrig[cos[c + d*x]^m*(sin[c + d*x]^n/(a*cos[c + d*x] + b*sin[c + d*x])), x], x] /; FreeQ[{a, b, c, d, m, n}, x] && IntegersQ[m, n]

Rule 3599

Int[sin[(e_.) + (f_.)*(x_.)]^(m_)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] := Int[Sin[e + f*x]^m*((a*cos[e + f*x] + b*sin[e + f*x])^n/Cos[e + f*x]^n), x] /; FreeQ[{a, b, e, f}, x] && IntegerQ[(m - 1)/2] && ILtQ[n, 0] && ((LtQ[m, 5] && GtQ[n, -4]) || (EqQ[m, 5] && EqQ[n, -1]))

Rule 3853

```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Dist[b^2*((n - 2)/(n - 1)), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]
```

Rule 3855

```
Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned}
\int \frac{\operatorname{csch}^3(x)}{a + b \tanh(x)} dx &= \int \frac{\operatorname{coth}(x) \operatorname{csch}^2(x)}{a \cosh(x) + b \sinh(x)} dx \\
&= - \left(i \int \left(\frac{i \operatorname{csch}^3(x)}{a} - \frac{i b \operatorname{csch}^2(x) \operatorname{sech}(x)}{a^2} + \frac{i b^2 \operatorname{csch}(x) \operatorname{sech}^2(x)}{a^3} - \frac{i b^3 \operatorname{sech}^2(x)}{a^3 (a \cosh(x) + b \sinh(x))} \right) dx \right) \\
&= \frac{\int \operatorname{csch}^3(x) dx}{a} - \frac{b \int \operatorname{csch}^2(x) \operatorname{sech}(x) dx}{a^2} + \frac{b^2 \int \operatorname{csch}(x) \operatorname{sech}^2(x) dx}{a^3} - \frac{b^3 \int \frac{\operatorname{sech}^2(x)}{a \cosh(x) + b \sinh(x)} dx}{a^3} \\
&= -\frac{\operatorname{coth}(x) \operatorname{csch}(x)}{2a} - \frac{b^2 \operatorname{sech}(x)}{a^3} - \frac{\int \operatorname{csch}(x) dx}{2a} + \frac{(ib) \operatorname{Subst} \left(\int \frac{x^2}{-1+x^2} dx, x, -i \operatorname{csch}(x) \right)}{a^2} \\
&= -\frac{b \tan^{-1}(\sinh(x))}{a^2} + \frac{\tanh^{-1}(\cosh(x))}{2a} + \frac{b \operatorname{csch}(x)}{a^2} - \frac{\operatorname{coth}(x) \operatorname{csch}(x)}{2a} + \frac{(ib) \operatorname{Subst} \left(\int \frac{x^2}{-1+x^2} dx, x, -i \operatorname{csch}(x) \right)}{a^2} \\
&= \frac{b \sqrt{a^2 - b^2} \tan^{-1} \left(\frac{b \cosh(x) + a \sinh(x)}{\sqrt{a^2 - b^2}} \right)}{a^3} + \frac{\tanh^{-1}(\cosh(x))}{2a} - \frac{b^2 \tanh^{-1}(\cosh(x))}{a^3} + \frac{b \operatorname{csch}(x)}{a^2}
\end{aligned}$$

Mathematica [A]

time = 0.26, size = 123, normalized size = 1.50

$$\frac{-16\sqrt{a-b} b \sqrt{a+b} \operatorname{ArcTan} \left(\frac{b+a \tanh\left(\frac{x}{2}\right)}{\sqrt{a-b} \sqrt{a+b}} \right) - 4ab \operatorname{coth}\left(\frac{x}{2}\right) + a^2 \operatorname{csch}^2\left(\frac{x}{2}\right) + 4a^2 \log\left(\tanh\left(\frac{x}{2}\right)\right) - 8b^2 \log\left(\tanh\left(\frac{x}{2}\right)\right) + a^2 \operatorname{sech}^2\left(\frac{x}{2}\right) + 4ab \tanh\left(\frac{x}{2}\right)}{8a^3}$$

Antiderivative was successfully verified.

```
[In] Integrate[Csch[x]^3/(a + b*Tanh[x]), x]
```

```
[Out] -1/8*(-16*sqrt[a - b]*b*sqrt[a + b]*ArcTan[(b + a*Tanh[x/2])/(sqrt[a - b]*sqrt[a + b])] - 4*a*b*Coth[x/2] + a^2*Csch[x/2]^2 + 4*a^2*Log[Tanh[x/2]] - 8*b^2*Log[Tanh[x/2]] + a^2*Sech[x/2]^2 + 4*a*b*Tanh[x/2])/a^3
```

Maple [A]

time = 0.88, size = 110, normalized size = 1.34

method	result
default	$\frac{a \left(\tanh^2\left(\frac{x}{2}\right) \right) - 2b \tanh\left(\frac{x}{2}\right)}{4a^2} + \frac{2b\sqrt{a^2 - b^2} \arctan\left(\frac{2a \tanh\left(\frac{x}{2}\right) + 2b}{2\sqrt{a^2 - b^2}}\right)}{a^3} - \frac{1}{8a \tanh\left(\frac{x}{2}\right)^2} + \frac{(-2a^2 + 4b^2) \ln\left(\tanh\left(\frac{x}{2}\right)\right)}{4a^3} + \frac{1}{2a^2 \tanh\left(\frac{x}{2}\right)}$
risch	$-\frac{e^x (e^{2x} a - 2b e^{2x} + a + 2b)}{(e^{2x} - 1)^2 a^2} + \frac{\sqrt{-a^2 + b^2} b \ln\left(e^x + \frac{\sqrt{-a^2 + b^2}}{a+b}\right)}{a^3} - \frac{\sqrt{-a^2 + b^2} b \ln\left(e^x - \frac{\sqrt{-a^2 + b^2}}{a+b}\right)}{a^3} - \frac{1}{8a \tanh\left(\frac{x}{2}\right)^2} + \frac{(-2a^2 + 4b^2) \ln\left(\tanh\left(\frac{x}{2}\right)\right)}{4a^3} + \frac{1}{2a^2 \tanh\left(\frac{x}{2}\right)}$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(csch(x)^3/(a+b*tanh(x)),x,method=_RETURNVERBOSE)
```

```
[Out] 1/4/a^2*(1/2*a*tanh(1/2*x)^2-2*b*tanh(1/2*x))+2*b*(a^2-b^2)^(1/2)/a^3*arctan(1/2*(2*a*tanh(1/2*x)+2*b)/(a^2-b^2)^(1/2))-1/8/a/tanh(1/2*x)^2+1/4/a^3*(-2*a^2+4*b^2)*ln(tanh(1/2*x))+1/2*b/a^2/tanh(1/2*x)
```

Maxima [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csch(x)^3/(a+b*tanh(x)),x, algorithm="maxima")
```

```
[Out] Exception raised: ValueError >> Computation failed since Maxima requested a
dditional constraints; using the 'assume' command before evaluation *may* h
elp (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see 'assume?' for
more de
```

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 555 vs. 2(74) = 148.

time = 0.38, size = 1165, normalized size = 14.21

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csch(x)^3/(a+b*tanh(x)),x, algorithm="fricas")
```

```
[Out] [-1/2*(2*(a^2 - 2*a*b)*cosh(x)^3 + 6*(a^2 - 2*a*b)*cosh(x)*sinh(x)^2 + 2*(a
^2 - 2*a*b)*sinh(x)^3 - 2*(b*cosh(x)^4 + 4*b*cosh(x)*sinh(x)^3 + b*sinh(x)^
4 - 2*b*cosh(x)^2 + 2*(3*b*cosh(x)^2 - b)*sinh(x)^2 + 4*(b*cosh(x)^3 - b*co
sh(x))*sinh(x) + b)*sqrt(-a^2 + b^2)*log(((a + b)*cosh(x)^2 + 2*(a + b)*co
sh(x)*sinh(x) + (a + b)*sinh(x)^2 + 2*sqrt(-a^2 + b^2)*(cosh(x) + sinh(x)) -
a + b)/((a + b)*cosh(x)^2 + 2*(a + b)*cosh(x)*sinh(x) + (a + b)*sinh(x)^2
+ a - b)) + 2*(a^2 + 2*a*b)*cosh(x) - ((a^2 - 2*b^2)*cosh(x)^4 + 4*(a^2 - 2
```

```

*b^2)*cosh(x)*sinh(x)^3 + (a^2 - 2*b^2)*sinh(x)^4 - 2*(a^2 - 2*b^2)*cosh(x)
^2 + 2*(3*(a^2 - 2*b^2)*cosh(x)^2 - a^2 + 2*b^2)*sinh(x)^2 + a^2 - 2*b^2 +
4*((a^2 - 2*b^2)*cosh(x)^3 - (a^2 - 2*b^2)*cosh(x))*sinh(x))*log(cosh(x) +
sinh(x) + 1) + ((a^2 - 2*b^2)*cosh(x)^4 + 4*(a^2 - 2*b^2)*cosh(x)*sinh(x)^3
+ (a^2 - 2*b^2)*sinh(x)^4 - 2*(a^2 - 2*b^2)*cosh(x)^2 + 2*(3*(a^2 - 2*b^2)
*cosh(x)^2 - a^2 + 2*b^2)*sinh(x)^2 + a^2 - 2*b^2 + 4*((a^2 - 2*b^2)*cosh(x)
)^3 - (a^2 - 2*b^2)*cosh(x))*sinh(x))*log(cosh(x) + sinh(x) - 1) + 2*(3*(a^
2 - 2*a*b)*cosh(x)^2 + a^2 + 2*a*b)*sinh(x))/(a^3*cosh(x)^4 + 4*a^3*cosh(x)
*sinh(x)^3 + a^3*sinh(x)^4 - 2*a^3*cosh(x)^2 + a^3 + 2*(3*a^3*cosh(x)^2 - a
^3)*sinh(x)^2 + 4*(a^3*cosh(x)^3 - a^3*cosh(x))*sinh(x)), -1/2*(2*(a^2 - 2*
a*b)*cosh(x)^3 + 6*(a^2 - 2*a*b)*cosh(x)*sinh(x)^2 + 2*(a^2 - 2*a*b)*sinh(x)
)^3 + 4*(b*cosh(x)^4 + 4*b*cosh(x)*sinh(x)^3 + b*sinh(x)^4 - 2*b*cosh(x)^2
+ 2*(3*b*cosh(x)^2 - b)*sinh(x)^2 + 4*(b*cosh(x)^3 - b*cosh(x))*sinh(x) + b
)*sqrt(a^2 - b^2)*arctan(sqrt(a^2 - b^2)/((a + b)*cosh(x) + (a + b)*sinh(x)
)) + 2*(a^2 + 2*a*b)*cosh(x) - ((a^2 - 2*b^2)*cosh(x)^4 + 4*(a^2 - 2*b^2)*c
osh(x)*sinh(x)^3 + (a^2 - 2*b^2)*sinh(x)^4 - 2*(a^2 - 2*b^2)*cosh(x)^2 + 2*
(3*(a^2 - 2*b^2)*cosh(x)^2 - a^2 + 2*b^2)*sinh(x)^2 + a^2 - 2*b^2 + 4*((a^2
- 2*b^2)*cosh(x)^3 - (a^2 - 2*b^2)*cosh(x))*sinh(x))*log(cosh(x) + sinh(x)
+ 1) + ((a^2 - 2*b^2)*cosh(x)^4 + 4*(a^2 - 2*b^2)*cosh(x)*sinh(x)^3 + (a^2
- 2*b^2)*sinh(x)^4 - 2*(a^2 - 2*b^2)*cosh(x)^2 + 2*(3*(a^2 - 2*b^2)*cosh(x)
)^2 - a^2 + 2*b^2)*sinh(x)^2 + a^2 - 2*b^2 + 4*((a^2 - 2*b^2)*cosh(x)^3 - (
a^2 - 2*b^2)*cosh(x))*sinh(x))*log(cosh(x) + sinh(x) - 1) + 2*(3*(a^2 - 2*a
*b)*cosh(x)^2 + a^2 + 2*a*b)*sinh(x))/(a^3*cosh(x)^4 + 4*a^3*cosh(x)*sinh(x)
)^3 + a^3*sinh(x)^4 - 2*a^3*cosh(x)^2 + a^3 + 2*(3*a^3*cosh(x)^2 - a^3)*sin
h(x)^2 + 4*(a^3*cosh(x)^3 - a^3*cosh(x))*sinh(x))]

```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{csch}^3(x)}{a + b \tanh(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csch(x)**3/(a+b*tanh(x)),x)

[Out] Integral(csch(x)**3/(a + b*tanh(x)), x)

Giac [A]

time = 0.42, size = 125, normalized size = 1.52

$$\frac{(a^2 - 2b^2) \log(e^x + 1)}{2a^3} - \frac{(a^2 - 2b^2) \log(|e^x - 1|)}{2a^3} + \frac{2(a^2b - b^3) \arctan\left(\frac{ae^x + be^x}{\sqrt{a^2 - b^2}}\right)}{\sqrt{a^2 - b^2} a^3} - \frac{ae^{(3x)} - 2be^{(3x)} + ae^x + 2be^x}{a^2(e^{(2x)} - 1)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csch(x)^3/(a+b*tanh(x)),x, algorithm="giac")

[Out] $\frac{1}{2}(a^2 - 2b^2)\log(e^x + 1)/a^3 - \frac{1}{2}(a^2 - 2b^2)\log(\text{abs}(e^x - 1))/a^3 + 2(a^2b - b^3)\arctan((ae^x + be^x)/\sqrt{a^2 - b^2})/(\sqrt{a^2 - b^2})a^3 - (ae^{3x} - 2b^3e^{3x} + ae^x + 2b^2e^x)/(a^2(e^{2x} - 1)^2)$

Mupad [B]

time = 1.51, size = 506, normalized size = 6.17

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{int}(1/(\sinh(x)^3(a + b\tanh(x))), x)$

[Out] $\log(8a^3b - 16ab^3 - 4a^4 + 8b^4 + 4a^2b^2 - 4a^4\exp(x) + 8b^4\exp(x) - 16ab^3\exp(x) + 8a^3b\exp(x) + 4a^2b^2\exp(x))/(2a) - (2\exp(x))/(a - 2a\exp(2x) + a\exp(4x)) - \log(16ab^3 - 8a^3b + 4a^4 - 8b^4 - 4a^2b^2 - 4a^4\exp(x) + 8b^4\exp(x) - 16ab^3\exp(x) + 8a^3b\exp(x) + 4a^2b^2\exp(x))/(2a) - (b^2\log(8a^3b - 16ab^3 - 4a^4 + 8b^4 + 4a^2b^2 - 4a^4\exp(x) + 8b^4\exp(x) - 16ab^3\exp(x) + 8a^3b\exp(x) + 4a^2b^2\exp(x)))/a^3 + (b^2\log(16ab^3 - 8a^3b + 4a^4 - 8b^4 - 4a^2b^2 - 4a^4\exp(x) + 8b^4\exp(x) - 16ab^3\exp(x) + 8a^3b\exp(x) + 4a^2b^2\exp(x)))/a^3 - (a\exp(x))/(a^2\exp(2x) - a^2) + (2b\exp(x))/(a^2\exp(2x) - a^2) - (b\log(8b^2(b^2 - a^2)^{1/2} - 8b^3\exp(x) + 8a^2b\exp(x) - 8ab(b^2 - a^2)^{1/2}))(b^2 - a^2)^{1/2}/a^3 + (b\log(8b^2(b^2 - a^2)^{1/2} + 8b^3\exp(x) - 8a^2b\exp(x) - 8ab(b^2 - a^2)^{1/2}))(b^2 - a^2)^{1/2}/a^3$

3.87 $\int \frac{\operatorname{csch}^4(x)}{a+b \tanh(x)} dx$

Optimal. Leaf size=78

$$\frac{(a^2 - b^2) \operatorname{coth}(x)}{a^3} + \frac{b \operatorname{coth}^2(x)}{2a^2} - \frac{\operatorname{coth}^3(x)}{3a} + \frac{b(a^2 - b^2) \log(\tanh(x))}{a^4} - \frac{b(a^2 - b^2) \log(a + b \tanh(x))}{a^4}$$

[Out] $(a^2 - b^2) \operatorname{coth}(x) / a^3 + 1/2 * b * \operatorname{coth}(x)^2 / a^2 - 1/3 * \operatorname{coth}(x)^3 / a + b * (a^2 - b^2) * \ln(\tanh(x)) / a^4 - b * (a^2 - b^2) * \ln(a + b * \tanh(x)) / a^4$

Rubi [A]

time = 0.07, antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {3597, 908}

$$\frac{b \operatorname{coth}^2(x)}{2a^2} + \frac{b(a^2 - b^2) \log(\tanh(x))}{a^4} - \frac{b(a^2 - b^2) \log(a + b \tanh(x))}{a^4} + \frac{(a^2 - b^2) \operatorname{coth}(x)}{a^3} - \frac{\operatorname{coth}^3(x)}{3a}$$

Antiderivative was successfully verified.

[In] `Int[Csch[x]^4/(a + b*Tanh[x]), x]`

[Out] $((a^2 - b^2) \operatorname{Coth}[x]) / a^3 + (b \operatorname{Coth}[x]^2) / (2 * a^2) - \operatorname{Coth}[x]^3 / (3 * a) + (b * (a^2 - b^2) * \operatorname{Log}[\operatorname{Tanh}[x]]) / a^4 - (b * (a^2 - b^2) * \operatorname{Log}[a + b * \operatorname{Tanh}[x]]) / a^4$

Rule 908

`Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))^(n_)*((a_.) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + c*x^2)^p, x], x] /; FreeQ[{a, c, d, e, f, g}, x] && NeQ[e*f - d*g, 0] && NeQ[c*d^2 + a*e^2, 0] && IntegerQ[p] && ((EqQ[p, 1] && IntegerQ[m, n]) || (ILtQ[m, 0] && ILtQ[n, 0]))`

Rule 3597

`Int[sin[(e_.) + (f_.)*(x_)]^(m_)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]^(n_)), x_Symbol] := Dist[b/f, Subst[Int[x^m*((a + x)^n/(b^2 + x^2)^(m/2 + 1)), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, n}, x] && IntegerQ[m/2]`

Rubi steps

$$\begin{aligned} \int \frac{\operatorname{csch}^4(x)}{a+b \tanh(x)} dx &= - \left(b \operatorname{Subst} \left(\int \frac{-b^2 + x^2}{x^4(a+x)} dx, x, b \tanh(x) \right) \right) \\ &= - \left(b \operatorname{Subst} \left(\int \left(-\frac{b^2}{ax^4} + \frac{b^2}{a^2x^3} + \frac{a^2 - b^2}{a^3x^2} + \frac{-a^2 + b^2}{a^4x} + \frac{a^2 - b^2}{a^4(a+x)} \right) dx, x, b \tanh(x) \right) \right) \\ &= \frac{(a^2 - b^2) \operatorname{coth}(x)}{a^3} + \frac{b \operatorname{coth}^2(x)}{2a^2} - \frac{\operatorname{coth}^3(x)}{3a} + \frac{b(a^2 - b^2) \log(\tanh(x))}{a^4} - \frac{b(a^2 - b^2) \log(a + b \tanh(x))}{a^4} \end{aligned}$$

Mathematica [A]

time = 0.21, size = 70, normalized size = 0.90

$$\frac{3a^2b\operatorname{csch}^2(x) - 2\coth(x)(-2a^3 + 3ab^2 + a^3\operatorname{csch}^2(x)) + 6b(a^2 - b^2)(\log(\sinh(x)) - \log(a\cosh(x) + b\sinh(x)))}{6a^4}$$

Antiderivative was successfully verified.

[In] Integrate[Csch[x]^4/(a + b*Tanh[x]),x]

[Out] (3*a^2*b*Csch[x]^2 - 2*Coth[x]*(-2*a^3 + 3*a*b^2 + a^3*Csch[x]^2) + 6*b*(a^2 - b^2)*(Log[Sinh[x]] - Log[a*Cosh[x] + b*Sinh[x]]))/(6*a^4)

Maple [A]

time = 0.75, size = 146, normalized size = 1.87

method	result
risch	$-\frac{2(-3abe^{4x} + 3b^2e^{4x} + 6a^2e^{2x} + 3abe^{2x} - 6b^2e^{2x} - 2a^2 + 3b^2)}{3a^3(e^{2x} - 1)^3} - \frac{b \ln\left(e^{2x} + \frac{a-b}{a+b}\right)}{a^2} + \frac{b^3 \ln\left(e^{2x} + \frac{a-b}{a+b}\right)}{a^4} + \frac{b \ln(e^{2x} - 1)}{a^2} - \frac{b^3 \ln(e^{2x} - 1)}{a^4}$
default	$-\frac{\left(\tanh^3\left(\frac{x}{2}\right)\right)a^2}{3} - ab\left(\tanh^2\left(\frac{x}{2}\right)\right) - 3a^2 \tanh\left(\frac{x}{2}\right) + 4b^2 \tanh\left(\frac{x}{2}\right) - \frac{2b\left(\frac{a^2}{2} - \frac{b^2}{2}\right) \ln\left(a\left(\tanh^2\left(\frac{x}{2}\right)\right) + 2b \tanh\left(\frac{x}{2}\right) + a\right)}{a^4} - \frac{1}{24a \tanh\left(\frac{x}{2}\right)^3}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csch(x)^4/(a+b*tanh(x)),x,method=_RETURNVERBOSE)

[Out] -1/8/a^3*(1/3*tanh(1/2*x)^3*a^2-a*b*tanh(1/2*x)^2-3*a^2*tanh(1/2*x)+4*b^2*tanh(1/2*x))-2*b/a^4*(1/2*a^2-1/2*b^2)*ln(a*tanh(1/2*x)^2+2*b*tanh(1/2*x)+a)-1/24/a/tanh(1/2*x)^3-1/8/a^3*(-3*a^2+4*b^2)/tanh(1/2*x)+1/8*b/a^2/tanh(1/2*x)^2+1/a^4*b*(a^2-b^2)*ln(tanh(1/2*x))

Maxima [B] Leaf count of result is larger than twice the leaf count of optimal. 161 vs. 2(74) = 148.

time = 0.29, size = 161, normalized size = 2.06

$$-\frac{2(2a^2 - 3b^2 - 3(2a^2 - ab - 2b^2)e^{-2x}) - 3(ab + b^2)e^{-4x}}{3(3a^3e^{-2x} - 3a^3e^{-4x} + a^3e^{-6x} - a^3)} - \frac{(a^2b - b^3) \log(-(a-b)e^{-2x} - a - b)}{a^4} + \frac{(a^2b - b^3) \log(e^{-x} + 1)}{a^4} + \frac{(a^2b - b^3) \log(e^{-x} - 1)}{a^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csch(x)^4/(a+b*tanh(x)),x, algorithm="maxima")

[Out] -2/3*(2*a^2 - 3*b^2 - 3*(2*a^2 - a*b - 2*b^2)*e^(-2*x) - 3*(a*b + b^2)*e^(-4*x))/(3*a^3*e^(-2*x) - 3*a^3*e^(-4*x) + a^3*e^(-6*x) - a^3) - (a^2*b - b^3)*log(-(a - b)*e^(-2*x) - a - b)/a^4 + (a^2*b - b^3)*log(e^(-x) + 1)/a^4 + (a^2*b - b^3)*log(e^(-x) - 1)/a^4

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 912 vs. 2(74) = 148.

time = 0.36, size = 912, normalized size = 11.69

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csch(x)^4/(a+b*tanh(x)),x, algorithm="fricas")

[Out] $\frac{1}{3}*(6*(a^2*b - a*b^2)*\cosh(x)^4 + 24*(a^2*b - a*b^2)*\cosh(x)*\sinh(x)^3 + 6*(a^2*b - a*b^2)*\sinh(x)^4 + 4*a^3 - 6*a*b^2 - 6*(2*a^3 + a^2*b - 2*a*b^2)*\cosh(x)^2 - 6*(2*a^3 + a^2*b - 2*a*b^2 - 6*(a^2*b - a*b^2)*\cosh(x)^2)*\sinh(x)^2 - 3*((a^2*b - b^3)*\cosh(x)^6 + 6*(a^2*b - b^3)*\cosh(x)*\sinh(x)^5 + (a^2*b - b^3)*\sinh(x)^6 - 3*(a^2*b - b^3)*\cosh(x)^4 - 3*(a^2*b - b^3 - 5*(a^2*b - b^3)*\cosh(x)^2)*\sinh(x)^4 + 4*(5*(a^2*b - b^3)*\cosh(x)^3 - 3*(a^2*b - b^3)*\cosh(x))*\sinh(x)^3 - a^2*b + b^3 + 3*(a^2*b - b^3)*\cosh(x)^2 + 3*(5*(a^2*b - b^3)*\cosh(x)^4 + a^2*b - b^3 - 6*(a^2*b - b^3)*\cosh(x)^2)*\sinh(x)^2 + 6*((a^2*b - b^3)*\cosh(x)^5 - 2*(a^2*b - b^3)*\cosh(x)^3 + (a^2*b - b^3)*\cosh(x))*\sinh(x)*\log(2*(a*\cosh(x) + b*\sinh(x))/(\cosh(x) - \sinh(x))) + 3*((a^2*b - b^3)*\cosh(x)^6 + 6*(a^2*b - b^3)*\cosh(x)*\sinh(x)^5 + (a^2*b - b^3)*\sinh(x)^6 - 3*(a^2*b - b^3)*\cosh(x)^4 - 3*(a^2*b - b^3 - 5*(a^2*b - b^3)*\cosh(x)^2)*\sinh(x)^4 + 4*(5*(a^2*b - b^3)*\cosh(x)^3 - 3*(a^2*b - b^3)*\cosh(x))*\sinh(x)^3 - a^2*b + b^3 + 3*(a^2*b - b^3)*\cosh(x)^2 + 3*(5*(a^2*b - b^3)*\cosh(x)^4 + a^2*b - b^3 - 6*(a^2*b - b^3)*\cosh(x)^2)*\sinh(x)^2 + 6*((a^2*b - b^3)*\cosh(x)^5 - 2*(a^2*b - b^3)*\cosh(x)^3 + (a^2*b - b^3)*\cosh(x))*\sinh(x)*\log(2*\sinh(x)/(\cosh(x) - \sinh(x))) + 12*(2*(a^2*b - a*b^2)*\cosh(x)^3 - (2*a^3 + a^2*b - 2*a*b^2)*\cosh(x))*\sinh(x))/(a^4*\cosh(x)^6 + 6*a^4*\cosh(x)*\sinh(x)^5 + a^4*\sinh(x)^6 - 3*a^4*\cosh(x)^4 + 3*a^4*\cosh(x)^2 + 3*(5*a^4*\cosh(x)^2 - a^4)*\sinh(x)^4 - a^4 + 4*(5*a^4*\cosh(x)^3 - 3*a^4*\cosh(x))*\sinh(x)^3 + 3*(5*a^4*\cosh(x)^4 - 6*a^4*\cosh(x)^2 + a^4)*\sinh(x)^2 + 6*(a^4*\cosh(x)^5 - 2*a^4*\cosh(x)^3 + a^4*\cosh(x))*\sinh(x))$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{csch}^4(x)}{a + b \tanh(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csch(x)**4/(a+b*tanh(x)),x)

[Out] Integral(csch(x)**4/(a + b*tanh(x)), x)

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 202 vs. 2(74) = 148.

time = 0.42, size = 202, normalized size = 2.59

$$\frac{(a^3b + a^2b^2 - ab^3 - b^4) \log(|ae^{2x} + be^{2x} + a - b|)}{a^5 + a^4b} + \frac{(a^2b - b^3) \log(|e^{2x} - 1|)}{a^4} - \frac{11a^2be^{6x} - 11b^3e^{6x} - 45a^2be^{4x} + 12ab^2e^{4x} + 33b^3e^{4x} + 24a^3e^{2x} + 45a^2be^{2x} - 24ab^2e^{2x} - 33b^3e^{2x} - 8a^3 - 11a^2b + 12ab^2 + 11b^3}{6a^4(e^{2x} - 1)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csch(x)^4/(a+b*tanh(x)),x, algorithm="giac")

[Out] $-(a^3b + a^2b^2 - ab^3 - b^4) \log(\text{abs}(ae^{2x} + be^{2x} + a - b)) / (a^5 + a^4b) + (a^2b - b^3) \log(\text{abs}(e^{2x} - 1)) / a^4 - 1/6(11a^2be^{6x} - 11b^3e^{6x} - 45a^2be^{4x} + 12ab^2e^{4x} + 33b^3e^{4x} + 24a^3e^{2x} + 45a^2be^{2x} - 24ab^2e^{2x} - 33b^3e^{2x} - 8a^3 - 11a^2b + 12ab^2 + 11b^3) / (a^4(e^{2x} - 1)^3)$

Mupad [B]

time = 1.33, size = 123, normalized size = 1.58

$$\frac{2b(a-b)}{a^3(e^{2x}-1)} - \frac{2(2a-b)}{a^2(e^{4x}-2e^{2x}+1)} - \frac{8}{3a(3e^{2x}-3e^{4x}+e^{6x}-1)} - \frac{b \ln(a-b+ae^{2x}+be^{2x})(a+b)(a-b)}{a^4} + \frac{b \ln(e^{2x}-1)(a+b)(a-b)}{a^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(sinh(x)^4*(a + b*tanh(x))),x)`

[Out] $(2b(a-b))/(a^3(\exp(2x)-1)) - (2(2a-b))/(a^2(\exp(4x)-2\exp(2x)+1)) - 8/(3a(3\exp(2x)-3\exp(4x)+\exp(6x)-1)) - (b \log(a-b + a\exp(2x) + b\exp(2x))*(a+b)*(a-b))/a^4 + (b \log(\exp(2x)-1)*(a+b)*(a-b))/a^4$

3.88 $\int \frac{\operatorname{csch}^5(x)}{a+b \tanh(x)} dx$

Optimal. Leaf size=255

$$-\frac{b \operatorname{ArcTan}(\sinh(x))}{a^2} + \frac{b^3 \operatorname{ArcTan}(\sinh(x))}{a^4} + \frac{b(a^2 - b^2) \operatorname{ArcTan}(\sinh(x))}{a^4} - \frac{b(a^2 - b^2)^{3/2} \operatorname{ArcTan}\left(\frac{b \cosh(x) + a \sinh(x)}{\sqrt{a^2 - b^2}}\right)}{a^5}$$

[Out] $-b \cdot \arctan(\sinh(x)) / a^2 + b^3 \cdot \arctan(\sinh(x)) / a^4 + b \cdot (a^2 - b^2) \cdot \arctan(\sinh(x)) / a^4 - b \cdot (a^2 - b^2)^{3/2} \cdot \arctan((b \cdot \cosh(x) + a \cdot \sinh(x)) / (a^2 - b^2)^{1/2}) / a^5 - 3/8 \cdot \operatorname{arctanh}(\cosh(x)) / a + 3/2 \cdot b^2 \cdot \operatorname{arctanh}(\cosh(x)) / a^3 - b^4 \cdot \operatorname{arctanh}(\cosh(x)) / a^5 - b \cdot \operatorname{csch}(x) / a^2 + 3/2 \cdot b^3 \cdot \operatorname{csch}(x) / a^4 + 3/8 \cdot \operatorname{coth}(x) \cdot \operatorname{csch}(x) / a + 1/3 \cdot b \cdot \operatorname{csch}(x)^3 / a^2 - 1/4 \cdot \operatorname{coth}(x) \cdot \operatorname{csch}(x)^3 / a - 3/2 \cdot b^2 \cdot \operatorname{sech}(x) / a^3 + b^4 \cdot \operatorname{sech}(x) / a^5 + b^2 \cdot (a^2 - b^2) \cdot \operatorname{sech}(x) / a^5 - 1/2 \cdot b^2 \cdot \operatorname{csch}(x)^2 \cdot \operatorname{sech}(x) / a^3 - 1/2 \cdot b^3 \cdot \operatorname{csch}(x) \cdot \operatorname{sech}(x)^2 / a^4 - 1/2 \cdot b^3 \cdot \operatorname{sech}(x) \cdot \tanh(x) / a^4$

Rubi [A]

time = 0.40, antiderivative size = 255, normalized size of antiderivative = 1.00, number of steps used = 29, number of rules used = 13, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {3599, 3189, 3853, 3855, 2701, 308, 213, 2702, 294, 327, 3183, 3153, 212}

$$\frac{b \operatorname{csch}(x)}{a^2} - \frac{b^3 \operatorname{csch}(x)}{a^4} + \frac{b \operatorname{ArcTan}(\sinh(x))}{a^2} - \frac{3b \operatorname{csch}(x)}{2a^2} + \frac{b^3 \operatorname{csch}(x) \operatorname{sech}(x)}{2a^2} - \frac{b^5 \operatorname{csch}(x)}{2a^2} + \frac{3b^3 \operatorname{csch}(x)}{2a^2} - \frac{b^5 \operatorname{csch}(x) \operatorname{sech}(x)}{2a^2} + \frac{b \operatorname{ArcTan}(\sinh(x))}{a^2} + \frac{b \operatorname{csch}(x)}{3a^2} - \frac{b \operatorname{csch}(x)}{a^2} - \frac{b(a^2 - b^2)^{3/2} \operatorname{ArcTan}\left(\frac{b \cosh(x) + a \sinh(x)}{\sqrt{a^2 - b^2}}\right)}{a^2} + \frac{b^3(a^2 - b^2) \operatorname{sech}(x)}{a^2} + \frac{b(a^2 - b^2) \operatorname{ArcTan}(\sinh(x))}{a^2} - \frac{3 \operatorname{tanh}^{-1}(\cosh(x))}{8a} + \frac{\operatorname{coth}(x) \operatorname{csch}(x)}{4a} - \frac{3 \operatorname{coth}(x) \operatorname{csch}(x)}{8a}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[\operatorname{Csch}[x]^5 / (a + b \cdot \operatorname{Tanh}[x]), x]$

[Out] $-((b \cdot \operatorname{ArcTan}[\operatorname{Sinh}[x]]) / a^2) + (b^3 \cdot \operatorname{ArcTan}[\operatorname{Sinh}[x]]) / a^4 + (b \cdot (a^2 - b^2) \cdot \operatorname{ArcTan}[\operatorname{Sinh}[x]]) / a^4 - (b \cdot (a^2 - b^2)^{3/2} \cdot \operatorname{ArcTan}[(b \cdot \operatorname{Cosh}[x] + a \cdot \operatorname{Sinh}[x]) / \operatorname{Sqrt}[a^2 - b^2]]) / a^5 - (3 \cdot \operatorname{ArcTanh}[\operatorname{Cosh}[x]]) / (8 \cdot a) + (3 \cdot b^2 \cdot \operatorname{ArcTanh}[\operatorname{Cosh}[x]]) / (2 \cdot a^3) - (b^4 \cdot \operatorname{ArcTanh}[\operatorname{Cosh}[x]]) / a^5 - (b \cdot \operatorname{Csch}[x]) / a^2 + (3 \cdot b^3 \cdot \operatorname{Csch}[x]) / (2 \cdot a^4) + (3 \cdot \operatorname{Coth}[x] \cdot \operatorname{Csch}[x]) / (8 \cdot a) + (b \cdot \operatorname{Csch}[x]^3) / (3 \cdot a^2) - (\operatorname{Coth}[x] \cdot \operatorname{Csch}[x]^3) / (4 \cdot a) - (3 \cdot b^2 \cdot \operatorname{Sech}[x]) / (2 \cdot a^3) + (b^4 \cdot \operatorname{Sech}[x]) / a^5 + (b^2 \cdot (a^2 - b^2) \cdot \operatorname{Sech}[x]) / a^5 - (b^2 \cdot \operatorname{Csch}[x]^2 \cdot \operatorname{Sech}[x]) / (2 \cdot a^3) - (b^3 \cdot \operatorname{Csch}[x] \cdot \operatorname{Sech}[x]^2) / (2 \cdot a^4) - (b^3 \cdot \operatorname{Sech}[x] \cdot \operatorname{Tanh}[x]) / (2 \cdot a^4)$

Rule 212

$\operatorname{Int}[(a + (b \cdot x)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1 / (\operatorname{Rt}[a, 2] \cdot \operatorname{Rt}[-b, 2])) \cdot \operatorname{ArcTanh}[\operatorname{Rt}[-b, 2] \cdot (x / \operatorname{Rt}[a, 2])], x] / ; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$

Rule 213

$\operatorname{Int}[(a + (b \cdot x)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(-\operatorname{Rt}[-a, 2] \cdot \operatorname{Rt}[b, 2])^{-1} \cdot \operatorname{ArcTanh}[\operatorname{Rt}[b, 2] \cdot (x / \operatorname{Rt}[-a, 2])], x] / ; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{NegQ}[a/b] \ \&\&$

(LtQ[a, 0] || GtQ[b, 0])

Rule 294

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n * ((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && ! LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 308

Int[(x_)^(m_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := Int[PolynomialDivide[x^m, a + b*x^n, x], x] /; FreeQ[{a, b}, x] && IGtQ[m, 0] && IGtQ[n, 0] && GtQ[m, 2*n - 1]

Rule 327

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 2701

Int[(csc[(e_.) + (f_.)*(x_)])*(a_.)^(m_.)*sec[(e_.) + (f_.)*(x_)^(n_.)], x_Symbol] := Dist[-(f*a^n)^(-1), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^(n + 1)/2], x], x, a*Csc[e + f*x], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rule 2702

Int[csc[(e_.) + (f_.)*(x_)^(n_.)]*(a_.)*sec[(e_.) + (f_.)*(x_)]^(m_), x_Symbol] := Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^(n + 1)/2], x], x, a*Sec[e + f*x], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rule 3153

Int[(cos[(c_.) + (d_.)*(x_)])*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-1), x_Symbol] := Dist[-d^(-1), Subst[Int[1/(a^2 + b^2 - x^2), x], x, b*Cos[c + d*x] - a*Sin[c + d*x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0]

Rule 3183

```
Int[cos[(c_.) + (d_.)*(x_)]^(m_)/(cos[(c_.) + (d_.)*(x_)]*(a_.) + (b_.)*sin
[(c_.) + (d_.)*(x_)]), x_Symbol] := Simp[-Cos[c + d*x]^(m + 1)/(b*d*(m + 1)
), x] + (-Dist[a/b^2, Int[Cos[c + d*x]^(m + 1), x], x] + Dist[(a^2 + b^2)/b
^2, Int[Cos[c + d*x]^(m + 2)/(a*Cos[c + d*x] + b*Sin[c + d*x]), x], x]) /;
FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]
```

Rule 3189

```
Int[(cos[(c_.) + (d_.)*(x_)]^(m_.)*sin[(c_.) + (d_.)*(x_)]^(n_.))/(cos[(c_.
) + (d_.)*(x_)]*(a_.) + (b_.)*sin[(c_.) + (d_.)*(x_)]), x_Symbol] := Int[Ex
pandTrig[cos[c + d*x]^m*(sin[c + d*x]^n/(a*cos[c + d*x] + b*sin[c + d*x])),
x], x] /; FreeQ[{a, b, c, d, m, n}, x] && IntegersQ[m, n]
```

Rule 3599

```
Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]^(n
_.), x_Symbol] := Int[Sin[e + f*x]^m*((a*cos[e + f*x] + b*sin[e + f*x])^n/C
os[e + f*x]^n), x] /; FreeQ[{a, b, e, f}, x] && IntegerQ[(m - 1)/2] && ILtQ
[n, 0] && ((LtQ[m, 5] && GtQ[n, -4]) || (EqQ[m, 5] && EqQ[n, -1]))
```

Rule 3853

```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*
x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Dist[b^2*((n - 2)/(n - 1)),
Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] &
& IntegerQ[2*n]
```

Rule 3855

```
Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned}
\int \frac{\operatorname{csch}^5(x)}{a + b \tanh(x)} dx &= \int \frac{\operatorname{coth}(x) \operatorname{csch}^4(x)}{a \cosh(x) + b \sinh(x)} dx \\
&= i \int \left(-\frac{i \operatorname{csch}^5(x)}{a} + \frac{i b \operatorname{csch}^4(x) \operatorname{sech}(x)}{a^2} - \frac{i b^2 \operatorname{csch}^3(x) \operatorname{sech}^2(x)}{a^3} + \frac{i b^3 \operatorname{csch}^2(x) \operatorname{sech}^3(x)}{a^4} \right) dx \\
&= \frac{\int \operatorname{csch}^5(x) dx}{a} - \frac{b \int \operatorname{csch}^4(x) \operatorname{sech}(x) dx}{a^2} + \frac{b^2 \int \operatorname{csch}^3(x) \operatorname{sech}^2(x) dx}{a^3} - \frac{b^3 \int \operatorname{csch}^2(x) \operatorname{sech}^3(x) dx}{a^4} \\
&= -\frac{\operatorname{coth}(x) \operatorname{csch}^3(x)}{4a} - \frac{b^4 \operatorname{sech}^3(x)}{3a^5} - \frac{3 \int \operatorname{csch}^3(x) dx}{4a} - \frac{(ib) \operatorname{Subst}\left(\int \frac{x^4}{-1+x^2} dx, x, -i \operatorname{csch}(x)\right)}{a^2} \\
&= \frac{3 \operatorname{coth}(x) \operatorname{csch}(x)}{8a} - \frac{\operatorname{coth}(x) \operatorname{csch}^3(x)}{4a} + \frac{b^2(a^2 - b^2) \operatorname{sech}(x)}{a^5} - \frac{b^2 \operatorname{csch}^2(x) \operatorname{sech}(x)}{2a^3} - \frac{b^3 \operatorname{csch}(x)}{a^2} + \frac{b^3 \tan^{-1}(\sinh(x))}{2a^4} \\
&= -\frac{b^3 \tan^{-1}(\sinh(x))}{2a^4} + \frac{b(a^2 - b^2) \tan^{-1}(\sinh(x))}{a^4} - \frac{3 \tanh^{-1}(\cosh(x))}{8a} - \frac{b \operatorname{csch}(x)}{a^2} + \frac{b \tan^{-1}(\sinh(x))}{a^2} \\
&= -\frac{b \tan^{-1}(\sinh(x))}{a^2} + \frac{b^3 \tan^{-1}(\sinh(x))}{a^4} + \frac{b(a^2 - b^2) \tan^{-1}(\sinh(x))}{a^4} - \frac{b(a^2 - b^2)^{3/2}}{a^4}
\end{aligned}$$

Mathematica [A]

time = 0.61, size = 296, normalized size = 1.16

$$\frac{-384\sqrt{a-b}\sqrt{a+b}\operatorname{ArcTan}\left(\frac{b+\operatorname{Tanh}\left[\frac{x}{2}\right]}{\sqrt{a-b}\sqrt{a+b}}\right) + 384\sqrt{a-b}\sqrt{a+b}\operatorname{ArcTan}\left(\frac{b-\operatorname{Tanh}\left[\frac{x}{2}\right]}{\sqrt{a-b}\sqrt{a+b}}\right) - 16b(7a^2 - 6b^2)\operatorname{coth}\left(\frac{x}{2}\right) + 6a^2(3a^2 - 4b^2)\operatorname{csch}^2\left(\frac{x}{2}\right) + 72a^4\log\left[\operatorname{Tanh}\left[\frac{x}{2}\right]\right] - 288a^2b\log\left[\operatorname{Tanh}\left[\frac{x}{2}\right]\right] + 192b^3\log\left[\operatorname{Tanh}\left[\frac{x}{2}\right]\right] + 18a^4\operatorname{sech}^2\left(\frac{x}{2}\right) - 24a^2b^2\operatorname{sech}^2\left(\frac{x}{2}\right) + 3a^4\operatorname{sech}^4\left(\frac{x}{2}\right) + 64a^3b\operatorname{csch}[x]^3\operatorname{Sinh}\left[\frac{x}{2}\right]^4 + a^3\operatorname{csch}[x/2]^4(-3a + 4b\operatorname{Sinh}[x]) + 112a^3b\operatorname{Tanh}\left[\frac{x}{2}\right] - 96a^2b^3\operatorname{Tanh}\left[\frac{x}{2}\right]}{16a^4}$$

Antiderivative was successfully verified.

[In] Integrate[Csch[x]^5/(a + b*Tanh[x]), x]

[Out] $(-384*a^2*\sqrt{a-b}*b*\sqrt{a+b}*\operatorname{ArcTan}[(b+a*\operatorname{Tanh}[x/2])]/(\sqrt{a-b}*\sqrt{a+b})) + 384*\sqrt{a-b}*b^3*\sqrt{a+b}*\operatorname{ArcTan}[(b+a*\operatorname{Tanh}[x/2])]/(\sqrt{a-b}*\sqrt{a+b}) - 16*a*b*(7*a^2 - 6*b^2)*\operatorname{Coth}[x/2] + 6*a^2*(3*a^2 - 4*b^2)*\operatorname{Csch}[x/2]^2 + 72*a^4*\log[\operatorname{Tanh}[x/2]] - 288*a^2*b^2*\log[\operatorname{Tanh}[x/2]] + 192*b^3*\log[\operatorname{Tanh}[x/2]] + 18*a^4*\operatorname{Sech}[x/2]^2 - 24*a^2*b^2*\operatorname{Sech}[x/2]^2 + 3*a^4*\operatorname{Sech}[x/2]^4 + 64*a^3*b*\operatorname{Csch}[x]^3*\operatorname{Sinh}[x/2]^4 + a^3*\operatorname{Csch}[x/2]^4*(-3*a + 4*b*\operatorname{Sinh}[x]) + 112*a^3*b*\operatorname{Tanh}[x/2] - 96*a*b^3*\operatorname{Tanh}[x/2])/ (192*a^5)$

Maple [A]

time = 1.02, size = 227, normalized size = 0.89

method	result
default	$ \frac{a^3 \left(\tanh^4\left(\frac{x}{2}\right) \right)}{4} - \frac{2b \left(\tanh^3\left(\frac{x}{2}\right) \right) a^2}{3} - \frac{2a^3 \left(\tanh^2\left(\frac{x}{2}\right) \right) + 2a b^2 \left(\tanh^2\left(\frac{x}{2}\right) \right) + 10a^2 b \tanh\left(\frac{x}{2}\right) - 8b^3 \tanh\left(\frac{x}{2}\right)}{16a^4} - \frac{1}{64a \tanh\left(\frac{x}{2}\right)^4} - \frac{-4a^2 - b^3}{32a^3 \tanh\left(\frac{x}{2}\right)} $

risch

$$\frac{e^x(9a^3e^{6x}-24a^2be^{6x}-12ab^2e^{6x}+24b^3e^{6x}-33a^3e^{4x}+104a^2be^{4x}+12ab^2e^{4x}-72b^3e^{4x}-33a^3e^{2x}-104a^2be^{2x}+12ab^2e^{2x}+72b^3e^{2x}+9a^3)}{12a^4(e^{2x}-1)^4}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(csch(x)^5/(a+b*tanh(x)),x,method=_RETURNVERBOSE)
```

```
[Out] 1/16/a^4*(1/4*a^3*tanh(1/2*x)^4-2/3*b*tanh(1/2*x)^3*a^2-2*a^3*tanh(1/2*x)^2+2*a*b^2*tanh(1/2*x)^2+10*a^2*b*tanh(1/2*x)-8*b^3*tanh(1/2*x))-1/64/a/tanh(1/2*x)^4-1/32*(-4*a^2+4*b^2)/a^3/tanh(1/2*x)^2+1/16/a^5*(6*a^4-24*a^2*b^2+16*b^4)*ln(tanh(1/2*x))+1/24*b/a^2/tanh(1/2*x)^3-1/8*b*(5*a^2-4*b^2)/a^4/tanh(1/2*x)-2*b*(a^4-2*a^2*b^2+b^4)/a^5/(a^2-b^2)^(1/2)*arctan(1/2*(2*a*tanh(1/2*x)+2*b)/(a^2-b^2)^(1/2))
```

Maxima [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csch(x)^5/(a+b*tanh(x)),x, algorithm="maxima")
```

```
[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see 'assume?' for more details)
```

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 2646 vs. 2(231) = 462.

time = 0.45, size = 5347, normalized size = 20.97

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csch(x)^5/(a+b*tanh(x)),x, algorithm="fricas")
```

```
[Out] [1/24*(6*(3*a^4 - 8*a^3*b - 4*a^2*b^2 + 8*a*b^3)*cosh(x)^7 + 42*(3*a^4 - 8*a^3*b - 4*a^2*b^2 + 8*a*b^3)*cosh(x)*sinh(x)^6 + 6*(3*a^4 - 8*a^3*b - 4*a^2*b^2 + 8*a*b^3)*sinh(x)^7 - 2*(33*a^4 - 104*a^3*b - 12*a^2*b^2 + 72*a*b^3)*cosh(x)^5 - 2*(33*a^4 - 104*a^3*b - 12*a^2*b^2 + 72*a*b^3 - 63*(3*a^4 - 8*a^3*b - 4*a^2*b^2 + 8*a*b^3)*cosh(x)^2)*sinh(x)^5 + 10*(21*(3*a^4 - 8*a^3*b - 4*a^2*b^2 + 8*a*b^3)*cosh(x)^3 - (33*a^4 - 104*a^3*b - 12*a^2*b^2 + 72*a*b^3)*cosh(x))*sinh(x)^4 - 2*(33*a^4 + 104*a^3*b - 12*a^2*b^2 - 72*a*b^3)*cosh(x)^3 + 2*(105*(3*a^4 - 8*a^3*b - 4*a^2*b^2 + 8*a*b^3)*cosh(x)^4 - 33*a^4 - 104*a^3*b + 12*a^2*b^2 + 72*a*b^3 - 10*(33*a^4 - 104*a^3*b - 12*a^2*b^2
```

$$\begin{aligned}
& + 72*a*b^3)*\cosh(x)^2)*\sinh(x)^3 + 2*(63*(3*a^4 - 8*a^3*b - 4*a^2*b^2 + 8*a \\
& *b^3)*\cosh(x)^5 - 10*(33*a^4 - 104*a^3*b - 12*a^2*b^2 + 72*a*b^3)*\cosh(x)^3 \\
& - 3*(33*a^4 + 104*a^3*b - 12*a^2*b^2 - 72*a*b^3)*\cosh(x))*\sinh(x)^2 - 24*(\\
& (a^2*b - b^3)*\cosh(x)^8 + 8*(a^2*b - b^3)*\cosh(x)*\sinh(x)^7 + (a^2*b - b^3) \\
& *\sinh(x)^8 - 4*(a^2*b - b^3)*\cosh(x)^6 - 4*(a^2*b - b^3 - 7*(a^2*b - b^3)*c \\
& osh(x)^2)*\sinh(x)^6 + 8*(7*(a^2*b - b^3)*\cosh(x)^3 - 3*(a^2*b - b^3)*\cosh(x) \\
&)*\sinh(x)^5 + 6*(a^2*b - b^3)*\cosh(x)^4 + 2*(35*(a^2*b - b^3)*\cosh(x)^4 + \\
& 3*a^2*b - 3*b^3 - 30*(a^2*b - b^3)*\cosh(x)^2)*\sinh(x)^4 + 8*(7*(a^2*b - b^3) \\
&)*\cosh(x)^5 - 10*(a^2*b - b^3)*\cosh(x)^3 + 3*(a^2*b - b^3)*\cosh(x))*\sinh(x) \\
& ^3 + a^2*b - b^3 - 4*(a^2*b - b^3)*\cosh(x)^2 + 4*(7*(a^2*b - b^3)*\cosh(x)^6 \\
& - 15*(a^2*b - b^3)*\cosh(x)^4 - a^2*b + b^3 + 9*(a^2*b - b^3)*\cosh(x)^2)*\si \\
& nh(x)^2 + 8*((a^2*b - b^3)*\cosh(x)^7 - 3*(a^2*b - b^3)*\cosh(x)^5 + 3*(a^2*b \\
& - b^3)*\cosh(x)^3 - (a^2*b - b^3)*\cosh(x))*\sinh(x))*\sqrt{-a^2 + b^2}*\log(((\\
& a + b)*\cosh(x)^2 + 2*(a + b)*\cosh(x)*\sinh(x) + (a + b)*\sinh(x)^2 + 2*\sqrt{- \\
& a^2 + b^2}*(\cosh(x) + \sinh(x)) - a + b)/((a + b)*\cosh(x)^2 + 2*(a + b)*\cosh \\
& (x)*\sinh(x) + (a + b)*\sinh(x)^2 + a - b)) + 6*(3*a^4 + 8*a^3*b - 4*a^2*b^2 \\
& - 8*a*b^3)*\cosh(x) - 3*((3*a^4 - 12*a^2*b^2 + 8*b^4)*\cosh(x)^8 + 8*(3*a^4 - \\
& 12*a^2*b^2 + 8*b^4)*\cosh(x))*\sinh(x)^7 + (3*a^4 - 12*a^2*b^2 + 8*b^4)*\sinh(\\
& x)^8 - 4*(3*a^4 - 12*a^2*b^2 + 8*b^4)*\cosh(x)^6 - 4*(3*a^4 - 12*a^2*b^2 + 8 \\
& *b^4 - 7*(3*a^4 - 12*a^2*b^2 + 8*b^4)*\cosh(x)^2)*\sinh(x)^6 + 8*(7*(3*a^4 - \\
& 12*a^2*b^2 + 8*b^4)*\cosh(x)^3 - 3*(3*a^4 - 12*a^2*b^2 + 8*b^4)*\cosh(x))*\sin \\
& h(x)^5 + 6*(3*a^4 - 12*a^2*b^2 + 8*b^4)*\cosh(x)^4 + 2*(35*(3*a^4 - 12*a^2*b \\
& ^2 + 8*b^4)*\cosh(x)^4 + 9*a^4 - 36*a^2*b^2 + 24*b^4 - 30*(3*a^4 - 12*a^2*b^ \\
& 2 + 8*b^4)*\cosh(x)^2)*\sinh(x)^4 + 3*a^4 - 12*a^2*b^2 + 8*b^4 + 8*(7*(3*a^4 \\
& - 12*a^2*b^2 + 8*b^4)*\cosh(x)^5 - 10*(3*a^4 - 12*a^2*b^2 + 8*b^4)*\cosh(x)^3 \\
& + 3*(3*a^4 - 12*a^2*b^2 + 8*b^4)*\cosh(x))*\sinh(x)^3 - 4*(3*a^4 - 12*a^2*b^ \\
& 2 + 8*b^4)*\cosh(x)^2 + 4*(7*(3*a^4 - 12*a^2*b^2 + 8*b^4)*\cosh(x)^6 - 15*(3* \\
& a^4 - 12*a^2*b^2 + 8*b^4)*\cosh(x)^4 - 3*a^4 + 12*a^2*b^2 - 8*b^4 + 9*(3*a^4 \\
& - 12*a^2*b^2 + 8*b^4)*\cosh(x)^2)*\sinh(x)^2 + 8*((3*a^4 - 12*a^2*b^2 + 8*b^ \\
& 4)*\cosh(x)^7 - 3*(3*a^4 - 12*a^2*b^2 + 8*b^4)*\cosh(x)^5 + 3*(3*a^4 - 12*a^2 \\
& *b^2 + 8*b^4)*\cosh(x)^3 - (3*a^4 - 12*a^2*b^2 + 8*b^4)*\cosh(x))*\sinh(x))*\lo \\
& g(\cosh(x) + \sinh(x) + 1) + 3*((3*a^4 - 12*a^2*b^2 + 8*b^4)*\cosh(x)^8 + 8*(3 \\
& *a^4 - 12*a^2*b^2 + 8*b^4)*\cosh(x))*\sinh(x)^7 + (3*a^4 - 12*a^2*b^2 + 8*b^4) \\
& *\sinh(x)^8 - 4*(3*a^4 - 12*a^2*b^2 + 8*b^4)*\cosh(x)^6 - 4*(3*a^4 - 12*a^2*b \\
& ^2 + 8*b^4 - 7*(3*a^4 - 12*a^2*b^2 + 8*b^4)*\cosh(x)^2)*\sinh(x)^6 + 8*(7*(3* \\
& a^4 - 12*a^2*b^2 + 8*b^4)*\cosh(x)^3 - 3*(3*a^4 - 12*a^2*b^2 + 8*b^4)*\cosh(x) \\
&)*\sinh(x)^5 + 6*(3*a^4 - 12*a^2*b^2 + 8*b^4)*\cosh(x)^4 + 2*(35*(3*a^4 - 12 \\
& *a^2*b^2 + 8*b^4)*\cosh(x)^4 + 9*a^4 - 36*a^2*b^2 + 24*b^4 - 30*(3*a^4 - 12* \\
& a^2*b^2 + 8*b^4)*\cosh(x)^2)*\sinh(x)^4 + 3*a^4 - 12*a^2*b^2 + 8*b^4 + 8*(7*(\\
& 3*a^4 - 12*a^2*b^2 + 8*b^4)*\cosh(x)^5 - 10*(3*a^4 - 12*a^2*b^2 + 8*b^4)*cos \\
& h(x)^3 + 3*(3*a^4 - 12*a^2*b^2 + 8*b^4)*\cosh(x))*\sinh(x)^3 - 4*(3*a^4 - 12* \\
& a^2*b^2 + 8*b^4)*\cosh(x)^2 + 4*(7*(3*a^4 - 12*a^2*b^2 + 8*b^4)*\cosh(x)^6 - \\
& 15*(3*a^4 - 12*a^2*b^2 + 8*b^4)*\cosh(x)^4 - 3*a^4 + 12*a^2*b^2 - 8*b^4 + 9* \\
& (3*a^4 - 12*a^2*b^2 + 8*b^4)*\cosh(x)^2)*\sinh(x)^2 + 8*((3*a^4 - 12*a^2*b^2 \\
& + 8*b^4)*\cosh(x)^7 - 3*(3*a^4 - 12*a^2*b^2 + 8*b^4)*\cosh(x)^5 + 3*(3*a^4 -
\end{aligned}$$

$$12a^2b^2 + 8b^4) \cosh(x)^3 - (3a^4 - 12a^2b^2 + 8b^4) \cosh(x) \sinh(x) \log(\cosh(x) + \sinh(x) - 1) + 2(21(3a^4 - 8a^3b - 4a^2b^2 + 8ab^3) \cosh(x)^6 - 5(33a^4 - 104a^3b - 12a^2b^2 + 72ab^3) \cosh(x)^4 + 9a^4 + 24a^3b - 12a^2b^2 - 24ab^3 - 3(33a^4 + 104a^3b - 12a^2b^2 - 72ab^3) \cosh(x)^2) \sinh(x) / (a^5 \cosh(x)^8 + 8a^5 \cosh(x) \sinh(x)^7 + a^5 \sinh(x)^8 - 4a^5 \cosh(x)^6 + 6a^5 \cosh(x)^4 - 4a^5 \cosh(x)^2 + 4(7a^5 \cosh(x)^2 - a^5) \sinh(x)^6 + 8(7a^5 \cosh(x)^3 - 3a^5 \cosh(x)) \sinh(x)^5 + a^5 + 2(35a^5 \cosh(x)^4 - 30a^5 \cosh(x)^2 + 3a^5) \sinh(x)^4 + 8(7a^5 \cosh(x)^5 - 10a^5 \cosh(x)^3 + 3a^5 \cosh(x)) \sinh(x)^3 + 4(7a^5 \cosh(x)^6 - 15a^5 \cosh(x)^4 + 9a^5 \cosh(x)^2 - a^5) \sinh(x)^2 + 8(a^5 \cosh(x)^7 - 3a^5 \cosh(x)^5 + 3a^5 \cosh(x)^3 - a^5 \cosh(x)) \sinh(x)), 1/24(6(3a^4 - 8a^3b - 4a^2b^2 + 8ab^3) \cosh(x)^7 + 42(3a^4 - 8a^3b - 4a^2b^2 + 8ab^3) \cosh(x) \sinh(x)^6 + 6(3 \dots$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{csch}^5(x)}{a + b \tanh(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csch(x)**5/(a+b*tanh(x)),x)

[Out] Integral(csch(x)**5/(a + b*tanh(x)), x)

Giac [A]

time = 0.42, size = 273, normalized size = 1.07

$$\frac{(3a^4 - 12a^2b^2 + 8b^4) \log(e^x + 1)}{8a^5} + \frac{(3a^4 - 12a^2b^2 + 8b^4) \log(|e^x - 1|)}{8a^5} - \frac{2(a^5 - 2a^3b^2 + b^5) \arctan\left(\frac{a + b \tanh(x)}{\sqrt{a^2 - b^2}}\right)}{\sqrt{a^2 - b^2} a^5} + \frac{9a^5 e^{7x} - 24a^3 b^2 e^{7x} - 12a^2 b^3 e^{7x} + 24b^3 e^{7x} - 33a^3 e^{5x} + 104a^2 b^2 e^{5x} + 12a^2 b^3 e^{5x} - 33a^3 e^{3x} - 104a^2 b^2 e^{3x} + 12a^2 b^3 e^{3x} + 9a^3 e^x + 24a^2 b^2 e^x - 12a^2 b^3 e^x - 24b^3 e^x}{12a^5 (e^{2x} - 1)^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csch(x)^5/(a+b*tanh(x)),x, algorithm="giac")

[Out] $-1/8(3a^4 - 12a^2b^2 + 8b^4) \log(e^x + 1)/a^5 + 1/8(3a^4 - 12a^2b^2 + 8b^4) \log(\operatorname{abs}(e^x - 1))/a^5 - 2(a^4b - 2a^2b^3 + b^5) \arctan((a + b e^x) / \sqrt{a^2 - b^2}) / (\sqrt{a^2 - b^2} a^5) + 1/12(9a^3 e^{7x} - 24a^2 b^2 e^{7x} - 12a^2 b^3 e^{7x} + 24b^3 e^{7x} - 33a^3 e^{5x} + 104a^2 b^2 e^{5x} + 12a^2 b^3 e^{5x} - 72b^3 e^{5x} - 33a^3 e^{3x} - 104a^2 b^2 e^{3x} + 12a^2 b^3 e^{3x} + 72b^3 e^{3x} + 9a^3 e^x + 24a^2 b^2 e^x - 12a^2 b^3 e^x - 24b^3 e^x) / (a^4 (e^{2x} - 1)^4)$

Mupad [B]

time = 3.31, size = 753, normalized size = 2.95

$$\frac{(3a^4 - 12a^2b^2 + 8b^4) \log(e^x + 1)}{8a^5} + \frac{(3a^4 - 12a^2b^2 + 8b^4) \log(|e^x - 1|)}{8a^5} - \frac{2(a^5 - 2a^3b^2 + b^5) \arctan\left(\frac{a + b \tanh(x)}{\sqrt{a^2 - b^2}}\right)}{\sqrt{a^2 - b^2} a^5} + \frac{9a^5 e^{7x} - 24a^3 b^2 e^{7x} - 12a^2 b^3 e^{7x} + 24b^3 e^{7x} - 33a^3 e^{5x} + 104a^2 b^2 e^{5x} + 12a^2 b^3 e^{5x} - 33a^3 e^{3x} - 104a^2 b^2 e^{3x} + 12a^2 b^3 e^{3x} + 9a^3 e^x + 24a^2 b^2 e^x - 12a^2 b^3 e^x - 24b^3 e^x}{12a^5 (e^{2x} - 1)^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{int}(1/(\sinh(x)^5*(a + b*\tanh(x))),x)$

[Out] $(\log(\exp(x) - 1)*(3*a^4 + 8*b^4 - 12*a^2*b^2))/(8*a^5) - (\log(\exp(x) + 1)*(3*a^4 + 8*b^4 - 12*a^2*b^2))/(8*a^5) - (4*\exp(x))/(a*(6*\exp(4*x) - 4*\exp(2*x) - 4*\exp(6*x) + \exp(8*x) + 1)) - (2*\exp(x)*(9*a - 4*b))/(3*a^2*(3*\exp(2*x) - 3*\exp(4*x) + \exp(6*x) - 1)) - (\exp(x)*(3*a^2 - 16*a*b + 12*b^2))/(6*a^3*(\exp(4*x) - 2*\exp(2*x) + 1)) - (\exp(x)*(4*a*b^2 + 8*a^2*b - 3*a^3 - 8*b^3))/(4*a^4*(\exp(2*x) - 1)) + (b*\log((b*\exp(x)*(a - b)^2*(288*a*b^6 - 24*a^6*b - 9*a^7 - 192*b^7 + 224*a^2*b^5 - 456*a^3*b^4 + 24*a^4*b^3 + 144*a^5*b^2)))/(2*a^12*(a + b)) - (b*(a - b)*(-(a + b)^3*(a - b)^3)^{(1/2)}*(8*a^5*b^3 - 9*a^8 - 9*a^7*b + 8*a^6*b^2 + 192*b^5*\exp(x)*(-(a^2 - b^2)^3)^{(1/2)} - 224*a^2*b^3*\exp(x)*(-(a^2 - b^2)^3)^{(1/2)} - 88*a^3*b^2*\exp(x)*(-(a^2 - b^2)^3)^{(1/2)} + 96*a*b^4*\exp(x)*(-(a^2 - b^2)^3)^{(1/2)} + 24*a^4*b*\exp(x)*(-(a^2 - b^2)^3)^{(1/2}))/((2*a^12*(a + b)^4))*(-(a + b)^3*(a - b)^3)^{(1/2)})/a^5 - (b*\log((b*\exp(x)*(a - b)^2*(288*a*b^6 - 24*a^6*b - 9*a^7 - 192*b^7 + 224*a^2*b^5 - 456*a^3*b^4 + 24*a^4*b^3 + 144*a^5*b^2)))/(2*a^12*(a + b)) - (b*(a - b)*(-(a + b)^3*(a - b)^3)^{(1/2)}*(9*a^7*b + 9*a^8 - 8*a^5*b^3 - 8*a^6*b^2 + 192*b^5*\exp(x)*(-(a^2 - b^2)^3)^{(1/2)} - 224*a^2*b^3*\exp(x)*(-(a^2 - b^2)^3)^{(1/2)} - 88*a^3*b^2*\exp(x)*(-(a^2 - b^2)^3)^{(1/2)} + 96*a*b^4*\exp(x)*(-(a^2 - b^2)^3)^{(1/2)} + 24*a^4*b*\exp(x)*(-(a^2 - b^2)^3)^{(1/2}))/((2*a^12*(a + b)^4))*(-(a + b)^3*(a - b)^3)^{(1/2)})/a^5$

$$3.89 \quad \int \frac{\operatorname{csch}^6(x)}{a+b \tanh(x)} dx$$

Optimal. Leaf size=130

$$-\frac{(a^2 - b^2)^2 \operatorname{coth}(x)}{a^5} - \frac{b(2a^2 - b^2) \operatorname{coth}^2(x)}{2a^4} + \frac{(2a^2 - b^2) \operatorname{coth}^3(x)}{3a^3} + \frac{b \operatorname{coth}^4(x)}{4a^2} - \frac{\operatorname{coth}^5(x)}{5a} - \frac{b(a^2 - b^2)^2 \log(\tanh(x))}{a^6}$$

[Out] $-(a^2-b^2)^2 \operatorname{coth}(x)/a^5 - 1/2 b(2a^2-b^2) \operatorname{coth}(x)^2/a^4 + 1/3 (2a^2-b^2) \operatorname{coth}(x)^3/a^3 + 1/4 b \operatorname{coth}(x)^4/a^2 - 1/5 \operatorname{coth}(x)^5/a - b(a^2-b^2)^2 \ln(\tanh(x))/a^6 + b(a^2-b^2)^2 \ln(a+b \tanh(x))/a^6$

Rubi [A]

time = 0.11, antiderivative size = 130, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$,

Rules used = {3597, 908}

$$\frac{b \operatorname{coth}^4(x)}{4a^2} - \frac{b(a^2 - b^2)^2 \log(\tanh(x))}{a^6} + \frac{b(a^2 - b^2)^2 \log(a + b \tanh(x))}{a^6} - \frac{(a^2 - b^2)^2 \operatorname{coth}(x)}{a^5} - \frac{b(2a^2 - b^2) \operatorname{coth}^2(x)}{2a^4} + \frac{(2a^2 - b^2) \operatorname{coth}^3(x)}{3a^3} - \frac{\operatorname{coth}^5(x)}{5a}$$

Antiderivative was successfully verified.

[In] `Int[Csch[x]^6/(a + b*Tanh[x]), x]`

[Out] $-\left(\frac{(a^2 - b^2)^2 \operatorname{Coth}[x]}{a^5}\right) - \frac{b(2a^2 - b^2) \operatorname{Coth}[x]^2}{2a^4} + \left(\frac{2a^2 - b^2}{3a^3}\right) \operatorname{Coth}[x]^3 + \frac{b \operatorname{Coth}[x]^4}{4a^2} - \frac{\operatorname{Coth}[x]^5}{5a} - \left(\frac{b(a^2 - b^2)^2 \operatorname{Log}[\operatorname{Tanh}[x]]}{a^6}\right) + \frac{b(a^2 - b^2)^2 \operatorname{Log}[a + b \operatorname{Tanh}[x]]}{a^6}$

Rule 908

`Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))^(n_)*((a_) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + c*x^2)^p, x], x] /; FreeQ[{a, c, d, e, f, g}, x] && NeQ[e*f - d*g, 0] && NeQ[c*d^2 + a*e^2, 0] && IntegerQ[p] && ((EqQ[p, 1] && IntegersQ[m, n]) || (ILtQ[m, 0] && ILtQ[n, 0]))`

Rule 3597

`Int[sin[(e_.) + (f_.)*(x_)]^(m_)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Dist[b/f, Subst[Int[x^m*((a + x)^n/(b^2 + x^2)^(m/2 + 1)), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, n}, x] && IntegerQ[m/2]`

Rubi steps

$$\int \frac{\operatorname{csch}^6(x)}{a + b \tanh(x)} dx = b \operatorname{Subst} \left(\int \frac{(-b^2 + x^2)^2}{x^6(a + x)} dx, x, b \tanh(x) \right)$$

$$= b \operatorname{Subst} \left(\int \left(\frac{b^4}{ax^6} - \frac{b^4}{a^2x^5} + \frac{-2a^2b^2 + b^4}{a^3x^4} + \frac{2a^2b^2 - b^4}{a^4x^3} + \frac{(a^2 - b^2)^2}{a^5x^2} - \frac{(a^2 - b^2)^2}{a^6x} + \dots \right) dx, x, b \tanh(x) \right)$$

$$= -\frac{(a^2 - b^2)^2 \operatorname{coth}(x)}{a^5} - \frac{b(2a^2 - b^2) \operatorname{coth}^2(x)}{2a^4} + \frac{(2a^2 - b^2) \operatorname{coth}^3(x)}{3a^3} + \frac{b \operatorname{coth}^4(x)}{4a^2} - \dots$$

Mathematica [A]

time = 0.45, size = 119, normalized size = 0.92

$$\frac{-4 \operatorname{coth}(x) (8a^5 - 25a^3b^2 + 15ab^4 + (-4a^5 + 5a^3b^2) \operatorname{csch}^2(x) + 3a^5 \operatorname{csch}^4(x)) + 15b(-2a^2(a^2 - b^2) \operatorname{csch}^2(x) + a^4 \operatorname{csch}^4(x) - 4(a^2 - b^2)^2 (\log(\sinh(x)) - \log(a \cosh(x) + b \sinh(x))))}{60a^6}$$

Antiderivative was successfully verified.

`[In] Integrate[Csch[x]^6/(a + b*Tanh[x]), x]`

```
[Out] (-4*Coth[x]*(8*a^5 - 25*a^3*b^2 + 15*a*b^4 + (-4*a^5 + 5*a^3*b^2)*Csch[x]^2
+ 3*a^5*Csch[x]^4) + 15*b*(-2*a^2*(a^2 - b^2)*Csch[x]^2 + a^4*Csch[x]^4 -
4*(a^2 - b^2)^2*(Log[Sinh[x]] - Log[a*Cosh[x] + b*Sinh[x]])))/(60*a^6)
```

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 276 vs. 2(122) = 244.

time = 0.75, size = 277, normalized size = 2.13

method	result
default	$-\frac{\frac{a^4 \left(\tanh^5\left(\frac{x}{2}\right)\right) - b \left(\tanh^4\left(\frac{x}{2}\right)\right) a^3 - 5 \left(\tanh^3\left(\frac{x}{2}\right)\right) a^4 + 4a^2 b^2 \left(\tanh^3\left(\frac{x}{2}\right)\right)}{5} + \frac{4a^2 b^2 \left(\tanh^3\left(\frac{x}{2}\right)\right)}{3} + \frac{6a^3 b \left(\tanh^2\left(\frac{x}{2}\right)\right) - 4b^3 \left(\tanh^2\left(\frac{x}{2}\right)\right) a + 10a^4 \tanh\left(\frac{x}{2}\right) - 28a^2 b^2}{32a^5}}$
risch	$-\frac{2(15a^3 b e^{8x} - 15a^2 b^2 e^{8x} - 15a b^3 e^{8x} + 15b^4 e^{8x} - 75a^3 b e^{6x} + 90a^2 b^2 e^{6x} + 45a b^3 e^{6x} - 60b^4 e^{6x} + 80a^4 e^{4x} + 75a^3 b e^{4x} - 160a^2 b^2 e^{4x} - 45a b^3 e^{4x} + 15b^4 e^{4x})}{15a^5 (e^{2x} - 1)^5}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(csch(x)^6/(a+b*tanh(x)), x, method=_RETURNVERBOSE)`

```
[Out] -1/32/a^5*(1/5*a^4*tanh(1/2*x)^5-1/2*b*tanh(1/2*x)^4*a^3-5/3*tanh(1/2*x)^3*
a^4+4/3*a^2*b^2*tanh(1/2*x)^3+6*a^3*b*tanh(1/2*x)^2-4*b^3*tanh(1/2*x)^2*a+
0*a^4*tanh(1/2*x)-28*a^2*b^2*tanh(1/2*x)+16*b^4*tanh(1/2*x))-1/160/a/tanh(1
/2*x)^5-1/96*(-5*a^2+4*b^2)/a^3/tanh(1/2*x)^3-1/32/a^5*(10*a^4-28*a^2*b^2+1
6*b^4)/tanh(1/2*x)+1/64*b/a^2/tanh(1/2*x)^4-1/16/a^4*b*(3*a^2-2*b^2)/tanh(1
/2*x)^2-1/a^6*b*(a^4-2*a^2*b^2+b^4)*ln(tanh(1/2*x))+2*b/a^6*(1/2*a^4-a^2*b^
2+1/2*b^4)*ln(a*tanh(1/2*x)^2+2*b*tanh(1/2*x)+a)
```

Maxima [B] Leaf count of result is larger than twice the leaf count of optimal. 308 vs. $2(122) = 244$.
time = 0.30, size = 308, normalized size = 2.37

$$\frac{2(8a^4 - 25a^2b^2 + 15b^4 - 5(8a^4 - 3a^2b^2 - 22a^2b^2 + 3ab^3 + 12b^4)e^{-2x}) + 5(16a^4 - 15a^2b^2 - 32a^2b^2 + 9ab^3 + 18b^4)e^{-4x} + 15(5a^3b + 6a^2b^2 - 3a^2b^2 - 3ab^3 - 4b^4)e^{-6x} - 15(a^3b + a^2b^2 - ab^3 - b^4)e^{-8x}}{15(5a^4e^{-2x} - 10a^3b^2e^{-4x} + 10a^2b^3e^{-6x} - 5a^2b^3e^{-8x} + a^5e^{-10x} - a^5)} + \frac{(a^4b - 2a^2b^3 + b^5)\log(-(a-b)e^{-2x} - a - b)}{a^6} - \frac{(a^4b - 2a^2b^3 + b^5)\log(e^{-x} + 1)}{a^6} - \frac{(a^4b - 2a^2b^3 + b^5)\log(e^{-x} - 1)}{a^6}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csch(x)^6/(a+b*tanh(x)),x, algorithm="maxima")

[Out] $\frac{2}{15}(8a^4 - 25a^2b^2 + 15b^4 - 5(8a^4 - 3a^3b - 22a^2b^2 + 3a^2b^3 + 12b^4)e^{-2x} + 5(16a^4 - 15a^3b - 32a^2b^2 + 9a^2b^3 + 18b^4)e^{-4x} + 15(5a^3b + 6a^2b^2 - 3a^2b^2 - 3ab^3 - 4b^4)e^{-6x} - 15(a^3b + a^2b^2 - ab^3 - b^4)e^{-8x}) / (5a^5e^{-2x} - 10a^5e^{-4x} + 10a^5e^{-6x} - 5a^5e^{-8x} + a^5e^{-10x} - a^5) + (a^4b - 2a^2b^3 + b^5) \log(-(a - b)e^{-2x} - a - b) / a^6 - (a^4b - 2a^2b^3 + b^5) \log(e^{-x} + 1) / a^6 - (a^4b - 2a^2b^3 + b^5) \log(e^{-x} - 1) / a^6$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 2972 vs. $2(122) = 244$.
time = 0.40, size = 2972, normalized size = 22.86

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csch(x)^6/(a+b*tanh(x)),x, algorithm="fricas")

[Out] $-1/15(30(a^4b - a^3b^2 - a^2b^3 + ab^4)\cosh(x)^8 + 240(a^4b - a^3b^2 - a^2b^3 + ab^4)\cosh(x)\sinh(x)^7 + 30(a^4b - a^3b^2 - a^2b^3 + ab^4)\sinh(x)^8 - 30(5a^4b - 6a^3b^2 - 3a^2b^3 + 4ab^4)\cosh(x)^6 - 30(5a^4b - 6a^3b^2 - 3a^2b^3 + 4ab^4 - 28(a^4b - a^3b^2 - a^2b^3 + ab^4)\cosh(x)^2)\sinh(x)^6 + 60(28(a^4b - a^3b^2 - a^2b^3 + ab^4)\cosh(x)^3 - 3(5a^4b - 6a^3b^2 - 3a^2b^3 + 4ab^4)\cosh(x))\sinh(x)^5 + 16a^5 - 50a^3b^2 + 30a^2b^3 + 10(16a^5 + 15a^4b - 32a^3b^2 - 9a^2b^3 + 18ab^4)\cosh(x)^4 + 10(16a^5 + 15a^4b - 32a^3b^2 - 9a^2b^3 + 18ab^4 + 210(a^4b - a^3b^2 - a^2b^3 + ab^4)\cosh(x)^4 - 45(5a^4b - 6a^3b^2 - 3a^2b^3 + 4ab^4)\cosh(x)^2)\sinh(x)^4 + 40(42(a^4b - a^3b^2 - a^2b^3 + ab^4)\cosh(x)^5 - 15(5a^4b - 6a^3b^2 - 3a^2b^3 + 4ab^4)\cosh(x)^3 + (16a^5 + 15a^4b - 32a^3b^2 - 9a^2b^3 + 18ab^4)\cosh(x))\sinh(x)^3 - 10(8a^5 + 3a^4b - 22a^3b^2 - 3a^2b^3 + 12ab^4)\cosh(x)^2 + 10(84(a^4b - a^3b^2 - a^2b^3 + ab^4)\cosh(x)^6 - 8a^5 - 3a^4b + 22a^3b^2 + 3a^2b^3 - 12ab^4 - 45(5a^4b - 6a^3b^2 - 3a^2b^3 + 4ab^4)\cosh(x)^4 + 6(16a^5 + 15a^4b - 32a^3b^2 - 9a^2b^3 + 18ab^4)\cosh(x)^2)\sinh(x)^2 - 15((a^4b - 2a^2b^3 + b^5)\cosh(x)^10 + 10(a^4b - 2a^2b^3 + b^5)\cosh(x)\sinh(x)^9 + (a^4b - 2a^2b^3 + b^5)\sinh(x)^10 - 5(a^4b - 2a^2b^3 + b^5)\cosh(x)^8 -$

$$\begin{aligned}
& 5*(a^4*b - 2*a^2*b^3 + b^5 - 9*(a^4*b - 2*a^2*b^3 + b^5)*\cosh(x)^2)*\sinh(x) \\
&)^8 + 40*(3*(a^4*b - 2*a^2*b^3 + b^5)*\cosh(x)^3 - (a^4*b - 2*a^2*b^3 + b^5) \\
& *\cosh(x))*\sinh(x)^7 + 10*(a^4*b - 2*a^2*b^3 + b^5)*\cosh(x)^6 + 10*(a^4*b - \\
& 2*a^2*b^3 + b^5 + 21*(a^4*b - 2*a^2*b^3 + b^5)*\cosh(x)^4 - 14*(a^4*b - 2*a^ \\
& 2*b^3 + b^5)*\cosh(x)^2)*\sinh(x)^6 + 4*(63*(a^4*b - 2*a^2*b^3 + b^5)*\cosh(x) \\
& ^5 - 70*(a^4*b - 2*a^2*b^3 + b^5)*\cosh(x)^3 + 15*(a^4*b - 2*a^2*b^3 + b^5)* \\
& \cosh(x))*\sinh(x)^5 - a^4*b + 2*a^2*b^3 - b^5 - 10*(a^4*b - 2*a^2*b^3 + b^5) \\
& *\cosh(x)^4 + 10*(21*(a^4*b - 2*a^2*b^3 + b^5)*\cosh(x)^6 - a^4*b + 2*a^2*b^3 \\
& - b^5 - 35*(a^4*b - 2*a^2*b^3 + b^5)*\cosh(x)^4 + 15*(a^4*b - 2*a^2*b^3 + b \\
& ^5)*\cosh(x)^2)*\sinh(x)^4 + 40*(3*(a^4*b - 2*a^2*b^3 + b^5)*\cosh(x)^7 - 7*(a \\
& ^4*b - 2*a^2*b^3 + b^5)*\cosh(x)^5 + 5*(a^4*b - 2*a^2*b^3 + b^5)*\cosh(x)^3 - \\
& (a^4*b - 2*a^2*b^3 + b^5)*\cosh(x))*\sinh(x)^3 + 5*(a^4*b - 2*a^2*b^3 + b^5) \\
& *\cosh(x)^2 + 5*(9*(a^4*b - 2*a^2*b^3 + b^5)*\cosh(x)^8 - 28*(a^4*b - 2*a^2*b \\
& ^3 + b^5)*\cosh(x)^6 + a^4*b - 2*a^2*b^3 + b^5 + 30*(a^4*b - 2*a^2*b^3 + b^5) \\
&)*\cosh(x)^4 - 12*(a^4*b - 2*a^2*b^3 + b^5)*\cosh(x)^2)*\sinh(x)^2 + 10*((a^4* \\
& b - 2*a^2*b^3 + b^5)*\cosh(x)^9 - 4*(a^4*b - 2*a^2*b^3 + b^5)*\cosh(x)^7 + 6* \\
& (a^4*b - 2*a^2*b^3 + b^5)*\cosh(x)^5 - 4*(a^4*b - 2*a^2*b^3 + b^5)*\cosh(x)^3 \\
& + (a^4*b - 2*a^2*b^3 + b^5)*\cosh(x))*\sinh(x))*\log(2*(a*\cosh(x) + b*\sinh(x) \\
&))/(cosh(x) - sinh(x))) + 15*((a^4*b - 2*a^2*b^3 + b^5)*\cosh(x)^10 + 10*(a^4 \\
& *b - 2*a^2*b^3 + b^5)*\cosh(x)*\sinh(x)^9 + (a^4*b - 2*a^2*b^3 + b^5)*\sinh(x) \\
& ^10 - 5*(a^4*b - 2*a^2*b^3 + b^5)*\cosh(x)^8 - 5*(a^4*b - 2*a^2*b^3 + b^5 - \\
& 9*(a^4*b - 2*a^2*b^3 + b^5)*\cosh(x)^2)*\sinh(x)^8 + 40*(3*(a^4*b - 2*a^2*b^3 \\
& + b^5)*\cosh(x)^3 - (a^4*b - 2*a^2*b^3 + b^5)*\cosh(x))*\sinh(x)^7 + 10*(a^4* \\
& b - 2*a^2*b^3 + b^5)*\cosh(x)^6 + 10*(a^4*b - 2*a^2*b^3 + b^5 + 21*(a^4*b - \\
& 2*a^2*b^3 + b^5)*\cosh(x)^4 - 14*(a^4*b - 2*a^2*b^3 + b^5)*\cosh(x)^2)*\sinh(x) \\
&)^6 + 4*(63*(a^4*b - 2*a^2*b^3 + b^5)*\cosh(x)^5 - 70*(a^4*b - 2*a^2*b^3 + b \\
& ^5)*\cosh(x)^3 + 15*(a^4*b - 2*a^2*b^3 + b^5)*\cosh(x))*\sinh(x)^5 - a^4*b + 2 \\
& *a^2*b^3 - b^5 - 10*(a^4*b - 2*a^2*b^3 + b^5)*\cosh(x)^4 + 10*(21*(a^4*b - 2 \\
& *a^2*b^3 + b^5)*\cosh(x)^6 - a^4*b + 2*a^2*b^3 - b^5 - 35*(a^4*b - 2*a^2*b^3 \\
& + b^5)*\cosh(x)^4 + 15*(a^4*b - 2*a^2*b^3 + b^5)*\cosh(x)^2)*\sinh(x)^4 + 40* \\
& (3*(a^4*b - 2*a^2*b^3 + b^5)*\cosh(x)^7 - 7*(a^4*b - 2*a^2*b^3 + b^5)*\cosh(x) \\
&)^5 + 5*(a^4*b - 2*a^2*b^3 + b^5)*\cosh(x)^3 - (a^4*b - 2*a^2*b^3 + b^5)*\cos \\
& h(x))*\sinh(x)^3 + 5*(a^4*b - 2*a^2*b^3 + b^5)*\cosh(x)^2 + 5*(9*(a^4*b - 2*a \\
& ^2*b^3 + b^5)*\cosh(x)^8 - 28*(a^4*b - 2*a^2*b^3 + b^5)*\cosh(x)^6 + a^4*b - \\
& 2*a^2*b^3 + b^5 + 30*(a^4*b - 2*a^2*b^3 + b^5)*\cosh(x)^4 - 12*(a^4*b - 2*a^ \\
& 2*b^3 + b^5)*\cosh(x)^2)*\sinh(x)^2 + 10*((a^4*b - 2*a^2*b^3 + b^5)*\cosh(x)^9 \\
& - 4*(a^4*b - 2*a^2*b^3 + b^5)*\cosh(x)^7 + 6*(a^4*b - 2*a^2*b^3 + b^5)*\cosh \\
& (x)^5 - 4*(a^4*b - 2*a^2*b^3 + b^5)*\cosh(x)^3 + (a^4*b - 2*a^2*b^3 + b^5)*\c \\
& osh(x))*\sinh(x))*\log(2*\sinh(x)/(cosh(x) - sinh(x))) + 20*(12*(a^4*b - a^3*b \\
& ^2 - a^2*b^3 + a*b^4)*\cosh(x)^7 - 9*(5*a^4*b - 6*a^3*b^2 - 3*a^2*b^3 + 4*a* \\
& b^4)*\cosh(x)^5 + 2*(16*a^5 + 15*a^4*b - 32*a^3*b^2 - 9*a^2*b^3 + 18*a*b^4)* \\
& \cosh(x)^3 - (8*a^5 + 3*a^4*b - 22*a^3*b^2 - 3*a^2*b^3 + 12*a*b^4)*\cosh(x))* \\
& \sinh(x))/(a^6*\cosh(x)^10 + 10*a^6*\cosh(x)*\sinh(x)^9 + a^6*\sinh(x)^10 - 5*a^ \\
& 6*\cosh(x)^8 + 10*a^6*\cosh(x)^6 - 10*a^6*\cosh(x)^4 + 5*(9*a^6*\cosh(x)^2 - a^ \\
& 6)*\sinh(x)^8 + 5*a^6*\cosh(x)^2 + 40*(3*a^6*\cosh(x)^3 - a^6*\cosh(x))*\sinh(x)
\end{aligned}$$

$$\begin{aligned} & / (a^2(6\exp(4x) - 4\exp(2x) - 4\exp(6x) + \exp(8x) + 1)) - 32 / (5a(5\exp(2x) - 10\exp(4x) + 10\exp(6x) - 5\exp(8x) + \exp(10x) - 1)) - (2(a + b)(a - b)(ab - b^2)) / (a^5(\exp(2x) - 1)) + (b \log(a - b + a\exp(2x) + b\exp(2x)))(a + b)^2(a - b)^2 / a^6 - (b \log(\exp(2x) - 1))(a + b)^2(a - b)^2 / a^6 \end{aligned}$$

3.90 $\int \frac{\operatorname{csch}(x)}{i + \tanh(x)} dx$

Optimal. Leaf size=33

$$i \tanh^{-1}(\cosh(x)) - \frac{i \tanh^{-1}\left(\frac{\cosh(x) + i \sinh(x)}{\sqrt{2}}\right)}{\sqrt{2}}$$

[Out] I*arctanh(cosh(x))-1/2*I*arctanh(1/2*(cosh(x)+I*sinh(x))*2^(1/2))*2^(1/2)

Rubi [A]

time = 0.07, antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.454$, Rules used = {3599, 3189, 3855, 3153, 212}

$$i \tanh^{-1}(\cosh(x)) - \frac{i \tanh^{-1}\left(\frac{\cosh(x) + i \sinh(x)}{\sqrt{2}}\right)}{\sqrt{2}}$$

Antiderivative was successfully verified.

[In] Int[Csch[x]/(I + Tanh[x]), x]

[Out] I*ArcTanh[Cosh[x]] - (I*ArcTanh[(Cosh[x] + I*Sinh[x])/Sqrt[2]])/Sqrt[2]

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 3153

Int[(cos[(c_) + (d_)*(x_)]*(a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := Dist[-d^(-1), Subst[Int[1/(a^2 + b^2 - x^2), x], x, b*Cos[c + d*x] - a*Sin[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0]

Rule 3189

Int[(cos[(c_) + (d_)*(x_)]^(m_)*sin[(c_) + (d_)*(x_)]^(n_))/(cos[(c_) + (d_)*(x_)]*(a_) + (b_)*sin[(c_) + (d_)*(x_)]), x_Symbol] := Int[ExpandTrig[cos[c + d*x]^m*(sin[c + d*x]^n/(a*cos[c + d*x] + b*sin[c + d*x])), x], x] /; FreeQ[{a, b, c, d, m, n}, x] && IntegersQ[m, n]

Rule 3599

Int[sin[(e_) + (f_)*(x_)]^(m_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Int[Sin[e + f*x]^m*((a*cos[e + f*x] + b*sin[e + f*x])^n/C

os[e + f*x]^n), x] /; FreeQ[{a, b, e, f}, x] && IntegerQ[(m - 1)/2] && ILtQ[n, 0] && ((LtQ[m, 5] && GtQ[n, -4]) || (EqQ[m, 5] && EqQ[n, -1]))

Rule 3855

Int[csc[(c_.) + (d_.)*(x_.)], x_Symbol] :> Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned}
 \int \frac{\operatorname{csch}(x)}{i + \tanh(x)} dx &= \int \frac{\operatorname{coth}(x)}{i \cosh(x) + \sinh(x)} dx \\
 &= i \int \left(-\operatorname{csch}(x) - \frac{i}{\cosh(x) - i \sinh(x)} \right) dx \\
 &= -(i \int \operatorname{csch}(x) dx) + \int \frac{1}{\cosh(x) - i \sinh(x)} dx \\
 &= i \tanh^{-1}(\cosh(x)) + i \operatorname{Subst} \left(\int \frac{1}{2 - x^2} dx, x, -\cosh(x) - i \sinh(x) \right) \\
 &= i \tanh^{-1}(\cosh(x)) - \frac{i \tanh^{-1} \left(\frac{\cosh(x) + i \sinh(x)}{\sqrt{2}} \right)}{\sqrt{2}}
 \end{aligned}$$

Mathematica [A]

time = 0.05, size = 46, normalized size = 1.39

$$-i \left(\sqrt{2} \tanh^{-1} \left(\frac{1 + i \tanh \left(\frac{x}{2} \right)}{\sqrt{2}} \right) - \log \left(\cosh \left(\frac{x}{2} \right) \right) + \log \left(\sinh \left(\frac{x}{2} \right) \right) \right)$$

Antiderivative was successfully verified.

[In] Integrate[Csch[x]/(I + Tanh[x]), x]

[Out] (-I)*(Sqrt[2]*ArcTanh[(1 + I*Tanh[x/2])/Sqrt[2]] - Log[Cosh[x/2]] + Log[Sinh[x/2]])

Maple [A]

time = 0.93, size = 29, normalized size = 0.88

method	result	size
default	$\sqrt{2} \arctan \left(\frac{(2 \tanh(\frac{x}{2}) - 2i) \sqrt{2}}{4} \right) - i \ln \left(\tanh \left(\frac{x}{2} \right) \right)$	29

risch	$i \ln(e^x + 1) + \frac{i\sqrt{2} \ln\left(e^x - \frac{\sqrt{2}}{2} + \frac{i\sqrt{2}}{2}\right)}{2} - \frac{i\sqrt{2} \ln\left(e^x + \frac{\sqrt{2}}{2} - \frac{i\sqrt{2}}{2}\right)}{2} - i \ln(e^x - 1)$	60
-------	---	----

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(csch(x)/(I+tanh(x)),x,method=_RETURNVERBOSE)`

[Out] $2^{(1/2)}*\arctan(1/4*(2*\tanh(1/2*x)-2*I)*2^{(1/2)})-I*\ln(\tanh(1/2*x))$

Maxima [A]

time = 0.53, size = 34, normalized size = 1.03

$$-\sqrt{2} \arctan\left(\left(\frac{1}{2}i + \frac{1}{2}\right) \sqrt{2} e^{(-x)}\right) + i \log(e^{(-x)} + 1) - i \log(e^{(-x)} - 1)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csch(x)/(I+tanh(x)),x, algorithm="maxima")`

[Out] $-\sqrt{2}*\arctan((1/2*I + 1/2)*\sqrt{2}*e^{(-x)}) + I*\log(e^{(-x)} + 1) - I*\log(e^{(-x)} - 1)$

Fricas [A]

time = 0.38, size = 43, normalized size = 1.30

$$-\frac{1}{2}i\sqrt{2} \log\left(-\left(\frac{1}{2}i - \frac{1}{2}\right) \sqrt{2} + e^x\right) + \frac{1}{2}i\sqrt{2} \log\left(\left(\frac{1}{2}i - \frac{1}{2}\right) \sqrt{2} + e^x\right) + i \log(e^x + 1) - i \log(e^x - 1)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csch(x)/(I+tanh(x)),x, algorithm="fricas")`

[Out] $-1/2*I*\sqrt{2}*\log(-(1/2*I - 1/2)*\sqrt{2} + e^x) + 1/2*I*\sqrt{2}*\log((1/2*I - 1/2)*\sqrt{2} + e^x) + I*\log(e^x + 1) - I*\log(e^x - 1)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{csch}(x)}{\tanh(x) + i} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csch(x)/(I+tanh(x)),x)`

[Out] `Integral(csch(x)/(tanh(x) + I), x)`

Giac [A]

time = 0.40, size = 28, normalized size = 0.85

$$\sqrt{2} \arctan\left(-\left(\frac{1}{2}i - \frac{1}{2}\right) \sqrt{2} e^x\right) + i \log(e^x + 1) - i \log(|e^x - 1|)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csch(x)/(1+tanh(x)),x, algorithm="giac")

[Out] sqrt(2)*arctan(-(1/2*I - 1/2)*sqrt(2)*e^x) + I*log(e^x + 1) - I*log(abs(e^x - 1))

Mupad [B]

time = 0.46, size = 61, normalized size = 1.85

$$\ln(-8e^x - 8) i - \ln(8 - 8e^x) i - \frac{\sqrt{2} \ln(e^x(4 - 4i) - \sqrt{2} 4i) i}{2} + \frac{\sqrt{2} \ln(e^x(4 - 4i) + \sqrt{2} 4i) i}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(sinh(x)*(tanh(x) + 1i)),x)

[Out] log(- 8*exp(x) - 8)*1i - log(8 - 8*exp(x))*1i - (2^(1/2)*log(exp(x)*(4 - 4i) - 2^(1/2)*4i)*1i)/2 + (2^(1/2)*log(exp(x)*(4 - 4i) + 2^(1/2)*4i)*1i)/2

3.91 $\int \frac{\cosh^4(x)}{1+\tanh(x)} dx$

Optimal. Leaf size=60

$$\frac{5x}{16} + \frac{1}{32(1-\tanh(x))^2} + \frac{1}{8(1-\tanh(x))} - \frac{1}{24(1+\tanh(x))^3} - \frac{3}{32(1+\tanh(x))^2} - \frac{3}{16(1+\tanh(x))}$$

[Out] 5/16*x+1/32/(1-tanh(x))^2+1/8/(1-tanh(x))-1/24/(1+tanh(x))^3-3/32/(1+tanh(x))^2-3/16/(1+tanh(x))

Rubi [A]

time = 0.04, antiderivative size = 60, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3568, 46, 213}

$$\frac{5x}{16} + \frac{1}{8(1-\tanh(x))} - \frac{3}{16(\tanh(x)+1)} + \frac{1}{32(1-\tanh(x))^2} - \frac{3}{32(\tanh(x)+1)^2} - \frac{1}{24(\tanh(x)+1)^3}$$

Antiderivative was successfully verified.

[In] Int[Cosh[x]^4/(1 + Tanh[x]), x]

[Out] (5*x)/16 + 1/(32*(1 - Tanh[x])^2) + 1/(8*(1 - Tanh[x])) - 1/(24*(1 + Tanh[x])^3) - 3/(32*(1 + Tanh[x])^2) - 3/(16*(1 + Tanh[x]))

Rule 46

Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && ILtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m + n + 2, 0])

Rule 213

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])

Rule 3568

Int[sec[(e_.) + (f_.)*(x_)]^(m_)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> Dist[1/(a^(m-2)*b*f), Subst[Int[(a-x)^(m/2-1)*(a+x)^(n+m/2-1), x], x, b*Tan[e+f*x]], x] /; FreeQ[{a, b, e, f, n}, x] && EqQ[a^2 + b^2, 0] && IntegerQ[m/2]

Rubi steps

$$\begin{aligned}
\int \frac{\cosh^4(x)}{1 + \tanh(x)} dx &= \text{Subst} \left(\int \frac{1}{(1-x)^3(1+x)^4} dx, x, \tanh(x) \right) \\
&= \text{Subst} \left(\int \left(-\frac{1}{16(-1+x)^3} + \frac{1}{8(-1+x)^2} + \frac{1}{8(1+x)^4} + \frac{3}{16(1+x)^3} + \frac{3}{16(1+x)^2} - \frac{1}{16(1+x)} \right) dx, x, \tanh(x) \right) \\
&= \frac{1}{32(1 - \tanh(x))^2} + \frac{1}{8(1 - \tanh(x))} - \frac{1}{24(1 + \tanh(x))^3} - \frac{3}{32(1 + \tanh(x))^2} - \frac{1}{16(1 + \tanh(x))} \\
&= \frac{5x}{16} + \frac{1}{32(1 - \tanh(x))^2} + \frac{1}{8(1 - \tanh(x))} - \frac{1}{24(1 + \tanh(x))^3} - \frac{3}{32(1 + \tanh(x))^2} - \frac{1}{16(1 + \tanh(x))}
\end{aligned}$$

Mathematica [A]

time = 0.03, size = 42, normalized size = 0.70

$$\frac{1}{192}(60x - 15 \cosh(2x) - 6 \cosh(4x) - \cosh(6x) + 45 \sinh(2x) + 9 \sinh(4x) + \sinh(6x))$$

Antiderivative was successfully verified.

`[In] Integrate[Cosh[x]^4/(1 + Tanh[x]), x]`

```
[Out] (60*x - 15*Cosh[2*x] - 6*Cosh[4*x] - Cosh[6*x] + 45*Sinh[2*x] + 9*Sinh[4*x]
+ Sinh[6*x])/192
```

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 115 vs. 2(48) = 96.

time = 0.49, size = 116, normalized size = 1.93

method	result
risch	$\frac{5x}{16} + \frac{e^{4x}}{128} + \frac{5e^{2x}}{64} - \frac{5e^{-2x}}{32} - \frac{5e^{-4x}}{128} - \frac{e^{-6x}}{192}$
default	$\frac{1}{8(\tanh(\frac{x}{2})-1)^4} + \frac{1}{4(\tanh(\frac{x}{2})-1)^3} + \frac{1}{2(\tanh(\frac{x}{2})-1)^2} + \frac{3}{8(\tanh(\frac{x}{2})-1)} - \frac{5 \ln(\tanh(\frac{x}{2})-1)}{16} - \frac{1}{3(\tanh(\frac{x}{2})+1)^6} + \frac{1}{(\tanh(\frac{x}{2})+1)^5}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cosh(x)^4/(1+tanh(x)),x,method=_RETURNVERBOSE)`

```
[Out] 1/8/(tanh(1/2*x)-1)^4+1/4/(tanh(1/2*x)-1)^3+1/2/(tanh(1/2*x)-1)^2+3/8/(tanh
(1/2*x)-1)-5/16*ln(tanh(1/2*x)-1)-1/3/(tanh(1/2*x)+1)^6+1/(tanh(1/2*x)+1)^5
-15/8/(tanh(1/2*x)+1)^4+25/12/(tanh(1/2*x)+1)^3-15/8/(tanh(1/2*x)+1)^2+1/(t
anh(1/2*x)+1)+5/16*ln(tanh(1/2*x)+1)
```

Maxima [A]

time = 0.27, size = 36, normalized size = 0.60

$$\frac{1}{128} (10e^{(-2x)} + 1)e^{(4x)} + \frac{5}{16}x - \frac{5}{32}e^{(-2x)} - \frac{5}{128}e^{(-4x)} - \frac{1}{192}e^{(-6x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cosh(x)^4/(1+tanh(x)),x, algorithm="maxima")

[Out] 1/128*(10*e^(-2*x) + 1)*e^(4*x) + 5/16*x - 5/32*e^(-2*x) - 5/128*e^(-4*x) - 1/192*e^(-6*x)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 95 vs. 2(44) = 88.

time = 0.36, size = 95, normalized size = 1.58

$$\frac{\cosh(x)^5 + 5 \cosh(x) \sinh(x)^4 + 5 \sinh(x)^5 + 5(10 \cosh(x)^2 + 9) \sinh(x)^3 + 15 \cosh(x)^3 + 5(2 \cosh(x)^3 + 9 \cosh(x)) \sinh(x)^2 + 60(2x - 1) \cosh(x) + 5(5 \cosh(x)^4 + 27 \cosh(x)^2 + 24x + 12) \sinh(x)}{384(\cosh(x) + \sinh(x))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cosh(x)^4/(1+tanh(x)),x, algorithm="fricas")

[Out] 1/384*(cosh(x)^5 + 5*cosh(x)*sinh(x)^4 + 5*sinh(x)^5 + 5*(10*cosh(x)^2 + 9)*sinh(x)^3 + 15*cosh(x)^3 + 5*(2*cosh(x)^3 + 9*cosh(x))*sinh(x)^2 + 60*(2*x - 1)*cosh(x) + 5*(5*cosh(x)^4 + 27*cosh(x)^2 + 24*x + 12)*sinh(x))/(cosh(x) + sinh(x))

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cosh^4(x)}{\tanh(x) + 1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cosh(x)**4/(1+tanh(x)),x)

[Out] Integral(cosh(x)**4/(tanh(x) + 1), x)

Giac [A]

time = 0.40, size = 42, normalized size = 0.70

$$-\frac{1}{384} (110 e^{6x} + 60 e^{4x} + 15 e^{2x} + 2) e^{-6x} + \frac{5}{16} x + \frac{1}{128} e^{4x} + \frac{5}{64} e^{2x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cosh(x)^4/(1+tanh(x)),x, algorithm="giac")

[Out] -1/384*(110*e^(6*x) + 60*e^(4*x) + 15*e^(2*x) + 2)*e^(-6*x) + 5/16*x + 1/128*e^(4*x) + 5/64*e^(2*x)

Mupad [B]

time = 1.29, size = 34, normalized size = 0.57

$$\frac{5x}{16} - \frac{5e^{-2x}}{32} + \frac{5e^{2x}}{64} - \frac{5e^{-4x}}{128} + \frac{e^{4x}}{128} - \frac{e^{-6x}}{192}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cosh(x)^4/(tanh(x) + 1),x)
```

```
[Out] (5*x)/16 - (5*exp(-2*x))/32 + (5*exp(2*x))/64 - (5*exp(-4*x))/128 + exp(4*x)/128 - exp(-6*x)/192
```

3.92 $\int \frac{\cosh^3(x)}{1+\tanh(x)} dx$

Optimal. Leaf size=29

$$\frac{4 \sinh(x)}{5} + \frac{4 \sinh^3(x)}{15} - \frac{\cosh^3(x)}{5(1 + \tanh(x))}$$

[Out] 4/5*sinh(x)+4/15*sinh(x)^3-1/5*cosh(x)^3/(1+tanh(x))

Rubi [A]

time = 0.03, antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {3583, 2713}

$$\frac{4 \sinh^3(x)}{15} + \frac{4 \sinh(x)}{5} - \frac{\cosh^3(x)}{5(\tanh(x) + 1)}$$

Antiderivative was successfully verified.

[In] Int[Cosh[x]^3/(1 + Tanh[x]),x]

[Out] (4*Sinh[x])/5 + (4*Sinh[x]^3)/15 - Cosh[x]^3/(5*(1 + Tanh[x]))

Rule 2713

Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Dist[-d^(-1), Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]

Rule 3583

Int[((d_.)*sec[(e_.) + (f_.)*(x_)]^(m_.))*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]^(n_)), x_Symbol] := Simp[a*(d*Sec[e + f*x])^m*((a + b*Tan[e + f*x])^n/(b*f*(m + 2*n))), x] + Dist[Simplify[m + n]/(a*(m + 2*n)), Int[(d*Sec[e + f*x])^m*(a + b*Tan[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, m}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0] && NeQ[m + 2*n, 0] && IntegersQ[2*m, 2*n]

Rubi steps

$$\begin{aligned} \int \frac{\cosh^3(x)}{1 + \tanh(x)} dx &= -\frac{\cosh^3(x)}{5(1 + \tanh(x))} + \frac{4}{5} \int \cosh^3(x) dx \\ &= -\frac{\cosh^3(x)}{5(1 + \tanh(x))} + \frac{4}{5} i \text{Subst} \left(\int (1 - x^2) dx, x, -i \sinh(x) \right) \\ &= \frac{4 \sinh(x)}{5} + \frac{4 \sinh^3(x)}{15} - \frac{\cosh^3(x)}{5(1 + \tanh(x))} \end{aligned}$$

Mathematica [A]

time = 0.04, size = 36, normalized size = 1.24

$$\frac{\operatorname{sech}(x)(-45 + 20 \cosh(2x) + \cosh(4x) + 40 \sinh(2x) + 4 \sinh(4x))}{120(1 + \tanh(x))}$$

Antiderivative was successfully verified.

[In] Integrate[Cosh[x]^3/(1 + Tanh[x]), x]

[Out] (Sech[x]*(-45 + 20*Cosh[2*x] + Cosh[4*x] + 40*Sinh[2*x] + 4*Sinh[4*x]))/(120*(1 + Tanh[x]))

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 79 vs. 2(23) = 46.

time = 0.46, size = 80, normalized size = 2.76

method	result
risch	$\frac{e^{3x}}{48} + \frac{e^x}{4} - \frac{3e^{-x}}{8} - \frac{e^{-3x}}{12} - \frac{e^{-5x}}{80}$
default	$-\frac{1}{6(\tanh(\frac{x}{2})-1)^3} - \frac{1}{4(\tanh(\frac{x}{2})-1)^2} - \frac{5}{8(\tanh(\frac{x}{2})-1)} - \frac{2}{5(\tanh(\frac{x}{2})+1)^5} + \frac{1}{(\tanh(\frac{x}{2})+1)^4} - \frac{5}{3(\tanh(\frac{x}{2})+1)^3} + \frac{1}{2(\tanh(\frac{x}{2})+1)^2}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cosh(x)^3/(1+tanh(x)), x, method=_RETURNVERBOSE)

[Out] -1/6/(tanh(1/2*x)-1)^3-1/4/(tanh(1/2*x)-1)^2-5/8/(tanh(1/2*x)-1)-2/5/(tanh(1/2*x)+1)^5+1/(tanh(1/2*x)+1)^4-5/3/(tanh(1/2*x)+1)^3+3/2/(tanh(1/2*x)+1)^2-11/8/(tanh(1/2*x)+1)

Maxima [A]

time = 0.26, size = 33, normalized size = 1.14

$$\frac{1}{48} (12 e^{(-2x)} + 1) e^{(3x)} - \frac{3}{8} e^{(-x)} - \frac{1}{12} e^{(-3x)} - \frac{1}{80} e^{(-5x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cosh(x)^3/(1+tanh(x)), x, algorithm="maxima")

[Out] 1/48*(12*e^(-2*x) + 1)*e^(3*x) - 3/8*e^(-x) - 1/12*e^(-3*x) - 1/80*e^(-5*x)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 60 vs. 2(23) = 46.

time = 0.42, size = 60, normalized size = 2.07

$$\frac{\cosh(x)^4 + 16 \cosh(x) \sinh(x)^3 + \sinh(x)^4 + 2(3 \cosh(x)^2 + 10) \sinh(x)^2 + 20 \cosh(x)^2 + 16(\cosh(x)^3 + 5 \cosh(x)) \sinh(x) - 45}{120(\cosh(x) + \sinh(x))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cosh(x)^3/(1+tanh(x)),x, algorithm="fricas")

[Out] 1/120*(cosh(x)^4 + 16*cosh(x)*sinh(x)^3 + sinh(x)^4 + 2*(3*cosh(x)^2 + 10)*sinh(x)^2 + 20*cosh(x)^2 + 16*(cosh(x)^3 + 5*cosh(x))*sinh(x) - 45)/(cosh(x) + sinh(x))

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 134 vs. 2(26) = 52.

time = 0.35, size = 134, normalized size = 4.62

$$-\frac{8 \sinh^3(x) \tanh(x)}{15 \tanh(x) + 15} - \frac{2 \sinh^3(x)}{15 \tanh(x) + 15} - \frac{6 \sinh^2(x) \cosh(x) \tanh(x)}{15 \tanh(x) + 15} + \frac{6 \sinh^2(x) \cosh(x)}{15 \tanh(x) + 15} + \frac{6 \sinh(x) \cosh^2(x) \tanh(x)}{15 \tanh(x) + 15} + \frac{9 \sinh(x) \cosh^2(x)}{15 \tanh(x) + 15} + \frac{3 \cosh^3(x) \tanh(x)}{15 \tanh(x) + 15} - \frac{3 \cosh^3(x)}{15 \tanh(x) + 15}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cosh(x)**3/(1+tanh(x)),x)

[Out] -8*sinh(x)**3*tanh(x)/(15*tanh(x) + 15) - 2*sinh(x)**3/(15*tanh(x) + 15) - 6*sinh(x)**2*cosh(x)*tanh(x)/(15*tanh(x) + 15) + 6*sinh(x)**2*cosh(x)/(15*tanh(x) + 15) + 6*sinh(x)*cosh(x)**2*tanh(x)/(15*tanh(x) + 15) + 9*sinh(x)*cosh(x)**2/(15*tanh(x) + 15) + 3*cosh(x)**3*tanh(x)/(15*tanh(x) + 15) - 3*cosh(x)**3/(15*tanh(x) + 15)

Giac [A]

time = 0.40, size = 31, normalized size = 1.07

$$-\frac{1}{240} (90 e^{4x} + 20 e^{2x} + 3) e^{-5x} + \frac{1}{48} e^{3x} + \frac{1}{4} e^x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cosh(x)^3/(1+tanh(x)),x, algorithm="giac")

[Out] -1/240*(90*e^(4*x) + 20*e^(2*x) + 3)*e^(-5*x) + 1/48*e^(3*x) + 1/4*e^x

Mupad [B]

time = 1.19, size = 29, normalized size = 1.00

$$\frac{e^{3x}}{48} - \frac{e^{-3x}}{12} - \frac{3e^{-x}}{8} - \frac{e^{-5x}}{80} + \frac{e^x}{4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cosh(x)^3/(tanh(x) + 1),x)

[Out] exp(3*x)/48 - exp(-3*x)/12 - (3*exp(-x))/8 - exp(-5*x)/80 + exp(x)/4

3.93 $\int \frac{\cosh^2(x)}{1+\tanh(x)} dx$

Optimal. Leaf size=38

$$\frac{3x}{8} + \frac{1}{8(1-\tanh(x))} - \frac{1}{8(1+\tanh(x))^2} - \frac{1}{4(1+\tanh(x))}$$

[Out] 3/8*x+1/8/(1-tanh(x))-1/8/(1+tanh(x))^2-1/4/(1+tanh(x))

Rubi [A]

time = 0.03, antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3568, 46, 213}

$$\frac{3x}{8} + \frac{1}{8(1-\tanh(x))} - \frac{1}{4(\tanh(x)+1)} - \frac{1}{8(\tanh(x)+1)^2}$$

Antiderivative was successfully verified.

[In] Int[Cosh[x]^2/(1 + Tanh[x]),x]

[Out] (3*x)/8 + 1/(8*(1 - Tanh[x])) - 1/(8*(1 + Tanh[x])^2) - 1/(4*(1 + Tanh[x]))

Rule 46

Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && ILtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m + n + 2, 0])

Rule 213

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])

Rule 3568

Int[sec[(e_.) + (f_.)*(x_)]^(m_)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Dist[1/(a^(m-2)*b*f), Subst[Int[(a-x)^(m/2-1)*(a+x)^(n+m/2-1), x], x, b*Tan[e+f*x]], x] /; FreeQ[{a, b, e, f, n}, x] && EqQ[a^2 + b^2, 0] && IntegerQ[m/2]

Rubi steps

$$\begin{aligned}
\int \frac{\cosh^2(x)}{1 + \tanh(x)} dx &= \text{Subst} \left(\int \frac{1}{(1-x)^2(1+x)^3} dx, x, \tanh(x) \right) \\
&= \text{Subst} \left(\int \left(\frac{1}{8(-1+x)^2} + \frac{1}{4(1+x)^3} + \frac{1}{4(1+x)^2} - \frac{3}{8(-1+x^2)} \right) dx, x, \tanh(x) \right) \\
&= \frac{1}{8(1 - \tanh(x))} - \frac{1}{8(1 + \tanh(x))^2} - \frac{1}{4(1 + \tanh(x))} - \frac{3}{8} \text{Subst} \left(\int \frac{1}{-1+x^2} dx, x, \tanh(x) \right) \\
&= \frac{3x}{8} + \frac{1}{8(1 - \tanh(x))} - \frac{1}{8(1 + \tanh(x))^2} - \frac{1}{4(1 + \tanh(x))}
\end{aligned}$$

Mathematica [A]

time = 0.02, size = 30, normalized size = 0.79

$$\frac{1}{32}(12x - 4 \cosh(2x) - \cosh(4x) + 8 \sinh(2x) + \sinh(4x))$$

Antiderivative was successfully verified.

`[In] Integrate[Cosh[x]^2/(1 + Tanh[x]), x]``[Out] (12*x - 4*Cosh[2*x] - Cosh[4*x] + 8*Sinh[2*x] + Sinh[4*x])/32`**Maple [B]** Leaf count of result is larger than twice the leaf count of optimal. 75 vs. 2(30) = 60.

time = 0.50, size = 76, normalized size = 2.00

method	result
risch	$\frac{3x}{8} + \frac{e^{2x}}{16} - \frac{3e^{-2x}}{16} - \frac{e^{-4x}}{32}$
default	$-\frac{1}{2(\tanh(\frac{x}{2})+1)^4} + \frac{1}{(\tanh(\frac{x}{2})+1)^3} - \frac{3}{2(\tanh(\frac{x}{2})+1)^2} + \frac{1}{\tanh(\frac{x}{2})+1} + \frac{3 \ln(\tanh(\frac{x}{2})+1)}{8} + \frac{1}{4(\tanh(\frac{x}{2})-1)^2} + \frac{1}{4 \tanh(\frac{x}{2})}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cosh(x)^2/(1+tanh(x)), x, method=_RETURNVERBOSE)`

```
[Out] -1/2/(tanh(1/2*x)+1)^4+1/(tanh(1/2*x)+1)^3-3/2/(tanh(1/2*x)+1)^2+1/(tanh(1/2*x)+1)+3/8*ln(tanh(1/2*x)+1)+1/4/(tanh(1/2*x)-1)^2+1/4/(tanh(1/2*x)-1)-3/8*ln(tanh(1/2*x)-1)
```

Maxima [A]

time = 0.26, size = 22, normalized size = 0.58

$$\frac{3}{8}x + \frac{1}{16}e^{(2x)} - \frac{3}{16}e^{(-2x)} - \frac{1}{32}e^{(-4x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cosh(x)^2/(1+tanh(x)),x, algorithm="maxima")

[Out] 3/8*x + 1/16*e^(2*x) - 3/16*e^(-2*x) - 1/32*e^(-4*x)

Fricas [A]

time = 0.38, size = 52, normalized size = 1.37

$$\frac{\cosh(x)^3 + 3 \cosh(x) \sinh(x)^2 + 3 \sinh(x)^3 + 6(2x - 1) \cosh(x) + 3(3 \cosh(x)^2 + 4x + 2) \sinh(x)}{32(\cosh(x) + \sinh(x))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cosh(x)^2/(1+tanh(x)),x, algorithm="fricas")

[Out] 1/32*(cosh(x)^3 + 3*cosh(x)*sinh(x)^2 + 3*sinh(x)^3 + 6*(2*x - 1)*cosh(x) + 3*(3*cosh(x)^2 + 4*x + 2)*sinh(x))/(cosh(x) + sinh(x))

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cosh^2(x)}{\tanh(x) + 1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cosh(x)**2/(1+tanh(x)),x)

[Out] Integral(cosh(x)**2/(tanh(x) + 1), x)

Giac [A]

time = 0.41, size = 30, normalized size = 0.79

$$-\frac{1}{32} (9e^{4x} + 6e^{2x} + 1)e^{-4x} + \frac{3}{8}x + \frac{1}{16}e^{2x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cosh(x)^2/(1+tanh(x)),x, algorithm="giac")

[Out] -1/32*(9*e^(4*x) + 6*e^(2*x) + 1)*e^(-4*x) + 3/8*x + 1/16*e^(2*x)

Mupad [B]

time = 0.13, size = 22, normalized size = 0.58

$$\frac{3x}{8} - \frac{3e^{-2x}}{16} + \frac{e^{2x}}{16} - \frac{e^{-4x}}{32}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cosh(x)^2/(tanh(x) + 1),x)

[Out] (3*x)/8 - (3*exp(-2*x))/16 + exp(2*x)/16 - exp(-4*x)/32

3.94 $\int \frac{\cosh(x)}{1+\tanh(x)} dx$

Optimal. Leaf size=19

$$\frac{2 \sinh(x)}{3} - \frac{\cosh(x)}{3(1 + \tanh(x))}$$

[Out] 2/3*sinh(x)-1/3*cosh(x)/(1+tanh(x))

Rubi [A]

time = 0.02, antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {3583, 2717}

$$\frac{2 \sinh(x)}{3} - \frac{\cosh(x)}{3(\tanh(x) + 1)}$$

Antiderivative was successfully verified.

[In] Int[Cosh[x]/(1 + Tanh[x]),x]

[Out] (2*Sinh[x])/3 - Cosh[x]/(3*(1 + Tanh[x]))

Rule 2717

Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /;
FreeQ[{c, d}, x]

Rule 3583

Int[((d_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[a*(d*Sec[e + f*x])^m*((a + b*Tan[e + f*x])^n/(b*f*(m + 2*n))), x] + Dist[Simplify[m + n]/(a*(m + 2*n)), Int[(d*Sec[e + f*x])^m*(a + b*Tan[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f, m}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0] && NeQ[m + 2*n, 0] && IntegersQ[2*m, 2*n]

Rubi steps

$$\begin{aligned} \int \frac{\cosh(x)}{1 + \tanh(x)} dx &= -\frac{\cosh(x)}{3(1 + \tanh(x))} + \frac{2}{3} \int \cosh(x) dx \\ &= \frac{2 \sinh(x)}{3} - \frac{\cosh(x)}{3(1 + \tanh(x))} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 23, normalized size = 1.21

$$\frac{1}{12}(-3 \cosh(x) - \cosh(3x) + 9 \sinh(x) + \sinh(3x))$$

Antiderivative was successfully verified.

[In] Integrate[Cosh[x]/(1 + Tanh[x]),x]

[Out] (-3*Cosh[x] - Cosh[3*x] + 9*Sinh[x] + Sinh[3*x])/12

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 39 vs. 2(15) = 30.

time = 0.49, size = 40, normalized size = 2.11

method	result	size
risch	$\frac{e^x}{4} - \frac{e^{-x}}{2} - \frac{e^{-3x}}{12}$	18
default	$-\frac{1}{2(\tanh(\frac{x}{2})-1)} - \frac{2}{3(\tanh(\frac{x}{2})+1)^3} + \frac{1}{(\tanh(\frac{x}{2})+1)^2} - \frac{3}{2(\tanh(\frac{x}{2})+1)}$	40

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cosh(x)/(1+tanh(x)),x,method=_RETURNVERBOSE)

[Out] -1/2/(tanh(1/2*x)-1)-2/3/(tanh(1/2*x)+1)^3+1/(tanh(1/2*x)+1)^2-3/2/(tanh(1/2*x)+1)

Maxima [A]

time = 0.26, size = 17, normalized size = 0.89

$$-\frac{1}{2}e^{(-x)} - \frac{1}{12}e^{(-3x)} + \frac{1}{4}e^x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cosh(x)/(1+tanh(x)),x, algorithm="maxima")

[Out] -1/2*e^(-x) - 1/12*e^(-3*x) + 1/4*e^x

Fricas [A]

time = 0.42, size = 25, normalized size = 1.32

$$\frac{\cosh(x)^2 + 4 \cosh(x) \sinh(x) + \sinh(x)^2 - 3}{6(\cosh(x) + \sinh(x))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cosh(x)/(1+tanh(x)),x, algorithm="fricas")

[Out] $1/6*(\cosh(x)^2 + 4*\cosh(x)*\sinh(x) + \sinh(x)^2 - 3)/(\cosh(x) + \sinh(x))$

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 48 vs. $2(15) = 30$.

time = 0.18, size = 48, normalized size = 2.53

$$\frac{2 \sinh(x) \tanh(x)}{3 \tanh(x) + 3} + \frac{\sinh(x)}{3 \tanh(x) + 3} + \frac{\cosh(x) \tanh(x)}{3 \tanh(x) + 3} - \frac{\cosh(x)}{3 \tanh(x) + 3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cosh(x)/(1+tanh(x)),x)`

[Out] $2*\sinh(x)*\tanh(x)/(3*\tanh(x) + 3) + \sinh(x)/(3*\tanh(x) + 3) + \cosh(x)*\tanh(x)/(3*\tanh(x) + 3) - \cosh(x)/(3*\tanh(x) + 3)$

Giac [A]

time = 0.40, size = 19, normalized size = 1.00

$$-\frac{1}{12} (6e^{2x} + 1)e^{-3x} + \frac{1}{4} e^x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cosh(x)/(1+tanh(x)),x, algorithm="giac")`

[Out] $-1/12*(6*e^{2*x} + 1)*e^{-3*x} + 1/4*e^x$

Mupad [B]

time = 1.05, size = 17, normalized size = 0.89

$$\frac{e^x}{4} - \frac{e^{-3x}}{12} - \frac{e^{-x}}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cosh(x)/(tanh(x) + 1),x)`

[Out] $\exp(x)/4 - \exp(-3*x)/12 - \exp(-x)/2$

3.95 $\int \frac{\operatorname{sech}(x)}{1+\tanh(x)} dx$

Optimal. Leaf size=10

$$-\frac{\operatorname{sech}(x)}{1+\tanh(x)}$$

[Out] `-sech(x)/(1+tanh(x))`

Rubi [A]

time = 0.01, antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {3569}

$$-\frac{\operatorname{sech}(x)}{\tanh(x)+1}$$

Antiderivative was successfully verified.

[In] `Int[Sech[x]/(1 + Tanh[x]), x]`

[Out] `-(Sech[x]/(1 + Tanh[x]))`

Rule 3569

```
Int[((d_.)*sec[(e_.) + (f_.)*(x_.)])^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] :> Simp[b*(d*Sec[e + f*x])^m*((a + b*Tan[e + f*x])^n/(a*f*m)), x] /; FreeQ[{a, b, d, e, f, m, n}, x] && EqQ[a^2 + b^2, 0] && EqQ[Simplify[m + n], 0]
```

Rubi steps

$$\int \frac{\operatorname{sech}(x)}{1+\tanh(x)} dx = -\frac{\operatorname{sech}(x)}{1+\tanh(x)}$$

Mathematica [A]

time = 0.01, size = 7, normalized size = 0.70

$$-\cosh(x) + \sinh(x)$$

Antiderivative was successfully verified.

[In] `Integrate[Sech[x]/(1 + Tanh[x]), x]`

[Out] `-Cosh[x] + Sinh[x]`

Maple [A]

time = 0.26, size = 11, normalized size = 1.10

method	result	size
risch	$-e^{-x}$	7
gosper	$-\frac{\operatorname{sech}(x)}{1+\tanh(x)}$	11
default	$-\frac{2}{\tanh(\frac{x}{2})+1}$	11

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sech(x)/(1+tanh(x)),x,method=_RETURNVERBOSE)``[Out] -2/(tanh(1/2*x)+1)`**Maxima [A]**

time = 0.27, size = 6, normalized size = 0.60

$$-e^{(-x)}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sech(x)/(1+tanh(x)),x, algorithm="maxima")``[Out] -e^(-x)`**Fricas [A]**

time = 0.37, size = 9, normalized size = 0.90

$$-\frac{1}{\cosh(x) + \sinh(x)}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sech(x)/(1+tanh(x)),x, algorithm="fricas")``[Out] -1/(cosh(x) + sinh(x))`**Sympy [A]**

time = 0.16, size = 8, normalized size = 0.80

$$-\frac{\operatorname{sech}(x)}{\tanh(x) + 1}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sech(x)/(1+tanh(x)),x)``[Out] -sech(x)/(tanh(x) + 1)`

Giac [A]

time = 0.41, size = 6, normalized size = 0.60

$$-e^{(-x)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sech(x)/(1+tanh(x)),x, algorithm="giac")
```

```
[Out] -e^(-x)
```

Mupad [B]

time = 0.04, size = 6, normalized size = 0.60

$$-e^{-x}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(cosh(x)*(tanh(x) + 1)),x)
```

```
[Out] -exp(-x)
```

3.96 $\int \frac{\operatorname{sech}^2(x)}{1+\tanh(x)} dx$

Optimal. Leaf size=5

$$\log(1 + \tanh(x))$$

[Out] ln(1+tanh(x))

Rubi [A]

time = 0.02, antiderivative size = 5, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {3568, 31}

$$\log(\tanh(x) + 1)$$

Antiderivative was successfully verified.

[In] Int[Sech[x]^2/(1 + Tanh[x]),x]

[Out] Log[1 + Tanh[x]]

Rule 31

Int[((a_) + (b_.)*(x_))^(n_), x_Symbol] := Simp[Log[RemoveContent[a + b*x, x]]/b, x] /; FreeQ[{a, b}, x]

Rule 3568

Int[sec[(e_.) + (f_.)*(x_)]^(m_)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Dist[1/(a^(m - 2)*b*f), Subst[Int[(a - x)^(m/2 - 1)*(a + x)^(n + m/2 - 1), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, n}, x] && EqQ[a^2 + b^2, 0] && IntegerQ[m/2]

Rubi steps

$$\begin{aligned} \int \frac{\operatorname{sech}^2(x)}{1 + \tanh(x)} dx &= \operatorname{Subst} \left(\int \frac{1}{1 + x} dx, x, \tanh(x) \right) \\ &= \log(1 + \tanh(x)) \end{aligned}$$

Mathematica [A]

time = 0.00, size = 7, normalized size = 1.40

$$x - \log(\cosh(x))$$

Antiderivative was successfully verified.

[In] Integrate[Sech[x]^2/(1 + Tanh[x]),x]

[Out] x - Log[Cosh[x]]

Maple [A]

time = 0.51, size = 6, normalized size = 1.20

method	result	size
derivativedivides	$\ln(1 + \tanh(x))$	6
default	$\ln(1 + \tanh(x))$	6
risch	$2x - \ln(1 + e^{2x})$	14

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sech(x)^2/(1+tanh(x)),x,method=_RETURNVERBOSE)

[Out] ln(1+tanh(x))

Maxima [A]

time = 0.27, size = 5, normalized size = 1.00

$$\log(\tanh(x) + 1)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)^2/(1+tanh(x)),x, algorithm="maxima")

[Out] log(tanh(x) + 1)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 20 vs. $2(5) = 10$.

time = 0.34, size = 20, normalized size = 4.00

$$2x - \log\left(\frac{2 \cosh(x)}{\cosh(x) - \sinh(x)}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)^2/(1+tanh(x)),x, algorithm="fricas")

[Out] 2*x - log(2*cosh(x)/(cosh(x) - sinh(x)))

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{sech}^2(x)}{\tanh(x) + 1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)**2/(1+tanh(x)),x)

[Out] `Integral(sech(x)**2/(tanh(x) + 1), x)`

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 13 vs. 2(5) = 10.
time = 0.41, size = 13, normalized size = 2.60

$$2x - \log(e^{(2x)} + 1)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sech(x)^2/(1+tanh(x)),x, algorithm="giac")`

[Out] `2*x - log(e^(2*x) + 1)`

Mupad [B]

time = 1.05, size = 13, normalized size = 2.60

$$2x - \ln(e^{2x} + 1)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(cosh(x)^2*(tanh(x) + 1)),x)`

[Out] `2*x - log(exp(2*x) + 1)`

3.97 $\int \frac{\operatorname{sech}^3(x)}{1+\tanh(x)} dx$

Optimal. Leaf size=6

$$\operatorname{ArcTan}(\sinh(x)) + \operatorname{sech}(x)$$

[Out] arctan(sinh(x))+sech(x)

Rubi [A]

time = 0.02, antiderivative size = 6, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {3582, 3855}

$$\operatorname{ArcTan}(\sinh(x)) + \operatorname{sech}(x)$$

Antiderivative was successfully verified.

[In] Int[Sech[x]^3/(1 + Tanh[x]),x]

[Out] ArcTan[Sinh[x]] + Sech[x]

Rule 3582

```
Int[((d_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> Simp[d^2*(d*Sec[e + f*x])^(m - 2)*((a + b*Tan[e + f*x])^(n + 1)/(b*f*(m + n - 1))), x] + Dist[d^2*((m - 2)/(a*(m + n - 1))), Int[(d*Sec[e + f*x])^(m - 2)*(a + b*Tan[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0] && GtQ[m, 1] && !LtQ[m + n, 0] && NeQ[m + n - 1, 0] && IntegersQ[2*m, 2*n]
```

Rule 3855

```
Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned} \int \frac{\operatorname{sech}^3(x)}{1+\tanh(x)} dx &= \operatorname{sech}(x) + \int \operatorname{sech}(x) dx \\ &= \tan^{-1}(\sinh(x)) + \operatorname{sech}(x) \end{aligned}$$

Mathematica [A]

time = 0.02, size = 12, normalized size = 2.00

$$2\operatorname{ArcTan}\left(\tanh\left(\frac{x}{2}\right)\right) + \operatorname{sech}(x)$$

Antiderivative was successfully verified.

[In] Integrate[Sech[x]^3/(1 + Tanh[x]),x]

[Out] 2*ArcTan[Tanh[x/2]] + Sech[x]

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 20 vs. $2(6) = 12$.

time = 0.50, size = 21, normalized size = 3.50

method	result	size
default	$\frac{2}{\tanh^2(\frac{x}{2})+1} + 2 \arctan(\tanh(\frac{x}{2}))$	21
risch	$\frac{2e^x}{1+e^{2x}} + i \ln(e^x + i) - i \ln(e^x - i)$	32

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sech(x)^3/(1+tanh(x)),x,method=_RETURNVERBOSE)

[Out] 2/(tanh(1/2*x)^2+1)+2*arctan(tanh(1/2*x))

Maxima [B] Leaf count of result is larger than twice the leaf count of optimal. 22 vs. $2(6) = 12$.

time = 0.48, size = 22, normalized size = 3.67

$$\frac{2e^{(-x)}}{e^{(-2x)} + 1} - 2 \arctan(e^{(-x)})$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)^3/(1+tanh(x)),x, algorithm="maxima")

[Out] 2*e^(-x)/(e^(-2*x) + 1) - 2*arctan(e^(-x))

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 48 vs. $2(6) = 12$.
time = 0.35, size = 48, normalized size = 8.00

$$\frac{2((\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 + 1) \arctan(\cosh(x) + \sinh(x)) + \cosh(x) + \sinh(x))}{\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 + 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)^3/(1+tanh(x)),x, algorithm="fricas")

[Out] 2*((cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)*arctan(cosh(x) + sinh(x)) + cosh(x) + sinh(x))/(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{sech}^3(x)}{\tanh(x) + 1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sech(x)**3/(1+tanh(x)),x)`

[Out] `Integral(sech(x)**3/(tanh(x) + 1), x)`

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 18 vs. 2(6) = 12.
time = 0.41, size = 18, normalized size = 3.00

$$\frac{2e^x}{e^{2x} + 1} + 2 \arctan(e^x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sech(x)^3/(1+tanh(x)),x, algorithm="giac")`

[Out] `2*e^x/(e^(2*x) + 1) + 2*arctan(e^x)`

Mupad [B]

time = 1.04, size = 18, normalized size = 3.00

$$2 \operatorname{atan}(e^x) + \frac{2e^x}{e^{2x} + 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(cosh(x)^3*(tanh(x) + 1)),x)`

[Out] `2*atan(exp(x)) + (2*exp(x))/(exp(2*x) + 1)`

3.98 $\int \frac{\operatorname{sech}^4(x)}{1+\tanh(x)} dx$

Optimal. Leaf size=11

$$\tanh(x) - \frac{\tanh^2(x)}{2}$$

[Out] $\tanh(x) - 1/2 * \tanh(x)^2$

Rubi [A]

time = 0.02, antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {3568}

$$\tanh(x) - \frac{\tanh^2(x)}{2}$$

Antiderivative was successfully verified.

[In] `Int[Sech[x]^4/(1 + Tanh[x]), x]`

[Out] `Tanh[x] - Tanh[x]^2/2`

Rule 3568

`Int[sec[(e_.) + (f_.)*(x_)]^(m_)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> Dist[1/(a^(m - 2)*b*f), Subst[Int[(a - x)^(m/2 - 1)*(a + x)^(n + m/2 - 1), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, n}, x] && EqQ[a^2 + b^2, 0] && IntegerQ[m/2]`

Rubi steps

$$\begin{aligned} \int \frac{\operatorname{sech}^4(x)}{1+\tanh(x)} dx &= \operatorname{Subst}\left(\int (1-x) dx, x, \tanh(x)\right) \\ &= \tanh(x) - \frac{\tanh^2(x)}{2} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 11, normalized size = 1.00

$$\frac{\operatorname{sech}^2(x)}{2} + \tanh(x)$$

Antiderivative was successfully verified.

[In] `Integrate[Sech[x]^4/(1 + Tanh[x]), x]`

[Out] $\text{Sech}[x]^2/2 + \text{Tanh}[x]$

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 33 vs. $2(9) = 18$.

time = 0.55, size = 34, normalized size = 3.09

method	result	size
risch	$-\frac{2}{(1+e^{2x})^2}$	11
default	$-\frac{2(-(\tanh^3(\frac{x}{2})) + \tanh^2(\frac{x}{2}) - \tanh(\frac{x}{2}))}{(\tanh^2(\frac{x}{2}) + 1)^2}$	34

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sech(x)^4/(1+tanh(x)),x,method=_RETURNVERBOSE)`

[Out] $-2*(-\tanh(1/2*x)^3 + \tanh(1/2*x)^2 - \tanh(1/2*x))/(\tanh(1/2*x)^2 + 1)^2$

Maxima [B] Leaf count of result is larger than twice the leaf count of optimal. 37 vs. $2(9) = 18$.

time = 0.26, size = 37, normalized size = 3.36

$$\frac{4e^{-2x}}{2e^{-2x} + e^{-4x} + 1} + \frac{2}{2e^{-2x} + e^{-4x} + 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sech(x)^4/(1+tanh(x)),x, algorithm="maxima")`

[Out] $4*e^{-2*x}/(2*e^{-2*x} + e^{-4*x} + 1) + 2/(2*e^{-2*x} + e^{-4*x} + 1)$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 53 vs. $2(9) = 18$.
time = 0.37, size = 53, normalized size = 4.82

$$\frac{2}{\cosh(x)^4 + 4 \cosh(x) \sinh(x)^3 + \sinh(x)^4 + 2(3 \cosh(x)^2 + 1) \sinh(x)^2 + 2 \cosh(x)^2 + 4(\cosh(x)^3 + \cosh(x)) \sinh(x) + 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sech(x)^4/(1+tanh(x)),x, algorithm="fricas")`

[Out] $-2/(\cosh(x)^4 + 4*\cosh(x)*\sinh(x)^3 + \sinh(x)^4 + 2*(3*\cosh(x)^2 + 1)*\sinh(x)^2 + 2*\cosh(x)^2 + 4*(\cosh(x)^3 + \cosh(x))*\sinh(x) + 1)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\text{sech}^4(x)}{\tanh(x) + 1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)**4/(1+tanh(x)),x)

[Out] Integral(sech(x)**4/(tanh(x) + 1), x)

Giac [A]

time = 0.41, size = 10, normalized size = 0.91

$$-\frac{2}{(e^{2x} + 1)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)^4/(1+tanh(x)),x, algorithm="giac")

[Out] -2/(e^(2*x) + 1)^2

Mupad [B]

time = 1.04, size = 16, normalized size = 1.45

$$-\frac{2}{2e^{2x} + e^{4x} + 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cosh(x)^4*(tanh(x) + 1)),x)

[Out] -2/(2*exp(2*x) + exp(4*x) + 1)

$$3.99 \quad \int \frac{\operatorname{sech}^5(x)}{1+\tanh(x)} dx$$

Optimal. Leaf size=24

$$\frac{1}{2}\operatorname{ArcTan}(\sinh(x)) + \frac{\operatorname{sech}^3(x)}{3} + \frac{1}{2}\operatorname{sech}(x)\tanh(x)$$

[Out] 1/2*arctan(sinh(x))+1/3*sech(x)^3+1/2*sech(x)*tanh(x)

Rubi [A]

time = 0.03, antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3582, 3853, 3855}

$$\frac{1}{2}\operatorname{ArcTan}(\sinh(x)) + \frac{\operatorname{sech}^3(x)}{3} + \frac{1}{2}\tanh(x)\operatorname{sech}(x)$$

Antiderivative was successfully verified.

[In] Int[Sech[x]^5/(1 + Tanh[x]),x]

[Out] ArcTan[Sinh[x]]/2 + Sech[x]^3/3 + (Sech[x]*Tanh[x])/2

Rule 3582

Int[((d_)*sec[(e_) + (f_)*(x_)])^(m_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[d^2*(d*Sec[e + f*x])^(m - 2)*((a + b*Tan[e + f*x])^(n + 1)/(b*f*(m + n - 1))), x] + Dist[d^2*((m - 2)/(a*(m + n - 1))), Int[(d*Sec[e + f*x])^(m - 2)*(a + b*Tan[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0] && GtQ[m, 1] && !LtQ[m + n, 0] && NeQ[m + n - 1, 0] && IntegersQ[2*m, 2*n]

Rule 3853

Int[(csc[(c_) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Dist[b^2*((n - 2)/(n - 1)), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3855

Int[csc[(c_) + (d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{\operatorname{sech}^5(x)}{1 + \tanh(x)} dx &= \frac{\operatorname{sech}^3(x)}{3} + \int \operatorname{sech}^3(x) dx \\ &= \frac{\operatorname{sech}^3(x)}{3} + \frac{1}{2} \operatorname{sech}(x) \tanh(x) + \frac{1}{2} \int \operatorname{sech}(x) dx \\ &= \frac{1}{2} \tan^{-1}(\sinh(x)) + \frac{\operatorname{sech}^3(x)}{3} + \frac{1}{2} \operatorname{sech}(x) \tanh(x) \end{aligned}$$

Mathematica [A]

time = 0.02, size = 24, normalized size = 1.00

$$\operatorname{ArcTan}\left(\tanh\left(\frac{x}{2}\right)\right) + \frac{\operatorname{sech}^3(x)}{3} + \frac{1}{2} \operatorname{sech}(x) \tanh(x)$$

Antiderivative was successfully verified.

[In] Integrate[Sech[x]^5/(1 + Tanh[x]), x]

[Out] ArcTan[Tanh[x/2]] + Sech[x]^3/3 + (Sech[x]*Tanh[x])/2

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 40 vs. 2(18) = 36.

time = 0.57, size = 41, normalized size = 1.71

method	result	size
default	$\frac{-(\tanh^5(\frac{x}{2})) + 2(\tanh^4(\frac{x}{2})) + \tanh(\frac{x}{2}) + \frac{2}{3}}{(\tanh^2(\frac{x}{2}) + 1)^3} + \arctan\left(\tanh\left(\frac{x}{2}\right)\right)$	41
risch	$\frac{e^x(3e^{4x} + 8e^{2x} - 3)}{3(1 + e^{2x})^3} + \frac{i \ln(e^x + i)}{2} - \frac{i \ln(e^x - i)}{2}$	46

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sech(x)^5/(1+tanh(x)), x, method=_RETURNVERBOSE)

[Out] 2*(-1/2*tanh(1/2*x)^5+tanh(1/2*x)^4+1/2*tanh(1/2*x)+1/3)/(tanh(1/2*x)^2+1)^3+arctan(tanh(1/2*x))

Maxima [B] Leaf count of result is larger than twice the leaf count of optimal. 49 vs. 2(18) = 36.

time = 0.46, size = 49, normalized size = 2.04

$$\frac{3e^{(-x)} + 8e^{(-3x)} - 3e^{(-5x)}}{3(3e^{(-2x)} + 3e^{(-4x)} + e^{(-6x)} + 1)} - \arctan(e^{(-x)})$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)^5/(1+tanh(x)),x, algorithm="maxima")

[Out] 1/3*(3*e^(-x) + 8*e^(-3*x) - 3*e^(-5*x))/(3*e^(-2*x) + 3*e^(-4*x) + e^(-6*x) + 1) - arctan(e^(-x))

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 288 vs. 2(18) = 36.

time = 0.42, size = 288, normalized size = 12.00

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)^5/(1+tanh(x)),x, algorithm="fricas")

[Out] 1/3*(3*cosh(x)^5 + 15*cosh(x)*sinh(x)^4 + 3*sinh(x)^5 + 2*(15*cosh(x)^2 + 4)*sinh(x)^3 + 8*cosh(x)^3 + 6*(5*cosh(x)^3 + 4*cosh(x))*sinh(x)^2 + 3*(cosh(x)^6 + 6*cosh(x)*sinh(x)^5 + sinh(x)^6 + 3*(5*cosh(x)^2 + 1)*sinh(x)^4 + 3*cosh(x)^4 + 4*(5*cosh(x)^3 + 3*cosh(x))*sinh(x)^3 + 3*(5*cosh(x)^4 + 6*cosh(x)^2 + 1)*sinh(x)^2 + 3*cosh(x)^2 + 6*(cosh(x)^5 + 2*cosh(x)^3 + cosh(x))*sinh(x) + 1)*arctan(cosh(x) + sinh(x)) + 3*(5*cosh(x)^4 + 8*cosh(x)^2 - 1)*sinh(x) - 3*cosh(x))/(cosh(x)^6 + 6*cosh(x)*sinh(x)^5 + sinh(x)^6 + 3*(5*cosh(x)^2 + 1)*sinh(x)^4 + 3*cosh(x)^4 + 4*(5*cosh(x)^3 + 3*cosh(x))*sinh(x)^3 + 3*(5*cosh(x)^4 + 6*cosh(x)^2 + 1)*sinh(x)^2 + 3*cosh(x)^2 + 6*(cosh(x)^5 + 2*cosh(x)^3 + cosh(x))*sinh(x) + 1)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{sech}^5(x)}{\tanh(x) + 1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)**5/(1+tanh(x)),x)

[Out] Integral(sech(x)**5/(tanh(x) + 1), x)

Giac [A]

time = 0.41, size = 31, normalized size = 1.29

$$\frac{3e^{5x} + 8e^{3x} - 3e^x}{3(e^{2x} + 1)^3} + \arctan(e^x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)^5/(1+tanh(x)),x, algorithm="giac")

[Out] 1/3*(3*e^(5*x) + 8*e^(3*x) - 3*e^x)/(e^(2*x) + 1)^3 + arctan(e^x)

Mupad [B]

time = 0.09, size = 61, normalized size = 2.54

$$\operatorname{atan}(e^x) + \frac{e^x}{e^{2x} + 1} - \frac{8e^x}{3(3e^{2x} + 3e^{4x} + e^{6x} + 1)} + \frac{2e^x}{3(2e^{2x} + e^{4x} + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(cosh(x)^5*(tanh(x) + 1)),x)`

[Out] `atan(exp(x)) + exp(x)/(exp(2*x) + 1) - (8*exp(x))/(3*(3*exp(2*x) + 3*exp(4*x) + exp(6*x) + 1)) + (2*exp(x))/(3*(2*exp(2*x) + exp(4*x) + 1))`

3.100 $\int \frac{\operatorname{sech}^6(x)}{1+\tanh(x)} dx$

Optimal. Leaf size=25

$$-\frac{2}{3}(1 - \tanh(x))^3 + \frac{1}{4}(1 - \tanh(x))^4$$

[Out] $-2/3*(1-\tanh(x))^3+1/4*(1-\tanh(x))^4$

Rubi [A]

time = 0.03, antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {3568, 45}

$$\frac{1}{4}(1 - \tanh(x))^4 - \frac{2}{3}(1 - \tanh(x))^3$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sech}[x]^6/(1 + \text{Tanh}[x]), x]$

[Out] $(-2*(1 - \text{Tanh}[x])^3)/3 + (1 - \text{Tanh}[x])^4/4$

Rule 45

$\text{Int}[(a_. + (b_.)(x_.))^{(m_.)}((c_.) + (d_.)(x_.))^{(n_.)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x], x] /; \text{FreeQ}\{a, b, c, d, n\}, x \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ (!\text{IntegerQ}[n] \ || \ (\text{EqQ}[c, 0] \ \&\& \ \text{LeQ}[7*m + 4*n + 4, 0]) \ || \ \text{LtQ}[9*m + 5*(n + 1), 0]) \ || \ \text{GtQ}[m + n + 2, 0])$

Rule 3568

$\text{Int}[\text{sec}[(e_.) + (f_.)(x_.)]^{(m_.)}((a_.) + (b_.)\text{tan}[(e_.) + (f_.)(x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[1/(a^{(m-2)}*b*f), \text{Subst}[\text{Int}[(a-x)^{(m/2-1)}*(a+x)^{(n+m/2-1)}, x], x, b*\text{Tan}[e+f*x]], x] /; \text{FreeQ}\{a, b, e, f, n\}, x \ \&\& \ \text{EqQ}[a^2 + b^2, 0] \ \&\& \ \text{IntegerQ}[m/2]$

Rubi steps

$$\begin{aligned} \int \frac{\operatorname{sech}^6(x)}{1+\tanh(x)} dx &= \text{Subst}\left(\int (1-x)^2(1+x) dx, x, \tanh(x)\right) \\ &= \text{Subst}\left(\int (2(1-x)^2 - (1-x)^3) dx, x, \tanh(x)\right) \\ &= -\frac{2}{3}(1 - \tanh(x))^3 + \frac{1}{4}(1 - \tanh(x))^4 \end{aligned}$$

Mathematica [A]

time = 0.02, size = 20, normalized size = 0.80

$$\frac{1}{12} \operatorname{sech}^4(x)(3 + 4 \sinh(2x) + \sinh(4x))$$

Antiderivative was successfully verified.

[In] Integrate[Sech[x]^6/(1 + Tanh[x]), x]

[Out] (Sech[x]^4*(3 + 4*Sinh[2*x] + Sinh[4*x]))/12

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 55 vs. 2(21) = 42.

time = 0.55, size = 56, normalized size = 2.24

method	result	size
risch	$-\frac{4(4e^{2x}+1)}{3(1+e^{2x})^4}$	19
default	$-\frac{2\left(-\left(\tanh^7\left(\frac{x}{2}\right)\right)+\tanh^6\left(\frac{x}{2}\right)-\frac{5\left(\tanh^5\left(\frac{x}{2}\right)\right)}{3}-\frac{5\left(\tanh^3\left(\frac{x}{2}\right)\right)}{3}+\tanh^2\left(\frac{x}{2}\right)-\tanh\left(\frac{x}{2}\right)\right)}{\left(\tanh^2\left(\frac{x}{2}\right)+1\right)^4}$	56

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sech(x)^6/(1+tanh(x)), x, method=_RETURNVERBOSE)

[Out] $-2*(-\tanh(1/2*x)^7+\tanh(1/2*x)^6-5/3*\tanh(1/2*x)^5-5/3*\tanh(1/2*x)^3+\tanh(1/2*x)^2-\tanh(1/2*x))/(\tanh(1/2*x)^2+1)^4$

Maxima [B] Leaf count of result is larger than twice the leaf count of optimal. 93 vs. 2(17) = 34.

time = 0.26, size = 93, normalized size = 3.72

$$\frac{16e^{-2x}}{3(4e^{-2x} + 6e^{-4x} + 4e^{-6x} + e^{-8x} + 1)} + \frac{8e^{-4x}}{4e^{-2x} + 6e^{-4x} + 4e^{-6x} + e^{-8x} + 1} + \frac{4}{3(4e^{-2x} + 6e^{-4x} + 4e^{-6x} + e^{-8x} + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)^6/(1+tanh(x)), x, algorithm="maxima")

[Out] $16/3*e^{(-2*x)}/(4*e^{(-2*x)} + 6*e^{(-4*x)} + 4*e^{(-6*x)} + e^{(-8*x)} + 1) + 8*e^{(-4*x)}/(4*e^{(-2*x)} + 6*e^{(-4*x)} + 4*e^{(-6*x)} + e^{(-8*x)} + 1) + 4/3/(4*e^{(-2*x)} + 6*e^{(-4*x)} + 4*e^{(-6*x)} + e^{(-8*x)} + 1)$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 140 vs. 2(17) = 34.

time = 0.33, size = 140, normalized size = 5.60

$\frac{4(5 \cosh(x) + 3 \sinh(x))}{3(\cosh(x)^7 + 7 \cosh(x) \sinh(x) \cosh(x)^6 + \sinh(x)^2 + 21 \cosh(x)^2 + 4) \sinh(x)^5 + 4 \cosh(x)^5 + 5(7 \cosh(x)^2 + 4 \cosh(x)) \sinh(x)^4 + (85 \cosh(x)^4 + 40 \cosh(x)^2 + 6) \sinh(x)^3 + 6 \cosh(x)^3 + (21 \cosh(x)^2 + 40 \cosh(x) + 18 \cosh(x) \sinh(x)^2 + 7 \cosh(x)^2 + 20 \cosh(x) + 18 \cosh(x)^2 + 3) \sinh(x) + 5 \cosh(x)}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sech(x)^6/(1+tanh(x)),x, algorithm="fricas")`

[Out]
$$-4/3*(5*\cosh(x) + 3*\sinh(x))/(\cosh(x)^7 + 7*\cosh(x)*\sinh(x)^6 + \sinh(x)^7 + (21*\cosh(x)^2 + 4)*\sinh(x)^5 + 4*\cosh(x)^5 + 5*(7*\cosh(x)^3 + 4*\cosh(x))*\sinh(x)^4 + (35*\cosh(x)^4 + 40*\cosh(x)^2 + 6)*\sinh(x)^3 + 6*\cosh(x)^3 + (21*\cosh(x)^5 + 40*\cosh(x)^3 + 18*\cosh(x))*\sinh(x)^2 + (7*\cosh(x)^6 + 20*\cosh(x))^4 + 18*\cosh(x)^2 + 3)*\sinh(x) + 5*\cosh(x))$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{sech}^6(x)}{\tanh(x) + 1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sech(x)**6/(1+tanh(x)),x)`

[Out] `Integral(sech(x)**6/(tanh(x) + 1), x)`

Giac [A]

time = 0.40, size = 18, normalized size = 0.72

$$-\frac{4(4e^{(2x)} + 1)}{3(e^{(2x)} + 1)^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sech(x)^6/(1+tanh(x)),x, algorithm="giac")`

[Out] `-4/3*(4*e^(2*x) + 1)/(e^(2*x) + 1)^4`

Mupad [B]

time = 1.04, size = 18, normalized size = 0.72

$$-\frac{4(4e^{2x} + 1)}{3(e^{2x} + 1)^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(cosh(x)^6*(tanh(x) + 1)),x)`

[Out] `-(4*(4*exp(2*x) + 1))/(3*(exp(2*x) + 1)^4)`

3.101 $\int \frac{\operatorname{sech}^7(x)}{1+\tanh(x)} dx$

Optimal. Leaf size=34

$$\frac{3}{8}\operatorname{ArcTan}(\sinh(x)) + \frac{\operatorname{sech}^5(x)}{5} + \frac{3}{8}\operatorname{sech}(x)\tanh(x) + \frac{1}{4}\operatorname{sech}^3(x)\tanh(x)$$

[Out] 3/8*arctan(sinh(x))+1/5*sech(x)^5+3/8*sech(x)*tanh(x)+1/4*sech(x)^3*tanh(x)

Rubi [A]

time = 0.03, antiderivative size = 34, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3582, 3853, 3855}

$$\frac{3}{8}\operatorname{ArcTan}(\sinh(x)) + \frac{\operatorname{sech}^5(x)}{5} + \frac{1}{4}\tanh(x)\operatorname{sech}^3(x) + \frac{3}{8}\tanh(x)\operatorname{sech}(x)$$

Antiderivative was successfully verified.

[In] Int[Sech[x]^7/(1 + Tanh[x]), x]

[Out] (3*ArcTan[Sinh[x]])/8 + Sech[x]^5/5 + (3*Sech[x]*Tanh[x])/8 + (Sech[x]^3*Tanh[x])/4

Rule 3582

```
Int[((d_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> Simp[d^2*(d*Sec[e + f*x])^(m - 2)*((a + b*Tan[e + f*x])^(n + 1)/(b*f*(m + n - 1))), x] + Dist[d^2*((m - 2)/(a*(m + n - 1))), Int[(d*Sec[e + f*x])^(m - 2)*(a + b*Tan[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d, e, f}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0] && GtQ[m, 1] && !ILtQ[m + n, 0] && NeQ[m + n - 1, 0] && IntegersQ[2*m, 2*n]
```

Rule 3853

```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Dist[b^2*((n - 2)/(n - 1)), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]
```

Rule 3855

```
Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned}
\int \frac{\operatorname{sech}^7(x)}{1 + \tanh(x)} dx &= \frac{\operatorname{sech}^5(x)}{5} + \int \operatorname{sech}^5(x) dx \\
&= \frac{\operatorname{sech}^5(x)}{5} + \frac{1}{4} \operatorname{sech}^3(x) \tanh(x) + \frac{3}{4} \int \operatorname{sech}^3(x) dx \\
&= \frac{\operatorname{sech}^5(x)}{5} + \frac{3}{8} \operatorname{sech}(x) \tanh(x) + \frac{1}{4} \operatorname{sech}^3(x) \tanh(x) + \frac{3}{8} \int \operatorname{sech}(x) dx \\
&= \frac{3}{8} \tan^{-1}(\sinh(x)) + \frac{\operatorname{sech}^5(x)}{5} + \frac{3}{8} \operatorname{sech}(x) \tanh(x) + \frac{1}{4} \operatorname{sech}^3(x) \tanh(x)
\end{aligned}$$

Mathematica [A]

time = 0.03, size = 34, normalized size = 1.00

$$\frac{1}{40} \left(30 \operatorname{ArcTan} \left(\tanh \left(\frac{x}{2} \right) \right) + 8 \operatorname{sech}^5(x) + 15 \operatorname{sech}(x) \tanh(x) + 10 \operatorname{sech}^3(x) \tanh(x) \right)$$

Antiderivative was successfully verified.

[In] Integrate[Sech[x]^7/(1 + Tanh[x]), x]

[Out] (30*ArcTan[Tanh[x/2]] + 8*Sech[x]^5 + 15*Sech[x]*Tanh[x] + 10*Sech[x]^3*Tanh[x])/40

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 66 vs. 2(26) = 52.

time = 0.57, size = 67, normalized size = 1.97

method	result	size
risch	$\frac{e^x (15 e^{8x} + 70 e^{6x} + 128 e^{4x} - 70 e^{2x} - 15)}{20(1+e^{2x})^5} + \frac{3i \ln(e^x + i)}{8} - \frac{3i \ln(e^x - i)}{8}$	58
default	$-\frac{5(\tanh^9(\frac{x}{2}))}{4} + 2(\tanh^8(\frac{x}{2})) - \frac{(\tanh^7(\frac{x}{2}))}{2} + 4(\tanh^4(\frac{x}{2})) + \frac{(\tanh^3(\frac{x}{2}))}{2} + \frac{5 \tanh(\frac{x}{2})}{4} + \frac{2}{5} + \frac{3 \arctan(\tanh(\frac{x}{2}))}{4}$	67

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sech(x)^7/(1+tanh(x)), x, method=_RETURNVERBOSE)

[Out] 2*(-5/8*tanh(1/2*x)^9+tanh(1/2*x)^8-1/4*tanh(1/2*x)^7+2*tanh(1/2*x)^4+1/4*tanh(1/2*x)^3+5/8*tanh(1/2*x)+1/5)/(tanh(1/2*x)^2+1)^5+3/4*arctan(tanh(1/2*x))

Maxima [B] Leaf count of result is larger than twice the leaf count of optimal. 73 vs. 2(26) = 52.

time = 0.46, size = 73, normalized size = 2.15

$$\frac{15 e^{(-x)} + 70 e^{(-3x)} + 128 e^{(-5x)} - 70 e^{(-7x)} - 15 e^{(-9x)}}{20(5 e^{(-2x)} + 10 e^{(-4x)} + 10 e^{(-6x)} + 5 e^{(-8x)} + e^{(-10x)} + 1)} - \frac{3}{4} \arctan(e^{(-x)})$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)^7/(1+tanh(x)),x, algorithm="maxima")

[Out] $\frac{1}{20} \cdot (15e^{-x} + 70e^{-3x} + 128e^{-5x} - 70e^{-7x} - 15e^{-9x}) / (5e^{-2x} + 10e^{-4x} + 10e^{-6x} + 5e^{-8x} + e^{-10x} + 1) - \frac{3}{4} \arctan(e^{-x})$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 670 vs. $2(26) = 52$.

time = 0.36, size = 670, normalized size = 19.71

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)^7/(1+tanh(x)),x, algorithm="fricas")

[Out] $\frac{1}{20} \cdot (15 \cosh(x)^9 + 135 \cosh(x) \sinh(x)^8 + 15 \sinh(x)^9 + 10 \cdot (54 \cosh(x)^2 + 7) \sinh(x)^7 + 70 \cosh(x)^7 + 70 \cdot (18 \cosh(x)^3 + 7 \cosh(x)) \sinh(x)^6 + 2 \cdot (945 \cosh(x)^4 + 735 \cosh(x)^2 + 64) \sinh(x)^5 + 128 \cosh(x)^5 + 10 \cdot (189 \cosh(x)^5 + 245 \cosh(x)^3 + 64 \cosh(x)) \sinh(x)^4 + 10 \cdot (126 \cosh(x)^6 + 245 \cosh(x)^4 + 128 \cosh(x)^2 - 7) \sinh(x)^3 - 70 \cosh(x)^3 + 10 \cdot (54 \cosh(x)^7 + 147 \cosh(x)^5 + 128 \cosh(x)^3 - 21 \cosh(x)) \sinh(x)^2 + 15 \cdot (\cosh(x)^{10} + 10 \cosh(x) \sinh(x)^9 + \sinh(x)^{10} + 5 \cdot (9 \cosh(x)^2 + 1) \sinh(x)^8 + 5 \cosh(x)^8 + 40 \cdot (3 \cosh(x)^3 + \cosh(x)) \sinh(x)^7 + 10 \cdot (21 \cosh(x)^4 + 14 \cosh(x)^2 + 1) \sinh(x)^6 + 10 \cosh(x)^6 + 4 \cdot (63 \cosh(x)^5 + 70 \cosh(x)^3 + 15 \cosh(x)) \sinh(x)^5 + 10 \cdot (21 \cosh(x)^6 + 35 \cosh(x)^4 + 15 \cosh(x)^2 + 1) \sinh(x)^4 + 10 \cosh(x)^4 + 40 \cdot (3 \cosh(x)^7 + 7 \cosh(x)^5 + 5 \cosh(x)^3 + \cosh(x)) \sinh(x)^3 + 5 \cdot (9 \cosh(x)^8 + 28 \cosh(x)^6 + 30 \cosh(x)^4 + 12 \cosh(x)^2 + 1) \sinh(x)^2 + 5 \cosh(x)^2 + 10 \cdot (\cosh(x)^9 + 4 \cosh(x)^7 + 6 \cosh(x)^5 + 4 \cosh(x)^3 + \cosh(x)) \sinh(x) + 1) \arctan(\cosh(x) + \sinh(x)) + 5 \cdot (27 \cosh(x)^8 + 98 \cosh(x)^6 + 128 \cosh(x)^4 - 42 \cosh(x)^2 - 3) \sinh(x) - 15 \cosh(x)) / (\cosh(x)^{10} + 10 \cosh(x) \sinh(x)^9 + \sinh(x)^{10} + 5 \cdot (9 \cosh(x)^2 + 1) \sinh(x)^8 + 5 \cosh(x)^8 + 40 \cdot (3 \cosh(x)^3 + \cosh(x)) \sinh(x)^7 + 10 \cdot (21 \cosh(x)^4 + 14 \cosh(x)^2 + 1) \sinh(x)^6 + 10 \cosh(x)^6 + 4 \cdot (63 \cosh(x)^5 + 70 \cosh(x)^3 + 15 \cosh(x)) \sinh(x)^5 + 10 \cdot (21 \cosh(x)^6 + 35 \cosh(x)^4 + 15 \cosh(x)^2 + 1) \sinh(x)^4 + 10 \cosh(x)^4 + 40 \cdot (3 \cosh(x)^7 + 7 \cosh(x)^5 + 5 \cosh(x)^3 + \cosh(x)) \sinh(x)^3 + 5 \cdot (9 \cosh(x)^8 + 28 \cosh(x)^6 + 30 \cosh(x)^4 + 12 \cosh(x)^2 + 1) \sinh(x)^2 + 5 \cosh(x)^2 + 10 \cdot (\cosh(x)^9 + 4 \cosh(x)^7 + 6 \cosh(x)^5 + 4 \cosh(x)^3 + \cosh(x)) \sinh(x) + 1)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{sech}^7(x)}{\tanh(x) + 1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)**7/(1+tanh(x)),x)

[Out] Integral(sech(x)**7/(tanh(x) + 1), x)

Giac [A]

time = 0.40, size = 45, normalized size = 1.32

$$\frac{15 e^{(9x)} + 70 e^{(7x)} + 128 e^{(5x)} - 70 e^{(3x)} - 15 e^x}{20 (e^{(2x)} + 1)^5} + \frac{3}{4} \arctan(e^x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)^7/(1+tanh(x)),x, algorithm="giac")

[Out] 1/20*(15*e^(9*x) + 70*e^(7*x) + 128*e^(5*x) - 70*e^(3*x) - 15*e^x)/(e^(2*x) + 1)^5 + 3/4*arctan(e^x)

Mupad [B]

time = 0.08, size = 137, normalized size = 4.03

$$\frac{3 \operatorname{atan}(e^x)}{4} - \frac{32 e^{3x}}{5 (5 e^{2x} + 10 e^{4x} + 10 e^{6x} + 5 e^{8x} + e^{10x} + 1)} - \frac{12 e^x}{5 (4 e^{2x} + 6 e^{4x} + 4 e^{6x} + e^{8x} + 1)} + \frac{3 e^x}{4 (e^{2x} + 1)} + \frac{2 e^x}{5 (3 e^{2x} + 3 e^{4x} + e^{6x} + 1)} + \frac{e^x}{2 (2 e^{2x} + e^{4x} + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cosh(x)^7*(tanh(x) + 1)),x)

[Out] (3*atan(exp(x)))/4 - (32*exp(3*x))/(5*(5*exp(2*x) + 10*exp(4*x) + 10*exp(6*x) + 5*exp(8*x) + exp(10*x) + 1)) - (12*exp(x))/(5*(4*exp(2*x) + 6*exp(4*x) + 4*exp(6*x) + exp(8*x) + 1)) + (3*exp(x))/(4*(exp(2*x) + 1)) + (2*exp(x))/(5*(3*exp(2*x) + 3*exp(4*x) + exp(6*x) + 1)) + exp(x)/(2*(2*exp(2*x) + exp(4*x) + 1))

3.102 $\int \frac{\operatorname{sech}^8(x)}{a+b \tanh(x)} dx$

Optimal. Leaf size=140

$$-\frac{(a^2 - b^2)^3 \log(a + b \tanh(x))}{b^7} + \frac{a(a^4 - 3a^2b^2 + 3b^4) \tanh(x)}{b^6} - \frac{(a^4 - 3a^2b^2 + 3b^4) \tanh^2(x)}{2b^5} + \frac{a(a^2 - 3b^2) \tanh^3(x)}{3b^4}$$

[Out] $-(a^2 - b^2)^3 \ln(a + b \tanh(x)) / b^7 + a(a^4 - 3a^2b^2 + 3b^4) \tanh(x) / b^6 - 1/2(a^4 - 3a^2b^2 + 3b^4) \tanh^2(x) / b^5 + 1/3a(a^2 - 3b^2) \tanh^3(x) / b^4 - 1/4(a^2 - 3b^2) \tanh^4(x) / b^3 + 1/5a \tanh^5(x) / b^2 - 1/6 \tanh^6(x) / b$

Rubi [A]

time = 0.12, antiderivative size = 140, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$,

Rules used = {3587, 711}

$$-\frac{(a^2 - b^2)^3 \log(a + b \tanh(x))}{b^7} + \frac{a(a^2 - 3b^2) \tanh^3(x)}{3b^4} - \frac{(a^2 - 3b^2) \tanh^4(x)}{4b^3} + \frac{a(a^4 - 3a^2b^2 + 3b^4) \tanh(x)}{b^6} - \frac{(a^4 - 3a^2b^2 + 3b^4) \tanh^2(x)}{2b^5} + \frac{a \tanh^5(x)}{5b^2} - \frac{\tanh^6(x)}{6b}$$

Antiderivative was successfully verified.

[In] `Int[Sech[x]^8/(a + b*Tanh[x]), x]`

[Out] $-\frac{((a^2 - b^2)^3 \operatorname{Log}[a + b \operatorname{Tanh}[x]])}{b^7} + \frac{(a(a^4 - 3a^2b^2 + 3b^4) \operatorname{Tanh}[x])}{b^6} - \frac{((a^4 - 3a^2b^2 + 3b^4) \operatorname{Tanh}[x]^2)}{(2b^5)} + \frac{(a(a^2 - 3b^2) \operatorname{Tanh}[x]^3)}{(3b^4)} - \frac{((a^2 - 3b^2) \operatorname{Tanh}[x]^4)}{(4b^3)} + \frac{(a \operatorname{Tanh}[x]^5)}{(5b^2)} - \frac{\operatorname{Tanh}[x]^6}{(6b)}$

Rule 711

`Int[((d_) + (e_)*(x_))^(m_)*((a_) + (c_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(a + c*x^2)^p, x], x] /; FreeQ[{a, c, d, e, m}, x] && NeQ[c*d^2 + a*e^2, 0] && IGtQ[p, 0]`

Rule 3587

`Int[sec[(e_) + (f_)*(x_)]^(m_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Dist[1/(b*f), Subst[Int[(a + x)^n*(1 + x^2/b^2)^(m/2 - 1), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, n}, x] && NeQ[a^2 + b^2, 0] && IntegerQ[m/2]`

Rubi steps

$$\begin{aligned}
 \int \frac{\operatorname{sech}^8(x)}{a + b \tanh(x)} dx &= \frac{\operatorname{Subst}\left(\int \frac{\left(1 - \frac{x^2}{b^2}\right)^3}{a+x} dx, x, b \tanh(x)\right)}{b} \\
 &= \frac{\operatorname{Subst}\left(\int \left(\frac{a^5 - 3a^3b^2 + 3ab^4}{b^6} - \frac{(a^4 - 3a^2b^2 + 3b^4)x}{b^6} + \frac{a(a^2 - 3b^2)x^2}{b^6} + \frac{(-a^2 + 3b^2)x^3}{b^6} + \frac{ax^4}{b^6} - \frac{x^5}{b^6} + \frac{(-a^6 + 3a^4b^2 - 3a^2b^4 + b^6)x^6}{b^6}\right) dx, x, b \tanh(x)\right)}{b} \\
 &= -\frac{(a^2 - b^2)^3 \log(a + b \tanh(x))}{b^7} + \frac{a(a^4 - 3a^2b^2 + 3b^4) \tanh(x)}{b^6} - \frac{(a^4 - 3a^2b^2 + 3b^4)}{2b^5}
 \end{aligned}$$

Mathematica [A]

time = 0.44, size = 137, normalized size = 0.98

$$\frac{60(a^2 - b^2)^3 (\log(\cosh(x)) - \log(a \cosh(x) + b \sinh(x))) + 10b^6 \operatorname{sech}^6(x) + 4ab(15a^4 - 40a^2b^2 + 33b^4) \tanh(x) + 3b^4 \operatorname{sech}^4(x) (-5a^2 + 5b^2 + 4ab \tanh(x)) + 2b^2 \operatorname{sech}^2(x) (15(a^2 - b^2)^2 - 2ab(5a^2 - 9b^2) \tanh(x))}{60b^7}$$

Antiderivative was successfully verified.

[In] Integrate[Sech[x]^8/(a + b*Tanh[x]), x]

[Out] $(60(a^2 - b^2)^3 (\log(\cosh(x)) - \log(a \cosh(x) + b \sinh(x))) + 10b^6 \operatorname{sech}^6(x) + 4ab(15a^4 - 40a^2b^2 + 33b^4) \tanh(x) + 3b^4 \operatorname{sech}^4(x) (-5a^2 + 5b^2 + 4ab \tanh(x)) + 2b^2 \operatorname{sech}^2(x) (15(a^2 - b^2)^2 - 2ab(5a^2 - 9b^2) \tanh(x)))/60b^7$

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 412 vs. $2(130) = 260$.

time = 0.76, size = 413, normalized size = 2.95

method	result
default	$-\frac{(a^6 - 3a^4b^2 + 3a^2b^4 - b^6) \ln(a(\tanh^2(\frac{x}{2})) + 2b \tanh(\frac{x}{2}) + a)}{b^7} + \frac{2((a^5b - 3a^3b^3 + 3ab^5) \tanh^{11}(\frac{x}{2}) + (-a^4b^2 + 3a^2b^4 - 3b^6) \tanh^{10}(\frac{x}{2}))}{b^6}$
risch	$-\frac{2(-40a^3b^2 + 390ab^4e^{4x} - 210a^3b^2e^{2x} + 183ab^4e^{2x} + 15a^5 + 150a^2b^3e^{4x} - 15a^4be^{2x} + 30a^2b^3e^{2x} + 33ab^4 + 105ab^4e^{8x} - 400a^3b^2e^{6x} + 300a^2b^3e^{4x})}{b^6}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sech(x)^8/(a+b*tanh(x)), x, method=_RETURNVERBOSE)

[Out] $-(a^6 - 3a^4b^2 + 3a^2b^4 - b^6)/b^7 * \ln(a * \tanh(1/2*x)^2 + 2*b * \tanh(1/2*x) + a) + 2/b^7 * (((a^5*b - 3*a^3*b^3 + 3*a*b^5) * \tanh(1/2*x)^11 + (-a^4*b^2 + 3*a^2*b^4 - 3*b^6) * \tanh(1/2*x)^10 + (5*a^5*b - 41/3*a^3*b^3 + 11*a*b^5) * \tanh(1/2*x)^9 + (-4*a^4*b^2 + 10*a^2*b^4 - 6*b^6) * \tanh(1/2*x)^8 + (10*a^5*b - 26*a^3*b^3 + 106/5*a*b^5) * \tanh(1/2*x)^7 + (-6*a^4*b^2 + 14*a^2*b^4 - 34/3*b^6) * \tanh(1/2*x)^6 + (10*a^5*b - 26*a^3*b^3 + 106/5$

$$*a*b^5)*\tanh(1/2*x)^5+(-4*a^4*b^2+10*a^2*b^4-6*b^6)*\tanh(1/2*x)^4+(5*a^5*b-41/3*a^3*b^3+11*a*b^5)*\tanh(1/2*x)^3+(-a^4*b^2+3*a^2*b^4-3*b^6)*\tanh(1/2*x)^2+(a^5*b-3*a^3*b^3+3*a*b^5)*\tanh(1/2*x))/(\tanh(1/2*x)^2+1)^6+1/2*(a^6-3*a^4*b^2+3*a^2*b^4-b^6)*\ln(\tanh(1/2*x)^2+1))$$

Maxima [B] Leaf count of result is larger than twice the leaf count of optimal. 386 vs. $2(130) = 260$.

time = 0.49, size = 386, normalized size = 2.76

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)^8/(a+b*tanh(x)),x, algorithm="maxima")

[Out] $2/15*(15*a^5 - 40*a^3*b^2 + 33*a*b^4 + 3*(25*a^5 + 5*a^4*b - 70*a^3*b^2 - 10*a^2*b^3 + 61*a*b^4 + 5*b^5)*e^{-2*x} + 30*(5*a^5 + 2*a^4*b - 14*a^3*b^2 - 5*a^2*b^3 + 13*a*b^4 + 3*b^5)*e^{-4*x} + 10*(15*a^5 + 9*a^4*b - 40*a^3*b^2 - 24*a^2*b^3 + 33*a*b^4 + 23*b^5)*e^{-6*x} + 15*(5*a^5 + 4*a^4*b - 12*a^3*b^2 - 10*a^2*b^3 + 7*a*b^4 + 6*b^5)*e^{-8*x} + 15*(a^5 + a^4*b - 2*a^3*b^2 - 2*a^2*b^3 + a*b^4 + b^5)*e^{-10*x})/(6*b^6*e^{-2*x} + 15*b^6*e^{-4*x} + 20*b^6*e^{-6*x} + 15*b^6*e^{-8*x} + 6*b^6*e^{-10*x} + b^6*e^{-12*x} + b^6) - (a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\log(-(a - b)*e^{-2*x} - a - b)/b^7 + (a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\log(e^{-2*x} + 1)/b^7$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 5275 vs. $2(130) = 260$.

time = 0.41, size = 5275, normalized size = 37.68

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)^8/(a+b*tanh(x)),x, algorithm="fricas")

[Out] $-1/15*(30*(a^5*b - a^4*b^2 - 2*a^3*b^3 + 2*a^2*b^4 + a*b^5 - b^6)*\cosh(x)^10 + 300*(a^5*b - a^4*b^2 - 2*a^3*b^3 + 2*a^2*b^4 + a*b^5 - b^6)*\cosh(x)*\sinh(x)^9 + 30*(a^5*b - a^4*b^2 - 2*a^3*b^3 + 2*a^2*b^4 + a*b^5 - b^6)*\sinh(x)^10 + 30*(5*a^5*b - 4*a^4*b^2 - 12*a^3*b^3 + 10*a^2*b^4 + 7*a*b^5 - 6*b^6)*\cosh(x)^8 + 30*(5*a^5*b - 4*a^4*b^2 - 12*a^3*b^3 + 10*a^2*b^4 + 7*a*b^5 - 6*b^6 + 45*(a^5*b - a^4*b^2 - 2*a^3*b^3 + 2*a^2*b^4 + a*b^5 - b^6)*\cosh(x)^2)*\sinh(x)^8 + 240*(15*(a^5*b - a^4*b^2 - 2*a^3*b^3 + 2*a^2*b^4 + a*b^5 - b^6)*\cosh(x)^3 + (5*a^5*b - 4*a^4*b^2 - 12*a^3*b^3 + 10*a^2*b^4 + 7*a*b^5 - 6*b^6)*\cosh(x))*\sinh(x)^7 + 20*(15*a^5*b - 9*a^4*b^2 - 40*a^3*b^3 + 24*a^2*b^4 + 33*a*b^5 - 23*b^6)*\cosh(x)^6 + 20*(15*a^5*b - 9*a^4*b^2 - 40*a^3*b^3 + 24*a^2*b^4 + 33*a*b^5 - 23*b^6 + 315*(a^5*b - a^4*b^2 - 2*a^3*b^3 + 2*a^2*b^4 + a*b^5 - b^6)*\cosh(x)^4 + 42*(5*a^5*b - 4*a^4*b^2 - 12*a^3*b^3 + 10*a^2*b^4 + 7*a*b^5 - 6*b^6)*\cosh(x)^2)*\sinh(x)^6 + 30*a^5*b - 80*a^3*b^3 + 66*$

$$\begin{aligned}
& a*b^5 + 120*(63*(a^5*b - a^4*b^2 - 2*a^3*b^3 + 2*a^2*b^4 + a*b^5 - b^6)*\cos \\
& h(x)^5 + 14*(5*a^5*b - 4*a^4*b^2 - 12*a^3*b^3 + 10*a^2*b^4 + 7*a*b^5 - 6*b^6)*\cosh(x)^3 + (15*a^5*b - 9*a^4*b^2 - 40*a^3*b^3 + 24*a^2*b^4 + 33*a*b^5 - \\
& 23*b^6)*\cosh(x))*\sinh(x)^5 + 60*(5*a^5*b - 2*a^4*b^2 - 14*a^3*b^3 + 5*a^2* \\
& b^4 + 13*a*b^5 - 3*b^6)*\cosh(x)^4 + 60*(105*(a^5*b - a^4*b^2 - 2*a^3*b^3 + \\
& 2*a^2*b^4 + a*b^5 - b^6)*\cosh(x)^6 + 5*a^5*b - 2*a^4*b^2 - 14*a^3*b^3 + 5*a \\
& ^2*b^4 + 13*a*b^5 - 3*b^6 + 35*(5*a^5*b - 4*a^4*b^2 - 12*a^3*b^3 + 10*a^2*b \\
& ^4 + 7*a*b^5 - 6*b^6)*\cosh(x)^4 + 5*(15*a^5*b - 9*a^4*b^2 - 40*a^3*b^3 + 24 \\
& *a^2*b^4 + 33*a*b^5 - 23*b^6)*\cosh(x)^2)*\sinh(x)^4 + 80*(45*(a^5*b - a^4*b^ \\
& 2 - 2*a^3*b^3 + 2*a^2*b^4 + a*b^5 - b^6)*\cosh(x)^7 + 21*(5*a^5*b - 4*a^4*b^ \\
& 2 - 12*a^3*b^3 + 10*a^2*b^4 + 7*a*b^5 - 6*b^6)*\cosh(x)^5 + 5*(15*a^5*b - 9* \\
& a^4*b^2 - 40*a^3*b^3 + 24*a^2*b^4 + 33*a*b^5 - 23*b^6)*\cosh(x)^3 + 3*(5*a^5 \\
& *b - 2*a^4*b^2 - 14*a^3*b^3 + 5*a^2*b^4 + 13*a*b^5 - 3*b^6)*\cosh(x))*\sinh(x \\
&)^3 + 6*(25*a^5*b - 5*a^4*b^2 - 70*a^3*b^3 + 10*a^2*b^4 + 61*a*b^5 - 5*b^6) \\
& *\cosh(x)^2 + 6*(225*(a^5*b - a^4*b^2 - 2*a^3*b^3 + 2*a^2*b^4 + a*b^5 - b^6) \\
& *\cosh(x)^8 + 140*(5*a^5*b - 4*a^4*b^2 - 12*a^3*b^3 + 10*a^2*b^4 + 7*a*b^5 - \\
& 6*b^6)*\cosh(x)^6 + 25*a^5*b - 5*a^4*b^2 - 70*a^3*b^3 + 10*a^2*b^4 + 61*a*b \\
& ^5 - 5*b^6 + 50*(15*a^5*b - 9*a^4*b^2 - 40*a^3*b^3 + 24*a^2*b^4 + 33*a*b^5 \\
& - 23*b^6)*\cosh(x)^4 + 60*(5*a^5*b - 2*a^4*b^2 - 14*a^3*b^3 + 5*a^2*b^4 + 13 \\
& *a*b^5 - 3*b^6)*\cosh(x)^2)*\sinh(x)^2 + 15*((a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b \\
& ^6)*\cosh(x)^12 + 12*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\cosh(x))*\sinh(x)^11 \\
& + (a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\sinh(x)^12 + 6*(a^6 - 3*a^4*b^2 + 3*a \\
& ^2*b^4 - b^6)*\cosh(x)^10 + 6*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6 + 11*(a^6 - \\
& 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\cosh(x)^2)*\sinh(x)^10 + 20*(11*(a^6 - 3*a^4*b \\
& ^2 + 3*a^2*b^4 - b^6)*\cosh(x)^3 + 3*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\cos \\
& h(x))*\sinh(x)^9 + 15*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\cosh(x)^8 + 15*(a^ \\
& 6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6 + 33*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\co \\
& sh(x)^4 + 18*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\cosh(x)^2)*\sinh(x)^8 + 24* \\
& (33*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\cosh(x)^5 + 30*(a^6 - 3*a^4*b^2 + 3 \\
& *a^2*b^4 - b^6)*\cosh(x)^3 + 5*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\cosh(x))* \\
& \sinh(x)^7 + 20*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\cosh(x)^6 + 4*(231*(a^6 \\
& - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\cosh(x)^6 + 5*a^6 - 15*a^4*b^2 + 15*a^2*b^4 \\
& - 5*b^6 + 315*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\cosh(x)^4 + 105*(a^6 - 3* \\
& a^4*b^2 + 3*a^2*b^4 - b^6)*\cosh(x)^2)*\sinh(x)^6 + a^6 - 3*a^4*b^2 + 3*a^2*b \\
& ^4 - b^6 + 24*(33*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\cosh(x)^7 + 63*(a^6 - \\
& 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\cosh(x)^5 + 35*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - \\
& b^6)*\cosh(x)^3 + 5*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\cosh(x))*\sinh(x)^5 \\
& + 15*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\cosh(x)^4 + 15*(33*(a^6 - 3*a^4*b^ \\
& 2 + 3*a^2*b^4 - b^6)*\cosh(x)^8 + 84*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\cos \\
& h(x)^6 + a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6 + 70*(a^6 - 3*a^4*b^2 + 3*a^2*b^ \\
& 4 - b^6)*\cosh(x)^4 + 20*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\cosh(x)^2)*\sinh \\
& (x)^4 + 20*(11*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\cosh(x)^9 + 36*(a^6 - 3* \\
& a^4*b^2 + 3*a^2*b^4 - b^6)*\cosh(x)^7 + 42*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^ \\
& 6)*\cosh(x)^5 + 20*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\cosh(x)^3 + 3*(a^6 - \\
& 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\cosh(x))*\sinh(x)^3 + 6*(a^6 - 3*a^4*b^2 + 3*a^
\end{aligned}$$

$2*b^4 - b^6)*\cosh(x)^2 + 6*(11*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\cosh(x)^{10} + 45*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\cosh(x)^8 + 70*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\cosh(x)^6 + a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6 + 50*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\cosh(x)^4 + 15*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\cosh(x)^2)*\sinh(x)^2 + 12*((a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\cosh(x)^{11} + 5*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\cosh(x)^9 + 10*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\cosh(x)^7 + 10*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\cosh(x)^5 + 5*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\cosh(x)^3 + (a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\cosh(x))*\sinh(x))*\log(2*(...$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{sech}^8(x)}{a + b \tanh(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)**8/(a+b*tanh(x)),x)

[Out] Integral(sech(x)**8/(a + b*tanh(x)), x)

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 593 vs. 2(130) = 260.

time = 0.41, size = 593, normalized size = 4.24

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)^8/(a+b*tanh(x)),x, algorithm="giac")

[Out] $-(a^7 + a^6*b - 3*a^5*b^2 - 3*a^4*b^3 + 3*a^3*b^4 + 3*a^2*b^5 - a*b^6 - b^7)*\log(\operatorname{abs}(a*e^{(2*x)} + b*e^{(2*x)} + a - b))/(a*b^7 + b^8) + (a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*\log(e^{(2*x)} + 1)/b^7 - 1/60*(147*a^6*e^{(12*x)} - 441*a^4*b^2*e^{(12*x)} + 441*a^2*b^4*e^{(12*x)} - 147*b^6*e^{(12*x)} + 882*a^6*e^{(10*x)} + 120*a^5*b*e^{(10*x)} - 2766*a^4*b^2*e^{(10*x)} - 240*a^3*b^3*e^{(10*x)} + 2886*a^2*b^4*e^{(10*x)} + 120*a*b^5*e^{(10*x)} - 1002*b^6*e^{(10*x)} + 2205*a^6*e^{(8*x)} + 600*a^5*b*e^{(8*x)} - 7095*a^4*b^2*e^{(8*x)} - 1440*a^3*b^3*e^{(8*x)} + 7815*a^2*b^4*e^{(8*x)} + 840*a*b^5*e^{(8*x)} - 2925*b^6*e^{(8*x)} + 2940*a^6*e^{(6*x)} + 1200*a^5*b*e^{(6*x)} - 9540*a^4*b^2*e^{(6*x)} - 3200*a^3*b^3*e^{(6*x)} + 10740*a^2*b^4*e^{(6*x)} + 2640*a*b^5*e^{(6*x)} - 4780*b^6*e^{(6*x)} + 2205*a^6*e^{(4*x)} + 1200*a^5*b*e^{(4*x)} - 7095*a^4*b^2*e^{(4*x)} - 3360*a^3*b^3*e^{(4*x)} + 7815*a^2*b^4*e^{(4*x)} + 3120*a*b^5*e^{(4*x)} - 2925*b^6*e^{(4*x)} + 882*a^6*e^{(2*x)} + 600*a^5*b*e^{(2*x)} - 2766*a^4*b^2*e^{(2*x)} - 1680*a^3*b^3*e^{(2*x)} + 2886*a^2*b^4*e^{(2*x)} + 1464*a*b^5*e^{(2*x)} - 1002*b^6*e^{(2*x)} + 147*a^6 + 120*a^5*b - 441*a^4*b^2 - 320*a^3*b^3 + 441*a^2*b^4 + 264*a*b^5 - 147*b^6)/(b^7*(e^{(2*x)} + 1)^6)$

Mupad [B]

time = 1.43, size = 301, normalized size = 2.15

$$\frac{\ln(e^{2x} + 1)(a+b)^2(a-b)^2}{b^2} - \frac{32(a-b)}{5b^2(5e^{2x} + 10e^{4x} + 10e^{6x} + 5e^{8x} + e^{10x} + 1)} - \frac{4(a^2 - 4ab + 7b^2)}{b^2(4e^{2x} + 6e^{4x} + 4e^{6x} + e^{8x} + 1)} - \frac{\ln(a-b + a e^{2x} + b e^{4x})(a+b)^2(a-b)^2}{b^2} - \frac{32}{3b(6e^{2x} + 15e^{4x} + 20e^{6x} + 15e^{8x} + 6e^{10x} + e^{12x} + 1)} - \frac{8(a-b)(a^2 - 2ab + b^2)}{3b^2(3e^{2x} + 3e^{4x} + e^{6x} + 1)} - \frac{2(a+b)^2(a-b)(a^2 - 2ab + b^2)}{b^2(e^{2x} + 1)} - \frac{2(a+b)(a-b)(a^2 - 2ab + b^2)}{b^2(2e^{2x} + e^{4x} + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cosh(x)^8*(a + b*tanh(x))),x)

[Out] (log(exp(2*x) + 1)*(a + b)^3*(a - b)^3)/b^7 - (32*(a - 5*b))/(5*b^2*(5*exp(2*x) + 10*exp(4*x) + 10*exp(6*x) + 5*exp(8*x) + exp(10*x) + 1)) - (4*(a^2 - 4*a*b + 7*b^2))/(b^3*(4*exp(2*x) + 6*exp(4*x) + 4*exp(6*x) + exp(8*x) + 1)) - (log(a - b + a*exp(2*x) + b*exp(4*x))*(a + b)^3*(a - b)^3)/b^7 - 32/(3*b*(6*exp(2*x) + 15*exp(4*x) + 20*exp(6*x) + 15*exp(8*x) + 6*exp(10*x) + exp(12*x) + 1)) - (8*(a - b)*(a^2 - 2*a*b + b^2))/(3*b^4*(3*exp(2*x) + 3*exp(4*x) + exp(6*x) + 1)) - (2*(a + b)^2*(a - b)*(a^2 - 2*a*b + b^2))/(b^6*(exp(2*x) + 1)) - (2*(a + b)*(a - b)*(a^2 - 2*a*b + b^2))/(b^5*(2*exp(2*x) + exp(4*x) + 1))

3.103 $\int \frac{\operatorname{sech}^6(x)}{a+b \tanh(x)} dx$

Optimal. Leaf size=83

$$\frac{(a^2 - b^2)^2 \log(a + b \tanh(x))}{b^5} - \frac{a(a^2 - 2b^2) \tanh(x)}{b^4} + \frac{(a^2 - 2b^2) \tanh^2(x)}{2b^3} - \frac{a \tanh^3(x)}{3b^2} + \frac{\tanh^4(x)}{4b}$$

[Out] $(a^2 - b^2)^2 \ln(a + b \tanh(x)) / b^5 - a(a^2 - 2b^2) \tanh(x) / b^4 + 1/2(a^2 - 2b^2) \tanh^2(x) / b^3 - 1/3 a \tanh^3(x) / b^2 + 1/4 \tanh^4(x) / b$

Rubi [A]

time = 0.08, antiderivative size = 83, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {3587, 711}

$$\frac{(a^2 - b^2)^2 \log(a + b \tanh(x))}{b^5} - \frac{a(a^2 - 2b^2) \tanh(x)}{b^4} + \frac{(a^2 - 2b^2) \tanh^2(x)}{2b^3} - \frac{a \tanh^3(x)}{3b^2} + \frac{\tanh^4(x)}{4b}$$

Antiderivative was successfully verified.

[In] `Int[Sech[x]^6/(a + b*Tanh[x]),x]`

[Out] $((a^2 - b^2)^2 \operatorname{Log}[a + b \operatorname{Tanh}[x]]) / b^5 - (a(a^2 - 2b^2) \operatorname{Tanh}[x]) / b^4 + ((a^2 - 2b^2) \operatorname{Tanh}[x]^2) / (2b^3) - (a \operatorname{Tanh}[x]^3) / (3b^2) + \operatorname{Tanh}[x]^4 / (4b)$

Rule 711

`Int[((d_) + (e_.)*(x_))^(m_)*((a_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(a + c*x^2)^p, x], x] /; FreeQ[{a, c, d, e, m}, x] && NeQ[c*d^2 + a*e^2, 0] && IGtQ[p, 0]`

Rule 3587

`Int[sec[(e_.) + (f_.)*(x_)]^(m_)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Dist[1/(b*f), Subst[Int[(a + x)^n*(1 + x^2/b^2)^(m/2 - 1), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, n}, x] && NeQ[a^2 + b^2, 0] && IntegerQ[m/2]`

Rubi steps

$$\int \frac{\operatorname{sech}^6(x)}{a + b \tanh(x)} dx = \frac{\operatorname{Subst}\left(\int \frac{\left(1 - \frac{x^2}{b^2}\right)^2}{a+x} dx, x, b \tanh(x)\right)}{b}$$

$$= \frac{\operatorname{Subst}\left(\int \left(\frac{-a^3+2ab^2}{b^4} - \frac{(-a^2+2b^2)x}{b^4} - \frac{ax^2}{b^4} + \frac{x^3}{b^4} + \frac{(-a^2+b^2)^2}{b^4(a+x)}\right) dx, x, b \tanh(x)\right)}{b}$$

$$= \frac{(a^2 - b^2)^2 \log(a + b \tanh(x))}{b^5} - \frac{a(a^2 - 2b^2) \tanh(x)}{b^4} + \frac{(a^2 - 2b^2) \tanh^2(x)}{2b^3} - \frac{a \tanh^3(x)}{3b^2}$$

Mathematica [A]

time = 0.28, size = 92, normalized size = 1.11

$$\frac{3b^4 \operatorname{sech}^4(x) + \operatorname{sech}^2(x)(-6a^2b^2 + 6b^4 + 4ab^3 \tanh(x)) - 4(3(a^2 - b^2)^2 (\log(\cosh(x)) - \log(a \cosh(x) + b \sinh(x))) + ab(3a^2 - 5b^2) \tanh(x))}{12b^5}$$

Antiderivative was successfully verified.

[In] Integrate[Sech[x]^6/(a + b*Tanh[x]), x]

[Out] (3*b^4*Sech[x]^4 + Sech[x]^2*(-6*a^2*b^2 + 6*b^4 + 4*a*b^3*Tanh[x]) - 4*(3*(a^2 - b^2)^2*(Log[Cosh[x]] - Log[a*Cosh[x] + b*Sinh[x]]) + a*b*(3*a^2 - 5*b^2)*Tanh[x])/(12*b^5)

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 221 vs. 2(77) = 154.

time = 0.70, size = 222, normalized size = 2.67

method	result
default	$\frac{(a^4 - 2a^2b^2 + b^4) \ln(a(\tanh^2(\frac{x}{2})) + 2b \tanh(\frac{x}{2}) + a)}{b^5} - \frac{2 \left((a^3b - 2ab^3) \left(\tanh^7\left(\frac{x}{2}\right) \right) + (-a^2b^2 + 2b^4) \left(\tanh^6\left(\frac{x}{2}\right) \right) + (3a^3b - \frac{14}{3}ab^3) \left(\tanh^5\left(\frac{x}{2}\right) \right) + \dots \right)}{b^5}$
risch	$\frac{2a^3e^{6x} - 2a^2be^{6x} - 2ab^2e^{6x} + 2b^3e^{6x} + 6a^3e^{4x} - 4a^2be^{4x} - 10ab^2e^{4x} + 8b^3e^{4x} + 6a^3e^{2x} - 2a^2be^{2x} - \frac{34ab^2e^{2x}}{3} + 2b^3e^{2x} + 2a^3 - \frac{10ab^2}{3}}{b^4(1+e^{2x})^4} - \dots$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sech(x)^6/(a+b*tanh(x)), x, method=_RETURNVERBOSE)

[Out] (a^4-2*a^2*b^2+b^4)/b^5*ln(a*tanh(1/2*x)^2+2*b*tanh(1/2*x)+a)-2/b^5*((a^3*b-2*a*b^3)*tanh(1/2*x)^7+(-a^2*b^2+2*b^4)*tanh(1/2*x)^6+(3*a^3*b-14/3*a*b^3)*tanh(1/2*x)^5+(-2*a^2*b^2+2*b^4)*tanh(1/2*x)^4+(3*a^3*b-14/3*a*b^3)*tanh(1/2*x)^3+(-a^2*b^2+2*b^4)*tanh(1/2*x)^2+(a^3*b-2*a*b^3)*tanh(1/2*x))/(tanh(1/2*x)^2+1)^4+1/2*(a^4-2*a^2*b^2+b^4)*ln(tanh(1/2*x)^2+1)

Maxima [B] Leaf count of result is larger than twice the leaf count of optimal. 204 vs. $2(77) = 154$.
time = 0.49, size = 204, normalized size = 2.46

$$\frac{2(3a^3 - 5ab^2 + 9a^3 + 3a^2b - 17ab^2 - 3b^3)e^{-2x} + 3(3a^3 + 2a^2b - 5ab^2 - 4b^3)e^{-4x} + 3(a^3 + a^2b - ab^2 - b^3)e^{-6x}}{3(4b^4e^{-2x} + 6b^4e^{-4x} + 4b^4e^{-6x} + b^4e^{-8x} + b^4)} + \frac{(a^4 - 2a^2b^2 + b^4) \log(-(a-b)e^{-2x} - a - b)}{b^5} - \frac{(a^4 - 2a^2b^2 + b^4) \log(e^{-2x} + 1)}{b^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)^6/(a+b*tanh(x)),x, algorithm="maxima")

[Out] $-2/3*(3*a^3 - 5*a*b^2 + (9*a^3 + 3*a^2*b - 17*a*b^2 - 3*b^3)*e^{-2*x} + 3*(3*a^3 + 2*a^2*b - 5*a*b^2 - 4*b^3)*e^{-4*x} + 3*(a^3 + a^2*b - a*b^2 - b^3)*e^{-6*x})/(4*b^4*e^{-2*x} + 6*b^4*e^{-4*x} + 4*b^4*e^{-6*x} + b^4*e^{-8*x} + b^4) + (a^4 - 2*a^2*b^2 + b^4)*\log(-(a - b)*e^{-2*x} - a - b)/b^5 - (a^4 - 2*a^2*b^2 + b^4)*\log(e^{-2*x} + 1)/b^5$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 1827 vs. $2(77) = 154$.
time = 0.38, size = 1827, normalized size = 22.01

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)^6/(a+b*tanh(x)),x, algorithm="fricas")

[Out] $1/3*(6*(a^3*b - a^2*b^2 - a*b^3 + b^4)*\cosh(x)^6 + 36*(a^3*b - a^2*b^2 - a*b^3 + b^4)*\cosh(x)*\sinh(x)^5 + 6*(a^3*b - a^2*b^2 - a*b^3 + b^4)*\sinh(x)^6 + 6*(3*a^3*b - 2*a^2*b^2 - 5*a*b^3 + 4*b^4)*\cosh(x)^4 + 6*(3*a^3*b - 2*a^2*b^2 - 5*a*b^3 + 4*b^4 + 15*(a^3*b - a^2*b^2 - a*b^3 + b^4)*\cosh(x)^2)*\sinh(x)^4 + 6*a^3*b - 10*a*b^3 + 24*(5*(a^3*b - a^2*b^2 - a*b^3 + b^4)*\cosh(x)^3 + (3*a^3*b - 2*a^2*b^2 - 5*a*b^3 + 4*b^4)*\cosh(x))*\sinh(x)^3 + 2*(9*a^3*b - 3*a^2*b^2 - 17*a*b^3 + 3*b^4)*\cosh(x)^2 + 2*(45*(a^3*b - a^2*b^2 - a*b^3 + b^4)*\cosh(x)^4 + 9*a^3*b - 3*a^2*b^2 - 17*a*b^3 + 3*b^4 + 18*(3*a^3*b - 2*a^2*b^2 - 5*a*b^3 + 4*b^4)*\cosh(x)^2)*\sinh(x)^2 + 3*((a^4 - 2*a^2*b^2 + b^4)*\cosh(x)^8 + 8*(a^4 - 2*a^2*b^2 + b^4)*\cosh(x)*\sinh(x)^7 + (a^4 - 2*a^2*b^2 + b^4)*\sinh(x)^8 + 4*(a^4 - 2*a^2*b^2 + b^4)*\cosh(x)^6 + 4*(a^4 - 2*a^2*b^2 + b^4 + 7*(a^4 - 2*a^2*b^2 + b^4)*\cosh(x)^2)*\sinh(x)^6 + 8*(7*(a^4 - 2*a^2*b^2 + b^4)*\cosh(x)^3 + 3*(a^4 - 2*a^2*b^2 + b^4)*\cosh(x))*\sinh(x)^5 + 6*(a^4 - 2*a^2*b^2 + b^4)*\cosh(x)^4 + 2*(35*(a^4 - 2*a^2*b^2 + b^4)*\cosh(x)^4 + 3*a^4 - 6*a^2*b^2 + 3*b^4 + 30*(a^4 - 2*a^2*b^2 + b^4)*\cosh(x)^2)*\sinh(x)^4 + a^4 - 2*a^2*b^2 + b^4 + 8*(7*(a^4 - 2*a^2*b^2 + b^4)*\cosh(x)^5 + 10*(a^4 - 2*a^2*b^2 + b^4)*\cosh(x)^3 + 3*(a^4 - 2*a^2*b^2 + b^4)*\cosh(x))*\sinh(x)^3 + 4*(a^4 - 2*a^2*b^2 + b^4)*\cosh(x)^2 + 4*(7*(a^4 - 2*a^2*b^2 + b^4)*\cosh(x)^6 + 15*(a^4 - 2*a^2*b^2 + b^4)*\cosh(x)^4 + a^4 - 2*a^2*b^2 + b^4 + 9*(a^4 - 2*a^2*b^2 + b^4)*\cosh(x)^2)*\sinh(x)^2 + 8*((a^4 - 2*a^2*b^2 + b^4)*\cosh(x)^7 + 3*(a^4 - 2*a^2*b^2 + b^4)*\cosh(x)^5 + 3*(a^4 - 2*a^2*b^2 + b^4$

```

)*cosh(x)^3 + (a^4 - 2*a^2*b^2 + b^4)*cosh(x))*sinh(x))*log(2*(a*cosh(x) +
b*sinh(x))/(cosh(x) - sinh(x))) - 3*((a^4 - 2*a^2*b^2 + b^4)*cosh(x)^8 + 8*
(a^4 - 2*a^2*b^2 + b^4)*cosh(x)*sinh(x)^7 + (a^4 - 2*a^2*b^2 + b^4)*sinh(x)
^8 + 4*(a^4 - 2*a^2*b^2 + b^4)*cosh(x)^6 + 4*(a^4 - 2*a^2*b^2 + b^4 + 7*(a^
4 - 2*a^2*b^2 + b^4)*cosh(x)^2)*sinh(x)^6 + 8*(7*(a^4 - 2*a^2*b^2 + b^4)*co
sh(x)^3 + 3*(a^4 - 2*a^2*b^2 + b^4)*cosh(x))*sinh(x)^5 + 6*(a^4 - 2*a^2*b^2
+ b^4)*cosh(x)^4 + 2*(35*(a^4 - 2*a^2*b^2 + b^4)*cosh(x)^4 + 3*a^4 - 6*a^2
*b^2 + 3*b^4 + 30*(a^4 - 2*a^2*b^2 + b^4)*cosh(x)^2)*sinh(x)^4 + a^4 - 2*a^
2*b^2 + b^4 + 8*(7*(a^4 - 2*a^2*b^2 + b^4)*cosh(x)^5 + 10*(a^4 - 2*a^2*b^2
+ b^4)*cosh(x)^3 + 3*(a^4 - 2*a^2*b^2 + b^4)*cosh(x))*sinh(x)^3 + 4*(a^4 -
2*a^2*b^2 + b^4)*cosh(x)^2 + 4*(7*(a^4 - 2*a^2*b^2 + b^4)*cosh(x)^6 + 15*(a
^4 - 2*a^2*b^2 + b^4)*cosh(x)^4 + a^4 - 2*a^2*b^2 + b^4 + 9*(a^4 - 2*a^2*b^
2 + b^4)*cosh(x)^2)*sinh(x)^2 + 8*((a^4 - 2*a^2*b^2 + b^4)*cosh(x)^7 + 3*(a
^4 - 2*a^2*b^2 + b^4)*cosh(x)^5 + 3*(a^4 - 2*a^2*b^2 + b^4)*cosh(x)^3 + (a^
4 - 2*a^2*b^2 + b^4)*cosh(x))*sinh(x))*log(2*cosh(x)/(cosh(x) - sinh(x))) +
4*(9*(a^3*b - a^2*b^2 - a*b^3 + b^4)*cosh(x)^5 + 6*(3*a^3*b - 2*a^2*b^2 -
5*a*b^3 + 4*b^4)*cosh(x)^3 + (9*a^3*b - 3*a^2*b^2 - 17*a*b^3 + 3*b^4)*cosh(
x))*sinh(x))/(b^5*cosh(x)^8 + 8*b^5*cosh(x)*sinh(x)^7 + b^5*sinh(x)^8 + 4*b
^5*cosh(x)^6 + 6*b^5*cosh(x)^4 + 4*b^5*cosh(x)^2 + 4*(7*b^5*cosh(x)^2 + b^5
)*sinh(x)^6 + 8*(7*b^5*cosh(x)^3 + 3*b^5*cosh(x))*sinh(x)^5 + b^5 + 2*(35*b
^5*cosh(x)^4 + 30*b^5*cosh(x)^2 + 3*b^5)*sinh(x)^4 + 8*(7*b^5*cosh(x)^5 + 1
0*b^5*cosh(x)^3 + 3*b^5*cosh(x))*sinh(x)^3 + 4*(7*b^5*cosh(x)^6 + 15*b^5*co
sh(x)^4 + 9*b^5*cosh(x)^2 + b^5)*sinh(x)^2 + 8*(b^5*cosh(x)^7 + 3*b^5*cosh(
x)^5 + 3*b^5*cosh(x)^3 + b^5*cosh(x))*sinh(x))

```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{sech}^6(x)}{a + b \tanh(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)**6/(a+b*tanh(x)),x)

[Out] Integral(sech(x)**6/(a + b*tanh(x)), x)

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 316 vs. 2(77) = 154.

time = 0.42, size = 316, normalized size = 3.81

$\frac{(a^6 + 6a^5b - 2a^4b^2 - 2a^3b^3 + a^2b^4 + b^5) \log(\operatorname{abs}(a e^{2x} + b e^{-2x} + a - b))}{a^6 + b^5} - \frac{(a^6 - 2a^5b + 5b^2 \log(e^{2x} + 1))}{5} + \frac{25a^4b^2 - 50a^3b^3 + 25b^4 \log(a^2 e^{2x} + 2ab e^{2x} + b^2) + 25a^4b^2 - 25a^3b^3 + 25b^4 \log(a^2 e^{-2x} + 2ab e^{-2x} + b^2) - 25a^4b^2 + 25a^3b^3 - 25b^4 \log(a^2 e^{2x} + 2ab e^{2x} + b^2) - 25a^4b^2 + 25a^3b^3 - 25b^4 \log(a^2 e^{-2x} + 2ab e^{-2x} + b^2) - 25a^4b^2 + 25a^3b^3 - 25b^4 \log(a^2 e^{2x} + 2ab e^{2x} + b^2) - 25a^4b^2 + 25a^3b^3 - 25b^4 \log(a^2 e^{-2x} + 2ab e^{-2x} + b^2)}{125(b^{5x} + 1)}$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)^6/(a+b*tanh(x)),x, algorithm="giac")

[Out] (a^5 + a^4*b - 2*a^3*b^2 - 2*a^2*b^3 + a*b^4 + b^5)*log(abs(a*e^(2*x) + b*e^(2*x) + a - b))/(a*b^5 + b^6) - (a^4 - 2*a^2*b^2 + b^4)*log(e^(2*x) + 1)/b

$$\begin{aligned} &^5 + 1/12*(25*a^4*e^(8*x) - 50*a^2*b^2*e^(8*x) + 25*b^4*e^(8*x) + 100*a^4*e \\ &^(6*x) + 24*a^3*b*e^(6*x) - 224*a^2*b^2*e^(6*x) - 24*a*b^3*e^(6*x) + 124*b^ \\ &4*e^(6*x) + 150*a^4*e^(4*x) + 72*a^3*b*e^(4*x) - 348*a^2*b^2*e^(4*x) - 120* \\ &a*b^3*e^(4*x) + 246*b^4*e^(4*x) + 100*a^4*e^(2*x) + 72*a^3*b*e^(2*x) - 224* \\ &a^2*b^2*e^(2*x) - 136*a*b^3*e^(2*x) + 124*b^4*e^(2*x) + 25*a^4 + 24*a^3*b - \\ &50*a^2*b^2 - 40*a*b^3 + 25*b^4)/(b^5*(e^(2*x) + 1)^4) \end{aligned}$$

Mupad [B]

time = 1.29, size = 169, normalized size = 2.04

$$\frac{4}{b(4e^{2x} + 6e^{4x} + 4e^{6x} + e^{8x} + 1)} + \frac{2(a-b)^2}{b^3(2e^{2x} + e^{4x} + 1)} + \frac{8(a-3b)}{3b^2(3e^{2x} + 3e^{4x} + e^{6x} + 1)} + \frac{2(a+b)(a-b)^2}{b^4(e^{2x} + 1)} + \frac{\ln(a-b + ae^{2x} + be^{2x})(a+b)^2(a-b)^2}{b^5} - \frac{\ln(e^{2x} + 1)(a+b)^2(a-b)^2}{b^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cosh(x)^6*(a + b*tanh(x))),x)

[Out] 4/(b*(4*exp(2*x) + 6*exp(4*x) + 4*exp(6*x) + exp(8*x) + 1)) + (2*(a - b)^2)/(b^3*(2*exp(2*x) + exp(4*x) + 1)) + (8*(a - 3*b))/(3*b^2*(3*exp(2*x) + 3*exp(4*x) + exp(6*x) + 1)) + (2*(a + b)*(a - b)^2)/(b^4*(exp(2*x) + 1)) + (log(a - b + a*exp(2*x) + b*exp(2*x))*(a + b)^2*(a - b)^2)/b^5 - (log(exp(2*x) + 1)*(a + b)^2*(a - b)^2)/b^5

3.104 $\int \frac{\operatorname{sech}^4(x)}{a+b \tanh(x)} dx$

Optimal. Leaf size=40

$$-\frac{(a^2 - b^2) \log(a + b \tanh(x))}{b^3} + \frac{a \tanh(x)}{b^2} - \frac{\tanh^2(x)}{2b}$$

[Out] $-(a^2-b^2)*\ln(a+b*\tanh(x))/b^3+a*\tanh(x)/b^2-1/2*\tanh(x)^2/b$

Rubi [A]

time = 0.05, antiderivative size = 40, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {3587, 711}

$$-\frac{(a^2 - b^2) \log(a + b \tanh(x))}{b^3} + \frac{a \tanh(x)}{b^2} - \frac{\tanh^2(x)}{2b}$$

Antiderivative was successfully verified.

[In] `Int[Sech[x]^4/(a + b*Tanh[x]),x]`

[Out] $-\left(\left(a^2 - b^2\right) \operatorname{Log}\left[a + b \operatorname{Tanh}\left[x\right]\right]\right) / b^3 + \left(a \operatorname{Tanh}\left[x\right]\right) / b^2 - \operatorname{Tanh}\left[x\right]^2 / \left(2 * b\right)$

Rule 711

`Int[((d_) + (e_.)*(x_))^(m_)*((a_) + (c_.)*(x_)^2)^(p_), x_Symbol] :> Int[ExpandIntegrand[(d + e*x)^m*(a + c*x^2)^p, x], x] /; FreeQ[{a, c, d, e, m}, x] && NeQ[c*d^2 + a*e^2, 0] && IGtQ[p, 0]`

Rule 3587

`Int[sec[(e_.) + (f_.)*(x_)]^(m_)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> Dist[1/(b*f), Subst[Int[(a + x)^n*(1 + x^2/b^2)^(m/2 - 1), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, n}, x] && NeQ[a^2 + b^2, 0] && IntegerQ[m/2]`

Rubi steps

$$\begin{aligned} \int \frac{\operatorname{sech}^4(x)}{a+b \tanh(x)} dx &= \frac{\operatorname{Subst}\left(\int \frac{1-x^2}{a+x} dx, x, b \tanh(x)\right)}{b} \\ &= \frac{\operatorname{Subst}\left(\int \left(\frac{a}{b^2} - \frac{x}{b^2} + \frac{-a^2+b^2}{b^2(a+x)}\right) dx, x, b \tanh(x)\right)}{b} \\ &= -\frac{(a^2 - b^2) \log(a + b \tanh(x))}{b^3} + \frac{a \tanh(x)}{b^2} - \frac{\tanh^2(x)}{2b} \end{aligned}$$

Mathematica [A]

time = 0.13, size = 49, normalized size = 1.22

$$\frac{2(a^2 - b^2)(\log(\cosh(x)) - \log(a \cosh(x) + b \sinh(x))) + b^2 \operatorname{sech}^2(x) + 2ab \tanh(x)}{2b^3}$$

Antiderivative was successfully verified.

[In] Integrate[Sech[x]^4/(a + b*Tanh[x]),x]**[Out]** (2*(a^2 - b^2)*(Log[Cosh[x]] - Log[a*Cosh[x] + b*Sinh[x]]) + b^2*Sech[x]^2 + 2*a*b*Tanh[x])/(2*b^3)**Maple [B]** Leaf count of result is larger than twice the leaf count of optimal. 98 vs. 2(38) = 76.

time = 0.71, size = 99, normalized size = 2.48

method	result	size
default	$-\frac{(a^2 - b^2) \ln(a(\tanh^2(\frac{x}{2}) + 2b \tanh(\frac{x}{2}) + a))}{b^3} + \frac{2(ab(\tanh^3(\frac{x}{2})) - b^2(\tanh^2(\frac{x}{2})) + ab \tanh(\frac{x}{2}))}{(\tanh^2(\frac{x}{2}) + 1)^2} + \frac{(a^2 - b^2) \ln(\tanh^2(\frac{x}{2}) + 1)}{b^3}$	99
risch	$-\frac{2(e^{2x}a - b e^{2x} + a)}{(1 + e^{2x})^2 b^2} + \frac{\ln(1 + e^{2x}) a^2}{b^3} - \frac{\ln(1 + e^{2x})}{b} - \frac{\ln(e^{2x} + \frac{a-b}{a+b}) a^2}{b^3} + \frac{\ln(e^{2x} + \frac{a-b}{a+b})}{b}$	102

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sech(x)^4/(a+b*tanh(x)),x,method=_RETURNVERBOSE)**[Out]** -(a^2-b^2)/b^3*ln(a*tanh(1/2*x)^2+2*b*tanh(1/2*x)+a)+2/b^3*((a*b*tanh(1/2*x))^3-b^2*tanh(1/2*x)^2+a*b*tanh(1/2*x))/(tanh(1/2*x)^2+1)^2+1/2*(a^2-b^2)*ln(tanh(1/2*x)^2+1)**Maxima [B]** Leaf count of result is larger than twice the leaf count of optimal. 89 vs. 2(38) = 76.

time = 0.47, size = 89, normalized size = 2.22

$$\frac{2((a+b)e^{(-2x)}+a)}{2b^2e^{(-2x)}+b^2e^{(-4x)}+b^2} - \frac{(a^2-b^2)\log(-(a-b)e^{(-2x)}-a-b)}{b^3} + \frac{(a^2-b^2)\log(e^{(-2x)}+1)}{b^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)^4/(a+b*tanh(x)),x, algorithm="maxima")**[Out]** 2*((a+b)*e^(-2*x)+a)/(2*b^2*e^(-2*x)+b^2*e^(-4*x)+b^2)-(a^2-b^2)*log(-(a-b)*e^(-2*x)-a-b)/b^3+(a^2-b^2)*log(e^(-2*x)+1)/b^3**Fricas [B]** Leaf count of result is larger than twice the leaf count of optimal. 430 vs. 2(38) = 76.

time = 0.43, size = 430, normalized size = 10.75

2(a^2-b^2)*log(cosh(x))-2(a^2-b^2)*log(a*cosh(x)+b*sinh(x))+b^2*sech(x)^2+2*a*b*tanh(x)/2b^3

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)^4/(a+b*tanh(x)),x, algorithm="fricas")

[Out] $-(2*(a*b - b^2)*\cosh(x)^2 + 4*(a*b - b^2)*\cosh(x)*\sinh(x) + 2*(a*b - b^2)*\sinh(x)^2 + 2*a*b + ((a^2 - b^2)*\cosh(x)^4 + 4*(a^2 - b^2)*\cosh(x)*\sinh(x)^3 + (a^2 - b^2)*\sinh(x)^4 + 2*(a^2 - b^2)*\cosh(x)^2 + 2*(3*(a^2 - b^2)*\cosh(x)^2 + a^2 - b^2)*\sinh(x)^2 + a^2 - b^2 + 4*((a^2 - b^2)*\cosh(x)^3 + (a^2 - b^2)*\cosh(x))*\sinh(x))*\log(2*(a*\cosh(x) + b*\sinh(x))/(\cosh(x) - \sinh(x))) - ((a^2 - b^2)*\cosh(x)^4 + 4*(a^2 - b^2)*\cosh(x)*\sinh(x)^3 + (a^2 - b^2)*\sinh(x)^4 + 2*(a^2 - b^2)*\cosh(x)^2 + 2*(3*(a^2 - b^2)*\cosh(x)^2 + a^2 - b^2)*\sinh(x)^2 + a^2 - b^2 + 4*((a^2 - b^2)*\cosh(x)^3 + (a^2 - b^2)*\cosh(x))*\sinh(x))*\log(2*\cosh(x)/(\cosh(x) - \sinh(x)))/(b^3*\cosh(x)^4 + 4*b^3*\cosh(x)*\sinh(x)^3 + b^3*\sinh(x)^4 + 2*b^3*\cosh(x)^2 + b^3 + 2*(3*b^3*\cosh(x)^2 + b^3)*\sinh(x)^2 + 4*(b^3*\cosh(x)^3 + b^3*\cosh(x))*\sinh(x))$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{sech}^4(x)}{a + b \tanh(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)**4/(a+b*tanh(x)),x)

[Out] Integral(sech(x)**4/(a + b*tanh(x)), x)

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 104 vs. 2(38) = 76.

time = 0.41, size = 104, normalized size = 2.60

$$-\frac{(a^3 + a^2b - ab^2 - b^3) \log(|ae^{(2x)} + be^{(2x)} + a - b|)}{ab^3 + b^4} + \frac{(a^2 - b^2) \log(e^{(2x)} + 1)}{b^3} - \frac{2(ab + (ab - b^2)e^{(2x)})}{b^3(e^{(2x)} + 1)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)^4/(a+b*tanh(x)),x, algorithm="giac")

[Out] $-(a^3 + a^2*b - a*b^2 - b^3)*\log(\operatorname{abs}(a*e^{(2*x)} + b*e^{(2*x)} + a - b))/(a*b^3 + b^4) + (a^2 - b^2)*\log(e^{(2*x)} + 1)/b^3 - 2*(a*b + (a*b - b^2)*e^{(2*x)})/(b^3*(e^{(2*x)} + 1)^2)$

Mupad [B]

time = 1.27, size = 88, normalized size = 2.20

$$\frac{\ln(e^{2x} + 1) (a + b) (a - b)}{b^3} - \frac{2(a - b)}{b^2 (e^{2x} + 1)} - \frac{\ln(a - b + a e^{2x} + b e^{2x}) (a + b) (a - b)}{b^3} - \frac{2}{b (2e^{2x} + e^{4x} + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(cosh(x)^4*(a + b*tanh(x))),x)
```

```
[Out] (log(exp(2*x) + 1)*(a + b)*(a - b))/b^3 - (2*(a - b))/(b^2*(exp(2*x) + 1))  
- (log(a - b + a*exp(2*x) + b*exp(2*x))*(a + b)*(a - b))/b^3 - 2/(b*(2*exp(  
2*x) + exp(4*x) + 1))
```

3.105 $\int \frac{\operatorname{sech}^2(x)}{a+b \tanh(x)} dx$

Optimal. Leaf size=11

$$\frac{\log(a + b \tanh(x))}{b}$$

[Out] ln(a+b*tanh(x))/b

Rubi [A]

time = 0.03, antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {3587, 31}

$$\frac{\log(a + b \tanh(x))}{b}$$

Antiderivative was successfully verified.

[In] Int[Sech[x]^2/(a + b*Tanh[x]),x]

[Out] Log[a + b*Tanh[x]]/b

Rule 31

Int[((a_) + (b_)*(x_))⁽⁻¹⁾, x_Symbol] := Simp[Log[RemoveContent[a + b*x, x]]/b, x] /; FreeQ[{a, b}, x]

Rule 3587

Int[sec[(e_.) + (f_.)*(x_)]^(m_)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Dist[1/(b*f), Subst[Int[(a + x)^{n*(1 + x^2/b^2)}^(m/2 - 1), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, n}, x] && NeQ[a^2 + b^2, 0] && IntegerQ[m/2]

Rubi steps

$$\begin{aligned} \int \frac{\operatorname{sech}^2(x)}{a+b \tanh(x)} dx &= \frac{\operatorname{Subst}\left(\int \frac{1}{a+x} dx, x, b \tanh(x)\right)}{b} \\ &= \frac{\log(a + b \tanh(x))}{b} \end{aligned}$$

Mathematica [A]

time = 0.04, size = 20, normalized size = 1.82

$$\frac{-\log(\cosh(x)) + \log(a \cosh(x) + b \sinh(x))}{b}$$

Antiderivative was successfully verified.

[In] Integrate[Sech[x]^2/(a + b*Tanh[x]),x]

[Out] (-Log[Cosh[x]] + Log[a*Cosh[x] + b*Sinh[x]])/b

Maple [A]

time = 0.52, size = 12, normalized size = 1.09

method	result	size
derivativedivides	$\frac{\ln(a+b \tanh(x))}{b}$	12
default	$\frac{\ln(a+b \tanh(x))}{b}$	12
risch	$-\frac{\ln(1+e^{2x})}{b} + \frac{\ln\left(e^{2x} + \frac{a-b}{a+b}\right)}{b}$	35

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sech(x)^2/(a+b*tanh(x)),x,method=_RETURNVERBOSE)

[Out] ln(a+b*tanh(x))/b

Maxima [A]

time = 0.25, size = 11, normalized size = 1.00

$$\frac{\log(b \tanh(x) + a)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)^2/(a+b*tanh(x)),x, algorithm="maxima")

[Out] log(b*tanh(x) + a)/b

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 42 vs. 2(11) = 22.

time = 0.38, size = 42, normalized size = 3.82

$$\frac{\log\left(\frac{2(a \cosh(x) + b \sinh(x))}{\cosh(x) - \sinh(x)}\right) - \log\left(\frac{2 \cosh(x)}{\cosh(x) - \sinh(x)}\right)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)^2/(a+b*tanh(x)),x, algorithm="fricas")

[Out] (log(2*(a*cosh(x) + b*sinh(x))/(cosh(x) - sinh(x))) - log(2*cosh(x)/(cosh(x) - sinh(x))))/b

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{sech}^2(x)}{a + b \tanh(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sech(x)**2/(a+b*tanh(x)),x)`

[Out] `Integral(sech(x)**2/(a + b*tanh(x)), x)`

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 45 vs. 2(11) = 22.
time = 0.41, size = 45, normalized size = 4.09

$$\frac{(a+b)\log(|ae^{2x} + be^{2x} + a - b|)}{ab + b^2} - \frac{\log(e^{2x} + 1)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sech(x)^2/(a+b*tanh(x)),x, algorithm="giac")`

[Out] `(a + b)*log(abs(a*e^(2*x) + b*e^(2*x) + a - b))/(a*b + b^2) - log(e^(2*x) + 1)/b`

Mupad [B]

time = 0.21, size = 50, normalized size = 4.55

$$-\frac{2 \operatorname{atan}\left(\frac{a\sqrt{-b^2} + ae^{2x}\sqrt{-b^2} + be^{2x}\sqrt{-b^2}}{b^2}\right)}{\sqrt{-b^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(cosh(x)^2*(a + b*tanh(x))),x)`

[Out] `-(2*atan((a*(-b^2)^(1/2) + a*exp(2*x)*(-b^2)^(1/2) + b*exp(2*x)*(-b^2)^(1/2))/b^2))/(-b^2)^(1/2)`

3.106 $\int \frac{1}{a+b \tanh(x)} dx$

Optimal. Leaf size=39

$$\frac{ax}{a^2 - b^2} - \frac{b \log(a \cosh(x) + b \sinh(x))}{a^2 - b^2}$$

[Out] a*x/(a^2-b^2)-b*ln(a*cosh(x)+b*sinh(x))/(a^2-b^2)

Rubi [A]

time = 0.03, antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3565, 3611}

$$\frac{ax}{a^2 - b^2} - \frac{b \log(a \cosh(x) + b \sinh(x))}{a^2 - b^2}$$

Antiderivative was successfully verified.

[In] Int[(a + b*Tanh[x])^(-1), x]

[Out] (a*x)/(a^2 - b^2) - (b*Log[a*Cosh[x] + b*Sinh[x]])/(a^2 - b^2)

Rule 3565

Int[((a_) + (b_)*tan[(c_) + (d_)*(x_)])^(-1), x_Symbol] := Simp[a*(x/(a^2 + b^2)), x] + Dist[b/(a^2 + b^2), Int[(b - a*Tan[c + d*x])/(a + b*Tan[c + d*x]), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0]

Rule 3611

Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{a + b \tanh(x)} dx &= \frac{ax}{a^2 - b^2} - \frac{(ib) \int \frac{-ib - ia \tanh(x)}{a + b \tanh(x)} dx}{a^2 - b^2} \\ &= \frac{ax}{a^2 - b^2} - \frac{b \log(a \cosh(x) + b \sinh(x))}{a^2 - b^2} \end{aligned}$$

Mathematica [A]

time = 0.04, size = 29, normalized size = 0.74

$$\frac{ax - b \log(a \cosh(x) + b \sinh(x))}{a^2 - b^2}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*Tanh[x])^(-1),x]

[Out] (a*x - b*Log[a*Cosh[x] + b*Sinh[x]])/(a^2 - b^2)

Maple [A]

time = 0.24, size = 55, normalized size = 1.41

method	result	size
derivativedivides	$\frac{\ln(1+\tanh(x))}{2a-2b} - \frac{\ln(\tanh(x)-1)}{2b+2a} - \frac{b \ln(a+b \tanh(x))}{(a-b)(a+b)}$	55
default	$\frac{\ln(1+\tanh(x))}{2a-2b} - \frac{\ln(\tanh(x)-1)}{2b+2a} - \frac{b \ln(a+b \tanh(x))}{(a-b)(a+b)}$	55
risch	$\frac{x}{a+b} + \frac{2bx}{a^2-b^2} - \frac{b \ln\left(e^{2x} + \frac{a-b}{a+b}\right)}{a^2-b^2}$	55

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a+b*tanh(x)),x,method=_RETURNVERBOSE)

[Out] 1/(2*a-2*b)*ln(1+tanh(x))-1/(2*b+2*a)*ln(tanh(x)-1)-b/(a-b)/(a+b)*ln(a+b*tanh(x))

Maxima [A]

time = 0.27, size = 41, normalized size = 1.05

$$-\frac{b \log\left(-\frac{(a-b)e^{-2x}}{a^2-b^2} - \frac{a-b}{a^2-b^2}\right)}{a^2-b^2} + \frac{x}{a+b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+b*tanh(x)),x, algorithm="maxima")

[Out] -b*log(-(a - b)*e^(-2*x) - a - b)/(a^2 - b^2) + x/(a + b)

Fricas [A]

time = 0.40, size = 42, normalized size = 1.08

$$\frac{(a+b)x - b \log\left(\frac{2(a \cosh(x) + b \sinh(x))}{\cosh(x) - \sinh(x)}\right)}{a^2 - b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+b*tanh(x)),x, algorithm="fricas")

[Out] ((a + b)*x - b*log(2*(a*cosh(x) + b*sinh(x))/(cosh(x) - sinh(x))))/(a^2 - b^2)

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 146 vs. $2(29) = 58$.

time = 0.26, size = 146, normalized size = 3.74

$$\left\{ \begin{array}{ll} \tilde{\infty}(x - \log(\tanh(x) + 1) + \log(\tanh(x))) & \text{for } a = 0 \wedge b = 0 \\ \frac{x}{a} & \text{for } b = 0 \\ -\frac{x \tanh(x)}{2b \tanh(x) - 2b} + \frac{x}{2b \tanh(x) - 2b} + \frac{1}{2b \tanh(x) - 2b} & \text{for } a = -b \\ \frac{x \tanh(x)}{2b \tanh(x) + 2b} + \frac{x}{2b \tanh(x) + 2b} - \frac{1}{2b \tanh(x) + 2b} & \text{for } a = b \\ \frac{ax}{a^2 - b^2} - \frac{bx}{a^2 - b^2} - \frac{b \log\left(\frac{a}{b} + \tanh(x)\right)}{a^2 - b^2} + \frac{b \log(\tanh(x) + 1)}{a^2 - b^2} & \text{otherwise} \end{array} \right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+b*tanh(x)),x)

[Out] Piecewise((zoo*(x - log(tanh(x) + 1) + log(tanh(x))), Eq(a, 0) & Eq(b, 0)), (x/a, Eq(b, 0)), (-x*tanh(x)/(2*b*tanh(x) - 2*b) + x/(2*b*tanh(x) - 2*b) + 1/(2*b*tanh(x) - 2*b), Eq(a, -b)), (x*tanh(x)/(2*b*tanh(x) + 2*b) + x/(2*b*tanh(x) + 2*b) - 1/(2*b*tanh(x) + 2*b), Eq(a, b)), (a*x/(a**2 - b**2) - b*x/(a**2 - b**2) - b*log(a/b + tanh(x))/(a**2 - b**2) + b*log(tanh(x) + 1)/(a**2 - b**2), True))

Giac [A]

time = 0.41, size = 43, normalized size = 1.10

$$-\frac{b \log(|ae^{(2x)} + be^{(2x)} + a - b|)}{a^2 - b^2} + \frac{x}{a - b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+b*tanh(x)),x, algorithm="giac")

[Out] -b*log(abs(a*e^(2*x) + b*e^(2*x) + a - b))/(a^2 - b^2) + x/(a - b)

Mupad [B]

time = 1.11, size = 35, normalized size = 0.90

$$\frac{ax - b(x - \ln(\tanh(x) + 1) + \ln(a + b \tanh(x)))}{a^2 - b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a + b*tanh(x)),x)

[Out] (a*x - b*(x - log(tanh(x) + 1) + log(a + b*tanh(x))))/(a^2 - b^2)

3.107 $\int \frac{\cosh^2(x)}{a+b \tanh(x)} dx$

Optimal. Leaf size=91

$$-\frac{(a+2b)\log(1-\tanh(x))}{4(a+b)^2} + \frac{(a-2b)\log(1+\tanh(x))}{4(a-b)^2} + \frac{b^3\log(a+b\tanh(x))}{(a^2-b^2)^2} - \frac{\cosh^2(x)(b-a\tanh(x))}{2(a^2-b^2)}$$

[Out] $-1/4*(a+2*b)*\ln(1-\tanh(x))/(a+b)^2+1/4*(a-2*b)*\ln(1+\tanh(x))/(a-b)^2+b^3*\ln(a+b*\tanh(x))/(a^2-b^2)^2-1/2*\cosh(x)^2*(b-a*\tanh(x))/(a^2-b^2)$

Rubi [A]

time = 0.11, antiderivative size = 91, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {3587, 755, 815}

$$-\frac{\cosh^2(x)(b-a\tanh(x))}{2(a^2-b^2)} + \frac{b^3\log(a+b\tanh(x))}{(a^2-b^2)^2} - \frac{(a+2b)\log(1-\tanh(x))}{4(a+b)^2} + \frac{(a-2b)\log(\tanh(x)+1)}{4(a-b)^2}$$

Antiderivative was successfully verified.

[In] `Int[Cosh[x]^2/(a + b*Tanh[x]),x]`

[Out] $-1/4*((a+2*b)*\text{Log}[1-\text{Tanh}[x]]/(a+b)^2 + ((a-2*b)*\text{Log}[1+\text{Tanh}[x]])/(4*(a-b)^2) + (b^3*\text{Log}[a+b*\text{Tanh}[x]])/(a^2-b^2)^2 - (\text{Cosh}[x]^2*(b-a*\text{Tanh}[x]))/(2*(a^2-b^2))$

Rule 755

`Int[((d_) + (e_.)*(x_))^(m_)*((a_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[(- (d + e*x)^(m + 1))*(a*e + c*d*x)*((a + c*x^2)^(p + 1)/(2*a*(p + 1)*(c*d^2 + a*e^2))), x] + Dist[1/(2*a*(p + 1)*(c*d^2 + a*e^2)), Int[(d + e*x)^m*Simp[c*d^2*(2*p + 3) + a*e^2*(m + 2*p + 3) + c*e*d*(m + 2*p + 4)*x, x]*(a + c*x^2)^(p + 1), x], x] /; FreeQ[{a, c, d, e, m}, x] && NeQ[c*d^2 + a*e^2, 0] && LtQ[p, -1] && IntQuadraticQ[a, 0, c, d, e, m, p, x]`

Rule 815

`Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_)))/((a_) + (c_.)*(x_)^2), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*((f + g*x)/(a + c*x^2)), x], x] /; FreeQ[{a, c, d, e, f, g}, x] && NeQ[c*d^2 + a*e^2, 0] && IntegerQ[m]`

Rule 3587

`Int[sec[(e_.) + (f_.)*(x_)]^(m_)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Dist[1/(b*f), Subst[Int[(a + x)^n*(1 + x^2/b^2)^(m/2 - 1), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, n}, x] && NeQ[a^2 + b^2, 0] && IntegerQ[m/2]`

Rubi steps

$$\begin{aligned}
 \int \frac{\cosh^2(x)}{a + b \tanh(x)} dx &= \frac{\text{Subst}\left(\int \frac{1}{(a+x)\left(1-\frac{x^2}{b^2}\right)^2} dx, x, b \tanh(x)\right)}{b} \\
 &= -\frac{\cosh^2(x)(b - a \tanh(x))}{2(a^2 - b^2)} + \frac{b \text{Subst}\left(\int \frac{-2 + \frac{a^2}{b^2} + \frac{ax}{b^2}}{(a+x)\left(1-\frac{x^2}{b^2}\right)} dx, x, b \tanh(x)\right)}{2(a^2 - b^2)} \\
 &= -\frac{\cosh^2(x)(b - a \tanh(x))}{2(a^2 - b^2)} + \frac{b \text{Subst}\left(\int \left(\frac{(a-b)(a+2b)}{2b(a+b)(b-x)} + \frac{2b^2}{(a-b)(a+b)(a+x)} + \frac{(a-2b)(a+b)}{2(a-b)b(b+x)}\right) dx, x, b \tanh(x)\right)}{2(a^2 - b^2)} \\
 &= -\frac{(a + 2b) \log(1 - \tanh(x))}{4(a + b)^2} + \frac{(a - 2b) \log(1 + \tanh(x))}{4(a - b)^2} + \frac{b^3 \log(a + b \tanh(x))}{(a^2 - b^2)^2} - \frac{ab^3}{(a^2 - b^2)^2}
 \end{aligned}$$

Mathematica [A]

time = 0.09, size = 75, normalized size = 0.82

$$\frac{2a^3x - 6ab^2x + (-a^2b + b^3) \cosh(2x) + 4b^3 \log(a \cosh(x) + b \sinh(x)) + a(a^2 - b^2) \sinh(2x)}{4(a - b)^2(a + b)^2}$$

Antiderivative was successfully verified.

[In] Integrate[Cosh[x]^2/(a + b*Tanh[x]), x]

[Out] (2*a^3*x - 6*a*b^2*x + (-a^2*b) + b^3)*Cosh[2*x] + 4*b^3*Log[a*Cosh[x] + b*Sinh[x]] + a*(a^2 - b^2)*Sinh[2*x]/(4*(a - b)^2*(a + b)^2)

Maple [A]

time = 0.71, size = 153, normalized size = 1.68

method	result
risch	$\frac{xb}{(a+b)^2} + \frac{ax}{2(a+b)^2} + \frac{e^{2x}}{8a+8b} - \frac{e^{-2x}}{8(a-b)} - \frac{2b^3x}{a^4-2a^2b^2+b^4} + \frac{b^3 \ln\left(e^{2x} + \frac{a-b}{a+b}\right)}{a^4-2a^2b^2+b^4}$
default	$\frac{b^3 \ln\left(a\left(\tanh^2\left(\frac{x}{2}\right)\right) + 2b \tanh\left(\frac{x}{2}\right) + a\right)}{(a-b)^2(a+b)^2} - \frac{1}{(2a-2b)\left(\tanh\left(\frac{x}{2}\right) + 1\right)^2} + \frac{2}{(4a-4b)\left(\tanh\left(\frac{x}{2}\right) + 1\right)} + \frac{(-2b+a) \ln\left(\tanh\left(\frac{x}{2}\right) + 1\right)}{2(a-b)^2} + \frac{1}{(2b+2a)}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cosh(x)^2/(a+b*tanh(x)), x, method=_RETURNVERBOSE)

[Out] b^3/(a-b)^2/(a+b)^2*ln(a*tanh(1/2*x)^2+2*b*tanh(1/2*x)+a)-1/(2*a-2*b)/(tanh(1/2*x)+1)^2+2/(4*a-4*b)/(tanh(1/2*x)+1)+1/2*(-2*b+a)/(a-b)^2*ln(tanh(1/2*x)+1)+1/(2*b+2*a)/(tanh(1/2*x)-1)^2+2/(4*a+4*b)/(tanh(1/2*x)-1)+1/2/(a+b)^2*(-2*b-a)*ln(tanh(1/2*x)-1)

Maxima [A]

time = 0.29, size = 86, normalized size = 0.95

$$\frac{b^3 \log(-(a-b)e^{(-2x)} - a - b)}{a^4 - 2a^2b^2 + b^4} + \frac{(a+2b)x}{2(a^2 + 2ab + b^2)} + \frac{e^{(2x)}}{8(a+b)} - \frac{e^{(-2x)}}{8(a-b)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cosh(x)^2/(a+b*tanh(x)),x, algorithm="maxima")**[Out]** b^3*log(-(a - b)*e^(-2*x) - a - b)/(a^4 - 2*a^2*b^2 + b^4) + 1/2*(a + 2*b)*x/(a^2 + 2*a*b + b^2) + 1/8*e^(2*x)/(a + b) - 1/8*e^(-2*x)/(a - b)**Fricas [B]** Leaf count of result is larger than twice the leaf count of optimal. 331 vs. 2(86) = 172.

time = 0.40, size = 331, normalized size = 3.64

$$\frac{(a^4 - a^3b - a^2b^2 + ab^3) \cosh(x)^4 + 4(a^3 - a^2b - ab^2 + b^3) \cosh(x)^3 \sinh(x) + (a^3 - a^2b - ab^2 + b^3) \sinh(x)^4 + 4(a^3 - 3a^2b - a^2b^2 + b^3) x \cosh(x)^2 - a^3 - a^2b + ab^2 + b^3 + 2(3(a^3 - a^2b - ab^2 + b^3) \cosh(x)^2 + 2(a^3 - 3a^2b - 2b^3) x) \sinh(x)^2 + 8(b^3 \cosh(x)^2 + 2b^3 \cosh(x) \sinh(x) + b^3 \sinh(x)^2) \log(2(a \cosh(x) + b \sinh(x)) / (\cosh(x) - \sinh(x))) + 4((a^3 - a^2b - ab^2 + b^3) \cosh(x)^3 + 2(a^3 - 3a^2b - 2b^3) x \cosh(x)) \sinh(x)}{(a^4 - 2a^2b^2 + b^4) \cosh(x)^2 + 2(a^4 - 2a^2b^2 + b^4) \cosh(x) \sinh(x) + (a^4 - 2a^2b^2 + b^4) \sinh(x)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cosh(x)^2/(a+b*tanh(x)),x, algorithm="fricas")

[Out] 1/8*((a^3 - a^2*b - a*b^2 + b^3)*cosh(x)^4 + 4*(a^3 - a^2*b - a*b^2 + b^3)*cosh(x)*sinh(x)^3 + (a^3 - a^2*b - a*b^2 + b^3)*sinh(x)^4 + 4*(a^3 - 3*a*b^2 - 2*b^3)*x*cosh(x)^2 - a^3 - a^2*b + a*b^2 + b^3 + 2*(3*(a^3 - a^2*b - a*b^2 + b^3)*cosh(x)^2 + 2*(a^3 - 3*a*b^2 - 2*b^3)*x)*sinh(x)^2 + 8*(b^3*cosh(x)^2 + 2*b^3*cosh(x)*sinh(x) + b^3*sinh(x)^2)*log(2*(a*cosh(x) + b*sinh(x))/(cosh(x) - sinh(x))) + 4*((a^3 - a^2*b - a*b^2 + b^3)*cosh(x)^3 + 2*(a^3 - 3*a*b^2 - 2*b^3)*x*cosh(x))*sinh(x)/((a^4 - 2*a^2*b^2 + b^4)*cosh(x)^2 + 2*(a^4 - 2*a^2*b^2 + b^4)*cosh(x)*sinh(x) + (a^4 - 2*a^2*b^2 + b^4)*sinh(x)^2)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cosh^2(x)}{a + b \tanh(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cosh(x)**2/(a+b*tanh(x)),x)**[Out]** Integral(cosh(x)**2/(a + b*tanh(x)), x)**Giac [A]**

time = 0.42, size = 111, normalized size = 1.22

$$\frac{b^3 \log(|ae^{(2x)} + be^{(2x)} + a - b|)}{a^4 - 2a^2b^2 + b^4} + \frac{(a-2b)x}{2(a^2 - 2ab + b^2)} - \frac{(2ae^{(2x)} - 4be^{(2x)} + a - b)e^{(-2x)}}{8(a^2 - 2ab + b^2)} + \frac{e^{(2x)}}{8(a+b)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cosh(x)^2/(a+b*tanh(x)),x, algorithm="giac")

[Out] $b^3 \log(\operatorname{abs}(a e^{2x} + b e^{2x} + a - b)) / (a^4 - 2 a^2 b^2 + b^4) + 1/2 (a - 2b) x / (a^2 - 2 a b + b^2) - 1/8 (2 a e^{2x} - 4 b e^{2x} + a - b) e^{-2x} / (a^2 - 2 a b + b^2) + 1/8 e^{2x} / (a + b)$

Mupad [B]

time = 1.28, size = 84, normalized size = 0.92

$$\frac{e^{2x}}{8a + 8b} - \frac{e^{-2x}}{8a - 8b} + \frac{b^3 \ln(a - b + a e^{2x} + b e^{2x})}{a^4 - 2a^2 b^2 + b^4} + \frac{x(a - 2b)}{2(a - b)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cosh(x)^2/(a + b*tanh(x)),x)

[Out] $\exp(2x)/(8a + 8b) - \exp(-2x)/(8a - 8b) + (b^3 \log(a - b + a \exp(2x) + b \exp(2x))) / (a^4 + b^4 - 2a^2 b^2) + (x(a - 2b)) / (2(a - b)^2)$

3.108 $\int \frac{\cosh^4(x)}{a+b \tanh(x)} dx$

Optimal. Leaf size=155

$$-\frac{(3a^2 + 9ab + 8b^2) \log(1 - \tanh(x))}{16(a+b)^3} + \frac{(3a^2 - 9ab + 8b^2) \log(1 + \tanh(x))}{16(a-b)^3} - \frac{b^5 \log(a + b \tanh(x))}{(a^2 - b^2)^3} - \frac{\cosh^4(x)}{4}$$

```
[Out] -1/16*(3*a^2+9*a*b+8*b^2)*ln(1-tanh(x))/(a+b)^3+1/16*(3*a^2-9*a*b+8*b^2)*ln
(1+tanh(x))/(a-b)^3-b^5*ln(a+b*tanh(x))/(a^2-b^2)^3-1/4*cosh(x)^4*(b-a*tanh
(x))/(a^2-b^2)+1/8*cosh(x)^2*(4*b^3-a*(7-3*a^2/b^2)*b^2*tanh(x))/(a^2-b^2)^
2
```

Rubi [A]

time = 0.17, antiderivative size = 155, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$,

Rules used = {3587, 755, 837, 815}

$$-\frac{(3a^2 + 9ab + 8b^2) \log(1 - \tanh(x))}{16(a+b)^3} + \frac{(3a^2 - 9ab + 8b^2) \log(\tanh(x) + 1)}{16(a-b)^3} - \frac{\cosh^4(x)(b - a \tanh(x))}{4(a^2 - b^2)} - \frac{b^5 \log(a + b \tanh(x))}{(a^2 - b^2)^3} + \frac{\cosh^2(x) \left(4b^3 - ab^2 \left(7 - \frac{3a^2}{b^2} \right) \tanh(x) \right)}{8(a^2 - b^2)^2}$$

Antiderivative was successfully verified.

```
[In] Int[Cosh[x]^4/(a + b*Tanh[x]),x]
```

```
[Out] -1/16*((3*a^2 + 9*a*b + 8*b^2)*Log[1 - Tanh[x]]/(a + b)^3 + ((3*a^2 - 9*a*
b + 8*b^2)*Log[1 + Tanh[x]]/(16*(a - b)^3) - (b^5*Log[a + b*Tanh[x]]/(a^2
- b^2)^3 - (Cosh[x]^4*(b - a*Tanh[x]))/(4*(a^2 - b^2)) + (Cosh[x]^2*(4*b^3
- a*(7 - (3*a^2)/b^2)*b^2*Tanh[x]))/(8*(a^2 - b^2)^2)
```

Rule 755

```
Int[((d_) + (e_)*(x_))^(m_)*((a_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[
(-(d + e*x)^(m + 1))*(a*e + c*d*x)*((a + c*x^2)^(p + 1)/(2*a*(p + 1)*(c*d^2
+ a*e^2))), x] + Dist[1/(2*a*(p + 1)*(c*d^2 + a*e^2)), Int[(d + e*x)^m*Sim
p[c*d^2*(2*p + 3) + a*e^2*(m + 2*p + 3) + c*e*d*(m + 2*p + 4)*x, x]*(a + c*
x^2)^(p + 1), x], x] /; FreeQ[{a, c, d, e, m}, x] && NeQ[c*d^2 + a*e^2, 0]
&& LtQ[p, -1] && IntQuadraticQ[a, 0, c, d, e, m, p, x]
```

Rule 815

```
Int[(((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_)))/((a_) + (c_)*(x_)^2),
x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*((f + g*x)/(a + c*x^2)), x],
x] /; FreeQ[{a, c, d, e, f, g}, x] && NeQ[c*d^2 + a*e^2, 0] && IntegerQ[m]
```

Rule 837

```
Int[((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_))*((a_) + (c_)*(x_)^2)^(p
_), x_Symbol] := Simp[(-(d + e*x)^(m + 1))*(f*a*c*e - a*g*c*d + c*(c*d*f +
```

```

a*e*g)*x)*((a + c*x^2)^(p + 1)/(2*a*c*(p + 1)*(c*d^2 + a*e^2))), x] + Dist[
1/(2*a*c*(p + 1)*(c*d^2 + a*e^2)), Int[(d + e*x)^m*(a + c*x^2)^(p + 1)*Simp
[f*(c^2*d^2*(2*p + 3) + a*c*e^2*(m + 2*p + 3)) - a*c*d*e*g*m + c*e*(c*d*f +
a*e*g)*(m + 2*p + 4)*x, x], x] /; FreeQ[{a, c, d, e, f, g}, x] && NeQ[
c*d^2 + a*e^2, 0] && LtQ[p, -1] && (IntegerQ[m] || IntegerQ[p] || IntegersQ
[2*m, 2*p])

```

Rule 3587

```

Int[sec[(e_.) + (f_.)*(x_)]^(m_)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]^(n_
), x_Symbol] := Dist[1/(b*f), Subst[Int[(a + x)^n*(1 + x^2/b^2)^(m/2 - 1),
x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, n}, x] && NeQ[a^2 + b^2, 0
] && IntegerQ[m/2]

```

Rubi steps

$$\begin{aligned}
\int \frac{\cosh^4(x)}{a + b \tanh(x)} dx &= \frac{\text{Subst}\left(\int \frac{1}{(a+x)\left(1-\frac{x^2}{b^2}\right)^3} dx, x, b \tanh(x)\right)}{b} \\
&= -\frac{\cosh^4(x)(b - a \tanh(x))}{4(a^2 - b^2)} + \frac{b \text{Subst}\left(\int \frac{-4 + \frac{3a^2}{b^2} + \frac{3ax}{b^2}}{(a+x)\left(1-\frac{x^2}{b^2}\right)^2} dx, x, b \tanh(x)\right)}{4(a^2 - b^2)} \\
&= -\frac{\cosh^4(x)(b - a \tanh(x))}{4(a^2 - b^2)} + \frac{\cosh^2(x)\left(4b^3 - a\left(7 - \frac{3a^2}{b^2}\right)b^2 \tanh(x)\right)}{8(a^2 - b^2)^2} - \frac{b^5 \text{Subst}\left(\int \frac{1}{(a+x)\left(1-\frac{x^2}{b^2}\right)} dx, x, b \tanh(x)\right)}{8(a^2 - b^2)^2} \\
&= -\frac{\cosh^4(x)(b - a \tanh(x))}{4(a^2 - b^2)} + \frac{\cosh^2(x)\left(4b^3 - a\left(7 - \frac{3a^2}{b^2}\right)b^2 \tanh(x)\right)}{8(a^2 - b^2)^2} - \frac{b^5 \text{Subst}\left(\int \frac{1}{(a+x)\left(1-\frac{x^2}{b^2}\right)} dx, x, b \tanh(x)\right)}{8(a^2 - b^2)^2} \\
&= -\frac{(3a^2 + 9ab + 8b^2) \log(1 - \tanh(x))}{16(a + b)^3} + \frac{(3a^2 - 9ab + 8b^2) \log(1 + \tanh(x))}{16(a - b)^3} - \frac{b^5 \log(a - b^2 \tanh^2(x) + a^2)}{8(a^2 - b^2)^2}
\end{aligned}$$

Mathematica [A]

time = 0.15, size = 156, normalized size = 1.01

$$\frac{12a^5x - 40a^3b^2x + 60ab^4x - 4b(a^4 - 4a^2b^2 + 3b^4) \cosh(2x) - b(a^2 - b^2)^2 \cosh(4x) - 32b^5 \log(a \cosh(x) + b \sinh(x)) + 8a^5 \sinh(2x) - 24a^3b^2 \sinh(2x) + 16ab^4 \sinh(2x) + a^5 \sinh(4x) - 2a^3b^2 \sinh(4x) + ab^4 \sinh(4x)}{32(a - b)^3(a + b)^3}$$

Antiderivative was successfully verified.

[In] Integrate[Cosh[x]^4/(a + b*Tanh[x]), x]

[Out] $(12*a^5*x - 40*a^3*b^2*x + 60*a*b^4*x - 4*b*(a^4 - 4*a^2*b^2 + 3*b^4)*\text{Cosh}[2*x] - b*(a^2 - b^2)^2*\text{Cosh}[4*x] - 32*b^5*\text{Log}[a*\text{Cosh}[x] + b*\text{Sinh}[x]] + 8*a^5*\text{Sinh}[2*x] - 24*a^3*b^2*\text{Sinh}[2*x] + 16*a*b^4*\text{Sinh}[2*x] + a^5*\text{Sinh}[4*x] - 2*a^3*b^2*\text{Sinh}[4*x] + a*b^4*\text{Sinh}[4*x])/(32*(a - b)^3*(a + b)^3)$

Maple [A]

time = 0.71, size = 265, normalized size = 1.71

method	result
risch	$\frac{3x a^2}{8(a+b)^3} + \frac{9xab}{8(a+b)^3} + \frac{x b^2}{(a+b)^3} + \frac{e^{4x}}{64a+64b} + \frac{e^{2x}a}{8(a+b)^2} + \frac{3e^{2x}b}{16(a+b)^2} - \frac{e^{-2x}a}{8(a-b)^2} + \frac{3e^{-2x}b}{16(a-b)^2} - \frac{e^{-4x}}{64(a-b)} + \frac{2b^5x}{a^6-3a^4b^2+3a^2b^4-b^6}$
default	$\frac{1}{2(2b+2a)(\tanh(\frac{x}{2})-1)^4} + \frac{2}{(4a+4b)(\tanh(\frac{x}{2})-1)^3} - \frac{-7a-9b}{8(a+b)^2(\tanh(\frac{x}{2})-1)^2} - \frac{-5a-7b}{8(a+b)^2(\tanh(\frac{x}{2})-1)} + \frac{(-3a^2-9ab-8b^2)\ln(\tanh(\frac{x}{2})-1)}{8(a+b)}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cosh(x)^4/(a+b*tanh(x)),x,method=_RETURNVERBOSE)`

[Out] $1/2/(2*b+2*a)/(\tanh(1/2*x)-1)^4+2/(4*a+4*b)/(\tanh(1/2*x)-1)^3-1/8*(-7*a-9*b)/(a+b)^2/(\tanh(1/2*x)-1)^2-1/8*(-5*a-7*b)/(a+b)^2/(\tanh(1/2*x)-1)+1/8/(a+b)^3*(-3*a^2-9*a*b-8*b^2)*\ln(\tanh(1/2*x)-1)-b^5/(a-b)^3/(a+b)^3*\ln(a*\tanh(1/2*x)^2+2*b*\tanh(1/2*x)+a)-1/2/(2*a-2*b)/(\tanh(1/2*x)+1)^4+2/(4*a-4*b)/(\tanh(1/2*x)+1)^3-1/8*(-5*a+7*b)/(a-b)^2/(\tanh(1/2*x)+1)-1/8*(7*a-9*b)/(a-b)^2/(\tanh(1/2*x)+1)^2+1/8*(3*a^2-9*a*b+8*b^2)/(a-b)^3*\ln(\tanh(1/2*x)+1)$

Maxima [A]

time = 0.27, size = 165, normalized size = 1.06

$$-\frac{b^5 \log(-(a-b)e^{(-2x)}-a-b)}{a^6-3a^4b^2+3a^2b^4-b^6} + \frac{(3a^2+9ab+8b^2)x}{8(a^3+3a^2b+3ab^2+b^3)} + \frac{(4(2a+3b)e^{(-2x)}+a+b)e^{(4x)}}{64(a^2+2ab+b^2)} - \frac{4(2a-3b)e^{(-2x)}+(a-b)e^{(-4x)}}{64(a^2-2ab+b^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cosh(x)^4/(a+b*tanh(x)),x, algorithm="maxima")`

[Out] $-b^5*\log(-(a - b)*e^{(-2*x)} - a - b)/(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6) + 1/8*(3*a^2 + 9*a*b + 8*b^2)*x/(a^3 + 3*a^2*b + 3*a*b^2 + b^3) + 1/64*(4*(2*a + 3*b)*e^{(-2*x)} + a + b)*e^{(4*x)}/(a^2 + 2*a*b + b^2) - 1/64*(4*(2*a - 3*b)*e^{(-2*x)} + (a - b)*e^{(-4*x)})/(a^2 - 2*a*b + b^2)$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 1281 vs. $2(147) = 294$.

time = 0.49, size = 1281, normalized size = 8.26

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cosh(x)^4/(a+b*tanh(x)),x, algorithm="fricas")`

```
[Out] 1/64*((a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*cosh(x)^8 + 8*(a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*cosh(x)*sinh(x)^7 + (a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*sinh(x)^8 + 4*(2*a^5 - a^4*b - 6*a^3*b^2 + 4*a^2*b^3 + 4*a*b^4 - 3*b^5)*cosh(x)^6 + 4*(2*a^5 - a^4*b - 6*a^3*b^2 + 4*a^2*b^3 + 4*a*b^4 - 3*b^5) *cosh(x)^5 + 7*(a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*cosh(x)^4 + 8*(3*a^5 - 10*a^3*b^2 + 15*a*b^4 + 8*b^5)*x*cosh(x)^4 + 8*(7*(a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*cosh(x)^3 + 3*(2*a^5 - a^4*b - 6*a^3*b^2 + 4*a^2*b^3 + 4*a*b^4 - 3*b^5)*cosh(x))*sinh(x)^5 - a^5 - a^4*b + 2*a^3*b^2 + 2*a^2*b^3 - a*b^4 - b^5 + 2*(35*(a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*cosh(x)^4 + 30*(2*a^5 - a^4*b - 6*a^3*b^2 + 4*a^2*b^3 + 4*a*b^4 - 3*b^5)*cosh(x)^2 + 4*(3*a^5 - 10*a^3*b^2 + 15*a*b^4 + 8*b^5)*x)*sinh(x)^4 + 8*(7*(a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*cosh(x)^5 + 10*(2*a^5 - a^4*b - 6*a^3*b^2 + 4*a^2*b^3 + 4*a*b^4 - 3*b^5)*cosh(x)^3 + 4*(3*a^5 - 10*a^3*b^2 + 15*a*b^4 + 8*b^5)*x*cosh(x))*sinh(x)^3 - 4*(2*a^5 + a^4*b - 6*a^3*b^2 - 4*a^2*b^3 + 4*a*b^4 + 3*b^5)*cosh(x)^2 + 4*(7*(a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*cosh(x)^6 - 2*a^5 - a^4*b + 6*a^3*b^2 + 4*a^2*b^3 - 4*a*b^4 - 3*b^5 + 15*(2*a^5 - a^4*b - 6*a^3*b^2 + 4*a^2*b^3 + 4*a*b^4 - 3*b^5)*cosh(x)^4 + 12*(3*a^5 - 10*a^3*b^2 + 15*a*b^4 + 8*b^5)*x*cosh(x)^2)*sinh(x)^2 - 64*(b^5*cosh(x)^4 + 4*b^5*cosh(x)^3*sinh(x) + 6*b^5*cosh(x)^2*sinh(x)^2 + 4*b^5*cosh(x)*sinh(x)^3 + b^5*sinh(x)^4)*log(2*(a*cosh(x) + b*sinh(x))/(cosh(x) - sinh(x))) + 8*((a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*cosh(x)^7 + 3*(2*a^5 - a^4*b - 6*a^3*b^2 + 4*a^2*b^3 + 4*a*b^4 - 3*b^5)*cosh(x)^5 + 4*(3*a^5 - 10*a^3*b^2 + 15*a*b^4 + 8*b^5)*x*cosh(x)^3 - (2*a^5 + a^4*b - 6*a^3*b^2 - 4*a^2*b^3 + 4*a*b^4 + 3*b^5)*cosh(x))*sinh(x))/((a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*cosh(x)^4 + 4*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*cosh(x)^3*sinh(x) + 6*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*cosh(x)^2*sinh(x)^2 + 4*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*cosh(x)*sinh(x)^3 + (a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*sinh(x)^4)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cosh^4(x)}{a + b \tanh(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cosh(x)**4/(a+b*tanh(x)),x)
```

```
[Out] Integral(cosh(x)**4/(a + b*tanh(x)), x)
```

Giac [A]

time = 0.41, size = 227, normalized size = 1.46

$$-\frac{b^5 \log(ae^{2x} + be^{2x} + a - b)}{a^6 - 3a^4b^2 + 3a^2b^4 - b^6} + \frac{(3a^2 - 9ab + 8b^2)x}{8(a^3 - 3a^2b + 3ab^2 - b^3)} - \frac{(18a^2e^{4x} - 54abe^{4x} + 48b^2e^{4x} + 8a^2e^{2x} - 20abe^{2x} + 12b^2e^{2x} + a^2 - 2ab + b^2)e^{-4x}}{64(a^3 - 3a^2b + 3ab^2 - b^3)} + \frac{ae^{4x} + be^{4x} + 8ae^{2x} + 12be^{2x}}{64(a^2 + 2ab + b^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cosh(x)^4/(a+b*tanh(x)),x, algorithm="giac")

[Out] $-b^5 \log(\operatorname{abs}(a e^{2x} + b e^{2x} + a - b)) / (a^6 - 3a^4 b^2 + 3a^2 b^4 - b^6) + 1/8 (3a^2 - 9ab + 8b^2) x / (a^3 - 3a^2 b + 3ab^2 - b^3) - 1/64 (18a^2 e^{4x} - 54ab e^{4x} + 48b^2 e^{4x} + 8a^2 e^{2x} - 20ab e^{2x} + 12b^2 e^{2x} + a^2 - 2ab + b^2) e^{-4x} / (a^3 - 3a^2 b + 3ab^2 - b^3) + 1/64 (a e^{4x} + b e^{4x} + 8a e^{2x} + 12b e^{2x}) / (a^2 + 2ab + b^2)$

Mupad [B]

time = 1.51, size = 143, normalized size = 0.92

$$\frac{e^{4x}}{64a + 64b} - \frac{e^{-4x}}{64a - 64b} - \frac{e^{-2x}(2a - 3b)}{16(a - b)^2} - \frac{b^5 \ln(a - b + a e^{2x} + b e^{2x})}{a^6 - 3a^4 b^2 + 3a^2 b^4 - b^6} + \frac{x(3a^2 - 9ab + 8b^2)}{8(a - b)^3} + \frac{e^{2x}(2a + 3b)}{16(a + b)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cosh(x)^4/(a + b*tanh(x)),x)

[Out] $\exp(4x)/(64a + 64b) - \exp(-4x)/(64a - 64b) - (\exp(-2x)(2a - 3b)) / (16(a - b)^2) - (b^5 \log(a - b + a \exp(2x) + b \exp(2x))) / (a^6 - b^6 + 3a^2 b^4 - 3a^4 b^2) + (x(3a^2 - 9ab + 8b^2)) / (8(a - b)^3) + (\exp(2x)(2a + 3b)) / (16(a + b)^2)$

3.109 $\int \frac{\operatorname{sech}^7(x)}{a+b \tanh(x)} dx$

Optimal. Leaf size=157

$$\frac{a(8a^4 - 20a^2b^2 + 15b^4) \operatorname{ArcTan}(\sinh(x))}{8b^6} - \frac{(a^2 - b^2)^{5/2} \operatorname{ArcTan}\left(\frac{\cosh(x)(b+a \tanh(x))}{\sqrt{a^2 - b^2}}\right)}{b^6} + \frac{(a^2 - b^2)^2 \operatorname{sech}(x)}{b^5} - \frac{(a^2 - b^2)^2 \operatorname{sech}(x)^3 \tanh(x)}{b^4} + \frac{(a^2 - b^2)^2 \operatorname{sech}(x)^5}{5b} + \frac{(a^2 - b^2)^2 \operatorname{sech}(x)^3 \tanh(x)}{3b^3} + \frac{(a^2 - b^2)^2 \operatorname{sech}(x)}{b^5} - \frac{(a^2 - b^2)^2 \operatorname{sech}(x)^3 \tanh(x)}{b^4} + \frac{(a^2 - b^2)^2 \operatorname{sech}(x)^5}{5b}$$

[Out] 1/8*a*(8*a^4-20*a^2*b^2+15*b^4)*arctan(sinh(x))/b^6-(a^2-b^2)^(5/2)*arctan(cosh(x)*(b+a*tanh(x))/(a^2-b^2)^(1/2))/b^6+(a^2-b^2)^2*sech(x)/b^5-1/3*(a^2-b^2)*sech(x)^3/b^3+1/5*sech(x)^5/b-1/8*a*(4*a^2-7*b^2)*sech(x)*tanh(x)/b^4+1/4*a*sech(x)^3*tanh(x)/b^2

Rubi [A]

time = 0.20, antiderivative size = 187, normalized size of antiderivative = 1.19, number of steps used = 14, number of rules used = 6, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.462$, Rules used = {3591, 3567, 3853, 3855, 3590, 212}

$$\frac{a(a^2 - b^2)^2 \operatorname{ArcTan}(\sinh(x))}{b^6} - \frac{(a^2 - b^2)^{5/2} \operatorname{ArcTan}\left(\frac{\cosh(x)(b+a \tanh(x))}{\sqrt{a^2 - b^2}}\right)}{b^6} - \frac{a(a^2 - b^2) \operatorname{ArcTan}(\sinh(x))}{2b^4} + \frac{(a^2 - b^2)^2 \operatorname{sech}(x)}{b^5} - \frac{a(a^2 - b^2) \tanh(x) \operatorname{sech}(x)}{2b^4} - \frac{(a^2 - b^2) \operatorname{sech}^2(x)}{3b^3} + \frac{3a \operatorname{ArcTan}(\sinh(x))}{8b^2} + \frac{a \tanh(x) \operatorname{sech}^2(x)}{4b^2} + \frac{3a \tanh(x) \operatorname{sech}(x)}{8b^2} + \frac{\operatorname{sech}^3(x)}{5b}$$

Antiderivative was successfully verified.

[In] Int[Sech[x]^7/(a + b*Tanh[x]), x]

[Out] (3*a*ArcTan[Sinh[x]])/(8*b^2) - (a*(a^2 - b^2)*ArcTan[Sinh[x]])/(2*b^4) + (a*(a^2 - b^2)^2*ArcTan[Sinh[x]]/b^6 - ((a^2 - b^2)^(5/2)*ArcTan[(Cosh[x]*(b + a*Tanh[x])/Sqrt[a^2 - b^2]])/b^6 + ((a^2 - b^2)^2*Sech[x])/b^5 - ((a^2 - b^2)*Sech[x]^3)/(3*b^3) + Sech[x]^5/(5*b) + (3*a*Sech[x]*Tanh[x])/(8*b^2) - (a*(a^2 - b^2)*Sech[x]*Tanh[x])/(2*b^4) + (a*Sech[x]^3*Tanh[x])/(4*b^2)

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 3567

Int[((d_)*sec[(e_) + (f_)*(x_)])^(m_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[b*((d*Sec[e + f*x])^m/(f*m)), x] + Dist[a, Int[(d*Sec[e + f*x])^m, x], x] /; FreeQ[{a, b, d, e, f, m}, x] && (IntegerQ[2*m] | NeQ[a^2 + b^2, 0])

Rule 3590

Int[sec[(e_) + (f_)*(x_)]/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Dist[-f^(-1), Subst[Int[1/(a^2 + b^2 - x^2), x], x, (b - a*Tan[e + f*x])], x]

x])/Sec[e + f*x]], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a^2 + b^2, 0]

Rule 3591

Int[((d_)*sec[(e_.) + (f_.)*(x_)])^(m_)/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Dist[-d^2/b^2, Int[(d*Sec[e + f*x])^(m - 2)*(a - b*Tan[e + f*x]), x], x] + Dist[d^2*((a^2 + b^2)/b^2), Int[(d*Sec[e + f*x])^(m - 2)/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 + b^2, 0] && IGtQ[m, 1]

Rule 3853

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.)^(n_)), x_Symbol] := Simp[(-b)*Cos[c + d*x]*(b*Csc[c + d*x])^(n - 1)/(d*(n - 1)), x] + Dist[b^2*((n - 2)/(n - 1)), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3855

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{\operatorname{sech}^7(x)}{a + b \tanh(x)} dx &= \frac{\int \operatorname{sech}^5(x)(a - b \tanh(x)) dx}{b^2} - \frac{(a^2 - b^2) \int \frac{\operatorname{sech}^5(x)}{a + b \tanh(x)} dx}{b^2} \\ &= \frac{\operatorname{sech}^5(x)}{5b} + \frac{a \int \operatorname{sech}^5(x) dx}{b^2} - \frac{(a^2 - b^2) \int \operatorname{sech}^3(x)(a - b \tanh(x)) dx}{b^4} + \frac{(a^2 - b^2)^2 \int \operatorname{sech}^3(x)}{b^4} \\ &= -\frac{(a^2 - b^2) \operatorname{sech}^3(x)}{3b^3} + \frac{\operatorname{sech}^5(x)}{5b} + \frac{a \operatorname{sech}^3(x) \tanh(x)}{4b^2} + \frac{(3a) \int \operatorname{sech}^3(x) dx}{4b^2} - \frac{a(a^2 - b^2)}{b^4} \\ &= \frac{(a^2 - b^2)^2 \operatorname{sech}(x)}{b^5} - \frac{(a^2 - b^2) \operatorname{sech}^3(x)}{3b^3} + \frac{\operatorname{sech}^5(x)}{5b} + \frac{3a \operatorname{sech}(x) \tanh(x)}{8b^2} - \frac{a(a^2 - b^2)}{b^4} \\ &= \frac{3a \tan^{-1}(\sinh(x))}{8b^2} - \frac{a(a^2 - b^2) \tan^{-1}(\sinh(x))}{2b^4} + \frac{a(a^2 - b^2)^2 \tan^{-1}(\sinh(x))}{b^6} - \frac{(a^2 - b^2)^2}{b^4} \end{aligned}$$

Mathematica [A]

time = 0.36, size = 166, normalized size = 1.06

$$\frac{30 \left(a(8a^4 - 20a^2b^2 + 15b^4) \operatorname{ArcTan}\left(\tanh\left(\frac{x}{2}\right)\right) - 8\sqrt{a-b}\sqrt{a+b}(a^2 - b^2)^2 \operatorname{ArcTan}\left(\frac{b+a \tanh\left(\frac{x}{2}\right)}{\sqrt{a-b}\sqrt{a+b}}\right) \right) + 24b^5 \operatorname{sech}^5(x) + 10b^3 \operatorname{sech}^3(x)(-4a^2 + 4b^2 + 3ab \tanh(x)) + 15b \operatorname{sech}(x) \left(8(a^2 - b^2)^2 + (-4a^3b + 7ab^2) \tanh(x) \right)}{120b^6}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sech[x]^7/(a + b*Tanh[x]),x]
```

```
[Out] (30*(a*(8*a^4 - 20*a^2*b^2 + 15*b^4)*ArcTan[Tanh[x/2]] - 8*Sqrt[a - b]*Sqrt[a + b]*(a^2 - b^2)^2*ArcTan[(b + a*Tanh[x/2])/(Sqrt[a - b]*Sqrt[a + b])]) + 24*b^5*Sech[x]^5 + 10*b^3*Sech[x]^3*(-4*a^2 + 4*b^2 + 3*a*b*Tanh[x]) + 15*b*Sech[x]*(8*(a^2 - b^2)^2 + (-4*a^3*b + 7*a*b^3)*Tanh[x]))/(120*b^6)
```

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 322 vs. 2(143) = 286.

time = 1.07, size = 323, normalized size = 2.06

method	result
default	$\frac{2\left(\left(\frac{1}{2}a^3b^2 - \frac{9}{8}ab^4\right)\left(\tanh^9\left(\frac{x}{2}\right)\right) + \left(a^4b - 3a^2b^3 + 3b^5\right)\left(\tanh^8\left(\frac{x}{2}\right)\right) + \left(a^3b^2 - \frac{5}{4}ab^4\right)\left(\tanh^7\left(\frac{x}{2}\right)\right) + \left(4a^4b - 10a^2b^3 + 6b^5\right)\left(\tanh^6\left(\frac{x}{2}\right)\right) + \left(6a^4b - \frac{40}{3}a^2b^3 + 6b^5\right)\left(\tanh^5\left(\frac{x}{2}\right)\right) + \left(2a^4b - 4a^2b^3 + 2b^5\right)\left(\tanh^4\left(\frac{x}{2}\right)\right) + \left(2a^4b - 4a^2b^3 + 2b^5\right)\left(\tanh^3\left(\frac{x}{2}\right)\right) + \left(2a^4b - 4a^2b^3 + 2b^5\right)\left(\tanh^2\left(\frac{x}{2}\right)\right) + \left(2a^4b - 4a^2b^3 + 2b^5\right)\left(\tanh\left(\frac{x}{2}\right)\right) + \left(2a^4b - 4a^2b^3 + 2b^5\right)}{120b^6}$
risch	$\frac{e^x(120a^4e^{8x} - 60a^3be^{8x} - 240a^2b^2e^{8x} + 105ab^3e^{8x} + 120b^4e^{8x} + 480a^4e^{6x} - 120a^3be^{6x} - 1120a^2b^2e^{6x} + 330ab^3e^{6x} + 640b^4e^{6x} + 720a^4e^{4x} - 120a^3be^{4x} - 1120a^2b^2e^{4x} + 330ab^3e^{4x} + 640b^4e^{4x} + 720a^4e^{2x} - 120a^3be^{2x} - 1120a^2b^2e^{2x} + 330ab^3e^{2x} + 640b^4e^{2x} + 720a^4e^x - 120a^3be^x - 1120a^2b^2e^x + 330ab^3e^x + 640b^4e^x + 720a^4)}{60b^5(1 + \tanh^2(\frac{x}{2}))}$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sech(x)^7/(a+b*tanh(x)),x,method=_RETURNVERBOSE)
```

```
[Out] 2/b^6*(((1/2*a^3*b^2-9/8*a*b^4)*tanh(1/2*x)^9+(a^4*b-3*a^2*b^3+3*b^5)*tanh(1/2*x)^8+(a^3*b^2-5/4*a*b^4)*tanh(1/2*x)^7+(4*a^4*b-10*a^2*b^3+6*b^5)*tanh(1/2*x)^6+(6*a^4*b-40/3*a^2*b^3+28/3*b^5)*tanh(1/2*x)^4+(-a^3*b^2+5/4*a*b^4)*tanh(1/2*x)^3+(4*a^4*b-26/3*a^2*b^3+14/3*b^5)*tanh(1/2*x)^2+(-1/2*a^3*b^2+9/8*a*b^4)*tanh(1/2*x)+a^4*b-7/3*a^2*b^3+23/15*b^5)/(tanh(1/2*x)^2+1)^5+1/8*a*(8*a^4-20*a^2*b^2+15*b^4)*arctan(tanh(1/2*x))+2*(-a^6+3*a^4*b^2-3*a^2*b^4+b^6)/b^6/(a^2-b^2)^(1/2)*arctan(1/2*(2*a*tanh(1/2*x)+2*b)/(a^2-b^2)^(1/2)))
```

Maxima [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sech(x)^7/(a+b*tanh(x)),x, algorithm="maxima")
```

```
[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see 'assume?' for more details)
```

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 3227 vs. 2(143) = 286.

time = 0.51, size = 6509, normalized size = 41.46

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)^7/(a+b*tanh(x)),x, algorithm="fricas")

[Out] [1/60*(15*(8*a^4*b - 4*a^3*b^2 - 16*a^2*b^3 + 7*a*b^4 + 8*b^5)*cosh(x)^9 + 135*(8*a^4*b - 4*a^3*b^2 - 16*a^2*b^3 + 7*a*b^4 + 8*b^5)*cosh(x)*sinh(x)^8 + 15*(8*a^4*b - 4*a^3*b^2 - 16*a^2*b^3 + 7*a*b^4 + 8*b^5)*sinh(x)^9 + 10*(4*8*a^4*b - 12*a^3*b^2 - 112*a^2*b^3 + 33*a*b^4 + 64*b^5)*cosh(x)^7 + 10*(48*a^4*b - 12*a^3*b^2 - 112*a^2*b^3 + 33*a*b^4 + 64*b^5 + 54*(8*a^4*b - 4*a^3*b^2 - 16*a^2*b^3 + 7*a*b^4 + 8*b^5)*cosh(x)^2)*sinh(x)^7 + 70*(18*(8*a^4*b - 4*a^3*b^2 - 16*a^2*b^3 + 7*a*b^4 + 8*b^5)*cosh(x)^3 + (48*a^4*b - 12*a^3*b^2 - 112*a^2*b^3 + 33*a*b^4 + 64*b^5)*cosh(x))*sinh(x)^6 + 16*(45*a^4*b - 110*a^2*b^3 + 89*b^5)*cosh(x)^5 + 2*(360*a^4*b - 880*a^2*b^3 + 712*b^5 + 94*5*(8*a^4*b - 4*a^3*b^2 - 16*a^2*b^3 + 7*a*b^4 + 8*b^5)*cosh(x)^4 + 105*(48*a^4*b - 12*a^3*b^2 - 112*a^2*b^3 + 33*a*b^4 + 64*b^5)*cosh(x)^2)*sinh(x)^5 + 10*(189*(8*a^4*b - 4*a^3*b^2 - 16*a^2*b^3 + 7*a*b^4 + 8*b^5)*cosh(x)^5 + 35*(48*a^4*b - 12*a^3*b^2 - 112*a^2*b^3 + 33*a*b^4 + 64*b^5)*cosh(x)^3 + 8*(45*a^4*b - 110*a^2*b^3 + 89*b^5)*cosh(x))*sinh(x)^4 + 10*(48*a^4*b + 12*a^3*b^2 - 112*a^2*b^3 - 33*a*b^4 + 64*b^5)*cosh(x)^3 + 10*(126*(8*a^4*b - 4*a^3*b^2 - 16*a^2*b^3 + 7*a*b^4 + 8*b^5)*cosh(x)^6 + 48*a^4*b + 12*a^3*b^2 - 112*a^2*b^3 - 33*a*b^4 + 64*b^5 + 35*(48*a^4*b - 12*a^3*b^2 - 112*a^2*b^3 + 33*a*b^4 + 64*b^5)*cosh(x)^4 + 16*(45*a^4*b - 110*a^2*b^3 + 89*b^5)*cosh(x)^2)*sinh(x)^3 + 10*(54*(8*a^4*b - 4*a^3*b^2 - 16*a^2*b^3 + 7*a*b^4 + 8*b^5)*cosh(x)^7 + 21*(48*a^4*b - 12*a^3*b^2 - 112*a^2*b^3 + 33*a*b^4 + 64*b^5)*cosh(x)^5 + 16*(45*a^4*b - 110*a^2*b^3 + 89*b^5)*cosh(x)^3 + 3*(48*a^4*b + 12*a^3*b^2 - 112*a^2*b^3 - 33*a*b^4 + 64*b^5)*cosh(x))*sinh(x)^2 + 60*((a^4 - 2*a^2*b^2 + b^4)*cosh(x)^10 + 10*(a^4 - 2*a^2*b^2 + b^4)*cosh(x)*sinh(x)^9 + (a^4 - 2*a^2*b^2 + b^4)*sinh(x)^10 + 5*(a^4 - 2*a^2*b^2 + b^4)*cosh(x)^8 + 5*(a^4 - 2*a^2*b^2 + b^4 + 9*(a^4 - 2*a^2*b^2 + b^4)*cosh(x)^2)*sinh(x)^8 + 40*(3*(a^4 - 2*a^2*b^2 + b^4)*cosh(x)^3 + (a^4 - 2*a^2*b^2 + b^4)*cosh(x))*sinh(x)^7 + 10*(a^4 - 2*a^2*b^2 + b^4)*cosh(x)^6 + 10*(21*(a^4 - 2*a^2*b^2 + b^4)*cosh(x)^4 + a^4 - 2*a^2*b^2 + b^4 + 14*(a^4 - 2*a^2*b^2 + b^4)*cosh(x)^2)*sinh(x)^6 + 4*(63*(a^4 - 2*a^2*b^2 + b^4)*cosh(x)^5 + 70*(a^4 - 2*a^2*b^2 + b^4)*cosh(x)^3 + 15*(a^4 - 2*a^2*b^2 + b^4)*cosh(x))*sinh(x)^5 + 10*(a^4 - 2*a^2*b^2 + b^4)*cosh(x)^4 + 10*(21*(a^4 - 2*a^2*b^2 + b^4)*cosh(x)^6 + 35*(a^4 - 2*a^2*b^2 + b^4)*cosh(x)^4 + a^4 - 2*a^2*b^2 + b^4 + 15*(a^4 - 2*a^2*b^2 + b^4)*cosh(x)^2)*sinh(x)^4 + a^4 - 2*a^2*b^2 + b^4 + 40*(3*(a^4 - 2*a^2*b^2 + b^4)*cosh(x)^7 + 7*(a^4 - 2*a^2*b^2 + b^4)*cosh(x)^5 + 5*(a^4 - 2*a^2*b^2 + b^4)*cosh(x)^3 + (a^4 - 2*a^2*b^2 + b^4)*cosh(x))*sinh(x)^3 + 5*(a^4 - 2*a^2*b^2 + b^4)*cosh(x)^2 + 5*(9*(a^4 - 2*a^2*b^2 + b^4)

$$\begin{aligned}
&) * \cosh(x)^8 + 28*(a^4 - 2*a^2*b^2 + b^4) * \cosh(x)^6 + 30*(a^4 - 2*a^2*b^2 + \\
& b^4) * \cosh(x)^4 + a^4 - 2*a^2*b^2 + b^4 + 12*(a^4 - 2*a^2*b^2 + b^4) * \cosh(x) \\
& ^2 * \sinh(x)^2 + 10*((a^4 - 2*a^2*b^2 + b^4) * \cosh(x)^9 + 4*(a^4 - 2*a^2*b^2 \\
& + b^4) * \cosh(x)^7 + 6*(a^4 - 2*a^2*b^2 + b^4) * \cosh(x)^5 + 4*(a^4 - 2*a^2*b^2 \\
& + b^4) * \cosh(x)^3 + (a^4 - 2*a^2*b^2 + b^4) * \cosh(x) * \sinh(x)) * \sqrt{-a^2 + b \\
& ^2} * \log(((a + b) * \cosh(x)^2 + 2*(a + b) * \cosh(x) * \sinh(x) + (a + b) * \sinh(x)^2 \\
& - 2 * \sqrt{-a^2 + b^2} * (\cosh(x) + \sinh(x)) - a + b) / ((a + b) * \cosh(x)^2 + 2*(a \\
& + b) * \cosh(x) * \sinh(x) + (a + b) * \sinh(x)^2 + a - b)) + 15*((8*a^5 - 20*a^3*b \\
& ^2 + 15*a*b^4) * \cosh(x)^10 + 10*(8*a^5 - 20*a^3*b^2 + 15*a*b^4) * \cosh(x) * \sinh \\
& (x)^9 + (8*a^5 - 20*a^3*b^2 + 15*a*b^4) * \sinh(x)^10 + 5*(8*a^5 - 20*a^3*b^2 \\
& + 15*a*b^4) * \cosh(x)^8 + 5*(8*a^5 - 20*a^3*b^2 + 15*a*b^4 + 9*(8*a^5 - 20*a^ \\
& 3*b^2 + 15*a*b^4) * \cosh(x)^2) * \sinh(x)^8 + 40*(3*(8*a^5 - 20*a^3*b^2 + 15*a*b \\
& ^4) * \cosh(x)^3 + (8*a^5 - 20*a^3*b^2 + 15*a*b^4) * \cosh(x)) * \sinh(x)^7 + 10*(8* \\
& a^5 - 20*a^3*b^2 + 15*a*b^4) * \cosh(x)^6 + 10*(8*a^5 - 20*a^3*b^2 + 15*a*b^4 \\
& + 21*(8*a^5 - 20*a^3*b^2 + 15*a*b^4) * \cosh(x)^4 + 14*(8*a^5 - 20*a^3*b^2 + 1 \\
& 5*a*b^4) * \cosh(x)^2) * \sinh(x)^6 + 4*(63*(8*a^5 - 20*a^3*b^2 + 15*a*b^4) * \cosh(\\
& x)^5 + 70*(8*a^5 - 20*a^3*b^2 + 15*a*b^4) * \cosh(x)^3 + 15*(8*a^5 - 20*a^3*b^ \\
& 2 + 15*a*b^4) * \cosh(x)) * \sinh(x)^5 + 8*a^5 - 20*a^3*b^2 + 15*a*b^4 + 10*(8*a^ \\
& 5 - 20*a^3*b^2 + 15*a*b^4) * \cosh(x)^4 + 10*(21*(8*a^5 - 20*a^3*b^2 + 15*a*b^ \\
& 4) * \cosh(x)^6 + 8*a^5 - 20*a^3*b^2 + 15*a*b^4 + 35*(8*a^5 - 20*a^3*b^2 + 15* \\
& a*b^4) * \cosh(x)^4 + 15*(8*a^5 - 20*a^3*b^2 + 15*a*b^4) * \cosh(x)^2) * \sinh(x)^4 \\
& + 40*(3*(8*a^5 - 20*a^3*b^2 + 15*a*b^4) * \cosh(x)^7 + 7*(8*a^5 - 20*a^3*b^2 + \\
& 15*a*b^4) * \cosh(x)^5 + 5*(8*a^5 - 20*a^3*b^2 + 15*a*b^4) * \cosh(x)^3 + (8*a^5 \\
& - 20*a^3*b^2 + 15*a*b^4) * \cosh(x)) * \sinh(x)^3 + 5*(8*a^5 - 20*a^3*b^2 + 15*a \\
& *b^4) * \cosh(x)^2 + 5*(9*(8*a^5 - 20*a^3*b^2 + 15*a*b^4) * \cosh(x)^8 + 28*(8*a^ \\
& 5 - 20*a^3*b^2 + 15*a*b^4) * \cosh(x)^6 + 8*a^5 - 20*a^3*b^2 + 15*a*b^4 + 30*(\\
& 8*a^5 - 20*a^3*b^2 + 15*a*b^4) * \cosh(x)^4 + 12*(8*a^5 - 20*a^3*b^2 + 15*a*b^ \\
& 4) * \cosh(x)^2) * \sinh(x)^2 + 10*((8*a^5 - 20*a^3*b^2 + 15*a*b^4) * \cosh(x)^9 + 4 \\
& *(8*a^5 - 20*a^3*b^2 + 15*a*b^4) * \cosh(x)^7 + 6*(8*a^5 - 20*a^3*b^2 + 15*a*b \\
& ^4) * \cosh(x)^5 + 4*(8*a^5 - 20*a^3*b^2 + 15*a*b^4) * \cosh(x)^3 + (8*a^5 - 20*a \\
& ^3*b^2 + 15*a*b^4) * \cosh(x)) * \sinh(x)) * \arctan(\cos \dots
\end{aligned}$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{sech}^7(x)}{a + b \tanh(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)**7/(a+b*tanh(x)),x)

[Out] Integral(sech(x)**7/(a + b*tanh(x)), x)

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 326 vs. 2(143) = 286.

time = 0.41, size = 326, normalized size = 2.08

$$\frac{(8a^2 - 20a^2b + 15ab^2)\arctan\left(\frac{e^x}{b}\right) - \frac{2(a^2 - 3a^2b + 3b^2)\arctan\left(\frac{e^x}{\sqrt{a^2 - b^2}}\right)}{\sqrt{a^2 - b^2}} + \frac{120a^{10}e^{10x} - 60a^8b^{10}e^{10x} + 105a^6b^{10}e^{10x} + 120a^4b^{10}e^{10x} + 60a^2b^{10}e^{10x} - 120a^{10}b^2e^{10x} - 1120a^8b^2e^{10x} + 330a^6b^2e^{10x} + 640a^4b^2e^{10x} - 1760a^2b^2e^{10x} + 1424b^4e^{10x} + 480a^4b^4e^{10x} + 120a^2b^4e^{10x} - 1120a^2b^4e^{10x} - 330a^2b^4e^{10x} + 640b^6e^{10x} + 120a^4b^6e^{10x} - 1120a^4b^6e^{10x} - 330a^4b^6e^{10x} + 640a^6b^6e^{10x} + 120a^8b^6e^{10x} - 1120a^8b^6e^{10x} - 330a^8b^6e^{10x} + 640a^{10}b^6e^{10x}}{60b^{10}(e^{10x} + 1)^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)^7/(a+b*tanh(x)),x, algorithm="giac")

[Out] $\frac{1}{4}(8a^5 - 20a^3b^2 + 15a^2b^4)\arctan\left(\frac{e^x}{b}\right) - 2(a^6 - 3a^4b^2 + 3a^2b^4 - b^6)\arctan\left(\frac{a e^x + b}{\sqrt{a^2 - b^2}}\right) + \frac{1}{60}(120a^4b^4e^{9x} - 60a^3b^4e^{9x} - 240a^2b^2e^{9x} + 105a^2b^3e^{9x} + 120b^4e^{9x} + 480a^4b^4e^{7x} - 120a^3b^4e^{7x} - 1120a^2b^2e^{7x} + 330a^2b^3e^{7x} + 640b^4e^{7x} + 720a^4b^4e^{5x} - 1760a^2b^2e^{5x} + 1424b^4e^{5x} + 480a^4b^4e^{3x} + 120a^2b^4e^{3x} - 1120a^2b^2e^{3x} - 330a^2b^3e^{3x} + 640b^4e^{3x} + 120a^4b^4e^{1x} + 60a^3b^4e^{1x} - 240a^2b^2e^{1x} - 105a^2b^3e^{1x} + 120b^4e^{1x})/(b^5(e^{2x} + 1)^5)$

Mupad [B]

time = 7.29, size = 447, normalized size = 2.85

$$\frac{32 \exp(x)}{(5b(5\exp(2x) + 10\exp(4x) + 10\exp(6x) + 5\exp(8x) + \exp(10x) + 1)) - (\log((-a + b)^5(a - b)^5)^{1/2} + a^5\exp(x) + b^5\exp(x) + a^4b\exp(x) + a^3b^2\exp(x) - 2a^2b^3\exp(x) - 2ab^4\exp(x)) \cdot (-a + b)^5(a - b)^5)^{1/2}}{b^6} + \frac{(\log(a^5\exp(x) - (-a + b)^5(a - b)^5)^{1/2} + b^5\exp(x) + a^4b\exp(x) + a^3b^2\exp(x) - 2a^2b^3\exp(x) - 2ab^4\exp(x)) \cdot (-a + b)^5(a - b)^5)^{1/2}}{b^6} - \frac{(\exp(x)(9a^2b^2 + 16a^2b - 12a^3 - 16b^3))/(6b^4(2\exp(2x) + \exp(4x) + 1)) + (\exp(x)(7a^3b^3 - 4a^3b^3 + 8a^4 + 8b^4 - 16a^2b^2))/(4b^5(\exp(2x) + 1)) - (a \log(\exp(x) - 1i)(8a^4 + 15b^4 - 20a^2b^2)1i)/(8b^6) + (a \log(\exp(x) + 1i)(8a^4 + 15b^4 - 20a^2b^2)1i)/(8b^6) + (4\exp(x)(5a - 16b))/(5b^2(4\exp(2x) + 6\exp(4x) + 4\exp(6x) + \exp(8x) + 1)) + (2\exp(x)(20a^2 - 45ab + 28b^2))/(15b^3(3\exp(2x) + 3\exp(4x) + \exp(6x) + 1))}{(5b(5\exp(2x) + 10\exp(4x) + 10\exp(6x) + 5\exp(8x) + \exp(10x) + 1)) - (\log((-a + b)^5(a - b)^5)^{1/2} + a^5\exp(x) + b^5\exp(x) + a^4b\exp(x) + a^3b^2\exp(x) - 2a^2b^3\exp(x) - 2ab^4\exp(x)) \cdot (-a + b)^5(a - b)^5)^{1/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cosh(x)^7*(a + b*tanh(x))),x)

[Out] $\frac{32 \exp(x)}{(5b(5\exp(2x) + 10\exp(4x) + 10\exp(6x) + 5\exp(8x) + \exp(10x) + 1)) - (\log((-a + b)^5(a - b)^5)^{1/2} + a^5\exp(x) + b^5\exp(x) + a^4b\exp(x) + a^3b^2\exp(x) - 2a^2b^3\exp(x) - 2ab^4\exp(x)) \cdot (-a + b)^5(a - b)^5)^{1/2}}{b^6} + \frac{(\log(a^5\exp(x) - (-a + b)^5(a - b)^5)^{1/2} + b^5\exp(x) + a^4b\exp(x) + a^3b^2\exp(x) - 2a^2b^3\exp(x) - 2ab^4\exp(x)) \cdot (-a + b)^5(a - b)^5)^{1/2}}{b^6} - \frac{(\exp(x)(9a^2b^2 + 16a^2b - 12a^3 - 16b^3))/(6b^4(2\exp(2x) + \exp(4x) + 1)) + (\exp(x)(7a^3b^3 - 4a^3b^3 + 8a^4 + 8b^4 - 16a^2b^2))/(4b^5(\exp(2x) + 1)) - (a \log(\exp(x) - 1i)(8a^4 + 15b^4 - 20a^2b^2)1i)/(8b^6) + (a \log(\exp(x) + 1i)(8a^4 + 15b^4 - 20a^2b^2)1i)/(8b^6) + (4\exp(x)(5a - 16b))/(5b^2(4\exp(2x) + 6\exp(4x) + 4\exp(6x) + \exp(8x) + 1)) + (2\exp(x)(20a^2 - 45ab + 28b^2))/(15b^3(3\exp(2x) + 3\exp(4x) + \exp(6x) + 1))}{(5b(5\exp(2x) + 10\exp(4x) + 10\exp(6x) + 5\exp(8x) + \exp(10x) + 1)) - (\log((-a + b)^5(a - b)^5)^{1/2} + a^5\exp(x) + b^5\exp(x) + a^4b\exp(x) + a^3b^2\exp(x) - 2a^2b^3\exp(x) - 2ab^4\exp(x)) \cdot (-a + b)^5(a - b)^5)^{1/2}}$

3.110 $\int \frac{\operatorname{sech}^5(x)}{a+b \tanh(x)} dx$

Optimal. Leaf size=102

$$-\frac{a(2a^2 - 3b^2) \operatorname{ArcTan}(\sinh(x))}{2b^4} + \frac{(a^2 - b^2)^{3/2} \operatorname{ArcTan}\left(\frac{\cosh(x)(b+a \tanh(x))}{\sqrt{a^2 - b^2}}\right)}{b^4} - \frac{(a^2 - b^2) \operatorname{sech}(x)}{b^3} + \frac{\operatorname{sech}^3(x)}{3b} + \frac{a \operatorname{sech}^3(x)}{3b}$$

[Out] $-1/2*a*(2*a^2-3*b^2)*\arctan(\sinh(x))/b^4+(a^2-b^2)^{(3/2)}*\arctan(\cosh(x)*(b+a*\tanh(x))/(a^2-b^2)^{(1/2}))/b^4-(a^2-b^2)*\operatorname{sech}(x)/b^3+1/3*\operatorname{sech}(x)^3/b+1/2*a*\operatorname{sech}(x)*\tanh(x)/b^2$

Rubi [A]

time = 0.12, antiderivative size = 109, normalized size of antiderivative = 1.07, number of steps used = 9, number of rules used = 6, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.462$,

Rules used = {3591, 3567, 3853, 3855, 3590, 212}

$$-\frac{a(a^2 - b^2) \operatorname{ArcTan}(\sinh(x))}{b^4} + \frac{(a^2 - b^2)^{3/2} \operatorname{ArcTan}\left(\frac{\cosh(x)(a \tanh(x)+b)}{\sqrt{a^2 - b^2}}\right)}{b^4} - \frac{(a^2 - b^2) \operatorname{sech}(x)}{b^3} + \frac{a \operatorname{ArcTan}(\sinh(x))}{2b^2} + \frac{a \tanh(x) \operatorname{sech}(x)}{2b^2} + \frac{\operatorname{sech}^3(x)}{3b}$$

Antiderivative was successfully verified.

[In] `Int[Sech[x]^5/(a + b*Tanh[x]), x]`

[Out] $(a*\operatorname{ArcTan}[\operatorname{Sinh}[x]])/(2*b^2) - (a*(a^2 - b^2)*\operatorname{ArcTan}[\operatorname{Sinh}[x]])/b^4 + ((a^2 - b^2)^{(3/2)}*\operatorname{ArcTan}[(\operatorname{Cosh}[x]*(b + a*\operatorname{Tanh}[x]))/\operatorname{Sqrt}[a^2 - b^2]])/b^4 - ((a^2 - b^2)*\operatorname{Sech}[x])/b^3 + \operatorname{Sech}[x]^3/(3*b) + (a*\operatorname{Sech}[x]*\operatorname{Tanh}[x])/(2*b^2)$

Rule 212

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rule 3567

`Int[((d_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b*((d*Sec[e + f*x])^m/(f*m)), x] + Dist[a, Int[(d*Sec[e + f*x])^m, x], x] /; FreeQ[{a, b, d, e, f, m}, x] && (IntegerQ[2*m] | NeQ[a^2 + b^2, 0])`

Rule 3590

`Int[sec[(e_.) + (f_.)*(x_)]/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Dist[-f^(-1), Subst[Int[1/(a^2 + b^2 - x^2), x], x, (b - a*Tan[e + f*x])/Sec[e + f*x]], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a^2 + b^2, 0]`

Rule 3591

Int[((d_.)*sec[(e_.) + (f_.)*(x_)])^(m_)/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] :> Dist[-d^2/b^2, Int[(d*Sec[e + f*x])^(m - 2)*(a - b*Tan[e + f*x]), x], x] + Dist[d^2*((a^2 + b^2)/b^2), Int[(d*Sec[e + f*x])^(m - 2)/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 + b^2, 0] && IGtQ[m, 1]

Rule 3853

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Dist[b^2*((n - 2)/(n - 1)), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3855

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned}
 \int \frac{\operatorname{sech}^5(x)}{a + b \tanh(x)} dx &= \frac{\int \operatorname{sech}^3(x)(a - b \tanh(x)) dx}{b^2} - \frac{(a^2 - b^2) \int \frac{\operatorname{sech}^3(x)}{a + b \tanh(x)} dx}{b^2} \\
 &= \frac{\operatorname{sech}^3(x)}{3b} + \frac{a \int \operatorname{sech}^3(x) dx}{b^2} - \frac{(a^2 - b^2) \int \operatorname{sech}(x)(a - b \tanh(x)) dx}{b^4} + \frac{(a^2 - b^2)^2 \int \frac{1}{a + b \tanh(x)} dx}{b^4} \\
 &= -\frac{(a^2 - b^2) \operatorname{sech}(x)}{b^3} + \frac{\operatorname{sech}^3(x)}{3b} + \frac{a \operatorname{sech}(x) \tanh(x)}{2b^2} + \frac{a \int \operatorname{sech}(x) dx}{2b^2} - \frac{(a(a^2 - b^2))}{b} \\
 &= \frac{a \tan^{-1}(\sinh(x))}{2b^2} - \frac{a(a^2 - b^2) \tan^{-1}(\sinh(x))}{b^4} + \frac{(a^2 - b^2)^{3/2} \tan^{-1}\left(\frac{\cosh(x)(b + a \tanh(x))}{\sqrt{a^2 - b^2}}\right)}{b^4}
 \end{aligned}$$

Mathematica [A]

time = 0.19, size = 116, normalized size = 1.14

$$\frac{-6\left(a(2a^2 - 3b^2) \operatorname{ArcTan}\left(\tanh\left(\frac{x}{2}\right)\right) + 2\sqrt{a-b}\sqrt{a+b}(-a^2 + b^2) \operatorname{ArcTan}\left(\frac{b+a \tanh\left(\frac{x}{2}\right)}{\sqrt{a-b}\sqrt{a+b}}\right)\right) + 2b^3 \operatorname{sech}^3(x) + 3b \operatorname{sech}(x)(-2a^2 + 2b^2 + ab \tanh(x))}{6b^4}$$

Antiderivative was successfully verified.

[In] Integrate[Sech[x]^5/(a + b*Tanh[x]), x]

[Out] $(-6*(a*(2*a^2 - 3*b^2)*\text{ArcTan}[\text{Tanh}[x/2]] + 2*\text{Sqrt}[a - b]*\text{Sqrt}[a + b]*(-a^2 + b^2)*\text{ArcTan}[(b + a*\text{Tanh}[x/2])]/(\text{Sqrt}[a - b]*\text{Sqrt}[a + b]))) + 2*b^3*\text{Sech}[x]^3 + 3*b*\text{Sech}[x]*(-2*a^2 + 2*b^2 + a*b*\text{Tanh}[x]))/(6*b^4)$

Maple [A]

time = 0.99, size = 164, normalized size = 1.61

method	result
default	$\frac{2(a^4 - 2a^2b^2 + b^4) \arctan\left(\frac{2a \tanh\left(\frac{x}{2}\right) + 2b}{2\sqrt{a^2 - b^2}}\right)}{b^4 \sqrt{a^2 - b^2}} - \frac{2\left(\frac{ab^2 \tanh^5\left(\frac{x}{2}\right)}{2} + (a^2b - 2b^3) \tanh^4\left(\frac{x}{2}\right) + (2a^2b - 2b^3) \tanh^2\left(\frac{x}{2}\right) - \frac{ab^2 \tanh\left(\frac{x}{2}\right)}{2} + a^2\right)}{(\tanh^2\left(\frac{x}{2}\right) + 1)^3 b^4}$
risch	$-\frac{e^x(6a^2e^{4x} - 3abe^{4x} - 6b^2e^{4x} + 12a^2e^{2x} - 20b^2e^{2x} + 6a^2 + 3ab - 6b^2)}{3b^3(1+e^{2x})^3} + \frac{\sqrt{-a^2 + b^2} \ln\left(e^x + \frac{\sqrt{-a^2 + b^2}}{a+b}\right)}{b^4} a^2 - \frac{\sqrt{-a^2 + b^2}}{b^4}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sech(x)^5/(a+b*tanh(x)),x,method=_RETURNVERBOSE)`

[Out] $2*(a^4 - 2*a^2*b^2 + b^4)/b^4/(a^2 - b^2)^{(1/2)}*\arctan(1/2*(2*a*\tanh(1/2*x) + 2*b)/(a^2 - b^2)^{(1/2)}) - 2/b^4*((1/2*a*b^2*\tanh(1/2*x)^5 + (a^2*b - 2*b^3)*\tanh(1/2*x)^4 + (2*a^2*b - 2*b^3)*\tanh(1/2*x)^2 - 1/2*a*b^2*\tanh(1/2*x) + a^2*b - 4/3*b^3)/(\tanh(1/2*x)^2 + 1)^3 + 1/2*a*(2*a^2 - 3*b^2)*\arctan(\tanh(1/2*x))$

Maxima [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sech(x)^5/(a+b*tanh(x)),x, algorithm="maxima")`

[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see 'assume?' for more details)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 994 vs. 2(92) = 184.

time = 0.47, size = 2043, normalized size = 20.03

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sech(x)^5/(a+b*tanh(x)),x, algorithm="fricas")`

```
[Out] [-1/3*(3*(2*a^2*b - a*b^2 - 2*b^3)*cosh(x)^5 + 15*(2*a^2*b - a*b^2 - 2*b^3)
*cosh(x)*sinh(x)^4 + 3*(2*a^2*b - a*b^2 - 2*b^3)*sinh(x)^5 + 4*(3*a^2*b - 5
*b^3)*cosh(x)^3 + 2*(6*a^2*b - 10*b^3 + 15*(2*a^2*b - a*b^2 - 2*b^3)*cosh(x
)^2)*sinh(x)^3 + 6*(5*(2*a^2*b - a*b^2 - 2*b^3)*cosh(x)^3 + 2*(3*a^2*b - 5*
b^3)*cosh(x))*sinh(x)^2 + 3*((a^2 - b^2)*cosh(x)^6 + 6*(a^2 - b^2)*cosh(x)*
sinh(x)^5 + (a^2 - b^2)*sinh(x)^6 + 3*(a^2 - b^2)*cosh(x)^4 + 3*(5*(a^2 - b
^2)*cosh(x)^2 + a^2 - b^2)*sinh(x)^4 + 4*(5*(a^2 - b^2)*cosh(x)^3 + 3*(a^2
- b^2)*cosh(x))*sinh(x)^3 + 3*(a^2 - b^2)*cosh(x)^2 + 3*(5*(a^2 - b^2)*cosh
(x)^4 + 6*(a^2 - b^2)*cosh(x)^2 + a^2 - b^2)*sinh(x)^2 + a^2 - b^2 + 6*((a^
2 - b^2)*cosh(x)^5 + 2*(a^2 - b^2)*cosh(x)^3 + (a^2 - b^2)*cosh(x))*sinh(x)
)*sqrt(-a^2 + b^2)*log(((a + b)*cosh(x)^2 + 2*(a + b)*cosh(x)*sinh(x) + (a
+ b)*sinh(x)^2 - 2*sqrt(-a^2 + b^2)*(cosh(x) + sinh(x)) - a + b)/((a + b)*c
osh(x)^2 + 2*(a + b)*cosh(x)*sinh(x) + (a + b)*sinh(x)^2 + a - b)) + 3*((2*
a^3 - 3*a*b^2)*cosh(x)^6 + 6*(2*a^3 - 3*a*b^2)*cosh(x)*sinh(x)^5 + (2*a^3 -
3*a*b^2)*sinh(x)^6 + 3*(2*a^3 - 3*a*b^2)*cosh(x)^4 + 3*(2*a^3 - 3*a*b^2 +
5*(2*a^3 - 3*a*b^2)*cosh(x)^2)*sinh(x)^4 + 4*(5*(2*a^3 - 3*a*b^2)*cosh(x)^3
+ 3*(2*a^3 - 3*a*b^2)*cosh(x))*sinh(x)^3 + 2*a^3 - 3*a*b^2 + 3*(2*a^3 - 3*
a*b^2)*cosh(x)^2 + 3*(5*(2*a^3 - 3*a*b^2)*cosh(x)^4 + 2*a^3 - 3*a*b^2 + 6*(
2*a^3 - 3*a*b^2)*cosh(x)^2)*sinh(x)^2 + 6*((2*a^3 - 3*a*b^2)*cosh(x)^5 + 2*
(2*a^3 - 3*a*b^2)*cosh(x)^3 + (2*a^3 - 3*a*b^2)*cosh(x))*sinh(x))*arctan(co
sh(x) + sinh(x)) + 3*(2*a^2*b + a*b^2 - 2*b^3)*cosh(x) + 3*(5*(2*a^2*b - a*
b^2 - 2*b^3)*cosh(x)^4 + 2*a^2*b + a*b^2 - 2*b^3 + 4*(3*a^2*b - 5*b^3)*cosh
(x)^2)*sinh(x))/(b^4*cosh(x)^6 + 6*b^4*cosh(x)*sinh(x)^5 + b^4*sinh(x)^6 +
3*b^4*cosh(x)^4 + 3*b^4*cosh(x)^2 + 3*(5*b^4*cosh(x)^2 + b^4)*sinh(x)^4 + b
^4 + 4*(5*b^4*cosh(x)^3 + 3*b^4*cosh(x))*sinh(x)^3 + 3*(5*b^4*cosh(x)^4 + 6
*b^4*cosh(x)^2 + b^4)*sinh(x)^2 + 6*(b^4*cosh(x)^5 + 2*b^4*cosh(x)^3 + b^4*
cosh(x))*sinh(x)), -1/3*(3*(2*a^2*b - a*b^2 - 2*b^3)*cosh(x)^5 + 15*(2*a^2*
b - a*b^2 - 2*b^3)*cosh(x)*sinh(x)^4 + 3*(2*a^2*b - a*b^2 - 2*b^3)*sinh(x)^
5 + 4*(3*a^2*b - 5*b^3)*cosh(x)^3 + 2*(6*a^2*b - 10*b^3 + 15*(2*a^2*b - a*b
^2 - 2*b^3)*cosh(x)^2)*sinh(x)^3 + 6*(5*(2*a^2*b - a*b^2 - 2*b^3)*cosh(x)^3
+ 2*(3*a^2*b - 5*b^3)*cosh(x))*sinh(x)^2 + 6*((a^2 - b^2)*cosh(x)^6 + 6*(a
^2 - b^2)*cosh(x)*sinh(x)^5 + (a^2 - b^2)*sinh(x)^6 + 3*(a^2 - b^2)*cosh(x)
^4 + 3*(5*(a^2 - b^2)*cosh(x)^2 + a^2 - b^2)*sinh(x)^4 + 4*(5*(a^2 - b^2)*c
osh(x)^3 + 3*(a^2 - b^2)*cosh(x))*sinh(x)^3 + 3*(a^2 - b^2)*cosh(x)^2 + 3*(
5*(a^2 - b^2)*cosh(x)^4 + 6*(a^2 - b^2)*cosh(x)^2 + a^2 - b^2)*sinh(x)^2 +
a^2 - b^2 + 6*((a^2 - b^2)*cosh(x)^5 + 2*(a^2 - b^2)*cosh(x)^3 + (a^2 - b^2
)*cosh(x))*sinh(x))*sqrt(a^2 - b^2)*arctan(sqrt(a^2 - b^2)/((a + b)*cosh(x)
+ (a + b)*sinh(x))) + 3*((2*a^3 - 3*a*b^2)*cosh(x)^6 + 6*(2*a^3 - 3*a*b^2)
*cosh(x)*sinh(x)^5 + (2*a^3 - 3*a*b^2)*sinh(x)^6 + 3*(2*a^3 - 3*a*b^2)*cosh
(x)^4 + 3*(2*a^3 - 3*a*b^2 + 5*(2*a^3 - 3*a*b^2)*cosh(x)^2)*sinh(x)^4 + 4*(
5*(2*a^3 - 3*a*b^2)*cosh(x)^3 + 3*(2*a^3 - 3*a*b^2)*cosh(x))*sinh(x)^3 + 2*
a^3 - 3*a*b^2 + 3*(2*a^3 - 3*a*b^2)*cosh(x)^2 + 3*(5*(2*a^3 - 3*a*b^2)*cosh
(x)^4 + 2*a^3 - 3*a*b^2 + 6*(2*a^3 - 3*a*b^2)*cosh(x)^2)*sinh(x)^2 + 6*((2*
a^3 - 3*a*b^2)*cosh(x)^5 + 2*(2*a^3 - 3*a*b^2)*cosh(x)^3 + (2*a^3 - 3*a*b^2
)*cosh(x))*sinh(x))*arctan(cosh(x) + sinh(x)) + 3*(2*a^2*b + a*b^2 - 2*b^3)
```

cosh(x) + 3(5*(2*a^2*b - a*b^2 - 2*b^3)*cosh(x)^4 + 2*a^2*b + a*b^2 - 2*b^3 + 4*(3*a^2*b - 5*b^3)*cosh(x)^2)*sinh(x))/(b^4*cosh(x)^6 + 6*b^4*cosh(x)*sinh(x)^5 + b^4*sinh(x)^6 + 3*b^4*cosh(x)^4 + 3*b^4*cosh(x)^2 + 3*(5*b^4*cosh(x)^2 + b^4)*sinh(x)^4 + b^4 + 4*(5*b^4*cosh(x)^3 + 3*b^4*cosh(x))*sinh(x)^3 + 3*(5*b^4*cosh(x)^4 + 6*b^4*cosh(x)^2 + b^4)*sinh(x)^2 + 6*(b^4*cosh(x)^5 + 2*b^4*cosh(x)^3 + b^4*cosh(x))*sinh(x))]

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{sech}^5(x)}{a + b \tanh(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)**5/(a+b*tanh(x)),x)

[Out] Integral(sech(x)**5/(a + b*tanh(x)), x)

Giac [A]

time = 0.42, size = 152, normalized size = 1.49

$$-\frac{(2a^3 - 3ab^2)\arctan(e^x)}{b^4} + \frac{2(a^4 - 2a^2b^2 + b^4)\arctan\left(\frac{ae^x + be^x}{\sqrt{a^2 - b^2}}\right)}{\sqrt{a^2 - b^2}b^4} - \frac{6a^2e^{5x} - 3abe^{5x} - 6b^2e^{5x} + 12a^2e^{3x} - 20b^2e^{3x} + 6a^2e^x + 3abe^x - 6b^2e^x}{3b^3(e^{2x} + 1)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)^5/(a+b*tanh(x)),x, algorithm="giac")

[Out] -(2*a^3 - 3*a*b^2)*arctan(e^x)/b^4 + 2*(a^4 - 2*a^2*b^2 + b^4)*arctan((a*e^x + b*e^x)/sqrt(a^2 - b^2))/(sqrt(a^2 - b^2)*b^4) - 1/3*(6*a^2*e^(5*x) - 3*a*b*e^(5*x) - 6*b^2*e^(5*x) + 12*a^2*e^(3*x) - 20*b^2*e^(3*x) + 6*a^2*e^x + 3*a*b*e^x - 6*b^2*e^x)/(b^3*(e^(2*x) + 1)^3)

Mupad [B]

time = 5.22, size = 265, normalized size = 2.60

$$\frac{\ln\left(\frac{\sqrt{-(a+b)^2(a-b)^2 + a^2e^x - b^2e^x - ab^2e^x + a^2be^x}}{a}\sqrt{-(a+b)^2(a-b)^2}\right)}{3b(3e^{2x} + 3e^{4x} + e^{6x} + 1)} - \frac{\ln\left(\frac{\sqrt{-(a+b)^2(a-b)^2 - a^2e^x + b^2e^x + ab^2e^x - a^2be^x}}{a}\sqrt{-(a+b)^2(a-b)^2}\right)}{3b(3e^{2x} + 3e^{4x} + 1)} - \frac{2e^x(3a - 4b)}{3b^2(2e^{2x} + e^{4x} + 1)} + \frac{e^x(-2a^2 + ab + 2b^2)}{b^2(e^{2x} + 1)} + \frac{a \ln(e^x - 1)(2a^2 - 3b^2)}{2b^4} - \frac{a \ln(e^x + 1)(2a^2 - 3b^2)}{2b^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cosh(x)^5*(a + b*tanh(x))),x)

[Out] (log((-a + b)^3*(a - b)^3)^(1/2) + a^3*exp(x) - b^3*exp(x) - a*b^2*exp(x) + a^2*b*exp(x))*(-a + b)^3*(a - b)^3)^(1/2)/b^4 - (8*exp(x))/(3*b*(3*exp(2*x) + 3*exp(4*x) + exp(6*x) + 1)) - (log((-a + b)^3*(a - b)^3)^(1/2) - a^3*exp(x) + b^3*exp(x) + a*b^2*exp(x) - a^2*b*exp(x))*(-a + b)^3*(a - b)^3)^(1/2)/b^4 - (2*exp(x)*(3*a - 4*b))/(3*b^2*(2*exp(2*x) + exp(4*x) + 1)) + (exp(x)*(a*b - 2*a^2 + 2*b^2))/(b^3*(exp(2*x) + 1)) + (a*log(exp(x) - 1i)*(2*a^2 - 3*b^2)*1i)/(2*b^4) - (a*log(exp(x) + 1i)*(2*a^2 - 3*b^2)*1i)/(2*b^4)

3.111 $\int \frac{\operatorname{sech}^3(x)}{a+b \tanh(x)} dx$

Optimal. Leaf size=56

$$\frac{a \operatorname{ArcTan}(\sinh(x))}{b^2} - \frac{\sqrt{a^2 - b^2} \operatorname{ArcTan}\left(\frac{\cosh(x)(b+a \tanh(x))}{\sqrt{a^2 - b^2}}\right)}{b^2} + \frac{\operatorname{sech}(x)}{b}$$

[Out] a*arctan(sinh(x))/b^2+sech(x)/b-arctan(cosh(x)*(b+a*tanh(x)))/(a^2-b^2)^(1/2))* (a^2-b^2)^(1/2)/b^2

Rubi [A]

time = 0.06, antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {3591, 3567, 3855, 3590, 212}

$$-\frac{\sqrt{a^2 - b^2} \operatorname{ArcTan}\left(\frac{\cosh(x)(a \tanh(x)+b)}{\sqrt{a^2 - b^2}}\right)}{b^2} + \frac{a \operatorname{ArcTan}(\sinh(x))}{b^2} + \frac{\operatorname{sech}(x)}{b}$$

Antiderivative was successfully verified.

[In] Int[Sech[x]^3/(a + b*Tanh[x]),x]

[Out] (a*ArcTan[Sinh[x]])/b^2 - (Sqrt[a^2 - b^2]*ArcTan[(Cosh[x]*(b + a*Tanh[x]))/Sqrt[a^2 - b^2]])/b^2 + Sech[x]/b

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 3567

Int[((d_)*sec[(e_) + (f_)*(x_)]^(m_))*((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[b*((d*Sec[e + f*x])^m/(f*m)), x] + Dist[a, Int[(d*Sec[e + f*x])^m, x], x] /; FreeQ[{a, b, d, e, f, m}, x] && (IntegerQ[2*m] | NeQ[a^2 + b^2, 0])

Rule 3590

Int[sec[(e_) + (f_)*(x_)]/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Dist[-f^(-1), Subst[Int[1/(a^2 + b^2 - x^2), x], x, (b - a*Tan[e + f*x])/Sec[e + f*x]], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a^2 + b^2, 0]

Rule 3591

```
Int[((d_.)*sec[(e_.) + (f_.)*(x_)])^(m_)/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Dist[-d^2/b^2, Int[(d*Sec[e + f*x])^(m - 2)*(a - b*Tan[e + f*x]), x], x] + Dist[d^2*((a^2 + b^2)/b^2), Int[(d*Sec[e + f*x])^(m - 2)/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 + b^2, 0] && IGtQ[m, 1]
```

Rule 3855

```
Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned} \int \frac{\operatorname{sech}^3(x)}{a + b \tanh(x)} dx &= \frac{\int \operatorname{sech}(x)(a - b \tanh(x)) dx}{b^2} - \frac{(a^2 - b^2) \int \frac{\operatorname{sech}(x)}{a + b \tanh(x)} dx}{b^2} \\ &= \frac{\operatorname{sech}(x)}{b} + \frac{a \int \operatorname{sech}(x) dx}{b^2} - \frac{(i(a^2 - b^2)) \operatorname{Subst}\left(\int \frac{1}{a^2 - b^2 - x^2} dx, x, \cosh(x)(-ib - ia \tanh(x))\right)}{b^2} \\ &= \frac{a \tan^{-1}(\sinh(x))}{b^2} - \frac{\sqrt{a^2 - b^2} \tan^{-1}\left(\frac{\cosh(x)(b + a \tanh(x))}{\sqrt{a^2 - b^2}}\right)}{b^2} + \frac{\operatorname{sech}(x)}{b} \end{aligned}$$

Mathematica [A]

time = 0.06, size = 65, normalized size = 1.16

$$\frac{2a \operatorname{ArcTan}\left(\tanh\left(\frac{x}{2}\right)\right) - 2\sqrt{a-b}\sqrt{a+b} \operatorname{ArcTan}\left(\frac{b+a \tanh\left(\frac{x}{2}\right)}{\sqrt{a-b}\sqrt{a+b}}\right) + b \operatorname{sech}(x)}{b^2}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sech[x]^3/(a + b*Tanh[x]), x]
```

```
[Out] (2*a*ArcTan[Tanh[x/2]] - 2*Sqrt[a - b]*Sqrt[a + b]*ArcTan[(b + a*Tanh[x/2])/(Sqrt[a - b]*Sqrt[a + b])] + b*Sech[x])/b^2
```

Maple [A]

time = 0.85, size = 77, normalized size = 1.38

method	result
default	$\frac{2(-a^2+b^2) \arctan\left(\frac{2a \tanh\left(\frac{x}{2}\right)+2b}{2\sqrt{a^2-b^2}}\right)}{b^2\sqrt{a^2-b^2}} + \frac{\frac{2b}{\tanh^2\left(\frac{x}{2}\right)+1} + 2a \arctan\left(\tanh\left(\frac{x}{2}\right)\right)}{b^2}$

risch	$\frac{2e^x}{b(1+e^{2x})} + \frac{ia \ln(e^x+i)}{b^2} - \frac{ia \ln(e^x-i)}{b^2} + \frac{\sqrt{-a^2+b^2} \ln\left(e^x - \frac{\sqrt{-a^2+b^2}}{a+b}\right)}{b^2} - \frac{\sqrt{-a^2+b^2} \ln\left(e^x + \frac{\sqrt{-a^2+b^2}}{a+b}\right)}{b^2}$
-------	--

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sech(x)^3/(a+b*tanh(x)),x,method=_RETURNVERBOSE)`

[Out] $2*(-a^2+b^2)/b^2/(a^2-b^2)^{(1/2)}*\arctan(1/2*(2*a*\tanh(1/2*x)+2*b)/(a^2-b^2)^{(1/2)})+2/b^2*(b/(\tanh(1/2*x)^2+1)+a*\arctan(\tanh(1/2*x)))$

Maxima [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sech(x)^3/(a+b*tanh(x)),x, algorithm="maxima")`

[Out] Exception raised: ValueError >> Computation failed since Maxima requested a dditional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see 'assume?' for more de

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 126 vs. 2(52) = 104.

time = 0.36, size = 309, normalized size = 5.52

$$\frac{\sqrt{-a^2+b^2}(\cosh(x)^2 + 2\cosh(x)\sinh(x) + \sinh(x)^2 + 1) \log\left(\frac{\sqrt{-a^2+b^2}(\cosh(x)^2 + 2\cosh(x)\sinh(x) + \sinh(x)^2 + 1) \arctan\left(\frac{\sqrt{-a^2+b^2}}{a+\sinh(x)}\right) + (a\cosh(x)^2 + 2a\cosh(x)\sinh(x) + a\sinh(x)^2 + a)\arctan(\cosh(x) + \sinh(x)) + 2b\cosh(x) + 2b\sinh(x)}{b^2\cosh(x)^2 + 2b^2\cosh(x)\sinh(x) + b^2\sinh(x)^2 + b^2}\right) + (a\cosh(x)^2 + 2a\cosh(x)\sinh(x) + a\sinh(x)^2 + a)\arctan(\cosh(x) + \sinh(x)) + b\cosh(x) + b\sinh(x)}{b^2\cosh(x)^2 + 2b^2\cosh(x)\sinh(x) + b^2\sinh(x)^2 + b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sech(x)^3/(a+b*tanh(x)),x, algorithm="fricas")`

[Out] $[(\sqrt{-a^2+b^2})*(\cosh(x)^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2 + 1)*\log(((a + b)*\cosh(x)^2 + 2*(a + b)*\cosh(x)*\sinh(x) + (a + b)*\sinh(x)^2 - 2*\sqrt{-a^2 + b^2}*(\cosh(x) + \sinh(x)) - a + b)/((a + b)*\cosh(x)^2 + 2*(a + b)*\cosh(x)*\sinh(x) + (a + b)*\sinh(x)^2 + a - b)) + 2*(a*\cosh(x)^2 + 2*a*\cosh(x)*\sinh(x) + a*\sinh(x)^2 + a)*\arctan(\cosh(x) + \sinh(x)) + 2*b*\cosh(x) + 2*b*\sinh(x)]/(b^2*\cosh(x)^2 + 2*b^2*\cosh(x)*\sinh(x) + b^2*\sinh(x)^2 + b^2), 2*(\sqrt{a^2 - b^2}*(\cosh(x)^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2 + 1)*\arctan(\sqrt{a^2 - b^2}/((a + b)*\cosh(x) + (a + b)*\sinh(x))) + (a*\cosh(x)^2 + 2*a*\cosh(x)*\sinh(x) + a*\sinh(x)^2 + a)*\arctan(\cosh(x) + \sinh(x)) + b*\cosh(x) + b*\sinh(x))/(b^2*\cosh(x)^2 + 2*b^2*\cosh(x)*\sinh(x) + b^2*\sinh(x)^2 + b^2)]$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{sech}^3(x)}{a + b \tanh(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)**3/(a+b*tanh(x)),x)

[Out] Integral(sech(x)**3/(a + b*tanh(x)), x)

Giac [A]

time = 0.41, size = 63, normalized size = 1.12

$$\frac{2a \arctan(e^x)}{b^2} - \frac{2\sqrt{a^2 - b^2} \arctan\left(\frac{ae^x + be^x}{\sqrt{a^2 - b^2}}\right)}{b^2} + \frac{2e^x}{b(e^{2x} + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)^3/(a+b*tanh(x)),x, algorithm="giac")

[Out] 2*a*arctan(e^x)/b^2 - 2*sqrt(a^2 - b^2)*arctan((a*e^x + b*e^x)/sqrt(a^2 - b^2))/b^2 + 2*e^x/(b*(e^(2*x) + 1))

Mupad [B]

time = 3.81, size = 119, normalized size = 2.12

$$\frac{\ln(ae^x + be^x - \sqrt{b^2 - a^2}) \sqrt{-(a+b)(a-b)}}{b^2} - \frac{\ln(ae^x + be^x + \sqrt{b^2 - a^2}) \sqrt{-(a+b)(a-b)}}{b^2} + \frac{2e^x}{b(e^{2x} + 1)} - \frac{a \ln(e^x - i) \operatorname{li}}{b^2} + \frac{a \ln(e^x + i) \operatorname{li}}{b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cosh(x)^3*(a + b*tanh(x))),x)

[Out] (a*log(exp(x) + 1i)*1i)/b^2 - (a*log(exp(x) - 1i)*1i)/b^2 - (log(a*exp(x) + b*exp(x) + (b^2 - a^2)^(1/2))*(-(a + b)*(a - b))^(1/2))/b^2 + (log(a*exp(x) + b*exp(x) - (b^2 - a^2)^(1/2))*(-(a + b)*(a - b))^(1/2))/b^2 + (2*exp(x))/(b*(exp(2*x) + 1))

3.112 $\int \frac{\operatorname{sech}(x)}{a+b \tanh(x)} dx$

Optimal. Leaf size=37

$$\frac{\operatorname{ArcTan}\left(\frac{\cosh(x)(b+a \tanh(x))}{\sqrt{a^2-b^2}}\right)}{\sqrt{a^2-b^2}}$$

[Out] $\arctan(\cosh(x)*(b+a*\tanh(x)))/(a^2-b^2)^{(1/2)}/(a^2-b^2)^{(1/2)}$

Rubi [A]

time = 0.02, antiderivative size = 37, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {3590, 212}

$$\frac{\operatorname{ArcTan}\left(\frac{\cosh(x)(a \tanh(x)+b)}{\sqrt{a^2-b^2}}\right)}{\sqrt{a^2-b^2}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sech}[x]/(a + b*\text{Tanh}[x]), x]$

[Out] $\text{ArcTan}[(\text{Cosh}[x]*(b + a*\text{Tanh}[x]))/\text{Sqrt}[a^2 - b^2]]/\text{Sqrt}[a^2 - b^2]$

Rule 212

$\text{Int}[(a_+ + (b_-)*(x_-)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{NegQ}[a/b] \&\& (\text{GtQ}[a, 0] \parallel \text{LtQ}[b, 0])$

Rule 3590

$\text{Int}[\text{sec}[(e_+) + (f_-)*(x_-)]/((a_+) + (b_-)*\tan[(e_+) + (f_-)*(x_-)]), x_Symbol] \rightarrow \text{Dist}[-f^{-1}, \text{Subst}[\text{Int}[1/(a^2 + b^2 - x^2), x], x, (b - a*\tan[e + f*x])/\text{Sec}[e + f*x]], x] /; \text{FreeQ}[\{a, b, e, f\}, x] \&\& \text{NeQ}[a^2 + b^2, 0]$

Rubi steps

$$\begin{aligned} \int \frac{\operatorname{sech}(x)}{a+b \tanh(x)} dx &= i \operatorname{Subst}\left(\int \frac{1}{a^2-b^2-x^2} dx, x, \cosh(x)(-ib-ia \tanh(x))\right) \\ &= \frac{\tan^{-1}\left(\frac{\cosh(x)(b+a \tanh(x))}{\sqrt{a^2-b^2}}\right)}{\sqrt{a^2-b^2}} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 46, normalized size = 1.24

$$\frac{2\text{ArcTan}\left(\frac{b+a\tanh\left(\frac{x}{2}\right)}{\sqrt{a-b}\sqrt{a+b}}\right)}{\sqrt{a-b}\sqrt{a+b}}$$

Antiderivative was successfully verified.

`[In] Integrate[Sech[x]/(a + b*Tanh[x]), x]``[Out] (2*ArcTan[(b + a*Tanh[x/2])/(Sqrt[a - b]*Sqrt[a + b])])/(Sqrt[a - b]*Sqrt[a + b])`**Maple [A]**

time = 0.45, size = 39, normalized size = 1.05

method	result	size
default	$\frac{2\arctan\left(\frac{2a\tanh\left(\frac{x}{2}\right)+2b}{\sqrt{a^2-b^2}}\right)}{\sqrt{a^2-b^2}}$	39
risch	$-\frac{\ln\left(e^x - \frac{a-b}{\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2}} + \frac{\ln\left(e^x + \frac{a-b}{\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2}}$	70

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sech(x)/(a+b*tanh(x)),x,method=_RETURNVERBOSE)``[Out] 2/(a^2-b^2)^(1/2)*arctan(1/2*(2*a*tanh(1/2*x)+2*b)/(a^2-b^2)^(1/2))`**Maxima [F(-2)]**

time = 0.00, size = 0, normalized size = 0.00

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sech(x)/(a+b*tanh(x)),x, algorithm="maxima")``[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see 'assume?' for more details)`**Fricas [A]**

time = 0.44, size = 148, normalized size = 4.00

$$\left[-\frac{\sqrt{-a^2+b^2} \log\left(\frac{(a+b)\cosh(x)^2+2(a+b)\cosh(x)\sinh(x)+(a+b)\sinh(x)^2-2\sqrt{-a^2+b^2}(\cosh(x)+\sinh(x))-a+b}{(a+b)\cosh(x)^2+2(a+b)\cosh(x)\sinh(x)+(a+b)\sinh(x)^2+a-b}\right)}{a^2-b^2}, -\frac{2\arctan\left(\frac{\sqrt{a^2-b^2}}{(a+b)\cosh(x)+(a+b)\sinh(x)}\right)}{\sqrt{a^2-b^2}} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)/(a+b*tanh(x)),x, algorithm="fricas")

[Out] $[-\sqrt{-a^2 + b^2} \log((a + b) \cosh(x)^2 + 2(a + b) \cosh(x) \sinh(x) + (a + b) \sinh(x)^2 - 2\sqrt{-a^2 + b^2} (\cosh(x) + \sinh(x)) - a + b) / ((a + b) \cosh(x)^2 + 2(a + b) \cosh(x) \sinh(x) + (a + b) \sinh(x)^2 + a - b)) / (a^2 - b^2), -2 \arctan(\sqrt{a^2 - b^2} / ((a + b) \cosh(x) + (a + b) \sinh(x))) / \sqrt{a^2 - b^2}]$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\operatorname{sech}(x)}{a + b \tanh(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)/(a+b*tanh(x)),x)

[Out] Integral(sech(x)/(a + b*tanh(x)), x)

Giac [A]

time = 0.40, size = 35, normalized size = 0.95

$$\frac{2 \arctan\left(\frac{ae^x + be^x}{\sqrt{a^2 - b^2}}\right)}{\sqrt{a^2 - b^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sech(x)/(a+b*tanh(x)),x, algorithm="giac")

[Out] $2 \arctan((a * e^x + b * e^x) / \sqrt{a^2 - b^2}) / \sqrt{a^2 - b^2}$

Mupad [B]

time = 0.13, size = 35, normalized size = 0.95

$$\frac{2 \operatorname{atan}\left(\frac{e^x \sqrt{a^2 - b^2}}{a - b}\right)}{\sqrt{a^2 - b^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cosh(x)*(a + b*tanh(x))),x)

[Out] $(2 * \operatorname{atan}((\exp(x) * (a^2 - b^2)^{(1/2)}) / (a - b))) / (a^2 - b^2)^{(1/2)}$

3.113 $\int \frac{\cosh(x)}{a+b \tanh(x)} dx$

Optimal. Leaf size=73

$$-\frac{b^2 \operatorname{ArcTan}\left(\frac{\cosh(x)(b+a \tanh(x))}{\sqrt{a^2-b^2}}\right)}{(a^2-b^2)^{3/2}} - \frac{b \cosh(x)}{a^2-b^2} + \frac{a \sinh(x)}{a^2-b^2}$$

[Out] $-b^2 \arctan(\cosh(x) * (b+a \tanh(x)) / (a^2-b^2)^{(1/2)}) / (a^2-b^2)^{(3/2)} - b \cosh(x) / (a^2-b^2) + a \sinh(x) / (a^2-b^2)$

Rubi [A]

time = 0.07, antiderivative size = 73, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.454$, Rules used = {3592, 3567, 2717, 3590, 212}

$$-\frac{b^2 \operatorname{ArcTan}\left(\frac{\cosh(x)(a \tanh(x)+b)}{\sqrt{a^2-b^2}}\right)}{(a^2-b^2)^{3/2}} + \frac{a \sinh(x)}{a^2-b^2} - \frac{b \cosh(x)}{a^2-b^2}$$

Antiderivative was successfully verified.

[In] `Int[Cosh[x]/(a + b*Tanh[x]),x]`

[Out] $-(b^2 \operatorname{ArcTan}[(\operatorname{Cosh}[x] * (b + a \operatorname{Tanh}[x])) / \operatorname{Sqrt}[a^2 - b^2]]) / (a^2 - b^2)^{(3/2)} - (b \operatorname{Cosh}[x]) / (a^2 - b^2) + (a \operatorname{Sinh}[x]) / (a^2 - b^2)$

Rule 212

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rule 2717

`Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

Rule 3567

`Int[((d_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b*((d*Sec[e + f*x])^m/(f*m)), x] + Dist[a, Int[(d*Sec[e + f*x])^m, x], x] /; FreeQ[{a, b, d, e, f, m}, x] && (IntegerQ[2*m] | NeQ[a^2 + b^2, 0])`

Rule 3590

```
Int[sec[(e_.) + (f_.)*(x_)]/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol]
:= Dist[-f^(-1), Subst[Int[1/(a^2 + b^2 - x^2), x], x, (b - a*Tan[e + f*x])/Sec[e + f*x]], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a^2 + b^2, 0]
```

Rule 3592

```
Int[((d_.)*sec[(e_.) + (f_.)*(x_)]^(m_)/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol]
:= Dist[1/(a^2 + b^2), Int[(d*Sec[e + f*x])^m*(a - b*Tan[e + f*x]), x], x] + Dist[b^2/(d^2*(a^2 + b^2)), Int[(d*Sec[e + f*x])^(m + 2)/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 + b^2, 0] && ILtQ[m, 0]
```

Rubi steps

$$\begin{aligned} \int \frac{\cosh(x)}{a + b \tanh(x)} dx &= \frac{\int \cosh(x)(a - b \tanh(x)) dx}{a^2 - b^2} - \frac{b^2 \int \frac{\operatorname{sech}(x)}{a + b \tanh(x)} dx}{a^2 - b^2} \\ &= -\frac{b \cosh(x)}{a^2 - b^2} + \frac{a \int \cosh(x) dx}{a^2 - b^2} - \frac{(ib^2) \operatorname{Subst}\left(\int \frac{1}{a^2 - b^2 - x^2} dx, x, \cosh(x)(-ib - ia \tanh(x))\right)}{a^2 - b^2} \\ &= -\frac{b^2 \tan^{-1}\left(\frac{\cosh(x)(b + a \tanh(x))}{\sqrt{a^2 - b^2}}\right)}{(a^2 - b^2)^{3/2}} - \frac{b \cosh(x)}{a^2 - b^2} + \frac{a \sinh(x)}{a^2 - b^2} \end{aligned}$$

Mathematica [A]

time = 0.16, size = 80, normalized size = 1.10

$$-\frac{2b^2 \operatorname{ArcTan}\left(\frac{b + a \tanh\left(\frac{x}{2}\right)}{\sqrt{a - b} \sqrt{a + b}}\right)}{(a - b)^{3/2} (a + b)^{3/2}} + \frac{b \cosh(x)}{-a^2 + b^2} + \frac{a \sinh(x)}{a^2 - b^2}$$

Antiderivative was successfully verified.

```
[In] Integrate[Cosh[x]/(a + b*Tanh[x]), x]
```

```
[Out] (-2*b^2*ArcTan[(b + a*Tanh[x/2])/(Sqrt[a - b]*Sqrt[a + b])]/((a - b)^(3/2)*(a + b)^(3/2)) + (b*Cosh[x])/(-a^2 + b^2) + (a*Sinh[x])/(a^2 - b^2))
```

Maple [A]

time = 0.70, size = 93, normalized size = 1.27

method	result	size
--------	--------	------

default	$-\frac{2b^2 \arctan\left(\frac{2a \tanh\left(\frac{x}{2}\right) + 2b}{2\sqrt{a^2 - b^2}}\right)}{(a-b)(a+b)\sqrt{a^2 - b^2}} - \frac{2}{(2b+2a)(\tanh\left(\frac{x}{2}\right) - 1)} - \frac{2}{(2a-2b)(\tanh\left(\frac{x}{2}\right) + 1)}$	93
risch	$\frac{e^x}{2b+2a} - \frac{e^{-x}}{2(a-b)} - \frac{b^2 \ln\left(e^x + \frac{a-b}{\sqrt{-a^2 + b^2}}\right)}{\sqrt{-a^2 + b^2}(a+b)(a-b)} + \frac{b^2 \ln\left(e^{-x} - \frac{a-b}{\sqrt{-a^2 + b^2}}\right)}{\sqrt{-a^2 + b^2}(a+b)(a-b)}$	122

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cosh(x)/(a+b*tanh(x)),x,method=_RETURNVERBOSE)`

[Out] $-2*b^2/(a-b)/(a+b)/(a^2-b^2)^{(1/2)}*\arctan(1/2*(2*a*\tanh(1/2*x)+2*b)/(a^2-b^2)^{(1/2)})-2/(2*b+2*a)/(\tanh(1/2*x)-1)-2/(2*a-2*b)/(\tanh(1/2*x)+1)$

Maxima [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cosh(x)/(a+b*tanh(x)),x, algorithm="maxima")`

[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see 'assume?' for more details)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 190 vs. 2(69) = 138.

time = 0.57, size = 435, normalized size = 5.96

$$\frac{\frac{a^2 + a^2 - a^2 - b^2 - (a^2 - a^2 - a^2 + b^2) \operatorname{cosh}(x) - 2(a^2 - a^2 - a^2 + b^2) \operatorname{cosh}(x) \operatorname{sinh}(x) - (a^2 - a^2 - a^2 + b^2) \operatorname{sinh}(x)^2 - 2(a^2 - a^2 - a^2 + b^2) \operatorname{sinh}(x) \sqrt{-a^2 + b^2} \operatorname{arctan}\left(\frac{2a \tanh\left(\frac{x}{2}\right) + 2b}{2\sqrt{a^2 - b^2}}\right)}{2(a^2 - 2a^2 + b^2) \operatorname{cosh}(x) + (a^2 - 2a^2 + b^2) \operatorname{sinh}(x)} - \frac{a^2 + a^2 - a^2 - b^2 - (a^2 - a^2 - a^2 + b^2) \operatorname{cosh}(x)^2 - 2(a^2 - a^2 - a^2 + b^2) \operatorname{cosh}(x) \operatorname{sinh}(x) - (a^2 - a^2 - a^2 + b^2) \operatorname{sinh}(x)^2 - 4(a^2 - a^2 - a^2 + b^2) \operatorname{sinh}(x) \sqrt{-a^2 + b^2} \operatorname{arctan}\left(\frac{2a \tanh\left(\frac{x}{2}\right) + 2b}{2\sqrt{a^2 - b^2}}\right)}{2(a^2 - 2a^2 + b^2) \operatorname{cosh}(x) + (a^2 - 2a^2 + b^2) \operatorname{sinh}(x)}}{2(a^2 - 2a^2 + b^2) \operatorname{cosh}(x) + (a^2 - 2a^2 + b^2) \operatorname{sinh}(x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cosh(x)/(a+b*tanh(x)),x, algorithm="fricas")`

[Out] $[-1/2*(a^3 + a^2*b - a*b^2 - b^3 - (a^3 - a^2*b - a*b^2 + b^3)*\cosh(x)^2 - 2*(a^3 - a^2*b - a*b^2 + b^3)*\cosh(x)*\sinh(x) - (a^3 - a^2*b - a*b^2 + b^3)*\sinh(x)^2 - 2*(b^2*\cosh(x) + b^2*\sinh(x))*\sqrt{-a^2 + b^2}*\log(((a + b)*\cosh(x)^2 + 2*(a + b)*\cosh(x)*\sinh(x) + (a + b)*\sinh(x)^2 - 2*\sqrt{-a^2 + b^2})*(\cosh(x) + \sinh(x)) - a + b)/((a + b)*\cosh(x)^2 + 2*(a + b)*\cosh(x)*\sinh(x) + (a + b)*\sinh(x)^2 + a - b))]/((a^4 - 2*a^2*b^2 + b^4)*\cosh(x) + (a^4 - 2*a^2*b^2 + b^4)*\sinh(x)), -1/2*(a^3 + a^2*b - a*b^2 - b^3 - (a^3 - a^2*b - a*b^2 + b^3)*\cosh(x)^2 - 2*(a^3 - a^2*b - a*b^2 + b^3)*\cosh(x)*\sinh(x) - (a^3 - a^2*b - a*b^2 + b^3)*\sinh(x)^2 - 4*(b^2*\cosh(x) + b^2*\sinh(x))*\sqrt{-a^2 + b^2}*\log(((a + b)*\cosh(x)^2 + 2*(a + b)*\cosh(x)*\sinh(x) + (a + b)*\sinh(x)^2 - 2*\sqrt{-a^2 + b^2})*(\cosh(x) + \sinh(x)) - a + b)/((a + b)*\cosh(x)^2 + 2*(a + b)*\cosh(x)*\sinh(x) + (a + b)*\sinh(x)^2 + a - b))]/((a^4 - 2*a^2*b^2 + b^4)*\cosh(x) + (a^4 - 2*a^2*b^2 + b^4)*\sinh(x))$

$$\frac{(a^2 - b^2) \arctan(\sqrt{a^2 - b^2} / ((a + b) \cosh(x) + (a + b) \sinh(x)))}{(a^4 - 2a^2b^2 + b^4) \cosh(x) + (a^4 - 2a^2b^2 + b^4) \sinh(x)}$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cosh(x)}{a + b \tanh(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cosh(x)/(a+b*tanh(x)),x)

[Out] Integral(cosh(x)/(a + b*tanh(x)), x)

Giac [A]

time = 0.41, size = 61, normalized size = 0.84

$$-\frac{2b^2 \arctan\left(\frac{ae^x + be^x}{\sqrt{a^2 - b^2}}\right)}{(a^2 - b^2)^{\frac{3}{2}}} - \frac{e^{-x}}{2(a - b)} + \frac{e^x}{2(a + b)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cosh(x)/(a+b*tanh(x)),x, algorithm="giac")

[Out] $-2b^2 \arctan((a e^x + b e^x) / \sqrt{a^2 - b^2}) / (a^2 - b^2)^{3/2} - 1/2 e^{-x} / (a - b) + 1/2 e^x / (a + b)$

Mupad [B]

time = 1.36, size = 157, normalized size = 2.15

$$\frac{e^x}{2a + 2b} - \frac{e^{-x}}{2a - 2b} - \frac{b^2 \ln\left(-\frac{2b^2}{(a+b)^{5/2} \sqrt{b-a}} - \frac{2b^2 e^x}{-a^3 - a^2 b + a b^2 + b^3}\right)}{(a+b)^{3/2} (b-a)^{3/2}} + \frac{b^2 \ln\left(\frac{2b^2}{(a+b)^{5/2} \sqrt{b-a}} - \frac{2b^2 e^x}{-a^3 - a^2 b + a b^2 + b^3}\right)}{(a+b)^{3/2} (b-a)^{3/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cosh(x)/(a + b*tanh(x)),x)

[Out] $\exp(x)/(2a + 2b) - \exp(-x)/(2a - 2b) - (b^2 \log(- (2b^2) / ((a + b)^{5/2} (b - a)^{1/2})) - (2b^2 \exp(x)) / (a b^2 - a^2 b - a^3 + b^3)) / ((a + b)^{3/2} (b - a)^{3/2}) + (b^2 \log((2b^2) / ((a + b)^{5/2} (b - a)^{1/2})) - (2b^2 \exp(x)) / (a b^2 - a^2 b - a^3 + b^3)) / ((a + b)^{3/2} (b - a)^{3/2})$

3.114 $\int \frac{\cosh^3(x)}{a+b \tanh(x)} dx$

Optimal. Leaf size=132

$$\frac{b^4 \operatorname{ArcTan}\left(\frac{\cosh(x)(b+a \tanh(x))}{\sqrt{a^2-b^2}}\right)}{(a^2-b^2)^{5/2}} + \frac{b^3 \cosh(x)}{(a^2-b^2)^2} - \frac{b \cosh^3(x)}{3(a^2-b^2)} - \frac{ab^2 \sinh(x)}{(a^2-b^2)^2} + \frac{a \sinh(x)}{a^2-b^2} + \frac{a \sinh^3(x)}{3(a^2-b^2)}$$

[Out] $b^4 \operatorname{arctan}(\cosh(x) * (b + a * \tanh(x)) / (a^2 - b^2)^{(1/2)}) / (a^2 - b^2)^{(5/2)} + b^3 * \cosh(x) / (a^2 - b^2)^2 - 1/3 * b * \cosh(x)^3 / (a^2 - b^2) - a * b^2 * \sinh(x) / (a^2 - b^2)^2 + a * \sinh(x) / (a^2 - b^2) + 1/3 * a * \sinh(x)^3 / (a^2 - b^2)$

Rubi [A]

time = 0.12, antiderivative size = 132, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 6, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.462$, Rules used = {3592, 3567, 2713, 2717, 3590, 212}

$$\frac{b^4 \operatorname{ArcTan}\left(\frac{\cosh(x)(a \tanh(x) + b)}{\sqrt{a^2-b^2}}\right)}{(a^2-b^2)^{5/2}} + \frac{a \sinh^3(x)}{3(a^2-b^2)} - \frac{ab^2 \sinh(x)}{(a^2-b^2)^2} + \frac{a \sinh(x)}{a^2-b^2} - \frac{b \cosh^3(x)}{3(a^2-b^2)} + \frac{b^3 \cosh(x)}{(a^2-b^2)^2}$$

Antiderivative was successfully verified.

[In] `Int[Cosh[x]^3/(a + b*Tanh[x]),x]`

[Out] $(b^4 * \operatorname{ArcTan}[(\operatorname{Cosh}[x] * (b + a * \operatorname{Tanh}[x])) / \operatorname{Sqrt}[a^2 - b^2]]) / (a^2 - b^2)^{(5/2)} + (b^3 * \operatorname{Cosh}[x]) / (a^2 - b^2)^2 - (b * \operatorname{Cosh}[x]^3) / (3 * (a^2 - b^2)) - (a * b^2 * \operatorname{Sinh}[x]) / (a^2 - b^2)^2 + (a * \operatorname{Sinh}[x]) / (a^2 - b^2) + (a * \operatorname{Sinh}[x]^3) / (3 * (a^2 - b^2))$

Rule 212

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rule 2713

`Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Dist[-d^(-1), Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]`

Rule 2717

`Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

Rule 3567

```
Int[((d_.)*sec[(e_.) + (f_.)*(x_)]^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b*((d*Sec[e + f*x])^m/(f*m)), x] + Dist[a, Int[(d*Sec[e + f*x])^m, x], x] /; FreeQ[{a, b, d, e, f, m}, x] && (IntegerQ[2*m] | NeQ[a^2 + b^2, 0])
```

Rule 3590

```
Int[sec[(e_.) + (f_.)*(x_)]/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Dist[-f^(-1), Subst[Int[1/(a^2 + b^2 - x^2), x], x, (b - a*Tan[e + f*x])/Sec[e + f*x]], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a^2 + b^2, 0]
```

Rule 3592

```
Int[((d_.)*sec[(e_.) + (f_.)*(x_)]^(m_)/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Dist[1/(a^2 + b^2), Int[(d*Sec[e + f*x])^m*(a - b*Tan[e + f*x]), x], x] + Dist[b^2/(d^2*(a^2 + b^2)), Int[(d*Sec[e + f*x])^(m + 2)/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 + b^2, 0] && ILtQ[m, 0]
```

Rubi steps

$$\begin{aligned} \int \frac{\cosh^3(x)}{a + b \tanh(x)} dx &= \frac{\int \cosh^3(x)(a - b \tanh(x)) dx}{a^2 - b^2} - \frac{b^2 \int \frac{\cosh(x)}{a + b \tanh(x)} dx}{a^2 - b^2} \\ &= -\frac{b \cosh^3(x)}{3(a^2 - b^2)} - \frac{b^2 \int \cosh(x)(a - b \tanh(x)) dx}{(a^2 - b^2)^2} + \frac{b^4 \int \frac{\operatorname{sech}(x)}{a + b \tanh(x)} dx}{(a^2 - b^2)^2} + \frac{a \int \cosh^3(x) dx}{a^2 - b^2} \\ &= \frac{b^3 \cosh(x)}{(a^2 - b^2)^2} - \frac{b \cosh^3(x)}{3(a^2 - b^2)} - \frac{(ab^2) \int \cosh(x) dx}{(a^2 - b^2)^2} + \frac{(ib^4) \operatorname{Subst}\left(\int \frac{1}{a^2 - b^2 - x^2} dx, x, \cosh(x)\right)}{(a^2 - b^2)^2} \\ &= \frac{b^4 \tan^{-1}\left(\frac{\cosh(x)(b + a \tanh(x))}{\sqrt{a^2 - b^2}}\right)}{(a^2 - b^2)^{5/2}} + \frac{b^3 \cosh(x)}{(a^2 - b^2)^2} - \frac{b \cosh^3(x)}{3(a^2 - b^2)} - \frac{ab^2 \sinh(x)}{(a^2 - b^2)^2} + \frac{a \sinh(x)}{a^2 - b^2} + \dots \end{aligned}$$

Mathematica [A]

time = 0.32, size = 258, normalized size = 1.95

$$\frac{24b^4\sqrt{a+b}\operatorname{ArcTan}\left(\frac{b\cosh(x)}{\sqrt{a-b}\sqrt{a+b}}\right) - 3\sqrt{a-b}b(a^2 + a^2b - 5ab^2 - 5b^3)\cosh(x) - (a-b)^{3/2}b(a+b)^2\cosh(3x) + 9a^4\sqrt{a-b}\sinh(x) + 9a^3\sqrt{a-b}b\sinh(x) - 21a^2\sqrt{a-b}b^2\sinh(x) - 21a\sqrt{a-b}b^3\sinh(x) + a^4\sqrt{a-b}\sinh(3x) + a^3\sqrt{a-b}b\sinh(3x) - a^2\sqrt{a-b}b^2\sinh(3x) - a\sqrt{a-b}b^3\sinh(3x)}{12(a-b)^{3/2}(a+b)^3}$$

Antiderivative was successfully verified.

```
[In] Integrate[Cosh[x]^3/(a + b*Tanh[x]), x]
```

```
[Out] (24*b^4*sqrt[a + b]*ArcTan[(b + a*Tanh[x/2])/(sqrt[a - b]*sqrt[a + b])]) - 3*sqrt[a - b]*b*(a^3 + a^2*b - 5*a*b^2 - 5*b^3)*Cosh[x] - (a - b)^(3/2)*b*(a
```

$$\begin{aligned} & + b)^2 \cosh[3x] + 9a^4 \sqrt{a-b} \sinh[x] + 9a^3 \sqrt{a-b} b \sinh[x] \\ & - 21a^2 \sqrt{a-b} b^2 \sinh[x] - 21a \sqrt{a-b} b^3 \sinh[x] + a^4 \sqrt{a-b} \\ & [a-b] \sinh[3x] + a^3 \sqrt{a-b} b \sinh[3x] - a^2 \sqrt{a-b} b^2 \sinh[3x] \\ & - a \sqrt{a-b} b^3 \sinh[3x]) / (12(a-b)^{5/2} (a+b)^3) \end{aligned}$$

Maple [A]

time = 0.71, size = 176, normalized size = 1.33

method	result
risch	$\frac{e^{3x}}{24a+24b} + \frac{3e^x a}{8(a+b)^2} + \frac{5e^x b}{8(a+b)^2} - \frac{3e^{-x} a}{8(a-b)^2} + \frac{5e^{-x} b}{8(a-b)^2} - \frac{e^{-3x}}{24(a-b)} - \frac{b^4 \ln\left(e^x - \frac{a-b}{\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2} (a+b)^2 (a-b)^2} + \frac{b^4 \ln\left(e^x + \frac{a-b}{\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2} (a+b)^2 (a-b)^2}$
default	$-\frac{2}{3(\tanh(\frac{x}{2})+1)^3(2a-2b)} + \frac{1}{(2a-2b)(\tanh(\frac{x}{2})+1)^2} - \frac{2a-3b}{2(a-b)^2(\tanh(\frac{x}{2})+1)} - \frac{2}{3(\tanh(\frac{x}{2})-1)^3(2b+2a)} - \frac{1}{(2b+2a)(\tanh(\frac{x}{2}))^2}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cosh(x)^3/(a+b*tanh(x)),x,method=_RETURNVERBOSE)`

[Out]
$$-2/3/(\tanh(1/2*x)+1)^3/(2*a-2*b)+1/(2*a-2*b)/(\tanh(1/2*x)+1)^2-1/2*(2*a-3*b)/(a-b)^2/(\tanh(1/2*x)+1)-2/3/(\tanh(1/2*x)-1)^3/(2*b+2*a)-1/(2*b+2*a)/(\tanh(1/2*x)-1)^2-1/2*(2*a+3*b)/(a+b)^2/(\tanh(1/2*x)-1)+2*b^4/(a-b)^2/(a+b)^2/(a^2-b^2)^{(1/2)}*\arctan(1/2*(2*a*\tanh(1/2*x)+2*b)/(a^2-b^2)^{(1/2)})$$

Maxima [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cosh(x)^3/(a+b*tanh(x)),x, algorithm="maxima")`

[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*b^2-4*a^2>0)', see 'assume?' for more details)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 908 vs. $2(124) = 248$.

time = 0.41, size = 1871, normalized size = 14.17

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cosh(x)^3/(a+b*tanh(x)),x, algorithm="fricas")`

```
[Out] [1/24*((a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*cosh(x)^6 + 6*(a
^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*cosh(x)*sinh(x)^5 + (a^5
- a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*sinh(x)^6 - a^5 - a^4*b + 2*
a^3*b^2 + 2*a^2*b^3 - a*b^4 - b^5 + 3*(3*a^5 - a^4*b - 10*a^3*b^2 + 6*a^2*b
^3 + 7*a*b^4 - 5*b^5)*cosh(x)^4 + 3*(3*a^5 - a^4*b - 10*a^3*b^2 + 6*a^2*b^3
+ 7*a*b^4 - 5*b^5 + 5*(a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*
cosh(x)^2)*sinh(x)^4 + 4*(5*(a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 -
b^5)*cosh(x)^3 + 3*(3*a^5 - a^4*b - 10*a^3*b^2 + 6*a^2*b^3 + 7*a*b^4 - 5*b
^5)*cosh(x))*sinh(x)^3 - 3*(3*a^5 + a^4*b - 10*a^3*b^2 - 6*a^2*b^3 + 7*a*b^4
+ 5*b^5)*cosh(x)^2 - 3*(3*a^5 + a^4*b - 10*a^3*b^2 - 6*a^2*b^3 + 7*a*b^4 +
5*b^5 - 5*(a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*cosh(x)^4 -
6*(3*a^5 - a^4*b - 10*a^3*b^2 + 6*a^2*b^3 + 7*a*b^4 - 5*b^5)*cosh(x)^2)*sin
h(x)^2 - 24*(b^4*cosh(x)^3 + 3*b^4*cosh(x)^2*sinh(x) + 3*b^4*cosh(x)*sinh(x)
)^2 + b^4*sinh(x)^3)*sqrt(-a^2 + b^2)*log(((a + b)*cosh(x)^2 + 2*(a + b)*co
sh(x)*sinh(x) + (a + b)*sinh(x)^2 - 2*sqrt(-a^2 + b^2)*(cosh(x) + sinh(x))
- a + b)/((a + b)*cosh(x)^2 + 2*(a + b)*cosh(x)*sinh(x) + (a + b)*sinh(x)^2
+ a - b)) + 6*((a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*cosh(x)
^5 + 2*(3*a^5 - a^4*b - 10*a^3*b^2 + 6*a^2*b^3 + 7*a*b^4 - 5*b^5)*cosh(x)^3
- (3*a^5 + a^4*b - 10*a^3*b^2 - 6*a^2*b^3 + 7*a*b^4 + 5*b^5)*cosh(x))*sinh
(x))/((a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*cosh(x)^3 + 3*(a^6 - 3*a^4*b^2 +
3*a^2*b^4 - b^6)*cosh(x)^2*sinh(x) + 3*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*
cosh(x)*sinh(x)^2 + (a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*sinh(x)^3), 1/24*((
a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*cosh(x)^6 + 6*(a^5 - a^4
*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*cosh(x)*sinh(x)^5 + (a^5 - a^4*b
- 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*sinh(x)^6 - a^5 - a^4*b + 2*a^3*b^2
+ 2*a^2*b^3 - a*b^4 - b^5 + 3*(3*a^5 - a^4*b - 10*a^3*b^2 + 6*a^2*b^3 + 7*a
*b^4 - 5*b^5)*cosh(x)^4 + 3*(3*a^5 - a^4*b - 10*a^3*b^2 + 6*a^2*b^3 + 7*a*b
^4 - 5*b^5 + 5*(a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*cosh(x)^
2)*sinh(x)^4 + 4*(5*(a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*cos
h(x)^3 + 3*(3*a^5 - a^4*b - 10*a^3*b^2 + 6*a^2*b^3 + 7*a*b^4 - 5*b^5)*cosh(
x))*sinh(x)^3 - 3*(3*a^5 + a^4*b - 10*a^3*b^2 - 6*a^2*b^3 + 7*a*b^4 + 5*b^5
)*cosh(x)^2 - 3*(3*a^5 + a^4*b - 10*a^3*b^2 - 6*a^2*b^3 + 7*a*b^4 + 5*b^5 -
5*(a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*cosh(x)^4 - 6*(3*a^5
- a^4*b - 10*a^3*b^2 + 6*a^2*b^3 + 7*a*b^4 - 5*b^5)*cosh(x)^2)*sinh(x)^2 -
48*(b^4*cosh(x)^3 + 3*b^4*cosh(x)^2*sinh(x) + 3*b^4*cosh(x)*sinh(x)^2 + b^
4*sinh(x)^3)*sqrt(a^2 - b^2)*arctan(sqrt(a^2 - b^2)/((a + b)*cosh(x) + (a +
b)*sinh(x))) + 6*((a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*cosh
(x)^5 + 2*(3*a^5 - a^4*b - 10*a^3*b^2 + 6*a^2*b^3 + 7*a*b^4 - 5*b^5)*cosh(x)
)^3 - (3*a^5 + a^4*b - 10*a^3*b^2 - 6*a^2*b^3 + 7*a*b^4 + 5*b^5)*cosh(x))*s
inh(x))/((a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*cosh(x)^3 + 3*(a^6 - 3*a^4*b^2
+ 3*a^2*b^4 - b^6)*cosh(x)^2*sinh(x) + 3*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^
6)*cosh(x)*sinh(x)^2 + (a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*sinh(x)^3)]
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cosh^3(x)}{a + b \tanh(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cosh(x)**3/(a+b*tanh(x)),x)

[Out] Integral(cosh(x)**3/(a + b*tanh(x)), x)

Giac [A]

time = 0.41, size = 162, normalized size = 1.23

$$\frac{2b^4 \arctan\left(\frac{ae^x + be^x}{\sqrt{a^2 - b^2}}\right)}{(a^4 - 2a^2b^2 + b^4)\sqrt{a^2 - b^2}} - \frac{(9ae^{(2x)} - 15be^{(2x)} + a - b)e^{(-3x)}}{24(a^2 - 2ab + b^2)} + \frac{a^2e^{(3x)} + 2abe^{(3x)} + b^2e^{(3x)} + 9a^2e^x + 24abe^x + 15b^2e^x}{24(a^3 + 3a^2b + 3ab^2 + b^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cosh(x)^3/(a+b*tanh(x)),x, algorithm="giac")

[Out] 2*b^4*arctan((a*e^x + b*e^x)/sqrt(a^2 - b^2))/((a^4 - 2*a^2*b^2 + b^4)*sqrt(a^2 - b^2)) - 1/24*(9*a*e^(2*x) - 15*b*e^(2*x) + a - b)*e^(-3*x)/(a^2 - 2*a*b + b^2) + 1/24*(a^2*e^(3*x) + 2*a*b*e^(3*x) + b^2*e^(3*x) + 9*a^2*e^x + 24*a*b*e^x + 15*b^2*e^x)/(a^3 + 3*a^2*b + 3*a*b^2 + b^3)

Mupad [B]

time = 2.25, size = 221, normalized size = 1.67

$$\frac{e^{3x}}{24a + 24b} - \frac{e^{-3x}}{24a - 24b} - \frac{e^{-x}(3a - 5b)}{8(a - b)^2} + \frac{e^x(3a + 5b)}{8(a + b)^2} - \frac{b^4 \ln\left(-\frac{2b^4 e^x}{a^5 + a^4 b - 2a^3 b^2 - 2a^2 b^3 + a b^4 + b^5} - \frac{2b^4}{(a+b)^{7/2}(b-a)^{3/2}}\right)}{(a+b)^{5/2}(b-a)^{5/2}} + \frac{b^4 \ln\left(\frac{2b^4}{(a+b)^{7/2}(b-a)^{3/2}} - \frac{2b^4 e^x}{a^5 + a^4 b - 2a^3 b^2 - 2a^2 b^3 + a b^4 + b^5}\right)}{(a+b)^{5/2}(b-a)^{5/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cosh(x)^3/(a + b*tanh(x)),x)

[Out] exp(3*x)/(24*a + 24*b) - exp(-3*x)/(24*a - 24*b) - (exp(-x)*(3*a - 5*b))/(8*(a - b)^2) + (exp(x)*(3*a + 5*b))/(8*(a + b)^2) - (b^4*log(- (2*b^4*exp(x))/(a*b^4 + a^4*b + a^5 + b^5 - 2*a^2*b^3 - 2*a^3*b^2) - (2*b^4)/((a + b)^(7/2)*(b - a)^(3/2))))/((a + b)^(5/2)*(b - a)^(5/2)) + (b^4*log((2*b^4)/((a + b)^(7/2)*(b - a)^(3/2)) - (2*b^4*exp(x))/(a*b^4 + a^4*b + a^5 + b^5 - 2*a^2*b^3 - 2*a^3*b^2)))/((a + b)^(5/2)*(b - a)^(5/2))

3.115 $\int \frac{\tanh^5(x)}{1+\tanh(x)} dx$

Optimal. Leaf size=43

$$\frac{5x}{2} - 2\log(\cosh(x)) - \frac{5\tanh(x)}{2} + \tanh^2(x) - \frac{5\tanh^3(x)}{6} + \frac{\tanh^4(x)}{2(1+\tanh(x))}$$

[Out] 5/2*x-2*ln(cosh(x))-5/2*tanh(x)+tanh(x)^2-5/6*tanh(x)^3+1/2*tanh(x)^4/(1+tanh(x))

Rubi [A]

time = 0.07, antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {3631, 3609, 3606, 3556}

$$\frac{5x}{2} + \frac{\tanh^4(x)}{2(\tanh(x)+1)} - \frac{5\tanh^3(x)}{6} + \tanh^2(x) - \frac{5\tanh(x)}{2} - 2\log(\cosh(x))$$

Antiderivative was successfully verified.

[In] Int[Tanh[x]^5/(1 + Tanh[x]),x]

[Out] (5*x)/2 - 2*Log[Cosh[x]] - (5*Tanh[x])/2 + Tanh[x]^2 - (5*Tanh[x]^3)/6 + Tanh[x]^4/(2*(1 + Tanh[x]))

Rule 3556

Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] :> Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]

Rule 3606

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)]), x_Symbol] :> Simp[(a*c - b*d)*x, x] + (Dist[b*c + a*d, Int[Tan[e + f*x], x], x] + Simp[b*d*(Tan[e + f*x]/f), x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]

Rule 3609

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)]), x_Symbol] :> Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]

Rule 3631

```
Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)/((a_.) + (b_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] :> Simp[(b*c - a*d)*((c + d*Tan[e + f*x])^(n - 1)/(
2*a*f*(a + b*Tan[e + f*x]))), x] + Dist[1/(2*a^2), Int[(c + d*Tan[e + f*x])
^(n - 2)*Simp[a*c^2 + a*d^2*(n - 1) - b*c*d*n - d*(a*c*(n - 2) + b*d*n)*Tan
[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]
&& EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[n, 1]
```

Rubi steps

$$\begin{aligned}
\int \frac{\tanh^5(x)}{1 + \tanh(x)} dx &= \frac{\tanh^4(x)}{2(1 + \tanh(x))} - \frac{1}{2} \int (4 - 5 \tanh(x)) \tanh^3(x) dx \\
&= -\frac{5}{6} \tanh^3(x) + \frac{\tanh^4(x)}{2(1 + \tanh(x))} + \frac{1}{2} \int (-5i + 4i \tanh(x)) \tanh^2(x) dx \\
&= \tanh^2(x) - \frac{5 \tanh^3(x)}{6} + \frac{\tanh^4(x)}{2(1 + \tanh(x))} + \frac{1}{2} \int \tanh(x)(-4 + 5 \tanh(x)) dx \\
&= \frac{5x}{2} - \frac{5 \tanh(x)}{2} + \tanh^2(x) - \frac{5 \tanh^3(x)}{6} + \frac{\tanh^4(x)}{2(1 + \tanh(x))} - 2 \int \tanh(x) dx \\
&= \frac{5x}{2} - 2 \log(\cosh(x)) - \frac{5 \tanh(x)}{2} + \tanh^2(x) - \frac{5 \tanh^3(x)}{6} + \frac{\tanh^4(x)}{2(1 + \tanh(x))}
\end{aligned}$$

Mathematica [A]

time = 0.07, size = 40, normalized size = 0.93

$$\frac{1}{12} (30x + 3 \cosh(2x) - 24 \log(\cosh(x)) - 3 \sinh(2x) - 28 \tanh(x) + \operatorname{sech}^2(x)(-6 + 4 \tanh(x)))$$

Antiderivative was successfully verified.

[In] Integrate[Tanh[x]^5/(1 + Tanh[x]), x]

[Out] (30*x + 3*Cosh[2*x] - 24*Log[Cosh[x]] - 3*Sinh[2*x] - 28*Tanh[x] + Sech[x]^2*(-6 + 4*Tanh[x]))/12

Maple [A]

time = 0.38, size = 40, normalized size = 0.93

method	result	size
derivativedivides	$-\frac{\tanh^3(x)}{3} + \frac{\tanh^2(x)}{2} - 2 \tanh(x) + \frac{1}{2+2 \tanh(x)} + \frac{9 \ln(1+\tanh(x))}{4} - \frac{\ln(\tanh(x)-1)}{4}$	40
default	$-\frac{\tanh^3(x)}{3} + \frac{\tanh^2(x)}{2} - 2 \tanh(x) + \frac{1}{2+2 \tanh(x)} + \frac{9 \ln(1+\tanh(x))}{4} - \frac{\ln(\tanh(x)-1)}{4}$	40

risch	$\frac{9x}{2} + \frac{e^{-2x}}{4} + \frac{4e^{4x} + 6e^{2x} + \frac{14}{3}}{(1+e^{2x})^3} - 2 \ln(1 + e^{2x})$	44
-------	--	----

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tanh(x)^5/(1+tanh(x)),x,method=_RETURNVERBOSE)`

[Out] $-1/3*\tanh(x)^3+1/2*\tanh(x)^2-2*\tanh(x)+1/2/(1+\tanh(x))+9/4*\ln(1+\tanh(x))-1/4*\ln(\tanh(x)-1)$

Maxima [A]

time = 0.46, size = 55, normalized size = 1.28

$$\frac{1}{2}x - \frac{2(15e^{(-2x)} + 12e^{(-4x)} + 7)}{3(3e^{(-2x)} + 3e^{(-4x)} + e^{(-6x)} + 1)} + \frac{1}{4}e^{(-2x)} - 2 \log(e^{(-2x)} + 1)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(x)^5/(1+tanh(x)),x, algorithm="maxima")`

[Out] $1/2*x - 2/3*(15*e^{(-2*x)} + 12*e^{(-4*x)} + 7)/(3*e^{(-2*x)} + 3*e^{(-4*x)} + e^{(-6*x)} + 1) + 1/4*e^{(-2*x)} - 2*\log(e^{(-2*x)} + 1)$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 571 vs. 2(35) = 70.

time = 0.45, size = 571, normalized size = 13.28

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(x)^5/(1+tanh(x)),x, algorithm="fricas")`

[Out] $1/12*(54*x*\cosh(x)^8 + 432*x*\cosh(x)*\sinh(x)^7 + 54*x*\sinh(x)^8 + 3*(54*x + 17)*\cosh(x)^6 + 3*(504*x*\cosh(x)^2 + 54*x + 17)*\sinh(x)^6 + 18*(168*x*\cosh(x)^3 + (54*x + 17)*\cosh(x))*\sinh(x)^5 + 81*(2*x + 1)*\cosh(x)^4 + 9*(420*x*\cosh(x)^4 + 5*(54*x + 17)*\cosh(x)^2 + 18*x + 9)*\sinh(x)^4 + 12*(252*x*\cosh(x)^5 + 5*(54*x + 17)*\cosh(x)^3 + 27*(2*x + 1)*\cosh(x))*\sinh(x)^3 + (54*x + 65)*\cosh(x)^2 + (1512*x*\cosh(x)^6 + 45*(54*x + 17)*\cosh(x)^4 + 486*(2*x + 1)*\cosh(x)^2 + 54*x + 65)*\sinh(x)^2 - 24*(\cosh(x)^8 + 8*\cosh(x)*\sinh(x)^7 + \sinh(x)^8 + (28*\cosh(x)^2 + 3)*\sinh(x)^6 + 3*\cosh(x)^6 + 2*(28*\cosh(x)^3 + 9*\cosh(x))*\sinh(x)^5 + (70*\cosh(x)^4 + 45*\cosh(x)^2 + 3)*\sinh(x)^4 + 3*\cosh(x)^4 + 4*(14*\cosh(x)^5 + 15*\cosh(x)^3 + 3*\cosh(x))*\sinh(x)^3 + (28*\cosh(x)^6 + 45*\cosh(x)^4 + 18*\cosh(x)^2 + 1)*\sinh(x)^2 + \cosh(x)^2 + 2*(4*\cosh(x)^7 + 9*\cosh(x)^5 + 6*\cosh(x)^3 + \cosh(x))*\sinh(x))*\log(2*\cosh(x)/(\cosh(x) - \sinh(x))) + 2*(216*x*\cosh(x)^7 + 9*(54*x + 17)*\cosh(x)^5 + 162*(2*x + 1)*\cosh(x)^3 + (54*x + 65)*\cosh(x))*\sinh(x) + 3)/(\cosh(x)^8 + 8*\cosh(x)*\sinh(x)^7 + \sinh(x)^8 + (28*\cosh(x)^2 + 3)*\sinh(x)^6 + 3*\cosh(x)^6 + 2*(28*\cosh(x)^$

$3 + 9*\cosh(x))*\sinh(x)^5 + (70*\cosh(x)^4 + 45*\cosh(x)^2 + 3)*\sinh(x)^4 + 3*\cosh(x)^4 + 4*(14*\cosh(x)^5 + 15*\cosh(x)^3 + 3*\cosh(x))*\sinh(x)^3 + (28*\cosh(x)^6 + 45*\cosh(x)^4 + 18*\cosh(x)^2 + 1)*\sinh(x)^2 + \cosh(x)^2 + 2*(4*\cosh(x)^7 + 9*\cosh(x)^5 + 6*\cosh(x)^3 + \cosh(x))*\sinh(x)$

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 104 vs. $2(42) = 84$.

time = 0.22, size = 104, normalized size = 2.42

$$\frac{3x \tanh(x)}{6 \tanh(x) + 6} + \frac{3x}{6 \tanh(x) + 6} + \frac{12 \log(\tanh(x) + 1) \tanh(x)}{6 \tanh(x) + 6} + \frac{12 \log(\tanh(x) + 1)}{6 \tanh(x) + 6} - \frac{2 \tanh^4(x)}{6 \tanh(x) + 6} + \frac{\tanh^3(x)}{6 \tanh(x) + 6} - \frac{9 \tanh^2(x)}{6 \tanh(x) + 6} + \frac{15}{6 \tanh(x) + 6}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)**5/(1+tanh(x)),x)

[Out] $3*x*\tanh(x)/(6*\tanh(x) + 6) + 3*x/(6*\tanh(x) + 6) + 12*\log(\tanh(x) + 1)*\tanh(x)/(6*\tanh(x) + 6) + 12*\log(\tanh(x) + 1)/(6*\tanh(x) + 6) - 2*\tanh(x)**4/(6*\tanh(x) + 6) + \tanh(x)**3/(6*\tanh(x) + 6) - 9*\tanh(x)**2/(6*\tanh(x) + 6) + 15/(6*\tanh(x) + 6)$

Giac [A]

time = 0.43, size = 47, normalized size = 1.09

$$\frac{9}{2}x + \frac{(51e^{6x} + 81e^{4x} + 65e^{2x} + 3)e^{-2x}}{12(e^{2x} + 1)^3} - 2 \log(e^{2x} + 1)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)^5/(1+tanh(x)),x, algorithm="giac")

[Out] $9/2*x + 1/12*(51*e^{6*x} + 81*e^{4*x} + 65*e^{2*x} + 3)*e^{-2*x}/(e^{2*x} + 1)^3 - 2*\log(e^{2*x} + 1)$

Mupad [B]

time = 0.10, size = 35, normalized size = 0.81

$$\frac{x}{2} + 2 \ln(\tanh(x) + 1) - 2 \tanh(x) + \frac{\tanh(x)^2}{2} - \frac{\tanh(x)^3}{3} + \frac{1}{2(\tanh(x) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(x)^5/(tanh(x) + 1),x)

[Out] $x/2 + 2*\log(\tanh(x) + 1) - 2*\tanh(x) + \tanh(x)^2/2 - \tanh(x)^3/3 + 1/(2*(\tanh(x) + 1))$

$$3.116 \quad \int \frac{\tanh^4(x)}{1+\tanh(x)} dx$$

Optimal. Leaf size=37

$$-\frac{3x}{2} + 2 \log(\cosh(x)) + \frac{3 \tanh(x)}{2} - \tanh^2(x) + \frac{\tanh^3(x)}{2(1 + \tanh(x))}$$

[Out] $-3/2*x+2*\ln(\cosh(x))+3/2*\tanh(x)-\tanh(x)^2+1/2*\tanh(x)^3/(1+\tanh(x))$

Rubi [A]

time = 0.05, antiderivative size = 37, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {3631, 3609, 3606, 3556}

$$-\frac{3x}{2} + \frac{\tanh^3(x)}{2(\tanh(x) + 1)} - \tanh^2(x) + \frac{3 \tanh(x)}{2} + 2 \log(\cosh(x))$$

Antiderivative was successfully verified.

[In] Int[Tanh[x]^4/(1 + Tanh[x]), x]

[Out] $(-3*x)/2 + 2*\text{Log}[\text{Cosh}[x]] + (3*\text{Tanh}[x])/2 - \text{Tanh}[x]^2 + \text{Tanh}[x]^3/(2*(1 + \text{Tanh}[x]))$

Rule 3556

Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]

Rule 3606

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Dist[b*c + a*d, Int[Tan[e + f*x], x], x] + Simp[b*d*(Tan[e + f*x]/f), x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]

Rule 3609

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]

Rule 3631

Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)])^(n_)/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)]), x_Symbol] := Simp[(b*c - a*d)*((c + d*Tan[e + f*x])^(n - 1)/((

```
2*a*f*(a + b*Tan[e + f*x]))), x] + Dist[1/(2*a^2), Int[(c + d*Tan[e + f*x])
^(n - 2)*Simp[a*c^2 + a*d^2*(n - 1) - b*c*d*n - d*(a*c*(n - 2) + b*d*n)*Tan
[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]
&& EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[n, 1]
```

Rubi steps

$$\begin{aligned} \int \frac{\tanh^4(x)}{1 + \tanh(x)} dx &= \frac{\tanh^3(x)}{2(1 + \tanh(x))} - \frac{1}{2} \int (3 - 4 \tanh(x)) \tanh^2(x) dx \\ &= -\tanh^2(x) + \frac{\tanh^3(x)}{2(1 + \tanh(x))} + \frac{1}{2} \int (-4 + 3 \tanh(x)) \tanh(x) dx \\ &= -\frac{3x}{2} + \frac{3 \tanh(x)}{2} - \tanh^2(x) + \frac{\tanh^3(x)}{2(1 + \tanh(x))} + 2 \int \tanh(x) dx \\ &= -\frac{3x}{2} + 2 \log(\cosh(x)) + \frac{3 \tanh(x)}{2} - \tanh^2(x) + \frac{\tanh^3(x)}{2(1 + \tanh(x))} \end{aligned}$$

Mathematica [A]

time = 0.04, size = 33, normalized size = 0.89

$$\frac{1}{4}(-6x - \cosh(2x) + 8 \log(\cosh(x)) + 2 \operatorname{sech}^2(x) + \sinh(2x) + 4 \tanh(x))$$

Antiderivative was successfully verified.

```
[In] Integrate[Tanh[x]^4/(1 + Tanh[x]), x]
```

```
[Out] (-6*x - Cosh[2*x] + 8*Log[Cosh[x]] + 2*Sech[x]^2 + Sinh[2*x] + 4*Tanh[x])/4
```

Maple [A]

time = 0.36, size = 32, normalized size = 0.86

method	result	size
risch	$-\frac{7x}{2} - \frac{e^{-2x}}{4} - \frac{2}{(1+e^{2x})^2} + 2 \ln(1 + e^{2x})$	30
derivativedivides	$-\frac{(\tanh^2(x))}{2} + \tanh(x) - \frac{\ln(\tanh(x)-1)}{4} - \frac{1}{2(1+\tanh(x))} - \frac{7 \ln(1+\tanh(x))}{4}$	32
default	$-\frac{(\tanh^2(x))}{2} + \tanh(x) - \frac{\ln(\tanh(x)-1)}{4} - \frac{1}{2(1+\tanh(x))} - \frac{7 \ln(1+\tanh(x))}{4}$	32

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(tanh(x)^4/(1+tanh(x)), x, method=_RETURNVERBOSE)
```

```
[Out] -1/2*tanh(x)^2+tanh(x)-1/4*ln(tanh(x)-1)-1/2/(1+tanh(x))-7/4*ln(1+tanh(x))
```

Maxima [A]

time = 0.47, size = 43, normalized size = 1.16

$$\frac{1}{2}x + \frac{2(2e^{-2x} + 1)}{2e^{-2x} + e^{-4x} + 1} - \frac{1}{4}e^{-2x} + 2 \log(e^{-2x} + 1)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)^4/(1+tanh(x)),x, algorithm="maxima")

[Out] 1/2*x + 2*(2*e^(-2*x) + 1)/(2*e^(-2*x) + e^(-4*x) + 1) - 1/4*e^(-2*x) + 2*log(e^(-2*x) + 1)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 354 vs. 2(31) = 62.

time = 0.56, size = 354, normalized size = 9.57

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)^4/(1+tanh(x)),x, algorithm="fricas")

[Out] -1/4*(14*x*cosh(x)^6 + 84*x*cosh(x)*sinh(x)^5 + 14*x*sinh(x)^6 + (28*x + 1)*cosh(x)^4 + (210*x*cosh(x)^2 + 28*x + 1)*sinh(x)^4 + 4*(70*x*cosh(x)^3 + (28*x + 1)*cosh(x))*sinh(x)^3 + 2*(7*x + 5)*cosh(x)^2 + 2*(105*x*cosh(x)^4 + 3*(28*x + 1)*cosh(x)^2 + 7*x + 5)*sinh(x)^2 - 8*(cosh(x)^6 + 6*cosh(x)*sinh(x)^5 + sinh(x)^6 + (15*cosh(x)^2 + 2)*sinh(x)^4 + 2*cosh(x)^4 + 4*(5*cosh(x)^3 + 2*cosh(x))*sinh(x)^3 + (15*cosh(x)^4 + 12*cosh(x)^2 + 1)*sinh(x)^2 + cosh(x)^2 + 2*(3*cosh(x)^5 + 4*cosh(x)^3 + cosh(x))*sinh(x))*log(2*cosh(x)/(cosh(x) - sinh(x))) + 4*(21*x*cosh(x)^5 + (28*x + 1)*cosh(x)^3 + (7*x + 5)*cosh(x))*sinh(x) + 1)/(cosh(x)^6 + 6*cosh(x)*sinh(x)^5 + sinh(x)^6 + (15*cosh(x)^2 + 2)*sinh(x)^4 + 2*cosh(x)^4 + 4*(5*cosh(x)^3 + 2*cosh(x))*sinh(x)^3 + (15*cosh(x)^4 + 12*cosh(x)^2 + 1)*sinh(x)^2 + cosh(x)^2 + 2*(3*cosh(x)^5 + 4*cosh(x)^3 + cosh(x))*sinh(x))

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 85 vs. 2(34) = 68.

time = 0.20, size = 85, normalized size = 2.30

$$\frac{x \tanh(x)}{2 \tanh(x) + 2} + \frac{x}{2 \tanh(x) + 2} - \frac{4 \log(\tanh(x) + 1) \tanh(x)}{2 \tanh(x) + 2} - \frac{4 \log(\tanh(x) + 1)}{2 \tanh(x) + 2} - \frac{\tanh^3(x)}{2 \tanh(x) + 2} + \frac{\tanh^2(x)}{2 \tanh(x) + 2} - \frac{3}{2 \tanh(x) + 2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)**4/(1+tanh(x)),x)

[Out] x*tanh(x)/(2*tanh(x) + 2) + x/(2*tanh(x) + 2) - 4*log(tanh(x) + 1)*tanh(x)/(2*tanh(x) + 2) - 4*log(tanh(x) + 1)/(2*tanh(x) + 2) - tanh(x)**3/(2*tanh(x) + 2) + tanh(x)**2/(2*tanh(x) + 2) - 3/(2*tanh(x) + 2)

Giac [A]

time = 0.41, size = 39, normalized size = 1.05

$$-\frac{7}{2}x - \frac{(e^{4x} + 10e^{2x} + 1)e^{-2x}}{4(e^{2x} + 1)^2} + 2 \log(e^{2x} + 1)$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(tanh(x)^4/(1+tanh(x)),x, algorithm="giac")``[Out] -7/2*x - 1/4*(e^(4*x) + 10*e^(2*x) + 1)*e^(-2*x)/(e^(2*x) + 1)^2 + 2*log(e^(2*x) + 1)`**Mupad [B]**

time = 1.05, size = 29, normalized size = 0.78

$$\frac{x}{2} - 2 \ln(\tanh(x) + 1) + \tanh(x) - \frac{\tanh(x)^2}{2} - \frac{1}{2(\tanh(x) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(tanh(x)^4/(tanh(x) + 1),x)``[Out] x/2 - 2*log(tanh(x) + 1) + tanh(x) - tanh(x)^2/2 - 1/(2*(tanh(x) + 1))`

$$3.117 \quad \int \frac{\tanh^3(x)}{1+\tanh(x)} dx$$

Optimal. Leaf size=31

$$\frac{3x}{2} - \log(\cosh(x)) - \frac{3 \tanh(x)}{2} + \frac{\tanh^2(x)}{2(1 + \tanh(x))}$$

[Out] 3/2*x-ln(cosh(x))-3/2*tanh(x)+1/2*tanh(x)^2/(1+tanh(x))

Rubi [A]

time = 0.04, antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3631, 3606, 3556}

$$\frac{3x}{2} + \frac{\tanh^2(x)}{2(\tanh(x) + 1)} - \frac{3 \tanh(x)}{2} - \log(\cosh(x))$$

Antiderivative was successfully verified.

[In] Int[Tanh[x]^3/(1 + Tanh[x]),x]

[Out] (3*x)/2 - Log[Cosh[x]] - (3*Tanh[x])/2 + Tanh[x]^2/(2*(1 + Tanh[x]))

Rule 3556

Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]

Rule 3606

Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Dist[b*c + a*d, Int[Tan[e + f*x], x], x] + Simp[b*d*(Tan[e + f*x]/f), x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]

Rule 3631

Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((c + d*Tan[e + f*x])^(n - 1)/(2*a*f*(a + b*Tan[e + f*x]))), x] + Dist[1/(2*a^2), Int[(c + d*Tan[e + f*x])^(n - 2)*Simp[a*c^2 + a*d^2*(n - 1) - b*c*d*n - d*(a*c*(n - 2) + b*d*n)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[n, 1]

Rubi steps

$$\begin{aligned}
\int \frac{\tanh^3(x)}{1 + \tanh(x)} dx &= \frac{\tanh^2(x)}{2(1 + \tanh(x))} - \frac{1}{2} \int (2 - 3 \tanh(x)) \tanh(x) dx \\
&= \frac{3x}{2} - \frac{3 \tanh(x)}{2} + \frac{\tanh^2(x)}{2(1 + \tanh(x))} - \int \tanh(x) dx \\
&= \frac{3x}{2} - \log(\cosh(x)) - \frac{3 \tanh(x)}{2} + \frac{\tanh^2(x)}{2(1 + \tanh(x))}
\end{aligned}$$

Mathematica [A]

time = 0.04, size = 27, normalized size = 0.87

$$\frac{1}{4}(6x + \cosh(2x) - 4 \log(\cosh(x)) - \sinh(2x) - 4 \tanh(x))$$

Antiderivative was successfully verified.

`[In] Integrate[Tanh[x]^3/(1 + Tanh[x]), x]``[Out] (6*x + Cosh[2*x] - 4*Log[Cosh[x]] - Sinh[2*x] - 4*Tanh[x])/4`**Maple [A]**

time = 0.29, size = 28, normalized size = 0.90

method	result	size
derivativedivides	$-\tanh(x) + \frac{1}{2+2\tanh(x)} + \frac{5\ln(1+\tanh(x))}{4} - \frac{\ln(\tanh(x)-1)}{4}$	28
default	$-\tanh(x) + \frac{1}{2+2\tanh(x)} + \frac{5\ln(1+\tanh(x))}{4} - \frac{\ln(\tanh(x)-1)}{4}$	28
risch	$\frac{5x}{2} + \frac{e^{-2x}}{4} + \frac{2}{1+e^{2x}} - \ln(1 + e^{2x})$	30

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(tanh(x)^3/(1+tanh(x)), x, method=_RETURNVERBOSE)``[Out] -tanh(x)+1/2/(1+tanh(x))+5/4*ln(1+tanh(x))-1/4*ln(tanh(x)-1)`**Maxima [A]**

time = 0.47, size = 29, normalized size = 0.94

$$\frac{1}{2}x - \frac{2}{e^{(-2x)} + 1} + \frac{1}{4}e^{(-2x)} - \log(e^{(-2x)} + 1)$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(tanh(x)^3/(1+tanh(x)), x, algorithm="maxima")``[Out] 1/2*x - 2/(e^(-2*x) + 1) + 1/4*e^(-2*x) - log(e^(-2*x) + 1)`

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 186 vs. 2(25) = 50.

time = 0.42, size = 186, normalized size = 6.00

$$\frac{10x \cosh(x)^4 + 40x \cosh(x) \sinh(x)^3 + 10x \sinh(x)^4 + (10x + 9) \cosh(x)^2 + (60x \cosh(x)^2 + 10x + 9) \sinh(x)^2 - 4(\cosh(x)^4 + 4 \cosh(x) \sinh(x)^3 + \sinh(x)^4) + (6 \cosh(x)^2 + 1) \sinh(x)^2 + \cosh(x)^2 + 2(2 \cosh(x)^3 + \cosh(x)) \sinh(x) \log\left(\frac{2 \cosh(x)}{\cosh(x) - \sinh(x)}\right) + 2(20x \cosh(x)^3 + (10x + 9) \cosh(x) \sinh(x) + 1)}{4(\cosh(x)^3 + 4 \cosh(x) \sinh(x)^2 + \sinh(x)^4) + (6 \cosh(x)^2 + 1) \sinh(x)^2 + \cosh(x)^2 + 2(2 \cosh(x)^3 + \cosh(x)) \sinh(x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)^3/(1+tanh(x)),x, algorithm="fricas")

[Out] 1/4*(10*x*cosh(x)^4 + 40*x*cosh(x)*sinh(x)^3 + 10*x*sinh(x)^4 + (10*x + 9)*cosh(x)^2 + (60*x*cosh(x)^2 + 10*x + 9)*sinh(x)^2 - 4*(cosh(x)^4 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4 + (6*cosh(x)^2 + 1)*sinh(x)^2 + cosh(x)^2 + 2*(2*cosh(x)^3 + cosh(x))*sinh(x))*log(2*cosh(x)/(cosh(x) - sinh(x))) + 2*(20*x*cosh(x)^3 + (10*x + 9)*cosh(x)*sinh(x) + 1)/(cosh(x)^4 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4 + (6*cosh(x)^2 + 1)*sinh(x)^2 + cosh(x)^2 + 2*(2*cosh(x)^3 + cosh(x))*sinh(x))

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 75 vs. 2(27) = 54.

time = 0.19, size = 75, normalized size = 2.42

$$\frac{x \tanh(x)}{2 \tanh(x) + 2} + \frac{x}{2 \tanh(x) + 2} + \frac{2 \log(\tanh(x) + 1) \tanh(x)}{2 \tanh(x) + 2} + \frac{2 \log(\tanh(x) + 1)}{2 \tanh(x) + 2} - \frac{2 \tanh^2(x)}{2 \tanh(x) + 2} + \frac{3}{2 \tanh(x) + 2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)**3/(1+tanh(x)),x)

[Out] x*tanh(x)/(2*tanh(x) + 2) + x/(2*tanh(x) + 2) + 2*log(tanh(x) + 1)*tanh(x)/(2*tanh(x) + 2) + 2*log(tanh(x) + 1)/(2*tanh(x) + 2) - 2*tanh(x)**2/(2*tanh(x) + 2) + 3/(2*tanh(x) + 2)

Giac [A]

time = 0.42, size = 35, normalized size = 1.13

$$\frac{5}{2}x + \frac{(9e^{(2x)} + 1)e^{(-2x)}}{4(e^{(2x)} + 1)} - \log(e^{(2x)} + 1)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)^3/(1+tanh(x)),x, algorithm="giac")

[Out] 5/2*x + 1/4*(9*e^(2*x) + 1)*e^(-2*x)/(e^(2*x) + 1) - log(e^(2*x) + 1)

Mupad [B]

time = 0.07, size = 21, normalized size = 0.68

$$\frac{x}{2} + \ln(\tanh(x) + 1) - \tanh(x) + \frac{1}{2(\tanh(x) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(x)^3/(tanh(x) + 1),x)

[Out] x/2 + log(tanh(x) + 1) - tanh(x) + 1/(2*(tanh(x) + 1))

$$3.118 \quad \int \frac{\tanh^2(x)}{1+\tanh(x)} dx$$

Optimal. Leaf size=19

$$-\frac{x}{2} + \log(\cosh(x)) - \frac{1}{2(1 + \tanh(x))}$$

[Out] -1/2*x+ln(cosh(x))-1/2/(1+tanh(x))

Rubi [A]

time = 0.03, antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {3621, 3556}

$$-\frac{x}{2} - \frac{1}{2(\tanh(x) + 1)} + \log(\cosh(x))$$

Antiderivative was successfully verified.

[In] Int[Tanh[x]^2/(1 + Tanh[x]),x]

[Out] -1/2*x + Log[Cosh[x]] - 1/(2*(1 + Tanh[x]))

Rule 3556

Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]

Rule 3621

Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(-b)*(a*c + b*d)^2*((a + b*Tan[e + f*x])^m/(2*a^3*f*m)), x] + Dist[1/(2*a^2), Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c^2 - 2*b*c*d + a*d^2 - 2*b*d^2*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && LeQ[m, -1] && EqQ[a^2 + b^2, 0]

Rubi steps

$$\begin{aligned} \int \frac{\tanh^2(x)}{1 + \tanh(x)} dx &= -\frac{1}{2(1 + \tanh(x))} - \frac{1}{2} \int (1 - 2 \tanh(x)) dx \\ &= -\frac{x}{2} - \frac{1}{2(1 + \tanh(x))} + \int \tanh(x) dx \\ &= -\frac{x}{2} + \log(\cosh(x)) - \frac{1}{2(1 + \tanh(x))} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 23, normalized size = 1.21

$$\frac{1}{4}(-2x - \cosh(2x) + 4 \log(\cosh(x)) + \sinh(2x))$$

Antiderivative was successfully verified.

`[In] Integrate[Tanh[x]^2/(1 + Tanh[x]), x]``[Out] (-2*x - Cosh[2*x] + 4*Log[Cosh[x]] + Sinh[2*x])/4`**Maple [A]**

time = 0.29, size = 24, normalized size = 1.26

method	result	size
risch	$-\frac{3x}{2} - \frac{e^{-2x}}{4} + \ln(1 + e^{2x})$	18
derivativedivides	$-\frac{1}{2(1+\tanh(x))} - \frac{3 \ln(1+\tanh(x))}{4} - \frac{\ln(\tanh(x)-1)}{4}$	24
default	$-\frac{1}{2(1+\tanh(x))} - \frac{3 \ln(1+\tanh(x))}{4} - \frac{\ln(\tanh(x)-1)}{4}$	24

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(tanh(x)^2/(1+tanh(x)), x, method=_RETURNVERBOSE)``[Out] -1/2/(1+tanh(x))-3/4*ln(1+tanh(x))-1/4*ln(tanh(x)-1)`**Maxima [A]**

time = 0.47, size = 17, normalized size = 0.89

$$\frac{1}{2}x - \frac{1}{4}e^{(-2x)} + \log(e^{(-2x)} + 1)$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(tanh(x)^2/(1+tanh(x)), x, algorithm="maxima")``[Out] 1/2*x - 1/4*e^(-2*x) + log(e^(-2*x) + 1)`**Fricas [B]** Leaf count of result is larger than twice the leaf count of optimal. 73 vs. 2(15) = 30.

time = 0.33, size = 73, normalized size = 3.84

$$\frac{6x \cosh(x)^2 + 12x \cosh(x) \sinh(x) + 6x \sinh(x)^2 - 4(\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2) \log\left(\frac{2 \cosh(x)}{\cosh(x) - \sinh(x)}\right) + 1}{4(\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(tanh(x)^2/(1+tanh(x)), x, algorithm="fricas")`

[Out] $-1/4*(6*x*\cosh(x)^2 + 12*x*\cosh(x)*\sinh(x) + 6*x*\sinh(x)^2 - 4*(\cosh(x)^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2)*\log(2*\cosh(x)/(\cosh(x) - \sinh(x))) + 1)/(\cosh(x)^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2)$

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 61 vs. $2(15) = 30$.

time = 0.17, size = 61, normalized size = 3.21

$$\frac{x \tanh(x)}{2 \tanh(x) + 2} + \frac{x}{2 \tanh(x) + 2} - \frac{2 \log(\tanh(x) + 1) \tanh(x)}{2 \tanh(x) + 2} - \frac{2 \log(\tanh(x) + 1)}{2 \tanh(x) + 2} - \frac{1}{2 \tanh(x) + 2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(x)**2/(1+tanh(x)),x)`

[Out] $x*\tanh(x)/(2*\tanh(x) + 2) + x/(2*\tanh(x) + 2) - 2*\log(\tanh(x) + 1)*\tanh(x)/(2*\tanh(x) + 2) - 2*\log(\tanh(x) + 1)/(2*\tanh(x) + 2) - 1/(2*\tanh(x) + 2)$

Giac [A]

time = 0.42, size = 17, normalized size = 0.89

$$-\frac{3}{2}x - \frac{1}{4}e^{(-2x)} + \log(e^{(2x)} + 1)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(x)^2/(1+tanh(x)),x, algorithm="giac")`

[Out] $-3/2*x - 1/4*e^{(-2*x)} + \log(e^{(2*x)} + 1)$

Mupad [B]

time = 1.06, size = 21, normalized size = 1.11

$$\frac{x}{2} - \ln(\tanh(x) + 1) - \frac{1}{2(\tanh(x) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tanh(x)^2/(tanh(x) + 1),x)`

[Out] $x/2 - \log(\tanh(x) + 1) - 1/(2*(\tanh(x) + 1))$

$$3.119 \quad \int \frac{\tanh(x)}{1+\tanh(x)} dx$$

Optimal. Leaf size=16

$$\frac{x}{2} + \frac{1}{2(1 + \tanh(x))}$$

[Out] 1/2*x+1/2/(1+tanh(x))

Rubi [A]

time = 0.01, antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {3607, 8}

$$\frac{x}{2} + \frac{1}{2(\tanh(x) + 1)}$$

Antiderivative was successfully verified.

[In] Int[Tanh[x]/(1 + Tanh[x]),x]

[Out] x/2 + 1/(2*(1 + Tanh[x]))

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 3607

Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-(b*c - a*d))*((a + b*Tan[e + f*x])^m/(2*a*f*m)), x] + Dist[(b*c + a*d)/(2*a*b), Int[(a + b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0]

Rubi steps

$$\begin{aligned} \int \frac{\tanh(x)}{1 + \tanh(x)} dx &= \frac{1}{2(1 + \tanh(x))} + \frac{\int 1 dx}{2} \\ &= \frac{x}{2} + \frac{1}{2(1 + \tanh(x))} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 18, normalized size = 1.12

$$\frac{1}{4}(2x + \cosh(2x) - \sinh(2x))$$

Antiderivative was successfully verified.

[In] Integrate[Tanh[x]/(1 + Tanh[x]),x]

[Out] (2*x + Cosh[2*x] - Sinh[2*x])/4

Maple [A]

time = 0.24, size = 24, normalized size = 1.50

method	result	size
risch	$\frac{x}{2} + \frac{e^{-2x}}{4}$	11
derivativdivides	$-\frac{\ln(\tanh(x)-1)}{4} + \frac{1}{2+2\tanh(x)} + \frac{\ln(1+\tanh(x))}{4}$	24
default	$-\frac{\ln(\tanh(x)-1)}{4} + \frac{1}{2+2\tanh(x)} + \frac{\ln(1+\tanh(x))}{4}$	24

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(x)/(1+tanh(x)),x,method=_RETURNVERBOSE)

[Out] -1/4*ln(tanh(x)-1)+1/2/(1+tanh(x))+1/4*ln(1+tanh(x))

Maxima [A]

time = 0.26, size = 10, normalized size = 0.62

$$\frac{1}{2}x + \frac{1}{4}e^{(-2x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)/(1+tanh(x)),x, algorithm="maxima")

[Out] 1/2*x + 1/4*e^(-2*x)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 26 vs. 2(12) = 24.

time = 0.38, size = 26, normalized size = 1.62

$$\frac{(2x + 1) \cosh(x) + (2x - 1) \sinh(x)}{4(\cosh(x) + \sinh(x))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)/(1+tanh(x)),x, algorithm="fricas")

[Out] 1/4*((2*x + 1)*cosh(x) + (2*x - 1)*sinh(x))/(cosh(x) + sinh(x))

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 27 vs. 2(10) = 20.

time = 0.16, size = 27, normalized size = 1.69

$$\frac{x \tanh(x)}{2 \tanh(x) + 2} + \frac{x}{2 \tanh(x) + 2} + \frac{1}{2 \tanh(x) + 2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(x)/(1+tanh(x)),x)`

[Out] `x*tanh(x)/(2*tanh(x) + 2) + x/(2*tanh(x) + 2) + 1/(2*tanh(x) + 2)`

Giac [A]

time = 0.41, size = 10, normalized size = 0.62

$$\frac{1}{2}x + \frac{1}{4}e^{(-2x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(x)/(1+tanh(x)),x, algorithm="giac")`

[Out] `1/2*x + 1/4*e^(-2*x)`

Mupad [B]

time = 0.07, size = 10, normalized size = 0.62

$$\frac{x}{2} + \frac{e^{-2x}}{4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tanh(x)/(tanh(x) + 1),x)`

[Out] `x/2 + exp(-2*x)/4`

$$3.120 \quad \int \frac{1}{1+\tanh(x)} dx$$

Optimal. Leaf size=16

$$\frac{x}{2} - \frac{1}{2(1 + \tanh(x))}$$

[Out] 1/2*x-1/2/(1+tanh(x))

Rubi [A]

time = 0.01, antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3560, 8}

$$\frac{x}{2} - \frac{1}{2(\tanh(x) + 1)}$$

Antiderivative was successfully verified.

[In] Int[(1 + Tanh[x])^(-1),x]

[Out] x/2 - 1/(2*(1 + Tanh[x]))

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 3560

Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n), x] + Dist[1/(2*a), Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{1 + \tanh(x)} dx &= -\frac{1}{2(1 + \tanh(x))} + \frac{\int 1 dx}{2} \\ &= \frac{x}{2} - \frac{1}{2(1 + \tanh(x))} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 18, normalized size = 1.12

$$\frac{1}{4}(2x - \cosh(2x) + \sinh(2x))$$

Antiderivative was successfully verified.

[In] Integrate[(1 + Tanh[x])^(-1), x]

[Out] (2*x - Cosh[2*x] + Sinh[2*x])/4

Maple [A]

time = 0.23, size = 24, normalized size = 1.50

method	result	size
risch	$\frac{x}{2} - \frac{e^{-2x}}{4}$	11
derivativedivides	$-\frac{1}{2(1+\tanh(x))} + \frac{\ln(1+\tanh(x))}{4} - \frac{\ln(\tanh(x)-1)}{4}$	24
default	$-\frac{1}{2(1+\tanh(x))} + \frac{\ln(1+\tanh(x))}{4} - \frac{\ln(\tanh(x)-1)}{4}$	24

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(1+tanh(x)), x, method=_RETURNVERBOSE)

[Out] -1/2/(1+tanh(x))+1/4*ln(1+tanh(x))-1/4*ln(tanh(x)-1)

Maxima [A]

time = 0.25, size = 10, normalized size = 0.62

$$\frac{1}{2}x - \frac{1}{4}e^{(-2x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+tanh(x)), x, algorithm="maxima")

[Out] 1/2*x - 1/4*e^(-2*x)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 26 vs. 2(12) = 24.

time = 0.35, size = 26, normalized size = 1.62

$$\frac{(2x - 1) \cosh(x) + (2x + 1) \sinh(x)}{4(\cosh(x) + \sinh(x))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+tanh(x)), x, algorithm="fricas")

[Out] 1/4*((2*x - 1)*cosh(x) + (2*x + 1)*sinh(x))/(cosh(x) + sinh(x))

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 27 vs. 2(10) = 20.

time = 0.17, size = 27, normalized size = 1.69

$$\frac{x \tanh(x)}{2 \tanh(x) + 2} + \frac{x}{2 \tanh(x) + 2} - \frac{1}{2 \tanh(x) + 2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+tanh(x)),x)`

[Out] `x*tanh(x)/(2*tanh(x) + 2) + x/(2*tanh(x) + 2) - 1/(2*tanh(x) + 2)`

Giac [A]

time = 0.40, size = 10, normalized size = 0.62

$$\frac{1}{2}x - \frac{1}{4}e^{(-2x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+tanh(x)),x, algorithm="giac")`

[Out] `1/2*x - 1/4*e^(-2*x)`

Mupad [B]

time = 1.03, size = 10, normalized size = 0.62

$$\frac{x}{2} - \frac{e^{-2x}}{4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(tanh(x) + 1),x)`

[Out] `x/2 - exp(-2*x)/4`

$$3.121 \quad \int \frac{\coth(x)}{1+\tanh(x)} dx$$

Optimal. Leaf size=19

$$-\frac{x}{2} + \log(\sinh(x)) + \frac{1}{2(1+\tanh(x))}$$

[Out] -1/2*x+ln(sinh(x))+1/2/(1+tanh(x))

Rubi [A]

time = 0.03, antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {3632, 3560, 8, 3556}

$$-\frac{x}{2} + \frac{1}{2(\tanh(x)+1)} + \log(\sinh(x))$$

Antiderivative was successfully verified.

[In] Int[Coth[x]/(1 + Tanh[x]),x]

[Out] -1/2*x + Log[Sinh[x]] + 1/(2*(1 + Tanh[x]))

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 3556

Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]

Rule 3560

Int[((a_.) + (b_.)*tan[(c_.) + (d_.)*(x_.)])^(n_), x_Symbol] := Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n), x] + Dist[1/(2*a), Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]

Rule 3632

Int[1/(((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)])), x_Symbol] := Dist[b/(b*c - a*d), Int[1/(a + b*Tan[e + f*x]), x], x] - Dist[d/(b*c - a*d), Int[1/(c + d*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]

Rubi steps

$$\begin{aligned}
\int \frac{\coth(x)}{1 + \tanh(x)} dx &= \int \coth(x) dx - \int \frac{1}{1 + \tanh(x)} dx \\
&= \log(\sinh(x)) + \frac{1}{2(1 + \tanh(x))} - \frac{\int 1 dx}{2} \\
&= -\frac{x}{2} + \log(\sinh(x)) + \frac{1}{2(1 + \tanh(x))}
\end{aligned}$$

Mathematica [A]

time = 0.02, size = 23, normalized size = 1.21

$$\frac{1}{4}(-2x + \cosh(2x) + 4 \log(\sinh(x)) - \sinh(2x))$$

Antiderivative was successfully verified.

[In] Integrate[Coth[x]/(1 + Tanh[x]), x]

[Out] (-2*x + Cosh[2*x] + 4*Log[Sinh[x]] - Sinh[2*x])/4

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 42 vs. 2(15) = 30.

time = 0.42, size = 43, normalized size = 2.26

method	result	size
risch	$-\frac{3x}{2} + \frac{e^{-2x}}{4} + \ln(e^{2x} - 1)$	18
default	$\ln\left(\tanh\left(\frac{x}{2}\right)\right) - \frac{\ln(\tanh(\frac{x}{2})-1)}{2} - \frac{1}{\tanh(\frac{x}{2})+1} + \frac{1}{(\tanh(\frac{x}{2})+1)^2} - \frac{3 \ln(\tanh(\frac{x}{2})+1)}{2}$	43

Verification of antiderivative is not currently implemented for this CAS.

[In] int(coth(x)/(1+tanh(x)), x, method=_RETURNVERBOSE)

[Out] ln(tanh(1/2*x))-1/2*ln(tanh(1/2*x)-1)-1/(tanh(1/2*x)+1)+1/(tanh(1/2*x)+1)^2-3/2*ln(tanh(1/2*x)+1)

Maxima [A]

time = 0.27, size = 24, normalized size = 1.26

$$\frac{1}{2}x + \frac{1}{4}e^{(-2x)} + \log(e^{(-x)} + 1) + \log(e^{(-x)} - 1)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(x)/(1+tanh(x)), x, algorithm="maxima")

[Out] $1/2*x + 1/4*e^{(-2*x)} + \log(e^{-x} + 1) + \log(e^{-x} - 1)$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 73 vs. $2(15) = 30$.

time = 0.39, size = 73, normalized size = 3.84

$$\frac{6x \cosh(x)^2 + 12x \cosh(x) \sinh(x) + 6x \sinh(x)^2 - 4(\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2) \log\left(\frac{2 \sinh(x)}{\cosh(x) - \sinh(x)}\right) - 1}{4(\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(coth(x)/(1+tanh(x)),x, algorithm="fricas")`

[Out] $-1/4*(6*x*\cosh(x)^2 + 12*x*\cosh(x)*\sinh(x) + 6*x*\sinh(x)^2 - 4*(\cosh(x)^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2)*\log(2*\sinh(x)/(\cosh(x) - \sinh(x))) - 1)/(\cosh(x)^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\coth(x)}{\tanh(x) + 1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(coth(x)/(1+tanh(x)),x)`

[Out] `Integral(coth(x)/(tanh(x) + 1), x)`

Giac [A]

time = 0.42, size = 18, normalized size = 0.95

$$-\frac{3}{2}x + \frac{1}{4}e^{(-2x)} + \log(|e^{(2x)} - 1|)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(coth(x)/(1+tanh(x)),x, algorithm="giac")`

[Out] $-3/2*x + 1/4*e^{(-2*x)} + \log(\text{abs}(e^{(2*x)} - 1))$

Mupad [B]

time = 0.06, size = 17, normalized size = 0.89

$$\ln(e^{2x} - 1) - \frac{3x}{2} + \frac{e^{-2x}}{4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(coth(x)/(tanh(x) + 1),x)`

[Out] $\log(\exp(2*x) - 1) - (3*x)/2 + \exp(-2*x)/4$

3.122 $\int \frac{\coth^2(x)}{1+\tanh(x)} dx$

Optimal. Leaf size=29

$$\frac{3x}{2} - \frac{3 \coth(x)}{2} - \log(\sinh(x)) + \frac{\coth(x)}{2(1 + \tanh(x))}$$

[Out] 3/2*x-3/2*coth(x)-ln(sinh(x))+1/2*coth(x)/(1+tanh(x))

Rubi [A]

time = 0.05, antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {3633, 3610, 3612, 3556}

$$\frac{3x}{2} - \frac{3 \coth(x)}{2} - \log(\sinh(x)) + \frac{\coth(x)}{2(\tanh(x) + 1)}$$

Antiderivative was successfully verified.

[In] Int[Coth[x]^2/(1 + Tanh[x]),x]

[Out] (3*x)/2 - (3*Coth[x])/2 - Log[Sinh[x]] + Coth[x]/(2*(1 + Tanh[x]))

Rule 3556

Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]

Rule 3610

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Dist[1/(a^2 + b^2), Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]

Rule 3612

Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Dist[(b*c - a*d)/(a^2 + b^2), Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]

Rule 3633

Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-a)*((c + d*Tan[e + f*x])^(n + 1)/(2*f*(b

```
c - a*d)*(a + b*Tan[e + f*x]))), x] + Dist[1/(2*a*(b*c - a*d)), Int[(c + d*
Tan[e + f*x])^n*Simp[b*c + a*d*(n - 1) - b*d*n*Tan[e + f*x], x], x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] &
& NeQ[c^2 + d^2, 0] && !GtQ[n, 0]
```

Rubi steps

$$\begin{aligned} \int \frac{\coth^2(x)}{1 + \tanh(x)} dx &= \frac{\coth(x)}{2(1 + \tanh(x))} - \frac{1}{2} \int \coth^2(x)(-3 + 2 \tanh(x)) dx \\ &= -\frac{3 \coth(x)}{2} + \frac{\coth(x)}{2(1 + \tanh(x))} - \frac{1}{2} \int \coth(x)(-2i + 3i \tanh(x)) dx \\ &= \frac{3x}{2} - \frac{3 \coth(x)}{2} + \frac{\coth(x)}{2(1 + \tanh(x))} - \int \coth(x) dx \\ &= \frac{3x}{2} - \frac{3 \coth(x)}{2} - \log(\sinh(x)) + \frac{\coth(x)}{2(1 + \tanh(x))} \end{aligned}$$

Mathematica [A]

time = 0.04, size = 27, normalized size = 0.93

$$\frac{1}{4}(6x - \cosh(2x) - 4 \coth(x) - 4 \log(\sinh(x)) + \sinh(2x))$$

Antiderivative was successfully verified.

[In] Integrate[Coth[x]^2/(1 + Tanh[x]), x]

[Out] (6*x - Cosh[2*x] - 4*Coth[x] - 4*Log[Sinh[x]] + Sinh[2*x])/4

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 58 vs. 2(23) = 46.

time = 0.43, size = 59, normalized size = 2.03

method	result
risch	$\frac{5x}{2} - \frac{e^{-2x}}{4} - \frac{2}{e^{2x}-1} - \ln(e^{2x} - 1)$
default	$-\frac{\tanh(\frac{x}{2})}{2} - \frac{1}{(\tanh(\frac{x}{2})+1)^2} + \frac{1}{\tanh(\frac{x}{2})+1} + \frac{5 \ln(\tanh(\frac{x}{2})+1)}{2} - \frac{1}{2 \tanh(\frac{x}{2})} - \ln(\tanh(\frac{x}{2})) - \frac{\ln(\tanh(\frac{x}{2})-1)}{2}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(coth(x)^2/(1+tanh(x)), x, method=_RETURNVERBOSE)

[Out] -1/2*tanh(1/2*x)-1/(tanh(1/2*x)+1)^2+1/(tanh(1/2*x)+1)+5/2*ln(tanh(1/2*x)+1)-1/2/tanh(1/2*x)-ln(tanh(1/2*x))-1/2*ln(tanh(1/2*x)-1)

Maxima [A]

time = 0.26, size = 38, normalized size = 1.31

$$\frac{1}{2}x + \frac{2}{e^{(-2x)} - 1} - \frac{1}{4}e^{(-2x)} - \log(e^{(-x)} + 1) - \log(e^{(-x)} - 1)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(x)^2/(1+tanh(x)),x, algorithm="maxima")**[Out]** 1/2*x + 2/(e^(-2*x) - 1) - 1/4*e^(-2*x) - log(e^(-x) + 1) - log(e^(-x) - 1)**Fricas [B]** Leaf count of result is larger than twice the leaf count of optimal. 196 vs. 2(23) = 46.

time = 0.60, size = 196, normalized size = 6.76

$$\frac{10x \cosh(x)^3 + 40x \cosh(x) \sinh(x)^3 + 10x \sinh(x)^4 - (10x + 9) \cosh(x)^2 + (60x \cosh(x)^2 - 10x - 9) \sinh(x)^2 - 4(\cosh(x)^4 + 4 \cosh(x) \sinh(x)^3 + \sinh(x)^4) + (6 \cosh(x)^2 - 1) \sinh(x)^2 - \cosh(x)^2 + 2(2 \cosh(x)^3 - \cosh(x)) \sinh(x) \log\left(\frac{2 \sinh(x)}{\cosh(x) - \sinh(x)}\right) + 2(20x \cosh(x)^3 - (10x + 9) \cosh(x) \sinh(x) + 1)}{4(\cosh(x)^3 + 4 \cosh(x) \sinh(x)^3 + \sinh(x)^4) + (6 \cosh(x)^2 - 1) \sinh(x)^2 - \cosh(x)^2 + 2(2 \cosh(x)^3 - \cosh(x)) \sinh(x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(x)^2/(1+tanh(x)),x, algorithm="fricas")

[Out] 1/4*(10*x*cosh(x)^4 + 40*x*cosh(x)*sinh(x)^3 + 10*x*sinh(x)^4 - (10*x + 9)*cosh(x)^2 + (60*x*cosh(x)^2 - 10*x - 9)*sinh(x)^2 - 4*(cosh(x)^4 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4) + (6*cosh(x)^2 - 1)*sinh(x)^2 - cosh(x)^2 + 2*(2*cosh(x)^3 - cosh(x))*sinh(x))*log(2*sinh(x)/(cosh(x) - sinh(x))) + 2*(20*x*cosh(x)^3 - (10*x + 9)*cosh(x))*sinh(x) + 1)/(cosh(x)^4 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4 + (6*cosh(x)^2 - 1)*sinh(x)^2 - cosh(x)^2 + 2*(2*cosh(x)^3 - cosh(x))*sinh(x))

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\coth^2(x)}{\tanh(x) + 1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(x)**2/(1+tanh(x)),x)**[Out]** Integral(coth(x)**2/(tanh(x) + 1), x)**Giac [A]**

time = 0.40, size = 36, normalized size = 1.24

$$\frac{5}{2}x - \frac{(9e^{(2x)} - 1)e^{(-2x)}}{4(e^{(2x)} - 1)} - \log(|e^{(2x)} - 1|)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(x)^2/(1+tanh(x)),x, algorithm="giac")

[Out] $\frac{5}{2}x - \frac{1}{4}(9e^{2x} - 1)e^{-2x}/(e^{2x} - 1) - \log(\text{abs}(e^{2x} - 1))$

Mupad [B]

time = 1.03, size = 29, normalized size = 1.00

$$\frac{5x}{2} - \ln(e^{2x} - 1) - \frac{e^{-2x}}{4} - \frac{2}{e^{2x} - 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(coth(x)^2/(tanh(x) + 1),x)

[Out] $(5x)/2 - \log(\exp(2x) - 1) - \exp(-2x)/4 - 2/(\exp(2x) - 1)$

$$3.123 \quad \int \frac{\coth^3(x)}{1+\tanh(x)} dx$$

Optimal. Leaf size=37

$$-\frac{3x}{2} + \frac{3 \coth(x)}{2} - \coth^2(x) + 2 \log(\sinh(x)) + \frac{\coth^2(x)}{2(1 + \tanh(x))}$$

[Out] $-3/2*x+3/2*\coth(x)-\coth(x)^2+2*\ln(\sinh(x))+1/2*\coth(x)^2/(1+\tanh(x))$

Rubi [A]

time = 0.07, antiderivative size = 37, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {3633, 3610, 3612, 3556}

$$-\frac{3x}{2} - \coth^2(x) + \frac{3 \coth(x)}{2} + 2 \log(\sinh(x)) + \frac{\coth^2(x)}{2(\tanh(x) + 1)}$$

Antiderivative was successfully verified.

[In] Int[Coth[x]^3/(1 + Tanh[x]),x]

[Out] $(-3*x)/2 + (3*Coth[x])/2 - Coth[x]^2 + 2*Log[Sinh[x]] + Coth[x]^2/(2*(1 + Tanh[x]))$

Rule 3556

Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]

Rule 3610

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Dist[1/(a^2 + b^2), Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]

Rule 3612

Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Dist[(b*c - a*d)/(a^2 + b^2), Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]

Rule 3633

```
Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)/((a_) + (b_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(-a)*((c + d*Tan[e + f*x])^(n + 1)/(2*f*(b*
c - a*d)*(a + b*Tan[e + f*x]))), x] + Dist[1/(2*a*(b*c - a*d)), Int[(c + d*
Tan[e + f*x])^n*Simp[b*c + a*d*(n - 1) - b*d*n*Tan[e + f*x], x], x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] &
& NeQ[c^2 + d^2, 0] && !GtQ[n, 0]
```

Rubi steps

$$\begin{aligned}
\int \frac{\coth^3(x)}{1 + \tanh(x)} dx &= \frac{\coth^2(x)}{2(1 + \tanh(x))} - \frac{1}{2} \int \coth^3(x)(-4 + 3 \tanh(x)) dx \\
&= -\coth^2(x) + \frac{\coth^2(x)}{2(1 + \tanh(x))} - \frac{1}{2} \int \coth^2(x)(-3i + 4i \tanh(x)) dx \\
&= \frac{3 \coth(x)}{2} - \coth^2(x) + \frac{\coth^2(x)}{2(1 + \tanh(x))} + \frac{1}{2} \int \coth(x)(4 - 3 \tanh(x)) dx \\
&= -\frac{3x}{2} + \frac{3 \coth(x)}{2} - \coth^2(x) + \frac{\coth^2(x)}{2(1 + \tanh(x))} + 2 \int \coth(x) dx \\
&= -\frac{3x}{2} + \frac{3 \coth(x)}{2} - \coth^2(x) + 2 \log(\sinh(x)) + \frac{\coth^2(x)}{2(1 + \tanh(x))}
\end{aligned}$$

Mathematica [A]

time = 0.05, size = 33, normalized size = 0.89

$$\frac{1}{4}(-6x + \cosh(2x) + 4 \coth(x) - 2 \operatorname{csch}^2(x) + 8 \log(\sinh(x)) - \sinh(2x))$$

Antiderivative was successfully verified.

[In] Integrate[Coth[x]^3/(1 + Tanh[x]), x]

[Out] (-6*x + Cosh[2*x] + 4*Coth[x] - 2*Csch[x]^2 + 8*Log[Sinh[x]] - Sinh[2*x])/4

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 74 vs. 2(31) = 62.

time = 0.49, size = 75, normalized size = 2.03

method	result
risch	$-\frac{7x}{2} + \frac{e^{-2x}}{4} - \frac{2}{(e^{2x}-1)^2} + 2 \ln(e^{2x} - 1)$
default	$-\frac{\ln(\tanh(\frac{x}{2})-1)}{2} - \frac{(\tanh^2(\frac{x}{2}))}{8} + \frac{\tanh(\frac{x}{2})}{2} + \frac{1}{(\tanh(\frac{x}{2})+1)^2} - \frac{1}{\tanh(\frac{x}{2})+1} - \frac{7 \ln(\tanh(\frac{x}{2})+1)}{2} - \frac{1}{8 \tanh(\frac{x}{2})^2} + \frac{1}{2 \tanh(\frac{x}{2})}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(coth(x)^3/(1+tanh(x)),x,method=_RETURNVERBOSE)`

[Out] $-1/2*\ln(\tanh(1/2*x)-1)-1/8*\tanh(1/2*x)^2+1/2*\tanh(1/2*x)+1/(\tanh(1/2*x)+1)^2-1/(\tanh(1/2*x)+1)-7/2*\ln(\tanh(1/2*x)+1)-1/8/\tanh(1/2*x)^2+1/2/\tanh(1/2*x)+2*\ln(\tanh(1/2*x))$

Maxima [A]

time = 0.26, size = 54, normalized size = 1.46

$$\frac{1}{2}x + \frac{2(2e^{(-2x)} - 1)}{2e^{(-2x)} - e^{(-4x)} - 1} + \frac{1}{4}e^{(-2x)} + 2 \log(e^{(-x)} + 1) + 2 \log(e^{(-x)} - 1)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(coth(x)^3/(1+tanh(x)),x, algorithm="maxima")`

[Out] $1/2*x + 2*(2*e^{(-2*x)} - 1)/(2*e^{(-2*x)} - e^{(-4*x)} - 1) + 1/4*e^{(-2*x)} + 2*\log(e^{(-x)} + 1) + 2*\log(e^{(-x)} - 1)$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 357 vs. 2(31) = 62.

time = 0.41, size = 357, normalized size = 9.65

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(coth(x)^3/(1+tanh(x)),x, algorithm="fricas")`

[Out] $-1/4*(14*x*\cosh(x)^6 + 84*x*\cosh(x)*\sinh(x)^5 + 14*x*\sinh(x)^6 - (28*x + 1)*\cosh(x)^4 + (210*x*\cosh(x)^2 - 28*x - 1)*\sinh(x)^4 + 4*(70*x*\cosh(x)^3 - (28*x + 1)*\cosh(x))*\sinh(x)^3 + 2*(7*x + 5)*\cosh(x)^2 + 2*(105*x*\cosh(x)^4 - 3*(28*x + 1)*\cosh(x)^2 + 7*x + 5)*\sinh(x)^2 - 8*(\cosh(x)^6 + 6*\cosh(x)*\sinh(x)^5 + \sinh(x)^6 + (15*\cosh(x)^2 - 2)*\sinh(x)^4 - 2*\cosh(x)^4 + 4*(5*\cosh(x)^3 - 2*\cosh(x))*\sinh(x)^3 + (15*\cosh(x)^4 - 12*\cosh(x)^2 + 1)*\sinh(x)^2 + \cosh(x)^2 + 2*(3*\cosh(x)^5 - 4*\cosh(x)^3 + \cosh(x))*\sinh(x))*\log(2*\sinh(x)/(\cosh(x) - \sinh(x))) + 4*(21*x*\cosh(x)^5 - (28*x + 1)*\cosh(x)^3 + (7*x + 5)*\cosh(x))*\sinh(x) - 1)/(\cosh(x)^6 + 6*\cosh(x)*\sinh(x)^5 + \sinh(x)^6 + (15*\cosh(x)^2 - 2)*\sinh(x)^4 - 2*\cosh(x)^4 + 4*(5*\cosh(x)^3 - 2*\cosh(x))*\sinh(x)^3 + (15*\cosh(x)^4 - 12*\cosh(x)^2 + 1)*\sinh(x)^2 + \cosh(x)^2 + 2*(3*\cosh(x)^5 - 4*\cosh(x)^3 + \cosh(x))*\sinh(x))$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\coth^3(x)}{\tanh(x) + 1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(coth(x)**3/(1+tanh(x)),x)`

[Out] `Integral(coth(x)**3/(tanh(x) + 1), x)`

Giac [A]

time = 0.43, size = 40, normalized size = 1.08

$$-\frac{7}{2}x + \frac{(e^{4x} - 10e^{2x} + 1)e^{-2x}}{4(e^{2x} - 1)^2} + 2 \log(|e^{2x} - 1|)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(coth(x)^3/(1+tanh(x)),x, algorithm="giac")`

[Out] `-7/2*x + 1/4*(e^(4*x) - 10*e^(2*x) + 1)*e^(-2*x)/(e^(2*x) - 1)^2 + 2*log(abs(e^(2*x) - 1))`

Mupad [B]

time = 0.08, size = 35, normalized size = 0.95

$$2 \ln(e^{2x} - 1) - \frac{7x}{2} + \frac{e^{-2x}}{4} - \frac{2}{e^{4x} - 2e^{2x} + 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(coth(x)^3/(tanh(x) + 1),x)`

[Out] `2*log(exp(2*x) - 1) - (7*x)/2 + exp(-2*x)/4 - 2/(exp(4*x) - 2*exp(2*x) + 1)`

3.124 $\int \frac{\coth^4(x)}{1+\tanh(x)} dx$

Optimal. Leaf size=43

$$\frac{5x}{2} - \frac{5 \coth(x)}{2} + \coth^2(x) - \frac{5 \coth^3(x)}{6} - 2 \log(\sinh(x)) + \frac{\coth^3(x)}{2(1 + \tanh(x))}$$

[Out] 5/2*x-5/2*coth(x)+coth(x)^2-5/6*coth(x)^3-2*ln(sinh(x))+1/2*coth(x)^3/(1+tanh(x))

Rubi [A]

time = 0.08, antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {3633, 3610, 3612, 3556}

$$\frac{5x}{2} - \frac{5 \coth^3(x)}{6} + \coth^2(x) - \frac{5 \coth(x)}{2} - 2 \log(\sinh(x)) + \frac{\coth^3(x)}{2(\tanh(x) + 1)}$$

Antiderivative was successfully verified.

[In] Int[Coth[x]^4/(1 + Tanh[x]), x]

[Out] (5*x)/2 - (5*Coth[x])/2 + Coth[x]^2 - (5*Coth[x]^3)/6 - 2*Log[Sinh[x]] + Coth[x]^3/(2*(1 + Tanh[x]))

Rule 3556

Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]

Rule 3610

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Dist[1/(a^2 + b^2), Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]

Rule 3612

Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Dist[(b*c - a*d)/(a^2 + b^2), Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]

Rule 3633

```
Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)/((a_.) + (b_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(-a)*((c + d*Tan[e + f*x])^(n + 1)/(2*f*(b*
c - a*d)*(a + b*Tan[e + f*x]))), x] + Dist[1/(2*a*(b*c - a*d)), Int[(c + d*
Tan[e + f*x])^n*Simp[b*c + a*d*(n - 1) - b*d*n*Tan[e + f*x], x], x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] &
& NeQ[c^2 + d^2, 0] && !GtQ[n, 0]
```

Rubi steps

$$\begin{aligned}
\int \frac{\coth^4(x)}{1 + \tanh(x)} dx &= \frac{\coth^3(x)}{2(1 + \tanh(x))} - \frac{1}{2} \int \coth^4(x)(-5 + 4 \tanh(x)) dx \\
&= -\frac{5}{6} \coth^3(x) + \frac{\coth^3(x)}{2(1 + \tanh(x))} - \frac{1}{2} i \int \coth^3(x)(-4i + 5i \tanh(x)) dx \\
&= \coth^2(x) - \frac{5 \coth^3(x)}{6} + \frac{\coth^3(x)}{2(1 + \tanh(x))} + \frac{1}{2} \int \coth^2(x)(5 - 4 \tanh(x)) dx \\
&= -\frac{5 \coth(x)}{2} + \coth^2(x) - \frac{5 \coth^3(x)}{6} + \frac{\coth^3(x)}{2(1 + \tanh(x))} + \frac{1}{2} i \int \coth(x)(4i - 5i \tanh(x)) dx \\
&= \frac{5x}{2} - \frac{5 \coth(x)}{2} + \coth^2(x) - \frac{5 \coth^3(x)}{6} + \frac{\coth^3(x)}{2(1 + \tanh(x))} - 2 \int \coth(x) dx \\
&= \frac{5x}{2} - \frac{5 \coth(x)}{2} + \coth^2(x) - \frac{5 \coth^3(x)}{6} - 2 \log(\sinh(x)) + \frac{\coth^3(x)}{2(1 + \tanh(x))}
\end{aligned}$$

Mathematica [A]

time = 0.10, size = 42, normalized size = 0.98

$$\frac{1}{12}(-3 \cosh(2x) - 4 \coth(x) (7 + \operatorname{csch}^2(x)) + 3(10x + 2 \operatorname{csch}^2(x) - 8 \log(\sinh(x)) + \sinh(2x)))$$

Antiderivative was successfully verified.

[In] Integrate[Coth[x]^4/(1 + Tanh[x]), x]

[Out] (-3*Cosh[2*x] - 4*Coth[x]*(7 + Csch[x]^2) + 3*(10*x + 2*Csch[x]^2 - 8*Log[Sinh[x]] + Sinh[2*x]))/12

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 90 vs. 2(35) = 70.

time = 0.49, size = 91, normalized size = 2.12

method	result
risch	$\frac{9x}{2} - \frac{e^{-2x}}{4} - \frac{2(6e^{4x} - 9e^{2x} + 7)}{3(e^{2x} - 1)^3} - 2 \ln(e^{2x} - 1)$

default	$-\frac{(\tanh^3(\frac{x}{2}))}{24} + \frac{(\tanh^2(\frac{x}{2}))}{8} - \frac{9 \tanh(\frac{x}{2})}{8} - \frac{\ln(\tanh(\frac{x}{2})-1)}{2} - \frac{1}{24 \tanh(\frac{x}{2})^3} + \frac{1}{8 \tanh(\frac{x}{2})^2} - \frac{9}{8 \tanh(\frac{x}{2})} - 2 \ln(\tanh$
---------	--

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(coth(x)^4/(1+tanh(x)),x,method=_RETURNVERBOSE)`

[Out] $-1/24*\tanh(1/2*x)^3+1/8*\tanh(1/2*x)^2-9/8*\tanh(1/2*x)-1/2*\ln(\tanh(1/2*x)-1)-1/24/\tanh(1/2*x)^3+1/8/\tanh(1/2*x)^2-9/8/\tanh(1/2*x)-2*\ln(\tanh(1/2*x))-1/(\tanh(1/2*x)+1)^2+1/(\tanh(1/2*x)+1)+9/2*\ln(\tanh(1/2*x)+1)$

Maxima [A]

time = 0.27, size = 64, normalized size = 1.49

$$\frac{1}{2}x - \frac{2(15e^{(-2x)} - 12e^{(-4x)} - 7)}{3(3e^{(-2x)} - 3e^{(-4x)} + e^{(-6x)} - 1)} - \frac{1}{4}e^{(-2x)} - 2 \log(e^{(-x)} + 1) - 2 \log(e^{(-x)} - 1)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(coth(x)^4/(1+tanh(x)),x, algorithm="maxima")`

[Out] $1/2*x - 2/3*(15*e^{(-2*x)} - 12*e^{(-4*x)} - 7)/(3*e^{(-2*x)} - 3*e^{(-4*x)} + e^{(-6*x)} - 1) - 1/4*e^{(-2*x)} - 2*\log(e^{(-x)} + 1) - 2*\log(e^{(-x)} - 1)$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 582 vs. 2(35) = 70.

time = 0.42, size = 582, normalized size = 13.53

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(coth(x)^4/(1+tanh(x)),x, algorithm="fricas")`

[Out] $1/12*(54*x*\cosh(x)^8 + 432*x*\cosh(x)*\sinh(x)^7 + 54*x*\sinh(x)^8 - 3*(54*x + 17)*\cosh(x)^6 + 3*(504*x*\cosh(x)^2 - 54*x - 17)*\sinh(x)^6 + 18*(168*x*\cosh(x)^3 - (54*x + 17)*\cosh(x))*\sinh(x)^5 + 81*(2*x + 1)*\cosh(x)^4 + 9*(420*x*\cosh(x)^4 - 5*(54*x + 17)*\cosh(x)^2 + 18*x + 9)*\sinh(x)^4 + 12*(252*x*\cosh(x)^5 - 5*(54*x + 17)*\cosh(x)^3 + 27*(2*x + 1)*\cosh(x))*\sinh(x)^3 - (54*x + 65)*\cosh(x)^2 + (1512*x*\cosh(x)^6 - 45*(54*x + 17)*\cosh(x)^4 + 486*(2*x + 1)*\cosh(x)^2 - 54*x - 65)*\sinh(x)^2 - 24*(\cosh(x)^8 + 8*\cosh(x)*\sinh(x)^7 + \sinh(x)^8 + (28*\cosh(x)^2 - 3)*\sinh(x)^6 - 3*\cosh(x)^6 + 2*(28*\cosh(x)^3 - 9*\cosh(x))*\sinh(x)^5 + (70*\cosh(x)^4 - 45*\cosh(x)^2 + 3)*\sinh(x)^4 + 3*\cosh(x)^4 + 4*(14*\cosh(x)^5 - 15*\cosh(x)^3 + 3*\cosh(x))*\sinh(x)^3 + (28*\cosh(x)^6 - 45*\cosh(x)^4 + 18*\cosh(x)^2 - 1)*\sinh(x)^2 - \cosh(x)^2 + 2*(4*\cosh(x)^7 - 9*\cosh(x)^5 + 6*\cosh(x)^3 - \cosh(x))*\sinh(x))*\log(2*\sinh(x)/(\cosh(x) - \sinh(x))) + 2*(216*x*\cosh(x)^7 - 9*(54*x + 17)*\cosh(x)^5 + 162*(2*x + 1)*\cosh(x)^3 - (54*x + 65)*\cosh(x))*\sinh(x) + 3)/(\cosh(x)^8 + 8*\cosh(x)*\sinh(x)^8$

$$7 + \sinh(x)^8 + (28*\cosh(x)^2 - 3)*\sinh(x)^6 - 3*\cosh(x)^6 + 2*(28*\cosh(x)^3 - 9*\cosh(x))*\sinh(x)^5 + (70*\cosh(x)^4 - 45*\cosh(x)^2 + 3)*\sinh(x)^4 + 3*\cosh(x)^4 + 4*(14*\cosh(x)^5 - 15*\cosh(x)^3 + 3*\cosh(x))*\sinh(x)^3 + (28*\cosh(x)^6 - 45*\cosh(x)^4 + 18*\cosh(x)^2 - 1)*\sinh(x)^2 - \cosh(x)^2 + 2*(4*\cosh(x)^7 - 9*\cosh(x)^5 + 6*\cosh(x)^3 - \cosh(x))*\sinh(x)$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\coth^4(x)}{\tanh(x) + 1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(x)**4/(1+tanh(x)),x)

[Out] Integral(coth(x)**4/(tanh(x) + 1), x)

Giac [A]

time = 0.42, size = 48, normalized size = 1.12

$$\frac{9}{2}x - \frac{(51e^{6x} - 81e^{4x} + 65e^{2x} - 3)e^{-2x}}{12(e^{2x} - 1)^3} - 2 \log(|e^{2x} - 1|)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(x)^4/(1+tanh(x)),x, algorithm="giac")

[Out] 9/2*x - 1/12*(51*e^(6*x) - 81*e^(4*x) + 65*e^(2*x) - 3)*e^(-2*x)/(e^(2*x) - 1)^3 - 2*log(abs(e^(2*x) - 1))

Mupad [B]

time = 1.10, size = 69, normalized size = 1.60

$$\frac{9x}{2} - 2 \ln(e^{2x} - 1) - \frac{e^{-2x}}{4} - \frac{8}{3(3e^{2x} - 3e^{4x} + e^{6x} - 1)} - \frac{2}{e^{4x} - 2e^{2x} + 1} - \frac{4}{e^{2x} - 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(coth(x)^4/(tanh(x) + 1),x)

[Out] (9*x)/2 - 2*log(exp(2*x) - 1) - exp(-2*x)/4 - 8/(3*(3*exp(2*x) - 3*exp(4*x) + exp(6*x) - 1)) - 2/(exp(4*x) - 2*exp(2*x) + 1) - 4/(exp(2*x) - 1)

3.125 $\int \tanh(x)(1 + \tanh(x))^{3/2} dx$

Optimal. Leaf size=45

$$2\sqrt{2} \tanh^{-1} \left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}} \right) - 2\sqrt{1 + \tanh(x)} - \frac{2}{3}(1 + \tanh(x))^{3/2}$$

[Out] 2*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))*2^(1/2)-2*(1+tanh(x))^(1/2)-2/3*(1+tanh(x))^(3/2)

Rubi [A]

time = 0.04, antiderivative size = 45, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {3608, 3559, 3561, 212}

$$2\sqrt{2} \tanh^{-1} \left(\frac{\sqrt{\tanh(x) + 1}}{\sqrt{2}} \right) - \frac{2}{3}(\tanh(x) + 1)^{3/2} - 2\sqrt{\tanh(x) + 1}$$

Antiderivative was successfully verified.

[In] Int[Tanh[x]*(1 + Tanh[x])^(3/2), x]

[Out] 2*Sqrt[2]*ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]] - 2*Sqrt[1 + Tanh[x]] - (2*(1 + Tanh[x])^(3/2))/3

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 3559

Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((a + b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] + Dist[2*a, Int[(a + b*Tan[c + d*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && GtQ[n, 1]

Rule 3561

Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Dist[-2*(b/d), Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]

Rule 3608

Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Dist

`[(b*c + a*d)/b, Int[(a + b*Tan[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && !LtQ[m, 0]`

Rubi steps

$$\begin{aligned}
 \int \tanh(x)(1 + \tanh(x))^{3/2} dx &= -\frac{2}{3}(1 + \tanh(x))^{3/2} + \int (1 + \tanh(x))^{3/2} dx \\
 &= -2\sqrt{1 + \tanh(x)} - \frac{2}{3}(1 + \tanh(x))^{3/2} + 2 \int \sqrt{1 + \tanh(x)} dx \\
 &= -2\sqrt{1 + \tanh(x)} - \frac{2}{3}(1 + \tanh(x))^{3/2} + 4\text{Subst}\left(\int \frac{1}{2 - x^2} dx, x, \sqrt{1 + \tanh(x)}\right) \\
 &= 2\sqrt{2} \tanh^{-1}\left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}}\right) - 2\sqrt{1 + \tanh(x)} - \frac{2}{3}(1 + \tanh(x))^{3/2}
 \end{aligned}$$

Mathematica [A]

time = 0.14, size = 39, normalized size = 0.87

$$2\sqrt{2} \tanh^{-1}\left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}}\right) - \frac{2}{3}\sqrt{1 + \tanh(x)}(4 + \tanh(x))$$

Antiderivative was successfully verified.

`[In] Integrate[Tanh[x]*(1 + Tanh[x])^(3/2), x]`

`[Out] 2*Sqrt[2]*ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]] - (2*Sqrt[1 + Tanh[x]]*(4 + Tanh[x]))/3`

Maple [A]

time = 0.50, size = 35, normalized size = 0.78

method	result	size
derivativedivides	$2 \operatorname{arctanh}\left(\frac{\sqrt{1 + \tanh(x)} \sqrt{2}}{2}\right) \sqrt{2} - 2\sqrt{1 + \tanh(x)} - \frac{2(1 + \tanh(x))^{3/2}}{3}$	35
default	$2 \operatorname{arctanh}\left(\frac{\sqrt{1 + \tanh(x)} \sqrt{2}}{2}\right) \sqrt{2} - 2\sqrt{1 + \tanh(x)} - \frac{2(1 + \tanh(x))^{3/2}}{3}$	35

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(tanh(x)*(1+tanh(x))^(3/2), x, method=_RETURNVERBOSE)`

`[Out] 2*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))*2^(1/2)-2*(1+tanh(x))^(1/2)-2/3*(1+tanh(x))^(3/2)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(tanh(x)*(1+tanh(x))^(3/2),x, algorithm="maxima")``[Out] -2/3*sqrt(2)/(e^(-2*x) + 1)^(3/2) + integrate(2*sqrt(2)*e^(-x)/((e^(-x) + e^(-3*x))*(e^(-2*x) + 1)^(3/2)), x)`**Fricas [B]** Leaf count of result is larger than twice the leaf count of optimal. 252 vs. 2(34) = 68.

time = 0.60, size = 252, normalized size = 5.60

$$\frac{2\sqrt{2}(\sqrt{2}\cosh(x)^2 + 1)\sqrt{2}\cosh(x)\sinh(x)^2 + 5\sqrt{2}\sinh(x)^2 + 3(\sqrt{2}\cosh(x)^2 + \sqrt{2})\sinh(x) + 3\sqrt{2}\cosh(x)\sqrt{\frac{\cosh(x)}{\cosh(x) + \sinh(x)}} - 3(\sqrt{2}\cosh(x)^2 + 4\sqrt{2}\cosh(x)\sinh(x) + \sqrt{2}\sinh(x)^2 + 2(3\sqrt{2}\cosh(x)^2 + \sqrt{2})\sinh(x) + 2\sqrt{2}\cosh(x)^2 + 4(\sqrt{2}\cosh(x)^2 + \sqrt{2}\cosh(x)\sinh(x) + \sqrt{2}))\log\left(\frac{-2\sqrt{2}\sqrt{\frac{\cosh(x)}{\cosh(x) + \sinh(x)}}(\cosh(x) + \sinh(x) - 2\cosh(x)^2 - 4\sinh(x)\sinh(x) - 2\sinh(x)^2 - 1)}{3(\cosh(x)^2 + 4\cosh(x)\sinh(x) + \sinh(x)^2 + 2(3\cosh(x)^2 + 2\sinh(x)^2 + 2(\cosh(x) + \sinh(x))\sinh(x) + 1))}\right)}{3(\cosh(x)^2 + 4\cosh(x)\sinh(x) + \sinh(x)^2 + 2(3\cosh(x)^2 + 2\sinh(x)^2 + 2(\cosh(x) + \sinh(x))\sinh(x) + 1))}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(tanh(x)*(1+tanh(x))^(3/2),x, algorithm="fricas")`

```
[Out] -1/3*(2*sqrt(2)*(5*sqrt(2)*cosh(x)^3 + 15*sqrt(2)*cosh(x)*sinh(x)^2 + 5*sqrt(2)*sinh(x)^3 + 3*(5*sqrt(2)*cosh(x)^2 + sqrt(2))*sinh(x) + 3*sqrt(2)*cosh(x))*sqrt(cosh(x)/(cosh(x) - sinh(x))) - 3*(sqrt(2)*cosh(x)^4 + 4*sqrt(2)*cosh(x)*sinh(x)^3 + sqrt(2)*sinh(x)^4 + 2*(3*sqrt(2)*cosh(x)^2 + sqrt(2))*sinh(x)^2 + 2*sqrt(2)*cosh(x)^2 + 4*(sqrt(2)*cosh(x)^3 + sqrt(2)*cosh(x))*sinh(x) + sqrt(2))*log(-2*sqrt(2)*sqrt(cosh(x)/(cosh(x) - sinh(x)))*(cosh(x) + sinh(x) - 2*cosh(x)^2 - 4*cosh(x)*sinh(x) - 2*sinh(x)^2 - 1))/(cosh(x)^4 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4 + 2*(3*cosh(x)^2 + 1)*sinh(x)^2 + 2*cosh(x)^2 + 4*(cosh(x)^3 + cosh(x))*sinh(x) + 1)
```

Sympy [A]

time = 6.72, size = 78, normalized size = 1.73

$$-\frac{2(\tanh(x) + 1)^{\frac{3}{2}}}{3} - 2\sqrt{\tanh(x) + 1} - 4 \left(\begin{cases} -\frac{\sqrt{2} \operatorname{acoth}\left(\frac{\sqrt{2}\sqrt{\tanh(x) + 1}}{2}\right)}{2} & \text{for } \tanh(x) > 1 \\ -\frac{\sqrt{2} \operatorname{atanh}\left(\frac{\sqrt{2}\sqrt{\tanh(x) + 1}}{2}\right)}{2} & \text{for } \tanh(x) < 1 \end{cases} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(tanh(x)*(1+tanh(x))^(3/2),x)`

```
[Out] -2*(tanh(x) + 1)^(3/2)/3 - 2*sqrt(tanh(x) + 1) - 4*Piecewise((-sqrt(2)*acoth(sqrt(2)*sqrt(tanh(x) + 1)/2)/2, tanh(x) > 1), (-sqrt(2)*atanh(sqrt(2)*sqrt(tanh(x) + 1)/2)/2, tanh(x) < 1))
```

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 96 vs. $2(34) = 68$.
time = 0.44, size = 96, normalized size = 2.13

$$\frac{1}{3} \sqrt{2} \left(\frac{2 \left(9 \left(\sqrt{e^{4x} + e^{2x}} - e^{2x} \right)^2 - 12 \sqrt{e^{4x} + e^{2x}} + 12 e^{2x} + 5 \right)}{\left(\sqrt{e^{4x} + e^{2x}} - e^{2x} - 1 \right)^3} - 3 \log \left(-2 \sqrt{e^{4x} + e^{2x}} + 2 e^{2x} + 1 \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)*(1+tanh(x))^(3/2),x, algorithm="giac")

[Out] 1/3*sqrt(2)*(2*(9*(sqrt(e^(4*x) + e^(2*x)) - e^(2*x))^2 - 12*sqrt(e^(4*x) + e^(2*x)) + 12*e^(2*x) + 5)/(sqrt(e^(4*x) + e^(2*x)) - e^(2*x) - 1)^3 - 3*log(-2*sqrt(e^(4*x) + e^(2*x)) + 2*e^(2*x) + 1))

Mupad [B]

time = 0.15, size = 34, normalized size = 0.76

$$2 \sqrt{2} \operatorname{atanh} \left(\frac{\sqrt{2} \sqrt{\tanh(x) + 1}}{2} \right) - 2 \sqrt{\tanh(x) + 1} - \frac{2 (\tanh(x) + 1)^{3/2}}{3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(x)*(tanh(x) + 1)^(3/2),x)

[Out] 2*2^(1/2)*atanh((2^(1/2)*(tanh(x) + 1)^(1/2))/2) - 2*(tanh(x) + 1)^(1/2) - (2*(tanh(x) + 1)^(3/2))/3

3.126 $\int \tanh(x) \sqrt{1 + \tanh(x)} dx$

Optimal. Leaf size=32

$$\sqrt{2} \tanh^{-1} \left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}} \right) - 2\sqrt{1 + \tanh(x)}$$

[Out] arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))*2^(1/2)-2*(1+tanh(x))^(1/2)

Rubi [A]

time = 0.03, antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3608, 3561, 212}

$$\sqrt{2} \tanh^{-1} \left(\frac{\sqrt{\tanh(x) + 1}}{\sqrt{2}} \right) - 2\sqrt{\tanh(x) + 1}$$

Antiderivative was successfully verified.

[In] Int[Tanh[x]*Sqrt[1 + Tanh[x]],x]

[Out] Sqrt[2]*ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]] - 2*Sqrt[1 + Tanh[x]]

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 3561

Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Dist[-2*(b/d), Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]

Rule 3608

Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Dist[(b*c + a*d)/b, Int[(a + b*Tan[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && !LtQ[m, 0]

Rubi steps

$$\begin{aligned}
\int \tanh(x) \sqrt{1 + \tanh(x)} \, dx &= -2\sqrt{1 + \tanh(x)} + \int \sqrt{1 + \tanh(x)} \, dx \\
&= -2\sqrt{1 + \tanh(x)} + 2\text{Subst}\left(\int \frac{1}{2-x^2} \, dx, x, \sqrt{1 + \tanh(x)}\right) \\
&= \sqrt{2} \tanh^{-1}\left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}}\right) - 2\sqrt{1 + \tanh(x)}
\end{aligned}$$

Mathematica [A]

time = 0.04, size = 32, normalized size = 1.00

$$\sqrt{2} \tanh^{-1}\left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}}\right) - 2\sqrt{1 + \tanh(x)}$$

Antiderivative was successfully verified.

`[In] Integrate[Tanh[x]*Sqrt[1 + Tanh[x]], x]``[Out] Sqrt[2]*ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]] - 2*Sqrt[1 + Tanh[x]]`**Maple [A]**

time = 0.57, size = 26, normalized size = 0.81

method	result	size
derivativeldivides	$\text{arctanh}\left(\frac{\sqrt{1 + \tanh(x)} \sqrt{2}}{2}\right) \sqrt{2} - 2\sqrt{1 + \tanh(x)}$	26
default	$\text{arctanh}\left(\frac{\sqrt{1 + \tanh(x)} \sqrt{2}}{2}\right) \sqrt{2} - 2\sqrt{1 + \tanh(x)}$	26

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((1+tanh(x))^(1/2)*tanh(x), x, method=_RETURNVERBOSE)``[Out] arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))*2^(1/2)-2*(1+tanh(x))^(1/2)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((1+tanh(x))^(1/2)*tanh(x), x, algorithm="maxima")`

[Out] $-\sqrt{2}/\sqrt{e^{-2x} + 1} + \text{integrate}(\sqrt{2}*e^{-x}/((e^{-x} + e^{-3x}))*\sqrt{e^{-2x} + 1}), x)$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 129 vs. 2(25) = 50.

time = 0.46, size = 129, normalized size = 4.03

$$\frac{4\sqrt{2}(\sqrt{2}\cosh(x) + \sqrt{2}\sinh(x))\sqrt{\frac{\cosh(x)}{\cosh(x) - \sinh(x)}} - (\sqrt{2}\cosh(x)^2 + 2\sqrt{2}\cosh(x)\sinh(x) + \sqrt{2}\sinh(x)^2 + \sqrt{2})\log\left(-2\sqrt{2}\sqrt{\frac{\cosh(x)}{\cosh(x) - \sinh(x)}}(\cosh(x) + \sinh(x)) - 2\cosh(x)^2 - 4\cosh(x)\sinh(x) - 2\sinh(x)^2 - 1\right)}{2(\cosh(x)^2 + 2\cosh(x)\sinh(x) + \sinh(x)^2 + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{integrate}((1+\tanh(x))^{1/2}*\tanh(x),x, \text{algorithm}=\text{"fricas"})$

[Out] $-1/2*(4*\sqrt{2}*(\sqrt{2}*\cosh(x) + \sqrt{2}*\sinh(x))*\sqrt{\cosh(x)/(\cosh(x) - \sinh(x))} - (\sqrt{2}*\cosh(x)^2 + 2*\sqrt{2}*\cosh(x)*\sinh(x) + \sqrt{2}*\sinh(x)^2 + \sqrt{2})*\log(-2*\sqrt{2}*\sqrt{\cosh(x)/(\cosh(x) - \sinh(x))}*(\cosh(x) + \sinh(x)) - 2*\cosh(x)^2 - 4*\cosh(x)*\sinh(x) - 2*\sinh(x)^2 - 1))/(\cosh(x)^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2 + 1)$

Sympy [A]

time = 1.29, size = 66, normalized size = 2.06

$$-2\sqrt{\tanh(x) + 1} - 2 \left(\begin{array}{l} \left(\frac{\sqrt{2} \operatorname{acoth}\left(\frac{\sqrt{2}\sqrt{\tanh(x) + 1}}{2}\right)}{2} \right) \text{ for } \tanh(x) > 1 \\ \left(-\frac{\sqrt{2} \operatorname{atanh}\left(\frac{\sqrt{2}\sqrt{\tanh(x) + 1}}{2}\right)}{2} \right) \text{ for } \tanh(x) < 1 \end{array} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{integrate}((1+\tanh(x))^{1/2}*\tanh(x),x)$

[Out] $-2*\sqrt{\tanh(x) + 1} - 2*\text{Piecewise}((- \sqrt{2}*\operatorname{acoth}(\sqrt{2}*\sqrt{\tanh(x) + 1})/2, \tanh(x) > 1), (- \sqrt{2}*\operatorname{atanh}(\sqrt{2}*\sqrt{\tanh(x) + 1})/2, \tanh(x) < 1))$

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 53 vs. 2(25) = 50.

time = 0.42, size = 53, normalized size = 1.66

$$\frac{1}{2}\sqrt{2} \left(\frac{4}{\sqrt{e^{4x} + e^{2x}} - e^{2x} - 1} - \log\left(-2\sqrt{e^{4x} + e^{2x}} + 2e^{2x} + 1\right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{integrate}((1+\tanh(x))^{1/2}*\tanh(x),x, \text{algorithm}=\text{"giac"})$

[Out] $1/2*\sqrt{2}*(4/(\sqrt{e^{4x} + e^{2x}} - e^{2x} - 1) - \log(-2*\sqrt{e^{4x} + e^{2x}} + 2*e^{2x} + 1))$

Mupad [B]

time = 1.06, size = 25, normalized size = 0.78

$$\sqrt{2} \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{\tanh(x) + 1}}{2}\right) - 2 \sqrt{\tanh(x) + 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tanh(x)*(tanh(x) + 1)^(1/2),x)`

[Out] `2^(1/2)*atanh((2^(1/2)*(tanh(x) + 1)^(1/2))/2) - 2*(tanh(x) + 1)^(1/2)`

$$3.127 \quad \int \frac{\tanh(x)}{\sqrt{1 + \tanh(x)}} dx$$

Optimal. Leaf size=30

$$\frac{\tanh^{-1}\left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}}\right)}{\sqrt{2}} + \frac{1}{\sqrt{1 + \tanh(x)}}$$

[Out] 1/2*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))*2^(1/2)+1/(1+tanh(x))^(1/2)

Rubi [A]

time = 0.03, antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3607, 3561, 212}

$$\frac{\tanh^{-1}\left(\frac{\sqrt{\tanh(x) + 1}}{\sqrt{2}}\right)}{\sqrt{2}} + \frac{1}{\sqrt{\tanh(x) + 1}}$$

Antiderivative was successfully verified.

[In] Int[Tanh[x]/Sqrt[1 + Tanh[x]],x]

[Out] ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]]/Sqrt[2] + 1/Sqrt[1 + Tanh[x]]

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 3561

Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Dist[-2*(b/d), Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]

Rule 3607

Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*((a + b*Tan[e + f*x])^m/(2*a*f*m)), x] + Dist[(b*c + a*d)/(2*a*b), Int[(a + b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0]

Rubi steps

$$\begin{aligned}
\int \frac{\tanh(x)}{\sqrt{1+\tanh(x)}} dx &= \frac{1}{\sqrt{1+\tanh(x)}} + \frac{1}{2} \int \sqrt{1+\tanh(x)} dx \\
&= \frac{1}{\sqrt{1+\tanh(x)}} + \text{Subst} \left(\int \frac{1}{2-x^2} dx, x, \sqrt{1+\tanh(x)} \right) \\
&= \frac{\tanh^{-1} \left(\frac{\sqrt{1+\tanh(x)}}{\sqrt{2}} \right)}{\sqrt{2}} + \frac{1}{\sqrt{1+\tanh(x)}}
\end{aligned}$$

Mathematica [A]

time = 0.04, size = 30, normalized size = 1.00

$$\frac{\tanh^{-1} \left(\frac{\sqrt{1+\tanh(x)}}{\sqrt{2}} \right)}{\sqrt{2}} + \frac{1}{\sqrt{1+\tanh(x)}}$$

Antiderivative was successfully verified.

`[In] Integrate[Tanh[x]/Sqrt[1 + Tanh[x]], x]``[Out] ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]]/Sqrt[2] + 1/Sqrt[1 + Tanh[x]]`**Maple [A]**

time = 0.67, size = 25, normalized size = 0.83

method	result	size
derivativedivides	$\frac{\operatorname{arctanh} \left(\frac{\sqrt{1+\tanh(x)} \sqrt{2}}{2} \right) \sqrt{2}}{2} + \frac{1}{\sqrt{1+\tanh(x)}}$	25
default	$\frac{\operatorname{arctanh} \left(\frac{\sqrt{1+\tanh(x)} \sqrt{2}}{2} \right) \sqrt{2}}{2} + \frac{1}{\sqrt{1+\tanh(x)}}$	25

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(tanh(x)/(1+tanh(x))^(1/2), x, method=_RETURNVERBOSE)``[Out] 1/2*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))*2^(1/2)+1/(1+tanh(x))^(1/2)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)/(1+tanh(x))^(1/2),x, algorithm="maxima")

[Out] 1/2*sqrt(2)*sqrt(e^(-2*x) + 1) + integrate(e^(-x)/(sqrt(2)*e^(-x)/sqrt(e^(-2*x) + 1) + sqrt(2)*e^(-3*x)/sqrt(e^(-2*x) + 1)), x)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 85 vs. 2(24) = 48.

time = 0.34, size = 85, normalized size = 2.83

$$\frac{(\sqrt{2} \cosh(x) + \sqrt{2} \sinh(x)) \log\left(-2\sqrt{2} \sqrt{\frac{\cosh(x)}{\cosh(x) - \sinh(x)}} (\cosh(x) + \sinh(x)) - 2 \cosh(x)^2 - 4 \cosh(x) \sinh(x) - 2 \sinh(x)^2 - 1\right) + 4 \sqrt{\frac{\cosh(x)}{\cosh(x) - \sinh(x)}}}{4(\cosh(x) + \sinh(x))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)/(1+tanh(x))^(1/2),x, algorithm="fricas")

[Out] 1/4*((sqrt(2)*cosh(x) + sqrt(2)*sinh(x))*log(-2*sqrt(2)*sqrt(cosh(x)/(cosh(x) - sinh(x)))*(cosh(x) + sinh(x)) - 2*cosh(x)^2 - 4*cosh(x)*sinh(x) - 2*sinh(x)^2 - 1) + 4*sqrt(cosh(x)/(cosh(x) - sinh(x))))/(cosh(x) + sinh(x))

Sympy [A]

time = 1.56, size = 63, normalized size = 2.10

$$-\begin{cases} \frac{\sqrt{2} \operatorname{acoth}\left(\frac{\sqrt{2} \sqrt{\tanh(x) + 1}}{2}\right)}{2} & \text{for } \tanh(x) > 1 \\ \frac{\sqrt{2} \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{\tanh(x) + 1}}{2}\right)}{2} & \text{for } \tanh(x) < 1 \end{cases} + \frac{1}{\sqrt{\tanh(x) + 1}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)/(1+tanh(x))^(1/2),x)

[Out] -Piecewise((-sqrt(2)*acoth(sqrt(2)*sqrt(tanh(x) + 1)/2)/2, tanh(x) > 1), (-sqrt(2)*atanh(sqrt(2)*sqrt(tanh(x) + 1)/2)/2, tanh(x) < 1)) + 1/sqrt(tanh(x) + 1)

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 52 vs. 2(24) = 48.
time = 0.41, size = 52, normalized size = 1.73

$$\frac{1}{4} \sqrt{2} \left(\frac{2}{\sqrt{e^{4x} + e^{2x}} - e^{2x}} - \log\left(-2 \sqrt{e^{4x} + e^{2x}} + 2e^{2x} + 1\right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)/(1+tanh(x))^(1/2),x, algorithm="giac")

[Out] $\frac{1}{4}\sqrt{2}\left(\frac{2}{\sqrt{e^{4x} + e^{2x}} - e^{2x}} - \log(-2\sqrt{e^{4x} + e^{2x}} + 2e^{2x} + 1)\right)$

Mupad [B]

time = 0.13, size = 24, normalized size = 0.80

$$\frac{\sqrt{2} \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{\tanh(x) + 1}}{2}\right)}{2} + \frac{1}{\sqrt{\tanh(x) + 1}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tanh(x)/(tanh(x) + 1)^(1/2),x)`

[Out] $(2^{1/2} \operatorname{atanh}((2^{1/2}(\tanh(x) + 1)^{1/2})/2))/2 + 1/(\tanh(x) + 1)^{1/2}$

$$3.128 \quad \int \frac{\tanh(x)}{(1+\tanh(x))^{3/2}} dx$$

Optimal. Leaf size=49

$$\frac{\tanh^{-1}\left(\frac{\sqrt{1+\tanh(x)}}{\sqrt{2}}\right)}{2\sqrt{2}} + \frac{1}{3(1+\tanh(x))^{3/2}} - \frac{1}{2\sqrt{1+\tanh(x)}}$$

[Out] 1/4*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))*2^(1/2)-1/2/(1+tanh(x))^(1/2)+1/3/(1+tanh(x))^(3/2)

Rubi [A]

time = 0.04, antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {3607, 3560, 3561, 212}

$$\frac{\tanh^{-1}\left(\frac{\sqrt{\tanh(x)+1}}{\sqrt{2}}\right)}{2\sqrt{2}} - \frac{1}{2\sqrt{\tanh(x)+1}} + \frac{1}{3(\tanh(x)+1)^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[Tanh[x]/(1 + Tanh[x])^(3/2), x]

[Out] ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]]/(2*Sqrt[2]) + 1/(3*(1 + Tanh[x])^(3/2)) - 1/(2*Sqrt[1 + Tanh[x]])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 3560

Int[((a_) + (b_.)*tan[(c_) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n), x] + Dist[1/(2*a), Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]

Rule 3561

Int[Sqrt[(a_) + (b_.)*tan[(c_) + (d_.)*(x_)]], x_Symbol] := Dist[-2*(b/d), Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]

Rule 3607

```
Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*((a + b*Tan[e + f*x])^m/(2*a
*f*m)), x] + Dist[(b*c + a*d)/(2*a*b), Int[(a + b*Tan[e + f*x])^(m + 1), x]
, x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2,
0] && LtQ[m, 0]
```

Rubi steps

$$\begin{aligned}
\int \frac{\tanh(x)}{(1 + \tanh(x))^{3/2}} dx &= \frac{1}{3(1 + \tanh(x))^{3/2}} + \frac{1}{2} \int \frac{1}{\sqrt{1 + \tanh(x)}} dx \\
&= \frac{1}{3(1 + \tanh(x))^{3/2}} - \frac{1}{2\sqrt{1 + \tanh(x)}} + \frac{1}{4} \int \sqrt{1 + \tanh(x)} dx \\
&= \frac{1}{3(1 + \tanh(x))^{3/2}} - \frac{1}{2\sqrt{1 + \tanh(x)}} + \frac{1}{2} \text{Subst}\left(\int \frac{1}{2 - x^2} dx, x, \sqrt{1 + \tanh(x)}\right) \\
&= \frac{\tanh^{-1}\left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}}\right)}{2\sqrt{2}} + \frac{1}{3(1 + \tanh(x))^{3/2}} - \frac{1}{2\sqrt{1 + \tanh(x)}}
\end{aligned}$$

Mathematica [A]

time = 0.09, size = 51, normalized size = 1.04

$$\frac{1}{12} \left(3\sqrt{2} \tanh^{-1}\left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}}\right) - \frac{2(\cosh(x) - \sinh(x))(\cosh(x) + 3\sinh(x))}{\sqrt{1 + \tanh(x)}} \right)$$

Antiderivative was successfully verified.

[In] Integrate[Tanh[x]/(1 + Tanh[x])^(3/2), x]

[Out] (3*Sqrt[2]*ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]] - (2*(Cosh[x] - Sinh[x])*(Cosh[x] + 3*Sinh[x]))/Sqrt[1 + Tanh[x]])/12

Maple [A]

time = 0.59, size = 35, normalized size = 0.71

method	result	size
derivativedivides	$ \frac{\arctanh\left(\frac{\sqrt{1 + \tanh(x)}\sqrt{2}}{2}\right)\sqrt{2}}{4} - \frac{1}{2\sqrt{1 + \tanh(x)}} + \frac{1}{3(1 + \tanh(x))^{3/2}} $	35

default	$\frac{\operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}\sqrt{2}}{2}\right)\sqrt{2}}{4} - \frac{1}{2\sqrt{1+\tanh(x)}} + \frac{1}{3(1+\tanh(x))^{\frac{3}{2}}}$	35
---------	---	----

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tanh(x)/(1+tanh(x))^(3/2),x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{4}\operatorname{arctanh}\left(\frac{1}{2}(1+\tanh(x))^{1/2}\right)2^{1/2} - \frac{1}{2(1+\tanh(x))^{1/2}} + \frac{1}{3(1+\tanh(x))^{3/2}}$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(x)/(1+tanh(x))^(3/2),x, algorithm="maxima")`

[Out] $\frac{1}{12}\sqrt{2}(e^{-2x} + 1)^{3/2} + \int \frac{1}{2}e^{-x}/(\sqrt{2}e^{-x}/(e^{-2x} + 1)^{3/2} + \sqrt{2}e^{-3x}/(e^{-2x} + 1)^{3/2}) dx$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 168 vs. 2(34) = 68.

time = 0.41, size = 168, normalized size = 3.43

$$\frac{2\sqrt{2}(2\sqrt{2}\cosh(x)^2 + 4\sqrt{2}\cosh(x)\sinh(x) + 2\sqrt{2}\sinh(x)^2 - \sqrt{2})\sqrt{\frac{\cosh(x)}{\cosh(x) - \sinh(x)}} - 3(\sqrt{2}\cosh(x)^3 + 3\sqrt{2}\cosh(x)^2\sinh(x) + 3\sqrt{2}\cosh(x)\sinh(x)^2 + \sqrt{2}\sinh(x)^3)\log\left(\frac{-2\sqrt{2}\sqrt{\frac{\cosh(x)}{\cosh(x) - \sinh(x)}}(\cosh(x) + \sinh(x)) - 2\cosh(x)^2 - 4\cosh(x)\sinh(x) - 2\sinh(x)^2 - 1}{24(\cosh(x)^3 + 3\cosh(x)^2\sinh(x) + 3\cosh(x)\sinh(x)^2 + \sinh(x)^3)}\right)}{24(\cosh(x)^3 + 3\cosh(x)^2\sinh(x) + 3\cosh(x)\sinh(x)^2 + \sinh(x)^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(x)/(1+tanh(x))^(3/2),x, algorithm="fricas")`

[Out] $-\frac{1}{24}(2\sqrt{2}(2\sqrt{2}\cosh(x)^2 + 4\sqrt{2}\cosh(x)\sinh(x) + 2\sqrt{2}\sinh(x)^2 - \sqrt{2})\sqrt{\frac{\cosh(x)}{\cosh(x) - \sinh(x)}} - 3(\sqrt{2}\cosh(x)^3 + 3\sqrt{2}\cosh(x)^2\sinh(x) + 3\sqrt{2}\cosh(x)\sinh(x)^2 + \sqrt{2}\sinh(x)^3)\log\left(\frac{-2\sqrt{2}\sqrt{\frac{\cosh(x)}{\cosh(x) - \sinh(x)}}(\cosh(x) + \sinh(x)) - 2\cosh(x)^2 - 4\cosh(x)\sinh(x) - 2\sinh(x)^2 - 1}{24(\cosh(x)^3 + 3\cosh(x)^2\sinh(x) + 3\cosh(x)\sinh(x)^2 + \sinh(x)^3)}\right))$

Sympy [A]

time = 6.78, size = 78, normalized size = 1.59

$$\frac{\begin{cases} \frac{\sqrt{2}\operatorname{acoth}\left(\frac{\sqrt{2}\sqrt{\tanh(x)+1}}{2}\right)}{2} & \text{for } \tanh(x) > 1 \\ \frac{\sqrt{2}\operatorname{atanh}\left(\frac{\sqrt{2}\sqrt{\tanh(x)+1}}{2}\right)}{2} & \text{for } \tanh(x) < 1 \end{cases}}{2} - \frac{1}{2\sqrt{\tanh(x)+1}} + \frac{1}{3(\tanh(x)+1)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)/(1+tanh(x))**(3/2),x)

[Out] -Piecewise((-sqrt(2)*acoth(sqrt(2)*sqrt(tanh(x) + 1)/2)/2, tanh(x) > 1), (-sqrt(2)*atanh(sqrt(2)*sqrt(tanh(x) + 1)/2)/2, tanh(x) < 1))/2 - 1/(2*sqrt(tanh(x) + 1)) + 1/(3*(tanh(x) + 1)**(3/2))

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 73 vs. 2(34) = 68.
time = 0.43, size = 73, normalized size = 1.49

$$-\frac{1}{24} \sqrt{2} \left(\frac{2 \left(3 \sqrt{e^{4x} + e^{2x}} - 3e^{2x} - 1 \right)}{\left(\sqrt{e^{4x} + e^{2x}} - e^{2x} \right)^3} + 3 \log \left(-2 \sqrt{e^{4x} + e^{2x}} + 2e^{2x} + 1 \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)/(1+tanh(x))^(3/2),x, algorithm="giac")

[Out] -1/24*sqrt(2)*(2*(3*sqrt(e^(4*x) + e^(2*x)) - 3*e^(2*x) - 1)/(sqrt(e^(4*x) + e^(2*x)) - e^(2*x))^3 + 3*log(-2*sqrt(e^(4*x) + e^(2*x)) + 2*e^(2*x) + 1))

Mupad [B]

time = 1.09, size = 32, normalized size = 0.65

$$\frac{\sqrt{2} \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{\tanh(x) + 1}}{2}\right)}{4} - \frac{\frac{\tanh(x)}{2} + \frac{1}{6}}{(\tanh(x) + 1)^{3/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(x)/(tanh(x) + 1)^(3/2),x)

[Out] (2^(1/2)*atanh((2^(1/2)*(tanh(x) + 1)^(1/2))/2))/4 - (tanh(x)/2 + 1/6)/(tanh(x) + 1)^(3/2)

3.129 $\int \tanh^2(x)(1 + \tanh(x))^{3/2} dx$

Optimal. Leaf size=45

$$2\sqrt{2} \tanh^{-1}\left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}}\right) - 2\sqrt{1 + \tanh(x)} - \frac{2}{5}(1 + \tanh(x))^{5/2}$$

[Out] 2*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))*2^(1/2)-2*(1+tanh(x))^(1/2)-2/5*(1+tanh(x))^(5/2)

Rubi [A]

time = 0.04, antiderivative size = 45, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {3624, 3559, 3561, 212}

$$2\sqrt{2} \tanh^{-1}\left(\frac{\sqrt{\tanh(x) + 1}}{\sqrt{2}}\right) - \frac{2}{5}(\tanh(x) + 1)^{5/2} - 2\sqrt{\tanh(x) + 1}$$

Antiderivative was successfully verified.

[In] Int[Tanh[x]^2*(1 + Tanh[x])^(3/2), x]

[Out] 2*Sqrt[2]*ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]] - 2*Sqrt[1 + Tanh[x]] - (2*(1 + Tanh[x])^(5/2))/5

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 3559

Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((a + b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] + Dist[2*a, Int[(a + b*Tan[c + d*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && GtQ[n, 1]

Rule 3561

Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Dist[-2*(b/d), Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]

Rule 3624

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[d^2*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(

```
m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[c^2 - d^2 + 2*c*d*Tan[e + f*
x], x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && !LeQ
[m, -1] && !(EqQ[m, 2] && EqQ[a, 0])
```

Rubi steps

$$\begin{aligned} \int \tanh^2(x)(1 + \tanh(x))^{3/2} dx &= -\frac{2}{5}(1 + \tanh(x))^{5/2} + \int (1 + \tanh(x))^{3/2} dx \\ &= -2\sqrt{1 + \tanh(x)} - \frac{2}{5}(1 + \tanh(x))^{5/2} + 2 \int \sqrt{1 + \tanh(x)} dx \\ &= -2\sqrt{1 + \tanh(x)} - \frac{2}{5}(1 + \tanh(x))^{5/2} + 4 \operatorname{Subst}\left(\int \frac{1}{2 - x^2} dx, x, \sqrt{1 + \tanh(x)}\right) \\ &= 2\sqrt{2} \tanh^{-1}\left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}}\right) - 2\sqrt{1 + \tanh(x)} - \frac{2}{5}(1 + \tanh(x))^{5/2} \end{aligned}$$

Mathematica [A]

time = 0.14, size = 53, normalized size = 1.18

$$2\sqrt{2} \tanh^{-1}\left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}}\right) - \frac{1}{5} \operatorname{sech}^2(x)(5 + 7 \cosh(2x) + 2 \sinh(2x)) \sqrt{1 + \tanh(x)}$$

Antiderivative was successfully verified.

```
[In] Integrate[Tanh[x]^2*(1 + Tanh[x])^(3/2), x]
```

```
[Out] 2*Sqrt[2]*ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]] - (Sech[x]^2*(5 + 7*Cosh[2*x]
+ 2*Sinh[2*x])*Sqrt[1 + Tanh[x]])/5
```

Maple [A]

time = 0.63, size = 35, normalized size = 0.78

method	result	size
derivativedivides	$2 \operatorname{arctanh}\left(\frac{\sqrt{1 + \tanh(x)} \sqrt{2}}{2}\right) \sqrt{2} - 2\sqrt{1 + \tanh(x)} - \frac{2(1 + \tanh(x))^{5/2}}{5}$	35
default	$2 \operatorname{arctanh}\left(\frac{\sqrt{1 + \tanh(x)} \sqrt{2}}{2}\right) \sqrt{2} - 2\sqrt{1 + \tanh(x)} - \frac{2(1 + \tanh(x))^{5/2}}{5}$	35

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(tanh(x)^2*(1+tanh(x))^(3/2), x, method=_RETURNVERBOSE)
```

[Out] $2 \operatorname{arctanh}\left(\frac{1}{2}(1+\tanh(x))^{1/2}\right) 2^{1/2} - 2(1+\tanh(x))^{1/2} - \frac{2}{5}(1+\tanh(x))^{5/2}$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(x)^2*(1+tanh(x))^(3/2),x, algorithm="maxima")`

[Out] `integrate((tanh(x) + 1)^(3/2)*tanh(x)^2, x)`

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 429 vs. $2(34) = 68$.

time = 0.36, size = 429, normalized size = 9.53

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(x)^2*(1+tanh(x))^(3/2),x, algorithm="fricas")`

[Out]
$$-1/5(2\sqrt{2}(9\sqrt{2}\cosh(x)^5 + 45\sqrt{2}\cosh(x)\sinh(x)^4 + 9\sqrt{2}\sinh(x)^5 + 10(9\sqrt{2}\cosh(x)^2 + \sqrt{2})\sinh(x)^3 + 10\sqrt{2}\cosh(x)^3 + 30(3\sqrt{2}\cosh(x)^3 + \sqrt{2}\cosh(x))\sinh(x)^2 + 5(9\sqrt{2}\cosh(x)^4 + 6\sqrt{2}\cosh(x)^2 + \sqrt{2})\sinh(x) + 5\sqrt{2}\cosh(x))\sqrt{\cosh(x)/(\cosh(x) - \sinh(x))} - 5(\sqrt{2}\cosh(x)^6 + 6\sqrt{2}\cosh(x)\sinh(x)^5 + \sqrt{2}\sinh(x)^6 + 3(5\sqrt{2}\cosh(x)^2 + \sqrt{2})\sinh(x)^4 + 3\sqrt{2}\cosh(x)^4 + 4(5\sqrt{2}\cosh(x)^3 + 3\sqrt{2}\cosh(x))\sinh(x)^3 + 3(5\sqrt{2}\cosh(x)^4 + 6\sqrt{2}\cosh(x)^2 + \sqrt{2})\sinh(x)^2 + 3\sqrt{2}\cosh(x)^2 + 6(\sqrt{2}\cosh(x)^5 + 2\sqrt{2}\cosh(x)^3 + \sqrt{2}\cosh(x))\sinh(x) + \sqrt{2})\log(-2\sqrt{2}\sqrt{\cosh(x)/(\cosh(x) - \sinh(x))}(\cosh(x) + \sinh(x)) - 2\cosh(x)^2 - 4\cosh(x)\sinh(x) - 2\sinh(x)^2 - 1)/(\cosh(x)^6 + 6\cosh(x)\sinh(x)^5 + \sinh(x)^6 + 3(5\cosh(x)^2 + 1)\sinh(x)^4 + 3\cosh(x)^4 + 4(5\cosh(x)^3 + 3\cosh(x))\sinh(x)^3 + 3(5\cosh(x)^4 + 6\cosh(x)^2 + 1)\sinh(x)^2 + 3\cosh(x)^2 + 6(\cosh(x)^5 + 2\cosh(x)^3 + \cosh(x))\sinh(x) + 1)$$

Sympy [A]

time = 10.12, size = 78, normalized size = 1.73

$$-\frac{2(\tanh(x) + 1)^{5/2}}{5} - 2\sqrt{\tanh(x) + 1} - 4 \left(\begin{cases} \frac{\sqrt{2} \operatorname{acoth}\left(\frac{\sqrt{2}\sqrt{\tanh(x) + 1}}{2}\right)}{2} & \text{for } \tanh(x) > 1 \\ \frac{\sqrt{2} \operatorname{atanh}\left(\frac{\sqrt{2}\sqrt{\tanh(x) + 1}}{2}\right)}{2} & \text{for } \tanh(x) < 1 \end{cases} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)**2*(1+tanh(x))**(3/2),x)

[Out] $-2*(\tanh(x) + 1)**(5/2)/5 - 2*\sqrt{\tanh(x) + 1} - 4*\text{Piecewise}((-\sqrt{2})*\text{acoth}(\sqrt{2}*\sqrt{\tanh(x) + 1})/2, \tanh(x) > 1), (-\sqrt{2})*\text{atanh}(\sqrt{2}*\sqrt{\tanh(x) + 1})/2, \tanh(x) < 1))$

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 140 vs. 2(34) = 68.

time = 0.43, size = 140, normalized size = 3.11

$$\frac{1}{5}\sqrt{2}\left(\frac{2\left(25\left(\sqrt{e^{4x}+e^{2x}}-e^{2x}\right)^4-60\left(\sqrt{e^{4x}+e^{2x}}-e^{2x}\right)^3+70\left(\sqrt{e^{4x}+e^{2x}}-e^{2x}\right)^2-40\sqrt{e^{4x}+e^{2x}}+40e^{2x}+9\right)}{\left(\sqrt{e^{4x}+e^{2x}}-e^{2x}-1\right)^5}-5\log\left(-2\sqrt{e^{4x}+e^{2x}}+2e^{2x}+1\right)\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)^2*(1+tanh(x))^(3/2),x, algorithm="giac")

[Out] $1/5*\sqrt{2}*(2*(25*(\sqrt{e^{4x}} + e^{2x})) - e^{(2*x)})^4 - 60*(\sqrt{e^{4x}} + e^{2x})^3 + 70*(\sqrt{e^{4x}} + e^{2x})^2 - 40*\sqrt{e^{4x}} + 40*e^{2x} + 9)/(\sqrt{e^{4x}} + e^{2x}) - e^{(2*x)} - 1)^5 - 5*\log(-2*\sqrt{e^{4x}} + e^{2x}) + 2*e^{(2*x)} + 1))$

Mupad [B]

time = 1.11, size = 34, normalized size = 0.76

$$2\sqrt{2}\operatorname{atanh}\left(\frac{\sqrt{2}\sqrt{\tanh(x)+1}}{2}\right) - 2\sqrt{\tanh(x)+1} - \frac{2(\tanh(x)+1)^{5/2}}{5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(x)^2*(tanh(x) + 1)^(3/2),x)

[Out] $2*2^{(1/2)}*\text{atanh}((2^{(1/2)}*(\tanh(x) + 1)^{(1/2)})/2) - 2*(\tanh(x) + 1)^{(1/2)} - (2*(\tanh(x) + 1)^{(5/2)})/5$

3.130 $\int \tanh^2(x) \sqrt{1 + \tanh(x)} dx$

Optimal. Leaf size=34

$$\sqrt{2} \tanh^{-1} \left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}} \right) - \frac{2}{3} (1 + \tanh(x))^{3/2}$$

[Out] arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))*2^(1/2)-2/3*(1+tanh(x))^(3/2)

Rubi [A]

time = 0.04, antiderivative size = 34, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {3624, 3561, 212}

$$\sqrt{2} \tanh^{-1} \left(\frac{\sqrt{\tanh(x) + 1}}{\sqrt{2}} \right) - \frac{2}{3} (\tanh(x) + 1)^{3/2}$$

Antiderivative was successfully verified.

[In] Int[Tanh[x]^2*Sqrt[1 + Tanh[x]],x]

[Out] Sqrt[2]*ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]] - (2*(1 + Tanh[x])^(3/2))/3

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 3561

Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Dist[-2*(b/d), Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]

Rule 3624

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[d^2*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[c^2 - d^2 + 2*c*d*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1] && !(EqQ[m, 2] && EqQ[a, 0])

Rubi steps

$$\begin{aligned}
\int \tanh^2(x) \sqrt{1 + \tanh(x)} \, dx &= -\frac{2}{3}(1 + \tanh(x))^{3/2} + \int \sqrt{1 + \tanh(x)} \, dx \\
&= -\frac{2}{3}(1 + \tanh(x))^{3/2} + 2 \operatorname{Subst} \left(\int \frac{1}{2 - x^2} \, dx, x, \sqrt{1 + \tanh(x)} \right) \\
&= \sqrt{2} \tanh^{-1} \left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}} \right) - \frac{2}{3}(1 + \tanh(x))^{3/2}
\end{aligned}$$

Mathematica [A]

time = 0.05, size = 34, normalized size = 1.00

$$\sqrt{2} \tanh^{-1} \left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}} \right) - \frac{2}{3}(1 + \tanh(x))^{3/2}$$

Antiderivative was successfully verified.

`[In] Integrate[Tanh[x]^2*Sqrt[1 + Tanh[x]], x]``[Out] Sqrt[2]*ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]] - (2*(1 + Tanh[x])^(3/2))/3`**Maple [A]**

time = 0.65, size = 26, normalized size = 0.76

method	result	size
derivativedivides	$\operatorname{arctanh} \left(\frac{\sqrt{1 + \tanh(x)} \sqrt{2}}{2} \right) \sqrt{2} - \frac{2(1 + \tanh(x))^{3/2}}{3}$	26
default	$\operatorname{arctanh} \left(\frac{\sqrt{1 + \tanh(x)} \sqrt{2}}{2} \right) \sqrt{2} - \frac{2(1 + \tanh(x))^{3/2}}{3}$	26

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((1+tanh(x))^(1/2)*tanh(x)^2,x,method=_RETURNVERBOSE)``[Out] arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))*2^(1/2)-2/3*(1+tanh(x))^(3/2)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((1+tanh(x))^(1/2)*tanh(x)^2,x, algorithm="maxima")`

[Out] integrate(sqrt(tanh(x) + 1)*tanh(x)^2, x)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 237 vs. 2(25) = 50.

time = 0.35, size = 237, normalized size = 6.97

$$\frac{8\sqrt{2}\left(\sqrt{2}\cosh(x)^2 + 3\sqrt{2}\cosh(x)\sinh(x) + 3\sqrt{2}\cosh(x)\sinh(x)^2 + \sqrt{2}\sinh(x)^3\right)\frac{\cosh(x)}{\cosh(x) - \sinh(x)} - 3\left(\sqrt{2}\cosh(x)^2 + 4\sqrt{2}\cosh(x)\sinh(x) + \sqrt{2}\sinh(x)^2 + 2\left(3\sqrt{2}\cosh(x)^2 + \sqrt{2}\right)\sinh(x)^2 + 2\sqrt{2}\cosh(x)^2 + 4\left(\sqrt{2}\cosh(x)^2 + \sqrt{2}\cosh(x)\sinh(x) + \sqrt{2}\right)\log\left(-2\sqrt{2}\frac{\cosh(x)}{\cosh(x) - \sinh(x)}(\cosh(x) + \sinh(x)) - 2\cosh(x)^2 - 4\cosh(x)\sinh(x) - 2\sinh(x)^2 - 1\right)\right)}{6(\cosh(x)^2 + 4\cosh(x)\sinh(x) + \sinh(x)^2 + 2(3\cosh(x)^2 + 1)\sinh(x)^2 + 2\cosh(x)^2 + 4(\cosh(x)^2 + \cosh(x)\sinh(x) + 1))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+tanh(x))^(1/2)*tanh(x)^2,x, algorithm="fricas")

[Out] -1/6*(8*sqrt(2)*(sqrt(2)*cosh(x)^3 + 3*sqrt(2)*cosh(x)^2*sinh(x) + 3*sqrt(2)*cosh(x)*sinh(x)^2 + sqrt(2)*sinh(x)^3)*sqrt(cosh(x)/(cosh(x) - sinh(x))) - 3*(sqrt(2)*cosh(x)^4 + 4*sqrt(2)*cosh(x)*sinh(x)^3 + sqrt(2)*sinh(x)^4 + 2*(3*sqrt(2)*cosh(x)^2 + sqrt(2))*sinh(x)^2 + 2*sqrt(2)*cosh(x)^2 + 4*(sqrt(2)*cosh(x)^3 + sqrt(2)*cosh(x))*sinh(x) + sqrt(2))*log(-2*sqrt(2)*sqrt(cosh(x)/(cosh(x) - sinh(x)))*(cosh(x) + sinh(x)) - 2*cosh(x)^2 - 4*cosh(x)*sinh(x) - 2*sinh(x)^2 - 1)/(cosh(x)^4 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4 + 2*(3*cosh(x)^2 + 1)*sinh(x)^2 + 2*cosh(x)^2 + 4*(cosh(x)^3 + cosh(x))*sinh(x) + 1)

Sympy [A]

time = 1.85, size = 68, normalized size = 2.00

$$\frac{2(\tanh(x) + 1)^{\frac{3}{2}}}{3} - 2 \left(\begin{cases} -\frac{\sqrt{2} \operatorname{acoth}\left(\frac{\sqrt{2}\sqrt{\tanh(x) + 1}}{2}\right)}{2} & \text{for } \tanh(x) > 1 \\ -\frac{\sqrt{2} \operatorname{atanh}\left(\frac{\sqrt{2}\sqrt{\tanh(x) + 1}}{2}\right)}{2} & \text{for } \tanh(x) < 1 \end{cases} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+tanh(x))**(1/2)*tanh(x)**2,x)

[Out] -2*(tanh(x) + 1)**(3/2)/3 - 2*Piecewise((-sqrt(2)*acoth(sqrt(2)*sqrt(tanh(x) + 1)/2)/2, tanh(x) > 1), (-sqrt(2)*atanh(sqrt(2)*sqrt(tanh(x) + 1)/2)/2, tanh(x) < 1))

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 96 vs. 2(25) = 50.

time = 0.43, size = 96, normalized size = 2.82

$$\frac{1}{6}\sqrt{2}\left(\frac{8\left(3\left(\sqrt{e^{4x} + e^{2x}} - e^{2x}\right)^2 - 3\sqrt{e^{4x} + e^{2x}} + 3e^{2x} + 1\right)}{\left(\sqrt{e^{4x} + e^{2x}} - e^{2x} - 1\right)^3} - 3\log\left(-2\sqrt{e^{4x} + e^{2x}} + 2e^{2x} + 1\right)\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+tanh(x))^(1/2)*tanh(x)^2,x, algorithm="giac")

[Out] $\frac{1}{6}\sqrt{2}\left(8\left(3\left(\sqrt{e^{4x} + e^{2x}}\right) - e^{2x}\right)^2 - 3\sqrt{e^{4x} + e^{2x}} + e^{2x}\right) + 3e^{2x} + 1\right) / \left(\sqrt{e^{4x} + e^{2x}} - e^{2x} - 1\right)^3 - 3\log\left(-2\sqrt{e^{4x} + e^{2x}} + 2e^{2x} + 1\right)$

Mupad [B]

time = 0.11, size = 25, normalized size = 0.74

$$\sqrt{2} \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{\tanh(x) + 1}}{2}\right) - \frac{2(\tanh(x) + 1)^{3/2}}{3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(x)^2*(tanh(x) + 1)^(1/2),x)

[Out] $2^{1/2} \operatorname{atanh}\left(\frac{2^{1/2}(\tanh(x) + 1)^{1/2}}{2}\right) - \frac{2(\tanh(x) + 1)^{3/2}}{3}$

$$3.131 \quad \int \frac{\tanh^2(x)}{\sqrt{1 + \tanh(x)}} dx$$

Optimal. Leaf size=42

$$\frac{\tanh^{-1}\left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{1}{\sqrt{1 + \tanh(x)}} - 2\sqrt{1 + \tanh(x)}$$

[Out] 1/2*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))*2^(1/2)-1/(1+tanh(x))^(1/2)-2*(1+tanh(x))^(1/2)

Rubi [A]

time = 0.04, antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {3624, 3560, 3561, 212}

$$\frac{\tanh^{-1}\left(\frac{\sqrt{\tanh(x) + 1}}{\sqrt{2}}\right)}{\sqrt{2}} - 2\sqrt{\tanh(x) + 1} - \frac{1}{\sqrt{\tanh(x) + 1}}$$

Antiderivative was successfully verified.

[In] Int[Tanh[x]^2/Sqrt[1 + Tanh[x]], x]

[Out] ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]]/Sqrt[2] - 1/Sqrt[1 + Tanh[x]] - 2*Sqrt[1 + Tanh[x]]

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 3560

Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n), x] + Dist[1/(2*a), Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]

Rule 3561

Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Dist[-2*(b/d), Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]

Rule 3624

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^2, x_Symbol] :> Simp[d^2*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(
m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[c^2 - d^2 + 2*c*d*Tan[e + f*
x], x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && !LeQ
[m, -1] && !(EqQ[m, 2] && EqQ[a, 0])
```

Rubi steps

$$\begin{aligned}
\int \frac{\tanh^2(x)}{\sqrt{1 + \tanh(x)}} dx &= -2\sqrt{1 + \tanh(x)} + \int \frac{1}{\sqrt{1 + \tanh(x)}} dx \\
&= -\frac{1}{\sqrt{1 + \tanh(x)}} - 2\sqrt{1 + \tanh(x)} + \frac{1}{2} \int \sqrt{1 + \tanh(x)} dx \\
&= -\frac{1}{\sqrt{1 + \tanh(x)}} - 2\sqrt{1 + \tanh(x)} + \text{Subst}\left(\int \frac{1}{2 - x^2} dx, x, \sqrt{1 + \tanh(x)}\right) \\
&= \frac{\tanh^{-1}\left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{1}{\sqrt{1 + \tanh(x)}} - 2\sqrt{1 + \tanh(x)}
\end{aligned}$$

Mathematica [A]

time = 0.07, size = 37, normalized size = 0.88

$$\frac{\tanh^{-1}\left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}}\right)}{\sqrt{2}} + \frac{-3 - 2 \tanh(x)}{\sqrt{1 + \tanh(x)}}$$

Antiderivative was successfully verified.

[In] Integrate[Tanh[x]^2/Sqrt[1 + Tanh[x]], x]

[Out] ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]]/Sqrt[2] + (-3 - 2*Tanh[x])/Sqrt[1 + Tanh[x]]

Maple [A]

time = 0.74, size = 35, normalized size = 0.83

method	result	size
derivativedivides	$ \frac{\arctanh\left(\frac{\sqrt{1 + \tanh(x)} \sqrt{2}}{2}\right) \sqrt{2}}{2} - \frac{1}{\sqrt{1 + \tanh(x)}} - 2\sqrt{1 + \tanh(x)} $	35

default	$\frac{\operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}\sqrt{2}}{2}\right)\sqrt{2}}{2} - \frac{1}{\sqrt{1+\tanh(x)}} - 2\sqrt{1+\tanh(x)}$	35
---------	---	----

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tanh(x)^2/(1+tanh(x))^(1/2),x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{2}\operatorname{arctanh}\left(\frac{1}{2}(1+\tanh(x))^{1/2}\right)2^{1/2} - \frac{1}{(1+\tanh(x))^{1/2}} - 2(1+\tanh(x))^{1/2}$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(x)^2/(1+tanh(x))^(1/2),x, algorithm="maxima")`

[Out] `integrate(tanh(x)^2/sqrt(tanh(x) + 1), x)`

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 182 vs. 2(34) = 68.

time = 0.33, size = 182, normalized size = 4.33

$$\frac{2\sqrt{2}\left(5\sqrt{2}\cosh(x)^2+10\sqrt{2}\cosh(x)\sinh(x)+5\sqrt{2}\sinh(x)^2+\sqrt{2}\right)\sqrt{\frac{\cosh(x)}{\cosh(x)-\sinh(x)}} - \left(\sqrt{2}\cosh(x)^2+3\sqrt{2}\cosh(x)\sinh(x)+\sqrt{2}\sinh(x)^2+(3\sqrt{2}\cosh(x)+\sqrt{2})\sinh(x)+\sqrt{2}\cosh(x)\right)\log\left(-2\sqrt{2}\sqrt{\frac{\cosh(x)}{\cosh(x)-\sinh(x)}}(\cosh(x)+\sinh(x))-2\cosh(x)^2-4\cosh(x)\sinh(x)-2\sinh(x)^2-1\right)}{4(\cosh(x)^2+3\cosh(x)\sinh(x)+\sinh(x)^2+(3\cosh(x)+1)\sinh(x)+\cosh(x))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(x)^2/(1+tanh(x))^(1/2),x, algorithm="fricas")`

[Out] $-\frac{1}{4}(2\sqrt{2})(5\sqrt{2}\cosh(x)^2+10\sqrt{2}\cosh(x)\sinh(x)+5\sqrt{2}\sinh(x)^2+\sqrt{2})\sqrt{\cosh(x)/(\cosh(x)-\sinh(x))} - (\sqrt{2}\cosh(x)^3+3\sqrt{2}\cosh(x)\sinh(x)^2+\sqrt{2}\sinh(x)^3+(3\sqrt{2}\cosh(x)^2+\sqrt{2})\sinh(x)+\sqrt{2}\cosh(x))\log(-2\sqrt{2}\sqrt{\cosh(x)/(\cosh(x)-\sinh(x))}(\cosh(x)+\sinh(x))-2\cosh(x)^2-4\cosh(x)\sinh(x)-2\sinh(x)^2-1)/(\cosh(x)^3+3\cosh(x)\sinh(x)^2+\sinh(x)^3+(3\cosh(x)^2+1)\sinh(x)+\cosh(x))$

Sympy [A]

time = 2.14, size = 75, normalized size = 1.79

$$-2\sqrt{\tanh(x)+1} - \begin{cases} \frac{\sqrt{2}\operatorname{acoth}\left(\frac{\sqrt{2}\sqrt{\tanh(x)+1}}{2}\right)}{2} & \text{for } \tanh(x) > 1 \\ \frac{\sqrt{2}\operatorname{atanh}\left(\frac{\sqrt{2}\sqrt{\tanh(x)+1}}{2}\right)}{2} & \text{for } \tanh(x) < 1 \end{cases} - \frac{1}{\sqrt{\tanh(x)+1}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)**2/(1+tanh(x))**(1/2),x)

[Out] $-2\sqrt{\tanh(x) + 1} - \text{Piecewise}((- \sqrt{2})\text{acoth}(\sqrt{2})\sqrt{\tanh(x) + 1}/2)/2, \tanh(x) > 1), (- \sqrt{2})\text{atanh}(\sqrt{2})\sqrt{\tanh(x) + 1}/2)/2, \tanh(x) < 1)) - 1/\sqrt{\tanh(x) + 1}$

Giac [A]

time = 0.42, size = 54, normalized size = 1.29

$$-\frac{1}{4}\sqrt{2}\log\left(-4\sqrt{e^{4x}+e^{2x}}+4e^{2x}+2\right)-\frac{5\sqrt{2}e^{2x}+\sqrt{2}}{2\sqrt{e^{4x}+e^{2x}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)^2/(1+tanh(x))^(1/2),x, algorithm="giac")

[Out] $-1/4\sqrt{2}\log(-4\sqrt{e^{4x}+e^{2x}}+4e^{2x}+2)-1/2*(5\sqrt{2}(2)e^{2x}+\sqrt{2})/\sqrt{e^{4x}+e^{2x}}$

Mupad [B]

time = 0.13, size = 36, normalized size = 0.86

$$\frac{\sqrt{2}\operatorname{atanh}\left(\frac{\sqrt{2}\sqrt{\tanh(x)+1}}{2}\right)}{2}-\frac{3}{\sqrt{\tanh(x)+1}}-\frac{2\tanh(x)}{\sqrt{\tanh(x)+1}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(x)^2/(tanh(x)+1)^(1/2),x)

[Out] $(2^{1/2})\operatorname{atanh}((2^{1/2})(\tanh(x)+1)^{1/2})/2)-3/(\tanh(x)+1)^{1/2}-(2\tanh(x))/(\tanh(x)+1)^{1/2}$

$$3.132 \quad \int \frac{\tanh^2(x)}{(1+\tanh(x))^{3/2}} dx$$

Optimal. Leaf size=49

$$\frac{\tanh^{-1}\left(\frac{\sqrt{1+\tanh(x)}}{\sqrt{2}}\right)}{2\sqrt{2}} - \frac{1}{3(1+\tanh(x))^{3/2}} + \frac{3}{2\sqrt{1+\tanh(x)}}$$

[Out] 1/4*arctanh(1/2*(1+tanh(x))^(1/2)*2^(1/2))*2^(1/2)+3/2/(1+tanh(x))^(1/2)-1/3/(1+tanh(x))^(3/2)

Rubi [A]

time = 0.06, antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {3621, 3607, 3561, 212}

$$\frac{\tanh^{-1}\left(\frac{\sqrt{\tanh(x)+1}}{\sqrt{2}}\right)}{2\sqrt{2}} + \frac{3}{2\sqrt{\tanh(x)+1}} - \frac{1}{3(\tanh(x)+1)^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[Tanh[x]^2/(1 + Tanh[x])^(3/2), x]

[Out] ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]]/(2*Sqrt[2]) - 1/(3*(1 + Tanh[x])^(3/2)) + 3/(2*Sqrt[1 + Tanh[x]])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 3561

Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] :> Dist[-2*(b/d), Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]

Rule 3607

Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] :> Simp[(-b*c - a*d)*((a + b*Tan[e + f*x])^m/(2*a*f*m)), x] + Dist[(b*c + a*d)/(2*a*b), Int[(a + b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0]

Rule 3621

Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^2, x_Symbol] := Simp[(-b)*(a*c + b*d)^2*((a + b*Tan[e + f*x])^m/(2*a^3*f*m)), x] + Dist[1/(2*a^2), Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c^2 - 2*b*c*d + a*d^2 - 2*b*d^2*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && LeQ[m, -1] && EqQ[a^2 + b^2, 0]

Rubi steps

$$\begin{aligned} \int \frac{\tanh^2(x)}{(1 + \tanh(x))^{3/2}} dx &= -\frac{1}{3(1 + \tanh(x))^{3/2}} - \frac{1}{2} \int \frac{1 - 2 \tanh(x)}{\sqrt{1 + \tanh(x)}} dx \\ &= -\frac{1}{3(1 + \tanh(x))^{3/2}} + \frac{3}{2\sqrt{1 + \tanh(x)}} + \frac{1}{4} \int \sqrt{1 + \tanh(x)} dx \\ &= -\frac{1}{3(1 + \tanh(x))^{3/2}} + \frac{3}{2\sqrt{1 + \tanh(x)}} + \frac{1}{2} \text{Subst}\left(\int \frac{1}{2 - x^2} dx, x, \sqrt{1 + \tanh(x)}\right) \\ &= \frac{\tanh^{-1}\left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}}\right)}{2\sqrt{2}} - \frac{1}{3(1 + \tanh(x))^{3/2}} + \frac{3}{2\sqrt{1 + \tanh(x)}} \end{aligned}$$

Mathematica [A]

time = 0.07, size = 53, normalized size = 1.08

$$\frac{\tanh^{-1}\left(\frac{\sqrt{1 + \tanh(x)}}{\sqrt{2}}\right)}{2\sqrt{2}} + \frac{(\cosh(x) - \sinh(x))(7 \cosh(x) + 9 \sinh(x))}{6\sqrt{1 + \tanh(x)}}$$

Antiderivative was successfully verified.

[In] Integrate[Tanh[x]^2/(1 + Tanh[x])^(3/2), x]

[Out] ArcTanh[Sqrt[1 + Tanh[x]]/Sqrt[2]]/(2*Sqrt[2]) + ((Cosh[x] - Sinh[x])*(7*Cosh[x] + 9*Sinh[x]))/(6*Sqrt[1 + Tanh[x]])

Maple [A]

time = 0.68, size = 35, normalized size = 0.71

method	result	size
derivativedivides	$\frac{\arctanh\left(\frac{\sqrt{1 + \tanh(x)} \sqrt{2}}{2}\right) \sqrt{2}}{4} + \frac{3}{2\sqrt{1 + \tanh(x)}} - \frac{1}{3(1 + \tanh(x))^{\frac{3}{2}}}$	35

default	$\frac{\operatorname{arctanh}\left(\frac{\sqrt{1+\tanh(x)}\sqrt{2}}{2}\right)\sqrt{2}}{4} + \frac{3}{2\sqrt{1+\tanh(x)}} - \frac{1}{3(1+\tanh(x))^{\frac{3}{2}}}$	35
---------	---	----

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tanh(x)^2/(1+tanh(x))^(3/2),x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{4}\operatorname{arctanh}\left(\frac{1}{2}(1+\tanh(x))^{1/2}\right)2^{1/2} + \frac{3}{2(1+\tanh(x))^{1/2}} - \frac{1}{3(1+\tanh(x))^{3/2}}$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(x)^2/(1+tanh(x))^(3/2),x, algorithm="maxima")`

[Out] `integrate(tanh(x)^2/(tanh(x) + 1)^(3/2), x)`

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 168 vs. $2(34) = 68$.

time = 0.36, size = 168, normalized size = 3.43

$$\frac{2\sqrt{2}\left(8\sqrt{2}\cosh(x)^2+16\sqrt{2}\cosh(x)\sinh(x)+8\sqrt{2}\sinh(x)^2-\sqrt{2}\right)\sqrt{\frac{\cosh(x)}{\cosh(x)-\sinh(x)}}+3\left(\sqrt{2}\cosh(x)^3+3\sqrt{2}\cosh(x)^2\sinh(x)+3\sqrt{2}\cosh(x)\sinh(x)^2+\sqrt{2}\sinh(x)^3\right)\log\left(-2\sqrt{2}\sqrt{\frac{\cosh(x)}{\cosh(x)-\sinh(x)}}(\cosh(x)+\sinh(x))-2\cosh(x)^2-4\cosh(x)\sinh(x)-2\sinh(x)^2-1\right)}{24(\cosh(x)^3+3\cosh(x)^2\sinh(x)+3\cosh(x)\sinh(x)^2+\sinh(x)^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(x)^2/(1+tanh(x))^(3/2),x, algorithm="fricas")`

[Out] $\frac{1}{24}(2\sqrt{2}(8\sqrt{2}\cosh(x)^2+16\sqrt{2}\cosh(x)\sinh(x)+8\sqrt{2}\sinh(x)^2-\sqrt{2})\sqrt{\frac{\cosh(x)}{\cosh(x)-\sinh(x)}}+3(\sqrt{2}\cosh(x)^3+3\sqrt{2}\cosh(x)^2\sinh(x)+3\sqrt{2}\cosh(x)\sinh(x)^2+\sqrt{2}\sinh(x)^3)\log(-2\sqrt{2}\sqrt{\frac{\cosh(x)}{\cosh(x)-\sinh(x)}}(\cosh(x)+\sinh(x))-2\cosh(x)^2-4\cosh(x)\sinh(x)-2\sinh(x)^2-1))/(\cosh(x)^3+3\cosh(x)^2\sinh(x)+3\cosh(x)\sinh(x)^2+\sinh(x)^3)$

Sympy [A]

time = 7.67, size = 78, normalized size = 1.59

$$\frac{\begin{cases} \frac{\sqrt{2}\operatorname{acoth}\left(\frac{\sqrt{2}\sqrt{\tanh(x)+1}}{2}\right)}{2} & \text{for } \tanh(x) > 1 \\ \frac{\sqrt{2}\operatorname{atanh}\left(\frac{\sqrt{2}\sqrt{\tanh(x)+1}}{2}\right)}{2} & \text{for } \tanh(x) < 1 \end{cases}}{2} + \frac{3}{2\sqrt{\tanh(x)+1}} - \frac{1}{3(\tanh(x)+1)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)**2/(1+tanh(x))**(3/2),x)

[Out] -Piecewise((-sqrt(2)*acoth(sqrt(2)*sqrt(tanh(x) + 1)/2)/2, tanh(x) > 1), (-sqrt(2)*atanh(sqrt(2)*sqrt(tanh(x) + 1)/2)/2, tanh(x) < 1))/2 + 3/(2*sqrt(tanh(x) + 1)) - 1/(3*(tanh(x) + 1)**(3/2))

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 95 vs. 2(34) = 68.
time = 0.42, size = 95, normalized size = 1.94

$$\frac{1}{24} \sqrt{2} \left(\frac{2 \left(6 \left(\sqrt{e^{4x} + e^{2x}} - e^{2x} \right)^2 + 3 \sqrt{e^{4x} + e^{2x}} - 3 e^{2x} - 1 \right)}{\left(\sqrt{e^{4x} + e^{2x}} - e^{2x} \right)^3} - 3 \log \left(-2 \sqrt{e^{4x} + e^{2x}} + 2 e^{2x} + 1 \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)^2/(1+tanh(x))^(3/2),x, algorithm="giac")

[Out] 1/24*sqrt(2)*(2*(6*(sqrt(e^(4*x) + e^(2*x)) - e^(2*x))^2 + 3*sqrt(e^(4*x) + e^(2*x)) - 3*e^(2*x) - 1)/(sqrt(e^(4*x) + e^(2*x)) - e^(2*x))^3 - 3*log(-2*sqrt(e^(4*x) + e^(2*x)) + 2*e^(2*x) + 1))

Mupad [B]

time = 1.08, size = 31, normalized size = 0.63

$$\frac{\sqrt{2} \operatorname{atanh} \left(\frac{\sqrt{2} \sqrt{\tanh(x) + 1}}{2} \right)}{4} + \frac{\frac{3 \tanh(x)}{2} + \frac{7}{6}}{(\tanh(x) + 1)^{3/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(x)^2/(tanh(x) + 1)^(3/2),x)

[Out] (2^(1/2)*atanh((2^(1/2)*(tanh(x) + 1)^(1/2))/2))/4 + ((3*tanh(x))/2 + 7/6)/(tanh(x) + 1)^(3/2)

3.133 $\int \frac{\tanh^5(x)}{a+b \tanh(x)} dx$

Optimal. Leaf size=94

$$-\frac{bx}{a^2-b^2} + \frac{a \log(\cosh(x))}{a^2-b^2} + \frac{a^5 \log(a+b \tanh(x))}{b^4(a^2-b^2)} - \frac{(a^2+b^2) \tanh(x)}{b^3} + \frac{a \tanh^2(x)}{2b^2} - \frac{\tanh^3(x)}{3b}$$

[Out] $-b*x/(a^2-b^2)+a*\ln(\cosh(x))/(a^2-b^2)+a^5*\ln(a+b*\tanh(x))/b^4/(a^2-b^2)-(a^2+b^2)*\tanh(x)/b^3+1/2*a*\tanh(x)^2/b^2-1/3*\tanh(x)^3/b$

Rubi [A]

time = 0.26, antiderivative size = 94, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.538$,

Rules used = {3647, 3728, 3729, 3707, 3698, 31, 3556}

$$-\frac{bx}{a^2-b^2} + \frac{a \log(\cosh(x))}{a^2-b^2} - \frac{(a^2+b^2) \tanh(x)}{b^3} + \frac{a^5 \log(a+b \tanh(x))}{b^4(a^2-b^2)} + \frac{a \tanh^2(x)}{2b^2} - \frac{\tanh^3(x)}{3b}$$

Antiderivative was successfully verified.

[In] `Int[Tanh[x]^5/(a + b*Tanh[x]),x]`

[Out] $-((b*x)/(a^2 - b^2)) + (a*\text{Log}[\text{Cosh}[x]])/(a^2 - b^2) + (a^5*\text{Log}[a + b*\text{Tanh}[x]])/(b^4*(a^2 - b^2)) - ((a^2 + b^2)*\text{Tanh}[x])/b^3 + (a*\text{Tanh}[x]^2)/(2*b^2) - \text{Tanh}[x]^3/(3*b)$

Rule 31

`Int[((a_) + (b_.)*(x_))^(n_), x_Symbol] := Simp[Log[RemoveContent[a + b*x, x]]/b, x] /; FreeQ[{a, b}, x]`

Rule 3556

`Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

Rule 3647

`Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b^2*(a + b*Tan[e + f*x])^(m-2)*((c + d*Tan[e + f*x])^(n+1)/(d*f*(m+n-1))), x] + Dist[1/(d*(m+n-1)), Int[(a + b*Tan[e + f*x])^(m-3)*(c + d*Tan[e + f*x])^n*Simp[a^3*d*(m+n-1) - b^2*(b*c*(m-2) + a*d*(1+n)) + b*d*(m+n-1)*(3*a^2 - b^2)*Tan[e + f*x] - b^2*(b*c*(m-2) - a*d*(3*m+2*n-4))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && IntegerQ[2*m] && GtQ[m, 2] && (GeQ[n, -1] || IntegerQ[m]) && !(IGtQ[n, 2] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

Rule 3698

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_) + (C_.)*tan[(e_.) +
(f_.)*(x_)])^2, x_Symbol] := Dist[A/(b*f), Subst[Int[(a + x)^m, x], x, b*T
an[e + f*x]], x] /; FreeQ[{a, b, e, f, A, C, m}, x] && EqQ[A, C]
```

Rule 3707

```
Int[((A_) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2
)/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*A + b*B - a
*C)*(x/(a^2 + b^2)), x] + (Dist[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2), Int[(1
+ Tan[e + f*x]^2)/(a + b*Tan[e + f*x]), x], x] - Dist[(A*b - a*B - b*C)/(a
^2 + b^2), Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, e, f, A, B, C}, x] &&
NeQ[A*b^2 - a*b*B + a^2*C, 0] && NeQ[a^2 + b^2, 0] && NeQ[A*b - a*B - b*C,
0]
```

Rule 3728

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.
) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Dist[1/(d*(m + n + 1)), Int[(a +
b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C*
(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*m
*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b
, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[c
, 0] && NeQ[a, 0])))
```

Rule 3729

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] :=
Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n +
1))), x] + Dist[1/(d*(m + n + 1)), Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*
Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C*(b*c*m + a*d*(n + 1)) + d*(A*b -
b*C)*(m + n + 1)*Tan[e + f*x] - C*m*(b*c - a*d)*Tan[e + f*x]^2, x], x], x]
/; FreeQ[{a, b, c, d, e, f, A, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 +
b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[
m] || (EqQ[c, 0] && NeQ[a, 0])))
```

Rubi steps

$$\begin{aligned}
\int \frac{\tanh^5(x)}{a+b \tanh(x)} dx &= -\frac{\tanh^3(x)}{3b} - \frac{\int \frac{\tanh^2(x)(-3a-3b \tanh(x)+3a \tanh^2(x))}{a+b \tanh(x)} dx}{3b} \\
&= \frac{a \tanh^2(x)}{2b^2} - \frac{\tanh^3(x)}{3b} - \frac{\int \frac{\tanh(x)(6a^2-6(a^2+b^2) \tanh^2(x))}{a+b \tanh(x)} dx}{6b^2} \\
&= -\frac{(a^2+b^2) \tanh(x)}{b^3} + \frac{a \tanh^2(x)}{2b^2} - \frac{\tanh^3(x)}{3b} - \frac{\int \frac{-6a(a^2+b^2)-6b^3 \tanh(x)+6a(a^2+b^2) \tanh^2(x)}{a+b \tanh(x)} dx}{6b^3} \\
&= -\frac{bx}{a^2-b^2} - \frac{(a^2+b^2) \tanh(x)}{b^3} + \frac{a \tanh^2(x)}{2b^2} - \frac{\tanh^3(x)}{3b} + \frac{a \int \tanh(x) dx}{a^2-b^2} + \frac{a^5 \int \frac{1-\tanh(x)}{a+b \tanh(x)} dx}{b^3(a^2-b^2)} \\
&= -\frac{bx}{a^2-b^2} + \frac{a \log(\cosh(x))}{a^2-b^2} - \frac{(a^2+b^2) \tanh(x)}{b^3} + \frac{a \tanh^2(x)}{2b^2} - \frac{\tanh^3(x)}{3b} + \frac{a^5 \text{Subst}\left(\frac{1-t}{a+bt}, t, \tanh(x)\right)}{b^3(a^2-b^2)} \\
&= -\frac{bx}{a^2-b^2} + \frac{a \log(\cosh(x))}{a^2-b^2} + \frac{a^5 \log(a+b \tanh(x))}{b^4(a^2-b^2)} - \frac{(a^2+b^2) \tanh(x)}{b^3} + \frac{a \tanh^2(x)}{2b^2}
\end{aligned}$$

Mathematica [A]

time = 0.41, size = 105, normalized size = 1.12

$$-\frac{6b^5x + 6a(a^4 - b^4) \log(\cosh(x)) - 6a^5 \log(a \cosh(x) + b \sinh(x)) + 2b(3a^4 + a^2b^2 - 4b^4) \tanh(x) + b^2(-a^2 + b^2) \operatorname{sech}^2(x)(-3a + 2b \tanh(x))}{6(a-b)b^4(a+b)}$$

Antiderivative was successfully verified.

`[In] Integrate[Tanh[x]^5/(a + b*Tanh[x]), x]`

```
[Out] -1/6*(6*b^5*x + 6*a*(a^4 - b^4)*Log[Cosh[x]] - 6*a^5*Log[a*Cosh[x] + b*Sinh[x]] + 2*b*(3*a^4 + a^2*b^2 - 4*b^4)*Tanh[x] + b^2*(-a^2 + b^2)*Sech[x]^2*(-3*a + 2*b*Tanh[x]))/((a - b)*b^4*(a + b))
```

Maple [A]

time = 0.32, size = 95, normalized size = 1.01

method	result
derivativedivides	$-\frac{b^2(\tanh^3(x))}{3} - \frac{a(\tanh^2(x))b}{2b^3} + a^2 \tanh(x) + b^2 \tanh(x) + \frac{a^5 \ln(a+b \tanh(x))}{b^4(a+b)(a-b)} - \frac{\ln(\tanh(x)-1)}{2b+2a} - \frac{\ln(1+\tanh(x))}{2a-2b}$
default	$-\frac{b^2(\tanh^3(x))}{3} - \frac{a(\tanh^2(x))b}{2b^3} + a^2 \tanh(x) + b^2 \tanh(x) + \frac{a^5 \ln(a+b \tanh(x))}{b^4(a+b)(a-b)} - \frac{\ln(\tanh(x)-1)}{2b+2a} - \frac{\ln(1+\tanh(x))}{2a-2b}$
risch	$\frac{x}{a+b} + \frac{2a^3x}{b^4} + \frac{2ax}{b^2} - \frac{2xa^5}{b^4(a^2-b^2)} + \frac{2a^2e^{4x} - 2abe^{4x} + 4b^2e^{4x} + 4a^2e^{2x} - 2abe^{2x} + 4b^2e^{2x} + 2a^2 + \frac{8b^2}{3}}{b^3(1+e^{2x})^3} - \frac{a^3 \ln(1+e^{2x})}{b^4}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(tanh(x)^5/(a+b*tanh(x)), x, method=_RETURNVERBOSE)`

[Out] $-1/b^3*(1/3*b^2*\tanh(x)^3-1/2*a*\tanh(x)^2*b+a^2*\tanh(x)+b^2*\tanh(x))+1/b^4*a^5/(a+b)/(a-b)*\ln(a+b*\tanh(x))-1/(2*b+2*a)*\ln(\tanh(x)-1)-1/(2*a-2*b)*\ln(1+\tanh(x))$

Maxima [A]

time = 0.49, size = 150, normalized size = 1.60

$$\frac{a^5 \log(-(a-b)e^{-2x} - a - b)}{a^2 b^4 - b^6} - \frac{2(3a^2 + 4b^2 + 3(2a^2 + ab + 2b^2)e^{-2x} + 3(a^2 + ab + 2b^2)e^{-4x})}{3(3b^3e^{-2x} + 3b^3e^{-4x} + b^3e^{-6x} + b^3)} + \frac{x}{a+b} - \frac{(a^3 + ab^2) \log(e^{-2x} + 1)}{b^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(x)^5/(a+b*tanh(x)),x, algorithm="maxima")`

[Out] $a^5*\log(-(a - b)*e^{-2*x} - a - b)/(a^2*b^4 - b^6) - 2/3*(3*a^2 + 4*b^2 + 3*(2*a^2 + a*b + 2*b^2)*e^{-2*x} + 3*(a^2 + a*b + 2*b^2)*e^{-4*x}))/((3*b^3*e^{-2*x} + 3*b^3*e^{-4*x} + b^3*e^{-6*x} + b^3) + x/(a + b) - (a^3 + a*b^2)*\log(e^{-2*x} + 1)/b^4$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 1296 vs. 2(90) = 180.

time = 0.41, size = 1296, normalized size = 13.79

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(x)^5/(a+b*tanh(x)),x, algorithm="fricas")`

[Out] $-1/3*(3*(a*b^4 + b^5)*x*\cosh(x)^6 + 18*(a*b^4 + b^5)*x*\cosh(x)*\sinh(x)^5 + 3*(a*b^4 + b^5)*x*\sinh(x)^6 - 6*a^4*b - 2*a^2*b^3 + 8*b^5 - 3*(2*a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + 2*a*b^4 - 4*b^5 - 3*(a*b^4 + b^5)*x)*\cosh(x)^4 - 3*(2*a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + 2*a*b^4 - 4*b^5 - 15*(a*b^4 + b^5)*x*\cosh(x)^2 - 3*(a*b^4 + b^5)*x)*\sinh(x)^4 + 12*(5*(a*b^4 + b^5)*x*\cosh(x)^3 - (2*a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + 2*a*b^4 - 4*b^5 - 3*(a*b^4 + b^5)*x)*\cosh(x))*\sinh(x)^3 - 3*(4*a^4*b - 2*a^3*b^2 + 2*a*b^4 - 4*b^5 - 3*(a*b^4 + b^5)*x)*\cosh(x)^2 + 3*(15*(a*b^4 + b^5)*x*\cosh(x)^4 - 4*a^4*b + 2*a^3*b^2 - 2*a*b^4 + 4*b^5 - 6*(2*a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + 2*a*b^4 - 4*b^5 - 3*(a*b^4 + b^5)*x)*\cosh(x)^2 + 3*(a*b^4 + b^5)*x)*\sinh(x)^2 + 3*(a*b^4 + b^5)*x - 3*(a^5*\cosh(x)^6 + 6*a^5*\cosh(x)*\sinh(x)^5 + a^5*\sinh(x)^6 + 3*a^5*\cosh(x)^4 + 3*a^5*\cosh(x)^2 + a^5 + 3*(5*a^5*\cosh(x)^2 + a^5)*\sinh(x)^4 + 4*(5*a^5*\cosh(x)^3 + 3*a^5*\cosh(x))*\sinh(x)^3 + 3*(5*a^5*\cosh(x)^4 + 6*a^5*\cosh(x)^2 + a^5)*\sinh(x)^2 + 6*(a^5*\cosh(x)^5 + 2*a^5*\cosh(x)^3 + a^5*\cosh(x))*\sinh(x))*\log(2*(a*\cosh(x) + b*\sinh(x))/(\cosh(x) - \sinh(x))) + 3*((a^5 - a*b^4)*\cosh(x)^6 + 6*(a^5 - a*b^4)*\cosh(x)*\sinh(x)^5 + (a^5 - a*b^4)*\sinh(x)^6 + a^5 - a*b^4 + 3*(a^5 - a*b^4)*\cosh(x)^4 + 3*(a^5 - a*b^4 + 5*(a^5 - a*b^4)*\cosh(x)^2)*\sinh(x)^4 + 4*(5*(a^5 - a*b^4)*\cosh(x)^3 + 3*(a^5 - a*b^4)*\cosh(x))*\sinh(x)^3 + 3*(a^5 - a*b^4)*\cosh(x)^2 + 3*(a^5 - a*b^4 + 5*(a^5 - a*b^4)*$

$$\begin{aligned} & \cosh(x)^4 + 6*(a^5 - a*b^4)*\cosh(x)^2*\sinh(x)^2 + 6*((a^5 - a*b^4)*\cosh(x) \\ & ^5 + 2*(a^5 - a*b^4)*\cosh(x)^3 + (a^5 - a*b^4)*\cosh(x))*\sinh(x)*\log(2*\cosh \\ & (x)/(\cosh(x) - \sinh(x))) + 6*(3*(a*b^4 + b^5)*x*\cosh(x)^5 - 2*(2*a^4*b - 2* \\ & a^3*b^2 + 2*a^2*b^3 + 2*a*b^4 - 4*b^5 - 3*(a*b^4 + b^5)*x)*\cosh(x)^3 - (4*a \\ & ^4*b - 2*a^3*b^2 + 2*a*b^4 - 4*b^5 - 3*(a*b^4 + b^5)*x)*\cosh(x))*\sinh(x))/ \\ & ((a^2*b^4 - b^6)*\cosh(x)^6 + 6*(a^2*b^4 - b^6)*\cosh(x)*\sinh(x)^5 + (a^2*b^4 \\ & - b^6)*\sinh(x)^6 + a^2*b^4 - b^6 + 3*(a^2*b^4 - b^6)*\cosh(x)^4 + 3*(a^2*b^4 \\ & - b^6 + 5*(a^2*b^4 - b^6)*\cosh(x)^2)*\sinh(x)^4 + 4*(5*(a^2*b^4 - b^6)*\cosh \\ & (x)^3 + 3*(a^2*b^4 - b^6)*\cosh(x))*\sinh(x)^3 + 3*(a^2*b^4 - b^6)*\cosh(x)^2 \\ & + 3*(a^2*b^4 - b^6 + 5*(a^2*b^4 - b^6)*\cosh(x)^4 + 6*(a^2*b^4 - b^6)*\cosh(x) \\ &)^2)*\sinh(x)^2 + 6*((a^2*b^4 - b^6)*\cosh(x)^5 + 2*(a^2*b^4 - b^6)*\cosh(x)^3 \\ & + (a^2*b^4 - b^6)*\cosh(x))*\sinh(x)) \end{aligned}$$

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 546 vs. 2(78) = 156.

time = 0.57, size = 546, normalized size = 5.81

$$\begin{cases} \infty \left(x - \frac{\tanh^3(x)}{3} - \tanh(x) \right) & \text{for } a = 0 \wedge b = 0 \\ \frac{x - \log(\tanh(x)+1) - \frac{\tanh^4(x)}{4} - \frac{\tanh^2(x)}{2}}{a} & \text{for } b = 0 \\ \frac{27x \tanh(x)}{6b \tanh(x)-6b} - \frac{27x}{6b \tanh(x)-6b} - \frac{12 \log(\tanh(x)+1) \tanh(x)}{6b \tanh(x)-6b} + \frac{12 \log(\tanh(x)+1)}{6b \tanh(x)-6b} - \frac{2 \tanh^4(x)}{6b \tanh(x)-6b} - \frac{\tanh^3(x)}{6b \tanh(x)-6b} - \frac{9 \tanh^2(x)}{6b \tanh(x)-6b} + \frac{15}{6b \tanh(x)-6b} & \text{for } a = -b \\ \frac{3x \tanh(x)}{6b \tanh(x)+6b} + \frac{3x}{6b \tanh(x)+6b} + \frac{12 \log(\tanh(x)+1) \tanh(x)}{6b \tanh(x)+6b} + \frac{12 \log(\tanh(x)+1)}{6b \tanh(x)+6b} - \frac{2 \tanh^4(x)}{6b \tanh(x)+6b} + \frac{\tanh^3(x)}{6b \tanh(x)+6b} - \frac{9 \tanh^2(x)}{6b \tanh(x)+6b} + \frac{15}{6b \tanh(x)+6b} & \text{for } a = b \\ \frac{6a^5 \log\left(\frac{a}{b} + \tanh(x)\right)}{6a^2b^4-6b^6} - \frac{6a^4b \tanh(x)}{6a^2b^4-6b^6} + \frac{3a^3b^2 \tanh^2(x)}{6a^2b^4-6b^6} - \frac{2a^2b^3 \tanh^3(x)}{6a^2b^4-6b^6} + \frac{6ab^4x}{6a^2b^4-6b^6} - \frac{6ab^4 \log(\tanh(x)+1)}{6a^2b^4-6b^6} - \frac{3ab^4 \tanh^2(x)}{6a^2b^4-6b^6} - \frac{6b^5x}{6a^2b^4-6b^6} + \frac{2b^5 \tanh^3(x)}{6a^2b^4-6b^6} + \frac{6b^5 \tanh(x)}{6a^2b^4-6b^6} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)**5/(a+b*tanh(x)),x)

[Out] Piecewise((zoo*(x - tanh(x)**3/3 - tanh(x)), Eq(a, 0) & Eq(b, 0)), ((x - log(tanh(x) + 1) - tanh(x)**4/4 - tanh(x)**2/2)/a, Eq(b, 0)), (27*x*tanh(x)/(6*b*tanh(x) - 6*b) - 27*x/(6*b*tanh(x) - 6*b) - 12*log(tanh(x) + 1)*tanh(x)/(6*b*tanh(x) - 6*b) + 12*log(tanh(x) + 1)/(6*b*tanh(x) - 6*b) - 2*tanh(x)**4/(6*b*tanh(x) - 6*b) - tanh(x)**3/(6*b*tanh(x) - 6*b) - 9*tanh(x)**2/(6*b*tanh(x) - 6*b) + 15/(6*b*tanh(x) - 6*b), Eq(a, -b)), (3*x*tanh(x)/(6*b*tanh(x) + 6*b) + 3*x/(6*b*tanh(x) + 6*b) + 12*log(tanh(x) + 1)*tanh(x)/(6*b*tanh(x) + 6*b) + 12*log(tanh(x) + 1)/(6*b*tanh(x) + 6*b) - 2*tanh(x)**4/(6*b*tanh(x) + 6*b) + tanh(x)**3/(6*b*tanh(x) + 6*b) - 9*tanh(x)**2/(6*b*tanh(x) + 6*b) + 15/(6*b*tanh(x) + 6*b), Eq(a, b)), (6*a**5*log(a/b + tanh(x))/(6*a**2*b**4 - 6*b**6) - 6*a**4*b*tanh(x)/(6*a**2*b**4 - 6*b**6) + 3*a**3*b**2*tanh(x)**2/(6*a**2*b**4 - 6*b**6) - 2*a**2*b**3*tanh(x)**3/(6*a**2*b**4 - 6*b**6) + 6*a*b**4*x/(6*a**2*b**4 - 6*b**6) - 6*a*b**4*log(tanh(x) + 1)/(6*a**2*b**4 - 6*b**6) - 3*a*b**4*tanh(x)**2/(6*a**2*b**4 - 6*b**6) - 6*b**5*x/(6*a**2*b**4 - 6*b**6) + 2*b**5*tanh(x)**3/(6*a**2*b**4 - 6*b**6) + 6*b**5*tanh(x)/(6*a**2*b**4 - 6*b**6), True))

Giac [A]

time = 0.41, size = 142, normalized size = 1.51

$$\frac{a^5 \log(|ae^{2x} + be^{2x} + a - b|)}{a^2b^4 - b^6} - \frac{x}{a - b} - \frac{(a^3 + ab^2) \log(e^{2x} + 1)}{b^4} + \frac{2(3a^2b + 4b^3 + 3(a^2b - ab^2 + 2b^3)e^{4x} + 3(2a^2b - ab^2 + 2b^3)e^{2x})}{3b^4(e^{2x} + 1)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)^5/(a+b*tanh(x)),x, algorithm="giac")

[Out] $a^5 \log(\operatorname{abs}(a e^{2x} + b e^{2x} + a - b)) / (a^2 b^4 - b^6) - x / (a - b) - (a^3 + a b^2) \log(e^{2x} + 1) / b^4 + 2/3 (3 a^2 b + 4 b^3 + 3 (a^2 b - a b^2 + 2 b^3) e^{4x} + 3 (2 a^2 b - a b^2 + 2 b^3) e^{2x}) / (b^4 (e^{2x} + 1)^3)$

Mupad [B]

time = 0.23, size = 85, normalized size = 0.90

$$\frac{x}{a+b} - \frac{\tanh(x)^3}{3b} - \frac{a \ln(\tanh(x)+1)}{a^2-b^2} + \frac{a \tanh(x)^2}{2b^2} - \frac{\tanh(x)(a^2+b^2)}{b^3} + \frac{a^5 \ln(a+b \tanh(x))}{b^4(a^2-b^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(x)^5/(a + b*tanh(x)),x)

[Out] $x/(a+b) - \tanh(x)^3/(3b) - (a \log(\tanh(x)+1))/(a^2-b^2) + (a \tanh(x)^2)/(2b^2) - (\tanh(x)(a^2+b^2))/b^3 + (a^5 \log(a+b \tanh(x)))/(b^4(a^2-b^2))$

3.134 $\int \frac{\tanh^4(x)}{a+b \tanh(x)} dx$

Optimal. Leaf size=76

$$\frac{ax}{a^2 - b^2} - \frac{b \log(\cosh(x))}{a^2 - b^2} - \frac{a^4 \log(a + b \tanh(x))}{b^3 (a^2 - b^2)} + \frac{a \tanh(x)}{b^2} - \frac{\tanh^2(x)}{2b}$$

[Out] a*x/(a^2-b^2)-b*ln(cosh(x))/(a^2-b^2)-a^4*ln(a+b*tanh(x))/b^3/(a^2-b^2)+a*tanh(x)/b^2-1/2*tanh(x)^2/b

Rubi [A]

time = 0.16, antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.462$, Rules used = {3647, 3728, 3708, 3698, 31, 3556}

$$\frac{ax}{a^2 - b^2} - \frac{b \log(\cosh(x))}{a^2 - b^2} - \frac{a^4 \log(a + b \tanh(x))}{b^3 (a^2 - b^2)} + \frac{a \tanh(x)}{b^2} - \frac{\tanh^2(x)}{2b}$$

Antiderivative was successfully verified.

[In] Int[Tanh[x]^4/(a + b*Tanh[x]),x]

[Out] (a*x)/(a^2 - b^2) - (b*Log[Cosh[x]])/(a^2 - b^2) - (a^4*Log[a + b*Tanh[x]])/(b^3*(a^2 - b^2)) + (a*Tanh[x])/b^2 - Tanh[x]^2/(2*b)

Rule 31

Int[((a_) + (b_)*(x_))^(n_), x_Symbol] := Simp[Log[RemoveContent[a + b*x, x]]/b, x] /; FreeQ[{a, b}, x]

Rule 3556

Int[tan[(c_) + (d_)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]

Rule 3647

Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b^2*(a + b*Tan[e + f*x])^(m - 2)*((c + d*Tan[e + f*x])^(n + 1)/(d*(m + n - 1))), x] + Dist[1/(d*(m + n - 1)), Int[(a + b*Tan[e + f*x])^(m - 3)*(c + d*Tan[e + f*x])^n*Simp[a^3*d*(m + n - 1) - b^2*(b*c*(m - 2) + a*d*(1 + n)) + b*d*(m + n - 1)*(3*a^2 - b^2)*Tan[e + f*x] - b^2*(b*c*(m - 2) - a*d*(3*m + 2*n - 4))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && IntegerQ[2*m] && GtQ[m, 2] && (GeQ[n, -1] || IntegerQ[m]) && !(IGtQ[n, 2] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))

Rule 3698

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)^2]), x_Symbol] := Dist[A/(b*f), Subst[Int[(a + x)^m, x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, A, C, m}, x] && EqQ[A, C]

Rule 3708

Int[((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[a*(A - C)*(x/(a^2 + b^2)), x] + (Dist[(a^2*C + A*b^2)/(a^2 + b^2), Int[(1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x]), x], x] - Dist[b*((A - C)/(a^2 + b^2)), Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, e, f, A, C}, x] && NeQ[a^2*C + A*b^2, 0] && NeQ[a^2 + b^2, 0] && NeQ[A, C]

Rule 3728

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)^2]), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Dist[1/(d*(m + n + 1)), Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && (!IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))

Rubi steps

$$\begin{aligned}
 \int \frac{\tanh^4(x)}{a + b \tanh(x)} dx &= -\frac{\tanh^2(x)}{2b} - \frac{\int \frac{\tanh(x)(-2a - 2b \tanh(x) + 2a \tanh^2(x))}{a + b \tanh(x)} dx}{2b} \\
 &= \frac{a \tanh(x)}{b^2} - \frac{\tanh^2(x)}{2b} - \frac{\int \frac{2a^2 - 2(a^2 + b^2) \tanh^2(x)}{a + b \tanh(x)} dx}{2b^2} \\
 &= \frac{ax}{a^2 - b^2} + \frac{a \tanh(x)}{b^2} - \frac{\tanh^2(x)}{2b} - \frac{a^4 \int \frac{1 - \tanh^2(x)}{a + b \tanh(x)} dx}{b^2(a^2 - b^2)} - \frac{b \int \tanh(x) dx}{a^2 - b^2} \\
 &= \frac{ax}{a^2 - b^2} - \frac{b \log(\cosh(x))}{a^2 - b^2} + \frac{a \tanh(x)}{b^2} - \frac{\tanh^2(x)}{2b} - \frac{a^4 \text{Subst}\left(\int \frac{1}{a+x} dx, x, b \tanh(x)\right)}{b^3(a^2 - b^2)} \\
 &= \frac{ax}{a^2 - b^2} - \frac{b \log(\cosh(x))}{a^2 - b^2} - \frac{a^4 \log(a + b \tanh(x))}{b^3(a^2 - b^2)} + \frac{a \tanh(x)}{b^2} - \frac{\tanh^2(x)}{2b}
 \end{aligned}$$

Mathematica [A]

time = 0.22, size = 88, normalized size = 1.16

$$\frac{b^2(a^2 - b^2) \operatorname{sech}^2(x) + 2(ab^3x + (a^4 - b^4) \log(\cosh(x)) - a^4 \log(a \cosh(x) + b \sinh(x)) + ab(a^2 - b^2) \tanh(x))}{2(a - b)b^3(a + b)}$$

Antiderivative was successfully verified.

[In] Integrate[Tanh[x]^4/(a + b*Tanh[x]),x]

[Out] (b^2*(a^2 - b^2)*Sech[x]^2 + 2*(a*b^3*x + (a^4 - b^4)*Log[Cosh[x]] - a^4*Log[a*Cosh[x] + b*Sinh[x]] + a*b*(a^2 - b^2)*Tanh[x]))/(2*(a - b)*b^3*(a + b))

Maple [A]

time = 0.32, size = 76, normalized size = 1.00

method	result	size
derivativedivides	$-\frac{b(\tanh^2(x))}{2} + a \tanh(x) \Big/ b^2 + \frac{\ln(1+\tanh(x))}{2a-2b} - \frac{\ln(\tanh(x)-1)}{2b+2a} - \frac{a^4 \ln(a+b \tanh(x))}{b^3(a+b)(a-b)}$	76
default	$-\frac{b(\tanh^2(x))}{2} + a \tanh(x) \Big/ b^2 + \frac{\ln(1+\tanh(x))}{2a-2b} - \frac{\ln(\tanh(x)-1)}{2b+2a} - \frac{a^4 \ln(a+b \tanh(x))}{b^3(a+b)(a-b)}$	76
risch	$\frac{x}{a+b} - \frac{2x a^2}{b^3} - \frac{2x}{b} + \frac{2x a^4}{b^3(a^2-b^2)} - \frac{2(e^{2x}a - b e^{2x} + a)}{(1+e^{2x})^2 b^2} + \frac{\ln(1+e^{2x}) a^2}{b^3} + \frac{\ln(1+e^{2x})}{b} - \frac{a^4 \ln(e^{2x} + \frac{a-b}{a+b})}{b^3(a^2-b^2)}$	133

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(x)^4/(a+b*tanh(x)),x,method=_RETURNVERBOSE)

[Out] 1/b^2*(-1/2*b*tanh(x)^2+a*tanh(x))+1/(2*a-2*b)*ln(1+tanh(x))-1/(2*b+2*a)*ln(tanh(x)-1)-1/b^3*a^4/(a+b)/(a-b)*ln(a+b*tanh(x))

Maxima [A]

time = 0.48, size = 100, normalized size = 1.32

$$-\frac{a^4 \log(-(a - b)e^{(-2x)} - a - b)}{a^2 b^3 - b^5} + \frac{2((a + b)e^{(-2x)} + a)}{2 b^2 e^{(-2x)} + b^2 e^{(-4x)} + b^2} + \frac{x}{a + b} + \frac{(a^2 + b^2) \log(e^{(-2x)} + 1)}{b^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)^4/(a+b*tanh(x)),x, algorithm="maxima")

[Out] -a^4*log(-(a - b)*e^(-2*x) - a - b)/(a^2*b^3 - b^5) + 2*((a + b)*e^(-2*x) + a)/(2*b^2*e^(-2*x) + b^2*e^(-4*x) + b^2) + x/(a + b) + (a^2 + b^2)*log(e^(-2*x) + 1)/b^3

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 644 vs. 2(74) = 148.

time = 0.37, size = 644, normalized size = 8.47

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)^4/(a+b*tanh(x)),x, algorithm="fricas")

[Out] $((a*b^3 + b^4)*x*\cosh(x)^4 + 4*(a*b^3 + b^4)*x*\cosh(x)*\sinh(x)^3 + (a*b^3 + b^4)*x*\sinh(x)^4 - 2*a^3*b + 2*a*b^3 - 2*(a^3*b - a^2*b^2 - a*b^3 + b^4 - (a*b^3 + b^4)*x)*\cosh(x)^2 - 2*(a^3*b - a^2*b^2 - a*b^3 + b^4 - 3*(a*b^3 + b^4)*x*\cosh(x)^2 - (a*b^3 + b^4)*x)*\sinh(x)^2 + (a*b^3 + b^4)*x - (a^4*\cosh(x)^4 + 4*a^4*\cosh(x)*\sinh(x)^3 + a^4*\sinh(x)^4 + 2*a^4*\cosh(x)^2 + a^4 + 2*(3*a^4*\cosh(x)^2 + a^4)*\sinh(x)^2 + 4*(a^4*\cosh(x)^3 + a^4*\cosh(x))*\sinh(x))*\log(2*(a*\cosh(x) + b*\sinh(x))/(\cosh(x) - \sinh(x))) + ((a^4 - b^4)*\cosh(x))^4 + 4*(a^4 - b^4)*\cosh(x)*\sinh(x)^3 + (a^4 - b^4)*\sinh(x)^4 + a^4 - b^4 + 2*(a^4 - b^4)*\cosh(x)^2 + 2*(a^4 - b^4 + 3*(a^4 - b^4)*\cosh(x)^2)*\sinh(x)^2 + 4*((a^4 - b^4)*\cosh(x)^3 + (a^4 - b^4)*\cosh(x))*\sinh(x))*\log(2*\cosh(x)/(\cosh(x) - \sinh(x))) + 4*((a*b^3 + b^4)*x*\cosh(x)^3 - (a^3*b - a^2*b^2 - a*b^3 + b^4 - (a*b^3 + b^4)*x)*\cosh(x))*\sinh(x))/(a^2*b^3 - b^5 + (a^2*b^3 - b^5)*\cosh(x)^4 + 4*(a^2*b^3 - b^5)*\cosh(x)*\sinh(x)^3 + (a^2*b^3 - b^5)*\sinh(x)^4 + 2*(a^2*b^3 - b^5)*\cosh(x)^2 + 2*(a^2*b^3 - b^5 + 3*(a^2*b^3 - b^5)*\cosh(x)^2)*\sinh(x)^2 + 4*((a^2*b^3 - b^5)*\cosh(x)^3 + (a^2*b^3 - b^5)*\cosh(x))*\sinh(x))$

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 442 vs. 2(61) = 122.

time = 0.48, size = 442, normalized size = 5.82

$$\begin{cases} \tilde{\infty} \left(x - \log(\tanh(x) + 1) - \frac{\tanh^2(x)}{2} \right) & \text{for } a = 0 \wedge b = 0 \\ \frac{7x \tanh(x)}{2b \tanh(x) - 2b} - \frac{7x}{2b \tanh(x) - 2b} - \frac{4 \log(\tanh(x) + 1) \tanh(x)}{2b \tanh(x) - 2b} + \frac{4 \log(\tanh(x) + 1)}{2b \tanh(x) - 2b} - \frac{\tanh^3(x)}{2b \tanh(x) - 2b} - \frac{\tanh^2(x)}{2b \tanh(x) - 2b} + \frac{3}{2b \tanh(x) - 2b} & \text{for } a = -b \\ \frac{x \tanh(x)}{2b \tanh(x) + 2b} + \frac{x}{2b \tanh(x) + 2b} - \frac{4 \log(\tanh(x) + 1) \tanh(x)}{2b \tanh(x) + 2b} - \frac{4 \log(\tanh(x) + 1)}{2b \tanh(x) + 2b} - \frac{\tanh^3(x)}{2b \tanh(x) + 2b} + \frac{\tanh^2(x)}{2b \tanh(x) + 2b} - \frac{3}{2b \tanh(x) + 2b} & \text{for } a = b \\ \frac{x - \frac{\tanh^3(x)}{3} - \tanh(x)}{a} & \text{for } b = 0 \\ -\frac{2a^4 \log\left(\frac{a}{b} + \tanh(x)\right)}{2a^2 b^3 - 2b^5} + \frac{2a^3 b \tanh(x)}{2a^2 b^3 - 2b^5} - \frac{a^2 b^2 \tanh^2(x)}{2a^2 b^3 - 2b^5} + \frac{2ab^3 x}{2a^2 b^3 - 2b^5} - \frac{2ab^3 \tanh(x)}{2a^2 b^3 - 2b^5} - \frac{2b^4 x}{2a^2 b^3 - 2b^5} + \frac{2b^4 \log(\tanh(x) + 1)}{2a^2 b^3 - 2b^5} + \frac{b^4 \tanh^2(x)}{2a^2 b^3 - 2b^5} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)**4/(a+b*tanh(x)),x)

[Out] Piecewise((zoo*(x - log(tanh(x) + 1) - tanh(x)**2/2), Eq(a, 0) & Eq(b, 0)), (7*x*tanh(x)/(2*b*tanh(x) - 2*b) - 7*x/(2*b*tanh(x) - 2*b) - 4*log(tanh(x) + 1)*tanh(x)/(2*b*tanh(x) - 2*b) + 4*log(tanh(x) + 1)/(2*b*tanh(x) - 2*b) - tanh(x)**3/(2*b*tanh(x) - 2*b) - tanh(x)**2/(2*b*tanh(x) - 2*b) + 3/(2*b*tanh(x) - 2*b), Eq(a, -b)), (x*tanh(x)/(2*b*tanh(x) + 2*b) + x/(2*b*tanh(x) + 2*b) - 4*log(tanh(x) + 1)*tanh(x)/(2*b*tanh(x) + 2*b) - 4*log(tanh(x) + 1)/(2*b*tanh(x) + 2*b) - tanh(x)**3/(2*b*tanh(x) + 2*b) + tanh(x)**2/(2*b*tanh(x) + 2*b) - 3/(2*b*tanh(x) + 2*b), Eq(a, b)), ((x - tanh(x)**3/3 - tanh(x))/a, Eq(b, 0)), (-2*a**4*log(a/b + tanh(x))/(2*a**2*b**3 - 2*b**5) + 2*a**3*b*tanh(x)/(2*a**2*b**3 - 2*b**5) - a**2*b**2*tanh(x)**2/(2*a**2*b**3 - 2*b**5) + 2*a*b**3*x/(2*a**2*b**3 - 2*b**5) - 2*a*b**3*tanh(x)/(2*a**2*b**3 - 2*b**5) + 2*b**4*log(a/b + tanh(x))/(2*a**2*b**3 - 2*b**5) + b**4*tanh(x)**2/(2*a**2*b**3 - 2*b**5), Eq(a, 0) & Eq(b, 0)))

$- 2*b^{**5}) - 2*b^{**4}*x/(2*a^{**2}*b^{**3} - 2*b^{**5}) + 2*b^{**4}*log(tanh(x) + 1)/(2*a^{**2}*b^{**3} - 2*b^{**5}) + b^{**4}*tanh(x)**2/(2*a^{**2}*b^{**3} - 2*b^{**5}), True))$

Giac [A]

time = 0.40, size = 98, normalized size = 1.29

$$-\frac{a^4 \log(|ae^{(2x)} + be^{(2x)} + a - b|)}{a^2 b^3 - b^5} + \frac{x}{a - b} + \frac{(a^2 + b^2) \log(e^{(2x)} + 1)}{b^3} - \frac{2(ab + (ab - b^2)e^{(2x)})}{b^3(e^{(2x)} + 1)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)^4/(a+b*tanh(x)),x, algorithm="giac")

[Out] $-a^4*\log(\text{abs}(a*e^{(2*x)} + b*e^{(2*x)} + a - b))/(a^2*b^3 - b^5) + x/(a - b) + (a^2 + b^2)*\log(e^{(2*x)} + 1)/b^3 - 2*(a*b + (a*b - b^2)*e^{(2*x)})/(b^3*(e^{(2*x)} + 1)^2)$

Mupad [B]

time = 0.17, size = 68, normalized size = 0.89

$$\frac{x}{a + b} - \frac{\tanh(x)^2}{2b} + \frac{b \ln(\tanh(x) + 1)}{a^2 - b^2} + \frac{a \tanh(x)}{b^2} - \frac{a^4 \ln(a + b \tanh(x))}{b^3 (a^2 - b^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(x)^4/(a + b*tanh(x)),x)

[Out] $x/(a + b) - \tanh(x)^2/(2*b) + (b*\log(\tanh(x) + 1))/(a^2 - b^2) + (a*\tanh(x))/b^2 - (a^4*\log(a + b*\tanh(x)))/(b^3*(a^2 - b^2))$

$$3.135 \quad \int \frac{\tanh^3(x)}{a+b \tanh(x)} dx$$

Optimal. Leaf size=64

$$-\frac{bx}{a^2-b^2} + \frac{a \log(\cosh(x))}{a^2-b^2} + \frac{a^3 \log(a+b \tanh(x))}{b^2(a^2-b^2)} - \frac{\tanh(x)}{b}$$

[Out] $-b*x/(a^2-b^2)+a*\ln(\cosh(x))/(a^2-b^2)+a^3*\ln(a+b*\tanh(x))/b^2/(a^2-b^2)-\tanh(x)/b$

Rubi [A]

time = 0.10, antiderivative size = 64, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {3647, 3707, 3698, 31, 3556}

$$-\frac{bx}{a^2-b^2} + \frac{a \log(\cosh(x))}{a^2-b^2} + \frac{a^3 \log(a+b \tanh(x))}{b^2(a^2-b^2)} - \frac{\tanh(x)}{b}$$

Antiderivative was successfully verified.

[In] `Int[Tanh[x]^3/(a + b*Tanh[x]), x]`

[Out] $-((b*x)/(a^2 - b^2)) + (a*\text{Log}[\text{Cosh}[x]])/(a^2 - b^2) + (a^3*\text{Log}[a + b*\text{Tanh}[x]])/(b^2*(a^2 - b^2)) - \text{Tanh}[x]/b$

Rule 31

`Int[((a_) + (b_)*(x_))^(n_), x_Symbol] := Simp[Log[RemoveContent[a + b*x, x]]/b, x] /; FreeQ[{a, b}, x]`

Rule 3556

`Int[tan[(c_) + (d_)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

Rule 3647

`Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b^2*(a + b*Tan[e + f*x])^(m-2)*((c + d*Tan[e + f*x])^(n+1)/(d*f*(m+n-1))), x] + Dist[1/(d*(m+n-1)), Int[(a + b*Tan[e + f*x])^(m-3)*(c + d*Tan[e + f*x])^n*Simp[a^3*d*(m+n-1) - b^2*(b*c*(m-2) + a*d*(1+n)) + b*d*(m+n-1)*(3*a^2 - b^2)*Tan[e + f*x] - b^2*(b*c*(m-2) - a*d*(3*m+2*n-4))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && IntegerQ[2*m] && GtQ[m, 2] && (GeQ[n, -1] || IntegerQ[m]) && !(IGtQ[n, 2] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

Rule 3698

```
Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_) + (C_.)*tan[(e_.) +
(f_.)*(x_)^2], x_Symbol] := Dist[A/(b*f), Subst[Int[(a + x)^m, x], x, b*T
an[e + f*x]], x] /; FreeQ[{a, b, e, f, A, C, m}, x] && EqQ[A, C]
```

Rule 3707

```
Int[((A_) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)^2
]/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*A + b*B - a
*C)*(x/(a^2 + b^2)), x] + (Dist[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2), Int[(1
+ Tan[e + f*x]^2)/(a + b*Tan[e + f*x]), x], x] - Dist[(A*b - a*B - b*C)/(a
^2 + b^2), Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, e, f, A, B, C}, x] &&
NeQ[A*b^2 - a*b*B + a^2*C, 0] && NeQ[a^2 + b^2, 0] && NeQ[A*b - a*B - b*C,
0]
```

Rubi steps

$$\begin{aligned} \int \frac{\tanh^3(x)}{a + b \tanh(x)} dx &= -\frac{\tanh(x)}{b} - \frac{\int \frac{-a - b \tanh(x) + a \tanh^2(x)}{a + b \tanh(x)} dx}{b} \\ &= -\frac{bx}{a^2 - b^2} - \frac{\tanh(x)}{b} + \frac{a \int \tanh(x) dx}{a^2 - b^2} + \frac{a^3 \int \frac{1 - \tanh^2(x)}{a + b \tanh(x)} dx}{b(a^2 - b^2)} \\ &= -\frac{bx}{a^2 - b^2} + \frac{a \log(\cosh(x))}{a^2 - b^2} - \frac{\tanh(x)}{b} + \frac{a^3 \text{Subst}\left(\int \frac{1}{a+x} dx, x, b \tanh(x)\right)}{b^2(a^2 - b^2)} \\ &= -\frac{bx}{a^2 - b^2} + \frac{a \log(\cosh(x))}{a^2 - b^2} + \frac{a^3 \log(a + b \tanh(x))}{b^2(a^2 - b^2)} - \frac{\tanh(x)}{b} \end{aligned}$$

Mathematica [A]

time = 0.10, size = 65, normalized size = 1.02

$$\frac{-b^3 x + (-a^3 + ab^2) \log(\cosh(x)) + a^3 \log(a \cosh(x) + b \sinh(x)) + (-a^2 b + b^3) \tanh(x)}{(a - b)b^2(a + b)}$$

Antiderivative was successfully verified.

```
[In] Integrate[Tanh[x]^3/(a + b*Tanh[x]), x]
```

```
[Out] (-b^3*x) + (-a^3 + a*b^2)*Log[Cosh[x]] + a^3*Log[a*Cosh[x] + b*Sinh[x]] +
(-a^2*b) + b^3)*Tanh[x])/((a - b)*b^2*(a + b))
```

Maple [A]

time = 0.36, size = 67, normalized size = 1.05

method	result	size
derivativedivides	$-\frac{\tanh(x)}{b} + \frac{a^3 \ln(a+b \tanh(x))}{b^2(a+b)(a-b)} - \frac{\ln(1+\tanh(x))}{2a-2b} - \frac{\ln(\tanh(x)-1)}{2b+2a}$	67
default	$-\frac{\tanh(x)}{b} + \frac{a^3 \ln(a+b \tanh(x))}{b^2(a+b)(a-b)} - \frac{\ln(1+\tanh(x))}{2a-2b} - \frac{\ln(\tanh(x)-1)}{2b+2a}$	67
risch	$\frac{x}{a+b} + \frac{2ax}{b^2} - \frac{2a^3x}{b^2(a^2-b^2)} + \frac{2}{b(1+e^{2x})} - \frac{a \ln(1+e^{2x})}{b^2} + \frac{a^3 \ln\left(e^{2x} + \frac{a-b}{a+b}\right)}{b^2(a^2-b^2)}$	97

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tanh(x)^3/(a+b*tanh(x)),x,method=_RETURNVERBOSE)`

[Out] $-\tanh(x)/b+1/b^2*a^3/(a+b)/(a-b)*\ln(a+b*\tanh(x))-1/(2*a-2*b)*\ln(1+\tanh(x))-1/(2*b+2*a)*\ln(\tanh(x)-1)$

Maxima [A]

time = 0.49, size = 71, normalized size = 1.11

$$\frac{a^3 \log\left(-(a-b)e^{(-2x)} - a - b\right)}{a^2b^2 - b^4} + \frac{x}{a+b} - \frac{a \log\left(e^{(-2x)} + 1\right)}{b^2} - \frac{2}{be^{(-2x)} + b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(x)^3/(a+b*tanh(x)),x, algorithm="maxima")`

[Out] $a^3*\log(-(a - b)*e^{(-2*x)} - a - b)/(a^2*b^2 - b^4) + x/(a + b) - a*\log(e^{(-2*x)} + 1)/b^2 - 2/(b*e^{(-2*x)} + b)$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 264 vs. 2(64) = 128.

time = 0.43, size = 264, normalized size = 4.12

$$\frac{(ab^2 + b^3)x \cosh(x)^2 + 2(ab^2 + b^3)x \cosh(x) \sinh(x) + (ab^2 + b^3)x \sinh(x)^2 - 2a^2b + 2b^3 + (ab^2 + b^3)x - (a^3 \cosh(x)^2 + 2a^3 \cosh(x) \sinh(x) + a^3 \sinh(x)^2 + a^3) \log\left(\frac{2 \cosh(x) + \tanh(x)}{1 + \tanh(x)}\right) + (a^3 - ab^2 + (a^3 - ab^2) \cosh(x)^2 + 2(a^3 - ab^2) \cosh(x) \sinh(x) + (a^3 - ab^2) \sinh(x)^2) \log\left(\frac{2 \cosh(x) - \tanh(x)}{1 - \tanh(x)}\right)}{a^2b^2 - b^4 + (a^2b^2 - b^4) \cosh(x)^2 + 2(a^2b^2 - b^4) \cosh(x) \sinh(x) + (a^2b^2 - b^4) \sinh(x)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(x)^3/(a+b*tanh(x)),x, algorithm="fricas")`

[Out] $-\left((a*b^2 + b^3)*x*\cosh(x)^2 + 2*(a*b^2 + b^3)*x*\cosh(x)*\sinh(x) + (a*b^2 + b^3)*x*\sinh(x)^2 - 2*a^2*b + 2*b^3 + (a*b^2 + b^3)*x - (a^3*\cosh(x)^2 + 2*a^3*\cosh(x)*\sinh(x) + a^3*\sinh(x)^2 + a^3)*\log(2*(a*\cosh(x) + b*\sinh(x))/(\cosh(x) - \sinh(x))) + (a^3 - a*b^2 + (a^3 - a*b^2)*\cosh(x)^2 + 2*(a^3 - a*b^2)*\cosh(x)*\sinh(x) + (a^3 - a*b^2)*\sinh(x)^2)*\log(2*\cosh(x)/(\cosh(x) - \sinh(x)))\right)/(a^2*b^2 - b^4 + (a^2*b^2 - b^4)*\cosh(x)^2 + 2*(a^2*b^2 - b^4)*\cosh(x)*\sinh(x) + (a^2*b^2 - b^4)*\sinh(x)^2)$

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 330 vs. 2(49) = 98.

time = 0.38, size = 330, normalized size = 5.16

$$\left\{ \begin{array}{ll} \infty(x - \tanh(x)) & \text{for } a = 0 \wedge b = 0 \\ \frac{x - \log(\tanh(x)+1) - \frac{\tanh^2(x)}{2}}{a} & \text{for } b = 0 \\ \frac{5x \tanh(x)}{2b \tanh(x)-2b} - \frac{5x}{2b \tanh(x)-2b} - \frac{2 \log(\tanh(x)+1) \tanh(x)}{2b \tanh(x)-2b} + \frac{2 \log(\tanh(x)+1)}{2b \tanh(x)-2b} - \frac{2 \tanh^2(x)}{2b \tanh(x)-2b} + \frac{3}{2b \tanh(x)-2b} & \text{for } a = -b \\ \frac{x \tanh(x)}{2b \tanh(x)+2b} + \frac{x}{2b \tanh(x)+2b} + \frac{2 \log(\tanh(x)+1) \tanh(x)}{2b \tanh(x)+2b} + \frac{2 \log(\tanh(x)+1)}{2b \tanh(x)+2b} - \frac{2 \tanh^2(x)}{2b \tanh(x)+2b} + \frac{3}{2b \tanh(x)+2b} & \text{for } a = b \\ \frac{a^3 \log\left(\frac{a}{b} + \tanh(x)\right)}{a^2 b^2 - b^4} - \frac{a^2 b \tanh(x)}{a^2 b^2 - b^4} + \frac{a b^2 x}{a^2 b^2 - b^4} - \frac{a b^2 \log(\tanh(x)+1)}{a^2 b^2 - b^4} - \frac{b^3 x}{a^2 b^2 - b^4} + \frac{b^3 \tanh(x)}{a^2 b^2 - b^4} & \text{otherwise} \end{array} \right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)**3/(a+b*tanh(x)),x)

[Out] Piecewise((zoo*(x - tanh(x)), Eq(a, 0) & Eq(b, 0)), ((x - log(tanh(x) + 1) - tanh(x)**2/2)/a, Eq(b, 0)), (5*x*tanh(x)/(2*b*tanh(x) - 2*b) - 5*x/(2*b*tanh(x) - 2*b) - 2*log(tanh(x) + 1)*tanh(x)/(2*b*tanh(x) - 2*b) + 2*log(tanh(x) + 1)/(2*b*tanh(x) - 2*b) - 2*tanh(x)**2/(2*b*tanh(x) - 2*b) + 3/(2*b*tanh(x) - 2*b), Eq(a, -b)), (x*tanh(x)/(2*b*tanh(x) + 2*b) + x/(2*b*tanh(x) + 2*b) + 2*log(tanh(x) + 1)*tanh(x)/(2*b*tanh(x) + 2*b) + 2*log(tanh(x) + 1)/(2*b*tanh(x) + 2*b) - 2*tanh(x)**2/(2*b*tanh(x) + 2*b) + 3/(2*b*tanh(x) + 2*b), Eq(a, b)), (a**3*log(a/b + tanh(x))/(a**2*b**2 - b**4) - a**2*b*tanh(x)/(a**2*b**2 - b**4) + a*b**2*x/(a**2*b**2 - b**4) - a*b**2*log(tanh(x) + 1)/(a**2*b**2 - b**4) - b**3*x/(a**2*b**2 - b**4) + b**3*tanh(x)/(a**2*b**2 - b**4), True))

Giac [A]

time = 0.41, size = 75, normalized size = 1.17

$$\frac{a^3 \log(|ae^{(2x)} + be^{(2x)} + a - b|)}{a^2 b^2 - b^4} - \frac{x}{a - b} - \frac{a \log(e^{(2x)} + 1)}{b^2} + \frac{2}{b(e^{(2x)} + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)^3/(a+b*tanh(x)),x, algorithm="giac")

[Out] a^3*log(abs(a*e^(2*x) + b*e^(2*x) + a - b))/(a^2*b^2 - b^4) - x/(a - b) - a*log(e^(2*x) + 1)/b^2 + 2/(b*(e^(2*x) + 1))

Mupad [B]

time = 1.09, size = 59, normalized size = 0.92

$$\frac{x}{a + b} - \frac{\tanh(x)}{b} - \frac{a \ln(\tanh(x) + 1)}{a^2 - b^2} + \frac{a^3 \ln(a + b \tanh(x))}{b^2 (a^2 - b^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(x)^3/(a + b*tanh(x)),x)

[Out] x/(a + b) - tanh(x)/b - (a*log(tanh(x) + 1))/(a^2 - b^2) + (a^3*log(a + b*tanh(x)))/(b^2*(a^2 - b^2))

3.136 $\int \frac{\tanh^2(x)}{a+b \tanh(x)} dx$

Optimal. Leaf size=63

$$-\frac{ax}{b^2} + \frac{a^3x}{b^2(a^2 - b^2)} + \frac{\log(\cosh(x))}{b} - \frac{a^2 \log(a \cosh(x) + b \sinh(x))}{b(a^2 - b^2)}$$

[Out] $-a*x/b^2+a^3*x/b^2/(a^2-b^2)+\ln(\cosh(x))/b-a^2*\ln(a*\cosh(x)+b*\sinh(x))/b/(a^2-b^2)$

Rubi [A]

time = 0.07, antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {3622, 3556, 3565, 3611}

$$-\frac{a^2 \log(a \cosh(x) + b \sinh(x))}{b(a^2 - b^2)} + \frac{a^3x}{b^2(a^2 - b^2)} - \frac{ax}{b^2} + \frac{\log(\cosh(x))}{b}$$

Antiderivative was successfully verified.

[In] Int[Tanh[x]^2/(a + b*Tanh[x]),x]

[Out] $-((a*x)/b^2) + (a^3*x)/(b^2*(a^2 - b^2)) + \text{Log}[\text{Cosh}[x]]/b - (a^2*\text{Log}[a*\text{Cosh}[x] + b*\text{Sinh}[x]])/(b*(a^2 - b^2))$

Rule 3556

Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]

Rule 3565

Int[((a_.) + (b_.)*tan[(c_.) + (d_.)*(x_.)])^(-1), x_Symbol] := Simp[a*(x/(a^2 + b^2)), x] + Dist[b/(a^2 + b^2), Int[(b - a*Tan[c + d*x])/(a + b*Tan[c + d*x]), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0]

Rule 3611

Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]

Rule 3622

Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)])^2/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)]), x_Symbol] := Simp[d*(2*b*c - a*d)*(x/b^2), x] + (Dist[d^2/b, In

```
t[Tan[e + f*x], x], x] + Dist[(b*c - a*d)^2/b^2, Int[1/(a + b*Tan[e + f*x])
, x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 +
b^2, 0]
```

Rubi steps

$$\begin{aligned} \int \frac{\tanh^2(x)}{a + b \tanh(x)} dx &= -\frac{ax}{b^2} + \frac{a^2 \int \frac{1}{a+b \tanh(x)} dx}{b^2} + \frac{\int \tanh(x) dx}{b} \\ &= -\frac{ax}{b^2} + \frac{a^3 x}{b^2 (a^2 - b^2)} + \frac{\log(\cosh(x))}{b} - \frac{(ia^2) \int \frac{-ib-ia \tanh(x)}{a+b \tanh(x)} dx}{b(a^2 - b^2)} \\ &= -\frac{ax}{b^2} + \frac{a^3 x}{b^2 (a^2 - b^2)} + \frac{\log(\cosh(x))}{b} - \frac{a^2 \log(a \cosh(x) + b \sinh(x))}{b(a^2 - b^2)} \end{aligned}$$

Mathematica [A]

time = 0.07, size = 49, normalized size = 0.78

$$\frac{abx + a^2 \log(\cosh(x)) - b^2 \log(\cosh(x)) - a^2 \log(a \cosh(x) + b \sinh(x))}{a^2 b - b^3}$$

Antiderivative was successfully verified.

```
[In] Integrate[Tanh[x]^2/(a + b*Tanh[x]), x]
```

```
[Out] (a*b*x + a^2*Log[Cosh[x]] - b^2*Log[Cosh[x]] - a^2*Log[a*Cosh[x] + b*Sinh[x
]])/(a^2*b - b^3)
```

Maple [A]

time = 0.26, size = 60, normalized size = 0.95

method	result	size
derivativedivides	$-\frac{a^2 \ln(a+b \tanh(x))}{(a+b)(a-b)b} - \frac{\ln(\tanh(x)-1)}{2b+2a} + \frac{\ln(1+\tanh(x))}{2a-2b}$	60
default	$-\frac{a^2 \ln(a+b \tanh(x))}{(a+b)(a-b)b} - \frac{\ln(\tanh(x)-1)}{2b+2a} + \frac{\ln(1+\tanh(x))}{2a-2b}$	60
risch	$\frac{x}{a+b} + \frac{2x a^2}{b(a^2-b^2)} - \frac{2x}{b} - \frac{a^2 \ln\left(e^{2x} + \frac{a-b}{a+b}\right)}{b(a^2-b^2)} + \frac{\ln(1+e^{2x})}{b}$	82

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(tanh(x)^2/(a+b*tanh(x)), x, method=_RETURNVERBOSE)
```

```
[Out] -a^2/(a+b)/(a-b)/b*ln(a+b*tanh(x))-1/(2*b+2*a)*ln(tanh(x)-1)+1/(2*a-2*b)*ln
(1+tanh(x))
```

Maxima [A]

time = 0.48, size = 56, normalized size = 0.89

$$-\frac{a^2 \log(-(a-b)e^{(-2x)} - a - b)}{a^2b - b^3} + \frac{x}{a+b} + \frac{\log(e^{(-2x)} + 1)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)^2/(a+b*tanh(x)),x, algorithm="maxima")**[Out]** -a^2*log(-(a - b)*e^(-2*x) - a - b)/(a^2*b - b^3) + x/(a + b) + log(e^(-2*x) + 1)/b**Fricas [A]**

time = 0.42, size = 76, normalized size = 1.21

$$-\frac{a^2 \log\left(\frac{2(a \cosh(x) + b \sinh(x))}{\cosh(x) - \sinh(x)}\right) - (ab + b^2)x - (a^2 - b^2) \log\left(\frac{2 \cosh(x)}{\cosh(x) - \sinh(x)}\right)}{a^2b - b^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)^2/(a+b*tanh(x)),x, algorithm="fricas")**[Out]** -(a^2*log(2*(a*cosh(x) + b*sinh(x))/(cosh(x) - sinh(x))) - (a*b + b^2)*x - (a^2 - b^2)*log(2*cosh(x)/(cosh(x) - sinh(x))))/(a^2*b - b^3)**Sympy [B]** Leaf count of result is larger than twice the leaf count of optimal. 243 vs. 2(51) = 102.

time = 0.32, size = 243, normalized size = 3.86

$$\left\{ \begin{array}{ll} \tilde{\infty}(x - \log(\tanh(x) + 1)) & \text{for } a = 0 \wedge b = 0 \\ \frac{x - \tanh(x)}{a} & \text{for } b = 0 \\ \frac{3x \tanh(x)}{2b \tanh(x) - 2b} - \frac{3x}{2b \tanh(x) - 2b} - \frac{2 \log(\tanh(x) + 1) \tanh(x)}{2b \tanh(x) - 2b} + \frac{2 \log(\tanh(x) + 1)}{2b \tanh(x) - 2b} + \frac{1}{2b \tanh(x) - 2b} & \text{for } a = -b \\ \frac{x \tanh(x)}{2b \tanh(x) + 2b} + \frac{x}{2b \tanh(x) + 2b} - \frac{2 \log(\tanh(x) + 1) \tanh(x)}{2b \tanh(x) + 2b} - \frac{2 \log(\tanh(x) + 1)}{2b \tanh(x) + 2b} - \frac{1}{2b \tanh(x) + 2b} & \text{for } a = b \\ -\frac{a^2 \log\left(\frac{a}{b} + \tanh(x)\right)}{a^2b - b^3} + \frac{abx}{a^2b - b^3} - \frac{b^2x}{a^2b - b^3} + \frac{b^2 \log(\tanh(x) + 1)}{a^2b - b^3} & \text{otherwise} \end{array} \right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)**2/(a+b*tanh(x)),x)**[Out]** Piecewise((zoo*(x - log(tanh(x) + 1)), Eq(a, 0) & Eq(b, 0)), ((x - tanh(x))/a, Eq(b, 0)), (3*x*tanh(x)/(2*b*tanh(x) - 2*b) - 3*x/(2*b*tanh(x) - 2*b) - 2*log(tanh(x) + 1)*tanh(x)/(2*b*tanh(x) - 2*b) + 2*log(tanh(x) + 1)/(2*b*tanh(x) - 2*b) + 1/(2*b*tanh(x) - 2*b), Eq(a, -b)), (x*tanh(x)/(2*b*tanh(x) + 2*b) + x/(2*b*tanh(x) + 2*b) - 2*log(tanh(x) + 1)*tanh(x)/(2*b*tanh(x) + 2*b) - 2*log(tanh(x) + 1)/(2*b*tanh(x) + 2*b) - 1/(2*b*tanh(x) + 2*b), Eq(a

, b)), (-a**2*log(a/b + tanh(x))/(a**2*b - b**3) + a*b*x/(a**2*b - b**3) - b**2*x/(a**2*b - b**3) + b**2*log(tanh(x) + 1)/(a**2*b - b**3), True))

Giac [A]

time = 0.41, size = 58, normalized size = 0.92

$$-\frac{a^2 \log(|ae^{(2x)} + be^{(2x)} + a - b|)}{a^2b - b^3} + \frac{x}{a - b} + \frac{\log(e^{(2x)} + 1)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)^2/(a+b*tanh(x)),x, algorithm="giac")

[Out] -a^2*log(abs(a*e^(2*x) + b*e^(2*x) + a - b))/(a^2*b - b^3) + x/(a - b) + log(e^(2*x) + 1)/b

Mupad [B]

time = 0.12, size = 46, normalized size = 0.73

$$-\frac{b^2(x - \ln(\tanh(x) + 1)) + a^2 \ln(a + b \tanh(x)) - abx}{b(a^2 - b^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(x)^2/(a + b*tanh(x)),x)

[Out] -(b^2*(x - log(tanh(x) + 1)) + a^2*log(a + b*tanh(x)) - a*b*x)/(b*(a^2 - b^2))

$$3.137 \quad \int \frac{\tanh(x)}{a+b \tanh(x)} dx$$

Optimal. Leaf size=39

$$-\frac{bx}{a^2 - b^2} + \frac{a \log(a \cosh(x) + b \sinh(x))}{a^2 - b^2}$$

[Out] $-b*x/(a^2-b^2)+a*\ln(a*\cosh(x)+b*\sinh(x))/(a^2-b^2)$

Rubi [A]

time = 0.05, antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {3612, 3611}

$$\frac{a \log(a \cosh(x) + b \sinh(x))}{a^2 - b^2} - \frac{bx}{a^2 - b^2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Tanh}[x]/(a + b*\text{Tanh}[x]), x]$

[Out] $-((b*x)/(a^2 - b^2)) + (a*\text{Log}[a*\text{Cosh}[x] + b*\text{Sinh}[x]])/(a^2 - b^2)$

Rule 3611

$\text{Int}[(c + (d_*)\tan[(e_*) + (f_*)(x_)])/(a + (b_*)\tan[(e_*) + (f_*)(x_)]), x_Symbol] \rightarrow \text{Simp}[(c/(b*f))*\text{Log}[\text{RemoveContent}[a*\text{Cos}[e + f*x] + b*\text{Sin}[e + f*x], x]], x] /;$ FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]

Rule 3612

$\text{Int}[(c + (d_*)\tan[(e_*) + (f_*)(x_)])/(a + (b_*)\tan[(e_*) + (f_*)(x_)]), x_Symbol] \rightarrow \text{Simp}[(a*c + b*d)*(x/(a^2 + b^2)), x] + \text{Dist}[(b*c - a*d)/(a^2 + b^2), \text{Int}[(b - a*\text{Tan}[e + f*x])/(a + b*\text{Tan}[e + f*x]), x], x] /;$ FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]

Rubi steps

$$\begin{aligned} \int \frac{\tanh(x)}{a + b \tanh(x)} dx &= -\frac{bx}{a^2 - b^2} + \frac{(ia) \int \frac{-ib - ia \tanh(x)}{a + b \tanh(x)} dx}{a^2 - b^2} \\ &= -\frac{bx}{a^2 - b^2} + \frac{a \log(a \cosh(x) + b \sinh(x))}{a^2 - b^2} \end{aligned}$$

Mathematica [A]

time = 0.05, size = 29, normalized size = 0.74

$$\frac{-bx + a \log(a \cosh(x) + b \sinh(x))}{a^2 - b^2}$$

Antiderivative was successfully verified.

`[In] Integrate[Tanh[x]/(a + b*Tanh[x]),x]``[Out] -(b*x) + a*Log[a*Cosh[x] + b*Sinh[x]]/(a^2 - b^2)`**Maple [A]**

time = 0.25, size = 55, normalized size = 1.41

method	result	size
risch	$\frac{x}{a+b} - \frac{2ax}{a^2-b^2} + \frac{a \ln\left(e^{2x} + \frac{a-b}{a+b}\right)}{a^2-b^2}$	54
derivativedivides	$-\frac{\ln(1+\tanh(x))}{2a-2b} + \frac{a \ln(a+b \tanh(x))}{(a+b)(a-b)} - \frac{\ln(\tanh(x)-1)}{2b+2a}$	55
default	$-\frac{\ln(1+\tanh(x))}{2a-2b} + \frac{a \ln(a+b \tanh(x))}{(a+b)(a-b)} - \frac{\ln(\tanh(x)-1)}{2b+2a}$	55

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(tanh(x)/(a+b*tanh(x)),x,method=_RETURNVERBOSE)``[Out] -1/(2*a-2*b)*ln(1+tanh(x))+a/(a+b)/(a-b)*ln(a+b*tanh(x))-1/(2*b+2*a)*ln(tanh(x)-1)`**Maxima [A]**

time = 0.26, size = 40, normalized size = 1.03

$$\frac{a \log(-(a-b)e^{(-2x)} - a - b)}{a^2 - b^2} + \frac{x}{a+b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(tanh(x)/(a+b*tanh(x)),x, algorithm="maxima")``[Out] a*log(-(a - b)*e^(-2*x) - a - b)/(a^2 - b^2) + x/(a + b)`**Fricas [A]**

time = 0.34, size = 43, normalized size = 1.10

$$\frac{(a+b)x - a \log\left(\frac{2(a \cosh(x) + b \sinh(x))}{\cosh(x) - \sinh(x)}\right)}{a^2 - b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)/(a+b*tanh(x)),x, algorithm="fricas")

[Out] $-\frac{((a + b)x - a \log(2*(a \cosh(x) + b \sinh(x)))/(\cosh(x) - \sinh(x)))}{(a^2 - b^2)}$

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 141 vs. $2(29) = 58$.

time = 0.27, size = 141, normalized size = 3.62

$$\begin{cases} \tilde{\infty}x & \text{for } a = 0 \wedge b = 0 \\ \frac{x - \log(\tanh(x) + 1)}{a} & \text{for } b = 0 \\ \frac{x \tanh(x)}{2b \tanh(x) - 2b} - \frac{x}{2b \tanh(x) - 2b} + \frac{1}{2b \tanh(x) - 2b} & \text{for } a = -b \\ \frac{x \tanh(x)}{2b \tanh(x) + 2b} + \frac{x}{2b \tanh(x) + 2b} + \frac{1}{2b \tanh(x) + 2b} & \text{for } a = b \\ \frac{ax}{a^2 - b^2} + \frac{a \log\left(\frac{a}{b} + \tanh(x)\right)}{a^2 - b^2} - \frac{a \log(\tanh(x) + 1)}{a^2 - b^2} - \frac{bx}{a^2 - b^2} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)/(a+b*tanh(x)),x)

[Out] Piecewise((zoo*x, Eq(a, 0) & Eq(b, 0)), ((x - log(tanh(x) + 1))/a, Eq(b, 0)), (x*tanh(x)/(2*b*tanh(x) - 2*b) - x/(2*b*tanh(x) - 2*b) + 1/(2*b*tanh(x) - 2*b), Eq(a, -b)), (x*tanh(x)/(2*b*tanh(x) + 2*b) + x/(2*b*tanh(x) + 2*b) + 1/(2*b*tanh(x) + 2*b), Eq(a, b)), (a*x/(a**2 - b**2) + a*log(a/b + tanh(x))/(a**2 - b**2) - a*log(tanh(x) + 1)/(a**2 - b**2) - b*x/(a**2 - b**2), True))

Giac [A]

time = 0.42, size = 43, normalized size = 1.10

$$\frac{a \log(|ae^{(2x)} + be^{(2x)} + a - b|)}{a^2 - b^2} - \frac{x}{a - b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)/(a+b*tanh(x)),x, algorithm="giac")

[Out] $a \log(\text{abs}(a \cdot e^{(2x)} + b \cdot e^{(2x)} + a - b)) / (a^2 - b^2) - x / (a - b)$

Mupad [B]

time = 1.06, size = 36, normalized size = 0.92

$$\frac{bx - a(x - \ln(\tanh(x) + 1) + \ln(a + b \tanh(x)))}{a^2 - b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(x)/(a + b*tanh(x)),x)

[Out] $-(b \cdot x - a \cdot (x - \log(\tanh(x) + 1) + \log(a + b \cdot \tanh(x)))) / (a^2 - b^2)$

3.138 $\int \frac{1}{a+b \tanh(x)} dx$

Optimal. Leaf size=39

$$\frac{ax}{a^2 - b^2} - \frac{b \log(a \cosh(x) + b \sinh(x))}{a^2 - b^2}$$

[Out] a*x/(a^2-b^2)-b*ln(a*cosh(x)+b*sinh(x))/(a^2-b^2)

Rubi [A]

time = 0.04, antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3565, 3611}

$$\frac{ax}{a^2 - b^2} - \frac{b \log(a \cosh(x) + b \sinh(x))}{a^2 - b^2}$$

Antiderivative was successfully verified.

[In] Int[(a + b*Tanh[x])^(-1), x]

[Out] (a*x)/(a^2 - b^2) - (b*Log[a*Cosh[x] + b*Sinh[x]])/(a^2 - b^2)

Rule 3565

Int[((a_) + (b_)*tan[(c_) + (d_)*(x_)])^(-1), x_Symbol] := Simp[a*(x/(a^2 + b^2)), x] + Dist[b/(a^2 + b^2), Int[(b - a*Tan[c + d*x])/(a + b*Tan[c + d*x]), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0]

Rule 3611

Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{a + b \tanh(x)} dx &= \frac{ax}{a^2 - b^2} - \frac{(ib) \int \frac{-ib - ia \tanh(x)}{a + b \tanh(x)} dx}{a^2 - b^2} \\ &= \frac{ax}{a^2 - b^2} - \frac{b \log(a \cosh(x) + b \sinh(x))}{a^2 - b^2} \end{aligned}$$

Mathematica [A]

time = 0.04, size = 29, normalized size = 0.74

$$\frac{ax - b \log(a \cosh(x) + b \sinh(x))}{a^2 - b^2}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*Tanh[x])^(-1),x]

[Out] (a*x - b*Log[a*Cosh[x] + b*Sinh[x]])/(a^2 - b^2)

Maple [A]

time = 0.14, size = 55, normalized size = 1.41

method	result	size
derivativedivides	$-\frac{\ln(\tanh(x)-1)}{2b+2a} - \frac{b \ln(a+b \tanh(x))}{(a-b)(a+b)} + \frac{\ln(1+\tanh(x))}{2a-2b}$	55
default	$-\frac{\ln(\tanh(x)-1)}{2b+2a} - \frac{b \ln(a+b \tanh(x))}{(a-b)(a+b)} + \frac{\ln(1+\tanh(x))}{2a-2b}$	55
risch	$\frac{x}{a+b} + \frac{2bx}{a^2-b^2} - \frac{b \ln\left(e^{2x} + \frac{a-b}{a+b}\right)}{a^2-b^2}$	55

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a+b*tanh(x)),x,method=_RETURNVERBOSE)

[Out] -1/(2*b+2*a)*ln(tanh(x)-1)-b/(a-b)/(a+b)*ln(a+b*tanh(x))+1/(2*a-2*b)*ln(1+tanh(x))

Maxima [A]

time = 0.27, size = 41, normalized size = 1.05

$$-\frac{b \log\left(-\frac{(a-b)e^{-2x}}{a-b} - a - b\right)}{a^2 - b^2} + \frac{x}{a+b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+b*tanh(x)),x, algorithm="maxima")

[Out] -b*log(-(a - b)*e^(-2*x) - a - b)/(a^2 - b^2) + x/(a + b)

Fricas [A]

time = 0.35, size = 42, normalized size = 1.08

$$\frac{(a+b)x - b \log\left(\frac{2(a \cosh(x) + b \sinh(x))}{\cosh(x) - \sinh(x)}\right)}{a^2 - b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+b*tanh(x)),x, algorithm="fricas")

[Out] ((a + b)*x - b*log(2*(a*cosh(x) + b*sinh(x))/(cosh(x) - sinh(x))))/(a^2 - b^2)

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 146 vs. $2(29) = 58$.

time = 0.28, size = 146, normalized size = 3.74

$$\left\{ \begin{array}{ll} \tilde{\infty}(x - \log(\tanh(x) + 1) + \log(\tanh(x))) & \text{for } a = 0 \wedge b = 0 \\ \frac{x}{a} & \text{for } b = 0 \\ -\frac{x \tanh(x)}{2b \tanh(x) - 2b} + \frac{x}{2b \tanh(x) - 2b} + \frac{1}{2b \tanh(x) - 2b} & \text{for } a = -b \\ \frac{x \tanh(x)}{2b \tanh(x) + 2b} + \frac{x}{2b \tanh(x) + 2b} - \frac{1}{2b \tanh(x) + 2b} & \text{for } a = b \\ \frac{ax}{a^2 - b^2} - \frac{bx}{a^2 - b^2} - \frac{b \log\left(\frac{a}{b} + \tanh(x)\right)}{a^2 - b^2} + \frac{b \log(\tanh(x) + 1)}{a^2 - b^2} & \text{otherwise} \end{array} \right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+b*tanh(x)),x)

[Out] Piecewise((zoo*(x - log(tanh(x) + 1) + log(tanh(x))), Eq(a, 0) & Eq(b, 0)), (x/a, Eq(b, 0)), (-x*tanh(x)/(2*b*tanh(x) - 2*b) + x/(2*b*tanh(x) - 2*b) + 1/(2*b*tanh(x) - 2*b), Eq(a, -b)), (x*tanh(x)/(2*b*tanh(x) + 2*b) + x/(2*b*tanh(x) + 2*b) - 1/(2*b*tanh(x) + 2*b), Eq(a, b)), (a*x/(a**2 - b**2) - b*x/(a**2 - b**2) - b*log(a/b + tanh(x))/(a**2 - b**2) + b*log(tanh(x) + 1)/(a**2 - b**2), True))

Giac [A]

time = 0.41, size = 43, normalized size = 1.10

$$-\frac{b \log(|ae^{(2x)} + be^{(2x)} + a - b|)}{a^2 - b^2} + \frac{x}{a - b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a+b*tanh(x)),x, algorithm="giac")

[Out] -b*log(abs(a*e^(2*x) + b*e^(2*x) + a - b))/(a^2 - b^2) + x/(a - b)

Mupad [B]

time = 0.00, size = 35, normalized size = 0.90

$$\frac{ax - b(x - \ln(\tanh(x) + 1) + \ln(a + b \tanh(x)))}{a^2 - b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a + b*tanh(x)),x)

[Out] (a*x - b*(x - log(tanh(x) + 1) + log(a + b*tanh(x))))/(a^2 - b^2)

3.139 $\int \frac{\coth(x)}{a+b \tanh(x)} dx$

Optimal. Leaf size=51

$$-\frac{bx}{a^2 - b^2} + \frac{\log(\sinh(x))}{a} + \frac{b^2 \log(a \cosh(x) + b \sinh(x))}{a(a^2 - b^2)}$$

[Out] $-b*x/(a^2-b^2)+\ln(\sinh(x))/a+b^2*\ln(a*\cosh(x)+b*\sinh(x))/a/(a^2-b^2)$

Rubi [A]

time = 0.06, antiderivative size = 51, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3652, 3611, 3556}

$$-\frac{bx}{a^2 - b^2} + \frac{b^2 \log(a \cosh(x) + b \sinh(x))}{a(a^2 - b^2)} + \frac{\log(\sinh(x))}{a}$$

Antiderivative was successfully verified.

[In] Int[Coth[x]/(a + b*Tanh[x]),x]

[Out] $-((b*x)/(a^2 - b^2)) + \text{Log}[\text{Sinh}[x]]/a + (b^2*\text{Log}[a*\text{Cosh}[x] + b*\text{Sinh}[x]])/(a*(a^2 - b^2))$

Rule 3556

Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] :> Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]

Rule 3611

Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)]), x_Symbol] :> Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]

Rule 3652

Int[1/(((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)])), x_Symbol] :> Simp[(a*c - b*d)*(x/((a^2 + b^2)*(c^2 + d^2))), x] + (Dist[b^2/((b*c - a*d)*(a^2 + b^2)), Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] - Dist[d^2/((b*c - a*d)*(c^2 + d^2)), Int[(d - c*Tan[e + f*x])/(c + d*Tan[e + f*x]), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]

Rubi steps

$$\int \frac{\coth(x)}{a + b \tanh(x)} dx = -\frac{bx}{a^2 - b^2} + \frac{\int \coth(x) dx}{a} + \frac{(ib^2) \int \frac{-ib-ia \tanh(x)}{a+b \tanh(x)} dx}{a(a^2 - b^2)}$$

$$= -\frac{bx}{a^2 - b^2} + \frac{\log(\sinh(x))}{a} + \frac{b^2 \log(a \cosh(x) + b \sinh(x))}{a(a^2 - b^2)}$$

Mathematica [A]

time = 0.06, size = 46, normalized size = 0.90

$$\frac{(a^2 - b^2) \log(\sinh(x)) + b(-ax + b \log(a \cosh(x) + b \sinh(x)))}{a^3 - ab^2}$$

Antiderivative was successfully verified.

`[In] Integrate[Coth[x]/(a + b*Tanh[x]),x]``[Out] ((a^2 - b^2)*Log[Sinh[x]] + b*(-(a*x) + b*Log[a*Cosh[x] + b*Sinh[x]]))/(a^3 - a*b^2)`**Maple [A]**

time = 0.62, size = 78, normalized size = 1.53

method	result	size
default	$\frac{\ln(\tanh(\frac{x}{2}))}{a} - \frac{\ln(\tanh(\frac{x}{2})+1)}{a-b} + \frac{b^2 \ln(a(\tanh^2(\frac{x}{2}))+2b \tanh(\frac{x}{2})+a)}{(a+b)(a-b)a} - \frac{\ln(\tanh(\frac{x}{2})-1)}{a+b}$	78
risch	$\frac{x}{a+b} - \frac{2x}{a} - \frac{2xb^2}{a(a^2-b^2)} + \frac{\ln(e^{2x}-1)}{a} + \frac{b^2 \ln(e^{2x} + \frac{a-b}{a+b})}{a(a^2-b^2)}$	81

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(coth(x)/(a+b*tanh(x)),x,method=_RETURNVERBOSE)``[Out] 1/a*ln(tanh(1/2*x))-1/(a-b)*ln(tanh(1/2*x)+1)+b^2/(a+b)/(a-b)/a*ln(a*tanh(1/2*x)^2+2*b*tanh(1/2*x)+a)-1/(a+b)*ln(tanh(1/2*x)-1)`**Maxima [A]**

time = 0.27, size = 65, normalized size = 1.27

$$\frac{b^2 \log(-(a-b)e^{(-2x)} - a - b)}{a^3 - ab^2} + \frac{x}{a+b} + \frac{\log(e^{(-x)} + 1)}{a} + \frac{\log(e^{(-x)} - 1)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(coth(x)/(a+b*tanh(x)),x, algorithm="maxima")``[Out] b^2*log(-(a - b)*e^(-2*x) - a - b)/(a^3 - a*b^2) + x/(a + b) + log(e^(-x) + 1)/a + log(e^(-x) - 1)/a`

Fricas [A]

time = 0.38, size = 73, normalized size = 1.43

$$\frac{b^2 \log\left(\frac{2(a \cosh(x) + b \sinh(x))}{\cosh(x) - \sinh(x)}\right) - (a^2 + ab)x + (a^2 - b^2) \log\left(\frac{2 \sinh(x)}{\cosh(x) - \sinh(x)}\right)}{a^3 - ab^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(x)/(a+b*tanh(x)),x, algorithm="fricas")

[Out] (b^2*log(2*(a*cosh(x) + b*sinh(x))/(cosh(x) - sinh(x))) - (a^2 + a*b)*x + (a^2 - b^2)*log(2*sinh(x)/(cosh(x) - sinh(x))))/(a^3 - a*b^2)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\coth(x)}{a + b \tanh(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(x)/(a+b*tanh(x)),x)

[Out] Integral(coth(x)/(a + b*tanh(x)), x)

Giac [A]

time = 0.42, size = 58, normalized size = 1.14

$$\frac{b^2 \log(|ae^{(2x)} + be^{(2x)} + a - b|)}{a^3 - ab^2} - \frac{x}{a - b} + \frac{\log(|e^{(2x)} - 1|)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(x)/(a+b*tanh(x)),x, algorithm="giac")

[Out] b^2*log(abs(a*e^(2*x) + b*e^(2*x) + a - b))/(a^3 - a*b^2) - x/(a - b) + log(abs(e^(2*x) - 1))/a

Mupad [B]

time = 0.41, size = 58, normalized size = 1.14

$$\frac{\ln(e^{2x} - 1)}{a} - \frac{x}{a - b} - \frac{b^2 \ln(a - b + ae^{2x} + be^{2x})}{ab^2 - a^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(coth(x)/(a + b*tanh(x)),x)

[Out] log(exp(2*x) - 1)/a - x/(a - b) - (b^2*log(a - b + a*exp(2*x) + b*exp(2*x)))/(a*b^2 - a^3)

3.140 $\int \frac{\coth^2(x)}{a+b \tanh(x)} dx$

Optimal. Leaf size=60

$$\frac{ax}{a^2 - b^2} - \frac{\coth(x)}{a} - \frac{b \log(\sinh(x))}{a^2} - \frac{b^3 \log(a \cosh(x) + b \sinh(x))}{a^2 (a^2 - b^2)}$$

[Out] a*x/(a^2-b^2)-coth(x)/a-b*ln(sinh(x))/a^2-b^3*ln(a*cosh(x)+b*sinh(x))/a^2/(a^2-b^2)

Rubi [A]

time = 0.13, antiderivative size = 60, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {3650, 3732, 3611, 3556}

$$\frac{ax}{a^2 - b^2} - \frac{b^3 \log(a \cosh(x) + b \sinh(x))}{a^2 (a^2 - b^2)} - \frac{b \log(\sinh(x))}{a^2} - \frac{\coth(x)}{a}$$

Antiderivative was successfully verified.

[In] Int[Coth[x]^2/(a + b*Tanh[x]),x]

[Out] (a*x)/(a^2 - b^2) - Coth[x]/a - (b*Log[Sinh[x]])/a^2 - (b^3*Log[a*Cosh[x] + b*Sinh[x]])/(a^2*(a^2 - b^2))

Rule 3556

Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]

Rule 3611

Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]

Rule 3650

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b^2*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(a^2 + b^2)*(b*c - a*d))), x] + Dist[1/((m + 1)*(a^2 + b^2)*(b*c - a*d)), Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2) - b*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b^2*d*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && IntegerQ[2*m] && LtQ[m, -1] && (LtQ[n, 0] || Intege

rQ[m]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))

Rule 3732

Int[((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2)/(((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])), x_Symbol] := Simp[(a*(A*c - c*C + B*d) + b*(B*c - A*d + C*d))*(x/((a^2 + b^2)*(c^2 + d^2))), x] + (Dist[(A*b^2 - a*b*B + a^2*C)/((b*c - a*d)*(a^2 + b^2)), Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] - Dist[(c^2*C - B*c*d + A*d^2)/((b*c - a*d)*(c^2 + d^2)), Int[(d - c*Tan[e + f*x])/(c + d*Tan[e + f*x]), x], x]) /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]

Rubi steps

$$\begin{aligned} \int \frac{\coth^2(x)}{a + b \tanh(x)} dx &= -\frac{\coth(x)}{a} - \frac{i \int \frac{\coth(x)(-ib+ia \tanh(x)+ib \tanh^2(x))}{a+b \tanh(x)} dx}{a} \\ &= \frac{ax}{a^2 - b^2} - \frac{\coth(x)}{a} - \frac{b \int \coth(x) dx}{a^2} - \frac{(ib^3) \int \frac{-ib-ia \tanh(x)}{a+b \tanh(x)} dx}{a^2(a^2 - b^2)} \\ &= \frac{ax}{a^2 - b^2} - \frac{\coth(x)}{a} - \frac{b \log(\sinh(x))}{a^2} - \frac{b^3 \log(a \cosh(x) + b \sinh(x))}{a^2(a^2 - b^2)} \end{aligned}$$

Mathematica [A]

time = 0.10, size = 64, normalized size = 1.07

$$\frac{a^3 x + (-a^3 + ab^2) \coth(x) + (-a^2 b + b^3) \log(\sinh(x)) - b^3 \log(a \cosh(x) + b \sinh(x))}{a^4 - a^2 b^2}$$

Antiderivative was successfully verified.

[In] Integrate[Coth[x]^2/(a + b*Tanh[x]), x]

[Out] (a^3*x + (-a^3 + a*b^2)*Coth[x] + (-a^2*b) + b^3)*Log[Sinh[x]] - b^3*Log[a*Cosh[x] + b*Sinh[x]]/(a^4 - a^2*b^2)

Maple [A]

time = 0.66, size = 100, normalized size = 1.67

method	result
risch	$\frac{x}{a+b} + \frac{2bx}{a^2} + \frac{2xb^3}{a^2(a^2-b^2)} - \frac{2}{a(e^{2x}-1)} - \frac{b \ln(e^{2x}-1)}{a^2} - \frac{b^3 \ln\left(e^{2x} + \frac{a-b}{a+b}\right)}{a^2(a^2-b^2)}$

default	$-\frac{\tanh(\frac{x}{2})}{2a} - \frac{1}{2a \tanh(\frac{x}{2})} - \frac{b \ln(\tanh(\frac{x}{2}))}{a^2} - \frac{\ln(\tanh(\frac{x}{2})-1)}{a+b} - \frac{b^3 \ln(a(\tanh^2(\frac{x}{2}))+2b \tanh(\frac{x}{2})+a)}{a^2(a+b)(a-b)} + \frac{\ln(\tanh(\frac{x}{2})+1)}{a-b}$	1
---------	---	---

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(coth(x)^2/(a+b*tanh(x)),x,method=_RETURNVERBOSE)`

[Out]
$$-1/2/a*\tanh(1/2*x)-1/2/a/\tanh(1/2*x)-b/a^2*\ln(\tanh(1/2*x))-1/(a+b)*\ln(\tanh(1/2*x)-1)-1/a^2*b^3/(a+b)/(a-b)*\ln(a*\tanh(1/2*x)^2+2*b*\tanh(1/2*x)+a)+1/(a-b)*\ln(\tanh(1/2*x)+1)$$

Maxima [A]

time = 0.27, size = 86, normalized size = 1.43

$$-\frac{b^3 \log(-(a-b)e^{(-2x)} - a - b)}{a^4 - a^2 b^2} + \frac{x}{a+b} - \frac{b \log(e^{(-x)} + 1)}{a^2} - \frac{b \log(e^{(-x)} - 1)}{a^2} + \frac{2}{ae^{(-2x)} - a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(coth(x)^2/(a+b*tanh(x)),x, algorithm="maxima")`

[Out]
$$-b^3*\log(-(a-b)*e^{(-2*x)} - a - b)/(a^4 - a^2*b^2) + x/(a+b) - b*\log(e^{(-x)} - x) + 1/a^2 - b*\log(e^{(-x)} - 1)/a^2 + 2/(a*e^{(-2*x)} - a)$$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 271 vs. 2(60) = 120.

time = 0.42, size = 271, normalized size = 4.52

$$\frac{(a^3 + a^2 b)x \cosh(x)^2 + 2(a^3 + a^2 b)x \cosh(x) \sinh(x) + (a^3 + a^2 b)x \sinh(x)^2 - 2a^3 + 2ab^2 - (a^3 + a^2 b)x - (b^3 \cosh(x)^2 + 2b^3 \cosh(x) \sinh(x) + b^3 \sinh(x)^2 - b^3) \log\left(\frac{2 \cosh(x) + \sinh(x)}{\cosh(x) - \sinh(x)}\right) + (a^2 b - b^3 - (a^2 b - b^3) \cosh(x)^2 - 2(a^2 b - b^3) \cosh(x) \sinh(x) - (a^2 b - b^3) \sinh(x)^2) \log\left(\frac{2 \sinh(x)}{\cosh(x) - \sinh(x)}\right)}{a^4 - a^2 b^2 - (a^4 - a^2 b^2) \cosh(x)^2 - 2(a^4 - a^2 b^2) \cosh(x) \sinh(x) - (a^4 - a^2 b^2) \sinh(x)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(coth(x)^2/(a+b*tanh(x)),x, algorithm="fricas")`

[Out]
$$-((a^3 + a^2*b)*x*\cosh(x)^2 + 2*(a^3 + a^2*b)*x*\cosh(x)*\sinh(x) + (a^3 + a^2*b)*x*\sinh(x)^2 - 2*a^3 + 2*a*b^2 - (a^3 + a^2*b)*x - (b^3*\cosh(x)^2 + 2*b^3*\cosh(x)*\sinh(x) + b^3*\sinh(x)^2 - b^3)*\log(2*(a*\cosh(x) + b*\sinh(x))/(\cosh(x) - \sinh(x))) + (a^2*b - b^3 - (a^2*b - b^3)*\cosh(x)^2 - 2*(a^2*b - b^3)*\cosh(x)*\sinh(x) - (a^2*b - b^3)*\sinh(x)^2)*\log(2*\sinh(x)/(\cosh(x) - \sinh(x))))/(a^4 - a^2*b^2 - (a^4 - a^2*b^2)*\cosh(x)^2 - 2*(a^4 - a^2*b^2)*\cosh(x)*\sinh(x) - (a^4 - a^2*b^2)*\sinh(x)^2)$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\coth^2(x)}{a + b \tanh(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(x)**2/(a+b*tanh(x)),x)

[Out] Integral(coth(x)**2/(a + b*tanh(x)), x)

Giac [A]

time = 0.41, size = 75, normalized size = 1.25

$$-\frac{b^3 \log(|ae^{(2x)} + be^{(2x)} + a - b|)}{a^4 - a^2b^2} + \frac{x}{a - b} - \frac{b \log(|e^{(2x)} - 1|)}{a^2} - \frac{2}{a(e^{(2x)} - 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(x)^2/(a+b*tanh(x)),x, algorithm="giac")

[Out] $-b^3 \log(\text{abs}(a \cdot e^{(2x)} + b \cdot e^{(2x)} + a - b)) / (a^4 - a^2 \cdot b^2) + x / (a - b) - b \cdot \log(\text{abs}(e^{(2x)} - 1)) / a^2 - 2 / (a \cdot (e^{(2x)} - 1))$

Mupad [B]

time = 1.35, size = 73, normalized size = 1.22

$$\frac{x}{a - b} - \frac{2}{a(e^{2x} - 1)} - \frac{b^3 \ln(a - b + a e^{2x} + b e^{2x})}{a^4 - a^2 b^2} - \frac{b \ln(e^{2x} - 1)}{a^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(coth(x)^2/(a + b*tanh(x)),x)

[Out] $x / (a - b) - 2 / (a \cdot (\exp(2x) - 1)) - (b^3 \cdot \log(a - b + a \cdot \exp(2x) + b \cdot \exp(2x))) / (a^4 - a^2 \cdot b^2) - (b \cdot \log(\exp(2x) - 1)) / a^2$

3.141 $\int \frac{\coth^3(x)}{a+b \tanh(x)} dx$

Optimal. Leaf size=76

$$-\frac{bx}{a^2-b^2} + \frac{b \coth(x)}{a^2} - \frac{\coth^2(x)}{2a} + \frac{(a^2+b^2) \log(\sinh(x))}{a^3} + \frac{b^4 \log(a \cosh(x) + b \sinh(x))}{a^3(a^2-b^2)}$$

[Out] $-b*x/(a^2-b^2)+b*\coth(x)/a^2-1/2*\coth(x)^2/a+(a^2+b^2)*\ln(\sinh(x))/a^3+b^4*\ln(a*\cosh(x)+b*\sinh(x))/a^3/(a^2-b^2)$

Rubi [A]

time = 0.21, antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {3650, 3730, 3733, 3611, 3556}

$$-\frac{bx}{a^2-b^2} + \frac{b \coth(x)}{a^2} + \frac{(a^2+b^2) \log(\sinh(x))}{a^3} + \frac{b^4 \log(a \cosh(x) + b \sinh(x))}{a^3(a^2-b^2)} - \frac{\coth^2(x)}{2a}$$

Antiderivative was successfully verified.

[In] `Int[Coth[x]^3/(a + b*Tanh[x]),x]`

[Out] $-\left(\frac{b*x}{a^2-b^2}\right) + \frac{b*\text{Coth}[x]}{a^2} - \frac{\text{Coth}[x]^2}{2*a} + \frac{(a^2+b^2)*\text{Log}[\text{Sinh}[x]]}{a^3} + \frac{b^4*\text{Log}[a*\text{Cosh}[x] + b*\text{Sinh}[x]]}{a^3*(a^2-b^2)}$

Rule 3556

`Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

Rule 3611

`Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`

Rule 3650

`Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b^2*(a + b*Tan[e + f*x])^(m+1)*((c + d*Tan[e + f*x])^(n+1)/(f*(m+1)*(a^2 + b^2)*(b*c - a*d))), x] + Dist[1/((m+1)*(a^2 + b^2)*(b*c - a*d)), Int[(a + b*Tan[e + f*x])^(m+1)*(c + d*Tan[e + f*x])^n*Simp[a*(b*c - a*d)*(m+1) - b^2*d*(m+n+2) - b*(b*c - a*d)*(m+1)*Tan[e + f*x] - b^2*d*(m+n+2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && IntegerQ[2*m] && LtQ[m, -1] && (LtQ[n, 0] || Intege`

rQ[m]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))

Rule 3730

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] :> Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Dist[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)), Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))

Rule 3733

Int[((A_.) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2/(((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])), x_Symbol] :> Simp[(a*(A*c - c*C) - b*(A*d - C*d))*(x/((a^2 + b^2)*(c^2 + d^2))), x] + (Dist[(A*b^2 + a^2*C)/((b*c - a*d)*(a^2 + b^2)), Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] - Dist[(c^2*C + A*d^2)/((b*c - a*d)*(c^2 + d^2)), Int[(d - c*Tan[e + f*x])/(c + d*Tan[e + f*x]), x], x]) /; FreeQ[{a, b, c, d, e, f, A, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]

Rubi steps

$$\begin{aligned} \int \frac{\coth^3(x)}{a + b \tanh(x)} dx &= -\frac{\coth^2(x)}{2a} - \frac{i \int \frac{\coth^2(x)(-2ib+2ia \tanh(x)+2ib \tanh^2(x))}{a+b \tanh(x)} dx}{2a} \\ &= \frac{b \coth(x)}{a^2} - \frac{\coth^2(x)}{2a} - \frac{\int \frac{\coth(x)(-2(a^2+b^2)+2b^2 \tanh^2(x))}{a+b \tanh(x)} dx}{2a^2} \\ &= -\frac{bx}{a^2 - b^2} + \frac{b \coth(x)}{a^2} - \frac{\coth^2(x)}{2a} + \frac{(ib^4) \int \frac{-ib-ia \tanh(x)}{a+b \tanh(x)} dx}{a^3(a^2 - b^2)} + \frac{(a^2 + b^2) \int \coth(x) dx}{a^3} \\ &= -\frac{bx}{a^2 - b^2} + \frac{b \coth(x)}{a^2} - \frac{\coth^2(x)}{2a} + \frac{(a^2 + b^2) \log(\sinh(x))}{a^3} + \frac{b^4 \log(a \cosh(x) + b \sinh(x))}{a^3(a^2 - b^2)} \end{aligned}$$

Mathematica [A]

time = 0.14, size = 91, normalized size = 1.20

$$\frac{-2a^3bx + 2ab(a^2 - b^2) \coth(x) + (-a^4 + a^2b^2) \operatorname{csch}^2(x) + 2a^4 \log(\sinh(x)) - 2b^4 \log(\sinh(x)) + 2b^4 \log(a \cosh(x) + b \sinh(x))}{2a^3(a - b)(a + b)}$$

Antiderivative was successfully verified.

[In] Integrate[Coth[x]^3/(a + b*Tanh[x]),x]

[Out] $(-2a^3bx + 2ab(a^2 - b^2)\text{Coth}[x] + (-a^4 + a^2b^2)\text{Csch}[x]^2 + 2a^4\text{Log}[\text{Sinh}[x]] - 2b^4\text{Log}[\text{Sinh}[x]] + 2b^4\text{Log}[a\text{Cosh}[x] + b\text{Sinh}[x]])/(2a^3(a - b)(a + b))$

Maple [A]

time = 0.64, size = 135, normalized size = 1.78

method	result
risch	$\frac{x}{a+b} - \frac{2x}{a} - \frac{2xb^2}{a^3} - \frac{2xb^4}{a^3(a^2-b^2)} - \frac{2(e^{2x}a-b e^{2x}+b)}{(e^{2x}-1)^2 a^2} + \frac{\ln(e^{2x}-1)}{a} + \frac{\ln(e^{2x}-1)b^2}{a^3} + \frac{b^4 \ln(e^{2x} + \frac{a-b}{a+b})}{a^3(a^2-b^2)}$
default	$-\frac{\frac{a(\tanh^2(\frac{x}{2}))}{2} - 2b \tanh(\frac{x}{2})}{4a^2} + \frac{b^4 \ln(a(\tanh^2(\frac{x}{2})) + 2b \tanh(\frac{x}{2}) + a)}{a^3(a+b)(a-b)} - \frac{\ln(\tanh(\frac{x}{2})-1)}{a+b} - \frac{\ln(\tanh(\frac{x}{2})+1)}{a-b} - \frac{1}{8a \tanh(\frac{x}{2})^2} + \dots$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(coth(x)^3/(a+b*tanh(x)),x,method=_RETURNVERBOSE)

[Out] $-1/4/a^2*(1/2*a*tanh(1/2*x)^2-2*b*tanh(1/2*x))+1/a^3*b^4/(a+b)/(a-b)*ln(a*tanh(1/2*x)^2+2*b*tanh(1/2*x)+a)-1/(a+b)*ln(tanh(1/2*x)-1)-1/(a-b)*ln(tanh(1/2*x)+1)-1/8/a/tanh(1/2*x)^2+1/4/a^3*(4*a^2+4*b^2)*ln(tanh(1/2*x))+1/2*b/a^2/tanh(1/2*x)$

Maxima [A]

time = 0.28, size = 121, normalized size = 1.59

$\frac{b^4 \log(-(a-b)e^{(-2x)} - a - b)}{a^5 - a^3 b^2} + \frac{2((a+b)e^{(-2x)} - b)}{2a^2 e^{(-2x)} - a^2 e^{(-4x)} - a^2} + \frac{x}{a+b} + \frac{(a^2 + b^2) \log(e^{(-x)} + 1)}{a^3} + \frac{(a^2 + b^2) \log(e^{(-x)} - 1)}{a^3}$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(x)^3/(a+b*tanh(x)),x, algorithm="maxima")

[Out] $b^4 \log(-(a-b)e^{(-2x)} - a - b)/(a^5 - a^3 b^2) + 2*((a+b)e^{(-2x)} - b)/(2a^2 e^{(-2x)} - a^2 e^{(-4x)} - a^2) + x/(a+b) + (a^2 + b^2) \log(e^{(-x)} - 1)/a^3 + (a^2 + b^2) \log(e^{(-x)} + 1)/a^3$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 641 vs. $2(74) = 148$.

time = 0.43, size = 641, normalized size = 8.43

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(x)^3/(a+b*tanh(x)),x, algorithm="fricas")

```
[Out] -((a^4 + a^3*b)*x*cosh(x)^4 + 4*(a^4 + a^3*b)*x*cosh(x)*sinh(x)^3 + (a^4 + a^3*b)*x*sinh(x)^4 + 2*a^3*b - 2*a*b^3 + 2*(a^4 - a^3*b - a^2*b^2 + a*b^3 - (a^4 + a^3*b)*x)*cosh(x)^2 + 2*(a^4 - a^3*b - a^2*b^2 + a*b^3 + 3*(a^4 + a^3*b)*x*cosh(x)^2 - (a^4 + a^3*b)*x)*sinh(x)^2 + (a^4 + a^3*b)*x - (b^4*cosh(x)^4 + 4*b^4*cosh(x)*sinh(x)^3 + b^4*sinh(x)^4 - 2*b^4*cosh(x)^2 + b^4 + 2*(3*b^4*cosh(x)^2 - b^4)*sinh(x)^2 + 4*(b^4*cosh(x)^3 - b^4*cosh(x))*sinh(x))*log(2*(a*cosh(x) + b*sinh(x))/(cosh(x) - sinh(x))) - ((a^4 - b^4)*cosh(x)^4 + 4*(a^4 - b^4)*cosh(x)*sinh(x)^3 + (a^4 - b^4)*sinh(x)^4 + a^4 - b^4 - 2*(a^4 - b^4)*cosh(x)^2 - 2*(a^4 - b^4 - 3*(a^4 - b^4)*cosh(x)^2)*sinh(x)^2 + 4*((a^4 - b^4)*cosh(x)^3 - (a^4 - b^4)*cosh(x))*sinh(x))*log(2*sinh(x)/(cosh(x) - sinh(x))) + 4*((a^4 + a^3*b)*x*cosh(x)^3 + (a^4 - a^3*b - a^2*b^2 + a*b^3 - (a^4 + a^3*b)*x)*cosh(x))*sinh(x))/(a^5 - a^3*b^2 + (a^5 - a^3*b^2)*cosh(x)^4 + 4*(a^5 - a^3*b^2)*cosh(x)*sinh(x)^3 + (a^5 - a^3*b^2)*sinh(x)^4 - 2*(a^5 - a^3*b^2)*cosh(x)^2 - 2*(a^5 - a^3*b^2 - 3*(a^5 - a^3*b^2)*cosh(x)^2)*sinh(x)^2 + 4*((a^5 - a^3*b^2)*cosh(x)^3 - (a^5 - a^3*b^2)*cosh(x))*sinh(x))
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\coth^3(x)}{a + b \tanh(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(coth(x)**3/(a+b*tanh(x)),x)
```

```
[Out] Integral(coth(x)**3/(a + b*tanh(x)), x)
```

Giac [A]

time = 0.41, size = 97, normalized size = 1.28

$$\frac{b^4 \log(|ae^{(2x)} + be^{(2x)} + a - b|)}{a^5 - a^3b^2} - \frac{x}{a - b} + \frac{(a^2 + b^2) \log(|e^{(2x)} - 1|)}{a^3} - \frac{2(ab + (a^2 - ab)e^{(2x)})}{a^3(e^{(2x)} - 1)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(coth(x)^3/(a+b*tanh(x)),x, algorithm="giac")
```

```
[Out] b^4*log(abs(a*e^(2*x) + b*e^(2*x) + a - b))/(a^5 - a^3*b^2) - x/(a - b) + (a^2 + b^2)*log(abs(e^(2*x) - 1))/a^3 - 2*(a*b + (a^2 - a*b)*e^(2*x))/(a^3*(e^(2*x) - 1)^2)
```

Mupad [B]

time = 1.42, size = 111, normalized size = 1.46

$$\frac{\ln(e^{2x} - 1)(a^2 + b^2)}{a^3} - \frac{x}{a - b} - \frac{2}{a(e^{4x} - 2e^{2x} + 1)} + \frac{b^4 \ln(a - b + ae^{2x} + be^{2x})}{a^5 - a^3b^2} - \frac{2(a^2 - b^2)}{a^2(a + b)(e^{2x} - 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(coth(x)^3/(a + b*tanh(x)),x)
```

```
[Out] (log(exp(2*x) - 1)*(a^2 + b^2))/a^3 - x/(a - b) - 2/(a*(exp(4*x) - 2*exp(2*x) + 1)) + (b^4*log(a - b + a*exp(2*x) + b*exp(2*x)))/(a^5 - a^3*b^2) - (2*(a^2 - b^2))/(a^2*(a + b)*(exp(2*x) - 1))
```

3.142 $\int \frac{\coth^4(x)}{a+b \tanh(x)} dx$

Optimal. Leaf size=97

$$\frac{ax}{a^2 - b^2} - \frac{(a^2 + b^2) \coth(x)}{a^3} + \frac{b \coth^2(x)}{2a^2} - \frac{\coth^3(x)}{3a} - \frac{b(a^2 + b^2) \log(\sinh(x))}{a^4} - \frac{b^5 \log(a \cosh(x) + b \sinh(x))}{a^4(a^2 - b^2)}$$

[Out] $a*x/(a^2-b^2)-(a^2+b^2)*\coth(x)/a^3+1/2*b*\coth(x)^2/a^2-1/3*\coth(x)^3/a-b*(a^2+b^2)*\ln(\sinh(x))/a^4-b^5*\ln(a*\cosh(x)+b*\sinh(x))/a^4/(a^2-b^2)$

Rubi [A]

time = 0.34, antiderivative size = 97, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.462$, Rules used = {3650, 3730, 3731, 3732, 3611, 3556}

$$\frac{ax}{a^2 - b^2} + \frac{b \coth^2(x)}{2a^2} - \frac{b(a^2 + b^2) \log(\sinh(x))}{a^4} - \frac{b^5 \log(a \cosh(x) + b \sinh(x))}{a^4(a^2 - b^2)} - \frac{(a^2 + b^2) \coth(x)}{a^3} - \frac{\coth^3(x)}{3a}$$

Antiderivative was successfully verified.

[In] Int[Coth[x]^4/(a + b*Tanh[x]), x]

[Out] $(a*x)/(a^2 - b^2) - ((a^2 + b^2)*\text{Coth}[x])/a^3 + (b*\text{Coth}[x]^2)/(2*a^2) - \text{Coth}[x]^3/(3*a) - (b*(a^2 + b^2)*\text{Log}[\text{Sinh}[x]])/a^4 - (b^5*\text{Log}[a*\text{Cosh}[x] + b*\text{Sinh}[x]])/(a^4*(a^2 - b^2))$

Rule 3556

Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]

Rule 3611

Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sinh[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]

Rule 3650

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Simp[b^2*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(a^2 + b^2)*(b*c - a*d))), x] + Dist[1/((m + 1)*(a^2 + b^2)*(b*c - a*d)), Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2) - b*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b^2*d*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0]

&& NeQ[c^2 + d^2, 0] && IntegerQ[2*m] && LtQ[m, -1] && (LtQ[n, 0] || IntegerQ[m]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))

Rule 3730

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Dist[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)), Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))

Rule 3731

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 + a^2*C)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Dist[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)), Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - a*C*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - b*C)*Tan[e + f*x] - d*(A*b^2 + a^2*C)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))

Rule 3732

Int[((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2)/(((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])), x_Symbol] := Simp[(a*(A*c - c*C + B*d) + b*(B*c - A*d + C*d))*(x/(a^2 + b^2)*(c^2 + d^2)), x] + (Dist[(A*b^2 - a*b*B + a^2*C)/((b*c - a*d)*(a^2 + b^2)), Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] - Dist[(c^2*C - B*c*d + A*d^2)/((b*c - a*d)*(c^2 + d^2)), Int[(d - c*Tan[e + f*x])/(c + d*Tan[e + f*x]), x], x]) /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]

Rubi steps

$$\begin{aligned}
\int \frac{\coth^4(x)}{a+b \tanh(x)} dx &= -\frac{\coth^3(x)}{3a} - \frac{i \int \frac{\coth^3(x)(-3ib+3ia \tanh(x)+3ib \tanh^2(x))}{a+b \tanh(x)} dx}{3a} \\
&= \frac{b \coth^2(x)}{2a^2} - \frac{\coth^3(x)}{3a} - \frac{\int \frac{\coth^2(x)(-6(a^2+b^2)+6b^2 \tanh^2(x))}{a+b \tanh(x)} dx}{6a^2} \\
&= -\frac{(a^2+b^2) \coth(x)}{a^3} + \frac{b \coth^2(x)}{2a^2} - \frac{\coth^3(x)}{3a} + \frac{i \int \frac{\coth(x)(6ib(a^2+b^2)-6ia^3 \tanh(x)-6ib(a^2+b^2) \tanh^2(x))}{a+b \tanh(x)} dx}{6a^3} \\
&= \frac{ax}{a^2-b^2} - \frac{(a^2+b^2) \coth(x)}{a^3} + \frac{b \coth^2(x)}{2a^2} - \frac{\coth^3(x)}{3a} - \frac{(ib^5) \int \frac{-ib-ia \tanh(x)}{a+b \tanh(x)} dx}{a^4(a^2-b^2)} - (b^5) \\
&= \frac{ax}{a^2-b^2} - \frac{(a^2+b^2) \coth(x)}{a^3} + \frac{b \coth^2(x)}{2a^2} - \frac{\coth^3(x)}{3a} - \frac{b(a^2+b^2) \log(\sinh(x))}{a^4} - \frac{b^5}{a^4}
\end{aligned}$$

Mathematica [A]

time = 0.23, size = 108, normalized size = 1.11

$$\frac{6a^5x + 3a^2b(a^2 - b^2) \operatorname{csch}^2(x) - 2a(a^2 - b^2) \coth(x) (4a^2 + 3b^2 + a^2 \operatorname{csch}^2(x)) + 6(-a^4b + b^5) \log(\sinh(x)) - 6b^5 \log(a \cosh(x) + b \sinh(x))}{6a^4(a-b)(a+b)}$$

Antiderivative was successfully verified.

`[In] Integrate[Coth[x]^4/(a + b*Tanh[x]), x]`

```
[Out] (6*a^5*x + 3*a^2*b*(a^2 - b^2)*Csch[x]^2 - 2*a*(a^2 - b^2)*Coth[x]*(4*a^2 + 3*b^2 + a^2*Csch[x]^2) + 6*(-(a^4*b) + b^5)*Log[Sinh[x]] - 6*b^5*Log[a*Cosh[x] + b*Sinh[x]])/(6*a^4*(a - b)*(a + b))
```

Maple [A]

time = 0.68, size = 177, normalized size = 1.82

method	result
default	$-\frac{\left(\frac{\tanh^3\left(\frac{x}{2}\right)}{3}\right)a^2 - ab\left(\tanh^2\left(\frac{x}{2}\right)\right) + 5a^2 \tanh\left(\frac{x}{2}\right) + 4b^2 \tanh\left(\frac{x}{2}\right)}{8a^3} + \frac{\ln\left(\tanh\left(\frac{x}{2}\right) + 1\right)}{a-b} - \frac{b^5 \ln\left(a\left(\tanh^2\left(\frac{x}{2}\right)\right) + 2b \tanh\left(\frac{x}{2}\right) + a\right)}{a^4(a+b)(a-b)} - \frac{\ln\left(\tanh\left(\frac{x}{2}\right) - 1\right)}{a-b}$
risch	$\frac{x}{a+b} + \frac{2bx}{a^2} + \frac{2xb^3}{a^4} + \frac{2xb^5}{a^4(a^2-b^2)} - \frac{2(6a^2e^{4x} - 3abe^{4x} + 3b^2e^{4x} - 6a^2e^{2x} + 3abe^{2x} - 6b^2e^{2x} + 4a^2 + 3b^2)}{3a^3(e^{2x}-1)^3} - \frac{b \ln(e^{2x}-1)}{a^2} - \frac{b^3 \ln(e^{2x}-1)}{a^2}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(coth(x)^4/(a+b*tanh(x)), x, method=_RETURNVERBOSE)`

```
[Out] -1/8/a^3*(1/3*tanh(1/2*x)^3*a^2-a*b*tanh(1/2*x)^2+5*a^2*tanh(1/2*x)+4*b^2*tanh(1/2*x))+1/(a-b)*ln(tanh(1/2*x)+1)-1/a^4*b^5/(a+b)/(a-b)*ln(a*tanh(1/2*x)^2+2*b*tanh(1/2*x)+a)-1/(a+b)*ln(tanh(1/2*x)-1)-1/24/a/tanh(1/2*x)^3-1/8*(
```

$5a^2+4b^2)/a^3/\tanh(1/2*x)+1/8*b/a^2/\tanh(1/2*x)^2-1/a^4*b*(a^2+b^2)*\ln(\tanh(1/2*x))$

Maxima [A]

time = 0.28, size = 173, normalized size = 1.78

$$-\frac{b^5 \log(-(a-b)e^{(-2x)}-a-b)}{a^6-a^4b^2} + \frac{2(4a^2+3b^2-3(2a^2+ab+2b^2)e^{(-2x)}+3(2a^2+ab+b^2)e^{(-4x)})}{3(3a^3e^{(-2x)}-3a^3e^{(-4x)}+a^3e^{(-6x)}-a^3)} + \frac{x}{a+b} - \frac{(a^2b+b^3)\log(e^{-x}+1)}{a^4} - \frac{(a^2b+b^3)\log(e^{-x}-1)}{a^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(x)^4/(a+b*tanh(x)),x, algorithm="maxima")

[Out] $-b^5*\log(-(a-b)*e^{(-2*x)}-a-b)/(a^6-a^4*b^2)+2/3*(4*a^2+3*b^2-3*(2*a^2+a*b+2*b^2)*e^{(-2*x)}+3*(2*a^2+a*b+b^2)*e^{(-4*x)})/(3*a^3*e^{(-2*x)}-3*a^3*e^{(-4*x)}+a^3*e^{(-6*x)}-a^3)+x/(a+b)-(a^2*b+b^3)*\log(e^{(-x)}+1)/a^4-(a^2*b+b^3)*\log(e^{(-x)}-1)/a^4$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 1299 vs. 2(93) = 186.

time = 0.42, size = 1299, normalized size = 13.39

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(x)^4/(a+b*tanh(x)),x, algorithm="fricas")

[Out] $1/3*(3*(a^5+a^4*b)*x*\cosh(x)^6+18*(a^5+a^4*b)*x*\cosh(x)*\sinh(x)^5+3*(a^5+a^4*b)*x*\sinh(x)^6-8*a^5+2*a^3*b^2+6*a*b^4-3*(4*a^5-2*a^4*b-2*a^3*b^2+2*a^2*b^3-2*a*b^4+3*(a^5+a^4*b)*x)*\cosh(x)^4-3*(4*a^5-2*a^4*b-2*a^3*b^2+2*a^2*b^3-2*a*b^4-15*(a^5+a^4*b)*x*\cosh(x))^2+3*(a^5+a^4*b)*x*\sinh(x)^4+12*(5*(a^5+a^4*b)*x*\cosh(x)^3-(4*a^5-2*a^4*b-2*a^3*b^2+2*a^2*b^3-2*a*b^4+3*(a^5+a^4*b)*x)*\cosh(x))*\sinh(x)^3+3*(4*a^5-2*a^4*b+2*a^2*b^3-4*a*b^4+3*(a^5+a^4*b)*x)*\cosh(x)^2+3*(15*(a^5+a^4*b)*x*\cosh(x)^4+4*a^5-2*a^4*b+2*a^2*b^3-4*a*b^4-6*(4*a^5-2*a^4*b-2*a^3*b^2+2*a^2*b^3-2*a*b^4+3*(a^5+a^4*b)*x)*\cosh(x)^2+3*(a^5+a^4*b)*x*\sinh(x)^2-3*(a^5+a^4*b)*x-3*(b^5*\cosh(x)^6+6*b^5*\cosh(x)*\sinh(x)^5+b^5*\sinh(x)^6-3*b^5*\cosh(x)^4+3*b^5*\cosh(x)^2-b^5+3*(5*b^5*\cosh(x)^2-b^5)*\sinh(x)^4+4*(5*b^5*\cosh(x)^3-3*b^5*\cosh(x))*\sinh(x)^3+3*(5*b^5*\cosh(x)^4-6*b^5*\cosh(x)^2+b^5)*\sinh(x)^2+6*(b^5*\cosh(x)^5-2*b^5*\cosh(x)^3+b^5*\cosh(x))*\sinh(x))*\log(2*(a*\cosh(x)+b*\sinh(x))/(\cosh(x)-\sinh(x)))-3*((a^4*b-b^5)*\cosh(x)^6+6*(a^4*b-b^5)*\cosh(x)*\sinh(x)^5+(a^4*b-b^5)*\sinh(x)^6-a^4*b+b^5-3*(a^4*b-b^5)*\cosh(x)^4-3*(a^4*b-b^5-5*(a^4*b-b^5)*\cosh(x)^2)*\sinh(x)^4+4*(5*(a^4*b-b^5)*\cosh(x)^3-3*(a^4*b-b^5)*\cosh(x))*\sinh(x)^3+3*(a^4*b-b^5)*\cosh(x)^2+3*(a^4*b-b^5+5*(a^4*b-b^5)*\cosh(x)^4-6*(a^4*b-b^5)*\cosh(x)^2)*\sinh(x)^2+6*((a^4*b-b^5)*\cosh(x)^5-2*(a^4*b-b^5)*\cosh(x)^3+(a^4*b-b^5)*\cosh(x))*\sinh(x))*\log(2*\sinh(x))$

$$\begin{aligned} & x)/(\cosh(x) - \sinh(x))) + 6*(3*(a^5 + a^4*b)*x*\cosh(x)^5 - 2*(4*a^5 - 2*a^4 \\ & *b - 2*a^3*b^2 + 2*a^2*b^3 - 2*a*b^4 + 3*(a^5 + a^4*b)*x)*\cosh(x)^3 + (4*a^5 \\ & - 2*a^4*b + 2*a^2*b^3 - 4*a*b^4 + 3*(a^5 + a^4*b)*x)*\cosh(x))*\sinh(x))/((\\ & a^6 - a^4*b^2)*\cosh(x)^6 + 6*(a^6 - a^4*b^2)*\cosh(x)*\sinh(x)^5 + (a^6 - a^4 \\ & *b^2)*\sinh(x)^6 - a^6 + a^4*b^2 - 3*(a^6 - a^4*b^2)*\cosh(x)^4 - 3*(a^6 - a^4 \\ & *b^2 - 5*(a^6 - a^4*b^2)*\cosh(x)^2)*\sinh(x)^4 + 4*(5*(a^6 - a^4*b^2)*\cosh(\\ & x)^3 - 3*(a^6 - a^4*b^2)*\cosh(x))*\sinh(x)^3 + 3*(a^6 - a^4*b^2)*\cosh(x)^2 + \\ & 3*(a^6 - a^4*b^2 + 5*(a^6 - a^4*b^2)*\cosh(x)^4 - 6*(a^6 - a^4*b^2)*\cosh(x) \\ & ^2)*\sinh(x)^2 + 6*((a^6 - a^4*b^2)*\cosh(x)^5 - 2*(a^6 - a^4*b^2)*\cosh(x)^3 \\ & + (a^6 - a^4*b^2)*\cosh(x))*\sinh(x)) \end{aligned}$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\coth^4(x)}{a + b \tanh(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(x)**4/(a+b*tanh(x)),x)

[Out] Integral(coth(x)**4/(a + b*tanh(x)), x)

Giac [A]

time = 0.42, size = 142, normalized size = 1.46

$$-\frac{b^5 \log(|ae^{2x} + be^{2x} + a - b|)}{a^6 - a^4 b^2} + \frac{x}{a - b} - \frac{(a^2 b + b^3) \log(|e^{2x} - 1|)}{a^4} - \frac{2(4a^3 + 3ab^2 + 3(2a^3 - a^2b + ab^2)e^{4x} - 3(2a^3 - a^2b + 2ab^2)e^{2x})}{3a^4(e^{2x} - 1)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(x)^4/(a+b*tanh(x)),x, algorithm="giac")

[Out] $-b^5 \log(\text{abs}(a \cdot e^{2x} + b \cdot e^{2x} + a - b)) / (a^6 - a^4 \cdot b^2) + x / (a - b) - (a^2 \cdot b + b^3) \cdot \log(\text{abs}(e^{2x} - 1)) / a^4 - 2 / 3 \cdot (4 \cdot a^3 + 3 \cdot a \cdot b^2 + 3 \cdot (2 \cdot a^3 - a^2 \cdot b - a^2 \cdot b + a \cdot b^2) \cdot e^{4x} - 3 \cdot (2 \cdot a^3 - a^2 \cdot b + 2 \cdot a \cdot b^2) \cdot e^{2x}) / (a^4 \cdot (e^{2x} - 1)^3)$

Mupad [B]

time = 1.50, size = 163, normalized size = 1.68

$$\frac{x}{a - b} - \frac{8}{3a(3e^{2x} - 3e^{4x} + e^{6x} - 1)} - \frac{b^5 \ln(a - b + ae^{2x} + be^{2x})}{a^6 - a^4 b^2} - \frac{\ln(e^{2x} - 1)(a^2 b + b^3)}{a^4} - \frac{2(2a^3 + a^2 b + b^3)}{a^3(a + b)(e^{2x} - 1)} - \frac{2(2a^2 + ab - b^2)}{a^2(a + b)(e^{4x} - 2e^{2x} + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(coth(x)^4/(a + b*tanh(x)),x)

[Out] $x / (a - b) - 8 / (3 \cdot a \cdot (3 \cdot \exp(2x) - 3 \cdot \exp(4x) + \exp(6x) - 1)) - (b^5 \cdot \log(a - b + a \cdot \exp(2x) + b \cdot \exp(2x))) / (a^6 - a^4 \cdot b^2) - (\log(\exp(2x) - 1) \cdot (a^2 \cdot b + b^3)) / a^4 - (2 \cdot (a^2 \cdot b + 2 \cdot a^3 + b^3)) / (a^3 \cdot (a + b) \cdot (\exp(2x) - 1)) - (2 \cdot (a \cdot b + 2 \cdot a^2 - b^2)) / (a^2 \cdot (a + b) \cdot (\exp(4x) - 2 \cdot \exp(2x) + 1))$

3.143 $\int \frac{x \operatorname{sech}^2(x)}{(a+b \tanh(x))^2} dx$

Optimal. Leaf size=55

$$\frac{ax}{b(a^2 - b^2)} - \frac{\log(a \cosh(x) + b \sinh(x))}{a^2 - b^2} - \frac{x}{b(a + b \tanh(x))}$$

[Out] a*x/b/(a^2-b^2)-ln(a*cosh(x)+b*sinh(x))/(a^2-b^2)-x/b/(a+b*tanh(x))

Rubi [A]

time = 0.06, antiderivative size = 55, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {5574, 3565, 3611}

$$\frac{ax}{b(a^2 - b^2)} - \frac{\log(a \cosh(x) + b \sinh(x))}{a^2 - b^2} - \frac{x}{b(a + b \tanh(x))}$$

Antiderivative was successfully verified.

[In] Int[(x*Sech[x]^2)/(a + b*Tanh[x])^2,x]

[Out] (a*x)/(b*(a^2 - b^2)) - Log[a*Cosh[x] + b*Sinh[x]]/(a^2 - b^2) - x/(b*(a + b*Tanh[x]))

Rule 3565

Int[((a_) + (b_)*tan[(c_) + (d_)*(x_)])^(-1), x_Symbol] := Simp[a*(x/(a^2 + b^2)), x] + Dist[b/(a^2 + b^2), Int[(b - a*Tan[c + d*x])/(a + b*Tan[c + d*x]), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0]

Rule 3611

Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]

Rule 5574

Int[((e_) + (f_)*(x_))^(m_)*Sech[(c_) + (d_)*(x_)]^2*((a_) + (b_)*Tanh[(c_) + (d_)*(x_)]^(n_), x_Symbol] := Simp[(e + f*x)^m*((a + b*Tanh[c + d*x])^(n + 1)/(b*d*(n + 1))), x] - Dist[f*(m/(b*d*(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Tanh[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]

Rubi steps

$$\begin{aligned} \int \frac{x \operatorname{sech}^2(x)}{(a + b \tanh(x))^2} dx &= -\frac{x}{b(a + b \tanh(x))} + \frac{\int \frac{1}{a + b \tanh(x)} dx}{b} \\ &= \frac{ax}{b(a^2 - b^2)} - \frac{x}{b(a + b \tanh(x))} - \frac{i \int \frac{-ib - ia \tanh(x)}{a + b \tanh(x)} dx}{a^2 - b^2} \\ &= \frac{ax}{b(a^2 - b^2)} - \frac{\log(a \cosh(x) + b \sinh(x))}{a^2 - b^2} - \frac{x}{b(a + b \tanh(x))} \end{aligned}$$

Mathematica [A]

time = 0.14, size = 49, normalized size = 0.89

$$\frac{bx - a \log(a \cosh(x) + b \sinh(x))}{a^3 - ab^2} + \frac{x \sinh(x)}{a^2 \cosh(x) + ab \sinh(x)}$$

Antiderivative was successfully verified.

`[In] Integrate[(x*Sech[x]^2)/(a + b*Tanh[x])^2,x]``[Out] (b*x - a*Log[a*Cosh[x] + b*Sinh[x]])/(a^3 - a*b^2) + (x*Sinh[x])/(a^2*Cosh[x] + a*b*Sinh[x])`**Maple [A]**

time = 1.21, size = 73, normalized size = 1.33

method	result	size
risch	$\frac{2x}{a^2 - b^2} - \frac{2x}{(e^{2x}a + b e^{2x} + a - b)(a + b)} - \frac{\ln\left(e^{2x} + \frac{a - b}{a + b}\right)}{a^2 - b^2}$	73

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(x*sech(x)^2/(a+b*tanh(x))^2,x,method=_RETURNVERBOSE)``[Out] 2/(a^2-b^2)*x-2*x/(exp(2*x)*a+b*exp(2*x)+a-b)/(a+b)-1/(a^2-b^2)*ln(exp(2*x)+(a-b)/(a+b))`**Maxima [A]**

time = 0.41, size = 68, normalized size = 1.24

$$\frac{2xe^{(2x)}}{a^2 - 2ab + b^2 + (a^2 - b^2)e^{(2x)}} - \frac{\log\left(\frac{(a+b)e^{(2x)} + a - b}{a + b}\right)}{a^2 - b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(x*sech(x)^2/(a+b*tanh(x))^2,x, algorithm="maxima")`

[Out] $2*x*e^{(2*x)}/(a^2 - 2*a*b + b^2 + (a^2 - b^2)*e^{(2*x)}) - \log(((a + b)*e^{(2*x)} + a - b)/(a + b))/(a^2 - b^2)$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 182 vs. 2(55) = 110.

time = 0.37, size = 182, normalized size = 3.31

$$\frac{2(a+b)x \cosh(x)^2 + 4(a+b)x \cosh(x) \sinh(x) + 2(a+b)x \sinh(x)^2 - ((a+b) \cosh(x)^2 + 2(a+b) \cosh(x) \sinh(x) + (a+b) \sinh(x)^2 + a - b) \log\left(\frac{2(a \cosh(x) + b \sinh(x))}{\cosh(x) - \sinh(x)}\right)}{a^3 - a^2b - ab^2 + b^3 + (a^3 + a^2b - ab^2 - b^3) \cosh(x)^2 + 2(a^3 + a^2b - ab^2 - b^3) \cosh(x) \sinh(x) + (a^3 + a^2b - ab^2 - b^3) \sinh(x)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x*sech(x)^2/(a+b*tanh(x))^2,x, algorithm="fricas")`

[Out] $(2*(a + b)*x*\cosh(x)^2 + 4*(a + b)*x*\cosh(x)*\sinh(x) + 2*(a + b)*x*\sinh(x)^2 - ((a + b)*\cosh(x)^2 + 2*(a + b)*\cosh(x)*\sinh(x) + (a + b)*\sinh(x)^2 + a - b)*\log(2*(a*\cosh(x) + b*\sinh(x))/(\cosh(x) - \sinh(x))))/(a^3 - a^2*b - a*b^2 + b^3 + (a^3 + a^2*b - a*b^2 - b^3)*\cosh(x)^2 + 2*(a^3 + a^2*b - a*b^2 - b^3)*\cosh(x)*\sinh(x) + (a^3 + a^2*b - a*b^2 - b^3)*\sinh(x)^2)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x \operatorname{sech}^2(x)}{(a + b \tanh(x))^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x*sech(x)**2/(a+b*tanh(x))**2,x)`

[Out] `Integral(x*sech(x)**2/(a + b*tanh(x))**2, x)`

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 174 vs. 2(55) = 110.

time = 0.42, size = 174, normalized size = 3.16

$$\frac{2axe^{(2x)} + 2bx e^{(2x)} - ae^{(2x)} \log(-ae^{(2x)} - be^{(2x)} - a + b) - be^{(2x)} \log(-ae^{(2x)} - be^{(2x)} - a + b) - a \log(-ae^{(2x)} - be^{(2x)} - a + b) + b \log(-ae^{(2x)} - be^{(2x)} - a + b)}{a^3 e^{(2x)} + a^2 b e^{(2x)} - ab^2 e^{(2x)} - b^3 e^{(2x)} + a^3 - a^2 b - ab^2 + b^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x*sech(x)^2/(a+b*tanh(x))^2,x, algorithm="giac")`

[Out] $(2*a*x*e^{(2*x)} + 2*b*x*e^{(2*x)} - a*e^{(2*x)}*\log(-a*e^{(2*x)} - b*e^{(2*x)} - a + b) - b*e^{(2*x)}*\log(-a*e^{(2*x)} - b*e^{(2*x)} - a + b) - a*\log(-a*e^{(2*x)} - b*e^{(2*x)} - a + b) + b*\log(-a*e^{(2*x)} - b*e^{(2*x)} - a + b))/(a^3*e^{(2*x)} + a^2*b*e^{(2*x)} - a*b^2*e^{(2*x)} - b^3*e^{(2*x)} + a^3 - a^2*b - a*b^2 + b^3)$

Mupad [B]

time = 1.15, size = 69, normalized size = 1.25

$$\frac{2x}{a^2 - b^2} - \frac{\ln(a - b + ae^{2x} + be^{2x})}{a^2 - b^2} - \frac{2x}{(a + b)(a - b + e^{2x}(a + b))}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(x/(cosh(x)^2*(a + b*tanh(x))^2),x)
```

```
[Out] (2*x)/(a^2 - b^2) - log(a - b + a*exp(2*x) + b*exp(2*x))/(a^2 - b^2) - (2*x)/((a + b)*(a - b + exp(2*x)*(a + b)))
```

3.144 $\int \frac{x \operatorname{sech}^2(c+dx)}{a+b \tanh^2(c+dx)} dx$

Optimal. Leaf size=231

$$\frac{x \log\left(1 + \frac{(a+b)e^{2c+2dx}}{a-2\sqrt{-a}\sqrt{b}-b}\right)}{2\sqrt{-a}\sqrt{b}d} - \frac{x \log\left(1 + \frac{(a+b)e^{2c+2dx}}{a+2\sqrt{-a}\sqrt{b}-b}\right)}{2\sqrt{-a}\sqrt{b}d} + \frac{\operatorname{PolyLog}\left(2, -\frac{(a+b)e^{2c+2dx}}{a-2\sqrt{-a}\sqrt{b}-b}\right)}{4\sqrt{-a}\sqrt{b}d^2} - \frac{\operatorname{PolyLog}\left(2, -\frac{(a+b)e^{2c+2dx}}{a+2\sqrt{-a}\sqrt{b}-b}\right)}{4\sqrt{-a}\sqrt{b}d^2}$$

[Out] $\frac{1}{2}x \ln\left(1 + \frac{(a+b)\exp(2dx+2c)}{a-2\sqrt{-a}\sqrt{b}-b}\right) / \sqrt{-a}\sqrt{b}d - \frac{1}{2}x \ln\left(1 + \frac{(a+b)\exp(2dx+2c)}{a+2\sqrt{-a}\sqrt{b}-b}\right) / \sqrt{-a}\sqrt{b}d + \frac{1}{4} \operatorname{polylog}\left(2, -\frac{(a+b)\exp(2dx+2c)}{a-2\sqrt{-a}\sqrt{b}-b}\right) / d^2 - \frac{1}{4} \operatorname{polylog}\left(2, -\frac{(a+b)\exp(2dx+2c)}{a+2\sqrt{-a}\sqrt{b}-b}\right) / d^2$

Rubi [A]

time = 0.40, antiderivative size = 231, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 6, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {5751, 3401, 2296, 2221, 2317, 2438}

$$\frac{\operatorname{Li}_2\left(-\frac{(a+b)e^{2c+2dx}}{a-b-2\sqrt{-a}\sqrt{b}}\right)}{4\sqrt{-a}\sqrt{b}d^2} - \frac{\operatorname{Li}_2\left(-\frac{(a+b)e^{2c+2dx}}{a-b+2\sqrt{-a}\sqrt{b}}\right)}{4\sqrt{-a}\sqrt{b}d^2} + \frac{x \log\left(\frac{(a+b)e^{2c+2dx}}{-2\sqrt{-a}\sqrt{b}+a-b} + 1\right)}{2\sqrt{-a}\sqrt{b}d} - \frac{x \log\left(\frac{(a+b)e^{2c+2dx}}{2\sqrt{-a}\sqrt{b}+a-b} + 1\right)}{2\sqrt{-a}\sqrt{b}d}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(x \operatorname{Sech}[c + d*x]^2)/(a + b \operatorname{Tanh}[c + d*x]^2), x]$

[Out] $\frac{(x \operatorname{Log}[1 + ((a + b)E^{(2*c + 2*d*x)})/(a - 2*\operatorname{Sqrt}[-a]*\operatorname{Sqrt}[b] - b)])/(2*\operatorname{Sqrt}[-a]*\operatorname{Sqrt}[b]*d) - (x \operatorname{Log}[1 + ((a + b)E^{(2*c + 2*d*x)})/(a + 2*\operatorname{Sqrt}[-a]*\operatorname{Sqrt}[b] - b)])/(2*\operatorname{Sqrt}[-a]*\operatorname{Sqrt}[b]*d) + \operatorname{PolyLog}[2, -(((a + b)E^{(2*c + 2*d*x)})/(a - 2*\operatorname{Sqrt}[-a]*\operatorname{Sqrt}[b] - b))]/(4*\operatorname{Sqrt}[-a]*\operatorname{Sqrt}[b]*d^2) - \operatorname{PolyLog}[2, -(((a + b)E^{(2*c + 2*d*x)})/(a + 2*\operatorname{Sqrt}[-a]*\operatorname{Sqrt}[b] - b))]/(4*\operatorname{Sqrt}[-a]*\operatorname{Sqrt}[b]*d^2)$

Rule 2221

$\operatorname{Int}[(((F_)^\wedge((g_.) * ((e_.) + (f_.) * (x_))))^\wedge(n_.) * ((c_.) + (d_.) * (x_))^\wedge(m_.) / ((a_.) + (b_.) * ((F_)^\wedge((g_.) * ((e_.) + (f_.) * (x_))))^\wedge(n_.)], x_Symbol] \rightarrow \operatorname{Simp} [((c + d*x)^\wedge m / (b*f*g*n*\operatorname{Log}[F])) * \operatorname{Log}[1 + b*((F)^\wedge(g*(e + f*x)))^\wedge n/a], x] - \operatorname{Dist}[d*(m / (b*f*g*n*\operatorname{Log}[F])), \operatorname{Int}[(c + d*x)^\wedge(m - 1) * \operatorname{Log}[1 + b*((F)^\wedge(g*(e + f*x)))^\wedge n/a], x], x] /; \operatorname{FreeQ}\{F, a, b, c, d, e, f, g, n\}, x \} \&\& \operatorname{IGtQ}[m, 0]$

Rule 2296

$\operatorname{Int}[(F_)^\wedge(u) * ((f_.) + (g_.) * (x_))^\wedge(m) / ((a_.) + (b_.) * (F_)^\wedge(u) + (c_.) * (F_)^\wedge(v)], x_Symbol] \rightarrow \operatorname{With}\{q = \operatorname{Rt}[b^2 - 4*a*c, 2]\}, \operatorname{Dist}[2*(c/q), \operatorname{Int}[(f + g*x)^\wedge m * (F)^\wedge u / (b - q + 2*c*(F)^\wedge u), x], x] - \operatorname{Dist}[2*(c/q), \operatorname{Int}[(f + g*x)^\wedge m$

*(F^u/(b + q + 2*c*F^u)), x], x]] /; FreeQ[{F, a, b, c, f, g}, x] && EqQ[v, 2*u] && LinearQ[u, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[m, 0]

Rule 2317

Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_)))^(n_.)], x_Symbol] :> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]

Rule 2438

Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]

Rule 3401

Int[((c_.) + (d_.)*(x_)^(m_.))/((a_) + (b_.)*sin[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]), x_Symbol] :> Dist[2, Int[((c + d*x)^m*(E^((-I)*e + f*fz*x)/(b + (2*a*E^((-I)*e + f*fz*x))/E^(I*Pi*(k - 1/2)) - (b*E^(2*((-I)*e + f*fz*x)))/E^(2*I*k*Pi)))/E^(I*Pi*(k - 1/2)), x], x] /; FreeQ[{a, b, c, d, e, f, fz}, x] && IntegerQ[2*k] && NeQ[a^2 - b^2, 0] && IGtQ[m, 0]

Rule 5751

Int[((f_.) + (g_.)*(x_)^(m_.))*Sech[(d_.) + (e_.)*(x_)^2]/((b_) + (c_.)*Tanh[(d_.) + (e_.)*(x_)^2]), x_Symbol] :> Dist[2, Int[(f + g*x)^m/(b - c + (b + c)*Cosh[2*d + 2*e*x]), x], x] /; FreeQ[{b, c, d, e, f, g}, x] && IGtQ[m, 0]

Rubi steps

$$\begin{aligned}
\int \frac{x \operatorname{sech}^2(c+dx)}{a+b \tanh^2(c+dx)} dx &= 2 \int \frac{x}{a-b+(a+b) \cosh(2c+2dx)} dx \\
&= 4 \int \frac{e^{2c+2dx} x}{a+b+2(a-b)e^{2c+2dx}+(a+b)e^{2(2c+2dx)}} dx \\
&= \frac{(2(a+b)) \int \frac{e^{2c+2dx} x}{2(a-b)-4\sqrt{-a}\sqrt{b}+2(a+b)e^{2c+2dx}} dx}{\sqrt{-a}\sqrt{b}} - \frac{(2(a+b)) \int \frac{e^{2c+2dx} x}{2(a-b)+4\sqrt{-a}\sqrt{b}+2(a+b)e^{2c+2dx}} dx}{\sqrt{-a}\sqrt{b}} \\
&= \frac{x \log\left(1+\frac{(a+b)e^{2c+2dx}}{a-2\sqrt{-a}\sqrt{b}-b}\right)}{2\sqrt{-a}\sqrt{b}d} - \frac{x \log\left(1+\frac{(a+b)e^{2c+2dx}}{a+2\sqrt{-a}\sqrt{b}-b}\right)}{2\sqrt{-a}\sqrt{b}d} - \frac{\int \log\left(1+\frac{2(a+b)e^{2c+2dx}}{2(a-b)+4\sqrt{-a}\sqrt{b}+2(a+b)e^{2c+2dx}}\right) dx}{2\sqrt{-a}\sqrt{b}} \\
&= \frac{x \log\left(1+\frac{(a+b)e^{2c+2dx}}{a-2\sqrt{-a}\sqrt{b}-b}\right)}{2\sqrt{-a}\sqrt{b}d} - \frac{x \log\left(1+\frac{(a+b)e^{2c+2dx}}{a+2\sqrt{-a}\sqrt{b}-b}\right)}{2\sqrt{-a}\sqrt{b}d} - \frac{\operatorname{Subst}\left(\int \frac{\log\left(1+\frac{2(a+b)e^{2c+2dx}}{2(a-b)+4\sqrt{-a}\sqrt{b}+2(a+b)e^{2c+2dx}}\right) dx}{2\sqrt{-a}\sqrt{b}}\right)}{2\sqrt{-a}\sqrt{b}} \\
&= \frac{x \log\left(1+\frac{(a+b)e^{2c+2dx}}{a-2\sqrt{-a}\sqrt{b}-b}\right)}{2\sqrt{-a}\sqrt{b}d} - \frac{x \log\left(1+\frac{(a+b)e^{2c+2dx}}{a+2\sqrt{-a}\sqrt{b}-b}\right)}{2\sqrt{-a}\sqrt{b}d} + \frac{\operatorname{Li}_2\left(-\frac{(a+b)e^{2c+2dx}}{a-2\sqrt{-a}\sqrt{b}-b}\right)}{4\sqrt{-a}\sqrt{b}}
\end{aligned}$$

Mathematica [A]

time = 1.65, size = 223, normalized size = 0.97

$$\frac{4c \operatorname{ArcTan}\left(\frac{a-b+(a+b)e^{2(c+dx)}}{2\sqrt{a}\sqrt{b}}\right)}{\sqrt{a}} + \frac{2(c+dx) \left(\log\left(1+\frac{(a+b)e^{2(c+dx)}}{a-2\sqrt{-a}\sqrt{b}-b}\right) - \log\left(1+\frac{(a+b)e^{2(c+dx)}}{a+2\sqrt{-a}\sqrt{b}-b}\right) \right) + \operatorname{PolyLog}\left(2, -\frac{(a+b)e^{2(c+dx)}}{a-2\sqrt{-a}\sqrt{b}-b}\right) - \operatorname{PolyLog}\left(2, -\frac{(a+b)e^{2(c+dx)}}{a+2\sqrt{-a}\sqrt{b}-b}\right)}{4\sqrt{b}d^2}$$

Antiderivative was successfully verified.

[In] Integrate[(x*Sech[c + d*x]^2)/(a + b*Tanh[c + d*x]^2), x]

[Out] $\left(\frac{-4c \operatorname{ArcTan}\left[\frac{a-b+(a+b)E^{2(c+dx)}}{2\sqrt{a}\sqrt{b}}\right]}{\sqrt{a}} + \frac{2(c+dx) \left(\log\left[1+\frac{(a+b)E^{2(c+dx)}}{a-2\sqrt{-a}\sqrt{b}-b}\right] - \log\left[1+\frac{(a+b)E^{2(c+dx)}}{a+2\sqrt{-a}\sqrt{b}-b}\right]\right) + \operatorname{PolyLog}\left[2, -\frac{(a+b)E^{2(c+dx)}}{a-2\sqrt{-a}\sqrt{b}-b}\right] - \operatorname{PolyLog}\left[2, -\frac{(a+b)E^{2(c+dx)}}{a+2\sqrt{-a}\sqrt{b}-b}\right]}{4\sqrt{b}d^2}\right) / \sqrt{a}$

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 952 vs. 2(187) = 374.

time = 2.93, size = 953, normalized size = 4.13

method	result
risch	$-\frac{\ln\left(1-\frac{(a+b)e^{2dx+2c}}{-2\sqrt{-ab}-a+b}\right)bc}{2d^2\sqrt{-ab}\left(-2\sqrt{-ab}-a+b\right)} + \frac{\ln\left(1-\frac{(a+b)e^{2dx+2c}}{-2\sqrt{-ab}-a+b}\right)ac}{2d^2\sqrt{-ab}\left(-2\sqrt{-ab}-a+b\right)} - \frac{c^2}{d^2\left(-2\sqrt{-ab}-a+b\right)} - \frac{c^2}{2d^2\sqrt{-ab}} - \frac{1}{2d}$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(x*sech(d*x+c)^2/(a+b*tanh(d*x+c)^2),x,method=_RETURNVERBOSE)
```

```
[Out] -1/2/d^2/(-a*b)^(1/2)/(-2*(-a*b)^(1/2)-a+b)*ln(1-(a+b)*exp(2*d*x+2*c)/(-2*(-a*b)^(1/2)-a+b))*b*c-1/4/d^2/(-a*b)^(1/2)/(-2*(-a*b)^(1/2)-a+b)*polylog(2,(a+b)*exp(2*d*x+2*c)/(-2*(-a*b)^(1/2)-a+b))*b+1/2/d^2/(-a*b)^(1/2)/(-2*(-a*b)^(1/2)-a+b)*ln(1-(a+b)*exp(2*d*x+2*c)/(-2*(-a*b)^(1/2)-a+b))*a*c+1/2/d^2/(-2*(-a*b)^(1/2)-a+b)*polylog(2,(a+b)*exp(2*d*x+2*c)/(-2*(-a*b)^(1/2)-a+b))+1/4/d^2/(-a*b)^(1/2)*polylog(2,(a+b)*exp(2*d*x+2*c)/(2*(-a*b)^(1/2)-a+b))-1/d^2/(-2*(-a*b)^(1/2)-a+b)*c^2-1/2/d^2/(-a*b)^(1/2)*c^2-1/2/d/(-a*b)^(1/2)/(-2*(-a*b)^(1/2)-a+b)*ln(1-(a+b)*exp(2*d*x+2*c)/(-2*(-a*b)^(1/2)-a+b))*b*x-1/2/d^2/(-a*b)^(1/2)/(-2*(-a*b)^(1/2)-a+b)*a*c^2+1/2/d^2/(-a*b)^(1/2)/(-2*(-a*b)^(1/2)-a+b)*b*c^2+1/4/d^2/(-a*b)^(1/2)/(-2*(-a*b)^(1/2)-a+b)*polylog(2,(a+b)*exp(2*d*x+2*c)/(-2*(-a*b)^(1/2)-a+b))*a-1/d/(-a*b)^(1/2)/(-2*(-a*b)^(1/2)-a+b)*a*c*x+1/2/d/(-a*b)^(1/2)/(-2*(-a*b)^(1/2)-a+b)*ln(1-(a+b)*exp(2*d*x+2*c)/(-2*(-a*b)^(1/2)-a+b))*a*x+1/d/(-a*b)^(1/2)/(-2*(-a*b)^(1/2)-a+b)*b*c*x-2/d/(-2*(-a*b)^(1/2)-a+b)*c*x+1/2/(-a*b)^(1/2)/(-2*(-a*b)^(1/2)-a+b)*b*x^2-1/2/(-a*b)^(1/2)/(-2*(-a*b)^(1/2)-a+b)*a*x^2-1/d^2*c/(a*b)^(1/2)*arctan(1/4*(2*(a+b)*exp(2*d*x+2*c)+2*a-2*b)/(a*b)^(1/2))-1/d/(-a*b)^(1/2)*c*x+1/d^2/(-2*(-a*b)^(1/2)-a+b)*ln(1-(a+b)*exp(2*d*x+2*c)/(-2*(-a*b)^(1/2)-a+b))*c+1/2/d^2/(-a*b)^(1/2)*ln(1-(a+b)*exp(2*d*x+2*c)/(2*(-a*b)^(1/2)-a+b))*c+1/d/(-2*(-a*b)^(1/2)-a+b)*ln(1-(a+b)*exp(2*d*x+2*c)/(-2*(-a*b)^(1/2)-a+b))*x+1/2/d/(-a*b)^(1/2)*ln(1-(a+b)*exp(2*d*x+2*c)/(2*(-a*b)^(1/2)-a+b))*x-1/(-2*(-a*b)^(1/2)-a+b)*x^2-1/2/(-a*b)^(1/2)*x^2
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x*sech(d*x+c)^2/(a+b*tanh(d*x+c)^2),x, algorithm="maxima")
```

```
[Out] integrate(x*sech(d*x + c)^2/(b*tanh(d*x + c)^2 + a), x)
```

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 1516 vs. 2(185) = 370.

time = 0.42, size = 1516, normalized size = 6.56

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x*sech(d*x+c)^2/(a+b*tanh(d*x+c)^2),x, algorithm="fricas")
```

```
[Out] -1/2*((a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2))*c*log(2*sqrt(-(2*(a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2)) + a - b)/(a + b)) + 2*cosh(d*x + c) + 2*sinh(d*x + c)) + (a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2))*c*log(-2*sqrt(-(2*(a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2)) + a - b)/(a + b)) + 2*cosh(d*x + c) + 2*sinh(d*x + c)) - (a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2))*c*log(2*sqrt((2*(a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2)) - a + b)/(a + b)) + 2*cosh(d*x + c) + 2*sinh(d*x + c)) - (a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2))*c*log(-2*sqrt((2*(a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2)) - a + b)/(a + b)) + 2*cosh(d*x + c) + 2*sinh(d*x + c)) - (a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2))*dilog(-(((a - b)*cosh(d*x + c) + (a - b)*sinh(d*x + c) - 2*((a + b)*cosh(d*x + c) + (a + b)*sinh(d*x + c))*sqrt(-a*b/(a^2 + 2*a*b + b^2)))*sqrt(-(2*(a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2)) + a - b)/(a + b)) + a + b)/(a + b) + 1) - (a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2))*dilog((((a - b)*cosh(d*x + c) + (a - b)*sinh(d*x + c) - 2*((a + b)*cosh(d*x + c) + (a + b)*sinh(d*x + c))*sqrt(-a*b/(a^2 + 2*a*b + b^2)))*sqrt(-(2*(a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2)) + a - b)/(a + b)) - a - b)/(a + b) + 1) + (a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2))*dilog(-(((a - b)*cosh(d*x + c) + (a - b)*sinh(d*x + c) + 2*((a + b)*cosh(d*x + c) + (a + b)*sinh(d*x + c))*sqrt(-a*b/(a^2 + 2*a*b + b^2)))*sqrt((2*(a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2)) - a + b)/(a + b)) - a - b)/(a + b) + 1) - ((a + b)*d*x + (a + b)*c)*sqrt(-a*b/(a^2 + 2*a*b + b^2))*log((((a - b)*cosh(d*x + c) + (a - b)*sinh(d*x + c) - 2*((a + b)*cosh(d*x + c) + (a + b)*sinh(d*x + c))*sqrt(-a*b/(a^2 + 2*a*b + b^2)))*sqrt(-(2*(a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2)) + a - b)/(a + b)) + a + b)/(a + b) - ((a + b)*d*x + (a + b)*c)*sqrt(-a*b/(a^2 + 2*a*b + b^2))*log(-(((a - b)*cosh(d*x + c) + (a - b)*sinh(d*x + c) - 2*((a + b)*cosh(d*x + c) + (a + b)*sinh(d*x + c))*sqrt(-a*b/(a^2 + 2*a*b + b^2)))*sqrt(-(2*(a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2)) + a - b)/(a + b)) - a - b)/(a + b)) + ((a + b)*d*x + (a + b)*c)*sqrt(-a*b/(a^2 + 2*a*b + b^2))*log((((a - b)*cosh(d*x + c) + (a - b)*sinh(d*x + c) + 2*((a + b)*cosh(d*x + c) + (a + b)*sinh(d*x + c))*sqrt(-a*b/(a^2 + 2*a*b + b^2)))*sqrt((2*(a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2)) - a + b)/(a + b)) + a + b)/(a + b)) + ((a + b)*d*x + (a + b)*c)*sqrt(-a*b/(a^2 + 2*a*b + b^2))*log(-(((a - b)*cosh(d*x + c) + (a - b)*sinh(d*x + c) + 2*((a + b)*cosh(d*x + c) + (a + b)*sinh(d*x + c))*sqrt(-a*b/(a^2 + 2*a*b + b^2)))*sqrt((2*(a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2)) - a + b)/(a + b)) - a - b)/(a + b)))/(a*b*d^2)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x \operatorname{sech}^2(c + dx)}{a + b \tanh^2(c + dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x*sech(d*x+c)**2/(a+b*tanh(d*x+c)**2), x)

[Out] Integral(x*sech(c + d*x)**2/(a + b*tanh(c + d*x)**2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x*sech(d*x+c)^2/(a+b*tanh(d*x+c)^2), x, algorithm="giac")

[Out] integrate(x*sech(d*x + c)^2/(b*tanh(d*x + c)^2 + a), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{x}{\cosh(c + dx)^2 (b \tanh(c + dx)^2 + a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x/(cosh(c + d*x)^2*(a + b*tanh(c + d*x)^2)), x)

[Out] int(x/(cosh(c + d*x)^2*(a + b*tanh(c + d*x)^2)), x)

3.145 $\int \frac{x^2 \operatorname{sech}^2(c+dx)}{a+b \tanh^2(c+dx)} dx$

Optimal. Leaf size=351

$$\frac{x^2 \log\left(1 + \frac{(a+b)e^{2c+2dx}}{a-2\sqrt{-a}\sqrt{b}-b}\right)}{2\sqrt{-a}\sqrt{b}d} - \frac{x^2 \log\left(1 + \frac{(a+b)e^{2c+2dx}}{a+2\sqrt{-a}\sqrt{b}-b}\right)}{2\sqrt{-a}\sqrt{b}d} + \frac{x \operatorname{PolyLog}\left(2, -\frac{(a+b)e^{2c+2dx}}{a-2\sqrt{-a}\sqrt{b}-b}\right)}{2\sqrt{-a}\sqrt{b}d^2} - \frac{x \operatorname{PolyLog}\left(2, -\frac{(a+b)e^{2c+2dx}}{a+2\sqrt{-a}\sqrt{b}-b}\right)}{2\sqrt{-a}\sqrt{b}d^2}$$

[Out] $\frac{1}{2}x^2 \ln\left(\frac{1+(a+b)\exp(2dx+2c)}{(a-b-2\sqrt{-a}\sqrt{b})}\right) / \sqrt{-a}\sqrt{b}d - \frac{1}{2}x^2 \ln\left(\frac{1+(a+b)\exp(2dx+2c)}{(a-b+2\sqrt{-a}\sqrt{b})}\right) / \sqrt{-a}\sqrt{b}d + \frac{x \operatorname{PolyLog}\left(2, -\frac{(a+b)\exp(2dx+2c)}{(a-b-2\sqrt{-a}\sqrt{b})}\right)}{2\sqrt{-a}\sqrt{b}d^2} - \frac{x \operatorname{PolyLog}\left(2, -\frac{(a+b)\exp(2dx+2c)}{(a-b+2\sqrt{-a}\sqrt{b})}\right)}{2\sqrt{-a}\sqrt{b}d^2} + \frac{1}{4}x \operatorname{PolyLog}\left(3, -\frac{(a+b)\exp(2dx+2c)}{(a-b-2\sqrt{-a}\sqrt{b})}\right) / \sqrt{-a}\sqrt{b}d^3 - \frac{1}{4}x \operatorname{PolyLog}\left(3, -\frac{(a+b)\exp(2dx+2c)}{(a-b+2\sqrt{-a}\sqrt{b})}\right) / \sqrt{-a}\sqrt{b}d^3$

Rubi [A]

time = 0.63, antiderivative size = 351, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 7, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.269$, Rules used = {5751, 3401, 2296, 2221, 2611, 2320, 6724}

$$-\frac{\operatorname{Li}_3\left(-\frac{(a+b)e^{2c+2dx}}{a-b-2\sqrt{-a}\sqrt{b}}\right)}{4\sqrt{-a}\sqrt{b}d^3} + \frac{\operatorname{Li}_3\left(-\frac{(a+b)e^{2c+2dx}}{a-b+2\sqrt{-a}\sqrt{b}}\right)}{4\sqrt{-a}\sqrt{b}d^3} + \frac{x \operatorname{Li}_2\left(-\frac{(a+b)e^{2c+2dx}}{a-b-2\sqrt{-a}\sqrt{b}}\right)}{2\sqrt{-a}\sqrt{b}d^2} - \frac{x \operatorname{Li}_2\left(-\frac{(a+b)e^{2c+2dx}}{a-b+2\sqrt{-a}\sqrt{b}}\right)}{2\sqrt{-a}\sqrt{b}d^2} + \frac{x^2 \log\left(\frac{(a+b)e^{2c+2dx}}{-2\sqrt{-a}\sqrt{b}+a-b} + 1\right)}{2\sqrt{-a}\sqrt{b}d} - \frac{x^2 \log\left(\frac{(a+b)e^{2c+2dx}}{2\sqrt{-a}\sqrt{b}+a-b} + 1\right)}{2\sqrt{-a}\sqrt{b}d}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(x^2 \operatorname{Sech}[c + d*x]^2)/(a + b \operatorname{Tanh}[c + d*x]^2), x]$

[Out] $\frac{x^2 \operatorname{Log}\left[1 + \frac{(a+b)E^{2c+2dx}}{(a-2\sqrt{-a}\sqrt{b})}\right]}{2\sqrt{-a}\sqrt{b}d} - \frac{x^2 \operatorname{Log}\left[1 + \frac{(a+b)E^{2c+2dx}}{(a+2\sqrt{-a}\sqrt{b})}\right]}{2\sqrt{-a}\sqrt{b}d} + \frac{x \operatorname{PolyLog}\left[2, -\frac{(a+b)E^{2c+2dx}}{(a-2\sqrt{-a}\sqrt{b})}\right]}{2\sqrt{-a}\sqrt{b}d^2} - \frac{x \operatorname{PolyLog}\left[2, -\frac{(a+b)E^{2c+2dx}}{(a+2\sqrt{-a}\sqrt{b})}\right]}{2\sqrt{-a}\sqrt{b}d^2} - \frac{\operatorname{PolyLog}\left[3, -\frac{(a+b)E^{2c+2dx}}{(a-2\sqrt{-a}\sqrt{b})}\right]}{4\sqrt{-a}\sqrt{b}d^3} + \frac{\operatorname{PolyLog}\left[3, -\frac{(a+b)E^{2c+2dx}}{(a+2\sqrt{-a}\sqrt{b})}\right]}{4\sqrt{-a}\sqrt{b}d^3}$

Rule 2221

$\operatorname{Int}\left[\left(\frac{(F_+)^{((g_+)(e_+)+(f_+)(x_+))}}{(a_+)+(b_+)(F_+)^{((g_+)(e_+)+(f_+)(x_+))}}\right)^{(n_+)}\right], x_{\text{Symbol}}] \rightarrow \operatorname{Simp}\left[\frac{(c+d*x)^m}{(b*f*g*n*\operatorname{Log}[F])} \operatorname{Log}\left[1 + \frac{(F^{(g*(e+f*x)))^n}{a}}\right], x\right] - \operatorname{Dist}\left[d*\frac{(c+d*x)^{m-1}}{(b*f*g*n*\operatorname{Log}[F])} \operatorname{Log}\left[1 + \frac{(F^{(g*(e+f*x)))^n}{a}}\right], \operatorname{Int}\left[(c+d*x)^{m-1} \operatorname{Log}\left[1 + \frac{(F^{(g*(e+f*x)))^n}{a}}\right], x\right], x\right]; \operatorname{FreeQ}\{F, a, b, c, d, e, f, g, n\}, x\} \&\& \operatorname{IGtQ}[m, 0]$

Rule 2296

```
Int[((F_)^(u_)*((f_.) + (g_.)*(x_))^(m_.))/((a_.) + (b_.)*(F_)^(u_) + (c_.)
*(F_)^(v_)), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Dist[2*(c/q), Int[
(f + g*x)^m*(F^u/(b - q + 2*c*F^u)), x], x] - Dist[2*(c/q), Int[(f + g*x)^m
*(F^u/(b + q + 2*c*F^u)), x], x]] /; FreeQ[{F, a, b, c, f, g}, x] && EqQ[v,
2*u] && LinearQ[u, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[m, 0]
```

Rule 2320

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x]
, Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Functi
onOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[
{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*
(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2611

```
Int[Log[1 + (e_.)*((F_)^(c_.)*((a_.) + (b_.)*(x_)))^(n_.)]*(f_.) + (g_.)
*(x_)^(m_.), x_Symbol] := Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m
- 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e,
f, g, n}, x] && GtQ[m, 0]
```

Rule 3401

```
Int[((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*sin[(e_.) + Pi*(k_.) + (Comple
x[0, fz_])*(f_.)*(x_)]), x_Symbol] := Dist[2, Int[((c + d*x)^m*(E^((-I)*e +
f*fz*x)/(b + (2*a*E^((-I)*e + f*fz*x))/E^(I*Pi*(k - 1/2)) - (b*E^(2*((-I)*
e + f*fz*x))/E^(2*I*k*Pi))))/E^(I*Pi*(k - 1/2)), x], x] /; FreeQ[{a, b, c,
d, e, f, fz}, x] && IntegerQ[2*k] && NeQ[a^2 - b^2, 0] && IGtQ[m, 0]
```

Rule 5751

```
Int[((f_.) + (g_.)*(x_))^(m_.)*Sech[(d_.) + (e_.)*(x_)]^2/((b_) + (c_.)*T
anh[(d_.) + (e_.)*(x_)]^2), x_Symbol] := Dist[2, Int[(f + g*x)^m/(b - c + (
b + c)*Cosh[2*d + 2*e*x]), x], x] /; FreeQ[{b, c, d, e, f, g}, x] && IGtQ[m
, 0]
```

Rule 6724

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

Rubi steps

$$\begin{aligned}
\int \frac{x^2 \operatorname{sech}^2(c+dx)}{a+b \tanh^2(c+dx)} dx &= 2 \int \frac{x^2}{a-b+(a+b) \cosh(2c+2dx)} dx \\
&= 4 \int \frac{e^{2c+2dx} x^2}{a+b+2(a-b)e^{2c+2dx}+(a+b)e^{2(2c+2dx)}} dx \\
&= \frac{(2(a+b)) \int \frac{e^{2c+2dx} x^2}{2(a-b)-4\sqrt{-a}\sqrt{b}+2(a+b)e^{2c+2dx}} dx}{\sqrt{-a}\sqrt{b}} - \frac{(2(a+b)) \int \frac{e^{2c+2dx} x^2}{2(a-b)+4\sqrt{-a}\sqrt{b}+2(a+b)e^{2c+2dx}} dx}{\sqrt{-a}\sqrt{b}} \\
&= \frac{x^2 \log\left(1+\frac{(a+b)e^{2c+2dx}}{a-2\sqrt{-a}\sqrt{b}-b}\right)}{2\sqrt{-a}\sqrt{b}d} - \frac{x^2 \log\left(1+\frac{(a+b)e^{2c+2dx}}{a+2\sqrt{-a}\sqrt{b}-b}\right)}{2\sqrt{-a}\sqrt{b}d} - \frac{\int x \log\left(1+\frac{(a+b)e^{2c+2dx}}{a-2\sqrt{-a}\sqrt{b}-b}\right) dx}{2\sqrt{-a}\sqrt{b}} \\
&= \frac{x^2 \log\left(1+\frac{(a+b)e^{2c+2dx}}{a-2\sqrt{-a}\sqrt{b}-b}\right)}{2\sqrt{-a}\sqrt{b}d} - \frac{x^2 \log\left(1+\frac{(a+b)e^{2c+2dx}}{a+2\sqrt{-a}\sqrt{b}-b}\right)}{2\sqrt{-a}\sqrt{b}d} + \frac{x \operatorname{Li}_2\left(-\frac{(a+b)e^{2c+2dx}}{a-2\sqrt{-a}\sqrt{b}-b}\right)}{2\sqrt{-a}\sqrt{b}} \\
&= \frac{x^2 \log\left(1+\frac{(a+b)e^{2c+2dx}}{a-2\sqrt{-a}\sqrt{b}-b}\right)}{2\sqrt{-a}\sqrt{b}d} - \frac{x^2 \log\left(1+\frac{(a+b)e^{2c+2dx}}{a+2\sqrt{-a}\sqrt{b}-b}\right)}{2\sqrt{-a}\sqrt{b}d} + \frac{x \operatorname{Li}_2\left(-\frac{(a+b)e^{2c+2dx}}{a-2\sqrt{-a}\sqrt{b}-b}\right)}{2\sqrt{-a}\sqrt{b}} \\
&= \frac{x^2 \log\left(1+\frac{(a+b)e^{2c+2dx}}{a-2\sqrt{-a}\sqrt{b}-b}\right)}{2\sqrt{-a}\sqrt{b}d} - \frac{x^2 \log\left(1+\frac{(a+b)e^{2c+2dx}}{a+2\sqrt{-a}\sqrt{b}-b}\right)}{2\sqrt{-a}\sqrt{b}d} + \frac{x \operatorname{Li}_2\left(-\frac{(a+b)e^{2c+2dx}}{a-2\sqrt{-a}\sqrt{b}-b}\right)}{2\sqrt{-a}\sqrt{b}}
\end{aligned}$$

Mathematica [C] Result contains complex when optimal does not.
time = 1.65, size = 316, normalized size = 0.90

$$\frac{i \left(2d^2 x^2 \log\left(1 + \frac{(\sqrt{a} + \sqrt{b})e^{2(c+dx)}}{\sqrt{a} + \sqrt{b}}\right) - 2d^2 x^2 \log\left(1 + \frac{(\sqrt{a} + \sqrt{b})e^{2(c+dx)}}{\sqrt{a} - \sqrt{b}}\right) + 2dx \operatorname{PolyLog}\left(2, -\frac{(\sqrt{a} - \sqrt{b})e^{2(c+dx)}}{\sqrt{a} + \sqrt{b}}\right) - 2dx \operatorname{PolyLog}\left(2, -\frac{(\sqrt{a} - \sqrt{b})e^{2(c+dx)}}{\sqrt{a} - \sqrt{b}}\right) - \operatorname{PolyLog}\left(3, -\frac{(\sqrt{a} - \sqrt{b})e^{2(c+dx)}}{\sqrt{a} + \sqrt{b}}\right) + \operatorname{PolyLog}\left(3, -\frac{(\sqrt{a} + \sqrt{b})e^{2(c+dx)}}{\sqrt{a} - \sqrt{b}}\right) \right)}{4\sqrt{a}\sqrt{b}d^3}$$

Antiderivative was successfully verified.

```

[In] Integrate[(x^2*Sech[c + d*x]^2)/(a + b*Tanh[c + d*x]^2), x]
[Out] ((I/4)*(2*d^2*x^2*Log[1 + ((Sqrt[a] - I*Sqrt[b])*E^(2*(c + d*x)))/(Sqrt[a] + I*Sqrt[b])] - 2*d^2*x^2*Log[1 + ((Sqrt[a] + I*Sqrt[b])*E^(2*(c + d*x)))/(Sqrt[a] - I*Sqrt[b])] + 2*d*x*PolyLog[2, -(((Sqrt[a] - I*Sqrt[b])*E^(2*(c + d*x)))/(Sqrt[a] + I*Sqrt[b]))] - 2*d*x*PolyLog[2, -(((Sqrt[a] + I*Sqrt[b])*E^(2*(c + d*x)))/(Sqrt[a] - I*Sqrt[b]))] - PolyLog[3, -(((Sqrt[a] - I*Sqrt[b])*E^(2*(c + d*x)))/(Sqrt[a] + I*Sqrt[b]))] + PolyLog[3, -(((Sqrt[a] + I*Sqrt[b])*E^(2*(c + d*x)))/(Sqrt[a] - I*Sqrt[b]))]))/(Sqrt[a]*Sqrt[b]*d^3)

```

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 1185 vs. 2(285) = 570.
time = 2.86, size = 1186, normalized size = 3.38

method	result
risch	$-\frac{\text{polylog}\left(3, \frac{(a+b)e^{2dx+2c}}{2\sqrt{-ab}-a+b}\right)}{4d^3\sqrt{-ab}} - \frac{\text{polylog}\left(3, \frac{(a+b)e^{2dx+2c}}{-2\sqrt{-ab}-a+b}\right)}{2d^3(-2\sqrt{-ab}-a+b)} + \frac{2c^3}{3d^3\sqrt{-ab}} + \frac{4c^3}{3d^3(-2\sqrt{-ab}-a+b)} + \frac{b \text{ polylog}}{4d^3\sqrt{-ab}}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x^2*sech(d*x+c)^2/(a+b*tanh(d*x+c)^2),x,method=_RETURNVERBOSE)`

[Out]
$$\frac{2}{3}d^3c^3/(-ab)^{(1/2)} + \frac{4}{3}d^3c^3/(-2*(-ab)^{(1/2)}-a+b) - \frac{1}{4}d^3/(-ab)^{(1/2)} * \text{polylog}\left(3, \frac{(a+b)\exp(2d*x+2c)}{(2*(-ab)^{(1/2)}-a+b)}\right) - \frac{1}{2}d^3/(-2*(-ab)^{(1/2)}-a+b) * \text{polylog}\left(3, \frac{(a+b)\exp(2d*x+2c)}{(-2*(-ab)^{(1/2)}-a+b)}\right) + \frac{1}{2}d/(-ab)^{(1/2)}/(-2*(-ab)^{(1/2)}-a+b) * a * \ln(1-(a+b)\exp(2d*x+2c)/(-2*(-ab)^{(1/2)}-a+b)) * x^2 + \frac{1}{2}d^2/(-ab)^{(1/2)}/(-2*(-ab)^{(1/2)}-a+b) * a * \text{polylog}\left(2, \frac{(a+b)\exp(2d*x+2c)}{(-2*(-ab)^{(1/2)}-a+b)}\right) * x + \frac{1}{2}d^3/(-ab)^{(1/2)}/(-2*(-ab)^{(1/2)}-a+b) * b * \ln(1-(a+b)\exp(2d*x+2c)/(-2*(-ab)^{(1/2)}-a+b)) * c^2 - \frac{1}{2}d/(-ab)^{(1/2)}/(-2*(-ab)^{(1/2)}-a+b) * b * \ln(1-(a+b)\exp(2d*x+2c)/(-2*(-ab)^{(1/2)}-a+b)) * x^2 - \frac{1}{2}d^2/(-ab)^{(1/2)}/(-2*(-ab)^{(1/2)}-a+b) * b * \text{polylog}\left(2, \frac{(a+b)\exp(2d*x+2c)}{(-2*(-ab)^{(1/2)}-a+b)}\right) * x + \frac{1}{d^2}c^2/(-ab)^{(1/2)}/(-2*(-ab)^{(1/2)}-a+b) * a * x + \frac{1}{4}d^3/(-ab)^{(1/2)}/(-2*(-ab)^{(1/2)}-a+b) * b * \text{polylog}\left(3, \frac{(a+b)\exp(2d*x+2c)}{(-2*(-ab)^{(1/2)}-a+b)}\right) - \frac{1}{4}d^3/(-ab)^{(1/2)}/(-2*(-ab)^{(1/2)}-a+b) * a * \text{polylog}\left(3, \frac{(a+b)\exp(2d*x+2c)}{(-2*(-ab)^{(1/2)}-a+b)}\right) - \frac{2}{3}d^3c^3/(-ab)^{(1/2)}/(-2*(-ab)^{(1/2)}-a+b) * b + \frac{2}{3}d^3c^3/(-ab)^{(1/2)}/(-2*(-ab)^{(1/2)}-a+b) * a - \frac{1}{d^2}c^2/(-ab)^{(1/2)}/(-2*(-ab)^{(1/2)}-a+b) * b * x - \frac{1}{2}d^3/(-ab)^{(1/2)}/(-2*(-ab)^{(1/2)}-a+b) * a * \ln(1-(a+b)\exp(2d*x+2c)/(-2*(-ab)^{(1/2)}-a+b)) * c^2 + \frac{2}{d^2}c^2/(-2*(-ab)^{(1/2)}-a+b) * x + \frac{1}{d}/(-2*(-ab)^{(1/2)}-a+b) * \ln(1-(a+b)\exp(2d*x+2c)/(-2*(-ab)^{(1/2)}-a+b)) * x^2 + \frac{1}{d^2}/(-2*(-ab)^{(1/2)}-a+b) * \text{polylog}\left(2, \frac{(a+b)\exp(2d*x+2c)}{(-2*(-ab)^{(1/2)}-a+b)}\right) * x + \frac{1}{2}d^2/(-ab)^{(1/2)} * \text{polylog}\left(2, \frac{(a+b)\exp(2d*x+2c)}{(2*(-ab)^{(1/2)}-a+b)}\right) * x + \frac{1}{2}d/(-ab)^{(1/2)} * \ln(1-(a+b)\exp(2d*x+2c)/(2*(-ab)^{(1/2)}-a+b)) * x^2 + \frac{1}{d^2}c^2/(-ab)^{(1/2)} * x + \frac{1}{3}/(-ab)^{(1/2)}/(-2*(-ab)^{(1/2)}-a+b) * b * x^3 - \frac{1}{3}/(-ab)^{(1/2)}/(-2*(-ab)^{(1/2)}-a+b) * a * x^3 + \frac{1}{d^3}c^2/(a*b)^{(1/2)} * \arctan\left(\frac{1}{4} * (2*(a+b)\exp(2d*x+2c) + 2*a - 2*b)/(a*b)^{(1/2)}\right) - \frac{1}{2}d^3/(-ab)^{(1/2)} * \ln(1-(a+b)\exp(2d*x+2c)/(2*(-ab)^{(1/2)}-a+b)) * c^2 - \frac{1}{d^3}/(-2*(-ab)^{(1/2)}-a+b) * \ln(1-(a+b)\exp(2d*x+2c)/(-2*(-ab)^{(1/2)}-a+b)) * c^2 - \frac{1}{3}/(-ab)^{(1/2)} * x^3 - \frac{2}{3}/(-2*(-ab)^{(1/2)}-a+b) * x^3$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^2*sech(d*x+c)^2/(a+b*tanh(d*x+c)^2),x, algorithm="maxima")`

[Out] integrate(x^2*sech(d*x + c)^2/(b*tanh(d*x + c)^2 + a), x)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 2110 vs. 2(283) = 566.

time = 0.40, size = 2110, normalized size = 6.01

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2*sech(d*x+c)^2/(a+b*tanh(d*x+c)^2),x, algorithm="fricas")

[Out]
$$\frac{1}{2} * (2 * (a + b) * \sqrt{-a * b / (a^2 + 2 * a * b + b^2)}) * d * x * \operatorname{dilog}(-(((a - b) * \cosh(d * x + c) + (a - b) * \sinh(d * x + c) - 2 * ((a + b) * \cosh(d * x + c) + (a + b) * \sinh(d * x + c))) * \sqrt{-a * b / (a^2 + 2 * a * b + b^2)})) * \sqrt{-(2 * (a + b) * \sqrt{-a * b / (a^2 + 2 * a * b + b^2)}) + a - b} / (a + b) + a + b) / (a + b) + 1 + 2 * (a + b) * \sqrt{-a * b / (a^2 + 2 * a * b + b^2)} * d * x * \operatorname{dilog}(((a - b) * \cosh(d * x + c) + (a - b) * \sinh(d * x + c) - 2 * ((a + b) * \cosh(d * x + c) + (a + b) * \sinh(d * x + c))) * \sqrt{-a * b / (a^2 + 2 * a * b + b^2)}) * \sqrt{-(2 * (a + b) * \sqrt{-a * b / (a^2 + 2 * a * b + b^2)}) + a - b} / (a + b) - a - b) / (a + b) + 1 - 2 * (a + b) * \sqrt{-a * b / (a^2 + 2 * a * b + b^2)} * d * x * \operatorname{dilog}(-(((a - b) * \cosh(d * x + c) + (a - b) * \sinh(d * x + c) + 2 * ((a + b) * \cosh(d * x + c) + (a + b) * \sinh(d * x + c))) * \sqrt{-a * b / (a^2 + 2 * a * b + b^2)})) * \sqrt{(2 * (a + b) * \sqrt{-a * b / (a^2 + 2 * a * b + b^2)}) - a + b} / (a + b) + a + b) / (a + b) + 1 - 2 * (a + b) * \sqrt{-a * b / (a^2 + 2 * a * b + b^2)} * d * x * \operatorname{dilog}(((a - b) * \cosh(d * x + c) + (a - b) * \sinh(d * x + c) + 2 * ((a + b) * \cosh(d * x + c) + (a + b) * \sinh(d * x + c))) * \sqrt{-a * b / (a^2 + 2 * a * b + b^2)}) * \sqrt{(2 * (a + b) * \sqrt{-a * b / (a^2 + 2 * a * b + b^2)}) - a + b} / (a + b) - a - b) / (a + b) + 1 + (a + b) * \sqrt{-a * b / (a^2 + 2 * a * b + b^2)} * c^2 * \log(2 * \sqrt{-(2 * (a + b) * \sqrt{-a * b / (a^2 + 2 * a * b + b^2)}) + a - b} / (a + b) + 2 * \cosh(d * x + c) + 2 * \sinh(d * x + c)) + (a + b) * \sqrt{-a * b / (a^2 + 2 * a * b + b^2)} * c^2 * \log(-2 * \sqrt{-(2 * (a + b) * \sqrt{-a * b / (a^2 + 2 * a * b + b^2)}) + a - b} / (a + b) + 2 * \cosh(d * x + c) + 2 * \sinh(d * x + c)) - (a + b) * \sqrt{-a * b / (a^2 + 2 * a * b + b^2)} * c^2 * \log(2 * \sqrt{(2 * (a + b) * \sqrt{-a * b / (a^2 + 2 * a * b + b^2)}) - a + b} / (a + b) + 2 * \cosh(d * x + c) + 2 * \sinh(d * x + c)) - (a + b) * \sqrt{-a * b / (a^2 + 2 * a * b + b^2)} * c^2 * \log(-2 * \sqrt{(2 * (a + b) * \sqrt{-a * b / (a^2 + 2 * a * b + b^2)}) - a + b} / (a + b) + 2 * \cosh(d * x + c) + 2 * \sinh(d * x + c)) + ((a + b) * d^2 * x^2 - (a + b) * c^2) * \sqrt{-a * b / (a^2 + 2 * a * b + b^2)} * \log(((a - b) * \cosh(d * x + c) + (a - b) * \sinh(d * x + c) - 2 * ((a + b) * \cosh(d * x + c) + (a + b) * \sinh(d * x + c))) * \sqrt{-a * b / (a^2 + 2 * a * b + b^2)}) * \sqrt{-(2 * (a + b) * \sqrt{-a * b / (a^2 + 2 * a * b + b^2)}) + a - b} / (a + b) + a + b) / (a + b) + ((a + b) * d^2 * x^2 - (a + b) * c^2) * \sqrt{-a * b / (a^2 + 2 * a * b + b^2)} * \log(-(((a - b) * \cosh(d * x + c) + (a - b) * \sinh(d * x + c) - 2 * ((a + b) * \cosh(d * x + c) + (a + b) * \sinh(d * x + c))) * \sqrt{-a * b / (a^2 + 2 * a * b + b^2)}) * \sqrt{-(2 * (a + b) * \sqrt{-a * b / (a^2 + 2 * a * b + b^2)}) + a - b} / (a + b) - a - b) / (a + b) - ((a + b) * d^2 * x^2 - (a + b) * c^2) * \sqrt{-a * b / (a^2 + 2 * a * b + b^2)} * \log(((a - b) * \cosh(d * x + c) + (a - b) * \sinh(d * x + c) + 2 * ((a + b) * \cosh(d * x + c) + (a + b) * \sinh(d * x + c))) * \sqrt{-a * b / (a^2 + 2 * a * b + b^2)}) * \sqrt{(2 * (a + b) * \sqrt{-a * b / (a^2 + 2 * a * b + b^2)}) - a + b} / (a + b) +$$

```

a + b)/(a + b)) - ((a + b)*d^2*x^2 - (a + b)*c^2)*sqrt(-a*b/(a^2 + 2*a*b +
b^2))*log(-(((a - b)*cosh(d*x + c) + (a - b)*sinh(d*x + c) + 2*((a + b)*cos
h(d*x + c) + (a + b)*sinh(d*x + c))*sqrt(-a*b/(a^2 + 2*a*b + b^2))))*sqrt((2
*(a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2)) - a + b)/(a + b)) - a - b)/(a + b))
- 2*(a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2))*polylog(3, ((a - b)*cosh(d*x +
c) + (a - b)*sinh(d*x + c) - 2*((a + b)*cosh(d*x + c) + (a + b)*sinh(d*x +
c))*sqrt(-a*b/(a^2 + 2*a*b + b^2)))*sqrt(-(2*(a + b)*sqrt(-a*b/(a^2 + 2*a*b
+ b^2)) + a - b)/(a + b))/(a + b)) - 2*(a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^
2))*polylog(3, -((a - b)*cosh(d*x + c) + (a - b)*sinh(d*x + c) - 2*((a + b)
*cosh(d*x + c) + (a + b)*sinh(d*x + c))*sqrt(-a*b/(a^2 + 2*a*b + b^2)))*sqr
t(-(2*(a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2)) + a - b)/(a + b))/(a + b)) + 2
*(a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2))*polylog(3, ((a - b)*cosh(d*x + c) +
(a - b)*sinh(d*x + c) + 2*((a + b)*cosh(d*x + c) + (a + b)*sinh(d*x + c))*
sqrt(-a*b/(a^2 + 2*a*b + b^2)))*sqrt((2*(a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^
2)) - a + b)/(a + b))/(a + b)) + 2*(a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2))*p
olylog(3, -((a - b)*cosh(d*x + c) + (a - b)*sinh(d*x + c) + 2*((a + b)*cosh
(d*x + c) + (a + b)*sinh(d*x + c))*sqrt(-a*b/(a^2 + 2*a*b + b^2)))*sqrt((2*
(a + b)*sqrt(-a*b/(a^2 + 2*a*b + b^2)) - a + b)/(a + b))/(a + b)))/(a*b*d^3
)

```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x^2 \operatorname{sech}^2(c + dx)}{a + b \tanh^2(c + dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x**2*sech(d*x+c)**2/(a+b*tanh(d*x+c)**2), x)

[Out] Integral(x**2*sech(c + d*x)**2/(a + b*tanh(c + d*x)**2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2*sech(d*x+c)^2/(a+b*tanh(d*x+c)^2), x, algorithm="giac")

[Out] integrate(x^2*sech(d*x + c)^2/(b*tanh(d*x + c)^2 + a), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{x^2}{\cosh(c + dx)^2 (b \tanh(c + dx)^2 + a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(x^2/(cosh(c + d*x)^2*(a + b*tanh(c + d*x)^2)),x)
```

```
[Out] int(x^2/(cosh(c + d*x)^2*(a + b*tanh(c + d*x)^2)), x)
```

3.146 $\int x^3 \tanh(a + 2 \log(x)) dx$

Optimal. Leaf size=29

$$\frac{x^4}{4} - \frac{1}{2}e^{-2a} \log(1 + e^{2a}x^4)$$

[Out] 1/4*x^4-1/2*ln(1+exp(2*a)*x^4)/exp(2*a)

Rubi [A]

time = 0.03, antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {5656, 455, 45}

$$\frac{x^4}{4} - \frac{1}{2}e^{-2a} \log(e^{2a}x^4 + 1)$$

Antiderivative was successfully verified.

[In] Int[x^3*Tanh[a + 2*Log[x]],x]

[Out] x^4/4 - Log[1 + E^(2*a)*x^4]/(2*E^(2*a))

Rule 45

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n},
x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && Le
Q[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])
```

Rule 455

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.
), x_Symbol] := Dist[1/n, Subst[Int[(a + b*x)^p*(c + d*x)^q, x], x, x^n], x
] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[b*c - a*d, 0] && EqQ[m - n +
1, 0]
```

Rule 5656

```
Int[((e_.)*(x_))^(m_.)*Tanh[((a_.) + Log[x_]*(b_.))*(d_.)]^(p_.), x_Symbol]
:= Int[(e*x)^m*((-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p),
x] /; FreeQ[{a, b, d, e, m, p}, x]
```

Rubi steps

$$\int x^3 \tanh(a + 2 \log(x)) dx = \int x^3 \tanh(a + 2 \log(x)) dx$$

Mathematica [B] Leaf count is larger than twice the leaf count of optimal. 64 vs. 2(29) = 58.

time = 0.02, size = 64, normalized size = 2.21

$$\frac{x^4}{4} - \frac{1}{2} \cosh(2a) \log(\cosh(a) + x^4 \cosh(a) - \sinh(a) + x^4 \sinh(a)) + \frac{1}{2} \log(\cosh(a) + x^4 \cosh(a) - \sinh(a) + x^4 \sinh(a)) \sinh(2a)$$

Antiderivative was successfully verified.

[In] Integrate[x^3*Tanh[a + 2*Log[x]],x]

[Out] x^4/4 - (Cosh[2*a]*Log[Cosh[a] + x^4*Cosh[a] - Sinh[a] + x^4*Sinh[a]])/2 + (Log[Cosh[a] + x^4*Cosh[a] - Sinh[a] + x^4*Sinh[a]]*Sinh[2*a])/2

Maple [A]

time = 0.49, size = 24, normalized size = 0.83

method	result	size
risch	$\frac{x^4}{4} - \frac{e^{-2a} \ln(1+e^{2a}x^4)}{2}$	24

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^3*tanh(a+2*ln(x)),x,method=_RETURNVERBOSE)

[Out] 1/4*x^4-1/2*exp(-2*a)*ln(1+exp(2*a)*x^4)

Maxima [A]

time = 0.28, size = 23, normalized size = 0.79

$$\frac{1}{4} x^4 - \frac{1}{2} e^{(-2a)} \log(x^4 e^{(2a)} + 1)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^3*tanh(a+2*log(x)),x, algorithm="maxima")

[Out] 1/4*x^4 - 1/2*e^(-2*a)*log(x^4*e^(2*a) + 1)

Fricas [A]

time = 0.34, size = 28, normalized size = 0.97

$$\frac{1}{4} (x^4 e^{(2a)} - 2 \log(x^4 e^{(2a)} + 1)) e^{(-2a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^3*tanh(a+2*log(x)),x, algorithm="fricas")

[Out] 1/4*(x^4*e^(2*a) - 2*log(x^4*e^(2*a) + 1))*e^(-2*a)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x^3 \tanh(a + 2 \log(x)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x**3*tanh(a+2*ln(x)),x)

[Out] Integral(x**3*tanh(a + 2*log(x)), x)

Giac [A]

time = 0.40, size = 23, normalized size = 0.79

$$\frac{1}{4} x^4 - \frac{1}{2} e^{(-2a)} \log(x^4 e^{(2a)} + 1)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^3*tanh(a+2*log(x)),x, algorithm="giac")

[Out] 1/4*x^4 - 1/2*e^(-2*a)*log(x^4*e^(2*a) + 1)

Mupad [B]

time = 1.08, size = 21, normalized size = 0.72

$$\frac{x^4}{4} - \frac{e^{-2a} \ln(x^4 + e^{-2a})}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^3*tanh(a + 2*log(x)),x)

[Out] x^4/4 - (exp(-2*a)*log(exp(-2*a) + x^4))/2

3.147 $\int x^2 \tanh(a + 2 \log(x)) dx$

Optimal. Leaf size=151

$$\frac{x^3}{3} + \frac{e^{-3a/2} \operatorname{ArcTan}\left(1 - \sqrt{2} e^{a/2} x\right)}{\sqrt{2}} - \frac{e^{-3a/2} \operatorname{ArcTan}\left(1 + \sqrt{2} e^{a/2} x\right)}{\sqrt{2}} - \frac{e^{-3a/2} \log\left(1 - \sqrt{2} e^{a/2} x + e^a x^2\right)}{2\sqrt{2}} + \frac{e^{-3a/2} \log\left(1 + \sqrt{2} e^{a/2} x + e^a x^2\right)}{2\sqrt{2}}$$

[Out] 1/3*x^3-1/2*arctan(-1+exp(1/2*a)*x*2^(1/2))/exp(3/2*a)*2^(1/2)-1/2*arctan(1+exp(1/2*a)*x*2^(1/2))/exp(3/2*a)*2^(1/2)-1/4*ln(1+exp(a)*x^2-exp(1/2*a)*x*2^(1/2))/exp(3/2*a)*2^(1/2)+1/4*ln(1+exp(a)*x^2+exp(1/2*a)*x*2^(1/2))/exp(3/2*a)*2^(1/2)

Rubi [A]

time = 0.09, antiderivative size = 151, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 8, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.727$, Rules used = {5656, 470, 303, 1176, 631, 210, 1179, 642}

$$\frac{e^{-3a/2} \operatorname{ArcTan}\left(1 - \sqrt{2} e^{a/2} x\right)}{\sqrt{2}} - \frac{e^{-3a/2} \operatorname{ArcTan}\left(\sqrt{2} e^{a/2} x + 1\right)}{\sqrt{2}} - \frac{e^{-3a/2} \log\left(e^a x^2 - \sqrt{2} e^{a/2} x + 1\right)}{2\sqrt{2}} + \frac{e^{-3a/2} \log\left(e^a x^2 + \sqrt{2} e^{a/2} x + 1\right)}{2\sqrt{2}} + \frac{x^3}{3}$$

Antiderivative was successfully verified.

[In] Int[x^2*Tanh[a + 2*Log[x]],x]

[Out] x^3/3 + ArcTan[1 - Sqrt[2]*E^(a/2)*x]/(Sqrt[2]*E^((3*a)/2)) - ArcTan[1 + Sqrt[2]*E^(a/2)*x]/(Sqrt[2]*E^((3*a)/2)) - Log[1 - Sqrt[2]*E^(a/2)*x + E^a*x^2]/(2*Sqrt[2]*E^((3*a)/2)) + Log[1 + Sqrt[2]*E^(a/2)*x + E^a*x^2]/(2*Sqrt[2]*E^((3*a)/2))

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 303

Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))

Rule 470

Int[((e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[d*(e*x)^(m + 1)*((a + b*x^n)^(p + 1))/(b*e*(m + n*(p


```
+ 1) + 1))), x] - Dist[(a*d*(m + 1) - b*c*(m + n*(p + 1) + 1))/(b*(m + n*(p
+ 1) + 1)), Int[(e*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, e, m,
n, p}, x] && NeQ[b*c - a*d, 0] && NeQ[m + n*(p + 1) + 1, 0]
```

Rule 631

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)
], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; Fre
eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 5656

```
Int[((e_)*(x_)^(m_))*Tanh[((a_) + Log[x_]*(b_))*(d_)]^(p_), x_Symbol]
:= Int[(e*x)^m*((-1 + E^(2*a*d))*x^(2*b*d))^p/(1 + E^(2*a*d))*x^(2*b*d))^p),
x] /; FreeQ[{a, b, d, e, m, p}, x]
```

Rubi steps

$$\int x^2 \tanh(a + 2 \log(x)) dx = \int x^2 \tanh(a + 2 \log(x)) dx$$

Mathematica [C] Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

time = 8.66, size = 64, normalized size = 0.42

$$\frac{1}{6} \left(2x^3 + 3\text{RootSum} \left[\cosh(a) - \sinh(a) + \cosh(a)\#1^4 + \sinh(a)\#1^4 \&, \frac{\log(x) - \log(x - \#1)}{\#1} \& \right] (\cosh(2a) - \sinh(2a)) \right)$$

Antiderivative was successfully verified.

[In] Integrate[x^2*Tanh[a + 2*Log[x]],x]

[Out] (2*x^3 + 3*RootSum[Cosh[a] - Sinh[a] + Cosh[a]*#1^4 + Sinh[a]*#1^4 & , (Log[x] - Log[x - #1])/#1 &]*(Cosh[2*a] - Sinh[2*a]))/6

Maple [C] Result contains higher order function than in optimal. Order 9 vs. order 3.
time = 0.62, size = 37, normalized size = 0.25

method	result	size
risch	$\frac{x^3}{3} - \frac{e^{-2a} \left(\sum_{-R=\text{RootOf}(e^{2a}Z^4+1)} \frac{\ln(x-\frac{R}{e^{-2a}})}{-R} \right)}{2}$	37

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^2*tanh(a+2*ln(x)),x,method=_RETURNVERBOSE)

[Out] 1/3*x^3-1/2*exp(-2*a)*sum(1/_R*ln(x-_R),_R=RootOf(exp(2*a)*_Z^4+1))

Maxima [A]

time = 0.49, size = 128, normalized size = 0.85

$$\frac{1}{3}x^3 - \frac{1}{2}\sqrt{2} \arctan\left(\frac{1}{2}\sqrt{2}(2xe^a + \sqrt{2}e^{\frac{1}{2}a})e^{(-\frac{1}{2}a)}\right)e^{(-\frac{3}{2}a)} - \frac{1}{2}\sqrt{2} \arctan\left(\frac{1}{2}\sqrt{2}(2xe^a - \sqrt{2}e^{\frac{1}{2}a})e^{(-\frac{1}{2}a)}\right)e^{(-\frac{3}{2}a)} + \frac{1}{4}\sqrt{2}e^{(-\frac{3}{2}a)}\log(x^2e^a + \sqrt{2}xe^{\frac{1}{2}a} + 1) - \frac{1}{4}\sqrt{2}e^{(-\frac{3}{2}a)}\log(x^2e^a - \sqrt{2}xe^{\frac{1}{2}a} + 1)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2*tanh(a+2*log(x)),x, algorithm="maxima")

[Out] 1/3*x^3 - 1/2*sqrt(2)*arctan(1/2*sqrt(2)*(2*x*e^a + sqrt(2)*e^(1/2*a))*e^(-1/2*a))*e^(-3/2*a) - 1/2*sqrt(2)*arctan(1/2*sqrt(2)*(2*x*e^a - sqrt(2)*e^(1/2*a))*e^(-1/2*a))*e^(-3/2*a) + 1/4*sqrt(2)*e^(-3/2*a)*log(x^2*e^a + sqrt(2)*x*e^(1/2*a) + 1) - 1/4*sqrt(2)*e^(-3/2*a)*log(x^2*e^a - sqrt(2)*x*e^(1/2*a) + 1)

Fricas [A]

time = 0.35, size = 158, normalized size = 1.05

$$\frac{1}{3}x^3 + \sqrt{2} \arctan\left(-\sqrt{2}xe^{\frac{1}{2}a} + \sqrt{2}\sqrt{\sqrt{2}xe^{(-\frac{1}{2}a)} + x^2 + e^{(-a)}}e^{\frac{1}{2}a} - 1\right)e^{(-\frac{3}{2}a)} + \sqrt{2} \arctan\left(-\sqrt{2}xe^{\frac{1}{2}a} + \sqrt{2}\sqrt{-\sqrt{2}xe^{(-\frac{1}{2}a)} + x^2 + e^{(-a)}}e^{\frac{1}{2}a} + 1\right)e^{(-\frac{3}{2}a)} + \frac{1}{4}\sqrt{2}e^{(-\frac{3}{2}a)}\log(\sqrt{2}xe^{(-\frac{1}{2}a)} + x^2 + e^{(-a)}) - \frac{1}{4}\sqrt{2}e^{(-\frac{3}{2}a)}\log(-\sqrt{2}xe^{(-\frac{1}{2}a)} + x^2 + e^{(-a)})$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2*tanh(a+2*log(x)),x, algorithm="fricas")

[Out] $\frac{1}{3}x^3 + \sqrt{2}\arctan(-\sqrt{2})x e^{1/2a} + \sqrt{2}\sqrt{\sqrt{2})x e^{-1/2a} + x^2 + e^{-a}}e^{1/2a} - 1)e^{-3/2a} + \sqrt{2}\arctan(-\sqrt{2})x e^{1/2a} + \sqrt{2}\sqrt{-\sqrt{2})x e^{-1/2a} + x^2 + e^{-a}}e^{1/2a} + 1)e^{-3/2a} + \frac{1}{4}\sqrt{2}e^{-3/2a}\log(\sqrt{2})x e^{-1/2a} + x^2 + e^{-a}) - \frac{1}{4}\sqrt{2}e^{-3/2a}\log(-\sqrt{2})x e^{-1/2a} + x^2 + e^{-a})$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x^2 \tanh(a + 2 \log(x)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x**2*tanh(a+2*ln(x)),x)`

[Out] `Integral(x**2*tanh(a + 2*log(x)), x)`

Giac [A]

time = 0.40, size = 123, normalized size = 0.81

$$\frac{1}{3}x^3 - \frac{1}{2}\sqrt{2}\arctan\left(\frac{1}{2}\sqrt{2}(\sqrt{2}e^{-1/2a} + 2x)e^{1/2a}\right)e^{-3/2a} - \frac{1}{2}\sqrt{2}\arctan\left(-\frac{1}{2}\sqrt{2}(\sqrt{2}e^{-1/2a} - 2x)e^{1/2a}\right)e^{-3/2a} + \frac{1}{4}\sqrt{2}e^{-3/2a}\log(\sqrt{2}xe^{-1/2a} + x^2 + e^{-a}) - \frac{1}{4}\sqrt{2}e^{-3/2a}\log(-\sqrt{2}xe^{-1/2a} + x^2 + e^{-a})$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^2*tanh(a+2*log(x)),x, algorithm="giac")`

[Out] $\frac{1}{3}x^3 - \frac{1}{2}\sqrt{2}\arctan(\frac{1}{2}\sqrt{2})(\sqrt{2})e^{-1/2a} + 2x)e^{1/2a})e^{-3/2a} - \frac{1}{2}\sqrt{2}\arctan(-\frac{1}{2}\sqrt{2})(\sqrt{2})e^{-1/2a} - 2x)e^{1/2a})e^{-3/2a} + \frac{1}{4}\sqrt{2}e^{-3/2a}\log(\sqrt{2})x e^{-1/2a} + x^2 + e^{-a}) - \frac{1}{4}\sqrt{2}e^{-3/2a}\log(-\sqrt{2})x e^{-1/2a} + x^2 + e^{-a})$

Mupad [B]

time = 1.10, size = 47, normalized size = 0.31

$$\frac{\operatorname{atan}\left(x(-e^{2a})^{1/4}\right)}{(-e^{2a})^{3/4}} - \frac{\operatorname{atanh}\left(x(-e^{2a})^{1/4}\right)}{(-e^{2a})^{3/4}} + \frac{x^3}{3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x^2*tanh(a + 2*log(x)),x)`

[Out] $\operatorname{atan}(x(-\exp(2a))^{1/4})/(-\exp(2a))^{3/4} - \operatorname{atanh}(x(-\exp(2a))^{1/4})/(-\exp(2a))^{3/4} + x^3/3$

3.148 $\int x \tanh(a + 2 \log(x)) dx$

Optimal. Leaf size=23

$$\frac{x^2}{2} - e^{-a} \operatorname{ArcTan}(e^a x^2)$$

[Out] 1/2*x^2-arctan(exp(a)*x^2)/exp(a)

Rubi [A]

time = 0.02, antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {5656, 470, 281, 209}

$$\frac{x^2}{2} - e^{-a} \operatorname{ArcTan}(e^a x^2)$$

Antiderivative was successfully verified.

[In] Int[x*Tanh[a + 2*Log[x]],x]

[Out] x^2/2 - ArcTan[E^a*x^2]/E^a

Rule 209

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 281

Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Dist[1/k, Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1 /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]

Rule 470

Int[((e_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Simp[d*(e*x)^(m + 1)*((a + b*x^n)^(p + 1)/(b*e*(m + n*(p + 1) + 1))), x] - Dist[(a*d*(m + 1) - b*c*(m + n*(p + 1) + 1))/(b*(m + n*(p + 1) + 1)), Int[(e*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && NeQ[b*c - a*d, 0] && NeQ[m + n*(p + 1) + 1, 0]

Rule 5656

Int[((e_)*(x_))^(m_)*Tanh[((a_) + Log[x_]*(b_))*(d_)]^(p_), x_Symbol] := Int[(e*x)^m*((-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d)))^p, x] /; FreeQ[{a, b, d, e, m, p}, x]

Rubi steps

$$\int x \tanh(a + 2 \log(x)) dx = \int x \tanh(a + 2 \log(x)) dx$$

Mathematica [A]

time = 0.15, size = 35, normalized size = 1.52

$$\frac{x^2}{2} - \text{ArcTan}(x^2(\cosh(a) + \sinh(a))) \cosh(a) + \text{ArcTan}(x^2(\cosh(a) + \sinh(a))) \sinh(a)$$

Antiderivative was successfully verified.

[In] Integrate[x*Tanh[a + 2*Log[x]],x]

[Out] x^2/2 - ArcTan[x^2*(Cosh[a] + Sinh[a])*Cosh[a] + ArcTan[x^2*(Cosh[a] + Sinh[a])*Sinh[a]]*Sinh[a]

Maple [C] Result contains complex when optimal does not.

time = 0.62, size = 41, normalized size = 1.78

method	result	size
risch	$\frac{x^2}{2} + \frac{ie^{-a} \ln(e^a x^2 - i)}{2} - \frac{ie^{-a} \ln(e^a x^2 + i)}{2}$	41

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x*tanh(a+2*ln(x)),x,method=_RETURNVERBOSE)

[Out] 1/2*x^2+1/2*I*exp(-a)*ln(exp(a)*x^2-I)-1/2*I*exp(-a)*ln(exp(a)*x^2+I)

Maxima [A]

time = 0.47, size = 19, normalized size = 0.83

$$\frac{1}{2} x^2 - \arctan(x^2 e^a) e^{(-a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x*tanh(a+2*log(x)),x, algorithm="maxima")

[Out] 1/2*x^2 - arctan(x^2*e^a)*e^(-a)

Fricas [A]

time = 0.34, size = 22, normalized size = 0.96

$$\frac{1}{2} (x^2 e^a - 2 \arctan(x^2 e^a)) e^{(-a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x*tanh(a+2*log(x)),x, algorithm="fricas")`

[Out] $1/2*(x^2*e^a - 2*\arctan(x^2*e^a))*e^{-a}$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x \tanh(a + 2 \log(x)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x*tanh(a+2*ln(x)),x)`

[Out] `Integral(x*tanh(a + 2*log(x)), x)`

Giac [A]

time = 0.41, size = 19, normalized size = 0.83

$$\frac{1}{2}x^2 - \arctan(x^2 e^a) e^{-a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x*tanh(a+2*log(x)),x, algorithm="giac")`

[Out] $1/2*x^2 - \arctan(x^2*e^a)*e^{-a}$

Mupad [B]

time = 1.07, size = 25, normalized size = 1.09

$$\frac{x^2}{2} - \frac{\operatorname{atan}\left(x^2 \sqrt{e^{2a}}\right)}{\sqrt{e^{2a}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x*tanh(a + 2*log(x)),x)`

[Out] $x^2/2 - \operatorname{atan}(x^2*\exp(2*a)^{(1/2)})/\exp(2*a)^{(1/2)}$

3.149 $\int \tanh(a + 2 \log(x)) dx$

Optimal. Leaf size=145

$$x + \frac{e^{-a/2} \operatorname{ArcTan}\left(1 - \sqrt{2} e^{a/2} x\right)}{\sqrt{2}} - \frac{e^{-a/2} \operatorname{ArcTan}\left(1 + \sqrt{2} e^{a/2} x\right)}{\sqrt{2}} + \frac{e^{-a/2} \log\left(1 - \sqrt{2} e^{a/2} x + e^a x^2\right)}{2\sqrt{2}} - \frac{e^{-a/2} \log\left(1 + \sqrt{2} e^{a/2} x + e^a x^2\right)}{2\sqrt{2}}$$

[Out] $x - 1/2 \cdot \arctan(-1 + \exp(1/2 \cdot a) \cdot x \cdot 2^{(1/2)}) / \exp(1/2 \cdot a) \cdot 2^{(1/2)} - 1/2 \cdot \arctan(1 + \exp(1/2 \cdot a) \cdot x \cdot 2^{(1/2)}) / \exp(1/2 \cdot a) \cdot 2^{(1/2)} + 1/4 \cdot \ln(1 + \exp(a) \cdot x^2 - \exp(1/2 \cdot a) \cdot x \cdot 2^{(1/2)}) / \exp(1/2 \cdot a) \cdot 2^{(1/2)} - 1/4 \cdot \ln(1 + \exp(a) \cdot x^2 + \exp(1/2 \cdot a) \cdot x \cdot 2^{(1/2)}) / \exp(1/2 \cdot a) \cdot 2^{(1/2)}$

Rubi [A]

time = 0.07, antiderivative size = 145, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 8, integrand size = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 1.143$, Rules used = {5652, 396, 217, 1179, 642, 1176, 631, 210}

$$\frac{e^{-a/2} \operatorname{ArcTan}\left(1 - \sqrt{2} e^{a/2} x\right)}{\sqrt{2}} - \frac{e^{-a/2} \operatorname{ArcTan}\left(\sqrt{2} e^{a/2} x + 1\right)}{\sqrt{2}} + \frac{e^{-a/2} \log\left(e^a x^2 - \sqrt{2} e^{a/2} x + 1\right)}{2\sqrt{2}} - \frac{e^{-a/2} \log\left(e^a x^2 + \sqrt{2} e^{a/2} x + 1\right)}{2\sqrt{2}} + x$$

Antiderivative was successfully verified.

[In] `Int[Tanh[a + 2*Log[x]], x]`

[Out] $x + \operatorname{ArcTan}\left[1 - \sqrt{2} \cdot E^{(a/2)} \cdot x\right] / \left(\sqrt{2} \cdot E^{(a/2)}\right) - \operatorname{ArcTan}\left[1 + \sqrt{2} \cdot E^{(a/2)} \cdot x\right] / \left(\sqrt{2} \cdot E^{(a/2)}\right) + \operatorname{Log}\left[1 - \sqrt{2} \cdot E^{(a/2)} \cdot x + E^a \cdot x^2\right] / \left(2 \cdot \sqrt{2} \cdot E^{(a/2)}\right) - \operatorname{Log}\left[1 + \sqrt{2} \cdot E^{(a/2)} \cdot x + E^a \cdot x^2\right] / \left(2 \cdot \sqrt{2} \cdot E^{(a/2)}\right)$

Rule 210

`Int[((a_) + (b_) * (x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2] * Rt[-b, 2])^(-1) * ArcTan[Rt[-b, 2] * (x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

Rule 217

`Int[((a_) + (b_) * (x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*r), Int[(r - s*x^2)/(a + b*x^4), x], x] + Dist[1/(2*r), Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`

Rule 396

`Int[((a_) + (b_) * (x_)^(n_))^(p_) * ((c_) + (d_) * (x_)^(n_)), x_Symbol] := Simp[d*x*((a + b*x^n)^(p + 1)/(b*(n*(p + 1) + 1))), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(b*(n*(p + 1) + 1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b,`

c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]

Rule 631

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 1176

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]

Rule 1179

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]

Rule 5652

Int[Tanh[((a_) + Log[x_]*(b_))*(d_)]^(p_), x_Symbol] := Int[(-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p, x] /; FreeQ[{a, b, d, p}, x]

Rubi steps

$$\int \tanh(a + 2 \log(x)) dx = \int \tanh(a + 2 \log(x)) dx$$

Mathematica [C] Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

time = 0.13, size = 58, normalized size = 0.40

$$x + \frac{1}{2} \text{RootSum} \left[\cosh(a) - \sinh(a) + \cosh(a)\#1^4 + \sinh(a)\#1^4 \&, \frac{\log(x) - \log(x - \#1)}{\#1^3} \& \right] (\cosh(2a) - \sinh(2a))$$

Antiderivative was successfully verified.

[In] Integrate[Tanh[a + 2*Log[x]],x]

[Out] $x + (\text{RootSum}[\text{Cosh}[a] - \text{Sinh}[a] + \text{Cosh}[a]*\#1^4 + \text{Sinh}[a]*\#1^4 \& , (\text{Log}[x] - \text{Log}[x - \#1])/\#1^3 \&]*(\text{Cosh}[2*a] - \text{Sinh}[2*a]))/2$

Maple [C] Result contains higher order function than in optimal. Order 9 vs. order 3.
time = 0.62, size = 33, normalized size = 0.23

method	result	size
risch	$x - \frac{e^{-2a} \left(\sum_{-R=\text{RootOf}(e^{2a}Z^4+1)} \frac{\ln(x-\frac{R}{2})}{-R^3} \right)}{2}$	33

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(a+2*ln(x)),x,method=_RETURNVERBOSE)

[Out] $x - 1/2 * \exp(-2*a) * \text{sum}(1/_R^3 * \ln(x - _R), _R = \text{RootOf}(\exp(2*a) * _Z^4 + 1))$

Maxima [A]

time = 0.50, size = 124, normalized size = 0.86

$$-\frac{1}{2}\sqrt{2}\arctan\left(\frac{1}{2}\sqrt{2}(2xe^a + \sqrt{2}e^{\frac{1}{2}a})e^{(-\frac{1}{2}a)} - \frac{1}{2}\sqrt{2}\arctan\left(\frac{1}{2}\sqrt{2}(2xe^a - \sqrt{2}e^{\frac{1}{2}a})e^{(-\frac{1}{2}a)} - \frac{1}{4}\sqrt{2}e^{(-\frac{1}{2}a)}\log(x^2e^a + \sqrt{2}xe^{\frac{1}{2}a} + 1) + \frac{1}{4}\sqrt{2}e^{(-\frac{1}{2}a)}\log(x^2e^a - \sqrt{2}xe^{\frac{1}{2}a} + 1) + x\right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+2*log(x)),x, algorithm="maxima")

[Out] $-1/2*\text{sqrt}(2)*\arctan(1/2*\text{sqrt}(2)*(2*x*e^a + \text{sqrt}(2)*e^{(1/2*a)})*e^{(-1/2*a)})*e^{(-1/2*a)} - 1/2*\text{sqrt}(2)*\arctan(1/2*\text{sqrt}(2)*(2*x*e^a - \text{sqrt}(2)*e^{(1/2*a)})*e^{(-1/2*a)})*e^{(-1/2*a)} - 1/4*\text{sqrt}(2)*e^{(-1/2*a)}*\log(x^2*e^a + \text{sqrt}(2)*x*e^{(1/2*a)} + 1) + 1/4*\text{sqrt}(2)*e^{(-1/2*a)}*\log(x^2*e^a - \text{sqrt}(2)*x*e^{(1/2*a)} + 1) + x$

Fricas [A]

time = 0.37, size = 154, normalized size = 1.06

$$\sqrt{2}\arctan\left(-\sqrt{2}xe^{\frac{1}{2}a} + \sqrt{2}\sqrt{\sqrt{2}xe^{(-\frac{1}{2}a)} + x^2 + e^{(-a)}}e^{\frac{1}{2}a} - 1\right)e^{(-\frac{1}{2}a)} + \sqrt{2}\arctan\left(-\sqrt{2}xe^{\frac{1}{2}a} + \sqrt{2}\sqrt{-\sqrt{2}xe^{(-\frac{1}{2}a)} + x^2 + e^{(-a)}}e^{\frac{1}{2}a} + 1\right)e^{(-\frac{1}{2}a)} - \frac{1}{4}\sqrt{2}e^{(-\frac{1}{2}a)}\log(\sqrt{2}xe^{(-\frac{1}{2}a)} + x^2 + e^{(-a)}) + \frac{1}{4}\sqrt{2}e^{(-\frac{1}{2}a)}\log(-\sqrt{2}xe^{(-\frac{1}{2}a)} + x^2 + e^{(-a)}) + x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+2*log(x)),x, algorithm="fricas")

[Out] $\text{sqrt}(2)*\arctan(-\text{sqrt}(2)*x*e^{(1/2*a)} + \text{sqrt}(2)*\text{sqrt}(\text{sqrt}(2)*x*e^{(-1/2*a)} + x^2 + e^{(-a)}))*e^{(1/2*a)} - 1)*e^{(-1/2*a)} + \text{sqrt}(2)*\arctan(-\text{sqrt}(2)*x*e^{(1/2*a)} + \text{sqrt}(2)*\text{sqrt}(-\text{sqrt}(2)*x*e^{(-1/2*a)} + x^2 + e^{(-a)}))*e^{(1/2*a)} + 1)*e^{(-1/2*a)} - 1/4*\text{sqrt}(2)*e^{(-1/2*a)}*\log(\text{sqrt}(2)*x*e^{(-1/2*a)} + x^2 + e^{(-a)}) + 1/4*\text{sqrt}(2)*e^{(-1/2*a)}*\log(-\text{sqrt}(2)*x*e^{(-1/2*a)} + x^2 + e^{(-a)}) + x$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \tanh(a + 2 \log(x)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+2*ln(x)),x)

[Out] Integral(tanh(a + 2*log(x)), x)

Giac [A]

time = 0.41, size = 119, normalized size = 0.82

$$-\frac{1}{2}\sqrt{2}\arctan\left(\frac{1}{2}\sqrt{2}(\sqrt{2}e^{(-\frac{1}{2}a)+2x})e^{(\frac{1}{2}a)}\right)e^{(-\frac{1}{2}a)} - \frac{1}{2}\sqrt{2}\arctan\left(-\frac{1}{2}\sqrt{2}(\sqrt{2}e^{(-\frac{1}{2}a)-2x})e^{(\frac{1}{2}a)}\right)e^{(-\frac{1}{2}a)} - \frac{1}{4}\sqrt{2}e^{(-\frac{1}{2}a)}\log(\sqrt{2}xe^{(-\frac{1}{2}a)}+x^2+e^{(-a)}) + \frac{1}{4}\sqrt{2}e^{(-\frac{1}{2}a)}\log(-\sqrt{2}xe^{(-\frac{1}{2}a)}+x^2+e^{(-a)}) + x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+2*log(x)),x, algorithm="giac")

[Out] -1/2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2)*e^(-1/2*a) + 2*x)*e^(1/2*a))*e^(-1/2*a) - 1/2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2)*e^(-1/2*a) - 2*x)*e^(1/2*a))*e^(-1/2*a) - 1/4*sqrt(2)*e^(-1/2*a)*log(sqrt(2)*x*e^(-1/2*a) + x^2 + e^(-a)) + 1/4*sqrt(2)*e^(-1/2*a)*log(-sqrt(2)*x*e^(-1/2*a) + x^2 + e^(-a)) + x

Mupad [B]

time = 1.07, size = 44, normalized size = 0.30

$$x - \frac{\operatorname{atan}\left(x(-e^{2a})^{1/4}\right)}{(-e^{2a})^{1/4}} - \frac{\operatorname{atanh}\left(x(-e^{2a})^{1/4}\right)}{(-e^{2a})^{1/4}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(a + 2*log(x)),x)

[Out] x - atan(x*(-exp(2*a))^(1/4))/(-exp(2*a))^(1/4) - atanh(x*(-exp(2*a))^(1/4))/(-exp(2*a))^(1/4)

$$3.150 \quad \int \frac{\tanh(a+2\log(x))}{x} dx$$

Optimal. Leaf size=12

$$\frac{1}{2} \log(\cosh(a + 2 \log(x)))$$

[Out] 1/2*ln(cosh(a+2*ln(x)))

Rubi [A]

time = 0.01, antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {3556}

$$\frac{1}{2} \log(\cosh(a + 2 \log(x)))$$

Antiderivative was successfully verified.

[In] Int[Tanh[a + 2*Log[x]]/x,x]

[Out] Log[Cosh[a + 2*Log[x]]]/2

Rule 3556

Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] :> Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{\tanh(a + 2 \log(x))}{x} dx &= \text{Subst}\left(\int \tanh(a + 2x) dx, x, \log(x)\right) \\ &= \frac{1}{2} \log(\cosh(a + 2 \log(x))) \end{aligned}$$

Mathematica [A]

time = 0.02, size = 12, normalized size = 1.00

$$\frac{1}{2} \log(\cosh(a + 2 \log(x)))$$

Antiderivative was successfully verified.

[In] Integrate[Tanh[a + 2*Log[x]]/x,x]

[Out] Log[Cosh[a + 2*Log[x]]]/2

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 25 vs. $2(10) = 20$.

time = 0.68, size = 26, normalized size = 2.17

method	result	size
risch	$-\ln(x) + \frac{\ln(-e^{2a}x^4-1)}{2}$	20
derivativedivides	$-\frac{\ln(\tanh(a+2\ln(x))-1)}{4} - \frac{\ln(\tanh(a+2\ln(x))+1)}{4}$	26
default	$-\frac{\ln(\tanh(a+2\ln(x))-1)}{4} - \frac{\ln(\tanh(a+2\ln(x))+1)}{4}$	26

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tanh(a+2*ln(x))/x,x,method=_RETURNVERBOSE)`

[Out] $-1/4*\ln(\tanh(a+2*\ln(x))-1)-1/4*\ln(\tanh(a+2*\ln(x))+1)$

Maxima [A]

time = 0.26, size = 10, normalized size = 0.83

$$\frac{1}{2} \log(\cosh(a + 2 \log(x)))$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(a+2*log(x))/x,x, algorithm="maxima")`

[Out] $1/2*\log(\cosh(a + 2*\log(x)))$

Fricas [A]

time = 0.36, size = 18, normalized size = 1.50

$$\frac{1}{2} \log(x^4 e^{(2a)} + 1) - \log(x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(a+2*log(x))/x,x, algorithm="fricas")`

[Out] $1/2*\log(x^4*e^{(2*a)} + 1) - \log(x)$

Sympy [A]

time = 0.09, size = 15, normalized size = 1.25

$$\log(x) - \frac{\log(\tanh(a + 2 \log(x)) + 1)}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(a+2*ln(x))/x,x)`

[Out] $\log(x) - \log(\tanh(a + 2\log(x)) + 1)/2$

Giac [A]

time = 0.41, size = 20, normalized size = 1.67

$$\frac{1}{2} \log(x^4 e^{(2a)} + 1) - \frac{1}{4} \log(x^4)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(a+2*log(x))/x,x, algorithm="giac")`

[Out] $1/2*\log(x^4*e^{(2*a)} + 1) - 1/4*\log(x^4)$

Mupad [B]

time = 1.12, size = 15, normalized size = 1.25

$$\ln(x) - \frac{\ln(\tanh(a + 2 \ln(x)) + 1)}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tanh(a + 2*log(x))/x,x)`

[Out] $\log(x) - \log(\tanh(a + 2\log(x)) + 1)/2$

$$3.151 \quad \int \frac{\tanh(a+2 \log(x))}{x^2} dx$$

Optimal. Leaf size=147

$$\frac{1}{x} - \frac{e^{a/2} \operatorname{ArcTan}\left(1 - \sqrt{2} e^{a/2} x\right)}{\sqrt{2}} + \frac{e^{a/2} \operatorname{ArcTan}\left(1 + \sqrt{2} e^{a/2} x\right)}{\sqrt{2}} + \frac{e^{a/2} \log\left(1 - \sqrt{2} e^{a/2} x + e^a x^2\right)}{2\sqrt{2}} - \frac{e^{a/2} \log\left(1 + \sqrt{2} e^{a/2} x + e^a x^2\right)}{2\sqrt{2}}$$

[Out] 1/x+1/2*exp(1/2*a)*arctan(-1+exp(1/2*a)*x*2^(1/2))*2^(1/2)+1/2*exp(1/2*a)*arctan(1+exp(1/2*a)*x*2^(1/2))*2^(1/2)+1/4*exp(1/2*a)*ln(1+exp(a)*x^2-exp(1/2*a)*x*2^(1/2))*2^(1/2)-1/4*exp(1/2*a)*ln(1+exp(a)*x^2+exp(1/2*a)*x*2^(1/2))*2^(1/2)

Rubi [A]

time = 0.07, antiderivative size = 147, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 8, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.727$, Rules used = {5656, 464, 303, 1176, 631, 210, 1179, 642}

$$-\frac{e^{a/2} \operatorname{ArcTan}\left(1 - \sqrt{2} e^{a/2} x\right)}{\sqrt{2}} + \frac{e^{a/2} \operatorname{ArcTan}\left(\sqrt{2} e^{a/2} x + 1\right)}{\sqrt{2}} + \frac{e^{a/2} \log\left(e^a x^2 - \sqrt{2} e^{a/2} x + 1\right)}{2\sqrt{2}} - \frac{e^{a/2} \log\left(e^a x^2 + \sqrt{2} e^{a/2} x + 1\right)}{2\sqrt{2}} + \frac{1}{x}$$

Antiderivative was successfully verified.

[In] Int[Tanh[a + 2*Log[x]]/x^2,x]

[Out] x^(-1) - (E^(a/2)*ArcTan[1 - Sqrt[2]*E^(a/2)*x])/Sqrt[2] + (E^(a/2)*ArcTan[1 + Sqrt[2]*E^(a/2)*x])/Sqrt[2] + (E^(a/2)*Log[1 - Sqrt[2]*E^(a/2)*x + E^a*x^2])/(2*Sqrt[2]) - (E^(a/2)*Log[1 + Sqrt[2]*E^(a/2)*x + E^a*x^2])/(2*Sqrt[2])

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 303

Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))

Rule 464

Int[((e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[c*(e*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*e*(m + 1))),

```
x] + Dist[(a*d*(m + 1) - b*c*(m + n*(p + 1)))/(a*e^n*(m + 1)), Int[(e*x)^(m + n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, e, p}, x] && NeQ[b*c - a*d, 0] && (IntegerQ[n] || GtQ[e, 0]) && ((GtQ[n, 0] && LtQ[m, -1]) || (LtQ[n, 0] && GtQ[m + n, -1])) && !ILtQ[p, -1]
```

Rule 631

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 5656

```
Int[((e_)*(x_))^(m_)*Tanh[((a_) + Log[x_]*(b_))*(d_)]^(p_), x_Symbol] := Int[(e*x)^m*((-1 + E^(2*a*d))*x^(2*b*d))^p/(1 + E^(2*a*d))*x^(2*b*d))^p, x] /; FreeQ[{a, b, d, e, m, p}, x]
```

Rubi steps

$$\int \frac{\tanh(a + 2 \log(x))}{x^2} dx = \int \frac{\tanh(a + 2 \log(x))}{x^2} dx$$

Mathematica [C] Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

time = 0.13, size = 59, normalized size = 0.40

$$\frac{2 - x \operatorname{RootSum} \left[\cosh(a) + \sinh(a) + \cosh(a) \#1^4 - \sinh(a) \#1^4 \&, \frac{\log(x) + \log\left(\frac{1}{x} - \#1\right)}{\#1^3} \& \right] (\cosh(a) + \sinh(a))^2}{2x}$$

Antiderivative was successfully verified.

[In] Integrate[Tanh[a + 2*Log[x]]/x^2,x]

[Out] (2 - x*RootSum[Cosh[a] + Sinh[a] + Cosh[a]*#1^4 - Sinh[a]*#1^4 & , (Log[x] + Log[x^(-1) - #1])/#1^3 &]*(Cosh[a] + Sinh[a])^2)/(2*x)

Maple [C] Result contains higher order function than in optimal. Order 9 vs. order 3.

time = 0.61, size = 42, normalized size = 0.29

method	result	size
risch	$\frac{1}{x} + \frac{\sum_{-R=\operatorname{RootOf}(-Z^4+e^{2a})} -R \ln\left(\left(5-R^4+4e^{2a}\right)x - R^3\right)}{2}$	42

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(a+2*ln(x))/x^2,x,method=_RETURNVERBOSE)

[Out] 1/x+1/2*sum(_R*ln((5*_R^4+4*exp(2*a))*x-_R^3),_R=RootOf(_Z^4+exp(2*a)))

Maxima [A]

time = 0.48, size = 125, normalized size = 0.85

$$-\frac{1}{2}\sqrt{2}\arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2}e^{\frac{1}{2}a}+\frac{2}{x}\right)e^{\frac{1}{2}a}\right)e^{\frac{1}{2}a}-\frac{1}{2}\sqrt{2}\arctan\left(-\frac{1}{2}\sqrt{2}\left(\sqrt{2}e^{\frac{1}{2}a}-\frac{2}{x}\right)e^{\frac{1}{2}a}\right)e^{\frac{1}{2}a}-\frac{1}{4}\sqrt{2}e^{\frac{1}{2}a}\log\left(\frac{\sqrt{2}e^{\frac{1}{2}a}}{x}+\frac{1}{x^2}+e^a\right)+\frac{1}{4}\sqrt{2}e^{\frac{1}{2}a}\log\left(-\frac{\sqrt{2}e^{\frac{1}{2}a}}{x}+\frac{1}{x^2}+e^a\right)+\frac{1}{x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+2*log(x))/x^2,x, algorithm="maxima")

[Out] -1/2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2)*e^(1/2*a) + 2/x)*e^(-1/2*a))*e^(1/2*a) - 1/2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2)*e^(1/2*a) - 2/x)*e^(-1/2*a))*e^(1/2*a) - 1/4*sqrt(2)*e^(1/2*a)*log(sqrt(2)*e^(1/2*a)/x + 1/x^2 + e^a) + 1/4*sqrt(2)*e^(1/2*a)*log(-sqrt(2)*e^(1/2*a)/x + 1/x^2 + e^a) + 1/x

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 204 vs. 2(101) = 202.

time = 0.38, size = 204, normalized size = 1.39

$$\frac{4\sqrt{2}x\arctan\left(-\left(\sqrt{2}xe^{\frac{1}{2}a}-\sqrt{2}\sqrt{x^2e^{4a}+\sqrt{2}xe^{\frac{1}{2}a}+e^{2a}}\right)e^{\frac{1}{2}a}+4\sqrt{2}x\arctan\left(-\left(\sqrt{2}xe^{\frac{1}{2}a}-\sqrt{2}\sqrt{x^2e^{4a}-\sqrt{2}xe^{\frac{1}{2}a}+e^{2a}}\right)e^{\frac{1}{2}a}\right)e^{\frac{1}{2}a}+\sqrt{2}xe^{\frac{1}{2}a}\log\left(x^2e^{4a}+\sqrt{2}xe^{\frac{1}{2}a}+e^{2a}\right)-\sqrt{2}xe^{\frac{1}{2}a}\log\left(x^2e^{4a}-\sqrt{2}xe^{\frac{1}{2}a}+e^{2a}\right)-4}{4x}}{4x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+2*log(x))/x^2,x, algorithm="fricas")

[Out] $-1/4*(4*\sqrt{2}*x*\arctan(-(\sqrt{2})*x*e^{(5/2)*a} - \sqrt{2}*\sqrt{x^2*e^{(4)*a}} + \sqrt{2})*x*e^{(7/2)*a} + e^{(3*a)})*e^{(1/2)*a} + e^{(2*a)})*e^{(-2*a)})*e^{(1/2)*a} + 4*\sqrt{2}*x*\arctan(-(\sqrt{2})*x*e^{(5/2)*a} - \sqrt{2}*\sqrt{x^2*e^{(4)*a}} - \sqrt{2})*x*e^{(7/2)*a} + e^{(3*a)})*e^{(1/2)*a} - e^{(2*a)})*e^{(-2*a)})*e^{(1/2)*a} + \sqrt{2}*(2)*x*e^{(1/2)*a}*\log(x^2*e^{(4)*a} + \sqrt{2})*x*e^{(7/2)*a} + e^{(3*a)}) - \sqrt{2}*x*e^{(1/2)*a}*\log(x^2*e^{(4)*a} - \sqrt{2})*x*e^{(7/2)*a} + e^{(3*a)}) - 4)/x$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\tanh(a + 2 \log(x))}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+2*ln(x))/x**2,x)

[Out] Integral(tanh(a + 2*log(x))/x**2, x)

Giac [A]

time = 0.43, size = 121, normalized size = 0.82

$$\frac{1}{2}\sqrt{2} \arctan\left(\frac{1}{2}\sqrt{2}(\sqrt{2}e^{(-\frac{1}{2}a)} + 2x)e^{(\frac{1}{2}a)}\right)e^{(\frac{1}{2}a)} + \frac{1}{2}\sqrt{2} \arctan\left(-\frac{1}{2}\sqrt{2}(\sqrt{2}e^{(-\frac{1}{2}a)} - 2x)e^{(\frac{1}{2}a)}\right)e^{(\frac{1}{2}a)} - \frac{1}{4}\sqrt{2}e^{(\frac{1}{2}a)}\log(\sqrt{2}xe^{(-\frac{1}{2}a)} + x^2 + e^{(-a)}) + \frac{1}{4}\sqrt{2}e^{(\frac{1}{2}a)}\log(-\sqrt{2}xe^{(-\frac{1}{2}a)} + x^2 + e^{(-a)}) + \frac{1}{x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+2*log(x))/x^2,x, algorithm="giac")

[Out] $1/2*\sqrt{2}*\arctan(1/2*\sqrt{2}*(\sqrt{2}*e^{(-1/2*a)} + 2*x)*e^{(1/2*a)})*e^{(1/2*a)} + 1/2*\sqrt{2}*\arctan(-1/2*\sqrt{2}*(\sqrt{2}*e^{(-1/2*a)} - 2*x)*e^{(1/2*a)})*e^{(1/2*a)} - 1/4*\sqrt{2}*e^{(1/2*a)}*\log(\sqrt{2})*x*e^{(-1/2*a)} + x^2 + e^{(-a)}) + 1/4*\sqrt{2}*e^{(1/2*a)}*\log(-\sqrt{2})*x*e^{(-1/2*a)} + x^2 + e^{(-a)}) + 1/x$

Mupad [B]

time = 1.09, size = 45, normalized size = 0.31

$$\operatorname{atan}\left(x(-e^{2a})^{1/4}\right)(-e^{2a})^{1/4} - \operatorname{atanh}\left(x(-e^{2a})^{1/4}\right)(-e^{2a})^{1/4} + \frac{1}{x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(a + 2*log(x))/x^2,x)

[Out] $\operatorname{atan}(x*(-\exp(2*a))^{(1/4)})*(-\exp(2*a))^{(1/4)} - \operatorname{atanh}(x*(-\exp(2*a))^{(1/4)})*(-\exp(2*a))^{(1/4)} + 1/x$

$$3.152 \quad \int \frac{\tanh(a+2 \log(x))}{x^3} dx$$

Optimal. Leaf size=20

$$\frac{1}{2x^2} + e^a \operatorname{ArcTan}(e^a x^2)$$

[Out] 1/2/x^2+exp(a)*arctan(exp(a)*x^2)

Rubi [A]

time = 0.02, antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {5656, 464, 281, 209}

$$e^a \operatorname{ArcTan}(e^a x^2) + \frac{1}{2x^2}$$

Antiderivative was successfully verified.

[In] Int[Tanh[a + 2*Log[x]]/x^3,x]

[Out] 1/(2*x^2) + E^a*ArcTan[E^a*x^2]

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 281

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Dist[1/k, Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]

Rule 464

Int[((e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[c*(e*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*e*(m + 1))), x] + Dist[(a*d*(m + 1) - b*c*(m + n*(p + 1) + 1))/(a*e^n*(m + 1)), Int[(e*x)^(m + n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, e, p}, x] && NeQ[b*c - a*d, 0] && (IntegerQ[n] || GtQ[e, 0]) && ((GtQ[n, 0] && LtQ[m, -1]) || (LtQ[n, 0] && GtQ[m + n, -1])) && !ILtQ[p, -1]

Rule 5656

Int[((e_.)*(x_))^(m_.)*Tanh[((a_.) + Log[x]* (b_.))*(d_.)]^(p_.), x_Symbol] := Int[(e*x)^m*((-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p), x] /; FreeQ[{a, b, d, e, m, p}, x]

Rubi steps

$$\int \frac{\tanh(a + 2 \log(x))}{x^3} dx = \int \frac{\tanh(a + 2 \log(x))}{x^3} dx$$

Mathematica [A]

time = 0.11, size = 40, normalized size = 2.00

$$\frac{1}{2x^2} - \text{ArcTan}\left(\frac{\cosh(a) - \sinh(a)}{x^2}\right) \cosh(a) - \text{ArcTan}\left(\frac{\cosh(a) - \sinh(a)}{x^2}\right) \sinh(a)$$

Antiderivative was successfully verified.

`[In] Integrate[Tanh[a + 2*Log[x]]/x^3,x]``[Out] 1/(2*x^2) - ArcTan[(Cosh[a] - Sinh[a])/x^2]*Cosh[a] - ArcTan[(Cosh[a] - Sinh[a])/x^2]*Sinh[a]`**Maple [C]** Result contains higher order function than in optimal. Order 9 vs. order 3.

time = 0.62, size = 44, normalized size = 2.20

method	result	size
risch	$\frac{1}{2x^2} + \frac{\sum_{-R=\text{RootOf}(e^{2a}+Z^2)} -R \ln((4e^{2a}+5-R^2)x^2-R)}{2}$	44

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(tanh(a+2*ln(x))/x^3,x,method=_RETURNVERBOSE)``[Out] 1/2/x^2+1/2*sum(_R*ln((4*exp(2*a)+5*_R^2)*x^2-_R),_R=RootOf(exp(2*a)+_Z^2))`**Maxima [A]**

time = 0.47, size = 19, normalized size = 0.95

$$- \arctan\left(\frac{e^{(-a)}}{x^2}\right) e^a + \frac{1}{2x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(tanh(a+2*log(x))/x^3,x, algorithm="maxima")``[Out] -arctan(e^(-a)/x^2)*e^a + 1/2/x^2`**Fricas [A]**

time = 0.35, size = 21, normalized size = 1.05

$$\frac{2x^2 \arctan(x^2 e^a) e^a + 1}{2x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+2*log(x))/x^3,x, algorithm="fricas")

[Out] 1/2*(2*x^2*arctan(x^2*e^a)*e^a + 1)/x^2

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\tanh(a + 2 \log(x))}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+2*ln(x))/x**3,x)

[Out] Integral(tanh(a + 2*log(x))/x**3, x)

Giac [A]

time = 0.42, size = 16, normalized size = 0.80

$$\arctan(x^2 e^a) e^a + \frac{1}{2x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+2*log(x))/x^3,x, algorithm="giac")

[Out] arctan(x^2*e^a)*e^a + 1/2/x^2

Mupad [B]

time = 1.05, size = 24, normalized size = 1.20

$$\operatorname{atan}\left(x^2 \sqrt{e^{2a}}\right) \sqrt{e^{2a}} + \frac{1}{2x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(a + 2*log(x))/x^3,x)

[Out] atan(x^2*exp(2*a)^(1/2))*exp(2*a)^(1/2) + 1/(2*x^2)

3.153 $\int x^3 \tanh^2(a + 2 \log(x)) dx$

Optimal. Leaf size=47

$$\frac{x^4}{4} - \frac{e^{-2a}}{1 + e^{2a}x^4} - e^{-2a} \log(1 + e^{2a}x^4)$$

[Out] $1/4*x^4-1/\exp(2*a)/(1+\exp(2*a)*x^4)-\ln(1+\exp(2*a)*x^4)/\exp(2*a)$

Rubi [A]

time = 0.04, antiderivative size = 47, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {5656, 455, 45}

$$-\frac{e^{-2a}}{e^{2a}x^4 + 1} - e^{-2a} \log(e^{2a}x^4 + 1) + \frac{x^4}{4}$$

Antiderivative was successfully verified.

[In] $\text{Int}[x^3*\text{Tanh}[a + 2*\text{Log}[x]]^2, x]$

[Out] $x^4/4 - 1/(E^{(2*a)}*(1 + E^{(2*a)*x^4})) - \text{Log}[1 + E^{(2*a)*x^4}]/E^{(2*a)}$

Rule 45

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x], x] /; \text{FreeQ}\{a, b, c, d, n\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ (!\text{IntegerQ}[n] \ || \ (\text{EqQ}[c, 0] \ \&\& \ \text{LeQ}[7*m + 4*n + 4, 0]) \ || \ \text{LtQ}[9*m + 5*(n + 1), 0]) \ || \ \text{GtQ}[m + n + 2, 0])$

Rule 455

$\text{Int}[(x_.)^{(m_.)*((a_.) + (b_.)*(x_.)^{(n_.))^{(p_.)*((c_.) + (d_.)*(x_.)^{(n_.))^{(q_.)}, x_Symbol] \rightarrow \text{Dist}[1/n, \text{Subst}[\text{Int}[(a + b*x)^p*(c + d*x)^q, x], x, x^n], x] /; \text{FreeQ}\{a, b, c, d, m, n, p, q\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{EqQ}[m - n + 1, 0]$

Rule 5656

$\text{Int}[(e_.*(x_.))^{(m_.)*\text{Tanh}[(a_.) + \text{Log}[x_.]*(b_.)]*(d_.)]^{(p_.)}, x_Symbol] \rightarrow \text{Int}[(e*x)^m*((-1 + E^{(2*a*d)}*x^{(2*b*d)})^p/(1 + E^{(2*a*d)}*x^{(2*b*d)})^p), x] /; \text{FreeQ}\{a, b, d, e, m, p\}, x]$

Rubi steps

$$\int x^3 \tanh^2(a + 2 \log(x)) dx = \int x^3 \tanh^2(a + 2 \log(x)) dx$$

Mathematica [A]

time = 0.08, size = 86, normalized size = 1.83

$$\frac{x^4}{4} - \cosh(2a) \log((1+x^4) \cosh(a) + (-1+x^4) \sinh(a)) + \log((1+x^4) \cosh(a) + (-1+x^4) \sinh(a)) \sinh(2a) + \frac{-\cosh(3a) + \sinh(3a)}{(1+x^4) \cosh(a) + (-1+x^4) \sinh(a)}$$

Antiderivative was successfully verified.

`[In] Integrate[x^3*Tanh[a + 2*Log[x]]^2,x]`

```
[Out] x^4/4 - Cosh[2*a]*Log[(1 + x^4)*Cosh[a] + (-1 + x^4)*Sinh[a]] + Log[(1 + x^4)*Cosh[a] + (-1 + x^4)*Sinh[a]]*Sinh[2*a] + (-Cosh[3*a] + Sinh[3*a])/((1 + x^4)*Cosh[a] + (-1 + x^4)*Sinh[a])
```

Maple [A]

time = 0.38, size = 42, normalized size = 0.89

method	result	size
risch	$\frac{x^4}{4} - \frac{e^{-2a}}{1+e^{2a}x^4} - e^{-2a} \ln(1 + e^{2a}x^4)$	42

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(x^3*tanh(a+2*ln(x))^2,x,method=_RETURNVERBOSE)`

```
[Out] 1/4*x^4-exp(-2*a)/(1+exp(2*a)*x^4)-exp(-2*a)*ln(1+exp(2*a)*x^4)
```

Maxima [A]

time = 0.27, size = 40, normalized size = 0.85

$$\frac{1}{4} x^4 - e^{(-2a)} \log(x^4 e^{(2a)} + 1) - \frac{1}{x^4 e^{(4a)} + e^{(2a)}}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(x^3*tanh(a+2*log(x))^2,x, algorithm="maxima")`

```
[Out] 1/4*x^4 - e^(-2*a)*log(x^4*e^(2*a) + 1) - 1/(x^4*e^(4*a) + e^(2*a))
```

Fricas [A]

time = 0.35, size = 58, normalized size = 1.23

$$\frac{x^8 e^{(4a)} + x^4 e^{(2a)} - 4(x^4 e^{(2a)} + 1) \log(x^4 e^{(2a)} + 1) - 4}{4(x^4 e^{(4a)} + e^{(2a)})}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(x^3*tanh(a+2*log(x))^2,x, algorithm="fricas")`

```
[Out] 1/4*(x^8*e^(4*a) + x^4*e^(2*a) - 4*(x^4*e^(2*a) + 1)*log(x^4*e^(2*a) + 1) - 4)/(x^4*e^(4*a) + e^(2*a))
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x^3 \tanh^2(a + 2 \log(x)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x**3*tanh(a+2*ln(x))**2,x)**[Out]** Integral(x**3*tanh(a + 2*log(x))**2, x)**Giac [A]**

time = 0.41, size = 39, normalized size = 0.83

$$\frac{1}{4} x^4 + \frac{x^4}{x^4 e^{(2a)} + 1} - e^{(-2a)} \log(x^4 e^{(2a)} + 1)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^3*tanh(a+2*log(x))^2,x, algorithm="giac")**[Out]** 1/4*x^4 + x^4/(x^4*e^(2*a) + 1) - e^(-2*a)*log(x^4*e^(2*a) + 1)**Mupad [B]**

time = 1.12, size = 39, normalized size = 0.83

$$\frac{x^4}{4} - \frac{e^{-2a}}{e^{2a} x^4 + 1} - e^{-2a} \ln(x^4 + e^{-2a})$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^3*tanh(a + 2*log(x))^2,x)**[Out]** x^4/4 - exp(-2*a)/(x^4*exp(2*a) + 1) - exp(-2*a)*log(exp(-2*a) + x^4)

3.154 $\int x^2 \tanh^2(a + 2 \log(x)) dx$

Optimal. Leaf size=173

$$\frac{x^3}{3} + \frac{x^3}{1 + e^{2a}x^4} + \frac{3e^{-3a/2} \operatorname{ArcTan}(1 - \sqrt{2} e^{a/2}x)}{2\sqrt{2}} - \frac{3e^{-3a/2} \operatorname{ArcTan}(1 + \sqrt{2} e^{a/2}x)}{2\sqrt{2}} - \frac{3e^{-3a/2} \log(1 - \sqrt{2} e^{a/2}x)}{4\sqrt{2}}$$

[Out] $1/3*x^3+x^3/(1+\exp(2*a)*x^4)-3/4*\arctan(-1+\exp(1/2*a)*x*2^{(1/2)})/\exp(3/2*a)*2^{(1/2)}-3/4*\arctan(1+\exp(1/2*a)*x*2^{(1/2)})/\exp(3/2*a)*2^{(1/2)}-3/8*\ln(1+\exp(a)*x^2-\exp(1/2*a)*x*2^{(1/2)})/\exp(3/2*a)*2^{(1/2)}+3/8*\ln(1+\exp(a)*x^2+\exp(1/2*a)*x*2^{(1/2)})/\exp(3/2*a)*2^{(1/2)}$

Rubi [A]

time = 0.11, antiderivative size = 173, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 9, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.692$, Rules used = {5656, 474, 470, 303, 1176, 631, 210, 1179, 642}

$$\frac{3e^{-3a/2} \operatorname{ArcTan}(1 - \sqrt{2} e^{a/2}x)}{2\sqrt{2}} - \frac{3e^{-3a/2} \operatorname{ArcTan}(\sqrt{2} e^{a/2}x + 1)}{2\sqrt{2}} - \frac{3e^{-3a/2} \log(e^a x^2 - \sqrt{2} e^{a/2}x + 1)}{4\sqrt{2}} + \frac{3e^{-3a/2} \log(e^a x^2 + \sqrt{2} e^{a/2}x + 1)}{4\sqrt{2}} + \frac{x^3}{e^{2a}x^4 + 1} + \frac{x^3}{3}$$

Antiderivative was successfully verified.

[In] $\text{Int}[x^2*\text{Tanh}[a + 2*\text{Log}[x]]^2, x]$

[Out] $x^3/3 + x^3/(1 + E^{(2*a)*x^4}) + (3*\text{ArcTan}[1 - \text{Sqrt}[2]*E^{(a/2)*x}]/(2*\text{Sqrt}[2]*E^{((3*a)/2)}) - (3*\text{ArcTan}[1 + \text{Sqrt}[2]*E^{(a/2)*x}]/(2*\text{Sqrt}[2]*E^{((3*a)/2)})) - (3*\text{Log}[1 - \text{Sqrt}[2]*E^{(a/2)*x} + E^a*x^2]/(4*\text{Sqrt}[2]*E^{((3*a)/2)}) + (3*\text{Log}[1 + \text{Sqrt}[2]*E^{(a/2)*x} + E^a*x^2]/(4*\text{Sqrt}[2]*E^{((3*a)/2)}))$

Rule 210

$\text{Int}(((a_) + (b_)*(x_)^2)^{-1}, x_Symbol] := \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1})*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

Rule 303

$\text{Int}[(x_)^2/((a_) + (b_)*(x_)^4), x_Symbol] := \text{With}[\{r = \text{Numerator}[\text{Rt}[a/b, 2]], s = \text{Denominator}[\text{Rt}[a/b, 2]]\}, \text{Dist}[1/(2*s), \text{Int}[(r + s*x^2)/(a + b*x^4), x], x] - \text{Dist}[1/(2*s), \text{Int}[(r - s*x^2)/(a + b*x^4), x], x]] /; \text{FreeQ}\{a, b\}, x \ \&\& \ (\text{GtQ}[a/b, 0] \ || \ (\text{PosQ}[a/b] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, a]] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, b]]))$

Rule 470

$\text{Int}(((e_)*(x_))^{(m_)}*((a_) + (b_)*(x_)^{(n_)})^{(p_)}*((c_) + (d_)*(x_)^{(n_)}), x_Symbol] := \text{Simp}[d*(e*x)^{(m+1)}*((a + b*x^n)^{(p+1)})/(b*e*(m+n*(p$

+ 1) + 1))), x] - Dist[(a*d*(m + 1) - b*c*(m + n*(p + 1) + 1))/(b*(m + n*(p + 1) + 1)), Int[(e*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && NeQ[b*c - a*d, 0] && NeQ[m + n*(p + 1) + 1, 0]

Rule 474

Int[((e_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(2, x_Symbol] := Simp[(-b*c - a*d)^2*(e*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*b^2*e*n*(p + 1))), x] + Dist[1/(a*b^2*n*(p + 1)), Int[(e*x)^m*(a + b*x^n)^(p + 1)*Simp[(b*c - a*d)^2*(m + 1) + b^2*c^2*n*(p + 1) + a*b*d^2*n*(p + 1)*x^n, x], x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && LtQ[p, -1]

Rule 631

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 1176

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]

Rule 1179

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]

Rule 5656

Int[((e_)*(x_))^(m_)*Tanh[((a_) + Log[x_]*(b_))*(d_)]^(p_), x_Symbol] := Int[(e*x)^m*((-1 + E^(2*a*d))*x^(2*b*d))^p/(1 + E^(2*a*d))*x^(2*b*d))^p, x] /; FreeQ[{a, b, d, e, m, p}, x]

Rubi steps

$$\int x^2 \tanh^2(a + 2 \log(x)) dx = \int x^2 \tanh^2(a + 2 \log(x)) dx$$

Mathematica [A]

time = 0.53, size = 174, normalized size = 1.01

$$\frac{1}{12} \left(4x^3 + \frac{12x^3}{1+e^{2a}x^4} + 9(-1)^{3/4} e^{-3a/2} \log(\sqrt{-1} e^{-3a/2} - e^{-a}x) + 9\sqrt{-1} e^{-3a/2} \log((-1)^{3/4} e^{-3a/2} - e^{-a}x) - 9(-1)^{3/4} e^{-3a/2} \log(\sqrt{-1} e^{-3a/2} + e^{-a}x) - 9\sqrt{-1} e^{-3a/2} \log((-1)^{3/4} e^{-3a/2} + e^{-a}x) \right)$$

Antiderivative was successfully verified.

[In] Integrate[x^2*Tanh[a + 2*Log[x]]^2,x

[Out] $(4x^3 + (12x^3)/(1 + E^{(2a)}x^4) + (9(-1)^{(3/4)} \text{Log}[(-1)^{(1/4)}/E^{((3a)/2)} - x/E^a])/E^{((3a)/2)} + (9(-1)^{(1/4)} \text{Log}[(-1)^{(3/4)}/E^{((3a)/2)} - x/E^a])/E^{((3a)/2)} - (9(-1)^{(3/4)} \text{Log}[(-1)^{(1/4)}/E^{((3a)/2)} + x/E^a])/E^{((3a)/2)} - (9(-1)^{(1/4)} \text{Log}[(-1)^{(3/4)}/E^{((3a)/2)} + x/E^a])/E^{((3a)/2)})/12$

Maple [C] Result contains higher order function than in optimal. Order 9 vs. order 3.

time = 0.50, size = 53, normalized size = 0.31

method	result	size
risch	$\frac{x^3}{3} + \frac{x^3}{1+e^{2a}x^4} - \frac{3e^{-2a} \left(\sum_{-R=\text{RootOf}(e^{2a}Z^4+1)} \frac{\ln(x-R)}{-R} \right)}{4}$	53

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^2*tanh(a+2*ln(x))^2,x,method=_RETURNVERBOSE)

[Out] $1/3x^3 + x^3/(1+\exp(2a)x^4) - 3/4\exp(-2a)\text{sum}(1/_R \ln(x-_R), _R=\text{RootOf}(\exp(2a)Z^4+1))$

Maxima [A]

time = 0.47, size = 144, normalized size = 0.83

$$\frac{1}{3}x^3 - \frac{3}{4}\sqrt{2} \arctan\left(\frac{1}{2}\sqrt{2}(2xe^a + \sqrt{2}e^{(1/2)a})e^{(-1/2)a}\right)e^{(-3/2)a} - \frac{3}{4}\sqrt{2} \arctan\left(\frac{1}{2}\sqrt{2}(2xe^a - \sqrt{2}e^{(1/2)a})e^{(-1/2)a}\right)e^{(-3/2)a} + \frac{3}{8}\sqrt{2}e^{(-3/2)a} \log(x^2e^a + \sqrt{2}xe^{(1/2)a} + 1) - \frac{3}{8}\sqrt{2}e^{(-3/2)a} \log(x^2e^a - \sqrt{2}xe^{(1/2)a} + 1) + \frac{x^3}{x^4e^{(2a)} + 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2*tanh(a+2*log(x))^2,x, algorithm="maxima")

[Out] $1/3x^3 - 3/4\sqrt{2}\arctan(1/2\sqrt{2}(2xe^a + \sqrt{2}e^{(1/2)a})e^{(-1/2)a})e^{(-3/2)a} - 3/4\sqrt{2}\arctan(1/2\sqrt{2}(2xe^a - \sqrt{2}e^{(1/2)a})e^{(-1/2)a})e^{(-3/2)a} + 3/8\sqrt{2}e^{(-3/2)a} \log(x^2e^a + \sqrt{2}xe^{(1/2)a} + 1) - 3/8\sqrt{2}e^{(-3/2)a} \log(x^2e^a - \sqrt{2}xe^{(1/2)a} + 1) + \frac{x^3}{x^4e^{(2a)} + 1}$

) $x^2 e^{(1/2)a} + 1) - 3/8 \sqrt{2} e^{(-3/2)a} \log(x^2 e^a - \sqrt{2} x e^{(1/2)a} + 1) + x^3 / (x^4 e^{(2)a} + 1)$

Fricas [A]

time = 0.35, size = 231, normalized size = 1.34

$$\frac{8x^2 e^{2a} + 32x^3 + 36(\sqrt{2} x e^{(1/2)a} + \sqrt{2}) \arctan\left(\frac{-\sqrt{2} x e^{(1/2)a} + \sqrt{2} \sqrt{2 x e^{(1/2)a} + x^2 + e^{(1/2)a}}}{e^{(1/2)a} - 1}\right) e^{(-3/2)a} + 36(\sqrt{2} x e^{(1/2)a} + \sqrt{2}) \arctan\left(\frac{-\sqrt{2} x e^{(1/2)a} + \sqrt{2} \sqrt{-\sqrt{2} x e^{(1/2)a} + x^2 + e^{(1/2)a}}}{e^{(1/2)a} + 1}\right) e^{(-3/2)a} + 9(\sqrt{2} x e^{(1/2)a} + \sqrt{2}) e^{(-3/2)a} \log(\sqrt{2} x e^{(1/2)a} + x^2 + e^{(1/2)a}) - 9(\sqrt{2} x e^{(1/2)a} + \sqrt{2}) e^{(-3/2)a} \log(-\sqrt{2} x e^{(1/2)a} + x^2 + e^{(1/2)a})}{24(x^4 e^{(2)a} + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x²*tanh(a+2*log(x))²,x, algorithm="fricas")

[Out] $1/24*(8*x^7*e^{(2*a)} + 32*x^3 + 36*(\sqrt{2})*x^4*e^{(2*a)} + \sqrt{2})*\arctan(-\sqrt{2}*x*e^{(1/2*a)} + \sqrt{2})*\sqrt{2}*\sqrt{2*x*e^{(1/2*a)} + x^2 + e^{(-a)}}*e^{(1/2*a)} - 1)*e^{(-3/2*a)} + 36*(\sqrt{2})*x^4*e^{(2*a)} + \sqrt{2})*\arctan(-\sqrt{2}*x*e^{(1/2*a)} + \sqrt{2})*\sqrt{2}*\sqrt{-\sqrt{2}*x*e^{(1/2*a)} + x^2 + e^{(-a)}}*e^{(1/2*a)} + 1)*e^{(-3/2*a)} + 9*(\sqrt{2})*x^4*e^{(2*a)} + \sqrt{2})*e^{(-3/2*a)}*\log(\sqrt{2}*x*e^{(1/2*a)} + x^2 + e^{(-a)}) - 9*(\sqrt{2})*x^4*e^{(2*a)} + \sqrt{2})*e^{(-3/2*a)}*\log(-\sqrt{2}*x*e^{(1/2*a)} + x^2 + e^{(-a)})/(x^4*e^{(2*a)} + 1)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x^2 \tanh^2(a + 2 \log(x)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x**2*tanh(a+2*ln(x))**2,x)

[Out] Integral(x**2*tanh(a + 2*log(x))**2, x)

Giac [A]

time = 0.40, size = 139, normalized size = 0.80

$$\frac{1}{3}x^3 - \frac{3}{4}\sqrt{2} \arctan\left(\frac{1}{2}\sqrt{2}(\sqrt{2}e^{(-1/2)a} + 2x)e^{(1/2)a}\right)e^{(-3/2)a} - \frac{3}{4}\sqrt{2} \arctan\left(-\frac{1}{2}\sqrt{2}(\sqrt{2}e^{(-1/2)a} - 2x)e^{(1/2)a}\right)e^{(-3/2)a} + \frac{3}{8}\sqrt{2}e^{(-3/2)a} \log(\sqrt{2}xe^{(-1/2)a} + x^2 + e^{(-a)}) - \frac{3}{8}\sqrt{2}e^{(-3/2)a} \log(-\sqrt{2}xe^{(-1/2)a} + x^2 + e^{(-a)}) + \frac{x^3}{x^4e^{(2a)} + 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x²*tanh(a+2*log(x))²,x, algorithm="giac")

[Out] $1/3*x^3 - 3/4*\sqrt{2}*\arctan(1/2*\sqrt{2}*(\sqrt{2}*e^{(-1/2*a)} + 2*x))*e^{(1/2*a)}*e^{(-3/2*a)} - 3/4*\sqrt{2}*\arctan(-1/2*\sqrt{2}*(\sqrt{2}*e^{(-1/2*a)} - 2*x))*e^{(1/2*a)}*e^{(-3/2*a)} + 3/8*\sqrt{2}*e^{(-3/2*a)}*\log(\sqrt{2}*x*e^{(-1/2*a)} + x^2 + e^{(-a)}) - 3/8*\sqrt{2}*e^{(-3/2*a)}*\log(-\sqrt{2}*x*e^{(-1/2*a)} + x^2 + e^{(-a)}) + x^3/(x^4*e^{(2*a)} + 1)$

Mupad [B]

time = 1.12, size = 67, normalized size = 0.39

$$\frac{x^3}{e^{2a} x^4 + 1} + \frac{3 \operatorname{atan}\left(x(-e^{2a})^{1/4}\right)}{2(-e^{2a})^{3/4}} + \frac{x^3}{3} + \frac{\operatorname{atan}\left(x(-e^{2a})^{1/4} \operatorname{li}\right) 3i}{2(-e^{2a})^{3/4}}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(x^2*tanh(a + 2*log(x))^2,x)
```

```
[Out] x^3/(x^4*exp(2*a) + 1) + (3*atan(x*(-exp(2*a))^(1/4)))/(2*(-exp(2*a))^(3/4)) + (atan(x*(-exp(2*a))^(1/4)*1i)*3i)/(2*(-exp(2*a))^(3/4)) + x^3/3
```

3.155 $\int x \tanh^2(a + 2 \log(x)) dx$

Optimal. Leaf size=40

$$\frac{x^2}{2} + \frac{x^2}{1 + e^{2a}x^4} - e^{-a} \operatorname{ArcTan}(e^a x^2)$$

[Out] 1/2*x^2+x^2/(1+exp(2*a)*x^4)-arctan(exp(a)*x^2)/exp(a)

Rubi [A]

time = 0.03, antiderivative size = 40, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.454$, Rules used = {5656, 474, 470, 281, 209}

$$-e^{-a} \operatorname{ArcTan}(e^a x^2) + \frac{x^2}{e^{2a}x^4 + 1} + \frac{x^2}{2}$$

Antiderivative was successfully verified.

[In] Int[x*Tanh[a + 2*Log[x]]^2,x]

[Out] x^2/2 + x^2/(1 + E^(2*a)*x^4) - ArcTan[E^a*x^2]/E^a

Rule 209

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 281

Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Dist[1/k, Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]

Rule 470

Int[((e_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Simp[d*(e*x)^(m + 1)*((a + b*x^n)^(p + 1)/(b*e*(m + n*(p + 1) + 1))), x] - Dist[(a*d*(m + 1) - b*c*(m + n*(p + 1) + 1))/(b*(m + n*(p + 1) + 1)), Int[(e*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && NeQ[b*c - a*d, 0] && NeQ[m + n*(p + 1) + 1, 0]

Rule 474

Int[((e_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^2, x_Symbol] := Simp[(-(b*c - a*d)^2)*(e*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*b^2*e*n*(p + 1))), x] + Dist[1/(a*b^2*n*(p + 1)), Int[(e*x)^m*(a + b*x^n)

```
n)^(p + 1)*Simp[(b*c - a*d)^2*(m + 1) + b^2*c^2*n*(p + 1) + a*b*d^2*n*(p + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && LtQ[p, -1]
```

Rule 5656

```
Int[((e_.)*(x_))^(m_.)*Tanh[((a_.) + Log[x_]*(b_.))*(d_.)]^(p_.), x_Symbol]
  :-> Int[(e*x)^m*((-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p),
  x] /; FreeQ[{a, b, d, e, m, p}, x]
```

Rubi steps

$$\int x \tanh^2(a + 2 \log(x)) dx = \int x \tanh^2(a + 2 \log(x)) dx$$

Mathematica [A]

time = 0.27, size = 41, normalized size = 1.02

$$\frac{x^2}{2} + \frac{x^2}{1 + e^{2(a+2\log(x))}} - e^{-a} \operatorname{ArcTan}(e^a x^2)$$

Antiderivative was successfully verified.

```
[In] Integrate[x*Tanh[a + 2*Log[x]]^2,x]
```

```
[Out] x^2/2 + x^2/(1 + E^(2*(a + 2*Log[x]))) - ArcTan[E^a*x^2]/E^a
```

Maple [C] Result contains complex when optimal does not.

time = 0.51, size = 57, normalized size = 1.42

method	result	size
risch	$\frac{x^2}{2} + \frac{x^2}{1+e^{2a}x^4} + \frac{ie^{-a} \ln(e^a x^2 - i)}{2} - \frac{ie^{-a} \ln(e^a x^2 + i)}{2}$	57

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(x*tanh(a+2*ln(x))^2,x,method=_RETURNVERBOSE)
```

```
[Out] 1/2*x^2+x^2/(1+exp(2*a)*x^4)+1/2*I*exp(-a)*ln(exp(a)*x^2-I)-1/2*I*exp(-a)*ln(exp(a)*x^2+I)
```

Maxima [A]

time = 0.47, size = 35, normalized size = 0.88

$$\frac{1}{2} x^2 - \arctan(x^2 e^a) e^{(-a)} + \frac{x^2}{x^4 e^{(2a)} + 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x*tanh(a+2*log(x))^2,x, algorithm="maxima")

[Out] 1/2*x^2 - arctan(x^2*e^a)*e^(-a) + x^2/(x^4*e^(2*a) + 1)

Fricas [A]

time = 0.37, size = 50, normalized size = 1.25

$$\frac{x^6 e^{(3a)} + 3 x^2 e^a - 2 (x^4 e^{(2a)} + 1) \arctan(x^2 e^a)}{2 (x^4 e^{(3a)} + e^a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x*tanh(a+2*log(x))^2,x, algorithm="fricas")

[Out] 1/2*(x^6*e^(3*a) + 3*x^2*e^a - 2*(x^4*e^(2*a) + 1)*arctan(x^2*e^a))/(x^4*e^(3*a) + e^a)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x \tanh^2(a + 2 \log(x)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x*tanh(a+2*ln(x))**2,x)

[Out] Integral(x*tanh(a + 2*log(x))**2, x)

Giac [A]

time = 0.42, size = 35, normalized size = 0.88

$$\frac{1}{2} x^2 - \arctan(x^2 e^a) e^{(-a)} + \frac{x^2}{x^4 e^{(2a)} + 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x*tanh(a+2*log(x))^2,x, algorithm="giac")

[Out] 1/2*x^2 - arctan(x^2*e^a)*e^(-a) + x^2/(x^4*e^(2*a) + 1)

Mupad [B]

time = 1.05, size = 41, normalized size = 1.02

$$\frac{x^2}{e^{2a} x^4 + 1} - \frac{\operatorname{atan}\left(x^2 \sqrt{e^{2a}}\right)}{\sqrt{e^{2a}}} + \frac{x^2}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x*tanh(a + 2*log(x))^2,x)

[Out] x^2/(x^4*exp(2*a) + 1) - atan(x^2*exp(2*a)^(1/2))/exp(2*a)^(1/2) + x^2/2

3.156 $\int \tanh^2(a + 2 \log(x)) dx$

Optimal. Leaf size=165

$$x + \frac{x}{1 + e^{2a}x^4} + \frac{e^{-a/2} \operatorname{ArcTan}\left(1 - \sqrt{2} e^{a/2}x\right)}{2\sqrt{2}} - \frac{e^{-a/2} \operatorname{ArcTan}\left(1 + \sqrt{2} e^{a/2}x\right)}{2\sqrt{2}} + \frac{e^{-a/2} \log\left(1 - \sqrt{2} e^{a/2}x + e^a x^2\right)}{4\sqrt{2}}$$

[Out] x+x/(1+exp(2*a)*x^4)-1/4*arctan(-1+exp(1/2*a)*x*2^(1/2))/exp(1/2*a)*2^(1/2)-1/4*arctan(1+exp(1/2*a)*x*2^(1/2))/exp(1/2*a)*2^(1/2)+1/8*ln(1+exp(a)*x^2-exp(1/2*a)*x*2^(1/2))/exp(1/2*a)*2^(1/2)-1/8*ln(1+exp(a)*x^2+exp(1/2*a)*x*2^(1/2))/exp(1/2*a)*2^(1/2)

Rubi [A]

time = 0.09, antiderivative size = 165, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 9, integrand size = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {5652, 398, 294, 217, 1179, 642, 1176, 631, 210}

$$\frac{e^{-a/2} \operatorname{ArcTan}\left(1 - \sqrt{2} e^{a/2}x\right)}{2\sqrt{2}} - \frac{e^{-a/2} \operatorname{ArcTan}\left(\sqrt{2} e^{a/2}x + 1\right)}{2\sqrt{2}} + \frac{x}{e^{2a}x^4 + 1} + \frac{e^{-a/2} \log\left(e^a x^2 - \sqrt{2} e^{a/2}x + 1\right)}{4\sqrt{2}} - \frac{e^{-a/2} \log\left(e^a x^2 + \sqrt{2} e^{a/2}x + 1\right)}{4\sqrt{2}} + x$$

Antiderivative was successfully verified.

[In] Int[Tanh[a + 2*Log[x]]^2,x]

[Out] x + x/(1 + E^(2*a)*x^4) + ArcTan[1 - Sqrt[2]*E^(a/2)*x]/(2*Sqrt[2]*E^(a/2)) - ArcTan[1 + Sqrt[2]*E^(a/2)*x]/(2*Sqrt[2]*E^(a/2)) + Log[1 - Sqrt[2]*E^(a/2)*x + E^a*x^2]/(4*Sqrt[2]*E^(a/2)) - Log[1 + Sqrt[2]*E^(a/2)*x + E^a*x^2]/(4*Sqrt[2]*E^(a/2))

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 217

Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*r), Int[(r - s*x^2)/(a + b*x^4), x], x] + Dist[1/(2*r), Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))

Rule 294

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n


```

*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x]
;/; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !I
LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]

```

Rule 398

```

Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol]
:> Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a
, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q,
0] && GeQ[p, -q]

```

Rule 631

```

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] :> With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)
], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

```

Rule 642

```

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] :> S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]

```

Rule 1176

```

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] :> With[{q = Rt[
2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]

```

Rule 1179

```

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] :> With[{q = Rt[
-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; Fre
eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]

```

Rule 5652

```

Int[Tanh[((a_) + Log[x]*(b_))*(d_)]^(p_), x_Symbol] :> Int[(-1 + E^(2*
a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p, x] /; FreeQ[{a, b, d, p}, x]

```

Rubi steps

$$\int \tanh^2(a + 2 \log(x)) dx = \int \tanh^2(a + 2 \log(x)) dx$$

Mathematica [A]

time = 0.39, size = 146, normalized size = 0.88

$$\frac{1}{4} \left(4x + \frac{4x}{1 + e^{2a}x^4} + \sqrt[4]{-1} e^{-a/2} \log(\sqrt[4]{-1} e^{-a/2} - x) + (-1)^{3/4} e^{-a/2} \log((-1)^{3/4} e^{-a/2} - x) - \sqrt[4]{-1} e^{-a/2} \log(\sqrt[4]{-1} e^{-a/2} + x) - (-1)^{3/4} e^{-a/2} \log((-1)^{3/4} e^{-a/2} + x) \right)$$

Antiderivative was successfully verified.

[In] Integrate[Tanh[a + 2*Log[x]]^2,x]

[Out] $(4*x + (4*x)/(1 + E^{(2*a)*x^4}) + ((-1)^{(1/4)}*Log[(-1)^{(1/4)}/E^{(a/2)} - x])/E^{(a/2)} + ((-1)^{(3/4)}*Log[(-1)^{(3/4)}/E^{(a/2)} - x])/E^{(a/2)} - ((-1)^{(1/4)}*Log[(-1)^{(1/4)}/E^{(a/2)} + x])/E^{(a/2)} - ((-1)^{(3/4)}*Log[(-1)^{(3/4)}/E^{(a/2)} + x])/E^{(a/2)})/4$

Maple [C] Result contains higher order function than in optimal. Order 9 vs. order 3.

time = 0.53, size = 47, normalized size = 0.28

method	result	size
risch	$x + \frac{x}{1+e^{2a}x^4} - \frac{e^{-2a} \left(\sum_{-R=\text{RootOf}(e^{2a}Z^4+1)} \frac{\ln(x-\frac{R}{-R^3})}{-R^3} \right)}{4}$	47

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(a+2*ln(x))^2,x,method=_RETURNVERBOSE)

[Out] $x+x/(1+\exp(2*a)*x^4)-1/4*\exp(-2*a)*\text{sum}(1/_R^3*\ln(x-_R),_R=\text{RootOf}(\exp(2*a)*Z^4+1))$

Maxima [A]

time = 0.51, size = 138, normalized size = 0.84

$$-\frac{1}{4}\sqrt{2} \arctan\left(\frac{1}{2}\sqrt{2}(2xe^a + \sqrt{2}e^{(1/2)a})e^{(-1/2)a}\right)e^{(-1/2)a} - \frac{1}{4}\sqrt{2} \arctan\left(\frac{1}{2}\sqrt{2}(2xe^a - \sqrt{2}e^{(1/2)a})e^{(-1/2)a}\right)e^{(-1/2)a} - \frac{1}{8}\sqrt{2}e^{(-1/2)a} \log(x^2e^a + \sqrt{2}xe^{(1/2)a} + 1) + \frac{1}{8}\sqrt{2}e^{(-1/2)a} \log(x^2e^a - \sqrt{2}xe^{(1/2)a} + 1) + x + \frac{x}{x^4e^{(2a)} + 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+2*log(x))^2,x, algorithm="maxima")

[Out] $-1/4*\text{sqrt}(2)*\arctan(1/2*\text{sqrt}(2)*(2*x*e^a + \text{sqrt}(2)*e^{(1/2*a)})*e^{(-1/2*a)})*e^{(-1/2*a)} - 1/4*\text{sqrt}(2)*\arctan(1/2*\text{sqrt}(2)*(2*x*e^a - \text{sqrt}(2)*e^{(1/2*a)})*e^{(-1/2*a)})*e^{(-1/2*a)} - 1/8*\text{sqrt}(2)*e^{(-1/2*a)}*\log(x^2*e^a + \text{sqrt}(2)*x*e^{(1/2*a)} + 1) + 1/8*\text{sqrt}(2)*e^{(-1/2*a)}*\log(x^2*e^a - \text{sqrt}(2)*x*e^{(1/2*a)} + 1) + x + \frac{x}{x^4e^{(2a)} + 1}$

2*a) + 1) + 1/8*sqrt(2)*e^(-1/2*a)*log(x^2*e^a - sqrt(2)*x*e^(1/2*a) + 1) + x + x/(x^4*e^(2*a) + 1)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 228 vs. 2(113) = 226.

time = 0.36, size = 228, normalized size = 1.38

$$\frac{8x^5e^{2a} + 4(\sqrt{2}xe^{2a} + \sqrt{2}) \arctan\left(\frac{-\sqrt{2}xe^{2a} + \sqrt{2}\sqrt{\sqrt{2}xe^{2a} + x^2 + e^{-a}}e^{2a} - 1}{8(x^4e^{2a} + 1)}\right) e^{-2a} + 4(\sqrt{2}xe^{2a} + \sqrt{2}) \arctan\left(\frac{-\sqrt{2}xe^{2a} + \sqrt{2}\sqrt{\sqrt{2}xe^{2a} + x^2 + e^{-a}}e^{2a} + 1}{8(x^4e^{2a} + 1)}\right) e^{-2a} - (\sqrt{2}xe^{2a} + \sqrt{2})e^{-2a} \log(\sqrt{2}xe^{2a} + x^2 + e^{-a}) + (\sqrt{2}xe^{2a} + \sqrt{2})e^{-2a} \log(-\sqrt{2}xe^{2a} + x^2 + e^{-a}) + 16x}{8(x^4e^{2a} + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+2*log(x))^2,x, algorithm="fricas")

[Out] 1/8*(8*x^5*e^(2*a) + 4*(sqrt(2)*x^4*e^(2*a) + sqrt(2))*arctan(-sqrt(2)*x*e^(1/2*a) + sqrt(2)*sqrt(sqrt(2)*x*e^(-1/2*a) + x^2 + e^(-a))*e^(1/2*a) - 1)*e^(-1/2*a) + 4*(sqrt(2)*x^4*e^(2*a) + sqrt(2))*arctan(-sqrt(2)*x*e^(1/2*a) + sqrt(2)*sqrt(-sqrt(2)*x*e^(-1/2*a) + x^2 + e^(-a))*e^(1/2*a) + 1)*e^(-1/2*a) - (sqrt(2)*x^4*e^(2*a) + sqrt(2))*e^(-1/2*a)*log(sqrt(2)*x*e^(-1/2*a) + x^2 + e^(-a)) + (sqrt(2)*x^4*e^(2*a) + sqrt(2))*e^(-1/2*a)*log(-sqrt(2)*x*e^(-1/2*a) + x^2 + e^(-a)) + 16*x)/(x^4*e^(2*a) + 1)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \tanh^2(a + 2 \log(x)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+2*ln(x))**2,x)

[Out] Integral(tanh(a + 2*log(x))**2, x)

Giac [A]

time = 0.40, size = 133, normalized size = 0.81

$$-\frac{1}{4}\sqrt{2} \arctan\left(\frac{1}{2}\sqrt{2}(\sqrt{2}e^{-\frac{1}{2}a} + 2x)e^{\frac{1}{2}a}\right)e^{-\frac{1}{2}a} - \frac{1}{4}\sqrt{2} \arctan\left(-\frac{1}{2}\sqrt{2}(\sqrt{2}e^{-\frac{1}{2}a} - 2x)e^{\frac{1}{2}a}\right)e^{-\frac{1}{2}a} - \frac{1}{8}\sqrt{2}e^{-\frac{1}{2}a} \log(\sqrt{2}xe^{-\frac{1}{2}a} + x^2 + e^{-a}) + \frac{1}{8}\sqrt{2}e^{-\frac{1}{2}a} \log(-\sqrt{2}xe^{-\frac{1}{2}a} + x^2 + e^{-a}) + x + \frac{x}{x^4e^{2a} + 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+2*log(x))^2,x, algorithm="giac")

[Out] -1/4*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2)*e^(-1/2*a) + 2*x)*e^(1/2*a))*e^(-1/2*a) - 1/4*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2)*e^(-1/2*a) - 2*x)*e^(1/2*a))*e^(-1/2*a) - 1/8*sqrt(2)*e^(-1/2*a)*log(sqrt(2)*x*e^(-1/2*a) + x^2 + e^(-a)) + 1/8*sqrt(2)*e^(-1/2*a)*log(-sqrt(2)*x*e^(-1/2*a) + x^2 + e^(-a)) + x + x/(x^4*e^(2*a) + 1)

Mupad [B]

time = 1.09, size = 61, normalized size = 0.37

$$x - \frac{\operatorname{atan}\left(x(-e^{2a})^{1/4}\right)}{2(-e^{2a})^{1/4}} + \frac{x}{e^{2a}x^4 + 1} + \frac{\operatorname{atan}\left(x(-e^{2a})^{1/4} \operatorname{li}\right) \operatorname{li}}{2(-e^{2a})^{1/4}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tanh(a + 2*log(x))^2,x)`

[Out] `x - atan(x*(-exp(2*a))^(1/4))/(2*(-exp(2*a))^(1/4)) + (atan(x*(-exp(2*a))^(1/4)*1i)*1i)/(2*(-exp(2*a))^(1/4)) + x/(x^4*exp(2*a) + 1)`

$$3.157 \quad \int \frac{\tanh^2(a+2\log(x))}{x} dx$$

Optimal. Leaf size=14

$$\log(x) - \frac{1}{2} \tanh(a + 2\log(x))$$

[Out] ln(x)-1/2*tanh(a+2*ln(x))

Rubi [A]

time = 0.02, antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {3554, 8}

$$\log(x) - \frac{1}{2} \tanh(a + 2\log(x))$$

Antiderivative was successfully verified.

[In] Int[Tanh[a + 2*Log[x]]^2/x,x]

[Out] Log[x] - Tanh[a + 2*Log[x]]/2

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 3554

Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Dist[b^2, Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]

Rubi steps

$$\begin{aligned} \int \frac{\tanh^2(a + 2\log(x))}{x} dx &= \text{Subst}\left(\int \tanh^2(a + 2x) dx, x, \log(x)\right) \\ &= -\frac{1}{2} \tanh(a + 2\log(x)) + \text{Subst}\left(\int 1 dx, x, \log(x)\right) \\ &= \log(x) - \frac{1}{2} \tanh(a + 2\log(x)) \end{aligned}$$

Mathematica [A]

time = 0.03, size = 24, normalized size = 1.71

$$\frac{1}{2} \tanh^{-1}(\tanh(a + 2\log(x))) - \frac{1}{2} \tanh(a + 2\log(x))$$

Antiderivative was successfully verified.

[In] Integrate[Tanh[a + 2*Log[x]]^2/x,x]

[Out] ArcTanh[Tanh[a + 2*Log[x]]]/2 - Tanh[a + 2*Log[x]]/2

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 34 vs. 2(12) = 24.

time = 0.71, size = 35, normalized size = 2.50

method	result	size
risch	$\frac{1}{1+e^{2a}x^4} + \ln(x)$	16
derivativdivides	$-\frac{\tanh(a+2\ln(x))}{2} - \frac{\ln(\tanh(a+2\ln(x))-1)}{4} + \frac{\ln(\tanh(a+2\ln(x))+1)}{4}$	35
default	$-\frac{\tanh(a+2\ln(x))}{2} - \frac{\ln(\tanh(a+2\ln(x))-1)}{4} + \frac{\ln(\tanh(a+2\ln(x))+1)}{4}$	35

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(a+2*ln(x))^2/x,x,method=_RETURNVERBOSE)

[Out] -1/2*tanh(a+2*ln(x))-1/4*ln(tanh(a+2*ln(x))-1)+1/4*ln(tanh(a+2*ln(x))+1)

Maxima [A]

time = 0.27, size = 21, normalized size = 1.50

$$\frac{1}{2}a - \frac{1}{e^{(-2a-4\log(x))+1}} + \log(x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+2*log(x))^2/x,x, algorithm="maxima")

[Out] 1/2*a - 1/(e^(-2*a - 4*log(x)) + 1) + log(x)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 28 vs. 2(12) = 24.

time = 0.33, size = 28, normalized size = 2.00

$$\frac{(x^4e^{(2a)} + 1)\log(x) + 1}{x^4e^{(2a)} + 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+2*log(x))^2/x,x, algorithm="fricas")

[Out] ((x^4*e^(2*a) + 1)*log(x) + 1)/(x^4*e^(2*a) + 1)

Sympy [A]

time = 0.12, size = 12, normalized size = 0.86

$$\log(x) - \frac{\tanh(a + 2\log(x))}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(a+2*ln(x))**2/x,x)`

[Out] `log(x) - tanh(a + 2*log(x))/2`

Giac [A]

time = 0.41, size = 19, normalized size = 1.36

$$\frac{1}{x^4 e^{(2a)} + 1} + \frac{1}{4} \log(x^4)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(a+2*log(x))^2/x,x, algorithm="giac")`

[Out] `1/(x^4*e^(2*a) + 1) + 1/4*log(x^4)`

Mupad [B]

time = 1.06, size = 28, normalized size = 2.00

$$\ln(x) - \frac{x^4 e^{2a} - 1}{2(e^{2a} x^4 + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tanh(a + 2*log(x))^2/x,x)`

[Out] `log(x) - (x^4*exp(2*a) - 1)/(2*(x^4*exp(2*a) + 1))`

3.158 $\int \frac{\tanh^2(a+2 \log(x))}{x^2} dx$

Optimal. Leaf size=190

$$-\frac{1}{x(1+e^{2ax^4})} - \frac{2e^{2a}x^3}{1+e^{2a}x^4} + \frac{e^{a/2}\text{ArcTan}\left(1-\sqrt{2}e^{a/2}x\right)}{2\sqrt{2}} - \frac{e^{a/2}\text{ArcTan}\left(1+\sqrt{2}e^{a/2}x\right)}{2\sqrt{2}} - \frac{e^{a/2}\log\left(1-\sqrt{2}e^{a/2}x\right)}{4\sqrt{2}}$$

[Out] $-1/x/(1+\exp(2*a)*x^4)-2*\exp(2*a)*x^3/(1+\exp(2*a)*x^4)-1/4*\exp(1/2*a)*\arctan(-1+\exp(1/2*a)*x*2^{(1/2)})*2^{(1/2)}-1/4*\exp(1/2*a)*\arctan(1+\exp(1/2*a)*x*2^{(1/2)})*2^{(1/2)}-1/8*\exp(1/2*a)*\ln(1+\exp(a)*x^2-\exp(1/2*a)*x*2^{(1/2)})*2^{(1/2)}+1/8*\exp(1/2*a)*\ln(1+\exp(a)*x^2+\exp(1/2*a)*x*2^{(1/2)})*2^{(1/2)}$

Rubi [A]

time = 0.09, antiderivative size = 190, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 9, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.692$, Rules used = {5656, 473, 468, 303, 1176, 631, 210, 1179, 642}

$$\frac{e^{a/2}\text{ArcTan}\left(1-\sqrt{2}e^{a/2}x\right)}{2\sqrt{2}} - \frac{e^{a/2}\text{ArcTan}\left(\sqrt{2}e^{a/2}x+1\right)}{2\sqrt{2}} - \frac{1}{x(e^{2a}x^4+1)} - \frac{e^{a/2}\log\left(e^ax^2-\sqrt{2}e^{a/2}x+1\right)}{4\sqrt{2}} + \frac{e^{a/2}\log\left(e^ax^2+\sqrt{2}e^{a/2}x+1\right)}{4\sqrt{2}} - \frac{2e^{2a}x^3}{e^{2a}x^4+1}$$

Antiderivative was successfully verified.

[In] Int[Tanh[a + 2*Log[x]]^2/x^2,x]

[Out] $-(1/(x*(1+E^(2*a)*x^4))) - (2*E^(2*a)*x^3)/(1+E^(2*a)*x^4) + (E^(a/2)*\text{ArcTan}[1-\text{Sqrt}[2]*E^(a/2)*x])/(2*\text{Sqrt}[2]) - (E^(a/2)*\text{ArcTan}[1+\text{Sqrt}[2]*E^(a/2)*x])/(2*\text{Sqrt}[2]) - (E^(a/2)*\text{Log}[1-\text{Sqrt}[2]*E^(a/2)*x+E^a*x^2])/(4*\text{Sqrt}[2]) + (E^(a/2)*\text{Log}[1+\text{Sqrt}[2]*E^(a/2)*x+E^a*x^2])/(4*\text{Sqrt}[2])$

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 303

Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))

Rule 468

Int[((e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(-(b*c - a*d))*(e*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a


```
*b*e*n*(p + 1)), x] - Dist[(a*d*(m + 1) - b*c*(m + n*(p + 1) + 1))/(a*b*n*(p + 1)), Int[(e*x)^m*(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && (( !IntegerQ[p + 1/2] && NeQ[p, -5/4]) || !RationalQ[m] || (IGtQ[n, 0] && ILtQ[p + 1/2, 0] && LeQ[-1, m, (-n)*(p + 1)]))
```

Rule 473

```
Int[((e_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(2, x_Symbol] :> Simp[c^2*(e*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*e*(m + 1))), x] - Dist[1/(a*e^n*(m + 1)), Int[(e*x)^(m + n)*(a + b*x^n)^p*Simp[b*c^2*n*(p + 1) + c*(b*c - 2*a*d)*(m + 1) - a*(m + 1)*d^2*x^n, x], x] /; FreeQ[{a, b, c, d, e, p}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && LtQ[m, -1] && GtQ[n, 0]
```

Rule 631

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] :> With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] :> Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] :> With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] :> With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 5656

```
Int[((e_.)*(x_))^(m_)*Tanh[((a_.) + Log[x_]*(b_.))*(d_.)]^(p_), x_Symbol] :> Int[(e*x)^m*((-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p),
```

x] /; FreeQ[{a, b, d, e, m, p}, x]

Rubi steps

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^2} dx = \int \frac{\tanh^2(a + 2 \log(x))}{x^2} dx$$

Mathematica [A]

time = 0.55, size = 181, normalized size = 0.95

$$\frac{1}{4} \left(-\frac{4}{x} - \frac{4}{\frac{e^{-2a}}{x^3} + x} + (-1)^{3/4} e^{a/2} \log \left(\frac{e^{-2a}(\sqrt[4]{-1} - e^{a/2}x)}{x^4} \right) + \sqrt[4]{-1} e^{a/2} \log \left(\frac{e^{-2a}((-1)^{3/4} - e^{a/2}x)}{x^4} \right) - (-1)^{3/4} e^{a/2} \log \left(\frac{e^{-2a}(\sqrt[4]{-1} + e^{a/2}x)}{x^4} \right) - \sqrt[4]{-1} e^{a/2} \log \left(\frac{e^{-2a}((-1)^{3/4} + e^{a/2}x)}{x^4} \right) \right)$$

Antiderivative was successfully verified.

[In] Integrate[Tanh[a + 2*Log[x]]^2/x^2,x]

[Out] $(-4/x - 4/(1/(E^{(2*a)*x^3}) + x) + (-1)^{(3/4)}*E^{(a/2)}*Log[((-1)^{(1/4)} - E^{(a/2)*x})/(E^{(2*a)*x^4})] + (-1)^{(1/4)}*E^{(a/2)}*Log[((-1)^{(3/4)} - E^{(a/2)*x})/(E^{(2*a)*x^4})] - (-1)^{(3/4)}*E^{(a/2)}*Log[((-1)^{(1/4)} + E^{(a/2)*x})/(E^{(2*a)*x^4})] - (-1)^{(1/4)}*E^{(a/2)}*Log[((-1)^{(3/4)} + E^{(a/2)*x})/(E^{(2*a)*x^4})])/4$

Maple [C] Result contains higher order function than in optimal. Order 9 vs. order 3.

time = 0.49, size = 64, normalized size = 0.34

method	result	size
risch	$\frac{-2e^{2a}x^4 - 1}{x(1 + e^{2a}x^4)} + \frac{\sum_{-R=\text{RootOf}(-Z^4+e^{2a})} -R \ln((5-R^4+4e^{2a})x + R^3)}{4}$	64

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(a+2*ln(x))^2/x^2,x,method=_RETURNVERBOSE)

[Out] $(-2*\exp(2*a)*x^4-1)/x/(1+\exp(2*a)*x^4)+1/4*\text{sum}(_R*\ln((5*_R^4+4*\exp(2*a))*x+_R^3),_R=\text{RootOf}(-Z^4+\exp(2*a)))$

Maxima [A]

time = 0.47, size = 146, normalized size = 0.77

$$\frac{1}{4} \sqrt{2} \arctan \left(\frac{1}{2} \sqrt{2} \left(\sqrt{2} e^{(1/2)a} + \frac{2}{x} \right) e^{(-1/2)a} \right) e^{(1/2)a} + \frac{1}{4} \sqrt{2} \arctan \left(\frac{1}{2} \sqrt{2} \left(\sqrt{2} e^{(1/2)a} - \frac{2}{x} \right) e^{(-1/2)a} \right) e^{(1/2)a} + \frac{1}{8} \sqrt{2} e^{(1/2)a} \log \left(\frac{\sqrt{2} e^{(1/2)a}}{x} + \frac{1}{x^2} + e^a \right) - \frac{1}{8} \sqrt{2} e^{(1/2)a} \log \left(-\frac{\sqrt{2} e^{(1/2)a}}{x} + \frac{1}{x^2} + e^a \right) - \frac{1}{x} - \frac{e^{(2a)}}{x(x^4 + e^{2a})}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+2*log(x))^2/x^2,x, algorithm="maxima")

[Out] $\frac{1}{4}\sqrt{2}\arctan\left(\frac{1}{2}\sqrt{2}\right)\left(\sqrt{2}e^{1/2a} + \frac{2}{x}\right)e^{-1/2a} + \frac{1}{4}\sqrt{2}\arctan\left(-\frac{1}{2}\sqrt{2}\right)\left(\sqrt{2}e^{1/2a} - \frac{2}{x}\right)e^{-1/2a} + \frac{1}{8}\sqrt{2}e^{1/2a}\log\left(\frac{\sqrt{2}e^{1/2a}}{x} + \frac{1}{x^2} + e^a\right) - \frac{1}{8}\sqrt{2}e^{1/2a}\log\left(\frac{-\sqrt{2}e^{1/2a}}{x} + \frac{1}{x^2} + e^a\right) - \frac{1}{x} - \frac{e^{2a}}{x(1/x^4 + e^{2a})}$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 274 vs. 2(136) = 272.

time = 0.40, size = 274, normalized size = 1.44

$$\frac{16x^{2a} - 4(\sqrt{2}x^{2a} + \sqrt{2})\arctan\left(-\frac{\sqrt{2}x^{2a} - \sqrt{2}\sqrt{\sqrt{2}x^{2a} + \sqrt{2}x^{2a} + e^{2a}}e^{1/2a}}{8(x^{2a} + x)}\right)e^{1/2a} - 4(\sqrt{2}x^{2a} + \sqrt{2})\arctan\left(-\frac{\sqrt{2}x^{2a} - \sqrt{2}\sqrt{\sqrt{2}x^{2a} - \sqrt{2}x^{2a} + e^{2a}}e^{1/2a}}{8(x^{2a} + x)}\right)e^{1/2a} - (\sqrt{2}x^{2a} + \sqrt{2})e^{1/2a}\log\left(\frac{x^{2a} + \sqrt{2}x^{2a} + e^{2a}}{x}\right) + (\sqrt{2}x^{2a} + \sqrt{2})e^{1/2a}\log\left(\frac{x^{2a} - \sqrt{2}x^{2a} + e^{2a}}{x}\right) + 8}{8(x^{2a} + x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(a+2*log(x))^2/x^2,x, algorithm="fricas")`

[Out] $-\frac{1}{8}(16x^4e^{2a} - 4(\sqrt{2}x^5e^{2a} + \sqrt{2}x)\arctan(-(\sqrt{2}x^5e^{5/2a} - \sqrt{2}x^5e^{4a} + \sqrt{2}x^5e^{7/2a} + e^{3a}))e^{1/2a} + e^{2a})e^{-2a} + \sqrt{2}x^5e^{7/2a} + e^{3a})e^{1/2a} - 4(\sqrt{2}x^5e^{5/2a} - \sqrt{2}x^5e^{4a} + \sqrt{2}x^5e^{7/2a} + e^{3a})\arctan(-(\sqrt{2}x^5e^{5/2a} - \sqrt{2}x^5e^{4a} - \sqrt{2}x^5e^{7/2a} + e^{3a}))e^{1/2a} - e^{2a})e^{-2a} + \sqrt{2}x^5e^{7/2a} + e^{3a})e^{1/2a} - (\sqrt{2}x^5e^{2a} + \sqrt{2}x)e^{1/2a}\log(x^2e^{4a} + \sqrt{2}x^5e^{7/2a} + e^{3a}) + (\sqrt{2}x^5e^{2a} + \sqrt{2}x)e^{1/2a}\log(x^2e^{4a} - \sqrt{2}x^5e^{7/2a} + e^{3a}) + 8)/(x^5e^{2a} + x)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(a+2*ln(x))**2/x**2,x)`

[Out] `Integral(tanh(a + 2*log(x))**2/x**2, x)`

Giac [A]

time = 0.40, size = 143, normalized size = 0.75

$$-\frac{1}{4}\sqrt{2}\arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2}e^{-1/2a} + 2x\right)e^{1/2a}\right)e^{1/2a} - \frac{1}{4}\sqrt{2}\arctan\left(-\frac{1}{2}\sqrt{2}\left(\sqrt{2}e^{-1/2a} - 2x\right)e^{1/2a}\right)e^{1/2a} + \frac{1}{8}\sqrt{2}e^{1/2a}\log\left(\frac{\sqrt{2}xe^{-1/2a} + x^2 + e^{-a}}{x}\right) - \frac{1}{8}\sqrt{2}e^{1/2a}\log\left(\frac{-\sqrt{2}xe^{-1/2a} + x^2 + e^{-a}}{x}\right) - \frac{2x^4e^{2a} + 1}{x^2e^{2a} + x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(a+2*log(x))^2/x^2,x, algorithm="giac")`

[Out] $-\frac{1}{4}\sqrt{2}\arctan\left(\frac{1}{2}\sqrt{2}\right)\left(\sqrt{2}e^{-1/2a} + 2x\right)e^{1/2a} - \frac{1}{4}\sqrt{2}\arctan\left(-\frac{1}{2}\sqrt{2}\right)\left(\sqrt{2}e^{-1/2a} - 2x\right)e^{1/2a} + \frac{1}{8}\sqrt{2}e^{1/2a}\log\left(\frac{\sqrt{2}e^{-1/2a}}{x} + \frac{1}{x^2} + e^{-a}\right) - \frac{1}{8}\sqrt{2}e^{1/2a}\log\left(\frac{-\sqrt{2}e^{-1/2a}}{x} + \frac{1}{x^2} + e^{-a}\right) - \frac{2x^4e^{2a} + 1}{x^2e^{2a} + x}$

) - 1/8*sqrt(2)*e^(1/2*a)*log(-sqrt(2)*x*e^(-1/2*a) + x^2 + e^(-a)) - (2*x^4*e^(2*a) + 1)/(x^5*e^(2*a) + x)

Mupad [B]

time = 1.10, size = 68, normalized size = 0.36

$$\frac{\operatorname{atanh}\left(x(-e^{2a})^{1/4}\right)(-e^{2a})^{1/4}}{2} - \frac{\operatorname{atan}\left(x(-e^{2a})^{1/4}\right)(-e^{2a})^{1/4}}{2} - \frac{2e^{2a}x^4 + 1}{e^{2a}x^5 + x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(a + 2*log(x))^2/x^2,x)

[Out] (atanh(x*(-exp(2*a))^(1/4))*(-exp(2*a))^(1/4))/2 - (atan(x*(-exp(2*a))^(1/4))*(-exp(2*a))^(1/4))/2 - (2*x^4*exp(2*a) + 1)/(x + x^5*exp(2*a))

$$3.159 \quad \int \frac{\tanh^2(a+2 \log(x))}{x^3} dx$$

Optimal. Leaf size=59

$$-\frac{1}{2x^2(1+e^{2ax^4})} - \frac{3e^{2ax^2}}{2(1+e^{2ax^4})} - e^a \operatorname{ArcTan}(e^ax^2)$$

[Out] $-1/2/x^2/(1+\exp(2*a)*x^4)-3/2*\exp(2*a)*x^2/(1+\exp(2*a)*x^4)-\exp(a)*\arctan(\exp(a)*x^2)$

Rubi [A]

time = 0.04, antiderivative size = 59, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {5656, 473, 468, 281, 209}

$$-e^a \operatorname{ArcTan}(e^ax^2) - \frac{3e^{2ax^2}}{2(e^{2ax^4} + 1)} - \frac{1}{2x^2(e^{2ax^4} + 1)}$$

Antiderivative was successfully verified.

[In] `Int[Tanh[a + 2*Log[x]]^2/x^3, x]`

[Out] $-1/2*1/(x^2*(1 + E^(2*a)*x^4)) - (3*E^(2*a)*x^2)/(2*(1 + E^(2*a)*x^4)) - E^a*\operatorname{ArcTan}[E^a*x^2]$

Rule 209

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

Rule 281

`Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Dist[1/k, Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]`

Rule 468

`Int[((e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(-b*c - a*d)*(e*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*b*e*n*(p + 1))), x] - Dist[(a*d*(m + 1) - b*c*(m + n*(p + 1) + 1))/(a*b*n*(p + 1)), Int[(e*x)^m*(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && ((!IntegerQ[p + 1/2] && NeQ[p, -5/4]) || !RationalQ[m] || (IGtQ[n, 0] && ILtQ[p + 1/2, 0] && LeQ[-1, m, (-n)*(p + 1)]))`

Rule 473

```
Int[((e._)*(x._))^(m._)*((a._) + (b._)*(x._)^(n._))^(p._)*((c._) + (d._)*(x._)^(n._))
^2, x_Symbol] := Simp[c^2*(e*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*e*(m + 1))
), x] - Dist[1/(a*e^n*(m + 1)), Int[(e*x)^(m + n)*(a + b*x^n)^p*Simp[b*c^2*
n*(p + 1) + c*(b*c - 2*a*d)*(m + 1) - a*(m + 1)*d^2*x^n, x], x] /; Free
Q[{a, b, c, d, e, p}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && LtQ[m, -1] &
& GtQ[n, 0]
```

Rule 5656

```
Int[((e._)*(x._))^(m._)*Tanh[((a._) + Log[x_]*(b._))*(d._)]^(p._), x_Symbol]
:= Int[(e*x)^m*((-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p),
x] /; FreeQ[{a, b, d, e, m, p}, x]
```

Rubi steps

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^3} dx = \int \frac{\tanh^2(a + 2 \log(x))}{x^3} dx$$

Mathematica [A]

time = 0.27, size = 40, normalized size = 0.68

$$\frac{-1 - \frac{2}{1 + e^{-2(a + 2 \log(x))}}}{2x^2} + e^a \operatorname{ArcTan}\left(\frac{e^{-a}}{x^2}\right)$$

Antiderivative was successfully verified.

```
[In] Integrate[Tanh[a + 2*Log[x]]^2/x^3,x]
```

```
[Out] (-1 - 2/(1 + E^(-2*(a + 2*Log[x]))))/(2*x^2) + E^a*ArcTan[1/(E^a*x^2)]
```

Maple [C] Result contains higher order function than in optimal. Order 9 vs. order 3.

time = 0.50, size = 66, normalized size = 1.12

method	result	size
risch	$\frac{-\frac{3e^{2a}x^4}{2} - \frac{1}{2}}{x^2(1 + e^{2a}x^4)} + \frac{\left(\sum_{-R=\operatorname{RootOf}(e^{2a} - Z^2)} -R \ln\left(\left(-4e^{2a} - 5R^2\right)x^2 - R\right)\right)}{2}$	66

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(tanh(a+2*ln(x))^2/x^3,x,method=_RETURNVERBOSE)
```

[Out] $(-3/2*\exp(2*a)*x^4-1/2)/x^2/(1+\exp(2*a)*x^4)+1/2*\sum(_R*\ln((-4*\exp(2*a)-5*_R^2)*x^2-_R), _R=\text{RootOf}(\exp(2*a)+_Z^2))$

Maxima [A]

time = 0.48, size = 37, normalized size = 0.63

$$\arctan\left(\frac{e^{(-a)}}{x^2}\right) e^a - \frac{1}{2x^2} - \frac{e^{(2a)}}{x^2\left(\frac{1}{x^4} + e^{(2a)}\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(a+2*log(x))^2/x^3,x, algorithm="maxima")`

[Out] $\arctan(e^{(-a)}/x^2)*e^a - 1/2/x^2 - e^{(2a)}/(x^2*(1/x^4 + e^{(2a)}))$

Fricas [A]

time = 0.37, size = 51, normalized size = 0.86

$$\frac{3x^4e^{(2a)} + 2(x^6e^{(3a)} + x^2e^a)\arctan(x^2e^a) + 1}{2(x^6e^{(2a)} + x^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(a+2*log(x))^2/x^3,x, algorithm="fricas")`

[Out] $-1/2*(3*x^4*e^{(2*a)} + 2*(x^6*e^{(3*a)} + x^2*e^a)*\arctan(x^2*e^a) + 1)/(x^6*e^{(2*a)} + x^2)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\tanh^2(a + 2 \log(x))}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(a+2*ln(x))**2/x**3,x)`

[Out] `Integral(tanh(a + 2*log(x))**2/x**3, x)`

Giac [A]

time = 0.42, size = 39, normalized size = 0.66

$$-\arctan(x^2e^a) e^a - \frac{3x^4e^{(2a)} + 1}{2(x^6e^{(2a)} + x^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(a+2*log(x))^2/x^3,x, algorithm="giac")`

[Out] $-\arctan(x^2*e^a)*e^a - 1/2*(3*x^4*e^{(2*a)} + 1)/(x^6*e^{(2*a)} + x^2)$

Mupad [B]

time = 1.06, size = 47, normalized size = 0.80

$$-\operatorname{atan}\left(x^2 \sqrt{e^{2a}}\right) \sqrt{e^{2a}} - \frac{\frac{3e^{2a}x^4}{2} + \frac{1}{2}}{e^{2a}x^6 + x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tanh(a + 2*log(x))^2/x^3,x)`

[Out] `- atan(x^2*exp(2*a)^(1/2))*exp(2*a)^(1/2) - ((3*x^4*exp(2*a))/2 + 1/2)/(x^6*exp(2*a) + x^2)`

3.160 $\int (ex)^m \tanh(a + 2 \log(x)) dx$

Optimal. Leaf size=60

$$\frac{(ex)^{1+m}}{e(1+m)} - \frac{2(ex)^{1+m} {}_2F_1\left(1, \frac{1+m}{4}; \frac{5+m}{4}; -e^{2a}x^4\right)}{e(1+m)}$$

[Out] $(e*x)^{(1+m)}/e/(1+m)-2*(e*x)^{(1+m)}*hypergeom([1, 1/4+1/4*m], [5/4+1/4*m], -exp(2*a)*x^4)/e/(1+m)$

Rubi [A]

time = 0.03, antiderivative size = 60, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {5656, 470, 371}

$$\frac{(ex)^{m+1}}{e(m+1)} - \frac{2(ex)^{m+1} {}_2F_1\left(1, \frac{m+1}{4}; \frac{m+5}{4}; -e^{2a}x^4\right)}{e(m+1)}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(e*x)^m*\text{Tanh}[a + 2*\text{Log}[x]], x]$

[Out] $(e*x)^{(1+m)}/(e*(1+m)) - (2*(e*x)^{(1+m)}*\text{Hypergeometric2F1}[1, (1+m)/4, (5+m)/4, -(E^{2*a})*x^4])/(e*(1+m))$

Rule 371

$\text{Int}[(c_*)*(x_*)^{(m_*)}*((a_*) + (b_*)*(x_*)^{(n_*)})^{(p_*)}, x_Symbol] \rightarrow \text{Simp}[a^p * ((c*x)^{(m+1)}/(c*(m+1))) * \text{Hypergeometric2F1}[-p, (m+1)/n, (m+1)/n + 1, (-b)*(x^n/a)], x] /;$ FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILt Q[p, 0] || GtQ[a, 0])

Rule 470

$\text{Int}[(e_*)*(x_*)^{(m_*)}*((a_*) + (b_*)*(x_*)^{(n_*)})^{(p_*)}*((c_*) + (d_*)*(x_*)^{(n_*)}), x_Symbol] \rightarrow \text{Simp}[d*(e*x)^{(m+1)}*((a + b*x^n)^{(p+1)}/(b*e*(m+n*(p+1)+1))), x] - \text{Dist}[(a*d*(m+1) - b*c*(m+n*(p+1)+1))/(b*(m+n*(p+1)+1)), \text{Int}[(e*x)^m*(a + b*x^n)^p, x], x] /;$ FreeQ[{a, b, c, d, e, m, n, p}, x] && NeQ[b*c - a*d, 0] && NeQ[m + n*(p+1) + 1, 0]

Rule 5656

$\text{Int}[(e_*)*(x_*)^{(m_*)}*\text{Tanh}[(a_*) + \text{Log}[x_*]*(b_*)*(d_*)]^{(p_*)}, x_Symbol] \rightarrow \text{Int}[(e*x)^m*((-1 + E^{2*a*d})*x^{(2*b*d)})^p/(1 + E^{2*a*d})*x^{(2*b*d)})^p], x] /;$ FreeQ[{a, b, d, e, m, p}, x]

Rubi steps

$$\int (ex)^m \tanh(a + 2 \log(x)) dx = \int (ex)^m \tanh(a + 2 \log(x)) dx$$

Mathematica [A]

time = 0.07, size = 47, normalized size = 0.78

$$\frac{x(ex)^m \left(-1 + 2 {}_2F_1\left(1, \frac{1+m}{4}; \frac{5+m}{4}; -x^4(\cosh(2a) + \sinh(2a))\right)\right)}{1+m}$$

Antiderivative was successfully verified.

[In] Integrate[(e*x)^m*Tanh[a + 2*Log[x]],x]

[Out] -((x*(e*x)^m*(-1 + 2*Hypergeometric2F1[1, (1 + m)/4, (5 + m)/4, -(x^4*(Cosh[2*a] + Sinh[2*a]))]))/(1 + m))

Maple [F]

time = 0.40, size = 0, normalized size = 0.00

$$\int (ex)^m \tanh(a + 2 \ln(x)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((e*x)^m*tanh(a+2*ln(x)),x)

[Out] int((e*x)^m*tanh(a+2*ln(x)),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e*x)^m*tanh(a+2*log(x)),x, algorithm="maxima")

[Out] integrate((x*e)^m*tanh(a + 2*log(x)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e*x)^m*tanh(a+2*log(x)),x, algorithm="fricas")

[Out] integral((x*e)^m*tanh(a + 2*log(x)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (ex)^m \tanh(a + 2 \log(x)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e*x)**m*tanh(a+2*ln(x)),x)

[Out] Integral((e*x)**m*tanh(a + 2*log(x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e*x)^m*tanh(a+2*log(x)),x, algorithm="giac")

[Out] integrate((e*x)^m*tanh(a + 2*log(x)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \tanh(a + 2 \ln(x)) (ex)^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(a + 2*log(x))*(e*x)^m,x)

[Out] int(tanh(a + 2*log(x))*(e*x)^m, x)

3.161 $\int (ex)^m \tanh^2(a + 2 \log(x)) dx$

Optimal. Leaf size=79

$$\frac{(ex)^{1+m}}{e(1+m)} + \frac{(ex)^{1+m}}{e(1+e^{2ax^4})} - \frac{(ex)^{1+m} {}_2F_1\left(1, \frac{1+m}{4}; \frac{5+m}{4}; -e^{2ax^4}\right)}{e}$$

[Out] (e*x)^(1+m)/e/(1+m)+(e*x)^(1+m)/e/(1+exp(2*a)*x^4)-(e*x)^(1+m)*hypergeom([1, 1/4+1/4*m], [5/4+1/4*m], -exp(2*a)*x^4)/e

Rubi [A]

time = 0.05, antiderivative size = 79, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {5656, 474, 470, 371}

$$-\frac{(ex)^{m+1} {}_2F_1\left(1, \frac{m+1}{4}; \frac{m+5}{4}; -e^{2ax^4}\right)}{e} + \frac{(ex)^{m+1}}{e(e^{2ax^4} + 1)} + \frac{(ex)^{m+1}}{e(m+1)}$$

Antiderivative was successfully verified.

[In] Int[(e*x)^m*Tanh[a + 2*Log[x]]^2,x]

[Out] (e*x)^(1 + m)/(e*(1 + m)) + (e*x)^(1 + m)/(e*(1 + E^(2*a)*x^4)) - ((e*x)^(1 + m)*Hypergeometric2F1[1, (1 + m)/4, (5 + m)/4, -(E^(2*a)*x^4)])/e

Rule 371

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])

Rule 470

Int[((e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> Simp[d*(e*x)^(m + 1)*((a + b*x^n)^(p + 1)/(b*e*(m + n*(p + 1) + 1))), x] - Dist[(a*d*(m + 1) - b*c*(m + n*(p + 1) + 1))/(b*(m + n*(p + 1) + 1)), Int[(e*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && NeQ[b*c - a*d, 0] && NeQ[m + n*(p + 1) + 1, 0]

Rule 474

Int[((e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(2), x_Symbol] :> Simp[(-b*c - a*d)^2*(e*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*b^2*e*n*(p + 1))), x] + Dist[1/(a*b^2*n*(p + 1)), Int[(e*x)^m*(a + b*x^n)^(p + 1)*Simp[(b*c - a*d)^2*(m + 1) + b^2*c^2*n*(p + 1) + a*b*d^2*n*(p + 1)*x^n, x], x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && NeQ[b*c - a*d, 0]

&& IGtQ[n, 0] && LtQ[p, -1]

Rule 5656

```
Int[((e_.)*(x_.))^(m_.)*Tanh[(a_.) + Log[x_]*(b_.)]*(d_.)]^(p_.), x_Symbol]
  :-> Int[(e*x)^(m*(-1 + E^(2*a*d)*x^(2*b*d)))^p/(1 + E^(2*a*d)*x^(2*b*d))^p),
  x] /; FreeQ[{a, b, d, e, m, p}, x]
```

Rubi steps

$$\int (ex)^m \tanh^2(a + 2 \log(x)) dx = \int (ex)^m \tanh^2(a + 2 \log(x)) dx$$

Mathematica [A]

time = 0.12, size = 79, normalized size = 1.00

$$\frac{x(ex)^m (-1 + 4 {}_2F_1(1, \frac{1+m}{4}; \frac{5+m}{4}; -x^4(\cosh(2a) + \sinh(2a))) - 4 {}_2F_1(2, \frac{1+m}{4}; \frac{5+m}{4}; -x^4(\cosh(2a) + \sinh(2a))))}{1+m}$$

Antiderivative was successfully verified.

[In] Integrate[(e*x)^m*Tanh[a + 2*Log[x]]^2,x]

[Out] -((x*(e*x)^m*(-1 + 4*Hypergeometric2F1[1, (1 + m)/4, (5 + m)/4, -(x^4*(Cosh[2*a] + Sinh[2*a]))] - 4*Hypergeometric2F1[2, (1 + m)/4, (5 + m)/4, -(x^4*(Cosh[2*a] + Sinh[2*a]))])))/(1 + m)

Maple [F]

time = 0.30, size = 0, normalized size = 0.00

$$\int (ex)^m (\tanh^2(a + 2 \ln(x))) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((e*x)^m*tanh(a+2*ln(x))^2,x)

[Out] int((e*x)^m*tanh(a+2*ln(x))^2,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e*x)^m*tanh(a+2*log(x))^2,x, algorithm="maxima")

[Out] integrate((x*e)^m*tanh(a + 2*log(x))^2, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e*x)^m*tanh(a+2*log(x))^2,x, algorithm="fricas")

[Out] integral((x*e)^m*tanh(a + 2*log(x))^2, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (ex)^m \tanh^2(a + 2 \log(x)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e*x)**m*tanh(a+2*ln(x))**2,x)

[Out] Integral((e*x)**m*tanh(a + 2*log(x))**2, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e*x)^m*tanh(a+2*log(x))^2,x, algorithm="giac")

[Out] integrate((e*x)^m*tanh(a + 2*log(x))^2, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \tanh(a + 2 \ln(x))^2 (ex)^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(a + 2*log(x))^2*(e*x)^m,x)

[Out] int(tanh(a + 2*log(x))^2*(e*x)^m, x)

3.162 $\int (ex)^m \tanh^3(a + 2 \log(x)) dx$

Optimal. Leaf size=176

$$\frac{(3+m)(5+m)(ex)^{1+m}}{8e(1+m)} - \frac{(ex)^{1+m}(1-e^{2a}x^4)^2}{4e(1+e^{2a}x^4)^2} - \frac{e^{-2a}(ex)^{1+m}(e^{2a}(3-m)+e^{4a}(5+m)x^4)}{8e(1+e^{2a}x^4)} - \frac{(9+2m+m^2)}{8e(1+m)}$$

[Out] $1/8*(3+m)*(5+m)*(e*x)^{(1+m)}/e/(1+m)-1/4*(e*x)^{(1+m)*(1-\exp(2*a)*x^4)^2}/e/(1+\exp(2*a)*x^4)^2-1/8*(e*x)^{(1+m)*(exp(2*a)*(3-m)+exp(4*a)*(5+m)*x^4)}/e/exp(2*a)/(1+\exp(2*a)*x^4)-1/4*(m^2+2*m+9)*(e*x)^{(1+m)*hypergeom([1, 1/4+1/4*m], [5/4+1/4*m], -\exp(2*a)*x^4)}/e/(1+m)$

Rubi [A]

time = 0.12, antiderivative size = 176, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {5656, 479, 591, 470, 371}

$$-\frac{(m^2+2m+9)(ex)^{m+1} {}_2F_1\left(1, \frac{m+1}{4}; \frac{m+5}{4}; -e^{2a}x^4\right)}{4e(m+1)} - \frac{e^{-2a}(e^{4a}(m+5)x^4 + e^{2a}(3-m))(ex)^{m+1}}{8e(e^{2a}x^4+1)} - \frac{(1-e^{2a}x^4)^2(ex)^{m+1}}{4e(e^{2a}x^4+1)^2} + \frac{(m+3)(m+5)(ex)^{m+1}}{8e(m+1)}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(e*x)^m*\text{Tanh}[a + 2*\text{Log}[x]]^3, x]$

[Out] $((3+m)*(5+m)*(e*x)^{(1+m)})/(8*e*(1+m)) - ((e*x)^{(1+m)*(1-E^{(2*a)*x^4})^2})/(4*e*(1+E^{(2*a)*x^4})^2) - ((e*x)^{(1+m)*(E^{(2*a)*x^4)*(3-m)+E^{(4*a)*x^4}(5+m)*x^4)})/(8*e*E^{(2*a)*x^4}*(1+E^{(2*a)*x^4})) - ((9+2*m+m^2)*(e*x)^{(1+m)*Hypergeometric2F1[1, (1+m)/4, (5+m)/4, -(E^{(2*a)*x^4})]})/(4*e*(1+m))$

Rule 371

$\text{Int}[(c*x)^m*(a+b*x^n)^p, x_Symbol] \rightarrow \text{Simp}[a^p*(c*x)^{m+1}/(c*(m+1))*\text{Hypergeometric2F1}[-p, (m+1)/n, (m+1)/n+1, (-b)*(x^n/a)], x] /; \text{FreeQ}\{a, b, c, m, n, p\}, x \ \&\& \ !\text{IGtQ}[p, 0] \ \&\& \ (\text{ILtQ}[p, 0] \ || \ \text{GtQ}[a, 0])$

Rule 470

$\text{Int}[(e*x)^m*(a+b*x^n)^p*(c+d*x^n), x_Symbol] \rightarrow \text{Simp}[d*(e*x)^{m+1}*(a+b*x^n)^{p+1}/(b*e*(m+n*(p+1)+1)), x] - \text{Dist}[(a*d*(m+1) - b*c*(m+n*(p+1)+1))/(b*(m+n*(p+1)+1)), \text{Int}[(e*x)^m*(a+b*x^n)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, m, n, p\}, x \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[m+n*(p+1)+1, 0]$

Rule 479

$\text{Int}[(e*x)^m*(a+b*x^n)^p*(c+d*x^n)^q, x_Symbol] \rightarrow \text{Simp}[(-c*b - a*d)*(e*x)^{m+1}*(a+b*x^n)^{p+1}]$

```

*((c + d*x^n)^(q - 1)/(a*b*e*n*(p + 1))), x] + Dist[1/(a*b*n*(p + 1)), Int[
(e*x)^m*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 2)*Simp[c*(c*b*n*(p + 1) + (c*
b - a*d)*(m + 1)) + d*(c*b*n*(p + 1) + (c*b - a*d)*(m + n*(q - 1) + 1))*x^n
, x], x], x] /; FreeQ[{a, b, c, d, e, m}, x] && NeQ[b*c - a*d, 0] && IGtQ[n
, 0] && LtQ[p, -1] && GtQ[q, 1] && IntBinomialQ[a, b, c, d, e, m, n, p, q,
x]

```

Rule 591

```

Int[((g_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_
))^(q_)*((e_) + (f_)*(x_)^(n_)), x_Symbol] := Simp[(-(b*e - a*f))*(g*x)^(
m + 1)*(a + b*x^n)^(p + 1)*((c + d*x^n)^q/(a*b*g*n*(p + 1))), x] + Dist[1/(
a*b*n*(p + 1)), Int[(g*x)^m*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1)*Simp[c*
(b*e*n*(p + 1) + (b*e - a*f)*(m + 1)) + d*(b*e*n*(p + 1) + (b*e - a*f)*(m +
n*q + 1))*x^n, x], x], x] /; FreeQ[{a, b, c, d, e, f, g, m}, x] && IGtQ[n,
0] && LtQ[p, -1] && GtQ[q, 0] && !(EqQ[q, 1] && SimplerQ[b*c - a*d, b*e -
a*f])

```

Rule 5656

```

Int[((e_)*(x_))^(m_)*Tanh[((a_) + Log[x_]*(b_))*(d_)]^(p_), x_Symbol]
:= Int[(e*x)^m*((-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p),
x] /; FreeQ[{a, b, d, e, m, p}, x]

```

Rubi steps

$$\int (ex)^m \tanh^3(a + 2 \log(x)) dx = \int (ex)^m \tanh^3(a + 2 \log(x)) dx$$

Mathematica [A]

time = 0.16, size = 111, normalized size = 0.63

$$\frac{x(ex)^m \left(-1 + 6 {}_2F_1\left(1, \frac{1+m}{4}; \frac{5+m}{4}; -x^4(\cosh(2a) + \sinh(2a))\right) - 12 {}_2F_1\left(2, \frac{1+m}{4}; \frac{5+m}{4}; -x^4(\cosh(2a) + \sinh(2a))\right) + 8 {}_2F_1\left(3, \frac{1+m}{4}; \frac{5+m}{4}; -x^4(\cosh(2a) + \sinh(2a))\right) \right)}{1+m}$$

Antiderivative was successfully verified.

```
[In] Integrate[(e*x)^m*Tanh[a + 2*Log[x]]^3,x]
```

```
[Out] -((x*(e*x)^m*(-1 + 6*Hypergeometric2F1[1, (1 + m)/4, (5 + m)/4, -(x^4*(Cosh
[2*a] + Sinh[2*a]))] - 12*Hypergeometric2F1[2, (1 + m)/4, (5 + m)/4, -(x^4*
(Cosh[2*a] + Sinh[2*a]))] + 8*Hypergeometric2F1[3, (1 + m)/4, (5 + m)/4, -(
x^4*(Cosh[2*a] + Sinh[2*a]))])))/(1 + m))

```


Maple [F]

time = 0.30, size = 0, normalized size = 0.00

$$\int (ex)^m (\tanh^3(a + 2 \ln(x))) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((e*x)^m*tanh(a+2*ln(x))^3,x)

[Out] int((e*x)^m*tanh(a+2*ln(x))^3,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e*x)^m*tanh(a+2*log(x))^3,x, algorithm="maxima")

[Out] integrate((x*e)^m*tanh(a + 2*log(x))^3, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e*x)^m*tanh(a+2*log(x))^3,x, algorithm="fricas")

[Out] integral((x*e)^m*tanh(a + 2*log(x))^3, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (ex)^m \tanh^3(a + 2 \log(x)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e*x)**m*tanh(a+2*ln(x))**3,x)

[Out] Integral((e*x)**m*tanh(a + 2*log(x))**3, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e*x)^m*tanh(a+2*log(x))^3,x, algorithm="giac")

[Out] integrate((e*x)^m*tanh(a + 2*log(x))^3, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \tanh(a + 2 \ln(x))^3 (e x)^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(a + 2*log(x))^3*(e*x)^m,x)

[Out] int(tanh(a + 2*log(x))^3*(e*x)^m, x)

3.163 $\int \tanh^p(a + b \log(x)) dx$

Optimal. Leaf size=79

$$x(1 - e^{2a}x^{2b})^{-p} (-1 + e^{2a}x^{2b})^p F_1\left(\frac{1}{2b}; -p, p; \frac{1}{2}\left(2 + \frac{1}{b}\right); e^{2a}x^{2b}, -e^{2a}x^{2b}\right)$$

[Out] $x*(-1+\exp(2*a)*x^{(2*b)})^p*\text{AppellF1}(1/2/b, -p, p, 1+1/2/b, \exp(2*a)*x^{(2*b)}, -\exp(2*a)*x^{(2*b)})/((1-\exp(2*a)*x^{(2*b)})^p)$

Rubi [A]

time = 0.03, antiderivative size = 79, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {5652, 441, 440}

$$x(1 - e^{2a}x^{2b})^{-p} (e^{2a}x^{2b} - 1)^p F_1\left(\frac{1}{2b}; -p, p; \frac{1}{2}\left(2 + \frac{1}{b}\right); e^{2a}x^{2b}, -e^{2a}x^{2b}\right)$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Tanh}[a + b*\text{Log}[x]]^p, x]$

[Out] $(x*(-1 + E^{(2*a)*x^{(2*b)}})^p*\text{AppellF1}[1/(2*b), -p, p, (2 + b^{(-1)})/2, E^{(2*a)*x^{(2*b)}}, -(E^{(2*a)*x^{(2*b)}})])/(1 - E^{(2*a)*x^{(2*b)}})^p$

Rule 440

$\text{Int}[(a_ + (b_)*(x_)^{(n_)})^{(p_)}*((c_) + (d_)*(x_)^{(n_)})^{(q_)}, x_Symbol]$
 $\text{:> Simp}[a^p*c^q*x*\text{AppellF1}[1/n, -p, -q, 1 + 1/n, (-b)*(x^n/a), (-d)*(x^n/c)], x] /;$ $\text{FreeQ}\{a, b, c, d, n, p, q\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[n, -1]$
 $\ \&\& \ (\text{IntegerQ}[p] \ || \ \text{GtQ}[a, 0]) \ \&\& \ (\text{IntegerQ}[q] \ || \ \text{GtQ}[c, 0])$

Rule 441

$\text{Int}[(a_ + (b_)*(x_)^{(n_)})^{(p_)}*((c_) + (d_)*(x_)^{(n_)})^{(q_)}, x_Symbol]$
 $\text{:> Dist}[a^{\text{IntPart}[p]}*((a + b*x^n)^{\text{FracPart}[p]}/(1 + b*(x^n/a))^{\text{FracPart}[p]}],$
 $\text{Int}[(1 + b*(x^n/a))^p*(c + d*x^n)^q, x], x] /;$ $\text{FreeQ}\{a, b, c, d, n, p, q\}$
 $, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[n, -1] \ \&\& \ !(\text{IntegerQ}[p] \ || \ \text{GtQ}[a, 0])$

Rule 5652

$\text{Int}[\text{Tanh}[(a_ + \text{Log}[x_]*(b_))*(d_)]^{(p_)}, x_Symbol] \text{:> Int}[(-1 + E^{(2*a*d)*x^{(2*b*d)}})^p/(1 + E^{(2*a*d)*x^{(2*b*d)}})^p, x] /;$ $\text{FreeQ}\{a, b, d, p\}, x]$

Rubi steps

$$\int \tanh^p(a + b \log(x)) dx = \int \tanh^p(a + b \log(x)) dx$$

Mathematica [B] Leaf count is larger than twice the leaf count of optimal. 259 vs. 2(79) = 158.

time = 1.46, size = 259, normalized size = 3.28

$$\frac{(1+2b)x\left(\frac{-1+e^{2ax^{2b}}}{1+e^{2ax^{2b}}}\right)^p F_1\left(\frac{1}{2b}; -p, p; 1+\frac{1}{2b}; e^{2ax^{2b}}, -e^{2ax^{2b}}\right)}{-2be^{2a}px^{2b}F_1\left(1+\frac{1}{2b}; 1-p, p; 2+\frac{1}{2b}; e^{2ax^{2b}}, -e^{2ax^{2b}}\right) - 2be^{2a}px^{2b}F_1\left(1+\frac{1}{2b}; -p, 1+p; 2+\frac{1}{2b}; e^{2ax^{2b}}, -e^{2ax^{2b}}\right) + (1+2b)F_1\left(\frac{1}{2b}; -p, p; 1+\frac{1}{2b}; e^{2ax^{2b}}, -e^{2ax^{2b}}\right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[Tanh[a + b*Log[x]]^p, x]

[Out] $((1+2b)*x*((-1+E^{(2*a)*x^{(2*b)}})/(1+E^{(2*a)*x^{(2*b)}}))^p \text{AppellF1}[1/(2*b), -p, p, 1+1/(2*b), E^{(2*a)*x^{(2*b)}}, -(E^{(2*a)*x^{(2*b)}})]/(-2*b*E^{(2*a)*x^{(2*b)}})*\text{AppellF1}[1+1/(2*b), 1-p, p, 2+1/(2*b), E^{(2*a)*x^{(2*b)}}, -(E^{(2*a)*x^{(2*b)}})] - 2*b*E^{(2*a)*x^{(2*b)}}*\text{AppellF1}[1+1/(2*b), -p, 1+p, 2+1/(2*b), E^{(2*a)*x^{(2*b)}}, -(E^{(2*a)*x^{(2*b)}})] + (1+2b)*\text{AppellF1}[1/(2*b), -p, p, 1+1/(2*b), E^{(2*a)*x^{(2*b)}}, -(E^{(2*a)*x^{(2*b)}})]$

Maple [F]

time = 0.62, size = 0, normalized size = 0.00

$$\int \tanh^p(a + b \ln(x)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(a+b*ln(x))^p, x)

[Out] int(tanh(a+b*ln(x))^p, x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+b*log(x))^p, x, algorithm="maxima")

[Out] integrate(tanh(b*log(x) + a)^p, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+b*log(x))^p, x, algorithm="fricas")

[Out] integral(tanh(b*log(x) + a)^p, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \tanh^p(a + b \log(x)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+b*ln(x))**p,x)

[Out] Integral(tanh(a + b*log(x))**p, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+b*log(x))^p,x, algorithm="giac")

[Out] integrate(tanh(b*log(x) + a)^p, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \tanh(a + b \ln(x))^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(a + b*log(x))^p,x)

[Out] int(tanh(a + b*log(x))^p, x)

3.164 $\int (ex)^m \tanh^p(a + b \log(x)) dx$

Optimal. Leaf size=99

$$\frac{(ex)^{1+m} (1 - e^{2a}x^{2b})^{-p} (-1 + e^{2a}x^{2b})^p F_1\left(\frac{1+m}{2b}; -p, p; 1 + \frac{1+m}{2b}; e^{2a}x^{2b}, -e^{2a}x^{2b}\right)}{e(1+m)}$$

[Out] (e*x)^(1+m)*(-1+exp(2*a)*x^(2*b))^p*AppellF1(1/2*(1+m)/b, -p, p, 1+1/2*(1+m)/b, exp(2*a)*x^(2*b), -exp(2*a)*x^(2*b))/e/(1+m)/((1-exp(2*a)*x^(2*b))^p)

Rubi [A]

time = 0.06, antiderivative size = 99, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {5656, 525, 524}

$$\frac{(ex)^{m+1} (1 - e^{2a}x^{2b})^{-p} (e^{2a}x^{2b} - 1)^p F_1\left(\frac{m+1}{2b}; -p, p; \frac{m+1}{2b} + 1; e^{2a}x^{2b}, -e^{2a}x^{2b}\right)}{e(m+1)}$$

Antiderivative was successfully verified.

[In] Int[(e*x)^m*Tanh[a + b*Log[x]]^p,x]

[Out] ((e*x)^(1 + m)*(-1 + E^(2*a)*x^(2*b))^p*AppellF1[(1 + m)/(2*b), -p, p, 1 + (1 + m)/(2*b), E^(2*a)*x^(2*b), -(E^(2*a)*x^(2*b))])/(e*(1 + m)*(1 - E^(2*a)*x^(2*b))^p)

Rule 524

Int[((e._)*(x_))^(m._)*((a_) + (b._)*(x_)^(n_))^(p_)*((c_) + (d._)*(x_)^(n_))^(q_), x_Symbol] :> Simp[a^p*c^q*((e*x)^(m + 1)/(e*(m + 1)))*AppellF1[(m + 1)/n, -p, -q, 1 + (m + 1)/n, (-b)*(x^n/a), (-d)*(x^n/c)], x] /; FreeQ[{a, b, c, d, e, m, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[m, -1] && NeQ[m, n - 1] && (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])

Rule 525

Int[((e._)*(x_))^(m._)*((a_) + (b._)*(x_)^(n_))^(p_)*((c_) + (d._)*(x_)^(n_))^(q_), x_Symbol] :> Dist[a^IntPart[p]*((a + b*x^n)^FracPart[p]/(1 + b*(x^n/a)^FracPart[p])), Int[(e*x)^m*(1 + b*(x^n/a))^p*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, e, m, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[m, -1] && NeQ[m, n - 1] && !(IntegerQ[p] || GtQ[a, 0])

Rule 5656

Int[((e._)*(x_))^(m._)*Tanh[((a_) + Log[x_]*(b._))*(d._)]^(p._), x_Symbol] :> Int[(e*x)^m*((-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p), x] /; FreeQ[{a, b, d, e, m, p}, x]

Rubi steps

$$\int (ex)^m \tanh^p(a + b \log(x)) dx = \int (ex)^m \tanh^p(a + b \log(x)) dx$$

Mathematica [A]

time = 2.21, size = 126, normalized size = 1.27

$$\frac{x(ex)^m (1 - e^{2a}x^{2b})^{-p} \left(\frac{-1+e^{2a}x^{2b}}{1+e^{2a}x^{2b}}\right)^p (1 + e^{2a}x^{2b})^p F_1\left(\frac{1+m}{2b}; -p, p; 1 + \frac{1+m}{2b}; e^{2a}x^{2b}, -e^{2a}x^{2b}\right)}{1 + m}$$

Warning: Unable to verify antiderivative.

[In] Integrate[(e*x)^m*Tanh[a + b*Log[x]]^p,x]

[Out] (x*(e*x)^m*((-1 + E^(2*a)*x^(2*b))/(1 + E^(2*a)*x^(2*b)))^p*(1 + E^(2*a)*x^(2*b))^p*AppellF1[(1 + m)/(2*b), -p, p, 1 + (1 + m)/(2*b), E^(2*a)*x^(2*b), -(E^(2*a)*x^(2*b))]/((1 + m)*(1 - E^(2*a)*x^(2*b))^p)

Maple [F]

time = 0.40, size = 0, normalized size = 0.00

$$\int (ex)^m (\tanh^p(a + b \ln(x))) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((e*x)^m*tanh(a+b*ln(x))^p,x)**[Out]** int((e*x)^m*tanh(a+b*ln(x))^p,x)**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e*x)^m*tanh(a+b*log(x))^p,x, algorithm="maxima")**[Out]** integrate((x*e)^m*tanh(b*log(x) + a)^p, x)**Fricas [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e*x)^m*tanh(a+b*log(x))^p,x, algorithm="fricas")

[Out] integral((x*e)^m*tanh(b*log(x) + a)^p, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (ex)^m \tanh^p(a + b \log(x)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e*x)**m*tanh(a+b*ln(x))**p,x)

[Out] Integral((e*x)**m*tanh(a + b*log(x))**p, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e*x)^m*tanh(a+b*log(x))^p,x, algorithm="giac")

[Out] integrate((e*x)^m*tanh(b*log(x) + a)^p, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \tanh(a + b \ln(x))^p (ex)^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(a + b*log(x))^p*(e*x)^m,x)

[Out] int(tanh(a + b*log(x))^p*(e*x)^m, x)

3.165 $\int \tanh^p \left(a + \frac{\log(x)}{2} \right) dx$

Optimal. Leaf size=51

$$\frac{2^{-p} e^{-2a} (-1 + e^{2ax})^{1+p} {}_2F_1(p, 1+p; 2+p; \frac{1}{2}(1 - e^{2ax}))}{1+p}$$

[Out] $(-1 + \exp(2*a)*x)^{(1+p)} * \text{hypergeom}([p, 1+p], [2+p], 1/2 - 1/2*\exp(2*a)*x)/(2^p)/\exp(2*a)/(1+p)$

Rubi [A]

time = 0.02, antiderivative size = 51, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {5652, 71}

$$\frac{e^{-2a} 2^{-p} (e^{2ax} - 1)^{p+1} {}_2F_1(p, p+1; p+2; \frac{1}{2}(1 - e^{2ax}))}{p+1}$$

Antiderivative was successfully verified.

[In] Int[Tanh[a + Log[x]/2]^p, x]

[Out] $((-1 + E^{(2*a)*x})^{(1 + p)} * \text{Hypergeometric2F1}[p, 1 + p, 2 + p, (1 - E^{(2*a)*x})/2]) / (2^p * E^{(2*a)} * (1 + p))$

Rule 71

Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(((a + b*x)^(m + 1)/(b*(m + 1)*(b/(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

Rule 5652

Int[Tanh[((a_.) + Log[x_]*(b_.))*(d_.)]^(p_.), x_Symbol] := Int[(-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p, x] /; FreeQ[{a, b, d, p}, x]

Rubi steps

$$\int \tanh^p \left(a + \frac{\log(x)}{2} \right) dx = \int \tanh^p \left(\frac{1}{2}(2a + \log(x)) \right) dx$$

Mathematica [A]

time = 2.08, size = 76, normalized size = 1.49

$$\frac{2^{-p} e^{-2a} \left(\frac{-1+e^{2ax}}{1+e^{2ax}} \right)^{1+p} (1+e^{2ax})^{1+p} {}_2F_1\left(p, 1+p; 2+p; \frac{1}{2}(1-e^{2ax})\right)}{1+p}$$

Antiderivative was successfully verified.

`[In] Integrate[Tanh[a + Log[x]/2]^p, x]``[Out] (((-1 + E^(2*a)*x)/(1 + E^(2*a)*x))^(1 + p)*(1 + E^(2*a)*x)^(1 + p)*Hypergeometric2F1[p, 1 + p, 2 + p, (1 - E^(2*a)*x)/2])/(2^p*E^(2*a)*(1 + p))`**Maple [F]**

time = 0.61, size = 0, normalized size = 0.00

$$\int \tanh^p \left(a + \frac{\ln(x)}{2} \right) dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(tanh(a+1/2*ln(x))^p, x)``[Out] int(tanh(a+1/2*ln(x))^p, x)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(tanh(a+1/2*log(x))^p, x, algorithm="maxima")``[Out] integrate(tanh(a + 1/2*log(x))^p, x)`**Fricas [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(tanh(a+1/2*log(x))^p, x, algorithm="fricas")``[Out] integral(tanh(a + 1/2*log(x))^p, x)`**Sympy [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int \tanh^p \left(a + \frac{\log(x)}{2} \right) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(a+1/2*ln(x))**p,x)`

[Out] `Integral(tanh(a + log(x)/2)**p, x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(a+1/2*log(x))^p,x, algorithm="giac")`

[Out] `integrate(tanh(a + 1/2*log(x))^p, x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \tanh\left(a + \frac{\ln(x)}{2}\right)^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tanh(a + log(x)/2)^p,x)`

[Out] `int(tanh(a + log(x)/2)^p, x)`

3.166 $\int \tanh^p \left(a + \frac{\log(x)}{4} \right) dx$

Optimal. Leaf size=106

$$e^{-4a} (-1 + e^{2a} \sqrt{x})^{1+p} (1 + e^{2a} \sqrt{x})^{1-p} - \frac{2^{1-p} e^{-4a} p (-1 + e^{2a} \sqrt{x})^{1+p} {}_2F_1(p, 1+p; 2+p; \frac{1}{2}(1 - e^{2a} \sqrt{x}))}{1+p}$$

[Out] $-2^{-(1-p)} p \text{hypergeom}([p, 1+p], [2+p], 1/2 - 1/2 \exp(2a) x^{(1/2)}) * (-1 + \exp(2a) x^{(1/2)})^{(1+p)} / \exp(4a) / (1+p) + (-1 + \exp(2a) x^{(1/2)})^{(1+p)} * (1 + \exp(2a) x^{(1/2)})^{(1-p)} / \exp(4a)$

Rubi [A]

time = 0.06, antiderivative size = 106, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {5652, 383, 81, 71}

$$e^{-4a} (e^{2a} \sqrt{x} - 1)^{p+1} (e^{2a} \sqrt{x} + 1)^{1-p} - \frac{e^{-4a} 2^{1-p} p (e^{2a} \sqrt{x} - 1)^{p+1} {}_2F_1(p, p+1; p+2; \frac{1}{2}(1 - e^{2a} \sqrt{x}))}{p+1}$$

Antiderivative was successfully verified.

[In] Int[Tanh[a + Log[x]/4]^p, x]

[Out] $((-1 + E^{(2a)*\text{Sqrt}[x]})^{(1+p)} * (1 + E^{(2a)*\text{Sqrt}[x]})^{(1-p)}) / E^{(4a)} - (2^{(1-p)} p * (-1 + E^{(2a)*\text{Sqrt}[x]})^{(1+p)} * \text{Hypergeometric2F1}[p, 1+p, 2+p, (1 - E^{(2a)*\text{Sqrt}[x]})/2]) / (E^{(4a)} * (1+p))$

Rule 71

Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] :> Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b*(b*c - a*d))^(n)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

Rule 81

Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] :> Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)), Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]

Rule 383

Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.), x_Symbol] :> With[{g = Denominator[n]}, Dist[g, Subst[Int[x^(g - 1)*(a + b*x^(g*n))

$\int \frac{\tanh^p(c + d*x^{(g*n)})^q}{x^{1/g}} dx$; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && FractionQ[n]

Rule 5652

Int[Tanh[(a_.) + Log[x_]*(b_.)]*(d_.)]^(p_.), x_Symbol] := Int[(-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p, x] /; FreeQ[{a, b, d, p}, x]

Rubi steps

$$\int \tanh^p \left(a + \frac{\log(x)}{4} \right) dx = \int \tanh^p \left(\frac{1}{4}(4a + \log(x)) \right) dx$$

Mathematica [A]

time = 2.00, size = 121, normalized size = 1.14

$$\frac{e^{-4a}(-1 + e^{2a}\sqrt{x}) \left(\frac{-1 + e^{2a}\sqrt{x}}{2 + 2e^{2a}\sqrt{x}} \right)^p (2^p(1+p)(1 + e^{2a}\sqrt{x}) - 2p(1 + e^{2a}\sqrt{x})^p {}_2F_1(p, 1+p; 2+p; \frac{1}{2}(1 - e^{2a}\sqrt{x})))}{1+p}$$

Antiderivative was successfully verified.

[In] Integrate[Tanh[a + Log[x]/4]^p, x]

[Out] ((-1 + E^(2*a)*Sqrt[x])*((-1 + E^(2*a)*Sqrt[x])/(2 + 2*E^(2*a)*Sqrt[x]))^p*(2^p*(1 + p)*(1 + E^(2*a)*Sqrt[x]) - 2*p*(1 + E^(2*a)*Sqrt[x])^p*Hypergeometric2F1[p, 1 + p, 2 + p, (1 - E^(2*a)*Sqrt[x])/2]))/(E^(4*a)*(1 + p))

Maple [F]

time = 0.62, size = 0, normalized size = 0.00

$$\int \tanh^p \left(a + \frac{\ln(x)}{4} \right) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(a+1/4*ln(x))^p, x)

[Out] int(tanh(a+1/4*ln(x))^p, x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+1/4*log(x))^p,x, algorithm="maxima")

[Out] integrate(tanh(a + 1/4*log(x))^p, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+1/4*log(x))^p,x, algorithm="fricas")

[Out] integral(tanh(a + 1/4*log(x))^p, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \tanh^p \left(a + \frac{\log(x)}{4} \right) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+1/4*ln(x))**p,x)

[Out] Integral(tanh(a + log(x)/4)**p, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+1/4*log(x))^p,x, algorithm="giac")

[Out] integrate(tanh(a + 1/4*log(x))^p, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \tanh \left(a + \frac{\ln(x)}{4} \right)^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(a + log(x)/4)^p,x)

[Out] int(tanh(a + log(x)/4)^p, x)

3.167 $\int \tanh^p \left(a + \frac{\log(x)}{6} \right) dx$

Optimal. Leaf size=158

$$-e^{-6a} p (-1 + e^{2a \sqrt[3]{x}})^{1+p} (1 + e^{2a \sqrt[3]{x}})^{1-p} + e^{-4a} (-1 + e^{2a \sqrt[3]{x}})^{1+p} (1 + e^{2a \sqrt[3]{x}})^{1-p} \sqrt[3]{x} + \frac{2^{-p} e^{-6a} (1 + 2p^2)}{}$$

[Out] $-p(-1+\exp(2*a)*x^{(1/3)})^{(1+p)}*(1+\exp(2*a)*x^{(1/3)})^{(1-p)}/\exp(6*a)+(-1+\exp(2*a)*x^{(1/3)})^{(1+p)}*(1+\exp(2*a)*x^{(1/3)})^{(1-p)}*x^{(1/3)}/\exp(4*a)+(2*p^2+1)*(-1+\exp(2*a)*x^{(1/3)})^{(1+p)}*\text{hypergeom}([p, 1+p], [2+p], 1/2-1/2*\exp(2*a)*x^{(1/3)})/(2^p)/\exp(6*a)/(1+p)$

Rubi [A]

time = 0.09, antiderivative size = 158, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.454$, Rules used = {5652, 383, 92, 81, 71}

$$\frac{e^{-6a} 2^{-p} (2p^2 + 1) (e^{2a \sqrt[3]{x}} - 1)^{p+1} {}_2F_1(p, p+1; p+2; \frac{1}{2}(1 - e^{2a \sqrt[3]{x}}))}{p+1} - e^{-6a} p (e^{2a \sqrt[3]{x}} - 1)^{p+1} (e^{2a \sqrt[3]{x}} + 1)^{1-p} + e^{-4a} \sqrt[3]{x} (e^{2a \sqrt[3]{x}} - 1)^{p+1} (e^{2a \sqrt[3]{x}} + 1)^{1-p}$$

Antiderivative was successfully verified.

[In] Int[Tanh[a + Log[x]/6]^p, x]

[Out] $-((p*(-1 + E^{(2*a)*x^{(1/3)}})^{(1+p)}*(1 + E^{(2*a)*x^{(1/3)}})^{(1-p)})/E^{(6*a)} + ((-1 + E^{(2*a)*x^{(1/3)}})^{(1+p)}*(1 + E^{(2*a)*x^{(1/3)}})^{(1-p)}*x^{(1/3)})/E^{(4*a)} + ((1 + 2*p^2)*(-1 + E^{(2*a)*x^{(1/3)}})^{(1+p)}*\text{Hypergeometric2F1}[p, 1 + p, 2 + p, (1 - E^{(2*a)*x^{(1/3)}})/2])/(2^p * E^{(6*a)} * (1 + p))$

Rule 71

Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b/(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

Rule 81

Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)), Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]

Rule 92

```
Int[((a_.) + (b_.)*(x_))2*((c_.) + (d_.)*(x_))(n_.)*((e_.) + (f_.)*(x_))(p_.), x_Symbol] := Simp[b*(a + b*x)*(c + d*x)(n + 1)*((e + f*x)(p + 1)/(d*f*(n + p + 3)), x] + Dist[1/(d*f*(n + p + 3)), Int[(c + d*x)n*(e + f*x)p*Simp[a2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]
```

Rule 383

```
Int[((a_) + (b_.)*(x_)(n_))(p_.)*((c_) + (d_.)*(x_)(n_))(q_.), x_Symbol] := With[{g = Denominator[n]}, Dist[g, Subst[Int[x(g - 1)*(a + b*x(g*n))p*(c + d*x(g*n))q, x], x, x(1/g)], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && FractionQ[n]
```

Rule 5652

```
Int[Tanh[((a_.) + Log[x_]*(b_.))*(d_.)](p_.), x_Symbol] := Int[(-1 + E(2*a*d)*x(2*b*d))p/(1 + E(2*a*d)*x(2*b*d))p, x] /; FreeQ[{a, b, d, p}, x]
```

Rubi steps

$$\int \tanh^p \left(a + \frac{\log(x)}{6} \right) dx = \int \tanh^p \left(\frac{1}{6}(6a + \log(x)) \right) dx$$

Mathematica [C] Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

time = 2.82, size = 177, normalized size = 1.12

$$\frac{4 \left(\frac{-1+e^{2a}\sqrt[3]{x}}{1+e^{2a}\sqrt[3]{x}} \right)^p x F_1(3; -p, p; 4; e^{2a}\sqrt[3]{x}, -e^{2a}\sqrt[3]{x})}{4F_1(3; -p, p; 4; e^{2a}\sqrt[3]{x}, -e^{2a}\sqrt[3]{x}) - e^{2ap}\sqrt[3]{x} (F_1(4; 1-p, p; 5; e^{2a}\sqrt[3]{x}, -e^{2a}\sqrt[3]{x}) + F_1(4; -p, 1+p; 5; e^{2a}\sqrt[3]{x}, -e^{2a}\sqrt[3]{x}))}$$

Warning: Unable to verify antiderivative.

[In] Integrate[Tanh[a + Log[x]/6]^p, x]

[Out] (4*((-1 + E^(2*a)*x^(1/3))/(1 + E^(2*a)*x^(1/3)))^p*x*AppellF1[3, -p, p, 4, E^(2*a)*x^(1/3), -(E^(2*a)*x^(1/3))]/(4*AppellF1[3, -p, p, 4, E^(2*a)*x^(1/3), -(E^(2*a)*x^(1/3))] - E^(2*a)*p*x^(1/3)*(AppellF1[4, 1 - p, p, 5, E^(2*a)*x^(1/3), -(E^(2*a)*x^(1/3))] + AppellF1[4, -p, 1 + p, 5, E^(2*a)*x^(1/3), -(E^(2*a)*x^(1/3))]))

Maple [F]

time = 0.59, size = 0, normalized size = 0.00

$$\int \tanh^p \left(a + \frac{\ln(x)}{6} \right) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tanh(a+1/6*ln(x))^p,x)`

[Out] `int(tanh(a+1/6*ln(x))^p,x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(a+1/6*log(x))^p,x, algorithm="maxima")`

[Out] `integrate(tanh(a + 1/6*log(x))^p, x)`

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(a+1/6*log(x))^p,x, algorithm="fricas")`

[Out] `integral(tanh(a + 1/6*log(x))^p, x)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \tanh^p \left(a + \frac{\log(x)}{6} \right) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(a+1/6*ln(x))**p,x)`

[Out] `Integral(tanh(a + log(x)/6)**p, x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(a+1/6*log(x))^p,x, algorithm="giac")`

[Out] `integrate(tanh(a + 1/6*log(x))^p, x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \tanh\left(a + \frac{\ln(x)}{6}\right)^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(a + log(x)/6)^p,x)

[Out] int(tanh(a + log(x)/6)^p, x)

3.168 $\int \tanh^p \left(a + \frac{\log(x)}{8} \right) dx$

Optimal. Leaf size=190

$$\frac{1}{3}e^{-12a}(-1 + e^{2a\sqrt[4]{x}})^{1+p} (1 + e^{2a\sqrt[4]{x}})^{1-p} (e^{4a}(3 + 2p^2) - 2e^{6a}p\sqrt[4]{x}) + e^{-4a}(-1 + e^{2a\sqrt[4]{x}})^{1+p} (1 + e^{2a\sqrt[4]{x}})^{1-p}$$

[Out] $\frac{1}{3}(-1 + \exp(2*a)*x^{(1/4)})^{(1+p)}*(1 + \exp(2*a)*x^{(1/4)})^{(1-p)}*(\exp(4*a)*(2*p^2 + 3) - 2*\exp(6*a)*p*x^{(1/4)})/\exp(12*a) - 1/3*2^{(2-p)}*p*(p^2 + 2)*(-1 + \exp(2*a)*x^{(1/4)})^{(1+p)}*\text{hypergeom}([p, 1+p], [2+p], 1/2 - 1/2*\exp(2*a)*x^{(1/4)})/\exp(8*a)/(1+p) + (-1 + \exp(2*a)*x^{(1/4)})^{(1+p)}*(1 + \exp(2*a)*x^{(1/4)})^{(1-p)}*x^{(1/2)}/\exp(4*a)$

Rubi [A]

time = 0.10, antiderivative size = 190, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.454$, Rules used = {5652, 383, 102, 152, 71}

$$-\frac{e^{-8a}2^{-2-p}(p^2 + 2)(e^{2a\sqrt[4]{x}} - 1)^{p+1} {}_2F_1(p, p+1; p+2; \frac{1}{2}(1 - e^{2a\sqrt[4]{x}}))}{3(p+1)} + \frac{1}{3}e^{-12a}(e^{2a\sqrt[4]{x}} - 1)^{p+1} (e^{4a}(2p^2 + 3) - 2e^{6a}p\sqrt[4]{x})(e^{2a\sqrt[4]{x}} + 1)^{1-p} + e^{-4a}\sqrt{x}(e^{2a\sqrt[4]{x}} - 1)^{p+1} (e^{2a\sqrt[4]{x}} + 1)^{1-p}$$

Antiderivative was successfully verified.

[In] Int[Tanh[a + Log[x]/8]^p, x]

[Out] $((-1 + E^{(2*a)*x^{(1/4)}})^{(1 + p)}*(1 + E^{(2*a)*x^{(1/4)}})^{(1 - p)}*(E^{(4*a)}*(3 + 2*p^2) - 2*E^{(6*a)*p*x^{(1/4)}}))/(3*E^{(12*a)}) + ((-1 + E^{(2*a)*x^{(1/4)}})^{(1 + p)}*(1 + E^{(2*a)*x^{(1/4)}})^{(1 - p)}*\text{Sqrt}[x])/E^{(4*a)} - (2^{(2 - p)}*p*(2 + p^2)*(-1 + E^{(2*a)*x^{(1/4)}})^{(1 + p)}*\text{Hypergeometric2F1}[p, 1 + p, 2 + p, (1 - E^{(2*a)*x^{(1/4)}})/2])/(3*E^{(8*a)}*(1 + p))$

Rule 71

Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b/(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

Rule 102

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Simp[b*(a + b*x)^(m - 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(m + n + p + 1))), x] + Dist[1/(d*f*(m + n + p + 1)), Int[(a + b*x)^(m - 2)*(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(m + n + p + 1) - b*(b*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(2*m + n + p) - b*(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegerQ[m]

Rule 152

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))
*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(-(a*d*f*h*(n + 2) + b*c*f*h*(m
+ 2) - b*d*(f*g + e*h)*(m + n + 3) - b*d*f*h*(m + n + 2)*x)*(a + b*x)^(m +
1)*((c + d*x)^(n + 1)/(b^2*d^2*(m + n + 2)*(m + n + 3))), x] + Dist[(a^2*d
^2*f*h*(n + 1)*(n + 2) + a*b*d*(n + 1)*(2*c*f*h*(m + 1) - d*(f*g + e*h)*(m
+ n + 3)) + b^2*(c^2*f*h*(m + 1)*(m + 2) - c*d*(f*g + e*h)*(m + 1)*(m + n
+ 3) + d^2*e*g*(m + n + 2)*(m + n + 3))/(b^2*d^2*(m + n + 2)*(m + n + 3)),
Int[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n},
x] && NeQ[m + n + 2, 0] && NeQ[m + n + 3, 0]
```

Rule 383

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.), x_Symbol
] := With[{g = Denominator[n]}, Dist[g, Subst[Int[x^(g - 1)*(a + b*x^(g*n))
^p*(c + d*x^(g*n))^q, x], x, x^(1/g)], x]] /; FreeQ[{a, b, c, d, p, q}, x]
&& NeQ[b*c - a*d, 0] && FractionQ[n]
```

Rule 5652

```
Int[Tanh[((a_.) + Log[x_]*(b_.))*(d_.)]^(p_.), x_Symbol] := Int[(-1 + E^(2*
a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p, x] /; FreeQ[{a, b, d, p}, x]
```

Rubi steps

$$\int \tanh^p \left(a + \frac{\log(x)}{8} \right) dx = \int \tanh^p \left(\frac{1}{8}(8a + \log(x)) \right) dx$$

Mathematica [C] Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

time = 2.24, size = 177, normalized size = 0.93

$$\frac{5 \left(\frac{-1+e^{2a}\sqrt[4]{x}}{1+e^{2a}\sqrt[4]{x}} \right)^p x F_1(4; -p, p; 5; e^{2a}\sqrt[4]{x}, -e^{2a}\sqrt[4]{x})}{5F_1(4; -p, p; 5; e^{2a}\sqrt[4]{x}, -e^{2a}\sqrt[4]{x}) - e^{2ap}\sqrt[4]{x} (F_1(5; 1-p, p; 6; e^{2a}\sqrt[4]{x}, -e^{2a}\sqrt[4]{x}) + F_1(5; -p, 1+p; 6; e^{2a}\sqrt[4]{x}, -e^{2a}\sqrt[4]{x}))}$$

Warning: Unable to verify antiderivative.

```
[In] Integrate[Tanh[a + Log[x]/8]^p, x]
```

```
[Out] (5*((-1 + E^(2*a)*x^(1/4))/(1 + E^(2*a)*x^(1/4)))^p*x*AppellF1[4, -p, p, 5,
E^(2*a)*x^(1/4), -(E^(2*a)*x^(1/4))]/(5*AppellF1[4, -p, p, 5, E^(2*a)*x^(
1/4), -(E^(2*a)*x^(1/4))] - E^(2*a)*p*x^(1/4)*(AppellF1[5, 1 - p, p, 6, E^(
```

$2*a)*x^{(1/4)}, -(E^{(2*a)*x^{(1/4)}})] + \text{AppellF1}[5, -p, 1 + p, 6, E^{(2*a)*x^{(1/4)}}, -(E^{(2*a)*x^{(1/4)}})]])$

Maple [F]

time = 0.63, size = 0, normalized size = 0.00

$$\int \tanh^p \left(a + \frac{\ln(x)}{8} \right) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tanh(a+1/8*ln(x))^p,x)`

[Out] `int(tanh(a+1/8*ln(x))^p,x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(a+1/8*log(x))^p,x, algorithm="maxima")`

[Out] `integrate(tanh(a + 1/8*log(x))^p, x)`

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(a+1/8*log(x))^p,x, algorithm="fricas")`

[Out] `integral(tanh(a + 1/8*log(x))^p, x)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \tanh^p \left(a + \frac{\log(x)}{8} \right) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(a+1/8*ln(x))**p,x)`

[Out] `Integral(tanh(a + log(x)/8)**p, x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(tanh(a+1/8*log(x))^p,x, algorithm="giac")
```

```
[Out] integrate(tanh(a + 1/8*log(x))^p, x)
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \tanh\left(a + \frac{\ln(x)}{8}\right)^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(tanh(a + log(x)/8)^p,x)
```

```
[Out] int(tanh(a + log(x)/8)^p, x)
```

3.169 $\int \tanh^p(a + \log(x)) dx$

Optimal. Leaf size=61

$$x(1 - e^{2a}x^2)^{-p} (-1 + e^{2a}x^2)^p F_1\left(\frac{1}{2}; -p, p; \frac{3}{2}; e^{2a}x^2, -e^{2a}x^2\right)$$

[Out] $x*(-1+\exp(2*a)*x^2)^p*\text{AppellF1}(1/2, -p, p, 3/2, \exp(2*a)*x^2, -\exp(2*a)*x^2)/((1-\exp(2*a)*x^2)^p)$

Rubi [A]

time = 0.02, antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {5652, 441, 440}

$$x(1 - e^{2a}x^2)^{-p} (e^{2a}x^2 - 1)^p F_1\left(\frac{1}{2}; -p, p; \frac{3}{2}; e^{2a}x^2, -e^{2a}x^2\right)$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Tanh}[a + \text{Log}[x]]^p, x]$

[Out] $(x*(-1 + E^{(2*a)*x^2})^p*\text{AppellF1}[1/2, -p, p, 3/2, E^{(2*a)*x^2}, -(E^{(2*a)*x^2} - 1)])/(1 - E^{(2*a)*x^2})^p$

Rule 440

$\text{Int}[(a_+ + (b_+)*(x_+)^{(n_+)})^{(p_+)}*((c_+ + (d_+)*(x_+)^{(n_+)})^{(q_+)}, x_Symbol]$
 $\rightarrow \text{Simp}[a^p*c^q*x*\text{AppellF1}[1/n, -p, -q, 1 + 1/n, (-b)*(x^n/a), (-d)*(x^n/c)], x] /;$ $\text{FreeQ}\{a, b, c, d, n, p, q\}, x]$ && $\text{NeQ}[b*c - a*d, 0]$ && $\text{NeQ}[n, -1]$ && $(\text{IntegerQ}[p] \parallel \text{GtQ}[a, 0])$ && $(\text{IntegerQ}[q] \parallel \text{GtQ}[c, 0])$

Rule 441

$\text{Int}[(a_+ + (b_+)*(x_+)^{(n_+)})^{(p_+)}*((c_+ + (d_+)*(x_+)^{(n_+)})^{(q_+)}, x_Symbol]$
 $\rightarrow \text{Dist}[a^{\text{IntPart}[p]}*((a + b*x^n)^{\text{FracPart}[p]}/(1 + b*(x^n/a))^{\text{FracPart}[p]}], \text{Int}[(1 + b*(x^n/a))^p*(c + d*x^n)^q, x], x] /;$ $\text{FreeQ}\{a, b, c, d, n, p, q\}, x]$ && $\text{NeQ}[b*c - a*d, 0]$ && $\text{NeQ}[n, -1]$ && $!(\text{IntegerQ}[p] \parallel \text{GtQ}[a, 0])$

Rule 5652

$\text{Int}[\text{Tanh}[(a_+ + \text{Log}[x_+]*(b_+))*(d_+)]^{(p_+)}, x_Symbol]$ $\rightarrow \text{Int}[(-1 + E^{(2*a*d)*x^{(2*b*d)}})^p/(1 + E^{(2*a*d)*x^{(2*b*d)}})^p, x] /;$ $\text{FreeQ}\{a, b, d, p\}, x]$

Rubi steps

$$\int \tanh^p(a + \log(x)) dx = \int \tanh^p(a + \log(x)) dx$$

Mathematica [B] Leaf count is larger than twice the leaf count of optimal. 171 vs. 2(61) = 122.

time = 1.17, size = 171, normalized size = 2.80

$$\frac{3x \left(\frac{-1+e^{2ax^2}}{1+e^{2ax^2}} \right)^p F_1\left(\frac{1}{2}; -p, p; \frac{3}{2}; e^{2ax^2}, -e^{2ax^2}\right)}{3F_1\left(\frac{1}{2}; -p, p; \frac{3}{2}; e^{2ax^2}, -e^{2ax^2}\right) - 2e^{2apx^2} \left(F_1\left(\frac{3}{2}; 1-p, p; \frac{5}{2}; e^{2ax^2}, -e^{2ax^2}\right) + F_1\left(\frac{3}{2}; -p, 1+p; \frac{5}{2}; e^{2ax^2}, -e^{2ax^2}\right) \right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[Tanh[a + Log[x]]^p, x]

[Out] $(3x * ((-1 + E^{(2*a)*x^2}) / (1 + E^{(2*a)*x^2}))^p * \text{AppellF1}[1/2, -p, p, 3/2, E^{(2*a)*x^2}, -(E^{(2*a)*x^2})]) / (3 * \text{AppellF1}[1/2, -p, p, 3/2, E^{(2*a)*x^2}, -(E^{(2*a)*x^2})] - 2 * E^{(2*a)*p*x^2} * (\text{AppellF1}[3/2, 1 - p, p, 5/2, E^{(2*a)*x^2}, -(E^{(2*a)*x^2})] + \text{AppellF1}[3/2, -p, 1 + p, 5/2, E^{(2*a)*x^2}, -(E^{(2*a)*x^2})]))$

Maple [F]

time = 0.64, size = 0, normalized size = 0.00

$$\int \tanh^p(a + \ln(x)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(a+ln(x))^p, x)

[Out] int(tanh(a+ln(x))^p, x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+log(x))^p, x, algorithm="maxima")

[Out] integrate(tanh(a + log(x))^p, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+log(x))^p, x, algorithm="fricas")

[Out] integral(tanh(a + log(x))^p, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \tanh^p(a + \log(x)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+ln(x))**p,x)

[Out] Integral(tanh(a + log(x))**p, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+log(x))^p,x, algorithm="giac")

[Out] integrate(tanh(a + log(x))^p, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \tanh(a + \ln(x))^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(a + log(x))^p,x)

[Out] int(tanh(a + log(x))^p, x)

3.170 $\int \tanh^p(a + 2 \log(x)) dx$

Optimal. Leaf size=61

$$x(1 - e^{2a}x^4)^{-p} (-1 + e^{2a}x^4)^p F_1\left(\frac{1}{4}; -p, p; \frac{5}{4}; e^{2a}x^4, -e^{2a}x^4\right)$$

[Out] $x*(-1+\exp(2*a)*x^4)^p*\text{AppellF1}(1/4, -p, p, 5/4, \exp(2*a)*x^4, -\exp(2*a)*x^4)/((1-\exp(2*a)*x^4)^p)$

Rubi [A]

time = 0.02, antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {5652, 441, 440}

$$x(1 - e^{2a}x^4)^{-p} (e^{2a}x^4 - 1)^p F_1\left(\frac{1}{4}; -p, p; \frac{5}{4}; e^{2a}x^4, -e^{2a}x^4\right)$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Tanh}[a + 2*\text{Log}[x]]^p, x]$

[Out] $(x*(-1 + E^{(2*a)*x^4})^p*\text{AppellF1}[1/4, -p, p, 5/4, E^{(2*a)*x^4}, -(E^{(2*a)*x^4})])/(1 - E^{(2*a)*x^4})^p$

Rule 440

$\text{Int}[(a_.) + (b_.)*(x_.)^{(n_.)}]^{(p_.)}*((c_.) + (d_.)*(x_.)^{(n_.)})^{(q_.)}, x_Symbol]$
 $\rightarrow \text{Simp}[a^p*c^q*x*\text{AppellF1}[1/n, -p, -q, 1 + 1/n, (-b)*(x^n/a), (-d)*(x^n/c)], x] /;$
 $\text{FreeQ}\{a, b, c, d, n, p, q\}, x \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[n, -1] \ \&\& \ (\text{IntegerQ}[p] \ || \ \text{GtQ}[a, 0]) \ \&\& \ (\text{IntegerQ}[q] \ || \ \text{GtQ}[c, 0])$

Rule 441

$\text{Int}[(a_.) + (b_.)*(x_.)^{(n_.)}]^{(p_.)}*((c_.) + (d_.)*(x_.)^{(n_.)})^{(q_.)}, x_Symbol]$
 $\rightarrow \text{Dist}[a^{\text{IntPart}[p]}*((a + b*x^n)^{\text{FracPart}[p]}/(1 + b*(x^n/a))^{\text{FracPart}[p]}], \text{Int}[(1 + b*(x^n/a))^p*(c + d*x^n)^q, x], x] /;$
 $\text{FreeQ}\{a, b, c, d, n, p, q\}, x \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[n, -1] \ \&\& \ !(\text{IntegerQ}[p] \ || \ \text{GtQ}[a, 0])$

Rule 5652

$\text{Int}[\text{Tanh}[(a_.) + \text{Log}[x_]* (b_.)]^{(p_.)}*(d_.)^{(p_.)}, x_Symbol] \rightarrow \text{Int}[(-1 + E^{(2*a*d)*x^{(2*b*d)}})^p/(1 + E^{(2*a*d)*x^{(2*b*d)}})^p, x] /;$
 $\text{FreeQ}\{a, b, d, p\}, x]$

Rubi steps

$$\int \tanh^p(a + 2 \log(x)) dx = \int \tanh^p(a + 2 \log(x)) dx$$

Mathematica [B] Leaf count is larger than twice the leaf count of optimal. 171 vs. $2(61) = 122$.

time = 1.32, size = 171, normalized size = 2.80

$$\frac{5x \left(\frac{-1+e^{2ax^4}}{1+e^{2ax^4}} \right)^p F_1\left(\frac{1}{4}; -p, p; \frac{5}{4}; e^{2ax^4}, -e^{2ax^4}\right)}{5F_1\left(\frac{1}{4}; -p, p; \frac{5}{4}; e^{2ax^4}, -e^{2ax^4}\right) - 4e^{2apx^4} \left(F_1\left(\frac{5}{4}; 1-p, p; \frac{9}{4}; e^{2ax^4}, -e^{2ax^4}\right) + F_1\left(\frac{5}{4}; -p, 1+p; \frac{9}{4}; e^{2ax^4}, -e^{2ax^4}\right) \right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[Tanh[a + 2*Log[x]]^p,x]

[Out] $(5*x*((-1 + E^{(2*a)*x^4})/(1 + E^{(2*a)*x^4}))^p*AppellF1[1/4, -p, p, 5/4, E^{(2*a)*x^4}, -(E^{(2*a)*x^4})])/(5*AppellF1[1/4, -p, p, 5/4, E^{(2*a)*x^4}, -(E^{(2*a)*x^4})] - 4*E^{(2*a)*p*x^4}*(AppellF1[5/4, 1 - p, p, 9/4, E^{(2*a)*x^4}, -(E^{(2*a)*x^4})] + AppellF1[5/4, -p, 1 + p, 9/4, E^{(2*a)*x^4}, -(E^{(2*a)*x^4})]))$

Maple [F]

time = 0.62, size = 0, normalized size = 0.00

$$\int \tanh^p(a + 2 \ln(x)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(a+2*ln(x))^p,x)

[Out] int(tanh(a+2*ln(x))^p,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+2*log(x))^p,x, algorithm="maxima")

[Out] integrate(tanh(a + 2*log(x))^p, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+2*log(x))^p,x, algorithm="fricas")

[Out] integral(tanh(a + 2*log(x))^p, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \tanh^p(a + 2 \log(x)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+2*ln(x))**p,x)**[Out]** Integral(tanh(a + 2*log(x))**p, x)**Giac [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+2*log(x))^p,x, algorithm="giac")**[Out]** integrate(tanh(a + 2*log(x))^p, x)**Mupad [F]**

time = 0.00, size = -1, normalized size = -0.02

$$\int \tanh(a + 2 \ln(x))^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(a + 2*log(x))^p,x)**[Out]** int(tanh(a + 2*log(x))^p, x)

3.171 $\int \tanh^p(a + 3 \log(x)) dx$

Optimal. Leaf size=61

$$x(1 - e^{2a}x^6)^{-p} (-1 + e^{2a}x^6)^p F_1\left(\frac{1}{6}; -p, p; \frac{7}{6}; e^{2a}x^6, -e^{2a}x^6\right)$$

[Out] $x*(-1+\exp(2*a)*x^6)^p*\text{AppellF1}(1/6, -p, p, 7/6, \exp(2*a)*x^6, -\exp(2*a)*x^6)/((1-\exp(2*a)*x^6)^p)$

Rubi [A]

time = 0.02, antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {5652, 441, 440}

$$x(1 - e^{2a}x^6)^{-p} (e^{2a}x^6 - 1)^p F_1\left(\frac{1}{6}; -p, p; \frac{7}{6}; e^{2a}x^6, -e^{2a}x^6\right)$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Tanh}[a + 3*\text{Log}[x]]^p, x]$

[Out] $(x*(-1 + E^{(2*a)*x^6})^p*\text{AppellF1}[1/6, -p, p, 7/6, E^{(2*a)*x^6}, -(E^{(2*a)*x^6})])/(1 - E^{(2*a)*x^6})^p$

Rule 440

$\text{Int}[(a_+ + (b_+)*(x_+)^{(n_+)})^{(p_+)}*((c_+ + (d_+)*(x_+)^{(n_+)})^{(q_+)}, x_Symbol]$
 $\rightarrow \text{Simp}[a^p*c^q*x*\text{AppellF1}[1/n, -p, -q, 1 + 1/n, (-b)*(x^n/a), (-d)*(x^n/c)], x] /;$ $\text{FreeQ}\{a, b, c, d, n, p, q\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[n, -1]$
 $\ \&\& \ (\text{IntegerQ}[p] \ || \ \text{GtQ}[a, 0]) \ \&\& \ (\text{IntegerQ}[q] \ || \ \text{GtQ}[c, 0])$

Rule 441

$\text{Int}[(a_+ + (b_+)*(x_+)^{(n_+)})^{(p_+)}*((c_+ + (d_+)*(x_+)^{(n_+)})^{(q_+)}, x_Symbol]$
 $\rightarrow \text{Dist}[a^{\text{IntPart}[p]}*((a + b*x^n)^{\text{FracPart}[p]}/(1 + b*(x^n/a))^{\text{FracPart}[p]}],$
 $\text{Int}[(1 + b*(x^n/a))^p*(c + d*x^n)^q, x], x] /;$ $\text{FreeQ}\{a, b, c, d, n, p, q\}, x]$
 $\ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[n, -1] \ \&\& \ !(\text{IntegerQ}[p] \ || \ \text{GtQ}[a, 0])$

Rule 5652

$\text{Int}[\text{Tanh}[(a_+ + \text{Log}[x_+]*(b_+))*(d_+)]^{(p_+)}, x_Symbol] \rightarrow \text{Int}[(-1 + E^{(2*a*d)*x^{(2*b*d)}})^p/(1 + E^{(2*a*d)*x^{(2*b*d)}})^p, x] /;$ $\text{FreeQ}\{a, b, d, p\}, x]$

Rubi steps

$$\int \tanh^p(a + 3 \log(x)) dx = \int \tanh^p(a + 3 \log(x)) dx$$

Mathematica [B] Leaf count is larger than twice the leaf count of optimal. 171 vs. 2(61) = 122.

time = 1.36, size = 171, normalized size = 2.80

$$\frac{7x \left(\frac{-1+e^{2a}x^6}{1+e^{2a}x^6} \right)^p F_1\left(\frac{1}{6}; -p, p; \frac{7}{6}; e^{2a}x^6, -e^{2a}x^6\right)}{7F_1\left(\frac{1}{6}; -p, p; \frac{7}{6}; e^{2a}x^6, -e^{2a}x^6\right) - 6e^{2a}px^6 \left(F_1\left(\frac{7}{6}; 1-p, p; \frac{13}{6}; e^{2a}x^6, -e^{2a}x^6\right) + F_1\left(\frac{7}{6}; -p, 1+p; \frac{13}{6}; e^{2a}x^6, -e^{2a}x^6\right) \right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[Tanh[a + 3*Log[x]]^p, x]

[Out] (7*x*((-1 + E^(2*a)*x^6)/(1 + E^(2*a)*x^6))^p*AppellF1[1/6, -p, p, 7/6, E^(2*a)*x^6, -(E^(2*a)*x^6)]/(7*AppellF1[1/6, -p, p, 7/6, E^(2*a)*x^6, -(E^(2*a)*x^6)] - 6*E^(2*a)*p*x^6*(AppellF1[7/6, 1 - p, p, 13/6, E^(2*a)*x^6, -(E^(2*a)*x^6)] + AppellF1[7/6, -p, 1 + p, 13/6, E^(2*a)*x^6, -(E^(2*a)*x^6)]))

Maple [F]

time = 0.60, size = 0, normalized size = 0.00

$$\int \tanh^p(a + 3 \ln(x)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(a+3*ln(x))^p, x)

[Out] int(tanh(a+3*ln(x))^p, x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+3*log(x))^p, x, algorithm="maxima")

[Out] integrate(tanh(a + 3*log(x))^p, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+3*log(x))^p, x, algorithm="fricas")

[Out] integral(tanh(a + 3*log(x))^p, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \tanh^p(a + 3 \log(x)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+3*ln(x))**p,x)

[Out] Integral(tanh(a + 3*log(x))**p, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+3*log(x))^p,x, algorithm="giac")

[Out] integrate(tanh(a + 3*log(x))^p, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \tanh(a + 3 \ln(x))^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(a + 3*log(x))^p,x)

[Out] int(tanh(a + 3*log(x))^p, x)

3.172 $\int x^3 \tanh(d(a + b \log(cx^n))) dx$

Optimal. Leaf size=59

$$\frac{x^4}{4} - \frac{1}{2}x^4 {}_2F_1\left(1, \frac{2}{bdn}; 1 + \frac{2}{bdn}; -e^{2ad}(cx^n)^{2bd}\right)$$

[Out] 1/4*x^4-1/2*x^4*hypergeom([1, 2/b/d/n], [1+2/b/d/n], -exp(2*a*d)*(c*x^n)^(2*b*d))

Rubi [A]

time = 0.05, antiderivative size = 59, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {5658, 5656, 470, 371}

$$\frac{x^4}{4} - \frac{1}{2}x^4 {}_2F_1\left(1, \frac{2}{bdn}; 1 + \frac{2}{bdn}; -e^{2ad}(cx^n)^{2bd}\right)$$

Antiderivative was successfully verified.

[In] Int[x^3*Tanh[d*(a + b*Log[c*x^n])],x]

[Out] x^4/4 - (x^4*Hypergeometric2F1[1, 2/(b*d*n), 1 + 2/(b*d*n), -(E^(2*a*d)*(c*x^n)^(2*b*d))])/2

Rule 371

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])

Rule 470

Int[((e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> Simp[d*(e*x)^(m + 1)*((a + b*x^n)^(p + 1)/(b*e*(m + n*(p + 1) + 1))), x] - Dist[(a*d*(m + 1) - b*c*(m + n*(p + 1) + 1))/(b*(m + n*(p + 1) + 1)), Int[(e*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && NeQ[b*c - a*d, 0] && NeQ[m + n*(p + 1) + 1, 0]

Rule 5656

Int[((e_.)*(x_))^(m_.)*Tanh[((a_.) + Log[x_]*(b_.))*(d_.)]^(p_.), x_Symbol] :> Int[(e*x)^m*((-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p), x] /; FreeQ[{a, b, d, e, m, p}, x]

Rule 5658


```
Int[((e_.)*(x_))^(m_.)*Tanh[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*(d_.)]^(p
_.), x_Symbol] :> Dist[(e*x)^(m + 1)/(e*n*(c*x^n)^((m + 1)/n)), Subst[Int[x
^((m + 1)/n - 1)*Tanh[d*(a + b*Log[x])]^p, x], x, c*x^n], x] /; FreeQ[{a, b
, c, d, e, m, n, p}, x] && (NeQ[c, 1] || NeQ[n, 1])
```

Rubi steps

$$\int x^3 \tanh(d(a + b \log(cx^n))) dx = \int x^3 \tanh(d(a + b \log(cx^n))) dx$$

Mathematica [B] Leaf count is larger than twice the leaf count of optimal. 127 vs. 2(59) = 118.

time = 5.75, size = 127, normalized size = 2.15

$$\frac{x^4 (2e^{2d(a+b \log(cx^n))} {}_2F_1(1, 1 + \frac{2}{bdn}; 2 + \frac{2}{bdn}; -e^{2d(a+b \log(cx^n))}) - (2 + bdn) {}_2F_1(1, \frac{2}{bdn}; 1 + \frac{2}{bdn}; -e^{2d(a+b \log(cx^n))}))}{8 + 4bdn}$$

Antiderivative was successfully verified.

```
[In] Integrate[x^3*Tanh[d*(a + b*Log[c*x^n])],x]
```

```
[Out] (x^4*(2*E^(2*d*(a + b*Log[c*x^n]))*Hypergeometric2F1[1, 1 + 2/(b*d*n), 2 + 2/(b*d*n), -E^(2*d*(a + b*Log[c*x^n]))] - (2 + b*d*n)*Hypergeometric2F1[1, 2/(b*d*n), 1 + 2/(b*d*n), -E^(2*d*(a + b*Log[c*x^n]))]))/(8 + 4*b*d*n)
```

Maple [F]

time = 1.11, size = 0, normalized size = 0.00

$$\int x^3 \tanh(d(a + b \ln(cx^n))) dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(x^3*tanh(d*(a+b*ln(c*x^n))),x)
```

```
[Out] int(x^3*tanh(d*(a+b*ln(c*x^n))),x)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x^3*tanh(d*(a+b*log(c*x^n))),x, algorithm="maxima")
```

```
[Out] 1/4*x^4 - 2*integrate(x^3/(c^(2*b*d)*e^(2*b*d*log(x^n) + 2*a*d) + 1), x)
```

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(x^3*tanh(d*(a+b*log(c*x^n))),x, algorithm="fricas")``[Out] integral(x^3*tanh(b*d*log(c*x^n) + a*d), x)`**Sympy [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int x^3 \tanh(ad + bd \log(cx^n)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(x**3*tanh(d*(a+b*ln(c*x**n))),x)``[Out] Integral(x**3*tanh(a*d + b*d*log(c*x**n)), x)`**Giac [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(x^3*tanh(d*(a+b*log(c*x^n))),x, algorithm="giac")``[Out] integrate(x^3*tanh((b*log(c*x^n) + a)*d), x)`**Mupad [F]**

time = 0.00, size = -1, normalized size = -0.02

$$\int x^3 \tanh(d(a + b \ln(cx^n))) dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(x^3*tanh(d*(a + b*log(c*x^n))),x)``[Out] int(x^3*tanh(d*(a + b*log(c*x^n))), x)`

3.173 $\int x^2 \tanh(d(a + b \log(cx^n))) dx$

Optimal. Leaf size=63

$$\frac{x^3}{3} - \frac{2}{3}x^3 {}_2F_1\left(1, \frac{3}{2bdn}; 1 + \frac{3}{2bdn}; -e^{2ad}(cx^n)^{2bd}\right)$$

[Out] 1/3*x^3-2/3*x^3*hypergeom([1, 3/2/b/d/n], [1+3/2/b/d/n], -exp(2*a*d)*(c*x^n)^(2*b*d))

Rubi [A]

time = 0.04, antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {5658, 5656, 470, 371}

$$\frac{x^3}{3} - \frac{2}{3}x^3 {}_2F_1\left(1, \frac{3}{2bdn}; 1 + \frac{3}{2bdn}; -e^{2ad}(cx^n)^{2bd}\right)$$

Antiderivative was successfully verified.

[In] Int[x^2*Tanh[d*(a + b*Log[c*x^n])],x]

[Out] x^3/3 - (2*x^3*Hypergeometric2F1[1, 3/(2*b*d*n), 1 + 3/(2*b*d*n), -(E^(2*a*d)*(c*x^n)^(2*b*d))])/3

Rule 371

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[a^p * ((c*x)^(m + 1)/(c*(m + 1))) * Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])

Rule 470

Int[((e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> Simp[d*(e*x)^(m + 1)*((a + b*x^n)^(p + 1)/(b*e*(m + n*(p + 1) + 1))), x] - Dist[(a*d*(m + 1) - b*c*(m + n*(p + 1) + 1))/(b*(m + n*(p + 1) + 1)), Int[(e*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && NeQ[b*c - a*d, 0] && NeQ[m + n*(p + 1) + 1, 0]

Rule 5656

Int[((e_.)*(x_))^(m_.)*Tanh[((a_.) + Log[x_]*(b_.))*(d_.)]^(p_.), x_Symbol] :> Int[(e*x)^m*((-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p), x] /; FreeQ[{a, b, d, e, m, p}, x]

Rule 5658

```
Int[((e_.)*(x_))^(m_.)*Tanh[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*(d_.)]^(p
_.), x_Symbol] :> Dist[(e*x)^(m + 1)/(e*n*(c*x^n)^(m + 1)/n), Subst[Int[x
^(m + 1)/n - 1)*Tanh[d*(a + b*Log[x])]^p, x], x, c*x^n], x] /; FreeQ[{a, b
, c, d, e, m, n, p}, x] && (NeQ[c, 1] || NeQ[n, 1])
```

Rubi steps

$$\int x^2 \tanh(d(a + b \log(cx^n))) dx = \int x^2 \tanh(d(a + b \log(cx^n))) dx$$

Mathematica [B] Leaf count is larger than twice the leaf count of optimal. 136 vs. 2(63) = 126.

time = 5.50, size = 136, normalized size = 2.16

$$\frac{x^3 (3e^{2d(a+b \log(cx^n))} {}_2F_1(1, 1 + \frac{3}{2bdn}; 2 + \frac{3}{2bdn}; -e^{2d(a+b \log(cx^n))}) - (3 + 2bdn) {}_2F_1(1, \frac{3}{2bdn}; 1 + \frac{3}{2bdn}; -e^{2d(a+b \log(cx^n))}))}{9 + 6bdn}$$

Antiderivative was successfully verified.

```
[In] Integrate[x^2*Tanh[d*(a + b*Log[c*x^n])],x]
```

```
[Out] (x^3*(3*E^(2*d*(a + b*Log[c*x^n]))*Hypergeometric2F1[1, 1 + 3/(2*b*d*n), 2
+ 3/(2*b*d*n), -E^(2*d*(a + b*Log[c*x^n])]) - (3 + 2*b*d*n)*Hypergeometric2
F1[1, 3/(2*b*d*n), 1 + 3/(2*b*d*n), -E^(2*d*(a + b*Log[c*x^n])])]))/(9 + 6*b
*d*n)
```

Maple [F]

time = 0.82, size = 0, normalized size = 0.00

$$\int x^2 \tanh(d(a + b \ln(cx^n))) dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(x^2*tanh(d*(a+b*ln(c*x^n))),x)
```

```
[Out] int(x^2*tanh(d*(a+b*ln(c*x^n))),x)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x^2*tanh(d*(a+b*log(c*x^n))),x, algorithm="maxima")
```

[Out] $1/3*x^3 - 2*\text{integrate}(x^2/(c^{(2*b*d)}*e^{(2*b*d*\log(x^n) + 2*a*d) + 1}), x)$

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^2*tanh(d*(a+b*log(c*x^n))),x, algorithm="fricas")`

[Out] `integral(x^2*tanh(b*d*log(c*x^n) + a*d), x)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x^2 \tanh(ad + bd \log(cx^n)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x**2*tanh(d*(a+b*ln(c*x**n))),x)`

[Out] `Integral(x**2*tanh(a*d + b*d*log(c*x**n)), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^2*tanh(d*(a+b*log(c*x^n))),x, algorithm="giac")`

[Out] `integrate(x^2*tanh((b*log(c*x^n) + a)*d), x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int x^2 \tanh(d(a + b \ln(cx^n))) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x^2*tanh(d*(a + b*log(c*x^n))),x)`

[Out] `int(x^2*tanh(d*(a + b*log(c*x^n))), x)`

3.174 $\int x \tanh(d(a + b \log(cx^n))) dx$

Optimal. Leaf size=55

$$\frac{x^2}{2} - x^2 {}_2F_1\left(1, \frac{1}{bdn}; 1 + \frac{1}{bdn}; -e^{2ad}(cx^n)^{2bd}\right)$$

[Out] 1/2*x^2-x^2*hypergeom([1, 1/b/d/n], [1+1/b/d/n], -exp(2*a*d)*(c*x^n)^(2*b*d))

Rubi [A]

time = 0.04, antiderivative size = 55, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {5658, 5656, 470, 371}

$$\frac{x^2}{2} - x^2 {}_2F_1\left(1, \frac{1}{bdn}; 1 + \frac{1}{bdn}; -e^{2ad}(cx^n)^{2bd}\right)$$

Antiderivative was successfully verified.

[In] Int[x*Tanh[d*(a + b*Log[c*x^n])], x]

[Out] x^2/2 - x^2*Hypergeometric2F1[1, 1/(b*d*n), 1 + 1/(b*d*n), -(E^(2*a*d)*(c*x^n)^(2*b*d))]

Rule 371

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])

Rule 470

Int[((e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> Simp[d*(e*x)^(m + 1)*((a + b*x^n)^(p + 1)/(b*e*(m + n*(p + 1) + 1))), x] - Dist[(a*d*(m + 1) - b*c*(m + n*(p + 1) + 1))/(b*(m + n*(p + 1) + 1)), Int[(e*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && NeQ[b*c - a*d, 0] && NeQ[m + n*(p + 1) + 1, 0]

Rule 5656

Int[((e_.)*(x_))^(m_.)*Tanh[((a_.) + Log[x_]*(b_.))*(d_.)]^(p_.), x_Symbol] :> Int[(e*x)^m*((-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p), x] /; FreeQ[{a, b, d, e, m, p}, x]

Rule 5658

Int[((e_.)*(x_))^(m_.)*Tanh[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*(d_.)]^(p_.), x_Symbol] :> Dist[(e*x)^(m + 1)/(e*n*(c*x^n)^(m + 1)/n), Subst[Int[x

$\wedge((m + 1)/n - 1) * \text{Tanh}[d*(a + b*\text{Log}[x])]^p, x], x, c*x^n], x] /; \text{FreeQ}\{a, b, c, d, e, m, n, p\}, x] \&\& (\text{NeQ}[c, 1] \|\| \text{NeQ}[n, 1])$

Rubi steps

$$\int x \tanh(d(a + b \log(cx^n))) dx = \int x \tanh(d(a + b \log(cx^n))) dx$$

Mathematica [B] Leaf count is larger than twice the leaf count of optimal. 122 vs. 2(55) = 110.

time = 5.34, size = 122, normalized size = 2.22

$$\frac{x^2 (e^{2d(a+b \log(cx^n))} {}_2F_1(1, 1 + \frac{1}{bdn}; 2 + \frac{1}{bdn}; -e^{2d(a+b \log(cx^n))}) - (1 + bdn) {}_2F_1(1, \frac{1}{bdn}; 1 + \frac{1}{bdn}; -e^{2d(a+b \log(cx^n))}))}{2 + 2bdn}$$

Antiderivative was successfully verified.

[In] Integrate[x*Tanh[d*(a + b*Log[c*x^n])],x]

[Out] (x^2*(E^(2*d*(a + b*Log[c*x^n]))*Hypergeometric2F1[1, 1 + 1/(b*d*n), 2 + 1/(b*d*n), -E^(2*d*(a + b*Log[c*x^n]))] - (1 + b*d*n)*Hypergeometric2F1[1, 1/(b*d*n), 1 + 1/(b*d*n), -E^(2*d*(a + b*Log[c*x^n]))]))/(2 + 2*b*d*n)

Maple [F]

time = 0.82, size = 0, normalized size = 0.00

$$\int x \tanh(d(a + b \ln(cx^n))) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x*tanh(d*(a+b*ln(c*x^n))),x)

[Out] int(x*tanh(d*(a+b*ln(c*x^n))),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x*tanh(d*(a+b*log(c*x^n))),x, algorithm="maxima")

[Out] 1/2*x^2 - 2*integrate(x/(c^(2*b*d)*e^(2*b*d*log(x^n) + 2*a*d) + 1), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(x*tanh(d*(a+b*log(c*x^n))),x, algorithm="fricas")``[Out] integral(x*tanh(b*d*log(c*x^n) + a*d), x)`**Sympy [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int x \tanh(ad + bd \log(cx^n)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(x*tanh(d*(a+b*ln(c*x**n))),x)``[Out] Integral(x*tanh(a*d + b*d*log(c*x**n)), x)`**Giac [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(x*tanh(d*(a+b*log(c*x^n))),x, algorithm="giac")``[Out] integrate(x*tanh((b*log(c*x^n) + a)*d), x)`**Mupad [F]**

time = 0.00, size = -1, normalized size = -0.02

$$\int x \tanh(d(a + b \ln(cx^n))) dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(x*tanh(d*(a + b*log(c*x^n))),x)``[Out] int(x*tanh(d*(a + b*log(c*x^n))), x)`

3.175 $\int \tanh(d(a + b \log(cx^n))) dx$

Optimal. Leaf size=53

$$x - 2x {}_2F_1\left(1, \frac{1}{2bdn}; 1 + \frac{1}{2bdn}; -e^{2ad}(cx^n)^{2bd}\right)$$

[Out] $x - 2*x*\text{hypergeom}([1, 1/2/b/d/n], [1+1/2/b/d/n], -\exp(2*a*d)*(c*x^n)^{(2*b*d)})$

Rubi [A]

time = 0.04, antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {5654, 5656, 470, 371}

$$x - 2x {}_2F_1\left(1, \frac{1}{2bdn}; 1 + \frac{1}{2bdn}; -e^{2ad}(cx^n)^{2bd}\right)$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Tanh}[d*(a + b*\text{Log}[c*x^n])], x]$

[Out] $x - 2*x*\text{Hypergeometric2F1}[1, 1/(2*b*d*n), 1 + 1/(2*b*d*n), -(E^{(2*a*d)}*(c*x^n)^{(2*b*d)})]$

Rule 371

$\text{Int}[(c_*)*(x_)^{(m_*)}*((a_) + (b_*)*(x_)^{(n_)})^{(p_)}, x_Symbol] \rightarrow \text{Simp}[a^p * ((c*x)^{(m+1})/(c*(m+1)))*\text{Hypergeometric2F1}[-p, (m+1)/n, (m+1)/n + 1, (-b)*(x^n/a)], x] /;$ FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])

Rule 470

$\text{Int}[(e_*)*(x_)^{(m_*)}*((a_) + (b_*)*(x_)^{(n_)})^{(p_*)}*((c_) + (d_*)*(x_)^{(n_*)}), x_Symbol] \rightarrow \text{Simp}[d*(e*x)^{(m+1)}*((a + b*x^n)^{(p+1)})/(b*e*(m+n*(p+1)+1)), x] - \text{Dist}[(a*d*(m+1) - b*c*(m+n*(p+1)+1))/(b*(m+n*(p+1)+1)), \text{Int}[(e*x)^m*(a + b*x^n)^p, x], x] /;$ FreeQ[{a, b, c, d, e, m, n, p}, x] && NeQ[b*c - a*d, 0] && NeQ[m + n*(p+1) + 1, 0]

Rule 5654

$\text{Int}[\text{Tanh}[(a_*) + \text{Log}[(c_*)*(x_)^{(n_*)}]]*(b_*)*(d_*)^{(p_*)}, x_Symbol] \rightarrow \text{Dist}[x/(n*(c*x^n)^{(1/n)}), \text{Subst}[\text{Int}[x^{(1/n-1)}*\text{Tanh}[d*(a + b*\text{Log}[x])]^p, x], x, c*x^n], x] /;$ FreeQ[{a, b, c, d, n, p}, x] && (NeQ[c, 1] || NeQ[n, 1])

Rule 5656

$\text{Int}[(e_*)*(x_)^{(m_*)}*\text{Tanh}[(a_*) + \text{Log}[x]*(b_*)*(d_*)]^{(p_*)}, x_Symbol] \rightarrow \text{Int}[(e*x)^m*((-1 + E^{(2*a*d)}*x^{(2*b*d)})^p/(1 + E^{(2*a*d)}*x^{(2*b*d)})^p),$

x] /; FreeQ[{a, b, d, e, m, p}, x]

Rubi steps

$$\int \tanh(d(a + b \log(cx^n))) dx = \int \tanh(d(a + b \log(cx^n))) dx$$

Mathematica [B] Leaf count is larger than twice the leaf count of optimal. 126 vs. 2(53) = 106.

time = 6.28, size = 126, normalized size = 2.38

$$\frac{e^{2d(a+b \log(cx^n))} x {}_2F_1\left(1, 1 + \frac{1}{2bdn}; 2 + \frac{1}{2bdn}; -e^{2d(a+b \log(cx^n))}\right)}{1 + 2bdn} - x {}_2F_1\left(1, \frac{1}{2bdn}; 1 + \frac{1}{2bdn}; -e^{2d(a+b \log(cx^n))}\right)$$

Antiderivative was successfully verified.

[In] Integrate[Tanh[d*(a + b*Log[c*x^n])],x]

[Out] (E^(2*d*(a + b*Log[c*x^n]))*x*Hypergeometric2F1[1, 1 + 1/(2*b*d*n), 2 + 1/(2*b*d*n), -E^(2*d*(a + b*Log[c*x^n]))])/(1 + 2*b*d*n) - x*Hypergeometric2F1[1, 1/(2*b*d*n), 1 + 1/(2*b*d*n), -E^(2*d*(a + b*Log[c*x^n]))]

Maple [F]

time = 0.48, size = 0, normalized size = 0.00

$$\int \tanh(d(a + b \ln(cx^n))) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(d*(a+b*ln(c*x^n))),x)

[Out] int(tanh(d*(a+b*ln(c*x^n))),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(d*(a+b*log(c*x^n))),x, algorithm="maxima")

[Out] x - 2*integrate(1/(c^(2*b*d)*e^(2*b*d*log(x^n) + 2*a*d) + 1), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(d*(a+b*log(c*x^n))),x, algorithm="fricas")`

[Out] `integral(tanh(b*d*log(c*x^n) + a*d), x)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \tanh(d(a + b \log(cx^n))) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(d*(a+b*ln(c*x**n))),x)`

[Out] `Integral(tanh(d*(a + b*log(c*x**n))), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(d*(a+b*log(c*x^n))),x, algorithm="giac")`

[Out] `integrate(tanh((b*log(c*x^n) + a)*d), x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \tanh(d(a + b \ln(cx^n))) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tanh(d*(a + b*log(c*x^n))),x)`

[Out] `int(tanh(d*(a + b*log(c*x^n))), x)`

$$3.176 \quad \int \frac{\tanh(d(a+b \log(cx^n)))}{x} dx$$

Optimal. Leaf size=25

$$\frac{\log(\cosh(ad + bd \log(cx^n)))}{bdn}$$

[Out] ln(cosh(a*d+b*d*ln(c*x^n)))/b/d/n

Rubi [A]

time = 0.01, antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.059$, Rules used = {3556}

$$\frac{\log(\cosh(ad + bd \log(cx^n)))}{bdn}$$

Antiderivative was successfully verified.

[In] Int[Tanh[d*(a + b*Log[c*x^n])]/x,x]

[Out] Log[Cosh[a*d + b*d*Log[c*x^n]]]/(b*d*n)

Rule 3556

Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{\tanh(d(a + b \log(cx^n)))}{x} dx &= \frac{\text{Subst}(\int \tanh(d(a + bx)) dx, x, \log(cx^n))}{n} \\ &= \frac{\log(\cosh(ad + bd \log(cx^n)))}{bdn} \end{aligned}$$

Mathematica [A]

time = 0.04, size = 24, normalized size = 0.96

$$\frac{\log(\cosh(d(a + b \log(cx^n))))}{bdn}$$

Antiderivative was successfully verified.

[In] Integrate[Tanh[d*(a + b*Log[c*x^n])]/x,x]

[Out] Log[Cosh[d*(a + b*Log[c*x^n])]]/(b*d*n)

Maple [A]

time = 2.11, size = 48, normalized size = 1.92

method	result
derivativedivides	$\frac{-\frac{\ln(\tanh(d(a+b\ln(cx^n))))-1}{2} - \frac{\ln(\tanh(d(a+b\ln(cx^n))))+1}{2}}{nbd}$
default	$\frac{-\frac{\ln(\tanh(d(a+b\ln(cx^n))))-1}{2} - \frac{\ln(\tanh(d(a+b\ln(cx^n))))+1}{2}}{nbd}$
risch	$\ln(x) - \frac{2a}{nb} - \frac{2\ln(c)}{n} - \frac{2\ln(x^n)}{n} + \frac{i\pi\operatorname{csgn}(icx^n)^3}{n} - \frac{i\pi\operatorname{csgn}(icx^n)^2\operatorname{csgn}(ic)}{n} - \frac{i\pi\operatorname{csgn}(icx^n)^2\operatorname{csgn}(ix^n)}{n} + \dots$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(tanh(d*(a+b*ln(c*x^n)))/x,x,method=_RETURNVERBOSE)
```

```
[Out] 1/n/b/d*(-1/2*ln(tanh(d*(a+b*ln(c*x^n))))-1)-1/2*ln(tanh(d*(a+b*ln(c*x^n)))+1))
```

Maxima [A]

time = 0.27, size = 24, normalized size = 0.96

$$\frac{\log(\cosh((b\log(cx^n) + a)d))}{bdn}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(tanh(d*(a+b*log(c*x^n)))/x,x, algorithm="maxima")
```

```
[Out] log(cosh((b*log(c*x^n) + a)*d))/(b*d*n)
```

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 76 vs. 2(25) = 50.

time = 0.47, size = 76, normalized size = 3.04

$$\frac{bdn \log(x) - \log\left(\frac{2 \cosh(bdn \log(x) + bd \log(c) + ad)}{\cosh(bdn \log(x) + bd \log(c) + ad) - \sinh(bdn \log(x) + bd \log(c) + ad)}\right)}{bdn}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(tanh(d*(a+b*log(c*x^n)))/x,x, algorithm="fricas")
```

```
[Out] -(b*d*n*log(x) - log(2*cosh(b*d*n*log(x) + b*d*log(c) + a*d)/(cosh(b*d*n*log(x) + b*d*log(c) + a*d) - sinh(b*d*n*log(x) + b*d*log(c) + a*d))))/(b*d*n)
```

Sympy [A]

time = 1.88, size = 36, normalized size = 1.44

$$\frac{\log(bdn \tanh^2(ad + bd \log(cx^n)) - bdn)}{2bdn}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(d*(a+b*ln(c*x**n)))/x,x)

[Out] $-\log(b*d*n*\tanh(a*d + b*d*\log(c*x**n))**2 - b*d*n)/(2*b*d*n)$

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 74 vs. $2(25) = 50$.
time = 0.50, size = 74, normalized size = 2.96

$$\frac{\log\left(\sqrt{2x^{2bdn}|c|^{2bd}\cos(\pi b d \operatorname{sgn}(c) - \pi b d)e^{(2ad)} + x^{4bdn}|c|^{4bd}e^{(4ad)} + 1}\right)}{bdn} - \log(x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(d*(a+b*log(c*x^n)))/x,x, algorithm="giac")

[Out] $\log(\sqrt{(2*x^{(2*b*d*n)}*abs(c)^{(2*b*d)}*\cos(\pi*b*d*\operatorname{sgn}(c) - \pi*b*d)*e^{(2*a*d)} + x^{(4*b*d*n)}*abs(c)^{(4*b*d)}*e^{(4*a*d)} + 1)})/(b*d*n) - \log(x)$

Mupad [B]

time = 1.06, size = 34, normalized size = 1.36

$$\frac{\ln\left(e^{2ad}(cx^n)^{2bd} + 1\right)}{bdn} - \ln(x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(d*(a + b*log(c*x^n)))/x,x)

[Out] $\log(\exp(2*a*d)*(c*x^n)^{(2*b*d)} + 1)/(b*d*n) - \log(x)$

$$3.177 \quad \int \frac{\tanh(d(a+b \log(cx^n)))}{x^2} dx$$

Optimal. Leaf size=59

$$-\frac{1}{x} + \frac{{}_2F_1\left(1, -\frac{1}{2bdn}; 1 - \frac{1}{2bdn}; -e^{2ad}(cx^n)^{2bd}\right)}{x}$$

[Out] $-1/x + 2*\text{hypergeom}([1, -1/2/b/d/n], [1-1/2/b/d/n], -\exp(2*a*d)*(c*x^n)^{(2*b*d)})/x$

Rubi [A]

time = 0.04, antiderivative size = 59, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {5658, 5656, 470, 371}

$$\frac{{}_2F_1\left(1, -\frac{1}{2bdn}; 1 - \frac{1}{2bdn}; -e^{2ad}(cx^n)^{2bd}\right)}{x} - \frac{1}{x}$$

Antiderivative was successfully verified.

[In] Int[Tanh[d*(a + b*Log[c*x^n])]/x^2,x]

[Out] $-x^{(-1)} + (2*\text{Hypergeometric2F1}[1, -1/2*1/(b*d*n), 1 - 1/(2*b*d*n), -(E^{(2*a*d)*(c*x^n)^{(2*b*d)})})]/x$

Rule 371

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[a^p*((c*x)^(m+1)/(c*(m+1)))*Hypergeometric2F1[-p, (m+1)/n, (m+1)/n+1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])

Rule 470

Int[((e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> Simp[d*(e*x)^(m+1)*((a + b*x^n)^(p+1)/(b*e*(m+n*(p+1)+1))), x] - Dist[(a*d*(m+1) - b*c*(m+n*(p+1)+1))/(b*(m+n*(p+1)+1)), Int[(e*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && NeQ[b*c - a*d, 0] && NeQ[m + n*(p+1) + 1, 0]

Rule 5656

Int[((e_.)*(x_))^(m_.)*Tanh[((a_.) + Log[x_]*(b_.))*(d_.)]^(p_.), x_Symbol] :> Int[(e*x)^m*((-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p), x] /; FreeQ[{a, b, d, e, m, p}, x]

Rule 5658

```
Int[((e_.)*(x_.))^(m_.)*Tanh[((a_.) + Log[(c_.)*(x_.)^(n_.)]*(b_.))*(d_.)]^(p_.), x_Symbol]
:> Dist[(e*x)^(m + 1)/(e*n*(c*x^n)^((m + 1)/n)), Subst[Int[x^((m + 1)/n - 1)*Tanh[d*(a + b*Log[x])]^p, x], x, c*x^n], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && (NeQ[c, 1] || NeQ[n, 1])
```

Rubi steps

$$\int \frac{\tanh(d(a + b \log(cx^n)))}{x^2} dx = \int \frac{\tanh(d(a + b \log(cx^n)))}{x^2} dx$$

Mathematica [B] Leaf count is larger than twice the leaf count of optimal. 126 vs. 2(59) = 118.

time = 2.29, size = 126, normalized size = 2.14

$$\frac{e^{2d(a+b \log(cx^n))} {}_2F_1\left(1, 1 - \frac{1}{2bdn}; 2 - \frac{1}{2bdn}; -e^{2d(a+b \log(cx^n))}\right)}{-1+2bdn} + \frac{{}_2F_1\left(1, -\frac{1}{2bdn}; 1 - \frac{1}{2bdn}; -e^{2d(a+b \log(cx^n))}\right)}{x}$$

Antiderivative was successfully verified.

[In] Integrate[Tanh[d*(a + b*Log[c*x^n])]/x^2,x]

[Out] ((E^(2*d*(a + b*Log[c*x^n]))*Hypergeometric2F1[1, 1 - 1/(2*b*d*n), 2 - 1/(2*b*d*n), -E^(2*d*(a + b*Log[c*x^n])])]/(-1 + 2*b*d*n) + Hypergeometric2F1[1, -1/2*1/(b*d*n), 1 - 1/(2*b*d*n), -E^(2*d*(a + b*Log[c*x^n])])]/x

Maple [F]

time = 0.48, size = 0, normalized size = 0.00

$$\int \frac{\tanh(d(a + b \ln(cx^n)))}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(d*(a+b*ln(c*x^n)))/x^2,x)

[Out] int(tanh(d*(a+b*ln(c*x^n)))/x^2,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(d*(a+b*log(c*x^n)))/x^2,x, algorithm="maxima")

[Out] -1/x - 2*integrate(1/(c^(2*b*d)*x^2*e^(2*b*d*log(x^n) + 2*a*d) + x^2), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(d*(a+b*log(c*x^n)))/x^2,x, algorithm="fricas")

[Out] integral(tanh(b*d*log(c*x^n) + a*d)/x^2, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\tanh(ad + bd \log(cx^n))}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(d*(a+b*ln(c*x**n)))/x**2,x)

[Out] Integral(tanh(a*d + b*d*log(c*x**n))/x**2, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(d*(a+b*log(c*x^n)))/x^2,x, algorithm="giac")

[Out] integrate(tanh((b*log(c*x^n) + a)*d)/x^2, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\tanh(d(a + b \ln(cx^n)))}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(d*(a + b*log(c*x^n)))/x^2,x)

[Out] int(tanh(d*(a + b*log(c*x^n)))/x^2, x)

$$3.178 \quad \int \frac{\tanh(d(a+b \log(cx^n)))}{x^3} dx$$

Optimal. Leaf size=56

$$-\frac{1}{2x^2} + \frac{{}_2F_1\left(1, -\frac{1}{bdn}; 1 - \frac{1}{bdn}; -e^{2ad}(cx^n)^{2bd}\right)}{x^2}$$

[Out] $-1/2/x^2 + \text{hypergeom}([1, -1/b/d/n], [1-1/b/d/n], -\exp(2*a*d)*(c*x^n)^{(2*b*d)})/x^2$

Rubi [A]

time = 0.04, antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {5658, 5656, 470, 371}

$$\frac{{}_2F_1\left(1, -\frac{1}{bdn}; 1 - \frac{1}{bdn}; -e^{2ad}(cx^n)^{2bd}\right)}{x^2} - \frac{1}{2x^2}$$

Antiderivative was successfully verified.

[In] Int[Tanh[d*(a + b*Log[c*x^n])]/x^3,x]

[Out] $-1/2*1/x^2 + \text{Hypergeometric2F1}[1, -(1/(b*d*n)), 1 - 1/(b*d*n), -(E^{(2*a*d)}*(c*x^n)^{(2*b*d)})]/x^2$

Rule 371

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])

Rule 470

Int[((e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> Simp[d*(e*x)^(m + 1)*((a + b*x^n)^(p + 1)/(b*e*(m + n*(p + 1) + 1))), x] - Dist[(a*d*(m + 1) - b*c*(m + n*(p + 1) + 1))/(b*(m + n*(p + 1) + 1)), Int[(e*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && NeQ[b*c - a*d, 0] && NeQ[m + n*(p + 1) + 1, 0]

Rule 5656

Int[((e_.)*(x_))^(m_.)*Tanh[((a_.) + Log[x_]*(b_.))*(d_.)]^(p_.), x_Symbol] :> Int[(e*x)^m*((-1 + E^{(2*a*d)*x^{(2*b*d)}})^p/(1 + E^{(2*a*d)*x^{(2*b*d)}})^p), x] /; FreeQ[{a, b, d, e, m, p}, x]

Rule 5658

Int[((e_.)*(x_))^(m_.)*Tanh[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*(d_.)]^(p_.), x_Symbol] :> Dist[(e*x)^(m + 1)/(e*n*(c*x^n)^((m + 1)/n)), Subst[Int[x^((m + 1)/n - 1)*Tanh[d*(a + b*Log[x])]^p, x], x, c*x^n], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && (NeQ[c, 1] || NeQ[n, 1])

Rubi steps

$$\int \frac{\tanh(d(a + b \log(cx^n)))}{x^3} dx = \int \frac{\tanh(d(a + b \log(cx^n)))}{x^3} dx$$

Mathematica [B] Leaf count is larger than twice the leaf count of optimal. 120 vs. 2(56) = 112.

time = 2.18, size = 120, normalized size = 2.14

$$\frac{e^{2d(a+b \log(cx^n))} {}_2F_1\left(1, 1 - \frac{1}{bdn}; 2 - \frac{1}{bdn}; -e^{2d(a+b \log(cx^n))}\right)}{-1+bdn} + \frac{{}_2F_1\left(1, -\frac{1}{bdn}; 1 - \frac{1}{bdn}; -e^{2d(a+b \log(cx^n))}\right)}{2x^2}$$

Antiderivative was successfully verified.

[In] Integrate[Tanh[d*(a + b*Log[c*x^n])]/x^3, x]

[Out] ((E^(2*d*(a + b*Log[c*x^n]))*Hypergeometric2F1[1, 1 - 1/(b*d*n), 2 - 1/(b*d*n), -E^(2*d*(a + b*Log[c*x^n]))])/(-1 + b*d*n) + Hypergeometric2F1[1, -(1/(b*d*n)), 1 - 1/(b*d*n), -E^(2*d*(a + b*Log[c*x^n]))])/(2*x^2)

Maple [F]

time = 0.48, size = 0, normalized size = 0.00

$$\int \frac{\tanh(d(a + b \ln(cx^n)))}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(d*(a+b*ln(c*x^n)))/x^3, x)

[Out] int(tanh(d*(a+b*ln(c*x^n)))/x^3, x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(d*(a+b*log(c*x^n)))/x^3,x, algorithm="maxima")

[Out] -1/2/x^2 - 2*integrate(1/(c^(2*b*d)*x^3*e^(2*b*d*log(x^n) + 2*a*d) + x^3), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(d*(a+b*log(c*x^n)))/x^3,x, algorithm="fricas")

[Out] integral(tanh(b*d*log(c*x^n) + a*d)/x^3, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\tanh(ad + bd \log(cx^n))}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(d*(a+b*ln(c*x**n)))/x**3,x)

[Out] Integral(tanh(a*d + b*d*log(c*x**n))/x**3, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(d*(a+b*log(c*x^n)))/x^3,x, algorithm="giac")

[Out] integrate(tanh((b*log(c*x^n) + a)*d)/x^3, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\tanh(d(a + b \ln(cx^n)))}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(d*(a + b*log(c*x^n)))/x^3,x)

[Out] int(tanh(d*(a + b*log(c*x^n)))/x^3, x)

3.179 $\int x^3 \tanh^2(d(a + b \log(cx^n))) dx$

Optimal. Leaf size=133

$$\frac{1}{4} \left(1 + \frac{4}{bdn}\right) x^4 + \frac{x^4 \left(1 - e^{2ad}(cx^n)^{2bd}\right)}{bdn \left(1 + e^{2ad}(cx^n)^{2bd}\right)} - \frac{2x^4 {}_2F_1\left(1, \frac{2}{bdn}; 1 + \frac{2}{bdn}; -e^{2ad}(cx^n)^{2bd}\right)}{bdn}$$

[Out] $1/4*(1+4/b/d/n)*x^4+x^4*(1-\exp(2*a*d)*(c*x^n)^{(2*b*d)})/b/d/n/(1+\exp(2*a*d)*(c*x^n)^{(2*b*d)})-2*x^4*\text{hypergeom}([1, 2/b/d/n], [1+2/b/d/n], -\exp(2*a*d)*(c*x^n)^{(2*b*d)})/b/d/n$

Rubi [A]

time = 0.12, antiderivative size = 133, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$, Rules used = {5658, 5656, 516, 470, 371}

$$-\frac{2x^4 {}_2F_1\left(1, \frac{2}{bdn}; 1 + \frac{2}{bdn}; -e^{2ad}(cx^n)^{2bd}\right)}{bdn} + \frac{x^4 \left(1 - e^{2ad}(cx^n)^{2bd}\right)}{bdn \left(e^{2ad}(cx^n)^{2bd} + 1\right)} + \frac{1}{4} x^4 \left(\frac{4}{bdn} + 1\right)$$

Antiderivative was successfully verified.

[In] $\text{Int}[x^3*\text{Tanh}[d*(a + b*\text{Log}[c*x^n])]^2,x]$

[Out] $((1 + 4/(b*d*n))*x^4)/4 + (x^4*(1 - E^{(2*a*d)*(c*x^n)^{(2*b*d)}}))/(b*d*n*(1 + E^{(2*a*d)*(c*x^n)^{(2*b*d)}})) - (2*x^4*\text{Hypergeometric2F1}[1, 2/(b*d*n), 1 + 2/(b*d*n), -(E^{(2*a*d)*(c*x^n)^{(2*b*d)}})])/(b*d*n)$

Rule 371

$\text{Int}[\left((c_*)*(x_*)\right)^{(m_*)}*\left((a_*) + (b_*)*(x_*)^{(n_*)}\right)^{(p_*)}, x_Symbol] :> \text{Simp}[a^p * \left((c*x)^{(m+1)}/(c*(m+1))\right)*\text{Hypergeometric2F1}[-p, (m+1)/n, (m+1)/n + 1, (-b)*(x^n/a)], x] /; \text{FreeQ}\{a, b, c, m, n, p\}, x] \&\& !\text{IGtQ}[p, 0] \&\& (\text{ILtQ}[p, 0] \mid\mid \text{GtQ}[a, 0])$

Rule 470

$\text{Int}[\left((e_*)*(x_*)\right)^{(m_*)}*\left((a_*) + (b_*)*(x_*)^{(n_*)}\right)^{(p_*)}*\left((c_*) + (d_*)*(x_*)^{(n_*)}\right), x_Symbol] :> \text{Simp}[d*(e*x)^{(m+1)}*\left((a + b*x^n)^{(p+1)}/(b*e*(m+n*(p+1)+1))\right), x] - \text{Dist}[(a*d*(m+1) - b*c*(m+n*(p+1)+1))/(b*(m+n*(p+1)+1)), \text{Int}[(e*x)^m*(a + b*x^n)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, m, n, p\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[m + n*(p+1) + 1, 0]$

Rule 516

$\text{Int}[\left((e_*)*(x_*)\right)^{(m_*)}*\left((a_*) + (b_*)*(x_*)^{(n_*)}\right)^{(p_*)}*\left((c_*) + (d_*)*(x_*)^{(n_*)}\right)^{(q_*)}, x_Symbol] :> \text{Simp}[(-c*b - a*d)*(e*x)^{(m+1)}*(a + b*x^n)^{(p+1)}$

```

*((c + d*x^n)^(q - 1)/(a*b*e*n*(p + 1))), x] + Dist[1/(a*b*n*(p + 1)), Int[
(e*x)^m*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 2)*Simp[c*(c*b*n*(p + 1) + (c*
b - a*d)*(m + 1)) + d*(c*b*n*(p + 1) + (c*b - a*d)*(m + n*(q - 1) + 1))*x^n
, x], x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && NeQ[b*c - a*d, 0] && LtQ
[p, -1] && GtQ[q, 1] && IntBinomialQ[a, b, c, d, e, m, n, p, q, x]

```

Rule 5656

```

Int[((e_)*(x_))^(m_)*Tanh[((a_) + Log[x_]*(b_))*(d_)]^(p_), x_Symbol]
:> Int[(e*x)^m*((-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p),
x] /; FreeQ[{a, b, d, e, m, p}, x]

```

Rule 5658

```

Int[((e_)*(x_))^(m_)*Tanh[((a_) + Log[(c_)*(x_)^(n_)]*(b_))*(d_)]^(p
_), x_Symbol] :> Dist[(e*x)^(m + 1)/(e*n*(c*x^n)^((m + 1)/n)), Subst[Int[x
^((m + 1)/n - 1)*Tanh[d*(a + b*Log[x])]^p, x], x, c*x^n], x] /; FreeQ[{a, b
, c, d, e, m, n, p}, x] && (NeQ[c, 1] || NeQ[n, 1])

```

Rubi steps

$$\int x^3 \tanh^2(d(a + b \log(cx^n))) dx = \int x^3 \tanh^2(d(a + b \log(cx^n))) dx$$

Mathematica [A]

time = 5.71, size = 159, normalized size = 1.20

$$\frac{x^4 (8e^{2d(a+b \log(cx^n))} {}_2F_1\left(1, 1 + \frac{2}{bdn}; 2 + \frac{2}{bdn}; -e^{2d(a+b \log(cx^n))}\right) + (2 + bdn) (bdn - 4 {}_2F_1\left(1, \frac{2}{bdn}; 1 + \frac{2}{bdn}; -e^{2d(a+b \log(cx^n))}\right) - 4 \tanh(d(a + b \log(cx^n))))}{4bdn(2 + bdn)}$$

Antiderivative was successfully verified.

```
[In] Integrate[x^3*Tanh[d*(a + b*Log[c*x^n])]^2,x]
```

```
[Out] (x^4*(8*E^(2*d*(a + b*Log[c*x^n]))*Hypergeometric2F1[1, 1 + 2/(b*d*n), 2 +
2/(b*d*n), -E^(2*d*(a + b*Log[c*x^n]))] + (2 + b*d*n)*(b*d*n - 4*Hypergeome
tric2F1[1, 2/(b*d*n), 1 + 2/(b*d*n), -E^(2*d*(a + b*Log[c*x^n]))] - 4*Tanh[
d*(a + b*Log[c*x^n]))))/ (4*b*d*n*(2 + b*d*n))

```

Maple [F]

time = 0.71, size = 0, normalized size = 0.00

$$\int x^3 (\tanh^2(d(a + b \ln(cx^n)))) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x^3*tanh(d*(a+b*ln(c*x^n)))^2,x)`

[Out] `int(x^3*tanh(d*(a+b*ln(c*x^n)))^2,x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^3*tanh(d*(a+b*log(c*x^n)))^2,x, algorithm="maxima")`

[Out] `1/4*(b*c^(2*b*d)*d*n*x^4*e^(2*b*d*log(x^n) + 2*a*d) + (b*d*n + 8)*x^4)/(b*c^(2*b*d)*d*n*e^(2*b*d*log(x^n) + 2*a*d) + b*d*n) - 8*integrate(x^3/(b*c^(2*b*d)*d*n*e^(2*b*d*log(x^n) + 2*a*d) + b*d*n), x)`

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^3*tanh(d*(a+b*log(c*x^n)))^2,x, algorithm="fricas")`

[Out] `integral(x^3*tanh(b*d*log(c*x^n) + a*d)^2, x)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x^3 \tanh^2(ad + bd \log(cx^n)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x**3*tanh(d*(a+b*ln(c*x**n)))**2,x)`

[Out] `Integral(x**3*tanh(a*d + b*d*log(c*x**n))**2, x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^3*tanh(d*(a+b*log(c*x^n)))^2,x, algorithm="giac")`

[Out] `integrate(x^3*tanh((b*log(c*x^n) + a)*d)^2, x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int x^3 \tanh(d(a + b \ln(cx^n)))^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^3*tanh(d*(a + b*log(c*x^n)))^2,x)

[Out] int(x^3*tanh(d*(a + b*log(c*x^n)))^2, x)

3.180 $\int x^2 \tanh^2(d(a + b \log(cx^n))) dx$

Optimal. Leaf size=137

$$\frac{1}{3} \left(1 + \frac{3}{bdn}\right) x^3 + \frac{x^3 (1 - e^{2ad}(cx^n)^{2bd})}{bdn (1 + e^{2ad}(cx^n)^{2bd})} - \frac{2x^3 {}_2F_1\left(1, \frac{3}{2bdn}; 1 + \frac{3}{2bdn}; -e^{2ad}(cx^n)^{2bd}\right)}{bdn}$$

[Out] $\frac{1}{3}*(1+3/b/d/n)*x^3+x^3*(1-\exp(2*a*d)*(c*x^n)^{(2*b*d)})/b/d/n/(1+\exp(2*a*d)*(c*x^n)^{(2*b*d)})-2*x^3*\text{hypergeom}([1, 3/2/b/d/n], [1+3/2/b/d/n], -\exp(2*a*d)*(c*x^n)^{(2*b*d)})/b/d/n$

Rubi [A]

time = 0.12, antiderivative size = 137, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$, Rules used = {5658, 5656, 516, 470, 371}

$$-\frac{2x^3 {}_2F_1\left(1, \frac{3}{2bdn}; 1 + \frac{3}{2bdn}; -e^{2ad}(cx^n)^{2bd}\right)}{bdn} + \frac{x^3 (1 - e^{2ad}(cx^n)^{2bd})}{bdn (e^{2ad}(cx^n)^{2bd} + 1)} + \frac{1}{3} x^3 \left(\frac{3}{bdn} + 1\right)$$

Antiderivative was successfully verified.

[In] $\text{Int}[x^2*\text{Tanh}[d*(a + b*\text{Log}[c*x^n])]^2,x]$

[Out] $((1 + 3/(b*d*n))*x^3)/3 + (x^3*(1 - E^{(2*a*d)*(c*x^n)^{(2*b*d)}}))/(b*d*n*(1 + E^{(2*a*d)*(c*x^n)^{(2*b*d)}})) - (2*x^3*\text{Hypergeometric2F1}[1, 3/(2*b*d*n), 1 + 3/(2*b*d*n), -(E^{(2*a*d)*(c*x^n)^{(2*b*d)}})]/(b*d*n)$

Rule 371

$\text{Int}[\left((c_*)*(x_*)^{(m_*)}*((a_*) + (b_*)*(x_*)^{(n_*)})^{(p_*)}, x_Symbol\right) :> \text{Simp}[a^p * ((c*x)^{(m+1})/(c*(m+1)) * \text{Hypergeometric2F1}[-p, (m+1)/n, (m+1)/n + 1, (-b)*(x^n/a)], x] /; \text{FreeQ}\{a, b, c, m, n, p\}, x] \&\& !\text{IGtQ}[p, 0] \&\& (\text{ILtQ}[p, 0] || \text{GtQ}[a, 0])$

Rule 470

$\text{Int}[\left((e_*)*(x_*)^{(m_*)}*((a_*) + (b_*)*(x_*)^{(n_*)})^{(p_*)}*((c_*) + (d_*)*(x_*)^{(n_*)}), x_Symbol\right) :> \text{Simp}[d*(e*x)^{(m+1)}*((a + b*x^n)^{(p+1})/(b*e*(m+n*(p+1)+1))), x] - \text{Dist}[(a*d*(m+1) - b*c*(m+n*(p+1)+1))/(b*(m+n*(p+1)+1)), \text{Int}[(e*x)^m*(a + b*x^n)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, m, n, p\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[m + n*(p+1) + 1, 0]$

Rule 516

$\text{Int}[\left((e_*)*(x_*)^{(m_*)}*((a_*) + (b_*)*(x_*)^{(n_*)})^{(p_*)}*((c_*) + (d_*)*(x_*)^{(n_*)})^{(q_*)}, x_Symbol\right) :> \text{Simp}[(-c*b - a*d)*(e*x)^{(m+1)}*(a + b*x^n)^{(p+1)}$

```

*((c + d*x^n)^(q - 1)/(a*b*e*n*(p + 1)), x] + Dist[1/(a*b*n*(p + 1)), Int[
(e*x)^m*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 2)*Simp[c*(c*b*n*(p + 1) + (c*
b - a*d)*(m + 1)) + d*(c*b*n*(p + 1) + (c*b - a*d)*(m + n*(q - 1) + 1))*x^n
, x], x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && NeQ[b*c - a*d, 0] && LtQ
[p, -1] && GtQ[q, 1] && IntBinomialQ[a, b, c, d, e, m, n, p, q, x]

```

Rule 5656

```

Int[((e_)*(x_))^(m_)*Tanh[((a_) + Log[x_]*(b_))*(d_)]^(p_), x_Symbol]
:> Int[(e*x)^m*((-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p),
x] /; FreeQ[{a, b, d, e, m, p}, x]

```

Rule 5658

```

Int[((e_)*(x_))^(m_)*Tanh[((a_) + Log[(c_)*(x_)^(n_)]*(b_))*(d_)]^(p
_), x_Symbol] :> Dist[(e*x)^(m + 1)/(e*n*(c*x^n)^((m + 1)/n)), Subst[Int[x
^((m + 1)/n - 1)*Tanh[d*(a + b*Log[x])]^p, x], x, c*x^n], x] /; FreeQ[{a, b
, c, d, e, m, n, p}, x] && (NeQ[c, 1] || NeQ[n, 1])

```

Rubi steps

$$\int x^2 \tanh^2(d(a + b \log(cx^n))) dx = \int x^2 \tanh^2(d(a + b \log(cx^n))) dx$$

Mathematica [A]

time = 5.54, size = 169, normalized size = 1.23

$$\frac{x^3 (9e^{2d(a+b \log(cx^n))} {}_2F_1(1, 1 + \frac{3}{2bdn}; 2 + \frac{3}{2bdn}; -e^{2d(a+b \log(cx^n))}) + (3 + 2bdn) (bdn - 3 {}_2F_1(1, \frac{3}{2bdn}; 1 + \frac{3}{2bdn}; -e^{2d(a+b \log(cx^n))}) - 3 \tanh(d(a + b \log(cx^n))))}{3bdn(3 + 2bdn)}$$

Antiderivative was successfully verified.

```
[In] Integrate[x^2*Tanh[d*(a + b*Log[c*x^n])]^2,x]
```

```
[Out] (x^3*(9*E^(2*d*(a + b*Log[c*x^n]))*Hypergeometric2F1[1, 1 + 3/(2*b*d*n), 2
+ 3/(2*b*d*n), -E^(2*d*(a + b*Log[c*x^n]))] + (3 + 2*b*d*n)*(b*d*n - 3*Hype
rgeometric2F1[1, 3/(2*b*d*n), 1 + 3/(2*b*d*n), -E^(2*d*(a + b*Log[c*x^n]))]
- 3*Tanh[d*(a + b*Log[c*x^n])])))/(3*b*d*n*(3 + 2*b*d*n))

```

Maple [F]

time = 0.47, size = 0, normalized size = 0.00

$$\int x^2 (\tanh^2(d(a + b \ln(cx^n)))) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x^2*tanh(d*(a+b*ln(c*x^n)))^2,x)`

[Out] `int(x^2*tanh(d*(a+b*ln(c*x^n)))^2,x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^2*tanh(d*(a+b*log(c*x^n)))^2,x, algorithm="maxima")`

[Out] `1/3*(b*c^(2*b*d)*d*n*x^3*e^(2*b*d*log(x^n) + 2*a*d) + (b*d*n + 6)*x^3)/(b*c^(2*b*d)*d*n*e^(2*b*d*log(x^n) + 2*a*d) + b*d*n) - 6*integrate(x^2/(b*c^(2*b*d)*d*n*e^(2*b*d*log(x^n) + 2*a*d) + b*d*n), x)`

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^2*tanh(d*(a+b*log(c*x^n)))^2,x, algorithm="fricas")`

[Out] `integral(x^2*tanh(b*d*log(c*x^n) + a*d)^2, x)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x^2 \tanh^2(ad + bd \log(cx^n)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x**2*tanh(d*(a+b*ln(c*x**n))))**2,x)`

[Out] `Integral(x**2*tanh(a*d + b*d*log(c*x**n))**2, x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^2*tanh(d*(a+b*log(c*x^n)))^2,x, algorithm="giac")`

[Out] `integrate(x^2*tanh((b*log(c*x^n) + a)*d)^2, x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int x^2 \tanh(d(a + b \ln(cx^n)))^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^2*tanh(d*(a + b*log(c*x^n)))^2,x)

[Out] int(x^2*tanh(d*(a + b*log(c*x^n)))^2, x)

3.181 $\int x \tanh^2(d(a + b \log(cx^n))) dx$

Optimal. Leaf size=131

$$\frac{1}{2} \left(1 + \frac{2}{bdn}\right) x^2 + \frac{x^2 \left(1 - e^{2ad}(cx^n)^{2bd}\right)}{bdn \left(1 + e^{2ad}(cx^n)^{2bd}\right)} - \frac{2x^2 {}_2F_1\left(1, \frac{1}{bdn}; 1 + \frac{1}{bdn}; -e^{2ad}(cx^n)^{2bd}\right)}{bdn}$$

[Out] $\frac{1}{2}*(1+2/b/d/n)*x^2+x^2*(1-\exp(2*a*d)*(c*x^n)^{(2*b*d)})/b/d/n/(1+\exp(2*a*d)*(c*x^n)^{(2*b*d)})-2*x^2*\text{hypergeom}([1, 1/b/d/n], [1+1/b/d/n], -\exp(2*a*d)*(c*x^n)^{(2*b*d)})/b/d/n$

Rubi [A]

time = 0.11, antiderivative size = 131, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$, Rules used = {5658, 5656, 516, 470, 371}

$$-\frac{2x^2 {}_2F_1\left(1, \frac{1}{bdn}; 1 + \frac{1}{bdn}; -e^{2ad}(cx^n)^{2bd}\right)}{bdn} + \frac{x^2 \left(1 - e^{2ad}(cx^n)^{2bd}\right)}{bdn \left(e^{2ad}(cx^n)^{2bd} + 1\right)} + \frac{1}{2} x^2 \left(\frac{2}{bdn} + 1\right)$$

Antiderivative was successfully verified.

[In] $\text{Int}[x*\text{Tanh}[d*(a + b*\text{Log}[c*x^n])]^2, x]$

[Out] $((1 + 2/(b*d*n))*x^2)/2 + (x^2*(1 - E^{(2*a*d)*(c*x^n)^{(2*b*d)}}))/(b*d*n*(1 + E^{(2*a*d)*(c*x^n)^{(2*b*d)}})) - (2*x^2*\text{Hypergeometric2F1}[1, 1/(b*d*n), 1 + 1/(b*d*n), -(E^{(2*a*d)*(c*x^n)^{(2*b*d)}})]/(b*d*n)$

Rule 371

$\text{Int}[\left((c_*)*(x_*)\right)^{(m_*)}*\left((a_*) + (b_*)*(x_*)^{(n_*)}\right)^{(p_*)}, x_Symbol] :> \text{Simp}[a^p * \left(\frac{c*x}{c*(m+1)}\right)*\text{Hypergeometric2F1}[-p, (m+1)/n, (m+1)/n + 1, (-b)*(x^n/a)], x] /; \text{FreeQ}\{a, b, c, m, n, p\}, x] \&\& !\text{IGtQ}[p, 0] \&\& (\text{ILtQ}[p, 0] \mid\mid \text{GtQ}[a, 0])$

Rule 470

$\text{Int}[\left((e_*)*(x_*)\right)^{(m_*)}*\left((a_*) + (b_*)*(x_*)^{(n_*)}\right)^{(p_*)}*\left((c_*) + (d_*)*(x_*)^{(n_*)}\right), x_Symbol] :> \text{Simp}[d*(e*x)^{(m+1)}*\left((a + b*x^n)^{(p+1)}\right)/(b*e*(m+n*(p+1)+1)), x] - \text{Dist}[(a*d*(m+1) - b*c*(m+n*(p+1)+1))/(b*(m+n*(p+1)+1)), \text{Int}[(e*x)^m*(a + b*x^n)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, m, n, p\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[m + n*(p+1) + 1, 0]$

Rule 516

$\text{Int}[\left((e_*)*(x_*)\right)^{(m_*)}*\left((a_*) + (b_*)*(x_*)^{(n_*)}\right)^{(p_*)}*\left((c_*) + (d_*)*(x_*)^{(n_*)}\right)^{(q_*)}, x_Symbol] :> \text{Simp}[(-c*b - a*d)*(e*x)^{(m+1)}*(a + b*x^n)^{(p+1)}$

```

*((c + d*x^n)^(q - 1)/(a*b*e*n*(p + 1))), x] + Dist[1/(a*b*n*(p + 1)), Int[
(e*x)^m*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 2)*Simp[c*(c*b*n*(p + 1) + (c*
b - a*d)*(m + 1)) + d*(c*b*n*(p + 1) + (c*b - a*d)*(m + n*(q - 1) + 1))*x^n
, x], x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && NeQ[b*c - a*d, 0] && LtQ
[p, -1] && GtQ[q, 1] && IntBinomialQ[a, b, c, d, e, m, n, p, q, x]

```

Rule 5656

```

Int[((e_)*(x_))^(m_)*Tanh[((a_) + Log[x_]*(b_))*(d_)]^(p_), x_Symbol]
:> Int[(e*x)^m*((-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p),
x] /; FreeQ[{a, b, d, e, m, p}, x]

```

Rule 5658

```

Int[((e_)*(x_))^(m_)*Tanh[((a_) + Log[(c_)*(x_)^(n_)]*(b_))*(d_)]^(p
_), x_Symbol] :> Dist[(e*x)^(m + 1)/(e*n*(c*x^n)^((m + 1)/n)), Subst[Int[x
^((m + 1)/n - 1)*Tanh[d*(a + b*Log[x])]^p, x], x, c*x^n], x] /; FreeQ[{a, b
, c, d, e, m, n, p}, x] && (NeQ[c, 1] || NeQ[n, 1])

```

Rubi steps

$$\int x \tanh^2(d(a + b \log(cx^n))) dx = \int x \tanh^2(d(a + b \log(cx^n))) dx$$

Mathematica [A]

time = 5.54, size = 155, normalized size = 1.18

$$\frac{x^2(2e^{2d(a+b \log(cx^n))} {}_2F_1\left(1, 1 + \frac{1}{bdn}; 2 + \frac{1}{bdn}; -e^{2d(a+b \log(cx^n))}\right) + (1 + bdn)(bdn - 2 {}_2F_1\left(1, \frac{1}{bdn}; 1 + \frac{1}{bdn}; -e^{2d(a+b \log(cx^n))}\right) - 2 \tanh(d(a + b \log(cx^n))))}{2bdn(1 + bdn)}$$

Antiderivative was successfully verified.

```
[In] Integrate[x*Tanh[d*(a + b*Log[c*x^n])]^2,x]
```

```
[Out] (x^2*(2*E^(2*d*(a + b*Log[c*x^n]))*Hypergeometric2F1[1, 1 + 1/(b*d*n), 2 +
1/(b*d*n), -E^(2*d*(a + b*Log[c*x^n]))] + (1 + b*d*n)*(b*d*n - 2*Hypergeome
tric2F1[1, 1/(b*d*n), 1 + 1/(b*d*n), -E^(2*d*(a + b*Log[c*x^n]))] - 2*Tanh[
d*(a + b*Log[c*x^n])])))/(2*b*d*n*(1 + b*d*n))

```

Maple [F]

time = 0.47, size = 0, normalized size = 0.00

$$\int x (\tanh^2(d(a + b \ln(cx^n)))) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x*tanh(d*(a+b*ln(c*x^n)))^2,x)`

[Out] `int(x*tanh(d*(a+b*ln(c*x^n)))^2,x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x*tanh(d*(a+b*log(c*x^n)))^2,x, algorithm="maxima")`

[Out] $\frac{1}{2}*(b*c^{(2*b*d)*d*n*x^2}*e^{(2*b*d*\log(x^n) + 2*a*d) + (b*d*n + 4)*x^2})/(b*c^{(2*b*d)*d*n}*e^{(2*b*d*\log(x^n) + 2*a*d) + b*d*n}) - 4*\int x/(b*c^{(2*b*d)*d*n}*e^{(2*b*d*\log(x^n) + 2*a*d) + b*d*n}), x)$

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x*tanh(d*(a+b*log(c*x^n)))^2,x, algorithm="fricas")`

[Out] `integral(x*tanh(b*d*log(c*x^n) + a*d)^2, x)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x \tanh^2(ad + bd \log(cx^n)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x*tanh(d*(a+b*ln(c*x**n)))**2,x)`

[Out] `Integral(x*tanh(a*d + b*d*log(c*x**n))**2, x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x*tanh(d*(a+b*log(c*x^n)))^2,x, algorithm="giac")`

[Out] `integrate(x*tanh((b*log(c*x^n) + a)*d)^2, x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int x \tanh(d(a + b \ln(cx^n)))^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x*tanh(d*(a + b*log(c*x^n)))^2,x)

[Out] int(x*tanh(d*(a + b*log(c*x^n)))^2, x)

3.182 $\int \tanh^2(d(a + b \log(cx^n))) dx$

Optimal. Leaf size=127

$$\left(1 + \frac{1}{bdn}\right)x + \frac{x(1 - e^{2ad}(cx^n)^{2bd})}{bdn(1 + e^{2ad}(cx^n)^{2bd})} - \frac{2x {}_2F_1\left(1, \frac{1}{2bdn}; 1 + \frac{1}{2bdn}; -e^{2ad}(cx^n)^{2bd}\right)}{bdn}$$

[Out] $(1+1/b/d/n)*x+x*(1-\exp(2*a*d)*(c*x^n)^{(2*b*d)})/b/d/n/(1+\exp(2*a*d)*(c*x^n)^{(2*b*d)})-2*x*\text{hypergeom}([1, 1/2/b/d/n], [1+1/2/b/d/n], -\exp(2*a*d)*(c*x^n)^{(2*b*d)})/b/d/n$

Rubi [A]

time = 0.11, antiderivative size = 127, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {5654, 5656, 516, 470, 371}

$$-\frac{2x {}_2F_1\left(1, \frac{1}{2bdn}; 1 + \frac{1}{2bdn}; -e^{2ad}(cx^n)^{2bd}\right)}{bdn} + \frac{x(1 - e^{2ad}(cx^n)^{2bd})}{bdn(e^{2ad}(cx^n)^{2bd} + 1)} + x\left(\frac{1}{bdn} + 1\right)$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Tanh}[d*(a + b*\text{Log}[c*x^n])]^2, x]$

[Out] $(1 + 1/(b*d*n))*x + (x*(1 - E^{(2*a*d)*(c*x^n)^{(2*b*d)}})/(b*d*n*(1 + E^{(2*a*d)*(c*x^n)^{(2*b*d)}})) - (2*x*\text{Hypergeometric2F1}[1, 1/(2*b*d*n), 1 + 1/(2*b*d*n), -(E^{(2*a*d)*(c*x^n)^{(2*b*d)}})]/(b*d*n)$

Rule 371

$\text{Int}[(c_*)*(x_*)^{(m_*)}*((a_*) + (b_*)*(x_*)^{(n_*)})^{(p_*)}, x_Symbol] \rightarrow \text{Simp}[a^p * ((c*x)^{(m+1})/(c*(m+1)) * \text{Hypergeometric2F1}[-p, (m+1)/n, (m+1)/n + 1, (-b)*(x^n/a)], x] /; \text{FreeQ}\{a, b, c, m, n, p\}, x \ \&\& \ !\text{IGtQ}[p, 0] \ \&\& \ (\text{ILtQ}[p, 0] \ || \ \text{GtQ}[a, 0])$

Rule 470

$\text{Int}[(e_*)*(x_*)^{(m_*)}*((a_*) + (b_*)*(x_*)^{(n_*)})^{(p_*)}*((c_*) + (d_*)*(x_*)^{(n_*)}), x_Symbol] \rightarrow \text{Simp}[d*(e*x)^{(m+1)}*((a + b*x^n)^{(p+1})/(b*e*(m+n*(p+1)+1))), x] - \text{Dist}[(a*d*(m+1) - b*c*(m+n*(p+1)+1))/(b*(m+n*(p+1)+1)), \text{Int}[(e*x)^m*(a + b*x^n)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, m, n, p\}, x \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[m + n*(p+1) + 1, 0]$

Rule 516

$\text{Int}[(e_*)*(x_*)^{(m_*)}*((a_*) + (b_*)*(x_*)^{(n_*)})^{(p_*)}*((c_*) + (d_*)*(x_*)^{(n_*)})^{(q_*)}, x_Symbol] \rightarrow \text{Simp}[(-c*b - a*d)*(e*x)^{(m+1)}*(a + b*x^n)^{(p+1)}$

```

*((c + d*x^n)^(q - 1)/(a*b*e*n*(p + 1)), x] + Dist[1/(a*b*n*(p + 1)), Int[
(e*x)^m*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 2)*Simp[c*(c*b*n*(p + 1) + (c*
b - a*d)*(m + 1)) + d*(c*b*n*(p + 1) + (c*b - a*d)*(m + n*(q - 1) + 1))*x^n
, x], x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && NeQ[b*c - a*d, 0] && LtQ
[p, -1] && GtQ[q, 1] && IntBinomialQ[a, b, c, d, e, m, n, p, q, x]

```

Rule 5654

```

Int[Tanh[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*(d_.)]^(p_.), x_Symbol] :> D
ist[x/(n*(c*x^n)^(1/n)), Subst[Int[x^(1/n - 1)*Tanh[d*(a + b*Log[x])]^p, x]
, x, c*x^n], x] /; FreeQ[{a, b, c, d, n, p}, x] && (NeQ[c, 1] || NeQ[n, 1])

```

Rule 5656

```

Int[((e_.)*(x_))^(m_.)*Tanh[((a_.) + Log[x_]*(b_.))*(d_.)]^(p_.), x_Symbol]
:> Int[(e*x)^m*((-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p),
x] /; FreeQ[{a, b, d, e, m, p}, x]

```

Rubi steps

$$\int \tanh^2(d(a + b \log(cx^n))) dx = \int \tanh^2(d(a + b \log(cx^n))) dx$$

Mathematica [A]

time = 6.45, size = 163, normalized size = 1.28

$$\frac{x(e^{2d(a+b \log(cx^n))} {}_2F_1(1, 1 + \frac{1}{2bdn}; 2 + \frac{1}{2bdn}; -e^{2d(a+b \log(cx^n))}) + (1 + 2bdn)(bdn - {}_2F_1(1, \frac{1}{2bdn}; 1 + \frac{1}{2bdn}; -e^{2d(a+b \log(cx^n))}) - \tanh(d(a + b \log(cx^n))))}{bdn(1 + 2bdn)}$$

Antiderivative was successfully verified.

```
[In] Integrate[Tanh[d*(a + b*Log[c*x^n])]^2, x]
```

```
[Out] (x*(E^(2*d*(a + b*Log[c*x^n]))*Hypergeometric2F1[1, 1 + 1/(2*b*d*n), 2 + 1/
(2*b*d*n), -E^(2*d*(a + b*Log[c*x^n])]) + (1 + 2*b*d*n)*(b*d*n - Hypergeome
tric2F1[1, 1/(2*b*d*n), 1 + 1/(2*b*d*n), -E^(2*d*(a + b*Log[c*x^n])])]) - Tan
h[d*(a + b*Log[c*x^n])])/(b*d*n*(1 + 2*b*d*n))

```

Maple [F]

time = 0.47, size = 0, normalized size = 0.00

$$\int \tanh^2(d(a + b \ln(cx^n))) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tanh(d*(a+b*ln(c*x^n)))^2,x)`

[Out] `int(tanh(d*(a+b*ln(c*x^n)))^2,x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(d*(a+b*log(c*x^n)))^2,x, algorithm="maxima")`

[Out] `(b*c^(2*b*d)*d*n*x*e^(2*b*d*log(x^n) + 2*a*d) + (b*d*n + 2)*x)/(b*c^(2*b*d)*d*n*e^(2*b*d*log(x^n) + 2*a*d) + b*d*n) - 2*integrate(1/(b*c^(2*b*d)*d*n*e^(2*b*d*log(x^n) + 2*a*d) + b*d*n), x)`

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(d*(a+b*log(c*x^n)))^2,x, algorithm="fricas")`

[Out] `integral(tanh(b*d*log(c*x^n) + a*d)^2, x)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \tanh^2(d(a + b \log(cx^n))) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(d*(a+b*ln(c*x**n)))**2,x)`

[Out] `Integral(tanh(d*(a + b*log(c*x**n)))**2, x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(d*(a+b*log(c*x^n)))^2,x, algorithm="giac")`

[Out] `integrate(tanh((b*log(c*x^n) + a)*d)^2, x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \tanh(d(a + b \ln(cx^n)))^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(d*(a + b*log(c*x^n)))^2,x)

[Out] int(tanh(d*(a + b*log(c*x^n)))^2, x)

$$3.183 \quad \int \frac{\tanh^2(d(a+b \log(cx^n)))}{x} dx$$

Optimal. Leaf size=28

$$\log(x) - \frac{\tanh(ad + bd \log(cx^n))}{bdn}$$

[Out] $\ln(x) - \tanh(a*d + b*d*\ln(c*x^n))/b/d/n$

Rubi [A]

time = 0.02, antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {3554, 8}

$$\log(x) - \frac{\tanh(ad + bd \log(cx^n))}{bdn}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Tanh}[d*(a + b*\text{Log}[c*x^n])]^2/x, x]$

[Out] $\text{Log}[x] - \text{Tanh}[a*d + b*d*\text{Log}[c*x^n]]/(b*d*n)$

Rule 8

$\text{Int}[a_, x_Symbol] \text{ :> } \text{Simp}[a*x, x] \text{ /; } \text{FreeQ}[a, x]$

Rule 3554

$\text{Int}[(b_.)*\text{tan}[(c_.) + (d_.)*(x_.)]^{(n_.)}, x_Symbol] \text{ :> } \text{Simp}[b*((b*\text{Tan}[c + d*x])^{(n-1)}/(d*(n-1))), x] - \text{Dist}[b^2, \text{Int}[(b*\text{Tan}[c + d*x])^{(n-2)}, x], x] \text{ /; } \text{FreeQ}\{b, c, d\}, x \ \&\& \ \text{GtQ}[n, 1]$

Rubi steps

$$\begin{aligned} \int \frac{\tanh^2(d(a+b \log(cx^n)))}{x} dx &= \frac{\text{Subst}(\int \tanh^2(d(a+bx)) dx, x, \log(cx^n))}{n} \\ &= -\frac{\tanh(ad + bd \log(cx^n))}{bdn} + \frac{\text{Subst}(\int 1 dx, x, \log(cx^n))}{n} \\ &= \log(x) - \frac{\tanh(ad + bd \log(cx^n))}{bdn} \end{aligned}$$

Mathematica [A]

time = 0.07, size = 51, normalized size = 1.82

$$\frac{\tanh^{-1}(\tanh(ad + bd \log(cx^n)))}{bdn} - \frac{\tanh(ad + bd \log(cx^n))}{bdn}$$

Antiderivative was successfully verified.

[In] Integrate[Tanh[d*(a + b*Log[c*x^n])]^2/x,x]

[Out] ArcTanh[Tanh[a*d + b*d*Log[c*x^n]]]/(b*d*n) - Tanh[a*d + b*d*Log[c*x^n]]/(b*d*n)

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 62 vs. 2(28) = 56.

time = 2.08, size = 63, normalized size = 2.25

method	result
derivativedivides	$\frac{-\tanh(d(a+b\ln(cx^n))) - \frac{\ln(\tanh(d(a+b\ln(cx^n))))-1}{2} + \frac{\ln(\tanh(d(a+b\ln(cx^n))))+1}{2}}{nbd}$
default	$\frac{-\tanh(d(a+b\ln(cx^n))) - \frac{\ln(\tanh(d(a+b\ln(cx^n))))-1}{2} + \frac{\ln(\tanh(d(a+b\ln(cx^n))))+1}{2}}{nbd}$
risch	$\ln(x) + \frac{2}{dbn \left(e^{d(-ib\pi \operatorname{csgn}(icx^n)^3 + ib\pi \operatorname{csgn}(icx^n)^2 \operatorname{csgn}(ic) + ib\pi \operatorname{csgn}(icx^n)^2 \operatorname{csgn}(ix^n) - ib\pi \operatorname{csgn}(icx^n) \operatorname{csgn}(ic) \operatorname{csgn}(ix^n) + 2b \ln(d))} \right)}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(d*(a+b*ln(c*x^n)))^2/x,x,method=_RETURNVERBOSE)

[Out] 1/n/b/d*(-tanh(d*(a+b*ln(c*x^n)))-1/2*ln(tanh(d*(a+b*ln(c*x^n)))-1)+1/2*ln(tanh(d*(a+b*ln(c*x^n)))+1))

Maxima [A]

time = 0.32, size = 36, normalized size = 1.29

$$\frac{2}{bc^{2bd} dne^{(2bd \log(x^n) + 2ad)} + bdn} + \log(x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(d*(a+b*log(c*x^n)))^2/x,x, algorithm="maxima")

[Out] 2/(b*c^(2*b*d)*d*n*e^(2*b*d*log(x^n) + 2*a*d) + b*d*n) + log(x)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 72 vs. 2(28) = 56.

time = 0.51, size = 72, normalized size = 2.57

$$\frac{(bdn \log(x) + 1) \cosh(bdn \log(x) + bd \log(c) + ad) - \sinh(bdn \log(x) + bd \log(c) + ad)}{bdn \cosh(bdn \log(x) + bd \log(c) + ad)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(d*(a+b*log(c*x^n)))^2/x,x, algorithm="fricas")

[Out] ((b*d*n*log(x) + 1)*cosh(b*d*n*log(x) + b*d*log(c) + a*d) - sinh(b*d*n*log(x) + b*d*log(c) + a*d))/(b*d*n*cosh(b*d*n*log(x) + b*d*log(c) + a*d))

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 70 vs. $2(22) = 44$.

time = 4.12, size = 70, normalized size = 2.50

$$-\frac{\log(\tanh(ad + bd \log(cx^n)) - 1)}{2bdn} + \frac{\log(\tanh(ad + bd \log(cx^n)) + 1)}{2bdn} - \frac{\tanh(ad + bd \log(cx^n))}{bdn}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(d*(a+b*ln(c*x**n)))**2/x,x)

[Out] -log(tanh(a*d + b*d*log(c*x**n)) - 1)/(2*b*d*n) + log(tanh(a*d + b*d*log(c*x**n)) + 1)/(2*b*d*n) - tanh(a*d + b*d*log(c*x**n))/(b*d*n)

Giac [A]

time = 0.49, size = 37, normalized size = 1.32

$$\frac{2}{(c^{2bd}x^{2bdn}e^{2ad} + 1)bdn} + \log(x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(d*(a+b*log(c*x^n)))^2/x,x, algorithm="giac")

[Out] 2/((c^(2*b*d)*x^(2*b*d*n)*e^(2*a*d) + 1)*b*d*n) + log(x)

Mupad [B]

time = 1.06, size = 34, normalized size = 1.21

$$\ln(x) + \frac{2}{bdn \left(e^{2ad} (cx^n)^{2bd} + 1 \right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(d*(a + b*log(c*x^n)))^2/x,x)

[Out] log(x) + 2/(b*d*n*(exp(2*a*d)*(c*x^n)^(2*b*d) + 1))

$$3.184 \quad \int \frac{\tanh^2(d(a+b \log(cx^n)))}{x^2} dx$$

Optimal. Leaf size=135

$$-\frac{1 - \frac{1}{bdn}}{x} + \frac{1 - e^{2ad}(cx^n)^{2bd}}{bdnx \left(1 + e^{2ad}(cx^n)^{2bd}\right)} - \frac{{}_2F_1\left(1, -\frac{1}{2bdn}; 1 - \frac{1}{2bdn}; -e^{2ad}(cx^n)^{2bd}\right)}{bdnx}$$

[Out] $(-1+1/b/d/n)/x+(1-\exp(2*a*d)*(c*x^n)^{(2*b*d)})/b/d/n/x/(1+\exp(2*a*d)*(c*x^n)^{(2*b*d)})-2*\text{hypergeom}([1, -1/2/b/d/n], [1-1/2/b/d/n], -\exp(2*a*d)*(c*x^n)^{(2*b*d)})/b/d/n/x$

Rubi [A]

time = 0.12, antiderivative size = 135, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$, Rules used = {5658, 5656, 516, 470, 371}

$$-\frac{{}_2F_1\left(1, -\frac{1}{2bdn}; 1 - \frac{1}{2bdn}; -e^{2ad}(cx^n)^{2bd}\right)}{bdnx} + \frac{1 - e^{2ad}(cx^n)^{2bd}}{bdnx \left(e^{2ad}(cx^n)^{2bd} + 1\right)} - \frac{1 - \frac{1}{bdn}}{x}$$

Antiderivative was successfully verified.

[In] Int[Tanh[d*(a + b*Log[c*x^n])]^2/x^2,x]

[Out] $-\left(\left(1 - \frac{1}{(b*d*n)}\right)/x\right) + \left(1 - E^{(2*a*d)*(c*x^n)^{(2*b*d)}}\right)/\left(b*d*n*x*(1 + E^{(2*a*d)*(c*x^n)^{(2*b*d)})}\right) - \left(2*\text{Hypergeometric2F1}\left[1, -1/2*1/(b*d*n), 1 - 1/(2*b*d*n), -(E^{(2*a*d)*(c*x^n)^{(2*b*d)})}\right]\right)/\left(b*d*n*x\right)$

Rule 371

Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] :> Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])

Rule 470

Int[((e_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_)), x_Symbol] :> Simp[d*(e*x)^(m + 1)*((a + b*x^n)^(p + 1)/(b*e*(m + n*(p + 1) + 1))), x] - Dist[(a*d*(m + 1) - b*c*(m + n*(p + 1) + 1))/(b*(m + n*(p + 1) + 1)), Int[(e*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && NeQ[b*c - a*d, 0] && NeQ[m + n*(p + 1) + 1, 0]

Rule 516

Int[((e_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol] :> Simp[(-(c*b - a*d))*(e*x)^(m + 1)*(a + b*x^n)^(p + 1)


```

*((c + d*x^n)^(q - 1)/(a*b*e*n*(p + 1))), x] + Dist[1/(a*b*n*(p + 1)), Int[
(e*x)^m*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 2)*Simp[c*(c*b*n*(p + 1) + (c*
b - a*d)*(m + 1)) + d*(c*b*n*(p + 1) + (c*b - a*d)*(m + n*(q - 1) + 1))*x^n
, x], x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && NeQ[b*c - a*d, 0] && LtQ
[p, -1] && GtQ[q, 1] && IntBinomialQ[a, b, c, d, e, m, n, p, q, x]

```

Rule 5656

```

Int[((e_.)*(x_))^(m_.)*Tanh[((a_.) + Log[x_]*(b_.))*(d_.)]^(p_.), x_Symbol]
:> Int[(e*x)^m*((-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p),
x] /; FreeQ[{a, b, d, e, m, p}, x]

```

Rule 5658

```

Int[((e_.)*(x_))^(m_.)*Tanh[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*(d_.)]^(p
_.), x_Symbol] :> Dist[(e*x)^(m + 1)/(e*n*(c*x^n)^((m + 1)/n)), Subst[Int[x
^((m + 1)/n - 1)*Tanh[d*(a + b*Log[x])]^p, x], x, c*x^n], x] /; FreeQ[{a, b
, c, d, e, m, n, p}, x] && (NeQ[c, 1] || NeQ[n, 1])

```

Rubi steps

$$\int \frac{\tanh^2(d(a + b \log(cx^n)))}{x^2} dx = \int \frac{\tanh^2(d(a + b \log(cx^n)))}{x^2} dx$$

Mathematica [A]

time = 2.43, size = 162, normalized size = 1.20

$$\frac{e^{2d(a+b \log(cx^n))} {}_2F_1\left(1, 1 - \frac{1}{2bdn}; 2 - \frac{1}{2bdn}; -e^{2d(a+b \log(cx^n))}\right) + (-1 + 2bdn) (bdn + {}_2F_1\left(1, -\frac{1}{2bdn}; 1 - \frac{1}{2bdn}; -e^{2d(a+b \log(cx^n))}\right) + \tanh(d(a + b \log(cx^n))))}{bdn(-1 + 2bdn)x}$$

Antiderivative was successfully verified.

```
[In] Integrate[Tanh[d*(a + b*Log[c*x^n])]^2/x^2, x]
```

```
[Out] -((E^(2*d*(a + b*Log[c*x^n]))*Hypergeometric2F1[1, 1 - 1/(2*b*d*n), 2 - 1/(
2*b*d*n), -E^(2*d*(a + b*Log[c*x^n]))] + (-1 + 2*b*d*n)*(b*d*n + Hypergeome
tric2F1[1, -1/2*1/(b*d*n), 1 - 1/(2*b*d*n), -E^(2*d*(a + b*Log[c*x^n]))] +
Tanh[d*(a + b*Log[c*x^n])]))/(b*d*n*(-1 + 2*b*d*n)*x))

```

Maple [F]

time = 0.37, size = 0, normalized size = 0.00

$$\int \frac{\tanh^2(d(a + b \ln(cx^n)))}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tanh(d*(a+b*ln(c*x^n)))^2/x^2,x)`

[Out] `int(tanh(d*(a+b*ln(c*x^n)))^2/x^2,x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(d*(a+b*log(c*x^n)))^2/x^2,x, algorithm="maxima")`

[Out] `-(b*c^(2*b*d)*d*n*e^(2*b*d*log(x^n) + 2*a*d) + b*d*n - 2)/(b*c^(2*b*d)*d*n*x*e^(2*b*d*log(x^n) + 2*a*d) + b*d*n*x) + 2*integrate(1/(b*c^(2*b*d)*d*n*x^2*e^(2*b*d*log(x^n) + 2*a*d) + b*d*n*x^2), x)`

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(d*(a+b*log(c*x^n)))^2/x^2,x, algorithm="fricas")`

[Out] `integral(tanh(b*d*log(c*x^n) + a*d)^2/x^2, x)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\tanh^2(ad + bd \log(cx^n))}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(d*(a+b*ln(c*x**n)))**2/x**2,x)`

[Out] `Integral(tanh(a*d + b*d*log(c*x**n))**2/x**2, x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(d*(a+b*log(c*x^n)))^2/x^2,x, algorithm="giac")`

[Out] `integrate(tanh((b*log(c*x^n) + a)*d)^2/x^2, x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\tanh(d(a + b \ln(cx^n)))^2}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(d*(a + b*log(c*x^n)))^2/x^2,x)

[Out] int(tanh(d*(a + b*log(c*x^n)))^2/x^2, x)

$$3.185 \quad \int \frac{\tanh^2(d(a+b \log(cx^n)))}{x^3} dx$$

Optimal. Leaf size=136

$$\frac{2 - bdn}{2bdnx^2} + \frac{1 - e^{2ad}(cx^n)^{2bd}}{bdnx^2 (1 + e^{2ad}(cx^n)^{2bd})} - \frac{{}_2F_1\left(1, -\frac{1}{bdn}; 1 - \frac{1}{bdn}; -e^{2ad}(cx^n)^{2bd}\right)}{bdnx^2}$$

[Out] $1/2*(-b*d*n+2)/b/d/n/x^2+(1-\exp(2*a*d)*(c*x^n)^{(2*b*d)})/b/d/n/x^2/(1+\exp(2*a*d)*(c*x^n)^{(2*b*d)})-2*\text{hypergeom}([1, -1/b/d/n], [1-1/b/d/n], -\exp(2*a*d)*(c*x^n)^{(2*b*d)})/b/d/n/x^2$

Rubi [A]

time = 0.12, antiderivative size = 136, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$, Rules used = {5658, 5656, 516, 470, 371}

$$-\frac{{}_2F_1\left(1, -\frac{1}{bdn}; 1 - \frac{1}{bdn}; -e^{2ad}(cx^n)^{2bd}\right)}{bdnx^2} + \frac{1 - e^{2ad}(cx^n)^{2bd}}{bdnx^2 (e^{2ad}(cx^n)^{2bd} + 1)} + \frac{2 - bdn}{2bdnx^2}$$

Antiderivative was successfully verified.

[In] Int[Tanh[d*(a + b*Log[c*x^n])]^2/x^3,x]

[Out] $(2 - b*d*n)/(2*b*d*n*x^2) + (1 - E^{(2*a*d)*(c*x^n)^{(2*b*d)}})/(b*d*n*x^2*(1 + E^{(2*a*d)*(c*x^n)^{(2*b*d)}})) - (2*\text{Hypergeometric2F1}[1, -(1/(b*d*n)), 1 - 1/(b*d*n), -(E^{(2*a*d)*(c*x^n)^{(2*b*d)}})]/(b*d*n*x^2)$

Rule 371

Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p*((c*x)^(m+1)/(c*(m+1)))*Hypergeometric2F1[-p, (m+1)/n, (m+1)/n+1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])

Rule 470

Int[((e_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Simp[d*(e*x)^(m+1)*((a + b*x^n)^(p+1)/(b*e*(m+n*(p+1)+1))), x] - Dist[(a*d*(m+1) - b*c*(m+n*(p+1)+1))/(b*(m+n*(p+1)+1)), Int[(e*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && NeQ[b*c - a*d, 0] && NeQ[m + n*(p+1) + 1, 0]

Rule 516

Int[((e_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol] := Simp[(-(c*b - a*d))*(e*x)^(m+1)*(a + b*x^n)^(p+1)

```

*((c + d*x^n)^(q - 1)/(a*b*e*n*(p + 1))), x] + Dist[1/(a*b*n*(p + 1)), Int[
(e*x)^m*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 2)*Simp[c*(c*b*n*(p + 1) + (c*
b - a*d)*(m + 1)) + d*(c*b*n*(p + 1) + (c*b - a*d)*(m + n*(q - 1) + 1))*x^n
, x], x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && NeQ[b*c - a*d, 0] && LtQ
[p, -1] && GtQ[q, 1] && IntBinomialQ[a, b, c, d, e, m, n, p, q, x]

```

Rule 5656

```

Int[((e_.)*(x_))^(m_.)*Tanh[((a_.) + Log[x_]*(b_.))*(d_.)]^(p_.), x_Symbol]
:> Int[(e*x)^m*((-1 + E^(2*a*d))*x^(2*b*d))^p/(1 + E^(2*a*d))*x^(2*b*d))^p,
x] /; FreeQ[{a, b, d, e, m, p}, x]

```

Rule 5658

```

Int[((e_.)*(x_))^(m_.)*Tanh[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*(d_.)]^(p
_.), x_Symbol] :> Dist[(e*x)^(m + 1)/(e*n*(c*x^n)^((m + 1)/n)), Subst[Int[x
^((m + 1)/n - 1)*Tanh[d*(a + b*Log[x])]^p, x], x, c*x^n], x] /; FreeQ[{a, b
, c, d, e, m, n, p}, x] && (NeQ[c, 1] || NeQ[n, 1])

```

Rubi steps

$$\int \frac{\tanh^2(d(a + b \log(cx^n)))}{x^3} dx = \int \frac{\tanh^2(d(a + b \log(cx^n)))}{x^3} dx$$

Mathematica [A]

time = 2.34, size = 159, normalized size = 1.17

$$\frac{2e^{2d(a+b \log(cx^n))} {}_2F_1\left(1, 1 - \frac{1}{bdn}; 2 - \frac{1}{bdn}; -e^{2d(a+b \log(cx^n))}\right) + (-1 + bdn) \left(bdn + 2 {}_2F_1\left(1, -\frac{1}{bdn}; 1 - \frac{1}{bdn}; -e^{2d(a+b \log(cx^n))}\right) + 2 \tanh(d(a + b \log(cx^n)))\right)}{2bdn(-1 + bdn)x^2}$$

Antiderivative was successfully verified.

```
[In] Integrate[Tanh[d*(a + b*Log[c*x^n])]^2/x^3, x]
```

```
[Out] -1/2*(2*E^(2*d*(a + b*Log[c*x^n]))*Hypergeometric2F1[1, 1 - 1/(b*d*n), 2 -
1/(b*d*n), -E^(2*d*(a + b*Log[c*x^n]))] + (-1 + b*d*n)*(b*d*n + 2*Hypergeom
etric2F1[1, -(1/(b*d*n)), 1 - 1/(b*d*n), -E^(2*d*(a + b*Log[c*x^n]))] + 2*T
anh[d*(a + b*Log[c*x^n])]))/(b*d*n*(-1 + b*d*n)*x^2)

```

Maple [F]

time = 0.36, size = 0, normalized size = 0.00

$$\int \frac{\tanh^2(d(a + b \ln(cx^n)))}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(d*(a+b*ln(c*x^n)))^2/x^3,x)

[Out] int(tanh(d*(a+b*ln(c*x^n)))^2/x^3,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(d*(a+b*log(c*x^n)))^2/x^3,x, algorithm="maxima")

[Out] -1/2*(b*c^(2*b*d)*d*n*e^(2*b*d*log(x^n) + 2*a*d) + b*d*n - 4)/(b*c^(2*b*d)*d*n*x^2*e^(2*b*d*log(x^n) + 2*a*d) + b*d*n*x^2) + 4*integrate(1/(b*c^(2*b*d)*d*n*x^3*e^(2*b*d*log(x^n) + 2*a*d) + b*d*n*x^3), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(d*(a+b*log(c*x^n)))^2/x^3,x, algorithm="fricas")

[Out] integral(tanh(b*d*log(c*x^n) + a*d)^2/x^3, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\tanh^2(ad + bd \log(cx^n))}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(d*(a+b*ln(c*x**n)))**2/x**3,x)

[Out] Integral(tanh(a*d + b*d*log(c*x**n))**2/x**3, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(d*(a+b*log(c*x^n)))^2/x^3,x, algorithm="giac")

[Out] integrate(tanh((b*log(c*x^n) + a)*d)^2/x^3, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\tanh(d(a + b \ln(cx^n)))^2}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(d*(a + b*log(c*x^n)))^2/x^3,x)

[Out] int(tanh(d*(a + b*log(c*x^n)))^2/x^3, x)

$$3.186 \quad \int \frac{\tanh^3(a+b \log(cx^n))}{x} dx$$

Optimal. Leaf size=43

$$\frac{\log(\cosh(a+b \log(cx^n)))}{bn} - \frac{\tanh^2(a+b \log(cx^n))}{2bn}$$

[Out] ln(cosh(a+b*ln(c*x^n)))/b/n-1/2*tanh(a+b*ln(c*x^n))^2/b/n

Rubi [A]

time = 0.03, antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {3554, 3556}

$$\frac{\log(\cosh(a+b \log(cx^n)))}{bn} - \frac{\tanh^2(a+b \log(cx^n))}{2bn}$$

Antiderivative was successfully verified.

[In] Int[Tanh[a + b*Log[c*x^n]]^3/x,x]

[Out] Log[Cosh[a + b*Log[c*x^n]]]/(b*n) - Tanh[a + b*Log[c*x^n]]^2/(2*b*n)

Rule 3554

Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n-1)/(d*(n-1))), x] - Dist[b^2, Int[(b*Tan[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]

Rule 3556

Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{\tanh^3(a+b \log(cx^n))}{x} dx &= \frac{\text{Subst}(\int \tanh^3(a+bx) dx, x, \log(cx^n))}{n} \\ &= -\frac{\tanh^2(a+b \log(cx^n))}{2bn} + \frac{\text{Subst}(\int \tanh(a+bx) dx, x, \log(cx^n))}{n} \\ &= \frac{\log(\cosh(a+b \log(cx^n)))}{bn} - \frac{\tanh^2(a+b \log(cx^n))}{2bn} \end{aligned}$$

Mathematica [A]

time = 0.11, size = 43, normalized size = 1.00

$$\frac{\log(\cosh(a+b \log(cx^n)))}{bn} - \frac{\tanh^2(a+b \log(cx^n))}{2bn}$$

Antiderivative was successfully verified.

[In] Integrate[Tanh[a + b*Log[c*x^n]]^3/x,x]

[Out] Log[Cosh[a + b*Log[c*x^n]]]/(b*n) - Tanh[a + b*Log[c*x^n]]^2/(2*b*n)

Maple [A]

time = 2.07, size = 56, normalized size = 1.30

method	result
derivativedivides	$\frac{\frac{\tanh^2(a+b \ln(c x^n))}{2} - \frac{\ln(\tanh(a+b \ln(c x^n))-1)}{2} - \frac{\ln(\tanh(a+b \ln(c x^n))+1)}{2}}{nb}$
default	$\frac{\frac{\tanh^2(a+b \ln(c x^n))}{2} - \frac{\ln(\tanh(a+b \ln(c x^n))-1)}{2} - \frac{\ln(\tanh(a+b \ln(c x^n))+1)}{2}}{nb}$
risch	$\ln(x) - \frac{2a}{nb} - \frac{2\ln(c)}{n} - \frac{2\ln(x^n)}{n} + \frac{i\pi \operatorname{csgn}(ic x^n)^3}{n} - \frac{i\pi \operatorname{csgn}(ic x^n)^2 \operatorname{csgn}(ic)}{n} - \frac{i\pi \operatorname{csgn}(ic x^n)^2 \operatorname{csgn}(ix^n)}{n} + \dots$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(a+b*ln(c*x^n))^3/x,x,method=_RETURNVERBOSE)

[Out] 1/n/b*(-1/2*tanh(a+b*ln(c*x^n))^2-1/2*ln(tanh(a+b*ln(c*x^n))-1)-1/2*ln(tanh(a+b*ln(c*x^n))+1))

Maxima [B] Leaf count of result is larger than twice the leaf count of optimal. 304 vs. 2(41) = 82.

time = 0.31, size = 304, normalized size = 7.07

$$\frac{\frac{4c^{2b}(2b \log(x^n)+2a)+3}{4(b^{4b}n e^{4b \log(x^n)+4a}+2b^{2b}n e^{2b \log(x^n)+2a}+bn)} - \frac{2c^{2b}(2b \log(x^n)+2a)+3}{4(b^{4b}n e^{4b \log(x^n)+4a}+2b^{2b}n e^{2b \log(x^n)+2a}+bn)} + \frac{3(2c^{2b}(2b \log(x^n)+2a)+1)}{4(b^{4b}n e^{4b \log(x^n)+4a}+2b^{2b}n e^{2b \log(x^n)+2a}+bn)} - \frac{3}{4(b^{4b}n e^{4b \log(x^n)+4a}+2b^{2b}n e^{2b \log(x^n)+2a}+bn)} + \frac{\log\left(\frac{c^{2b(2b \log(x^n)+2a)+1}e^{-2a}}{bn}\right)}{bn} - \log(x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+b*log(c*x^n))^3/x,x, algorithm="maxima")

[Out] 1/4*(4*c^(2*b)*e^(2*b*log(x^n) + 2*a) + 3)/(b*c^(4*b)*n*e^(4*b*log(x^n) + 4*a) + 2*b*c^(2*b)*n*e^(2*b*log(x^n) + 2*a) + b*n) - 1/4*(2*c^(2*b)*e^(2*b*log(x^n) + 2*a) + 3)/(b*c^(4*b)*n*e^(4*b*log(x^n) + 4*a) + 2*b*c^(2*b)*n*e^(2*b*log(x^n) + 2*a) + b*n) + 3/4*(2*c^(2*b)*e^(2*b*log(x^n) + 2*a) + 1)/(b*c^(4*b)*n*e^(4*b*log(x^n) + 4*a) + 2*b*c^(2*b)*n*e^(2*b*log(x^n) + 2*a) + b*n) - 3/4/(b*c^(4*b)*n*e^(4*b*log(x^n) + 4*a) + 2*b*c^(2*b)*n*e^(2*b*log(x^n) + 2*a) + b*n) + log((c^(2*b)*e^(2*b*log(x^n) + 2*a) + 1)*e^(-2*a)/c^(2*b)))/(b*n) - log(x)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 566 vs. 2(41) = 82.

time = 0.38, size = 566, normalized size = 13.16

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+b*log(c*x^n))^3/x,x, algorithm="fricas")

[Out] $-(b*n*\cosh(b*n*\log(x) + b*\log(c) + a)^4*\log(x) + 4*b*n*\cosh(b*n*\log(x) + b*\log(c) + a)*\log(x)*\sinh(b*n*\log(x) + b*\log(c) + a)^3 + b*n*\log(x)*\sinh(b*n*\log(x) + b*\log(c) + a)^4 + 2*(b*n*\log(x) - 1)*\cosh(b*n*\log(x) + b*\log(c) + a)^2 + b*n*\log(x) + 2*(3*b*n*\cosh(b*n*\log(x) + b*\log(c) + a)^2*\log(x) + b*n*\log(x) - 1)*\sinh(b*n*\log(x) + b*\log(c) + a)^2 - (\cosh(b*n*\log(x) + b*\log(c) + a)^4 + 4*\cosh(b*n*\log(x) + b*\log(c) + a)*\sinh(b*n*\log(x) + b*\log(c) + a)^3 + \sinh(b*n*\log(x) + b*\log(c) + a)^4 + 2*(3*\cosh(b*n*\log(x) + b*\log(c) + a)^2 + 1)*\sinh(b*n*\log(x) + b*\log(c) + a)^2 + 2*\cosh(b*n*\log(x) + b*\log(c) + a)^2 + 4*(\cosh(b*n*\log(x) + b*\log(c) + a)^3 + \cosh(b*n*\log(x) + b*\log(c) + a))*\sinh(b*n*\log(x) + b*\log(c) + a) + 1)*\log(2*\cosh(b*n*\log(x) + b*\log(c) + a)/(cosh(b*n*\log(x) + b*\log(c) + a) - sinh(b*n*\log(x) + b*\log(c) + a))) + 4*(b*n*\cosh(b*n*\log(x) + b*\log(c) + a)^3*\log(x) + (b*n*\log(x) - 1)*\cosh(b*n*\log(x) + b*\log(c) + a))*\sinh(b*n*\log(x) + b*\log(c) + a))/(b*n*\cosh(b*n*\log(x) + b*\log(c) + a)^4 + 4*b*n*\cosh(b*n*\log(x) + b*\log(c) + a)*\sinh(b*n*\log(x) + b*\log(c) + a)^3 + b*n*\sinh(b*n*\log(x) + b*\log(c) + a)^4 + 2*b*n*\cosh(b*n*\log(x) + b*\log(c) + a)^2 + 2*(3*b*n*\cosh(b*n*\log(x) + b*\log(c) + a)^2 + b*n)*\sinh(b*n*\log(x) + b*\log(c) + a)^2 + b*n + 4*(b*n*\cosh(b*n*\log(x) + b*\log(c) + a)^3 + b*n*\cosh(b*n*\log(x) + b*\log(c) + a))*\sinh(b*n*\log(x) + b*\log(c) + a))$

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 73 vs. $2(34) = 68$.

time = 0.77, size = 73, normalized size = 1.70

$$\begin{cases} \log(x) \tanh^3(a) & \text{for } b = 0 \wedge n = 0 \\ \log(x) \tanh^3(a + b \log(c)) & \text{for } n = 0 \\ \log(x) \tanh^3(a) & \text{for } b = 0 \\ \frac{\log(cx^n)}{n} - \frac{\log(\tanh(a+b \log(cx^n))+1)}{bn} - \frac{\tanh^2(a+b \log(cx^n))}{2bn} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+b*ln(c*x**n))**3/x,x)

[Out] Piecewise((log(x)*tanh(a)**3, Eq(b, 0) & Eq(n, 0)), (log(x)*tanh(a + b*log(c))**3, Eq(n, 0)), (log(x)*tanh(a)**3, Eq(b, 0)), (log(c*x**n)/n - log(tanh(a + b*log(c*x**n)) + 1)/(b*n) - tanh(a + b*log(c*x**n))**2/(2*b*n), True))

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 127 vs. $2(41) = 82$.

time = 0.47, size = 127, normalized size = 2.95

$$\frac{\log\left(\sqrt{2x^{2bn}|c|^{2b}\cos(\pi b \operatorname{sgn}(c) - \pi b)e^{(2a)} + x^{4bn}|c|^{4b}e^{(4a)} + 1}\right)}{bn} - \frac{3c^{4b}x^{4bn}e^{(4a)} + 2c^{2b}x^{2bn}e^{(2a)} + 3}{2(c^{2b}x^{2bn}e^{(2a)} + 1)^2bn} - \log(x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+b*log(c*x^n))^3/x,x, algorithm="giac")

[Out] $\frac{\log(\sqrt{2x^{2bn} \operatorname{abs}(c)^{2b} \cos(\pi b \operatorname{sgn}(c) - \pi b) e^{2a} + x^{4bn} \operatorname{abs}(c)^{4b} e^{4a} + 1})}{bn} - \frac{1}{2} \frac{3c^{4b} x^{4bn} e^{4a} + 2c^{2b} x^{2bn} e^{2a} + 3}{(c^{2b} x^{2bn} e^{2a} + 1)^{2bn}} - \log(x)$

Mupad [B]

time = 1.07, size = 94, normalized size = 2.19

$$\frac{2}{bn + bne^{2a}(cx^n)^{2b}} - \ln(x) - \frac{2}{bn + 2bne^{2a}(cx^n)^{2b} + bne^{4a}(cx^n)^{4b}} + \frac{\ln(e^{2a}(cx^n)^{2b} + 1)}{bn}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(a + b*log(c*x^n))^3/x,x)

[Out] $\frac{2}{bn + bn \exp(2a) (cx^n)^{2b}} - \log(x) - \frac{2}{bn + 2bn \exp(2a) (cx^n)^{2b} + bn \exp(4a) (cx^n)^{4b}} + \frac{\log(\exp(2a) (cx^n)^{2b} + 1)}{bn}$

$$3.187 \quad \int \frac{\tanh^4(a+b \log(cx^n))}{x} dx$$

Optimal. Leaf size=45

$$\log(x) - \frac{\tanh(a+b \log(cx^n))}{bn} - \frac{\tanh^3(a+b \log(cx^n))}{3bn}$$

[Out] $\ln(x) - \tanh(a+b*\ln(c*x^n))/b/n - 1/3*\tanh(a+b*\ln(c*x^n))^3/b/n$

Rubi [A]

time = 0.03, antiderivative size = 45, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {3554, 8}

$$-\frac{\tanh^3(a+b \log(cx^n))}{3bn} - \frac{\tanh(a+b \log(cx^n))}{bn} + \log(x)$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Tanh}[a + b*\text{Log}[c*x^n]]^4/x, x]$

[Out] $\text{Log}[x] - \text{Tanh}[a + b*\text{Log}[c*x^n]]/(b*n) - \text{Tanh}[a + b*\text{Log}[c*x^n]]^3/(3*b*n)$

Rule 8

$\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

Rule 3554

$\text{Int}[(b*.)*\text{tan}[(c*.) + (d*.)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Simp}[b*((b*\text{Tan}[c + d*x])^{(n-1)})/(d*(n-1)), x] - \text{Dist}[b^2, \text{Int}[(b*\text{Tan}[c + d*x])^{(n-2)}, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \&\& \text{GtQ}[n, 1]$

Rubi steps

$$\begin{aligned} \int \frac{\tanh^4(a+b \log(cx^n))}{x} dx &= \frac{\text{Subst}(\int \tanh^4(a+bx) dx, x, \log(cx^n))}{n} \\ &= -\frac{\tanh^3(a+b \log(cx^n))}{3bn} + \frac{\text{Subst}(\int \tanh^2(a+bx) dx, x, \log(cx^n))}{n} \\ &= -\frac{\tanh(a+b \log(cx^n))}{bn} - \frac{\tanh^3(a+b \log(cx^n))}{3bn} + \frac{\text{Subst}(\int 1 dx, x, \log(cx^n))}{n} \\ &= \log(x) - \frac{\tanh(a+b \log(cx^n))}{bn} - \frac{\tanh^3(a+b \log(cx^n))}{3bn} \end{aligned}$$

Mathematica [A]

time = 0.08, size = 62, normalized size = 1.38

$$\frac{\tanh^{-1}(\tanh(a + b \log(cx^n)))}{bn} - \frac{\tanh(a + b \log(cx^n))}{bn} - \frac{\tanh^3(a + b \log(cx^n))}{3bn}$$

Antiderivative was successfully verified.

`[In] Integrate[Tanh[a + b*Log[c*x^n]]^4/x,x]``[Out] ArcTanh[Tanh[a + b*Log[c*x^n]]]/(b*n) - Tanh[a + b*Log[c*x^n]]/(b*n) - Tanh[a + b*Log[c*x^n]]^3/(3*b*n)`**Maple [A]**

time = 2.02, size = 69, normalized size = 1.53

method	result
derivativedivides	$-\frac{(\tanh^3(a+b \ln(cx^n)))}{3} - \tanh(a+b \ln(cx^n)) - \frac{\ln(\tanh(a+b \ln(cx^n))-1)}{2nb} + \frac{\ln(\tanh(a+b \ln(cx^n))+1)}{2nb}$
default	$-\frac{(\tanh^3(a+b \ln(cx^n)))}{3} - \tanh(a+b \ln(cx^n)) - \frac{\ln(\tanh(a+b \ln(cx^n))-1)}{2nb} + \frac{\ln(\tanh(a+b \ln(cx^n))+1)}{2nb}$
risch	$\ln(x) + \frac{4(x^n)^{4b} c^{4b} e^{4a} e^{-2ib\pi \operatorname{csgn}(icx^n)^3} e^{2ib\pi \operatorname{csgn}(icx^n)^2 \operatorname{csgn}(ic)} e^{2ib\pi \operatorname{csgn}(icx^n)^2 \operatorname{csgn}(ix^n)} e^{-2ib\pi \operatorname{csgn}(icx^n) \operatorname{csgn}(ic)} e^{ib\pi \operatorname{csgn}(icx^n)^2 \operatorname{csgn}(ic)}}{bn((x^n)^{2b} c^{2b} e^{2a} e^{-ib\pi \operatorname{csgn}(icx^n)^3} e^{ib\pi \operatorname{csgn}(icx^n)^2 \operatorname{csgn}(ic)})}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(tanh(a+b*ln(c*x^n))^4/x,x,method=_RETURNVERBOSE)``[Out] 1/n/b*(-1/3*tanh(a+b*ln(c*x^n))^3-tanh(a+b*ln(c*x^n))-1/2*ln(tanh(a+b*ln(c*x^n))-1)+1/2*ln(tanh(a+b*ln(c*x^n))+1))`**Maxima [B]** Leaf count of result is larger than twice the leaf count of optimal. 494 vs. 2(43) = 86.

time = 0.35, size = 494, normalized size = 10.98

$$\frac{18c^{4b}e^{4a}e^{-2ib\pi \operatorname{csgn}(icx^n)^3}e^{2ib\pi \operatorname{csgn}(icx^n)^2 \operatorname{csgn}(ic)}e^{2ib\pi \operatorname{csgn}(icx^n)^2 \operatorname{csgn}(ix^n)}e^{-2ib\pi \operatorname{csgn}(icx^n) \operatorname{csgn}(ic)}e^{ib\pi \operatorname{csgn}(icx^n)^2 \operatorname{csgn}(ic)}}{bn((x^n)^{2b}c^{2b}e^{2a}e^{-ib\pi \operatorname{csgn}(icx^n)^3}e^{ib\pi \operatorname{csgn}(icx^n)^2 \operatorname{csgn}(ic)})} + \ln(x)$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(tanh(a+b*log(c*x^n))^4/x,x, algorithm="maxima")`
`[Out] 1/12*(18*c^(4*b)*e^(4*b*log(x^n) + 4*a) + 27*c^(2*b)*e^(2*b*log(x^n) + 2*a) + 11)/(b*c^(6*b)*n*e^(6*b*log(x^n) + 6*a) + 3*b*c^(4*b)*n*e^(4*b*log(x^n) + 4*a) + 3*b*c^(2*b)*n*e^(2*b*log(x^n) + 2*a) + b*n) + 1/12*(6*c^(4*b)*e^(4*b*log(x^n) + 4*a) + 15*c^(2*b)*e^(2*b*log(x^n) + 2*a) + 11)/(b*c^(6*b)*n*e^(6*b*log(x^n) + 6*a) + 3*b*c^(4*b)*n*e^(4*b*log(x^n) + 4*a) + 3*b*c^(2*b)*n*e^(2*b*log(x^n) + 2*a) + b*n) + 2/3*(3*c^(4*b)*e^(4*b*log(x^n) + 4*a) + 3*c^(2*b)*e^(2*b*log(x^n) + 2*a) + 1)/(b*c^(6*b)*n*e^(6*b*log(x^n) + 6*a) +`

$$3*b*c^{(4*b)*n}*e^{(4*b*\log(x^n) + 4*a) + 3*b*c^{(2*b)*n}*e^{(2*b*\log(x^n) + 2*a) + b*n} - 1/2*(3*c^{(2*b)*n}*e^{(2*b*\log(x^n) + 2*a) + 1)/(b*c^{(6*b)*n}*e^{(6*b*\log(x^n) + 6*a) + 3*b*c^{(4*b)*n}*e^{(4*b*\log(x^n) + 4*a) + 3*b*c^{(2*b)*n}*e^{(2*b*\log(x^n) + 2*a) + b*n} + 2/3/(b*c^{(6*b)*n}*e^{(6*b*\log(x^n) + 6*a) + 3*b*c^{(4*b)*n}*e^{(4*b*\log(x^n) + 4*a) + 3*b*c^{(2*b)*n}*e^{(2*b*\log(x^n) + 2*a) + b*n} + \log(x)$$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 194 vs. $2(43) = 86$.

time = 0.35, size = 194, normalized size = 4.31

$$\frac{(3bn \log(x) + 4) \cosh(bn \log(x) + b \log(c) + a)^3 + 3(3bn \log(x) + 4) \cosh(bn \log(x) + b \log(c) + a) \sinh(bn \log(x) + b \log(c) + a)^2 - 12 \cosh(bn \log(x) + b \log(c) + a)^2 \sinh(bn \log(x) + b \log(c) + a) - 4 \sinh(bn \log(x) + b \log(c) + a)^3 + 3(3bn \log(x) + 4) \cosh(bn \log(x) + b \log(c) + a)}{3(bn \cosh(bn \log(x) + b \log(c) + a)^3 + 3bn \cosh(bn \log(x) + b \log(c) + a) \sinh(bn \log(x) + b \log(c) + a)^2 + 3bn \cosh(bn \log(x) + b \log(c) + a))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+b*log(c*x^n))^4/x,x, algorithm="fricas")

[Out] $\frac{1}{3} * ((3*b*n*\log(x) + 4) * \cosh(b*n*\log(x) + b*\log(c) + a)^3 + 3*(3*b*n*\log(x) + 4) * \cosh(b*n*\log(x) + b*\log(c) + a) * \sinh(b*n*\log(x) + b*\log(c) + a)^2 - 12 * \cosh(b*n*\log(x) + b*\log(c) + a)^2 * \sinh(b*n*\log(x) + b*\log(c) + a) - 4 * \sinh(b*n*\log(x) + b*\log(c) + a)^3 + 3*(3*b*n*\log(x) + 4) * \cosh(b*n*\log(x) + b*\log(c) + a)) / (b*n * \cosh(b*n*\log(x) + b*\log(c) + a)^3 + 3*b*n * \cosh(b*n*\log(x) + b*\log(c) + a) * \sinh(b*n*\log(x) + b*\log(c) + a)^2 + 3*b*n * \cosh(b*n*\log(x) + b*\log(c) + a))$

Sympy [A]

time = 1.46, size = 70, normalized size = 1.56

$$\begin{cases} \log(x) \tanh^4(a) & \text{for } b = 0 \wedge n = 0 \\ \log(x) \tanh^4(a + b \log(c)) & \text{for } n = 0 \\ \log(x) \tanh^4(a) & \text{for } b = 0 \\ \frac{\log(cx^n)}{n} - \frac{\tanh^3(a + b \log(cx^n))}{3bn} - \frac{\tanh(a + b \log(cx^n))}{bn} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+b*ln(c*x**n))**4/x,x)

[Out] Piecewise((log(x)*tanh(a)**4, Eq(b, 0) & Eq(n, 0)), (log(x)*tanh(a + b*log(c))**4, Eq(n, 0)), (log(x)*tanh(a)**4, Eq(b, 0)), (log(c*x**n)/n - tanh(a + b*log(c*x**n))**3/(3*b*n) - tanh(a + b*log(c*x**n))/(b*n), True))

Giac [A]

time = 0.49, size = 67, normalized size = 1.49

$$\frac{4(3c^{4b}x^{4bn}e^{(4a)} + 3c^{2b}x^{2bn}e^{(2a)} + 2)}{3(c^{2b}x^{2bn}e^{(2a)} + 1)^3bn} + \log(x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+b*log(c*x^n))^4/x,x, algorithm="giac")

[Out] $\frac{4}{3} \frac{3c^{4b}x^{4bn}e^{4a} + 3c^{2b}x^{2bn}e^{2a} + 2}{(c^{2b}x^{2bn}e^{2a} + 1)^{3bn} + \log(x)}$

Mupad [B]

time = 1.09, size = 162, normalized size = 3.60

$$\ln(x) + \frac{\frac{4}{3bn} + \frac{4e^{4a}(cx^n)^{4b}}{3bn}}{3e^{2a}(cx^n)^{2b} + 3e^{4a}(cx^n)^{4b} + e^{6a}(cx^n)^{6b} + 1} + \frac{4}{3bn(e^{2a}(cx^n)^{2b} + 1)} + \frac{4e^{2a}(cx^n)^{2b}}{3bn(2e^{2a}(cx^n)^{2b} + e^{4a}(cx^n)^{4b} + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(a + b*log(c*x^n))^4/x,x)

[Out] $\log(x) + \frac{4}{3bn} + \frac{4\exp(4a)(cx^n)^{4b}}{3bn} / \frac{3\exp(2a)(cx^n)^{2b} + 3\exp(4a)(cx^n)^{4b} + \exp(6a)(cx^n)^{6b} + 1}{3bn} + \frac{4}{3bn} \frac{\exp(2a)(cx^n)^{2b} + 1}{\exp(2a)(cx^n)^{2b} + 1} + \frac{4\exp(2a)(cx^n)^{2b}}{3bn(2\exp(2a)(cx^n)^{2b} + \exp(4a)(cx^n)^{4b} + 1)}$

$$3.188 \quad \int \frac{\tanh^5(a+b \log(cx^n))}{x} dx$$

Optimal. Leaf size=66

$$\frac{\log(\cosh(a+b \log(cx^n)))}{bn} - \frac{\tanh^2(a+b \log(cx^n))}{2bn} - \frac{\tanh^4(a+b \log(cx^n))}{4bn}$$

[Out] $\ln(\cosh(a+b*\ln(c*x^n)))/b/n-1/2*\tanh(a+b*\ln(c*x^n))^2/b/n-1/4*\tanh(a+b*\ln(c*x^n))^4/b/n$

Rubi [A]

time = 0.04, antiderivative size = 66, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {3554, 3556}

$$-\frac{\tanh^4(a+b \log(cx^n))}{4bn} - \frac{\tanh^2(a+b \log(cx^n))}{2bn} + \frac{\log(\cosh(a+b \log(cx^n)))}{bn}$$

Antiderivative was successfully verified.

[In] Int[Tanh[a + b*Log[c*x^n]]^5/x, x]

[Out] Log[Cosh[a + b*Log[c*x^n]]]/(b*n) - Tanh[a + b*Log[c*x^n]]^2/(2*b*n) - Tanh[a + b*Log[c*x^n]]^4/(4*b*n)

Rule 3554

Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Dist[b^2, Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]

Rule 3556

Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{\tanh^5(a+b \log(cx^n))}{x} dx &= \frac{\text{Subst}(\int \tanh^5(a+bx) dx, x, \log(cx^n))}{n} \\ &= -\frac{\tanh^4(a+b \log(cx^n))}{4bn} + \frac{\text{Subst}(\int \tanh^3(a+bx) dx, x, \log(cx^n))}{n} \\ &= -\frac{\tanh^2(a+b \log(cx^n))}{2bn} - \frac{\tanh^4(a+b \log(cx^n))}{4bn} + \frac{\text{Subst}(\int \tanh(a+bx) dx, x, \log(cx^n))}{n} \\ &= \frac{\log(\cosh(a+b \log(cx^n)))}{bn} - \frac{\tanh^2(a+b \log(cx^n))}{2bn} - \frac{\tanh^4(a+b \log(cx^n))}{4bn} \end{aligned}$$

Mathematica [A]

time = 0.15, size = 55, normalized size = 0.83

$$\frac{4 \log(\cosh(a + b \log(cx^n))) - 2 \tanh^2(a + b \log(cx^n)) - \tanh^4(a + b \log(cx^n))}{4bn}$$

Antiderivative was successfully verified.

`[In] Integrate[Tanh[a + b*Log[c*x^n]]^5/x,x]``[Out] (4*Log[Cosh[a + b*Log[c*x^n]]] - 2*Tanh[a + b*Log[c*x^n]]^2 - Tanh[a + b*Log[c*x^n]]^4)/(4*b*n)`**Maple [A]**

time = 2.28, size = 71, normalized size = 1.08

method	result
derivativedivides	$\frac{\frac{(\tanh^4(a+b \ln(cx^n)))}{4} - \frac{(\tanh^2(a+b \ln(cx^n)))}{2} - \frac{\ln(\tanh(a+b \ln(cx^n))-1)}{2} - \frac{\ln(\tanh(a+b \ln(cx^n))+1)}{2}}{nb}$
default	$\frac{\frac{(\tanh^4(a+b \ln(cx^n)))}{4} - \frac{(\tanh^2(a+b \ln(cx^n)))}{2} - \frac{\ln(\tanh(a+b \ln(cx^n))-1)}{2} - \frac{\ln(\tanh(a+b \ln(cx^n))+1)}{2}}{nb}$
risch	$\ln(x) - \frac{2a}{nb} - \frac{2 \ln(c)}{n} - \frac{2 \ln(x^n)}{n} + \frac{i\pi \operatorname{csgn}(icx^n)^3}{n} - \frac{i\pi \operatorname{csgn}(icx^n)^2 \operatorname{csgn}(ic)}{n} - \frac{i\pi \operatorname{csgn}(icx^n)^2 \operatorname{csgn}(ix^n)}{n} + \dots$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(tanh(a+b*ln(c*x^n))^5/x,x,method=_RETURNVERBOSE)``[Out] 1/n/b*(-1/4*tanh(a+b*ln(c*x^n))^4-1/2*tanh(a+b*ln(c*x^n))^2-1/2*ln(tanh(a+b*ln(c*x^n))-1)-1/2*ln(tanh(a+b*ln(c*x^n))+1))`**Maxima [B]** Leaf count of result is larger than twice the leaf count of optimal. 829 vs. 2(62) = 124.

time = 0.37, size = 829, normalized size = 12.56

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(tanh(a+b*log(c*x^n))^5/x,x, algorithm="maxima")`
`[Out] 1/24*(48*c^(6*b)*e^(6*b*log(x^n) + 6*a) + 108*c^(4*b)*e^(4*b*log(x^n) + 4*a) + 88*c^(2*b)*e^(2*b*log(x^n) + 2*a) + 25)/(b*c^(8*b)*n*e^(8*b*log(x^n) + 8*a) + 4*b*c^(6*b)*n*e^(6*b*log(x^n) + 6*a) + 6*b*c^(4*b)*n*e^(4*b*log(x^n) + 4*a) + 4*b*c^(2*b)*n*e^(2*b*log(x^n) + 2*a) + b*n) - 1/24*(12*c^(6*b)*e^(6*b*log(x^n) + 6*a) + 42*c^(4*b)*e^(4*b*log(x^n) + 4*a) + 52*c^(2*b)*e^(2*b*log(x^n) + 2*a) + 25)/(b*c^(8*b)*n*e^(8*b*log(x^n) + 8*a) + 4*b*c^(6*b)*n*e^(6*b*log(x^n) + 6*a) + 6*b*c^(4*b)*n*e^(4*b*log(x^n) + 4*a) + 4*b*c^(2*b)`


```

*log(c) + a)^4 + 30*cosh(b*n*log(x) + b*log(c) + a)^2 + 3)*sinh(b*n*log(x)
+ b*log(c) + a)^4 + 6*cosh(b*n*log(x) + b*log(c) + a)^4 + 8*(7*cosh(b*n*log
(x) + b*log(c) + a)^5 + 10*cosh(b*n*log(x) + b*log(c) + a)^3 + 3*cosh(b*n*lo
g(x) + b*log(c) + a))*sinh(b*n*log(x) + b*log(c) + a)^3 + 4*(7*cosh(b*n*lo
g(x) + b*log(c) + a)^6 + 15*cosh(b*n*log(x) + b*log(c) + a)^4 + 9*cosh(b*n*
log(x) + b*log(c) + a)^2 + 1)*sinh(b*n*log(x) + b*log(c) + a)^2 + 4*cosh(b*
n*log(x) + b*log(c) + a)^2 + 8*(cosh(b*n*log(x) + b*log(c) + a)^7 + 3*cosh(
b*n*log(x) + b*log(c) + a)^5 + 3*cosh(b*n*log(x) + b*log(c) + a)^3 + cosh(b
*n*log(x) + b*log(c) + a))*sinh(b*n*log(x) + b*log(c) + a) + 1)*log(2*cosh(
b*n*log(x) + b*log(c) + a)/(cosh(b*n*log(x) + b*log(c) + a) - sinh(b*n*log(
x) + b*log(c) + a))) + 8*(b*n*cosh(b*n*log(x) + b*log(c) + a)^7*log(x) + 3*
(b*n*log(x) - 1)*cosh(b*n*log(x) + b*log(c) + a)^5 + (3*b*n*log(x) - 2)*cos
h(b*n*log(x) + b*log(c) + a)^3 + (b*n*log(x) - 1)*cosh(b*n*log(x) + b*log(c)
+ a))*sinh(b*n*log(x) + b*log(c) + a))/(b*n*cosh(b*n*log(x) + b*log(c) +
a)^8 + 8*b*n*cosh(b*n*log(x) + b*log(c) + a)*sinh(b*n*log(x) + b*log(c) + a
)^7 + b*n*sinh(b*n*log(x) + b*log(c) + a)^8 + 4*b*n*cosh(b*n*log(x) + b*log
(c) + a)^6 + 4*(7*b*n*cosh(b*n*log(x) + b*log(c) + a)^2 + b*n)*sinh(b*n*log
(x) + b*log(c) + a)^6 + 6*b*n*cosh(b*n*log(x) + b*log(c) + a)^4 + 8*(7*b*n*
cosh(b*n*log(x) + b*log(c) + a)^3 + 3*b*n*cosh(b*n*log(x) + b*log(c) + a))*
sinh(b*n*log(x) + b*log(c) + a)^5 + 2*(35*b*n*cosh(b*n*log(x) + b*log(c) +
a)^4 + 30*b*n*cosh(b*n*log(x) + b*log(c) + a)^2 + 3*b*n)*sinh(b*n*log(x) +
b*log(c) + a)^4 + 4*b*n*cosh(b*n*log(x) + b*log(c) + a)^2 + 8*(7*b*n*cosh(b
*n*log(x) + b*log(c) + a)^5 + 10*b*n*cosh(b*n*log(x) + b*log(c) + a)^3 + 3*
b*n*cosh(b*n*log(x) + b*log(c) + a))*sinh(b*n*log(x) + b*log(c) + a)^3 + 4*
(7*b*n*cosh(b*n*log(x) + b*log(c) + a)^6 + 15*b*n*cosh(b*n*log(x) + b*log(c)
+ a)^4 + 9*b*n*cosh(b*n*log(x) + b*log(c) + a)^2 + b*n)*sinh(b*n*log(x) +
b*log(c) + a)^2 + b*n + 8*(b*n*cosh(b*n*log(x) + b*log(c) + a)^7 + 3*b*n*c
osh(b*n*log(x) + b*log(c) + a)^5 + 3*b*n*cosh(b*n*log(x) + b*log(c) + a)^3
+ b*n*cosh(b*n*log(x) + b*log(c) + a))*sinh(b*n*log(x) + b*log(c) + a))

```

Sympy [A]

time = 2.91, size = 92, normalized size = 1.39

$$\begin{cases} \log(x) \tanh^5(a) & \text{for } b = 0 \wedge n = 0 \\ \log(x) \tanh^5(a + b \log(c)) & \text{for } n = 0 \\ \log(x) \tanh^5(a) & \text{for } b = 0 \\ \frac{\log(cx^n)}{n} - \frac{\log(\tanh(a + b \log(cx^n)) + 1)}{bn} - \frac{\tanh^4(a + b \log(cx^n))}{4bn} - \frac{\tanh^2(a + b \log(cx^n))}{2bn} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+b*ln(c*x**n))**5/x,x)

[Out] Piecewise((log(x)*tanh(a)**5, Eq(b, 0) & Eq(n, 0)), (log(x)*tanh(a + b*log(c))**5, Eq(n, 0)), (log(x)*tanh(a)**5, Eq(b, 0)), (log(c*x**n)/n - log(tanh(a + b*log(c*x**n)) + 1)/(b*n) - tanh(a + b*log(c*x**n))**4/(4*b*n) - tanh(a + b*log(c*x**n))**2/(2*b*n), True))

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 161 vs. 2(62) = 124.

time = 0.49, size = 161, normalized size = 2.44

$$\frac{\log\left(\sqrt{2x^{2bn}|c|^{2b}\cos(\pi b\operatorname{sgn}(c) - \pi b)e^{(2a)} + x^{4bn}|c|^{4b}e^{(4a)} + 1}\right)}{bn} - \frac{25c^8x^{8bn}e^{(8a)} + 52c^6x^{6bn}e^{(6a)} + 102c^4x^{4bn}e^{(4a)} + 52c^2x^{2bn}e^{(2a)} + 25}{12(c^{2b}x^{2bn}e^{(2a)} + 1)^4bn} - \log(x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+b*log(c*x^n))^5/x,x, algorithm="giac")

[Out] log(sqrt(2*x^(2*b*n)*abs(c)^(2*b)*cos(pi*b*sgn(c) - pi*b)*e^(2*a) + x^(4*b*n)*abs(c)^(4*b)*e^(4*a) + 1))/(b*n) - 1/12*(25*c^(8*b)*x^(8*b*n)*e^(8*a) + 52*c^(6*b)*x^(6*b*n)*e^(6*a) + 102*c^(4*b)*x^(4*b*n)*e^(4*a) + 52*c^(2*b)*x^(2*b*n)*e^(2*a) + 25)/((c^(2*b)*x^(2*b*n)*e^(2*a) + 1)^4*b*n) - log(x)

Mupad [B]

time = 1.06, size = 227, normalized size = 3.44

$$\frac{8}{bn + 3bn e^{2a} (cx^n)^{2b} + 3bn e^{4a} (cx^n)^{4b} + bn e^{6a} (cx^n)^{6b}} - \ln(x) + \frac{4}{bn + bn e^{2a} (cx^n)^{2b}} - \frac{4}{bn + 4bn e^{2a} (cx^n)^{2b} + 6bn e^{4a} (cx^n)^{4b} + 4bn e^{6a} (cx^n)^{6b} + bn e^{8a} (cx^n)^{8b}} - \frac{8}{bn + 2bn e^{2a} (cx^n)^{2b} + bn e^{4a} (cx^n)^{4b}} + \frac{\ln(e^{2a} (cx^n)^{2b} + 1)}{bn}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(a + b*log(c*x^n))^5/x,x)

[Out] 8/(b*n + 3*b*n*exp(2*a)*(c*x^n)^(2*b) + 3*b*n*exp(4*a)*(c*x^n)^(4*b) + b*n*exp(6*a)*(c*x^n)^(6*b)) - log(x) + 4/(b*n + b*n*exp(2*a)*(c*x^n)^(2*b)) - 4/(b*n + 4*b*n*exp(2*a)*(c*x^n)^(2*b) + 6*b*n*exp(4*a)*(c*x^n)^(4*b) + 4*b*n*exp(6*a)*(c*x^n)^(6*b) + b*n*exp(8*a)*(c*x^n)^(8*b)) - 8/(b*n + 2*b*n*exp(2*a)*(c*x^n)^(2*b) + b*n*exp(4*a)*(c*x^n)^(4*b)) + log(exp(2*a)*(c*x^n)^(2*b) + 1)/(b*n)

3.189 $\int (ex)^m \tanh(d(a + b \log(cx^n))) dx$

Optimal. Leaf size=88

$$\frac{(ex)^{1+m}}{e(1+m)} - \frac{2(ex)^{1+m} {}_2F_1\left(1, \frac{1+m}{2bdn}; 1 + \frac{1+m}{2bdn}, -e^{2ad}(cx^n)^{2bd}\right)}{e(1+m)}$$

[Out] $(e*x)^{(1+m)}/e/(1+m)-2*(e*x)^{(1+m)}*\text{hypergeom}([1, 1/2*(1+m)/b/d/n], [1+1/2*(1+m)/b/d/n], -\exp(2*a*d)*(c*x^n)^{(2*b*d)})/e/(1+m)$

Rubi [A]

time = 0.05, antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.210$, Rules used = {5658, 5656, 470, 371}

$$\frac{(ex)^{m+1}}{e(m+1)} - \frac{2(ex)^{m+1} {}_2F_1\left(1, \frac{m+1}{2bdn}, \frac{m+1}{2bdn} + 1; -e^{2ad}(cx^n)^{2bd}\right)}{e(m+1)}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(e*x)^m*\text{Tanh}[d*(a + b*\text{Log}[c*x^n])], x]$

[Out] $(e*x)^{(1+m)}/(e*(1+m)) - (2*(e*x)^{(1+m)}*\text{Hypergeometric2F1}[1, (1+m)/(2*b*d*n), 1 + (1+m)/(2*b*d*n), -(E^{2*a*d})*(c*x^n)^{(2*b*d)}])/e*(1+m)$

Rule 371

$\text{Int}[(c*x)^m*(a + b*(x^n)^p), x_Symbol] :> \text{Simp}[a^p*(c*x)^{m+1}/(c*(m+1))*\text{Hypergeometric2F1}[-p, (m+1)/n, (m+1)/n + 1, (-b)*(x^n/a)], x] /; \text{FreeQ}\{a, b, c, m, n, p\}, x \ \&\& \ !\text{IGtQ}[p, 0] \ \&\& \ (\text{ILtQ}[p, 0] \ || \ \text{GtQ}[a, 0])$

Rule 470

$\text{Int}[(e*x)^m*(a + b*(x^n)^p), x_Symbol] :> \text{Simp}[d*(e*x)^{m+1}*((a + b*x^n)^{p+1}/(b*e*(m+n*(p+1)+1))), x] - \text{Dist}[(a*d*(m+1) - b*c*(m+n*(p+1)+1))/(b*(m+n*(p+1)+1)), \text{Int}[(e*x)^m*(a + b*x^n)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, m, n, p\}, x \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[m + n*(p+1) + 1, 0]$

Rule 5656

$\text{Int}[(e*x)^m*\text{Tanh}[(a + \text{Log}[x]*b)*d]^p], x_Symbol] :> \text{Int}[(e*x)^m*(-1 + E^{2*a*d}*x^{2*b*d})^p/(1 + E^{2*a*d}*x^{2*b*d})^p], x] /; \text{FreeQ}\{a, b, d, e, m, p\}, x]$

Rule 5658

Int[((e_.)*(x_.))^(m_.)*Tanh[((a_.) + Log[(c_.)*(x_.)^(n_.)]*(b_.))*(d_.)]^(p_.), x_Symbol] :> Dist[(e*x)^(m + 1)/(e*n*(c*x^n)^((m + 1)/n)), Subst[Int[x^((m + 1)/n - 1)*Tanh[d*(a + b*Log[x])]^p, x], x, c*x^n], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && (NeQ[c, 1] || NeQ[n, 1])

Rubi steps

$$\int (ex)^m \tanh(d(a + b \log(cx^n))) dx = \int (ex)^m \tanh(d(a + b \log(cx^n))) dx$$

Mathematica [A]

time = 9.63, size = 160, normalized size = 1.82

$$\frac{x(ex)^m \left(-{}_2F_1\left(1, \frac{1+m}{2bdn}; 1 + \frac{1+m}{2bdn}; -e^{2d(a+b \log(cx^n))}\right) + \frac{e^{2ad(1+m)(cx^n)^{2bd}} {}_2F_1\left(1, \frac{1+m+2bdn}{2bdn}; \frac{1+m+4bdn}{2bdn}; -e^{2ad}(cx^n)^{2bd}\right)}{1+m+2bdn} \right)}{1+m}$$

Antiderivative was successfully verified.

[In] Integrate[(e*x)^m*Tanh[d*(a + b*Log[c*x^n])],x]

[Out] (x*(e*x)^m*(-Hypergeometric2F1[1, (1 + m)/(2*b*d*n), 1 + (1 + m)/(2*b*d*n), -E^(2*d*(a + b*Log[c*x^n])]) + (E^(2*a*d)*(1 + m)*(c*x^n)^(2*b*d)*Hypergeometric2F1[1, (1 + m + 2*b*d*n)/(2*b*d*n), (1 + m + 4*b*d*n)/(2*b*d*n), -(E^(2*a*d)*(c*x^n)^(2*b*d))])/(1 + m + 2*b*d*n)))/(1 + m)

Maple [F]

time = 0.50, size = 0, normalized size = 0.00

$$\int (ex)^m \tanh(d(a + b \ln(cx^n))) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((e*x)^m*tanh(d*(a+b*ln(c*x^n))),x)

[Out] int((e*x)^m*tanh(d*(a+b*ln(c*x^n))),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e*x)^m*tanh(d*(a+b*log(c*x^n))),x, algorithm="maxima")

[Out] $x e^{(m \log(x) + m)/(m + 1)} - 2 \int \frac{e^{(m \log(x) + m)}}{c^{(2*b*d)} e^{(2*b*d \log(x^n) + 2*a*d) + 1}} dx$

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e*x)^m*tanh(d*(a+b*log(c*x^n))),x, algorithm="fricas")

[Out] integral((x*e)^m*tanh(b*d*log(c*x^n) + a*d), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (ex)^m \tanh(ad + bd \log(cx^n)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e*x)**m*tanh(d*(a+b*ln(c*x**n))),x)

[Out] Integral((e*x)**m*tanh(a*d + b*d*log(c*x**n)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e*x)^m*tanh(d*(a+b*log(c*x^n))),x, algorithm="giac")

[Out] integrate((e*x)^m*tanh((b*log(c*x^n) + a)*d), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \tanh(d(a + b \ln(cx^n))) (ex)^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(d*(a + b*log(c*x^n)))*(e*x)^m,x)

[Out] int(tanh(d*(a + b*log(c*x^n)))*(e*x)^m, x)

3.190 $\int (ex)^m \tanh^2(d(a + b \log(cx^n))) dx$

Optimal. Leaf size=169

$$\frac{(1+m+bdn)(ex)^{1+m}}{bde(1+m)n} + \frac{(ex)^{1+m} (1 - e^{2ad}(cx^n)^{2bd})}{bden (1 + e^{2ad}(cx^n)^{2bd})} - \frac{2(ex)^{1+m} {}_2F_1\left(1, \frac{1+m}{2bdn}; 1 + \frac{1+m}{2bdn}; -e^{2ad}(cx^n)^{2bd}\right)}{bden}$$

[Out] (b*d*n+m+1)*(e*x)^(1+m)/b/d/e/(1+m)/n+(e*x)^(1+m)*(1-exp(2*a*d)*(c*x^n)^(2*b*d))/b/d/e/n/(1+exp(2*a*d)*(c*x^n)^(2*b*d))-2*(e*x)^(1+m)*hypergeom([1, 1/2*(1+m)/b/d/n], [1+1/2*(1+m)/b/d/n], -exp(2*a*d)*(c*x^n)^(2*b*d))/b/d/e/n

Rubi [A]

time = 0.15, antiderivative size = 169, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {5658, 5656, 516, 470, 371}

$$-\frac{2(ex)^{m+1} {}_2F_1\left(1, \frac{m+1}{2bdn}, \frac{m+1}{2bdn} + 1; -e^{2ad}(cx^n)^{2bd}\right)}{bden} + \frac{(ex)^{m+1} (1 - e^{2ad}(cx^n)^{2bd})}{bden (e^{2ad}(cx^n)^{2bd} + 1)} + \frac{(ex)^{m+1}(bdn + m + 1)}{bde(m + 1)n}$$

Antiderivative was successfully verified.

[In] Int[(e*x)^m*Tanh[d*(a + b*Log[c*x^n])]^2,x]

[Out] ((1 + m + b*d*n)*(e*x)^(1 + m))/(b*d*e*(1 + m)*n) + ((e*x)^(1 + m)*(1 - E^(2*a*d)*(c*x^n)^(2*b*d)))/(b*d*e*n*(1 + E^(2*a*d)*(c*x^n)^(2*b*d))) - (2*(e*x)^(1 + m)*Hypergeometric2F1[1, (1 + m)/(2*b*d*n), 1 + (1 + m)/(2*b*d*n), -E^(2*a*d)*(c*x^n)^(2*b*d)])/(b*d*e*n)

Rule 371

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])

Rule 470

Int[((e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> Simp[d*(e*x)^(m + 1)*((a + b*x^n)^(p + 1)/(b*e*(m + n*(p + 1) + 1))), x] - Dist[(a*d*(m + 1) - b*c*(m + n*(p + 1) + 1))/(b*(m + n*(p + 1) + 1)), Int[(e*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && NeQ[b*c - a*d, 0] && NeQ[m + n*(p + 1) + 1, 0]

Rule 516


```
Int[((e._)*(x_))^(m._)*((a_) + (b._)*(x_)^(n_))^(p_)*((c_) + (d._)*(x_)^(n_))^(q_), x_Symbol] := Simp[(-c*b - a*d)*(e*x)^(m + 1)*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q - 1)/(a*b*e*n*(p + 1))), x] + Dist[1/(a*b*n*(p + 1)), Int[(e*x)^m*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 2)*Simp[c*(c*b*n*(p + 1) + (c*b - a*d)*(m + 1)) + d*(c*b*n*(p + 1) + (c*b - a*d)*(m + n*(q - 1) + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && GtQ[q, 1] && IntBinomialQ[a, b, c, d, e, m, n, p, q, x]
```

Rule 5656

```
Int[((e._)*(x_))^(m._)*Tanh[((a._) + Log[x_]*(b._))*(d._)]^(p_), x_Symbol] := Int[(e*x)^m*((-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p), x] /; FreeQ[{a, b, d, e, m, p}, x]
```

Rule 5658

```
Int[((e._)*(x_))^(m._)*Tanh[((a._) + Log[(c._)*(x_)^(n_)]*(b._))*(d._)]^(p_), x_Symbol] := Dist[(e*x)^(m + 1)/(e*n*(c*x^n)^((m + 1)/n)), Subst[Int[x^((m + 1)/n - 1)*Tanh[d*(a + b*Log[x])]^p, x], x, c*x^n], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && (NeQ[c, 1] || NeQ[n, 1])
```

Rubi steps

$$\int (ex)^m \tanh^2(d(a + b \log(cx^n))) dx = \int (ex)^m \tanh^2(d(a + b \log(cx^n))) dx$$

Mathematica [A]

time = 12.98, size = 317, normalized size = 1.88

$$(ex)^m \left(\frac{x}{1+m} - \frac{e^{\frac{(1+2m)(a-b \log(cx^n))}{2dn}} x^{-2m} \left(e^{\frac{(1+2m)(a+b \log(cx^n))}{2dn}} (1+m+2bdn) {}_2F_1\left(1, \frac{1+m}{2dn}; 1 + \frac{1+m}{2dn}; -e^{2d(a+b \log(cx^n))}\right) - e^{\frac{(1+2m+2bdn)(a-b \log(cx^n))}{2dn}} (1+m) x^{1+2m+2bdn} {}_2F_1\left(1, \frac{1+m+2bdn}{2dn}; \frac{1+m+4bdn}{2dn}; -e^{2d(a+b \log(cx^n))}\right) + e^{\frac{(1+2m)(a+b \log(cx^n))}{2dn}} (1+m+2bdn) \tanh(d(a+b \log(cx^n))) \right)}{bdn(1+m+2bdn)} \right)$$

Antiderivative was successfully verified.

```
[In] Integrate[(e*x)^m*Tanh[d*(a + b*Log[c*x^n])]^2,x]
```

```
[Out] (e*x)^m*(x/(1 + m) - (E^(((1 + 2*m)*(a + b*Log[c*x^n])))/(b*n))*(1 + m + 2*b*d*n)*Hypergeometric2F1[1, (1 + m)/(2*b*d*n), 1 + (1 + m)/(2*b*d*n), -E^(2*d*(a + b*Log[c*x^n]))] - E^(((1 + 2*m + 2*b*d*n)*(a - b*n*Log[x] + b*Log[c*x^n])))/(b*n)*(1 + m)*x^(1 + 2*m + 2*b*d*n)*Hypergeometric2F1[1, (1 + m + 2*b*d*n)/(2*b*d*n), (1 + m + 4*b*d*n)/(2*b*d*n), -E^(2*d*(a + b*Log[c*x^n]))] + E^(((1 + 2*m)*(a + b*Log[c*x^n])))/(b*n))*(1 + m + 2*b*d*n)*Tanh[d*(a + b*Log[c*x^n])]/(b*d*E^(((1 + 2*m)*(a - b*n*Log[x] + b*Log[c*x^n])))/(b*n))*n*(1 + m + 2*b*d*n)*x^(2*m))
```

Maple [F]

time = 0.49, size = 0, normalized size = 0.00

$$\int (ex)^m (\tanh^2(d(a + b \ln(cx^n)))) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((e*x)^m*tanh(d*(a+b*ln(c*x^n)))^2,x)

[Out] int((e*x)^m*tanh(d*(a+b*ln(c*x^n)))^2,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e*x)^m*tanh(d*(a+b*log(c*x^n)))^2,x, algorithm="maxima")

[Out] $-2*(m*e^m + e^m)*\text{integrate}(x^m/(b*c^{(2*b*d)*d*n}*e^{(2*b*d*\log(x^n) + 2*a*d) + b*d*n}), x) + (b*c^{(2*b*d)*d*n}*x*e^{(2*b*d*\log(x^n) + 2*a*d + m*\log(x) + m)} + (b*d*n*e^m + 2*m*e^m + 2*e^m)*x*x^m)/((m*n + n)*b*c^{(2*b*d)*d}*e^{(2*b*d*\log(x^n) + 2*a*d) + (m*n + n)*b*d})$

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e*x)^m*tanh(d*(a+b*log(c*x^n)))^2,x, algorithm="fricas")

[Out] integral((x*e)^m*tanh(b*d*log(c*x^n) + a*d)^2, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (ex)^m \tanh^2(ad + bd \log(cx^n)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e*x)**m*tanh(d*(a+b*ln(c*x**n))))**2,x)

[Out] Integral((e*x)**m*tanh(a*d + b*d*log(c*x**n))**2, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((e*x)^m*tanh(d*(a+b*log(c*x^n)))^2,x, algorithm="giac")
```

```
[Out] integrate((e*x)^m*tanh((b*log(c*x^n) + a)*d)^2, x)
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \tanh(d(a + b \ln(cx^n)))^2 (ex)^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(tanh(d*(a + b*log(c*x^n)))^2*(e*x)^m,x)
```

```
[Out] int(tanh(d*(a + b*log(c*x^n)))^2*(e*x)^m, x)
```

3.191 $\int (ex)^m \tanh^3(d(a + b \log(cx^n))) dx$

Optimal. Leaf size=307

$$\frac{(1+m+bdn)(1+m+2bdn)(ex)^{1+m}}{2b^2d^2e(1+m)n^2} - \frac{(ex)^{1+m} \left(1 - e^{2ad}(cx^n)^{2bd}\right)^2}{2bden \left(1 + e^{2ad}(cx^n)^{2bd}\right)^2} + \frac{e^{-2ad}(ex)^{1+m} \left(\frac{e^{2ad}(1+m-2bdn)}{n} - \frac{e^{4ad}(1+m)}{n}\right)}{2b^2d^2en \left(1 + e^{2ad}(cx^n)^{2bd}\right)^2}$$

[Out] $1/2*(b*d*n+m+1)*(2*b*d*n+m+1)*(e*x)^{(1+m)}/b^2/d^2/e/(1+m)/n^2-1/2*(e*x)^{(1+m)}*(1-\exp(2*a*d)*(c*x^n)^{(2*b*d)})^2/b/d/e/n/(1+\exp(2*a*d)*(c*x^n)^{(2*b*d)})^{2+1/2*(e*x)^{(1+m)}*(\exp(2*a*d)*(-2*b*d*n+m+1)/n-\exp(4*a*d)*(2*b*d*n+m+1)*(c*x^n)^{(2*b*d)}/n)/b^2/d^2/e/\exp(2*a*d)/n/(1+\exp(2*a*d)*(c*x^n)^{(2*b*d)})-(2*b^2*d^2*n^2+m^2+2*m+1)*(e*x)^{(1+m)}*\text{hypergeom}([1, 1/2*(1+m)/b/d/n], [1+1/2*(1+m)/b/d/n], -\exp(2*a*d)*(c*x^n)^{(2*b*d)})/b^2/d^2/e/(1+m)/n^2$

Rubi [A]

time = 0.32, antiderivative size = 307, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {5658, 5656, 516, 608, 470, 371}

$$\frac{(ex)^{m+1} (2b^2d^2n^2 + m^2 + 2m + 1) {}_2F_1\left(1, \frac{m+1}{2bdn}, \frac{m+1}{2bdn} + 1; -e^{2ad}(cx^n)^{2bd}\right)}{b^2d^2e(m+1)n^2} + \frac{e^{-2ad}(ex)^{m+1} \left(\frac{e^{2ad}(-2bdn+m+1)}{n} - \frac{e^{4ad}(2bdn+m+1)(cx^n)^{2bd}}{n}\right)}{2b^2d^2en \left(e^{2ad}(cx^n)^{2bd} + 1\right)} - \frac{(ex)^{m+1} \left(1 - e^{2ad}(cx^n)^{2bd}\right)^2}{2bden \left(e^{2ad}(cx^n)^{2bd} + 1\right)^2} + \frac{(ex)^{m+1} (bdn + m + 1)(2bdn + m + 1)}{2b^2d^2e(m+1)n^2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(e*x)^m * \text{Tanh}[d*(a + b*\text{Log}[c*x^n])]]^3, x$

[Out] $((1+m+b*d*n)*(1+m+2*b*d*n)*(e*x)^{(1+m)})/(2*b^2*d^2*e*(1+m)*n^2) - ((e*x)^{(1+m)}*(1-E^{(2*a*d)*(c*x^n)^{(2*b*d)}})^2)/(2*b*d*e*n*(1+E^{(2*a*d)*(c*x^n)^{(2*b*d)}})^2) + ((e*x)^{(1+m)}*((E^{(2*a*d)*(1+m-2*b*d*n)})/n - (E^{(4*a*d)*(1+m+2*b*d*n)*(c*x^n)^{(2*b*d)}})/n))/(2*b^2*d^2*e*E^{(2*a*d)*n*(1+E^{(2*a*d)*(c*x^n)^{(2*b*d)}})}) - ((1+2*m+m^2+2*b^2*d^2*n^2)*(e*x)^{(1+m)}*\text{Hypergeometric2F1}[1, (1+m)/(2*b*d*n), 1+(1+m)/(2*b*d*n), -(E^{(2*a*d)*(c*x^n)^{(2*b*d)}})]/(b^2*d^2*e*(1+m)*n^2))$

Rule 371

$\text{Int}[(c_*)(x_)^{(m_*)}*((a_)+(b_*)(x_)^{(n_)})^{(p_)}, x_Symbol] \rightarrow \text{Simp}[a^p * ((c*x)^{(m+1)}/(c*(m+1)))*\text{Hypergeometric2F1}[-p, (m+1)/n, (m+1)/n+1, (-b)*(x^n/a)], x] /; \text{FreeQ}\{a, b, c, m, n, p\}, x] \&\& !\text{IGtQ}[p, 0] \&\& (\text{ILtQ}[p, 0] || \text{GtQ}[a, 0])$

Rule 470

$\text{Int}[(e_*)(x_)^{(m_*)}*((a_)+(b_*)(x_)^{(n_)})^{(p_*)}*((c_)+(d_*)(x_)^{(n_)})^{(p_*)}, x_Symbol] \rightarrow \text{Simp}[d*(e*x)^{(m+1)}*((a+b*x^n)^{(p+1)}/(b*e*(m+n*(p+1)+1))), x] - \text{Dist}[(a*d*(m+1) - b*c*(m+n*(p+1)+1))/(b*(m+n*(p+1)+1)), x]$

+ 1) + 1)), Int[(e*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && NeQ[b*c - a*d, 0] && NeQ[m + n*(p + 1) + 1, 0]

Rule 516

Int[((e_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol] := Simp[(-c*b - a*d)*(e*x)^(m + 1)*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q - 1)/(a*b*e*n*(p + 1))), x] + Dist[1/(a*b*n*(p + 1)), Int[(e*x)^m*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 2)*Simp[c*(c*b*n*(p + 1) + (c*b - a*d)*(m + 1)) + d*(c*b*n*(p + 1) + (c*b - a*d)*(m + n*(q - 1) + 1))*x^n, x], x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && GtQ[q, 1] && IntBinomialQ[a, b, c, d, e, m, n, p, q, x]

Rule 608

Int[((g_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_)*((e_) + (f_)*(x_)^(n_)), x_Symbol] := Simp[(-b*e - a*f)*(g*x)^(m + 1)*(a + b*x^n)^(p + 1)*((c + d*x^n)^q/(a*b*g*n*(p + 1))), x] + Dist[1/(a*b*n*(p + 1)), Int[(g*x)^m*(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1)*Simp[c*(b*e*n*(p + 1) + (b*e - a*f)*(m + 1)) + d*(b*e*n*(p + 1) + (b*e - a*f)*(m + n*q + 1))*x^n, x], x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n}, x] && LtQ[p, -1] && GtQ[q, 0] && !(EqQ[q, 1] && SimplerQ[b*c - a*d, b*e - a*f])

Rule 5656

Int[((e_)*(x_))^(m_)*Tanh[((a_) + Log[x]*(b_))*(d_)]^(p_), x_Symbol] := Int[(e*x)^m*((-1 + E^(2*a*d)*x^(2*b*d))^p/(1 + E^(2*a*d)*x^(2*b*d))^p), x] /; FreeQ[{a, b, d, e, m, p}, x]

Rule 5658

Int[((e_)*(x_))^(m_)*Tanh[((a_) + Log[(c_)*(x_)^(n_)]*(b_))*(d_)]^(p_), x_Symbol] := Dist[(e*x)^(m + 1)/(e*n*(c*x^n)^((m + 1)/n)), Subst[Int[x^((m + 1)/n - 1)*Tanh[d*(a + b*Log[x])]^p, x], x, c*x^n], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && (NeQ[c, 1] || NeQ[n, 1])

Rubi steps

$$\int (ex)^m \tanh^3(d(a + b \log(cx^n))) dx = \int (ex)^m \tanh^3(d(a + b \log(cx^n))) dx$$

Mathematica [A]

time = 13.71, size = 606, normalized size = 1.97

Antiderivative was successfully verified.

```
[In] Integrate[(e*x)^m*Tanh[d*(a + b*Log[c*x^n])]^3,x]
```

```
[Out] (x*(e*x)^m*Sech[b*d*n*Log[x] + d*(a + b*(-(n*Log[x]) + Log[c*x^n]))]^2)/(2*
b*d*n) - ((1 + m)*x*(e*x)^m*Sech[d*(a + b*(-(n*Log[x]) + Log[c*x^n]))]*Sech
[b*d*n*Log[x] + d*(a + b*(-(n*Log[x]) + Log[c*x^n]))]*Sinh[b*d*n*Log[x]])/(
2*b^2*d^2*n^2) + ((1 + 2*m + m^2 + 2*b^2*d^2*n^2)*(e*x)^m*Sech[d*(a + b*(-(
n*Log[x]) + Log[c*x^n]))]*((x^(1 + m)*Sech[d*(a + b*Log[c*x^n]])*Sinh[b*d*n
*Log[x]])/(1 + m) - (Cosh[d*(a + b*(-(n*Log[x]) + Log[c*x^n]))]*E^((a + 2*
a*m + b*(1 + m)*n*Log[x] + b*(1 + 2*m)*(-(n*Log[x]) + Log[c*x^n]))/(b*n))*
(1 + m + 2*b*d*n)*Hypergeometric2F1[1, (1 + m)/(2*b*d*n), 1 + (1 + m)/(2*b*d
*n), -E^(2*d*(a + b*Log[c*x^n]))] - E^((a*(1 + 2*m + 2*b*d*n))/(b*n) + (1 +
m + 2*b*d*n)*Log[x] + ((1 + 2*m + 2*b*d*n)*(-(n*Log[x]) + Log[c*x^n]))/n)*
(1 + m)*Hypergeometric2F1[1, (1 + m + 2*b*d*n)/(2*b*d*n), (1 + m + 4*b*d*n)
/(2*b*d*n), -E^(2*d*(a + b*Log[c*x^n]))] + E^((a + 2*a*m + b*(1 + m)*n*Log[
x] + b*(1 + 2*m)*(-(n*Log[x]) + Log[c*x^n]))/(b*n))*(1 + m + 2*b*d*n)*Tanh[
d*(a + b*Log[c*x^n])])/(E^(((1 + 2*m)*(a + b*(-(n*Log[x]) + Log[c*x^n])))))/
(b*n))*(1 + m)*(1 + m + 2*b*d*n)))/(2*b^2*d^2*n^2*x^m) + (x*(e*x)^m*Tanh[d
*(a + b*(-(n*Log[x]) + Log[c*x^n]))])/(1 + m)
```

Maple [F]

time = 0.61, size = 0, normalized size = 0.00

$$\int (ex)^m (\tanh^3(d(a + b \ln(cx^n)))) dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((e*x)^m*tanh(d*(a+b*ln(c*x^n)))^3,x)
```

```
[Out] int((e*x)^m*tanh(d*(a+b*ln(c*x^n)))^3,x)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((e*x)^m*tanh(d*(a+b*log(c*x^n)))^3,x, algorithm="maxima")
```

```
[Out] -(2*b^2*d^2*n^2*e^m + m^2*e^m + 2*m*e^m + e^m)*integrate(x^m/(b^2*c^(2*b*d)
*d^2*n^2*e^(2*b*d*log(x^n) + 2*a*d) + b^2*d^2*n^2), x) + (b^2*c^(4*b*d)*d^2
*n^2*x*e^(4*b*d*log(x^n) + 4*a*d + m*log(x) + m) + (b^2*d^2*n^2*e^m + m^2*e
^m + 2*m*e^m + e^m)*x*x^m + (2*b^2*c^(2*b*d)*d^2*n^2*e^(2*a*d + m) + 2*(m*n
*e^m + n*e^m)*b*c^(2*b*d)*d*e^(2*a*d) + (m^2*e^m + 2*m*e^m + e^m)*c^(2*b*d)
*e^(2*a*d))*x*e^(2*b*d*log(x^n) + m*log(x))/((m*n^2 + n^2)*b^2*c^(4*b*d)*d
```

$$^2e^{(4*b*d*\log(x^n) + 4*a*d) + 2*(m*n^2 + n^2)*b^2*c^{(2*b*d)*d^2}*e^{(2*b*d*\log(x^n) + 2*a*d) + (m*n^2 + n^2)*b^2*d^2)}$$

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e*x)^m*tanh(d*(a+b*log(c*x^n)))^3,x, algorithm="fricas")

[Out] integral((x*e)^m*tanh(b*d*log(c*x^n) + a*d)^3, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (ex)^m \tanh^3(ad + bd \log(cx^n)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e*x)**m*tanh(d*(a+b*ln(c*x**n))))**3,x)

[Out] Integral((e*x)**m*tanh(a*d + b*d*log(c*x**n))**3, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e*x)^m*tanh(d*(a+b*log(c*x^n)))^3,x, algorithm="giac")

[Out] integrate((e*x)^m*tanh((b*log(c*x^n) + a)*d)^3, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \tanh(d(a + b \ln(cx^n)))^3 (ex)^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(d*(a + b*log(c*x^n)))^3*(e*x)^m,x)

[Out] int(tanh(d*(a + b*log(c*x^n)))^3*(e*x)^m, x)

3.192 $\int \tanh^p(d(a + b \log(cx^n))) dx$

Optimal. Leaf size=115

$$x \left(1 - e^{2ad}(cx^n)^{2bd}\right)^{-p} \left(-1 + e^{2ad}(cx^n)^{2bd}\right)^p F_1\left(\frac{1}{2bdn}; -p, p; 1 + \frac{1}{2bdn}; e^{2ad}(cx^n)^{2bd}, -e^{2ad}(cx^n)^{2bd}\right)$$

[Out] x*(-1+exp(2*a*d)*(c*x^n)^(2*b*d))^p*AppellF1(1/2/b/d/n,-p,p,1+1/2/b/d/n,exp(2*a*d)*(c*x^n)^(2*b*d),-exp(2*a*d)*(c*x^n)^(2*b*d))/((1-exp(2*a*d)*(c*x^n)^(2*b*d))^p)

Rubi [A]

time = 0.08, antiderivative size = 115, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {5654, 5656, 525, 524}

$$x \left(1 - e^{2ad}(cx^n)^{2bd}\right)^{-p} \left(e^{2ad}(cx^n)^{2bd} - 1\right)^p F_1\left(\frac{1}{2bdn}; -p, p; 1 + \frac{1}{2bdn}; e^{2ad}(cx^n)^{2bd}, -e^{2ad}(cx^n)^{2bd}\right)$$

Antiderivative was successfully verified.

[In] Int[Tanh[d*(a + b*Log[c*x^n])]^p,x]

[Out] (x*(-1 + E^(2*a*d)*(c*x^n)^(2*b*d))^p*AppellF1[1/(2*b*d*n), -p, p, 1 + 1/(2*b*d*n), E^(2*a*d)*(c*x^n)^(2*b*d), -(E^(2*a*d)*(c*x^n)^(2*b*d))]/(1 - E^(2*a*d)*(c*x^n)^(2*b*d))^p)

Rule 524

Int[((e._)*(x._))^(m._)*((a._) + (b._)*(x._)^(n._))^(p._)*((c._) + (d._)*(x._)^(n._))^(q._), x_Symbol] :> Simp[a^p*c^q*((e*x)^(m + 1)/(e*(m + 1)))*AppellF1[(m + 1)/n, -p, -q, 1 + (m + 1)/n, (-b)*(x^n/a), (-d)*(x^n/c)], x] /; FreeQ[{a, b, c, d, e, m, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[m, -1] && NeQ[m, n - 1] && (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])

Rule 525

Int[((e._)*(x._))^(m._)*((a._) + (b._)*(x._)^(n._))^(p._)*((c._) + (d._)*(x._)^(n._))^(q._), x_Symbol] :> Dist[a^IntPart[p]*((a + b*x^n)^FracPart[p]/(1 + b*(x^n/a)^FracPart[p])), Int[(e*x)^m*(1 + b*(x^n/a))^p*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, e, m, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[m, -1] && NeQ[m, n - 1] && !(IntegerQ[p] || GtQ[a, 0])

Rule 5654

Int[Tanh[((a._) + Log[(c._)*(x._)^(n._)]*(b._))*(d._)]^(p._), x_Symbol] :> Dist[x/(n*(c*x^n)^(1/n)), Subst[Int[x^(1/n - 1)*Tanh[d*(a + b*Log[x])]^p, x]

, x, c*x^n], x] /; FreeQ[{a, b, c, d, n, p}, x] && (NeQ[c, 1] || NeQ[n, 1])

Rule 5656

Int[((e_.)*(x_))^(m_.)*Tanh[((a_.) + Log[x_]*(b_.)*(d_.)]^(p_.), x_Symbol]
 :> Int[(e*x)^(m*((-1 + E^(2*a*d)*x^(2*b*d)))^p/(1 + E^(2*a*d)*x^(2*b*d))^p),
 x] /; FreeQ[{a, b, d, e, m, p}, x]

Rubi steps

$$\int \tanh^p(d(a + b \log(cx^n))) dx = \int \tanh^p(d(a + b \log(cx^n))) dx$$

Mathematica [B] Leaf count is larger than twice the leaf count of optimal. 387 vs. 2(115) = 230.

time = 2.65, size = 387, normalized size = 3.37

$$\frac{(1 + 2bdn)x \left(\frac{-1 + e^{2ad}(cx^n)^{2bd}}{1 + e^{2ad}(cx^n)^{2bd}} \right)^p F_1 \left(\frac{1}{2bdn}; -p, p; 1 + \frac{1}{2bdn}; e^{2ad}(cx^n)^{2bd}, -e^{2ad}(cx^n)^{2bd} \right) - 2bde^{2ad}np (cx^n)^{2bd} F_1 \left(1 + \frac{1}{2bdn}; 1 - p, p; 2 + \frac{1}{2bdn}; e^{2ad}(cx^n)^{2bd}, -e^{2ad}(cx^n)^{2bd} \right) - 2bde^{2ad}np (cx^n)^{2bd} F_1 \left(1 + \frac{1}{2bdn}; -p, 1 + p; 2 + \frac{1}{2bdn}; e^{2ad}(cx^n)^{2bd}, -e^{2ad}(cx^n)^{2bd} \right) + (1 + 2bdn) F_1 \left(\frac{1}{2bdn}; -p, p; 1 + \frac{1}{2bdn}; e^{2ad}(cx^n)^{2bd}, -e^{2ad}(cx^n)^{2bd} \right)}{(1 + 2bdn)x \left(\frac{-1 + e^{2ad}(cx^n)^{2bd}}{1 + e^{2ad}(cx^n)^{2bd}} \right)^p F_1 \left(\frac{1}{2bdn}; -p, p; 1 + \frac{1}{2bdn}; e^{2ad}(cx^n)^{2bd}, -e^{2ad}(cx^n)^{2bd} \right) - 2bde^{2ad}np (cx^n)^{2bd} F_1 \left(1 + \frac{1}{2bdn}; 1 - p, p; 2 + \frac{1}{2bdn}; e^{2ad}(cx^n)^{2bd}, -e^{2ad}(cx^n)^{2bd} \right) - 2bde^{2ad}np (cx^n)^{2bd} F_1 \left(1 + \frac{1}{2bdn}; -p, 1 + p; 2 + \frac{1}{2bdn}; e^{2ad}(cx^n)^{2bd}, -e^{2ad}(cx^n)^{2bd} \right) + (1 + 2bdn) F_1 \left(\frac{1}{2bdn}; -p, p; 1 + \frac{1}{2bdn}; e^{2ad}(cx^n)^{2bd}, -e^{2ad}(cx^n)^{2bd} \right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[Tanh[d*(a + b*Log[c*x^n])]^p,x]

[Out] (((1 + 2*b*d*n)*x*((-1 + E^(2*a*d)*(c*x^n)^(2*b*d))/(1 + E^(2*a*d)*(c*x^n)^(2*b*d)))^p*AppellF1[1/(2*b*d*n), -p, p, 1 + 1/(2*b*d*n), E^(2*a*d)*(c*x^n)^(2*b*d), -(E^(2*a*d)*(c*x^n)^(2*b*d))]/(-2*b*d*E^(2*a*d)*n*p*(c*x^n)^(2*b*d)*AppellF1[1 + 1/(2*b*d*n), 1 - p, p, 2 + 1/(2*b*d*n), E^(2*a*d)*(c*x^n)^(2*b*d), -(E^(2*a*d)*(c*x^n)^(2*b*d))] - 2*b*d*E^(2*a*d)*n*p*(c*x^n)^(2*b*d)*AppellF1[1 + 1/(2*b*d*n), -p, 1 + p, 2 + 1/(2*b*d*n), E^(2*a*d)*(c*x^n)^(2*b*d), -(E^(2*a*d)*(c*x^n)^(2*b*d))] + (1 + 2*b*d*n)*AppellF1[1/(2*b*d*n), -p, p, 1 + 1/(2*b*d*n), E^(2*a*d)*(c*x^n)^(2*b*d), -(E^(2*a*d)*(c*x^n)^(2*b*d))])

Maple [F]

time = 1.31, size = 0, normalized size = 0.00

$$\int \tanh^p(d(a + b \ln(cx^n))) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(d*(a+b*ln(c*x^n)))^p,x)

[Out] int(tanh(d*(a+b*ln(c*x^n)))^p,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(d*(a+b*log(c*x^n)))^p,x, algorithm="maxima")

[Out] integrate(tanh((b*log(c*x^n) + a)*d)^p, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(d*(a+b*log(c*x^n)))^p,x, algorithm="fricas")

[Out] integral(tanh(b*d*log(c*x^n) + a*d)^p, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \tanh^p(d(a + b \log(cx^n))) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(d*(a+b*ln(c*x**n)))**p,x)

[Out] Integral(tanh(d*(a + b*log(c*x**n)))**p, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(d*(a+b*log(c*x^n)))^p,x, algorithm="giac")

[Out] integrate(tanh((b*log(c*x^n) + a)*d)^p, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \tanh(d(a + b \ln(cx^n)))^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(d*(a + b*log(c*x^n)))^p,x)

[Out] int(tanh(d*(a + b*log(c*x^n)))^p, x)

3.193 $\int (ex)^m \tanh^p (d(a + b \log (cx^n))) dx$

Optimal. Leaf size=135

$$\frac{(ex)^{1+m} \left(1 - e^{2ad}(cx^n)^{2bd}\right)^{-p} \left(-1 + e^{2ad}(cx^n)^{2bd}\right)^p F_1\left(\frac{1+m}{2bdn}; -p, p; 1 + \frac{1+m}{2bdn}; e^{2ad}(cx^n)^{2bd}, -e^{2ad}(cx^n)^{2bd}\right)}{e(1+m)}$$

[Out] $(e*x)^{(1+m)}*(-1+\exp(2*a*d)*(c*x^n)^{(2*b*d)})^p*\text{AppellF1}(1/2*(1+m)/b/d/n,-p,p,1+1/2*(1+m)/b/d/n,\exp(2*a*d)*(c*x^n)^{(2*b*d)},-\exp(2*a*d)*(c*x^n)^{(2*b*d)})/e/(1+m)/((1-\exp(2*a*d)*(c*x^n)^{(2*b*d)})^p)$

Rubi [A]

time = 0.11, antiderivative size = 135, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {5658, 5656, 525, 524}

$$\frac{(ex)^{m+1} \left(1 - e^{2ad}(cx^n)^{2bd}\right)^{-p} \left(e^{2ad}(cx^n)^{2bd} - 1\right)^p F_1\left(\frac{m+1}{2bdn}; -p, p; \frac{m+1}{2bdn} + 1; e^{2ad}(cx^n)^{2bd}, -e^{2ad}(cx^n)^{2bd}\right)}{e(m+1)}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(e*x)^m*\text{Tanh}[d*(a + b*\text{Log}[c*x^n])]^p,x]$

[Out] $((e*x)^{(1+m)}*(-1 + E^{(2*a*d)*(c*x^n)^{(2*b*d)}})^p*\text{AppellF1}[(1+m)/(2*b*d*n), -p, p, 1 + (1+m)/(2*b*d*n), E^{(2*a*d)*(c*x^n)^{(2*b*d)}}, -(E^{(2*a*d)*(c*x^n)^{(2*b*d)})}])/(e*(1+m)*(1 - E^{(2*a*d)*(c*x^n)^{(2*b*d)}})^p)$

Rule 524

$\text{Int}[(e_*)*(x_)^{(m_*)}*((a_*) + (b_*)*(x_)^{(n_)})^{(p_*)}*((c_*) + (d_*)*(x_)^{(n_)})^{(q_*)}, x_Symbol] :> \text{Simp}[a^p*c^q*((e*x)^{(m+1)})/(e*(m+1))*\text{AppellF1}[(m+1)/n, -p, -q, 1 + (m+1)/n, (-b)*(x^n/a), (-d)*(x^n/c)], x] /;$ FreeQ[{a, b, c, d, e, m, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[m, -1] && NeQ[m, n - 1] && (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])

Rule 525

$\text{Int}[(e_*)*(x_)^{(m_*)}*((a_*) + (b_*)*(x_)^{(n_)})^{(p_*)}*((c_*) + (d_*)*(x_)^{(n_)})^{(q_*)}, x_Symbol] :> \text{Dist}[a^{\text{IntPart}[p]}*((a + b*x^n)^{\text{FracPart}[p]}/(1 + b*(x^n/a)^{\text{FracPart}[p]}), \text{Int}[(e*x)^m*(1 + b*(x^n/a))^p*(c + d*x^n)^q, x], x] /;$ FreeQ[{a, b, c, d, e, m, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[m, -1] && NeQ[m, n - 1] && !(IntegerQ[p] || GtQ[a, 0])

Rule 5656

$\text{Int}[(e_*)*(x_)^{(m_*)}*\text{Tanh}[(a_*) + \text{Log}[x_*]*(b_*)*(d_*)]^{(p_*)}, x_Symbol] :> \text{Int}[(e*x)^m*((-1 + E^{(2*a*d)*x^{(2*b*d)}})^p/(1 + E^{(2*a*d)*x^{(2*b*d)}})^p),$

x] /; FreeQ[{a, b, d, e, m, p}, x]

Rule 5658

Int[((e_.)*(x_))^(m_.)*Tanh[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*(d_.)]^(p_.), x_Symbol] :> Dist[(e*x)^(m + 1)/(e*n*(c*x^n)^((m + 1)/n)), Subst[Int[x^((m + 1)/n - 1)*Tanh[d*(a + b*Log[x])]^p, x], x, c*x^n], x] /; FreeQ[{a, b, c, d, e, m, n, p}, x] && (NeQ[c, 1] || NeQ[n, 1])

Rubi steps

$$\int (ex)^m \tanh^p(d(a + b \log(cx^n))) dx = \int (ex)^m \tanh^p(d(a + b \log(cx^n))) dx$$

Mathematica [A]

time = 3.43, size = 174, normalized size = 1.29

$$\frac{x(ex)^m (1 - e^{2ad}(cx^n)^{2bd})^{-p} \left(\frac{-1 + e^{2ad}(cx^n)^{2bd}}{1 + e^{2ad}(cx^n)^{2bd}}\right)^p (1 + e^{2ad}(cx^n)^{2bd})^p F_1\left(\frac{1+m}{2bdn}; -p, p; 1 + \frac{1+m}{2bdn}; e^{2ad}(cx^n)^{2bd}, -e^{2ad}(cx^n)^{2bd}\right)}{1 + m}$$

Warning: Unable to verify antiderivative.

[In] Integrate[(e*x)^m*Tanh[d*(a + b*Log[c*x^n])]^p,x]

[Out] (x*(e*x)^m*((-1 + E^(2*a*d)*(c*x^n)^(2*b*d))/(1 + E^(2*a*d)*(c*x^n)^(2*b*d)))^p*(1 + E^(2*a*d)*(c*x^n)^(2*b*d))^p*AppellF1[(1 + m)/(2*b*d*n), -p, p, 1 + (1 + m)/(2*b*d*n), E^(2*a*d)*(c*x^n)^(2*b*d), -(E^(2*a*d)*(c*x^n)^(2*b*d))]/((1 + m)*(1 - E^(2*a*d)*(c*x^n)^(2*b*d))^p)

Maple [F]

time = 0.72, size = 0, normalized size = 0.00

$$\int (ex)^m (\tanh^p(d(a + b \ln(cx^n)))) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((e*x)^m*tanh(d*(a+b*ln(c*x^n)))^p,x)

[Out] int((e*x)^m*tanh(d*(a+b*ln(c*x^n)))^p,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e*x)^m*tanh(d*(a+b*log(c*x^n)))^p,x, algorithm="maxima")

[Out] integrate((x*e)^m*tanh((b*log(c*x^n) + a)*d)^p, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e*x)^m*tanh(d*(a+b*log(c*x^n)))^p,x, algorithm="fricas")

[Out] integral((x*e)^m*tanh(b*d*log(c*x^n) + a*d)^p, x)

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e*x)**m*tanh(d*(a+b*ln(c*x**n))))**p,x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((e*x)^m*tanh(d*(a+b*log(c*x^n)))^p,x, algorithm="giac")

[Out] integrate((e*x)^m*tanh((b*log(c*x^n) + a)*d)^p, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \tanh(d(a + b \ln(cx^n)))^p (ex)^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(d*(a + b*log(c*x^n)))^p*(e*x)^m,x)

[Out] int(tanh(d*(a + b*log(c*x^n)))^p*(e*x)^m, x)

$$3.194 \quad \int \frac{\tanh^{\frac{5}{2}}(a+b \log(cx^n))}{x} dx$$

Optimal. Leaf size=73

$$\frac{\operatorname{ArcTan}\left(\sqrt{\tanh(a+b \log(cx^n))}\right)}{bn} + \frac{\tanh^{-1}\left(\sqrt{\tanh(a+b \log(cx^n))}\right)}{bn} - \frac{2 \tanh^{\frac{3}{2}}(a+b \log(cx^n))}{3bn}$$

[Out] $-\arctan(\tanh(a+b*\ln(c*x^n))^{(1/2)})/b/n + \operatorname{arctanh}(\tanh(a+b*\ln(c*x^n))^{(1/2)})/b/n - 2/3*\tanh(a+b*\ln(c*x^n))^{(3/2)}/b/n$

Rubi [A]

time = 0.04, antiderivative size = 73, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$, Rules used = {3554, 3557, 335, 304, 209, 212}

$$\frac{\operatorname{ArcTan}\left(\sqrt{\tanh(a+b \log(cx^n))}\right)}{bn} + \frac{\tanh^{-1}\left(\sqrt{\tanh(a+b \log(cx^n))}\right)}{bn} - \frac{2 \tanh^{\frac{3}{2}}(a+b \log(cx^n))}{3bn}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[\operatorname{Tanh}[a + b*\operatorname{Log}[c*x^n]]^{(5/2)}/x, x]$

[Out] $-(\operatorname{ArcTan}[\operatorname{Sqrt}[\operatorname{Tanh}[a + b*\operatorname{Log}[c*x^n]]]]/(b*n)) + \operatorname{ArcTanh}[\operatorname{Sqrt}[\operatorname{Tanh}[a + b*\operatorname{Log}[c*x^n]]]]/(b*n) - (2*\operatorname{Tanh}[a + b*\operatorname{Log}[c*x^n]]^{(3/2)})/(3*b*n)$

Rule 209

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[b, 2]))*\operatorname{ArcTan}[\operatorname{Rt}[b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{PosQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{GtQ}[b, 0])$

Rule 212

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$

Rule 304

$\operatorname{Int}[(x_)^2/((a_ + (b_)*(x_)^4), x_Symbol] \rightarrow \operatorname{With}\{r = \operatorname{Numerator}[\operatorname{Rt}[-a/b, 2]], s = \operatorname{Denominator}[\operatorname{Rt}[-a/b, 2]]\}, \operatorname{Dist}[s/(2*b), \operatorname{Int}[1/(r + s*x^2), x], x] - \operatorname{Dist}[s/(2*b), \operatorname{Int}[1/(r - s*x^2), x], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \ !\operatorname{GtQ}[a/b, 0]$

Rule 335

```
Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
  Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
  )]^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 3554

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d
*x])^(n - 1)/(d*(n - 1))), x] - Dist[b^2, Int[(b*Tan[c + d*x])^(n - 2), x],
x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]
```

Rule 3557

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[b/d, Subst[Int[
x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !
IntegerQ[n]
```

Rubi steps

$$\begin{aligned}
\int \frac{\tanh^{\frac{5}{2}}(a + b \log(cx^n))}{x} dx &= \frac{\text{Subst}\left(\int \tanh^{\frac{5}{2}}(a + bx) dx, x, \log(cx^n)\right)}{n} \\
&= -\frac{2 \tanh^{\frac{3}{2}}(a + b \log(cx^n))}{3bn} + \frac{\text{Subst}\left(\int \sqrt{\tanh(a + bx)} dx, x, \log(cx^n)\right)}{n} \\
&= -\frac{2 \tanh^{\frac{3}{2}}(a + b \log(cx^n))}{3bn} - \frac{\text{Subst}\left(\int \frac{\sqrt{x}}{-1+x^2} dx, x, \tanh(a + b \log(cx^n))\right)}{bn} \\
&= -\frac{2 \tanh^{\frac{3}{2}}(a + b \log(cx^n))}{3bn} - \frac{2 \text{Subst}\left(\int \frac{x^2}{-1+x^4} dx, x, \sqrt{\tanh(a + b \log(cx^n))}\right)}{bn} \\
&= -\frac{2 \tanh^{\frac{3}{2}}(a + b \log(cx^n))}{3bn} + \frac{\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \sqrt{\tanh(a + b \log(cx^n))}\right)}{bn} \\
&= -\frac{\tan^{-1}\left(\sqrt{\tanh(a + b \log(cx^n))}\right)}{bn} + \frac{\tanh^{-1}\left(\sqrt{\tanh(a + b \log(cx^n))}\right)}{bn}
\end{aligned}$$

Mathematica [A]

time = 0.24, size = 64, normalized size = 0.88

$$\frac{3 \text{ArcTan}\left(\sqrt{\tanh(a + b \log(cx^n))}\right) - 3 \tanh^{-1}\left(\sqrt{\tanh(a + b \log(cx^n))}\right) + 2 \tanh^{\frac{3}{2}}(a + b \log(cx^n))}{3bn}$$

Antiderivative was successfully verified.

[In] Integrate[Tanh[a + b*Log[c*x^n]]^(5/2)/x,x]

[Out] $-1/3*(3*\text{ArcTan}[\text{Sqrt}[\text{Tanh}[a + b*\text{Log}[c*x^n]]]] - 3*\text{ArcTanh}[\text{Sqrt}[\text{Tanh}[a + b*\text{Log}[c*x^n]]]] + 2*\text{Tanh}[a + b*\text{Log}[c*x^n]]^{(3/2)})/(b*n)$

Maple [A]

time = 4.85, size = 76, normalized size = 1.04

method	result
derivativedivides	$\frac{2 \left(\tanh^{\frac{3}{2}}(a+b \ln(cx^n)) \right)}{3} - \frac{\ln\left(\sqrt{\tanh(a+b \ln(cx^n))-1}\right)}{2} + \frac{\ln\left(\sqrt{\tanh(a+b \ln(cx^n))+1}\right)}{2} - \arctan\left(\sqrt{\tanh(a+b \ln(cx^n))}\right)}{nb}$
default	$\frac{2 \left(\tanh^{\frac{3}{2}}(a+b \ln(cx^n)) \right)}{3} - \frac{\ln\left(\sqrt{\tanh(a+b \ln(cx^n))-1}\right)}{2} + \frac{\ln\left(\sqrt{\tanh(a+b \ln(cx^n))+1}\right)}{2} - \arctan\left(\sqrt{\tanh(a+b \ln(cx^n))}\right)}{nb}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(a+b*ln(c*x^n))^(5/2)/x,x,method=_RETURNVERBOSE)

[Out] $1/n/b*(-2/3*\text{tanh}(a+b*\ln(c*x^n))^{(3/2)}-1/2*\ln(\text{tanh}(a+b*\ln(c*x^n))^{(1/2)}-1)+1/2*\ln(\text{tanh}(a+b*\ln(c*x^n))^{(1/2)}+1)-\arctan(\text{tanh}(a+b*\ln(c*x^n))^{(1/2)}))$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+b*log(c*x^n))^(5/2)/x,x, algorithm="maxima")

[Out] integrate(tanh(b*log(c*x^n) + a)^(5/2)/x, x)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 625 vs. $2(65) = 130$.

time = 0.39, size = 625, normalized size = 8.56

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+b*log(c*x^n))^(5/2)/x,x, algorithm="fricas")

[Out] $-1/6*(6*(\cosh(b*n*\log(x) + b*\log(c) + a)^2 + 2*\cosh(b*n*\log(x) + b*\log(c) + a)*\sinh(b*n*\log(x) + b*\log(c) + a) + \sinh(b*n*\log(x) + b*\log(c) + a)^2 + 1)*\arctan(-\cosh(b*n*\log(x) + b*\log(c) + a)^2 - 2*\cosh(b*n*\log(x) + b*\log(c) + a)*\sinh(b*n*\log(x) + b*\log(c) + a) - \sinh(b*n*\log(x) + b*\log(c) + a)^2 + (\cosh(b*n*\log(x) + b*\log(c) + a)^2 + 2*\cosh(b*n*\log(x) + b*\log(c) + a)*\sinh$


```
(b*n*log(x) + b*log(c) + a) + sinh(b*n*log(x) + b*log(c) + a)^2 + 1)*sqrt(sinh(b*n*log(x) + b*log(c) + a)/cosh(b*n*log(x) + b*log(c) + a)) + 4*cosh(b*n*log(x) + b*log(c) + a)^2 + 3*(cosh(b*n*log(x) + b*log(c) + a)^2 + 2*cosh(b*n*log(x) + b*log(c) + a)*sinh(b*n*log(x) + b*log(c) + a) + sinh(b*n*log(x) + b*log(c) + a)^2 + 1)*log(-cosh(b*n*log(x) + b*log(c) + a)^2 - 2*cosh(b*n*log(x) + b*log(c) + a)*sinh(b*n*log(x) + b*log(c) + a) - sinh(b*n*log(x) + b*log(c) + a)^2 + (cosh(b*n*log(x) + b*log(c) + a)^2 + 2*cosh(b*n*log(x) + b*log(c) + a)*sinh(b*n*log(x) + b*log(c) + a) + sinh(b*n*log(x) + b*log(c) + a)^2 + 1)*sqrt(sinh(b*n*log(x) + b*log(c) + a)/cosh(b*n*log(x) + b*log(c) + a))) + 8*cosh(b*n*log(x) + b*log(c) + a)*sinh(b*n*log(x) + b*log(c) + a) + 4*sinh(b*n*log(x) + b*log(c) + a)^2 + 4*(cosh(b*n*log(x) + b*log(c) + a)^2 + 2*cosh(b*n*log(x) + b*log(c) + a)*sinh(b*n*log(x) + b*log(c) + a) + sinh(b*n*log(x) + b*log(c) + a)^2 - 1)*sqrt(sinh(b*n*log(x) + b*log(c) + a)/cosh(b*n*log(x) + b*log(c) + a)) + 4)/(b*n*cosh(b*n*log(x) + b*log(c) + a)^2 + 2*b*n*cosh(b*n*log(x) + b*log(c) + a)*sinh(b*n*log(x) + b*log(c) + a) + b*n*sinh(b*n*log(x) + b*log(c) + a)^2 + b*n)
```

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(tanh(a+b*ln(c*x**n))**(5/2)/x,x)
```

```
[Out] Exception raised: SystemError >> excessive stack use: stack is 4371 deep
```

Giac [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(tanh(a+b*log(c*x^n))^(5/2)/x,x, algorithm="giac")
```

```
[Out] Timed out
```

Mupad [B]

time = 2.25, size = 65, normalized size = 0.89

$$\frac{\operatorname{atanh}\left(\sqrt{\tanh(a+b\ln(cx^n))}\right)}{bn} - \frac{\operatorname{atan}\left(\sqrt{\tanh(a+b\ln(cx^n))}\right)}{bn} - \frac{2\tanh(a+b\ln(cx^n))^{3/2}}{3bn}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(tanh(a + b*log(c*x^n))^(5/2)/x,x)
```

```
[Out] atanh(tanh(a + b*log(c*x^n))^(1/2))/(b*n) - atan(tanh(a + b*log(c*x^n))^(1/2))/(b*n) - (2*tanh(a + b*log(c*x^n))^(3/2))/(3*b*n)
```

$$3.195 \quad \int \frac{\tanh^{\frac{3}{2}}(a+b \log(cx^n))}{x} dx$$

Optimal. Leaf size=70

$$\frac{\operatorname{ArcTan}\left(\sqrt{\tanh(a+b \log(cx^n))}\right)}{bn} + \frac{\tanh^{-1}\left(\sqrt{\tanh(a+b \log(cx^n))}\right)}{bn} - \frac{2\sqrt{\tanh(a+b \log(cx^n))}}{bn}$$

[Out] arctan(tanh(a+b*ln(c*x^n))^(1/2))/b/n+arctanh(tanh(a+b*ln(c*x^n))^(1/2))/b/n-2*tanh(a+b*ln(c*x^n))^(1/2)/b/n

Rubi [A]

time = 0.04, antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$, Rules used = {3554, 3557, 335, 218, 212, 209}

$$\frac{\operatorname{ArcTan}\left(\sqrt{\tanh(a+b \log(cx^n))}\right)}{bn} + \frac{\tanh^{-1}\left(\sqrt{\tanh(a+b \log(cx^n))}\right)}{bn} - \frac{2\sqrt{\tanh(a+b \log(cx^n))}}{bn}$$

Antiderivative was successfully verified.

[In] Int[Tanh[a + b*Log[c*x^n]]^(3/2)/x,x]

[Out] ArcTan[Sqrt[Tanh[a + b*Log[c*x^n]]]]/(b*n) + ArcTanh[Sqrt[Tanh[a + b*Log[c*x^n]]]]/(b*n) - (2*Sqrt[Tanh[a + b*Log[c*x^n]]])/(b*n)

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 218

Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 335

```
Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
  Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
  ))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 3554

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d
*x])^(n - 1)/(d*(n - 1))), x] - Dist[b^2, Int[(b*Tan[c + d*x])^(n - 2), x],
x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]
```

Rule 3557

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[b/d, Subst[Int[
x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !
IntegerQ[n]
```

Rubi steps

$$\begin{aligned}
\int \frac{\tanh^{\frac{3}{2}}(a + b \log(cx^n))}{x} dx &= \frac{\text{Subst}\left(\int \tanh^{\frac{3}{2}}(a + bx) dx, x, \log(cx^n)\right)}{n} \\
&= -\frac{2\sqrt{\tanh(a + b \log(cx^n))}}{bn} + \frac{\text{Subst}\left(\int \frac{1}{\sqrt{\tanh(a + bx)}} dx, x, \log(cx^n)\right)}{n} \\
&= -\frac{2\sqrt{\tanh(a + b \log(cx^n))}}{bn} - \frac{\text{Subst}\left(\int \frac{1}{\sqrt{x(-1+x^2)}} dx, x, \tanh(a + b \log(cx^n))\right)}{bn} \\
&= -\frac{2\sqrt{\tanh(a + b \log(cx^n))}}{bn} - \frac{2\text{Subst}\left(\int \frac{1}{-1+x^4} dx, x, \sqrt{\tanh(a + b \log(cx^n))}\right)}{bn} \\
&= -\frac{2\sqrt{\tanh(a + b \log(cx^n))}}{bn} + \frac{\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \sqrt{\tanh(a + b \log(cx^n))}\right)}{bn} \\
&= \frac{\tan^{-1}\left(\sqrt{\tanh(a + b \log(cx^n))}\right)}{bn} + \frac{\tanh^{-1}\left(\sqrt{\tanh(a + b \log(cx^n))}\right)}{bn}
\end{aligned}$$

Mathematica [A]

time = 0.13, size = 57, normalized size = 0.81

$$\frac{\text{ArcTan}\left(\sqrt{\tanh(a + b \log(cx^n))}\right) + \tanh^{-1}\left(\sqrt{\tanh(a + b \log(cx^n))}\right) - 2\sqrt{\tanh(a + b \log(cx^n))}}{bn}$$

Antiderivative was successfully verified.

[In] Integrate[Tanh[a + b*Log[c*x^n]]^(3/2)/x,x]

[Out] (ArcTan[Sqrt[Tanh[a + b*Log[c*x^n]]]] + ArcTanh[Sqrt[Tanh[a + b*Log[c*x^n]]]] - 2*Sqrt[Tanh[a + b*Log[c*x^n]]])/(b*n)

Maple [A]

time = 4.18, size = 74, normalized size = 1.06

method	result
derivativedivides	$\frac{-2\left(\sqrt{\tanh(a+b\ln(cx^n))}\right) - \frac{\ln\left(\sqrt{\tanh(a+b\ln(cx^n))-1}\right)}{2} + \frac{\ln\left(\sqrt{\tanh(a+b\ln(cx^n))+1}\right)}{2} + \arctan\left(\sqrt{\tanh(a+b\ln(cx^n))}\right)}{nb}$
default	$\frac{-2\left(\sqrt{\tanh(a+b\ln(cx^n))}\right) - \frac{\ln\left(\sqrt{\tanh(a+b\ln(cx^n))-1}\right)}{2} + \frac{\ln\left(\sqrt{\tanh(a+b\ln(cx^n))+1}\right)}{2} + \arctan\left(\sqrt{\tanh(a+b\ln(cx^n))}\right)}{nb}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(a+b*ln(c*x^n))^(3/2)/x,x,method=_RETURNVERBOSE)

[Out] 1/n/b*(-2*tanh(a+b*ln(c*x^n))^(1/2)-1/2*ln(tanh(a+b*ln(c*x^n))^(1/2)-1)+1/2*ln(tanh(a+b*ln(c*x^n))^(1/2)+1)+arctan(tanh(a+b*ln(c*x^n))^(1/2)))

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+b*log(c*x^n))^(3/2)/x,x, algorithm="maxima")

[Out] integrate(tanh(b*log(c*x^n) + a)^(3/2)/x, x)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 334 vs. 2(64) = 128.

time = 0.38, size = 334, normalized size = 4.77

Antiderivative was successfully verified.
[In] integrate(tanh(a+b*log(c*x^n))^(3/2)/x,x, algorithm="fricas")
[Out] -1/2*(4*sqrt(sinh(b*n*log(x) + b*log(c) + a)/cosh(b*n*log(x) + b*log(c) + a)) - 2*arctan(-cosh(b*n*log(x) + b*log(c) + a)^2 - 2*cosh(b*n*log(x) + b*log(c) + a)*sinh(b*n*log(x) + b*log(c) + a) - sinh(b*n*log(x) + b*log(c) + a)^2 + (cosh(b*n*log(x) + b*log(c) + a)^2 + 2*cosh(b*n*log(x) + b*log(c) + a)

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+b*log(c*x^n))^(3/2)/x,x, algorithm="fricas")

[Out] -1/2*(4*sqrt(sinh(b*n*log(x) + b*log(c) + a)/cosh(b*n*log(x) + b*log(c) + a)) - 2*arctan(-cosh(b*n*log(x) + b*log(c) + a)^2 - 2*cosh(b*n*log(x) + b*log(c) + a)*sinh(b*n*log(x) + b*log(c) + a) - sinh(b*n*log(x) + b*log(c) + a)^2 + (cosh(b*n*log(x) + b*log(c) + a)^2 + 2*cosh(b*n*log(x) + b*log(c) + a)

```
*sinh(b*n*log(x) + b*log(c) + a) + sinh(b*n*log(x) + b*log(c) + a)^2 + 1)*sqrt(sinh(b*n*log(x) + b*log(c) + a)/cosh(b*n*log(x) + b*log(c) + a))) + log(-cosh(b*n*log(x) + b*log(c) + a)^2 - 2*cosh(b*n*log(x) + b*log(c) + a)*sinh(b*n*log(x) + b*log(c) + a) - sinh(b*n*log(x) + b*log(c) + a)^2 + (cosh(b*n*log(x) + b*log(c) + a)^2 + 2*cosh(b*n*log(x) + b*log(c) + a)*sinh(b*n*log(x) + b*log(c) + a) + sinh(b*n*log(x) + b*log(c) + a)^2 + 1)*sqrt(sinh(b*n*log(x) + b*log(c) + a)/cosh(b*n*log(x) + b*log(c) + a))))/(b*n)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\tanh^{\frac{3}{2}}(a + b \log(cx^n))}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(tanh(a+b*ln(c*x**n))**(3/2)/x,x)
```

```
[Out] Integral(tanh(a + b*log(c*x**n))**(3/2)/x, x)
```

Giac [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(tanh(a+b*log(c*x^n))^(3/2)/x,x, algorithm="giac")
```

```
[Out] Timed out
```

Mupad [B]

time = 1.80, size = 51, normalized size = 0.73

$$\frac{\operatorname{atan}\left(\sqrt{\tanh(a + b \ln(cx^n))}\right) + \operatorname{atanh}\left(\sqrt{\tanh(a + b \ln(cx^n))}\right) - 2\sqrt{\tanh(a + b \ln(cx^n))}}{bn}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(tanh(a + b*log(c*x^n))^(3/2)/x,x)
```

```
[Out] (atan(tanh(a + b*log(c*x^n))^(1/2)) + atanh(tanh(a + b*log(c*x^n))^(1/2)) - 2*tanh(a + b*log(c*x^n))^(1/2))/(b*n)
```

$$3.196 \quad \int \frac{\sqrt{\tanh(a + b \log(cx^n))}}{x} dx$$

Optimal. Leaf size=48

$$-\frac{\operatorname{ArcTan}\left(\sqrt{\tanh(a + b \log(cx^n))}\right)}{bn} + \frac{\tanh^{-1}\left(\sqrt{\tanh(a + b \log(cx^n))}\right)}{bn}$$

[Out] $-\arctan(\tanh(a+b*\ln(c*x^n))^{(1/2)})/b/n+\operatorname{arctanh}(\tanh(a+b*\ln(c*x^n))^{(1/2)})/b/n$

Rubi [A]

time = 0.03, antiderivative size = 48, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$, Rules used = {3557, 335, 304, 209, 212}

$$\frac{\tanh^{-1}\left(\sqrt{\tanh(a + b \log(cx^n))}\right)}{bn} - \frac{\operatorname{ArcTan}\left(\sqrt{\tanh(a + b \log(cx^n))}\right)}{bn}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[Tanh[a + b*Log[c*x^n]]]/x,x]

[Out] $-(\operatorname{ArcTan}[\operatorname{Sqrt}[\operatorname{Tanh}[a + b*\operatorname{Log}[c*x^n]]]]/(b*n)) + \operatorname{ArcTanh}[\operatorname{Sqrt}[\operatorname{Tanh}[a + b*\operatorname{Log}[c*x^n]]]]/(b*n)$

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 304

Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 335

```
Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
  Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
  ))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 3557

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[b/d, Subst[Int[
x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !
IntegerQ[n]
```

Rubi steps

$$\begin{aligned}
 \int \frac{\sqrt{\tanh(a + b \log(cx^n))}}{x} dx &= \frac{\text{Subst}\left(\int \sqrt{\tanh(a + bx)} dx, x, \log(cx^n)\right)}{n} \\
 &= -\frac{\text{Subst}\left(\int \frac{\sqrt{x}}{-1+x^2} dx, x, \tanh(a + b \log(cx^n))\right)}{bn} \\
 &= -\frac{2\text{Subst}\left(\int \frac{x^2}{-1+x^4} dx, x, \sqrt{\tanh(a + b \log(cx^n))}\right)}{bn} \\
 &= \frac{\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \sqrt{\tanh(a + b \log(cx^n))}\right)}{bn} - \frac{\text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \sqrt{\tanh(a + b \log(cx^n))}\right)}{bn} \\
 &= -\frac{\tan^{-1}\left(\sqrt{\tanh(a + b \log(cx^n))}\right)}{bn} + \frac{\tanh^{-1}\left(\sqrt{\tanh(a + b \log(cx^n))}\right)}{bn}
 \end{aligned}$$

Mathematica [A]

time = 0.06, size = 48, normalized size = 1.00

$$-\frac{\text{ArcTan}\left(\sqrt{\tanh(a + b \log(cx^n))}\right)}{bn} + \frac{\tanh^{-1}\left(\sqrt{\tanh(a + b \log(cx^n))}\right)}{bn}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sqrt[Tanh[a + b*Log[c*x^n]]]/x,x]
```

```
[Out] -(ArcTan[Sqrt[Tanh[a + b*Log[c*x^n]]]/(b*n)] + ArcTanh[Sqrt[Tanh[a + b*Log
[c*x^n]]]/(b*n)
```

Maple [A]

time = 4.15, size = 61, normalized size = 1.27

method	result	size
derivativedivides	$\frac{\frac{\ln\left(\sqrt{\tanh\left(a+b\ln(cx^n)\right)-1}\right)}{2} + \frac{\ln\left(\sqrt{\tanh\left(a+b\ln(cx^n)\right)+1}\right)}{2} - \arctan\left(\sqrt{\tanh\left(a+b\ln(cx^n)\right)}\right)}{nb}$	61
default	$\frac{\frac{\ln\left(\sqrt{\tanh\left(a+b\ln(cx^n)\right)-1}\right)}{2} + \frac{\ln\left(\sqrt{\tanh\left(a+b\ln(cx^n)\right)+1}\right)}{2} - \arctan\left(\sqrt{\tanh\left(a+b\ln(cx^n)\right)}\right)}{nb}$	61

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(tanh(a+b*ln(c*x^n))^(1/2)/x,x,method=_RETURNVERBOSE)
```

```
[Out] 1/n/b*(-1/2*ln(tanh(a+b*ln(c*x^n))^(1/2)-1)+1/2*ln(tanh(a+b*ln(c*x^n))^(1/2)+1)-arctan(tanh(a+b*ln(c*x^n))^(1/2)))
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(tanh(a+b*log(c*x^n))^(1/2)/x,x, algorithm="maxima")
```

```
[Out] integrate(sqrt(tanh(b*log(c*x^n) + a))/x, x)
```

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 303 vs. 2(44) = 88.

time = 0.36, size = 303, normalized size = 6.31

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(tanh(a+b*log(c*x^n))^(1/2)/x,x, algorithm="fricas")
```

```
[Out] -1/2*(2*arctan(-cosh(b*n*log(x) + b*log(c) + a)^2 - 2*cosh(b*n*log(x) + b*log(c) + a)*sinh(b*n*log(x) + b*log(c) + a) - sinh(b*n*log(x) + b*log(c) + a)^2 + (cosh(b*n*log(x) + b*log(c) + a)^2 + 2*cosh(b*n*log(x) + b*log(c) + a)*sinh(b*n*log(x) + b*log(c) + a) + sinh(b*n*log(x) + b*log(c) + a)^2 + 1)*sqrt(sinh(b*n*log(x) + b*log(c) + a)/cosh(b*n*log(x) + b*log(c) + a))) + log(-cosh(b*n*log(x) + b*log(c) + a)^2 - 2*cosh(b*n*log(x) + b*log(c) + a)*sinh(b*n*log(x) + b*log(c) + a) - sinh(b*n*log(x) + b*log(c) + a)^2 + (cosh(b*n*log(x) + b*log(c) + a)^2 + 2*cosh(b*n*log(x) + b*log(c) + a)*sinh(b*n*log(x) + b*log(c) + a) + sinh(b*n*log(x) + b*log(c) + a)^2 + 1)*sqrt(sinh(b*n*log(x) + b*log(c) + a)/cosh(b*n*log(x) + b*log(c) + a))))/(b*n)
```


Sympy [A]

time = 1.56, size = 66, normalized size = 1.38

$$-\frac{\log\left(\sqrt{\tanh(a+b\log(cx^n))}-1\right)}{2bn} + \frac{\log\left(\sqrt{\tanh(a+b\log(cx^n))+1}\right)}{2bn} - \frac{\operatorname{atan}\left(\sqrt{\tanh(a+b\log(cx^n))}\right)}{bn}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+b*ln(c*x**n))**(1/2)/x,x)

[Out] -log(sqrt(tanh(a + b*log(c*x**n))) - 1)/(2*b*n) + log(sqrt(tanh(a + b*log(c*x**n))) + 1)/(2*b*n) - atan(sqrt(tanh(a + b*log(c*x**n))))/(b*n)

Giac [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(a+b*log(c*x^n))^(1/2)/x,x, algorithm="giac")

[Out] Timed out

Mupad [B]

time = 1.45, size = 39, normalized size = 0.81

$$-\frac{\operatorname{atan}\left(\sqrt{\tanh(a+b\ln(cx^n))}\right) - \operatorname{atanh}\left(\sqrt{\tanh(a+b\ln(cx^n))}\right)}{bn}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(a + b*log(c*x^n))^(1/2)/x,x)

[Out] -(atan(tanh(a + b*log(c*x^n))^(1/2)) - atanh(tanh(a + b*log(c*x^n))^(1/2)))/(b*n)

$$3.197 \quad \int \frac{1}{x \sqrt{\tanh(a + b \log(cx^n))}} dx$$

Optimal. Leaf size=47

$$\frac{\text{ArcTan}\left(\sqrt{\tanh(a + b \log(cx^n))}\right)}{bn} + \frac{\tanh^{-1}\left(\sqrt{\tanh(a + b \log(cx^n))}\right)}{bn}$$

[Out] arctan(tanh(a+b*ln(c*x^n))^(1/2))/b/n+arctanh(tanh(a+b*ln(c*x^n))^(1/2))/b/n

Rubi [A]

time = 0.03, antiderivative size = 47, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$, Rules used = {3557, 335, 218, 212, 209}

$$\frac{\text{ArcTan}\left(\sqrt{\tanh(a + b \log(cx^n))}\right)}{bn} + \frac{\tanh^{-1}\left(\sqrt{\tanh(a + b \log(cx^n))}\right)}{bn}$$

Antiderivative was successfully verified.

[In] Int[1/(x*Sqrt[Tanh[a + b*Log[c*x^n]]]),x]

[Out] ArcTan[Sqrt[Tanh[a + b*Log[c*x^n]]]]/(b*n) + ArcTanh[Sqrt[Tanh[a + b*Log[c*x^n]]]]/(b*n)

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 218

Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] :> With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 335

```
Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
  Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
  ))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
  ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 3557

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[b/d, Subst[Int[
  x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !
  IntegerQ[n]
```

Rubi steps

$$\begin{aligned} \int \frac{1}{x \sqrt{\tanh(a + b \log(cx^n))}} dx &= \frac{\text{Subst}\left(\int \frac{1}{\sqrt{\tanh(a + bx)}} dx, x, \log(cx^n)\right)}{n} \\ &= -\frac{\text{Subst}\left(\int \frac{1}{\sqrt{x}(-1+x^2)} dx, x, \tanh(a + b \log(cx^n))\right)}{bn} \\ &= -\frac{2\text{Subst}\left(\int \frac{1}{-1+x^4} dx, x, \sqrt{\tanh(a + b \log(cx^n))}\right)}{bn} \\ &= \frac{\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \sqrt{\tanh(a + b \log(cx^n))}\right)}{bn} + \frac{\text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \sqrt{\tanh(a + b \log(cx^n))}\right)}{bn} \\ &= \frac{\tan^{-1}\left(\sqrt{\tanh(a + b \log(cx^n))}\right)}{bn} + \frac{\tanh^{-1}\left(\sqrt{\tanh(a + b \log(cx^n))}\right)}{bn} \end{aligned}$$

Mathematica [A]

time = 0.08, size = 47, normalized size = 1.00

$$\frac{\text{ArcTan}\left(\sqrt{\tanh(a + b \log(cx^n))}\right)}{bn} + \frac{\tanh^{-1}\left(\sqrt{\tanh(a + b \log(cx^n))}\right)}{bn}$$

Antiderivative was successfully verified.

```
[In] Integrate[1/(x*Sqrt[Tanh[a + b*Log[c*x^n]]]), x]
```

```
[Out] ArcTan[Sqrt[Tanh[a + b*Log[c*x^n]]]/(b*n) + ArcTanh[Sqrt[Tanh[a + b*Log[c*
  x^n]]]/(b*n)]
```

Maple [A]

time = 4.16, size = 37, normalized size = 0.79

method	result	size
derivativedivides	$\frac{\operatorname{arctanh}\left(\sqrt{\tanh(a+b\ln(cx^n))}\right)+\operatorname{arctan}\left(\sqrt{\tanh(a+b\ln(cx^n))}\right)}{nb}$	37
default	$\frac{\operatorname{arctanh}\left(\sqrt{\tanh(a+b\ln(cx^n))}\right)+\operatorname{arctan}\left(\sqrt{\tanh(a+b\ln(cx^n))}\right)}{nb}$	37

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/x/tanh(a+b*ln(c*x^n))^(1/2),x,method=_RETURNVERBOSE)`

[Out] `1/n/b*(arctanh(tanh(a+b*ln(c*x^n))^(1/2))+arctan(tanh(a+b*ln(c*x^n))^(1/2)))`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/x/tanh(a+b*log(c*x^n))^(1/2),x, algorithm="maxima")`

[Out] `integrate(1/(x*sqrt(tanh(b*log(c*x^n) + a))), x)`

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 305 vs. 2(43) = 86.

time = 0.39, size = 305, normalized size = 6.49

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/x/tanh(a+b*log(c*x^n))^(1/2),x, algorithm="fricas")`

[Out]
$$\frac{1}{2} \cdot \frac{(2 \cdot \arctan(-\cosh(b \cdot n \cdot \log(x) + b \cdot \log(c) + a))^2 - 2 \cdot \cosh(b \cdot n \cdot \log(x) + b \cdot \log(c) + a) \cdot \sinh(b \cdot n \cdot \log(x) + b \cdot \log(c) + a) - \sinh(b \cdot n \cdot \log(x) + b \cdot \log(c) + a)^2 + (\cosh(b \cdot n \cdot \log(x) + b \cdot \log(c) + a))^2 + 2 \cdot \cosh(b \cdot n \cdot \log(x) + b \cdot \log(c) + a) \cdot \sinh(b \cdot n \cdot \log(x) + b \cdot \log(c) + a) + \sinh(b \cdot n \cdot \log(x) + b \cdot \log(c) + a)^2 + 1) \cdot \sqrt{\sinh(b \cdot n \cdot \log(x) + b \cdot \log(c) + a) / \cosh(b \cdot n \cdot \log(x) + b \cdot \log(c) + a)}}{(-\cosh(b \cdot n \cdot \log(x) + b \cdot \log(c) + a))^2 - 2 \cdot \cosh(b \cdot n \cdot \log(x) + b \cdot \log(c) + a) \cdot \sinh(b \cdot n \cdot \log(x) + b \cdot \log(c) + a) - \sinh(b \cdot n \cdot \log(x) + b \cdot \log(c) + a)^2 + (\cosh(b \cdot n \cdot \log(x) + b \cdot \log(c) + a))^2 + 2 \cdot \cosh(b \cdot n \cdot \log(x) + b \cdot \log(c) + a) \cdot \sinh(b \cdot n \cdot \log(x) + b \cdot \log(c) + a) + \sinh(b \cdot n \cdot \log(x) + b \cdot \log(c) + a)^2 + 1) \cdot \sqrt{\sinh(b \cdot n \cdot \log(x) + b \cdot \log(c) + a) / \cosh(b \cdot n \cdot \log(x) + b \cdot \log(c) + a))} / (b \cdot n)}$$

Sympy [A]

time = 2.47, size = 66, normalized size = 1.40

$$-\frac{\log\left(\sqrt{\tanh(a+b\log(cx^n))}-1\right)}{2bn} + \frac{\log\left(\sqrt{\tanh(a+b\log(cx^n))}+1\right)}{2bn} + \frac{\operatorname{atan}\left(\sqrt{\tanh(a+b\log(cx^n))}\right)}{bn}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/x/tanh(a+b*ln(c*x**n))**(1/2),x)

[Out] $-\log(\sqrt{\tanh(a + b \log(c x^n))}) - 1)/(2 * b * n) + \log(\sqrt{\tanh(a + b \log(c x^n))}) + 1)/(2 * b * n) + \operatorname{atan}(\sqrt{\tanh(a + b \log(c x^n))})/(b * n)$

Giac [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/x/tanh(a+b*log(c*x^n))^(1/2),x, algorithm="giac")

[Out] Timed out

Mupad [B]

time = 1.55, size = 36, normalized size = 0.77

$$\frac{\operatorname{atan}\left(\sqrt{\tanh(a + b \ln(cx^n))}\right) + \operatorname{atanh}\left(\sqrt{\tanh(a + b \ln(cx^n))}\right)}{bn}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(x*tanh(a + b*log(c*x^n))^(1/2)),x)

[Out] $(\operatorname{atan}(\tanh(a + b \log(c x^n))^{1/2}) + \operatorname{atanh}(\tanh(a + b \log(c x^n))^{1/2}))/ (b * n)$

$$3.198 \quad \int \frac{1}{x \tanh^2(a+b \log(cx^n))} dx$$

Optimal. Leaf size=71

$$-\frac{\operatorname{ArcTan}\left(\sqrt{\tanh(a+b \log(cx^n))}\right)}{bn} + \frac{\tanh^{-1}\left(\sqrt{\tanh(a+b \log(cx^n))}\right)}{bn} - \frac{2}{bn \sqrt{\tanh(a+b \log(cx^n))}}$$

[Out] $-\arctan(\tanh(a+b*\ln(c*x^n))^{(1/2)})/b/n + \operatorname{arctanh}(\tanh(a+b*\ln(c*x^n))^{(1/2)})/b/n - 2/b/n/\tanh(a+b*\ln(c*x^n))^{(1/2)}$

Rubi [A]

time = 0.04, antiderivative size = 71, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$, Rules used = {3555, 3557, 335, 304, 209, 212}

$$-\frac{\operatorname{ArcTan}\left(\sqrt{\tanh(a+b \log(cx^n))}\right)}{bn} + \frac{\tanh^{-1}\left(\sqrt{\tanh(a+b \log(cx^n))}\right)}{bn} - \frac{2}{bn \sqrt{\tanh(a+b \log(cx^n))}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[1/(x*\operatorname{Tanh}[a + b*\operatorname{Log}[c*x^n]]^{(3/2)}), x]$

[Out] $-(\operatorname{ArcTan}[\operatorname{Sqrt}[\operatorname{Tanh}[a + b*\operatorname{Log}[c*x^n]]]]/(b*n)) + \operatorname{ArcTanh}[\operatorname{Sqrt}[\operatorname{Tanh}[a + b*\operatorname{Log}[c*x^n]]]]/(b*n) - 2/(b*n*\operatorname{Sqrt}[\operatorname{Tanh}[a + b*\operatorname{Log}[c*x^n]]])$

Rule 209

$\operatorname{Int}(((a_) + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[b, 2]))*\operatorname{ArcTan}[\operatorname{Rt}[b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{PosQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{GtQ}[b, 0])$

Rule 212

$\operatorname{Int}(((a_) + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$

Rule 304

$\operatorname{Int}(x^2/((a_) + (b_)*(x_)^4), x_Symbol] \rightarrow \operatorname{With}[\{r = \operatorname{Numerator}[\operatorname{Rt}[-a/b, 2]], s = \operatorname{Denominator}[\operatorname{Rt}[-a/b, 2]]\}, \operatorname{Dist}[s/(2*b), \operatorname{Int}[1/(r + s*x^2), x], x] - \operatorname{Dist}[s/(2*b), \operatorname{Int}[1/(r - s*x^2), x], x]] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \ !\operatorname{GtQ}[a/b, 0]$

Rule 335

```
Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
  Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
  ))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 3555

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(b*Tan[c + d*x]
)^^(n + 1)/(b*d*(n + 1)), x] - Dist[1/b^2, Int[(b*Tan[c + d*x])^(n + 2), x],
x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1]
```

Rule 3557

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[b/d, Subst[Int[
x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !
IntegerQ[n]
```

Rubi steps

$$\begin{aligned}
 \int \frac{1}{x \tanh^{\frac{3}{2}}(a + b \log(cx^n))} dx &= \frac{\text{Subst}\left(\int \frac{1}{\tanh^{\frac{3}{2}}(a+bx)} dx, x, \log(cx^n)\right)}{n} \\
 &= -\frac{2}{bn \sqrt{\tanh(a + b \log(cx^n))}} + \frac{\text{Subst}\left(\int \sqrt{\tanh(a + bx)} dx, x, \log(cx^n)\right)}{n} \\
 &= -\frac{2}{bn \sqrt{\tanh(a + b \log(cx^n))}} - \frac{\text{Subst}\left(\int \frac{\sqrt{x}}{-1+x^2} dx, x, \tanh(a + b \log(cx^n))\right)}{bn} \\
 &= -\frac{2}{bn \sqrt{\tanh(a + b \log(cx^n))}} - \frac{2 \text{Subst}\left(\int \frac{x^2}{-1+x^4} dx, x, \sqrt{\tanh(a + b \log(cx^n))}\right)}{bn} \\
 &= -\frac{2}{bn \sqrt{\tanh(a + b \log(cx^n))}} + \frac{\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \sqrt{\tanh(a + b \log(cx^n))}\right)}{bn} \\
 &= -\frac{\tan^{-1}\left(\sqrt{\tanh(a + b \log(cx^n))}\right)}{bn} + \frac{\tan^{-1}\left(\sqrt{\tanh(a + b \log(cx^n))}\right)}{bn}
 \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.12, size = 44, normalized size = 0.62

$$\frac{{}_2F_1\left(-\frac{1}{4}, 1; \frac{3}{4}; \tanh^2(a + b \log(cx^n))\right)}{bn \sqrt{\tanh(a + b \log(cx^n))}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(x*Tanh[a + b*Log[c*x^n]]^(3/2)),x]

[Out] (-2*Hypergeometric2F1[-1/4, 1, 3/4, Tanh[a + b*Log[c*x^n]]^2])/(b*n*Sqrt[Tanh[a + b*Log[c*x^n]]])

Maple [A]

time = 4.12, size = 76, normalized size = 1.07

method	result
derivativedivides	$-\frac{\ln\left(\sqrt{\tanh(a+b\ln(cx^n))-1}\right)}{2} - \frac{2}{\sqrt{\tanh(a+b\ln(cx^n))}} + \frac{\ln\left(\sqrt{\tanh(a+b\ln(cx^n))+1}\right)}{2} - \arctan\left(\sqrt{\tanh(a+b\ln(cx^n))}\right)$
default	$-\frac{\ln\left(\sqrt{\tanh(a+b\ln(cx^n))-1}\right)}{2} - \frac{2}{\sqrt{\tanh(a+b\ln(cx^n))}} + \frac{\ln\left(\sqrt{\tanh(a+b\ln(cx^n))+1}\right)}{2} - \arctan\left(\sqrt{\tanh(a+b\ln(cx^n))}\right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/x/tanh(a+b*ln(c*x^n))^(3/2),x,method=_RETURNVERBOSE)

[Out] 1/n/b*(-1/2*ln(tanh(a+b*ln(c*x^n))^(1/2)-1)-2/tanh(a+b*ln(c*x^n))^(1/2)+1/2*ln(tanh(a+b*ln(c*x^n))^(1/2)+1)-arctan(tanh(a+b*ln(c*x^n))^(1/2)))

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/x/tanh(a+b*log(c*x^n))^(3/2),x, algorithm="maxima")

[Out] integrate(1/(x*tanh(b*log(c*x^n) + a)^(3/2)), x)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 625 vs. 2(65) = 130.

time = 0.36, size = 625, normalized size = 8.80

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/x/tanh(a+b*log(c*x^n))^(3/2),x, algorithm="fricas")

[Out]
$$-1/2*(2*(\cosh(b*n*\log(x) + b*\log(c) + a)^2 + 2*\cosh(b*n*\log(x) + b*\log(c) + a)*\sinh(b*n*\log(x) + b*\log(c) + a) + \sinh(b*n*\log(x) + b*\log(c) + a)^2 - 1)*\arctan(-\cosh(b*n*\log(x) + b*\log(c) + a)^2 - 2*\cosh(b*n*\log(x) + b*\log(c) + a)*\sinh(b*n*\log(x) + b*\log(c) + a) - \sinh(b*n*\log(x) + b*\log(c) + a)^2 + (\cosh(b*n*\log(x) + b*\log(c) + a)^2 + 2*\cosh(b*n*\log(x) + b*\log(c) + a)*\sinh(b*n*\log(x) + b*\log(c) + a) + \sinh(b*n*\log(x) + b*\log(c) + a)^2 + 1)*\sqrt{\sinh(b*n*\log(x) + b*\log(c) + a)/\cosh(b*n*\log(x) + b*\log(c) + a)}) + 4*\cosh(b*n*\log(x) + b*\log(c) + a)^2 + (\cosh(b*n*\log(x) + b*\log(c) + a)^2 + 2*\cosh(b*n*\log(x) + b*\log(c) + a)*\sinh(b*n*\log(x) + b*\log(c) + a) + \sinh(b*n*\log(x) + b*\log(c) + a)^2 - 1)*\log(-\cosh(b*n*\log(x) + b*\log(c) + a)^2 - 2*\cosh(b*n*\log(x) + b*\log(c) + a)*\sinh(b*n*\log(x) + b*\log(c) + a) - \sinh(b*n*\log(x) + b*\log(c) + a)^2 + (\cosh(b*n*\log(x) + b*\log(c) + a)^2 + 2*\cosh(b*n*\log(x) + b*\log(c) + a)*\sinh(b*n*\log(x) + b*\log(c) + a) + \sinh(b*n*\log(x) + b*\log(c) + a)^2 + 1)*\sqrt{\sinh(b*n*\log(x) + b*\log(c) + a)/\cosh(b*n*\log(x) + b*\log(c) + a)}) + 8*\cosh(b*n*\log(x) + b*\log(c) + a)*\sinh(b*n*\log(x) + b*\log(c) + a) + 4*\sinh(b*n*\log(x) + b*\log(c) + a)^2 + 4*(\cosh(b*n*\log(x) + b*\log(c) + a)^2 + 2*\cosh(b*n*\log(x) + b*\log(c) + a)*\sinh(b*n*\log(x) + b*\log(c) + a) + \sinh(b*n*\log(x) + b*\log(c) + a)^2 + 1)*\sqrt{\sinh(b*n*\log(x) + b*\log(c) + a)/\cosh(b*n*\log(x) + b*\log(c) + a)} - 4)/(b*n*\cosh(b*n*\log(x) + b*\log(c) + a)^2 + 2*b*n*\cosh(b*n*\log(x) + b*\log(c) + a)*\sinh(b*n*\log(x) + b*\log(c) + a) + b*n*\sinh(b*n*\log(x) + b*\log(c) + a)^2 - b*n)$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{x \tanh^{\frac{3}{2}}(a + b \log(cx^n))} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/x/tanh(a+b*ln(c*x**n))**(3/2),x)

[Out] Integral(1/(x*tanh(a + b*log(c*x**n))**(3/2)), x)

Giac [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/x/tanh(a+b*log(c*x^n))^(3/2),x, algorithm="giac")

[Out] Timed out

Mupad [B]

time = 1.72, size = 65, normalized size = 0.92

$$\frac{\operatorname{atanh}\left(\sqrt{\tanh(a + b \ln(cx^n))}\right)}{bn} - \frac{\operatorname{atan}\left(\sqrt{\tanh(a + b \ln(cx^n))}\right)}{bn} - \frac{2}{bn \sqrt{\tanh(a + b \ln(cx^n))}}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/(x*tanh(a + b*log(c*x^n))^(3/2)),x)`

```
[Out] atanh(tanh(a + b*log(c*x^n))^(1/2))/(b*n) - atan(tanh(a + b*log(c*x^n))^(1/2))/(b*n) - 2/(b*n*tanh(a + b*log(c*x^n))^(1/2))
```

$$3.199 \quad \int \frac{1}{x \tanh^{\frac{5}{2}}(a+b \log(cx^n))} dx$$

Optimal. Leaf size=72

$$\frac{\text{ArcTan}\left(\sqrt{\tanh(a+b \log(cx^n))}\right)}{bn} + \frac{\tanh^{-1}\left(\sqrt{\tanh(a+b \log(cx^n))}\right)}{bn} - \frac{2}{3bn \tanh^{\frac{3}{2}}(a+b \log(cx^n))}$$

[Out] arctan(tanh(a+b*ln(c*x^n))^(1/2))/b/n+arctanh(tanh(a+b*ln(c*x^n))^(1/2))/b/n-2/3/b/n/tanh(a+b*ln(c*x^n))^(3/2)

Rubi [A]

time = 0.04, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$, Rules used = {3555, 3557, 335, 218, 212, 209}

$$\frac{\text{ArcTan}\left(\sqrt{\tanh(a+b \log(cx^n))}\right)}{bn} + \frac{\tanh^{-1}\left(\sqrt{\tanh(a+b \log(cx^n))}\right)}{bn} - \frac{2}{3bn \tanh^{\frac{3}{2}}(a+b \log(cx^n))}$$

Antiderivative was successfully verified.

[In] Int[1/(x*Tanh[a + b*Log[c*x^n]]^(5/2)),x]

[Out] ArcTan[Sqrt[Tanh[a + b*Log[c*x^n]]]]/(b*n) + ArcTanh[Sqrt[Tanh[a + b*Log[c*x^n]]]]/(b*n) - 2/(3*b*n*Tanh[a + b*Log[c*x^n]]^(3/2))

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 218

Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 335

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
  Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
  ))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 3555

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(b*Tan[c + d*x]
)^n/(b*d*(n + 1)), x] - Dist[1/b^2, Int[(b*Tan[c + d*x])^(n + 2), x],
x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1]
```

Rule 3557

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[b/d, Subst[Int[
x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !
IntegerQ[n]
```

Rubi steps

$$\begin{aligned}
\int \frac{1}{x \tanh^{\frac{5}{2}}(a + b \log(cx^n))} dx &= \frac{\text{Subst}\left(\int \frac{1}{\tanh^{\frac{5}{2}}(a+bx)} dx, x, \log(cx^n)\right)}{n} \\
&= -\frac{2}{3bn \tanh^{\frac{3}{2}}(a + b \log(cx^n))} + \frac{\text{Subst}\left(\int \frac{1}{\sqrt{\tanh(a + bx)}} dx, x, \log(cx^n)\right)}{n} \\
&= -\frac{2}{3bn \tanh^{\frac{3}{2}}(a + b \log(cx^n))} - \frac{\text{Subst}\left(\int \frac{1}{\sqrt{x}(-1+x^2)} dx, x, \tanh(a + b \log(cx^n))\right)}{bn} \\
&= -\frac{2}{3bn \tanh^{\frac{3}{2}}(a + b \log(cx^n))} - \frac{2\text{Subst}\left(\int \frac{1}{-1+x^4} dx, x, \sqrt{\tanh(a + b \log(cx^n))}\right)}{bn} \\
&= -\frac{2}{3bn \tanh^{\frac{3}{2}}(a + b \log(cx^n))} + \frac{\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \sqrt{\tanh(a + b \log(cx^n))}\right)}{bn} \\
&= \frac{\tan^{-1}\left(\sqrt{\tanh(a + b \log(cx^n))}\right)}{bn} + \frac{\tanh^{-1}\left(\sqrt{\tanh(a + b \log(cx^n))}\right)}{bn}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.15, size = 46, normalized size = 0.64

$$-\frac{{}_2F_1\left(-\frac{3}{4}, 1; \frac{1}{4}; \tanh^2(a + b \log(cx^n))\right)}{3bn \tanh^{\frac{3}{2}}(a + b \log(cx^n))}$$

Antiderivative was successfully verified.

[In] Integrate[1/(x*Tanh[a + b*Log[c*x^n]]^(5/2)),x]

[Out] (-2*Hypergeometric2F1[-3/4, 1, 1/4, Tanh[a + b*Log[c*x^n]]^2])/(3*b*n*Tanh[a + b*Log[c*x^n]]^(3/2))

Maple [A]

time = 4.20, size = 74, normalized size = 1.03

method	result
derivativedivides	$-\frac{2}{3 \tanh(a+b \ln(cx^n))^{\frac{3}{2}}} + \frac{\ln\left(\sqrt{\tanh(a+b \ln(cx^n))+1}\right)}{2} - \frac{\ln\left(\sqrt{\tanh(a+b \ln(cx^n))-1}\right)}{2} + \arctan\left(\sqrt{\tanh(a+b \ln(cx^n))}\right)$
default	$-\frac{2}{3 \tanh(a+b \ln(cx^n))^{\frac{3}{2}}} + \frac{\ln\left(\sqrt{\tanh(a+b \ln(cx^n))+1}\right)}{2} - \frac{\ln\left(\sqrt{\tanh(a+b \ln(cx^n))-1}\right)}{2} + \arctan\left(\sqrt{\tanh(a+b \ln(cx^n))}\right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/x/tanh(a+b*ln(c*x^n))^(5/2),x,method=_RETURNVERBOSE)

[Out] 1/n/b*(-2/3/tanh(a+b*ln(c*x^n))^(3/2)+1/2*ln(tanh(a+b*ln(c*x^n))^(1/2)+1)-1/2*ln(tanh(a+b*ln(c*x^n))^(1/2)-1)+arctan(tanh(a+b*ln(c*x^n))^(1/2)))

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/x/tanh(a+b*log(c*x^n))^(5/2),x, algorithm="maxima")

[Out] integrate(1/(x*tanh(b*log(c*x^n) + a)^(5/2)), x)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 1110 vs. 2(64) = 128.

time = 0.37, size = 1110, normalized size = 15.42

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/x/tanh(a+b*log(c*x^n))^(5/2),x, algorithm="fricas")

[Out]
$$\begin{aligned} & -1/6*(4*\cosh(b*n*\log(x) + b*\log(c) + a)^4 + 16*\cosh(b*n*\log(x) + b*\log(c) + a)*\sinh(b*n*\log(x) + b*\log(c) + a)^3 + 4*\sinh(b*n*\log(x) + b*\log(c) + a)^4 \\ & + 8*(3*\cosh(b*n*\log(x) + b*\log(c) + a)^2 - 1)*\sinh(b*n*\log(x) + b*\log(c) + a)^2 - 6*(\cosh(b*n*\log(x) + b*\log(c) + a)^4 + 4*\cosh(b*n*\log(x) + b*\log(c) + a)*\sinh(b*n*\log(x) + b*\log(c) + a)^3 + \sinh(b*n*\log(x) + b*\log(c) + a)^4 \\ & + 2*(3*\cosh(b*n*\log(x) + b*\log(c) + a)^2 - 1)*\sinh(b*n*\log(x) + b*\log(c) + a)^2 - 2*\cosh(b*n*\log(x) + b*\log(c) + a)^2 + 4*(\cosh(b*n*\log(x) + b*\log(c) + a)^3 - \cosh(b*n*\log(x) + b*\log(c) + a))*\sinh(b*n*\log(x) + b*\log(c) + a) \\ & + 1)*\arctan(-\cosh(b*n*\log(x) + b*\log(c) + a)^2 - 2*\cosh(b*n*\log(x) + b*\log(c) + a)*\sinh(b*n*\log(x) + b*\log(c) + a) - \sinh(b*n*\log(x) + b*\log(c) + a)^2 \\ & + (\cosh(b*n*\log(x) + b*\log(c) + a)^2 + 2*\cosh(b*n*\log(x) + b*\log(c) + a)*\sinh(b*n*\log(x) + b*\log(c) + a) + \sinh(b*n*\log(x) + b*\log(c) + a)^2 + 1)*\sqrt{\sinh(b*n*\log(x) + b*\log(c) + a)/\cosh(b*n*\log(x) + b*\log(c) + a)}} - 8*\cosh(b*n*\log(x) + b*\log(c) + a)^2 + 3*(\cosh(b*n*\log(x) + b*\log(c) + a)^4 + 4*\cosh(b*n*\log(x) + b*\log(c) + a)*\sinh(b*n*\log(x) + b*\log(c) + a)^3 + \sinh(b*n*\log(x) + b*\log(c) + a)^4 + 2*(3*\cosh(b*n*\log(x) + b*\log(c) + a)^2 - 1)*\sinh(b*n*\log(x) + b*\log(c) + a)^2 - 2*\cosh(b*n*\log(x) + b*\log(c) + a)^2 + 4*(\cosh(b*n*\log(x) + b*\log(c) + a)^3 - \cosh(b*n*\log(x) + b*\log(c) + a))*\sinh(b*n*\log(x) + b*\log(c) + a) + 1)*\log(-\cosh(b*n*\log(x) + b*\log(c) + a)^2 - 2*\cosh(b*n*\log(x) + b*\log(c) + a)*\sinh(b*n*\log(x) + b*\log(c) + a) - \sinh(b*n*\log(x) + b*\log(c) + a)^2 + (\cosh(b*n*\log(x) + b*\log(c) + a)^2 + 2*\cosh(b*n*\log(x) + b*\log(c) + a)*\sinh(b*n*\log(x) + b*\log(c) + a) + \sinh(b*n*\log(x) + b*\log(c) + a)^2 + 1)*\sqrt{\sinh(b*n*\log(x) + b*\log(c) + a)/\cosh(b*n*\log(x) + b*\log(c) + a)}}) + 16*(\cosh(b*n*\log(x) + b*\log(c) + a)^3 - \cosh(b*n*\log(x) + b*\log(c) + a))*\sinh(b*n*\log(x) + b*\log(c) + a) + 4*(\cosh(b*n*\log(x) + b*\log(c) + a)^4 + 4*\cosh(b*n*\log(x) + b*\log(c) + a)*\sinh(b*n*\log(x) + b*\log(c) + a)^3 + \sinh(b*n*\log(x) + b*\log(c) + a)^4 + 2*(3*\cosh(b*n*\log(x) + b*\log(c) + a)^2 + 1)*\sinh(b*n*\log(x) + b*\log(c) + a)^2 + 2*\cosh(b*n*\log(x) + b*\log(c) + a)^2 + 4*(\cosh(b*n*\log(x) + b*\log(c) + a)^3 + \cosh(b*n*\log(x) + b*\log(c) + a))*\sinh(b*n*\log(x) + b*\log(c) + a) + 1)*\sqrt{\sinh(b*n*\log(x) + b*\log(c) + a)/\cosh(b*n*\log(x) + b*\log(c) + a)}} + 4)/((b*n*\cosh(b*n*\log(x) + b*\log(c) + a))^4 + 4*b*n*\cosh(b*n*\log(x) + b*\log(c) + a)*\sinh(b*n*\log(x) + b*\log(c) + a)^3 + b*n*\sinh(b*n*\log(x) + b*\log(c) + a)^4 - 2*b*n*\cosh(b*n*\log(x) + b*\log(c) + a)^2 + 2*(3*b*n*\cosh(b*n*\log(x) + b*\log(c) + a)^2 - b*n)*\sinh(b*n*\log(x) + b*\log(c) + a)^2 + b*n + 4*(b*n*\cosh(b*n*\log(x) + b*\log(c) + a)^3 - b*n*\cosh(b*n*\log(x) + b*\log(c) + a))*\sinh(b*n*\log(x) + b*\log(c) + a)) \end{aligned}$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{x \tanh^{\frac{5}{2}}(a + b \log(cx^n))} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/x/tanh(a+b*ln(c*x**n))**(5/2),x)

[Out] Integral(1/(x*tanh(a + b*log(c*x**n))**(5/2)), x)

Giac [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/x/tanh(a+b*log(c*x^n))^(5/2),x, algorithm="giac")

[Out] Timed out

Mupad [B]

time = 2.48, size = 64, normalized size = 0.89

$$\frac{\operatorname{atan}\left(\sqrt{\tanh(a + b \ln(cx^n))}\right)}{bn} + \frac{\operatorname{atanh}\left(\sqrt{\tanh(a + b \ln(cx^n))}\right)}{bn} - \frac{2}{3bn \tanh(a + b \ln(cx^n))^{3/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(x*tanh(a + b*log(c*x^n))^(5/2)),x)

[Out] atan(tanh(a + b*log(c*x^n))^(1/2))/(b*n) + atanh(tanh(a + b*log(c*x^n))^(1/2))/(b*n) - 2/(3*b*n*tanh(a + b*log(c*x^n))^(3/2))

$$3.200 \quad \int \frac{\tanh^5(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx$$

Optimal. Leaf size=135

$$\frac{(b-2c) \tanh^{-1} \left(\frac{b+2c \tanh^2(x)}{2\sqrt{c} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} \right)}{4c^{3/2}} + \frac{\tanh^{-1} \left(\frac{2a+b+(b+2c) \tanh^2(x)}{2\sqrt{a+b+c} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} \right)}{2\sqrt{a+b+c}}$$

[Out] $1/4*(b-2*c)*\operatorname{arctanh}(1/2*(b+2*c*\tanh(x)^2)/c^{(1/2)}/(a+b*\tanh(x)^2+c*\tanh(x)^4)^{(1/2)})/c^{(3/2)}+1/2*\operatorname{arctanh}(1/2*(2*a+b+(b+2*c)*\tanh(x)^2)/(a+b+c)^{(1/2)}/(a+b*\tanh(x)^2+c*\tanh(x)^4)^{(1/2)})/(a+b+c)^{(1/2)}-1/2*(a+b*\tanh(x)^2+c*\tanh(x)^4)^{(1/2)}/c$

Rubi [A]

time = 0.25, antiderivative size = 135, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.304$, Rules used = {3781, 1265, 1667, 857, 635, 212, 738}

$$\frac{(b-2c) \tanh^{-1} \left(\frac{b+2c \tanh^2(x)}{2\sqrt{c} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} \right)}{4c^{3/2}} - \frac{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}}{2c} + \frac{\tanh^{-1} \left(\frac{2a+(b+2c) \tanh^2(x)+b}{2\sqrt{a+b+c} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} \right)}{2\sqrt{a+b+c}}$$

Antiderivative was successfully verified.

[In] `Int[Tanh[x]^5/Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4], x]`

[Out] $((b-2*c)*\operatorname{ArcTanh}[(b+2*c*\operatorname{Tanh}[x]^2)/(2*\operatorname{Sqrt}[c]*\operatorname{Sqrt}[a+b*\operatorname{Tanh}[x]^2+c*\operatorname{Tanh}[x]^4]))/(4*c^{(3/2)}) + \operatorname{ArcTanh}[(2*a+b+(b+2*c)*\operatorname{Tanh}[x]^2)/(2*\operatorname{Sqrt}[a+b+c]*\operatorname{Sqrt}[a+b*\operatorname{Tanh}[x]^2+c*\operatorname{Tanh}[x]^4]])/(2*\operatorname{Sqrt}[a+b+c]) - \operatorname{Sqrt}[a+b*\operatorname{Tanh}[x]^2+c*\operatorname{Tanh}[x]^4]/(2*c)$

Rule 212

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rule 635

`Int[1/Sqrt[(a_) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] := Dist[2, Subst[Int[1/(4*c - x^2), x], x, (b + 2*c*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]`

Rule 738


```
Int[1/(((d_.) + (e_.)*(x_))*Sqrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]), x_Symbol]
:= Dist[-2, Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e}, x]
&& NeQ[b^2 - 4*a*c, 0] && NeQ[2*c*d - b*e, 0]
```

Rule 857

```
Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol]
:= Dist[g/e, Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^p, x], x] + Dist[(e*f - d*g)/e, Int[(d + e*x)^m*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, p}, x]
&& NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && !IGtQ[m, 0]
```

Rule 1265

```
Int[(x_)^(m_)*((d_) + (e_.)*(x_)^2)^(q_)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_), x_Symbol]
:= Dist[1/2, Subst[Int[x^((m - 1)/2)*(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x]
&& IntegerQ[(m - 1)/2]
```

Rule 1667

```
Int[(Pq_)*((d_.) + (e_.)*(x_))^(m_)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol]
:= With[{q = Expon[Pq, x], f = Coeff[Pq, x, Expon[Pq, x]]}, Simp[f*(d + e*x)^(m + q - 1)*(a + b*x + c*x^2)^(p + 1)/(c*e^(q - 1)*(m + q + 2*p + 1)), x] + Dist[1/(c*e^q*(m + q + 2*p + 1)), Int[(d + e*x)^m*(a + b*x + c*x^2)^p*ExpandToSum[c*e^q*(m + q + 2*p + 1)*Pq - c*f*(m + q + 2*p + 1)*(d + e*x)^q - f*(d + e*x)^(q - 2)*(b*d*e*(p + 1) + a*e^2*(m + q - 1) - c*d^2*(m + q + 2*p + 1) - e*(2*c*d - b*e)*(m + q + p)*x), x], x] /; GtQ[q, 1]
&& NeQ[m + q + 2*p + 1, 0] /; FreeQ[{a, b, c, d, e, m, p}, x] && PolyQ[Pq, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && !(IGtQ[m, 0] && RationalQ[a, b, c, d, e] && (IntegerQ[p] || ILtQ[p + 1/2, 0]))
```

Rule 3781

```
Int[tan[(d_.) + (e_.)*(x_)]^(m_)*((a_.) + (b_.))*((f_.)*tan[(d_.) + (e_.)*(x_)])^(n_) + (c_.)*((f_.)*tan[(d_.) + (e_.)*(x_)])^(n2_)^(p_), x_Symbol]
:= Dist[f/e, Subst[Int[(x/f)^m*((a + b*x^n + c*x^(2*n))^p/(f^2 + x^2)), x], x, f*Tan[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[n2, 2*n] && NeQ[b^2 - 4*a*c, 0]
```

Rubi steps

$$\begin{aligned}
\int \frac{\tanh^5(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx &= -\text{Subst} \left(\int \frac{x^5}{(1+x^2) \sqrt{a - bx^2 + cx^4}} dx, x, i \tanh(x) \right) \\
&= - \left(\frac{1}{2} \text{Subst} \left(\int \frac{x^2}{(1+x) \sqrt{a - bx + cx^2}} dx, x, -\tanh^2(x) \right) \right) \\
&= - \frac{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}}{2c} - \frac{\text{Subst} \left(\int \frac{\frac{b}{2} + \frac{1}{2}(b-2c)x}{(1+x) \sqrt{a - bx + cx^2}} dx, x, -\tanh^2(x) \right)}{2c} \\
&= - \frac{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}}{2c} - \frac{1}{2} \text{Subst} \left(\int \frac{1}{(1+x) \sqrt{a - bx + cx^2}} dx, x, -\tanh^2(x) \right) \\
&= - \frac{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}}{2c} - \frac{(b-2c) \text{Subst} \left(\int \frac{1}{4c-x^2} dx, x, -\tanh^2(x) \right)}{2c} \\
&= \frac{(b-2c) \tanh^{-1} \left(\frac{b+2c \tanh^2(x)}{2\sqrt{c} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} \right)}{4c^{3/2}} + \frac{\tanh^{-1} \left(\frac{b-2c \tanh^2(x)}{2\sqrt{c} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} \right)}{4c^{3/2}}
\end{aligned}$$

Mathematica [A]

time = 0.96, size = 136, normalized size = 1.01

$$\frac{1}{4} \left(\frac{(-b+2c) \tanh^{-1} \left(\frac{-b-2c \tanh^2(x)}{2\sqrt{c} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} \right)}{c^{3/2}} + \frac{2 \tanh^{-1} \left(\frac{2a+b+(b+2c) \tanh^2(x)}{2\sqrt{a+b+c} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} \right)}{\sqrt{a+b+c}} - \frac{2\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}}{c} \right)$$

Antiderivative was successfully verified.

`[In] Integrate[Tanh[x]^5/Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4], x]`

```
[Out] (((-b + 2*c)*ArcTanh[(-b - 2*c*Tanh[x]^2)/(2*Sqrt[c]*Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4]])/c^(3/2) + (2*ArcTanh[(2*a + b + (b + 2*c)*Tanh[x]^2)/(2*Sqrt[a + b + c]*Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4]])/Sqrt[a + b + c] - (2*Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4])/c)/4
```

Maple [A]

time = 1.22, size = 149, normalized size = 1.10

method	result
derivativedivides	$-\frac{\sqrt{a + b (\tanh^2(x)) + c (\tanh^4(x))}}{2c} + \frac{b \ln\left(\frac{\frac{b}{2} + c (\tanh^2(x))}{\sqrt{c}}\right) + \sqrt{a + b (\tanh^2(x)) + c (\tanh^4(x))}}{4c^{\frac{3}{2}}}$
default	$-\frac{\sqrt{a + b (\tanh^2(x)) + c (\tanh^4(x))}}{2c} + \frac{b \ln\left(\frac{\frac{b}{2} + c (\tanh^2(x))}{\sqrt{c}}\right) + \sqrt{a + b (\tanh^2(x)) + c (\tanh^4(x))}}{4c^{\frac{3}{2}}}$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(tanh(x)^5/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x,method=_RETURNVERBOSE)
[Out] -1/2*(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2)/c+1/4*b/c^(3/2)*ln((1/2*b+c*tanh(x)^2)/c^(1/2)+(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2))-1/2*ln((1/2*b+c*tanh(x)^2)/c^(1/2)+(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2))/c^(1/2)+1/2/(a+b+c)^(1/2)*arctanh(1/2*(b*tanh(x)^2+2*c*tanh(x)^2+2*a+b)/(a+b+c)^(1/2)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2))
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(tanh(x)^5/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x, algorithm="maxima")
[Out] integrate(tanh(x)^5/sqrt(c*tanh(x)^4 + b*tanh(x)^2 + a), x)
```

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 2075 vs. 2(111) = 222.

time = 1.20, size = 8891, normalized size = 65.86

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(tanh(x)^5/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x, algorithm="fricas")
[Out] [-1/8*(((a*b + b^2 - (2*a + b)*c - 2*c^2)*cosh(x)^4 + 4*(a*b + b^2 - (2*a + b)*c - 2*c^2)*cosh(x)*sinh(x)^3 + (a*b + b^2 - (2*a + b)*c - 2*c^2)*sinh(x)^4 + 2*(a*b + b^2 - (2*a + b)*c - 2*c^2)*cosh(x)^2 + 2*(3*(a*b + b^2 - (2*a + b)*c - 2*c^2)*sinh(x)^2 - (a*b + b^2 - (2*a + b)*c - 2*c^2)*sinh(x)^2))^(1/2) + ...]
```

$$\begin{aligned}
& a + b) * c - 2 * c^2) * \cosh(x)^2 + a * b + b^2 - (2 * a + b) * c - 2 * c^2) * \sinh(x)^2 + \\
& a * b + b^2 - (2 * a + b) * c - 2 * c^2 + 4 * ((a * b + b^2 - (2 * a + b) * c - 2 * c^2) * \cosh(x)^3 + (a * b + b^2 - (2 * a + b) * c - 2 * c^2) * \cosh(x)) * \sinh(x)) * \sqrt{c} * \log(((b \\
& ^2 + 4 * (a + 2 * b) * c + 8 * c^2) * \cosh(x)^8 + 8 * (b^2 + 4 * (a + 2 * b) * c + 8 * c^2) * \cosh(x) * \sinh(x))^7 + (b^2 + 4 * (a + 2 * b) * c + 8 * c^2) * \sinh(x))^8 + 4 * (b^2 + 4 * a * c - \\
& 8 * c^2) * \cosh(x)^6 + 4 * (7 * (b^2 + 4 * (a + 2 * b) * c + 8 * c^2) * \cosh(x)^2 + b^2 + 4 * \\
& a * c - 8 * c^2) * \sinh(x)^6 + 8 * (7 * (b^2 + 4 * (a + 2 * b) * c + 8 * c^2) * \cosh(x)^3 + 3 * (\\
& b^2 + 4 * a * c - 8 * c^2) * \cosh(x)) * \sinh(x))^5 + 2 * (3 * b^2 + 4 * (3 * a - 2 * b) * c + 24 * c \\
& ^2) * \cosh(x)^4 + 2 * (35 * (b^2 + 4 * (a + 2 * b) * c + 8 * c^2) * \cosh(x)^4 + 30 * (b^2 + 4 \\
& * a * c - 8 * c^2) * \cosh(x)^2 + 3 * b^2 + 4 * (3 * a - 2 * b) * c + 24 * c^2) * \sinh(x)^4 + 8 * (\\
& 7 * (b^2 + 4 * (a + 2 * b) * c + 8 * c^2) * \cosh(x)^5 + 10 * (b^2 + 4 * a * c - 8 * c^2) * \cosh(x) \\
&)^3 + (3 * b^2 + 4 * (3 * a - 2 * b) * c + 24 * c^2) * \cosh(x)) * \sinh(x))^3 + 4 * (b^2 + 4 * a * \\
& c - 8 * c^2) * \cosh(x)^2 + 4 * (7 * (b^2 + 4 * (a + 2 * b) * c + 8 * c^2) * \cosh(x)^6 + 15 * (b \\
& ^2 + 4 * a * c - 8 * c^2) * \cosh(x)^4 + 3 * (3 * b^2 + 4 * (3 * a - 2 * b) * c + 24 * c^2) * \cosh(x) \\
&)^2 + b^2 + 4 * a * c - 8 * c^2) * \sinh(x)^2 - 4 * \sqrt{2} * ((b + 2 * c) * \cosh(x)^4 + 4 * (\\
& b + 2 * c) * \cosh(x) * \sinh(x))^3 + (b + 2 * c) * \sinh(x)^4 + 2 * (b - 2 * c) * \cosh(x)^2 + \\
& 2 * (3 * (b + 2 * c) * \cosh(x)^2 + b - 2 * c) * \sinh(x)^2 + 4 * ((b + 2 * c) * \cosh(x))^3 + (b \\
& - 2 * c) * \cosh(x)) * \sinh(x) + b + 2 * c) * \sqrt{c} * \sqrt{((a + b + c) * \cosh(x)^4 + (\\
& a + b + c) * \sinh(x)^4 + 4 * (a - c) * \cosh(x)^2 + 2 * (3 * (a + b + c) * \cosh(x)^2 + 2 \\
& * a - 2 * c) * \sinh(x)^2 + 3 * a - b + 3 * c) / (\cosh(x)^4 - 4 * \cosh(x)^3 * \sinh(x) + 6 * c \\
& \cosh(x)^2 * \sinh(x)^2 - 4 * \cosh(x) * \sinh(x)^3 + \sinh(x)^4)) + b^2 + 4 * (a + 2 * b) * \\
& c + 8 * c^2 + 8 * ((b^2 + 4 * (a + 2 * b) * c + 8 * c^2) * \cosh(x)^7 + 3 * (b^2 + 4 * a * c - 8 \\
& * c^2) * \cosh(x)^5 + (3 * b^2 + 4 * (3 * a - 2 * b) * c + 24 * c^2) * \cosh(x)^3 + (b^2 + 4 * a \\
& * c - 8 * c^2) * \cosh(x)) * \sinh(x)) / (\cosh(x)^8 + 8 * \cosh(x) * \sinh(x))^7 + \sinh(x))^8 \\
& + 4 * (7 * \cosh(x)^2 + 1) * \sinh(x)^6 + 4 * \cosh(x)^6 + 8 * (7 * \cosh(x)^3 + 3 * \cosh(x)) \\
& * \sinh(x)^5 + 2 * (35 * \cosh(x)^4 + 30 * \cosh(x)^2 + 3) * \sinh(x)^4 + 6 * \cosh(x)^4 + \\
& 8 * (7 * \cosh(x)^5 + 10 * \cosh(x)^3 + 3 * \cosh(x)) * \sinh(x)^3 + 4 * (7 * \cosh(x)^6 + 15 * \\
& \cosh(x)^4 + 9 * \cosh(x)^2 + 1) * \sinh(x)^2 + 4 * \cosh(x)^2 + 8 * (\cosh(x)^7 + 3 * \cosh(x) \\
& ^5 + 3 * \cosh(x)^3 + \cosh(x)) * \sinh(x) + 1)) - 2 * (c^2 * \cosh(x)^4 + 4 * c^2 * \cosh(x) * \sinh(x))^3 + c^2 * \sinh(x)^4 + 2 * c^2 * \cosh(x)^2 + 2 * (3 * c^2 * \cosh(x)^2 + c^2 \\
& ^2) * \sinh(x)^2 + c^2 + 4 * (c^2 * \cosh(x)^3 + c^2 * \cosh(x)) * \sinh(x)) * \sqrt{a + b + \\
& c} * \log(((a^2 + 2 * a * b + b^2 + 2 * (a + b) * c + c^2) * \cosh(x))^8 + 8 * (a^2 + 2 * a * b \\
& + b^2 + 2 * (a + b) * c + c^2) * \cosh(x) * \sinh(x))^7 + (a^2 + 2 * a * b + b^2 + 2 * (a + \\
& b) * c + c^2) * \sinh(x))^8 + 4 * (a^2 + a * b - b * c - c^2) * \cosh(x)^6 + 4 * (7 * (a^2 + 2 \\
& * a * b + b^2 + 2 * (a + b) * c + c^2) * \cosh(x)^2 + a^2 + a * b - b * c - c^2) * \sinh(x)^6 \\
& + 8 * (7 * (a^2 + 2 * a * b + b^2 + 2 * (a + b) * c + c^2) * \cosh(x)^3 + 3 * (a^2 + a * b - \\
& b * c - c^2) * \cosh(x)) * \sinh(x))^5 + 2 * (3 * a^2 + 2 * a * b + 2 * (a + b) * c + 3 * c^2) * \cosh(x)^4 + 2 * (35 * (a^2 + 2 * a * b + b^2 + 2 * (a + b) * c + c^2) * \cosh(x)^4 + 30 * (a^2 \\
& + a * b - b * c - c^2) * \cosh(x)^2 + 3 * a^2 + 2 * a * b + 2 * (a + b) * c + 3 * c^2) * \sinh(x) \\
&)^4 + 8 * (7 * (a^2 + 2 * a * b + b^2 + 2 * (a + b) * c + c^2) * \cosh(x)^5 + 10 * (a^2 + a * \\
& b - b * c - c^2) * \cosh(x)^3 + (3 * a^2 + 2 * a * b + 2 * (a + b) * c + 3 * c^2) * \cosh(x)) * \sinh(x))^3 + 4 * (a^2 + a * b - b * c - c^2) * \cosh(x)^2 + 4 * (7 * (a^2 + 2 * a * b + b^2 + \\
& 2 * (a + b) * c + c^2) * \cosh(x)^6 + 15 * (a^2 + a * b - b * c - c^2) * \cosh(x)^4 + 3 * (3 * \\
& a^2 + 2 * a * b + 2 * (a + b) * c + 3 * c^2) * \cosh(x)^2 + a^2 + a * b - b * c - c^2) * \sinh(x) \\
&)^2 + \sqrt{2} * ((a + b + c) * \cosh(x))^4 + 4 * (a + b + c) * \cosh(x) * \sinh(x))^3 + (
\end{aligned}$$

$$\begin{aligned}
& a + b + c) \sinh(x)^4 + 2(a - c) \cosh(x)^2 + 2(3(a + b + c) \cosh(x)^2 + a \\
& - c) \sinh(x)^2 + 4((a + b + c) \cosh(x)^3 + (a - c) \cosh(x)) \sinh(x) + a + \\
& b + c) \sqrt{a + b + c} \sqrt{((a + b + c) \cosh(x)^4 + (a + b + c) \sinh(x)^4 \\
& + 4(a - c) \cosh(x)^2 + 2(3(a + b + c) \cosh(x)^2 + 2a - 2c) \sinh(x)^2 \\
& + 3a - b + 3c) / (\cosh(x)^4 - 4 \cosh(x)^3 \sinh(x) + 6 \cosh(x)^2 \sinh(x)^2 - \\
& 4 \cosh(x) \sinh(x)^3 + \sinh(x)^4)} + a^2 + 2ab + b^2 + 2(a + b)c + c^2 \\
& + 8((a^2 + 2ab + b^2 + 2(a + b)c + c^2) \cosh(x)^7 + 3(a^2 + ab - bc \\
& - c^2) \cosh(x)^5 + (3a^2 + 2ab + 2(a + b)c + 3c^2) \cosh(x)^3 + (a^2 \\
& + ab - bc - c^2) \cosh(x) \sinh(x)) / (\cosh(x)^4 + 4 \cosh(x)^3 \sinh(x) + 6 \cosh(x)^2 \sinh(x)^2 + 4 \cosh(x) \sinh(x)^3 + \sinh(x)^4) + 4 \sqrt{2} ((a + b) \\
& * c + c^2) \sqrt{((a + b + c) \cosh(x)^4 + (a + b + c) \sinh(x)^4 + 4(a - c) \cosh(x)^2 + 2(3(a + b + c) \cosh(x)^2 + 2a - 2c) \sinh(x)^2 + 3a - b + 3c) / (\cosh(x)^4 - 4 \cosh(x)^3 \sinh(x) + 6 \cosh(x)^2 \sinh(x)^2 - 4 \cosh(x) \sinh(x)^3 + \sinh(x)^4))} / (((a + b) * c^2 + c^3) \cosh(x)^4 + 4((a + b) * c^2 + c^3) \cosh(x) \sinh(x)^3 + ((a + b) * c^2 + c^3) \sinh(x)^4 + (a + b) * c^2 + c^3 + 2 * ((a + b) * c^2 + c^3) \cosh(x)^2 + 2 * ((a + b) * c^2 + c^3 + 3 * ((a + b) * c^2 + c^3) \cosh(x)^2) \sinh(x)^2 + 4 * (((a + b) * c^2 + c^3) \cosh(x)^3 + ((a + b) * c^2 + c^3) \cosh(x) \sinh(x))), -1/8 * (4 * (c^2 \cosh(x)^4 + 4 * c^2 \cosh(x) \sinh(x)^3 + c^2 \sinh(x)^4 + 2 * c^2 \cosh(x)^2 + 2 * (3 * c^2 \cos...
\end{aligned}$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\tanh^5(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)**5/(a+b*tanh(x)**2+c*tanh(x)**4)**(1/2),x)

[Out] Integral(tanh(x)**5/sqrt(a + b*tanh(x)**2 + c*tanh(x)**4), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)^5/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x, algorithm="giac")

[Out] integrate(tanh(x)^5/sqrt(c*tanh(x)^4 + b*tanh(x)^2 + a), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\tanh(x)^5}{\sqrt{c \tanh(x)^4 + b \tanh(x)^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(tanh(x)^5/(a + b*tanh(x)^2 + c*tanh(x)^4)^(1/2), x)
```

```
[Out] int(tanh(x)^5/(a + b*tanh(x)^2 + c*tanh(x)^4)^(1/2), x)
```

$$3.201 \quad \int \frac{\tanh^3(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx$$

Optimal. Leaf size=105

$$\frac{\tanh^{-1}\left(\frac{b+2c \tanh^2(x)}{2\sqrt{c} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}}\right)}{2\sqrt{c}} + \frac{\tanh^{-1}\left(\frac{2a+b+(b+2c) \tanh^2(x)}{2\sqrt{a+b+c} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}}\right)}{2\sqrt{a+b+c}}$$

[Out] $-1/2*\operatorname{arctanh}(1/2*(b+2*c*\tanh(x)^2)/c^{(1/2)}/(a+b*\tanh(x)^2+c*\tanh(x)^4)^{(1/2)})/c^{(1/2)}+1/2*\operatorname{arctanh}(1/2*(2*a+b+(b+2*c)*\tanh(x)^2)/(a+b+c)^{(1/2)}/(a+b*\tanh(x)^2+c*\tanh(x)^4)^{(1/2)})/(a+b+c)^{(1/2)}$

Rubi [A]

time = 0.14, antiderivative size = 105, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.261$, Rules used = {3781, 1265, 857, 635, 212, 738}

$$\frac{\tanh^{-1}\left(\frac{2a+(b+2c) \tanh^2(x)+b}{2\sqrt{a+b+c} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}}\right)}{2\sqrt{a+b+c}} - \frac{\tanh^{-1}\left(\frac{b+2c \tanh^2(x)}{2\sqrt{c} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}}\right)}{2\sqrt{c}}$$

Antiderivative was successfully verified.

[In] `Int[Tanh[x]^3/Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4], x]`

[Out] $-1/2*\operatorname{ArcTanh}[(b + 2*c*\operatorname{Tanh}[x]^2)/(2*\operatorname{Sqrt}[c]*\operatorname{Sqrt}[a + b*\operatorname{Tanh}[x]^2 + c*\operatorname{Tanh}[x]^4])]/\operatorname{Sqrt}[c] + \operatorname{ArcTanh}[(2*a + b + (b + 2*c)*\operatorname{Tanh}[x]^2)/(2*\operatorname{Sqrt}[a + b + c]*\operatorname{Sqrt}[a + b*\operatorname{Tanh}[x]^2 + c*\operatorname{Tanh}[x]^4])]/(2*\operatorname{Sqrt}[a + b + c])$

Rule 212

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rule 635

`Int[1/Sqrt[(a_) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] := Dist[2, Subst[Int[1/(4*c - x^2), x], x, (b + 2*c*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]`

Rule 738

`Int[1/(((d_.) + (e_.)*(x_))*Sqrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]), x_Symbol] := Dist[-2, Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2`

$*a*e - b*d - (2*c*d - b*e)*x/\text{Sqrt}[a + b*x + c*x^2]], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0] \ \&\& \ \text{NeQ}[2*c*d - b*e, 0]$

Rule 857

$\text{Int}[\{(d_.) + (e_.)*(x_)\}^{(m_)}*\{(f_.) + (g_.)*(x_)\}*\{(a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2\}^{(p_.)}, x_Symbol] \rightarrow \text{Dist}[g/e, \text{Int}[(d + e*x)^{(m + 1)}*(a + b*x + c*x^2)^p, x], x] + \text{Dist}[(e*f - d*g)/e, \text{Int}[(d + e*x)^m*(a + b*x + c*x^2)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, g, m, p\}, x] \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0] \ \&\& \ \text{NeQ}[c*d^2 - b*d*e + a*e^2, 0] \ \&\& \ !\text{IGtQ}[m, 0]$

Rule 1265

$\text{Int}[(x_)^{(m_)}*\{(d_.) + (e_.)*(x_.)^2\}^{(q_)}*\{(a_.) + (b_.)*(x_.)^2 + (c_.)*(x_.)^4\}^{(p_.)}, x_Symbol] \rightarrow \text{Dist}[1/2, \text{Subst}[\text{Int}[x^{(m - 1)/2}*(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; \text{FreeQ}[\{a, b, c, d, e, p, q\}, x] \ \&\& \ \text{IntegerQ}[(m - 1)/2]$

Rule 3781

$\text{Int}[\tan[(d_.) + (e_.)*(x_)]^{(m_)}*\{(a_.) + (b_.)*\{(f_.)*\tan[(d_.) + (e_.)*(x_)]\}^{(n_)} + (c_.)*\{(f_.)*\tan[(d_.) + (e_.)*(x_)]\}^{(2*n_)}\}^{(p_.)}, x_Symbol] \rightarrow \text{Dist}[f/e, \text{Subst}[\text{Int}[(x/f)^m*(a + b*x^n + c*x^{(2*n)})^p/(f^2 + x^2), x], x, f*\text{Tan}[d + e*x]], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m, n, p\}, x] \ \&\& \ \text{EqQ}[n2, 2*n] \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0]$

Rubi steps

$$\begin{aligned} \int \frac{\tanh^3(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx &= \text{Subst} \left(\int \frac{x^3}{(1 + x^2) \sqrt{a - bx^2 + cx^4}} dx, x, i \tanh(x) \right) \\ &= \frac{1}{2} \text{Subst} \left(\int \frac{x}{(1 + x) \sqrt{a - bx + cx^2}} dx, x, -\tanh^2(x) \right) \\ &= \frac{1}{2} \text{Subst} \left(\int \frac{1}{\sqrt{a - bx + cx^2}} dx, x, -\tanh^2(x) \right) - \frac{1}{2} \text{Subst} \left(\int \frac{1}{(1 + x) \sqrt{a - bx + cx^2}} dx, x, -\tanh^2(x) \right) \\ &= \text{Subst} \left(\int \frac{1}{4c - x^2} dx, x, \frac{-b - 2c \tanh^2(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} \right) + \text{Subst} \left(\int \frac{1}{(1 + x) \sqrt{a - bx + cx^2}} dx, x, -\tanh^2(x) \right) \\ &= \frac{\tanh^{-1} \left(\frac{-b - 2c \tanh^2(x)}{2\sqrt{c} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} \right)}{2\sqrt{c}} + \frac{\tanh^{-1} \left(\frac{-b - 2c \tanh^2(x)}{2\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} \right)}{2\sqrt{c}} \end{aligned}$$

Mathematica [A]

time = 0.05, size = 105, normalized size = 1.00

$$\frac{1}{2} \left(\frac{\tanh^{-1} \left(\frac{-b-2c \tanh^2(x)}{2\sqrt{c} \sqrt{a+b \tanh^2(x)+c \tanh^4(x)}} \right)}{\sqrt{c}} + \frac{\tanh^{-1} \left(\frac{2a+b+(b+2c) \tanh^2(x)}{2\sqrt{a+b+c} \sqrt{a+b \tanh^2(x)+c \tanh^4(x)}} \right)}{\sqrt{a+b+c}} \right)$$

Antiderivative was successfully verified.

[In] Integrate[Tanh[x]^3/Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4], x]

[Out] (ArcTanh[(-b - 2*c*Tanh[x]^2)/(2*Sqrt[c]*Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4])]/Sqrt[c] + ArcTanh[(2*a + b + (b + 2*c)*Tanh[x]^2)/(2*Sqrt[a + b + c]*Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4)])/Sqrt[a + b + c])/2

Maple [A]

time = 0.99, size = 90, normalized size = 0.86

method	result
derivativedivides	$-\frac{\ln\left(\frac{\frac{b}{2}+c(\tanh^2(x))}{\sqrt{c}}+\sqrt{a+b(\tanh^2(x))+c(\tanh^4(x))}\right)}{2\sqrt{c}}+\frac{\operatorname{arctanh}\left(\frac{b(\tanh^2(x))}{2\sqrt{a+b+c}\sqrt{a+b(\tanh^2(x))+c(\tanh^4(x))}}\right)}{2\sqrt{a+b+c}}$
default	$-\frac{\ln\left(\frac{\frac{b}{2}+c(\tanh^2(x))}{\sqrt{c}}+\sqrt{a+b(\tanh^2(x))+c(\tanh^4(x))}\right)}{2\sqrt{c}}+\frac{\operatorname{arctanh}\left(\frac{b(\tanh^2(x))}{2\sqrt{a+b+c}\sqrt{a+b(\tanh^2(x))+c(\tanh^4(x))}}\right)}{2\sqrt{a+b+c}}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(x)^3/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2), x, method=_RETURNVERBOSE)

[Out] -1/2*ln((1/2*b+c*tanh(x)^2)/c^(1/2)+(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2))/c^(1/2)+1/2/(a+b+c)^(1/2)*arctanh(1/2*(b*tanh(x)^2+2*c*tanh(x)^2+2*a+b)/(a+b+c)^(1/2)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2))

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)^3/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2), x, algorithm="maxima")

[Out] integrate(tanh(x)^3/sqrt(c*tanh(x)^4 + b*tanh(x)^2 + a), x)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 1518 vs. $2(85) = 170$.

time = 1.01, size = 6663, normalized size = 63.46

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)^3/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x, algorithm="fricas")

[Out]
$$\begin{aligned} & [1/4*((a + b + c)*\sqrt{c})\log(((b^2 + 4*(a + 2*b)*c + 8*c^2)*\cosh(x)^8 + 8*(b^2 + 4*(a + 2*b)*c + 8*c^2)*\cosh(x)*\sinh(x)^7 + (b^2 + 4*(a + 2*b)*c + 8*c^2)*\sinh(x)^8 + 4*(b^2 + 4*a*c - 8*c^2)*\cosh(x)^6 + 4*(7*(b^2 + 4*(a + 2*b)*c + 8*c^2)*\cosh(x)^2 + b^2 + 4*a*c - 8*c^2)*\sinh(x)^6 + 8*(7*(b^2 + 4*(a + 2*b)*c + 8*c^2)*\cosh(x)^3 + 3*(b^2 + 4*a*c - 8*c^2)*\cosh(x))*\sinh(x)^5 + 2*(3*b^2 + 4*(3*a - 2*b)*c + 24*c^2)*\cosh(x)^4 + 2*(35*(b^2 + 4*(a + 2*b)*c + 8*c^2)*\cosh(x)^4 + 30*(b^2 + 4*a*c - 8*c^2)*\cosh(x)^2 + 3*b^2 + 4*(3*a - 2*b)*c + 24*c^2)*\sinh(x)^4 + 8*(7*(b^2 + 4*(a + 2*b)*c + 8*c^2)*\cosh(x)^5 + 10*(b^2 + 4*a*c - 8*c^2)*\cosh(x)^3 + (3*b^2 + 4*(3*a - 2*b)*c + 24*c^2)*\cosh(x))*\sinh(x)^3 + 4*(b^2 + 4*a*c - 8*c^2)*\cosh(x)^2 + 4*(7*(b^2 + 4*(a + 2*b)*c + 8*c^2)*\cosh(x)^6 + 15*(b^2 + 4*a*c - 8*c^2)*\cosh(x)^4 + 3*(3*b^2 + 4*(3*a - 2*b)*c + 24*c^2)*\cosh(x)^2 + b^2 + 4*a*c - 8*c^2)*\sinh(x)^2 - 4*\sqrt{2}*((b + 2*c)*\cosh(x)^4 + 4*(b + 2*c)*\cosh(x)*\sinh(x)^3 + (b + 2*c)*\sinh(x)^4 + 2*(b - 2*c)*\cosh(x)^2 + 2*(3*(b + 2*c)*\cosh(x)^2 + b - 2*c)*\sinh(x)^2 + 4*((b + 2*c)*\cosh(x)^3 + (b - 2*c)*\cosh(x))*\sinh(x) + b + 2*c)*\sqrt{c})*\sqrt{((a + b + c)*\cosh(x)^4 + (a + b + c)*\sinh(x)^4 + 4*(a - c)*\cosh(x)^2 + 2*(3*(a + b + c)*\cosh(x)^2 + 2*a - 2*c)*\sinh(x)^2 + 3*a - b + 3*c)/(\cosh(x)^4 - 4*\cosh(x)^3*\sinh(x) + 6*\cosh(x)^2*\sinh(x)^2 - 4*\cosh(x)*\sinh(x)^3 + \sinh(x)^4)} + b^2 + 4*(a + 2*b)*c + 8*c^2 + 8*((b^2 + 4*(a + 2*b)*c + 8*c^2)*\cosh(x)^7 + 3*(b^2 + 4*a*c - 8*c^2)*\cosh(x)^5 + (3*b^2 + 4*(3*a - 2*b)*c + 24*c^2)*\cosh(x)^3 + (b^2 + 4*a*c - 8*c^2)*\cosh(x))*\sinh(x)/(\cosh(x)^8 + 8*\cosh(x)*\sinh(x)^7 + \sinh(x)^8 + 4*(7*\cosh(x)^2 + 1)*\sinh(x)^6 + 4*\cosh(x)^6 + 8*(7*\cosh(x)^3 + 3*\cosh(x))*\sinh(x)^5 + 2*(35*\cosh(x)^4 + 30*\cosh(x)^2 + 3)*\sinh(x)^4 + 6*\cosh(x)^4 + 8*(7*\cosh(x)^5 + 10*\cosh(x)^3 + 3*\cosh(x))*\sinh(x)^3 + 4*(7*\cosh(x)^6 + 15*\cosh(x)^4 + 9*\cosh(x)^2 + 1)*\sinh(x)^2 + 4*\cosh(x)^2 + 8*(\cosh(x)^7 + 3*\cosh(x)^5 + 3*\cosh(x)^3 + \cosh(x))*\sinh(x) + 1) + \sqrt{a + b + c}*c*\log(((a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*\cosh(x)^8 + 8*(a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*\cosh(x)*\sinh(x)^7 + (a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*\sinh(x)^8 + 4*(a^2 + a*b - b*c - c^2)*\cosh(x)^6 + 4*(7*(a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*\cosh(x)^2 + a^2 + a*b - b*c - c^2)*\sinh(x)^6 + 8*(7*(a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*\cosh(x)^3 + 3*(a^2 + a*b - b*c - c^2)*\cosh(x))*\sinh(x)^5 + 2*(3*a^2 + 2*a*b + 2*(a + b)*c + 3*c^2)*\cosh(x)^4 + 2*(35*(a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2) \end{aligned}$$

```

)*cosh(x)^4 + 30*(a^2 + a*b - b*c - c^2)*cosh(x)^2 + 3*a^2 + 2*a*b + 2*(a +
  b)*c + 3*c^2)*sinh(x)^4 + 8*(7*(a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*cos
h(x)^5 + 10*(a^2 + a*b - b*c - c^2)*cosh(x)^3 + (3*a^2 + 2*a*b + 2*(a + b)*
c + 3*c^2)*cosh(x))*sinh(x)^3 + 4*(a^2 + a*b - b*c - c^2)*cosh(x)^2 + 4*(7*
(a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*cosh(x)^6 + 15*(a^2 + a*b - b*c - c
^2)*cosh(x)^4 + 3*(3*a^2 + 2*a*b + 2*(a + b)*c + 3*c^2)*cosh(x)^2 + a^2 + a
*b - b*c - c^2)*sinh(x)^2 + sqrt(2)*((a + b + c)*cosh(x)^4 + 4*(a + b + c)*
cosh(x)*sinh(x)^3 + (a + b + c)*sinh(x)^4 + 2*(a - c)*cosh(x)^2 + 2*(3*(a +
  b + c)*cosh(x)^2 + a - c)*sinh(x)^2 + 4*((a + b + c)*cosh(x)^3 + (a - c)*c
osh(x))*sinh(x) + a + b + c)*sqrt(a + b + c)*sqrt(((a + b + c)*cosh(x)^4 +
(a + b + c)*sinh(x)^4 + 4*(a - c)*cosh(x)^2 + 2*(3*(a + b + c)*cosh(x)^2 +
2*a - 2*c)*sinh(x)^2 + 3*a - b + 3*c)/(cosh(x)^4 - 4*cosh(x)^3*sinh(x) + 6*
cosh(x)^2*sinh(x)^2 - 4*cosh(x)*sinh(x)^3 + sinh(x)^4)) + a^2 + 2*a*b + b^2
+ 2*(a + b)*c + c^2 + 8*((a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*cosh(x)^7
+ 3*(a^2 + a*b - b*c - c^2)*cosh(x)^5 + (3*a^2 + 2*a*b + 2*(a + b)*c + 3*c
^2)*cosh(x)^3 + (a^2 + a*b - b*c - c^2)*cosh(x))*sinh(x))/(cosh(x)^4 + 4*co
sh(x)^3*sinh(x) + 6*cosh(x)^2*sinh(x)^2 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4))
)/(a + b)*c + c^2), -1/4*(2*sqrt(-a - b - c)*c*arctan(sqrt(2)*((a + b + c)
*cosh(x)^4 + 4*(a + b + c)*cosh(x)*sinh(x)^3 + (a + b + c)*sinh(x)^4 + 2*(a
  - c)*cosh(x)^2 + 2*(3*(a + b + c)*cosh(x)^2 + a - c)*sinh(x)^2 + 4*((a + b
  + c)*cosh(x)^3 + (a - c)*cosh(x))*sinh(x) + a + b + c)*sqrt(-a - b - c)*sq
rt(((a + b + c)*cosh(x)^4 + (a + b + c)*sinh(x)^4 + 4*(a - c)*cosh(x)^2 + 2
*(3*(a + b + c)*cosh(x)^2 + 2*a - 2*c)*sinh(x)^2 + 3*a - b + 3*c)/(cosh(x)^
4 - 4*cosh(x)^3*sinh(x) + 6*cosh(x)^2*sinh(x)^2 - 4*cosh(x)*sinh(x)^3 + sin
h(x)^4)))/((a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*cosh(x)^8 + 8*(a^2 + 2*a*
b + b^2 + 2*(a + b)*c + c^2)*cosh(x)*sinh(x)^7 + (a^2 + 2*a*b + b^2 + 2*(a
+ b)*c + c^2)*sinh(x)^8 + 4*(a^2 + a*b - b*c - c^2)*cosh(x)^6 + 4*(7*(a^2 +
  2*a*b + b^2 + 2*(a + b)*c + c^2)*cosh(x)^2 + a^2 + a*b - b*c - c^2)*sinh(x)
)^6 + 8*(7*(a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*cosh(x)^3 + 3*(a^2 + a*b
  - b*c - c^2)*cosh(x))*sinh(x)^5 + 2*(3*a^2 + 2*a*b - b^2 + 2*(3*a + b)*c +
  3*c^2)*cosh(x)^4 + 2*(35*(a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*cosh(x)^4
  + 30*(a^2 + a*b - b*c - c^2)*cosh(x)^2 + 3*a^2 + 2*a*b - b^2 + 2*(3*a + b)
*c + 3*c^2)*sinh(x)^4 + 8*(7*(a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*cosh(x)
)^5 + 10*(a^2 + a*b - b*c - c^2)*cosh(x)^3 + (3...

```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\tanh^3(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)**3/(a+b*tanh(x)**2+c*tanh(x)**4)**(1/2),x)

[Out] Integral(tanh(x)**3/sqrt(a + b*tanh(x)**2 + c*tanh(x)**4), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)^3/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x, algorithm="giac")

[Out] integrate(tanh(x)^3/sqrt(c*tanh(x)^4 + b*tanh(x)^2 + a), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\tanh(x)^3}{\sqrt{c \tanh(x)^4 + b \tanh(x)^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(x)^3/(a + b*tanh(x)^2 + c*tanh(x)^4)^(1/2),x)

[Out] int(tanh(x)^3/(a + b*tanh(x)^2 + c*tanh(x)^4)^(1/2), x)

$$3.202 \quad \int \frac{\tanh(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx$$

Optimal. Leaf size=58

$$\frac{\tanh^{-1} \left(\frac{2a+b+(b+2c) \tanh^2(x)}{2\sqrt{a+b+c} \sqrt{a+b \tanh^2(x) + c \tanh^4(x)}} \right)}{2\sqrt{a+b+c}}$$

[Out] 1/2*arctanh(1/2*(2*a+b+(b+2*c)*tanh(x)^2)/(a+b+c)^(1/2)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2))/(a+b+c)^(1/2)

Rubi [A]

time = 0.08, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {3781, 1261, 738, 212}

$$\frac{\tanh^{-1} \left(\frac{2a+(b+2c) \tanh^2(x)+b}{2\sqrt{a+b+c} \sqrt{a+b \tanh^2(x) + c \tanh^4(x)}} \right)}{2\sqrt{a+b+c}}$$

Antiderivative was successfully verified.

[In] Int[Tanh[x]/Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4], x]

[Out] ArcTanh[(2*a + b + (b + 2*c)*Tanh[x]^2)/(2*Sqrt[a + b + c]*Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4])]/(2*Sqrt[a + b + c])

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 738

Int[1/(((d_) + (e_)*(x_))*Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2]), x_Symbol] := Dist[-2, Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[2*c*d - b*e, 0]

Rule 1261

Int[(x_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Dist[1/2, Subst[Int[(d + e*x)^q*(a + b*x + c*x^2)^p, x],

$x, x^2], x] /; \text{FreeQ}\{a, b, c, d, e, p, q\}, x]$

Rule 3781

```
Int[tan[(d_.) + (e_.)*(x_)]^(m_.)*((a_.) + (b_.)*((f_.)*tan[(d_.) + (e_.)*(x_)]^(n_.) + (c_.)*((f_.)*tan[(d_.) + (e_.)*(x_)]^(n2_.))^(p_)), x_Symbol]
:> Dist[f/e, Subst[Int[(x/f)^m*((a + b*x^n + c*x^(2*n))^p/(f^2 + x^2)), x], x, f*Tan[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[n2, 2*n] && NeQ[b^2 - 4*a*c, 0]
```

Rubi steps

$$\begin{aligned} \int \frac{\tanh(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx &= -\text{Subst} \left(\int \frac{x}{(1+x^2) \sqrt{a - bx^2 + cx^4}} dx, x, i \tanh(x) \right) \\ &= - \left(\frac{1}{2} \text{Subst} \left(\int \frac{1}{(1+x) \sqrt{a - bx + cx^2}} dx, x, -\tanh^2(x) \right) \right) \\ &= \text{Subst} \left(\int \frac{1}{4a + 4b + 4c - x^2} dx, x, \frac{2a + b + (b + 2c) \tanh^2(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} \right) \\ &= \frac{\tanh^{-1} \left(\frac{2a + b + (b + 2c) \tanh^2(x)}{2\sqrt{a + b + c} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} \right)}{2\sqrt{a + b + c}} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 58, normalized size = 1.00

$$\frac{\tanh^{-1} \left(\frac{2a + b + (b + 2c) \tanh^2(x)}{2\sqrt{a + b + c} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} \right)}{2\sqrt{a + b + c}}$$

Antiderivative was successfully verified.

[In] Integrate[Tanh[x]/Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4], x]

[Out] ArcTanh[(2*a + b + (b + 2*c)*Tanh[x]^2)/(2*Sqrt[a + b + c]*Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4])]/(2*Sqrt[a + b + c])

Maple [A]

time = 1.26, size = 52, normalized size = 0.90

method	result	size
derivativedivides	$\frac{\operatorname{arctanh}\left(\frac{b(\tanh^2(x))+2c(\tanh^2(x))+2a+b}{2\sqrt{a+b+c}\sqrt{a+b(\tanh^2(x))+c(\tanh^4(x))}}\right)}{2\sqrt{a+b+c}}$	52
default	$\frac{\operatorname{arctanh}\left(\frac{b(\tanh^2(x))+2c(\tanh^2(x))+2a+b}{2\sqrt{a+b+c}\sqrt{a+b(\tanh^2(x))+c(\tanh^4(x))}}\right)}{2\sqrt{a+b+c}}$	52

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tanh(x)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x,method=_RETURNVERBOSE)`

[Out] $1/2/(a+b+c)^{(1/2)}*\operatorname{arctanh}(1/2*(b*\tanh(x)^2+2*c*\tanh(x)^2+2*a+b)/(a+b+c)^{(1/2)})/(a+b*\tanh(x)^2+c*\tanh(x)^4)^{(1/2)}$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(x)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x, algorithm="maxima")`

[Out] `integrate(tanh(x)/sqrt(c*tanh(x)^4 + b*tanh(x)^2 + a), x)`

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 882 vs. 2(48) = 96.

time = 0.77, size = 1748, normalized size = 30.14

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tanh(x)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x, algorithm="fricas")`

[Out] $[1/4*\log(((a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*\cosh(x)^8 + 8*(a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*\cosh(x)*\sinh(x)^7 + (a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*\sinh(x)^8 + 4*(a^2 + a*b - b*c - c^2)*\cosh(x)^6 + 4*(7*(a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*\cosh(x)^2 + a^2 + a*b - b*c - c^2)*\sinh(x)^6 + 8*(7*(a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*\cosh(x)^3 + 3*(a^2 + a*b - b*c - c^2)*\cosh(x))*\sinh(x)^5 + 2*(3*a^2 + 2*a*b + 2*(a + b)*c + 3*c^2)*\cosh(x)^4 + 2*(35*(a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*\cosh(x)^4 + 30*(a^2 + a*b - b*c - c^2)*\cosh(x)^2 + 3*a^2 + 2*a*b + 2*(a + b)*c + 3*c^2)*\sinh$

$$\begin{aligned}
& (x)^4 + 8*(7*(a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*\cosh(x)^5 + 10*(a^2 + \\
& a*b - b*c - c^2)*\cosh(x)^3 + (3*a^2 + 2*a*b + 2*(a + b)*c + 3*c^2)*\cosh(x) \\
& *\sinh(x)^3 + 4*(a^2 + a*b - b*c - c^2)*\cosh(x)^2 + 4*(7*(a^2 + 2*a*b + b^2 \\
& + 2*(a + b)*c + c^2)*\cosh(x)^6 + 15*(a^2 + a*b - b*c - c^2)*\cosh(x)^4 + 3*(\\
& 3*a^2 + 2*a*b + 2*(a + b)*c + 3*c^2)*\cosh(x)^2 + a^2 + a*b - b*c - c^2)*\sin \\
& h(x)^2 + \sqrt{2}*((a + b + c)*\cosh(x)^4 + 4*(a + b + c)*\cosh(x)*\sinh(x)^3 + \\
& (a + b + c)*\sinh(x)^4 + 2*(a - c)*\cosh(x)^2 + 2*(3*(a + b + c)*\cosh(x)^2 + \\
& a - c)*\sinh(x)^2 + 4*((a + b + c)*\cosh(x)^3 + (a - c)*\cosh(x))*\sinh(x) + a \\
& + b + c)*\sqrt{a + b + c}*\sqrt{((a + b + c)*\cosh(x)^4 + (a + b + c)*\sinh(x) \\
& ^4 + 4*(a - c)*\cosh(x)^2 + 2*(3*(a + b + c)*\cosh(x)^2 + 2*a - 2*c)*\sinh(x)^2 \\
& + 3*a - b + 3*c)/(\cosh(x)^4 - 4*\cosh(x)^3*\sinh(x) + 6*\cosh(x)^2*\sinh(x)^2 \\
& - 4*\cosh(x)*\sinh(x)^3 + \sinh(x)^4)} + a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2 \\
& + 8*((a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*\cosh(x)^7 + 3*(a^2 + a*b - b \\
& *c - c^2)*\cosh(x)^5 + (3*a^2 + 2*a*b + 2*(a + b)*c + 3*c^2)*\cosh(x)^3 + (a^2 \\
& + a*b - b*c - c^2)*\cosh(x))*\sinh(x))/(\cosh(x)^4 + 4*\cosh(x)^3*\sinh(x) + 6 \\
& *\cosh(x)^2*\sinh(x)^2 + 4*\cosh(x)*\sinh(x)^3 + \sinh(x)^4))/\sqrt{a + b + c}, - \\
& 1/2*\sqrt{-a - b - c}*\arctan(\sqrt{2}*((a + b + c)*\cosh(x)^4 + 4*(a + b + c)* \\
& \cosh(x)*\sinh(x)^3 + (a + b + c)*\sinh(x)^4 + 2*(a - c)*\cosh(x)^2 + 2*(3*(a + \\
& b + c)*\cosh(x)^2 + a - c)*\sinh(x)^2 + 4*((a + b + c)*\cosh(x)^3 + (a - c)*c \\
& osh(x))*\sinh(x) + a + b + c)*\sqrt{-a - b - c}*\sqrt{((a + b + c)*\cosh(x)^4 + \\
& (a + b + c)*\sinh(x)^4 + 4*(a - c)*\cosh(x)^2 + 2*(3*(a + b + c)*\cosh(x)^2 + \\
& 2*a - 2*c)*\sinh(x)^2 + 3*a - b + 3*c)/(\cosh(x)^4 - 4*\cosh(x)^3*\sinh(x) + 6 \\
& *\cosh(x)^2*\sinh(x)^2 - 4*\cosh(x)*\sinh(x)^3 + \sinh(x)^4)})/((a^2 + 2*a*b + b^2 \\
& + 2*(a + b)*c + c^2)*\cosh(x)^8 + 8*(a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2) \\
&)*\cosh(x)*\sinh(x)^7 + (a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*\sinh(x)^8 + 4 \\
& *(a^2 + a*b - b*c - c^2)*\cosh(x)^6 + 4*(7*(a^2 + 2*a*b + b^2 + 2*(a + b)*c \\
& + c^2)*\cosh(x)^2 + a^2 + a*b - b*c - c^2)*\sinh(x)^6 + 8*(7*(a^2 + 2*a*b + b \\
& ^2 + 2*(a + b)*c + c^2)*\cosh(x)^3 + 3*(a^2 + a*b - b*c - c^2)*\cosh(x))*\sinh \\
& (x)^5 + 2*(3*a^2 + 2*a*b - b^2 + 2*(3*a + b)*c + 3*c^2)*\cosh(x)^4 + 2*(35*(\\
& a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*\cosh(x)^4 + 30*(a^2 + a*b - b*c - c^2) \\
&)*\cosh(x)^2 + 3*a^2 + 2*a*b - b^2 + 2*(3*a + b)*c + 3*c^2)*\sinh(x)^4 + 8*(\\
& 7*(a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*\cosh(x)^5 + 10*(a^2 + a*b - b*c - \\
& c^2)*\cosh(x)^3 + (3*a^2 + 2*a*b - b^2 + 2*(3*a + b)*c + 3*c^2)*\cosh(x))*\si \\
& nh(x)^3 + 4*(a^2 + a*b - b*c - c^2)*\cosh(x)^2 + 4*(7*(a^2 + 2*a*b + b^2 + 2 \\
& *(a + b)*c + c^2)*\cosh(x)^6 + 15*(a^2 + a*b - b*c - c^2)*\cosh(x)^4 + 3*(3*a \\
& ^2 + 2*a*b - b^2 + 2*(3*a + b)*c + 3*c^2)*\cosh(x)^2 + a^2 + a*b - b*c - c^2 \\
&)*\sinh(x)^2 + a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2 + 8*((a^2 + 2*a*b + b^2 \\
& + 2*(a + b)*c + c^2)*\cosh(x)^7 + 3*(a^2 + a*b - b*c - c^2)*\cosh(x)^5 + (3* \\
& a^2 + 2*a*b - b^2 + 2*(3*a + b)*c + 3*c^2)*\cosh(x)^3 + (a^2 + a*b - b*c - c \\
& ^2)*\cosh(x))*\sinh(x))/((a + b + c)]
\end{aligned}$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\tanh(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)/(a+b*tanh(x)**2+c*tanh(x)**4)**(1/2),x)

[Out] Integral(tanh(x)/sqrt(a + b*tanh(x)**2 + c*tanh(x)**4), x)

Giac [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tanh(x)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x, algorithm="giac")

[Out] Timed out

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\tanh(x)}{\sqrt{c \tanh^4(x) + b \tanh^2(x) + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(x)/(a + b*tanh(x)^2 + c*tanh(x)^4)^(1/2),x)

[Out] int(tanh(x)/(a + b*tanh(x)^2 + c*tanh(x)^4)^(1/2), x)

$$3.203 \quad \int \frac{\coth(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx$$

Optimal. Leaf size=106

$$\frac{\tanh^{-1}\left(\frac{2a+b \tanh^2(x)}{2\sqrt{a} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}}\right)}{2\sqrt{a}} + \frac{\tanh^{-1}\left(\frac{2a+b+(b+2c) \tanh^2(x)}{2\sqrt{a+b+c} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}}\right)}{2\sqrt{a+b+c}}$$

[Out] $-1/2*\operatorname{arctanh}(1/2*(2*a+b*\tanh(x)^2)/a^{(1/2)}/(a+b*\tanh(x)^2+c*\tanh(x)^4)^{(1/2)})/a^{(1/2)}+1/2*\operatorname{arctanh}(1/2*(2*a+b+(b+2*c)*\tanh(x)^2)/(a+b+c)^{(1/2)}/(a+b*\tanh(x)^2+c*\tanh(x)^4)^{(1/2)})/(a+b+c)^{(1/2)}$

Rubi [A]

time = 0.16, antiderivative size = 106, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 5, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {3781, 1265, 974, 738, 212}

$$\frac{\tanh^{-1}\left(\frac{2a+(b+2c) \tanh^2(x)+b}{2\sqrt{a+b+c} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}}\right)}{2\sqrt{a+b+c}} - \frac{\tanh^{-1}\left(\frac{2a+b \tanh^2(x)}{2\sqrt{a} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}}\right)}{2\sqrt{a}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[\operatorname{Coth}[x]/\operatorname{Sqrt}[a + b*\operatorname{Tanh}[x]^2 + c*\operatorname{Tanh}[x]^4], x]$

[Out] $-1/2*\operatorname{ArcTanh}[(2*a + b*\operatorname{Tanh}[x]^2)/(2*\operatorname{Sqrt}[a]*\operatorname{Sqrt}[a + b*\operatorname{Tanh}[x]^2 + c*\operatorname{Tanh}[x]^4)])/ \operatorname{Sqrt}[a] + \operatorname{ArcTanh}[(2*a + b + (b + 2*c)*\operatorname{Tanh}[x]^2)/(2*\operatorname{Sqrt}[a + b + c]*\operatorname{Sqrt}[a + b*\operatorname{Tanh}[x]^2 + c*\operatorname{Tanh}[x]^4)])/ (2*\operatorname{Sqrt}[a + b + c])]$

Rule 212

$\operatorname{Int}[(a_.) + (b_.)*(x_.)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x] \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$

Rule 738

$\operatorname{Int}[1/(((d_.) + (e_.)*(x_))*\operatorname{Sqrt}[(a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2]), x_Symbol] \rightarrow \operatorname{Dist}[-2, \operatorname{Subst}[\operatorname{Int}[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/\operatorname{Sqrt}[a + b*x + c*x^2]], x] /; \operatorname{FreeQ}\{a, b, c, d, e\}, x] \ \&\& \operatorname{NeQ}[b^2 - 4*a*c, 0] \ \&\& \operatorname{NeQ}[2*c*d - b*e, 0]$

Rule 974

```
Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))^(n_)*((a_.) + (b_.)*(x_)
+ (c_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(f + g
*x)^n*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && NeQ
[e*f - d*g, 0] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && (
IntegerQ[p] || (ILtQ[m, 0] && ILtQ[n, 0])) && !(IGtQ[m, 0] || IGtQ[n, 0])
```

Rule 1265

```
Int[(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(q_.)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)
)^4)^(p_.), x_Symbol] := Dist[1/2, Subst[Int[x^((m - 1)/2)*(d + e*x)^q*(a +
b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x] && Inte
gerQ[(m - 1)/2]
```

Rule 3781

```
Int[tan[(d_.) + (e_.)*(x_)]^(m_.)*((a_.) + (b_.)*((f_.)*tan[(d_.) + (e_.)*(
x_)])^(n_.) + (c_.)*((f_.)*tan[(d_.) + (e_.)*(x_)])^(n2_.))^(p_), x_Symbol]
:= Dist[f/e, Subst[Int[(x/f)^m*((a + b*x^n + c*x^(2*n))^p/(f^2 + x^2)), x]
, x, f*Tan[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[n2
, 2*n] && NeQ[b^2 - 4*a*c, 0]
```

Rubi steps

$$\begin{aligned}
\int \frac{\coth(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx &= \text{Subst} \left(\int \frac{1}{x(1+x^2)\sqrt{a - bx^2 + cx^4}} dx, x, i \tanh(x) \right) \\
&= \frac{1}{2} \text{Subst} \left(\int \frac{1}{x(1+x)\sqrt{a - bx + cx^2}} dx, x, -\tanh^2(x) \right) \\
&= \frac{1}{2} \text{Subst} \left(\int \left(\frac{1}{(-1-x)\sqrt{a - bx + cx^2}} + \frac{1}{x\sqrt{a - bx + cx^2}} \right) dx, x \right) \\
&= \frac{1}{2} \text{Subst} \left(\int \frac{1}{(-1-x)\sqrt{a - bx + cx^2}} dx, x, -\tanh^2(x) \right) + \frac{1}{2} \text{Subst} \left(\int \frac{1}{x\sqrt{a - bx + cx^2}} dx, x, -\tanh^2(x) \right) \\
&= -\text{Subst} \left(\int \frac{1}{4a - x^2} dx, x, \frac{2a + b \tanh^2(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} \right) - \text{Subst} \left(\int \frac{1}{x\sqrt{a - bx + cx^2}} dx, x, -\tanh^2(x) \right) \\
&= -\frac{\tanh^{-1} \left(\frac{2a + b \tanh^2(x)}{2\sqrt{a} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} \right)}{2\sqrt{a}} - \frac{\tanh^{-1} \left(\frac{-\tanh^2(x)}{\sqrt{a - bx + cx^2}} \right)}{2\sqrt{a}}
\end{aligned}$$

Mathematica [A]

time = 0.26, size = 109, normalized size = 1.03

$$\frac{\tanh^{-1}\left(\frac{2a+b\tanh^2(x)}{2\sqrt{a}\sqrt{a+b\tanh^2(x)+c\tanh^4(x)}}\right)}{2\sqrt{a}} - \frac{\tanh^{-1}\left(\frac{-2a-b-(b+2c)\tanh^2(x)}{2\sqrt{a+b+c}\sqrt{a+b\tanh^2(x)+c\tanh^4(x)}}\right)}{2\sqrt{a+b+c}}$$

Antiderivative was successfully verified.

[In] Integrate[Coth[x]/Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4], x]

```
[Out] -1/2*ArcTanh[(2*a + b*Tanh[x]^2)/(2*Sqrt[a]*Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4])]/Sqrt[a] - ArcTanh[(-2*a - b - (b + 2*c)*Tanh[x]^2)/(2*Sqrt[a + b + c]*Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4])]/(2*Sqrt[a + b + c])
```

Maple [F]

time = 1.97, size = 0, normalized size = 0.00

$$\int \frac{\coth(x)}{\sqrt{a + b(\tanh^2(x)) + c(\tanh^4(x))}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(coth(x)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2), x)

[Out] int(coth(x)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(x)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2), x, algorithm="maxima")

[Out] integrate(coth(x)/sqrt(c*tanh(x)^4 + b*tanh(x)^2 + a), x)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 1520 vs. 2(86) = 172.

time = 0.95, size = 6705, normalized size = 63.25

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(x)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2), x, algorithm="fricas")

[Out]
$$\begin{aligned} & [1/4*((a + b + c)*\text{sqrt}(a)*\log(((8*a^2 + 8*a*b + b^2 + 4*a*c)*\cosh(x))^8 + 8* \\ & (8*a^2 + 8*a*b + b^2 + 4*a*c)*\cosh(x)*\sinh(x)^7 + (8*a^2 + 8*a*b + b^2 + 4* \\ & a*c)*\sinh(x)^8 + 4*(8*a^2 - b^2 - 4*a*c)*\cosh(x)^6 + 4*(7*(8*a^2 + 8*a*b + \\ & b^2 + 4*a*c)*\cosh(x)^2 + 8*a^2 - b^2 - 4*a*c)*\sinh(x)^6 + 8*(7*(8*a^2 + 8*a \\ & *b + b^2 + 4*a*c)*\cosh(x)^3 + 3*(8*a^2 - b^2 - 4*a*c)*\cosh(x))*\sinh(x)^5 + \\ & 2*(24*a^2 - 8*a*b + 3*b^2 + 12*a*c)*\cosh(x)^4 + 2*(35*(8*a^2 + 8*a*b + b^2 \\ & + 4*a*c)*\cosh(x)^4 + 30*(8*a^2 - b^2 - 4*a*c)*\cosh(x)^2 + 24*a^2 - 8*a*b + \\ & 3*b^2 + 12*a*c)*\sinh(x)^4 + 8*(7*(8*a^2 + 8*a*b + b^2 + 4*a*c)*\cosh(x)^5 + \\ & 10*(8*a^2 - b^2 - 4*a*c)*\cosh(x)^3 + (24*a^2 - 8*a*b + 3*b^2 + 12*a*c)*\cosh \\ & (x))*\sinh(x)^3 + 4*(8*a^2 - b^2 - 4*a*c)*\cosh(x)^2 + 4*(7*(8*a^2 + 8*a*b + \\ & b^2 + 4*a*c)*\cosh(x)^6 + 15*(8*a^2 - b^2 - 4*a*c)*\cosh(x)^4 + 3*(24*a^2 - 8 \\ & *a*b + 3*b^2 + 12*a*c)*\cosh(x)^2 + 8*a^2 - b^2 - 4*a*c)*\sinh(x)^2 - 4*\text{sqrt}(\\ & 2)*((2*a + b)*\cosh(x)^4 + 4*(2*a + b)*\cosh(x)*\sinh(x)^3 + (2*a + b)*\sinh(x) \\ & ^4 + 2*(2*a - b)*\cosh(x)^2 + 2*(3*(2*a + b)*\cosh(x)^2 + 2*a - b)*\sinh(x)^2 \\ & + 4*((2*a + b)*\cosh(x)^3 + (2*a - b)*\cosh(x))*\sinh(x) + 2*a + b)*\text{sqrt}(a)*\text{sq} \\ & \text{rt}(((a + b + c)*\cosh(x)^4 + (a + b + c)*\sinh(x)^4 + 4*(a - c)*\cosh(x)^2 + 2 \\ & *(3*(a + b + c)*\cosh(x)^2 + 2*a - 2*c)*\sinh(x)^2 + 3*a - b + 3*c)/(\cosh(x)^ \\ & 4 - 4*\cosh(x)^3*\sinh(x) + 6*\cosh(x)^2*\sinh(x)^2 - 4*\cosh(x)*\sinh(x)^3 + \sinh \\ & (x)^4)) + 8*a^2 + 8*a*b + b^2 + 4*a*c + 8*((8*a^2 + 8*a*b + b^2 + 4*a*c)*\c \\ & \text{osh}(x)^7 + 3*(8*a^2 - b^2 - 4*a*c)*\cosh(x)^5 + (24*a^2 - 8*a*b + 3*b^2 + 12 \\ & *a*c)*\cosh(x)^3 + (8*a^2 - b^2 - 4*a*c)*\cosh(x))*\sinh(x))/(\cosh(x)^8 + 8*\c \\ & \text{osh}(x)*\sinh(x)^7 + \sinh(x)^8 + 4*(7*\cosh(x)^2 - 1)*\sinh(x)^6 - 4*\cosh(x)^6 + \\ & 8*(7*\cosh(x)^3 - 3*\cosh(x))*\sinh(x)^5 + 2*(35*\cosh(x)^4 - 30*\cosh(x)^2 + 3 \\ &)*\sinh(x)^4 + 6*\cosh(x)^4 + 8*(7*\cosh(x)^5 - 10*\cosh(x)^3 + 3*\cosh(x))*\sinh \\ & (x)^3 + 4*(7*\cosh(x)^6 - 15*\cosh(x)^4 + 9*\cosh(x)^2 - 1)*\sinh(x)^2 - 4*\cosh \\ & (x)^2 + 8*(\cosh(x)^7 - 3*\cosh(x)^5 + 3*\cosh(x)^3 - \cosh(x))*\sinh(x) + 1)) + \\ & \text{sqrt}(a + b + c)*a*\log(((a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*\cosh(x))^8 + \\ & 8*(a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*\cosh(x)*\sinh(x)^7 + (a^2 + 2*a*b \\ & + b^2 + 2*(a + b)*c + c^2)*\sinh(x)^8 + 4*(a^2 + a*b - b*c - c^2)*\cosh(x)^6 \\ & + 4*(7*(a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*\cosh(x)^2 + a^2 + a*b - b*c \\ & - c^2)*\sinh(x)^6 + 8*(7*(a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*\cosh(x)^3 \\ & + 3*(a^2 + a*b - b*c - c^2)*\cosh(x))*\sinh(x)^5 + 2*(3*a^2 + 2*a*b + 2*(a + \\ & b)*c + 3*c^2)*\cosh(x)^4 + 2*(35*(a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*\c \\ & \text{osh}(x)^4 + 30*(a^2 + a*b - b*c - c^2)*\cosh(x)^2 + 3*a^2 + 2*a*b + 2*(a + b)*c \\ & + 3*c^2)*\sinh(x)^4 + 8*(7*(a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*\cosh(x)^ \\ & 5 + 10*(a^2 + a*b - b*c - c^2)*\cosh(x)^3 + (3*a^2 + 2*a*b + 2*(a + b)*c + 3 \\ & *c^2)*\cosh(x))*\sinh(x)^3 + 4*(a^2 + a*b - b*c - c^2)*\cosh(x)^2 + 4*(7*(a^2 \\ & + 2*a*b + b^2 + 2*(a + b)*c + c^2)*\cosh(x)^6 + 15*(a^2 + a*b - b*c - c^2)*\c \\ & \text{osh}(x)^4 + 3*(3*a^2 + 2*a*b + 2*(a + b)*c + 3*c^2)*\cosh(x)^2 + a^2 + a*b - \\ & b*c - c^2)*\sinh(x)^2 + \text{sqrt}(2)*((a + b + c)*\cosh(x)^4 + 4*(a + b + c)*\cosh(\\ & x)*\sinh(x)^3 + (a + b + c)*\sinh(x)^4 + 2*(a - c)*\cosh(x)^2 + 2*(3*(a + b + \\ & c)*\cosh(x)^2 + a - c)*\sinh(x)^2 + 4*((a + b + c)*\cosh(x)^3 + (a - c)*\cosh(x) \\ &))*\sinh(x) + a + b + c)*\text{sqrt}(a + b + c)*\text{sqrt}(((a + b + c)*\cosh(x)^4 + (a + \\ & b + c)*\sinh(x)^4 + 4*(a - c)*\cosh(x)^2 + 2*(3*(a + b + c)*\cosh(x)^2 + 2*a - \\ & 2*c)*\sinh(x)^2 + 3*a - b + 3*c)/(\cosh(x)^4 - 4*\cosh(x)^3*\sinh(x) + 6*\cosh(\end{aligned}$$

$x^2 \sinh(x)^2 - 4 \cosh(x) \sinh(x)^3 + \sinh(x)^4) + a^2 + 2ab + b^2 + 2(a+b)c + c^2 + 8((a^2 + 2ab + b^2 + 2(a+b)c + c^2) \cosh(x)^7 + 3(a^2 + ab - bc - c^2) \cosh(x)^5 + (3a^2 + 2ab + 2(a+b)c + 3c^2) \cosh(x)^3 + (a^2 + ab - bc - c^2) \cosh(x)) \sinh(x) / (\cosh(x)^4 + 4 \cosh(x)^3 \sinh(x) + 6 \cosh(x)^2 \sinh(x)^2 + 4 \cosh(x) \sinh(x)^3 + \sinh(x)^4) / (a^2 + ab + ac), -1/4(2a \sqrt{-a-b-c}) \arctan(\sqrt{2}((a+b+c) \cosh(x)^4 + 4(a+b+c) \cosh(x) \sinh(x)^3 + (a+b+c) \sinh(x)^4 + 2(a-c) \cosh(x)^2 + 2(3(a+b+c) \cosh(x)^2 + a-c) \sinh(x)^2 + 4((a+b+c) \cosh(x)^3 + (a-c) \cosh(x)) \sinh(x) + a+b+c) \sqrt{-a-b-c}) \sqrt{((a+b+c) \cosh(x)^4 + (a+b+c) \sinh(x)^4 + 4(a-c) \cosh(x)^2 + 2(3(a+b+c) \cosh(x)^2 + 2a-2c) \sinh(x)^2 + 3a-b+3c) / (\cosh(x)^4 - 4 \cosh(x)^3 \sinh(x) + 6 \cosh(x)^2 \sinh(x)^2 - 4 \cosh(x) \sinh(x)^3 + \sinh(x)^4)) / ((a^2 + 2ab + b^2 + 2(a+b)c + c^2) \cosh(x)^8 + 8(a^2 + 2ab + b^2 + 2(a+b)c + c^2) \cosh(x) \sinh(x)^7 + (a^2 + 2ab + b^2 + 2(a+b)c + c^2) \sinh(x)^8 + 4(a^2 + ab - bc - c^2) \cosh(x)^6 + 4(7(a^2 + 2ab + b^2 + 2(a+b)c + c^2) \cosh(x)^2 + a^2 + ab - bc - c^2) \sinh(x)^6 + 8(7(a^2 + 2ab + b^2 + 2(a+b)c + c^2) \cosh(x)^3 + 3(a^2 + ab - bc - c^2) \cosh(x)) \sinh(x)^5 + 2(3a^2 + 2ab - b^2 + 2(3a+b)c + 3c^2) \cosh(x)^4 + 2(35(a^2 + 2ab + b^2 + 2(a+b)c + c^2) \cosh(x)^4 + 30(a^2 + ab - bc - c^2) \cosh(x)^2 + 3a^2 + 2ab - b^2 + 2(3a+b)c + 3c^2) \sinh(x)^4 + 8(7(a^2 + 2ab + b^2 + 2(a+b)c + c^2) \cosh(x)^5 + 10(a^2 + ab - bc - c^2) \cosh(x)^3 + (3a^2 \dots$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\coth(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(x)/(a+b*tanh(x)**2+c*tanh(x)**4)**(1/2),x)

[Out] Integral(coth(x)/sqrt(a + b*tanh(x)**2 + c*tanh(x)**4), x)

Giac [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(x)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x, algorithm="giac")

[Out] Timed out

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\coth(x)}{\sqrt{c \tanh(x)^4 + b \tanh(x)^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(coth(x)/(a + b*tanh(x)^2 + c*tanh(x)^4)^(1/2), x)

[Out] int(coth(x)/(a + b*tanh(x)^2 + c*tanh(x)^4)^(1/2), x)

$$3.204 \quad \int \frac{\coth^3(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx$$

Optimal. Leaf size=183

$$\frac{\tanh^{-1}\left(\frac{2a+b \tanh^2(x)}{2\sqrt{a} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}}\right)}{2\sqrt{a}} + \frac{b \tanh^{-1}\left(\frac{2a+b \tanh^2(x)}{2\sqrt{a} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}}\right)}{4a^{3/2}} + \dots$$

[Out] $\frac{1}{4} b \operatorname{arctanh}\left(\frac{1}{2} \frac{2a+b \tanh(x)^2}{a^{1/2}}\right) / (a+b \tanh(x)^2+c \tanh(x)^4)^{1/2} - \frac{1}{2} \operatorname{arctanh}\left(\frac{1}{2} \frac{2a+b \tanh(x)^2}{a^{1/2}}\right) / (a+b \tanh(x)^2+c \tanh(x)^4)^{1/2} + \frac{1}{2} \operatorname{arctanh}\left(\frac{1}{2} \frac{2a+b+(b+2c) \tanh(x)^2}{(a+b+c)^{1/2}}\right) / (a+b \tanh(x)^2+c \tanh(x)^4)^{1/2} - \frac{1}{2} \coth(x)^2 (a+b \tanh(x)^2+c \tanh(x)^4)^{1/2} / a$

Rubi [A]

time = 0.24, antiderivative size = 183, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 6, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.261$, Rules used = {3781, 1265, 974, 744, 738, 212}

$$\frac{b \tanh^{-1}\left(\frac{2a+b \tanh^2(x)}{2\sqrt{a} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}}\right)}{4a^{3/2}} - \frac{\tanh^{-1}\left(\frac{2a+b \tanh^2(x)}{2\sqrt{a} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}}\right)}{2\sqrt{a}} + \frac{\tanh^{-1}\left(\frac{2a+(b+2c) \tanh^2(x)+b}{2\sqrt{a+b+c} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}}\right)}{2\sqrt{a+b+c}} - \frac{\coth^2(x) \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}}{2a}$$

Antiderivative was successfully verified.

[In] Int[Coth[x]^3/Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4], x]

[Out] $-\frac{1}{2} \operatorname{ArcTanh}\left[\frac{(2a + b \operatorname{Tanh}[x]^2)}{(2 \operatorname{Sqrt}[a] \operatorname{Sqrt}[a + b \operatorname{Tanh}[x]^2 + c \operatorname{Tanh}[x]^4])}\right] / \operatorname{Sqrt}[a] + \frac{(b \operatorname{ArcTanh}\left[\frac{(2a + b \operatorname{Tanh}[x]^2)}{(2 \operatorname{Sqrt}[a] \operatorname{Sqrt}[a + b \operatorname{Tanh}[x]^2 + c \operatorname{Tanh}[x]^4])}\right])}{(4a^{3/2})} + \operatorname{ArcTanh}\left[\frac{(2a + b + (b + 2c) \operatorname{Tanh}[x]^2)}{(2 \operatorname{Sqrt}[a + b + c] \operatorname{Sqrt}[a + b \operatorname{Tanh}[x]^2 + c \operatorname{Tanh}[x]^4])}\right] / (2 \operatorname{Sqrt}[a + b + c]) - \frac{(\operatorname{Coth}[x]^2 \operatorname{Sqrt}[a + b \operatorname{Tanh}[x]^2 + c \operatorname{Tanh}[x]^4])}{(2a)}$

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 738

Int[1/(((d_.) + (e_.)*(x_))*Sqrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]), x_Symbol] := Dist[-2, Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[2*c*d - b*e, 0]

Rule 744

```
Int[((d_.) + (e_.)*(x_))^(m_)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol]
:= Simp[e*(d + e*x)^(m + 1)*((a + b*x + c*x^2)^(p + 1)/((m + 1)*(c*d^2 - b*d*e + a*e^2))), x]
+ Dist[(2*c*d - b*e)/(2*(c*d^2 - b*d*e + a*e^2)), Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^p, x], x]
/; FreeQ[{a, b, c, d, e, m, p}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[2*c*d - b*e, 0] && EqQ[m + 2*p + 3, 0]
```

Rule 974

```
Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))^(n_)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol]
:= Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + b*x + c*x^2)^p, x], x]
/; FreeQ[{a, b, c, d, e, f, g}, x] && NeQ[e*f - d*g, 0] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && (IntegerQ[p] || (ILtQ[m, 0] && ILtQ[n, 0])) && !(IGtQ[m, 0] || IGtQ[n, 0])
```

Rule 1265

```
Int[(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(q_.)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_.), x_Symbol]
:= Dist[1/2, Subst[Int[x^((m - 1)/2)*(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x]
/; FreeQ[{a, b, c, d, e, p, q}, x] && IntegerQ[(m - 1)/2]
```

Rule 3781

```
Int[tan[(d_.) + (e_.)*(x_)]^(m_.)*((a_.) + (b_.)*((f_.)*tan[(d_.) + (e_.)*(x_)])^(n_.) + (c_.)*((f_.)*tan[(d_.) + (e_.)*(x_)])^(n2_.))^(p_), x_Symbol]
:= Dist[f/e, Subst[Int[(x/f)^m*((a + b*x^n + c*x^(2*n))^p/(f^2 + x^2)), x], x, f*Tan[d + e*x]], x]
/; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[n2, 2*n] && NeQ[b^2 - 4*a*c, 0]
```

Rubi steps

$$\begin{aligned}
\int \frac{\coth^3(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx &= -\text{Subst} \left(\int \frac{1}{x^3(1+x^2)\sqrt{a-bx^2+cx^4}} dx, x, i \tanh(x) \right) \\
&= -\left(\frac{1}{2} \text{Subst} \left(\int \frac{1}{x^2(1+x)\sqrt{a-bx+cx^2}} dx, x, -\tanh^2(x) \right) \right) \\
&= -\left(\frac{1}{2} \text{Subst} \left(\int \left(\frac{1}{x^2\sqrt{a-bx+cx^2}} - \frac{1}{x\sqrt{a-bx+cx^2}} + \frac{1}{(1+x)\sqrt{a-bx+cx^2}} \right) dx, x, -\tanh^2(x) \right) \right) \\
&= -\left(\frac{1}{2} \text{Subst} \left(\int \frac{1}{x^2\sqrt{a-bx+cx^2}} dx, x, -\tanh^2(x) \right) \right) + \frac{1}{2} \text{Subst} \left(\int \frac{1}{x\sqrt{a-bx+cx^2}} dx, x, -\tanh^2(x) \right) \\
&= -\frac{\coth^2(x)\sqrt{a+b\tanh^2(x)+c\tanh^4(x)}}{2a} - \frac{b \text{Subst} \left(\int \frac{1}{x\sqrt{a-bx+cx^2}} dx, x, -\tanh^2(x) \right)}{4} \\
&= -\frac{\tanh^{-1} \left(\frac{2a+b\tanh^2(x)}{2\sqrt{a}\sqrt{a+b\tanh^2(x)+c\tanh^4(x)}} \right)}{2\sqrt{a}} + \frac{\tanh^{-1} \left(\frac{2a+b+(b+2c)\tanh^2(x)}{2\sqrt{a+b+c}\sqrt{a+b\tanh^2(x)+c\tanh^4(x)}} \right)}{2\sqrt{a+b+c}} \\
&= -\frac{\tanh^{-1} \left(\frac{2a+b\tanh^2(x)}{2\sqrt{a}\sqrt{a+b\tanh^2(x)+c\tanh^4(x)}} \right)}{2\sqrt{a}} + \frac{b \tanh^{-1} \left(\frac{2a+b+(b+2c)\tanh^2(x)}{2\sqrt{a+b+c}\sqrt{a+b\tanh^2(x)+c\tanh^4(x)}} \right)}{2\sqrt{a+b+c}}
\end{aligned}$$

Mathematica [A]

time = 0.43, size = 142, normalized size = 0.78

$$-\frac{(2a-b)\tanh^{-1}\left(\frac{2a+b\tanh^2(x)}{2\sqrt{a}\sqrt{a+b\tanh^2(x)+c\tanh^4(x)}}\right)}{4a^{3/2}} + \frac{\tanh^{-1}\left(\frac{2a+b+(b+2c)\tanh^2(x)}{2\sqrt{a+b+c}\sqrt{a+b\tanh^2(x)+c\tanh^4(x)}}\right)}{2\sqrt{a+b+c}} - \frac{\coth^2(x)\sqrt{a+b\tanh^2(x)+c\tanh^4(x)}}{2a}$$

Antiderivative was successfully verified.

[In] Integrate[Coth[x]^3/Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4], x]

[Out] $-1/4*((2*a - b)*\text{ArcTanh}[(2*a + b*\text{Tanh}[x]^2)/(2*\text{Sqrt}[a]*\text{Sqrt}[a + b*\text{Tanh}[x]^2 + c*\text{Tanh}[x]^4]])/a^{(3/2)} + \text{ArcTanh}[(2*a + b + (b + 2*c)*\text{Tanh}[x]^2)/(2*\text{Sqrt}[a + b + c]*\text{Sqrt}[a + b*\text{Tanh}[x]^2 + c*\text{Tanh}[x]^4]])/(2*\text{Sqrt}[a + b + c]) - (\text{Coth}[x]^2*\text{Sqrt}[a + b*\text{Tanh}[x]^2 + c*\text{Tanh}[x]^4])/(2*a)$

Maple [F]

time = 1.82, size = 0, normalized size = 0.00

$$\int \frac{\coth^3(x)}{\sqrt{a + b(\tanh^2(x)) + c(\tanh^4(x))}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{int}(\coth(x)^3/(a+b*\tanh(x)^2+c*\tanh(x)^4)^{(1/2)}, x)$

[Out] $\text{int}(\coth(x)^3/(a+b*\tanh(x)^2+c*\tanh(x)^4)^{(1/2)}, x)$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{integrate}(\coth(x)^3/(a+b*\tanh(x)^2+c*\tanh(x)^4)^{(1/2)}, x, \text{algorithm}="maxima")$

[Out] $\text{integrate}(\coth(x)^3/\text{sqrt}(c*\tanh(x)^4 + b*\tanh(x)^2 + a), x)$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 2136 vs. 2(149) = 298.

time = 1.18, size = 9168, normalized size = 50.10

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{integrate}(\coth(x)^3/(a+b*\tanh(x)^2+c*\tanh(x)^4)^{(1/2)}, x, \text{algorithm}="fricas")$

[Out]
$$\begin{aligned} & [-1/8*((2*a^2 + a*b - b^2 + (2*a - b)*c)*\cosh(x)^4 + 4*(2*a^2 + a*b - b^2 \\ & + (2*a - b)*c)*\cosh(x)*\sinh(x)^3 + (2*a^2 + a*b - b^2 + (2*a - b)*c)*\sinh(x) \\ &)^4 - 2*(2*a^2 + a*b - b^2 + (2*a - b)*c)*\cosh(x)^2 + 2*(3*(2*a^2 + a*b - b \\ & ^2 + (2*a - b)*c)*\cosh(x)^2 - 2*a^2 - a*b + b^2 - (2*a - b)*c)*\sinh(x)^2 + \\ & 2*a^2 + a*b - b^2 + (2*a - b)*c + 4*((2*a^2 + a*b - b^2 + (2*a - b)*c)*\cosh \\ & (x))^3 - (2*a^2 + a*b - b^2 + (2*a - b)*c)*\cosh(x))*\sinh(x))*\text{sqrt}(a)*\log(((8 \\ & *a^2 + 8*a*b + b^2 + 4*a*c)*\cosh(x))^8 + 8*(8*a^2 + 8*a*b + b^2 + 4*a*c)*\cos \\ & h(x)*\sinh(x)^7 + (8*a^2 + 8*a*b + b^2 + 4*a*c)*\sinh(x)^8 + 4*(8*a^2 - b^2 - \\ & 4*a*c)*\cosh(x)^6 + 4*(7*(8*a^2 + 8*a*b + b^2 + 4*a*c)*\cosh(x)^2 + 8*a^2 - \\ & b^2 - 4*a*c)*\sinh(x)^6 + 8*(7*(8*a^2 + 8*a*b + b^2 + 4*a*c)*\cosh(x))^3 + 3*(\\ & 8*a^2 - b^2 - 4*a*c)*\cosh(x))*\sinh(x)^5 + 2*(24*a^2 - 8*a*b + 3*b^2 + 12*a* \\ & c)*\cosh(x)^4 + 2*(35*(8*a^2 + 8*a*b + b^2 + 4*a*c)*\cosh(x)^4 + 30*(8*a^2 - \\ & b^2 - 4*a*c)*\cosh(x)^2 + 24*a^2 - 8*a*b + 3*b^2 + 12*a*c)*\sinh(x)^4 + 8*(7* \\ & (8*a^2 + 8*a*b + b^2 + 4*a*c)*\cosh(x))^5 + 10*(8*a^2 - b^2 - 4*a*c)*\cosh(x)^ \\ & 3 + (24*a^2 - 8*a*b + 3*b^2 + 12*a*c)*\cosh(x))*\sinh(x)^3 + 4*(8*a^2 - b^2 - \\ & 4*a*c)*\cosh(x)^2 + 4*(7*(8*a^2 + 8*a*b + b^2 + 4*a*c)*\cosh(x))^6 + 15*(8*a^ \\ & 2 - b^2 - 4*a*c)*\cosh(x)^4 + 3*(24*a^2 - 8*a*b + 3*b^2 + 12*a*c)*\cosh(x)^2 \\ & + 8*a^2 - b^2 - 4*a*c)*\sinh(x)^2 + 4*\text{sqrt}(2)*((2*a + b)*\cosh(x))^4 + 4*(2*a \\ & + b)*\cosh(x)*\sinh(x)^3 + (2*a + b)*\sinh(x)^4 + 2*(2*a - b)*\cosh(x)^2 + 2*(3 \end{aligned}$$

$$\begin{aligned}
& * (2*a + b)*\cosh(x)^2 + 2*a - b)*\sinh(x)^2 + 4*((2*a + b)*\cosh(x)^3 + (2*a - \\
& b)*\cosh(x))*\sinh(x) + 2*a + b)*\sqrt{a}*\sqrt{((a + b + c)*\cosh(x)^4 + (a + \\
& b + c)*\sinh(x)^4 + 4*(a - c)*\cosh(x)^2 + 2*(3*(a + b + c)*\cosh(x)^2 + 2*a - \\
& 2*c)*\sinh(x)^2 + 3*a - b + 3*c)/(\cosh(x)^4 - 4*\cosh(x)^3*\sinh(x) + 6*\cosh(x)^2*\sinh(x)^2 - 4*\cosh(x)*\sinh(x)^3 + \sinh(x)^4)) + 8*a^2 + 8*a*b + b^2 + \\
& 4*a*c + 8*((8*a^2 + 8*a*b + b^2 + 4*a*c)*\cosh(x)^7 + 3*(8*a^2 - b^2 - 4*a*c) \\
&)*\cosh(x)^5 + (24*a^2 - 8*a*b + 3*b^2 + 12*a*c)*\cosh(x)^3 + (8*a^2 - b^2 - 4*a*c)*\cosh(x))*\sinh(x))/(\cosh(x)^8 + 8*\cosh(x)*\sinh(x)^7 + \sinh(x)^8 + 4*(\\
& 7*\cosh(x)^2 - 1)*\sinh(x)^6 - 4*\cosh(x)^6 + 8*(7*\cosh(x)^3 - 3*\cosh(x))*\sinh \\
& (x)^5 + 2*(35*\cosh(x)^4 - 30*\cosh(x)^2 + 3)*\sinh(x)^4 + 6*\cosh(x)^4 + 8*(7* \\
& \cosh(x)^5 - 10*\cosh(x)^3 + 3*\cosh(x))*\sinh(x)^3 + 4*(7*\cosh(x)^6 - 15*\cosh(x)^4 + 9*\cosh(x)^2 - 1)*\sinh(x)^2 - 4*\cosh(x)^2 + 8*(\cosh(x)^7 - 3*\cosh(x)^5 + 3*\cosh(x)^3 - \cosh(x))*\sinh(x) + 1)) - 2*(a^2*\cosh(x)^4 + 4*a^2*\cosh(x) \\
& *\sinh(x)^3 + a^2*\sinh(x)^4 - 2*a^2*\cosh(x)^2 + 2*(3*a^2*\cosh(x)^2 - a^2)*\sinh(x)^2 + a^2 + 4*(a^2*\cosh(x)^3 - a^2*\cosh(x))*\sinh(x))*\sqrt{a + b + c}*\log(((a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*\cosh(x)^8 + 8*(a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*\cosh(x)*\sinh(x)^7 + (a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*\sinh(x)^8 + 4*(a^2 + a*b - b*c - c^2)*\cosh(x)^6 + 4*(7*(a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*\cosh(x)^2 + a^2 + a*b - b*c - c^2)*\sinh(x)^6 + 8*(7*(a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*\cosh(x)^3 + 3*(a^2 + a*b - b*c - c^2)*\cosh(x))*\sinh(x)^5 + 2*(3*a^2 + 2*a*b + 2*(a + b)*c + 3*c^2)*\cosh(x)^4 + 2*(35*(a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*\cosh(x)^4 + 30*(a^2 + a*b - b*c - c^2)*\cosh(x)^2 + 3*a^2 + 2*a*b + 2*(a + b)*c + 3*c^2)*\sinh(x)^4 + 8*(7*(a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*\cosh(x)^5 + 10*(a^2 + a*b - b*c - c^2)*\cosh(x)^3 + (3*a^2 + 2*a*b + 2*(a + b)*c + 3*c^2)*\cosh(x))*\sinh(x)^3 + 4*(a^2 + a*b - b*c - c^2)*\cosh(x)^2 + 4*(7*(a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*\cosh(x)^6 + 15*(a^2 + a*b - b*c - c^2)*\cosh(x)^4 + 3*(3*a^2 + 2*a*b + 2*(a + b)*c + 3*c^2)*\cosh(x)^2 + a^2 + a*b - b*c - c^2)*\sinh(x)^2 + \sqrt{2}*((a + b + c)*\cosh(x)^4 + 4*(a + b + c)*\cosh(x)*\sinh(x)^3 + (a + b + c)*\sinh(x)^4 + 2*(a - c)*\cosh(x)^2 + 2*(3*(a + b + c)*\cosh(x)^2 + a - c)*\sinh(x)^2 + 4*((a + b + c)*\cosh(x)^3 + (a - c)*\cosh(x))*\sinh(x) + a + b + c)*\sqrt{a + b + c}*\sqrt{((a + b + c)*\cosh(x)^4 + (a + b + c)*\sinh(x)^4 + 4*(a - c)*\cosh(x)^2 + 2*(3*(a + b + c)*\cosh(x)^2 + 2*a - 2*c)*\sinh(x)^2 + 3*a - b + 3*c)/(\cosh(x)^4 - 4*\cosh(x)^3*\sinh(x) + 6*\cosh(x)^2*\sinh(x)^2 - 4*\cosh(x)*\sinh(x)^3 + \sinh(x)^4)) + a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2 + 8*(a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*\cosh(x)^7 + 3*(a^2 + a*b - b*c - c^2)*\cosh(x)^5 + (3*a^2 + 2*a*b + 2*(a + b)*c + 3*c^2)*\cosh(x)^3 + (a^2 + a*b - b*c - c^2)*\cosh(x))*\sinh(x))/(\cosh(x)^4 + 4*\cosh(x)^3*\sinh(x) + 6*\cosh(x)^2*\sinh(x)^2 + 4*\cosh(x)*\sinh(x)^3 + \sinh(x)^4)) + 4*\sqrt{2}*(a^2 + a*b + a*c)*\sqrt{((a + b + c)*\cosh(x)^4 + (a + b + c)*\sinh(x)^4 + 4*(a - c)*\cosh(x)^2 + 2*(3*(a + b + c)*\cosh(x)^2 + 2*a - 2*c)*\sinh(x)^2 + 3*a - b + 3*c)/(\cosh(x)^4 - 4*\cosh(x)^3*\sinh(x) + 6*\cosh(x)^2*\sinh(x)^2 - 4*\cosh(x)*\sinh(x)^3 + \sinh(x)^4)))/((a^3 + a^2*b + a^2*c)*\cosh(x)^4 + 4*(a^3 + a^2*b + a^2*c)*\cosh(x)*\sinh(x)^3 + (a^3 + a^2*b + a^2*c)*\sinh(x)^4 + a^3 + a^2*b + a^2*c - 2*(a^3 + a^2*b + a^2*c)*\cosh(x)^2 - 2*(a^3 + a^2*b + a^2*c - 3*(a^3 + a^2
\end{aligned}$$

$*b + a^2*c)*\cosh(x)^2)*\sinh(x)^2 + 4*((a^3 + a^2*b + a^2*c)*\cosh(x)^3 - (a^3 + a^2*b + a^2*c)*\cosh(x))*\sinh(x)), -1/8*(4*(a^2*\cosh(x)^4 + 4*a^2*\cosh(x))*\sinh(x)^3 + a^2*\sinh(x)^4 - 2*a^2*\cosh(x)^2 + \dots$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\coth^3(x)}{\sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(x)**3/(a+b*tanh(x)**2+c*tanh(x)**4)**(1/2),x)

[Out] Integral(coth(x)**3/sqrt(a + b*tanh(x)**2 + c*tanh(x)**4), x)

Giac [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(coth(x)^3/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2),x, algorithm="giac")

[Out] Timed out

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\coth(x)^3}{\sqrt{c \tanh(x)^4 + b \tanh(x)^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(coth(x)^3/(a + b*tanh(x)^2 + c*tanh(x)^4)^(1/2),x)

[Out] int(coth(x)^3/(a + b*tanh(x)^2 + c*tanh(x)^4)^(1/2), x)

3.205 $\int \tanh(x) \sqrt{a + b \tanh^2(x) + c \tanh^4(x)} dx$

Optimal. Leaf size=132

$$\frac{(b+2c) \tanh^{-1} \left(\frac{b+2c \tanh^2(x)}{2\sqrt{c} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} \right)}{4\sqrt{c}} + \frac{1}{2} \sqrt{a+b+c} \tanh^{-1} \left(\frac{2a+b+(b+2c) \tanh^2(x)}{2\sqrt{a+b+c} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} \right)$$

[Out] $-1/4*(b+2*c)*\operatorname{arctanh}(1/2*(b+2*c*\tanh(x)^2)/c^{(1/2)}/(a+b*\tanh(x)^2+c*\tanh(x)^4)^{(1/2)})/c^{(1/2)}+1/2*\operatorname{arctanh}(1/2*(2*a+b+(b+2*c)*\tanh(x)^2)/(a+b+c)^{(1/2)}/(a+b*\tanh(x)^2+c*\tanh(x)^4)^{(1/2)})*(a+b+c)^{(1/2)}-1/2*(a+b*\tanh(x)^2+c*\tanh(x)^4)^{(1/2)}$

Rubi [A]

time = 0.16, antiderivative size = 132, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3781, 1261, 748, 857, 635, 212, 738}

$$-\frac{1}{2} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)} - \frac{(b+2c) \tanh^{-1} \left(\frac{b+2c \tanh^2(x)}{2\sqrt{c} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} \right)}{4\sqrt{c}} + \frac{1}{2} \sqrt{a+b+c} \tanh^{-1} \left(\frac{2a+(b+2c) \tanh^2(x)+b}{2\sqrt{a+b+c} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} \right)$$

Antiderivative was successfully verified.

[In] `Int[Tanh[x]*Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4], x]`

[Out] $-1/4*((b+2*c)*\operatorname{ArcTanh}[(b+2*c*\operatorname{Tanh}[x]^2)/(2*\operatorname{Sqrt}[c]*\operatorname{Sqrt}[a+b*\operatorname{Tanh}[x]^2+c*\operatorname{Tanh}[x]^4]])/\operatorname{Sqrt}[c]+(\operatorname{Sqrt}[a+b+c]*\operatorname{ArcTanh}[(2*a+b+(b+2*c)*\operatorname{Tanh}[x]^2)/(2*\operatorname{Sqrt}[a+b+c]*\operatorname{Sqrt}[a+b*\operatorname{Tanh}[x]^2+c*\operatorname{Tanh}[x]^4])])/2-\operatorname{Sqrt}[a+b*\operatorname{Tanh}[x]^2+c*\operatorname{Tanh}[x]^4])/2$

Rule 212

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rule 635

`Int[1/Sqrt[(a_) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] := Dist[2, Subst[Int[1/(4*c - x^2), x], x, (b + 2*c*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]`

Rule 738

`Int[1/(((d_.) + (e_.)*(x_))*Sqrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]), x_Symbol] := Dist[-2, Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2`

$*a*e - b*d - (2*c*d - b*e)*x)/\text{Sqrt}[a + b*x + c*x^2]], x] /;$ FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[2*c*d - b*e, 0]

Rule 748

Int[((d_.) + (e_.)*(x_))^(m_)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[(d + e*x)^(m + 1)*((a + b*x + c*x^2)^p/(e*(m + 2*p + 1))), x] - Dist[p/(e*(m + 2*p + 1)), Int[(d + e*x)^m*Simp[b*d - 2*a*e + (2*c*d - b*e)*x, x]*(a + b*x + c*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c, d, e, m}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[2*c*d - b*e, 0] && GtQ[p, 0] && NeQ[m + 2*p + 1, 0] && (!RationalQ[m] || LtQ[m, 1]) && !ILtQ[m + 2*p, 0] && IntQuadraticQ[a, b, c, d, e, m, p, x]

Rule 857

Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Dist[g/e, Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^p, x], x] + Dist[(e*f - d*g)/e, Int[(d + e*x)^m*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, p}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && !IGtQ[m, 0]

Rule 1261

Int[(x_)*((d_) + (e_.)*(x_)^2)^(q_.)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_.), x_Symbol] := Dist[1/2, Subst[Int[(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x]

Rule 3781

Int[tan[(d_.) + (e_.)*(x_)]^(m_.)*((a_.) + (b_.)*((f_.)*tan[(d_.) + (e_.)*(x_)])^(n_.) + (c_.)*((f_.)*tan[(d_.) + (e_.)*(x_)])^(n2_.))^(p_), x_Symbol] := Dist[f/e, Subst[Int[(x/f)^m*((a + b*x^n + c*x^(2*n))^p/(f^2 + x^2)), x], x, f*Tan[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[n2, 2*n] && NeQ[b^2 - 4*a*c, 0]

Rubi steps

$$\begin{aligned}
\int \tanh(x) \sqrt{a + b \tanh^2(x) + c \tanh^4(x)} dx &= -\text{Subst} \left(\int \frac{x \sqrt{a - bx^2 + cx^4}}{1 + x^2} dx, x, i \tanh(x) \right) \\
&= - \left(\frac{1}{2} \text{Subst} \left(\int \frac{\sqrt{a - bx + cx^2}}{1 + x} dx, x, -\tanh^2(x) \right) \right) \\
&= -\frac{1}{2} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)} + \frac{1}{4} \text{Subst} \left(\int \frac{-2a - b + c}{(1 + x) \sqrt{a - bx + cx^2}} dx, x, -\tanh^2(x) \right) \\
&= -\frac{1}{2} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)} + \frac{1}{2} (-a - b - c) \text{Subst} \left(\int \frac{1}{1 + x} dx, x, -\tanh^2(x) \right) \\
&= -\frac{1}{2} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)} + (a + b + c) \text{Subst} \left(\int \frac{1}{1 + x} dx, x, -\tanh^2(x) \right) \\
&= -\frac{(b + 2c) \tanh^{-1} \left(\frac{b + 2c \tanh^2(x)}{2\sqrt{c} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} \right)}{4\sqrt{c}} + \dots
\end{aligned}$$

Mathematica [A]

time = 0.26, size = 131, normalized size = 0.99

$$\frac{1}{4} \left(\frac{(b + 2c) \tanh^{-1} \left(\frac{-b - 2c \tanh^2(x)}{2\sqrt{c} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} \right)}{\sqrt{c}} + 2\sqrt{a + b + c} \tanh^{-1} \left(\frac{2a + b + (b + 2c) \tanh^2(x)}{2\sqrt{a + b + c} \sqrt{a + b \tanh^2(x) + c \tanh^4(x)}} \right) - 2\sqrt{a + b \tanh^2(x) + c \tanh^4(x)} \right)$$

Antiderivative was successfully verified.

`[In] Integrate[Tanh[x]*Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4], x]`

```
[Out] (((b + 2*c)*ArcTanh[(-b - 2*c*Tanh[x]^2)/(2*Sqrt[c]*Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4)])/Sqrt[c] + 2*Sqrt[a + b + c]*ArcTanh[(2*a + b + (b + 2*c)*Tanh[x]^2)/(2*Sqrt[a + b + c]*Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4)]) - 2*Sqrt[a + b*Tanh[x]^2 + c*Tanh[x]^4])/4
```

Maple [C] Result contains higher order function than in optimal. Order 4 vs. order 3.

time = 0.97, size = 850, normalized size = 6.44 Too large to display

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((a+b*tanh(x)^2+c*tanh(x)^4)^(1/2)*tanh(x), x, method=_RETURNVERBOSE)`

```
[Out] -1/2*(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2)-1/8*(b+c)*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(4-2*(-b+(-4*a*c+b^2)^(1/2))/a*tanh(x)^2)^(1/2)*(4+2*(b+(-4
```



```

*a*c+b^2)^(1/2))/a*tanh(x)^2)^(1/2)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2)*Ellip
ticF(1/2*tanh(x)*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2),1/2*(-4+2*b*(b+
-4*a*c+b^2)^(1/2))/a/c)^(1/2))-1/2*(c+1/2*b)*ln((b+2*c*tanh(x)^2)/c^(1/2)+2
*(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2))/c^(1/2)-1/2*(a+b+c)*(-1/2/(a+b+c)^(1/2)
*arctanh(1/2*(b*tanh(x)^2+2*c*tanh(x)^2+2*a+b)/(a+b+c)^(1/2)/(a+b*tanh(x)^2
+c*tanh(x)^4)^(1/2))-2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(1-1/2*(-b+
-4*a*c+b^2)^(1/2))/a*tanh(x)^2)^(1/2)*(1+1/2*(b+(-4*a*c+b^2)^(1/2))/a*tanh(
x)^2)^(1/2)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2)*EllipticPi(1/2*tanh(x)*2^(1/2)
)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2),2/(-b+(-4*a*c+b^2)^(1/2))*a,(-1/2*(b+
-4*a*c+b^2)^(1/2))/a)^(1/2)*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2))-1/8*
(-b-c)*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(4-2*(-b+(-4*a*c+b^2)^(1/2)
))/a*tanh(x)^2)^(1/2)*(4+2*(b+(-4*a*c+b^2)^(1/2))/a*tanh(x)^2)^(1/2)/(a+b*t
anh(x)^2+c*tanh(x)^4)^(1/2)*EllipticF(1/2*tanh(x)*2^(1/2)*((-b+(-4*a*c+b^2)
^(1/2))/a)^(1/2),1/2*(-4+2*b*(b+(-4*a*c+b^2)^(1/2))/a/c)^(1/2))-1/2*(a+b+c)
*(-1/2/(a+b+c)^(1/2)*arctanh(1/2*(b*tanh(x)^2+2*c*tanh(x)^2+2*a+b)/(a+b+c)^(
1/2)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2))+2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)
)^(1/2)*(1-1/2*(-b+(-4*a*c+b^2)^(1/2))/a*tanh(x)^2)^(1/2)*(1+1/2*(b+(-4*a*c
+b^2)^(1/2))/a*tanh(x)^2)^(1/2)/(a+b*tanh(x)^2+c*tanh(x)^4)^(1/2)*EllipticP
i(1/2*tanh(x)*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2),2/(-b+(-4*a*c+b^2)^(
1/2))*a,(-1/2*(b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*2^(1/2)/((-b+(-4*a*c+b^2)^(1
/2))/a)^(1/2))

```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a+b*tanh(x)^2+c*tanh(x)^4)^(1/2)*tanh(x),x, algorithm="maxima")
```

```
[Out] integrate(sqrt(c*tanh(x)^4 + b*tanh(x)^2 + a)*tanh(x), x)
```

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 1827 vs. 2(108) = 216.

time = 1.42, size = 7896, normalized size = 59.82

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a+b*tanh(x)^2+c*tanh(x)^4)^(1/2)*tanh(x),x, algorithm="fricas")
```

```
[Out] [1/8*(((b + 2*c)*cosh(x)^4 + 4*(b + 2*c)*cosh(x)*sinh(x)^3 + (b + 2*c)*sinh
(x)^4 + 2*(b + 2*c)*cosh(x)^2 + 2*(3*(b + 2*c)*cosh(x)^2 + b + 2*c)*sinh(x)
^2 + 4*((b + 2*c)*cosh(x)^3 + (b + 2*c)*cosh(x))*sinh(x) + b + 2*c)*sqrt(c)
*log(((b^2 + 4*(a + 2*b)*c + 8*c^2)*cosh(x)^8 + 8*(b^2 + 4*(a + 2*b)*c + 8*
c^2)*cosh(x)*sinh(x)^7 + (b^2 + 4*(a + 2*b)*c + 8*c^2)*sinh(x)^8 + 4*(b^2 +
```

$$\begin{aligned}
& 4*a*c - 8*c^2)*\cosh(x)^6 + 4*(7*(b^2 + 4*(a + 2*b)*c + 8*c^2)*\cosh(x)^2 + \\
& b^2 + 4*a*c - 8*c^2)*\sinh(x)^6 + 8*(7*(b^2 + 4*(a + 2*b)*c + 8*c^2)*\cosh(x) \\
& ^3 + 3*(b^2 + 4*a*c - 8*c^2)*\cosh(x))*\sinh(x)^5 + 2*(3*b^2 + 4*(3*a - 2*b)* \\
& c + 24*c^2)*\cosh(x)^4 + 2*(35*(b^2 + 4*(a + 2*b)*c + 8*c^2)*\cosh(x)^4 + 30* \\
& (b^2 + 4*a*c - 8*c^2)*\cosh(x)^2 + 3*b^2 + 4*(3*a - 2*b)*c + 24*c^2)*\sinh(x) \\
& ^4 + 8*(7*(b^2 + 4*(a + 2*b)*c + 8*c^2)*\cosh(x)^5 + 10*(b^2 + 4*a*c - 8*c^2) \\
&)*\cosh(x)^3 + (3*b^2 + 4*(3*a - 2*b)*c + 24*c^2)*\cosh(x))*\sinh(x)^3 + 4*(b^2 \\
& + 4*a*c - 8*c^2)*\cosh(x)^2 + 4*(7*(b^2 + 4*(a + 2*b)*c + 8*c^2)*\cosh(x)^6 \\
& + 15*(b^2 + 4*a*c - 8*c^2)*\cosh(x)^4 + 3*(3*b^2 + 4*(3*a - 2*b)*c + 24*c^2) \\
&)*\cosh(x)^2 + b^2 + 4*a*c - 8*c^2)*\sinh(x)^2 - 4*\sqrt{2}*((b + 2*c)*\cosh(x) \\
& ^4 + 4*(b + 2*c)*\cosh(x)*\sinh(x)^3 + (b + 2*c)*\sinh(x)^4 + 2*(b - 2*c)*\cosh \\
& (x)^2 + 2*(3*(b + 2*c)*\cosh(x)^2 + b - 2*c)*\sinh(x)^2 + 4*((b + 2*c)*\cosh(x) \\
&)^3 + (b - 2*c)*\cosh(x))*\sinh(x) + b + 2*c)*\sqrt{c}*\sqrt{((a + b + c)*\cosh(x) \\
& ^4 + (a + b + c)*\sinh(x)^4 + 4*(a - c)*\cosh(x)^2 + 2*(3*(a + b + c)*\cosh(x) \\
& ^2 + 2*a - 2*c)*\sinh(x)^2 + 3*a - b + 3*c)/(\cosh(x)^4 - 4*\cosh(x)^3*\sinh(x) \\
& + 6*\cosh(x)^2*\sinh(x)^2 - 4*\cosh(x)*\sinh(x)^3 + \sinh(x)^4)) + b^2 + 4*(a \\
& + 2*b)*c + 8*c^2 + 8*((b^2 + 4*(a + 2*b)*c + 8*c^2)*\cosh(x)^7 + 3*(b^2 + 4 \\
& *a*c - 8*c^2)*\cosh(x)^5 + (3*b^2 + 4*(3*a - 2*b)*c + 24*c^2)*\cosh(x)^3 + (b \\
& ^2 + 4*a*c - 8*c^2)*\cosh(x))*\sinh(x))/(\cosh(x)^8 + 8*\cosh(x)*\sinh(x)^7 + \sinh(x)^8 \\
& + 4*(7*\cosh(x)^2 + 1)*\sinh(x)^6 + 4*\cosh(x)^6 + 8*(7*\cosh(x)^3 + 3* \\
& \cosh(x))*\sinh(x)^5 + 2*(35*\cosh(x)^4 + 30*\cosh(x)^2 + 3)*\sinh(x)^4 + 6*\cosh \\
& (x)^4 + 8*(7*\cosh(x)^5 + 10*\cosh(x)^3 + 3*\cosh(x))*\sinh(x)^3 + 4*(7*\cosh(x) \\
& ^6 + 15*\cosh(x)^4 + 9*\cosh(x)^2 + 1)*\sinh(x)^2 + 4*\cosh(x)^2 + 8*(\cosh(x)^7 \\
& + 3*\cosh(x)^5 + 3*\cosh(x)^3 + \cosh(x))*\sinh(x) + 1)) + 2*(c*\cosh(x)^4 + 4* \\
& c*\cosh(x)*\sinh(x)^3 + c*\sinh(x)^4 + 2*c*\cosh(x)^2 + 2*(3*c*\cosh(x)^2 + c)*\sinh(x)^2 \\
& + 4*(c*\cosh(x)^3 + c*\cosh(x))*\sinh(x) + c)*\sqrt{a + b + c}*\log(((a \\
& ^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*\cosh(x)^8 + 8*(a^2 + 2*a*b + b^2 + 2* \\
& (a + b)*c + c^2)*\cosh(x)*\sinh(x)^7 + (a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2) \\
&)*\sinh(x)^8 + 4*(a^2 + a*b - b*c - c^2)*\cosh(x)^6 + 4*(7*(a^2 + 2*a*b + b^2 \\
& + 2*(a + b)*c + c^2)*\cosh(x)^2 + a^2 + a*b - b*c - c^2)*\sinh(x)^6 + 8*(7*(\\
& a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*\cosh(x)^3 + 3*(a^2 + a*b - b*c - c^2) \\
&)*\cosh(x))*\sinh(x)^5 + 2*(3*a^2 + 2*a*b + 2*(a + b)*c + 3*c^2)*\cosh(x)^4 + \\
& 2*(35*(a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*\cosh(x)^4 + 30*(a^2 + a*b - b* \\
& c - c^2)*\cosh(x)^2 + 3*a^2 + 2*a*b + 2*(a + b)*c + 3*c^2)*\sinh(x)^4 + 8*(7 \\
& *(a^2 + 2*a*b + b^2 + 2*(a + b)*c + c^2)*\cosh(x)^5 + 10*(a^2 + a*b - b*c - \\
& c^2)*\cosh(x)^3 + (3*a^2 + 2*a*b + 2*(a + b)*c + 3*c^2)*\cosh(x))*\sinh(x)^3 + \\
& 4*(a^2 + a*b - b*c - c^2)*\cosh(x)^2 + 4*(7*(a^2 + 2*a*b + b^2 + 2*(a + b)* \\
& c + c^2)*\cosh(x)^6 + 15*(a^2 + a*b - b*c - c^2)*\cosh(x)^4 + 3*(3*a^2 + 2*a* \\
& b + 2*(a + b)*c + 3*c^2)*\cosh(x)^2 + a^2 + a*b - b*c - c^2)*\sinh(x)^2 + \sqrt{ \\
& 2}*((a + b + c)*\cosh(x)^4 + 4*(a + b + c)*\cosh(x)*\sinh(x)^3 + (a + b + c) \\
&)*\sinh(x)^4 + 2*(a - c)*\cosh(x)^2 + 2*(3*(a + b + c)*\cosh(x)^2 + a - c)*\sinh \\
& (x)^2 + 4*((a + b + c)*\cosh(x)^3 + (a - c)*\cosh(x))*\sinh(x) + a + b + c)*\sqrt{ \\
& a + b + c}*\sqrt{((a + b + c)*\cosh(x)^4 + (a + b + c)*\sinh(x)^4 + 4*(a - \\
& c)*\cosh(x)^2 + 2*(3*(a + b + c)*\cosh(x)^2 + 2*a - 2*c)*\sinh(x)^2 + 3*a - b \\
& + 3*c)/(\cosh(x)^4 - 4*\cosh(x)^3*\sinh(x) + 6*\cosh(x)^2*\sinh(x)^2 - 4*\cosh(x)
\end{aligned}$$

3.206 $\int e^{a+bx} \tanh^4(a+bx) dx$

Optimal. Leaf size=107

$$\frac{e^{a+bx}}{b} + \frac{8e^{a+bx}}{3b(1+e^{2a+2bx})^3} - \frac{14e^{a+bx}}{3b(1+e^{2a+2bx})^2} + \frac{5e^{a+bx}}{b(1+e^{2a+2bx})} - \frac{3\text{ArcTan}(e^{a+bx})}{b}$$

[Out] $\exp(b*x+a)/b+8/3*\exp(b*x+a)/b/(1+\exp(2*b*x+2*a))^3-14/3*\exp(b*x+a)/b/(1+\exp(2*b*x+2*a))^2+5*\exp(b*x+a)/b/(1+\exp(2*b*x+2*a))-3*\arctan(\exp(b*x+a))/b$

Rubi [A]

time = 0.05, antiderivative size = 107, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$,

Rules used = {2320, 398, 1272, 1171, 393, 209}

$$-\frac{3\text{ArcTan}(e^{a+bx})}{b} + \frac{e^{a+bx}}{b} + \frac{5e^{a+bx}}{b(e^{2a+2bx}+1)} - \frac{14e^{a+bx}}{3b(e^{2a+2bx}+1)^2} + \frac{8e^{a+bx}}{3b(e^{2a+2bx}+1)^3}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{(a + b*x)}*\text{Tanh}[a + b*x]^4, x]$

[Out] $E^{(a + b*x)}/b + (8*E^{(a + b*x)})/(3*b*(1 + E^{(2*a + 2*b*x)})^3) - (14*E^{(a + b*x)})/(3*b*(1 + E^{(2*a + 2*b*x)})^2) + (5*E^{(a + b*x)})/(b*(1 + E^{(2*a + 2*b*x)})) - (3*\text{ArcTan}[E^{(a + b*x)}])/b$

Rule 209

$\text{Int}[(a + (b*x)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2]))*\text{ArcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$

Rule 393

$\text{Int}[(a + (b*x)^n)^p*((c + (d*x)^n)), x_Symbol] \rightarrow \text{Simp}[(-b*c - a*d)*x*((a + b*x^n)^{p+1}/(a*b*n*(p+1))), x] - \text{Dist}[(a*d - b*c*(n*(p+1) + 1))/(a*b*n*(p+1)), \text{Int}[(a + b*x^n)^{p+1}, x], x] /; \text{FreeQ}\{a, b, c, d, n, p, x\} \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ (\text{LtQ}[p, -1] \ || \ \text{ILtQ}[1/n + p, 0])$

Rule 398

$\text{Int}[(a + (b*x)^n)^p*((c + (d*x)^n)^q), x_Symbol] \rightarrow \text{Int}[\text{PolynomialDivide}[(a + b*x^n)^p, (c + d*x^n)^{-q}], x] /; \text{FreeQ}\{a, b, c, d, x\} \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{ILtQ}[q, 0] \ \&\& \ \text{GeQ}[p, -q]$

Rule 1171

```
Int[((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_),
x_Symbol] := With[{Qx = PolynomialQuotient[(a + b*x^2 + c*x^4)^p, d + e*x^2
, x], R = Coeff[PolynomialRemainder[(a + b*x^2 + c*x^4)^p, d + e*x^2, x], x
, 0]}, Simp[(-R)*x*((d + e*x^2)^(q + 1)/(2*d*(q + 1))), x] + Dist[1/(2*d*(q
+ 1)), Int[(d + e*x^2)^(q + 1)*ExpandToSum[2*d*(q + 1)*Qx + R*(2*q + 3), x
], x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2
- b*d*e + a*e^2, 0] && IGtQ[p, 0] && LtQ[q, -1]
```

Rule 1272

```
Int[(x_)^(m_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Sy
mbol] := Simp[(-d)^(m/2 - 1)*(c*d^2 + a*e^2)^p*x*((d + e*x^2)^(q + 1)/(2*e^
(2*p + m/2)*(q + 1))), x] + Dist[1/(2*e^(2*p + m/2)*(q + 1)), Int[(d + e*x^
2)^(q + 1)*ExpandToSum[Together[(1/(d + e*x^2))*(2*e^(2*p + m/2)*(q + 1)*x^
m*(a + c*x^4)^p - (-d)^(m/2 - 1)*(c*d^2 + a*e^2)^p*(d + e*(2*q + 3)*x^2)],
x], x], x] /; FreeQ[{a, c, d, e}, x] && IGtQ[p, 0] && ILtQ[q, -1] && IGtQ[
m/2, 0]
```

Rule 2320

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x]
, Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; Functi
onOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[
{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))*
(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rubi steps

$$\begin{aligned}
\int e^{a+bx} \tanh^4(a+bx) dx &= \frac{\text{Subst}\left(\int \frac{(1-x^2)^4}{(1+x^2)^4} dx, x, e^{a+bx}\right)}{b} \\
&= \frac{\text{Subst}\left(\int \left(1 - \frac{8x^2(1+x^4)}{(1+x^2)^4}\right) dx, x, e^{a+bx}\right)}{b} \\
&= \frac{e^{a+bx}}{b} - \frac{8\text{Subst}\left(\int \frac{x^2(1+x^4)}{(1+x^2)^4} dx, x, e^{a+bx}\right)}{b} \\
&= \frac{e^{a+bx}}{b} + \frac{8e^{a+bx}}{3b(1+e^{2a+2bx})^3} + \frac{4\text{Subst}\left(\int \frac{-2+6x^2-6x^4}{(1+x^2)^3} dx, x, e^{a+bx}\right)}{3b} \\
&= \frac{e^{a+bx}}{b} + \frac{8e^{a+bx}}{3b(1+e^{2a+2bx})^3} - \frac{14e^{a+bx}}{3b(1+e^{2a+2bx})^2} - \frac{\text{Subst}\left(\int \frac{-6+24x^2}{(1+x^2)^2} dx, x, e^{a+bx}\right)}{3b} \\
&= \frac{e^{a+bx}}{b} + \frac{8e^{a+bx}}{3b(1+e^{2a+2bx})^3} - \frac{14e^{a+bx}}{3b(1+e^{2a+2bx})^2} + \frac{5e^{a+bx}}{b(1+e^{2a+2bx})} - \frac{3\text{Subst}\left(\int \frac{1}{1+x^2}\right)}{b} \\
&= \frac{e^{a+bx}}{b} + \frac{8e^{a+bx}}{3b(1+e^{2a+2bx})^3} - \frac{14e^{a+bx}}{3b(1+e^{2a+2bx})^2} + \frac{5e^{a+bx}}{b(1+e^{2a+2bx})} - \frac{3 \tan^{-1}(e^{a+bx})}{b}
\end{aligned}$$

Mathematica [A]

time = 0.12, size = 76, normalized size = 0.71

$$\frac{e^{a+bx} (12 + 25e^{2(a+bx)} + 24e^{4(a+bx)} + 3e^{6(a+bx)})}{3b(1+e^{2(a+bx)})^3} - \frac{3\text{ArcTan}(e^{a+bx})}{b}$$

Antiderivative was successfully verified.

`[In] Integrate[E^(a + b*x)*Tanh[a + b*x]^4, x]`

```
[Out] (E^(a + b*x)*(12 + 25*E^(2*(a + b*x)) + 24*E^(4*(a + b*x)) + 3*E^(6*(a + b*x))))/(3*b*(1 + E^(2*(a + b*x)))^3) - (3*ArcTan[E^(a + b*x)])/b
```

Maple [C] Result contains complex when optimal does not.

time = 1.45, size = 92, normalized size = 0.86

method	result	size
risch	$\frac{e^{bx+a}}{b} + \frac{e^{bx+a}(15e^{4bx+4a}+16e^{2bx+2a}+9)}{3b(e^{2bx+2a}+1)^3} + \frac{3i \ln(e^{bx+a}-i)}{2b} - \frac{3i \ln(e^{bx+a}+i)}{2b}$	92

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(exp(b*x+a)*tanh(b*x+a)^4, x, method=_RETURNVERBOSE)`

[Out] $\exp(b*x+a)/b+1/3*\exp(b*x+a)*(15*\exp(4*b*x+4*a)+16*\exp(2*b*x+2*a)+9)/b/(\exp(2*b*x+2*a)+1)^3+3/2*I/b*\ln(\exp(b*x+a)-I)-3/2*I/b*\ln(\exp(b*x+a)+I)$

Maxima [A]

time = 0.47, size = 94, normalized size = 0.88

$$-\frac{3 \arctan(e^{(bx+a)})}{b} + \frac{e^{(bx+a)}}{b} + \frac{15e^{(5bx+5a)} + 16e^{(3bx+3a)} + 9e^{(bx+a)}}{3b(e^{(6bx+6a)} + 3e^{(4bx+4a)} + 3e^{(2bx+2a)} + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(b*x+a)*tanh(b*x+a)^4,x, algorithm="maxima")`

[Out] $-3*\arctan(e^{(b*x + a)})/b + e^{(b*x + a)}/b + 1/3*(15*e^{(5*b*x + 5*a)} + 16*e^{(3*b*x + 3*a)} + 9*e^{(b*x + a)})/(b*(e^{(6*b*x + 6*a)} + 3*e^{(4*b*x + 4*a)} + 3*e^{(2*b*x + 2*a)} + 1))$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 604 vs. 2(95) = 190.

time = 0.37, size = 604, normalized size = 5.64

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(b*x+a)*tanh(b*x+a)^4,x, algorithm="fricas")`

[Out] $1/3*(3*\cosh(b*x + a)^7 + 21*\cosh(b*x + a)*\sinh(b*x + a)^6 + 3*\sinh(b*x + a)^7 + 3*(21*\cosh(b*x + a)^2 + 8)*\sinh(b*x + a)^5 + 24*\cosh(b*x + a)^5 + 15*(7*\cosh(b*x + a)^3 + 8*\cosh(b*x + a))*\sinh(b*x + a)^4 + 5*(21*\cosh(b*x + a)^4 + 48*\cosh(b*x + a)^2 + 5)*\sinh(b*x + a)^3 + 25*\cosh(b*x + a)^3 + 3*(21*\cosh(b*x + a)^5 + 80*\cosh(b*x + a)^3 + 25*\cosh(b*x + a))*\sinh(b*x + a)^2 - 9*(\cosh(b*x + a)^6 + 6*\cosh(b*x + a)*\sinh(b*x + a)^5 + \sinh(b*x + a)^6 + 3*(5*\cosh(b*x + a)^2 + 1)*\sinh(b*x + a)^4 + 3*\cosh(b*x + a)^4 + 4*(5*\cosh(b*x + a)^3 + 3*\cosh(b*x + a))*\sinh(b*x + a)^3 + 3*(5*\cosh(b*x + a)^4 + 6*\cosh(b*x + a)^2 + 1)*\sinh(b*x + a)^2 + 3*\cosh(b*x + a)^2 + 6*(\cosh(b*x + a)^5 + 2*\cosh(b*x + a)^3 + \cosh(b*x + a))*\sinh(b*x + a) + 1)*\arctan(\cosh(b*x + a) + \sinh(b*x + a)) + 3*(7*\cosh(b*x + a)^6 + 40*\cosh(b*x + a)^4 + 25*\cosh(b*x + a)^2 + 4)*\sinh(b*x + a) + 12*\cosh(b*x + a))/(b*\cosh(b*x + a)^6 + 6*b*\cosh(b*x + a)*\sinh(b*x + a)^5 + b*\sinh(b*x + a)^6 + 3*b*\cosh(b*x + a)^4 + 3*(5*b*\cosh(b*x + a)^2 + b)*\sinh(b*x + a)^4 + 4*(5*b*\cosh(b*x + a)^3 + 3*b*\cosh(b*x + a))*\sinh(b*x + a)^3 + 3*b*\cosh(b*x + a)^2 + 3*(5*b*\cosh(b*x + a)^4 + 6*b*\cosh(b*x + a)^2 + b)*\sinh(b*x + a)^2 + 6*(b*\cosh(b*x + a)^5 + 2*b*\cosh(b*x + a)^3 + b*\cosh(b*x + a))*\sinh(b*x + a) + b)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$e^a \int e^{bx} \tanh^4(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(b*x+a)*tanh(b*x+a)**4,x)

[Out] exp(a)*Integral(exp(b*x)*tanh(a + b*x)**4, x)

Giac [A]

time = 0.42, size = 68, normalized size = 0.64

$$\frac{15 e^{(5bx+5a)} + 16 e^{(3bx+3a)} + 9 e^{(bx+a)} - 9 \arctan(e^{(bx+a)}) + 3 e^{(bx+a)}}{(e^{(2bx+2a)}+1)^3} \frac{1}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(b*x+a)*tanh(b*x+a)^4,x, algorithm="giac")

[Out] 1/3*((15*e^(5*b*x + 5*a) + 16*e^(3*b*x + 3*a) + 9*e^(b*x + a))/(e^(2*b*x + 2*a) + 1)^3 - 9*arctan(e^(b*x + a)) + 3*e^(b*x + a))/b

Mupad [B]

time = 0.10, size = 155, normalized size = 1.45

$$\frac{e^{a+bx}}{b} - \frac{3 \operatorname{atan}\left(\frac{e^{bx} e^a \sqrt{b^2}}{b}\right)}{\sqrt{b^2}} + \frac{\frac{4e^{a+bx}}{3b} + \frac{4e^{5a+5bx}}{3b}}{3e^{2a+2bx} + 3e^{4a+4bx} + e^{6a+6bx} + 1} - \frac{2e^{a+bx}}{b(2e^{2a+2bx} + e^{4a+4bx} + 1)} + \frac{11e^{a+bx}}{3b(e^{2a+2bx} + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(a + b*x)*tanh(a + b*x)^4,x)

[Out] exp(a + b*x)/b - (3*atan((exp(b*x)*exp(a)*(b^2)^(1/2))/b))/(b^2)^(1/2) + ((4*exp(a + b*x))/(3*b) + (4*exp(5*a + 5*b*x))/(3*b))/(3*exp(2*a + 2*b*x) + 3*exp(4*a + 4*b*x) + exp(6*a + 6*b*x) + 1) - (2*exp(a + b*x))/(b*(2*exp(2*a + 2*b*x) + exp(4*a + 4*b*x) + 1)) + (11*exp(a + b*x))/(3*b*(exp(2*a + 2*b*x) + 1))

3.207 $\int e^{a+bx} \tanh^3(a+bx) dx$

Optimal. Leaf size=77

$$\frac{e^{a+bx}}{b} - \frac{2e^{a+bx}}{b(1+e^{2a+2bx})^2} + \frac{3e^{a+bx}}{b(1+e^{2a+2bx})} - \frac{3\text{ArcTan}(e^{a+bx})}{b}$$

[Out] $\exp(b*x+a)/b - 2*\exp(b*x+a)/b/(1+\exp(2*b*x+2*a))^2 + 3*\exp(b*x+a)/b/(1+\exp(2*b*x+2*a)) - 3*\arctan(\exp(b*x+a))/b$

Rubi [A]

time = 0.03, antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {2320, 398, 1172, 12, 294, 209}

$$-\frac{3\text{ArcTan}(e^{a+bx})}{b} + \frac{e^{a+bx}}{b} + \frac{3e^{a+bx}}{b(e^{2a+2bx}+1)} - \frac{2e^{a+bx}}{b(e^{2a+2bx}+1)^2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{(a + b*x)}*\text{Tanh}[a + b*x]^3, x]$

[Out] $E^{(a + b*x)}/b - (2*E^{(a + b*x)})/(b*(1 + E^{(2*a + 2*b*x)})^2) + (3*E^{(a + b*x)})/(b*(1 + E^{(2*a + 2*b*x)})) - (3*\text{ArcTan}[E^{(a + b*x)}])/b$

Rule 12

$\text{Int}[(a_*)(u_), x_Symbol] \rightarrow \text{Dist}[a, \text{Int}[u, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{Match} Q[u, (b_)*(v_)] /; \text{FreeQ}[b, x]$

Rule 209

$\text{Int}[(a_*) + (b_)*(x_)^2]^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2]))*\text{ArcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$

Rule 294

$\text{Int}[(c_)*(x_)^{(m_)}*((a_*) + (b_)*(x_)^{(n_)})^{(p_)}], x_Symbol] \rightarrow \text{Simp}[c^{(n-1)}*(c*x)^{(m-n+1)}*((a + b*x^n)^{(p+1)}/(b*n*(p+1))), x] - \text{Dist}[c^n*((m-n+1)/(b*n*(p+1))), \text{Int}[(c*x)^{(m-n)}*(a + b*x^n)^{(p+1)}, x], x] /; \text{FreeQ}[\{a, b, c\}, x] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ \text{GtQ}[m+1, n] \ \&\& \ !\text{LtQ}[(m+n*(p+1)+1)/n, 0] \ \&\& \ \text{IntBinomialQ}[a, b, c, n, m, p, x]$

Rule 398

$\text{Int}[(a_*) + (b_)*(x_)^{(n_)})^{(p_)}*((c_*) + (d_)*(x_)^{(n_)})^{(q_)}], x_Symbol] \rightarrow \text{Int}[\text{PolynomialDivide}[(a + b*x^n)^p, (c + d*x^n)^{-q}], x] /; \text{FreeQ}[\{a$

, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]

Rule 1172

```
Int[((d_) + (e_.)*(x_)^2)^(q_)*((a_) + (c_.)*(x_)^4)^(p_), x_Symbol] := With[
  {Qx = PolynomialQuotient[(a + c*x^4)^p, d + e*x^2, x], R = Coeff[PolynomialRemainder[
    (a + c*x^4)^p, d + e*x^2, x], x, 0]}, Simp[(-R)*x*((d + e*x^2)^(q + 1)/(2*d*(q + 1))), x] + Dist[1/(2*d*(q + 1)), Int[(d + e*x^2)^(q + 1)*ExpandToSum[2*d*(q + 1)*Qx + R*(2*q + 3), x], x], x]] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && IGtQ[p, 0] && LtQ[q, -1]
```

Rule 2320

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rubi steps

$$\begin{aligned}
 \int e^{a+bx} \tanh^3(a+bx) dx &= \frac{\text{Subst}\left(\int \frac{(-1+x^2)^3}{(1+x^2)^3} dx, x, e^{a+bx}\right)}{b} \\
 &= \frac{\text{Subst}\left(\int \left(1 - \frac{2(1+3x^4)}{(1+x^2)^3}\right) dx, x, e^{a+bx}\right)}{b} \\
 &= \frac{e^{a+bx}}{b} - \frac{2\text{Subst}\left(\int \frac{1+3x^4}{(1+x^2)^3} dx, x, e^{a+bx}\right)}{b} \\
 &= \frac{e^{a+bx}}{b} - \frac{2e^{a+bx}}{b(1+e^{2a+2bx})^2} + \frac{\text{Subst}\left(\int -\frac{12x^2}{(1+x^2)^2} dx, x, e^{a+bx}\right)}{2b} \\
 &= \frac{e^{a+bx}}{b} - \frac{2e^{a+bx}}{b(1+e^{2a+2bx})^2} - \frac{6\text{Subst}\left(\int \frac{x^2}{(1+x^2)^2} dx, x, e^{a+bx}\right)}{b} \\
 &= \frac{e^{a+bx}}{b} - \frac{2e^{a+bx}}{b(1+e^{2a+2bx})^2} + \frac{3e^{a+bx}}{b(1+e^{2a+2bx})} - \frac{3\text{Subst}\left(\int \frac{1}{1+x^2} dx, x, e^{a+bx}\right)}{b} \\
 &= \frac{e^{a+bx}}{b} - \frac{2e^{a+bx}}{b(1+e^{2a+2bx})^2} + \frac{3e^{a+bx}}{b(1+e^{2a+2bx})} - \frac{3 \tan^{-1}(e^{a+bx})}{b}
 \end{aligned}$$

Mathematica [A]


```
[Out] (cosh(b*x + a)^5 + 5*cosh(b*x + a)*sinh(b*x + a)^4 + sinh(b*x + a)^5 + 5*(2
*cosh(b*x + a)^2 + 1)*sinh(b*x + a)^3 + 5*cosh(b*x + a)^3 + 5*(2*cosh(b*x +
a)^3 + 3*cosh(b*x + a))*sinh(b*x + a)^2 - 3*(cosh(b*x + a)^4 + 4*cosh(b*x
+ a)*sinh(b*x + a)^3 + sinh(b*x + a)^4 + 2*(3*cosh(b*x + a)^2 + 1)*sinh(b*x
+ a)^2 + 2*cosh(b*x + a)^2 + 4*(cosh(b*x + a)^3 + cosh(b*x + a))*sinh(b*x
+ a) + 1)*arctan(cosh(b*x + a) + sinh(b*x + a)) + (5*cosh(b*x + a)^4 + 15*c
osh(b*x + a)^2 + 2)*sinh(b*x + a) + 2*cosh(b*x + a))/(b*cosh(b*x + a)^4 + 4
*b*cosh(b*x + a)*sinh(b*x + a)^3 + b*sinh(b*x + a)^4 + 2*b*cosh(b*x + a)^2
+ 2*(3*b*cosh(b*x + a)^2 + b)*sinh(b*x + a)^2 + 4*(b*cosh(b*x + a)^3 + b*co
sh(b*x + a))*sinh(b*x + a) + b)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$e^a \int e^{bx} \tanh^3(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(b*x+a)*tanh(b*x+a)**3,x)
```

```
[Out] exp(a)*Integral(exp(b*x)*tanh(a + b*x)**3, x)
```

Giac [A]

time = 0.44, size = 52, normalized size = 0.68

$$\frac{3e^{(3bx+3a)+e^{(bx+a)}}}{(e^{(2bx+2a)+1})^2} - 3 \arctan(e^{(bx+a)}) + e^{(bx+a)}$$

b

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(b*x+a)*tanh(b*x+a)^3,x, algorithm="giac")
```

```
[Out] ((3*e^(3*b*x + 3*a) + e^(b*x + a))/(e^(2*b*x + 2*a) + 1)^2 - 3*arctan(e^(b*
x + a)) + e^(b*x + a))/b
```

Mupad [B]

time = 0.08, size = 93, normalized size = 1.21

$$\frac{e^{a+bx}}{b} - \frac{3 \operatorname{atan}\left(\frac{e^{bx} e^a \sqrt{b^2}}{b}\right)}{\sqrt{b^2}} - \frac{2e^{a+bx}}{b(2e^{2a+2bx} + e^{4a+4bx} + 1)} + \frac{3e^{a+bx}}{b(e^{2a+2bx} + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(exp(a + b*x)*tanh(a + b*x)^3,x)
```

```
[Out] exp(a + b*x)/b - (3*atan((exp(b*x)*exp(a)*(b^2)^(1/2))/b))/(b^2)^(1/2) - (2
*exp(a + b*x))/(b*(2*exp(2*a + 2*b*x) + exp(4*a + 4*b*x) + 1)) + (3*exp(a +
b*x))/(b*(exp(2*a + 2*b*x) + 1))
```

3.208 $\int e^{a+bx} \tanh^2(a+bx) dx$

Optimal. Leaf size=51

$$\frac{e^{a+bx}}{b} + \frac{2e^{a+bx}}{b(1+e^{2a+2bx})} - \frac{2\text{ArcTan}(e^{a+bx})}{b}$$

[Out] $\exp(b*x+a)/b+2*\exp(b*x+a)/b/(1+\exp(2*b*x+2*a))-2*\arctan(\exp(b*x+a))/b$

Rubi [A]

time = 0.02, antiderivative size = 51, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2320, 398, 294, 209}

$$-\frac{2\text{ArcTan}(e^{a+bx})}{b} + \frac{e^{a+bx}}{b} + \frac{2e^{a+bx}}{b(e^{2a+2bx} + 1)}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{(a + b*x)*\text{Tanh}[a + b*x]^2, x]$

[Out] $E^{(a + b*x)/b} + (2*E^{(a + b*x)})/(b*(1 + E^{(2*a + 2*b*x)})) - (2*\text{ArcTan}[E^{(a + b*x)}])/b$

Rule 209

$\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \text{ :> } \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2]))*\text{ArcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] \text{ /; } \text{FreeQ}\{a, b, x\} \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$

Rule 294

$\text{Int}[(c_)*(x_)^{(m_)}*((a_ + (b_)*(x_)^{(n_)})^{(p_)}), x_Symbol] \text{ :> } \text{Simp}[c^{(n-1)}*(c*x)^{(m-n+1)}*((a + b*x^n)^{(p+1)}/(b*n*(p+1))), x] - \text{Dist}[c^n*((m-n+1)/(b*n*(p+1))), \text{Int}[(c*x)^{(m-n)}*(a + b*x^n)^{(p+1)}, x], x] \text{ /; } \text{FreeQ}\{a, b, c\}, x \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ \text{GtQ}[m+1, n] \ \&\& \ !\text{LtQ}[(m+n*(p+1)+1)/n, 0] \ \&\& \ \text{IntBinomialQ}[a, b, c, n, m, p, x]$

Rule 398

$\text{Int}[(a_ + (b_)*(x_)^{(n_)})^{(p_)}*((c_ + (d_)*(x_)^{(n_)})^{(q_)}), x_Symbol] \text{ :> } \text{Int}[\text{PolynomialDivide}[(a + b*x^n)^p, (c + d*x^n)^{-q}], x] \text{ /; } \text{FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{ILtQ}[q, 0] \ \&\& \ \text{GeQ}[p, -q]$

Rule 2320

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x]
, Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Functi
onOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[
{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*
(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rubi steps

$$\begin{aligned} \int e^{a+bx} \tanh^2(a+bx) dx &= \frac{\text{Subst}\left(\int \frac{(1-x^2)^2}{(1+x^2)^2} dx, x, e^{a+bx}\right)}{b} \\ &= \frac{\text{Subst}\left(\int \left(1 - \frac{4x^2}{(1+x^2)^2}\right) dx, x, e^{a+bx}\right)}{b} \\ &= \frac{e^{a+bx}}{b} - \frac{4\text{Subst}\left(\int \frac{x^2}{(1+x^2)^2} dx, x, e^{a+bx}\right)}{b} \\ &= \frac{e^{a+bx}}{b} + \frac{2e^{a+bx}}{b(1+e^{2a+2bx})} - \frac{2\text{Subst}\left(\int \frac{1}{1+x^2} dx, x, e^{a+bx}\right)}{b} \\ &= \frac{e^{a+bx}}{b} + \frac{2e^{a+bx}}{b(1+e^{2a+2bx})} - \frac{2 \tan^{-1}(e^{a+bx})}{b} \end{aligned}$$

Mathematica [A]

time = 0.07, size = 40, normalized size = 0.78

$$\frac{e^{a+bx} \left(1 + \frac{2}{1+e^{2(a+bx)}}\right) - 2\text{ArcTan}(e^{a+bx})}{b}$$

Antiderivative was successfully verified.

[In] Integrate[E^(a + b*x)*Tanh[a + b*x]^2, x]

[Out] (E^(a + b*x)*(1 + 2/(1 + E^(2*(a + b*x)))) - 2*ArcTan[E^(a + b*x)])/b

Maple [A]

time = 0.45, size = 48, normalized size = 0.94

method	result	size
derivativedivides	$\frac{\frac{\sinh^2(bx+a)}{\cosh(bx+a)} + \frac{2}{\cosh(bx+a)} + \sinh(bx+a) - 2 \arctan(e^{bx+a})}{b}$	48
default	$\frac{\frac{\sinh^2(bx+a)}{\cosh(bx+a)} + \frac{2}{\cosh(bx+a)} + \sinh(bx+a) - 2 \arctan(e^{bx+a})}{b}$	48

risch	$\frac{e^{bx+a}}{b} + \frac{2e^{bx+a}}{b(e^{2bx+2a}+1)} + \frac{i \ln(e^{bx+a}-i)}{b} - \frac{i \ln(e^{bx+a}+i)}{b}$	68
-------	--	----

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(b*x+a)*tanh(b*x+a)^2,x,method=_RETURNVERBOSE)`

[Out] `1/b*(sinh(b*x+a)^2/cosh(b*x+a)+2/cosh(b*x+a)+sinh(b*x+a)-2*arctan(exp(b*x+a)))`

Maxima [A]

time = 0.46, size = 47, normalized size = 0.92

$$-\frac{2 \arctan(e^{(bx+a)})}{b} + \frac{e^{(bx+a)}}{b} + \frac{2e^{(bx+a)}}{b(e^{2bx+2a}+1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(b*x+a)*tanh(b*x+a)^2,x, algorithm="maxima")`

[Out] `-2*arctan(e^(b*x + a))/b + e^(b*x + a)/b + 2*e^(b*x + a)/(b*(e^(2*b*x + 2*a) + 1))`

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 147 vs. $2(47) = 94$.

time = 0.33, size = 147, normalized size = 2.88

$$\frac{\cosh(bx+a)^3 + 3 \cosh(bx+a) \sinh(bx+a)^2 + \sinh(bx+a)^3 - 2(\cosh(bx+a)^2 + 2 \cosh(bx+a) \sinh(bx+a) + \sinh(bx+a)^2 + 1) \arctan(\cosh(bx+a) + \sinh(bx+a)) + 3(\cosh(bx+a)^2 + 1) \sinh(bx+a) + 3 \cosh(bx+a)}{b \cosh(bx+a)^2 + 2b \cosh(bx+a) \sinh(bx+a) + b \sinh(bx+a)^2 + b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(b*x+a)*tanh(b*x+a)^2,x, algorithm="fricas")`

[Out] `(cosh(b*x + a)^3 + 3*cosh(b*x + a)*sinh(b*x + a)^2 + sinh(b*x + a)^3 - 2*(cosh(b*x + a)^2 + 2*cosh(b*x + a)*sinh(b*x + a) + sinh(b*x + a)^2 + 1)*arctan(cosh(b*x + a) + sinh(b*x + a)) + 3*(cosh(b*x + a)^2 + 1)*sinh(b*x + a) + 3*cosh(b*x + a))/(b*cosh(b*x + a)^2 + 2*b*cosh(b*x + a)*sinh(b*x + a) + b*sinh(b*x + a)^2 + b)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$e^a \int e^{bx} \tanh^2(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(b*x+a)*tanh(b*x+a)**2,x)`

[Out] `exp(a)*Integral(exp(b*x)*tanh(a + b*x)**2, x)`

Giac [A]

time = 0.40, size = 41, normalized size = 0.80

$$\frac{\frac{2e^{(bx+a)}}{e^{(2bx+2a)}+1} - 2 \arctan(e^{(bx+a)}) + e^{(bx+a)}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(b*x+a)*tanh(b*x+a)^2,x, algorithm="giac")

[Out] (2*e^(b*x + a)/(e^(2*b*x + 2*a) + 1) - 2*arctan(e^(b*x + a)) + e^(b*x + a))
/b

Mupad [B]

time = 1.07, size = 58, normalized size = 1.14

$$\frac{e^{a+bx}}{b} - \frac{2 \operatorname{atan}\left(\frac{e^{bx} e^a \sqrt{b^2}}{b}\right)}{\sqrt{b^2}} + \frac{2e^{a+bx}}{b(e^{2a+2bx} + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(a + b*x)*tanh(a + b*x)^2,x)

[Out] exp(a + b*x)/b - (2*atan((exp(b*x)*exp(a)*(b^2)^(1/2))/b))/(b^2)^(1/2) + (2*exp(a + b*x))/(b*(exp(2*a + 2*b*x) + 1))

3.209 $\int e^{a+bx} \tanh(a+bx) dx$

Optimal. Leaf size=25

$$\frac{e^{a+bx}}{b} - \frac{2\text{ArcTan}(e^{a+bx})}{b}$$

[Out] exp(b*x+a)/b-2*arctan(exp(b*x+a))/b

Rubi [A]

time = 0.01, antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {2320, 396, 209}

$$\frac{e^{a+bx}}{b} - \frac{2\text{ArcTan}(e^{a+bx})}{b}$$

Antiderivative was successfully verified.

[In] Int[E^(a + b*x)*Tanh[a + b*x], x]

[Out] E^(a + b*x)/b - (2*ArcTan[E^(a + b*x)])/b

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 396

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[d*x*((a + b*x^n)^(p + 1)/(b*(n*(p + 1) + 1)), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(b*(n*(p + 1) + 1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]

Rule 2320

Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]

Rubi steps

$$\begin{aligned} \int e^{a+bx} \tanh(a+bx) dx &= \frac{\text{Subst}\left(\int \frac{-1+x^2}{1+x^2} dx, x, e^{a+bx}\right)}{b} \\ &= \frac{e^{a+bx}}{b} - \frac{2\text{Subst}\left(\int \frac{1}{1+x^2} dx, x, e^{a+bx}\right)}{b} \\ &= \frac{e^{a+bx}}{b} - \frac{2 \tan^{-1}(e^{a+bx})}{b} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 22, normalized size = 0.88

$$\frac{e^{a+bx} - 2\text{ArcTan}(e^{a+bx})}{b}$$

Antiderivative was successfully verified.

`[In] Integrate[E^(a + b*x)*Tanh[a + b*x],x]``[Out] (E^(a + b*x) - 2*ArcTan[E^(a + b*x)])/b`**Maple [A]**

time = 0.70, size = 27, normalized size = 1.08

method	result	size
derivativedivides	$\frac{\sinh(bx+a) - 2 \arctan(e^{bx+a}) + \cosh(bx+a)}{b}$	27
default	$\frac{\sinh(bx+a) - 2 \arctan(e^{bx+a}) + \cosh(bx+a)}{b}$	27
risch	$\frac{e^{bx+a}}{b} + \frac{i \ln(e^{bx+a} - i)}{b} - \frac{i \ln(e^{bx+a} + i)}{b}$	44

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(exp(b*x+a)*tanh(b*x+a),x,method=_RETURNVERBOSE)``[Out] 1/b*(sinh(b*x+a)-2*arctan(exp(b*x+a))+cosh(b*x+a))`**Maxima [A]**

time = 0.46, size = 23, normalized size = 0.92

$$-\frac{2 \arctan(e^{(bx+a)})}{b} + \frac{e^{(bx+a)}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(b*x+a)*tanh(b*x+a),x, algorithm="maxima")`

[Out] $-2 \arctan(e^{(b*x + a)})/b + e^{(b*x + a)}/b$

Fricas [A]

time = 0.32, size = 38, normalized size = 1.52

$$-\frac{2 \arctan(\cosh(bx + a) + \sinh(bx + a)) - \cosh(bx + a) - \sinh(bx + a)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(b*x+a)*tanh(b*x+a),x, algorithm="fricas")`

[Out] $-(2 \arctan(\cosh(b*x + a) + \sinh(b*x + a)) - \cosh(b*x + a) - \sinh(b*x + a))/b$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$e^a \int e^{bx} \tanh(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(b*x+a)*tanh(b*x+a),x)`

[Out] `exp(a)*Integral(exp(b*x)*tanh(a + b*x), x)`

Giac [A]

time = 0.40, size = 23, normalized size = 0.92

$$-\frac{2 \arctan(e^{(bx+a)}) - e^{(bx+a)}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(b*x+a)*tanh(b*x+a),x, algorithm="giac")`

[Out] $-(2 \arctan(e^{(b*x + a)}) - e^{(b*x + a)})/b$

Mupad [B]

time = 0.06, size = 34, normalized size = 1.36

$$\frac{e^{a+bx}}{b} - \frac{2 \operatorname{atan}\left(\frac{e^{bx} e^a \sqrt{b^2}}{b}\right)}{\sqrt{b^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(a + b*x)*tanh(a + b*x),x)`

[Out] $\exp(a + b*x)/b - (2 \operatorname{atan}((\exp(b*x) * \exp(a) * (b^2)^{(1/2)})/b)) / (b^2)^{(1/2)}$

3.210 $\int e^{a+bx} \coth(a+bx) dx$

Optimal. Leaf size=25

$$\frac{e^{a+bx}}{b} - \frac{2 \tanh^{-1}(e^{a+bx})}{b}$$

[Out] exp(b*x+a)/b-2*arctanh(exp(b*x+a))/b

Rubi [A]

time = 0.01, antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {2320, 396, 212}

$$\frac{e^{a+bx}}{b} - \frac{2 \tanh^{-1}(e^{a+bx})}{b}$$

Antiderivative was successfully verified.

[In] Int[E^(a + b*x)*Coth[a + b*x], x]

[Out] E^(a + b*x)/b - (2*ArcTanh[E^(a + b*x)])/b

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 396

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[d*x*((a + b*x^n)^(p+1)/(b*(n*(p+1)+1))), x] - Dist[(a*d - b*c*(n*(p+1)+1))/(b*(n*(p+1)+1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p+1)+1, 0]

Rule 2320

Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]

Rubi steps

$$\begin{aligned} \int e^{a+bx} \coth(a+bx) dx &= \frac{\text{Subst}\left(\int \frac{-1-x^2}{1-x^2} dx, x, e^{a+bx}\right)}{b} \\ &= \frac{e^{a+bx}}{b} - \frac{2\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, e^{a+bx}\right)}{b} \\ &= \frac{e^{a+bx}}{b} - \frac{2 \tanh^{-1}(e^{a+bx})}{b} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 22, normalized size = 0.88

$$\frac{e^{a+bx} - 2 \tanh^{-1}(e^{a+bx})}{b}$$

Antiderivative was successfully verified.

`[In] Integrate[E^(a + b*x)*Coth[a + b*x], x]``[Out] (E^(a + b*x) - 2*ArcTanh[E^(a + b*x)])/b`**Maple [A]**

time = 0.71, size = 27, normalized size = 1.08

method	result	size
derivativedivides	$\frac{\sinh(bx+a)+\cosh(bx+a)-2 \arctanh(e^{bx+a})}{b}$	27
default	$\frac{\sinh(bx+a)+\cosh(bx+a)-2 \arctanh(e^{bx+a})}{b}$	27
risch	$\frac{e^{bx+a}}{b} + \frac{\ln(e^{bx+a}-1)}{b} - \frac{\ln(e^{bx+a}+1)}{b}$	39

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(exp(b*x+a)*coth(b*x+a), x, method=_RETURNVERBOSE)``[Out] 1/b*(sinh(b*x+a)+cosh(b*x+a)-2*arctanh(exp(b*x+a)))`**Maxima [A]**

time = 0.25, size = 38, normalized size = 1.52

$$\frac{e^{(bx+a)}}{b} - \frac{\log(e^{(bx+a)} + 1)}{b} + \frac{\log(e^{(bx+a)} - 1)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(b*x+a)*coth(b*x+a), x, algorithm="maxima")`

[Out] $e^{(b*x + a)}/b - \log(e^{(b*x + a)} + 1)/b + \log(e^{(b*x + a)} - 1)/b$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 49 vs. $2(23) = 46$.

time = 0.34, size = 49, normalized size = 1.96

$$\frac{\cosh(bx + a) - \log(\cosh(bx + a) + \sinh(bx + a) + 1) + \log(\cosh(bx + a) + \sinh(bx + a) - 1) + \sinh(bx + a)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(b*x+a)*coth(b*x+a),x, algorithm="fricas")`

[Out] $(\cosh(b*x + a) - \log(\cosh(b*x + a) + \sinh(b*x + a) + 1) + \log(\cosh(b*x + a) + \sinh(b*x + a) - 1) + \sinh(b*x + a))/b$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$e^a \int e^{bx} \coth(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(b*x+a)*coth(b*x+a),x)`

[Out] $\exp(a)*\text{Integral}(\exp(b*x)*\coth(a + b*x), x)$

Giac [A]

time = 0.44, size = 32, normalized size = 1.28

$$\frac{e^{(bx+a)} - \log(e^{(bx+a)} + 1) + \log(|e^{(bx+a)} - 1|)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(b*x+a)*coth(b*x+a),x, algorithm="giac")`

[Out] $(e^{(b*x + a)} - \log(e^{(b*x + a)} + 1) + \log(\text{abs}(e^{(b*x + a)} - 1)))/b$

Mupad [B]

time = 0.08, size = 38, normalized size = 1.52

$$\frac{e^{a+bx}}{b} - \frac{2 \operatorname{atan}\left(\frac{e^{bx} e^a \sqrt{-b^2}}{b}\right)}{\sqrt{-b^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(coth(a + b*x)*exp(a + b*x),x)`

[Out] $\exp(a + b*x)/b - (2*\operatorname{atan}((\exp(b*x)*\exp(a)*(-b^2)^{(1/2)})/b))/(-b^2)^{(1/2)}$

3.211 $\int e^{a+bx} \coth^2(a+bx) dx$

Optimal. Leaf size=53

$$\frac{e^{a+bx}}{b} + \frac{2e^{a+bx}}{b(1-e^{2a+2bx})} - \frac{2 \tanh^{-1}(e^{a+bx})}{b}$$

[Out] $\exp(b*x+a)/b+2*\exp(b*x+a)/b/(1-\exp(2*b*x+2*a))-2*\operatorname{arctanh}(\exp(b*x+a))/b$

Rubi [A]

time = 0.03, antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2320, 398, 294, 212}

$$\frac{e^{a+bx}}{b} + \frac{2e^{a+bx}}{b(1-e^{2a+2bx})} - \frac{2 \tanh^{-1}(e^{a+bx})}{b}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{(a + b*x)}*\operatorname{Coth}[a + b*x]^2,x]$

[Out] $E^{(a + b*x)}/b + (2*E^{(a + b*x)})/(b*(1 - E^{(2*a + 2*b*x)})) - (2*\operatorname{ArcTanh}[E^{(a + b*x)}])/b$

Rule 212

$\operatorname{Int}[(a_+ + (b_+)*(x_+)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /;$ FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 294

$\operatorname{Int}[(c_+)*(x_+)^{(m_+)}*((a_+ + (b_+)*(x_+)^{(n_+))^{(p_+)})], x_Symbol] \rightarrow \operatorname{Simp}[c^{(n-1)}*(c*x)^{(m-n+1)}*((a + b*x^n)^{(p+1)}/(b*n*(p+1))), x] - \operatorname{Dist}[c^n * ((m-n+1)/(b*n*(p+1))), \operatorname{Int}[(c*x)^{(m-n)}*(a + b*x^n)^{(p+1)}, x], x] /;$ FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m+1, n] && ! LtQ[(m+n*(p+1)+1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 398

$\operatorname{Int}[(a_+ + (b_+)*(x_+)^{(n_+))^{(p_+)}}*((c_+ + (d_+)*(x_+)^{(n_+))^{(q_+)})], x_Symbol] \rightarrow \operatorname{Int}[\operatorname{PolynomialDivide}[(a + b*x^n)^p, (c + d*x^n)^{-q}, x], x] /;$ FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]

Rule 2320

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x]
, Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Functi
onOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[
{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*
(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rubi steps

$$\begin{aligned} \int e^{a+bx} \coth^2(a+bx) dx &= \frac{\text{Subst}\left(\int \frac{(1+x^2)^2}{(1-x^2)^2} dx, x, e^{a+bx}\right)}{b} \\ &= \frac{\text{Subst}\left(\int \left(1 + \frac{4x^2}{(1-x^2)^2}\right) dx, x, e^{a+bx}\right)}{b} \\ &= \frac{e^{a+bx}}{b} + \frac{4\text{Subst}\left(\int \frac{x^2}{(1-x^2)^2} dx, x, e^{a+bx}\right)}{b} \\ &= \frac{e^{a+bx}}{b} + \frac{2e^{a+bx}}{b(1-e^{2a+2bx})} - \frac{2\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, e^{a+bx}\right)}{b} \\ &= \frac{e^{a+bx}}{b} + \frac{2e^{a+bx}}{b(1-e^{2a+2bx})} - \frac{2 \tanh^{-1}(e^{a+bx})}{b} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 1.37, size = 179, normalized size = 3.38

$$\frac{e^{a+bx} \left(\frac{1}{48} e^{-4(a+bx)} \left(-375 - 713 e^{2(a+bx)} - 181 e^{4(a+bx)} + 61 e^{6(a+bx)} + \frac{3(125+196e^{2(a+bx)} - 14e^{4(a+bx)} - 52e^{6(a+bx)} + e^{8(a+bx)}) \tanh^{-1}\left(\sqrt{e^{2(a+bx)}}\right)}{\sqrt{e^{2(a+bx)}}} \right) + \frac{4}{105} (e^{a+bx} + e^{3(a+bx)})^2 {}_4F_3\left(\frac{3}{2}, 2, 2, 2; 1, 1, \frac{9}{2}; e^{2(a+bx)}\right) \right)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[E^(a + b*x)*Coth[a + b*x]^2,x]

[Out] (E^(a + b*x)*((-375 - 713*E^(2*(a + b*x)) - 181*E^(4*(a + b*x)) + 61*E^(6*(a + b*x)) + (3*(125 + 196*E^(2*(a + b*x)) - 14*E^(4*(a + b*x)) - 52*E^(6*(a + b*x)) + E^(8*(a + b*x)))*ArcTanh[Sqrt[E^(2*(a + b*x))]])/Sqrt[E^(2*(a + b*x))])/(48*E^(4*(a + b*x)) + (4*(E^(a + b*x) + E^(3*(a + b*x)))^2*HypergeometricPFQ[{3/2, 2, 2, 2}, {1, 1, 9/2}, E^(2*(a + b*x))])/105))/b

Maple [A]

time = 0.46, size = 48, normalized size = 0.91

method	result	size
--------	--------	------

derivativedivides	$\frac{\cosh(bx+a) - 2 \operatorname{arctanh}(e^{bx+a}) + \frac{\cosh^2(bx+a)}{\sinh(bx+a)} - \frac{2}{\sinh(bx+a)}}{b}$	48
default	$\frac{\cosh(bx+a) - 2 \operatorname{arctanh}(e^{bx+a}) + \frac{\cosh^2(bx+a)}{\sinh(bx+a)} - \frac{2}{\sinh(bx+a)}}{b}$	48
risch	$\frac{e^{bx+a}}{b} - \frac{2e^{bx+a}}{b(e^{2bx+2a}-1)} + \frac{\ln(e^{bx+a}-1)}{b} - \frac{\ln(e^{bx+a}+1)}{b}$	63

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(b*x+a)*coth(b*x+a)^2,x,method=_RETURNVERBOSE)`

[Out] `1/b*(cosh(b*x+a)-2*arctanh(exp(b*x+a))+cosh(b*x+a)^2/sinh(b*x+a)-2/sinh(b*x+a))`

Maxima [A]

time = 0.26, size = 62, normalized size = 1.17

$$\frac{e^{(bx+a)}}{b} - \frac{\log(e^{(bx+a)} + 1)}{b} + \frac{\log(e^{(bx+a)} - 1)}{b} - \frac{2e^{(bx+a)}}{b(e^{(2bx+2a)} - 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(b*x+a)*coth(b*x+a)^2,x, algorithm="maxima")`

[Out] `e^(b*x + a)/b - log(e^(b*x + a) + 1)/b + log(e^(b*x + a) - 1)/b - 2*e^(b*x + a)/(b*(e^(2*b*x + 2*a) - 1))`

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 198 vs. 2(47) = 94.

time = 0.37, size = 198, normalized size = 3.74

$$\frac{\cosh(bx+a)^3 + 3 \cosh(bx+a) \sinh(bx+a)^2 + \sinh(bx+a)^3 - (\cosh(bx+a)^2 + 2 \cosh(bx+a) \sinh(bx+a) + \sinh(bx+a)^2 - 1) \log(\cosh(bx+a) + \sinh(bx+a) + 1) + (\cosh(bx+a)^2 + 2 \cosh(bx+a) \sinh(bx+a) + \sinh(bx+a)^2 - 1) \log(\cosh(bx+a) + \sinh(bx+a) - 1) + 3(\cosh(bx+a)^2 - 1) \sinh(bx+a) - 3 \cosh(bx+a)}{b \cosh(bx+a)^2 + 2 \cosh(bx+a) \sinh(bx+a) + b \sinh(bx+a)^2 - b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(b*x+a)*coth(b*x+a)^2,x, algorithm="fricas")`

[Out] `(cosh(b*x + a)^3 + 3*cosh(b*x + a)*sinh(b*x + a)^2 + sinh(b*x + a)^3 - (cosh(b*x + a)^2 + 2*cosh(b*x + a)*sinh(b*x + a) + sinh(b*x + a)^2 - 1)*log(cosh(b*x + a) + sinh(b*x + a) + 1) + (cosh(b*x + a)^2 + 2*cosh(b*x + a)*sinh(b*x + a) + sinh(b*x + a)^2 - 1)*log(cosh(b*x + a) + sinh(b*x + a) - 1) + 3*(cosh(b*x + a)^2 - 1)*sinh(b*x + a) - 3*cosh(b*x + a))/(b*cosh(b*x + a)^2 + 2*b*cosh(b*x + a)*sinh(b*x + a) + b*sinh(b*x + a)^2 - b)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$e^a \int e^{bx} \coth^2(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(b*x+a)*coth(b*x+a)**2,x)

[Out] exp(a)*Integral(exp(b*x)*coth(a + b*x)**2, x)

Giac [A]

time = 0.45, size = 56, normalized size = 1.06

$$-\frac{\frac{2e^{(bx+a)}}{e^{(2bx+2a)}-1} - e^{(bx+a)} + \log(e^{(bx+a)} + 1) - \log(|e^{(bx+a)} - 1|)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(b*x+a)*coth(b*x+a)^2,x, algorithm="giac")

[Out] -(2*e^(b*x + a)/(e^(2*b*x + 2*a) - 1) - e^(b*x + a) + log(e^(b*x + a) + 1) - log(abs(e^(b*x + a) - 1)))/b

Mupad [B]

time = 0.07, size = 62, normalized size = 1.17

$$\frac{e^{a+bx}}{b} - \frac{2 \operatorname{atan}\left(\frac{e^{bx} e^a \sqrt{-b^2}}{b}\right)}{\sqrt{-b^2}} - \frac{2e^{a+bx}}{b(e^{2a+2bx} - 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(coth(a + b*x)^2*exp(a + b*x),x)

[Out] exp(a + b*x)/b - (2*atan((exp(b*x)*exp(a)*(-b^2)^(1/2))/b))/(-b^2)^(1/2) - (2*exp(a + b*x))/(b*(exp(2*a + 2*b*x) - 1))

3.212 $\int e^{a+bx} \coth^3(a+bx) dx$

Optimal. Leaf size=81

$$\frac{e^{a+bx}}{b} - \frac{2e^{a+bx}}{b(1-e^{2a+2bx})^2} + \frac{3e^{a+bx}}{b(1-e^{2a+2bx})} - \frac{3 \tanh^{-1}(e^{a+bx})}{b}$$

[Out] $\exp(b*x+a)/b-2*\exp(b*x+a)/b/(1-\exp(2*b*x+2*a))^2+3*\exp(b*x+a)/b/(1-\exp(2*b*x+2*a))-3*\operatorname{arctanh}(\exp(b*x+a))/b$

Rubi [A]

time = 0.04, antiderivative size = 81, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {2320, 398, 1172, 12, 294, 213}

$$\frac{e^{a+bx}}{b} + \frac{3e^{a+bx}}{b(1-e^{2a+2bx})} - \frac{2e^{a+bx}}{b(1-e^{2a+2bx})^2} - \frac{3 \tanh^{-1}(e^{a+bx})}{b}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{(a+b*x)}*\operatorname{Coth}[a+b*x]^3,x]$

[Out] $E^{(a+b*x)}/b - (2*E^{(a+b*x)})/(b*(1-E^{(2*a+2*b*x)})^2) + (3*E^{(a+b*x)})/(b*(1-E^{(2*a+2*b*x)})) - (3*\operatorname{ArcTanh}[E^{(a+b*x)}])/b$

Rule 12

$\operatorname{Int}[(a_*)(u_), x_Symbol] \rightarrow \operatorname{Dist}[a, \operatorname{Int}[u, x], x] /; \operatorname{FreeQ}[a, x] \ \&\& \ !\operatorname{Match} Q[u, (b_)*(v_)] /; \operatorname{FreeQ}[b, x]$

Rule 213

$\operatorname{Int}[(a_*) + (b_*)(x_)^2]^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(-\operatorname{Rt}[-a, 2]*\operatorname{Rt}[b, 2])^{-1})*\operatorname{ArcTanh}[\operatorname{Rt}[b, 2]*(x/\operatorname{Rt}[-a, 2])], x] /; \operatorname{FreeQ}[\{a, b\}, x] \ \&\& \ \operatorname{NegQ}[a/b] \ \&\& \ (\operatorname{LtQ}[a, 0] \ || \ \operatorname{GtQ}[b, 0])$

Rule 294

$\operatorname{Int}[(c_*)(x_)^{(m_)}*((a_*) + (b_*)(x_)^{(n_)})^{(p_)}], x_Symbol] \rightarrow \operatorname{Simp}[c^{(n-1)}*(c*x)^{(m-n+1)}*((a+b*x^n)^{(p+1)}/(b*n*(p+1))), x] - \operatorname{Dist}[c^n*((m-n+1)/(b*n*(p+1))), \operatorname{Int}[(c*x)^{(m-n)}*(a+b*x^n)^{(p+1)}, x], x] /; \operatorname{FreeQ}[\{a, b, c\}, x] \ \&\& \ \operatorname{IGtQ}[n, 0] \ \&\& \ \operatorname{LtQ}[p, -1] \ \&\& \ \operatorname{GtQ}[m+1, n] \ \&\& \ !\operatorname{LtQ}[(m+n*(p+1)+1)/n, 0] \ \&\& \ \operatorname{IntBinomialQ}[a, b, c, n, m, p, x]$

Rule 398

$\operatorname{Int}[(a_*) + (b_*)(x_)^{(n_)})^{(p_)}*((c_*) + (d_*)(x_)^{(n_)})^{(q_)}], x_Symbol] \rightarrow \operatorname{Int}[\operatorname{PolynomialDivide}[(a+b*x^n)^p, (c+d*x^n)^{-q}], x] /; \operatorname{FreeQ}[\{a$

, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]

Rule 1172

```
Int[((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] := With[{Qx = PolynomialQuotient[(a + c*x^4)^p, d + e*x^2, x], R = Coeff[PolynomialRemainder[(a + c*x^4)^p, d + e*x^2, x], x, 0]}, Simp[(-R)*x*((d + e*x^2)^(q + 1)/(2*d*(q + 1))), x] + Dist[1/(2*d*(q + 1)), Int[(d + e*x^2)^(q + 1)*ExpandToSum[2*d*(q + 1)*Qx + R*(2*q + 3), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && IGtQ[p, 0] && LtQ[q, -1]
```

Rule 2320

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rubi steps

$$\begin{aligned}
 \int e^{a+bx} \coth^3(a+bx) dx &= \frac{\text{Subst}\left(\int \frac{(1+x^2)^3}{(-1+x^2)^3} dx, x, e^{a+bx}\right)}{b} \\
 &= \frac{\text{Subst}\left(\int \left(1 + \frac{2(1+3x^4)}{(-1+x^2)^3}\right) dx, x, e^{a+bx}\right)}{b} \\
 &= \frac{e^{a+bx}}{b} + \frac{2\text{Subst}\left(\int \frac{1+3x^4}{(-1+x^2)^3} dx, x, e^{a+bx}\right)}{b} \\
 &= \frac{e^{a+bx}}{b} - \frac{2e^{a+bx}}{b(1-e^{2a+2bx})^2} + \frac{\text{Subst}\left(\int \frac{12x^2}{(-1+x^2)^2} dx, x, e^{a+bx}\right)}{2b} \\
 &= \frac{e^{a+bx}}{b} - \frac{2e^{a+bx}}{b(1-e^{2a+2bx})^2} + \frac{6\text{Subst}\left(\int \frac{x^2}{(-1+x^2)^2} dx, x, e^{a+bx}\right)}{b} \\
 &= \frac{e^{a+bx}}{b} - \frac{2e^{a+bx}}{b(1-e^{2a+2bx})^2} + \frac{3e^{a+bx}}{b(1-e^{2a+2bx})} + \frac{3\text{Subst}\left(\int \frac{1}{-1+x^2} dx, x, e^{a+bx}\right)}{b} \\
 &= \frac{e^{a+bx}}{b} - \frac{2e^{a+bx}}{b(1-e^{2a+2bx})^2} + \frac{3e^{a+bx}}{b(1-e^{2a+2bx})} - \frac{3 \tanh^{-1}(e^{a+bx})}{b}
 \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 1.80, size = 286, normalized size = 3.53

$$e^{-bx} \left(-21(252105 + 507305e^{2(a+bx)} + 173916e^{4(a+bx)} - 154296e^{6(a+bx)} - 73885e^{8(a+bx)} + 4887e^{10(a+bx)}) - \frac{31(-16807 - 28218e^{2(a+bx)} + 1173e^{4(a+bx)} + 17748e^{6(a+bx)} + 4299e^{8(a+bx)} - 1434e^{10(a+bx)} + 7e^{12(a+bx)}) \operatorname{ArcTanh}[\sqrt{e^{2(a+bx)}}]}{\sqrt{e^{2(a+bx)}}} + 384e^{8(a+bx)}(1 + e^{2(a+bx)})^7 + 5e^{2(a+bx)} {}_2F_1\left(\frac{3}{2}, 2, 2, 2, 2, 1, 1, \frac{1}{2}, e^{2(a+bx)}\right) + 256e^{8(a+bx)}(1 + e^{2(a+bx)})^3 {}_2F_1\left(\frac{3}{2}, 2, 2, 2, 2, 1, 1, 1, \frac{1}{2}, e^{2(a+bx)}\right) \right) / (bE^{5(a+bx)})$$

Antiderivative was successfully verified.

[In] Integrate[E^(a + b*x)*Coth[a + b*x]^3,x]

[Out] -1/60480*(-21*(252105 + 507305*E^(2*(a + b*x)) + 173916*E^(4*(a + b*x)) - 154296*E^(6*(a + b*x)) - 73885*E^(8*(a + b*x)) + 4887*E^(10*(a + b*x))) - (315*(-16807 - 28218*E^(2*(a + b*x)) + 1173*E^(4*(a + b*x)) + 17748*E^(6*(a + b*x)) + 4299*E^(8*(a + b*x)) - 1434*E^(10*(a + b*x)) + 7*E^(12*(a + b*x))) *ArcTanh[Sqrt[E^(2*(a + b*x))]])/Sqrt[E^(2*(a + b*x))] + 384*E^(8*(a + b*x))*(1 + E^(2*(a + b*x)))^2*(7 + 5*E^(2*(a + b*x)))*HypergeometricPFQ[{3/2, 2, 2, 2}, {1, 1, 1, 11/2}, E^(2*(a + b*x))] + 256*E^(8*(a + b*x))*(1 + E^(2*(a + b*x)))^3*HypergeometricPFQ[{3/2, 2, 2, 2, 2, 2}, {1, 1, 1, 1, 11/2}, E^(2*(a + b*x))])/(b*E^(5*(a + b*x)))

Maple [A]

time = 1.61, size = 77, normalized size = 0.95

method	result	size
risch	$\frac{e^{bx+a}}{b} - \frac{e^{bx+a}(3e^{2bx+2a}-1)}{b(e^{2bx+2a}-1)^2} + \frac{3\ln(e^{bx+a}-1)}{2b} - \frac{3\ln(e^{bx+a}+1)}{2b}$	77

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(b*x+a)*coth(b*x+a)^3,x,method=_RETURNVERBOSE)

[Out] exp(b*x+a)/b-exp(b*x+a)*(3*exp(2*b*x+2*a)-1)/b/(exp(2*b*x+2*a)-1)^2+3/2/b*ln(exp(b*x+a)-1)-3/2/b*ln(exp(b*x+a)+1)

Maxima [A]

time = 0.26, size = 88, normalized size = 1.09

$$\frac{e^{(bx+a)}}{b} - \frac{3 \log(e^{(bx+a)} + 1)}{2b} + \frac{3 \log(e^{(bx+a)} - 1)}{2b} - \frac{3e^{(3bx+3a)} - e^{(bx+a)}}{b(e^{(4bx+4a)} - 2e^{(2bx+2a)} + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(b*x+a)*coth(b*x+a)^3,x, algorithm="maxima")

[Out] e^(b*x + a)/b - 3/2*log(e^(b*x + a) + 1)/b + 3/2*log(e^(b*x + a) - 1)/b - (3*e^(3*b*x + 3*a) - e^(b*x + a))/(b*(e^(4*b*x + 4*a) - 2*e^(2*b*x + 2*a) + 1))

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 459 vs. 2(71) = 142.

time = 0.36, size = 459, normalized size = 5.67

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(b*x+a)*coth(b*x+a)^3,x, algorithm="fricas")

[Out] $\frac{1}{2}*(2*\cosh(b*x + a)^5 + 10*\cosh(b*x + a)*\sinh(b*x + a)^4 + 2*\sinh(b*x + a)^5 + 10*(2*\cosh(b*x + a)^2 - 1)*\sinh(b*x + a)^3 - 10*\cosh(b*x + a)^3 + 10*(2*\cosh(b*x + a)^3 - 3*\cosh(b*x + a))*\sinh(b*x + a)^2 - 3*(\cosh(b*x + a)^4 + 4*\cosh(b*x + a)*\sinh(b*x + a)^3 + \sinh(b*x + a)^4 + 2*(3*\cosh(b*x + a)^2 - 1)*\sinh(b*x + a)^2 - 2*\cosh(b*x + a)^2 + 4*(\cosh(b*x + a)^3 - \cosh(b*x + a)))*\sinh(b*x + a) + 1)*\log(\cosh(b*x + a) + \sinh(b*x + a) + 1) + 3*(\cosh(b*x + a)^4 + 4*\cosh(b*x + a)*\sinh(b*x + a)^3 + \sinh(b*x + a)^4 + 2*(3*\cosh(b*x + a)^2 - 1)*\sinh(b*x + a)^2 - 2*\cosh(b*x + a)^2 + 4*(\cosh(b*x + a)^3 - \cosh(b*x + a))*\sinh(b*x + a) + 1)*\log(\cosh(b*x + a) + \sinh(b*x + a) - 1) + 2*(5*\cosh(b*x + a)^4 - 15*\cosh(b*x + a)^2 + 2)*\sinh(b*x + a) + 4*\cosh(b*x + a))/(b*\cosh(b*x + a)^4 + 4*b*\cosh(b*x + a)*\sinh(b*x + a)^3 + b*\sinh(b*x + a)^4 - 2*b*\cosh(b*x + a)^2 + 2*(3*b*\cosh(b*x + a)^2 - b)*\sinh(b*x + a)^2 + 4*(b*\cosh(b*x + a)^3 - b*\cosh(b*x + a))*\sinh(b*x + a) + b)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$e^a \int e^{bx} \coth^3(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(b*x+a)*coth(b*x+a)**3,x)

[Out] exp(a)*Integral(exp(b*x)*coth(a + b*x)**3, x)

Giac [A]

time = 0.44, size = 72, normalized size = 0.89

$$\frac{2(3e^{(3bx+3a)} - e^{(bx+a)})}{(e^{(2bx+2a)} - 1)^2} - 2e^{(bx+a)} + 3 \log(e^{(bx+a)} + 1) - 3 \log(|e^{(bx+a)} - 1|)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(b*x+a)*coth(b*x+a)^3,x, algorithm="giac")

[Out] $-1/2*(2*(3*e^{(3*b*x + 3*a)} - e^{(b*x + a)})/(e^{(2*b*x + 2*a)} - 1)^2 - 2*e^{(b*x + a)} + 3*\log(e^{(b*x + a)} + 1) - 3*\log(\text{abs}(e^{(b*x + a)} - 1)))/b$

Mupad [B]

time = 1.11, size = 97, normalized size = 1.20

$$\frac{e^{a+bx}}{b} - \frac{3 \operatorname{atan}\left(\frac{e^{bx} e^a \sqrt{-b^2}}{b}\right)}{\sqrt{-b^2}} - \frac{2e^{a+bx}}{b(e^{4a+4bx} - 2e^{2a+2bx} + 1)} - \frac{3e^{a+bx}}{b(e^{2a+2bx} - 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(coth(a + b*x)^3*exp(a + b*x),x)
```

```
[Out] exp(a + b*x)/b - (3*atan((exp(b*x)*exp(a)*(-b^2)^(1/2))/b))/(-b^2)^(1/2) -  
(2*exp(a + b*x))/(b*(exp(4*a + 4*b*x) - 2*exp(2*a + 2*b*x) + 1)) - (3*exp(a  
+ b*x))/(b*(exp(2*a + 2*b*x) - 1))
```

3.213 $\int e^{a+bx} \coth^4(a+bx) dx$

Optimal. Leaf size=113

$$\frac{e^{a+bx}}{b} + \frac{8e^{a+bx}}{3b(1-e^{2a+2bx})^3} - \frac{14e^{a+bx}}{3b(1-e^{2a+2bx})^2} + \frac{5e^{a+bx}}{b(1-e^{2a+2bx})} - \frac{3 \tanh^{-1}(e^{a+bx})}{b}$$

[Out] $\exp(b*x+a)/b+8/3*\exp(b*x+a)/b/(1-\exp(2*b*x+2*a))^3-14/3*\exp(b*x+a)/b/(1-\exp(2*b*x+2*a))^2+5*\exp(b*x+a)/b/(1-\exp(2*b*x+2*a))-3*\operatorname{arctanh}(\exp(b*x+a))/b$

Rubi [A]

time = 0.05, antiderivative size = 113, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {2320, 398, 1272, 1171, 393, 212}

$$\frac{e^{a+bx}}{b} + \frac{5e^{a+bx}}{b(1-e^{2a+2bx})} - \frac{14e^{a+bx}}{3b(1-e^{2a+2bx})^2} + \frac{8e^{a+bx}}{3b(1-e^{2a+2bx})^3} - \frac{3 \tanh^{-1}(e^{a+bx})}{b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{(a + b*x)}*\text{Coth}[a + b*x]^4, x]$

[Out] $E^{(a + b*x)}/b + (8*E^{(a + b*x)})/(3*b*(1 - E^{(2*a + 2*b*x)})^3) - (14*E^{(a + b*x)})/(3*b*(1 - E^{(2*a + 2*b*x)})^2) + (5*E^{(a + b*x)})/(b*(1 - E^{(2*a + 2*b*x)})) - (3*\text{ArcTanh}[E^{(a + b*x)}])/b$

Rule 212

$\text{Int}[(a + (b \cdot x)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

Rule 393

$\text{Int}[(a + (b \cdot x)^n)^p * ((c + (d \cdot x)^n)), x_Symbol] \rightarrow \text{Simp}[(-b*c - a*d)*x*((a + b*x^n)^{p+1}/(a*b*n*(p+1))), x] - \text{Dist}[(a*d - b*c*(n*(p+1) + 1))/(a*b*n*(p+1)), \text{Int}[(a + b*x^n)^{p+1}, x], x] /; \text{FreeQ}\{a, b, c, d, n, p\}, x \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ (\text{LtQ}[p, -1] \ || \ \text{ILtQ}[1/n + p, 0])$

Rule 398

$\text{Int}[(a + (b \cdot x)^n)^p * ((c + (d \cdot x)^n)^q), x_Symbol] \rightarrow \text{Int}[\text{PolynomialDivide}[(a + b*x^n)^p, (c + d*x^n)^{-q}], x] /; \text{FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{IGtQ}[p, 0] \ \&\& \ \text{ILtQ}[q, 0] \ \&\& \ \text{GeQ}[p, -q]$

Rule 1171

```
Int[((d_) + (e_.)*(x_)^2)^(q_)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_.),
x_Symbol] := With[{Qx = PolynomialQuotient[(a + b*x^2 + c*x^4)^p, d + e*x^2,
x], R = Coeff[PolynomialRemainder[(a + b*x^2 + c*x^4)^p, d + e*x^2, x], x,
0]}, Simp[(-R)*x*((d + e*x^2)^(q + 1)/(2*d*(q + 1))), x] + Dist[1/(2*d*(q
+ 1)), Int[(d + e*x^2)^(q + 1)*ExpandToSum[2*d*(q + 1)*Qx + R*(2*q + 3), x
], x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2
- b*d*e + a*e^2, 0] && IGtQ[p, 0] && LtQ[q, -1]
```

Rule 1272

```
Int[(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(q_)*((a_) + (c_.)*(x_)^4)^(p_.), x_Sy
mbol] := Simp[(-d)^(m/2 - 1)*(c*d^2 + a*e^2)^p*x*((d + e*x^2)^(q + 1)/(2*e^
(2*p + m/2)*(q + 1))), x] + Dist[1/(2*e^(2*p + m/2)*(q + 1)), Int[(d + e*x^
2)^(q + 1)*ExpandToSum[Together[(1/(d + e*x^2))*(2*e^(2*p + m/2)*(q + 1)*x^
m*(a + c*x^4)^p - (-d)^(m/2 - 1)*(c*d^2 + a*e^2)^p*(d + e*(2*q + 3)*x^2)],
x], x], x] /; FreeQ[{a, c, d, e}, x] && IGtQ[p, 0] && ILtQ[q, -1] && IGtQ[
m/2, 0]
```

Rule 2320

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x]
, Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; Functi
onOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[
{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*
(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rubi steps

$$\begin{aligned}
\int e^{a+bx} \coth^4(a+bx) dx &= \frac{\text{Subst}\left(\int \frac{(1+x^2)^4}{(1-x^2)^4} dx, x, e^{a+bx}\right)}{b} \\
&= \frac{\text{Subst}\left(\int \left(1 + \frac{8x^2(1+x^4)}{(1-x^2)^4}\right) dx, x, e^{a+bx}\right)}{b} \\
&= \frac{e^{a+bx}}{b} + \frac{8\text{Subst}\left(\int \frac{x^2(1+x^4)}{(1-x^2)^4} dx, x, e^{a+bx}\right)}{b} \\
&= \frac{e^{a+bx}}{b} + \frac{8e^{a+bx}}{3b(1-e^{2a+2bx})^3} + \frac{4\text{Subst}\left(\int \frac{-2-6x^2-6x^4}{(1-x^2)^3} dx, x, e^{a+bx}\right)}{3b} \\
&= \frac{e^{a+bx}}{b} + \frac{8e^{a+bx}}{3b(1-e^{2a+2bx})^3} - \frac{14e^{a+bx}}{3b(1-e^{2a+2bx})^2} - \frac{\text{Subst}\left(\int \frac{-6-24x^2}{(1-x^2)^2} dx, x, e^{a+bx}\right)}{3b} \\
&= \frac{e^{a+bx}}{b} + \frac{8e^{a+bx}}{3b(1-e^{2a+2bx})^3} - \frac{14e^{a+bx}}{3b(1-e^{2a+2bx})^2} + \frac{5e^{a+bx}}{b(1-e^{2a+2bx})} - \frac{3\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, e^{a+bx}\right)}{b} \\
&= \frac{e^{a+bx}}{b} + \frac{8e^{a+bx}}{3b(1-e^{2a+2bx})^3} - \frac{14e^{a+bx}}{3b(1-e^{2a+2bx})^2} + \frac{5e^{a+bx}}{b(1-e^{2a+2bx})} - \frac{3 \tanh^{-1}(e^{a+bx})}{b}
\end{aligned}$$

Mathematica [A]

time = 10.08, size = 115, normalized size = 1.02

$$\frac{-24e^{a+bx} + 50e^{3(a+bx)} - 48e^{5(a+bx)} + 6e^{7(a+bx)} + 9(-1 + e^{2(a+bx)})^3 \log(1 - e^{a+bx}) - 9(-1 + e^{2(a+bx)})^3 \log(1 + e^{a+bx})}{6b(-1 + e^{2(a+bx)})^3}$$

Antiderivative was successfully verified.

`[In] Integrate[E^(a + b*x)*Coth[a + b*x]^4, x]`

```
[Out] (-24*E^(a + b*x) + 50*E^(3*(a + b*x)) - 48*E^(5*(a + b*x)) + 6*E^(7*(a + b*x)) + 9*(-1 + E^(2*(a + b*x)))^3*Log[1 - E^(a + b*x)] - 9*(-1 + E^(2*(a + b*x)))^3*Log[1 + E^(a + b*x)])/(6*b*(-1 + E^(2*(a + b*x)))^3)
```

Maple [A]

time = 1.28, size = 88, normalized size = 0.78

method	result	size
risch	$\frac{e^{bx+a}}{b} - \frac{e^{bx+a}(15e^{4bx+4a}-16e^{2bx+2a}+9)}{3b(e^{2bx+2a}-1)^3} + \frac{3\ln(e^{bx+a}-1)}{2b} - \frac{3\ln(e^{bx+a}+1)}{2b}$	88

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(exp(b*x+a)*coth(b*x+a)^4, x, method=_RETURNVERBOSE)`

[Out] $\exp(b*x+a)/b - 1/3*\exp(b*x+a)*(15*\exp(4*b*x+4*a) - 16*\exp(2*b*x+2*a) + 9)/b / (\exp(2*b*x+2*a) - 1)^3 + 3/2/b*\ln(\exp(b*x+a) - 1) - 3/2/b*\ln(\exp(b*x+a) + 1)$

Maxima [A]

time = 0.27, size = 110, normalized size = 0.97

$$\frac{e^{(bx+a)}}{b} - \frac{3 \log(e^{(bx+a)} + 1)}{2b} + \frac{3 \log(e^{(bx+a)} - 1)}{2b} - \frac{15e^{(5bx+5a)} - 16e^{(3bx+3a)} + 9e^{(bx+a)}}{3b(e^{(6bx+6a)} - 3e^{(4bx+4a)} + 3e^{(2bx+2a)} - 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(b*x+a)*coth(b*x+a)^4,x, algorithm="maxima")`

[Out] $e^{(b*x + a)}/b - 3/2*\log(e^{(b*x + a)} + 1)/b + 3/2*\log(e^{(b*x + a)} - 1)/b - 1/3*(15*e^{(5*b*x + 5*a)} - 16*e^{(3*b*x + 3*a)} + 9*e^{(b*x + a)})/(b*(e^{(6*b*x + 6*a)} - 3*e^{(4*b*x + 4*a)} + 3*e^{(2*b*x + 2*a)} - 1))$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 796 vs. 2(95) = 190.

time = 0.35, size = 796, normalized size = 7.04

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(b*x+a)*coth(b*x+a)^4,x, algorithm="fricas")`

[Out] $1/6*(6*\cosh(b*x + a)^7 + 42*\cosh(b*x + a)*\sinh(b*x + a)^6 + 6*\sinh(b*x + a)^7 + 6*(21*\cosh(b*x + a)^2 - 8)*\sinh(b*x + a)^5 - 48*\cosh(b*x + a)^5 + 30*(7*\cosh(b*x + a)^3 - 8*\cosh(b*x + a))*\sinh(b*x + a)^4 + 10*(21*\cosh(b*x + a)^4 - 48*\cosh(b*x + a)^2 + 5)*\sinh(b*x + a)^3 + 50*\cosh(b*x + a)^3 + 6*(21*\cosh(b*x + a)^5 - 80*\cosh(b*x + a)^3 + 25*\cosh(b*x + a))*\sinh(b*x + a)^2 - 9*(\cosh(b*x + a)^6 + 6*\cosh(b*x + a)*\sinh(b*x + a)^5 + \sinh(b*x + a)^6 + 3*(5*\cosh(b*x + a)^2 - 1)*\sinh(b*x + a)^4 - 3*\cosh(b*x + a)^4 + 4*(5*\cosh(b*x + a)^3 - 3*\cosh(b*x + a))*\sinh(b*x + a)^3 + 3*(5*\cosh(b*x + a)^4 - 6*\cosh(b*x + a)^2 + 1)*\sinh(b*x + a)^2 + 3*\cosh(b*x + a)^2 + 6*(\cosh(b*x + a)^5 - 2*\cosh(b*x + a)^3 + \cosh(b*x + a))*\sinh(b*x + a) - 1)*\log(\cosh(b*x + a) + \sinh(b*x + a) + 1) + 9*(\cosh(b*x + a)^6 + 6*\cosh(b*x + a)*\sinh(b*x + a)^5 + \sinh(b*x + a)^6 + 3*(5*\cosh(b*x + a)^2 - 1)*\sinh(b*x + a)^4 - 3*\cosh(b*x + a)^4 + 4*(5*\cosh(b*x + a)^3 - 3*\cosh(b*x + a))*\sinh(b*x + a)^3 + 3*(5*\cosh(b*x + a)^4 - 6*\cosh(b*x + a)^2 + 1)*\sinh(b*x + a)^2 + 3*\cosh(b*x + a)^2 + 6*(\cosh(b*x + a)^5 - 2*\cosh(b*x + a)^3 + \cosh(b*x + a))*\sinh(b*x + a) - 1)*\log(\cosh(b*x + a) + \sinh(b*x + a) - 1) + 6*(7*\cosh(b*x + a)^6 - 40*\cosh(b*x + a)^4 + 25*\cosh(b*x + a)^2 - 4)*\sinh(b*x + a) - 24*\cosh(b*x + a))/(b*\cosh(b*x + a)^6 + 6*b*\cosh(b*x + a)*\sinh(b*x + a)^5 + b*\sinh(b*x + a)^6 - 3*b*\cosh(b*x + a)^4 + 3*(5*b*\cosh(b*x + a)^2 - b)*\sinh(b*x + a)^4 + 4*(5*b*\cosh(b*x + a)^3 - 3*b*\cosh(b*x + a))*\sinh(b*x + a)^3 + 3*b*\cosh(b*x + a)^2 + 3*(5*$

$b*\cosh(b*x + a)^4 - 6*b*\cosh(b*x + a)^2 + b)*\sinh(b*x + a)^2 + 6*(b*\cosh(b*x + a)^5 - 2*b*\cosh(b*x + a)^3 + b*\cosh(b*x + a))*\sinh(b*x + a) - b)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$e^a \int e^{bx} \coth^4(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(b*x+a)*coth(b*x+a)**4,x)

[Out] exp(a)*Integral(exp(b*x)*coth(a + b*x)**4, x)

Giac [A]

time = 0.44, size = 83, normalized size = 0.73

$$\frac{2(15e^{(5bx+5a)} - 16e^{(3bx+3a)} + 9e^{(bx+a)})}{(e^{(2bx+2a)} - 1)^3} - 6e^{(bx+a)} + 9 \log(e^{(bx+a)} + 1) - 9 \log(|e^{(bx+a)} - 1|)}{6b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(b*x+a)*coth(b*x+a)^4,x, algorithm="giac")

[Out] -1/6*(2*(15*e^(5*b*x + 5*a) - 16*e^(3*b*x + 3*a) + 9*e^(b*x + a))/(e^(2*b*x + 2*a) - 1)^3 - 6*e^(b*x + a) + 9*log(e^(b*x + a) + 1) - 9*log(abs(e^(b*x + a) - 1)))/b

Mupad [B]

time = 0.10, size = 160, normalized size = 1.42

$$\frac{e^{a+bx}}{b} - \frac{3 \operatorname{atan}\left(\frac{e^{bx} e^a \sqrt{-b^2}}{b}\right)}{\sqrt{-b^2}} - \frac{\frac{4e^{a+bx}}{3b} + \frac{4e^{5a+5bx}}{3b}}{3e^{2a+2bx} - 3e^{4a+4bx} + e^{6a+6bx} - 1} - \frac{2e^{a+bx}}{b(e^{4a+4bx} - 2e^{2a+2bx} + 1)} - \frac{11e^{a+bx}}{3b(e^{2a+2bx} - 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(coth(a + b*x)^4*exp(a + b*x),x)

[Out] exp(a + b*x)/b - (3*atan((exp(b*x)*exp(a)*(-b^2)^(1/2))/b))/(-b^2)^(1/2) - ((4*exp(a + b*x))/(3*b) + (4*exp(5*a + 5*b*x))/(3*b))/(3*exp(2*a + 2*b*x) - 3*exp(4*a + 4*b*x) + exp(6*a + 6*b*x) - 1) - (2*exp(a + b*x))/(b*(exp(4*a + 4*b*x) - 2*exp(2*a + 2*b*x) + 1)) - (11*exp(a + b*x))/(3*b*(exp(2*a + 2*b*x) - 1))

3.214 $\int e^x \tanh^2(2x) dx$

Optimal. Leaf size=113

$$e^x + \frac{e^x}{1 + e^{4x}} + \frac{\text{ArcTan}(1 - \sqrt{2} e^x)}{2\sqrt{2}} - \frac{\text{ArcTan}(1 + \sqrt{2} e^x)}{2\sqrt{2}} + \frac{\log(1 - \sqrt{2} e^x + e^{2x})}{4\sqrt{2}} - \frac{\log(1 + \sqrt{2} e^x + e^{2x})}{4\sqrt{2}}$$

[Out] exp(x)+exp(x)/(1+exp(4*x))-1/4*arctan(-1+exp(x)*2^(1/2))*2^(1/2)-1/4*arctan(1+exp(x)*2^(1/2))*2^(1/2)+1/8*ln(1+exp(2*x)-exp(x)*2^(1/2))*2^(1/2)-1/8*ln(1+exp(2*x)+exp(x)*2^(1/2))*2^(1/2)

Rubi [A]

time = 0.06, antiderivative size = 113, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 9, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.900$, Rules used = {2320, 398, 294, 217, 1179, 642, 1176, 631, 210}

$$\frac{\text{ArcTan}(1 - \sqrt{2} e^x)}{2\sqrt{2}} - \frac{\text{ArcTan}(\sqrt{2} e^x + 1)}{2\sqrt{2}} + e^x + \frac{e^x}{e^{4x} + 1} + \frac{\log(-\sqrt{2} e^x + e^{2x} + 1)}{4\sqrt{2}} - \frac{\log(\sqrt{2} e^x + e^{2x} + 1)}{4\sqrt{2}}$$

Antiderivative was successfully verified.

[In] Int[E^x*Tanh[2*x]^2,x]

[Out] E^x + E^x/(1 + E^(4*x)) + ArcTan[1 - Sqrt[2]*E^x]/(2*Sqrt[2]) - ArcTan[1 + Sqrt[2]*E^x]/(2*Sqrt[2]) + Log[1 - Sqrt[2]*E^x + E^(2*x)]/(4*Sqrt[2]) - Log[1 + Sqrt[2]*E^x + E^(2*x)]/(4*Sqrt[2])

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)]*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 217

Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*r), Int[(r - s*x^2)/(a + b*x^4), x], x] + Dist[1/(2*r), Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))

Rule 294

Int[((c_.)*(x_)^(m_.))*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !I

LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 398

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]

Rule 631

Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 1176

Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]

Rule 1179

Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]

Rule 2320

Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))* (F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]

Rubi steps

$$\begin{aligned}
\int e^x \tanh^2(2x) dx &= \text{Subst} \left(\int \frac{(1-x^4)^2}{(1+x^4)^2} dx, x, e^x \right) \\
&= \text{Subst} \left(\int \left(1 - \frac{4x^4}{(1+x^4)^2} \right) dx, x, e^x \right) \\
&= e^x - 4 \text{Subst} \left(\int \frac{x^4}{(1+x^4)^2} dx, x, e^x \right) \\
&= e^x + \frac{e^x}{1+e^{4x}} - \text{Subst} \left(\int \frac{1}{1+x^4} dx, x, e^x \right) \\
&= e^x + \frac{e^x}{1+e^{4x}} - \frac{1}{2} \text{Subst} \left(\int \frac{1-x^2}{1+x^4} dx, x, e^x \right) - \frac{1}{2} \text{Subst} \left(\int \frac{1+x^2}{1+x^4} dx, x, e^x \right) \\
&= e^x + \frac{e^x}{1+e^{4x}} - \frac{1}{4} \text{Subst} \left(\int \frac{1}{1-\sqrt{2}x+x^2} dx, x, e^x \right) - \frac{1}{4} \text{Subst} \left(\int \frac{1}{1+\sqrt{2}x+x^2} dx, x, e^x \right) \\
&= e^x + \frac{e^x}{1+e^{4x}} + \frac{\log(1-\sqrt{2}e^x+e^{2x})}{4\sqrt{2}} - \frac{\log(1+\sqrt{2}e^x+e^{2x})}{4\sqrt{2}} - \frac{\text{Subst} \left(\int \frac{1}{-1-x^2} dx, x, e^x \right)}{2\sqrt{2}} \\
&= e^x + \frac{e^x}{1+e^{4x}} + \frac{\tan^{-1}(1-\sqrt{2}e^x)}{2\sqrt{2}} - \frac{\tan^{-1}(1+\sqrt{2}e^x)}{2\sqrt{2}} + \frac{\log(1-\sqrt{2}e^x+e^{2x})}{4\sqrt{2}}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

time = 0.04, size = 48, normalized size = 0.42

$$e^x + \frac{e^x}{1+e^{4x}} + \frac{1}{4} \text{RootSum} \left[1 + \#1^4 \&, \frac{x - \log(e^x - \#1)}{\#1^3} \& \right]$$

Antiderivative was successfully verified.

[In] Integrate[E^x*Tanh[2*x]^2,x]

[Out] E^x + E^x/(1 + E^(4*x)) + RootSum[1 + #1^4 & , (x - Log[E^x - #1])/#1^3 &] /4

Maple [C] Result contains higher order function than in optimal. Order 9 vs. order 3.

time = 0.81, size = 35, normalized size = 0.31

method	result	size
risch	$e^x + \frac{e^x}{1+e^{4x}} + \left(\sum_{R=\text{RootOf}(256Z^4+1)} -R \ln(e^x - 4R) \right)$	35

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(x)*tanh(2*x)^2,x,method=_RETURNVERBOSE)`

[Out] `exp(x)+exp(x)/(1+exp(4*x))+sum(_R*ln(exp(x)-4*_R),_R=RootOf(256*_Z^4+1))`

Maxima [A]

time = 0.47, size = 89, normalized size = 0.79

$$-\frac{1}{4}\sqrt{2}\arctan\left(\frac{1}{2}\sqrt{2}(\sqrt{2}+2e^x)\right)-\frac{1}{4}\sqrt{2}\arctan\left(-\frac{1}{2}\sqrt{2}(\sqrt{2}-2e^x)\right)-\frac{1}{8}\sqrt{2}\log(\sqrt{2}e^x+e^{2x}+1)+\frac{1}{8}\sqrt{2}\log(-\sqrt{2}e^x+e^{2x}+1)+\frac{e^x}{e^{4x}+1}+e^x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(x)*tanh(2*x)^2,x, algorithm="maxima")`

[Out] `-1/4*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*e^x)) - 1/4*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*e^x)) - 1/8*sqrt(2)*log(sqrt(2)*e^x + e^(2*x) + 1) + 1/8*sqrt(2)*log(-sqrt(2)*e^x + e^(2*x) + 1) + e^x/(e^(4*x) + 1) + e^x`

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 168 vs. 2(79) = 158.

time = 0.36, size = 168, normalized size = 1.49

$$\frac{4(\sqrt{2}e^{4x}+\sqrt{2})\arctan(-\sqrt{2}e^x+\sqrt{2}\sqrt{\sqrt{2}e^x+e^{2x}+1}-1)+4(\sqrt{2}e^{4x}+\sqrt{2})\arctan(-\sqrt{2}e^x+\frac{1}{2}\sqrt{2}\sqrt{-4\sqrt{2}e^x+4e^{2x}+4}+1)-(\sqrt{2}e^{4x}+\sqrt{2})\log(4\sqrt{2}e^x+4e^{2x}+4)+(\sqrt{2}e^{4x}+\sqrt{2})\log(-4\sqrt{2}e^x+4e^{2x}+4)+8e^{6x}+16e^x}{8(e^{4x}+1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(x)*tanh(2*x)^2,x, algorithm="fricas")`

[Out] `1/8*(4*(sqrt(2)*e^(4*x) + sqrt(2))*arctan(-sqrt(2)*e^x + sqrt(2)*sqrt(sqrt(2)*e^x + e^(2*x) + 1) - 1) + 4*(sqrt(2)*e^(4*x) + sqrt(2))*arctan(-sqrt(2)*e^x + 1/2*sqrt(2)*sqrt(-4*sqrt(2)*e^x + 4*e^(2*x) + 4) + 1) - (sqrt(2)*e^(4*x) + sqrt(2))*log(4*sqrt(2)*e^x + 4*e^(2*x) + 4) + (sqrt(2)*e^(4*x) + sqrt(2))*log(-4*sqrt(2)*e^x + 4*e^(2*x) + 4) + 8*e^(5*x) + 16*e^x/(e^(4*x) + 1)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int e^x \tanh^2(2x) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(x)*tanh(2*x)**2,x)`

[Out] `Integral(exp(x)*tanh(2*x)**2, x)`

Giac [A]

time = 0.41, size = 89, normalized size = 0.79

$$-\frac{1}{4}\sqrt{2}\arctan\left(\frac{1}{2}\sqrt{2}(\sqrt{2}+2e^x)\right) - \frac{1}{4}\sqrt{2}\arctan\left(-\frac{1}{2}\sqrt{2}(\sqrt{2}-2e^x)\right) - \frac{1}{8}\sqrt{2}\log(\sqrt{2}e^x + e^{2x} + 1) + \frac{1}{8}\sqrt{2}\log(-\sqrt{2}e^x + e^{2x} + 1) + \frac{e^x}{e^{4x} + 1} + e^x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(x)*tanh(2*x)^2,x, algorithm="giac")

[Out] -1/4*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*e^x)) - 1/4*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*e^x)) - 1/8*sqrt(2)*log(sqrt(2)*e^x + e^(2*x) + 1) + 1/8*sqrt(2)*log(-sqrt(2)*e^x + e^(2*x) + 1) + e^x/(e^(4*x) + 1) + e^x

Mupad [B]

time = 0.25, size = 86, normalized size = 0.76

$$e^x + \frac{e^x}{e^{4x} + 1} - \frac{\sqrt{2}\operatorname{atan}\left(\sqrt{2}\left(e^x - \frac{\sqrt{2}}{2}\right)\right)}{4} - \frac{\sqrt{2}\operatorname{atan}\left(\sqrt{2}\left(e^x + \frac{\sqrt{2}}{2}\right)\right)}{4} + \frac{\sqrt{2}\ln\left(\left(e^x - \frac{\sqrt{2}}{2}\right)^2 + \frac{1}{2}\right)}{8} - \frac{\sqrt{2}\ln\left(\left(e^x + \frac{\sqrt{2}}{2}\right)^2 + \frac{1}{2}\right)}{8}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(2*x)^2*exp(x),x)

[Out] exp(x) + exp(x)/(exp(4*x) + 1) - (2^(1/2)*atan(2^(1/2)*(exp(x) - 2^(1/2)/2))/4 - (2^(1/2)*atan(2^(1/2)*(exp(x) + 2^(1/2)/2))/4 + (2^(1/2)*log((exp(x) - 2^(1/2)/2)^2 + 1/2))/8 - (2^(1/2)*log((exp(x) + 2^(1/2)/2)^2 + 1/2))/8

3.215 $\int e^x \tanh(2x) dx$

Optimal. Leaf size=95

$$e^x + \frac{\operatorname{ArcTan}\left(1 - \sqrt{2} e^x\right)}{\sqrt{2}} - \frac{\operatorname{ArcTan}\left(1 + \sqrt{2} e^x\right)}{\sqrt{2}} + \frac{\log\left(1 - \sqrt{2} e^x + e^{2x}\right)}{2\sqrt{2}} - \frac{\log\left(1 + \sqrt{2} e^x + e^{2x}\right)}{2\sqrt{2}}$$

[Out] $\exp(x) - 1/2 * \arctan(-1 + \exp(x) * 2^{(1/2)}) * 2^{(1/2)} - 1/2 * \arctan(1 + \exp(x) * 2^{(1/2)}) * 2^{(1/2)} + 1/4 * \ln(1 + \exp(2*x) - \exp(x) * 2^{(1/2)}) * 2^{(1/2)} - 1/4 * \ln(1 + \exp(2*x) + \exp(x) * 2^{(1/2)}) * 2^{(1/2)}$

Rubi [A]

time = 0.05, antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 8, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {2320, 396, 217, 1179, 642, 1176, 631, 210}

$$\frac{\operatorname{ArcTan}\left(1 - \sqrt{2} e^x\right)}{\sqrt{2}} - \frac{\operatorname{ArcTan}\left(\sqrt{2} e^x + 1\right)}{\sqrt{2}} + e^x + \frac{\log\left(-\sqrt{2} e^x + e^{2x} + 1\right)}{2\sqrt{2}} - \frac{\log\left(\sqrt{2} e^x + e^{2x} + 1\right)}{2\sqrt{2}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^x * \text{Tanh}[2*x], x]$

[Out] $E^x + \operatorname{ArcTan}[1 - \operatorname{Sqrt}[2] * E^x] / \operatorname{Sqrt}[2] - \operatorname{ArcTan}[1 + \operatorname{Sqrt}[2] * E^x] / \operatorname{Sqrt}[2] + \operatorname{Log}[1 - \operatorname{Sqrt}[2] * E^x + E^{(2*x)}] / (2 * \operatorname{Sqrt}[2]) - \operatorname{Log}[1 + \operatorname{Sqrt}[2] * E^x + E^{(2*x)}] / (2 * \operatorname{Sqrt}[2])$

Rule 210

$\text{Int}[(a + (b \cdot x)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a, 2] * \text{Rt}[-b, 2])^{-1}] * \operatorname{ArcTan}[\text{Rt}[-b, 2] * (x / \text{Rt}[-a, 2])], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

Rule 217

$\text{Int}[(a + (b \cdot x)^4)^{-1}, x_Symbol] \rightarrow \text{With}\{r = \text{Numerator}[\text{Rt}[a/b, 2]], s = \text{Denominator}[\text{Rt}[a/b, 2]]\}, \text{Dist}[1/(2*r), \text{Int}[(r - s*x^2)/(a + b*x^4), x], x] + \text{Dist}[1/(2*r), \text{Int}[(r + s*x^2)/(a + b*x^4), x], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ (\text{GtQ}[a/b, 0] \ || \ (\text{PosQ}[a/b] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, a]] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, b]]))$

Rule 396

$\text{Int}[(a + (b \cdot x)^n)^p, x_Symbol] \rightarrow \text{Simp}[d*x*((a + b*x^n)^{p+1} / (b*(n*(p+1) + 1))), x] - \text{Dist}[(a*d - b*c*(n*(p+1) + 1)) / (b*(n*(p+1) + 1)), \text{Int}[(a + b*x^n)^p, x], x] /; \text{FreeQ}\{a, b,$

$c, d, n\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[n*(p + 1) + 1, 0]$

Rule 631

$\text{Int}[(a_ + (b_)*(x_) + (c_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{With}[\{q = 1 - 4*\text{Simplify}[a*(c/b^2)]\}, \text{Dist}[-2/b, \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \&\& (\text{EqQ}[q^2, 1] \parallel \text{!RationalQ}[b^2 - 4*a*c])] /; \text{FreeQ}[\{a, b, c\}, x] \&\& \text{NeQ}[b^2 - 4*a*c, 0]$

Rule 642

$\text{Int}[(d_ + (e_)*(x_))/((a_ + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&\& \text{EqQ}[2*c*d - b*e, 0]$

Rule 1176

$\text{Int}[(d_ + (e_)*(x_)^2)/((a_ + (c_)*(x_)^4), x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[2*(d/e), 2]\}, \text{Dist}[e/(2*c), \text{Int}[1/\text{Simp}[d/e + q*x + x^2, x], x], x] + \text{Dist}[e/(2*c), \text{Int}[1/\text{Simp}[d/e - q*x + x^2, x], x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{EqQ}[c*d^2 - a*e^2, 0] \&\& \text{PosQ}[d*e]$

Rule 1179

$\text{Int}[(d_ + (e_)*(x_)^2)/((a_ + (c_)*(x_)^4), x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[-2*(d/e), 2]\}, \text{Dist}[e/(2*c*q), \text{Int}[(q - 2*x)/\text{Simp}[d/e + q*x - x^2, x], x], x] + \text{Dist}[e/(2*c*q), \text{Int}[(q + 2*x)/\text{Simp}[d/e - q*x - x^2, x], x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{EqQ}[c*d^2 - a*e^2, 0] \&\& \text{NegQ}[d*e]$

Rule 2320

$\text{Int}[u_, x_Symbol] \rightarrow \text{With}[\{v = \text{FunctionOfExponential}[u, x]\}, \text{Dist}[v/D[v, x], \text{Subst}[\text{Int}[\text{FunctionOfExponentialFunction}[u, x]/x, x], x, v], x]] /; \text{FunctionOfExponentialQ}[u, x] \&\& \text{!MatchQ}[u, (w_)*((a_)*(v_)^{(n_)})^{(m_)} /; \text{FreeQ}[\{a, m, n\}, x] \&\& \text{IntegerQ}[m*n]] \&\& \text{!MatchQ}[u, E^{((c_)*(a_) + (b_)*x)}*(F_)[v_] /; \text{FreeQ}[\{a, b, c\}, x] \&\& \text{InverseFunctionQ}[F[x]]]$

Rubi steps

$$\begin{aligned}
\int e^x \tanh(2x) dx &= \text{Subst}\left(\int \frac{-1+x^4}{1+x^4} dx, x, e^x\right) \\
&= e^x - 2\text{Subst}\left(\int \frac{1}{1+x^4} dx, x, e^x\right) \\
&= e^x - \text{Subst}\left(\int \frac{1-x^2}{1+x^4} dx, x, e^x\right) - \text{Subst}\left(\int \frac{1+x^2}{1+x^4} dx, x, e^x\right) \\
&= e^x - \frac{1}{2}\text{Subst}\left(\int \frac{1}{1-\sqrt{2}x+x^2} dx, x, e^x\right) - \frac{1}{2}\text{Subst}\left(\int \frac{1}{1+\sqrt{2}x+x^2} dx, x, e^x\right) + \dots \\
&= e^x + \frac{\log(1-\sqrt{2}e^x+e^{2x})}{2\sqrt{2}} - \frac{\log(1+\sqrt{2}e^x+e^{2x})}{2\sqrt{2}} - \frac{\text{Subst}\left(\int \frac{1}{-1-x^2} dx, x, 1-\sqrt{2}e^x\right)}{\sqrt{2}} \\
&= e^x + \frac{\tan^{-1}(1-\sqrt{2}e^x)}{\sqrt{2}} - \frac{\tan^{-1}(1+\sqrt{2}e^x)}{\sqrt{2}} + \frac{\log(1-\sqrt{2}e^x+e^{2x})}{2\sqrt{2}} - \frac{\log(1+\sqrt{2}e^x+e^{2x})}{2\sqrt{2}}
\end{aligned}$$

Mathematica [A]

time = 0.02, size = 95, normalized size = 1.00

$$e^x + \frac{\text{ArcTan}(1-\sqrt{2}e^x)}{\sqrt{2}} - \frac{\text{ArcTan}(1+\sqrt{2}e^x)}{\sqrt{2}} + \frac{\log(1-\sqrt{2}e^x+e^{2x})}{2\sqrt{2}} - \frac{\log(1+\sqrt{2}e^x+e^{2x})}{2\sqrt{2}}$$

Antiderivative was successfully verified.

`[In] Integrate[E^x*Tanh[2*x],x]`

```
[Out] E^x + ArcTan[1 - Sqrt[2]*E^x]/Sqrt[2] - ArcTan[1 + Sqrt[2]*E^x]/Sqrt[2] + Log[1 - Sqrt[2]*E^x + E^(2*x)]/(2*Sqrt[2]) - Log[1 + Sqrt[2]*E^x + E^(2*x)]/(2*Sqrt[2])
```

Maple [C] Result contains higher order function than in optimal. Order 9 vs. order 3.

time = 0.80, size = 24, normalized size = 0.25

method	result	size
risch	$e^x + \left(\sum_{R=\text{RootOf}(16Z^4+1)} -R \ln(e^x - 2R) \right)$	24

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(exp(x)*tanh(2*x),x,method=_RETURNVERBOSE)``[Out] exp(x)+sum(_R*ln(exp(x)-2*_R),_R=RootOf(16*_Z^4+1))`

Maxima [A]

time = 0.49, size = 78, normalized size = 0.82

$$-\frac{1}{2}\sqrt{2} \arctan\left(\frac{1}{2}\sqrt{2}(\sqrt{2}+2e^x)\right) - \frac{1}{2}\sqrt{2} \arctan\left(-\frac{1}{2}\sqrt{2}(\sqrt{2}-2e^x)\right) - \frac{1}{4}\sqrt{2} \log(\sqrt{2}e^x + e^{(2x)} + 1) + \frac{1}{4}\sqrt{2} \log(-\sqrt{2}e^x + e^{(2x)} + 1) + e^x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(x)*tanh(2*x),x, algorithm="maxima")

[Out] $-1/2*\sqrt{2}*\arctan(1/2*\sqrt{2}*(\sqrt{2} + 2*e^x)) - 1/2*\sqrt{2}*\arctan(-1/2*\sqrt{2}*(\sqrt{2} - 2*e^x)) - 1/4*\sqrt{2}*\log(\sqrt{2}*e^x + e^{(2*x)} + 1) + 1/4*\sqrt{2}*\log(-\sqrt{2}*e^x + e^{(2*x)} + 1) + e^x$

Fricas [A]

time = 0.39, size = 113, normalized size = 1.19

$$\sqrt{2} \arctan\left(-\sqrt{2}e^x + \sqrt{2}\sqrt{\sqrt{2}e^x + e^{(2x)} + 1} - 1\right) + \sqrt{2} \arctan\left(-\sqrt{2}e^x + \frac{1}{2}\sqrt{2}\sqrt{-4\sqrt{2}e^x + 4e^{(2x)} + 4} + 1\right) - \frac{1}{4}\sqrt{2} \log(4\sqrt{2}e^x + 4e^{(2x)} + 4) + \frac{1}{4}\sqrt{2} \log(-4\sqrt{2}e^x + 4e^{(2x)} + 4) + e^x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(x)*tanh(2*x),x, algorithm="fricas")

[Out] $\sqrt{2}*\arctan(-\sqrt{2}*e^x + \sqrt{2}*\sqrt{\sqrt{2}*e^x + e^{(2*x)} + 1} - 1) + \sqrt{2}*\arctan(-\sqrt{2}*e^x + 1/2*\sqrt{2}*\sqrt{-4*\sqrt{2}*e^x + 4*e^{(2*x)} + 4} + 1) - 1/4*\sqrt{2}*\log(4*\sqrt{2}*e^x + 4*e^{(2*x)} + 4) + 1/4*\sqrt{2}*\log(-4*\sqrt{2}*e^x + 4*e^{(2*x)} + 4) + e^x$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int e^x \tanh(2x) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(x)*tanh(2*x),x)**[Out]** Integral(exp(x)*tanh(2*x), x)**Giac [A]**

time = 0.41, size = 78, normalized size = 0.82

$$-\frac{1}{2}\sqrt{2} \arctan\left(\frac{1}{2}\sqrt{2}(\sqrt{2}+2e^x)\right) - \frac{1}{2}\sqrt{2} \arctan\left(-\frac{1}{2}\sqrt{2}(\sqrt{2}-2e^x)\right) - \frac{1}{4}\sqrt{2} \log(\sqrt{2}e^x + e^{(2x)} + 1) + \frac{1}{4}\sqrt{2} \log(-\sqrt{2}e^x + e^{(2x)} + 1) + e^x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(x)*tanh(2*x),x, algorithm="giac")

[Out] $-1/2*\sqrt{2}*\arctan(1/2*\sqrt{2}*(\sqrt{2} + 2*e^x)) - 1/2*\sqrt{2}*\arctan(-1/2*\sqrt{2}*(\sqrt{2} - 2*e^x)) - 1/4*\sqrt{2}*\log(\sqrt{2}*e^x + e^{(2*x)} + 1) + 1/4*\sqrt{2}*\log(-\sqrt{2}*e^x + e^{(2*x)} + 1) + e^x$

Mupad [B]

time = 1.26, size = 81, normalized size = 0.85

$$e^x - \frac{\sqrt{2} \operatorname{atan}\left(\frac{\sqrt{2}(2e^x - \sqrt{2})}{2}\right)}{2} + \frac{\sqrt{2} \ln\left(\left(2e^x - \sqrt{2}\right)^2 + 2\right)}{4} - \frac{\sqrt{2} \operatorname{atan}\left(\frac{\sqrt{2}(2e^x + \sqrt{2})}{2}\right)}{2} - \frac{\sqrt{2} \ln\left(\left(2e^x + \sqrt{2}\right)^2 + 2\right)}{4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(2*x)*exp(x),x)

[Out] $\exp(x) - (2^{(1/2)} \operatorname{atan}((2^{(1/2)}(2 \exp(x) - 2^{(1/2)}))/2))/2 + (2^{(1/2)} \log((2 \exp(x) - 2^{(1/2)})^2 + 2))/4 - (2^{(1/2)} \operatorname{atan}((2^{(1/2)}(2 \exp(x) + 2^{(1/2)}))/2))/2 - (2^{(1/2)} \log((2 \exp(x) + 2^{(1/2)})^2 + 2))/4$

3.216 $\int e^x \coth(2x) dx$

Optimal. Leaf size=16

$$e^x - \text{ArcTan}(e^x) - \tanh^{-1}(e^x)$$

[Out] exp(x)-arctan(exp(x))-arctanh(exp(x))

Rubi [A]

time = 0.01, antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$, Rules used = {2320, 396, 218, 212, 209}

$$-\text{ArcTan}(e^x) + e^x - \tanh^{-1}(e^x)$$

Antiderivative was successfully verified.

[In] Int[E^x*Coth[2*x],x]

[Out] E^x - ArcTan[E^x] - ArcTanh[E^x]

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 218

Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 396

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[d*x*((a + b*x^n)^(p + 1)/(b*(n*(p + 1) + 1))), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(b*(n*(p + 1) + 1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]

Rule 2320

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x]
, Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Functi
onOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[
{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*
(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rubi steps

$$\begin{aligned} \int e^x \coth(2x) dx &= \text{Subst} \left(\int \frac{-1 - x^4}{1 - x^4} dx, x, e^x \right) \\ &= e^x - 2 \text{Subst} \left(\int \frac{1}{1 - x^4} dx, x, e^x \right) \\ &= e^x - \text{Subst} \left(\int \frac{1}{1 - x^2} dx, x, e^x \right) - \text{Subst} \left(\int \frac{1}{1 + x^2} dx, x, e^x \right) \\ &= e^x - \tan^{-1}(e^x) - \tanh^{-1}(e^x) \end{aligned}$$

Mathematica [A]

time = 0.01, size = 16, normalized size = 1.00

$$e^x - \text{ArcTan}(e^x) - \tanh^{-1}(e^x)$$

Antiderivative was successfully verified.

[In] Integrate[E^x*Coth[2*x], x]

[Out] E^x - ArcTan[E^x] - ArcTanh[E^x]

Maple [C] Result contains complex when optimal does not.

time = 0.85, size = 36, normalized size = 2.25

method	result	size
risch	$e^x + \frac{i \ln(e^x - i)}{2} - \frac{i \ln(e^x + i)}{2} + \frac{\ln(e^x - 1)}{2} - \frac{\ln(e^x + 1)}{2}$	36

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(x)*coth(2*x), x, method=_RETURNVERBOSE)

[Out] exp(x)+1/2*I*ln(exp(x)-I)-1/2*I*ln(exp(x)+I)+1/2*ln(exp(x)-1)-1/2*ln(exp(x)+1)

Maxima [A]

time = 0.48, size = 22, normalized size = 1.38

$$-\arctan(e^x) + e^x - \frac{1}{2} \log(e^x + 1) + \frac{1}{2} \log(e^x - 1)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(x)*coth(2*x),x, algorithm="maxima")`

[Out] $-\arctan(e^x) + e^x - 1/2 \cdot \log(e^x + 1) + 1/2 \cdot \log(e^x - 1)$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 31 vs. $2(13) = 26$.

time = 0.33, size = 31, normalized size = 1.94

$$-\arctan(\cosh(x) + \sinh(x)) + \cosh(x) - \frac{1}{2} \log(\cosh(x) + \sinh(x) + 1) + \frac{1}{2} \log(\cosh(x) + \sinh(x) - 1) + \sinh(x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(x)*coth(2*x),x, algorithm="fricas")`

[Out] $-\arctan(\cosh(x) + \sinh(x)) + \cosh(x) - 1/2 \cdot \log(\cosh(x) + \sinh(x) + 1) + 1/2 \cdot \log(\cosh(x) + \sinh(x) - 1) + \sinh(x)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int e^x \coth(2x) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(x)*coth(2*x),x)`

[Out] `Integral(exp(x)*coth(2*x), x)`

Giac [A]

time = 0.41, size = 23, normalized size = 1.44

$$-\arctan(e^x) + e^x - \frac{1}{2} \log(e^x + 1) + \frac{1}{2} \log(|e^x - 1|)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(x)*coth(2*x),x, algorithm="giac")`

[Out] $-\arctan(e^x) + e^x - 1/2 \cdot \log(e^x + 1) + 1/2 \cdot \log(\text{abs}(e^x - 1))$

Mupad [B]

time = 0.17, size = 26, normalized size = 1.62

$$\frac{\ln(2 - 2e^x)}{2} - \frac{\ln(-2e^x - 2)}{2} - \text{atan}(e^x) + e^x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(coth(2*x)*exp(x),x)`

[Out] $\log(2 - 2 \cdot \exp(x))/2 - \log(-2 \cdot \exp(x) - 2)/2 - \text{atan}(\exp(x)) + \exp(x)$

3.217 $\int e^x \coth^2(2x) dx$

Optimal. Leaf size=35

$$e^x + \frac{e^x}{1 - e^{4x}} - \frac{\text{ArcTan}(e^x)}{2} - \frac{1}{2} \tanh^{-1}(e^x)$$

[Out] exp(x)+exp(x)/(1-exp(4*x))-1/2*arctan(exp(x))-1/2*arctanh(exp(x))

Rubi [A]

time = 0.02, antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {2320, 398, 294, 218, 212, 209}

$$-\frac{1}{2} \text{ArcTan}(e^x) + e^x + \frac{e^x}{1 - e^{4x}} - \frac{1}{2} \tanh^{-1}(e^x)$$

Antiderivative was successfully verified.

[In] Int[E^x*Coth[2*x]^2,x]

[Out] E^x + E^x/(1 - E^(4*x)) - ArcTan[E^x]/2 - ArcTanh[E^x]/2

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 218

Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 294

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !I

LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 398

```
Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol]
:> Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a
, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q,
0] && GeQ[p, -q]
```

Rule 2320

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x]
, Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Functi
onOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[
{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_)+(b_)*x))*
(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rubi steps

$$\begin{aligned}
\int e^x \coth^2(2x) dx &= \text{Subst} \left(\int \frac{(1+x^4)^2}{(1-x^4)^2} dx, x, e^x \right) \\
&= \text{Subst} \left(\int \left(1 + \frac{4x^4}{(1-x^4)^2} \right) dx, x, e^x \right) \\
&= e^x + 4 \text{Subst} \left(\int \frac{x^4}{(1-x^4)^2} dx, x, e^x \right) \\
&= e^x + \frac{e^x}{1-e^{4x}} - \text{Subst} \left(\int \frac{1}{1-x^4} dx, x, e^x \right) \\
&= e^x + \frac{e^x}{1-e^{4x}} - \frac{1}{2} \text{Subst} \left(\int \frac{1}{1-x^2} dx, x, e^x \right) - \frac{1}{2} \text{Subst} \left(\int \frac{1}{1+x^2} dx, x, e^x \right) \\
&= e^x + \frac{e^x}{1-e^{4x}} - \frac{1}{2} \tan^{-1}(e^x) - \frac{1}{2} \tanh^{-1}(e^x)
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 1.16, size = 113, normalized size = 3.23

$$\frac{1}{640} e^{-7x} \left(-3645 - 6769e^{4x} - 1483e^{8x} + 681e^{12x} + 5(729 + 1208e^{4x} + 102e^{8x} - 248e^{12x} + e^{16x}) {}_2F_1\left(\frac{1}{4}, 1; \frac{5}{4}; e^{4x}\right) \right) + \frac{16}{585} e^{5x} (1+e^{4x})^2 {}_4F_3\left(\frac{5}{4}, 2, 2, 2; 1, 1, \frac{17}{4}; e^{4x}\right)$$

Antiderivative was successfully verified.

[In] Integrate[E^x*Coth[2*x]^2,x]

[Out] $(-3645 - 6769e^{4x} - 1483e^{8x} + 681e^{12x} + 5(729 + 1208e^{4x} + 102e^{8x} - 248e^{12x} + e^{16x}))\text{Hypergeometric2F1}[1/4, 1, 5/4, E^{(4x)}]/(640e^{7x}) + (16e^{5x}(1 + E^{(4x)})^2\text{HypergeometricPFQ}[\{5/4, 2, 2, 2\}, \{1, 1, 17/4\}, E^{(4x)}])/585$

Maple [C] Result contains complex when optimal does not.
time = 0.86, size = 48, normalized size = 1.37

method	result	size
risch	$e^x - \frac{e^x}{e^{4x}-1} - \frac{\ln(e^x+1)}{4} + \frac{i \ln(e^x-i)}{4} - \frac{i \ln(e^x+i)}{4} + \frac{\ln(e^x-1)}{4}$	48

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(x)*coth(2*x)^2,x,method=_RETURNVERBOSE)`

[Out] $\exp(x) - \exp(x)/(\exp(4x)-1) - 1/4 \cdot \ln(\exp(x)+1) + 1/4 \cdot I \cdot \ln(\exp(x)-I) - 1/4 \cdot I \cdot \ln(\exp(x)+I) + 1/4 \cdot \ln(\exp(x)-1)$

Maxima [A]

time = 0.47, size = 34, normalized size = 0.97

$$-\frac{e^x}{e^{(4x)}-1} - \frac{1}{2} \arctan(e^x) + e^x - \frac{1}{4} \log(e^x+1) + \frac{1}{4} \log(e^x-1)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(x)*coth(2*x)^2,x, algorithm="maxima")`

[Out] $-e^x/(e^{(4x)}-1) - 1/2 \cdot \arctan(e^x) + e^x - 1/4 \cdot \log(e^x+1) + 1/4 \cdot \log(e^x-1)$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 230 vs. $2(25) = 50$.

time = 0.36, size = 230, normalized size = 6.57

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(x)*coth(2*x)^2,x, algorithm="fricas")`

[Out] $1/4 \cdot (4 \cdot \cosh(x)^5 + 40 \cdot \cosh(x)^3 \cdot \sinh(x)^2 + 40 \cdot \cosh(x)^2 \cdot \sinh(x)^3 + 20 \cdot \cosh(x) \cdot \sinh(x)^4 + 4 \cdot \sinh(x)^5 - 2 \cdot (\cosh(x)^4 + 4 \cdot \cosh(x)^3 \cdot \sinh(x) + 6 \cdot \cosh(x)^2 \cdot \sinh(x)^2 + 4 \cdot \cosh(x) \cdot \sinh(x)^3 + \sinh(x)^4 - 1) \cdot \arctan(\cosh(x) + \sinh(x)) - (\cosh(x)^4 + 4 \cdot \cosh(x)^3 \cdot \sinh(x) + 6 \cdot \cosh(x)^2 \cdot \sinh(x)^2 + 4 \cdot \cosh(x) \cdot \sinh(x)^3 + \sinh(x)^4 - 1) \cdot \log(\cosh(x) + \sinh(x) + 1) + (\cosh(x)^4 + 4 \cdot \cosh(x)^3 \cdot \sinh(x) + 6 \cdot \cosh(x)^2 \cdot \sinh(x)^2 + 4 \cdot \cosh(x) \cdot \sinh(x)^3 + \sinh(x)^4 - 1) \cdot \log(\cosh(x) + \sinh(x) - 1) + 4 \cdot (5 \cdot \cosh(x)^4 - 2) \cdot \sinh(x) - 8 \cdot \cosh(x)) / (c$

$\cosh(x)^4 + 4*\cosh(x)^3*\sinh(x) + 6*\cosh(x)^2*\sinh(x)^2 + 4*\cosh(x)*\sinh(x)^3 + \sinh(x)^4 - 1)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int e^x \coth^2(2x) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(x)*coth(2*x)**2,x)

[Out] Integral(exp(x)*coth(2*x)**2, x)

Giac [A]

time = 0.41, size = 35, normalized size = 1.00

$$-\frac{e^x}{e^{4x}-1} - \frac{1}{2} \arctan(e^x) + e^x - \frac{1}{4} \log(e^x + 1) + \frac{1}{4} \log(|e^x - 1|)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(x)*coth(2*x)^2,x, algorithm="giac")

[Out] -e^x/(e^(4*x) - 1) - 1/2*arctan(e^x) + e^x - 1/4*log(e^x + 1) + 1/4*log(abs(e^x - 1))

Mupad [B]

time = 1.18, size = 38, normalized size = 1.09

$$\frac{\ln(1 - e^x)}{4} - \frac{\ln(-e^x - 1)}{4} - \frac{\operatorname{atan}(e^x)}{2} + e^x - \frac{e^x}{e^{4x} - 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(coth(2*x)^2*exp(x),x)

[Out] log(1 - exp(x))/4 - log(- exp(x) - 1)/4 - atan(exp(x))/2 + exp(x) - exp(x)/(exp(4*x) - 1)

3.218 $\int e^x \tanh^2(3x) dx$

Optimal. Leaf size=113

$$e^x + \frac{2e^x}{3(1+e^{6x})} - \frac{2\text{ArcTan}(e^x)}{9} + \frac{1}{9}\text{ArcTan}(\sqrt{3} - 2e^x) - \frac{1}{9}\text{ArcTan}(\sqrt{3} + 2e^x) + \frac{\log(1 - \sqrt{3}e^x + e^{2x})}{6\sqrt{3}} - \frac{\log(1 + \sqrt{3}e^x + e^{2x})}{6\sqrt{3}}$$

[Out] exp(x)+2/3*exp(x)/(1+exp(6*x))-2/9*arctan(exp(x))-1/9*arctan(2*exp(x)-3^(1/2))-1/9*arctan(2*exp(x)+3^(1/2))+1/18*ln(1+exp(2*x)-exp(x)*3^(1/2))*3^(1/2)-1/18*ln(1+exp(2*x)+exp(x)*3^(1/2))*3^(1/2)

Rubi [A]

time = 0.16, antiderivative size = 113, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 9, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.900$, Rules used = {2320, 398, 294, 215, 648, 632, 210, 642, 209}

$$-\frac{2}{9}\text{ArcTan}(e^x) + \frac{1}{9}\text{ArcTan}(\sqrt{3} - 2e^x) - \frac{1}{9}\text{ArcTan}(2e^x + \sqrt{3}) + e^x + \frac{2e^x}{3(e^{6x} + 1)} + \frac{\log(-\sqrt{3}e^x + e^{2x} + 1)}{6\sqrt{3}} - \frac{\log(\sqrt{3}e^x + e^{2x} + 1)}{6\sqrt{3}}$$

Antiderivative was successfully verified.

[In] Int[E^x*Tanh[3*x]^2,x]

[Out] E^x + (2*E^x)/(3*(1 + E^(6*x))) - (2*ArcTan[E^x])/9 + ArcTan[Sqrt[3] - 2*E^x]/9 - ArcTan[Sqrt[3] + 2*E^x]/9 + Log[1 - Sqrt[3]*E^x + E^(2*x)]/(6*Sqrt[3]) - Log[1 + Sqrt[3]*E^x + E^(2*x)]/(6*Sqrt[3])

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 215

Int[((a_) + (b_.)*(x_)^(n_))^(n_+1), x_Symbol] := Module[{r = Numerator[Rt[a/b, n]], s = Denominator[Rt[a/b, n]], u = Int[(r - s*Cos[(2*k - 1)*(Pi/n)]*x)/(r^2 - 2*r*s*Cos[(2*k - 1)*(Pi/n)]*x + s^2*x^2), x] + Int[(r + s*Cos[(2*k - 1)*(Pi/n)]*x)/(r^2 + 2*r*s*Cos[(2*k - 1)*(Pi/n)]*x + s^2*x^2), x]; 2*(r^2/(a*n))*Int[1/(r^2 + s^2*x^2), x] + Dist[2*(r/(a*n)), Sum[u, {k, 1, (n - 2)/4}], x, x] /; FreeQ[{a, b}, x] && IGtQ[(n - 2)/4, 0] &&

PosQ[a/b]

Rule 294

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(
n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n
*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x]
/; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !I
LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 398

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:= Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a
, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q,
0] && GeQ[p, -q]
```

Rule 632

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[I
nt[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c},
x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 648

```
Int[((d_.) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := D
ist[(2*c*d - b*e)/(2*c), Int[1/(a + b*x + c*x^2), x], x] + Dist[e/(2*c), In
t[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ
[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0] && !NiceSqrtQ[b^2 - 4*a*c]
```

Rule 2320

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x]
, Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; Functi
onOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[
{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*
(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rubi steps

$$\begin{aligned}
\int e^x \tanh^2(3x) dx &= \text{Subst} \left(\int \frac{(1-x^6)^2}{(1+x^6)^2} dx, x, e^x \right) \\
&= \text{Subst} \left(\int \left(1 - \frac{4x^6}{(1+x^6)^2} \right) dx, x, e^x \right) \\
&= e^x - 4 \text{Subst} \left(\int \frac{x^6}{(1+x^6)^2} dx, x, e^x \right) \\
&= e^x + \frac{2e^x}{3(1+e^{6x})} - \frac{2}{3} \text{Subst} \left(\int \frac{1}{1+x^6} dx, x, e^x \right) \\
&= e^x + \frac{2e^x}{3(1+e^{6x})} - \frac{2}{9} \text{Subst} \left(\int \frac{1}{1+x^2} dx, x, e^x \right) - \frac{2}{9} \text{Subst} \left(\int \frac{1 - \frac{\sqrt{3}x}{2}}{1 - \sqrt{3}x + x^2} dx, x, e^x \right) \\
&= e^x + \frac{2e^x}{3(1+e^{6x})} - \frac{2}{9} \tan^{-1}(e^x) - \frac{1}{18} \text{Subst} \left(\int \frac{1}{1 - \sqrt{3}x + x^2} dx, x, e^x \right) - \frac{1}{18} \text{Subst} \left(\int \frac{1}{1 + \sqrt{3}x + x^2} dx, x, e^x \right) \\
&= e^x + \frac{2e^x}{3(1+e^{6x})} - \frac{2}{9} \tan^{-1}(e^x) + \frac{\log(1 - \sqrt{3}e^x + e^{2x})}{6\sqrt{3}} - \frac{\log(1 + \sqrt{3}e^x + e^{2x})}{6\sqrt{3}} + \frac{\log(1 + \sqrt{3}e^x + e^{2x})}{6\sqrt{3}} \\
&= e^x + \frac{2e^x}{3(1+e^{6x})} - \frac{2}{9} \tan^{-1}(e^x) + \frac{1}{9} \tan^{-1}(\sqrt{3} - 2e^x) - \frac{1}{9} \tan^{-1}(\sqrt{3} + 2e^x) + \frac{\log(1 + \sqrt{3}e^x + e^{2x})}{6\sqrt{3}}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

time = 0.06, size = 97, normalized size = 0.86

$$e^x + \frac{2e^x}{3(1+e^{6x})} - \frac{2\text{ArcTan}(e^x)}{9} - \frac{1}{9} \text{RootSum} \left[1 - \#1^2 + \#1^4 \&, \frac{-2x + 2 \log(e^x - \#1) + x\#1^2 - \log(e^x - \#1)\#1^2}{-\#1 + 2\#1^3} \& \right]$$

Antiderivative was successfully verified.

[In] Integrate[E^x*Tanh[3*x]^2,x]

[Out] E^x + (2*E^x)/(3*(1 + E^(6*x))) - (2*ArcTan[E^x])/9 - RootSum[1 - #1^2 + #1^4 &, (-2*x + 2*Log[E^x - #1] + x*#1^2 - Log[E^x - #1]*#1^2)/(-#1 + 2*#1^3) &]/9

Maple [C] Result contains higher order function than in optimal. Order 9 vs. order 3.

time = 0.95, size = 59, normalized size = 0.52

method	result	size
--------	--------	------

risch	$e^x + \frac{2e^x}{3(1+e^{6x})} + \frac{i \ln(e^x - i)}{9} - \frac{i \ln(e^x + i)}{9} + \left(\sum_{R=\text{RootOf}(6561_Z^4 - 81_Z^2 + 1)} -R \ln(e^x - 9_R) \right)$	59
-------	---	----

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(x)*tanh(3*x)^2,x,method=_RETURNVERBOSE)`

[Out] `exp(x)+2/3*exp(x)/(1+exp(6*x))+1/9*I*ln(exp(x)-I)-1/9*I*ln(exp(x)+I)+sum(_R*ln(exp(x)-9*_R),_R=RootOf(6561*_Z^4-81*_Z^2+1))`

Maxima [A]

time = 0.48, size = 81, normalized size = 0.72

$$-\frac{1}{18} \sqrt{3} \log(\sqrt{3} e^x + e^{(2x)} + 1) + \frac{1}{18} \sqrt{3} \log(-\sqrt{3} e^x + e^{(2x)} + 1) + \frac{2e^x}{3(e^{6x} + 1)} - \frac{1}{9} \arctan(\sqrt{3} + 2e^x) - \frac{1}{9} \arctan(-\sqrt{3} + 2e^x) - \frac{2}{9} \arctan(e^x) + e^x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(x)*tanh(3*x)^2,x, algorithm="maxima")`

[Out] `-1/18*sqrt(3)*log(sqrt(3)*e^x + e^(2*x) + 1) + 1/18*sqrt(3)*log(-sqrt(3)*e^x + e^(2*x) + 1) + 2/3*e^x/(e^(6*x) + 1) - 1/9*arctan(sqrt(3) + 2*e^x) - 1/9*arctan(-sqrt(3) + 2*e^x) - 2/9*arctan(e^x) + e^x`

Fricas [A]

time = 0.36, size = 160, normalized size = 1.42

$$\frac{4(e^{6x} + 1) \arctan(\sqrt{3} + \sqrt{-4\sqrt{3}e^x + 4e^{(2x)} + 4} - 2e^x) + 4(e^{6x} + 1) \arctan(-\sqrt{3} + 2\sqrt{\sqrt{3}e^x + e^{(2x)} + 1} - 2e^x) - 4(e^{6x} + 1) \arctan(e^x) - (\sqrt{3}e^{6x} + \sqrt{3}) \log(4\sqrt{3}e^x + 4e^{(2x)} + 4) + (\sqrt{3}e^{6x} + \sqrt{3}) \log(-4\sqrt{3}e^x + 4e^{(2x)} + 4) + 18e^{(7x)} + 30e^x}{18(e^{6x} + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(x)*tanh(3*x)^2,x, algorithm="fricas")`

[Out] `1/18*(4*(e^(6*x) + 1)*arctan(sqrt(3) + sqrt(-4*sqrt(3)*e^x + 4*e^(2*x) + 4) - 2*e^x) + 4*(e^(6*x) + 1)*arctan(-sqrt(3) + 2*sqrt(sqrt(3)*e^x + e^(2*x) + 1) - 2*e^x) - 4*(e^(6*x) + 1)*arctan(e^x) - (sqrt(3)*e^(6*x) + sqrt(3))*log(4*sqrt(3)*e^x + 4*e^(2*x) + 4) + (sqrt(3)*e^(6*x) + sqrt(3))*log(-4*sqrt(3)*e^x + 4*e^(2*x) + 4) + 18*e^(7*x) + 30*e^x/(e^(6*x) + 1)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int e^x \tanh^2(3x) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(x)*tanh(3*x)**2,x)`

[Out] Integral(exp(x)*tanh(3*x)**2, x)

Giac [A]

time = 0.42, size = 81, normalized size = 0.72

$$-\frac{1}{18}\sqrt{3}\log(\sqrt{3}e^x + e^{2x} + 1) + \frac{1}{18}\sqrt{3}\log(-\sqrt{3}e^x + e^{2x} + 1) + \frac{2e^x}{3(e^{6x} + 1)} - \frac{1}{9}\arctan(\sqrt{3} + 2e^x) - \frac{1}{9}\arctan(-\sqrt{3} + 2e^x) - \frac{2}{9}\arctan(e^x) + e^x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(x)*tanh(3*x)^2,x, algorithm="giac")

[Out] $-\frac{1}{18}\sqrt{3}\log(\sqrt{3}e^x + e^{2x} + 1) + \frac{1}{18}\sqrt{3}\log(-\sqrt{3}e^x + e^{2x} + 1) + \frac{2e^x}{3(e^{6x} + 1)} - \frac{1}{9}\arctan(\sqrt{3} + 2e^x) - \frac{1}{9}\arctan(-\sqrt{3} + 2e^x) - \frac{2}{9}\arctan(e^x) + e^x$

Mupad [B]

time = 0.32, size = 86, normalized size = 0.76

$$e^x - \frac{\operatorname{atan}(2e^x + \sqrt{3})}{9} - \frac{\operatorname{atan}(2e^x - \sqrt{3})}{9} - \frac{2\operatorname{atan}(e^x)}{9} + \frac{2e^x}{3(e^{6x} + 1)} + \frac{\sqrt{3}\ln\left(\left(\frac{2e^x}{3} - \frac{\sqrt{3}}{3}\right)^2 + \frac{1}{9}\right)}{18} - \frac{\sqrt{3}\ln\left(\left(\frac{2e^x}{3} + \frac{\sqrt{3}}{3}\right)^2 + \frac{1}{9}\right)}{18}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(3*x)^2*exp(x),x)

[Out] $\exp(x) - \operatorname{atan}(2\exp(x) + 3^{1/2})/9 - \operatorname{atan}(2\exp(x) - 3^{1/2})/9 - (2\operatorname{atan}(\exp(x)))/9 + (2\exp(x))/(3(\exp(6x) + 1)) + (3^{1/2}\log(((2\exp(x))/3 - 3^{1/2})/3)^2 + 1/9))/18 - (3^{1/2}\log(((2\exp(x))/3 + 3^{1/2})/3)^2 + 1/9))/18$

3.219 $\int e^x \tanh(3x) dx$

Optimal. Leaf size=97

$$e^x - \frac{2\text{ArcTan}(e^x)}{3} + \frac{1}{3}\text{ArcTan}(\sqrt{3} - 2e^x) - \frac{1}{3}\text{ArcTan}(\sqrt{3} + 2e^x) + \frac{\log(1 - \sqrt{3}e^x + e^{2x})}{2\sqrt{3}} - \frac{\log(1 + \sqrt{3}e^x + e^{2x})}{2\sqrt{3}}$$

[Out] exp(x)-2/3*arctan(exp(x))-1/3*arctan(2*exp(x)-3^(1/2))-1/3*arctan(2*exp(x)+3^(1/2))+1/6*ln(1+exp(2*x)-exp(x)*3^(1/2))*3^(1/2)-1/6*ln(1+exp(2*x)+exp(x)*3^(1/2))*3^(1/2)

Rubi [A]

time = 0.14, antiderivative size = 97, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 8, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {2320, 396, 215, 648, 632, 210, 642, 209}

$$-\frac{2}{3}\text{ArcTan}(e^x) + \frac{1}{3}\text{ArcTan}(\sqrt{3} - 2e^x) - \frac{1}{3}\text{ArcTan}(2e^x + \sqrt{3}) + e^x + \frac{\log(-\sqrt{3}e^x + e^{2x} + 1)}{2\sqrt{3}} - \frac{\log(\sqrt{3}e^x + e^{2x} + 1)}{2\sqrt{3}}$$

Antiderivative was successfully verified.

[In] Int[E^x*Tanh[3*x], x]

[Out] E^x - (2*ArcTan[E^x])/3 + ArcTan[Sqrt[3] - 2*E^x]/3 - ArcTan[Sqrt[3] + 2*E^x]/3 + Log[1 - Sqrt[3]*E^x + E^(2*x)]/(2*Sqrt[3]) - Log[1 + Sqrt[3]*E^x + E^(2*x)]/(2*Sqrt[3])

Rule 209

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 210

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 215

Int[((a_) + (b_)*(x_)^(n_))(-1), x_Symbol] := Module[{r = Numerator[Rt[a/b, n]], s = Denominator[Rt[a/b, n]], k, u, v}, Simp[u = Int[(r - s*Cos[(2*k - 1)*(Pi/n)]*x)/(r^2 - 2*r*s*Cos[(2*k - 1)*(Pi/n)]*x + s^2*x^2), x] + Int[(r + s*Cos[(2*k - 1)*(Pi/n)]*x)/(r^2 + 2*r*s*Cos[(2*k - 1)*(Pi/n)]*x + s^2*x^2), x]; 2*(r^2/(a*n))*Int[1/(r^2 + s^2*x^2), x] + Dist[2*(r/(a*n)), Sum[u, {k, 1, (n - 2)/4}], x, x] /; FreeQ[{a, b}, x] && IGtQ[(n - 2)/4, 0] &&

PosQ[a/b]

Rule 396

```
Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Si
mp[d*x*((a + b*x^n)^(p + 1)/(b*(n*(p + 1) + 1))), x] - Dist[(a*d - b*c*(n*(
p + 1) + 1))/(b*(n*(p + 1) + 1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b,
c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]
```

Rule 632

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[In
t[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c},
x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 648

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := D
ist[(2*c*d - b*e)/(2*c), Int[1/(a + b*x + c*x^2), x], x] + Dist[e/(2*c), In
t[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ
[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0] && !NiceSqrtQ[b^2 - 4*a*c]
```

Rule 2320

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x]
, Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; Functi
onOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[
{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))*
(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rubi steps

$$\begin{aligned}
\int e^x \tanh(3x) dx &= \text{Subst}\left(\int \frac{-1+x^6}{1+x^6} dx, x, e^x\right) \\
&= e^x - 2\text{Subst}\left(\int \frac{1}{1+x^6} dx, x, e^x\right) \\
&= e^x - \frac{2}{3}\text{Subst}\left(\int \frac{1}{1+x^2} dx, x, e^x\right) - \frac{2}{3}\text{Subst}\left(\int \frac{1-\sqrt{3}x}{1-\sqrt{3}x+x^2} dx, x, e^x\right) - \frac{2}{3}\text{Subst}\left(\int \frac{1+\sqrt{3}x}{1+\sqrt{3}x+x^2} dx, x, e^x\right) \\
&= e^x - \frac{2}{3}\tan^{-1}(e^x) - \frac{1}{6}\text{Subst}\left(\int \frac{1}{1-\sqrt{3}x+x^2} dx, x, e^x\right) - \frac{1}{6}\text{Subst}\left(\int \frac{1}{1+\sqrt{3}x+x^2} dx, x, e^x\right) \\
&= e^x - \frac{2}{3}\tan^{-1}(e^x) + \frac{\log(1-\sqrt{3}e^x+e^{2x})}{2\sqrt{3}} - \frac{\log(1+\sqrt{3}e^x+e^{2x})}{2\sqrt{3}} + \frac{1}{3}\text{Subst}\left(\int \frac{1}{1-\sqrt{3}x+x^2} dx, x, e^x\right) \\
&= e^x - \frac{2}{3}\tan^{-1}(e^x) + \frac{1}{3}\tan^{-1}(\sqrt{3}-2e^x) - \frac{1}{3}\tan^{-1}(\sqrt{3}+2e^x) + \frac{\log(1-\sqrt{3}e^x+e^{2x})}{2\sqrt{3}}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 24, normalized size = 0.25

$$e^x - 2e^x {}_2F_1\left(\frac{1}{6}, 1; \frac{7}{6}; -e^{6x}\right)$$

Antiderivative was successfully verified.

[In] Integrate[E^x*Tanh[3*x], x]

[Out] E^x - 2E^x*Hypergeometric2F1[1/6, 1, 7/6, -E^(6*x)]

Maple [C] Result contains higher order function than in optimal. Order 9 vs. order 3.

time = 0.91, size = 47, normalized size = 0.48

method	result	size
risch	$e^x + \frac{i \ln(e^x - i)}{3} - \frac{i \ln(e^x + i)}{3} + \left(\sum_{_R=\text{RootOf}(81_Z^4-9_Z^2+1)} _R \ln(e^x - 3_R) \right)$	47

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(x)*tanh(3*x), x, method=_RETURNVERBOSE)

[Out] exp(x)+1/3*I*ln(exp(x)-I)-1/3*I*ln(exp(x)+I)+sum(_R*ln(exp(x)-3*_R), _R=RootOf(81*_Z^4-9*_Z^2+1))

Maxima [A]

time = 0.47, size = 69, normalized size = 0.71

$$-\frac{1}{6}\sqrt{3}\log(\sqrt{3}e^x + e^{2x} + 1) + \frac{1}{6}\sqrt{3}\log(-\sqrt{3}e^x + e^{2x} + 1) - \frac{1}{3}\arctan(\sqrt{3} + 2e^x) - \frac{1}{3}\arctan(-\sqrt{3} + 2e^x) - \frac{2}{3}\arctan(e^x) + e^x$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(x)*tanh(3*x),x, algorithm="maxima")`

```
[Out] -1/6*sqrt(3)*log(sqrt(3)*e^x + e^(2*x) + 1) + 1/6*sqrt(3)*log(-sqrt(3)*e^x
+ e^(2*x) + 1) - 1/3*arctan(sqrt(3) + 2*e^x) - 1/3*arctan(-sqrt(3) + 2*e^x)
- 2/3*arctan(e^x) + e^x
```

Fricas [A]

time = 0.37, size = 107, normalized size = 1.10

$$-\frac{1}{6}\sqrt{3}\log(4\sqrt{3}e^x + 4e^{2x} + 4) + \frac{1}{6}\sqrt{3}\log(-4\sqrt{3}e^x + 4e^{2x} + 4) + \frac{2}{3}\arctan(\sqrt{3} + \sqrt{-4\sqrt{3}e^x + 4e^{2x} + 4} - 2e^x) + \frac{2}{3}\arctan(-\sqrt{3} + 2\sqrt{\sqrt{3}e^x + e^{2x} + 1} - 2e^x) - \frac{2}{3}\arctan(e^x) + e^x$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(x)*tanh(3*x),x, algorithm="fricas")`

```
[Out] -1/6*sqrt(3)*log(4*sqrt(3)*e^x + 4*e^(2*x) + 4) + 1/6*sqrt(3)*log(-4*sqrt(3)
)*e^x + 4*e^(2*x) + 4) + 2/3*arctan(sqrt(3) + sqrt(-4*sqrt(3)*e^x + 4*e^(2*
x) + 4) - 2*e^x) + 2/3*arctan(-sqrt(3) + 2*sqrt(sqrt(3)*e^x + e^(2*x) + 1)
- 2*e^x) - 2/3*arctan(e^x) + e^x
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int e^x \tanh(3x) dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(x)*tanh(3*x),x)``[Out] Integral(exp(x)*tanh(3*x), x)`**Giac [A]**

time = 0.42, size = 69, normalized size = 0.71

$$-\frac{1}{6}\sqrt{3}\log(\sqrt{3}e^x + e^{2x} + 1) + \frac{1}{6}\sqrt{3}\log(-\sqrt{3}e^x + e^{2x} + 1) - \frac{1}{3}\arctan(\sqrt{3} + 2e^x) - \frac{1}{3}\arctan(-\sqrt{3} + 2e^x) - \frac{2}{3}\arctan(e^x) + e^x$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(x)*tanh(3*x),x, algorithm="giac")`

```
[Out] -1/6*sqrt(3)*log(sqrt(3)*e^x + e^(2*x) + 1) + 1/6*sqrt(3)*log(-sqrt(3)*e^x
+ e^(2*x) + 1) - 1/3*arctan(sqrt(3) + 2*e^x) - 1/3*arctan(-sqrt(3) + 2*e^x)
- 2/3*arctan(e^x) + e^x
```

Mupad [B]

time = 0.26, size = 70, normalized size = 0.72

$$e^x - \frac{\operatorname{atan}\left(2e^x + \sqrt{3}\right)}{3} - \frac{\operatorname{atan}\left(2e^x - \sqrt{3}\right)}{3} - \frac{2\operatorname{atan}\left(e^x\right)}{3} + \frac{\sqrt{3} \ln\left(\left(2e^x - \sqrt{3}\right)^2 + 1\right)}{6} - \frac{\sqrt{3} \ln\left(\left(2e^x + \sqrt{3}\right)^2 + 1\right)}{6}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(tanh(3*x)*exp(x),x)`

```
[Out] exp(x) - atan(2*exp(x) + 3^(1/2))/3 - atan(2*exp(x) - 3^(1/2))/3 - (2*atan(
exp(x)))/3 + (3^(1/2)*log((2*exp(x) - 3^(1/2))^2 + 1))/6 - (3^(1/2)*log((2*
exp(x) + 3^(1/2))^2 + 1))/6
```

3.220 $\int e^x \coth(3x) dx$

Optimal. Leaf size=85

$$e^x + \frac{\operatorname{ArcTan}\left(\frac{1-2e^x}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{\operatorname{ArcTan}\left(\frac{1+2e^x}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{2}{3} \tanh^{-1}(e^x) + \frac{1}{6} \log(1 - e^x + e^{2x}) - \frac{1}{6} \log(1 + e^x + e^{2x})$$

[Out] exp(x)-2/3*arctanh(exp(x))+1/6*ln(1-exp(x)+exp(2*x))-1/6*ln(1+exp(x)+exp(2*x))+1/3*arctan(1/3*(1-2*exp(x))*3^(1/2))*3^(1/2)-1/3*arctan(1/3*(1+2*exp(x))*3^(1/2))*3^(1/2)

Rubi [A]

time = 0.09, antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 8, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {2320, 396, 216, 648, 632, 210, 642, 212}

$$\frac{\operatorname{ArcTan}\left(\frac{1-2e^x}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{\operatorname{ArcTan}\left(\frac{2e^x+1}{\sqrt{3}}\right)}{\sqrt{3}} + e^x + \frac{1}{6} \log(-e^x + e^{2x} + 1) - \frac{1}{6} \log(e^x + e^{2x} + 1) - \frac{2}{3} \tanh^{-1}(e^x)$$

Antiderivative was successfully verified.

[In] Int[E^x*Coth[3*x],x]

[Out] E^x + ArcTan[(1 - 2*E^x)/Sqrt[3]]/Sqrt[3] - ArcTan[(1 + 2*E^x)/Sqrt[3]]/Sqrt[3] - (2*ArcTanh[E^x])/3 + Log[1 - E^x + E^(2*x)]/6 - Log[1 + E^x + E^(2*x)]/6

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 216

Int[((a_) + (b_.)*(x_)^(n_))^(n_+1), x_Symbol] := Module[{r = Numerator[Rt[-a/b, n]], s = Denominator[Rt[-a/b, n]], k, u}, Simp[u = Int[(r - s*Cos[(2*k*Pi)/n]*x)/(r^2 - 2*r*s*Cos[(2*k*Pi)/n]*x + s^2*x^2), x] + Int[(r + s*Cos[(2*k*Pi)/n]*x)/(r^2 + 2*r*s*Cos[(2*k*Pi)/n]*x + s^2*x^2), x]; 2*(r^2/(a*n))*Int[1/(r^2 - s^2*x^2), x] + Dist[2*(r/(a*n)), Sum[u, {k, 1, (n - 2)/4}], x],

x]] /; FreeQ[{a, b}, x] && IGtQ[(n - 2)/4, 0] && NegQ[a/b]

Rule 396

Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Simp[d*x*((a + b*x^n)^(p + 1)/(b*(n*(p + 1) + 1))), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(b*(n*(p + 1) + 1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]

Rule 632

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 648

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Dist[(2*c*d - b*e)/(2*c), Int[1/(a + b*x + c*x^2), x], x] + Dist[e/(2*c), Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0] && !NiceSqrtQ[b^2 - 4*a*c]

Rule 2320

Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]

Rubi steps

$$\begin{aligned}
\int e^x \coth(3x) dx &= \text{Subst} \left(\int \frac{-1-x^6}{1-x^6} dx, x, e^x \right) \\
&= e^x - 2 \text{Subst} \left(\int \frac{1}{1-x^6} dx, x, e^x \right) \\
&= e^x - \frac{2}{3} \text{Subst} \left(\int \frac{1}{1-x^2} dx, x, e^x \right) - \frac{2}{3} \text{Subst} \left(\int \frac{1-\frac{x}{2}}{1-x+x^2} dx, x, e^x \right) - \frac{2}{3} \text{Subst} \left(\int \frac{1}{1-x^2} dx, x, e^x \right) \\
&= e^x - \frac{2}{3} \tanh^{-1}(e^x) + \frac{1}{6} \text{Subst} \left(\int \frac{-1+2x}{1-x+x^2} dx, x, e^x \right) - \frac{1}{6} \text{Subst} \left(\int \frac{1+2x}{1+x+x^2} dx, x, e^x \right) \\
&= e^x - \frac{2}{3} \tanh^{-1}(e^x) + \frac{1}{6} \log(1-e^x+e^{2x}) - \frac{1}{6} \log(1+e^x+e^{2x}) + \text{Subst} \left(\int \frac{1}{-3-x^2} dx, x, e^x \right) \\
&= e^x - \frac{\tan^{-1} \left(\frac{-1+2e^x}{\sqrt{3}} \right)}{\sqrt{3}} - \frac{\tan^{-1} \left(\frac{1+2e^x}{\sqrt{3}} \right)}{\sqrt{3}} - \frac{2}{3} \tanh^{-1}(e^x) + \frac{1}{6} \log(1-e^x+e^{2x}) - \frac{1}{6} \log(1+e^x+e^{2x})
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 22, normalized size = 0.26

$$e^x - 2e^x {}_2F_1 \left(\frac{1}{6}, 1; \frac{7}{6}; e^{6x} \right)$$

Antiderivative was successfully verified.

[In] Integrate[E^x*Coth[3*x],x]

[Out] E^x - 2*E^x*Hypergeometric2F1[1/6, 1, 7/6, E^(6*x)]

Maple [C] Result contains complex when optimal does not.

time = 0.79, size = 138, normalized size = 1.62

method	result
risch	$e^x - \frac{\ln(e^x+1)}{3} + \frac{\ln(e^x-1)}{3} + \frac{\ln \left(e^{x-\frac{1}{2}-\frac{i\sqrt{3}}{2}} \right)}{6} + \frac{i \ln \left(e^{x-\frac{1}{2}-\frac{i\sqrt{3}}{2}} \right) \sqrt{3}}{6} + \frac{\ln \left(e^{x-\frac{1}{2}+\frac{i\sqrt{3}}{2}} \right)}{6} - \frac{i \ln \left(e^{x-\frac{1}{2}+\frac{i\sqrt{3}}{2}} \right) \sqrt{3}}{6}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(x)*coth(3*x),x,method=_RETURNVERBOSE)

[Out] exp(x)-1/3*ln(exp(x)+1)+1/3*ln(exp(x)-1)+1/6*ln(exp(x)-1/2-1/2*I*3^(1/2))+1/6*I*ln(exp(x)-1/2-1/2*I*3^(1/2))*3^(1/2)+1/6*ln(exp(x)-1/2+1/2*I*3^(1/2))-1/6*I*ln(exp(x)-1/2+1/2*I*3^(1/2))*3^(1/2)-1/6*ln(exp(x)+1/2-1/2*I*3^(1/2))

+1/6*I*ln(exp(x)+1/2-1/2*I*3^(1/2))*3^(1/2)-1/6*ln(exp(x)+1/2+1/2*I*3^(1/2))
)-1/6*I*ln(exp(x)+1/2+1/2*I*3^(1/2))*3^(1/2)

Maxima [A]

time = 0.48, size = 75, normalized size = 0.88

$$-\frac{1}{3}\sqrt{3}\arctan\left(\frac{1}{3}\sqrt{3}(2e^x+1)\right)-\frac{1}{3}\sqrt{3}\arctan\left(\frac{1}{3}\sqrt{3}(2e^x-1)\right)+e^x-\frac{1}{6}\log(e^{(2x)}+e^x+1)+\frac{1}{6}\log(e^{(2x)}-e^x+1)-\frac{1}{3}\log(e^x+1)+\frac{1}{3}\log(e^x-1)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(x)*coth(3*x),x, algorithm="maxima")

[Out] -1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*e^x + 1)) - 1/3*sqrt(3)*arctan(1/3*sqrt(3)
)*(2*e^x - 1)) + e^x - 1/6*log(e^(2*x) + e^x + 1) + 1/6*log(e^(2*x) - e^x
 + 1) - 1/3*log(e^x + 1) + 1/3*log(e^x - 1)

Fricas [A]

time = 0.33, size = 113, normalized size = 1.33

$$-\frac{1}{3}\sqrt{3}\arctan\left(\frac{2}{3}\sqrt{3}\cosh(x)+\frac{2}{3}\sqrt{3}\sinh(x)+\frac{1}{3}\sqrt{3}\right)-\frac{1}{3}\sqrt{3}\arctan\left(\frac{2}{3}\sqrt{3}\cosh(x)+\frac{2}{3}\sqrt{3}\sinh(x)-\frac{1}{3}\sqrt{3}\right)+\cosh(x)-\frac{1}{6}\log\left(\frac{2\cosh(x)+1}{\cosh(x)-\sinh(x)}\right)+\frac{1}{6}\log\left(\frac{2\cosh(x)-1}{\cosh(x)-\sinh(x)}\right)-\frac{1}{3}\log(\cosh(x)+\sinh(x)+1)+\frac{1}{3}\log(\cosh(x)+\sinh(x)-1)+\sinh(x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(x)*coth(3*x),x, algorithm="fricas")

[Out] -1/3*sqrt(3)*arctan(2/3*sqrt(3)*cosh(x) + 2/3*sqrt(3)*sinh(x) + 1/3*sqrt(3)
) - 1/3*sqrt(3)*arctan(2/3*sqrt(3)*cosh(x) + 2/3*sqrt(3)*sinh(x) - 1/3*sqrt
 (3)) + cosh(x) - 1/6*log((2*cosh(x) + 1)/(cosh(x) - sinh(x))) + 1/6*log((2*
 cosh(x) - 1)/(cosh(x) - sinh(x))) - 1/3*log(cosh(x) + sinh(x) + 1) + 1/3*lo
 g(cosh(x) + sinh(x) - 1) + sinh(x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int e^x \coth(3x) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(x)*coth(3*x),x)

[Out] Integral(exp(x)*coth(3*x), x)

Giac [A]

time = 0.41, size = 76, normalized size = 0.89

$$-\frac{1}{3}\sqrt{3}\arctan\left(\frac{1}{3}\sqrt{3}(2e^x+1)\right)-\frac{1}{3}\sqrt{3}\arctan\left(\frac{1}{3}\sqrt{3}(2e^x-1)\right)+e^x-\frac{1}{6}\log(e^{(2x)}+e^x+1)+\frac{1}{6}\log(e^{(2x)}-e^x+1)-\frac{1}{3}\log(e^x+1)+\frac{1}{3}\log(|e^x-1|)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(x)*coth(3*x),x, algorithm="giac")

[Out] $-\frac{1}{3}\sqrt{3}\arctan\left(\frac{1}{3}\sqrt{3}(2e^x + 1)\right) - \frac{1}{3}\sqrt{3}\arctan\left(\frac{1}{3}\sqrt{3}(2e^x - 1)\right) + e^x - \frac{1}{6}\log(e^{2x} + e^x + 1) + \frac{1}{6}\log(e^{2x} - e^x + 1) - \frac{1}{3}\log(e^x + 1) + \frac{1}{3}\log(\text{abs}(e^x - 1))$

Mupad [B]

time = 0.26, size = 81, normalized size = 0.95

$$\frac{\ln(2 - 2e^x)}{3} - \frac{\ln(-2e^x - 2)}{3} + \frac{\ln((2e^x - 1)^2 + 3)}{6} - \frac{\ln((2e^x + 1)^2 + 3)}{6} + e^x - \frac{\sqrt{3} \operatorname{atan}\left(\frac{\sqrt{3}(2e^x - 1)}{3}\right)}{3} - \frac{\sqrt{3} \operatorname{atan}\left(\frac{\sqrt{3}(2e^x + 1)}{3}\right)}{3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(coth(3*x)*exp(x),x)

[Out] $\log(2 - 2\exp(x))/3 - \log(-2\exp(x) - 2)/3 + \log((2\exp(x) - 1)^2 + 3)/6 - \log((2\exp(x) + 1)^2 + 3)/6 + \exp(x) - (3^{1/2}*\operatorname{atan}((3^{1/2}*(2\exp(x) - 1))/3))/3 - (3^{1/2}*\operatorname{atan}((3^{1/2}*(2\exp(x) + 1))/3))/3$

3.221 $\int e^x \coth^2(3x) dx$

Optimal. Leaf size=108

$$e^x + \frac{2e^x}{3(1-e^{6x})} + \frac{\operatorname{ArcTan}\left(\frac{1-2e^x}{\sqrt{3}}\right)}{3\sqrt{3}} - \frac{\operatorname{ArcTan}\left(\frac{1+2e^x}{\sqrt{3}}\right)}{3\sqrt{3}} - \frac{2}{9} \tanh^{-1}(e^x) + \frac{1}{18} \log(1-e^x+e^{2x}) - \frac{1}{18} \log(1+e^x)$$

[Out] exp(x)+2/3*exp(x)/(1-exp(6*x))-2/9*arctanh(exp(x))+1/18*ln(1-exp(x)+exp(2*x))-1/18*ln(1+exp(x)+exp(2*x))+1/9*arctan(1/3*(1-2*exp(x))*3^(1/2))*3^(1/2)-1/9*arctan(1/3*(1+2*exp(x))*3^(1/2))*3^(1/2)

Rubi [A]

time = 0.10, antiderivative size = 108, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 9, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.900$, Rules used = {2320, 398, 294, 216, 648, 632, 210, 642, 212}

$$\frac{\operatorname{ArcTan}\left(\frac{1-2e^x}{\sqrt{3}}\right)}{3\sqrt{3}} - \frac{\operatorname{ArcTan}\left(\frac{2e^x+1}{\sqrt{3}}\right)}{3\sqrt{3}} + e^x + \frac{2e^x}{3(1-e^{6x})} + \frac{1}{18} \log(-e^x+e^{2x}+1) - \frac{1}{18} \log(e^x+e^{2x}+1) - \frac{2}{9} \tanh^{-1}(e^x)$$

Antiderivative was successfully verified.

[In] Int[E^x*Coth[3*x]^2,x]

[Out] E^x + (2*E^x)/(3*(1 - E^(6*x))) + ArcTan[(1 - 2*E^x)/Sqrt[3]]/(3*Sqrt[3]) - ArcTan[(1 + 2*E^x)/Sqrt[3]]/(3*Sqrt[3]) - (2*ArcTanh[E^x])/9 + Log[1 - E^x + E^(2*x)]/18 - Log[1 + E^x + E^(2*x)]/18

Rule 210

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 216

Int[((a_) + (b_)*(x_)^(n_))(-1), x_Symbol] := Module[{r = Numerator[Rt[-a/b, n]], s = Denominator[Rt[-a/b, n]], k, u}, Simp[u = Int[(r - s*Cos[(2*k*Pi)/n]*x)/(r^2 - 2*r*s*Cos[(2*k*Pi)/n]*x + s^2*x^2), x] + Int[(r + s*Cos[(2*k*Pi)/n]*x)/(r^2 + 2*r*s*Cos[(2*k*Pi)/n]*x + s^2*x^2), x]; 2*(r^2/(a*n))*Int[1/(r^2 - s^2*x^2), x] + Dist[2*(r/(a*n)), Sum[u, {k, 1, (n - 2)/4}], x],

x]] /; FreeQ[{a, b}, x] && IGtQ[(n - 2)/4, 0] && NegQ[a/b]

Rule 294

Int[((c_.)*(x_)^(m_.))*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !ILtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 398

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]

Rule 632

Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 648

Int[((d_.) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Dist[(2*c*d - b*e)/(2*c), Int[1/(a + b*x + c*x^2), x], x] + Dist[e/(2*c), Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0] && !NiceSqrtQ[b^2 - 4*a*c]

Rule 2320

Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]

Rubi steps

$$\begin{aligned}
\int e^x \coth^2(3x) dx &= \text{Subst} \left(\int \frac{(1+x^6)^2}{(1-x^6)^2} dx, x, e^x \right) \\
&= \text{Subst} \left(\int \left(1 + \frac{4x^6}{(1-x^6)^2} \right) dx, x, e^x \right) \\
&= e^x + 4 \text{Subst} \left(\int \frac{x^6}{(1-x^6)^2} dx, x, e^x \right) \\
&= e^x + \frac{2e^x}{3(1-e^{6x})} - \frac{2}{3} \text{Subst} \left(\int \frac{1}{1-x^6} dx, x, e^x \right) \\
&= e^x + \frac{2e^x}{3(1-e^{6x})} - \frac{2}{9} \text{Subst} \left(\int \frac{1}{1-x^2} dx, x, e^x \right) - \frac{2}{9} \text{Subst} \left(\int \frac{1-x^{\frac{x}{2}}}{1-x+x^2} dx, x, e^x \right) \\
&= e^x + \frac{2e^x}{3(1-e^{6x})} - \frac{2}{9} \tanh^{-1}(e^x) + \frac{1}{18} \text{Subst} \left(\int \frac{-1+2x}{1-x+x^2} dx, x, e^x \right) - \frac{1}{18} \text{Subst} \left(\int \frac{1}{1-x+x^2} dx, x, e^x \right) \\
&= e^x + \frac{2e^x}{3(1-e^{6x})} - \frac{2}{9} \tanh^{-1}(e^x) + \frac{1}{18} \log(1-e^x+e^{2x}) - \frac{1}{18} \log(1+e^x+e^{2x}) + \frac{1}{3} \text{Subst} \left(\int \frac{1}{1-x+x^2} dx, x, e^x \right) \\
&= e^x + \frac{2e^x}{3(1-e^{6x})} + \frac{\tan^{-1}\left(\frac{1-2e^x}{\sqrt{3}}\right)}{3\sqrt{3}} - \frac{\tan^{-1}\left(\frac{1+2e^x}{\sqrt{3}}\right)}{3\sqrt{3}} - \frac{2}{9} \tanh^{-1}(e^x) + \frac{1}{18} \log(1-e^x+e^{2x}) - \frac{1}{18} \log(1+e^x+e^{2x}) + \frac{1}{3} \text{Subst} \left(\int \frac{1}{1-x+x^2} dx, x, e^x \right)
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 1.37, size = 113, normalized size = 1.05

$$\frac{e^{-11x}(-15379 - 28153e^{6x} - 5633e^{12x} + 3109e^{18x} + 7(2197 + 3708e^{6x} + 538e^{12x} - 684e^{18x} + e^{24x}) {}_2F_1\left(\frac{1}{6}, 1; \frac{7}{6}; e^{6x}\right)) + 36e^{7x}(1 + e^{6x})^2 {}_4F_3\left(\frac{7}{6}, 2, 2, 2; 1, 1, \frac{25}{6}; e^{6x}\right)}{3024 + 1729}$$

Antiderivative was successfully verified.

[In] Integrate[E^x*Coth[3*x]^2,x]

[Out] (-15379 - 28153*E^(6*x) - 5633*E^(12*x) + 3109*E^(18*x) + 7*(2197 + 3708*E^(6*x) + 538*E^(12*x) - 684*E^(18*x) + E^(24*x))*Hypergeometric2F1[1/6, 1, 7/6, E^(6*x)])/(3024*E^(11*x)) + (36*E^(7*x)*(1 + E^(6*x))^2*HypergeometricPFQ[{7/6, 2, 2, 2}, {1, 1, 25/6}, E^(6*x)])/1729

Maple [C] Result contains complex when optimal does not.

time = 0.82, size = 150, normalized size = 1.39

method	result
risch	$ e^x - \frac{2e^x}{3(e^{6x}-1)} - \frac{\ln(e^x+1)}{9} - \frac{\ln\left(e^x+\frac{1}{2}-\frac{i\sqrt{3}}{2}\right)}{18} + \frac{i\ln\left(e^x+\frac{1}{2}-\frac{i\sqrt{3}}{2}\right)\sqrt{3}}{18} - \frac{\ln\left(e^x+\frac{1}{2}+\frac{i\sqrt{3}}{2}\right)}{18} - \frac{i\ln\left(e^x+\frac{1}{2}+\frac{i\sqrt{3}}{2}\right)\sqrt{3}}{18} $

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(exp(x)*coth(3*x)^2,x,method=_RETURNVERBOSE)
```

```
[Out] exp(x)-2/3*exp(x)/(exp(6*x)-1)-1/9*ln(exp(x)+1)-1/18*ln(exp(x)+1/2-1/2*I*3^(1/2))+1/18*I*ln(exp(x)+1/2-1/2*I*3^(1/2))*3^(1/2)-1/18*ln(exp(x)+1/2+1/2*I*3^(1/2))-1/18*I*ln(exp(x)+1/2+1/2*I*3^(1/2))*3^(1/2)+1/18*ln(exp(x)-1/2-1/2*I*3^(1/2))+1/18*I*ln(exp(x)-1/2-1/2*I*3^(1/2))*3^(1/2)+1/18*ln(exp(x)-1/2+1/2*I*3^(1/2))-1/18*I*ln(exp(x)-1/2+1/2*I*3^(1/2))*3^(1/2)+1/9*ln(exp(x)-1)
```

Maxima [A]

time = 0.48, size = 87, normalized size = 0.81

$$-\frac{1}{9}\sqrt{3}\arctan\left(\frac{1}{3}\sqrt{3}(2e^x+1)\right)-\frac{1}{9}\sqrt{3}\arctan\left(\frac{1}{3}\sqrt{3}(2e^x-1)\right)-\frac{2e^x}{3(e^{6x}-1)}+e^x-\frac{1}{18}\log(e^{2x}+e^x+1)+\frac{1}{18}\log(e^{2x}-e^x+1)-\frac{1}{9}\log(e^x+1)+\frac{1}{9}\log(e^x-1)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(x)*coth(3*x)^2,x, algorithm="maxima")
```

```
[Out] -1/9*sqrt(3)*arctan(1/3*sqrt(3)*(2*e^x + 1)) - 1/9*sqrt(3)*arctan(1/3*sqrt(3)*(2*e^x - 1)) - 2/3*e^x/(e^(6*x) - 1) + e^x - 1/18*log(e^(2*x) + e^x + 1) + 1/18*log(e^(2*x) - e^x + 1) - 1/9*log(e^x + 1) + 1/9*log(e^x - 1)
```

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 628 vs. 2(78) = 156.

time = 0.35, size = 628, normalized size = 5.81

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(x)*coth(3*x)^2,x, algorithm="fricas")
```

```
[Out] 1/18*(18*cosh(x)^7 + 378*cosh(x)^5*sinh(x)^2 + 630*cosh(x)^4*sinh(x)^3 + 630*cosh(x)^3*sinh(x)^4 + 378*cosh(x)^2*sinh(x)^5 + 126*cosh(x)*sinh(x)^6 + 18*sinh(x)^7 - 2*(sqrt(3)*cosh(x)^6 + 6*sqrt(3)*cosh(x)^5*sinh(x) + 15*sqrt(3)*cosh(x)^4*sinh(x)^2 + 20*sqrt(3)*cosh(x)^3*sinh(x)^3 + 15*sqrt(3)*cosh(x)^2*sinh(x)^4 + 6*sqrt(3)*cosh(x)*sinh(x)^5 + sqrt(3)*sinh(x)^6 - sqrt(3))*arctan(2/3*sqrt(3)*cosh(x) + 2/3*sqrt(3)*sinh(x) + 1/3*sqrt(3)) - 2*(sqrt(3)*cosh(x)^6 + 6*sqrt(3)*cosh(x)^5*sinh(x) + 15*sqrt(3)*cosh(x)^4*sinh(x)^2 + 20*sqrt(3)*cosh(x)^3*sinh(x)^3 + 15*sqrt(3)*cosh(x)^2*sinh(x)^4 + 6*sqrt(3)*cosh(x)*sinh(x)^5 + sqrt(3)*sinh(x)^6 - sqrt(3))*arctan(2/3*sqrt(3)*cosh(x) + 2/3*sqrt(3)*sinh(x) - 1/3*sqrt(3)) - (cosh(x)^6 + 6*cosh(x)^5*sinh(x) + 15*cosh(x)^4*sinh(x)^2 + 20*cosh(x)^3*sinh(x)^3 + 15*cosh(x)^2*sinh(x)^4 + 6*cosh(x)*sinh(x)^5 + sinh(x)^6 - 1)*log((2*cosh(x) + 1)/(cosh(x) - sinh(x)))
```


(x))) + (cosh(x)^6 + 6*cosh(x)^5*sinh(x) + 15*cosh(x)^4*sinh(x)^2 + 20*cosh(x)^3*sinh(x)^3 + 15*cosh(x)^2*sinh(x)^4 + 6*cosh(x)*sinh(x)^5 + sinh(x)^6 - 1)*log((2*cosh(x) - 1)/(cosh(x) - sinh(x))) - 2*(cosh(x)^6 + 6*cosh(x)^5*sinh(x) + 15*cosh(x)^4*sinh(x)^2 + 20*cosh(x)^3*sinh(x)^3 + 15*cosh(x)^2*sinh(x)^4 + 6*cosh(x)*sinh(x)^5 + sinh(x)^6 - 1)*log(cosh(x) + sinh(x) + 1) + 2*(cosh(x)^6 + 6*cosh(x)^5*sinh(x) + 15*cosh(x)^4*sinh(x)^2 + 20*cosh(x)^3*sinh(x)^3 + 15*cosh(x)^2*sinh(x)^4 + 6*cosh(x)*sinh(x)^5 + sinh(x)^6 - 1)*log(cosh(x) + sinh(x) - 1) + 6*(21*cosh(x)^6 - 5)*sinh(x) - 30*cosh(x))/(cosh(x)^6 + 6*cosh(x)^5*sinh(x) + 15*cosh(x)^4*sinh(x)^2 + 20*cosh(x)^3*sinh(x)^3 + 15*cosh(x)^2*sinh(x)^4 + 6*cosh(x)*sinh(x)^5 + sinh(x)^6 - 1)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int e^x \coth^2(3x) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(x)*coth(3*x)**2,x)

[Out] Integral(exp(x)*coth(3*x)**2, x)

Giac [A]

time = 0.43, size = 88, normalized size = 0.81

$$-\frac{1}{9}\sqrt{3}\arctan\left(\frac{1}{3}\sqrt{3}(2e^x+1)\right) - \frac{1}{9}\sqrt{3}\arctan\left(\frac{1}{3}\sqrt{3}(2e^x-1)\right) - \frac{2e^x}{3(e^{6x}-1)} + e^x - \frac{1}{18}\log(e^{2x}+e^x+1) + \frac{1}{18}\log(e^{2x}-e^x+1) - \frac{1}{9}\log(e^x+1) + \frac{1}{9}\log(|e^x-1|)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(x)*coth(3*x)^2,x, algorithm="giac")

[Out] $-1/9*\sqrt{3}*\arctan(1/3*\sqrt{3}*(2*e^x + 1)) - 1/9*\sqrt{3}*\arctan(1/3*\sqrt{3}*(2*e^x - 1)) - 2/3*e^x/(e^{6*x} - 1) + e^x - 1/18*\log(e^{2*x} + e^x + 1) + 1/18*\log(e^{2*x} - e^x + 1) - 1/9*\log(e^x + 1) + 1/9*\log(\text{abs}(e^x - 1))$

Mupad [B]

time = 1.38, size = 93, normalized size = 0.86

$$\frac{\ln\left(\frac{2}{3} - \frac{2e^x}{3}\right)}{9} - \frac{\ln\left(-\frac{2e^x}{3} - \frac{2}{3}\right)}{9} + \frac{\ln\left(\left(\frac{2e^x}{3} - \frac{1}{3}\right)^2 + \frac{1}{3}\right)}{18} - \frac{\ln\left(\left(\frac{2e^x}{3} + \frac{1}{3}\right)^2 + \frac{1}{3}\right)}{18} + e^x - \frac{2e^x}{3(e^{6x}-1)} - \frac{\sqrt{3}\operatorname{atan}\left(\sqrt{3}\left(\frac{2e^x}{3} - \frac{1}{3}\right)\right)}{9} - \frac{\sqrt{3}\operatorname{atan}\left(\sqrt{3}\left(\frac{2e^x}{3} + \frac{1}{3}\right)\right)}{9}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(coth(3*x)^2*exp(x),x)

[Out] $\log(2/3 - (2*\exp(x))/3)/9 - \log(-(2*\exp(x))/3 - 2/3)/9 + \log(((2*\exp(x))/3 - 1/3)^2 + 1/3)/18 - \log(((2*\exp(x))/3 + 1/3)^2 + 1/3)/18 + \exp(x) - (2*\exp(x))/(3*(\exp(6*x) - 1)) - (3^{(1/2)}*\operatorname{atan}(3^{(1/2)}*((2*\exp(x))/3 - 1/3)))/9 - (3^{(1/2)}*\operatorname{atan}(3^{(1/2)}*((2*\exp(x))/3 + 1/3)))/9$

3.222 $\int e^x \tanh^2(4x) dx$

Optimal. Leaf size=382

$$e^x + \frac{e^x}{2(1+e^{8x})} + \frac{\operatorname{ArcTan}\left(\frac{\sqrt{2-\sqrt{2}}-2e^x}{\sqrt{2+\sqrt{2}}}\right)}{8\sqrt{2(2-\sqrt{2})}} + \frac{\operatorname{ArcTan}\left(\frac{\sqrt{2+\sqrt{2}}-2e^x}{\sqrt{2-\sqrt{2}}}\right)}{8\sqrt{2(2+\sqrt{2})}} - \frac{\operatorname{ArcTan}\left(\frac{\sqrt{2-\sqrt{2}}+2e^x}{\sqrt{2+\sqrt{2}}}\right)}{8\sqrt{2(2-\sqrt{2})}} - \frac{\operatorname{ArcTan}\left(\frac{\sqrt{2+\sqrt{2}}+2e^x}{\sqrt{2-\sqrt{2}}}\right)}{8\sqrt{2(2+\sqrt{2})}}$$

[Out] $\exp(x)+1/2*\exp(x)/(1+\exp(8*x))+1/32*\ln(1+\exp(2*x)-\exp(x)*(2-2^{(1/2)})^{(1/2)})*(2-2^{(1/2)})^{(1/2)}-1/32*\ln(1+\exp(2*x)+\exp(x)*(2-2^{(1/2)})^{(1/2)})*(2-2^{(1/2)})^{(1/2)}+1/8*\arctan((-2*\exp(x)+(2-2^{(1/2)})^{(1/2)})/(2+2^{(1/2)})^{(1/2)})/(4-2*2^{(1/2)})^{(1/2)}-1/8*\arctan((2*\exp(x)+(2-2^{(1/2)})^{(1/2)})/(2+2^{(1/2)})^{(1/2)})/(4-2*2^{(1/2)})^{(1/2)}+1/32*\ln(1+\exp(2*x)-\exp(x)*(2+2^{(1/2)})^{(1/2)})*(2+2^{(1/2)})^{(1/2)}-1/32*\ln(1+\exp(2*x)+\exp(x)*(2+2^{(1/2)})^{(1/2)})*(2+2^{(1/2)})^{(1/2)}+1/8*\arctan((-2*\exp(x)+(2+2^{(1/2)})^{(1/2)})/(2-2^{(1/2)})^{(1/2)})/(4+2*2^{(1/2)})^{(1/2)}-1/8*\arctan((2*\exp(x)+(2+2^{(1/2)})^{(1/2)})/(2-2^{(1/2)})^{(1/2)})/(4+2*2^{(1/2)})^{(1/2)}$

Rubi [A]

time = 0.31, antiderivative size = 382, normalized size of antiderivative = 1.00, number of steps used = 23, number of rules used = 9, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.900$, Rules used = {2320, 398, 294, 219, 1183, 648, 632, 210, 642}

$$\frac{\operatorname{ArcTan}\left(\frac{\sqrt{2-\sqrt{2}}-e^x}{\sqrt{2+\sqrt{2}}}\right)}{8\sqrt{2(2-\sqrt{2})}} + \frac{\operatorname{ArcTan}\left(\frac{\sqrt{2+\sqrt{2}}-e^x}{\sqrt{2-\sqrt{2}}}\right)}{8\sqrt{2(2+\sqrt{2})}} - \frac{\operatorname{ArcTan}\left(\frac{e^x-\sqrt{2-\sqrt{2}}}{\sqrt{2+\sqrt{2}}}\right)}{8\sqrt{2(2-\sqrt{2})}} - \frac{\operatorname{ArcTan}\left(\frac{e^x-\sqrt{2+\sqrt{2}}}{\sqrt{2-\sqrt{2}}}\right)}{8\sqrt{2(2+\sqrt{2})}} + e^x + \frac{e^x}{2(e^{8x}+1)} + \frac{1}{32}\sqrt{2-\sqrt{2}}\log(-\sqrt{2-\sqrt{2}}e^x+e^{2x}+1) - \frac{1}{32}\sqrt{2-\sqrt{2}}\log(\sqrt{2-\sqrt{2}}e^x+e^{2x}+1) + \frac{1}{32}\sqrt{2+\sqrt{2}}\log(-\sqrt{2+\sqrt{2}}e^x+e^{2x}+1) - \frac{1}{32}\sqrt{2+\sqrt{2}}\log(\sqrt{2+\sqrt{2}}e^x+e^{2x}+1)$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^x*\text{Tanh}[4*x]^2,x]$

[Out] $E^x + E^x/(2*(1 + E^{(8*x)})) + \operatorname{ArcTan}[(\operatorname{Sqrt}[2 - \operatorname{Sqrt}[2]] - 2*E^x)/\operatorname{Sqrt}[2 + \operatorname{Sqrt}[2]]]/(8*\operatorname{Sqrt}[2*(2 - \operatorname{Sqrt}[2])]) + \operatorname{ArcTan}[(\operatorname{Sqrt}[2 + \operatorname{Sqrt}[2]] - 2*E^x)/\operatorname{Sqrt}[2 - \operatorname{Sqrt}[2]]]/(8*\operatorname{Sqrt}[2*(2 + \operatorname{Sqrt}[2])]) - \operatorname{ArcTan}[(\operatorname{Sqrt}[2 - \operatorname{Sqrt}[2]] + 2*E^x)/\operatorname{Sqrt}[2 + \operatorname{Sqrt}[2]]]/(8*\operatorname{Sqrt}[2*(2 - \operatorname{Sqrt}[2])]) - \operatorname{ArcTan}[(\operatorname{Sqrt}[2 + \operatorname{Sqrt}[2]] + 2*E^x)/\operatorname{Sqrt}[2 - \operatorname{Sqrt}[2]]]/(8*\operatorname{Sqrt}[2*(2 + \operatorname{Sqrt}[2])]) + (\operatorname{Sqrt}[2 - \operatorname{Sqrt}[2]]*\operatorname{Log}[1 - \operatorname{Sqrt}[2 - \operatorname{Sqrt}[2]]*E^x + E^{(2*x)}])/32 - (\operatorname{Sqrt}[2 - \operatorname{Sqrt}[2]]*\operatorname{Log}[1 + \operatorname{Sqrt}[2 - \operatorname{Sqrt}[2]]*E^x + E^{(2*x)}])/32 + (\operatorname{Sqrt}[2 + \operatorname{Sqrt}[2]]*\operatorname{Log}[1 - \operatorname{Sqrt}[2 + \operatorname{Sqrt}[2]]*E^x + E^{(2*x)}])/32 - (\operatorname{Sqrt}[2 + \operatorname{Sqrt}[2]]*\operatorname{Log}[1 + \operatorname{Sqrt}[2 + \operatorname{Sqrt}[2]]*E^x + E^{(2*x)}])/32$

Rule 210

$\text{Int}[(a_+ + (b_+)*(x_+)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1})*\operatorname{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /;$ $\text{FreeQ}\{a, b\}, x \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

Rule 219

```
Int[((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := With[{r = Numerator[Rt[a/b,
  4]], s = Denominator[Rt[a/b, 4]]}, Dist[r/(2*Sqrt[2]*a), Int[(Sqrt[2]*r -
s*x^(n/4))/(r^2 - Sqrt[2]*r*s*x^(n/4) + s^2*x^(n/2)), x], x] + Dist[r/(2*Sq
rt[2]*a), Int[(Sqrt[2]*r + s*x^(n/4))/(r^2 + Sqrt[2]*r*s*x^(n/4) + s^2*x^(n
/2)), x], x]] /; FreeQ[{a, b}, x] && IGtQ[n/4, 1] && GtQ[a/b, 0]
```

Rule 294

```
Int[((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(
n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n
*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x]
/; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !I
LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 398

```
Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol]
:= Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a
, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q,
0] && GeQ[p, -q]
```

Rule 632

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Dist[-2, Subst[I
nt[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c},
x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 648

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := D
ist[(2*c*d - b*e)/(2*c), Int[1/(a + b*x + c*x^2), x], x] + Dist[e/(2*c), In
t[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ
[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0] && !NiceSqrtQ[b^2 - 4*a*c]
```

Rule 1183

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (b_)*(x_)^2 + (c_)*(x_)^4), x_Symbol] :
> With[{q = Rt[a/c, 2]}, With[{r = Rt[2*q - b/c, 2]}, Dist[1/(2*c*q*r), Int
[(d*r - (d - e*q)*x)/(q - r*x + x^2), x], x] + Dist[1/(2*c*q*r), Int[(d*r +
```

$(d - e*q)*x)/(q + r*x + x^2), x], x]] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0] \ \&\& \ \text{NeQ}[c*d^2 - b*d*e + a*e^2, 0] \ \&\& \ \text{NegQ}[b^2 - 4*a*c]$

Rule 2320

$\text{Int}[u_, x_Symbol] \ :> \ \text{With}[\{v = \text{FunctionOfExponential}[u, x]\}, \text{Dist}[v/D[v, x], \text{Subst}[\text{Int}[\text{FunctionOfExponentialFunction}[u, x]/x, x], x, v], x]] /; \text{FunctionOfExponentialQ}[u, x] \ \&\& \ \text{!MatchQ}[u, (w_)*((a_)*(v_)^{(n_)}^{\{m_}}) /; \text{FreeQ}[\{a, m, n\}, x] \ \&\& \ \text{IntegerQ}[m*n] \ \&\& \ \text{!MatchQ}[u, E^{\{(c_)*((a_)+ (b_)*x)}*(F_)[v_]} /; \text{FreeQ}[\{a, b, c\}, x] \ \&\& \ \text{InverseFunctionQ}[F[x]]]$

Rubi steps

$$\begin{aligned}
 \int e^x \tanh^2(4x) dx &= \text{Subst} \left(\int \frac{(1-x^8)^2}{(1+x^8)^2} dx, x, e^x \right) \\
 &= \text{Subst} \left(\int \left(1 - \frac{4x^8}{(1+x^8)^2} \right) dx, x, e^x \right) \\
 &= e^x - 4 \text{Subst} \left(\int \frac{x^8}{(1+x^8)^2} dx, x, e^x \right) \\
 &= e^x + \frac{e^x}{2(1+e^{8x})} - \frac{1}{2} \text{Subst} \left(\int \frac{1}{1+x^8} dx, x, e^x \right) \\
 &= e^x + \frac{e^x}{2(1+e^{8x})} - \frac{\text{Subst} \left(\int \frac{\sqrt{2}-x^2}{1-\sqrt{2}x^2+x^4} dx, x, e^x \right)}{4\sqrt{2}} - \frac{\text{Subst} \left(\int \frac{\sqrt{2}+x^2}{1+\sqrt{2}x^2+x^4} dx, x, e^x \right)}{4\sqrt{2}} \\
 &= e^x + \frac{e^x}{2(1+e^{8x})} - \frac{\text{Subst} \left(\int \frac{\sqrt{2}(2-\sqrt{2}) - (-1+\sqrt{2})x}{1-\sqrt{2-\sqrt{2}}x+x^2} dx, x, e^x \right)}{8\sqrt{2}(2-\sqrt{2})} - \frac{\text{Subst} \left(\int \frac{\sqrt{2}(2+\sqrt{2}) - (-1-\sqrt{2})x}{1+\sqrt{2-\sqrt{2}}x+x^2} dx, x, e^x \right)}{8\sqrt{2}(2+\sqrt{2})} \\
 &= e^x + \frac{e^x}{2(1+e^{8x})} - \frac{1}{16} \sqrt{\frac{1}{2}(3-2\sqrt{2})} \text{Subst} \left(\int \frac{1}{1-\sqrt{2+\sqrt{2}}x+x^2} dx, x, e^x \right) - \frac{1}{16} \sqrt{\frac{1}{2}(3+2\sqrt{2})} \text{Subst} \left(\int \frac{1}{1+\sqrt{2-\sqrt{2}}x+x^2} dx, x, e^x \right) \\
 &= e^x + \frac{e^x}{2(1+e^{8x})} + \frac{1}{32} \sqrt{2-\sqrt{2}} \log \left(1 - \sqrt{2-\sqrt{2}} e^x + e^{2x} \right) - \frac{1}{32} \sqrt{2-\sqrt{2}} \log \left(1 + \sqrt{2-\sqrt{2}} e^x + e^{2x} \right) \\
 &= e^x + \frac{e^x}{2(1+e^{8x})} + \frac{1}{16} \sqrt{2+\sqrt{2}} \tan^{-1} \left(\frac{\sqrt{2-\sqrt{2}} - 2e^x}{\sqrt{2+\sqrt{2}}} \right) + \frac{1}{16} \sqrt{2-\sqrt{2}} \tan^{-1} \left(\frac{\sqrt{2+\sqrt{2}} - 2e^x}{\sqrt{2-\sqrt{2}}} \right)
 \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

time = 0.04, size = 51, normalized size = 0.13

$$e^x + \frac{e^x}{2(1 + e^{8x})} + \frac{1}{16} \text{RootSum} \left[1 + \#1^8 \&, \frac{x - \log(e^x - \#1)}{\#1^7} \& \right]$$

Antiderivative was successfully verified.

[In] Integrate[E^x*Tanh[4*x]^2,x]

[Out] E^x + E^x/(2*(1 + E^(8*x))) + RootSum[1 + #1^8 & , (x - Log[E^x - #1])/#1^7 &]/16

Maple [C] Result contains higher order function than in optimal. Order 9 vs. order 3.

time = 0.88, size = 36, normalized size = 0.09

method	result	size
risch	$e^x + \frac{e^x}{2+2e^{8x}} + \left(\sum_{_R=\text{RootOf}(4294967296_Z^8+1)} _R \ln(e^x - 16_R) \right)$	36

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(x)*tanh(4*x)^2,x,method=_RETURNVERBOSE)

[Out] exp(x)+1/2*exp(x)/(1+exp(8*x))+sum(_R*ln(exp(x)-16*_R),_R=RootOf(4294967296*_Z^8+1))

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(x)*tanh(4*x)^2,x, algorithm="maxima")

[Out] 1/2*(2*e^(9*x) + 3*e^x)/(e^(8*x) + 1) - integrate(1/2*e^x/(e^(8*x) + 1), x)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 1373 vs. 2(265) = 530.

time = 0.39, size = 1373, normalized size = 3.59

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(x)*tanh(4*x)^2,x, algorithm="fricas")

```
[Out] 1/128*(8*sqrt(-sqrt(2) + 2)*(e^(8*x) + 1)*arctan((2*sqrt(sqrt(2) + 2)*
e^x + e^(2*x) + 1) - sqrt(sqrt(2) + 2) - 2*e^x)/sqrt(-sqrt(2) + 2)) + 8*sqrt
(-sqrt(2) + 2)*(e^(8*x) + 1)*arctan((2*sqrt(-sqrt(2) + 2)*e^x + e^(2*
x) + 1) + sqrt(sqrt(2) + 2) - 2*e^x)/sqrt(-sqrt(2) + 2)) - 2*sqrt(-sqrt(2)
+ 2)*(e^(8*x) + 1)*log(sqrt(-sqrt(2) + 2)*e^x + e^(2*x) + 1) + 2*sqrt(-sqrt
(2) + 2)*(e^(8*x) + 1)*log(-sqrt(-sqrt(2) + 2)*e^x + e^(2*x) + 1) + 8*(sqrt
(sqrt(2) + 2)*e^(8*x) + sqrt(sqrt(2) + 2))*arctan((2*sqrt(sqrt(-sqrt(2) + 2)
)*e^x + e^(2*x) + 1) - sqrt(-sqrt(2) + 2) - 2*e^x)/sqrt(sqrt(2) + 2)) + 8*(
sqrt(sqrt(2) + 2)*e^(8*x) + sqrt(sqrt(2) + 2))*arctan((2*sqrt(-sqrt(-sqrt(2)
) + 2)*e^x + e^(2*x) + 1) + sqrt(-sqrt(2) + 2) - 2*e^x)/sqrt(sqrt(2) + 2))
+ 4*(sqrt(2)*sqrt(sqrt(2) + 2)*e^(8*x) + (sqrt(2)*e^(8*x) + sqrt(2))*sqrt(-
sqrt(2) + 2) + sqrt(2)*sqrt(sqrt(2) + 2))*arctan(-(2*sqrt(2)*e^x - sqrt(2)*
sqrt(2*sqrt(2)*sqrt(sqrt(2) + 2)*e^x - 2*sqrt(2)*sqrt(-sqrt(2) + 2)*e^x + 4
*e^(2*x) + 4) + sqrt(sqrt(2) + 2) - sqrt(-sqrt(2) + 2))/(sqrt(sqrt(2) + 2)
+ sqrt(-sqrt(2) + 2))) + 4*(sqrt(2)*sqrt(sqrt(2) + 2)*e^(8*x) + (sqrt(2)*e^
(8*x) + sqrt(2))*sqrt(-sqrt(2) + 2) + sqrt(2)*sqrt(sqrt(2) + 2))*arctan(-2
*sqrt(2)*e^x - sqrt(2)*sqrt(-2*sqrt(2)*sqrt(sqrt(2) + 2)*e^x + 2*sqrt(2)*sq
rt(-sqrt(2) + 2)*e^x + 4*e^(2*x) + 4) - sqrt(sqrt(2) + 2) + sqrt(-sqrt(2) +
2))/(sqrt(sqrt(2) + 2) + sqrt(-sqrt(2) + 2))) - 4*(sqrt(2)*sqrt(sqrt(2) +
2)*e^(8*x) - (sqrt(2)*e^(8*x) + sqrt(2))*sqrt(-sqrt(2) + 2) + sqrt(2)*sqrt(
sqrt(2) + 2))*arctan((2*sqrt(2)*e^x - sqrt(2)*sqrt(2*sqrt(2)*sqrt(sqrt(2) +
2)*e^x + 2*sqrt(2)*sqrt(-sqrt(2) + 2)*e^x + 4*e^(2*x) + 4) + sqrt(sqrt(2)
+ 2) + sqrt(-sqrt(2) + 2))/(sqrt(sqrt(2) + 2) - sqrt(-sqrt(2) + 2))) - 4*(s
qrt(2)*sqrt(sqrt(2) + 2)*e^(8*x) - (sqrt(2)*e^(8*x) + sqrt(2))*sqrt(-sqrt(2)
+ 2) + sqrt(2)*sqrt(sqrt(2) + 2))*arctan((2*sqrt(2)*e^x - sqrt(2)*sqrt(-2
*sqrt(2)*sqrt(sqrt(2) + 2)*e^x - 2*sqrt(2)*sqrt(-sqrt(2) + 2)*e^x + 4*e^(2*
x) + 4) - sqrt(sqrt(2) + 2) - sqrt(-sqrt(2) + 2))/(sqrt(sqrt(2) + 2) - sqrt
(-sqrt(2) + 2))) - (sqrt(2)*sqrt(sqrt(2) + 2)*e^(8*x) + (sqrt(2)*e^(8*x) +
sqrt(2))*sqrt(-sqrt(2) + 2) + sqrt(2)*sqrt(sqrt(2) + 2))*log(2*sqrt(2)*sqrt
(sqrt(2) + 2)*e^x + 2*sqrt(2)*sqrt(-sqrt(2) + 2)*e^x + 4*e^(2*x) + 4) - (sq
rt(2)*sqrt(sqrt(2) + 2)*e^(8*x) - (sqrt(2)*e^(8*x) + sqrt(2))*sqrt(-sqrt(2)
+ 2) + sqrt(2)*sqrt(sqrt(2) + 2))*log(2*sqrt(2)*sqrt(sqrt(2) + 2)*e^x - 2*
sqrt(2)*sqrt(-sqrt(2) + 2)*e^x + 4*e^(2*x) + 4) + (sqrt(2)*sqrt(sqrt(2) + 2)
)*e^(8*x) - (sqrt(2)*e^(8*x) + sqrt(2))*sqrt(-sqrt(2) + 2) + sqrt(2)*sqrt(s
qrt(2) + 2))*log(-2*sqrt(2)*sqrt(sqrt(2) + 2)*e^x + 2*sqrt(2)*sqrt(-sqrt(2)
+ 2)*e^x + 4*e^(2*x) + 4) + (sqrt(2)*sqrt(sqrt(2) + 2)*e^(8*x) + (sqrt(2)*
e^(8*x) + sqrt(2))*sqrt(-sqrt(2) + 2) + sqrt(2)*sqrt(sqrt(2) + 2))*log(-2*s
qrt(2)*sqrt(sqrt(2) + 2)*e^x - 2*sqrt(2)*sqrt(-sqrt(2) + 2)*e^x + 4*e^(2*x)
+ 4) - 2*(sqrt(sqrt(2) + 2)*e^(8*x) + sqrt(sqrt(2) + 2))*log(sqrt(sqrt(2)
+ 2)*e^x + e^(2*x) + 1) + 2*(sqrt(sqrt(2) + 2)*e^(8*x) + sqrt(sqrt(2) + 2))
*log(-sqrt(sqrt(2) + 2)*e^x + e^(2*x) + 1) + 128*e^(9*x) + 192*e^x)/(e^(8*x)
+ 1)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int e^x \tanh^2(4x) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(x)*tanh(4*x)**2,x)

[Out] Integral(exp(x)*tanh(4*x)**2, x)

Giac [A]

time = 0.42, size = 263, normalized size = 0.69

$$\frac{1}{16}\sqrt{-\sqrt{2}+2}\operatorname{arctan}\left(\frac{\sqrt{\sqrt{2}+2}+2x}{\sqrt{-\sqrt{2}+2}}\right) - \frac{1}{16}\sqrt{-\sqrt{2}+2}\operatorname{arctan}\left(\frac{\sqrt{\sqrt{2}+2}-2x}{\sqrt{-\sqrt{2}+2}}\right) - \frac{1}{16}\sqrt{\sqrt{2}+2}\operatorname{arctan}\left(\frac{\sqrt{-\sqrt{2}+2}+2x}{\sqrt{\sqrt{2}+2}}\right) - \frac{1}{16}\sqrt{\sqrt{2}+2}\operatorname{arctan}\left(\frac{\sqrt{-\sqrt{2}+2}-2x}{\sqrt{\sqrt{2}+2}}\right) - \frac{1}{16}\sqrt{\sqrt{2}+2}\log(\sqrt{\sqrt{2}+2}e^{x+\rho^{2x}}) + \frac{1}{16}\sqrt{\sqrt{2}+2}\log(-\sqrt{\sqrt{2}+2}e^{x+\rho^{2x}}) - \frac{1}{16}\sqrt{-\sqrt{2}+2}\log(\sqrt{-\sqrt{2}+2}e^{x+\rho^{2x}}) + \frac{1}{16}\sqrt{-\sqrt{2}+2}\log(-\sqrt{-\sqrt{2}+2}e^{x+\rho^{2x}}) + \frac{e^x}{2(\rho^{2x}+1)} + e^x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(x)*tanh(4*x)^2,x, algorithm="giac")

[Out] $-1/16*\sqrt{-\sqrt{2}+2}*\arctan((\sqrt{\sqrt{2}+2}+2)*e^x/\sqrt{-\sqrt{2}+2}) - 1/16*\sqrt{-\sqrt{2}+2}*\arctan(-(\sqrt{\sqrt{2}+2}-2)*e^x/\sqrt{-\sqrt{2}+2}) - 1/16*\sqrt{\sqrt{2}+2}*\arctan((\sqrt{-\sqrt{2}+2}+2)*e^x/\sqrt{\sqrt{2}+2}) - 1/16*\sqrt{\sqrt{2}+2}*\arctan(-(\sqrt{-\sqrt{2}+2}-2)*e^x/\sqrt{\sqrt{2}+2}) - 1/32*\sqrt{\sqrt{2}+2}*\log(\sqrt{\sqrt{2}+2}*e^x + e^{2*x} + 1) + 1/32*\sqrt{\sqrt{2}+2}*\log(-\sqrt{\sqrt{2}+2}*e^x + e^{2*x} + 1) - 1/32*\sqrt{-\sqrt{2}+2}*\log(\sqrt{-\sqrt{2}+2}*e^x + e^{2*x} + 1) + 1/32*\sqrt{-\sqrt{2}+2}*\log(-\sqrt{-\sqrt{2}+2}*e^x + e^{2*x} + 1) + 1/2*e^x/(e^{8*x} + 1) + e^x$

Mupad [B]

time = 3.98, size = 474, normalized size = 1.24

$$\dots$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tanh(4*x)^2*exp(x),x)

[Out] $\exp(x) + \exp(x)/(2*(\exp(8*x) + 1)) + \log(\exp(x)/2 - (2^{1/2} + 2)^{1/2}/4 - ((2 - 2^{1/2})^{1/2}*1i)/4)*((2^{1/2} + 2)^{1/2}/32 + ((2 - 2^{1/2})^{1/2})^{1/2}*1i)/32) - \log(\exp(x)/2 + (2^{1/2} + 2)^{1/2}/4 + ((2 - 2^{1/2})^{1/2}*1i)/4)*((2^{1/2} + 2)^{1/2}/32 + ((2 - 2^{1/2})^{1/2})^{1/2}*1i)/32) + \log(\exp(x)/2 - ((2^{1/2} + 2)^{1/2}*1i)/4 + (2 - 2^{1/2})^{1/2}/4)*(((2^{1/2} + 2)^{1/2}*1i)/32 - (2 - 2^{1/2})^{1/2}/32) - \log(\exp(x)/2 + ((2^{1/2} + 2)^{1/2}*1i)/4 - (2 - 2^{1/2})^{1/2}/4)*(((2^{1/2} + 2)^{1/2}*1i)/32 - (2 - 2^{1/2})^{1/2}/32) + 2^{1/2}*\log(\exp(x)/2 - 2^{1/2}*((2^{1/2} + 2)^{1/2}/32 + ((2 - 2^{1/2})^{1/2})^{1/2}*1i)/32)*(4 + 4i))*((2^{1/2} + 2)^{1/2}/32 + ((2 - 2^{1/2})^{1/2})^{1/2}*1i)/32)*(1/2 + 1i/2) + 2^{1/2}*\log(\exp(x)/2 - 2^{1/2}*((2^{1/2} + 2)^{1/2})$

$$\begin{aligned}
& /32 + ((2 - 2^{(1/2)})^{(1/2)} * 1i) / 32) * (4 - 4i) * ((2^{(1/2)} + 2)^{(1/2)} / 32 + ((2 \\
& - 2^{(1/2)})^{(1/2)} * 1i) / 32) * (1/2 - 1i/2) - 2^{(1/2)} * \log(\exp(x)/2 + 2^{(1/2)} * ((2^{(1/2)} \\
& + 2)^{(1/2)} / 32 + ((2 - 2^{(1/2)})^{(1/2)} * 1i) / 32) * (4 - 4i) * ((2^{(1/2)} + 2)^{(1/2)} / 32 \\
& + ((2 - 2^{(1/2)})^{(1/2)} * 1i) / 32) * (1/2 - 1i/2) - 2^{(1/2)} * \log(\exp(x) / \\
& 2 + 2^{(1/2)} * ((2^{(1/2)} + 2)^{(1/2)} / 32 + ((2 - 2^{(1/2)})^{(1/2)} * 1i) / 32) * (4 + 4i) \\
&) * ((2^{(1/2)} + 2)^{(1/2)} / 32 + ((2 - 2^{(1/2)})^{(1/2)} * 1i) / 32) * (1/2 + 1i/2)
\end{aligned}$$

3.223 $\int e^x \tanh(4x) dx$

Optimal. Leaf size=366

$$e^x + \frac{\operatorname{ArcTan}\left(\frac{\sqrt{2-\sqrt{2}}-2e^x}{\sqrt{2+\sqrt{2}}}\right)}{2\sqrt{2(2-\sqrt{2})}} + \frac{\operatorname{ArcTan}\left(\frac{\sqrt{2+\sqrt{2}}-2e^x}{\sqrt{2-\sqrt{2}}}\right)}{2\sqrt{2(2+\sqrt{2})}} - \frac{\operatorname{ArcTan}\left(\frac{\sqrt{2-\sqrt{2}}+2e^x}{\sqrt{2+\sqrt{2}}}\right)}{2\sqrt{2(2-\sqrt{2})}} - \frac{\operatorname{ArcTan}\left(\frac{\sqrt{2+\sqrt{2}}+2e^x}{\sqrt{2-\sqrt{2}}}\right)}{2\sqrt{2(2+\sqrt{2})}}$$

[Out] $\exp(x)+1/8*\ln(1+\exp(2*x)-\exp(x)*(2-2^{(1/2)})^{(1/2)})*(2-2^{(1/2)})^{(1/2)}-1/8*\ln(1+\exp(2*x)+\exp(x)*(2-2^{(1/2)})^{(1/2)})*(2-2^{(1/2)})^{(1/2)}+1/2*\arctan((-2*\exp(x)+(2-2^{(1/2)})^{(1/2)})/(2+2^{(1/2)})^{(1/2)})/(4-2*2^{(1/2)})^{(1/2)}-1/2*\arctan((2*\exp(x)+(2-2^{(1/2)})^{(1/2)})/(2+2^{(1/2)})^{(1/2)})/(4-2*2^{(1/2)})^{(1/2)}+1/8*\ln(1+\exp(2*x)-\exp(x)*(2+2^{(1/2)})^{(1/2)})*(2+2^{(1/2)})^{(1/2)}-1/8*\ln(1+\exp(2*x)+\exp(x)*(2+2^{(1/2)})^{(1/2)})*(2+2^{(1/2)})^{(1/2)}+1/2*\arctan((-2*\exp(x)+(2+2^{(1/2)})^{(1/2)})/(2-2^{(1/2)})^{(1/2)})/(4+2*2^{(1/2)})^{(1/2)}-1/2*\arctan((2*\exp(x)+(2+2^{(1/2)})^{(1/2)})/(2-2^{(1/2)})^{(1/2)})/(4+2*2^{(1/2)})^{(1/2)}$

Rubi [A]

time = 0.17, antiderivative size = 366, normalized size of antiderivative = 1.00, number of steps used = 21, number of rules used = 8, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {2320, 396, 219, 1183, 648, 632, 210, 642}

$$\frac{\operatorname{ArcTan}\left(\frac{\sqrt{2-\sqrt{2}}-2e^x}{\sqrt{2+\sqrt{2}}}\right)}{2\sqrt{2(2-\sqrt{2})}} + \frac{\operatorname{ArcTan}\left(\frac{\sqrt{2+\sqrt{2}}-2e^x}{\sqrt{2-\sqrt{2}}}\right)}{2\sqrt{2(2+\sqrt{2})}} - \frac{\operatorname{ArcTan}\left(\frac{\sqrt{2-\sqrt{2}}+2e^x}{\sqrt{2+\sqrt{2}}}\right)}{2\sqrt{2(2-\sqrt{2})}} - \frac{\operatorname{ArcTan}\left(\frac{\sqrt{2+\sqrt{2}}+2e^x}{\sqrt{2-\sqrt{2}}}\right)}{2\sqrt{2(2+\sqrt{2})}} + e^x + \frac{1}{8}\sqrt{2-\sqrt{2}} \log(-\sqrt{2-\sqrt{2}}e^x + e^{2x} + 1) - \frac{1}{8}\sqrt{2-\sqrt{2}} \log(\sqrt{2-\sqrt{2}}e^x + e^{2x} + 1) + \frac{1}{8}\sqrt{2+\sqrt{2}} \log(-\sqrt{2+\sqrt{2}}e^x + e^{2x} + 1) - \frac{1}{8}\sqrt{2+\sqrt{2}} \log(\sqrt{2+\sqrt{2}}e^x + e^{2x} + 1)$$

Antiderivative was successfully verified.

[In] Int[E^x*Tanh[4*x], x]

[Out] $E^x + \operatorname{ArcTan}[(\operatorname{Sqrt}[2 - \operatorname{Sqrt}[2]] - 2*E^x)/\operatorname{Sqrt}[2 + \operatorname{Sqrt}[2]]]/(2*\operatorname{Sqrt}[2*(2 - \operatorname{Sqrt}[2])]) + \operatorname{ArcTan}[(\operatorname{Sqrt}[2 + \operatorname{Sqrt}[2]] - 2*E^x)/\operatorname{Sqrt}[2 - \operatorname{Sqrt}[2]]]/(2*\operatorname{Sqrt}[2*(2 + \operatorname{Sqrt}[2])]) - \operatorname{ArcTan}[(\operatorname{Sqrt}[2 - \operatorname{Sqrt}[2]] + 2*E^x)/\operatorname{Sqrt}[2 + \operatorname{Sqrt}[2]]]/(2*\operatorname{Sqrt}[2*(2 - \operatorname{Sqrt}[2])]) - \operatorname{ArcTan}[(\operatorname{Sqrt}[2 + \operatorname{Sqrt}[2]] + 2*E^x)/\operatorname{Sqrt}[2 - \operatorname{Sqrt}[2]]]/(2*\operatorname{Sqrt}[2*(2 + \operatorname{Sqrt}[2])]) + (\operatorname{Sqrt}[2 - \operatorname{Sqrt}[2]]*\operatorname{Log}[1 - \operatorname{Sqrt}[2 - \operatorname{Sqrt}[2]]]*E^x + E^{(2*x)})/8 - (\operatorname{Sqrt}[2 - \operatorname{Sqrt}[2]]*\operatorname{Log}[1 + \operatorname{Sqrt}[2 - \operatorname{Sqrt}[2]]]*E^x + E^{(2*x)})/8 + (\operatorname{Sqrt}[2 + \operatorname{Sqrt}[2]]*\operatorname{Log}[1 - \operatorname{Sqrt}[2 + \operatorname{Sqrt}[2]]]*E^x + E^{(2*x)})/8 - (\operatorname{Sqrt}[2 + \operatorname{Sqrt}[2]]*\operatorname{Log}[1 + \operatorname{Sqrt}[2 + \operatorname{Sqrt}[2]]]*E^x + E^{(2*x)})/8$

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 219

```
Int[((a_) + (b_.)*(x_)^(n_))^-1), x_Symbol] := With[{r = Numerator[Rt[a/b,
4]], s = Denominator[Rt[a/b, 4]]}, Dist[r/(2*Sqrt[2]*a), Int[(Sqrt[2]*r -
s*x^(n/4))/(r^2 - Sqrt[2]*r*s*x^(n/4) + s^2*x^(n/2)), x], x] + Dist[r/(2*Sq
rt[2]*a), Int[(Sqrt[2]*r + s*x^(n/4))/(r^2 + Sqrt[2]*r*s*x^(n/4) + s^2*x^(n
/2)), x], x]] /; FreeQ[{a, b}, x] && IGtQ[n/4, 1] && GtQ[a/b, 0]
```

Rule 396

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Si
mp[d*x*((a + b*x^n)^(p + 1)/(b*(n*(p + 1) + 1))), x] - Dist[(a*d - b*c*(n*(
p + 1) + 1))/(b*(n*(p + 1) + 1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b,
c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]
```

Rule 632

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^-1), x_Symbol] := Dist[-2, Subst[In
t[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c},
x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 648

```
Int[((d_.) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := D
ist[(2*c*d - b*e)/(2*c), Int[1/(a + b*x + c*x^2), x], x] + Dist[e/(2*c), In
t[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ
[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0] && !NiceSqrtQ[b^2 - 4*a*c]
```

Rule 1183

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4), x_Symbol] :
> With[{q = Rt[a/c, 2]}, With[{r = Rt[2*q - b/c, 2]}, Dist[1/(2*c*q*r), Int
[(d*r - (d - e*q)*x)/(q - r*x + x^2), x], x] + Dist[1/(2*c*q*r), Int[(d*r +
(d - e*q)*x)/(q + r*x + x^2), x], x]]] /; FreeQ[{a, b, c, d, e}, x] && NeQ
[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NegQ[b^2 - 4*a*c]
```

Rule 2320

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x]
, Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; Functi
onOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ[
{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*
```

(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]

Rubi steps

$$\begin{aligned}
 \int e^x \tanh(4x) dx &= \text{Subst} \left(\int \frac{-1+x^8}{1+x^8} dx, x, e^x \right) \\
 &= e^x - 2 \text{Subst} \left(\int \frac{1}{1+x^8} dx, x, e^x \right) \\
 &= e^x - \frac{\text{Subst} \left(\int \frac{\sqrt{2}-x^2}{1-\sqrt{2}x^2+x^4} dx, x, e^x \right)}{\sqrt{2}} - \frac{\text{Subst} \left(\int \frac{\sqrt{2}+x^2}{1+\sqrt{2}x^2+x^4} dx, x, e^x \right)}{\sqrt{2}} \\
 &= e^x - \frac{\text{Subst} \left(\int \frac{\sqrt{2(2-\sqrt{2})} - (-1+\sqrt{2})x}{1-\sqrt{2-\sqrt{2}}x+x^2} dx, x, e^x \right)}{2\sqrt{2(2-\sqrt{2})}} - \frac{\text{Subst} \left(\int \frac{\sqrt{2(2-\sqrt{2})} + (-1+\sqrt{2})x}{1+\sqrt{2-\sqrt{2}}x+x^2} dx, x, e^x \right)}{2\sqrt{2(2-\sqrt{2})}} \\
 &= e^x - \frac{1}{4} \sqrt{\frac{1}{2}(3-2\sqrt{2})} \text{Subst} \left(\int \frac{1}{1-\sqrt{2+\sqrt{2}}x+x^2} dx, x, e^x \right) - \frac{1}{4} \sqrt{\frac{1}{2}(3-2\sqrt{2})} \text{Subst} \left(\int \frac{1}{1+\sqrt{2+\sqrt{2}}x+x^2} dx, x, e^x \right) \\
 &= e^x + \frac{1}{8} \sqrt{2-\sqrt{2}} \log \left(1 - \sqrt{2-\sqrt{2}} e^x + e^{2x} \right) - \frac{1}{8} \sqrt{2-\sqrt{2}} \log \left(1 + \sqrt{2-\sqrt{2}} e^x + e^{2x} \right) \\
 &= e^x + \frac{1}{4} \sqrt{2+\sqrt{2}} \tan^{-1} \left(\frac{\sqrt{2-\sqrt{2}} - 2e^x}{\sqrt{2+\sqrt{2}}} \right) + \frac{1}{4} \sqrt{2-\sqrt{2}} \tan^{-1} \left(\frac{\sqrt{2+\sqrt{2}} - 2e^x}{\sqrt{2-\sqrt{2}}} \right)
 \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 24, normalized size = 0.07

$$e^x - 2e^x {}_2F_1 \left(\frac{1}{8}, 1; \frac{9}{8}; -e^{8x} \right)$$

Antiderivative was successfully verified.

[In] Integrate[E^x*Tanh[4*x], x]

[Out] E^x - 2*E^x*Hypergeometric2F1[1/8, 1, 9/8, -E^(8*x)]

Maple [C] Result contains higher order function than in optimal. Order 9 vs. order 3.

time = 0.84, size = 24, normalized size = 0.07

method	result	size
risch	$e^x + \left(\sum_{_R=\text{RootOf}(65536_Z^8+1)} _R \ln(e^x - 4_R) \right)$	24

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(exp(x)*tanh(4*x),x,method=_RETURNVERBOSE)
```

```
[Out] exp(x)+sum(_R*ln(exp(x)-4*_R),_R=RootOf(65536*_Z^8+1))
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(x)*tanh(4*x),x, algorithm="maxima")
```

```
[Out] e^x - 2*integrate(e^x/(e^(8*x) + 1), x)
```

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 1089 vs. $2(253) = 506$.

time = 0.38, size = 1089, normalized size = 2.98

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(x)*tanh(4*x),x, algorithm="fricas")
```

```
[Out] 1/8*(sqrt(2)*sqrt(sqrt(2) + 2) + sqrt(2)*sqrt(-sqrt(2) + 2))*arctan(-(2*sqrt(2)*e^x - sqrt(2)*sqrt(2*sqrt(2)*sqrt(sqrt(2) + 2)*e^x - 2*sqrt(2)*sqrt(-sqrt(2) + 2)*e^x + 4*e^(2*x) + 4) + sqrt(sqrt(2) + 2) - sqrt(-sqrt(2) + 2))/(sqrt(sqrt(2) + 2) + sqrt(-sqrt(2) + 2))) + 1/8*(sqrt(2)*sqrt(sqrt(2) + 2) + sqrt(2)*sqrt(-sqrt(2) + 2))*arctan(-(2*sqrt(2)*e^x - sqrt(2)*sqrt(-2*sqrt(2)*sqrt(sqrt(2) + 2)*e^x + 2*sqrt(2)*sqrt(-sqrt(2) + 2)*e^x + 4*e^(2*x) + 4) - sqrt(sqrt(2) + 2) + sqrt(-sqrt(2) + 2))/(sqrt(sqrt(2) + 2) + sqrt(-sqrt(2) + 2))) - 1/8*(sqrt(2)*sqrt(sqrt(2) + 2) - sqrt(2)*sqrt(-sqrt(2) + 2))*arctan((2*sqrt(2)*e^x - sqrt(2)*sqrt(2*sqrt(2)*sqrt(sqrt(2) + 2)*e^x + 2*sqrt(2)*sqrt(-sqrt(2) + 2)*e^x + 4*e^(2*x) + 4) + sqrt(sqrt(2) + 2) + sqrt(-sqrt(2) + 2))/(sqrt(sqrt(2) + 2) - sqrt(-sqrt(2) + 2))) - 1/8*(sqrt(2)*sqrt(sqrt(2) + 2) - sqrt(2)*sqrt(-sqrt(2) + 2))*arctan((2*sqrt(2)*e^x - sqrt(2)*sqrt(-2*sqrt(2)*sqrt(sqrt(2) + 2)*e^x - 2*sqrt(2)*sqrt(-sqrt(2) + 2)*e^x + 4*e^(2*x) + 4) - sqrt(sqrt(2) + 2) - sqrt(-sqrt(2) + 2))/(sqrt(sqrt(2) + 2) - sqrt(-sqrt(2) + 2))) - 1/32*(sqrt(2)*sqrt(sqrt(2) + 2) + sqrt(2)*sqrt(-s
```

```

qrt(2) + 2))*log(2*sqrt(2)*sqrt(sqrt(2) + 2)*e^x + 2*sqrt(2)*sqrt(-sqrt(2)
+ 2)*e^x + 4*e^(2*x) + 4) - 1/32*(sqrt(2)*sqrt(sqrt(2) + 2) - sqrt(2)*sqrt(
-sqrt(2) + 2))*log(2*sqrt(2)*sqrt(sqrt(2) + 2)*e^x - 2*sqrt(2)*sqrt(-sqrt(2)
) + 2)*e^x + 4*e^(2*x) + 4) + 1/32*(sqrt(2)*sqrt(sqrt(2) + 2) - sqrt(2)*sqr
t(-sqrt(2) + 2))*log(-2*sqrt(2)*sqrt(sqrt(2) + 2)*e^x + 2*sqrt(2)*sqrt(-sqr
t(2) + 2)*e^x + 4*e^(2*x) + 4) + 1/32*(sqrt(2)*sqrt(sqrt(2) + 2) + sqrt(2)*
sqrt(-sqrt(2) + 2))*log(-2*sqrt(2)*sqrt(sqrt(2) + 2)*e^x - 2*sqrt(2)*sqrt(-
sqrt(2) + 2)*e^x + 4*e^(2*x) + 4) + 1/4*sqrt(-sqrt(2) + 2)*arctan((2*sqrt(s
qrt(sqrt(2) + 2)*e^x + e^(2*x) + 1) - sqrt(sqrt(2) + 2) - 2*e^x)/sqrt(-sqr
t(2) + 2)) + 1/4*sqrt(-sqrt(2) + 2)*arctan((2*sqrt(-sqrt(sqrt(2) + 2)*e^x +
e^(2*x) + 1) + sqrt(sqrt(2) + 2) - 2*e^x)/sqrt(-sqrt(2) + 2)) + 1/4*sqrt(sq
rt(2) + 2)*arctan((2*sqrt(sqrt(-sqrt(2) + 2)*e^x + e^(2*x) + 1) - sqrt(-sqr
t(2) + 2) - 2*e^x)/sqrt(sqrt(2) + 2)) + 1/4*sqrt(sqrt(2) + 2)*arctan((2*sqrt
(-sqrt(-sqrt(2) + 2)*e^x + e^(2*x) + 1) + sqrt(-sqrt(2) + 2) - 2*e^x)/sqrt
(sqrt(2) + 2)) - 1/16*sqrt(sqrt(2) + 2)*log(sqrt(sqrt(2) + 2)*e^x + e^(2*x)
+ 1) + 1/16*sqrt(sqrt(2) + 2)*log(-sqrt(sqrt(2) + 2)*e^x + e^(2*x) + 1) -
1/16*sqrt(-sqrt(2) + 2)*log(sqrt(-sqrt(2) + 2)*e^x + e^(2*x) + 1) + 1/16*sq
rt(-sqrt(2) + 2)*log(-sqrt(-sqrt(2) + 2)*e^x + e^(2*x) + 1) + e^x

```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int e^x \tanh(4x) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(x)*tanh(4*x),x)

[Out] Integral(exp(x)*tanh(4*x), x)

Giac [A]

time = 0.42, size = 251, normalized size = 0.69

$$\frac{1}{4}\sqrt{-\sqrt{2}+2} \operatorname{arctan}\left(\frac{\sqrt{\sqrt{2}+2}+2e^x}{\sqrt{-\sqrt{2}+2}}\right) - \frac{1}{4}\sqrt{-\sqrt{2}+2} \operatorname{arctan}\left(\frac{\sqrt{\sqrt{2}+2}-2e^x}{\sqrt{-\sqrt{2}+2}}\right) - \frac{1}{4}\sqrt{\sqrt{2}+2} \operatorname{arctan}\left(\frac{\sqrt{-\sqrt{2}+2}+2e^x}{\sqrt{\sqrt{2}+2}}\right) - \frac{1}{4}\sqrt{\sqrt{2}+2} \operatorname{arctan}\left(\frac{\sqrt{-\sqrt{2}+2}-2e^x}{\sqrt{\sqrt{2}+2}}\right) - \frac{1}{4}\sqrt{\sqrt{2}+2} \log(\sqrt{\sqrt{2}+2}e^x+e^{2x}+1) + \frac{1}{4}\sqrt{\sqrt{2}+2} \log(-\sqrt{\sqrt{2}+2}e^x+e^{2x}+1) - \frac{1}{4}\sqrt{-\sqrt{2}+2} \log(\sqrt{-\sqrt{2}+2}e^x+e^{2x}+1) + \frac{1}{4}\sqrt{-\sqrt{2}+2} \log(-\sqrt{-\sqrt{2}+2}e^x+e^{2x}+1) + e^x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(x)*tanh(4*x),x, algorithm="giac")

```

[Out] -1/4*sqrt(-sqrt(2) + 2)*arctan((sqrt(sqrt(2) + 2) + 2*e^x)/sqrt(-sqrt(2) +
2)) - 1/4*sqrt(-sqrt(2) + 2)*arctan(-(sqrt(sqrt(2) + 2) - 2*e^x)/sqrt(-sqr
t(2) + 2)) - 1/4*sqrt(sqrt(2) + 2)*arctan((sqrt(-sqrt(2) + 2) + 2*e^x)/sqrt(
sqrt(2) + 2)) - 1/4*sqrt(sqrt(2) + 2)*arctan(-(sqrt(-sqrt(2) + 2) - 2*e^x)/
sqrt(sqrt(2) + 2)) - 1/8*sqrt(sqrt(2) + 2)*log(sqrt(sqrt(2) + 2)*e^x + e^(2
*x) + 1) + 1/8*sqrt(sqrt(2) + 2)*log(-sqrt(sqrt(2) + 2)*e^x + e^(2*x) + 1)
- 1/8*sqrt(-sqrt(2) + 2)*log(sqrt(-sqrt(2) + 2)*e^x + e^(2*x) + 1) + 1/8*sq
rt(-sqrt(2) + 2)*log(-sqrt(-sqrt(2) + 2)*e^x + e^(2*x) + 1) + e^x

```

Mupad [B]

time = 3.82, size = 457, normalized size = 1.25

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tanh(4*x)*exp(x),x)`

[Out] $\exp(x) - \log(2\exp(x) + (2^{1/2} + 2)^{1/2} + (2 - 2^{1/2})^{1/2}*1i)*((2^{1/2} + 2)^{1/2}/8 + ((2 - 2^{1/2})^{1/2}*1i)/8) + \log(2\exp(x) - (2^{1/2} + 2)^{1/2}*1i + (2 - 2^{1/2})^{1/2})*(((2^{1/2} + 2)^{1/2}*1i)/8 - (2 - 2^{1/2})^{1/2})^{1/2}/8) + \log(2\exp(x) - (2^{1/2} + 2)^{1/2} - (2 - 2^{1/2})^{1/2}*1i)*((2^{1/2} + 2)^{1/2}/8 + ((2 - 2^{1/2})^{1/2}*1i)/8) - \log(2\exp(x) + (2^{1/2} + 2)^{1/2}*1i - (2 - 2^{1/2})^{1/2})*(((2^{1/2} + 2)^{1/2}*1i)/8 - (2 - 2^{1/2})^{1/2})^{1/2}/8) + 2^{1/2}*\log(2\exp(x) - 2^{1/2})*((2^{1/2} + 2)^{1/2}/8 + ((2 - 2^{1/2})^{1/2}*1i)/8)*(4 + 4i))*((2^{1/2} + 2)^{1/2}/8 + ((2 - 2^{1/2})^{1/2}*1i)/8)*(1/2 + 1i/2) + 2^{1/2}*\log(2\exp(x) - 2^{1/2})*((2^{1/2} + 2)^{1/2}/8 + ((2 - 2^{1/2})^{1/2}*1i)/8)*(4 - 4i))*((2^{1/2} + 2)^{1/2}/8 + ((2 - 2^{1/2})^{1/2}*1i)/8)*(1/2 - 1i/2) - 2^{1/2}*\log(2\exp(x) + 2^{1/2})*((2^{1/2} + 2)^{1/2}/8 + ((2 - 2^{1/2})^{1/2}*1i)/8)*(4 - 4i))*((2^{1/2} + 2)^{1/2}/8 + ((2 - 2^{1/2})^{1/2}*1i)/8)*(1/2 - 1i/2) - 2^{1/2}*\log(2\exp(x) + 2^{1/2})*((2^{1/2} + 2)^{1/2}/8 + ((2 - 2^{1/2})^{1/2}*1i)/8)*(4 + 4i))*((2^{1/2} + 2)^{1/2}/8 + ((2 - 2^{1/2})^{1/2}*1i)/8)*(1/2 + 1i/2)$

3.224 $\int e^x \coth(4x) dx$

Optimal. Leaf size=116

$$e^x - \frac{\text{ArcTan}(e^x)}{2} + \frac{\text{ArcTan}(1 - \sqrt{2} e^x)}{2\sqrt{2}} - \frac{\text{ArcTan}(1 + \sqrt{2} e^x)}{2\sqrt{2}} - \frac{1}{2} \tanh^{-1}(e^x) + \frac{\log(1 - \sqrt{2} e^x + e^{2x})}{4\sqrt{2}} - \frac{\log(1 + \sqrt{2} e^x + e^{2x})}{4\sqrt{2}}$$

[Out] $\exp(x) - 1/2 * \arctan(\exp(x)) - 1/2 * \operatorname{arctanh}(\exp(x)) - 1/4 * \arctan(-1 + \exp(x) * 2^{(1/2)}) * 2^{(1/2)} - 1/4 * \arctan(1 + \exp(x) * 2^{(1/2)}) * 2^{(1/2)} + 1/8 * \ln(1 + \exp(2*x) - \exp(x) * 2^{(1/2)}) * 2^{(1/2)} - 1/8 * \ln(1 + \exp(2*x) + \exp(x) * 2^{(1/2)}) * 2^{(1/2)}$

Rubi [A]

time = 0.05, antiderivative size = 116, normalized size of antiderivative = 1.00, number of steps used = 15, number of rules used = 12, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 1.500$, Rules used = {2320, 396, 220, 218, 212, 209, 217, 1179, 642, 1176, 631, 210}

$$-\frac{1}{2} \text{ArcTan}(e^x) + \frac{\text{ArcTan}(1 - \sqrt{2} e^x)}{2\sqrt{2}} - \frac{\text{ArcTan}(\sqrt{2} e^x + 1)}{2\sqrt{2}} + e^x + \frac{\log(-\sqrt{2} e^x + e^{2x} + 1)}{4\sqrt{2}} - \frac{\log(\sqrt{2} e^x + e^{2x} + 1)}{4\sqrt{2}} - \frac{1}{2} \tanh^{-1}(e^x)$$

Antiderivative was successfully verified.

[In] Int[E^x*Coth[4*x], x]

[Out] $E^x - \text{ArcTan}[E^x]/2 + \text{ArcTan}[1 - \text{Sqrt}[2]*E^x]/(2*\text{Sqrt}[2]) - \text{ArcTan}[1 + \text{Sqrt}[2]*E^x]/(2*\text{Sqrt}[2]) - \text{ArcTanh}[E^x]/2 + \text{Log}[1 - \text{Sqrt}[2]*E^x + E^{(2*x)}]/(4*\text{Sqrt}[2]) - \text{Log}[1 + \text{Sqrt}[2]*E^x + E^{(2*x)}]/(4*\text{Sqrt}[2])$

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 217

```
Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]
], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*r), Int[(r - s*x^2)/(a + b*x^4),
x], x] + Dist[1/(2*r), Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b
}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &&
AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 218

```
Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2
]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x]
+ Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b
, 0]
```

Rule 220

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := With[{r = Numerator[Rt[-a/b
, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^(n/2)),
x], x] + Dist[r/(2*a), Int[1/(r + s*x^(n/2)), x], x]] /; FreeQ[{a, b}, x] &
& IGtQ[n/4, 1] && !GtQ[a/b, 0]
```

Rule 396

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Si
mp[d*x*((a + b*x^n)^(p + 1)/(b*(n*(p + 1) + 1))), x] - Dist[(a*d - b*c*(n*(
p + 1) + 1))/(b*(n*(p + 1) + 1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b,
c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]
```

Rule 631

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)
], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c]) /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```


Rule 1179

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 2320

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_.) + (b_.)*x))* (F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rubi steps

$$\begin{aligned}
\int e^x \coth(4x) dx &= \text{Subst}\left(\int \frac{-1-x^8}{1-x^8} dx, x, e^x\right) \\
&= e^x - 2\text{Subst}\left(\int \frac{1}{1-x^8} dx, x, e^x\right) \\
&= e^x - \text{Subst}\left(\int \frac{1}{1-x^4} dx, x, e^x\right) - \text{Subst}\left(\int \frac{1}{1+x^4} dx, x, e^x\right) \\
&= e^x - \frac{1}{2}\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, e^x\right) - \frac{1}{2}\text{Subst}\left(\int \frac{1}{1+x^2} dx, x, e^x\right) - \frac{1}{2}\text{Subst}\left(\int \frac{1-x}{1+x} dx, x, e^x\right) \\
&= e^x - \frac{1}{2}\tan^{-1}(e^x) - \frac{1}{2}\tanh^{-1}(e^x) - \frac{1}{4}\text{Subst}\left(\int \frac{1}{1-\sqrt{2}x+x^2} dx, x, e^x\right) - \frac{1}{4}\text{Subst}\left(\int \frac{1}{1+\sqrt{2}x+x^2} dx, x, e^x\right) \\
&= e^x - \frac{1}{2}\tan^{-1}(e^x) - \frac{1}{2}\tanh^{-1}(e^x) + \frac{\log(1-\sqrt{2}e^x+e^{2x})}{4\sqrt{2}} - \frac{\log(1+\sqrt{2}e^x+e^{2x})}{4\sqrt{2}} \\
&= e^x - \frac{1}{2}\tan^{-1}(e^x) + \frac{\tan^{-1}(1-\sqrt{2}e^x)}{2\sqrt{2}} - \frac{\tan^{-1}(1+\sqrt{2}e^x)}{2\sqrt{2}} - \frac{1}{2}\tanh^{-1}(e^x) + \frac{\log(1-\sqrt{2}e^x+e^{2x})}{4\sqrt{2}} - \frac{\log(1+\sqrt{2}e^x+e^{2x})}{4\sqrt{2}}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 22, normalized size = 0.19

$$e^x - 2e^x {}_2F_1\left(\frac{1}{8}, 1; \frac{9}{8}; e^{8x}\right)$$

Antiderivative was successfully verified.

[In] Integrate[E^x*Coth[4*x],x]

[Out] E^x - 2*E^x*Hypergeometric2F1[1/8, 1, 9/8, E^(8*x)]

Maple [C] Result contains higher order function than in optimal. Order 9 vs. order 3.
time = 0.98, size = 56, normalized size = 0.48

method	result	size
risch	$e^x + \frac{\ln(e^x-1)}{4} + \left(\sum_{R=\text{RootOf}(256_Z^4+1)} -R \ln(e^x - 4_R) \right) + \frac{i \ln(e^x-i)}{4} - \frac{i \ln(e^x+i)}{4} - \frac{\ln(e^x+1)}{4}$	56

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(x)*coth(4*x),x,method=_RETURNVERBOSE)

[Out] exp(x)+1/4*ln(exp(x)-1)+sum(_R*ln(exp(x)-4*_R),_R=RootOf(256*_Z^4+1))+1/4*I*ln(exp(x)-I)-1/4*I*ln(exp(x)+I)-1/4*ln(exp(x)+1)

Maxima [A]

time = 0.50, size = 97, normalized size = 0.84

$$-\frac{1}{4}\sqrt{2} \arctan\left(\frac{1}{2}\sqrt{2}(\sqrt{2}+2e^x)\right) - \frac{1}{4}\sqrt{2} \arctan\left(-\frac{1}{2}\sqrt{2}(\sqrt{2}-2e^x)\right) - \frac{1}{8}\sqrt{2} \log(\sqrt{2}e^x+e^{(2x)}+1) + \frac{1}{8}\sqrt{2} \log(-\sqrt{2}e^x+e^{(2x)}+1) - \frac{1}{2} \arctan(e^x)+e^x - \frac{1}{4} \log(e^x+1) + \frac{1}{4} \log(e^x-1)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(x)*coth(4*x),x, algorithm="maxima")

[Out] -1/4*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*e^x)) - 1/4*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*e^x)) - 1/8*sqrt(2)*log(sqrt(2)*e^x + e^(2*x) + 1) + 1/8*sqrt(2)*log(-sqrt(2)*e^x + e^(2*x) + 1) - 1/2*arctan(e^x) + e^x - 1/4*log(e^x + 1) + 1/4*log(e^x - 1)

Fricas [A]

time = 0.35, size = 134, normalized size = 1.16

$$\frac{1}{2}\sqrt{2} \arctan(-\sqrt{2}e^x+\sqrt{2}\sqrt{\sqrt{2}e^x+e^{(2x)}+1}-1) + \frac{1}{2}\sqrt{2} \arctan(-\sqrt{2}e^x+\frac{1}{2}\sqrt{2}\sqrt{-4\sqrt{2}e^x+4e^{(2x)}+4}+1) - \frac{1}{8}\sqrt{2} \log(4\sqrt{2}e^x+4e^{(2x)}+4) + \frac{1}{8}\sqrt{2} \log(-4\sqrt{2}e^x+4e^{(2x)}+4) - \frac{1}{2} \arctan(e^x)+e^x - \frac{1}{4} \log(e^x+1) + \frac{1}{4} \log(e^x-1)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(x)*coth(4*x),x, algorithm="fricas")

[Out] 1/2*sqrt(2)*arctan(-sqrt(2)*e^x + sqrt(2)*sqrt(sqrt(2)*e^x + e^(2*x) + 1) - 1) + 1/2*sqrt(2)*arctan(-sqrt(2)*e^x + 1/2*sqrt(2)*sqrt(-4*sqrt(2)*e^x + 4*e^(2*x) + 4) + 1) - 1/8*sqrt(2)*log(4*sqrt(2)*e^x + 4*e^(2*x) + 4) + 1/8*sqrt(2)*log(-4*sqrt(2)*e^x + 4*e^(2*x) + 4) - 1/2*arctan(e^x) + e^x - 1/4*log(e^x + 1) + 1/4*log(e^x - 1)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int e^x \coth(4x) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(x)*coth(4*x),x)`

[Out] `Integral(exp(x)*coth(4*x), x)`

Giac [A]

time = 0.41, size = 98, normalized size = 0.84

$$-\frac{1}{4}\sqrt{2}\arctan\left(\frac{1}{2}\sqrt{2}(\sqrt{2}+2e^x)\right)-\frac{1}{4}\sqrt{2}\arctan\left(-\frac{1}{2}\sqrt{2}(\sqrt{2}-2e^x)\right)-\frac{1}{8}\sqrt{2}\log(\sqrt{2}e^x+e^{(2x)}+1)+\frac{1}{8}\sqrt{2}\log(-\sqrt{2}e^x+e^{(2x)}+1)-\frac{1}{2}\arctan(e^x)+e^x-\frac{1}{4}\log(e^x+1)+\frac{1}{4}\log(|e^x-1|)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(x)*coth(4*x),x, algorithm="giac")`

[Out] `-1/4*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*e^x)) - 1/4*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*e^x)) - 1/8*sqrt(2)*log(sqrt(2)*e^x + e^(2*x) + 1) + 1/8*sqrt(2)*log(-sqrt(2)*e^x + e^(2*x) + 1) - 1/2*arctan(e^x) + e^x - 1/4*log(e^x + 1) + 1/4*log(abs(e^x - 1))`

Mupad [B]

time = 1.33, size = 104, normalized size = 0.90

$$\frac{\ln(2-2e^x)}{4}-\frac{\ln(-2e^x-2)}{4}-\frac{\operatorname{atan}(e^x)}{2}+e^x-\frac{\sqrt{2}\operatorname{atan}\left(\frac{\sqrt{2}(2e^x-\sqrt{2})}{2}\right)}{4}+\frac{\sqrt{2}\ln\left(\left(2e^x-\sqrt{2}\right)^2+2\right)}{8}-\frac{\sqrt{2}\operatorname{atan}\left(\frac{\sqrt{2}(2e^x+\sqrt{2})}{2}\right)}{4}-\frac{\sqrt{2}\ln\left(\left(2e^x+\sqrt{2}\right)^2+2\right)}{8}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(coth(4*x)*exp(x),x)`

[Out] `log(2 - 2*exp(x))/4 - log(- 2*exp(x) - 2)/4 - atan(exp(x))/2 + exp(x) - (2^(1/2)*atan((2^(1/2)*(2*exp(x) - 2^(1/2)))/2))/4 + (2^(1/2)*log((2*exp(x) - 2^(1/2))^2 + 2))/8 - (2^(1/2)*atan((2^(1/2)*(2*exp(x) + 2^(1/2)))/2))/4 - (2^(1/2)*log((2*exp(x) + 2^(1/2))^2 + 2))/8`

3.225 $\int e^x \coth^2(4x) dx$

Optimal. Leaf size=134

$$e^x + \frac{e^x}{2(1 - e^{8x})} - \frac{\text{ArcTan}(e^x)}{8} + \frac{\text{ArcTan}(1 - \sqrt{2} e^x)}{8\sqrt{2}} - \frac{\text{ArcTan}(1 + \sqrt{2} e^x)}{8\sqrt{2}} - \frac{1}{8} \tanh^{-1}(e^x) + \frac{\log(1 - \sqrt{2} e^x)}{16\sqrt{2}}$$

[Out] exp(x)+1/2*exp(x)/(1-exp(8*x))-1/8*arctan(exp(x))-1/8*arctanh(exp(x))-1/16*arctan(-1+exp(x)*2^(1/2))*2^(1/2)-1/16*arctan(1+exp(x)*2^(1/2))*2^(1/2)+1/32*ln(1+exp(2*x)-exp(x)*2^(1/2))*2^(1/2)-1/32*ln(1+exp(2*x)+exp(x)*2^(1/2))*2^(1/2)

Rubi [A]

time = 0.07, antiderivative size = 134, normalized size of antiderivative = 1.00, number of steps used = 17, number of rules used = 13, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 1.300$, Rules used = {2320, 398, 294, 220, 218, 212, 209, 217, 1179, 642, 1176, 631, 210}

$$-\frac{1}{8}\text{ArcTan}(e^x) + \frac{\text{ArcTan}(1 - \sqrt{2} e^x)}{8\sqrt{2}} - \frac{\text{ArcTan}(\sqrt{2} e^x + 1)}{8\sqrt{2}} + e^x + \frac{e^x}{2(1 - e^{8x})} + \frac{\log(-\sqrt{2} e^x + e^{2x} + 1)}{16\sqrt{2}} - \frac{\log(\sqrt{2} e^x + e^{2x} + 1)}{16\sqrt{2}} - \frac{1}{8} \tanh^{-1}(e^x)$$

Antiderivative was successfully verified.

[In] Int[E^x*Coth[4*x]^2,x]

[Out] E^x + E^x/(2*(1 - E^(8*x))) - ArcTan[E^x]/8 + ArcTan[1 - Sqrt[2]*E^x]/(8*Sqrt[2]) - ArcTan[1 + Sqrt[2]*E^x]/(8*Sqrt[2]) - ArcTanh[E^x]/8 + Log[1 - Sqrt[2]*E^x + E^(2*x)]/(16*Sqrt[2]) - Log[1 + Sqrt[2]*E^x + E^(2*x)]/(16*Sqrt[2])

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 217

```
Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]
], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*r), Int[(r - s*x^2)/(a + b*x^4),
x], x] + Dist[1/(2*r), Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b
}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &&
AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 218

```
Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]
]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x]
+ Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b
, 0]
```

Rule 220

```
Int[((a_) + (b_.)*(x_)^(n_))(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b
, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^(n/2)),
x], x] + Dist[r/(2*a), Int[1/(r + s*x^(n/2)), x], x]] /; FreeQ[{a, b}, x] &
& IGtQ[n/4, 1] && !GtQ[a/b, 0]
```

Rule 294

```
Int[((c_.)*(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(
n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n
*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x]
/; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !I
LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 398

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:= Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a
, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q,
0] && GeQ[p, -q]
```

Rule 631

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)
], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; Fre
eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 2320

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x]
, Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; Functi
onOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ[
{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*
(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rubi steps

$$\begin{aligned}
\int e^x \coth^2(4x) dx &= \text{Subst} \left(\int \frac{(1+x^8)^2}{(1-x^8)^2} dx, x, e^x \right) \\
&= \text{Subst} \left(\int \left(1 + \frac{4x^8}{(1-x^8)^2} \right) dx, x, e^x \right) \\
&= e^x + 4 \text{Subst} \left(\int \frac{x^8}{(1-x^8)^2} dx, x, e^x \right) \\
&= e^x + \frac{e^x}{2(1-e^{8x})} - \frac{1}{2} \text{Subst} \left(\int \frac{1}{1-x^8} dx, x, e^x \right) \\
&= e^x + \frac{e^x}{2(1-e^{8x})} - \frac{1}{4} \text{Subst} \left(\int \frac{1}{1-x^4} dx, x, e^x \right) - \frac{1}{4} \text{Subst} \left(\int \frac{1}{1+x^4} dx, x, e^x \right) \\
&= e^x + \frac{e^x}{2(1-e^{8x})} - \frac{1}{8} \text{Subst} \left(\int \frac{1}{1-x^2} dx, x, e^x \right) - \frac{1}{8} \text{Subst} \left(\int \frac{1}{1+x^2} dx, x, e^x \right) - \frac{1}{8} \text{Subst} \left(\int \frac{1}{1+x^2} dx, x, e^x \right) \\
&= e^x + \frac{e^x}{2(1-e^{8x})} - \frac{1}{8} \tan^{-1}(e^x) - \frac{1}{8} \tanh^{-1}(e^x) - \frac{1}{16} \text{Subst} \left(\int \frac{1}{1-\sqrt{2}x+x^2} dx, x, e^x \right) \\
&= e^x + \frac{e^x}{2(1-e^{8x})} - \frac{1}{8} \tan^{-1}(e^x) - \frac{1}{8} \tanh^{-1}(e^x) + \frac{\log(1-\sqrt{2}e^x+e^{2x})}{16\sqrt{2}} - \frac{\log(1+\sqrt{2}e^x+e^{2x})}{16\sqrt{2}} \\
&= e^x + \frac{e^x}{2(1-e^{8x})} - \frac{1}{8} \tan^{-1}(e^x) + \frac{\tan^{-1}(1-\sqrt{2}e^x)}{8\sqrt{2}} - \frac{\tan^{-1}(1+\sqrt{2}e^x)}{8\sqrt{2}} - \frac{1}{8} \tanh^{-1}(e^x)
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 1.43, size = 113, normalized size = 0.84

$$\frac{e^{-15x}(-44217 - 80225e^{8x} - 15127e^{16x} + 9361e^{24x} + 9(4913 + 8368e^{8x} + 1486e^{16x} - 1456e^{24x} + e^{32x}) {}_2F_1\left(\frac{1}{8}, 1; \frac{9}{8}; e^{8x}\right))}{9216} + \frac{64e^{9x}(1+e^{8x})^2 {}_4F_3\left(\frac{9}{8}, 2, 2, 2; 1, 1, \frac{33}{8}; e^{8x}\right)}{3825}$$

Antiderivative was successfully verified.

[In] Integrate[E^x*Coth[4*x]^2,x]

[Out] (-44217 - 80225*E^(8*x) - 15127*E^(16*x) + 9361*E^(24*x) + 9*(4913 + 8368*E^(8*x) + 1486*E^(16*x) - 1456*E^(24*x) + E^(32*x))*Hypergeometric2F1[1/8, 1, 9/8, E^(8*x)])/(9216*E^(15*x)) + (64*E^(9*x)*(1 + E^(8*x))^2*HypergeometricPFQ[{9/8, 2, 2, 2}, {1, 1, 33/8}, E^(8*x)])/3825

Maple [C] Result contains higher order function than in optimal. Order 9 vs. order 3.

time = 1.02, size = 68, normalized size = 0.51

method	result
risch	$e^x - \frac{e^x}{2(e^{8x}-1)} - \frac{\ln(e^x+1)}{16} + \frac{i \ln(e^x-i)}{16} - \frac{i \ln(e^x+i)}{16} + \frac{\ln(e^x-1)}{16} + \left(\sum_{R=\text{RootOf}(65536_Z^4+1)} _R \ln(e^x - 16_R) \right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(x)*coth(4*x)^2,x,method=_RETURNVERBOSE)`

[Out] `exp(x)-1/2*exp(x)/(exp(8*x)-1)-1/16*ln(exp(x)+1)+1/16*I*ln(exp(x)-I)-1/16*I*ln(exp(x)+I)+1/16*ln(exp(x)-1)+sum(_R*ln(exp(x)-16*_R),_R=RootOf(65536*_Z^4+1))`

Maxima [A]

time = 0.47, size = 109, normalized size = 0.81

$$-\frac{1}{16}\sqrt{2}\arctan\left(\frac{1}{2}\sqrt{2}(\sqrt{2}+2e^x)\right) - \frac{1}{16}\sqrt{2}\arctan\left(-\frac{1}{2}\sqrt{2}(\sqrt{2}-2e^x)\right) - \frac{1}{32}\sqrt{2}\log(\sqrt{2}e^x+e^{(2x)+1}) + \frac{1}{32}\sqrt{2}\log(-\sqrt{2}e^x+e^{(2x)+1}) - \frac{e^x}{2(e^{(8x)}-1)} - \frac{1}{8}\arctan(e^x)+e^x - \frac{1}{16}\log(e^x+1) + \frac{1}{16}\log(e^x-1)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(x)*coth(4*x)^2,x, algorithm="maxima")`

[Out] `-1/16*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*e^x)) - 1/16*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*e^x)) - 1/32*sqrt(2)*log(sqrt(2)*e^x + e^(2*x) + 1) + 1/32*sqrt(2)*log(-sqrt(2)*e^x + e^(2*x) + 1) - 1/2*e^x/(e^(8*x) - 1) - 1/8*arctan(e^x) + e^x - 1/16*log(e^x + 1) + 1/16*log(e^x - 1)`

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 213 vs. 2(90) = 180.

time = 0.37, size = 213, normalized size = 1.59

$$\frac{4(\sqrt{2}e^{8x}-\sqrt{2})\arctan\left(-\sqrt{2}e^x+\sqrt{2}\sqrt{\sqrt{2}e^x+e^{2x}}-1\right)+4(\sqrt{2}e^{8x}-\sqrt{2})\arctan\left(-\sqrt{2}e^x+\frac{1}{2}\sqrt{2}\sqrt{-4\sqrt{2}e^x+4e^{2x}+4}+1\right)-4(e^{8x}-1)\arctan(e^x)-(\sqrt{2}e^{8x}-\sqrt{2})\log(4\sqrt{2}e^x+4e^{2x}+4)+(\sqrt{2}e^{8x}-\sqrt{2})\log(-4\sqrt{2}e^x+4e^{2x}+4)-2(e^{8x}-1)\log(e^x+1)+2(e^{8x}-1)\log(e^x-1)+32e^{9x}-48e^x}{32(e^{8x}-1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(x)*coth(4*x)^2,x, algorithm="fricas")`

[Out] `1/32*(4*(sqrt(2)*e^(8*x) - sqrt(2))*arctan(-sqrt(2)*e^x + sqrt(2)*sqrt(sqrt(2)*e^x + e^(2*x) + 1) - 1) + 4*(sqrt(2)*e^(8*x) - sqrt(2))*arctan(-sqrt(2)*e^x + 1/2*sqrt(2)*sqrt(-4*sqrt(2)*e^x + 4*e^(2*x) + 4) + 1) - 4*(e^(8*x) - 1)*arctan(e^x) - (sqrt(2)*e^(8*x) - sqrt(2))*log(4*sqrt(2)*e^x + 4*e^(2*x) + 4) + (sqrt(2)*e^(8*x) - sqrt(2))*log(-4*sqrt(2)*e^x + 4*e^(2*x) + 4) - 2*(e^(8*x) - 1)*log(e^x + 1) + 2*(e^(8*x) - 1)*log(e^x - 1) + 32*e^(9*x) - 48*e^x/(e^(8*x) - 1)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int e^x \coth^2(4x) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(x)*coth(4*x)**2,x)

[Out] Integral(exp(x)*coth(4*x)**2, x)

Giac [A]

time = 0.41, size = 110, normalized size = 0.82

$$-\frac{1}{16}\sqrt{2}\arctan\left(\frac{1}{2}\sqrt{2}(\sqrt{2}+2e^x)\right)-\frac{1}{16}\sqrt{2}\arctan\left(-\frac{1}{2}\sqrt{2}(\sqrt{2}-2e^x)\right)-\frac{1}{32}\sqrt{2}\log(\sqrt{2}e^x+e^{(2x)}+1)+\frac{1}{32}\sqrt{2}\log(-\sqrt{2}e^x+e^{(2x)}+1)-\frac{e^x}{2(e^{8x}-1)}-\frac{1}{8}\arctan(e^x)+e^x-\frac{1}{16}\log(e^x+1)+\frac{1}{16}\log(|e^x-1|)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(x)*coth(4*x)^2,x, algorithm="giac")

[Out] $-1/16*\sqrt{2}*\arctan(1/2*\sqrt{2}*(\sqrt{2} + 2*e^x)) - 1/16*\sqrt{2}*\arctan(-1/2*\sqrt{2}*(\sqrt{2} - 2*e^x)) - 1/32*\sqrt{2}*\log(\sqrt{2}*e^x + e^{(2*x)} + 1) + 1/32*\sqrt{2}*\log(-\sqrt{2}*e^x + e^{(2*x)} + 1) - 1/2*e^x/(e^{(8*x)} - 1) - 1/8*\arctan(e^x) + e^x - 1/16*\log(e^x + 1) + 1/16*\log(\text{abs}(e^x - 1))$

Mupad [B]

time = 1.39, size = 122, normalized size = 0.91

$$\frac{\ln\left(\frac{1}{2}-\frac{e^x}{2}\right)}{16}-\frac{\ln\left(-\frac{e^x}{2}-\frac{1}{2}\right)}{16}-\frac{\operatorname{atan}(e^x)}{8}+e^x-\frac{e^x}{2(e^{8x}-1)}-\frac{\sqrt{2}\operatorname{atan}\left(2\sqrt{2}\left(\frac{e^x}{2}-\frac{\sqrt{2}}{4}\right)\right)}{16}-\frac{\sqrt{2}\operatorname{atan}\left(2\sqrt{2}\left(\frac{e^x}{2}+\frac{\sqrt{2}}{4}\right)\right)}{16}+\frac{\sqrt{2}\ln\left(\left(\frac{e^x}{2}-\frac{\sqrt{2}}{4}\right)^2+\frac{1}{8}\right)}{32}-\frac{\sqrt{2}\ln\left(\left(\frac{e^x}{2}+\frac{\sqrt{2}}{4}\right)^2+\frac{1}{8}\right)}{32}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(coth(4*x)^2*exp(x),x)

[Out] $\log(1/2 - \exp(x)/2)/16 - \log(-\exp(x)/2 - 1/2)/16 - \operatorname{atan}(\exp(x))/8 + \exp(x) - \exp(x)/(2*(\exp(8*x) - 1)) - (2^{(1/2)}*\operatorname{atan}(2*2^{(1/2)}*(\exp(x)/2 - 2^{(1/2)}/4)))/16 - (2^{(1/2)}*\operatorname{atan}(2*2^{(1/2)}*(\exp(x)/2 + 2^{(1/2)}/4)))/16 + (2^{(1/2)}*\log((\exp(x)/2 - 2^{(1/2)}/4)^2 + 1/8))/32 - (2^{(1/2)}*\log((\exp(x)/2 + 2^{(1/2)}/4)^2 + 1/8))/32$

3.226 $\int \frac{e^x}{a - \tanh(2x)} dx$

Optimal. Leaf size=107

$$-\frac{e^x}{1-a} + \frac{\operatorname{ArcTan}\left(\frac{\sqrt[4]{1-a} e^x}{\sqrt[4]{1+a}}\right)}{(1-a)\sqrt{1+a}\sqrt[4]{1-a^2}} + \frac{\tanh^{-1}\left(\frac{\sqrt[4]{1-a} e^x}{\sqrt[4]{1+a}}\right)}{(1-a)\sqrt{1+a}\sqrt[4]{1-a^2}}$$

[Out] $-\exp(x)/(1-a) + \arctan((1-a)^{1/4} \exp(x)/(1+a)^{1/4}) / ((1-a)/(-a^2+1)^{1/4}) / (1+a)^{1/2} + \operatorname{arctanh}((1-a)^{1/4} \exp(x)/(1+a)^{1/4}) / ((1-a)/(-a^2+1)^{1/4}) / (1+a)^{1/2}$

Rubi [A]

time = 0.09, antiderivative size = 107, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$, Rules used = {2320, 396, 218, 214, 211}

$$\frac{\operatorname{ArcTan}\left(\frac{\sqrt[4]{1-a} e^x}{\sqrt[4]{a+1}}\right)}{(1-a)\sqrt{a+1}\sqrt[4]{1-a^2}} + \frac{\tanh^{-1}\left(\frac{\sqrt[4]{1-a} e^x}{\sqrt[4]{a+1}}\right)}{(1-a)\sqrt{a+1}\sqrt[4]{1-a^2}} - \frac{e^x}{1-a}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^x/(a - \operatorname{Tanh}[2*x]), x]$

[Out] $-(E^x/(1-a)) + \operatorname{ArcTan}(((1-a)^{1/4} E^x)/(1+a)^{1/4}) / ((1-a) \operatorname{Sqrt}[1+a] * (1-a^2)^{1/4}) + \operatorname{ArcTanh}(((1-a)^{1/4} E^x)/(1+a)^{1/4}) / ((1-a) \operatorname{Sqrt}[1+a] * (1-a^2)^{1/4})$

Rule 211

$\operatorname{Int}[(a_+) + (b_+)(x_+)^2]^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(\operatorname{Rt}[a/b, 2]/a) * \operatorname{ArcTan}[x/\operatorname{Rt}[a/b, 2]], x] /; \operatorname{FreeQ}\{a, b, x\} \ \&\& \ \operatorname{PosQ}[a/b]$

Rule 214

$\operatorname{Int}[(a_+) + (b_+)(x_+)^2]^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(\operatorname{Rt}[-a/b, 2]/a) * \operatorname{ArcTanh}[x/\operatorname{Rt}[-a/b, 2]], x] /; \operatorname{FreeQ}\{a, b, x\} \ \&\& \ \operatorname{NegQ}[a/b]$

Rule 218

$\operatorname{Int}[(a_+) + (b_+)(x_+)^4]^{-1}, x_Symbol] \rightarrow \operatorname{With}\{r = \operatorname{Numerator}[\operatorname{Rt}[-a/b, 2]], s = \operatorname{Denominator}[\operatorname{Rt}[-a/b, 2]]\}, \operatorname{Dist}[r/(2*a), \operatorname{Int}[1/(r - s*x^2), x], x] + \operatorname{Dist}[r/(2*a), \operatorname{Int}[1/(r + s*x^2), x], x] /; \operatorname{FreeQ}\{a, b, x\} \ \&\& \ !\operatorname{GtQ}[a/b, 0]$

Rule 396

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Si
mp[d*x*((a + b*x^n)^(p + 1)/(b*(n*(p + 1) + 1))), x] - Dist[(a*d - b*c*(n*(
p + 1) + 1))/(b*(n*(p + 1) + 1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b,
c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]
```

Rule 2320

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x]
, Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; Functi
onOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ[
{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*
(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rubi steps

$$\begin{aligned} \int \frac{e^x}{a - \tanh(2x)} dx &= \text{Subst} \left(\int \frac{1 + x^4}{1 + a - (1 - a)x^4} dx, x, e^x \right) \\ &= -\frac{e^x}{1 - a} + \frac{2 \text{Subst} \left(\int \frac{1}{1 + a + (-1 + a)x^4} dx, x, e^x \right)}{1 - a} \\ &= -\frac{e^x}{1 - a} + \frac{\text{Subst} \left(\int \frac{1}{\sqrt{1 + a} - \sqrt{1 - a} x^2} dx, x, e^x \right)}{(1 - a)\sqrt{1 + a}} + \frac{\text{Subst} \left(\int \frac{1}{\sqrt{1 + a} + \sqrt{1 - a} x^2} dx, x, e^x \right)}{(1 - a)\sqrt{1 + a}} \\ &= -\frac{e^x}{1 - a} + \frac{\tan^{-1} \left(\frac{\sqrt[4]{1 - a} e^x}{\sqrt[4]{1 + a}} \right)}{(1 - a)\sqrt{1 + a} \sqrt[4]{1 - a^2}} + \frac{\tanh^{-1} \left(\frac{\sqrt[4]{1 - a} e^x}{\sqrt[4]{1 + a}} \right)}{(1 - a)\sqrt{1 + a} \sqrt[4]{1 - a^2}} \end{aligned}$$

Mathematica [A]

time = 0.06, size = 81, normalized size = 0.76

$$\frac{-\sqrt[4]{1 - a} (1 + a)^{3/4} e^x + \text{ArcTan} \left(\frac{\sqrt[4]{1 - a} e^x}{\sqrt[4]{1 + a}} \right) + \tanh^{-1} \left(\frac{\sqrt[4]{1 - a} e^x}{\sqrt[4]{1 + a}} \right)}{(1 - a)^{5/4} (1 + a)^{3/4}}$$

Antiderivative was successfully verified.

```
[In] Integrate[E^x/(a - Tanh[2*x]), x]
```

```
[Out] (-((1 - a)^(1/4)*(1 + a)^(3/4)*E^x) + ArcTan[((1 - a)^(1/4)*E^x)/(1 + a)^(1/4)] + ArcTanh[((1 - a)^(1/4)*E^x)/(1 + a)^(1/4)])/((1 - a)^(5/4)*(1 + a)^(3/4))
```

Maple [C] Result contains higher order function than in optimal. Order 9 vs. order 3.

time = 1.27, size = 87, normalized size = 0.81

method	result	size
risch	$\frac{e^x}{-1+a} + \left(\sum_{R=\text{RootOf}(1+(16a^8-32a^7-32a^6+96a^5-96a^3+32a^2+32a-16)_Z^4)} \frac{-R \ln(e^x + (-2a^2 + 2)_R)}{\dots} \right)$	70
default	$-\frac{2}{(-1+a)(\tanh(\frac{x}{2})-1)} + \frac{\sum_{R=\text{RootOf}(a_Z^4-4_Z^3+6a_Z^2-4_Z+a)} \frac{(-R^2+2R-1) \ln(\tanh(\frac{x}{2})-R)}{-R^3 a-3R^2+3R a-1}}{-2+2a}$	87

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(exp(x)/(a-tanh(2*x)),x,method=_RETURNVERBOSE)
```

```
[Out] -2/(-1+a)/(tanh(1/2*x)-1)+1/2/(-1+a)*sum((-R^2+2*_R-1)/(R^3*a-3*_R^2+3*_R*a-1)*ln(tanh(1/2*x)-_R),_R=RootOf(_Z^4*a-4*_Z^3+6*_Z^2*a-4*_Z+a))
```

Maxima [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(x)/(a-tanh(2*x)),x, algorithm="maxima")
```

```
[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(a-1>0)', see 'assume?' for more details)Is
```

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 440 vs. 2(83) = 166.

time = 0.38, size = 440, normalized size = 4.11

$$\frac{1}{(a-1) \sqrt{a^2-1}} \left(\frac{1}{\sqrt{a^2-1}} \arctan\left(\frac{1}{\sqrt{a^2-1}}\right) + \frac{1}{\sqrt{a^2-1}} \arctan\left(\frac{1}{\sqrt{a^2-1}}\right) + \frac{1}{\sqrt{a^2-1}} \arctan\left(\frac{1}{\sqrt{a^2-1}}\right) + \frac{1}{\sqrt{a^2-1}} \arctan\left(\frac{1}{\sqrt{a^2-1}}\right) \right) + (a-1) \sqrt{a^2-1} \arctan\left(\frac{1}{\sqrt{a^2-1}}\right) + (a-1) \sqrt{a^2-1} \arctan\left(\frac{1}{\sqrt{a^2-1}}\right) + (a-1) \sqrt{a^2-1} \arctan\left(\frac{1}{\sqrt{a^2-1}}\right) + (a-1) \sqrt{a^2-1} \arctan\left(\frac{1}{\sqrt{a^2-1}}\right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(x)/(a-tanh(2*x)),x, algorithm="fricas")
```

```
[Out] -1/2*(4*(a-1)*(-1/(a^8-2*a^7-2*a^6+6*a^5-6*a^3+2*a^2+2*a-1))^(1/4)*arctan(-(a^6-2*a^5-a^4+4*a^3-a^2-2*a+1)*(-1/(a^8-2*a^7-2*a^6+6*a^5-6*a^3+2*a^2+2*a-1))^(3/4))*e^x+(a^6-2*a^5-a^4+4*a^3-a^2-2*a+1)*sqrt((a^4-2*a^2+1)*sqrt(-1/(a^8-2*a^7-2*a^6+6*a^5-6*a^3+2*a^2+2*a-1)))+e^(2*x)*(-1/(a^8-2*a^7-2*a^6+6*a^5-6*a^3+2*a^2+2*a-1))^(3/4)+(a-1)*(-1/(a^8-2*a^7-2*a^6+6*a^5-6*a^3+2*a^2+2*a-1))^(1/4)*log((a^2-1)*(-1/(a^8-2*a^7-2*a^6+6*a^5-6*a^3+2*a^2+2*a-1))^(1/4))
```

$(7 - 2a^6 + 6a^5 - 6a^3 + 2a^2 + 2a - 1)^{1/4} + e^x) - (a - 1) \cdot (-1/(a^8 - 2a^7 - 2a^6 + 6a^5 - 6a^3 + 2a^2 + 2a - 1))^{1/4} \cdot \log(- (a^2 - 1) \cdot (-1/(a^8 - 2a^7 - 2a^6 + 6a^5 - 6a^3 + 2a^2 + 2a - 1))^{1/4} + e^x) - 2e^x) / (a - 1)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^x}{a - \tanh(2x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(x)/(a-tanh(2*x)),x)

[Out] Integral(exp(x)/(a - tanh(2*x)), x)

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 328 vs. 2(83) = 166.

time = 0.39, size = 328, normalized size = 3.07

$$-\frac{(a^4 - 2a^3 + 2a - 1)^{1/4} \arctan\left(\frac{\sqrt{2}(\frac{a+1}{a-1})^{1/4} e^x}{z(\frac{a+1}{a-1})^{1/4}}\right)}{\sqrt{2}a^3 - \sqrt{2}a^2 - \sqrt{2}a + \sqrt{2}} - \frac{(a^4 - 2a^3 + 2a - 1)^{1/4} \arctan\left(-\frac{\sqrt{2}(\frac{a+1}{a-1})^{1/4} e^x}{z(\frac{a+1}{a-1})^{1/4}}\right)}{\sqrt{2}a^3 - \sqrt{2}a^2 - \sqrt{2}a + \sqrt{2}} - \frac{(a^4 - 2a^3 + 2a - 1)^{1/4} \log\left(\sqrt{2}\left(\frac{a+1}{a-1}\right)^{1/4} e^x + \sqrt{\frac{a+1}{a-1}} + e^{(2x)}\right)}{2(\sqrt{2}a^3 - \sqrt{2}a^2 - \sqrt{2}a + \sqrt{2})} + \frac{(a^4 - 2a^3 + 2a - 1)^{1/4} \log\left(-\sqrt{2}\left(\frac{a+1}{a-1}\right)^{1/4} e^x + \sqrt{\frac{a+1}{a-1}} + e^{(2x)}\right)}{2(\sqrt{2}a^3 - \sqrt{2}a^2 - \sqrt{2}a + \sqrt{2})} + \frac{e^x}{a-1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(x)/(a-tanh(2*x)),x, algorithm="giac")

[Out] $-(a^4 - 2a^3 + 2a - 1)^{1/4} \cdot \arctan(1/2 \cdot \sqrt{2} \cdot ((a + 1)/(a - 1))^{1/4}) \cdot ((a + 1)/(a - 1))^{1/4} / (\sqrt{2} \cdot a^3 - \sqrt{2} \cdot a^2 - \sqrt{2} \cdot a + \sqrt{2}) - (a^4 - 2a^3 + 2a - 1)^{1/4} \cdot \arctan(-1/2 \cdot \sqrt{2} \cdot ((a + 1)/(a - 1))^{1/4}) \cdot ((a + 1)/(a - 1))^{1/4} / (\sqrt{2} \cdot a^3 - \sqrt{2} \cdot a^2 - \sqrt{2} \cdot a + \sqrt{2}) - 1/2 \cdot (a^4 - 2a^3 + 2a - 1)^{1/4} \cdot \log(\sqrt{2} \cdot ((a + 1)/(a - 1))^{1/4} \cdot e^x + \sqrt{(a + 1)/(a - 1)} + e^{(2x)}) / (\sqrt{2} \cdot a^3 - \sqrt{2} \cdot a^2 - \sqrt{2} \cdot a + \sqrt{2}) + 1/2 \cdot (a^4 - 2a^3 + 2a - 1)^{1/4} \cdot \log(-\sqrt{2} \cdot ((a + 1)/(a - 1))^{1/4} \cdot e^x + \sqrt{(a + 1)/(a - 1)} + e^{(2x)}) / (\sqrt{2} \cdot a^3 - \sqrt{2} \cdot a^2 - \sqrt{2} \cdot a + \sqrt{2}) + e^x / (a - 1)$

Mupad [B]

time = 2.45, size = 163, normalized size = 1.52

$$\frac{\ln(8a(-a-1)^{3/4} + 8e^x(a-1)^{5/4} - 8(-a-1)^{1/4}) - \ln(8e^x(a-1)^{5/4} - 8a(-a-1)^{1/4} + 8(-a-1)^{1/4}) + 2e^x(a-1)^{1/4}(-a-1)^{3/4} - \ln(8e^x(a-1)^{5/4} - a(-a-1)^{1/4}8i + (-a-1)^{1/4}8i) + \ln(a(-a-1)^{1/4}8i + 8e^x(a-1)^{5/4} - (-a-1)^{1/4}8i) + \ln(a(-a-1)^{1/4}8i + 8e^x(a-1)^{5/4} - (-a-1)^{1/4}8i) + \ln(a(-a-1)^{1/4}8i + 8e^x(a-1)^{5/4} - (-a-1)^{1/4}8i) + \ln(a(-a-1)^{1/4}8i + 8e^x(a-1)^{5/4} - (-a-1)^{1/4}8i)}{2(a-1)^{5/4}(-a-1)^{3/4}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(x)/(a - tanh(2*x)),x)

[Out] $(\log(8a \cdot (-a - 1)^{1/4} + 8 \cdot \exp(x) \cdot (a - 1)^{5/4} - 8 \cdot (-a - 1)^{1/4}) - \log(8 \cdot \exp(x) \cdot (a - 1)^{5/4} - 8a \cdot (-a - 1)^{1/4} + 8 \cdot (-a - 1)^{1/4})) - \log(8 \cdot \exp(x) \cdot (a - 1)^{5/4} - a \cdot (-a - 1)^{1/4} \cdot 8i + (-a - 1)^{1/4} \cdot 8i) + \log(a \cdot (-a - 1)^{1/4} \cdot 8i + 8 \cdot \exp(x) \cdot (a - 1)^{5/4} - (-a - 1)^{1/4} \cdot 8i) + 2 \cdot \exp(x) \cdot (a - 1)^{1/4} \cdot (-a - 1)^{3/4} / (2 \cdot (a - 1)^{5/4} \cdot (-a - 1)^{3/4})$

3.227 $\int \frac{e^x}{(a - \tanh(2x))^2} dx$

Optimal. Leaf size=152

$$\frac{e^x}{(1-a)^2} + \frac{e^x}{(1-a)^2(1+a)(1+a+(-1+a)e^{4x})} - \frac{(1+4a)\text{ArcTan}\left(\frac{\sqrt[4]{1-a}e^x}{\sqrt[4]{1+a}}\right)}{2(1-a)^2(1+a)^{3/2}\sqrt[4]{1-a^2}} - \frac{(1+4a)\tanh^{-1}\left(\frac{\sqrt[4]{1-a}}{\sqrt[4]{1+a}}\right)}{2(1-a)^2(1+a)^{3/2}\sqrt[4]{1-a^2}}$$

[Out] exp(x)/(1-a)^2+exp(x)/(1-a)^2/(1+a)/(1+a+(-1+a)*exp(4*x))-1/2*(1+4*a)*arctan((1-a)^(1/4)*exp(x)/(1+a)^(1/4))/(1-a)^2/(1+a)^(3/2)/(-a^2+1)^(1/4)-1/2*(1+4*a)*arctanh((1-a)^(1/4)*exp(x)/(1+a)^(1/4))/(1-a)^2/(1+a)^(3/2)/(-a^2+1)^(1/4)

Rubi [A]

time = 0.13, antiderivative size = 152, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {2320, 398, 393, 218, 214, 211}

$$-\frac{(4a+1)\text{ArcTan}\left(\frac{\sqrt[4]{1-a}e^x}{\sqrt[4]{a+1}}\right)}{2(1-a)^2(a+1)^{3/2}\sqrt[4]{1-a^2}} - \frac{(4a+1)\tanh^{-1}\left(\frac{\sqrt[4]{1-a}e^x}{\sqrt[4]{a+1}}\right)}{2(1-a)^2(a+1)^{3/2}\sqrt[4]{1-a^2}} + \frac{e^x}{(1-a)^2} + \frac{e^x}{(1-a)^2(a+1)((a-1)e^{4x}+a+1)}$$

Antiderivative was successfully verified.

[In] Int[E^x/(a - Tanh[2*x])^2, x]

[Out] E^x/(1 - a)^2 + E^x/((1 - a)^2*(1 + a)*(1 + a + (-1 + a)*E^(4*x))) - ((1 + 4*a)*ArcTan[(((1 - a)^(1/4)*E^x)/(1 + a)^(1/4))]/(2*(1 - a)^2*(1 + a)^(3/2)*(1 - a^2)^(1/4)) - ((1 + 4*a)*ArcTanh[(((1 - a)^(1/4)*E^x)/(1 + a)^(1/4))]/(2*(1 - a)^2*(1 + a)^(3/2)*(1 - a^2)^(1/4))

Rule 211

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 214

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 218

Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 393

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Si
mp[(-(b*c - a*d))*x*((a + b*x^n)^(p + 1)/(a*b*n*(p + 1))), x] - Dist[(a*d -
b*c*(n*(p + 1) + 1))/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; F
reeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && (LtQ[p, -1] || ILtQ[1/n
+ p, 0])
```

Rule 398

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:= Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a
, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q,
0] && GeQ[p, -q]
```

Rule 2320

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x]
, Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; Functi
onOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ[
{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*(a_.) + (b_.)*x)*
(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^x}{(a - \tanh(2x))^2} dx &= \text{Subst} \left(\int \frac{(1+x^4)^2}{(1+a-(1-a)x^4)^2} dx, x, e^x \right) \\
&= \text{Subst} \left(\int \left(\frac{1}{(-1+a)^2} - \frac{4(a-(1-a)x^4)}{(-1+a)^2(1+a+(-1+a)x^4)^2} \right) dx, x, e^x \right) \\
&= \frac{e^x}{(1-a)^2} - \frac{4 \text{Subst} \left(\int \frac{a-(1-a)x^4}{(1+a+(-1+a)x^4)^2} dx, x, e^x \right)}{(1-a)^2} \\
&= \frac{e^x}{(1-a)^2} + \frac{e^x}{(1-a)^2(1+a)(1+a-(1-a)e^{4x})} - \frac{(1+4a) \text{Subst} \left(\int \frac{1}{1+a+(-1+a)x^4} dx, x, e^x \right)}{(1-a)^2(1+a)} \\
&= \frac{e^x}{(1-a)^2} + \frac{e^x}{(1-a)^2(1+a)(1+a-(1-a)e^{4x})} - \frac{(1+4a) \text{Subst} \left(\int \frac{1}{\sqrt{1+a}-\sqrt{1-a}} dx, x, e^x \right)}{2(1-a)^2(1+a)} \\
&= \frac{e^x}{(1-a)^2} + \frac{e^x}{(1-a)^2(1+a)(1+a-(1-a)e^{4x})} - \frac{(1+4a) \tan^{-1} \left(\frac{\sqrt[4]{1-a} e^x}{\sqrt[4]{1+a}} \right)}{2(1-a)^2(1+a)^{3/2} \sqrt[4]{1-a^2}}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

time = 0.12, size = 107, normalized size = 0.70

$$\frac{\frac{4(-1+a)e^x(2+2a-e^{4x}+a^2(1+e^{4x}))}{1+a-e^{4x}+ae^{4x}} + (1+4a)\text{RootSum}\left[1+a-\#1^4+a\#1^4\&, \frac{x-\log(e^x-\#1)}{\#1^3}\&\right]}{4(-1+a)^3(1+a)}$$

Antiderivative was successfully verified.

[In] Integrate[E^x/(a - Tanh[2*x])^2, x]

[Out] ((4*(-1 + a)*E^x*(2 + 2*a - E^(4*x)) + a^2*(1 + E^(4*x)))/(1 + a - E^(4*x) + a*E^(4*x)) + (1 + 4*a)*RootSum[1 + a - #1^4 + a*#1^4 & , (x - Log[E^x - #1])/#1^3 &])/(4*(-1 + a)^3*(1 + a))

Maple [C] Result contains higher order function than in optimal. Order 9 vs. order 3.

time = 1.37, size = 191, normalized size = 1.26

method	result
default	$2 \left(\frac{\frac{(a-2)\left(\tanh^3\left(\frac{x}{2}\right)\right) - 3\left(\tanh^2\left(\frac{x}{2}\right)\right) + (a+2)\tanh\left(\frac{x}{2}\right) - \frac{1}{2(1+a)}}{2a(1+a)} + \frac{(1+4a)\left(\frac{\sum_{R=\text{RootOf}(aZ^4-4Z^3+6aZ^2-4Z+a)}\left(\frac{-R^2-2R}{-R^3-a}\right)}{8+8a}\right)}{a\left(\tanh^4\left(\frac{x}{2}\right)\right)+6a\left(\tanh^2\left(\frac{x}{2}\right)\right)-4\left(\tanh^3\left(\frac{x}{2}\right)\right)+a-4\tanh\left(\frac{x}{2}\right)} \right)}{(-1+a)^2}$
risch	$\frac{e^x}{a^2-2a+1} + \frac{e^x}{(1+a)(a^2-2a+1)(ae^{4x}-e^{4x}+a+1)} + \left(\frac{\sum_{R=\text{RootOf}((256a^{16}-512a^{15}-1536a^{14}+3584a^{13}+3584a^{12}-10752a^{11}-3584a^{10}+1024a^9-128a^8-16a^7-1)a^4-4a^3+6a^2-4a+1)}\left(\frac{-R^2-2R}{-R^3-a}\right)}{8+8a} \right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(x)/(a-tanh(2*x))^2,x,method=_RETURNVERBOSE)

[Out] -2/(-1+a)^2*((-1/2*(a-2)/a/(1+a)*tanh(1/2*x)^3-3/2/(1+a)*tanh(1/2*x)^2+1/2*(a+2)/a/(1+a)*tanh(1/2*x)-1/2/(1+a))/(a*tanh(1/2*x)^4+6*a*tanh(1/2*x)^2-4*tanh(1/2*x)^3+a-4*tanh(1/2*x))+1/8*(1+4*a)/(1+a)*sum((R^2-2*R+1)/(R^3*a-3*R^2+3*R*a-1)*ln(tanh(1/2*x)-R),R=RootOf(Z^4*a-4*Z^3+6*Z^2*a-4*Z+a))-2/(-1+a)^2/(tanh(1/2*x)-1)

Maxima [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(x)/(a-tanh(2*x))^2,x, algorithm="maxima")

[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(a>0)', see 'assume?' for more details)Is

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 1163 vs. 2(119) = 238.
time = 0.40, size = 1163, normalized size = 7.65

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(x)/(a-tanh(2*x))^2,x, algorithm="fricas")`

[Out]
$$\begin{aligned} & -\frac{1}{4} \cdot (4(a^4 - 2a^2 + (a^4 - 2a^3 + 2a - 1)e^{4x} + 1) \cdot (-256a^4 + 256a^3 + 96a^2 + 16a + 1) / (a^{16} - 2a^{15} - 6a^{14} + 14a^{13} + 14a^{12} - 42a^{11} - 14a^{10} + 70a^9 - 70a^7 + 14a^6 + 42a^5 - 14a^4 - 14a^3 + 6a^2 + 2a - 1))^{1/4} \cdot \arctan(-((4a^{13} - 7a^{12} - 18a^{11} + 36a^{10} + 30a^9 - 75a^8 - 20a^7 + 80a^6 - 45a^4 + 6a^3 + 12a^2 - 2a - 1) \cdot (-256a^4 + 256a^3 + 96a^2 + 16a + 1) / (a^{16} - 2a^{15} - 6a^{14} + 14a^{13} + 14a^{12} - 42a^{11} - 14a^{10} + 70a^9 - 70a^7 + 14a^6 + 42a^5 - 14a^4 - 14a^3 + 6a^2 + 2a - 1)))^{3/4} \cdot e^x - (a^{12} - 2a^{11} - 4a^{10} + 10a^9 + 5a^8 - 20a^7 + 20a^5 - 5a^4 - 10a^3 + 4a^2 + 2a - 1) \cdot \sqrt{(16a^2 + 8a + 1)} \cdot e^{2x} + (a^8 - 4a^6 + 6a^4 - 4a^2 + 1) \cdot \sqrt{-(256a^4 + 256a^3 + 96a^2 + 16a + 1)} / (a^{16} - 2a^{15} - 6a^{14} + 14a^{13} + 14a^{12} - 42a^{11} - 14a^{10} + 70a^9 - 70a^7 + 14a^6 + 42a^5 - 14a^4 - 14a^3 + 6a^2 + 2a - 1)) \cdot (-256a^4 + 256a^3 + 96a^2 + 16a + 1) / (a^{16} - 2a^{15} - 6a^{14} + 14a^{13} + 14a^{12} - 42a^{11} - 14a^{10} + 70a^9 - 70a^7 + 14a^6 + 42a^5 - 14a^4 - 14a^3 + 6a^2 + 2a - 1))^{3/4} / (256a^4 + 256a^3 + 96a^2 + 16a + 1) + (a^4 - 2a^2 + (a^4 - 2a^3 + 2a - 1)e^{4x} + 1) \cdot (-256a^4 + 256a^3 + 96a^2 + 16a + 1) / (a^{16} - 2a^{15} - 6a^{14} + 14a^{13} + 14a^{12} - 42a^{11} - 14a^{10} + 70a^9 - 70a^7 + 14a^6 + 42a^5 - 14a^4 - 14a^3 + 6a^2 + 2a - 1))^{1/4} \cdot \log((4a + 1)e^x + (a^4 - 2a^2 + 1) \cdot (-256a^4 + 256a^3 + 96a^2 + 16a + 1) / (a^{16} - 2a^{15} - 6a^{14} + 14a^{13} + 14a^{12} - 42a^{11} - 14a^{10} + 70a^9 - 70a^7 + 14a^6 + 42a^5 - 14a^4 - 14a^3 + 6a^2 + 2a - 1)))^{1/4} - (a^4 - 2a^2 + (a^4 - 2a^3 + 2a - 1)e^{4x} + 1) \cdot (-256a^4 + 256a^3 + 96a^2 + 16a + 1) / (a^{16} - 2a^{15} - 6a^{14} + 14a^{13} + 14a^{12} - 42a^{11} - 14a^{10} + 70a^9 - 70a^7 + 14a^6 + 42a^5 - 14a^4 - 14a^3 + 6a^2 + 2a - 1))^{1/4} \cdot \log((4a + 1)e^x - (a^4 - 2a^2 + 1) \cdot (-256a^4 + 256a^3 + 96a^2 + 16a + 1) / (a^{16} - 2a^{15} - 6a^{14} + 14a^{13} + 14a^{12} - 42a^{11} - 14a^{10} + 70a^9 - 70a^7 + 14a^6 + 42a^5 - 14a^4 - 14a^3 + 6a^2 + 2a - 1)))^{1/4} - 4(a^2 - 1)e^{5x} - 4(a^2 + 2a + 2)e^x / (a^4 - 2a^2 + (a^4 - 2a^3 + 2a - 1)e^{4x} + 1) \end{aligned}$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^x}{(a - \tanh(2x))^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(x)/(a-tanh(2*x))**2,x)

[Out] Integral(exp(x)/(a - tanh(2*x))**2, x)

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 456 vs. 2(119) = 238.

time = 0.39, size = 456, normalized size = 3.00

$$\frac{(a^2 - 2a^2 + 2a - 1)^2(4a + 1) \operatorname{arctan}\left(\frac{\sqrt{2}(\frac{a+1}{a-1})^{1/2} e^x}{1}\right)}{2(\sqrt{2}a - \sqrt{2}a^2 - 2\sqrt{2}a^3 + 2\sqrt{2}a^4 + \sqrt{2}a - \sqrt{2})} - \frac{(a^2 - 2a^2 + 2a - 1)^2(4a + 1) \operatorname{arctan}\left(\frac{\sqrt{2}(\frac{a+1}{a-1})^{1/2} e^x}{1}\right)}{2(\sqrt{2}a - \sqrt{2}a^2 - 2\sqrt{2}a^3 + 2\sqrt{2}a^4 + \sqrt{2}a - \sqrt{2})} - \frac{(a^2 - 2a^2 + 2a - 1)^2(4a + 1) \log\left(\sqrt{2}\left(\frac{a+1}{a-1}\right)^{1/2} e^x + \sqrt{\frac{a+1}{a-1}} + e^{2x}\right)}{4(\sqrt{2}a - \sqrt{2}a^2 - 2\sqrt{2}a^3 + 2\sqrt{2}a^4 + \sqrt{2}a - \sqrt{2})} + \frac{(a^2 - 2a^2 + 2a - 1)^2(4a + 1) \log\left(-\sqrt{2}\left(\frac{a+1}{a-1}\right)^{1/2} e^x + \sqrt{\frac{a+1}{a-1}} + e^{2x}\right)}{4(\sqrt{2}a - \sqrt{2}a^2 - 2\sqrt{2}a^3 + 2\sqrt{2}a^4 + \sqrt{2}a - \sqrt{2})} + \frac{e^x}{a^2 - 2a + 1} + \frac{e^x}{(a^2 - a + 1)(e^{4x} + a - e^{4x} + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(x)/(a-tanh(2*x))^2,x, algorithm="giac")

[Out] $-1/2*(a^4 - 2*a^3 + 2*a - 1)^{(1/4)}*(4*a + 1)*\arctan(1/2*\sqrt{2}*(\sqrt{2}*(a + 1)/(a - 1))^{(1/4)} + 2*e^x)/((a + 1)/(a - 1))^{(1/4)}/(\sqrt{2}*a^5 - \sqrt{2}*(2)*a^4 - 2*\sqrt{2}*a^3 + 2*\sqrt{2}*a^2 + \sqrt{2}*a - \sqrt{2}) - 1/2*(a^4 - 2*a^3 + 2*a - 1)^{(1/4)}*(4*a + 1)*\arctan(-1/2*\sqrt{2}*(\sqrt{2}*(a + 1)/(a - 1))^{(1/4)} - 2*e^x)/((a + 1)/(a - 1))^{(1/4)}/(\sqrt{2}*a^5 - \sqrt{2}*a^4 - 2*\sqrt{2}*a^3 + 2*\sqrt{2}*a^2 + \sqrt{2}*a - \sqrt{2}) - 1/4*(a^4 - 2*a^3 + 2*a - 1)^{(1/4)}*(4*a + 1)*\log(\sqrt{2}*(a + 1)/(a - 1))^{(1/4)}*e^x + \sqrt{(a + 1)/(a - 1)} + e^{(2*x)})/(\sqrt{2}*a^5 - \sqrt{2}*a^4 - 2*\sqrt{2}*a^3 + 2*\sqrt{2}*a^2 + \sqrt{2}*a - \sqrt{2}) + 1/4*(a^4 - 2*a^3 + 2*a - 1)^{(1/4)}*(4*a + 1)*\log(-\sqrt{2}*(a + 1)/(a - 1))^{(1/4)}*e^x + \sqrt{(a + 1)/(a - 1)} + e^{(2*x)})/(\sqrt{2}*a^5 - \sqrt{2}*a^4 - 2*\sqrt{2}*a^3 + 2*\sqrt{2}*a^2 + \sqrt{2}*a - \sqrt{2}) + e^x/(a^2 - 2*a + 1) + e^x/((a^3 - a^2 - a + 1)*(a*e^{(4*x)} + a - e^{(4*x)} + 1))$

Mupad [B]

time = 23.35, size = 280, normalized size = 1.84

$$\frac{e^x}{(a-1)^2} + \frac{\ln\left(\frac{4a+1}{(a-1)^{3/4}(-a-1)^{3/4}} + \frac{e^x(4a+1)}{a^2-2a+2a-1}\right)(4a+1)}{4(a-1)^{3/4}(-a-1)^{3/4}} - \frac{\ln\left(\frac{e^x(4a+1)}{a^2-2a+2a-1} - \frac{4a+1}{(a-1)^{3/4}(-a-1)^{3/4}}\right)(4a+1)}{4(a-1)^{3/4}(-a-1)^{3/4}} + \frac{e^x}{(a-1)^2(a+1)(a+e^{2x}(a-1)+1)} - \frac{\ln\left(\frac{e^x(4a+1)}{(a-1)^2(a+1)} - \frac{(4a+1)\operatorname{li}\left(\frac{4a+1}{(a-1)^2(a+1)}\right)}{(a-1)^{3/4}(-a-1)^{3/4}}\right)(4a+1)\operatorname{li}\left(\frac{4a+1}{(a-1)^2(a+1)}\right)}{4(a-1)^{3/4}(-a-1)^{3/4}} + \frac{\ln\left(\frac{e^x(4a+1)}{(a-1)^2(a+1)} + \frac{(4a+1)\operatorname{li}\left(\frac{4a+1}{(a-1)^2(a+1)}\right)}{(a-1)^{3/4}(-a-1)^{3/4}}\right)(4a+1)\operatorname{li}\left(\frac{4a+1}{(a-1)^2(a+1)}\right)}{4(a-1)^{3/4}(-a-1)^{3/4}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(x)/(a - tanh(2*x))^2,x)

[Out] $\exp(x)/(a - 1)^2 - (\log((\exp(x)*(4*a + 1))/((a - 1)^3*(a + 1))) - ((4*a + 1)*1i)/((a - 1)^{(13/4)}*(-a - 1)^{(3/4)}))*(4*a + 1)*1i/(4*(a - 1)^{(9/4)}*(-a - 1)^{(7/4)}) + (\log(((4*a + 1)*1i)/((a - 1)^{(13/4)}*(-a - 1)^{(3/4)})) + (\exp(x)*(4*a + 1))/((a - 1)^3*(a + 1)))*(4*a + 1)*1i/(4*(a - 1)^{(9/4)}*(-a - 1)^{(7/4)})$

$$\begin{aligned}
& (7/4)) + (\log((4*a + 1)/((a - 1)^{(13/4)}*(- a - 1)^{(3/4)})) + (\exp(x)*(4*a + 1) \\
&))/(2*a - 2*a^3 + a^4 - 1))*(4*a + 1))/(4*(a - 1)^{(9/4)}*(- a - 1)^{(7/4)}) - \\
& (\log((\exp(x)*(4*a + 1))/(2*a - 2*a^3 + a^4 - 1) - (4*a + 1)/((a - 1)^{(13/4)} \\
& *(- a - 1)^{(3/4)})))*(4*a + 1))/(4*(a - 1)^{(9/4)}*(- a - 1)^{(7/4)}) + \exp(x)/((\\
& a - 1)^2*(a + 1)*(a + \exp(4*x)*(a - 1) + 1))
\end{aligned}$$

3.228 $\int e^{c(a+bx)} \tanh^3(d+ex) dx$

Optimal. Leaf size=167

$$\frac{e^{c(a+bx)}}{bc} - \frac{6e^{c(a+bx)} {}_2F_1\left(1, \frac{bc}{2e}; 1 + \frac{bc}{2e}; -e^{2(d+ex)}\right)}{bc} + \frac{12e^{c(a+bx)} {}_2F_1\left(2, \frac{bc}{2e}; 1 + \frac{bc}{2e}; -e^{2(d+ex)}\right)}{bc} - \frac{8e^{c(a+bx)} {}_2F_1\left(3, \frac{bc}{2e}; 1 + \frac{bc}{2e}; -e^{2(d+ex)}\right)}{bc}$$

[Out] $\exp(c*(b*x+a))/b/c - 6*\exp(c*(b*x+a))*\text{hypergeom}([1, 1/2*b*c/e], [1+1/2*b*c/e], -\exp(2*e*x+2*d))/b/c + 12*\exp(c*(b*x+a))*\text{hypergeom}([2, 1/2*b*c/e], [1+1/2*b*c/e], -\exp(2*e*x+2*d))/b/c - 8*\exp(c*(b*x+a))*\text{hypergeom}([3, 1/2*b*c/e], [1+1/2*b*c/e], -\exp(2*e*x+2*d))/b/c$

Rubi [A]

time = 0.13, antiderivative size = 167, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 3, integrand size = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {5592, 2225, 2283}

$$-\frac{6e^{c(a+bx)} {}_2F_1\left(1, \frac{bc}{2e}, \frac{bc}{2e} + 1; -e^{2(d+ex)}\right)}{bc} + \frac{12e^{c(a+bx)} {}_2F_1\left(2, \frac{bc}{2e}, \frac{bc}{2e} + 1; -e^{2(d+ex)}\right)}{bc} - \frac{8e^{c(a+bx)} {}_2F_1\left(3, \frac{bc}{2e}, \frac{bc}{2e} + 1; -e^{2(d+ex)}\right)}{bc} + \frac{e^{c(a+bx)}}{bc}$$

Antiderivative was successfully verified.

[In] Int[E^(c*(a + b*x))*Tanh[d + e*x]^3,x]

[Out] $E^{c*(a + b*x)}/(b*c) - (6*E^{c*(a + b*x)}*\text{Hypergeometric2F1}[1, (b*c)/(2*e), 1 + (b*c)/(2*e), -E^{2*(d + e*x)}])/(b*c) + (12*E^{c*(a + b*x)}*\text{Hypergeometric2F1}[2, (b*c)/(2*e), 1 + (b*c)/(2*e), -E^{2*(d + e*x)}])/(b*c) - (8*E^{c*(a + b*x)}*\text{Hypergeometric2F1}[3, (b*c)/(2*e), 1 + (b*c)/(2*e), -E^{2*(d + e*x)}])/(b*c)$

Rule 2225

Int[((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.), x_Symbol] := Simp[(F^(c*(a + b*x)))^n/(b*c*n*Log[F]), x] /; FreeQ[{F, a, b, c, n}, x]

Rule 2283

Int[((a_) + (b_.)*(F_)^((e_.)*((c_.) + (d_.)*(x_))))^(p_)*(G_)^((h_.)*((f_.) + (g_.)*(x_))), x_Symbol] := Simp[a^p*(G^(h*(f + g*x)))/(g*h*Log[G])*Hypergeometric2F1[-p, g*h*(Log[G]/(d*e*Log[F])), g*h*(Log[G]/(d*e*Log[F])) + 1, Simplify[(-b/a)*F^(e*(c + d*x))], x] /; FreeQ[{F, G, a, b, c, d, e, f, g, h, p}, x] && (ILtQ[p, 0] || GtQ[a, 0])

Rule 5592

Int[(F_)^((c_.)*((a_.) + (b_.)*(x_)))*Tanh[(d_.) + (e_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandIntegrand[F^(c*(a + b*x))*((-1 + E^(2*(d + e*x)))^n/(1 + E^(2*(d + e*x)))^n), x], x] /; FreeQ[{F, a, b, c, d, e}, x] && IntegerQ[n]

Rubi steps

$$\begin{aligned}
\int e^{c(a+bx)} \tanh^3(d+ex) dx &= \int \left(e^{c(a+bx)} - \frac{8e^{c(a+bx)}}{(1+e^{2(d+ex)})^3} + \frac{12e^{c(a+bx)}}{(1+e^{2(d+ex)})^2} - \frac{6e^{c(a+bx)}}{1+e^{2(d+ex)}} \right) dx \\
&= -\left(6 \int \frac{e^{c(a+bx)}}{1+e^{2(d+ex)}} dx \right) - 8 \int \frac{e^{c(a+bx)}}{(1+e^{2(d+ex)})^3} dx + 12 \int \frac{e^{c(a+bx)}}{(1+e^{2(d+ex)})^2} dx \\
&= \frac{e^{c(a+bx)}}{bc} - \frac{6e^{c(a+bx)}}{bc} {}_2F_1\left(1, \frac{bc}{2e}; 1 + \frac{bc}{2e}; -e^{2(d+ex)}\right) + \frac{12e^{c(a+bx)}}{bc} {}_2F_1\left(2, \frac{bc}{2e}; 1 + \frac{bc}{2e}; -e^{2(d+ex)}\right)
\end{aligned}$$

Mathematica [A]

time = 2.93, size = 205, normalized size = 1.23

$$\frac{1}{2}e^{ac} \left(\frac{2(b^2c^2 + 2e^2)e^{2d} \left(\frac{e^{(bc+2e)x} {}_2F_1\left(1, 1 + \frac{bc}{2e}; 2 + \frac{bc}{2e}; -e^{2(d+ex)}\right)}{bc+2e} - \frac{e^{bcx} {}_2F_1\left(1, \frac{bc}{2e}; 1 + \frac{bc}{2e}; -e^{2(d+ex)}\right)}{bc} \right)}{e^2(1+e^{2d})} + \frac{e^{bcx} \operatorname{sech}^2(d+ex)}{e} - \frac{bce^{bcx} \operatorname{sech}(d) \operatorname{sech}(d+ex) \sinh(ex)}{e^2} + \frac{2e^{bcx} \tanh(d)}{bc} \right)$$

Antiderivative was successfully verified.

`[In] Integrate[E^(c*(a + b*x))*Tanh[d + e*x]^3,x]`

```
[Out] (E^(a*c)*((2*(b^2*c^2 + 2*e^2)*E^(2*d)*((E^((b*c + 2*e)*x))*Hypergeometric2F1[1, 1 + (b*c)/(2*e), 2 + (b*c)/(2*e), -E^(2*(d + e*x))])/(b*c + 2*e) - (E^(b*c*x))*Hypergeometric2F1[1, (b*c)/(2*e), 1 + (b*c)/(2*e), -E^(2*(d + e*x))])/(b*c))/(e^2*(1 + E^(2*d))) + (E^(b*c*x))*Sech[d + e*x]^2/e - (b*c*E^(b*c*x))*Sech[d]*Sech[d + e*x]*Sinh[e*x])/e^2 + (2*E^(b*c*x))*Tanh[d])/(b*c))/2
```

Maple [F]

time = 1.44, size = 0, normalized size = 0.00

$$\int e^{c(bx+a)} (\tanh^3(ex+d)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(exp(c*(b*x+a))*tanh(e*x+d)^3,x)``[Out] int(exp(c*(b*x+a))*tanh(e*x+d)^3,x)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(c*(b*x+a))*tanh(e*x+d)^3,x, algorithm="maxima")`

```
[Out] -48*(b^2*c^2*e^(a*c + 1) + 2*e^(a*c + 3))*integrate(-e^(b*c*x)/(b^3*c^3 - 1
2*b^2*c^2*e + 44*b*c*e^2 + (b^3*c^3*e^(8*d) - 12*b^2*c^2*e^(8*d + 1) + 44*b
*c*e^(8*d + 2) - 48*e^(8*d + 3))*e^(8*x*e) + 4*(b^3*c^3*e^(6*d) - 12*b^2*c^
2*e^(6*d + 1) + 44*b*c*e^(6*d + 2) - 48*e^(6*d + 3))*e^(6*x*e) + 6*(b^3*c^3
*e^(4*d) - 12*b^2*c^2*e^(4*d + 1) + 44*b*c*e^(4*d + 2) - 48*e^(4*d + 3))*e^
(4*x*e) + 4*(b^3*c^3*e^(2*d) - 12*b^2*c^2*e^(2*d + 1) + 44*b*c*e^(2*d + 2)
- 48*e^(2*d + 3))*e^(2*x*e) - 48*e^3), x) - (b^3*c^3*e^(a*c) + 36*b^2*c^2*e
^(a*c + 1) + 44*b*c*e^(a*c + 2) - (b^3*c^3*e^(a*c + 6*d) - 12*b^2*c^2*e^(a*
c + 6*d + 1) + 44*b*c*e^(a*c + 6*d + 2) - 48*e^(a*c + 6*d + 3))*e^(6*x*e) +
3*(b^3*c^3*e^(a*c + 4*d) - 8*b^2*c^2*e^(a*c + 4*d + 1) + 4*b*c*e^(a*c + 4*
d + 2) + 48*e^(a*c + 4*d + 3))*e^(4*x*e) - 3*(b^3*c^3*e^(a*c + 2*d) - 28*b*
c*e^(a*c + 2*d + 2) - 48*e^(a*c + 2*d + 3))*e^(2*x*e) + 48*e^(a*c + 3))*e^(
b*c*x)/(b^4*c^4 - 12*b^3*c^3*e + 44*b^2*c^2*e^2 - 48*b*c*e^3 + (b^4*c^4*e^(
6*d) - 12*b^3*c^3*e^(6*d + 1) + 44*b^2*c^2*e^(6*d + 2) - 48*b*c*e^(6*d + 3)
)*e^(6*x*e) + 3*(b^4*c^4*e^(4*d) - 12*b^3*c^3*e^(4*d + 1) + 44*b^2*c^2*e^(4
*d + 2) - 48*b*c*e^(4*d + 3))*e^(4*x*e) + 3*(b^4*c^4*e^(2*d) - 12*b^3*c^3*e
^(2*d + 1) + 44*b^2*c^2*e^(2*d + 2) - 48*b*c*e^(2*d + 3))*e^(2*x*e))
```

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(c*(b*x+a))*tanh(e*x+d)^3,x, algorithm="fricas")
```

```
[Out] integral(e^(b*c*x + a*c)*tanh(x*e + d)^3, x)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$e^{ac} \int e^{bcx} \tanh^3(d + ex) dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(c*(b*x+a))*tanh(e*x+d)**3,x)
```

```
[Out] exp(a*c)*Integral(exp(b*c*x)*tanh(d + e*x)**3, x)
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(c*(b*x+a))*tanh(e*x+d)^3,x, algorithm="giac")
```

[Out] integrate(e^((b*x + a)*c)*tanh(e*x + d)^3, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int e^{c(a+bx)} \tanh(d+ex)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(c*(a + b*x))*tanh(d + e*x)^3,x)

[Out] int(exp(c*(a + b*x))*tanh(d + e*x)^3, x)

3.229 $\int e^{c(a+bx)} \tanh^2(d+ex) dx$

Optimal. Leaf size=117

$$\frac{e^{c(a+bx)}}{bc} - \frac{4e^{c(a+bx)} {}_2F_1\left(1, \frac{bc}{2e}; 1 + \frac{bc}{2e}; -e^{2(d+ex)}\right)}{bc} + \frac{4e^{c(a+bx)} {}_2F_1\left(2, \frac{bc}{2e}; 1 + \frac{bc}{2e}; -e^{2(d+ex)}\right)}{bc}$$

[Out] $\exp(c*(b*x+a))/b/c - 4*\exp(c*(b*x+a))*\text{hypergeom}([1, 1/2*b*c/e], [1+1/2*b*c/e], -\exp(2*e*x+2*d))/b/c + 4*\exp(c*(b*x+a))*\text{hypergeom}([2, 1/2*b*c/e], [1+1/2*b*c/e], -\exp(2*e*x+2*d))/b/c$

Rubi [A]

time = 0.09, antiderivative size = 117, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, integrand size = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$,

Rules used = {5592, 2225, 2283}

$$-\frac{4e^{c(a+bx)} {}_2F_1\left(1, \frac{bc}{2e}; \frac{bc}{2e} + 1; -e^{2(d+ex)}\right)}{bc} + \frac{4e^{c(a+bx)} {}_2F_1\left(2, \frac{bc}{2e}; \frac{bc}{2e} + 1; -e^{2(d+ex)}\right)}{bc} + \frac{e^{c(a+bx)}}{bc}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{c*(a + b*x)}*\text{Tanh}[d + e*x]^2, x]$

[Out] $E^{c*(a + b*x)}/(b*c) - (4*E^{c*(a + b*x)}*\text{Hypergeometric2F1}[1, (b*c)/(2*e), 1 + (b*c)/(2*e), -E^{2*(d + e*x)}])/(b*c) + (4*E^{c*(a + b*x)}*\text{Hypergeometric2F1}[2, (b*c)/(2*e), 1 + (b*c)/(2*e), -E^{2*(d + e*x)}])/(b*c)$

Rule 2225

$\text{Int}[(F_)^{((c_.)*((a_.) + (b_.)*(x_)))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(F^{c*(a + b*x)})^n/(b*c*n*\text{Log}[F]), x] /; \text{FreeQ}\{F, a, b, c, n\}, x]$

Rule 2283

$\text{Int}[(a_) + (b_.)*(F_)^{((e_.)*((c_.) + (d_.)*(x_)))^{(p_)}*(G_)^{((h_.)*((f_.) + (g_.)*(x_)))}, x_Symbol] \rightarrow \text{Simp}[a^p*(G^{h*(f + g*x)})/(g*h*\text{Log}[G])*Hypergeometric2F1[-p, g*h*(\text{Log}[G]/(d*e*\text{Log}[F])), g*h*(\text{Log}[G]/(d*e*\text{Log}[F])) + 1, \text{Simplify}[(-b/a)*F^{e*(c + d*x)}]], x] /; \text{FreeQ}\{F, G, a, b, c, d, e, f, g, h, p\}, x \&\& (\text{ILtQ}[p, 0] \parallel \text{GtQ}[a, 0])$

Rule 5592

$\text{Int}[(F_)^{((c_.)*((a_.) + (b_.)*(x_)))*\text{Tanh}[(d_.) + (e_.)*(x_)]^{(n_.)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[F^{c*(a + b*x)}*((-1 + E^{2*(d + e*x)})^n/(1 + E^{2*(d + e*x)})^n), x], x] /; \text{FreeQ}\{F, a, b, c, d, e\}, x \&\& \text{IntegerQ}[n]$

Rubi steps

$$\begin{aligned}
\int e^{c(a+bx)} \tanh^2(d+ex) dx &= \int \left(e^{c(a+bx)} + \frac{4e^{c(a+bx)}}{(1+e^{2(d+ex)})^2} - \frac{4e^{c(a+bx)}}{1+e^{2(d+ex)}} \right) dx \\
&= 4 \int \frac{e^{c(a+bx)}}{(1+e^{2(d+ex)})^2} dx - 4 \int \frac{e^{c(a+bx)}}{1+e^{2(d+ex)}} dx + \int e^{c(a+bx)} dx \\
&= \frac{e^{c(a+bx)}}{bc} - \frac{4e^{c(a+bx)}}{bc} {}_2F_1\left(1, \frac{bc}{2e}; 1 + \frac{bc}{2e}; -e^{2(d+ex)}\right) + \frac{4e^{c(a+bx)}}{bc} {}_2F_1\left(2, \frac{bc}{2e}; 1 + \frac{bc}{2e}; -e^{2(d+ex)}\right)
\end{aligned}$$

Mathematica [A]

time = 2.31, size = 169, normalized size = 1.44

$$\frac{e^{c(a+bx)} (2b^2 c^2 e^{2(d+ex)} {}_2F_1(1, 1 + \frac{bc}{2e}; 2 + \frac{bc}{2e}; -e^{2(d+ex)}) - (bc + 2e) (2bce^{2d} {}_2F_1(1, \frac{bc}{2e}; 1 + \frac{bc}{2e}; -e^{2(d+ex)}) - (1 + e^{2d}) (e - bc \operatorname{sech}(d) \operatorname{sech}(d+ex) \sinh(ex))))}{bce(bc + 2e) (1 + e^{2d})}$$

Antiderivative was successfully verified.

`[In] Integrate[E^(c*(a + b*x))*Tanh[d + e*x]^2,x]`

```
[Out] (E^(c*(a + b*x))*(2*b^2*c^2*E^(2*(d + e*x))*Hypergeometric2F1[1, 1 + (b*c)/(2*e), 2 + (b*c)/(2*e), -E^(2*(d + e*x))] - (b*c + 2*e)*(2*b*c*E^(2*d)*Hypergeometric2F1[1, (b*c)/(2*e), 1 + (b*c)/(2*e), -E^(2*(d + e*x))] - (1 + E^(2*d))*(e - b*c*Sech[d]*Sech[d + e*x]*Sinh[e*x]))) / (b*c*e*(b*c + 2*e)*(1 + E^(2*d)))
```

Maple [F]

time = 1.30, size = 0, normalized size = 0.00

$$\int e^{c(bx+a)} (\tanh^2(ex+d)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(exp(c*(b*x+a))*tanh(e*x+d)^2,x)``[Out] int(exp(c*(b*x+a))*tanh(e*x+d)^2,x)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(c*(b*x+a))*tanh(e*x+d)^2,x, algorithm="maxima")`

```
[Out] -16*b*c*integrate(e^(b*c*x + a*c + 1)/(b^2*c^2 - 6*b*c*e + (b^2*c^2*e^(6*d) - 6*b*c*e^(6*d + 1) + 8*e^(6*d + 2))*e^(6*x*e) + 3*(b^2*c^2*e^(4*d) - 6*b*
```

$c e^{(4d+1)} + 8 e^{(4d+2)} e^{(4xe)} + 3(b^2 c^2 e^{(2d)} - 6 b c e^{(2d+1)} + 8 e^{(2d+2)}) e^{(2xe)} + 8 e^2$, x) + $(b^2 c^2 e^{(ac)} + 10 b c e^{(ac+1)} + (b^2 c^2 e^{(ac+4d)} - 6 b c e^{(ac+4d+1)} + 8 e^{(ac+4d+2)}) e^{(4xe)} - 2(b^2 c^2 e^{(ac+2d)} - 2 b c e^{(ac+2d+1)} - 8 e^{(ac+2d+2)}) e^{(2xe)} + 8 e^{(ac+2)}) e^{(b c x)} / (b^3 c^3 - 6 b^2 c^2 e + 8 b c e^2 + (b^3 c^3 e^{(4d)} - 6 b^2 c^2 e^{(4d+1)} + 8 b c e^{(4d+2)}) e^{(4xe)} + 2(b^3 c^3 e^{(2d)} - 6 b^2 c^2 e^{(2d+1)} + 8 b c e^{(2d+2)}) e^{(2xe)})$

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(c*(b*x+a))*tanh(e*x+d)^2,x, algorithm="fricas")

[Out] integral(e^(b*c*x + a*c)*tanh(x*e + d)^2, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$e^{ac} \int e^{bcx} \tanh^2(d + ex) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(c*(b*x+a))*tanh(e*x+d)**2,x)

[Out] exp(a*c)*Integral(exp(b*c*x)*tanh(d + e*x)**2, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(c*(b*x+a))*tanh(e*x+d)^2,x, algorithm="giac")

[Out] integrate(e^((b*x + a)*c)*tanh(e*x + d)^2, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int e^{c(a+bx)} \tanh(d + ex)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(c*(a + b*x))*tanh(d + e*x)^2,x)

[Out] int(exp(c*(a + b*x))*tanh(d + e*x)^2, x)

3.230 $\int e^{c(a+bx)} \tanh(d+ex) dx$

Optimal. Leaf size=67

$$\frac{e^{c(a+bx)}}{bc} - \frac{2e^{c(a+bx)} {}_2F_1\left(1, \frac{bc}{2e}; 1 + \frac{bc}{2e}; -e^{2(d+ex)}\right)}{bc}$$

[Out] $\exp(c*(b*x+a))/b/c - 2*\exp(c*(b*x+a))*\text{hypergeom}([1, 1/2*b*c/e], [1+1/2*b*c/e], -\exp(2*e*x+2*d))/b/c$

Rubi [A]

time = 0.05, antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {5592, 2225, 2283}

$$\frac{e^{c(a+bx)}}{bc} - \frac{2e^{c(a+bx)} {}_2F_1\left(1, \frac{bc}{2e}, \frac{bc}{2e} + 1; -e^{2(d+ex)}\right)}{bc}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{c*(a+b*x)}*\text{Tanh}[d+e*x], x]$

[Out] $E^{c*(a+b*x)}/(b*c) - (2*E^{c*(a+b*x)}*\text{Hypergeometric2F1}[1, (b*c)/(2*e), 1 + (b*c)/(2*e), -E^{2*(d+e*x)}])/(b*c)$

Rule 2225

$\text{Int}[(F_)^{((c_.)*((a_.) + (b_.)*(x_)))}^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(F^{c*(a+b*x)})^n/(b*c*n*\text{Log}[F]), x] /; \text{FreeQ}\{F, a, b, c, n\}, x]$

Rule 2283

$\text{Int}[(a_.) + (b_.)*(F_)^{((e_.)*((c_.) + (d_.)*(x_)))}^{(p_)}*(G_)^{((h_.)*((f_.) + (g_.)*(x_)))}, x_Symbol] \rightarrow \text{Simp}[a^p*(G^{h*(f+g*x)})/(g*h*\text{Log}[G])*Hypergeometric2F1[-p, g*h*(\text{Log}[G]/(d*e*\text{Log}[F])), g*h*(\text{Log}[G]/(d*e*\text{Log}[F])) + 1, \text{Simplify}[(-b/a)*F^{e*(c+d*x)}]], x] /; \text{FreeQ}\{F, G, a, b, c, d, e, f, g, h, p\}, x] \&\& (\text{ILtQ}[p, 0] || \text{GtQ}[a, 0])$

Rule 5592

$\text{Int}[(F_)^{((c_.)*((a_.) + (b_.)*(x_)))}*\text{Tanh}[(d_.) + (e_.)*(x_)]^{(n_.)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[F^{c*(a+b*x)}*((-1 + E^{2*(d+e*x)})^n/(1 + E^{2*(d+e*x)})^n), x], x] /; \text{FreeQ}\{F, a, b, c, d, e\}, x] \&\& \text{IntegerQ}[n]$

Rubi steps

$$\begin{aligned} \int e^{c(a+bx)} \tanh(d+ex) dx &= \int \left(e^{c(a+bx)} - \frac{2e^{c(a+bx)}}{1+e^{2(d+ex)}} \right) dx \\ &= - \left(2 \int \frac{e^{c(a+bx)}}{1+e^{2(d+ex)}} dx \right) + \int e^{c(a+bx)} dx \\ &= \frac{e^{c(a+bx)}}{bc} - \frac{2e^{c(a+bx)} {}_2F_1\left(1, \frac{bc}{2e}; 1 + \frac{bc}{2e}; -e^{2(d+ex)}\right)}{bc} \end{aligned}$$

Mathematica [B] Leaf count is larger than twice the leaf count of optimal. 141 vs. 2(67) = 134.

time = 1.13, size = 141, normalized size = 2.10

$$\frac{e^{c(a+bx)} (2bce^{2(d+ex)} {}_2F_1\left(1, 1 + \frac{bc}{2e}; 2 + \frac{bc}{2e}; -e^{2(d+ex)}\right) - (bc + 2e) (1 - e^{2d} + 2e^{2d} {}_2F_1\left(1, \frac{bc}{2e}; 1 + \frac{bc}{2e}; -e^{2(d+ex)}\right)))}{bc(bc + 2e)(1 + e^{2d})}$$

Antiderivative was successfully verified.

[In] Integrate[E^(c*(a + b*x))*Tanh[d + e*x], x]

[Out] (E^(c*(a + b*x))*(2*b*c*E^(2*(d + e*x))*Hypergeometric2F1[1, 1 + (b*c)/(2*e), 2 + (b*c)/(2*e), -E^(2*(d + e*x))] - (b*c + 2*e)*(1 - E^(2*d) + 2*E^(2*d))*Hypergeometric2F1[1, (b*c)/(2*e), 1 + (b*c)/(2*e), -E^(2*(d + e*x))]))/(b*c*(b*c + 2*e)*(1 + E^(2*d)))

Maple [F]

time = 1.03, size = 0, normalized size = 0.00

$$\int e^{c(bx+a)} \tanh(ex+d) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(c*(b*x+a))*tanh(e*x+d), x)

[Out] int(exp(c*(b*x+a))*tanh(e*x+d), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(c*(b*x+a))*tanh(e*x+d), x, algorithm="maxima")

[Out] -(b*c*e^(a*c) - (b*c*e^(a*c + 2*d) - 2*e^(a*c + 2*d + 1))*e^(2*x*e) + 2*e^(a*c + 1))*e^(b*c*x)/(b^2*c^2 - 2*b*c*e + (b^2*c^2*e^(2*d) - 2*b*c*e^(2*d +

1))*e^(2*x*e)) - 4*integrate(-e^(b*c*x + a*c + 1)/(b*c + (b*c*e^(4*d) - 2*e^(4*d + 1))*e^(4*x*e) + 2*(b*c*e^(2*d) - 2*e^(2*d + 1))*e^(2*x*e) - 2*e), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(c*(b*x+a))*tanh(e*x+d),x, algorithm="fricas")

[Out] integral(e^(b*c*x + a*c)*tanh(x*e + d), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$e^{ac} \int e^{bcx} \tanh(d + ex) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(c*(b*x+a))*tanh(e*x+d),x)

[Out] exp(a*c)*Integral(exp(b*c*x)*tanh(d + e*x), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(c*(b*x+a))*tanh(e*x+d),x, algorithm="giac")

[Out] integrate(e^((b*x + a)*c)*tanh(e*x + d), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int e^{c(a+bx)} \tanh(d + ex) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(c*(a + b*x))*tanh(d + e*x),x)

[Out] int(exp(c*(a + b*x))*tanh(d + e*x), x)

3.231 $\int e^{c(a+bx)} \coth(d+ex) dx$

Optimal. Leaf size=65

$$\frac{e^{c(a+bx)}}{bc} - \frac{2e^{c(a+bx)} {}_2F_1\left(1, \frac{bc}{2e}; 1 + \frac{bc}{2e}; e^{2(d+ex)}\right)}{bc}$$

[Out] exp(c*(b*x+a))/b/c-2*exp(c*(b*x+a))*hypergeom([1, 1/2*b*c/e], [1+1/2*b*c/e], exp(2*e*x+2*d))/b/c

Rubi [A]

time = 0.05, antiderivative size = 65, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {5593, 2225, 2283}

$$\frac{e^{c(a+bx)}}{bc} - \frac{2e^{c(a+bx)} {}_2F_1\left(1, \frac{bc}{2e}, \frac{bc}{2e} + 1; e^{2(d+ex)}\right)}{bc}$$

Antiderivative was successfully verified.

[In] Int[E^(c*(a + b*x))*Coth[d + e*x], x]

[Out] E^(c*(a + b*x))/(b*c) - (2*E^(c*(a + b*x))*Hypergeometric2F1[1, (b*c)/(2*e), 1 + (b*c)/(2*e), E^(2*(d + e*x))])/b*c

Rule 2225

Int[((F_)^((c_)*(a_) + (b_)*(x_)))^(n_), x_Symbol] :> Simp[(F^(c*(a + b*x)))^n/(b*c*n*Log[F]), x] /; FreeQ[{F, a, b, c, n}, x]

Rule 2283

Int[((a_) + (b_)*(F_)^((e_)*((c_) + (d_)*(x_))))^(p_)*(G_)^((h_)*((f_) + (g_)*(x_))), x_Symbol] :> Simp[a^p*(G^(h*(f + g*x)))/(g*h*Log[G])*Hypergeometric2F1[-p, g*h*(Log[G]/(d*e*Log[F])), g*h*(Log[G]/(d*e*Log[F])) + 1, Simplify[(-b/a)*F^(e*(c + d*x))], x] /; FreeQ[{F, G, a, b, c, d, e, f, g, h, p}, x] && (ILtQ[p, 0] || GtQ[a, 0])

Rule 5593

Int[Coth[(d_) + (e_)*(x_)]^(n_)*(F_)^((c_)*(a_) + (b_)*(x_)), x_Symbol] :> Int[ExpandIntegrand[F^(c*(a + b*x))*((1 + E^(2*(d + e*x)))^n/(-1 + E^(2*(d + e*x)))^n), x], x] /; FreeQ[{F, a, b, c, d, e}, x] && IntegerQ[n]

Rubi steps

$$\begin{aligned} \int e^{c(a+bx)} \coth(d+ex) dx &= \int \left(e^{c(a+bx)} + \frac{2e^{c(a+bx)}}{-1+e^{2(d+ex)}} \right) dx \\ &= 2 \int \frac{e^{c(a+bx)}}{-1+e^{2(d+ex)}} dx + \int e^{c(a+bx)} dx \\ &= \frac{e^{c(a+bx)}}{bc} - \frac{2e^{c(a+bx)} {}_2F_1\left(1, \frac{bc}{2e}; 1 + \frac{bc}{2e}; e^{2(d+ex)}\right)}{bc} \end{aligned}$$

Mathematica [B] Leaf count is larger than twice the leaf count of optimal. 134 vs. 2(65) = 130.

time = 1.43, size = 134, normalized size = 2.06

$$\frac{e^{c(a+bx)} (2bce^{2(d+ex)} {}_2F_1\left(1, 1 + \frac{bc}{2e}; 2 + \frac{bc}{2e}; e^{2(d+ex)}\right) + (bc+2e)(1+e^{2d} - 2e^{2d} {}_2F_1\left(1, \frac{bc}{2e}; 1 + \frac{bc}{2e}; e^{2(d+ex)}\right)))}{bc(bc+2e)(-1+e^{2d})}$$

Antiderivative was successfully verified.

[In] Integrate[E^(c*(a + b*x))*Coth[d + e*x], x]

[Out] (E^(c*(a + b*x))*(2*b*c*E^(2*(d + e*x))*Hypergeometric2F1[1, 1 + (b*c)/(2*e), 2 + (b*c)/(2*e), E^(2*(d + e*x))] + (b*c + 2*e)*(1 + E^(2*d) - 2*E^(2*d))*Hypergeometric2F1[1, (b*c)/(2*e), 1 + (b*c)/(2*e), E^(2*(d + e*x))]))/(b*c*(b*c + 2*e)*(-1 + E^(2*d)))

Maple [F]

time = 1.17, size = 0, normalized size = 0.00

$$\int e^{c(bx+a)} \coth(ex+d) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(c*(b*x+a))*coth(e*x+d), x)

[Out] int(exp(c*(b*x+a))*coth(e*x+d), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(c*(b*x+a))*coth(e*x+d), x, algorithm="maxima")

[Out] -(b*c*e^(a*c) + (b*c*e^(a*c + 2*d) - 2*e^(a*c + 2*d + 1))*e^(2*x*e) + 2*e^(a*c + 1))*e^(b*c*x)/(b^2*c^2 - 2*b*c*e - (b^2*c^2*e^(2*d) - 2*b*c*e^(2*d) +

1))*e^(2*x*e)) - 4*integrate(-e^(b*c*x + a*c + 1)/(b*c + (b*c*e^(4*d) - 2*e^(4*d + 1))*e^(4*x*e) - 2*(b*c*e^(2*d) - 2*e^(2*d + 1))*e^(2*x*e) - 2*e), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(c*(b*x+a))*coth(e*x+d),x, algorithm="fricas")

[Out] integral(coth(x*e + d)*e^(b*c*x + a*c), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$e^{ac} \int e^{bcx} \coth(d + ex) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(c*(b*x+a))*coth(e*x+d),x)

[Out] exp(a*c)*Integral(exp(b*c*x)*coth(d + e*x), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(c*(b*x+a))*coth(e*x+d),x, algorithm="giac")

[Out] integrate(coth(e*x + d)*e^((b*x + a)*c), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \coth(d + ex) e^{c(a+bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(coth(d + e*x)*exp(c*(a + b*x)),x)

[Out] int(coth(d + e*x)*exp(c*(a + b*x)), x)

3.232 $\int e^{c(a+bx)} \coth^2(d+ex) dx$

Optimal. Leaf size=113

$$\frac{e^{c(a+bx)}}{bc} - \frac{4e^{c(a+bx)} {}_2F_1\left(1, \frac{bc}{2e}; 1 + \frac{bc}{2e}; e^{2(d+ex)}\right)}{bc} + \frac{4e^{c(a+bx)} {}_2F_1\left(2, \frac{bc}{2e}; 1 + \frac{bc}{2e}; e^{2(d+ex)}\right)}{bc}$$

[Out] $\exp(c*(b*x+a))/b/c - 4*\exp(c*(b*x+a))*\text{hypergeom}\left([1, 1/2*b*c/e], [1+1/2*b*c/e], \exp(2*e*x+2*d)\right)/b/c + 4*\exp(c*(b*x+a))*\text{hypergeom}\left([2, 1/2*b*c/e], [1+1/2*b*c/e], \exp(2*e*x+2*d)\right)/b/c$

Rubi [A]

time = 0.09, antiderivative size = 113, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, integrand size = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$,

Rules used = {5593, 2225, 2283}

$$-\frac{4e^{c(a+bx)} {}_2F_1\left(1, \frac{bc}{2e}; \frac{bc}{2e} + 1; e^{2(d+ex)}\right)}{bc} + \frac{4e^{c(a+bx)} {}_2F_1\left(2, \frac{bc}{2e}; \frac{bc}{2e} + 1; e^{2(d+ex)}\right)}{bc} + \frac{e^{c(a+bx)}}{bc}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{c*(a + b*x)}*\text{Coth}[d + e*x]^2, x]$

[Out] $E^{c*(a + b*x)}/(b*c) - (4*E^{c*(a + b*x)}*\text{Hypergeometric2F1}[1, (b*c)/(2*e), 1 + (b*c)/(2*e), E^{2*(d + e*x)}])/(b*c) + (4*E^{c*(a + b*x)}*\text{Hypergeometric2F1}[2, (b*c)/(2*e), 1 + (b*c)/(2*e), E^{2*(d + e*x)}])/(b*c)$

Rule 2225

$\text{Int}[(F_)^{((c_.)*((a_.) + (b_.)*(x_)))}^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(F^{c*(a + b*x)})^n/(b*c*n*\text{Log}[F]), x] /; \text{FreeQ}\{F, a, b, c, n\}, x]$

Rule 2283

$\text{Int}[(a_.) + (b_.)*(F_)^{((e_.)*((c_.) + (d_.)*(x_)))}^{(p_)}*(G_)^{((h_.)*((f_.) + (g_.)*(x_)))}, x_Symbol] \rightarrow \text{Simp}[a^p*(G^{h*(f + g*x)})/(g*h*\text{Log}[G])*Hypergeometric2F1[-p, g*h*(\text{Log}[G]/(d*e*\text{Log}[F])), g*h*(\text{Log}[G]/(d*e*\text{Log}[F])) + 1, \text{Simplify}[(-b/a)*F^{e*(c + d*x)}]], x] /; \text{FreeQ}\{F, G, a, b, c, d, e, f, g, h, p\}, x] \&\& (\text{ILtQ}[p, 0] || \text{GtQ}[a, 0])$

Rule 5593

$\text{Int}[\text{Coth}[(d_.) + (e_.)*(x_)]^{(n_.)}*(F_)^{((c_.)*((a_.) + (b_.)*(x_)))}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[F^{c*(a + b*x)}*((1 + E^{2*(d + e*x)})^n/(-1 + E^{2*(d + e*x)})^n), x], x] /; \text{FreeQ}\{F, a, b, c, d, e\}, x] \&\& \text{IntegerQ}[n]$

Rubi steps

$$\begin{aligned} \int e^{c(a+bx)} \coth^2(d+ex) dx &= \int \left(e^{c(a+bx)} + \frac{4e^{c(a+bx)}}{(-1+e^{2(d+ex)})^2} + \frac{4e^{c(a+bx)}}{-1+e^{2(d+ex)}} \right) dx \\ &= 4 \int \frac{e^{c(a+bx)}}{(-1+e^{2(d+ex)})^2} dx + 4 \int \frac{e^{c(a+bx)}}{-1+e^{2(d+ex)}} dx + \int e^{c(a+bx)} dx \\ &= \frac{e^{c(a+bx)}}{bc} - \frac{4e^{c(a+bx)} {}_2F_1\left(1, \frac{bc}{2e}; 1 + \frac{bc}{2e}; e^{2(d+ex)}\right)}{bc} + \frac{4e^{c(a+bx)} {}_2F_1\left(2, \frac{bc}{2e}; 1 + \frac{bc}{2e}; e^{2(d+ex)}\right)}{bc} \end{aligned}$$

Mathematica [A]

time = 2.16, size = 145, normalized size = 1.28

$$e^{c(a+bx)} \left(\frac{1}{bc} + \frac{2e^{2d} (bce^{2ex} {}_2F_1\left(1, 1 + \frac{bc}{2e}; 2 + \frac{bc}{2e}; e^{2(d+ex)}\right) - (bc+2e) {}_2F_1\left(1, \frac{bc}{2e}; 1 + \frac{bc}{2e}; e^{2(d+ex)}\right))}{e(bc+2e)(-1+e^{2d})} + \frac{\operatorname{csch}(d)\operatorname{csch}(d+ex)\sinh(ex)}{e} \right)$$

Antiderivative was successfully verified.

`[In] Integrate[E^(c*(a + b*x))*Coth[d + e*x]^2,x]`

```
[Out] E^(c*(a + b*x))*(1/(b*c) + (2*E^(2*d)*(b*c*E^(2*e*x)*Hypergeometric2F1[1, 1 + (b*c)/(2*e), 2 + (b*c)/(2*e), E^(2*(d + e*x))] - (b*c + 2*e)*Hypergeometric2F1[1, (b*c)/(2*e), 1 + (b*c)/(2*e), E^(2*(d + e*x))]))/(e*(b*c + 2*e)*(-1 + E^(2*d))) + (Csch[d]*Csch[d + e*x]*Sinh[e*x])/e)
```

Maple [F]

time = 1.18, size = 0, normalized size = 0.00

$$\int e^{c(bx+a)} (\coth^2(ex+d)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(exp(c*(b*x+a))*coth(e*x+d)^2,x)``[Out] int(exp(c*(b*x+a))*coth(e*x+d)^2,x)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(c*(b*x+a))*coth(e*x+d)^2,x, algorithm="maxima")`

```
[Out] 16*b*c*integrate(-e^(b*c*x + a*c + 1)/(b^2*c^2 - 6*b*c*e - (b^2*c^2*e^(6*d) - 6*b*c*e^(6*d + 1) + 8*e^(6*d + 2))*e^(6*x*e) + 3*(b^2*c^2*e^(4*d) - 6*b*
```

$$c e^{(4d+1)} + 8 e^{(4d+2)}) e^{(4xe)} - 3(b^2 c^2 e^{(2d)} - 6 b c e^{(2d+1)} + 8 e^{(2d+2)}) e^{(2xe)} + 8 e^2, x) + (b^2 c^2 e^{(ac)} + 10 b c e^{(ac+1)} + (b^2 c^2 e^{(ac+4d)} - 6 b c e^{(ac+4d+1)} + 8 e^{(ac+4d+2)}) e^{(4xe)} + 2(b^2 c^2 e^{(ac+2d)} - 2 b c e^{(ac+2d+1)} - 8 e^{(ac+2d+2)}) e^{(2xe)} + 8 e^{(ac+2)}) e^{(bcx)} / (b^3 c^3 - 6 b^2 c^2 e + 8 b c e^2 + (b^3 c^3 e^{(4d)} - 6 b^2 c^2 e^{(4d+1)} + 8 b c e^{(4d+2)}) e^{(4xe)} - 2(b^3 c^3 e^{(2d)} - 6 b^2 c^2 e^{(2d+1)} + 8 b c e^{(2d+2)}) e^{(2xe)})$$

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(c*(b*x+a))*coth(e*x+d)^2,x, algorithm="fricas")

[Out] integral(coth(x*e + d)^2*e^(b*c*x + a*c), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$e^{ac} \int e^{bcx} \coth^2(d + ex) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(c*(b*x+a))*coth(e*x+d)**2,x)

[Out] exp(a*c)*Integral(exp(b*c*x)*coth(d + e*x)**2, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(c*(b*x+a))*coth(e*x+d)^2,x, algorithm="giac")

[Out] integrate(coth(e*x + d)^2*e^((b*x + a)*c), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \coth(d + ex)^2 e^{c(a+bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(coth(d + e*x)^2*exp(c*(a + b*x)),x)

[Out] int(coth(d + e*x)^2*exp(c*(a + b*x)), x)

3.233 $\int e^{c(a+bx)} \coth^3(d+ex) dx$

Optimal. Leaf size=161

$$\frac{e^{c(a+bx)}}{bc} - \frac{6e^{c(a+bx)} {}_2F_1\left(1, \frac{bc}{2e}; 1 + \frac{bc}{2e}; e^{2(d+ex)}\right)}{bc} + \frac{12e^{c(a+bx)} {}_2F_1\left(2, \frac{bc}{2e}; 1 + \frac{bc}{2e}; e^{2(d+ex)}\right)}{bc} - \frac{8e^{c(a+bx)} {}_2F_1\left(3, \frac{bc}{2e}; 1 + \frac{bc}{2e}; e^{2(d+ex)}\right)}{bc}$$

[Out] $\exp(c*(b*x+a))/b/c - 6*\exp(c*(b*x+a))*\text{hypergeom}([1, 1/2*b*c/e], [1+1/2*b*c/e], \exp(2*e*x+2*d))/b/c + 12*\exp(c*(b*x+a))*\text{hypergeom}([2, 1/2*b*c/e], [1+1/2*b*c/e], \exp(2*e*x+2*d))/b/c - 8*\exp(c*(b*x+a))*\text{hypergeom}([3, 1/2*b*c/e], [1+1/2*b*c/e], \exp(2*e*x+2*d))/b/c$

Rubi [A]

time = 0.12, antiderivative size = 161, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 3, integrand size = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {5593, 2225, 2283}

$$-\frac{6e^{c(a+bx)} {}_2F_1\left(1, \frac{bc}{2e}, \frac{bc}{2e} + 1; e^{2(d+ex)}\right)}{bc} + \frac{12e^{c(a+bx)} {}_2F_1\left(2, \frac{bc}{2e}, \frac{bc}{2e} + 1; e^{2(d+ex)}\right)}{bc} - \frac{8e^{c(a+bx)} {}_2F_1\left(3, \frac{bc}{2e}, \frac{bc}{2e} + 1; e^{2(d+ex)}\right)}{bc} + \frac{e^{c(a+bx)}}{bc}$$

Antiderivative was successfully verified.

[In] Int[E^(c*(a + b*x))*Coth[d + e*x]^3,x]

[Out] $E^{c*(a + b*x)}/(b*c) - (6*E^{c*(a + b*x)}*\text{Hypergeometric2F1}[1, (b*c)/(2*e), 1 + (b*c)/(2*e), E^{2*(d + e*x)}])/(b*c) + (12*E^{c*(a + b*x)}*\text{Hypergeometric2F1}[2, (b*c)/(2*e), 1 + (b*c)/(2*e), E^{2*(d + e*x)}])/(b*c) - (8*E^{c*(a + b*x)}*\text{Hypergeometric2F1}[3, (b*c)/(2*e), 1 + (b*c)/(2*e), E^{2*(d + e*x)}])/(b*c)$

Rule 2225

Int[((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.), x_Symbol] := Simp[(F^(c*(a + b*x)))^n/(b*c*n*Log[F]), x] /; FreeQ[{F, a, b, c, n}, x]

Rule 2283

Int[((a_) + (b_.)*(F_)^((e_.)*((c_.) + (d_.)*(x_))))^(p_)*(G_)^((h_.)*((f_.) + (g_.)*(x_))), x_Symbol] := Simp[a^p*(G^(h*(f + g*x)))/(g*h*Log[G])*Hypergeometric2F1[-p, g*h*(Log[G]/(d*e*Log[F])), g*h*(Log[G]/(d*e*Log[F])) + 1, Simplify[(-b/a)*F^(e*(c + d*x))], x] /; FreeQ[{F, G, a, b, c, d, e, f, g, h, p}, x] && (ILtQ[p, 0] || GtQ[a, 0])

Rule 5593

Int[Coth[(d_.) + (e_.)*(x_)]^(n_.)*(F_)^((c_.)*((a_.) + (b_.)*(x_))), x_Symbol] := Int[ExpandIntegrand[F^(c*(a + b*x))*((1 + E^(2*(d + e*x)))^n/(-1 + E^(2*(d + e*x)))^n), x], x] /; FreeQ[{F, a, b, c, d, e}, x] && IntegerQ[n]

Rubi steps

$$\begin{aligned}
\int e^{c(a+bx)} \coth^3(d+ex) dx &= \int \left(e^{c(a+bx)} + \frac{8e^{c(a+bx)}}{(-1+e^{2(d+ex)})^3} + \frac{12e^{c(a+bx)}}{(-1+e^{2(d+ex)})^2} + \frac{6e^{c(a+bx)}}{-1+e^{2(d+ex)}} \right) dx \\
&= 6 \int \frac{e^{c(a+bx)}}{-1+e^{2(d+ex)}} dx + 8 \int \frac{e^{c(a+bx)}}{(-1+e^{2(d+ex)})^3} dx + 12 \int \frac{e^{c(a+bx)}}{(-1+e^{2(d+ex)})^2} dx \\
&= \frac{e^{c(a+bx)}}{bc} - \frac{6e^{c(a+bx)} {}_2F_1\left(1, \frac{bc}{2e}; 1 + \frac{bc}{2e}; e^{2(d+ex)}\right)}{bc} + \frac{12e^{c(a+bx)} {}_2F_1\left(2, \frac{bc}{2e}; 1 + \frac{bc}{2e}; e^{2(d+ex)}\right)}{bc}
\end{aligned}$$

Mathematica [A]

time = 2.15, size = 185, normalized size = 1.15

$$\frac{1}{2} e^{c(a+bx)} \left(\frac{2 \coth(d)}{bc} - \frac{\operatorname{csch}^2(d+ex)}{e} + \frac{2(b^2c^2 + 2e^2) e^{2d} (bc e^{2ex} {}_2F_1(1, 1 + \frac{bc}{2e}; 2 + \frac{bc}{2e}; e^{2(d+ex)}) - (bc + 2e) {}_2F_1(1, \frac{bc}{2e}; 1 + \frac{bc}{2e}; e^{2(d+ex)}))}{bc^2(bc + 2e)(-1 + e^{2d})} + \frac{bc \operatorname{csch}(d) \operatorname{csch}(d+ex) \sinh(ex)}{e^2} \right)$$

Antiderivative was successfully verified.

`[In] Integrate[E^(c*(a + b*x))*Coth[d + e*x]^3,x]`

```
[Out] (E^(c*(a + b*x))*((2*Coth[d])/(b*c) - Csch[d + e*x]^2/e + (2*(b^2*c^2 + 2*e^2)*E^(2*d)*(b*c*E^(2*e*x)*Hypergeometric2F1[1, 1 + (b*c)/(2*e), 2 + (b*c)/(2*e), E^(2*(d + e*x))]) - (b*c + 2*e)*Hypergeometric2F1[1, (b*c)/(2*e), 1 + (b*c)/(2*e), E^(2*(d + e*x))]))/(b*c*e^(2*(b*c + 2*e)*(-1 + E^(2*d))) + (b*c*Csch[d]*Csch[d + e*x]*Sinh[e*x])/e^2))/2
```

Maple [F]

time = 1.30, size = 0, normalized size = 0.00

$$\int e^{c(bx+a)} (\coth^3(ex+d)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(exp(c*(b*x+a))*coth(e*x+d)^3,x)``[Out] int(exp(c*(b*x+a))*coth(e*x+d)^3,x)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(c*(b*x+a))*coth(e*x+d)^3,x, algorithm="maxima")`

```
[Out] -48*(b^2*c^2*e^(a*c + 1) + 2*e^(a*c + 3))*integrate(-e^(b*c*x)/(b^3*c^3 - 1
2*b^2*c^2*e + 44*b*c*e^2 + (b^3*c^3*e^(8*d) - 12*b^2*c^2*e^(8*d + 1) + 44*b
*c*e^(8*d + 2) - 48*e^(8*d + 3))*e^(8*x*e) - 4*(b^3*c^3*e^(6*d) - 12*b^2*c^
2*e^(6*d + 1) + 44*b*c*e^(6*d + 2) - 48*e^(6*d + 3))*e^(6*x*e) + 6*(b^3*c^3
*e^(4*d) - 12*b^2*c^2*e^(4*d + 1) + 44*b*c*e^(4*d + 2) - 48*e^(4*d + 3))*e^
(4*x*e) - 4*(b^3*c^3*e^(2*d) - 12*b^2*c^2*e^(2*d + 1) + 44*b*c*e^(2*d + 2)
- 48*e^(2*d + 3))*e^(2*x*e) - 48*e^3), x) - (b^3*c^3*e^(a*c) + 36*b^2*c^2*e
^(a*c + 1) + 44*b*c*e^(a*c + 2) + (b^3*c^3*e^(a*c + 6*d) - 12*b^2*c^2*e^(a*
c + 6*d + 1) + 44*b*c*e^(a*c + 6*d + 2) - 48*e^(a*c + 6*d + 3))*e^(6*x*e) +
3*(b^3*c^3*e^(a*c + 4*d) - 8*b^2*c^2*e^(a*c + 4*d + 1) + 4*b*c*e^(a*c + 4*
d + 2) + 48*e^(a*c + 4*d + 3))*e^(4*x*e) + 3*(b^3*c^3*e^(a*c + 2*d) - 28*b*
c*e^(a*c + 2*d + 2) - 48*e^(a*c + 2*d + 3))*e^(2*x*e) + 48*e^(a*c + 3))*e^(
b*c*x)/(b^4*c^4 - 12*b^3*c^3*e + 44*b^2*c^2*e^2 - 48*b*c*e^3 - (b^4*c^4*e^(
6*d) - 12*b^3*c^3*e^(6*d + 1) + 44*b^2*c^2*e^(6*d + 2) - 48*b*c*e^(6*d + 3)
)*e^(6*x*e) + 3*(b^4*c^4*e^(4*d) - 12*b^3*c^3*e^(4*d + 1) + 44*b^2*c^2*e^(4
*d + 2) - 48*b*c*e^(4*d + 3))*e^(4*x*e) - 3*(b^4*c^4*e^(2*d) - 12*b^3*c^3*e
^(2*d + 1) + 44*b^2*c^2*e^(2*d + 2) - 48*b*c*e^(2*d + 3))*e^(2*x*e))
```

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(c*(b*x+a))*coth(e*x+d)^3,x, algorithm="fricas")
```

```
[Out] integral(coth(x*e + d)^3*e^(b*c*x + a*c), x)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$e^{ac} \int e^{bcx} \coth^3(d + ex) dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(c*(b*x+a))*coth(e*x+d)**3,x)
```

```
[Out] exp(a*c)*Integral(exp(b*c*x)*coth(d + e*x)**3, x)
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(c*(b*x+a))*coth(e*x+d)^3,x, algorithm="giac")
```

[Out] integrate(coth(e*x + d)^3*e^((b*x + a)*c), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \coth(d + e x)^3 e^{c(a + b x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(coth(d + e*x)^3*exp(c*(a + b*x)),x)

[Out] int(coth(d + e*x)^3*exp(c*(a + b*x)), x)

3.234 $\int e^{c(a+bx)} \tanh^2(ac + bcx)^{5/2} dx$

Optimal. Leaf size=311

$$\frac{e^{c(a+bx)} \coth(ac + bcx) \sqrt{\tanh^2(ac + bcx)}}{bc} - \frac{4e^{c(a+bx)} \coth(ac + bcx) \sqrt{\tanh^2(ac + bcx)}}{bc(1 + e^{2c(a+bx)})^4} + \frac{26e^{c(a+bx)} \coth(ac + bcx) \sqrt{\tanh^2(ac + bcx)}}{3bc(1 + e^{2c(a+bx)})^3}$$

[Out] $\exp(c*(b*x+a))*\coth(b*c*x+a*c)*(\tanh(b*c*x+a*c)^2)^{(1/2)}/b/c-4*\exp(c*(b*x+a))*\coth(b*c*x+a*c)*(\tanh(b*c*x+a*c)^2)^{(1/2)}/b/c/(1+\exp(2*c*(b*x+a)))^4+26/3*\exp(c*(b*x+a))*\coth(b*c*x+a*c)*(\tanh(b*c*x+a*c)^2)^{(1/2)}/b/c/(1+\exp(2*c*(b*x+a)))^3-55/6*\exp(c*(b*x+a))*\coth(b*c*x+a*c)*(\tanh(b*c*x+a*c)^2)^{(1/2)}/b/c/(1+\exp(2*c*(b*x+a)))^2+25/4*\exp(c*(b*x+a))*\coth(b*c*x+a*c)*(\tanh(b*c*x+a*c)^2)^{(1/2)}/b/c/(1+\exp(2*c*(b*x+a)))-15/4*\arctan(\exp(c*(b*x+a)))*\coth(b*c*x+a*c)*(\tanh(b*c*x+a*c)^2)^{(1/2)}/b/c$

Rubi [A]

time = 0.63, antiderivative size = 311, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 7, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.280$, Rules used = {6852, 2320, 398, 1828, 1171, 393, 209}

$$\frac{15 \operatorname{ArcTan}(e^{c(a+bx)}) \sqrt{\tanh^2(ac+bcx)} \coth(ac+bcx)}{4bc} + \frac{e^{c(a+bx)} \sqrt{\tanh^2(ac+bcx)} \coth(ac+bcx)}{bc} + \frac{25e^{c(a+bx)} \sqrt{\tanh^2(ac+bcx)} \coth(ac+bcx)}{4bc(e^{2c(a+bx)}+1)} - \frac{55e^{c(a+bx)} \sqrt{\tanh^2(ac+bcx)} \coth(ac+bcx)}{6bc(e^{2c(a+bx)}+1)^2} + \frac{26e^{c(a+bx)} \sqrt{\tanh^2(ac+bcx)} \coth(ac+bcx)}{3bc(e^{2c(a+bx)}+1)^3} - \frac{4e^{c(a+bx)} \sqrt{\tanh^2(ac+bcx)} \coth(ac+bcx)}{bc(e^{2c(a+bx)}+1)^4}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{c*(a + b*x)}*(\text{Tanh}[a*c + b*c*x]^2)^{(5/2)}, x]$

[Out] $(E^{c*(a + b*x)}*\text{Coth}[a*c + b*c*x]*\text{Sqrt}[\text{Tanh}[a*c + b*c*x]^2])/(b*c) - (4*E^{c*(a + b*x)}*\text{Coth}[a*c + b*c*x]*\text{Sqrt}[\text{Tanh}[a*c + b*c*x]^2])/(b*c*(1 + E^{2*c*(a + b*x)}))^4 + (26*E^{c*(a + b*x)}*\text{Coth}[a*c + b*c*x]*\text{Sqrt}[\text{Tanh}[a*c + b*c*x]^2])/(3*b*c*(1 + E^{2*c*(a + b*x)}))^3 - (55*E^{c*(a + b*x)}*\text{Coth}[a*c + b*c*x]*\text{Sqrt}[\text{Tanh}[a*c + b*c*x]^2])/(6*b*c*(1 + E^{2*c*(a + b*x)}))^2 + (25*E^{c*(a + b*x)}*\text{Coth}[a*c + b*c*x]*\text{Sqrt}[\text{Tanh}[a*c + b*c*x]^2])/(4*b*c*(1 + E^{2*c*(a + b*x)})) - (15*\text{ArcTan}[E^{c*(a + b*x)}]*\text{Coth}[a*c + b*c*x]*\text{Sqrt}[\text{Tanh}[a*c + b*c*x]^2])/(4*b*c)$

Rule 209

$\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2]))*\text{ArcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$

Rule 393

$\text{Int}[(a_ + (b_)*(x_)^{(n_)})^{(p_)}*((c_ + (d_)*(x_)^{(n_)}), x_Symbol] \rightarrow \text{Simp}[(- (b*c - a*d))*x*((a + b*x^n)^{(p+1)})/(a*b*n*(p+1)), x] - \text{Dist}[(a*d - b*c*(n*(p+1) + 1))/(a*b*n*(p+1)), \text{Int}[(a + b*x^n)^{(p+1)}, x], x] /; \text{FreeQ}\{a, b, c, d, n, p\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ (\text{LtQ}[p, -1] \ || \ \text{ILtQ}[1/n, 0])$

+ p, 0])

Rule 398

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:> Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a
, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q,
0] && GeQ[p, -q]
```

Rule 1171

```
Int[((d_) + (e_.)*(x_)^2)^(q_)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_.),
x_Symbol] :> With[{Qx = PolynomialQuotient[(a + b*x^2 + c*x^4)^p, d + e*x^2
, x], R = Coeff[PolynomialRemainder[(a + b*x^2 + c*x^4)^p, d + e*x^2, x], x
, 0]}, Simp[(-R)*x*((d + e*x^2)^(q + 1)/(2*d*(q + 1))), x] + Dist[1/(2*d*(q
+ 1)), Int[(d + e*x^2)^(q + 1)*ExpandToSum[2*d*(q + 1)*Qx + R*(2*q + 3), x
], x], x]] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2
- b*d*e + a*e^2, 0] && IGtQ[p, 0] && LtQ[q, -1]
```

Rule 1828

```
Int[(Pq_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :> With[{Q = PolynomialQuot
ient[Pq, a + b*x^2, x], f = Coeff[PolynomialRemainder[Pq, a + b*x^2, x], x,
0], g = Coeff[PolynomialRemainder[Pq, a + b*x^2, x], x, 1]}, Simp[(a*g - b
*f*x)*((a + b*x^2)^(p + 1)/(2*a*b*(p + 1))), x] + Dist[1/(2*a*(p + 1)), Int
[(a + b*x^2)^(p + 1)*ExpandToSum[2*a*(p + 1)*Q + f*(2*p + 3), x], x], x]] /
; FreeQ[{a, b}, x] && PolyQ[Pq, x] && LtQ[p, -1]
```

Rule 2320

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x]
, Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Functi
onOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[
{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*
(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]
```

Rule 6852

```
Int[(u_.)*((a_.)*(v_)^(m_.))^(p_), x_Symbol] :> Dist[a^IntPart[p]*((a*v^m)^
FracPart[p]/v^(m*FracPart[p])), Int[u*v^(m*p), x], x] /; FreeQ[{a, m, p}, x
] && !IntegerQ[p] && !FreeQ[v, x] && !(EqQ[a, 1] && EqQ[m, 1]) && !(EqQ
[v, x] && EqQ[m, 1])
```

Rubi steps

$$\begin{aligned}
\int e^{c(a+bx)} \tanh^2(ac+bcx)^{5/2} dx &= \left(\coth(ac+bcx) \sqrt{\tanh^2(ac+bcx)} \right) \int e^{c(a+bx)} \tanh^5(ac+bcx) dx \\
&= \frac{\left(\coth(ac+bcx) \sqrt{\tanh^2(ac+bcx)} \right) \text{Subst} \left(\int \frac{(-1+x^2)^5}{(1+x^2)^5} dx, x, e^{c(a+bx)} \right)}{bc} \\
&= \frac{\left(\coth(ac+bcx) \sqrt{\tanh^2(ac+bcx)} \right) \text{Subst} \left(\int \left(1 - \frac{2(1+10x^4+5x^8)}{(1+x^2)^5} \right) dx, x, \right)}{bc} \\
&= \frac{e^{c(a+bx)} \coth(ac+bcx) \sqrt{\tanh^2(ac+bcx)}}{bc} - \frac{\left(2 \coth(ac+bcx) \sqrt{\tanh^2(ac+bcx)} \right)}{bc(1+e^{2c(a+bx)})} \\
&= \frac{e^{c(a+bx)} \coth(ac+bcx) \sqrt{\tanh^2(ac+bcx)}}{bc} - \frac{4e^{c(a+bx)} \coth(ac+bcx) \sqrt{\tanh^2(ac+bcx)}}{bc(1+e^{2c(a+bx)})} \\
&= \frac{e^{c(a+bx)} \coth(ac+bcx) \sqrt{\tanh^2(ac+bcx)}}{bc} - \frac{4e^{c(a+bx)} \coth(ac+bcx) \sqrt{\tanh^2(ac+bcx)}}{bc(1+e^{2c(a+bx)})} \\
&= \frac{e^{c(a+bx)} \coth(ac+bcx) \sqrt{\tanh^2(ac+bcx)}}{bc} - \frac{4e^{c(a+bx)} \coth(ac+bcx) \sqrt{\tanh^2(ac+bcx)}}{bc(1+e^{2c(a+bx)})} \\
&= \frac{e^{c(a+bx)} \coth(ac+bcx) \sqrt{\tanh^2(ac+bcx)}}{bc} - \frac{4e^{c(a+bx)} \coth(ac+bcx) \sqrt{\tanh^2(ac+bcx)}}{bc(1+e^{2c(a+bx)})} \\
&= \frac{e^{c(a+bx)} \coth(ac+bcx) \sqrt{\tanh^2(ac+bcx)}}{bc} - \frac{4e^{c(a+bx)} \coth(ac+bcx) \sqrt{\tanh^2(ac+bcx)}}{bc(1+e^{2c(a+bx)})} \\
&= \frac{e^{c(a+bx)} \coth(ac+bcx) \sqrt{\tanh^2(ac+bcx)}}{bc} - \frac{4e^{c(a+bx)} \coth(ac+bcx) \sqrt{\tanh^2(ac+bcx)}}{bc(1+e^{2c(a+bx)})}
\end{aligned}$$

Mathematica [A]

time = 0.17, size = 133, normalized size = 0.43

$$\frac{\left(e^{c(a+bx)} (33 + 157e^{2c(a+bx)} + 187e^{4c(a+bx)} + 123e^{6c(a+bx)} + 12e^{8c(a+bx)}) - 45(1 + e^{2c(a+bx)})^4 \text{ArcTan}(e^{c(a+bx)}) \right) \coth(c(a+bx)) \sqrt{\tanh^2(c(a+bx))}}{12bc(1 + e^{2c(a+bx)})^4}$$

Antiderivative was successfully verified.

[In] Integrate[E^(c*(a + b*x))*(Tanh[a*c + b*c*x]^2)^(5/2), x]

[Out] ((E^(c*(a + b*x))*(33 + 157*E^(2*c*(a + b*x)) + 187*E^(4*c*(a + b*x)) + 123*E^(6*c*(a + b*x)) + 12*E^(8*c*(a + b*x))) - 45*(1 + E^(2*c*(a + b*x)))^4*ArcTan[E^(c*(a + b*x))])*Coth[c*(a + b*x)]*Sqrt[Tanh[c*(a + b*x)]^2])/(12*b*c*(1 + E^(2*c*(a + b*x)))^4)

Maple [C] Result contains complex when optimal does not.

time = 4.82, size = 324, normalized size = 1.04

method	result
risch	$\frac{(1+e^{2c(bx+a)})\sqrt{\frac{(e^{2c(bx+a)}-1)^2}{(1+e^{2c(bx+a)})^2}}e^{c(bx+a)}}{(e^{2c(bx+a)}-1)bc} + \frac{\sqrt{\frac{(e^{2c(bx+a)}-1)^2}{(1+e^{2c(bx+a)})^2}}e^{c(bx+a)}(75e^{6c(bx+a)}+115e^{4c(bx+a)}+109e^{2c(bx+a)}+21)}{12(e^{2c(bx+a)}-1)(1+e^{2c(bx+a)})^3bc} +$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(c*(b*x+a))*(tanh(b*c*x+a*c)^2)^(5/2),x,method=_RETURNVERBOSE)`

[Out] $1/(\exp(2*c*(b*x+a))-1)*(1+\exp(2*c*(b*x+a)))*((\exp(2*c*(b*x+a))-1)^2/(1+\exp(2*c*(b*x+a)))^2)^{(1/2)}*\exp(c*(b*x+a))/b/c+1/12/(\exp(2*c*(b*x+a))-1)/(1+\exp(2*c*(b*x+a)))^3*((\exp(2*c*(b*x+a))-1)^2/(1+\exp(2*c*(b*x+a)))^2)^{(1/2)}*\exp(c*(b*x+a))*(75*\exp(6*c*(b*x+a))+115*\exp(4*c*(b*x+a))+109*\exp(2*c*(b*x+a))+21)/b/c+15/8*I/(\exp(2*c*(b*x+a))-1)*(1+\exp(2*c*(b*x+a)))*((\exp(2*c*(b*x+a))-1)^2/(1+\exp(2*c*(b*x+a)))^2)^{(1/2)}/c/b*\ln(\exp(c*(b*x+a))-I)-15/8*I/(\exp(2*c*(b*x+a))-1)*(1+\exp(2*c*(b*x+a)))*((\exp(2*c*(b*x+a))-1)^2/(1+\exp(2*c*(b*x+a)))^2)^{(1/2)}/c/b*\ln(\exp(c*(b*x+a))+I)$

Maxima [A]

time = 0.48, size = 145, normalized size = 0.47

$$\frac{15 \arctan(e^{(bcx+ac)})}{4bc} + \frac{12e^{(9bcx+9ac)} + 123e^{(7bcx+7ac)} + 187e^{(5bcx+5ac)} + 157e^{(3bcx+3ac)} + 33e^{(bcx+ac)}}{12bc(e^{(8bcx+8ac)} + 4e^{(6bcx+6ac)} + 6e^{(4bcx+4ac)} + 4e^{(2bcx+2ac)} + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(c*(b*x+a))*(tanh(b*c*x+a*c)^2)^(5/2),x, algorithm="maxima")`

[Out] $-15/4*\arctan(e^{(b*c*x + a*c)})/(b*c) + 1/12*(12*e^{(9*b*c*x + 9*a*c)} + 123*e^{(7*b*c*x + 7*a*c)} + 187*e^{(5*b*c*x + 5*a*c)} + 157*e^{(3*b*c*x + 3*a*c)} + 33*e^{(b*c*x + a*c)})/(b*c*(e^{(8*b*c*x + 8*a*c)} + 4*e^{(6*b*c*x + 6*a*c)} + 6*e^{(4*b*c*x + 4*a*c)} + 4*e^{(2*b*c*x + 2*a*c)} + 1))$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 1226 vs. 2(281) = 562.

time = 0.36, size = 1226, normalized size = 3.94

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(c*(b*x+a))*(tanh(b*c*x+a*c)^2)^(5/2),x, algorithm="fricas")`

[Out] $1/12*(12*\cosh(b*c*x + a*c)^9 + 108*\cosh(b*c*x + a*c)*\sinh(b*c*x + a*c)^8 + 12*\sinh(b*c*x + a*c)^9 + 3*(144*\cosh(b*c*x + a*c)^2 + 41)*\sinh(b*c*x + a*c)$

$$\begin{aligned} &^7 + 123*\cosh(b*c*x + a*c)^7 + 21*(48*\cosh(b*c*x + a*c)^3 + 41*\cosh(b*c*x + \\ &a*c))*\sinh(b*c*x + a*c)^6 + (1512*\cosh(b*c*x + a*c)^4 + 2583*\cosh(b*c*x + \\ &a*c)^2 + 187)*\sinh(b*c*x + a*c)^5 + 187*\cosh(b*c*x + a*c)^5 + (1512*\cosh(b* \\ &c*x + a*c)^5 + 4305*\cosh(b*c*x + a*c)^3 + 935*\cosh(b*c*x + a*c))*\sinh(b*c*x \\ &+ a*c)^4 + (1008*\cosh(b*c*x + a*c)^6 + 4305*\cosh(b*c*x + a*c)^4 + 1870*cos \\ &h(b*c*x + a*c)^2 + 157)*\sinh(b*c*x + a*c)^3 + 157*\cosh(b*c*x + a*c)^3 + (43 \\ &2*\cosh(b*c*x + a*c)^7 + 2583*\cosh(b*c*x + a*c)^5 + 1870*\cosh(b*c*x + a*c)^3 \\ &+ 471*\cosh(b*c*x + a*c))*\sinh(b*c*x + a*c)^2 - 45*(\cosh(b*c*x + a*c)^8 + 8 \\ &*\cosh(b*c*x + a*c)*\sinh(b*c*x + a*c)^7 + \sinh(b*c*x + a*c)^8 + 4*(7*\cosh(b* \\ &c*x + a*c)^2 + 1)*\sinh(b*c*x + a*c)^6 + 4*\cosh(b*c*x + a*c)^6 + 8*(7*\cosh(b \\ &c*x + a*c)^3 + 3*\cosh(b*c*x + a*c))*\sinh(b*c*x + a*c)^5 + 2*(35*\cosh(b*c*x \\ &+ a*c)^4 + 30*\cosh(b*c*x + a*c)^2 + 3)*\sinh(b*c*x + a*c)^4 + 6*\cosh(b*c*x \\ &+ a*c)^4 + 8*(7*\cosh(b*c*x + a*c)^5 + 10*\cosh(b*c*x + a*c)^3 + 3*\cosh(b*c*x \\ &+ a*c))*\sinh(b*c*x + a*c)^3 + 4*(7*\cosh(b*c*x + a*c)^6 + 15*\cosh(b*c*x + a \\ &c)^4 + 9*\cosh(b*c*x + a*c)^2 + 1)*\sinh(b*c*x + a*c)^2 + 4*\cosh(b*c*x + a*c \\ &)^2 + 8*(\cosh(b*c*x + a*c)^7 + 3*\cosh(b*c*x + a*c)^5 + 3*\cosh(b*c*x + a*c)^ \\ &3 + \cosh(b*c*x + a*c))*\sinh(b*c*x + a*c) + 1)*\arctan(\cosh(b*c*x + a*c) + si \\ &nh(b*c*x + a*c)) + (108*\cosh(b*c*x + a*c)^8 + 861*\cosh(b*c*x + a*c)^6 + 935 \\ &*\cosh(b*c*x + a*c)^4 + 471*\cosh(b*c*x + a*c)^2 + 33)*\sinh(b*c*x + a*c) + 33 \\ &*\cosh(b*c*x + a*c))/(b*c*\cosh(b*c*x + a*c)^8 + 8*b*c*\cosh(b*c*x + a*c)*\sinh \\ &(b*c*x + a*c)^7 + b*c*\sinh(b*c*x + a*c)^8 + 4*b*c*\cosh(b*c*x + a*c)^6 + 4*(\\ &7*b*c*\cosh(b*c*x + a*c)^2 + b*c)*\sinh(b*c*x + a*c)^6 + 6*b*c*\cosh(b*c*x + \\ &a*c)^4 + 8*(7*b*c*\cosh(b*c*x + a*c)^3 + 3*b*c*\cosh(b*c*x + a*c))*\sinh(b*c*x \\ &+ a*c)^5 + 2*(35*b*c*\cosh(b*c*x + a*c)^4 + 30*b*c*\cosh(b*c*x + a*c)^2 + 3*b \\ &c)*\sinh(b*c*x + a*c)^4 + 4*b*c*\cosh(b*c*x + a*c)^2 + 8*(7*b*c*\cosh(b*c*x + \\ &a*c)^5 + 10*b*c*\cosh(b*c*x + a*c)^3 + 3*b*c*\cosh(b*c*x + a*c))*\sinh(b*c*x \\ &+ a*c)^3 + 4*(7*b*c*\cosh(b*c*x + a*c)^6 + 15*b*c*\cosh(b*c*x + a*c)^4 + 9*b \\ &c*\cosh(b*c*x + a*c)^2 + b*c)*\sinh(b*c*x + a*c)^2 + b*c + 8*(b*c*\cosh(b*c*x \\ &+ a*c)^7 + 3*b*c*\cosh(b*c*x + a*c)^5 + 3*b*c*\cosh(b*c*x + a*c)^3 + b*c*\cosh \\ &(b*c*x + a*c))*\sinh(b*c*x + a*c)) \end{aligned}$$

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(c*(b*x+a))*(tanh(b*c*x+a*c)**2)**(5/2), x)

[Out] Exception raised: SystemError >> excessive stack use: stack is 6191 deep

Giac [A]

time = 0.45, size = 184, normalized size = 0.59

$$45 \arctan(e^{(bcx+ac)}) \operatorname{sgn}(e^{(2bcx+2ac)} - 1) - 12 e^{(bcx+ac)} \operatorname{sgn}(e^{(2bcx+2ac)} - 1) - \frac{75 e^{(7bcx+7ac)} \operatorname{sgn}(e^{(2bcx+2ac)} - 1) + 115 e^{(5bcx+5ac)} \operatorname{sgn}(e^{(2bcx+2ac)} - 1) + 109 e^{(3bcx+3ac)} \operatorname{sgn}(e^{(2bcx+2ac)} - 1) + 21 e^{(bcx+ac)} \operatorname{sgn}(e^{(2bcx+2ac)} - 1)}{(e^{(2bcx+2ac)} + 1)^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(c*(b*x+a))*(tanh(b*c*x+a*c)^2)^(5/2),x, algorithm="giac")

[Out]
$$-1/12*(45*\arctan(e^{(b*c*x + a*c)})*\operatorname{sgn}(e^{(2*b*c*x + 2*a*c)} - 1) - 12*e^{(b*c*x + a*c)}*\operatorname{sgn}(e^{(2*b*c*x + 2*a*c)} - 1) - (75*e^{(7*b*c*x + 7*a*c)}*\operatorname{sgn}(e^{(2*b*c*x + 2*a*c)} - 1) + 115*e^{(5*b*c*x + 5*a*c)}*\operatorname{sgn}(e^{(2*b*c*x + 2*a*c)} - 1) + 109*e^{(3*b*c*x + 3*a*c)}*\operatorname{sgn}(e^{(2*b*c*x + 2*a*c)} - 1) + 21*e^{(b*c*x + a*c)}*\operatorname{sgn}(e^{(2*b*c*x + 2*a*c)} - 1)))/(e^{(2*b*c*x + 2*a*c)} + 1)^4/(b*c)$$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int e^{c(a+bx)} (\tanh(ac + bcx)^2)^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(c*(a + b*x))*(tanh(a*c + b*c*x)^2)^(5/2),x)

[Out] int(exp(c*(a + b*x))*(tanh(a*c + b*c*x)^2)^(5/2), x)

3.235 $\int e^{c(a+bx)} \tanh^2(ac + bcx)^{3/2} dx$

Optimal. Leaf size=193

$$\frac{e^{c(a+bx)} \coth(ac + bcx) \sqrt{\tanh^2(ac + bcx)}}{bc} - \frac{2e^{c(a+bx)} \coth(ac + bcx) \sqrt{\tanh^2(ac + bcx)}}{bc(1 + e^{2c(a+bx)})^2} + \frac{3e^{c(a+bx)} \coth(ac + bcx) \sqrt{\tanh^2(ac + bcx)}}{bc(1 + e^{2c(a+bx)})^2}$$

[Out] exp(c*(b*x+a))*coth(b*c*x+a*c)*(tanh(b*c*x+a*c)^2)^(1/2)/b/c-2*exp(c*(b*x+a))*coth(b*c*x+a*c)*(tanh(b*c*x+a*c)^2)^(1/2)/b/c/(1+exp(2*c*(b*x+a)))^2+3*exp(c*(b*x+a))*coth(b*c*x+a*c)*(tanh(b*c*x+a*c)^2)^(1/2)/b/c/(1+exp(2*c*(b*x+a)))^2-3*arctan(exp(c*(b*x+a)))*coth(b*c*x+a*c)*(tanh(b*c*x+a*c)^2)^(1/2)/b/c

Rubi [A]

time = 0.20, antiderivative size = 193, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.280$, Rules used = {6852, 2320, 398, 1172, 12, 294, 209}

$$-\frac{3\text{ArcTan}(e^{c(a+bx)}) \sqrt{\tanh^2(ac + bcx)} \coth(ac + bcx)}{bc} + \frac{e^{c(a+bx)} \sqrt{\tanh^2(ac + bcx)} \coth(ac + bcx)}{bc} + \frac{3e^{c(a+bx)} \sqrt{\tanh^2(ac + bcx)} \coth(ac + bcx)}{bc(e^{2c(a+bx)} + 1)} - \frac{2e^{c(a+bx)} \sqrt{\tanh^2(ac + bcx)} \coth(ac + bcx)}{bc(e^{2c(a+bx)} + 1)^2}$$

Antiderivative was successfully verified.

[In] Int[E^(c*(a + b*x))*(Tanh[a*c + b*c*x]^2)^(3/2), x]

[Out] (E^(c*(a + b*x))*Coth[a*c + b*c*x]*Sqrt[Tanh[a*c + b*c*x]^2])/(b*c) - (2*E^(c*(a + b*x))*Coth[a*c + b*c*x]*Sqrt[Tanh[a*c + b*c*x]^2])/(b*c*(1 + E^(2*c*(a + b*x)))) + (3*E^(c*(a + b*x))*Coth[a*c + b*c*x]*Sqrt[Tanh[a*c + b*c*x]^2])/(b*c*(1 + E^(2*c*(a + b*x)))) - (3*ArcTan[E^(c*(a + b*x))]*Coth[a*c + b*c*x]*Sqrt[Tanh[a*c + b*c*x]^2])/(b*c)

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 294

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x]

```

/; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !I
LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]

```

Rule 398

```

Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol]
:> Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a
, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q,
0] && GeQ[p, -q]

```

Rule 1172

```

Int[((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] :> Wi
th[{Qx = PolynomialQuotient[(a + c*x^4)^p, d + e*x^2, x], R = Coeff[Polynom
ialRemainder[(a + c*x^4)^p, d + e*x^2, x], x, 0]}, Simp[(-R)*x*((d + e*x^2)
^(q + 1)/(2*d*(q + 1))), x] + Dist[1/(2*d*(q + 1)), Int[(d + e*x^2)^(q + 1)
*ExpandToSum[2*d*(q + 1)*Qx + R*(2*q + 3), x], x], x]] /; FreeQ[{a, c, d, e
}, x] && NeQ[c*d^2 + a*e^2, 0] && IGtQ[p, 0] && LtQ[q, -1]

```

Rule 2320

```

Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x]
, Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Functi
onOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[
{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))*
(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]

```

Rule 6852

```

Int[(u_)*((a_)*(v_)^(m_))^(p_), x_Symbol] :> Dist[a^IntPart[p]*((a*v^m)^
FracPart[p]/v^(m*FracPart[p])), Int[u*v^(m*p), x], x] /; FreeQ[{a, m, p}, x
] && !IntegerQ[p] && !FreeQ[v, x] && !(EqQ[a, 1] && EqQ[m, 1]) && !(EqQ
[v, x] && EqQ[m, 1])

```

Rubi steps

$$\begin{aligned}
\int e^{c(a+bx)} \tanh^2(ac+bcx)^{3/2} dx &= \left(\coth(ac+bcx) \sqrt{\tanh^2(ac+bcx)} \right) \int e^{c(a+bx)} \tanh^3(ac+bcx) dx \\
&= \frac{\left(\coth(ac+bcx) \sqrt{\tanh^2(ac+bcx)} \right) \text{Subst}\left(\int \frac{(-1+x^2)^3}{(1+x^2)^3} dx, x, e^{c(a+bx)}\right)}{bc} \\
&= \frac{\left(\coth(ac+bcx) \sqrt{\tanh^2(ac+bcx)} \right) \text{Subst}\left(\int \left(1 - \frac{2(1+3x^4)}{(1+x^2)^3}\right) dx, x, e^{c(a+bx)}\right)}{bc} \\
&= \frac{e^{c(a+bx)} \coth(ac+bcx) \sqrt{\tanh^2(ac+bcx)}}{bc} - \frac{\left(2 \coth(ac+bcx) \sqrt{\tanh^2(ac+bcx)}\right)}{bc} \\
&= \frac{e^{c(a+bx)} \coth(ac+bcx) \sqrt{\tanh^2(ac+bcx)}}{bc} - \frac{2e^{c(a+bx)} \coth(ac+bcx) \sqrt{\tanh^2(ac+bcx)}}{bc(1+e^{2c(a+bx)})} \\
&= \frac{e^{c(a+bx)} \coth(ac+bcx) \sqrt{\tanh^2(ac+bcx)}}{bc} - \frac{2e^{c(a+bx)} \coth(ac+bcx) \sqrt{\tanh^2(ac+bcx)}}{bc(1+e^{2c(a+bx)})} \\
&= \frac{e^{c(a+bx)} \coth(ac+bcx) \sqrt{\tanh^2(ac+bcx)}}{bc} - \frac{2e^{c(a+bx)} \coth(ac+bcx) \sqrt{\tanh^2(ac+bcx)}}{bc(1+e^{2c(a+bx)})} \\
&= \frac{e^{c(a+bx)} \coth(ac+bcx) \sqrt{\tanh^2(ac+bcx)}}{bc} - \frac{2e^{c(a+bx)} \coth(ac+bcx) \sqrt{\tanh^2(ac+bcx)}}{bc(1+e^{2c(a+bx)})}
\end{aligned}$$

Mathematica [A]

time = 0.12, size = 104, normalized size = 0.54

$$\frac{\left(e^{c(a+bx)}(2+5e^{2c(a+bx)}+e^{4c(a+bx)})-3(1+e^{2c(a+bx)})^2 \text{ArcTan}(e^{c(a+bx)})\right) \coth(c(a+bx)) \sqrt{\tanh^2(c(a+bx))}}{bc(1+e^{2c(a+bx)})^2}$$

Antiderivative was successfully verified.

[In] Integrate[E^(c*(a + b*x))*(Tanh[a*c + b*c*x]^2)^(3/2), x]

[Out] ((E^(c*(a + b*x))*(2 + 5*E^(2*c*(a + b*x)) + E^(4*c*(a + b*x))) - 3*(1 + E^(2*c*(a + b*x))))^2*ArcTan[E^(c*(a + b*x))]*Coth[c*(a + b*x)]*Sqrt[Tanh[c*(a + b*x)]^2])/(b*c*(1 + E^(2*c*(a + b*x)))^2)

Maple [C] Result contains complex when optimal does not.

time = 4.72, size = 301, normalized size = 1.56

method	result
risch	$\frac{(1+e^{2c(bx+a)})\sqrt{\frac{(e^{2c(bx+a)}-1)^2}{(1+e^{2c(bx+a)})^2}}e^{c(bx+a)}}{(e^{2c(bx+a)}-1)bc} + \frac{\sqrt{\frac{(e^{2c(bx+a)}-1)^2}{(1+e^{2c(bx+a)})^2}}e^{c(bx+a)}(3e^{2c(bx+a)}+1)}{(e^{2c(bx+a)}-1)(1+e^{2c(bx+a)})bc} + \frac{3i(1+e^{2c(bx+a)})\sqrt{\frac{(e^{2c(bx+a)}-1)^2}{(1+e^{2c(bx+a)})^2}}}{2(e^{2c(bx+a)}-1)bc}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(c*(b*x+a))*(tanh(b*c*x+a*c)^2)^(3/2),x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{(\exp(2c(bx+a))-1)(1+\exp(2c(bx+a)))} \left(\frac{(\exp(2c(bx+a))-1)^2}{(1+\exp(2c(bx+a)))^2} \right)^{1/2} \frac{\exp(c(bx+a))}{b/c+1} + \frac{(\exp(2c(bx+a))-1)^2}{(1+\exp(2c(bx+a)))^2} \left(\frac{\exp(c(bx+a))}{b/c+3/2} I(\exp(2c(bx+a))-1)(1+\exp(2c(bx+a))) \right) \left(\frac{(\exp(2c(bx+a))-1)^2}{(1+\exp(2c(bx+a)))^2} \right)^{1/2} \frac{\exp(c(bx+a))}{c/b \ln(\exp(c(bx+a))+I)} - \frac{3}{2} I(\exp(2c(bx+a))-1)(1+\exp(2c(bx+a))) \left(\frac{(\exp(2c(bx+a))-1)^2}{(1+\exp(2c(bx+a)))^2} \right)^{1/2} \frac{\exp(c(bx+a))}{c/b \ln(\exp(c(bx+a))+I)}$

Maxima [A]

time = 0.48, size = 90, normalized size = 0.47

$$-\frac{3 \arctan(e^{(bcx+ac)})}{bc} + \frac{e^{(5bcx+5ac)} + 5e^{(3bcx+3ac)} + 2e^{(bcx+ac)}}{bc(e^{(4bcx+4ac)} + 2e^{(2bcx+2ac)} + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(c*(b*x+a))*(tanh(b*c*x+a*c)^2)^(3/2),x, algorithm="maxima")`

[Out] $-3 \arctan(e^{(b*c*x + a*c)})/(b*c) + (e^{(5*b*c*x + 5*a*c)} + 5e^{(3*b*c*x + 3*a*c)} + 2e^{(b*c*x + a*c)})/(b*c*(e^{(4*b*c*x + 4*a*c)} + 2e^{(2*b*c*x + 2*a*c)} + 1))$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 458 vs. 2(179) = 358.

time = 0.35, size = 458, normalized size = 2.37

sinh(x) = (e^x - e^-x)/2, cosh(x) = (e^x + e^-x)/2, tanh(x) = (e^x - e^-x)/(e^x + e^-x), coth(x) = (e^x + e^-x)/(e^x - e^-x), sech(x) = 2/(e^x + e^-x), csch(x) = 2/(e^x - e^-x), sinh(2x) = 2sinh(x)cosh(x), cosh(2x) = cosh^2(x) + sinh^2(x), tanh(2x) = 2tanh(x)/(1-tanh^2(x)), coth(2x) = (coth^2(x) + 1)/(2coth(x)), sech(2x) = 2sech(x)cosh(x)/(cosh^2(x) + sinh^2(x)), csch(2x) = csch(x)sinh(x)/(cosh^2(x) + sinh^2(x)), sinh(x)sinh(y) = (cosh(x+y) - cosh(x-y))/2, cosh(x)cosh(y) = (cosh(x+y) + cosh(x-y))/2, sinh(x)cosh(y) = (sinh(x+y) + sinh(x-y))/2, cosh(x)sinh(y) = (sinh(x+y) - sinh(x-y))/2, sinh(x)sinh(y) = (cosh(x+y) - cosh(x-y))/2, cosh(x)cosh(y) = (cosh(x+y) + cosh(x-y))/2, tanh(x)sinh(y) = (sinh(x+y) - sinh(x-y))/(cosh(x)cosh(y)), coth(x)cosh(y) = (sinh(x+y) + sinh(x-y))/(cosh(x)cosh(y)), sech(x)sinh(y) = (sinh(x+y) - sinh(x-y))/(cosh(x)cosh(y)), csch(x)cosh(y) = (sinh(x+y) + sinh(x-y))/(cosh(x)cosh(y)), sinh(x)coth(y) = (sinh(x+y) - sinh(x-y))/(sinh(x)cosh(y)), cosh(x)coth(y) = (sinh(x+y) + sinh(x-y))/(sinh(x)cosh(y)), sech(x)coth(y) = (sinh(x+y) - sinh(x-y))/(sinh(x)cosh(y)), csch(x)coth(y) = (sinh(x+y) + sinh(x-y))/(sinh(x)cosh(y)), sinh(x)sinh(y) = (cosh(x+y) - cosh(x-y))/2, cosh(x)cosh(y) = (cosh(x+y) + cosh(x-y))/2, tanh(x)sinh(y) = (sinh(x+y) - sinh(x-y))/(cosh(x)cosh(y)), coth(x)cosh(y) = (sinh(x+y) + sinh(x-y))/(cosh(x)cosh(y)), sech(x)sinh(y) = (sinh(x+y) - sinh(x-y))/(cosh(x)cosh(y)), csch(x)cosh(y) = (sinh(x+y) + sinh(x-y))/(cosh(x)cosh(y)), sinh(x)coth(y) = (sinh(x+y) - sinh(x-y))/(sinh(x)cosh(y)), cosh(x)coth(y) = (sinh(x+y) + sinh(x-y))/(sinh(x)cosh(y)), sech(x)coth(y) = (sinh(x+y) - sinh(x-y))/(sinh(x)cosh(y)), csch(x)coth(y) = (sinh(x+y) + sinh(x-y))/(sinh(x)cosh(y))

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(c*(b*x+a))*(tanh(b*c*x+a*c)^2)^(3/2),x, algorithm="fricas")`

[Out] $(\cosh(b*c*x + a*c))^5 + 5 \cosh(b*c*x + a*c) \sinh(b*c*x + a*c)^4 + \sinh(b*c*x + a*c)^5 + 5(2 \cosh(b*c*x + a*c)^2 + 1) \sinh(b*c*x + a*c)^3 + 5 \cosh(b*c*x + a*c)^3 + 5(2 \cosh(b*c*x + a*c)^3 + 3 \cosh(b*c*x + a*c)) \sinh(b*c*x + a*c)^2 - 3(\cosh(b*c*x + a*c)^4 + 4 \cosh(b*c*x + a*c) \sinh(b*c*x + a*c)^3 + \sinh(b*c*x + a*c)^4 + 2(3 \cosh(b*c*x + a*c)^2 + 1) \sinh(b*c*x + a*c)^2 + 2 \cosh(b*c*x + a*c)^2 + 4(\cosh(b*c*x + a*c)^3 + \cosh(b*c*x + a*c)) \sinh(b*c$

```
*x + a*c) + 1)*arctan(cosh(b*c*x + a*c) + sinh(b*c*x + a*c)) + (5*cosh(b*c*x + a*c)^4 + 15*cosh(b*c*x + a*c)^2 + 2)*sinh(b*c*x + a*c) + 2*cosh(b*c*x + a*c))/(b*c*cosh(b*c*x + a*c)^4 + 4*b*c*cosh(b*c*x + a*c)*sinh(b*c*x + a*c)^3 + b*c*sinh(b*c*x + a*c)^4 + 2*b*c*cosh(b*c*x + a*c)^2 + 2*(3*b*c*cosh(b*c*x + a*c)^2 + b*c)*sinh(b*c*x + a*c)^2 + b*c + 4*(b*c*cosh(b*c*x + a*c)^3 + b*c*cosh(b*c*x + a*c))*sinh(b*c*x + a*c))
```

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(c*(b*x+a))*(tanh(b*c*x+a*c)**2)**(3/2), x)
```

[Out] Timed out

Giac [A]

time = 0.43, size = 129, normalized size = 0.67

$$\frac{3 \arctan(e^{(bcx+ac)}) \operatorname{sgn}(e^{(2bcx+2ac)} - 1) - e^{(bcx+ac)} \operatorname{sgn}(e^{(2bcx+2ac)} - 1) - \frac{3 e^{(3bcx+3ac)} \operatorname{sgn}(e^{(2bcx+2ac)} - 1) + e^{(bcx+ac)} \operatorname{sgn}(e^{(2bcx+2ac)} - 1)}{(e^{(2bcx+2ac)} + 1)^2}}{bc}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(c*(b*x+a))*(tanh(b*c*x+a*c)^2)^(3/2), x, algorithm="giac")
```

```
[Out] -(3*arctan(e^(b*c*x + a*c))*sgn(e^(2*b*c*x + 2*a*c) - 1) - e^(b*c*x + a*c)*sgn(e^(2*b*c*x + 2*a*c) - 1) - (3*e^(3*b*c*x + 3*a*c)*sgn(e^(2*b*c*x + 2*a*c) - 1) + e^(b*c*x + a*c)*sgn(e^(2*b*c*x + 2*a*c) - 1))/(e^(2*b*c*x + 2*a*c) + 1)^2)/(b*c)
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int e^{c(a+bx)} (\tanh(ac + bcx)^2)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(exp(c*(a + b*x))*(tanh(a*c + b*c*x)^2)^(3/2), x)
```

```
[Out] int(exp(c*(a + b*x))*(tanh(a*c + b*c*x)^2)^(3/2), x)
```

3.236 $\int e^{c(a+bx)} \sqrt{\tanh^2(ac+bcx)} dx$

Optimal. Leaf size=83

$$\frac{e^{c(a+bx)} \coth(ac+bcx) \sqrt{\tanh^2(ac+bcx)}}{bc} - \frac{2 \operatorname{ArcTan}(e^{c(a+bx)}) \coth(ac+bcx) \sqrt{\tanh^2(ac+bcx)}}{bc}$$

[Out] $\exp(c*(b*x+a))*\coth(b*c*x+a*c)*(\tanh(b*c*x+a*c)^2)^{(1/2)}/b/c-2*\arctan(\exp(c*(b*x+a)))*\coth(b*c*x+a*c)*(\tanh(b*c*x+a*c)^2)^{(1/2)}/b/c$

Rubi [A]

time = 0.11, antiderivative size = 83, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$, Rules used = {6852, 2320, 396, 209}

$$\frac{e^{c(a+bx)} \sqrt{\tanh^2(ac+bcx)} \coth(ac+bcx)}{bc} - \frac{2 \operatorname{ArcTan}(e^{c(a+bx)}) \sqrt{\tanh^2(ac+bcx)} \coth(ac+bcx)}{bc}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{(c*(a + b*x))*Sqrt[Tanh[a*c + b*c*x]^2]}, x]$

[Out] $(E^{(c*(a + b*x))*Coth[a*c + b*c*x]*Sqrt[Tanh[a*c + b*c*x]^2]}/(b*c) - (2*ArcTan[E^{(c*(a + b*x))*Coth[a*c + b*c*x]*Sqrt[Tanh[a*c + b*c*x]^2]}/(b*c)$

Rule 209

$\text{Int}[(a + (b \cdot x)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2]))* \text{ArcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$

Rule 396

$\text{Int}[(a + (b \cdot x)^{n_1})^{p_1} * ((c + (d \cdot x)^{n_2}))^{p_2}, x_Symbol] \rightarrow \text{Simp}[d*x*((a + b*x^n)^{(p+1)}/(b*(n*(p+1)+1))), x] - \text{Dist}[(a*d - b*c*(n*(p+1)+1))/(b*(n*(p+1)+1)), \text{Int}[(a + b*x^n)^p, x], x] /; \text{FreeQ}\{a, b, c, d, n\}, x\} \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[n*(p+1)+1, 0]$

Rule 2320

$\text{Int}[u, x_Symbol] \rightarrow \text{With}\{v = \text{FunctionOfExponential}[u, x]\}, \text{Dist}[v/D[v, x], \text{Subst}[\text{Int}[\text{FunctionOfExponentialFunction}[u, x]/x, x], x, v], x] /; \text{FunctionOfExponentialQ}[u, x] \ \&\& \ !\text{MatchQ}[u, (w_)*((a_)*(v_)^{(n_)})^{(m_)} /; \text{FreeQ}\{a, m, n\}, x\} \ \&\& \ \text{IntegerQ}[m*n] \ \&\& \ !\text{MatchQ}[u, E^{((c_)*((a_)+(b_)*x))}*(F_)] [v_] /; \text{FreeQ}\{a, b, c\}, x\} \ \&\& \ \text{InverseFunctionQ}[F[x]]$

Rule 6852

```
Int[(u_.)*((a_.)*(v_)^(m_.))^(p_), x_Symbol] := Dist[a^IntPart[p]*((a*v^m)^
FracPart[p]/v^(m*FracPart[p])), Int[u*v^(m*p), x], x] /; FreeQ[{a, m, p}, x
] && !IntegerQ[p] && !FreeQ[v, x] && !(EqQ[a, 1] && EqQ[m, 1]) && !(EqQ
[v, x] && EqQ[m, 1])
```

Rubi steps

$$\begin{aligned} \int e^{c(a+bx)} \sqrt{\tanh^2(ac+bcx)} dx &= \left(\coth(ac+bcx) \sqrt{\tanh^2(ac+bcx)} \right) \int e^{c(a+bx)} \tanh(ac+bcx) dx \\ &= \frac{\left(\coth(ac+bcx) \sqrt{\tanh^2(ac+bcx)} \right) \text{Subst}\left(\int \frac{-1+x^2}{1+x^2} dx, x, e^{c(a+bx)}\right)}{bc} \\ &= \frac{e^{c(a+bx)} \coth(ac+bcx) \sqrt{\tanh^2(ac+bcx)}}{bc} - \frac{\left(2 \coth(ac+bcx) \sqrt{\tanh^2(ac+bcx)}\right)}{bc} \\ &= \frac{e^{c(a+bx)} \coth(ac+bcx) \sqrt{\tanh^2(ac+bcx)}}{bc} - \frac{2 \tan^{-1}\left(e^{c(a+bx)}\right) \coth(ac+bcx)}{bc} \end{aligned}$$

Mathematica [A]

time = 0.04, size = 51, normalized size = 0.61

$$\frac{\left(e^{c(a+bx)} - 2 \text{ArcTan}\left(e^{c(a+bx)}\right)\right) \coth(c(a+bx)) \sqrt{\tanh^2(c(a+bx))}}{bc}$$

Antiderivative was successfully verified.

[In] Integrate[E^(c*(a + b*x))*Sqrt[Tanh[a*c + b*c*x]^2], x]

[Out] ((E^(c*(a + b*x)) - 2*ArcTan[E^(c*(a + b*x))])*Coth[c*(a + b*x)]*Sqrt[Tanh[c*(a + b*x)]^2])/(b*c)

Maple [C] Result contains complex when optimal does not.

time = 4.91, size = 218, normalized size = 2.63

method	result
risch	$\frac{(1+e^{2c(bx+a)}) \sqrt{\frac{(e^{2c(bx+a)}-1)^2}{(1+e^{2c(bx+a)})^2}} e^{c(bx+a)}}{(e^{2c(bx+a)}-1)bc} + \frac{i(1+e^{2c(bx+a)}) \sqrt{\frac{(e^{2c(bx+a)}-1)^2}{(1+e^{2c(bx+a)})^2}} \ln(e^{c(bx+a)}-i)}{(e^{2c(bx+a)}-1)cb} - \frac{i(1+e^{2c(bx+a)}) \sqrt{\frac{(e^{2c(bx+a)}-1)^2}{(1+e^{2c(bx+a)})^2}}}{(e^{2c(bx+a)}-1)cb}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(c*(b*x+a))*(tanh(b*c*x+a*c)^2)^(1/2), x, method=_RETURNVERBOSE)

[Out] $\frac{1}{(\exp(2cx+a)-1)(1+\exp(2cx+a))} \left(\frac{(\exp(2cx+a)-1)^2}{(1+\exp(2cx+a))^2} \right)^{1/2} \exp(cx+a)/b/c + I \left(\frac{(\exp(2cx+a)-1)^2}{(1+\exp(2cx+a))^2} \right)^{1/2} / (\exp(2cx+a)-1)(1+\exp(2cx+a))/c/b \ln(\exp(cx+a)-1) - I \left(\frac{(\exp(2cx+a)-1)^2}{(1+\exp(2cx+a))^2} \right)^{1/2} / (\exp(2cx+a)-1)(1+\exp(2cx+a))/c/b \ln(\exp(cx+a)+1)$

Maxima [A]

time = 0.48, size = 35, normalized size = 0.42

$$-\frac{2 \arctan(e^{(bcx+ac)})}{bc} + \frac{e^{(bcx+ac)}}{bc}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(c*(b*x+a))*(tanh(b*c*x+a*c)^2)^(1/2),x, algorithm="maxima")`

[Out] $-2 \arctan(e^{(b*c*x + a*c)})/(b*c) + e^{(b*c*x + a*c)}/(b*c)$

Fricas [A]

time = 0.33, size = 53, normalized size = 0.64

$$-\frac{2 \arctan(\cosh(bc x + ac) + \sinh(bc x + ac)) - \cosh(bc x + ac) - \sinh(bc x + ac)}{bc}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(c*(b*x+a))*(tanh(b*c*x+a*c)^2)^(1/2),x, algorithm="fricas")`

[Out] $-(2 \arctan(\cosh(b*c*x + a*c) + \sinh(b*c*x + a*c)) - \cosh(b*c*x + a*c) - \sinh(b*c*x + a*c))/(b*c)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$e^{ac} \int \sqrt{\tanh^2(ac + bcx)} e^{bcx} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(c*(b*x+a))*(tanh(b*c*x+a*c)**2)**(1/2),x)`

[Out] $\exp(a*c) * \text{Integral}(\sqrt{\tanh(a*c + b*c*x)**2} * \exp(b*c*x), x)$

Giac [A]

time = 0.41, size = 60, normalized size = 0.72

$$-\frac{2 \arctan(e^{(bcx+ac)}) \operatorname{sgn}(e^{(2bcx+2ac)} - 1) - e^{(bcx+ac)} \operatorname{sgn}(e^{(2bcx+2ac)} - 1)}{bc}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(c*(b*x+a))*(tanh(b*c*x+a*c)^2)^(1/2),x, algorithm="giac")

[Out] $-(2*\arctan(e^{(b*c*x + a*c)})*\operatorname{sgn}(e^{(2*b*c*x + 2*a*c)} - 1) - e^{(b*c*x + a*c)}*\operatorname{sgn}(e^{(2*b*c*x + 2*a*c)} - 1))/(b*c)$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int e^{c(a+bx)} \sqrt{\tanh(ac+bcx)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(c*(a + b*x))*(tanh(a*c + b*c*x)^2)^(1/2),x)

[Out] int(exp(c*(a + b*x))*(tanh(a*c + b*c*x)^2)^(1/2), x)

$$3.237 \quad \int \frac{e^{c(a+bx)}}{\sqrt{\tanh^2(ac+bcx)}} dx$$

Optimal. Leaf size=83

$$\frac{e^{c(a+bx)} \tanh(ac+bcx)}{bc\sqrt{\tanh^2(ac+bcx)}} - \frac{2 \tanh^{-1}(e^{c(a+bx)}) \tanh(ac+bcx)}{bc\sqrt{\tanh^2(ac+bcx)}}$$

[Out] exp(c*(b*x+a))*tanh(b*c*x+a*c)/b/c/(tanh(b*c*x+a*c)^2)^(1/2)-2*arctanh(exp(c*(b*x+a))*tanh(b*c*x+a*c)/b/c/(tanh(b*c*x+a*c)^2)^(1/2)

Rubi [A]

time = 0.14, antiderivative size = 83, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$, Rules used = {6852, 2320, 396, 212}

$$\frac{e^{c(a+bx)} \tanh(ac+bcx)}{bc\sqrt{\tanh^2(ac+bcx)}} - \frac{2 \tanh^{-1}(e^{c(a+bx)}) \tanh(ac+bcx)}{bc\sqrt{\tanh^2(ac+bcx)}}$$

Antiderivative was successfully verified.

[In] Int[E^(c*(a + b*x))/Sqrt[Tanh[a*c + b*c*x]^2], x]

[Out] (E^(c*(a + b*x))*Tanh[a*c + b*c*x])/(b*c*Sqrt[Tanh[a*c + b*c*x]^2]) - (2*ArcTanh[E^(c*(a + b*x))]*Tanh[a*c + b*c*x])/(b*c*Sqrt[Tanh[a*c + b*c*x]^2])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 396

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[d*x*((a + b*x^n)^(p + 1)/(b*(n*(p + 1) + 1))), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(b*(n*(p + 1) + 1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]

Rule 2320

Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*

(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]

Rule 6852

Int[(u_)*((a_)*(v_)^(m_.))^p_], x_Symbol] := Dist[a^IntPart[p]*((a*v^m)^FracPart[p]/v^(m*FracPart[p])), Int[u*v^(m*p), x], x] /; FreeQ[{a, m, p}, x] && !IntegerQ[p] && !FreeQ[v, x] && !(EqQ[a, 1] && EqQ[m, 1]) && !(EqQ[v, x] && EqQ[m, 1])

Rubi steps

$$\begin{aligned} \int \frac{e^{c(a+bx)}}{\sqrt{\tanh^2(ac+bcx)}} dx &= \frac{\tanh(ac+bcx) \int e^{c(a+bx)} \coth(ac+bcx) dx}{\sqrt{\tanh^2(ac+bcx)}} \\ &= \frac{\tanh(ac+bcx) \text{Subst}\left(\int \frac{-1-x^2}{1-x^2} dx, x, e^{c(a+bx)}\right)}{bc\sqrt{\tanh^2(ac+bcx)}} \\ &= \frac{e^{c(a+bx)} \tanh(ac+bcx)}{bc\sqrt{\tanh^2(ac+bcx)}} - \frac{(2 \tanh(ac+bcx)) \text{Subst}\left(\int \frac{1}{1-x^2} dx, x, e^{c(a+bx)}\right)}{bc\sqrt{\tanh^2(ac+bcx)}} \\ &= \frac{e^{c(a+bx)} \tanh(ac+bcx)}{bc\sqrt{\tanh^2(ac+bcx)}} - \frac{2 \tanh^{-1}(e^{c(a+bx)}) \tanh(ac+bcx)}{bc\sqrt{\tanh^2(ac+bcx)}} \end{aligned}$$

Mathematica [A]

time = 0.10, size = 51, normalized size = 0.61

$$\frac{(e^{c(a+bx)} - 2 \tanh^{-1}(e^{c(a+bx)})) \tanh(c(a+bx))}{bc\sqrt{\tanh^2(c(a+bx))}}$$

Antiderivative was successfully verified.

[In] Integrate[E^(c*(a + b*x))/Sqrt[Tanh[a*c + b*c*x]^2], x]

[Out] ((E^(c*(a + b*x)) - 2*ArcTanh[E^(c*(a + b*x))])*Tanh[c*(a + b*x)])/(b*c*Sqrt[Tanh[c*(a + b*x)]^2])

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 212 vs. 2(77) = 154.

time = 7.08, size = 213, normalized size = 2.57

method	result	size
--------	--------	------

risch	$\frac{(e^{2c(bx+a)}-1)e^{c(bx+a)}}{\sqrt{\frac{(e^{2c(bx+a)}-1)^2}{(1+e^{2c(bx+a)})^2}}(1+e^{2c(bx+a)})bc} + \frac{(e^{2c(bx+a)}-1)\ln(e^{c(bx+a)}-1)}{\sqrt{\frac{(e^{2c(bx+a)}-1)^2}{(1+e^{2c(bx+a)})^2}}(1+e^{2c(bx+a)})bc} - \frac{(e^{2c(bx+a)}-1)\ln(e^{c(bx+a)}+1)}{\sqrt{\frac{(e^{2c(bx+a)}-1)^2}{(1+e^{2c(bx+a)})^2}}(1+e^{2c(bx+a)})bc}$	213
-------	--	-----

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(c*(b*x+a))/(tanh(b*c*x+a*c)^2)^(1/2),x,method=_RETURNVERBOSE)`

[Out] $1/((\exp(2*c*(b*x+a))-1)^2/(1+\exp(2*c*(b*x+a)))^2)^(1/2)/(1+\exp(2*c*(b*x+a)))*(\exp(2*c*(b*x+a))-1)*\exp(c*(b*x+a))/b/c+1/((\exp(2*c*(b*x+a))-1)^2/(1+\exp(2*c*(b*x+a)))^2)^(1/2)/(1+\exp(2*c*(b*x+a)))*(\exp(2*c*(b*x+a))-1)/b/c*\ln(\exp(c*(b*x+a))-1)-1/((\exp(2*c*(b*x+a))-1)^2/(1+\exp(2*c*(b*x+a)))^2)^(1/2)/(1+\exp(2*c*(b*x+a)))*(\exp(2*c*(b*x+a))-1)/b/c*\ln(\exp(c*(b*x+a))+1)$

Maxima [A]

time = 0.48, size = 56, normalized size = 0.67

$$\frac{e^{(bcx+ac)}}{bc} - \frac{\log(e^{(bcx+ac)} + 1)}{bc} + \frac{\log(e^{(bcx+ac)} - 1)}{bc}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(c*(b*x+a))/(tanh(b*c*x+a*c)^2)^(1/2),x, algorithm="maxima")`

[Out] $e^{(b*c*x + a*c)}/(b*c) - \log(e^{(b*c*x + a*c)} + 1)/(b*c) + \log(e^{(b*c*x + a*c)} - 1)/(b*c)$

Fricas [A]

time = 0.34, size = 70, normalized size = 0.84

$$\frac{\cosh(bc x + ac) - \log(\cosh(bc x + ac) + \sinh(bc x + ac) + 1) + \log(\cosh(bc x + ac) + \sinh(bc x + ac) - 1) + \sinh(bc x + ac)}{bc}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(c*(b*x+a))/(tanh(b*c*x+a*c)^2)^(1/2),x, algorithm="fricas")`

[Out] $(\cosh(b*c*x + a*c) - \log(\cosh(b*c*x + a*c) + \sinh(b*c*x + a*c) + 1) + \log(\cosh(b*c*x + a*c) + \sinh(b*c*x + a*c) - 1) + \sinh(b*c*x + a*c))/(b*c)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$e^{ac} \int \frac{e^{bcx}}{\sqrt{\tanh^2(ac + bcx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(c*(b*x+a))/(tanh(b*c*x+a*c)**2)**(1/2),x)`

[Out] $\exp(a*c)*\text{Integral}(\exp(b*c*x)/\sqrt{\tanh(a*c + b*c*x)**2}, x)$

Giac [A]

time = 0.43, size = 88, normalized size = 1.06

$$\frac{e^{(bcx+ac)} \operatorname{sgn}(e^{(2bcx+2ac)} - 1) - \log(e^{(bcx+ac)} + 1) \operatorname{sgn}(e^{(2bcx+2ac)} - 1) + \log(|e^{(bcx+ac)} - 1|) \operatorname{sgn}(e^{(2bcx+2ac)} - 1)}{bc}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{integrate}(\exp(c*(b*x+a))/(\tanh(b*c*x+a*c)^2)^{(1/2)}, x, \text{algorithm}="giac")$

[Out] $(e^{(b*c*x + a*c)} * \operatorname{sgn}(e^{(2*b*c*x + 2*a*c)} - 1) - \log(e^{(b*c*x + a*c)} + 1) * \operatorname{sgn}(e^{(2*b*c*x + 2*a*c)} - 1) + \log(\operatorname{abs}(e^{(b*c*x + a*c)} - 1)) * \operatorname{sgn}(e^{(2*b*c*x + 2*a*c)} - 1)) / (b*c)$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{e^{c(a+bx)}}{\sqrt{\tanh(ac + bcx)^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{int}(\exp(c*(a + b*x))/(\tanh(a*c + b*c*x)^2)^{(1/2)}, x)$

[Out] $\text{int}(\exp(c*(a + b*x))/(\tanh(a*c + b*c*x)^2)^{(1/2)}, x)$

$$3.238 \quad \int \frac{e^{c(a+bx)}}{\tanh^2(ac+bcx)^{3/2}} dx$$

Optimal. Leaf size=197

$$\frac{e^{c(a+bx)} \tanh(ac+bcx)}{bc \sqrt{\tanh^2(ac+bcx)}} - \frac{2e^{c(a+bx)} \tanh(ac+bcx)}{bc(1-e^{2c(a+bx)})^2 \sqrt{\tanh^2(ac+bcx)}} + \frac{3e^{c(a+bx)} \tanh(ac+bcx)}{bc(1-e^{2c(a+bx)}) \sqrt{\tanh^2(ac+bcx)}} - 3 \tanh^{-1} \left(\frac{e^{c(a+bx)} \tanh(ac+bcx)}{bc \sqrt{\tanh^2(ac+bcx)}} \right)$$

[Out] exp(c*(b*x+a))*tanh(b*c*x+a*c)/b/c/(tanh(b*c*x+a*c)^2)^(1/2)-2*exp(c*(b*x+a))*tanh(b*c*x+a*c)/b/c/(1-exp(2*c*(b*x+a)))^2/(tanh(b*c*x+a*c)^2)^(1/2)+3*exp(c*(b*x+a))*tanh(b*c*x+a*c)/b/c/(1-exp(2*c*(b*x+a)))/(tanh(b*c*x+a*c)^2)^(1/2)-3*arctanh(exp(c*(b*x+a))*tanh(b*c*x+a*c)/b/c/(tanh(b*c*x+a*c)^2)^(1/2))

Rubi [A]

time = 0.60, antiderivative size = 197, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.280$, Rules used = {6852, 2320, 398, 1172, 12, 294, 213}

$$\frac{e^{c(a+bx)} \tanh(ac+bcx)}{bc \sqrt{\tanh^2(ac+bcx)}} + \frac{3e^{c(a+bx)} \tanh(ac+bcx)}{bc(1-e^{2c(a+bx)}) \sqrt{\tanh^2(ac+bcx)}} - \frac{2e^{c(a+bx)} \tanh(ac+bcx)}{bc(1-e^{2c(a+bx)})^2 \sqrt{\tanh^2(ac+bcx)}} - \frac{3 \tanh^{-1} \left(\frac{e^{c(a+bx)} \tanh(ac+bcx)}{bc \sqrt{\tanh^2(ac+bcx)}} \right)}{bc \sqrt{\tanh^2(ac+bcx)}}$$

Antiderivative was successfully verified.

[In] Int[E^(c*(a + b*x))/(Tanh[a*c + b*c*x]^2)^(3/2), x]

[Out] (E^(c*(a + b*x))*Tanh[a*c + b*c*x])/(b*c*Sqrt[Tanh[a*c + b*c*x]^2]) - (2*E^(c*(a + b*x))*Tanh[a*c + b*c*x])/(b*c*(1 - E^(2*c*(a + b*x)))^2*Sqrt[Tanh[a*c + b*c*x]^2]) + (3*E^(c*(a + b*x))*Tanh[a*c + b*c*x])/(b*c*(1 - E^(2*c*(a + b*x))))*Sqrt[Tanh[a*c + b*c*x]^2]) - (3*ArcTanh[E^(c*(a + b*x))*Tanh[a*c + b*c*x])/(b*c*Sqrt[Tanh[a*c + b*c*x]^2])

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 213

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])

Rule 294

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n-1)*(c*x)^(m-n+1)*((a + b*x^n)^(p+1)/(b*n*(p+1))), x] - Dist[c^n

```
*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x]
;/; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !I
LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 398

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:> Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a
, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q,
0] && GeQ[p, -q]
```

Rule 1172

```
Int[((d_) + (e_.)*(x_)^2)^(q_)*((a_) + (c_.)*(x_)^4)^(p_), x_Symbol] :> Wi
th[{Qx = PolynomialQuotient[(a + c*x^4)^p, d + e*x^2, x], R = Coeff[Polynom
ialRemainder[(a + c*x^4)^p, d + e*x^2, x], x, 0]}, Simp[(-R)*x*((d + e*x^2)
^(q + 1)/(2*d*(q + 1))), x] + Dist[1/(2*d*(q + 1)), Int[(d + e*x^2)^(q + 1)
*ExpandToSum[2*d*(q + 1)*Qx + R*(2*q + 3), x], x], x] /; FreeQ[{a, c, d, e
}, x] && NeQ[c*d^2 + a*e^2, 0] && IGtQ[p, 0] && LtQ[q, -1]
```

Rule 2320

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x]
, Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; Functi
onOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[
{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*
(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 6852

```
Int[(u_.)*((a_.)*(v_)^(m_.))^(p_), x_Symbol] :> Dist[a^IntPart[p]*((a*v^m)^
FracPart[p]/v^(m*FracPart[p])), Int[u*v^(m*p), x], x] /; FreeQ[{a, m, p}, x
] && !IntegerQ[p] && !FreeQ[v, x] && !(EqQ[a, 1] && EqQ[m, 1]) && !(EqQ
[v, x] && EqQ[m, 1])
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{c(a+bx)}}{\tanh^2(ac+bcx)^{3/2}} dx &= \frac{\tanh(ac+bcx) \int e^{c(a+bx)} \coth^3(ac+bcx) dx}{\sqrt{\tanh^2(ac+bcx)}} \\
&= \frac{\tanh(ac+bcx) \text{Subst}\left(\int \frac{(1+x^2)^3}{(-1+x^2)^3} dx, x, e^{c(a+bx)}\right)}{bc\sqrt{\tanh^2(ac+bcx)}} \\
&= \frac{\tanh(ac+bcx) \text{Subst}\left(\int \left(1 + \frac{2(1+3x^4)}{(-1+x^2)^3}\right) dx, x, e^{c(a+bx)}\right)}{bc\sqrt{\tanh^2(ac+bcx)}} \\
&= \frac{e^{c(a+bx)} \tanh(ac+bcx)}{bc\sqrt{\tanh^2(ac+bcx)}} + \frac{(2 \tanh(ac+bcx)) \text{Subst}\left(\int \frac{1+3x^4}{(-1+x^2)^3} dx, x, e^{c(a+bx)}\right)}{bc\sqrt{\tanh^2(ac+bcx)}} \\
&= \frac{e^{c(a+bx)} \tanh(ac+bcx)}{bc\sqrt{\tanh^2(ac+bcx)}} - \frac{2e^{c(a+bx)} \tanh(ac+bcx)}{bc(1-e^{2c(a+bx)})^2 \sqrt{\tanh^2(ac+bcx)}} + \frac{\tanh(ac+bcx)}{bc} \\
&= \frac{e^{c(a+bx)} \tanh(ac+bcx)}{bc\sqrt{\tanh^2(ac+bcx)}} - \frac{2e^{c(a+bx)} \tanh(ac+bcx)}{bc(1-e^{2c(a+bx)})^2 \sqrt{\tanh^2(ac+bcx)}} + \frac{(6 \tanh(ac+bcx))}{bc} \\
&= \frac{e^{c(a+bx)} \tanh(ac+bcx)}{bc\sqrt{\tanh^2(ac+bcx)}} - \frac{2e^{c(a+bx)} \tanh(ac+bcx)}{bc(1-e^{2c(a+bx)})^2 \sqrt{\tanh^2(ac+bcx)}} + \frac{3e^{c(a+bx)}}{bc(1-e^{2c(a+bx)})} \\
&= \frac{e^{c(a+bx)} \tanh(ac+bcx)}{bc\sqrt{\tanh^2(ac+bcx)}} - \frac{2e^{c(a+bx)} \tanh(ac+bcx)}{bc(1-e^{2c(a+bx)})^2 \sqrt{\tanh^2(ac+bcx)}} + \frac{3e^{c(a+bx)}}{bc(1-e^{2c(a+bx)})}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 7.26, size = 334, normalized size = 1.70

$$\frac{e^{-bcx} \left(-21(252105 + 507305e^{2c(a+bx)} + 173916e^{4c(a+bx)} - 154296e^{6c(a+bx)} - 73885e^{8c(a+bx)} + 4887e^{10c(a+bx)}) - \frac{315(-16807 - 28218e^{2c(a+bx)} + 1173e^{4c(a+bx)} + 4299e^{6c(a+bx)} - 1434e^{8c(a+bx)} + 7e^{12c(a+bx)})}{\sqrt{\tanh^2(ac+bcx)}} + \frac{256e^{4c(a+bx)}(1 + e^{2c(a+bx)})^2}{bc} {}_2F_1\left(\frac{3}{2}, 2, 2, 2, 1, 1, \frac{1}{2}, e^{2c(a+bx)}\right) + \frac{256e^{6c(a+bx)}(1 + e^{2c(a+bx)})^2}{bc} {}_2F_1\left(\frac{3}{2}, 2, 2, 2, 2, 1, 1, \frac{1}{2}, e^{2c(a+bx)}\right) \right) \tanh^3(c(a+bx))}{60480c \tanh^3(c(a+bx))^{3/2}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[E^(c*(a + b*x))/(Tanh[a*c + b*c*x]^2)^(3/2), x]

[Out] -1/60480*((-21*(252105 + 507305*E^(2*c*(a + b*x)) + 173916*E^(4*c*(a + b*x)) - 154296*E^(6*c*(a + b*x)) - 73885*E^(8*c*(a + b*x)) + 4887*E^(10*c*(a + b*x))) - (315*(-16807 - 28218*E^(2*c*(a + b*x)) + 1173*E^(4*c*(a + b*x)) + 17748*E^(6*c*(a + b*x)) + 4299*E^(8*c*(a + b*x)) - 1434*E^(10*c*(a + b*x)) + 7*E^(12*c*(a + b*x)))*ArcTanh[Sqrt[E^(2*c*(a + b*x))]])/Sqrt[E^(2*c*(a + b*x))]

$b*x))]] + 384*E^{(8*c*(a + b*x))*(1 + E^{(2*c*(a + b*x)))^2*(7 + 5*E^{(2*c*(a + b*x))})}*HypergeometricPFQ[\{3/2, 2, 2, 2, 2\}, \{1, 1, 1, 11/2\}, E^{(2*c*(a + b*x))}] + 256*E^{(8*c*(a + b*x))*(1 + E^{(2*c*(a + b*x)))^3}*HypergeometricPFQ[\{3/2, 2, 2, 2, 2, 2\}, \{1, 1, 1, 1, 11/2\}, E^{(2*c*(a + b*x))}]]*Tanh[c*(a + b*x)]^3/(b*c*E^{(5*c*(a + b*x))*(Tanh[c*(a + b*x)]^2)^{(3/2)}}$

Maple [A]

time = 6.77, size = 298, normalized size = 1.51

method	result
risch	$\frac{(e^{2c(bx+a)}-1)e^{c(bx+a)}}{\sqrt{\frac{(e^{2c(bx+a)}-1)^2}{(1+e^{2c(bx+a)})^2}}(1+e^{2c(bx+a)})bc} - \frac{e^{c(bx+a)}(3e^{2c(bx+a)}-1)}{(e^{2c(bx+a)}-1)(1+e^{2c(bx+a)})\sqrt{\frac{(e^{2c(bx+a)}-1)^2}{(1+e^{2c(bx+a)})^2}}bc} + \frac{3(e^{2c(bx+a)}-1)\ln(e^{c(bx+a)}-1)}{2\sqrt{\frac{(e^{2c(bx+a)}-1)^2}{(1+e^{2c(bx+a)})^2}}(1+e^{2c(bx+a)})}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(c*(b*x+a))/(tanh(b*c*x+a*c)^2)^(3/2),x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{((\exp(2c*(b*x+a))-1)^2/(1+\exp(2c*(b*x+a)))^2)^{(1/2)/(1+\exp(2c*(b*x+a)))}*(\exp(2c*(b*x+a))-1)*\exp(c*(b*x+a))/b/c-1/(\exp(2c*(b*x+a))-1)/(1+\exp(2c*(b*x+a)))}/((\exp(2c*(b*x+a))-1)^2/(1+\exp(2c*(b*x+a)))^2)^{(1/2)*\exp(c*(b*x+a))}*(3*\exp(2c*(b*x+a))-1)/b/c+3/2/((\exp(2c*(b*x+a))-1)^2/(1+\exp(2c*(b*x+a)))^2)^{(1/2)/(1+\exp(2c*(b*x+a)))}*(\exp(2c*(b*x+a))-1)/b/c*\ln(\exp(c*(b*x+a))-1)-3/2/((\exp(2c*(b*x+a))-1)^2/(1+\exp(2c*(b*x+a)))^2)^{(1/2)/(1+\exp(2c*(b*x+a)))}*(\exp(2c*(b*x+a))-1)/b/c*\ln(\exp(c*(b*x+a))+1)}$

Maxima [A]

time = 0.49, size = 112, normalized size = 0.57

$$-\frac{3 \log(e^{(bcx+ac)} + 1)}{2bc} + \frac{3 \log(e^{(bcx+ac)} - 1)}{2bc} + \frac{e^{(5bcx+5ac)} - 5e^{(3bcx+3ac)} + 2e^{(bcx+ac)}}{bc(e^{(4bcx+4ac)} - 2e^{(2bcx+2ac)} + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(c*(b*x+a))/(tanh(b*c*x+a*c)^2)^(3/2),x, algorithm="maxima")`

[Out] $-3/2*\log(e^{(b*c*x + a*c)} + 1)/(b*c) + 3/2*\log(e^{(b*c*x + a*c)} - 1)/(b*c) + (e^{(5*b*c*x + 5*a*c)} - 5*e^{(3*b*c*x + 3*a*c)} + 2*e^{(b*c*x + a*c)})/(b*c*(e^{(4*b*c*x + 4*a*c)} - 2*e^{(2*b*c*x + 2*a*c)} + 1))$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 613 vs. 2(179) = 358.

time = 0.36, size = 613, normalized size = 3.11

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(c*(b*x+a))/(tanh(b*c*x+a*c)^2)^(3/2),x, algorithm="fricas")
[Out] 1/2*(2*cosh(b*c*x + a*c)^5 + 10*cosh(b*c*x + a*c)*sinh(b*c*x + a*c)^4 + 2*s
inh(b*c*x + a*c)^5 + 10*(2*cosh(b*c*x + a*c)^2 - 1)*sinh(b*c*x + a*c)^3 - 1
0*cosh(b*c*x + a*c)^3 + 10*(2*cosh(b*c*x + a*c)^3 - 3*cosh(b*c*x + a*c))*si
nh(b*c*x + a*c)^2 - 3*(cosh(b*c*x + a*c)^4 + 4*cosh(b*c*x + a*c)*sinh(b*c*x
+ a*c)^3 + sinh(b*c*x + a*c)^4 + 2*(3*cosh(b*c*x + a*c)^2 - 1)*sinh(b*c*x
+ a*c)^2 - 2*cosh(b*c*x + a*c)^2 + 4*(cosh(b*c*x + a*c)^3 - cosh(b*c*x + a*
c))*sinh(b*c*x + a*c) + 1)*log(cosh(b*c*x + a*c) + sinh(b*c*x + a*c) + 1) +
3*(cosh(b*c*x + a*c)^4 + 4*cosh(b*c*x + a*c)*sinh(b*c*x + a*c)^3 + sinh(b*
c*x + a*c)^4 + 2*(3*cosh(b*c*x + a*c)^2 - 1)*sinh(b*c*x + a*c)^2 - 2*cosh(b
*c*x + a*c)^2 + 4*(cosh(b*c*x + a*c)^3 - cosh(b*c*x + a*c))*sinh(b*c*x + a*
c) + 1)*log(cosh(b*c*x + a*c) + sinh(b*c*x + a*c) - 1) + 2*(5*cosh(b*c*x +
a*c)^4 - 15*cosh(b*c*x + a*c)^2 + 2)*sinh(b*c*x + a*c) + 4*cosh(b*c*x + a*c
))/(b*c*cosh(b*c*x + a*c)^4 + 4*b*c*cosh(b*c*x + a*c)*sinh(b*c*x + a*c)^3 +
b*c*sinh(b*c*x + a*c)^4 - 2*b*c*cosh(b*c*x + a*c)^2 + 2*(3*b*c*cosh(b*c*x
+ a*c)^2 - b*c)*sinh(b*c*x + a*c)^2 + b*c + 4*(b*c*cosh(b*c*x + a*c)^3 - b*
c*cosh(b*c*x + a*c))*sinh(b*c*x + a*c))
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$e^{ac} \int \frac{e^{bcx}}{(\tanh^2(ac + bcx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(c*(b*x+a))/(tanh(b*c*x+a*c)**2)**(3/2),x)
```

```
[Out] exp(a*c)*Integral(exp(b*c*x)/(tanh(a*c + b*c*x)**2)**(3/2), x)
```

Giac [A]

time = 0.49, size = 161, normalized size = 0.82

$$\frac{2e^{(bcx+ac)}\operatorname{sgn}(e^{(2bcx+2ac)}-1) - 3\log(e^{(bcx+ac)}+1)\operatorname{sgn}(e^{(2bcx+2ac)}-1) + 3\log(|e^{(bcx+ac)}-1|)\operatorname{sgn}(e^{(2bcx+2ac)}-1) - \frac{2(3e^{(3bcx+3ac)}\operatorname{sgn}(e^{(2bcx+2ac)}-1) - e^{(bcx+ac)}\operatorname{sgn}(e^{(2bcx+2ac)}-1))}{(e^{(2bcx+2ac)}-1)^2}}{2bc}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(c*(b*x+a))/(tanh(b*c*x+a*c)^2)^(3/2),x, algorithm="giac")
```

```
[Out] 1/2*(2*e^(b*c*x + a*c)*sgn(e^(2*b*c*x + 2*a*c) - 1) - 3*log(e^(b*c*x + a*c)
+ 1)*sgn(e^(2*b*c*x + 2*a*c) - 1) + 3*log(abs(e^(b*c*x + a*c) - 1))*sgn(e^
(2*b*c*x + 2*a*c) - 1) - 2*(3*e^(3*b*c*x + 3*a*c)*sgn(e^(2*b*c*x + 2*a*c) -
1) - e^(b*c*x + a*c)*sgn(e^(2*b*c*x + 2*a*c) - 1))/(e^(2*b*c*x + 2*a*c) -
1)^2)/(b*c)
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{e^{c(a+bx)}}{(\tanh(ac + bcx)^2)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(exp(c*(a + b*x))/(tanh(a*c + b*c*x)^2)^(3/2), x)
```

```
[Out] int(exp(c*(a + b*x))/(tanh(a*c + b*c*x)^2)^(3/2), x)
```


$$3.239 \quad \int \frac{e^{c(a+bx)}}{\tanh^2(ac+bcx)^{5/2}} dx$$

Optimal. Leaf size=319

$$\frac{e^{c(a+bx)} \tanh(ac+bcx)}{bc \sqrt{\tanh^2(ac+bcx)}} - \frac{4e^{c(a+bx)} \tanh(ac+bcx)}{bc(1-e^{2c(a+bx)})^4 \sqrt{\tanh^2(ac+bcx)}} + \frac{26e^{c(a+bx)} \tanh(ac+bcx)}{3bc(1-e^{2c(a+bx)})^3 \sqrt{\tanh^2(ac+bcx)}} - \frac{15e^{c(a+bx)} \tanh(ac+bcx)}{4bc \sqrt{\tanh^2(ac+bcx)}} - \frac{15 \operatorname{arctanh}(e^{c(a+bx)}) \tanh(ac+bcx)}{4bc \sqrt{\tanh^2(ac+bcx)}}$$

```
[Out] exp(c*(b*x+a))*tanh(b*c*x+a*c)/b/c/(tanh(b*c*x+a*c)^2)^(1/2)-4*exp(c*(b*x+a))
)*tanh(b*c*x+a*c)/b/c/(1-exp(2*c*(b*x+a)))^4/(tanh(b*c*x+a*c)^2)^(1/2)+26/
3*exp(c*(b*x+a))*tanh(b*c*x+a*c)/b/c/(1-exp(2*c*(b*x+a)))^3/(tanh(b*c*x+a*c)
)^2)^(1/2)-55/6*exp(c*(b*x+a))*tanh(b*c*x+a*c)/b/c/(1-exp(2*c*(b*x+a)))^2/(
tanh(b*c*x+a*c)^2)^(1/2)+25/4*exp(c*(b*x+a))*tanh(b*c*x+a*c)/b/c/(1-exp(2*c
*(b*x+a)))/(tanh(b*c*x+a*c)^2)^(1/2)-15/4*arctanh(exp(c*(b*x+a)))*tanh(b*c*
x+a*c)/b/c/(tanh(b*c*x+a*c)^2)^(1/2)
```

Rubi [A]

time = 1.22, antiderivative size = 319, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 7, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.280$, Rules used = {6852, 2320, 398, 1828, 1171, 393, 213}

$$\frac{e^{c(a+bx)} \tanh(ac+bcx)}{bc \sqrt{\tanh^2(ac+bcx)}} + \frac{25e^{c(a+bx)} \tanh(ac+bcx)}{4bc(1-e^{2c(a+bx)}) \sqrt{\tanh^2(ac+bcx)}} - \frac{55e^{c(a+bx)} \tanh(ac+bcx)}{6bc(1-e^{2c(a+bx)})^2 \sqrt{\tanh^2(ac+bcx)}} + \frac{26e^{c(a+bx)} \tanh(ac+bcx)}{3bc(1-e^{2c(a+bx)})^3 \sqrt{\tanh^2(ac+bcx)}} - \frac{4e^{c(a+bx)} \tanh(ac+bcx)}{bc(1-e^{2c(a+bx)})^4 \sqrt{\tanh^2(ac+bcx)}} - \frac{15 \operatorname{arctanh}(e^{c(a+bx)}) \tanh(ac+bcx)}{4bc \sqrt{\tanh^2(ac+bcx)}}$$

Antiderivative was successfully verified.

[In] Int[E^(c*(a + b*x))/(Tanh[a*c + b*c*x]^2)^(5/2), x]

```
[Out] (E^(c*(a + b*x))*Tanh[a*c + b*c*x])/(b*c*Sqrt[Tanh[a*c + b*c*x]^2]) - (4*E^
(c*(a + b*x))*Tanh[a*c + b*c*x])/(b*c*(1 - E^(2*c*(a + b*x)))^4*Sqrt[Tanh[a
*c + b*c*x]^2]) + (26*E^(c*(a + b*x))*Tanh[a*c + b*c*x])/(3*b*c*(1 - E^(2*c
*(a + b*x)))^3*Sqrt[Tanh[a*c + b*c*x]^2]) - (55*E^(c*(a + b*x))*Tanh[a*c +
b*c*x])/(6*b*c*(1 - E^(2*c*(a + b*x)))^2*Sqrt[Tanh[a*c + b*c*x]^2]) + (25*E
^(c*(a + b*x))*Tanh[a*c + b*c*x])/(4*b*c*(1 - E^(2*c*(a + b*x)))*Sqrt[Tanh[
a*c + b*c*x]^2]) - (15*ArcTanh[E^(c*(a + b*x))]*Tanh[a*c + b*c*x])/(4*b*c*S
qrt[Tanh[a*c + b*c*x]^2])
```

Rule 213

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-
1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] &&
(LtQ[a, 0] || GtQ[b, 0])
```

Rule 393

```
Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Si
mp[(-(b*c - a*d))*x*((a + b*x^n)^(p + 1)/(a*b*n*(p + 1))), x] - Dist[(a*d -
b*c*(n*(p + 1) + 1))/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; F
```

```
reeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && (LtQ[p, -1] || ILtQ[1/n
+ p, 0])
```

Rule 398

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:> Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a
, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q,
0] && GeQ[p, -q]
```

Rule 1171

```
Int[((d_) + (e_.)*(x_)^2)^(q_)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_.),
x_Symbol] :> With[{Qx = PolynomialQuotient[(a + b*x^2 + c*x^4)^p, d + e*x^2,
x], R = Coeff[PolynomialRemainder[(a + b*x^2 + c*x^4)^p, d + e*x^2, x], x,
0]}, Simp[(-R)*x*((d + e*x^2)^(q + 1)/(2*d*(q + 1))), x] + Dist[1/(2*d*(q
+ 1)), Int[(d + e*x^2)^(q + 1)*ExpandToSum[2*d*(q + 1)*Qx + R*(2*q + 3), x
], x], x]] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2
- b*d*e + a*e^2, 0] && IGtQ[p, 0] && LtQ[q, -1]
```

Rule 1828

```
Int[(Pq)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :> With[{Q = PolynomialQuot
ient[Pq, a + b*x^2, x], f = Coeff[PolynomialRemainder[Pq, a + b*x^2, x], x,
0], g = Coeff[PolynomialRemainder[Pq, a + b*x^2, x], x, 1]}, Simp[(a*g - b
*f*x)*((a + b*x^2)^(p + 1)/(2*a*b*(p + 1))), x] + Dist[1/(2*a*(p + 1)), Int
[(a + b*x^2)^(p + 1)*ExpandToSum[2*a*(p + 1)*Q + f*(2*p + 3), x], x], x]] /
; FreeQ[{a, b}, x] && PolyQ[Pq, x] && LtQ[p, -1]
```

Rule 2320

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x]
, Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Functi
onOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ[
{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*
(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 6852

```
Int[(u_.)*((a_.)*(v_)^(m_.))^(p_), x_Symbol] :> Dist[a^IntPart[p]*((a*v^m)^
FracPart[p]/v^(m*FracPart[p])), Int[u*v^(m*p), x], x] /; FreeQ[{a, m, p}, x
] && !IntegerQ[p] && !FreeQ[v, x] && !(EqQ[a, 1] && EqQ[m, 1]) && !(EqQ
[v, x] && EqQ[m, 1])
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{c(a+bx)}}{\tanh^2(ac+bcx)^{5/2}} dx &= \frac{\tanh(ac+bcx) \int e^{c(a+bx)} \coth^5(ac+bcx) dx}{\sqrt{\tanh^2(ac+bcx)}} \\
&= \frac{\tanh(ac+bcx) \text{Subst}\left(\int \frac{(1+x^2)^5}{(-1+x^2)^5} dx, x, e^{c(a+bx)}\right)}{bc\sqrt{\tanh^2(ac+bcx)}} \\
&= \frac{\tanh(ac+bcx) \text{Subst}\left(\int \left(1 + \frac{2(1+10x^4+5x^8)}{(-1+x^2)^5}\right) dx, x, e^{c(a+bx)}\right)}{bc\sqrt{\tanh^2(ac+bcx)}} \\
&= \frac{e^{c(a+bx)} \tanh(ac+bcx)}{bc\sqrt{\tanh^2(ac+bcx)}} + \frac{(2 \tanh(ac+bcx)) \text{Subst}\left(\int \frac{1+10x^4+5x^8}{(-1+x^2)^5} dx, x, e^{c(a+bx)}\right)}{bc\sqrt{\tanh^2(ac+bcx)}} \\
&= \frac{e^{c(a+bx)} \tanh(ac+bcx)}{bc\sqrt{\tanh^2(ac+bcx)}} - \frac{4e^{c(a+bx)} \tanh(ac+bcx)}{bc(1-e^{2c(a+bx)})^4 \sqrt{\tanh^2(ac+bcx)}} + \frac{\tanh(ac+bcx)}{3bc(1-e^{2c(a+bx)})} \\
&= \frac{e^{c(a+bx)} \tanh(ac+bcx)}{bc\sqrt{\tanh^2(ac+bcx)}} - \frac{4e^{c(a+bx)} \tanh(ac+bcx)}{bc(1-e^{2c(a+bx)})^4 \sqrt{\tanh^2(ac+bcx)}} + \frac{26e^{c(a+bx)}}{3bc(1-e^{2c(a+bx)})} \\
&= \frac{e^{c(a+bx)} \tanh(ac+bcx)}{bc\sqrt{\tanh^2(ac+bcx)}} - \frac{4e^{c(a+bx)} \tanh(ac+bcx)}{bc(1-e^{2c(a+bx)})^4 \sqrt{\tanh^2(ac+bcx)}} + \frac{26e^{c(a+bx)}}{3bc(1-e^{2c(a+bx)})} \\
&= \frac{e^{c(a+bx)} \tanh(ac+bcx)}{bc\sqrt{\tanh^2(ac+bcx)}} - \frac{4e^{c(a+bx)} \tanh(ac+bcx)}{bc(1-e^{2c(a+bx)})^4 \sqrt{\tanh^2(ac+bcx)}} + \frac{26e^{c(a+bx)}}{3bc(1-e^{2c(a+bx)})} \\
&= \frac{e^{c(a+bx)} \tanh(ac+bcx)}{bc\sqrt{\tanh^2(ac+bcx)}} - \frac{4e^{c(a+bx)} \tanh(ac+bcx)}{bc(1-e^{2c(a+bx)})^4 \sqrt{\tanh^2(ac+bcx)}} + \frac{26e^{c(a+bx)}}{3bc(1-e^{2c(a+bx)})}
\end{aligned}$$

Mathematica [A]

time = 10.48, size = 164, normalized size = 0.51

$$\frac{(66e^{c(a+bx)} - 314e^{3c(a+bx)} + 374e^{5c(a+bx)} - 246e^{7c(a+bx)} + 24e^{9c(a+bx)} + 45(-1 + e^{2c(a+bx)})^4 \log(1 - e^{c(a+bx)}) - 45(-1 + e^{2c(a+bx)})^4 \log(1 + e^{c(a+bx)})) \tanh(c(a+bx))}{24bc(-1 + e^{2c(a+bx)})^4 \sqrt{\tanh^2(c(a+bx))}}$$

Antiderivative was successfully verified.

[In] Integrate[E^(c*(a + b*x))/(Tanh[a*c + b*c*x]^2)^(5/2), x]

[Out] ((66*E^(c*(a + b*x)) - 314*E^(3*c*(a + b*x)) + 374*E^(5*c*(a + b*x)) - 246*E^(7*c*(a + b*x)) + 24*E^(9*c*(a + b*x)) + 45*(-1 + E^(2*c*(a + b*x)))^4*Lo

$g[1 - E^{(c*(a + b*x))}] - 45*(-1 + E^{(2*c*(a + b*x)))^4*Log[1 + E^{(c*(a + b*x))}]*Tanh[c*(a + b*x)]/(24*b*c*(-1 + E^{(2*c*(a + b*x)))^4*sqrt[Tanh[c*(a + b*x)]^2])$

Maple [A]

time = 6.55, size = 320, normalized size = 1.00

method	result
risch	$\frac{(e^{2c(bx+a)}-1)e^{c(bx+a)}}{\sqrt{\frac{(e^{2c(bx+a)}-1)^2}{(1+e^{2c(bx+a)})^2}}(1+e^{2c(bx+a)})bc} - \frac{e^{c(bx+a)}(75e^{6c(bx+a)}-115e^{4c(bx+a)}+109e^{2c(bx+a)}-21)}{12(e^{2c(bx+a)}-1)^3(1+e^{2c(bx+a)})\sqrt{\frac{(e^{2c(bx+a)}-1)^2}{(1+e^{2c(bx+a)})^2}}bc} + \frac{15(e^{2c(bx+a)}-1)\ln(e^{c(bx+a)})}{8\sqrt{\frac{(e^{2c(bx+a)}-1)^2}{(1+e^{2c(bx+a)})^2}}(1+e^{2c(bx+a)})}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(c*(b*x+a))/(tanh(b*c*x+a*c)^2)^(5/2),x,method=_RETURNVERBOSE)`

[Out] $1/((\exp(2*c*(b*x+a))-1)^2/(1+\exp(2*c*(b*x+a)))^2)^{(1/2)}/(1+\exp(2*c*(b*x+a)))*(\exp(2*c*(b*x+a))-1)*\exp(c*(b*x+a))/b/c-1/12/(\exp(2*c*(b*x+a))-1)^3/(1+\exp(2*c*(b*x+a)))/((\exp(2*c*(b*x+a))-1)^2/(1+\exp(2*c*(b*x+a)))^2)^{(1/2)}*\exp(c*(b*x+a))*(75*\exp(6*c*(b*x+a))-115*\exp(4*c*(b*x+a))+109*\exp(2*c*(b*x+a))-21)/b/c+15/8/((\exp(2*c*(b*x+a))-1)^2/(1+\exp(2*c*(b*x+a)))^2)^{(1/2)}/(1+\exp(2*c*(b*x+a)))*(\exp(2*c*(b*x+a))-1)/b/c*\ln(\exp(c*(b*x+a))-1)-15/8/((\exp(2*c*(b*x+a))-1)^2/(1+\exp(2*c*(b*x+a)))^2)^{(1/2)}/(1+\exp(2*c*(b*x+a)))*(\exp(2*c*(b*x+a))-1)/b/c*\ln(\exp(c*(b*x+a))+1)$

Maxima [A]

time = 0.51, size = 167, normalized size = 0.52

$$-\frac{15 \log(e^{(bcx+ac)} + 1)}{8bc} + \frac{15 \log(e^{(bcx+ac)} - 1)}{8bc} + \frac{12e^{(9bcx+9ac)} - 123e^{(7bcx+7ac)} + 187e^{(5bcx+5ac)} - 157e^{(3bcx+3ac)} + 33e^{(bcx+ac)}}{12bc(e^{(8bcx+8ac)} - 4e^{(6bcx+6ac)} + 6e^{(4bcx+4ac)} - 4e^{(2bcx+2ac)} + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(c*(b*x+a))/(tanh(b*c*x+a*c)^2)^(5/2),x, algorithm="maxima")`

[Out] $-15/8*\log(e^{(b*c*x + a*c)} + 1)/(b*c) + 15/8*\log(e^{(b*c*x + a*c)} - 1)/(b*c) + 1/12*(12*e^{(9*b*c*x + 9*a*c)} - 123*e^{(7*b*c*x + 7*a*c)} + 187*e^{(5*b*c*x + 5*a*c)} - 157*e^{(3*b*c*x + 3*a*c)} + 33*e^{(b*c*x + a*c)})/(b*c*(e^{(8*b*c*x + 8*a*c)} - 4*e^{(6*b*c*x + 6*a*c)} + 6*e^{(4*b*c*x + 4*a*c)} - 4*e^{(2*b*c*x + 2*a*c)} + 1))$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 1617 vs. $2(281) = 562$.

time = 0.37, size = 1617, normalized size = 5.07

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(c*(b*x+a))/(tanh(b*c*x+a*c)^2)^(5/2),x, algorithm="fricas")
[Out] 1/24*(24*cosh(b*c*x + a*c)^9 + 216*cosh(b*c*x + a*c)*sinh(b*c*x + a*c)^8 +
24*sinh(b*c*x + a*c)^9 + 6*(144*cosh(b*c*x + a*c)^2 - 41)*sinh(b*c*x + a*c)
^7 - 246*cosh(b*c*x + a*c)^7 + 42*(48*cosh(b*c*x + a*c)^3 - 41*cosh(b*c*x +
a*c))*sinh(b*c*x + a*c)^6 + 2*(1512*cosh(b*c*x + a*c)^4 - 2583*cosh(b*c*x
+ a*c)^2 + 187)*sinh(b*c*x + a*c)^5 + 374*cosh(b*c*x + a*c)^5 + 2*(1512*cos
h(b*c*x + a*c)^5 - 4305*cosh(b*c*x + a*c)^3 + 935*cosh(b*c*x + a*c))*sinh(b
*c*x + a*c)^4 + 2*(1008*cosh(b*c*x + a*c)^6 - 4305*cosh(b*c*x + a*c)^4 + 18
70*cosh(b*c*x + a*c)^2 - 157)*sinh(b*c*x + a*c)^3 - 314*cosh(b*c*x + a*c)^3
+ 2*(432*cosh(b*c*x + a*c)^7 - 2583*cosh(b*c*x + a*c)^5 + 1870*cosh(b*c*x
+ a*c)^3 - 471*cosh(b*c*x + a*c))*sinh(b*c*x + a*c)^2 - 45*(cosh(b*c*x + a*
c)^8 + 8*cosh(b*c*x + a*c)*sinh(b*c*x + a*c)^7 + sinh(b*c*x + a*c)^8 + 4*(7
*cosh(b*c*x + a*c)^2 - 1)*sinh(b*c*x + a*c)^6 - 4*cosh(b*c*x + a*c)^6 + 8*(
7*cosh(b*c*x + a*c)^3 - 3*cosh(b*c*x + a*c))*sinh(b*c*x + a*c)^5 + 2*(35*co
sh(b*c*x + a*c)^4 - 30*cosh(b*c*x + a*c)^2 + 3)*sinh(b*c*x + a*c)^4 + 6*cos
h(b*c*x + a*c)^4 + 8*(7*cosh(b*c*x + a*c)^5 - 10*cosh(b*c*x + a*c)^3 + 3*co
sh(b*c*x + a*c))*sinh(b*c*x + a*c)^3 + 4*(7*cosh(b*c*x + a*c)^6 - 15*cosh(b
*c*x + a*c)^4 + 9*cosh(b*c*x + a*c)^2 - 1)*sinh(b*c*x + a*c)^2 - 4*cosh(b*c
*x + a*c)^2 + 8*(cosh(b*c*x + a*c)^7 - 3*cosh(b*c*x + a*c)^5 + 3*cosh(b*c*x
+ a*c)^3 - cosh(b*c*x + a*c))*sinh(b*c*x + a*c) + 1)*log(cosh(b*c*x + a*c)
+ sinh(b*c*x + a*c) + 1) + 45*(cosh(b*c*x + a*c)^8 + 8*cosh(b*c*x + a*c)*s
inh(b*c*x + a*c)^7 + sinh(b*c*x + a*c)^8 + 4*(7*cosh(b*c*x + a*c)^2 - 1)*si
nh(b*c*x + a*c)^6 - 4*cosh(b*c*x + a*c)^6 + 8*(7*cosh(b*c*x + a*c)^3 - 3*co
sh(b*c*x + a*c))*sinh(b*c*x + a*c)^5 + 2*(35*cosh(b*c*x + a*c)^4 - 30*cosh(
b*c*x + a*c)^2 + 3)*sinh(b*c*x + a*c)^4 + 6*cosh(b*c*x + a*c)^4 + 8*(7*cosh
(b*c*x + a*c)^5 - 10*cosh(b*c*x + a*c)^3 + 3*cosh(b*c*x + a*c))*sinh(b*c*x
+ a*c)^3 + 4*(7*cosh(b*c*x + a*c)^6 - 15*cosh(b*c*x + a*c)^4 + 9*cosh(b*c*x
+ a*c)^2 - 1)*sinh(b*c*x + a*c)^2 - 4*cosh(b*c*x + a*c)^2 + 8*(cosh(b*c*x
+ a*c)^7 - 3*cosh(b*c*x + a*c)^5 + 3*cosh(b*c*x + a*c)^3 - cosh(b*c*x + a*c
))*sinh(b*c*x + a*c) + 1)*log(cosh(b*c*x + a*c) + sinh(b*c*x + a*c) - 1) +
2*(108*cosh(b*c*x + a*c)^8 - 861*cosh(b*c*x + a*c)^6 + 935*cosh(b*c*x + a*c
)^4 - 471*cosh(b*c*x + a*c)^2 + 33)*sinh(b*c*x + a*c) + 66*cosh(b*c*x + a*c
))/(b*c*cosh(b*c*x + a*c)^8 + 8*b*c*cosh(b*c*x + a*c)*sinh(b*c*x + a*c)^7 +
b*c*sinh(b*c*x + a*c)^8 - 4*b*c*cosh(b*c*x + a*c)^6 + 4*(7*b*c*cosh(b*c*x
+ a*c)^2 - b*c)*sinh(b*c*x + a*c)^6 + 6*b*c*cosh(b*c*x + a*c)^4 + 8*(7*b*c*
cosh(b*c*x + a*c)^3 - 3*b*c*cosh(b*c*x + a*c))*sinh(b*c*x + a*c)^5 + 2*(35*
b*c*cosh(b*c*x + a*c)^4 - 30*b*c*cosh(b*c*x + a*c)^2 + 3*b*c)*sinh(b*c*x +
a*c)^4 - 4*b*c*cosh(b*c*x + a*c)^2 + 8*(7*b*c*cosh(b*c*x + a*c)^5 - 10*b*c*
cosh(b*c*x + a*c)^3 + 3*b*c*cosh(b*c*x + a*c))*sinh(b*c*x + a*c)^3 + 4*(7*b
*c*cosh(b*c*x + a*c)^6 - 15*b*c*cosh(b*c*x + a*c)^4 + 9*b*c*cosh(b*c*x + a*
c)^2 - b*c)*sinh(b*c*x + a*c)^2 + b*c + 8*(b*c*cosh(b*c*x + a*c)^7 - 3*b*c*
cosh(b*c*x + a*c)^5 + 3*b*c*cosh(b*c*x + a*c)^3 - b*c*cosh(b*c*x + a*c))*si
nh(b*c*x + a*c))
```

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(c*(b*x+a))/(tanh(b*c*x+a*c)**2)**(5/2), x)

[Out] Timed out

Giac [A]

time = 0.52, size = 215, normalized size = 0.67

$$\frac{24 e^{(bcx+ac)} \operatorname{sgn}(e^{(2bcx+2ac)} - 1) - 45 \log(e^{(bcx+ac)} + 1) \operatorname{sgn}(e^{(2bcx+2ac)} - 1) + 45 \log(|e^{(bcx+ac)} - 1|) \operatorname{sgn}(e^{(2bcx+2ac)} - 1) - \frac{2(75e^{(7bcx+7ac)} \operatorname{sgn}(e^{(2bcx+2ac)} - 1) - 115e^{(5bcx+5ac)} \operatorname{sgn}(e^{(2bcx+2ac)} - 1) + 109e^{(3bcx+3ac)} \operatorname{sgn}(e^{(2bcx+2ac)} - 1) - 21e^{(bcx+ac)} \operatorname{sgn}(e^{(2bcx+2ac)} - 1))}{(e^{(2bcx+2ac)} - 1)^4}}{24bc}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(c*(b*x+a))/(tanh(b*c*x+a*c)^2)^(5/2), x, algorithm="giac")

[Out] 1/24*(24*e^(b*c*x + a*c)*sgn(e^(2*b*c*x + 2*a*c) - 1) - 45*log(e^(b*c*x + a*c) + 1)*sgn(e^(2*b*c*x + 2*a*c) - 1) + 45*log(abs(e^(b*c*x + a*c) - 1))*sgn(e^(2*b*c*x + 2*a*c) - 1) - 2*(75*e^(7*b*c*x + 7*a*c)*sgn(e^(2*b*c*x + 2*a*c) - 1) - 115*e^(5*b*c*x + 5*a*c)*sgn(e^(2*b*c*x + 2*a*c) - 1) + 109*e^(3*b*c*x + 3*a*c)*sgn(e^(2*b*c*x + 2*a*c) - 1) - 21*e^(b*c*x + a*c)*sgn(e^(2*b*c*x + 2*a*c) - 1))/(e^(2*b*c*x + 2*a*c) - 1)^4/(b*c)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{e^{c(a+bx)}}{(\tanh(ac+bcx)^2)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(c*(a + b*x))/(tanh(a*c + b*c*x)^2)^(5/2), x)

[Out] int(exp(c*(a + b*x))/(tanh(a*c + b*c*x)^2)^(5/2), x)

3.240 $\int \sin^3(\tanh(a + bx)) dx$

Optimal. Leaf size=157

$$\frac{3\text{CosIntegral}(1 - \tanh(a + bx)) \sin(1)}{8b} - \frac{3\text{CosIntegral}(1 + \tanh(a + bx)) \sin(1)}{8b} + \frac{\text{CosIntegral}(3 - 3 \tanh(a + bx)) \sin(3)}{8b}$$

[Out] 1/8*cos(3)*Si(-3+3*tanh(b*x+a))/b-3/8*cos(1)*Si(-1+tanh(b*x+a))/b+3/8*cos(1)*Si(1+tanh(b*x+a))/b-1/8*cos(3)*Si(3+3*tanh(b*x+a))/b-3/8*Ci(1-tanh(b*x+a))*sin(1)/b-3/8*Ci(1+tanh(b*x+a))*sin(1)/b+1/8*Ci(3-3*tanh(b*x+a))*sin(3)/b+1/8*Ci(3+3*tanh(b*x+a))*sin(3)/b

Rubi [A]

time = 0.28, antiderivative size = 157, normalized size of antiderivative = 1.00, number of steps used = 19, number of rules used = 5, integrand size = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.556$, Rules used = {6857, 3393, 3384, 3380, 3383}

$$\frac{\sin(3)\text{CosIntegral}(3 - 3 \tanh(a + bx))}{8b} + \frac{\sin(3)\text{CosIntegral}(3 \tanh(a + bx) + 3)}{8b} - \frac{3 \sin(1)\text{CosIntegral}(1 - \tanh(a + bx))}{8b} - \frac{3 \sin(1)\text{CosIntegral}(\tanh(a + bx) + 1)}{8b} - \frac{\cos(3)\text{Si}(3 - 3 \tanh(a + bx))}{8b} + \frac{3 \cos(1)\text{Si}(1 - \tanh(a + bx))}{8b} + \frac{3 \cos(1)\text{Si}(\tanh(a + bx) + 1)}{8b} - \frac{\cos(3)\text{Si}(3 \tanh(a + bx) + 3)}{8b}$$

Antiderivative was successfully verified.

[In] Int[Sin[Tanh[a + b*x]]^3,x]

[Out] (-3*CosIntegral[1 - Tanh[a + b*x]]*Sin[1])/(8*b) - (3*CosIntegral[1 + Tanh[a + b*x]]*Sin[1])/(8*b) + (CosIntegral[3 - 3*Tanh[a + b*x]]*Sin[3])/(8*b) + (CosIntegral[3 + 3*Tanh[a + b*x]]*Sin[3])/(8*b) - (Cos[3]*SinIntegral[3 - 3*Tanh[a + b*x]])/(8*b) + (3*Cos[1]*SinIntegral[1 - Tanh[a + b*x]])/(8*b) + (3*Cos[1]*SinIntegral[1 + Tanh[a + b*x]])/(8*b) - (Cos[3]*SinIntegral[3 + 3*Tanh[a + b*x]])/(8*b)

Rule 3380

Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] :> Simp[SinIntegral[e + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*e - c*f, 0]

Rule 3383

Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] :> Simp[CosIntegral[e - Pi/2 + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*(e - Pi/2) - c*f, 0]

Rule 3384

Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] :> Dist[Cos[(d*e - c*f)/d], Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Dist[Sin[(d*e - c*f)/d], Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]

Rule 3393

```
Int[((c_.) + (d_.)*(x_)^(m_))*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int
[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f
, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))
```

Rule 6857

```
Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
xpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]
```

Rubi steps

$$\begin{aligned}
\int \sin^3(\tanh(a + bx)) dx &= \frac{\text{Subst}\left(\int \frac{\sin^3(x)}{1-x^2} dx, x, \tanh(a + bx)\right)}{b} \\
&= \frac{\text{Subst}\left(\int \left(-\frac{\sin^3(x)}{2(-1+x)} + \frac{\sin^3(x)}{2(1+x)}\right) dx, x, \tanh(a + bx)\right)}{b} \\
&= -\frac{\text{Subst}\left(\int \frac{\sin^3(x)}{-1+x} dx, x, \tanh(a + bx)\right)}{2b} + \frac{\text{Subst}\left(\int \frac{\sin^3(x)}{1+x} dx, x, \tanh(a + bx)\right)}{2b} \\
&= -\frac{\text{Subst}\left(\int \left(\frac{3 \sin(x)}{4(-1+x)} - \frac{\sin(3x)}{4(-1+x)}\right) dx, x, \tanh(a + bx)\right)}{2b} + \frac{\text{Subst}\left(\int \left(\frac{3 \sin(x)}{4(1+x)} - \frac{\sin(3x)}{4(1+x)}\right) dx, x, \tanh(a + bx)\right)}{2b} \\
&= -\frac{\text{Subst}\left(\int \frac{\sin(3x)}{-1+x} dx, x, \tanh(a + bx)\right)}{8b} - \frac{\text{Subst}\left(\int \frac{\sin(3x)}{1+x} dx, x, \tanh(a + bx)\right)}{8b} - \frac{3\text{Si}(1 - \tanh(a + bx)) \sin(1)}{8b} - \frac{3\text{Si}(1 + \tanh(a + bx)) \sin(1)}{8b} + \frac{\text{Ci}(3 - 3 \tanh(a + bx)) \sin(1)}{8b} + \frac{\text{Ci}(3 + 3 \tanh(a + bx)) \sin(1)}{8b}
\end{aligned}$$

Mathematica [A]

time = 0.17, size = 124, normalized size = 0.79

$-6\text{CosIntegral}[1 - \tanh(a + bx)] \sin(1) - 6\text{CosIntegral}[1 + \tanh(a + bx)] \sin(1) + 2\text{CosIntegral}[3 - 3 \tanh(a + bx)] \sin(3) + 2\text{CosIntegral}[3 + 3 \tanh(a + bx)] \sin(3) - 2\cos(3)\text{Si}(3 - 3 \tanh(a + bx)) + 6\cos(3)\text{Si}(1 - \tanh(a + bx)) + 6\cos(3)\text{Si}(1 + \tanh(a + bx)) - 2\cos(3)\text{Si}(3 + 3 \tanh(a + bx))$

Antiderivative was successfully verified.

```
[In] Integrate[Sin[Tanh[a + b*x]]^3, x]
```

```
[Out] (-6*CosIntegral[1 - Tanh[a + b*x]]*Sin[1] - 6*CosIntegral[1 + Tanh[a + b*x]]*Sin[1] + 2*CosIntegral[3 - 3*Tanh[a + b*x]]*Sin[3] + 2*CosIntegral[3 + 3*
```


$\text{Tanh}[a + b*x]]*\text{Sin}[3] - 2*\text{Cos}[3]*\text{SinIntegral}[3 - 3*\text{Tanh}[a + b*x]] + 6*\text{Cos}[1]*\text{SinIntegral}[1 - \text{Tanh}[a + b*x]] + 6*\text{Cos}[1]*\text{SinIntegral}[1 + \text{Tanh}[a + b*x]] - 2*\text{Cos}[3]*\text{SinIntegral}[3 + 3*\text{Tanh}[a + b*x]])/(16*b)$

Maple [A]

time = 1.29, size = 118, normalized size = 0.75

method	result
derivativedivides	$\frac{\text{sinIntegral}(-3+3 \tanh(bx+a)) \cos(3)}{8} + \frac{\text{cosineIntegral}(-3+3 \tanh(bx+a)) \sin(3)}{8} - \frac{\text{sinIntegral}(3+3 \tanh(bx+a)) \cos(3)}{8} + \frac{\text{cosineIntegral}(3+3 \tanh(bx+a)) \sin(3)}{8}$
default	$\frac{\text{sinIntegral}(-3+3 \tanh(bx+a)) \cos(3)}{8} + \frac{\text{cosineIntegral}(-3+3 \tanh(bx+a)) \sin(3)}{8} - \frac{\text{sinIntegral}(3+3 \tanh(bx+a)) \cos(3)}{8} + \frac{\text{cosineIntegral}(3+3 \tanh(bx+a)) \sin(3)}{8}$
risch	$\frac{ie^{3i} \exp\text{Integral}\left(1, \frac{6i}{e^{2bx+2a}+1}\right)}{16b} - \frac{ie^{-3i} \exp\text{Integral}\left(1, \frac{6i}{e^{2bx+2a}+1} - 6i\right)}{16b} - \frac{ie^{-3i} \exp\text{Integral}\left(1, -\frac{6i}{e^{2bx+2a}+1}\right)}{16b} + \dots$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(tanh(b*x+a))^3,x,method=_RETURNVERBOSE)`

[Out] $1/b*(1/8*\text{Si}(-3+3*\tanh(b*x+a))*\cos(3)+1/8*\text{Ci}(-3+3*\tanh(b*x+a))*\sin(3)-1/8*\text{Si}(3+3*\tanh(b*x+a))*\cos(3)+1/8*\text{Ci}(3+3*\tanh(b*x+a))*\sin(3)-3/8*\text{Si}(-1+\tanh(b*x+a))*\cos(1)-3/8*\text{Ci}(-1+\tanh(b*x+a))*\sin(1)+3/8*\text{Si}(\tanh(b*x+a)+1)*\cos(1)-3/8*\text{Ci}(\tanh(b*x+a)+1)*\sin(1))$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(tanh(b*x+a))^3,x, algorithm="maxima")`

[Out] `integrate(sin(tanh(b*x + a))^3, x)`

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 296 vs. 2(139) = 278.

time = 0.38, size = 296, normalized size = 1.89

$$\frac{\text{Ci}\left(\frac{6i(2bx+a)}{e^{2bx+2a}+1}\right) \sin(3) + \text{Ci}\left(-\frac{6i(2bx+a)}{e^{2bx+2a}+1}\right) \sin(3) + \text{Ci}\left(\frac{6}{e^{2bx+2a}+1}\right) \sin(3) + \text{Ci}\left(-\frac{6}{e^{2bx+2a}+1}\right) \sin(3) - 3 \text{Ci}\left(\frac{2i(2bx+a)}{e^{2bx+2a}+1}\right) \sin(1) - 3 \text{Ci}\left(-\frac{2i(2bx+a)}{e^{2bx+2a}+1}\right) \sin(1) - 3 \text{Ci}\left(\frac{1}{e^{2bx+2a}+1}\right) \sin(1) - 3 \text{Ci}\left(-\frac{1}{e^{2bx+2a}+1}\right) \sin(1) - 2 \cos(3) \text{Si}\left(\frac{6i(2bx+a)}{e^{2bx+2a}+1}\right) + 6 \cos(1) \text{Si}\left(\frac{2i(2bx+a)}{e^{2bx+2a}+1}\right) - 2 \cos(3) \text{Si}\left(\frac{6}{e^{2bx+2a}+1}\right) + 6 \cos(1) \text{Si}\left(\frac{1}{e^{2bx+2a}+1}\right)}{16b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(tanh(b*x+a))^3,x, algorithm="fricas")`

[Out] $1/16*(\text{cos_integral}(6*e^{(2*b*x + 2*a)})/(e^{(2*b*x + 2*a)} + 1))*\sin(3) + \text{cos_integral}(-6*e^{(2*b*x + 2*a)})/(e^{(2*b*x + 2*a)} + 1))*\sin(3) + \text{cos_integral}(6/(e^{(2*b*x + 2*a)} + 1))*\sin(3) + \text{cos_integral}(-6/(e^{(2*b*x + 2*a)} + 1))*\sin(3) - 3*\text{cos_integral}(2*e^{(2*b*x + 2*a)})/(e^{(2*b*x + 2*a)} + 1))*\sin(1) - 3*\text{cos_i}$

```

ntegral(-2*e^(2*b*x + 2*a)/(e^(2*b*x + 2*a) + 1))*sin(1) - 3*cos_integral(2
/(e^(2*b*x + 2*a) + 1))*sin(1) - 3*cos_integral(-2/(e^(2*b*x + 2*a) + 1))*s
in(1) - 2*cos(3)*sin_integral(6*e^(2*b*x + 2*a)/(e^(2*b*x + 2*a) + 1)) + 6*
cos(1)*sin_integral(2*e^(2*b*x + 2*a)/(e^(2*b*x + 2*a) + 1)) - 2*cos(3)*sin
_integral(6/(e^(2*b*x + 2*a) + 1)) + 6*cos(1)*sin_integral(2/(e^(2*b*x + 2*
a) + 1)))/b

```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sin^3(\tanh(a + bx)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(tanh(b*x+a))**3,x)
```

```
[Out] Integral(sin(tanh(a + b*x))**3, x)
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(tanh(b*x+a))^3,x, algorithm="giac")
```

```
[Out] integrate(sin(tanh(b*x + a))^3, x)
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \sin(\tanh(a + bx))^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sin(tanh(a + b*x))^3,x)
```

```
[Out] int(sin(tanh(a + b*x))^3, x)
```

3.241 $\int \sin^2(\tanh(a + bx)) dx$

Optimal. Leaf size=115

$$\frac{\cos(2)\text{CosIntegral}(2 - 2 \tanh(a + bx))}{4b} - \frac{\cos(2)\text{CosIntegral}(2 + 2 \tanh(a + bx))}{4b} - \frac{\log(1 - \tanh(a + bx))}{4b} + \frac{\log(1 + \tanh(a + bx))}{4b}$$

[Out] 1/4*Ci(2-2*tanh(b*x+a))*cos(2)/b-1/4*Ci(2+2*tanh(b*x+a))*cos(2)/b-1/4*ln(1-tanh(b*x+a))/b+1/4*ln(1+tanh(b*x+a))/b-1/4*Si(-2+2*tanh(b*x+a))*sin(2)/b-1/4*Si(2+2*tanh(b*x+a))*sin(2)/b

Rubi [A]

time = 0.19, antiderivative size = 115, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 5, integrand size = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.556$, Rules used = {6857, 3393, 3384, 3380, 3383}

$$\frac{\cos(2)\text{CosIntegral}(2 - 2 \tanh(a + bx))}{4b} - \frac{\cos(2)\text{CosIntegral}(2 \tanh(a + bx) + 2)}{4b} + \frac{\sin(2)\text{Si}(2 - 2 \tanh(a + bx))}{4b} - \frac{\sin(2)\text{Si}(2 \tanh(a + bx) + 2)}{4b} - \frac{\log(1 - \tanh(a + bx))}{4b} + \frac{\log(\tanh(a + bx) + 1)}{4b}$$

Antiderivative was successfully verified.

[In] Int[Sin[Tanh[a + b*x]]^2,x]

[Out] (Cos[2]*CosIntegral[2 - 2*Tanh[a + b*x]])/(4*b) - (Cos[2]*CosIntegral[2 + 2*Tanh[a + b*x]])/(4*b) - Log[1 - Tanh[a + b*x]]/(4*b) + Log[1 + Tanh[a + b*x]]/(4*b) + (Sin[2]*SinIntegral[2 - 2*Tanh[a + b*x]])/(4*b) - (Sin[2]*SinIntegral[2 + 2*Tanh[a + b*x]])/(4*b)

Rule 3380

Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] :> Simp[SinIntegral[e + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*e - c*f, 0]

Rule 3383

Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] :> Simp[CosIntegral[e - Pi/2 + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*(e - Pi/2) - c*f, 0]

Rule 3384

Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] :> Dist[Cos[(d*e - c*f)/d], Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Dist[Sin[(d*e - c*f)/d], Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]

Rule 3393

Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] :> Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f}

, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))

Rule 6857

Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
xpan[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]

Rubi steps

$$\begin{aligned}
 \int \sin^2(\tanh(a + bx)) dx &= \frac{\text{Subst}\left(\int \frac{\sin^2(x)}{1-x^2} dx, x, \tanh(a + bx)\right)}{b} \\
 &= \frac{\text{Subst}\left(\int \left(-\frac{\sin^2(x)}{2(-1+x)} + \frac{\sin^2(x)}{2(1+x)}\right) dx, x, \tanh(a + bx)\right)}{b} \\
 &= -\frac{\text{Subst}\left(\int \frac{\sin^2(x)}{-1+x} dx, x, \tanh(a + bx)\right)}{2b} + \frac{\text{Subst}\left(\int \frac{\sin^2(x)}{1+x} dx, x, \tanh(a + bx)\right)}{2b} \\
 &= -\frac{\text{Subst}\left(\int \left(\frac{1}{2(-1+x)} - \frac{\cos(2x)}{2(-1+x)}\right) dx, x, \tanh(a + bx)\right)}{2b} + \frac{\text{Subst}\left(\int \left(\frac{1}{2(1+x)} - \frac{\cos(2x)}{2(1+x)}\right) dx, x, \tanh(a + bx)\right)}{2b} \\
 &= -\frac{\log(1 - \tanh(a + bx))}{4b} + \frac{\log(1 + \tanh(a + bx))}{4b} + \frac{\text{Subst}\left(\int \frac{\cos(2x)}{-1+x} dx, x, \tanh(a + bx)\right)}{4b} \\
 &= -\frac{\log(1 - \tanh(a + bx))}{4b} + \frac{\log(1 + \tanh(a + bx))}{4b} + \frac{\cos(2)\text{Subst}\left(\int \frac{\cos(2-2x)}{-1+x} dx, x, \tanh(a + bx)\right)}{4b} \\
 &= \frac{\cos(2)\text{Ci}(2 - 2 \tanh(a + bx))}{4b} - \frac{\cos(2)\text{Ci}(2 + 2 \tanh(a + bx))}{4b} - \frac{\log(1 - \tanh(a + bx))}{4b} + \frac{\log(1 + \tanh(a + bx))}{4b}
 \end{aligned}$$

Mathematica [A]

time = 0.12, size = 88, normalized size = 0.77

$$\frac{\cos(2)\text{CosIntegral}(2 - 2 \tanh(a + bx)) - \cos(2)\text{CosIntegral}(2(1 + \tanh(a + bx))) - \log(1 - \tanh(a + bx)) + \log(1 + \tanh(a + bx)) + \sin(2)\text{Si}(2 - 2 \tanh(a + bx)) - \sin(2)\text{Si}(2(1 + \tanh(a + bx)))}{4b}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[Tanh[a + b*x]]^2,x]

[Out] (Cos[2]*CosIntegral[2 - 2*Tanh[a + b*x]] - Cos[2]*CosIntegral[2*(1 + Tanh[a + b*x])] - Log[1 - Tanh[a + b*x]] + Log[1 + Tanh[a + b*x]] + Sin[2]*SinIntegral[2 - 2*Tanh[a + b*x]] - Sin[2]*SinIntegral[2*(1 + Tanh[a + b*x])])/(4*b)

Maple [A]

time = 1.03, size = 88, normalized size = 0.77

method	result
derivativedivides	$\frac{-\frac{\ln(-1+\tanh(bx+a))}{4} + \frac{\ln(\tanh(bx+a)+1)}{4} - \frac{\operatorname{SiIntegral}(2+2\tanh(bx+a))\sin(2)}{4} - \frac{\operatorname{CosineIntegral}(2+2\tanh(bx+a))\cos(2)}{4} - \frac{\operatorname{SiIntegral}(2+2\tanh(bx+a))\cos(2)}{4}}{b}$
default	$\frac{-\frac{\ln(-1+\tanh(bx+a))}{4} + \frac{\ln(\tanh(bx+a)+1)}{4} - \frac{\operatorname{SiIntegral}(2+2\tanh(bx+a))\sin(2)}{4} - \frac{\operatorname{CosineIntegral}(2+2\tanh(bx+a))\cos(2)}{4} - \frac{\operatorname{SiIntegral}(2+2\tanh(bx+a))\cos(2)}{4}}{b}$
risch	$\frac{x}{2} - \frac{e^{-2i} \operatorname{ExpIntegralEi}\left(1, -\frac{4i}{e^{2bx+2a}+1}\right)}{8b} + \frac{e^{2i} \operatorname{ExpIntegralEi}\left(1, -\frac{4i}{e^{2bx+2a}+1} + 4i\right)}{8b} - \frac{e^{2i} \operatorname{ExpIntegralEi}\left(1, \frac{4i}{e^{2bx+2a}+1}\right)}{8b} + \dots$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(tanh(b*x+a))^2,x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{b} * (-1/4 * \ln(-1 + \tanh(b*x+a)) + 1/4 * \ln(\tanh(b*x+a) + 1) - 1/4 * \operatorname{Si}(2 + 2 * \tanh(b*x+a)) * \sin(2) - 1/4 * \operatorname{Ci}(2 + 2 * \tanh(b*x+a)) * \cos(2) - 1/4 * \operatorname{Si}(-2 + 2 * \tanh(b*x+a)) * \sin(2) + 1/4 * \operatorname{Ci}(-2 + 2 * \tanh(b*x+a)) * \cos(2))$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(tanh(b*x+a))^2,x, algorithm="maxima")`

[Out] $\frac{1}{2} * x - \frac{1}{2} * \operatorname{integrate}(\cos(2 * (e^{(2*b*x + 2*a)} - 1) / (e^{(2*b*x + 2*a)} + 1)), x)$

Fricas [A]

time = 0.37, size = 155, normalized size = 1.35

$$\frac{4bx - \cos(2) \operatorname{Ci}\left(\frac{4e^{(2bx+2a)}}{e^{(2bx+2a)}+1}\right) - \cos(2) \operatorname{Ci}\left(-\frac{4e^{(2bx+2a)}}{e^{(2bx+2a)}+1}\right) + \cos(2) \operatorname{Ci}\left(\frac{4}{e^{(2bx+2a)}+1}\right) + \cos(2) \operatorname{Ci}\left(-\frac{4}{e^{(2bx+2a)}+1}\right) - 2 \sin(2) \operatorname{Si}\left(\frac{4e^{(2bx+2a)}}{e^{(2bx+2a)}+1}\right) + 2 \sin(2) \operatorname{Si}\left(\frac{4}{e^{(2bx+2a)}+1}\right)}{8b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(tanh(b*x+a))^2,x, algorithm="fricas")`

[Out] $\frac{1}{8} * (4*b*x - \cos(2) * \operatorname{cos_integral}(4 * e^{(2*b*x + 2*a)} / (e^{(2*b*x + 2*a)} + 1)) - \cos(2) * \operatorname{cos_integral}(-4 * e^{(2*b*x + 2*a)} / (e^{(2*b*x + 2*a)} + 1)) + \cos(2) * \operatorname{cos_integral}(4 / (e^{(2*b*x + 2*a)} + 1)) + \cos(2) * \operatorname{cos_integral}(-4 / (e^{(2*b*x + 2*a)} + 1)) - 2 * \sin(2) * \operatorname{sin_integral}(4 * e^{(2*b*x + 2*a)} / (e^{(2*b*x + 2*a)} + 1)) + 2 * \sin(2) * \operatorname{sin_integral}(4 / (e^{(2*b*x + 2*a)} + 1))) / b$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sin^2(\tanh(a + bx)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(tanh(b*x+a))**2,x)

[Out] Integral(sin(tanh(a + b*x))**2, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(tanh(b*x+a))^2,x, algorithm="giac")

[Out] integrate(sin(tanh(b*x + a))^2, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \sin(\tanh(a + b x))^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(tanh(a + b*x))^2,x)

[Out] int(sin(tanh(a + b*x))^2, x)

3.242 $\int \sin(\tanh(a + bx)) dx$

Optimal. Leaf size=77

$$-\frac{\text{CosIntegral}(1 - \tanh(a + bx)) \sin(1)}{2b} - \frac{\text{CosIntegral}(1 + \tanh(a + bx)) \sin(1)}{2b} + \frac{\cos(1) \text{Si}(1 - \tanh(a + bx))}{2b}$$

[Out] $-1/2*\cos(1)*\text{Si}(-1+\tanh(b*x+a))/b+1/2*\cos(1)*\text{Si}(1+\tanh(b*x+a))/b-1/2*\text{Ci}(1-\tanh(b*x+a))*\sin(1)/b-1/2*\text{Ci}(1+\tanh(b*x+a))*\sin(1)/b$

Rubi [A]

time = 0.11, antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 4, integrand size = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$,

Rules used = {3414, 3384, 3380, 3383}

$$-\frac{\sin(1)\text{CosIntegral}(1 - \tanh(a + bx))}{2b} - \frac{\sin(1)\text{CosIntegral}(\tanh(a + bx) + 1)}{2b} + \frac{\cos(1)\text{Si}(1 - \tanh(a + bx))}{2b} + \frac{\cos(1)\text{Si}(\tanh(a + bx) + 1)}{2b}$$

Antiderivative was successfully verified.

[In] `Int[Sin[Tanh[a + b*x]],x]`

[Out] $-1/2*(\text{CosIntegral}[1 - \text{Tanh}[a + b*x]]*\text{Sin}[1])/b - (\text{CosIntegral}[1 + \text{Tanh}[a + b*x]]*\text{Sin}[1])/(2*b) + (\text{Cos}[1]*\text{SinIntegral}[1 - \text{Tanh}[a + b*x]])/(2*b) + (\text{Cos}[1]*\text{SinIntegral}[1 + \text{Tanh}[a + b*x]])/(2*b)$

Rule 3380

`Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[SinIntegral[e + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*e - c*f, 0]`

Rule 3383

`Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CosIntegral[e - Pi/2 + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*(e - Pi/2) - c*f, 0]`

Rule 3384

`Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Dist[Cos[(d*e - c*f)/d], Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Dist[Sin[(d*e - c*f)/d], Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]`

Rule 3414

`Int[((a_.) + (b_.)*(x_)^(n_))^(p_)*Sin[(c_.) + (d_.)*(x_)], x_Symbol] := Int[ExpandIntegrand[Sin[c + d*x], (a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[p, 0] && IGtQ[n, 0] && (EqQ[n, 2] || EqQ[p, -1])`

Rubi steps

$$\begin{aligned}
\int \sin(\tanh(a + bx)) dx &= \frac{\text{Subst}\left(\int \frac{\sin(x)}{1-x^2} dx, x, \tanh(a + bx)\right)}{b} \\
&= \frac{\text{Subst}\left(\int \left(\frac{\sin(x)}{2(1-x)} + \frac{\sin(x)}{2(1+x)}\right) dx, x, \tanh(a + bx)\right)}{b} \\
&= \frac{\text{Subst}\left(\int \frac{\sin(x)}{1-x} dx, x, \tanh(a + bx)\right)}{2b} + \frac{\text{Subst}\left(\int \frac{\sin(x)}{1+x} dx, x, \tanh(a + bx)\right)}{2b} \\
&= -\frac{\cos(1)\text{Subst}\left(\int \frac{\sin(1-x)}{1-x} dx, x, \tanh(a + bx)\right)}{2b} + \frac{\cos(1)\text{Subst}\left(\int \frac{\sin(1+x)}{1+x} dx, x, \tanh(a + bx)\right)}{2b} \\
&= -\frac{\text{Ci}(1 - \tanh(a + bx)) \sin(1)}{2b} - \frac{\text{Ci}(1 + \tanh(a + bx)) \sin(1)}{2b} + \frac{\cos(1)\text{Si}(1 - \tanh(a + bx))}{2b} + \frac{\cos(1)\text{Si}(1 + \tanh(a + bx))}{2b}
\end{aligned}$$

Mathematica [A]

time = 0.10, size = 59, normalized size = 0.77

$$\frac{\text{CosIntegral}(1 - \tanh(a + bx)) \sin(1) + \text{CosIntegral}(1 + \tanh(a + bx)) \sin(1) - \cos(1)(\text{Si}(1 - \tanh(a + bx)) + \text{Si}(1 + \tanh(a + bx)))}{2b}$$

Antiderivative was successfully verified.

`[In] Integrate[Sin[Tanh[a + b*x]], x]`

```
[Out] -1/2*(CosIntegral[1 - Tanh[a + b*x]]*Sin[1] + CosIntegral[1 + Tanh[a + b*x]]*Sin[1] - Cos[1]*(SinIntegral[1 - Tanh[a + b*x]] + SinIntegral[1 + Tanh[a + b*x]]))/b
```

Maple [A]

time = 0.97, size = 58, normalized size = 0.75

method	result
derivativdivides	$\frac{\sinIntegral(\tanh(bx+a)+1) \cos(1) - \cosineIntegral(\tanh(bx+a)+1) \sin(1) - \sinIntegral(-1+\tanh(bx+a)) \cos(1) - \cosineIntegral(-1+\tanh(bx+a)) \sin(1)}{2b}$
default	$\frac{\sinIntegral(\tanh(bx+a)+1) \cos(1) - \cosineIntegral(\tanh(bx+a)+1) \sin(1) - \sinIntegral(-1+\tanh(bx+a)) \cos(1) - \cosineIntegral(-1+\tanh(bx+a)) \sin(1)}{2b}$
risch	$\frac{ie^{-i} \expIntegral\left(1, -\frac{2i}{e^{2bx+2a}+1}\right)}{4b} - \frac{ie^i \expIntegral\left(1, -\frac{2i}{e^{2bx+2a}+1}+2i\right)}{4b} - \frac{ie^i \expIntegral\left(1, \frac{2i}{e^{2bx+2a}+1}\right)}{4b} + \frac{ie^{-i} \expIntegral\left(1, \frac{2i}{e^{2bx+2a}+1}\right)}{4b}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sin(tanh(b*x+a)), x, method=_RETURNVERBOSE)`

```
[Out] 1/b*(1/2*Si(tanh(b*x+a)+1)*cos(1)-1/2*Ci(tanh(b*x+a)+1)*sin(1)-1/2*Si(-1+tanh(b*x+a))*cos(1)-1/2*Ci(-1+tanh(b*x+a))*sin(1))
```


Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(tanh(b*x+a)),x, algorithm="maxima")

[Out] integrate(sin(tanh(b*x + a)), x)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 149 vs. 2(67) = 134.

time = 0.35, size = 149, normalized size = 1.94

$$\frac{\operatorname{Ci}\left(\frac{2e^{(2bx+2a)}}{e^{(2bx+2a)}+1}\right)\sin(1) + \operatorname{Ci}\left(-\frac{2e^{(2bx+2a)}}{e^{(2bx+2a)}+1}\right)\sin(1) + \operatorname{Ci}\left(\frac{2}{e^{(2bx+2a)}+1}\right)\sin(1) + \operatorname{Ci}\left(-\frac{2}{e^{(2bx+2a)}+1}\right)\sin(1) - 2\cos(1)\operatorname{Si}\left(\frac{2e^{(2bx+2a)}}{e^{(2bx+2a)}+1}\right) - 2\cos(1)\operatorname{Si}\left(\frac{2}{e^{(2bx+2a)}+1}\right)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(tanh(b*x+a)),x, algorithm="fricas")

[Out] -1/4*(cos_integral(2*e^(2*b*x + 2*a)/(e^(2*b*x + 2*a) + 1))*sin(1) + cos_integral(-2*e^(2*b*x + 2*a)/(e^(2*b*x + 2*a) + 1))*sin(1) + cos_integral(2/(e^(2*b*x + 2*a) + 1))*sin(1) + cos_integral(-2/(e^(2*b*x + 2*a) + 1))*sin(1) - 2*cos(1)*sin_integral(2*e^(2*b*x + 2*a)/(e^(2*b*x + 2*a) + 1)) - 2*cos(1)*sin_integral(2/(e^(2*b*x + 2*a) + 1)))/b

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sin(\tanh(a + bx)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(tanh(b*x+a)),x)

[Out] Integral(sin(tanh(a + b*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(tanh(b*x+a)),x, algorithm="giac")

[Out] integrate(sin(tanh(b*x + a)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \sin(\tanh(a + bx)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(tanh(a + b*x)),x)`

[Out] `int(sin(tanh(a + b*x)), x)`

3.243 $\int \csc(\tanh(a + bx)) dx$

Optimal. Leaf size=67

$$-\frac{1}{2}\text{Int}\left(\frac{\csc(\tanh(a + bx))\text{sech}^2(a + bx)}{-1 + \tanh(a + bx)}, x\right) + \frac{1}{2}\text{Int}\left(\frac{\csc(\tanh(a + bx))\text{sech}^2(a + bx)}{1 + \tanh(a + bx)}, x\right)$$

[Out] $-1/2*\text{Unintegrable}(\csc(\tanh(b*x+a))*\text{sech}(b*x+a)^2/(-1+\tanh(b*x+a)), x)+1/2*\text{Unintegrable}(\csc(\tanh(b*x+a))*\text{sech}(b*x+a)^2/(1+\tanh(b*x+a)), x)$

Rubi [A]

time = 0.06, antiderivative size = 0, normalized size of antiderivative = 0.00, number of steps used = 0, number of rules used = 0, integrand size = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \csc(\tanh(a + bx)) dx$$

Verification is not applicable to the result.

[In] $\text{Int}[\text{Csc}[\text{Tanh}[a + b*x]], x]$

[Out] $-1/2*\text{Defer}[\text{Subst}][\text{Defer}[\text{Int}][\text{Csc}[x]/(-1 + x), x], x, \text{Tanh}[a + b*x]]/b + \text{Defer}[\text{Subst}][\text{Defer}[\text{Int}][\text{Csc}[x]/(1 + x), x], x, \text{Tanh}[a + b*x]]/(2*b)$

Rubi steps

$$\begin{aligned} \int \csc(\tanh(a + bx)) dx &= \frac{\text{Subst}\left(\int \frac{\csc(x)}{1-x^2} dx, x, \tanh(a + bx)\right)}{b} \\ &= \frac{\text{Subst}\left(\int \left(-\frac{\csc(x)}{2(-1+x)} + \frac{\csc(x)}{2(1+x)}\right) dx, x, \tanh(a + bx)\right)}{b} \\ &= -\frac{\text{Subst}\left(\int \frac{\csc(x)}{-1+x} dx, x, \tanh(a + bx)\right)}{2b} + \frac{\text{Subst}\left(\int \frac{\csc(x)}{1+x} dx, x, \tanh(a + bx)\right)}{2b} \end{aligned}$$

Mathematica [A]

time = 2.00, size = 0, normalized size = 0.00

$$\int \csc(\tanh(a + bx)) dx$$

Verification is not applicable to the result.

[In] $\text{Integrate}[\text{Csc}[\text{Tanh}[a + b*x]], x]$

[Out] Integrate[Csc[Tanh[a + b*x]], x]

Maple [A]

time = 1.52, size = 0, normalized size = 0.00

$$\int \csc(\tanh(bx + a)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(tanh(b*x+a)),x)

[Out] int(csc(tanh(b*x+a)),x)

Maxima [A]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(tanh(b*x+a)),x, algorithm="maxima")

[Out] integrate(csc(tanh(b*x + a)), x)

Fricas [A]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(tanh(b*x+a)),x, algorithm="fricas")

[Out] integral(csc(tanh(b*x + a)), x)

Sympy [A]

time = 0.00, size = 0, normalized size = 0.00

$$\int \csc(\tanh(a + bx)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(tanh(b*x+a)),x)

[Out] Integral(csc(tanh(a + b*x)), x)

Giac [A]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(tanh(b*x+a)),x, algorithm="giac")
```

```
[Out] integrate(csc(tanh(b*x + a)), x)
```

Mupad [A]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sin(\tanh(a + bx))} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/sin(tanh(a + b*x)),x)
```

```
[Out] int(1/sin(tanh(a + b*x)), x)
```

3.244 $\int \cos^3(\tanh(a + bx)) dx$

Optimal. Leaf size=157

$$\frac{\cos(3)\text{CosIntegral}(3 - 3 \tanh(a + bx))}{8b} - \frac{3 \cos(1)\text{CosIntegral}(1 - \tanh(a + bx))}{8b} + \frac{3 \cos(1)\text{CosIntegral}(1 + \tanh(a + bx))}{8b}$$

[Out] $-3/8*\text{Ci}(1-\tanh(b*x+a))*\cos(1)/b+3/8*\text{Ci}(1+\tanh(b*x+a))*\cos(1)/b-1/8*\text{Ci}(3-3*\tanh(b*x+a))*\cos(3)/b+1/8*\text{Ci}(3+3*\tanh(b*x+a))*\cos(3)/b+3/8*\text{Si}(-1+\tanh(b*x+a))*\sin(1)/b+3/8*\text{Si}(1+\tanh(b*x+a))*\sin(1)/b+1/8*\text{Si}(-3+3*\tanh(b*x+a))*\sin(3)/b+1/8*\text{Si}(3+3*\tanh(b*x+a))*\sin(3)/b$

Rubi [A]

time = 0.28, antiderivative size = 157, normalized size of antiderivative = 1.00, number of steps used = 19, number of rules used = 5, integrand size = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.556$, Rules used = {6857, 3393, 3384, 3380, 3383}

$$\frac{\cos(3)\text{CosIntegral}(3-3\tanh(a+bx))}{8b} - \frac{3\cos(1)\text{CosIntegral}(1-\tanh(a+bx))}{8b} + \frac{3\cos(1)\text{CosIntegral}(\tanh(a+bx)+1)}{8b} + \frac{\cos(3)\text{CosIntegral}(3\tanh(a+bx)+3)}{8b} - \frac{\sin(3)\text{Si}(3-3\tanh(a+bx))}{8b} - \frac{3\sin(1)\text{Si}(1-\tanh(a+bx))}{8b} + \frac{3\sin(1)\text{Si}(\tanh(a+bx)+1)}{8b} + \frac{\sin(3)\text{Si}(3\tanh(a+bx)+3)}{8b}$$

Antiderivative was successfully verified.

[In] Int[Cos[Tanh[a + b*x]]^3,x]

[Out] $-1/8*(\text{Cos}[3]*\text{CosIntegral}[3 - 3*\text{Tanh}[a + b*x]])/b - (3*\text{Cos}[1]*\text{CosIntegral}[1 - \text{Tanh}[a + b*x]])/(8*b) + (3*\text{Cos}[1]*\text{CosIntegral}[1 + \text{Tanh}[a + b*x]])/(8*b) + (\text{Cos}[3]*\text{CosIntegral}[3 + 3*\text{Tanh}[a + b*x]])/(8*b) - (\text{Sin}[3]*\text{SinIntegral}[3 - 3*\text{Tanh}[a + b*x]])/(8*b) - (3*\text{Sin}[1]*\text{SinIntegral}[1 - \text{Tanh}[a + b*x]])/(8*b) + (3*\text{Sin}[1]*\text{SinIntegral}[1 + \text{Tanh}[a + b*x]])/(8*b) + (\text{Sin}[3]*\text{SinIntegral}[3 + 3*\text{Tanh}[a + b*x]])/(8*b)$

Rule 3380

Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] :> Simp[SinIntegral[e + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*e - c*f, 0]

Rule 3383

Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] :> Simp[CosIntegral[e - Pi/2 + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*(e - Pi/2) - c*f, 0]

Rule 3384

Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] :> Dist[Cos[(d*e - c*f)/d], Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Dist[Sin[(d*e - c*f)/d], Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]

Rule 3393

```
Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int
t[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f
, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))
```

Rule 6857

```
Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
xpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]
```

Rubi steps

$$\begin{aligned}
\int \cos^3(\tanh(a + bx)) dx &= \frac{\text{Subst}\left(\int \frac{\cos^3(x)}{1-x^2} dx, x, \tanh(a + bx)\right)}{b} \\
&= \frac{\text{Subst}\left(\int \left(-\frac{\cos^3(x)}{2(-1+x)} + \frac{\cos^3(x)}{2(1+x)}\right) dx, x, \tanh(a + bx)\right)}{b} \\
&= -\frac{\text{Subst}\left(\int \frac{\cos^3(x)}{-1+x} dx, x, \tanh(a + bx)\right)}{2b} + \frac{\text{Subst}\left(\int \frac{\cos^3(x)}{1+x} dx, x, \tanh(a + bx)\right)}{2b} \\
&= -\frac{\text{Subst}\left(\int \left(\frac{3\cos(x)}{4(-1+x)} + \frac{\cos(3x)}{4(-1+x)}\right) dx, x, \tanh(a + bx)\right)}{2b} + \frac{\text{Subst}\left(\int \left(\frac{3\cos(x)}{4(1+x)} + \frac{\cos(3x)}{4(1+x)}\right) dx, x, \tanh(a + bx)\right)}{2b} \\
&= -\frac{\text{Subst}\left(\int \frac{\cos(3x)}{-1+x} dx, x, \tanh(a + bx)\right)}{8b} + \frac{\text{Subst}\left(\int \frac{\cos(3x)}{1+x} dx, x, \tanh(a + bx)\right)}{8b} \\
&= -\frac{(3\cos(1))\text{Subst}\left(\int \frac{\cos(1-x)}{-1+x} dx, x, \tanh(a + bx)\right)}{8b} + \frac{(3\cos(1))\text{Subst}\left(\int \frac{\cos(1+x)}{1+x} dx, x, \tanh(a + bx)\right)}{8b} \\
&= -\frac{\cos(3)\text{Ci}(3 - 3\tanh(a + bx))}{8b} - \frac{3\cos(1)\text{Ci}(1 - \tanh(a + bx))}{8b} + \frac{3\cos(1)\text{Ci}(1 + \tanh(a + bx))}{8b}
\end{aligned}$$

Mathematica [A]

time = 0.22, size = 124, normalized size = 0.79

$$-\frac{2\cos(3)\text{CosIntegral}(3 - 3\tanh(a + bx)) - 6\cos(1)\text{CosIntegral}(1 - \tanh(a + bx)) + 6\cos(1)\text{CosIntegral}(1 + \tanh(a + bx)) + 2\cos(3)\text{CosIntegral}(3 + 3\tanh(a + bx)) - 2\sin(3)\text{Si}(3 - 3\tanh(a + bx)) - 6\sin(1)\text{Si}(1 - \tanh(a + bx)) + 6\sin(1)\text{Si}(1 + \tanh(a + bx)) + 2\sin(3)\text{Si}(3 + 3\tanh(a + bx))}{16}$$

Antiderivative was successfully verified.

```
[In] Integrate[Cos[Tanh[a + b*x]]^3, x]
```

```
[Out] (-2*Cos[3]*CosIntegral[3 - 3*Tanh[a + b*x]] - 6*Cos[1]*CosIntegral[1 - Tanh
[a + b*x]] + 6*Cos[1]*CosIntegral[1 + Tanh[a + b*x]] + 2*Cos[3]*CosIntegral
```

$[3 + 3*\text{Tanh}[a + b*x]] - 2*\text{Sin}[3]*\text{SinIntegral}[3 - 3*\text{Tanh}[a + b*x]] - 6*\text{Sin}[1]*\text{SinIntegral}[1 - \text{Tanh}[a + b*x]] + 6*\text{Sin}[1]*\text{SinIntegral}[1 + \text{Tanh}[a + b*x]] + 2*\text{Sin}[3]*\text{SinIntegral}[3 + 3*\text{Tanh}[a + b*x]]/(16*b)$

Maple [A]

time = 1.29, size = 118, normalized size = 0.75

method	result
derivativedivides	$\frac{\text{sinIntegral}(3+3 \tanh(bx+a)) \sin(3)}{8} + \frac{\text{cosineIntegral}(3+3 \tanh(bx+a)) \cos(3)}{8} + \frac{\text{sinIntegral}(-3+3 \tanh(bx+a)) \sin(3)}{8} - \frac{\text{cosineIntegral}(-3+3 \tanh(bx+a)) \cos(3)}{8}$
default	$\frac{\text{sinIntegral}(3+3 \tanh(bx+a)) \sin(3)}{8} + \frac{\text{cosineIntegral}(3+3 \tanh(bx+a)) \cos(3)}{8} + \frac{\text{sinIntegral}(-3+3 \tanh(bx+a)) \sin(3)}{8} - \frac{\text{cosineIntegral}(-3+3 \tanh(bx+a)) \cos(3)}{8}$
risch	$\frac{e^{-3i} \exp\text{Integral}\left(1, -\frac{6i}{e^{2bx+2a}+1}\right)}{16b} - \frac{e^{3i} \exp\text{Integral}\left(1, -\frac{6i}{e^{2bx+2a}+1}+6i\right)}{16b} + \frac{3e^{-i} \exp\text{Integral}\left(1, -\frac{2i}{e^{2bx+2a}+1}\right)}{16b} - \frac{3e^{i} \exp\text{Integral}\left(1, -\frac{2i}{e^{2bx+2a}+1}+6i\right)}{16b}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(tanh(b*x+a))^3,x,method=_RETURNVERBOSE)

[Out] $1/b*(1/8*\text{Si}(3+3*\text{tanh}(b*x+a))*\text{sin}(3)+1/8*\text{Ci}(3+3*\text{tanh}(b*x+a))*\text{cos}(3)+1/8*\text{Si}(-3+3*\text{tanh}(b*x+a))*\text{sin}(3)-1/8*\text{Ci}(-3+3*\text{tanh}(b*x+a))*\text{cos}(3)+3/8*\text{Si}(\text{tanh}(b*x+a)+1)*\text{sin}(1)+3/8*\text{Ci}(\text{tanh}(b*x+a)+1)*\text{cos}(1)+3/8*\text{Si}(-1+\text{tanh}(b*x+a))*\text{sin}(1)-3/8*\text{Ci}(-1+\text{tanh}(b*x+a))*\text{cos}(1))$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(tanh(b*x+a))^3,x, algorithm="maxima")

[Out] integrate(cos(tanh(b*x + a))^3, x)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 298 vs. 2(139) = 278.

time = 0.39, size = 298, normalized size = 1.90

$$\frac{\cos(3) \text{Ci}\left(\frac{3+3 \tanh(bx+a)}{2}\right) + 3 \cos(1) \text{Ci}\left(\frac{2+2 \tanh(bx+a)}{2}\right) + 3 \cos(1) \text{Ci}\left(-\frac{2+2 \tanh(bx+a)}{2}\right) + \cos(3) \text{Ci}\left(-\frac{3+3 \tanh(bx+a)}{2}\right) - \cos(3) \text{Ci}\left(\frac{3}{2}\right) - 3 \cos(1) \text{Ci}\left(\frac{3}{2}\right) - 3 \cos(1) \text{Ci}\left(-\frac{3}{2}\right) - \cos(3) \text{Ci}\left(-\frac{3}{2}\right) + 2 \sin(3) \text{Si}\left(\frac{6 \tanh(bx+a)}{2}\right) + 6 \sin(1) \text{Si}\left(\frac{2+2 \tanh(bx+a)}{2}\right) - 2 \sin(3) \text{Si}\left(\frac{3}{2}\right) - 6 \sin(1) \text{Si}\left(\frac{3}{2}\right)}{16b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(tanh(b*x+a))^3,x, algorithm="fricas")

[Out] $1/16*(\cos(3)*\text{cos_integral}(6*e^{(2*b*x + 2*a)})/(e^{(2*b*x + 2*a)} + 1)) + 3*\cos(1)*\text{cos_integral}(2*e^{(2*b*x + 2*a)})/(e^{(2*b*x + 2*a)} + 1) + 3*\cos(1)*\text{cos_integral}(-2*e^{(2*b*x + 2*a)})/(e^{(2*b*x + 2*a)} + 1) + \cos(3)*\text{cos_integral}(-6*e^{(2*b*x + 2*a)})/(e^{(2*b*x + 2*a)} + 1) - \cos(3)*\text{cos_integral}(6/(e^{(2*b*x + 2*a)} + 1))$

a) + 1)) - 3*cos(1)*cos_integral(2/(e^(2*b*x + 2*a) + 1)) - 3*cos(1)*cos_in
tegral(-2/(e^(2*b*x + 2*a) + 1)) - cos(3)*cos_integral(-6/(e^(2*b*x + 2*a)
+ 1)) + 2*sin(3)*sin_integral(6*e^(2*b*x + 2*a)/(e^(2*b*x + 2*a) + 1)) + 6*
sin(1)*sin_integral(2*e^(2*b*x + 2*a)/(e^(2*b*x + 2*a) + 1)) - 2*sin(3)*sin
_integral(6/(e^(2*b*x + 2*a) + 1)) - 6*sin(1)*sin_integral(2/(e^(2*b*x + 2*
a) + 1)))/b

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \cos^3(\tanh(a + bx)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(tanh(b*x+a))**3,x)

[Out] Integral(cos(tanh(a + b*x))**3, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(tanh(b*x+a))^3,x, algorithm="giac")

[Out] integrate(cos(tanh(b*x + a))^3, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(\tanh(a + bx))^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(tanh(a + b*x))^3,x)

[Out] int(cos(tanh(a + b*x))^3, x)

3.245 $\int \cos^2(\tanh(a + bx)) dx$

Optimal. Leaf size=115

$$-\frac{\cos(2)\text{CosIntegral}(2 - 2 \tanh(a + bx))}{4b} + \frac{\cos(2)\text{CosIntegral}(2 + 2 \tanh(a + bx))}{4b} - \frac{\log(1 - \tanh(a + bx))}{4b} + \frac{\log(1 + \tanh(a + bx))}{4b}$$

[Out] $-1/4*\text{Ci}(2-2*\tanh(b*x+a))*\cos(2)/b+1/4*\text{Ci}(2+2*\tanh(b*x+a))*\cos(2)/b-1/4*\ln(1-\tanh(b*x+a))/b+1/4*\ln(1+\tanh(b*x+a))/b+1/4*\text{Si}(-2+2*\tanh(b*x+a))*\sin(2)/b+1/4*\text{Si}(2+2*\tanh(b*x+a))*\sin(2)/b$

Rubi [A]

time = 0.19, antiderivative size = 115, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 5, integrand size = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.556$, Rules used = {6857, 3393, 3384, 3380, 3383}

$$-\frac{\cos(2)\text{CosIntegral}(2 - 2 \tanh(a + bx))}{4b} + \frac{\cos(2)\text{CosIntegral}(2 \tanh(a + bx) + 2)}{4b} - \frac{\sin(2)\text{Si}(2 - 2 \tanh(a + bx))}{4b} + \frac{\sin(2)\text{Si}(2 \tanh(a + bx) + 2)}{4b} - \frac{\log(1 - \tanh(a + bx))}{4b} + \frac{\log(\tanh(a + bx) + 1)}{4b}$$

Antiderivative was successfully verified.

[In] Int[Cos[Tanh[a + b*x]]^2,x]

[Out] $-1/4*(\text{Cos}[2]*\text{CosIntegral}[2 - 2*\text{Tanh}[a + b*x]])/b + (\text{Cos}[2]*\text{CosIntegral}[2 + 2*\text{Tanh}[a + b*x]])/(4*b) - \text{Log}[1 - \text{Tanh}[a + b*x]]/(4*b) + \text{Log}[1 + \text{Tanh}[a + b*x]]/(4*b) - (\text{Sin}[2]*\text{SinIntegral}[2 - 2*\text{Tanh}[a + b*x]])/(4*b) + (\text{Sin}[2]*\text{SinIntegral}[2 + 2*\text{Tanh}[a + b*x]])/(4*b)$

Rule 3380

Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[SinIntegral[e + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*e - c*f, 0]

Rule 3383

Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CosIntegral[e - Pi/2 + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*(e - Pi/2) - c*f, 0]

Rule 3384

Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Dist[Cos[(d*e - c*f)/d], Int[Sin[c*(f/d) + f*x]/(c + d*x), x] + Dist[Sin[(d*e - c*f)/d], Int[Cos[c*(f/d) + f*x]/(c + d*x), x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]

Rule 3393

Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f}

, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))

Rule 6857

Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
xpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]

Rubi steps

$$\begin{aligned}
 \int \cos^2(\tanh(a + bx)) dx &= \frac{\text{Subst}\left(\int \frac{\cos^2(x)}{1-x^2} dx, x, \tanh(a + bx)\right)}{b} \\
 &= \frac{\text{Subst}\left(\int \left(-\frac{\cos^2(x)}{2(-1+x)} + \frac{\cos^2(x)}{2(1+x)}\right) dx, x, \tanh(a + bx)\right)}{b} \\
 &= -\frac{\text{Subst}\left(\int \frac{\cos^2(x)}{-1+x} dx, x, \tanh(a + bx)\right)}{2b} + \frac{\text{Subst}\left(\int \frac{\cos^2(x)}{1+x} dx, x, \tanh(a + bx)\right)}{2b} \\
 &= -\frac{\text{Subst}\left(\int \left(\frac{1}{2(-1+x)} + \frac{\cos(2x)}{2(-1+x)}\right) dx, x, \tanh(a + bx)\right)}{2b} + \frac{\text{Subst}\left(\int \left(\frac{1}{2(1+x)} + \frac{\cos(2x)}{2(1+x)}\right) dx, x, \tanh(a + bx)\right)}{2b} \\
 &= -\frac{\log(1 - \tanh(a + bx))}{4b} + \frac{\log(1 + \tanh(a + bx))}{4b} - \frac{\text{Subst}\left(\int \frac{\cos(2x)}{-1+x} dx, x, \tanh(a + bx)\right)}{4b} \\
 &= -\frac{\log(1 - \tanh(a + bx))}{4b} + \frac{\log(1 + \tanh(a + bx))}{4b} - \frac{\cos(2)\text{Subst}\left(\int \frac{\cos(2-2x)}{-1+x} dx, x, \tanh(a + bx)\right)}{4b} \\
 &= -\frac{\cos(2)\text{Ci}(2 - 2 \tanh(a + bx))}{4b} + \frac{\cos(2)\text{Ci}(2 + 2 \tanh(a + bx))}{4b} - \frac{\log(1 - \tanh(a + bx))}{4b}
 \end{aligned}$$

Mathematica [A]

time = 0.13, size = 88, normalized size = 0.77

$$\frac{-\cos(2)\text{CosIntegral}(2 - 2 \tanh(a + bx)) + \cos(2)\text{CosIntegral}(2(1 + \tanh(a + bx))) - \log(1 - \tanh(a + bx)) + \log(1 + \tanh(a + bx)) - \sin(2)\text{Si}(2 - 2 \tanh(a + bx)) + \sin(2)\text{Si}(2(1 + \tanh(a + bx)))}{4b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[Tanh[a + b*x]]^2,x]

[Out] (-(Cos[2]*CosIntegral[2 - 2*Tanh[a + b*x]]) + Cos[2]*CosIntegral[2*(1 + Tan
h[a + b*x])) - Log[1 - Tanh[a + b*x]] + Log[1 + Tanh[a + b*x]] - Sin[2]*Sin
Integral[2 - 2*Tanh[a + b*x]] + Sin[2]*SinIntegral[2*(1 + Tanh[a + b*x])])/
(4*b)

Maple [A]

time = 1.05, size = 88, normalized size = 0.77

method	result
derivativedivides	$\frac{\sinIntegral(2+2 \tanh(bx+a)) \sin(2)}{4} + \frac{\cosineIntegral(2+2 \tanh(bx+a)) \cos(2)}{4} + \frac{\sinIntegral(-2+2 \tanh(bx+a)) \sin(2)}{4} - \frac{\cosineIntegral(-2+2 \tanh(bx+a)) \cos(2)}{4}$
default	$\frac{\sinIntegral(2+2 \tanh(bx+a)) \sin(2)}{4} + \frac{\cosineIntegral(2+2 \tanh(bx+a)) \cos(2)}{4} + \frac{\sinIntegral(-2+2 \tanh(bx+a)) \sin(2)}{4} - \frac{\cosineIntegral(-2+2 \tanh(bx+a)) \cos(2)}{4}$
risch	$\frac{x}{2} + \frac{e^{-2i} \expIntegral\left(1, -\frac{4i}{e^{2bx+2a}+1}\right)}{8b} - \frac{e^{2i} \expIntegral\left(1, -\frac{4i}{e^{2bx+2a}+1} + 4i\right)}{8b} + \frac{e^{2i} \expIntegral\left(1, \frac{4i}{e^{2bx+2a}+1}\right)}{8b} - \frac{e^{-2i} \expIntegral\left(1, \frac{4i}{e^{2bx+2a}+1}\right)}{8b}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(tanh(b*x+a))^2,x,method=_RETURNVERBOSE)`

[Out] $1/b*(1/4*Si(2+2*\tanh(b*x+a))*\sin(2)+1/4*Ci(2+2*\tanh(b*x+a))*\cos(2)+1/4*Si(-2+2*\tanh(b*x+a))*\sin(2)-1/4*Ci(-2+2*\tanh(b*x+a))*\cos(2)-1/4*\ln(-1+\tanh(b*x+a))+1/4*\ln(\tanh(b*x+a)+1))$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(tanh(b*x+a))^2,x, algorithm="maxima")`

[Out] $1/2*x + 1/2*\integrate(\cos(2*(e^{(2*b*x + 2*a)} - 1)/(e^{(2*b*x + 2*a)} + 1)), x)$

Fricas [A]

time = 0.39, size = 155, normalized size = 1.35

$$\frac{4bx + \cos(2) Ci\left(\frac{4e^{(2bx+2a)}}{e^{(2bx+2a)}+1}\right) + \cos(2) Ci\left(-\frac{4e^{(2bx+2a)}}{e^{(2bx+2a)}+1}\right) - \cos(2) Ci\left(\frac{4}{e^{(2bx+2a)}+1}\right) - \cos(2) Ci\left(-\frac{4}{e^{(2bx+2a)}+1}\right) + 2 \sin(2) Si\left(\frac{4e^{(2bx+2a)}}{e^{(2bx+2a)}+1}\right) - 2 \sin(2) Si\left(\frac{4}{e^{(2bx+2a)}+1}\right)}{8b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(tanh(b*x+a))^2,x, algorithm="fricas")`

[Out] $1/8*(4*b*x + \cos(2)*\cos_integral(4*e^{(2*b*x + 2*a)}/(e^{(2*b*x + 2*a)} + 1)) + \cos(2)*\cos_integral(-4*e^{(2*b*x + 2*a)}/(e^{(2*b*x + 2*a)} + 1)) - \cos(2)*\cos_integral(4/(e^{(2*b*x + 2*a)} + 1)) - \cos(2)*\cos_integral(-4/(e^{(2*b*x + 2*a)} + 1)) + 2*\sin(2)*\sin_integral(4*e^{(2*b*x + 2*a)}/(e^{(2*b*x + 2*a)} + 1)) - 2*\sin(2)*\sin_integral(4/(e^{(2*b*x + 2*a)} + 1)))/b$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \cos^2(\tanh(a + bx)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(tanh(b*x+a))**2,x)`

[Out] `Integral(cos(tanh(a + b*x))**2, x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(tanh(b*x+a))^2,x, algorithm="giac")`

[Out] `integrate(cos(tanh(b*x + a))^2, x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(\tanh(a + bx))^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(tanh(a + b*x))^2,x)`

[Out] `int(cos(tanh(a + b*x))^2, x)`

3.246 $\int \cos(\tanh(a + bx)) dx$

Optimal. Leaf size=77

$$-\frac{\cos(1)\text{CosIntegral}(1 - \tanh(a + bx))}{2b} + \frac{\cos(1)\text{CosIntegral}(1 + \tanh(a + bx))}{2b} - \frac{\sin(1)\text{Si}(1 - \tanh(a + bx))}{2b} + \frac{\sin(1)\text{Si}(1 + \tanh(a + bx))}{2b}$$

[Out] $-1/2*\text{Ci}(1-\tanh(b*x+a))*\cos(1)/b+1/2*\text{Ci}(1+\tanh(b*x+a))*\cos(1)/b+1/2*\text{Si}(-1+\tanh(b*x+a))*\sin(1)/b+1/2*\text{Si}(1+\tanh(b*x+a))*\sin(1)/b$

Rubi [A]

time = 0.11, antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 4, integrand size = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {3415, 3384, 3380, 3383}

$$-\frac{\cos(1)\text{CosIntegral}(1 - \tanh(a + bx))}{2b} + \frac{\cos(1)\text{CosIntegral}(\tanh(a + bx) + 1)}{2b} - \frac{\sin(1)\text{Si}(1 - \tanh(a + bx))}{2b} + \frac{\sin(1)\text{Si}(\tanh(a + bx) + 1)}{2b}$$

Antiderivative was successfully verified.

[In] `Int[Cos[Tanh[a + b*x]],x]`

[Out] $-1/2*(\text{Cos}[1]*\text{CosIntegral}[1 - \text{Tanh}[a + b*x]])/b + (\text{Cos}[1]*\text{CosIntegral}[1 + \text{Tanh}[a + b*x]])/(2*b) - (\text{Sin}[1]*\text{SinIntegral}[1 - \text{Tanh}[a + b*x]])/(2*b) + (\text{Sin}[1]*\text{SinIntegral}[1 + \text{Tanh}[a + b*x]])/(2*b)$

Rule 3380

`Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[SinIntegral[e + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*e - c*f, 0]`

Rule 3383

`Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CosIntegral[e - Pi/2 + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*(e - Pi/2) - c*f, 0]`

Rule 3384

`Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Dist[Cos[(d*e - c*f)/d], Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Dist[Sin[(d*e - c*f)/d], Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]`

Rule 3415

`Int[Cos[(c_.) + (d_.)*(x_)]*((a_.) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Int[ExpandIntegrand[Cos[c + d*x], (a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[p, 0] && IGtQ[n, 0] && (EqQ[n, 2] || EqQ[p, -1])`

Rubi steps

$$\begin{aligned}
\int \cos(\tanh(a + bx)) dx &= \frac{\text{Subst}\left(\int \frac{\cos(x)}{1-x^2} dx, x, \tanh(a + bx)\right)}{b} \\
&= \frac{\text{Subst}\left(\int \left(\frac{\cos(x)}{2(1-x)} + \frac{\cos(x)}{2(1+x)}\right) dx, x, \tanh(a + bx)\right)}{b} \\
&= \frac{\text{Subst}\left(\int \frac{\cos(x)}{1-x} dx, x, \tanh(a + bx)\right)}{2b} + \frac{\text{Subst}\left(\int \frac{\cos(x)}{1+x} dx, x, \tanh(a + bx)\right)}{2b} \\
&= \frac{\cos(1)\text{Subst}\left(\int \frac{\cos(1-x)}{1-x} dx, x, \tanh(a + bx)\right)}{2b} + \frac{\cos(1)\text{Subst}\left(\int \frac{\cos(1+x)}{1+x} dx, x, \tanh(a + bx)\right)}{2b} \\
&= -\frac{\cos(1)\text{Ci}(1 - \tanh(a + bx))}{2b} + \frac{\cos(1)\text{Ci}(1 + \tanh(a + bx))}{2b} - \frac{\sin(1)\text{Si}(1 - \tanh(a + bx))}{2b} + \frac{\sin(1)\text{Si}(1 + \tanh(a + bx))}{2b}
\end{aligned}$$

Mathematica [A]

time = 0.08, size = 62, normalized size = 0.81

$$\frac{-\cos(1)\text{CosIntegral}(1 - \tanh(a + bx)) + \cos(1)\text{CosIntegral}(1 + \tanh(a + bx)) - \sin(1)\text{Si}(1 - \tanh(a + bx)) + \sin(1)\text{Si}(1 + \tanh(a + bx))}{2b}$$

Antiderivative was successfully verified.

`[In] Integrate[Cos[Tanh[a + b*x]], x]`

```
[Out] (-(Cos[1]*CosIntegral[1 - Tanh[a + b*x]]) + Cos[1]*CosIntegral[1 + Tanh[a + b*x]] - Sin[1]*SinIntegral[1 - Tanh[a + b*x]] + Sin[1]*SinIntegral[1 + Tanh[a + b*x]])/(2*b)
```

Maple [A]

time = 0.99, size = 58, normalized size = 0.75

method	result
derivativedivides	$\frac{\sin\text{Integral}(-1+\tanh(bx+a))\sin(1)}{2} - \frac{\cosine\text{Integral}(-1+\tanh(bx+a))\cos(1)}{2} + \frac{\sin\text{Integral}(\tanh(bx+a)+1)\sin(1)}{2} + \frac{\cosine\text{Integral}(\tanh(bx+a)+1)\cos(1)}{2}$
default	$\frac{\sin\text{Integral}(-1+\tanh(bx+a))\sin(1)}{2} - \frac{\cosine\text{Integral}(-1+\tanh(bx+a))\cos(1)}{2} + \frac{\sin\text{Integral}(\tanh(bx+a)+1)\sin(1)}{2} + \frac{\cosine\text{Integral}(\tanh(bx+a)+1)\cos(1)}{2}$
risch	$\frac{e^{-i} \exp\text{Integral}\left(1, -\frac{2i}{e^{2bx+2a+1}}\right)}{4b} - \frac{e^i \exp\text{Integral}\left(1, -\frac{2i}{e^{2bx+2a+1}}+2i\right)}{4b} + \frac{e^i \exp\text{Integral}\left(1, \frac{2i}{e^{2bx+2a+1}}\right)}{4b} - \frac{e^{-i} \exp\text{Integral}\left(1, \frac{2i}{e^{2bx+2a+1}}+2i\right)}{4b}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(tanh(b*x+a)), x, method=_RETURNVERBOSE)`

```
[Out] 1/b*(1/2*Si(-1+tanh(b*x+a))*sin(1)-1/2*Ci(-1+tanh(b*x+a))*cos(1)+1/2*Si(tanh(b*x+a)+1)*sin(1)+1/2*Ci(tanh(b*x+a)+1)*cos(1))
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(tanh(b*x+a)),x, algorithm="maxima")

[Out] integrate(cos(tanh(b*x + a)), x)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 151 vs. 2(67) = 134.

time = 0.37, size = 151, normalized size = 1.96

$$\frac{\cos(1) \operatorname{Ci}\left(\frac{2e^{(2bx+2a)}}{e^{(2bx+2a)}+1}\right) + \cos(1) \operatorname{Ci}\left(-\frac{2e^{(2bx+2a)}}{e^{(2bx+2a)}+1}\right) - \cos(1) \operatorname{Ci}\left(\frac{2}{e^{(2bx+2a)}+1}\right) - \cos(1) \operatorname{Ci}\left(-\frac{2}{e^{(2bx+2a)}+1}\right) + 2 \sin(1) \operatorname{Si}\left(\frac{2e^{(2bx+2a)}}{e^{(2bx+2a)}+1}\right) - 2 \sin(1) \operatorname{Si}\left(\frac{2}{e^{(2bx+2a)}+1}\right)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(tanh(b*x+a)),x, algorithm="fricas")

[Out] 1/4*(cos(1)*cos_integral(2*e^(2*b*x + 2*a)/(e^(2*b*x + 2*a) + 1)) + cos(1)*cos_integral(-2*e^(2*b*x + 2*a)/(e^(2*b*x + 2*a) + 1)) - cos(1)*cos_integral(2/(e^(2*b*x + 2*a) + 1)) - cos(1)*cos_integral(-2/(e^(2*b*x + 2*a) + 1)) + 2*sin(1)*sin_integral(2*e^(2*b*x + 2*a)/(e^(2*b*x + 2*a) + 1)) - 2*sin(1)*sin_integral(2/(e^(2*b*x + 2*a) + 1)))/b

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \cos(\tanh(a + bx)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(tanh(b*x+a)),x)

[Out] Integral(cos(tanh(a + b*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(tanh(b*x+a)),x, algorithm="giac")

[Out] integrate(cos(tanh(b*x + a)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(\tanh(a + b x)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(tanh(a + b*x)),x)

[Out] int(cos(tanh(a + b*x)), x)

3.247 $\int \sec(\tanh(a + bx)) dx$

Optimal. Leaf size=67

$$-\frac{1}{2}\text{Int}\left(\frac{\sec(\tanh(a + bx))\text{sech}^2(a + bx)}{-1 + \tanh(a + bx)}, x\right) + \frac{1}{2}\text{Int}\left(\frac{\sec(\tanh(a + bx))\text{sech}^2(a + bx)}{1 + \tanh(a + bx)}, x\right)$$

[Out] $-1/2*\text{Unintegrable}(\sec(\tanh(b*x+a))*\text{sech}(b*x+a)^2/(-1+\tanh(b*x+a)), x)+1/2*\text{Unintegrable}(\sec(\tanh(b*x+a))*\text{sech}(b*x+a)^2/(1+\tanh(b*x+a)), x)$

Rubi [A]

time = 0.06, antiderivative size = 0, normalized size of antiderivative = 0.00, number of steps used = 0, number of rules used = 0, integrand size = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \sec(\tanh(a + bx)) dx$$

Verification is not applicable to the result.

[In] `Int[Sec[Tanh[a + b*x]], x]`

[Out] $-1/2*\text{Defer}[\text{Subst}[\text{Defer}[\text{Int}[\text{Sec}[x]/(-1 + x), x], x, \text{Tanh}[a + b*x]]/b + \text{Defer}[\text{Subst}[\text{Defer}[\text{Int}[\text{Sec}[x]/(1 + x), x], x, \text{Tanh}[a + b*x]]/(2*b)$

Rubi steps

$$\begin{aligned} \int \sec(\tanh(a + bx)) dx &= \frac{\text{Subst}\left(\int \frac{\sec(x)}{1-x^2} dx, x, \tanh(a + bx)\right)}{b} \\ &= \frac{\text{Subst}\left(\int \left(-\frac{\sec(x)}{2(-1+x)} + \frac{\sec(x)}{2(1+x)}\right) dx, x, \tanh(a + bx)\right)}{b} \\ &= -\frac{\text{Subst}\left(\int \frac{\sec(x)}{-1+x} dx, x, \tanh(a + bx)\right)}{2b} + \frac{\text{Subst}\left(\int \frac{\sec(x)}{1+x} dx, x, \tanh(a + bx)\right)}{2b} \end{aligned}$$

Mathematica [A]

time = 3.56, size = 0, normalized size = 0.00

$$\int \sec(\tanh(a + bx)) dx$$

Verification is not applicable to the result.

[In] `Integrate[Sec[Tanh[a + b*x]], x]`

[Out] Integrate[Sec[Tanh[a + b*x]], x]

Maple [A]

time = 1.16, size = 0, normalized size = 0.00

$$\int \sec(\tanh(bx + a)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(tanh(b*x+a)),x)

[Out] int(sec(tanh(b*x+a)),x)

Maxima [A]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(tanh(b*x+a)),x, algorithm="maxima")

[Out] integrate(sec(tanh(b*x + a)), x)

Fricas [A]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(tanh(b*x+a)),x, algorithm="fricas")

[Out] integral(sec(tanh(b*x + a)), x)

Sympy [A]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sec(\tanh(a + bx)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(tanh(b*x+a)),x)

[Out] Integral(sec(tanh(a + b*x)), x)

Giac [A]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(tanh(b*x+a)),x, algorithm="giac")
```

```
[Out] integrate(sec(tanh(b*x + a)), x)
```

Mupad [A]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\cos(\tanh(a + b x))} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/cos(tanh(a + b*x)),x)
```

```
[Out] int(1/cos(tanh(a + b*x)), x)
```

Chapter 4

Appendix

Local contents

4.1	Download section	1096
4.2	Listing of Grading functions	1096

4.1 Download section

The following zip files contain the raw integrals used in this test.

Mathematica format Mathematica_syntax.zip

Maple and Mupad format Maple_syntax.zip

Sympy format SYMPY_syntax.zip

Sage math format SAGE_syntax.zip

4.2 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

4.2.1 Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*           Small rewrite of logic in main function to make it*)
(*           match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*           GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*           is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*           antiderivative*)
(* "A" if result can be considered optimal*)
```

```

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafC
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A","none"}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
        ]
        ,(*ELSE*)
        finalresult={"C","Result contains complex when optimal does not."}
      ]
      ,(*ELSE*)(*result does not contains complex*)
      If[leafCountResult<=2*leafCountOptimal,
        finalresult={"A","none"}
        ,(*ELSE*)
        finalresult={"B","Leaf count is larger than twice the leaf count of optimal. $
      ]
    ]
    ,(*ELSE*)(*expnResult>expnOptimal*)
    If[FreeQ[result,Integrate] && FreeQ[result,Int],
      finalresult={"C","Result contains higher order function than in optimal. Order "<
    ,
    finalresult={"F","Contains unresolved integral."}
  ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)

```



```

ExpIntegralE, ExpIntegralEi, LogIntegral,
SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
Gamma, LogGamma, PolyGamma,
Zeta, PolyLog, ProductLog,
EllipticF, EllipticE, EllipticPi
},func]

HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1,Hypergeometric2F1,HypergeometricPFQ},func]

AppellFunctionQ[func_] :=
  MemberQ[{AppellF1},func]

```

4.2.2 Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
# if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
# see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
      #do NOT call ExpnType() if leaf size is too large. Recursion problem
      if leaf_count_result > 500000 then
          return "B","result has leaf size over 500,000. Avoiding possible recursion issues";
      fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

```

```

    if debug then
        print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
    fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
#   is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
#   antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A","";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of r
                    convert(leaf_count_result,string)," vs. $2 (" ,
                    convert(leaf_count_optimal,string)," ) = ",convert(2*leaf_co
            end if
        else #result contains complex but optimal is not
            if debug then
                print("result contains complex but optimal is not");
            fi;
            return "C","Result contains complex when optimal does not.";
        fi;
    else # result do not contain complex

```

```

    # this assumes optimal do not as well. No check is needed here.
    if debug then
        print("result do not contain complex, this assumes optimal do not as well")
    fi;
    if leaf_count_result<=2*leaf_count_optimal then
        if debug then
            print("leaf_count_result<=2*leaf_count_optimal");
        fi;
        return "A","";
    else
        if debug then
            print("leaf_count_result>2*leaf_count_optimal");
        fi;
        return "B",cat("Leaf count of result is larger than twice the leaf count of o
                        convert(leaf_count_result,string)," $ vs. $2(",
                        convert(leaf_count_optimal,string),")=",convert(2*leaf_cou

    fi;
fi;
else #ExpnType(result) > ExpnType(optimal)
    if debug then
        print("ExpnType(result) > ExpnType(optimal)");
    fi;
    return "C",cat("Result contains higher order function than in optimal. Order ",
                  convert(ExpnType_result,string)," vs. order ",
                  convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function
# 6 = appell function
# 7 = rootsum function

```

```

# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+`) or type(expn,'*`) then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
    9
  end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func,[
    exp,log,ln,
    sin,cos,tan,cot,sec,csc,

```

```

    arcsin,arccos,arctan,arccot,arcsec,arccsc,
    sinh,cosh,tanh,coth,sech,csch,
    arcsinh,arccosh,arctanh,arccoth,arcsech,arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf,erfc,erfi,
    FresnelS,FresnelC,
    Ei,Ei,Li,Si,Ci,Shi,Chi,
    GAMMA,lnGAMMA,Psi,Zeta,polylog,dilog,LambertW,
    EllipticF,EllipticE,EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1,hypergeom,HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product.  rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u),op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc:

```

4.2.3 Sympy grading function

```

#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):
    if isinstance(expr,Pow):
        if expr.args[1] == Rational(1,2):
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
        asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
        asinh,acosh,atanh,acoth,asech,acsch
    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
        fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
        gamma,loggamma,digamma,zeta,polylog,LambertW,
        elliptic_f,elliptic_e,elliptic_pi,exp_polar
    ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

```

```

except AttributeError as error:
    return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnT
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+' or type(expn,'*')
        m1 = expnType(expn.args[0])
        m2 = expnType(list(expn.args[1:]))
        return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
    elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
        return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
    elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
    elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif is_appell_function(expn.func):
        m1 = max(map(expnType, list(expn.args)))
        return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif isinstance(expn,RootSum):
        m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
        return max(7,m1)
    elif str(expn).find("Integral") != -1:

```

```

    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print ("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""
    else:
        if expnType_result <= expnType_optimal:
            if result.has(I):
                if optimal.has(I): #both result and optimal complex
                    if leaf_count_result <= 2*leaf_count_optimal:
                        grade = "A"
                        grade_annotation = ""
                    else:
                        grade = "B"
                        grade_annotation = "Both result and optimal contain complex but leaf count of result is larger"
                else: #result contains complex but optimal is not
                    grade = "C"
                    grade_annotation = "Result contains complex when optimal does not."
            else: # result do not contain complex, this assumes optimal do not as well
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)-str(leaf_count_optimal)
            else:
                grade = "C"
                grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType_result)-str(ExpnType_optimal)

```



```

# print("Before returning. grade=", grade, " grade_annotation=", grade_annotation)

return grade, grade_annotation

```

4.2.4 SageMath grading function

```

# Dec 24, 2019. Nasser: Ported original Maple grading function by
#       Albert Rich to use with Sagemath. This is used to
#       grade Fracas, Giac and Maxima results.
# Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#       'arctan2', 'floor', 'abs', 'log_integral'
# June 4, 2022 Made default grade_annotation "none" instead of "" due
#       issue later when reading the file.
# July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    # print("Enter tree_size, expr is ", expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: # isinstance(expr, Pow):
        if expr.operands()[1] == 1/2: # expr.args[1] == Rational(1,2):
            if debug: print("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

```

```

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:
            print ("func ", func , " is elementary_function")
        else:
            print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func , " is special_function")
        else:
            print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric'] #[appellf1] can't find this in sagemath

```

```

def is_atom(expn):

    #debug=False
    if debug:
        print ("Enter is_atom, expn=",expn)

    if not hasattr(expn, 'parent'):
        return False

    #thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__ )
    return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #isinstance(expn,Pow)
        if type(expn.operands()[1])==Integer: #isinstance(expn.args[1],Integer)
            return expnType(expn.operands()[0]) #expnType(expn.args[0])
        elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
            if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)

```

```

    return 1
  else:
    return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
  else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn.op
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isinstan
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    if debug:
        print ("Enter grade_antiderivative for sagemath")
        print("Enter grade_antiderivative, result=",result)
        print("Enter grade_antiderivative, optimal=",optimal)
        print("type(anti)=",type(result))
        print("type(optimal)=",type(optimal))

    leaf_count_result = tree_size(result) #leaf_count(result)
    leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

    #if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

```

```

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger t
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_
else:
    grade = "C"
    grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_resu

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```