


A Solution Manual For

Differential equations and linear algebra, 4th ed., Edwards and Penney



Nasser M. Abbasi

March 3, 2024

Contents

1	Section 5.2, Higher-Order Linear Differential Equations. General solutions of Linear Equations. Page 288	2
2	Section 5.3, Higher-Order Linear Differential Equations. Homogeneous Equations with Constant Coefficients. Page 300	10
3	Section 7.2, Matrices and Linear systems. Page 384	39
4	Section 7.3, The eigenvalue method for linear systems. Page 395	42
5	Section 7.6, Multiple Eigenvalue Solutions. Examples. Page 437	104
6	Section 7.6, Multiple Eigenvalue Solutions. Page 451	112
7	Chapter 11 Power series methods. Section 11.1 Introduction and Review of power series. Page 615	172
8	Chapter 11 Power series methods. Section 11.2 Power series solutions. Page 624	198

1 Section 5.2, Higher-Order Linear Differential Equations. General solutions of Linear Equations. Page 288

1.1	problem problem 38	3
1.2	problem problem 39	4
1.3	problem problem 40	5
1.4	problem problem 41	6
1.5	problem problem 42	7
1.6	problem problem 43	8
1.7	problem problem 44	9

1.1 problem problem 38

Internal problem ID [278]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 5.2, Higher-Order Linear Differential Equations. General solutions of Linear Equations. Page 288

Problem number: problem 38.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler], [_2nd_order, _linear, '_with_symmetry_[0,F`

$$x^2y'' + y'x - 9y = 0$$

Given that one solution of the ode is

$$y_1 = x^3$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve([x^2*diff(y(x),x$2)+x*diff(y(x),x)-9*y(x)=0,x^3],y(x), singsol=all)
```

$$y(x) = \frac{c_1}{x^3} + c_2x^3$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 18

```
DSolve[x^2*y''[x]+x*y'[x]-9*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_2x^6 + c_1}{x^3}$$

1.2 problem problem 39

Internal problem ID [279]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 5.2, Higher-Order Linear Differential Equations. General solutions of Linear Equations. Page 288

Problem number: problem 39.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$4y'' - 4y' + y = 0$$

Given that one solution of the ode is

$$y_1 = e^{\frac{x}{2}}$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 18

```
dsolve([4*diff(y(x),x$2)-4*diff(y(x),x)+y(x)=0,exp(x/2)],y(x), singsol=all)
```

$$y(x) = c_1 e^{\frac{x}{2}} + c_2 e^{\frac{x}{2}} x$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 20

```
DSolve[4*y''[x]-4*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{x/2}(c_2 x + c_1)$$

1.3 problem problem 40

Internal problem ID [280]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 5.2, Higher-Order Linear Differential Equations. General solutions of Linear Equations. Page 288

Problem number: problem 40.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' - x(2+x)y' + (2+x)y = 0$$

Given that one solution of the ode is

$$y_1 = x$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 13

```
dsolve([x^2*diff(y(x),x$2)-x*(x+2)*diff(y(x),x)+(x+2)*y(x)=0,x],y(x), singsol=all)
```

$$y(x) = c_1 x + c_2 x e^x$$

✓ Solution by Mathematica

Time used: 0.029 (sec). Leaf size: 16

```
DSolve[x^2*y''[x]-x*(x+2)*y'[x]+(x+2)*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x(c_2 e^x + c_1)$$

1.4 problem problem 41

Internal problem ID [281]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 5.2, Higher-Order Linear Differential Equations. General solutions of Linear Equations. Page 288

Problem number: problem 41.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x + 1)y'' - (2 + x)y' + y = 0$$

Given that one solution of the ode is

$$y_1 = e^x$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve([(x+1)*diff(y(x),x$2)-(x+2)*diff(y(x),x)+y(x)=0,exp(x)],y(x), singsol=all)
```

$$y(x) = c_1(2 + x) + e^x c_2$$

✓ Solution by Mathematica

Time used: 0.143 (sec). Leaf size: 29

```
DSolve[(x+1)*y'[x]-(x+2)*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_1 e^{x+1} - 2c_2(x+2)}{\sqrt{2e}}$$

1.5 problem problem 42

Internal problem ID [282]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 5.2, Higher-Order Linear Differential Equations. General solutions of Linear Equations. Page 288

Problem number: problem 42.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type [Gegenbauer]

$$(-x^2 + 1)y'' + 2y'x - 2y = 0$$

Given that one solution of the ode is

$$y_1 = x$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve([(1-x^2)*diff(y(x),x$2)+2*x*diff(y(x),x)-2*y(x)=0,x],y(x), singsol=all)
```

$$y(x) = c_1x + c_2(x^2 + 1)$$

✓ Solution by Mathematica

Time used: 0.08 (sec). Leaf size: 39

```
DSolve[(1-x^2)*y'[x]+2*x*y'[x]-2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\sqrt{x^2 - 1}(c_1(x - 1)^2 + c_2x)}{\sqrt{1 - x^2}}$$

1.6 problem problem 43

Internal problem ID [283]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 5.2, Higher-Order Linear Differential Equations. General solutions of Linear Equations. Page 288

Problem number: problem 43.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type [Gegenbauer]

$$(-x^2 + 1)y'' - 2y'x + 2y = 0$$

Given that one solution of the ode is

$$y_1 = x$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve([(1-x^2)*diff(y(x),x$2)-2*x*diff(y(x),x)+2*y(x)=0,x],y(x), singsol=all)
```

$$y(x) = c_1x + c_2\left(-\frac{x \ln(x+1)}{2} + \frac{x \ln(x-1)}{2} + 1\right)$$

✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 33

```
DSolve[(1-x^2)*y'[x]-2*x*y'[x]+2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1x - \frac{1}{2}c_2(x \log(1-x) - x \log(x+1) + 2)$$

1.7 problem problem 44

Internal problem ID [284]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 5.2, Higher-Order Linear Differential Equations. General solutions of Linear Equations. Page 288

Problem number: problem 44.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2 y'' + y' x + \left(x^2 - \frac{1}{4}\right) y = 0$$

Given that one solution of the ode is

$$y_1 = \frac{\cos(x)}{\sqrt{x}}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 19

```
dsolve([x^2*diff(y(x),x$2)+x*diff(y(x),x)+(x^2-1/4)*y(x)=0,x^(-1/2)*cos(x)],y(x), singsol=all)
```

$$y(x) = \frac{c_1 \sin(x)}{\sqrt{x}} + \frac{c_2 \cos(x)}{\sqrt{x}}$$

✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 33

```
DSolve[(1-x^2)*y'[x]-2*x*y'[x]+2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 x - \frac{1}{2} c_2 (x \log(1-x) - x \log(x+1) + 2)$$

2 Section 5.3, Higher-Order Linear Differential Equations. Homogeneous Equations with Constant Coefficients. Page 300

2.1	problem problem 10	11
2.2	problem problem 11	12
2.3	problem problem 12	13
2.4	problem problem 13	14
2.5	problem problem 14	15
2.6	problem problem 15	16
2.7	problem problem 16	17
2.8	problem problem 17	18
2.9	problem problem 18	19
2.10	problem problem 19	20
2.11	problem problem 20	21
2.12	problem problem 24	22
2.13	problem problem 25	23
2.14	problem problem 26	24
2.15	problem problem 27	25
2.16	problem problem 28	26
2.17	problem problem 29	27
2.18	problem problem 30	28
2.19	problem problem 31	29
2.20	problem problem 32	30
2.21	problem problem 38	31
2.22	problem problem 48	32
2.23	problem problem 49	33
2.24	problem problem 54	34
2.25	problem problem 55	35
2.26	problem problem 56	36
2.27	problem problem 57	37
2.28	problem problem 58	38

2.1 problem problem 10

Internal problem ID [285]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 5.3, Higher-Order Linear Differential Equations. Homogeneous Equations with Constant Coefficients. Page 300

Problem number: problem 10.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$5y'''' + 3y''' = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 20

```
dsolve(5*diff(y(x),x$4)+3*diff(y(x),x$3)=0,y(x), singsol=all)
```

$$y(x) = c_1 + xc_2 + c_3x^2 + c_4e^{-\frac{3x}{5}}$$

✓ Solution by Mathematica

Time used: 0.037 (sec). Leaf size: 30

```
DSolve[5*y''''[x]+3*y'''[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{125}{27}c_1e^{-3x/5} + x(c_4x + c_3) + c_2$$

2.2 problem problem 11

Internal problem ID [286]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 5.3, Higher-Order Linear Differential Equations. Homogeneous Equations with Constant Coefficients. Page 300

Problem number: problem 11.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' - 8y''' + 16y'' = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 22

```
dsolve(diff(y(x),x$4)-8*diff(y(x),x$3)+16*diff(y(x),x$2)=0,y(x), singsol=all)
```

$$y(x) = c_1 + xc_2 + c_3e^{4x} + c_4e^{4x}x$$

✓ Solution by Mathematica

Time used: 0.081 (sec). Leaf size: 34

```
DSolve[y''''[x]-8*y'''[x]+16*y''[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{32}e^{4x}(c_2(2x - 1) + 2c_1) + c_4x + c_3$$

2.3 problem problem 12

Internal problem ID [287]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 5.3, Higher-Order Linear Differential Equations. Homogeneous Equations with Constant Coefficients. Page 300

Problem number: problem 12.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' - 3y''' + 3y'' - y' = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 22

```
dsolve(diff(y(x),x$4)-3*diff(y(x),x$3)+3*diff(y(x),x$2)-diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = c_1 + e^x c_2 + c_3 e^x x + c_4 e^x x^2$$

✓ Solution by Mathematica

Time used: 0.028 (sec). Leaf size: 32

```
DSolve[y''''[x]-3*y'''[x]+3*y''[x]-y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x (c_3(x^2 - 2x + 2) + c_2(x - 1) + c_1) + c_4$$

2.4 problem problem 13

Internal problem ID [288]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 5.3, Higher-Order Linear Differential Equations. Homogeneous Equations with Constant Coefficients. Page 300

Problem number: problem 13.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$9y''' + 12y'' + 4y' = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(9*diff(y(x),x$3)+12*diff(y(x),x$2)+4*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = c_1 + c_2 e^{-\frac{2x}{3}} + c_3 e^{-\frac{2x}{3}} x$$

✓ Solution by Mathematica

Time used: 0.047 (sec). Leaf size: 32

```
DSolve[9*y'''[x]+12*y''[x]+4*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_3 - \frac{3}{4} e^{-2x/3} (c_2(2x + 3) + 2c_1)$$

2.5 problem problem 14

Internal problem ID [289]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 5.3, Higher-Order Linear Differential Equations. Homogeneous Equations with Constant Coefficients. Page 300

Problem number: problem 14.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' + 3y'' - 4y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve(diff(y(x),x$4)+3*diff(y(x),x$2)-4*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^x c_1 + e^{-x} c_2 + c_3 \sin(2x) + c_4 \cos(2x)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 34

```
DSolve[y''''[x]+3*y''[x]-4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_3 e^{-x} + c_4 e^x + c_1 \cos(2x) + c_2 \sin(2x)$$

2.6 problem problem 15

Internal problem ID [290]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 5.3, Higher-Order Linear Differential Equations. Homogeneous Equations with Constant Coefficients. Page 300

Problem number: problem 15.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' - 16y'' + 16y = 0$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 75

```
dsolve(diff(y(x),x$4)-16*diff(y(x),x$2)+16*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{(-\sqrt{3}\sqrt{2}-\sqrt{2})x} + c_2 e^{(\sqrt{3}\sqrt{2}+\sqrt{2})x} + c_3 e^{(-\sqrt{3}\sqrt{2}+\sqrt{2})x} + c_4 e^{(\sqrt{3}\sqrt{2}-\sqrt{2})x}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 86

```
DSolve[y''''[x]-16*y''[x]+16*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{2\sqrt{2-\sqrt{3}}x} + c_2 e^{-2\sqrt{2-\sqrt{3}}x} + c_3 e^{2\sqrt{2+\sqrt{3}}x} + c_4 e^{-2\sqrt{2+\sqrt{3}}x}$$

2.7 problem problem 16

Internal problem ID [291]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 5.3, Higher-Order Linear Differential Equations. Homogeneous Equations with Constant Coefficients. Page 300

Problem number: problem 16.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' + 18y'' + 81y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 31

```
dsolve(diff(y(x),x$4)+18*diff(y(x),x$2)+81*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 \sin(3x) + c_2 \cos(3x) + c_3 \sin(3x)x + c_4 \cos(3x)x$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 30

```
DSolve[y''''[x]+18*y''[x]+81*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow (c_2x + c_1) \cos(3x) + (c_4x + c_3) \sin(3x)$$

2.8 problem problem 17

Internal problem ID [292]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 5.3, Higher-Order Linear Differential Equations. Homogeneous Equations with Constant Coefficients. Page 300

Problem number: problem 17.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$6y'''' + 11y'' + 4y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 41

```
dsolve(6*diff(y(x),x$4)+11*diff(y(x),x$2)+4*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 \sin\left(\frac{\sqrt{2}x}{2}\right) + c_2 \cos\left(\frac{\sqrt{2}x}{2}\right) + c_3 \sin\left(\frac{2\sqrt{3}x}{3}\right) + c_4 \cos\left(\frac{2\sqrt{3}x}{3}\right)$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 94

```
DSolve[y''''[x]+11*y''[x]+4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_3 \cos\left(\sqrt{\frac{1}{2}(11 - \sqrt{105})}x\right) + c_1 \cos\left(\sqrt{\frac{1}{2}(11 + \sqrt{105})}x\right) \\ + c_4 \sin\left(\sqrt{\frac{1}{2}(11 - \sqrt{105})}x\right) + c_2 \sin\left(\sqrt{\frac{1}{2}(11 + \sqrt{105})}x\right)$$

2.9 problem problem 18

Internal problem ID [293]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 5.3, Higher-Order Linear Differential Equations. Homogeneous Equations with Constant Coefficients. Page 300

Problem number: problem 18.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' - 16y = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 29

```
dsolve(diff(y(x),x$4)=16*y(x),y(x), singsol=all)
```

$$y(x) = c_1 e^{2x} + c_2 e^{-2x} + c_3 \sin(2x) + c_4 \cos(2x)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 36

```
DSolve[y''''[x]==16*y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{2x} + c_3 e^{-2x} + c_2 \cos(2x) + c_4 \sin(2x)$$

2.10 problem problem 19

Internal problem ID [294]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 5.3, Higher-Order Linear Differential Equations. Homogeneous Equations with Constant Coefficients. Page 300

Problem number: problem 19.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' + y'' - y' - y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 22

```
dsolve(diff(y(x),x$3)+diff(y(x),x$2)-diff(y(x),x)-y(x)=0,y(x), singsol=all)
```

$$y(x) = e^x c_1 + e^{-x} c_2 + c_3 e^{-x} x$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 26

```
DSolve[y'''[x]+y''[x]-y'[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x} (c_2 x + c_3 e^{2x} + c_1)$$

2.11 problem problem 20

Internal problem ID [295]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 5.3, Higher-Order Linear Differential Equations. Homogeneous Equations with Constant Coefficients. Page 300

Problem number: problem 20.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' + 2y''' + 3y'' + 2y' + y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 59

```
dsolve(diff(y(x),x$4)+2*diff(y(x),x$3)+3*diff(y(x),x$2)+2*diff(y(x),x)+y(x)=0,y(x), singsol=
```

$$y(x) = c_1 e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) + c_2 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) \\ + c_3 e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) x + c_4 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) x$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 52

```
DSolve[y''''[x]+2*y'''[x]+3*y''[x]+2*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x/2} \left((c_4 x + c_3) \cos\left(\frac{\sqrt{3}x}{2}\right) + (c_2 x + c_1) \sin\left(\frac{\sqrt{3}x}{2}\right) \right)$$

2.12 problem problem 24

Internal problem ID [296]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 5.3, Higher-Order Linear Differential Equations. Homogeneous Equations with Constant Coefficients. Page 300

Problem number: problem 24.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$2y''' - 3y'' - 2y' = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = -1, y''(0) = 3]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 18

```
dsolve([2*diff(y(x),x$3)-3*diff(y(x),x$2)-2*diff(y(x),x)=0,y(0) = 1, D(y)(0) = -1, (D@@2)(y)
```

$$y(x) = -\frac{7}{2} + \frac{e^{2x}}{2} + 4e^{-\frac{x}{2}}$$

✓ Solution by Mathematica

Time used: 0.351 (sec). Leaf size: 70

```
DSolve[{2*y'''[x]-3*y''[x]-3*y'[x]==0,{y[0]==1,y'[0]==-1,y''[0]==3}},y[x],x,IncludeSingularS
```

$$y(x) \rightarrow \frac{1}{66}e^{-\frac{1}{4}(\sqrt{33}-3)x} \left((99 - 13\sqrt{33}) e^{\frac{\sqrt{33}x}{2}} - 132e^{\frac{1}{4}(\sqrt{33}-3)x} + 99 + 13\sqrt{33} \right)$$

2.13 problem problem 25

Internal problem ID [297]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 5.3, Higher-Order Linear Differential Equations. Homogeneous Equations with Constant Coefficients. Page 300

Problem number: problem 25.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$3y''' + 2y'' = 0$$

With initial conditions

$$[y(0) = -1, y'(0) = 0, y''(0) = 1]$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 15

```
dsolve([3*diff(y(x),x$3)+2*diff(y(x),x$2)=0,y(0) = -1, D(y)(0) = 0, (D@@2)(y)(0) = 1],y(x),
```

$$y(x) = -\frac{13}{4} + \frac{3x}{2} + \frac{9e^{-\frac{2x}{3}}}{4}$$

✓ Solution by Mathematica

Time used: 0.038 (sec). Leaf size: 23

```
DSolve[{3*y'''[x]+2*y''[x]==0,{y[0]==1,y'[0]==-1,y''[0]==3}},y[x],x,IncludeSingularSolutions
```

$$y(x) \rightarrow \frac{1}{4}(14x + 27e^{-2x/3} - 23)$$

2.14 problem problem 26

Internal problem ID [298]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 5.3, Higher-Order Linear Differential Equations. Homogeneous Equations with Constant Coefficients. Page 300

Problem number: problem 26.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' + 10y'' + 25y' = 0$$

With initial conditions

$$[y(0) = 3, y'(0) = 4, y''(0) = 5]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 19

```
dsolve([diff(y(x),x$3)+10*diff(y(x),x$2)+25*diff(y(x),x)=0,y(0) = 3, D(y)(0) = 4, (D@@2)(y)(0) = 5],y(x),x)
```

$$y(x) = \frac{24}{5} - \frac{9e^{-5x}}{5} - 5e^{-5x}x$$

✓ Solution by Mathematica

Time used: 0.045 (sec). Leaf size: 26

```
DSolve[{y'''[x]+10*y''[x]+25*y'[x]==0,{y[0]==3,y'[0]==4,y''[0]==5}},y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \frac{1}{5}e^{-5x}(-25x + 24e^{5x} - 9)$$

2.15 problem problem 27

Internal problem ID [299]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 5.3, Higher-Order Linear Differential Equations. Homogeneous Equations with Constant Coefficients. Page 300

Problem number: problem 27.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' + 3y'' - 4y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 22

```
dsolve(diff(y(x),x$3)+3*diff(y(x),x$2)-4*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^x c_1 + c_2 e^{-2x} + c_3 e^{-2x} x$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 26

```
DSolve[y'''[x]+3*y''[x]-4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2x} (c_2 x + c_3 e^{3x} + c_1)$$

2.16 problem problem 28

Internal problem ID [300]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 5.3, Higher-Order Linear Differential Equations. Homogeneous Equations with Constant Coefficients. Page 300

Problem number: problem 28.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$2y''' - y'' - 5y' - 2y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(2*diff(y(x),x$3)-diff(y(x),x$2)-5*diff(y(x),x)-2*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{2x} + c_2 e^{-\frac{x}{2}} + c_3 e^{-x}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 32

```
DSolve[2*y'''[x]-y''[x]-5*y'[x]-2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x}(c_1 e^{x/2} + c_3 e^{3x} + c_2)$$

2.17 problem problem 29

Internal problem ID [301]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 5.3, Higher-Order Linear Differential Equations. Homogeneous Equations with Constant Coefficients. Page 300

Problem number: problem 29.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' + 27y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 37

```
dsolve(diff(y(x),x$3)+27*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{-3x} + c_2 e^{\frac{3x}{2}} \sin\left(\frac{3\sqrt{3}x}{2}\right) + c_3 e^{\frac{3x}{2}} \cos\left(\frac{3\sqrt{3}x}{2}\right)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 56

```
DSolve[y'''[x]+27*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-3x} \left(c_3 e^{9x/2} \cos\left(\frac{3\sqrt{3}x}{2}\right) + c_2 e^{9x/2} \sin\left(\frac{3\sqrt{3}x}{2}\right) + c_1 \right)$$

2.18 problem problem 30

Internal problem ID [302]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 5.3, Higher-Order Linear Differential Equations. Homogeneous Equations with Constant Coefficients. Page 300

Problem number: problem 30.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' - y''' + y'' - 3y' - 6y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 33

```
dsolve(diff(y(x),x$4)-diff(y(x),x$3)+diff(y(x),x$2)-3*diff(y(x),x)-6*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{2x} + e^{-x} c_2 + c_3 \sin(\sqrt{3}x) + c_4 \cos(\sqrt{3}x)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 44

```
DSolve[y''''[x]-y'''[x]+y''[x]-3*y'[x]-6*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_3 e^{-x} + c_4 e^{2x} + c_1 \cos(\sqrt{3}x) + c_2 \sin(\sqrt{3}x)$$

2.19 problem problem 31

Internal problem ID [303]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 5.3, Higher-Order Linear Differential Equations. Homogeneous Equations with Constant Coefficients. Page 300

Problem number: problem 31.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' + 3y'' + 4y' - 8y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
dsolve(diff(y(x),x$3)+3*diff(y(x),x$2)+4*diff(y(x),x)-8*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^x c_1 + c_2 e^{-2x} \sin(2x) + c_3 e^{-2x} \cos(2x)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 34

```
DSolve[y'''[x]+3*y''[x]+4*y'[x]-8*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2x} (c_3 e^{3x} + c_2 \cos(2x) + c_1 \sin(2x))$$

2.20 problem problem 32

Internal problem ID [304]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 5.3, Higher-Order Linear Differential Equations. Homogeneous Equations with Constant Coefficients. Page 300

Problem number: problem 32.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' + y''' - 3y'' - 5y' - 2y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 33

```
dsolve(diff(y(x),x$4)+diff(y(x),x$3)-3*diff(y(x),x$2)-5*diff(y(x),x)-2*y(x)=0,y(x), singsol=
```

$$y(x) = c_1 e^{2x} + e^{-x} c_2 + c_3 e^{-x} x + c_4 e^{-x} x^2$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 32

```
DSolve[y''''[x]+y'''[x]-3*y''[x]-5*y'[x]-2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x} (c_3 x^2 + c_2 x + c_4 e^{3x} + c_1)$$

2.21 problem problem 38

Internal problem ID [305]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 5.3, Higher-Order Linear Differential Equations. Homogeneous Equations with Constant Coefficients. Page 300

Problem number: problem 38.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' - 5y'' + 100y' - 500y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 10, y''(0) = 250]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 17

```
dsolve([diff(y(x),x$3)-5*diff(y(x),x$2)+100*diff(y(x),x)-500*y(x)=0,y(0) = 0, D(y)(0) = 10,
```

$$y(x) = 2e^{5x} - 2\cos(10x)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 19

```
DSolve[{y'''[x]-5*y''[x]+100*y'[x]-500*y[x]==0,{y[0]==0,y'[0]==10,y''[0]==250}},y[x],x,Inclu
```

$$y(x) \rightarrow 2(e^{5x} - \cos(10x))$$

2.22 problem problem 48

Internal problem ID [306]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 5.3, Higher-Order Linear Differential Equations. Homogeneous Equations with Constant Coefficients. Page 300

Problem number: problem 48.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_x]]`

$$y''' - y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 0, y''(0) = 0]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 22

```
dsolve([diff(y(x),x$3)=y(x),y(0) = 1, D(y)(0) = 0, (D@@2)(y)(0) = 0],y(x), singsol=all)
```

$$y(x) = \frac{e^x}{3} + \frac{2e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 33

```
DSolve[{y'''[x]==y[x],{y[0]==1,y'[0]==0,y''[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{3} \left(e^x + 2e^{-x/2} \cos\left(\frac{\sqrt{3}x}{2}\right) \right)$$

2.23 problem problem 49

Internal problem ID [307]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 5.3, Higher-Order Linear Differential Equations. Homogeneous Equations with Constant Coefficients. Page 300

Problem number: problem 49.

ODE order: 4.

ODE degree: 1.

CAS Maple gives this as type `[[_high_order, _missing_x]]`

$$y'''' - y''' - y'' - y' - 2y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 0, y''(0) = 0, y'''(0) = 15]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 23

```
dsolve([diff(y(x),x$4)=diff(y(x),x$3)+diff(y(x),x$2)+diff(y(x),x)+2*y(x),y(0) = 0, D(y)(0) =
```

$$y(x) = e^{2x} - \frac{5e^{-x}}{2} - \frac{9\sin(x)}{2} + \frac{3\cos(x)}{2}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 33

```
DSolve[{y''''[x]==y[x],{y[0]==1,y'[0]==0,y''[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{3} \left(e^x + 2e^{-x/2} \cos \left(\frac{\sqrt{3}x}{2} \right) \right)$$

2.24 problem problem 54

Internal problem ID [308]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 5.3, Higher-Order Linear Differential Equations. Homogeneous Equations with Constant Coefficients. Page 300

Problem number: problem 54.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_y]]`

$$x^3 y''' + 6x^2 y'' + 4y'x = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(x^3*diff(y(x),x$3)+6*x^2*diff(y(x),x$2)+4*x*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = c_1 + c_2 \ln(x) + \frac{c_3}{x^3}$$

✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 22

```
DSolve[x^3*y'''[x]+6*x^2*y''[x]+4*x*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{c_1}{3x^3} + c_2 \log(x) + c_3$$

2.25 problem problem 55

Internal problem ID [309]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 5.3, Higher-Order Linear Differential Equations. Homogeneous Equations with Constant Coefficients. Page 300

Problem number: problem 55.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_y]]`

$$x^3 y''' - x^2 y'' + y' x = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 18

```
dsolve(x^3*diff(y(x),x$3)-x^2*diff(y(x),x$2)+x*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = c_1 + x^2 c_2 + c_3 x^2 \ln(x)$$

✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 35

```
DSolve[x^3*y'''[x]-x^2*y''[x]+x*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{4}(2c_1 - c_2)x^2 + \frac{1}{2}c_2 x^2 \log(x) + c_3$$

2.26 problem problem 56

Internal problem ID [310]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 5.3, Higher-Order Linear Differential Equations. Homogeneous Equations with Constant Coefficients. Page 300

Problem number: problem 56.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_y]]`

$$x^3 y''' + 3x^2 y'' + y' x = 0$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 16

```
dsolve(x^3*diff(y(x),x$3)+3*x^2*diff(y(x),x$2)+x*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = c_3 \ln(x)^2 + c_2 \ln(x) + c_1$$

✓ Solution by Mathematica

Time used: 0.019 (sec). Leaf size: 23

```
DSolve[x^3*y'''[x]+3*x^2*y''[x]+x*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}c_2 \log^2(x) + c_1 \log(x) + c_3$$

2.27 problem problem 57

Internal problem ID [311]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 5.3, Higher-Order Linear Differential Equations. Homogeneous Equations with Constant Coefficients. Page 300

Problem number: problem 57.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _missing_y]]`

$$x^3 y''' - 3x^2 y'' + y' x = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(x^3*diff(y(x),x$3)-3*x^2*diff(y(x),x$2)+x*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = c_1 + c_2 x^{3+\sqrt{3}} + c_3 x^{3-\sqrt{3}}$$

✓ Solution by Mathematica

Time used: 0.136 (sec). Leaf size: 54

```
DSolve[x^3*y'''[x]-3*x^2*y''[x]+x*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_2 x^{3+\sqrt{3}}}{3+\sqrt{3}} + \frac{c_1 x^{3-\sqrt{3}}}{3-\sqrt{3}} + c_3$$

2.28 problem problem 58

Internal problem ID [312]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 5.3, Higher-Order Linear Differential Equations. Homogeneous Equations with Constant Coefficients. Page 300

Problem number: problem 58.

ODE order: 3.

ODE degree: 1.

CAS Maple gives this as type `[[_3rd_order, _exact, _linear, _homogeneous]]`

$$x^3 y''' + 6x^2 y'' + 7y'x + y = 0$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(x^3*diff(y(x),x$3)+6*x^2*diff(y(x),x$2)+7*x*diff(y(x),x)+y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{c_1}{x} + \frac{c_2 \ln(x)}{x} + \frac{c_3 \ln(x)^2}{x}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 24

```
DSolve[x^3*y'''[x]+6*x^2*y''[x]+7*x*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_3 \log^2(x) + c_2 \log(x) + c_1}{x}$$

3 Section 7.2, Matrices and Linear systems. Page 384

3.1	problem problem 13	40
3.2	problem problem 14	41

3.1 problem problem 13

Internal problem ID [313]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.2, Matrices and Linear systems. Page 384

Problem number: problem 13.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= 6x_1(t) \\x_2'(t) &= -3x_1(t) - x_2(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 28

```
dsolve([diff(x__1(t),t)=4*x__1(t)+2*x__1(t),diff(x__2(t),t)=-3*x__1(t)-x__2(t)],[x__1(t), x__2(t)])
```

$$x_1(t) = -\frac{7c_2e^{6t}}{3}$$

$$x_2(t) = e^{-t}c_1 + c_2e^{6t}$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 56

```
DSolve[{x1'[t]==4*x1[t]+2*x2[t],x2'[t]==-3*x1[t]-x2[t]},{x1[t],x2[t]},t,IncludeSingularSolutions->True]
```

$$x1(t) \rightarrow e^t(c_1(3e^t - 2) + 2c_2(e^t - 1))$$

$$x2(t) \rightarrow e^t(c_2(3 - 2e^t) - 3c_1(e^t - 1))$$

3.2 problem problem 14

Internal problem ID [314]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.2, Matrices and Linear systems. Page 384

Problem number: problem 14.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = -3x_1(t) + 2x_2(t)$$

$$x_2'(t) = -3x_1(t) + 4x_2(t)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 36

```
dsolve([diff(x__1(t),t)=-3*x__1(t)+2*x__2(t),diff(x__2(t),t)=-3*x__1(t)+4*x__2(t)], [x__1(t),
```

$$x_1(t) = 2c_1e^{-2t} + \frac{c_2e^{3t}}{3}$$

$$x_2(t) = c_1e^{-2t} + c_2e^{3t}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 73

```
DSolve[{x1'[t]==-3*x1[t]+2*x2[t],x2'[t]==-3*x1[t]+4*x2[t]},{x1[t],x2[t]},t,IncludeSingularSo
```

$$x_1(t) \rightarrow \frac{1}{5}e^{-2t}(2c_2(e^{5t} - 1) - c_1(e^{5t} - 6))$$

$$x_2(t) \rightarrow \frac{1}{5}e^{-2t}(c_2(6e^{5t} - 1) - 3c_1(e^{5t} - 1))$$

4 Section 7.3, The eigenvalue method for linear systems. Page 395

4.1	problem problem 1	44
4.2	problem problem 2	45
4.3	problem problem 3	46
4.4	problem problem 4	47
4.5	problem problem 5	48
4.6	problem problem 6	49
4.7	problem problem 7	50
4.8	problem problem 8	51
4.9	problem problem 9	52
4.10	problem problem 10	53
4.11	problem problem 11	54
4.12	problem problem 12	55
4.13	problem problem 13	56
4.14	problem problem 14	57
4.15	problem problem 15	58
4.16	problem problem 16	59
4.17	problem problem 17	60
4.18	problem problem 18	62
4.19	problem problem 19	64
4.20	problem problem 20	66
4.21	problem problem 21	68
4.22	problem problem 22	70
4.23	problem problem 23	71
4.24	problem problem 24	72
4.25	problem problem 25	74
4.26	problem problem 26	76
4.27	problem problem 38	78
4.28	problem problem 39	80
4.29	problem problem 40	82
4.30	problem problem 41	84
4.31	problem problem 42	86
4.32	problem problem 43	88
4.33	problem problem 44	90
4.34	problem problem 45	92
4.35	problem problem 46	94
4.36	problem problem 47	96

4.37	problem	problem 48	98
4.38	problem	problem 49	100
4.39	problem	problem 50	102

4.1 problem problem 1

Internal problem ID [315]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 1.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = x_1(t) + 2x_2(t)$$

$$x_2'(t) = 2x_1(t) + x_2(t)$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 35

```
dsolve([diff(x__1(t),t)=x__1(t)+2*x__2(t),diff(x__2(t),t)=2*x__1(t)+x__2(t)],[x__1(t), x__2(t)
```

$$x_1(t) = -e^{-t}c_1 + c_2e^{3t}$$

$$x_2(t) = e^{-t}c_1 + c_2e^{3t}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 68

```
DSolve[{x1'[t]==x1[t]+2*x2[t],x2'[t]==2*x1[t]+x2[t]},{x1[t],x2[t]},t,IncludeSingularSolution
```

$$x1(t) \rightarrow \frac{1}{2}e^{-t}(c_1(e^{4t} + 1) + c_2(e^{4t} - 1))$$

$$x2(t) \rightarrow \frac{1}{2}e^{-t}(c_1(e^{4t} - 1) + c_2(e^{4t} + 1))$$

4.2 problem problem 2

Internal problem ID [316]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 2.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 2x_1(t) + 3x_2(t)$$

$$x_2'(t) = 2x_1(t) + x_2(t)$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 36

```
dsolve([diff(x__1(t),t)=2*x__1(t)+3*x__2(t),diff(x__2(t),t)=2*x__1(t)+x__2(t)],[x__1(t), x__2(t)]
```

$$x_1(t) = \frac{3c_1 e^{4t}}{2} - c_2 e^{-t}$$

$$x_2(t) = c_1 e^{4t} + c_2 e^{-t}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 74

```
DSolve[{x1'[t]==2*x1[t]+3*x2[t],x2'[t]==2*x1[t]+x2[t]},{x1[t],x2[t]},t,IncludeSingularSolutions->True]
```

$$x_1(t) \rightarrow \frac{1}{5} e^{-t} (c_1 (3e^{5t} + 2) + 3c_2 (e^{5t} - 1))$$

$$x_2(t) \rightarrow \frac{1}{5} e^{-t} (2c_1 (e^{5t} - 1) + c_2 (2e^{5t} + 3))$$

4.3 problem problem 3

Internal problem ID [317]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 3.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 3x_1(t) + 4x_2(t)$$

$$x_2'(t) = 3x_1(t) + 2x_2(t)$$

With initial conditions

$$[x_1(0) = 1, x_2(0) = 1]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 34

```
dsolve([diff(x__1(t),t) = 3*x__1(t)+4*x__2(t), diff(x__2(t),t) = 3*x__1(t)+2*x__2(t), x__1(0)=1, x__2(0)=1])
```

$$x_1(t) = -\frac{e^{-t}}{7} + \frac{8e^{6t}}{7}$$

$$x_2(t) = \frac{e^{-t}}{7} + \frac{6e^{6t}}{7}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 44

```
DSolve[{x1'[t]==3*x1[t]+4*x2[t], x2'[t]==3*x1[t]+2*x2[t]}, {x1[0]==1, x2[0]==1}, {x1[t], x2[t]}, t]
```

$$x_1(t) \rightarrow \frac{1}{7}e^{-t}(8e^{7t} - 1)$$

$$x_2(t) \rightarrow \frac{1}{7}e^{-t}(6e^{7t} + 1)$$

4.4 problem problem 4

Internal problem ID [318]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 4.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 4x_1(t) + x_2(t)$$

$$x_2'(t) = 6x_1(t) - x_2(t)$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 35

```
dsolve([diff(x__1(t),t)=4*x__1(t)+x__2(t),diff(x__2(t),t)=6*x__1(t)-x__2(t)],[x__1(t), x__2(t)
```

$$x_1(t) = -\frac{c_1 e^{-2t}}{6} + c_2 e^{5t}$$

$$x_2(t) = c_1 e^{-2t} + c_2 e^{5t}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 71

```
DSolve[{x1'[t]==4*x1[t]+x2[t],x2'[t]==6*x1[t]-x2[t]},{x1[t],x2[t]},t,IncludeSingularSolution
```

$$x_1(t) \rightarrow \frac{1}{7} e^{-2t} (c_1 (6e^{7t} + 1) + c_2 (e^{7t} - 1))$$

$$x_2(t) \rightarrow \frac{1}{7} e^{-2t} (6c_1 (e^{7t} - 1) + c_2 (e^{7t} + 6))$$

4.5 problem problem 5

Internal problem ID [319]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 5.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 6x_1(t) - 7x_2(t)$$

$$x_2'(t) = x_1(t) - 2x_2(t)$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 35

```
dsolve([diff(x__1(t),t)=6*x__1(t)-7*x__2(t),diff(x__2(t),t)=x__1(t)-2*x__2(t)],[x__1(t), x__2(t)])
```

$$x_1(t) = 7c_1e^{5t} + c_2e^{-t}$$

$$x_2(t) = c_1e^{5t} + c_2e^{-t}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 72

```
DSolve[{x1'[t]==6*x1[t]-7*x2[t],x2'[t]==x1[t]-2*x2[t]},{x1[t],x2[t]},t,IncludeSingularSolutions->True]
```

$$x_1(t) \rightarrow \frac{1}{6}e^{-t}(c_1(7e^{6t} - 1) - 7c_2(e^{6t} - 1))$$

$$x_2(t) \rightarrow \frac{1}{6}e^{-t}(c_1(e^{6t} - 1) - c_2(e^{6t} - 7))$$

4.6 problem problem 6

Internal problem ID [320]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 6.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= 9x_1(t) + 5x_2(t) \\x_2'(t) &= -6x_1(t) - 2x_2(t)\end{aligned}$$

With initial conditions

$$[x_1(0) = 1, x_2(0) = 0]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 34

```
dsolve([diff(x__1(t),t) = 9*x__1(t)+5*x__2(t), diff(x__2(t),t) = -6*x__1(t)-2*x__2(t), x__1(0)=1, x__2(0)=0])
```

$$x_1(t) = 6e^{4t} - 5e^{3t}$$

$$x_2(t) = -6e^{4t} + 6e^{3t}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 33

```
DSolve[{x1'[t]==9*x1[t]+5*x2[t],x2'[t]==-6*x1[t]-2*x2[t]},{x1[0]==1,x2[0]==0},{x1[t],x2[t]},t]
```

$$x1(t) \rightarrow e^{3t}(6e^t - 5)$$

$$x2(t) \rightarrow -6e^{3t}(e^t - 1)$$

4.7 problem problem 7

Internal problem ID [321]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 7.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = -3x_1(t) + 4x_2(t)$$

$$x_2'(t) = 6x_1(t) - 5x_2(t)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 31

```
dsolve([diff(x__1(t),t)=-3*x__1(t)+4*x__2(t),diff(x__2(t),t)=6*x__1(t)-5*x__2(t)],[x__1(t),
```

$$x_1(t) = -\frac{2c_1e^{-9t}}{3} + c_2e^t$$

$$x_2(t) = c_1e^{-9t} + c_2e^t$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 74

```
DSolve[{x1'[t]==-3*x1[t]+4*x2[t],x2'[t]==6*x1[t]-5*x2[t]},{x1[t],x2[t]},t,IncludeSingularSol
```

$$x_1(t) \rightarrow \frac{1}{5}e^{-9t}(c_1(3e^{10t} + 2) + 2c_2(e^{10t} - 1))$$

$$x_2(t) \rightarrow \frac{1}{5}e^{-9t}(3c_1(e^{10t} - 1) + c_2(2e^{10t} + 3))$$

4.8 problem problem 8

Internal problem ID [322]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 8.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = x_1(t) - 5x_2(t)$$

$$x_2'(t) = x_1(t) - x_2(t)$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 48

```
dsolve([diff(x__1(t),t)=x__1(t)-5*x__2(t),diff(x__2(t),t)=x__1(t)-x__2(t)],[x__1(t), x__2(t)]
```

$$x_1(t) = 2c_1 \cos(2t) - 2c_2 \sin(2t) + c_1 \sin(2t) + c_2 \cos(2t)$$

$$x_2(t) = c_1 \sin(2t) + c_2 \cos(2t)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 48

```
DSolve[{x1'[t]==x1[t]-5*x2[t],x2'[t]==x1[t]-x2[t]},{x1[t],x2[t]},t,IncludeSingularSolutions
```

$$x1(t) \rightarrow c_1 \cos(2t) + (c_1 - 5c_2) \sin(t) \cos(t)$$

$$x2(t) \rightarrow c_2 \cos(2t) + (c_1 - c_2) \sin(t) \cos(t)$$

4.9 problem problem 9

Internal problem ID [323]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 9.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 2x_1(t) - 5x_2(t)$$

$$x_2'(t) = 4x_1(t) - 2x_2(t)$$

With initial conditions

$$[x_1(0) = 2, x_2(0) = 3]$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 34

```
dsolve([diff(x__1(t),t) = 2*x__1(t)-5*x__2(t), diff(x__2(t),t) = 4*x__1(t)-2*x__2(t), x__1(0)=2, x__2(0)=3])
```

$$x_1(t) = 2 \cos(4t) - \frac{11 \sin(4t)}{4}$$

$$x_2(t) = \frac{\sin(4t)}{2} + 3 \cos(4t)$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 34

```
DSolve[{x1'[t]==x1[t]-5*x2[t],x2'[t]==x1[t]-x2[t]},{x1[0]==2,x2[0]==3},{x1[t],x2[t]},t,IncludeSingularFunctions->False]
```

$$x1(t) \rightarrow 2 \cos(2t) - 13 \sin(t) \cos(t)$$

$$x2(t) \rightarrow 3 \cos(2t) - \sin(t) \cos(t)$$

4.10 problem problem 10

Internal problem ID [324]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 10.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = -3x_1(t) - 2x_2(t)$$

$$x_2'(t) = 9x_1(t) + 3x_2(t)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 50

```
dsolve([diff(x__1(t),t)=-3*x__1(t)-2*x__2(t),diff(x__2(t),t)=9*x__1(t)+3*x__2(t)],[x__1(t),
```

$$x_1(t) = \frac{c_1 \cos(3t)}{3} - \frac{c_2 \sin(3t)}{3} - \frac{c_1 \sin(3t)}{3} - \frac{c_2 \cos(3t)}{3}$$

$$x_2(t) = c_1 \sin(3t) + c_2 \cos(3t)$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 53

```
DSolve[{x1'[t]==-3*x1[t]-2*x2[t],x2'[t]==9*x1[t]+3*x2[t]},{x1[t],x2[t]},t,IncludeSingularSol
```

$$x_1(t) \rightarrow c_1 \cos(3t) - \frac{1}{3}(3c_1 + 2c_2) \sin(3t)$$

$$x_2(t) \rightarrow c_2 \cos(3t) + (3c_1 + c_2) \sin(3t)$$

4.11 problem problem 11

Internal problem ID [325]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 11.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = x_1(t) - 2x_2(t)$$

$$x_2'(t) = 2x_1(t) + x_2(t)$$

With initial conditions

$$[x_1(0) = 0, x_2(0) = 4]$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 24

```
dsolve([diff(x__1(t),t) = x__1(t)-2*x__2(t), diff(x__2(t),t) = 2*x__1(t)+x__2(t), x__1(0) =
```

$$x_1(t) = -4e^t \sin(2t)$$

$$x_2(t) = 4e^t \cos(2t)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 26

```
DSolve[{x1'[t]==x1[t]-2*x2[t],x2'[t]==2*x1[t]+x2[t]},{x1[0]==0,x2[0]==4},{x1[t],x2[t]},t,Inc
```

$$x1(t) \rightarrow -4e^t \sin(2t)$$

$$x2(t) \rightarrow 4e^t \cos(2t)$$

4.12 problem problem 12

Internal problem ID [326]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 12.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = x_1(t) - 5x_2(t)$$

$$x_2'(t) = x_1(t) + 3x_2(t)$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 60

```
dsolve([diff(x__1(t),t)=x__1(t)-5*x__2(t),diff(x__2(t),t)=x__1(t)+3*x__2(t)],[x__1(t), x__2(t)
```

$$x_1(t) = e^{2t}(2c_1 \cos(2t) - c_2 \cos(2t) - c_1 \sin(2t) - 2c_2 \sin(2t))$$

$$x_2(t) = e^{2t}(c_1 \sin(2t) + c_2 \cos(2t))$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 67

```
DSolve[{x1'[t]==x1[t]-5*x2[t],x2'[t]==x1[t]+3*x2[t]},{x1[t],x2[t]},t,IncludeSingularSolution
```

$$x1(t) \rightarrow \frac{1}{2}e^{2t}(2c_1 \cos(2t) - (c_1 + 5c_2) \sin(2t))$$

$$x2(t) \rightarrow \frac{1}{2}e^{2t}(2c_2 \cos(2t) + (c_1 + c_2) \sin(2t))$$

4.13 problem problem 13

Internal problem ID [327]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 13.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 5x_1(t) - 9x_2(t)$$

$$x_2'(t) = 2x_1(t) - x_2(t)$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 58

```
dsolve([diff(x__1(t),t)=5*x__1(t)-9*x__2(t),diff(x__2(t),t)=2*x__1(t)-x__2(t)], [x__1(t), x__2(t)])
```

$$x_1(t) = \frac{3e^{2t}(c_1 \cos(3t) + c_2 \cos(3t) + c_1 \sin(3t) - c_2 \sin(3t))}{2}$$

$$x_2(t) = e^{2t}(c_1 \sin(3t) + c_2 \cos(3t))$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 66

```
DSolve[{x1'[t]==5*x1[t]-9*x2[t],x2'[t]==2*x1[t]-x2[t]},{x1[t],x2[t]},t,IncludeSingularSolutions->True]
```

$$x1(t) \rightarrow e^{2t}(c_1 \cos(3t) + (c_1 - 3c_2) \sin(3t))$$

$$x2(t) \rightarrow \frac{1}{3}e^{2t}(3c_2 \cos(3t) + (2c_1 - 3c_2) \sin(3t))$$

4.14 problem problem 14

Internal problem ID [328]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 14.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 3x_1(t) - 4x_2(t)$$

$$x_2'(t) = 4x_1(t) + 3x_2(t)$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 45

```
dsolve([diff(x__1(t),t)=3*x__1(t)-4*x__2(t),diff(x__2(t),t)=4*x__1(t)+3*x__2(t)],[x__1(t), x
```

$$x_1(t) = e^{3t}(c_1 \cos(4t) - c_2 \sin(4t))$$

$$x_2(t) = e^{3t}(c_1 \sin(4t) + c_2 \cos(4t))$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 51

```
DSolve[{x1'[t]==3*x1[t]-4*x2[t],x2'[t]==4*x1[t]+3*x2[t]},{x1[t],x2[t]},t,IncludeSingularSolu
```

$$x1(t) \rightarrow e^{3t}(c_1 \cos(4t) - c_2 \sin(4t))$$

$$x2(t) \rightarrow e^{3t}(c_2 \cos(4t) + c_1 \sin(4t))$$

4.15 problem problem 15

Internal problem ID [329]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 15.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 7x_1(t) - 5x_2(t)$$

$$x_2'(t) = 4x_1(t) + 3x_2(t)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 59

```
dsolve([diff(x__1(t),t)=7*x__1(t)-5*x__2(t),diff(x__2(t),t)=4*x__1(t)+3*x__2(t)],[x__1(t), x
```

$$x_1(t) = \frac{e^{5t}(2c_1 \cos(4t) - 2c_2 \sin(4t) + c_1 \sin(4t) + c_2 \cos(4t))}{2}$$

$$x_2(t) = e^{5t}(c_1 \sin(4t) + c_2 \cos(4t))$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 72

```
DSolve[{x1'[t]==7*x1[t]-5*x2[t],x2'[t]==4*x1[t]+3*x2[t]},{x1[t],x2[t]},t,IncludeSingularSolu
```

$$x1(t) \rightarrow \frac{1}{4}e^{5t}(4c_1 \cos(4t) + (2c_1 - 5c_2) \sin(4t))$$

$$x2(t) \rightarrow \frac{1}{2}e^{5t}(2c_2 \cos(4t) + (2c_1 - c_2) \sin(4t))$$

4.16 problem problem 16

Internal problem ID [330]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 16.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = -50x_1(t) + 20x_2(t)$$

$$x_2'(t) = 100x_1(t) - 60x_2(t)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 36

```
dsolve([diff(x__1(t),t)=-50*x__1(t)+20*x__2(t),diff(x__2(t),t)=100*x__1(t)-60*x__2(t)], [x__1
```

$$x_1(t) = -\frac{2c_1e^{-100t}}{5} + \frac{c_2e^{-10t}}{2}$$

$$x_2(t) = c_1e^{-100t} + c_2e^{-10t}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 74

```
DSolve[{x1'[t]==-50*x1[t]+20*x2[t],x2'[t]==100*x1[t]-60*x2[t]},{x1[t],x2[t]},t,IncludeSingular
```

$$x_1(t) \rightarrow \frac{1}{9}e^{-100t}(c_1(5e^{90t} + 4) + 2c_2(e^{90t} - 1))$$

$$x_2(t) \rightarrow \frac{1}{9}e^{-100t}(10c_1(e^{90t} - 1) + c_2(4e^{90t} + 5))$$

4.17 problem problem 17

Internal problem ID [331]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 17.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 4x_1(t) + x_2(t) + 4x_3(t)$$

$$x_2'(t) = x_1(t) + 7x_2(t) + x_3(t)$$

$$x_3'(t) = 4x_1(t) + x_2(t) + 4x_3(t)$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 55

```
dsolve([diff(x__1(t),t)=4*x__1(t)+x__2(t)+4*x__3(t),diff(x__2(t),t)=x__1(t)+7*x__2(t)+x__3(t),diff(x__3(t),t)=4*x__1(t)+x__2(t)+4*x__3(t))
```

$$x_1(t) = c_2 e^{9t} + c_3 e^{6t} - c_1$$

$$x_2(t) = c_2 e^{9t} - 2c_3 e^{6t}$$

$$x_3(t) = c_1 + c_2 e^{9t} + c_3 e^{6t}$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 158

```
DSolve[{x1'[t]==4*x1[t]+x2[t]+4*x3[t],x2'[t]==x1[t]+7*x2[t]+x3[t],x3'[t]==4*x1[t]+x2[t]+4*x3[t]}
```

$$x1(t) \rightarrow \frac{1}{6}(c_1(e^{6t} + 2e^{9t} + 3) + (e^{3t} - 1)(3c_3e^{3t} + 2(c_2 + c_3)e^{6t} + 3c_3))$$

$$x2(t) \rightarrow \frac{1}{3}e^{6t}(c_1(e^{3t} - 1) + c_2(e^{3t} + 2) + c_3(e^{3t} - 1))$$

$$x3(t) \rightarrow \frac{1}{6}(c_1(e^{6t} + 2e^{9t} - 3) + (c_3 - 2c_2)e^{6t} + 2(c_2 + c_3)e^{9t} + 3c_3)$$

4.18 problem problem 18

Internal problem ID [332]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 18.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = x_1(t) + 2x_2(t) + 2x_3(t)$$

$$x_2'(t) = 2x_1(t) + 7x_2(t) + x_3(t)$$

$$x_3'(t) = 2x_1(t) + x_2(t) + 7x_3(t)$$

✓ Solution by Maple

Time used: 0.281 (sec). Leaf size: 51

```
dsolve([diff(x__1(t),t)=x__1(t)+2*x__2(t)+2*x__3(t),diff(x__2(t),t)=2*x__1(t)+7*x__2(t)+x__3
```

$$x_1(t) = \frac{c_2 e^{9t}}{2} - 4c_1$$

$$x_2(t) = c_2 e^{9t} - c_3 e^{6t} + c_1$$

$$x_3(t) = c_1 + c_2 e^{9t} + c_3 e^{6t}$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 148

```
DSolve[{x1'[t]==x1[t]+2*x2[t]+2*x3[t],x2'[t]==2*x1[t]+7*x2[t]+x3[t],x3'[t]==2*x1[t]+x2[t]+7*x3[t]}
```

$$x1(t) \rightarrow \frac{1}{9}(c_1(e^{9t} + 8) + 2(c_2 + c_3)(e^{9t} - 1))$$

$$x2(t) \rightarrow \frac{1}{18}(4c_1(e^{9t} - 1) + c_2(9e^{6t} + 8e^{9t} + 1) + c_3(-9e^{6t} + 8e^{9t} + 1))$$

$$x3(t) \rightarrow \frac{1}{18}(4c_1(e^{9t} - 1) + c_2(-9e^{6t} + 8e^{9t} + 1) + c_3(9e^{6t} + 8e^{9t} + 1))$$

4.19 problem problem 19

Internal problem ID [333]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 19.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 4x_1(t) + x_2(t) + x_3(t)$$

$$x_2'(t) = x_1(t) + 4x_2(t) + x_3(t)$$

$$x_3'(t) = x_1(t) + x_2(t) + 4x_3(t)$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 64

```
dsolve([diff(x__1(t),t)=4*x__1(t)+1*x__2(t)+1*x__3(t),diff(x__2(t),t)=1*x__1(t)+4*x__2(t)+1*
```

$$x_1(t) = -2c_2e^{3t} + c_3e^{6t} - c_1e^{3t}$$

$$x_2(t) = c_2e^{3t} + c_3e^{6t} + c_1e^{3t}$$

$$x_3(t) = c_2e^{3t} + c_3e^{6t}$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 124

```
DSolve[{x1'[t]==4*x1[t]+1*x2[t]+1*x3[t],x2'[t]==1*x1[t]+4*x2[t]+1*x3[t],x3'[t]==1*x1[t]+1*x2
```

$$x1(t) \rightarrow \frac{1}{3}e^{3t}(c_1(e^{3t} + 2) + (c_2 + c_3)(e^{3t} - 1))$$

$$x2(t) \rightarrow \frac{1}{3}e^{3t}(c_1(e^{3t} - 1) + c_2(e^{3t} + 2) + c_3(e^{3t} - 1))$$

$$x3(t) \rightarrow \frac{1}{3}e^{3t}(c_1(e^{3t} - 1) + c_2(e^{3t} - 1) + c_3(e^{3t} + 2))$$

4.20 problem problem 20

Internal problem ID [334]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 20.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 5x_1(t) + x_2(t) + 3x_3(t)$$

$$x_2'(t) = x_1(t) + 7x_2(t) + x_3(t)$$

$$x_3'(t) = 3x_1(t) + x_2(t) + 5x_3(t)$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 64

```
dsolve([diff(x__1(t),t)=5*x__1(t)+1*x__2(t)+3*x__3(t),diff(x__2(t),t)=1*x__1(t)+7*x__2(t)+1*
```

$$x_1(t) = c_1 e^{9t} - c_2 e^{2t} + c_3 e^{6t}$$

$$x_2(t) = c_1 e^{9t} - 2c_3 e^{6t}$$

$$x_3(t) = c_1 e^{9t} + c_2 e^{2t} + c_3 e^{6t}$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 163

```
DSolve[{x1'[t]==5*x1[t]+1*x2[t]+3*x3[t],x2'[t]==1*x1[t]+7*x2[t]+1*x3[t],x3'[t]==3*x1[t]+1*x2[t]}
```

$$x1(t) \rightarrow \frac{1}{6}e^{2t}(c_1(e^{4t} + 2e^{7t} + 3) + (c_3 - 2c_2)e^{4t} + 2(c_2 + c_3)e^{7t} - 3c_3)$$

$$x2(t) \rightarrow \frac{1}{3}e^{6t}(c_1(e^{3t} - 1) + c_2(e^{3t} + 2) + c_3(e^{3t} - 1))$$

$$x3(t) \rightarrow \frac{1}{6}e^{2t}(c_1(e^{4t} + 2e^{7t} - 3) + (c_3 - 2c_2)e^{4t} + 2(c_2 + c_3)e^{7t} + 3c_3)$$

4.21 problem problem 21

Internal problem ID [335]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 21.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 5x_1(t) - 6x_3(t)$$

$$x_2'(t) = 2x_1(t) - x_2(t) - 2x_3(t)$$

$$x_3'(t) = 4x_1(t) - 2x_2(t) - 4x_3(t)$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 54

```
dsolve([diff(x__1(t),t)=5*x__1(t)+0*x__2(t)-6*x__3(t),diff(x__2(t),t)=2*x__1(t)-1*x__2(t)-2*
```

$$x_1(t) = \frac{3c_2e^t}{2} + c_3e^{-t} + \frac{6c_1}{5}$$

$$x_2(t) = \frac{c_2e^t}{2} + \frac{c_3e^{-t}}{2} + \frac{2c_1}{5}$$

$$x_3(t) = c_1 + c_2e^t + c_3e^{-t}$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 139

```
DSolve[{x1'[t]==5*x1[t]+0*x2[t]-6*x3[t],x2'[t]==2*x1[t]-1*x2[t]-2*x3[t],x3'[t]==4*x1[t]-2*x2[t]}
```

$$x1(t) \rightarrow e^{-t}(c_1(3e^{2t} - 2) + 6(e^t - 1)(c_2(e^t - 1) - c_3e^t))$$

$$x2(t) \rightarrow e^{-t}(c_1(e^{2t} - 1) + c_2(-4e^t + 2e^{2t} + 3) - 2c_3e^t(e^t - 1))$$

$$x3(t) \rightarrow -2(c_1 - 3c_2)e^{-t} + 2(c_1 + 2c_2 - 2c_3)e^t + 5(c_3 - 2c_2)$$

4.22 problem problem 22

Internal problem ID [336]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 22.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 3x_1(t) + 2x_2(t) + 2x_3(t)$$

$$x_2'(t) = -5x_1(t) - 4x_2(t) - 2x_3(t)$$

$$x_3'(t) = 5x_1(t) + 5x_2(t) + 3x_3(t)$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 55

```
dsolve([diff(x__1(t),t)=3*x__1(t)+2*x__2(t)+2*x__3(t),diff(x__2(t),t)=-5*x__1(t)-4*x__2(t)-2
```

$$x_1(t) = c_3 e^{3t} - c_1 e^t$$

$$x_2(t) = -e^{-2t} c_2 - c_3 e^{3t} + c_1 e^t$$

$$x_3(t) = e^{-2t} c_2 + c_3 e^{3t}$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 123

```
DSolve[{x1'[t]==3*x1[t]+2*x2[t]+2*x3[t],x2'[t]==-5*x1[t]-4*x2[t]-2*x3[t],x3'[t]==5*x1[t]+5*x
```

$$x1(t) \rightarrow e^t((c_1 + c_2 + c_3)e^{2t} - c_2 - c_3)$$

$$x2(t) \rightarrow e^{-2t}(-(c_1(e^{5t} - 1)) + c_2(e^{3t} - e^{5t} + 1) - c_3 e^{3t}(e^{2t} - 1))$$

$$x3(t) \rightarrow e^{-2t}(c_1(e^{5t} - 1) + c_2(e^{5t} - 1) + c_3 e^{5t})$$

4.23 problem problem 23

Internal problem ID [337]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 23.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= 3x_1(t) + x_2(t) + x_3(t) \\x_2'(t) &= -5x_1(t) - 3x_2(t) - x_3(t) \\x_3'(t) &= 5x_1(t) + 5x_2(t) + 3x_3(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.046 (sec). Leaf size: 59

```
dsolve([diff(x__1(t),t)=3*x__1(t)+1*x__2(t)+1*x__3(t),diff(x__2(t),t)=-5*x__1(t)-3*x__2(t)-1
```

$$x_1(t) = c_3e^{3t} - c_1e^{2t}$$

$$x_2(t) = -e^{-2t}c_2 - c_3e^{3t} + c_1e^{2t}$$

$$x_3(t) = e^{-2t}c_2 + c_3e^{3t}$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 121

```
DSolve[{x1'[t]==3*x1[t]+1*x2[t]+1*x3[t],x2'[t]==-5*x1[t]-3*x2[t]-1*x3[t],x3'[t]==5*x1[t]+5*x
```

$$x1(t) \rightarrow e^{2t}((c_1 + c_2 + c_3)e^t - c_2 - c_3)$$

$$x2(t) \rightarrow e^{-2t}(-(c_1(e^{5t} - 1)) + c_2(e^{4t} - e^{5t} + 1) - c_3e^{4t}(e^t - 1))$$

$$x3(t) \rightarrow e^{-2t}(c_1(e^{5t} - 1) + c_2(e^{5t} - 1) + c_3e^{5t})$$

4.24 problem problem 24

Internal problem ID [338]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 24.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 2x_1(t) + x_2(t) - x_3(t)$$

$$x_2'(t) = -4x_1(t) - 3x_2(t) - x_3(t)$$

$$x_3'(t) = 4x_1(t) + 4x_2(t) + 2x_3(t)$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 77

```
dsolve([diff(x__1(t),t)=2*x__1(t)+1*x__2(t)-1*x__3(t),diff(x__2(t),t)=-4*x__1(t)-3*x__2(t)-1*x__3(t),diff(x__3(t),t)=4*x__1(t)+4*x__2(t)+2*x__3(t)),x__1(t),x__2(t),x__3(t))
```

$$x_1(t) = \frac{c_2 \cos(2t)}{2} - \frac{c_3 \sin(2t)}{2} + \frac{c_2 \sin(2t)}{2} + \frac{c_3 \cos(2t)}{2} - c_1 e^t$$

$$x_2(t) = -c_2 \sin(2t) - c_3 \cos(2t) + c_1 e^t$$

$$x_3(t) = c_2 \sin(2t) + c_3 \cos(2t)$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 103

```
DSolve[{x1'[t]==2*x1[t]+1*x2[t]-1*x3[t],x2'[t]==-4*x1[t]-3*x2[t]-1*x3[t],x3'[t]==4*x1[t]+4*x
```

$$x1(t) \rightarrow (c_2 + c_3) (-e^t) + (c_1 + c_2 + c_3) \cos(2t) + (c_1 + c_2) \sin(2t)$$

$$x2(t) \rightarrow (c_2 + c_3)e^t - c_3 \cos(2t) - (2c_1 + 2c_2 + c_3) \sin(2t)$$

$$x3(t) \rightarrow c_3 \cos(2t) + (2c_1 + 2c_2 + c_3) \sin(2t)$$

4.25 problem problem 25

Internal problem ID [339]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 25.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 5x_1(t) + 5x_2(t) + 2x_3(t)$$

$$x_2'(t) = -6x_1(t) - 6x_2(t) - 5x_3(t)$$

$$x_3'(t) = 6x_1(t) + 6x_2(t) + 5x_3(t)$$

✓ Solution by Maple

Time used: 0.046 (sec). Leaf size: 101

```
dsolve([diff(x__1(t),t)=5*x__1(t)+5*x__2(t)+2*x__3(t),diff(x__2(t),t)=-6*x__1(t)-6*x__2(t)-5
```

$$x_1(t) = \frac{c_2 e^{2t} \sin(3t)}{2} + \frac{c_2 e^{2t} \cos(3t)}{2} + \frac{c_3 e^{2t} \cos(3t)}{2} - \frac{c_3 e^{2t} \sin(3t)}{2} - c_1$$

$$x_2(t) = -c_2 e^{2t} \sin(3t) - c_3 e^{2t} \cos(3t) + c_1$$

$$x_3(t) = e^{2t}(c_2 \sin(3t) + c_3 \cos(3t))$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 122

```
DSolve[{x1'[t]==5*x1[t]+5*x2[t]+2*x3[t],x2'[t]==-6*x1[t]-6*x2[t]-5*x3[t],x3'[t]==6*x1[t]+6*x
```

$$x1(t) \rightarrow (c_1 + c_2 + c_3)e^{2t} \cos(3t) + (c_1 + c_2)e^{2t} \sin(3t) - c_2 - c_3$$

$$x2(t) \rightarrow -c_3e^{2t} \cos(3t) - (2c_1 + 2c_2 + c_3)e^{2t} \sin(3t) + c_2 + c_3$$

$$x3(t) \rightarrow e^{2t}(c_3 \cos(3t) + (2c_1 + 2c_2 + c_3) \sin(3t))$$

4.26 problem problem 26

Internal problem ID [340]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 26.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 3x_1(t) + x_3(t)$$

$$x_2'(t) = 9x_1(t) - x_2(t) + 2x_3(t)$$

$$x_3'(t) = -9x_1(t) + 4x_2(t) - x_3(t)$$

With initial conditions

$$[x_1(0) = 0, x_2(0) = 0, x_3(0) = 17]$$

✓ Solution by Maple

Time used: 0.125 (sec). Leaf size: 64

```
dsolve([diff(x__1(t),t) = 3*x__1(t)+x__3(t), diff(x__2(t),t) = 9*x__1(t)-x__2(t)+2*x__3(t),
```

$$x_1(t) = e^{-t} \sin(t) - 4e^{-t} \cos(t) + 4e^{3t}$$

$$x_2(t) = -9e^{-t} \cos(t) - 2e^{-t} \sin(t) + 9e^{3t}$$

$$x_3(t) = 17e^{-t} \cos(t)$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 62

```
DSolve[{x1'[t]==3*x1[t]+0*x2[t]+1*x3[t],x2'[t]==9*x1[t]-1*x2[t]+2*x3[t],x3'[t]==-9*x1[t]+4*x
```

$$x1(t) \rightarrow e^{-t}(4e^{4t} + \sin(t) - 4 \cos(t))$$

$$x2(t) \rightarrow e^{-t}(9e^{4t} - 2 \sin(t) - 9 \cos(t))$$

$$x3(t) \rightarrow 17e^{-t} \cos(t)$$

4.27 problem problem 38

Internal problem ID [341]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 38.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = x_1(t)$$

$$x_2'(t) = 2x_1(t) + 2x_2(t)$$

$$x_3'(t) = 3x_2(t) + 3x_3(t)$$

$$x_4'(t) = 4x_3(t) + 4x_4(t)$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 75

```
dsolve([diff(x__1(t),t)=x__1(t)+0*x__2(t)+0*x__3(t)+0*x__4(t),diff(x__2(t),t)=2*x__1(t)+2*x__2(t),diff(x__3(t),t)=3*x__2(t)+3*x__3(t),diff(x__4(t),t)=4*x__3(t)+4*x__4(t)),t)
```

$$x_1(t) = -\frac{c_1 e^t}{4}$$

$$x_2(t) = \frac{c_1 e^t}{2} + \frac{c_2 e^{2t}}{6}$$

$$x_3(t) = -\frac{3c_1 e^t}{4} - \frac{c_2 e^{2t}}{2} - \frac{c_4 e^{3t}}{4}$$

$$x_4(t) = c_1 e^t + c_2 e^{2t} + c_3 e^{4t} + c_4 e^{3t}$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 128

```
DSolve[{x1'[t]==1*x1[t]+0*x2[t]+0*x3[t]+0*x4[t],x2'[t]==2*x1[t]+2*x2[t]+0*x3[t]+0*x4[t],x3'[t]==3*x1[t]+3*x2[t]+3*x3[t]+0*x4[t],x4'[t]==4*x1[t]+4*x2[t]+4*x3[t]+4*x4[t]}
```

$$x1(t) \rightarrow c_1 e^t$$

$$x2(t) \rightarrow e^t (2c_1 (e^t - 1) + c_2 e^t)$$

$$x3(t) \rightarrow e^t (3c_1 (e^t - 1)^2 + e^t (3c_2 (e^t - 1) + c_3 e^t))$$

$$x4(t) \rightarrow e^t (4c_1 (e^t - 1)^3 + e^t (6c_2 (e^t - 1)^2 + e^t (4c_3 (e^t - 1) + c_4 e^t)))$$

4.28 problem problem 39

Internal problem ID [342]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 39.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = -2x_1(t) + 9x_4(t)$$

$$x_2'(t) = 4x_1(t) + 2x_2(t) - 10x_4(t)$$

$$x_3'(t) = -x_3(t) + 8x_4(t)$$

$$x_4'(t) = x_4(t)$$

✓ Solution by Maple

Time used: 0.094 (sec). Leaf size: 61

```
dsolve([diff(x__1(t),t)=-2*x__1(t)+0*x__2(t)+0*x__3(t)+9*x__4(t),diff(x__2(t),t)=4*x__1(t)+2
```

$$x_1(t) = -c_1e^{-2t} + 3c_4e^t$$

$$x_2(t) = c_2e^{2t} + c_1e^{-2t} - 2c_4e^t$$

$$x_3(t) = 4c_4e^t + c_3e^{-t}$$

$$x_4(t) = c_4e^t$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 103

```
DSolve[{x1'[t]==-2*x1[t]+0*x2[t]+0*x3[t]+9*x4[t],x2'[t]==4*x1[t]+2*x2[t]+0*x3[t]-10*x4[t],x3
```

$$x1(t) \rightarrow e^{-2t}(3c_4(e^{3t} - 1) + c_1)$$

$$x2(t) \rightarrow e^{-2t}(c_1(e^{4t} - 1) + (c_2 - c_4)e^{4t} - 2c_4e^{3t} + 3c_4)$$

$$x3(t) \rightarrow e^{-t}(4c_4(e^{2t} - 1) + c_3)$$

$$x4(t) \rightarrow c_4e^t$$

4.29 problem problem 40

Internal problem ID [343]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 40.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 2x_1(t)$$

$$x_2'(t) = -21x_1(t) - 5x_2(t) - 27x_3(t) - 9x_4(t)$$

$$x_3'(t) = 5x_3(t)$$

$$x_4'(t) = -21x_3(t) - 2x_4(t)$$

✓ Solution by Maple

Time used: 0.11 (sec). Leaf size: 61

```
dsolve([diff(x__1(t),t)=2*x__1(t)+0*x__2(t)+0*x__3(t)+0*x__4(t),diff(x__2(t),t)=-21*x__1(t)-
```

$$x_1(t) = -\frac{c_2 e^{2t}}{3}$$

$$x_2(t) = c_2 e^{2t} + e^{-5t} c_1 - 3c_3 e^{-2t}$$

$$x_3(t) = -\frac{c_4 e^{5t}}{3}$$

$$x_4(t) = c_3 e^{-2t} + c_4 e^{5t}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 86

```
DSolve[{x1'[t]==2*x1[t]+0*x2[t]+0*x3[t]+0*x4[t],x2'[t]==-21*x1[t]-5*x2[t]-27*x3[t]-9*x4[t],x3'[t]==-21*x1[t]-5*x2[t]-27*x3[t]-9*x4[t],x4'[t]==-21*x1[t]-5*x2[t]-27*x3[t]-9*x4[t]}
```

$$x1(t) \rightarrow c_1 e^{2t}$$

$$x2(t) \rightarrow e^{-5t}(-3c_1(e^{7t} - 1) - 3(3c_3 + c_4)(e^{3t} - 1) + c_2)$$

$$x3(t) \rightarrow c_3 e^{5t}$$

$$x4(t) \rightarrow e^{-2t}(c_4 - 3c_3(e^{7t} - 1))$$

4.30 problem problem 41

Internal problem ID [344]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 41.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= 4x_1(t) + x_2(t) + x_3(t) + 7x_4(t) \\x_2'(t) &= x_1(t) + 4x_2(t) + 10x_3(t) + x_4(t) \\x_3'(t) &= x_1(t) + 10x_2(t) + 4x_3(t) + x_4(t) \\x_4'(t) &= 7x_1(t) + x_2(t) + x_3(t) + 4x_4(t)\end{aligned}$$

With initial conditions

$$[x_1(0) = 3, x_2(0) = 1, x_3(0) = 1, x_4(0) = 3]$$

✓ Solution by Maple

Time used: 0.094 (sec). Leaf size: 62

```
dsolve([diff(x__1(t),t) = 4*x__1(t)+x__2(t)+x__3(t)+7*x__4(t), diff(x__2(t),t) = x__1(t)+4*x__2(t)+10*x__3(t)+x__4(t), diff(x__3(t),t) = x__1(t)+10*x__2(t)+4*x__3(t)+x__4(t), diff(x__4(t),t) = 7*x__1(t)+x__2(t)+x__3(t)+4*x__4(t)], [x__1(t), x__2(t), x__3(t), x__4(t)])
```

$$x_1(t) = e^{15t} + 2e^{10t}$$

$$x_2(t) = 2e^{15t} - e^{10t}$$

$$x_3(t) = 2e^{15t} - e^{10t}$$

$$x_4(t) = e^{15t} + 2e^{10t}$$

✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 70

```
DSolve[{x1'[t]==4*x1[t]+1*x2[t]+1*x3[t]+7*x4[t],x2'[t]==1*x1[t]+4*x2[t]+10*x3[t]+1*x4[t],x3'
```

$$x1(t) \rightarrow e^{10t}(e^{5t} + 2)$$

$$x2(t) \rightarrow e^{10t}(2e^{5t} - 1)$$

$$x3(t) \rightarrow e^{10t}(2e^{5t} - 1)$$

$$x4(t) \rightarrow e^{10t}(e^{5t} + 2)$$

4.31 problem problem 42

Internal problem ID [345]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 42.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = -40x_1(t) - 12x_2(t) + 54x_3(t)$$

$$x_2'(t) = 35x_1(t) + 13x_2(t) - 46x_3(t)$$

$$x_3'(t) = -25x_1(t) - 7x_2(t) + 34x_3(t)$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 59

```
dsolve([diff(x__1(t),t)=-40*x__1(t)-12*x__2(t)+54*x__3(t),diff(x__2(t),t)=35*x__1(t)+13*x__2
```

$$x_1(t) = c_2e^{2t} + 2c_3e^{5t} + \frac{3c_1}{2}$$

$$x_2(t) = c_2e^{2t} - 3c_3e^{5t} - \frac{c_1}{2}$$

$$x_3(t) = c_1 + c_2e^{2t} + c_3e^{5t}$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 181

```
DSolve[{x1'[t]==-40*x1[t]-12*x2[t]+54*x3[t],x2'[t]==35*x1[t]+13*x2[t]-46*x3[t],x3'[t]==-25*x
```

$$x1(t) \rightarrow c_1(-5e^{2t} - 6e^{5t} + 12) - c_2(e^{2t} + 2e^{5t} - 3) + c_3(7e^{2t} + 8e^{5t} - 15)$$

$$x2(t) \rightarrow c_1(-5e^{2t} + 9e^{5t} - 4) + c_2(-e^{2t} + 3e^{5t} - 1) + c_3(7e^{2t} - 12e^{5t} + 5)$$

$$x3(t) \rightarrow c_1(-5e^{2t} - 3e^{5t} + 8) - c_2(e^{2t} + e^{5t} - 2) + c_3(7e^{2t} + 4e^{5t} - 10)$$

4.32 problem problem 43

Internal problem ID [346]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 43.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = -20x_1(t) + 11x_2(t) + 13x_3(t)$$

$$x_2'(t) = 12x_1(t) - x_2(t) - 7x_3(t)$$

$$x_3'(t) = -48x_1(t) + 21x_2(t) + 31x_3(t)$$

✓ Solution by Maple

Time used: 0.046 (sec). Leaf size: 72

```
dsolve([diff(x__1(t),t)=-20*x__1(t)+11*x__2(t)+13*x__3(t),diff(x__2(t),t)=12*x__1(t)-1*x__2(t)
```

$$x_1(t) = \frac{3c_1e^{-2t}}{5} + c_2e^{4t} + \frac{c_3e^{8t}}{3}$$

$$x_2(t) = -\frac{c_1e^{-2t}}{5} + c_2e^{4t} - \frac{c_3e^{8t}}{3}$$

$$x_3(t) = c_1e^{-2t} + c_2e^{4t} + c_3e^{8t}$$

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 554

`DSolve[{x1'[t]==20*x1[t]+11*x2[t]+13*x3[t],x2'[t]==12*x1[t]-1*x2[t]-7*x3[t],x3'[t]==-48*x1[t]`

$$\begin{aligned}
 x_1(t) &\rightarrow c_2 \text{RootSum} \left[\#1^3 - 50\#1^2 + 1208\#1 - 4576\&, \frac{11\#1e^{\#1t} - 68e^{\#1t}}{3\#1^2 - 100\#1 + 1208}\& \right] \\
 &+ c_3 \text{RootSum} \left[\#1^3 - 50\#1^2 + 1208\#1 - 4576\&, \frac{13\#1e^{\#1t} - 64e^{\#1t}}{3\#1^2 - 100\#1 + 1208}\& \right] \\
 &+ c_1 \text{RootSum} \left[\#1^3 - 50\#1^2 + 1208\#1 \right. \\
 &\quad \left. - 4576\&, \frac{\#1^2e^{\#1t} - 30\#1e^{\#1t} + 116e^{\#1t}}{3\#1^2 - 100\#1 + 1208}\& \right] \\
 x_2(t) &\rightarrow 12c_1 \text{RootSum} \left[\#1^3 - 50\#1^2 + 1208\#1 - 4576\&, \frac{\#1e^{\#1t} - 3e^{\#1t}}{3\#1^2 - 100\#1 + 1208}\& \right] \\
 &- c_3 \text{RootSum} \left[\#1^3 - 50\#1^2 + 1208\#1 - 4576\&, \frac{7\#1e^{\#1t} - 296e^{\#1t}}{3\#1^2 - 100\#1 + 1208}\& \right] \\
 &+ c_2 \text{RootSum} \left[\#1^3 - 50\#1^2 + 1208\#1 \right. \\
 &\quad \left. - 4576\&, \frac{\#1^2e^{\#1t} - 51\#1e^{\#1t} + 1244e^{\#1t}}{3\#1^2 - 100\#1 + 1208}\& \right] \\
 x_3(t) &\rightarrow -12c_1 \text{RootSum} \left[\#1^3 - 50\#1^2 + 1208\#1 - 4576\&, \frac{4\#1e^{\#1t} - 17e^{\#1t}}{3\#1^2 - 100\#1 + 1208}\& \right] \\
 &+ 3c_2 \text{RootSum} \left[\#1^3 - 50\#1^2 + 1208\#1 - 4576\&, \frac{7\#1e^{\#1t} - 316e^{\#1t}}{3\#1^2 - 100\#1 + 1208}\& \right] \\
 &+ c_3 \text{RootSum} \left[\#1^3 - 50\#1^2 + 1208\#1 \right. \\
 &\quad \left. - 4576\&, \frac{\#1^2e^{\#1t} - 19\#1e^{\#1t} - 152e^{\#1t}}{3\#1^2 - 100\#1 + 1208}\& \right]
 \end{aligned}$$

4.33 problem problem 44

Internal problem ID [347]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 44.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 147x_1(t) + 23x_2(t) - 202x_3(t)$$

$$x_2'(t) = -90x_1(t) - 9x_2(t) + 129x_3(t)$$

$$x_3'(t) = 90x_1(t) + 15x_2(t) - 123x_3(t)$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 74

```
dsolve([diff(x__1(t),t)=147*x__1(t)+23*x__2(t)-202*x__3(t),diff(x__2(t),t)=-90*x__1(t)-9*x__
```

$$x_1(t) = \frac{5c_1e^{12t}}{3} + \frac{3c_2e^{-3t}}{2} + \frac{7c_3e^{6t}}{5}$$

$$x_2(t) = -c_1e^{12t} - c_2e^{-3t} + \frac{c_3e^{6t}}{5}$$

$$x_3(t) = c_1e^{12t} + c_2e^{-3t} + c_3e^{6t}$$

✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 188

```
DSolve[{x1'[t]==147*x1[t]+23*x2[t]-202*x3[t],x2'[t]==-90*x1[t]-9*x2[t]+129*x3[t],x3'[t]==90*
```

$$x1(t) \rightarrow \frac{1}{6}e^{-3t}(6c_1(10e^{15t} - 9) + c_2(7e^{9t} + 5e^{15t} - 12) - c_3(-7e^{9t} + 85e^{15t} - 78))$$

$$x2(t) \rightarrow \frac{1}{6}e^{-3t}(-36c_1(e^{15t} - 1) + c_2(e^{9t} - 3e^{15t} + 8) + c_3(e^{9t} + 51e^{15t} - 52))$$

$$x3(t) \rightarrow \frac{1}{6}e^{-3t}(36c_1(e^{15t} - 1) + c_2(5e^{9t} + 3e^{15t} - 8) - c_3(-5e^{9t} + 51e^{15t} - 52))$$

4.34 problem problem 45

Internal problem ID [348]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 45.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 9x_1(t) - 7x_2(t) - 5x_3(t)$$

$$x_2'(t) = -12x_1(t) + 7x_2(t) + 11x_3(t) + 9x_4(t)$$

$$x_3'(t) = 24x_1(t) - 17x_2(t) - 19x_3(t) - 9x_4(t)$$

$$x_4'(t) = -18x_1(t) + 13x_2(t) + 17x_3(t) + 9x_4(t)$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 104

```
dsolve([diff(x__1(t),t)=9*x__1(t)-7*x__2(t)-5*x__3(t)+0*x__4(t),diff(x__2(t),t)=-12*x__1(t)+
```

$$x_1(t) = -c_2e^{-3t} + 2c_3e^{3t} - c_4e^{6t} + c_1$$

$$x_2(t) = -c_2e^{-3t} + c_3e^{3t} + c_4e^{6t} + 2c_1$$

$$x_3(t) = -c_2e^{-3t} + c_3e^{3t} - 2c_4e^{6t} - c_1$$

$$x_4(t) = c_1 + c_2e^{-3t} + c_3e^{3t} + c_4e^{6t}$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 430

`DSolve[{x1'[t]==9*x1[t]-7*x2[t]-5*x3[t]+0*x4[t],x2'[t]==-12*x1[t]+7*x2[t]+11*x3[t]+9*x4[t],x`

$$x1(t) \rightarrow \frac{1}{3}e^{-3t} \left(c_1(6e^{3t} - 6e^{6t} + 6e^{9t} - 3) - (e^{3t} - 1) (c_2(4e^{6t} + 3) + c_3(-3e^{3t} + 5e^{6t} + 3) + 3c_4e^{3t}(e^{3t} - 1)) \right)$$

$$x2(t) \rightarrow \frac{1}{3}e^{-3t} \left(-3c_1(-4e^{3t} + e^{6t} + 2e^{9t} + 1) + c_2(-6e^{3t} + 2e^{6t} + 4e^{9t} + 3) + (e^{3t} - 1) (c_3(9e^{3t} + 5e^{6t} - 3) + 3c_4e^{3t}(e^{3t} + 2)) \right)$$

$$x3(t) \rightarrow c_1(-e^{-3t} - e^{3t} + 4e^{6t} - 2) + c_2 \left(e^{-3t} + \frac{2e^{3t}}{3} - \frac{8e^{6t}}{3} + 1 \right) + c_3e^{-3t} + \frac{4}{3}c_3e^{3t} - \frac{10}{3}c_3e^{6t} + c_4e^{3t} - 2c_4e^{6t} + 2c_3 + c_4$$

$$x4(t) \rightarrow \frac{1}{3} \left(c_1(3e^{-3t} - 3e^{3t} - 6e^{6t} + 6) + c_2(-3e^{-3t} + 2e^{3t} + 4e^{6t} - 3) - 3c_3e^{-3t} + 4c_3e^{3t} + 5c_3e^{6t} + 3c_4e^{3t} + 3c_4e^{6t} - 6c_3 - 3c_4 \right)$$

4.35 problem problem 46

Internal problem ID [349]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 46.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 13x_1(t) - 42x_2(t) + 106x_3(t) + 139x_4(t)$$

$$x_2'(t) = 2x_1(t) - 16x_2(t) + 52x_3(t) + 70x_4(t)$$

$$x_3'(t) = x_1(t) + 6x_2(t) - 20x_3(t) - 31x_4(t)$$

$$x_4'(t) = -x_1(t) - 6x_2(t) + 22x_3(t) + 33x_4(t)$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 124

```
dsolve([diff(x__1(t),t)=13*x__1(t)-42*x__2(t)+106*x__3(t)+139*x__4(t),diff(x__2(t),t)=2*x__1
```

$$x_1(t) = -c_1e^{2t} + c_2e^{4t} + 3c_3e^{-4t} - c_4e^{8t}$$

$$x_2(t) = -2c_1e^{2t} + c_2e^{4t} + 2c_3e^{-4t} + \frac{2c_4e^{8t}}{3}$$

$$x_3(t) = -2c_1e^{2t} - c_2e^{4t} - c_3e^{-4t} - c_4e^{8t}$$

$$x_4(t) = c_1e^{2t} + c_2e^{4t} + c_3e^{-4t} + c_4e^{8t}$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 449

```
DSolve[{x1'[t]==13*x1[t]-42*x2[t]+106*x3[t]+139*x4[t],x2'[t]==2*x1[t]-16*x2[t]+52*x3[t]+70*x4[t]}
```

$$x1(t) \rightarrow \frac{1}{4}e^{-4t} (c_1(4e^{8t} + 3e^{12t} - 3) - 6c_2(2e^{8t} + e^{12t} - 3) + 4c_3e^{6t} + 32c_3e^{8t} + 12c_3e^{12t} + 4c_4e^{6t} + 44c_4e^{8t} + 15c_4e^{12t} - 48c_3 - 63c_4)$$

$$x2(t) \rightarrow \frac{1}{2}e^{-4t} (-c_1(-2e^{8t} + e^{12t} + 1)) + 2c_2(-3e^{8t} + e^{12t} + 3) + 4c_3e^{6t} + 16c_3e^{8t} - 4c_3e^{12t} + 4c_4e^{6t} + 22c_4e^{8t} - 5c_4e^{12t} - 16c_3 - 21c_4)$$

$$x3(t) \rightarrow \frac{1}{4}e^{-4t} (c_1(-4e^{8t} + 3e^{12t} + 1) - 6c_2(-2e^{8t} + e^{12t} + 1) + 8c_3e^{6t} - 32c_3e^{8t} + 12c_3e^{12t} + 8c_4e^{6t} - 44c_4e^{8t} + 15c_4e^{12t} + 16c_3 + 21c_4)$$

$$x4(t) \rightarrow \frac{1}{4}e^{-4t} (c_1(4e^{8t} - 3e^{12t} - 1) + 6c_2(-2e^{8t} + e^{12t} + 1) - 4c_3e^{6t} + 32c_3e^{8t} - 12c_3e^{12t} - 4c_4e^{6t} + 44c_4e^{8t} - 15c_4e^{12t} - 16c_3 - 21c_4)$$

4.36 problem problem 47

Internal problem ID [350]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 47.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= 23x_1(t) - 18x_2(t) - 16x_3(t) \\x_2'(t) &= -8x_1(t) + 6x_2(t) + 7x_3(t) + 9x_4(t) \\x_3'(t) &= 34x_1(t) - 27x_2(t) - 26x_3(t) - 9x_4(t) \\x_4'(t) &= -26x_1(t) + 21x_2(t) + 25x_3(t) + 12x_4(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 122

```
dsolve([diff(x__1(t),t)=23*x__1(t)-18*x__2(t)-16*x__3(t)+0*x__4(t),diff(x__2(t),t)=-8*x__1(t)
```

$$x_1(t) = -c_1e^{9t} - 2c_2e^{-3t} + c_3e^{3t} + 2c_4e^{6t}$$

$$x_2(t) = c_1e^{9t} - 2c_2e^{-3t} + 2c_3e^{3t} + c_4e^{6t}$$

$$x_3(t) = -2c_1e^{9t} - c_2e^{-3t} - c_3e^{3t} + c_4e^{6t}$$

$$x_4(t) = c_1e^{9t} + c_2e^{-3t} + c_3e^{3t} + c_4e^{6t}$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 469

```
DSolve[{x1'[t]==23*x1[t]-18*x2[t]-16*x3[t]+0*x4[t],x2'[t]==-8*x1[t]+6*x2[t]+7*x3[t]+9*x4[t],
```

$$x1(t) \rightarrow \frac{1}{3}e^{-3t}(c_1(9e^{6t} - 8e^{9t} + 8e^{12t} - 6) - (e^{3t} - 1)(6c_2(e^{3t} + e^{9t} + 1) + c_3(6e^{3t} - 3e^{6t} + 7e^{9t} + 6) + 3c_4e^{6t}(e^{3t} - 1)))$$

$$x2(t) \rightarrow \frac{1}{3}e^{-3t}(-2c_1(-9e^{6t} + 2e^{9t} + 4e^{12t} + 3) + 3c_2(-4e^{6t} + e^{9t} + 2e^{12t} + 2) + (e^{3t} - 1)(c_3(-6e^{3t} + 12e^{6t} + 7e^{9t} - 6) + 3c_4e^{6t}(e^{3t} + 2)))$$

$$x3(t) \rightarrow \frac{1}{3}e^{-3t}(c_1(-9e^{6t} - 4e^{9t} + 16e^{12t} - 3) + 3c_2(2e^{6t} + e^{9t} - 4e^{12t} + 1) + 9c_3e^{6t} + 5c_3e^{9t} - 14c_3e^{12t} + 3c_4e^{6t} + 3c_4e^{9t} - 6c_4e^{12t} + 3c_3)$$

$$x4(t) \rightarrow \frac{1}{3}e^{-3t}(c_1(9e^{6t} - 4e^{9t} - 8e^{12t} + 3) + 3c_2(-2e^{6t} + e^{9t} + 2e^{12t} - 1) - 9c_3e^{6t} + 5c_3e^{9t} + 7c_3e^{12t} - 3c_4e^{6t} + 3c_4e^{9t} + 3c_4e^{12t} - 3c_3)$$

4.37 problem problem 48

Internal problem ID [351]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 48.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 47x_1(t) - 8x_2(t) + 5x_3(t) - 5x_4(t)$$

$$x_2'(t) = -10x_1(t) + 32x_2(t) + 18x_3(t) - 2x_4(t)$$

$$x_3'(t) = 139x_1(t) - 40x_2(t) - 167x_3(t) - 121x_4(t)$$

$$x_4'(t) = -232x_1(t) + 64x_2(t) + 360x_3(t) + 248x_4(t)$$

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 125

```
dsolve([diff(x__1(t),t)=47*x__1(t)-8*x__2(t)+5*x__3(t)-5*x__4(t),diff(x__2(t),t)=-10*x__1(t)
```

$$x_1(t) = \frac{3c_1e^{48t}}{2} + \frac{c_2e^{16t}}{2} - \frac{c_3e^{64t}}{3} - 2c_4e^{32t}$$

$$x_2(t) = -\frac{c_1e^{48t}}{2} + c_2e^{16t} - \frac{c_3e^{64t}}{3} - 5c_4e^{32t}$$

$$x_3(t) = \frac{c_1e^{48t}}{2} - \frac{c_2e^{16t}}{2} - \frac{2c_3e^{64t}}{3} - c_4e^{32t}$$

$$x_4(t) = c_1e^{48t} + c_2e^{16t} + c_3e^{64t} + c_4e^{32t}$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 448

```
DSolve[{x1'[t]==47*x1[t]-8*x2[t]+5*x3[t]-5*x4[t],x2'[t]==-10*x1[t]+32*x2[t]+18*x3[t]-2*x4[t]}
```

$$x1(t) \rightarrow \frac{1}{16}e^{16t}(c_1(-38e^{16t} - 6e^{32t} + 27e^{48t} + 33) - (e^{16t} - 1)(8c_2(e^{16t} + e^{32t} - 1) + c_3(9e^{16t} + 39e^{32t} - 53) + c_4(7e^{16t} + 25e^{32t} - 27)))$$

$$x2(t) \rightarrow \frac{1}{16}e^{16t}(c_1(-95e^{16t} + 2e^{32t} + 27e^{48t} + 66) - 8c_2(-5e^{16t} + e^{48t} + 2) - (e^{16t} - 1)(c_3(49e^{16t} + 39e^{32t} - 106) + c_4(31e^{16t} + 25e^{32t} - 54)))$$

$$x3(t) \rightarrow \frac{1}{16}e^{16t}(c_1(-19e^{16t} - 2e^{32t} + 54e^{48t} - 33) + 8c_2(e^{16t} - 2e^{48t} + 1) + 31c_3e^{16t} + 10c_3e^{32t} - 78c_3e^{48t} + 17c_4e^{16t} + 6c_4e^{32t} - 50c_4e^{48t} + 53c_3 + 27c_4)$$

$$x4(t) \rightarrow -\frac{1}{16}e^{16t}(c_1(-19e^{16t} + 4e^{32t} + 81e^{48t} - 66) + 8c_2(e^{16t} - 3e^{48t} + 2) + 31c_3e^{16t} - 20c_3e^{32t} - 117c_3e^{48t} + 17c_4e^{16t} - 12c_4e^{32t} - 75c_4e^{48t} + 106c_3 + 54c_4)$$

4.38 problem problem 49

Internal problem ID [352]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 49.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 139x_1(t) - 14x_2(t) - 52x_3(t) - 14x_4(t) + 28x_5(t)$$

$$x_2'(t) = -22x_1(t) + 5x_2(t) + 7x_3(t) + 8x_4(t) - 7x_5(t)$$

$$x_3'(t) = 370x_1(t) - 38x_2(t) - 139x_3(t) - 38x_4(t) + 76x_5(t)$$

$$x_4'(t) = 152x_1(t) - 16x_2(t) - 59x_3(t) - 13x_4(t) + 35x_5(t)$$

$$x_5'(t) = 95x_1(t) - 10x_2(t) - 38x_3(t) - 7x_4(t) + 23x_5(t)$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 130

```
dsolve([diff(x__1(t),t)=139*x__1(t)-14*x__2(t)-52*x__3(t)-14*x__4(t)+28*x__5(t),diff(x__2(t)
```

$$x_1(t) = 2c_2e^{9t} + c_3e^{-3t} + c_4e^{3t}$$

$$x_2(t) = 7c_4e^{3t} + c_5e^{6t} + 3c_1$$

$$x_3(t) = 5c_2e^{9t} + 3c_3e^{-3t} + c_4e^{3t}$$

$$x_4(t) = 2c_2e^{9t} + c_3e^{-3t} + c_4e^{3t} + c_5e^{6t} - c_1$$

$$x_5(t) = c_1 + c_2e^{9t} + c_3e^{-3t} + c_4e^{3t} + c_5e^{6t}$$

✓ Solution by Mathematica

Time used: 0.052 (sec). Leaf size: 2676

```
DSolve[{x1'[t]==139*x1[t]-14*x2[t]-52*x3[t]-14*x4[t]+28*x5[t],x2'[t]==-22*x1[t]+5*x2[t]+7*x3
```

Too large to display

4.39 problem problem 50

Internal problem ID [353]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.3, The eigenvalue method for linear systems. Page 395

Problem number: problem 50.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 9x_1(t) + 13x_2(t) - 13x_6(t)$$

$$x_2'(t) = -14x_1(t) + 19x_2(t) - 10x_3(t) - 20x_4(t) + 10x_5(t) + 4x_6(t)$$

$$x_3'(t) = -30x_1(t) + 12x_2(t) - 7x_3(t) - 30x_4(t) + 12x_5(t) + 18x_6(t)$$

$$x_4'(t) = -12x_1(t) + 10x_2(t) - 10x_3(t) - 9x_4(t) + 10x_5(t) + 2x_6(t)$$

$$x_5'(t) = 6x_1(t) + 9x_2(t) + 6x_4(t) + 5x_5(t) - 15x_6(t)$$

$$x_6'(t) = -14x_1(t) + 23x_2(t) - 10x_3(t) - 20x_4(t) + 10x_5(t)$$

✓ Solution by Maple

Time used: 0.172 (sec). Leaf size: 135

```
dsolve([diff(x__1(t),t)=9*x__1(t)+13*x__2(t)+0*x__3(t)+0*x__4(t)+0*x__5(t)-13*x__6(t),diff(x
```

$$x_1(t) = c_3 e^{9t} + c_5 e^{-4t}$$

$$x_2(t) = c_3 e^{9t} + c_4 e^{3t} + c_6 e^{-7t}$$

$$x_3(t) = c_6 e^{-7t} + c_2 e^{5t} - e^{11t} c_1$$

$$x_4(t) = e^{11t} c_1 + c_4 e^{3t} + c_6 e^{-7t}$$

$$x_5(t) = c_2 e^{5t} + e^{11t} c_1 + c_5 e^{-4t}$$

$$x_6(t) = c_3 e^{9t} + c_4 e^{3t} + c_5 e^{-4t} + c_6 e^{-7t}$$

✓ Solution by Mathematica

Time used: 0.107 (sec). Leaf size: 1882

```
DSolve[{x1'[t]==9*x1[t]+13*x2[t]-13*x6[t],x2'[t]==-14*x1[t]+19*x2[t]-10*x3[t]-20*x4[t]+10*x5
```

Too large to display

5 Section 7.6, Multiple Eigenvalue Solutions.

Examples. Page 437

5.1	problem Example 1	105
5.2	problem Example 3	107
5.3	problem Example 4	108
5.4	problem Example 6	110

5.1 problem Example 1

Internal problem ID [354]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.6, Multiple Eigenvalue Solutions. Examples. Page 437

Problem number: Example 1.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 9x_1(t) + 4x_2(t)$$

$$x_2'(t) = -6x_1(t) - x_2(t)$$

$$x_3'(t) = 6x_1(t) + 4x_2(t) + 3x_3(t)$$

✓ Solution by Maple

Time used: 0.046 (sec). Leaf size: 66

```
dsolve([diff(x__1(t),t)=9*x__1(t)+4*x__2(t)+0*x__3(t),diff(x__2(t),t)=-6*x__1(t)-1*x__2(t)+0
```

$$x_1(t) = c_2 e^{5t} + \frac{2c_3 e^{3t}}{3} - \frac{2c_1 e^{3t}}{3}$$

$$x_2(t) = -c_2 e^{5t} - c_3 e^{3t} + c_1 e^{3t}$$

$$x_3(t) = c_2 e^{5t} + c_3 e^{3t}$$

✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 113

```
DSolve[{x1'[t]==9*x1[t]+4*x2[t]+0*x3[t],x2'[t]==-6*x1[t]-1*x2[t]+0*x3[t],x3'[t]==6*x1[t]+4*x3[t]}
```

$$x1(t) \rightarrow e^{3t}(c_1(3e^{2t} - 2) + 2c_2(e^{2t} - 1))$$

$$x2(t) \rightarrow -e^{3t}(3c_1(e^{2t} - 1) + c_2(2e^{2t} - 3))$$

$$x3(t) \rightarrow \int_1^t 3x(K[1])dK[1] + \frac{6}{5}c_1(e^{5t} - 1) + \frac{4}{5}c_2(e^{5t} - 1) + c_3$$

5.2 problem Example 3

Internal problem ID [355]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.6, Multiple Eigenvalue Solutions. Examples. Page 437

Problem number: Example 3.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= x_1(t) - 3x_2(t) \\x_2'(t) &= 3x_1(t) + 7x_2(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 35

```
dsolve([diff(x__1(t),t)=1*x__1(t)-3*x__2(t),diff(x__2(t),t)=3*x__1(t)+7*x__2(t)],[x__1(t), x
```

$$x_1(t) = -\frac{e^{4t}(3c_2t + 3c_1 - c_2)}{3}$$

$$x_2(t) = e^{4t}(c_2t + c_1)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 46

```
DSolve[{x1'[t]==1*x1[t]-3*x2[t],x2'[t]==3*x1[t]+7*x2[t]},{x1[t],x2[t]},t,IncludeSingularSolu
```

$$x1(t) \rightarrow -e^{4t}(c_1(3t - 1) + 3c_2t)$$

$$x2(t) \rightarrow e^{4t}(3(c_1 + c_2)t + c_2)$$

5.3 problem Example 4

Internal problem ID [356]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.6, Multiple Eigenvalue Solutions. Examples. Page 437

Problem number: Example 4.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = x_2(t) + 2x_3(t)$$

$$x_2'(t) = -5x_1(t) - 3x_2(t) - 7x_3(t)$$

$$x_3'(t) = x_1(t)$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 75

```
dsolve([diff(x__1(t),t)=0*x__1(t)+1*x__2(t)+2*x__3(t),diff(x__2(t),t)=-5*x__1(t)-3*x__2(t)-7*x__3(t),diff(x__3(t),t)=x__1(t)),x__1(t)=c1,x__2(t)=c2,x__3(t)=c3)
```

$$x_1(t) = -e^{-t}(c_3t^2 + c_2t - 2c_3t + c_1 - c_2)$$

$$x_2(t) = -e^{-t}(c_3t^2 + c_2t + 4c_3t + c_1 + 2c_2 - 2c_3)$$

$$x_3(t) = e^{-t}(c_3t^2 + c_2t + c_1)$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 134

```
DSolve[{x1'[t]==0*x1[t]+1*x2[t]+2*x3[t],x2'[t]==-5*x1[t]-3*x2[t]-7*x3[t],x3'[t]==1*x1[t]+0*x2[t]+0*x3[t]}
```

$$x1(t) \rightarrow \frac{1}{2}e^{-t}(c_1(-2t^2 + 2t + 2) - c_2(t - 2)t + c_3(4 - 3t)t)$$

$$x2(t) \rightarrow \frac{1}{2}e^{-t}(-((2c_1 + c_2 + 3c_3)t^2) - 2(5c_1 + 2c_2 + 7c_3)t + 2c_2)$$

$$x3(t) \rightarrow \frac{1}{2}e^{-t}((2c_1 + c_2 + 3c_3)t^2 + 2(c_1 + c_3)t + 2c_3)$$

5.4 problem Example 6

Internal problem ID [357]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.6, Multiple Eigenvalue Solutions. Examples. Page 437

Problem number: Example 6.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = x_3(t)$$

$$x_2'(t) = x_4(t)$$

$$x_3'(t) = -2x_1(t) + 2x_2(t) - 3x_3(t) + x_4(t)$$

$$x_4'(t) = 2x_1(t) - 2x_2(t) + x_3(t) - 3x_4(t)$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 87

```
dsolve([diff(x__1(t),t)=0*x__1(t)+0*x__2(t)+1*x__3(t)+0*x__4(t),diff(x__2(t),t)=0*x__1(t)+0*
```

$$x_1(t) = \frac{c_4 e^{-2t} t}{2} - \frac{e^{-2t} c_2}{2} + \frac{c_3 e^{-2t}}{2} + \frac{c_4 e^{-2t}}{4} + c_1$$

$$x_2(t) = \left(\left(-\frac{t}{2} - \frac{1}{4} \right) c_4 - \frac{c_3}{2} \right) e^{-2t} + c_1$$

$$x_3(t) = e^{-2t} (-c_4 t + c_2 - c_3)$$

$$x_4(t) = e^{-2t} (c_4 t + c_3)$$

✓ Solution by Mathematica

Time used: 0.06 (sec). Leaf size: 210

```
DSolve[{x1'[t]==0*x1[t]+0*x2[t]+1*x3[t]+0*x4[t],x2'[t]==0*x1[t]+0*x2[t]+0*x3[t]+1*x4[t],x3'[t]==0*x1[t]+0*x2[t]+0*x3[t]+0*x4[t],x4'[t]==0*x1[t]+0*x2[t]+0*x3[t]+0*x4[t]},x1,x2,x3,x4,t]
```

$$x1(t) \rightarrow \frac{1}{4}e^{-2t}(2c_1(2t + e^{2t} + 1) + 2c_2(-2t + e^{2t} - 1) + c_3e^{2t} + 2c_3t + c_4e^{2t} - 2c_4t - c_3 - c_4)$$

$$x2(t) \rightarrow \frac{1}{4}e^{-2t}(2c_1(-2t + e^{2t} - 1) + 2c_2(2t + e^{2t} + 1) + c_3e^{2t} - 2c_3t + c_4e^{2t} + 2c_4t - c_3 - c_4)$$

$$x3(t) \rightarrow e^{-2t}((-2c_1 + 2c_2 - c_3 + c_4)t + c_3)$$

$$x4(t) \rightarrow e^{-2t}((2c_1 - 2c_2 + c_3 - c_4)t + c_4)$$

6 Section 7.6, Multiple Eigenvalue Solutions. Page 451

6.1	problem problem 1	113
6.2	problem problem 2	114
6.3	problem problem 3	115
6.4	problem problem 4	116
6.5	problem problem 5	117
6.6	problem problem 6	118
6.7	problem problem 7	119
6.8	problem problem 8	120
6.9	problem problem 9	121
6.10	problem problem 10	123
6.11	problem problem 11	125
6.12	problem problem 12	127
6.13	problem problem 13	129
6.14	problem problem 14	131
6.15	problem problem 15	133
6.16	problem problem 16	134
6.17	problem problem 17	136
6.18	problem problem 18	138
6.19	problem problem 19	140
6.20	problem problem 20	142
6.21	problem problem 21	144
6.22	problem problem 22	146
6.23	problem problem 23	148
6.24	problem problem 24	150
6.25	problem problem 25	152
6.26	problem problem 26	154
6.27	problem problem 27	156
6.28	problem problem 28	158
6.29	problem problem 29	160
6.30	problem problem 30	162
6.31	problem problem 31	164
6.32	problem problem 32	166
6.33	problem problem 33	168
6.34	problem problem 34	170

6.1 problem problem 1

Internal problem ID [358]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.6, Multiple Eigenvalue Solutions. Page 451

Problem number: problem 1.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = -2x_1(t) + x_2(t)$$

$$x_2'(t) = -x_1(t) - 4x_2(t)$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 30

```
dsolve([diff(x__1(t),t)=-2*x__1(t)+1*x__2(t),diff(x__2(t),t)=-1*x__1(t)-4*x__2(t)],[x__1(t),
```

$$x_1(t) = -e^{-3t}(c_2t + c_1 + c_2)$$

$$x_2(t) = e^{-3t}(c_2t + c_1)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 42

```
DSolve[{x1'[t]==-2*x1[t]+1*x2[t],x2'[t]==-1*x1[t]-4*x2[t]},{x1[t],x2[t]},t,IncludeSingularSo
```

$$x1(t) \rightarrow e^{-3t}(c_1(t+1) + c_2t)$$

$$x2(t) \rightarrow e^{-3t}(c_2 - (c_1 + c_2)t)$$

6.2 problem problem 2

Internal problem ID [359]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.6, Multiple Eigenvalue Solutions. Page 451

Problem number: problem 2.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 3x_1(t) - x_2(t)$$

$$x_2'(t) = x_1(t) + x_2(t)$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 29

```
dsolve([diff(x__1(t),t)=3*x__1(t)-1*x__2(t),diff(x__2(t),t)=1*x__1(t)+1*x__2(t)],[x__1(t), x
```

$$x_1(t) = e^{2t}(c_2t + c_1 + c_2)$$

$$x_2(t) = e^{2t}(c_2t + c_1)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 44

```
DSolve[{x1'[t]==3*x1[t]-1*x2[t],x2'[t]==1*x1[t]+1*x2[t]},{x1[t],x2[t]},t,IncludeSingularSolu
```

$$x1(t) \rightarrow e^{2t}(c_1(t + 1) - c_2t)$$

$$x2(t) \rightarrow e^{2t}((c_1 - c_2)t + c_2)$$

6.3 problem problem 3

Internal problem ID [360]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.6, Multiple Eigenvalue Solutions. Page 451

Problem number: problem 3.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = x_1(t) - 2x_2(t)$$

$$x_2'(t) = 2x_1(t) + 5x_2(t)$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 35

```
dsolve([diff(x__1(t),t)=1*x__1(t)-2*x__2(t),diff(x__2(t),t)=2*x__1(t)+5*x__2(t)],[x__1(t), x
```

$$x_1(t) = -\frac{e^{3t}(2c_2t + 2c_1 - c_2)}{2}$$

$$x_2(t) = e^{3t}(c_2t + c_1)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 46

```
DSolve[{x1'[t]==1*x1[t]-2*x2[t],x2'[t]==2*x1[t]+5*x2[t]},{x1[t],x2[t]},t,IncludeSingularSolu
```

$$x1(t) \rightarrow -e^{3t}(c_1(2t - 1) + 2c_2t)$$

$$x2(t) \rightarrow e^{3t}(2(c_1 + c_2)t + c_2)$$

6.4 problem problem 4

Internal problem ID [361]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.6, Multiple Eigenvalue Solutions. Page 451

Problem number: problem 4.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 3x_1(t) - x_2(t)$$

$$x_2'(t) = x_1(t) + 5x_2(t)$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 32

```
dsolve([diff(x__1(t),t)=3*x__1(t)-1*x__2(t),diff(x__2(t),t)=1*x__1(t)+5*x__2(t)],[x__1(t), x
```

$$x_1(t) = -e^{4t}(c_2t + c_1 - c_2)$$

$$x_2(t) = e^{4t}(c_2t + c_1)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 42

```
DSolve[{x1'[t]==3*x1[t]-1*x2[t],x2'[t]==1*x1[t]+5*x2[t]},{x1[t],x2[t]},t,IncludeSingularSolu
```

$$x1(t) \rightarrow -e^{4t}(c_1(t - 1) + c_2t)$$

$$x2(t) \rightarrow e^{4t}((c_1 + c_2)t + c_2)$$

6.5 problem problem 5

Internal problem ID [362]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.6, Multiple Eigenvalue Solutions. Page 451

Problem number: problem 5.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= 7x_1(t) + x_2(t) \\x_2'(t) &= -4x_1(t) + 3x_2(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 33

```
dsolve([diff(x__1(t),t)=7*x__1(t)+1*x__2(t),diff(x__2(t),t)=-4*x__1(t)+3*x__2(t)],[x__1(t),
```

$$x_1(t) = -\frac{e^{5t}(2c_2t + 2c_1 + c_2)}{4}$$

$$x_2(t) = e^{5t}(c_2t + c_1)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 45

```
DSolve[{x1'[t]==7*x1[t]+1*x2[t],x2'[t]==-4*x1[t]+3*x2[t]},{x1[t],x2[t]},t,IncludeSingularSol
```

$$x1(t) \rightarrow e^{5t}(2c_1t + c_2t + c_1)$$

$$x2(t) \rightarrow e^{5t}(c_2 - 2(2c_1 + c_2)t)$$

6.6 problem problem 6

Internal problem ID [363]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.6, Multiple Eigenvalue Solutions. Page 451

Problem number: problem 6.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = x_1(t) - 4x_2(t)$$

$$x_2'(t) = 4x_1(t) + 9x_2(t)$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 35

```
dsolve([diff(x__1(t),t)=1*x__1(t)-4*x__2(t),diff(x__2(t),t)=4*x__1(t)+9*x__2(t)],[x__1(t), x
```

$$x_1(t) = -\frac{e^{5t}(4c_2t + 4c_1 - c_2)}{4}$$

$$x_2(t) = e^{5t}(c_2t + c_1)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 46

```
DSolve[{x1'[t]==1*x1[t]-4*x2[t],x2'[t]==4*x1[t]+9*x2[t]},{x1[t],x2[t]},t,IncludeSingularSolu
```

$$x1(t) \rightarrow -e^{5t}(c_1(4t - 1) + 4c_2t)$$

$$x2(t) \rightarrow e^{5t}(4(c_1 + c_2)t + c_2)$$

6.7 problem problem 7

Internal problem ID [364]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.6, Multiple Eigenvalue Solutions. Page 451

Problem number: problem 7.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= 2x_1(t) \\x_2'(t) &= -7x_1(t) + 9x_2(t) + 7x_3(t) \\x_3'(t) &= 2x_3(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 38

```
dsolve([diff(x__1(t),t)=2*x__1(t)+0*x__2(t)+0*x__3(t),diff(x__2(t),t)=-7*x__1(t)+9*x__2(t)+7*x__3(t),diff(x__3(t),t)=2*x__3(t)),x__1(t),x__2(t),x__3(t))
```

$$x_1(t) = e^{2t}(c_2 + c_3)$$

$$x_2(t) = c_1 e^{9t} + c_2 e^{2t}$$

$$x_3(t) = c_3 e^{2t}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 60

```
DSolve[{x1'[t]==2*x1[t]+0*x2[t]+0*x3[t],x2'[t]==-7*x1[t]+9*x2[t]+7*x3[t],x3'[t]==0*x1[t]+0*x2[t]+2*x3[t]},x1[t],x2[t],x3[t]]
```

$$x1(t) \rightarrow c_1 e^{2t}$$

$$x2(t) \rightarrow e^{2t}(-c_1(e^{7t} - 1)) + (c_2 + c_3)e^{7t} - c_3$$

$$x3(t) \rightarrow c_3 e^{2t}$$

6.8 problem problem 8

Internal problem ID [365]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.6, Multiple Eigenvalue Solutions. Page 451

Problem number: problem 8.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 25x_1(t) + 12x_2(t)$$

$$x_2'(t) = -18x_1(t) - 5x_2(t)$$

$$x_3'(t) = 6x_1(t) + 6x_2(t) + 13x_3(t)$$

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 67

```
dsolve([diff(x__1(t),t)=25*x__1(t)+12*x__2(t)+0*x__3(t),diff(x__2(t),t)=-18*x__1(t)-5*x__2(t)
```

$$x_1(t) = 2c_3e^{7t} + 3c_2e^{13t} - e^{13t}c_1$$

$$x_2(t) = -3c_2e^{13t} - 3c_3e^{7t} + e^{13t}c_1$$

$$x_3(t) = c_2e^{13t} + c_3e^{7t}$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 107

```
DSolve[{x1'[t]==25*x1[t]+12*x2[t]+0*x3[t],x2'[t]==-18*x1[t]-5*x2[t]+0*x3[t],x3'[t]==6*x1[t]+
```

$$x1(t) \rightarrow e^{7t}(c_1(3e^{6t} - 2) + 2c_2(e^{6t} - 1))$$

$$x2(t) \rightarrow -e^{7t}(3c_1(e^{6t} - 1) + c_2(2e^{6t} - 3))$$

$$x3(t) \rightarrow e^{7t}(c_1(e^{6t} - 1) + c_2(e^{6t} - 1) + c_3e^{6t})$$

6.9 problem problem 9

Internal problem ID [366]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.6, Multiple Eigenvalue Solutions. Page 451

Problem number: problem 9.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = -19x_1(t) + 12x_2(t) + 84x_3(t)$$

$$x_2'(t) = 5x_2(t)$$

$$x_3'(t) = -8x_1(t) + 4x_2(t) + 33x_3(t)$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 52

```
dsolve([diff(x__1(t),t)=-19*x__1(t)+12*x__2(t)+84*x__3(t),diff(x__2(t),t)=0*x__1(t)+5*x__2(t),diff(x__3(t),t)=-8*x__1(t)+4*x__2(t)+33*x__3(t)),x__1(t),x__2(t),x__3(t))
```

$$x_1(t) = 3c_2e^{9t} + \frac{7c_3e^{5t}}{2} + \frac{c_1e^{5t}}{2}$$

$$x_2(t) = c_1e^{5t}$$

$$x_3(t) = c_2e^{9t} + c_3e^{5t}$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 94

```
DSolve[{x1'[t]==-19*x1[t]+12*x2[t]+84*x3[t],x2'[t]==0*x1[t]+5*x2[t]+0*x3[t],x3'[t]==-8*x1[t]
```

$$x1(t) \rightarrow e^{5t}(c_1(7 - 6e^{4t}) + 3(c_2 + 7c_3)(e^{4t} - 1))$$

$$x2(t) \rightarrow c_2 e^{5t}$$

$$x3(t) \rightarrow e^{5t}(-2c_1(e^{4t} - 1) + c_2(e^{4t} - 1) + c_3(7e^{4t} - 6))$$

6.10 problem problem 10

Internal problem ID [367]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.6, Multiple Eigenvalue Solutions. Page 451

Problem number: problem 10.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = -13x_1(t) + 40x_2(t) - 48x_3(t)$$

$$x_2'(t) = -8x_1(t) + 23x_2(t) - 24x_3(t)$$

$$x_3'(t) = 3x_3(t)$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 52

```
dsolve([diff(x__1(t),t)=-13*x__1(t)+40*x__2(t)-48*x__3(t),diff(x__2(t),t)=-8*x__1(t)+23*x__2
```

$$x_1(t) = \frac{5c_1e^{3t}}{2} + 2c_2e^{7t} - 3c_3e^{3t}$$

$$x_2(t) = c_1e^{3t} + c_2e^{7t}$$

$$x_3(t) = c_3e^{3t}$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 98

```
DSolve[{x1'[t]==-13*x1[t]+40*x2[t]-48*x3[t],x2'[t]==-8*x1[t]+23*x2[t]-24*x3[t],x3'[t]==0*x1[t]
```

$$x1(t) \rightarrow e^{3t}(c_1(5 - 4e^{4t}) + 2(5c_2 - 6c_3)(e^{4t} - 1))$$

$$x2(t) \rightarrow -e^{3t}(2c_1(e^{4t} - 1) + c_2(4 - 5e^{4t}) + 6c_3(e^{4t} - 1))$$

$$x3(t) \rightarrow c_3e^{3t}$$

6.11 problem problem 11

Internal problem ID [368]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.6, Multiple Eigenvalue Solutions. Page 451

Problem number: problem 11.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = -3x_1(t) - 4x_3(t)$$

$$x_2'(t) = -x_1(t) - x_2(t) - x_3(t)$$

$$x_3'(t) = x_1(t) + x_3(t)$$

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 61

```
dsolve([diff(x__1(t),t)=-3*x__1(t)+0*x__2(t)-4*x__3(t),diff(x__2(t),t)=-1*x__1(t)-1*x__2(t)-
```

$$x_1(t) = -e^{-t}(2c_3t + 2c_2 - c_3)$$

$$x_2(t) = \frac{(c_3t^2 + 2c_2t - 2c_3t + 2c_1)e^{-t}}{2}$$

$$x_3(t) = e^{-t}(c_3t + c_2)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 81

```
DSolve[{x1'[t]==-3*x1[t]+0*x2[t]-4*x3[t],x2'[t]==-1*x1[t]-1*x2[t]-1*x3[t],x3'[t]==1*x1[t]+0*
```

$$x1(t) \rightarrow e^{-t}(-2c_1t - 4c_3t + c_1)$$

$$x2(t) \rightarrow \frac{1}{2}e^{-t}((c_1 + 2c_3)t^2 - 2(c_1 + c_3)t + 2c_2)$$

$$x3(t) \rightarrow e^{-t}((c_1 + 2c_3)t + c_3)$$

6.12 problem problem 12

Internal problem ID [369]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.6, Multiple Eigenvalue Solutions. Page 451

Problem number: problem 12.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = -x_1(t) + x_3(t)$$

$$x_2'(t) = -x_2(t) + x_3(t)$$

$$x_3'(t) = x_1(t) - x_2(t) - x_3(t)$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 62

```
dsolve([diff(x__1(t),t)=-1*x__1(t)+0*x__2(t)+1*x__3(t),diff(x__2(t),t)=0*x__1(t)-1*x__2(t)+1
```

$$x_1(t) = \frac{e^{-t}(c_3 t^2 + 2c_2 t + 2c_1 + 2c_3)}{2}$$

$$x_2(t) = \frac{(c_3 t^2 + 2c_2 t + 2c_1) e^{-t}}{2}$$

$$x_3(t) = e^{-t}(c_3 t + c_2)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 89

```
DSolve[{x1'[t]==-1*x1[t]+0*x2[t]+1*x3[t],x2'[t]==0*x1[t]-1*x2[t]+1*x3[t],x3'[t]==1*x1[t]-1*x
```

$$x1(t) \rightarrow \frac{1}{2}e^{-t}(c_1(t^2 + 2) + t(2c_3 - c_2t))$$

$$x2(t) \rightarrow \frac{1}{2}e^{-t}((c_1 - c_2)t^2 + 2c_3t + 2c_2)$$

$$x3(t) \rightarrow e^{-t}((c_1 - c_2)t + c_3)$$

6.13 problem problem 13

Internal problem ID [370]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.6, Multiple Eigenvalue Solutions. Page 451

Problem number: problem 13.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = -x_1(t) + x_3(t)$$

$$x_2'(t) = x_2(t) - 4x_3(t)$$

$$x_3'(t) = x_2(t) - 3x_3(t)$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 54

```
dsolve([diff(x__1(t),t)=-1*x__1(t)+0*x__2(t)+1*x__3(t),diff(x__2(t),t)=0*x__1(t)+1*x__2(t)-4
```

$$x_1(t) = \frac{(c_3 t^2 + 2c_2 t + 2c_1) e^{-t}}{2}$$

$$x_2(t) = e^{-t}(2c_3 t + 2c_2 + c_3)$$

$$x_3(t) = e^{-t}(c_3 t + c_2)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 78

```
DSolve[{x1'[t]==-1*x1[t]+0*x2[t]+1*x3[t],x2'[t]==0*x1[t]+1*x2[t]-4*x3[t],x3'[t]==0*x1[t]+1*x
```

$$x1(t) \rightarrow \frac{1}{2}e^{-t}(t((c_2 - 2c_3)t + 2c_3) + 2c_1)$$

$$x2(t) \rightarrow e^{-t}(2c_2t - 4c_3t + c_2)$$

$$x3(t) \rightarrow e^{-t}((c_2 - 2c_3)t + c_3)$$

6.14 problem problem 14

Internal problem ID [371]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.6, Multiple Eigenvalue Solutions. Page 451

Problem number: problem 14.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = x_3(t)$$

$$x_2'(t) = -5x_1(t) - x_2(t) - 5x_3(t)$$

$$x_3'(t) = 4x_1(t) + x_2(t) - 2x_3(t)$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 79

```
dsolve([diff(x__1(t),t)=0*x__1(t)+0*x__2(t)+1*x__3(t),diff(x__2(t),t)=-5*x__1(t)-1*x__2(t)-5
```

$$x_1(t) = -e^{-t}(c_3t^2 + c_2t + 2c_3t + c_1 + c_2 + 2c_3)$$

$$x_2(t) = e^{-t}(5c_3t^2 + 5c_2t + 10c_3t + 5c_1 + 5c_2 + 8c_3)$$

$$x_3(t) = e^{-t}(c_3t^2 + c_2t + c_1)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 119

```
DSolve[{x1'[t]==0*x1[t]+0*x2[t]+1*x3[t],x2'[t]==-5*x1[t]-1*x2[t]-5*x3[t],x3'[t]==4*x1[t]+1*x
```

$$x1(t) \rightarrow \frac{1}{2}e^{-t}(c_1(5t^2 + 2t + 2) + t(c_2t + 2c_3))$$

$$x2(t) \rightarrow \frac{1}{2}e^{-t}(-5(5c_1 + c_2)t^2 - 10(c_1 + c_3)t + 2c_2)$$

$$x3(t) \rightarrow \frac{1}{2}e^{-t}(-((5c_1 + c_2)t^2) + 2(4c_1 + c_2 - c_3)t + 2c_3)$$

6.15 problem problem 15

Internal problem ID [372]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.6, Multiple Eigenvalue Solutions. Page 451

Problem number: problem 15.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = -2x_1(t) - 9x_2(t)$$

$$x_2'(t) = x_1(t) + 4x_2(t)$$

$$x_3'(t) = x_1(t) + 3x_2(t) + x_3(t)$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 46

```
dsolve([diff(x__1(t),t)=-2*x__1(t)-9*x__2(t)-0*x__3(t),diff(x__2(t),t)=1*x__1(t)+4*x__2(t)-0
```

$$x_1(t) = -e^t(3c_3t + 3c_1 + 3c_2 - c_3)$$

$$x_2(t) = e^t(c_3t + c_1 + c_2)$$

$$x_3(t) = e^t(c_3t + c_2)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 62

```
DSolve[{x1'[t]==-2*x1[t]-9*x2[t]-0*x3[t],x2'[t]==1*x1[t]+4*x2[t]-0*x3[t],x3'[t]==1*x1[t]+3*x
```

$$x_1(t) \rightarrow -e^t(c_1(3t - 1) + 9c_2t)$$

$$x_2(t) \rightarrow e^t((c_1 + 3c_2)t + c_2)$$

$$x_3(t) \rightarrow e^t((c_1 + 3c_2)t + c_3)$$

6.16 problem problem 16

Internal problem ID [373]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.6, Multiple Eigenvalue Solutions. Page 451

Problem number: problem 16.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = x_1(t)$$

$$x_2'(t) = -2x_1(t) - 2x_2(t) - 3x_3(t)$$

$$x_3'(t) = 2x_1(t) + 3x_2(t) + 4x_3(t)$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 42

```
dsolve([diff(x__1(t),t)=1*x__1(t)+0*x__2(t)-0*x__3(t),diff(x__2(t),t)=-2*x__1(t)-2*x__2(t)-3
```

$$x_1(t) = -\frac{e^t(3c_1 - c_3)}{2}$$

$$x_2(t) = e^t(-c_3t + c_1 - c_2)$$

$$x_3(t) = e^t(c_3t + c_2)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 57

```
DSolve[{x1'[t]==1*x1[t]+0*x2[t]-0*x3[t],x2'[t]==-2*x1[t]-2*x2[t]-3*x3[t],x3'[t]==2*x1[t]+3*x
```

$$x1(t) \rightarrow c_1 e^t$$

$$x2(t) \rightarrow e^t(-2c_1 t - 3(c_2 + c_3)t + c_2)$$

$$x3(t) \rightarrow e^t(2c_1 t + 3(c_2 + c_3)t + c_3)$$

6.17 problem problem 17

Internal problem ID [374]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.6, Multiple Eigenvalue Solutions. Page 451

Problem number: problem 17.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = x_1(t)$$

$$x_2'(t) = 18x_1(t) + 7x_2(t) + 4x_3(t)$$

$$x_3'(t) = -27x_1(t) - 9x_2(t) - 5x_3(t)$$

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 43

```
dsolve([diff(x__1(t),t)=1*x__1(t)+0*x__2(t)-0*x__3(t),diff(x__2(t),t)=18*x__1(t)+7*x__2(t)+4
```

$$x_1(t) = -\frac{e^t(9c_1 + c_3)}{27}$$

$$x_2(t) = \frac{e^t(-2c_3t + 3c_1 - 2c_2)}{3}$$

$$x_3(t) = e^t(c_3t + c_2)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 63

```
DSolve[{x1'[t]==1*x1[t]+0*x2[t]-0*x3[t],x2'[t]==18*x1[t]+7*x2[t]+4*x3[t],x3'[t]==-27*x1[t]-9
```

$$x1(t) \rightarrow c_1 e^t$$

$$x2(t) \rightarrow e^t(2(9c_1 + 3c_2 + 2c_3)t + c_2)$$

$$x3(t) \rightarrow e^t(c_3 - 3(9c_1 + 3c_2 + 2c_3)t)$$

6.18 problem problem 18

Internal problem ID [375]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.6, Multiple Eigenvalue Solutions. Page 451

Problem number: problem 18.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = x_1(t)$$

$$x_2'(t) = x_1(t) + 3x_2(t) + x_3(t)$$

$$x_3'(t) = -2x_1(t) - 4x_2(t) - x_3(t)$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 43

```
dsolve([diff(x__1(t),t)=1*x__1(t)+0*x__2(t)-0*x__3(t),diff(x__2(t),t)=1*x__1(t)+3*x__2(t)+1*
```

$$x_1(t) = -\frac{e^t(4c_1 + c_3)}{2}$$

$$x_2(t) = \frac{e^t(-c_3t + 2c_1 - c_2)}{2}$$

$$x_3(t) = e^t(c_3t + c_2)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 54

```
DSolve[{x1'[t]==1*x1[t]+0*x2[t]-0*x3[t],x2'[t]==1*x1[t]+3*x2[t]+1*x3[t],x3'[t]==-2*x1[t]-4*x2[t]-4*x3[t]}
```

$$x1(t) \rightarrow c_1 e^t$$

$$x2(t) \rightarrow e^t((c_1 + 2c_2 + c_3)t + c_2)$$

$$x3(t) \rightarrow e^t(c_3 - 2(c_1 + 2c_2 + c_3)t)$$

6.19 problem problem 19

Internal problem ID [376]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.6, Multiple Eigenvalue Solutions. Page 451

Problem number: problem 19.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = x_1(t) - 4x_2(t) - 2x_4(t)$$

$$x_2'(t) = x_2(t)$$

$$x_3'(t) = 6x_1(t) - 12x_2(t) - x_3(t) - 6x_4(t)$$

$$x_4'(t) = -4x_2(t) - x_4(t)$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 53

```
dsolve([diff(x__1(t),t)=1*x__1(t)-4*x__2(t)+0*x__3(t)-2*x__4(t),diff(x__2(t),t)=0*x__1(t)+1*
```

$$x_1(t) = \frac{c_1 e^t}{3} + c_4 e^{-t}$$

$$x_2(t) = -\frac{c_3 e^t}{2}$$

$$x_3(t) = c_1 e^t + c_2 e^{-t}$$

$$x_4(t) = c_3 e^t + c_4 e^{-t}$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 114

```
DSolve[{x1'[t]==1*x1[t]-4*x2[t]+0*x3[t]-2*x4[t],x2'[t]==0*x1[t]+1*x2[t]+0*x3[t]+0*x4[t],x3'[t]==0*x1[t]+0*x2[t]+1*x3[t]-2*x4[t],x4'[t]==0*x1[t]+0*x2[t]+0*x3[t]+1*x4[t]}
```

$$x1(t) \rightarrow e^{-t}((c_1 - 2c_2 - c_4)e^{2t} + 2c_2 + c_4)$$

$$x2(t) \rightarrow c_2 e^t$$

$$x3(t) \rightarrow e^{-t}(3c_1(e^{2t} - 1) - 6c_2(e^{2t} - 1) - 3c_4 e^{2t} + c_3 + 3c_4)$$

$$x4(t) \rightarrow e^{-t}(c_4 - 2c_2(e^{2t} - 1))$$

6.20 problem problem 20

Internal problem ID [377]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.6, Multiple Eigenvalue Solutions. Page 451

Problem number: problem 20.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 2x_1(t) + x_2(t) + x_4(t)$$

$$x_2'(t) = 2x_2(t) + x_3(t)$$

$$x_3'(t) = 2x_3(t) + x_4(t)$$

$$x_4'(t) = 2x_4(t)$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 78

```
dsolve([diff(x__1(t),t)=2*x__1(t)+1*x__2(t)+0*x__3(t)+1*x__4(t),diff(x__2(t),t)=0*x__1(t)+2*
```

$$x_1(t) = \frac{(c_4 t^3 + 3c_3 t^2 + 6c_2 t + 6c_4 t + 6c_1) e^{2t}}{6}$$

$$x_2(t) = \frac{(c_4 t^2 + 2c_3 t + 2c_2) e^{2t}}{2}$$

$$x_3(t) = (c_4 t + c_3) e^{2t}$$

$$x_4(t) = c_4 e^{2t}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 96

```
DSolve[{x1'[t]==2*x1[t]+1*x2[t]+0*x3[t]+1*x4[t],x2'[t]==0*x1[t]+2*x2[t]+1*x3[t]+0*x4[t],x3'[t]==0*x1[t]+0*x2[t]+0*x3[t]+0*x4[t],x4'[t]==0*x1[t]+0*x2[t]+0*x3[t]+0*x4[t]}
```

$$x1(t) \rightarrow \frac{1}{6}e^{2t}(t(c_4t^2 + 3c_3t + 6c_2 + 6c_4) + 6c_1)$$

$$x2(t) \rightarrow \frac{1}{2}e^{2t}(t(c_4t + 2c_3) + 2c_2)$$

$$x3(t) \rightarrow e^{2t}(c_4t + c_3)$$

$$x4(t) \rightarrow c_4e^{2t}$$

6.21 problem problem 21

Internal problem ID [378]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.6, Multiple Eigenvalue Solutions. Page 451

Problem number: problem 21.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = -x_1(t) - 4x_2(t)$$

$$x_2'(t) = x_1(t) + 3x_2(t)$$

$$x_3'(t) = x_1(t) + 2x_2(t) + x_3(t)$$

$$x_4'(t) = x_2(t) + x_4(t)$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 59

```
dsolve([diff(x__1(t),t)=-1*x__1(t)-4*x__2(t)+0*x__3(t)+0*x__4(t),diff(x__2(t),t)=1*x__1(t)+3
```

$$x_1(t) = -2e^t(2c_4t + c_3 - c_4)$$

$$x_2(t) = e^t(2c_4t + c_3)$$

$$x_3(t) = e^t(2c_4t + c_1 + c_3)$$

$$x_4(t) = e^t(c_4t^2 + c_3t + c_2)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 91

```
DSolve[{x1'[t]==-1*x1[t]-4*x2[t]+0*x3[t]+0*x4[t],x2'[t]==1*x1[t]+3*x2[t]+0*x3[t]+0*x4[t],x3'
```

$$x1(t) \rightarrow -e^t(c_1(2t - 1) + 4c_2t)$$

$$x2(t) \rightarrow e^t((c_1 + 2c_2)t + c_2)$$

$$x3(t) \rightarrow e^t((c_1 + 2c_2)t + c_3)$$

$$x4(t) \rightarrow \frac{1}{2}e^t(c_1t^2 + 2c_2(t + 1)t + 2c_4)$$

6.22 problem problem 22

Internal problem ID [379]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.6, Multiple Eigenvalue Solutions. Page 451

Problem number: problem 22.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = x_1(t) + 3x_2(t) + 7x_3(t)$$

$$x_2'(t) = -x_2(t) - 4x_3(t)$$

$$x_3'(t) = x_2(t) + 3x_3(t)$$

$$x_4'(t) = -6x_2(t) - 14x_3(t) + x_4(t)$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 73

```
dsolve([diff(x__1(t),t)=1*x__1(t)+3*x__2(t)+7*x__3(t)+0*x__4(t),diff(x__2(t),t)=0*x__1(t)-1*
```

$$x_1(t) = \frac{e^t(-c_4t^2 - c_3t + 2c_1 - c_2)}{2}$$

$$x_2(t) = e^t(2c_4t + c_3 - 7c_4)$$

$$x_3(t) = -\frac{e^t(2c_4t + c_3 - 6c_4)}{2}$$

$$x_4(t) = e^t(c_4t^2 + c_3t + c_2)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 99

```
DSolve[{x1'[t]==1*x1[t]+3*x2[t]+7*x3[t]+0*x4[t],x2'[t]==0*x1[t]-1*x2[t]-4*x3[t]+0*x4[t],x3'[t]==0*x1[t]+0*x2[t]+0*x3[t]+0*x4[t],x4'[t]==0*x1[t]+0*x2[t]+0*x3[t]+0*x4[t]}
```

$$x1(t) \rightarrow \frac{1}{2}e^t(c_2t(t+6) + 2c_3t(t+7) + 2c_1)$$

$$x2(t) \rightarrow -e^t(c_2(2t-1) + 4c_3t)$$

$$x3(t) \rightarrow e^t((c_2 + 2c_3)t + c_3)$$

$$x4(t) \rightarrow e^t(c_2(-t)(t+6) - 2c_3t(t+7) + c_4)$$

6.23 problem problem 23

Internal problem ID [380]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.6, Multiple Eigenvalue Solutions. Page 451

Problem number: problem 23.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 39x_1(t) + 8x_2(t) - 16x_3(t)$$

$$x_2'(t) = -36x_1(t) - 5x_2(t) + 16x_3(t)$$

$$x_3'(t) = 72x_1(t) + 16x_2(t) - 29x_3(t)$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 67

```
dsolve([diff(x__1(t),t)=39*x__1(t)+8*x__2(t)-16*x__3(t),diff(x__2(t),t)=-36*x__1(t)-5*x__2(t)
```

$$x_1(t) = \frac{c_2 e^{-t}}{2} + \frac{5c_3 e^{3t}}{9} - \frac{2c_1 e^{3t}}{9}$$

$$x_2(t) = -\frac{c_2 e^{-t}}{2} - \frac{c_3 e^{3t}}{2} + c_1 e^{3t}$$

$$x_3(t) = c_2 e^{-t} + c_3 e^{3t}$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 127

```
DSolve[{x1'[t]==39*x1[t]+8*x2[t]-16*x3[t],x2'[t]==-36*x1[t]-5*x2[t]+16*x3[t],x3'[t]==72*x1[t]
```

$$x1(t) \rightarrow e^{-t}(c_1(10e^{4t} - 9) + 2(c_2 - 2c_3)(e^{4t} - 1))$$

$$x2(t) \rightarrow e^{-t}(-9c_1(e^{4t} - 1) - c_2(e^{4t} - 2) + 4c_3(e^{4t} - 1))$$

$$x3(t) \rightarrow e^{-t}(18c_1(e^{4t} - 1) + 4c_2(e^{4t} - 1) + c_3(8 - 7e^{4t}))$$

6.24 problem problem 24

Internal problem ID [381]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.6, Multiple Eigenvalue Solutions. Page 451

Problem number: problem 24.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 28x_1(t) + 50x_2(t) + 100x_3(t)$$

$$x_2'(t) = 15x_1(t) + 33x_2(t) + 60x_3(t)$$

$$x_3'(t) = -15x_1(t) - 30x_2(t) - 57x_3(t)$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 67

```
dsolve([diff(x__1(t),t)=28*x__1(t)+50*x__2(t)+100*x__3(t),diff(x__2(t),t)=15*x__1(t)+33*x__2
```

$$x_1(t) = -\frac{5e^{-2t}c_2}{3} - 2c_3e^{3t} - 2c_1e^{3t}$$

$$x_2(t) = -e^{-2t}c_2 - c_3e^{3t} + c_1e^{3t}$$

$$x_3(t) = e^{-2t}c_2 + c_3e^{3t}$$

✓ Solution by Mathematica

Time used: 0.046 (sec). Leaf size: 229

```
DSolve[{x1'[t]==28*x1[t]+50*x2[t]+100*x3[t],x2'[t]==15*x1[t]+33*x2[t]+60*x3[t],x3'[t]==-15*x
```

$$x1(t) \rightarrow \frac{1}{57}e^{t/2} \left(19(3c_1 - 5c_2)e^{5t/2} + 95c_2 \cos\left(\frac{5\sqrt{95}t}{2}\right) + \sqrt{95}(6c_1 + 13c_2 + 24c_3) \sin\left(\frac{5\sqrt{95}t}{2}\right) \right)$$

$$x2(t) \rightarrow \frac{1}{95}e^{t/2} \left(95c_2 \cos\left(\frac{5\sqrt{95}t}{2}\right) + \sqrt{95}(6c_1 + 13c_2 + 24c_3) \sin\left(\frac{5\sqrt{95}t}{2}\right) \right)$$

$$x3(t) \rightarrow \frac{e^{t/2} \left(95(3c_1 - 5c_2)e^{5t/2} - 95(3c_1 - 5c_2 + 12c_3) \cos\left(\frac{5\sqrt{95}t}{2}\right) + \sqrt{95}(69c_1 + 197c_2 + 276c_3) \sin\left(\frac{5\sqrt{95}t}{2}\right) \right)}{1140}$$

6.25 problem problem 25

Internal problem ID [382]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.6, Multiple Eigenvalue Solutions. Page 451

Problem number: problem 25.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = -2x_1(t) + 17x_2(t) + 4x_3(t)$$

$$x_2'(t) = -x_1(t) + 6x_2(t) + x_3(t)$$

$$x_3'(t) = x_2(t) + 2x_3(t)$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 62

```
dsolve([diff(x__1(t),t)=-2*x__1(t)+17*x__2(t)+4*x__3(t),diff(x__2(t),t)=-1*x__1(t)+6*x__2(t)
```

$$x_1(t) = e^{2t}(c_3t^2 + c_2t + 8c_3t + c_1 + 4c_2 - 2c_3)$$

$$x_2(t) = e^{2t}(2c_3t + c_2)$$

$$x_3(t) = e^{2t}(c_3t^2 + c_2t + c_1)$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 105

```
DSolve[{x1'[t]==-2*x1[t]+17*x2[t]+4*x3[t],x2'[t]==-1*x1[t]+6*x2[t]+1*x3[t],x3'[t]==0*x1[t]+1
```

$$x1(t) \rightarrow \frac{1}{2}e^{2t}(-c_1(t^2 + 8t - 2)) + c_2t(4t + 34) + c_3t(t + 8)$$

$$x2(t) \rightarrow e^{2t}((-c_1 + 4c_2 + c_3)t + c_2)$$

$$x3(t) \rightarrow \frac{1}{2}e^{2t}((-c_1 + 4c_2 + c_3)t^2 + 2c_2t + 2c_3)$$

6.26 problem problem 26

Internal problem ID [383]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.6, Multiple Eigenvalue Solutions. Page 451

Problem number: problem 26.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 5x_1(t) - x_2(t) + x_3(t)$$

$$x_2'(t) = x_1(t) + 3x_2(t)$$

$$x_3'(t) = -3x_1(t) + 2x_2(t) + x_3(t)$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 65

```
dsolve([diff(x__1(t),t)=5*x__1(t)-1*x__2(t)+1*x__3(t),diff(x__2(t),t)=1*x__1(t)+3*x__2(t)+0*
```

$$x_1(t) = e^{3t}(2c_3t + c_2 + 4c_3)$$

$$x_2(t) = e^{3t}(c_3t^2 + c_2t + 4c_3t + c_1 + 2c_2 + 6c_3)$$

$$x_3(t) = e^{3t}(c_3t^2 + c_2t + c_1)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 105

```
DSolve[{x1'[t]==5*x1[t]-1*x2[t]+1*x3[t],x2'[t]==1*x1[t]+3*x2[t]+0*x3[t],x3'[t]==-3*x1[t]+2*x
```

$$x1(t) \rightarrow e^{3t}(2c_1t - c_2t + c_3t + c_1)$$

$$x2(t) \rightarrow \frac{1}{2}e^{3t}((2c_1 - c_2 + c_3)t^2 + 2c_1t + 2c_2)$$

$$x3(t) \rightarrow \frac{1}{2}e^{3t}(c_3(t^2 - 4t + 2) + 2c_1(t - 3)t - c_2(t - 4)t)$$

6.27 problem problem 27

Internal problem ID [384]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.6, Multiple Eigenvalue Solutions. Page 451

Problem number: problem 27.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = -3x_1(t) + 5x_2(t) - 5x_3(t)$$

$$x_2'(t) = 3x_1(t) - x_2(t) + 3x_3(t)$$

$$x_3'(t) = 8x_1(t) - 8x_2(t) + 10x_3(t)$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 56

```
dsolve([diff(x__1(t),t)=-3*x__1(t)+5*x__2(t)-5*x__3(t),diff(x__2(t),t)=3*x__1(t)-1*x__2(t)+3
```

$$x_1(t) = \frac{e^{2t}(-5c_3t + 8c_1 - 5c_2 + c_3)}{8}$$

$$x_2(t) = \frac{e^{2t}(3c_3t + 8c_1 + 3c_2)}{8}$$

$$x_3(t) = e^{2t}(c_3t + c_2)$$

✓ Solution by Mathematica

Time used: 0.034 (sec). Leaf size: 174

```
DSolve[{x1'[t]==-3*x1[t]+5*x2[t]-5*x3[t],x2'[t]==4*x1[t]-1*x2[t]+4*x3[t],x3'[t]==8*x1[t]-8*x
```

$$x_1(t) \rightarrow \frac{1}{3}e^{2t} \left(-5(c_1 + c_3) \cos(\sqrt{3}t) - 5\sqrt{3}(c_1 - c_2 + c_3) \sin(\sqrt{3}t) + 8c_1 + 5c_3 \right)$$

$$x_2(t) \rightarrow \frac{1}{3}e^{2t} \left(3c_2 \cos(\sqrt{3}t) + \sqrt{3}(4c_1 - 3c_2 + 4c_3) \sin(\sqrt{3}t) \right)$$

$$x_3(t) \rightarrow \frac{1}{3}e^{2t} \left(8(c_1 + c_3) \cos(\sqrt{3}t) + 8\sqrt{3}(c_1 - c_2 + c_3) \sin(\sqrt{3}t) - 8c_1 - 5c_3 \right)$$

6.28 problem problem 28

Internal problem ID [385]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.6, Multiple Eigenvalue Solutions. Page 451

Problem number: problem 28.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = -15x_1(t) - 7x_2(t) + 4x_3(t)$$

$$x_2'(t) = 34x_1(t) + 16x_2(t) - 11x_3(t)$$

$$x_3'(t) = 17x_1(t) + 7x_2(t) + 5x_3(t)$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 75

```
dsolve([diff(x__1(t),t)=-15*x__1(t)-7*x__2(t)+4*x__3(t),diff(x__2(t),t)=34*x__1(t)+16*x__2(t)
```

$$x_1(t) = -\frac{e^{2t}(-119c_3t^2 - 238c_2t + 34c_3t + 14c_1 + 6c_2 - 2c_3)}{34}$$

$$x_2(t) = \frac{(-17c_3t^2 - 34c_2t + 4c_3t + 2c_1)e^{2t}}{2}$$

$$x_3(t) = e^{2t}(c_3t + c_2)$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 124

```
DSolve[{x1'[t]==-15*x1[t]-7*x2[t]+4*x3[t],x2'[t]==34*x1[t]+16*x2[t]-11*x3[t],x3'[t]==17*x1[t]
```

$$x1(t) \rightarrow \frac{1}{2}e^{2t}(c_1(119t^2 - 34t + 2) + 7c_2t(7t - 2) + c_3t(21t + 8))$$

$$x2(t) \rightarrow -\frac{1}{2}e^{2t}(17(17c_1 + 7c_2 + 3c_3)t^2 + (-68c_1 - 28c_2 + 22c_3)t - 2c_2)$$

$$x3(t) \rightarrow e^{2t}((17c_1 + 7c_2 + 3c_3)t + c_3)$$

6.29 problem problem 29

Internal problem ID [386]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.6, Multiple Eigenvalue Solutions. Page 451

Problem number: problem 29.

ODE order: 1.

ODE degree: 1.

Solve

$$\begin{aligned}x_1'(t) &= -x_1(t) + x_2(t) + x_3(t) - 2x_4(t) \\x_2'(t) &= 7x_1(t) - 4x_2(t) - 6x_3(t) + 11x_4(t) \\x_3'(t) &= 5x_1(t) - x_2(t) + x_3(t) + 3x_4(t) \\x_4'(t) &= 6x_1(t) - 2x_2(t) - 2x_3(t) + 6x_4(t)\end{aligned}$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 119

```
dsolve([diff(x__1(t),t)=-1*x__1(t)+1*x__2(t)+1*x__3(t)-2*x__4(t),diff(x__2(t),t)=7*x__1(t)-4
```

$$x_1(t) = -\frac{e^{-t}(c_4 t + c_3)}{2}$$

$$x_2(t) = -c_2 e^{2t} t + \frac{3c_4 e^{-t} t}{2} - c_1 e^{2t} + 2c_2 e^{2t} + \frac{3c_3 e^{-t}}{2} - \frac{c_4 e^{-t}}{2}$$

$$x_3(t) = c_2 e^{2t} t + c_1 e^{2t} + \frac{c_4 e^{-t} t}{2} + \frac{c_3 e^{-t}}{2}$$

$$x_4(t) = c_2 e^{2t} + c_3 e^{-t} + c_4 e^{-t} t$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 196

```
DSolve[{x1'[t]==-1*x1[t]+1*x2[t]+1*x3[t]-2*x4[t],x2'[t]==7*x1[t]-4*x2[t]-6*x3[t]+11*x4[t],x3
```

$$x1(t) \rightarrow e^{-t}((c_2 + c_3 - 2c_4)t + c_1)$$

$$x2(t) \rightarrow e^{-t}(c_1(e^{3t}(3-2t)-3) - 3c_2t - c_3e^{3t} - 3c_3t + 2c_4e^{3t} - c_4e^{3t}t + 6c_4t + c_2 + c_3 - 2c_4)$$

$$x3(t) \rightarrow e^{-t}(c_1(e^{3t}(2t+1)-1) + c_3e^{3t} - t(-c_4(e^{3t}+2) + c_2 + c_3))$$

$$x4(t) \rightarrow e^{-t}(2c_1(e^{3t}-1) - 2(c_2 + c_3 - 2c_4)t + c_4e^{3t})$$

6.30 problem problem 30

Internal problem ID [387]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.6, Multiple Eigenvalue Solutions. Page 451

Problem number: problem 30.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 2x_1(t) + x_2(t) - 2x_3(t) + x_4(t)$$

$$x_2'(t) = 3x_2(t) - 5x_3(t) + 3x_4(t)$$

$$x_3'(t) = -13x_2(t) + 22x_3(t) - 12x_4(t)$$

$$x_4'(t) = -27x_2(t) + 45x_3(t) - 25x_4(t)$$

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 95

```
dsolve([diff(x__1(t),t)=2*x__1(t)+1*x__2(t)-2*x__3(t)+1*x__4(t),diff(x__2(t),t)=0*x__1(t)+3*
```

$$x_1(t) = \frac{(-c_2 t + 5c_1) e^{2t}}{5}$$

$$x_2(t) = -\frac{e^{-t}(3c_4 t + 3c_3 + 2c_4)}{9}$$

$$x_3(t) = \frac{3c_2 e^{2t}}{5} + \frac{c_3 e^{-t}}{3} + \frac{c_4 e^{-t} t}{3} - \frac{c_4 e^{-t}}{9}$$

$$x_4(t) = c_2 e^{2t} + c_3 e^{-t} + c_4 e^{-t} t$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 161

```
DSolve[{x1'[t]==2*x1[t]+1*x2[t]-2*x3[t]+1*x4[t],x2'[t]==0*x1[t]+3*x2[t]-5*x3[t]+3*x4[t],x3'[t]==-2*x1[t]+1*x2[t]-3*x3[t]+2*x4[t],x4'[t]==1*x1[t]-1*x2[t]+2*x3[t]-1*x4[t]},x1[t],x2[t],x3[t],x4[t],t]
```

$$x1(t) \rightarrow e^{2t}((c_2 - 2c_3 + c_4)t + c_1)$$

$$x2(t) \rightarrow e^{-t}(4c_2t - 5c_3t + 3c_4t + c_2)$$

$$x3(t) \rightarrow e^{-t}(c_2(-4t - 3e^{3t} + 3) + c_3(5t + 6e^{3t} - 5) - 3c_4(t + e^{3t} - 1))$$

$$x4(t) \rightarrow e^{-t}(c_2(-12t - 5e^{3t} + 5) + 5c_3(3t + 2e^{3t} - 2) - c_4(9t + 5e^{3t} - 6))$$

6.31 problem problem 31

Internal problem ID [388]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.6, Multiple Eigenvalue Solutions. Page 451

Problem number: problem 31.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 35x_1(t) - 12x_2(t) + 4x_3(t) + 30x_4(t)$$

$$x_2'(t) = 22x_1(t) - 8x_2(t) + 3x_3(t) + 19x_4(t)$$

$$x_3'(t) = -10x_1(t) + 3x_2(t) - 9x_4(t)$$

$$x_4'(t) = -27x_1(t) + 9x_2(t) - 3x_3(t) - 23x_4(t)$$

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 112

```
dsolve([diff(x__1(t),t)=35*x__1(t)-12*x__2(t)+4*x__3(t)+30*x__4(t),diff(x__2(t),t)=22*x__1(t)
```

$$x_1(t) = -\frac{e^t(6c_4t^2 + 6c_3t + 2c_4t + 6c_2 + c_3 - c_4)}{6}$$

$$x_2(t) = \frac{e^t(-2c_4t^2 - 2c_3t - 10c_4t + 4c_1 - 2c_2 - 5c_3 + 6c_4)}{12}$$

$$x_3(t) = \frac{e^t(6c_4t^2 + 6c_3t - 2c_4t + 12c_1 + 6c_2 - c_3)}{12}$$

$$x_4(t) = e^t(c_4t^2 + c_3t + c_2)$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 207

```
DSolve[{x1'[t]==35*x1[t]-12*x2[t]+4*x3[t]+30*x4[t],x2'[t]==22*x1[t]-8*x2[t]+3*x3[t]+19*x4[t]}
```

$$x1(t) \rightarrow e^t(c_1(21t^2 + 34t + 1) - 3c_2t(3t + 4) + c_3t(3t + 4) + 6c_4t(3t + 5))$$

$$x2(t) \rightarrow \frac{1}{2}e^t((7c_1 - 3c_2 + c_3 + 6c_4)t^2 + 2(22c_1 - 9c_2 + 3c_3 + 19c_4)t + 2c_2)$$

$$x3(t) \rightarrow \frac{1}{2}e^t(-3(7c_1 - 3c_2 + c_3 + 6c_4)t^2 - 2(10c_1 - 3c_2 + c_3 + 9c_4)t + 2c_3)$$

$$x4(t) \rightarrow e^t(-3(7c_1 - 3c_2 + c_3 + 6c_4)t^2 - 3(9c_1 - 3c_2 + c_3 + 8c_4)t + c_4)$$

6.32 problem problem 32

Internal problem ID [389]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.6, Multiple Eigenvalue Solutions. Page 451

Problem number: problem 32.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 11x_1(t) - x_2(t) + 26x_3(t) + 6x_4(t) - 3x_5(t)$$

$$x_2'(t) = 3x_2(t)$$

$$x_3'(t) = -9x_1(t) - 24x_3(t) - 6x_4(t) + 3x_5(t)$$

$$x_4'(t) = 3x_1(t) + 9x_3(t) + 5x_4(t) - x_5(t)$$

$$x_5'(t) = -48x_1(t) - 3x_2(t) - 138x_3(t) - 30x_4(t) + 18x_5(t)$$

✓ Solution by Maple

Time used: 0.094 (sec). Leaf size: 116

```
dsolve([diff(x__1(t),t)=11*x__1(t)-1*x__2(t)+26*x__3(t)+6*x__4(t)-3*x__5(t),diff(x__2(t),t)=
```

$$x_1(t) = 8c_2e^{2t} + \frac{25c_3e^{3t}}{3} - 3c_1e^{3t} + \frac{c_4e^{2t}}{3} + \frac{c_5e^{3t}}{3}$$

$$x_2(t) = \frac{e^{3t}(6c_1 - 16c_3 - c_5)}{3}$$

$$x_3(t) = -3c_2e^{2t} - 3c_3e^{3t} + c_1e^{3t}$$

$$x_4(t) = c_2e^{2t} + c_3e^{3t}$$

$$x_5(t) = c_4e^{2t} + c_5e^{3t}$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 211

```
DSolve[{x1'[t]==11*x1[t]-1*x2[t]+26*x3[t]+6*x4[t]-3*x5[t],x2'[t]==0*x1[t]+3*x2[t],x3'[t]==-9
```

$$x1(t) \rightarrow e^{2t}(c_1(9e^t - 8) - (c_2 - 26c_3 - 6c_4 + 3c_5)(e^t - 1))$$

$$x2(t) \rightarrow c_2 e^{3t}$$

$$x3(t) \rightarrow -e^{2t}(9c_1(e^t - 1) + c_3(26e^t - 27) + 3(2c_4 - c_5)(e^t - 1))$$

$$x4(t) \rightarrow e^{2t}(3c_1(e^t - 1) + 9c_3(e^t - 1) + 3c_4e^t - c_5e^t - 2c_4 + c_5)$$

$$x5(t) \rightarrow -e^{2t}(48c_1(e^t - 1) + 3c_2(e^t - 1) + 138c_3e^t + 30c_4e^t - 16c_5e^t - 138c_3 - 30c_4 + 15c_5)$$

6.33 problem problem 33

Internal problem ID [390]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.6, Multiple Eigenvalue Solutions. Page 451

Problem number: problem 33.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 3x_1(t) - 4x_2(t) + x_3(t)$$

$$x_2'(t) = 4x_1(t) + 3x_2(t) + x_4(t)$$

$$x_3'(t) = 3x_3(t) - 4x_4(t)$$

$$x_4'(t) = 4x_3(t) + 3x_4(t)$$

✓ Solution by Maple

Time used: 0.75 (sec). Leaf size: 139

```
dsolve([diff(x__1(t),t)=3*x__1(t)-4*x__2(t)+1*x__3(t)+0*x__4(t),diff(x__2(t),t)=4*x__1(t)+3*
```

$$x_1(t) = \frac{e^{3t}(4 \cos(4t) c_3 t - 4 \sin(4t) c_4 t + 4c_2 \cos(4t) - c_4 \cos(4t) - 4c_1 \sin(4t))}{4}$$

$$x_2(t) = \frac{e^{3t}(4 \cos(4t) c_4 t + 4 \sin(4t) c_3 t + 4c_1 \cos(4t) + 4c_2 \sin(4t) - \sin(4t) c_4)}{4}$$

$$x_3(t) = e^{3t}(\cos(4t) c_3 - \sin(4t) c_4)$$

$$x_4(t) = e^{3t}(c_4 \cos(4t) + c_3 \sin(4t))$$

✓ Solution by Mathematica

Time used: 0.1 (sec). Leaf size: 120

```
DSolve[{x1'[t]==3*x1[t]-4*x2[t]+1*x3[t]+0*x4[t],x2'[t]==4*x1[t]+3*x2[t]+0*x3[t]+1*x4[t],x3'[t]==-4*x1[t]-3*x2[t]+0*x3[t]+0*x4[t],x4'[t]==1*x1[t]-3*x2[t]+0*x3[t]+0*x4[t]},x1,x2,x3,x4,t]
```

$$x1(t) \rightarrow e^{3t}((c_3 t + c_1) \cos(4t) - (c_4 t + c_2) \sin(4t))$$

$$x2(t) \rightarrow e^{3t}((c_4 t + c_2) \cos(4t) + (c_3 t + c_1) \sin(4t))$$

$$x3(t) \rightarrow e^{3t}(c_3 \cos(4t) - c_4 \sin(4t))$$

$$x4(t) \rightarrow e^{3t}(c_4 \cos(4t) + c_3 \sin(4t))$$

6.34 problem problem 34

Internal problem ID [391]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Section 7.6, Multiple Eigenvalue Solutions. Page 451

Problem number: problem 34.

ODE order: 1.

ODE degree: 1.

Solve

$$x_1'(t) = 2x_1(t) - 8x_3(t) - 3x_4(t)$$

$$x_2'(t) = -18x_1(t) - x_2(t)$$

$$x_3'(t) = -9x_1(t) - 3x_2(t) - 25x_3(t) - 9x_4(t)$$

$$x_4'(t) = 33x_1(t) + 10x_2(t) + 90x_3(t) + 32x_4(t)$$

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 208

```
dsolve([diff(x__1(t),t)=2*x__1(t)+0*x__2(t)-8*x__3(t)-3*x__4(t),diff(x__2(t),t)=-18*x__1(t)-
```

$$x_1(t) = e^{2t}(\cos(3t) c_3 t - \sin(3t) c_4 t + c_1 \cos(3t) - 3 \cos(3t) c_4 - c_2 \sin(3t) - 3c_3 \sin(3t))$$

$$x_2(t) = -e^{2t}(3 \cos(3t) c_3 t + 3 \cos(3t) c_4 t + 3 \sin(3t) c_3 t - 3 \sin(3t) c_4 t + 3c_1 \cos(3t) + 3c_2 \cos(3t) + 9c_3 \cos(3t) - 10 \cos(3t) c_4 + 3c_1 \sin(3t) - 3c_2 \sin(3t) - 10c_3 \sin(3t) - 9 \sin(3t) c_4)$$

$$x_3(t) = e^{2t}(c_3 \cos(3t) - \sin(3t) c_4)$$

$$x_4(t) = e^{2t}(\cos(3t) c_4 t + \sin(3t) c_3 t + c_2 \cos(3t) + c_1 \sin(3t))$$

✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 482

`DSolve[{x1'[t]==2*x1[t]+0*x2[t]-8*x3[t]-3*x4[t],x2'[t]==-18*x1[t]-1*x2[t]+0*x3[t]+0*x4[t],x3`

$$x1(t) \rightarrow \frac{1}{2}e^{(2-3i)t} (c_1(e^{6it}(1+3it) - 3it + 1) + i(3c_3 + c_4)(-1 + e^{6it}) + t(ic_2(-1 + e^{6it}) + c_3((1+9i)e^{6it} + (1-9i)) + 3ic_4(-1 + e^{6it})))$$

$$x2(t) \rightarrow -\frac{1}{2}e^{(2-3i)t} (c_1((9-9i)t + e^{6it}((9+9i)t - 3i) + 3i) + c_2((3-3i)t + e^{6it}(-1 + (3+3i)t) - 1) + 10ic_3e^{6it} + (30+24i)c_3e^{6it}t + (30-24i)c_3t + 3ic_4e^{6it} + (9+9i)c_4e^{6it}t + (9-9i)c_4t - 10ic_3 - 3ic_4)$$

$$x3(t) \rightarrow \frac{1}{2}e^{(2-3i)t} (3ic_1(-1 + e^{6it}) + ic_2(-1 + e^{6it}) + (1+9i)c_3e^{6it} + 3ic_4e^{6it} + (1-9i)c_3 - 3ic_4)$$

$$x4(t) \rightarrow \frac{1}{2}e^{(2-3i)t} (c_1(3t + e^{6it}(3t - 10i) + 10i) + c_2(t + e^{6it}(t - 3i) + 3i) - 27ic_3e^{6it} + (9-i)c_3e^{6it}t + (9+i)c_3t + (1-9i)c_4e^{6it} + 3c_4e^{6it}t + 3c_4t + 27ic_3 + (1+9i)c_4)$$

**7 Chapter 11 Power series methods. Section 11.1
Introduction and Review of power series. Page
615**

7.1	problem problem 1	173
7.2	problem problem 2	174
7.3	problem problem 3	175
7.4	problem problem 4	176
7.5	problem problem 5	177
7.6	problem problem 6	178
7.7	problem problem 7	179
7.8	problem problem 8	180
7.9	problem problem 9	181
7.10	problem problem 10	182
7.11	problem problem 11	183
7.12	problem problem 12	184
7.13	problem problem 13	185
7.14	problem problem 14	186
7.15	problem problem 15	187
7.16	problem problem 16	188
7.17	problem problem 17	189
7.18	problem problem 18	190
7.19	problem problem 19	191
7.20	problem problem 20	192
7.21	problem problem 21	193
7.22	problem problem 22	194
7.23	problem problem 23	195
7.24	problem problem 26(a)	197

7.1 problem problem 1

Internal problem ID [392]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.1 Introduction and Review of power series. Page 615

Problem number: problem 1.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$y' - y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
Order:=6;  
dsolve(diff(y(x),x)=y(x),y(x),type='series',x=0);
```

$$y(x) = \left(1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5\right) y(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 37

```
AsymptoticDSolveValue[y'[x]==y[x],y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{x^5}{120} + \frac{x^4}{24} + \frac{x^3}{6} + \frac{x^2}{2} + x + 1 \right)$$

7.2 problem problem 2

Internal problem ID [393]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.1 Introduction and Review of power series. Page 615

Problem number: problem 2.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_quadrature]`

$$y' - 4y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 36

```
Order:=6;  
dsolve(diff(y(x),x)=4*y(x),y(x),type='series',x=0);
```

$$y(x) = \left(1 + 4x + 8x^2 + \frac{32}{3}x^3 + \frac{32}{3}x^4 + \frac{128}{15}x^5\right)y(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 37

```
AsymptoticDSolveValue[y'[x]==4*y[x],y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{128x^5}{15} + \frac{32x^4}{3} + \frac{32x^3}{3} + 8x^2 + 4x + 1 \right)$$

7.3 problem problem 3

Internal problem ID [394]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.1 Introduction and Review of power series. Page 615

Problem number: problem 3.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [_quadrature]

$$2y' + 3y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 36

```
Order:=6;  
dsolve(2*diff(y(x),x)+3*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{3}{2}x + \frac{9}{8}x^2 - \frac{9}{16}x^3 + \frac{27}{128}x^4 - \frac{81}{1280}x^5\right) y(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 41

```
AsymptoticDSolveValue[2*y'[x]+3*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(-\frac{81x^5}{1280} + \frac{27x^4}{128} - \frac{9x^3}{16} + \frac{9x^2}{8} - \frac{3x}{2} + 1 \right)$$

7.4 problem problem 4

Internal problem ID [395]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.1 Introduction and Review of power series. Page 615

Problem number: problem 4.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' + 2yx = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
Order:=6;  
dsolve(diff(y(x),x)+2*x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - x^2 + \frac{1}{2}x^4\right) y(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 20

```
AsymptoticDSolveValue[y'[x]+2*x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{x^4}{2} - x^2 + 1\right)$$

7.5 problem problem 5

Internal problem ID [396]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.1 Introduction and Review of power series. Page 615

Problem number: problem 5.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y' - x^2y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
Order:=6;  
dsolve(diff(y(x),x)=x^2*y(x),y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{x^3}{3}\right)y(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 15

```
AsymptoticDSolveValue[y'[x]==x^2*y[x],y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{x^3}{3} + 1\right)$$

7.6 problem problem 6

Internal problem ID [397]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.1 Introduction and Review of power series. Page 615

Problem number: problem 6.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$(-2 + x)y' + y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 36

```
Order:=6;  
dsolve((x-2)*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{1}{2}x + \frac{1}{4}x^2 + \frac{1}{8}x^3 + \frac{1}{16}x^4 + \frac{1}{32}x^5\right)y(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 41

```
AsymptoticDSolveValue[(x-2)*y'[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{x^5}{32} + \frac{x^4}{16} + \frac{x^3}{8} + \frac{x^2}{4} + \frac{x}{2} + 1 \right)$$

7.7 problem problem 7

Internal problem ID [398]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.1 Introduction and Review of power series. Page 615

Problem number: problem 7.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [`_separable`]

$$(2x - 1)y' + 2y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 36

```
Order:=6;  
dsolve((2*x-1)*diff(y(x),x)+2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (32x^5 + 16x^4 + 8x^3 + 4x^2 + 2x + 1)y(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 31

```
AsymptoticDSolveValue[(2*x-1)*y'[x]+2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1(32x^5 + 16x^4 + 8x^3 + 4x^2 + 2x + 1)$$

7.8 problem problem 8

Internal problem ID [399]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.1 Introduction and Review of power series. Page 615

Problem number: problem 8.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$2(x+1)y' - y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 36

```
Order:=6;  
dsolve(2*(x+1)*diff(y(x),x)=y(x),y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{1}{2}x - \frac{1}{8}x^2 + \frac{1}{16}x^3 - \frac{5}{128}x^4 + \frac{7}{256}x^5\right)y(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 41

```
AsymptoticDSolveValue[2*(x+1)*y'[x]==y[x],y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{7x^5}{256} - \frac{5x^4}{128} + \frac{x^3}{16} - \frac{x^2}{8} + \frac{x}{2} + 1 \right)$$

7.9 problem problem 9

Internal problem ID [400]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.1 Introduction and Review of power series. Page 615

Problem number: problem 9.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$(x - 1)y' + 2y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 36

```
Order:=6;  
dsolve((x-1)*diff(y(x),x)+2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (6x^5 + 5x^4 + 4x^3 + 3x^2 + 2x + 1)y(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 31

```
AsymptoticDSolveValue[(x-1)*y'[x]+2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1(6x^5 + 5x^4 + 4x^3 + 3x^2 + 2x + 1)$$

7.10 problem problem 10

Internal problem ID [401]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.1 Introduction and Review of power series. Page 615

Problem number: problem 10.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$2(x-1)y' - 3y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 36

```
Order:=6;  
dsolve(2*(x-1)*diff(y(x),x)=3*y(x),y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{3}{2}x + \frac{3}{8}x^2 + \frac{1}{16}x^3 + \frac{3}{128}x^4 + \frac{3}{256}x^5\right) y(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 41

```
AsymptoticDSolveValue[2*(x-1)*y'[x]==3*y[x],y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{3x^5}{256} + \frac{3x^4}{128} + \frac{x^3}{16} + \frac{3x^2}{8} - \frac{3x}{2} + 1 \right)$$

7.11 problem problem 11

Internal problem ID [402]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.1 Introduction and Review of power series. Page 615

Problem number: problem 11.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
Order:=6;  
dsolve(diff(y(x),x$2)=y(x),y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{1}{2}x^2 + \frac{1}{24}x^4\right) y(0) + \left(x + \frac{1}{6}x^3 + \frac{1}{120}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 42

```
AsymptoticDSolveValue[y''[x]==y[x],y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{x^5}{120} + \frac{x^3}{6} + x \right) + c_1 \left(\frac{x^4}{24} + \frac{x^2}{2} + 1 \right)$$

7.12 problem problem 12

Internal problem ID [403]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.1 Introduction and Review of power series. Page 615

Problem number: problem 12.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 4y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 34

```
Order:=6;  
dsolve(diff(y(x),x$2)=4*y(x),y(x),type='series',x=0);
```

$$y(x) = \left(1 + 2x^2 + \frac{2}{3}x^4\right) y(0) + \left(x + \frac{2}{3}x^3 + \frac{2}{15}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 40

```
AsymptoticDSolveValue[y''[x]==4*y[x],y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{2x^5}{15} + \frac{2x^3}{3} + x \right) + c_1 \left(\frac{2x^4}{3} + 2x^2 + 1 \right)$$

7.13 problem problem 13

Internal problem ID [404]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.1 Introduction and Review of power series. Page 615

Problem number: problem 13.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 9y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
Order:=6;  
dsolve(diff(y(x),x$2)+9*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{9}{2}x^2 + \frac{27}{8}x^4\right) y(0) + \left(x - \frac{3}{2}x^3 + \frac{27}{40}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 42

```
AsymptoticDSolveValue[y''[x]+9*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{27x^5}{40} - \frac{3x^3}{2} + x \right) + c_1 \left(\frac{27x^4}{8} - \frac{9x^2}{2} + 1 \right)$$

7.14 problem problem 14

Internal problem ID [405]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.1 Introduction and Review of power series. Page 615

Problem number: problem 14.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + y = x$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 38

```
Order:=6;  
dsolve(diff(y(x),x$2)+y(x)=x,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{2}x^2 + \frac{1}{24}x^4\right)y(0) + \left(x - \frac{1}{6}x^3 + \frac{1}{120}x^5\right)D(y)(0) + \frac{x^3}{6} - \frac{x^5}{120} + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.043 (sec). Leaf size: 56

```
AsymptoticDSolveValue[y''[x]+y[x]==x,y[x],{x,0,5}]
```

$$y(x) \rightarrow -\frac{x^5}{120} + \frac{x^3}{6} + c_2\left(\frac{x^5}{120} - \frac{x^3}{6} + x\right) + c_1\left(\frac{x^4}{24} - \frac{x^2}{2} + 1\right)$$

7.15 problem problem 15

Internal problem ID [406]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.1 Introduction and Review of power series. Page 615

Problem number: problem 15.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$y + y'x = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
Order:=6;  
dsolve(x*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{c_1}{x} + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 9

```
AsymptoticDSolveValue[x*y'[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{c_1}{x}$$

7.16 problem problem 16

Internal problem ID [407]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.1 Introduction and Review of power series. Page 615

Problem number: problem 16.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$2y'x - y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
Order:=6;  
dsolve(2*x*diff(y(x),x)=y(x),y(x),type='series',x=0);
```

$$y(x) = c_1\sqrt{x} + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 11

```
AsymptoticDSolveValue[2*x*y'[x]==y[x],y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1\sqrt{x}$$

7.17 problem problem 17

Internal problem ID [408]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.1 Introduction and Review of power series. Page 615

Problem number: problem 17.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [`_separable`]

$$y'x^2 + y = 0$$

With the expansion point for the power series method at $x = 0$.

 Solution by Maple

```
Order:=6;  
dsolve(x^2*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

No solution found

 Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 11

```
AsymptoticDSolveValue[x^2*y'[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 e^{\frac{1}{x}}$$

7.18 problem problem 18

Internal problem ID [409]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.1 Introduction and Review of power series. Page 615

Problem number: problem 18.

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type [separable]

$$x^3 y' - 2y = 0$$

With the expansion point for the power series method at $x = 0$.

 Solution by Maple

```
Order:=6;  
dsolve(x^3*diff(y(x),x)=2*y(x),y(x),type='series',x=0);
```

No solution found

 Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 13

```
AsymptoticDSolveValue[x^3*y'[x]==2*y[x],y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 e^{-\frac{1}{x^2}}$$

7.19 problem problem 19

Internal problem ID [410]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.1 Introduction and Review of power series. Page 615

Problem number: problem 19.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + 4y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 3]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
Order:=6;  
dsolve([diff(y(x),x$2)+4*y(x)=0,y(0) = 0, D(y)(0) = 3],y(x),type='series',x=0);
```

$$y(x) = 3x - 2x^3 + \frac{2}{5}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 19

```
AsymptoticDSolveValue[{y'[x]+4*y[x]==0,{y[0]==0,y'[0]==3}},y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{2x^5}{5} - 2x^3 + 3x$$

7.20 problem problem 20

Internal problem ID [411]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.1 Introduction and Review of power series. Page 615

Problem number: problem 20.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 4y = 0$$

With initial conditions

$$[y(0) = 2, y'(0) = 0]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
Order:=6;  
dsolve([diff(y(x),x$2)-4*y(x)=0,y(0) = 2, D(y)(0) = 0],y(x),type='series',x=0);
```

$$y(x) = 2 + 4x^2 + \frac{4}{3}x^4 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 17

```
AsymptoticDSolveValue[{y'[x]-4*y[x]==0,{y[0]==2,y'[0]==0}},y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{4x^4}{3} + 4x^2 + 2$$

7.21 problem problem 21

Internal problem ID [412]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.1 Introduction and Review of power series. Page 615

Problem number: problem 21.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' - 2y' + y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 1]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
Order:=6;
```

```
dsolve([diff(y(x),x$2)-2*diff(y(x),x)+y(x)=0,y(0) = 0, D(y)(0) = 1],y(x),type='series',x=0);
```

$$y(x) = x + x^2 + \frac{1}{2}x^3 + \frac{1}{6}x^4 + \frac{1}{24}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 29

```
AsymptoticDSolveValue[{y'[x]-2*y'[x]+y[x]==0,{y[0]==0,y'[0]==1}},y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{x^5}{24} + \frac{x^4}{6} + \frac{x^3}{2} + x^2 + x$$

7.22 problem problem 22

Internal problem ID [413]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.1 Introduction and Review of power series. Page 615

Problem number: problem 22.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_x]]`

$$y'' + y' - 2y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = -2]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
Order:=6;
```

```
dsolve([diff(y(x),x$2)+diff(y(x),x)-2*y(x)=0,y(0) = 1, D(y)(0) = -2],y(x),type='series',x=0)
```

$$y(x) = 1 - 2x + 2x^2 - \frac{4}{3}x^3 + \frac{2}{3}x^4 - \frac{4}{15}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 34

```
AsymptoticDSolveValue[{y'[x]+y'[x]-2*y[x]==0,{y[0]==1,y'[0]==-2}},y[x],{x,0,5}]
```

$$y(x) \rightarrow -\frac{4x^5}{15} + \frac{2x^4}{3} - \frac{4x^3}{3} + 2x^2 - 2x + 1$$

7.23 problem problem 23

Internal problem ID [414]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.1 Introduction and Review of power series. Page 615

Problem number: problem 23.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$x^2y'' + y'x^2 + y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 907

```
Order:=6;
```

```
dsolve(x^2*diff(y(x),x$2)+x^2*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$\begin{aligned} y(x) = & \sqrt{x} \left(c_2 x^{\frac{i\sqrt{3}}{2}} \left(1 - \frac{1}{2}x + \frac{i\sqrt{3} + 3}{8i\sqrt{3} + 16}x^2 + \frac{-i\sqrt{3} - 5}{48i\sqrt{3} + 96}x^3 \right. \right. \\ & + \frac{1}{384} \frac{(i\sqrt{3} + 5)(i\sqrt{3} + 7)}{(i\sqrt{3} + 4)(i\sqrt{3} + 2)}x^4 - \frac{1}{3840} \frac{(i\sqrt{3} + 7)(i\sqrt{3} + 9)}{(i\sqrt{3} + 4)(i\sqrt{3} + 2)}x^5 + O(x^6) \left. \right) \\ & + c_1 x^{-\frac{i\sqrt{3}}{2}} \left(1 - \frac{1}{2}x + \frac{i\sqrt{3} - 3}{8i\sqrt{3} - 16}x^2 + \frac{-i\sqrt{3} + 5}{48i\sqrt{3} - 96}x^3 \right. \\ & \left. \left. + \frac{1}{384} \frac{(i\sqrt{3} - 5)(i\sqrt{3} - 7)}{(i\sqrt{3} - 4)(i\sqrt{3} - 2)}x^4 - \frac{1}{3840} \frac{(i\sqrt{3} - 7)(i\sqrt{3} - 9)}{(i\sqrt{3} - 4)(i\sqrt{3} - 2)}x^5 + O(x^6) \right) \right) \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 886

AsymptoticDSolveValue[x^2*y''[x]+x^2*y'[x]+y[x]==0,y[x],{x,0,5}]

$$\begin{aligned}
 & y(x) \\
 \rightarrow & \left(\frac{(-1)^{2/3} (1 - (-1)^{2/3}) (2 - (-1)^{2/3}) (3 - (-1)^{2/3}) (4 - (-1)^{2/3})}{(1 - (-1)^{2/3}) (1 - (-1)^{2/3}) (1 + (1 - (-1)^{2/3}) (2 - (-1)^{2/3})) (1 + (2 - (-1)^{2/3}) (3 - (-1)^{2/3})) (1 + (3 - (-1)^{2/3}) (4 - (-1)^{2/3}))} \right. \\
 & - \frac{(-1)^{2/3} (1 - (-1)^{2/3}) (2 - (-1)^{2/3}) (3 - (-1)^{2/3}) x^4}{(1 - (-1)^{2/3}) (1 - (-1)^{2/3}) (1 + (1 - (-1)^{2/3}) (2 - (-1)^{2/3})) (1 + (2 - (-1)^{2/3}) (3 - (-1)^{2/3})) (1 + (3 - (-1)^{2/3}) (4 - (-1)^{2/3}))} \\
 & + \frac{(-1)^{2/3} (1 - (-1)^{2/3}) (2 - (-1)^{2/3}) x^3}{(1 - (-1)^{2/3}) (1 - (-1)^{2/3}) (1 + (1 - (-1)^{2/3}) (2 - (-1)^{2/3})) (1 + (2 - (-1)^{2/3}) (3 - (-1)^{2/3}))} \\
 & - \frac{(-1)^{2/3} (1 - (-1)^{2/3}) x^2}{(1 - (-1)^{2/3}) (1 - (-1)^{2/3}) (1 + (1 - (-1)^{2/3}) (2 - (-1)^{2/3}))} \\
 & \left. + \frac{(-1)^{2/3} x}{1 - (-1)^{2/3} (1 - (-1)^{2/3})} \right) \\
 & + 1 \Big) c_1 x^{-(-1)^{2/3}} + \left(- \frac{\sqrt[3]{-1} (1 + \sqrt[3]{-1}) (2 + \sqrt[3]{-1}) (3 + \sqrt[3]{-1})}{(1 + \sqrt[3]{-1}) (1 + \sqrt[3]{-1}) (1 + (1 + \sqrt[3]{-1}) (2 + \sqrt[3]{-1})) (1 + (2 + \sqrt[3]{-1}) (3 + \sqrt[3]{-1}))} \right)
 \end{aligned}$$

7.24 problem problem 26(a)

Internal problem ID [415]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.1 Introduction and Review of power series. Page 615

Problem number: problem 26(a).

ODE order: 1.

ODE degree: 1.

CAS Maple gives this as type `[_quadrature]`

$$y' - y^2 = 1$$

With initial conditions

$$[y(0) = 0]$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 6

```
dsolve([diff(y(x),x)=1+y(x)^2,y(0) = 0],y(x), singsol=all)
```

$$y(x) = \tan(x)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 7

```
DSolve[{y'[x]==1+y[x]^2,{y[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \tan(x)$$

8 Chapter 11 Power series methods. Section 11.2 Power series solutions. Page 624

8.1	problem problem 1	199
8.2	problem problem 2	200
8.3	problem problem 3	201
8.4	problem problem 4	202
8.5	problem problem 5	203
8.6	problem problem 6	204
8.7	problem problem 7	205
8.8	problem problem 8	206
8.9	problem problem 9	207
8.10	problem problem 10	208
8.11	problem problem 11	209
8.12	problem problem 12	210
8.13	problem problem 13	211
8.14	problem problem 14	212
8.15	problem problem 15	213
8.16	problem problem 16	214
8.17	problem problem 17	215
8.18	problem problem 18	216
8.19	problem problem 19	217
8.20	problem problem 20	218
8.21	problem problem 21	219
8.22	problem problem 22	220
8.23	problem problem 23	221
8.24	problem problem 24	222
8.25	problem problem 25	223
8.26	problem problem 26	224
8.27	problem problem 27	225
8.28	problem problem 28	226
8.29	problem problem 29	227
8.30	problem problem 30	228
8.31	problem problem 33	229
8.32	problem problem 34	230

8.1 problem problem 1

Internal problem ID [416]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.2 Power series solutions. Page 624

Problem number: problem 1.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$(x^2 - 1)y'' + 4y'x + 2y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
Order:=6;  
dsolve((x^2-1)*diff(y(x),x$2)+4*x*diff(y(x),x)+2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (x^4 + x^2 + 1)y(0) + (x^5 + x^3 + x)D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 26

```
AsymptoticDSolveValue[(x^2-1)*y''[x]+4*x*y'[x]+2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2(x^5 + x^3 + x) + c_1(x^4 + x^2 + 1)$$

8.2 problem problem 2

Internal problem ID [417]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.2 Power series solutions. Page 624

Problem number: problem 2.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$(x^2 + 2)y'' + 4y'x + 2y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
Order:=6;  
dsolve((x^2+2)*diff(y(x),x$2)+4*x*diff(y(x),x)+2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{2}x^2 + \frac{1}{4}x^4\right) y(0) + \left(x - \frac{1}{2}x^3 + \frac{1}{4}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 68

```
AsymptoticDSolveValue[(x^2+2)*y''[x]+4*y'[x]+2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(-\frac{x^5}{30} - \frac{x^4}{12} + \frac{x^3}{3} - \frac{x^2}{2} + 1 \right) + c_2 \left(-\frac{x^5}{15} - \frac{x^4}{12} + \frac{x^3}{2} - x^2 + x \right)$$

8.3 problem problem 3

Internal problem ID [418]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.2 Power series solutions. Page 624

Problem number: problem 3.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$y'' + y'x + y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
Order:=6;  
dsolve(diff(y(x),x$2)+x*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{2}x^2 + \frac{1}{8}x^4\right)y(0) + \left(x - \frac{1}{3}x^3 + \frac{1}{15}x^5\right)D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 42

```
AsymptoticDSolveValue[y'[x]+x*y'[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{x^5}{15} - \frac{x^3}{3} + x \right) + c_1 \left(\frac{x^4}{8} - \frac{x^2}{2} + 1 \right)$$

8.4 problem problem 4

Internal problem ID [419]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.2 Power series solutions. Page 624

Problem number: problem 4.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$(x^2 + 1)y'' + 6y'x + 4y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
Order:=6;  
dsolve((x^2+1)*diff(y(x),x$2)+6*x*diff(y(x),x)+4*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (3x^4 - 2x^2 + 1)y(0) + \left(x - \frac{5}{3}x^3 + \frac{7}{3}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 60

```
AsymptoticDSolveValue[(x^2+1)*y''[x]+6*y'[x]+4*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1(4x^5 - 5x^4 + 4x^3 - 2x^2 + 1) + c_2\left(\frac{77x^5}{15} - \frac{13x^4}{2} + \frac{16x^3}{3} - 3x^2 + x\right)$$

8.5 problem problem 5

Internal problem ID [420]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.2 Power series solutions. Page 624

Problem number: problem 5.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _missing_y]]`

$$(x^2 + 1)y'' + 2y'x = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 24

```
Order:=6;  
dsolve((x^2+1)*diff(y(x),x$2)+2*x*diff(y(x),x)=0,y(x),type='series',x=0);
```

$$y(x) = y(0) + \left(x - \frac{1}{3}x^3 + \frac{1}{5}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 25

```
AsymptoticDSolveValue[(x^2-3)*y''[x]+2*x*y'[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{x^5}{45} + \frac{x^3}{9} + x \right) + c_1$$

8.6 problem problem 6

Internal problem ID [421]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.2 Power series solutions. Page 624

Problem number: problem 6.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type [Gegenbauer]

$$(x^2 - 1)y'' - 6y'x + 12y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 25

```
Order:=6;
```

```
dsolve((x^2-1)*diff(y(x),x$2)-6*x*diff(y(x),x)+12*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (x^4 + 6x^2 + 1)y(0) + (x^3 + x)D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 25

```
AsymptoticDSolveValue[(x^2-1)*y''[x]-6*x*y'[x]+12*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2(x^3 + x) + c_1(x^4 + 6x^2 + 1)$$

8.7 problem problem 7

Internal problem ID [422]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.2 Power series solutions. Page 624

Problem number: problem 7.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x^2 + 3)y'' - 7y'x + 16y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
Order:=6;
```

```
dsolve((x^2+3)*diff(y(x),x$2)-7*x*diff(y(x),x)+16*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{8}{3}x^2 + \frac{8}{27}x^4\right)y(0) + \left(x - \frac{1}{2}x^3 + \frac{1}{120}x^5\right)D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 42

```
AsymptoticDSolveValue[(x^2+3)*y''[x]-7*x*y'[x]+16*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{x^5}{120} - \frac{x^3}{2} + x \right) + c_1 \left(\frac{8x^4}{27} - \frac{8x^2}{3} + 1 \right)$$

8.8 problem problem 8

Internal problem ID [423]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.2 Power series solutions. Page 624

Problem number: problem 8.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries], [_2nd_order, _linear,`

$$(-x^2 + 2)y'' - y'x + 16y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 34

```
Order:=6;
dsolve((2-x^2)*diff(y(x),x$2)-x*diff(y(x),x)+16*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (2x^4 - 4x^2 + 1)y(0) + \left(x - \frac{5}{4}x^3 + \frac{7}{32}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 38

```
AsymptoticDSolveValue[(2-x^2)*y''[x]-x*y'[x]+16*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{7x^5}{32} - \frac{5x^3}{4} + x \right) + c_1 (2x^4 - 4x^2 + 1)$$

8.9 problem problem 9

Internal problem ID [424]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.2 Power series solutions. Page 624

Problem number: problem 9.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type [Gegenbauer]

$$(x^2 - 1)y'' + 8y'x + 12y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
Order:=6;
```

```
dsolve((x^2-1)*diff(y(x),x$2)+8*x*diff(y(x),x)+12*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (15x^4 + 6x^2 + 1)y(0) + \left(x + \frac{10}{3}x^3 + 7x^5\right)D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 36

```
AsymptoticDSolveValue[(x^2-1)*y''[x]+8*x*y'[x]+12*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(7x^5 + \frac{10x^3}{3} + x\right) + c_1(15x^4 + 6x^2 + 1)$$

8.10 problem problem 10

Internal problem ID [425]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.2 Power series solutions. Page 624

Problem number: problem 10.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$3y'' + y'x - 4y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
Order:=6;  
dsolve(3*diff(y(x),x$2)+x*diff(y(x),x)-4*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{2}{3}x^2 + \frac{1}{27}x^4\right) y(0) + \left(x + \frac{1}{6}x^3 + \frac{1}{360}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 42

```
AsymptoticDSolveValue[3*y''[x]+x*y'[x]-4*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{x^5}{360} + \frac{x^3}{6} + x \right) + c_1 \left(\frac{x^4}{27} + \frac{2x^2}{3} + 1 \right)$$

8.11 problem problem 11

Internal problem ID [426]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.2 Power series solutions. Page 624

Problem number: problem 11.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$5y'' - 2y'x + 10y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
Order:=6;  
dsolve(5*diff(y(x),x$2)-2*x*diff(y(x),x)+10*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - x^2 + \frac{1}{10}x^4\right)y(0) + \left(\frac{4}{375}x^5 - \frac{4}{15}x^3 + x\right)D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 40

```
AsymptoticDSolveValue[5*y''[x]-2*x*y'[x]+10*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{4x^5}{375} - \frac{4x^3}{15} + x \right) + c_1 \left(\frac{x^4}{10} - x^2 + 1 \right)$$

8.12 problem problem 12

Internal problem ID [427]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.2 Power series solutions. Page 624

Problem number: problem 12.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - y'x^2 - 3yx = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 24

```
Order:=6;  
dsolve(diff(y(x),x$2)-x^2*diff(y(x),x)-3*x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{x^3}{2}\right) y(0) + \left(x + \frac{1}{3}x^4\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 28

```
AsymptoticDSolveValue[y'[x]-x^2*y'[x]-3*x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{x^4}{3} + x \right) + c_1 \left(\frac{x^3}{2} + 1 \right)$$

8.13 problem problem 13

Internal problem ID [428]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.2 Power series solutions. Page 624

Problem number: problem 13.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$y'' + y'x^2 + 2yx = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
Order:=6;  
dsolve(diff(y(x),x$2)+x^2*diff(y(x),x)+2*x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{x^3}{3}\right) y(0) + \left(x - \frac{1}{4}x^4\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 28

```
AsymptoticDSolveValue[y'[x]+x^2*y'[x]+2*x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(x - \frac{x^4}{4}\right) + c_1 \left(1 - \frac{x^3}{3}\right)$$

8.14 problem problem 14

Internal problem ID [429]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.2 Power series solutions. Page 624

Problem number: problem 14.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$y'' + yx = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
Order:=6;  
dsolve(diff(y(x),x$2)+x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{x^3}{6}\right) y(0) + \left(x - \frac{1}{12}x^4\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 28

```
AsymptoticDSolveValue[y''[x]+x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(x - \frac{x^4}{12}\right) + c_1 \left(1 - \frac{x^3}{6}\right)$$

8.15 problem problem 15

Internal problem ID [430]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.2 Power series solutions. Page 624

Problem number: problem 15.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$y'' + x^2y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
Order:=6;  
dsolve(diff(y(x),x$2)+x^2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{x^4}{12}\right) y(0) + \left(x - \frac{1}{20}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 28

```
AsymptoticDSolveValue[y'[x]+x^2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(x - \frac{x^5}{20}\right) + c_1 \left(1 - \frac{x^4}{12}\right)$$

8.16 problem problem 16

Internal problem ID [431]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.2 Power series solutions. Page 624

Problem number: problem 16.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x^2 + 1)y'' + 2y'x - 2y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 1]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 5

```
Order:=6;
dsolve([(1+x^2)*diff(y(x),x$2)+2*x*diff(y(x),x)-2*y(x)=0,y(0) = 0, D(y)(0) = 1],y(x),type='s
```

$$y(x) = x$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 4

```
AsymptoticDSolveValue[{{(1+x^2)*y'[x]+2*x*y'[x]-2*y[x]==0,{y[0]==0,y'[0]==1}},y[x],{x,0,5}]
```

$$y(x) \rightarrow x$$

8.17 problem problem 17

Internal problem ID [432]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.2 Power series solutions. Page 624

Problem number: problem 17.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + y'x - 2y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 0]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
Order:=6;
```

```
dsolve([diff(y(x),x$2)+x*diff(y(x),x)-2*y(x)=0,y(0) = 1, D(y)(0) = 0],y(x),type='series',x=0)
```

$$y(x) = x^2 + 1$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 19

```
AsymptoticDSolveValue[{y''[x]+x*y'[x]-2*y[x]==0,{y[0]==0,y'[0]==1}},y[x],{x,0,5}]
```

$$y(x) \rightarrow -\frac{x^5}{120} + \frac{x^3}{6} + x$$

8.18 problem problem 18

Internal problem ID [433]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.2 Power series solutions. Page 624

Problem number: problem 18.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$y'' + (x - 1)y' + y = 0$$

With initial conditions

$$[y(1) = 2, y'(1) = 0]$$

With the expansion point for the power series method at $x = 1$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
Order:=6;  
dsolve([diff(y(x),x$2)+(x-1)*diff(y(x),x)+y(x)=0,y(1) = 2, D(y)(1) = 0],y(x),type='series',x
```

$$y(x) = 2 - (x - 1)^2 + \frac{1}{4}(x - 1)^4 + O((x - 1)^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 21

```
AsymptoticDSolveValue[{y'[x]+(x-1)*y'[x]+y[x]==0,{y[1]==2,y'[1]==0}},y[x],{x,1,5}]
```

$$y(x) \rightarrow \frac{1}{4}(x - 1)^4 - (x - 1)^2 + 2$$

8.19 problem problem 19

Internal problem ID [434]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.2 Power series solutions. Page 624

Problem number: problem 19.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$(-x^2 + 2x)y'' - 6(x - 1)y' - 4y = 0$$

With initial conditions

$$[y(1) = 0, y'(1) = 1]$$

With the expansion point for the power series method at $x = 1$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
Order:=6;
dsolve([(2*x-x^2)*diff(y(x),x$2)-6*(x-1)*diff(y(x),x)-4*y(x)=0,y(1) = 0, D(y)(1) = 1],y(x),t
```

$$y(x) = (x - 1) + \frac{5}{3}(x - 1)^3 + \frac{7}{3}(x - 1)^5 + O((x - 1)^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 24

```
AsymptoticDSolveValue[{{(2*x-x^2)*y'[x]-6*(x-1)*y'[x]-4*y[x]==0,{y[1]==0,y'[1]==1}},y[x],{x,
```

$$y(x) \rightarrow \frac{7}{3}(x - 1)^5 + \frac{5}{3}(x - 1)^3 + x - 1$$

8.20 problem problem 20

Internal problem ID [435]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.2 Power series solutions. Page 624

Problem number: problem 20.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x^2 - 6x + 10) y'' - 4(x - 3) y' + 6y = 0$$

With initial conditions

$$[y(3) = 2, y'(3) = 0]$$

With the expansion point for the power series method at $x = 3$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
Order:=6;
dsolve([(x^2-6*x+10)*diff(y(x),x$2)-4*(x-3)*diff(y(x),x)+6*y(x)=0,y(3) = 2, D(y)(3) = 0],y(x)
```

$$y(x) = -6x^2 + 36x - 52$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 12

```
AsymptoticDSolveValue[{(x^2-6*x+10)*y''[x]-4*(x-3)*y'[x]+6*y[x]==0,{y[3]==2,y'[3]==0}},y[x],
```

$$y(x) \rightarrow 2 - 6(x - 3)^2$$

8.21 problem problem 21

Internal problem ID [436]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.2 Power series solutions. Page 624

Problem number: problem 21.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _exact, _linear, _homogeneous]]`

$$(4x^2 + 16x + 17)y'' - 8y = 0$$

With initial conditions

$$[y(-2) = 1, y'(-2) = 0]$$

With the expansion point for the power series method at $x = -2$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
Order:=6;
dsolve([(4*x^2+16*x+17)*diff(y(x),x$2)=8*y(x),y(-2) = 1, D(y)(-2) = 0],y(x),type='series',x=
```

$$y(x) = 4x^2 + 16x + 17$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 12

```
AsymptoticDSolveValue[{(4*x^2+16*x+17)*y'[x]==8*y[x],{y[-2]==1,y'[-2]==0}},y[x],{x,-2,5}]
```

$$y(x) \rightarrow 4(x + 2)^2 + 1$$

8.22 problem problem 22

Internal problem ID [437]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.2 Power series solutions. Page 624

Problem number: problem 22.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x^2 + 6x)y'' + (3x + 9)y' - 3y = 0$$

With initial conditions

$$[y(-3) = 1, y'(-3) = 0]$$

With the expansion point for the power series method at $x = -3$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

Order:=6;

```
dsolve([(x^2+6*x)*diff(y(x),x$2)+(3*x+9)*diff(y(x),x)-3*y(x)=0,y(-3) = 1, D(y)(-3) = 0],y(x))
```

$$y(x) = 1 - \frac{1}{6}(x + 3)^2 - \frac{5}{648}(x + 3)^4 + O((x + 3)^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 23

```
AsymptoticDSolveValue[{{(x^2+6*x)*y''[x]+(3*x+9)*y'[x]-3*y[x]==0,{y[-3]==1,y'[-3]==0}},y[x],{
```

$$y(x) \rightarrow -\frac{5}{648}(x + 3)^4 - \frac{1}{6}(x + 3)^2 + 1$$

8.23 problem problem 23

Internal problem ID [438]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.2 Power series solutions. Page 624

Problem number: problem 23.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + (x + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 49

```
Order:=6;  
dsolve(diff(y(x),x$2)+(1+x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{30}x^5\right) y(0) \\ + \left(x - \frac{1}{6}x^3 - \frac{1}{12}x^4 + \frac{1}{120}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 63

```
AsymptoticDSolveValue[y''[x]+(1+x)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{x^5}{120} - \frac{x^4}{12} - \frac{x^3}{6} + x \right) + c_1 \left(\frac{x^5}{30} + \frac{x^4}{24} - \frac{x^3}{6} - \frac{x^2}{2} + 1 \right)$$

8.24 problem problem 24

Internal problem ID [439]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.2 Power series solutions. Page 624

Problem number: problem 24.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x^2 - 1)y'' + 2y'x + 2yx = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 39

```
Order:=6;  
dsolve((x^2-1)*diff(y(x),x$2)+2*x*diff(y(x),x)+2*x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{1}{3}x^3 + \frac{1}{5}x^5\right)y(0) + \left(x + \frac{1}{3}x^3 + \frac{1}{6}x^4 + \frac{1}{5}x^5\right)D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 49

```
AsymptoticDSolveValue[(x^2+1)*y''[x]+2*x*y'[x]+2*x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{x^5}{5} - \frac{x^3}{3} + 1 \right) + c_2 \left(\frac{x^5}{5} - \frac{x^4}{6} - \frac{x^3}{3} + x \right)$$

8.25 problem problem 25

Internal problem ID [440]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.2 Power series solutions. Page 624

Problem number: problem 25.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + y'x^2 + x^2y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
Order:=6;  
dsolve(diff(y(x),x$2)+x^2*diff(y(x),x)+x^2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{x^4}{12}\right) y(0) + \left(x - \frac{1}{12}x^4 - \frac{1}{20}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 35

```
AsymptoticDSolveValue[y''[x]+x^2*y'[x]+x^2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(1 - \frac{x^4}{12}\right) + c_2 \left(-\frac{x^5}{20} - \frac{x^4}{12} + x\right)$$

8.26 problem problem 26

Internal problem ID [441]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.2 Power series solutions. Page 624

Problem number: problem 26.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$(x^3 + 1) y'' + yx^4 = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
Order:=6;  
dsolve((1+x^3)*diff(y(x),x$2)+x^4*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = y(0) + D(y)(0)x + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 10

```
AsymptoticDSolveValue[(1+x^3)*y''[x]+x^4*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2x + c_1$$

8.27 problem problem 27

Internal problem ID [442]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.2 Power series solutions. Page 624

Problem number: problem 27.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + y'x + y(2x^2 + 1) = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = -1]$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
Order:=6;  
dsolve([diff(y(x),x$2)+x*diff(y(x),x)+(2*x^2+1)*y(x)=0,y(0) = 1, D(y)(0) = -1],y(x),type='se
```

$$y(x) = 1 - x - \frac{1}{2}x^2 + \frac{1}{3}x^3 - \frac{1}{24}x^4 + \frac{1}{30}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 49

```
AsymptoticDSolveValue[{{(x^2+1)*y'[x]+2*x*y'[x]+2*x*y[x]==0,{}},y[x]},{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(\frac{x^5}{5} - \frac{x^3}{3} + 1 \right) + c_2 \left(\frac{x^5}{5} - \frac{x^4}{6} - \frac{x^3}{3} + x \right)$$

8.28 problem problem 28

Internal problem ID [443]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.2 Power series solutions. Page 624

Problem number: problem 28.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' + y e^{-x} = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 44

```
Order:=6;  
dsolve(diff(y(x),x$2)+exp(-x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{40}x^5\right) y(0) + \left(x - \frac{1}{6}x^3 + \frac{1}{12}x^4 - \frac{1}{60}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 56

```
AsymptoticDSolveValue[y''[x]+Exp[-x]*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(-\frac{x^5}{60} + \frac{x^4}{12} - \frac{x^3}{6} + x \right) + c_1 \left(-\frac{x^5}{40} + \frac{x^3}{6} - \frac{x^2}{2} + 1 \right)$$

8.29 problem problem 29

Internal problem ID [444]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.2 Power series solutions. Page 624

Problem number: problem 29.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$\cos(x)y'' + y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 29

```
Order:=6;  
dsolve(cos(x)*diff(y(x),x$2)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{x^2}{2}\right) y(0) + \left(x - \frac{1}{6}x^3 - \frac{1}{60}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 35

```
AsymptoticDSolveValue[Cos[x]*y''[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(1 - \frac{x^2}{2}\right) + c_2 \left(-\frac{x^5}{60} - \frac{x^3}{6} + x\right)$$

8.30 problem problem 30

Internal problem ID [445]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.2 Power series solutions. Page 624

Problem number: problem 30.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type [Lienard]

$$xy'' + \sin(x)y' + yx = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 44

```
Order:=6;  
dsolve(x*diff(y(x),x$2)+sin(x)*diff(y(x),x)+x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{60}x^5\right)y(0) + \left(x - \frac{1}{2}x^2 + \frac{1}{18}x^4 - \frac{7}{360}x^5\right)D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 56

```
AsymptoticDSolveValue[x*y''[x]+Sin[x]*y'[x]+x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(-\frac{7x^5}{360} + \frac{x^4}{18} - \frac{x^2}{2} + x \right) + c_1 \left(-\frac{x^5}{60} + \frac{x^3}{6} - \frac{x^2}{2} + 1 \right)$$

8.31 problem problem 33

Internal problem ID [446]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.2 Power series solutions. Page 624

Problem number: problem 33.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_2nd_order, _with_linear_symmetries]]`

$$y'' - 2y'x + 2\alpha y = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 63

```
Order:=6;  
dsolve(diff(y(x),x$2)-2*x*diff(y(x),x)+2*alpha*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \alpha x^2 + \frac{\alpha(\alpha - 2)x^4}{6}\right) y(0) + \left(x - \frac{(\alpha - 1)x^3}{3} + \frac{(\alpha^2 - 4\alpha + 3)x^5}{30}\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 78

```
AsymptoticDSolveValue[y''[x]-2*x*y'[x]+2*\[Alpha]*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{\alpha^2 x^5}{30} - \frac{2\alpha x^5}{15} + \frac{x^5}{10} - \frac{\alpha x^3}{3} + \frac{x^3}{3} + x \right) + c_1 \left(\frac{\alpha^2 x^4}{6} - \frac{\alpha x^4}{3} - \alpha x^2 + 1 \right)$$

8.32 problem problem 34

Internal problem ID [447]

Book: Differential equations and linear algebra, 4th ed., Edwards and Penney

Section: Chapter 11 Power series methods. Section 11.2 Power series solutions. Page 624

Problem number: problem 34.

ODE order: 2.

ODE degree: 1.

CAS Maple gives this as type `[[_Emden, _Fowler]]`

$$y'' - yx = 0$$

With the expansion point for the power series method at $x = 0$.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
Order:=6;  
dsolve(diff(y(x),x$2)=x*y(x),y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{x^3}{6}\right) y(0) + \left(x + \frac{1}{12}x^4\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 28

```
AsymptoticDSolveValue[y''[x]==x*y[x],y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{x^4}{12} + x \right) + c_1 \left(\frac{x^3}{6} + 1 \right)$$