


A Solution Manual For

**An introduction to Ordinary Differential
Equations. Earl A. Coddington. Dover.
NY 1961**



Nasser M. Abbasi

May 15, 2024

Contents

1	Chapter 1.3 Introduction– Linear equations of First Order. Page 38	3
2	Chapter 1.6 Introduction– Linear equations of First Order. Page 41	105
3	Chapter 1. Introduction– Linear equations of First Order. Page 45	214
4	Chapter 2. Linear equations with constant coefficients. Page 52	332
5	Chapter 2. Linear equations with constant coefficients. Page 59	462
6	Chapter 2. Linear equations with constant coefficients. Page 69	504
7	Chapter 2. Linear equations with constant coefficients. Page 74	635
8	Chapter 2. Linear equations with constant coefficients. Page 79	674
9	Chapter 2. Linear equations with constant coefficients. Page 83	693
10	Chapter 2. Linear equations with constant coefficients. Page 89	751
11	Chapter 2. Linear equations with constant coefficients. Page 93	812
12	Chapter 3. Linear equations with variable coefficients. Page 108	899
13	Chapter 3. Linear equations with variable coefficients. Page 121	991
14	Chapter 3. Linear equations with variable coefficients. Page 124	1025
15	Chapter 3. Linear equations with variable coefficients. Page 130	1045
16	Chapter 4. Linear equations with Regular Singular Points. Page 149	161
17	Chapter 4. Linear equations with Regular Singular Points. Page 154	332
18	Chapter 4. Linear equations with Regular Singular Points. Page 159	469
19	Chapter 4. Linear equations with Regular Singular Points. Page 166	509
20	Chapter 4. Linear equations with Regular Singular Points. Page 182	657
21	Chapter 5. Existence and uniqueness of solutions to first order equations. Page 190	1669

22	Chapter 5. Existence and uniqueness of solutions to first order equations. Page 198	1806
23	Chapter 6. Existence and uniqueness of solutions to systems and nth order equations. Page 238	1882
24	Chapter 6. Existence and uniqueness of solutions to systems and nth order equations. Page 250	1986
25	Chapter 6. Existence and uniqueness of solutions to systems and nth order equations. Page 254	2014

1 Chapter 1.3 Introduction– Linear equations of First Order. Page 38

1.1	problem 1 (a)	4
1.2	problem 1 (b)	7
1.3	problem 1 (d)	23
1.4	problem 2 (a)	27
1.5	problem 2 (b)	42
1.6	problem 2 (c)	55
1.7	problem 2 (f)	65
1.8	problem 2 (h)	75
1.9	problem 3(a)	83
1.10	problem 4(a)	86
1.11	problem 5(a)	102

1.1 problem 1 (a)

1.1.1 Solving as quadrature ode	4
1.1.2 Maple step by step solution	5

Internal problem ID [5912]

Internal file name [OUTPUT/5160_Sunday_June_05_2022_03_26_32_PM_48421319/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.3 Introduction– Linear equations of First Order. Page 38

Problem number: 1 (a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

`[_quadrature]`

$$y' = e^{3x} + \sin(x)$$

1.1.1 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned} y &= \int e^{3x} + \sin(x) \, dx \\ &= \frac{e^{3x}}{3} - \cos(x) + c_1 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{e^{3x}}{3} - \cos(x) + c_1 \tag{1}$$




Figure 1: Slope field plot

Verification of solutions

$$y = \frac{e^{3x}}{3} - \cos(x) + c_1$$

Verified OK.

1.1.2 Maple step by step solution

Let's solve

$$y' = e^{3x} + \sin(x)$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Integrate both sides with respect to x

$$\int y' dx = \int (e^{3x} + \sin(x)) dx + c_1$$

- Evaluate integral

$$y = \frac{e^{3x}}{3} - \cos(x) + c_1$$

- Solve for y

$$y = \frac{e^{3x}}{3} - \cos(x) + c_1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve(diff(y(x),x)=exp(3*x)+sin(x),y(x), singsol=all)
```

$$y(x) = \frac{e^{3x}}{3} - \cos(x) + c_1$$

✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 21

```
DSolve[y'[x]==Exp[3*x]+Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{3x}}{3} - \cos(x) + c_1$$

1.2 problem 1 (b)

1.2.1	Solving as second order ode quadrature ode	7
1.2.2	Solving as second order linear constant coeff ode	8
1.2.3	Solving as second order integrable as is ode	11
1.2.4	Solving as second order ode missing y ode	12
1.2.5	Solving using Kovacic algorithm	14
1.2.6	Solving as exact linear second order ode ode	19
1.2.7	Maple step by step solution	21

Internal problem ID [5913]

Internal file name [OUTPUT/5161_Sunday_June_05_2022_03_26_33_PM_10950098/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.3 Introduction– Linear equations of First Order. Page 38

Problem number: 1 (b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "exact linear second order ode", "second_order_integrable_as_is", "second_order_ode_missing_y", "second_order_ode_quadrature", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _quadrature]]
```

$$y'' = x + 2$$

1.2.1 Solving as second order ode quadrature ode

Integrating once gives

$$y' = \frac{1}{2}x^2 + 2x + c_1$$

Integrating again gives

$$y = \frac{1}{6}x^3 + x^2 + c_1x + c_2$$

Summary

The solution(s) found are the following

$$y = \frac{1}{6}x^3 + x^2 + c_1x + c_2 \quad (1)$$




Figure 2: Slope field plot

Verification of solutions

$$y = \frac{1}{6}x^3 + x^2 + c_1x + c_2$$

Verified OK.

1.2.2 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 0, f(x) = x + 2$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 0$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 0$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{(0)^2 - (4)(1)(0)} \\ &= 0 \end{aligned}$$

Hence this is the case of a double root $\lambda_{1,2} = 0$. Therefore the solution is

$$y = c_1 1 + c_2 x \quad (1)$$

Therefore the homogeneous solution y_h is

$$y_h = c_2 x + c_1$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$1 + x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, x\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x, x^2\}]$$

Since x is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x^2, x^3\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_2x^3 + A_1x^2$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$6xA_2 + 2A_1 = x + 2$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = 1, A_2 = \frac{1}{6} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{1}{6}x^3 + x^2$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_2x + c_1) + \left(\frac{1}{6}x^3 + x^2 \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_2x + c_1 + \frac{1}{6}x^3 + x^2 \tag{1}$$




Figure 3: Slope field plot

Verification of solutions

$$y = c_2x + c_1 + \frac{1}{6}x^3 + x^2$$

Verified OK.

1.2.3 Solving as second order integrable as is ode

Integrating both sides of the ODE w.r.t x gives

$$\int y'' dx = \int (x + 2) dx$$

$$y' = \frac{1}{2}x^2 + 2x + c_1$$

Which is now solved for y . Integrating both sides gives

$$y = \int \frac{1}{2}x^2 + 2x + c_1 dx$$

$$= \frac{1}{6}x^3 + x^2 + c_1x + c_2$$

Summary

The solution(s) found are the following

$$y = \frac{1}{6}x^3 + x^2 + c_1x + c_2 \quad (1)$$




Figure 4: Slope field plot

Verification of solutions

$$y = \frac{1}{6}x^3 + x^2 + c_1x + c_2$$

Verified OK.

1.2.4 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x) - x - 2 = 0$$

Which is now solve for $p(x)$ as first order ode. Integrating both sides gives

$$\begin{aligned} p(x) &= \int x + 2 \, dx \\ &= \frac{1}{2}x^2 + 2x + c_1 \end{aligned}$$

Since $p = y'$ then the new first order ode to solve is

$$y' = \frac{1}{2}x^2 + 2x + c_1$$

Integrating both sides gives

$$\begin{aligned} y &= \int \frac{1}{2}x^2 + 2x + c_1 \, dx \\ &= \frac{1}{6}x^3 + x^2 + c_1x + c_2 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{1}{6}x^3 + x^2 + c_1x + c_2 \tag{1}$$




Figure 5: Slope field plot

Verification of solutions

$$y = \frac{1}{6}x^3 + x^2 + c_1x + c_2$$

Verified OK.

1.2.5 Solving using Kovacic algorithm

Writing the ode as

$$y'' = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 0 \quad (3)$$

$$C = 0$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$r = \frac{s}{t} \quad (5)$$
$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \quad (6)$$

Comparing the above to (5) shows that

$$s = 0$$

$$t = 1$$

Therefore eq. (4) becomes

$$z''(x) = 0 \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned} y_1 &= z_1 \\ &= 1 \end{aligned}$$

Which simplifies to

$$y_1 = 1$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= 1 \int \frac{1}{1} dx \\ &= 1(x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1(1) + c_2(1(x)) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_2x + c_1$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$1 + x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, x\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x, x^2\}]$$

Since x is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x^2, x^3\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_2x^3 + A_1x^2$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$6xA_2 + 2A_1 = x + 2$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = 1, A_2 = \frac{1}{6} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{1}{6}x^3 + x^2$$

Therefore the general solution is

$$\begin{aligned}y &= y_h + y_p \\ &= (c_2x + c_1) + \left(\frac{1}{6}x^3 + x^2\right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_2x + c_1 + \frac{1}{6}x^3 + x^2 \quad (1)$$




Figure 6: Slope field plot

Verification of solutions

$$y = c_2x + c_1 + \frac{1}{6}x^3 + x^2$$

Verified OK.

1.2.6 Solving as exact linear second order ode

An ode of the form

$$p(x)y'' + q(x)y' + r(x)y = s(x)$$

is exact if

$$p''(x) - q'(x) + r(x) = 0 \tag{1}$$

For the given ode we have

$$\begin{aligned} p(x) &= 1 \\ q(x) &= 0 \\ r(x) &= 0 \\ s(x) &= x + 2 \end{aligned}$$

Hence

$$\begin{aligned} p''(x) &= 0 \\ q'(x) &= 0 \end{aligned}$$

Therefore (1) becomes

$$0 - (0) + (0) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(x)y' + (q(x) - p'(x))y)' = s(x)$$

Integrating gives

$$p(x)y' + (q(x) - p'(x))y = \int s(x) dx$$

Substituting the above values for p, q, r, s gives

$$y' = \int x + 2 dx$$

We now have a first order ode to solve which is

$$y' = \frac{1}{2}x^2 + 2x + c_1$$

Integrating both sides gives

$$\begin{aligned}y &= \int \frac{1}{2}x^2 + 2x + c_1 \, dx \\ &= \frac{1}{6}x^3 + x^2 + c_1x + c_2\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{1}{6}x^3 + x^2 + c_1x + c_2 \quad (1)$$




Figure 7: Slope field plot

Verification of solutions

$$y = \frac{1}{6}x^3 + x^2 + c_1x + c_2$$

Verified OK.

1.2.7 Maple step by step solution

Let's solve

$$y'' = x + 2$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{0})}{2}$$

- Roots of the characteristic polynomial

$$r = 0$$

- 1st solution of the homogeneous ODE

$$y_1(x) = 1$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = x$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 + c_2 x + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = x + 2 \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} 1 & x \\ 0 & 1 \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\left(\int x(x+2) dx\right) + x\left(\int (x+2) dx\right)$$

- Compute integrals

$$y_p(x) = \frac{1}{6}x^3 + x^2$$

- Substitute particular solution into general solution to ODE

$$y = c_2x + c_1 + \frac{1}{6}x^3 + x^2$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)=2+x,y(x), singsol=all)
```

$$y(x) = \frac{1}{6}x^3 + x^2 + c_1x + c_2$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 22

```
DSolve[y''[x]==2+x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^3}{6} + x^2 + c_2x + c_1$$

1.3 problem 1 (d)

Internal problem ID [5914]

Internal file name [OUTPUT/5162_Sunday_June_05_2022_03_26_34_PM_30795132/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.3 Introduction– Linear equations of First Order. Page 38

Problem number: 1 (d).

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_3rd_order , _quadrature]]
```

$$y''' = x^2$$

This is higher order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE And y_p is a particular solution to the nonhomogeneous ODE. y_h is the solution to

$$y''' = 0$$

The characteristic equation is

$$\lambda^3 = 0$$

The roots of the above equation are

$$\lambda_1 = 0$$

$$\lambda_2 = 0$$

$$\lambda_3 = 0$$

Therefore the homogeneous solution is

$$y_h(x) = c_3x^2 + c_2x + c_1$$

The fundamental set of solutions for the homogeneous solution are the following

$$\begin{aligned}y_1 &= 1 \\y_2 &= x \\y_3 &= x^2\end{aligned}$$

Now the particular solution to the given ODE is found

$$y''' = x^2$$

The particular solution is found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x^2$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x, x^2\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, x, x^2\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x, x^2, x^3\}]$$

Since x is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x^2, x^3, x^4\}]$$

Since x^2 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x^3, x^4, x^5\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_3x^5 + A_2x^4 + A_1x^3$$

The unknowns $\{A_1, A_2, A_3\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$60x^2A_3 + 24xA_2 + 6A_1 = x^2$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = 0, A_2 = 0, A_3 = \frac{1}{60} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{x^5}{60}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_3x^2 + c_2x + c_1) + \left(\frac{x^5}{60} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_3x^2 + c_2x + c_1 + \frac{1}{60}x^5 \quad (1)$$

Verification of solutions

$$y = c_3x^2 + c_2x + c_1 + \frac{1}{60}x^5$$

Verified OK.

Maple trace

```
`Methods for third order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve(diff(y(x),x$3)=x^2,y(x), singsol=all)
```

$$y(x) = \frac{1}{60}x^5 + \frac{1}{2}c_1x^2 + c_2x + c_3$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 25

```
DSolve[y'''[x]==x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^5}{60} + c_3x^2 + c_2x + c_1$$

1.4 problem 2 (a)

1.4.1	Solving as separable ode	27
1.4.2	Solving as linear ode	29
1.4.3	Solving as homogeneousTypeD2 ode	30
1.4.4	Solving as first order ode lie symmetry lookup ode	32
1.4.5	Solving as exact ode	36
1.4.6	Maple step by step solution	40

Internal problem ID [5915]

Internal file name [OUTPUT/5163_Sunday_June_05_2022_03_26_35_PM_81840713/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.3 Introduction– Linear equations of First Order. Page 38

Problem number: 2 (a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "linear", "separable", "homogeneousTypeD2", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_separable]

$$y' + y \cos(x) = 0$$

1.4.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= -y \cos(x)\end{aligned}$$

Where $f(x) = -\cos(x)$ and $g(y) = y$. Integrating both sides gives

$$\begin{aligned}\frac{1}{y} dy &= -\cos(x) dx \\ \int \frac{1}{y} dy &= \int -\cos(x) dx \\ \ln(y) &= -\sin(x) + c_1 \\ y &= e^{-\sin(x)+c_1} \\ &= c_1 e^{-\sin(x)}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-\sin(x)} \tag{1}$$




Figure 8: Slope field plot

Verification of solutions

$$y = c_1 e^{-\sin(x)}$$

Verified OK.

1.4.2 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = \cos(x)$$

$$q(x) = 0$$

Hence the ode is

$$y' + y \cos(x) = 0$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int \cos(x) dx} \\ &= e^{\sin(x)}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx} \mu y &= 0 \\ \frac{d}{dx} (e^{\sin(x)} y) &= 0\end{aligned}$$

Integrating gives

$$e^{\sin(x)} y = c_1$$

Dividing both sides by the integrating factor $\mu = e^{\sin(x)}$ results in

$$y = c_1 e^{-\sin(x)}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-\sin(x)} \tag{1}$$




Figure 9: Slope field plot

Verification of solutions

$$y = c_1 e^{-\sin(x)}$$

Verified OK.

1.4.3 Solving as homogeneousTypeD2 ode

Using the change of variables $y = u(x)x$ on the above ode results in new ode in $u(x)$

$$u'(x)x + u(x) + u(x)x \cos(x) = 0$$

In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{u(\cos(x)x + 1)}{x} \end{aligned}$$

Where $f(x) = -\frac{\cos(x)x+1}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u} du &= -\frac{\cos(x)x+1}{x} dx \\ \int \frac{1}{u} du &= \int -\frac{\cos(x)x+1}{x} dx \\ \ln(u) &= -\sin(x) - \ln(x) + c_2 \\ u &= e^{-\sin(x)-\ln(x)+c_2} \\ &= c_2 e^{-\sin(x)-\ln(x)}\end{aligned}$$

Which simplifies to

$$u(x) = \frac{c_2 e^{-\sin(x)}}{x}$$

Therefore the solution y is

$$\begin{aligned}y &= xu \\ &= c_2 e^{-\sin(x)}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_2 e^{-\sin(x)} \tag{1}$$




Figure 10: Slope field plot

Verification of solutions

$$y = c_2 e^{-\sin(x)}$$

Verified OK.

1.4.4 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$\begin{aligned}y' &= -y \cos(x) \\y' &= \omega(x, y)\end{aligned}$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 4: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= e^{-\sin(x)}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^{-\sin(x)}} dy \end{aligned}$$

Which results in

$$S = e^{\sin(x)} y$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -y \cos(x)$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \cos(x) e^{\sin(x)} y \\ S_y &= e^{\sin(x)} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by

integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \tag{4}$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$e^{\sin(x)}y = c_1$$



Which simplifies to

$$e^{\sin(x)}y = c_1$$

Which gives

$$y = c_1 e^{-\sin(x)}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -y \cos(x)$ 	$R = x$ $S = e^{\sin(x)}y$	$\frac{dS}{dR} = 0$ 

Summary

The solution(s) found are the following

$$y = c_1 e^{-\sin(x)} \tag{1}$$




Figure 11: Slope field plot

Verification of solutions

$$y = c_1 e^{-\sin(x)}$$

Verified OK.

1.4.5 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}\left(-\frac{1}{y}\right) dy &= (\cos(x)) dx \\ (-\cos(x)) dx + \left(-\frac{1}{y}\right) dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -\cos(x) \\ N(x, y) &= -\frac{1}{y}\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-\cos(x)) \\ &= 0\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(-\frac{1}{y} \right) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \tag{1}$$

$$\frac{\partial \phi}{\partial y} = N \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\cos(x) dx \\ \phi &= -\sin(x) + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = -\frac{1}{y}$. Therefore equation (4) becomes

$$-\frac{1}{y} = 0 + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = -\frac{1}{y}$$

Integrating the above w.r.t y gives

$$\begin{aligned}\int f'(y) dy &= \int \left(-\frac{1}{y} \right) dy \\ f(y) &= -\ln(y) + c_1\end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\sin(x) - \ln(y) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\sin(x) - \ln(y)$$

The solution becomes

$$y = e^{-\sin(x)-c_1}$$

Summary

The solution(s) found are the following

$$y = e^{-\sin(x)-c_1} \tag{1}$$




Figure 12: Slope field plot

Verification of solutions

$$y = e^{-\sin(x)-c_1}$$

Verified OK.

1.4.6 Maple step by step solution

Let's solve

$$y' + y \cos(x) = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y} = -\cos(x)$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y} dx = \int -\cos(x) dx + c_1$$

- Evaluate integral

$$\ln(y) = -\sin(x) + c_1$$

- Solve for y

$$y = e^{-\sin(x)+c_1}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve(diff(y(x),x)+cos(x)*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{-\sin(x)}$$

✓ Solution by Mathematica

Time used: 0.029 (sec). Leaf size: 19

```
DSolve[y'[x]+Cos[x]*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-\sin(x)}$$

$$y(x) \rightarrow 0$$

1.5 problem 2 (b)

1.5.1	Solving as linear ode	42
1.5.2	Solving as first order ode lie symmetry lookup ode	44
1.5.3	Solving as exact ode	48
1.5.4	Maple step by step solution	52

Internal problem ID [5916]

Internal file name [OUTPUT/5164_Sunday_June_05_2022_03_26_36_PM_70038094/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.3 Introduction– Linear equations of First Order. Page 38

Problem number: 2 (b).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"linear", "exactWithIntegrationFactor", "first_order_ode_lie_symmetry_lookup"**

Maple gives the following as the ode type

`[_linear]`

$$y' + y \cos(x) = \cos(x) \sin(x)$$

1.5.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = \cos(x)$$
$$q(x) = \frac{\sin(2x)}{2}$$

Hence the ode is

$$y' + y \cos(x) = \frac{\sin(2x)}{2}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int \cos(x) dx} \\ &= e^{\sin(x)}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu) \left(\frac{\sin(2x)}{2} \right) \\ \frac{d}{dx}(e^{\sin(x)} y) &= (e^{\sin(x)}) \left(\frac{\sin(2x)}{2} \right) \\ d(e^{\sin(x)} y) &= \left(\frac{\sin(2x) e^{\sin(x)}}{2} \right) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}e^{\sin(x)} y &= \int \frac{\sin(2x) e^{\sin(x)}}{2} dx \\ e^{\sin(x)} y &= \sin(x) e^{\sin(x)} - e^{\sin(x)} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{\sin(x)}$ results in

$$y = e^{-\sin(x)} (\sin(x) e^{\sin(x)} - e^{\sin(x)}) + c_1 e^{-\sin(x)}$$

which simplifies to

$$y = \sin(x) - 1 + c_1 e^{-\sin(x)}$$

Summary

The solution(s) found are the following

$$y = \sin(x) - 1 + c_1 e^{-\sin(x)} \tag{1}$$




Figure 13: Slope field plot

Verification of solutions

$$y = \sin(x) - 1 + c_1 e^{-\sin(x)}$$

Verified OK.

1.5.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = -y \cos(x) + \cos(x) \sin(x)$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 7: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= e^{-\sin(x)}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^{-\sin(x)}} dy \end{aligned}$$

Which results in

$$S = e^{\sin(x)} y$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -y \cos(x) + \cos(x) \sin(x)$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \cos(x) e^{\sin(x)} y \\ S_y &= e^{\sin(x)} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \cos(x) e^{\sin(x)} \sin(x) \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \cos(R) e^{\sin(R)} \sin(R)$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by

integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 + e^{\sin(R)}(-1 + \sin(R)) \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$e^{\sin(x)}y = e^{\sin(x)}(-1 + \sin(x)) + c_1$$



Which simplifies to

$$e^{\sin(x)}y = e^{\sin(x)}(-1 + \sin(x)) + c_1$$

Which gives

$$y = e^{-\sin(x)}(\sin(x)e^{\sin(x)} - e^{\sin(x)} + c_1)$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -y \cos(x) + \cos(x) \sin(x)$ 	$R = x$ $S = e^{\sin(x)}y$	$\frac{dS}{dR} = \cos(R) e^{\sin(R)} \sin(R)$ 

Summary

The solution(s) found are the following

$$y = e^{-\sin(x)}(\sin(x)e^{\sin(x)} - e^{\sin(x)} + c_1) \quad (1)$$




Figure 14: Slope field plot

Verification of solutions

$$y = e^{-\sin(x)} (\sin(x) e^{\sin(x)} - e^{\sin(x)} + c_1)$$

Verified OK.

1.5.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}dy &= (-y \cos(x) + \cos(x) \sin(x)) dx \\ (y \cos(x) - \cos(x) \sin(x)) dx + dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= y \cos(x) - \cos(x) \sin(x) \\ N(x, y) &= 1\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(y \cos(x) - \cos(x) \sin(x)) \\ &= \cos(x)\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(1) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= 1((\cos(x)) - (0)) \\ &= \cos(x) \end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int A dx} \\ &= e^{\int \cos(x) dx} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^{\sin(x)} \\ &= e^{\sin(x)} \end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} for now so not to confuse them with the original M and N .

$$\begin{aligned} \bar{M} &= \mu M \\ &= e^{\sin(x)}(y \cos(x) - \cos(x) \sin(x)) \\ &= \cos(x) (-\sin(x) + y) e^{\sin(x)} \end{aligned}$$

And

$$\begin{aligned} \bar{N} &= \mu N \\ &= e^{\sin(x)}(1) \\ &= e^{\sin(x)} \end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned} \bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ (\cos(x) (-\sin(x) + y) e^{\sin(x)} + (e^{\sin(x)}) \frac{dy}{dx} &= 0 \end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\int \frac{\partial \phi}{\partial x} dx = \int \overline{M} dx$$

$$\int \frac{\partial \phi}{\partial x} dx = \int \cos(x) (-\sin(x) + y) e^{\sin(x)} dx$$

$$\phi = (y - \sin(x) + 1) e^{\sin(x)} + f(y) \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = e^{\sin(x)} + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = e^{\sin(x)}$. Therefore equation (4) becomes

$$e^{\sin(x)} = e^{\sin(x)} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = (y - \sin(x) + 1) e^{\sin(x)} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = (y - \sin(x) + 1) e^{\sin(x)}$$

The solution becomes

$$y = e^{-\sin(x)} (\sin(x) e^{\sin(x)} - e^{\sin(x)} + c_1)$$

Summary

The solution(s) found are the following

$$y = e^{-\sin(x)} (\sin(x) e^{\sin(x)} - e^{\sin(x)} + c_1) \quad (1)$$




Figure 15: Slope field plot

Verification of solutions

$$y = e^{-\sin(x)} (\sin(x) e^{\sin(x)} - e^{\sin(x)} + c_1)$$

Verified OK.

1.5.4 Maple step by step solution

Let's solve

$$y' + y \cos(x) = \cos(x) \sin(x)$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = -y \cos(x) + \cos(x) \sin(x)$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' + y \cos(x) = \cos(x) \sin(x)$$
- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) (y' + y \cos(x)) = \mu(x) \cos(x) \sin(x)$$
- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x) y)$

$$\mu(x) (y' + y \cos(x)) = \mu'(x) y + \mu(x) y'$$
- Isolate $\mu'(x)$

$$\mu'(x) = \mu(x) \cos(x)$$
- Solve to find the integrating factor

$$\mu(x) = e^{\sin(x)}$$
- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x) y) \right) dx = \int \mu(x) \cos(x) \sin(x) dx + c_1$$
- Evaluate the integral on the lhs

$$\mu(x) y = \int \mu(x) \cos(x) \sin(x) dx + c_1$$
- Solve for y

$$y = \frac{\int \mu(x) \cos(x) \sin(x) dx + c_1}{\mu(x)}$$
- Substitute $\mu(x) = e^{\sin(x)}$

$$y = \frac{\int \cos(x) e^{\sin(x)} \sin(x) dx + c_1}{e^{\sin(x)}}$$
- Evaluate the integrals on the rhs

$$y = \frac{\sin(x) e^{\sin(x)} - e^{\sin(x)} + c_1}{e^{\sin(x)}}$$
- Simplify

$$y = \sin(x) - 1 + c_1 e^{-\sin(x)}$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(diff(y(x),x)+cos(x)*y(x)=sin(x)*cos(x),y(x), singsol=all)
```

$$y(x) = \sin(x) - 1 + c_1 e^{-\sin(x)}$$

✓ Solution by Mathematica

Time used: 0.049 (sec). Leaf size: 18

```
DSolve[y'[x]+Cos[x]*y[x]==Sin[x]*Cos[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \sin(x) + c_1 e^{-\sin(x)} - 1$$

1.6 problem 2 (c)

1.6.1	Solving as second order linear constant coeff ode	55
1.6.2	Solving as second order ode can be made integrable ode	57
1.6.3	Solving using Kovacic algorithm	59
1.6.4	Maple step by step solution	63

Internal problem ID [5917]

Internal file name [OUTPUT/5165_Sunday_June_05_2022_03_26_38_PM_74591855/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.3 Introduction– Linear equations of First Order. Page 38

Problem number: 2 (c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "second_order_ode_can_be_made_integrable"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' - y = 0$$

1.6.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = -1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 1 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = -1$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(-1)} \\ &= \pm 1\end{aligned}$$

Hence

$$\lambda_1 = +1$$

$$\lambda_2 = -1$$

Which simplifies to

$$\lambda_1 = 1$$

$$\lambda_2 = -1$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(1)x} + c_2 e^{(-1)x}$$

Or

$$y = c_1 e^x + c_2 e^{-x}$$

Summary

The solution(s) found are the following

$$y = c_1 e^x + c_2 e^{-x} \tag{1}$$




Figure 16: Slope field plot

Verification of solutions

$$y = c_1 e^x + c_2 e^{-x}$$

Verified OK.

1.6.2 Solving as second order ode can be made integrable ode

Multiplying the ode by y' gives

$$y' y'' - y' y = 0$$

Integrating the above w.r.t x gives

$$\int (y' y'' - y' y) dx = 0$$

$$\frac{y'^2}{2} - \frac{y^2}{2} = c_2$$

Which is now solved for y . Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \sqrt{y^2 + 2c_1} \tag{1}$$

$$y' = -\sqrt{y^2 + 2c_1} \tag{2}$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int \frac{1}{\sqrt{y^2 + 2c_1}} dy = \int dx$$
$$\ln \left(y + \sqrt{y^2 + 2c_1} \right) = c_2 + x$$

Raising both side to exponential gives

$$y + \sqrt{y^2 + 2c_1} = e^{c_2+x}$$

Which simplifies to

$$y + \sqrt{y^2 + 2c_1} = c_3 e^x$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{1}{\sqrt{y^2 + 2c_1}} dy = \int dx$$
$$-\ln \left(y + \sqrt{y^2 + 2c_1} \right) = x + c_4$$

Raising both side to exponential gives

$$\frac{1}{y + \sqrt{y^2 + 2c_1}} = e^{x+c_4}$$

Which simplifies to

$$\frac{1}{y + \sqrt{y^2 + 2c_1}} = c_5 e^x$$

Summary

The solution(s) found are the following

$$y = \frac{(e^{2x} c_3^2 - 2c_1) e^{-x}}{2c_3} \quad (1)$$

$$y = -\frac{(2c_1 c_5^2 e^{2x} - 1) e^{-x}}{2c_5} \quad (2)$$




Figure 17: Slope field plot

Verification of solutions

$$y = \frac{(e^{2x}c_3^2 - 2c_1)e^{-x}}{2c_3}$$

Verified OK.

$$y = -\frac{(2c_1c_5^2e^{2x} - 1)e^{-x}}{2c_5}$$

Verified OK.

1.6.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' - y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 0 \tag{3}$$

$$C = -1$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{1}{1} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 10: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-x}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}
 y_1 &= z_1 \\
 &= e^{-x}
 \end{aligned}$$

Which simplifies to

$$y_1 = e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= e^{-x} \int \frac{1}{e^{-2x}} dx \\ &= e^{-x} \left(\frac{e^{2x}}{2} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-x}) + c_2 \left(e^{-x} \left(\frac{e^{2x}}{2} \right) \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-x} + \frac{c_2 e^x}{2} \tag{1}$$




Figure 18: Slope field plot

Verification of solutions

$$y = c_1 e^{-x} + \frac{c_2 e^x}{2}$$

Verified OK.

1.6.4 Maple step by step solution

Let's solve

$$y'' - y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 - 1 = 0$$

- Factor the characteristic polynomial

$$(r - 1)(r + 1) = 0$$

- Roots of the characteristic polynomial

- $r = (-1, 1)$
 - 1st solution of the ODE
 $y_1(x) = e^{-x}$
 - 2nd solution of the ODE
 $y_2(x) = e^x$
 - General solution of the ODE
 $y = c_1 y_1(x) + c_2 y_2(x)$
 - Substitute in solutions
 $y = c_1 e^{-x} + c_2 e^x$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(diff(y(x),x$2)-y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{-x} + e^x c_2$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 20

```
DSolve[y''[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^x + c_2 e^{-x}$$

1.7 problem 2 (f)

1.7.1	Solving as second order linear constant coeff ode	65
1.7.2	Solving as second order ode can be made integrable ode	67
1.7.3	Solving using Kovacic algorithm	69
1.7.4	Maple step by step solution	73

Internal problem ID [5918]

Internal file name [OUTPUT/5166_Sunday_June_05_2022_03_26_38_PM_21565163/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.3 Introduction– Linear equations of First Order. Page 38

Problem number: 2 (f).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "second_order_ode_can_be_made_integrable"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' + 4y = 0$$

1.7.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 4$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 4e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 4 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 4$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(4)} \\ &= \pm 2i\end{aligned}$$

Hence

$$\lambda_1 = +2i$$

$$\lambda_2 = -2i$$

Which simplifies to

$$\lambda_1 = 2i$$

$$\lambda_2 = -2i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 2$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0(c_1 \cos(2x) + c_2 \sin(2x))$$

Or

$$y = c_1 \cos(2x) + c_2 \sin(2x)$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(2x) + c_2 \sin(2x) \tag{1}$$




Figure 19: Slope field plot

Verification of solutions

$$y = c_1 \cos(2x) + c_2 \sin(2x)$$

Verified OK.

1.7.2 Solving as second order ode can be made integrable ode

Multiplying the ode by y' gives

$$y'y'' + 4y'y = 0$$

Integrating the above w.r.t x gives

$$\int (y'y'' + 4y'y) dx = 0$$

$$\frac{y'^2}{2} + 2y^2 = c_2$$

Which is now solved for y . Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \sqrt{-4y^2 + 2c_1} \tag{1}$$

$$y' = -\sqrt{-4y^2 + 2c_1} \tag{2}$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int \frac{1}{\sqrt{-4y^2 + 2c_1}} dy = \int dx$$
$$\frac{\arctan\left(\frac{2y}{\sqrt{-4y^2 + 2c_1}}\right)}{2} = c_2 + x$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{1}{\sqrt{-4y^2 + 2c_1}} dy = \int dx$$
$$-\frac{\arctan\left(\frac{2y}{\sqrt{-4y^2 + 2c_1}}\right)}{2} = x + c_3$$

Summary

The solution(s) found are the following

$$\frac{\arctan\left(\frac{2y}{\sqrt{-4y^2 + 2c_1}}\right)}{2} = c_2 + x \quad (1)$$

$$-\frac{\arctan\left(\frac{2y}{\sqrt{-4y^2 + 2c_1}}\right)}{2} = x + c_3 \quad (2)$$




Figure 20: Slope field plot

Verification of solutions

$$\frac{\arctan\left(\frac{2y}{\sqrt{-4y^2+2c_1}}\right)}{2} = c_2 + x$$

Verified OK.

$$-\frac{\arctan\left(\frac{2y}{\sqrt{-4y^2+2c_1}}\right)}{2} = x + c_3$$

Verified OK.

1.7.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 4y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= 1 \\B &= 0 \\C &= 4\end{aligned}\tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x)\tag{4}$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\&= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned}\tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-4}{1}\tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= -4 \\t &= 1\end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -4z(x)\tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 12: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -4$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(2x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}
 y_1 &= z_1 \\
 &= \cos(2x)
 \end{aligned}$$

Which simplifies to

$$y_1 = \cos(2x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(2x) \int \frac{1}{\cos(2x)^2} dx \\ &= \cos(2x) \left(\frac{\tan(2x)}{2} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(2x)) + c_2 \left(\cos(2x) \left(\frac{\tan(2x)}{2} \right) \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(2x) + \frac{c_2 \sin(2x)}{2} \tag{1}$$




Figure 21: Slope field plot

Verification of solutions

$$y = c_1 \cos(2x) + \frac{c_2 \sin(2x)}{2}$$

Verified OK.

1.7.4 Maple step by step solution

Let's solve

$$y'' + 4y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 + 4 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-16})}{2}$$

- Roots of the characteristic polynomial

$$r = (-2I, 2I)$$

- 1st solution of the ODE
 $y_1(x) = \cos(2x)$
- 2nd solution of the ODE
 $y_2(x) = \sin(2x)$
- General solution of the ODE
 $y = c_1 y_1(x) + c_2 y_2(x)$
- Substitute in solutions
 $y = c_1 \cos(2x) + c_2 \sin(2x)$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)+4*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 \sin(2x) + c_2 \cos(2x)$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 20

```
DSolve[y''[x]+4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos(2x) + c_2 \sin(2x)$$

1.8 problem 2 (h)

1.8.1	Solving as second order linear constant coeff ode	75
1.8.2	Solving as second order ode can be made integrable ode	77
1.8.3	Solving using Kovacic algorithm	78
1.8.4	Maple step by step solution	81

Internal problem ID [5919]

Internal file name [OUTPUT/5167_Sunday_June_05_2022_03_26_39_PM_73850312/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.3 Introduction– Linear equations of First Order. Page 38

Problem number: 2 (h).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "second_order_ode_can_be_made_integrable"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' + k^2y = 0$$

1.8.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = k^2$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + k^2 e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$k^2 + \lambda^2 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = k^2$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(k^2)} \\ &= \pm \sqrt{-k^2}\end{aligned}$$

Hence

$$\begin{aligned}\lambda_1 &= +\sqrt{-k^2} \\ \lambda_2 &= -\sqrt{-k^2}\end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= \sqrt{-k^2} \\ \lambda_2 &= -\sqrt{-k^2}\end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned}y &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ y &= c_1 e^{(\sqrt{-k^2})x} + c_2 e^{(-\sqrt{-k^2})x}\end{aligned}$$

Or

$$y = c_1 e^{\sqrt{-k^2}x} + c_2 e^{-\sqrt{-k^2}x}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{\sqrt{-k^2}x} + c_2 e^{-\sqrt{-k^2}x} \quad (1)$$

Verification of solutions

$$y = c_1 e^{\sqrt{-k^2}x} + c_2 e^{-\sqrt{-k^2}x}$$

Verified OK.

1.8.2 Solving as second order ode can be made integrable ode

Multiplying the ode by y' gives

$$y'y'' + k^2y'y = 0$$

Integrating the above w.r.t x gives

$$\int (y'y'' + k^2y'y) dx = 0$$
$$\frac{y'^2}{2} + \frac{y^2k^2}{2} = c_2$$

Which is now solved for y . Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \sqrt{-y^2k^2 + 2c_1} \quad (1)$$

$$y' = -\sqrt{-y^2k^2 + 2c_1} \quad (2)$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int \frac{1}{\sqrt{-y^2k^2 + 2c_1}} dy = \int dx$$
$$\frac{\arctan\left(\frac{\sqrt{k^2}y}{\sqrt{-y^2k^2+2c_1}}\right)}{\sqrt{k^2}} = c_2 + x$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{1}{\sqrt{-y^2k^2 + 2c_1}} dy = \int dx$$
$$-\frac{\arctan\left(\frac{\sqrt{k^2}y}{\sqrt{-y^2k^2+2c_1}}\right)}{\sqrt{k^2}} = x + c_3$$

Summary

The solution(s) found are the following

$$\frac{\arctan\left(\frac{\sqrt{k^2}y}{\sqrt{-y^2k^2+2c_1}}\right)}{\sqrt{k^2}} = c_2 + x \quad (1)$$

$$-\frac{\arctan\left(\frac{\sqrt{k^2}y}{\sqrt{-y^2k^2+2c_1}}\right)}{\sqrt{k^2}} = x + c_3 \quad (2)$$

Verification of solutions

$$\frac{\arctan\left(\frac{\sqrt{k^2}y}{\sqrt{-y^2k^2+2c_1}}\right)}{\sqrt{k^2}} = c_2 + x$$

Verified OK.

$$-\frac{\arctan\left(\frac{\sqrt{k^2}y}{\sqrt{-y^2k^2+2c_1}}\right)}{\sqrt{k^2}} = x + c_3$$

Verified OK.

1.8.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + k^2y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= k^2 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-k^2}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -k^2 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = (-k^2) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 14: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -k^2$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{\sqrt{-k^2} x}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned} y_1 &= z_1 \\ &= e^{\sqrt{-k^2} x} \end{aligned}$$

Which simplifies to

$$y_1 = e^{\sqrt{-k^2} x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= e^{\sqrt{-k^2} x} \int \frac{1}{e^{2\sqrt{-k^2} x}} dx \\ &= e^{\sqrt{-k^2} x} \left(\frac{\sqrt{-k^2} e^{-2\sqrt{-k^2} x}}{2k^2} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}
y &= c_1 y_1 + c_2 y_2 \\
&= c_1 \left(e^{\sqrt{-k^2} x} \right) + c_2 \left(e^{\sqrt{-k^2} x} \left(\frac{\sqrt{-k^2} e^{-2\sqrt{-k^2} x}}{2k^2} \right) \right)
\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{\sqrt{-k^2} x} + \frac{c_2 \sqrt{-k^2} e^{-\sqrt{-k^2} x}}{2k^2} \quad (1)$$

Verification of solutions

$$y = c_1 e^{\sqrt{-k^2} x} + \frac{c_2 \sqrt{-k^2} e^{-\sqrt{-k^2} x}}{2k^2}$$

Verified OK.

1.8.4 Maple step by step solution

Let's solve

$$y'' + k^2 y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$k^2 + r^2 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4k^2})}{2}$$

- Roots of the characteristic polynomial

$$r = (\sqrt{-k^2}, -\sqrt{-k^2})$$

- 1st solution of the ODE

$$y_1(x) = e^{\sqrt{-k^2} x}$$

- 2nd solution of the ODE

$$y_2(x) = e^{-\sqrt{-k^2} x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = c_1 e^{\sqrt{-k^2} x} + c_2 e^{-\sqrt{-k^2} x}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)+k^2*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 \sin(kx) + c_2 \cos(kx)$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 20

```
DSolve[y''[x]+k^2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos(kx) + c_2 \sin(kx)$$

1.9 problem 3(a)

1.9.1 Solving as quadrature ode	83
1.9.2 Maple step by step solution	84

Internal problem ID [5920]

Internal file name [OUTPUT/5168_Sunday_June_05_2022_03_26_40_PM_97624927/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.3 Introduction– Linear equations of First Order. Page 38

Problem number: 3(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$y' + 5y = 2$$

1.9.1 Solving as quadrature ode

Integrating both sides gives

$$\int \frac{1}{-5y + 2} dy = \int dx$$
$$-\frac{\ln(-5y + 2)}{5} = x + c_1$$

Raising both side to exponential gives

$$\frac{1}{(-5y + 2)^{\frac{1}{5}}} = e^{x+c_1}$$

Which simplifies to

$$\frac{1}{(-5y + 2)^{\frac{1}{5}}} = c_2 e^x$$

Summary

The solution(s) found are the following

$$y = -\frac{e^{-5x}}{5c_2^5} + \frac{2}{5} \quad (1)$$




Figure 22: Slope field plot

Verification of solutions

$$y = -\frac{e^{-5x}}{5c_2^5} + \frac{2}{5}$$

Verified OK.

1.9.2 Maple step by step solution

Let's solve

$$y' + 5y = 2$$

- Highest derivative means the order of the ODE is 1
- y'
- Separate variables

$$\frac{y'}{-5y+2} = 1$$

- Integrate both sides with respect to x

$$\int \frac{y'}{-5y+2} dx = \int 1 dx + c_1$$

- Evaluate integral

$$-\frac{\ln(-5y+2)}{5} = x + c_1$$

- Solve for y

$$y = -\frac{e^{-5x-5c_1}}{5} + \frac{2}{5}$$

Maple trace

```
`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)+5*y(x)=2,y(x), singsol=all)
```

$$y(x) = \frac{2}{5} + e^{-5x}c_1$$

✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 24

```
DSolve[y'[x]+5*y[x]==2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2}{5} + c_1 e^{-5x}$$

$$y(x) \rightarrow \frac{2}{5}$$

1.10 problem 4(a)

1.10.1 Solving as second order ode quadrature ode	86
1.10.2 Solving as second order linear constant coeff ode	87
1.10.3 Solving as second order integrable as is ode	90
1.10.4 Solving as second order ode missing y ode	91
1.10.5 Solving using Kovacic algorithm	93
1.10.6 Solving as exact linear second order ode ode	98
1.10.7 Maple step by step solution	100

Internal problem ID [5921]

Internal file name [OUTPUT/5169_Sunday_June_05_2022_03_26_41_PM_27157766/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.3 Introduction– Linear equations of First Order. Page 38

Problem number: 4(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "exact linear second order ode", "second_order_integrable_as_is", "second_order_ode_missing_y", "second_order_ode_quadrature", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _quadrature]]
```

$$y'' = 3x + 1$$

1.10.1 Solving as second order ode quadrature ode

Integrating once gives

$$y' = \frac{3}{2}x^2 + x + c_1$$

Integrating again gives

$$y = \frac{1}{2}x^3 + \frac{1}{2}x^2 + c_1x + c_2$$

Summary

The solution(s) found are the following

$$y = \frac{1}{2}x^3 + \frac{1}{2}x^2 + c_1x + c_2 \quad (1)$$




Figure 23: Slope field plot

Verification of solutions

$$y = \frac{1}{2}x^3 + \frac{1}{2}x^2 + c_1x + c_2$$

Verified OK.

1.10.2 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 0, f(x) = 3x + 1$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 0$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 0$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{(0)^2 - (4)(1)(0)} \\ &= 0 \end{aligned}$$

Hence this is the case of a double root $\lambda_{1,2} = 0$. Therefore the solution is

$$y = c_1 1 + c_2 x \quad (1)$$

Therefore the homogeneous solution y_h is

$$y_h = c_2 x + c_1$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$1 + x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, x\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x, x^2\}]$$

Since x is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x^2, x^3\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_2x^3 + A_1x^2$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$6xA_2 + 2A_1 = 3x + 1$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{2}, A_2 = \frac{1}{2} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{1}{2}x^3 + \frac{1}{2}x^2$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_2x + c_1) + \left(\frac{1}{2}x^3 + \frac{1}{2}x^2 \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_2x + c_1 + \frac{1}{2}x^3 + \frac{1}{2}x^2 \tag{1}$$




Figure 24: Slope field plot

Verification of solutions

$$y = c_2x + c_1 + \frac{1}{2}x^3 + \frac{1}{2}x^2$$

Verified OK.

1.10.3 Solving as second order integrable as is ode

Integrating both sides of the ODE w.r.t x gives

$$\int y'' dx = \int (3x + 1) dx$$

$$y' = \frac{3}{2}x^2 + x + c_1$$

Which is now solved for y . Integrating both sides gives

$$y = \int \frac{3}{2}x^2 + x + c_1 dx$$

$$= \frac{1}{2}x^3 + \frac{1}{2}x^2 + c_1x + c_2$$

Summary

The solution(s) found are the following

$$y = \frac{1}{2}x^3 + \frac{1}{2}x^2 + c_1x + c_2 \quad (1)$$




Figure 25: Slope field plot

Verification of solutions

$$y = \frac{1}{2}x^3 + \frac{1}{2}x^2 + c_1x + c_2$$

Verified OK.

1.10.4 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x) - 3x - 1 = 0$$

Which is now solve for $p(x)$ as first order ode. Integrating both sides gives

$$\begin{aligned} p(x) &= \int 3x + 1 \, dx \\ &= \frac{3}{2}x^2 + x + c_1 \end{aligned}$$

Since $p = y'$ then the new first order ode to solve is

$$y' = \frac{3}{2}x^2 + x + c_1$$

Integrating both sides gives

$$\begin{aligned} y &= \int \frac{3}{2}x^2 + x + c_1 \, dx \\ &= \frac{1}{2}x^3 + \frac{1}{2}x^2 + c_1x + c_2 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{1}{2}x^3 + \frac{1}{2}x^2 + c_1x + c_2 \tag{1}$$




Figure 26: Slope field plot

Verification of solutions

$$y = \frac{1}{2}x^3 + \frac{1}{2}x^2 + c_1x + c_2$$

Verified OK.

1.10.5 Solving using Kovacic algorithm

Writing the ode as

$$y'' = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 0 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 0 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 0 \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 17: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned} y_1 &= z_1 \\ &= 1 \end{aligned}$$

Which simplifies to

$$y_1 = 1$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= 1 \int \frac{1}{1} dx \\ &= 1(x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1(1) + c_2(1(x)) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_2x + c_1$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$1 + x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, x\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x, x^2\}]$$

Since x is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x^2, x^3\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_2x^3 + A_1x^2$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$6xA_2 + 2A_1 = 3x + 1$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{2}, A_2 = \frac{1}{2} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{1}{2}x^3 + \frac{1}{2}x^2$$

Therefore the general solution is

$$\begin{aligned}y &= y_h + y_p \\ &= (c_2x + c_1) + \left(\frac{1}{2}x^3 + \frac{1}{2}x^2\right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_2x + c_1 + \frac{1}{2}x^3 + \frac{1}{2}x^2 \quad (1)$$




Figure 27: Slope field plot

Verification of solutions

$$y = c_2x + c_1 + \frac{1}{2}x^3 + \frac{1}{2}x^2$$

Verified OK.

1.10.6 Solving as exact linear second order ode

An ode of the form

$$p(x) y'' + q(x) y' + r(x) y = s(x)$$

is exact if

$$p''(x) - q'(x) + r(x) = 0 \tag{1}$$

For the given ode we have

$$\begin{aligned} p(x) &= 1 \\ q(x) &= 0 \\ r(x) &= 0 \\ s(x) &= 3x + 1 \end{aligned}$$

Hence

$$\begin{aligned} p''(x) &= 0 \\ q'(x) &= 0 \end{aligned}$$

Therefore (1) becomes

$$0 - (0) + (0) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(x) y' + (q(x) - p'(x)) y)' = s(x)$$

Integrating gives

$$p(x) y' + (q(x) - p'(x)) y = \int s(x) dx$$

Substituting the above values for p, q, r, s gives

$$y' = \int 3x + 1 dx$$

We now have a first order ode to solve which is

$$y' = \frac{3}{2}x^2 + x + c_1$$

Integrating both sides gives

$$\begin{aligned}y &= \int \frac{3}{2}x^2 + x + c_1 \, dx \\ &= \frac{1}{2}x^3 + \frac{1}{2}x^2 + c_1x + c_2\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{1}{2}x^3 + \frac{1}{2}x^2 + c_1x + c_2 \quad (1)$$




Figure 28: Slope field plot

Verification of solutions

$$y = \frac{1}{2}x^3 + \frac{1}{2}x^2 + c_1x + c_2$$

Verified OK.

1.10.7 Maple step by step solution

Let's solve

$$y'' = 3x + 1$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{0})}{2}$$

- Roots of the characteristic polynomial

$$r = 0$$

- 1st solution of the homogeneous ODE

$$y_1(x) = 1$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = x$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 + c_2 x + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = 3x + 1 \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} 1 & x \\ 0 & 1 \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\left(\int (3x^2 + x) dx\right) + x\left(\int (3x + 1) dx\right)$$

- Compute integrals

$$y_p(x) = \frac{x^2(1+x)}{2}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 + c_2x + \frac{x^2(1+x)}{2}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
<- quadrature successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(diff(y(x),x$2)=3*x+1,y(x), singsol=all)
```

$$y(x) = \frac{1}{2}x^3 + \frac{1}{2}x^2 + c_1x + c_2$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 25

```
DSolve[y''[x]==3*x+1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}(x^3 + x^2 + 2c_2x + 2c_1)$$

1.11 problem 5(a)

1.11.1 Solving as quadrature ode 102

1.11.2 Maple step by step solution 103

Internal problem ID [5922]

Internal file name [OUTPUT/5170_Sunday_June_05_2022_03_26_42_PM_62334749/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.3 Introduction– Linear equations of First Order. Page 38

Problem number: 5(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$y' - yk = 0$$

1.11.1 Solving as quadrature ode

Integrating both sides gives

$$\int \frac{1}{yk} dy = \int dx$$
$$\frac{\ln(y)}{k} = x + c_1$$

Raising both side to exponential gives

$$e^{\frac{\ln(y)}{k}} = e^{x+c_1}$$

Which simplifies to

$$y^{\frac{1}{k}} = c_2 e^x$$

Summary

The solution(s) found are the following

$$y = (c_2 e^x)^k \tag{1}$$

Verification of solutions

$$y = (c_2 e^x)^k$$

Verified OK.

1.11.2 Maple step by step solution

Let's solve

$$y' - yk = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y} = k$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y} dx = \int k dx + c_1$$

- Evaluate integral

$$\ln(y) = kx + c_1$$

- Solve for y

$$y = e^{kx+c_1}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```


✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 10

```
dsolve(diff(y(x),x)=k*y(x),y(x), singsol=all)
```

$$y(x) = c_1 e^{kx}$$

✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 18

```
DSolve[y'[x]==k*y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{kx}$$

$$y(x) \rightarrow 0$$

2 Chapter 1.6 Introduction– Linear equations of First Order. Page 41

2.1	problem 1(a)	106
2.2	problem 1(b)	109
2.3	problem 1(c)	122
2.4	problem 1(d)	135
2.5	problem 1(e)	148
2.6	problem 2	161
2.7	problem 3	174
2.8	problem 4	177
2.9	problem 5	190
2.10	problem 7	203

2.1 problem 1(a)

2.1.1 Solving as quadrature ode	106
2.1.2 Maple step by step solution	107

Internal problem ID [5923]

Internal file name [OUTPUT/5171_Sunday_June_05_2022_03_26_43_PM_81309885/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.6 Introduction– Linear equations of First Order. Page 41

Problem number: 1(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$y' - 2y = 1$$

2.1.1 Solving as quadrature ode

Integrating both sides gives

$$\int \frac{1}{1+2y} dy = \int dx$$
$$\frac{\ln(1+2y)}{2} = x + c_1$$

Raising both side to exponential gives

$$\sqrt{1+2y} = e^{x+c_1}$$

Which simplifies to

$$\sqrt{1+2y} = c_2 e^x$$

Summary

The solution(s) found are the following

$$y = \frac{e^{2x} c_2^2}{2} - \frac{1}{2} \tag{1}$$




Figure 29: Slope field plot

Verification of solutions

$$y = \frac{e^{2x}c_2^2}{2} - \frac{1}{2}$$

Verified OK.

2.1.2 Maple step by step solution

Let's solve

$$y' - 2y = 1$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{2y+1} = 1$$

- Integrate both sides with respect to x

$$\int \frac{y'}{2y+1} dx = \int 1 dx + c_1$$

- Evaluate integral

$$\frac{\ln(2y+1)}{2} = x + c_1$$

- Solve for y

$$y = \frac{e^{2c_1+2x}}{2} - \frac{1}{2}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)-2*y(x)=1,y(x), singsol=all)
```

$$y(x) = -\frac{1}{2} + e^{2x}c_1$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 24

```
DSolve[y'[x]-2*y[x]==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{2} + c_1 e^{2x}$$
$$y(x) \rightarrow -\frac{1}{2}$$

2.2 problem 1(b)

2.2.1	Solving as linear ode	109
2.2.2	Solving as first order ode lie symmetry lookup ode	111
2.2.3	Solving as exact ode	115
2.2.4	Maple step by step solution	119

Internal problem ID [5924]

Internal file name [OUTPUT/5172_Sunday_June_05_2022_03_26_44_PM_38055412/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.6 Introduction– Linear equations of First Order. Page 41

Problem number: 1(b).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"linear", "exactWithIntegrationFactor", "first_order_ode_lie_symmetry_lookup"**

Maple gives the following as the ode type

```
[[_linear, `class A`]]
```

$$y + y' = e^x$$

2.2.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = 1$$

$$q(x) = e^x$$

Hence the ode is

$$y + y' = e^x$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int 1 dx} \\ &= e^x\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu)(e^x) \\ \frac{d}{dx}(y e^x) &= (e^x)(e^x) \\ d(y e^x) &= e^{2x} dx\end{aligned}$$

Integrating gives

$$\begin{aligned}y e^x &= \int e^{2x} dx \\ y e^x &= \frac{e^{2x}}{2} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^x$ results in

$$y = \frac{e^{-x}e^{2x}}{2} + c_1e^{-x}$$

which simplifies to

$$y = c_1e^{-x} + \frac{e^x}{2}$$

Summary

The solution(s) found are the following

$$y = c_1e^{-x} + \frac{e^x}{2} \tag{1}$$




Figure 30: Slope field plot

Verification of solutions

$$y = c_1 e^{-x} + \frac{e^x}{2}$$

Verified OK.

2.2.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$\begin{aligned} y' &= -y + e^x \\ y' &= \omega(x, y) \end{aligned}$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 21: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= e^{-x}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^{-x}} dy \end{aligned}$$

Which results in

$$S = y e^x$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -y + e^x$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= y e^x \\ S_y &= e^x \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = e^{2x} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = e^{2R}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by

integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \frac{e^{2R}}{2} + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$e^x y = \frac{e^{2x}}{2} + c_1$$



Which simplifies to

$$e^x y = \frac{e^{2x}}{2} + c_1$$

Which gives

$$y = \frac{(e^{2x} + 2c_1) e^{-x}}{2}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -y + e^x$ 	$R = x$ $S = y e^x$	$\frac{dS}{dR} = e^{2R}$ 

Summary

The solution(s) found are the following

$$y = \frac{(e^{2x} + 2c_1) e^{-x}}{2} \quad (1)$$




Figure 31: Slope field plot

Verification of solutions

$$y = \frac{(e^{2x} + 2c_1) e^{-x}}{2}$$

Verified OK.

2.2.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}dy &= (-y + e^x) dx \\ (y - e^x) dx + dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= y - e^x \\ N(x, y) &= 1\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(y - e^x) \\ &= 1\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(1) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= 1((1) - (0)) \\ &= 1 \end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int A dx} \\ &= e^{\int 1 dx} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^x \\ &= e^x \end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} for now so not to confuse them with the original M and N .

$$\begin{aligned} \bar{M} &= \mu M \\ &= e^x(y - e^x) \\ &= (y - e^x) e^x \end{aligned}$$

And

$$\begin{aligned} \bar{N} &= \mu N \\ &= e^x(1) \\ &= e^x \end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned} \bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ ((y - e^x) e^x) + (e^x) \frac{dy}{dx} &= 0 \end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \overline{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int (y - e^x) e^x dx \\ \phi &= -\frac{e^{2x}}{2} + y e^x + f(y)\end{aligned}\quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = e^x + f'(y)\quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = e^x$. Therefore equation (4) becomes

$$e^x = e^x + f'(y)\quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = -\frac{e^{2x}}{2} + y e^x + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\frac{e^{2x}}{2} + y e^x$$

The solution becomes

$$y = \frac{(e^{2x} + 2c_1) e^{-x}}{2}$$

Summary

The solution(s) found are the following

$$y = \frac{(e^{2x} + 2c_1) e^{-x}}{2} \quad (1)$$




Figure 32: Slope field plot

Verification of solutions

$$y = \frac{(e^{2x} + 2c_1) e^{-x}}{2}$$

Verified OK.

2.2.4 Maple step by step solution

Let's solve

$$y + y' = e^x$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = -y + e^x$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y + y' = e^x$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) (y + y') = \mu(x) e^x$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x) y)$

$$\mu(x) (y + y') = \mu'(x) y + \mu(x) y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = \mu(x)$$

- Solve to find the integrating factor

$$\mu(x) = e^x$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x) y) \right) dx = \int \mu(x) e^x dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x) y = \int \mu(x) e^x dx + c_1$$

- Solve for y

$$y = \frac{\int \mu(x) e^x dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = e^x$

$$y = \frac{\int (e^x)^2 dx + c_1}{e^x}$$

- Evaluate the integrals on the rhs

$$y = \frac{\frac{(e^x)^2}{2} + c_1}{e^x}$$

- Simplify

$$y = c_1 e^{-x} + \frac{e^x}{2}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(diff(y(x),x)+y(x)=exp(x),y(x), singsol=all)
```

$$y(x) = \frac{e^x}{2} + c_1 e^{-x}$$

✓ Solution by Mathematica

Time used: 0.04 (sec). Leaf size: 21

```
DSolve[y'[x]+y[x]==Exp[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^x}{2} + c_1 e^{-x}$$

2.3 problem 1(c)

2.3.1	Solving as linear ode	122
2.3.2	Solving as first order ode lie symmetry lookup ode	124
2.3.3	Solving as exact ode	128
2.3.4	Maple step by step solution	132

Internal problem ID [5925]

Internal file name [OUTPUT/5173_Sunday_June_05_2022_03_26_45_PM_39439636/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.6 Introduction– Linear equations of First Order. Page 41

Problem number: 1(c).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"linear", "exactWithIntegrationFactor", "first_order_ode_lie_symmetry_lookup"**

Maple gives the following as the ode type

```
[[_linear, `class A`]]
```

$$y' - 2y = x^2 + x$$

2.3.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -2$$

$$q(x) = x^2 + x$$

Hence the ode is

$$y' - 2y = x^2 + x$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int (-2) dx} \\ &= e^{-2x}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu)(x^2 + x) \\ \frac{d}{dx}(y e^{-2x}) &= (e^{-2x})(x^2 + x) \\ d(y e^{-2x}) &= (x(1 + x) e^{-2x}) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}y e^{-2x} &= \int x(1 + x) e^{-2x} dx \\ y e^{-2x} &= -\frac{(x^2 + 2x + 1) e^{-2x}}{2} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{-2x}$ results in

$$y = -\frac{e^{2x}(x^2 + 2x + 1) e^{-2x}}{2} + c_1 e^{2x}$$

which simplifies to

$$y = c_1 e^{2x} - \frac{(1 + x)^2}{2}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{2x} - \frac{(1 + x)^2}{2} \tag{1}$$




Figure 33: Slope field plot

Verification of solutions

$$y = c_1 e^{2x} - \frac{(1+x)^2}{2}$$

Verified OK.

2.3.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$\begin{aligned} y' &= x^2 + x + 2y \\ y' &= \omega(x, y) \end{aligned}$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 24: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= e^{2x}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^{2x}} dy \end{aligned}$$

Which results in

$$S = y e^{-2x}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = x^2 + x + 2y$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= -2y e^{-2x} \\ S_y &= e^{-2x} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = x(1 + x) e^{-2x} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = R(1 + R) e^{-2R}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by

integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = -\frac{(R^2 + 2R + 1)e^{-2R}}{2} + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$e^{-2x}y = -\frac{(x^2 + 2x + 1)e^{-2x}}{2} + c_1$$



Which simplifies to

$$e^{-2x}y = -\frac{(x^2 + 2x + 1)e^{-2x}}{2} + c_1$$

Which gives

$$y = -\frac{(x^2e^{-2x} + 2xe^{-2x} + e^{-2x} - 2c_1)e^{2x}}{2}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = x^2 + x + 2y$ 	$R = x$ $S = ye^{-2x}$	$\frac{dS}{dR} = R(1 + R)e^{-2R}$ 

Summary

The solution(s) found are the following

$$y = -\frac{(x^2e^{-2x} + 2xe^{-2x} + e^{-2x} - 2c_1)e^{2x}}{2} \quad (1)$$




Figure 34: Slope field plot

Verification of solutions

$$y = -\frac{(x^2 e^{-2x} + 2x e^{-2x} + e^{-2x} - 2c_1) e^{2x}}{2}$$

Verified OK.

2.3.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}dy &= (x^2 + x + 2y) dx \\ (-x^2 - x - 2y) dx + dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -x^2 - x - 2y \\ N(x, y) &= 1\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-x^2 - x - 2y) \\ &= -2\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(1) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= 1((-2) - (0)) \\ &= -2 \end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int A dx} \\ &= e^{\int -2 dx} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^{-2x} \\ &= e^{-2x} \end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} for now so not to confuse them with the original M and N .

$$\begin{aligned} \bar{M} &= \mu M \\ &= e^{-2x}(-x^2 - x - 2y) \\ &= -e^{-2x}(x^2 + x + 2y) \end{aligned}$$

And

$$\begin{aligned} \bar{N} &= \mu N \\ &= e^{-2x}(1) \\ &= e^{-2x} \end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned} \bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ (-e^{-2x}(x^2 + x + 2y)) + (e^{-2x}) \frac{dy}{dx} &= 0 \end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -e^{-2x}(x^2 + x + 2y) dx \\ \phi &= \frac{(x^2 + 2x + 2y + 1)e^{-2x}}{2} + f(y)\end{aligned}\quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = e^{-2x} + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = e^{-2x}$. Therefore equation (4) becomes

$$e^{-2x} = e^{-2x} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = \frac{(x^2 + 2x + 2y + 1)e^{-2x}}{2} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \frac{(x^2 + 2x + 2y + 1)e^{-2x}}{2}$$

The solution becomes

$$y = -\frac{(x^2 e^{-2x} + 2x e^{-2x} + e^{-2x} - 2c_1) e^{2x}}{2}$$

Summary

The solution(s) found are the following

$$y = -\frac{(x^2 e^{-2x} + 2x e^{-2x} + e^{-2x} - 2c_1) e^{2x}}{2} \quad (1)$$




Figure 35: Slope field plot

Verification of solutions

$$y = -\frac{(x^2 e^{-2x} + 2x e^{-2x} + e^{-2x} - 2c_1) e^{2x}}{2}$$

Verified OK.

2.3.4 Maple step by step solution

Let's solve

$$y' - 2y = x^2 + x$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = 2y + x^2 + x$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' - 2y = x^2 + x$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) (y' - 2y) = \mu(x) (x^2 + x)$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x) y)$

$$\mu(x) (y' - 2y) = \mu'(x) y + \mu(x) y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = -2\mu(x)$$

- Solve to find the integrating factor

$$\mu(x) = e^{-2x}$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x) y) \right) dx = \int \mu(x) (x^2 + x) dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x) y = \int \mu(x) (x^2 + x) dx + c_1$$

- Solve for y

$$y = \frac{\int \mu(x)(x^2+x)dx+c_1}{\mu(x)}$$

- Substitute $\mu(x) = e^{-2x}$

$$y = \frac{\int (x^2+x)e^{-2x} dx+c_1}{e^{-2x}}$$

- Evaluate the integrals on the rhs

$$y = \frac{-\frac{e^{-2x}(1+x)^2}{2}+c_1}{e^{-2x}}$$

- Simplify

$$y = c_1 e^{2x} - \frac{(1+x)^2}{2}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(diff(y(x),x)-2*y(x)=x^2+x,y(x), singsol=all)
```

$$y(x) = e^{2x}c_1 - \frac{(x+1)^2}{2}$$

✓ Solution by Mathematica

Time used: 0.073 (sec). Leaf size: 23

```
DSolve[y'[x]-2*y[x]==x^2+x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{2}(x+1)^2 + c_1e^{2x}$$

2.4 problem 1(d)

2.4.1	Solving as linear ode	135
2.4.2	Solving as first order ode lie symmetry lookup ode	137
2.4.3	Solving as exact ode	141
2.4.4	Maple step by step solution	146

Internal problem ID [5926]

Internal file name [OUTPUT/5174_Sunday_June_05_2022_03_26_47_PM_14465740/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.6 Introduction– Linear equations of First Order. Page 41

Problem number: 1(d).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"linear", "exactWithIntegrationFactor", "first_order_ode_lie_symmetry_lookup"**

Maple gives the following as the ode type

```
[[_linear, `class A`]]
```

$$y + 3y' = 2e^{-x}$$

2.4.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = \frac{1}{3}$$
$$q(x) = \frac{2e^{-x}}{3}$$

Hence the ode is

$$y' + \frac{y}{3} = \frac{2e^{-x}}{3}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int \frac{1}{3} dx} \\ &= e^{\frac{x}{3}}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu) \left(\frac{2e^{-x}}{3} \right) \\ \frac{d}{dx}(e^{\frac{x}{3}} y) &= (e^{\frac{x}{3}}) \left(\frac{2e^{-x}}{3} \right) \\ d(e^{\frac{x}{3}} y) &= \left(\frac{2e^{-\frac{2x}{3}}}{3} \right) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}e^{\frac{x}{3}} y &= \int \frac{2e^{-\frac{2x}{3}}}{3} dx \\ e^{\frac{x}{3}} y &= -e^{-\frac{2x}{3}} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{\frac{x}{3}}$ results in

$$y = -e^{-\frac{x}{3}} e^{-\frac{2x}{3}} + c_1 e^{-\frac{x}{3}}$$

which simplifies to

$$y = -e^{-x} + c_1 e^{-\frac{x}{3}}$$

Summary

The solution(s) found are the following

$$y = -e^{-x} + c_1 e^{-\frac{x}{3}} \tag{1}$$




Figure 36: Slope field plot

Verification of solutions

$$y = -e^{-x} + c_1 e^{-\frac{x}{3}}$$

Verified OK.

2.4.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = -\frac{y}{3} + \frac{2e^{-x}}{3}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 27: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= e^{-\frac{x}{3}}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^{-\frac{x}{3}}} dy \end{aligned}$$

Which results in

$$S = e^{\frac{x}{3}} y$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -\frac{y}{3} + \frac{2e^{-x}}{3}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{e^{\frac{x}{3}} y}{3} \\ S_y &= e^{\frac{x}{3}} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \frac{2e^{-\frac{2x}{3}}}{3} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \frac{2e^{-\frac{2R}{3}}}{3}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = -e^{-\frac{2R}{3}} + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$e^{\frac{x}{3}} y = -e^{-\frac{2x}{3}} + c_1$$



Which simplifies to

$$e^{\frac{x}{3}} y = -e^{-\frac{2x}{3}} + c_1$$

Which gives

$$y = -\left(e^{-\frac{2x}{3}} - c_1\right) e^{-\frac{x}{3}}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -\frac{y}{3} + \frac{2e^{-x}}{3}$ 	$R = x$ $S = e^{\frac{x}{3}} y$	$\frac{dS}{dR} = \frac{2e^{-\frac{2R}{3}}}{3}$ 

Summary

The solution(s) found are the following

$$y = -\left(e^{-\frac{2x}{3}} - c_1\right) e^{-\frac{x}{3}} \quad (1)$$




Figure 37: Slope field plot

Verification of solutions

$$y = -\left(e^{-\frac{2x}{3}} - c_1\right) e^{-\frac{x}{3}}$$

Verified OK.

2.4.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}(3) dy &= (-y + 2e^{-x}) dx \\ (y - 2e^{-x}) dx + (3) dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= y - 2e^{-x} \\ N(x, y) &= 3\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(y - 2e^{-x}) \\ &= 1\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(3) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= \frac{1}{3} ((1) - (0)) \\ &= \frac{1}{3} \end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int A dx} \\ &= e^{\int \frac{1}{3} dx} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^{\frac{x}{3}} \\ &= e^{\frac{x}{3}} \end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \overline{M} and \overline{N} for now so not to confuse them with the original M and N .

$$\begin{aligned} \overline{M} &= \mu M \\ &= e^{\frac{x}{3}} (y - 2e^{-x}) \\ &= (ye^x - 2) e^{-\frac{2x}{3}} \end{aligned}$$

And

$$\begin{aligned} \overline{N} &= \mu N \\ &= e^{\frac{x}{3}} (3) \\ &= 3e^{\frac{x}{3}} \end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned} \overline{M} + \overline{N} \frac{dy}{dx} &= 0 \\ \left((ye^x - 2) e^{-\frac{2x}{3}} \right) + (3e^{\frac{x}{3}}) \frac{dy}{dx} &= 0 \end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \overline{M} \quad (1)$$

$$\frac{\partial \phi}{\partial y} = \overline{N} \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned} \int \frac{\partial \phi}{\partial x} dx &= \int \overline{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int (y e^x - 2) e^{-\frac{2x}{3}} dx \\ \phi &= 3(y e^x + 1) e^{-\frac{2x}{3}} + f(y) \end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\begin{aligned} \frac{\partial \phi}{\partial y} &= 3 e^x e^{-\frac{2x}{3}} + f'(y) \\ &= 3 e^{\frac{x}{3}} + f'(y) \end{aligned} \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = 3 e^{\frac{x}{3}}$. Therefore equation (4) becomes

$$3 e^{\frac{x}{3}} = 3 e^{\frac{x}{3}} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = 3(y e^x + 1) e^{-\frac{2x}{3}} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = 3(y e^x + 1) e^{-\frac{2x}{3}}$$

The solution becomes

$$y = -\frac{\left(3e^{-\frac{2x}{3}} - c_1\right) e^{-x} e^{\frac{2x}{3}}}{3}$$

Summary

The solution(s) found are the following

$$y = -\frac{\left(3e^{-\frac{2x}{3}} - c_1\right) e^{-x} e^{\frac{2x}{3}}}{3} \quad (1)$$




Figure 38: Slope field plot

Verification of solutions

$$y = -\frac{\left(3e^{-\frac{2x}{3}} - c_1\right) e^{-x} e^{\frac{2x}{3}}}{3}$$

Verified OK.

2.4.4 Maple step by step solution

Let's solve

$$y + 3y' = 2e^{-x}$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = -\frac{y}{3} + \frac{2e^{-x}}{3}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' + \frac{y}{3} = \frac{2e^{-x}}{3}$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) \left(y' + \frac{y}{3} \right) = \frac{2\mu(x)e^{-x}}{3}$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x)y)$

$$\mu(x) \left(y' + \frac{y}{3} \right) = \mu'(x)y + \mu(x)y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = \frac{\mu(x)}{3}$$

- Solve to find the integrating factor

$$\mu(x) = e^{\frac{x}{3}}$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x)y) \right) dx = \int \frac{2\mu(x)e^{-x}}{3} dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x)y = \int \frac{2\mu(x)e^{-x}}{3} dx + c_1$$

- Solve for y

$$y = \frac{\int \frac{2\mu(x)e^{-x}}{3} dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = e^{\frac{x}{3}}$

$$y = \frac{\int \frac{2e^{-x}e^{\frac{x}{3}}}{3} dx + c_1}{e^{\frac{x}{3}}}$$

- Evaluate the integrals on the rhs

$$y = \frac{-e^{-\frac{2x}{3}} + c_1}{e^{\frac{x}{3}}}$$

- Simplify

$$y = \left(-e^{-\frac{2x}{3}} + c_1\right) e^{-\frac{x}{3}}$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(3*diff(y(x),x)+y(x)=2*exp(-x),y(x), singsol=all)
```

$$y(x) = -e^{-x} + e^{-\frac{x}{3}} c_1$$

✓ Solution by Mathematica

Time used: 0.051 (sec). Leaf size: 23

```
DSolve[3*y'[x]+y[x]==2*Exp[-x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x} (-1 + c_1 e^{2x/3})$$

2.5 problem 1(e)

2.5.1	Solving as linear ode	148
2.5.2	Solving as first order ode lie symmetry lookup ode	150
2.5.3	Solving as exact ode	154
2.5.4	Maple step by step solution	158

Internal problem ID [5927]

Internal file name [OUTPUT/5175_Sunday_June_05_2022_03_26_48_PM_51247357/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.6 Introduction– Linear equations of First Order. Page 41

Problem number: 1(e).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"linear", "exactWithIntegrationFactor", "first_order_ode_lie_symmetry_lookup"**

Maple gives the following as the ode type

```
[[_linear, `class A`]]
```

$$y' + 3y = e^{ix}$$

2.5.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = 3$$

$$q(x) = e^{ix}$$

Hence the ode is

$$y' + 3y = e^{ix}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int 3dx} \\ &= e^{3x}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu) (e^{ix}) \\ \frac{d}{dx}(e^{3x}y) &= (e^{3x}) (e^{ix}) \\ d(e^{3x}y) &= e^{(3+i)x} dx\end{aligned}$$

Integrating gives

$$\begin{aligned}e^{3x}y &= \int e^{(3+i)x} dx \\ e^{3x}y &= \left(\frac{3}{10} - \frac{i}{10}\right) e^{(3+i)x} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{3x}$ results in

$$y = \left(\frac{3}{10} - \frac{i}{10}\right) e^{-3x} e^{(3+i)x} + c_1 e^{-3x}$$

which simplifies to

$$y = \left(\frac{3}{10} - \frac{i}{10}\right) e^{ix} + c_1 e^{-3x}$$

Summary

The solution(s) found are the following

$$y = \left(\frac{3}{10} - \frac{i}{10}\right) e^{ix} + c_1 e^{-3x} \quad (1)$$




Figure 39: Slope field plot

Verification of solutions

$$y = \left(\frac{3}{10} - \frac{i}{10} \right) e^{ix} + c_1 e^{-3x}$$

Verified OK.

2.5.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$\begin{aligned} y' &= -3y + e^{ix} \\ y' &= \omega(x, y) \end{aligned}$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 30: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= e^{-3x}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^{-3x}} dy \end{aligned}$$

Which results in

$$S = e^{3x}y$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -3y + e^{ix}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= 3e^{3x}y \\ S_y &= e^{3x} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = e^{(3+i)x} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = e^{(3+i)R}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by

integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \left(\frac{3}{10} - \frac{i}{10} \right) e^{(3+i)R} + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$e^{3x}y = \left(\frac{3}{10} - \frac{i}{10} \right) e^{(3+i)x} + c_1$$



Which simplifies to

$$e^{3x}y = \left(\frac{3}{10} - \frac{i}{10} \right) e^{(3+i)x} + c_1$$

Which gives

$$y = \left(\frac{3}{10} - \frac{i}{10} \right) (ic_1 + e^{(3+i)x} + 3c_1) e^{-3x}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -3y + e^{ix}$ 	$R = x$ $S = e^{3x}y$	$\frac{dS}{dR} = e^{(3+i)R}$ 

Summary

The solution(s) found are the following

$$y = \left(\frac{3}{10} - \frac{i}{10} \right) (ic_1 + e^{(3+i)x} + 3c_1) e^{-3x} \quad (1)$$




Figure 40: Slope field plot

Verification of solutions

$$y = \left(\frac{3}{10} - \frac{i}{10} \right) (ic_1 + e^{(3+i)x} + 3c_1) e^{-3x}$$

Verified OK.

2.5.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned}dy &= (-3y + e^{ix}) dx \\ (3y - e^{ix}) dx + dy &= 0\end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= 3y - e^{ix} \\ N(x, y) &= 1\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(3y - e^{ix}) \\ &= 3\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(1) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= 1((3) - (0)) \\ &= 3 \end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int A dx} \\ &= e^{\int 3 dx} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^{3x} \\ &= e^{3x} \end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} for now so not to confuse them with the original M and N .

$$\begin{aligned} \bar{M} &= \mu M \\ &= e^{3x}(3y - e^{ix}) \\ &= (3y - e^{ix}) e^{3x} \end{aligned}$$

And

$$\begin{aligned} \bar{N} &= \mu N \\ &= e^{3x}(1) \\ &= e^{3x} \end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned} \bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ ((3y - e^{ix}) e^{3x}) + (e^{3x}) \frac{dy}{dx} &= 0 \end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int (3y - e^{ix}) e^{3x} dx \\ \phi &= \int^x (3y - e^{i-a}) e^{3-a} d_a + f(y)\end{aligned}\quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = e^{3x} + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = e^{3x}$. Therefore equation (4) becomes

$$e^{3x} = e^{3x} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = \int^x (3y - e^{i-a}) e^{3-a} d_a + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \int^x (3y - e^{i-a}) e^{3-a} d_a$$

Summary

The solution(s) found are the following

$$\int^x (3y - e^{i-a}) e^{3-a} d_a = c_1 \quad (1)$$




Figure 41: Slope field plot

Verification of solutions

$$\int^x (3y - e^{i-a}) e^{3-a} d_a = c_1$$

Verified OK.

2.5.4 Maple step by step solution

Let's solve

$$y' + 3y = e^{Ix}$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = -3y + e^{Ix}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' + 3y = e^{Ix}$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) (y' + 3y) = \mu(x) e^{Ix}$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x) y)$

$$\mu(x) (y' + 3y) = \mu'(x) y + \mu(x) y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = 3\mu(x)$$

- Solve to find the integrating factor

$$\mu(x) = e^{3x}$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x) y) \right) dx = \int \mu(x) e^{Ix} dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x) y = \int \mu(x) e^{Ix} dx + c_1$$

- Solve for y

$$y = \frac{\int \mu(x) e^{Ix} dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = e^{3x}$

$$y = \frac{\int e^{Ix} e^{3x} dx + c_1}{e^{3x}}$$

- Evaluate the integrals on the rhs

$$y = \frac{\left(\frac{3}{10} - \frac{1}{10} \right) e^{Ix+3x} + c_1}{e^{3x}}$$

- Simplify

$$y = -\frac{e^{-3x} ((-3+I)e^{(3+I)x} - 10c_1)}{10}$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```


✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
dsolve(diff(y(x),x)+3*y(x)=exp(I*x),y(x), singsol=all)
```

$$y(x) = -\frac{e^{-3x}((-3+i)e^{(3+i)x} - 10c_1)}{10}$$

✓ Solution by Mathematica

Time used: 0.054 (sec). Leaf size: 29

```
DSolve[y'[x]+3*y[x]==Exp[I*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \left(\frac{3}{10} - \frac{i}{10}\right) e^{ix} + c_1 e^{-3x}$$

2.6 problem 2

2.6.1	Solving as linear ode	161
2.6.2	Solving as first order ode lie symmetry lookup ode	163
2.6.3	Solving as exact ode	167
2.6.4	Maple step by step solution	171

Internal problem ID [5928]

Internal file name [OUTPUT/5176_Sunday_June_05_2022_03_26_49_PM_19687612/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.6 Introduction– Linear equations of First Order. Page 41

Problem number: 2.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"linear", "exactWithIntegrationFactor", "first_order_ode_lie_symmetry_lookup"**

Maple gives the following as the ode type

```
[[_linear, `class A`]]
```

$$y' + iy = x$$

2.6.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = i$$

$$q(x) = x$$

Hence the ode is

$$y' + iy = x$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int idx} \\ &= e^{ix}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu)(x) \\ \frac{d}{dx}(y e^{ix}) &= (e^{ix})(x) \\ d(y e^{ix}) &= (x e^{ix}) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}y e^{ix} &= \int x e^{ix} dx \\ y e^{ix} &= -(ix - 1) e^{ix} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{ix}$ results in

$$y = -e^{-ix}(ix - 1) e^{ix} + c_1 e^{-ix}$$

which simplifies to

$$y = -ix + 1 + c_1 e^{-ix}$$

Summary

The solution(s) found are the following

$$y = -ix + 1 + c_1 e^{-ix} \tag{1}$$




Figure 42: Slope field plot

Verification of solutions

$$y = -ix + 1 + c_1 e^{-ix}$$

Verified OK.

2.6.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = -iy + x$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \tag{A}$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 33: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= e^{-ix}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^{-ix}} dy \end{aligned}$$

Which results in

$$S = y e^{ix}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -iy + x$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= iy e^{ix} \\ S_y &= e^{ix} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = x e^{ix} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = R e^{iR}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by

integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = -(iR - 1) e^{iR} + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$e^{ix} y = -(ix - 1) e^{ix} + c_1$$



Which simplifies to

$$(ix + y - 1) e^{ix} - c_1 = 0$$

Which gives

$$y = -i(i e^{ix} + x e^{ix} + i c_1) e^{-ix}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -iy + x$ 	$R = x$ $S = y e^{ix}$	$\frac{dS}{dR} = R e^{iR}$ 

Summary

The solution(s) found are the following

$$y = -i(i e^{ix} + x e^{ix} + i c_1) e^{-ix} \quad (1)$$




Figure 43: Slope field plot

Verification of solutions

$$y = -i(i e^{ix} + x e^{ix} + ic_1) e^{-ix}$$

Verified OK.

2.6.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}dy &= (-iy + x) dx \\ (iy - x) dx + dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= iy - x \\ N(x, y) &= 1\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(iy - x) \\ &= i\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(1) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= 1((i) - (0)) \\ &= i \end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int A dx} \\ &= e^{\int i dx} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^{ix} \\ &= e^{ix} \end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} for now so not to confuse them with the original M and N .

$$\begin{aligned} \bar{M} &= \mu M \\ &= e^{ix}(iy - x) \\ &= (iy - x) e^{ix} \end{aligned}$$

And

$$\begin{aligned} \bar{N} &= \mu N \\ &= e^{ix}(1) \\ &= e^{ix} \end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned} \bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ ((iy - x) e^{ix}) + (e^{ix}) \frac{dy}{dx} &= 0 \end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int (iy - x) e^{ix} dx \\ \phi &= \int^x (iy - a) e^{i-a} d_a + f(y)\end{aligned}\quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = e^{ix} + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = e^{ix}$. Therefore equation (4) becomes

$$e^{ix} = e^{ix} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = \int^x (iy - a) e^{i-a} d_a + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \int^x (iy - a) e^{i-a} d_a$$

Summary

The solution(s) found are the following

$$\int^x (iy - a) e^{i-a} d_a = c_1 \quad (1)$$




Figure 44: Slope field plot

Verification of solutions

$$\int^x (iy - a) e^{i-a} d_a = c_1$$

Verified OK.

2.6.4 Maple step by step solution

Let's solve

$$y' + Iy = x$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = -Iy + x$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' + Iy = x$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) (y' + Iy) = \mu(x) x$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x) y)$

$$\mu(x) (y' + Iy) = \mu'(x) y + \mu(x) y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = I\mu(x)$$

- Solve to find the integrating factor

$$\mu(x) = e^{Ix}$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x) y) \right) dx = \int \mu(x) x dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x) y = \int \mu(x) x dx + c_1$$

- Solve for y

$$y = \frac{\int \mu(x) x dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = e^{Ix}$

$$y = \frac{\int x e^{Ix} dx + c_1}{e^{Ix}}$$

- Evaluate the integrals on the rhs

$$y = \frac{-(Ix-1)e^{Ix} + c_1}{e^{Ix}}$$

- Simplify

$$y = -Ix + 1 + c_1 e^{-Ix}$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x)+I*y(x)=x,y(x), singsol=all)
```

$$y(x) = -ix + 1 + e^{-ix}c_1$$

✓ Solution by Mathematica

Time used: 0.028 (sec). Leaf size: 22

```
DSolve[y'[x]+I*y[x]==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -ix + c_1e^{-ix} + 1$$

2.7 problem 3

2.7.1 Solving as quadrature ode	174
2.7.2 Maple step by step solution	175

Internal problem ID [5929]

Internal file name [OUTPUT/5177_Sunday_June_05_2022_03_26_50_PM_84062245/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.6 Introduction– Linear equations of First Order. Page 41

Problem number: 3.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"quadrature"**

Maple gives the following as the ode type

`[_quadrature]`

$$Ly' + Ry = E$$

2.7.1 Solving as quadrature ode

Integrating both sides gives

$$\int \frac{L}{-Ry + E} dy = \int dx$$
$$-\frac{L \ln(-Ry + E)}{R} = x + c_1$$

Raising both side to exponential gives

$$e^{-\frac{L \ln(-Ry + E)}{R}} = e^{x+c_1}$$

Which simplifies to

$$(-Ry + E)^{-\frac{L}{R}} = c_2 e^x$$

Summary

The solution(s) found are the following

$$y = -\frac{(c_2 e^x)^{-\frac{R}{L}} - E}{R} \tag{1}$$

Verification of solutions

$$y = -\frac{(c_2 e^x)^{-\frac{R}{L}} - E}{R}$$

Verified OK.

2.7.2 Maple step by step solution

Let's solve

$$Ly' + Ry = E$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{-Ry+E} = \frac{1}{L}$$

- Integrate both sides with respect to x

$$\int \frac{y'}{-Ry+E} dx = \int \frac{1}{L} dx + c_1$$

- Evaluate integral

$$-\frac{\ln(-Ry+E)}{R} = \frac{x}{L} + c_1$$

- Solve for y

$$y = \frac{-e^{-\frac{R(Lc_1+x)}{L}} + E}{R}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```


✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(L*diff(y(x),x)+R*y(x)=E,y(x), singsol=all)
```

$$y(x) = \frac{e^{-\frac{Rx}{L}} c_1 R + E}{R}$$

✓ Solution by Mathematica

Time used: 0.056 (sec). Leaf size: 23

```
DSolve[L*y'[x]+R*y[x]==E0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{E0 - E0e^{-\frac{Rx}{L}}}{R}$$

2.8 problem 4

2.8.1	Existence and uniqueness analysis	177
2.8.2	Solving as linear ode	178
2.8.3	Solving as first order ode lie symmetry lookup ode	179
2.8.4	Solving as exact ode	183
2.8.5	Maple step by step solution	187

Internal problem ID [5930]

Internal file name [OUTPUT/5178_Sunday_June_05_2022_03_26_51_PM_53986323/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.6 Introduction– Linear equations of First Order. Page 41

Problem number: 4.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"linear", "exactWithIntegrationFactor", "first_order_ode_lie_symmetry_lookup"**

Maple gives the following as the ode type

```
[[_linear, `class A`]]
```

$$Ly' + Ry = E \sin(\omega x)$$

With initial conditions

$$[y(0) = 0]$$

2.8.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = \frac{R}{L}$$
$$q(x) = \frac{E \sin(\omega x)}{L}$$

Hence the ode is

$$y' + \frac{Ry}{L} = \frac{E \sin(\omega x)}{L}$$

The domain of $p(x) = \frac{R}{L}$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = \frac{E \sin(\omega x)}{L}$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

2.8.2 Solving as linear ode

Entering Linear first order ODE solver. The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int \frac{R}{L} dx} \\ &= e^{\frac{Rx}{L}} \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx}(\mu y) &= (\mu) \left(\frac{E \sin(\omega x)}{L} \right) \\ \frac{d}{dx} \left(e^{\frac{Rx}{L}} y \right) &= \left(e^{\frac{Rx}{L}} \right) \left(\frac{E \sin(\omega x)}{L} \right) \\ d \left(e^{\frac{Rx}{L}} y \right) &= \left(\frac{E \sin(\omega x) e^{\frac{Rx}{L}}}{L} \right) dx \end{aligned}$$

Integrating gives

$$\begin{aligned} e^{\frac{Rx}{L}} y &= \int \frac{E \sin(\omega x) e^{\frac{Rx}{L}}}{L} dx \\ e^{\frac{Rx}{L}} y &= \frac{E \left(-\frac{\omega e^{\frac{Rx}{L}} \cos(\omega x)}{\frac{R^2}{L^2} + \omega^2} + \frac{R e^{\frac{Rx}{L}} \sin(\omega x)}{L(\frac{R^2}{L^2} + \omega^2)} \right)}{L} + c_1 \end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{\frac{Rx}{L}}$ results in

$$y = \frac{e^{-\frac{Rx}{L}} E \left(-\frac{\omega e^{\frac{Rx}{L}} \cos(\omega x)}{\frac{R^2}{L^2} + \omega^2} + \frac{R e^{\frac{Rx}{L}} \sin(\omega x)}{L(\frac{R^2}{L^2} + \omega^2)} \right)}{L} + c_1 e^{-\frac{Rx}{L}}$$

which simplifies to

$$y = \frac{c_1(\omega^2 L^2 + R^2) e^{-\frac{Rx}{L}} - E(L \cos(\omega x) \omega - \sin(\omega x) R)}{\omega^2 L^2 + R^2}$$

Initial conditions are used to solve for c_1 . Substituting $x = 0$ and $y = 0$ in the above solution gives an equation to solve for the constant of integration.

$$0 = \frac{L^2 c_1 \omega^2 - EL\omega + R^2 c_1}{\omega^2 L^2 + R^2}$$

$$c_1 = \frac{EL\omega}{\omega^2 L^2 + R^2}$$

Substituting c_1 found above in the general solution gives

$$y = \frac{-EL\omega \cos(\omega x) + EL e^{-\frac{Rx}{L}} \omega + ER \sin(\omega x)}{\omega^2 L^2 + R^2}$$

Summary

The solution(s) found are the following

$$y = \frac{-EL\omega \cos(\omega x) + EL e^{-\frac{Rx}{L}} \omega + ER \sin(\omega x)}{\omega^2 L^2 + R^2} \quad (1)$$

Verification of solutions

$$y = \frac{-EL\omega \cos(\omega x) + EL e^{-\frac{Rx}{L}} \omega + ER \sin(\omega x)}{\omega^2 L^2 + R^2}$$

Verified OK.

2.8.3 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{-Ry + E \sin(\omega x)}{L}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (A)$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 37: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= e^{-\frac{Rx}{L}}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^{-\frac{Rx}{L}}} dy \end{aligned}$$

Which results in

$$S = e^{\frac{Rx}{L}} y$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{-Ry + E \sin(\omega x)}{L}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{R e^{\frac{Rx}{L}} y}{L} \\ S_y &= e^{\frac{Rx}{L}} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \frac{E \sin(\omega x) e^{\frac{Rx}{L}}}{L} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \frac{E \sin(\omega R) e^{\frac{RR}{L}}}{L}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = -\frac{-c_1(\omega^2 L^2 + R^2) + E e^{\frac{Rr}{L}} (L \cos(\omega R) \omega - \sin(\omega R) R)}{\omega^2 L^2 + R^2} \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$e^{\frac{Rx}{L}} y = -\frac{-c_1(\omega^2 L^2 + R^2) + E e^{\frac{Rx}{L}} (L \cos(\omega x) \omega - \sin(\omega x) R)}{\omega^2 L^2 + R^2}$$

Which simplifies to

$$e^{\frac{Rx}{L}} y = -\frac{-c_1(\omega^2 L^2 + R^2) + E e^{\frac{Rx}{L}} (L \cos(\omega x) \omega - \sin(\omega x) R)}{\omega^2 L^2 + R^2}$$

Which gives

$$y = -\frac{e^{-\frac{Rx}{L}} \left(E \omega \cos(\omega x) e^{\frac{Rx}{L}} L - L^2 c_1 \omega^2 - E \sin(\omega x) R e^{\frac{Rx}{L}} - R^2 c_1 \right)}{\omega^2 L^2 + R^2}$$

Initial conditions are used to solve for c_1 . Substituting $x = 0$ and $y = 0$ in the above solution gives an equation to solve for the constant of integration.

$$0 = \frac{L^2 c_1 \omega^2 - EL\omega + R^2 c_1}{\omega^2 L^2 + R^2}$$

$$c_1 = \frac{EL\omega}{\omega^2 L^2 + R^2}$$

Substituting c_1 found above in the general solution gives

$$y = \frac{-EL \cos(\omega x) e^{\frac{Rx}{L}} e^{-\frac{Rx}{L}} \omega + E \sin(\omega x) e^{\frac{Rx}{L}} e^{-\frac{Rx}{L}} R + EL e^{-\frac{Rx}{L}} \omega}{\omega^2 L^2 + R^2}$$

Summary

The solution(s) found are the following

$$y = \frac{-EL \cos(\omega x) e^{\frac{Rx}{L}} e^{-\frac{Rx}{L}} \omega + E \sin(\omega x) e^{\frac{Rx}{L}} e^{-\frac{Rx}{L}} R + EL e^{-\frac{Rx}{L}} \omega}{\omega^2 L^2 + R^2} \quad (1)$$

Verification of solutions

$$y = \frac{-EL \cos(\omega x) e^{\frac{Rx}{L}} e^{-\frac{Rx}{L}} \omega + E \sin(\omega x) e^{\frac{Rx}{L}} e^{-\frac{Rx}{L}} R + EL e^{-\frac{Rx}{L}} \omega}{\omega^2 L^2 + R^2}$$

Verified OK.

2.8.4 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} (L) dy &= (-Ry + E \sin(\omega x)) dx \\ (Ry - E \sin(\omega x)) dx + (L) dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= Ry - E \sin(\omega x) \\ N(x, y) &= L \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(Ry - E \sin(\omega x)) \\ &= R\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(L) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned}A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= \frac{1}{L} ((R) - (0)) \\ &= \frac{R}{L}\end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int A dx} \\ &= e^{\int \frac{R}{L} dx}\end{aligned}$$

The result of integrating gives

$$\begin{aligned}\mu &= e^{\frac{Rx}{L}} \\ &= e^{\frac{Rx}{L}}\end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} for now so not to confuse them with the original M and N .

$$\begin{aligned}\bar{M} &= \mu M \\ &= e^{\frac{Rx}{L}} (Ry - E \sin(\omega x)) \\ &= -e^{\frac{Rx}{L}} (-Ry + E \sin(\omega x))\end{aligned}$$

And

$$\begin{aligned}\bar{N} &= \mu N \\ &= e^{\frac{Rx}{L}}(L) \\ &= L e^{\frac{Rx}{L}}\end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned}\bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ \left(-e^{\frac{Rx}{L}}(-Ry + E \sin(\omega x))\right) + \left(L e^{\frac{Rx}{L}}\right) \frac{dy}{dx} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \quad (1)$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -e^{\frac{Rx}{L}}(-Ry + E \sin(\omega x)) dx \\ \phi &= \frac{(EL\omega \cos(\omega x) - ER \sin(\omega x) + y(\omega^2 L^2 + R^2)) e^{\frac{Rx}{L}} L}{\omega^2 L^2 + R^2} + f(y) \quad (3)\end{aligned}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = L e^{\frac{Rx}{L}} + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = L e^{\frac{Rx}{L}}$. Therefore equation (4) becomes

$$L e^{\frac{Rx}{L}} = L e^{\frac{Rx}{L}} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = \frac{(EL\omega \cos(\omega x) - ER \sin(\omega x) + y(\omega^2 L^2 + R^2)) e^{\frac{Rx}{L}} L}{\omega^2 L^2 + R^2} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \frac{(EL\omega \cos(\omega x) - ER \sin(\omega x) + y(\omega^2 L^2 + R^2)) e^{\frac{Rx}{L}} L}{\omega^2 L^2 + R^2}$$

The solution becomes

$$y = -\frac{(E L^2 \cos(\omega x) e^{\frac{Rx}{L}} \omega - EL \sin(\omega x) e^{\frac{Rx}{L}} R - L^2 c_1 \omega^2 - R^2 c_1) e^{-\frac{Rx}{L}}}{(\omega^2 L^2 + R^2) L}$$

Initial conditions are used to solve for c_1 . Substituting $x = 0$ and $y = 0$ in the above solution gives an equation to solve for the constant of integration.

$$0 = \frac{L^2 c_1 \omega^2 - E L^2 \omega + R^2 c_1}{L^3 \omega^2 + L R^2}$$

$$c_1 = \frac{E L^2 \omega}{\omega^2 L^2 + R^2}$$

Substituting c_1 found above in the general solution gives

$$y = \frac{-EL \cos(\omega x) e^{\frac{Rx}{L}} e^{-\frac{Rx}{L}} \omega + E \sin(\omega x) e^{\frac{Rx}{L}} e^{-\frac{Rx}{L}} R + EL e^{-\frac{Rx}{L}} \omega}{\omega^2 L^2 + R^2}$$

Summary

The solution(s) found are the following

$$y = \frac{-EL \cos(\omega x) e^{\frac{Rx}{L}} e^{-\frac{Rx}{L}} \omega + E \sin(\omega x) e^{\frac{Rx}{L}} e^{-\frac{Rx}{L}} R + EL e^{-\frac{Rx}{L}} \omega}{\omega^2 L^2 + R^2} \quad (1)$$

Verification of solutions

$$y = \frac{-EL \cos(\omega x) e^{\frac{Rx}{L}} e^{-\frac{Rx}{L}} \omega + E \sin(\omega x) e^{\frac{Rx}{L}} e^{-\frac{Rx}{L}} R + EL e^{-\frac{Rx}{L}} \omega}{\omega^2 L^2 + R^2}$$

Verified OK.

2.8.5 Maple step by step solution

Let's solve

$$[Ly' + Ry = E \sin(\omega x), y(0) = 0]$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = -\frac{Ry}{L} + \frac{E \sin(\omega x)}{L}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' + \frac{Ry}{L} = \frac{E \sin(\omega x)}{L}$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) \left(y' + \frac{Ry}{L} \right) = \frac{\mu(x)E \sin(\omega x)}{L}$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x)y)$

$$\mu(x) \left(y' + \frac{Ry}{L} \right) = \mu'(x)y + \mu(x)y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = \frac{\mu(x)R}{L}$$

- Solve to find the integrating factor

$$\mu(x) = e^{\frac{Rx}{L}}$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x)y) \right) dx = \int \frac{\mu(x)E \sin(\omega x)}{L} dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x)y = \int \frac{\mu(x)E \sin(\omega x)}{L} dx + c_1$$

- Solve for y

$$y = \frac{\int \frac{\mu(x)E \sin(\omega x)}{L} dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = e^{\frac{Rx}{L}}$

$$y = \frac{\int \frac{E \sin(\omega x) e^{\frac{Rx}{L}}}{L} dx + c_1}{e^{\frac{Rx}{L}}}$$

- Evaluate the integrals on the rhs

$$y = \frac{E \left(-\omega e^{-\frac{Rx}{L}} \frac{\cos(\omega x)}{L^2 + \omega^2} + R e^{-\frac{Rx}{L}} \frac{\sin(\omega x)}{L(L^2 + \omega^2)} \right)}{e^{-\frac{Rx}{L}}} + c_1$$

- Simplify

$$y = \frac{c_1(\omega^2 L^2 + R^2) e^{-\frac{Rx}{L}} - E(L \cos(\omega x) \omega - \sin(\omega x) R)}{\omega^2 L^2 + R^2}$$

- Use initial condition $y(0) = 0$

$$0 = \frac{c_1(\omega^2 L^2 + R^2) - EL\omega}{\omega^2 L^2 + R^2}$$

- Solve for c_1

$$c_1 = \frac{EL\omega}{\omega^2 L^2 + R^2}$$

- Substitute $c_1 = \frac{EL\omega}{\omega^2 L^2 + R^2}$ into general solution and simplify

$$y = -\frac{E \left(L \cos(\omega x) \omega - L e^{-\frac{Rx}{L}} \omega - \sin(\omega x) R \right)}{\omega^2 L^2 + R^2}$$

- Solution to the IVP

$$y = -\frac{E \left(L \cos(\omega x) \omega - L e^{-\frac{Rx}{L}} \omega - \sin(\omega x) R \right)}{\omega^2 L^2 + R^2}$$

Maple trace

```
`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 45

```
dsolve([L*dif(y(x),x)+R*y(x)=E*sin(omega*x),y(0) = 0],y(x), singsol=all)
```

$$y(x) = \frac{E \left(e^{-\frac{Rx}{L}} L \omega - L \cos(\omega x) \omega + \sin(\omega x) R \right)}{\omega^2 L^2 + R^2}$$

✓ Solution by Mathematica

Time used: 0.115 (sec). Leaf size: 47

```
DSolve[{L*y'[x]+R*y[x]==E0*Sin[\[Omega]*x],{y[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{E_0 \left(L \omega e^{-\frac{Rx}{L}} - L \omega \cos(x\omega) + R \sin(x\omega) \right)}{L^2 \omega^2 + R^2}$$

2.9 problem 5

2.9.1	Existence and uniqueness analysis	190
2.9.2	Solving as linear ode	191
2.9.3	Solving as first order ode lie symmetry lookup ode	192
2.9.4	Solving as exact ode	196
2.9.5	Maple step by step solution	200

Internal problem ID [5931]

Internal file name [OUTPUT/5179_Sunday_June_05_2022_03_26_53_PM_63615461/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.6 Introduction– Linear equations of First Order. Page 41

Problem number: 5.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**linear**", "**exactWithIntegrationFactor**", "**first_order_ode_lie_symmetry_lookup**"

Maple gives the following as the ode type

```
[[_linear, `class A`]]
```

$$Ly' + Ry = E e^{i\omega x}$$

With initial conditions

$$[y(0) = 0]$$

2.9.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = \frac{R}{L}$$
$$q(x) = \frac{E e^{i\omega x}}{L}$$

Hence the ode is

$$y' + \frac{Ry}{L} = \frac{E e^{i\omega x}}{L}$$

The domain of $p(x) = \frac{R}{L}$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = \frac{E e^{i\omega x}}{L}$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

2.9.2 Solving as linear ode

Entering Linear first order ODE solver. The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int \frac{R}{L} dx} \\ &= e^{\frac{Rx}{L}} \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx}(\mu y) &= (\mu) \left(\frac{E e^{i\omega x}}{L} \right) \\ \frac{d}{dx} \left(e^{\frac{Rx}{L}} y \right) &= \left(e^{\frac{Rx}{L}} \right) \left(\frac{E e^{i\omega x}}{L} \right) \\ d \left(e^{\frac{Rx}{L}} y \right) &= \left(\frac{E e^{\frac{x(iL\omega + R)}{L}}}{L} \right) dx \end{aligned}$$

Integrating gives

$$\begin{aligned} e^{\frac{Rx}{L}} y &= \int \frac{E e^{\frac{x(iL\omega + R)}{L}}}{L} dx \\ e^{\frac{Rx}{L}} y &= \frac{E e^{\frac{x(iL\omega + R)}{L}}}{iL\omega + R} + c_1 \end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{\frac{Rx}{L}}$ results in

$$y = \frac{e^{-\frac{Rx}{L}} E e^{\frac{x(iL\omega + R)}{L}}}{iL\omega + R} + c_1 e^{-\frac{Rx}{L}}$$

which simplifies to

$$y = \frac{e^{-\frac{Rx}{L}} \left(E e^{\frac{x(iL\omega+R)}{L}} + (iL\omega + R) c_1 \right)}{iL\omega + R}$$

Initial conditions are used to solve for c_1 . Substituting $x = 0$ and $y = 0$ in the above solution gives an equation to solve for the constant of integration.

$$0 = \frac{Lc_1\omega i + Rc_1 + E}{iL\omega + R}$$

$$c_1 = -\frac{E}{iL\omega + R}$$

Substituting c_1 found above in the general solution gives

$$y = \frac{e^{-\frac{Rx}{L}} E e^{\frac{x(iL\omega+R)}{L}} - E e^{-\frac{Rx}{L}}}{iL\omega + R}$$

Summary

The solution(s) found are the following

$$y = \frac{e^{-\frac{Rx}{L}} E e^{\frac{x(iL\omega+R)}{L}} - E e^{-\frac{Rx}{L}}}{iL\omega + R} \quad (1)$$

Verification of solutions

$$y = \frac{e^{-\frac{Rx}{L}} E e^{\frac{x(iL\omega+R)}{L}} - E e^{-\frac{Rx}{L}}}{iL\omega + R}$$

Verified OK.

2.9.3 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{-Ry + E e^{i\omega x}}{L}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (A)$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 40: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= e^{-\frac{Rx}{L}}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right)S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^{-\frac{Rx}{L}}} dy \end{aligned}$$

Which results in

$$S = e^{\frac{Rx}{L}} y$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{-Ry + E e^{i\omega x}}{L}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{R e^{\frac{Rx}{L}} y}{L} \\ S_y &= e^{\frac{Rx}{L}} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \frac{E e^{\frac{x(iL\omega+R)}{L}}}{L} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \frac{E e^{\frac{R(iL\omega+R)}{L}}}{L}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \frac{E e^{\frac{R(iL\omega+R)}{L}}}{iL\omega + R} + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$e^{\frac{Rx}{L}} y = \frac{E e^{\frac{x(iL\omega+R)}{L}}}{iL\omega + R} + c_1$$

Which simplifies to

$$e^{\frac{Rx}{L}} y = \frac{E e^{\frac{x(iL\omega+R)}{L}}}{iL\omega + R} + c_1$$

Which gives

$$y = \frac{e^{-\frac{Rx}{L}} \left(Lc_1\omega i + E e^{\frac{x(iL\omega+R)}{L}} + Rc_1 \right)}{iL\omega + R}$$

Initial conditions are used to solve for c_1 . Substituting $x = 0$ and $y = 0$ in the above solution gives an equation to solve for the constant of integration.

$$0 = \frac{Lc_1\omega i + Rc_1 + E}{iL\omega + R}$$

$$c_1 = -\frac{E}{iL\omega + R}$$

Substituting c_1 found above in the general solution gives

$$y = \frac{e^{-\frac{Rx}{L}} E e^{\frac{x(iL\omega+R)}{L}} - E e^{-\frac{Rx}{L}}}{iL\omega + R}$$

Summary

The solution(s) found are the following

$$y = \frac{e^{-\frac{Rx}{L}} E e^{\frac{x(iL\omega+R)}{L}} - E e^{-\frac{Rx}{L}}}{iL\omega + R} \quad (1)$$

Verification of solutions

$$y = \frac{e^{-\frac{Rx}{L}} E e^{\frac{x(iL\omega+R)}{L}} - E e^{-\frac{Rx}{L}}}{iL\omega + R}$$

Verified OK.

2.9.4 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} (L) dy &= (-Ry + E e^{i\omega x}) dx \\ (Ry - E e^{i\omega x}) dx + (L) dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= Ry - E e^{i\omega x} \\ N(x, y) &= L \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(Ry - E e^{i\omega x}) \\ &= R\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(L) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned}A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= \frac{1}{L} ((R) - (0)) \\ &= \frac{R}{L}\end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int A dx} \\ &= e^{\int \frac{R}{L} dx}\end{aligned}$$

The result of integrating gives

$$\begin{aligned}\mu &= e^{\frac{Rx}{L}} \\ &= e^{\frac{Rx}{L}}\end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \overline{M} and \overline{N} for now so not to confuse them with the original M and N .

$$\begin{aligned}\overline{M} &= \mu M \\ &= e^{\frac{Rx}{L}} (Ry - E e^{i\omega x}) \\ &= -e^{\frac{Rx}{L}} (-Ry + E e^{i\omega x})\end{aligned}$$

And

$$\begin{aligned}\bar{N} &= \mu N \\ &= e^{\frac{Rx}{L}} (L) \\ &= L e^{\frac{Rx}{L}}\end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned}\bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ \left(-e^{\frac{Rx}{L}} (-Ry + E e^{i\omega x})\right) + \left(L e^{\frac{Rx}{L}}\right) \frac{dy}{dx} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \quad (1)$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -e^{\frac{Rx}{L}} (-Ry + E e^{i\omega x}) dx \\ \phi &= \int_0^x -e^{\frac{R}{L}a} (-Ry + E e^{i\omega -a}) d_a + f(y)\end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = \int_0^x e^{\frac{R}{L}a} R d_a + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = L e^{\frac{Rx}{L}}$. Therefore equation (4) becomes

$$L e^{\frac{Rx}{L}} = R \left(\int_0^x e^{\frac{R}{L}a} d_a \right) + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = L e^{\frac{Rx}{L}} - R \left(\int_0^x e^{\frac{R-a}{L} d_{-a}} \right)$$

Integrating the above w.r.t y gives

$$\int f'(y) dy = \int \left(L e^{\frac{Rx}{L}} - R \left(\int_0^x e^{\frac{R-a}{L} d_{-a}} \right) \right) dy$$

$$f(y) = \left(L e^{\frac{Rx}{L}} - R \left(\int_0^x e^{\frac{R-a}{L} d_{-a}} \right) \right) y + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = \int_0^x -e^{\frac{R-a}{L}} (-Ry + E e^{i\omega-a}) d_{-a} + \left(L e^{\frac{Rx}{L}} - R \left(\int_0^x e^{\frac{R-a}{L} d_{-a}} \right) \right) y + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \int_0^x -e^{\frac{R-a}{L}} (-Ry + E e^{i\omega-a}) d_{-a} + \left(L e^{\frac{Rx}{L}} - R \left(\int_0^x e^{\frac{R-a}{L} d_{-a}} \right) \right) y$$

The solution becomes

$$y = \frac{\left(E \left(\int_0^x e^{\frac{R-a}{L}} e^{i\omega-a} d_{-a} \right) + c_1 \right) e^{-\frac{Rx}{L}}}{L}$$

Initial conditions are used to solve for c_1 . Substituting $x = 0$ and $y = 0$ in the above solution gives an equation to solve for the constant of integration.

$$0 = \frac{c_1}{L}$$

$$c_1 = 0$$

Substituting c_1 found above in the general solution gives

$$y = \frac{E \left(\int_0^x e^{\frac{-a(iL\omega+R)}{L}} d_{-a} \right) e^{-\frac{Rx}{L}}}{L}$$

Summary

The solution(s) found are the following

$$y = \frac{E \left(\int_0^x e^{-\frac{a(iL\omega+R)}{L}} d_a \right) e^{-\frac{Rx}{L}}}{L} \quad (1)$$

Verification of solutions

$$y = \frac{E \left(\int_0^x e^{-\frac{a(iL\omega+R)}{L}} d_a \right) e^{-\frac{Rx}{L}}}{L}$$

Verified OK.

2.9.5 Maple step by step solution

Let's solve

$$[Ly' + Ry = E e^{i\omega x}, y(0) = 0]$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = -\frac{Ry}{L} + \frac{E e^{i\omega x}}{L}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' + \frac{Ry}{L} = \frac{E e^{i\omega x}}{L}$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) \left(y' + \frac{Ry}{L} \right) = \frac{\mu(x) E e^{i\omega x}}{L}$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x) y)$

$$\mu(x) \left(y' + \frac{Ry}{L} \right) = \mu'(x) y + \mu(x) y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = \frac{\mu(x) R}{L}$$

- Solve to find the integrating factor

$$\mu(x) = e^{\frac{Rx}{L}}$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x) y) \right) dx = \int \frac{\mu(x) E e^{i\omega x}}{L} dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x)y = \int \frac{\mu(x)Ee^{I\omega x}}{L} dx + c_1$$

- Solve for y

$$y = \frac{\int \frac{\mu(x)Ee^{I\omega x}}{L} dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = e^{\frac{Rx}{L}}$

$$y = \frac{\int \frac{Ee^{I\omega x}e^{\frac{Rx}{L}}}{L} dx + c_1}{e^{\frac{Rx}{L}}}$$

- Evaluate the integrals on the rhs

$$y = \frac{\frac{Ee^{I\omega x + \frac{Rx}{L}}}{IL\omega + R} + c_1}{e^{\frac{Rx}{L}}}$$

- Simplify

$$y = \frac{e^{-\frac{Rx}{L}} \left(Ee^{\frac{x(IL\omega + R)}{L}} + (IL\omega + R)c_1 \right)}{IL\omega + R}$$

- Use initial condition $y(0) = 0$

$$0 = \frac{E + (IL\omega + R)c_1}{IL\omega + R}$$

- Solve for c_1

$$c_1 = -\frac{E}{IL\omega + R}$$

- Substitute $c_1 = -\frac{E}{IL\omega + R}$ into general solution and simplify

$$y = \frac{E \left(e^{\frac{x(IL\omega + R)}{L}} - 1 \right) e^{-\frac{Rx}{L}}}{IL\omega + R}$$

- Solution to the IVP

$$y = \frac{E \left(e^{\frac{x(IL\omega + R)}{L}} - 1 \right) e^{-\frac{Rx}{L}}}{IL\omega + R}$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 38

```
dsolve([L*diff(y(x),x)+R*y(x)=E*exp(I*omega*x),y(0) = 0],y(x), singsol=all)
```

$$y(x) = \frac{E \left(e^{\frac{x(iL\omega + R)}{L}} - 1 \right) e^{-\frac{Rx}{L}}}{iL\omega + R}$$

✓ Solution by Mathematica

Time used: 0.101 (sec). Leaf size: 43

```
DSolve[{L*y'[x]+R*y[x]==E0*Exp[I*\[Omega]*x],{y[0]==0}},y[x],x,IncludeSingularSolutions -> T
```

$$y(x) \rightarrow \frac{E0 e^{-\frac{Rx}{L}} \left(-1 + e^{\frac{x(R+iL\omega)}{L}} \right)}{R + iL\omega}$$

2.10 problem 7

2.10.1 Solving as linear ode	203
2.10.2 Solving as first order ode lie symmetry lookup ode	205
2.10.3 Solving as exact ode	208
2.10.4 Maple step by step solution	211

Internal problem ID [5932]

Internal file name [OUTPUT/5180_Sunday_June_05_2022_03_26_54_PM_11645007/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1.6 Introduction– Linear equations of First Order. Page 41

Problem number: 7.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"linear", "exactWithIntegrationFactor", "first_order_ode_lie_symmetry_lookup"**

Maple gives the following as the ode type

```
[[_linear, `class A`]]
```

$$y' + ya = b(x)$$

2.10.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = a$$

$$q(x) = b(x)$$

Hence the ode is

$$y' + ya = b(x)$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int adx} \\ &= e^{ax}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu)(b(x)) \\ \frac{d}{dx}(e^{ax}y) &= (e^{ax})(b(x)) \\ d(e^{ax}y) &= (b(x)e^{ax}) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}e^{ax}y &= \int b(x)e^{ax} dx \\ e^{ax}y &= \int b(x)e^{ax} dx + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{ax}$ results in

$$y = e^{-ax} \left(\int b(x)e^{ax} dx \right) + c_1 e^{-ax}$$

which simplifies to

$$y = e^{-ax} \left(\int b(x)e^{ax} dx + c_1 \right)$$

Summary

The solution(s) found are the following

$$y = e^{-ax} \left(\int b(x)e^{ax} dx + c_1 \right) \tag{1}$$

Verification of solutions

$$y = e^{-ax} \left(\int b(x)e^{ax} dx + c_1 \right)$$

Verified OK.

2.10.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = -ya + b(x)$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 43: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= e^{-ax}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^{-ax}} dy\end{aligned}$$

Which results in

$$S = e^{ax}y$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y}\tag{2}$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -ya + b(x)$$

Evaluating all the partial derivatives gives

$$\begin{aligned}R_x &= 1 \\ R_y &= 0 \\ S_x &= a e^{ax}y \\ S_y &= e^{ax}\end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = b(x) e^{ax} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = b(R) e^{aR}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \int b(R) e^{aR} dR + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$e^{ax} y = \int b(x) e^{ax} dx + c_1$$

Which simplifies to

$$e^{ax} y = \int b(x) e^{ax} dx + c_1$$

Which gives

$$y = e^{-ax} \left(\int b(x) e^{ax} dx + c_1 \right)$$

Summary

The solution(s) found are the following

$$y = e^{-ax} \left(\int b(x) e^{ax} dx + c_1 \right) \quad (1)$$

Verification of solutions

$$y = e^{-ax} \left(\int b(x) e^{ax} dx + c_1 \right)$$

Verified OK.

2.10.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial\phi}{\partial x} + \frac{\partial\phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} dy &= (-ya + b(x)) dx \\ (ya - b(x)) dx + dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= ya - b(x) \\ N(x, y) &= 1 \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(ya - b(x)) \\ &= a\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(1) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned}A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= 1((a) - (0)) \\ &= a\end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int A dx} \\ &= e^{\int a dx}\end{aligned}$$

The result of integrating gives

$$\begin{aligned}\mu &= e^{ax} \\ &= e^{ax}\end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} for now so not to confuse them with the original M and N .

$$\begin{aligned}\bar{M} &= \mu M \\ &= e^{ax}(ya - b(x)) \\ &= (ya - b(x)) e^{ax}\end{aligned}$$

And

$$\begin{aligned}\bar{N} &= \mu N \\ &= e^{ax}(1) \\ &= e^{ax}\end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned}\bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ ((ya - b(x))e^{ax}) + (e^{ax}) \frac{dy}{dx} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \quad (1)$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int (ya - b(x)) e^{ax} dx \\ \phi &= \int^x (ya - b(x)) e^{ax} dx + f(y)\end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = e^{ax} + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = e^{ax}$. Therefore equation (4) becomes

$$e^{ax} = e^{ax} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = \int^x (ya - b(a)) e^{a-a} d_a + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \int^x (ya - b(a)) e^{a-a} d_a$$

Summary

The solution(s) found are the following

$$\int^x (ya - b(a)) e^{a-a} d_a = c_1 \quad (1)$$

Verification of solutions

$$\int^x (ya - b(a)) e^{a-a} d_a = c_1$$

Verified OK.

2.10.4 Maple step by step solution

Let's solve

$$y' + ya = b(x)$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = -ya + b(x)$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' + ya = b(x)$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) (y' + ya) = \mu(x) b(x)$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x)y)$

$$\mu(x)(y' + ya) = \mu'(x)y + \mu(x)y'$$
- Isolate $\mu'(x)$

$$\mu'(x) = \mu(x)a$$
- Solve to find the integrating factor

$$\mu(x) = e^{ax}$$
- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x)y) \right) dx = \int \mu(x)b(x) dx + c_1$$
- Evaluate the integral on the lhs

$$\mu(x)y = \int \mu(x)b(x) dx + c_1$$
- Solve for y

$$y = \frac{\int \mu(x)b(x)dx + c_1}{\mu(x)}$$
- Substitute $\mu(x) = e^{ax}$

$$y = \frac{\int b(x)e^{ax} dx + c_1}{e^{ax}}$$
- Simplify

$$y = e^{-ax} \left(\int b(x) e^{ax} dx + c_1 \right)$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 21

```
dsolve(diff(y(x),x)+a*y(x)=b(x),y(x), singsol=all)
```

$$y(x) = \left(\int b(x) e^{ax} dx + c_1 \right) e^{-ax}$$

✓ Solution by Mathematica

Time used: 0.048 (sec). Leaf size: 32

```
DSolve[y'[x]+a*y[x]==b[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-ax} \left(\int_1^x e^{aK[1]} b(K[1]) dK[1] + c_1 \right)$$

3 Chapter 1. Introduction– Linear equations of First Order. Page 45

3.1	problem 1(a)	215
3.2	problem 1(b)	228
3.3	problem 1(c)	242
3.4	problem 1(d)	255
3.5	problem 1(e)	268
3.6	problem 2	281
3.7	problem 3	295
3.8	problem 8	309
3.9	problem 14(a)	320
3.10	problem 14(b)	324
3.11	problem 14(b)	328

3.1 problem 1(a)

3.1.1	Solving as separable ode	215
3.1.2	Solving as linear ode	217
3.1.3	Solving as first order ode lie symmetry lookup ode	218
3.1.4	Solving as exact ode	222
3.1.5	Maple step by step solution	226

Internal problem ID [5933]

Internal file name [OUTPUT/5181_Sunday_June_05_2022_03_26_55_PM_83246739/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1. Introduction– Linear equations of First Order. Page 45

Problem number: 1(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "linear", "separable", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_separable]

$$2xy + y' = x$$

3.1.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= x(-2y + 1)\end{aligned}$$

Where $f(x) = x$ and $g(y) = -2y + 1$. Integrating both sides gives

$$\begin{aligned}\frac{1}{-2y + 1} dy &= x dx \\ \int \frac{1}{-2y + 1} dy &= \int x dx\end{aligned}$$

$$-\frac{\ln(-2y+1)}{2} = \frac{x^2}{2} + c_1$$

Raising both side to exponential gives

$$\frac{1}{\sqrt{-2y+1}} = e^{\frac{x^2}{2}+c_1}$$

Which simplifies to

$$\frac{1}{\sqrt{-2y+1}} = c_2 e^{\frac{x^2}{2}}$$

Summary

The solution(s) found are the following

$$y = \frac{\left(c_2^2 e^{x^2+2c_1} - 1\right) e^{-x^2-2c_1}}{2c_2^2} \quad (1)$$




Figure 45: Slope field plot

Verification of solutions

$$y = \frac{\left(c_2^2 e^{x^2+2c_1} - 1\right) e^{-x^2-2c_1}}{2c_2^2}$$

Verified OK.

3.1.2 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = 2x$$

$$q(x) = x$$

Hence the ode is

$$2xy + y' = x$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int 2x dx} \\ &= e^{x^2}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu)(x) \\ \frac{d}{dx}(e^{x^2} y) &= (e^{x^2})(x) \\ d(e^{x^2} y) &= (x e^{x^2}) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}e^{x^2} y &= \int x e^{x^2} dx \\ e^{x^2} y &= \frac{e^{x^2}}{2} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{x^2}$ results in

$$y = \frac{e^{-x^2} e^{x^2}}{2} + c_1 e^{-x^2}$$

which simplifies to

$$y = \frac{1}{2} + c_1 e^{-x^2}$$

Summary

The solution(s) found are the following

$$y = \frac{1}{2} + c_1 e^{-x^2} \quad (1)$$




Figure 46: Slope field plot

Verification of solutions

$$y = \frac{1}{2} + c_1 e^{-x^2}$$

Verified OK.

3.1.3 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = -2xy + x$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (A)$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 46: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= e^{-x^2}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^{-x^2}} dy \end{aligned}$$

Which results in

$$S = e^{x^2} y$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -2xy + x$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= 2x e^{x^2} y \\ S_y &= e^{x^2} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = x e^{x^2} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = R e^{R^2}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \frac{e^{R^2}}{2} + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$y e^{x^2} = \frac{e^{x^2}}{2} + c_1$$



Which simplifies to

$$y e^{x^2} = \frac{e^{x^2}}{2} + c_1$$

Which gives

$$y = \frac{(e^{x^2} + 2c_1) e^{-x^2}}{2}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -2xy + x$ 	$R = x$ $S = e^{x^2} y$	$\frac{dS}{dR} = R e^{R^2}$ 

Summary

The solution(s) found are the following

$$y = \frac{(e^{x^2} + 2c_1) e^{-x^2}}{2} \quad (1)$$




Figure 47: Slope field plot

Verification of solutions

$$y = \frac{(e^{x^2} + 2c_1) e^{-x^2}}{2}$$

Verified OK.

3.1.4 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (A)$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the

ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial\phi}{\partial x} + \frac{\partial\phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} \left(\frac{1}{-2y+1}\right) dy &= (x) dx \\ (-x) dx + \left(\frac{1}{-2y+1}\right) dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -x \\ N(x, y) &= \frac{1}{-2y+1} \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-x) \\ &= 0\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}\left(\frac{1}{-2y+1}\right) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \tag{1}$$

$$\frac{\partial \phi}{\partial y} = N \tag{2}$$

Integrating (1) w.r.t. x gives

$$\int \frac{\partial \phi}{\partial x} dx = \int M dx$$

$$\int \frac{\partial \phi}{\partial x} dx = \int -x dx$$

$$\phi = -\frac{x^2}{2} + f(y) \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{1}{-2y+1}$. Therefore equation (4) becomes

$$\frac{1}{-2y+1} = 0 + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = -\frac{1}{2y-1}$$

Integrating the above w.r.t y gives

$$\int f'(y) dy = \int \left(-\frac{1}{2y-1} \right) dy$$
$$f(y) = -\frac{\ln(2y-1)}{2} + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\frac{x^2}{2} - \frac{\ln(2y-1)}{2} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\frac{x^2}{2} - \frac{\ln(2y-1)}{2}$$

The solution becomes

$$y = \frac{e^{-x^2-2c_1}}{2} + \frac{1}{2}$$

Summary

The solution(s) found are the following

$$y = \frac{e^{-x^2-2c_1}}{2} + \frac{1}{2} \tag{1}$$




Figure 48: Slope field plot

Verification of solutions

$$y = \frac{e^{-x^2-2c_1}}{2} + \frac{1}{2}$$

Verified OK.

3.1.5 Maple step by step solution

Let's solve

$$2xy + y' = x$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{2y-1} = -x$$

- Integrate both sides with respect to x

$$\int \frac{y'}{2y-1} dx = \int -x dx + c_1$$

- Evaluate integral

$$\frac{\ln(2y-1)}{2} = -\frac{x^2}{2} + c_1$$

- Solve for y

$$y = \frac{1}{2} + \frac{e^{-x^2+2c_1}}{2}$$

Maple trace

```
`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(diff(y(x),x)+2*x*y(x)=x,y(x), singsol=all)
```

$$y(x) = \frac{1}{2} + e^{-x^2} c_1$$

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 26

```
DSolve[y'[x]+2*x*y[x]==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} + c_1 e^{-x^2}$$

$$y(x) \rightarrow \frac{1}{2}$$

3.2 problem 1(b)

3.2.1	Solving as linear ode	228
3.2.2	Solving as differentialType ode	230
3.2.3	Solving as first order ode lie symmetry lookup ode	232
3.2.4	Solving as exact ode	236
3.2.5	Maple step by step solution	240

Internal problem ID [5934]

Internal file name [OUTPUT/5182_Sunday_June_05_2022_03_26_56_PM_5862443/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1. Introduction– Linear equations of First Order. Page 45

Problem number: 1(b).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "linear", "differentialType", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_linear]

$$xy' + y = 3x^3 - 1$$

3.2.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = \frac{1}{x}$$
$$q(x) = \frac{3x^3 - 1}{x}$$

Hence the ode is

$$y' + \frac{y}{x} = \frac{3x^3 - 1}{x}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int \frac{1}{x} dx} \\ &= x\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu) \left(\frac{3x^3 - 1}{x} \right) \\ \frac{d}{dx}(xy) &= (x) \left(\frac{3x^3 - 1}{x} \right) \\ d(xy) &= (3x^3 - 1) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}xy &= \int 3x^3 - 1 dx \\ xy &= \frac{3}{4}x^4 - x + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = x$ results in

$$y = \frac{\frac{3}{4}x^4 - x}{x} + \frac{c_1}{x}$$

which simplifies to

$$y = \frac{3x^4 + 4c_1 - 4x}{4x}$$

Summary

The solution(s) found are the following

$$y = \frac{3x^4 + 4c_1 - 4x}{4x} \tag{1}$$




Figure 49: Slope field plot

Verification of solutions

$$y = \frac{3x^4 + 4c_1 - 4x}{4x}$$

Verified OK.

3.2.2 Solving as differentialType ode

Writing the ode as

$$y' = \frac{-y + 3x^3 - 1}{x} \tag{1}$$

Which becomes

$$0 = (-x) dy + (3x^3 - y - 1) dx \tag{2}$$

But the RHS is complete differential because

$$(-x) dy + (3x^3 - y - 1) dx = d\left(\frac{3}{4}x^4 - xy - x\right)$$

Hence (2) becomes

$$0 = d\left(\frac{3}{4}x^4 - xy - x\right)$$

Integrating both sides gives gives these solutions

$$y = \frac{3x^4 + 4c_1 - 4x}{4x} + c_1$$

Summary

The solution(s) found are the following

$$y = \frac{3x^4 + 4c_1 - 4x}{4x} + c_1 \tag{1}$$




Figure 50: Slope field plot

Verification of solutions

$$y = \frac{3x^4 + 4c_1 - 4x}{4x} + c_1$$

Verified OK.

3.2.3 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = -\frac{-3x^3 + y + 1}{x}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 49: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= \frac{1}{x}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{1}{x}} dy\end{aligned}$$

Which results in

$$S = xy$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y}\tag{2}$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -\frac{-3x^3 + y + 1}{x}$$

Evaluating all the partial derivatives gives

$$R_x = 1$$

$$R_y = 0$$

$$S_x = y$$

$$S_y = x$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 3x^3 - 1 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 3R^3 - 1$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \frac{3}{4}R^4 - R + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$xy = \frac{3}{4}x^4 - x + c_1$$



Which simplifies to

$$xy = \frac{3}{4}x^4 - x + c_1$$

Which gives

$$y = \frac{3x^4 + 4c_1 - 4x}{4x}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -\frac{-3x^3+y+1}{x}$ 	$R = x$ $S = xy$	$\frac{dS}{dR} = 3R^3 - 1$ 

Summary

The solution(s) found are the following

$$y = \frac{3x^4 + 4c_1 - 4x}{4x} \tag{1}$$




Figure 51: Slope field plot

Verification of solutions

$$y = \frac{3x^4 + 4c_1 - 4x}{4x}$$

Verified OK.

3.2.4 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}(x) dy &= (3x^3 - y - 1) dx \\ (-3x^3 + y + 1) dx + (x) dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -3x^3 + y + 1 \\ N(x, y) &= x\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-3x^3 + y + 1) \\ &= 1\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(x) \\ &= 1\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\int \frac{\partial \phi}{\partial x} dx = \int M dx$$

$$\int \frac{\partial \phi}{\partial x} dx = \int -3x^3 + y + 1 dx$$

$$\phi = -\frac{3}{4}x^4 + xy + x + f(y) \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = x + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = x$. Therefore equation (4) becomes

$$x = x + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = -\frac{3}{4}x^4 + xy + x + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\frac{3}{4}x^4 + xy + x$$

The solution becomes

$$y = \frac{3x^4 + 4c_1 - 4x}{4x}$$

Summary

The solution(s) found are the following

$$y = \frac{3x^4 + 4c_1 - 4x}{4x} \tag{1}$$




Figure 52: Slope field plot

Verification of solutions

$$y = \frac{3x^4 + 4c_1 - 4x}{4x}$$

Verified OK.

3.2.5 Maple step by step solution

Let's solve

$$xy' + y = 3x^3 - 1$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = -\frac{y}{x} + \frac{3x^3-1}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' + \frac{y}{x} = \frac{3x^3-1}{x}$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) \left(y' + \frac{y}{x} \right) = \frac{\mu(x)(3x^3-1)}{x}$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x)y)$

$$\mu(x) \left(y' + \frac{y}{x} \right) = \mu'(x)y + \mu(x)y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = \frac{\mu(x)}{x}$$

- Solve to find the integrating factor

$$\mu(x) = x$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x)y) \right) dx = \int \frac{\mu(x)(3x^3-1)}{x} dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x)y = \int \frac{\mu(x)(3x^3-1)}{x} dx + c_1$$

- Solve for y

$$y = \frac{\int \frac{\mu(x)(3x^3-1)}{x} dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = x$

$$y = \frac{\int (3x^3-1) dx + c_1}{x}$$

- Evaluate the integrals on the rhs

$$y = \frac{\frac{3}{4}x^4 - x + c_1}{x}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(x*diff(y(x),x)+y(x)=3*x^3-1,y(x), singsol=all)
```

$$y(x) = \frac{\frac{3}{4}x^4 - x + c_1}{x}$$

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 20

```
DSolve[x*y'[x]+y[x]==3*x^3-1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{3x^3}{4} + \frac{c_1}{x} - 1$$

3.3 problem 1(c)

3.3.1	Solving as separable ode	242
3.3.2	Solving as linear ode	244
3.3.3	Solving as first order ode lie symmetry lookup ode	245
3.3.4	Solving as exact ode	249
3.3.5	Maple step by step solution	253

Internal problem ID [5935]

Internal file name [OUTPUT/5183_Sunday_June_05_2022_03_26_57_PM_84736558/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1. Introduction– Linear equations of First Order. Page 45

Problem number: 1(c).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "linear", "separable", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_separable]

$$y' + e^x y = 3 e^x$$

3.3.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= f(x)g(y) \\ &= e^x(3 - y) \end{aligned}$$

Where $f(x) = e^x$ and $g(y) = 3 - y$. Integrating both sides gives

$$\begin{aligned} \frac{1}{3 - y} dy &= e^x dx \\ \int \frac{1}{3 - y} dy &= \int e^x dx \\ -\ln(-3 + y) &= e^x + c_1 \end{aligned}$$

Raising both side to exponential gives

$$\frac{1}{-3 + y} = e^{e^x + c_1}$$

Which simplifies to

$$\frac{1}{-3 + y} = c_2 e^{e^x}$$

Summary

The solution(s) found are the following

$$y = \frac{(3c_2 e^{e^x + c_1} + 1) e^{-e^x - c_1}}{c_2} \quad (1)$$




Figure 53: Slope field plot

Verification of solutions

$$y = \frac{(3c_2 e^{e^x + c_1} + 1) e^{-e^x - c_1}}{c_2}$$

Verified OK.

3.3.2 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$\begin{aligned}p(x) &= e^x \\q(x) &= 3e^x\end{aligned}$$

Hence the ode is

$$y' + e^x y = 3e^x$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int e^x dx} \\ &= e^{e^x}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu)(3e^x) \\ \frac{d}{dx}(e^{e^x} y) &= (e^{e^x})(3e^x) \\ d(e^{e^x} y) &= (3e^{e^x+x}) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}e^{e^x} y &= \int 3e^{e^x+x} dx \\ e^{e^x} y &= 3e^{e^x} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{e^x}$ results in

$$y = 3e^{-e^x} e^{e^x} + c_1 e^{-e^x}$$

which simplifies to

$$y = 3 + c_1 e^{-e^x}$$

Summary

The solution(s) found are the following

$$y = 3 + c_1 e^{-e^x} \tag{1}$$




Figure 54: Slope field plot

Verification of solutions

$$y = 3 + c_1 e^{-e^x}$$

Verified OK.

3.3.3 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = -y e^x + 3 e^x$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 52: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= e^{-e^x}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^{-e^x}} dy \end{aligned}$$

Which results in

$$S = e^{e^x} y$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -y e^x + 3 e^x$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= y e^{e^x+x} \\ S_y &= e^{e^x} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 3 e^{e^x+x} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 3 e^{e^R+R}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by

integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = 3 e^{e^R} + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$e^{e^x} y = 3 e^{e^x} + c_1$$



Which simplifies to

$$e^{e^x} y = 3 e^{e^x} + c_1$$

Which gives

$$y = (3 e^{e^x} + c_1) e^{-e^x}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -y e^x + 3 e^x$ 	$R = x$ $S = e^{e^x} y$	$\frac{dS}{dR} = 3 e^{e^R + R}$ 

Summary

The solution(s) found are the following

$$y = (3 e^{e^x} + c_1) e^{-e^x} \quad (1)$$




Figure 55: Slope field plot

Verification of solutions

$$y = (3e^{e^x} + c_1)e^{-e^x}$$

Verified OK.

3.3.4 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial\phi}{\partial x} + \frac{\partial\phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}\left(\frac{1}{3-y}\right) dy &= (e^x) dx \\ (-e^x) dx + \left(\frac{1}{3-y}\right) dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -e^x \\ N(x, y) &= \frac{1}{3-y}\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-e^x) \\ &= 0\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{1}{3-y} \right) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -e^x dx \\ \phi &= -e^x + f(y)\end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{1}{3-y}$. Therefore equation (4) becomes

$$\frac{1}{3-y} = 0 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = -\frac{1}{-3+y}$$

Integrating the above w.r.t y gives

$$\begin{aligned}\int f'(y) dy &= \int \left(-\frac{1}{-3+y} \right) dy \\ f(y) &= -\ln(-3+y) + c_1\end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -e^x - \ln(-3 + y) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -e^x - \ln(-3 + y)$$

The solution becomes

$$y = e^{-e^x - c_1} + 3$$

Summary

The solution(s) found are the following

$$y = e^{-e^x - c_1} + 3 \tag{1}$$




Figure 56: Slope field plot

Verification of solutions

$$y = e^{-e^x - c_1} + 3$$

Verified OK.

3.3.5 Maple step by step solution

Let's solve

$$y' + e^x y = 3e^x$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y-3} = -e^x$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y-3} dx = \int -e^x dx + c_1$$

- Evaluate integral

$$\ln(y-3) = -e^x + c_1$$

- Solve for y

$$y = e^{-e^x + c_1} + 3$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(diff(y(x),x)+exp(x)*y(x)=3*exp(x),y(x), singsol=all)
```

$$y(x) = 3 + e^{-e^x} c_1$$

✓ Solution by Mathematica

Time used: 0.057 (sec). Leaf size: 22

```
DSolve[y'[x]+Exp[x]*y[x]==3*Exp[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 3 + c_1 e^{-e^x}$$

$$y(x) \rightarrow 3$$

3.4 problem 1(d)

3.4.1	Solving as linear ode	255
3.4.2	Solving as first order ode lie symmetry lookup ode	257
3.4.3	Solving as exact ode	261
3.4.4	Maple step by step solution	265

Internal problem ID [5936]

Internal file name [OUTPUT/5184_Sunday_June_05_2022_03_26_59_PM_8251530/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1. Introduction– Linear equations of First Order. Page 45

Problem number: 1(d).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**linear**", "**exactWithIntegrationFactor**", "**first_order_ode_lie_symmetry_lookup**"

Maple gives the following as the ode type

[_linear]

$$y' - y \tan(x) = e^{\sin(x)}$$

3.4.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -\tan(x)$$

$$q(x) = e^{\sin(x)}$$

Hence the ode is

$$y' - y \tan(x) = e^{\sin(x)}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int -\tan(x)dx} \\ &= \cos(x)\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu) (e^{\sin(x)}) \\ \frac{d}{dx}(y \cos(x)) &= (\cos(x)) (e^{\sin(x)}) \\ d(y \cos(x)) &= (\cos(x) e^{\sin(x)}) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}y \cos(x) &= \int \cos(x) e^{\sin(x)} dx \\ y \cos(x) &= e^{\sin(x)} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = \cos(x)$ results in

$$y = e^{\sin(x)} \sec(x) + c_1 \sec(x)$$

which simplifies to

$$y = \sec(x) (e^{\sin(x)} + c_1)$$

Summary

The solution(s) found are the following

$$y = \sec(x) (e^{\sin(x)} + c_1) \tag{1}$$




Figure 57: Slope field plot

Verification of solutions

$$y = \sec(x) (e^{\sin(x)} + c_1)$$

Verified OK.

3.4.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = y \tan(x) + e^{\sin(x)}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 55: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= \frac{1}{\cos(x)}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right)S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{1}{\cos(x)}} dy \end{aligned}$$

Which results in

$$S = y \cos(x)$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = y \tan(x) + e^{\sin(x)}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= -y \sin(x) \\ S_y &= \cos(x) \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \cos(x) e^{\sin(x)} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \cos(R) e^{\sin(R)}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by

integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = e^{\sin(R)} + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$y \cos(x) = e^{\sin(x)} + c_1$$



Which simplifies to

$$y \cos(x) = e^{\sin(x)} + c_1$$

Which gives

$$y = \frac{e^{\sin(x)} + c_1}{\cos(x)}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = y \tan(x) + e^{\sin(x)}$ 	$R = x$ $S = y \cos(x)$	$\frac{dS}{dR} = \cos(R) e^{\sin(R)}$ 

Summary

The solution(s) found are the following

$$y = \frac{e^{\sin(x)} + c_1}{\cos(x)} \quad (1)$$




Figure 58: Slope field plot

Verification of solutions

$$y = \frac{e^{\sin(x)} + c_1}{\cos(x)}$$

Verified OK.

3.4.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned}dy &= (y \tan(x) + e^{\sin(x)}) dx \\ (-y \tan(x) - e^{\sin(x)}) dx + dy &= 0\end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -y \tan(x) - e^{\sin(x)} \\ N(x, y) &= 1\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} (-y \tan(x) - e^{\sin(x)}) \\ &= -\tan(x)\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} (1) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= 1((- \tan(x)) - (0)) \\ &= -\tan(x) \end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int A \, dx} \\ &= e^{\int -\tan(x) \, dx} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^{\ln(\cos(x))} \\ &= \cos(x) \end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} for now so not to confuse them with the original M and N .

$$\begin{aligned} \bar{M} &= \mu M \\ &= \cos(x) (-y \tan(x) - e^{\sin(x)}) \\ &= -y \sin(x) - \cos(x) e^{\sin(x)} \end{aligned}$$

And

$$\begin{aligned} \bar{N} &= \mu N \\ &= \cos(x) (1) \\ &= \cos(x) \end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned} \bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ (-y \sin(x) - \cos(x) e^{\sin(x)}) + (\cos(x)) \frac{dy}{dx} &= 0 \end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \overline{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -y \sin(x) - \cos(x) e^{\sin(x)} dx \\ \phi &= y \cos(x) - e^{\sin(x)} + f(y)\end{aligned}\quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = \cos(x) + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \cos(x)$. Therefore equation (4) becomes

$$\cos(x) = \cos(x) + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = y \cos(x) - e^{\sin(x)} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = y \cos(x) - e^{\sin(x)}$$

The solution becomes

$$y = \frac{e^{\sin(x)} + c_1}{\cos(x)}$$

Summary

The solution(s) found are the following

$$y = \frac{e^{\sin(x)} + c_1}{\cos(x)} \quad (1)$$




Figure 59: Slope field plot

Verification of solutions

$$y = \frac{e^{\sin(x)} + c_1}{\cos(x)}$$

Verified OK.

3.4.4 Maple step by step solution

Let's solve

$$y' - y \tan(x) = e^{\sin(x)}$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = y \tan(x) + e^{\sin(x)}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' - y \tan(x) = e^{\sin(x)}$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) (y' - y \tan(x)) = \mu(x) e^{\sin(x)}$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x) y)$

$$\mu(x) (y' - y \tan(x)) = \mu'(x) y + \mu(x) y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = -\mu(x) \tan(x)$$

- Solve to find the integrating factor

$$\mu(x) = \cos(x)$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x) y) \right) dx = \int \mu(x) e^{\sin(x)} dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x) y = \int \mu(x) e^{\sin(x)} dx + c_1$$

- Solve for y

$$y = \frac{\int \mu(x) e^{\sin(x)} dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = \cos(x)$

$$y = \frac{\int \cos(x) e^{\sin(x)} dx + c_1}{\cos(x)}$$

- Evaluate the integrals on the rhs

$$y = \frac{e^{\sin(x)} + c_1}{\cos(x)}$$

- Simplify

$$y = \sec(x) (e^{\sin(x)} + c_1)$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)-tan(x)*y(x)=exp(sin(x)),y(x), singsol=all)
```

$$y(x) = \sec(x) (e^{\sin(x)} + c_1)$$

✓ Solution by Mathematica

Time used: 0.149 (sec). Leaf size: 15

```
DSolve[y'[x]-Tan[x]*y[x]==Exp[Sin[x]],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \sec(x) (e^{\sin(x)} + c_1)$$

3.5 problem 1(e)

3.5.1	Solving as linear ode	268
3.5.2	Solving as first order ode lie symmetry lookup ode	270
3.5.3	Solving as exact ode	274
3.5.4	Maple step by step solution	279

Internal problem ID [5937]

Internal file name [OUTPUT/5185_Sunday_June_05_2022_03_27_00_PM_16717071/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1. Introduction– Linear equations of First Order. Page 45

Problem number: 1(e).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"linear", "exactWithIntegrationFactor", "first_order_ode_lie_symmetry_lookup"**

Maple gives the following as the ode type

`[_linear]`

$$2xy + y' = x e^{-x^2}$$

3.5.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = 2x$$
$$q(x) = x e^{-x^2}$$

Hence the ode is

$$2xy + y' = x e^{-x^2}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int 2x dx} \\ &= e^{x^2}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu) (x e^{-x^2}) \\ \frac{d}{dx}(e^{x^2} y) &= (e^{x^2}) (x e^{-x^2}) \\ d(e^{x^2} y) &= x dx\end{aligned}$$

Integrating gives

$$\begin{aligned}e^{x^2} y &= \int x dx \\ e^{x^2} y &= \frac{x^2}{2} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{x^2}$ results in

$$y = \frac{x^2 e^{-x^2}}{2} + c_1 e^{-x^2}$$

which simplifies to

$$y = e^{-x^2} \left(\frac{x^2}{2} + c_1 \right)$$

Summary

The solution(s) found are the following

$$y = e^{-x^2} \left(\frac{x^2}{2} + c_1 \right) \tag{1}$$




Figure 60: Slope field plot

Verification of solutions

$$y = e^{-x^2} \left(\frac{x^2}{2} + c_1 \right)$$

Verified OK.

3.5.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = -2xy + x e^{-x^2}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 58: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= e^{-x^2}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right)S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^{-x^2}} dy \end{aligned}$$

Which results in

$$S = e^{x^2} y$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -2xy + x e^{-x^2}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= 2x e^{x^2} y \\ S_y &= e^{x^2} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = x \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = R$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by

integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \frac{R^2}{2} + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$y e^{x^2} = \frac{x^2}{2} + c_1$$



Which simplifies to

$$y e^{x^2} = \frac{x^2}{2} + c_1$$

Which gives

$$y = \frac{e^{-x^2}(x^2 + 2c_1)}{2}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -2xy + x e^{-x^2}$ 	$R = x$ $S = e^{x^2} y$	$\frac{dS}{dR} = R$ 

Summary

The solution(s) found are the following

$$y = \frac{e^{-x^2}(x^2 + 2c_1)}{2} \quad (1)$$




Figure 61: Slope field plot

Verification of solutions

$$y = \frac{e^{-x^2}(x^2 + 2c_1)}{2}$$

Verified OK.

3.5.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}dy &= \left(-2xy + x e^{-x^2}\right) dx \\ \left(2xy - x e^{-x^2}\right) dx + dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= 2xy - x e^{-x^2} \\ N(x, y) &= 1\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(2xy - x e^{-x^2}\right) \\ &= 2x\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} (1) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= 1((2x) - (0)) \\ &= 2x \end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int A \, dx} \\ &= e^{\int 2x \, dx} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^{x^2} \\ &= e^{x^2} \end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} for now so not to confuse them with the original M and N .

$$\begin{aligned} \bar{M} &= \mu M \\ &= e^{x^2} (2xy - x e^{-x^2}) \\ &= 2x e^{x^2} y - x \end{aligned}$$

And

$$\begin{aligned} \bar{N} &= \mu N \\ &= e^{x^2} (1) \\ &= e^{x^2} \end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned} \bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ (2x e^{x^2} y - x) + (e^{x^2}) \frac{dy}{dx} &= 0 \end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \overline{M} \quad (1)$$

$$\frac{\partial \phi}{\partial y} = \overline{N} \quad (2)$$

Integrating (1) w.r.t. x gives

$$\int \frac{\partial \phi}{\partial x} dx = \int \overline{M} dx$$
$$\int \frac{\partial \phi}{\partial x} dx = \int 2x e^{x^2} y - x dx$$
$$\phi = -\frac{x^2}{2} + e^{x^2} y + f(y) \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = e^{x^2} + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = e^{x^2}$. Therefore equation (4) becomes

$$e^{x^2} = e^{x^2} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = -\frac{x^2}{2} + e^{x^2} y + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\frac{x^2}{2} + e^{x^2} y$$

The solution becomes

$$y = \frac{e^{-x^2}(x^2 + 2c_1)}{2}$$

Summary

The solution(s) found are the following

$$y = \frac{e^{-x^2}(x^2 + 2c_1)}{2} \tag{1}$$




Figure 62: Slope field plot

Verification of solutions

$$y = \frac{e^{-x^2}(x^2 + 2c_1)}{2}$$

Verified OK.

3.5.4 Maple step by step solution

Let's solve

$$2xy + y' = x e^{-x^2}$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = -2xy + x e^{-x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$2xy + y' = x e^{-x^2}$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) (2xy + y') = \mu(x) x e^{-x^2}$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x) y)$

$$\mu(x) (2xy + y') = \mu'(x) y + \mu(x) y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = 2\mu(x) x$$

- Solve to find the integrating factor

$$\mu(x) = e^{x^2}$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x) y) \right) dx = \int \mu(x) x e^{-x^2} dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x) y = \int \mu(x) x e^{-x^2} dx + c_1$$

- Solve for y

$$y = \frac{\int \mu(x) x e^{-x^2} dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = e^{x^2}$

$$y = \frac{\int x e^{-x^2} e^{x^2} dx + c_1}{e^{x^2}}$$

- Evaluate the integrals on the rhs

$$y = \frac{\frac{x^2}{2} + c_1}{e^{x^2}}$$

- Simplify

$$y = \frac{e^{-x^2}(x^2 + 2c_1)}{2}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(diff(y(x),x)+2*x*y(x)=x*exp(-x^2),y(x), singsol=all)
```

$$y(x) = \frac{(x^2 + 2c_1)e^{-x^2}}{2}$$

✓ Solution by Mathematica

Time used: 0.057 (sec). Leaf size: 24

```
DSolve[y'[x]+2*x*y[x]==x*Exp[-x^2],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}e^{-x^2}(x^2 + 2c_1)$$

3.6 problem 2

3.6.1	Existence and uniqueness analysis	281
3.6.2	Solving as linear ode	282
3.6.3	Solving as first order ode lie symmetry lookup ode	284
3.6.4	Solving as exact ode	288
3.6.5	Maple step by step solution	292

Internal problem ID [5938]

Internal file name [OUTPUT/5186_Sunday_June_05_2022_03_27_01_PM_41049433/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1. Introduction– Linear equations of First Order. Page 45

Problem number: 2.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**linear**", "**exactWithIntegrationFactor**", "**first_order_ode_lie_symmetry_lookup**"

Maple gives the following as the ode type

`[_linear]`

$$y' + y \cos(x) = e^{-\sin(x)}$$

With initial conditions

$$[y(\pi) = \pi]$$

3.6.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = \cos(x)$$

$$q(x) = e^{-\sin(x)}$$

Hence the ode is

$$y' + y \cos(x) = e^{-\sin(x)}$$

The domain of $p(x) = \cos(x)$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = \pi$ is inside this domain. The domain of $q(x) = e^{-\sin(x)}$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = \pi$ is also inside this domain. Hence solution exists and is unique.

3.6.2 Solving as linear ode

Entering Linear first order ODE solver. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int \cos(x) dx} \\ &= e^{\sin(x)}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu) (e^{-\sin(x)}) \\ \frac{d}{dx}(e^{\sin(x)} y) &= (e^{\sin(x)}) (e^{-\sin(x)}) \\ d(e^{\sin(x)} y) &= dx\end{aligned}$$

Integrating gives

$$\begin{aligned}e^{\sin(x)} y &= \int dx \\ e^{\sin(x)} y &= x + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{\sin(x)}$ results in

$$y = e^{-\sin(x)} x + c_1 e^{-\sin(x)}$$

which simplifies to

$$y = e^{-\sin(x)}(x + c_1)$$

Initial conditions are used to solve for c_1 . Substituting $x = \pi$ and $y = \pi$ in the above solution gives an equation to solve for the constant of integration.

$$\pi = \pi + c_1$$

$$c_1 = 0$$


Substituting c_1 found above in the general solution gives

$$y = e^{-\sin(x)}x$$


Summary

The solution(s) found are the following

$$y = e^{-\sin(x)}x \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = e^{-\sin(x)}x$$

Verified OK.

3.6.3 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = -y \cos(x) + e^{-\sin(x)}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 61: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= e^{-\sin(x)}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^{-\sin(x)}} dy\end{aligned}$$

Which results in

$$S = e^{\sin(x)} y$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y}\tag{2}$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -y \cos(x) + e^{-\sin(x)}$$

Evaluating all the partial derivatives gives

$$\begin{aligned}R_x &= 1 \\ R_y &= 0 \\ S_x &= \cos(x) e^{\sin(x)} y \\ S_y &= e^{\sin(x)}\end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 1 \tag{2A}$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 1$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = R + c_1 \tag{4}$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$e^{\sin(x)}y = x + c_1$$



Which simplifies to

$$e^{\sin(x)}y = x + c_1$$

Which gives

$$y = e^{-\sin(x)}(x + c_1)$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -y \cos(x) + e^{-\sin(x)}$ 	$R = x$ $S = e^{\sin(x)}y$	$\frac{dS}{dR} = 1$ 

Initial conditions are used to solve for c_1 . Substituting $x = \pi$ and $y = \pi$ in the above solution gives an equation to solve for the constant of integration.

$$\pi = \pi + c_1$$

$$c_1 = 0$$


Substituting c_1 found above in the general solution gives

$$y = e^{-\sin(x)}x$$


Summary

The solution(s) found are the following

$$y = e^{-\sin(x)}x \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = e^{-\sin(x)} x$$

Verified OK.

3.6.4 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned} dy &= (-y \cos(x) + e^{-\sin(x)}) dx \\ (y \cos(x) - e^{-\sin(x)}) dx + dy &= 0 \end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= y \cos(x) - e^{-\sin(x)} \\ N(x, y) &= 1 \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} (y \cos(x) - e^{-\sin(x)}) \\ &= \cos(x) \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} (1) \\ &= 0 \end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= 1((\cos(x)) - (0)) \\ &= \cos(x) \end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int A dx} \\ &= e^{\int \cos(x) dx}\end{aligned}$$

The result of integrating gives

$$\begin{aligned}\mu &= e^{\sin(x)} \\ &= e^{\sin(x)}\end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} for now so not to confuse them with the original M and N .

$$\begin{aligned}\bar{M} &= \mu M \\ &= e^{\sin(x)}(y \cos(x) - e^{-\sin(x)}) \\ &= \cos(x) e^{\sin(x)}y - 1\end{aligned}$$

And

$$\begin{aligned}\bar{N} &= \mu N \\ &= e^{\sin(x)}(1) \\ &= e^{\sin(x)}\end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned}\bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ (\cos(x) e^{\sin(x)}y - 1) + (e^{\sin(x)}) \frac{dy}{dx} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int \cos(x) e^{\sin(x)}y - 1 dx \\ \phi &= -x + e^{\sin(x)}y + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = e^{\sin(x)} + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = e^{\sin(x)}$. Therefore equation (4) becomes

$$e^{\sin(x)} = e^{\sin(x)} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = -x + e^{\sin(x)}y + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -x + e^{\sin(x)}y$$

The solution becomes

$$y = e^{-\sin(x)}(x + c_1)$$

Initial conditions are used to solve for c_1 . Substituting $x = \pi$ and $y = \pi$ in the above solution gives an equation to solve for the constant of integration.

$$\pi = \pi + c_1$$

$$c_1 = 0$$


Substituting c_1 found above in the general solution gives

$$y = e^{-\sin(x)}x$$


Summary

The solution(s) found are the following

$$y = e^{-\sin(x)} x \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = e^{-\sin(x)} x$$

Verified OK.

3.6.5 Maple step by step solution

Let's solve

$$[y' + y \cos(x) = e^{-\sin(x)}, y(\pi) = \pi]$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = -y \cos(x) + e^{-\sin(x)}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' + y \cos(x) = e^{-\sin(x)}$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) (y' + y \cos(x)) = \mu(x) e^{-\sin(x)}$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x) y)$

$$\mu(x) (y' + y \cos(x)) = \mu'(x) y + \mu(x) y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = \mu(x) \cos(x)$$

- Solve to find the integrating factor

$$\mu(x) = e^{\sin(x)}$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x) y) \right) dx = \int \mu(x) e^{-\sin(x)} dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x) y = \int \mu(x) e^{-\sin(x)} dx + c_1$$

- Solve for y

$$y = \frac{\int \mu(x) e^{-\sin(x)} dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = e^{\sin(x)}$

$$y = \frac{\int e^{-\sin(x)} e^{\sin(x)} dx + c_1}{e^{\sin(x)}}$$

- Evaluate the integrals on the rhs

$$y = \frac{x + c_1}{e^{\sin(x)}}$$

- Simplify

$$y = e^{-\sin(x)}(x + c_1)$$

- Use initial condition $y(\pi) = \pi$

$$\pi = \pi + c_1$$

- Solve for c_1

$$c_1 = 0$$

- Substitute $c_1 = 0$ into general solution and simplify

$$y = e^{-\sin(x)} x$$

- Solution to the IVP

$$y = e^{-\sin(x)} x$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 11

```
dsolve([diff(y(x),x)+cos(x)*y(x)=exp(-sin(x)),y(Pi) = Pi],y(x), singsol=all)
```

$$y(x) = e^{-\sin(x)}x$$

✓ Solution by Mathematica

Time used: 0.138 (sec). Leaf size: 13

```
DSolve[{y'[x]+Cos[x]*y[x]==Exp[-Sin[x]],{y[Pi]==Pi}},y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow xe^{-\sin(x)}$$

3.7 problem 3

3.7.1	Solving as linear ode	295
3.7.2	Solving as differentialType ode	297
3.7.3	Solving as first order ode lie symmetry lookup ode	299
3.7.4	Solving as exact ode	303
3.7.5	Maple step by step solution	307

Internal problem ID [5939]

Internal file name [OUTPUT/5187_Sunday_June_05_2022_03_27_03_PM_39784597/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1. Introduction– Linear equations of First Order. Page 45

Problem number: 3.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "linear", "differentialType", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_linear]

$$x^2y' + 2xy = 1$$

3.7.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = \frac{2}{x}$$
$$q(x) = \frac{1}{x^2}$$

Hence the ode is

$$y' + \frac{2y}{x} = \frac{1}{x^2}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int \frac{2}{x} dx} \\ &= x^2\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu) \left(\frac{1}{x^2} \right) \\ \frac{d}{dx}(y x^2) &= (x^2) \left(\frac{1}{x^2} \right) \\ d(y x^2) &= dx\end{aligned}$$

Integrating gives

$$\begin{aligned}y x^2 &= \int dx \\ y x^2 &= x + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = x^2$ results in

$$y = \frac{1}{x} + \frac{c_1}{x^2}$$

which simplifies to

$$y = \frac{x + c_1}{x^2}$$

Summary

The solution(s) found are the following

$$y = \frac{x + c_1}{x^2} \tag{1}$$




Figure 66: Slope field plot

Verification of solutions

$$y = \frac{x + c_1}{x^2}$$

Verified OK.

3.7.2 Solving as differentialType ode

Writing the ode as

$$y' = \frac{-2xy + 1}{x^2} \tag{1}$$

Which becomes

$$0 = (-x^2) dy + (-2xy + 1) dx \tag{2}$$

But the RHS is complete differential because

$$(-x^2) dy + (-2xy + 1) dx = d(-y x^2 + x)$$

Hence (2) becomes

$$0 = d(-y x^2 + x)$$

Integrating both sides gives gives these solutions

$$y = \frac{x + c_1}{x^2} + c_1$$

Summary

The solution(s) found are the following

$$y = \frac{x + c_1}{x^2} + c_1 \tag{1}$$




Figure 67: Slope field plot

Verification of solutions

$$y = \frac{x + c_1}{x^2} + c_1$$

Verified OK.

3.7.3 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = -\frac{2xy - 1}{x^2}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 64: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= \frac{1}{x^2}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{1}{x^2}} dy\end{aligned}$$

Which results in

$$S = y x^2$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y}\tag{2}$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -\frac{2xy - 1}{x^2}$$

Evaluating all the partial derivatives gives

$$\begin{aligned}R_x &= 1 \\R_y &= 0 \\S_x &= 2xy \\S_y &= x^2\end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 1 \tag{2A}$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 1$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = R + c_1 \tag{4}$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$yx^2 = x + c_1$$



Which simplifies to

$$yx^2 = x + c_1$$

Which gives

$$y = \frac{x + c_1}{x^2}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -\frac{2xy-1}{x^2}$ 	$R = x$ $S = yx^2$	$\frac{dS}{dR} = 1$ 

Summary

The solution(s) found are the following

$$y = \frac{x + c_1}{x^2} \tag{1}$$




Figure 68: Slope field plot

Verification of solutions

$$y = \frac{x + c_1}{x^2}$$

Verified OK.

3.7.4 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned}(x^2) dy &= (-2xy + 1) dx \\ (2xy - 1) dx + (x^2) dy &= 0\end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= 2xy - 1 \\ N(x, y) &= x^2\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(2xy - 1) \\ &= 2x\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(x^2) \\ &= 2x\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\int \frac{\partial \phi}{\partial x} dx = \int M dx$$

$$\int \frac{\partial \phi}{\partial x} dx = \int 2xy - 1 dx$$

$$\phi = yx^2 - x + f(y) \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = x^2 + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = x^2$. Therefore equation (4) becomes

$$x^2 = x^2 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = yx^2 - x + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = yx^2 - x$$

The solution becomes

$$y = \frac{x + c_1}{x^2}$$

Summary

The solution(s) found are the following

$$y = \frac{x + c_1}{x^2} \tag{1}$$




Figure 69: Slope field plot

Verification of solutions

$$y = \frac{x + c_1}{x^2}$$

Verified OK.

3.7.5 Maple step by step solution

Let's solve

$$x^2 y' + 2xy = 1$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = -\frac{2y}{x} + \frac{1}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' + \frac{2y}{x} = \frac{1}{x^2}$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) \left(y' + \frac{2y}{x} \right) = \frac{\mu(x)}{x^2}$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x)y)$

$$\mu(x) \left(y' + \frac{2y}{x} \right) = \mu'(x)y + \mu(x)y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = \frac{2\mu(x)}{x}$$

- Solve to find the integrating factor

$$\mu(x) = x^2$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x)y) \right) dx = \int \frac{\mu(x)}{x^2} dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x)y = \int \frac{\mu(x)}{x^2} dx + c_1$$

- Solve for y

$$y = \frac{\int \frac{\mu(x)}{x^2} dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = x^2$

$$y = \frac{\int 1 dx + c_1}{x^2}$$

- Evaluate the integrals on the rhs

$$y = \frac{x + c_1}{x^2}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve(x^2*diff(y(x),x)+2*x*y(x)=1,y(x), singsol=all)
```

$$y(x) = \frac{x + c_1}{x^2}$$

✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 13

```
DSolve[x^2*y'[x]+2*x*y[x]==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x + c_1}{x^2}$$

3.8 problem 8

3.8.1	Solving as linear ode	309
3.8.2	Solving as first order ode lie symmetry lookup ode	311
3.8.3	Solving as exact ode	314
3.8.4	Maple step by step solution	317

Internal problem ID [5940]

Internal file name [OUTPUT/5188_Sunday_June_05_2022_03_27_04_PM_44227942/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1. Introduction– Linear equations of First Order. Page 45

Problem number: 8.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "linear", "exactWithIntegrationFactor", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

```
[[_linear, `class A`]]
```

$$y' + 2y = b(x)$$

3.8.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = 2$$

$$q(x) = b(x)$$

Hence the ode is

$$y' + 2y = b(x)$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int 2dx} \\ &= e^{2x}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu) (b(x)) \\ \frac{d}{dx}(e^{2x}y) &= (e^{2x}) (b(x)) \\ d(e^{2x}y) &= (b(x) e^{2x}) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}e^{2x}y &= \int b(x) e^{2x} dx \\ e^{2x}y &= \int b(x) e^{2x} dx + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{2x}$ results in

$$y = e^{-2x} \left(\int b(x) e^{2x} dx \right) + c_1 e^{-2x}$$

which simplifies to

$$y = e^{-2x} \left(\int b(x) e^{2x} dx + c_1 \right)$$

Summary

The solution(s) found are the following

$$y = e^{-2x} \left(\int b(x) e^{2x} dx + c_1 \right) \tag{1}$$

Verification of solutions

$$y = e^{-2x} \left(\int b(x) e^{2x} dx + c_1 \right)$$

Verified OK.

3.8.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = -2y + b(x)$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 67: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= e^{-2x}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^{-2x}} dy\end{aligned}$$

Which results in

$$S = e^{2x}y$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y}\tag{2}$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -2y + b(x)$$

Evaluating all the partial derivatives gives

$$\begin{aligned}R_x &= 1 \\ R_y &= 0 \\ S_x &= 2e^{2x}y \\ S_y &= e^{2x}\end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = b(x) e^{2x} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = b(R) e^{2R}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \int b(R) e^{2R} dR + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$e^{2x} y = \int b(x) e^{2x} dx + c_1$$

Which simplifies to

$$e^{2x} y = \int b(x) e^{2x} dx + c_1$$

Which gives

$$y = e^{-2x} \left(\int b(x) e^{2x} dx + c_1 \right)$$

Summary

The solution(s) found are the following

$$y = e^{-2x} \left(\int b(x) e^{2x} dx + c_1 \right) \quad (1)$$

Verification of solutions

$$y = e^{-2x} \left(\int b(x) e^{2x} dx + c_1 \right)$$

Verified OK.

3.8.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial\phi}{\partial x} + \frac{\partial\phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} dy &= (-2y + b(x)) dx \\ (2y - b(x)) dx + dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= 2y - b(x) \\ N(x, y) &= 1 \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(2y - b(x)) \\ &= 2\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(1) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned}A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= 1((2) - (0)) \\ &= 2\end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int A dx} \\ &= e^{\int 2 dx}\end{aligned}$$

The result of integrating gives

$$\begin{aligned}\mu &= e^{2x} \\ &= e^{2x}\end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} for now so not to confuse them with the original M and N .

$$\begin{aligned}\bar{M} &= \mu M \\ &= e^{2x}(2y - b(x)) \\ &= (2y - b(x))e^{2x}\end{aligned}$$

And

$$\begin{aligned}\bar{N} &= \mu N \\ &= e^{2x}(1) \\ &= e^{2x}\end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned}\bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ ((2y - b(x)) e^{2x}) + (e^{2x}) \frac{dy}{dx} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int (2y - b(x)) e^{2x} dx \\ \phi &= \int^x (2y - b(x)) e^{2x} dx + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = e^{2x} + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = e^{2x}$. Therefore equation (4) becomes

$$e^{2x} = e^{2x} + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = \int^x (2y - b(a)) e^{2-a} da + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \int^x (2y - b(a)) e^{2-a} da$$

Summary

The solution(s) found are the following

$$\int^x (2y - b(a)) e^{2-a} da = c_1 \quad (1)$$

Verification of solutions

$$\int^x (2y - b(a)) e^{2-a} da = c_1$$

Verified OK.

3.8.4 Maple step by step solution

Let's solve

$$y' + 2y = b(x)$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = -2y + b(x)$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' + 2y = b(x)$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) (y' + 2y) = \mu(x) b(x)$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x)y)$

$$\mu(x)(y' + 2y) = \mu'(x)y + \mu(x)y'$$
- Isolate $\mu'(x)$

$$\mu'(x) = 2\mu(x)$$
- Solve to find the integrating factor

$$\mu(x) = e^{2x}$$
- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x)y) \right) dx = \int \mu(x)b(x) dx + c_1$$
- Evaluate the integral on the lhs

$$\mu(x)y = \int \mu(x)b(x) dx + c_1$$
- Solve for y

$$y = \frac{\int \mu(x)b(x)dx + c_1}{\mu(x)}$$
- Substitute $\mu(x) = e^{2x}$

$$y = \frac{\int b(x)e^{2x} dx + c_1}{e^{2x}}$$
- Simplify

$$y = e^{-2x} \left(\int b(x) e^{2x} dx + c_1 \right)$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 20

```
dsolve(diff(y(x),x)+2*y(x)=b(x),y(x), singsol=all)
```

$$y(x) = \left(\int b(x) e^{2x} dx + c_1 \right) e^{-2x}$$

✓ Solution by Mathematica

Time used: 0.045 (sec). Leaf size: 31

```
DSolve[y'[x]+2*y[x]==b[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2x} \left(\int_1^x e^{2K[1]} b(K[1]) dK[1] + c_1 \right)$$

3.9 problem 14(a)

3.9.1	Existence and uniqueness analysis	320
3.9.2	Solving as quadrature ode	321
3.9.3	Maple step by step solution	322

Internal problem ID [5941]

Internal file name [OUTPUT/5189_Sunday_June_05_2022_03_27_05_PM_11378483/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1. Introduction– Linear equations of First Order. Page 45

Problem number: 14(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

`[_quadrature]`

$$y' - y = 1$$

With initial conditions

$$[y(0) = 0]$$

3.9.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -1$$

$$q(x) = 1$$

Hence the ode is

$$y' - y = 1$$

The domain of $p(x) = -1$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = 1$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

3.9.2 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned}\int \frac{1}{1+y} dy &= \int dx \\ \ln(1+y) &= x + c_1\end{aligned}$$

Raising both side to exponential gives

$$1 + y = e^{x+c_1}$$

Which simplifies to

$$1 + y = c_2 e^x$$

Initial conditions are used to solve for c_2 . Substituting $x = 0$ and $y = 0$ in the above solution gives an equation to solve for the constant of integration.

$$0 = -1 + c_2$$

$$c_2 = 1$$


Substituting c_2 found above in the general solution gives

$$y = e^x - 1$$


Summary

The solution(s) found are the following

$$y = e^x - 1 \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = e^x - 1$$

Verified OK.

3.9.3 Maple step by step solution

Let's solve

$$[y' - y = 1, y(0) = 0]$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{1+y} = 1$$

- Integrate both sides with respect to x

$$\int \frac{y'}{1+y} dx = \int 1 dx + c_1$$

- Evaluate integral

$$\ln(1+y) = x + c_1$$

- Solve for y

$$y = e^{x+c_1} - 1$$

- Use initial condition $y(0) = 0$
 $0 = e^{c_1} - 1$
- Solve for c_1
 $c_1 = 0$
- Substitute $c_1 = 0$ into general solution and simplify
 $y = e^x - 1$
- Solution to the IVP
 $y = e^x - 1$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 8

```
dsolve([diff(y(x),x)=1+y(x),y(0) = 0],y(x), singsol=all)
```

$$y(x) = -1 + e^x$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 10

```
DSolve[{y'[x]==1+y[x],{y[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x - 1$$

3.10 problem 14(b)

3.10.1 Existence and uniqueness analysis	324
3.10.2 Solving as quadrature ode	325
3.10.3 Maple step by step solution	326

Internal problem ID [5942]

Internal file name [OUTPUT/5190_Sunday_June_05_2022_03_27_07_PM_65145226/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1. Introduction– Linear equations of First Order. Page 45

Problem number: 14(b).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

`[_quadrature]`

$$y' - y^2 = 1$$

With initial conditions

$$[y(0) = 0]$$

3.10.1 Existence and uniqueness analysis

This is non linear first order ODE. In canonical form it is written as

$$\begin{aligned} y' &= f(x, y) \\ &= y^2 + 1 \end{aligned}$$

The y domain of $f(x, y)$ when $x = 0$ is

$$\{-\infty < y < \infty\}$$

And the point $y_0 = 0$ is inside this domain. Now we will look at the continuity of

$$\begin{aligned} \frac{\partial f}{\partial y} &= \frac{\partial}{\partial y}(y^2 + 1) \\ &= 2y \end{aligned}$$

The y domain of $\frac{\partial f}{\partial y}$ when $x = 0$ is

$$\{-\infty < y < \infty\}$$

And the point $y_0 = 0$ is inside this domain. Therefore solution exists and is unique.

3.10.2 Solving as quadrature ode

Integrating both sides gives

$$\int \frac{1}{y^2 + 1} dy = x + c_1$$
$$\arctan(y) = x + c_1$$

Solving for y gives these solutions

$$y_1 = \tan(x + c_1)$$

Initial conditions are used to solve for c_1 . Substituting $x = 0$ and $y = 0$ in the above solution gives an equation to solve for the constant of integration.

$$0 = \tan(c_1)$$

$$c_1 = 0$$


Substituting c_1 found above in the general solution gives

$$y = \tan(x)$$


Summary

The solution(s) found are the following

$$y = \tan(x) \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \tan(x)$$

Verified OK.

3.10.3 Maple step by step solution

Let's solve

$$[y' - y^2 = 1, y(0) = 0]$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{1+y^2} = 1$$

- Integrate both sides with respect to x

$$\int \frac{y'}{1+y^2} dx = \int 1 dx + c_1$$

- Evaluate integral

$$\arctan(y) = x + c_1$$

- Solve for y

$$y = \tan(x + c_1)$$

- Use initial condition $y(0) = 0$
 $0 = \tan(c_1)$
- Solve for c_1
 $c_1 = 0$
- Substitute $c_1 = 0$ into general solution and simplify
 $y = \tan(x)$
- Solution to the IVP
 $y = \tan(x)$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
<- separable successful`

```

✓ Solution by Maple

Time used: 0.046 (sec). Leaf size: 6

```
dsolve([diff(y(x),x)=1+y(x)^2,y(0) = 0],y(x), singsol=all)
```

$$y(x) = \tan(x)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 7

```
DSolve[{y'[x]==1+y[x]^2,{y[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \tan(x)$$

3.11 problem 14(b)

3.11.1 Existence and uniqueness analysis	328
3.11.2 Solving as quadrature ode	329
3.11.3 Maple step by step solution	330

Internal problem ID [5943]

Internal file name [OUTPUT/5191_Sunday_June_05_2022_03_27_08_PM_70795525/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 1. Introduction– Linear equations of First Order. Page 45

Problem number: 14(b).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

`[_quadrature]`

$$y' - y^2 = 1$$

With initial conditions

$$[y(0) = 0]$$

3.11.1 Existence and uniqueness analysis

This is non linear first order ODE. In canonical form it is written as

$$\begin{aligned}y' &= f(x, y) \\ &= y^2 + 1\end{aligned}$$

The y domain of $f(x, y)$ when $x = 0$ is

$$\{-\infty < y < \infty\}$$

And the point $y_0 = 0$ is inside this domain. Now we will look at the continuity of

$$\begin{aligned}\frac{\partial f}{\partial y} &= \frac{\partial}{\partial y}(y^2 + 1) \\ &= 2y\end{aligned}$$

The y domain of $\frac{\partial f}{\partial y}$ when $x = 0$ is

$$\{-\infty < y < \infty\}$$

And the point $y_0 = 0$ is inside this domain. Therefore solution exists and is unique.

3.11.2 Solving as quadrature ode

Integrating both sides gives

$$\int \frac{1}{y^2 + 1} dy = x + c_1$$
$$\arctan(y) = x + c_1$$

Solving for y gives these solutions

$$y_1 = \tan(x + c_1)$$

Initial conditions are used to solve for c_1 . Substituting $x = 0$ and $y = 0$ in the above solution gives an equation to solve for the constant of integration.

$$0 = \tan(c_1)$$

$$c_1 = 0$$


Substituting c_1 found above in the general solution gives

$$y = \tan(x)$$


Summary

The solution(s) found are the following

$$y = \tan(x) \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \tan(x)$$

Verified OK.

3.11.3 Maple step by step solution

Let's solve

$$[y' - y^2 = 1, y(0) = 0]$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{1+y^2} = 1$$

- Integrate both sides with respect to x

$$\int \frac{y'}{1+y^2} dx = \int 1 dx + c_1$$

- Evaluate integral

$$\arctan(y) = x + c_1$$

- Solve for y

$$y = \tan(x + c_1)$$

- Use initial condition $y(0) = 0$
 $0 = \tan(c_1)$
- Solve for c_1
 $c_1 = 0$
- Substitute $c_1 = 0$ into general solution and simplify
 $y = \tan(x)$
- Solution to the IVP
 $y = \tan(x)$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
<- separable successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 6

```
dsolve([diff(y(x),x)=1+y(x)^2,y(0) = 0],y(x), singsol=all)
```

$$y(x) = \tan(x)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 7

```
DSolve[{y'[x]==1+y[x]^2,{y[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \tan(x)$$

4 Chapter 2. Linear equations with constant coefficients. Page 52

4.1	problem 1(a)	333
4.2	problem 1(b)	343
4.3	problem 1(c)	354
4.4	problem 1(d)	364
4.5	problem 1(e)	378
4.6	problem 1(f)	386
4.7	problem 1(g)	394
4.8	problem 2(a)	402
4.9	problem 2(b)	412
4.10	problem 3(a)	422
4.11	problem 3(b)	432
4.12	problem 3(c)	441
4.13	problem 3(d)	452

4.1 problem 1(a)

4.1.1	Solving as second order linear constant coeff ode	333
4.1.2	Solving as second order ode can be made integrable ode	335
4.1.3	Solving using Kovacic algorithm	337
4.1.4	Maple step by step solution	341

Internal problem ID [5944]

Internal file name [OUTPUT/5192_Sunday_June_05_2022_03_27_10_PM_82633137/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 52

Problem number: 1(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "second_order_ode_can_be_made_integrable"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' - 4y = 0$$

4.1.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = -4$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 4 e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 4 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = -4$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(-4)} \\ &= \pm 2\end{aligned}$$

Hence

$$\lambda_1 = +2$$

$$\lambda_2 = -2$$

Which simplifies to

$$\lambda_1 = 2$$

$$\lambda_2 = -2$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(2)x} + c_2 e^{(-2)x}$$

Or

$$y = c_1 e^{2x} + c_2 e^{-2x}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{2x} + c_2 e^{-2x} \tag{1}$$




Figure 73: Slope field plot

Verification of solutions

$$y = c_1 e^{2x} + c_2 e^{-2x}$$

Verified OK.

4.1.2 Solving as second order ode can be made integrable ode

Multiplying the ode by y' gives

$$y' y'' - 4y' y = 0$$

Integrating the above w.r.t x gives

$$\int (y' y'' - 4y' y) dx = 0$$

$$\frac{y'^2}{2} - 2y^2 = c_2$$

Which is now solved for y . Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \sqrt{4y^2 + 2c_1} \tag{1}$$

$$y' = -\sqrt{4y^2 + 2c_1} \tag{2}$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int \frac{1}{\sqrt{4y^2 + 2c_1}} dy = \int dx$$

$$\frac{\ln(y\sqrt{4} + \sqrt{4y^2 + 2c_1})\sqrt{4}}{4} = c_2 + x$$

Raising both side to exponential gives

$$e^{\frac{\ln(y\sqrt{4} + \sqrt{4y^2 + 2c_1})\sqrt{4}}{4}} = e^{c_2 + x}$$

Which simplifies to

$$\sqrt{2y + \sqrt{4y^2 + 2c_1}} = c_3 e^x$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{1}{\sqrt{4y^2 + 2c_1}} dy = \int dx$$

$$-\frac{\ln(y\sqrt{4} + \sqrt{4y^2 + 2c_1})\sqrt{4}}{4} = x + c_4$$

Raising both side to exponential gives

$$e^{-\frac{\ln(y\sqrt{4} + \sqrt{4y^2 + 2c_1})\sqrt{4}}{4}} = e^{x + c_4}$$

Which simplifies to

$$\frac{1}{\sqrt{2y + \sqrt{4y^2 + 2c_1}}} = c_5 e^x$$

Summary

The solution(s) found are the following

$$y = \frac{(e^{4x} c_3^4 - 2c_1) e^{-2x}}{4c_3^2} \quad (1)$$

$$y = -\frac{(2c_1 c_5^4 e^{4x} - 1) e^{-2x}}{4c_5^2} \quad (2)$$




Figure 74: Slope field plot

Verification of solutions

$$y = \frac{(e^{4x}c_3^4 - 2c_1) e^{-2x}}{4c_3^2}$$

Verified OK.

$$y = -\frac{(2c_1c_5^4e^{4x} - 1) e^{-2x}}{4c_5^2}$$

Verified OK.

4.1.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 4y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= 1 \\B &= 0 \\C &= -4\end{aligned}\tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x)\tag{4}$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\&= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned}\tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{4}{1}\tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= 4 \\t &= 1\end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 4z(x)\tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 73: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 4$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-2x}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}
 y_1 &= z_1 \\
 &= e^{-2x}
 \end{aligned}$$

Which simplifies to

$$y_1 = e^{-2x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= e^{-2x} \int \frac{1}{e^{-4x}} dx \\ &= e^{-2x} \left(\frac{e^{4x}}{4} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-2x}) + c_2 \left(e^{-2x} \left(\frac{e^{4x}}{4} \right) \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-2x} + \frac{c_2 e^{2x}}{4} \tag{1}$$




Figure 75: Slope field plot

Verification of solutions

$$y = c_1 e^{-2x} + \frac{c_2 e^{2x}}{4}$$

Verified OK.

4.1.4 Maple step by step solution

Let's solve

$$y'' - 4y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 - 4 = 0$$

- Factor the characteristic polynomial

$$(r - 2)(r + 2) = 0$$

- Roots of the characteristic polynomial

- $r = (-2, 2)$
- 1st solution of the ODE
 $y_1(x) = e^{-2x}$
 - 2nd solution of the ODE
 $y_2(x) = e^{2x}$
 - General solution of the ODE
 $y = c_1y_1(x) + c_2y_2(x)$
 - Substitute in solutions
 $y = c_1e^{-2x} + c_2e^{2x}$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)-4*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{2x}c_1 + c_2e^{-2x}$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 22

```
DSolve[y''[x]-4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2x}(c_1e^{4x} + c_2)$$

4.2 problem 1(b)

4.2.1	Solving as second order linear constant coeff ode	343
4.2.2	Solving as second order ode can be made integrable ode	345
4.2.3	Solving using Kovacic algorithm	347
4.2.4	Maple step by step solution	351

Internal problem ID [5945]

Internal file name [OUTPUT/5193_Sunday_June_05_2022_03_27_11_PM_74597554/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 52

Problem number: 1(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "second_order_ode_can_be_made_integrable"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$3y'' + 2y = 0$$

4.2.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 3, B = 0, C = 2$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$3\lambda^2 e^{\lambda x} + 2e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$3\lambda^2 + 2 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 3, B = 0, C = 2$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{0}{(2)(3)} \pm \frac{1}{(2)(3)} \sqrt{0^2 - (4)(3)(2)} \\ &= \pm \frac{i\sqrt{6}}{3}\end{aligned}$$

Hence

$$\begin{aligned}\lambda_1 &= +\frac{i\sqrt{6}}{3} \\ \lambda_2 &= -\frac{i\sqrt{6}}{3}\end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= \frac{i\sqrt{6}}{3} \\ \lambda_2 &= -\frac{i\sqrt{6}}{3}\end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = \frac{\sqrt{6}}{3}$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 \left(c_1 \cos \left(\frac{x\sqrt{6}}{3} \right) + c_2 \sin \left(\frac{x\sqrt{6}}{3} \right) \right)$$

Or

$$y = c_1 \cos \left(\frac{x\sqrt{6}}{3} \right) + c_2 \sin \left(\frac{x\sqrt{6}}{3} \right)$$

Summary

The solution(s) found are the following

$$y = c_1 \cos \left(\frac{x\sqrt{6}}{3} \right) + c_2 \sin \left(\frac{x\sqrt{6}}{3} \right) \quad (1)$$




Figure 76: Slope field plot

Verification of solutions

$$y = c_1 \cos\left(\frac{x\sqrt{6}}{3}\right) + c_2 \sin\left(\frac{x\sqrt{6}}{3}\right)$$

Verified OK.

4.2.2 Solving as second order ode can be made integrable ode

Multiplying the ode by y' gives

$$3y'y'' + 2y'y = 0$$

Integrating the above w.r.t x gives

$$\int (3y'y'' + 2y'y) dx = 0$$

$$\frac{3y'^2}{2} + y^2 = c_2$$

Which is now solved for y . Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \frac{\sqrt{-6y^2 + 6c_1}}{3} \quad (1)$$

$$y' = -\frac{\sqrt{-6y^2 + 6c_1}}{3} \quad (2)$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int \frac{3}{\sqrt{-6y^2 + 6c_1}} dy = \int dx$$

$$\frac{\sqrt{6} \arctan\left(\frac{\sqrt{6}y}{\sqrt{-6y^2 + 6c_1}}\right)}{2} = c_2 + x$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{3}{\sqrt{-6y^2 + 6c_1}} dy = \int dx$$

$$-\frac{\sqrt{6} \arctan\left(\frac{\sqrt{6}y}{\sqrt{-6y^2 + 6c_1}}\right)}{2} = x + c_3$$

Summary

The solution(s) found are the following

$$\frac{\sqrt{6} \arctan\left(\frac{\sqrt{6}y}{\sqrt{-6y^2 + 6c_1}}\right)}{2} = c_2 + x \quad (1)$$

$$-\frac{\sqrt{6} \arctan\left(\frac{\sqrt{6}y}{\sqrt{-6y^2 + 6c_1}}\right)}{2} = x + c_3 \quad (2)$$




Figure 77: Slope field plot

Verification of solutions

$$\frac{\sqrt{6} \arctan\left(\frac{\sqrt{6}y}{\sqrt{-6y^2+6c_1}}\right)}{2} = c_2 + x$$

Verified OK.

$$-\frac{\sqrt{6} \arctan\left(\frac{\sqrt{6}y}{\sqrt{-6y^2+6c_1}}\right)}{2} = x + c_3$$

Verified OK.

4.2.3 Solving using Kovacic algorithm

Writing the ode as

$$3y'' + 2y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 3 \\ B &= 0 \\ C &= 2 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-2}{3} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -2 \\ t &= 3 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -\frac{2z(x)}{3} \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 75: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -\frac{2}{3}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos\left(\frac{x\sqrt{6}}{3}\right)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}
 y_1 &= z_1 \\
 &= \cos\left(\frac{x\sqrt{6}}{3}\right)
 \end{aligned}$$

Which simplifies to

$$y_1 = \cos\left(\frac{x\sqrt{6}}{3}\right)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos\left(\frac{x\sqrt{6}}{3}\right) \int \frac{1}{\cos\left(\frac{x\sqrt{6}}{3}\right)^2} dx \\ &= \cos\left(\frac{x\sqrt{6}}{3}\right) \left(\frac{\sqrt{6} \tan\left(\frac{x\sqrt{6}}{3}\right)}{2}\right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(\cos\left(\frac{x\sqrt{6}}{3}\right)\right) + c_2 \left(\cos\left(\frac{x\sqrt{6}}{3}\right) \left(\frac{\sqrt{6} \tan\left(\frac{x\sqrt{6}}{3}\right)}{2}\right)\right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \cos\left(\frac{x\sqrt{6}}{3}\right) + \frac{c_2 \sqrt{6} \sin\left(\frac{x\sqrt{6}}{3}\right)}{2} \quad (1)$$




Figure 78: Slope field plot

Verification of solutions

$$y = c_1 \cos\left(\frac{x\sqrt{6}}{3}\right) + \frac{c_2\sqrt{6} \sin\left(\frac{x\sqrt{6}}{3}\right)}{2}$$

Verified OK.

4.2.4 Maple step by step solution

Let's solve

$$3y'' + 2y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{2y}{3}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{2y}{3} = 0$$

- Characteristic polynomial of ODE

$$r^2 + \frac{2}{3} = 0$$
- Use quadratic formula to solve for r

$$r = \frac{0 \pm \left(\sqrt{-\frac{8}{3}}\right)}{2}$$
- Roots of the characteristic polynomial

$$r = \left(-\frac{1}{3}\sqrt{6}, \frac{1}{3}\sqrt{6}\right)$$
- 1st solution of the ODE

$$y_1(x) = \cos\left(\frac{x\sqrt{6}}{3}\right)$$
- 2nd solution of the ODE

$$y_2(x) = \sin\left(\frac{x\sqrt{6}}{3}\right)$$
- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$
- Substitute in solutions

$$y = c_1 \cos\left(\frac{x\sqrt{6}}{3}\right) + c_2 \sin\left(\frac{x\sqrt{6}}{3}\right)$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(3*diff(y(x),x$2)+2*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 \sin\left(\frac{\sqrt{6}x}{3}\right) + c_2 \cos\left(\frac{\sqrt{6}x}{3}\right)$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 32

```
DSolve[3*y''[x]+2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos\left(\sqrt{\frac{2}{3}}x\right) + c_2 \sin\left(\sqrt{\frac{2}{3}}x\right)$$

4.3 problem 1(c)

4.3.1	Solving as second order linear constant coeff ode	354
4.3.2	Solving as second order ode can be made integrable ode	356
4.3.3	Solving using Kovacic algorithm	358
4.3.4	Maple step by step solution	362

Internal problem ID [5946]

Internal file name [OUTPUT/5194_Sunday_June_05_2022_03_27_12_PM_22379450/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 52

Problem number: 1(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "second_order_ode_can_be_made_integrable"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' + 16y = 0$$

4.3.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 16$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 16 e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 16 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 16$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(16)} \\ &= \pm 4i\end{aligned}$$

Hence

$$\lambda_1 = +4i$$

$$\lambda_2 = -4i$$

Which simplifies to

$$\lambda_1 = 4i$$

$$\lambda_2 = -4i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 4$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0(c_1 \cos(4x) + c_2 \sin(4x))$$

Or

$$y = c_1 \cos(4x) + c_2 \sin(4x)$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(4x) + c_2 \sin(4x) \tag{1}$$




Figure 79: Slope field plot

Verification of solutions

$$y = c_1 \cos(4x) + c_2 \sin(4x)$$

Verified OK.

4.3.2 Solving as second order ode can be made integrable ode

Multiplying the ode by y' gives

$$y'y'' + 16y'y = 0$$

Integrating the above w.r.t x gives

$$\int (y'y'' + 16y'y) dx = 0$$

$$\frac{y'^2}{2} + 8y^2 = c_2$$

Which is now solved for y . Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \sqrt{-16y^2 + 2c_1} \quad (1)$$

$$y' = -\sqrt{-16y^2 + 2c_1} \quad (2)$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int \frac{1}{\sqrt{-16y^2 + 2c_1}} dy = \int dx$$
$$\frac{\arctan\left(\frac{4y}{\sqrt{-16y^2 + 2c_1}}\right)}{4} = c_2 + x$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{1}{\sqrt{-16y^2 + 2c_1}} dy = \int dx$$
$$-\frac{\arctan\left(\frac{4y}{\sqrt{-16y^2 + 2c_1}}\right)}{4} = x + c_3$$

Summary

The solution(s) found are the following

$$\frac{\arctan\left(\frac{4y}{\sqrt{-16y^2 + 2c_1}}\right)}{4} = c_2 + x \quad (1)$$

$$-\frac{\arctan\left(\frac{4y}{\sqrt{-16y^2 + 2c_1}}\right)}{4} = x + c_3 \quad (2)$$




Figure 80: Slope field plot

Verification of solutions

$$\frac{\arctan\left(\frac{4y}{\sqrt{-16y^2+2c_1}}\right)}{4} = c_2 + x$$

Verified OK.

$$-\frac{\arctan\left(\frac{4y}{\sqrt{-16y^2+2c_1}}\right)}{4} = x + c_3$$

Verified OK.

4.3.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 16y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 16 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-16}{1} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -16 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -16z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 77: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -16$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(4x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}
 y_1 &= z_1 \\
 &= \cos(4x)
 \end{aligned}$$

Which simplifies to

$$y_1 = \cos(4x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(4x) \int \frac{1}{\cos(4x)^2} dx \\ &= \cos(4x) \left(\frac{\tan(4x)}{4} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1(\cos(4x)) + c_2 \left(\cos(4x) \left(\frac{\tan(4x)}{4} \right) \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(4x) + \frac{c_2 \sin(4x)}{4} \tag{1}$$




Figure 81: Slope field plot

Verification of solutions

$$y = c_1 \cos(4x) + \frac{c_2 \sin(4x)}{4}$$

Verified OK.

4.3.4 Maple step by step solution

Let's solve

$$y'' + 16y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 + 16 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-64})}{2}$$

- Roots of the characteristic polynomial

$$r = (-4I, 4I)$$

- 1st solution of the ODE
 $y_1(x) = \cos(4x)$
- 2nd solution of the ODE
 $y_2(x) = \sin(4x)$
- General solution of the ODE
 $y = c_1 y_1(x) + c_2 y_2(x)$
- Substitute in solutions
 $y = c_1 \cos(4x) + c_2 \sin(4x)$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)+16*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 \sin(4x) + c_2 \cos(4x)$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 20

```
DSolve[y''[x]+16*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos(4x) + c_2 \sin(4x)$$

4.4 problem 1(d)

4.4.1	Solving as second order ode quadrature ode	364
4.4.2	Solving as second order linear constant coeff ode	365
4.4.3	Solving as second order ode can be made integrable ode	367
4.4.4	Solving as second order integrable as is ode	368
4.4.5	Solving as second order ode missing y ode	369
4.4.6	Solving using Kovacic algorithm	371
4.4.7	Solving as exact linear second order ode ode	374
4.4.8	Maple step by step solution	376

Internal problem ID [5947]

Internal file name [OUTPUT/5195_Sunday_June_05_2022_03_27_13_PM_90127351/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 52

Problem number: 1(d).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "exact linear second order ode", "second_order_integrable_as_is", "second_order_ode_missing_y", "second_order_ode_quadrature", "second_order_linear_constant_coeff", "second_order_ode_can_be_made_integrable"

Maple gives the following as the ode type

```
[[_2nd_order , _quadrature]]
```

$$y'' = 0$$

4.4.1 Solving as second order ode quadrature ode

Integrating twice gives the solution

$$y = c_1x + c_2$$

Summary

The solution(s) found are the following

$$y = c_1x + c_2 \tag{1}$$




Figure 82: Slope field plot

Verification of solutions

$$y = c_1x + c_2$$

Verified OK.

4.4.2 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 0$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 0$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{(0)^2 - (4)(1)(0)} \\ &= 0\end{aligned}$$

Hence this is the case of a double root $\lambda_{1,2} = 0$. Therefore the solution is

$$y = c_1 1 + c_2 x \quad (1)$$

Summary

The solution(s) found are the following

$$y = c_2 x + c_1 \quad (1)$$




Figure 83: Slope field plot

Verification of solutions

$$y = c_2 x + c_1$$

Verified OK.

4.4.3 Solving as second order ode can be made integrable ode

Multiplying the ode by y' gives

$$y'y'' = 0$$

Integrating the above w.r.t x gives

$$\int y'y'' dx = 0$$
$$\frac{y'^2}{2} = c_2$$

Which is now solved for y . Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \sqrt{2} \sqrt{c_1} \quad (1)$$

$$y' = -\sqrt{2} \sqrt{c_1} \quad (2)$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$y = \int \sqrt{2} \sqrt{c_1} dx$$
$$= x\sqrt{2} \sqrt{c_1} + c_2$$

Solving equation (2)

Integrating both sides gives

$$y = \int -\sqrt{2} \sqrt{c_1} dx$$
$$= -x\sqrt{2} \sqrt{c_1} + c_3$$

Summary

The solution(s) found are the following

$$y = x\sqrt{2} \sqrt{c_1} + c_2 \quad (1)$$

$$y = -x\sqrt{2} \sqrt{c_1} + c_3 \quad (2)$$




Figure 84: Slope field plot

Verification of solutions

$$y = x\sqrt{2}\sqrt{c_1} + c_2$$

Verified OK.

$$y = -x\sqrt{2}\sqrt{c_1} + c_3$$

Verified OK.

4.4.4 Solving as second order integrable as is ode

Integrating both sides of the ODE w.r.t x gives

$$\int y'' dx = 0$$

$$y' = c_1$$

Which is now solved for y . Integrating both sides gives

$$y = \int c_1 dx$$

$$= c_1x + c_2$$

Summary

The solution(s) found are the following

$$y = c_1x + c_2 \quad (1)$$




Figure 85: Slope field plot

Verification of solutions

$$y = c_1x + c_2$$

Verified OK.

4.4.5 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x) = 0$$

Which is now solve for $p(x)$ as first order ode. Integrating both sides gives

$$\begin{aligned} p(x) &= \int 0 \, dx \\ &= c_1 \end{aligned}$$

Since $p = y'$ then the new first order ode to solve is

$$y' = c_1$$

Integrating both sides gives

$$\begin{aligned} y &= \int c_1 \, dx \\ &= c_1 x + c_2 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 x + c_2 \tag{1}$$




Figure 86: Slope field plot

Verification of solutions

$$y = c_1 x + c_2$$

Verified OK.

4.4.6 Solving using Kovacic algorithm

Writing the ode as

$$y'' = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 0 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 0 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 0 \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 79: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}y_1 &= z_1 \\ &= 1\end{aligned}$$

Which simplifies to

$$y_1 = 1$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned}y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= 1 \int \frac{1}{1} dx \\ &= 1(x)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1(1) + c_2(1(x))\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_2 x + c_1 \tag{1}$$




Figure 87: Slope field plot

Verification of solutions

$$y = c_2x + c_1$$

Verified OK.

4.4.7 Solving as exact linear second order ode ode

An ode of the form

$$p(x)y'' + q(x)y' + r(x)y = s(x)$$

is exact if

$$p''(x) - q'(x) + r(x) = 0 \tag{1}$$

For the given ode we have

$$\begin{aligned} p(x) &= 1 \\ q(x) &= 0 \\ r(x) &= 0 \\ s(x) &= 0 \end{aligned}$$

Hence

$$\begin{aligned}p''(x) &= 0 \\q'(x) &= 0\end{aligned}$$

Therefore (1) becomes

$$0 - (0) + (0) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(x) y' + (q(x) - p'(x)) y)' = s(x)$$

Integrating gives

$$p(x) y' + (q(x) - p'(x)) y = \int s(x) dx$$

Substituting the above values for p, q, r, s gives

$$y' = c_1$$

We now have a first order ode to solve which is

$$y' = c_1$$

Integrating both sides gives

$$\begin{aligned}y &= \int c_1 dx \\ &= c_1 x + c_2\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 x + c_2 \tag{1}$$




Figure 88: Slope field plot

Verification of solutions

$$y = c_1x + c_2$$

Verified OK.

4.4.8 Maple step by step solution

Let's solve

$$y'' = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{0})}{2}$$

- Roots of the characteristic polynomial

$$r = 0$$

- 1st solution of the ODE

$$y_1(x) = 1$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = x$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = c_2 x + c_1$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
dsolve(diff(y(x),x$2)=0,y(x), singsol=all)
```

$$y(x) = c_1 x + c_2$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 12

```
DSolve[y''[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 x + c_1$$

4.5 problem 1(e)

4.5.1	Solving as second order linear constant coeff ode	378
4.5.2	Solving using Kovacic algorithm	380
4.5.3	Maple step by step solution	384

Internal problem ID [5948]

Internal file name [OUTPUT/5196_Sunday_June_05_2022_03_27_14_PM_86840818/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 52

Problem number: 1(e).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' + 2iy' + y = 0$$

4.5.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 2i, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 2i\lambda e^{\lambda x} + e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 2i\lambda + 1 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 2i, C = 1$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{-2i}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{2i^2 - (4)(1)(1)} \\ &= -i \pm i\sqrt{2}\end{aligned}$$

Hence

$$\begin{aligned}\lambda_1 &= -i + i\sqrt{2} \\ \lambda_2 &= -i - i\sqrt{2}\end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= i(\sqrt{2} - 1) \\ \lambda_2 &= -i(1 + \sqrt{2})\end{aligned}$$

The roots are complex but they are not conjugate of each others. Hence simplification using Euler relation is not possible here. Therefore the final solution is

$$\begin{aligned}y &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ &= c_1 e^{i(\sqrt{2}-1)x} + c_2 e^{-i(1+\sqrt{2})x}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{i(\sqrt{2}-1)x} + c_2 e^{-i(1+\sqrt{2})x} \quad (1)$$




Figure 89: Slope field plot

Verification of solutions

$$y = c_1 e^{i(\sqrt{2}-1)x} + c_2 e^{-i(1+\sqrt{2})x}$$

Verified OK.

4.5.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 2iy' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 2i \\ C &= 1 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-2}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -2 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -2z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 81: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -2$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x\sqrt{2})$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{2i}{1} dx} \\ &= z_1 e^{-ix} \\ &= z_1 (e^{-ix}) \end{aligned}$$

Which simplifies to

$$y_1 = \cos(x\sqrt{2}) e^{-ix}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{2i}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-2ix}}{(y_1)^2} dx \\ &= y_1 \left(\frac{\sqrt{2} \tan(x\sqrt{2})}{2} \right) \end{aligned}$$

Therefore the solution is

$$y = c_1 y_1 + c_2 y_2$$

$$= c_1 \left(\cos(x\sqrt{2}) e^{-ix} \right) + c_2 \left(\cos(x\sqrt{2}) e^{-ix} \left(\frac{\sqrt{2} \tan(x\sqrt{2})}{2} \right) \right)$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(x\sqrt{2}) e^{-ix} + \frac{c_2 e^{-ix} \sqrt{2} \sin(x\sqrt{2})}{2} \quad (1)$$




Figure 90: Slope field plot

Verification of solutions

$$y = c_1 \cos(x\sqrt{2}) e^{-ix} + \frac{c_2 e^{-ix} \sqrt{2} \sin(x\sqrt{2})}{2}$$

Verified OK.

4.5.3 Maple step by step solution

Let's solve

$$y'' + 2Iy' + y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 + 2Ir + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-2I) \pm (\sqrt{-8})}{2}$$

- Roots of the characteristic polynomial

$$r = (-I - I\sqrt{2}, -I + I\sqrt{2})$$

- 1st solution of the ODE

$$y_1(x) = \cos((1 + \sqrt{2})x)$$

- 2nd solution of the ODE

$$y_2(x) = \sin((1 + \sqrt{2})x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = c_1 \cos((1 + \sqrt{2})x) + c_2 \sin((1 + \sqrt{2})x)$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve(diff(y(x),x$2)+2*I*diff(y(x),x)+y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{-ix} \left(c_1 \sin(\sqrt{2}x) + c_2 \cos(\sqrt{2}x) \right)$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 38

```
DSolve[y''[x]+2*I*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-i(1+\sqrt{2})x} \left(c_2 e^{2i\sqrt{2}x} + c_1 \right)$$

4.6 problem 1(f)

4.6.1	Solving as second order linear constant coeff ode	386
4.6.2	Solving using Kovacic algorithm	388
4.6.3	Maple step by step solution	392

Internal problem ID [5949]

Internal file name [OUTPUT/5197_Sunday_June_05_2022_03_27_15_PM_94509966/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 52

Problem number: 1(f).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' - 4y' + 5y = 0$$

4.6.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -4, C = 5$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 4\lambda e^{\lambda x} + 5 e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 4\lambda + 5 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -4, C = 5$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{4}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-4^2 - (4)(1)(5)} \\ &= 2 \pm i\end{aligned}$$

Hence

$$\lambda_1 = 2 + i$$

$$\lambda_2 = 2 - i$$

Which simplifies to

$$\lambda_1 = 2 + i$$

$$\lambda_2 = 2 - i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 2$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^{2x}(\cos(x) c_1 + c_2 \sin(x))$$

Summary

The solution(s) found are the following

$$y = e^{2x}(\cos(x) c_1 + c_2 \sin(x)) \tag{1}$$




Figure 91: Slope field plot

Verification of solutions

$$y = e^{2x}(\cos(x) c_1 + c_2 \sin(x))$$

Verified OK.

4.6.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 4y' + 5y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -4 \\ C &= 5 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 83: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-4}{1} dx} \\ &= z_1 e^{2x} \\ &= z_1 (e^{2x}) \end{aligned}$$

Which simplifies to

$$y_1 = \cos(x) e^{2x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-4}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{4x}}{(y_1)^2} dx \\ &= y_1 (\tan(x)) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(x) e^{2x}) + c_2 (\cos(x) e^{2x} (\tan(x)))\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(x) e^{2x} + c_2 \sin(x) e^{2x} \quad (1)$$




Figure 92: Slope field plot

Verification of solutions

$$y = c_1 \cos(x) e^{2x} + c_2 \sin(x) e^{2x}$$

Verified OK.

4.6.3 Maple step by step solution

Let's solve

$$y'' - 4y' + 5y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 - 4r + 5 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{4 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (2 - I, 2 + I)$$

- 1st solution of the ODE

$$y_1(x) = \cos(x) e^{2x}$$

- 2nd solution of the ODE

$$y_2(x) = \sin(x) e^{2x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = c_1 \cos(x) e^{2x} + c_2 \sin(x) e^{2x}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(diff(y(x),x$2)-4*diff(y(x),x)+5*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{2x}(c_1 \sin(x) + \cos(x) c_2)$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 22

```
DSolve[y''[x]-4*y'[x]+5*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{2x}(c_2 \cos(x) + c_1 \sin(x))$$

4.7 problem 1(g)

4.7.1 Solving as second order linear constant coeff ode	394
4.7.2 Solving using Kovacic algorithm	396
4.7.3 Maple step by step solution	400

Internal problem ID [5950]

Internal file name [OUTPUT/5198_Sunday_June_05_2022_03_27_16_PM_19777031/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 52

Problem number: 1(g).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"kovacic", "second_order_linear_constant_coeff"**

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' + (-1 + 3i)y' - 3iy = 0$$

4.7.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -1 + 3i, C = -3i$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + (-1 + 3i)\lambda e^{\lambda x} - 3ie^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + (-1 + 3i)\lambda - 3i = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -1 + 3i, C = -3i$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{1 - 3i}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-1 + 3i^2 - (4)(1)(-3i)} \\ &= \frac{1}{2} - \frac{3i}{2} \pm \frac{1}{2} + \frac{3i}{2}\end{aligned}$$

Hence

$$\begin{aligned}\lambda_1 &= \frac{1}{2} - \frac{3i}{2} + \frac{1}{2} + \frac{3i}{2} \\ \lambda_2 &= \frac{1}{2} - \frac{3i}{2} - \frac{1}{2} + \frac{3i}{2}\end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= 1 \\ \lambda_2 &= -3i\end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned}y &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ y &= c_1 e^{(1)x} + c_2 e^{(-3i)x}\end{aligned}$$

Or

$$y = c_1 e^x + e^{-3ix} c_2$$

Summary

The solution(s) found are the following

$$y = c_1 e^x + e^{-3ix} c_2 \tag{1}$$




Figure 93: Slope field plot

Verification of solutions

$$y = c_1 e^x + e^{-3ix} c_2$$

Verified OK.

4.7.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + (-1 + 3i)y' - 3iy = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -1 + 3i \\ C &= -3i \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-4 + 3i}{2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -4 + 3i \\ t &= 2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(-2 + \frac{3i}{2}\right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 85: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -2 + \frac{3i}{2}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{\left(\frac{1}{2} + \frac{3i}{2}\right)x}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-1+3i}{1} dx} \\ &= z_1 e^{\left(\frac{1}{2} - \frac{3i}{2}\right)x} \\ &= z_1 \left(e^{\left(\frac{1}{2} - \frac{3i}{2}\right)x} \right) \end{aligned}$$

Which simplifies to

$$y_1 = e^x$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-1+3i}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{(1-3i)x}}{(y_1)^2} dx \\ &= y_1 \left(\left(-\frac{1}{10} + \frac{3i}{10} \right) e^{(-1-3i)x} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1(e^x) + c_2\left(e^x\left(\left(-\frac{1}{10} + \frac{3i}{10}\right)e^{(-1-3i)x}\right)\right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^x + \left(-\frac{1}{10} + \frac{3i}{10}\right) c_2 e^{-3ix} \quad (1)$$




Figure 94: Slope field plot

Verification of solutions

$$y = c_1 e^x + \left(-\frac{1}{10} + \frac{3i}{10}\right) c_2 e^{-3ix}$$

Verified OK.

4.7.3 Maple step by step solution

Let's solve

$$y'' + (-1 + 3I)y' - 3Iy = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 + (-1 + 3I)r - 3I = 0$$

- Factor the characteristic polynomial

$$(r - 1)(r + 3I) = 0$$

- Roots of the characteristic polynomial

$$r = (1, -3I)$$

- 1st solution of the ODE

$$y_1(x) = e^x$$

- 2nd solution of the ODE

$$y_2(x) = 0$$

- General solution of the ODE

$$y = c_1y_1(x) + c_2y_2(x)$$

- Substitute in solutions

$$y = c_1e^x$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve(diff(y(x),x$2)+(3*I-1)*diff(y(x),x)-3*I*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^x c_1 + c_2 e^{-3ix}$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 22

```
DSolve[y''[x]+(3*I-1)*y'[x]-3*I*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-3ix} + c_2 e^x$$

4.8 problem 2(a)

4.8.1	Existence and uniqueness analysis	402
4.8.2	Solving as second order linear constant coeff ode	403
4.8.3	Solving using Kovacic algorithm	405
4.8.4	Maple step by step solution	410

Internal problem ID [5951]

Internal file name [OUTPUT/5199_Sunday_June_05_2022_03_27_18_PM_96449350/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 52

Problem number: 2(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"kovacic", "second_order_linear_constant_coeff"**

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' + y' - 6y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 0]$$

4.8.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = 1$$

$$q(x) = -6$$

$$F = 0$$

Hence the ode is

$$y'' + y' - 6y = 0$$

The domain of $p(x) = 1$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = -6$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

4.8.2 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 1, C = -6$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + \lambda e^{\lambda x} - 6 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + \lambda - 6 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 1, C = -6$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{1^2 - (4)(1)(-6)} \\ &= -\frac{1}{2} \pm \frac{5}{2} \end{aligned}$$

Hence

$$\lambda_1 = -\frac{1}{2} + \frac{5}{2}$$

$$\lambda_2 = -\frac{1}{2} - \frac{5}{2}$$

Which simplifies to

$$\lambda_1 = 2$$

$$\lambda_2 = -3$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(2)x} + c_2 e^{(-3)x}$$

Or

$$y = c_1 e^{2x} + c_2 e^{-3x}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 e^{2x} + c_2 e^{-3x} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 0$ in the above gives

$$1 = c_1 + c_2 \quad (1A)$$

Taking derivative of the solution gives

$$y' = 2c_1 e^{2x} - 3c_2 e^{-3x}$$

substituting $y' = 0$ and $x = 0$ in the above gives

$$0 = 2c_1 - 3c_2 \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = \frac{3}{5}$$

$$c_2 = \frac{2}{5}$$


Substituting these values back in above solution results in

$$y = \frac{3 e^{2x}}{5} + \frac{2 e^{-3x}}{5}$$


Summary

The solution(s) found are the following

$$y = \frac{3e^{2x}}{5} + \frac{2e^{-3x}}{5} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{3e^{2x}}{5} + \frac{2e^{-3x}}{5}$$

Verified OK.

4.8.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y' - 6y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 1$$

$$C = -6$$

(3)

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{25}{4} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 25 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \frac{25z(x)}{4} \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 87: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{25}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{5x}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}
 y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\
 &= z_1 e^{-\int \frac{1}{2} \frac{1}{1} dx} \\
 &= z_1 e^{-\frac{x}{2}} \\
 &= z_1 \left(e^{-\frac{x}{2}} \right)
 \end{aligned}$$

Which simplifies to

$$y_1 = e^{-3x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{1}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-x}}{(y_1)^2} dx \\ &= y_1 \left(\frac{e^{5x}}{5} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-3x}) + c_2 \left(e^{-3x} \left(\frac{e^{5x}}{5} \right) \right) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 e^{-3x} + \frac{c_2 e^{2x}}{5} \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 0$ in the above gives

$$1 = c_1 + \frac{c_2}{5} \tag{1A}$$

Taking derivative of the solution gives

$$y' = -3c_1 e^{-3x} + \frac{2c_2 e^{2x}}{5}$$

substituting $y' = 0$ and $x = 0$ in the above gives

$$0 = -3c_1 + \frac{2c_2}{5} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = \frac{2}{5}$$
$$c_2 = 3$$


Substituting these values back in above solution results in

$$y = \frac{3e^{2x}}{5} + \frac{2e^{-3x}}{5}$$


Summary

The solution(s) found are the following

$$y = \frac{3e^{2x}}{5} + \frac{2e^{-3x}}{5} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{3e^{2x}}{5} + \frac{2e^{-3x}}{5}$$

Verified OK.

4.8.4 Maple step by step solution

Let's solve

$$\left[y'' + y' - 6y = 0, y(0) = 1, y' \Big|_{\{x=0\}} = 0 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 + r - 6 = 0$$

- Factor the characteristic polynomial

$$(r + 3)(r - 2) = 0$$

- Roots of the characteristic polynomial

$$r = (-3, 2)$$

- 1st solution of the ODE

$$y_1(x) = e^{-3x}$$

- 2nd solution of the ODE

$$y_2(x) = e^{2x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = c_1 e^{-3x} + c_2 e^{2x}$$

- Check validity of solution $y = c_1 e^{-3x} + c_2 e^{2x}$

- Use initial condition $y(0) = 1$

$$1 = c_1 + c_2$$

- Compute derivative of the solution

$$y' = -3c_1 e^{-3x} + 2c_2 e^{2x}$$

- Use the initial condition $y' \Big|_{\{x=0\}} = 0$

$$0 = -3c_1 + 2c_2$$

- Solve for c_1 and c_2

$$\left\{ c_1 = \frac{2}{5}, c_2 = \frac{3}{5} \right\}$$

- Substitute constant values into general solution and simplify

$$y = \frac{(3e^{5x}+2)e^{-3x}}{5}$$

- Solution to the IVP

$$y = \frac{(3e^{5x}+2)e^{-3x}}{5}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 18

```
dsolve([diff(y(x),x$2)+diff(y(x),x)-6*y(x)=0,y(0) = 1, D(y)(0) = 0],y(x), singsol=all)
```

$$y(x) = \frac{(3e^{5x} + 2)e^{-3x}}{5}$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 23

```
DSolve[{y''[x]+y'[x]-6*y[x]==0,{y[0]==1,y'[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{5}e^{-3x}(3e^{5x} + 2)$$

4.9 problem 2(b)

4.9.1	Existence and uniqueness analysis	412
4.9.2	Solving as second order linear constant coeff ode	413
4.9.3	Solving using Kovacic algorithm	415
4.9.4	Maple step by step solution	420

Internal problem ID [5952]

Internal file name [OUTPUT/5200_Sunday_June_05_2022_03_27_19_PM_9050429/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 52

Problem number: 2(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"kovacic", "second_order_linear_constant_coeff"**

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x]]
```

$$y'' + y' - 6y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 1]$$

4.9.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = 1$$

$$q(x) = -6$$

$$F = 0$$

Hence the ode is

$$y'' + y' - 6y = 0$$

The domain of $p(x) = 1$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = -6$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

4.9.2 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 1, C = -6$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + \lambda e^{\lambda x} - 6 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + \lambda - 6 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 1, C = -6$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{1^2 - (4)(1)(-6)} \\ &= -\frac{1}{2} \pm \frac{5}{2} \end{aligned}$$

Hence

$$\lambda_1 = -\frac{1}{2} + \frac{5}{2}$$

$$\lambda_2 = -\frac{1}{2} - \frac{5}{2}$$

Which simplifies to

$$\lambda_1 = 2$$

$$\lambda_2 = -3$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(2)x} + c_2 e^{(-3)x}$$

Or

$$y = c_1 e^{2x} + c_2 e^{-3x}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 e^{2x} + c_2 e^{-3x} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_1 + c_2 \quad (1A)$$

Taking derivative of the solution gives

$$y' = 2c_1 e^{2x} - 3c_2 e^{-3x}$$

substituting $y' = 1$ and $x = 0$ in the above gives

$$1 = 2c_1 - 3c_2 \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = \frac{1}{5}$$

$$c_2 = -\frac{1}{5}$$

Substituting these values back in above solution results in

$$y = \frac{e^{2x}}{5} - \frac{e^{-3x}}{5}$$


Which simplifies to

$$y = \frac{(e^{5x} - 1)e^{-3x}}{5}$$


Summary

The solution(s) found are the following

$$y = \frac{(e^{5x} - 1)e^{-3x}}{5} \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{(e^{5x} - 1)e^{-3x}}{5}$$

Verified OK.

4.9.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y' - 6y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= 1 \\B &= 1 \\C &= -6\end{aligned}\tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x)\tag{4}$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\&= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned}\tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{25}{4}\tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= 25 \\t &= 4\end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \frac{25z(x)}{4}\tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 89: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{25}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{5x}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}
 y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\
 &= z_1 e^{-\int \frac{1}{2} \frac{1}{1} dx} \\
 &= z_1 e^{-\frac{x}{2}} \\
 &= z_1 \left(e^{-\frac{x}{2}} \right)
 \end{aligned}$$

Which simplifies to

$$y_1 = e^{-3x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{1}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-x}}{(y_1)^2} dx \\ &= y_1 \left(\frac{e^{5x}}{5} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-3x}) + c_2 \left(e^{-3x} \left(\frac{e^{5x}}{5} \right) \right) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 e^{-3x} + \frac{c_2 e^{2x}}{5} \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_1 + \frac{c_2}{5} \tag{1A}$$

Taking derivative of the solution gives

$$y' = -3c_1 e^{-3x} + \frac{2c_2 e^{2x}}{5}$$

substituting $y' = 1$ and $x = 0$ in the above gives

$$1 = -3c_1 + \frac{2c_2}{5} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = -\frac{1}{5}$$
$$c_2 = 1$$

Substituting these values back in above solution results in

$$y = \frac{e^{2x}}{5} - \frac{e^{-3x}}{5}$$


Which simplifies to

$$y = \frac{(e^{5x} - 1)e^{-3x}}{5}$$


Summary

The solution(s) found are the following

$$y = \frac{(e^{5x} - 1)e^{-3x}}{5} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{(e^{5x} - 1)e^{-3x}}{5}$$

Verified OK.

4.9.4 Maple step by step solution

Let's solve

$$\left[y'' + y' - 6y = 0, y(0) = 0, y' \Big|_{\{x=0\}} = 1 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 + r - 6 = 0$$

- Factor the characteristic polynomial

$$(r + 3)(r - 2) = 0$$

- Roots of the characteristic polynomial

$$r = (-3, 2)$$

- 1st solution of the ODE

$$y_1(x) = e^{-3x}$$

- 2nd solution of the ODE

$$y_2(x) = e^{2x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = c_1 e^{-3x} + c_2 e^{2x}$$

- Check validity of solution $y = c_1 e^{-3x} + c_2 e^{2x}$

- Use initial condition $y(0) = 0$

$$0 = c_1 + c_2$$

- Compute derivative of the solution

$$y' = -3c_1 e^{-3x} + 2c_2 e^{2x}$$

- Use the initial condition $y' \Big|_{\{x=0\}} = 1$

$$1 = -3c_1 + 2c_2$$

- Solve for c_1 and c_2

$$\left\{ c_1 = -\frac{1}{5}, c_2 = \frac{1}{5} \right\}$$

- Substitute constant values into general solution and simplify

$$y = \frac{(e^{5x}-1)e^{-3x}}{5}$$

- Solution to the IVP

$$y = \frac{(e^{5x}-1)e^{-3x}}{5}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 16

```
dsolve([diff(y(x),x$2)+diff(y(x),x)-6*y(x)=0,y(0) = 0, D(y)(0) = 1],y(x), singsol=all)
```

$$y(x) = \frac{(e^{5x} - 1)e^{-3x}}{5}$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 21

```
DSolve[{y''[x]+y'[x]-6*y[x]==0,{y[0]==0,y'[0]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{5}e^{-3x}(e^{5x} - 1)$$

4.10 problem 3(a)

4.10.1 Solving as second order linear constant coeff ode	422
4.10.2 Solving as second order ode can be made integrable ode	425
4.10.3 Solving using Kovacic algorithm	426
4.10.4 Maple step by step solution	430

Internal problem ID [5953]

Internal file name [OUTPUT/5201_Sunday_June_05_2022_03_27_20_PM_43362679/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 52

Problem number: 3(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "second_order_ode_can_be_made_integrable"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x]]
```

$$y'' + y = 0$$

With initial conditions

$$\left[y(0) = 1, y\left(\frac{\pi}{2}\right) = 2 \right]$$

4.10.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +i \\ \lambda_2 &= -i \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= i \\ \lambda_2 &= -i \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (\cos(x) c_1 + c_2 \sin(x))$$

Or

$$y = \cos(x) c_1 + c_2 \sin(x)$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = \cos(x) c_1 + c_2 \sin(x) \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 2$ and $x = \frac{\pi}{2}$ in the above gives

$$2 = c_2 \quad (1A)$$

substituting $y = 1$ and $x = 0$ in the above gives

$$1 = c_1 \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 1$$

$$c_2 = 2$$

Substituting these values back in above solution results in

$$y = 2 \sin(x) + \cos(x)$$

Summary

The solution(s) found are the following

$$y = 2 \sin(x) + \cos(x) \quad (1)$$




Figure 99: Solution plot

Verification of solutions

$$y = 2 \sin(x) + \cos(x)$$

Verified OK.

4.10.2 Solving as second order ode can be made integrable ode

Multiplying the ode by y' gives

$$y'y'' + y'y = 0$$

Integrating the above w.r.t x gives

$$\int (y'y'' + y'y) dx = 0$$
$$\frac{y'^2}{2} + \frac{y^2}{2} = c_2$$

Which is now solved for y . Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \sqrt{-y^2 + 2c_1} \quad (1)$$

$$y' = -\sqrt{-y^2 + 2c_1} \quad (2)$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int \frac{1}{\sqrt{-y^2 + 2c_1}} dy = \int dx$$
$$\arctan\left(\frac{y}{\sqrt{-y^2 + 2c_1}}\right) = c_2 + x$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{1}{\sqrt{-y^2 + 2c_1}} dy = \int dx$$
$$-\arctan\left(\frac{y}{\sqrt{-y^2 + 2c_1}}\right) = x + c_3$$

Initial conditions are used to solve for the constants of integration.

Looking at the First solution

$$\arctan\left(\frac{y}{\sqrt{-y^2 + 2c_1}}\right) = c_2 + x \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 2$ and $x = \frac{\pi}{2}$ in the above gives

$$\arctan\left(\frac{2}{\sqrt{-4 + 2c_1}}\right) = c_2 + \frac{\pi}{2} \quad (1A)$$

substituting $y = 1$ and $x = 0$ in the above gives

$$\arctan\left(\frac{1}{\sqrt{-1 + 2c_1}}\right) = c_2 \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. There is no solution for the constants of integrations. This solution is removed.

Looking at the Second solution

$$-\arctan\left(\frac{y}{\sqrt{-y^2 + 2c_1}}\right) = x + c_3 \quad (2)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 2$ and $x = \frac{\pi}{2}$ in the above gives

$$-\arctan\left(\frac{2}{\sqrt{-4 + 2c_1}}\right) = \frac{\pi}{2} + c_3 \quad (1A)$$

substituting $y = 1$ and $x = 0$ in the above gives

$$-\arctan\left(\frac{1}{\sqrt{-1 + 2c_1}}\right) = c_3 \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_3\}$. There is no solution for the constants of integrations. This solution is removed.

Verification of solutions N/A

4.10.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 1 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 91: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}
 y_1 &= z_1 \\
 &= \cos(x)
 \end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x)) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(x)) + c_2 (\cos(x) (\tan(x))) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = \cos(x) c_1 + c_2 \sin(x) \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 2$ and $x = \frac{\pi}{2}$ in the above gives

$$2 = c_2 \tag{1A}$$

substituting $y = 1$ and $x = 0$ in the above gives

$$1 = c_1 \tag{2A}$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 1$$

$$c_2 = 2$$

Substituting these values back in above solution results in

$$y = 2 \sin(x) + \cos(x)$$

Summary

The solution(s) found are the following

$$y = 2 \sin(x) + \cos(x) \tag{1}$$




Figure 100: Solution plot

Verification of solutions

$$y = 2 \sin(x) + \cos(x)$$

Verified OK.

4.10.4 Maple step by step solution

Let's solve

$$[y'' + y = 0, y(0) = 1, y(\frac{\pi}{2}) = 2]$$

- Highest derivative means the order of the ODE is 2

y''

- Characteristic polynomial of ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm \sqrt{-4}}{2}$$
- Roots of the characteristic polynomial

$$r = (-I, I)$$
- 1st solution of the ODE

$$y_1(x) = \cos(x)$$
- 2nd solution of the ODE

$$y_2(x) = \sin(x)$$
- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$
- Substitute in solutions

$$y = \cos(x) c_1 + c_2 \sin(x)$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 11

```
dsolve([diff(y(x),x$2)+y(x)=0,y(0) = 1, y(1/2*Pi) = 2],y(x), singsol=all)
```

$$y(x) = 2 \sin(x) + \cos(x)$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 12

```
DSolve[{y'[x]+y[x]==0,{y[0]==1,y[Pi/2]==2}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2 \sin(x) + \cos(x)$$

4.11 problem 3(b)

4.11.1 Solving as second order linear constant coeff ode	432
4.11.2 Solving as second order ode can be made integrable ode	434
4.11.3 Solving using Kovacic algorithm	436
4.11.4 Maple step by step solution	439

Internal problem ID [5954]

Internal file name [OUTPUT/5202_Sunday_June_05_2022_03_27_21_PM_88438460/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 52

Problem number: 3(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "second_order_ode_can_be_made_integrable"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x]]
```

$$y'' + y = 0$$

With initial conditions

$$[y(0) = 0, y(\pi) = 0]$$

4.11.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +i \\ \lambda_2 &= -i \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= i \\ \lambda_2 &= -i \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (\cos(x) c_1 + c_2 \sin(x))$$

Or

$$y = \cos(x) c_1 + c_2 \sin(x)$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = \cos(x) c_1 + c_2 \sin(x) \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = \pi$ in the above gives

$$0 = -c_1 \quad (1A)$$

substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_1 \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 0$$

Substituting these values back in above solution results in

$$y = c_2 \sin(x)$$

Summary

The solution(s) found are the following

$$y = c_2 \sin(x) \quad (1)$$

Verification of solutions

$$y = c_2 \sin(x)$$

Verified OK.

4.11.2 Solving as second order ode can be made integrable ode

Multiplying the ode by y' gives

$$y'y'' + y'y = 0$$

Integrating the above w.r.t x gives

$$\int (y'y'' + y'y) dx = 0$$

$$\frac{y'^2}{2} + \frac{y^2}{2} = c_2$$

Which is now solved for y . Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \sqrt{-y^2 + 2c_1} \quad (1)$$

$$y' = -\sqrt{-y^2 + 2c_1} \quad (2)$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int \frac{1}{\sqrt{-y^2 + 2c_1}} dy = \int dx$$
$$\arctan\left(\frac{y}{\sqrt{-y^2 + 2c_1}}\right) = c_2 + x$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{1}{\sqrt{-y^2 + 2c_1}} dy = \int dx$$
$$-\arctan\left(\frac{y}{\sqrt{-y^2 + 2c_1}}\right) = x + c_3$$

Initial conditions are used to solve for the constants of integration.

Looking at the First solution

$$\arctan\left(\frac{y}{\sqrt{-y^2 + 2c_1}}\right) = c_2 + x \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = \pi$ in the above gives

$$0 = c_2 + \pi \quad (1A)$$

substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_2 \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. There is no solution for the constants of integrations. This solution is removed.

Looking at the Second solution

$$-\arctan\left(\frac{y}{\sqrt{-y^2 + 2c_1}}\right) = x + c_3 \quad (2)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = \pi$ in the above gives

$$0 = \pi + c_3 \quad (1A)$$

substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_3 \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_3\}$. There is no solution for the constants of integrations. This solution is removed.

Verification of solutions N/A

4.11.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 1 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \quad (6)$$

Comparing the above to (5) shows that

$$s = -1$$

$$t = 1$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 93: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned} y_1 &= z_1 \\ &= \cos(x) \end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x)) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(x)) + c_2 (\cos(x) (\tan(x))) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = \cos(x) c_1 + c_2 \sin(x) \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = \pi$ in the above gives

$$0 = -c_1 \quad (1A)$$

substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_1 \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 0$$

Substituting these values back in above solution results in

$$y = c_2 \sin(x)$$

Summary

The solution(s) found are the following

$$y = c_2 \sin(x) \quad (1)$$

Verification of solutions

$$y = c_2 \sin(x)$$

Verified OK.

4.11.4 Maple step by step solution

Let's solve

$$[y'' + y = 0, y(0) = 0, y(\pi) = 0]$$

- Highest derivative means the order of the ODE is 2
 y''
- Characteristic polynomial of ODE
 $r^2 + 1 = 0$
- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (-I, I)$$

- 1st solution of the ODE

$$y_1(x) = \cos(x)$$

- 2nd solution of the ODE

$$y_2(x) = \sin(x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = \cos(x) c_1 + c_2 \sin(x)$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 8

```
dsolve([diff(y(x),x$2)+y(x)=0,y(0) = 0, y(Pi) = 0],y(x), singsol=all)
```

$$y(x) = c_1 \sin(x)$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 10

```
DSolve[{y'[x]+y[x]==0,{y[0]==0,y[Pi]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \sin(x)$$

4.12 problem 3(c)

4.12.1 Solving as second order linear constant coeff ode	441
4.12.2 Solving as second order ode can be made integrable ode	443
4.12.3 Solving using Kovacic algorithm	445
4.12.4 Maple step by step solution	449

Internal problem ID [5955]

Internal file name [OUTPUT/5203_Sunday_June_05_2022_03_27_23_PM_76068316/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 52

Problem number: 3(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "second_order_ode_can_be_made_integrable"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x]]
```

$$y'' + y = 0$$

With initial conditions

$$\left[y(0) = 0, y' \left(\frac{\pi}{2} \right) = 0 \right]$$

4.12.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +i \\ \lambda_2 &= -i \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= i \\ \lambda_2 &= -i \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (\cos(x) c_1 + c_2 \sin(x))$$

Or

$$y = \cos(x) c_1 + c_2 \sin(x)$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = \cos(x) c_1 + c_2 \sin(x) \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_1 \tag{1A}$$

Taking derivative of the solution gives

$$y' = -\sin(x) c_1 + c_2 \cos(x)$$

substituting $y' = 0$ and $x = \frac{\pi}{2}$ in the above gives

$$0 = -c_1 \tag{2A}$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 0$$

Substituting these values back in above solution results in

$$y = c_2 \sin(x)$$

Summary

The solution(s) found are the following

$$y = c_2 \sin(x) \tag{1}$$

Verification of solutions

$$y = c_2 \sin(x)$$

Verified OK.

4.12.2 Solving as second order ode can be made integrable ode

Multiplying the ode by y' gives

$$y'y'' + y'y' = 0$$

Integrating the above w.r.t x gives

$$\int (y'y'' + y'y') dx = 0$$

$$\frac{y'^2}{2} + \frac{y^2}{2} = c_2$$

Which is now solved for y . Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \sqrt{-y^2 + 2c_1} \quad (1)$$

$$y' = -\sqrt{-y^2 + 2c_1} \quad (2)$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int \frac{1}{\sqrt{-y^2 + 2c_1}} dy = \int dx$$

$$\arctan\left(\frac{y}{\sqrt{-y^2 + 2c_1}}\right) = c_2 + x$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{1}{\sqrt{-y^2 + 2c_1}} dy = \int dx$$

$$-\arctan\left(\frac{y}{\sqrt{-y^2 + 2c_1}}\right) = x + c_3$$

Initial conditions are used to solve for the constants of integration.

Looking at the First solution

$$\arctan\left(\frac{y}{\sqrt{-y^2 + 2c_1}}\right) = c_2 + x \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_2 \quad (1A)$$

Taking derivative of the solution gives

$$y' = (\tan(c_2 + x)^2 + 1) \sqrt{2} \sqrt{\frac{c_1}{\tan(c_2 + x)^2 + 1}} - \frac{\tan(c_2 + x)^2 \sqrt{2} c_1}{\sqrt{\frac{c_1}{\tan(c_2 + x)^2 + 1}} (\tan(c_2 + x)^2 + 1)}$$

substituting $y' = 0$ and $x = \frac{\pi}{2}$ in the above gives

$$0 = \frac{\sin(c_2)^2 c_1 \sqrt{2}}{\sqrt{c_1 \sin(c_2)^2}} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. There is no solution for the constants of integrations. This solution is removed.

Looking at the Second solution

$$-\arctan\left(\frac{y}{\sqrt{-y^2 + 2c_1}}\right) = x + c_3 \quad (2)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_3 \quad (1A)$$

Taking derivative of the solution gives

$$y' = -(\tan(x + c_3)^2 + 1) \sqrt{2} \sqrt{\frac{c_1}{\tan(x + c_3)^2 + 1}} + \frac{\tan(x + c_3)^2 \sqrt{2} c_1}{\sqrt{\frac{c_1}{\tan(x + c_3)^2 + 1}} (\tan(x + c_3)^2 + 1)}$$

substituting $y' = 0$ and $x = \frac{\pi}{2}$ in the above gives

$$0 = -\frac{\sin(c_3)^2 c_1 \sqrt{2}}{\sqrt{c_1 \sin(c_3)^2}} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_3\}$. There is no solution for the constants of integrations. This solution is removed.

Verification of solutions N/A

4.12.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 1 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 95: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}
 y_1 &= z_1 \\
 &= \cos(x)
 \end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x)) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(x)) + c_2 (\cos(x) (\tan(x))) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = \cos(x) c_1 + c_2 \sin(x) \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_1 \tag{1A}$$

Taking derivative of the solution gives

$$y' = -\sin(x) c_1 + c_2 \cos(x)$$

substituting $y' = 0$ and $x = \frac{\pi}{2}$ in the above gives

$$0 = -c_1 \tag{2A}$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 0$$

Substituting these values back in above solution results in

$$y = c_2 \sin(x)$$

Summary

The solution(s) found are the following

$$y = c_2 \sin(x) \tag{1}$$

Verification of solutions

$$y = c_2 \sin(x)$$

Verified OK.

4.12.4 Maple step by step solution

Let's solve

$$\left[y'' + y = 0, y(0) = 0, y' \Big|_{\{x=\frac{\pi}{2}\}} = 0 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (-I, I)$$

- 1st solution of the ODE

$$y_1(x) = \cos(x)$$

- 2nd solution of the ODE

$$y_2(x) = \sin(x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = \cos(x) c_1 + c_2 \sin(x)$$

- Check validity of solution $y = \cos(x) c_1 + c_2 \sin(x)$

- Use initial condition $y(0) = 0$

$$0 = c_1$$

- Compute derivative of the solution

$$y' = -\sin(x) c_1 + c_2 \cos(x)$$

- Use the initial condition $y' \Big|_{\{x=\frac{\pi}{2}\}} = 0$

$$0 = -c_1$$

- Solve for c_1 and c_2

$$\{c_1 = 0, c_2 = c_2\}$$

- Substitute constant values into general solution and simplify

$$y = c_2 \sin(x)$$

- Solution to the IVP

$$y = c_2 \sin(x)$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 8

```
dsolve([diff(y(x),x$2)+y(x)=0,y(0) = 0, D(y)(1/2*Pi) = 0],y(x), singsol=all)
```

$$y(x) = c_1 \sin(x)$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 10

```
DSolve[{y'[x]+y[x]==0,{y[0]==0,y'[Pi/2]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \sin(x)$$

4.13 problem 3(d)

4.13.1 Solving as second order linear constant coeff ode	452
4.13.2 Solving as second order ode can be made integrable ode	455
4.13.3 Solving using Kovacic algorithm	456
4.13.4 Maple step by step solution	460

Internal problem ID [5956]

Internal file name [OUTPUT/5204_Sunday_June_05_2022_03_27_24_PM_9716775/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 52

Problem number: 3(d).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "second_order_ode_can_be_made_integrable"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x]]
```

$$y'' + y = 0$$

With initial conditions

$$\left[y(0) = 0, y\left(\frac{\pi}{2}\right) = 0 \right]$$

4.13.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +i \\ \lambda_2 &= -i \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= i \\ \lambda_2 &= -i \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (\cos(x) c_1 + c_2 \sin(x))$$

Or

$$y = \cos(x) c_1 + c_2 \sin(x)$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = \cos(x) c_1 + c_2 \sin(x) \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = \frac{\pi}{2}$ in the above gives

$$0 = c_2 \tag{1A}$$

substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_1 \tag{2A}$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 0$$

$$c_2 = 0$$

Substituting these values back in above solution results in

$$y = 0$$

Summary

The solution(s) found are the following

$$y = 0 \tag{1}$$




Figure 101: Solution plot

Verification of solutions

$$y = 0$$

Verified OK.

4.13.2 Solving as second order ode can be made integrable ode

Multiplying the ode by y' gives

$$y'y'' + y'y = 0$$

Integrating the above w.r.t x gives

$$\int (y'y'' + y'y) dx = 0$$
$$\frac{y'^2}{2} + \frac{y^2}{2} = c_2$$

Which is now solved for y . Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \sqrt{-y^2 + 2c_1} \quad (1)$$

$$y' = -\sqrt{-y^2 + 2c_1} \quad (2)$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int \frac{1}{\sqrt{-y^2 + 2c_1}} dy = \int dx$$
$$\arctan\left(\frac{y}{\sqrt{-y^2 + 2c_1}}\right) = c_2 + x$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{1}{\sqrt{-y^2 + 2c_1}} dy = \int dx$$
$$-\arctan\left(\frac{y}{\sqrt{-y^2 + 2c_1}}\right) = x + c_3$$

Initial conditions are used to solve for the constants of integration.

Looking at the First solution

$$\arctan\left(\frac{y}{\sqrt{-y^2 + 2c_1}}\right) = c_2 + x \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = \frac{\pi}{2}$ in the above gives

$$0 = c_2 + \frac{\pi}{2} \quad (1A)$$

substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_2 \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. There is no solution for the constants of integrations. This solution is removed.

Looking at the Second solution

$$-\arctan\left(\frac{y}{\sqrt{-y^2 + 2c_1}}\right) = x + c_3 \quad (2)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = \frac{\pi}{2}$ in the above gives

$$0 = \frac{\pi}{2} + c_3 \quad (1A)$$

substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_3 \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_3\}$. There is no solution for the constants of integrations. This solution is removed.

Verification of solutions N/A

4.13.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 1 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 97: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}
 y_1 &= z_1 \\
 &= \cos(x)
 \end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x)) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(x)) + c_2 (\cos(x) (\tan(x))) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = \cos(x) c_1 + c_2 \sin(x) \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = \frac{\pi}{2}$ in the above gives

$$0 = c_2 \tag{1A}$$

substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_1 \tag{2A}$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 0$$

$$c_2 = 0$$

Substituting these values back in above solution results in

$$y = 0$$

Summary

The solution(s) found are the following

$$y = 0 \tag{1}$$




Figure 102: Solution plot

Verification of solutions

$$y = 0$$

Verified OK.

4.13.4 Maple step by step solution

Let's solve

$$[y'' + y = 0, y(0) = 0, y(\frac{\pi}{2}) = 0]$$

- Highest derivative means the order of the ODE is 2
 y''
- Characteristic polynomial of ODE
 $r^2 + 1 = 0$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$
- Roots of the characteristic polynomial

$$r = (-I, I)$$
- 1st solution of the ODE

$$y_1(x) = \cos(x)$$
- 2nd solution of the ODE

$$y_2(x) = \sin(x)$$
- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$
- Substitute in solutions

$$y = \cos(x) c_1 + c_2 \sin(x)$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 5

```
dsolve([diff(y(x),x$2)+y(x)=0,y(0) = 0, y(1/2*Pi) = 0],y(x), singsol=all)
```

$$y(x) = 0$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 6

```
DSolve[{y'[x]+y[x]==0,{y[0]==0,y[Pi/2]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 0$$

5 Chapter 2. Linear equations with constant coefficients. Page 59

5.1	problem 1(a)	463
5.2	problem 1(b)	473
5.3	problem 1(c)	483
5.4	problem 1(d)	491

5.1 problem 1(a)

5.1.1	Existence and uniqueness analysis	463
5.1.2	Solving as second order linear constant coeff ode	464
5.1.3	Solving using Kovacic algorithm	466
5.1.4	Maple step by step solution	471

Internal problem ID [5957]

Internal file name [OUTPUT/5205_Sunday_June_05_2022_03_27_25_PM_76562253/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 59

Problem number: 1(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"kovacic", "second_order_linear_constant_coeff"**

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x]]
```

$$y'' - 2y' - 3y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 1]$$

5.1.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = -2$$

$$q(x) = -3$$

$$F = 0$$

Hence the ode is

$$y'' - 2y' - 3y = 0$$

The domain of $p(x) = -2$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = -3$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

5.1.2 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -2, C = -3$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 2\lambda e^{\lambda x} - 3e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 2\lambda - 3 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -2, C = -3$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{2}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-2^2 - (4)(1)(-3)} \\ &= 1 \pm 2 \end{aligned}$$

Hence

$$\lambda_1 = 1 + 2$$

$$\lambda_2 = 1 - 2$$

Which simplifies to

$$\lambda_1 = 3$$

$$\lambda_2 = -1$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(3)x} + c_2 e^{(-1)x}$$

Or

$$y = e^{3x} c_1 + c_2 e^{-x}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = e^{3x} c_1 + c_2 e^{-x} \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_1 + c_2 \tag{1A}$$

Taking derivative of the solution gives

$$y' = 3 e^{3x} c_1 - c_2 e^{-x}$$

substituting $y' = 1$ and $x = 0$ in the above gives

$$1 = 3c_1 - c_2 \tag{2A}$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = \frac{1}{4}$$

$$c_2 = -\frac{1}{4}$$


Substituting these values back in above solution results in

$$y = \frac{e^{3x}}{4} - \frac{e^{-x}}{4}$$


Summary

The solution(s) found are the following

$$y = \frac{e^{3x}}{4} - \frac{e^{-x}}{4} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{e^{3x}}{4} - \frac{e^{-x}}{4}$$

Verified OK.

5.1.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 2y' - 3y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = -2$$

$$C = -3$$

(3)

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{4}{1} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 4 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 4z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 99: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 4$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-2x}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}
 y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\
 &= z_1 e^{-\int \frac{1}{2} \frac{-2}{1} dx} \\
 &= z_1 e^x \\
 &= z_1 (e^x)
 \end{aligned}$$

Which simplifies to

$$y_1 = e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{2}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{2x}}{(y_1)^2} dx \\ &= y_1 \left(\frac{e^{4x}}{4} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-x}) + c_2 \left(e^{-x} \left(\frac{e^{4x}}{4} \right) \right) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 e^{-x} + \frac{c_2 e^{3x}}{4} \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_1 + \frac{c_2}{4} \tag{1A}$$

Taking derivative of the solution gives

$$y' = -c_1 e^{-x} + \frac{3c_2 e^{3x}}{4}$$

substituting $y' = 1$ and $x = 0$ in the above gives

$$1 = -c_1 + \frac{3c_2}{4} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = -\frac{1}{4}$$

$$c_2 = 1$$


Substituting these values back in above solution results in

$$y = \frac{e^{3x}}{4} - \frac{e^{-x}}{4}$$


Summary

The solution(s) found are the following

$$y = \frac{e^{3x}}{4} - \frac{e^{-x}}{4} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{e^{3x}}{4} - \frac{e^{-x}}{4}$$

Verified OK.

5.1.4 Maple step by step solution

Let's solve

$$\left[y'' - 2y' - 3y = 0, y(0) = 0, y' \Big|_{\{x=0\}} = 1 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 - 2r - 3 = 0$$

- Factor the characteristic polynomial

$$(r + 1)(r - 3) = 0$$

- Roots of the characteristic polynomial

$$r = (-1, 3)$$

- 1st solution of the ODE

$$y_1(x) = e^{-x}$$

- 2nd solution of the ODE

$$y_2(x) = e^{3x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = c_1 e^{-x} + c_2 e^{3x}$$

- Check validity of solution $y = c_1 e^{-x} + c_2 e^{3x}$

- Use initial condition $y(0) = 0$

$$0 = c_1 + c_2$$

- Compute derivative of the solution

$$y' = -c_1 e^{-x} + 3c_2 e^{3x}$$

- Use the initial condition $y' \Big|_{\{x=0\}} = 1$

$$1 = -c_1 + 3c_2$$

- Solve for c_1 and c_2

$$\left\{ c_1 = -\frac{1}{4}, c_2 = \frac{1}{4} \right\}$$

- Substitute constant values into general solution and simplify

$$y = \frac{e^{3x}}{4} - \frac{e^{-x}}{4}$$

- Solution to the IVP

$$y = \frac{e^{3x}}{4} - \frac{e^{-x}}{4}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 17

```
dsolve([diff(y(x),x$2)-2*diff(y(x),x)-3*y(x)=0,y(0) = 0, D(y)(0) = 1],y(x), singsol=all)
```

$$y(x) = \frac{e^{3x}}{4} - \frac{e^{-x}}{4}$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 21

```
DSolve[{y''[x]-2*y'[x]-3*y[x]==0,{y[0]==0,y'[0]==1}},y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow \frac{1}{4}e^{-x}(e^{4x} - 1)$$

5.2 problem 1(b)

5.2.1	Existence and uniqueness analysis	473
5.2.2	Solving as second order linear constant coeff ode	474
5.2.3	Solving using Kovacic algorithm	477
5.2.4	Maple step by step solution	481

Internal problem ID [5958]

Internal file name [OUTPUT/5206_Sunday_June_05_2022_03_27_26_PM_81244003/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 59

Problem number: 1(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"kovacic", "second_order_linear_constant_coeff"**

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x]]
```

$$y'' + (1 + 4i)y' + y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

5.2.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = 1 + 4i$$

$$q(x) = 1$$

$$F = 0$$

Hence the ode is

$$y'' + (1 + 4i)y' + y = 0$$

The domain of $p(x) = 1 + 4i$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = 1$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

5.2.2 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 1 + 4i, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + (1 + 4i)\lambda e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + (1 + 4i)\lambda + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 1 + 4i, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-1 - 4i}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{1 + 4i^2 - (4)(1)(1)} \\ &= -\frac{1}{2} - 2i \pm \frac{\sqrt{-19 + 8i}}{2} \end{aligned}$$

Hence

$$\lambda_1 = -\frac{1}{2} - 2i + \frac{\sqrt{-19 + 8i}}{2}$$

$$\lambda_2 = -\frac{1}{2} - 2i - \frac{\sqrt{-19 + 8i}}{2}$$

Which simplifies to

$$\lambda_1 = -\frac{1}{2} - 2i + \frac{\sqrt{-19 + 8i}}{2}$$

$$\lambda_2 = -\frac{1}{2} - 2i - \frac{\sqrt{-19 + 8i}}{2}$$

The roots are complex but they are not conjugate of each others. Hence simplification using Euler relation is not possible here. Therefore the final solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$= c_1 e^{\left(-\frac{1}{2} - 2i + \frac{\sqrt{-19 + 8i}}{2}\right)x} + c_2 e^{\left(-\frac{1}{2} - 2i - \frac{\sqrt{-19 + 8i}}{2}\right)x}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 e^{\left(-\frac{1}{2} - 2i + \frac{\sqrt{-19 + 8i}}{2}\right)x} + c_2 e^{\left(-\frac{1}{2} - 2i - \frac{\sqrt{-19 + 8i}}{2}\right)x} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_1 + c_2 \quad (1A)$$

Taking derivative of the solution gives

$$y' = c_1 \left(-\frac{1}{2} - 2i + \frac{\sqrt{-19 + 8i}}{2}\right) e^{\left(-\frac{1}{2} - 2i + \frac{\sqrt{-19 + 8i}}{2}\right)x} + c_2 \left(-\frac{1}{2} - 2i - \frac{\sqrt{-19 + 8i}}{2}\right) e^{\left(-\frac{1}{2} - 2i - \frac{\sqrt{-19 + 8i}}{2}\right)x}$$

substituting $y' = 0$ and $x = 0$ in the above gives

$$0 = \frac{(c_1 - c_2)\sqrt{-19 + 8i}}{2} + \left(-\frac{1}{2} - 2i\right) c_1 + \left(-\frac{1}{2} - 2i\right) c_2 \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 0$$

$$c_2 = 0$$


Substituting these values back in above solution results in

$$y = 0$$


Summary

The solution(s) found are the following

$$y = 0 \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = 0$$

Verified OK.

5.2.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + (1 + 4i)y' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 1 + 4i \\ C &= 1 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-19 + 8i}{4} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -19 + 8i \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(-\frac{19}{4} + 2i\right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 101: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -\frac{19}{4} + 2i$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{\frac{x\sqrt{-19+8i}}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{1+4i}{1} dx} \end{aligned}$$

$$\begin{aligned}
&= z_1 e^{(-\frac{1}{2}-2i)x} \\
&= z_1 \left(e^{(-\frac{1}{2}-2i)x} \right)
\end{aligned}$$

Which simplifies to

$$y_1 = e^{-\frac{(1+4i-\sqrt{-19+8i})x}{2}}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}
y_2 &= y_1 \int \frac{e^{\int -\frac{1+4i}{1} dx}}{(y_1)^2} dx \\
&= y_1 \int \frac{e^{(-1-4i)x}}{(y_1)^2} dx \\
&= y_1 \left(\left(\frac{19}{425} + \frac{8i}{425} \right) \sqrt{-19+8i} e^{-x\sqrt{-19+8i}} \right)
\end{aligned}$$

Therefore the solution is

$$\begin{aligned}
y &= c_1 y_1 + c_2 y_2 \\
&= c_1 \left(e^{-\frac{(1+4i-\sqrt{-19+8i})x}{2}} \right) + c_2 \left(e^{-\frac{(1+4i-\sqrt{-19+8i})x}{2}} \left(\left(\frac{19}{425} + \frac{8i}{425} \right) \sqrt{-19+8i} e^{-x\sqrt{-19+8i}} \right) \right)
\end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 e^{-\frac{(1+4i-\sqrt{-19+8i})x}{2}} + \left(\frac{19}{425} + \frac{8i}{425} \right) c_2 \sqrt{-19+8i} e^{-\frac{(1+4i+\sqrt{-19+8i})x}{2}} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_1 + \left(\frac{19}{425} + \frac{8i}{425} \right) \sqrt{-19+8i} c_2 \quad (1A)$$

Taking derivative of the solution gives

$$y' = c_1 \left(-\frac{1}{2} - 2i + \frac{\sqrt{-19 + 8i}}{2} \right) e^{-\frac{(1+4i-\sqrt{-19+8i})x}{2}} + \left(\frac{19}{425} + \frac{8i}{425} \right) c_2 \sqrt{-19 + 8i} \left(-\frac{1}{2} - 2i - \frac{\sqrt{-19 + 8i}}{2} \right)$$

substituting $y' = 0$ and $x = 0$ in the above gives

$$0 = \frac{(425c_1 + (13 - 84i)c_2)\sqrt{-19 + 8i}}{850} + \left(-\frac{1}{2} - 2i \right) c_1 + \frac{c_2}{2} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 0$$

$$c_2 = 0$$


Substituting these values back in above solution results in

$$y = 0$$


Summary

The solution(s) found are the following

$$y = 0 \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = 0$$

Verified OK.

5.2.4 Maple step by step solution

Let's solve

$$\left[y'' + (1 + 4I) y' + y = 0, y(0) = 0, y' \Big|_{\{x=0\}} = 0 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 + (1 + 4I) r + 1 = 0$$

- Factor the characteristic polynomial

$$4Ir + r^2 + r + 1 = 0$$

- Roots of the characteristic polynomial

$$r = \left(-\frac{1}{2} - 2I + \frac{\sqrt{-19+8I}}{2}, -\frac{1}{2} - 2I - \frac{\sqrt{-19+8I}}{2} \right)$$

- 1st solution of the ODE

$$y_1(x) = e^{\left(-\frac{1}{2} + \frac{\sqrt{-38+10\sqrt{17}}}{4}\right)x} \cos\left(\left(-2 + \frac{\sqrt{38+10\sqrt{17}}}{4}\right)x\right)$$

- 2nd solution of the ODE

$$y_2(x) = e^{\left(-\frac{1}{2} + \frac{\sqrt{-38+10\sqrt{17}}}{4}\right)x} \sin\left(\left(-2 + \frac{\sqrt{38+10\sqrt{17}}}{4}\right)x\right)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = c_1 e^{\left(-\frac{1}{2} + \frac{\sqrt{-38+10\sqrt{17}}}{4}\right)x} \cos\left(\left(-2 + \frac{\sqrt{38+10\sqrt{17}}}{4}\right)x\right) + c_2 e^{\left(-\frac{1}{2} + \frac{\sqrt{-38+10\sqrt{17}}}{4}\right)x} \sin\left(\left(-2 + \frac{\sqrt{38+10\sqrt{17}}}{4}\right)x\right)$$

- Check validity of solution $y = c_1 e^{\left(-\frac{1}{2} + \frac{\sqrt{-38+10\sqrt{17}}}{4}\right)x} \cos\left(\left(-2 + \frac{\sqrt{38+10\sqrt{17}}}{4}\right)x\right) + c_2 e^{\left(-\frac{1}{2} + \frac{\sqrt{-38+10\sqrt{17}}}{4}\right)x} \sin\left(\left(-2 + \frac{\sqrt{38+10\sqrt{17}}}{4}\right)x\right)$

- Use initial condition $y(0) = 0$

$$0 = c_1$$

- Compute derivative of the solution

$$y' = c_1 \left(-\frac{1}{2} + \frac{\sqrt{-38+10\sqrt{17}}}{4}\right) e^{\left(-\frac{1}{2} + \frac{\sqrt{-38+10\sqrt{17}}}{4}\right)x} \cos\left(\left(-2 + \frac{\sqrt{38+10\sqrt{17}}}{4}\right)x\right) - c_1 e^{\left(-\frac{1}{2} + \frac{\sqrt{-38+10\sqrt{17}}}{4}\right)x} \sin\left(\left(-2 + \frac{\sqrt{38+10\sqrt{17}}}{4}\right)x\right)$$

- Use the initial condition $y' \Big|_{\{x=0\}} = 0$

$$0 = c_1 \left(-\frac{1}{2} + \frac{\sqrt{-38+10\sqrt{17}}}{4} \right) + c_2 \left(-2 + \frac{\sqrt{38+10\sqrt{17}}}{4} \right)$$
- Solve for c_1 and c_2

$$\{c_1 = 0, c_2 = 0\}$$
- Substitute constant values into general solution and simplify
$$y = 0$$
- Solution to the IVP
$$y = 0$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.156 (sec). Leaf size: 5

```
dsolve([diff(y(x),x$2)+(4*I+1)*diff(y(x),x)+y(x)=0,y(0) = 0, D(y)(0) = 0],y(x), singsol=all)
```

$$y(x) = 0$$

✓ Solution by Mathematica

Time used: 0.054 (sec). Leaf size: 6

```
DSolve[{y''[x]+(4*I+1)*y'[x]+y[x]==0,{y[0]==0,y'[0]==0}},y[x],x,IncludeSingularSolutions ->
```

$$y(x) \rightarrow 0$$

5.3 problem 1(c)

5.3.1	Existence and uniqueness analysis	483
5.3.2	Solving as second order linear constant coeff ode	484
5.3.3	Solving using Kovacic algorithm	486

Internal problem ID [5959]

Internal file name [OUTPUT/5207_Sunday_June_05_2022_03_27_28_PM_34164133/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 59

Problem number: 1(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' + (-1 + 3i)y' - 3iy = 0$$

With initial conditions

$$[y(0) = 2, y'(0) = 0]$$

5.3.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = -1 + 3i$$

$$q(x) = -3i$$

$$F = 0$$

Hence the ode is

$$y'' + (-1 + 3i)y' - 3iy = 0$$

The domain of $p(x) = -1 + 3i$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = -3i$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

5.3.2 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -1 + 3i, C = -3i$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + (-1 + 3i)\lambda e^{\lambda x} - 3ie^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + (-1 + 3i)\lambda - 3i = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -1 + 3i, C = -3i$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{1 - 3i}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-1 + 3i^2 - (4)(1)(-3i)} \\ &= \frac{1}{2} - \frac{3i}{2} \pm \frac{1}{2} + \frac{3i}{2} \end{aligned}$$

Hence

$$\lambda_1 = \frac{1}{2} - \frac{3i}{2} + \frac{1}{2} + \frac{3i}{2}$$

$$\lambda_2 = \frac{1}{2} - \frac{3i}{2} - \frac{1}{2} + \frac{3i}{2}$$

Which simplifies to

$$\lambda_1 = 1$$

$$\lambda_2 = -3i$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(1)x} + c_2 e^{(-3i)x}$$

Or

$$y = c_1 e^x + e^{-3ix} c_2$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 e^x + e^{-3ix} c_2 \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 2$ and $x = 0$ in the above gives

$$2 = c_1 + c_2 \tag{1A}$$

Taking derivative of the solution gives

$$y' = c_1 e^x - 3ie^{-3ix} c_2$$

substituting $y' = 0$ and $x = 0$ in the above gives

$$0 = -3c_2 i + c_1 \tag{2A}$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = \frac{9}{5} + \frac{3i}{5}$$

$$c_2 = \frac{1}{5} - \frac{3i}{5}$$

Substituting these values back in above solution results in

$$y = \frac{9 e^x}{5} + \frac{3ie^x}{5} + \frac{e^{-3ix}}{5} - \frac{3ie^{-3ix}}{5}$$

Summary

The solution(s) found are the following

$$y = \left(\frac{1}{5} - \frac{3i}{5}\right) e^{-3ix} + \left(\frac{9}{5} + \frac{3i}{5}\right) e^x \quad (1)$$

Verification of solutions

$$y = \left(\frac{1}{5} - \frac{3i}{5}\right) e^{-3ix} + \left(\frac{9}{5} + \frac{3i}{5}\right) e^x$$

Verified OK.

5.3.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + (-1 + 3i)y' - 3iy = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -1 + 3i \\ C &= -3i \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-4 + 3i}{2} \quad (6)$$

Comparing the above to (5) shows that

$$s = -4 + 3i$$

$$t = 2$$

Therefore eq. (4) becomes

$$z''(x) = \left(-2 + \frac{3i}{2}\right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 103: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$O(\infty) = \deg(t) - \deg(s)$$

$$= 0 - 0$$

$$= 0$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case

one are met. Therefore

$$L = [1]$$

Since $r = -2 + \frac{3i}{2}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{\left(\frac{1}{2} + \frac{3i}{2}\right)x}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-1+3i}{1} dx} \\ &= z_1 e^{\left(\frac{1}{2} - \frac{3i}{2}\right)x} \\ &= z_1 \left(e^{\left(\frac{1}{2} - \frac{3i}{2}\right)x} \right) \end{aligned}$$

Which simplifies to

$$y_1 = e^x$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-1+3i}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{(1-3i)x}}{(y_1)^2} dx \\ &= y_1 \left(\left(-\frac{1}{10} + \frac{3i}{10} \right) e^{(-1-3i)x} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^x) + c_2 \left(e^x \left(\left(-\frac{1}{10} + \frac{3i}{10} \right) e^{(-1-3i)x} \right) \right) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 e^x + \left(-\frac{1}{10} + \frac{3i}{10}\right) c_2 e^{-3ix} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 2$ and $x = 0$ in the above gives

$$2 = c_1 + \left(-\frac{1}{10} + \frac{3i}{10}\right) c_2 \quad (1A)$$

Taking derivative of the solution gives

$$y' = c_1 e^x + \left(\frac{9}{10} + \frac{3i}{10}\right) c_2 e^{-3ix}$$

substituting $y' = 0$ and $x = 0$ in the above gives

$$0 = c_1 + \left(\frac{9}{10} + \frac{3i}{10}\right) c_2 \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$\begin{aligned} c_1 &= \frac{9}{5} + \frac{3i}{5} \\ c_2 &= -2 \end{aligned}$$

Substituting these values back in above solution results in

$$y = \frac{9e^x}{5} + \frac{3ie^x}{5} + \frac{e^{-3ix}}{5} - \frac{3ie^{-3ix}}{5}$$

Summary

The solution(s) found are the following

$$y = \left(\frac{1}{5} - \frac{3i}{5}\right) e^{-3ix} + \left(\frac{9}{5} + \frac{3i}{5}\right) e^x \quad (1)$$

Verification of solutions

$$y = \left(\frac{1}{5} - \frac{3i}{5}\right) e^{-3ix} + \left(\frac{9}{5} + \frac{3i}{5}\right) e^x$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 20

```
dsolve([diff(y(x),x$2)+(3*I-1)*diff(y(x),x)-3*I*y(x)=0,y(0) = 2, D(y)(0) = 0],y(x), singsol=
```

$$y(x) = \left(\frac{9}{5} + \frac{3i}{5}\right) e^x + \left(\frac{1}{5} - \frac{3i}{5}\right) e^{-3ix}$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 31

```
DSolve[{y'[x]+(3*I-1)*y'[x]-3*I*y[x]==0,{y[0]==2,y'[0]==0}},y[x],x,IncludeSingularSolutions
```

$$y(x) \rightarrow \frac{1}{5} e^{-3ix} ((9 + 3i)e^{(1+3i)x} + (1 - 3i))$$

5.4 problem 1(d)

5.4.1	Existence and uniqueness analysis	491
5.4.2	Solving as second order linear constant coeff ode	492
5.4.3	Solving as second order ode can be made integrable ode	495
5.4.4	Solving using Kovacic algorithm	497
5.4.5	Maple step by step solution	501

Internal problem ID [5960]

Internal file name [OUTPUT/5208_Sunday_June_05_2022_03_27_29_PM_32949906/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 59

Problem number: 1(d).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"kovacic", "second_order_linear_constant_coeff", "second_order_ode_can_be_made_integrable"**

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' + 10y = 0$$

With initial conditions

$$[y(0) = \pi, y'(0) = \pi^2]$$

5.4.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = 0$$

$$q(x) = 10$$

$$F = 0$$

Hence the ode is

$$y'' + 10y = 0$$

The domain of $p(x) = 0$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = 10$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

5.4.2 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 10$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 10 e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 10 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 10$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(10)} \\ &= \pm i\sqrt{10} \end{aligned}$$

Hence

$$\lambda_1 = +i\sqrt{10}$$

$$\lambda_2 = -i\sqrt{10}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= i\sqrt{10} \\ \lambda_2 &= -i\sqrt{10}\end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = \sqrt{10}$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0(c_1 \cos(\sqrt{10}x) + c_2 \sin(\sqrt{10}x))$$

Or

$$y = c_1 \cos(\sqrt{10}x) + c_2 \sin(\sqrt{10}x)$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 \cos(\sqrt{10}x) + c_2 \sin(\sqrt{10}x) \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = \pi$ and $x = 0$ in the above gives

$$\pi = c_1 \tag{1A}$$

Taking derivative of the solution gives

$$y' = -c_1\sqrt{10} \sin(\sqrt{10}x) + c_2\sqrt{10} \cos(\sqrt{10}x)$$

substituting $y' = \pi^2$ and $x = 0$ in the above gives

$$\pi^2 = c_2\sqrt{10} \tag{2A}$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = \pi$$

$$c_2 = \frac{\sqrt{10}\pi^2}{10}$$


Substituting these values back in above solution results in

$$y = \cos(\sqrt{10}x)\pi + \frac{\sin(\sqrt{10}x)\sqrt{10}\pi^2}{10}$$


Summary

The solution(s) found are the following

$$y = \cos(\sqrt{10}x)\pi + \frac{\sin(\sqrt{10}x)\sqrt{10}\pi^2}{10} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \cos(\sqrt{10}x)\pi + \frac{\sin(\sqrt{10}x)\sqrt{10}\pi^2}{10}$$

Verified OK.

5.4.3 Solving as second order ode can be made integrable ode

Multiplying the ode by y' gives

$$y'y'' + 10y'y = 0$$

Integrating the above w.r.t x gives

$$\int (y'y'' + 10y'y) dx = 0$$
$$\frac{y'^2}{2} + 5y^2 = c_2$$

Which is now solved for y . Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \sqrt{-10y^2 + 2c_1} \quad (1)$$

$$y' = -\sqrt{-10y^2 + 2c_1} \quad (2)$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int \frac{1}{\sqrt{-10y^2 + 2c_1}} dy = \int dx$$
$$\frac{\sqrt{10} \arctan\left(\frac{\sqrt{10}y}{\sqrt{-10y^2 + 2c_1}}\right)}{10} = c_2 + x$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{1}{\sqrt{-10y^2 + 2c_1}} dy = \int dx$$
$$-\frac{\sqrt{10} \arctan\left(\frac{\sqrt{10}y}{\sqrt{-10y^2 + 2c_1}}\right)}{10} = x + c_3$$

Initial conditions are used to solve for the constants of integration.

Looking at the First solution

$$\frac{\sqrt{10} \arctan \left(\frac{\sqrt{10} y}{\sqrt{-10y^2+2c_1}} \right)}{10} = c_2 + x \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = \pi$ and $x = 0$ in the above gives

$$\frac{\arctan \left(\frac{\sqrt{5} \pi}{\sqrt{-5\pi^2+c_1}} \right) \sqrt{10}}{10} = c_2 \quad (1A)$$

Taking derivative of the solution gives

$$y' = \left(\tan \left((c_2 + x) \sqrt{10} \right)^2 + 1 \right) \sqrt{2} \sqrt{\frac{c_1}{\tan \left((c_2 + x) \sqrt{10} \right)^2 + 1}} - \frac{\tan \left((c_2 + x) \sqrt{10} \right)^2 \sqrt{2} c_1}{\sqrt{\frac{c_1}{\tan \left((c_2 + x) \sqrt{10} \right)^2 + 1}} \left(\tan \left((c_2 + x) \sqrt{10} \right) \right)}$$

substituting $y' = \pi^2$ and $x = 0$ in the above gives

$$\pi^2 = \frac{\cos \left(c_2 \sqrt{10} \right)^2 c_1 \sqrt{2}}{\sqrt{c_1 \cos \left(c_2 \sqrt{10} \right)^2}} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = \frac{1}{2} \pi^4 + 5\pi^2$$

$$c_2 = \frac{\arctan \left(\frac{\sqrt{10}}{\pi} \right) \sqrt{10}}{10}$$

Substituting these values back in above solution results in

$$\frac{\arctan \left(\frac{\sqrt{10} y}{\sqrt{-10y^2+\pi^4+10\pi^2}} \right) \sqrt{10}}{10} = \frac{\arctan \left(\frac{\sqrt{10}}{\pi} \right) \sqrt{10}}{10} + x$$

Looking at the Second solution

$$-\frac{\sqrt{10} \arctan \left(\frac{\sqrt{10} y}{\sqrt{-10y^2+2c_1}} \right)}{10} = x + c_3 \quad (2)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = \pi$ and $x = 0$ in the above gives

$$-\frac{\arctan \left(\frac{\sqrt{5} \pi}{\sqrt{-5\pi^2+c_1}} \right) \sqrt{10}}{10} = c_3 \quad (1A)$$

Taking derivative of the solution gives

$$y' = -\left(\tan\left((x+c_3)\sqrt{10}\right)^2 + 1\right)\sqrt{2}\sqrt{\frac{c_1}{\tan\left((x+c_3)\sqrt{10}\right)^2 + 1}} + \frac{\tan\left((x+c_3)\sqrt{10}\right)^2\sqrt{2}c_1}{\sqrt{\frac{c_1}{\tan\left((x+c_3)\sqrt{10}\right)^2 + 1}}\left(\tan\left((x+c_3)\sqrt{10}\right)\sqrt{10}\right)}$$

substituting $y' = \pi^2$ and $x = 0$ in the above gives

$$\pi^2 = -\frac{\cos\left(c_3\sqrt{10}\right)^2 c_1\sqrt{2}}{\sqrt{c_1 \cos\left(c_3\sqrt{10}\right)^2}} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_3\}$. There is no solution for the constants of integrations. This solution is removed.

Summary

The solution(s) found are the following

$$\frac{\arctan\left(\frac{\sqrt{10}y}{\sqrt{-10y^2+\pi^4+10\pi^2}}\right)\sqrt{10}}{10} = \frac{\arctan\left(\frac{\sqrt{10}}{\pi}\right)\sqrt{10}}{10} + x \quad (1)$$

Verification of solutions

$$\frac{\arctan\left(\frac{\sqrt{10}y}{\sqrt{-10y^2+\pi^4+10\pi^2}}\right)\sqrt{10}}{10} = \frac{\arctan\left(\frac{\sqrt{10}}{\pi}\right)\sqrt{10}}{10} + x$$

Verified OK.

5.4.4 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 10y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 10 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} = \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-10}{1} \tag{6}$$

Comparing the above to (5) shows that

$$s = -10$$

$$t = 1$$

Therefore eq. (4) becomes

$$z''(x) = -10z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 104: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -10$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(\sqrt{10}x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned} y_1 &= z_1 \\ &= \cos(\sqrt{10}x) \end{aligned}$$

Which simplifies to

$$y_1 = \cos(\sqrt{10}x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(\sqrt{10}x) \int \frac{1}{\cos^2(\sqrt{10}x)} dx \\ &= \cos(\sqrt{10}x) \left(\frac{\sqrt{10} \tan(\sqrt{10}x)}{10} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}
 y &= c_1 y_1 + c_2 y_2 \\
 &= c_1 \left(\cos \left(\sqrt{10} x \right) \right) + c_2 \left(\cos \left(\sqrt{10} x \right) \left(\frac{\sqrt{10} \tan \left(\sqrt{10} x \right)}{10} \right) \right)
 \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 \cos \left(\sqrt{10} x \right) + \frac{c_2 \sqrt{10} \sin \left(\sqrt{10} x \right)}{10} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = \pi$ and $x = 0$ in the above gives

$$\pi = c_1 \quad (1A)$$

Taking derivative of the solution gives

$$y' = -c_1 \sqrt{10} \sin \left(\sqrt{10} x \right) + c_2 \cos \left(\sqrt{10} x \right)$$

substituting $y' = \pi^2$ and $x = 0$ in the above gives

$$\pi^2 = c_2 \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = \pi$$

$$c_2 = \pi^2$$


Substituting these values back in above solution results in

$$y = \cos \left(\sqrt{10} x \right) \pi + \frac{\sin \left(\sqrt{10} x \right) \sqrt{10} \pi^2}{10}$$


Summary

The solution(s) found are the following

$$y = \cos \left(\sqrt{10} x \right) \pi + \frac{\sin \left(\sqrt{10} x \right) \sqrt{10} \pi^2}{10} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \cos(\sqrt{10}x)\pi + \frac{\sin(\sqrt{10}x)\sqrt{10}\pi^2}{10}$$

Verified OK.

5.4.5 Maple step by step solution

Let's solve

$$\left[y'' + 10y = 0, y(0) = \pi, y'|_{\{x=0\}} = \pi^2 \right]$$

- Highest derivative means the order of the ODE is 2
- y''
- Characteristic polynomial of ODE
- $r^2 + 10 = 0$
- Use quadratic formula to solve for r
- $r = \frac{0 \pm (\sqrt{-40})}{2}$
- Roots of the characteristic polynomial
- $r = (-I\sqrt{10}, I\sqrt{10})$
- 1st solution of the ODE

$$y_1(x) = \cos(\sqrt{10}x)$$

- 2nd solution of the ODE

$$y_2(x) = \sin(\sqrt{10}x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = c_1 \cos(\sqrt{10}x) + c_2 \sin(\sqrt{10}x)$$

- Check validity of solution $y = c_1 \cos(\sqrt{10}x) + c_2 \sin(\sqrt{10}x)$

- Use initial condition $y(0) = \pi$

$$\pi = c_1$$

- Compute derivative of the solution

$$y' = -c_1 \sqrt{10} \sin(\sqrt{10}x) + c_2 \sqrt{10} \cos(\sqrt{10}x)$$

- Use the initial condition $y'|_{\{x=0\}} = \pi^2$

$$\pi^2 = c_2 \sqrt{10}$$

- Solve for c_1 and c_2

$$\left\{ c_1 = \pi, c_2 = \frac{\sqrt{10} \pi^2}{10} \right\}$$

- Substitute constant values into general solution and simplify

$$y = \frac{\pi(\sin(\sqrt{10}x)\sqrt{10}\pi + 10\cos(\sqrt{10}x))}{10}$$

- Solution to the IVP

$$y = \frac{\pi(\sin(\sqrt{10}x)\sqrt{10}\pi + 10\cos(\sqrt{10}x))}{10}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 27

```
dsolve([diff(y(x),x$2)+10*y(x)=0,y(0) = Pi, D(y)(0) = Pi^2],y(x), singsol=all)
```

$$y(x) = \frac{\pi(\pi\sqrt{10} \sin(\sqrt{10}x) + 10 \cos(\sqrt{10}x))}{10}$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 33

```
DSolve[{y'[x]+10*y[x]==0,{y[0]==Pi,y'[0]==Pi^2}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\pi^2 \sin(\sqrt{10}x)}{\sqrt{10}} + \pi \cos(\sqrt{10}x)$$

6 Chapter 2. Linear equations with constant coefficients. Page 69

6.1	problem 1(a)	505
6.2	problem 1(b)	516
6.3	problem 1(c)	527
6.4	problem 1(d)	540
6.5	problem 1(e)	551
6.6	problem 1(f)	562
6.7	problem 1(g)	573
6.8	problem 1(h)	584
6.9	problem 1(i)	597
6.10	problem 1(j)	608
6.11	problem 4(c)	619

6.1 problem 1(a)

6.1.1 Solving as second order linear constant coeff ode	505
6.1.2 Solving using Kovacic algorithm	508
6.1.3 Maple step by step solution	513

Internal problem ID [5961]

Internal file name [OUTPUT/5209_Sunday_June_05_2022_03_27_31_PM_85388453/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 69

Problem number: 1(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"kovacic", "second_order_linear_constant_coeff"**

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + 4y = \cos(x)$$

6.1.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 4, f(x) = \cos(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 4y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 4$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 4 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 4 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 4$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(4)} \\ &= \pm 2i \end{aligned}$$

Hence

$$\lambda_1 = +2i$$

$$\lambda_2 = -2i$$

Which simplifies to

$$\lambda_1 = 2i$$

$$\lambda_2 = -2i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 2$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (c_1 \cos(2x) + c_2 \sin(2x))$$

Or

$$y = c_1 \cos(2x) + c_2 \sin(2x)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \cos(2x) + c_2 \sin(2x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\cos(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(2x), \sin(2x)\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(x) + A_2 \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$3A_1 \cos(x) + 3A_2 \sin(x) = \cos(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{3}, A_2 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{\cos(x)}{3}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(2x) + c_2 \sin(2x)) + \left(\frac{\cos(x)}{3} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(2x) + c_2 \sin(2x) + \frac{\cos(x)}{3} \quad (1)$$




Figure 109: Slope field plot

Verification of solutions

$$y = c_1 \cos(2x) + c_2 \sin(2x) + \frac{\cos(x)}{3}$$

Verified OK.

6.1.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 4y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= 1 \\B &= 0 \\C &= 4\end{aligned}\tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x)\tag{4}$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\&= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned}\tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-4}{1}\tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= -4 \\t &= 1\end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -4z(x)\tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 106: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -4$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(2x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}
 y_1 &= z_1 \\
 &= \cos(2x)
 \end{aligned}$$

Which simplifies to

$$y_1 = \cos(2x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(2x) \int \frac{1}{\cos(2x)^2} dx \\ &= \cos(2x) \left(\frac{\tan(2x)}{2} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(2x)) + c_2 \left(\cos(2x) \left(\frac{\tan(2x)}{2} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$.
 y_h is the solution to

$$y'' + 4y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 \cos(2x) + \frac{c_2 \sin(2x)}{2}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\cos(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ \frac{\sin(2x)}{2}, \cos(2x) \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(x) + A_2 \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$3A_1 \cos(x) + 3A_2 \sin(x) = \cos(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{3}, A_2 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{\cos(x)}{3}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 \cos(2x) + \frac{c_2 \sin(2x)}{2} \right) + \left(\frac{\cos(x)}{3} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(2x) + \frac{c_2 \sin(2x)}{2} + \frac{\cos(x)}{3} \quad (1)$$




Figure 110: Slope field plot

Verification of solutions

$$y = c_1 \cos(2x) + \frac{c_2 \sin(2x)}{2} + \frac{\cos(x)}{3}$$

Verified OK.

6.1.3 Maple step by step solution

Let's solve

$$y'' + 4y = \cos(x)$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 4 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-16})}{2}$$

- Roots of the characteristic polynomial

$$r = (-2I, 2I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(2x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(2x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 \cos(2x) + c_2 \sin(2x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \cos(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(2x) & \sin(2x) \\ -2\sin(2x) & 2\cos(2x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 2$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{\cos(2x) \left(\int 4 \cos(x)^2 \sin(x) dx \right)}{4} + \frac{\sin(2x) \left(\int (\cos(x) + \cos(3x)) dx \right)}{4}$$

- Compute integrals

$$y_p(x) = \frac{\cos(x)}{3}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 \cos(2x) + c_2 \sin(2x) + \frac{\cos(x)}{3}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(diff(y(x),x$2)+4*y(x)=cos(x),y(x), singsol=all)
```

$$y(x) = \sin(2x)c_2 + \cos(2x)c_1 + \frac{\cos(x)}{3}$$

✓ Solution by Mathematica

Time used: 0.032 (sec). Leaf size: 26

```
DSolve[y''[x]+4*y[x]==Cos[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\cos(x)}{3} + c_1 \cos(2x) + c_2 \sin(2x)$$

6.2 problem 1(b)

6.2.1	Solving as second order linear constant coeff ode	516
6.2.2	Solving using Kovacic algorithm	520
6.2.3	Maple step by step solution	525

Internal problem ID [5962]

Internal file name [OUTPUT/5210_Sunday_June_05_2022_03_27_32_PM_59574836/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 69

Problem number: 1(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + 9y = \sin(3x)$$

6.2.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 9, f(x) = \sin(3x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 9y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 9$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 9 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 9 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 9$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(9)} \\ &= \pm 3i \end{aligned}$$

Hence

$$\lambda_1 = +3i$$

$$\lambda_2 = -3i$$

Which simplifies to

$$\lambda_1 = 3i$$

$$\lambda_2 = -3i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 3$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (c_1 \cos(3x) + c_2 \sin(3x))$$

Or

$$y = c_1 \cos(3x) + c_2 \sin(3x)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \cos(3x) + c_2 \sin(3x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(3x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(3x), \sin(3x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(3x), \sin(3x)\}$$

Since $\cos(3x)$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x \cos(3x), x \sin(3x)\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x \cos(3x) + A_2 x \sin(3x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-6A_1 \sin(3x) + 6A_2 \cos(3x) = \sin(3x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{6}, A_2 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{x \cos(3x)}{6}$$

Therefore the general solution is

$$\begin{aligned}y &= y_h + y_p \\ &= (c_1 \cos(3x) + c_2 \sin(3x)) + \left(-\frac{x \cos(3x)}{6}\right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(3x) + c_2 \sin(3x) - \frac{x \cos(3x)}{6} \quad (1)$$




Figure 111: Slope field plot

Verification of solutions

$$y = c_1 \cos(3x) + c_2 \sin(3x) - \frac{x \cos(3x)}{6}$$

Verified OK.

6.2.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 9y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 9 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-9}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -9 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -9z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 108: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -9$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(3x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}y_1 &= z_1 \\ &= \cos(3x)\end{aligned}$$

Which simplifies to

$$y_1 = \cos(3x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned}y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(3x) \int \frac{1}{\cos(3x)^2} dx \\ &= \cos(3x) \left(\frac{\tan(3x)}{3} \right)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1(\cos(3x)) + c_2 \left(\cos(3x) \left(\frac{\tan(3x)}{3} \right) \right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$.
 y_h is the solution to

$$y'' + 9y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 \cos(3x) + \frac{c_2 \sin(3x)}{3}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(3x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(3x), \sin(3x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ \frac{\sin(3x)}{3}, \cos(3x) \right\}$$

Since $\cos(3x)$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x \cos(3x), x \sin(3x)\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x \cos(3x) + A_2 x \sin(3x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-6A_1 \sin(3x) + 6A_2 \cos(3x) = \sin(3x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{6}, A_2 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{x \cos(3x)}{6}$$

Therefore the general solution is

$$y = y_h + y_p \\ = \left(c_1 \cos(3x) + \frac{c_2 \sin(3x)}{3} \right) + \left(-\frac{x \cos(3x)}{6} \right)$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(3x) + \frac{c_2 \sin(3x)}{3} - \frac{x \cos(3x)}{6} \quad (1)$$




Figure 112: Slope field plot

Verification of solutions

$$y = c_1 \cos(3x) + \frac{c_2 \sin(3x)}{3} - \frac{x \cos(3x)}{6}$$

Verified OK.

6.2.3 Maple step by step solution

Let's solve

$$y'' + 9y = \sin(3x)$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 9 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-36})}{2}$$

- Roots of the characteristic polynomial

$$r = (-3I, 3I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(3x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(3x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 \cos(3x) + c_2 \sin(3x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \sin(3x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(3x) & \sin(3x) \\ -3\sin(3x) & 3\cos(3x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 3$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{\cos(3x)(\int \sin(3x)^2 dx)}{3} + \frac{\sin(3x)(\int \sin(6x) dx)}{6}$$

- Compute integrals

$$y_p(x) = \frac{\sin(3x)}{36} - \frac{x \cos(3x)}{6}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 \cos(3x) + c_2 \sin(3x) + \frac{\sin(3x)}{36} - \frac{x \cos(3x)}{6}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 24

```
dsolve(diff(y(x),x$2)+9*y(x)=sin(3*x),y(x), singsol=all)
```

$$y(x) = \frac{(-x + 6c_1) \cos(3x)}{6} + \sin(3x) c_2$$

✓ Solution by Mathematica

Time used: 0.036 (sec). Leaf size: 33

```
DSolve[y''[x]+9*y[x]==Sin[3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \left(-\frac{x}{6} + c_1\right) \cos(3x) + \frac{1}{36}(1 + 36c_2) \sin(3x)$$

6.3 problem 1(c)

6.3.1	Solving as second order linear constant coeff ode	527
6.3.2	Solving using Kovacic algorithm	532
6.3.3	Maple step by step solution	537

Internal problem ID [5963]

Internal file name [OUTPUT/5211_Sunday_June_05_2022_03_27_34_PM_71151919/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 69

Problem number: 1(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + y = \tan(x)$$

6.3.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 1, f(x) = \tan(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\lambda_1 = +i$$

$$\lambda_2 = -i$$

Which simplifies to

$$\lambda_1 = i$$

$$\lambda_2 = -i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (\cos(x) c_1 + c_2 \sin(x))$$

Or

$$y = \cos(x) c_1 + c_2 \sin(x)$$

Therefore the homogeneous solution y_h is

$$y_h = \cos(x) c_1 + c_2 \sin(x)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \cos(x)$$

$$y_2 = \sin(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ \frac{d}{dx}(\cos(x)) & \frac{d}{dx}(\sin(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{vmatrix}$$

Therefore

$$W = (\cos(x))(\cos(x)) - (\sin(x))(-\sin(x))$$

Which simplifies to

$$W = \cos(x)^2 + \sin(x)^2$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\sin(x) \tan(x)}{1} dx$$

Which simplifies to

$$u_1 = - \int \sin(x) \tan(x) dx$$

Hence

$$u_1 = \sin(x) - \ln(\sec(x) + \tan(x))$$

And Eq. (3) becomes

$$u_2 = \int \frac{\tan(x) \cos(x)}{1} dx$$

Which simplifies to

$$u_2 = \int \sin(x) dx$$

Hence

$$u_2 = -\cos(x)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = (\sin(x) - \ln(\sec(x) + \tan(x))) \cos(x) - \cos(x) \sin(x)$$

Which simplifies to

$$y_p(x) = -\cos(x) \ln(\sec(x) + \tan(x))$$

Therefore the general solution is

$$\begin{aligned}y &= y_h + y_p \\ &= (\cos(x) c_1 + c_2 \sin(x)) + (-\cos(x) \ln(\sec(x) + \tan(x)))\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \cos(x) c_1 + c_2 \sin(x) - \cos(x) \ln(\sec(x) + \tan(x)) \quad (1)$$




Figure 113: Slope field plot

Verification of solutions

$$y = \cos(x) c_1 + c_2 \sin(x) - \cos(x) \ln(\sec(x) + \tan(x))$$

Verified OK.

6.3.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 1 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 110: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}y_1 &= z_1 \\ &= \cos(x)\end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned}y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x))\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(x)) + c_2 (\cos(x) (\tan(x)))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = \cos(x) c_1 + c_2 \sin(x)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \cos(x)$$

$$y_2 = \sin(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ \frac{d}{dx}(\cos(x)) & \frac{d}{dx}(\sin(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{vmatrix}$$

Therefore

$$W = (\cos(x))(\cos(x)) - (\sin(x))(-\sin(x))$$

Which simplifies to

$$W = \cos(x)^2 + \sin(x)^2$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\sin(x) \tan(x)}{1} dx$$

Which simplifies to

$$u_1 = - \int \sin(x) \tan(x) dx$$

Hence

$$u_1 = \sin(x) - \ln(\sec(x) + \tan(x))$$

And Eq. (3) becomes

$$u_2 = \int \frac{\tan(x) \cos(x)}{1} dx$$

Which simplifies to

$$u_2 = \int \sin(x) dx$$

Hence

$$u_2 = -\cos(x)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = (\sin(x) - \ln(\sec(x) + \tan(x))) \cos(x) - \cos(x) \sin(x)$$

Which simplifies to

$$y_p(x) = -\cos(x) \ln(\sec(x) + \tan(x))$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (\cos(x) c_1 + c_2 \sin(x)) + (-\cos(x) \ln(\sec(x) + \tan(x))) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \cos(x) c_1 + c_2 \sin(x) - \cos(x) \ln(\sec(x) + \tan(x)) \quad (1)$$




Figure 114: Slope field plot

Verification of solutions

$$y = \cos(x) c_1 + c_2 \sin(x) - \cos(x) \ln(\sec(x) + \tan(x))$$

Verified OK.

6.3.3 Maple step by step solution

Let's solve

$$y'' + y = \tan(x)$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$
- Roots of the characteristic polynomial

$$r = (-I, I)$$
- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x)$$
- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(x)$$
- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$
- Substitute in solutions of the homogeneous ODE

$$y = \cos(x) c_1 + c_2 \sin(x) + y_p(x)$$
- Find a particular solution $y_p(x)$ of the ODE
 - Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \tan(x) \right]$$
 - Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix}$$
 - Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$
 - Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\cos(x) \left(\int \sin(x) \tan(x) dx \right) + \sin(x) \left(\int \sin(x) dx \right)$$
 - Compute integrals

$$y_p(x) = -\cos(x) \ln(\sec(x) + \tan(x))$$
- Substitute particular solution into general solution to ODE

$$y = \cos(x) c_1 + c_2 \sin(x) - \cos(x) \ln(\sec(x) + \tan(x))$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 23

```
dsolve(diff(y(x),x$2)+y(x)=tan(x),y(x), singsol=all)
```

$$y(x) = \sin(x) c_2 + \cos(x) c_1 - \cos(x) \ln(\sec(x) + \tan(x))$$

✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 23

```
DSolve[y''[x]+y[x]==Tan[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \cos(x)(-\operatorname{arctanh}(\sin(x))) + c_1 \cos(x) + c_2 \sin(x)$$

6.4 problem 1(d)

6.4.1	Solving as second order linear constant coeff ode	540
6.4.2	Solving using Kovacic algorithm	543
6.4.3	Maple step by step solution	548

Internal problem ID [5964]

Internal file name [OUTPUT/5212_Sunday_June_05_2022_03_27_35_PM_73300072/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 69

Problem number: 1(d).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"kovacic", "second_order_linear_constant_coeff"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' + 2iy' + y = x$$

6.4.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 2i, C = 1, f(x) = x$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 2iy' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 2i, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 2i\lambda e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 2i\lambda + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 2i, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-2i}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{2i^2 - (4)(1)(1)} \\ &= -i \pm i\sqrt{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= -i + i\sqrt{2} \\ \lambda_2 &= -i - i\sqrt{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= i(\sqrt{2} - 1) \\ \lambda_2 &= -i(1 + \sqrt{2}) \end{aligned}$$

The roots are complex but they are not conjugate of each others. Hence simplification using Euler relation is not possible here. Therefore the final solution is

$$\begin{aligned} y &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ &= c_1 e^{i(\sqrt{2}-1)x} + c_2 e^{-i(1+\sqrt{2})x} \end{aligned}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^{i(\sqrt{2}-1)x} + c_2 e^{-i(1+\sqrt{2})x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{i(\sqrt{2}-1)x}, e^{-i(1+\sqrt{2})x} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_2x + A_1$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_2x + 2iA_2 + A_1 = x$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = -2i, A_2 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -2i + x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 e^{i(\sqrt{2}-1)x} + c_2 e^{-i(1+\sqrt{2})x} \right) + (-2i + x) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{i(\sqrt{2}-1)x} + c_2 e^{-i(1+\sqrt{2})x} - 2i + x \quad (1)$$




Figure 115: Slope field plot

Verification of solutions

$$y = c_1 e^{i(\sqrt{2}-1)x} + c_2 e^{-i(1+\sqrt{2})x} - 2i + x$$

Verified OK.

6.4.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 2iy' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 2i \\ C &= 1 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-2}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -2 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -2z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 112: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -2$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x\sqrt{2})$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{2i}{1} dx} \\ &= z_1 e^{-ix} \\ &= z_1 (e^{-ix}) \end{aligned}$$

Which simplifies to

$$y_1 = \cos(x\sqrt{2}) e^{-ix}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{2i}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-2ix}}{(y_1)^2} dx \\ &= y_1 \left(\frac{\sqrt{2} \tan(x\sqrt{2})}{2} \right) \end{aligned}$$

Therefore the solution is

$$y = c_1 y_1 + c_2 y_2$$

$$= c_1 \left(\cos(x\sqrt{2}) e^{-ix} \right) + c_2 \left(\cos(x\sqrt{2}) e^{-ix} \left(\frac{\sqrt{2} \tan(x\sqrt{2})}{2} \right) \right)$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 2iy' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 \cos(x\sqrt{2}) e^{-ix} + \frac{c_2 e^{-ix} \sqrt{2} \sin(x\sqrt{2})}{2}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ \cos(x\sqrt{2}) e^{-ix}, \frac{e^{-ix} \sqrt{2} \sin(x\sqrt{2})}{2} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_2 x + A_1$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_2x + 2iA_2 + A_1 = x$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = -2i, A_2 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -2i + x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 \cos(x\sqrt{2}) e^{-ix} + \frac{c_2 e^{-ix} \sqrt{2} \sin(x\sqrt{2})}{2} \right) + (-2i + x) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(x\sqrt{2}) e^{-ix} + \frac{c_2 e^{-ix} \sqrt{2} \sin(x\sqrt{2})}{2} - 2i + x \quad (1)$$




Figure 116: Slope field plot

Verification of solutions

$$y = c_1 \cos(x\sqrt{2}) e^{-ix} + \frac{c_2 e^{-ix} \sqrt{2} \sin(x\sqrt{2})}{2} - 2i + x$$

Verified OK.

6.4.3 Maple step by step solution

Let's solve

$$y'' + 2Iy' + y = x$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 2Ir + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-2I) \pm (\sqrt{-8})}{2}$$

- Roots of the characteristic polynomial

$$r = (-1 - i\sqrt{2}, -1 + i\sqrt{2})$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos((1 + \sqrt{2})x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin((1 + \sqrt{2})x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 \cos((1 + \sqrt{2})x) + c_2 \sin((1 + \sqrt{2})x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = x \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos((1 + \sqrt{2})x) & \sin((1 + \sqrt{2})x) \\ -(1 + \sqrt{2})\sin((1 + \sqrt{2})x) & (1 + \sqrt{2})\cos((1 + \sqrt{2})x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1 + \sqrt{2}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = \frac{-\cos((1 + \sqrt{2})x) \left(\int \sin((1 + \sqrt{2})x) x dx \right) + \sin((1 + \sqrt{2})x) \left(\int \cos((1 + \sqrt{2})x) x dx \right)}{1 + \sqrt{2}}$$

- Compute integrals

$$y_p(x) = \frac{x}{(1 + \sqrt{2})^2}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 \cos((1 + \sqrt{2})x) + c_2 \sin((1 + \sqrt{2})x) + \frac{x}{(1 + \sqrt{2})^2}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 34

```
dsolve(diff(y(x),x$2)+2*I*diff(y(x),x)+y(x)=x,y(x), singsol=all)
```

$$y(x) = e^{-ix} \sin(\sqrt{2}x) c_2 + e^{-ix} \cos(\sqrt{2}x) c_1 + x - 2i$$

✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 44

```
DSolve[y''[x]+2*I*y'[x]+y[x]==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x + c_1 e^{-i(1+\sqrt{2})x} + c_2 e^{i(\sqrt{2}-1)x} - 2i$$

6.5 problem 1(e)

6.5.1	Solving as second order linear constant coeff ode	551
6.5.2	Solving using Kovacic algorithm	554
6.5.3	Maple step by step solution	559

Internal problem ID [5965]

Internal file name [OUTPUT/5213_Sunday_June_05_2022_03_27_37_PM_28932921/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 69

Problem number: 1(e).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"kovacic", "second_order_linear_constant_coeff"**

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' - 4y' + 5y = 3e^{-x} + 2x^2$$

6.5.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = -4, C = 5, f(x) = 3e^{-x} + 2x^2$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 4y' + 5y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -4, C = 5$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 4\lambda e^{\lambda x} + 5 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 4\lambda + 5 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -4, C = 5$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{4}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-4^2 - (4)(1)(5)} \\ &= 2 \pm i \end{aligned}$$

Hence

$$\lambda_1 = 2 + i$$

$$\lambda_2 = 2 - i$$

Which simplifies to

$$\lambda_1 = 2 + i$$

$$\lambda_2 = 2 - i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 2$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^{2x} (\cos(x) c_1 + c_2 \sin(x))$$

Therefore the homogeneous solution y_h is

$$y_h = e^{2x} (\cos(x) c_1 + c_2 \sin(x))$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$3e^{-x} + 2x^2$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{-x}\}, \{1, x, x^2\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x)e^{2x}, \sin(x)e^{2x}\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1e^{-x} + A_2 + A_3x + A_4x^2$$

The unknowns $\{A_1, A_2, A_3, A_4\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$10A_1e^{-x} + 2A_4 - 4A_3 - 8A_4x + 5A_2 + 5A_3x + 5A_4x^2 = 3e^{-x} + 2x^2$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{3}{10}, A_2 = \frac{44}{125}, A_3 = \frac{16}{25}, A_4 = \frac{2}{5} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{3e^{-x}}{10} + \frac{44}{125} + \frac{16x}{25} + \frac{2x^2}{5}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (e^{2x}(\cos(x)c_1 + c_2 \sin(x))) + \left(\frac{3e^{-x}}{10} + \frac{44}{125} + \frac{16x}{25} + \frac{2x^2}{5} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = e^{2x}(\cos(x) c_1 + c_2 \sin(x)) + \frac{3e^{-x}}{10} + \frac{44}{125} + \frac{16x}{25} + \frac{2x^2}{5} \quad (1)$$




Figure 117: Slope field plot

Verification of solutions

$$y = e^{2x}(\cos(x) c_1 + c_2 \sin(x)) + \frac{3e^{-x}}{10} + \frac{44}{125} + \frac{16x}{25} + \frac{2x^2}{5}$$

Verified OK.

6.5.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 4y' + 5y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= 1 \\B &= -4 \\C &= 5\end{aligned}\tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x)\tag{4}$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\&= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned}\tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1}\tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= -1 \\t &= 1\end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x)\tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 114: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}
 y_1 &= z_1 e^{\int -\frac{1}{2} dx} \\
 &= z_1 e^{-\int \frac{1}{2} dx} \\
 &= z_1 e^{-x} \\
 &= z_1 (e^{-x})
 \end{aligned}$$

Which simplifies to

$$y_1 = \cos(x) e^{2x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{4}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{4x}}{(y_1)^2} dx \\ &= y_1(\tan(x)) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1(\cos(x) e^{2x}) + c_2(\cos(x) e^{2x}(\tan(x))) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 4y' + 5y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 \cos(x) e^{2x} + c_2 \sin(x) e^{2x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$3e^{-x} + 2x^2$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{-x}\}, \{1, x, x^2\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x) e^{2x}, \sin(x) e^{2x}\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^{-x} + A_2 + A_3 x + A_4 x^2$$

The unknowns $\{A_1, A_2, A_3, A_4\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$10A_1 e^{-x} + 2A_4 - 4A_3 - 8A_4 x + 5A_2 + 5A_3 x + 5A_4 x^2 = 3e^{-x} + 2x^2$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{3}{10}, A_2 = \frac{44}{125}, A_3 = \frac{16}{25}, A_4 = \frac{2}{5} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{3e^{-x}}{10} + \frac{44}{125} + \frac{16x}{25} + \frac{2x^2}{5}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(x) e^{2x} + c_2 \sin(x) e^{2x}) + \left(\frac{3e^{-x}}{10} + \frac{44}{125} + \frac{16x}{25} + \frac{2x^2}{5} \right) \end{aligned}$$

Which simplifies to

$$y = e^{2x}(\cos(x) c_1 + c_2 \sin(x)) + \frac{3e^{-x}}{10} + \frac{44}{125} + \frac{16x}{25} + \frac{2x^2}{5}$$

Summary

The solution(s) found are the following

$$y = e^{2x}(\cos(x) c_1 + c_2 \sin(x)) + \frac{3e^{-x}}{10} + \frac{44}{125} + \frac{16x}{25} + \frac{2x^2}{5} \quad (1)$$




Figure 118: Slope field plot

Verification of solutions

$$y = e^{2x}(\cos(x) c_1 + c_2 \sin(x)) + \frac{3e^{-x}}{10} + \frac{44}{125} + \frac{16x}{25} + \frac{2x^2}{5}$$

Verified OK.

6.5.3 Maple step by step solution

Let's solve

$$y'' - 4y' + 5y = 3e^{-x} + 2x^2$$

- Highest derivative means the order of the ODE is 2
- y''
- Characteristic polynomial of homogeneous ODE

$$r^2 - 4r + 5 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{4 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (2 - I, 2 + I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x) e^{2x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(x) e^{2x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = e^{2x} \cos(x) c_1 + e^{2x} \sin(x) c_2 + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = 3e^{-x} + 2x^2 \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) e^{2x} & \sin(x) e^{2x} \\ -\sin(x) e^{2x} + 2 \cos(x) e^{2x} & \cos(x) e^{2x} + 2 \sin(x) e^{2x} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = e^{4x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -e^{2x} (\cos(x) (\int \sin(x) e^{-2x} (3e^{-x} + 2x^2) dx)) - \sin(x) (\int \cos(x) e^{-2x} (3e^{-x} + 2x^2) dx)$$

- Compute integrals

$$y_p(x) = \frac{3e^{-x}}{10} + \frac{44}{125} + \frac{16x}{25} + \frac{2x^2}{5}$$

- Substitute particular solution into general solution to ODE

$$y = e^{2x} \cos(x) c_1 + e^{2x} \sin(x) c_2 + \frac{2x^2}{5} + \frac{3e^{-x}}{10} + \frac{16x}{25} + \frac{44}{125}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 36

```
dsolve(diff(y(x),x$2)-4*diff(y(x),x)+5*y(x)=3*exp(-x)+2*x^2,y(x), singsol=all)
```

$$y(x) = e^{2x} \sin(x) c_2 + e^{2x} \cos(x) c_1 + \frac{3e^{-x}}{10} + \frac{2x^2}{5} + \frac{16x}{25} + \frac{44}{125}$$

✓ Solution by Mathematica

Time used: 0.316 (sec). Leaf size: 47

```
DSolve[y''[x]-4*y'[x]+5*y[x]==3*Exp[-x]+2*x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{250}(100x^2 + 160x + 75e^{-x} + 88) + c_2 e^{2x} \cos(x) + c_1 e^{2x} \sin(x)$$

6.6 problem 1(f)

6.6.1	Solving as second order linear constant coeff ode	562
6.6.2	Solving using Kovacic algorithm	565
6.6.3	Maple step by step solution	570

Internal problem ID [5966]

Internal file name [OUTPUT/5214_Sunday_June_05_2022_03_27_39_PM_49247672/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 69

Problem number: 1(f).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"kovacic", "second_order_linear_constant_coeff"**

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' - 7y' + 6y = \sin(x)$$

6.6.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = -7, C = 6, f(x) = \sin(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 7y' + 6y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -7, C = 6$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 7\lambda e^{\lambda x} + 6 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 7\lambda + 6 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -7, C = 6$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{7}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-7^2 - (4)(1)(6)} \\ &= \frac{7}{2} \pm \frac{5}{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= \frac{7}{2} + \frac{5}{2} \\ \lambda_2 &= \frac{7}{2} - \frac{5}{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= 6 \\ \lambda_2 &= 1 \end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned} y &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ y &= c_1 e^{(6)x} + c_2 e^{(1)x} \end{aligned}$$

Or

$$y = c_1 e^{6x} + c_2 e^x$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^{6x} + c_2 e^x$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^x, e^{6x}\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(x) + A_2 \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$5A_1 \cos(x) + 5A_2 \sin(x) + 7A_1 \sin(x) - 7A_2 \cos(x) = \sin(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{7}{74}, A_2 = \frac{5}{74} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{7 \cos(x)}{74} + \frac{5 \sin(x)}{74}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^{6x} + c_2 e^x) + \left(\frac{7 \cos(x)}{74} + \frac{5 \sin(x)}{74} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{6x} + c_2 e^x + \frac{7 \cos(x)}{74} + \frac{5 \sin(x)}{74} \quad (1)$$




Figure 119: Slope field plot

Verification of solutions

$$y = c_1 e^{6x} + c_2 e^x + \frac{7 \cos(x)}{74} + \frac{5 \sin(x)}{74}$$

Verified OK.

6.6.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 7y' + 6y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -7 \\ C &= 6 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{25}{4} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 25 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \frac{25z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 116: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{25}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{5x}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-7}{1} dx} \\ &= z_1 e^{\frac{7x}{2}} \\ &= z_1 \left(e^{\frac{7x}{2}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = e^x$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-7}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{7x}}{(y_1)^2} dx \\ &= y_1 \left(\frac{e^{5x}}{5} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1(e^x) + c_2\left(e^x\left(\frac{e^{5x}}{5}\right)\right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 7y' + 6y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^x + \frac{c_2 e^{6x}}{5}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{\frac{e^{6x}}{5}, e^x\right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(x) + A_2 \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$5A_1 \cos(x) + 5A_2 \sin(x) + 7A_1 \sin(x) - 7A_2 \cos(x) = \sin(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{7}{74}, A_2 = \frac{5}{74} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{7 \cos(x)}{74} + \frac{5 \sin(x)}{74}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 e^x + \frac{c_2 e^{6x}}{5} \right) + \left(\frac{7 \cos(x)}{74} + \frac{5 \sin(x)}{74} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^x + \frac{c_2 e^{6x}}{5} + \frac{7 \cos(x)}{74} + \frac{5 \sin(x)}{74} \quad (1)$$




Figure 120: Slope field plot

Verification of solutions

$$y = c_1 e^x + \frac{c_2 e^{6x}}{5} + \frac{7 \cos(x)}{74} + \frac{5 \sin(x)}{74}$$

Verified OK.

6.6.3 Maple step by step solution

Let's solve

$$y'' - 7y' + 6y = \sin(x)$$

- Highest derivative means the order of the ODE is 2
- y''
- Characteristic polynomial of homogeneous ODE
- $r^2 - 7r + 6 = 0$
- Factor the characteristic polynomial
- $(r - 1)(r - 6) = 0$
- Roots of the characteristic polynomial

$$r = (1, 6)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^x$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{6x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^x + c_2 e^{6x} + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \sin(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^x & e^{6x} \\ e^x & 6e^{6x} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 5e^{7x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{e^x \left(\int e^{-x} \sin(x) dx \right)}{5} + \frac{e^{6x} \left(\int \sin(x) e^{-6x} dx \right)}{5}$$

- Compute integrals

$$y_p(x) = \frac{7 \cos(x)}{74} + \frac{5 \sin(x)}{74}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^x + c_2 e^{6x} + \frac{7 \cos(x)}{74} + \frac{5 \sin(x)}{74}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(diff(y(x),x$2)-7*diff(y(x),x)+6*y(x)=sin(x),y(x), singsol=all)
```

$$y(x) = e^{6x}c_2 + e^x c_1 + \frac{7 \cos(x)}{74} + \frac{5 \sin(x)}{74}$$

✓ Solution by Mathematica

Time used: 0.063 (sec). Leaf size: 32

```
DSolve[y''[x]-7*y'[x]+6*y[x]==Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{5 \sin(x)}{74} + \frac{7 \cos(x)}{74} + c_1 e^x + c_2 e^{6x}$$

6.7 problem 1(g)

6.7.1	Solving as second order linear constant coeff ode	573
6.7.2	Solving using Kovacic algorithm	577
6.7.3	Maple step by step solution	581

Internal problem ID [5967]

Internal file name [OUTPUT/5215_Sunday_June_05_2022_03_27_40_PM_60845922/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 69

Problem number: 1(g).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + y = 2 \sin(x) \sin(2x)$$

6.7.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 1, f(x) = \cos(x) - \cos(3x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\lambda_1 = +i$$

$$\lambda_2 = -i$$

Which simplifies to

$$\lambda_1 = i$$

$$\lambda_2 = -i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (\cos(x) c_1 + c_2 \sin(x))$$

Or

$$y = \cos(x) c_1 + c_2 \sin(x)$$

Therefore the homogeneous solution y_h is

$$y_h = \cos(x) c_1 + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\cos(x) - \cos(3x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}, \{\cos(3x), \sin(3x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since $\cos(x)$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{\cos(x)x, \sin(x)x\}, \{\cos(3x), \sin(3x)\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 \cos(x)x + A_2 \sin(x)x + A_3 \cos(3x) + A_4 \sin(3x)$$

The unknowns $\{A_1, A_2, A_3, A_4\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-2A_1 \sin(x) + 2A_2 \cos(x) - 8A_3 \cos(3x) - 8A_4 \sin(3x) = \cos(x) - \cos(3x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = 0, A_2 = \frac{1}{2}, A_3 = \frac{1}{8}, A_4 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{\sin(x)x}{2} + \frac{\cos(3x)}{8}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (\cos(x) c_1 + c_2 \sin(x)) + \left(\frac{\sin(x) x}{2} + \frac{\cos(3x)}{8} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \cos(x) c_1 + c_2 \sin(x) + \frac{\sin(x) x}{2} + \frac{\cos(3x)}{8} \quad (1)$$




Figure 121: Slope field plot

Verification of solutions

$$y = \cos(x) c_1 + c_2 \sin(x) + \frac{\sin(x) x}{2} + \frac{\cos(3x)}{8}$$

Verified OK.

6.7.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 1 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 118: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}y_1 &= z_1 \\ &= \cos(x)\end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned}y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x))\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(x)) + c_2 (\cos(x) (\tan(x)))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = \cos(x) c_1 + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$4 \cos(x) \sin(x)^2$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}, \{\cos(3x), \sin(3x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since $\cos(x)$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{\cos(x)x, \sin(x)x\}, \{\cos(3x), \sin(3x)\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 \cos(x)x + A_2 \sin(x)x + A_3 \cos(3x) + A_4 \sin(3x)$$

The unknowns $\{A_1, A_2, A_3, A_4\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-2A_1 \sin(x) + 2A_2 \cos(x) - 8A_3 \cos(3x) - 8A_4 \sin(3x) = \cos(x) - \cos(3x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = 0, A_2 = \frac{1}{2}, A_3 = \frac{1}{8}, A_4 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{\sin(x)x}{2} + \frac{\cos(3x)}{8}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (\cos(x)c_1 + c_2 \sin(x)) + \left(\frac{\sin(x)x}{2} + \frac{\cos(3x)}{8} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \cos(x) c_1 + c_2 \sin(x) + \frac{\sin(x) x}{2} + \frac{\cos(3x)}{8} \quad (1)$$




Figure 122: Slope field plot

Verification of solutions

$$y = \cos(x) c_1 + c_2 \sin(x) + \frac{\sin(x) x}{2} + \frac{\cos(3x)}{8}$$

Verified OK.

6.7.3 Maple step by step solution

Let's solve

$$y'' + y = \cos(x) - \cos(3x)$$

- Highest derivative means the order of the ODE is 2
- y''
- Characteristic polynomial of homogeneous ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (-I, I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = \cos(x) c_1 + c_2 \sin(x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x), y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x), y_2(x))} dx \right), f(x) = \cos(x) - \cos(3x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -4 \cos(x) \left(\int \sin(x)^3 \cos(x) dx \right) + \frac{\sin(x) \left(\int (1 - \cos(4x)) dx \right)}{2}$$

- Compute integrals

$$y_p(x) = \frac{\sin(x)(-\cos(x)\sin(x)+x)}{2}$$

- Substitute particular solution into general solution to ODE

$$y = \cos(x) c_1 + c_2 \sin(x) + \frac{\sin(x)(-\cos(x)\sin(x)+x)}{2}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(diff(y(x),x$2)+y(x)=2*sin(x)*sin(2*x),y(x), singsol=all)
```

$$y(x) = -\frac{\cos(x) \sin(x)^2}{2} + \frac{(2c_2 + x) \sin(x)}{2} + \cos(x) c_1$$

✓ Solution by Mathematica

Time used: 0.033 (sec). Leaf size: 33

```
DSolve[y''[x]+y[x]==2*Sin[x]*Sin[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{8}(\cos(3x) + (-1 + 8c_1) \cos(x) + 4(x + 2c_2) \sin(x))$$

6.8 problem 1(h)

6.8.1	Solving as second order linear constant coeff ode	584
6.8.2	Solving using Kovacic algorithm	588
6.8.3	Maple step by step solution	594

Internal problem ID [5968]

Internal file name [OUTPUT/5216_Sunday_June_05_2022_03_27_42_PM_83336233/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 69

Problem number: 1(h).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + y = \sec(x)$$

6.8.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 1, f(x) = \sec(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\lambda_1 = +i$$

$$\lambda_2 = -i$$

Which simplifies to

$$\lambda_1 = i$$

$$\lambda_2 = -i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (\cos(x) c_1 + c_2 \sin(x))$$

Or

$$y = \cos(x) c_1 + c_2 \sin(x)$$

Therefore the homogeneous solution y_h is

$$y_h = \cos(x) c_1 + c_2 \sin(x)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \cos(x)$$

$$y_2 = \sin(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ \frac{d}{dx}(\cos(x)) & \frac{d}{dx}(\sin(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{vmatrix}$$

Therefore

$$W = (\cos(x))(\cos(x)) - (\sin(x))(-\sin(x))$$

Which simplifies to

$$W = \cos(x)^2 + \sin(x)^2$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\sin(x) \sec(x)}{1} dx$$

Which simplifies to

$$u_1 = - \int \tan(x) dx$$

Hence

$$u_1 = \ln(\cos(x))$$

And Eq. (3) becomes

$$u_2 = \int \frac{\sec(x) \cos(x)}{1} dx$$

Which simplifies to

$$u_2 = \int 1 dx$$

Hence

$$u_2 = x$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \ln(\cos(x)) \cos(x) + \sin(x) x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (\cos(x) c_1 + c_2 \sin(x)) + (\ln(\cos(x)) \cos(x) + \sin(x) x) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \cos(x) c_1 + c_2 \sin(x) + \ln(\cos(x)) \cos(x) + \sin(x) x \quad (1)$$




Figure 123: Slope field plot

Verification of solutions

$$y = \cos(x) c_1 + c_2 \sin(x) + \ln(\cos(x)) \cos(x) + \sin(x) x$$

Verified OK.

6.8.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 0 \quad (3)$$

$$C = 1$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 120: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}
 y_1 &= z_1 \\
 &= \cos(x)
 \end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x)) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1(\cos(x)) + c_2(\cos(x) (\tan(x))) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = \cos(x) c_1 + c_2 \sin(x)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of

parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \cos(x)$$

$$y_2 = \sin(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ \frac{d}{dx}(\cos(x)) & \frac{d}{dx}(\sin(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{vmatrix}$$

Therefore

$$W = (\cos(x))(\cos(x)) - (\sin(x))(-\sin(x))$$

Which simplifies to

$$W = \cos(x)^2 + \sin(x)^2$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\sin(x) \sec(x)}{1} dx$$

Which simplifies to

$$u_1 = - \int \tan(x) dx$$

Hence

$$u_1 = \ln(\cos(x))$$

And Eq. (3) becomes

$$u_2 = \int \frac{\sec(x) \cos(x)}{1} dx$$

Which simplifies to

$$u_2 = \int 1 dx$$

Hence

$$u_2 = x$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \ln(\cos(x)) \cos(x) + \sin(x) x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (\cos(x) c_1 + c_2 \sin(x)) + (\ln(\cos(x)) \cos(x) + \sin(x) x) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \cos(x) c_1 + c_2 \sin(x) + \ln(\cos(x)) \cos(x) + \sin(x) x \quad (1)$$




Figure 124: Slope field plot

Verification of solutions

$$y = \cos(x) c_1 + c_2 \sin(x) + \ln(\cos(x)) \cos(x) + \sin(x) x$$

Verified OK.

6.8.3 Maple step by step solution

Let's solve

$$y'' + y = \sec(x)$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (-I, I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = \cos(x) c_1 + c_2 \sin(x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \sec(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\cos(x) \left(\int \tan(x) dx \right) + \sin(x) \left(\int 1 dx \right)$$

- Compute integrals

$$y_p(x) = \ln(\cos(x)) \cos(x) + \sin(x) x$$

- Substitute particular solution into general solution to ODE

$$y = \cos(x) c_1 + c_2 \sin(x) + \ln(\cos(x)) \cos(x) + \sin(x) x$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 22

```
dsolve(diff(y(x),x$2)+y(x)=sec(x),y(x), singsol=all)
```

$$y(x) = -\ln(\sec(x)) \cos(x) + \cos(x) c_1 + \sin(x) (x + c_2)$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 22

```
DSolve[y''[x]+y[x]==Sec[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow (x + c_2) \sin(x) + \cos(x) (\log(\cos(x)) + c_1)$$

6.9 problem 1(i)

6.9.1	Solving as second order linear constant coeff ode	597
6.9.2	Solving using Kovacic algorithm	600
6.9.3	Maple step by step solution	605

Internal problem ID [5969]

Internal file name [OUTPUT/5217_Sunday_June_05_2022_03_27_43_PM_95613046/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 69

Problem number: 1(i).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$4y'' - y = e^x$$

6.9.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 4, B = 0, C = -1, f(x) = e^x$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$4y'' - y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 4, B = 0, C = -1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$4\lambda^2 e^{\lambda x} - e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$4\lambda^2 - 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 4, B = 0, C = -1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(4)} \pm \frac{1}{(2)(4)} \sqrt{0^2 - (4)(4)(-1)} \\ &= \pm \frac{1}{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +\frac{1}{2} \\ \lambda_2 &= -\frac{1}{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= \frac{1}{2} \\ \lambda_2 &= -\frac{1}{2} \end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned} y &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ y &= c_1 e^{(\frac{1}{2})x} + c_2 e^{(-\frac{1}{2})x} \end{aligned}$$

Or

$$y = c_1 e^{\frac{x}{2}} + c_2 e^{-\frac{x}{2}}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^{\frac{x}{2}} + c_2 e^{-\frac{x}{2}}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$e^x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[e^x]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^{-\frac{x}{2}}, e^{\frac{x}{2}}\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^x$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$3A_1 e^x = e^x$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{3} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{e^x}{3}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^{\frac{x}{2}} + c_2 e^{-\frac{x}{2}}) + \left(\frac{e^x}{3} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{\frac{x}{2}} + c_2 e^{-\frac{x}{2}} + \frac{e^x}{3} \quad (1)$$




Figure 125: Slope field plot

Verification of solutions

$$y = c_1 e^{\frac{x}{2}} + c_2 e^{-\frac{x}{2}} + \frac{e^x}{3}$$

Verified OK.

6.9.2 Solving using Kovacic algorithm

Writing the ode as

$$4y'' - y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 4 \\ B &= 0 \\ C &= -1 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{1}{4} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 1 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \frac{z(x)}{4} \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 122: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{1}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{x}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned} y_1 &= z_1 \\ &= e^{-\frac{x}{2}} \end{aligned}$$

Which simplifies to

$$y_1 = e^{-\frac{x}{2}}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= e^{-\frac{x}{2}} \int \frac{1}{e^{-x}} dx \\ &= e^{-\frac{x}{2}} (e^x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-\frac{x}{2}}) + c_2 (e^{-\frac{x}{2}} (e^x))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$4y'' - y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-\frac{x}{2}} + c_2 e^{\frac{x}{2}}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$e^x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^{-\frac{x}{2}}, e^{\frac{x}{2}}\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^x$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$3A_1 e^x = e^x$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{3} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{e^x}{3}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^{-\frac{x}{2}} + c_2 e^{\frac{x}{2}}) + \left(\frac{e^x}{3} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-\frac{x}{2}} + c_2 e^{\frac{x}{2}} + \frac{e^x}{3} \tag{1}$$




Figure 126: Slope field plot

Verification of solutions

$$y = c_1 e^{-\frac{x}{2}} + c_2 e^{\frac{x}{2}} + \frac{e^x}{3}$$

Verified OK.

6.9.3 Maple step by step solution

Let's solve

$$4y'' - y = e^x$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = \frac{y}{4} + \frac{e^x}{4}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - \frac{y}{4} = \frac{e^x}{4}$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - \frac{1}{4} = 0$$

- Factor the characteristic polynomial

$$\frac{(2r-1)(2r+1)}{4} = 0$$

- Roots of the characteristic polynomial

$$r = \left(-\frac{1}{2}, \frac{1}{2}\right)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-\frac{x}{2}}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{\frac{x}{2}}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{-\frac{x}{2}} + c_2 e^{\frac{x}{2}} + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \frac{e^x}{4} \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-\frac{x}{2}} & e^{\frac{x}{2}} \\ -\frac{e^{-\frac{x}{2}}}{2} & \frac{e^{\frac{x}{2}}}{2} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{e^{-\frac{x}{2}} \left(\int e^{\frac{3x}{2}} dx \right)}{4} + \frac{e^{\frac{x}{2}} \left(\int e^{\frac{x}{2}} dx \right)}{4}$$

- Compute integrals

$$y_p(x) = \frac{e^x}{3}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{-\frac{x}{2}} + c_2 e^{\frac{x}{2}} + \frac{e^x}{3}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 21

```
dsolve(4*diff(y(x),x$2)-y(x)=exp(x),y(x), singsol=all)
```

$$y(x) = e^{\frac{x}{2}} c_2 + c_1 e^{-\frac{x}{2}} + \frac{e^x}{3}$$

✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 33

```
DSolve[4*y''[x]-y[x]==Exp[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^x}{3} + c_1 e^{x/2} + c_2 e^{-x/2}$$

6.10 problem 1(j)

6.10.1 Solving as second order linear constant coeff ode	608
6.10.2 Solving using Kovacic algorithm	611
6.10.3 Maple step by step solution	616

Internal problem ID [5970]

Internal file name [OUTPUT/5218_Sunday_June_05_2022_03_27_45_PM_87728478/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 69

Problem number: 1(j).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"kovacic", "second_order_linear_constant_coeff"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$6y'' + 5y' - 6y = x$$

6.10.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 6, B = 5, C = -6, f(x) = x$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$6y'' + 5y' - 6y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 6, B = 5, C = -6$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$6\lambda^2 e^{\lambda x} + 5\lambda e^{\lambda x} - 6e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$6\lambda^2 + 5\lambda - 6 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 6, B = 5, C = -6$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-5}{(2)(6)} \pm \frac{1}{(2)(6)} \sqrt{5^2 - (4)(6)(-6)} \\ &= -\frac{5}{12} \pm \frac{13}{12} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= -\frac{5}{12} + \frac{13}{12} \\ \lambda_2 &= -\frac{5}{12} - \frac{13}{12} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= \frac{2}{3} \\ \lambda_2 &= -\frac{3}{2} \end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned} y &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ y &= c_1 e^{(\frac{2}{3})x} + c_2 e^{(-\frac{3}{2})x} \end{aligned}$$

Or

$$y = e^{\frac{2x}{3}} c_1 + c_2 e^{-\frac{3x}{2}}$$

Therefore the homogeneous solution y_h is

$$y_h = e^{\frac{2x}{3}} c_1 + c_2 e^{-\frac{3x}{2}}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{-\frac{3x}{2}}, e^{\frac{2x}{3}} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_2x + A_1$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-6A_2x - 6A_1 + 5A_2 = x$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{5}{36}, A_2 = -\frac{1}{6} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{x}{6} - \frac{5}{36}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(e^{\frac{2x}{3}} c_1 + c_2 e^{-\frac{3x}{2}} \right) + \left(-\frac{x}{6} - \frac{5}{36} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = e^{\frac{2x}{3}} c_1 + c_2 e^{-\frac{3x}{2}} - \frac{x}{6} - \frac{5}{36} \quad (1)$$




Figure 127: Slope field plot

Verification of solutions

$$y = e^{\frac{2x}{3}} c_1 + c_2 e^{-\frac{3x}{2}} - \frac{x}{6} - \frac{5}{36}$$

Verified OK.

6.10.2 Solving using Kovacic algorithm

Writing the ode as

$$6y'' + 5y' - 6y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 6 \\ B &= 5 \\ C &= -6 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{169}{144} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 169 \\ t &= 144 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \frac{169z(x)}{144} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 124: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{169}{144}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{13x}{12}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{5}{6} dx} \\ &= z_1 e^{-\frac{5x}{12}} \\ &= z_1 \left(e^{-\frac{5x}{12}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-\frac{3x}{2}}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{5}{6} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-\frac{5x}{6}}}{(y_1)^2} dx \\ &= y_1 \left(\frac{6 e^{\frac{13x}{6}}}{13} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(e^{-\frac{3x}{2}} \right) + c_2 \left(e^{-\frac{3x}{2}} \left(\frac{6 e^{\frac{13x}{6}}}{13} \right) \right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$6y'' + 5y' - 6y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-\frac{3x}{2}} + \frac{6c_2 e^{\frac{2x}{3}}}{13}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ \frac{6 e^{\frac{2x}{3}}}{13}, e^{-\frac{3x}{2}} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_2 x + A_1$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-6A_2x - 6A_1 + 5A_2 = x$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{5}{36}, A_2 = -\frac{1}{6} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{x}{6} - \frac{5}{36}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 e^{-\frac{3x}{2}} + \frac{6c_2 e^{\frac{2x}{3}}}{13} \right) + \left(-\frac{x}{6} - \frac{5}{36} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-\frac{3x}{2}} + \frac{6c_2 e^{\frac{2x}{3}}}{13} - \frac{x}{6} - \frac{5}{36} \quad (1)$$




Figure 128: Slope field plot

Verification of solutions

$$y = c_1 e^{-\frac{3x}{2}} + \frac{6c_2 e^{\frac{2x}{3}}}{13} - \frac{x}{6} - \frac{5}{36}$$

Verified OK.

6.10.3 Maple step by step solution

Let's solve

$$6y'' + 5y' - 6y = x$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{5y'}{6} + y + \frac{x}{6}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{5y'}{6} - y = \frac{x}{6}$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + \frac{5}{6}r - 1 = 0$$

- Factor the characteristic polynomial

$$\frac{(2r+3)(3r-2)}{6} = 0$$

- Roots of the characteristic polynomial

$$r = \left(-\frac{3}{2}, \frac{2}{3}\right)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-\frac{3x}{2}}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{\frac{2x}{3}}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{-\frac{3x}{2}} + c_2 e^{\frac{2x}{3}} + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \frac{x}{6} \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-\frac{3x}{2}} & e^{\frac{2x}{3}} \\ -\frac{3}{2}e^{-\frac{3x}{2}} & \frac{2}{3}e^{\frac{2x}{3}} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = \frac{13e^{-\frac{5x}{6}}}{6}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = \frac{\left(e^{\frac{13x}{6}} \left(\int x e^{-\frac{2x}{3}} dx \right) - \left(\int x e^{\frac{3x}{2}} dx \right) \right) e^{-\frac{3x}{2}}}{13}$$

- Compute integrals

$$y_p(x) = -\frac{x}{6} - \frac{5}{36}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{-\frac{3x}{2}} + c_2 e^{\frac{2x}{3}} - \frac{x}{6} - \frac{5}{36}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
dsolve(6*diff(y(x),x$2)+5*diff(y(x),x)-6*y(x)=x,y(x), singsol=all)
```

$$y(x) = -\frac{\left(\left(x + \frac{5}{6}\right) e^{\frac{3x}{2}} - 6 e^{\frac{13x}{6}} c_2 - 6c_1\right) e^{-\frac{3x}{2}}}{6}$$

✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 34

```
DSolve[6*y''[x]+5*y'[x]-6*y[x]==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{x}{6} + c_1 e^{2x/3} + c_2 e^{-3x/2} - \frac{5}{36}$$

6.11 problem 4(c)

6.11.1 Existence and uniqueness analysis	619
6.11.2 Solving as second order linear constant coeff ode	620
6.11.3 Solving using Kovacic algorithm	625
6.11.4 Maple step by step solution	632

Internal problem ID [5971]

Internal file name [OUTPUT/5219_Sunday_June_05_2022_03_27_46_PM_50415/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 69

Problem number: 4(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"kovacic", "second_order_linear_constant_coeff"**

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + \omega^2 y = A \cos(\omega x)$$

With initial conditions

$$[y(0) = 0, y'(0) = 1]$$

6.11.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = 0$$

$$q(x) = \omega^2$$

$$F = A \cos(\omega x)$$

Hence the ode is

$$y'' + \omega^2 y = A \cos(\omega x)$$

The domain of $p(x) = 0$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = \omega^2$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. The domain of $F = A \cos(\omega x)$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

6.11.2 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = \omega^2, f(x) = A \cos(\omega x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + \omega^2 y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = \omega^2$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + \omega^2 e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + \omega^2 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = \omega^2$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(\omega^2)} \\ &= \pm \sqrt{-\omega^2} \end{aligned}$$

Hence

$$\lambda_1 = +\sqrt{-\omega^2}$$

$$\lambda_2 = -\sqrt{-\omega^2}$$

Which simplifies to

$$\lambda_1 = \sqrt{-\omega^2}$$

$$\lambda_2 = -\sqrt{-\omega^2}$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(\sqrt{-\omega^2})x} + c_2 e^{(-\sqrt{-\omega^2})x}$$

Or

$$y = c_1 e^{\sqrt{-\omega^2} x} + c_2 e^{-\sqrt{-\omega^2} x}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^{\sqrt{-\omega^2} x} + c_2 e^{-\sqrt{-\omega^2} x}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = e^{\sqrt{-\omega^2} x}$$

$$y_2 = e^{-\sqrt{-\omega^2} x}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^{\sqrt{-\omega^2} x} & e^{-\sqrt{-\omega^2} x} \\ \frac{d}{dx} \left(e^{\sqrt{-\omega^2} x} \right) & \frac{d}{dx} \left(e^{-\sqrt{-\omega^2} x} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^{\sqrt{-\omega^2} x} & e^{-\sqrt{-\omega^2} x} \\ \sqrt{-\omega^2} e^{\sqrt{-\omega^2} x} & -\sqrt{-\omega^2} e^{-\sqrt{-\omega^2} x} \end{vmatrix}$$

Therefore

$$W = \left(e^{\sqrt{-\omega^2} x} \right) \left(-\sqrt{-\omega^2} e^{-\sqrt{-\omega^2} x} \right) - \left(e^{-\sqrt{-\omega^2} x} \right) \left(\sqrt{-\omega^2} e^{\sqrt{-\omega^2} x} \right)$$

Which simplifies to

$$W = -2 e^{\sqrt{-\omega^2} x} \sqrt{-\omega^2} e^{-\sqrt{-\omega^2} x}$$

Which simplifies to

$$W = -2\sqrt{-\omega^2}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{e^{-\sqrt{-\omega^2} x} A \cos(\omega x)}{-2\sqrt{-\omega^2}} dx$$

Which simplifies to

$$u_1 = - \int - \frac{e^{-\sqrt{-\omega^2} x} A \cos(\omega x)}{2\sqrt{-\omega^2}} dx$$

Hence

$$u_1 = \frac{-\frac{A e^{-\sqrt{-\omega^2} x}}{4\omega} - \frac{A x e^{-\sqrt{-\omega^2} x} \tan\left(\frac{\omega x}{2}\right)}{2} + \frac{A \sqrt{-\omega^2} x e^{-\sqrt{-\omega^2} x}}{4\omega} + \frac{A e^{-\sqrt{-\omega^2} x} \tan\left(\frac{\omega x}{2}\right)^2}{4\omega} - \frac{A \sqrt{-\omega^2} x e^{-\sqrt{-\omega^2} x} \tan\left(\frac{\omega x}{2}\right)^2}{4\omega}}{\omega \left(1 + \tan\left(\frac{\omega x}{2}\right)^2\right)}$$

And Eq. (3) becomes

$$u_2 = \int \frac{e^{\sqrt{-\omega^2} x} A \cos(\omega x)}{-2\sqrt{-\omega^2}} dx$$

Which simplifies to

$$u_2 = \int - \frac{e^{\sqrt{-\omega^2} x} A \cos(\omega x)}{2\sqrt{-\omega^2}} dx$$

Hence

$$u_2 = \frac{\frac{A e^{\sqrt{-\omega^2} x}}{4\omega} + \frac{A x e^{\sqrt{-\omega^2} x} \tan\left(\frac{\omega x}{2}\right)}{2} - \frac{A e^{\sqrt{-\omega^2} x} \tan\left(\frac{\omega x}{2}\right)^2}{4\omega} + \frac{A \sqrt{-\omega^2} x e^{\sqrt{-\omega^2} x}}{4\omega} - \frac{A \sqrt{-\omega^2} x e^{\sqrt{-\omega^2} x} \tan\left(\frac{\omega x}{2}\right)^2}{4\omega}}{\omega \left(1 + \tan\left(\frac{\omega x}{2}\right)^2\right)}$$

Which simplifies to

$$u_1 = \frac{A e^{-\sqrt{-\omega^2} x} (-\sqrt{-\omega^2} x \cos(\omega x) + x\omega \sin(\omega x) + \cos(\omega x))}{4\omega^2}$$

$$u_2 = \frac{A e^{\sqrt{-\omega^2} x} (\sqrt{-\omega^2} x \cos(\omega x) + x\omega \sin(\omega x) + \cos(\omega x))}{4\omega^2}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{A e^{-\sqrt{-\omega^2} x} (-\sqrt{-\omega^2} x \cos(\omega x) + x\omega \sin(\omega x) + \cos(\omega x)) e^{\sqrt{-\omega^2} x}}{4\omega^2} + \frac{A e^{\sqrt{-\omega^2} x} (\sqrt{-\omega^2} x \cos(\omega x) + x\omega \sin(\omega x) + \cos(\omega x)) e^{-\sqrt{-\omega^2} x}}{4\omega^2}$$

Which simplifies to

$$y_p(x) = \frac{A(x\omega \sin(\omega x) + \cos(\omega x))}{2\omega^2}$$

Therefore the general solution is

$$y = y_h + y_p = (c_1 e^{\sqrt{-\omega^2} x} + c_2 e^{-\sqrt{-\omega^2} x}) + \left(\frac{A(x\omega \sin(\omega x) + \cos(\omega x))}{2\omega^2} \right)$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 e^{\sqrt{-\omega^2} x} + c_2 e^{-\sqrt{-\omega^2} x} + \frac{A(x\omega \sin(\omega x) + \cos(\omega x))}{2\omega^2} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = \frac{(2c_1 + 2c_2)\omega^2 + A}{2\omega^2} \quad (1A)$$

Taking derivative of the solution gives

$$y' = c_1 \sqrt{-\omega^2} e^{\sqrt{-\omega^2} x} - c_2 \sqrt{-\omega^2} e^{-\sqrt{-\omega^2} x} + \frac{Ax \cos(\omega x)}{2}$$

substituting $y' = 1$ and $x = 0$ in the above gives

$$1 = (c_1 - c_2) \sqrt{-\omega^2} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = -\frac{A + 2\sqrt{-\omega^2}}{4\omega^2}$$

$$c_2 = -\frac{A - 2\sqrt{-\omega^2}}{4\omega^2}$$

Substituting these values back in above solution results in

$$y = \frac{2A \sin(\omega x) \omega x + 2A \cos(\omega x) - A e^{\sqrt{-\omega^2} x} - A e^{-\sqrt{-\omega^2} x} - 2\sqrt{-\omega^2} e^{\sqrt{-\omega^2} x} + 2\sqrt{-\omega^2} e^{-\sqrt{-\omega^2} x}}{4\omega^2}$$

Which simplifies to

$$y = \frac{(-A + 2\sqrt{-\omega^2}) e^{-\sqrt{-\omega^2} x} + (-A - 2\sqrt{-\omega^2}) e^{\sqrt{-\omega^2} x} + 2A(x\omega \sin(\omega x) + \cos(\omega x))}{4\omega^2}$$

Summary

The solution(s) found are the following

$$y = \frac{(-A + 2\sqrt{-\omega^2}) e^{-\sqrt{-\omega^2} x} + (-A - 2\sqrt{-\omega^2}) e^{\sqrt{-\omega^2} x} + 2A(x\omega \sin(\omega x) + \cos(\omega x))}{4\omega^2} \quad (1)$$

Verification of solutions

$$y = \frac{(-A + 2\sqrt{-\omega^2}) e^{-\sqrt{-\omega^2} x} + (-A - 2\sqrt{-\omega^2}) e^{\sqrt{-\omega^2} x} + 2A(x\omega \sin(\omega x) + \cos(\omega x))}{4\omega^2}$$

Verified OK.

6.11.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + \omega^2 y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 0 \quad (3)$$

$$C = \omega^2$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$r = \frac{s}{t} \tag{5}$$

$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-\omega^2}{1} \tag{6}$$

Comparing the above to (5) shows that

$$s = -\omega^2$$

$$t = 1$$

Therefore eq. (4) becomes

$$z''(x) = (-\omega^2) z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 126: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -\omega^2$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{\sqrt{-\omega^2} x}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned} y_1 &= z_1 \\ &= e^{\sqrt{-\omega^2} x} \end{aligned}$$

Which simplifies to

$$y_1 = e^{\sqrt{-\omega^2} x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= e^{\sqrt{-\omega^2} x} \int \frac{1}{e^{2\sqrt{-\omega^2} x}} dx \\ &= e^{\sqrt{-\omega^2} x} \left(\frac{\sqrt{-\omega^2} e^{-2\sqrt{-\omega^2} x}}{2\omega^2} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(e^{\sqrt{-\omega^2} x} \right) + c_2 \left(e^{\sqrt{-\omega^2} x} \left(\frac{\sqrt{-\omega^2} e^{-2\sqrt{-\omega^2} x}}{2\omega^2} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + \omega^2 y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{\sqrt{-\omega^2} x} + \frac{c_2 e^{-\sqrt{-\omega^2} x} \sqrt{-\omega^2}}{2\omega^2}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned} y_1 &= e^{\sqrt{-\omega^2} x} \\ y_2 &= \frac{e^{-\sqrt{-\omega^2} x} \sqrt{-\omega^2}}{2\omega^2} \end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^{\sqrt{-\omega^2} x} & \frac{e^{-\sqrt{-\omega^2} x} \sqrt{-\omega^2}}{2\omega^2} \\ \frac{d}{dx} \left(e^{\sqrt{-\omega^2} x} \right) & \frac{d}{dx} \left(\frac{e^{-\sqrt{-\omega^2} x} \sqrt{-\omega^2}}{2\omega^2} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^{\sqrt{-\omega^2} x} & \frac{e^{-\sqrt{-\omega^2} x} \sqrt{-\omega^2}}{2\omega^2} \\ \sqrt{-\omega^2} e^{\sqrt{-\omega^2} x} & \frac{e^{-\sqrt{-\omega^2} x}}{2} \end{vmatrix}$$

Therefore

$$W = \left(e^{\sqrt{-\omega^2} x} \right) \left(\frac{e^{-\sqrt{-\omega^2} x}}{2} \right) - \left(\frac{e^{-\sqrt{-\omega^2} x} \sqrt{-\omega^2}}{2\omega^2} \right) \left(\sqrt{-\omega^2} e^{\sqrt{-\omega^2} x} \right)$$

Which simplifies to

$$W = e^{\sqrt{-\omega^2} x} e^{-\sqrt{-\omega^2} x}$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{e^{-\sqrt{-\omega^2} x} \sqrt{-\omega^2} A \cos(\omega x)}{\frac{2\omega^2}{1}} dx$$

Which simplifies to

$$u_1 = - \int \frac{e^{-\sqrt{-\omega^2} x} \sqrt{-\omega^2} A \cos(\omega x)}{2\omega^2} dx$$

Hence

$$u_1 = \frac{-\frac{A e^{-\sqrt{-\omega^2} x}}{4\omega} - \frac{A x e^{-\sqrt{-\omega^2} x} \tan\left(\frac{\omega x}{2}\right)}{2} + \frac{A \sqrt{-\omega^2} x e^{-\sqrt{-\omega^2} x}}{4\omega} + \frac{A e^{-\sqrt{-\omega^2} x} \tan\left(\frac{\omega x}{2}\right)^2}{4\omega} - \frac{A \sqrt{-\omega^2} x e^{-\sqrt{-\omega^2} x} \tan\left(\frac{\omega x}{2}\right)^2}{4\omega}}{\omega \left(1 + \tan\left(\frac{\omega x}{2}\right)^2 \right)}$$

And Eq. (3) becomes

$$u_2 = \int \frac{e^{\sqrt{-\omega^2} x} A \cos(\omega x)}{1} dx$$

Which simplifies to

$$u_2 = \int e^{\sqrt{-\omega^2} x} A \cos(\omega x) dx$$

Hence

$$u_2 = \frac{\frac{A e^{\sqrt{-\omega^2} x} \tan\left(\frac{\omega x}{2}\right)}{\omega} + \frac{A x e^{\sqrt{-\omega^2} x}}{2} - \frac{A x e^{\sqrt{-\omega^2} x} \tan\left(\frac{\omega x}{2}\right)^2}{2} - \frac{A \sqrt{-\omega^2} x e^{\sqrt{-\omega^2} x} \tan\left(\frac{\omega x}{2}\right)}{\omega}}{1 + \tan\left(\frac{\omega x}{2}\right)^2}$$

Which simplifies to

$$u_1 = \frac{A e^{-\sqrt{-\omega^2} x} (-\sqrt{-\omega^2} x \cos(\omega x) + x \omega \sin(\omega x) + \cos(\omega x))}{4\omega^2}$$

$$u_2 = \frac{A e^{\sqrt{-\omega^2} x} (\omega x \cos(\omega x) - \sqrt{-\omega^2} x \sin(\omega x) + \sin(\omega x))}{2\omega}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{A e^{-\sqrt{-\omega^2} x} (-\sqrt{-\omega^2} x \cos(\omega x) + x \omega \sin(\omega x) + \cos(\omega x)) e^{\sqrt{-\omega^2} x}}{4\omega^2} + \frac{A e^{\sqrt{-\omega^2} x} (\omega x \cos(\omega x) - \sqrt{-\omega^2} x \sin(\omega x) + \sin(\omega x)) e^{-\sqrt{-\omega^2} x} \sqrt{-\omega^2}}{4\omega^3}$$

Which simplifies to

$$y_p(x) = \frac{A(2 \sin(\omega x) \omega^2 x + \omega \cos(\omega x) + \sqrt{-\omega^2} \sin(\omega x))}{4\omega^3}$$

Therefore the general solution is

$$y = y_h + y_p$$

$$= \left(c_1 e^{\sqrt{-\omega^2} x} + \frac{c_2 e^{-\sqrt{-\omega^2} x} \sqrt{-\omega^2}}{2\omega^2} \right) + \left(\frac{A(2 \sin(\omega x) \omega^2 x + \omega \cos(\omega x) + \sqrt{-\omega^2} \sin(\omega x))}{4\omega^3} \right)$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 e^{\sqrt{-\omega^2} x} + \frac{c_2 e^{-\sqrt{-\omega^2} x} \sqrt{-\omega^2}}{2\omega^2} + \frac{A(2 \sin(\omega x) \omega^2 x + \omega \cos(\omega x) + \sqrt{-\omega^2} \sin(\omega x))}{4\omega^3} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = \frac{4c_1\omega^2 + 2\sqrt{-\omega^2} c_2 + A}{4\omega^2} \quad (1A)$$

Taking derivative of the solution gives

$$y' = c_1 \sqrt{-\omega^2} e^{\sqrt{-\omega^2} x} + \frac{c_2 e^{-\sqrt{-\omega^2} x}}{2} + \frac{A(2\omega^3 \cos(\omega x) x + \sin(\omega x) \omega^2 + \sqrt{-\omega^2} \omega \cos(\omega x))}{4\omega^3}$$

substituting $y' = 1$ and $x = 0$ in the above gives

$$1 = \frac{(4c_1\omega^2 + A) \sqrt{-\omega^2} + 2c_2\omega^2}{4\omega^2} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = -\frac{A + 2\sqrt{-\omega^2}}{4\omega^2}$$

$$c_2 = 1$$

Substituting these values back in above solution results in

$$y = \frac{2A \sin(\omega x) \omega^2 x + A \cos(\omega x) \omega + A \sin(\omega x) \sqrt{-\omega^2} - e^{\sqrt{-\omega^2} x} A \omega - 2 e^{\sqrt{-\omega^2} x} \sqrt{-\omega^2} \omega + 2 e^{-\sqrt{-\omega^2} x} \sqrt{-\omega^2}}{4\omega^3}$$

Which simplifies to

$$y = \frac{2 e^{-\sqrt{-\omega^2} x} \sqrt{-\omega^2} \omega - \omega(A + 2\sqrt{-\omega^2}) e^{\sqrt{-\omega^2} x} + A \sin(\omega x) \sqrt{-\omega^2} + 2\omega \left(x\omega \sin(\omega x) + \frac{\cos(\omega x)}{2} \right) A}{4\omega^3}$$

Summary

The solution(s) found are the following

$$y = \frac{2 e^{-\sqrt{-\omega^2} x} \sqrt{-\omega^2} \omega - \omega(A + 2\sqrt{-\omega^2}) e^{\sqrt{-\omega^2} x} + A \sin(\omega x) \sqrt{-\omega^2} + 2\omega \left(x\omega \sin(\omega x) + \frac{\cos(\omega x)}{2} \right) A}{4\omega^3} \quad (1)$$

Verification of solutions

y

$$= \frac{2e^{-\sqrt{-\omega^2}x} \sqrt{-\omega^2} \omega - \omega(A + 2\sqrt{-\omega^2}) e^{\sqrt{-\omega^2}x} + A \sin(\omega x) \sqrt{-\omega^2} + 2\omega \left(x\omega \sin(\omega x) + \frac{\cos(\omega x)}{2} \right) A}{4\omega^3}$$

Verified OK.

6.11.4 Maple step by step solution

Let's solve

$$\left[y'' + \omega^2 y = A \cos(\omega x), y(0) = 0, y' \Big|_{\{x=0\}} = 1 \right]$$

- Highest derivative means the order of the ODE is 2

y''

- Characteristic polynomial of homogeneous ODE

$$\omega^2 + r^2 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4\omega^2})}{2}$$

- Roots of the characteristic polynomial

$$r = (\sqrt{-\omega^2}, -\sqrt{-\omega^2})$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{\sqrt{-\omega^2}x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{-\sqrt{-\omega^2}x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{\sqrt{-\omega^2}x} + c_2 e^{-\sqrt{-\omega^2}x} + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = A \cos(\omega x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{\sqrt{-\omega^2} x} & e^{-\sqrt{-\omega^2} x} \\ \sqrt{-\omega^2} e^{\sqrt{-\omega^2} x} & -\sqrt{-\omega^2} e^{-\sqrt{-\omega^2} x} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = -2\sqrt{-\omega^2}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = \frac{A(e^{\sqrt{-\omega^2} x} (\int e^{-\sqrt{-\omega^2} x} \cos(\omega x) dx) - e^{-\sqrt{-\omega^2} x} (\int e^{\sqrt{-\omega^2} x} \cos(\omega x) dx))}{2\sqrt{-\omega^2}}$$

- Compute integrals

$$y_p(x) = \frac{(\sin(\omega x)(2\sqrt{-\omega^2} x - 1)\omega + \cos(\omega x)\sqrt{-\omega^2})A}{4\sqrt{-\omega^2}\omega^2}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{\sqrt{-\omega^2} x} + c_2 e^{-\sqrt{-\omega^2} x} + \frac{(\sin(\omega x)(2\sqrt{-\omega^2} x - 1)\omega + \cos(\omega x)\sqrt{-\omega^2})A}{4\sqrt{-\omega^2}\omega^2}$$

- Check validity of solution $y = c_1 e^{\sqrt{-\omega^2} x} + c_2 e^{-\sqrt{-\omega^2} x} + \frac{(\sin(\omega x)(2\sqrt{-\omega^2} x - 1)\omega + \cos(\omega x)\sqrt{-\omega^2})A}{4\sqrt{-\omega^2}\omega^2}$

- Use initial condition $y(0) = 0$

$$0 = c_1 + c_2 + \frac{A}{4\omega^2}$$

- Compute derivative of the solution

$$y' = c_1 \sqrt{-\omega^2} e^{\sqrt{-\omega^2} x} - c_2 \sqrt{-\omega^2} e^{-\sqrt{-\omega^2} x} + \frac{(\omega^2 \cos(\omega x)(2\sqrt{-\omega^2} x - 1) + \sin(\omega x)\sqrt{-\omega^2} \omega)A}{4\sqrt{-\omega^2}\omega^2}$$

- Use the initial condition $y'|_{\{x=0\}} = 1$

$$1 = \sqrt{-\omega^2} c_1 - \sqrt{-\omega^2} c_2 - \frac{A}{4\sqrt{-\omega^2}}$$

- Solve for c_1 and c_2

$$\left\{ c_1 = -\frac{A+2\sqrt{-\omega^2}}{4\omega^2}, c_2 = \frac{\sqrt{-\omega^2}}{2\omega^2} \right\}$$

- Substitute constant values into general solution and simplify

$$y = \frac{2A \sin(\omega x)\sqrt{-\omega^2} \omega x + A \cos(\omega x)\sqrt{-\omega^2} - A\omega \sin(\omega x) - e^{\sqrt{-\omega^2} x} \sqrt{-\omega^2} A + 2e^{\sqrt{-\omega^2} x} \omega^2 - 2\omega^2 e^{-\sqrt{-\omega^2} x}}{4\sqrt{-\omega^2}\omega^2}$$

- Solution to the IVP

$$y = \frac{2A \sin(\omega x)\sqrt{-\omega^2} \omega x + A \cos(\omega x)\sqrt{-\omega^2} - A\omega \sin(\omega x) - e^{\sqrt{-\omega^2} x} \sqrt{-\omega^2} A + 2e^{\sqrt{-\omega^2} x} \omega^2 - 2\omega^2 e^{-\sqrt{-\omega^2} x}}{4\sqrt{-\omega^2}\omega^2}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 18

```
dsolve([diff(y(x),x$2)+omega^2*y(x)=A*cos(omega*x),y(0) = 0, D(y)(0) = 1],y(x), singsol=all)
```

$$y(x) = \frac{\sin(\omega x) \left(1 + \frac{Ax}{2}\right)}{\omega}$$

✓ Solution by Mathematica

Time used: 0.058 (sec). Leaf size: 21

```
DSolve[{y''[x]+\[Omega]^2*y[x]==A*Cos\[Omega]*x},{y[0]==0,y'[0]==1},y[x],x,IncludeSingular
```

$$y(x) \rightarrow \frac{(Ax + 2) \sin(x\omega)}{2\omega}$$

7 Chapter 2. Linear equations with constant coefficients. Page 74

7.1	problem 4(a)	636
7.2	problem 4(b)	641
7.3	problem 4(c)	648
7.4	problem 4(d)	653
7.5	problem 4(f)	655
7.6	problem 4(g)	661
7.7	problem 4(h)	667
7.8	problem 4(i)	672

7.1 problem 4(a)

7.1.1 Maple step by step solution 637

Internal problem ID [5972]

Internal file name [OUTPUT/5220_Sunday_June_05_2022_03_27_49_PM_89684670/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 74

Problem number: 4(a).

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_3rd_order , _missing_x]]
```

$$y''' - 8y = 0$$

The characteristic equation is

$$\lambda^3 - 8 = 0$$

The roots of the above equation are

$$\lambda_1 = 2$$

$$\lambda_2 = i\sqrt{3} - 1$$

$$\lambda_3 = -i\sqrt{3} - 1$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 e^{2x} + e^{(i\sqrt{3}-1)x} c_2 + e^{(-i\sqrt{3}-1)x} c_3$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = e^{2x}$$

$$y_2 = e^{(i\sqrt{3}-1)x}$$

$$y_3 = e^{(-i\sqrt{3}-1)x}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{2x} + e^{(i\sqrt{3}-1)x} c_2 + e^{(-i\sqrt{3}-1)x} c_3 \quad (1)$$

Verification of solutions

$$y = c_1 e^{2x} + e^{(i\sqrt{3}-1)x} c_2 + e^{(-i\sqrt{3}-1)x} c_3$$

Verified OK.

7.1.1 Maple step by step solution

Let's solve

$$y''' - 8y = 0$$

- Highest derivative means the order of the ODE is 3

$$y'''$$

- Convert linear ODE into a system of first order ODEs

- Define new variable $y_1(x)$

$$y_1(x) = y$$

- Define new variable $y_2(x)$

$$y_2(x) = y'$$

- Define new variable $y_3(x)$

$$y_3(x) = y''$$

- Isolate for $y_3'(x)$ using original ODE

$$y_3'(x) = 8y_1(x)$$

Convert linear ODE into a system of first order ODEs

$$[y_2(x) = y_1'(x), y_3(x) = y_2'(x), y_3'(x) = 8y_1(x)]$$

- Define vector

$$\vec{y}(x) = \begin{bmatrix} y_1(x) \\ y_2(x) \\ y_3(x) \end{bmatrix}$$

- System to solve

$$\vec{y}'(x) = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 8 & 0 & 0 \end{bmatrix} \cdot \vec{y}(x)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 8 & 0 & 0 \end{bmatrix}$$

- Rewrite the system as

$$\vec{y}'(x) = A \cdot \vec{y}(x)$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[2, \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right], \left[-i\sqrt{3} - 1, \begin{bmatrix} \frac{1}{(-i\sqrt{3}-1)^2} \\ \frac{1}{-i\sqrt{3}-1} \\ 1 \end{bmatrix} \right], \left[i\sqrt{3} - 1, \begin{bmatrix} \frac{1}{(i\sqrt{3}-1)^2} \\ \frac{1}{i\sqrt{3}-1} \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[2, \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_1 = e^{2x} \cdot \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix}$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$\left[-i\sqrt{3} - 1, \begin{bmatrix} \frac{1}{(-i\sqrt{3}-1)^2} \\ \frac{1}{-i\sqrt{3}-1} \\ 1 \end{bmatrix} \right]$$

- Solution from eigenpair

$$e^{(-I\sqrt{3}-1)x} \cdot \begin{bmatrix} \frac{1}{(-I\sqrt{3}-1)^2} \\ \frac{1}{-I\sqrt{3}-1} \\ 1 \end{bmatrix}$$

- Use Euler identity to write solution in terms of sin and cos

$$e^{-x} \cdot (\cos(\sqrt{3}x) - I \sin(\sqrt{3}x)) \cdot \begin{bmatrix} \frac{1}{(-I\sqrt{3}-1)^2} \\ \frac{1}{-I\sqrt{3}-1} \\ 1 \end{bmatrix}$$

- Simplify expression

$$e^{-x} \cdot \begin{bmatrix} \frac{\cos(\sqrt{3}x) - I \sin(\sqrt{3}x)}{(-I\sqrt{3}-1)^2} \\ \frac{\cos(\sqrt{3}x) - I \sin(\sqrt{3}x)}{-I\sqrt{3}-1} \\ \cos(\sqrt{3}x) - I \sin(\sqrt{3}x) \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\vec{y}_2(x) = e^{-x} \cdot \begin{bmatrix} -\frac{\cos(\sqrt{3}x)}{8} - \frac{\sqrt{3} \sin(\sqrt{3}x)}{8} \\ -\frac{\cos(\sqrt{3}x)}{4} + \frac{\sqrt{3} \sin(\sqrt{3}x)}{4} \\ \cos(\sqrt{3}x) \end{bmatrix}, \vec{y}_3(x) = e^{-x} \cdot \begin{bmatrix} -\frac{\sqrt{3} \cos(\sqrt{3}x)}{8} + \frac{\sin(\sqrt{3}x)}{8} \\ \frac{\sqrt{3} \cos(\sqrt{3}x)}{4} + \frac{\sin(\sqrt{3}x)}{4} \\ -\sin(\sqrt{3}x) \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{y} = c_1 \vec{y}_1 + c_2 \vec{y}_2(x) + c_3 \vec{y}_3(x)$$

- Substitute solutions into the general solution

$$\vec{y} = c_1 e^{2x} \cdot \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} + c_2 e^{-x} \cdot \begin{bmatrix} -\frac{\cos(\sqrt{3}x)}{8} - \frac{\sqrt{3} \sin(\sqrt{3}x)}{8} \\ -\frac{\cos(\sqrt{3}x)}{4} + \frac{\sqrt{3} \sin(\sqrt{3}x)}{4} \\ \cos(\sqrt{3}x) \end{bmatrix} + c_3 e^{-x} \cdot \begin{bmatrix} -\frac{\sqrt{3} \cos(\sqrt{3}x)}{8} + \frac{\sin(\sqrt{3}x)}{8} \\ \frac{\sqrt{3} \cos(\sqrt{3}x)}{4} + \frac{\sin(\sqrt{3}x)}{4} \\ -\sin(\sqrt{3}x) \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y = -\frac{e^{-x}(c_3\sqrt{3}+c_2)\cos(\sqrt{3}x)}{8} - \frac{e^{-x}(\sqrt{3}c_2-c_3)\sin(\sqrt{3}x)}{8} + \frac{c_1 e^{2x}}{4}$$

Maple trace

```
`Methods for third order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 35

```
dsolve(diff(y(x),x$3)-8*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{2x}c_1 + c_2e^{-x} \sin(\sqrt{3}x) + c_3e^{-x} \cos(\sqrt{3}x)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 42

```
DSolve[y'''[x]-8*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x} \left(c_1 e^{3x} + c_2 \cos(\sqrt{3}x) + c_3 \sin(\sqrt{3}x) \right)$$

7.2 problem 4(b)

7.2.1 Maple step by step solution 642

Internal problem ID [5973]

Internal file name [OUTPUT/5221_Sunday_June_05_2022_03_27_50_PM_1305271/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 74

Problem number: 4(b).

ODE order: 4.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

[[_high_order , _missing_x]]

$$y'''' + 16y = 0$$

The characteristic equation is

$$\lambda^4 + 16 = 0$$

The roots of the above equation are

$$\begin{aligned}\lambda_1 &= \sqrt{2} + i\sqrt{2} \\ \lambda_2 &= -\sqrt{2} + i\sqrt{2} \\ \lambda_3 &= -\sqrt{2} - i\sqrt{2} \\ \lambda_4 &= -i\sqrt{2} + \sqrt{2}\end{aligned}$$

Therefore the homogeneous solution is

$$y_h(x) = e^{(-i\sqrt{2}+\sqrt{2})x} c_1 + e^{(-\sqrt{2}+i\sqrt{2})x} c_2 + e^{(-\sqrt{2}-i\sqrt{2})x} c_3 + e^{(\sqrt{2}+i\sqrt{2})x} c_4$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = e^{(-i\sqrt{2}+\sqrt{2})x}$$

$$y_2 = e^{(-\sqrt{2}+i\sqrt{2})x}$$

$$y_3 = e^{(-\sqrt{2}-i\sqrt{2})x}$$

$$y_4 = e^{(\sqrt{2}+i\sqrt{2})x}$$

Summary

The solution(s) found are the following

$$y = e^{(-i\sqrt{2}+\sqrt{2})x} c_1 + e^{(-\sqrt{2}+i\sqrt{2})x} c_2 + e^{(-\sqrt{2}-i\sqrt{2})x} c_3 + e^{(\sqrt{2}+i\sqrt{2})x} c_4 \quad (1)$$

Verification of solutions

$$y = e^{(-i\sqrt{2}+\sqrt{2})x} c_1 + e^{(-\sqrt{2}+i\sqrt{2})x} c_2 + e^{(-\sqrt{2}-i\sqrt{2})x} c_3 + e^{(\sqrt{2}+i\sqrt{2})x} c_4$$

Verified OK.

7.2.1 Maple step by step solution

Let's solve

$$y'''' + 16y = 0$$

- Highest derivative means the order of the ODE is 4

$$y''''$$

- Convert linear ODE into a system of first order ODEs

- Define new variable $y_1(x)$

$$y_1(x) = y$$

- Define new variable $y_2(x)$

$$y_2(x) = y'$$

- Define new variable $y_3(x)$

$$y_3(x) = y''$$

- Define new variable $y_4(x)$

$$y_4(x) = y'''$$

- Isolate for $y_4'(x)$ using original ODE

$$y_4'(x) = -16y_1(x)$$

Convert linear ODE into a system of first order ODEs

$$[y_2(x) = y_1'(x), y_3(x) = y_2'(x), y_4(x) = y_3'(x), y_4'(x) = -16y_1(x)]$$

- Define vector

$$\vec{y}(x) = \begin{bmatrix} y_1(x) \\ y_2(x) \\ y_3(x) \\ y_4(x) \end{bmatrix}$$

- System to solve

$$\vec{y}'(x) = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -16 & 0 & 0 & 0 \end{bmatrix} \cdot \vec{y}(x)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -16 & 0 & 0 & 0 \end{bmatrix}$$

- Rewrite the system as

$$\vec{y}'(x) = A \cdot \vec{y}(x)$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[-\sqrt{2} - I\sqrt{2}, \begin{bmatrix} \frac{1}{(-\sqrt{2}-I\sqrt{2})^3} \\ \frac{1}{(-\sqrt{2}-I\sqrt{2})^2} \\ \frac{1}{-\sqrt{2}-I\sqrt{2}} \\ 1 \end{bmatrix} \right], \left[-\sqrt{2} + I\sqrt{2}, \begin{bmatrix} \frac{1}{(-\sqrt{2}+I\sqrt{2})^3} \\ \frac{1}{(-\sqrt{2}+I\sqrt{2})^2} \\ \frac{1}{-\sqrt{2}+I\sqrt{2}} \\ 1 \end{bmatrix} \right], \left[\sqrt{2} + I\sqrt{2}, \begin{bmatrix} \frac{1}{(\sqrt{2}+I\sqrt{2})^3} \\ \frac{1}{(\sqrt{2}+I\sqrt{2})^2} \\ \frac{1}{\sqrt{2}+I\sqrt{2}} \\ 1 \end{bmatrix} \right] \right]$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$\begin{bmatrix} -\sqrt{2} - I\sqrt{2}, & \begin{bmatrix} \frac{1}{(-\sqrt{2}-I\sqrt{2})^3} \\ \frac{1}{(-\sqrt{2}-I\sqrt{2})^2} \\ \frac{1}{-\sqrt{2}-I\sqrt{2}} \\ 1 \end{bmatrix} \end{bmatrix}$$

- Solution from eigenpair

$$e^{(-\sqrt{2}-I\sqrt{2})x} \cdot \begin{bmatrix} \frac{1}{(-\sqrt{2}-I\sqrt{2})^3} \\ \frac{1}{(-\sqrt{2}-I\sqrt{2})^2} \\ \frac{1}{-\sqrt{2}-I\sqrt{2}} \\ 1 \end{bmatrix}$$

- Use Euler identity to write solution in terms of sin and cos

$$e^{-x\sqrt{2}} \cdot (\cos(x\sqrt{2}) - I \sin(x\sqrt{2})) \cdot \begin{bmatrix} \frac{1}{(-\sqrt{2}-I\sqrt{2})^3} \\ \frac{1}{(-\sqrt{2}-I\sqrt{2})^2} \\ \frac{1}{-\sqrt{2}-I\sqrt{2}} \\ 1 \end{bmatrix}$$

- Simplify expression

$$e^{-x\sqrt{2}} \cdot \begin{bmatrix} \frac{\cos(x\sqrt{2}) - I \sin(x\sqrt{2})}{(-\sqrt{2}-I\sqrt{2})^3} \\ \frac{\cos(x\sqrt{2}) - I \sin(x\sqrt{2})}{(-\sqrt{2}-I\sqrt{2})^2} \\ \frac{\cos(x\sqrt{2}) - I \sin(x\sqrt{2})}{-\sqrt{2}-I\sqrt{2}} \\ \cos(x\sqrt{2}) - I \sin(x\sqrt{2}) \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\vec{y}_1(x) = e^{-x\sqrt{2}} \cdot \begin{bmatrix} \frac{\cos(x\sqrt{2})\sqrt{2}}{16} + \frac{\sin(x\sqrt{2})\sqrt{2}}{16} \\ -\frac{\sin(x\sqrt{2})}{4} \\ -\frac{\cos(x\sqrt{2})\sqrt{2}}{4} + \frac{\sin(x\sqrt{2})\sqrt{2}}{4} \\ \cos(x\sqrt{2}) \end{bmatrix}, \vec{y}_2(x) = e^{-x\sqrt{2}} \cdot \begin{bmatrix} \frac{\cos(x\sqrt{2})\sqrt{2}}{16} - \frac{\sin(x\sqrt{2})\sqrt{2}}{16} \\ -\frac{\cos(x\sqrt{2})}{4} \\ \frac{\cos(x\sqrt{2})\sqrt{2}}{4} + \frac{\sin(x\sqrt{2})\sqrt{2}}{4} \\ -\sin(x\sqrt{2}) \end{bmatrix}$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$\sqrt{2} + I\sqrt{2}, \begin{bmatrix} \frac{1}{(\sqrt{2}+I\sqrt{2})^3} \\ \frac{1}{(\sqrt{2}+I\sqrt{2})^2} \\ \frac{1}{\sqrt{2}+I\sqrt{2}} \\ 1 \end{bmatrix}$$

- Solution from eigenpair

$$e^{(\sqrt{2}+I\sqrt{2})x} \cdot \begin{bmatrix} \frac{1}{(\sqrt{2}+I\sqrt{2})^3} \\ \frac{1}{(\sqrt{2}+I\sqrt{2})^2} \\ \frac{1}{\sqrt{2}+I\sqrt{2}} \\ 1 \end{bmatrix}$$

- Use Euler identity to write solution in terms of sin and cos

$$e^{x\sqrt{2}} \cdot (\cos(x\sqrt{2}) + I \sin(x\sqrt{2})) \cdot \begin{bmatrix} \frac{1}{(\sqrt{2}+I\sqrt{2})^3} \\ \frac{1}{(\sqrt{2}+I\sqrt{2})^2} \\ \frac{1}{\sqrt{2}+I\sqrt{2}} \\ 1 \end{bmatrix}$$

- Simplify expression

$$e^{x\sqrt{2}} \cdot \begin{bmatrix} \frac{\cos(x\sqrt{2}) + I \sin(x\sqrt{2})}{(\sqrt{2} + I\sqrt{2})^3} \\ \frac{\cos(x\sqrt{2}) + I \sin(x\sqrt{2})}{(\sqrt{2} + I\sqrt{2})^2} \\ \frac{\cos(x\sqrt{2}) + I \sin(x\sqrt{2})}{\sqrt{2} + I\sqrt{2}} \\ \cos(x\sqrt{2}) + I \sin(x\sqrt{2}) \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\vec{y}_3(x) = e^{x\sqrt{2}} \cdot \begin{bmatrix} -\frac{\cos(x\sqrt{2})\sqrt{2}}{16} + \frac{\sin(x\sqrt{2})\sqrt{2}}{16} \\ \frac{\sin(x\sqrt{2})}{4} \\ \frac{\cos(x\sqrt{2})\sqrt{2}}{4} + \frac{\sin(x\sqrt{2})\sqrt{2}}{4} \\ \cos(x\sqrt{2}) \end{bmatrix}, \vec{y}_4(x) = e^{x\sqrt{2}} \cdot \begin{bmatrix} -\frac{\cos(x\sqrt{2})\sqrt{2}}{16} - \frac{\sin(x\sqrt{2})\sqrt{2}}{16} \\ -\frac{\cos(x\sqrt{2})}{4} \\ -\frac{\cos(x\sqrt{2})\sqrt{2}}{4} + \frac{\sin(x\sqrt{2})\sqrt{2}}{4} \\ \sin(x\sqrt{2}) \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{y} = c_1 \vec{y}_1(x) + c_2 \vec{y}_2(x) + c_3 \vec{y}_3(x) + c_4 \vec{y}_4(x)$$

- Substitute solutions into the general solution

$$\vec{y} = c_1 e^{-x\sqrt{2}} \cdot \begin{bmatrix} \frac{\cos(x\sqrt{2})\sqrt{2}}{16} + \frac{\sin(x\sqrt{2})\sqrt{2}}{16} \\ -\frac{\sin(x\sqrt{2})}{4} \\ -\frac{\cos(x\sqrt{2})\sqrt{2}}{4} + \frac{\sin(x\sqrt{2})\sqrt{2}}{4} \\ \cos(x\sqrt{2}) \end{bmatrix} + c_2 e^{-x\sqrt{2}} \cdot \begin{bmatrix} \frac{\cos(x\sqrt{2})\sqrt{2}}{16} - \frac{\sin(x\sqrt{2})\sqrt{2}}{16} \\ -\frac{\cos(x\sqrt{2})}{4} \\ \frac{\cos(x\sqrt{2})\sqrt{2}}{4} + \frac{\sin(x\sqrt{2})\sqrt{2}}{4} \\ -\sin(x\sqrt{2}) \end{bmatrix} + e^{x\sqrt{2}} c_3 \cdot$$

- First component of the vector is the solution to the ODE

$$y = \frac{\sqrt{2} \left((c_1 + c_2) \cos(x\sqrt{2}) + \sin(x\sqrt{2}) (c_1 - c_2) \right) e^{-x\sqrt{2}} - \left((c_3 + c_4) \cos(x\sqrt{2}) - \sin(x\sqrt{2}) (c_3 - c_4) \right) e^{x\sqrt{2}}}{16}$$

Maple trace

```
`Methods for high order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 65

```
dsolve(diff(y(x),x$4)+16*y(x)=0,y(x), singsol=all)
```

$$y(x) = -c_1 e^{-\sqrt{2}x} \sin(\sqrt{2}x) - c_2 e^{\sqrt{2}x} \sin(\sqrt{2}x) \\ + c_3 e^{-\sqrt{2}x} \cos(\sqrt{2}x) + c_4 e^{\sqrt{2}x} \cos(\sqrt{2}x)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 67

```
DSolve[y''''[x]+16*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-\sqrt{2}x} \left((c_1 e^{2\sqrt{2}x} + c_2) \cos(\sqrt{2}x) + (c_4 e^{2\sqrt{2}x} + c_3) \sin(\sqrt{2}x) \right)$$

7.3 problem 4(c)

7.3.1 Maple step by step solution 649

Internal problem ID [5974]

Internal file name [OUTPUT/5222_Sunday_June_05_2022_03_27_51_PM_4576512/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 74

Problem number: 4(c).

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_3rd_order , _missing_x]]
```

$$y''' - 5y'' + 6y' = 0$$

The characteristic equation is

$$\lambda^3 - 5\lambda^2 + 6\lambda = 0$$

The roots of the above equation are

$$\lambda_1 = 0$$

$$\lambda_2 = 3$$

$$\lambda_3 = 2$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 + c_2e^{2x} + c_3e^{3x}$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = 1$$

$$y_2 = e^{2x}$$

$$y_3 = e^{3x}$$

Summary

The solution(s) found are the following

$$y = c_1 + c_2e^{2x} + c_3e^{3x} \quad (1)$$

Verification of solutions

$$y = c_1 + c_2e^{2x} + c_3e^{3x}$$

Verified OK.

7.3.1 Maple step by step solution

Let's solve

$$y''' - 5y'' + 6y' = 0$$

- Highest derivative means the order of the ODE is 3
 y'''

□ Convert linear ODE into a system of first order ODEs

- Define new variable $y_1(x)$

$$y_1(x) = y$$

- Define new variable $y_2(x)$

$$y_2(x) = y'$$

- Define new variable $y_3(x)$

$$y_3(x) = y''$$

- Isolate for $y_3'(x)$ using original ODE

$$y_3'(x) = 5y_3(x) - 6y_2(x)$$

Convert linear ODE into a system of first order ODEs

$$[y_2(x) = y_1'(x), y_3(x) = y_2'(x), y_3'(x) = 5y_3(x) - 6y_2(x)]$$

- Define vector

$$\vec{y}(x) = \begin{bmatrix} y_1(x) \\ y_2(x) \\ y_3(x) \end{bmatrix}$$

- System to solve

$$\vec{y}'(x) = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & -6 & 5 \end{bmatrix} \cdot \vec{y}(x)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & -6 & 5 \end{bmatrix}$$

- Rewrite the system as

$$\vec{y}'(x) = A \cdot \vec{y}(x)$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[0, \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \right], \left[2, \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right], \left[3, \begin{bmatrix} \frac{1}{9} \\ \frac{1}{3} \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[0, \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

- Consider eigenpair

$$\left[2, \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_2 = e^{2x} \cdot \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[3, \begin{bmatrix} \frac{1}{9} \\ \frac{1}{3} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_3 = e^{3x} \cdot \begin{bmatrix} \frac{1}{9} \\ \frac{1}{3} \\ 1 \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{y} = c_1 \vec{y}_1 + c_2 \vec{y}_2 + c_3 \vec{y}_3$$

- Substitute solutions into the general solution

$$\vec{y} = c_2 e^{2x} \cdot \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} + c_3 e^{3x} \cdot \begin{bmatrix} \frac{1}{9} \\ \frac{1}{3} \\ 1 \end{bmatrix} + \begin{bmatrix} c_1 \\ 0 \\ 0 \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y = \frac{c_2 e^{2x}}{4} + \frac{c_3 e^{3x}}{9} + c_1$$

Maple trace

```

`Methods for third order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(diff(y(x),x$3)-5*diff(y(x),x$2)+6*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = c_1 + c_2 e^{2x} + c_3 e^{3x}$$

✓ Solution by Mathematica

Time used: 0.033 (sec). Leaf size: 30

```
DSolve[y'''[x]-5*y''[x]+6*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}c_1 e^{2x} + \frac{1}{3}c_2 e^{3x} + c_3$$

7.4 problem 4(d)

Internal problem ID [5975]

Internal file name [OUTPUT/5223_Sunday_June_05_2022_03_27_52_PM_23069737/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 74

Problem number: 4(d).

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_3rd_order , _missing_x]]
```

$$y''' - iy'' + 4y' - 4iy = 0$$

The characteristic equation is

$$\lambda^3 - i\lambda^2 + 4\lambda - 4i = 0$$

The roots of the above equation are

$$\lambda_1 = -2i$$

$$\lambda_2 = 2i$$

$$\lambda_3 = i$$

Therefore the homogeneous solution is

$$y_h(x) = e^{-2ix}c_1 + e^{2ix}c_2 + e^{ix}c_3$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = e^{-2ix}$$

$$y_2 = e^{2ix}$$

$$y_3 = e^{ix}$$

Summary

The solution(s) found are the following

$$y = e^{-2ix} c_1 + e^{2ix} c_2 + e^{ix} c_3 \quad (1)$$

Verification of solutions

$$y = e^{-2ix} c_1 + e^{2ix} c_2 + e^{ix} c_3$$

Verified OK.

Maple trace

```
`Methods for third order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 26

```
dsolve(diff(y(x),x$3)-I*diff(y(x),x$2)+4*diff(y(x),x)-4*I*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{ix} + c_2 e^{2ix} + c_3 e^{-2ix}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 36

```
DSolve[y'''[x]-I*y''[x]+4*y'[x]-4*I*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2ix} (c_2 e^{4ix} + c_3 e^{3ix} + c_1)$$

7.5 problem 4(f)

7.5.1 Maple step by step solution 656

Internal problem ID [5976]

Internal file name [OUTPUT/5224_Sunday_June_05_2022_03_27_53_PM_58330443/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 74

Problem number: 4(f).

ODE order: 4.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_high_order , _missing_x]]
```

$$y'''' + 5y'' + 4y = 0$$

The characteristic equation is

$$\lambda^4 + 5\lambda^2 + 4 = 0$$

The roots of the above equation are

$$\lambda_1 = 2i$$

$$\lambda_2 = -2i$$

$$\lambda_3 = i$$

$$\lambda_4 = -i$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 e^{-ix} + e^{-2ix} c_2 + e^{2ix} c_3 + e^{ix} c_4$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = e^{-ix}$$

$$y_2 = e^{-2ix}$$

$$y_3 = e^{2ix}$$

$$y_4 = e^{ix}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-ix} + e^{-2ix} c_2 + e^{2ix} c_3 + e^{ix} c_4 \quad (1)$$

Verification of solutions

$$y = c_1 e^{-ix} + e^{-2ix} c_2 + e^{2ix} c_3 + e^{ix} c_4$$

Verified OK.

7.5.1 Maple step by step solution

Let's solve

$$y'''' + 5y'' + 4y = 0$$

- Highest derivative means the order of the ODE is 4

$$y''''$$

- Convert linear ODE into a system of first order ODEs

- Define new variable $y_1(x)$

$$y_1(x) = y$$

- Define new variable $y_2(x)$

$$y_2(x) = y'$$

- Define new variable $y_3(x)$

$$y_3(x) = y''$$

- Define new variable $y_4(x)$

$$y_4(x) = y'''$$

- Isolate for $y_4'(x)$ using original ODE

$$y_4'(x) = -5y_3(x) - 4y_1(x)$$

Convert linear ODE into a system of first order ODEs

$$[y_2(x) = y_1'(x), y_3(x) = y_2'(x), y_4(x) = y_3'(x), y_4'(x) = -5y_3(x) - 4y_1(x)]$$

- Define vector

$$\vec{y}(x) = \begin{bmatrix} y_1(x) \\ y_2(x) \\ y_3(x) \\ y_4(x) \end{bmatrix}$$

- System to solve

$$\vec{y}'(x) = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -4 & 0 & -5 & 0 \end{bmatrix} \cdot \vec{y}(x)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -4 & 0 & -5 & 0 \end{bmatrix}$$

- Rewrite the system as

$$\vec{y}'(x) = A \cdot \vec{y}(x)$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[-2I, \begin{bmatrix} -\frac{1}{8} \\ -\frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right], \left[-I, \begin{bmatrix} -1 \\ -1 \\ 1 \\ 1 \end{bmatrix} \right], \left[I, \begin{bmatrix} 1 \\ -1 \\ -1 \\ 1 \end{bmatrix} \right], \left[2I, \begin{bmatrix} \frac{1}{8} \\ -\frac{1}{4} \\ -\frac{1}{2} \\ 1 \end{bmatrix} \right] \right]$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$\left[-2I, \begin{bmatrix} -\frac{1}{8} \\ -\frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right]$$

- Solution from eigenpair

$$e^{-2Ix} \cdot \begin{bmatrix} -\frac{I}{8} \\ -\frac{1}{4} \\ \frac{I}{2} \\ 1 \end{bmatrix}$$

- Use Euler identity to write solution in terms of sin and cos

$$(\cos(2x) - I \sin(2x)) \cdot \begin{bmatrix} -\frac{I}{8} \\ -\frac{1}{4} \\ \frac{I}{2} \\ 1 \end{bmatrix}$$

- Simplify expression

$$\begin{bmatrix} -\frac{I}{8}(\cos(2x) - I \sin(2x)) \\ -\frac{\cos(2x)}{4} + \frac{I \sin(2x)}{4} \\ \frac{I}{2}(\cos(2x) - I \sin(2x)) \\ \cos(2x) - I \sin(2x) \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\vec{y}_1(x) = \begin{bmatrix} -\frac{\sin(2x)}{8} \\ -\frac{\cos(2x)}{4} \\ \frac{\sin(2x)}{2} \\ \cos(2x) \end{bmatrix}, \vec{y}_2(x) = \begin{bmatrix} -\frac{\cos(2x)}{8} \\ \frac{\sin(2x)}{4} \\ \frac{\cos(2x)}{2} \\ -\sin(2x) \end{bmatrix}$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$-I, \begin{bmatrix} -I \\ -1 \\ I \\ 1 \end{bmatrix}$$

- Solution from eigenpair

$$e^{-Ix} \cdot \begin{bmatrix} -I \\ -1 \\ I \\ 1 \end{bmatrix}$$

- Use Euler identity to write solution in terms of sin and cos

$$(\cos(x) - I \sin(x)) \cdot \begin{bmatrix} -I \\ -1 \\ I \\ 1 \end{bmatrix}$$

- Simplify expression

$$\begin{bmatrix} -I(\cos(x) - I \sin(x)) \\ -\cos(x) + I \sin(x) \\ I(\cos(x) - I \sin(x)) \\ \cos(x) - I \sin(x) \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\vec{y}_3(x) = \begin{bmatrix} -\sin(x) \\ -\cos(x) \\ \sin(x) \\ \cos(x) \end{bmatrix}, \vec{y}_4(x) = \begin{bmatrix} -\cos(x) \\ \sin(x) \\ \cos(x) \\ -\sin(x) \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{y} = c_1 \vec{y}_1(x) + c_2 \vec{y}_2(x) + c_3 \vec{y}_3(x) + c_4 \vec{y}_4(x)$$

- Substitute solutions into the general solution

$$\vec{y} = \begin{bmatrix} -c_4 \cos(x) - c_3 \sin(x) - \frac{c_2 \cos(2x)}{8} - \frac{\sin(2x)c_1}{8} \\ c_4 \sin(x) - c_3 \cos(x) + \frac{c_2 \sin(2x)}{4} - \frac{c_1 \cos(2x)}{4} \\ c_4 \cos(x) + c_3 \sin(x) + \frac{c_2 \cos(2x)}{2} + \frac{\sin(2x)c_1}{2} \\ -c_4 \sin(x) + c_3 \cos(x) - c_2 \sin(2x) + c_1 \cos(2x) \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y = -c_4 \cos(x) - c_3 \sin(x) - \frac{c_2 \cos(2x)}{8} - \frac{\sin(2x)c_1}{8}$$

Maple trace

```
`Methods for high order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
dsolve(diff(y(x),x$4)+5*diff(y(x),x$2)+4*y(x)=0,y(x), singsol=all)
```

$$y(x) = 2c_2 \cos(x)^2 + (2c_1 \sin(x) + c_4) \cos(x) + c_3 \sin(x) - c_2$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 30

```
DSolve[y''''[x]+5*y''[x]+4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos(2x) + c_4 \sin(x) + \cos(x)(2c_2 \sin(x) + c_3)$$

7.6 problem 4(g)

7.6.1 Maple step by step solution 662

Internal problem ID [5977]

Internal file name [OUTPUT/5225_Sunday_June_05_2022_03_27_55_PM_82770425/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 74

Problem number: 4(g).

ODE order: 4.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_high_order , _missing_x]]
```

$$y'''' - 16y = 0$$

The characteristic equation is

$$\lambda^4 - 16 = 0$$

The roots of the above equation are

$$\lambda_1 = 2$$

$$\lambda_2 = -2$$

$$\lambda_3 = 2i$$

$$\lambda_4 = -2i$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 e^{-2x} + c_2 e^{2x} + e^{-2ix} c_3 + e^{2ix} c_4$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = e^{-2x}$$

$$y_2 = e^{2x}$$

$$y_3 = e^{-2ix}$$

$$y_4 = e^{2ix}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-2x} + c_2 e^{2x} + e^{-2ix} c_3 + e^{2ix} c_4 \quad (1)$$

Verification of solutions

$$y = c_1 e^{-2x} + c_2 e^{2x} + e^{-2ix} c_3 + e^{2ix} c_4$$

Verified OK.

7.6.1 Maple step by step solution

Let's solve

$$y'''' - 16y = 0$$

- Highest derivative means the order of the ODE is 4

$$y''''$$

- Convert linear ODE into a system of first order ODEs

- Define new variable $y_1(x)$

$$y_1(x) = y$$

- Define new variable $y_2(x)$

$$y_2(x) = y'$$

- Define new variable $y_3(x)$

$$y_3(x) = y''$$

- Define new variable $y_4(x)$

$$y_4(x) = y'''$$

- Isolate for $y_4'(x)$ using original ODE

$$y_4'(x) = 16y_1(x)$$

Convert linear ODE into a system of first order ODEs

$$[y_2(x) = y_1'(x), y_3(x) = y_2'(x), y_4(x) = y_3'(x), y_4'(x) = 16y_1(x)]$$

- Define vector

$$\vec{y}(x) = \begin{bmatrix} y_1(x) \\ y_2(x) \\ y_3(x) \\ y_4(x) \end{bmatrix}$$

- System to solve

$$\vec{y}'(x) = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 16 & 0 & 0 & 0 \end{bmatrix} \cdot \vec{y}(x)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 16 & 0 & 0 & 0 \end{bmatrix}$$

- Rewrite the system as

$$\vec{y}'(x) = A \cdot \vec{y}(x)$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[-2, \begin{bmatrix} -\frac{1}{8} \\ \frac{1}{4} \\ -\frac{1}{2} \\ 1 \end{bmatrix} \right], \left[2, \begin{bmatrix} \frac{1}{8} \\ \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right], \left[-2\mathbf{I}, \begin{bmatrix} -\frac{1}{8} \\ -\frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right], \left[2\mathbf{I}, \begin{bmatrix} \frac{1}{8} \\ -\frac{1}{4} \\ -\frac{1}{2} \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[-2, \begin{bmatrix} -\frac{1}{8} \\ \frac{1}{4} \\ -\frac{1}{2} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_1 = e^{-2x} \cdot \begin{bmatrix} -\frac{1}{8} \\ \frac{1}{4} \\ -\frac{1}{2} \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[2, \begin{bmatrix} \frac{1}{8} \\ \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_2 = e^{2x} \cdot \begin{bmatrix} \frac{1}{8} \\ \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix}$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$\left[-2I, \begin{bmatrix} -\frac{1}{8} \\ -\frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right]$$

- Solution from eigenpair

$$e^{-2Ix} \cdot \begin{bmatrix} -\frac{1}{8} \\ -\frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix}$$

- Use Euler identity to write solution in terms of sin and cos

$$(\cos(2x) - I \sin(2x)) \cdot \begin{bmatrix} -\frac{I}{8} \\ -\frac{1}{4} \\ \frac{I}{2} \\ 1 \end{bmatrix}$$

- Simplify expression

$$\begin{bmatrix} -\frac{I}{8}(\cos(2x) - I \sin(2x)) \\ -\frac{\cos(2x)}{4} + \frac{I \sin(2x)}{4} \\ \frac{I}{2}(\cos(2x) - I \sin(2x)) \\ \cos(2x) - I \sin(2x) \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\vec{y}_3(x) = \begin{bmatrix} -\frac{\sin(2x)}{8} \\ -\frac{\cos(2x)}{4} \\ \frac{\sin(2x)}{2} \\ \cos(2x) \end{bmatrix}, \vec{y}_4(x) = \begin{bmatrix} -\frac{\cos(2x)}{8} \\ \frac{\sin(2x)}{4} \\ \frac{\cos(2x)}{2} \\ -\sin(2x) \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{y} = c_1 \vec{y}_1 + c_2 \vec{y}_2 + c_3 \vec{y}_3(x) + c_4 \vec{y}_4(x)$$

- Substitute solutions into the general solution

$$\vec{y} = c_1 e^{-2x} \cdot \begin{bmatrix} -\frac{1}{8} \\ \frac{1}{4} \\ -\frac{1}{2} \\ 1 \end{bmatrix} + c_2 e^{2x} \cdot \begin{bmatrix} \frac{1}{8} \\ \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} + \begin{bmatrix} -\frac{c_3 \sin(2x)}{8} - \frac{c_4 \cos(2x)}{8} \\ -\frac{c_3 \cos(2x)}{4} + \frac{c_4 \sin(2x)}{4} \\ \frac{c_3 \sin(2x)}{2} + \frac{c_4 \cos(2x)}{2} \\ c_3 \cos(2x) - c_4 \sin(2x) \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y = -\frac{c_1 e^{-2x}}{8} + \frac{c_2 e^{2x}}{8} - \frac{c_4 \cos(2x)}{8} - \frac{c_3 \sin(2x)}{8}$$

Maple trace

```
`Methods for high order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
dsolve(diff(y(x),x$4)-16*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{2x}c_1 + c_2e^{-2x} + c_3 \sin(2x) + c_4 \cos(2x)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 36

```
DSolve[y''''[x]-16*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1e^{2x} + c_3e^{-2x} + c_2 \cos(2x) + c_4 \sin(2x)$$

7.7 problem 4(h)

7.7.1 Maple step by step solution 668

Internal problem ID [5978]

Internal file name [OUTPUT/5226_Sunday_June_05_2022_03_27_56_PM_99880841/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 74

Problem number: 4(h).

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_3rd_order , _missing_x]]
```

$$y''' - 3y' - 2y = 0$$

The characteristic equation is

$$\lambda^3 - 3\lambda - 2 = 0$$

The roots of the above equation are

$$\lambda_1 = 2$$

$$\lambda_2 = -1$$

$$\lambda_3 = -1$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 e^{-x} + x e^{-x} c_2 + e^{2x} c_3$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = e^{-x}$$

$$y_2 = x e^{-x}$$

$$y_3 = e^{2x}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-x} + x e^{-x} c_2 + e^{2x} c_3 \quad (1)$$

Verification of solutions

$$y = c_1 e^{-x} + x e^{-x} c_2 + e^{2x} c_3$$

Verified OK.

7.7.1 Maple step by step solution

Let's solve

$$y''' - 3y' - 2y = 0$$

- Highest derivative means the order of the ODE is 3

$$y'''$$

- Convert linear ODE into a system of first order ODEs

- Define new variable $y_1(x)$

$$y_1(x) = y$$

- Define new variable $y_2(x)$

$$y_2(x) = y'$$

- Define new variable $y_3(x)$

$$y_3(x) = y''$$

- Isolate for $y_3'(x)$ using original ODE

$$y_3'(x) = 3y_2(x) + 2y_1(x)$$

Convert linear ODE into a system of first order ODEs

$$[y_2(x) = y_1'(x), y_3(x) = y_2'(x), y_3'(x) = 3y_2(x) + 2y_1(x)]$$

- Define vector

$$\vec{y}(x) = \begin{bmatrix} y_1(x) \\ y_2(x) \\ y_3(x) \end{bmatrix}$$

- System to solve

$$\vec{y}'(x) = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 2 & 3 & 0 \end{bmatrix} \cdot \vec{y}(x)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 2 & 3 & 0 \end{bmatrix}$$

- Rewrite the system as

$$\vec{y}'(x) = A \cdot \vec{y}(x)$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[-1, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right], \left[-1, \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \right], \left[2, \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair, with eigenvalue of algebraic multiplicity 2

$$\left[-1, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right]$$

- First solution from eigenvalue -1

$$\vec{y}_1(x) = e^{-x} \cdot \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}$$

- Form of the 2nd homogeneous solution where \vec{p} is to be solved for, $\lambda = -1$ is the eigenvalue, a

$$\vec{y}_2(x) = e^{\lambda x} (x\vec{v} + \vec{p})$$

- Note that the x multiplying \vec{v} makes this solution linearly independent to the 1st solution obtained

- Substitute $\vec{y}_2(x)$ into the homogeneous system

$$\lambda e^{\lambda x} (x\vec{v} + \vec{p}) + e^{\lambda x} \vec{v} = (e^{\lambda x} A) \cdot (x\vec{v} + \vec{p})$$

- Use the fact that \vec{v} is an eigenvector of A

$$\lambda e^{\lambda x} (x\vec{v} + \vec{p}) + e^{\lambda x} \vec{v} = e^{\lambda x} (\lambda x\vec{v} + A \cdot \vec{p})$$

- Simplify equation

$$\lambda \vec{p} + \vec{v} = A \cdot \vec{p}$$

- Make use of the identity matrix I

$$(\lambda \cdot I) \cdot \vec{p} + \vec{v} = A \cdot \vec{p}$$

- Condition \vec{p} must meet for $\vec{y}_2(x)$ to be a solution to the homogeneous system

$$(A - \lambda \cdot I) \cdot \vec{p} = \vec{v}$$

- Choose \vec{p} to use in the second solution to the homogeneous system from eigenvalue -1

$$\left(\begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 2 & 3 & 0 \end{bmatrix} - (-1) \cdot \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right) \cdot \vec{p} = \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}$$

- Choice of \vec{p}

$$\vec{p} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

- Second solution from eigenvalue -1

$$\vec{y}_2(x) = e^{-x} \cdot \left(x \cdot \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} + \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \right)$$

- Consider eigenpair

$$\left[2, \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_3 = e^{2x} \cdot \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{y} = c_1 \vec{y}_1(x) + c_2 \vec{y}_2(x) + c_3 \vec{y}_3$$

- Substitute solutions into the general solution

$$\vec{y} = c_1 e^{-x} \cdot \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} + c_2 e^{-x} \cdot \left(x \cdot \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} + \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \right) + e^{2x} c_3 \cdot \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y = (c_2(1+x) + c_1) e^{-x} + \frac{e^{2x} c_3}{4}$$

Maple trace

```
`Methods for third order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(diff(y(x),x$3)-3*diff(y(x),x)-2*y(x)=0,y(x), singsol=all)
```

$$y(x) = (c_3 x + c_2) e^{-x} + e^{2x} c_1$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 26

```
DSolve[y'''[x]-3*y'[x]-2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x} (c_2 x + c_3 e^{3x} + c_1)$$

7.8 problem 4(i)

Internal problem ID [5979]

Internal file name [OUTPUT/5227_Sunday_June_05_2022_03_27_57_PM_31923992/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 74

Problem number: 4(i).

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_3rd_order , _missing_x]]
```

$$y''' - 3iy'' - 3y' + iy = 0$$

The characteristic equation is

$$\lambda^3 - 3i\lambda^2 - 3\lambda + i = 0$$

The roots of the above equation are

$$\lambda_1 = i$$

$$\lambda_2 = i$$

$$\lambda_3 = i$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 e^{ix} + e^{ix} c_2 x + x^2 e^{ix} c_3$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = e^{ix}$$

$$y_2 = x e^{ix}$$

$$y_3 = x^2 e^{ix}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{ix} + e^{ix} c_2 x + x^2 e^{ix} c_3 \quad (1)$$

Verification of solutions

$$y = c_1 e^{ix} + e^{ix} c_2 x + x^2 e^{ix} c_3$$

Verified OK.

Maple trace

```
`Methods for third order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve(diff(y(x),x$3)-3*I*diff(y(x),x$2)-3*diff(y(x),x)+I*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{ix}(c_3 x^2 + c_2 x + c_1)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 25

```
DSolve[y'''[x]-3*I*y''[x]-3*y'[x]+I*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{ix}(x(c_3 x + c_2) + c_1)$$

8 Chapter 2. Linear equations with constant coefficients. Page 79

8.1	problem 1(c)	675
8.2	problem 2(c)	682

8.1 problem 1(c)

8.1.1 Maple step by step solution 677

Internal problem ID [5980]

Internal file name [OUTPUT/5228_Sunday_June_05_2022_03_27_58_PM_63047303/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 79

Problem number: 1(c).

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_3rd_order , _missing_x]]
```

$$y''' - 4y' = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 1, y''(0) = 0]$$

The characteristic equation is

$$\lambda^3 - 4\lambda = 0$$

The roots of the above equation are

$$\lambda_1 = 0$$

$$\lambda_2 = 2$$

$$\lambda_3 = -2$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 + c_2 e^{-2x} + e^{2x} c_3$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = 1$$

$$y_2 = e^{-2x}$$

$$y_3 = e^{2x}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 + c_2 e^{-2x} + e^{2x} c_3 \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_1 + c_2 + c_3 \quad (1A)$$

Taking derivative of the solution gives

$$y' = -2c_2 e^{-2x} + 2e^{2x} c_3$$

substituting $y' = 1$ and $x = 0$ in the above gives

$$1 = -2c_2 + 2c_3 \quad (2A)$$

Taking two derivatives of the solution gives

$$y'' = 4c_2 e^{-2x} + 4e^{2x} c_3$$

substituting $y'' = 0$ and $x = 0$ in the above gives

$$0 = 4c_2 + 4c_3 \quad (3A)$$

Equations {1A,2A,3A} are now solved for $\{c_1, c_2, c_3\}$. Solving for the constants gives

$$\begin{aligned} c_1 &= 0 \\ c_2 &= -\frac{1}{4} \\ c_3 &= \frac{1}{4} \end{aligned}$$

Substituting these values back in above solution results in

$$y = -\frac{e^{-2x}}{4} + \frac{e^{2x}}{4}$$

Summary

The solution(s) found are the following

$$y = -\frac{e^{-2x}}{4} + \frac{e^{2x}}{4} \quad (1)$$




Figure 129: Solution plot

Verification of solutions

$$y = -\frac{e^{-2x}}{4} + \frac{e^{2x}}{4}$$

Verified OK.

8.1.1 Maple step by step solution

Let's solve

$$\left[y''' - 4y' = 0, y(0) = 0, y'|_{\{x=0\}} = 1, y''|_{\{x=0\}} = 0 \right]$$

- Highest derivative means the order of the ODE is 3
 y'''
- Convert linear ODE into a system of first order ODEs
 - Define new variable $y_1(x)$
 $y_1(x) = y$
 - Define new variable $y_2(x)$
 $y_2(x) = y'$
 - Define new variable $y_3(x)$

$$y_3(x) = y''$$

- Isolate for $y_3'(x)$ using original ODE

$$y_3'(x) = 4y_2(x)$$

Convert linear ODE into a system of first order ODEs

$$[y_2(x) = y_1'(x), y_3(x) = y_2'(x), y_3'(x) = 4y_2(x)]$$

- Define vector

$$\vec{y}(x) = \begin{bmatrix} y_1(x) \\ y_2(x) \\ y_3(x) \end{bmatrix}$$

- System to solve

$$\vec{y}'(x) = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 4 & 0 \end{bmatrix} \cdot \vec{y}(x)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 4 & 0 \end{bmatrix}$$

- Rewrite the system as

$$\vec{y}'(x) = A \cdot \vec{y}(x)$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[-2, \begin{bmatrix} \frac{1}{4} \\ -\frac{1}{2} \\ 1 \end{bmatrix} \right], \left[0, \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \right], \left[2, \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[-2, \begin{bmatrix} \frac{1}{4} \\ -\frac{1}{2} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_1 = e^{-2x} \cdot \begin{bmatrix} \frac{1}{4} \\ -\frac{1}{2} \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[0, \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_2 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

- Consider eigenpair

$$\left[2, \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_3 = e^{2x} \cdot \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{y} = c_1 \vec{y}_1 + c_2 \vec{y}_2 + c_3 \vec{y}_3$$

- Substitute solutions into the general solution

$$\vec{y} = c_1 e^{-2x} \cdot \begin{bmatrix} \frac{1}{4} \\ -\frac{1}{2} \\ 1 \end{bmatrix} + e^{2x} c_3 \cdot \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} + \begin{bmatrix} c_2 \\ 0 \\ 0 \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y = \frac{c_1 e^{-2x}}{4} + \frac{e^{2x} c_3}{4} + c_2$$

- Use the initial condition $y(0) = 0$

$$0 = \frac{c_1}{4} + \frac{c_3}{4} + c_2$$

- Calculate the 1st derivative of the solution

$$y' = -\frac{c_1 e^{-2x}}{2} + \frac{e^{2x} c_3}{2}$$

- Use the initial condition $y'|_{\{x=0\}} = 1$

$$1 = -\frac{c_1}{2} + \frac{c_3}{2}$$

- Calculate the 2nd derivative of the solution

$$y'' = c_1 e^{-2x} + e^{2x} c_3$$

- Use the initial condition $y''|_{\{x=0\}} = 0$

$$0 = c_1 + c_3$$

- Solve for the unknown coefficients

$$\{c_1 = -1, c_2 = 0, c_3 = 1\}$$

- Solution to the IVP

$$y = -\frac{e^{-2x}}{4} + \frac{e^{2x}}{4}$$

Maple trace

```

`Methods for third order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 17

```
dsolve([diff(y(x),x$3)-4*diff(y(x),x)=0,y(0) = 0, D(y)(0) = 1, (D@@2)(y)(0) = 0],y(x), sings
```

$$y(x) = \frac{e^{2x}}{4} - \frac{e^{-2x}}{4}$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 69

```
DSolve[{y'''[x]-4*y[x]==0,{y[0]==0,y'[0]==1,y''[0]==0}},y[x],x,IncludeSingularSolutions -> T
```

$$y(x) \rightarrow \frac{e^{-\frac{x}{\sqrt[3]{2}}} \left(e^{\frac{3x}{\sqrt[3]{2}}} + \sqrt{3} \sin \left(\frac{\sqrt{3}x}{\sqrt[3]{2}} \right) - \cos \left(\frac{\sqrt{3}x}{\sqrt[3]{2}} \right) \right)}{3 \cdot 2^{2/3}}$$

8.2 problem 2(c)

8.2.1 Maple step by step solution 685

Internal problem ID [5981]

Internal file name [OUTPUT/5229_Sunday_June_05_2022_03_28_00_PM_96542066/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 79

Problem number: 2(c).

ODE order: 5.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_high_order , _missing_x]]
```

$$y^{(5)} - y'''' - y' + y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 0, y''(0) = 0, y'''(0) = 0, y''''(0) = 0]$$

The characteristic equation is

$$\lambda^5 - \lambda^4 - \lambda + 1 = 0$$

The roots of the above equation are

$$\lambda_1 = -1$$

$$\lambda_2 = i$$

$$\lambda_3 = -i$$

$$\lambda_4 = 1$$

$$\lambda_5 = 1$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 e^{-x} + c_2 e^x + x e^x c_3 + e^{-ix} c_4 + e^{ix} c_5$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = e^{-x}$$

$$y_2 = e^x$$

$$y_3 = x e^x$$

$$y_4 = e^{-ix}$$

$$y_5 = e^{ix}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 e^{-x} + c_2 e^x + x e^x c_3 + e^{-ix} c_4 + e^{ix} c_5 \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 0$ in the above gives

$$1 = c_1 + c_2 + c_4 + c_5 \quad (1A)$$

Taking derivative of the solution gives

$$y' = -c_1 e^{-x} + c_2 e^x + c_3 e^x + x e^x c_3 - i e^{-ix} c_4 + i e^{ix} c_5$$

substituting $y' = 0$ and $x = 0$ in the above gives

$$0 = -c_4 i + c_5 i - c_1 + c_2 + c_3 \quad (2A)$$

Taking two derivatives of the solution gives

$$y'' = c_1 e^{-x} + c_2 e^x + 2c_3 e^x + x e^x c_3 - e^{-ix} c_4 - e^{ix} c_5$$

substituting $y'' = 0$ and $x = 0$ in the above gives

$$0 = c_1 + c_2 + 2c_3 - c_4 - c_5 \quad (3A)$$

Taking three derivatives of the solution gives

$$y''' = -c_1 e^{-x} + c_2 e^x + 3c_3 e^x + x e^x c_3 + i e^{-ix} c_4 - i e^{ix} c_5$$

substituting $y''' = 0$ and $x = 0$ in the above gives

$$0 = c_4 i - c_5 i - c_1 + c_2 + 3c_3 \quad (4A)$$

Taking four derivatives of the solution gives

$$y'''' = c_1 e^{-x} + c_2 e^x + 4c_3 e^x + x e^x c_3 + e^{-ix} c_4 + e^{ix} c_5$$

substituting $y'''' = 0$ and $x = 0$ in the above gives

$$0 = c_1 + c_2 + 4c_3 + c_4 + c_5 \quad (5A)$$

Equations {1A,2A,3A,4A,5A} are now solved for $\{c_1, c_2, c_3, c_4, c_5\}$. Solving for the constants gives

$$\begin{aligned} c_1 &= \frac{1}{8} \\ c_2 &= \frac{5}{8} \\ c_3 &= -\frac{1}{4} \\ c_4 &= \frac{1}{8} - \frac{i}{8} \\ c_5 &= \frac{1}{8} + \frac{i}{8} \end{aligned}$$

Substituting these values back in above solution results in

$$y = \frac{e^{-x}}{8} + \frac{5e^x}{8} - \frac{x e^x}{4} + \frac{\cos(x)}{4} - \frac{\sin(x)}{4}$$

Summary

The solution(s) found are the following

$$y = \frac{e^{-x}}{8} + \frac{5e^x}{8} - \frac{x e^x}{4} + \frac{\cos(x)}{4} - \frac{\sin(x)}{4} \quad (1)$$




Figure 130: Solution plot

Verification of solutions

$$y = \frac{e^{-x}}{8} + \frac{5e^x}{8} - \frac{x e^x}{4} + \frac{\cos(x)}{4} - \frac{\sin(x)}{4}$$

Verified OK.

8.2.1 Maple step by step solution

Let's solve

$$\left[y^{(5)} - y'''' - y' + y = 0, y(0) = 1, y'|_{\{x=0\}} = 0, y''|_{\{x=0\}} = 0, y'''|_{\{x=0\}} = 0, y''''|_{\{x=0\}} = 0 \right]$$

- Highest derivative means the order of the ODE is 5
 $y^{(5)}$
- Convert linear ODE into a system of first order ODEs
 - Define new variable $y_1(x)$
 $y_1(x) = y$
 - Define new variable $y_2(x)$
 $y_2(x) = y'$
 - Define new variable $y_3(x)$

$$y_3(x) = y''$$

- Define new variable $y_4(x)$

$$y_4(x) = y'''$$

- Define new variable $y_5(x)$

$$y_5(x) = y''''$$

- Isolate for $y_5'(x)$ using original ODE

$$y_5'(x) = y_5(x) + y_2(x) - y_1(x)$$

Convert linear ODE into a system of first order ODEs

$$[y_2(x) = y_1'(x), y_3(x) = y_2'(x), y_4(x) = y_3'(x), y_5(x) = y_4'(x), y_5'(x) = y_5(x) + y_2(x) - y_1(x)]$$

- Define vector

$$\vec{y}(x) = \begin{bmatrix} y_1(x) \\ y_2(x) \\ y_3(x) \\ y_4(x) \\ y_5(x) \end{bmatrix}$$

- System to solve

$$\vec{y}'(x) = \begin{bmatrix} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ -1 & 1 & 0 & 0 & 1 \end{bmatrix} \cdot \vec{y}(x)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ -1 & 1 & 0 & 0 & 1 \end{bmatrix}$$

- Rewrite the system as

$$\vec{y}'(x) = A \cdot \vec{y}(x)$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[\begin{array}{c} 1 \\ -1 \end{array} \right], \left[\begin{array}{c} 1 \\ -1 \\ 1 \\ 1 \end{array} \right] \right], \left[1, \left[\begin{array}{c} 1 \\ 1 \\ 1 \\ 1 \end{array} \right] \right], \left[1, \left[\begin{array}{c} 0 \\ 0 \\ 0 \\ 0 \end{array} \right] \right], \left[-I, \left[\begin{array}{c} 1 \\ -1 \\ I \\ 1 \end{array} \right] \right], \left[I, \left[\begin{array}{c} 1 \\ I \\ -1 \\ 1 \end{array} \right] \right]$$

- Consider eigenpair

$$\left[\begin{array}{c} -1, \\ 1 \\ -1 \\ 1 \end{array} \right], \left[\begin{array}{c} 1 \\ -1 \\ 1 \\ 1 \end{array} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_1 = e^{-x} \cdot \left[\begin{array}{c} 1 \\ -1 \\ 1 \\ -1 \\ 1 \end{array} \right]$$

- Consider eigenpair, with eigenvalue of algebraic multiplicity 2

$$\left[1, \left[\begin{array}{c} 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{array} \right] \right]$$

- First solution from eigenvalue 1

$$\vec{y}_2(x) = e^x \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}$$

- Form of the 2nd homogeneous solution where \vec{p} is to be solved for, $\lambda = 1$ is the eigenvalue, and

$$\vec{y}_3(x) = e^{\lambda x} (x\vec{v} + \vec{p})$$

- Note that the x multiplying \vec{v} makes this solution linearly independent to the 1st solution obtained

- Substitute $\vec{y}_3(x)$ into the homogeneous system

$$\lambda e^{\lambda x} (x\vec{v} + \vec{p}) + e^{\lambda x} \vec{v} = (e^{\lambda x} A) \cdot (x\vec{v} + \vec{p})$$

- Use the fact that \vec{v} is an eigenvector of A

$$\lambda e^{\lambda x} (x\vec{v} + \vec{p}) + e^{\lambda x} \vec{v} = e^{\lambda x} (\lambda x\vec{v} + A \cdot \vec{p})$$

- Simplify equation

$$\lambda \vec{p} + \vec{v} = A \cdot \vec{p}$$

- Make use of the identity matrix I

$$(\lambda \cdot I) \cdot \vec{p} + \vec{v} = A \cdot \vec{p}$$

- Condition \vec{p} must meet for $\vec{y}_3(x)$ to be a solution to the homogeneous system

$$(A - \lambda \cdot I) \cdot \vec{p} = \vec{v}$$

- Choose \vec{p} to use in the second solution to the homogeneous system from eigenvalue 1

$$\left(\begin{bmatrix} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ -1 & 1 & 0 & 0 & 1 \end{bmatrix} - 1 \cdot \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix} \right) \cdot \vec{p} = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}$$

- Choice of \vec{p}

$$\vec{p} = \begin{bmatrix} -1 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

- Second solution from eigenvalue 1

$$\vec{y}_3(x) = e^x \cdot \left(x \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} + \begin{bmatrix} -1 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} \right)$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$\left[-I, \begin{bmatrix} 1 \\ -I \\ -1 \\ I \\ 1 \end{bmatrix} \right]$$

- Solution from eigenpair

$$e^{-Ix} \cdot \begin{bmatrix} 1 \\ -I \\ -1 \\ I \\ 1 \end{bmatrix}$$

- Use Euler identity to write solution in terms of sin and cos

$$(\cos(x) - I \sin(x)) \cdot \begin{bmatrix} 1 \\ -I \\ -1 \\ I \\ 1 \end{bmatrix}$$

- Simplify expression

$$\begin{bmatrix} \cos(x) - I \sin(x) \\ -I(\cos(x) - I \sin(x)) \\ -\cos(x) + I \sin(x) \\ I(\cos(x) - I \sin(x)) \\ \cos(x) - I \sin(x) \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\vec{y}_4(x) = \begin{bmatrix} \cos(x) \\ -\sin(x) \\ -\cos(x) \\ \sin(x) \\ \cos(x) \end{bmatrix}, \vec{y}_5(x) = \begin{bmatrix} -\sin(x) \\ -\cos(x) \\ \sin(x) \\ \cos(x) \\ -\sin(x) \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{y} = c_1 \vec{y}_1 + c_2 \vec{y}_2(x) + c_3 \vec{y}_3(x) + c_4 \vec{y}_4(x) + c_5 \vec{y}_5(x)$$

- Substitute solutions into the general solution

$$\vec{y} = c_1 e^{-x} \cdot \begin{bmatrix} 1 \\ -1 \\ 1 \\ -1 \\ 1 \end{bmatrix} + c_2 e^x \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} + c_3 e^x \cdot \left(x \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} + \begin{bmatrix} -1 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} \right) + \begin{bmatrix} c_4 \cos(x) - c_5 \sin(x) \\ -c_4 \sin(x) - c_5 \cos(x) \\ -c_4 \cos(x) + c_5 \sin(x) \\ c_4 \sin(x) + c_5 \cos(x) \\ c_4 \cos(x) - c_5 \sin(x) \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y = c_1 e^{-x} + (c_3(x-1) + c_2) e^x + c_4 \cos(x) - c_5 \sin(x)$$

- Use the initial condition $y(0) = 1$

$$1 = c_1 - c_3 + c_2 + c_4$$

- Calculate the 1st derivative of the solution

$$y' = -c_1 e^{-x} + c_3 e^x + (c_3(x-1) + c_2) e^x - c_4 \sin(x) - c_5 \cos(x)$$
- Use the initial condition $y' \Big|_{\{x=0\}} = 0$

$$0 = -c_1 + c_2 - c_5$$
- Calculate the 2nd derivative of the solution

$$y'' = c_1 e^{-x} + 2c_3 e^x + (c_3(x-1) + c_2) e^x - c_4 \cos(x) + c_5 \sin(x)$$
- Use the initial condition $y'' \Big|_{\{x=0\}} = 0$

$$0 = c_1 + c_3 + c_2 - c_4$$
- Calculate the 3rd derivative of the solution

$$y''' = -c_1 e^{-x} + 3c_3 e^x + (c_3(x-1) + c_2) e^x + c_4 \sin(x) + c_5 \cos(x)$$
- Use the initial condition $y''' \Big|_{\{x=0\}} = 0$

$$0 = -c_1 + 2c_3 + c_2 + c_5$$
- Calculate the 4th derivative of the solution

$$y'''' = c_1 e^{-x} + 4c_3 e^x + (c_3(x-1) + c_2) e^x + c_4 \cos(x) - c_5 \sin(x)$$
- Use the initial condition $y'''' \Big|_{\{x=0\}} = 0$

$$0 = c_1 + 3c_3 + c_2 + c_4$$
- Solve for the unknown coefficients

$$\left\{ c_1 = \frac{1}{8}, c_2 = \frac{3}{8}, c_3 = -\frac{1}{4}, c_4 = \frac{1}{4}, c_5 = \frac{1}{4} \right\}$$
- Solution to the IVP

$$y = \frac{e^{-x}}{8} + \frac{(-2x+5)e^x}{8} + \frac{\cos(x)}{4} - \frac{\sin(x)}{4}$$

Maple trace

```

`Methods for high order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 28

```
dsolve([diff(y(x),x$5)-diff(y(x),x$4)-diff(y(x),x)+y(x)=0,y(0) = 1, D(y)(0) = 0, (D@@2)(y)(0)
```

$$y(x) = \frac{e^{-x}}{8} + \frac{(-2x + 5)e^x}{8} + \frac{\cos(x)}{4} - \frac{\sin(x)}{4}$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 34

```
DSolve[{y''''[x]-y''''[x]-y'[x]+y[x]==0,{y[0]==1,y'[0]==0,y''[0]==0,y'''[0]==0,y''''[0]==0}
```

$$y(x) \rightarrow \frac{1}{8}(-2e^x x + e^{-x} + 5e^x - 2\sin(x) + 2\cos(x))$$

9 Chapter 2. Linear equations with constant coefficients. Page 83

9.1	problem 1(a)	694
9.2	problem 1(b)	704
9.3	problem 1(c)	714
9.4	problem 1(d)	720
9.5	problem 1(e)	723
9.6	problem 2	729
9.7	problem 3(a)	736
9.8	problem 3(b)	738
9.9	problem 5(b)	747

9.1 problem 1(a)

9.1.1	Solving as second order linear constant coeff ode	694
9.1.2	Solving as second order ode can be made integrable ode	696
9.1.3	Solving using Kovacic algorithm	698
9.1.4	Maple step by step solution	702

Internal problem ID [5982]

Internal file name [OUTPUT/5230_Sunday_June_05_2022_03_28_02_PM_72058856/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 83

Problem number: 1(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "second_order_ode_can_be_made_integrable"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' + y = 0$$

9.1.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i\end{aligned}$$

Hence

$$\lambda_1 = +i$$

$$\lambda_2 = -i$$

Which simplifies to

$$\lambda_1 = i$$

$$\lambda_2 = -i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0(\cos(x) c_1 + c_2 \sin(x))$$

Or

$$y = \cos(x) c_1 + c_2 \sin(x)$$

Summary

The solution(s) found are the following

$$y = \cos(x) c_1 + c_2 \sin(x) \tag{1}$$




Figure 131: Slope field plot

Verification of solutions

$$y = \cos(x) c_1 + c_2 \sin(x)$$

Verified OK.

9.1.2 Solving as second order ode can be made integrable ode

Multiplying the ode by y' gives

$$y' y'' + y' y = 0$$

Integrating the above w.r.t x gives

$$\int (y' y'' + y' y) dx = 0$$

$$\frac{y'^2}{2} + \frac{y^2}{2} = c_2$$

Which is now solved for y . Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \sqrt{-y^2 + 2c_1} \tag{1}$$

$$y' = -\sqrt{-y^2 + 2c_1} \tag{2}$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int \frac{1}{\sqrt{-y^2 + 2c_1}} dy = \int dx$$
$$\arctan\left(\frac{y}{\sqrt{-y^2 + 2c_1}}\right) = c_2 + x$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{1}{\sqrt{-y^2 + 2c_1}} dy = \int dx$$
$$-\arctan\left(\frac{y}{\sqrt{-y^2 + 2c_1}}\right) = x + c_3$$

Summary

The solution(s) found are the following

$$\arctan\left(\frac{y}{\sqrt{-y^2 + 2c_1}}\right) = c_2 + x \tag{1}$$

$$-\arctan\left(\frac{y}{\sqrt{-y^2 + 2c_1}}\right) = x + c_3 \tag{2}$$




Figure 132: Slope field plot

Verification of solutions

$$\arctan\left(\frac{y}{\sqrt{-y^2 + 2c_1}}\right) = c_2 + x$$

Verified OK.

$$-\arctan\left(\frac{y}{\sqrt{-y^2 + 2c_1}}\right) = x + c_3$$

Verified OK.

9.1.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= 1 \\B &= 0 \\C &= 1\end{aligned}\tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x)\tag{4}$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\&= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned}\tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1}\tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= -1 \\t &= 1\end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x)\tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 136: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}
 y_1 &= z_1 \\
 &= \cos(x)
 \end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x)) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1(\cos(x)) + c_2(\cos(x) (\tan(x))) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \cos(x) c_1 + c_2 \sin(x) \tag{1}$$




Figure 133: Slope field plot

Verification of solutions

$$y = \cos(x) c_1 + c_2 \sin(x)$$

Verified OK.

9.1.4 Maple step by step solution

Let's solve

$$y'' + y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

- $r = (-I, I)$
- 1st solution of the ODE
 $y_1(x) = \cos(x)$
- 2nd solution of the ODE
 $y_2(x) = \sin(x)$
- General solution of the ODE
 $y = c_1 y_1(x) + c_2 y_2(x)$
- Substitute in solutions
 $y = \cos(x) c_1 + c_2 \sin(x)$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(diff(y(x),x$2)+y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 \sin(x) + \cos(x) c_2$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 16

```
DSolve[y''[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos(x) + c_2 \sin(x)$$

9.2 problem 1(b)

9.2.1	Solving as second order linear constant coeff ode	704
9.2.2	Solving as second order ode can be made integrable ode	706
9.2.3	Solving using Kovacic algorithm	708
9.2.4	Maple step by step solution	712

Internal problem ID [5983]

Internal file name [OUTPUT/5231_Sunday_June_05_2022_03_28_03_PM_32615447/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 83

Problem number: 1(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "second_order_ode_can_be_made_integrable"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' - y = 0$$

9.2.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = -1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 1 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = -1$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(-1)} \\ &= \pm 1\end{aligned}$$

Hence

$$\lambda_1 = +1$$

$$\lambda_2 = -1$$

Which simplifies to

$$\lambda_1 = 1$$

$$\lambda_2 = -1$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(1)x} + c_2 e^{(-1)x}$$

Or

$$y = c_1 e^x + c_2 e^{-x}$$

Summary

The solution(s) found are the following

$$y = c_1 e^x + c_2 e^{-x} \tag{1}$$




Figure 134: Slope field plot

Verification of solutions

$$y = c_1 e^x + c_2 e^{-x}$$

Verified OK.

9.2.2 Solving as second order ode can be made integrable ode

Multiplying the ode by y' gives

$$y' y'' - y' y = 0$$

Integrating the above w.r.t x gives

$$\int (y' y'' - y' y) dx = 0$$

$$\frac{y'^2}{2} - \frac{y^2}{2} = c_2$$

Which is now solved for y . Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \sqrt{y^2 + 2c_1} \tag{1}$$

$$y' = -\sqrt{y^2 + 2c_1} \tag{2}$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int \frac{1}{\sqrt{y^2 + 2c_1}} dy = \int dx$$
$$\ln(y + \sqrt{y^2 + 2c_1}) = c_2 + x$$

Raising both side to exponential gives

$$y + \sqrt{y^2 + 2c_1} = e^{c_2+x}$$

Which simplifies to

$$y + \sqrt{y^2 + 2c_1} = c_3 e^x$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{1}{\sqrt{y^2 + 2c_1}} dy = \int dx$$
$$-\ln(y + \sqrt{y^2 + 2c_1}) = x + c_4$$

Raising both side to exponential gives

$$\frac{1}{y + \sqrt{y^2 + 2c_1}} = e^{x+c_4}$$

Which simplifies to

$$\frac{1}{y + \sqrt{y^2 + 2c_1}} = c_5 e^x$$

Summary

The solution(s) found are the following

$$y = \frac{(e^{2x} c_3^2 - 2c_1) e^{-x}}{2c_3} \tag{1}$$

$$y = -\frac{(2c_1 c_5^2 e^{2x} - 1) e^{-x}}{2c_5} \tag{2}$$




Figure 135: Slope field plot

Verification of solutions

$$y = \frac{(e^{2x}c_3^2 - 2c_1)e^{-x}}{2c_3}$$

Verified OK.

$$y = -\frac{(2c_1c_5^2e^{2x} - 1)e^{-x}}{2c_5}$$

Verified OK.

9.2.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' - y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 0 \tag{3}$$

$$C = -1$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{1}{1} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 138: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-x}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}
 y_1 &= z_1 \\
 &= e^{-x}
 \end{aligned}$$

Which simplifies to

$$y_1 = e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= e^{-x} \int \frac{1}{e^{-2x}} dx \\ &= e^{-x} \left(\frac{e^{2x}}{2} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-x}) + c_2 \left(e^{-x} \left(\frac{e^{2x}}{2} \right) \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-x} + \frac{c_2 e^x}{2} \tag{1}$$




Figure 136: Slope field plot

Verification of solutions

$$y = c_1 e^{-x} + \frac{c_2 e^x}{2}$$

Verified OK.

9.2.4 Maple step by step solution

Let's solve

$$y'' - y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 - 1 = 0$$

- Factor the characteristic polynomial

$$(r - 1)(r + 1) = 0$$

- Roots of the characteristic polynomial

- $r = (-1, 1)$
 - 1st solution of the ODE
 $y_1(x) = e^{-x}$
 - 2nd solution of the ODE
 $y_2(x) = e^x$
 - General solution of the ODE
 $y = c_1 y_1(x) + c_2 y_2(x)$
 - Substitute in solutions
 $y = c_1 e^{-x} + c_2 e^x$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
dsolve(diff(y(x),x$2)-y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{-x} + e^x c_2$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 20

```
DSolve[y''[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^x + c_2 e^{-x}$$

9.3 problem 1(c)

9.3.1 Maple step by step solution 715

Internal problem ID [5984]

Internal file name [OUTPUT/5232_Sunday_June_05_2022_03_28_04_PM_84836869/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 83

Problem number: 1(c).

ODE order: 4.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_high_order , _missing_x]]
```

$$y'''' - y = 0$$

The characteristic equation is

$$\lambda^4 - 1 = 0$$

The roots of the above equation are

$$\lambda_1 = 1$$

$$\lambda_2 = -1$$

$$\lambda_3 = i$$

$$\lambda_4 = -i$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 e^{-x} + c_2 e^x + e^{-ix} c_3 + e^{ix} c_4$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = e^{-x}$$

$$y_2 = e^x$$

$$y_3 = e^{-ix}$$

$$y_4 = e^{ix}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-x} + c_2 e^x + e^{-ix} c_3 + e^{ix} c_4 \quad (1)$$

Verification of solutions

$$y = c_1 e^{-x} + c_2 e^x + e^{-ix} c_3 + e^{ix} c_4$$

Verified OK.

9.3.1 Maple step by step solution

Let's solve

$$y'''' - y = 0$$

- Highest derivative means the order of the ODE is 4

$$y''''$$

- Convert linear ODE into a system of first order ODEs

- Define new variable $y_1(x)$

$$y_1(x) = y$$

- Define new variable $y_2(x)$

$$y_2(x) = y'$$

- Define new variable $y_3(x)$

$$y_3(x) = y''$$

- Define new variable $y_4(x)$

$$y_4(x) = y'''$$

- Isolate for $y_4'(x)$ using original ODE

$$y_4'(x) = y_1(x)$$

Convert linear ODE into a system of first order ODEs

$$[y_2(x) = y_1'(x), y_3(x) = y_2'(x), y_4(x) = y_3'(x), y_4'(x) = y_1(x)]$$

- Define vector

$$\vec{y}(x) = \begin{bmatrix} y_1(x) \\ y_2(x) \\ y_3(x) \\ y_4(x) \end{bmatrix}$$

- System to solve

$$\vec{y}'(x) = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \end{bmatrix} \cdot \vec{y}(x)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \end{bmatrix}$$

- Rewrite the system as

$$\vec{y}'(x) = A \cdot \vec{y}(x)$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[-1, \begin{bmatrix} -1 \\ 1 \\ -1 \\ 1 \end{bmatrix} \right] \right], \left[\left[1, \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} \right] \right], \left[\left[-I, \begin{bmatrix} -I \\ -1 \\ I \\ 1 \end{bmatrix} \right] \right], \left[\left[I, \begin{bmatrix} I \\ -1 \\ -I \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[\left[-1, \begin{bmatrix} -1 \\ 1 \\ -1 \\ 1 \end{bmatrix} \right] \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_1 = e^{-x} \cdot \begin{bmatrix} -1 \\ 1 \\ -1 \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[1, \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_2 = e^x \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$\left[-I, \begin{bmatrix} -I \\ -1 \\ I \\ 1 \end{bmatrix} \right]$$

- Solution from eigenpair

$$e^{-Ix} \cdot \begin{bmatrix} -I \\ -1 \\ I \\ 1 \end{bmatrix}$$

- Use Euler identity to write solution in terms of sin and cos

$$(\cos(x) - I \sin(x)) \cdot \begin{bmatrix} -I \\ -1 \\ I \\ 1 \end{bmatrix}$$

- Simplify expression

$$\begin{bmatrix} -I(\cos(x) - I \sin(x)) \\ -\cos(x) + I \sin(x) \\ I(\cos(x) - I \sin(x)) \\ \cos(x) - I \sin(x) \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\vec{y}_3(x) = \begin{bmatrix} -\sin(x) \\ -\cos(x) \\ \sin(x) \\ \cos(x) \end{bmatrix}, \vec{y}_4(x) = \begin{bmatrix} -\cos(x) \\ \sin(x) \\ \cos(x) \\ -\sin(x) \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{y} = c_1 \vec{y}_1 + c_2 \vec{y}_2 + c_3 \vec{y}_3(x) + c_4 \vec{y}_4(x)$$

- Substitute solutions into the general solution

$$\vec{y} = c_1 e^{-x} \cdot \begin{bmatrix} -1 \\ 1 \\ -1 \\ 1 \end{bmatrix} + c_2 e^x \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} + \begin{bmatrix} -c_3 \sin(x) - c_4 \cos(x) \\ -c_3 \cos(x) + c_4 \sin(x) \\ c_3 \sin(x) + c_4 \cos(x) \\ c_3 \cos(x) - c_4 \sin(x) \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y = -c_1 e^{-x} + c_2 e^x - c_4 \cos(x) - c_3 \sin(x)$$

Maple trace

```
`Methods for high order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(diff(y(x),x$4)-y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{-x} + e^x c_2 + c_3 \sin(x) + c_4 \cos(x)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 30

```
DSolve[y''''[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^x + c_3 e^{-x} + c_2 \cos(x) + c_4 \sin(x)$$

9.4 problem 1(d)

Internal problem ID [5985]

Internal file name [OUTPUT/5233_Sunday_June_05_2022_03_28_05_PM_69631145/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 83

Problem number: 1(d).

ODE order: 5.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_high_order, _missing_x]]
```

$$y^{(5)} + 2y = 0$$

The characteristic equation is

$$\lambda^5 + 2 = 0$$

The roots of the above equation are

$$\lambda_1 = \cos\left(\frac{\pi}{5}\right) 2^{\frac{1}{5}} + i \sin\left(\frac{\pi}{5}\right) 2^{\frac{1}{5}}$$

$$\lambda_2 = \left(\left(\frac{\sqrt{5}}{4} - \frac{1}{4} \right) \cos\left(\frac{\pi}{5}\right) - \frac{\sqrt{2} \sqrt{5 + \sqrt{5}} \sin\left(\frac{\pi}{5}\right)}{4} \right) 2^{\frac{1}{5}} + i \left(\frac{\sqrt{2} \sqrt{5 + \sqrt{5}} \cos\left(\frac{\pi}{5}\right)}{4} + \left(\frac{\sqrt{5}}{4} - \frac{1}{4} \right) \sin\left(\frac{\pi}{5}\right) \right) 2^{\frac{1}{5}}$$

$$\lambda_3 = \left(\left(-\frac{\sqrt{5}}{4} - \frac{1}{4} \right) \cos\left(\frac{\pi}{5}\right) - \frac{\sqrt{2} \sqrt{5 - \sqrt{5}} \sin\left(\frac{\pi}{5}\right)}{4} \right) 2^{\frac{1}{5}} + i \left(\frac{\sqrt{2} \sqrt{5 - \sqrt{5}} \cos\left(\frac{\pi}{5}\right)}{4} + \left(-\frac{\sqrt{5}}{4} - \frac{1}{4} \right) \sin\left(\frac{\pi}{5}\right) \right) 2^{\frac{1}{5}}$$

$$\lambda_4 = \left(\left(-\frac{\sqrt{5}}{4} - \frac{1}{4} \right) \cos\left(\frac{\pi}{5}\right) + \frac{\sqrt{2} \sqrt{5 - \sqrt{5}} \sin\left(\frac{\pi}{5}\right)}{4} \right) 2^{\frac{1}{5}} + i \left(-\frac{\sqrt{2} \sqrt{5 - \sqrt{5}} \cos\left(\frac{\pi}{5}\right)}{4} + \left(-\frac{\sqrt{5}}{4} - \frac{1}{4} \right) \sin\left(\frac{\pi}{5}\right) \right) 2^{\frac{1}{5}}$$

$$\lambda_5 = \left(\left(\frac{\sqrt{5}}{4} - \frac{1}{4} \right) \cos\left(\frac{\pi}{5}\right) + \frac{\sqrt{2} \sqrt{5 + \sqrt{5}} \sin\left(\frac{\pi}{5}\right)}{4} \right) 2^{\frac{1}{5}} + i \left(-\frac{\sqrt{2} \sqrt{5 + \sqrt{5}} \cos\left(\frac{\pi}{5}\right)}{4} + \left(\frac{\sqrt{5}}{4} - \frac{1}{4} \right) \sin\left(\frac{\pi}{5}\right) \right) 2^{\frac{1}{5}}$$

Therefore the homogeneous solution is

$$y_h(x) = e^{\left(\left(\left(\frac{\sqrt{5}}{4} - \frac{1}{4} \right) \cos\left(\frac{\pi}{5}\right) - \frac{\sqrt{2} \sqrt{5 + \sqrt{5}} \sin\left(\frac{\pi}{5}\right)}{4} \right) 2^{\frac{1}{5}} + i \left(\frac{\sqrt{2} \sqrt{5 + \sqrt{5}} \cos\left(\frac{\pi}{5}\right)}{4} + \left(\frac{\sqrt{5}}{4} - \frac{1}{4} \right) \sin\left(\frac{\pi}{5}\right) \right) 2^{\frac{1}{5}} \right) x} c_1 + e^{\left(\left(\left(-\frac{\sqrt{5}}{4} - \frac{1}{4} \right) \cos\left(\frac{\pi}{5}\right) - \frac{\sqrt{2} \sqrt{5 - \sqrt{5}} \sin\left(\frac{\pi}{5}\right)}{4} \right) 2^{\frac{1}{5}} + i \left(\frac{\sqrt{2} \sqrt{5 - \sqrt{5}} \cos\left(\frac{\pi}{5}\right)}{4} + \left(-\frac{\sqrt{5}}{4} - \frac{1}{4} \right) \sin\left(\frac{\pi}{5}\right) \right) 2^{\frac{1}{5}} \right) x} c_2 + e^{\left(\left(\left(-\frac{\sqrt{5}}{4} - \frac{1}{4} \right) \cos\left(\frac{\pi}{5}\right) + \frac{\sqrt{2} \sqrt{5 - \sqrt{5}} \sin\left(\frac{\pi}{5}\right)}{4} \right) 2^{\frac{1}{5}} + i \left(-\frac{\sqrt{2} \sqrt{5 - \sqrt{5}} \cos\left(\frac{\pi}{5}\right)}{4} + \left(-\frac{\sqrt{5}}{4} - \frac{1}{4} \right) \sin\left(\frac{\pi}{5}\right) \right) 2^{\frac{1}{5}} \right) x} c_3 + e^{\left(\left(\left(\frac{\sqrt{5}}{4} - \frac{1}{4} \right) \cos\left(\frac{\pi}{5}\right) + \frac{\sqrt{2} \sqrt{5 + \sqrt{5}} \sin\left(\frac{\pi}{5}\right)}{4} \right) 2^{\frac{1}{5}} + i \left(-\frac{\sqrt{2} \sqrt{5 + \sqrt{5}} \cos\left(\frac{\pi}{5}\right)}{4} + \left(\frac{\sqrt{5}}{4} - \frac{1}{4} \right) \sin\left(\frac{\pi}{5}\right) \right) 2^{\frac{1}{5}} \right) x} c_4 + e^{\left(\left(\left(\frac{\sqrt{5}}{4} - \frac{1}{4} \right) \cos\left(\frac{\pi}{5}\right) - \frac{\sqrt{2} \sqrt{5 + \sqrt{5}} \sin\left(\frac{\pi}{5}\right)}{4} \right) 2^{\frac{1}{5}} + i \left(\frac{\sqrt{2} \sqrt{5 + \sqrt{5}} \cos\left(\frac{\pi}{5}\right)}{4} + \left(\frac{\sqrt{5}}{4} - \frac{1}{4} \right) \sin\left(\frac{\pi}{5}\right) \right) 2^{\frac{1}{5}} \right) x} c_5$$

Maple trace

```
`Methods for high order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 140

```
dsolve(diff(y(x),x$5)+2*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{\frac{(-i2^{\frac{7}{10}}\sqrt{5-\sqrt{5}}+2^{\frac{1}{5}}\sqrt{5}+2^{\frac{1}{5}})x}{4}} + c_2 e^{-\frac{x(i(\sqrt{5}+1)2^{\frac{7}{10}}\sqrt{5-\sqrt{5}}+22^{\frac{1}{5}}(\sqrt{5}-1))}{8}} \\ + c_3 e^{-2^{\frac{1}{5}}x} + c_4 e^{\frac{(i(\sqrt{5}+1)2^{\frac{7}{10}}\sqrt{5-\sqrt{5}}-22^{\frac{1}{5}}(\sqrt{5}-1))x}{8}} + c_5 e^{2^{\frac{1}{5}}(\cos(\frac{\pi}{5})+i\sin(\frac{\pi}{5}))x}$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 180

```
DSolve[y'''''[x]+2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-\frac{(\sqrt{5}-1)x}{2 \cdot 2^{4/5}}} \left(c_5 e^{\frac{(\sqrt{5}-5)x}{2 \cdot 2^{4/5}}} \right. \\ \left. + c_3 e^{\frac{\sqrt{5}x}{2^{4/5}}} \cos\left(\frac{\sqrt{5-\sqrt{5}x}}{2 \cdot 2^{3/10}}\right) + c_4 \cos\left(\frac{\sqrt{5+\sqrt{5}x}}{2 \cdot 2^{3/10}}\right) + c_2 e^{\frac{\sqrt{5}x}{2^{4/5}}} \sin\left(\frac{\sqrt{5-\sqrt{5}x}}{2 \cdot 2^{3/10}}\right) + c_1 \sin\left(\frac{\sqrt{5+\sqrt{5}x}}{2 \cdot 2^{3/10}}\right) \right)$$

9.5 problem 1(e)

9.5.1 Maple step by step solution 724

Internal problem ID [5986]

Internal file name [OUTPUT/5234_Sunday_June_05_2022_03_28_06_PM_17994693/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 83

Problem number: 1(e).

ODE order: 4.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_high_order , _missing_x]]
```

$$y'''' - 5y'' + 4y = 0$$

The characteristic equation is

$$\lambda^4 - 5\lambda^2 + 4 = 0$$

The roots of the above equation are

$$\lambda_1 = 2$$

$$\lambda_2 = -2$$

$$\lambda_3 = 1$$

$$\lambda_4 = -1$$

Therefore the homogeneous solution is

$$y_h(x) = c_1e^{-x} + c_2e^{-2x} + c_3e^x + e^{2x}c_4$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = e^{-x}$$

$$y_2 = e^{-2x}$$

$$y_3 = e^x$$

$$y_4 = e^{2x}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-x} + c_2 e^{-2x} + c_3 e^x + e^{2x} c_4 \quad (1)$$

Verification of solutions

$$y = c_1 e^{-x} + c_2 e^{-2x} + c_3 e^x + e^{2x} c_4$$

Verified OK.

9.5.1 Maple step by step solution

Let's solve

$$y'''' - 5y'' + 4y = 0$$

- Highest derivative means the order of the ODE is 4

$$y''''$$

- Convert linear ODE into a system of first order ODEs

- Define new variable $y_1(x)$

$$y_1(x) = y$$

- Define new variable $y_2(x)$

$$y_2(x) = y'$$

- Define new variable $y_3(x)$

$$y_3(x) = y''$$

- Define new variable $y_4(x)$

$$y_4(x) = y'''$$

- Isolate for $y_4'(x)$ using original ODE

$$y_4'(x) = 5y_3(x) - 4y_1(x)$$

Convert linear ODE into a system of first order ODEs

$$[y_2(x) = y_1'(x), y_3(x) = y_2'(x), y_4(x) = y_3'(x), y_4'(x) = 5y_3(x) - 4y_1(x)]$$

- Define vector

$$\vec{y}(x) = \begin{bmatrix} y_1(x) \\ y_2(x) \\ y_3(x) \\ y_4(x) \end{bmatrix}$$

- System to solve

$$\vec{y}'(x) = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -4 & 0 & 5 & 0 \end{bmatrix} \cdot \vec{y}(x)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -4 & 0 & 5 & 0 \end{bmatrix}$$

- Rewrite the system as

$$\vec{y}'(x) = A \cdot \vec{y}(x)$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[-2, \begin{bmatrix} -\frac{1}{8} \\ \frac{1}{4} \\ -\frac{1}{2} \\ 1 \end{bmatrix} \right], \left[-1, \begin{bmatrix} -1 \\ 1 \\ -1 \\ 1 \end{bmatrix} \right], \left[1, \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} \right], \left[2, \begin{bmatrix} \frac{1}{8} \\ \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[-2, \begin{bmatrix} -\frac{1}{8} \\ \frac{1}{4} \\ -\frac{1}{2} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_1 = e^{-2x} \cdot \begin{bmatrix} -\frac{1}{8} \\ \frac{1}{4} \\ -\frac{1}{2} \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[-1, \begin{bmatrix} -1 \\ 1 \\ -1 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_2 = e^{-x} \cdot \begin{bmatrix} -1 \\ 1 \\ -1 \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[1, \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_3 = e^x \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[2, \begin{bmatrix} \frac{1}{8} \\ \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_4 = e^{2x} \cdot \begin{bmatrix} \frac{1}{8} \\ \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{y} = c_1 \vec{y}_1 + c_2 \vec{y}_2 + c_3 \vec{y}_3 + c_4 \vec{y}_4$$

- Substitute solutions into the general solution

$$\vec{y} = c_1 e^{-2x} \cdot \begin{bmatrix} -\frac{1}{8} \\ \frac{1}{4} \\ -\frac{1}{2} \\ 1 \end{bmatrix} + c_2 e^{-x} \cdot \begin{bmatrix} -1 \\ 1 \\ -1 \\ 1 \end{bmatrix} + c_3 e^x \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} + e^{2x} c_4 \cdot \begin{bmatrix} \frac{1}{8} \\ \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y = -\frac{(-e^{4x}c_4 - 8c_3e^{3x} + 8c_2e^x + c_1)e^{-2x}}{8}$$

Maple trace

```

`Methods for high order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```


✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 27

```
dsolve(diff(y(x),x$4)-5*diff(y(x),x$2)+4*y(x)=0,y(x), singsol=all)
```

$$y(x) = (e^{4x}c_1 + c_4e^{3x} + e^xc_2 + c_3) e^{-2x}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 35

```
DSolve[y''''[x]-5*y''[x]+4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2x} (c_2e^x + e^{3x}(c_4e^x + c_3) + c_1)$$

9.6 problem 2

9.6.1 Maple step by step solution 731

Internal problem ID [5987]

Internal file name [OUTPUT/5235_Sunday_June_05_2022_03_28_07_PM_72595949/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 83

Problem number: 2.

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_3rd_order , _missing_x]]
```

$$y''' + y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 1, y''(0) = 0]$$

The characteristic equation is

$$\lambda^3 + 1 = 0$$

The roots of the above equation are

$$\begin{aligned}\lambda_1 &= -1 \\ \lambda_2 &= \frac{1}{2} - \frac{i\sqrt{3}}{2} \\ \lambda_3 &= \frac{1}{2} + \frac{i\sqrt{3}}{2}\end{aligned}$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 e^{-x} + e^{\left(\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)x} c_2 + e^{\left(\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)x} c_3$$

The fundamental set of solutions for the homogeneous solution are the following

$$\begin{aligned}y_1 &= e^{-x} \\y_2 &= e^{\left(\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)x} \\y_3 &= e^{\left(\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)x}\end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 e^{-x} + e^{\left(\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)x} c_2 + e^{\left(\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)x} c_3 \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_1 + c_2 + c_3 \quad (1A)$$

Taking derivative of the solution gives

$$y' = -c_1 e^{-x} + \left(\frac{1}{2} + \frac{i\sqrt{3}}{2}\right) e^{\left(\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)x} c_2 + \left(\frac{1}{2} - \frac{i\sqrt{3}}{2}\right) e^{\left(\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)x} c_3$$

substituting $y' = 1$ and $x = 0$ in the above gives

$$1 = \frac{i(c_2 - c_3)\sqrt{3}}{2} - c_1 + \frac{c_2}{2} + \frac{c_3}{2} \quad (2A)$$

Taking two derivatives of the solution gives

$$y'' = c_1 e^{-x} + \left(\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)^2 e^{\left(\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)x} c_2 + \left(\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)^2 e^{\left(\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)x} c_3$$

substituting $y'' = 0$ and $x = 0$ in the above gives

$$0 = \frac{i(c_2 - c_3)\sqrt{3}}{2} + c_1 - \frac{c_2}{2} - \frac{c_3}{2} \quad (3A)$$

Equations {1A,2A,3A} are now solved for $\{c_1, c_2, c_3\}$. Solving for the constants gives

$$\begin{aligned}c_1 &= -\frac{1}{3} \\c_2 &= -\frac{(-\sqrt{3} + 3i)\sqrt{3}}{18} \\c_3 &= \frac{\sqrt{3}(\sqrt{3} + 3i)}{18}\end{aligned}$$

Substituting these values back in above solution results in

$$y = -\frac{e^{-x}}{3} - \frac{i\sqrt{3}e^{\frac{(1+i\sqrt{3})x}{2}}}{6} + \frac{e^{\frac{(1+i\sqrt{3})x}{2}}}{6} + \frac{i\sqrt{3}e^{-\frac{(i\sqrt{3}-1)x}{2}}}{6} + \frac{e^{-\frac{(i\sqrt{3}-1)x}{2}}}{6}$$

Summary

The solution(s) found are the following

$$y = \frac{(1+i\sqrt{3})e^{-\frac{(i\sqrt{3}-1)x}{2}}}{6} - \frac{i\sqrt{3}e^{\frac{(1+i\sqrt{3})x}{2}}}{6} - \frac{e^{-x}}{3} + \frac{e^{\frac{(1+i\sqrt{3})x}{2}}}{6} \quad (1)$$

Verification of solutions

$$y = \frac{(1+i\sqrt{3})e^{-\frac{(i\sqrt{3}-1)x}{2}}}{6} - \frac{i\sqrt{3}e^{\frac{(1+i\sqrt{3})x}{2}}}{6} - \frac{e^{-x}}{3} + \frac{e^{\frac{(1+i\sqrt{3})x}{2}}}{6}$$

Verified OK.

9.6.1 Maple step by step solution

Let's solve

$$\left[y''' + y = 0, y(0) = 0, y'|_{\{x=0\}} = 1, y''|_{\{x=0\}} = 0 \right]$$

- Highest derivative means the order of the ODE is 3

y'''

- Convert linear ODE into a system of first order ODEs

- Define new variable $y_1(x)$

$$y_1(x) = y$$

- Define new variable $y_2(x)$

$$y_2(x) = y'$$

- Define new variable $y_3(x)$

$$y_3(x) = y''$$

- Isolate for $y'_3(x)$ using original ODE

$$y'_3(x) = -y_1(x)$$

Convert linear ODE into a system of first order ODEs

$$[y_2(x) = y'_1(x), y_3(x) = y'_2(x), y'_3(x) = -y_1(x)]$$

- Define vector

$$\vec{y}(x) = \begin{bmatrix} y_1(x) \\ y_2(x) \\ y_3(x) \end{bmatrix}$$

- System to solve

$$\vec{y}'(x) = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ -1 & 0 & 0 \end{bmatrix} \cdot \vec{y}(x)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ -1 & 0 & 0 \end{bmatrix}$$

- Rewrite the system as

$$\vec{y}'(x) = A \cdot \vec{y}(x)$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[-1, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right], \left[\frac{1}{2} - \frac{i\sqrt{3}}{2}, \begin{bmatrix} \frac{1}{\left(\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)^2} \\ \frac{1}{\frac{1}{2} - \frac{i\sqrt{3}}{2}} \\ 1 \end{bmatrix} \right], \left[\frac{1}{2} + \frac{i\sqrt{3}}{2}, \begin{bmatrix} \frac{1}{\left(\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)^2} \\ \frac{1}{\frac{1}{2} + \frac{i\sqrt{3}}{2}} \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[-1, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_1 = e^{-x} \cdot \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$\begin{bmatrix} \frac{1}{2} - \frac{i\sqrt{3}}{2}, & \begin{bmatrix} \frac{1}{\left(\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)^2} \\ \frac{1}{\frac{1}{2} - \frac{i\sqrt{3}}{2}} \\ 1 \end{bmatrix} \end{bmatrix}$$

- Solution from eigenpair

$$e^{\left(\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)x} \cdot \begin{bmatrix} \frac{1}{\left(\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)^2} \\ \frac{1}{\frac{1}{2} - \frac{i\sqrt{3}}{2}} \\ 1 \end{bmatrix}$$

- Use Euler identity to write solution in terms of sin and cos

$$e^{\frac{x}{2}} \cdot \left(\cos\left(\frac{\sqrt{3}x}{2}\right) - i \sin\left(\frac{\sqrt{3}x}{2}\right) \right) \cdot \begin{bmatrix} \frac{1}{\left(\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)^2} \\ \frac{1}{\frac{1}{2} - \frac{i\sqrt{3}}{2}} \\ 1 \end{bmatrix}$$

- Simplify expression

$$e^{\frac{x}{2}} \cdot \begin{bmatrix} \frac{\cos\left(\frac{\sqrt{3}x}{2}\right) - i \sin\left(\frac{\sqrt{3}x}{2}\right)}{\left(\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)^2} \\ \frac{\cos\left(\frac{\sqrt{3}x}{2}\right) - i \sin\left(\frac{\sqrt{3}x}{2}\right)}{\frac{1}{2} - \frac{i\sqrt{3}}{2}} \\ \cos\left(\frac{\sqrt{3}x}{2}\right) - i \sin\left(\frac{\sqrt{3}x}{2}\right) \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\vec{y}_2(x) = e^{\frac{x}{2}} \cdot \begin{bmatrix} -\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} \\ \frac{\cos\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} \\ \cos\left(\frac{\sqrt{3}x}{2}\right) \end{bmatrix}, \vec{y}_3(x) = e^{\frac{x}{2}} \cdot \begin{bmatrix} \frac{\cos\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)}{2} \\ \frac{\cos\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} - \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)}{2} \\ -\sin\left(\frac{\sqrt{3}x}{2}\right) \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{y} = c_1 \vec{y}_1 + c_2 \vec{y}_2(x) + c_3 \vec{y}_3(x)$$

- Substitute solutions into the general solution

$$\vec{y} = c_1 e^{-x} \cdot \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} + c_2 e^{\frac{x}{2}} \cdot \begin{bmatrix} -\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} \\ \frac{\cos\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} \\ \cos\left(\frac{\sqrt{3}x}{2}\right) \end{bmatrix} + c_3 e^{\frac{x}{2}} \cdot \begin{bmatrix} \frac{\cos\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)}{2} \\ \frac{\cos\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} - \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)}{2} \\ -\sin\left(\frac{\sqrt{3}x}{2}\right) \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y = \left(-\frac{e^{\frac{3x}{2}}(-c_3\sqrt{3}+c_2)\cos\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{e^{\frac{3x}{2}}(\sqrt{3}c_2+c_3)\sin\left(\frac{\sqrt{3}x}{2}\right)}{2} + c_1 \right) e^{-x}$$

- Use the initial condition $y(0) = 0$

$$0 = \frac{c_3\sqrt{3}}{2} - \frac{c_2}{2} + c_1$$

- Calculate the 1st derivative of the solution

$$y' = \left(-\frac{3e^{\frac{3x}{2}}(-c_3\sqrt{3}+c_2)\cos\left(\frac{\sqrt{3}x}{2}\right)}{4} + \frac{e^{\frac{3x}{2}}(-c_3\sqrt{3}+c_2)\sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{4} + \frac{3e^{\frac{3x}{2}}(\sqrt{3}c_2+c_3)\sin\left(\frac{\sqrt{3}x}{2}\right)}{4} + \frac{e^{\frac{3x}{2}}(\sqrt{3}c_2+c_3)\cos\left(\frac{\sqrt{3}x}{2}\right)}{4} - c_1 \right) e^{-x}$$

- Use the initial condition $y'|_{\{x=0\}} = 1$

$$1 = \frac{c_3\sqrt{3}}{4} - \frac{c_2}{4} + \frac{(\sqrt{3}c_2+c_3)\sqrt{3}}{4} - c_1$$

- Calculate the 2nd derivative of the solution

$$y'' = \left(-\frac{3e^{\frac{3x}{2}}(-c_3\sqrt{3}+c_2)\cos\left(\frac{\sqrt{3}x}{2}\right)}{4} + \frac{3e^{\frac{3x}{2}}(-c_3\sqrt{3}+c_2)\sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{4} + \frac{3e^{\frac{3x}{2}}(\sqrt{3}c_2+c_3)\sin\left(\frac{\sqrt{3}x}{2}\right)}{4} + \frac{3e^{\frac{3x}{2}}(\sqrt{3}c_2+c_3)\cos\left(\frac{\sqrt{3}x}{2}\right)}{4} - c_1 \right) e^{-x}$$

- Use the initial condition $y''|_{\{x=0\}} = 0$

$$0 = -\frac{c_3\sqrt{3}}{4} + \frac{c_2}{4} + \frac{(\sqrt{3}c_2+c_3)\sqrt{3}}{4} + c_1$$

- Solve for the unknown coefficients

$$\left\{ c_1 = -\frac{1}{3}, c_2 = \frac{1}{3}, c_3 = \frac{\sqrt{3}}{3} \right\}$$

- Solution to the IVP

$$y = \frac{e^{-x}\left(\sqrt{3}e^{\frac{3x}{2}}\sin\left(\frac{\sqrt{3}x}{2}\right) + e^{\frac{3x}{2}}\cos\left(\frac{\sqrt{3}x}{2}\right) - 1\right)}{3}$$

Maple trace

```
`Methods for third order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 39

```
dsolve([diff(y(x),x$3)+y(x)=0,y(0) = 0, D(y)(0) = 1, (D@@2)(y)(0) = 0],y(x), singsol=all)
```

$$y(x) = \frac{\left(e^{\frac{3x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) \sqrt{3} + e^{\frac{3x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) - 1 \right) e^{-x}}{3}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 59

```
DSolve[{y'''[x]+y[x]==0,{y[0]==0,y'[0]==1,y''[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{3} e^{-x} \left(\sqrt{3} e^{3x/2} \sin\left(\frac{\sqrt{3}x}{2}\right) + e^{3x/2} \cos\left(\frac{\sqrt{3}x}{2}\right) - 1 \right)$$

9.7 problem 3(a)

Internal problem ID [5988]

Internal file name [OUTPUT/5236_Sunday_June_05_2022_03_28_09_PM_26492945/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 83

Problem number: 3(a).

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_3rd_order , _missing_x]]
```

$$y''' - iy'' + y' - iy = 0$$

The characteristic equation is

$$\lambda^3 - i\lambda^2 + \lambda - i = 0$$

The roots of the above equation are

$$\lambda_1 = -i$$

$$\lambda_2 = i$$

$$\lambda_3 = i$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 e^{-ix} + c_2 e^{ix} + x e^{ix} c_3$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = e^{-ix}$$

$$y_2 = e^{ix}$$

$$y_3 = x e^{ix}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-ix} + c_2 e^{ix} + x e^{ix} c_3 \quad (1)$$

Verification of solutions

$$y = c_1 e^{-ix} + c_2 e^{ix} + x e^{ix} c_3$$

Verified OK.

Maple trace

```
`Methods for third order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(diff(y(x),x$3)-I*diff(y(x),x$2)+diff(y(x),x)-I*y(x)=0,y(x), singsol=all)
```

$$y(x) = (c_3 x + c_2) e^{ix} + e^{-ix} c_1$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 31

```
DSolve[y'''[x]-I*y''[x]+y'[x]-I*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-ix} (e^{2ix} (c_3 x + c_2) + c_1)$$

9.8 problem 3(b)

9.8.1	Solving as second order linear constant coeff ode	738
9.8.2	Solving as linear second order ode solved by an integrating factor ode	740
9.8.3	Solving using Kovacic algorithm	741
9.8.4	Maple step by step solution	745

Internal problem ID [5989]

Internal file name [OUTPUT/5237_Sunday_June_05_2022_03_28_10_PM_90187819/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 83

Problem number: 3(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "linear_second_order_ode_solved_by_an_integrating_factor"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' - 2iy' - y = 0$$

9.8.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -2i, C = -1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 2i\lambda e^{\lambda x} - e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 2i\lambda - 1 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -2i, C = -1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{2i}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{(-2i)^2 - (4)(1)(-1)} \\ &= i \end{aligned}$$

Hence this is the case of a double root $\lambda_{1,2} = -i$. Therefore the solution is

$$y = c_1 e^{ix} + c_2 x e^{ix} \quad (1)$$

Summary

The solution(s) found are the following

$$y = c_1 e^{ix} + e^{ix} c_2 x \quad (1)$$




Figure 137: Slope field plot

Verification of solutions

$$y = c_1 e^{ix} + e^{ix} c_2 x$$

Verified OK.

9.8.2 Solving as linear second order ode solved by an integrating factor ode

The ode satisfies this form

$$y'' + p(x)y' + \frac{(p(x)^2 + p'(x))y}{2} = f(x)$$

Where $p(x) = -2i$. Therefore, there is an integrating factor given by

$$\begin{aligned}M(x) &= e^{\frac{1}{2} \int p \, dx} \\ &= e^{\int -2i \, dx} \\ &= e^{-ix}\end{aligned}$$

Multiplying both sides of the ODE by the integrating factor $M(x)$ makes the left side of the ODE a complete differential

$$\begin{aligned}(M(x)y)'' &= 0 \\ (ye^{-ix})'' &= 0\end{aligned}$$

Integrating once gives

$$(ye^{-ix})' = c_1$$

Integrating again gives

$$(ye^{-ix}) = c_1x + c_2$$

Hence the solution is

$$y = \frac{c_1x + c_2}{e^{-ix}}$$

Or

$$y = c_1x e^{ix} + c_2 e^{ix}$$

Summary

The solution(s) found are the following

$$y = c_1x e^{ix} + c_2 e^{ix} \tag{1}$$




Figure 138: Slope field plot

Verification of solutions

$$y = c_1 x e^{ix} + c_2 e^{ix}$$

Verified OK.

9.8.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 2iy' - y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = -2i \tag{3}$$

$$C = -1$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 0 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 0 \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 143: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-2i}{1} dx} \\ &= z_1 e^{ix} \\ &= z_1 (e^{ix}) \end{aligned}$$

Which simplifies to

$$y_1 = e^{ix}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-2i}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{2ix}}{(y_1)^2} dx \\ &= y_1(x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{ix}) + c_2 (e^{ix}(x))\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{ix} + e^{ix} c_2 x \tag{1}$$




Figure 139: Slope field plot

Verification of solutions

$$y = c_1 e^{ix} + e^{ix} c_2 x$$

Verified OK.

9.8.4 Maple step by step solution

Let's solve

$$y'' - 2Iy' - y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 - 2Ir - 1 = 0$$

- Factor the characteristic polynomial

$$(-r + I)^2 = 0$$

- Root of the characteristic polynomial

$$r = I$$

- 1st solution of the ODE

$$y_1(x) = e^{Ix}$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = x e^{Ix}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = c_1 e^{Ix} + e^{Ix} c_2 x$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
dsolve(diff(y(x),x$2)-2*I*diff(y(x),x)-y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{ix}(c_2x + c_1)$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 20

```
DSolve[y''[x]-2*I*y'[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{ix}(c_2x + c_1)$$

9.9 problem 5(b)

Internal problem ID [5990]

Internal file name [OUTPUT/5238_Sunday_June_05_2022_03_28_11_PM_31224304/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 83

Problem number: 5(b).

ODE order: 4.

ODE degree: 1.

The type(s) of ODE detected by this program : "**unknown**"

Maple gives the following as the ode type

```
[[_high_order , _missing_x]]
```

Unable to solve or complete the solution.

$$y'''' - k^4 y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 0, y(1) = 0, y'(1) = 0]$$

The characteristic equation is

$$-k^4 + \lambda^4 = 0$$

The roots of the above equation are

$$\lambda_1 = k$$

$$\lambda_2 = -k$$

$$\lambda_3 = ik$$

$$\lambda_4 = -ik$$

Therefore the homogeneous solution is

$$y_h(x) = e^{ikx} c_1 + e^{kx} c_2 + e^{-ikx} c_3 + e^{-kx} c_4$$

The fundamental set of solutions for the homogeneous solution are the following

$$\begin{aligned}y_1 &= e^{ikx} \\y_2 &= e^{kx} \\y_3 &= e^{-ikx} \\y_4 &= e^{-kx}\end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = e^{ikx}c_1 + e^{kx}c_2 + e^{-ikx}c_3 + e^{-kx}c_4 \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 1$ in the above gives

$$0 = e^{ik}c_1 + e^k c_2 + e^{-ik}c_3 + e^{-k}c_4 \quad (1A)$$

substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_1 + c_2 + c_3 + c_4 \quad (2A)$$

Taking derivative of the solution gives

$$y' = ik e^{ikx}c_1 + k e^{kx}c_2 - ik e^{-ikx}c_3 - k e^{-kx}c_4$$

substituting $y' = 0$ and $x = 1$ in the above gives

$$0 = (ie^{ik}c_1 + e^k c_2 - ie^{-ik}c_3 - e^{-k}c_4) k \quad (3A)$$

Taking derivative of the solution gives

$$y' = ik e^{ikx}c_1 + k e^{kx}c_2 - ik e^{-ikx}c_3 - k e^{-kx}c_4$$

substituting $y' = 0$ and $x = 0$ in the above gives

$$0 = (c_1i - c_3i + c_2 - c_4) k \quad (4A)$$

Equations {1A,2A,3A,4A} are now solved for $\{c_1, c_2, c_3, c_4\}$. Solving for the constants gives

$$\begin{aligned}c_1 &= 0 \\c_2 &= 0 \\c_3 &= 0 \\c_4 &= 0\end{aligned}$$

Substituting these values back in above solution results in

$$y = 0$$

Summary

The solution(s) found are the following

$$y = 0 \tag{1}$$




Figure 140: Solution plot

Verification of solutions

$$y = 0$$

Verified OK.

Maple trace

```
`Methods for high order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.094 (sec). Leaf size: 5

```
dsolve([diff(y(x),x$4)-k^4*y(x)=0,y(0) = 0, D(y)(0) = 0, y(1) = 0, D(y)(1) = 0],y(x), singso
```

$$y(x) = 0$$

✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 6

```
DSolve[{y''''[x]-k^4*y[x]==0,{y[0]==0,y[1]==0,y'[0]==0,y'[1]==0}},y[x],x,IncludeSingularSolu
```

$$y(x) \rightarrow 0$$

10 Chapter 2. Linear equations with constant coefficients. Page 89

10.1	problem 1(a)	752
10.2	problem 1(b)	760
10.3	problem 1(c)	771
10.4	problem 1(d)	781
10.5	problem 1(e)	785
10.6	problem 1(f)	796

10.1 problem 1(a)

10.1.1 Maple step by step solution 754

Internal problem ID [5991]

Internal file name [OUTPUT/5239_Sunday_June_05_2022_03_28_13_PM_86747064/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 89

Problem number: 1(a).

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_3rd_order , _with_linear_symmetries]]
```

$$y''' - y = x$$

This is higher order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE And y_p is a particular solution to the nonhomogeneous ODE. y_h is the solution to

$$y''' - y = 0$$

The characteristic equation is

$$\lambda^3 - 1 = 0$$

The roots of the above equation are

$$\lambda_1 = 1$$

$$\lambda_2 = -\frac{1}{2} - \frac{i\sqrt{3}}{2}$$

$$\lambda_3 = -\frac{1}{2} + \frac{i\sqrt{3}}{2}$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 e^x + e^{\left(-\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)x} c_2 + e^{\left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)x} c_3$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = e^x$$

$$y_2 = e^{\left(-\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)x}$$

$$y_3 = e^{\left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)x}$$

Now the particular solution to the given ODE is found

$$y''' - y = x$$

The particular solution is found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^x, e^{\left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)x}, e^{\left(-\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)x} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_2 x + A_1$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-A_2 x - A_1 = x$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 0, A_2 = -1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 e^x + e^{\left(-\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)x} c_2 + e^{\left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)x} c_3 \right) + (-x) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^x + e^{\left(-\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)x} c_2 + e^{\left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)x} c_3 - x \quad (1)$$

Verification of solutions

$$y = c_1 e^x + e^{\left(-\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)x} c_2 + e^{\left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)x} c_3 - x$$

Verified OK.

10.1.1 Maple step by step solution

Let's solve

$$y''' - y = x$$

- Highest derivative means the order of the ODE is 3

$$y'''$$

- Convert linear ODE into a system of first order ODEs

- Define new variable $y_1(x)$

$$y_1(x) = y$$

- Define new variable $y_2(x)$

$$y_2(x) = y'$$

- Define new variable $y_3(x)$

$$y_3(x) = y''$$

- Isolate for $y_3'(x)$ using original ODE

$$y_3'(x) = x + y_1(x)$$

Convert linear ODE into a system of first order ODEs

$$[y_2(x) = y_1'(x), y_3(x) = y_2'(x), y_3'(x) = x + y_1(x)]$$

- Define vector

$$\vec{y}(x) = \begin{bmatrix} y_1(x) \\ y_2(x) \\ y_3(x) \end{bmatrix}$$

- System to solve

$$\vec{y}'(x) = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix} \cdot \vec{y}(x) + \begin{bmatrix} 0 \\ 0 \\ x \end{bmatrix}$$

- Define the forcing function

$$\vec{f}(x) = \begin{bmatrix} 0 \\ 0 \\ x \end{bmatrix}$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}$$

- Rewrite the system as

$$\vec{y}'(x) = A \cdot \vec{y}(x) + \vec{f}$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right], \left[-\frac{1}{2} - \frac{i\sqrt{3}}{2}, \begin{bmatrix} \frac{1}{\left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)^2} \\ \frac{1}{-\frac{1}{2} - \frac{i\sqrt{3}}{2}} \\ 1 \end{bmatrix} \right], \left[-\frac{1}{2} + \frac{i\sqrt{3}}{2}, \begin{bmatrix} \frac{1}{\left(-\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)^2} \\ \frac{1}{-\frac{1}{2} + \frac{i\sqrt{3}}{2}} \\ 1 \end{bmatrix} \right]$$

- Consider eigenpair

$$\left[1, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_1 = e^x \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$\left[-\frac{1}{2} - \frac{i\sqrt{3}}{2}, \begin{bmatrix} \frac{1}{\left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)^2} \\ \frac{1}{-\frac{1}{2} - \frac{i\sqrt{3}}{2}} \\ 1 \end{bmatrix} \right]$$

- Solution from eigenpair

$$e^{\left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)x} \cdot \begin{bmatrix} \frac{1}{\left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)^2} \\ \frac{1}{-\frac{1}{2} - \frac{i\sqrt{3}}{2}} \\ 1 \end{bmatrix}$$

- Use Euler identity to write solution in terms of sin and cos

$$e^{-\frac{x}{2}} \cdot \left(\cos\left(\frac{\sqrt{3}x}{2}\right) - i \sin\left(\frac{\sqrt{3}x}{2}\right) \right) \cdot \begin{bmatrix} \frac{1}{\left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)^2} \\ \frac{1}{-\frac{1}{2} - \frac{i\sqrt{3}}{2}} \\ 1 \end{bmatrix}$$

- Simplify expression

$$e^{-\frac{x}{2}} \cdot \begin{bmatrix} \frac{\cos\left(\frac{\sqrt{3}x}{2}\right) - i \sin\left(\frac{\sqrt{3}x}{2}\right)}{\left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)^2} \\ \frac{\cos\left(\frac{\sqrt{3}x}{2}\right) - i \sin\left(\frac{\sqrt{3}x}{2}\right)}{-\frac{1}{2} - \frac{i\sqrt{3}}{2}} \\ \cos\left(\frac{\sqrt{3}x}{2}\right) - i \sin\left(\frac{\sqrt{3}x}{2}\right) \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\vec{y}_2(x) = e^{-\frac{x}{2}} \cdot \begin{bmatrix} -\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)}{2} - \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} \\ -\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} \\ \cos\left(\frac{\sqrt{3}x}{2}\right) \end{bmatrix}, \vec{y}_3(x) = e^{-\frac{x}{2}} \cdot \begin{bmatrix} -\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)}{2} \\ \frac{\cos\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)}{2} \\ -\sin\left(\frac{\sqrt{3}x}{2}\right) \end{bmatrix}$$

- General solution of the system of ODEs can be written in terms of the particular solution $\vec{y}_p(x)$
 $\vec{y}(x) = c_1\vec{y}_1 + c_2\vec{y}_2(x) + c_3\vec{y}_3(x) + \vec{y}_p(x)$

□ Fundamental matrix

- Let $\phi(x)$ be the matrix whose columns are the independent solutions of the homogeneous system

$$\phi(x) = \begin{bmatrix} e^x & e^{-\frac{x}{2}} \left(-\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)}{2} - \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} \right) & e^{-\frac{x}{2}} \left(-\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)}{2} \right) \\ e^x & e^{-\frac{x}{2}} \left(-\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} \right) & e^{-\frac{x}{2}} \left(\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)}{2} \right) \\ e^x & e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) & -e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) \end{bmatrix}$$

- The fundamental matrix, $\Phi(x)$ is a normalized version of $\phi(x)$ satisfying $\Phi(0) = I$ where I is the identity matrix
 $\Phi(x) = \phi(x) \cdot \frac{1}{\phi(0)}$

- Substitute the value of $\phi(x)$ and $\phi(0)$

$$\Phi(x) = \begin{bmatrix} e^x & e^{-\frac{x}{2}} \left(-\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)}{2} - \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} \right) & e^{-\frac{x}{2}} \left(-\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)}{2} \right) \\ e^x & e^{-\frac{x}{2}} \left(-\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} \right) & e^{-\frac{x}{2}} \left(\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)}{2} \right) \\ e^x & e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) & -e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) \end{bmatrix} \cdot \begin{bmatrix} 1 & & \\ 1 & -\frac{1}{2} & -\frac{\sqrt{3}}{2} \\ 1 & -\frac{1}{2} & \frac{\sqrt{3}}{2} \\ 1 & 1 & 0 \end{bmatrix}$$

- Evaluate and simplify to get the fundamental matrix

$$\Phi(x) = \begin{bmatrix} \frac{e^x}{3} + \frac{2e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} & \frac{e^x}{3} - \frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} + \frac{e^{-\frac{x}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3} & \frac{e^x}{3} - \frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} \\ \frac{e^x}{3} - \frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} - \frac{e^{-\frac{x}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3} & \frac{e^x}{3} + \frac{2e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} & \frac{e^x}{3} - \frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} \\ \frac{e^x}{3} - \frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} + \frac{e^{-\frac{x}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3} & \frac{e^x}{3} - \frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} - \frac{e^{-\frac{x}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3} & \frac{e^x}{3} + \frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} \end{bmatrix}$$

□ Find a particular solution of the system of ODEs using variation of parameters

- Let the particular solution be the fundamental matrix multiplied by $\vec{v}(x)$ and solve for $\vec{v}(x)$

$$\vec{y}_p(x) = \Phi(x) \cdot \vec{v}(x)$$

- Take the derivative of the particular solution

$$\vec{y}'_p(x) = \Phi'(x) \cdot \vec{v}(x) + \Phi(x) \cdot \vec{v}'(x)$$

- Substitute particular solution and its derivative into the system of ODEs

$$\Phi'(x) \cdot \vec{v}(x) + \Phi(x) \cdot \vec{v}'(x) = A \cdot \Phi(x) \cdot \vec{v}(x) + \vec{f}(x)$$

- The fundamental matrix has columns that are solutions to the homogeneous system so its derivative is

$$A \cdot \Phi(x) \cdot \vec{v}(x) + \Phi(x) \cdot \vec{v}'(x) = A \cdot \Phi(x) \cdot \vec{v}(x) + \vec{f}(x)$$

- Cancel like terms

$$\Phi(x) \cdot \vec{v}'(x) = \vec{f}(x)$$

- Multiply by the inverse of the fundamental matrix

$$\vec{v}'(x) = \frac{1}{\Phi(x)} \cdot \vec{f}(x)$$

- Integrate to solve for $\vec{v}(x)$

$$\vec{v}(x) = \int_0^x \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds$$

- Plug $\vec{v}(x)$ into the equation for the particular solution

$$\vec{y}_p(x) = \Phi(x) \cdot \left(\int_0^x \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds \right)$$

- Plug in the fundamental matrix and the forcing function and compute

$$\vec{y}_p(x) = \begin{bmatrix} \frac{e^{-\frac{x}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3} - x + \frac{e^x}{3} - \frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} \\ \frac{e^x}{3} + \frac{2e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} - 1 \\ -\frac{e^{-\frac{x}{2}} \left(-e^{\frac{3x}{2}} + \sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3} + \cos\left(\frac{\sqrt{3}x}{2}\right)\right)}{3} \end{bmatrix}$$

- Plug particular solution back into general solution

$$\vec{y}(x) = c_1 \vec{y}_1 + c_2 \vec{y}_2(x) + c_3 \vec{y}_3(x) + \begin{bmatrix} \frac{e^{-\frac{x}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3} - x + \frac{e^x}{3} - \frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} \\ \frac{e^x}{3} + \frac{2e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} - 1 \\ -\frac{e^{-\frac{x}{2}} \left(-e^{\frac{3x}{2}} + \sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3} + \cos\left(\frac{\sqrt{3}x}{2}\right)\right)}{3} \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y = -\frac{(c_3 \sqrt{3} + c_2 + \frac{2}{3})e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{2} - \frac{((c_2 - \frac{2}{3})\sqrt{3} - c_3)e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{(6c_1 + 2)e^x}{6} - x$$

Maple trace

```
`Methods for third order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 3; linear nonhomogeneous with symmetry [0,1]  
trying high order linear exact nonhomogeneous  
trying differential order: 3; missing the dependent variable  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 38

```
dsolve(diff(y(x),x$3)-y(x)=x,y(x), singsol=all)
```

$$y(x) = -x + e^x c_1 + c_2 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + c_3 e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 57

```
DSolve[y'''[x]-y[x]==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x + c_1 e^x + c_2 e^{-x/2} \cos\left(\frac{\sqrt{3}x}{2}\right) + c_3 e^{-x/2} \sin\left(\frac{\sqrt{3}x}{2}\right)$$

10.2 problem 1(b)

10.2.1 Maple step by step solution 765

Internal problem ID [5992]

Internal file name [OUTPUT/5240_Sunday_June_05_2022_03_28_15_PM_82884444/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 89

Problem number: 1(b).

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_3rd_order , _with_linear_symmetries]]
```

$$y''' - 8y = e^{ix}$$

This is higher order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE And y_p is a particular solution to the nonhomogeneous ODE. y_h is the solution to

$$y''' - 8y = 0$$

The characteristic equation is

$$\lambda^3 - 8 = 0$$

The roots of the above equation are

$$\lambda_1 = 2$$

$$\lambda_2 = i\sqrt{3} - 1$$

$$\lambda_3 = -i\sqrt{3} - 1$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 e^{2x} + e^{(i\sqrt{3}-1)x} c_2 + e^{(-i\sqrt{3}-1)x} c_3$$

The fundamental set of solutions for the homogeneous solution are the following

$$\begin{aligned} y_1 &= e^{2x} \\ y_2 &= e^{(i\sqrt{3}-1)x} \\ y_3 &= e^{(-i\sqrt{3}-1)x} \end{aligned}$$

Now the particular solution to the given ODE is found

$$y''' - 8y = e^{ix}$$

Let the particular solution be

$$y_p = U_1 y_1 + U_2 y_2 + U_3 y_3$$

Where y_i are the basis solutions found above for the homogeneous solution y_h and $U_i(x)$ are functions to be determined as follows

$$U_i = (-1)^{n-i} \int \frac{F(x)W_i(x)}{aW(x)} dx$$

Where $W(x)$ is the Wronskian and $W_i(x)$ is the Wronskian that results after deleting the last row and the i -th column of the determinant and n is the order of the ODE or equivalently, the number of basis solutions, and a is the coefficient of the leading derivative in the ODE, and $F(x)$ is the RHS of the ODE. Therefore, the first step is to find the Wronskian $W(x)$. This is given by

$$W(x) = \begin{vmatrix} y_1 & y_2 & y_3 \\ y_1' & y_2' & y_3' \\ y_1'' & y_2'' & y_3'' \end{vmatrix}$$

Substituting the fundamental set of solutions y_i found above in the Wronskian gives

$$\begin{aligned} W &= \begin{vmatrix} e^{2x} & e^{(i\sqrt{3}-1)x} & e^{-(1+i\sqrt{3})x} \\ 2e^{2x} & (i\sqrt{3}-1)e^{(i\sqrt{3}-1)x} & (-i\sqrt{3}-1)e^{-(1+i\sqrt{3})x} \\ 4e^{2x} & (i\sqrt{3}-1)^2 e^{(i\sqrt{3}-1)x} & (1+i\sqrt{3})^2 e^{-(1+i\sqrt{3})x} \end{vmatrix} \\ |W| &= -24ie^{2x}\sqrt{3}e^{-(1+i\sqrt{3})x}e^{(i\sqrt{3}-1)x} \end{aligned}$$

The determinant simplifies to

$$|W| = -24i\sqrt{3}$$

Now we determine W_i for each U_i .

$$\begin{aligned} W_1(x) &= \det \begin{bmatrix} e^{(i\sqrt{3}-1)x} & e^{-(1+i\sqrt{3})x} \\ (i\sqrt{3}-1)e^{(i\sqrt{3}-1)x} & (-i\sqrt{3}-1)e^{-(1+i\sqrt{3})x} \end{bmatrix} \\ &= -2i\sqrt{3}e^{-2x} \end{aligned}$$

$$\begin{aligned} W_2(x) &= \det \begin{bmatrix} e^{2x} & e^{-(1+i\sqrt{3})x} \\ 2e^{2x} & (-i\sqrt{3}-1)e^{-(1+i\sqrt{3})x} \end{bmatrix} \\ &= -e^{-(i\sqrt{3}-1)x}(i\sqrt{3}+3) \end{aligned}$$

$$\begin{aligned} W_3(x) &= \det \begin{bmatrix} e^{2x} & e^{(i\sqrt{3}-1)x} \\ 2e^{2x} & (i\sqrt{3}-1)e^{(i\sqrt{3}-1)x} \end{bmatrix} \\ &= e^{(1+i\sqrt{3})x}(i\sqrt{3}-3) \end{aligned}$$

Now we are ready to evaluate each $U_i(x)$.

$$\begin{aligned} U_1 &= (-1)^{3-1} \int \frac{F(x)W_1(x)}{aW(x)} dx \\ &= (-1)^2 \int \frac{(e^{ix})(-2i\sqrt{3}e^{-2x})}{(1)(-24i\sqrt{3})} dx \\ &= \int \frac{-2ie^{ix}\sqrt{3}e^{-2x}}{-24i\sqrt{3}} dx \\ &= \int \left(\frac{e^{(-2+i)x}}{12} \right) dx \\ &= \left(-\frac{1}{30} - \frac{i}{60} \right) e^{(-2+i)x} \end{aligned}$$

$$\begin{aligned}
U_2 &= (-1)^{3-2} \int \frac{F(x)W_2(x)}{aW(x)} dx \\
&= (-1)^1 \int \frac{(e^{ix}) \left(-e^{-(i\sqrt{3}-1)x} (i\sqrt{3} + 3) \right)}{(1) (-24i\sqrt{3})} dx \\
&= - \int \frac{-e^{ix} e^{-(i\sqrt{3}-1)x} (i\sqrt{3} + 3)}{-24i\sqrt{3}} dx \\
&= - \int \left(-\frac{(-\sqrt{3} + 3i) \sqrt{3} e^{(-i\sqrt{3}+1+i)x}}{72} \right) dx \\
&= \frac{(3i\sqrt{3} + 5 + i + 2\sqrt{3}) (-\sqrt{3} + 3i) \sqrt{3} e^{-x(i\sqrt{3}-1-i)}}{936} \\
&= \frac{(3i\sqrt{3} + 5 + i + 2\sqrt{3}) (-\sqrt{3} + 3i) \sqrt{3} e^{-x(i\sqrt{3}-1-i)}}{936}
\end{aligned}$$

$$\begin{aligned}
U_3 &= (-1)^{3-3} \int \frac{F(x)W_3(x)}{aW(x)} dx \\
&= (-1)^0 \int \frac{(e^{ix}) \left(e^{(1+i\sqrt{3})x} (i\sqrt{3} - 3) \right)}{(1) (-24i\sqrt{3})} dx \\
&= \int \frac{e^{ix} e^{(1+i\sqrt{3})x} (i\sqrt{3} - 3)}{-24i\sqrt{3}} dx \\
&= \int \left(-\frac{\sqrt{3} e^{x(i\sqrt{3}+1+i)} (\sqrt{3} + 3i)}{72} \right) dx \\
&= \frac{(3i\sqrt{3} + 2\sqrt{3} - 5 - i) \sqrt{3} e^{x(i\sqrt{3}+1+i)} (\sqrt{3} + 3i)}{936}
\end{aligned}$$

Now that all the U_i functions have been determined, the particular solution is found from

$$y_p = U_1 y_1 + U_2 y_2 + U_3 y_3$$

Hence

$$\begin{aligned}
 y_p &= \left(\left(-\frac{1}{30} - \frac{i}{60} \right) e^{(-2+i)x} \right) (e^{2x}) \\
 &+ \left(\frac{(3i\sqrt{3} + 5 + i + 2\sqrt{3})(-\sqrt{3} + 3i)\sqrt{3}e^{-x(i\sqrt{3}-1-i)}}{936} \right) \left(e^{(i\sqrt{3}-1)x} \right) \\
 &+ \left(\frac{(3i\sqrt{3} + 2\sqrt{3} - 5 - i)\sqrt{3}e^{x(i\sqrt{3}+1+i)}(\sqrt{3} + 3i)}{936} \right) \left(e^{(-i\sqrt{3}-1)x} \right)
 \end{aligned}$$

Therefore the particular solution is

$$y_p = \left(-\frac{8}{65} + \frac{i}{65} \right) e^{ix}$$

Which simplifies to

$$y_p = \left(-\frac{8}{65} + \frac{i}{65} \right) \cos(x) + \left(-\frac{1}{65} - \frac{8i}{65} \right) \sin(x)$$

Therefore the general solution is

$$\begin{aligned}
 y &= y_h + y_p \\
 &= \left(c_1 e^{2x} + e^{(i\sqrt{3}-1)x} c_2 + e^{(-i\sqrt{3}-1)x} c_3 \right) + \left(\left(-\frac{8}{65} + \frac{i}{65} \right) \cos(x) + \left(-\frac{1}{65} - \frac{8i}{65} \right) \sin(x) \right)
 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{2x} + e^{(i\sqrt{3}-1)x} c_2 + e^{(-i\sqrt{3}-1)x} c_3 + \left(-\frac{8}{65} + \frac{i}{65} \right) \cos(x) + \left(-\frac{1}{65} - \frac{8i}{65} \right) \sin(x)$$

Verification of solutions

$$y = c_1 e^{2x} + e^{(i\sqrt{3}-1)x} c_2 + e^{(-i\sqrt{3}-1)x} c_3 + \left(-\frac{8}{65} + \frac{i}{65} \right) \cos(x) + \left(-\frac{1}{65} - \frac{8i}{65} \right) \sin(x)$$

Verified OK.

10.2.1 Maple step by step solution

Let's solve

$$y''' - 8y = e^{Ix}$$

- Highest derivative means the order of the ODE is 3

$$y'''$$

- Convert linear ODE into a system of first order ODEs

- Define new variable $y_1(x)$

$$y_1(x) = y$$

- Define new variable $y_2(x)$

$$y_2(x) = y'$$

- Define new variable $y_3(x)$

$$y_3(x) = y''$$

- Isolate for $y_3'(x)$ using original ODE

$$y_3'(x) = e^{Ix} + 8y_1(x)$$

Convert linear ODE into a system of first order ODEs

$$[y_2(x) = y_1'(x), y_3(x) = y_2'(x), y_3'(x) = e^{Ix} + 8y_1(x)]$$

- Define vector

$$\vec{y}(x) = \begin{bmatrix} y_1(x) \\ y_2(x) \\ y_3(x) \end{bmatrix}$$

- System to solve

$$\vec{y}'(x) = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 8 & 0 & 0 \end{bmatrix} \cdot \vec{y}(x) + \begin{bmatrix} 0 \\ 0 \\ e^{Ix} \end{bmatrix}$$

- Define the forcing function

$$\vec{f}(x) = \begin{bmatrix} 0 \\ 0 \\ e^{Ix} \end{bmatrix}$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 8 & 0 & 0 \end{bmatrix}$$

- Rewrite the system as

$$\vec{y}'(x) = A \cdot \vec{y}(x) + \vec{f}$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[\left[2, \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right], \left[-I\sqrt{3} - 1, \begin{bmatrix} \frac{1}{(-I\sqrt{3}-1)^2} \\ \frac{1}{-I\sqrt{3}-1} \\ 1 \end{bmatrix} \right], \left[I\sqrt{3} - 1, \begin{bmatrix} \frac{1}{(I\sqrt{3}-1)^2} \\ \frac{1}{I\sqrt{3}-1} \\ 1 \end{bmatrix} \right] \right] \right]$$

- Consider eigenpair

$$\left[\left[\begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right] \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_1 = e^{2x} \cdot \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix}$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$\left[\left[-I\sqrt{3} - 1, \begin{bmatrix} \frac{1}{(-I\sqrt{3}-1)^2} \\ \frac{1}{-I\sqrt{3}-1} \\ 1 \end{bmatrix} \right] \right]$$

- Solution from eigenpair

$$e^{(-I\sqrt{3}-1)x} \cdot \begin{bmatrix} \frac{1}{(-I\sqrt{3}-1)^2} \\ \frac{1}{-I\sqrt{3}-1} \\ 1 \end{bmatrix}$$

- Use Euler identity to write solution in terms of sin and cos

$$e^{-x} \cdot (\cos(\sqrt{3}x) - I \sin(\sqrt{3}x)) \cdot \begin{bmatrix} \frac{1}{(-I\sqrt{3}-1)^2} \\ \frac{1}{-I\sqrt{3}-1} \\ 1 \end{bmatrix}$$

- Simplify expression

$$e^{-x} \cdot \begin{bmatrix} \frac{\cos(\sqrt{3}x) - I \sin(\sqrt{3}x)}{(-I\sqrt{3}-1)^2} \\ \frac{\cos(\sqrt{3}x) - I \sin(\sqrt{3}x)}{-I\sqrt{3}-1} \\ \cos(\sqrt{3}x) - I \sin(\sqrt{3}x) \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\begin{bmatrix} \vec{y}_2(x) = e^{-x} \cdot \begin{bmatrix} -\frac{\cos(\sqrt{3}x)}{8} - \frac{\sqrt{3} \sin(\sqrt{3}x)}{8} \\ -\frac{\cos(\sqrt{3}x)}{4} + \frac{\sqrt{3} \sin(\sqrt{3}x)}{4} \\ \cos(\sqrt{3}x) \end{bmatrix}, \vec{y}_3(x) = e^{-x} \cdot \begin{bmatrix} -\frac{\sqrt{3} \cos(\sqrt{3}x)}{8} + \frac{\sin(\sqrt{3}x)}{8} \\ \frac{\sqrt{3} \cos(\sqrt{3}x)}{4} + \frac{\sin(\sqrt{3}x)}{4} \\ -\sin(\sqrt{3}x) \end{bmatrix} \end{bmatrix}$$

- General solution of the system of ODEs can be written in terms of the particular solution $\vec{y}_p(x)$
 $\vec{y}(x) = c_1 \vec{y}_1 + c_2 \vec{y}_2(x) + c_3 \vec{y}_3(x) + \vec{y}_p(x)$

□ Fundamental matrix

- Let $\phi(x)$ be the matrix whose columns are the independent solutions of the homogeneous system

$$\phi(x) = \begin{bmatrix} \frac{e^{2x}}{4} & e^{-x} \left(-\frac{\cos(\sqrt{3}x)}{8} - \frac{\sqrt{3} \sin(\sqrt{3}x)}{8} \right) & e^{-x} \left(-\frac{\sqrt{3} \cos(\sqrt{3}x)}{8} + \frac{\sin(\sqrt{3}x)}{8} \right) \\ \frac{e^{2x}}{2} & e^{-x} \left(-\frac{\cos(\sqrt{3}x)}{4} + \frac{\sqrt{3} \sin(\sqrt{3}x)}{4} \right) & e^{-x} \left(\frac{\sqrt{3} \cos(\sqrt{3}x)}{4} + \frac{\sin(\sqrt{3}x)}{4} \right) \\ e^{2x} & e^{-x} \cos(\sqrt{3}x) & -e^{-x} \sin(\sqrt{3}x) \end{bmatrix}$$

- The fundamental matrix, $\Phi(x)$ is a normalized version of $\phi(x)$ satisfying $\Phi(0) = I$ where I is the identity matrix

$$\Phi(x) = \phi(x) \cdot \frac{1}{\phi(0)}$$

- Substitute the value of $\phi(x)$ and $\phi(0)$

$$\Phi(x) = \begin{bmatrix} \frac{e^{2x}}{4} & e^{-x} \left(-\frac{\cos(\sqrt{3}x)}{8} - \frac{\sqrt{3} \sin(\sqrt{3}x)}{8} \right) & e^{-x} \left(-\frac{\sqrt{3} \cos(\sqrt{3}x)}{8} + \frac{\sin(\sqrt{3}x)}{8} \right) \\ \frac{e^{2x}}{2} & e^{-x} \left(-\frac{\cos(\sqrt{3}x)}{4} + \frac{\sqrt{3} \sin(\sqrt{3}x)}{4} \right) & e^{-x} \left(\frac{\sqrt{3} \cos(\sqrt{3}x)}{4} + \frac{\sin(\sqrt{3}x)}{4} \right) \\ e^{2x} & e^{-x} \cos(\sqrt{3}x) & -e^{-x} \sin(\sqrt{3}x) \end{bmatrix} \cdot \frac{1}{\begin{bmatrix} \frac{1}{4} & -\frac{1}{8} & - \\ \frac{1}{2} & -\frac{1}{4} & \\ 1 & 1 & \end{bmatrix}}$$

- Evaluate and simplify to get the fundamental matrix

$$\Phi(x) = \begin{bmatrix} \frac{e^{2x}}{3} + \frac{2e^{-x} \cos(\sqrt{3}x)}{3} & \frac{e^{-x} \sin(\sqrt{3}x)\sqrt{3}}{6} - \frac{e^{-x} \cos(\sqrt{3}x)}{6} + \frac{e^{2x}}{6} & -\frac{e^{-x} \sin(\sqrt{3}x)}{6} \\ \frac{2e^{2x}}{3} - \frac{2e^{-x} \cos(\sqrt{3}x)}{3} - \frac{2e^{-x} \sin(\sqrt{3}x)\sqrt{3}}{3} & \frac{e^{2x}}{3} + \frac{2e^{-x} \cos(\sqrt{3}x)}{3} & \frac{e^{-x} \sin(\sqrt{3}x)}{6} \\ \frac{4e^{2x}}{3} - \frac{4e^{-x} \cos(\sqrt{3}x)}{3} + \frac{4e^{-x} \sin(\sqrt{3}x)\sqrt{3}}{3} & \frac{2e^{2x}}{3} - \frac{2e^{-x} \cos(\sqrt{3}x)}{3} - \frac{2e^{-x} \sin(\sqrt{3}x)\sqrt{3}}{3} & \frac{e^{-x} \sin(\sqrt{3}x)}{6} \end{bmatrix}$$

- Find a particular solution of the system of ODEs using variation of parameters

- Let the particular solution be the fundamental matrix multiplied by $\vec{v}(x)$ and solve for $\vec{v}(x)$

$$\vec{y}_p(x) = \Phi(x) \cdot \vec{v}(x)$$

- Take the derivative of the particular solution

$$\vec{y}'_p(x) = \Phi'(x) \cdot \vec{v}(x) + \Phi(x) \cdot \vec{v}'(x)$$

- Substitute particular solution and its derivative into the system of ODEs

$$\Phi'(x) \cdot \vec{v}(x) + \Phi(x) \cdot \vec{v}'(x) = A \cdot \Phi(x) \cdot \vec{v}(x) + \vec{f}(x)$$

- The fundamental matrix has columns that are solutions to the homogeneous system so its derivative is $A \cdot \Phi(x)$

$$A \cdot \Phi(x) \cdot \vec{v}(x) + \Phi(x) \cdot \vec{v}'(x) = A \cdot \Phi(x) \cdot \vec{v}(x) + \vec{f}(x)$$

- Cancel like terms

$$\Phi(x) \cdot \vec{v}'(x) = \vec{f}(x)$$

- Multiply by the inverse of the fundamental matrix

$$\vec{v}'(x) = \frac{1}{\Phi(x)} \cdot \vec{f}(x)$$

- Integrate to solve for $\vec{v}(x)$

$$\vec{v}(x) = \int_0^x \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds$$

- Plug $\vec{v}(x)$ into the equation for the particular solution

$$\vec{y}_p(x) = \Phi(x) \cdot \left(\int_0^x \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds \right)$$

- Plug in the fundamental matrix and the forcing function and compute

$$\vec{y}_p(x) = \begin{bmatrix} \left(\frac{7}{78} - \frac{51}{156}\right) e^{-x} \cos(\sqrt{3}x) + \left(\frac{1}{78} + \frac{1}{52}\right) e^{-x} \sqrt{3} \sin(\sqrt{3}x) + \left(-\frac{8}{65} + \frac{1}{65}\right) e^{Ix} + \left(\frac{1}{30} + \frac{1}{60}\right) e^{2x} \\ \left(-\frac{2}{39} + \frac{71}{78}\right) e^{-x} \cos(\sqrt{3}x) + \left(-\frac{4}{39} + \frac{1}{78}\right) e^{-x} \sqrt{3} \sin(\sqrt{3}x) - \left(\frac{1}{65} + \frac{81}{65}\right) e^{Ix} + \left(\frac{1}{15} + \frac{1}{30}\right) e^{2x} \\ -\left(\frac{10}{39} + \frac{21}{39}\right) e^{-x} \cos(\sqrt{3}x) + \left(\frac{2}{13} - \frac{41}{39}\right) e^{-x} \sqrt{3} \sin(\sqrt{3}x) + \left(\frac{8}{65} - \frac{1}{65}\right) e^{Ix} + \left(\frac{2}{15} + \frac{1}{15}\right) e^{2x} \end{bmatrix}$$

- Plug particular solution back into general solution

$$\vec{y}(x) = c_1 \vec{y}_1 + c_2 \vec{y}_2(x) + c_3 \vec{y}_3(x) + \begin{bmatrix} \left(\frac{7}{78} - \frac{51}{156}\right) e^{-x} \cos(\sqrt{3}x) + \left(\frac{1}{78} + \frac{1}{52}\right) e^{-x} \sqrt{3} \sin(\sqrt{3}x) \\ \left(-\frac{2}{39} + \frac{71}{78}\right) e^{-x} \cos(\sqrt{3}x) + \left(-\frac{4}{39} + \frac{1}{78}\right) e^{-x} \sqrt{3} \sin(\sqrt{3}x) \\ -\left(\frac{10}{39} + \frac{21}{39}\right) e^{-x} \cos(\sqrt{3}x) + \left(\frac{2}{13} - \frac{41}{39}\right) e^{-x} \sqrt{3} \sin(\sqrt{3}x) \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y = \frac{5\left(-\frac{39c_3\sqrt{3}}{10} + \frac{14}{5} - 1 - \frac{39c_2}{10}\right) e^{-x} \cos(\sqrt{3}x)}{156} + \frac{\left(\left(\frac{2}{3} + 1 - \frac{13c_2}{2}\right)\sqrt{3} + \frac{13c_3}{2}\right) e^{-x} \sin(\sqrt{3}x)}{52} + \left(-\frac{8}{65} + \frac{1}{65}\right) e^{Ix} + \frac{e^{2x}(2+1)}{60}$$

Maple trace

```
`Methods for third order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 3; linear nonhomogeneous with symmetry [0,1]
trying high order linear exact nonhomogeneous
trying differential order: 3; missing the dependent variable
checking if the LODE has constant coefficients
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 44

```
dsolve(diff(y(x),x$3)-8*y(x)=exp(I*x),y(x), singsol=all)
```

$$y(x) = \left(-\frac{8}{65} + \frac{i}{65}\right) e^{ix} + e^{2x} c_1 + c_2 e^{-x} \cos(\sqrt{3}x) + c_3 e^{-x} \sin(\sqrt{3}x)$$

✓ Solution by Mathematica

Time used: 0.472 (sec). Leaf size: 59

```
DSolve[y'''[x]-8*y[x]==Exp[I*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{65} e^{-x} \left(-(8-i)e^{(1+i)x} + 65c_1 e^{3x} + 65c_2 \cos(\sqrt{3}x) + 65c_3 \sin(\sqrt{3}x) \right)$$

10.3 problem 1(c)

10.3.1 Maple step by step solution 773

Internal problem ID [5993]

Internal file name [OUTPUT/5241_Sunday_June_05_2022_03_28_16_PM_76447078/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 89

Problem number: 1(c).

ODE order: 4.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_high_order , _linear , _nonhomogeneous]]
```

$$y'''' + 16y = \cos(x)$$

This is higher order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE And y_p is a particular solution to the nonhomogeneous ODE. y_h is the solution to

$$y'''' + 16y = 0$$

The characteristic equation is

$$\lambda^4 + 16 = 0$$

The roots of the above equation are

$$\lambda_1 = \sqrt{2} + i\sqrt{2}$$

$$\lambda_2 = -\sqrt{2} + i\sqrt{2}$$

$$\lambda_3 = -\sqrt{2} - i\sqrt{2}$$

$$\lambda_4 = -i\sqrt{2} + \sqrt{2}$$

Therefore the homogeneous solution is

$$y_h(x) = e^{(-i\sqrt{2}+\sqrt{2})x} c_1 + e^{(-\sqrt{2}+i\sqrt{2})x} c_2 + e^{(-\sqrt{2}-i\sqrt{2})x} c_3 + e^{(\sqrt{2}+i\sqrt{2})x} c_4$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = e^{(-i\sqrt{2}+\sqrt{2})x}$$

$$y_2 = e^{(-\sqrt{2}+i\sqrt{2})x}$$

$$y_3 = e^{(-\sqrt{2}-i\sqrt{2})x}$$

$$y_4 = e^{(\sqrt{2}+i\sqrt{2})x}$$

Now the particular solution to the given ODE is found

$$y'''' + 16y = \cos(x)$$

The particular solution is found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\cos(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{(-\sqrt{2}-i\sqrt{2})x}, e^{(-\sqrt{2}+i\sqrt{2})x}, e^{(\sqrt{2}+i\sqrt{2})x}, e^{(-i\sqrt{2}+\sqrt{2})x} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(x) + A_2 \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$17A_1 \cos(x) + 17A_2 \sin(x) = \cos(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{17}, A_2 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{\cos(x)}{17}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(e^{(-i\sqrt{2}+\sqrt{2})x} c_1 + e^{(-\sqrt{2}+i\sqrt{2})x} c_2 + e^{(-\sqrt{2}-i\sqrt{2})x} c_3 + e^{(\sqrt{2}+i\sqrt{2})x} c_4 \right) + \left(\frac{\cos(x)}{17} \right) \end{aligned}$$

Which simplifies to

$$y = e^{(1-i)\sqrt{2}x} c_1 + e^{(-1+i)\sqrt{2}x} c_2 + e^{(-1-i)\sqrt{2}x} c_3 + e^{(1+i)\sqrt{2}x} c_4 + \frac{\cos(x)}{17}$$

Summary

The solution(s) found are the following

$$y = e^{(1-i)\sqrt{2}x} c_1 + e^{(-1+i)\sqrt{2}x} c_2 + e^{(-1-i)\sqrt{2}x} c_3 + e^{(1+i)\sqrt{2}x} c_4 + \frac{\cos(x)}{17} \quad (1)$$

Verification of solutions

$$y = e^{(1-i)\sqrt{2}x} c_1 + e^{(-1+i)\sqrt{2}x} c_2 + e^{(-1-i)\sqrt{2}x} c_3 + e^{(1+i)\sqrt{2}x} c_4 + \frac{\cos(x)}{17}$$

Verified OK.

10.3.1 Maple step by step solution

Let's solve

$$y'''' + 16y = \cos(x)$$

- Highest derivative means the order of the ODE is 4

$$y''''$$

- Convert linear ODE into a system of first order ODEs

- Define new variable $y_1(x)$

$$y_1(x) = y$$

- Define new variable $y_2(x)$

$$y_2(x) = y'$$

- Define new variable $y_3(x)$

$$y_3(x) = y''$$

- Define new variable $y_4(x)$

$$y_4(x) = y'''$$

- Isolate for $y_4'(x)$ using original ODE

$$y_4'(x) = \cos(x) - 16y_1(x)$$

Convert linear ODE into a system of first order ODEs

$$[y_2(x) = y_1'(x), y_3(x) = y_2'(x), y_4(x) = y_3'(x), y_4'(x) = \cos(x) - 16y_1(x)]$$

- Define vector

$$\vec{y}(x) = \begin{bmatrix} y_1(x) \\ y_2(x) \\ y_3(x) \\ y_4(x) \end{bmatrix}$$

- System to solve

$$\vec{y}'(x) = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -16 & 0 & 0 & 0 \end{bmatrix} \cdot \vec{y}(x) + \begin{bmatrix} 0 \\ 0 \\ 0 \\ \cos(x) \end{bmatrix}$$

- Define the forcing function

$$\vec{f}(x) = \begin{bmatrix} 0 \\ 0 \\ 0 \\ \cos(x) \end{bmatrix}$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -16 & 0 & 0 & 0 \end{bmatrix}$$

- Rewrite the system as

$$\vec{y}'(x) = A \cdot \vec{y}(x) + \vec{f}$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\begin{array}{c} \left[\begin{array}{c} -\sqrt{2} - I\sqrt{2}, \\ \frac{1}{(-\sqrt{2}-I\sqrt{2})^3} \\ \frac{1}{(-\sqrt{2}-I\sqrt{2})^2} \\ \frac{1}{-\sqrt{2}-I\sqrt{2}} \\ 1 \end{array} \right] \\ \left[\begin{array}{c} -\sqrt{2} + I\sqrt{2}, \\ \frac{1}{(-\sqrt{2}+I\sqrt{2})^3} \\ \frac{1}{(-\sqrt{2}+I\sqrt{2})^2} \\ \frac{1}{-\sqrt{2}+I\sqrt{2}} \\ 1 \end{array} \right] \\ \left[\begin{array}{c} \sqrt{2} + I\sqrt{2}, \\ \frac{1}{(\sqrt{2}+I\sqrt{2})^3} \\ \frac{1}{(\sqrt{2}+I\sqrt{2})^2} \\ \frac{1}{\sqrt{2}+I\sqrt{2}} \\ 1 \end{array} \right] \end{array} \right]$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$\left[\begin{array}{c} \left[\begin{array}{c} -\sqrt{2} - I\sqrt{2}, \\ \frac{1}{(-\sqrt{2}-I\sqrt{2})^3} \\ \frac{1}{(-\sqrt{2}-I\sqrt{2})^2} \\ \frac{1}{-\sqrt{2}-I\sqrt{2}} \\ 1 \end{array} \right] \end{array} \right]$$

- Solution from eigenpair

$$e^{(-\sqrt{2}-I\sqrt{2})x} \cdot \left[\begin{array}{c} \frac{1}{(-\sqrt{2}-I\sqrt{2})^3} \\ \frac{1}{(-\sqrt{2}-I\sqrt{2})^2} \\ \frac{1}{-\sqrt{2}-I\sqrt{2}} \\ 1 \end{array} \right]$$

- Use Euler identity to write solution in terms of sin and cos

$$e^{-x\sqrt{2}} \cdot (\cos(x\sqrt{2}) - I \sin(x\sqrt{2})) \cdot \begin{bmatrix} \frac{1}{(-\sqrt{2}-I\sqrt{2})^3} \\ \frac{1}{(-\sqrt{2}-I\sqrt{2})^2} \\ \frac{1}{-\sqrt{2}-I\sqrt{2}} \\ 1 \end{bmatrix}$$

- Simplify expression

$$e^{-x\sqrt{2}} \cdot \begin{bmatrix} \frac{\cos(x\sqrt{2}) - I \sin(x\sqrt{2})}{(-\sqrt{2}-I\sqrt{2})^3} \\ \frac{\cos(x\sqrt{2}) - I \sin(x\sqrt{2})}{(-\sqrt{2}-I\sqrt{2})^2} \\ \frac{\cos(x\sqrt{2}) - I \sin(x\sqrt{2})}{-\sqrt{2}-I\sqrt{2}} \\ \cos(x\sqrt{2}) - I \sin(x\sqrt{2}) \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\vec{y}_1(x) = e^{-x\sqrt{2}} \cdot \begin{bmatrix} \frac{\cos(x\sqrt{2})\sqrt{2}}{16} + \frac{\sin(x\sqrt{2})\sqrt{2}}{16} \\ -\frac{\sin(x\sqrt{2})}{4} \\ -\frac{\cos(x\sqrt{2})\sqrt{2}}{4} + \frac{\sin(x\sqrt{2})\sqrt{2}}{4} \\ \cos(x\sqrt{2}) \end{bmatrix}, \vec{y}_2(x) = e^{-x\sqrt{2}} \cdot \begin{bmatrix} \frac{\cos(x\sqrt{2})\sqrt{2}}{16} - \frac{\sin(x\sqrt{2})\sqrt{2}}{16} \\ -\frac{\cos(x\sqrt{2})}{4} \\ \frac{\cos(x\sqrt{2})\sqrt{2}}{4} + \frac{\sin(x\sqrt{2})\sqrt{2}}{4} \\ -\sin(x\sqrt{2}) \end{bmatrix}$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$\sqrt{2} + I\sqrt{2}, \begin{bmatrix} \frac{1}{(\sqrt{2}+I\sqrt{2})^3} \\ \frac{1}{(\sqrt{2}+I\sqrt{2})^2} \\ \frac{1}{\sqrt{2}+I\sqrt{2}} \\ 1 \end{bmatrix}$$

- Solution from eigenpair

$$e^{(\sqrt{2}+I\sqrt{2})x} \cdot \begin{bmatrix} \frac{1}{(\sqrt{2}+I\sqrt{2})^3} \\ \frac{1}{(\sqrt{2}+I\sqrt{2})^2} \\ \frac{1}{\sqrt{2}+I\sqrt{2}} \\ 1 \end{bmatrix}$$

- Use Euler identity to write solution in terms of sin and cos

$$e^{x\sqrt{2}} \cdot (\cos(x\sqrt{2}) + I \sin(x\sqrt{2})) \cdot \begin{bmatrix} \frac{1}{(\sqrt{2}+I\sqrt{2})^3} \\ \frac{1}{(\sqrt{2}+I\sqrt{2})^2} \\ \frac{1}{\sqrt{2}+I\sqrt{2}} \\ 1 \end{bmatrix}$$

- Simplify expression

$$e^{x\sqrt{2}} \cdot \begin{bmatrix} \frac{\cos(x\sqrt{2}) + I \sin(x\sqrt{2})}{(\sqrt{2}+I\sqrt{2})^3} \\ \frac{\cos(x\sqrt{2}) + I \sin(x\sqrt{2})}{(\sqrt{2}+I\sqrt{2})^2} \\ \frac{\cos(x\sqrt{2}) + I \sin(x\sqrt{2})}{\sqrt{2}+I\sqrt{2}} \\ \cos(x\sqrt{2}) + I \sin(x\sqrt{2}) \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\vec{y}_3(x) = e^{x\sqrt{2}} \cdot \begin{bmatrix} -\frac{\cos(x\sqrt{2})\sqrt{2}}{16} + \frac{\sin(x\sqrt{2})\sqrt{2}}{16} \\ \frac{\sin(x\sqrt{2})}{4} \\ \frac{\cos(x\sqrt{2})\sqrt{2}}{4} + \frac{\sin(x\sqrt{2})\sqrt{2}}{4} \\ \cos(x\sqrt{2}) \end{bmatrix}, \vec{y}_4(x) = e^{x\sqrt{2}} \cdot \begin{bmatrix} -\frac{\cos(x\sqrt{2})\sqrt{2}}{16} - \frac{\sin(x\sqrt{2})\sqrt{2}}{16} \\ -\frac{\cos(x\sqrt{2})}{4} \\ -\frac{\cos(x\sqrt{2})\sqrt{2}}{4} + \frac{\sin(x\sqrt{2})\sqrt{2}}{4} \\ \sin(x\sqrt{2}) \end{bmatrix}$$

- General solution of the system of ODEs can be written in terms of the particular solution $\vec{y}_p(x)$

$$\vec{y}(x) = c_1 \vec{y}_1(x) + c_2 \vec{y}_2(x) + c_3 \vec{y}_3(x) + c_4 \vec{y}_4(x) + \vec{y}_p(x)$$

- Fundamental matrix

- Let $\phi(x)$ be the matrix whose columns are the independent solutions of the homogeneous system

$$\phi(x) = \begin{bmatrix} e^{-x\sqrt{2}} \left(\frac{\cos(x\sqrt{2})\sqrt{2}}{16} + \frac{\sin(x\sqrt{2})\sqrt{2}}{16} \right) & e^{-x\sqrt{2}} \left(\frac{\cos(x\sqrt{2})\sqrt{2}}{16} - \frac{\sin(x\sqrt{2})\sqrt{2}}{16} \right) & e^{x\sqrt{2}} \left(-\frac{\cos(x\sqrt{2})\sqrt{2}}{16} \right) \\ -\frac{e^{-x\sqrt{2}} \sin(x\sqrt{2})}{4} & -\frac{e^{-x\sqrt{2}} \cos(x\sqrt{2})}{4} & \frac{e^{x\sqrt{2}} \sin(x\sqrt{2})}{4} \\ e^{-x\sqrt{2}} \left(-\frac{\cos(x\sqrt{2})\sqrt{2}}{4} + \frac{\sin(x\sqrt{2})\sqrt{2}}{4} \right) & e^{-x\sqrt{2}} \left(\frac{\cos(x\sqrt{2})\sqrt{2}}{4} + \frac{\sin(x\sqrt{2})\sqrt{2}}{4} \right) & e^{x\sqrt{2}} \left(\frac{\cos(x\sqrt{2})\sqrt{2}}{4} \right) \\ e^{-x\sqrt{2}} \cos(x\sqrt{2}) & -e^{-x\sqrt{2}} \sin(x\sqrt{2}) & e^{x\sqrt{2}} \cos(x\sqrt{2}) \end{bmatrix}$$

- The fundamental matrix, $\Phi(x)$ is a normalized version of $\phi(x)$ satisfying $\Phi(0) = I$ where I is the identity matrix.
 $\Phi(x) = \phi(x) \cdot \frac{1}{\phi(0)}$

- Substitute the value of $\phi(x)$ and $\phi(0)$

$$\Phi(x) = \begin{bmatrix} e^{-x\sqrt{2}} \left(\frac{\cos(x\sqrt{2})\sqrt{2}}{16} + \frac{\sin(x\sqrt{2})\sqrt{2}}{16} \right) & e^{-x\sqrt{2}} \left(\frac{\cos(x\sqrt{2})\sqrt{2}}{16} - \frac{\sin(x\sqrt{2})\sqrt{2}}{16} \right) & e^{x\sqrt{2}} \left(-\frac{\cos(x\sqrt{2})\sqrt{2}}{16} \right) \\ -\frac{e^{-x\sqrt{2}} \sin(x\sqrt{2})}{4} & -\frac{e^{-x\sqrt{2}} \cos(x\sqrt{2})}{4} & \frac{e^{x\sqrt{2}} \sin(x\sqrt{2})}{4} \\ e^{-x\sqrt{2}} \left(-\frac{\cos(x\sqrt{2})\sqrt{2}}{4} + \frac{\sin(x\sqrt{2})\sqrt{2}}{4} \right) & e^{-x\sqrt{2}} \left(\frac{\cos(x\sqrt{2})\sqrt{2}}{4} + \frac{\sin(x\sqrt{2})\sqrt{2}}{4} \right) & e^{x\sqrt{2}} \left(\frac{\cos(x\sqrt{2})\sqrt{2}}{4} \right) \\ e^{-x\sqrt{2}} \cos(x\sqrt{2}) & -e^{-x\sqrt{2}} \sin(x\sqrt{2}) & e^{x\sqrt{2}} \cos(x\sqrt{2}) \end{bmatrix}$$

- Evaluate and simplify to get the fundamental matrix

$$\Phi(x) = \begin{bmatrix} \frac{\cos(x\sqrt{2})(e^{-x\sqrt{2}}+e^{x\sqrt{2}})}{2} & \frac{\sqrt{2}(e^{-x\sqrt{2}}-e^{x\sqrt{2}})}{2} \\ \frac{\sqrt{2}((- \cos(x\sqrt{2})-\sin(x\sqrt{2}))e^{-x\sqrt{2}}+e^{x\sqrt{2}}(\cos(x\sqrt{2})-\sin(x\sqrt{2})))}{2} & \frac{\sqrt{2}(e^{-x\sqrt{2}}+e^{x\sqrt{2}})}{2} \\ 2 \sin(x\sqrt{2})(e^{-x\sqrt{2}}-e^{x\sqrt{2}}) & \frac{\sqrt{2}((- \cos(x\sqrt{2})+\sin(x\sqrt{2}))e^{-x\sqrt{2}}+e^{x\sqrt{2}}(\cos(x\sqrt{2})+\sin(x\sqrt{2})))}{2} \\ -2\sqrt{2}(e^{-x\sqrt{2}}(-\cos(x\sqrt{2})+\sin(x\sqrt{2}))+e^{x\sqrt{2}}(\cos(x\sqrt{2})+\sin(x\sqrt{2}))) & \frac{\sqrt{2}(e^{-x\sqrt{2}}-e^{x\sqrt{2}})}{2} \end{bmatrix}$$

- Find a particular solution of the system of ODEs using variation of parameters

- Let the particular solution be the fundamental matrix multiplied by $\vec{v}(x)$ and solve for $\vec{v}(x)$

$$\vec{y}_p(x) = \Phi(x) \cdot \vec{v}(x)$$

- Take the derivative of the particular solution

$$\vec{y}'_p(x) = \Phi'(x) \cdot \vec{v}(x) + \Phi(x) \cdot \vec{v}'(x)$$

- Substitute particular solution and its derivative into the system of ODEs

$$\Phi'(x) \cdot \vec{v}(x) + \Phi(x) \cdot \vec{v}'(x) = A \cdot \Phi(x) \cdot \vec{v}(x) + \vec{f}(x)$$

- The fundamental matrix has columns that are solutions to the homogeneous system so its derivative is $A \cdot \Phi(x)$

$$A \cdot \Phi(x) \cdot \vec{v}(x) + \Phi(x) \cdot \vec{v}'(x) = A \cdot \Phi(x) \cdot \vec{v}(x) + \vec{f}(x)$$

- Cancel like terms

$$\Phi(x) \cdot \vec{v}'(x) = \vec{f}(x)$$

- Multiply by the inverse of the fundamental matrix

$$\vec{v}'(x) = \frac{1}{\Phi(x)} \cdot \vec{f}(x)$$

- Integrate to solve for $\vec{v}(x)$

$$\vec{v}(x) = \int_0^x \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds$$

- Plug $\vec{v}(x)$ into the equation for the particular solution

$$\vec{y}_p(x) = \Phi(x) \cdot \left(\int_0^x \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds \right)$$

- Plug in the fundamental matrix and the forcing function and compute

$$\vec{y}_p(x) = \begin{bmatrix} \frac{e^{-x\sqrt{2}} \left((-4 \cos(x\sqrt{2}) + \sin(x\sqrt{2})) e^{2x\sqrt{2}} + 8 \cos(x\sqrt{2}) - 4 \cos(x\sqrt{2}) - \sin(x\sqrt{2}) \right)}{136} \\ \frac{3 \left(\cos(x\sqrt{2}) + \frac{5 \sin(x\sqrt{2})}{3} \right) \sqrt{2} e^{-x\sqrt{2}}}{136} - \frac{3 e^{x\sqrt{2}} \cos(x\sqrt{2}) \sqrt{2}}{136} + \frac{5 e^{x\sqrt{2}} \sin(x\sqrt{2}) \sqrt{2}}{136} - \frac{\sin(x)}{17} \\ - \frac{\left(\left(-\frac{\cos(x\sqrt{2})}{2} - 2 \sin(x\sqrt{2}) \right) e^{2x\sqrt{2}} + \cos(x\sqrt{2}) e^{x\sqrt{2}} - \frac{\cos(x\sqrt{2})}{2} + 2 \sin(x\sqrt{2}) \right) e^{-x\sqrt{2}}}{17} \\ - \frac{5\sqrt{2} \left(\cos(x\sqrt{2}) - \frac{3 \sin(x\sqrt{2})}{5} \right) e^{-x\sqrt{2}}}{34} + \frac{5 e^{x\sqrt{2}} \cos(x\sqrt{2}) \sqrt{2}}{34} + \frac{3 e^{x\sqrt{2}} \sin(x\sqrt{2}) \sqrt{2}}{34} + \frac{\sin(x)}{17} \end{bmatrix}$$

- Plug particular solution back into general solution

$$\vec{y}(x) = c_1 \vec{y}_1(x) + c_2 \vec{y}_2(x) + c_3 \vec{y}_3(x) + c_4 \vec{y}_4(x) + \begin{bmatrix} \frac{e^{-x\sqrt{2}} \left((-4 \cos(x\sqrt{2}) + \sin(x\sqrt{2})) e^{2x\sqrt{2}} + 8 \cos(x\sqrt{2}) - 4 \cos(x\sqrt{2}) - \sin(x\sqrt{2}) \right)}{136} \\ \frac{3 \left(\cos(x\sqrt{2}) + \frac{5 \sin(x\sqrt{2})}{3} \right) \sqrt{2} e^{-x\sqrt{2}}}{136} - \frac{3 e^{x\sqrt{2}} \cos(x\sqrt{2}) \sqrt{2}}{136} \\ - \frac{\left(\left(-\frac{\cos(x\sqrt{2})}{2} - 2 \sin(x\sqrt{2}) \right) e^{2x\sqrt{2}} + \cos(x\sqrt{2}) e^{x\sqrt{2}} - \frac{\cos(x\sqrt{2})}{2} + 2 \sin(x\sqrt{2}) \right) e^{-x\sqrt{2}}}{17} \\ - \frac{5\sqrt{2} \left(\cos(x\sqrt{2}) - \frac{3 \sin(x\sqrt{2})}{5} \right) e^{-x\sqrt{2}}}{34} + \frac{5 e^{x\sqrt{2}} \cos(x\sqrt{2}) \sqrt{2}}{34} \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y = \frac{e^{-x\sqrt{2}} \left(\left(\left(-\frac{8}{17} + (-c_4 - c_3)\sqrt{2} \right) \cos(x\sqrt{2}) + \sin(x\sqrt{2}) \left(\frac{2}{17} + (c_3 - c_4)\sqrt{2} \right) \right) e^{2x\sqrt{2}} + \left(-\frac{8}{17} + (c_1 + c_2)\sqrt{2} \right) \cos(x\sqrt{2}) + \left(-\frac{2}{17} + (c_1 - c_2)\sqrt{2} \right) \sin(x\sqrt{2}) \right)}{16}$$

Maple trace

```

`Methods for high order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 4; linear nonhomogeneous with symmetry [0,1]
trying high order linear exact nonhomogeneous
trying differential order: 4; missing the dependent variable
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 67

```
dsolve(diff(y(x),x$4)+16*y(x)=cos(x),y(x), singsol=all)
```

$$y(x) = c_4 e^{-\sqrt{2}x} \sin(\sqrt{2}x) + c_2 e^{\sqrt{2}x} \sin(\sqrt{2}x) + c_3 e^{-\sqrt{2}x} \cos(\sqrt{2}x) + c_1 e^{\sqrt{2}x} \cos(\sqrt{2}x) + \frac{\cos(x)}{17}$$

✓ Solution by Mathematica

Time used: 0.762 (sec). Leaf size: 74

```
DSolve[y''''[x]+16*y[x]==Cos[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\cos(x)}{17} + e^{-\sqrt{2}x} \left((c_1 e^{2\sqrt{2}x} + c_2) \cos(\sqrt{2}x) + (c_4 e^{2\sqrt{2}x} + c_3) \sin(\sqrt{2}x) \right)$$

10.4 problem 1(d)

Internal problem ID [5994]

Internal file name [OUTPUT/5242_Sunday_June_05_2022_03_28_18_PM_84877747/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 89

Problem number: 1(d).

ODE order: 4.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_high_order , _with_linear_symmetries]]
```

$$y'''' - 4y'''' + 6y'' - 4y' + y = e^x$$

This is higher order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE And y_p is a particular solution to the nonhomogeneous ODE. y_h is the solution to

$$y'''' - 4y'''' + 6y'' - 4y' + y = 0$$

The characteristic equation is

$$\lambda^4 - 4\lambda^3 + 6\lambda^2 - 4\lambda + 1 = 0$$

The roots of the above equation are

$$\lambda_1 = 1$$

$$\lambda_2 = 1$$

$$\lambda_3 = 1$$

$$\lambda_4 = 1$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 e^x + c_2 x e^x + x^2 e^x c_3 + x^3 e^x c_4$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = e^x$$

$$y_2 = x e^x$$

$$y_3 = x^2 e^x$$

$$y_4 = x^3 e^x$$

Now the particular solution to the given ODE is found

$$y'''' - 4y''' + 6y'' - 4y' + y = e^x$$

The particular solution is found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$e^x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[{\{e^x\}}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{x^3 e^x, x e^x, x^2 e^x, e^x\}$$

Since e^x is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[{\{x e^x\}}]$$

Since $x e^x$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[{\{x^2 e^x\}}]$$

Since $x^2 e^x$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[{\{x^3 e^x\}}]$$

Since $x^3 e^x$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[{\{x^4 e^x\}}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x^4 e^x$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$24A_1 e^x = e^x$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{24} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{x^4 e^x}{24}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^x + c_2 x e^x + x^2 e^x c_3 + x^3 e^x c_4) + \left(\frac{x^4 e^x}{24} \right) \end{aligned}$$

Which simplifies to

$$y = e^x (c_4 x^3 + c_3 x^2 + c_2 x + c_1) + \frac{x^4 e^x}{24}$$

Summary

The solution(s) found are the following

$$y = e^x (c_4 x^3 + c_3 x^2 + c_2 x + c_1) + \frac{x^4 e^x}{24} \quad (1)$$

Verification of solutions

$$y = e^x (c_4 x^3 + c_3 x^2 + c_2 x + c_1) + \frac{x^4 e^x}{24}$$

Verified OK.

Maple trace

```
`Methods for high order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 4; linear nonhomogeneous with symmetry [0,1]  
trying high order linear exact nonhomogeneous  
trying differential order: 4; missing the dependent variable  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve(diff(y(x),x$4)-4*diff(y(x),x$3)+6*diff(y(x),x$2)-4*diff(y(x),x)+y(x)=exp(x),y(x), sin
```

$$y(x) = e^x \left(\frac{1}{24}x^4 + c_1 + c_2x + c_3x^2 + x^3c_4 \right)$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 39

```
DSolve[y''''[x]-4*y'''[x]+6*y''[x]-4*y'[x]+y[x]==Exp[x],y[x],x,IncludeSingularSolutions -> T
```

$$y(x) \rightarrow \frac{1}{24}e^x(x^4 + 24c_4x^3 + 24c_3x^2 + 24c_2x + 24c_1)$$

10.5 problem 1(e)

10.5.1 Maple step by step solution 790

Internal problem ID [5995]

Internal file name [OUTPUT/5243_Sunday_June_05_2022_03_28_20_PM_18327027/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 89

Problem number: 1(e).

ODE order: 4.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_high_order , _linear , _nonhomogeneous]]
```

$$y'''' - y = \cos(x)$$

This is higher order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE And y_p is a particular solution to the nonhomogeneous ODE. y_h is the solution to

$$y'''' - y = 0$$

The characteristic equation is

$$\lambda^4 - 1 = 0$$

The roots of the above equation are

$$\lambda_1 = 1$$

$$\lambda_2 = -1$$

$$\lambda_3 = i$$

$$\lambda_4 = -i$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 e^{-x} + c_2 e^x + e^{-ix} c_3 + e^{ix} c_4$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = e^{-x}$$

$$y_2 = e^x$$

$$y_3 = e^{-ix}$$

$$y_4 = e^{ix}$$

Now the particular solution to the given ODE is found

$$y'''' - y = \cos(x)$$

Let the particular solution be

$$y_p = U_1 y_1 + U_2 y_2 + U_3 y_3 + U_4 y_4$$

Where y_i are the basis solutions found above for the homogeneous solution y_h and $U_i(x)$ are functions to be determined as follows

$$U_i = (-1)^{n-i} \int \frac{F(x)W_i(x)}{aW(x)} dx$$

Where $W(x)$ is the Wronskian and $W_i(x)$ is the Wronskian that results after deleting the last row and the i -th column of the determinant and n is the order of the ODE or equivalently, the number of basis solutions, and a is the coefficient of the leading derivative in the ODE, and $F(x)$ is the RHS of the ODE. Therefore, the first step is to find the Wronskian $W(x)$. This is given by

$$W(x) = \begin{vmatrix} y_1 & y_2 & y_3 & y_4 \\ y_1' & y_2' & y_3' & y_4' \\ y_1'' & y_2'' & y_3'' & y_4'' \\ y_1''' & y_2''' & y_3''' & y_4''' \end{vmatrix}$$

Substituting the fundamental set of solutions y_i found above in the Wronskian gives

$$W = \begin{bmatrix} e^{-x} & e^x & e^{-ix} & e^{ix} \\ -e^{-x} & e^x & -ie^{-ix} & ie^{ix} \\ e^{-x} & e^x & -e^{-ix} & -e^{ix} \\ -e^{-x} & e^x & ie^{-ix} & -ie^{ix} \end{bmatrix}$$

$$|W| = 16ie^{-x}e^xe^{-ix}e^{ix}$$

The determinant simplifies to

$$|W| = 16i$$

Now we determine W_i for each U_i .

$$\begin{aligned} W_1(x) &= \det \begin{bmatrix} e^x & e^{-ix} & e^{ix} \\ e^x & -ie^{-ix} & ie^{ix} \\ e^x & -e^{-ix} & -e^{ix} \end{bmatrix} \\ &= 4ie^x \end{aligned}$$

$$\begin{aligned} W_2(x) &= \det \begin{bmatrix} e^{-x} & e^{-ix} & e^{ix} \\ -e^{-x} & -ie^{-ix} & ie^{ix} \\ e^{-x} & -e^{-ix} & -e^{ix} \end{bmatrix} \\ &= 4ie^{-x} \end{aligned}$$

$$\begin{aligned} W_3(x) &= \det \begin{bmatrix} e^{-x} & e^x & e^{ix} \\ -e^{-x} & e^x & ie^{ix} \\ e^{-x} & e^x & -e^{ix} \end{bmatrix} \\ &= -4e^{ix} \end{aligned}$$

$$\begin{aligned} W_4(x) &= \det \begin{bmatrix} e^{-x} & e^x & e^{-ix} \\ -e^{-x} & e^x & -ie^{-ix} \\ e^{-x} & e^x & -e^{-ix} \end{bmatrix} \\ &= -4e^{-ix} \end{aligned}$$

Now we are ready to evaluate each $U_i(x)$.

$$\begin{aligned} U_1 &= (-1)^{4-1} \int \frac{F(x)W_1(x)}{aW(x)} dx \\ &= (-1)^3 \int \frac{(\cos(x))(4ie^x)}{(1)(16i)} dx \\ &= - \int \frac{4i \cos(x) e^x}{16i} dx \\ &= - \int \left(\frac{\cos(x) e^x}{4} \right) dx \\ &= - \frac{\cos(x) e^x}{8} - \frac{\sin(x) e^x}{8} \end{aligned}$$

$$\begin{aligned}
U_2 &= (-1)^{4-2} \int \frac{F(x)W_2(x)}{aW(x)} dx \\
&= (-1)^2 \int \frac{(\cos(x))(4ie^{-x})}{(1)(16i)} dx \\
&= \int \frac{4i \cos(x) e^{-x}}{16i} dx \\
&= \int \left(\frac{e^{-x} \cos(x)}{4} \right) dx \\
&= -\frac{e^{-x} \cos(x)}{8} + \frac{e^{-x} \sin(x)}{8}
\end{aligned}$$

$$\begin{aligned}
U_3 &= (-1)^{4-3} \int \frac{F(x)W_3(x)}{aW(x)} dx \\
&= (-1)^1 \int \frac{(\cos(x))(-4e^{ix})}{(1)(16i)} dx \\
&= - \int \frac{-4 \cos(x) e^{ix}}{16i} dx \\
&= - \int \left(\frac{i \cos(x) e^{ix}}{4} \right) dx \\
&= -\frac{ix}{8} - \frac{e^{2ix}}{16}
\end{aligned}$$

$$\begin{aligned}
U_4 &= (-1)^{4-4} \int \frac{F(x)W_4(x)}{aW(x)} dx \\
&= (-1)^0 \int \frac{(\cos(x))(-4e^{-ix})}{(1)(16i)} dx \\
&= \int \frac{-4 \cos(x) e^{-ix}}{16i} dx \\
&= \int \left(\frac{i \cos(x) e^{-ix}}{4} \right) dx \\
&= \int \frac{i \cos(x) e^{-ix}}{4} dx
\end{aligned}$$

Now that all the U_i functions have been determined, the particular solution is found from

$$y_p = U_1y_1 + U_2y_2 + U_3y_3 + U_4y_4$$

Hence

$$\begin{aligned}y_p &= \left(-\frac{\cos(x) e^x}{8} - \frac{\sin(x) e^x}{8} \right) (e^{-x}) \\&+ \left(-\frac{e^{-x} \cos(x)}{8} + \frac{e^{-x} \sin(x)}{8} \right) (e^x) \\&+ \left(-\frac{ix}{8} - \frac{e^{2ix}}{16} \right) (e^{-ix}) \\&+ \left(\int \frac{i \cos(x) e^{-ix}}{4} dx \right) (e^{ix})\end{aligned}$$

Therefore the particular solution is

$$y_p = -\frac{5 \cos(x)}{16} + \frac{(i - 4x) \sin(x)}{16}$$

Which simplifies to

$$y_p = -\frac{5 \cos(x)}{16} + \frac{(i - 4x) \sin(x)}{16}$$

Therefore the general solution is

$$\begin{aligned}y &= y_h + y_p \\&= (c_1 e^{-x} + c_2 e^x + e^{-ix} c_3 + e^{ix} c_4) + \left(-\frac{5 \cos(x)}{16} + \frac{(i - 4x) \sin(x)}{16} \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-x} + c_2 e^x + e^{-ix} c_3 + e^{ix} c_4 - \frac{5 \cos(x)}{16} + \frac{(i - 4x) \sin(x)}{16} \quad (1)$$

Verification of solutions

$$y = c_1 e^{-x} + c_2 e^x + e^{-ix} c_3 + e^{ix} c_4 - \frac{5 \cos(x)}{16} + \frac{(i - 4x) \sin(x)}{16}$$

Verified OK.

10.5.1 Maple step by step solution

Let's solve

$$y'''' - y = \cos(x)$$

- Highest derivative means the order of the ODE is 4

$$y''''$$

- Convert linear ODE into a system of first order ODEs

- Define new variable $y_1(x)$

$$y_1(x) = y$$

- Define new variable $y_2(x)$

$$y_2(x) = y'$$

- Define new variable $y_3(x)$

$$y_3(x) = y''$$

- Define new variable $y_4(x)$

$$y_4(x) = y'''$$

- Isolate for $y_4'(x)$ using original ODE

$$y_4'(x) = \cos(x) + y_1(x)$$

Convert linear ODE into a system of first order ODEs

$$[y_2(x) = y_1'(x), y_3(x) = y_2'(x), y_4(x) = y_3'(x), y_4'(x) = \cos(x) + y_1(x)]$$

- Define vector

$$\vec{y}(x) = \begin{bmatrix} y_1(x) \\ y_2(x) \\ y_3(x) \\ y_4(x) \end{bmatrix}$$

- System to solve

$$\vec{y}'(x) = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \end{bmatrix} \cdot \vec{y}(x) + \begin{bmatrix} 0 \\ 0 \\ 0 \\ \cos(x) \end{bmatrix}$$

- Define the forcing function

$$\vec{f}(x) = \begin{bmatrix} 0 \\ 0 \\ 0 \\ \cos(x) \end{bmatrix}$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \end{bmatrix}$$

- Rewrite the system as

$$\vec{y}'(x) = A \cdot \vec{y}(x) + \vec{f}$$

- To solve the system, find the eigenvalues and eigenvectors of A
- Eigenpairs of A

$$\left[\left[\begin{bmatrix} -1 \\ 1 \\ -1 \\ 1 \end{bmatrix} \right], \left[\begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} \right], \left[-I, \begin{bmatrix} -I \\ -1 \\ I \\ 1 \end{bmatrix} \right], \left[I, \begin{bmatrix} I \\ -1 \\ -I \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[-1, \begin{bmatrix} -1 \\ 1 \\ -1 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_1 = e^{-x} \cdot \begin{bmatrix} -1 \\ 1 \\ -1 \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[1, \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_2 = e^x \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$\left[-I, \begin{bmatrix} -I \\ -1 \\ I \\ 1 \end{bmatrix} \right]$$

- Solution from eigenpair

$$e^{-Ix} \cdot \begin{bmatrix} -I \\ -1 \\ I \\ 1 \end{bmatrix}$$

- Use Euler identity to write solution in terms of sin and cos

$$(\cos(x) - I \sin(x)) \cdot \begin{bmatrix} -I \\ -1 \\ I \\ 1 \end{bmatrix}$$

- Simplify expression

$$\begin{bmatrix} -I(\cos(x) - I \sin(x)) \\ -\cos(x) + I \sin(x) \\ I(\cos(x) - I \sin(x)) \\ \cos(x) - I \sin(x) \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\vec{y}_3(x) = \begin{bmatrix} -\sin(x) \\ -\cos(x) \\ \sin(x) \\ \cos(x) \end{bmatrix}, \vec{y}_4(x) = \begin{bmatrix} -\cos(x) \\ \sin(x) \\ \cos(x) \\ -\sin(x) \end{bmatrix}$$

- General solution of the system of ODEs can be written in terms of the particular solution $\vec{y}_p(x)$

$$\vec{y}(x) = c_1 \vec{y}_1 + c_2 \vec{y}_2 + c_3 \vec{y}_3(x) + c_4 \vec{y}_4(x) + \vec{y}_p(x)$$

□

Fundamental matrix

- Let $\phi(x)$ be the matrix whose columns are the independent solutions of the homogeneous system

$$\phi(x) = \begin{bmatrix} -e^{-x} & e^x & -\sin(x) & -\cos(x) \\ e^{-x} & e^x & -\cos(x) & \sin(x) \\ -e^{-x} & e^x & \sin(x) & \cos(x) \\ e^{-x} & e^x & \cos(x) & -\sin(x) \end{bmatrix}$$

- The fundamental matrix, $\Phi(x)$ is a normalized version of $\phi(x)$ satisfying $\Phi(0) = I$ where I is the identity matrix

$$\Phi(x) = \phi(x) \cdot \frac{1}{\phi(0)}$$

- Substitute the value of $\phi(x)$ and $\phi(0)$

$$\Phi(x) = \begin{bmatrix} -e^{-x} & e^x & -\sin(x) & -\cos(x) \\ e^{-x} & e^x & -\cos(x) & \sin(x) \\ -e^{-x} & e^x & \sin(x) & \cos(x) \\ e^{-x} & e^x & \cos(x) & -\sin(x) \end{bmatrix} \cdot \frac{1}{\begin{bmatrix} -1 & 1 & 0 & -1 \\ 1 & 1 & -1 & 0 \\ -1 & 1 & 0 & 1 \\ 1 & 1 & 1 & 0 \end{bmatrix}}$$

- Evaluate and simplify to get the fundamental matrix

$$\Phi(x) = \begin{bmatrix} \frac{\cos(x)}{2} + \frac{e^{-x}}{4} + \frac{e^x}{4} & -\frac{e^{-x}}{4} + \frac{e^x}{4} + \frac{\sin(x)}{2} & \frac{e^{-x}}{4} + \frac{e^x}{4} - \frac{\cos(x)}{2} & -\frac{e^{-x}}{4} + \frac{e^x}{4} - \frac{\sin(x)}{2} \\ -\frac{e^{-x}}{4} + \frac{e^x}{4} - \frac{\sin(x)}{2} & \frac{\cos(x)}{2} + \frac{e^{-x}}{4} + \frac{e^x}{4} & -\frac{e^{-x}}{4} + \frac{e^x}{4} + \frac{\sin(x)}{2} & \frac{e^{-x}}{4} + \frac{e^x}{4} - \frac{\cos(x)}{2} \\ \frac{e^{-x}}{4} + \frac{e^x}{4} - \frac{\cos(x)}{2} & -\frac{e^{-x}}{4} + \frac{e^x}{4} - \frac{\sin(x)}{2} & \frac{\cos(x)}{2} + \frac{e^{-x}}{4} + \frac{e^x}{4} & -\frac{e^{-x}}{4} + \frac{e^x}{4} + \frac{\sin(x)}{2} \\ -\frac{e^{-x}}{4} + \frac{e^x}{4} + \frac{\sin(x)}{2} & \frac{e^{-x}}{4} + \frac{e^x}{4} - \frac{\cos(x)}{2} & -\frac{e^{-x}}{4} + \frac{e^x}{4} - \frac{\sin(x)}{2} & \frac{\cos(x)}{2} + \frac{e^{-x}}{4} + \frac{e^x}{4} \end{bmatrix}$$

□

Find a particular solution of the system of ODEs using variation of parameters

- Let the particular solution be the fundamental matrix multiplied by $\vec{v}(x)$ and solve for $\vec{v}(x)$

$$\vec{y}_p(x) = \Phi(x) \cdot \vec{v}(x)$$

- Take the derivative of the particular solution

$$\vec{y}_p'(x) = \Phi'(x) \cdot \vec{v}(x) + \Phi(x) \cdot \vec{v}'(x)$$

- Substitute particular solution and its derivative into the system of ODEs

$$\Phi'(x) \cdot \vec{v}(x) + \Phi(x) \cdot \vec{v}'(x) = A \cdot \Phi(x) \cdot \vec{v}(x) + \vec{f}(x)$$

- The fundamental matrix has columns that are solutions to the homogeneous system so its derivative is $A \cdot \Phi(x)$

$$A \cdot \Phi(x) \cdot \vec{v}(x) + \Phi(x) \cdot \vec{v}'(x) = A \cdot \Phi(x) \cdot \vec{v}(x) + \vec{f}(x)$$

- Cancel like terms

$$\Phi(x) \cdot \vec{v}'(x) = \vec{f}(x)$$

- Multiply by the inverse of the fundamental matrix

$$\vec{v}'(x) = \frac{1}{\Phi(x)} \cdot \vec{f}(x)$$

- Integrate to solve for $\vec{v}(x)$

$$\vec{v}(x) = \int_0^x \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds$$

- Plug $\vec{v}(x)$ into the equation for the particular solution

$$\vec{y}_p(x) = \Phi(x) \cdot \left(\int_0^x \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds \right)$$

- Plug in the fundamental matrix and the forcing function and compute

$$\vec{y}_p(x) = \begin{bmatrix} \frac{e^{-x}}{8} + \frac{e^x}{8} - \frac{\cos(x)}{4} - \frac{\sin(x)x}{4} \\ -\frac{\cos(x)x}{4} - \frac{e^{-x}}{8} + \frac{e^x}{8} \\ \frac{e^{-x}}{8} + \frac{e^x}{8} - \frac{\cos(x)}{4} + \frac{\sin(x)x}{4} \\ \frac{\sin(x)}{2} + \frac{\cos(x)x}{4} - \frac{e^{-x}}{8} + \frac{e^x}{8} \end{bmatrix}$$

- Plug particular solution back into general solution

$$\vec{y}(x) = c_1 \vec{y}_1 + c_2 \vec{y}_2 + c_3 \vec{y}_3(x) + c_4 \vec{y}_4(x) + \begin{bmatrix} \frac{e^{-x}}{8} + \frac{e^x}{8} - \frac{\cos(x)}{4} - \frac{\sin(x)x}{4} \\ -\frac{\cos(x)x}{4} - \frac{e^{-x}}{8} + \frac{e^x}{8} \\ \frac{e^{-x}}{8} + \frac{e^x}{8} - \frac{\cos(x)}{4} + \frac{\sin(x)x}{4} \\ \frac{\sin(x)}{2} + \frac{\cos(x)x}{4} - \frac{e^{-x}}{8} + \frac{e^x}{8} \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y = -c_1 e^{-x} + c_2 e^x + \frac{e^{-x}}{8} + \frac{e^x}{8} - \frac{\cos(x)}{4} - \frac{\sin(x)x}{4} - c_4 \cos(x) - c_3 \sin(x)$$

Maple trace

```

`Methods for high order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 4; linear nonhomogeneous with symmetry [0,1]
trying high order linear exact nonhomogeneous
trying differential order: 4; missing the dependent variable
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 35

```
dsolve(diff(y(x),x$4)-y(x)=cos(x),y(x), singsol=all)
```

$$y(x) = c_4 e^{-x} + \frac{(4c_1 - 1) \cos(x)}{4} + \frac{(-x + 4c_3) \sin(x)}{4} + e^x c_2$$

✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 40

```
DSolve[y''''[x]-y[x]==Cos[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^x + c_3 e^{-x} + \left(-\frac{1}{2} + c_2\right) \cos(x) + \left(-\frac{x}{4} + c_4\right) \sin(x)$$

10.6 problem 1(f)

10.6.1 Solving as second order linear constant coeff ode	796
10.6.2 Solving as linear second order ode solved by an integrating factor ode	801
10.6.3 Solving using Kovacic algorithm	802
10.6.4 Maple step by step solution	809

Internal problem ID [5996]

Internal file name [OUTPUT/5244_Sunday_June_05_2022_03_28_22_PM_60486184/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 89

Problem number: 1(f).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "linear_second_order_ode_solved_by_an_integrating_factor"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' - 2iy' - y = e^{ix} - 2e^{-ix}$$

10.6.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = -2i, C = -1, f(x) = -\cos(x) + 3i \sin(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 2iy' - y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -2i, C = -1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 2i\lambda e^{\lambda x} - e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 2i\lambda - 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -2i, C = -1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{2i}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{(-2i)^2 - (4)(1)(-1)} \\ &= i \end{aligned}$$

Hence this is the case of a double root $\lambda_{1,2} = -i$. Therefore the solution is

$$y = c_1 e^{ix} + c_2 x e^{ix} \quad (1)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^{ix} + e^{ix} c_2 x$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned} y_1 &= e^{ix} \\ y_2 &= x e^{ix} \end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^{ix} & x e^{ix} \\ \frac{d}{dx}(e^{ix}) & \frac{d}{dx}(x e^{ix}) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^{ix} & x e^{ix} \\ ie^{ix} & e^{ix} + ie^{ix}x \end{vmatrix}$$

Therefore

$$W = (e^{ix})(e^{ix} + ie^{ix}x) - (x e^{ix})(ie^{ix})$$

Which simplifies to

$$W = e^{2ix}$$

Which simplifies to

$$W = e^{2ix}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x e^{ix}(-\cos(x) + 3i \sin(x))}{e^{2ix}} dx$$

Which simplifies to

$$u_1 = - \int (-\cos(x) + 3i \sin(x)) x e^{-ix} dx$$

Hence

$$\begin{aligned}
 u_1 &= \frac{\frac{e^{-ix}}{4} - \frac{e^{-ix} \tan(\frac{x}{2})^2}{4} + \frac{x^2 e^{-ix}}{4} + \frac{x e^{-ix} \tan(\frac{x}{2})}{2} - \frac{x^2 e^{-ix} \tan(\frac{x}{2})^2}{4} + \frac{ix e^{-ix}}{4} - \frac{ix e^{-ix} \tan(\frac{x}{2})^2}{4} + \frac{ix^2 e^{-ix} \tan(\frac{x}{2})}{2}}{1 + \tan(\frac{x}{2})^2} \\
 &= \frac{3i \left(\frac{ie^{-ix}}{4} - \frac{ie^{-ix} \tan(\frac{x}{2})^2}{4} - \frac{x e^{-ix}}{4} + \frac{x e^{-ix} \tan(\frac{x}{2})^2}{4} + \frac{x^2 e^{-ix} \tan(\frac{x}{2})}{2} - \frac{ix^2 e^{-ix}}{4} + \frac{ix e^{-ix} \tan(\frac{x}{2})}{2} + \frac{ix^2 e^{-ix} \tan(\frac{x}{2})^2}{4} \right)}{1 + \tan(\frac{x}{2})^2}
 \end{aligned}$$

And Eq. (3) becomes

$$u_2 = \int \frac{e^{ix}(-\cos(x) + 3i \sin(x))}{e^{2ix}} dx$$

Which simplifies to

$$u_2 = \int (-\cos(x) + 3i \sin(x)) e^{-ix} dx$$

Hence

$$\begin{aligned}
 u_2 &= -\frac{e^{-ix} \tan(\frac{x}{2}) + ix e^{-ix} \tan(\frac{x}{2}) + \frac{x e^{-ix}}{2} - \frac{x e^{-ix} \tan(\frac{x}{2})^2}{2}}{1 + \tan(\frac{x}{2})^2} \\
 &+ \frac{3i \left(x e^{-ix} \tan(\frac{x}{2}) + i e^{-ix} \tan(\frac{x}{2}) - \frac{ix e^{-ix}}{2} + \frac{ix e^{-ix} \tan(\frac{x}{2})^2}{2} \right)}{1 + \tan(\frac{x}{2})^2}
 \end{aligned}$$

Which simplifies to

$$\begin{aligned}
 u_1 &= -\frac{e^{-ix}((x^2 - 2ix - 2) \cos(x) + \sin(x) x(ix - 2))}{2} \\
 u_2 &= e^{-ix}(i \sin(x) x + \cos(x) x - 2 \sin(x))
 \end{aligned}$$

Therefore the particular solution, from equation (1) is

$$\begin{aligned}
 y_p(x) &= -\frac{e^{-ix}((x^2 - 2ix - 2) \cos(x) + \sin(x) x(ix - 2)) e^{ix}}{2} \\
 &+ e^{-ix}(i \sin(x) x + \cos(x) x - 2 \sin(x)) x e^{ix}
 \end{aligned}$$

Which simplifies to

$$y_p(x) = \frac{(x^2 + 2ix + 2) \cos(x)}{2} + \frac{\sin(x) x(ix - 2)}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^{ix} + e^{ix} c_2 x) + \left(\frac{(x^2 + 2ix + 2) \cos(x)}{2} + \frac{\sin(x) x(ix - 2)}{2} \right) \end{aligned}$$

Which simplifies to

$$y = e^{ix}(c_2 x + c_1) + \frac{(x^2 + 2ix + 2) \cos(x)}{2} + \frac{\sin(x) x(ix - 2)}{2}$$

Summary

The solution(s) found are the following

$$y = e^{ix}(c_2 x + c_1) + \frac{(x^2 + 2ix + 2) \cos(x)}{2} + \frac{\sin(x) x(ix - 2)}{2} \quad (1)$$




Figure 141: Slope field plot

Verification of solutions

$$y = e^{ix}(c_2x + c_1) + \frac{(x^2 + 2ix + 2) \cos(x)}{2} + \frac{\sin(x) x(ix - 2)}{2}$$

Verified OK.

10.6.2 Solving as linear second order ode solved by an integrating factor ode

The ode satisfies this form

$$y'' + p(x)y' + \frac{(p(x)^2 + p'(x))y}{2} = f(x)$$

Where $p(x) = -2i$. Therefore, there is an integrating factor given by

$$\begin{aligned} M(x) &= e^{\frac{1}{2} \int p dx} \\ &= e^{\int -2i dx} \\ &= e^{-ix} \end{aligned}$$

Multiplying both sides of the ODE by the integrating factor $M(x)$ makes the left side of the ODE a complete differential

$$\begin{aligned} (M(x)y)'' &= (-\cos(x) + 3i \sin(x)) e^{-ix} \\ (y e^{-ix})'' &= (-\cos(x) + 3i \sin(x)) e^{-ix} \end{aligned}$$

Integrating once gives

$$(y e^{-ix})' = e^{-ix}(i \sin(x) x + \cos(x) x - 2 \sin(x)) + c_1$$

Integrating again gives

$$(y e^{-ix}) = -\frac{1}{2} + \frac{e^{-2ix}}{2} + \frac{x^2}{2} + x(c_1 + i) + c_2$$

Hence the solution is

$$y = \frac{-\frac{1}{2} + \frac{e^{-2ix}}{2} + \frac{x^2}{2} + x(c_1 + i) + c_2}{e^{-ix}}$$

Or

$$y = c_1 x e^{ix} + \frac{x^2 e^{ix}}{2} + c_2 e^{ix} + i e^{ix} x - \frac{e^{ix}}{2} + \frac{e^{-ix}}{2}$$

Summary

The solution(s) found are the following

$$y = c_1 x e^{ix} + \frac{x^2 e^{ix}}{2} + c_2 e^{ix} + i e^{ix} x - \frac{e^{ix}}{2} + \frac{e^{-ix}}{2} \quad (1)$$




Figure 142: Slope field plot

Verification of solutions

$$y = c_1 x e^{ix} + \frac{x^2 e^{ix}}{2} + c_2 e^{ix} + i e^{ix} x - \frac{e^{ix}}{2} + \frac{e^{-ix}}{2}$$

Verified OK.

10.6.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 2iy' - y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= 1 \\B &= -2i \\C &= -1\end{aligned}\tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x)\tag{4}$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\&= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned}\tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1}\tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= 0 \\t &= 1\end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 0\tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 149: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - -\infty \\
 &= \infty
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}
 y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\
 &= z_1 e^{-\int \frac{1}{2} \frac{-2i}{1} dx} \\
 &= z_1 e^{ix} \\
 &= z_1 (e^{ix})
 \end{aligned}$$

Which simplifies to

$$y_1 = e^{ix}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{2i}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{2ix}}{(y_1)^2} dx \\ &= y_1(x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1(e^{ix}) + c_2(e^{ix}(x)) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 2iy' - y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{ix} + e^{ix} c_2 x$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of

parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = e^{ix}$$

$$y_2 = x e^{ix}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^{ix} & x e^{ix} \\ \frac{d}{dx}(e^{ix}) & \frac{d}{dx}(x e^{ix}) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^{ix} & x e^{ix} \\ ie^{ix} & e^{ix} + ie^{ix} x \end{vmatrix}$$

Therefore

$$W = (e^{ix})(e^{ix} + ie^{ix} x) - (x e^{ix})(ie^{ix})$$

Which simplifies to

$$W = e^{2ix}$$

Which simplifies to

$$W = e^{2ix}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x e^{ix} (-\cos(x) + 3i \sin(x))}{e^{2ix}} dx$$

Which simplifies to

$$u_1 = - \int (-\cos(x) + 3i \sin(x)) x e^{-ix} dx$$

Hence

$$\begin{aligned} u_1 &= \frac{e^{-ix}}{4} - \frac{e^{-ix} \tan(\frac{x}{2})^2}{4} + \frac{x^2 e^{-ix}}{4} + \frac{x e^{-ix} \tan(\frac{x}{2})}{2} - \frac{x^2 e^{-ix} \tan(\frac{x}{2})^2}{4} + \frac{ix e^{-ix}}{4} - \frac{ix e^{-ix} \tan(\frac{x}{2})^2}{4} + \frac{ix^2 e^{-ix} \tan(\frac{x}{2})}{2} \\ &= \frac{1 + \tan(\frac{x}{2})^2}{1 + \tan(\frac{x}{2})^2} \\ &= \frac{3i \left(\frac{ie^{-ix}}{4} - \frac{ie^{-ix} \tan(\frac{x}{2})^2}{4} - \frac{x e^{-ix}}{4} + \frac{x e^{-ix} \tan(\frac{x}{2})^2}{4} + \frac{x^2 e^{-ix} \tan(\frac{x}{2})}{2} - \frac{ix^2 e^{-ix}}{4} + \frac{ix e^{-ix} \tan(\frac{x}{2})}{2} + \frac{ix^2 e^{-ix} \tan(\frac{x}{2})^2}{4} \right)}{1 + \tan(\frac{x}{2})^2} \end{aligned}$$

And Eq. (3) becomes

$$u_2 = \int \frac{e^{ix} (-\cos(x) + 3i \sin(x))}{e^{2ix}} dx$$

Which simplifies to

$$u_2 = \int (-\cos(x) + 3i \sin(x)) e^{-ix} dx$$

Hence

$$\begin{aligned} u_2 &= - \frac{e^{-ix} \tan(\frac{x}{2}) + ix e^{-ix} \tan(\frac{x}{2}) + \frac{x e^{-ix}}{2} - \frac{x e^{-ix} \tan(\frac{x}{2})^2}{2}}{1 + \tan(\frac{x}{2})^2} \\ &+ \frac{3i \left(x e^{-ix} \tan(\frac{x}{2}) + i e^{-ix} \tan(\frac{x}{2}) - \frac{ix e^{-ix}}{2} + \frac{ix e^{-ix} \tan(\frac{x}{2})^2}{2} \right)}{1 + \tan(\frac{x}{2})^2} \end{aligned}$$

Which simplifies to

$$u_1 = - \frac{e^{-ix} ((x^2 - 2ix - 2) \cos(x) + \sin(x) x (ix - 2))}{2}$$

$$u_2 = e^{-ix} (i \sin(x) x + \cos(x) x - 2 \sin(x))$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{e^{-ix}((x^2 - 2ix - 2) \cos(x) + \sin(x) x(ix - 2)) e^{ix}}{2} + e^{-ix}(i \sin(x) x + \cos(x) x - 2 \sin(x)) x e^{ix}$$

Which simplifies to

$$y_p(x) = \frac{(x^2 + 2ix + 2) \cos(x)}{2} + \frac{\sin(x) x(ix - 2)}{2}$$

Therefore the general solution is

$$y = y_h + y_p \\ = (c_1 e^{ix} + e^{ix} c_2 x) + \left(\frac{(x^2 + 2ix + 2) \cos(x)}{2} + \frac{\sin(x) x(ix - 2)}{2} \right)$$

Which simplifies to

$$y = e^{ix}(c_2 x + c_1) + \frac{(x^2 + 2ix + 2) \cos(x)}{2} + \frac{\sin(x) x(ix - 2)}{2}$$

Summary

The solution(s) found are the following

$$y = e^{ix}(c_2 x + c_1) + \frac{(x^2 + 2ix + 2) \cos(x)}{2} + \frac{\sin(x) x(ix - 2)}{2} \quad (1)$$




Figure 143: Slope field plot

Verification of solutions

$$y = e^{ix}(c_2x + c_1) + \frac{(x^2 + 2ix + 2) \cos(x)}{2} + \frac{\sin(x) x(ix - 2)}{2}$$

Verified OK.

10.6.4 Maple step by step solution

Let's solve

$$y'' - 2Iy' - y = -\cos(x) + 3I\sin(x)$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - 2Ir - 1 = 0$$

- Factor the characteristic polynomial

$$(-r + I)^2 = 0$$

- Root of the characteristic polynomial

$$r = 1$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{Ix}$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = x e^{Ix}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{Ix} + e^{Ix} c_2 x + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = -\cos(x) + 3I \sin(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{Ix} & x e^{Ix} \\ I e^{Ix} & e^{Ix} + I x e^{Ix} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = e^{2Ix}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = e^{Ix} \left(\int (\cos(x) - 3I \sin(x)) x e^{-Ix} dx - x \left(\int (\cos(x) - 3I \sin(x)) e^{-Ix} dx \right) \right)$$

- Compute integrals

$$y_p(x) = \frac{(2Ix+x^2+2)\cos(x)}{2} + \frac{\sin(x)x(Ix-2)}{2}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{Ix} + e^{Ix} c_2 x + \frac{(2Ix+x^2+2)\cos(x)}{2} + \frac{\sin(x)x(Ix-2)}{2}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 58

```
dsolve(diff(y(x),x$2)-2*I*diff(y(x),x)-y(x)=exp(I*x)-2*exp(-I*x),y(x), singsol=all)
```

$$y(x) = -1 + \cos\left(\frac{x}{2}\right)^2 (x^2 + 2ix + 2) + \sin\left(\frac{x}{2}\right) x(ix - 2) \cos\left(\frac{x}{2}\right) + (c_1x + c_2)e^{ix} - ix - \frac{x^2}{2}$$

✓ Solution by Mathematica

Time used: 0.177 (sec). Leaf size: 39

```
DSolve[y''[x]-2*I*y'[x]-y[x]==Exp[I*x]-2*Exp[-I*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}e^{-ix}(1 + e^{2ix}(x^2 + 2c_2x + 2c_1))$$

11 Chapter 2. Linear equations with constant coefficients. Page 93

11.1	problem 1(a)	813
11.2	problem 1(b)	824
11.3	problem 1(c)	835
11.4	problem 1(d)	847
11.5	problem 1(e)	858
11.6	problem 1(f)	869
11.7	problem 1(g)	880
11.8	problem 1(h)	891
11.9	problem 1(i)	895

11.1 problem 1(a)

11.1.1 Solving as second order linear constant coeff ode	813
11.1.2 Solving using Kovacic algorithm	816
11.1.3 Maple step by step solution	821

Internal problem ID [5997]

Internal file name [OUTPUT/5245_Sunday_June_05_2022_03_28_25_PM_22707088/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 93

Problem number: 1(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + 4y = \cos(x)$$

11.1.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 4, f(x) = \cos(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 4y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 4$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 4 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 4 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 4$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(4)} \\ &= \pm 2i \end{aligned}$$

Hence

$$\lambda_1 = +2i$$

$$\lambda_2 = -2i$$

Which simplifies to

$$\lambda_1 = 2i$$

$$\lambda_2 = -2i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 2$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (c_1 \cos(2x) + c_2 \sin(2x))$$

Or

$$y = c_1 \cos(2x) + c_2 \sin(2x)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \cos(2x) + c_2 \sin(2x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\cos(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(2x), \sin(2x)\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(x) + A_2 \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$3A_1 \cos(x) + 3A_2 \sin(x) = \cos(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{3}, A_2 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{\cos(x)}{3}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(2x) + c_2 \sin(2x)) + \left(\frac{\cos(x)}{3} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(2x) + c_2 \sin(2x) + \frac{\cos(x)}{3} \quad (1)$$




Figure 144: Slope field plot

Verification of solutions

$$y = c_1 \cos(2x) + c_2 \sin(2x) + \frac{\cos(x)}{3}$$

Verified OK.

11.1.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 4y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= 1 \\B &= 0 \\C &= 4\end{aligned}\tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x)\tag{4}$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\&= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned}\tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-4}{1}\tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= -4 \\t &= 1\end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -4z(x)\tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 151: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -4$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(2x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}
 y_1 &= z_1 \\
 &= \cos(2x)
 \end{aligned}$$

Which simplifies to

$$y_1 = \cos(2x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(2x) \int \frac{1}{\cos(2x)^2} dx \\ &= \cos(2x) \left(\frac{\tan(2x)}{2} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(2x)) + c_2 \left(\cos(2x) \left(\frac{\tan(2x)}{2} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 4y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 \cos(2x) + \frac{c_2 \sin(2x)}{2}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\cos(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ \frac{\sin(2x)}{2}, \cos(2x) \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(x) + A_2 \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$3A_1 \cos(x) + 3A_2 \sin(x) = \cos(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{3}, A_2 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{\cos(x)}{3}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 \cos(2x) + \frac{c_2 \sin(2x)}{2} \right) + \left(\frac{\cos(x)}{3} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(2x) + \frac{c_2 \sin(2x)}{2} + \frac{\cos(x)}{3} \quad (1)$$




Figure 145: Slope field plot

Verification of solutions

$$y = c_1 \cos(2x) + \frac{c_2 \sin(2x)}{2} + \frac{\cos(x)}{3}$$

Verified OK.

11.1.3 Maple step by step solution

Let's solve

$$y'' + 4y = \cos(x)$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 4 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-16})}{2}$$

- Roots of the characteristic polynomial

$$r = (-2I, 2I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(2x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(2x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 \cos(2x) + c_2 \sin(2x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \cos(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(2x) & \sin(2x) \\ -2\sin(2x) & 2\cos(2x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 2$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{\cos(2x) \left(\int 4 \cos(x)^2 \sin(x) dx \right)}{4} + \frac{\sin(2x) \left(\int (\cos(x) + \cos(3x)) dx \right)}{4}$$

- Compute integrals

$$y_p(x) = \frac{\cos(x)}{3}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 \cos(2x) + c_2 \sin(2x) + \frac{\cos(x)}{3}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 21

```
dsolve(diff(y(x),x$2)+4*y(x)=cos(x),y(x), singsol=all)
```

$$y(x) = \sin(2x)c_2 + \cos(2x)c_1 + \frac{\cos(x)}{3}$$

✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 26

```
DSolve[y''[x]+4*y[x]==Cos[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\cos(x)}{3} + c_1 \cos(2x) + c_2 \sin(2x)$$

11.2 problem 1(b)

11.2.1 Solving as second order linear constant coeff ode	824
11.2.2 Solving using Kovacic algorithm	828
11.2.3 Maple step by step solution	833

Internal problem ID [5998]

Internal file name [OUTPUT/5246_Sunday_June_05_2022_03_28_26_PM_54439359/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 93

Problem number: 1(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + 4y = \sin(2x)$$

11.2.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 4, f(x) = \sin(2x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 4y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 4$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 4 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 4 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 4$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(4)} \\ &= \pm 2i \end{aligned}$$

Hence

$$\lambda_1 = +2i$$

$$\lambda_2 = -2i$$

Which simplifies to

$$\lambda_1 = 2i$$

$$\lambda_2 = -2i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 2$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (c_1 \cos(2x) + c_2 \sin(2x))$$

Or

$$y = c_1 \cos(2x) + c_2 \sin(2x)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \cos(2x) + c_2 \sin(2x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(2x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(2x), \sin(2x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(2x), \sin(2x)\}$$

Since $\cos(2x)$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x \cos(2x), x \sin(2x)\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x \cos(2x) + A_2 x \sin(2x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-4A_1 \sin(2x) + 4A_2 \cos(2x) = \sin(2x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{4}, A_2 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{x \cos(2x)}{4}$$

Therefore the general solution is

$$\begin{aligned}y &= y_h + y_p \\ &= (c_1 \cos(2x) + c_2 \sin(2x)) + \left(-\frac{x \cos(2x)}{4}\right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(2x) + c_2 \sin(2x) - \frac{x \cos(2x)}{4} \quad (1)$$




Figure 146: Slope field plot

Verification of solutions

$$y = c_1 \cos(2x) + c_2 \sin(2x) - \frac{x \cos(2x)}{4}$$

Verified OK.

11.2.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 4y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 4 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-4}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -4 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -4z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 153: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -4$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(2x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}y_1 &= z_1 \\ &= \cos(2x)\end{aligned}$$

Which simplifies to

$$y_1 = \cos(2x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned}y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(2x) \int \frac{1}{\cos(2x)^2} dx \\ &= \cos(2x) \left(\frac{\tan(2x)}{2} \right)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1(\cos(2x)) + c_2 \left(\cos(2x) \left(\frac{\tan(2x)}{2} \right) \right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$.
 y_h is the solution to

$$y'' + 4y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 \cos(2x) + \frac{c_2 \sin(2x)}{2}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(2x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(2x), \sin(2x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ \frac{\sin(2x)}{2}, \cos(2x) \right\}$$

Since $\cos(2x)$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x \cos(2x), x \sin(2x)\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x \cos(2x) + A_2 x \sin(2x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-4A_1 \sin(2x) + 4A_2 \cos(2x) = \sin(2x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{4}, A_2 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{x \cos(2x)}{4}$$

Therefore the general solution is

$$\begin{aligned}y &= y_h + y_p \\ &= \left(c_1 \cos(2x) + \frac{c_2 \sin(2x)}{2} \right) + \left(-\frac{x \cos(2x)}{4} \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(2x) + \frac{c_2 \sin(2x)}{2} - \frac{x \cos(2x)}{4} \quad (1)$$




Figure 147: Slope field plot

Verification of solutions

$$y = c_1 \cos(2x) + \frac{c_2 \sin(2x)}{2} - \frac{x \cos(2x)}{4}$$

Verified OK.

11.2.3 Maple step by step solution

Let's solve

$$y'' + 4y = \sin(2x)$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 4 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-16})}{2}$$

- Roots of the characteristic polynomial

$$r = (-2i, 2i)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(2x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(2x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 \cos(2x) + c_2 \sin(2x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \sin(2x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(2x) & \sin(2x) \\ -2\sin(2x) & 2\cos(2x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 2$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{\cos(2x)\left(\int \sin(2x)^2 dx\right)}{2} + \frac{\sin(2x)\left(\int \sin(4x) dx\right)}{4}$$

- Compute integrals

$$y_p(x) = \frac{\sin(2x)}{16} - \frac{x \cos(2x)}{4}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 \cos(2x) + c_2 \sin(2x) + \frac{\sin(2x)}{16} - \frac{x \cos(2x)}{4}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
dsolve(diff(y(x),x$2)+4*y(x)=sin(2*x),y(x), singsol=all)
```

$$y(x) = \frac{(-x + 4c_1) \cos(2x)}{4} + \sin(2x) c_2$$

✓ Solution by Mathematica

Time used: 0.03 (sec). Leaf size: 33

```
DSolve[y''[x]+4*y[x]==Sin[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \left(-\frac{x}{4} + c_1\right) \cos(2x) + \frac{1}{8}(1 + 16c_2) \sin(x) \cos(x)$$

11.3 problem 1(c)

11.3.1 Solving as second order linear constant coeff ode	835
11.3.2 Solving using Kovacic algorithm	838
11.3.3 Maple step by step solution	845

Internal problem ID [5999]

Internal file name [OUTPUT/5247_Sunday_June_05_2022_03_28_28_PM_81105194/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 93

Problem number: 1(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' - 4y = 3e^{2x} + 4e^{-x}$$

11.3.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = -4, f(x) = 3e^{2x} + 4e^{-x}$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 4y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = -4$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 4 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 4 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = -4$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(-4)} \\ &= \pm 2 \end{aligned}$$

Hence

$$\lambda_1 = +2$$

$$\lambda_2 = -2$$

Which simplifies to

$$\lambda_1 = 2$$

$$\lambda_2 = -2$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(2)x} + c_2 e^{(-2)x}$$

Or

$$y = c_1 e^{2x} + c_2 e^{-2x}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^{2x} + c_2 e^{-2x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$3e^{2x} + 4e^{-x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{-x}\}, \{e^{2x}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^{-2x}, e^{2x}\}$$

Since e^{2x} is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{e^{-x}\}, \{e^{2x}x\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1e^{-x} + A_2e^{2x}x$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-3A_1e^{-x} + 4A_2e^{2x} = 3e^{2x} + 4e^{-x}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{4}{3}, A_2 = \frac{3}{4} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{4e^{-x}}{3} + \frac{3e^{2x}x}{4}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1e^{2x} + c_2e^{-2x}) + \left(-\frac{4e^{-x}}{3} + \frac{3e^{2x}x}{4} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{2x} + c_2 e^{-2x} - \frac{4e^{-x}}{3} + \frac{3e^{2x}x}{4} \quad (1)$$




Figure 148: Slope field plot

Verification of solutions

$$y = c_1 e^{2x} + c_2 e^{-2x} - \frac{4e^{-x}}{3} + \frac{3e^{2x}x}{4}$$

Verified OK.

11.3.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 4y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= 1 \\B &= 0 \\C &= -4\end{aligned}\tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x)\tag{4}$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\&= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned}\tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{4}{1}\tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= 4 \\t &= 1\end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 4z(x)\tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 155: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 4$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-2x}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}
 y_1 &= z_1 \\
 &= e^{-2x}
 \end{aligned}$$

Which simplifies to

$$y_1 = e^{-2x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= e^{-2x} \int \frac{1}{e^{-4x}} dx \\ &= e^{-2x} \left(\frac{e^{4x}}{4} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-2x}) + c_2 \left(e^{-2x} \left(\frac{e^{4x}}{4} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 4y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-2x} + \frac{c_2 e^{2x}}{4}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of

parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = e^{-2x}$$

$$y_2 = \frac{e^{2x}}{4}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^{-2x} & \frac{e^{2x}}{4} \\ \frac{d}{dx}(e^{-2x}) & \frac{d}{dx}\left(\frac{e^{2x}}{4}\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^{-2x} & \frac{e^{2x}}{4} \\ -2e^{-2x} & \frac{e^{2x}}{2} \end{vmatrix}$$

Therefore

$$W = (e^{-2x}) \left(\frac{e^{2x}}{2}\right) - \left(\frac{e^{2x}}{4}\right) (-2e^{-2x})$$

Which simplifies to

$$W = e^{-2x} e^{2x}$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{e^{2x}(3e^{2x} + 4e^{-x})}{1} dx$$

Which simplifies to

$$u_1 = - \int \left(\frac{3e^{4x}}{4} + e^x \right) dx$$

Hence

$$u_1 = -e^x - \frac{3e^{4x}}{16}$$

And Eq. (3) becomes

$$u_2 = \int \frac{e^{-2x}(3e^{2x} + 4e^{-x})}{1} dx$$

Which simplifies to

$$u_2 = \int (3 + 4e^{-3x}) dx$$

Hence

$$u_2 = 3x - \frac{4e^{-3x}}{3}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(-e^x - \frac{3e^{4x}}{16} \right) e^{-2x} + \frac{\left(3x - \frac{4e^{-3x}}{3} \right) e^{2x}}{4}$$

Which simplifies to

$$y_p(x) = \frac{3(4x - 1)e^{2x}}{16} - \frac{4e^{-x}}{3}$$

Therefore the general solution is

$$y = y_h + y_p \\ = \left(c_1 e^{-2x} + \frac{c_2 e^{2x}}{4} \right) + \left(\frac{3(4x - 1) e^{2x}}{16} - \frac{4 e^{-x}}{3} \right)$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-2x} + \frac{c_2 e^{2x}}{4} + \frac{3(4x - 1) e^{2x}}{16} - \frac{4 e^{-x}}{3} \quad (1)$$




Figure 149: Slope field plot

Verification of solutions

$$y = c_1 e^{-2x} + \frac{c_2 e^{2x}}{4} + \frac{3(4x - 1) e^{2x}}{16} - \frac{4 e^{-x}}{3}$$

Verified OK.

11.3.3 Maple step by step solution

Let's solve

$$y'' - 4y = 3e^{2x} + 4e^{-x}$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - 4 = 0$$

- Factor the characteristic polynomial

$$(r - 2)(r + 2) = 0$$

- Roots of the characteristic polynomial

$$r = (-2, 2)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-2x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{2x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{-2x} + c_2 e^{2x} + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = 3e^{2x} + 4e^{-x} \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-2x} & e^{2x} \\ -2e^{-2x} & 2e^{2x} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 4$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{e^{-2x}(\int(3e^{4x}+4e^x)dx)}{4} + \frac{e^{2x}(\int(3+4e^{-3x})dx)}{4}$$

- Compute integrals

$$y_p(x) = \frac{3(4x-1)e^{2x}}{16} - \frac{4e^{-x}}{3}$$

- Substitute particular solution into general solution to ODE

$$y = c_1e^{-2x} + c_2e^{2x} + \frac{3(4x-1)e^{2x}}{16} - \frac{4e^{-x}}{3}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 29

```
dsolve(diff(y(x),x$2)-4*y(x)=3*exp(2*x)+4*exp(-x),y(x), singsol=all)
```

$$y(x) = e^{-2x} \left(\frac{(12x + 16c_2 - 3)e^{4x}}{16} + c_1 - \frac{4e^x}{3} \right)$$

✓ Solution by Mathematica

Time used: 0.345 (sec). Leaf size: 86

```
DSolve[y''[x]-4*y[x]==3*exp[2*x]+4*Exp[-x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2x} \left(e^{4x} \int_1^x \frac{1}{4} e^{-3K[1]} (3e^{K[1]} \exp(2K[1]) + 4) dK[1] + \int_1^x -\frac{1}{4} e^{K[2]} (3e^{K[2]} \exp(2K[2]) + 4) dK[2] + c_1 e^{4x} + c_2 \right)$$

11.4 problem 1(d)

11.4.1 Solving as second order linear constant coeff ode	847
11.4.2 Solving using Kovacic algorithm	850
11.4.3 Maple step by step solution	855

Internal problem ID [6000]

Internal file name [OUTPUT/5248_Sunday_June_05_2022_03_28_29_PM_25528176/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 93

Problem number: 1(d).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' - y' - 2y = x^2 + \cos(x)$$

11.4.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = -1, C = -2, f(x) = x^2 + \cos(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - y' - 2y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -1, C = -2$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - \lambda e^{\lambda x} - 2e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - \lambda - 2 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -1, C = -2$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-1^2 - (4)(1)(-2)} \\ &= \frac{1}{2} \pm \frac{3}{2} \end{aligned}$$

Hence

$$\lambda_1 = \frac{1}{2} + \frac{3}{2}$$

$$\lambda_2 = \frac{1}{2} - \frac{3}{2}$$

Which simplifies to

$$\lambda_1 = 2$$

$$\lambda_2 = -1$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(2)x} + c_2 e^{(-1)x}$$

Or

$$y = c_1 e^{2x} + c_2 e^{-x}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^{2x} + c_2 e^{-x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x^2 + \cos(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}, \{1, x, x^2\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^{-x}, e^{2x}\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(x) + A_2 \sin(x) + A_3 + A_4 x + A_5 x^2$$

The unknowns $\{A_1, A_2, A_3, A_4, A_5\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$\begin{aligned} -3A_1 \cos(x) - 3A_2 \sin(x) + 2A_5 + A_1 \sin(x) - A_2 \cos(x) \\ - A_4 - 2A_5 x - 2A_3 - 2A_4 x - 2A_5 x^2 = x^2 + \cos(x) \end{aligned}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{3}{10}, A_2 = -\frac{1}{10}, A_3 = -\frac{3}{4}, A_4 = \frac{1}{2}, A_5 = -\frac{1}{2} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{3 \cos(x)}{10} - \frac{\sin(x)}{10} - \frac{3}{4} + \frac{x}{2} - \frac{x^2}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^{2x} + c_2 e^{-x}) + \left(-\frac{3 \cos(x)}{10} - \frac{\sin(x)}{10} - \frac{3}{4} + \frac{x}{2} - \frac{x^2}{2} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{2x} + c_2 e^{-x} - \frac{3 \cos(x)}{10} - \frac{\sin(x)}{10} - \frac{3}{4} + \frac{x}{2} - \frac{x^2}{2} \quad (1)$$




Figure 150: Slope field plot

Verification of solutions

$$y = c_1 e^{2x} + c_2 e^{-x} - \frac{3 \cos(x)}{10} - \frac{\sin(x)}{10} - \frac{3}{4} + \frac{x}{2} - \frac{x^2}{2}$$

Verified OK.

11.4.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' - y' - 2y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= 1 \\B &= -1 \\C &= -2\end{aligned}\tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x)\tag{4}$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\&= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned}\tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{9}{4}\tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= 9 \\t &= 4\end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \frac{9z(x)}{4}\tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 157: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{9}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{3x}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}
 y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\
 &= z_1 e^{-\int \frac{1}{2} \frac{-1}{1} dx} \\
 &= z_1 e^{\frac{x}{2}} \\
 &= z_1 \left(e^{\frac{x}{2}} \right)
 \end{aligned}$$

Which simplifies to

$$y_1 = e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-1}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^x}{(y_1)^2} dx \\ &= y_1 \left(\frac{e^{3x}}{3} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-x}) + c_2 \left(e^{-x} \left(\frac{e^{3x}}{3} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - y' - 2y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-x} + \frac{c_2 e^{2x}}{3}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x^2 + \cos(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}, \{1, x, x^2\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ \frac{e^{2x}}{3}, e^{-x} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(x) + A_2 \sin(x) + A_3 + A_4 x + A_5 x^2$$

The unknowns $\{A_1, A_2, A_3, A_4, A_5\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$\begin{aligned} -3A_1 \cos(x) - 3A_2 \sin(x) + 2A_5 + A_1 \sin(x) - A_2 \cos(x) \\ - A_4 - 2A_5 x - 2A_3 - 2A_4 x - 2A_5 x^2 = x^2 + \cos(x) \end{aligned}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{3}{10}, A_2 = -\frac{1}{10}, A_3 = -\frac{3}{4}, A_4 = \frac{1}{2}, A_5 = -\frac{1}{2} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{3 \cos(x)}{10} - \frac{\sin(x)}{10} - \frac{3}{4} + \frac{x}{2} - \frac{x^2}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 e^{-x} + \frac{c_2 e^{2x}}{3} \right) + \left(-\frac{3 \cos(x)}{10} - \frac{\sin(x)}{10} - \frac{3}{4} + \frac{x}{2} - \frac{x^2}{2} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-x} + \frac{c_2 e^{2x}}{3} - \frac{3 \cos(x)}{10} - \frac{\sin(x)}{10} - \frac{3}{4} + \frac{x}{2} - \frac{x^2}{2} \quad (1)$$




Figure 151: Slope field plot

Verification of solutions

$$y = c_1 e^{-x} + \frac{c_2 e^{2x}}{3} - \frac{3 \cos(x)}{10} - \frac{\sin(x)}{10} - \frac{3}{4} + \frac{x}{2} - \frac{x^2}{2}$$

Verified OK.

11.4.3 Maple step by step solution

Let's solve

$$y'' - y' - 2y = x^2 + \cos(x)$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - r - 2 = 0$$

- Factor the characteristic polynomial

$$(r + 1)(r - 2) = 0$$

- Roots of the characteristic polynomial

$$r = (-1, 2)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{2x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{-x} + c_2 e^{2x} + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = x^2 + \cos(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-x} & e^{2x} \\ -e^{-x} & 2e^{2x} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 3e^x$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{e^{-x} \left(\int e^x (x^2 + \cos(x)) dx \right)}{3} + \frac{e^{2x} \left(\int e^{-2x} (x^2 + \cos(x)) dx \right)}{3}$$

- Compute integrals

$$y_p(x) = -\frac{3 \cos(x)}{10} - \frac{\sin(x)}{10} - \frac{3}{4} + \frac{x}{2} - \frac{x^2}{2}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{-x} + c_2 e^{2x} - \frac{3 \cos(x)}{10} - \frac{\sin(x)}{10} - \frac{3}{4} + \frac{x}{2} - \frac{x^2}{2}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
dsolve(diff(y(x),x$2)-diff(y(x),x)-2*y(x)=x^2+cos(x),y(x), singsol=all)
```

$$y(x) = c_2 e^{2x} + c_1 e^{-x} - \frac{x^2}{2} - \frac{3 \cos(x)}{10} - \frac{\sin(x)}{10} + \frac{x}{2} - \frac{3}{4}$$

✓ Solution by Mathematica

Time used: 0.142 (sec). Leaf size: 44

```
DSolve[y''[x]-y'[x]-2*y[x]==x^2+Cos[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{20}(-10x^2 + 10x - 2 \sin(x) - 6 \cos(x) - 15) + c_1 e^{-x} + c_2 e^{2x}$$

11.5 problem 1(e)

11.5.1 Solving as second order linear constant coeff ode	858
11.5.2 Solving using Kovacic algorithm	861
11.5.3 Maple step by step solution	866

Internal problem ID [6001]

Internal file name [OUTPUT/5249_Sunday_June_05_2022_03_28_31_PM_29344672/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 93

Problem number: 1(e).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + 9y = x^2e^{3x}$$

11.5.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 9, f(x) = x^2e^{3x}$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 9y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 9$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 9 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 9 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 9$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(9)} \\ &= \pm 3i \end{aligned}$$

Hence

$$\lambda_1 = +3i$$

$$\lambda_2 = -3i$$

Which simplifies to

$$\lambda_1 = 3i$$

$$\lambda_2 = -3i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 3$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (c_1 \cos(3x) + c_2 \sin(3x))$$

Or

$$y = c_1 \cos(3x) + c_2 \sin(3x)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \cos(3x) + c_2 \sin(3x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x^2 e^{3x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{x^2 e^{3x}, e^{3x} x, e^{3x}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(3x), \sin(3x)\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 x^2 e^{3x} + A_2 e^{3x} x + A_3 e^{3x}$$

The unknowns $\{A_1, A_2, A_3\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$2A_1 e^{3x} + 12A_1 x e^{3x} + 18A_1 x^2 e^{3x} + 18A_2 e^{3x} x + 6A_2 e^{3x} + 18A_3 e^{3x} = x^2 e^{3x}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{18}, A_2 = -\frac{1}{27}, A_3 = \frac{1}{162} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{x^2 e^{3x}}{18} - \frac{e^{3x} x}{27} + \frac{e^{3x}}{162}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(3x) + c_2 \sin(3x)) + \left(\frac{x^2 e^{3x}}{18} - \frac{e^{3x} x}{27} + \frac{e^{3x}}{162} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(3x) + c_2 \sin(3x) + \frac{x^2 e^{3x}}{18} - \frac{e^{3x} x}{27} + \frac{e^{3x}}{162} \quad (1)$$




Figure 152: Slope field plot

Verification of solutions

$$y = c_1 \cos(3x) + c_2 \sin(3x) + \frac{x^2 e^{3x}}{18} - \frac{e^{3x} x}{27} + \frac{e^{3x}}{162}$$

Verified OK.

11.5.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 9y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= 1 \\B &= 0 \\C &= 9\end{aligned}\tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x)\tag{4}$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\&= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned}\tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-9}{1}\tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= -9 \\t &= 1\end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -9z(x)\tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 159: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -9$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(3x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}
 y_1 &= z_1 \\
 &= \cos(3x)
 \end{aligned}$$

Which simplifies to

$$y_1 = \cos(3x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(3x) \int \frac{1}{\cos(3x)^2} dx \\ &= \cos(3x) \left(\frac{\tan(3x)}{3} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(3x)) + c_2 \left(\cos(3x) \left(\frac{\tan(3x)}{3} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 9y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 \cos(3x) + \frac{c_2 \sin(3x)}{3}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x^2 e^{3x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$\{x^2 e^{3x}, e^{3x} x, e^{3x}\}$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ \frac{\sin(3x)}{3}, \cos(3x) \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 x^2 e^{3x} + A_2 e^{3x} x + A_3 e^{3x}$$

The unknowns $\{A_1, A_2, A_3\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$2A_1 e^{3x} + 12A_1 x e^{3x} + 18A_1 x^2 e^{3x} + 18A_2 e^{3x} x + 6A_2 e^{3x} + 18A_3 e^{3x} = x^2 e^{3x}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{18}, A_2 = -\frac{1}{27}, A_3 = \frac{1}{162} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{x^2 e^{3x}}{18} - \frac{e^{3x} x}{27} + \frac{e^{3x}}{162}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 \cos(3x) + \frac{c_2 \sin(3x)}{3} \right) + \left(\frac{x^2 e^{3x}}{18} - \frac{e^{3x} x}{27} + \frac{e^{3x}}{162} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(3x) + \frac{c_2 \sin(3x)}{3} + \frac{x^2 e^{3x}}{18} - \frac{e^{3x} x}{27} + \frac{e^{3x}}{162} \quad (1)$$




Figure 153: Slope field plot

Verification of solutions

$$y = c_1 \cos(3x) + \frac{c_2 \sin(3x)}{3} + \frac{x^2 e^{3x}}{18} - \frac{e^{3x} x}{27} + \frac{e^{3x}}{162}$$

Verified OK.

11.5.3 Maple step by step solution

Let's solve

$$y'' + 9y = x^2 e^{3x}$$

- Highest derivative means the order of the ODE is 2
- y''
- Characteristic polynomial of homogeneous ODE

$$r^2 + 9 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-36})}{2}$$

- Roots of the characteristic polynomial

$$r = (-3I, 3I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(3x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(3x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 \cos(3x) + c_2 \sin(3x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = x^2 e^{3x} \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(3x) & \sin(3x) \\ -3 \sin(3x) & 3 \cos(3x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 3$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{\cos(3x) \left(\int \sin(3x) x^2 e^{3x} dx \right)}{3} + \frac{\sin(3x) \left(\int \cos(3x) x^2 e^{3x} dx \right)}{3}$$

- Compute integrals

$$y_p(x) = \frac{e^{3x}(3x-1)^2}{162}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 \cos(3x) + c_2 \sin(3x) + \frac{e^{3x}(3x-1)^2}{162}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 28

```
dsolve(diff(y(x),x$2)+9*y(x)=x^2*exp(3*x),y(x), singsol=all)
```

$$y(x) = \frac{\left(x - \frac{1}{3}\right)^2 e^{3x}}{18} + \cos(3x) c_1 + \sin(3x) c_2$$

✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 36

```
DSolve[y''[x]+9*y[x]==x^2*Exp[3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{162} e^{3x} (1 - 3x)^2 + c_1 \cos(3x) + c_2 \sin(3x)$$

11.6 problem 1(f)

11.6.1 Solving as second order linear constant coeff ode	869
11.6.2 Solving using Kovacic algorithm	873
11.6.3 Maple step by step solution	877

Internal problem ID [6002]

Internal file name [OUTPUT/5250_Sunday_June_05_2022_03_28_33_PM_86921640/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 93

Problem number: 1(f).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + y = x e^x \cos(2x)$$

11.6.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 1, f(x) = x e^x \cos(2x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\lambda_1 = +i$$

$$\lambda_2 = -i$$

Which simplifies to

$$\lambda_1 = i$$

$$\lambda_2 = -i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (\cos(x) c_1 + c_2 \sin(x))$$

Or

$$y = \cos(x) c_1 + c_2 \sin(x)$$

Therefore the homogeneous solution y_h is

$$y_h = \cos(x) c_1 + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x e^x \cos(2x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$\{e^x \cos(2x), e^x \sin(2x), x e^x \cos(2x), x e^x \sin(2x)\}$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^x \cos(2x) + A_2 e^x \sin(2x) + A_3 x e^x \cos(2x) + A_4 x e^x \sin(2x)$$

The unknowns $\{A_1, A_2, A_3, A_4\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$\begin{aligned} & -2A_1 e^x \cos(2x) - 4A_1 e^x \sin(2x) - 2A_2 e^x \sin(2x) + 4A_2 e^x \cos(2x) + 2A_3 e^x \cos(2x) \\ & - 4A_3 e^x \sin(2x) - 2A_3 x e^x \cos(2x) - 4A_3 x e^x \sin(2x) + 2A_4 e^x \sin(2x) \\ & + 4A_4 e^x \cos(2x) - 2A_4 x e^x \sin(2x) + 4A_4 x e^x \cos(2x) = x e^x \cos(2x) \end{aligned}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{11}{50}, A_2 = -\frac{1}{25}, A_3 = -\frac{1}{10}, A_4 = \frac{1}{5} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{11 e^x \cos(2x)}{50} - \frac{e^x \sin(2x)}{25} - \frac{x e^x \cos(2x)}{10} + \frac{x e^x \sin(2x)}{5}$$

Therefore the general solution is

$$y = y_h + y_p$$

$$= (\cos(x) c_1 + c_2 \sin(x)) + \left(\frac{11 e^x \cos(2x)}{50} - \frac{e^x \sin(2x)}{25} - \frac{x e^x \cos(2x)}{10} + \frac{x e^x \sin(2x)}{5} \right)$$

Summary

The solution(s) found are the following

$$y = \cos(x) c_1 + c_2 \sin(x) + \frac{11 e^x \cos(2x)}{50} - \frac{e^x \sin(2x)}{25} - \frac{x e^x \cos(2x)}{10} + \frac{x e^x \sin(2x)}{5} \quad (1)$$




Figure 154: Slope field plot

Verification of solutions

$$y = \cos(x) c_1 + c_2 \sin(x) + \frac{11 e^x \cos(2x)}{50} - \frac{e^x \sin(2x)}{25} - \frac{x e^x \cos(2x)}{10} + \frac{x e^x \sin(2x)}{5}$$

Verified OK.

11.6.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 1 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 161: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}y_1 &= z_1 \\ &= \cos(x)\end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned}y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x))\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(x)) + c_2 (\cos(x) (\tan(x)))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = \cos(x) c_1 + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x e^x \cos(2x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$\{e^x \cos(2x), e^x \sin(2x), x e^x \cos(2x), x e^x \sin(2x)\}$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^x \cos(2x) + A_2 e^x \sin(2x) + A_3 x e^x \cos(2x) + A_4 x e^x \sin(2x)$$

The unknowns $\{A_1, A_2, A_3, A_4\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$\begin{aligned} & -2A_1 e^x \cos(2x) - 4A_1 e^x \sin(2x) - 2A_2 e^x \sin(2x) + 4A_2 e^x \cos(2x) + 2A_3 e^x \cos(2x) \\ & - 4A_3 e^x \sin(2x) - 2A_3 x e^x \cos(2x) - 4A_3 x e^x \sin(2x) + 2A_4 e^x \sin(2x) \\ & + 4A_4 e^x \cos(2x) - 2A_4 x e^x \sin(2x) + 4A_4 x e^x \cos(2x) = x e^x \cos(2x) \end{aligned}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{11}{50}, A_2 = -\frac{1}{25}, A_3 = -\frac{1}{10}, A_4 = \frac{1}{5} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{11 e^x \cos(2x)}{50} - \frac{e^x \sin(2x)}{25} - \frac{x e^x \cos(2x)}{10} + \frac{x e^x \sin(2x)}{5}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (\cos(x) c_1 + c_2 \sin(x)) + \left(\frac{11 e^x \cos(2x)}{50} - \frac{e^x \sin(2x)}{25} - \frac{x e^x \cos(2x)}{10} + \frac{x e^x \sin(2x)}{5} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \cos(x) c_1 + c_2 \sin(x) + \frac{11 e^x \cos(2x)}{50} - \frac{e^x \sin(2x)}{25} - \frac{x e^x \cos(2x)}{10} + \frac{x e^x \sin(2x)}{5} \quad (1)$$




Figure 155: Slope field plot

Verification of solutions

$$y = \cos(x) c_1 + c_2 \sin(x) + \frac{11 e^x \cos(2x)}{50} - \frac{e^x \sin(2x)}{25} - \frac{x e^x \cos(2x)}{10} + \frac{x e^x \sin(2x)}{5}$$

Verified OK.

11.6.3 Maple step by step solution

Let's solve

$$y'' + y = x e^x \cos(2x)$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (-I, I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = \cos(x) c_1 + c_2 \sin(x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x), y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x), y_2(x))} dx \right), f(x) = x e^x \cos(2x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\cos(x) \left(\int \sin(x) x e^x \cos(2x) dx \right) + \sin(x) \left(\int \cos(x) x e^x \cos(2x) dx \right)$$

- Compute integrals

$$y_p(x) = -\frac{e^x \left(\left(x - \frac{11}{5} \right) \cos(x)^2 - 2 \sin(x) \left(-\frac{1}{5} + x \right) \cos(x) - \frac{x}{2} + \frac{11}{10} \right)}{5}$$

- Substitute particular solution into general solution to ODE

$$y = \cos(x) c_1 + c_2 \sin(x) - \frac{e^x \left(\left(x - \frac{11}{5} \right) \cos(x)^2 - 2 \sin(x) \left(-\frac{1}{5} + x \right) \cos(x) - \frac{x}{2} + \frac{11}{10} \right)}{5}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 42

```
dsolve(diff(y(x),x$2)+y(x)=x*exp(x)*cos(2*x),y(x), singsol=all)
```

$$y(x) = \frac{((-10x + 22) \cos(x))^2 + (20x - 4) \sin(x) \cos(x) + 5x - 11) e^x}{50} + \cos(x) c_1 + \sin(x) c_2$$

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 45

```
DSolve[y''[x]+y[x]==x*Exp[x]*Cos[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{50}e^x(2(1 - 5x) \sin(2x) + (5x - 11) \cos(2x)) + c_1 \cos(x) + c_2 \sin(x)$$

11.7 problem 1(g)

11.7.1 Solving as second order linear constant coeff ode	880
11.7.2 Solving using Kovacic algorithm	883
11.7.3 Maple step by step solution	888

Internal problem ID [6003]

Internal file name [OUTPUT/5251_Sunday_June_05_2022_03_28_34_PM_39858621/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 93

Problem number: 1(g).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"kovacic", "second_order_linear_constant_coeff"**

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + iy' + 2y = 2 \cosh(2x) + e^{-2x}$$

11.7.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = i, C = 2, f(x) = 2e^{-2x} + e^{2x}$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + iy' + 2y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = i, C = 2$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + i\lambda e^{\lambda x} + 2e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + i\lambda + 2 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = i, C = 2$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-i}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{i^2 - (4)(1)(2)} \\ &= -\frac{i}{2} \pm \frac{3i}{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= -\frac{i}{2} + \frac{3i}{2} \\ \lambda_2 &= -\frac{i}{2} - \frac{3i}{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= i \\ \lambda_2 &= -2i \end{aligned}$$

The roots are complex but they are not conjugate of each others. Hence simplification using Euler relation is not possible here. Therefore the final solution is

$$\begin{aligned} y &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ &= c_1 e^{ix} + c_2 e^{-2ix} \end{aligned}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^{ix} + e^{-2ix} c_2$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$2e^{-2x} + e^{2x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{-2x}\}, \{e^{2x}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^{ix}, e^{-2ix}\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^{-2x} + A_2 e^{2x}$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$6A_1 e^{-2x} + 6A_2 e^{2x} + i(-2A_1 e^{-2x} + 2A_2 e^{2x}) = 2e^{-2x} + e^{2x}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{3}{10} + \frac{i}{10}, A_2 = \frac{3}{20} - \frac{i}{20} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \left(\frac{3}{10} + \frac{i}{10} \right) e^{-2x} + \left(\frac{3}{20} - \frac{i}{20} \right) e^{2x}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^{ix} + e^{-2ix} c_2) + \left(\left(\frac{3}{10} + \frac{i}{10} \right) e^{-2x} + \left(\frac{3}{20} - \frac{i}{20} \right) e^{2x} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{ix} + e^{-2ix} c_2 + \left(\frac{3}{10} + \frac{i}{10} \right) e^{-2x} + \left(\frac{3}{20} - \frac{i}{20} \right) e^{2x} \quad (1)$$




Figure 156: Slope field plot

Verification of solutions

$$y = c_1 e^{ix} + e^{-2ix} c_2 + \left(\frac{3}{10} + \frac{i}{10} \right) e^{-2x} + \left(\frac{3}{20} - \frac{i}{20} \right) e^{2x}$$

Verified OK.

11.7.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + iy' + 2y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = i \tag{3}$$

$$C = 2$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-9}{4} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -9 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -\frac{9z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 163: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -\frac{9}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos\left(\frac{3x}{2}\right)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{i}{1} dx} \\ &= z_1 e^{-\frac{ix}{2}} \\ &= z_1 \left(e^{-\frac{ix}{2}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = \cos\left(\frac{3x}{2}\right) e^{-\frac{ix}{2}}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{i}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-ix}}{(y_1)^2} dx \\ &= y_1 \left(\frac{2 \tan\left(\frac{3x}{2}\right)}{3} \right) \end{aligned}$$

Therefore the solution is

$$y = c_1 y_1 + c_2 y_2$$

$$= c_1 \left(\cos \left(\frac{3x}{2} \right) e^{-\frac{ix}{2}} \right) + c_2 \left(\cos \left(\frac{3x}{2} \right) e^{-\frac{ix}{2}} \left(\frac{2 \tan \left(\frac{3x}{2} \right)}{3} \right) \right)$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + iy' + 2y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 \cos \left(\frac{3x}{2} \right) e^{-\frac{ix}{2}} + \frac{2c_2 e^{-\frac{ix}{2}} \sin \left(\frac{3x}{2} \right)}{3}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$2e^{-2x} + e^{2x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{-2x}\}, \{e^{2x}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ \cos \left(\frac{3x}{2} \right) e^{-\frac{ix}{2}}, \frac{2e^{-\frac{ix}{2}} \sin \left(\frac{3x}{2} \right)}{3} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^{-2x} + A_2 e^{2x}$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$6A_1e^{-2x} + 6A_2e^{2x} + i(-2A_1e^{-2x} + 2A_2e^{2x}) = 2e^{-2x} + e^{2x}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{3}{10} + \frac{i}{10}, A_2 = \frac{3}{20} - \frac{i}{20} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \left(\frac{3}{10} + \frac{i}{10} \right) e^{-2x} + \left(\frac{3}{20} - \frac{i}{20} \right) e^{2x}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 \cos\left(\frac{3x}{2}\right) e^{-\frac{ix}{2}} + \frac{2c_2 e^{-\frac{ix}{2}} \sin\left(\frac{3x}{2}\right)}{3} \right) + \left(\left(\frac{3}{10} + \frac{i}{10} \right) e^{-2x} + \left(\frac{3}{20} - \frac{i}{20} \right) e^{2x} \right) \end{aligned}$$

Which simplifies to

$$y = \frac{e^{-\frac{ix}{2}} (3c_1 \cos\left(\frac{3x}{2}\right) + 2c_2 \sin\left(\frac{3x}{2}\right))}{3} + \left(\frac{3}{10} + \frac{i}{10} \right) e^{-2x} + \left(\frac{3}{20} - \frac{i}{20} \right) e^{2x}$$

Summary

The solution(s) found are the following

$$y = \frac{e^{-\frac{ix}{2}} (3c_1 \cos\left(\frac{3x}{2}\right) + 2c_2 \sin\left(\frac{3x}{2}\right))}{3} + \left(\frac{3}{10} + \frac{i}{10} \right) e^{-2x} + \left(\frac{3}{20} - \frac{i}{20} \right) e^{2x} \quad (1)$$




Figure 157: Slope field plot

Verification of solutions

$$y = \frac{e^{-\frac{ix}{2}} \left(3c_1 \cos\left(\frac{3x}{2}\right) + 2c_2 \sin\left(\frac{3x}{2}\right) \right)}{3} + \left(\frac{3}{10} + \frac{i}{10} \right) e^{-2x} + \left(\frac{3}{20} - \frac{i}{20} \right) e^{2x}$$

Verified OK.

11.7.3 Maple step by step solution

Let's solve

$$y'' + Iy' + 2y = 2e^{-2x} + e^{2x}$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + Ir + 2 = 0$$

- Factor the characteristic polynomial

$$-(-r + I)(r + 2I) = 0$$

- Roots of the characteristic polynomial

$$r = (-2I, I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(2x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(2x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 \cos(2x) + c_2 \sin(2x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = 2e^{-2x} + e^{2x} \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(2x) & \sin(2x) \\ -2\sin(2x) & 2\cos(2x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 2$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{\cos(2x)(\int \sin(2x)(2e^{-2x}+e^{2x})dx)}{2} + \frac{\sin(2x)(\int \cos(2x)(2e^{-2x}+e^{2x})dx)}{2}$$

- Compute integrals

$$y_p(x) = \frac{e^{-2x}}{4} + \frac{e^{2x}}{8}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 \cos(2x) + c_2 \sin(2x) + \frac{e^{-2x}}{4} + \frac{e^{2x}}{8}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 35

```
dsolve(diff(y(x),x$2)+I*diff(y(x),x)+2*y(x)=2*cosh(2*x)+exp(-2*x),y(x), singsol=all)
```

$$y(x) = c_2 e^{ix} + e^{-2ix} c_1 + \left(\frac{3}{10} + \frac{i}{10}\right) e^{-2x} + \left(\frac{3}{20} - \frac{i}{20}\right) e^{2x}$$

✓ Solution by Mathematica

Time used: 0.157 (sec). Leaf size: 48

```
DSolve[y''[x]+I*y'[x]+2*y[x]==2*Cosh[2*x]+Exp[-2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{20} e^{-2x} ((3-i)e^{4x} + (6+2i)) + c_1 e^{-2ix} + c_2 e^{ix}$$

11.8 problem 1(h)

Internal problem ID [6004]

Internal file name [OUTPUT/5252_Sunday_June_05_2022_03_28_36_PM_69622299/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 93

Problem number: 1(h).

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_3rd_order , _quadrature]]
```

$$y''' = x^2 + e^{-x} \sin(x)$$

This is higher order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE And y_p is a particular solution to the nonhomogeneous ODE. y_h is the solution to

$$y''' = 0$$

The characteristic equation is

$$\lambda^3 = 0$$

The roots of the above equation are

$$\lambda_1 = 0$$

$$\lambda_2 = 0$$

$$\lambda_3 = 0$$

Therefore the homogeneous solution is

$$y_h(x) = c_3x^2 + c_2x + c_1$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = 1$$

$$y_2 = x$$

$$y_3 = x^2$$

Now the particular solution to the given ODE is found

$$y''' = x^2 + e^{-x} \sin(x)$$

The particular solution is found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x^2 + e^{-x} \sin(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{-x} \cos(x), e^{-x} \sin(x)\}, \{1, x, x^2\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, x, x^2\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{e^{-x} \cos(x), e^{-x} \sin(x)\}, \{x, x^2, x^3\}]$$

Since x is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{e^{-x} \cos(x), e^{-x} \sin(x)\}, \{x^2, x^3, x^4\}]$$

Since x^2 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{e^{-x} \cos(x), e^{-x} \sin(x)\}, \{x^3, x^4, x^5\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 e^{-x} \cos(x) + A_2 e^{-x} \sin(x) + A_3 x^3 + A_4 x^4 + A_5 x^5$$

The unknowns $\{A_1, A_2, A_3, A_4, A_5\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$\begin{aligned} & -2A_1e^{-x}\sin(x) + 2A_1e^{-x}\cos(x) + 2A_2e^{-x}\cos(x) \\ & + 2A_2e^{-x}\sin(x) + 6A_3 + 24A_4x + 60A_5x^2 = x^2 + e^{-x}\sin(x) \end{aligned}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{4}, A_2 = \frac{1}{4}, A_3 = 0, A_4 = 0, A_5 = \frac{1}{60} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{e^{-x}\cos(x)}{4} + \frac{e^{-x}\sin(x)}{4} + \frac{x^5}{60}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_3x^2 + c_2x + c_1) + \left(-\frac{e^{-x}\cos(x)}{4} + \frac{e^{-x}\sin(x)}{4} + \frac{x^5}{60} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_3x^2 + c_2x + c_1 - \frac{e^{-x}\cos(x)}{4} + \frac{e^{-x}\sin(x)}{4} + \frac{x^5}{60} \quad (1)$$

Verification of solutions

$$y = c_3x^2 + c_2x + c_1 - \frac{e^{-x}\cos(x)}{4} + \frac{e^{-x}\sin(x)}{4} + \frac{x^5}{60}$$

Verified OK.

Maple trace

```
`Methods for third order ODEs:
--- Trying classification methods ---
trying a quadrature
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 33

```
dsolve(diff(y(x),x$3)=x^2+exp(-x)*sin(x),y(x), singsol=all)
```

$$y(x) = \frac{e^{-x}(-\cos(x) + \sin(x))}{4} + \frac{x^5}{60} + \frac{c_1 x^2}{2} + c_2 x + c_3$$

✓ Solution by Mathematica

Time used: 0.114 (sec). Leaf size: 47

```
DSolve[y'''[x]==x^2+Exp[-x]*Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^5}{60} + c_3 x^2 + \frac{1}{4} e^{-x} \sin(x) - \frac{1}{4} e^{-x} \cos(x) + c_2 x + c_1$$

11.9 problem 1(i)

Internal problem ID [6005]

Internal file name [OUTPUT/5253_Sunday_June_05_2022_03_28_38_PM_63402563/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 2. Linear equations with constant coefficients. Page 93

Problem number: 1(i).

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_3rd_order , _linear , _nonhomogeneous]]
```

$$y''' + 3y'' + 3y' + y = x^2e^{-x}$$

This is higher order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE And y_p is a particular solution to the nonhomogeneous ODE. y_h is the solution to

$$y''' + 3y'' + 3y' + y = 0$$

The characteristic equation is

$$\lambda^3 + 3\lambda^2 + 3\lambda + 1 = 0$$

The roots of the above equation are

$$\lambda_1 = -1$$

$$\lambda_2 = -1$$

$$\lambda_3 = -1$$

Therefore the homogeneous solution is

$$y_h(x) = c_1e^{-x} + xe^{-x}c_2 + c_3x^2e^{-x}$$

The fundamental set of solutions for the homogeneous solution are the following

$$\begin{aligned}y_1 &= e^{-x} \\y_2 &= x e^{-x} \\y_3 &= x^2 e^{-x}\end{aligned}$$

Now the particular solution to the given ODE is found

$$y''' + 3y'' + 3y' + y = x^2 e^{-x}$$

The particular solution is found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x^2 e^{-x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[{\{x e^{-x}, x^2 e^{-x}, e^{-x}\}}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{x e^{-x}, x^2 e^{-x}, e^{-x}\}$$

Since e^{-x} is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[{\{x e^{-x}, x^2 e^{-x}, x^3 e^{-x}\}}]$$

Since $x e^{-x}$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[{\{x^2 e^{-x}, x^3 e^{-x}, x^4 e^{-x}\}}]$$

Since $x^2 e^{-x}$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[{\{x^3 e^{-x}, x^4 e^{-x}, x^5 e^{-x}\}}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x^3 e^{-x} + A_2 x^4 e^{-x} + A_3 x^5 e^{-x}$$

The unknowns $\{A_1, A_2, A_3\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$24A_2x e^{-x} + 60A_3x^2e^{-x} + 6A_1e^{-x} = x^2e^{-x}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = 0, A_2 = 0, A_3 = \frac{1}{60} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{x^5e^{-x}}{60}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1e^{-x} + xe^{-x}c_2 + c_3x^2e^{-x}) + \left(\frac{x^5e^{-x}}{60} \right) \end{aligned}$$

Which simplifies to

$$y = e^{-x}(c_3x^2 + c_2x + c_1) + \frac{x^5e^{-x}}{60}$$

Summary

The solution(s) found are the following

$$y = e^{-x}(c_3x^2 + c_2x + c_1) + \frac{x^5e^{-x}}{60} \quad (1)$$

Verification of solutions

$$y = e^{-x}(c_3x^2 + c_2x + c_1) + \frac{x^5e^{-x}}{60}$$

Verified OK.

Maple trace

```
`Methods for third order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 3; linear nonhomogeneous with symmetry [0,1]  
trying high order linear exact nonhomogeneous  
trying differential order: 3; missing the dependent variable  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 24

```
dsolve(diff(y(x),x$3)+3*diff(y(x),x$2)+3*diff(y(x),x)+y(x)=x^2*exp(-x),y(x), singsol=all)
```

$$y(x) = e^{-x} \left(\frac{1}{60}x^5 + c_1 + c_2x + c_3x^2 \right)$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 34

```
DSolve[y'''[x]+3*y''[x]+3*y'[x]+y[x]==x^2*Exp[-x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{60}e^{-x}(x^5 + 60c_3x^2 + 60c_2x + 60c_1)$$

12 Chapter 3. Linear equations with variable coefficients. Page 108

12.1	problem 1(c.1)	900
12.2	problem 1(c.2)	935
12.3	problem 2	970

12.1 problem 1(c.1)

12.1.1 Existence and uniqueness analysis	901
12.1.2 Solving as second order euler ode ode	901
12.1.3 Solving as second order change of variable on x method 2 ode .	904
12.1.4 Solving as second order change of variable on x method 1 ode .	907
12.1.5 Solving as second order change of variable on y method 2 ode .	910
12.1.6 Solving as second order integrable as is ode	914
12.1.7 Solving as second order ode non constant coeff transformation on B ode	916
12.1.8 Solving as type second_order_integrable_as_is (not using ABC version)	919
12.1.9 Solving using Kovacic algorithm	922
12.1.10 Solving as exact linear second order ode ode	928
12.1.11 Maple step by step solution	932

Internal problem ID [6006]

Internal file name [OUTPUT/5254_Sunday_June_05_2022_03_28_40_PM_67813326/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY
1961

Section: Chapter 3. Linear equations with variable coefficients. Page 108

Problem number: 1(c.1).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_euler_ode", "exact linear second order ode", "second_order_integrable_as_is", "second_order_change_of_variable_on_x_method_1", "second_order_change_of_variable_on_x_method_2", "second_order_change_of_variable_on_y_method_2", "second_order_ode_non_constant_coeff_transformation_on_B"

Maple gives the following as the ode type

```
[[_2nd_order, _exact, _linear, _homogeneous]]
```

$$y'' + \frac{y'}{x} - \frac{y}{x^2} = 0$$

With initial conditions

$$[y(1) = 1, y'(1) = 0]$$

The ode can be written as

$$x^2y'' + xy' - y = 0$$

Which shows it is a Euler ODE.

12.1.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$\begin{aligned} p(x) &= \frac{1}{x} \\ q(x) &= -\frac{1}{x^2} \\ F &= 0 \end{aligned}$$

Hence the ode is

$$y'' + \frac{y'}{x} - \frac{y}{x^2} = 0$$

The domain of $p(x) = \frac{1}{x}$ is

$$\{x < 0 \vee 0 < x\}$$

And the point $x_0 = 1$ is inside this domain. The domain of $q(x) = -\frac{1}{x^2}$ is

$$\{x < 0 \vee 0 < x\}$$

And the point $x_0 = 1$ is also inside this domain. Hence solution exists and is unique.

12.1.2 Solving as second order euler ode ode

This is Euler second order ODE. Let the solution be $y = x^r$, then $y' = rx^{r-1}$ and $y'' = r(r-1)x^{r-2}$. Substituting these back into the given ODE gives

$$x^2(r(r-1))x^{r-2} + xx^{r-1} - x^r = 0$$

Simplifying gives

$$r(r - 1)x^r + rx^r - x^r = 0$$

Since $x^r \neq 0$ then dividing throughout by x^r gives

$$r(r - 1) + r - 1 = 0$$

Or

$$r^2 - 1 = 0 \tag{1}$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = -1$$

$$r_2 = 1$$

Since the roots are real and distinct, then the general solution is

$$y = c_1y_1 + c_2y_2$$

Where $y_1 = x^{r_1}$ and $y_2 = x^{r_2}$. Hence

$$y = \frac{c_1}{x} + c_2x$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = \frac{c_1}{x} + c_2x \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 1$ in the above gives

$$1 = c_1 + c_2 \tag{1A}$$

Taking derivative of the solution gives

$$y' = -\frac{c_1}{x^2} + c_2$$

substituting $y' = 0$ and $x = 1$ in the above gives

$$0 = -c_1 + c_2 \tag{2A}$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = \frac{1}{2}$$
$$c_2 = \frac{1}{2}$$

Substituting these values back in above solution results in

$$y = \frac{x^2 + 1}{2x}$$

Summary

The solution(s) found are the following

$$y = \frac{x^2 + 1}{2x} \tag{1}$$




Figure 158: Solution plot

Verification of solutions

$$y = \frac{x^2 + 1}{2x}$$

Verified OK.

12.1.3 Solving as second order change of variable on x method 2 ode

In normal form the ode

$$x^2 y'' + xy' - y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = -\frac{1}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x)\tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned} \tau &= \int e^{-(\int p(x)dx)} dx \\ &= \int e^{-(\int \frac{1}{x} dx)} dx \\ &= \int e^{-\ln(x)} dx \\ &= \int \frac{1}{x} dx \\ &= \ln(x) \end{aligned} \quad (6)$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned} q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\ &= \frac{-\frac{1}{x^2}}{\frac{1}{x^2}} \\ &= -1 \end{aligned} \tag{7}$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned} \frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) - y(\tau) &= 0 \end{aligned}$$

The above ode is now solved for $y(\tau)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(\tau) + By'(\tau) + Cy(\tau) = 0$$

Where in the above $A = 1, B = 0, C = -1$. Let the solution be $y(\tau) = e^{\lambda\tau}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda\tau} - e^{\lambda\tau} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda\tau}$ gives

$$\lambda^2 - 1 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = -1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(-1)} \\ &= \pm 1 \end{aligned}$$

Hence

$$\lambda_1 = +1$$

$$\lambda_2 = -1$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= 1 \\ \lambda_2 &= -1\end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned}y(\tau) &= c_1 e^{\lambda_1 \tau} + c_2 e^{\lambda_2 \tau} \\ y(\tau) &= c_1 e^{(1)\tau} + c_2 e^{(-1)\tau}\end{aligned}$$

Or

$$y(\tau) = c_1 e^{\tau} + c_2 e^{-\tau}$$

The above solution is now transformed back to y using (6) which results in

$$y = \frac{c_1 x^2 + c_2}{x}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = \frac{c_1 x^2 + c_2}{x} \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 1$ in the above gives

$$1 = c_1 + c_2 \tag{1A}$$

Taking derivative of the solution gives

$$y' = 2c_1 - \frac{c_1 x^2 + c_2}{x^2}$$

substituting $y' = 0$ and $x = 1$ in the above gives

$$0 = c_1 - c_2 \tag{2A}$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$\begin{aligned}c_1 &= \frac{1}{2} \\ c_2 &= \frac{1}{2}\end{aligned}$$

Substituting these values back in above solution results in

$$y = \frac{x^2 + 1}{2x}$$

Summary

The solution(s) found are the following

$$y = \frac{x^2 + 1}{2x} \tag{1}$$




Figure 159: Solution plot

Verification of solutions

$$y = \frac{x^2 + 1}{2x}$$

Verified OK.

12.1.4 Solving as second order change of variable on x method 1 ode

In normal form the ode

$$x^2 y'' + xy' - y = 0 \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \tag{2}$$

Where

$$p(x) = \frac{1}{x}$$

$$q(x) = -\frac{1}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) results

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $q_1 = c^2$ where c is some constant. Therefore from (5)

$$\tau' = \frac{1}{c}\sqrt{q}$$

$$= \frac{\sqrt{-\frac{1}{x^2}}}{c} \quad (6)$$

$$\tau'' = \frac{1}{c\sqrt{-\frac{1}{x^2}}x^3}$$

Substituting the above into (4) results in

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2}$$

$$= \frac{\frac{1}{c\sqrt{-\frac{1}{x^2}}x^3} + \frac{1}{x}\frac{\sqrt{-\frac{1}{x^2}}}{c}}{\left(\frac{\sqrt{-\frac{1}{x^2}}}{c}\right)^2}$$

$$= 0$$

Therefore ode (3) now becomes

$$y(\tau)'' + p_1y(\tau)' + q_1y(\tau) = 0$$

$$\frac{d^2}{d\tau^2}y(\tau) + c^2y(\tau) = 0 \quad (7)$$

The above ode is now solved for $y(\tau)$. Since the ode is now constant coefficients, it can be easily solved to give

$$y(\tau) = c_1 \cos(c\tau) + c_2 \sin(c\tau)$$

Now from (6)

$$\begin{aligned} \tau &= \int \frac{1}{c} \sqrt{q} dx \\ &= \frac{\int \sqrt{-\frac{1}{x^2}} dx}{c} \\ &= \frac{\sqrt{-\frac{1}{x^2}} x \ln(x)}{c} \end{aligned}$$

Substituting the above into the solution obtained gives

$$y = \frac{(ic_2 + c_1)x^2 - ic_2 + c_1}{2x}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = \frac{(ic_2 + c_1)x^2 - ic_2 + c_1}{2x} \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 1$ in the above gives

$$1 = c_1 \tag{1A}$$

Taking derivative of the solution gives

$$y' = ic_2 + c_1 - \frac{(ic_2 + c_1)x^2 - ic_2 + c_1}{2x^2}$$

substituting $y' = 0$ and $x = 1$ in the above gives

$$0 = ic_2 \tag{2A}$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 1$$

$$c_2 = 0$$

Substituting these values back in above solution results in

$$y = \frac{x^2 + 1}{2x}$$

Summary

The solution(s) found are the following

$$y = \frac{x^2 + 1}{2x} \tag{1}$$




Figure 160: Solution plot

Verification of solutions

$$y = \frac{x^2 + 1}{2x}$$

Verified OK.

12.1.5 Solving as second order change of variable on y method 2 ode

In normal form the ode

$$x^2 y'' + xy' - y = 0 \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \tag{2}$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = -\frac{1}{x^2}$$

Applying change of variables on the dependent variable $y = v(x)x^n$ to (2) gives the following ode where the dependent variables is $v(x)$ and not y .

$$v''(x) + \left(\frac{2n}{x} + p\right)v'(x) + \left(\frac{n(n-1)}{x^2} + \frac{np}{x} + q\right)v(x) = 0 \quad (3)$$

Let the coefficient of $v(x)$ above be zero. Hence

$$\frac{n(n-1)}{x^2} + \frac{np}{x} + q = 0 \quad (4)$$

Substituting the earlier values found for $p(x)$ and $q(x)$ into (4) gives

$$\frac{n(n-1)}{x^2} + \frac{n}{x^2} - \frac{1}{x^2} = 0 \quad (5)$$

Solving (5) for n gives

$$n = 1 \quad (6)$$

Substituting this value in (3) gives

$$v''(x) + \frac{3v'(x)}{x} = 0$$
$$v''(x) + \frac{3v'(x)}{x} = 0 \quad (7)$$

Using the substitution

$$u(x) = v'(x)$$

Then (7) becomes

$$u'(x) + \frac{3u(x)}{x} = 0 \quad (8)$$

The above is now solved for $u(x)$. In canonical form the ODE is

$$\begin{aligned}u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{3u}{x}\end{aligned}$$

Where $f(x) = -\frac{3}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u} du &= -\frac{3}{x} dx \\ \int \frac{1}{u} du &= \int -\frac{3}{x} dx \\ \ln(u) &= -3 \ln(x) + c_1 \\ u &= e^{-3 \ln(x) + c_1} \\ &= \frac{c_1}{x^3}\end{aligned}$$

Now that $u(x)$ is known, then

$$\begin{aligned}v'(x) &= u(x) \\ v(x) &= \int u(x) dx + c_2 \\ &= -\frac{c_1}{2x^2} + c_2\end{aligned}$$

Hence

$$\begin{aligned}y &= v(x) x^n \\ &= \left(-\frac{c_1}{2x^2} + c_2\right) x \\ &= \left(-\frac{c_1}{2x^2} + c_2\right) x\end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = \left(-\frac{c_1}{2x^2} + c_2\right) x \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 1$ in the above gives

$$1 = -\frac{c_1}{2} + c_2 \tag{1A}$$

Taking derivative of the solution gives

$$y' = \frac{c_1}{2x^2} + c_2$$

substituting $y' = 0$ and $x = 1$ in the above gives

$$0 = \frac{c_1}{2} + c_2 \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = -1$$
$$c_2 = \frac{1}{2}$$

Substituting these values back in above solution results in

$$y = \frac{x^2 + 1}{2x}$$

Summary

The solution(s) found are the following

$$y = \frac{x^2 + 1}{2x} \quad (1)$$




Figure 161: Solution plot

Verification of solutions

$$y = \frac{x^2 + 1}{2x}$$

Verified OK.

12.1.6 Solving as second order integrable as is ode

Integrating both sides of the ODE w.r.t x gives

$$\int (x^2 y'' + x y' - y) dx = 0$$
$$x^2 y' - xy = c_1$$

Which is now solved for y .

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -\frac{1}{x}$$
$$q(x) = \frac{c_1}{x^2}$$

Hence the ode is

$$y' - \frac{y}{x} = \frac{c_1}{x^2}$$

The integrating factor μ is

$$\mu = e^{\int -\frac{1}{x} dx}$$
$$= \frac{1}{x}$$

The ode becomes

$$\frac{d}{dx}(\mu y) = (\mu) \left(\frac{c_1}{x^2} \right)$$
$$\frac{d}{dx} \left(\frac{y}{x} \right) = \left(\frac{1}{x} \right) \left(\frac{c_1}{x^2} \right)$$
$$d \left(\frac{y}{x} \right) = \left(\frac{c_1}{x^3} \right) dx$$

Integrating gives

$$\frac{y}{x} = \int \frac{c_1}{x^3} dx$$
$$\frac{y}{x} = -\frac{c_1}{2x^2} + c_2$$

Dividing both sides by the integrating factor $\mu = \frac{1}{x}$ results in

$$y = -\frac{c_1}{2x} + c_2x$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = -\frac{c_1}{2x} + c_2x \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 1$ in the above gives

$$1 = -\frac{c_1}{2} + c_2 \quad (1A)$$

Taking derivative of the solution gives

$$y' = \frac{c_1}{2x^2} + c_2$$

substituting $y' = 0$ and $x = 1$ in the above gives

$$0 = \frac{c_1}{2} + c_2 \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$\begin{aligned} c_1 &= -1 \\ c_2 &= \frac{1}{2} \end{aligned}$$

Substituting these values back in above solution results in

$$y = \frac{x^2 + 1}{2x}$$

Summary

The solution(s) found are the following

$$y = \frac{x^2 + 1}{2x} \quad (1)$$




Figure 162: Solution plot

Verification of solutions

$$y = \frac{x^2 + 1}{2x}$$

Verified OK.

12.1.7 Solving as second order ode non constant coeff transformation on B ode

Given an ode of the form

$$Ay'' + By' + Cy = F(x)$$

This method reduces the order ode the ODE by one by applying the transformation

$$y = Bv$$

This results in

$$\begin{aligned} y' &= B'v + v'B \\ y'' &= B''v + B'v' + v''B + v'B' \\ &= v''B + 2v' + B' + B''v \end{aligned}$$

And now the original ode becomes

$$\begin{aligned} A(v''B + 2v'B' + B''v) + B(B'v + v'B) + CBv &= 0 \\ ABv'' + (2AB' + B^2)v' + (AB'' + BB' + CB)v &= 0 \end{aligned} \tag{1}$$

If the term $AB'' + BB' + CB$ is zero, then this method works and can be used to solve

$$ABv'' + (2AB' + B^2)v' = 0$$

By Using $u = v'$ which reduces the order of the above ode to one. The new ode is

$$ABu' + (2AB' + B^2)u = 0$$

The above ode is first order ode which is solved for u . Now a new ode $v' = u$ is solved for v as first order ode. Then the final solution is obtain from $y = Bv$.

This method works only if the term $AB'' + BB' + CB$ is zero. The given ODE shows that

$$A = x^2$$

$$B = x$$

$$C = -1$$

$$F = 0$$

The above shows that for this ode

$$\begin{aligned} AB'' + BB' + CB &= (x^2)(0) + (x)(1) + (-1)(x) \\ &= 0 \end{aligned}$$

Hence the ode in v given in (1) now simplifies to

$$x^3v'' + (3x^2)v' = 0$$

Now by applying $v' = u$ the above becomes

$$x^2(u'(x)x + 3u(x)) = 0$$

Which is now solved for u . In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{3u}{x} \end{aligned}$$

Where $f(x) = -\frac{3}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned} \frac{1}{u} du &= -\frac{3}{x} dx \\ \int \frac{1}{u} du &= \int -\frac{3}{x} dx \\ \ln(u) &= -3 \ln(x) + c_1 \\ u &= e^{-3 \ln(x) + c_1} \\ &= \frac{c_1}{x^3} \end{aligned}$$

The ode for v now becomes

$$\begin{aligned}v' &= u \\ &= \frac{c_1}{x^3}\end{aligned}$$

Which is now solved for v . Integrating both sides gives

$$\begin{aligned}v(x) &= \int \frac{c_1}{x^3} dx \\ &= -\frac{c_1}{2x^2} + c_2\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y(x) &= Bv \\ &= (x) \left(-\frac{c_1}{2x^2} + c_2 \right) \\ &= \left(-\frac{c_1}{2x^2} + c_2 \right) x\end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = \left(-\frac{c_1}{2x^2} + c_2 \right) x \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 1$ in the above gives

$$1 = -\frac{c_1}{2} + c_2 \quad (1A)$$

Taking derivative of the solution gives

$$y' = \frac{c_1}{2x^2} + c_2$$

substituting $y' = 0$ and $x = 1$ in the above gives

$$0 = \frac{c_1}{2} + c_2 \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$\begin{aligned}c_1 &= -1 \\ c_2 &= \frac{1}{2}\end{aligned}$$

Substituting these values back in above solution results in

$$y = \frac{x^2 + 1}{2x}$$

Summary

The solution(s) found are the following

$$y = \frac{x^2 + 1}{2x} \tag{1}$$




Figure 163: Solution plot

Verification of solutions

$$y = \frac{x^2 + 1}{2x}$$

Verified OK.

12.1.8 Solving as type second_order_integrable_as_is (not using ABC version)

Writing the ode as

$$x^2 y'' + xy' - y = 0$$

Integrating both sides of the ODE w.r.t x gives

$$\int (x^2 y'' + x y' - y) dx = 0$$
$$x^2 y' - xy = c_1$$

Which is now solved for y .

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -\frac{1}{x}$$
$$q(x) = \frac{c_1}{x^2}$$

Hence the ode is

$$y' - \frac{y}{x} = \frac{c_1}{x^2}$$

The integrating factor μ is

$$\mu = e^{\int -\frac{1}{x} dx}$$
$$= \frac{1}{x}$$

The ode becomes

$$\frac{d}{dx}(\mu y) = (\mu) \left(\frac{c_1}{x^2} \right)$$
$$\frac{d}{dx} \left(\frac{y}{x} \right) = \left(\frac{1}{x} \right) \left(\frac{c_1}{x^2} \right)$$
$$d \left(\frac{y}{x} \right) = \left(\frac{c_1}{x^3} \right) dx$$

Integrating gives

$$\frac{y}{x} = \int \frac{c_1}{x^3} dx$$
$$\frac{y}{x} = -\frac{c_1}{2x^2} + c_2$$

Dividing both sides by the integrating factor $\mu = \frac{1}{x}$ results in

$$y = -\frac{c_1}{2x} + c_2 x$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = -\frac{c_1}{2x} + c_2x \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 1$ in the above gives

$$1 = -\frac{c_1}{2} + c_2 \quad (1A)$$

Taking derivative of the solution gives

$$y' = \frac{c_1}{2x^2} + c_2$$

substituting $y' = 0$ and $x = 1$ in the above gives

$$0 = \frac{c_1}{2} + c_2 \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$\begin{aligned} c_1 &= -1 \\ c_2 &= \frac{1}{2} \end{aligned}$$

Substituting these values back in above solution results in

$$y = \frac{x^2 + 1}{2x}$$

Summary

The solution(s) found are the following

$$y = \frac{x^2 + 1}{2x} \quad (1)$$




Figure 164: Solution plot

Verification of solutions

$$y = \frac{x^2 + 1}{2x}$$

Verified OK.

12.1.9 Solving using Kovacic algorithm

Writing the ode as

$$x^2 y'' + x y' - y = 0 \tag{1}$$

$$A y'' + B y' + C y = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= x^2 \\ B &= x \\ C &= -1 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = y e^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = r z(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} \tag{5}$$

$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{3}{4x^2} \tag{6}$$

Comparing the above to (5) shows that

$$s = 3$$

$$t = 4x^2$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{3}{4x^2} \right) z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 165: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = \frac{3}{4x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = \frac{3}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{3}{2} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = -\frac{1}{2} \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = \frac{3}{4x^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = \frac{3}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{3}{2} \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = -\frac{1}{2} \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{3}{4x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	$\frac{3}{2}$	$-\frac{1}{2}$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	$\frac{3}{2}$	$-\frac{1}{2}$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = -\frac{1}{2}$ then

$$\begin{aligned} d &= \alpha_\infty^- - (\alpha_{c_1}^-) \\ &= -\frac{1}{2} - \left(-\frac{1}{2}\right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{x - c_1} \right) + (-)[\sqrt{r}]_\infty \\ &= -\frac{1}{2x} + (-)(0) \\ &= -\frac{1}{2x} \\ &= -\frac{1}{2x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$(0) + 2\left(-\frac{1}{2x}\right)(0) + \left(\left(\frac{1}{2x^2}\right) + \left(-\frac{1}{2x}\right)^2 - \left(\frac{3}{4x^2}\right)\right) = 0$$

$$0 = 0$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= pe^{\int \omega dx} \\ &= e^{\int -\frac{1}{2x} dx} \\ &= \frac{1}{\sqrt{x}} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{x}{x^2} dx} \\ &= z_1 e^{-\frac{\ln(x)}{2}} \\ &= z_1 \left(\frac{1}{\sqrt{x}}\right) \end{aligned}$$

Which simplifies to

$$y_1 = \frac{1}{x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{x}{x^2} dx}}{(y_1)^2} dx \\&= y_1 \int \frac{e^{-\ln(x)}}{(y_1)^2} dx \\&= y_1 \left(\frac{x^2}{2} \right)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\&= c_1 \left(\frac{1}{x} \right) + c_2 \left(\frac{1}{x} \left(\frac{x^2}{2} \right) \right)\end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = \frac{c_1}{x} + \frac{c_2 x}{2} \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 1$ in the above gives

$$1 = c_1 + \frac{c_2}{2} \tag{1A}$$

Taking derivative of the solution gives

$$y' = -\frac{c_1}{x^2} + \frac{c_2}{2}$$

substituting $y' = 0$ and $x = 1$ in the above gives

$$0 = -c_1 + \frac{c_2}{2} \tag{2A}$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$\begin{aligned}c_1 &= \frac{1}{2} \\c_2 &= 1\end{aligned}$$

Substituting these values back in above solution results in

$$y = \frac{x^2 + 1}{2x}$$

Summary

The solution(s) found are the following

$$y = \frac{x^2 + 1}{2x} \tag{1}$$




Figure 165: Solution plot

Verification of solutions

$$y = \frac{x^2 + 1}{2x}$$

Verified OK.

12.1.10 Solving as exact linear second order ode

An ode of the form

$$p(x)y'' + q(x)y' + r(x)y = s(x)$$

is exact if

$$p''(x) - q'(x) + r(x) = 0 \tag{1}$$

For the given ode we have

$$\begin{aligned}p(x) &= x^2 \\q(x) &= x \\r(x) &= -1 \\s(x) &= 0\end{aligned}$$

Hence

$$\begin{aligned}p''(x) &= 2 \\q'(x) &= 1\end{aligned}$$

Therefore (1) becomes

$$2 - (1) + (-1) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(x)y' + (q(x) - p'(x))y)' = s(x)$$

Integrating gives

$$p(x)y' + (q(x) - p'(x))y = \int s(x) dx$$

Substituting the above values for p, q, r, s gives

$$x^2y' - xy = c_1$$

We now have a first order ode to solve which is

$$x^2y' - xy = c_1$$

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$\begin{aligned}p(x) &= -\frac{1}{x} \\q(x) &= \frac{c_1}{x^2}\end{aligned}$$

Hence the ode is

$$y' - \frac{y}{x} = \frac{c_1}{x^2}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int -\frac{1}{x} dx} \\ &= \frac{1}{x}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu) \left(\frac{c_1}{x^2}\right) \\ \frac{d}{dx}\left(\frac{y}{x}\right) &= \left(\frac{1}{x}\right) \left(\frac{c_1}{x^2}\right) \\ d\left(\frac{y}{x}\right) &= \left(\frac{c_1}{x^3}\right) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}\frac{y}{x} &= \int \frac{c_1}{x^3} dx \\ \frac{y}{x} &= -\frac{c_1}{2x^2} + c_2\end{aligned}$$

Dividing both sides by the integrating factor $\mu = \frac{1}{x}$ results in

$$y = -\frac{c_1}{2x} + c_2x$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = -\frac{c_1}{2x} + c_2x \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 1$ in the above gives

$$1 = -\frac{c_1}{2} + c_2 \tag{1A}$$

Taking derivative of the solution gives

$$y' = \frac{c_1}{2x^2} + c_2$$

substituting $y' = 0$ and $x = 1$ in the above gives

$$0 = \frac{c_1}{2} + c_2 \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$\begin{aligned} c_1 &= -1 \\ c_2 &= \frac{1}{2} \end{aligned}$$

Substituting these values back in above solution results in

$$y = \frac{x^2 + 1}{2x}$$

Summary

The solution(s) found are the following

$$y = \frac{x^2 + 1}{2x} \quad (1)$$




Figure 166: Solution plot

Verification of solutions

$$y = \frac{x^2 + 1}{2x}$$

Verified OK.

12.1.11 Maple step by step solution

Let's solve

$$\left[x^2 y'' + x y' - y = 0, y(1) = 1, y' \Big|_{\{x=1\}} = 0 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{y'}{x} + \frac{y}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{y'}{x} - \frac{y}{x^2} = 0$$

- Multiply by denominators of the ODE

$$x^2 y'' + x y' - y = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x , using the chain rule

$$y' = \left(\frac{d}{dt} y(t) \right) t'(x)$$

- Compute derivative

$$y' = \frac{\frac{d}{dt} y(t)}{x}$$

- Calculate the 2nd derivative of y with respect to x , using the chain rule

$$y'' = \left(\frac{d^2}{dt^2} y(t) \right) t'(x)^2 + t''(x) \left(\frac{d}{dt} y(t) \right)$$

- Compute derivative

$$y'' = \frac{\frac{d^2}{dt^2} y(t)}{x^2} - \frac{\frac{d}{dt} y(t)}{x^2}$$

Substitute the change of variables back into the ODE

$$x^2 \left(\frac{\frac{d^2}{dt^2} y(t)}{x^2} - \frac{\frac{d}{dt} y(t)}{x^2} \right) + \frac{d}{dt} y(t) - y(t) = 0$$

- Simplify

$$\frac{d^2}{dt^2} y(t) - y(t) = 0$$

- Characteristic polynomial of ODE

$$r^2 - 1 = 0$$

- Factor the characteristic polynomial

$$(r - 1)(r + 1) = 0$$

- Roots of the characteristic polynomial

$$r = (-1, 1)$$

- 1st solution of the ODE

$$y_1(t) = e^{-t}$$

- 2nd solution of the ODE

$$y_2(t) = e^t$$

- General solution of the ODE

$$y(t) = c_1 y_1(t) + c_2 y_2(t)$$

- Substitute in solutions

$$y(t) = c_1 e^{-t} + c_2 e^t$$

- Change variables back using $t = \ln(x)$

$$y = \frac{c_1}{x} + c_2 x$$

- Simplify

$$y = \frac{c_1}{x} + c_2 x$$

- Check validity of solution $y = \frac{c_1}{x} + c_2 x$

- Use initial condition $y(1) = 1$

$$1 = c_1 + c_2$$

- Compute derivative of the solution

$$y' = -\frac{c_1}{x^2} + c_2$$

- Use the initial condition $y' \Big|_{\{x=1\}} = 0$

$$0 = -c_1 + c_2$$

- Solve for c_1 and c_2

$$\left\{ c_1 = \frac{1}{2}, c_2 = \frac{1}{2} \right\}$$

- Substitute constant values into general solution and simplify

$$y = \frac{x}{2} + \frac{1}{2x}$$

- Solution to the IVP

$$y = \frac{x}{2} + \frac{1}{2x}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
<- LODE of Euler type successful`

```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 13

```
dsolve([diff(y(x),x$2)+1/x*diff(y(x),x)-1/x^2*y(x)=0,y(1) = 1, D(y)(1) = 0],y(x), singsol=all)
```

$$y(x) = \frac{1}{2x} + \frac{x}{2}$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 17

```
DSolve[{y'[x]+1/x*y'[x]-1/x^2*y[x]==0,{y[1]==1,y'[1]==0}},y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \frac{x^2 + 1}{2x}$$

12.2 problem 1(c.2)

12.2.1 Existence and uniqueness analysis	936
12.2.2 Solving as second order euler ode ode	936
12.2.3 Solving as second order change of variable on x method 2 ode .	939
12.2.4 Solving as second order change of variable on x method 1 ode .	942
12.2.5 Solving as second order change of variable on y method 2 ode .	945
12.2.6 Solving as second order integrable as is ode	949
12.2.7 Solving as second order ode non constant coeff transformation on B ode	951
12.2.8 Solving as type second_order_integrable_as_is (not using ABC version)	954
12.2.9 Solving using Kovacic algorithm	957
12.2.10 Solving as exact linear second order ode ode	963
12.2.11 Maple step by step solution	967

Internal problem ID [6007]

Internal file name [OUTPUT/5255_Sunday_June_05_2022_03_28_42_PM_16790640/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY
1961

Section: Chapter 3. Linear equations with variable coefficients. Page 108

Problem number: 1(c.2).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_euler_ode", "exact linear second order ode", "second_order_integrable_as_is", "second_order_change_of_variable_on_x_method_1", "second_order_change_of_variable_on_x_method_2", "second_order_change_of_variable_on_y_method_2", "second_order_ode_non_constant_coeff_transformation_on_B"

Maple gives the following as the ode type

[[_2nd_order, _exact, _linear, _homogeneous]]

$$y'' + \frac{y'}{x} - \frac{y}{x^2} = 0$$

With initial conditions

$$[y(1) = 0, y'(1) = 1]$$

The ode can be written as

$$x^2y'' + xy' - y = 0$$

Which shows it is a Euler ODE.

12.2.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$\begin{aligned} p(x) &= \frac{1}{x} \\ q(x) &= -\frac{1}{x^2} \\ F &= 0 \end{aligned}$$

Hence the ode is

$$y'' + \frac{y'}{x} - \frac{y}{x^2} = 0$$

The domain of $p(x) = \frac{1}{x}$ is

$$\{x < 0 \vee 0 < x\}$$

And the point $x_0 = 1$ is inside this domain. The domain of $q(x) = -\frac{1}{x^2}$ is

$$\{x < 0 \vee 0 < x\}$$

And the point $x_0 = 1$ is also inside this domain. Hence solution exists and is unique.

12.2.2 Solving as second order euler ode ode

This is Euler second order ODE. Let the solution be $y = x^r$, then $y' = rx^{r-1}$ and $y'' = r(r-1)x^{r-2}$. Substituting these back into the given ODE gives

$$x^2(r(r-1))x^{r-2} + xx^{r-1} - x^r = 0$$

Simplifying gives

$$r(r - 1)x^r + rx^r - x^r = 0$$

Since $x^r \neq 0$ then dividing throughout by x^r gives

$$r(r - 1) + r - 1 = 0$$

Or

$$r^2 - 1 = 0 \tag{1}$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = -1$$

$$r_2 = 1$$

Since the roots are real and distinct, then the general solution is

$$y = c_1y_1 + c_2y_2$$

Where $y_1 = x^{r_1}$ and $y_2 = x^{r_2}$. Hence

$$y = \frac{c_1}{x} + c_2x$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = \frac{c_1}{x} + c_2x \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 1$ in the above gives

$$0 = c_1 + c_2 \tag{1A}$$

Taking derivative of the solution gives

$$y' = -\frac{c_1}{x^2} + c_2$$

substituting $y' = 1$ and $x = 1$ in the above gives

$$1 = -c_1 + c_2 \tag{2A}$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = -\frac{1}{2}$$
$$c_2 = \frac{1}{2}$$

Substituting these values back in above solution results in

$$y = \frac{x^2 - 1}{2x}$$

Summary

The solution(s) found are the following

$$y = \frac{x^2 - 1}{2x} \tag{1}$$




Figure 167: Solution plot

Verification of solutions

$$y = \frac{x^2 - 1}{2x}$$

Verified OK.

12.2.3 Solving as second order change of variable on x method 2 ode

In normal form the ode

$$x^2 y'' + xy' - y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = -\frac{1}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x)\tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned} \tau &= \int e^{-(\int p(x)dx)} dx \\ &= \int e^{-(\int \frac{1}{x} dx)} dx \\ &= \int e^{-\ln(x)} dx \\ &= \int \frac{1}{x} dx \\ &= \ln(x) \end{aligned} \quad (6)$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned} q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\ &= \frac{-\frac{1}{x^2}}{\frac{1}{x^2}} \\ &= -1 \end{aligned} \tag{7}$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned} \frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) - y(\tau) &= 0 \end{aligned}$$

The above ode is now solved for $y(\tau)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(\tau) + By'(\tau) + Cy(\tau) = 0$$

Where in the above $A = 1, B = 0, C = -1$. Let the solution be $y(\tau) = e^{\lambda\tau}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda\tau} - e^{\lambda\tau} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda\tau}$ gives

$$\lambda^2 - 1 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = -1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(-1)} \\ &= \pm 1 \end{aligned}$$

Hence

$$\lambda_1 = +1$$

$$\lambda_2 = -1$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= 1 \\ \lambda_2 &= -1\end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned}y(\tau) &= c_1 e^{\lambda_1 \tau} + c_2 e^{\lambda_2 \tau} \\ y(\tau) &= c_1 e^{(1)\tau} + c_2 e^{(-1)\tau}\end{aligned}$$

Or

$$y(\tau) = c_1 e^{\tau} + c_2 e^{-\tau}$$

The above solution is now transformed back to y using (6) which results in

$$y = \frac{c_1 x^2 + c_2}{x}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = \frac{c_1 x^2 + c_2}{x} \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 1$ in the above gives

$$0 = c_1 + c_2 \tag{1A}$$

Taking derivative of the solution gives

$$y' = 2c_1 - \frac{c_1 x^2 + c_2}{x^2}$$

substituting $y' = 1$ and $x = 1$ in the above gives

$$1 = c_1 - c_2 \tag{2A}$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$\begin{aligned}c_1 &= \frac{1}{2} \\ c_2 &= -\frac{1}{2}\end{aligned}$$

Substituting these values back in above solution results in

$$y = \frac{x^2 - 1}{2x}$$

Summary

The solution(s) found are the following

$$y = \frac{x^2 - 1}{2x} \tag{1}$$




Figure 168: Solution plot

Verification of solutions

$$y = \frac{x^2 - 1}{2x}$$

Verified OK.

12.2.4 Solving as second order change of variable on x method 1 ode

In normal form the ode

$$x^2 y'' + xy' - y = 0 \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \tag{2}$$

Where

$$p(x) = \frac{1}{x}$$

$$q(x) = -\frac{1}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) results

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $q_1 = c^2$ where c is some constant. Therefore from (5)

$$\tau' = \frac{1}{c}\sqrt{q}$$

$$= \frac{\sqrt{-\frac{1}{x^2}}}{c} \quad (6)$$

$$\tau'' = \frac{1}{c\sqrt{-\frac{1}{x^2}}x^3}$$

Substituting the above into (4) results in

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2}$$

$$= \frac{\frac{1}{c\sqrt{-\frac{1}{x^2}}x^3} + \frac{1}{x}\frac{\sqrt{-\frac{1}{x^2}}}{c}}{\left(\frac{\sqrt{-\frac{1}{x^2}}}{c}\right)^2}$$

$$= 0$$

Therefore ode (3) now becomes

$$y(\tau)'' + p_1y(\tau)' + q_1y(\tau) = 0$$

$$\frac{d^2}{d\tau^2}y(\tau) + c^2y(\tau) = 0 \quad (7)$$

The above ode is now solved for $y(\tau)$. Since the ode is now constant coefficients, it can be easily solved to give

$$y(\tau) = c_1 \cos(c\tau) + c_2 \sin(c\tau)$$

Now from (6)

$$\begin{aligned} \tau &= \int \frac{1}{c} \sqrt{q} dx \\ &= \frac{\int \sqrt{-\frac{1}{x^2}} dx}{c} \\ &= \frac{\sqrt{-\frac{1}{x^2}} x \ln(x)}{c} \end{aligned}$$

Substituting the above into the solution obtained gives

$$y = \frac{(ic_2 + c_1)x^2 - ic_2 + c_1}{2x}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = \frac{(ic_2 + c_1)x^2 - ic_2 + c_1}{2x} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 1$ in the above gives

$$0 = c_1 \quad (1A)$$

Taking derivative of the solution gives

$$y' = ic_2 + c_1 - \frac{(ic_2 + c_1)x^2 - ic_2 + c_1}{2x^2}$$

substituting $y' = 1$ and $x = 1$ in the above gives

$$1 = ic_2 \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$\begin{aligned} c_1 &= 0 \\ c_2 &= -i \end{aligned}$$

Substituting these values back in above solution results in

$$y = \frac{x^2 - 1}{2x}$$

Summary

The solution(s) found are the following

$$y = \frac{x^2 - 1}{2x} \tag{1}$$




Figure 169: Solution plot

Verification of solutions

$$y = \frac{x^2 - 1}{2x}$$

Verified OK.

12.2.5 Solving as second order change of variable on y method 2 ode

In normal form the ode

$$x^2 y'' + xy' - y = 0 \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \tag{2}$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = -\frac{1}{x^2}$$

Applying change of variables on the dependent variable $y = v(x)x^n$ to (2) gives the following ode where the dependent variables is $v(x)$ and not y .

$$v''(x) + \left(\frac{2n}{x} + p\right)v'(x) + \left(\frac{n(n-1)}{x^2} + \frac{np}{x} + q\right)v(x) = 0 \quad (3)$$

Let the coefficient of $v(x)$ above be zero. Hence

$$\frac{n(n-1)}{x^2} + \frac{np}{x} + q = 0 \quad (4)$$

Substituting the earlier values found for $p(x)$ and $q(x)$ into (4) gives

$$\frac{n(n-1)}{x^2} + \frac{n}{x^2} - \frac{1}{x^2} = 0 \quad (5)$$

Solving (5) for n gives

$$n = 1 \quad (6)$$

Substituting this value in (3) gives

$$v''(x) + \frac{3v'(x)}{x} = 0$$
$$v''(x) + \frac{3v'(x)}{x} = 0 \quad (7)$$

Using the substitution

$$u(x) = v'(x)$$

Then (7) becomes

$$u'(x) + \frac{3u(x)}{x} = 0 \quad (8)$$

The above is now solved for $u(x)$. In canonical form the ODE is

$$\begin{aligned}u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{3u}{x}\end{aligned}$$

Where $f(x) = -\frac{3}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u} du &= -\frac{3}{x} dx \\ \int \frac{1}{u} du &= \int -\frac{3}{x} dx \\ \ln(u) &= -3 \ln(x) + c_1 \\ u &= e^{-3 \ln(x) + c_1} \\ &= \frac{c_1}{x^3}\end{aligned}$$

Now that $u(x)$ is known, then

$$\begin{aligned}v'(x) &= u(x) \\ v(x) &= \int u(x) dx + c_2 \\ &= -\frac{c_1}{2x^2} + c_2\end{aligned}$$

Hence

$$\begin{aligned}y &= v(x) x^n \\ &= \left(-\frac{c_1}{2x^2} + c_2\right) x \\ &= \left(-\frac{c_1}{2x^2} + c_2\right) x\end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = \left(-\frac{c_1}{2x^2} + c_2\right) x \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 1$ in the above gives

$$0 = -\frac{c_1}{2} + c_2 \tag{1A}$$

Taking derivative of the solution gives

$$y' = \frac{c_1}{2x^2} + c_2$$

substituting $y' = 1$ and $x = 1$ in the above gives

$$1 = \frac{c_1}{2} + c_2 \tag{2A}$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 1$$

$$c_2 = \frac{1}{2}$$

Substituting these values back in above solution results in

$$y = \frac{x^2 - 1}{2x}$$

Summary

The solution(s) found are the following

$$y = \frac{x^2 - 1}{2x} \tag{1}$$




Figure 170: Solution plot

Verification of solutions

$$y = \frac{x^2 - 1}{2x}$$

Verified OK.

12.2.6 Solving as second order integrable as is ode

Integrating both sides of the ODE w.r.t x gives

$$\int (x^2 y'' + x y' - y) dx = 0$$
$$x^2 y' - xy = c_1$$

Which is now solved for y .

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -\frac{1}{x}$$
$$q(x) = \frac{c_1}{x^2}$$

Hence the ode is

$$y' - \frac{y}{x} = \frac{c_1}{x^2}$$

The integrating factor μ is

$$\mu = e^{\int -\frac{1}{x} dx}$$
$$= \frac{1}{x}$$

The ode becomes

$$\frac{d}{dx}(\mu y) = (\mu) \left(\frac{c_1}{x^2} \right)$$
$$\frac{d}{dx} \left(\frac{y}{x} \right) = \left(\frac{1}{x} \right) \left(\frac{c_1}{x^2} \right)$$
$$d \left(\frac{y}{x} \right) = \left(\frac{c_1}{x^3} \right) dx$$

Integrating gives

$$\frac{y}{x} = \int \frac{c_1}{x^3} dx$$
$$\frac{y}{x} = -\frac{c_1}{2x^2} + c_2$$

Dividing both sides by the integrating factor $\mu = \frac{1}{x}$ results in

$$y = -\frac{c_1}{2x} + c_2x$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = -\frac{c_1}{2x} + c_2x \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 1$ in the above gives

$$0 = -\frac{c_1}{2} + c_2 \quad (1A)$$

Taking derivative of the solution gives

$$y' = \frac{c_1}{2x^2} + c_2$$

substituting $y' = 1$ and $x = 1$ in the above gives

$$1 = \frac{c_1}{2} + c_2 \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$\begin{aligned} c_1 &= 1 \\ c_2 &= \frac{1}{2} \end{aligned}$$

Substituting these values back in above solution results in

$$y = \frac{x^2 - 1}{2x}$$

Summary

The solution(s) found are the following

$$y = \frac{x^2 - 1}{2x} \quad (1)$$




Figure 171: Solution plot

Verification of solutions

$$y = \frac{x^2 - 1}{2x}$$

Verified OK.

12.2.7 Solving as second order ode non constant coeff transformation on B ode

Given an ode of the form

$$Ay'' + By' + Cy = F(x)$$

This method reduces the order ode the ODE by one by applying the transformation

$$y = Bv$$

This results in

$$\begin{aligned} y' &= B'v + v'B \\ y'' &= B''v + B'v' + v''B + v'B' \\ &= v''B + 2v' + B' + B''v \end{aligned}$$

And now the original ode becomes

$$\begin{aligned} A(v''B + 2v'B' + B''v) + B(B'v + v'B) + CBv &= 0 \\ ABv'' + (2AB' + B^2)v' + (AB'' + BB' + CB)v &= 0 \end{aligned} \tag{1}$$

If the term $AB'' + BB' + CB$ is zero, then this method works and can be used to solve

$$ABv'' + (2AB' + B^2)v' = 0$$

By Using $u = v'$ which reduces the order of the above ode to one. The new ode is

$$ABu' + (2AB' + B^2)u = 0$$

The above ode is first order ode which is solved for u . Now a new ode $v' = u$ is solved for v as first order ode. Then the final solution is obtain from $y = Bv$.

This method works only if the term $AB'' + BB' + CB$ is zero. The given ODE shows that

$$A = x^2$$

$$B = x$$

$$C = -1$$

$$F = 0$$

The above shows that for this ode

$$\begin{aligned} AB'' + BB' + CB &= (x^2)(0) + (x)(1) + (-1)(x) \\ &= 0 \end{aligned}$$

Hence the ode in v given in (1) now simplifies to

$$x^3v'' + (3x^2)v' = 0$$

Now by applying $v' = u$ the above becomes

$$x^2(u'(x)x + 3u(x)) = 0$$

Which is now solved for u . In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{3u}{x} \end{aligned}$$

Where $f(x) = -\frac{3}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned} \frac{1}{u} du &= -\frac{3}{x} dx \\ \int \frac{1}{u} du &= \int -\frac{3}{x} dx \\ \ln(u) &= -3 \ln(x) + c_1 \\ u &= e^{-3 \ln(x) + c_1} \\ &= \frac{c_1}{x^3} \end{aligned}$$

The ode for v now becomes

$$\begin{aligned}v' &= u \\ &= \frac{c_1}{x^3}\end{aligned}$$

Which is now solved for v . Integrating both sides gives

$$\begin{aligned}v(x) &= \int \frac{c_1}{x^3} dx \\ &= -\frac{c_1}{2x^2} + c_2\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y(x) &= Bv \\ &= (x) \left(-\frac{c_1}{2x^2} + c_2 \right) \\ &= \left(-\frac{c_1}{2x^2} + c_2 \right) x\end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = \left(-\frac{c_1}{2x^2} + c_2 \right) x \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 1$ in the above gives

$$0 = -\frac{c_1}{2} + c_2 \quad (1A)$$

Taking derivative of the solution gives

$$y' = \frac{c_1}{2x^2} + c_2$$

substituting $y' = 1$ and $x = 1$ in the above gives

$$1 = \frac{c_1}{2} + c_2 \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$\begin{aligned}c_1 &= 1 \\ c_2 &= \frac{1}{2}\end{aligned}$$

Substituting these values back in above solution results in

$$y = \frac{x^2 - 1}{2x}$$

Summary

The solution(s) found are the following

$$y = \frac{x^2 - 1}{2x} \tag{1}$$




Figure 172: Solution plot

Verification of solutions

$$y = \frac{x^2 - 1}{2x}$$

Verified OK.

12.2.8 Solving as type second_order_integrable_as_is (not using ABC version)

Writing the ode as

$$x^2 y'' + xy' - y = 0$$

Integrating both sides of the ODE w.r.t x gives

$$\int (x^2 y'' + x y' - y) dx = 0$$
$$x^2 y' - xy = c_1$$

Which is now solved for y .

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -\frac{1}{x}$$
$$q(x) = \frac{c_1}{x^2}$$

Hence the ode is

$$y' - \frac{y}{x} = \frac{c_1}{x^2}$$

The integrating factor μ is

$$\mu = e^{\int -\frac{1}{x} dx}$$
$$= \frac{1}{x}$$

The ode becomes

$$\frac{d}{dx}(\mu y) = (\mu) \left(\frac{c_1}{x^2} \right)$$
$$\frac{d}{dx} \left(\frac{y}{x} \right) = \left(\frac{1}{x} \right) \left(\frac{c_1}{x^2} \right)$$
$$d \left(\frac{y}{x} \right) = \left(\frac{c_1}{x^3} \right) dx$$

Integrating gives

$$\frac{y}{x} = \int \frac{c_1}{x^3} dx$$
$$\frac{y}{x} = -\frac{c_1}{2x^2} + c_2$$

Dividing both sides by the integrating factor $\mu = \frac{1}{x}$ results in

$$y = -\frac{c_1}{2x} + c_2 x$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = -\frac{c_1}{2x} + c_2x \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 1$ in the above gives

$$0 = -\frac{c_1}{2} + c_2 \quad (1A)$$

Taking derivative of the solution gives

$$y' = \frac{c_1}{2x^2} + c_2$$

substituting $y' = 1$ and $x = 1$ in the above gives

$$1 = \frac{c_1}{2} + c_2 \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$\begin{aligned} c_1 &= 1 \\ c_2 &= \frac{1}{2} \end{aligned}$$

Substituting these values back in above solution results in

$$y = \frac{x^2 - 1}{2x}$$

Summary

The solution(s) found are the following

$$y = \frac{x^2 - 1}{2x} \quad (1)$$




Figure 173: Solution plot

Verification of solutions

$$y = \frac{x^2 - 1}{2x}$$

Verified OK.

12.2.9 Solving using Kovacic algorithm

Writing the ode as

$$x^2 y'' + xy' - y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= x^2 \\ B &= x \\ C &= -1 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} \tag{5}$$

$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{3}{4x^2} \tag{6}$$

Comparing the above to (5) shows that

$$s = 3$$

$$t = 4x^2$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{3}{4x^2} \right) z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 167: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = \frac{3}{4x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = \frac{3}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{3}{2} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = -\frac{1}{2} \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = \frac{3}{4x^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = \frac{3}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{3}{2} \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = -\frac{1}{2} \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{3}{4x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	$\frac{3}{2}$	$-\frac{1}{2}$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	$\frac{3}{2}$	$-\frac{1}{2}$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = -\frac{1}{2}$ then

$$\begin{aligned} d &= \alpha_\infty^- - (\alpha_{c_1}^-) \\ &= -\frac{1}{2} - \left(-\frac{1}{2}\right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{x - c_1} \right) + (-)[\sqrt{r}]_\infty \\ &= -\frac{1}{2x} + (-)(0) \\ &= -\frac{1}{2x} \\ &= -\frac{1}{2x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$(0) + 2\left(-\frac{1}{2x}\right)(0) + \left(\left(\frac{1}{2x^2}\right) + \left(-\frac{1}{2x}\right)^2 - \left(\frac{3}{4x^2}\right)\right) = 0$$

$$0 = 0$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= pe^{\int \omega dx} \\ &= e^{\int -\frac{1}{2x} dx} \\ &= \frac{1}{\sqrt{x}} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{x}{x^2} dx} \\ &= z_1 e^{-\frac{\ln(x)}{2}} \\ &= z_1 \left(\frac{1}{\sqrt{x}}\right) \end{aligned}$$

Which simplifies to

$$y_1 = \frac{1}{x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{x}{x^2} dx}}{(y_1)^2} dx \\&= y_1 \int \frac{e^{-\ln(x)}}{(y_1)^2} dx \\&= y_1 \left(\frac{x^2}{2} \right)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\&= c_1 \left(\frac{1}{x} \right) + c_2 \left(\frac{1}{x} \left(\frac{x^2}{2} \right) \right)\end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = \frac{c_1}{x} + \frac{c_2 x}{2} \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 1$ in the above gives

$$0 = c_1 + \frac{c_2}{2} \tag{1A}$$

Taking derivative of the solution gives

$$y' = -\frac{c_1}{x^2} + \frac{c_2}{2}$$

substituting $y' = 1$ and $x = 1$ in the above gives

$$1 = -c_1 + \frac{c_2}{2} \tag{2A}$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$\begin{aligned}c_1 &= -\frac{1}{2} \\c_2 &= 1\end{aligned}$$

Substituting these values back in above solution results in

$$y = \frac{x^2 - 1}{2x}$$

Summary

The solution(s) found are the following

$$y = \frac{x^2 - 1}{2x} \tag{1}$$




Figure 174: Solution plot

Verification of solutions

$$y = \frac{x^2 - 1}{2x}$$

Verified OK.

12.2.10 Solving as exact linear second order ode

An ode of the form

$$p(x) y'' + q(x) y' + r(x) y = s(x)$$

is exact if

$$p''(x) - q'(x) + r(x) = 0 \tag{1}$$

For the given ode we have

$$\begin{aligned}p(x) &= x^2 \\q(x) &= x \\r(x) &= -1 \\s(x) &= 0\end{aligned}$$

Hence

$$\begin{aligned}p''(x) &= 2 \\q'(x) &= 1\end{aligned}$$

Therefore (1) becomes

$$2 - (1) + (-1) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(x)y' + (q(x) - p'(x))y)' = s(x)$$

Integrating gives

$$p(x)y' + (q(x) - p'(x))y = \int s(x) dx$$

Substituting the above values for p, q, r, s gives

$$x^2y' - xy = c_1$$

We now have a first order ode to solve which is

$$x^2y' - xy = c_1$$

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$\begin{aligned}p(x) &= -\frac{1}{x} \\q(x) &= \frac{c_1}{x^2}\end{aligned}$$

Hence the ode is

$$y' - \frac{y}{x} = \frac{c_1}{x^2}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int -\frac{1}{x} dx} \\ &= \frac{1}{x}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu) \left(\frac{c_1}{x^2}\right) \\ \frac{d}{dx}\left(\frac{y}{x}\right) &= \left(\frac{1}{x}\right) \left(\frac{c_1}{x^2}\right) \\ d\left(\frac{y}{x}\right) &= \left(\frac{c_1}{x^3}\right) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}\frac{y}{x} &= \int \frac{c_1}{x^3} dx \\ \frac{y}{x} &= -\frac{c_1}{2x^2} + c_2\end{aligned}$$

Dividing both sides by the integrating factor $\mu = \frac{1}{x}$ results in

$$y = -\frac{c_1}{2x} + c_2x$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = -\frac{c_1}{2x} + c_2x \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 1$ in the above gives

$$0 = -\frac{c_1}{2} + c_2 \tag{1A}$$

Taking derivative of the solution gives

$$y' = \frac{c_1}{2x^2} + c_2$$

substituting $y' = 1$ and $x = 1$ in the above gives

$$1 = \frac{c_1}{2} + c_2 \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$\begin{aligned} c_1 &= 1 \\ c_2 &= \frac{1}{2} \end{aligned}$$

Substituting these values back in above solution results in

$$y = \frac{x^2 - 1}{2x}$$

Summary

The solution(s) found are the following

$$y = \frac{x^2 - 1}{2x} \quad (1)$$




Figure 175: Solution plot

Verification of solutions

$$y = \frac{x^2 - 1}{2x}$$

Verified OK.

12.2.11 Maple step by step solution

Let's solve

$$\left[x^2 y'' + xy' - y = 0, y(1) = 0, y'|_{\{x=1\}} = 1 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{y'}{x} + \frac{y}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{y'}{x} - \frac{y}{x^2} = 0$$

- Multiply by denominators of the ODE

$$x^2 y'' + xy' - y = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x , using the chain rule

$$y' = \left(\frac{d}{dt} y(t) \right) t'(x)$$

- Compute derivative

$$y' = \frac{\frac{d}{dt} y(t)}{x}$$

- Calculate the 2nd derivative of y with respect to x , using the chain rule

$$y'' = \left(\frac{d^2}{dt^2} y(t) \right) t'(x)^2 + t''(x) \left(\frac{d}{dt} y(t) \right)$$

- Compute derivative

$$y'' = \frac{\frac{d^2}{dt^2} y(t)}{x^2} - \frac{\frac{d}{dt} y(t)}{x^2}$$

Substitute the change of variables back into the ODE

$$x^2 \left(\frac{\frac{d^2}{dt^2} y(t)}{x^2} - \frac{\frac{d}{dt} y(t)}{x^2} \right) + \frac{d}{dt} y(t) - y(t) = 0$$

- Simplify

$$\frac{d^2}{dt^2} y(t) - y(t) = 0$$

- Characteristic polynomial of ODE

$$r^2 - 1 = 0$$

- Factor the characteristic polynomial

$$(r - 1)(r + 1) = 0$$

- Roots of the characteristic polynomial

$$r = (-1, 1)$$

- 1st solution of the ODE

$$y_1(t) = e^{-t}$$

- 2nd solution of the ODE

$$y_2(t) = e^t$$

- General solution of the ODE

$$y(t) = c_1 y_1(t) + c_2 y_2(t)$$

- Substitute in solutions

$$y(t) = c_1 e^{-t} + c_2 e^t$$

- Change variables back using $t = \ln(x)$

$$y = \frac{c_1}{x} + c_2 x$$

- Simplify

$$y = \frac{c_1}{x} + c_2 x$$

- Check validity of solution $y = \frac{c_1}{x} + c_2 x$

- Use initial condition $y(1) = 0$

$$0 = c_1 + c_2$$

- Compute derivative of the solution

$$y' = -\frac{c_1}{x^2} + c_2$$

- Use the initial condition $y' \Big|_{\{x=1\}} = 1$

$$1 = -c_1 + c_2$$

- Solve for c_1 and c_2

$$\left\{ c_1 = -\frac{1}{2}, c_2 = \frac{1}{2} \right\}$$

- Substitute constant values into general solution and simplify

$$y = \frac{x}{2} - \frac{1}{2x}$$

- Solution to the IVP

$$y = \frac{x}{2} - \frac{1}{2x}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
<- LODE of Euler type successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 13

```
dsolve([diff(y(x),x$2)+1/x*diff(y(x),x)-1/x^2*y(x)=0,y(1) = 0, D(y)(1) = 1],y(x), singsol=all)
```

$$y(x) = -\frac{1}{2x} + \frac{x}{2}$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 17

```
DSolve[{y'[x]+1/x*y'[x]-1/x^2*y[x]==0,{y[1]==0,y'[1]==1}},y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \frac{x^2 - 1}{2x}$$

12.3 problem 2

12.3.1 Solving as second order change of variable on x method 2 ode .	971
12.3.2 Solving as second order change of variable on x method 1 ode .	973
12.3.3 Solving as second order integrable as is ode	975
12.3.4 Solving as second order ode non constant coeff transformation on B ode	977
12.3.5 Solving as type second_order_integrable_as_is (not using ABC version)	979
12.3.6 Solving using Kovacic algorithm	981
12.3.7 Solving as exact linear second order ode ode	986
12.3.8 Maple step by step solution	988

Internal problem ID [6008]

Internal file name [OUTPUT/5256_Sunday_June_05_2022_03_28_43_PM_3532610/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 108

Problem number: 2.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "exact linear second order ode", "second_order_integrable_as_is", "second_order_change_of_variable_on_x_method_1", "second_order_change_of_variable_on_x_method_2", "second_order_ode_non_constant_coeff_transformation_on_B"

Maple gives the following as the ode type

```
[[_2nd_order , _exact , _linear , _homogeneous]]
```

$$(3x - 1)^2 y'' + (9x - 3) y' - 9y = 0$$

12.3.1 Solving as second order change of variable on x method 2 ode

In normal form the ode

$$9\left(-\frac{1}{3} + x\right)^2 y'' + (9x - 3)y' - 9y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{9x - 3}{9\left(-\frac{1}{3} + x\right)^2}$$
$$q(x) = -\frac{9}{(3x - 1)^2}$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x)\tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned} \tau &= \int e^{-(\int p(x)dx)} dx \\ &= \int e^{-\left(\int \frac{9x-3}{9\left(-\frac{1}{3}+x\right)^2} dx\right)} dx \\ &= \int e^{-\ln(3x-1)} dx \\ &= \int \frac{1}{3x-1} dx \\ &= \frac{\ln(3x-1)}{3} \end{aligned} \quad (6)$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned} q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\ &= \frac{-\frac{9}{(3x-1)^2}}{\frac{1}{(3x-1)^2}} \\ &= -9 \end{aligned} \tag{7}$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned} \frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) - 9y(\tau) &= 0 \end{aligned}$$

The above ode is now solved for $y(\tau)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(\tau) + By'(\tau) + Cy(\tau) = 0$$

Where in the above $A = 1, B = 0, C = -9$. Let the solution be $y(\tau) = e^{\lambda\tau}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda\tau} - 9 e^{\lambda\tau} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda\tau}$ gives

$$\lambda^2 - 9 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = -9$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(-9)} \\ &= \pm 3 \end{aligned}$$

Hence

$$\lambda_1 = +3$$

$$\lambda_2 = -3$$

Which simplifies to

$$\lambda_1 = 3$$

$$\lambda_2 = -3$$

Since roots are real and distinct, then the solution is

$$y(\tau) = c_1 e^{\lambda_1 \tau} + c_2 e^{\lambda_2 \tau}$$

$$y(\tau) = c_1 e^{(3)\tau} + c_2 e^{(-3)\tau}$$

Or

$$y(\tau) = c_1 e^{3\tau} + c_2 e^{-3\tau}$$

The above solution is now transformed back to y using (6) which results in

$$y = \frac{9\left(-\frac{1}{3} + x\right)^2 c_1 + c_2}{3x - 1}$$

Summary

The solution(s) found are the following

$$y = \frac{9\left(-\frac{1}{3} + x\right)^2 c_1 + c_2}{3x - 1} \quad (1)$$

Verification of solutions

$$y = \frac{9\left(-\frac{1}{3} + x\right)^2 c_1 + c_2}{3x - 1}$$

Verified OK.

12.3.2 Solving as second order change of variable on x method 1 ode

In normal form the ode

$$9\left(-\frac{1}{3} + x\right)^2 y'' + (9x - 3)y' - 9y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{3}{3x-1}$$

$$q(x) = -\frac{9}{(3x-1)^2}$$

Applying change of variables $\tau = g(x)$ to (2) results

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $q_1 = c^2$ where c is some constant. Therefore from (5)

$$\tau' = \frac{1}{c}\sqrt{q}$$

$$= \frac{3\sqrt{-\frac{1}{(3x-1)^2}}}{c} \quad (6)$$

$$\tau'' = \frac{9}{c\sqrt{-\frac{1}{(3x-1)^2}}(3x-1)^3}$$

Substituting the above into (4) results in

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2}$$

$$= \frac{\frac{9}{c\sqrt{-\frac{1}{(3x-1)^2}}(3x-1)^3} + \frac{3}{3x-1}\frac{3\sqrt{-\frac{1}{(3x-1)^2}}}{c}}{\left(\frac{3\sqrt{-\frac{1}{(3x-1)^2}}}{c}\right)^2}$$

$$= 0$$

Therefore ode (3) now becomes

$$y(\tau)'' + p_1y(\tau)' + q_1y(\tau) = 0$$

$$\frac{d^2}{d\tau^2}y(\tau) + c^2y(\tau) = 0 \quad (7)$$

The above ode is now solved for $y(\tau)$. Since the ode is now constant coefficients, it can be easily solved to give

$$y(\tau) = c_1 \cos(c\tau) + c_2 \sin(c\tau)$$

Now from (6)

$$\begin{aligned} \tau &= \int \frac{1}{c} \sqrt{q} dx \\ &= \frac{\int 3 \sqrt{-\frac{1}{(3x-1)^2}} dx}{c} \\ &= \frac{\sqrt{-\frac{1}{(3x-1)^2}} (3x-1) \ln(3x-1)}{c} \end{aligned}$$

Substituting the above into the solution obtained gives

$$y = \frac{(6c_1x - 2c_1) \cosh(\ln(3x-1)) + 9ic_2x(x - \frac{2}{3})}{6x-2}$$

Summary

The solution(s) found are the following

$$y = \frac{(6c_1x - 2c_1) \left(\frac{3x}{2} - \frac{1}{2} + \frac{1}{6x-2}\right) + 9ic_2x(x - \frac{2}{3})}{6x-2} \quad (1)$$

Verification of solutions

$$y = \frac{(6c_1x - 2c_1) \left(\frac{3x}{2} - \frac{1}{2} + \frac{1}{6x-2}\right) + 9ic_2x(x - \frac{2}{3})}{6x-2}$$

Verified OK.

12.3.3 Solving as second order integrable as is ode

Integrating both sides of the ODE w.r.t x gives

$$\begin{aligned} \int \left(9 \left(-\frac{1}{3} + x \right)^2 y'' + (9x-3) y' - 9y \right) dx &= 0 \\ (-9x+3)y + (9x^2-6x+1)y' &= c_1 \end{aligned}$$

Which is now solved for y .

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -\frac{3}{3x-1}$$
$$q(x) = \frac{c_1}{9\left(-\frac{1}{3} + x\right)^2}$$

Hence the ode is

$$y' - \frac{3y}{3x-1} = \frac{c_1}{9\left(-\frac{1}{3} + x\right)^2}$$

The integrating factor μ is

$$\mu = e^{\int -\frac{3}{3x-1} dx}$$
$$= \frac{1}{3x-1}$$

The ode becomes

$$\frac{d}{dx}(\mu y) = (\mu) \left(\frac{c_1}{9\left(-\frac{1}{3} + x\right)^2} \right)$$
$$\frac{d}{dx} \left(\frac{y}{3x-1} \right) = \left(\frac{1}{3x-1} \right) \left(\frac{c_1}{9\left(-\frac{1}{3} + x\right)^2} \right)$$
$$d \left(\frac{y}{3x-1} \right) = \left(\frac{c_1}{(3x-1)^3} \right) dx$$

Integrating gives

$$\frac{y}{3x-1} = \int \frac{c_1}{(3x-1)^3} dx$$
$$\frac{y}{3x-1} = -\frac{c_1}{6(3x-1)^2} + c_2$$

Dividing both sides by the integrating factor $\mu = \frac{1}{3x-1}$ results in

$$y = -\frac{c_1}{6(3x-1)} + c_2(3x-1)$$

Summary

The solution(s) found are the following

$$y = -\frac{c_1}{6(3x-1)} + c_2(3x-1) \quad (1)$$

Verification of solutions

$$y = -\frac{c_1}{6(3x-1)} + c_2(3x-1)$$

Verified OK.

12.3.4 Solving as second order ode non constant coeff transformation on B ode

Given an ode of the form

$$Ay'' + By' + Cy = F(x)$$

This method reduces the order ode the ODE by one by applying the transformation

$$y = Bv$$

This results in

$$\begin{aligned}y' &= B'v + v'B \\y'' &= B''v + B'v' + v''B + v'B' \\&= v''B + 2v' + B' + B''v\end{aligned}$$

And now the original ode becomes

$$\begin{aligned}A(v''B + 2v'B' + B''v) + B(B'v + v'B) + CBv &= 0 \\ABv'' + (2AB' + B^2)v' + (AB'' + BB' + CB)v &= 0\end{aligned}\tag{1}$$

If the term $AB'' + BB' + CB$ is zero, then this method works and can be used to solve

$$ABv'' + (2AB' + B^2)v' = 0$$

By Using $u = v'$ which reduces the order of the above ode to one. The new ode is

$$ABu' + (2AB' + B^2)u = 0$$

The above ode is first order ode which is solved for u . Now a new ode $v' = u$ is solved for v as first order ode. Then the final solution is obtain from $y = Bv$.

This method works only if the term $AB'' + BB' + CB$ is zero. The given ODE shows that

$$\begin{aligned}A &= 9\left(-\frac{1}{3} + x\right)^2 \\B &= 9x - 3 \\C &= -9 \\F &= 0\end{aligned}$$

The above shows that for this ode

$$\begin{aligned} AB'' + BB' + CB &= \left(9\left(-\frac{1}{3} + x\right)^2\right)(0) + (9x - 3)(9) + (-9)(9x - 3) \\ &= 0 \end{aligned}$$

Hence the ode in v given in (1) now simplifies to

$$3(3x - 1)^3 v'' + (27(3x - 1)^2) v' = 0$$

Now by applying $v' = u$ the above becomes

$$3(3x - 1)^3 u'(x) + 27(3x - 1)^2 u(x) = 0$$

Which is now solved for u . In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{9u}{3x - 1} \end{aligned}$$

Where $f(x) = -\frac{9}{3x-1}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned} \frac{1}{u} du &= -\frac{9}{3x - 1} dx \\ \int \frac{1}{u} du &= \int -\frac{9}{3x - 1} dx \\ \ln(u) &= -3 \ln(3x - 1) + c_1 \\ u &= e^{-3 \ln(3x-1)+c_1} \\ &= \frac{c_1}{(3x - 1)^3} \end{aligned}$$

The ode for v now becomes

$$\begin{aligned} v' &= u \\ &= \frac{c_1}{(3x - 1)^3} \end{aligned}$$

Which is now solved for v . Integrating both sides gives

$$\begin{aligned} v(x) &= \int \frac{c_1}{(3x - 1)^3} dx \\ &= -\frac{c_1}{6(3x - 1)^2} + c_2 \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y(x) &= Bv \\&= (9x - 3) \left(-\frac{c_1}{6(3x - 1)^2} + c_2 \right) \\&= \frac{54\left(-\frac{1}{3} + x\right)^2 c_2 - c_1}{6x - 2}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{54\left(-\frac{1}{3} + x\right)^2 c_2 - c_1}{6x - 2} \quad (1)$$

Verification of solutions

$$y = \frac{54\left(-\frac{1}{3} + x\right)^2 c_2 - c_1}{6x - 2}$$

Verified OK.

12.3.5 Solving as type second_order_integrable_as_is (not using ABC version)

Writing the ode as

$$9\left(-\frac{1}{3} + x\right)^2 y'' + (9x - 3)y' - 9y = 0$$

Integrating both sides of the ODE w.r.t x gives

$$\begin{aligned}\int \left(9\left(-\frac{1}{3} + x\right)^2 y'' + (9x - 3)y' - 9y \right) dx &= 0 \\(-9x + 3)y + (9x^2 - 6x + 1)y' &= c_1\end{aligned}$$

Which is now solved for y .

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$\begin{aligned}p(x) &= -\frac{3}{3x - 1} \\q(x) &= \frac{c_1}{9\left(-\frac{1}{3} + x\right)^2}\end{aligned}$$

Hence the ode is

$$y' - \frac{3y}{3x-1} = \frac{c_1}{9\left(-\frac{1}{3} + x\right)^2}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int -\frac{3}{3x-1} dx} \\ &= \frac{1}{3x-1}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu) \left(\frac{c_1}{9\left(-\frac{1}{3} + x\right)^2} \right) \\ \frac{d}{dx} \left(\frac{y}{3x-1} \right) &= \left(\frac{1}{3x-1} \right) \left(\frac{c_1}{9\left(-\frac{1}{3} + x\right)^2} \right) \\ d \left(\frac{y}{3x-1} \right) &= \left(\frac{c_1}{(3x-1)^3} \right) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}\frac{y}{3x-1} &= \int \frac{c_1}{(3x-1)^3} dx \\ \frac{y}{3x-1} &= -\frac{c_1}{6(3x-1)^2} + c_2\end{aligned}$$

Dividing both sides by the integrating factor $\mu = \frac{1}{3x-1}$ results in

$$y = -\frac{c_1}{6(3x-1)} + c_2(3x-1)$$

Summary

The solution(s) found are the following

$$y = -\frac{c_1}{6(3x-1)} + c_2(3x-1) \tag{1}$$

Verification of solutions

$$y = -\frac{c_1}{6(3x-1)} + c_2(3x-1)$$

Verified OK.

12.3.6 Solving using Kovacic algorithm

Writing the ode as

$$9\left(-\frac{1}{3} + x\right)^2 y'' + (9x - 3)y' - 9y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 9\left(-\frac{1}{3} + x\right)^2 \\ B &= 9x - 3 \\ C &= -9 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{27}{4(3x - 1)^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 27 \\ t &= 4(3x - 1)^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{27}{4(3x - 1)^2}\right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 169: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4(3x - 1)^2$. There is a pole at $x = \frac{1}{3}$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = \frac{3}{4\left(-\frac{1}{3} + x\right)^2}$$

For the pole at $x = \frac{1}{3}$ let b be the coefficient of $\frac{1}{\left(-\frac{1}{3} + x\right)^2}$ in the partial fractions decomposition of r given above. Therefore $b = \frac{3}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{3}{2} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = -\frac{1}{2} \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = \frac{27}{4(3x - 1)^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = \frac{3}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{3}{2} \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = -\frac{1}{2} \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{27}{4(3x - 1)^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
$\frac{1}{3}$	2	0	$\frac{3}{2}$	$-\frac{1}{2}$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	$\frac{3}{2}$	$-\frac{1}{2}$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to

determine possible non negative integer d from these using

$$d = \alpha_{\infty}^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_{∞}^{\pm} . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_{\infty}^{-} = -\frac{1}{2}$ then

$$\begin{aligned} d &= \alpha_{\infty}^{-} - (\alpha_{c_1}^{-}) \\ &= -\frac{1}{2} - \left(-\frac{1}{2}\right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_{\infty}$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^{-}}{x - c_1} \right) + (-)[\sqrt{r}]_{\infty} \\ &= -\frac{1}{2\left(-\frac{1}{3} + x\right)} + (-)(0) \\ &= -\frac{1}{2\left(-\frac{1}{3} + x\right)} \\ &= -\frac{3}{6x - 2} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$(0) + 2\left(-\frac{1}{2\left(-\frac{1}{3} + x\right)}\right)(0) + \left(\left(\frac{1}{2\left(-\frac{1}{3} + x\right)^2}\right) + \left(-\frac{1}{2\left(-\frac{1}{3} + x\right)}\right)^2 - \left(\frac{27}{4(3x - 1)^2}\right)\right) = 0$$

$0 = 0$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= pe^{\int \omega dx} \\ &= e^{\int -\frac{1}{2(-\frac{1}{3}+x)} dx} \\ &= \frac{1}{\sqrt{3x-1}} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{9x-3}{9(-\frac{1}{3}+x)^2} dx} \\ &= z_1 e^{-\frac{\ln(3x-1)}{2}} \\ &= z_1 \left(\frac{1}{\sqrt{3x-1}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = \frac{1}{3x-1}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{9x-3}{9(-\frac{1}{3}+x)^2} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-\ln(3x-1)}}{(y_1)^2} dx \\ &= y_1 \left(\frac{3}{2} x^2 - x \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}
y &= c_1 y_1 + c_2 y_2 \\
&= c_1 \left(\frac{1}{3x-1} \right) + c_2 \left(\frac{1}{3x-1} \left(\frac{3}{2} x^2 - x \right) \right)
\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1}{3x-1} + \frac{c_2 x(3x-2)}{6x-2} \quad (1)$$

Verification of solutions

$$y = \frac{c_1}{3x-1} + \frac{c_2 x(3x-2)}{6x-2}$$

Verified OK.

12.3.7 Solving as exact linear second order ode

An ode of the form

$$p(x) y'' + q(x) y' + r(x) y = s(x)$$

is exact if

$$p''(x) - q'(x) + r(x) = 0 \quad (1)$$

For the given ode we have

$$p(x) = 9 \left(-\frac{1}{3} + x \right)^2$$

$$q(x) = 9x - 3$$

$$r(x) = -9$$

$$s(x) = 0$$

Hence

$$p''(x) = 18$$

$$q'(x) = 9$$

Therefore (1) becomes

$$18 - (9) + (-9) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(x)y' + (q(x) - p'(x))y)' = s(x)$$

Integrating gives

$$p(x)y' + (q(x) - p'(x))y = \int s(x) dx$$

Substituting the above values for p, q, r, s gives

$$9\left(-\frac{1}{3} + x\right)^2 y' + (-9x + 3)y = c_1$$

We now have a first order ode to solve which is

$$9\left(-\frac{1}{3} + x\right)^2 y' + (-9x + 3)y = c_1$$

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -\frac{3}{3x-1}$$
$$q(x) = \frac{c_1}{(3x-1)^2}$$

Hence the ode is

$$y' - \frac{3y}{3x-1} = \frac{c_1}{(3x-1)^2}$$

The integrating factor μ is

$$\mu = e^{\int -\frac{3}{3x-1} dx}$$
$$= \frac{1}{3x-1}$$

The ode becomes

$$\frac{d}{dx}(\mu y) = (\mu) \left(\frac{c_1}{(3x-1)^2} \right)$$
$$\frac{d}{dx} \left(\frac{y}{3x-1} \right) = \left(\frac{1}{3x-1} \right) \left(\frac{c_1}{(3x-1)^2} \right)$$
$$d \left(\frac{y}{3x-1} \right) = \left(\frac{c_1}{(3x-1)^3} \right) dx$$

Integrating gives

$$\frac{y}{3x-1} = \int \frac{c_1}{(3x-1)^3} dx$$
$$\frac{y}{3x-1} = -\frac{c_1}{6(3x-1)^2} + c_2$$

Dividing both sides by the integrating factor $\mu = \frac{1}{3x-1}$ results in

$$y = -\frac{c_1}{6(3x-1)} + c_2(3x-1)$$

Summary

The solution(s) found are the following

$$y = -\frac{c_1}{6(3x-1)} + c_2(3x-1) \quad (1)$$

Verification of solutions

$$y = -\frac{c_1}{6(3x-1)} + c_2(3x-1)$$

Verified OK.

12.3.8 Maple step by step solution

Let's solve

$$9\left(-\frac{1}{3} + x\right)^2 y'' + (9x-3)y' - 9y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{3y'}{3x-1} + \frac{9y}{(3x-1)^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{3y'}{3x-1} - \frac{9y}{(3x-1)^2} = 0$$

- Check to see if $x_0 = \frac{1}{3}$ is a regular singular point

- Define functions

$$\left[P_2(x) = \frac{3}{3x-1}, P_3(x) = -\frac{9}{(3x-1)^2} \right]$$

- $\left(-\frac{1}{3} + x\right) \cdot P_2(x)$ is analytic at $x = \frac{1}{3}$

$$\left(\left(-\frac{1}{3} + x\right) \cdot P_2(x) \right) \Big|_{x=\frac{1}{3}} = 1$$

- $\left(-\frac{1}{3} + x\right)^2 \cdot P_3(x)$ is analytic at $x = \frac{1}{3}$

$$\left(\left(-\frac{1}{3} + x\right)^2 \cdot P_3(x) \right) \Big|_{x=\frac{1}{3}} = -1$$

- $x = \frac{1}{3}$ is a regular singular point

Check to see if $x_0 = \frac{1}{3}$ is a regular singular point

$$x_0 = \frac{1}{3}$$

- Multiply by denominators

$$(3x - 1)^2 y'' + (9x - 3) y' - 9y = 0$$

- Change variables using $x = u + \frac{1}{3}$ so that the regular singular point is at $u = 0$

$$9u^2 \left(\frac{d^2}{du^2} y(u) \right) + 9u \left(\frac{d}{du} y(u) \right) - 9y(u) = 0$$

- Assume series solution for $y(u)$

$$y(u) = \sum_{k=0}^{\infty} a_k u^{k+r}$$

- Rewrite DE with series expansions

- Convert $u \cdot \left(\frac{d}{du} y(u) \right)$ to series expansion

$$u \cdot \left(\frac{d}{du} y(u) \right) = \sum_{k=0}^{\infty} a_k (k+r) u^{k+r}$$

- Convert $u^2 \cdot \left(\frac{d^2}{du^2} y(u) \right)$ to series expansion

$$u^2 \cdot \left(\frac{d^2}{du^2} y(u) \right) = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) u^{k+r}$$

Rewrite DE with series expansions

$$\sum_{k=0}^{\infty} 9a_k (k+r+1)(k+r-1) u^{k+r} = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$r = 0$$

- Each term in the series must be 0, giving the recursion relation

$$(9k^2 - 9) a_k = 0$$

- Recursion relation that defines series solution to ODE

$$a_k = 0$$

- Recursion relation for $r = 0$
 $a_k = 0$
- Solution for $r = 0$
$$\left[y(u) = \sum_{k=0}^{\infty} a_k u^k, a_k = 0 \right]$$
- Revert the change of variables $u = -\frac{1}{3} + x$
$$\left[y = \sum_{k=0}^{\infty} a_k \left(-\frac{1}{3} + x\right)^k, a_k = 0 \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
<- LODE of Euler type successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
dsolve((3*x-1)^2*diff(y(x),x$2)+(9*x-3)*diff(y(x),x)-9*y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{9\left(x - \frac{1}{3}\right)^2 c_2 + 9c_1}{9x - 3}$$

✓ Solution by Mathematica

Time used: 0.031 (sec). Leaf size: 39

```
DSolve[(3*x-1)^2*y'[x]+(9*x-3)*y'[x]-9*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_1(-9x^2 + 6x - 2) - 3ic_2x(3x - 2)}{6x - 2}$$

13 Chapter 3. Linear equations with variable coefficients. Page 121

13.1	problem 1(a)	992
13.2	problem 1(b)	996
13.3	problem 1(c)	1000
13.4	problem 1(d)	1004
13.5	problem 1(e)	1009
13.6	problem 1(f)	1014
13.7	problem 2	1018

13.1 problem 1(a)

13.1.1 Maple step by step solution 993

Internal problem ID [6009]

Internal file name [OUTPUT/5257_Sunday_June_05_2022_03_28_44_PM_50954804/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 121

Problem number: 1(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "reduction_of_order", "second_order_euler_ode", "second_order_change_of_variable_on_x_method_1", "second_order_change_of_variable_on_x_method_2", "second_order_change_of_variable_on_y_method_2"

Maple gives the following as the ode type

```
[[_Emden, _Fowler]]
```

$$x^2y'' - 7xy' + 15y = 0$$

Given that one solution of the ode is

$$y_1 = x^3$$

Given one basis solution $y_1(x)$, then the second basis solution is given by

$$y_2(x) = y_1 \left(\int \frac{e^{-\int p dx}}{y_1^2} dx \right)$$

Where $p(x)$ is the coefficient of y' when the ode is written in the normal form

$$y'' + p(x)y' + q(x)y = f(x)$$

Looking at the ode to solve shows that

$$p(x) = -\frac{7}{x}$$

Therefore

$$y_2(x) = x^3 \left(\int \frac{e^{-\left(\int -\frac{7}{x} dx\right)}}{x^6} dx \right)$$

$$y_2(x) = x^3 \int \frac{x^7}{x^6} dx$$

$$y_2(x) = x^3 \left(\int x dx \right)$$

$$y_2(x) = \frac{x^5}{2}$$

Hence the solution is

$$\begin{aligned} y &= c_1 y_1(x) + c_2 y_2(x) \\ &= c_1 x^3 + \frac{1}{2} c_2 x^5 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 x^3 + \frac{1}{2} c_2 x^5 \quad (1)$$

Verification of solutions

$$y = c_1 x^3 + \frac{1}{2} c_2 x^5$$

Verified OK.

13.1.1 Maple step by step solution

Let's solve

$$x^2 y'' - 7xy' + 15y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = \frac{7y'}{x} - \frac{15y}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - \frac{7y'}{x} + \frac{15y}{x^2} = 0$$

- Multiply by denominators of the ODE

$$x^2 y'' - 7xy' + 15y = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x , using the chain rule

$$y' = \left(\frac{d}{dt} y(t) \right) t'(x)$$

- Compute derivative

$$y' = \frac{\frac{d}{dt} y(t)}{x}$$

- Calculate the 2nd derivative of y with respect to x , using the chain rule

$$y'' = \left(\frac{d^2}{dt^2} y(t) \right) t'(x)^2 + t''(x) \left(\frac{d}{dt} y(t) \right)$$

- Compute derivative

$$y'' = \frac{\frac{d^2}{dt^2} y(t)}{x^2} - \frac{\frac{d}{dt} y(t)}{x^2}$$

Substitute the change of variables back into the ODE

$$x^2 \left(\frac{\frac{d^2}{dt^2} y(t)}{x^2} - \frac{\frac{d}{dt} y(t)}{x^2} \right) - 7 \frac{d}{dt} y(t) + 15y(t) = 0$$

- Simplify

$$\frac{d^2}{dt^2} y(t) - 8 \frac{d}{dt} y(t) + 15y(t) = 0$$

- Characteristic polynomial of ODE

$$r^2 - 8r + 15 = 0$$

- Factor the characteristic polynomial

$$(r - 3)(r - 5) = 0$$

- Roots of the characteristic polynomial

$$r = (3, 5)$$

- 1st solution of the ODE

$$y_1(t) = e^{3t}$$

- 2nd solution of the ODE

$$y_2(t) = e^{5t}$$

- General solution of the ODE

$$y(t) = c_1 y_1(t) + c_2 y_2(t)$$

- Substitute in solutions

$$y(t) = c_1 e^{3t} + c_2 e^{5t}$$

- Change variables back using $t = \ln(x)$

$$y = c_2 x^5 + c_1 x^3$$

- Simplify

$$y = x^3(c_2 x^2 + c_1)$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
<- LODE of Euler type successful`

```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 15

```
dsolve([x^2*diff(y(x),x$2)-7*x*diff(y(x),x)+15*y(x)=0,x^3],singsol=all)
```

$$y(x) = x^3(c_1 x^2 + c_2)$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 18

```
DSolve[x^2*y''[x]-7*x*y'[x]+15*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^3(c_2 x^2 + c_1)$$

13.2 problem 1(b)

13.2.1 Maple step by step solution 997

Internal problem ID [6010]

Internal file name [OUTPUT/5258_Sunday_June_05_2022_03_28_46_PM_84153657/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 121

Problem number: 1(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "reduction_of_order", "second_order_euler_ode", "second_order_change_of_variable_on_x_method_1", "second_order_change_of_variable_on_x_method_2", "second_order_change_of_variable_on_y_method_2", "second_order_ode_non_constant_coeff_transformation_on_B"

Maple gives the following as the ode type

```
[[_Emden , _Fowler]]
```

$$x^2y'' - xy' + y = 0$$

Given that one solution of the ode is

$$y_1 = x$$

Given one basis solution $y_1(x)$, then the second basis solution is given by

$$y_2(x) = y_1 \left(\int \frac{e^{-\int p dx}}{y_1^2} dx \right)$$

Where $p(x)$ is the coefficient of y' when the ode is written in the normal form

$$y'' + p(x)y' + q(x)y = f(x)$$

Looking at the ode to solve shows that

$$p(x) = -\frac{1}{x}$$

Therefore

$$y_2(x) = x \left(\int \frac{e^{-(\int -\frac{1}{x} dx)}}{x^2} dx \right)$$

$$y_2(x) = x \int \frac{x}{x^2}, dx$$

$$y_2(x) = \left(\int \frac{1}{x} dx \right) x$$

$$y_2(x) = \ln(x) x$$

Hence the solution is

$$\begin{aligned} y &= c_1 y_1(x) + c_2 y_2(x) \\ &= c_1 x + c_2 \ln(x) x \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 x + c_2 \ln(x) x \tag{1}$$

Verification of solutions

$$y = c_1 x + c_2 \ln(x) x$$

Verified OK.

13.2.1 Maple step by step solution

Let's solve

$$x^2 y'' - xy' + y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = \frac{y'}{x} - \frac{y}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - \frac{y'}{x} + \frac{y}{x^2} = 0$$

- Multiply by denominators of the ODE

$$x^2 y'' - xy' + y = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x , using the chain rule

$$y' = \left(\frac{d}{dt} y(t) \right) t'(x)$$

- Compute derivative

$$y' = \frac{\frac{d}{dt} y(t)}{x}$$

- Calculate the 2nd derivative of y with respect to x , using the chain rule

$$y'' = \left(\frac{d^2}{dt^2} y(t) \right) t'(x)^2 + t''(x) \left(\frac{d}{dt} y(t) \right)$$

- Compute derivative

$$y'' = \frac{\frac{d^2}{dt^2} y(t)}{x^2} - \frac{\frac{d}{dt} y(t)}{x^2}$$

Substitute the change of variables back into the ODE

$$x^2 \left(\frac{\frac{d^2}{dt^2} y(t)}{x^2} - \frac{\frac{d}{dt} y(t)}{x^2} \right) - \frac{d}{dt} y(t) + y(t) = 0$$

- Simplify

$$\frac{d^2}{dt^2} y(t) - 2 \frac{d}{dt} y(t) + y(t) = 0$$

- Characteristic polynomial of ODE

$$r^2 - 2r + 1 = 0$$

- Factor the characteristic polynomial

$$(r - 1)^2 = 0$$

- Root of the characteristic polynomial

$$r = 1$$

- 1st solution of the ODE

$$y_1(t) = e^t$$

- Repeated root, multiply $y_1(t)$ by t to ensure linear independence

$$y_2(t) = t e^t$$

- General solution of the ODE

$$y(t) = c_1 y_1(t) + c_2 y_2(t)$$

- Substitute in solutions

$$y(t) = c_1 e^t + c_2 t e^t$$

- Change variables back using $t = \ln(x)$

$$y = c_1 x + c_2 \ln(x) x$$

- Simplify

$$y = x(c_1 + c_2 \ln(x))$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
<- LODE of Euler type successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve([x^2*diff(y(x),x$2)-x*diff(y(x),x)+y(x)=0,x],singsol=all)
```

$$y(x) = x(c_2 \ln(x) + c_1)$$

✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 15

```
DSolve[x^2*y''[x]-x*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x(c_2 \log(x) + c_1)$$

13.3 problem 1(c)

13.3.1 Maple step by step solution 1001

Internal problem ID [6011]

Internal file name [OUTPUT/5259_Sunday_June_05_2022_03_28_47_PM_54326568/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 121

Problem number: 1(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "reduction_of_order", "second_order_change_of_variable_on_y_method_1", "linear_second_order_ode_solved_by_an_integrating_factor"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - 4xy' + (4x^2 - 2)y = 0$$

Given that one solution of the ode is

$$y_1 = e^{x^2}$$

Given one basis solution $y_1(x)$, then the second basis solution is given by

$$y_2(x) = y_1 \left(\int \frac{e^{-\int p dx}}{y_1^2} dx \right)$$

Where $p(x)$ is the coefficient of y' when the ode is written in the normal form

$$y'' + p(x)y' + q(x)y = f(x)$$

Looking at the ode to solve shows that

$$p(x) = -4x$$

Therefore

$$y_2(x) = e^{x^2} \left(\int e^{-(\int -4x dx)} e^{-2x^2} dx \right)$$

$$y_2(x) = e^{x^2} \int \frac{e^{2x^2}}{e^{2x^2}} dx$$

$$y_2(x) = e^{x^2} \left(\int 1 dx \right)$$

$$y_2(x) = x e^{x^2}$$

Hence the solution is

$$\begin{aligned} y &= c_1 y_1(x) + c_2 y_2(x) \\ &= c_1 e^{x^2} + c_2 x e^{x^2} \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{x^2} + c_2 x e^{x^2} \quad (1)$$

Verification of solutions

$$y = c_1 e^{x^2} + c_2 x e^{x^2}$$

Verified OK.

13.3.1 Maple step by step solution

Let's solve

$$y'' - 4xy' + (4x^2 - 2)y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^k$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 0..2$

$$x^m \cdot y = \sum_{k=\max(0,-m)}^{\infty} a_k x^{k+m}$$

- Shift index using $k \rightarrow k - m$

$$x^m \cdot y = \sum_{k=\max(0, -m)+m}^{\infty} a_{k-m} x^k$$

- Convert $x \cdot y'$ to series expansion

$$x \cdot y' = \sum_{k=0}^{\infty} a_k k x^k$$

- Convert y'' to series expansion

$$y'' = \sum_{k=2}^{\infty} a_k k(k-1) x^{k-2}$$

- Shift index using $k \rightarrow k + 2$

$$y'' = \sum_{k=0}^{\infty} a_{k+2} (k+2)(k+1) x^k$$

Rewrite ODE with series expansions

$$2a_2 - 2a_0 + (6a_3 - 6a_1)x + \left(\sum_{k=2}^{\infty} (a_{k+2}(k+2)(k+1) - 2a_k(2k+1) + 4a_{k-2}) x^k \right) = 0$$

- The coefficients of each power of x must be 0

$$[2a_2 - 2a_0 = 0, 6a_3 - 6a_1 = 0]$$

- Solve for the dependent coefficient(s)

$$\{a_2 = a_0, a_3 = a_1\}$$

- Each term in the series must be 0, giving the recursion relation

$$(k^2 + 3k + 2) a_{k+2} - 4a_k k - 2a_k + 4a_{k-2} = 0$$

- Shift index using $k \rightarrow k + 2$

$$((k+2)^2 + 3k + 8) a_{k+4} - 4a_{k+2}(k+2) - 2a_{k+2} + 4a_k = 0$$

- Recursion relation that defines the series solution to the ODE

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+4} = \frac{2(2ka_{k+2} - 2a_k + 5a_{k-2})}{k^2 + 7k + 12}, a_2 = a_0, a_3 = a_1 \right]$$

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacics algorithm  
  A Liouvillian solution exists  
  Reducible group (found an exponential solution)  
<- Kovacics algorithm successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve([diff(y(x),x$2)-4*x*diff(y(x),x)+(4*x^2-2)*y(x)=0,exp(x^2)],singsol=all)
```

$$y(x) = e^{x^2}(c_2x + c_1)$$

✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 18

```
DSolve[y''[x]-4*x*y'[x]+(4*x^2-2)*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{x^2}(c_2x + c_1)$$

13.4 problem 1(d)

13.4.1 Maple step by step solution 1005

Internal problem ID [6012]

Internal file name [OUTPUT/5260_Sunday_June_05_2022_03_28_48_PM_65912661/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 121

Problem number: 1(d).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "reduction_of_order", "second_order_ode_non_constant_coeff_transformation_on_B"

Maple gives the following as the ode type

[_Laguerre]

$$xy'' - (1+x)y' + y = 0$$

Given that one solution of the ode is

$$y_1 = e^x$$

Given one basis solution $y_1(x)$, then the second basis solution is given by

$$y_2(x) = y_1 \left(\int \frac{e^{-\int p dx}}{y_1^2} dx \right)$$

Where $p(x)$ is the coefficient of y' when the ode is written in the normal form

$$y'' + p(x)y' + q(x)y = f(x)$$

Looking at the ode to solve shows that

$$p(x) = \frac{-1-x}{x}$$

Therefore

$$y_2(x) = e^x \left(\int e^{-\left(\int \frac{-1-x}{x} dx\right)} e^{-2x} dx \right)$$

$$y_2(x) = e^x \int \frac{e^{x+\ln(x)}}{e^{2x}} dx$$

$$y_2(x) = e^x \left(\int x e^{-x} dx \right)$$

$$y_2(x) = -e^x(1+x)e^{-x}$$

Hence the solution is

$$\begin{aligned} y &= c_1 y_1(x) + c_2 y_2(x) \\ &= c_1 e^x - c_2 e^x(1+x)e^{-x} \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^x - c_2 e^x(1+x)e^{-x} \quad (1)$$

Verification of solutions

$$y = c_1 e^x - c_2 e^x(1+x)e^{-x}$$

Verified OK.

13.4.1 Maple step by step solution

Let's solve

$$y''x + (-1-x)y' + y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{y}{x} + \frac{(1+x)y'}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - \frac{(1+x)y'}{x} + \frac{y}{x} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$[P_2(x) = -\frac{1+x}{x}, P_3(x) = \frac{1}{x}]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = -1$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 0$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$y''x + (-1 - x)y' + y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y'$ to series expansion for $m = 0..1$

$$x^m \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1+m}$$

- Shift index using $k \rightarrow k+1-m$

$$x^m \cdot y' = \sum_{k=-1+m}^{\infty} a_{k+1-m} (k+1-m+r) x^{k+r}$$

- Convert $x \cdot y''$ to series expansion

$$x \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r-1}$$

- Shift index using $k \rightarrow k+1$

$$x \cdot y'' = \sum_{k=-1}^{\infty} a_{k+1} (k+1+r)(k+r) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0 r(-2+r) x^{-1+r} + \left(\sum_{k=0}^{\infty} (a_{k+1} (k+1+r)(k+r-1) - a_k (k+r-1)) x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$r(-2 + r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \{0, 2\}$$

- Each term in the series must be 0, giving the recursion relation

$$(k + r - 1)(a_{k+1}(k + 1 + r) - a_k) = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+1} = \frac{a_k}{k+1+r}$$

- Recursion relation for $r = 0$

$$a_{k+1} = \frac{a_k}{k+1}$$

- Solution for $r = 0$

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+1} = \frac{a_k}{k+1} \right]$$

- Recursion relation for $r = 2$

$$a_{k+1} = \frac{a_k}{k+3}$$

- Solution for $r = 2$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+2}, a_{k+1} = \frac{a_k}{k+3} \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^k \right) + \left(\sum_{k=0}^{\infty} b_k x^{k+2} \right), a_{k+1} = \frac{a_k}{k+1}, b_{k+1} = \frac{b_k}{k+3} \right]$$

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacics algorithm  
  A Liouvillian solution exists  
  Reducible group (found an exponential solution)  
  Reducible group (found another exponential solution)  
<- Kovacics algorithm successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve([x*diff(y(x),x$2)-(x+1)*diff(y(x),x)+y(x)=0,exp(x)],singsol=all)
```

$$y(x) = e^x c_2 + c_1 x + c_1$$

✓ Solution by Mathematica

Time used: 0.029 (sec). Leaf size: 19

```
DSolve[x*y''[x]-(x+1)*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^x - c_2(x + 1)$$

13.5 problem 1(e)

13.5.1 Maple step by step solution 1010

Internal problem ID [6013]

Internal file name [OUTPUT/5261_Sunday_June_05_2022_03_28_49_PM_68004065/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 121

Problem number: 1(e).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "reduction_of_order", "second_order_change_of_variable_on_y_method_2", "second_order_ode_non_constant_coeff_transformation_on_B"

Maple gives the following as the ode type

[_Gegenbauer]

$$(-x^2 + 1)y'' - 2xy' + 2y = 0$$

Given that one solution of the ode is

$$y_1 = x$$

Given one basis solution $y_1(x)$, then the second basis solution is given by

$$y_2(x) = y_1 \left(\int \frac{e^{-\int p dx}}{y_1^2} dx \right)$$

Where $p(x)$ is the coefficient of y' when the ode is written in the normal form

$$y'' + p(x)y' + q(x)y = f(x)$$

Looking at the ode to solve shows that

$$p(x) = -\frac{2x}{-x^2 + 1}$$

Therefore

$$y_2(x) = x \left(\int \frac{e^{-\left(\int -\frac{2x}{x^2+1} dx\right)}}{x^2} dx \right)$$

$$y_2(x) = x \int \frac{e^{-\ln(x-1)-\ln(1+x)}}{x^2} dx$$

$$y_2(x) = x \left(\int \frac{1}{x^2(x^2-1)} dx \right)$$

$$y_2(x) = x \left(-\frac{\ln(1+x)}{2} + \frac{\ln(x-1)}{2} + \frac{1}{x} \right)$$

Hence the solution is

$$\begin{aligned} y &= c_1 y_1(x) + c_2 y_2(x) \\ &= c_1 x + c_2 x \left(-\frac{\ln(1+x)}{2} + \frac{\ln(x-1)}{2} + \frac{1}{x} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 x + c_2 x \left(-\frac{\ln(1+x)}{2} + \frac{\ln(x-1)}{2} + \frac{1}{x} \right) \quad (1)$$

Verification of solutions

$$y = c_1 x + c_2 x \left(-\frac{\ln(1+x)}{2} + \frac{\ln(x-1)}{2} + \frac{1}{x} \right)$$

Verified OK.

13.5.1 Maple step by step solution

Let's solve

$$(-x^2 + 1)y'' - 2xy' + 2y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{2xy'}{x^2-1} + \frac{2y}{x^2-1}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{2xy'}{x^2-1} - \frac{2y}{x^2-1} = 0$$

- Check to see if x_0 is a regular singular point

- Define functions

$$[P_2(x) = \frac{2x}{x^2-1}, P_3(x) = -\frac{2}{x^2-1}]$$

- $(1+x) \cdot P_2(x)$ is analytic at $x = -1$

$$((1+x) \cdot P_2(x)) \Big|_{x=-1} = 1$$

- $(1+x)^2 \cdot P_3(x)$ is analytic at $x = -1$

$$((1+x)^2 \cdot P_3(x)) \Big|_{x=-1} = 0$$

- $x = -1$ is a regular singular point

Check to see if x_0 is a regular singular point

$$x_0 = -1$$

- Multiply by denominators

$$(x^2 - 1)y'' + 2xy' - 2y = 0$$

- Change variables using $x = u - 1$ so that the regular singular point is at $u = 0$

$$(u^2 - 2u) \left(\frac{d^2}{du^2} y(u) \right) + (2u - 2) \left(\frac{d}{du} y(u) \right) - 2y(u) = 0$$

- Assume series solution for $y(u)$

$$y(u) = \sum_{k=0}^{\infty} a_k u^{k+r}$$

- Rewrite ODE with series expansions

- Convert $u^m \cdot \left(\frac{d}{du} y(u) \right)$ to series expansion for $m = 0..1$

$$u^m \cdot \left(\frac{d}{du} y(u) \right) = \sum_{k=0}^{\infty} a_k (k+r) u^{k+r-1+m}$$

- Shift index using $k \rightarrow k + 1 - m$

$$u^m \cdot \left(\frac{d}{du} y(u) \right) = \sum_{k=-1+m}^{\infty} a_{k+1-m} (k+1-m+r) u^{k+r}$$

- Convert $u^m \cdot \left(\frac{d^2}{du^2} y(u) \right)$ to series expansion for $m = 1..2$

$$u^m \cdot \left(\frac{d^2}{du^2} y(u) \right) = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) u^{k+r-2+m}$$

- Shift index using $k \rightarrow k + 2 - m$

$$u^m \cdot \left(\frac{d^2}{du^2} y(u) \right) = \sum_{k=-2+m}^{\infty} a_{k+2-m} (k+2-m+r) (k+1-m+r) u^{k+r}$$

Rewrite ODE with series expansions

$$-2a_0 r^2 u^{-1+r} + \left(\sum_{k=0}^{\infty} (-2a_{k+1} (k+1+r)^2 + a_k (k+r+2) (k+r-1)) u^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$-2r^2 = 0$$

- Values of r that satisfy the indicial equation

$$r = 0$$

- Each term in the series must be 0, giving the recursion relation

$$-2a_{k+1} (k+1)^2 + a_k (k+2) (k-1) = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+1} = \frac{a_k (k+2)(k-1)}{2(k+1)^2}$$

- Recursion relation for $r = 0$; series terminates at $k = 1$

$$a_{k+1} = \frac{a_k (k+2)(k-1)}{2(k+1)^2}$$

- Apply recursion relation for $k = 0$

$$a_1 = -a_0$$

- Terminating series solution of the ODE for $r = 0$. Use reduction of order to find the second li

$$y(u) = a_0 \cdot (-u + 1)$$

- Revert the change of variables $u = 1 + x$

$$[y = -a_0 x]$$

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacics algorithm  
  A Liouvillian solution exists  
  Reducible group (found an exponential solution)  
  Group is reducible, not completely reducible  
<- Kovacics algorithm successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve([(1-x^2)*diff(y(x),x$2)-2*x*diff(y(x),x)+2*y(x)=0,x],singsol=all)
```

$$y(x) = -\frac{c_2 \ln(x+1)x}{2} + \frac{c_2 \ln(x-1)x}{2} + c_1x + c_2$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 33

```
DSolve[(1-x^2)*y'[x]-2*x*y'[x]+2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1x - \frac{1}{2}c_2(x \log(1-x) - x \log(x+1) + 2)$$

13.6 problem 1(f)

13.6.1 Maple step by step solution 1016

Internal problem ID [6014]

Internal file name [OUTPUT/5262_Sunday_June_05_2022_03_28_50_PM_9372561/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 121

Problem number: 1(f).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "reduction_of_order", "second_order_change_of_variable_on_y_method_2", "second_order_ode_non_constant_coeff_transformation_on_B"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - 2xy' + 2y = 0$$

Given that one solution of the ode is

$$y_1 = x$$

Given one basis solution $y_1(x)$, then the second basis solution is given by

$$y_2(x) = y_1 \left(\int \frac{e^{-\int p dx}}{y_1^2} dx \right)$$

Where $p(x)$ is the coefficient of y' when the ode is written in the normal form

$$y'' + p(x)y' + q(x)y = f(x)$$

Looking at the ode to solve shows that

$$p(x) = -2x$$

Therefore

$$y_2(x) = x \left(\int \frac{e^{-(\int -2x dx)}}{x^2} dx \right)$$

$$y_2(x) = x \int \frac{e^{x^2}}{x^2} dx$$

$$y_2(x) = x \left(\int \frac{e^{x^2}}{x^2} dx \right)$$

$$y_2(x) = x \left(-\frac{e^{x^2}}{x} + \sqrt{\pi} \operatorname{erfi}(x) \right)$$

Hence the solution is

$$\begin{aligned} y &= c_1 y_1(x) + c_2 y_2(x) \\ &= c_1 x + c_2 x \left(-\frac{e^{x^2}}{x} + \sqrt{\pi} \operatorname{erfi}(x) \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 x + c_2 x \left(-\frac{e^{x^2}}{x} + \sqrt{\pi} \operatorname{erfi}(x) \right) \quad (1)$$

Verification of solutions

$$y = c_1 x + c_2 x \left(-\frac{e^{x^2}}{x} + \sqrt{\pi} \operatorname{erfi}(x) \right)$$

Verified OK.

13.6.1 Maple step by step solution

Let's solve

$$y'' - 2xy' + 2y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^k$$

- Rewrite DE with series expansions

- Convert $x \cdot y'$ to series expansion

$$x \cdot y' = \sum_{k=0}^{\infty} a_k k x^k$$

- Convert y'' to series expansion

$$y'' = \sum_{k=2}^{\infty} a_k k(k-1) x^{k-2}$$

- Shift index using $k- > k+2$

$$y'' = \sum_{k=0}^{\infty} a_{k+2}(k+2)(k+1) x^k$$

Rewrite DE with series expansions

$$\sum_{k=0}^{\infty} (a_{k+2}(k+2)(k+1) - 2a_k(k-1)) x^k = 0$$

- Each term in the series must be 0, giving the recursion relation

$$(k^2 + 3k + 2) a_{k+2} - 2a_k(k-1) = 0$$

- Recursion relation that defines the series solution to the ODE

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+2} = \frac{2a_k(k-1)}{k^2+3k+2} \right]$$

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacics algorithm  
  A Liouvillian solution exists  
  Reducible group (found an exponential solution)  
  Group is reducible, not completely reducible  
<- Kovacics algorithm successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 23

```
dsolve([diff(y(x),x$2)-2*x*diff(y(x),x)+2*y(x)=0,x],singsol=all)
```

$$y(x) = e^{x^2} c_2 + x(-\sqrt{\pi} c_2 \operatorname{erfi}(x) + c_1)$$

✓ Solution by Mathematica

Time used: 0.034 (sec). Leaf size: 43

```
DSolve[y''[x]-2*x*y'[x]+2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sqrt{\pi} c_2 \sqrt{x^2} \operatorname{erfi}(\sqrt{x^2}) + c_2 e^{x^2} + 2c_1 x$$

13.7 problem 2

13.7.1 Maple step by step solution 1020

Internal problem ID [6015]

Internal file name [OUTPUT/5263_Sunday_June_05_2022_03_28_52_PM_74182912/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 121

Problem number: 2.

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_ODE_non_constant_coefficients_of_type_Euler**"

Maple gives the following as the ode type

```
[[_3rd_order , _with_linear_symmetries]]
```

$$x^3y''' - 3x^2y'' + 6xy' - 6y = 0$$

This is Euler ODE of higher order. Let $y = x^\lambda$. Hence

$$y' = \lambda x^{\lambda-1}$$

$$y'' = \lambda(\lambda - 1) x^{\lambda-2}$$

$$y''' = \lambda(\lambda - 1)(\lambda - 2) x^{\lambda-3}$$

Substituting these back into

$$x^3y''' - 3x^2y'' + 6xy' - 6y = 0$$

gives

$$6x\lambda x^{\lambda-1} - 3x^2\lambda(\lambda - 1) x^{\lambda-2} + x^3\lambda(\lambda - 1)(\lambda - 2) x^{\lambda-3} - 6x^\lambda = 0$$

Which simplifies to

$$6\lambda x^\lambda - 3\lambda(\lambda - 1) x^\lambda + \lambda(\lambda - 1)(\lambda - 2) x^\lambda - 6x^\lambda = 0$$

And since $x^\lambda \neq 0$ then dividing through by x^λ , the above becomes

$$6\lambda - 3\lambda(\lambda - 1) + \lambda(\lambda - 1)(\lambda - 2) - 6 = 0$$

Simplifying gives the characteristic equation as

$$\lambda^3 - 6\lambda^2 + 11\lambda - 6 = 0$$

Solving the above gives the following roots

$$\lambda_1 = 1$$

$$\lambda_2 = 2$$

$$\lambda_3 = 3$$

This table summarises the result

root	multiplicity	type of root
1	1	real root
2	1	real root
3	1	real root

The solution is generated by going over the above table. For each real root λ of multiplicity one generates a c_1x^λ basis solution. Each real root of multiplicity two, generates c_1x^λ and $c_2x^\lambda \ln(x)$ basis solutions. Each real root of multiplicity three, generates c_1x^λ and $c_2x^\lambda \ln(x)$ and $c_3x^\lambda \ln(x)^2$ basis solutions, and so on. Each complex root $\alpha \pm i\beta$ of multiplicity one generates $x^\alpha(c_1 \cos(\beta \ln(x)) + c_2 \sin(\beta \ln(x)))$ basis solutions. And each complex root $\alpha \pm i\beta$ of multiplicity two generates $\ln(x) x^\alpha(c_1 \cos(\beta \ln(x)) + c_2 \sin(\beta \ln(x)))$ basis solutions. And each complex root $\alpha \pm i\beta$ of multiplicity three generates $\ln(x)^2 x^\alpha(c_1 \cos(\beta \ln(x)) + c_2 \sin(\beta \ln(x)))$ basis solutions. And so on. Using the above show that the solution is

$$y = c_3x^3 + c_2x^2 + c_1x$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = x$$

$$y_2 = x^2$$

$$y_3 = x^3$$

Summary

The solution(s) found are the following

$$y = c_3x^3 + c_2x^2 + c_1x \tag{1}$$

Verification of solutions

$$y = c_3x^3 + c_2x^2 + c_1x$$

Verified OK.

13.7.1 Maple step by step solution

Let's solve

$$x^3y''' - 3x^2y'' + 6xy' - 6y = 0$$

- Highest derivative means the order of the ODE is 3

$$y'''$$

- Isolate 3rd derivative

$$y''' = \frac{6y}{x^3} + \frac{3(y''x - 2y')}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y''' - \frac{3y''}{x} + \frac{6y'}{x^2} - \frac{6y}{x^3} = 0$$

- Multiply by denominators of the ODE

$$x^3y''' - 3x^2y'' + 6xy' - 6y = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x , using the chain rule

$$y' = \left(\frac{d}{dt}y(t)\right) t'(x)$$

- Compute derivative

$$y' = \frac{\frac{d}{dt}y(t)}{x}$$

- Calculate the 2nd derivative of y with respect to x , using the chain rule

$$y'' = \left(\frac{d^2}{dt^2}y(t)\right) t'(x)^2 + t''(x) \left(\frac{d}{dt}y(t)\right)$$

- Compute derivative

$$y'' = \frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2}$$

- Calculate the 3rd derivative of y with respect to x , using the chain rule

$$y''' = \left(\frac{d^3}{dt^3}y(t)\right) t'(x)^3 + 3t'(x)t''(x) \left(\frac{d^2}{dt^2}y(t)\right) + t'''(x) \left(\frac{d}{dt}y(t)\right)$$

- Compute derivative

$$y''' = \frac{\frac{d^3}{dt^3}y(t)}{x^3} - \frac{3\left(\frac{d^2}{dt^2}y(t)\right)}{x^3} + \frac{2\left(\frac{d}{dt}y(t)\right)}{x^3}$$

Substitute the change of variables back into the ODE

$$x^3 \left(\frac{\frac{d^3}{dt^3}y(t)}{x^3} - \frac{3\left(\frac{d^2}{dt^2}y(t)\right)}{x^3} + \frac{2\left(\frac{d}{dt}y(t)\right)}{x^3} \right) - 3x^2 \left(\frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2} \right) + 6\frac{d}{dt}y(t) - 6y(t) = 0$$

- Simplify

$$\frac{d^3}{dt^3}y(t) - 6\frac{d^2}{dt^2}y(t) + 11\frac{d}{dt}y(t) - 6y(t) = 0$$

- Convert linear ODE into a system of first order ODEs

- Define new variable $y_1(t)$

$$y_1(t) = y(t)$$

- Define new variable $y_2(t)$

$$y_2(t) = \frac{d}{dt}y(t)$$

- Define new variable $y_3(t)$

$$y_3(t) = \frac{d^2}{dt^2}y(t)$$

- Isolate for $\frac{d}{dt}y_3(t)$ using original ODE

$$\frac{d}{dt}y_3(t) = 6y_3(t) - 11y_2(t) + 6y_1(t)$$

Convert linear ODE into a system of first order ODEs

$$\left[y_2(t) = \frac{d}{dt}y_1(t), y_3(t) = \frac{d}{dt}y_2(t), \frac{d}{dt}y_3(t) = 6y_3(t) - 11y_2(t) + 6y_1(t) \right]$$

- Define vector

$$\vec{y}(t) = \begin{bmatrix} y_1(t) \\ y_2(t) \\ y_3(t) \end{bmatrix}$$

- System to solve

$$\frac{d}{dt}\vec{y}(t) = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 6 & -11 & 6 \end{bmatrix} \cdot \vec{y}(t)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 6 & -11 & 6 \end{bmatrix}$$

- Rewrite the system as

$$\frac{d}{dt} \vec{y}(t) = A \cdot \vec{y}(t)$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[1, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right], \left[2, \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right], \left[3, \begin{bmatrix} \frac{1}{9} \\ \frac{1}{3} \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[1, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_1 = e^t \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[2, \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_2 = e^{2t} \cdot \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[3, \begin{bmatrix} \frac{1}{9} \\ \frac{1}{3} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_3 = e^{3t} \cdot \begin{bmatrix} \frac{1}{9} \\ \frac{1}{3} \\ 1 \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{y} = c_1 \vec{y}_1 + c_2 \vec{y}_2 + c_3 \vec{y}_3$$

- Substitute solutions into the general solution

$$\vec{y} = c_1 e^t \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + c_2 e^{2t} \cdot \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} + c_3 e^{3t} \cdot \begin{bmatrix} \frac{1}{9} \\ \frac{1}{3} \\ 1 \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y(t) = c_1 e^t + \frac{c_2 e^{2t}}{4} + \frac{c_3 e^{3t}}{9}$$

- Change variables back using $t = \ln(x)$

$$y = c_1 x + \frac{1}{4} c_2 x^2 + \frac{1}{9} c_3 x^3$$

Maple trace

```

`Methods for third order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
<- LODE of Euler type successful`

```


✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve([x^3*diff(y(x),x$3)-3*x^2*diff(y(x),x$2)+6*x*diff(y(x),x)-6*y(x)=0,x],singsol=all)
```

$$y(x) = x(c_2x^2 + c_1x + c_3)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 19

```
DSolve[x^3*y'''[x]-3*x^2*y''[x]+6*x*y'[x]-6*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x(x(c_3x + c_2) + c_1)$$

14 Chapter 3. Linear equations with variable coefficients. Page 124

14.1 problem 1	1026
14.2 problem 2	1030
14.3 problem 3	1034

14.1 problem 1

14.1.1 Maple step by step solution 1027

Internal problem ID [6016]

Internal file name [OUTPUT/5264_Sunday_June_05_2022_03_28_53_PM_30888276/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 124

Problem number: 1.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "reduction_of_order", "second_order_euler_ode", "exact linear second order ode", "second_order_integrable_as_is"

Maple gives the following as the ode type

```
[[_2nd_order , _exact , _linear , _homogeneous]]
```

$$x^2y'' - 2y = 0$$

Given that one solution of the ode is

$$y_1 = x^2$$

Given one basis solution $y_1(x)$, then the second basis solution is given by

$$y_2(x) = y_1 \left(\int \frac{e^{-\int p dx}}{y_1^2} dx \right)$$

Where $p(x)$ is the coefficient of y' when the ode is written in the normal form

$$y'' + p(x)y' + q(x)y = f(x)$$

Looking at the ode to solve shows that

$$p(x) = 0$$

Therefore

$$y_2(x) = x^2 \left(\int \frac{e^{-(\int 0 dx)}}{x^4} dx \right)$$

$$y_2(x) = x^2 \int \frac{1}{x^4} dx$$

$$y_2(x) = x^2 \left(\int \frac{1}{x^4} dx \right)$$

$$y_2(x) = -\frac{1}{3x}$$

Hence the solution is

$$\begin{aligned} y &= c_1 y_1(x) + c_2 y_2(x) \\ &= c_1 x^2 - \frac{c_2}{3x} \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 x^2 - \frac{c_2}{3x} \tag{1}$$

Verification of solutions

$$y = c_1 x^2 - \frac{c_2}{3x}$$

Verified OK.

14.1.1 Maple step by step solution

Let's solve

$$x^2 y'' - 2y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = \frac{2y}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - \frac{2y}{x^2} = 0$$

- Multiply by denominators of the ODE

$$x^2 y'' - 2y = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x , using the chain rule

$$y' = \left(\frac{d}{dt} y(t) \right) t'(x)$$

- Compute derivative

$$y' = \frac{\frac{d}{dt} y(t)}{x}$$

- Calculate the 2nd derivative of y with respect to x , using the chain rule

$$y'' = \left(\frac{d^2}{dt^2} y(t) \right) t'(x)^2 + t''(x) \left(\frac{d}{dt} y(t) \right)$$

- Compute derivative

$$y'' = \frac{\frac{d^2}{dt^2} y(t)}{x^2} - \frac{\frac{d}{dt} y(t)}{x^2}$$

Substitute the change of variables back into the ODE

$$x^2 \left(\frac{\frac{d^2}{dt^2} y(t)}{x^2} - \frac{\frac{d}{dt} y(t)}{x^2} \right) - 2y(t) = 0$$

- Simplify

$$\frac{d^2}{dt^2} y(t) - \frac{d}{dt} y(t) - 2y(t) = 0$$

- Characteristic polynomial of ODE

$$r^2 - r - 2 = 0$$

- Factor the characteristic polynomial

$$(r + 1)(r - 2) = 0$$

- Roots of the characteristic polynomial

$$r = (-1, 2)$$

- 1st solution of the ODE

$$y_1(t) = e^{-t}$$

- 2nd solution of the ODE

$$y_2(t) = e^{2t}$$

- General solution of the ODE

$$y(t) = c_1 y_1(t) + c_2 y_2(t)$$

- Substitute in solutions

$$y(t) = c_1 e^{-t} + c_2 e^{2t}$$

- Change variables back using $t = \ln(x)$

$$y = \frac{c_1}{x} + c_2 x^2$$

- Simplify

$$y = \frac{c_1}{x} + c_2 x^2$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
<- LODE of Euler type successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve([x^2*diff(y(x),x$2)-2*y(x)=0,x^2],singsol=all)
```

$$y(x) = \frac{c_1 x^3 + c_2}{x}$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 18

```
DSolve[x^2*y''[x]-2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_2 x^3 + c_1}{x}$$

14.2 problem 2

14.2.1 Maple step by step solution 1031

Internal problem ID [6017]

Internal file name [OUTPUT/5265_Sunday_June_05_2022_03_28_54_PM_20156416/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 124

Problem number: 2.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "reduction_of_order", "second_order_euler_ode", "second_order_change_of_variable_on_x_method_1", "second_order_change_of_variable_on_x_method_2", "second_order_change_of_variable_on_y_method_2", "second_order_ode_non_constant_coeff_transformation_on_B"

Maple gives the following as the ode type

```
[[_Emden , _Fowler]]
```

$$x^2y'' - xy' + y = 0$$

Given that one solution of the ode is

$$y_1 = x$$

Given one basis solution $y_1(x)$, then the second basis solution is given by

$$y_2(x) = y_1 \left(\int \frac{e^{-\int p dx}}{y_1^2} dx \right)$$

Where $p(x)$ is the coefficient of y' when the ode is written in the normal form

$$y'' + p(x)y' + q(x)y = f(x)$$

Looking at the ode to solve shows that

$$p(x) = -\frac{1}{x}$$

Therefore

$$y_2(x) = x \left(\int \frac{e^{-(\int -\frac{1}{x} dx)}}{x^2} dx \right)$$

$$y_2(x) = x \int \frac{x}{x^2} dx$$

$$y_2(x) = \left(\int \frac{1}{x} dx \right) x$$

$$y_2(x) = \ln(x) x$$

Hence the solution is

$$\begin{aligned} y &= c_1 y_1(x) + c_2 y_2(x) \\ &= c_1 x + c_2 \ln(x) x \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 x + c_2 \ln(x) x \tag{1}$$

Verification of solutions

$$y = c_1 x + c_2 \ln(x) x$$

Verified OK.

14.2.1 Maple step by step solution

Let's solve

$$x^2 y'' - xy' + y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = \frac{y'}{x} - \frac{y}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - \frac{y'}{x} + \frac{y}{x^2} = 0$$

- Multiply by denominators of the ODE

$$x^2 y'' - xy' + y = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x , using the chain rule

$$y' = \left(\frac{d}{dt} y(t) \right) t'(x)$$

- Compute derivative

$$y' = \frac{\frac{d}{dt} y(t)}{x}$$

- Calculate the 2nd derivative of y with respect to x , using the chain rule

$$y'' = \left(\frac{d^2}{dt^2} y(t) \right) t'(x)^2 + t''(x) \left(\frac{d}{dt} y(t) \right)$$

- Compute derivative

$$y'' = \frac{\frac{d^2}{dt^2} y(t)}{x^2} - \frac{\frac{d}{dt} y(t)}{x^2}$$

Substitute the change of variables back into the ODE

$$x^2 \left(\frac{\frac{d^2}{dt^2} y(t)}{x^2} - \frac{\frac{d}{dt} y(t)}{x^2} \right) - \frac{d}{dt} y(t) + y(t) = 0$$

- Simplify

$$\frac{d^2}{dt^2} y(t) - 2 \frac{d}{dt} y(t) + y(t) = 0$$

- Characteristic polynomial of ODE

$$r^2 - 2r + 1 = 0$$

- Factor the characteristic polynomial

$$(r - 1)^2 = 0$$

- Root of the characteristic polynomial

$$r = 1$$

- 1st solution of the ODE

$$y_1(t) = e^t$$

- Repeated root, multiply $y_1(t)$ by t to ensure linear independence

$$y_2(t) = t e^t$$

- General solution of the ODE

$$y(t) = c_1 y_1(t) + c_2 y_2(t)$$

- Substitute in solutions

$$y(t) = c_1 e^t + c_2 t e^t$$

- Change variables back using $t = \ln(x)$

$$y = c_1 x + c_2 \ln(x) x$$

- Simplify

$$y = x(c_1 + c_2 \ln(x))$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
<- LODE of Euler type successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve([x^2*diff(y(x),x$2)-x*diff(y(x),x)+y(x)=0,x],singsol=all)
```

$$y(x) = x(c_2 \ln(x) + c_1)$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 15

```
DSolve[x^2*y''[x]-x*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x(c_2 \log(x) + c_1)$$

14.3 problem 3

14.3.1 Solving as second order change of variable on y method 1 ode .	1034
14.3.2 Solving as second order bessel ode ode	1037
14.3.3 Solving using Kovacic algorithm	1038
14.3.4 Maple step by step solution	1041

Internal problem ID [6018]

Internal file name [OUTPUT/5266_Sunday_June_05_2022_03_28_55_PM_15202253/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 124

Problem number: 3.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_bessel_ode", "second_order_change_of_variable_on_y_method_1"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$x^2y'' + 4xy' + y(x^2 + 2) = 0$$

14.3.1 Solving as second order change of variable on y method 1 ode

In normal form the given ode is written as

$$y'' + p(x)y' + q(x)y = 0 \tag{2}$$

Where

$$p(x) = \frac{4}{x}$$
$$q(x) = \frac{x^2 + 2}{x^2}$$

Calculating the Liouville ode invariant Q given by

$$\begin{aligned}
 Q &= q - \frac{p'}{2} - \frac{p^2}{4} \\
 &= \frac{x^2 + 2}{x^2} - \frac{\left(\frac{4}{x}\right)'}{2} - \frac{\left(\frac{4}{x}\right)^2}{4} \\
 &= \frac{x^2 + 2}{x^2} - \frac{\left(-\frac{4}{x^2}\right)}{2} - \frac{\left(\frac{16}{x^2}\right)}{4} \\
 &= \frac{x^2 + 2}{x^2} - \left(-\frac{2}{x^2}\right) - \frac{4}{x^2} \\
 &= 1
 \end{aligned}$$

Since the Liouville ode invariant does not depend on the independent variable x then the transformation

$$y = v(x) z(x) \tag{3}$$

is used to change the original ode to a constant coefficients ode in v . In (3) the term $z(x)$ is given by

$$\begin{aligned}
 z(x) &= e^{-\left(\int \frac{v(x)}{2} dx\right)} \\
 &= e^{-\int \frac{4}{x} dx} \\
 &= \frac{1}{x^2}
 \end{aligned} \tag{5}$$

Hence (3) becomes

$$y = \frac{v(x)}{x^2} \tag{4}$$

Applying this change of variable to the original ode results in

$$v''(x) + v(x) = 0$$

Which is now solved for $v(x)$ This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Av''(x) + Bv'(x) + Cv(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $v(x) = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\lambda_1 = +i$$

$$\lambda_2 = -i$$

Which simplifies to

$$\lambda_1 = i$$

$$\lambda_2 = -i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$v(x) = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$v(x) = e^0 (\cos(x) c_1 + c_2 \sin(x))$$

Or

$$v(x) = \cos(x) c_1 + c_2 \sin(x)$$

Now that $v(x)$ is known, then

$$\begin{aligned} y &= v(x) z(x) \\ &= (\cos(x) c_1 + c_2 \sin(x)) (z(x)) \end{aligned} \quad (7)$$

But from (5)

$$z(x) = \frac{1}{x^2}$$

Hence (7) becomes

$$y = \frac{\cos(x) c_1 + c_2 \sin(x)}{x^2}$$

Summary

The solution(s) found are the following

$$y = \frac{\cos(x) c_1 + c_2 \sin(x)}{x^2} \quad (1)$$

Verification of solutions

$$y = \frac{\cos(x) c_1 + c_2 \sin(x)}{x^2}$$

Verified OK.

14.3.2 Solving as second order bessel ode ode

Writing the ode as

$$x^2 y'' + 4xy' + y(x^2 + 2) = 0 \quad (1)$$

Bessel ode has the form

$$x^2 y'' + xy' + (-n^2 + x^2) y = 0 \quad (2)$$

The generalized form of Bessel ode is given by Bowman (1958) as the following

$$x^2 y'' + (1 - 2\alpha) xy' + (\beta^2 \gamma^2 x^{2\gamma} - n^2 \gamma^2 + \alpha^2) y = 0 \quad (3)$$

With the standard solution

$$y = x^\alpha (c_1 \text{BesselJ}(n, \beta x^\gamma) + c_2 \text{BesselY}(n, \beta x^\gamma)) \quad (4)$$

Comparing (3) to (1) and solving for α, β, n, γ gives

$$\alpha = -\frac{3}{2}$$

$$\beta = 1$$

$$n = -\frac{1}{2}$$

$$\gamma = 1$$

Substituting all the above into (4) gives the solution as

$$y = \frac{c_1\sqrt{2} \cos(x)}{x^2\sqrt{\pi}} + \frac{c_2\sqrt{2} \sin(x)}{x^2\sqrt{\pi}}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1\sqrt{2} \cos(x)}{x^2\sqrt{\pi}} + \frac{c_2\sqrt{2} \sin(x)}{x^2\sqrt{\pi}} \quad (1)$$

Verification of solutions

$$y = \frac{c_1\sqrt{2} \cos(x)}{x^2\sqrt{\pi}} + \frac{c_2\sqrt{2} \sin(x)}{x^2\sqrt{\pi}}$$

Verified OK.

14.3.3 Solving using Kovacic algorithm

Writing the ode as

$$x^2y'' + 4xy' + y(x^2 + 2) = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= x^2 \\ B &= 4x \\ C &= x^2 + 2 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 180: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{4x}{x^2} dx} \\ &= z_1 e^{-2 \ln(x)} \\ &= z_1 \left(\frac{1}{x^2} \right) \end{aligned}$$

Which simplifies to

$$y_1 = \frac{\cos(x)}{x^2}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{4x}{x^2} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-4 \ln(x)}}{(y_1)^2} dx \\ &= y_1 (\tan(x)) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}
y &= c_1 y_1 + c_2 y_2 \\
&= c_1 \left(\frac{\cos(x)}{x^2} \right) + c_2 \left(\frac{\cos(x)}{x^2} (\tan(x)) \right)
\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1 \cos(x)}{x^2} + \frac{c_2 \sin(x)}{x^2} \quad (1)$$

Verification of solutions

$$y = \frac{c_1 \cos(x)}{x^2} + \frac{c_2 \sin(x)}{x^2}$$

Verified OK.

14.3.4 Maple step by step solution

Let's solve

$$x^2 y'' + 4xy' + y(x^2 + 2) = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{(x^2+2)y}{x^2} - \frac{4y'}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{4y'}{x} + \frac{(x^2+2)y}{x^2} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$\left[P_2(x) = \frac{4}{x}, P_3(x) = \frac{x^2+2}{x^2} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 4$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 2$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$x^2 y'' + 4xy' + y(x^2 + 2) = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 0..2$

$$x^m \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

- Shift index using $k \rightarrow k - m$

$$x^m \cdot y = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

- Convert $x \cdot y'$ to series expansion

$$x \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r}$$

- Convert $x^2 \cdot y''$ to series expansion

$$x^2 \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(2+r)(1+r)x^r + a_1(3+r)(2+r)x^{1+r} + \left(\sum_{k=2}^{\infty} (a_k(k+r+2)(k+r+1) + a_{k-2}) x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$(2+r)(1+r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \{-2, -1\}$$

- Each term must be 0

$$a_1(3+r)(2+r) = 0$$

- Solve for the dependent coefficient(s)

$$a_1 = 0$$

- Each term in the series must be 0, giving the recursion relation

$$a_k(k+r+2)(k+r+1) + a_{k-2} = 0$$

- Shift index using $k \rightarrow k+2$

$$a_{k+2}(k+4+r)(k+3+r) + a_k = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+2} = -\frac{a_k}{(k+4+r)(k+3+r)}$$

- Recursion relation for $r = -2$

$$a_{k+2} = -\frac{a_k}{(k+2)(k+1)}$$

- Solution for $r = -2$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k-2}, a_{k+2} = -\frac{a_k}{(k+2)(k+1)}, a_1 = 0 \right]$$

- Recursion relation for $r = -1$

$$a_{k+2} = -\frac{a_k}{(k+3)(k+2)}$$

- Solution for $r = -1$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k-1}, a_{k+2} = -\frac{a_k}{(k+3)(k+2)}, a_1 = 0 \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^{k-2} \right) + \left(\sum_{k=0}^{\infty} b_k x^{k-1} \right), a_{k+2} = -\frac{a_k}{(k+2)(k+1)}, a_1 = 0, b_{k+2} = -\frac{b_k}{(k+3)(k+2)}, b_1 = 0 \right]$$

Maple trace Kovacic algorithm successful

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Group is reducible or imprimitive
<- Kovacics algorithm successful`

```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 17

```
dsolve(x^2*diff(y(x),x$2)+4*x*diff(y(x),x)+(2+x^2)*y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{c_1 \sin(x) + \cos(x) c_2}{x^2}$$

✓ Solution by Mathematica

Time used: 0.03 (sec). Leaf size: 37

```
DSolve[x^2*y''[x]+4*x*y'[x]+(2+x^2)*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2c_1 e^{-ix} - ic_2 e^{ix}}{2x^2}$$

15 Chapter 3. Linear equations with variable coefficients. Page 130

15.1	problem 1(a)	1046
15.2	problem 1(b)	1055
15.3	problem 1(c)	1065
15.4	problem 1(d)	1074
15.5	problem 1(e)	1084
15.6	problem 2	1094
15.7	problem 3	1106
15.8	problem 4	1116
15.9	problem 5	1128
15.10	problem 6	1131
15.11	problem 7	1143
15.12	problem 8	1158

15.1 problem 1(a)

15.1.1 Maple step by step solution 1053

Internal problem ID [6019]

Internal file name [OUTPUT/5267_Sunday_June_05_2022_03_28_57_PM_36183594/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 130

Problem number: 1(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second order series method. Ordinary point", "second_order_change_of_variable_on_y_method_2", "second order series method. Taylor series method", "second_order_ode_non_constant_coeff_transformation_on_B"

Maple gives the following as the ode type

[_Hermite]

$$y'' - xy' + y = 0$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using

Taylor series gives

$$\begin{aligned}
 y(x) &= y(x_0) + (x - x_0) y'(x_0) + \frac{(x - x_0)^2}{2} y''(x_0) + \frac{(x - x_0)^3}{3!} y'''(x_0) + \dots \\
 &= y_0 + x y'_0 + \frac{x^2}{2} f|_{x_0, y_0, y'_0} + \frac{x^3}{3!} f'|_{x_0, y_0, y'_0} + \dots \\
 &= y_0 + x y'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0}
 \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \tag{1}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \tag{242}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \tag{243}$$

$$\begin{aligned}
 \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \tag{2}
 \end{aligned}$$

$$\begin{aligned}
 \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \tag{3}
 \end{aligned}$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \quad (4)$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \quad (5)$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \\ &\vdots \\ F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \quad (6)$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \quad (7)$$

To find $y(x)$ series solution around $x = 0$. Hence

$$\begin{aligned}
 F_0 &= -y + xy' \\
 F_1 &= \frac{dF_0}{dx} \\
 &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \\
 &= (-y + xy')x \\
 F_2 &= \frac{dF_1}{dx} \\
 &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y}y' + \frac{\partial F_1}{\partial y'}F_1 \\
 &= (x^2 + 1)(-y + xy') \\
 F_3 &= \frac{dF_2}{dx} \\
 &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y}y' + \frac{\partial F_2}{\partial y'}F_2 \\
 &= x(x^2 + 3)(-y + xy') \\
 F_4 &= \frac{dF_3}{dx} \\
 &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y}y' + \frac{\partial F_3}{\partial y'}F_3 \\
 &= (-y + xy')(x^4 + 6x^2 + 3)
 \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and $y'(0) = y'(0)$ gives

$$\begin{aligned}
 F_0 &= -y(0) \\
 F_1 &= 0 \\
 F_2 &= -y(0) \\
 F_3 &= 0 \\
 F_4 &= -3y(0)
 \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$y = \left(1 - \frac{1}{2}x^2 - \frac{1}{24}x^4 - \frac{1}{240}x^6\right)y(0) + xy'(0) + O(x^6)$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard

power series Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$y' = \sum_{n=1}^{\infty} n a_n x^{n-1}$$

$$y'' = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}$$

Substituting the above back into the ode gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = - \left(\sum_{n=0}^{\infty} a_n x^n \right) + x \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) \quad (1)$$

Which simplifies to

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \sum_{n=1}^{\infty} (-n x^n a_n) + \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n \right) + \sum_{n=1}^{\infty} (-n x^n a_n) + \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0 \quad (3)$$

$n = 0$ gives

$$2a_2 + a_0 = 0$$

$$a_2 = -\frac{a_0}{2}$$

For $1 \leq n$, the recurrence equation is

$$(n + 2) a_{n+2}(n + 1) - na_n + a_n = 0 \quad (4)$$

Solving for a_{n+2} , gives

$$a_{n+2} = \frac{a_n(n - 1)}{(n + 2)(n + 1)} \quad (5)$$

For $n = 1$ the recurrence equation gives

$$6a_3 = 0$$

Which after substituting the earlier terms found becomes

$$a_3 = 0$$

For $n = 2$ the recurrence equation gives

$$12a_4 - a_2 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = -\frac{a_0}{24}$$

For $n = 3$ the recurrence equation gives

$$20a_5 - 2a_3 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = 0$$

For $n = 4$ the recurrence equation gives

$$30a_6 - 3a_4 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = -\frac{a_0}{240}$$

For $n = 5$ the recurrence equation gives

$$42a_7 - 4a_5 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = 0$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + a_1 x - \frac{1}{2} a_0 x^2 - \frac{1}{24} a_0 x^4 + \dots$$

Collecting terms, the solution becomes

$$y = \left(1 - \frac{1}{2}x^2 - \frac{1}{24}x^4\right) a_0 + a_1 x + O(x^6) \quad (3)$$

At $x = 0$ the solution above becomes

$$y = \left(1 - \frac{1}{2}x^2 - \frac{1}{24}x^4\right) c_1 + c_2 x + O(x^6)$$

Summary

The solution(s) found are the following

$$y = \left(1 - \frac{1}{2}x^2 - \frac{1}{24}x^4 - \frac{1}{240}x^6\right) y(0) + xy'(0) + O(x^6) \quad (1)$$

$$y = \left(1 - \frac{1}{2}x^2 - \frac{1}{24}x^4\right) c_1 + c_2 x + O(x^6) \quad (2)$$

Verification of solutions

$$y = \left(1 - \frac{1}{2}x^2 - \frac{1}{24}x^4 - \frac{1}{240}x^6\right) y(0) + xy'(0) + O(x^6)$$

Verified OK.

$$y = \left(1 - \frac{1}{2}x^2 - \frac{1}{24}x^4\right) c_1 + c_2 x + O(x^6)$$

Verified OK.

15.1.1 Maple step by step solution

Let's solve

$$y'' = -y + xy'$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - xy' + y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^k$$

- Rewrite DE with series expansions

- Convert $x \cdot y'$ to series expansion

$$x \cdot y' = \sum_{k=0}^{\infty} a_k k x^k$$

- Convert y'' to series expansion

$$y'' = \sum_{k=2}^{\infty} a_k k(k-1) x^{k-2}$$

- Shift index using $k- > k+2$

$$y'' = \sum_{k=0}^{\infty} a_{k+2}(k+2)(k+1) x^k$$

Rewrite DE with series expansions

$$\sum_{k=0}^{\infty} (a_{k+2}(k+2)(k+1) - a_k(k-1)) x^k = 0$$

- Each term in the series must be 0, giving the recursion relation

$$(k^2 + 3k + 2) a_{k+2} - a_k(k-1) = 0$$

- Recursion relation that defines the series solution to the ODE

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+2} = \frac{a_k(k-1)}{k^2+3k+2} \right]$$

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacics algorithm  
  A Liouvillian solution exists  
  Reducible group (found an exponential solution)  
  Group is reducible, not completely reducible  
<- Kovacics algorithm successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
Order:=6;  
dsolve(diff(y(x),x$2)-x*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{2}x^2 - \frac{1}{24}x^4\right)y(0) + D(y)(0)x + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 27

```
AsymptoticDSolveValue[y''[x]-x*y'[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_1 \left(-\frac{x^4}{24} - \frac{x^2}{2} + 1\right) + c_2 x$$

15.2 problem 1(b)

15.2.1 Maple step by step solution 1062

Internal problem ID [6020]

Internal file name [OUTPUT/5268_Sunday_June_05_2022_03_28_58_PM_87853364/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 130

Problem number: 1(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second order series method. Ordinary point", "second order series method. Taylor series method"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' + 3x^2y' - xy = 0$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using Taylor series gives

$$\begin{aligned} y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \cdots \\ &= y_0 + xy'_0 + \frac{x^2}{2}f|_{x_0, y_0, y'_0} + \frac{x^3}{3!}f'|_{x_0, y_0, y'_0} + \cdots \\ &= y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0} \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \quad (1)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \quad (245)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \quad (246)$$

$$\begin{aligned} \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \end{aligned} \quad (2)$$

$$\begin{aligned} \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \end{aligned} \quad (3)$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \quad (4)$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \quad (5)$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \\ &\vdots \\ F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \quad (6)$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \quad (7)$$

To find $y(x)$ series solution around $x = 0$. Hence

$$F_0 = -3x^2y' + xy$$

$$\begin{aligned} F_1 &= \frac{dF_0}{dx} \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \\ &= 9y'x^4 - 3yx^3 - 5xy' + y \end{aligned}$$

$$\begin{aligned} F_2 &= \frac{dF_1}{dx} \\ &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y}y' + \frac{\partial F_1}{\partial y'}F_1 \\ &= -27y'x^6 + 9yx^5 + 48y'x^3 - 14yx^2 - 4y' \end{aligned}$$

$$\begin{aligned} F_3 &= \frac{dF_2}{dx} \\ &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y}y' + \frac{\partial F_2}{\partial y'}F_2 \\ &= (81x^8 - 297x^5 + 142x^2)y' - 27yx\left(x^6 - \frac{31}{9}x^3 + \frac{32}{27}\right) \end{aligned}$$

$$\begin{aligned} F_4 &= \frac{dF_3}{dx} \\ &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y}y' + \frac{\partial F_3}{\partial y'}F_3 \\ &= (-243x^{10} + 1512x^7 - 1818x^4 + 252x)y' + 81\left(x^9 - 6x^6 + \frac{514}{81}x^3 - \frac{32}{81}\right)y \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and $y'(0) = y'(0)$ gives

$$\begin{aligned} F_0 &= 0 \\ F_1 &= y(0) \\ F_2 &= -4y'(0) \\ F_3 &= 0 \\ F_4 &= -32y(0) \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$y = \left(1 + \frac{1}{6}x^3 - \frac{2}{45}x^6\right)y(0) + \left(x - \frac{1}{6}x^4\right)y'(0) + O(x^6)$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard power series. Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$y' = \sum_{n=1}^{\infty} n a_n x^{n-1}$$

$$y'' = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}$$

Substituting the above back into the ode gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = -3x^2 \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) + x \left(\sum_{n=0}^{\infty} a_n x^n \right) \quad (1)$$

Which simplifies to

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \left(\sum_{n=1}^{\infty} 3n x^{1+n} a_n \right) + \sum_{n=0}^{\infty} (-x^{1+n} a_n) = 0 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2) a_{n+2} (1+n) x^n$$

$$\sum_{n=1}^{\infty} 3n x^{1+n} a_n = \sum_{n=2}^{\infty} 3(n-1) a_{n-1} x^n$$

$$\sum_{n=0}^{\infty} (-x^{1+n} a_n) = \sum_{n=1}^{\infty} (-a_{n-1} x^n)$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (1+n) x^n \right) + \left(\sum_{n=2}^{\infty} 3(n-1) a_{n-1} x^n \right) + \sum_{n=1}^{\infty} (-a_{n-1} x^n) = 0 \quad (3)$$

$n = 1$ gives

$$6a_3 - a_0 = 0$$

Which after substituting earlier equations, simplifies to

$$a_3 = \frac{a_0}{6}$$

For $2 \leq n$, the recurrence equation is

$$(n + 2) a_{n+2}(1 + n) + 3(n - 1) a_{n-1} - a_{n-1} = 0 \quad (4)$$

Solving for a_{n+2} , gives

$$a_{n+2} = -\frac{a_{n-1}(3n - 4)}{(n + 2)(1 + n)} \quad (5)$$

For $n = 2$ the recurrence equation gives

$$12a_4 + 2a_1 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = -\frac{a_1}{6}$$

For $n = 3$ the recurrence equation gives

$$20a_5 + 5a_2 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = 0$$

For $n = 4$ the recurrence equation gives

$$30a_6 + 8a_3 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = -\frac{2a_0}{45}$$

For $n = 5$ the recurrence equation gives

$$42a_7 + 11a_4 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = \frac{11a_1}{252}$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + a_1 x + \frac{1}{6} a_0 x^3 - \frac{1}{6} a_1 x^4 + \dots$$

Collecting terms, the solution becomes

$$y = \left(1 + \frac{x^3}{6}\right) a_0 + \left(x - \frac{1}{6} x^4\right) a_1 + O(x^6) \quad (3)$$

At $x = 0$ the solution above becomes

$$y = \left(1 + \frac{x^3}{6}\right) c_1 + \left(x - \frac{1}{6} x^4\right) c_2 + O(x^6)$$

Summary

The solution(s) found are the following

$$y = \left(1 + \frac{1}{6} x^3 - \frac{2}{45} x^6\right) y(0) + \left(x - \frac{1}{6} x^4\right) y'(0) + O(x^6) \quad (1)$$

$$y = \left(1 + \frac{x^3}{6}\right) c_1 + \left(x - \frac{1}{6} x^4\right) c_2 + O(x^6) \quad (2)$$

Verification of solutions

$$y = \left(1 + \frac{1}{6} x^3 - \frac{2}{45} x^6\right) y(0) + \left(x - \frac{1}{6} x^4\right) y'(0) + O(x^6)$$

Verified OK.

$$y = \left(1 + \frac{x^3}{6}\right) c_1 + \left(x - \frac{1}{6} x^4\right) c_2 + O(x^6)$$

Verified OK.

15.2.1 Maple step by step solution

Let's solve

$$y'' = -3x^2y' + xy$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + 3x^2y' - xy = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^k$$

- Rewrite ODE with series expansions

- Convert $x \cdot y$ to series expansion

$$x \cdot y = \sum_{k=0}^{\infty} a_k x^{k+1}$$

- Shift index using $k- > k - 1$

$$x \cdot y = \sum_{k=1}^{\infty} a_{k-1} x^k$$

- Convert $x^2 \cdot y'$ to series expansion

$$x^2 \cdot y' = \sum_{k=0}^{\infty} a_k k x^{k+1}$$

- Shift index using $k- > k - 1$

$$x^2 \cdot y' = \sum_{k=1}^{\infty} a_{k-1} (k-1) x^k$$

- Convert y'' to series expansion

$$y'' = \sum_{k=2}^{\infty} a_k k(k-1) x^{k-2}$$

- Shift index using $k- > k + 2$

$$y'' = \sum_{k=0}^{\infty} a_{k+2} (k+2)(k+1) x^k$$

Rewrite ODE with series expansions

$$2a_2 + \left(\sum_{k=1}^{\infty} (a_{k+2}(k+2)(k+1) + a_{k-1}(3k-4)) x^k \right) = 0$$

- Each term must be 0
 $2a_2 = 0$
- Each term in the series must be 0, giving the recursion relation
 $(k^2 + 3k + 2) a_{k+2} + 3a_{k-1}k - 4a_{k-1} = 0$
- Shift index using $k \rightarrow k + 1$
 $((k + 1)^2 + 3k + 5) a_{k+3} + 3a_k(k + 1) - 4a_k = 0$
- Recursion relation that defines the series solution to the ODE
$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+3} = -\frac{a_k(3k-1)}{k^2+5k+6}, 2a_2 = 0 \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
  -> Bessel
  -> elliptic
  -> Legendre
  -> Kummer
      -> hyper3: Equivalence to 1F1 under a power @ Moebius
      <- hyper3 successful: received ODE is equivalent to the 1F1 ODE
      <- Kummer successful
<- special function solution successful`

```


✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
Order:=6;  
dsolve(diff(y(x),x$2)+3*x^2*diff(y(x),x)-x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{x^3}{6}\right) y(0) + \left(x - \frac{1}{6}x^4\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 28

```
AsymptoticDSolveValue[y''[x]+3*x^2*y'[x]-x*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(x - \frac{x^4}{6}\right) + c_1 \left(\frac{x^3}{6} + 1\right)$$

15.3 problem 1(c)

15.3.1 Maple step by step solution 1071

Internal problem ID [6021]

Internal file name [OUTPUT/5269_Sunday_June_05_2022_03_29_00_PM_31831911/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 130

Problem number: 1(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_bessel_ode", "second order series method. Ordinary point", "second order series method. Taylor series method"

Maple gives the following as the ode type

```
[[_Emden, _Fowler]]
```

$$y'' - yx^2 = 0$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using Taylor series gives

$$\begin{aligned} y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \cdots \\ &= y_0 + xy'_0 + \frac{x^2}{2}f|_{x_0, y_0, y'_0} + \frac{x^3}{3!}f'|_{x_0, y_0, y'_0} + \cdots \\ &= y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0} \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \quad (1)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \quad (248)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \quad (249)$$

$$\begin{aligned} \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \end{aligned} \quad (2)$$

$$\begin{aligned} \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \end{aligned} \quad (3)$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \quad (4)$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \quad (5)$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \\ &\vdots \\ F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \quad (6)$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \quad (7)$$

To find $y(x)$ series solution around $x = 0$. Hence

$$\begin{aligned}
 F_0 &= yx^2 \\
 F_1 &= \frac{dF_0}{dx} \\
 &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \\
 &= x(xy' + 2y) \\
 F_2 &= \frac{dF_1}{dx} \\
 &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y}y' + \frac{\partial F_1}{\partial y'}F_1 \\
 &= yx^4 + 4xy' + 2y \\
 F_3 &= \frac{dF_2}{dx} \\
 &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y}y' + \frac{\partial F_2}{\partial y'}F_2 \\
 &= y'x^4 + 8yx^3 + 6y' \\
 F_4 &= \frac{dF_3}{dx} \\
 &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y}y' + \frac{\partial F_3}{\partial y'}F_3 \\
 &= 12y'x^3 + x^2y(x^4 + 30)
 \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and $y'(0) = y'(0)$ gives

$$\begin{aligned}
 F_0 &= 0 \\
 F_1 &= 0 \\
 F_2 &= 2y(0) \\
 F_3 &= 6y'(0) \\
 F_4 &= 0
 \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$y = \left(1 + \frac{x^4}{12}\right)y(0) + \left(x + \frac{1}{20}x^5\right)y'(0) + O(x^6)$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard

power series Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$y' = \sum_{n=1}^{\infty} n a_n x^{n-1}$$

$$y'' = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}$$

Substituting the above back into the ode gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \left(\sum_{n=0}^{\infty} a_n x^n \right) x^2 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \sum_{n=0}^{\infty} (-x^{n+2} a_n) = 0 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n$$

$$\sum_{n=0}^{\infty} (-x^{n+2} a_n) = \sum_{n=2}^{\infty} (-a_{n-2} x^n)$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n \right) + \sum_{n=2}^{\infty} (-a_{n-2} x^n) = 0 \quad (3)$$

For $2 \leq n$, the recurrence equation is

$$(n+2) a_{n+2} (n+1) - a_{n-2} = 0 \quad (4)$$

Solving for a_{n+2} , gives

$$a_{n+2} = \frac{a_{n-2}}{(n+2)(n+1)} \quad (5)$$

For $n = 2$ the recurrence equation gives

$$12a_4 - a_0 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = \frac{a_0}{12}$$

For $n = 3$ the recurrence equation gives

$$20a_5 - a_1 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = \frac{a_1}{20}$$

For $n = 4$ the recurrence equation gives

$$30a_6 - a_2 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = 0$$

For $n = 5$ the recurrence equation gives

$$42a_7 - a_3 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = 0$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + a_1x + \frac{1}{12}a_0x^4 + \frac{1}{20}a_1x^5 + \dots$$

Collecting terms, the solution becomes

$$y = \left(1 + \frac{x^4}{12}\right) a_0 + \left(x + \frac{1}{20}x^5\right) a_1 + O(x^6) \quad (3)$$

At $x = 0$ the solution above becomes

$$y = \left(1 + \frac{x^4}{12}\right) c_1 + \left(x + \frac{1}{20}x^5\right) c_2 + O(x^6)$$

Summary

The solution(s) found are the following

$$y = \left(1 + \frac{x^4}{12}\right) y(0) + \left(x + \frac{1}{20}x^5\right) y'(0) + O(x^6) \quad (1)$$

$$y = \left(1 + \frac{x^4}{12}\right) c_1 + \left(x + \frac{1}{20}x^5\right) c_2 + O(x^6) \quad (2)$$

Verification of solutions

$$y = \left(1 + \frac{x^4}{12}\right) y(0) + \left(x + \frac{1}{20}x^5\right) y'(0) + O(x^6)$$

Verified OK.

$$y = \left(1 + \frac{x^4}{12}\right) c_1 + \left(x + \frac{1}{20}x^5\right) c_2 + O(x^6)$$

Verified OK.

15.3.1 Maple step by step solution

Let's solve

$$y'' = yx^2$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - yx^2 = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^k$$

- Rewrite ODE with series expansions

- Convert $x^2 \cdot y$ to series expansion

$$x^2 \cdot y = \sum_{k=0}^{\infty} a_k x^{k+2}$$

- Shift index using $k \rightarrow k - 2$

$$x^2 \cdot y = \sum_{k=2}^{\infty} a_{k-2} x^k$$

- Convert y'' to series expansion

$$y'' = \sum_{k=2}^{\infty} a_k k(k-1) x^{k-2}$$

- Shift index using $k \rightarrow k + 2$

$$y'' = \sum_{k=0}^{\infty} a_{k+2} (k+2)(k+1) x^k$$

Rewrite ODE with series expansions

$$6a_3x + 2a_2 + \left(\sum_{k=2}^{\infty} (a_{k+2}(k+2)(k+1) - a_{k-2}) x^k \right) = 0$$

- The coefficients of each power of x must be 0

$$[2a_2 = 0, 6a_3 = 0]$$

- Solve for the dependent coefficient(s)

$$\{a_2 = 0, a_3 = 0\}$$

- Each term in the series must be 0, giving the recursion relation

$$(k^2 + 3k + 2) a_{k+2} - a_{k-2} = 0$$

- Shift index using $k \rightarrow k + 2$

$$((k+2)^2 + 3k + 8) a_{k+4} - a_k = 0$$

- Recursion relation that defines the series solution to the ODE

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+4} = \frac{a_k}{k^2 + 7k + 12}, a_2 = 0, a_3 = 0 \right]$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacic's algorithm  
<- No Liouvillian solutions exist  
-> Trying a solution in terms of special functions:  
  -> Bessel  
  <- Bessel successful  
<- special function solution successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
Order:=6;  
dsolve(diff(y(x),x$2)-x^2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{x^4}{12}\right) y(0) + \left(x + \frac{1}{20}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 28

```
AsymptoticDSolveValue[y''[x]-x^2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{x^5}{20} + x\right) + c_1 \left(\frac{x^4}{12} + 1\right)$$

15.4 problem 1(d)

15.4.1 Maple step by step solution 1081

Internal problem ID [6022]

Internal file name [OUTPUT/5270_Sunday_June_05_2022_03_29_01_PM_84068285/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 130

Problem number: 1(d).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Ordinary point", "second order series method. Taylor series method"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' + y'x^3 + yx^2 = 0$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using Taylor series gives

$$\begin{aligned} y(x) &= y(x_0) + (x - x_0) y'(x_0) + \frac{(x - x_0)^2}{2} y''(x_0) + \frac{(x - x_0)^3}{3!} y'''(x_0) + \cdots \\ &= y_0 + x y'_0 + \frac{x^2}{2} f|_{x_0, y_0, y'_0} + \frac{x^3}{3!} f'|_{x_0, y_0, y'_0} + \cdots \\ &= y_0 + x y'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0} \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \quad (1)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \quad (251)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \quad (252)$$

$$\begin{aligned} \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \end{aligned} \quad (2)$$

$$\begin{aligned} \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \end{aligned} \quad (3)$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \quad (4)$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \end{aligned} \quad (5)$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \\ &\vdots \\ F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \quad (6)$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \quad (7)$$

To find $y(x)$ series solution around $x = 0$. Hence

$$\begin{aligned}
 F_0 &= -y'x^3 - yx^2 \\
 F_1 &= \frac{dF_0}{dx} \\
 &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \\
 &= ((x^5 - 4x)y' + y(x^4 - 2))x \\
 F_2 &= \frac{dF_1}{dx} \\
 &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y}y' + \frac{\partial F_1}{\partial y'}F_1 \\
 &= (-x^9 + 11x^5 - 10x)y' - y(x^8 - 9x^4 + 2) \\
 F_3 &= \frac{dF_2}{dx} \\
 &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y}y' + \frac{\partial F_2}{\partial y'}F_2 \\
 &= (x^{12} - 21x^8 + 74x^4 - 12)y' + yx^3(x^8 - 19x^4 + 46) \\
 F_4 &= \frac{dF_3}{dx} \\
 &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y}y' + \frac{\partial F_3}{\partial y'}F_3 \\
 &= -((x^{13} - 34x^9 + 261x^5 - 354x)y' + y(x^{12} - 32x^8 + 207x^4 - 150))x^2
 \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and $y'(0) = y'(0)$ gives

$$\begin{aligned}
 F_0 &= 0 \\
 F_1 &= 0 \\
 F_2 &= -2y(0) \\
 F_3 &= -12y'(0) \\
 F_4 &= 0
 \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$y = \left(1 - \frac{x^4}{12}\right)y(0) + \left(x - \frac{1}{10}x^5\right)y'(0) + O(x^6)$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard

power series Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$y' = \sum_{n=1}^{\infty} n a_n x^{n-1}$$

$$y'' = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}$$

Substituting the above back into the ode gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = - \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) x^3 - \left(\sum_{n=0}^{\infty} a_n x^n \right) x^2 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \left(\sum_{n=1}^{\infty} n x^{2+n} a_n \right) + \left(\sum_{n=0}^{\infty} x^{2+n} a_n \right) = 0 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (2+n) a_{2+n} (1+n) x^n$$

$$\sum_{n=1}^{\infty} n x^{2+n} a_n = \sum_{n=3}^{\infty} (n-2) a_{n-2} x^n$$

$$\sum_{n=0}^{\infty} x^{2+n} a_n = \sum_{n=2}^{\infty} a_{n-2} x^n$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} (2+n) a_{2+n} (1+n) x^n \right) + \left(\sum_{n=3}^{\infty} (n-2) a_{n-2} x^n \right) + \left(\sum_{n=2}^{\infty} a_{n-2} x^n \right) = 0 \quad (3)$$

$n = 2$ gives

$$12a_4 + a_0 = 0$$

Which after substituting earlier equations, simplifies to

$$a_4 = -\frac{a_0}{12}$$

For $3 \leq n$, the recurrence equation is

$$(2 + n)a_{2+n}(1 + n) + (n - 2)a_{n-2} + a_{n-2} = 0 \quad (4)$$

Solving for a_{2+n} , gives

$$a_{2+n} = -\frac{a_{n-2}(n - 1)}{(2 + n)(1 + n)} \quad (5)$$

For $n = 3$ the recurrence equation gives

$$20a_5 + 2a_1 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = -\frac{a_1}{10}$$

For $n = 4$ the recurrence equation gives

$$30a_6 + 3a_2 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = 0$$

For $n = 5$ the recurrence equation gives

$$42a_7 + 4a_3 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = 0$$

And so on. Therefore the solution is

$$\begin{aligned}y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots\end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + a_1 x - \frac{1}{12} a_0 x^4 - \frac{1}{10} a_1 x^5 + \dots$$

Collecting terms, the solution becomes

$$y = \left(1 - \frac{x^4}{12}\right) a_0 + \left(x - \frac{1}{10} x^5\right) a_1 + O(x^6) \quad (3)$$

At $x = 0$ the solution above becomes

$$y = \left(1 - \frac{x^4}{12}\right) c_1 + \left(x - \frac{1}{10} x^5\right) c_2 + O(x^6)$$

Summary

The solution(s) found are the following

$$y = \left(1 - \frac{x^4}{12}\right) y(0) + \left(x - \frac{1}{10} x^5\right) y'(0) + O(x^6) \quad (1)$$

$$y = \left(1 - \frac{x^4}{12}\right) c_1 + \left(x - \frac{1}{10} x^5\right) c_2 + O(x^6) \quad (2)$$

Verification of solutions

$$y = \left(1 - \frac{x^4}{12}\right) y(0) + \left(x - \frac{1}{10} x^5\right) y'(0) + O(x^6)$$

Verified OK.

$$y = \left(1 - \frac{x^4}{12}\right) c_1 + \left(x - \frac{1}{10} x^5\right) c_2 + O(x^6)$$

Verified OK.

15.4.1 Maple step by step solution

Let's solve

$$y'' = -y'x^3 - yx^2$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + y'x^3 + yx^2 = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^k$$

- Rewrite ODE with series expansions

- Convert $x^2 \cdot y$ to series expansion

$$x^2 \cdot y = \sum_{k=0}^{\infty} a_k x^{k+2}$$

- Shift index using $k- > k - 2$

$$x^2 \cdot y = \sum_{k=2}^{\infty} a_{k-2} x^k$$

- Convert $x^3 \cdot y'$ to series expansion

$$x^3 \cdot y' = \sum_{k=0}^{\infty} a_k k x^{k+2}$$

- Shift index using $k- > k - 2$

$$x^3 \cdot y' = \sum_{k=2}^{\infty} a_{k-2} (k-2) x^k$$

- Convert y'' to series expansion

$$y'' = \sum_{k=2}^{\infty} a_k k(k-1) x^{k-2}$$

- Shift index using $k- > k + 2$

$$y'' = \sum_{k=0}^{\infty} a_{k+2} (k+2)(k+1) x^k$$

Rewrite ODE with series expansions

$$6a_3x + 2a_2 + \left(\sum_{k=2}^{\infty} (a_{k+2}(k+2)(k+1) + a_{k-2}(k-1)) x^k \right) = 0$$

- The coefficients of each power of x must be 0
 $[2a_2 = 0, 6a_3 = 0]$
- Solve for the dependent coefficient(s)
 $\{a_2 = 0, a_3 = 0\}$
- Each term in the series must be 0, giving the recursion relation
 $(k^2 + 3k + 2) a_{k+2} + a_{k-2}(k - 1) = 0$
- Shift index using $k \rightarrow k + 2$
 $((k + 2)^2 + 3k + 8) a_{k+4} + a_k(k + 1) = 0$
- Recursion relation that defines the series solution to the ODE

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+4} = -\frac{a_k(k+1)}{k^2+7k+12}, a_2 = 0, a_3 = 0 \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
-> Bessel
-> elliptic
-> Legendre
-> Kummer
    -> hyper3: Equivalence to 1F1 under a power @ Moebius
    <- hyper3 successful: received ODE is equivalent to the 1F1 ODE
    <- Kummer successful
<- special function solution successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
Order:=6;  
dsolve(diff(y(x),x$2)+x^3*diff(y(x),x)+x^2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{x^4}{12}\right) y(0) + \left(x - \frac{1}{10}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 28

```
AsymptoticDSolveValue[y''[x]+x^3*y'[x]+x^2*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(x - \frac{x^5}{10}\right) + c_1 \left(1 - \frac{x^4}{12}\right)$$

15.5 problem 1(e)

15.5.1 Maple step by step solution 1091

Internal problem ID [6023]

Internal file name [OUTPUT/5271_Sunday_June_05_2022_03_29_03_PM_57526431/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 130

Problem number: 1(e).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_linear_constant_coeff**", "**second_order_ode_can_be_made_integrable**", "**second order series method. Ordinary point**", "**second order series method. Taylor series method**"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' + y = 0$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using Taylor series gives

$$\begin{aligned} y(x) &= y(x_0) + (x - x_0) y'(x_0) + \frac{(x - x_0)^2}{2} y''(x_0) + \frac{(x - x_0)^3}{3!} y'''(x_0) + \cdots \\ &= y_0 + x y'_0 + \frac{x^2}{2} f|_{x_0, y_0, y'_0} + \frac{x^3}{3!} f'|_{x_0, y_0, y'_0} + \cdots \\ &= y_0 + x y'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0} \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \quad (1)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \quad (254)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \quad (255)$$

$$\begin{aligned} \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \end{aligned} \quad (2)$$

$$\begin{aligned} \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \end{aligned} \quad (3)$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \quad (4)$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \quad (5)$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \end{aligned}$$

⋮

$$\begin{aligned} F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \quad (6)$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \quad (7)$$

To find $y(x)$ series solution around $x = 0$. Hence

$$\begin{aligned}
 F_0 &= -y \\
 F_1 &= \frac{dF_0}{dx} \\
 &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \\
 &= -y' \\
 F_2 &= \frac{dF_1}{dx} \\
 &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} y' + \frac{\partial F_1}{\partial y'} F_1 \\
 &= y \\
 F_3 &= \frac{dF_2}{dx} \\
 &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} y' + \frac{\partial F_2}{\partial y'} F_2 \\
 &= y' \\
 F_4 &= \frac{dF_3}{dx} \\
 &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} y' + \frac{\partial F_3}{\partial y'} F_3 \\
 &= -y
 \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and $y'(0) = y'(0)$ gives

$$\begin{aligned}
 F_0 &= -y(0) \\
 F_1 &= -y'(0) \\
 F_2 &= y(0) \\
 F_3 &= y'(0) \\
 F_4 &= -y(0)
 \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$y = \left(1 - \frac{1}{2}x^2 + \frac{1}{24}x^4 - \frac{1}{720}x^6\right) y(0) + \left(x - \frac{1}{6}x^3 + \frac{1}{120}x^5\right) y'(0) + O(x^6)$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard

power series Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$y' = \sum_{n=1}^{\infty} n a_n x^{n-1}$$

$$y'' = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}$$

Substituting the above back into the ode gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = - \left(\sum_{n=0}^{\infty} a_n x^n \right) \quad (1)$$

Which simplifies to

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n \right) + \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0 \quad (3)$$

For $0 \leq n$, the recurrence equation is

$$(n+2) a_{n+2} (n+1) + a_n = 0 \quad (4)$$

Solving for a_{n+2} , gives

$$a_{n+2} = -\frac{a_n}{(n+2)(n+1)} \quad (5)$$

For $n = 0$ the recurrence equation gives

$$2a_2 + a_0 = 0$$

Which after substituting the earlier terms found becomes

$$a_2 = -\frac{a_0}{2}$$

For $n = 1$ the recurrence equation gives

$$6a_3 + a_1 = 0$$

Which after substituting the earlier terms found becomes

$$a_3 = -\frac{a_1}{6}$$

For $n = 2$ the recurrence equation gives

$$12a_4 + a_2 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = \frac{a_0}{24}$$

For $n = 3$ the recurrence equation gives

$$20a_5 + a_3 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = \frac{a_1}{120}$$

For $n = 4$ the recurrence equation gives

$$30a_6 + a_4 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = -\frac{a_0}{720}$$

For $n = 5$ the recurrence equation gives

$$42a_7 + a_5 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = -\frac{a_1}{5040}$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + a_1 x - \frac{1}{2} a_0 x^2 - \frac{1}{6} a_1 x^3 + \frac{1}{24} a_0 x^4 + \frac{1}{120} a_1 x^5 + \dots$$

Collecting terms, the solution becomes

$$y = \left(1 - \frac{1}{2}x^2 + \frac{1}{24}x^4\right) a_0 + \left(x - \frac{1}{6}x^3 + \frac{1}{120}x^5\right) a_1 + O(x^6) \quad (3)$$

At $x = 0$ the solution above becomes

$$y = \left(1 - \frac{1}{2}x^2 + \frac{1}{24}x^4\right) c_1 + \left(x - \frac{1}{6}x^3 + \frac{1}{120}x^5\right) c_2 + O(x^6)$$

Summary

The solution(s) found are the following

$$y = \left(1 - \frac{1}{2}x^2 + \frac{1}{24}x^4 - \frac{1}{720}x^6\right) y(0) + \left(x - \frac{1}{6}x^3 + \frac{1}{120}x^5\right) y'(0) + O(x^6) \quad (1)$$

$$y = \left(1 - \frac{1}{2}x^2 + \frac{1}{24}x^4\right) c_1 + \left(x - \frac{1}{6}x^3 + \frac{1}{120}x^5\right) c_2 + O(x^6) \quad (2)$$




Figure 176: Slope field plot

Verification of solutions

$$y = \left(1 - \frac{1}{2}x^2 + \frac{1}{24}x^4 - \frac{1}{720}x^6\right) y(0) + \left(x - \frac{1}{6}x^3 + \frac{1}{120}x^5\right) y'(0) + O(x^6)$$

Verified OK.

$$y = \left(1 - \frac{1}{2}x^2 + \frac{1}{24}x^4\right) c_1 + \left(x - \frac{1}{6}x^3 + \frac{1}{120}x^5\right) c_2 + O(x^6)$$

Verified OK.

15.5.1 Maple step by step solution

Let's solve

$$y'' = -y$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + y = 0$$

- Characteristic polynomial of ODE
 $r^2 + 1 = 0$
- Use quadratic formula to solve for r
 $r = \frac{0 \pm (\sqrt{-4})}{2}$
- Roots of the characteristic polynomial
 $r = (-I, I)$
- 1st solution of the ODE
 $y_1(x) = \cos(x)$
- 2nd solution of the ODE
 $y_2(x) = \sin(x)$
- General solution of the ODE
 $y = c_1 y_1(x) + c_2 y_2(x)$
- Substitute in solutions
 $y = \cos(x) c_1 + c_2 \sin(x)$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```

Order:=6;
dsolve(diff(y(x),x$2)+y(x)=0,y(x),type='series',x=0);

```

$$y(x) = \left(1 - \frac{1}{2}x^2 + \frac{1}{24}x^4\right) y(0) + \left(x - \frac{1}{6}x^3 + \frac{1}{120}x^5\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 42

```
AsymptoticDSolveValue[y''[x]+y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{x^5}{120} - \frac{x^3}{6} + x \right) + c_1 \left(\frac{x^4}{24} - \frac{x^2}{2} + 1 \right)$$

15.6 problem 2

- 15.6.1 Existence and uniqueness analysis 1094
- 15.6.2 Maple step by step solution 1102

Internal problem ID [6024]

Internal file name [OUTPUT/5272_Sunday_June_05_2022_03_29_04_PM_58711402/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 130

Problem number: 2.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second order series method. Ordinary point", "second_order_change_of_variable_on_y_method_2", "second order series method. Taylor series method"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' + (x - 1)^2 y' - (x - 1)y = 0$$

With initial conditions

$$[y(1) = 1, y'(1) = 0]$$

With the expansion point for the power series method at $x = 1$.

15.6.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = (x - 1)^2$$

$$q(x) = 1 - x$$

$$F = 0$$

Hence the ode is

$$y'' + (x - 1)^2 y' + (1 - x)y = 0$$

The domain of $p(x) = (x - 1)^2$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 1$ is inside this domain. The domain of $q(x) = 1 - x$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 1$ is also inside this domain. Hence solution exists and is unique.

The ode does not have its expansion point at $x = 0$, therefore to simplify the computation of power series expansion, change of variable is made on the independent variable to shift the initial conditions and the expansion point back to zero. The new ode is then solved more easily since the expansion point is now at zero. The solution converted back to the original independent variable. Let

$$t = x - 1$$

The ode is converted to be in terms of the new independent variable t . This results in

$$\frac{d^2}{dt^2}y(t) + \left(\frac{d}{dt}y(t)\right)t^2 - ty(t) = 0$$

With its expansion point and initial conditions now at $t = 0$. With initial conditions now becoming

$$\begin{aligned}y(0) &= 1 \\y'(0) &= 0\end{aligned}$$

The transformed ODE is now solved. Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the

case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using Taylor series gives

$$\begin{aligned}
 y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \cdots \\
 &= y_0 + xy'_0 + \frac{x^2}{2}f|_{x_0, y_0, y'_0} + \frac{x^3}{3!}f'|_{x_0, y_0, y'_0} + \cdots \\
 &= y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0}
 \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \tag{1}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \tag{257}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \tag{258}$$

$$\begin{aligned}
 \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \tag{2}
 \end{aligned}$$

$$\begin{aligned}
 \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \tag{3}
 \end{aligned}$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \tag{4}$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \tag{5}$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \end{aligned}$$

⋮

$$\begin{aligned} F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \tag{6}$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \tag{7}$$

To find $y(x)$ series solution around $x = 0$. Hence

$$\begin{aligned}
 F_0 &= -\left(\frac{d}{dt}y(t)\right)t^2 + ty(t) \\
 F_1 &= \frac{dF_0}{dt} \\
 &= \frac{\partial F_0}{\partial t} + \frac{\partial F_0}{\partial y} \frac{d}{dt}y(t) + \frac{\partial F_0}{\partial \frac{d}{dt}y(t)} F_0 \\
 &= \left(\frac{d}{dt}y(t)\right)t^4 - y(t)t^3 - t\left(\frac{d}{dt}y(t)\right) + y(t) \\
 F_2 &= \frac{dF_1}{dt} \\
 &= \frac{\partial F_1}{\partial t} + \frac{\partial F_1}{\partial y} \frac{d}{dt}y(t) + \frac{\partial F_1}{\partial \frac{d}{dt}y(t)} F_1 \\
 &= -t^2(t^3 - 4)\left(t\left(\frac{d}{dt}y(t)\right) - y(t)\right) \\
 F_3 &= \frac{dF_2}{dt} \\
 &= \frac{\partial F_2}{\partial t} + \frac{\partial F_2}{\partial y} \frac{d}{dt}y(t) + \frac{\partial F_2}{\partial \frac{d}{dt}y(t)} F_2 \\
 &= (t^8 - 9t^5 + 8t^2)\left(\frac{d}{dt}y(t)\right) + (-t^7 + 9t^4 - 8t)y(t) \\
 F_4 &= \frac{dF_3}{dt} \\
 &= \frac{\partial F_3}{\partial t} + \frac{\partial F_3}{\partial y} \frac{d}{dt}y(t) + \frac{\partial F_3}{\partial \frac{d}{dt}y(t)} F_3 \\
 &= -(t^9 - 16t^6 + 44t^3 - 8)\left(t\left(\frac{d}{dt}y(t)\right) - y(t)\right)
 \end{aligned}$$

And so on. Evaluating all the above at initial conditions $t = 0$ and $y(0) = 1$ and $y'(0) = 0$ gives

$$\begin{aligned}
 F_0 &= 0 \\
 F_1 &= 1 \\
 F_2 &= 0 \\
 F_3 &= 0 \\
 F_4 &= -8
 \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$y(t) = 1 + \frac{t^3}{6} - \frac{t^6}{90} + O(t^6)$$

$$y(t) = 1 + \frac{t^3}{6} - \frac{t^6}{90} + O(t^6)$$

Since the expansion point $t = 0$ is an ordinary, we can also solve this using standard power series. Let the solution be represented as power series of the form

$$y(t) = \sum_{n=0}^{\infty} a_n t^n$$

Then

$$\begin{aligned} \frac{d}{dt}y(t) &= \sum_{n=1}^{\infty} n a_n t^{n-1} \\ \frac{d^2}{dt^2}y(t) &= \sum_{n=2}^{\infty} n(n-1) a_n t^{n-2} \end{aligned}$$

Substituting the above back into the ode gives

$$\sum_{n=2}^{\infty} n(n-1) a_n t^{n-2} = - \left(\sum_{n=1}^{\infty} n a_n t^{n-1} \right) t^2 + t \left(\sum_{n=0}^{\infty} a_n t^n \right) \quad (1)$$

Which simplifies to

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n t^{n-2} \right) + \left(\sum_{n=1}^{\infty} n t^{1+n} a_n \right) + \sum_{n=0}^{\infty} (-t^{1+n} a_n) = 0 \quad (2)$$

The next step is to make all powers of t be n in each summation term. Going over each summation term above with power of t in it which is not already t^n and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=2}^{\infty} n(n-1) a_n t^{n-2} &= \sum_{n=0}^{\infty} (n+2) a_{n+2} (1+n) t^n \\ \sum_{n=1}^{\infty} n t^{1+n} a_n &= \sum_{n=2}^{\infty} (n-1) a_{n-1} t^n \\ \sum_{n=0}^{\infty} (-t^{1+n} a_n) &= \sum_{n=1}^{\infty} (-a_{n-1} t^n) \end{aligned}$$

Substituting all the above in Eq (2) gives the following equation where now all powers of t are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (1+n) t^n \right) + \left(\sum_{n=2}^{\infty} (n-1) a_{n-1} t^n \right) + \sum_{n=1}^{\infty} (-a_{n-1} t^n) = 0 \quad (3)$$

$n = 1$ gives

$$6a_3 - a_0 = 0$$

Which after substituting earlier equations, simplifies to

$$a_3 = \frac{a_0}{6}$$

For $2 \leq n$, the recurrence equation is

$$(n + 2) a_{n+2}(1 + n) + (n - 1) a_{n-1} - a_{n-1} = 0 \quad (4)$$

Solving for a_{n+2} , gives

$$a_{n+2} = -\frac{a_{n-1}(n - 2)}{(n + 2)(1 + n)} \quad (5)$$

For $n = 2$ the recurrence equation gives

$$12a_4 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = 0$$

For $n = 3$ the recurrence equation gives

$$20a_5 + a_2 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = 0$$

For $n = 4$ the recurrence equation gives

$$30a_6 + 2a_3 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = -\frac{a_0}{90}$$

For $n = 5$ the recurrence equation gives

$$42a_7 + 3a_4 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = 0$$

And so on. Therefore the solution is

$$\begin{aligned} y(t) &= \sum_{n=0}^{\infty} a_n t^n \\ &= a_3 t^3 + a_2 t^2 + a_1 t + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y(t) = a_0 + a_1 t + \frac{1}{6} a_0 t^3 + \dots$$

Collecting terms, the solution becomes

$$y(t) = \left(1 + \frac{t^3}{6}\right) a_0 + a_1 t + O(t^6) \quad (3)$$

At $t = 0$ the solution above becomes

$$y(t) = \left(1 + \frac{t^3}{6}\right) c_1 + c_2 t + O(t^6)$$

$$y(t) = 1 + \frac{t^3}{6} + O(t^6)$$

Replacing t in the above with the original independent variable x using $t = x - 1$ results in

$$y = 1 + \frac{(x-1)^3}{6} - \frac{(x-1)^6}{90} + O((x-1)^6)$$

Summary

The solution(s) found are the following

$$y = 1 + \frac{(x-1)^3}{6} - \frac{(x-1)^6}{90} + O((x-1)^6) \quad (1)$$

Verification of solutions

$$y = 1 + \frac{(x-1)^3}{6} - \frac{(x-1)^6}{90} + O((x-1)^6)$$

Verified OK.

15.6.2 Maple step by step solution

Let's solve

$$\left[y'' + (x-1)^2 y' + (1-x)y = 0, y(1) = 1, y'|_{\{x=1\}} = 0 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^k$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 0..1$

$$x^m \cdot y = \sum_{k=\max(0,-m)}^{\infty} a_k x^{k+m}$$

- Shift index using $k- > k - m$

$$x^m \cdot y = \sum_{k=\max(0,-m)+m}^{\infty} a_{k-m} x^k$$

- Convert $x^m \cdot y'$ to series expansion for $m = 0..2$

$$x^m \cdot y' = \sum_{k=\max(0,1-m)}^{\infty} a_k k x^{k-1+m}$$

- Shift index using $k- > k + 1 - m$

$$x^m \cdot y' = \sum_{k=\max(0,1-m)+m-1}^{\infty} a_{k+1-m} (k+1-m) x^k$$

- Convert y'' to series expansion

$$y'' = \sum_{k=2}^{\infty} a_k k(k-1) x^{k-2}$$

- Shift index using $k- > k + 2$

$$y'' = \sum_{k=0}^{\infty} a_{k+2} (k+2)(k+1) x^k$$

Rewrite ODE with series expansions

$$2a_2 + a_1 + a_0 + \left(\sum_{k=1}^{\infty} (a_{k+2}(k+2)(k+1) + a_{k+1}(k+1) - a_k(2k-1) + a_{k-1}(k-2)) x^k \right) = 0$$

- Each term must be 0

$$2a_2 + a_1 + a_0 = 0$$

- Each term in the series must be 0, giving the recursion relation

$$k^2 a_{k+2} + (-2a_k + a_{k-1} + a_{k+1} + 3a_{k+2}) k + a_k - 2a_{k-1} + a_{k+1} + 2a_{k+2} = 0$$

- Shift index using $k \rightarrow k+1$

$$(k+1)^2 a_{k+3} + (-2a_{k+1} + a_k + a_{k+2} + 3a_{k+3}) (k+1) + a_{k+1} - 2a_k + a_{k+2} + 2a_{k+3} = 0$$

- Recursion relation that defines the series solution to the ODE

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+3} = -\frac{a_k k - 2a_{k+1} k + k a_{k+2} - a_k - a_{k+1} + 2a_{k+2}}{k^2 + 5k + 6}, 2a_2 + a_1 + a_0 = 0 \right]$$

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
  A Liouvillian solution exists
  Reducible group (found an exponential solution)
  Group is reducible, not completely reducible
  Solution has integrals. Trying a special function solution free of integrals...
-> Trying a solution in terms of special functions:
  -> Bessel
  -> elliptic
  -> Legendre
  -> Kummer
    -> hyper3: Equivalence to 1F1 under a power @ Moebius
  -> hypergeometric
    -> heuristic approach
    -> hyper3: Equivalence to 2F1, 1F1 or 0F1 under a power @ Moebius
  -> Mathieu
    -> Equivalence to the rational form of Mathieu ODE under a power @ Moebius
  -> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
    <- Heun successful: received ODE is equivalent to the HeunT ODE, case c = 0
  Special function solution also has integrals. Returning default Liouvillian solution.
<- Kovacics algorithm successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
Order:=6;
dsolve([diff(y(x),x$2)+(x-1)^2*diff(y(x),x)-(x-1)*y(x)=0,y(1) = 1, D(y)(1) = 0],y(x),type='s'
```

$$y(x) = 1 + \frac{1}{6}(x-1)^3 + O((x-1)^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 14

```
AsymptoticDSolveValue[{y'[x]+(x-1)^2*y'[x]-(x-1)*y[x]==0,{y[1]==1,y'[1]==0}},y[x],{x,1,5}]
```

$$y(x) \rightarrow \frac{1}{6}(x-1)^3 + 1$$

15.7 problem 3

15.7.1 Existence and uniqueness analysis 1106

Internal problem ID [6025]

Internal file name [OUTPUT/5273_Sunday_June_05_2022_03_29_08_PM_63537463/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 130

Problem number: 3.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Ordinary point", "second order series method. Taylor series method"**

Maple gives the following as the ode type

[[_Emden , _Fowler]]

$$(x^2 + 1) y'' + y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 1]$$

With the expansion point for the power series method at $x = 0$.

15.7.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$\begin{aligned} p(x) &= 0 \\ q(x) &= \frac{1}{x^2 + 1} \\ F &= 0 \end{aligned}$$

Hence the ode is

$$y'' + \frac{y}{x^2 + 1} = 0$$

The domain of $p(x) = 0$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = \frac{1}{x^2+1}$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using Taylor series gives

$$\begin{aligned} y(x) &= y(x_0) + (x - x_0) y'(x_0) + \frac{(x - x_0)^2}{2} y''(x_0) + \frac{(x - x_0)^3}{3!} y'''(x_0) + \dots \\ &= y_0 + x y'_0 + \frac{x^2}{2} f|_{x_0, y_0, y'_0} + \frac{x^3}{3!} f'|_{x_0, y_0, y'_0} + \dots \\ &= y_0 + x y'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \left. \frac{d^n f}{dx^n} \right|_{x_0, y_0, y'_0} \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \quad (1)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \quad (260)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \quad (261)$$

$$\begin{aligned} \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \end{aligned} \quad (2)$$

$$\begin{aligned} \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \end{aligned} \quad (3)$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \quad (4)$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \quad (5)$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \\ &\vdots \\ F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \quad (6)$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \quad (7)$$

To find $y(x)$ series solution around $x = 0$. Hence

$$\begin{aligned}
 F_0 &= -\frac{y}{x^2 + 1} \\
 F_1 &= \frac{dF_0}{dx} \\
 &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \\
 &= \frac{-x^2 y' + 2xy - y'}{(x^2 + 1)^2} \\
 F_2 &= \frac{dF_1}{dx} \\
 &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} y' + \frac{\partial F_1}{\partial y'} F_1 \\
 &= \frac{4y'x^3 - 5yx^2 + 4xy' + 3y}{(x^2 + 1)^3} \\
 F_3 &= \frac{dF_2}{dx} \\
 &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} y' + \frac{\partial F_2}{\partial y'} F_2 \\
 &= \frac{(-17x^4 - 10x^2 + 7)y' + 16yx(x^2 - 2)}{(x^2 + 1)^4} \\
 F_4 &= \frac{dF_3}{dx} \\
 &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} y' + \frac{\partial F_3}{\partial y'} F_3 \\
 &= \frac{(84x^5 - 24x^3 - 108x)y' + (-63x^4 + 282x^2 - 39)y}{(x^2 + 1)^5}
 \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = 0$ and $y'(0) = 1$ gives

$$\begin{aligned}
 F_0 &= 0 \\
 F_1 &= -1 \\
 F_2 &= 0 \\
 F_3 &= 7 \\
 F_4 &= 0
 \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$y = x - \frac{x^3}{6} + \frac{7x^5}{120} + O(x^6)$$

$$y = x - \frac{x^3}{6} + \frac{7x^5}{120} + O(x^6)$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard power series The ode is normalized to be

$$(x^2 + 1) y'' + y = 0$$

Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$y' = \sum_{n=1}^{\infty} n a_n x^{n-1}$$

$$y'' = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}$$

Substituting the above back into the ode gives

$$(x^2 + 1) \left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=2}^{\infty} x^n a_n n(n-1) \right) + \left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=2}^{\infty} x^n a_n n(n-1) \right) + \left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n \right) + \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0 \quad (3)$$

$n = 0$ gives

$$2a_2 + a_0 = 0$$

$$a_2 = -\frac{a_0}{2}$$

$n = 1$ gives

$$6a_3 + a_1 = 0$$

Which after substituting earlier equations, simplifies to

$$a_3 = -\frac{a_1}{6}$$

For $2 \leq n$, the recurrence equation is

$$na_n(n-1) + (n+2)a_{n+2}(n+1) + a_n = 0 \quad (4)$$

Solving for a_{n+2} , gives

$$a_{n+2} = -\frac{a_n(n^2 - n + 1)}{(n+2)(n+1)} \quad (5)$$

For $n = 2$ the recurrence equation gives

$$3a_2 + 12a_4 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = \frac{a_0}{8}$$

For $n = 3$ the recurrence equation gives

$$7a_3 + 20a_5 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = \frac{7a_1}{120}$$

For $n = 4$ the recurrence equation gives

$$13a_4 + 30a_6 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = -\frac{13a_0}{240}$$

For $n = 5$ the recurrence equation gives

$$21a_5 + 42a_7 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = -\frac{7a_1}{240}$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + a_1 x - \frac{1}{2} a_0 x^2 - \frac{1}{6} a_1 x^3 + \frac{1}{8} a_0 x^4 + \frac{7}{120} a_1 x^5 + \dots$$

Collecting terms, the solution becomes

$$y = \left(1 - \frac{1}{2}x^2 + \frac{1}{8}x^4\right) a_0 + \left(x - \frac{1}{6}x^3 + \frac{7}{120}x^5\right) a_1 + O(x^6) \quad (3)$$

At $x = 0$ the solution above becomes

$$y = \left(1 - \frac{1}{2}x^2 + \frac{1}{8}x^4\right) c_1 + \left(x - \frac{1}{6}x^3 + \frac{7}{120}x^5\right) c_2 + O(x^6)$$

$$y = x - \frac{x^3}{6} + \frac{7x^5}{120} + O(x^6)$$

Summary

The solution(s) found are the following

$$y = x - \frac{x^3}{6} + \frac{7x^5}{120} + O(x^6) \quad (1)$$

$$y = x - \frac{x^3}{6} + \frac{7x^5}{120} + O(x^6) \quad (2)$$

Verification of solutions

$$y = x - \frac{x^3}{6} + \frac{7x^5}{120} + O(x^6)$$

Verified OK.

$$y = x - \frac{x^3}{6} + \frac{7x^5}{120} + O(x^6)$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
  -> Bessel
  -> elliptic
  -> Legendre
  -> Whittaker
    -> hyper3: Equivalence to 1F1 under a power @ Moebius
  -> hypergeometric
    -> heuristic approach
    -> hyper3: Equivalence to 2F1, 1F1 or 0F1 under a power @ Moebius
      <- hyper3 successful: received ODE is equivalent to the 2F1 ODE
      <- hypergeometric successful
<- special function solution successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
Order:=6;  
dsolve([(1+x^2)*diff(y(x),x$2)+y(x)=0,y(0) = 0, D(y)(0) = 1],y(x),type='series',x=0);
```

$$y(x) = x - \frac{1}{6}x^3 + \frac{7}{120}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 19

```
AsymptoticDSolveValue[{(1+x^2)*y''[x]+y[x]==0,{y[0]==0,y'[0]==1}},y[x],{x,0,5}]
```

$$y(x) \rightarrow \frac{7x^5}{120} - \frac{x^3}{6} + x$$

15.8 problem 4

15.8.1 Existence and uniqueness analysis 1116

Internal problem ID [6026]

Internal file name [OUTPUT/5274_Sunday_June_05_2022_03_29_11_PM_66482120/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 130

Problem number: 4.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_bessel_ode_form_A**", "**second order series method. Ordinary point**", "**second order series method. Taylor series method**"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' + e^x y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 0]$$

With the expansion point for the power series method at $x = 0$.

15.8.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = 0$$

$$q(x) = e^x$$

$$F = 0$$

Hence the ode is

$$y'' + e^x y = 0$$

The domain of $p(x) = 0$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = e^x$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using Taylor series gives

$$\begin{aligned} y(x) &= y(x_0) + (x - x_0) y'(x_0) + \frac{(x - x_0)^2}{2} y''(x_0) + \frac{(x - x_0)^3}{3!} y'''(x_0) + \dots \\ &= y_0 + x y'_0 + \frac{x^2}{2} f|_{x_0, y_0, y'_0} + \frac{x^3}{3!} f'|_{x_0, y_0, y'_0} + \dots \\ &= y_0 + x y'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \left. \frac{d^n f}{dx^n} \right|_{x_0, y_0, y'_0} \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \quad (1)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \quad (263)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \quad (264)$$

$$\begin{aligned} \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \end{aligned} \quad (2)$$

$$\begin{aligned} \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \end{aligned} \quad (3)$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \quad (4)$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \end{aligned} \quad (5)$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \\ &\vdots \\ F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \quad (6)$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \quad (7)$$

To find $y(x)$ series solution around $x = 0$. Hence

$$\begin{aligned}
 F_0 &= -e^x y \\
 F_1 &= \frac{dF_0}{dx} \\
 &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \\
 &= -e^x (y + y') \\
 F_2 &= \frac{dF_1}{dx} \\
 &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} y' + \frac{\partial F_1}{\partial y'} F_1 \\
 &= e^x (-2y' + (e^x - 1)y) \\
 F_3 &= \frac{dF_2}{dx} \\
 &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} y' + \frac{\partial F_2}{\partial y'} F_2 \\
 &= (4y + y') e^{2x} - e^x (y + 3y') \\
 F_4 &= \frac{dF_3}{dx} \\
 &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} y' + \frac{\partial F_3}{\partial y'} F_3 \\
 &= (11y + 6y') e^{2x} - e^{3x} y - e^x (4y' + y)
 \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = 1$ and $y'(0) = 0$ gives

$$\begin{aligned}
 F_0 &= -1 \\
 F_1 &= -1 \\
 F_2 &= 0 \\
 F_3 &= 3 \\
 F_4 &= 9
 \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$y = 1 - \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^5}{40} + \frac{x^6}{80} + O(x^6)$$

$$y = 1 - \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^5}{40} + \frac{x^6}{80} + O(x^6)$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard power series. Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$y' = \sum_{n=1}^{\infty} n a_n x^{n-1}$$

$$y'' = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}$$

Substituting the above back into the ode gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = -e^x \left(\sum_{n=0}^{\infty} a_n x^n \right) \quad (1)$$

Expanding e^x as Taylor series around $x = 0$ and keeping only the first 6 terms gives

$$e^x = 1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{720}x^6 + \dots$$

$$= 1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{720}x^6$$

Hence the ODE in Eq (1) becomes

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \left(1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{720}x^6 \right) \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0$$

Expanding the second term in (1) gives

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + 1 \cdot \left(\sum_{n=0}^{\infty} a_n x^n \right) + x \cdot \left(\sum_{n=0}^{\infty} a_n x^n \right) + \frac{x^2}{2} \cdot \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$+ \frac{x^3}{6} \cdot \left(\sum_{n=0}^{\infty} a_n x^n \right) + \frac{x^4}{24} \cdot \left(\sum_{n=0}^{\infty} a_n x^n \right) + \frac{x^5}{120} \cdot \left(\sum_{n=0}^{\infty} a_n x^n \right) + \frac{x^6}{720} \cdot \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \left(\sum_{n=0}^{\infty} a_n x^n \right) + \left(\sum_{n=0}^{\infty} x^{1+n} a_n \right) + \left(\sum_{n=0}^{\infty} \frac{x^{n+2} a_n}{2} \right) \\ & + \left(\sum_{n=0}^{\infty} \frac{x^{n+3} a_n}{6} \right) + \left(\sum_{n=0}^{\infty} \frac{x^{n+4} a_n}{24} \right) + \left(\sum_{n=0}^{\infty} \frac{x^{n+5} a_n}{120} \right) + \left(\sum_{n=0}^{\infty} \frac{x^{n+6} a_n}{720} \right) = 0 \end{aligned} \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} &= \sum_{n=0}^{\infty} (n+2) a_{n+2} (1+n) x^n \\ \sum_{n=0}^{\infty} x^{1+n} a_n &= \sum_{n=1}^{\infty} a_{n-1} x^n \\ \sum_{n=0}^{\infty} \frac{x^{n+2} a_n}{2} &= \sum_{n=2}^{\infty} \frac{a_{n-2} x^n}{2} \\ \sum_{n=0}^{\infty} \frac{x^{n+3} a_n}{6} &= \sum_{n=3}^{\infty} \frac{a_{n-3} x^n}{6} \\ \sum_{n=0}^{\infty} \frac{x^{n+4} a_n}{24} &= \sum_{n=4}^{\infty} \frac{a_{n-4} x^n}{24} \\ \sum_{n=0}^{\infty} \frac{x^{n+5} a_n}{120} &= \sum_{n=5}^{\infty} \frac{a_{n-5} x^n}{120} \\ \sum_{n=0}^{\infty} \frac{x^{n+6} a_n}{720} &= \sum_{n=6}^{\infty} \frac{a_{n-6} x^n}{720} \end{aligned}$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (1+n) x^n \right) + \left(\sum_{n=0}^{\infty} a_n x^n \right) + \left(\sum_{n=1}^{\infty} a_{n-1} x^n \right) + \left(\sum_{n=2}^{\infty} \frac{a_{n-2} x^n}{2} \right) \\ & + \left(\sum_{n=3}^{\infty} \frac{a_{n-3} x^n}{6} \right) + \left(\sum_{n=4}^{\infty} \frac{a_{n-4} x^n}{24} \right) + \left(\sum_{n=5}^{\infty} \frac{a_{n-5} x^n}{120} \right) + \left(\sum_{n=6}^{\infty} \frac{a_{n-6} x^n}{720} \right) = 0 \end{aligned} \quad (3)$$

$n = 0$ gives

$$2a_2 + a_0 = 0$$

$$a_2 = -\frac{a_0}{2}$$

$n = 1$ gives

$$6a_3 + a_1 + a_0 = 0$$

Which after substituting earlier equations, simplifies to

$$a_3 = -\frac{a_0}{6} - \frac{a_1}{6}$$

$n = 2$ gives

$$12a_4 + a_2 + a_1 + \frac{a_0}{2} = 0$$

Which after substituting earlier equations, simplifies to

$$a_4 = -\frac{a_1}{12}$$

$n = 3$ gives

$$20a_5 + a_3 + a_2 + \frac{a_1}{2} + \frac{a_0}{6} = 0$$

Which after substituting earlier equations, simplifies to

$$a_5 = \frac{a_0}{40} - \frac{a_1}{60}$$

$n = 4$ gives

$$30a_6 + a_4 + a_3 + \frac{a_2}{2} + \frac{a_1}{6} + \frac{a_0}{24} = 0$$

Which after substituting earlier equations, simplifies to

$$a_6 = \frac{a_0}{80} + \frac{a_1}{360}$$

$n = 5$ gives

$$42a_7 + a_5 + a_4 + \frac{a_3}{2} + \frac{a_2}{6} + \frac{a_1}{24} + \frac{a_0}{120} = 0$$

Which after substituting earlier equations, simplifies to

$$a_7 = \frac{a_0}{315} + \frac{17a_1}{5040}$$

For $6 \leq n$, the recurrence equation is

$$(n+2)a_{n+2}(1+n) + a_n + a_{n-1} + \frac{a_{n-2}}{2} + \frac{a_{n-3}}{6} + \frac{a_{n-4}}{24} + \frac{a_{n-5}}{120} + \frac{a_{n-6}}{720} = 0 \quad (4)$$

Solving for a_{n+2} , gives

$$\begin{aligned} a_{n+2} &= -\frac{720a_n + 720a_{n-1} + 360a_{n-2} + 120a_{n-3} + 30a_{n-4} + 6a_{n-5} + a_{n-6}}{720(n+2)(1+n)} \\ (5) \quad &= -\frac{a_n}{(n+2)(1+n)} - \frac{a_{n-1}}{(n+2)(1+n)} - \frac{a_{n-2}}{120(n+2)(1+n)} \\ &\quad - \frac{a_{n-3}}{24(n+2)(1+n)} - \frac{a_{n-4}}{6(n+2)(1+n)} - \frac{a_{n-5}}{2(n+2)(1+n)} - \frac{a_{n-6}}{(n+2)(1+n)} \end{aligned}$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + a_1 x - \frac{a_0 x^2}{2} + \left(-\frac{a_0}{6} - \frac{a_1}{6}\right) x^3 - \frac{a_1 x^4}{12} + \left(\frac{a_0}{40} - \frac{a_1}{60}\right) x^5 + \dots$$

Collecting terms, the solution becomes

$$y = \left(1 - \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{40}x^5\right) a_0 + \left(x - \frac{1}{6}x^3 - \frac{1}{12}x^4 - \frac{1}{60}x^5\right) a_1 + O(x^6) \quad (3)$$

At $x = 0$ the solution above becomes

$$y = \left(1 - \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{40}x^5\right) c_1 + \left(x - \frac{1}{6}x^3 - \frac{1}{12}x^4 - \frac{1}{60}x^5\right) c_2 + O(x^6)$$

$$y = 1 - \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^5}{40} + O(x^6)$$

Summary

The solution(s) found are the following

$$y = 1 - \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^5}{40} + \frac{x^6}{80} + O(x^6) \quad (1)$$

$$y = 1 - \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^5}{40} + O(x^6) \quad (2)$$

Verification of solutions

$$y = 1 - \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^5}{40} + \frac{x^6}{80} + O(x^6)$$

Verified OK.

$$y = 1 - \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^5}{40} + O(x^6)$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) * 2F1([a
-> Trying changes of variables to rationalize or make the ODE simpler
    trying a quadrature
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exists
-> Trying a solution in terms of special functions:
    -> Bessel
    <- Bessel successful
<- special function solution successful
Change of variables used:
    [x = ln(t)]
Linear ODE actually solved:
    u(t)+diff(u(t),t)+t*diff(diff(u(t),t),t) = 0
<- change of variables successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
Order:=6;
dsolve([diff(y(x),x$2)+exp(x)*y(x)=0,y(0) = 1, D(y)(0) = 0],y(x),type='series',x=0);
```

$$y(x) = 1 - \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{40}x^5 + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 56

```
AsymptoticDSolveValue[{y'[x]+Exp[x]*y[x]==0,{}},y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(-\frac{x^5}{60} - \frac{x^4}{12} - \frac{x^3}{6} + x \right) + c_1 \left(\frac{x^5}{40} - \frac{x^3}{6} - \frac{x^2}{2} + 1 \right)$$

15.9 problem 5

15.9.1 Maple step by step solution 1128

Internal problem ID [6027]

Internal file name [OUTPUT/5275_Sunday_June_05_2022_03_29_13_PM_38106072/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 130

Problem number: 5.

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : **"unknown"**

Maple gives the following as the ode type

```
[[_3rd_order , _with_linear_symmetries]]
```

Unable to solve or complete the solution.

$$y''' - xy = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 0, y''(0) = 0]$$

Unable to solve this ODE. Initial conditions are used to solve for the constants of integration.

15.9.1 Maple step by step solution

Let's solve

$$\left[y''' - xy = 0, y(0) = 1, y'|_{\{x=0\}} = 0, y''|_{\{x=0\}} = 0 \right]$$

- Highest derivative means the order of the ODE is 3
 y'''
- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^k$$

□ Rewrite ODE with series expansions

- Convert $x \cdot y$ to series expansion

$$x \cdot y = \sum_{k=0}^{\infty} a_k x^{k+1}$$

- Shift index using $k \rightarrow k - 1$

$$x \cdot y = \sum_{k=1}^{\infty} a_{k-1} x^k$$

- Convert y''' to series expansion

$$y''' = \sum_{k=3}^{\infty} a_k k(k-1)(k-2) x^{k-3}$$

- Shift index using $k \rightarrow k + 3$

$$y''' = \sum_{k=0}^{\infty} a_{k+3} (k+3)(k+2)(k+1) x^k$$

Rewrite ODE with series expansions

$$6a_3 + \left(\sum_{k=1}^{\infty} (a_{k+3}(k+3)(k+2)(k+1) - a_{k-1}) x^k \right) = 0$$

- Each term must be 0

$$6a_3 = 0$$

- Each term in the series must be 0, giving the recursion relation

$$(k^3 + 6k^2 + 11k + 6) a_{k+3} - a_{k-1} = 0$$

- Shift index using $k \rightarrow k + 1$

$$((k+1)^3 + 6(k+1)^2 + 11k + 17) a_{k+4} - a_k = 0$$

- Recursion relation that defines the series solution to the ODE

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+4} = \frac{a_k}{k^3 + 9k^2 + 26k + 24}, 6a_3 = 0 \right]$$

Maple trace

```
`Methods for third order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying high order exact linear fully integrable  
trying to convert to a linear ODE with constant coefficients  
trying differential order: 3; missing the dependent variable  
trying Louvillian solutions for 3rd order ODEs, imprimitive case  
-> pFq: Equivalence to the 3F2 or one of its 3 confluent cases under a power @ Moebius  
<- pFq successful: received ODE is equivalent to the 0F2 ODE, case c = 0`
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 14

```
dsolve([diff(y(x),x$3)-x*y(x)=0,y(0) = 1, D(y)(0) = 0, (D@@2)(y)(0) = 0],y(x), singsol=all)
```

$$y(x) = \text{hypergeom}\left(\left[\right], \left[\frac{1}{2}, \frac{3}{4}\right], \frac{x^4}{64}\right)$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 21

```
DSolve[{y'''[x]-x*y[x]==0,{y[0]==1,y'[0]==0,y''[0]==0}},y[x],x,IncludeSingularSolutions -> T
```

$$y(x) \rightarrow {}_0F_2\left(\left[\right]; \frac{1}{2}, \frac{3}{4}; \frac{x^4}{64}\right)$$

15.10 problem 6

15.10.1 Maple step by step solution 1139

Internal problem ID [6028]

Internal file name [OUTPUT/5276_Sunday_June_05_2022_03_29_14_PM_70586326/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 130

Problem number: 6.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Ordinary point", "second order series method. Taylor series method"**

Maple gives the following as the ode type

[_Gegenbauer]

$$(-x^2 + 1) y'' - 2xy' + \alpha(\alpha + 1) y = 0$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using Taylor series gives

$$\begin{aligned} y(x) &= y(x_0) + (x - x_0) y'(x_0) + \frac{(x - x_0)^2}{2} y''(x_0) + \frac{(x - x_0)^3}{3!} y'''(x_0) + \cdots \\ &= y_0 + xy'_0 + \frac{x^2}{2} f|_{x_0, y_0, y'_0} + \frac{x^3}{3!} f'|_{x_0, y_0, y'_0} + \cdots \\ &= y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0} \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \quad (1)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \quad (266)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \quad (267)$$

$$\begin{aligned} \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \end{aligned} \quad (2)$$

$$\begin{aligned} \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \end{aligned} \quad (3)$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \quad (4)$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \end{aligned} \quad (5)$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \end{aligned}$$

⋮

$$\begin{aligned} F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \quad (6)$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \quad (7)$$

To find $y(x)$ series solution around $x = 0$. Hence

$$F_0 = \frac{\alpha^2 y + \alpha y - 2xy'}{x^2 - 1}$$

$$\begin{aligned} F_1 &= \frac{dF_0}{dx} \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \\ &= \frac{(x^2 \alpha^2 + x^2 \alpha - \alpha^2 + 6x^2 - \alpha + 2) y' - 4\alpha xy(\alpha + 1)}{(x^2 - 1)^2} \end{aligned}$$

$$\begin{aligned} F_2 &= \frac{dF_1}{dx} \\ &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} y' + \frac{\partial F_1}{\partial y'} F_1 \\ &= \frac{8 \left(((\alpha^2 + \alpha + 3) x^2 - \alpha^2 - \alpha + 3) xy' - \frac{y\alpha((\alpha^2 + \alpha + 18)x^2 - \alpha^2 - \alpha + 6)(\alpha + 1)}{8} \right) (x - 1)(1 + x)}{(x^2 - 1)^4} \end{aligned}$$

$$\begin{aligned} F_3 &= \frac{dF_2}{dx} \\ &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} y' + \frac{\partial F_2}{\partial y'} F_2 \\ &= \frac{(x - 1)(1 + x) \left((\alpha^4 + 2\alpha^3 + 59\alpha^2 + 58\alpha + 120) x^4 + (-2\alpha^4 - 4\alpha^3 - 46\alpha^2 - 44\alpha + 240) x^2 + \alpha^4 + \right)}{(x^2 - 1)^5} \end{aligned}$$

$$\begin{aligned} F_4 &= \frac{dF_3}{dx} \\ &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} y' + \frac{\partial F_3}{\partial y'} F_3 \\ &= \frac{18 \left(\left(40 + (x^4 - 2x^2 + 1) \alpha^4 + 2(x^4 - 2x^2 + 1) \alpha^3 + \left(-\frac{26}{3}x^2 - 17 + \frac{77}{3}x^4 \right) \alpha^2 + 2 \left(-9 - \frac{10}{3}x^2 + \frac{37}{3}x \right) \alpha \right) \right)}{(x^2 - 1)^6} \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and $y'(0) = y'(0)$ gives

$$F_0 = -y(0) \alpha(\alpha + 1)$$

$$F_1 = -y'(0) \alpha^2 - y'(0) \alpha + 2y'(0)$$

$$F_2 = y(0) \alpha^4 + 2y(0) \alpha^3 - 5y(0) \alpha^2 - 6y(0) \alpha$$

$$F_3 = y'(0) \alpha^4 + 2y'(0) \alpha^3 - 13y'(0) \alpha^2 - 14y'(0) \alpha + 24y'(0)$$

$$F_4 = -y(0) \alpha^6 - 3y(0) \alpha^5 + 23y(0) \alpha^4 + 51y(0) \alpha^3 - 94y(0) \alpha^2 - 120y(0) \alpha$$

Substituting all the above in (7) and simplifying gives the solution as

$$\begin{aligned}
 y = & \left(1 - \frac{1}{2}x^2\alpha^2 - \frac{1}{2}x^2\alpha + \frac{1}{24}\alpha^4x^4 + \frac{1}{12}\alpha^3x^4 - \frac{5}{24}\alpha^2x^4 - \frac{1}{4}\alpha x^4 - \frac{1}{720}x^6\alpha^6 - \frac{1}{240}x^6\alpha^5 \right. \\
 & \left. + \frac{23}{720}x^6\alpha^4 + \frac{17}{240}x^6\alpha^3 - \frac{47}{360}x^6\alpha^2 - \frac{1}{6}x^6\alpha \right) y(0) \\
 & + \left(x - \frac{1}{6}\alpha^2x^3 - \frac{1}{6}\alpha x^3 + \frac{1}{3}x^3 + \frac{1}{120}x^5\alpha^4 + \frac{1}{60}x^5\alpha^3 - \frac{13}{120}x^5\alpha^2 - \frac{7}{60}x^5\alpha + \frac{1}{5}x^5 \right) y'(0) \\
 & + O(x^6)
 \end{aligned}$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard power series The ode is normalized to be

$$(-x^2 + 1)y'' - 2xy' + (\alpha^2 + \alpha)y = 0$$

Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$\begin{aligned}
 y' &= \sum_{n=1}^{\infty} n a_n x^{n-1} \\
 y'' &= \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}
 \end{aligned}$$

Substituting the above back into the ode gives

$$(-x^2 + 1) \left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) - 2x \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) + (\alpha^2 + \alpha) \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0 \quad (1)$$

Which simplifies to

$$\begin{aligned}
 & \sum_{n=2}^{\infty} (-x^n a_n n(n-1)) + \left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) \\
 & + \sum_{n=1}^{\infty} (-2n a_n x^n) + \left(\sum_{n=0}^{\infty} (\alpha^2 + \alpha) a_n x^n \right) = 0
 \end{aligned} \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\begin{aligned} \sum_{n=2}^{\infty} (-x^n a_n n(n-1)) + \left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n \right) \\ + \sum_{n=1}^{\infty} (-2n a_n x^n) + \left(\sum_{n=0}^{\infty} (\alpha^2 + \alpha) a_n x^n \right) = 0 \end{aligned} \quad (3)$$

$n = 0$ gives

$$2a_2 + a_0\alpha(\alpha + 1) = 0$$

$$a_2 = -\frac{1}{2}a_0\alpha^2 - \frac{1}{2}a_0\alpha$$

$n = 1$ gives

$$6a_3 - 2a_1 + a_1\alpha(\alpha + 1) = 0$$

Which after substituting earlier equations, simplifies to

$$a_3 = -\frac{1}{6}a_1\alpha^2 - \frac{1}{6}a_1\alpha + \frac{1}{3}a_1$$

For $2 \leq n$, the recurrence equation is

$$-na_n(n-1) + (n+2) a_{n+2}(n+1) - 2na_n + a_n\alpha(\alpha + 1) = 0 \quad (4)$$

Solving for a_{n+2} , gives

$$a_{n+2} = -\frac{a_n(\alpha^2 - n^2 + \alpha - n)}{(n+2)(n+1)} \quad (5)$$

For $n = 2$ the recurrence equation gives

$$-6a_2 + 12a_4 + a_2\alpha(\alpha + 1) = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = -\frac{5}{24}a_0\alpha^2 - \frac{1}{4}a_0\alpha + \frac{1}{24}a_0\alpha^4 + \frac{1}{12}a_0\alpha^3$$

For $n = 3$ the recurrence equation gives

$$-12a_3 + 20a_5 + a_3\alpha(\alpha + 1) = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = -\frac{13}{120}a_1\alpha^2 - \frac{7}{60}a_1\alpha + \frac{1}{5}a_1 + \frac{1}{120}a_1\alpha^4 + \frac{1}{60}a_1\alpha^3$$

For $n = 4$ the recurrence equation gives

$$-20a_4 + 30a_6 + a_4\alpha(\alpha + 1) = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = -\frac{47}{360}a_0\alpha^2 - \frac{1}{6}a_0\alpha + \frac{23}{720}a_0\alpha^4 + \frac{17}{240}a_0\alpha^3 - \frac{1}{720}a_0\alpha^6 - \frac{1}{240}a_0\alpha^5$$

For $n = 5$ the recurrence equation gives

$$-30a_5 + 42a_7 + a_5\alpha(\alpha + 1) = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = -\frac{5}{63}a_1\alpha^2 - \frac{37}{420}a_1\alpha + \frac{1}{7}a_1 + \frac{41}{5040}a_1\alpha^4 + \frac{29}{1680}a_1\alpha^3 - \frac{1}{5040}a_1\alpha^6 - \frac{1}{1680}a_1\alpha^5$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$\begin{aligned} y = & a_0 + a_1x + \left(-\frac{1}{2}a_0\alpha^2 - \frac{1}{2}a_0\alpha\right)x^2 + \left(-\frac{1}{6}a_1\alpha^2 - \frac{1}{6}a_1\alpha + \frac{1}{3}a_1\right)x^3 \\ & + \left(-\frac{5}{24}a_0\alpha^2 - \frac{1}{4}a_0\alpha + \frac{1}{24}a_0\alpha^4 + \frac{1}{12}a_0\alpha^3\right)x^4 \\ & + \left(-\frac{13}{120}a_1\alpha^2 - \frac{7}{60}a_1\alpha + \frac{1}{5}a_1 + \frac{1}{120}a_1\alpha^4 + \frac{1}{60}a_1\alpha^3\right)x^5 + \dots \end{aligned}$$

Collecting terms, the solution becomes

$$\begin{aligned} y = & \left(1 + \left(-\frac{1}{2}\alpha^2 - \frac{1}{2}\alpha\right)x^2 + \left(-\frac{5}{24}\alpha^2 - \frac{1}{4}\alpha + \frac{1}{24}\alpha^4 + \frac{1}{12}\alpha^3\right)x^4\right)a_0 + \left(x \right. \\ & \left. + \left(-\frac{1}{6}\alpha^2 - \frac{1}{6}\alpha + \frac{1}{3}\right)x^3 + \left(-\frac{13}{120}\alpha^2 - \frac{7}{60}\alpha + \frac{1}{5} + \frac{1}{120}\alpha^4 + \frac{1}{60}\alpha^3\right)x^5\right)a_1 + O(x^6) \end{aligned} \quad (3)$$

At $x = 0$ the solution above becomes

$$\begin{aligned} y = & \left(1 + \left(-\frac{1}{2}\alpha^2 - \frac{1}{2}\alpha\right)x^2 + \left(-\frac{5}{24}\alpha^2 - \frac{1}{4}\alpha + \frac{1}{24}\alpha^4 + \frac{1}{12}\alpha^3\right)x^4\right)c_1 + \left(x \right. \\ & \left. + \left(-\frac{1}{6}\alpha^2 - \frac{1}{6}\alpha + \frac{1}{3}\right)x^3 + \left(-\frac{13}{120}\alpha^2 - \frac{7}{60}\alpha + \frac{1}{5} + \frac{1}{120}\alpha^4 + \frac{1}{60}\alpha^3\right)x^5\right)c_2 + O(x^6) \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned} y = & \left(1 - \frac{1}{2}x^2\alpha^2 - \frac{1}{2}x^2\alpha + \frac{1}{24}\alpha^4x^4 + \frac{1}{12}\alpha^3x^4 - \frac{5}{24}\alpha^2x^4 - \frac{1}{4}\alpha x^4 - \frac{1}{720}x^6\alpha^6 - \frac{1}{240}x^6\alpha^5 \right. \\ & \left. + \frac{23}{720}x^6\alpha^4 + \frac{17}{240}x^6\alpha^3 - \frac{47}{360}x^6\alpha^2 - \frac{1}{6}x^6\alpha\right)y(0) + \left(x - \frac{1}{6}\alpha^2x^3 - \frac{1}{6}\alpha x^3 + \frac{1}{3}x^3 \right. \\ & \left. + \frac{1}{120}x^5\alpha^4 + \frac{1}{60}x^5\alpha^3 - \frac{13}{120}x^5\alpha^2 - \frac{7}{60}x^5\alpha + \frac{1}{5}x^5\right)y'(0) + O(x^6) \end{aligned}$$

$$\begin{aligned} y = & \left(1 + \left(-\frac{1}{2}\alpha^2 - \frac{1}{2}\alpha\right)x^2 + \left(-\frac{5}{24}\alpha^2 - \frac{1}{4}\alpha + \frac{1}{24}\alpha^4 + \frac{1}{12}\alpha^3\right)x^4\right)c_1 \\ & + \left(x + \left(-\frac{1}{6}\alpha^2 - \frac{1}{6}\alpha + \frac{1}{3}\right)x^3 + \left(-\frac{13}{120}\alpha^2 - \frac{7}{60}\alpha + \frac{1}{5} + \frac{1}{120}\alpha^4 + \frac{1}{60}\alpha^3\right)x^5\right)c_2 \\ & + O(x^6) \end{aligned}$$

Verification of solutions

$$y = \left(1 - \frac{1}{2}x^2\alpha^2 - \frac{1}{2}x^2\alpha + \frac{1}{24}\alpha^4x^4 + \frac{1}{12}\alpha^3x^4 - \frac{5}{24}\alpha^2x^4 - \frac{1}{4}\alpha x^4 - \frac{1}{720}x^6\alpha^6 - \frac{1}{240}x^6\alpha^5 + \frac{23}{720}x^6\alpha^4 + \frac{17}{240}x^6\alpha^3 - \frac{47}{360}x^6\alpha^2 - \frac{1}{6}x^6\alpha\right) y(0) + \left(x - \frac{1}{6}\alpha^2x^3 - \frac{1}{6}\alpha x^3 + \frac{1}{3}x^3 + \frac{1}{120}x^5\alpha^4 + \frac{1}{60}x^5\alpha^3 - \frac{13}{120}x^5\alpha^2 - \frac{7}{60}x^5\alpha + \frac{1}{5}x^5\right) y'(0) + O(x^6)$$

Verified OK.

$$y = \left(1 + \left(-\frac{1}{2}\alpha^2 - \frac{1}{2}\alpha\right)x^2 + \left(-\frac{5}{24}\alpha^2 - \frac{1}{4}\alpha + \frac{1}{24}\alpha^4 + \frac{1}{12}\alpha^3\right)x^4\right) c_1 + \left(x + \left(-\frac{1}{6}\alpha^2 - \frac{1}{6}\alpha + \frac{1}{3}\right)x^3 + \left(-\frac{13}{120}\alpha^2 - \frac{7}{60}\alpha + \frac{1}{5} + \frac{1}{120}\alpha^4 + \frac{1}{60}\alpha^3\right)x^5\right) c_2 + O(x^6)$$

Verified OK.

15.10.1 Maple step by step solution

Let's solve

$$(-x^2 + 1)y'' - 2xy' + (\alpha^2 + \alpha)y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = \frac{\alpha(\alpha+1)y}{x^2-1} - \frac{2xy'}{x^2-1}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{2xy'}{x^2-1} - \frac{\alpha(\alpha+1)y}{x^2-1} = 0$$

- Check to see if x_0 is a regular singular point

- Define functions

$$\left[P_2(x) = \frac{2x}{x^2-1}, P_3(x) = -\frac{\alpha(\alpha+1)}{x^2-1} \right]$$

- $(1+x) \cdot P_2(x)$ is analytic at $x = -1$

$$\left. ((1+x) \cdot P_2(x)) \right|_{x=-1} = 1$$

- $(1+x)^2 \cdot P_3(x)$ is analytic at $x = -1$

$$((1+x)^2 \cdot P_3(x)) \Big|_{x=-1} = 0$$

- $x = -1$ is a regular singular point

Check to see if x_0 is a regular singular point

$$x_0 = -1$$

- Multiply by denominators

$$(x^2 - 1)y'' + 2xy' - \alpha(\alpha + 1)y = 0$$

- Change variables using $x = u - 1$ so that the regular singular point is at $u = 0$

$$(u^2 - 2u) \left(\frac{d^2}{du^2} y(u) \right) + (2u - 2) \left(\frac{d}{du} y(u) \right) + (-\alpha^2 - \alpha) y(u) = 0$$

- Assume series solution for $y(u)$

$$y(u) = \sum_{k=0}^{\infty} a_k u^{k+r}$$

- Rewrite ODE with series expansions

- Convert $u^m \cdot \left(\frac{d}{du} y(u) \right)$ to series expansion for $m = 0..1$

$$u^m \cdot \left(\frac{d}{du} y(u) \right) = \sum_{k=0}^{\infty} a_k (k+r) u^{k+r-1+m}$$

- Shift index using $k \rightarrow k + 1 - m$

$$u^m \cdot \left(\frac{d}{du} y(u) \right) = \sum_{k=-1+m}^{\infty} a_{k+1-m} (k+1-m+r) u^{k+r}$$

- Convert $u^m \cdot \left(\frac{d^2}{du^2} y(u) \right)$ to series expansion for $m = 1..2$

$$u^m \cdot \left(\frac{d^2}{du^2} y(u) \right) = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) u^{k+r-2+m}$$

- Shift index using $k \rightarrow k + 2 - m$

$$u^m \cdot \left(\frac{d^2}{du^2} y(u) \right) = \sum_{k=-2+m}^{\infty} a_{k+2-m} (k+2-m+r)(k+1-m+r) u^{k+r}$$

Rewrite ODE with series expansions

$$-2a_0 r^2 u^{-1+r} + \left(\sum_{k=0}^{\infty} (-2a_{k+1} (k+1+r)^2 - a_k (r+1+k+\alpha)(-r-k+\alpha)) u^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$-2r^2 = 0$$

- Values of r that satisfy the indicial equation

$$r = 0$$

- Each term in the series must be 0, giving the recursion relation

$$-2a_{k+1}(k+1)^2 + a_k(1+k+\alpha)(k-\alpha) = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+1} = -\frac{a_k(1+k+\alpha)(-k+\alpha)}{2(k+1)^2}$$

- Recursion relation for $r = 0$

$$a_{k+1} = -\frac{a_k(1+k+\alpha)(-k+\alpha)}{2(k+1)^2}$$

- Solution for $r = 0$

$$\left[y(u) = \sum_{k=0}^{\infty} a_k u^k, a_{k+1} = -\frac{a_k(1+k+\alpha)(-k+\alpha)}{2(k+1)^2} \right]$$

- Revert the change of variables $u = 1 + x$

$$\left[y = \sum_{k=0}^{\infty} a_k (1+x)^k, a_{k+1} = -\frac{a_k(1+k+\alpha)(-k+\alpha)}{2(k+1)^2} \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
  -> Bessel
  -> elliptic
  -> Legendre
  <- Legendre successful
<- special function solution successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 101

Order:=6;

```
dsolve((1-x^2)*diff(y(x),x$2)-2*x*diff(y(x),x)+alpha*(alpha+1)*y(x)=0,y(x),type='series',x=0
```

$$y(x) = \left(1 - \frac{\alpha(\alpha+1)x^2}{2} + \frac{\alpha(\alpha^3+2\alpha^2-5\alpha-6)x^4}{24}\right) y(0) \\ + \left(x - \frac{(\alpha^2+\alpha-2)x^3}{6} + \frac{(\alpha^4+2\alpha^3-13\alpha^2-14\alpha+24)x^5}{120}\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 127

```
AsymptoticDSolveValue[(1-x^2)*y'[x]-2*x*y'[x]+\[Alpha]*(\[Alpha]+1)*y[x]==0,y[x],{x,0,5}]
```

$$y(x) \rightarrow c_2 \left(\frac{1}{60}(-\alpha^2 - \alpha)x^5 - \frac{1}{120}(-\alpha^2 - \alpha)(\alpha^2 + \alpha)x^5 - \frac{1}{10}(\alpha^2 + \alpha)x^5 + \frac{x^5}{5} \right. \\ \left. - \frac{1}{6}(\alpha^2 + \alpha)x^3 + \frac{x^3}{3} + x \right) + c_1 \left(\frac{1}{24}(\alpha^2 + \alpha)^2 x^4 - \frac{1}{4}(\alpha^2 + \alpha)x^4 - \frac{1}{2}(\alpha^2 + \alpha)x^2 + 1 \right)$$

15.11 problem 7

- 15.11.1 Solving as second order change of variable on x method 2 ode . 1143
- 15.11.2 Solving as second order change of variable on x method 1 ode . 1146
- 15.11.3 Solving using Kovacic algorithm 1148
- 15.11.4 Maple step by step solution 1154

Internal problem ID [6029]

Internal file name [OUTPUT/5277_Sunday_June_05_2022_03_29_16_PM_49957946/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 130

Problem number: 7.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_change_of_variable_on_x_method_1", "second_order_change_of_variable_on_x_method_2"

Maple gives the following as the ode type

```
[_Gegenbauer , [_2nd_order , _linear , ` _with_symmetry_[0,F(x)]`]]
```

$$(-x^2 + 1) y'' - xy' + \alpha^2 y = 0$$

15.11.1 Solving as second order change of variable on x method 2 ode

In normal form the ode

$$(-x^2 + 1) y'' - xy' + \alpha^2 y = 0 \tag{1}$$

Becomes

$$y'' + p(x) y' + q(x) y = 0 \tag{2}$$

Where

$$p(x) = \frac{x}{x^2 - 1}$$
$$q(x) = \frac{\alpha^2}{-x^2 + 1}$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x)\tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned} \tau &= \int e^{-(\int p(x)dx)} dx \\ &= \int e^{-\left(\int \frac{x}{x^2-1} dx\right)} dx \\ &= \int e^{-\frac{\ln(x-1)}{2} - \frac{\ln(1+x)}{2}} dx \\ &= \int \frac{1}{\sqrt{x-1}\sqrt{1+x}} dx \\ &= \frac{\sqrt{(x-1)(1+x)} \ln(x + \sqrt{x^2-1})}{\sqrt{x-1}\sqrt{1+x}} \end{aligned} \quad (6)$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned} q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\ &= \frac{\alpha^2}{\frac{-x^2+1}{1}} \\ &= -\alpha^2 \end{aligned} \quad (7)$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned} \frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) - \alpha^2y(\tau) &= 0 \end{aligned}$$

The above ode is now solved for $y(\tau)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(\tau) + By'(\tau) + Cy(\tau) = 0$$

Where in the above $A = 1, B = 0, C = -\alpha^2$. Let the solution be $y(\tau) = e^{\lambda\tau}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda\tau} - \alpha^2 e^{\lambda\tau} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda\tau}$ gives

$$-\alpha^2 + \lambda^2 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = -\alpha^2$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(-\alpha^2)} \\ &= \pm \sqrt{\alpha^2} \end{aligned}$$

Hence

$$\lambda_1 = +\sqrt{\alpha^2}$$

$$\lambda_2 = -\sqrt{\alpha^2}$$

Which simplifies to

$$\lambda_1 = \sqrt{\alpha^2}$$

$$\lambda_2 = -\sqrt{\alpha^2}$$

Since roots are real and distinct, then the solution is

$$y(\tau) = c_1 e^{\lambda_1 \tau} + c_2 e^{\lambda_2 \tau}$$

$$y(\tau) = c_1 e^{(\sqrt{\alpha^2})\tau} + c_2 e^{(-\sqrt{\alpha^2})\tau}$$

Or

$$y(\tau) = c_1 e^{\sqrt{\alpha^2} \tau} + c_2 e^{-\sqrt{\alpha^2} \tau}$$

The above solution is now transformed back to y using (6) which results in

$$y = c_1 \left(x + \sqrt{x^2 - 1} \right)^\alpha + c_2 \left(x + \sqrt{x^2 - 1} \right)^{-\alpha}$$

Summary

The solution(s) found are the following

$$y = c_1 \left(x + \sqrt{x^2 - 1} \right)^\alpha + c_2 \left(x + \sqrt{x^2 - 1} \right)^{-\alpha} \quad (1)$$

Verification of solutions

$$y = c_1 \left(x + \sqrt{x^2 - 1} \right)^\alpha + c_2 \left(x + \sqrt{x^2 - 1} \right)^{-\alpha}$$

Verified OK.

15.11.2 Solving as second order change of variable on x method 1 ode

In normal form the ode

$$(-x^2 + 1) y'' - xy' + \alpha^2 y = 0 \quad (1)$$

Becomes

$$y'' + p(x) y' + q(x) y = 0 \quad (2)$$

Where

$$p(x) = \frac{x}{x^2 - 1}$$

$$q(x) = -\frac{\alpha^2}{x^2 - 1}$$

Applying change of variables $\tau = g(x)$ to (2) results

$$\frac{d^2}{d\tau^2} y(\tau) + p_1 \left(\frac{d}{d\tau} y(\tau) \right) + q_1 y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x) \tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $q_1 = c^2$ where c is some constant. Therefore from (5)

$$\begin{aligned}\tau' &= \frac{1}{c}\sqrt{q} \\ &= \frac{\sqrt{-\frac{\alpha^2}{x^2-1}}}{c} \\ \tau'' &= \frac{\alpha^2 x}{c\sqrt{-\frac{\alpha^2}{x^2-1}}(x^2-1)^2}\end{aligned}\tag{6}$$

Substituting the above into (4) results in

$$\begin{aligned}p_1(\tau) &= \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \\ &= \frac{\frac{\alpha^2 x}{c\sqrt{-\frac{\alpha^2}{x^2-1}}(x^2-1)^2} + \frac{x}{x^2-1}\frac{\sqrt{-\frac{\alpha^2}{x^2-1}}}{c}}{\left(\frac{\sqrt{-\frac{\alpha^2}{x^2-1}}}{c}\right)^2} \\ &= 0\end{aligned}$$

Therefore ode (3) now becomes

$$\begin{aligned}y(\tau)'' + p_1 y(\tau)' + q_1 y(\tau) &= 0 \\ \frac{d^2}{d\tau^2} y(\tau) + c^2 y(\tau) &= 0\end{aligned}\tag{7}$$

The above ode is now solved for $y(\tau)$. Since the ode is now constant coefficients, it can be easily solved to give

$$y(\tau) = c_1 \cos(c\tau) + c_2 \sin(c\tau)$$

Now from (6)

$$\begin{aligned}\tau &= \int \frac{1}{c}\sqrt{q} dx \\ &= \frac{\int \sqrt{-\frac{\alpha^2}{x^2-1}} dx}{c} \\ &= \frac{\sqrt{-\frac{\alpha^2}{x^2-1}} \sqrt{x^2-1} \ln(x + \sqrt{x^2-1})}{c}\end{aligned}$$

Substituting the above into the solution obtained gives

$$y = c_1 \cos \left(\alpha \sqrt{-\frac{1}{x^2 - 1}} \sqrt{x^2 - 1} \ln (x + \sqrt{x^2 - 1}) \right) \\ + c_2 \sin \left(\alpha \sqrt{-\frac{1}{x^2 - 1}} \sqrt{x^2 - 1} \ln (x + \sqrt{x^2 - 1}) \right)$$

Summary

The solution(s) found are the following

$$y = c_1 \cos \left(\alpha \sqrt{-\frac{1}{x^2 - 1}} \sqrt{x^2 - 1} \ln (x + \sqrt{x^2 - 1}) \right) \\ + c_2 \sin \left(\alpha \sqrt{-\frac{1}{x^2 - 1}} \sqrt{x^2 - 1} \ln (x + \sqrt{x^2 - 1}) \right) \quad (1)$$

Verification of solutions

$$y = c_1 \cos \left(\alpha \sqrt{-\frac{1}{x^2 - 1}} \sqrt{x^2 - 1} \ln (x + \sqrt{x^2 - 1}) \right) \\ + c_2 \sin \left(\alpha \sqrt{-\frac{1}{x^2 - 1}} \sqrt{x^2 - 1} \ln (x + \sqrt{x^2 - 1}) \right)$$

Verified OK.

15.11.3 Solving using Kovacic algorithm

Writing the ode as

$$(-x^2 + 1) y'' - xy' + \alpha^2 y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$A = -x^2 + 1 \\ B = -x \\ C = \alpha^2 \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{4x^2\alpha^2 - 4\alpha^2 - x^2 - 2}{4(x^2 - 1)^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 4x^2\alpha^2 - 4\alpha^2 - x^2 - 2 \\ t &= 4(x^2 - 1)^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{4x^2\alpha^2 - 4\alpha^2 - x^2 - 2}{4(x^2 - 1)^2} \right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 190: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 4 - 2 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4(x^2 - 1)^2$. There is a pole at $x = 1$ of order 2. There is a pole at $x = -1$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Unable to find solution using case one

Attempting to find a solution using case $n = 2$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = -\frac{3}{16(x-1)^2} + \frac{\frac{1}{16} + \frac{\alpha^2}{2}}{x-1} - \frac{3}{16(1+x)^2} + \frac{-\frac{\alpha^2}{2} - \frac{1}{16}}{1+x}$$

For the pole at $x = 1$ let b be the coefficient of $\frac{1}{(x-1)^2}$ in the partial fractions decomposition of r given above. Therefore $b = -\frac{3}{16}$. Hence

$$\begin{aligned} E_c &= \{2, 2 + 2\sqrt{1 + 4b}, 2 - 2\sqrt{1 + 4b}\} \\ &= \{1, 2, 3\} \end{aligned}$$

For the pole at $x = -1$ let b be the coefficient of $\frac{1}{(1+x)^2}$ in the partial fractions decomposition of r given above. Therefore $b = -\frac{3}{16}$. Hence

$$\begin{aligned} E_c &= \{2, 2 + 2\sqrt{1 + 4b}, 2 - 2\sqrt{1 + 4b}\} \\ &= \{1, 2, 3\} \end{aligned}$$

Since the order of r at ∞ is 2 then let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = \frac{4x^2\alpha^2 - 4\alpha^2 - x^2 - 2}{4(x^2 - 1)^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = 1$. Hence

$$\begin{aligned} E_\infty &= \{2, 2 + 2\sqrt{1 + 4b}, 2 - 2\sqrt{1 + 4b}\} \\ &= \{2\} \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ for case 2 of Kovacic algorithm.

pole c location	pole order	E_c
1	2	{1, 2, 3}
-1	2	{1, 2, 3}

Order of r at ∞	E_∞
2	{2}

Using the family $\{e_1, e_2, \dots, e_\infty\}$ given by

$$e_1 = 1, e_2 = 1, e_\infty = 2$$

Gives a non negative integer d (the degree of the polynomial $p(x)$), which is generated using

$$\begin{aligned} d &= \frac{1}{2} \left(e_\infty - \sum_{c \in \Gamma} e_c \right) \\ &= \frac{1}{2} (2 - (1 + (1))) \\ &= 0 \end{aligned}$$

We now form the following rational function

$$\begin{aligned} \theta &= \frac{1}{2} \sum_{c \in \Gamma} \frac{e_c}{x - c} \\ &= \frac{1}{2} \left(\frac{1}{(x - (1))} + \frac{1}{(x - (-1))} \right) \\ &= \frac{1}{2x - 2} + \frac{1}{2 + 2x} \end{aligned}$$

Now we search for a monic polynomial $p(x)$ of degree $d = 0$ such that

$$p''' + 3\theta p'' + (3\theta^2 + 3\theta' - 4r)p' + (\theta'' + 3\theta\theta' + \theta^3 - 4r\theta - 2r')p = 0 \quad (1A)$$

Since $d = 0$, then letting

$$p = 1 \quad (2A)$$

Substituting p and θ into Eq. (1A) gives

$$0 = 0$$

And solving for p gives

$$p = 1$$

Now that $p(x)$ is found let

$$\begin{aligned}\phi &= \theta + \frac{p'}{p} \\ &= \frac{1}{2x-2} + \frac{1}{2+2x}\end{aligned}$$

Let ω be the solution of

$$\omega^2 - \phi\omega + \left(\frac{1}{2}\phi' + \frac{1}{2}\phi^2 - r\right) = 0$$

Substituting the values for ϕ and r into the above equation gives

$$w^2 - \left(\frac{1}{2x-2} + \frac{1}{2+2x}\right)w + \frac{-4x^2\alpha^2 + 4\alpha^2 + x^2}{4(x^2-1)^2} = 0$$

Solving for ω gives

$$\omega = \frac{x + 2\alpha\sqrt{x^2-1}}{2(x-1)(1+x)}$$

Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned}z_1(x) &= e^{\int \omega dx} \\ &= e^{\int \frac{x+2\alpha\sqrt{x^2-1}}{2(x-1)(1+x)} dx} \\ &= (x^2-1)^{\frac{1}{4}} \left(x + \sqrt{x^2-1}\right)^\alpha\end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-x}{-x^2+1} dx} \\ &= z_1 e^{-\frac{\ln(x-1)}{4} - \frac{\ln(1+x)}{4}} \\ &= z_1 \left(\frac{1}{(x-1)^{\frac{1}{4}} (1+x)^{\frac{1}{4}}}\right)\end{aligned}$$

Which simplifies to

$$y_1 = \frac{(x^2 - 1)^{\frac{1}{4}} (x + \sqrt{x^2 - 1})^\alpha}{(x - 1)^{\frac{1}{4}} (1 + x)^{\frac{1}{4}}}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-x}{-x^2+1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-\frac{\ln(x-1)}{2} - \frac{\ln(1+x)}{2}}}{(y_1)^2} dx \\ &= y_1 \left(-\frac{(x + \sqrt{x^2 - 1})^{-2\alpha}}{2\alpha} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(\frac{(x^2 - 1)^{\frac{1}{4}} (x + \sqrt{x^2 - 1})^\alpha}{(x - 1)^{\frac{1}{4}} (1 + x)^{\frac{1}{4}}} \right) \\ &\quad + c_2 \left(\frac{(x^2 - 1)^{\frac{1}{4}} (x + \sqrt{x^2 - 1})^\alpha}{(x - 1)^{\frac{1}{4}} (1 + x)^{\frac{1}{4}}} \left(-\frac{(x + \sqrt{x^2 - 1})^{-2\alpha}}{2\alpha} \right) \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1 (x^2 - 1)^{\frac{1}{4}} (x + \sqrt{x^2 - 1})^\alpha}{(x - 1)^{\frac{1}{4}} (1 + x)^{\frac{1}{4}}} - \frac{c_2 (x^2 - 1)^{\frac{1}{4}} (x + \sqrt{x^2 - 1})^{-\alpha}}{2\alpha (x - 1)^{\frac{1}{4}} (1 + x)^{\frac{1}{4}}} \quad (1)$$

Verification of solutions

$$y = \frac{c_1 (x^2 - 1)^{\frac{1}{4}} (x + \sqrt{x^2 - 1})^\alpha}{(x - 1)^{\frac{1}{4}} (1 + x)^{\frac{1}{4}}} - \frac{c_2 (x^2 - 1)^{\frac{1}{4}} (x + \sqrt{x^2 - 1})^{-\alpha}}{2\alpha (x - 1)^{\frac{1}{4}} (1 + x)^{\frac{1}{4}}}$$

Verified OK.

15.11.4 Maple step by step solution

Let's solve

$$(-x^2 + 1)y'' - xy' + \alpha^2 y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{xy'}{x^2-1} + \frac{\alpha^2 y}{x^2-1}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{xy'}{x^2-1} - \frac{\alpha^2 y}{x^2-1} = 0$$

- Check to see if x_0 is a regular singular point

- o Define functions

$$\left[P_2(x) = \frac{x}{x^2-1}, P_3(x) = -\frac{\alpha^2}{x^2-1} \right]$$

- o $(1+x) \cdot P_2(x)$ is analytic at $x = -1$

$$\left. ((1+x) \cdot P_2(x)) \right|_{x=-1} = \frac{1}{2}$$

- o $(1+x)^2 \cdot P_3(x)$ is analytic at $x = -1$

$$\left. ((1+x)^2 \cdot P_3(x)) \right|_{x=-1} = 0$$

- o $x = -1$ is a regular singular point

Check to see if x_0 is a regular singular point

$$x_0 = -1$$

- Multiply by denominators

$$(x^2 - 1)y'' + xy' - \alpha^2 y = 0$$

- Change variables using $x = u - 1$ so that the regular singular point is at $u = 0$

$$(u^2 - 2u) \left(\frac{d^2}{du^2} y(u) \right) + (u - 1) \left(\frac{d}{du} y(u) \right) - \alpha^2 y(u) = 0$$

- Assume series solution for $y(u)$

$$y(u) = \sum_{k=0}^{\infty} a_k u^{k+r}$$

- Rewrite ODE with series expansions

- Convert $u^m \cdot \left(\frac{d}{du}y(u)\right)$ to series expansion for $m = 0..1$

$$u^m \cdot \left(\frac{d}{du}y(u)\right) = \sum_{k=0}^{\infty} a_k(k+r) u^{k+r-1+m}$$

- Shift index using $k \rightarrow k+1-m$

$$u^m \cdot \left(\frac{d}{du}y(u)\right) = \sum_{k=-1+m}^{\infty} a_{k+1-m}(k+1-m+r) u^{k+r}$$

- Convert $u^m \cdot \left(\frac{d^2}{du^2}y(u)\right)$ to series expansion for $m = 1..2$

$$u^m \cdot \left(\frac{d^2}{du^2}y(u)\right) = \sum_{k=0}^{\infty} a_k(k+r)(k+r-1) u^{k+r-2+m}$$

- Shift index using $k \rightarrow k+2-m$

$$u^m \cdot \left(\frac{d^2}{du^2}y(u)\right) = \sum_{k=-2+m}^{\infty} a_{k+2-m}(k+2-m+r)(k+1-m+r) u^{k+r}$$

Rewrite ODE with series expansions

$$-a_0r(-1+2r)u^{-1+r} + \left(\sum_{k=0}^{\infty} (-a_{1+k}(1+k+r)(1+2k+2r) - a_k(\alpha+k+r)(\alpha-k-r)) u^{k+r}\right)$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$-r(-1+2r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \left\{0, \frac{1}{2}\right\}$$

- Each term in the series must be 0, giving the recursion relation

$$-2\left(\frac{1}{2} + k + r\right)(1+k+r)a_{1+k} + a_k(\alpha+k+r)(k+r-\alpha) = 0$$

- Recursion relation that defines series solution to ODE

$$a_{1+k} = -\frac{a_k(\alpha+k+r)(\alpha-k-r)}{(1+2k+2r)(1+k+r)}$$

- Recursion relation for $r = 0$

$$a_{1+k} = -\frac{a_k(\alpha+k)(\alpha-k)}{(1+2k)(1+k)}$$

- Solution for $r = 0$

$$\left[y(u) = \sum_{k=0}^{\infty} a_k u^k, a_{1+k} = -\frac{a_k(\alpha+k)(\alpha-k)}{(1+2k)(1+k)} \right]$$

- Revert the change of variables $u = 1+x$

$$\left[y = \sum_{k=0}^{\infty} a_k(1+x)^k, a_{1+k} = -\frac{a_k(\alpha+k)(\alpha-k)}{(1+2k)(1+k)} \right]$$

- Recursion relation for $r = \frac{1}{2}$

$$a_{1+k} = -\frac{a_k(\alpha+k+\frac{1}{2})(\alpha-k-\frac{1}{2})}{(2+2k)(\frac{3}{2}+k)}$$

- Solution for $r = \frac{1}{2}$

$$\left[y(u) = \sum_{k=0}^{\infty} a_k u^{k+\frac{1}{2}}, a_{1+k} = -\frac{a_k(\alpha+k+\frac{1}{2})(\alpha-k-\frac{1}{2})}{(2+2k)(\frac{3}{2}+k)} \right]$$

- Revert the change of variables $u = 1 + x$

$$\left[y = \sum_{k=0}^{\infty} a_k (1+x)^{k+\frac{1}{2}}, a_{1+k} = -\frac{a_k(\alpha+k+\frac{1}{2})(\alpha-k-\frac{1}{2})}{(2+2k)(\frac{3}{2}+k)} \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k (1+x)^k \right) + \left(\sum_{k=0}^{\infty} b_k (1+x)^{k+\frac{1}{2}} \right), a_{k+1} = -\frac{a_k(\alpha+k)(\alpha-k)}{(1+2k)(k+1)}, b_{k+1} = -\frac{b_k(\alpha+k+\frac{1}{2})(\alpha-k-\frac{1}{2})}{(2k+2)(\frac{3}{2}+k)} \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
<- linear_1 successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 33

```
dsolve((1-x^2)*diff(y(x),x$2)-x*diff(y(x),x)+alpha^2*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 \left(x + \sqrt{x^2 - 1} \right)^{-\alpha} + c_2 \left(x + \sqrt{x^2 - 1} \right)^{\alpha}$$

✓ Solution by Mathematica

Time used: 0.089 (sec). Leaf size: 91

```
DSolve[(1-x^2)*y'[x]-x*y'[x]+\[Alpha]^2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cosh \left(\frac{1}{2} \alpha \left(\log \left(1 - \frac{x}{\sqrt{x^2 - 1}} \right) - \log \left(\frac{x}{\sqrt{x^2 - 1}} + 1 \right) \right) \right) - i c_2 \sinh \left(\frac{1}{2} \alpha \left(\log \left(1 - \frac{x}{\sqrt{x^2 - 1}} \right) - \log \left(\frac{x}{\sqrt{x^2 - 1}} + 1 \right) \right) \right)$$

15.12 problem 8

15.12.1 Maple step by step solution 1158

Internal problem ID [6030]

Internal file name [OUTPUT/5278_Sunday_June_05_2022_03_29_18_PM_68325152/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 3. Linear equations with variable coefficients. Page 130

Problem number: 8.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**unknown**"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

Unable to solve or complete the solution.

$$y'' - 2xy' + 2\alpha y = 0$$

15.12.1 Maple step by step solution

Let's solve

$$y'' - 2xy' + 2\alpha y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^k$$

- Rewrite DE with series expansions

- Convert $x \cdot y'$ to series expansion

$$x \cdot y' = \sum_{k=0}^{\infty} a_k k x^k$$

- Convert y'' to series expansion

$$y'' = \sum_{k=2}^{\infty} a_k k(k-1) x^{k-2}$$

- Shift index using $k \rightarrow k+2$

$$y'' = \sum_{k=0}^{\infty} a_{k+2}(k+2)(k+1)x^k$$

Rewrite DE with series expansions

$$\sum_{k=0}^{\infty} (a_{k+2}(k+2)(k+1) + 2a_k(\alpha - k)) x^k = 0$$

- Each term in the series must be 0, giving the recursion relation

$$(k^2 + 3k + 2) a_{k+2} - 2a_k(k - \alpha) = 0$$
- Recursion relation that defines the series solution to the ODE

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+2} = -\frac{2a_k(\alpha - k)}{k^2 + 3k + 2} \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
  -> Bessel
  -> elliptic
  -> Legendre
  -> Kummer
    -> hyper3: Equivalence to 1F1 under a power @ Moebius
    <- hyper3 successful: received ODE is equivalent to the 1F1 ODE
  <- Kummer successful
<- special function solution successful`

```


✓ Solution by Maple

Time used: 0.079 (sec). Leaf size: 31

```
dsolve(diff(y(x),x$2)-2*x*diff(y(x),x)+2*alpha*y(x)=0,y(x), singsol=all)
```

$$y(x) = x \left(\text{KummerM} \left(\frac{1}{2} - \frac{\alpha}{2}, \frac{3}{2}, x^2 \right) c_1 + \text{KummerU} \left(\frac{1}{2} - \frac{\alpha}{2}, \frac{3}{2}, x^2 \right) c_2 \right)$$

✓ Solution by Mathematica

Time used: 0.062 (sec). Leaf size: 91

```
DSolve[(1-x^2)*y'[x]-x*y'[x]+\[Alpha]^2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cosh \left(\frac{1}{2} \alpha \left(\log \left(1 - \frac{x}{\sqrt{x^2 - 1}} \right) - \log \left(\frac{x}{\sqrt{x^2 - 1}} + 1 \right) \right) \right) \\ - i c_2 \sinh \left(\frac{1}{2} \alpha \left(\log \left(1 - \frac{x}{\sqrt{x^2 - 1}} \right) - \log \left(\frac{x}{\sqrt{x^2 - 1}} + 1 \right) \right) \right)$$

16 Chapter 4. Linear equations with Regular Singular Points. Page 149

16.1	problem 1(a)	1162
16.2	problem 1(b)	1176
16.3	problem 1(c)	1193
16.4	problem 1(d)	1209
16.5	problem 1(e)	1237
16.6	problem 2(a)	1244
16.7	problem 2(b)	1272
16.8	problem 2(c)	1289
16.9	problem 2(d)	1304

16.1 problem 1(a)

- 16.1.1 Solving as second order euler ode ode 1162
- 16.1.2 Solving as second order change of variable on x method 2 ode . 1163
- 16.1.3 Solving as second order change of variable on y method 2 ode . 1166
- 16.1.4 Solving using Kovacic algorithm 1168
- 16.1.5 Maple step by step solution 1173

Internal problem ID [6031]

Internal file name [OUTPUT/5279_Sunday_June_05_2022_03_29_20_PM_47389931/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 149

Problem number: 1(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_euler_ode", "second_order_change_of_variable_on_x_method_2", "second_order_change_of_variable_on_y_method_2"

Maple gives the following as the ode type

```
[[_Emden, _Fowler], [_2nd_order, _linear, `_with_symmetry_[0,F(x)]`]]
```

$$x^2y'' + 2xy' - 6y = 0$$

16.1.1 Solving as second order euler ode ode

This is Euler second order ODE. Let the solution be $y = x^r$, then $y' = rx^{r-1}$ and $y'' = r(r-1)x^{r-2}$. Substituting these back into the given ODE gives

$$x^2(r(r-1))x^{r-2} + 2rxr^{r-1} - 6x^r = 0$$

Simplifying gives

$$r(r-1)x^r + 2rx^r - 6x^r = 0$$

Since $x^r \neq 0$ then dividing throughout by x^r gives

$$r(r-1) + 2r - 6 = 0$$

Or

$$r^2 + r - 6 = 0 \quad (1)$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = -3$$

$$r_2 = 2$$

Since the roots are real and distinct, then the general solution is

$$y = c_1 y_1 + c_2 y_2$$

Where $y_1 = x^{r_1}$ and $y_2 = x^{r_2}$. Hence

$$y = \frac{c_1}{x^3} + c_2 x^2$$

Summary

The solution(s) found are the following

$$y = \frac{c_1}{x^3} + c_2 x^2 \quad (1)$$

Verification of solutions

$$y = \frac{c_1}{x^3} + c_2 x^2$$

Verified OK.

16.1.2 Solving as second order change of variable on x method 2 ode

In normal form the ode

$$x^2 y'' + 2xy' - 6y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{2}{x}$$
$$q(x) = -\frac{6}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x)\tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned} \tau &= \int e^{-(\int p(x)dx)} dx \\ &= \int e^{-(\int \frac{2}{x} dx)} dx \\ &= \int e^{-2\ln(x)} dx \\ &= \int \frac{1}{x^2} dx \\ &= -\frac{1}{x} \end{aligned} \quad (6)$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned} q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\ &= \frac{-\frac{6}{x^2}}{\frac{1}{x^4}} \\ &= -6x^2 \end{aligned} \quad (7)$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned} \frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) - 6x^2y(\tau) &= 0 \end{aligned}$$

But in terms of τ

$$-6x^2 = -\frac{6}{\tau^2}$$

Hence the above ode becomes

$$\frac{d^2}{d\tau^2}y(\tau) - \frac{6y(\tau)}{\tau^2} = 0$$

The above ode is now solved for $y(\tau)$. The ode can be written as

$$\left(\frac{d^2}{d\tau^2}y(\tau)\right)\tau^2 - 6y(\tau) = 0$$

Which shows it is a Euler ODE. This is Euler second order ODE. Let the solution be $y(\tau) = \tau^r$, then $y' = r\tau^{r-1}$ and $y'' = r(r-1)\tau^{r-2}$. Substituting these back into the given ODE gives

$$\tau^2(r(r-1))\tau^{r-2} + 0r\tau^{r-1} - 6\tau^r = 0$$

Simplifying gives

$$r(r-1)\tau^r + 0\tau^r - 6\tau^r = 0$$

Since $\tau^r \neq 0$ then dividing throughout by τ^r gives

$$r(r-1) + 0 - 6 = 0$$

Or

$$r^2 - r - 6 = 0 \tag{1}$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = -2$$

$$r_2 = 3$$

Since the roots are real and distinct, then the general solution is

$$y(\tau) = c_1y_1 + c_2y_2$$

Where $y_1 = \tau^{r_1}$ and $y_2 = \tau^{r_2}$. Hence

$$y(\tau) = \frac{c_1}{\tau^2} + c_2\tau^3$$

The above solution is now transformed back to y using (6) which results in

$$y = \frac{c_1x^5 - c_2}{x^3}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1 x^5 - c_2}{x^3} \quad (1)$$

Verification of solutions

$$y = \frac{c_1 x^5 - c_2}{x^3}$$

Verified OK.

16.1.3 Solving as second order change of variable on y method 2 ode

In normal form the ode

$$x^2 y'' + 2xy' - 6y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{2}{x}$$
$$q(x) = -\frac{6}{x^2}$$

Applying change of variables on the dependent variable $y = v(x)x^n$ to (2) gives the following ode where the dependent variables is $v(x)$ and not y .

$$v''(x) + \left(\frac{2n}{x} + p\right)v'(x) + \left(\frac{n(n-1)}{x^2} + \frac{np}{x} + q\right)v(x) = 0 \quad (3)$$

Let the coefficient of $v(x)$ above be zero. Hence

$$\frac{n(n-1)}{x^2} + \frac{np}{x} + q = 0 \quad (4)$$

Substituting the earlier values found for $p(x)$ and $q(x)$ into (4) gives

$$\frac{n(n-1)}{x^2} + \frac{2n}{x^2} - \frac{6}{x^2} = 0 \quad (5)$$

Solving (5) for n gives

$$n = 2 \quad (6)$$

Substituting this value in (3) gives

$$\begin{aligned}v''(x) + \frac{6v'(x)}{x} &= 0 \\v''(x) + \frac{6v'(x)}{x} &= 0\end{aligned}\tag{7}$$

Using the substitution

$$u(x) = v'(x)$$

Then (7) becomes

$$u'(x) + \frac{6u(x)}{x} = 0\tag{8}$$

The above is now solved for $u(x)$. In canonical form the ODE is

$$\begin{aligned}u' &= F(x, u) \\&= f(x)g(u) \\&= -\frac{6u}{x}\end{aligned}$$

Where $f(x) = -\frac{6}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u} du &= -\frac{6}{x} dx \\ \int \frac{1}{u} du &= \int -\frac{6}{x} dx \\ \ln(u) &= -6 \ln(x) + c_1 \\ u &= e^{-6 \ln(x) + c_1} \\ &= \frac{c_1}{x^6}\end{aligned}$$

Now that $u(x)$ is known, then

$$\begin{aligned}v'(x) &= u(x) \\ v(x) &= \int u(x) dx + c_2 \\ &= -\frac{c_1}{5x^5} + c_2\end{aligned}$$

Hence

$$\begin{aligned}y &= v(x) x^n \\ &= \left(-\frac{c_1}{5x^5} + c_2\right) x^2 \\ &= \frac{5c_2x^5 - c_1}{5x^3}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \left(-\frac{c_1}{5x^5} + c_2\right) x^2 \quad (1)$$

Verification of solutions

$$y = \left(-\frac{c_1}{5x^5} + c_2\right) x^2$$

Verified OK.

16.1.4 Solving using Kovacic algorithm

Writing the ode as

$$x^2y'' + 2xy' - 6y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= x^2 \\ B &= 2x \\ C &= -6\end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{6}{x^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 6 \\ t &= x^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{6}{x^2}\right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 193: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = \frac{6}{x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = 6$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = 3 \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = -2 \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = \frac{6}{x^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = 6$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = 3 \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = -2 \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{6}{x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	3	-2

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	3	-2

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = -2$ then

$$\begin{aligned} d &= \alpha_\infty^- - (\alpha_{c_1}^-) \\ &= -2 - (-2) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{x - c_1} \right) + (-)[\sqrt{r}]_\infty \\ &= -\frac{2}{x} + (-)(0) \\ &= -\frac{2}{x} \\ &= -\frac{2}{x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$(0) + 2\left(-\frac{2}{x}\right)(0) + \left(\left(\frac{2}{x^2}\right) + \left(-\frac{2}{x}\right)^2 - \left(\frac{6}{x^2}\right)\right) = 0$$
$$0 = 0$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$z_1(x) = pe^{\int \omega dx}$$
$$= e^{\int -\frac{2}{x} dx}$$
$$= \frac{1}{x^2}$$

The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$
$$= z_1 e^{-\int \frac{1}{2} \frac{2x}{x^2} dx}$$
$$= z_1 e^{-\ln(x)}$$
$$= z_1 \left(\frac{1}{x}\right)$$

Which simplifies to

$$y_1 = \frac{1}{x^3}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$y_2 = y_1 \int \frac{e^{\int -\frac{2x}{x^2} dx}}{(y_1)^2} dx$$
$$= y_1 \int \frac{e^{-2\ln(x)}}{(y_1)^2} dx$$
$$= y_1 \left(\frac{x^5}{5}\right)$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(\frac{1}{x^3} \right) + c_2 \left(\frac{1}{x^3} \left(\frac{x^5}{5} \right) \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1}{x^3} + \frac{c_2 x^2}{5} \quad (1)$$

Verification of solutions

$$y = \frac{c_1}{x^3} + \frac{c_2 x^2}{5}$$

Verified OK.

16.1.5 Maple step by step solution

Let's solve

$$x^2 y'' + 2xy' - 6y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{2y'}{x} + \frac{6y}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{2y'}{x} - \frac{6y}{x^2} = 0$$

- Multiply by denominators of the ODE

$$x^2 y'' + 2xy' - 6y = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x , using the chain rule

$$y' = \left(\frac{d}{dt} y(t) \right) t'(x)$$

- Compute derivative

$$y' = \frac{\frac{d}{dt}y(t)}{x}$$

- Calculate the 2nd derivative of y with respect to x , using the chain rule

$$y'' = \left(\frac{d^2}{dt^2}y(t) \right) t'(x)^2 + t''(x) \left(\frac{d}{dt}y(t) \right)$$

- Compute derivative

$$y'' = \frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2}$$

Substitute the change of variables back into the ODE

$$x^2 \left(\frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2} \right) + 2 \frac{d}{dt}y(t) - 6y(t) = 0$$

- Simplify

$$\frac{d^2}{dt^2}y(t) + \frac{d}{dt}y(t) - 6y(t) = 0$$

- Characteristic polynomial of ODE

$$r^2 + r - 6 = 0$$

- Factor the characteristic polynomial

$$(r + 3)(r - 2) = 0$$

- Roots of the characteristic polynomial

$$r = (-3, 2)$$

- 1st solution of the ODE

$$y_1(t) = e^{-3t}$$

- 2nd solution of the ODE

$$y_2(t) = e^{2t}$$

- General solution of the ODE

$$y(t) = c_1 y_1(t) + c_2 y_2(t)$$

- Substitute in solutions

$$y(t) = c_1 e^{-3t} + c_2 e^{2t}$$

- Change variables back using $t = \ln(x)$

$$y = \frac{c_1}{x^3} + c_2 x^2$$

- Simplify

$$y = \frac{c_1}{x^3} + c_2 x^2$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
<- LODE of Euler type successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
dsolve(x^2*diff(y(x),x$2)+2*x*diff(y(x),x)-6*y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{c_1 x^5 + c_2}{x^3}$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 18

```
DSolve[x^2*y''[x]+2*x*y'[x]-6*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_2 x^5 + c_1}{x^3}$$

16.2 problem 1(b)

16.2.1 Solving as second order euler ode ode	1176
16.2.2 Solving as second order change of variable on x method 2 ode .	1177
16.2.3 Solving as second order change of variable on y method 2 ode .	1180
16.2.4 Solving as second order ode non constant coeff transformation on B ode	1182
16.2.5 Solving using Kovacic algorithm	1185
16.2.6 Maple step by step solution	1190

Internal problem ID [6032]

Internal file name [OUTPUT/5280_Sunday_June_05_2022_03_29_22_PM_91398161/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 149

Problem number: 1(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_euler_ode", "second_order_change_of_variable_on_x_method_2", "second_order_change_of_variable_on_y_method_2", "second_order_ode_non_constant_coeff_transformation_on_B"

Maple gives the following as the ode type

```
[[_Emden, _Fowler]]
```

$$2x^2y'' + xy' - y = 0$$

16.2.1 Solving as second order euler ode ode

This is Euler second order ODE. Let the solution be $y = x^r$, then $y' = rx^{r-1}$ and $y'' = r(r-1)x^{r-2}$. Substituting these back into the given ODE gives

$$2x^2(r(r-1))x^{r-2} + rrx^{r-1} - x^r = 0$$

Simplifying gives

$$2r(r-1)x^r + rx^r - x^r = 0$$

Since $x^r \neq 0$ then dividing throughout by x^r gives

$$2r(r - 1) + r - 1 = 0$$

Or

$$2r^2 - r - 1 = 0 \quad (1)$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$\begin{aligned} r_1 &= 1 \\ r_2 &= -\frac{1}{2} \end{aligned}$$

Since the roots are real and distinct, then the general solution is

$$y = c_1 y_1 + c_2 y_2$$

Where $y_1 = x^{r_1}$ and $y_2 = x^{r_2}$. Hence

$$y = c_1 x + \frac{c_2}{\sqrt{x}}$$

Summary

The solution(s) found are the following

$$y = c_1 x + \frac{c_2}{\sqrt{x}} \quad (1)$$

Verification of solutions

$$y = c_1 x + \frac{c_2}{\sqrt{x}}$$

Verified OK.

16.2.2 Solving as second order change of variable on x method 2 ode

In normal form the ode

$$2x^2 y'' + xy' - y = 0 \quad (1)$$

Becomes

$$y'' + p(x) y' + q(x) y = 0 \quad (2)$$

Where

$$p(x) = \frac{1}{2x}$$
$$q(x) = -\frac{1}{2x^2}$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x)\tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned} \tau &= \int e^{-(\int p(x)dx)} dx \\ &= \int e^{-(\int \frac{1}{2x} dx)} dx \\ &= \int e^{-\frac{\ln(x)}{2}} dx \\ &= \int \frac{1}{\sqrt{x}} dx \\ &= 2\sqrt{x} \end{aligned} \quad (6)$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned} q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\ &= \frac{-\frac{1}{2x^2}}{\frac{1}{x}} \\ &= -\frac{1}{2x} \end{aligned} \quad (7)$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned}\frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) - \frac{y(\tau)}{2x} &= 0\end{aligned}$$

But in terms of τ

$$-\frac{1}{2x} = -\frac{2}{\tau^2}$$

Hence the above ode becomes

$$\frac{d^2}{d\tau^2}y(\tau) - \frac{2y(\tau)}{\tau^2} = 0$$

The above ode is now solved for $y(\tau)$. The ode can be written as

$$\left(\frac{d^2}{d\tau^2}y(\tau)\right)\tau^2 - 2y(\tau) = 0$$

Which shows it is a Euler ODE. This is Euler second order ODE. Let the solution be $y(\tau) = \tau^r$, then $y' = r\tau^{r-1}$ and $y'' = r(r-1)\tau^{r-2}$. Substituting these back into the given ODE gives

$$\tau^2(r(r-1))\tau^{r-2} + 0r\tau^{r-1} - 2\tau^r = 0$$

Simplifying gives

$$r(r-1)\tau^r + 0\tau^r - 2\tau^r = 0$$

Since $\tau^r \neq 0$ then dividing throughout by τ^r gives

$$r(r-1) + 0 - 2 = 0$$

Or

$$r^2 - r - 2 = 0 \tag{1}$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = -1$$

$$r_2 = 2$$

Since the roots are real and distinct, then the general solution is

$$y(\tau) = c_1y_1 + c_2y_2$$

Where $y_1 = \tau^{r_1}$ and $y_2 = \tau^{r_2}$. Hence

$$y(\tau) = \frac{c_1}{\tau} + c_2\tau^2$$

The above solution is now transformed back to y using (6) which results in

$$y = \frac{c_1}{2\sqrt{x}} + 4c_2x$$

Summary

The solution(s) found are the following

$$y = \frac{c_1}{2\sqrt{x}} + 4c_2x \quad (1)$$

Verification of solutions

$$y = \frac{c_1}{2\sqrt{x}} + 4c_2x$$

Verified OK.

16.2.3 Solving as second order change of variable on y method 2 ode

In normal form the ode

$$2x^2y'' + xy' - y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{1}{2x}$$
$$q(x) = -\frac{1}{2x^2}$$

Applying change of variables on the dependent variable $y = v(x)x^n$ to (2) gives the following ode where the dependent variables is $v(x)$ and not y .

$$v''(x) + \left(\frac{2n}{x} + p\right)v'(x) + \left(\frac{n(n-1)}{x^2} + \frac{np}{x} + q\right)v(x) = 0 \quad (3)$$

Let the coefficient of $v(x)$ above be zero. Hence

$$\frac{n(n-1)}{x^2} + \frac{np}{x} + q = 0 \quad (4)$$

Substituting the earlier values found for $p(x)$ and $q(x)$ into (4) gives

$$\frac{n(n-1)}{x^2} + \frac{n}{2x^2} - \frac{1}{2x^2} = 0 \quad (5)$$

Solving (5) for n gives

$$n = 1 \quad (6)$$

Substituting this value in (3) gives

$$\begin{aligned} v''(x) + \frac{5v'(x)}{2x} &= 0 \\ v''(x) + \frac{5v'(x)}{2x} &= 0 \end{aligned} \quad (7)$$

Using the substitution

$$u(x) = v'(x)$$

Then (7) becomes

$$u'(x) + \frac{5u(x)}{2x} = 0 \quad (8)$$

The above is now solved for $u(x)$. In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{5u}{2x} \end{aligned}$$

Where $f(x) = -\frac{5}{2x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned} \frac{1}{u} du &= -\frac{5}{2x} dx \\ \int \frac{1}{u} du &= \int -\frac{5}{2x} dx \\ \ln(u) &= -\frac{5 \ln(x)}{2} + c_1 \\ u &= e^{-\frac{5 \ln(x)}{2} + c_1} \\ &= \frac{c_1}{x^{\frac{5}{2}}} \end{aligned}$$

Now that $u(x)$ is known, then

$$\begin{aligned}v'(x) &= u(x) \\v(x) &= \int u(x) dx + c_2 \\&= -\frac{2c_1}{3x^{\frac{3}{2}}} + c_2\end{aligned}$$

Hence

$$\begin{aligned}y &= v(x) x^n \\&= \left(-\frac{2c_1}{3x^{\frac{3}{2}}} + c_2\right) x \\&= \left(-\frac{2c_1}{3x^{\frac{3}{2}}} + c_2\right) x\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \left(-\frac{2c_1}{3x^{\frac{3}{2}}} + c_2\right) x \quad (1)$$

Verification of solutions

$$y = \left(-\frac{2c_1}{3x^{\frac{3}{2}}} + c_2\right) x$$

Verified OK.

16.2.4 Solving as second order ode non constant coeff transformation on B ode

Given an ode of the form

$$Ay'' + By' + Cy = F(x)$$

This method reduces the order ode the ODE by one by applying the transformation

$$y = Bv$$

This results in

$$\begin{aligned}y' &= B'v + v'B \\y'' &= B''v + B'v' + v''B + v'B' \\&= v''B + 2v' + B' + B''v\end{aligned}$$

And now the original ode becomes

$$\begin{aligned} A(v''B + 2v'B' + B''v) + B(B'v + v'B) + CBv &= 0 \\ ABv'' + (2AB' + B^2)v' + (AB'' + BB' + CB)v &= 0 \end{aligned} \quad (1)$$

If the term $AB'' + BB' + CB$ is zero, then this method works and can be used to solve

$$ABv'' + (2AB' + B^2)v' = 0$$

By Using $u = v'$ which reduces the order of the above ode to one. The new ode is

$$ABu' + (2AB' + B^2)u = 0$$

The above ode is first order ode which is solved for u . Now a new ode $v' = u$ is solved for v as first order ode. Then the final solution is obtain from $y = Bv$.

This method works only if the term $AB'' + BB' + CB$ is zero. The given ODE shows that

$$A = 2x^2$$

$$B = x$$

$$C = -1$$

$$F = 0$$

The above shows that for this ode

$$\begin{aligned} AB'' + BB' + CB &= (2x^2)(0) + (x)(1) + (-1)(x) \\ &= 0 \end{aligned}$$

Hence the ode in v given in (1) now simplifies to

$$2x^3v'' + (5x^2)v' = 0$$

Now by applying $v' = u$ the above becomes

$$2x^3u'(x) + 5x^2u(x) = 0$$

Which is now solved for u . In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{5u}{2x} \end{aligned}$$

Where $f(x) = -\frac{5}{2x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u} du &= -\frac{5}{2x} dx \\ \int \frac{1}{u} du &= \int -\frac{5}{2x} dx \\ \ln(u) &= -\frac{5 \ln(x)}{2} + c_1 \\ u &= e^{-\frac{5 \ln(x)}{2} + c_1} \\ &= \frac{c_1}{x^{\frac{5}{2}}}\end{aligned}$$

The ode for v now becomes

$$\begin{aligned}v' &= u \\ &= \frac{c_1}{x^{\frac{5}{2}}}\end{aligned}$$

Which is now solved for v . Integrating both sides gives

$$\begin{aligned}v(x) &= \int \frac{c_1}{x^{\frac{5}{2}}} dx \\ &= -\frac{2c_1}{3x^{\frac{3}{2}}} + c_2\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y(x) &= Bv \\ &= (x) \left(-\frac{2c_1}{3x^{\frac{3}{2}}} + c_2 \right) \\ &= \left(-\frac{2c_1}{3x^{\frac{3}{2}}} + c_2 \right) x\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \left(-\frac{2c_1}{3x^{\frac{3}{2}}} + c_2 \right) x \tag{1}$$

Verification of solutions

$$y = \left(-\frac{2c_1}{3x^{\frac{3}{2}}} + c_2 \right) x$$

Verified OK.

16.2.5 Solving using Kovacic algorithm

Writing the ode as

$$2x^2y'' + xy' - y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 2x^2 \\ B &= x \\ C &= -1 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{5}{16x^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 5 \\ t &= 16x^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{5}{16x^2} \right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 195: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 16x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = \frac{5}{16x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = \frac{5}{16}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{5}{4} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = -\frac{1}{4} \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ , which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = \frac{5}{16x^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = \frac{5}{16}$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{5}{4} \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = -\frac{1}{4} \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{5}{16x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	$\frac{5}{4}$	$-\frac{1}{4}$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	$\frac{5}{4}$	$-\frac{1}{4}$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω .

Trying $\alpha_{\infty}^{-} = -\frac{1}{4}$ then

$$\begin{aligned} d &= \alpha_{\infty}^{-} - (\alpha_{c_1}^{-}) \\ &= -\frac{1}{4} - \left(-\frac{1}{4}\right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_{\infty}$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^{-}}{x - c_1} \right) + (-)[\sqrt{r}]_{\infty} \\ &= -\frac{1}{4x} + (-)(0) \\ &= -\frac{1}{4x} \\ &= -\frac{1}{4x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2\left(-\frac{1}{4x}\right)(0) + \left(\left(\frac{1}{4x^2}\right) + \left(-\frac{1}{4x}\right)^2 - \left(\frac{5}{16x^2}\right)\right) &= 0 \\ 0 &= 0 \end{aligned}$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= pe^{\int \omega dx} \\ &= e^{\int -\frac{1}{4x} dx} \\ &= \frac{1}{x^{\frac{1}{4}}} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\&= z_1 e^{-\int \frac{1}{2} \frac{x}{2x^2} dx} \\&= z_1 e^{-\frac{\ln(x)}{4}} \\&= z_1 \left(\frac{1}{x^{\frac{1}{4}}} \right)\end{aligned}$$

Which simplifies to

$$y_1 = \frac{1}{\sqrt{x}}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{x}{2x^2} dx}}{(y_1)^2} dx \\&= y_1 \int \frac{e^{-\frac{\ln(x)}{2}}}{(y_1)^2} dx \\&= y_1 \left(\frac{2x^{\frac{3}{2}}}{3} \right)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\&= c_1 \left(\frac{1}{\sqrt{x}} \right) + c_2 \left(\frac{1}{\sqrt{x}} \left(\frac{2x^{\frac{3}{2}}}{3} \right) \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1}{\sqrt{x}} + \frac{2c_2 x}{3} \tag{1}$$

Verification of solutions

$$y = \frac{c_1}{\sqrt{x}} + \frac{2c_2x}{3}$$

Verified OK.

16.2.6 Maple step by step solution

Let's solve

$$2x^2y'' + xy' - y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{y'}{2x} + \frac{y}{2x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{y'}{2x} - \frac{y}{2x^2} = 0$$

- Multiply by denominators of the ODE

$$2x^2y'' + xy' - y = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x , using the chain rule

$$y' = \left(\frac{d}{dt}y(t)\right) t'(x)$$

- Compute derivative

$$y' = \frac{\frac{d}{dt}y(t)}{x}$$

- Calculate the 2nd derivative of y with respect to x , using the chain rule

$$y'' = \left(\frac{d^2}{dt^2}y(t)\right) t'(x)^2 + t''(x) \left(\frac{d}{dt}y(t)\right)$$

- Compute derivative

$$y'' = \frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2}$$

Substitute the change of variables back into the ODE

$$2x^2 \left(\frac{\frac{d^2}{dt^2} y(t)}{x^2} - \frac{\frac{d}{dt} y(t)}{x^2} \right) + \frac{d}{dt} y(t) - y(t) = 0$$

- Simplify

$$2 \frac{d^2}{dt^2} y(t) - \frac{d}{dt} y(t) - y(t) = 0$$

- Isolate 2nd derivative

$$\frac{d^2}{dt^2} y(t) = \frac{\frac{d}{dt} y(t)}{2} + \frac{y(t)}{2}$$

- Group terms with $y(t)$ on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$\frac{d^2}{dt^2} y(t) - \frac{\frac{d}{dt} y(t)}{2} - \frac{y(t)}{2} = 0$$

- Characteristic polynomial of ODE

$$r^2 - \frac{1}{2}r - \frac{1}{2} = 0$$

- Factor the characteristic polynomial

$$\frac{(2r+1)(r-1)}{2} = 0$$

- Roots of the characteristic polynomial

$$r = \left(1, -\frac{1}{2} \right)$$

- 1st solution of the ODE

$$y_1(t) = e^t$$

- 2nd solution of the ODE

$$y_2(t) = e^{-\frac{t}{2}}$$

- General solution of the ODE

$$y(t) = c_1 y_1(t) + c_2 y_2(t)$$

- Substitute in solutions

$$y(t) = c_1 e^t + c_2 e^{-\frac{t}{2}}$$

- Change variables back using $t = \ln(x)$

$$y = c_1 x + \frac{c_2}{\sqrt{x}}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
<- LODE of Euler type successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(2*x^2*diff(y(x),x$2)+x*diff(y(x),x)-y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1x + \frac{c_2}{\sqrt{x}}$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 18

```
DSolve[2*x^2*y''[x]+x*y'[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_1}{\sqrt{x}} + c_2x$$

16.3 problem 1(c)

16.3.1 Solving as second order euler ode ode	1193
16.3.2 Solving as second order change of variable on x method 2 ode .	1194
16.3.3 Solving as second order change of variable on x method 1 ode .	1197
16.3.4 Solving as second order change of variable on y method 2 ode .	1199
16.3.5 Solving using Kovacic algorithm	1201
16.3.6 Maple step by step solution	1206

Internal problem ID [6033]

Internal file name [OUTPUT/5281_Sunday_June_05_2022_03_29_23_PM_90527720/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 149

Problem number: 1(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_euler_ode", "second_order_change_of_variable_on_x_method_1", "second_order_change_of_variable_on_x_method_2", "second_order_change_of_variable_on_y_method_2"

Maple gives the following as the ode type

```
[[_Emden, _Fowler], [_2nd_order, _linear, `_with_symmetry_[0,F(x)]`]]
```

$$x^2y'' + xy' - 4y = 0$$

16.3.1 Solving as second order euler ode ode

This is Euler second order ODE. Let the solution be $y = x^r$, then $y' = rx^{r-1}$ and $y'' = r(r-1)x^{r-2}$. Substituting these back into the given ODE gives

$$x^2(r(r-1))x^{r-2} + xrx^{r-1} - 4x^r = 0$$

Simplifying gives

$$r(r-1)x^r + rx^r - 4x^r = 0$$

Since $x^r \neq 0$ then dividing throughout by x^r gives

$$r(r - 1) + r - 4 = 0$$

Or

$$r^2 - 4 = 0 \tag{1}$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = -2$$

$$r_2 = 2$$

Since the roots are real and distinct, then the general solution is

$$y = c_1y_1 + c_2y_2$$

Where $y_1 = x^{r_1}$ and $y_2 = x^{r_2}$. Hence

$$y = \frac{c_1}{x^2} + c_2x^2$$

Summary

The solution(s) found are the following

$$y = \frac{c_1}{x^2} + c_2x^2 \tag{1}$$

Verification of solutions

$$y = \frac{c_1}{x^2} + c_2x^2$$

Verified OK.

16.3.2 Solving as second order change of variable on x method 2 ode

In normal form the ode

$$x^2y'' + xy' - 4y = 0 \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \tag{2}$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = -\frac{4}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x)\tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned} \tau &= \int e^{-(\int p(x)dx)} dx \\ &= \int e^{-(\int \frac{1}{x}dx)} dx \\ &= \int e^{-\ln(x)} dx \\ &= \int \frac{1}{x} dx \\ &= \ln(x) \end{aligned} \quad (6)$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned} q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\ &= \frac{-\frac{4}{x^2}}{\frac{1}{x^2}} \\ &= -4 \end{aligned} \quad (7)$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned} \frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) - 4y(\tau) &= 0 \end{aligned}$$

The above ode is now solved for $y(\tau)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(\tau) + By'(\tau) + Cy(\tau) = 0$$

Where in the above $A = 1, B = 0, C = -4$. Let the solution be $y(\tau) = e^{\lambda\tau}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda\tau} - 4 e^{\lambda\tau} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda\tau}$ gives

$$\lambda^2 - 4 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = -4$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(-4)} \\ &= \pm 2 \end{aligned}$$

Hence

$$\lambda_1 = +2$$

$$\lambda_2 = -2$$

Which simplifies to

$$\lambda_1 = 2$$

$$\lambda_2 = -2$$

Since roots are real and distinct, then the solution is

$$y(\tau) = c_1 e^{\lambda_1 \tau} + c_2 e^{\lambda_2 \tau}$$

$$y(\tau) = c_1 e^{(2)\tau} + c_2 e^{(-2)\tau}$$

Or

$$y(\tau) = c_1 e^{2\tau} + c_2 e^{-2\tau}$$

The above solution is now transformed back to y using (6) which results in

$$y = \frac{c_1 x^4 + c_2}{x^2}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1 x^4 + c_2}{x^2} \quad (1)$$

Verification of solutions

$$y = \frac{c_1 x^4 + c_2}{x^2}$$

Verified OK.

16.3.3 Solving as second order change of variable on x method 1 ode

In normal form the ode

$$x^2 y'' + xy' - 4y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = -\frac{4}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) results

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $q_1 = c^2$ where c is some constant. Therefore from (5)

$$\begin{aligned}\tau' &= \frac{1}{c}\sqrt{q} \\ &= \frac{2\sqrt{-\frac{1}{x^2}}}{c} \\ \tau'' &= \frac{2}{c\sqrt{-\frac{1}{x^2}}x^3}\end{aligned}\tag{6}$$

Substituting the above into (4) results in

$$\begin{aligned}p_1(\tau) &= \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \\ &= \frac{\frac{2}{c\sqrt{-\frac{1}{x^2}}x^3} + \frac{1}{x}\frac{2\sqrt{-\frac{1}{x^2}}}{c}}{\left(\frac{2\sqrt{-\frac{1}{x^2}}}{c}\right)^2} \\ &= 0\end{aligned}$$

Therefore ode (3) now becomes

$$\begin{aligned}y(\tau)'' + p_1y(\tau)' + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) + c^2y(\tau) &= 0\end{aligned}\tag{7}$$

The above ode is now solved for $y(\tau)$. Since the ode is now constant coefficients, it can be easily solved to give

$$y(\tau) = c_1 \cos(c\tau) + c_2 \sin(c\tau)$$

Now from (6)

$$\begin{aligned}\tau &= \int \frac{1}{c}\sqrt{q} dx \\ &= \frac{\int 2\sqrt{-\frac{1}{x^2}} dx}{c} \\ &= \frac{2\sqrt{-\frac{1}{x^2}}x \ln(x)}{c}\end{aligned}$$

Substituting the above into the solution obtained gives

$$y = c_1 \cosh(2 \ln(x)) + ic_2 \sinh(2 \ln(x))$$

Summary

The solution(s) found are the following

$$y = c_1 \cosh(2 \ln(x)) + ic_2 \sinh(2 \ln(x)) \quad (1)$$

Verification of solutions

$$y = c_1 \cosh(2 \ln(x)) + ic_2 \sinh(2 \ln(x))$$

Verified OK.

16.3.4 Solving as second order change of variable on y method 2 ode

In normal form the ode

$$x^2 y'' + xy' - 4y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = -\frac{4}{x^2}$$

Applying change of variables on the dependent variable $y = v(x)x^n$ to (2) gives the following ode where the dependent variables is $v(x)$ and not y .

$$v''(x) + \left(\frac{2n}{x} + p\right)v'(x) + \left(\frac{n(n-1)}{x^2} + \frac{np}{x} + q\right)v(x) = 0 \quad (3)$$

Let the coefficient of $v(x)$ above be zero. Hence

$$\frac{n(n-1)}{x^2} + \frac{np}{x} + q = 0 \quad (4)$$

Substituting the earlier values found for $p(x)$ and $q(x)$ into (4) gives

$$\frac{n(n-1)}{x^2} + \frac{n}{x^2} - \frac{4}{x^2} = 0 \quad (5)$$

Solving (5) for n gives

$$n = 2 \quad (6)$$

Substituting this value in (3) gives

$$\begin{aligned}v''(x) + \frac{5v'(x)}{x} &= 0 \\v''(x) + \frac{5v'(x)}{x} &= 0\end{aligned}\tag{7}$$

Using the substitution

$$u(x) = v'(x)$$

Then (7) becomes

$$u'(x) + \frac{5u(x)}{x} = 0\tag{8}$$

The above is now solved for $u(x)$. In canonical form the ODE is

$$\begin{aligned}u' &= F(x, u) \\&= f(x)g(u) \\&= -\frac{5u}{x}\end{aligned}$$

Where $f(x) = -\frac{5}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u} du &= -\frac{5}{x} dx \\ \int \frac{1}{u} du &= \int -\frac{5}{x} dx \\ \ln(u) &= -5 \ln(x) + c_1 \\ u &= e^{-5 \ln(x) + c_1} \\ &= \frac{c_1}{x^5}\end{aligned}$$

Now that $u(x)$ is known, then

$$\begin{aligned}v'(x) &= u(x) \\ v(x) &= \int u(x) dx + c_2 \\ &= -\frac{c_1}{4x^4} + c_2\end{aligned}$$

Hence

$$\begin{aligned}y &= v(x) x^n \\ &= \left(-\frac{c_1}{4x^4} + c_2\right) x^2 \\ &= \frac{4c_2x^4 - c_1}{4x^2}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \left(-\frac{c_1}{4x^4} + c_2\right) x^2 \quad (1)$$

Verification of solutions

$$y = \left(-\frac{c_1}{4x^4} + c_2\right) x^2$$

Verified OK.

16.3.5 Solving using Kovacic algorithm

Writing the ode as

$$x^2y'' + xy' - 4y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= x^2 \\ B &= x \\ C &= -4\end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{15}{4x^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 15 \\ t &= 4x^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{15}{4x^2} \right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 197: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = \frac{15}{4x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = \frac{15}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{5}{2} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = -\frac{3}{2} \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = \frac{15}{4x^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = \frac{15}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{5}{2} \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = -\frac{3}{2} \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{15}{4x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	$\frac{5}{2}$	$-\frac{3}{2}$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	$\frac{5}{2}$	$-\frac{3}{2}$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = -\frac{3}{2}$ then

$$\begin{aligned} d &= \alpha_\infty^- - (\alpha_{c_1}^-) \\ &= -\frac{3}{2} - \left(-\frac{3}{2}\right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c) [\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty) [\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= \left((-) [\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{x - c_1} \right) + (-) [\sqrt{r}]_\infty \\ &= -\frac{3}{2x} + (-) (0) \\ &= -\frac{3}{2x} \\ &= -\frac{3}{2x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$(0) + 2\left(-\frac{3}{2x}\right)(0) + \left(\left(\frac{3}{2x^2}\right) + \left(-\frac{3}{2x}\right)^2 - \left(\frac{15}{4x^2}\right)\right) = 0$$
$$0 = 0$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$z_1(x) = pe^{\int \omega dx}$$
$$= e^{\int -\frac{3}{2x} dx}$$
$$= \frac{1}{x^{\frac{3}{2}}}$$

The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$
$$= z_1 e^{-\int \frac{1}{2} \frac{x}{x^2} dx}$$
$$= z_1 e^{-\frac{\ln(x)}{2}}$$
$$= z_1 \left(\frac{1}{\sqrt{x}}\right)$$

Which simplifies to

$$y_1 = \frac{1}{x^2}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$y_2 = y_1 \int \frac{e^{\int -\frac{x}{x^2} dx}}{(y_1)^2} dx$$
$$= y_1 \int \frac{e^{-\ln(x)}}{(y_1)^2} dx$$
$$= y_1 \left(\frac{x^4}{4}\right)$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(\frac{1}{x^2} \right) + c_2 \left(\frac{1}{x^2} \left(\frac{x^4}{4} \right) \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1}{x^2} + \frac{c_2 x^2}{4} \quad (1)$$

Verification of solutions

$$y = \frac{c_1}{x^2} + \frac{c_2 x^2}{4}$$

Verified OK.

16.3.6 Maple step by step solution

Let's solve

$$x^2 y'' + xy' - 4y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{y'}{x} + \frac{4y}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{y'}{x} - \frac{4y}{x^2} = 0$$

- Multiply by denominators of the ODE

$$x^2 y'' + xy' - 4y = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x , using the chain rule

$$y' = \left(\frac{d}{dt} y(t) \right) t'(x)$$

- Compute derivative

$$y' = \frac{\frac{d}{dt}y(t)}{x}$$

- Calculate the 2nd derivative of y with respect to x , using the chain rule

$$y'' = \left(\frac{d^2}{dt^2}y(t) \right) t'(x)^2 + t''(x) \left(\frac{d}{dt}y(t) \right)$$

- Compute derivative

$$y'' = \frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2}$$

Substitute the change of variables back into the ODE

$$x^2 \left(\frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2} \right) + \frac{d}{dt}y(t) - 4y(t) = 0$$

- Simplify

$$\frac{d^2}{dt^2}y(t) - 4y(t) = 0$$

- Characteristic polynomial of ODE

$$r^2 - 4 = 0$$

- Factor the characteristic polynomial

$$(r - 2)(r + 2) = 0$$

- Roots of the characteristic polynomial

$$r = (-2, 2)$$

- 1st solution of the ODE

$$y_1(t) = e^{-2t}$$

- 2nd solution of the ODE

$$y_2(t) = e^{2t}$$

- General solution of the ODE

$$y(t) = c_1 y_1(t) + c_2 y_2(t)$$

- Substitute in solutions

$$y(t) = c_1 e^{-2t} + c_2 e^{2t}$$

- Change variables back using $t = \ln(x)$

$$y = \frac{c_1}{x^2} + c_2 x^2$$

- Simplify

$$y = \frac{c_1}{x^2} + c_2 x^2$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
<- LODE of Euler type successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 15

```
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)-4*y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{c_1 x^4 + c_2}{x^2}$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 18

```
DSolve[x^2*y''[x]+x*y'[x]-4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_2 x^4 + c_1}{x^2}$$

16.4 problem 1(d)

- 16.4.1 Solving as second order euler ode 1209
- 16.4.2 Solving as second order change of variable on x method 2 ode . 1213
- 16.4.3 Solving as second order change of variable on x method 1 ode . 1218
- 16.4.4 Solving as second order change of variable on y method 2 ode . 1223
- 16.4.5 Solving using Kovacic algorithm 1228

Internal problem ID [6034]

Internal file name [OUTPUT/5282_Sunday_June_05_2022_03_29_24_PM_28936060/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 149

Problem number: 1(d).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_euler_ode", "second_order_change_of_variable_on_x_method_1", "second_order_change_of_variable_on_x_method_2", "second_order_change_of_variable_on_y_method_2"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$x^2y'' - 5xy' + 9y = x^2$$

16.4.1 Solving as second order euler ode ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = x^2, B = -5x, C = 9, f(x) = x^2$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$x^2y'' - 5xy' + 9y = 0$$

This is Euler second order ODE. Let the solution be $y = x^r$, then $y' = rx^{r-1}$ and $y'' = r(r-1)x^{r-2}$. Substituting these back into the given ODE gives

$$x^2(r(r-1))x^{r-2} - 5rxr^{r-1} + 9x^r = 0$$

Simplifying gives

$$r(r-1)x^r - 5rx^r + 9x^r = 0$$

Since $x^r \neq 0$ then dividing throughout by x^r gives

$$r(r-1) - 5r + 9 = 0$$

Or

$$r^2 - 6r + 9 = 0 \tag{1}$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = 3$$

$$r_2 = 3$$

Since the roots are equal, then the general solution is

$$y = c_1y_1 + c_2y_2$$

Where $y_1 = x^r$ and $y_2 = x^r \ln(x)$. Hence

$$y = c_1x^3 + c_2x^3 \ln(x)$$

Next, we find the particular solution to the ODE

$$x^2y'' - 5xy' + 9y = x^2$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1y_1 + u_2y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = x^3$$

$$y_2 = x^3 \ln(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} x^3 & x^3 \ln(x) \\ \frac{d}{dx}(x^3) & \frac{d}{dx}(x^3 \ln(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} x^3 & x^3 \ln(x) \\ 3x^2 & 3 \ln(x) x^2 + x^2 \end{vmatrix}$$

Therefore

$$W = (x^3) (3 \ln(x) x^2 + x^2) - (x^3 \ln(x)) (3x^2)$$

Which simplifies to

$$W = x^5$$

Which simplifies to

$$W = x^5$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^5 \ln(x)}{x^7} dx$$

Which simplifies to

$$u_1 = - \int \frac{\ln(x)}{x^2} dx$$

Hence

$$u_1 = \frac{\ln(x)}{x} + \frac{1}{x}$$

And Eq. (3) becomes

$$u_2 = \int \frac{x^5}{x^7} dx$$

Which simplifies to

$$u_2 = \int \frac{1}{x^2} dx$$

Hence

$$u_2 = -\frac{1}{x}$$

Which simplifies to

$$u_1 = \frac{1 + \ln(x)}{x}$$

$$u_2 = -\frac{1}{x}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = (1 + \ln(x)) x^2 - \ln(x) x^2$$

Which simplifies to

$$y_p(x) = x^2$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= x^2 + c_1 x^3 + c_2 x^3 \ln(x) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = x^2 + c_1 x^3 + c_2 x^3 \ln(x) \tag{1}$$

Verification of solutions

$$y = x^2 + c_1 x^3 + c_2 x^3 \ln(x)$$

Verified OK.

16.4.2 Solving as second order change of variable on x method 2 ode

This is second order non-homogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$x^2y'' - 5xy' + 9y = 0$$

In normal form the ode

$$x^2y'' - 5xy' + 9y = 0 \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \tag{2}$$

Where

$$p(x) = -\frac{5}{x}$$
$$q(x) = \frac{9}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \tag{3}$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \tag{4}$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \tag{5}$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x)\tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned}
 \tau &= \int e^{-(\int p(x)dx)} dx \\
 &= \int e^{-(\int -\frac{5}{x} dx)} dx \\
 &= \int e^{5\ln(x)} dx \\
 &= \int x^5 dx \\
 &= \frac{x^6}{6}
 \end{aligned} \tag{6}$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned}
 q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\
 &= \frac{\frac{9}{x^2}}{x^{10}} \\
 &= \frac{9}{x^{12}}
 \end{aligned} \tag{7}$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned}
 \frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) &= 0 \\
 \frac{d^2}{d\tau^2}y(\tau) + \frac{9y(\tau)}{x^{12}} &= 0
 \end{aligned}$$

But in terms of τ

$$\frac{9}{x^{12}} = \frac{1}{4\tau^2}$$

Hence the above ode becomes

$$\frac{d^2}{d\tau^2}y(\tau) + \frac{y(\tau)}{4\tau^2} = 0$$

The above ode is now solved for $y(\tau)$. The ode can be written as

$$4 \left(\frac{d^2}{d\tau^2}y(\tau) \right) \tau^2 + y(\tau) = 0$$

Which shows it is a Euler ODE. This is Euler second order ODE. Let the solution be $y(\tau) = \tau^r$, then $y' = r\tau^{r-1}$ and $y'' = r(r-1)\tau^{r-2}$. Substituting these back into the given ODE gives

$$4\tau^2(r(r-1))\tau^{r-2} + 0r\tau^{r-1} + \tau^r = 0$$

Simplifying gives

$$4r(r - 1) \tau^r + 0 \tau^r + \tau^r = 0$$

Since $\tau^r \neq 0$ then dividing throughout by τ^r gives

$$4r(r - 1) + 0 + 1 = 0$$

Or

$$4r^2 - 4r + 1 = 0 \tag{1}$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = \frac{1}{2}$$
$$r_2 = \frac{1}{2}$$

Since the roots are equal, then the general solution is

$$y(\tau) = c_1 y_1 + c_2 y_2$$

Where $y_1 = \tau^r$ and $y_2 = \tau^r \ln(\tau)$. Hence

$$y(\tau) = c_1 \sqrt{\tau} + c_2 \sqrt{\tau} \ln(\tau)$$

The above solution is now transformed back to y using (6) which results in

$$y = \frac{\sqrt{6} \sqrt{x^6} (c_1 + c_2 \ln(x^6) - c_2 \ln(3) - c_2 \ln(2))}{6}$$

Therefore the homogeneous solution y_h is

$$y_h = \frac{\sqrt{6} \sqrt{x^6} (c_1 + c_2 \ln(x^6) - c_2 \ln(3) - c_2 \ln(2))}{6}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \sqrt{x^6}$$

$$y_2 = \frac{\sqrt{6} \sqrt{x^6} \ln(x^6)}{6} - \frac{\sqrt{6} \sqrt{x^6} \ln(3)}{6} - \frac{\sqrt{6} \sqrt{x^6} \ln(2)}{6}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \sqrt{x^6} & \frac{\sqrt{6} \sqrt{x^6} \ln(x^6)}{6} - \frac{\sqrt{6} \sqrt{x^6} \ln(3)}{6} - \frac{\sqrt{6} \sqrt{x^6} \ln(2)}{6} \\ \frac{d}{dx}(\sqrt{x^6}) & \frac{d}{dx} \left(\frac{\sqrt{6} \sqrt{x^6} \ln(x^6)}{6} - \frac{\sqrt{6} \sqrt{x^6} \ln(3)}{6} - \frac{\sqrt{6} \sqrt{x^6} \ln(2)}{6} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \sqrt{x^6} & \frac{\sqrt{6} \sqrt{x^6} \ln(x^6)}{6} - \frac{\sqrt{6} \sqrt{x^6} \ln(3)}{6} - \frac{\sqrt{6} \sqrt{x^6} \ln(2)}{6} \\ \frac{3x^5}{\sqrt{x^6}} & \frac{\sqrt{6} \ln(x^6)x^5}{2\sqrt{x^6}} + \frac{\sqrt{6} \sqrt{x^6}}{x} - \frac{\sqrt{6} \ln(3)x^5}{2\sqrt{x^6}} - \frac{\sqrt{6} \ln(2)x^5}{2\sqrt{x^6}} \end{vmatrix}$$

Therefore

$$W = \left(\sqrt{x^6} \right) \left(\frac{\sqrt{6} \ln(x^6) x^5}{2\sqrt{x^6}} + \frac{\sqrt{6} \sqrt{x^6}}{x} - \frac{\sqrt{6} \ln(3) x^5}{2\sqrt{x^6}} - \frac{\sqrt{6} \ln(2) x^5}{2\sqrt{x^6}} \right) - \left(\frac{\sqrt{6} \sqrt{x^6} \ln(x^6)}{6} - \frac{\sqrt{6} \sqrt{x^6} \ln(3)}{6} - \frac{\sqrt{6} \sqrt{x^6} \ln(2)}{6} \right) \left(\frac{3x^5}{\sqrt{x^6}} \right)$$

Which simplifies to

$$W = \sqrt{6} x^5$$

Which simplifies to

$$W = \sqrt{6} x^5$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\left(\frac{\sqrt{6} \sqrt{x^6} \ln(x^6)}{6} - \frac{\sqrt{6} \sqrt{x^6} \ln(3)}{6} - \frac{\sqrt{6} \sqrt{x^6} \ln(2)}{6} \right) x^2}{x^7 \sqrt{6}} dx$$

Which simplifies for $0 < x$ to

$$u_1 = - \int \frac{6 \ln(x) - \ln(3) - \ln(2)}{6x^2} dx$$

Hence

$$u_1 = - \frac{-6 \ln(x) - 6 + \ln(3) + \ln(2)}{6x}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\sqrt{x^6} x^2}{x^7 \sqrt{6}} dx$$

Which simplifies for $0 < x$ to

$$u_2 = \int \frac{\sqrt{6}}{6x^2} dx$$

Hence

$$u_2 = - \frac{\sqrt{6}}{6x}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = - \frac{(-6 \ln(x) - 6 + \ln(3) + \ln(2)) \sqrt{x^6}}{6x} - \frac{\sqrt{6} \left(\frac{\sqrt{6} \sqrt{x^6} \ln(x^6)}{6} - \frac{\sqrt{6} \sqrt{x^6} \ln(3)}{6} - \frac{\sqrt{6} \sqrt{x^6} \ln(2)}{6} \right)}{6x}$$

Which simplifies to

$$y_p(x) = x^2$$

Therefore the general solution is

$$y = y_h + y_p \\ = \left(\frac{\sqrt{6} \sqrt{x^6} (c_1 + c_2 \ln(x^6)) - c_2 \ln(3) - c_2 \ln(2)}{6} \right) + (x^2)$$

Summary

The solution(s) found are the following

$$y = \frac{\sqrt{6} \sqrt{x^6} (c_1 + c_2 \ln(x^6)) - c_2 \ln(3) - c_2 \ln(2)}{6} + x^2 \quad (1)$$

Verification of solutions

$$y = \frac{\sqrt{6} \sqrt{x^6} (c_1 + c_2 \ln(x^6)) - c_2 \ln(3) - c_2 \ln(2)}{6} + x^2$$

Verified OK. $\{0 < x\}$

16.4.3 Solving as second order change of variable on x method 1 ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = x^2, B = -5x, C = 9, f(x) = x^2$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$x^2 y'' - 5xy' + 9y = 0$$

In normal form the ode

$$x^2 y'' - 5xy' + 9y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = -\frac{5}{x} \\ q(x) = \frac{9}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) results

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $q_1 = c^2$ where c is some constant. Therefore from (5)

$$\begin{aligned} \tau' &= \frac{1}{c}\sqrt{q} \\ &= \frac{3\sqrt{\frac{1}{x^2}}}{c} \\ \tau'' &= -\frac{3}{c\sqrt{\frac{1}{x^2}}x^3} \end{aligned} \quad (6)$$

Substituting the above into (4) results in

$$\begin{aligned} p_1(\tau) &= \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \\ &= \frac{-\frac{3}{c\sqrt{\frac{1}{x^2}}x^3} - \frac{5}{x}\frac{3\sqrt{\frac{1}{x^2}}}{c}}{\left(\frac{3\sqrt{\frac{1}{x^2}}}{c}\right)^2} \\ &= -2c \end{aligned}$$

Therefore ode (3) now becomes

$$\begin{aligned} y(\tau)'' + p_1y(\tau)' + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) - 2c\left(\frac{d}{d\tau}y(\tau)\right) + c^2y(\tau) &= 0 \end{aligned} \quad (7)$$

The above ode is now solved for $y(\tau)$. Since the ode is now constant coefficients, it can be easily solved to give

$$y(\tau) = e^{c\tau}c_1$$

Now from (6)

$$\begin{aligned}\tau &= \int \frac{1}{c} \sqrt{q} dx \\ &= \frac{\int 3\sqrt{\frac{1}{x^2}} dx}{c} \\ &= \frac{3\sqrt{\frac{1}{x^2}} x \ln(x)}{c}\end{aligned}$$

Substituting the above into the solution obtained gives

$$y = c_1 x^3$$

Now the particular solution to this ODE is found

$$x^2 y'' - 5xy' + 9y = x^2$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned}y_1 &= \sqrt{x^6} \\ y_2 &= \frac{\sqrt{6} \sqrt{x^6} \ln(x^6)}{6} - \frac{\sqrt{6} \sqrt{x^6} \ln(3)}{6} - \frac{\sqrt{6} \sqrt{x^6} \ln(2)}{6}\end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \sqrt{x^6} & \frac{\sqrt{6}\sqrt{x^6}\ln(x^6)}{6} - \frac{\sqrt{6}\sqrt{x^6}\ln(3)}{6} - \frac{\sqrt{6}\sqrt{x^6}\ln(2)}{6} \\ \frac{d}{dx}(\sqrt{x^6}) & \frac{d}{dx}\left(\frac{\sqrt{6}\sqrt{x^6}\ln(x^6)}{6} - \frac{\sqrt{6}\sqrt{x^6}\ln(3)}{6} - \frac{\sqrt{6}\sqrt{x^6}\ln(2)}{6}\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \sqrt{x^6} & \frac{\sqrt{6}\sqrt{x^6}\ln(x^6)}{6} - \frac{\sqrt{6}\sqrt{x^6}\ln(3)}{6} - \frac{\sqrt{6}\sqrt{x^6}\ln(2)}{6} \\ \frac{3x^5}{\sqrt{x^6}} & \frac{\sqrt{6}\ln(x^6)x^5}{2\sqrt{x^6}} + \frac{\sqrt{6}\sqrt{x^6}}{x} - \frac{\sqrt{6}\ln(3)x^5}{2\sqrt{x^6}} - \frac{\sqrt{6}\ln(2)x^5}{2\sqrt{x^6}} \end{vmatrix}$$

Therefore

$$W = (\sqrt{x^6}) \left(\frac{\sqrt{6}\ln(x^6)x^5}{2\sqrt{x^6}} + \frac{\sqrt{6}\sqrt{x^6}}{x} - \frac{\sqrt{6}\ln(3)x^5}{2\sqrt{x^6}} - \frac{\sqrt{6}\ln(2)x^5}{2\sqrt{x^6}} \right) - \left(\frac{\sqrt{6}\sqrt{x^6}\ln(x^6)}{6} - \frac{\sqrt{6}\sqrt{x^6}\ln(3)}{6} - \frac{\sqrt{6}\sqrt{x^6}\ln(2)}{6} \right) \left(\frac{3x^5}{\sqrt{x^6}} \right)$$

Which simplifies to

$$W = \sqrt{6}x^5$$

Which simplifies to

$$W = \sqrt{6}x^5$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\left(\frac{\sqrt{6}\sqrt{x^6}\ln(x^6)}{6} - \frac{\sqrt{6}\sqrt{x^6}\ln(3)}{6} - \frac{\sqrt{6}\sqrt{x^6}\ln(2)}{6} \right) x^2}{x^7\sqrt{6}} dx$$

Which simplifies for $0 < x$ to

$$u_1 = - \int \frac{6\ln(x) - \ln(3) - \ln(2)}{6x^2} dx$$

Hence

$$u_1 = -\frac{-6 \ln(x) - 6 + \ln(3) + \ln(2)}{6x}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\sqrt{x^6} x^2}{x^7 \sqrt{6}} dx$$

Which simplifies for $0 < x$ to

$$u_2 = \int \frac{\sqrt{6}}{6x^2} dx$$

Hence

$$u_2 = -\frac{\sqrt{6}}{6x}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{(-6 \ln(x) - 6 + \ln(3) + \ln(2)) \sqrt{x^6}}{6x} - \frac{\sqrt{6} \left(\frac{\sqrt{6} \sqrt{x^6} \ln(x^6)}{6} - \frac{\sqrt{6} \sqrt{x^6} \ln(3)}{6} - \frac{\sqrt{6} \sqrt{x^6} \ln(2)}{6} \right)}{6x}$$

Which simplifies to

$$y_p(x) = x^2$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 x^3) + (x^2) \\ &= c_1 x^3 + x^2 \end{aligned}$$

Which simplifies to

$$y = c_1 x^3 + x^2$$

Summary

The solution(s) found are the following

$$y = c_1 x^3 + x^2 \tag{1}$$

Verification of solutions

$$y = c_1x^3 + x^2$$

Verified OK. $\{0 < x\}$

16.4.4 Solving as second order change of variable on y method 2 ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = x^2, B = -5x, C = 9, f(x) = x^2$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$x^2y'' - 5xy' + 9y = 0$$

In normal form the ode

$$x^2y'' - 5xy' + 9y = 0 \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \tag{2}$$

Where

$$p(x) = -\frac{5}{x}$$
$$q(x) = \frac{9}{x^2}$$

Applying change of variables on the dependent variable $y = v(x)x^n$ to (2) gives the following ode where the dependent variables is $v(x)$ and not y .

$$v''(x) + \left(\frac{2n}{x} + p\right)v'(x) + \left(\frac{n(n-1)}{x^2} + \frac{np}{x} + q\right)v(x) = 0 \tag{3}$$

Let the coefficient of $v(x)$ above be zero. Hence

$$\frac{n(n-1)}{x^2} + \frac{np}{x} + q = 0 \tag{4}$$

Substituting the earlier values found for $p(x)$ and $q(x)$ into (4) gives

$$\frac{n(n-1)}{x^2} - \frac{5n}{x^2} + \frac{9}{x^2} = 0 \quad (5)$$

Solving (5) for n gives

$$n = 3 \quad (6)$$

Substituting this value in (3) gives

$$\begin{aligned} v''(x) + \frac{v'(x)}{x} &= 0 \\ v''(x) + \frac{v'(x)}{x} &= 0 \end{aligned} \quad (7)$$

Using the substitution

$$u(x) = v'(x)$$

Then (7) becomes

$$u'(x) + \frac{u(x)}{x} = 0 \quad (8)$$

The above is now solved for $u(x)$. In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{u}{x} \end{aligned}$$

Where $f(x) = -\frac{1}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned} \frac{1}{u} du &= -\frac{1}{x} dx \\ \int \frac{1}{u} du &= \int -\frac{1}{x} dx \\ \ln(u) &= -\ln(x) + c_1 \\ u &= e^{-\ln(x)+c_1} \\ &= \frac{c_1}{x} \end{aligned}$$

Now that $u(x)$ is known, then

$$\begin{aligned}v'(x) &= u(x) \\v(x) &= \int u(x) dx + c_2 \\&= c_1 \ln(x) + c_2\end{aligned}$$

Hence

$$\begin{aligned}y &= v(x) x^n \\&= (c_1 \ln(x) + c_2) x^3 \\&= (c_1 \ln(x) + c_2) x^3\end{aligned}$$

Now the particular solution to this ODE is found

$$x^2 y'' - 5xy' + 9y = x^2$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned}y_1 &= x^3 \\y_2 &= x^3 \ln(x)\end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} x^3 & x^3 \ln(x) \\ \frac{d}{dx}(x^3) & \frac{d}{dx}(x^3 \ln(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} x^3 & x^3 \ln(x) \\ 3x^2 & 3 \ln(x) x^2 + x^2 \end{vmatrix}$$

Therefore

$$W = (x^3) (3 \ln(x) x^2 + x^2) - (x^3 \ln(x)) (3x^2)$$

Which simplifies to

$$W = x^5$$

Which simplifies to

$$W = x^5$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^5 \ln(x)}{x^7} dx$$

Which simplifies to

$$u_1 = - \int \frac{\ln(x)}{x^2} dx$$

Hence

$$u_1 = - \frac{-1 - \ln(x)}{x}$$

And Eq. (3) becomes

$$u_2 = \int \frac{x^5}{x^7} dx$$

Which simplifies to

$$u_2 = \int \frac{1}{x^2} dx$$

Hence

$$u_2 = -\frac{1}{x}$$

Which simplifies to

$$u_1 = \frac{1 + \ln(x)}{x}$$

$$u_2 = -\frac{1}{x}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = (1 + \ln(x)) x^2 - \ln(x) x^2$$

Which simplifies to

$$y_p(x) = x^2$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= ((c_1 \ln(x) + c_2) x^3) + (x^2) \\ &= x^2 + (c_1 \ln(x) + c_2) x^3 \end{aligned}$$

Which simplifies to

$$y = x^2 + (c_1 \ln(x) + c_2) x^3$$

Summary

The solution(s) found are the following

$$y = x^2 + (c_1 \ln(x) + c_2) x^3 \tag{1}$$

Verification of solutions

$$y = x^2 + (c_1 \ln(x) + c_2) x^3$$

Verified OK. $\{0 < x\}$

16.4.5 Solving using Kovacic algorithm

Writing the ode as

$$x^2y'' - 5xy' + 9y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= x^2 \\ B &= -5x \\ C &= 9 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{4x^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 4x^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(-\frac{1}{4x^2}\right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 199: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = -\frac{1}{4x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = -\frac{1}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{1}{2} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{2} \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ , which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = -\frac{1}{4x^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = -\frac{1}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{1}{2} \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{2} \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = -\frac{1}{4x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	$\frac{1}{2}$	$\frac{1}{2}$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	$\frac{1}{2}$	$\frac{1}{2}$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω .

Trying $\alpha_{\infty}^{-} = \frac{1}{2}$ then

$$\begin{aligned} d &= \alpha_{\infty}^{-} - (\alpha_{c_1}^{-}) \\ &= \frac{1}{2} - \left(\frac{1}{2}\right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_{\infty}$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^{-}}{x - c_1} \right) + (-)[\sqrt{r}]_{\infty} \\ &= \frac{1}{2x} + (-)(0) \\ &= \frac{1}{2x} \\ &= \frac{1}{2x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r)p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2\left(\frac{1}{2x}\right)(0) + \left(\left(-\frac{1}{2x^2}\right) + \left(\frac{1}{2x}\right)^2 - \left(-\frac{1}{4x^2}\right) \right) &= 0 \\ 0 &= 0 \end{aligned}$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= pe^{\int \omega dx} \\ &= e^{\int \frac{1}{2x} dx} \\ &= \sqrt{x} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\&= z_1 e^{-\int \frac{1}{2} \frac{-5x}{x^2} dx} \\&= z_1 e^{\frac{5 \ln(x)}{2}} \\&= z_1 \left(x^{\frac{5}{2}} \right)\end{aligned}$$

Which simplifies to

$$y_1 = x^3$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{-5x}{x^2} dx}}{(y_1)^2} dx \\&= y_1 \int \frac{e^{5 \ln(x)}}{(y_1)^2} dx \\&= y_1 (\ln(x))\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\&= c_1 (x^3) + c_2 (x^3 (\ln(x)))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$.
 y_h is the solution to

$$x^2 y'' - 5x y' + 9y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1x^3 + c_2x^3 \ln(x)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1y_1 + u_2y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = x^3$$

$$y_2 = x^3 \ln(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} x^3 & x^3 \ln(x) \\ \frac{d}{dx}(x^3) & \frac{d}{dx}(x^3 \ln(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} x^3 & x^3 \ln(x) \\ 3x^2 & 3 \ln(x) x^2 + x^2 \end{vmatrix}$$

Therefore

$$W = (x^3) (3 \ln(x) x^2 + x^2) - (x^3 \ln(x)) (3x^2)$$

Which simplifies to

$$W = x^5$$

Which simplifies to

$$W = x^5$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^5 \ln(x)}{x^7} dx$$

Which simplifies to

$$u_1 = - \int \frac{\ln(x)}{x^2} dx$$

Hence

$$u_1 = - \frac{-1 - \ln(x)}{x}$$

And Eq. (3) becomes

$$u_2 = \int \frac{x^5}{x^7} dx$$

Which simplifies to

$$u_2 = \int \frac{1}{x^2} dx$$

Hence

$$u_2 = -\frac{1}{x}$$

Which simplifies to

$$u_1 = \frac{1 + \ln(x)}{x}$$

$$u_2 = -\frac{1}{x}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = (1 + \ln(x)) x^2 - \ln(x) x^2$$

Which simplifies to

$$y_p(x) = x^2$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 x^3 + c_2 x^3 \ln(x)) + (x^2) \end{aligned}$$

Which simplifies to

$$y = x^3(c_1 + c_2 \ln(x)) + x^2$$

Summary

The solution(s) found are the following

$$y = x^3(c_1 + c_2 \ln(x)) + x^2 \quad (1)$$

Verification of solutions

$$y = x^3(c_1 + c_2 \ln(x)) + x^2$$

Verified OK. $\{0 < x\}$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    <- LODE of Euler type successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(x^2*diff(y(x),x$2)-5*x*diff(y(x),x)+9*y(x)=x^2,y(x), singsol=all)
```

$$y(x) = x^2(\ln(x) c_1 x + c_2 x + 1)$$

✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 22

```
DSolve[x^2*y''[x]-5*x*y'[x]+9*y[x]==x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^2(c_1x + 3c_2x \log(x) + 1)$$

16.5 problem 1(e)

16.5.1 Maple step by step solution 1239

Internal problem ID [6035]

Internal file name [OUTPUT/5283_Sunday_June_05_2022_03_29_26_PM_70302027/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 149

Problem number: 1(e).

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_ODE_non_constant_coefficients_of_type_Euler**"

Maple gives the following as the ode type

```
[[_3rd_order , _exact , _linear , _homogeneous]]
```

$$x^3y''' + 2x^2y'' - xy' + y = 0$$

This is Euler ODE of higher order. Let $y = x^\lambda$. Hence

$$y' = \lambda x^{\lambda-1}$$

$$y'' = \lambda(\lambda - 1) x^{\lambda-2}$$

$$y''' = \lambda(\lambda - 1)(\lambda - 2) x^{\lambda-3}$$

Substituting these back into

$$x^3y''' + 2x^2y'' - xy' + y = 0$$

gives

$$-x\lambda x^{\lambda-1} + 2x^2\lambda(\lambda - 1) x^{\lambda-2} + x^3\lambda(\lambda - 1)(\lambda - 2) x^{\lambda-3} + x^\lambda = 0$$

Which simplifies to

$$-\lambda x^\lambda + 2\lambda(\lambda - 1) x^\lambda + \lambda(\lambda - 1)(\lambda - 2) x^\lambda + x^\lambda = 0$$

And since $x^\lambda \neq 0$ then dividing through by x^λ , the above becomes

$$-\lambda + 2\lambda(\lambda - 1) + \lambda(\lambda - 1)(\lambda - 2) + 1 = 0$$

Simplifying gives the characteristic equation as

$$(\lambda + 1)(\lambda - 1)^2 = 0$$

Solving the above gives the following roots

$$\lambda_1 = -1$$

$$\lambda_2 = 1$$

$$\lambda_3 = 1$$

This table summarises the result

root	multiplicity	type of root
-1	1	real root
1	2	real root

The solution is generated by going over the above table. For each real root λ of multiplicity one generates a c_1x^λ basis solution. Each real root of multiplicity two, generates c_1x^λ and $c_2x^\lambda \ln(x)$ basis solutions. Each real root of multiplicity three, generates c_1x^λ and $c_2x^\lambda \ln(x)$ and $c_3x^\lambda \ln(x)^2$ basis solutions, and so on. Each complex root $\alpha \pm i\beta$ of multiplicity one generates $x^\alpha(c_1 \cos(\beta \ln(x)) + c_2 \sin(\beta \ln(x)))$ basis solutions. And each complex root $\alpha \pm i\beta$ of multiplicity two generates $\ln(x) x^\alpha(c_1 \cos(\beta \ln(x)) + c_2 \sin(\beta \ln(x)))$ basis solutions. And each complex root $\alpha \pm i\beta$ of multiplicity three generates $\ln(x)^2 x^\alpha(c_1 \cos(\beta \ln(x)) + c_2 \sin(\beta \ln(x)))$ basis solutions. And so on. Using the above show that the solution is

$$y = \frac{c_1}{x} + c_2x + c_3 \ln(x) x$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = \frac{1}{x}$$

$$y_2 = x$$

$$y_3 = \ln(x) x$$

Summary

The solution(s) found are the following

$$y = \frac{c_1}{x} + c_2x + c_3 \ln(x) x \tag{1}$$

Verification of solutions

$$y = \frac{c_1}{x} + c_2x + c_3 \ln(x) x$$

Verified OK.

16.5.1 Maple step by step solution

Let's solve

$$x^3y''' + 2x^2y'' - xy' + y = 0$$

- Highest derivative means the order of the ODE is 3

$$y'''$$

- Isolate 3rd derivative

$$y''' = -\frac{y}{x^3} - \frac{2y''x - y'}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y''' + \frac{2y''}{x} - \frac{y'}{x^2} + \frac{y}{x^3} = 0$$

- Multiply by denominators of the ODE

$$x^3y''' + 2x^2y'' - xy' + y = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x , using the chain rule

$$y' = \left(\frac{d}{dt}y(t)\right) t'(x)$$

- Compute derivative

$$y' = \frac{\frac{d}{dt}y(t)}{x}$$

- Calculate the 2nd derivative of y with respect to x , using the chain rule

$$y'' = \left(\frac{d^2}{dt^2}y(t)\right) t'(x)^2 + t''(x) \left(\frac{d}{dt}y(t)\right)$$

- Compute derivative

$$y'' = \frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2}$$

- Calculate the 3rd derivative of y with respect to x , using the chain rule

$$y''' = \left(\frac{d^3}{dt^3}y(t)\right) t'(x)^3 + 3t'(x)t''(x) \left(\frac{d^2}{dt^2}y(t)\right) + t'''(x) \left(\frac{d}{dt}y(t)\right)$$

- Compute derivative

$$y''' = \frac{\frac{d^3}{dt^3}y(t)}{x^3} - \frac{3\left(\frac{d^2}{dt^2}y(t)\right)}{x^3} + \frac{2\left(\frac{d}{dt}y(t)\right)}{x^3}$$

Substitute the change of variables back into the ODE

$$x^3 \left(\frac{\frac{d^3}{dt^3}y(t)}{x^3} - \frac{3\left(\frac{d^2}{dt^2}y(t)\right)}{x^3} + \frac{2\left(\frac{d}{dt}y(t)\right)}{x^3} \right) + 2x^2 \left(\frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2} \right) - \frac{d}{dt}y(t) + y(t) = 0$$

- Simplify

$$\frac{d^3}{dt^3}y(t) - \frac{d^2}{dt^2}y(t) - \frac{d}{dt}y(t) + y(t) = 0$$

- Convert linear ODE into a system of first order ODEs

- Define new variable $y_1(t)$

$$y_1(t) = y(t)$$

- Define new variable $y_2(t)$

$$y_2(t) = \frac{d}{dt}y(t)$$

- Define new variable $y_3(t)$

$$y_3(t) = \frac{d^2}{dt^2}y(t)$$

- Isolate for $\frac{d}{dt}y_3(t)$ using original ODE

$$\frac{d}{dt}y_3(t) = y_3(t) + y_2(t) - y_1(t)$$

Convert linear ODE into a system of first order ODEs

$$\left[y_2(t) = \frac{d}{dt}y_1(t), y_3(t) = \frac{d}{dt}y_2(t), \frac{d}{dt}y_3(t) = y_3(t) + y_2(t) - y_1(t) \right]$$

- Define vector

$$\vec{y}(t) = \begin{bmatrix} y_1(t) \\ y_2(t) \\ y_3(t) \end{bmatrix}$$

- System to solve

$$\frac{d}{dt}\vec{y}(t) = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ -1 & 1 & 1 \end{bmatrix} \cdot \vec{y}(t)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ -1 & 1 & 1 \end{bmatrix}$$

- Rewrite the system as

$$\frac{d}{dt}\vec{y}(t) = A \cdot \vec{y}(t)$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[-1, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right], \left[1, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right], \left[1, \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[-1, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_1 = e^{-t} \cdot \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}$$

- Consider eigenpair, with eigenvalue of algebraic multiplicity 2

$$\left[1, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right]$$

- First solution from eigenvalue 1

$$\vec{y}_2(t) = e^t \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

- Form of the 2nd homogeneous solution where \vec{p} is to be solved for, $\lambda = 1$ is the eigenvalue, and

$$\vec{y}_3(t) = e^{\lambda t} (t\vec{v} + \vec{p})$$

- Note that the t multiplying \vec{v} makes this solution linearly independent to the 1st solution obtained

- Substitute $\vec{y}_3(t)$ into the homogeneous system

$$\lambda e^{\lambda t} (t\vec{v} + \vec{p}) + e^{\lambda t} \vec{v} = (e^{\lambda t} A) \cdot (t\vec{v} + \vec{p})$$

- Use the fact that \vec{v} is an eigenvector of A

$$\lambda e^{\lambda t} (t\vec{v} + \vec{p}) + e^{\lambda t} \vec{v} = e^{\lambda t} (\lambda t\vec{v} + A \cdot \vec{p})$$

- Simplify equation

$$\lambda \vec{p} + \vec{v} = A \cdot \vec{p}$$

- Make use of the identity matrix I

$$(\lambda \cdot I) \cdot \vec{p} + \vec{v} = A \cdot \vec{p}$$

- Condition \vec{p} must meet for $\vec{y}_3(t)$ to be a solution to the homogeneous system

$$(A - \lambda \cdot I) \cdot \vec{p} = -\vec{v}$$

- Choose \vec{p} to use in the second solution to the homogeneous system from eigenvalue 1

$$\left(\begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ -1 & 1 & 1 \end{bmatrix} - 1 \cdot \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right) \cdot \vec{p} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

- Choice of \vec{p}

$$\vec{p} = \begin{bmatrix} -1 \\ 0 \\ 0 \end{bmatrix}$$

- Second solution from eigenvalue 1

$$\vec{y}_3(t) = e^t \cdot \left(t \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + \begin{bmatrix} -1 \\ 0 \\ 0 \end{bmatrix} \right)$$

- General solution to the system of ODEs

$$\vec{y} = c_1 \vec{y}_1 + c_2 \vec{y}_2(t) + c_3 \vec{y}_3(t)$$

- Substitute solutions into the general solution

$$\vec{y} = c_1 e^{-t} \cdot \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} + c_2 e^t \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + c_3 e^t \cdot \left(t \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + \begin{bmatrix} -1 \\ 0 \\ 0 \end{bmatrix} \right)$$

- First component of the vector is the solution to the ODE

$$y(t) = c_1 e^{-t} + e^t((t-1)c_3 + c_2)$$

- Change variables back using $t = \ln(x)$

$$y = \frac{c_1}{x} + x((\ln(x) - 1)c_3 + c_2)$$

- Simplify

$$y = c_3 \ln(x) x + c_2 x - c_3 x + \frac{c_1}{x}$$

Maple trace

```

`Methods for third order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
<- LODE of Euler type successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 22

```
dsolve(x^3*diff(y(x),x$3)+2*x^2*diff(y(x),x$2)-x*diff(y(x),x)+y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{c_3 \ln(x) x^2 + c_2 x^2 + c_1}{x}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 22

```
DSolve[x^3*y'''[x]+2*x^2*y''[x]-x*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_1}{x} + c_2 x + c_3 x \log(x)$$

16.6 problem 2(a)

16.6.1 Solving as second order euler ode ode	1244
16.6.2 Solving as second order change of variable on x method 2 ode .	1248
16.6.3 Solving as second order change of variable on x method 1 ode .	1253
16.6.4 Solving as second order change of variable on y method 2 ode .	1258
16.6.5 Solving using Kovacic algorithm	1263

Internal problem ID [6036]

Internal file name [OUTPUT/5284_Sunday_June_05_2022_03_29_27_PM_41308682/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 149

Problem number: 2(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_euler_ode", "second_order_change_of_variable_on_x_method_1", "second_order_change_of_variable_on_x_method_2", "second_order_change_of_variable_on_y_method_2"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$x^2y'' + xy' + 4y = 1$$

16.6.1 Solving as second order euler ode ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = x^2, B = x, C = 4, f(x) = 1$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$x^2y'' + xy' + 4y = 0$$

This is Euler second order ODE. Let the solution be $y = x^r$, then $y' = rx^{r-1}$ and $y'' = r(r-1)x^{r-2}$. Substituting these back into the given ODE gives

$$x^2(r(r-1))x^{r-2} + xrx^{r-1} + 4x^r = 0$$

Simplifying gives

$$r(r-1)x^r + rx^r + 4x^r = 0$$

Since $x^r \neq 0$ then dividing throughout by x^r gives

$$r(r-1) + r + 4 = 0$$

Or

$$r^2 + 4 = 0 \tag{1}$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = -2i$$

$$r_2 = 2i$$

The roots are complex conjugate of each others. Let the roots be

$$r_1 = \alpha + i\beta$$

$$r_2 = \alpha - i\beta$$

Where in this case $\alpha = 0$ and $\beta = -2$. Hence the solution becomes

$$\begin{aligned} y &= c_1x^{r_1} + c_2x^{r_2} \\ &= c_1x^{\alpha+i\beta} + c_2x^{\alpha-i\beta} \\ &= x^\alpha (c_1x^{i\beta} + c_2x^{-i\beta}) \\ &= x^\alpha (c_1e^{\ln(x^{i\beta})} + c_2e^{\ln(x^{-i\beta})}) \\ &= x^\alpha (c_1e^{i(\beta \ln x)} + c_2e^{-i(\beta \ln x)}) \end{aligned}$$

Using the values for $\alpha = 0$, $\beta = -2$, the above becomes

$$y = x^0 (c_1e^{-2i \ln(x)} + c_2e^{2i \ln(x)})$$

Using Euler relation, the expression $c_1e^{iA} + c_2e^{-iA}$ is transformed to $c_1 \cos A + c_1 \sin A$ where the constants are free to change. Applying this to the above result gives

$$y = c_1 \cos (2 \ln (x)) + c_2 \sin (2 \ln (x))$$

Next, we find the particular solution to the ODE

$$x^2 y'' + xy' + 4y = 1$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \cos(2 \ln(x))$$

$$y_2 = -\sin(2 \ln(x))$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(2 \ln(x)) & -\sin(2 \ln(x)) \\ \frac{d}{dx}(\cos(2 \ln(x))) & \frac{d}{dx}(-\sin(2 \ln(x))) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(2 \ln(x)) & -\sin(2 \ln(x)) \\ -\frac{2 \sin(2 \ln(x))}{x} & -\frac{2 \cos(2 \ln(x))}{x} \end{vmatrix}$$

Therefore

$$W = (\cos(2 \ln(x))) \left(-\frac{2 \cos(2 \ln(x))}{x} \right) - (-\sin(2 \ln(x))) \left(-\frac{2 \sin(2 \ln(x))}{x} \right)$$

Which simplifies to

$$W = -\frac{2(\cos(2 \ln(x))^2 + \sin(2 \ln(x))^2)}{x}$$

Which simplifies to

$$W = -\frac{2}{x}$$

Therefore Eq. (2) becomes

$$u_1 = -\int \frac{-\sin(2 \ln(x))}{-2x} dx$$

Which simplifies to

$$u_1 = -\int \frac{\sin(2 \ln(x))}{2x} dx$$

Hence

$$u_1 = \frac{\cos(2 \ln(x))}{4}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\cos(2 \ln(x))}{-2x} dx$$

Which simplifies to

$$u_2 = \int -\frac{\cos(2 \ln(x))}{2x} dx$$

Hence

$$u_2 = -\frac{\sin(2 \ln(x))}{4}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{\cos(2 \ln(x))^2}{4} + \frac{\sin(2 \ln(x))^2}{4}$$

Which simplifies to

$$y_p(x) = \frac{1}{4}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \frac{1}{4} + c_1 \cos(2 \ln(x)) + c_2 \sin(2 \ln(x)) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{1}{4} + c_1 \cos(2 \ln(x)) + c_2 \sin(2 \ln(x)) \quad (1)$$

Verification of solutions

$$y = \frac{1}{4} + c_1 \cos(2 \ln(x)) + c_2 \sin(2 \ln(x))$$

Verified OK.

16.6.2 Solving as second order change of variable on x method 2 ode

This is second order non-homogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$x^2 y'' + xy' + 4y = 0$$

In normal form the ode

$$x^2 y'' + xy' + 4y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$\begin{aligned} p(x) &= \frac{1}{x} \\ q(x) &= \frac{4}{x^2} \end{aligned}$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x)\tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned} \tau &= \int e^{-(\int p(x)dx)} dx \\ &= \int e^{-(\int \frac{1}{x} dx)} dx \\ &= \int e^{-\ln(x)} dx \\ &= \int \frac{1}{x} dx \\ &= \ln(x) \end{aligned} \quad (6)$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned} q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\ &= \frac{4}{x^2} \\ &= \frac{1}{x^2} \\ &= 4 \end{aligned} \quad (7)$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned} \frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) + 4y(\tau) &= 0 \end{aligned}$$

The above ode is now solved for $y(\tau)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(\tau) + By'(\tau) + Cy(\tau) = 0$$

Where in the above $A = 1, B = 0, C = 4$. Let the solution be $y(\tau) = e^{\lambda\tau}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda\tau} + 4 e^{\lambda\tau} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda\tau}$ gives

$$\lambda^2 + 4 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 4$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(4)} \\ &= \pm 2i \end{aligned}$$

Hence

$$\lambda_1 = +2i$$

$$\lambda_2 = -2i$$

Which simplifies to

$$\lambda_1 = 2i$$

$$\lambda_2 = -2i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 2$. Therefore the final solution, when using Euler relation, can be written as

$$y(\tau) = e^{\alpha\tau} (c_1 \cos(\beta\tau) + c_2 \sin(\beta\tau))$$

Which becomes

$$y(\tau) = e^0 (c_1 \cos(2\tau) + c_2 \sin(2\tau))$$

Or

$$y(\tau) = c_1 \cos(2\tau) + c_2 \sin(2\tau)$$

The above solution is now transformed back to y using (6) which results in

$$y = c_1 \cos(2 \ln(x)) + c_2 \sin(2 \ln(x))$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \cos(2 \ln(x)) + c_2 \sin(2 \ln(x))$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \sin(\ln(x)) \cos(\ln(x))$$

$$y_2 = 2 \cos(\ln(x))^2 - 1$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \sin(\ln(x)) \cos(\ln(x)) & 2 \cos(\ln(x))^2 - 1 \\ \frac{d}{dx}(\sin(\ln(x)) \cos(\ln(x))) & \frac{d}{dx}(2 \cos(\ln(x))^2 - 1) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \sin(\ln(x)) \cos(\ln(x)) & 2 \cos(\ln(x))^2 - 1 \\ \frac{\cos(\ln(x))^2}{x} - \frac{\sin(\ln(x))^2}{x} & -\frac{4 \cos(\ln(x)) \sin(\ln(x))}{x} \end{vmatrix}$$

Therefore

$$W = (\sin(\ln(x)) \cos(\ln(x))) \left(-\frac{4 \cos(\ln(x)) \sin(\ln(x))}{x} \right) - (2 \cos(\ln(x))^2 - 1) \left(\frac{\cos(\ln(x))^2}{x} - \frac{\sin(\ln(x))^2}{x} \right)$$

Which simplifies to

$$W = -\frac{2 \sin(\ln(x))^2 \cos(\ln(x))^2 + 2 \cos(\ln(x))^4 + \sin(\ln(x))^2 - \cos(\ln(x))^2}{x}$$

Which simplifies to

$$W = -\frac{1}{x}$$

Therefore Eq. (2) becomes

$$u_1 = -\int \frac{2 \cos(\ln(x))^2 - 1}{-x} dx$$

Which simplifies to

$$u_1 = -\int -\frac{\cos(2 \ln(x))}{x} dx$$

Hence

$$u_1 = \frac{\sin(2 \ln(x))}{2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\sin(\ln(x)) \cos(\ln(x))}{-x} dx$$

Which simplifies to

$$u_2 = \int -\frac{\sin(2 \ln(x))}{2x} dx$$

Hence

$$u_2 = \frac{\cos(2 \ln(x))}{4}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{\sin(2 \ln(x)) \sin(\ln(x)) \cos(\ln(x))}{2} + \frac{\cos(2 \ln(x)) (2 \cos(\ln(x))^2 - 1)}{4}$$

Which simplifies to

$$y_p(x) = \frac{1}{4}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(2 \ln(x)) + c_2 \sin(2 \ln(x))) + \left(\frac{1}{4}\right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{1}{4} + c_1 \cos(2 \ln(x)) + c_2 \sin(2 \ln(x)) \quad (1)$$

Verification of solutions

$$y = \frac{1}{4} + c_1 \cos(2 \ln(x)) + c_2 \sin(2 \ln(x))$$

Verified OK.

16.6.3 Solving as second order change of variable on x method 1 ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = x^2, B = x, C = 4, f(x) = 1$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$x^2 y'' + xy' + 4y = 0$$

In normal form the ode

$$x^2 y'' + x y' + 4y = 0 \quad (1)$$

Becomes

$$y'' + p(x) y' + q(x) y = 0 \quad (2)$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = \frac{4}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) results

$$\frac{d^2}{d\tau^2} y(\tau) + p_1 \left(\frac{d}{d\tau} y(\tau) \right) + q_1 y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x) \tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $q_1 = c^2$ where c is some constant. Therefore from (5)

$$\tau' = \frac{1}{c} \sqrt{q}$$
$$= \frac{2\sqrt{\frac{1}{x^2}}}{c} \quad (6)$$

$$\tau'' = -\frac{2}{c\sqrt{\frac{1}{x^2}} x^3}$$

Substituting the above into (4) results in

$$p_1(\tau) = \frac{\tau''(x) + p(x) \tau'(x)}{\tau'(x)^2}$$
$$= \frac{-\frac{2}{c\sqrt{\frac{1}{x^2}} x^3} + \frac{1}{x} \frac{2\sqrt{\frac{1}{x^2}}}{c}}{\left(\frac{2\sqrt{\frac{1}{x^2}}}{c}\right)^2}$$
$$= 0$$

Therefore ode (3) now becomes

$$\begin{aligned} y(\tau)'' + p_1 y(\tau)' + q_1 y(\tau) &= 0 \\ \frac{d^2}{d\tau^2} y(\tau) + c^2 y(\tau) &= 0 \end{aligned} \quad (7)$$

The above ode is now solved for $y(\tau)$. Since the ode is now constant coefficients, it can be easily solved to give

$$y(\tau) = c_1 \cos(c\tau) + c_2 \sin(c\tau)$$

Now from (6)

$$\begin{aligned} \tau &= \int \frac{1}{c} \sqrt{q} dx \\ &= \frac{\int 2\sqrt{\frac{1}{x^2}} dx}{c} \\ &= \frac{2\sqrt{\frac{1}{x^2}} x \ln(x)}{c} \end{aligned}$$

Substituting the above into the solution obtained gives

$$y = c_1 \cos(2 \ln(x)) + c_2 \sin(2 \ln(x))$$

Now the particular solution to this ODE is found

$$x^2 y'' + xy' + 4y = 1$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned} y_1 &= \sin(\ln(x)) \cos(\ln(x)) \\ y_2 &= 2 \cos(\ln(x))^2 - 1 \end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \sin(\ln(x)) \cos(\ln(x)) & 2 \cos(\ln(x))^2 - 1 \\ \frac{d}{dx}(\sin(\ln(x)) \cos(\ln(x))) & \frac{d}{dx}(2 \cos(\ln(x))^2 - 1) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \sin(\ln(x)) \cos(\ln(x)) & 2 \cos(\ln(x))^2 - 1 \\ \frac{\cos(\ln(x))^2}{x} - \frac{\sin(\ln(x))^2}{x} & -\frac{4 \cos(\ln(x)) \sin(\ln(x))}{x} \end{vmatrix}$$

Therefore

$$\begin{aligned} W &= (\sin(\ln(x)) \cos(\ln(x))) \left(-\frac{4 \cos(\ln(x)) \sin(\ln(x))}{x} \right) \\ &\quad - (2 \cos(\ln(x))^2 - 1) \left(\frac{\cos(\ln(x))^2}{x} - \frac{\sin(\ln(x))^2}{x} \right) \end{aligned}$$

Which simplifies to

$$W = -\frac{2 \sin(\ln(x))^2 \cos(\ln(x))^2 + 2 \cos(\ln(x))^4 + \sin(\ln(x))^2 - \cos(\ln(x))^2}{x}$$

Which simplifies to

$$W = -\frac{1}{x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{2 \cos(\ln(x))^2 - 1}{-x} dx$$

Which simplifies to

$$u_1 = - \int -\frac{\cos(2 \ln(x))}{x} dx$$

Hence

$$u_1 = \frac{\sin(2 \ln(x))}{2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\sin(\ln(x)) \cos(\ln(x))}{-x} dx$$

Which simplifies to

$$u_2 = \int -\frac{\sin(2 \ln(x))}{2x} dx$$

Hence

$$u_2 = \frac{\cos(2 \ln(x))}{4}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{\sin(2 \ln(x)) \sin(\ln(x)) \cos(\ln(x))}{2} + \frac{\cos(2 \ln(x)) (2 \cos(\ln(x))^2 - 1)}{4}$$

Which simplifies to

$$y_p(x) = \frac{1}{4}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(2 \ln(x)) + c_2 \sin(2 \ln(x))) + \left(\frac{1}{4}\right) \\ &= \frac{1}{4} + c_1 \cos(2 \ln(x)) + c_2 \sin(2 \ln(x)) \end{aligned}$$

Which simplifies to

$$y = \frac{1}{4} + c_1 \cos(2 \ln(x)) + c_2 \sin(2 \ln(x))$$

Summary

The solution(s) found are the following

$$y = \frac{1}{4} + c_1 \cos(2 \ln(x)) + c_2 \sin(2 \ln(x)) \quad (1)$$

Verification of solutions

$$y = \frac{1}{4} + c_1 \cos(2 \ln(x)) + c_2 \sin(2 \ln(x))$$

Verified OK.

16.6.4 Solving as second order change of variable on y method 2 ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = x^2$, $B = x$, $C = 4$, $f(x) = 1$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$x^2 y'' + xy' + 4y = 0$$

In normal form the ode

$$x^2 y'' + xy' + 4y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = \frac{4}{x^2}$$

Applying change of variables on the dependent variable $y = v(x)x^n$ to (2) gives the following ode where the dependent variables is $v(x)$ and not y .

$$v''(x) + \left(\frac{2n}{x} + p\right)v'(x) + \left(\frac{n(n-1)}{x^2} + \frac{np}{x} + q\right)v(x) = 0 \quad (3)$$

Let the coefficient of $v(x)$ above be zero. Hence

$$\frac{n(n-1)}{x^2} + \frac{np}{x} + q = 0 \quad (4)$$

Substituting the earlier values found for $p(x)$ and $q(x)$ into (4) gives

$$\frac{n(n-1)}{x^2} + \frac{n}{x^2} + \frac{4}{x^2} = 0 \quad (5)$$

Solving (5) for n gives

$$n = 2i \quad (6)$$

Substituting this value in (3) gives

$$\begin{aligned} v''(x) + \left(\frac{4i}{x} + \frac{1}{x} \right) v'(x) &= 0 \\ v''(x) + \frac{(1+4i)v'(x)}{x} &= 0 \end{aligned} \quad (7)$$

Using the substitution

$$u(x) = v'(x)$$

Then (7) becomes

$$u'(x) + \frac{(1+4i)u(x)}{x} = 0 \quad (8)$$

The above is now solved for $u(x)$. In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= \frac{(-1-4i)u}{x} \end{aligned}$$

Where $f(x) = \frac{-1-4i}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned} \frac{1}{u} du &= \frac{-1-4i}{x} dx \\ \int \frac{1}{u} du &= \int \frac{-1-4i}{x} dx \\ \ln(u) &= (-1-4i) \ln(x) + c_1 \\ u &= e^{(-1-4i) \ln(x) + c_1} \\ &= c_1 e^{(-1-4i) \ln(x)} \end{aligned}$$

Which simplifies to

$$u(x) = \frac{c_1 x^{-4i}}{x}$$

Now that $u(x)$ is known, then

$$\begin{aligned}v'(x) &= u(x) \\v(x) &= \int u(x) dx + c_2 \\&= \frac{ic_1 x^{-4i}}{4} + c_2\end{aligned}$$

Hence

$$\begin{aligned}y &= v(x) x^n \\&= \left(\frac{ic_1 x^{-4i}}{4} + c_2 \right) x^{2i} \\&= x^{2i} c_2 + \frac{ix^{-2i} c_1}{4}\end{aligned}$$

Now the particular solution to this ODE is found

$$x^2 y'' + xy' + 4y = 1$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned}y_1 &= x^{2i} \\y_2 &= x^{2i} x^{-4i}\end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} x^{2i} & x^{2i}x^{-4i} \\ \frac{d}{dx}(x^{2i}) & \frac{d}{dx}(x^{2i}x^{-4i}) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} x^{2i} & x^{2i}x^{-4i} \\ \frac{2ix^{2i}}{x} & -\frac{2ix^{2i}x^{-4i}}{x} \end{vmatrix}$$

Therefore

$$W = (x^{2i}) \left(-\frac{2ix^{2i}x^{-4i}}{x} \right) - (x^{2i}x^{-4i}) \left(\frac{2ix^{2i}}{x} \right)$$

Which simplifies to

$$W = -\frac{4ix^{4i}x^{-4i}}{x}$$

Which simplifies to

$$W = -\frac{4i}{x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^{2i}x^{-4i}}{-4ix} dx$$

Which simplifies to

$$u_1 = - \int \frac{ix^{-1-2i}}{4} dx$$

Hence

$$u_1 = \text{undefined}$$

And Eq. (3) becomes

$$u_2 = \int \frac{x^{2i}}{-4ix} dx$$

Which simplifies to

$$u_2 = \int \frac{ix^{-1+2i}}{4} dx$$

Hence

$$u_2 = \text{undefined}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \text{undefined } x^{2i} + \text{undefined } x^{2i} x^{-4i}$$

Which simplifies to

$$y_p(x) = \text{undefined } (x^{2i} + x^{-2i})$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(\left(\frac{ic_1 x^{-4i}}{4} + c_2 \right) x^{2i} \right) + (\text{undefined } (x^{2i} + x^{-2i})) \\ &= \text{undefined } (x^{2i} + x^{-2i}) + \left(\frac{ic_1 x^{-4i}}{4} + c_2 \right) x^{2i} \end{aligned}$$

Which simplifies to

$$y = \frac{(ic_1 + \text{undefined}) x^{-2i}}{4} + (\text{undefined} + c_2) x^{2i}$$

Summary

The solution(s) found are the following

$$y = \frac{(ic_1 + \text{undefined}) x^{-2i}}{4} + (\text{undefined} + c_2) x^{2i} \quad (1)$$

Verification of solutions

$$y = \frac{(ic_1 + \text{undefined}) x^{-2i}}{4} + (\text{undefined} + c_2) x^{2i}$$

Verified OK.

16.6.5 Solving using Kovacic algorithm

Writing the ode as

$$x^2 y'' + xy' + 4y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= x^2 \\ B &= x \\ C &= 4 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-17}{4x^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -17 \\ t &= 4x^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(-\frac{17}{4x^2}\right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 201: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = -\frac{17}{4x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = -\frac{17}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{1}{2} + 2i \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{2} - 2i \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = -\frac{17}{4x^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = -\frac{17}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{1}{2} + 2i \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{2} - 2i \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = -\frac{17}{4x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	$\frac{1}{2} + 2i$	$\frac{1}{2} - 2i$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	$\frac{1}{2} + 2i$	$\frac{1}{2} - 2i$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to

determine possible non negative integer d from these using

$$d = \alpha_{\infty}^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_{∞}^{\pm} . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_{\infty}^{-} = \frac{1}{2} - 2i$ then

$$\begin{aligned} d &= \alpha_{\infty}^{-} - (\alpha_{c_1}^{-}) \\ &= \frac{1}{2} - 2i - \left(\frac{1}{2} - 2i \right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_{\infty}$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^{-}}{x - c_1} \right) + (-)[\sqrt{r}]_{\infty} \\ &= \frac{\frac{1}{2} - 2i}{x} + (-)(0) \\ &= \frac{\frac{1}{2} - 2i}{x} \\ &= \frac{\frac{1}{2} - 2i}{x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \tag{1A}$$

Let

$$p(x) = 1 \tag{2A}$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2 \left(\frac{\frac{1}{2} - 2i}{x} \right) (0) + \left(\left(\frac{-\frac{1}{2} + 2i}{x^2} \right) + \left(\frac{\frac{1}{2} - 2i}{x} \right)^2 - \left(-\frac{17}{4x^2} \right) \right) &= 0 \\ 0 &= 0 \end{aligned}$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= pe^{\int \omega dx} \\ &= e^{\int \frac{1}{2} - 2i dx} \\ &= x^{\frac{1}{2} - 2i} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{x}{x^2} dx} \\ &= z_1 e^{-\frac{\ln(x)}{2}} \\ &= z_1 \left(\frac{1}{\sqrt{x}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = x^{-2i}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{x}{x^2} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-\ln(x)}}{(y_1)^2} dx \\ &= y_1 \left(-\frac{ix^{4i}}{4} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (x^{-2i}) + c_2 \left(x^{-2i} \left(-\frac{ix^{4i}}{4} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$x^2y'' + xy' + 4y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = x^{-2i}c_1 - \frac{ic_2x^{2i}}{4}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1y_1 + u_2y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = x^{-2i}$$

$$y_2 = -\frac{ix^{2i}}{4}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = -\int \frac{y_2f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} x^{-2i} & -\frac{ix^{2i}}{4} \\ \frac{d}{dx}(x^{-2i}) & \frac{d}{dx}\left(-\frac{ix^{2i}}{4}\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} x^{-2i} & -\frac{ix^{2i}}{4} \\ -\frac{2ix^{-2i}}{x} & \frac{x^{2i}}{2x} \end{vmatrix}$$

Therefore

$$W = (x^{-2i}) \left(\frac{x^{2i}}{2x} \right) - \left(-\frac{ix^{2i}}{4} \right) \left(-\frac{2ix^{-2i}}{x} \right)$$

Which simplifies to

$$W = \frac{x^{2i}x^{-2i}}{x}$$

Which simplifies to

$$W = \frac{1}{x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{-\frac{ix^{2i}}{4}}{x} dx$$

Which simplifies to

$$u_1 = - \int -\frac{ix^{-1+2i}}{4} dx$$

Hence

$$u_1 = \text{undefined}$$

And Eq. (3) becomes

$$u_2 = \int \frac{x^{-2i}}{x} dx$$

Which simplifies to

$$u_2 = \int x^{-1-2i} dx$$

Hence

$$u_2 = \text{undefined}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{1}{4} x^{-2i} - i \frac{1}{4} x^{2i}$$

Which simplifies to

$$y_p(x) = \frac{1}{4} (ix^{2i} + x^{-2i})$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(x^{-2i} c_1 - \frac{ic_2 x^{2i}}{4} \right) + \frac{1}{4} (ix^{2i} + x^{-2i}) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = x^{-2i} c_1 - \frac{ic_2 x^{2i}}{4} + \frac{1}{4} (ix^{2i} + x^{-2i}) \quad (1)$$

Verification of solutions

$$y = x^{-2i} c_1 - \frac{ic_2 x^{2i}}{4} + \frac{1}{4} (ix^{2i} + x^{-2i})$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    <- LODE of Euler type successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 20

```
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)+4*y(x)=1,y(x), singsol=all)
```

$$y(x) = \sin(2 \ln(x)) c_2 + \cos(2 \ln(x)) c_1 + \frac{1}{4}$$

✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 25

```
DSolve[x^2*y''[x]+x*y'[x]+4*y[x]==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos(2 \log(x)) + c_2 \sin(2 \log(x)) + \frac{1}{4}$$

16.7 problem 2(b)

16.7.1 Solving as second order euler ode ode	1272
16.7.2 Solving as second order change of variable on x method 2 ode .	1274
16.7.3 Solving as second order change of variable on x method 1 ode .	1277
16.7.4 Solving as second order change of variable on y method 2 ode .	1279
16.7.5 Solving using Kovacic algorithm	1281
16.7.6 Maple step by step solution	1286

Internal problem ID [6037]

Internal file name [OUTPUT/5285_Sunday_June_05_2022_03_29_29_PM_62290384/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 149

Problem number: 2(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_euler_ode", "second_order_change_of_variable_on_x_method_1", "second_order_change_of_variable_on_x_method_2", "second_order_change_of_variable_on_y_method_2"

Maple gives the following as the ode type

```
[[_Emden, _Fowler]]
```

$$x^2y'' - 3xy' + 5y = 0$$

16.7.1 Solving as second order euler ode ode

This is Euler second order ODE. Let the solution be $y = x^r$, then $y' = rx^{r-1}$ and $y'' = r(r-1)x^{r-2}$. Substituting these back into the given ODE gives

$$x^2(r(r-1))x^{r-2} - 3rx^{r-1} + 5x^r = 0$$

Simplifying gives

$$r(r-1)x^r - 3rx^r + 5x^r = 0$$

Since $x^r \neq 0$ then dividing throughout by x^r gives

$$r(r-1) - 3r + 5 = 0$$

Or

$$r^2 - 4r + 5 = 0 \quad (1)$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = 2 - i$$

$$r_2 = 2 + i$$

The roots are complex conjugate of each others. Let the roots be

$$r_1 = \alpha + i\beta$$

$$r_2 = \alpha - i\beta$$

Where in this case $\alpha = 2$ and $\beta = -1$. Hence the solution becomes

$$\begin{aligned} y &= c_1 x^{r_1} + c_2 x^{r_2} \\ &= c_1 x^{\alpha+i\beta} + c_2 x^{\alpha-i\beta} \\ &= x^\alpha (c_1 x^{i\beta} + c_2 x^{-i\beta}) \\ &= x^\alpha (c_1 e^{\ln(x^{i\beta})} + c_2 e^{\ln(x^{-i\beta})}) \\ &= x^\alpha (c_1 e^{i(\beta \ln x)} + c_2 e^{-i(\beta \ln x)}) \end{aligned}$$

Using the values for $\alpha = 2, \beta = -1$, the above becomes

$$y = x^2 (c_1 e^{-i \ln(x)} + c_2 e^{i \ln(x)})$$

Using Euler relation, the expression $c_1 e^{iA} + c_2 e^{-iA}$ is transformed to $c_1 \cos A + c_1 \sin A$ where the constants are free to change. Applying this to the above result gives

$$y = x^2 (c_1 \cos (\ln (x)) + c_2 \sin (\ln (x)))$$

Summary

The solution(s) found are the following

$$y = x^2 (c_1 \cos (\ln (x)) + c_2 \sin (\ln (x))) \quad (1)$$

Verification of solutions

$$y = x^2 (c_1 \cos (\ln (x)) + c_2 \sin (\ln (x)))$$

Verified OK.

16.7.2 Solving as second order change of variable on x method 2 ode

In normal form the ode

$$x^2 y'' - 3xy' + 5y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = -\frac{3}{x}$$
$$q(x) = \frac{5}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x)\tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned} \tau &= \int e^{-(\int p(x)dx)} dx \\ &= \int e^{-(\int -\frac{3}{x}dx)} dx \\ &= \int e^{3\ln(x)} dx \\ &= \int x^3 dx \\ &= \frac{x^4}{4} \end{aligned} \quad (6)$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned} q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\ &= \frac{\frac{5}{x^2}}{x^6} \\ &= \frac{5}{x^8} \end{aligned} \tag{7}$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned} \frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) + \frac{5y(\tau)}{x^8} &= 0 \end{aligned}$$

But in terms of τ

$$\frac{5}{x^8} = \frac{5}{16\tau^2}$$

Hence the above ode becomes

$$\frac{d^2}{d\tau^2}y(\tau) + \frac{5y(\tau)}{16\tau^2} = 0$$

The above ode is now solved for $y(\tau)$. The ode can be written as

$$16 \left(\frac{d^2}{d\tau^2}y(\tau) \right) \tau^2 + 5y(\tau) = 0$$

Which shows it is a Euler ODE. This is Euler second order ODE. Let the solution be $y(\tau) = \tau^r$, then $y' = r\tau^{r-1}$ and $y'' = r(r-1)\tau^{r-2}$. Substituting these back into the given ODE gives

$$16\tau^2(r(r-1))\tau^{r-2} + 0r\tau^{r-1} + 5\tau^r = 0$$

Simplifying gives

$$16r(r-1)\tau^r + 0\tau^r + 5\tau^r = 0$$

Since $\tau^r \neq 0$ then dividing throughout by τ^r gives

$$16r(r-1) + 0 + 5 = 0$$

Or

$$16r^2 - 16r + 5 = 0 \tag{1}$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = \frac{1}{2} - \frac{i}{4}$$

$$r_2 = \frac{1}{2} + \frac{i}{4}$$

The roots are complex conjugate of each others. Let the roots be

$$r_1 = \alpha + i\beta$$

$$r_2 = \alpha - i\beta$$

Where in this case $\alpha = \frac{1}{2}$ and $\beta = -\frac{1}{4}$. Hence the solution becomes

$$\begin{aligned} y(\tau) &= c_1\tau^{r_1} + c_2\tau^{r_2} \\ &= c_1\tau^{\alpha+i\beta} + c_2\tau^{\alpha-i\beta} \\ &= \tau^\alpha(c_1\tau^{i\beta} + c_2\tau^{-i\beta}) \\ &= \tau^\alpha(c_1e^{\ln(\tau^{i\beta})} + c_2e^{\ln(\tau^{-i\beta})}) \\ &= \tau^\alpha(c_1e^{i(\beta \ln \tau)} + c_2e^{-i(\beta \ln \tau)}) \end{aligned}$$

Using the values for $\alpha = \frac{1}{2}, \beta = -\frac{1}{4}$, the above becomes

$$y(\tau) = \tau^{\frac{1}{2}} \left(c_1 e^{-\frac{i \ln(\tau)}{4}} + c_2 e^{\frac{i \ln(\tau)}{4}} \right)$$

Using Euler relation, the expression $c_1 e^{iA} + c_2 e^{-iA}$ is transformed to $c_1 \cos A + c_1 \sin A$ where the constants are free to change. Applying this to the above result gives

$$y(\tau) = \sqrt{\tau} \left(c_1 \cos \left(\frac{\ln(\tau)}{4} \right) + c_2 \sin \left(\frac{\ln(\tau)}{4} \right) \right)$$

The above solution is now transformed back to y using (6) which results in

$$y = \frac{\left(c_1 \cos \left(-\frac{\ln(2)}{2} + \ln(x) \right) + c_2 \sin \left(-\frac{\ln(2)}{2} + \ln(x) \right) \right) x^2}{2}$$

Summary

The solution(s) found are the following

$$y = \frac{\left(c_1 \cos \left(-\frac{\ln(2)}{2} + \ln(x) \right) + c_2 \sin \left(-\frac{\ln(2)}{2} + \ln(x) \right) \right) x^2}{2} \quad (1)$$

Verification of solutions

$$y = \frac{\left(c_1 \cos \left(-\frac{\ln(2)}{2} + \ln(x) \right) + c_2 \sin \left(-\frac{\ln(2)}{2} + \ln(x) \right) \right) x^2}{2}$$

Verified OK.

16.7.3 Solving as second order change of variable on x method 1 ode

In normal form the ode

$$x^2 y'' - 3xy' + 5y = 0 \quad (1)$$

Becomes

$$y'' + p(x) y' + q(x) y = 0 \quad (2)$$

Where

$$p(x) = -\frac{3}{x}$$

$$q(x) = \frac{5}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) results

$$\frac{d^2}{d\tau^2} y(\tau) + p_1 \left(\frac{d}{d\tau} y(\tau) \right) + q_1 y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x) \tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $q_1 = c^2$ where c is some constant. Therefore from (5)

$$\begin{aligned} \tau' &= \frac{1}{c} \sqrt{q} \\ &= \frac{\sqrt{5} \sqrt{\frac{1}{x^2}}}{c} \end{aligned} \quad (6)$$

$$\tau'' = -\frac{\sqrt{5}}{c \sqrt{\frac{1}{x^2}} x^3}$$

Substituting the above into (4) results in

$$\begin{aligned}
 p_1(\tau) &= \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \\
 &= \frac{-\frac{\sqrt{5}}{c\sqrt{\frac{1}{x^2}}x^3} - \frac{3}{x}\frac{\sqrt{5}\sqrt{\frac{1}{x^2}}}{c}}{\left(\frac{\sqrt{5}\sqrt{\frac{1}{x^2}}}{c}\right)^2} \\
 &= -\frac{4c\sqrt{5}}{5}
 \end{aligned}$$

Therefore ode (3) now becomes

$$\begin{aligned}
 y(\tau)'' + p_1y(\tau)' + q_1y(\tau) &= 0 \\
 \frac{d^2}{d\tau^2}y(\tau) - \frac{4c\sqrt{5}}{5}\left(\frac{d}{d\tau}y(\tau)\right) + c^2y(\tau) &= 0 \tag{7}
 \end{aligned}$$

The above ode is now solved for $y(\tau)$. Since the ode is now constant coefficients, it can be easily solved to give

$$y(\tau) = e^{\frac{2\sqrt{5}c\tau}{5}} \left(c_1 \cos\left(\frac{\sqrt{5}c\tau}{5}\right) + c_2 \sin\left(\frac{\sqrt{5}c\tau}{5}\right) \right)$$

Now from (6)

$$\begin{aligned}
 \tau &= \int \frac{1}{c}\sqrt{q} dx \\
 &= \frac{\int \sqrt{5}\sqrt{\frac{1}{x^2}} dx}{c} \\
 &= \frac{\sqrt{5}\sqrt{\frac{1}{x^2}} x \ln(x)}{c}
 \end{aligned}$$

Substituting the above into the solution obtained gives

$$y = x^2(c_1 \cos(\ln(x)) + c_2 \sin(\ln(x)))$$

Summary

The solution(s) found are the following

$$y = x^2(c_1 \cos(\ln(x)) + c_2 \sin(\ln(x))) \tag{1}$$

Verification of solutions

$$y = x^2(c_1 \cos(\ln(x)) + c_2 \sin(\ln(x)))$$

Verified OK.

16.7.4 Solving as second order change of variable on y method 2 ode

In normal form the ode

$$x^2 y'' - 3xy' + 5y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = -\frac{3}{x}$$
$$q(x) = \frac{5}{x^2}$$

Applying change of variables on the dependent variable $y = v(x)x^n$ to (2) gives the following ode where the dependent variables is $v(x)$ and not y .

$$v''(x) + \left(\frac{2n}{x} + p\right)v'(x) + \left(\frac{n(n-1)}{x^2} + \frac{np}{x} + q\right)v(x) = 0 \quad (3)$$

Let the coefficient of $v(x)$ above be zero. Hence

$$\frac{n(n-1)}{x^2} + \frac{np}{x} + q = 0 \quad (4)$$

Substituting the earlier values found for $p(x)$ and $q(x)$ into (4) gives

$$\frac{n(n-1)}{x^2} - \frac{3n}{x^2} + \frac{5}{x^2} = 0 \quad (5)$$

Solving (5) for n gives

$$n = 2 + i \quad (6)$$

Substituting this value in (3) gives

$$v''(x) + \left(\frac{4+2i}{x} - \frac{3}{x}\right)v'(x) = 0$$
$$v''(x) + \frac{(1+2i)v'(x)}{x} = 0 \quad (7)$$

Using the substitution

$$u(x) = v'(x)$$

Then (7) becomes

$$u'(x) + \frac{(1 + 2i)u(x)}{x} = 0 \quad (8)$$

The above is now solved for $u(x)$. In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= \frac{(-1 - 2i)u}{x} \end{aligned}$$

Where $f(x) = \frac{-1-2i}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned} \frac{1}{u} du &= \frac{-1 - 2i}{x} dx \\ \int \frac{1}{u} du &= \int \frac{-1 - 2i}{x} dx \\ \ln(u) &= (-1 - 2i) \ln(x) + c_1 \\ u &= e^{(-1-2i)\ln(x)+c_1} \\ &= c_1 e^{(-1-2i)\ln(x)} \end{aligned}$$

Which simplifies to

$$u(x) = \frac{c_1 x^{-2i}}{x}$$

Now that $u(x)$ is known, then

$$\begin{aligned} v'(x) &= u(x) \\ v(x) &= \int u(x) dx + c_2 \\ &= \frac{ix^{-2i}c_1}{2} + c_2 \end{aligned}$$

Hence

$$\begin{aligned} y &= v(x) x^n \\ &= \left(\frac{ix^{-2i}c_1}{2} + c_2 \right) x^{2+i} \\ &= c_2 x^{2+i} + \frac{ic_1 x^{2-i}}{2} \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \left(\frac{ix^{-2i}c_1}{2} + c_2 \right) x^{2+i} \quad (1)$$

Verification of solutions

$$y = \left(\frac{ix^{-2i}c_1}{2} + c_2 \right) x^{2+i}$$

Verified OK.

16.7.5 Solving using Kovacic algorithm

Writing the ode as

$$x^2y'' - 3xy' + 5y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= x^2 \\ B &= -3x \\ C &= 5 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-5}{4x^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -5 \\ t &= 4x^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(-\frac{5}{4x^2}\right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 202: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole

larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = -\frac{5}{4x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = -\frac{5}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{1}{2} + i \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{2} - i \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = -\frac{5}{4x^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = -\frac{5}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{1}{2} + i \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{2} - i \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = -\frac{5}{4x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	$\frac{1}{2} + i$	$\frac{1}{2} - i$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	$\frac{1}{2} + i$	$\frac{1}{2} - i$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = \frac{1}{2} - i$ then

$$\begin{aligned} d &= \alpha_\infty^- - (\alpha_{c_1}^-) \\ &= \frac{1}{2} - i - \left(\frac{1}{2} - i \right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c) [\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty) [\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= \left((-) [\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{x - c_1} \right) + (-) [\sqrt{r}]_\infty \\ &= \frac{\frac{1}{2} - i}{x} + (-) (0) \\ &= \frac{\frac{1}{2} - i}{x} \\ &= \frac{\frac{1}{2} - i}{x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$(0) + 2\left(\frac{\frac{1}{2} - i}{x}\right)(0) + \left(\left(\frac{-\frac{1}{2} + i}{x^2}\right) + \left(\frac{\frac{1}{2} - i}{x}\right)^2 - \left(-\frac{5}{4x^2}\right)\right) = 0$$

$$0 = 0$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$z_1(x) = pe^{\int \omega dx}$$

$$= e^{\int \frac{\frac{1}{2} - i}{x} dx}$$

$$= x^{\frac{1}{2} - i}$$

The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

$$= z_1 e^{-\int \frac{1}{2} \frac{-3x}{x^2} dx}$$

$$= z_1 e^{\frac{3 \ln(x)}{2}}$$

$$= z_1 \left(x^{\frac{3}{2}}\right)$$

Which simplifies to

$$y_1 = x^{2-i}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$y_2 = y_1 \int \frac{e^{\int -\frac{-3x}{x^2} dx}}{(y_1)^2} dx$$

$$= y_1 \int \frac{e^{3 \ln(x)}}{(y_1)^2} dx$$

$$= y_1 \left(-\frac{ix^{2i}}{2}\right)$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (x^{2-i}) + c_2 \left(x^{2-i} \left(-\frac{i x^{2i}}{2} \right) \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 x^{2-i} - \frac{i c_2 x^{2+i}}{2} \quad (1)$$

Verification of solutions

$$y = c_1 x^{2-i} - \frac{i c_2 x^{2+i}}{2}$$

Verified OK.

16.7.6 Maple step by step solution

Let's solve

$$x^2 y'' - 3xy' + 5y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = \frac{3y'}{x} - \frac{5y}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - \frac{3y'}{x} + \frac{5y}{x^2} = 0$$

- Multiply by denominators of the ODE

$$x^2 y'' - 3xy' + 5y = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x , using the chain rule

$$y' = \left(\frac{d}{dt} y(t) \right) t'(x)$$

- Compute derivative

$$y' = \frac{d}{dt} y(t)$$

- Calculate the 2nd derivative of y with respect to x , using the chain rule

$$y'' = \left(\frac{d^2}{dt^2} y(t) \right) t'(x)^2 + t''(x) \left(\frac{d}{dt} y(t) \right)$$

- Compute derivative

$$y'' = \frac{\frac{d^2}{dt^2} y(t)}{x^2} - \frac{\frac{d}{dt} y(t)}{x^2}$$

Substitute the change of variables back into the ODE

$$x^2 \left(\frac{\frac{d^2}{dt^2} y(t)}{x^2} - \frac{\frac{d}{dt} y(t)}{x^2} \right) - 3 \frac{d}{dt} y(t) + 5y(t) = 0$$

- Simplify

$$\frac{d^2}{dt^2} y(t) - 4 \frac{d}{dt} y(t) + 5y(t) = 0$$

- Characteristic polynomial of ODE

$$r^2 - 4r + 5 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{4 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (2 - I, 2 + I)$$

- 1st solution of the ODE

$$y_1(t) = e^{2t} \cos(t)$$

- 2nd solution of the ODE

$$y_2(t) = e^{2t} \sin(t)$$

- General solution of the ODE

$$y(t) = c_1 y_1(t) + c_2 y_2(t)$$

- Substitute in solutions

$$y(t) = c_1 e^{2t} \cos(t) + c_2 e^{2t} \sin(t)$$

- Change variables back using $t = \ln(x)$

$$y = c_1 x^2 \cos(\ln(x)) + c_2 x^2 \sin(\ln(x))$$

- Simplify

$$y = x^2 (c_1 \cos(\ln(x)) + c_2 \sin(\ln(x)))$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
<- LODE of Euler type successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(x^2*diff(y(x),x$2)-3*x*diff(y(x),x)+5*y(x)=0,y(x), singsol=all)
```

$$y(x) = x^2(c_1 \sin(\ln(x)) + c_2 \cos(\ln(x)))$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 22

```
DSolve[x^2*y''[x]-3*x*y'[x]+5*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^2(c_2 \cos(\log(x)) + c_1 \sin(\log(x)))$$

16.8 problem 2(c)

16.8.1 Solving as second order euler ode ode	1289
16.8.2 Solving as second order change of variable on x method 2 ode .	1290
16.8.3 Solving as second order change of variable on y method 2 ode .	1293
16.8.4 Solving using Kovacic algorithm	1296
16.8.5 Maple step by step solution	1301

Internal problem ID [6038]

Internal file name [OUTPUT/5286_Sunday_June_05_2022_03_29_30_PM_70934068/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 149

Problem number: 2(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_euler_ode", "second_order_change_of_variable_on_x_method_2", "second_order_change_of_variable_on_y_method_2"

Maple gives the following as the ode type

```
[[_Emden, _Fowler]]
```

$$x^2y'' + (-2 - i)xy' + 3iy = 0$$

16.8.1 Solving as second order euler ode ode

This is Euler second order ODE. Let the solution be $y = x^r$, then $y' = rx^{r-1}$ and $y'' = r(r-1)x^{r-2}$. Substituting these back into the given ODE gives

$$x^2(r(r-1))x^{r-2}(-2-i)rx^{r-1} + 3ix^r = 0$$

Simplifying gives

$$r(r-1)x^r(-2-i)rx^r + 3ix^r = 0$$

Since $x^r \neq 0$ then dividing throughout by x^r gives

$$r(r-1)(-2-i)r + 3i = 0$$

Or

$$3i + r^2 + (-3 - i)r = 0 \quad (1)$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$\begin{aligned} r_1 &= 3 \\ r_2 &= i \end{aligned}$$

Since the roots are real and distinct, then the general solution is

$$y = c_1 y_1 + c_2 y_2$$

Where $y_1 = x^{r_1}$ and $y_2 = x^{r_2}$. Hence

$$y = c_1 x^3 + c_2 x^i$$

Summary

The solution(s) found are the following

$$y = c_1 x^3 + c_2 x^i \quad (1)$$

Verification of solutions

$$y = c_1 x^3 + c_2 x^i$$

Verified OK.

16.8.2 Solving as second order change of variable on x method 2 ode

In normal form the ode

$$x^2 y'' + (-2 - i)xy' + 3iy = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$\begin{aligned} p(x) &= \frac{-2 - i}{x} \\ q(x) &= \frac{3i}{x^2} \end{aligned}$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x)\tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned} \tau &= \int e^{-(\int p(x)dx)} dx \\ &= \int e^{-\left(\int \frac{-2-i}{x} dx\right)} dx \\ &= \int e^{(2+i)\ln(x)} dx \\ &= \int x^{2+i} dx \\ &= \left(\frac{3}{10} - \frac{i}{10}\right) x^{3+i} \end{aligned} \quad (6)$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned} q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\ &= \frac{\frac{3i}{x^2}}{x^{4+2i}} \\ &= 3ix^{-6-2i} \end{aligned} \quad (7)$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned} \frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) + 3ix^{-6-2i}y(\tau) &= 0 \end{aligned}$$

But in terms of τ

$$3ix^{-6-2i} = \frac{9}{50} + \frac{6i}{25} \tau^2$$

Hence the above ode becomes

$$\frac{d^2}{d\tau^2}y(\tau) + \frac{\left(\frac{9}{50} + \frac{6i}{25}\right)y(\tau)}{\tau^2} = 0$$

The above ode is now solved for $y(\tau)$. The ode can be written as

$$50\left(\frac{d^2}{d\tau^2}y(\tau)\right)\tau^2 + (9 + 12i)y(\tau) = 0$$

Which shows it is a Euler ODE. This is Euler second order ODE. Let the solution be $y(\tau) = \tau^r$, then $y' = r\tau^{r-1}$ and $y'' = r(r-1)\tau^{r-2}$. Substituting these back into the given ODE gives

$$50\tau^2(r(r-1))\tau^{r-2} + 0r\tau^{r-1} + (9 + 12i)\tau^r = 0$$

Simplifying gives

$$50r(r-1)\tau^r + 0\tau^r + (9 + 12i)\tau^r = 0$$

Since $\tau^r \neq 0$ then dividing throughout by τ^r gives

$$50r(r-1) + 0 + 9 + 12i = 0$$

Or

$$50r^2 - 50r + 12i + 9 = 0 \tag{1}$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$\begin{aligned} r_1 &= \frac{1}{10} + \frac{3i}{10} \\ r_2 &= \frac{9}{10} - \frac{3i}{10} \end{aligned}$$

The roots are complex conjugate of each others. Let the roots be

$$\begin{aligned} r_1 &= \alpha + i\beta \\ r_2 &= \alpha - i\beta \end{aligned}$$

Where in this case $\alpha = \frac{1}{10}$ and $\beta = \frac{3}{10}$. Hence the solution becomes

$$\begin{aligned} y(\tau) &= c_1\tau^{r_1} + c_2\tau^{r_2} \\ &= c_1\tau^{\alpha+i\beta} + c_2\tau^{\alpha-i\beta} \\ &= \tau^\alpha(c_1\tau^{i\beta} + c_2\tau^{-i\beta}) \\ &= \tau^\alpha\left(c_1e^{\ln(\tau^{i\beta})} + c_2e^{\ln(\tau^{-i\beta})}\right) \\ &= \tau^\alpha\left(c_1e^{i(\beta \ln \tau)} + c_2e^{-i(\beta \ln \tau)}\right) \end{aligned}$$

Using the values for $\alpha = \frac{1}{10}, \beta = \frac{3}{10}$, the above becomes

$$y(\tau) = \tau^{\frac{1}{10}} \left(c_1 e^{\frac{3i \ln(\tau)}{10}} + c_2 e^{-\frac{3i \ln(\tau)}{10}} \right)$$

Using Euler relation, the expression $c_1 e^{iA} + c_2 e^{-iA}$ is transformed to $c_1 \cos A + c_1 \sin A$ where the constants are free to change. Applying this to the above result gives

$$y(\tau) = \tau^{\frac{1}{10}} \left(c_1 \cos \left(\frac{3 \ln(\tau)}{10} \right) + c_2 \sin \left(\frac{3 \ln(\tau)}{10} \right) \right)$$

The above solution is now transformed back to y using (6) which results in

$$y = \frac{10^{\frac{4}{5}} ((30 - 10i) x^{3+i})^{\frac{1}{10}} \left(c_1 \cos \left(\frac{3 \ln \left(\left(\frac{3}{10} - \frac{i}{10} \right) x^{3+i} \right)}{10} \right) + c_2 \sin \left(\frac{3 \ln \left(\left(\frac{3}{10} - \frac{i}{10} \right) x^{3+i} \right)}{10} \right) \right)}{10}$$

Summary

The solution(s) found are the following

$$y = \frac{10^{\frac{4}{5}} ((30 - 10i) x^{3+i})^{\frac{1}{10}} \left(c_1 \cos \left(\frac{3 \ln \left(\left(\frac{3}{10} - \frac{i}{10} \right) x^{3+i} \right)}{10} \right) + c_2 \sin \left(\frac{3 \ln \left(\left(\frac{3}{10} - \frac{i}{10} \right) x^{3+i} \right)}{10} \right) \right)}{10} \quad (1)$$

Verification of solutions

$$y = \frac{10^{\frac{4}{5}} ((30 - 10i) x^{3+i})^{\frac{1}{10}} \left(c_1 \cos \left(\frac{3 \ln \left(\left(\frac{3}{10} - \frac{i}{10} \right) x^{3+i} \right)}{10} \right) + c_2 \sin \left(\frac{3 \ln \left(\left(\frac{3}{10} - \frac{i}{10} \right) x^{3+i} \right)}{10} \right) \right)}{10}$$

Verified OK.

16.8.3 Solving as second order change of variable on y method 2 ode

In normal form the ode

$$x^2 y'' + (-2 - i) xy' + 3iy = 0 \quad (1)$$

Becomes

$$y'' + p(x) y' + q(x) y = 0 \quad (2)$$

Where

$$p(x) = \frac{-2 - i}{x}$$

$$q(x) = \frac{3i}{x^2}$$

Applying change of variables on the dependent variable $y = v(x)x^n$ to (2) gives the following ode where the dependent variables is $v(x)$ and not y .

$$v''(x) + \left(\frac{2n}{x} + p\right)v'(x) + \left(\frac{n(n-1)}{x^2} + \frac{np}{x} + q\right)v(x) = 0 \quad (3)$$

Let the coefficient of $v(x)$ above be zero. Hence

$$\frac{n(n-1)}{x^2} + \frac{np}{x} + q = 0 \quad (4)$$

Substituting the earlier values found for $p(x)$ and $q(x)$ into (4) gives

$$\frac{n(n-1)}{x^2} + \frac{(-2-i)n}{x^2} + \frac{3i}{x^2} = 0 \quad (5)$$

Solving (5) for n gives

$$n = 3 \quad (6)$$

Substituting this value in (3) gives

$$\begin{aligned} v''(x) + \left(\frac{6}{x} + \frac{-2-i}{x}\right)v'(x) &= 0 \\ v''(x) + \frac{(4-i)v'(x)}{x} &= 0 \end{aligned} \quad (7)$$

Using the substitution

$$u(x) = v'(x)$$

Then (7) becomes

$$u'(x) + \frac{(4-i)u(x)}{x} = 0 \quad (8)$$

The above is now solved for $u(x)$. In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= \frac{(-4+i)u}{x} \end{aligned}$$

Where $f(x) = \frac{-4+i}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u} du &= \frac{-4+i}{x} dx \\ \int \frac{1}{u} du &= \int \frac{-4+i}{x} dx \\ \ln(u) &= (-4+i) \ln(x) + c_1 \\ u &= e^{(-4+i) \ln(x) + c_1} \\ &= c_1 e^{(-4+i) \ln(x)}\end{aligned}$$

Which simplifies to

$$u(x) = \frac{c_1 x^i}{x^4}$$

Now that $u(x)$ is known, then

$$\begin{aligned}v'(x) &= u(x) \\ v(x) &= \int u(x) dx + c_2 \\ &= \left(-\frac{3}{10} - \frac{i}{10}\right) c_1 x^{-3+i} + c_2\end{aligned}$$

Hence

$$\begin{aligned}y &= v(x) x^n \\ &= \left(\left(-\frac{3}{10} - \frac{i}{10}\right) c_1 x^{-3+i} + c_2\right) x^3 \\ &= \frac{x^3((3+i) c_1 x^{-3+i} - 10c_2)}{10}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \left(\left(-\frac{3}{10} - \frac{i}{10}\right) c_1 x^{-3+i} + c_2\right) x^3 \quad (1)$$

Verification of solutions

$$y = \left(\left(-\frac{3}{10} - \frac{i}{10}\right) c_1 x^{-3+i} + c_2\right) x^3$$

Verified OK.

16.8.4 Solving using Kovacic algorithm

Writing the ode as

$$x^2 y'' + (-2 - i)xy' + 3iy = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= x^2 \\ B &= (-2 - i)x \\ C &= 3i \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{7 - 6i}{4x^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 7 - 6i \\ t &= 4x^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{\frac{7}{4} - \frac{3i}{2}}{x^2} \right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 204: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = \frac{\frac{7}{4} - \frac{3i}{2}}{x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = \frac{7}{4} - \frac{3i}{2}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = 2 - \frac{i}{2} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = -1 + \frac{i}{2} \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = \frac{\frac{7}{4} - \frac{3i}{2}}{x^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = \frac{7}{4} - \frac{3i}{2}$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = 2 - \frac{i}{2} \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = -1 + \frac{i}{2} \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{\frac{7}{4} - \frac{3i}{2}}{x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	$2 - \frac{i}{2}$	$-1 + \frac{i}{2}$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	$2 - \frac{i}{2}$	$-1 + \frac{i}{2}$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω .

Trying $\alpha_{\infty}^{-} = -1 + \frac{i}{2}$ then

$$\begin{aligned} d &= \alpha_{\infty}^{-} - (\alpha_{c_1}^{-}) \\ &= -1 + \frac{i}{2} - \left(-1 + \frac{i}{2}\right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_{\infty}$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^{-}}{x - c_1} \right) + (-)[\sqrt{r}]_{\infty} \\ &= \frac{-1 + \frac{i}{2}}{x} + (-)(0) \\ &= \frac{-1 + \frac{i}{2}}{x} \\ &= \frac{-1 + \frac{i}{2}}{x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r)p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2\left(\frac{-1 + \frac{i}{2}}{x}\right)(0) + \left(\left(\frac{1 - \frac{i}{2}}{x^2}\right) + \left(\frac{-1 + \frac{i}{2}}{x}\right)^2 - \left(\frac{7}{4} - \frac{3i}{2}\right)\right) &= 0 \\ 0 &= 0 \end{aligned}$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= pe^{\int \omega dx} \\ &= e^{\int \frac{-1 + \frac{i}{2}}{x} dx} \\ &= x^{-1 + \frac{i}{2}} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\&= z_1 e^{-\int \frac{1}{2} \frac{(-2-i)x}{x^2} dx} \\&= z_1 e^{(1+\frac{i}{2}) \ln(x)} \\&= z_1 \left(x^{1+\frac{i}{2}} \right)\end{aligned}$$

Which simplifies to

$$y_1 = x^i$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{(-2-i)x}{x^2} dx}}{(y_1)^2} dx \\&= y_1 \int \frac{e^{(2+i) \ln(x)}}{(y_1)^2} dx \\&= y_1 \left(\left(\frac{3}{10} + \frac{i}{10} \right) x^{3-i} \right)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\&= c_1 (x^i) + c_2 \left(x^i \left(\left(\frac{3}{10} + \frac{i}{10} \right) x^{3-i} \right) \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 x^i + \left(\frac{3}{10} + \frac{i}{10} \right) c_2 x^3 \quad (1)$$

Verification of solutions

$$y = c_1 x^i + \left(\frac{3}{10} + \frac{i}{10} \right) c_2 x^3$$

Verified OK.

16.8.5 Maple step by step solution

Let's solve

$$x^2 y'' - (2 + I) xy' + 3 Iy = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{3Iy}{x^2} + \frac{(2+I)y'}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - \frac{(2+I)y'}{x} + \frac{3Iy}{x^2} = 0$$

- Multiply by denominators of the ODE

$$x^2 y'' - (2 + I) xy' + 3 Iy = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x , using the chain rule

$$y' = \left(\frac{d}{dt}y(t)\right) t'(x)$$

- Compute derivative

$$y' = \frac{\frac{d}{dt}y(t)}{x}$$

- Calculate the 2nd derivative of y with respect to x , using the chain rule

$$y'' = \left(\frac{d^2}{dt^2}y(t)\right) t'(x)^2 + t''(x) \left(\frac{d}{dt}y(t)\right)$$

- Compute derivative

$$y'' = \frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2}$$

Substitute the change of variables back into the ODE

$$x^2 \left(\frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2}\right) - (2 + I) \left(\frac{d}{dt}y(t)\right) + 3 Iy(t) = 0$$

- Simplify

$$\frac{d^2}{dt^2}y(t) - (3 + I) \left(\frac{d}{dt}y(t)\right) + 3 Iy(t) = 0$$

- Characteristic polynomial of ODE

$$r^2 - (3 + I)r + 3 I = 0$$

- Factor the characteristic polynomial
 $-(r - 3)(-r + 1) = 0$
- Roots of the characteristic polynomial
 $r = (3, 1)$
- 1st solution of the ODE
 $y_1(t) = e^{3t}$
- 2nd solution of the ODE
 $y_2(t) = 0$
- General solution of the ODE
 $y(t) = c_1 y_1(t) + c_2 y_2(t)$
- Substitute in solutions
 $y(t) = c_1 e^{3t}$
- Change variables back using $t = \ln(x)$
 $y = c_1 x^3$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
<- LODE of Euler type successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve(x^2*diff(y(x),x$2)-(2+1)*x*diff(y(x),x)+3*1*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 x^3 + c_2 x^1$$

✓ Solution by Mathematica

Time used: 0.041 (sec). Leaf size: 20

```
DSolve[x^2*y'[x]-(2+I)*x*y'[x]+3*I*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 x^i + c_2 x^3$$

16.9 problem 2(d)

16.9.1 Solving as second order euler ode ode	1304
16.9.2 Solving as second order change of variable on x method 2 ode .	1308
16.9.3 Solving as second order change of variable on x method 1 ode .	1313
16.9.4 Solving as second order change of variable on y method 2 ode .	1318
16.9.5 Solving using Kovacic algorithm	1323

Internal problem ID [6039]

Internal file name [OUTPUT/5287_Sunday_June_05_2022_03_29_32_PM_10541990/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 149

Problem number: 2(d).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_euler_ode", "second_order_change_of_variable_on_x_method_1", "second_order_change_of_variable_on_x_method_2", "second_order_change_of_variable_on_y_method_2"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$x^2y'' + xy' - 4\pi y = x$$

16.9.1 Solving as second order euler ode ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = x^2, B = x, C = -4\pi, f(x) = x$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$x^2y'' + xy' - 4\pi y = 0$$

This is Euler second order ODE. Let the solution be $y = x^r$, then $y' = rx^{r-1}$ and $y'' = r(r-1)x^{r-2}$. Substituting these back into the given ODE gives

$$x^2(r(r-1))x^{r-2} + xrx^{r-1} - 4\pi x^r = 0$$

Simplifying gives

$$r(r-1)x^r + rx^r - 4\pi x^r = 0$$

Since $x^r \neq 0$ then dividing throughout by x^r gives

$$r(r-1) + r - 4\pi = 0$$

Or

$$r^2 - 4\pi = 0 \tag{1}$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$\begin{aligned} r_1 &= -2\sqrt{\pi} \\ r_2 &= 2\sqrt{\pi} \end{aligned}$$

Since the roots are real and distinct, then the general solution is

$$y = c_1y_1 + c_2y_2$$

Where $y_1 = x^{r_1}$ and $y_2 = x^{r_2}$. Hence

$$y = c_1x^{-2\sqrt{\pi}} + c_2x^{2\sqrt{\pi}}$$

Next, we find the particular solution to the ODE

$$x^2y'' + xy' - 4\pi y = x$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1y_1 + u_2y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned} y_1 &= x^{-2\sqrt{\pi}} \\ y_2 &= x^{2\sqrt{\pi}} \end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} x^{-2\sqrt{\pi}} & x^{2\sqrt{\pi}} \\ \frac{d}{dx}(x^{-2\sqrt{\pi}}) & \frac{d}{dx}(x^{2\sqrt{\pi}}) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} x^{-2\sqrt{\pi}} & x^{2\sqrt{\pi}} \\ -\frac{2x^{-2\sqrt{\pi}}\sqrt{\pi}}{x} & \frac{2x^{2\sqrt{\pi}}\sqrt{\pi}}{x} \end{vmatrix}$$

Therefore

$$W = (x^{-2\sqrt{\pi}}) \left(\frac{2x^{2\sqrt{\pi}}\sqrt{\pi}}{x} \right) - (x^{2\sqrt{\pi}}) \left(-\frac{2x^{-2\sqrt{\pi}}\sqrt{\pi}}{x} \right)$$

Which simplifies to

$$W = \frac{4x^{-2\sqrt{\pi}}x^{2\sqrt{\pi}}\sqrt{\pi}}{x}$$

Which simplifies to

$$W = \frac{4\sqrt{\pi}}{x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x x^{2\sqrt{\pi}}}{4\sqrt{\pi} x} dx$$

Which simplifies to

$$u_1 = - \int \frac{x^{2\sqrt{\pi}}}{4\sqrt{\pi}} dx$$

Hence

$$u_1 = -\frac{x^{1+2\sqrt{\pi}}}{4\sqrt{\pi} (1 + 2\sqrt{\pi})}$$

And Eq. (3) becomes

$$u_2 = \int \frac{x^{-2\sqrt{\pi}} x}{4\sqrt{\pi} x} dx$$

Which simplifies to

$$u_2 = \int \frac{x^{-2\sqrt{\pi}}}{4\sqrt{\pi}} dx$$

Hence

$$u_2 = -\frac{x^{-2\sqrt{\pi}+1}}{4\sqrt{\pi} (2\sqrt{\pi} - 1)}$$

Which simplifies to

$$u_1 = -\frac{x^{1+2\sqrt{\pi}}}{\sqrt{\pi} (4 + 8\sqrt{\pi})}$$
$$u_2 = -\frac{x^{-2\sqrt{\pi}+1}}{\sqrt{\pi} (8\sqrt{\pi} - 4)}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{x^{1+2\sqrt{\pi}} x^{-2\sqrt{\pi}}}{\sqrt{\pi} (4 + 8\sqrt{\pi})} - \frac{x^{-2\sqrt{\pi}+1} x^{2\sqrt{\pi}}}{\sqrt{\pi} (8\sqrt{\pi} - 4)}$$

Which simplifies to

$$y_p(x) = -\frac{x}{4\pi - 1}$$

Therefore the general solution is

$$y = y_h + y_p$$
$$= -\frac{x}{4\pi - 1} + c_1 x^{-2\sqrt{\pi}} + c_2 x^{2\sqrt{\pi}}$$

Summary

The solution(s) found are the following

$$y = -\frac{x}{4\pi - 1} + c_1x^{-2\sqrt{\pi}} + c_2x^{2\sqrt{\pi}} \quad (1)$$

Verification of solutions

$$y = -\frac{x}{4\pi - 1} + c_1x^{-2\sqrt{\pi}} + c_2x^{2\sqrt{\pi}}$$

Verified OK.

16.9.2 Solving as second order change of variable on x method 2 ode

This is second order non-homogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$.
 y_h is the solution to

$$x^2y'' + xy' - 4\pi y = 0$$

In normal form the ode

$$x^2y'' + xy' - 4\pi y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = -\frac{4\pi}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x) \tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned} \tau &= \int e^{-(\int p(x)dx)} dx \\ &= \int e^{-(\int \frac{1}{x} dx)} dx \\ &= \int e^{-\ln(x)} dx \\ &= \int \frac{1}{x} dx \\ &= \ln(x) \end{aligned} \tag{6}$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned} q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\ &= \frac{\frac{4\pi}{x^2}}{\frac{1}{x^2}} \\ &= -4\pi \end{aligned} \tag{7}$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned} \frac{d^2}{d\tau^2} y(\tau) + q_1 y(\tau) &= 0 \\ \frac{d^2}{d\tau^2} y(\tau) - 4\pi y(\tau) &= 0 \end{aligned}$$

The above ode is now solved for $y(\tau)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(\tau) + By'(\tau) + Cy(\tau) = 0$$

Where in the above $A = 1, B = 0, C = -4\pi$. Let the solution be $y(\tau) = e^{\lambda\tau}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda\tau} - 4\pi e^{\lambda\tau} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda\tau}$ gives

$$\lambda^2 - 4\pi = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = -4\pi$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(-4\pi)} \\ &= \pm 2\sqrt{\pi} \end{aligned}$$

Hence

$$\lambda_1 = +2\sqrt{\pi}$$

$$\lambda_2 = -2\sqrt{\pi}$$

Which simplifies to

$$\lambda_1 = 2\sqrt{\pi}$$

$$\lambda_2 = -2\sqrt{\pi}$$

Since roots are real and distinct, then the solution is

$$y(\tau) = c_1 e^{\lambda_1 \tau} + c_2 e^{\lambda_2 \tau}$$

$$y(\tau) = c_1 e^{(2\sqrt{\pi})\tau} + c_2 e^{(-2\sqrt{\pi})\tau}$$

Or

$$y(\tau) = c_1 e^{2\sqrt{\pi}\tau} + c_2 e^{-2\sqrt{\pi}\tau}$$

The above solution is now transformed back to y using (6) which results in

$$y = c_1 x^{2\sqrt{\pi}} + c_2 x^{-2\sqrt{\pi}}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 x^{2\sqrt{\pi}} + c_2 x^{-2\sqrt{\pi}}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of

parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = x^{-2\sqrt{\pi}}$$

$$y_2 = x^{2\sqrt{\pi}}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} x^{-2\sqrt{\pi}} & x^{2\sqrt{\pi}} \\ \frac{d}{dx}(x^{-2\sqrt{\pi}}) & \frac{d}{dx}(x^{2\sqrt{\pi}}) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} x^{-2\sqrt{\pi}} & x^{2\sqrt{\pi}} \\ -\frac{2x^{-2\sqrt{\pi}}\sqrt{\pi}}{x} & \frac{2x^{2\sqrt{\pi}}\sqrt{\pi}}{x} \end{vmatrix}$$

Therefore

$$W = (x^{-2\sqrt{\pi}}) \left(\frac{2x^{2\sqrt{\pi}}\sqrt{\pi}}{x} \right) - (x^{2\sqrt{\pi}}) \left(-\frac{2x^{-2\sqrt{\pi}}\sqrt{\pi}}{x} \right)$$

Which simplifies to

$$W = \frac{4\sqrt{\pi}}{x}$$

Which simplifies to

$$W = \frac{4\sqrt{\pi}}{x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x x^{2\sqrt{\pi}}}{4\sqrt{\pi} x} dx$$

Which simplifies to

$$u_1 = - \int \frac{x^{2\sqrt{\pi}}}{4\sqrt{\pi}} dx$$

Hence

$$u_1 = - \frac{x^{1+2\sqrt{\pi}}}{4\sqrt{\pi} (1 + 2\sqrt{\pi})}$$

And Eq. (3) becomes

$$u_2 = \int \frac{x^{-2\sqrt{\pi}} x}{4\sqrt{\pi} x} dx$$

Which simplifies to

$$u_2 = \int \frac{x^{-2\sqrt{\pi}}}{4\sqrt{\pi}} dx$$

Hence

$$u_2 = - \frac{x^{-2\sqrt{\pi}+1}}{4\sqrt{\pi} (2\sqrt{\pi} - 1)}$$

Which simplifies to

$$u_1 = - \frac{x^{1+2\sqrt{\pi}}}{\sqrt{\pi} (4 + 8\sqrt{\pi})}$$

$$u_2 = - \frac{x^{-2\sqrt{\pi}+1}}{\sqrt{\pi} (8\sqrt{\pi} - 4)}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = - \frac{x^{1+2\sqrt{\pi}} x^{-2\sqrt{\pi}}}{\sqrt{\pi} (4 + 8\sqrt{\pi})} - \frac{x^{-2\sqrt{\pi}+1} x^{2\sqrt{\pi}}}{\sqrt{\pi} (8\sqrt{\pi} - 4)}$$

Which simplifies to

$$y_p(x) = -\frac{x}{4\pi - 1}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 x^{2\sqrt{\pi}} + c_2 x^{-2\sqrt{\pi}} \right) + \left(-\frac{x}{4\pi - 1} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 x^{2\sqrt{\pi}} + c_2 x^{-2\sqrt{\pi}} - \frac{x}{4\pi - 1} \quad (1)$$

Verification of solutions

$$y = c_1 x^{2\sqrt{\pi}} + c_2 x^{-2\sqrt{\pi}} - \frac{x}{4\pi - 1}$$

Verified OK.

16.9.3 Solving as second order change of variable on x method 1 ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = x^2$, $B = x$, $C = -4\pi$, $f(x) = x$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$x^2 y'' + xy' - 4\pi y = 0$$

In normal form the ode

$$x^2 y'' + xy' - 4\pi y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{1}{x}$$

$$q(x) = -\frac{4\pi}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) results

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $q_1 = c^2$ where c is some constant. Therefore from (5)

$$\tau' = \frac{1}{c}\sqrt{q}$$

$$= \frac{2\sqrt{-\frac{\pi}{x^2}}}{c} \quad (6)$$

$$\tau'' = \frac{2\pi}{c\sqrt{-\frac{\pi}{x^2}}x^3}$$

Substituting the above into (4) results in

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2}$$

$$= \frac{\frac{2\pi}{c\sqrt{-\frac{\pi}{x^2}}x^3} + \frac{1}{x}\frac{2\sqrt{-\frac{\pi}{x^2}}}{c}}{\left(\frac{2\sqrt{-\frac{\pi}{x^2}}}{c}\right)^2}$$

$$= 0$$

Therefore ode (3) now becomes

$$y(\tau)'' + p_1y(\tau)' + q_1y(\tau) = 0$$

$$\frac{d^2}{d\tau^2}y(\tau) + c^2y(\tau) = 0 \quad (7)$$

The above ode is now solved for $y(\tau)$. Since the ode is now constant coefficients, it can be easily solved to give

$$y(\tau) = c_1 \cos(c\tau) + c_2 \sin(c\tau)$$

Now from (6)

$$\begin{aligned} \tau &= \int \frac{1}{c} \sqrt{q} dx \\ &= \frac{\int 2\sqrt{-\frac{\pi}{x^2}} dx}{c} \\ &= \frac{2\sqrt{-\frac{\pi}{x^2}} x \ln(x)}{c} \end{aligned}$$

Substituting the above into the solution obtained gives

$$y = c_1 \cosh(2\sqrt{\pi} \ln(x)) + ic_2 \sinh(2\sqrt{\pi} \ln(x))$$

Now the particular solution to this ODE is found

$$x^2 y'' + xy' - 4\pi y = x$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned} y_1 &= x^{-2\sqrt{\pi}} \\ y_2 &= x^{2\sqrt{\pi}} \end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} x^{-2\sqrt{\pi}} & x^{2\sqrt{\pi}} \\ \frac{d}{dx}(x^{-2\sqrt{\pi}}) & \frac{d}{dx}(x^{2\sqrt{\pi}}) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} x^{-2\sqrt{\pi}} & x^{2\sqrt{\pi}} \\ -\frac{2x^{-2\sqrt{\pi}}\sqrt{\pi}}{x} & \frac{2x^{2\sqrt{\pi}}\sqrt{\pi}}{x} \end{vmatrix}$$

Therefore

$$W = (x^{-2\sqrt{\pi}}) \left(\frac{2x^{2\sqrt{\pi}}\sqrt{\pi}}{x} \right) - (x^{2\sqrt{\pi}}) \left(-\frac{2x^{-2\sqrt{\pi}}\sqrt{\pi}}{x} \right)$$

Which simplifies to

$$W = \frac{4\sqrt{\pi}}{x}$$

Which simplifies to

$$W = \frac{4\sqrt{\pi}}{x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x x^{2\sqrt{\pi}}}{4\sqrt{\pi} x} dx$$

Which simplifies to

$$u_1 = - \int \frac{x^{2\sqrt{\pi}}}{4\sqrt{\pi}} dx$$

Hence

$$u_1 = - \frac{x^{1+2\sqrt{\pi}}}{4\sqrt{\pi} (1 + 2\sqrt{\pi})}$$

And Eq. (3) becomes

$$u_2 = \int \frac{x^{-2\sqrt{\pi}} x}{4\sqrt{\pi} x} dx$$

Which simplifies to

$$u_2 = \int \frac{x^{-2\sqrt{\pi}}}{4\sqrt{\pi}} dx$$

Hence

$$u_2 = -\frac{x^{-2\sqrt{\pi}+1}}{4\sqrt{\pi} (2\sqrt{\pi} - 1)}$$

Which simplifies to

$$u_1 = -\frac{x^{1+2\sqrt{\pi}}}{\sqrt{\pi} (4 + 8\sqrt{\pi})}$$
$$u_2 = -\frac{x^{-2\sqrt{\pi}+1}}{\sqrt{\pi} (8\sqrt{\pi} - 4)}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{x^{1+2\sqrt{\pi}}x^{-2\sqrt{\pi}}}{\sqrt{\pi} (4 + 8\sqrt{\pi})} - \frac{x^{-2\sqrt{\pi}+1}x^{2\sqrt{\pi}}}{\sqrt{\pi} (8\sqrt{\pi} - 4)}$$

Which simplifies to

$$y_p(x) = -\frac{x}{4\pi - 1}$$

Therefore the general solution is

$$y = y_h + y_p$$
$$= (c_1 \cosh (2\sqrt{\pi} \ln (x)) + ic_2 \sinh (2\sqrt{\pi} \ln (x))) + \left(-\frac{x}{4\pi - 1} \right)$$
$$= -\frac{x}{4\pi - 1} + c_1 \cosh (2\sqrt{\pi} \ln (x)) + ic_2 \sinh (2\sqrt{\pi} \ln (x))$$

Which simplifies to

$$y = -\frac{x}{4\pi - 1} + c_1 \cosh (2\sqrt{\pi} \ln (x)) + ic_2 \sinh (2\sqrt{\pi} \ln (x))$$

Summary

The solution(s) found are the following

$$y = -\frac{x}{4\pi - 1} + c_1 \cosh (2\sqrt{\pi} \ln (x)) + ic_2 \sinh (2\sqrt{\pi} \ln (x)) \quad (1)$$

Verification of solutions

$$y = -\frac{x}{4\pi - 1} + c_1 \cosh(2\sqrt{\pi} \ln(x)) + ic_2 \sinh(2\sqrt{\pi} \ln(x))$$

Verified OK.

16.9.4 Solving as second order change of variable on y method 2 ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = x^2$, $B = x$, $C = -4\pi$, $f(x) = x$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$x^2y'' + xy' - 4\pi y = 0$$

In normal form the ode

$$x^2y'' + xy' - 4\pi y = 0 \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \tag{2}$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = -\frac{4\pi}{x^2}$$

Applying change of variables on the dependent variable $y = v(x)x^n$ to (2) gives the following ode where the dependent variables is $v(x)$ and not y .

$$v''(x) + \left(\frac{2n}{x} + p\right)v'(x) + \left(\frac{n(n-1)}{x^2} + \frac{np}{x} + q\right)v(x) = 0 \tag{3}$$

Let the coefficient of $v(x)$ above be zero. Hence

$$\frac{n(n-1)}{x^2} + \frac{np}{x} + q = 0 \tag{4}$$

Substituting the earlier values found for $p(x)$ and $q(x)$ into (4) gives

$$\frac{n(n-1)}{x^2} + \frac{n}{x^2} - \frac{4\pi}{x^2} = 0 \quad (5)$$

Solving (5) for n gives

$$n = 2\sqrt{\pi} \quad (6)$$

Substituting this value in (3) gives

$$\begin{aligned} v''(x) + \left(\frac{4\sqrt{\pi}}{x} + \frac{1}{x} \right) v'(x) &= 0 \\ v''(x) + \frac{(4\sqrt{\pi} + 1) v'(x)}{x} &= 0 \end{aligned} \quad (7)$$

Using the substitution

$$u(x) = v'(x)$$

Then (7) becomes

$$u'(x) + \frac{(4\sqrt{\pi} + 1) u(x)}{x} = 0 \quad (8)$$

The above is now solved for $u(x)$. In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= \frac{(-4\sqrt{\pi} - 1) u}{x} \end{aligned}$$

Where $f(x) = \frac{-4\sqrt{\pi}-1}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned} \frac{1}{u} du &= \frac{-4\sqrt{\pi} - 1}{x} dx \\ \int \frac{1}{u} du &= \int \frac{-4\sqrt{\pi} - 1}{x} dx \\ \ln(u) &= (-4\sqrt{\pi} - 1) \ln(x) + c_1 \\ u &= e^{(-4\sqrt{\pi}-1) \ln(x)+c_1} \\ &= c_1 e^{(-4\sqrt{\pi}-1) \ln(x)} \end{aligned}$$

Which simplifies to

$$u(x) = \frac{c_1 x^{-4\sqrt{\pi}}}{x}$$

Now that $u(x)$ is known, then

$$\begin{aligned}v'(x) &= u(x) \\v(x) &= \int u(x) dx + c_2 \\&= -\frac{c_1 x^{-4\sqrt{\pi}}}{4\sqrt{\pi}} + c_2\end{aligned}$$

Hence

$$\begin{aligned}y &= v(x) x^n \\&= \left(-\frac{c_1 x^{-4\sqrt{\pi}}}{4\sqrt{\pi}} + c_2 \right) x^{2\sqrt{\pi}} \\&= -\frac{4x^{2\sqrt{\pi}} c_2 \sqrt{\pi} + c_1 x^{-2\sqrt{\pi}}}{4\sqrt{\pi}}\end{aligned}$$

Now the particular solution to this ODE is found

$$x^2 y'' + x y' - 4\pi y = x$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned}y_1 &= x^{-2\sqrt{\pi}} \\y_2 &= x^{2\sqrt{\pi}}\end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} x^{-2\sqrt{\pi}} & x^{2\sqrt{\pi}} \\ \frac{d}{dx}(x^{-2\sqrt{\pi}}) & \frac{d}{dx}(x^{2\sqrt{\pi}}) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} x^{-2\sqrt{\pi}} & x^{2\sqrt{\pi}} \\ -\frac{2x^{-2\sqrt{\pi}}\sqrt{\pi}}{x} & \frac{2x^{2\sqrt{\pi}}\sqrt{\pi}}{x} \end{vmatrix}$$

Therefore

$$W = (x^{-2\sqrt{\pi}}) \left(\frac{2x^{2\sqrt{\pi}}\sqrt{\pi}}{x} \right) - (x^{2\sqrt{\pi}}) \left(-\frac{2x^{-2\sqrt{\pi}}\sqrt{\pi}}{x} \right)$$

Which simplifies to

$$W = \frac{4\sqrt{\pi}}{x}$$

Which simplifies to

$$W = \frac{4\sqrt{\pi}}{x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x x^{2\sqrt{\pi}}}{4\sqrt{\pi} x} dx$$

Which simplifies to

$$u_1 = - \int \frac{x^{2\sqrt{\pi}}}{4\sqrt{\pi}} dx$$

Hence

$$u_1 = -\frac{x^{1+2\sqrt{\pi}}}{4\sqrt{\pi} (1 + 2\sqrt{\pi})}$$

And Eq. (3) becomes

$$u_2 = \int \frac{x^{-2\sqrt{\pi}} x}{4\sqrt{\pi} x} dx$$

Which simplifies to

$$u_2 = \int \frac{x^{-2\sqrt{\pi}}}{4\sqrt{\pi}} dx$$

Hence

$$u_2 = -\frac{x^{-2\sqrt{\pi}+1}}{4\sqrt{\pi} (2\sqrt{\pi} - 1)}$$

Which simplifies to

$$u_1 = -\frac{x^{1+2\sqrt{\pi}}}{\sqrt{\pi} (4 + 8\sqrt{\pi})}$$
$$u_2 = -\frac{x^{-2\sqrt{\pi}+1}}{\sqrt{\pi} (8\sqrt{\pi} - 4)}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{x^{1+2\sqrt{\pi}} x^{-2\sqrt{\pi}}}{\sqrt{\pi} (4 + 8\sqrt{\pi})} - \frac{x^{-2\sqrt{\pi}+1} x^{2\sqrt{\pi}}}{\sqrt{\pi} (8\sqrt{\pi} - 4)}$$

Which simplifies to

$$y_p(x) = -\frac{x}{4\pi - 1}$$

Therefore the general solution is

$$y = y_h + y_p$$
$$= \left(\left(-\frac{c_1 x^{-4\sqrt{\pi}}}{4\sqrt{\pi}} + c_2 \right) x^{2\sqrt{\pi}} \right) + \left(-\frac{x}{4\pi - 1} \right)$$
$$= -\frac{x}{4\pi - 1} + \left(-\frac{c_1 x^{-4\sqrt{\pi}}}{4\sqrt{\pi}} + c_2 \right) x^{2\sqrt{\pi}}$$

Which simplifies to

$$y = -\frac{x}{4\pi - 1} + \left(-\frac{c_1 x^{-4\sqrt{\pi}}}{4\sqrt{\pi}} + c_2 \right) x^{2\sqrt{\pi}}$$

Summary

The solution(s) found are the following

$$y = -\frac{x}{4\pi - 1} + \left(-\frac{c_1 x^{-4\sqrt{\pi}}}{4\sqrt{\pi}} + c_2 \right) x^{2\sqrt{\pi}} \quad (1)$$

Verification of solutions

$$y = -\frac{x}{4\pi - 1} + \left(-\frac{c_1 x^{-4\sqrt{\pi}}}{4\sqrt{\pi}} + c_2 \right) x^{2\sqrt{\pi}}$$

Verified OK.

16.9.5 Solving using Kovacic algorithm

Writing the ode as

$$x^2 y'' + x y' - 4\pi y = 0 \quad (1)$$

$$A y'' + B y' + C y = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= x^2 \\ B &= x \\ C &= -4\pi \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = y e^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = r z(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1 + 16\pi}{4x^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 + 16\pi \\ t &= 4x^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{-1 + 16\pi}{4x^2} \right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 206: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = \frac{-\frac{1}{4} + 4\pi}{x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = -\frac{1}{4} + 4\pi$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{1}{2} + 2\sqrt{\pi} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{2} - 2\sqrt{\pi} \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ , which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = \frac{-1 + 16\pi}{4x^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = -\frac{1}{4} + 4\pi$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{1}{2} + 2\sqrt{\pi} \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{2} - 2\sqrt{\pi} \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{-1 + 16\pi}{4x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	$\frac{1}{2} + 2\sqrt{\pi}$	$\frac{1}{2} - 2\sqrt{\pi}$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	$\frac{1}{2} + 2\sqrt{\pi}$	$\frac{1}{2} - 2\sqrt{\pi}$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = \frac{1}{2} - 2\sqrt{\pi}$ then

$$\begin{aligned} d &= \alpha_\infty^- - (\alpha_{c_1}^-) \\ &= \frac{1}{2} - 2\sqrt{\pi} - \left(\frac{1}{2} - 2\sqrt{\pi} \right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c) [\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty) [\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= \left((-) [\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{x - c_1} \right) + (-) [\sqrt{r}]_\infty \\ &= \frac{\frac{1}{2} - 2\sqrt{\pi}}{x} + (-) (0) \\ &= \frac{\frac{1}{2} - 2\sqrt{\pi}}{x} \\ &= \frac{1 - 4\sqrt{\pi}}{2x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \tag{1A}$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$(0) + 2\left(\frac{\frac{1}{2} - 2\sqrt{\pi}}{x}\right) (0) + \left(\left(-\frac{\frac{1}{2} - 2\sqrt{\pi}}{x^2}\right) + \left(\frac{\frac{1}{2} - 2\sqrt{\pi}}{x}\right)^2 - \left(\frac{-1 + 16\pi}{4x^2}\right)\right) = 0$$
$$0 = 0$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= pe^{\int \omega dx} \\ &= e^{\int \frac{\frac{1}{2} - 2\sqrt{\pi}}{x} dx} \\ &= x^{\frac{1}{2} - 2\sqrt{\pi}} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{x}{x^2} dx} \\ &= z_1 e^{-\frac{\ln(x)}{2}} \\ &= z_1 \left(\frac{1}{\sqrt{x}}\right) \end{aligned}$$

Which simplifies to

$$y_1 = x^{-2\sqrt{\pi}}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{x}{x^2} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-\ln(x)}}{(y_1)^2} dx \\ &= y_1 \left(\frac{x^{4\sqrt{\pi}}}{4\sqrt{\pi}}\right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(x^{-2\sqrt{\pi}} \right) + c_2 \left(x^{-2\sqrt{\pi}} \left(\frac{x^{4\sqrt{\pi}}}{4\sqrt{\pi}} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$x^2 y'' + x y' - 4\pi y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 x^{-2\sqrt{\pi}} + \frac{c_2 x^{2\sqrt{\pi}}}{4\sqrt{\pi}}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned} y_1 &= x^{-2\sqrt{\pi}} \\ y_2 &= \frac{x^{2\sqrt{\pi}}}{4\sqrt{\pi}} \end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} x^{-2\sqrt{\pi}} & \frac{x^{2\sqrt{\pi}}}{4\sqrt{\pi}} \\ \frac{d}{dx}(x^{-2\sqrt{\pi}}) & \frac{d}{dx}\left(\frac{x^{2\sqrt{\pi}}}{4\sqrt{\pi}}\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} x^{-2\sqrt{\pi}} & \frac{x^{2\sqrt{\pi}}}{4\sqrt{\pi}} \\ -\frac{2x^{-2\sqrt{\pi}}\sqrt{\pi}}{x} & \frac{x^{2\sqrt{\pi}}}{2x} \end{vmatrix}$$

Therefore

$$W = (x^{-2\sqrt{\pi}}) \left(\frac{x^{2\sqrt{\pi}}}{2x}\right) - \left(\frac{x^{2\sqrt{\pi}}}{4\sqrt{\pi}}\right) \left(-\frac{2x^{-2\sqrt{\pi}}\sqrt{\pi}}{x}\right)$$

Which simplifies to

$$W = \frac{x^{2\sqrt{\pi}}x^{-2\sqrt{\pi}}}{x}$$

Which simplifies to

$$W = \frac{1}{x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\frac{x^{2\sqrt{\pi}}x}{4\sqrt{\pi}}}{x} dx$$

Which simplifies to

$$u_1 = - \int \frac{x^{2\sqrt{\pi}}}{4\sqrt{\pi}} dx$$

Hence

$$u_1 = - \frac{x^{1+2\sqrt{\pi}}}{4\sqrt{\pi} (1 + 2\sqrt{\pi})}$$

And Eq. (3) becomes

$$u_2 = \int \frac{x^{-2\sqrt{\pi}}x}{x} dx$$

Which simplifies to

$$u_2 = \int x^{-2\sqrt{\pi}} dx$$

Hence

$$u_2 = -\frac{x^{-2\sqrt{\pi}+1}}{2\sqrt{\pi}-1}$$

Which simplifies to

$$u_1 = -\frac{x^{1+2\sqrt{\pi}}}{\sqrt{\pi}(4+8\sqrt{\pi})}$$
$$u_2 = -\frac{x^{-2\sqrt{\pi}+1}}{2\sqrt{\pi}-1}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{x^{1+2\sqrt{\pi}}x^{-2\sqrt{\pi}}}{\sqrt{\pi}(4+8\sqrt{\pi})} - \frac{x^{-2\sqrt{\pi}+1}x^{2\sqrt{\pi}}}{4(2\sqrt{\pi}-1)\sqrt{\pi}}$$

Which simplifies to

$$y_p(x) = -\frac{x}{4\pi-1}$$

Therefore the general solution is

$$y = y_h + y_p$$
$$= \left(c_1 x^{-2\sqrt{\pi}} + \frac{c_2 x^{2\sqrt{\pi}}}{4\sqrt{\pi}} \right) + \left(-\frac{x}{4\pi-1} \right)$$

Summary

The solution(s) found are the following

$$y = c_1 x^{-2\sqrt{\pi}} + \frac{c_2 x^{2\sqrt{\pi}}}{4\sqrt{\pi}} - \frac{x}{4\pi-1} \quad (1)$$

Verification of solutions

$$y = c_1 x^{-2\sqrt{\pi}} + \frac{c_2 x^{2\sqrt{\pi}}}{4\sqrt{\pi}} - \frac{x}{4\pi-1}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    checking if the LODE is of Euler type  
    <- LODE of Euler type successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 44

```
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)-4*Pi*y(x)=x,y(x), singsol=all)
```

$$y(x) = \frac{c_2(4\pi - 1)x^{-2\sqrt{\pi}} + c_1(4\pi - 1)x^{2\sqrt{\pi}} - x}{4\pi - 1}$$

✓ Solution by Mathematica

Time used: 0.042 (sec). Leaf size: 39

```
DSolve[x^2*y'[x]+x*y'[x]-4*Pi*y[x]==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 x^{2\sqrt{\pi}} + c_1 x^{-2\sqrt{\pi}} + \frac{x}{1 - 4\pi}$$

17 Chapter 4. Linear equations with Regular Singular Points. Page 154

17.1 problem 1(a)	1333
17.2 problem 1(b)	1350
17.3 problem 1(c)	1365
17.4 problem 1(d)	1369
17.5 problem 1(e)	1386
17.6 problem 1(f)	1399
17.7 problem 1(g)	1417
17.8 problem 2(b)	1427
17.9 problem 2(c)	1442
17.10 problem 2(d)	1458

17.1 problem 1(a)

17.1.1 Maple step by step solution 1346

Internal problem ID [6040]

Internal file name [OUTPUT/5288_Sunday_June_05_2022_03_29_33_PM_87366917/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 154

Problem number: 1(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference is integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$x^2y'' + (x^2 + x)y' - y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2y'' + (x^2 + x)y' - y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{1+x}{x}$$
$$q(x) = -\frac{1}{x^2}$$

Table 207: Table $p(x), q(x)$ singularities.

$p(x) = \frac{1+x}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = -\frac{1}{x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2 y'' + (x^2 + x) y' - y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned}
 & x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\
 & + (x^2 + x) \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) - \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0
 \end{aligned} \tag{1}$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{1+n+r} a_n (n+r) \right) \\ & + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) \right) + \sum_{n=0}^{\infty} (-a_n x^{n+r}) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} x^{1+n+r} a_n (n+r) = \sum_{n=1}^{\infty} a_{n-1} (n+r-1) x^{n+r}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=1}^{\infty} a_{n-1} (n+r-1) x^{n+r} \right) \\ & + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) \right) + \sum_{n=0}^{\infty} (-a_n x^{n+r}) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r} a_n (n+r) (n+r-1) + x^{n+r} a_n (n+r) - a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$x^r a_0 r(-1+r) + x^r a_0 r - a_0 x^r = 0$$

Or

$$(x^r r(-1+r) + x^r r - x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(r^2 - 1) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$r^2 - 1 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 1 \\ r_2 &= -1 \end{aligned}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(r^2 - 1) x^r = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = 2$ is an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= C y_1(x) \ln(x) + x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= x \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= C y_1(x) \ln(x) + \frac{\sum_{n=0}^{\infty} b_n x^n}{x} \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^{1+n} \\ y_2(x) &= C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n-1} \right) \end{aligned}$$

Where C above can be zero. We start by finding y_1 . Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) + a_{n-1}(n+r-1) + a_n(n+r) - a_n = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{a_{n-1}}{1+n+r} \quad (4)$$

Which for the root $r = 1$ becomes

$$a_n = -\frac{a_{n-1}}{2+n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 1$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = -\frac{1}{2+r}$$

Which for the root $r = 1$ becomes

$$a_1 = -\frac{1}{3}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{2+r}$	$-\frac{1}{3}$

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{(2+r)(3+r)}$$

Which for the root $r = 1$ becomes

$$a_2 = \frac{1}{12}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{2+r}$	$-\frac{1}{3}$
a_2	$\frac{1}{(2+r)(3+r)}$	$\frac{1}{12}$

For $n = 3$, using the above recursive equation gives

$$a_3 = -\frac{1}{(3+r)(2+r)(4+r)}$$

Which for the root $r = 1$ becomes

$$a_3 = -\frac{1}{60}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{2+r}$	$-\frac{1}{3}$
a_2	$\frac{1}{(2+r)(3+r)}$	$\frac{1}{12}$
a_3	$-\frac{1}{(3+r)(2+r)(4+r)}$	$-\frac{1}{60}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{(2+r)(4+r)(3+r)(5+r)}$$

Which for the root $r = 1$ becomes

$$a_4 = \frac{1}{360}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{2+r}$	$-\frac{1}{3}$
a_2	$\frac{1}{(2+r)(3+r)}$	$\frac{1}{12}$
a_3	$-\frac{1}{(3+r)(2+r)(4+r)}$	$-\frac{1}{60}$
a_4	$\frac{1}{(2+r)(4+r)(3+r)(5+r)}$	$\frac{1}{360}$

For $n = 5$, using the above recursive equation gives

$$a_5 = -\frac{1}{(2+r)(3+r)(5+r)(6+r)(4+r)}$$

Which for the root $r = 1$ becomes

$$a_5 = -\frac{1}{2520}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{2+r}$	$-\frac{1}{3}$
a_2	$\frac{1}{(2+r)(3+r)}$	$\frac{1}{12}$
a_3	$-\frac{1}{(3+r)(2+r)(4+r)}$	$-\frac{1}{60}$
a_4	$\frac{1}{(2+r)(4+r)(3+r)(5+r)}$	$\frac{1}{360}$
a_5	$-\frac{1}{(2+r)(3+r)(5+r)(6+r)(4+r)}$	$-\frac{1}{2520}$

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{1}{(2+r)(3+r)(6+r)(4+r)(5+r)(7+r)}$$

Which for the root $r = 1$ becomes

$$a_6 = \frac{1}{20160}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{2+r}$	$-\frac{1}{3}$
a_2	$\frac{1}{(2+r)(3+r)}$	$\frac{1}{12}$
a_3	$-\frac{1}{(3+r)(2+r)(4+r)}$	$-\frac{1}{60}$
a_4	$\frac{1}{(2+r)(4+r)(3+r)(5+r)}$	$\frac{1}{360}$
a_5	$-\frac{1}{(2+r)(3+r)(5+r)(6+r)(4+r)}$	$-\frac{1}{2520}$
a_6	$\frac{1}{(2+r)(3+r)(6+r)(4+r)(5+r)(7+r)}$	$\frac{1}{20160}$

For $n = 7$, using the above recursive equation gives

$$a_7 = -\frac{1}{(2+r)(3+r)(4+r)(5+r)(7+r)(8+r)(6+r)}$$

Which for the root $r = 1$ becomes

$$a_7 = -\frac{1}{181440}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{2+r}$	$-\frac{1}{3}$
a_2	$\frac{1}{(2+r)(3+r)}$	$\frac{1}{12}$
a_3	$-\frac{1}{(3+r)(2+r)(4+r)}$	$-\frac{1}{60}$
a_4	$\frac{1}{(2+r)(4+r)(3+r)(5+r)}$	$\frac{1}{360}$
a_5	$-\frac{1}{(2+r)(3+r)(5+r)(6+r)(4+r)}$	$-\frac{1}{2520}$
a_6	$\frac{1}{(2+r)(3+r)(6+r)(4+r)(5+r)(7+r)}$	$\frac{1}{20160}$
a_7	$-\frac{1}{(2+r)(3+r)(4+r)(5+r)(7+r)(8+r)(6+r)}$	$-\frac{1}{181440}$

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots) \\ &= x \left(1 - \frac{x}{3} + \frac{x^2}{12} - \frac{x^3}{60} + \frac{x^4}{360} - \frac{x^5}{2520} + \frac{x^6}{20160} - \frac{x^7}{181440} + O(x^8) \right) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Let

$$r_1 - r_2 = N$$

Where N is positive integer which is the difference between the two roots. r_1 is taken as the larger root. Hence for this problem we have $N = 2$. Now we need to determine if C is zero or not. This is done by finding $\lim_{r \rightarrow r_2} a_2(r)$. If this limit exists, then $C = 0$, else we need to keep the log term and $C \neq 0$. The above table shows that

$$\begin{aligned} a_N &= a_2 \\ &= \frac{1}{(2+r)(3+r)} \end{aligned}$$

Therefore

$$\begin{aligned} \lim_{r \rightarrow r_2} \frac{1}{(2+r)(3+r)} &= \lim_{r \rightarrow -1} \frac{1}{(2+r)(3+r)} \\ &= \frac{1}{2} \end{aligned}$$

The limit is $\frac{1}{2}$. Since the limit exists then the log term is not needed and we can set $C = 0$. Therefore the second solution has the form

$$\begin{aligned} y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+r} \\ &= \sum_{n=0}^{\infty} b_n x^{n-1} \end{aligned}$$

Eq (3) derived above is used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. For $1 \leq n$ the recursive equation is

$$b_n(n+r)(n+r-1) + b_{n-1}(n+r-1) + b_n(n+r) - b_n = 0 \quad (4)$$

Which for the root $r = -1$ becomes

$$b_n(n-1)(n-2) + b_{n-1}(n-2) + b_n(n-1) - b_n = 0 \quad (4A)$$

Solving for b_n from the recursive equation (4) gives

$$b_n = -\frac{b_{n-1}}{1+n+r} \quad (5)$$

Which for the root $r = -1$ becomes

$$b_n = -\frac{b_{n-1}}{n} \quad (6)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = -1$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1

For $n = 1$, using the above recursive equation gives

$$b_1 = -\frac{1}{2+r}$$

Which for the root $r = -1$ becomes

$$b_1 = -1$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{1}{2+r}$	-1

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{1}{(2+r)(3+r)}$$

Which for the root $r = -1$ becomes

$$b_2 = \frac{1}{2}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{1}{2+r}$	-1
b_2	$\frac{1}{(2+r)(3+r)}$	$\frac{1}{2}$

For $n = 3$, using the above recursive equation gives

$$b_3 = -\frac{1}{(3+r)(2+r)(4+r)}$$

Which for the root $r = -1$ becomes

$$b_3 = -\frac{1}{6}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{1}{2+r}$	-1
b_2	$\frac{1}{(2+r)(3+r)}$	$\frac{1}{2}$
b_3	$-\frac{1}{(3+r)(2+r)(4+r)}$	$-\frac{1}{6}$

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{1}{(2+r)(4+r)(3+r)(5+r)}$$

Which for the root $r = -1$ becomes

$$b_4 = \frac{1}{24}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{1}{2+r}$	-1
b_2	$\frac{1}{(2+r)(3+r)}$	$\frac{1}{2}$
b_3	$-\frac{1}{(3+r)(2+r)(4+r)}$	$-\frac{1}{6}$
b_4	$\frac{1}{(2+r)(4+r)(3+r)(5+r)}$	$\frac{1}{24}$

For $n = 5$, using the above recursive equation gives

$$b_5 = -\frac{1}{(2+r)(3+r)(5+r)(6+r)(4+r)}$$

Which for the root $r = -1$ becomes

$$b_5 = -\frac{1}{120}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{1}{2+r}$	-1
b_2	$\frac{1}{(2+r)(3+r)}$	$\frac{1}{2}$
b_3	$-\frac{1}{(3+r)(2+r)(4+r)}$	$-\frac{1}{6}$
b_4	$\frac{1}{(2+r)(4+r)(3+r)(5+r)}$	$\frac{1}{24}$
b_5	$-\frac{1}{(2+r)(3+r)(5+r)(6+r)(4+r)}$	$-\frac{1}{120}$

For $n = 6$, using the above recursive equation gives

$$b_6 = \frac{1}{(2+r)(3+r)(6+r)(4+r)(5+r)(7+r)}$$

Which for the root $r = -1$ becomes

$$b_6 = \frac{1}{720}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{1}{2+r}$	-1
b_2	$\frac{1}{(2+r)(3+r)}$	$\frac{1}{2}$
b_3	$-\frac{1}{(3+r)(2+r)(4+r)}$	$-\frac{1}{6}$
b_4	$\frac{1}{(2+r)(4+r)(3+r)(5+r)}$	$\frac{1}{24}$
b_5	$-\frac{1}{(2+r)(3+r)(5+r)(6+r)(4+r)}$	$-\frac{1}{120}$
b_6	$\frac{1}{(2+r)(3+r)(6+r)(4+r)(5+r)(7+r)}$	$\frac{1}{720}$

For $n = 7$, using the above recursive equation gives

$$b_7 = -\frac{1}{(2+r)(3+r)(4+r)(5+r)(7+r)(8+r)(6+r)}$$

Which for the root $r = -1$ becomes

$$b_7 = -\frac{1}{5040}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{1}{2+r}$	-1
b_2	$\frac{1}{(2+r)(3+r)}$	$\frac{1}{2}$
b_3	$-\frac{1}{(3+r)(2+r)(4+r)}$	$-\frac{1}{6}$
b_4	$\frac{1}{(2+r)(4+r)(3+r)(5+r)}$	$\frac{1}{24}$
b_5	$-\frac{1}{(2+r)(3+r)(5+r)(6+r)(4+r)}$	$-\frac{1}{120}$
b_6	$\frac{1}{(2+r)(3+r)(6+r)(4+r)(5+r)(7+r)}$	$\frac{1}{720}$
b_7	$-\frac{1}{(2+r)(3+r)(4+r)(5+r)(7+r)(8+r)(6+r)}$	$-\frac{1}{5040}$

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned}
 y_2(x) &= x(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 + b_7x^7 + b_8x^8 \dots) \\
 &= \frac{1 - x + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + \frac{x^6}{720} - \frac{x^7}{5040} + O(x^8)}{x}
 \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 y_h(x) &= c_1y_1(x) + c_2y_2(x) \\
 &= c_1x \left(1 - \frac{x}{3} + \frac{x^2}{12} - \frac{x^3}{60} + \frac{x^4}{360} - \frac{x^5}{2520} + \frac{x^6}{20160} - \frac{x^7}{181440} + O(x^8) \right) \\
 &\quad + \frac{c_2 \left(1 - x + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + \frac{x^6}{720} - \frac{x^7}{5040} + O(x^8) \right)}{x}
 \end{aligned}$$

Hence the final solution is

$$\begin{aligned}
 y &= y_h \\
 &= c_1x \left(1 - \frac{x}{3} + \frac{x^2}{12} - \frac{x^3}{60} + \frac{x^4}{360} - \frac{x^5}{2520} + \frac{x^6}{20160} - \frac{x^7}{181440} + O(x^8) \right) \\
 &\quad + \frac{c_2 \left(1 - x + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + \frac{x^6}{720} - \frac{x^7}{5040} + O(x^8) \right)}{x}
 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 x \left(1 - \frac{x}{3} + \frac{x^2}{12} - \frac{x^3}{60} + \frac{x^4}{360} - \frac{x^5}{2520} + \frac{x^6}{20160} - \frac{x^7}{181440} + O(x^8) \right) + \frac{c_2 \left(1 - x + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + \frac{x^6}{720} - \frac{x^7}{5040} + O(x^8) \right)}{x} \quad (1)$$

Verification of solutions

$$y = c_1 x \left(1 - \frac{x}{3} + \frac{x^2}{12} - \frac{x^3}{60} + \frac{x^4}{360} - \frac{x^5}{2520} + \frac{x^6}{20160} - \frac{x^7}{181440} + O(x^8) \right) + \frac{c_2 \left(1 - x + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + \frac{x^6}{720} - \frac{x^7}{5040} + O(x^8) \right)}{x}$$

Verified OK.

17.1.1 Maple step by step solution

Let's solve

$$x^2 y'' + (x^2 + x) y' - y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = \frac{y}{x^2} - \frac{(1+x)y'}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{(1+x)y'}{x} - \frac{y}{x^2} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$\left[P_2(x) = \frac{1+x}{x}, P_3(x) = -\frac{1}{x^2} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 1$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = -1$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$x^2 y'' + x(1+x)y' - y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y'$ to series expansion for $m = 1..2$

$$x^m \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1+m}$$

- Shift index using $k \rightarrow k+1-m$

$$x^m \cdot y' = \sum_{k=-1+m}^{\infty} a_{k+1-m} (k+1-m+r) x^{k+r}$$

- Convert $x^2 \cdot y''$ to series expansion

$$x^2 \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(1+r)(-1+r)x^r + \left(\sum_{k=1}^{\infty} (a_k(k+r+1)(k+r-1) + a_{k-1}(k+r-1)) x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$(1+r)(-1+r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \{-1, 1\}$$

- Each term in the series must be 0, giving the recursion relation

$$(k+r-1)(a_k(k+r+1) + a_{k-1}) = 0$$

- Shift index using $k \rightarrow k+1$

$$(k+r)(a_{k+1}(k+2+r) + a_k) = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+1} = -\frac{a_k}{k+2+r}$$

- Recursion relation for $r = -1$

$$a_{k+1} = -\frac{a_k}{k+1}$$

- Solution for $r = -1$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k-1}, a_{k+1} = -\frac{a_k}{k+1} \right]$$

- Recursion relation for $r = 1$

$$a_{k+1} = -\frac{a_k}{k+3}$$

- Solution for $r = 1$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+1}, a_{k+1} = -\frac{a_k}{k+3} \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^{k-1} \right) + \left(\sum_{k=0}^{\infty} b_k x^{k+1} \right), a_{k+1} = -\frac{a_k}{k+1}, b_{k+1} = -\frac{b_k}{k+3} \right]$$

Maple trace **Kovacic algorithm successful**

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Reducible group (found an exponential solution)
    Reducible group (found another exponential solution)
<- Kovacics algorithm successful`

```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 53

Order:=8;

```
dsolve(x^2*diff(y(x),x$2)+(x+x^2)*diff(y(x),x)-y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x \left(1 - \frac{1}{3}x + \frac{1}{12}x^2 - \frac{1}{60}x^3 + \frac{1}{360}x^4 - \frac{1}{2520}x^5 + \frac{1}{20160}x^6 - \frac{1}{181440}x^7 + O(x^8) \right) \\ + \frac{c_2 \left(-2 + 2x - x^2 + \frac{1}{3}x^3 - \frac{1}{12}x^4 + \frac{1}{60}x^5 - \frac{1}{360}x^6 + \frac{1}{2520}x^7 + O(x^8) \right)}{x}$$

✓ Solution by Mathematica

Time used: 0.039 (sec). Leaf size: 92

```
AsymptoticDSolveValue[x^2*y''[x]+(x+x^2)*y'[x]-y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(\frac{x^5}{720} - \frac{x^4}{120} + \frac{x^3}{24} - \frac{x^2}{6} + \frac{x}{2} + \frac{1}{x} - 1 \right) \\ + c_2 \left(\frac{x^7}{20160} - \frac{x^6}{2520} + \frac{x^5}{360} - \frac{x^4}{60} + \frac{x^3}{12} - \frac{x^2}{3} + x \right)$$

17.2 problem 1(b)

Internal problem ID [6041]

Internal file name [OUTPUT/5289_Sunday_June_05_2022_03_29_37_PM_51446362/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 154

Problem number: 1(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference is integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$3x^2y'' + y'x^6 + 2xy = 0$$

With the expansion point for the power series method at $x = 0$.

The ODE is

$$3x^2y'' + y'x^6 + 2xy = 0$$

Or

$$x(y'x^5 + 3y''x + 2y) = 0$$

For $x \neq 0$ the above simplifies to

$$y'x^5 + 3y''x + 2y = 0$$

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$3x^2y'' + y'x^6 + 2xy = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{x^4}{3}$$

$$q(x) = \frac{2}{3x}$$

Table 209: Table $p(x), q(x)$ singularities.

$p(x) = \frac{x^4}{3}$	
singularity	type
$x = \infty$	“regular”
$x = -\infty$	“regular”

$q(x) = \frac{2}{3x}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[\infty, -\infty, 0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$3x^2y'' + y'x^6 + 2xy = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned}
& 3x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\
& + \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) x^6 + 2x \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0
\end{aligned} \tag{1}$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} 3x^{n+r} a_n (n+r)(n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{5+n+r} a_n (n+r) \right) + \left(\sum_{n=0}^{\infty} 2x^{1+n+r} a_n \right) = 0 \tag{2A}$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\begin{aligned}
\sum_{n=0}^{\infty} x^{5+n+r} a_n (n+r) &= \sum_{n=5}^{\infty} a_{n-5} (n-5+r) x^{n+r} \\
\sum_{n=0}^{\infty} 2x^{1+n+r} a_n &= \sum_{n=1}^{\infty} 2a_{n-1} x^{n+r}
\end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\left(\sum_{n=0}^{\infty} 3x^{n+r} a_n (n+r)(n+r-1) \right) + \left(\sum_{n=5}^{\infty} a_{n-5} (n-5+r) x^{n+r} \right) + \left(\sum_{n=1}^{\infty} 2a_{n-1} x^{n+r} \right) = 0 \tag{2B}$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$3x^{n+r} a_n (n+r)(n+r-1) = 0$$

When $n=0$ the above becomes

$$3x^r a_0 r(-1+r) = 0$$

Or

$$3x^r a_0 r(-1 + r) = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$3x^r r(-1 + r) = 0$$

Since the above is true for all x then the indicial equation becomes

$$3r(-1 + r) = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = 1$$

$$r_2 = 0$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$3x^r r(-1 + r) = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = 1$ is an integer, then we can construct two linearly independent solutions

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = C y_1(x) \ln(x) + x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = x \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{1+n}$$

$$y_2(x) = C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Where C above can be zero. We start by finding y_1 . Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = -\frac{2}{3r(1+r)}$$

Substituting $n = 2$ in Eq. (2B) gives

$$a_2 = \frac{4}{9r(1+r)^2(2+r)}$$

Substituting $n = 3$ in Eq. (2B) gives

$$a_3 = -\frac{8}{27r(1+r)^2(2+r)^2(3+r)}$$

Substituting $n = 4$ in Eq. (2B) gives

$$a_4 = \frac{16}{81r(1+r)^2(2+r)^2(3+r)^2(4+r)}$$

For $5 \leq n$ the recursive equation is

$$3a_n(n+r)(n+r-1) + a_{n-5}(n-5+r) + 2a_{n-1} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{na_{n-5} + ra_{n-5} - 5a_{n-5} + 2a_{n-1}}{3(n+r)(n+r-1)} \quad (4)$$

Which for the root $r = 1$ becomes

$$a_n = \frac{-na_{n-5} + 4a_{n-5} - 2a_{n-1}}{3(1+n)n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 1$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{2}{3r(1+r)}$	$-\frac{1}{3}$
a_2	$\frac{4}{9r(1+r)^2(2+r)}$	$\frac{1}{27}$
a_3	$-\frac{8}{27r(1+r)^2(2+r)^2(3+r)}$	$-\frac{1}{486}$
a_4	$\frac{16}{81r(1+r)^2(2+r)^2(3+r)^2(4+r)}$	$\frac{1}{14580}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{-81r^9 - 1296r^8 - 8586r^7 - 30456r^6 - 62289r^5 - 73224r^4 - 45684r^3 - 11664r^2 - 32}{243r(1+r)^2(2+r)^2(3+r)^2(4+r)^2(5+r)}$$

Which for the root $r = 1$ becomes

$$a_5 = -\frac{7291}{656100}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{2}{3r(1+r)}$	$-\frac{1}{3}$
a_2	$\frac{4}{9r(1+r)^2(2+r)}$	$\frac{1}{27}$
a_3	$-\frac{8}{27r(1+r)^2(2+r)^2(3+r)}$	$-\frac{1}{486}$
a_4	$\frac{16}{81r(1+r)^2(2+r)^2(3+r)^2(4+r)}$	$\frac{1}{14580}$
a_5	$\frac{-81r^9 - 1296r^8 - 8586r^7 - 30456r^6 - 62289r^5 - 73224r^4 - 45684r^3 - 11664r^2 - 32}{243r(1+r)^2(2+r)^2(3+r)^2(4+r)^2(5+r)}$	$-\frac{7291}{656100}$

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{\frac{4}{9}r^9 + \frac{82}{9}r^8 + \frac{752}{9}r^7 + \frac{4052}{9}r^6 + \frac{14084}{9}r^5 + \frac{32498}{9}r^4 + \frac{49288}{9}r^3 + \frac{46888}{9}r^2 + \frac{8384}{3}r + \frac{466624}{729}}{r(1+r)^2(2+r)^2(3+r)^2(4+r)^2(5+r)^2(6+r)}$$

Which for the root $r = 1$ becomes

$$a_6 = \frac{225991}{41334300}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{2}{3r(1+r)}$	$-\frac{1}{3}$
a_2	$\frac{4}{9r(1+r)^2(2+r)}$	$\frac{1}{27}$
a_3	$-\frac{8}{27r(1+r)^2(2+r)^2(3+r)}$	$-\frac{1}{486}$
a_4	$\frac{16}{81r(1+r)^2(2+r)^2(3+r)^2(4+r)}$	$\frac{1}{14580}$
a_5	$\frac{-81r^9 - 1296r^8 - 8586r^7 - 30456r^6 - 62289r^5 - 73224r^4 - 45684r^3 - 11664r^2 - 32}{243r(1+r)^2(2+r)^2(3+r)^2(4+r)^2(5+r)}$	$-\frac{7291}{656100}$
a_6	$\frac{\frac{4}{9}r^9 + \frac{82}{9}r^8 + \frac{752}{9}r^7 + \frac{4052}{9}r^6 + \frac{14084}{9}r^5 + \frac{32498}{9}r^4 + \frac{49288}{9}r^3 + \frac{46888}{9}r^2 + \frac{8384}{3}r + \frac{466624}{729}}{r(1+r)^2(2+r)^2(3+r)^2(4+r)^2(5+r)^2(6+r)}$	$\frac{225991}{41334300}$

For $n = 7$, using the above recursive equation gives

$$a_7 = \frac{-\frac{4}{9}r^9 - \frac{100}{9}r^8 - \frac{392}{3}r^7 - \frac{25400}{27}r^6 - \frac{13516}{3}r^5 - \frac{396140}{27}r^4 - \frac{289024}{9}r^3 - \frac{1219760}{27}r^2 - \frac{331648}{9}r - \frac{28926848}{2187}}{r(1+r)^2(2+r)^2(3+r)^2(4+r)^2(5+r)^2(6+r)^2(7+r)}$$

Which for the root $r = 1$ becomes

$$a_7 = -\frac{2522341}{3472081200}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{2}{3r(1+r)}$	$-\frac{1}{3}$
a_2	$\frac{4}{9r(1+r)^2(2+r)}$	$\frac{1}{27}$
a_3	$-\frac{8}{27r(1+r)^2(2+r)^2(3+r)}$	$-\frac{1}{486}$
a_4	$\frac{16}{81r(1+r)^2(2+r)^2(3+r)^2(4+r)}$	$\frac{1}{14580}$
a_5	$\frac{-81r^9 - 1296r^8 - 8586r^7 - 30456r^6 - 62289r^5 - 73224r^4 - 45684r^3 - 11664r^2 - 32}{243r(1+r)^2(2+r)^2(3+r)^2(4+r)^2(5+r)}$	$-\frac{7291}{656100}$
a_6	$\frac{\frac{4}{9}r^9 + \frac{82}{9}r^8 + \frac{752}{9}r^7 + \frac{4052}{9}r^6 + \frac{14084}{9}r^5 + \frac{32498}{9}r^4 + \frac{49288}{9}r^3 + \frac{46888}{9}r^2 + \frac{8384}{3}r + \frac{466624}{729}}{r(1+r)^2(2+r)^2(3+r)^2(4+r)^2(5+r)^2(6+r)}$	$\frac{225991}{41334300}$
a_7	$\frac{-\frac{4}{9}r^9 - \frac{100}{9}r^8 - \frac{392}{3}r^7 - \frac{25400}{27}r^6 - \frac{13516}{3}r^5 - \frac{396140}{27}r^4 - \frac{289024}{9}r^3 - \frac{1219760}{27}r^2 - \frac{331648}{9}r - \frac{28926848}{2187}}{r(1+r)^2(2+r)^2(3+r)^2(4+r)^2(5+r)^2(6+r)^2(7+r)}$	$-\frac{2522341}{3472081200}$

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots) \\ &= x\left(1 - \frac{x}{3} + \frac{x^2}{27} - \frac{x^3}{486} + \frac{x^4}{14580} - \frac{7291x^5}{656100} + \frac{225991x^6}{41334300} - \frac{2522341x^7}{3472081200} + O(x^8)\right) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Let

$$r_1 - r_2 = N$$

Where N is positive integer which is the difference between the two roots. r_1 is taken as the larger root. Hence for this problem we have $N = 1$. Now we need to determine if C is zero or not. This is done by finding $\lim_{r \rightarrow r_2} a_1(r)$. If this limit exists, then $C = 0$, else we need to keep the log term and $C \neq 0$. The above table shows that

$$\begin{aligned} a_N &= a_1 \\ &= -\frac{2}{3r(1+r)} \end{aligned}$$

Therefore

$$\begin{aligned}\lim_{r \rightarrow r_2} -\frac{2}{3r(1+r)} &= \lim_{r \rightarrow 0} -\frac{2}{3r(1+r)} \\ &= \text{undefined}\end{aligned}$$

Since the limit does not exist then the log term is needed. Therefore the second solution has the form

$$y_2(x) = Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right)$$

Therefore

$$\begin{aligned}\frac{d}{dx} y_2(x) &= Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2}(n+r_2)}{x} \right) \\ &= Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} x^{-1+n+r_2} b_n (n+r_2) \right) \\ \frac{d^2}{dx^2} y_2(x) &= Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} \\ &\quad + \sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2}(n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2}(n+r_2)}{x^2} \right) \\ &= Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} + \left(\sum_{n=0}^{\infty} x^{-2+n+r_2} b_n (n+r_2) (-1+n+r_2) \right)\end{aligned}$$

Substituting these back into the given ode $3x^2 y'' + y' x^6 + 2xy = 0$ gives

$$\begin{aligned}3x^2 &\left(Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} \right. \\ &\quad \left. + \sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2}(n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2}(n+r_2)}{x^2} \right) \right) \\ &+ \left(Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2}(n+r_2)}{x} \right) \right) x^6 \\ &+ 2x \left(Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) \right) = 0\end{aligned}$$

Which can be written as

$$\begin{aligned} & \left((y_1'(x) x^6 + 3y_1''(x) x^2 + 2y_1(x) x) \ln(x) + 3x^2 \left(\frac{2y_1'(x)}{x} - \frac{y_1(x)}{x^2} \right) \right. \\ & \left. + y_1(x) x^5 \right) C + \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2} (n+r_2)}{x} \right) x^6 \\ & + 3x^2 \left(\sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2} (n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2} (n+r_2)}{x^2} \right) \right) + 2x \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) = 0 \end{aligned} \quad (7)$$

But since $y_1(x)$ is a solution to the ode, then

$$y_1'(x) x^6 + 3y_1''(x) x^2 + 2y_1(x) x = 0$$

Eq (7) simplifies to

$$\begin{aligned} & \left(3x^2 \left(\frac{2y_1'(x)}{x} - \frac{y_1(x)}{x^2} \right) + y_1(x) x^5 \right) C + \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2} (n+r_2)}{x} \right) x^6 \\ & + 3x^2 \left(\sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2} (n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2} (n+r_2)}{x^2} \right) \right) + 2x \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) = 0 \end{aligned} \quad (8)$$

Substituting $y_1 = \sum_{n=0}^{\infty} a_n x^{n+r_1}$ into the above gives

$$\begin{aligned} & \left(6 \left(\sum_{n=0}^{\infty} x^{-1+n+r_1} a_n (n+r_1) \right) x + (x^5 - 3) \left(\sum_{n=0}^{\infty} a_n x^{n+r_1} \right) \right) C \\ & + \left(\sum_{n=0}^{\infty} x^{-1+n+r_2} b_n (n+r_2) \right) x^6 \\ & + 3 \left(\sum_{n=0}^{\infty} x^{-2+n+r_2} b_n (n+r_2) (-1+n+r_2) \right) x^2 + 2x \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) = 0 \end{aligned} \quad (9)$$

Since $r_1 = 1$ and $r_2 = 0$ then the above becomes

$$\begin{aligned} & \left(6 \left(\sum_{n=0}^{\infty} x^n a_n (1+n) \right) x + (x^5 - 3) \left(\sum_{n=0}^{\infty} a_n x^{1+n} \right) \right) C \\ & + \left(\sum_{n=0}^{\infty} x^{n-1} b_n n \right) x^6 + 3 \left(\sum_{n=0}^{\infty} x^{-2+n} b_n n (n-1) \right) x^2 + 2x \left(\sum_{n=0}^{\infty} b_n x^n \right) = 0 \end{aligned} \quad (10)$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 6C x^{1+n} a_n (1+n) \right) + \left(\sum_{n=0}^{\infty} C x^{n+6} a_n \right) + \sum_{n=0}^{\infty} (-3C x^{1+n} a_n) \\ & + \left(\sum_{n=0}^{\infty} n x^{5+n} b_n \right) + \left(\sum_{n=0}^{\infty} 3n x^n b_n (n-1) \right) + \left(\sum_{n=0}^{\infty} 2x^{1+n} b_n \right) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=0}^{\infty} 6C x^{1+n} a_n (1+n) &= \sum_{n=1}^{\infty} 6C a_{n-1} n x^n \\ \sum_{n=0}^{\infty} C x^{n+6} a_n &= \sum_{n=6}^{\infty} C a_{n-6} x^n \\ \sum_{n=0}^{\infty} (-3C x^{1+n} a_n) &= \sum_{n=1}^{\infty} (-3C a_{n-1} x^n) \\ \sum_{n=0}^{\infty} n x^{5+n} b_n &= \sum_{n=5}^{\infty} (n-5) b_{n-5} x^n \\ \sum_{n=0}^{\infty} 2x^{1+n} b_n &= \sum_{n=1}^{\infty} 2b_{n-1} x^n \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to n .

$$\begin{aligned} & \left(\sum_{n=1}^{\infty} 6C a_{n-1} n x^n \right) + \left(\sum_{n=6}^{\infty} C a_{n-6} x^n \right) + \sum_{n=1}^{\infty} (-3C a_{n-1} x^n) \\ & + \left(\sum_{n=5}^{\infty} (n-5) b_{n-5} x^n \right) + \left(\sum_{n=0}^{\infty} 3n x^n b_n (n-1) \right) + \left(\sum_{n=1}^{\infty} 2b_{n-1} x^n \right) = 0 \end{aligned} \quad (2B)$$

For $n = 0$ in Eq. (2B), we choose arbitrary value for b_0 as $b_0 = 1$. For $n = N$, where $N = 1$ which is the difference between the two roots, we are free to choose $b_1 = 0$. Hence for $n = 1$, Eq (2B) gives

$$3C + 2 = 0$$

Which is solved for C . Solving for C gives

$$C = -\frac{2}{3}$$

For $n = 2$, Eq (2B) gives

$$9Ca_1 + 2b_1 + 6b_2 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$6b_2 + 2 = 0$$

Solving the above for b_2 gives

$$b_2 = -\frac{1}{3}$$

For $n = 3$, Eq (2B) gives

$$15Ca_2 + 2b_2 + 18b_3 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$18b_3 - \frac{28}{27} = 0$$

Solving the above for b_3 gives

$$b_3 = \frac{14}{243}$$

For $n = 4$, Eq (2B) gives

$$21Ca_3 + 2b_3 + 36b_4 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$36b_4 + \frac{35}{243} = 0$$

Solving the above for b_4 gives

$$b_4 = -\frac{35}{8748}$$

For $n = 5$, Eq (2B) gives

$$27Ca_4 + 2b_4 + 60b_5 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$60b_5 - \frac{101}{10935} = 0$$

Solving the above for b_5 gives

$$b_5 = \frac{101}{656100}$$

For $n = 6$, Eq (2B) gives

$$(a_0 + 33a_5)C + b_1 + 2b_5 + 90b_6 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$-\frac{69199}{164025} + 90b_6 = 0$$

Solving the above for b_6 gives

$$b_6 = \frac{69199}{14762250}$$

For $n = 7$, Eq (2B) gives

$$(a_1 + 39a_6)C + 2b_2 + 2b_6 + 126b_7 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$-\frac{19882543}{34445250} + 126b_7 = 0$$

Solving the above for b_7 gives

$$b_7 = \frac{19882543}{4340101500}$$

Now that we found all b_n and C , we can calculate the second solution from

$$y_2(x) = Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right)$$

Using the above value found for $C = -\frac{2}{3}$ and all b_n , then the second solution becomes

$$y_2(x) = -\frac{2}{3} \left(x \left(1 - \frac{x}{3} + \frac{x^2}{27} - \frac{x^3}{486} + \frac{x^4}{14580} - \frac{7291x^5}{656100} + \frac{225991x^6}{41334300} - \frac{2522341x^7}{3472081200} + O(x^8) \right) \right) \ln(x) + 1 - \frac{x^2}{3} + \frac{14x^3}{243} - \frac{35x^4}{8748} + \frac{101x^5}{656100} + \frac{69199x^6}{14762250} + \frac{19882543x^7}{4340101500} + O(x^8)$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 y_1(x) + c_2 y_2(x)$$

$$= c_1 x \left(1 - \frac{x}{3} + \frac{x^2}{27} - \frac{x^3}{486} + \frac{x^4}{14580} - \frac{7291x^5}{656100} + \frac{225991x^6}{41334300} - \frac{2522341x^7}{3472081200} + O(x^8) \right) + c_2 \left(-\frac{2}{3} \left(x \left(1 - \frac{x}{3} + \frac{x^2}{27} - \frac{x^3}{486} + \frac{x^4}{14580} - \frac{7291x^5}{656100} + \frac{225991x^6}{41334300} - \frac{2522341x^7}{3472081200} + O(x^8) \right) \right) \ln(x) + 1 - \frac{x^2}{3} + \frac{14x^3}{243} - \frac{35x^4}{8748} + \frac{101x^5}{656100} + \frac{69199x^6}{14762250} + \frac{19882543x^7}{4340101500} + O(x^8) \right)$$

Hence the final solution is

$$\begin{aligned}
 y &= y_h \\
 &= c_1 x \left(1 - \frac{x}{3} + \frac{x^2}{27} - \frac{x^3}{486} + \frac{x^4}{14580} - \frac{7291x^5}{656100} + \frac{225991x^6}{41334300} - \frac{2522341x^7}{3472081200} + O(x^8) \right) \\
 &\quad + c_2 \left(-\frac{2x \left(1 - \frac{x}{3} + \frac{x^2}{27} - \frac{x^3}{486} + \frac{x^4}{14580} - \frac{7291x^5}{656100} + \frac{225991x^6}{41334300} - \frac{2522341x^7}{3472081200} + O(x^8) \right) \ln(x)}{3} \right. \\
 &\quad \left. + 1 - \frac{x^2}{3} + \frac{14x^3}{243} - \frac{35x^4}{8748} + \frac{101x^5}{656100} + \frac{69199x^6}{14762250} + \frac{19882543x^7}{4340101500} + O(x^8) \right)
 \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}
 y &= c_1 x \left(1 - \frac{x}{3} + \frac{x^2}{27} - \frac{x^3}{486} + \frac{x^4}{14580} - \frac{7291x^5}{656100} + \frac{225991x^6}{41334300} - \frac{2522341x^7}{3472081200} + O(x^8) \right) \\
 &\quad + c_2 \left(-\frac{2x \left(1 - \frac{x}{3} + \frac{x^2}{27} - \frac{x^3}{486} + \frac{x^4}{14580} - \frac{7291x^5}{656100} + \frac{225991x^6}{41334300} - \frac{2522341x^7}{3472081200} + O(x^8) \right) \ln(x)}{3} \right. \\
 &\quad \left. + 1 - \frac{x^2}{3} + \frac{14x^3}{243} - \frac{35x^4}{8748} + \frac{101x^5}{656100} + \frac{69199x^6}{14762250} + \frac{19882543x^7}{4340101500} + O(x^8) \right)
 \end{aligned} \tag{1}$$

Verification of solutions

$$\begin{aligned}
 y &= c_1 x \left(1 - \frac{x}{3} + \frac{x^2}{27} - \frac{x^3}{486} + \frac{x^4}{14580} - \frac{7291x^5}{656100} + \frac{225991x^6}{41334300} - \frac{2522341x^7}{3472081200} + O(x^8) \right) \\
 &\quad + c_2 \left(-\frac{2x \left(1 - \frac{x}{3} + \frac{x^2}{27} - \frac{x^3}{486} + \frac{x^4}{14580} - \frac{7291x^5}{656100} + \frac{225991x^6}{41334300} - \frac{2522341x^7}{3472081200} + O(x^8) \right) \ln(x)}{3} \right. \\
 &\quad \left. + 1 - \frac{x^2}{3} + \frac{14x^3}{243} - \frac{35x^4}{8748} + \frac{101x^5}{656100} + \frac{69199x^6}{14762250} + \frac{19882543x^7}{4340101500} + O(x^8) \right)
 \end{aligned}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
  -> Bessel
  -> elliptic
  -> Legendre
  -> Kummer
    -> hyper3: Equivalence to 1F1 under a power @ Moebius
  -> hypergeometric
    -> heuristic approach
    -> hyper3: Equivalence to 2F1, 1F1 or 0F1 under a power @ Moebius
  -> Mathieu
    -> Equivalence to the rational form of Mathieu ODE under a power @ Moebius
trying a solution in terms of MeijerG functions
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form  $r_0(x) * Y + r_1(x) * Y$  where  $Y = \exp(\int(r(x), dx)) * 2F1([a$ 
  trying a symmetry of the form [xi=0, eta=F(x)]
  trying differential order: 2; exact nonlinear
  trying symmetries linear in x and y(x)
  trying to convert to a linear ODE with constant coefficients
  trying 2nd order, integrating factor of the form  $\mu(x,y)$ 
  -> Trying a solution in terms of special functions:
    -> Bessel
    -> elliptic
    -> Legendre
    -> Kummer
      -> hyper3: Equivalence to 1F1 under a power @ Moebius
    -> hypergeometric
      -> heuristic approach
      -> hyper3: Equivalence to 2F1, 1F1 or 0F1 under a power @ Moebius
    -> Mathieu
      -> Equivalence to the rational form of Mathieu ODE under a power @ Moebius
  trying 2nd order exact linear
  trying symmetries linear in x and y(x)
  trying to convert to a linear ODE with constant coefficients
  trying to convert to an ODE of Bessel type
trying to convert to an ODE of Bessel type
-> trying reduction of order to Bessel
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 70

Order:=8;

dsolve(3*x^2*diff(y(x),x\$2)+x^6*diff(y(x),x)+2*x*y(x)=0,y(x),type='series',x=0);

$$y(x) = c_1 x \left(1 - \frac{1}{3}x + \frac{1}{27}x^2 - \frac{1}{486}x^3 + \frac{1}{14580}x^4 - \frac{7291}{656100}x^5 + \frac{225991}{41334300}x^6 - \frac{2522341}{3472081200}x^7 + O(x^8) \right) + c_2 \left(\ln(x) \left(-\frac{2}{3}x + \frac{2}{9}x^2 - \frac{2}{81}x^3 + \frac{1}{729}x^4 - \frac{1}{21870}x^5 + \frac{7291}{984150}x^6 - \frac{225991}{62001450}x^7 + O(x^8) \right) + \left(1 - \frac{1}{3}x^2 + \frac{14}{243}x^3 - \frac{35}{8748}x^4 + \frac{101}{656100}x^5 + \frac{69199}{14762250}x^6 + \frac{19882543}{4340101500}x^7 + O(x^8) \right) \right)$$

✓ Solution by Mathematica

Time used: 0.044 (sec). Leaf size: 121

AsymptoticDSolveValue[3*x^2*y'[x]+x^6*y'[x]+2*x*y[x]==0,y[x],{x,0,7}]

$$y(x) \rightarrow c_1 \left(\frac{x(7291x^5 - 45x^4 + 1350x^3 - 24300x^2 + 218700x - 656100) \log(x)}{984150} + \frac{-80332x^6 + 5895x^5 - 158625x^4 + 2430000x^3 - 16402500x^2 + 19683000x + 29524500}{29524500} \right) + c_2 \left(\frac{225991x^7}{41334300} - \frac{7291x^6}{656100} + \frac{x^5}{14580} - \frac{x^4}{486} + \frac{x^3}{27} - \frac{x^2}{3} + x \right)$$

17.3 problem 1(c)

Internal problem ID [6042]

Internal file name [OUTPUT/5290_Sunday_June_05_2022_03_29_41_PM_43044529/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 154

Problem number: 1(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Irregular singular point"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

Unable to solve or complete the solution.

$$x^2y'' - 5y' + 3yx^2 = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2y'' - 5y' + 3yx^2 = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{5}{x^2}$$
$$q(x) = 3$$

Table 210: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{5}{x^2}$	
singularity	type
$x = 0$	“irregular”

$q(x) = 3$	
singularity	type

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : []

Irregular singular points : $[0, \infty]$

Since $x = 0$ is not an ordinary point, then we will now check if it is a regular singular point. Unable to solve since $x = 0$ is not regular singular point. Terminating.

Verification of solutions N/A

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
  -> Bessel
  -> elliptic
  -> Legendre
  -> Kummer
    -> hyper3: Equivalence to 1F1 under a power @ Moebius
  -> hypergeometric
    -> heuristic approach
    -> hyper3: Equivalence to 2F1, 1F1 or 0F1 under a power @ Moebius
  -> Mathieu
    -> Equivalence to the rational form of Mathieu ODE under a power @ Moebius
trying a solution in terms of MeijerG functions
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
<- Heun successful: received ODE is equivalent to the HeunD ODE, case c = 0`
```

X Solution by Maple

```
Order:=8;
dsolve(x^2*diff(y(x),x$2)-5*diff(y(x),x)+3*x^2*y(x)=0,y(x),type='series',x=0);
```

No solution found

✓ Solution by Mathematica

Time used: 0.033 (sec). Leaf size: 106

```
AsymptoticDSolveValue[x^2*y''[x]-5*y'[x]+3*x^2*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(\frac{339x^7}{8750} + \frac{49x^6}{1250} + \frac{18x^5}{625} + \frac{3x^4}{50} + \frac{x^3}{5} + 1 \right) + c_2 e^{-5/x} \left(-\frac{302083x^7}{218750} + \frac{5243x^6}{6250} - \frac{357x^5}{625} + \frac{113x^4}{250} - \frac{49x^3}{125} + \frac{6x^2}{25} - \frac{2x}{5} + 1 \right) x^2$$

17.4 problem 1(d)

17.4.1 Maple step by step solution 1382

Internal problem ID [6043]

Internal file name [OUTPUT/5291_Sunday_June_05_2022_03_29_43_PM_77397404/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 154

Problem number: 1(d).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference is integer"**

Maple gives the following as the ode type

[[_Emden , _Fowler]]

$$y''x + 4y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$y''x + 4y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = 0$$

$$q(x) = \frac{4}{x}$$

Table 211: Table $p(x), q(x)$ singularities.

$p(x) = 0$	
singularity	type

$q(x) = \frac{4}{x}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$y''x + 4y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) x + 4 \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} x^{n+r-1} a_n (n+r)(n+r-1) \right) + \left(\sum_{n=0}^{\infty} 4a_n x^{n+r} \right) = 0 \quad (2A)$$

The next step is to make all powers of x be $n + r - 1$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r-1} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} 4a_n x^{n+r} = \sum_{n=1}^{\infty} 4a_{n-1} x^{n+r-1}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n + r - 1$.

$$\left(\sum_{n=0}^{\infty} x^{n+r-1} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=1}^{\infty} 4a_{n-1} x^{n+r-1} \right) = 0 \quad (2B)$$

The indicial equation is obtained from $n = 0$. From Eq (2B) this gives

$$x^{n+r-1} a_n (n+r) (n+r-1) = 0$$

When $n = 0$ the above becomes

$$x^{-1+r} a_0 r (-1+r) = 0$$

Or

$$x^{-1+r} a_0 r (-1+r) = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$x^{-1+r} r (-1+r) = 0$$

Since the above is true for all x then the indicial equation becomes

$$r(-1+r) = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = 1$$

$$r_2 = 0$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$x^{-1+r} r (-1+r) = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = 1$ is an integer, then we can construct two linearly independent solutions

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = C y_1(x) \ln(x) + x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = x \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+1}$$

$$y_2(x) = C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Where C above can be zero. We start by finding y_1 . Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) + 4a_{n-1} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{4a_{n-1}}{(n+r)(n+r-1)} \quad (4)$$

Which for the root $r = 1$ becomes

$$a_n = -\frac{4a_{n-1}}{(n+1)n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 1$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = -\frac{4}{(1+r)r}$$

Which for the root $r = 1$ becomes

$$a_1 = -2$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{4}{(1+r)r}$	-2

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{16}{r(1+r)^2(2+r)}$$

Which for the root $r = 1$ becomes

$$a_2 = \frac{4}{3}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{4}{(1+r)r}$	-2
a_2	$\frac{16}{r(1+r)^2(2+r)}$	$\frac{4}{3}$

For $n = 3$, using the above recursive equation gives

$$a_3 = -\frac{64}{r(1+r)^2(2+r)^2(3+r)}$$

Which for the root $r = 1$ becomes

$$a_3 = -\frac{4}{9}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{4}{(1+r)r}$	-2
a_2	$\frac{16}{r(1+r)^2(2+r)}$	$\frac{4}{3}$
a_3	$-\frac{64}{r(1+r)^2(2+r)^2(3+r)}$	$-\frac{4}{9}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{256}{r(1+r)^2(2+r)^2(3+r)^2(4+r)}$$

Which for the root $r = 1$ becomes

$$a_4 = \frac{4}{45}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{4}{(1+r)r}$	-2
a_2	$\frac{16}{r(1+r)^2(2+r)}$	$\frac{4}{3}$
a_3	$-\frac{64}{r(1+r)^2(2+r)^2(3+r)}$	$-\frac{4}{9}$
a_4	$\frac{256}{r(1+r)^2(2+r)^2(3+r)^2(4+r)}$	$\frac{4}{45}$

For $n = 5$, using the above recursive equation gives

$$a_5 = -\frac{1024}{r(1+r)^2(2+r)^2(3+r)^2(4+r)^2(5+r)}$$

Which for the root $r = 1$ becomes

$$a_5 = -\frac{8}{675}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{4}{(1+r)r}$	-2
a_2	$\frac{16}{r(1+r)^2(2+r)}$	$\frac{4}{3}$
a_3	$-\frac{64}{r(1+r)^2(2+r)^2(3+r)}$	$-\frac{4}{9}$
a_4	$\frac{256}{r(1+r)^2(2+r)^2(3+r)^2(4+r)}$	$\frac{4}{45}$
a_5	$-\frac{1024}{r(1+r)^2(2+r)^2(3+r)^2(4+r)^2(5+r)}$	$-\frac{8}{675}$

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{4096}{r(1+r)^2(2+r)^2(3+r)^2(4+r)^2(5+r)^2(6+r)}$$

Which for the root $r = 1$ becomes

$$a_6 = \frac{16}{14175}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{4}{(1+r)r}$	-2
a_2	$\frac{16}{r(1+r)^2(2+r)}$	$\frac{4}{3}$
a_3	$-\frac{64}{r(1+r)^2(2+r)^2(3+r)}$	$-\frac{4}{9}$
a_4	$\frac{256}{r(1+r)^2(2+r)^2(3+r)^2(4+r)}$	$\frac{4}{45}$
a_5	$-\frac{1024}{r(1+r)^2(2+r)^2(3+r)^2(4+r)^2(5+r)}$	$-\frac{8}{675}$
a_6	$\frac{4096}{r(1+r)^2(2+r)^2(3+r)^2(4+r)^2(5+r)^2(6+r)}$	$\frac{16}{14175}$

For $n = 7$, using the above recursive equation gives

$$a_7 = -\frac{16384}{r(1+r)^2(2+r)^2(3+r)^2(4+r)^2(5+r)^2(6+r)^2(7+r)}$$

Which for the root $r = 1$ becomes

$$a_7 = -\frac{8}{99225}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{4}{(1+r)r}$	-2
a_2	$\frac{16}{r(1+r)^2(2+r)}$	$\frac{4}{3}$
a_3	$-\frac{64}{r(1+r)^2(2+r)^2(3+r)}$	$-\frac{4}{9}$
a_4	$\frac{256}{r(1+r)^2(2+r)^2(3+r)^2(4+r)}$	$\frac{4}{45}$
a_5	$-\frac{1024}{r(1+r)^2(2+r)^2(3+r)^2(4+r)^2(5+r)}$	$-\frac{8}{675}$
a_6	$\frac{4096}{r(1+r)^2(2+r)^2(3+r)^2(4+r)^2(5+r)^2(6+r)}$	$\frac{16}{14175}$
a_7	$-\frac{16384}{r(1+r)^2(2+r)^2(3+r)^2(4+r)^2(5+r)^2(6+r)^2(7+r)}$	$-\frac{8}{99225}$

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned}
 y_1(x) &= x(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots) \\
 &= x\left(1 - 2x + \frac{4x^2}{3} - \frac{4x^3}{9} + \frac{4x^4}{45} - \frac{8x^5}{675} + \frac{16x^6}{14175} - \frac{8x^7}{99225} + O(x^8)\right)
 \end{aligned}$$

Now the second solution $y_2(x)$ is found. Let

$$r_1 - r_2 = N$$

Where N is positive integer which is the difference between the two roots. r_1 is taken as the larger root. Hence for this problem we have $N = 1$. Now we need to determine if C is zero or not. This is done by finding $\lim_{r \rightarrow r_2} a_1(r)$. If this limit exists, then $C = 0$, else we need to keep the log term and $C \neq 0$. The above table shows that

$$\begin{aligned}
 a_N &= a_1 \\
 &= -\frac{4}{(1+r)r}
 \end{aligned}$$

Therefore

$$\begin{aligned}
 \lim_{r \rightarrow r_2} -\frac{4}{(1+r)r} &= \lim_{r \rightarrow 0} -\frac{4}{(1+r)r} \\
 &= \text{undefined}
 \end{aligned}$$

Since the limit does not exist then the log term is needed. Therefore the second solution has the form

$$y_2(x) = Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right)$$

Therefore

$$\begin{aligned} \frac{d}{dx} y_2(x) &= Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2} (n+r_2)}{x} \right) \\ &= Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} x^{-1+n+r_2} b_n (n+r_2) \right) \\ \frac{d^2}{dx^2} y_2(x) &= Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} \\ &\quad + \sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2} (n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2} (n+r_2)}{x^2} \right) \\ &= Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} + \left(\sum_{n=0}^{\infty} x^{-2+n+r_2} b_n (n+r_2) (-1+n+r_2) \right) \end{aligned}$$

Substituting these back into the given ode $y''x + 4y = 0$ gives

$$\begin{aligned} &\left(Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} \right. \\ &\quad \left. + \sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2} (n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2} (n+r_2)}{x^2} \right) \right) x \\ &\quad + 4Cy_1(x) \ln(x) + 4 \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) = 0 \end{aligned}$$

Which can be written as

$$\begin{aligned} &\left((y_1''(x)x + 4y_1(x)) \ln(x) + \left(\frac{2y_1'(x)}{x} - \frac{y_1(x)}{x^2} \right) x \right) C \\ &\quad + \left(\sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2} (n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2} (n+r_2)}{x^2} \right) \right) x + 4 \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) = 0 \end{aligned} \tag{7}$$

But since $y_1(x)$ is a solution to the ode, then

$$y_1''(x)x + 4y_1(x) = 0$$

Eq (7) simplifies to

$$\begin{aligned} & \left(\frac{2y_1'(x)}{x} - \frac{y_1(x)}{x^2} \right) xC + \left(\sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2} (n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2} (n+r_2)}{x^2} \right) \right) x \\ & + 4 \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) = 0 \end{aligned} \quad (8)$$

Substituting $y_1 = \sum_{n=0}^{\infty} a_n x^{n+r_1}$ into the above gives

$$\begin{aligned} & \frac{\left(2 \left(\sum_{n=0}^{\infty} x^{-1+n+r_1} a_n (n+r_1) \right) x - \left(\sum_{n=0}^{\infty} a_n x^{n+r_1} \right) \right) C}{x} \\ & + \frac{\left(\sum_{n=0}^{\infty} x^{-2+n+r_2} b_n (n+r_2) (-1+n+r_2) \right) x^2 + 4 \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) x}{x} = 0 \end{aligned} \quad (9)$$

Since $r_1 = 1$ and $r_2 = 0$ then the above becomes

$$\begin{aligned} & \frac{\left(2 \left(\sum_{n=0}^{\infty} x^n a_n (n+1) \right) x - \left(\sum_{n=0}^{\infty} a_n x^{n+1} \right) \right) C}{x} \\ & + \frac{\left(\sum_{n=0}^{\infty} x^{-2+n} b_n n (n-1) \right) x^2 + 4 \left(\sum_{n=0}^{\infty} b_n x^n \right) x}{x} = 0 \end{aligned} \quad (10)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} 2C x^n a_n (n+1) \right) + \sum_{n=0}^{\infty} (-C x^n a_n) + \left(\sum_{n=0}^{\infty} n x^{n-1} b_n (n-1) \right) + \left(\sum_{n=0}^{\infty} 4b_n x^n \right) = 0 \quad (2A)$$

The next step is to make all powers of x be $n-1$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n-1} and

adjusting the power and the corresponding index gives

$$\begin{aligned}\sum_{n=0}^{\infty} 2C x^n a_n (n+1) &= \sum_{n=1}^{\infty} 2C a_{n-1} n x^{n-1} \\ \sum_{n=0}^{\infty} (-C x^n a_n) &= \sum_{n=1}^{\infty} (-C a_{n-1} x^{n-1}) \\ \sum_{n=0}^{\infty} 4b_n x^n &= \sum_{n=1}^{\infty} 4b_{n-1} x^{n-1}\end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n - 1$.

$$\begin{aligned}\left(\sum_{n=1}^{\infty} 2C a_{n-1} n x^{n-1}\right) + \sum_{n=1}^{\infty} (-C a_{n-1} x^{n-1}) \\ + \left(\sum_{n=0}^{\infty} n x^{n-1} b_n (n-1)\right) + \left(\sum_{n=1}^{\infty} 4b_{n-1} x^{n-1}\right) = 0\end{aligned}\tag{2B}$$

For $n = 0$ in Eq. (2B), we choose arbitrary value for b_0 as $b_0 = 1$. For $n = N$, where $N = 1$ which is the difference between the two roots, we are free to choose $b_1 = 0$. Hence for $n = 1$, Eq (2B) gives

$$C + 4 = 0$$

Which is solved for C . Solving for C gives

$$C = -4$$

For $n = 2$, Eq (2B) gives

$$3C a_1 + 4b_1 + 2b_2 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$2b_2 + 24 = 0$$

Solving the above for b_2 gives

$$b_2 = -12$$

For $n = 3$, Eq (2B) gives

$$5C a_2 + 4b_2 + 6b_3 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$6b_3 - \frac{224}{3} = 0$$

Solving the above for b_3 gives

$$b_3 = \frac{112}{9}$$

For $n = 4$, Eq (2B) gives

$$7Ca_3 + 4b_3 + 12b_4 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$12b_4 + \frac{560}{9} = 0$$

Solving the above for b_4 gives

$$b_4 = -\frac{140}{27}$$

For $n = 5$, Eq (2B) gives

$$9Ca_4 + 4b_4 + 20b_5 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$20b_5 - \frac{3232}{135} = 0$$

Solving the above for b_5 gives

$$b_5 = \frac{808}{675}$$

For $n = 6$, Eq (2B) gives

$$11Ca_5 + 4b_5 + 30b_6 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$30b_6 + \frac{3584}{675} = 0$$

Solving the above for b_6 gives

$$b_6 = -\frac{1792}{10125}$$

For $n = 7$, Eq (2B) gives

$$13Ca_6 + 4b_6 + 42b_7 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$42b_7 - \frac{18112}{23625} = 0$$

Solving the above for b_7 gives

$$b_7 = \frac{9056}{496125}$$

Now that we found all b_n and C , we can calculate the second solution from

$$y_2(x) = Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right)$$

Using the above value found for $C = -4$ and all b_n , then the second solution becomes

$$\begin{aligned} y_2(x) = & (-4) \left(x \left(1 - 2x + \frac{4x^2}{3} - \frac{4x^3}{9} + \frac{4x^4}{45} - \frac{8x^5}{675} + \frac{16x^6}{14175} - \frac{8x^7}{99225} + O(x^8) \right) \right) \ln(x) \\ & + 1 - 12x^2 + \frac{112x^3}{9} - \frac{140x^4}{27} + \frac{808x^5}{675} - \frac{1792x^6}{10125} + \frac{9056x^7}{496125} + O(x^8) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) = & c_1 y_1(x) + c_2 y_2(x) \\ = & c_1 x \left(1 - 2x + \frac{4x^2}{3} - \frac{4x^3}{9} + \frac{4x^4}{45} - \frac{8x^5}{675} + \frac{16x^6}{14175} - \frac{8x^7}{99225} + O(x^8) \right) \\ & + c_2 \left((-4) \left(x \left(1 - 2x + \frac{4x^2}{3} - \frac{4x^3}{9} + \frac{4x^4}{45} - \frac{8x^5}{675} + \frac{16x^6}{14175} - \frac{8x^7}{99225} + O(x^8) \right) \right) \ln(x) \right. \\ & \left. + 1 - 12x^2 + \frac{112x^3}{9} - \frac{140x^4}{27} + \frac{808x^5}{675} - \frac{1792x^6}{10125} + \frac{9056x^7}{496125} + O(x^8) \right) \end{aligned}$$

Hence the final solution is

$$\begin{aligned} y = & y_h \\ = & c_1 x \left(1 - 2x + \frac{4x^2}{3} - \frac{4x^3}{9} + \frac{4x^4}{45} - \frac{8x^5}{675} + \frac{16x^6}{14175} - \frac{8x^7}{99225} + O(x^8) \right) \\ & + c_2 \left(-4x \left(1 - 2x + \frac{4x^2}{3} - \frac{4x^3}{9} + \frac{4x^4}{45} - \frac{8x^5}{675} + \frac{16x^6}{14175} - \frac{8x^7}{99225} + O(x^8) \right) \ln(x) \right. \\ & \left. + 1 - 12x^2 + \frac{112x^3}{9} - \frac{140x^4}{27} + \frac{808x^5}{675} - \frac{1792x^6}{10125} + \frac{9056x^7}{496125} + O(x^8) \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 x \left(1 - 2x + \frac{4x^2}{3} - \frac{4x^3}{9} + \frac{4x^4}{45} - \frac{8x^5}{675} + \frac{16x^6}{14175} - \frac{8x^7}{99225} + O(x^8) \right) \\ + c_2 \left(-4x \left(1 - 2x + \frac{4x^2}{3} - \frac{4x^3}{9} + \frac{4x^4}{45} - \frac{8x^5}{675} + \frac{16x^6}{14175} - \frac{8x^7}{99225} + O(x^8) \right) \ln(x) \right. \\ \left. + 1 - 12x^2 + \frac{112x^3}{9} - \frac{140x^4}{27} + \frac{808x^5}{675} - \frac{1792x^6}{10125} + \frac{9056x^7}{496125} + O(x^8) \right)$$

Verification of solutions

$$y = c_1 x \left(1 - 2x + \frac{4x^2}{3} - \frac{4x^3}{9} + \frac{4x^4}{45} - \frac{8x^5}{675} + \frac{16x^6}{14175} - \frac{8x^7}{99225} + O(x^8) \right) \\ + c_2 \left(-4x \left(1 - 2x + \frac{4x^2}{3} - \frac{4x^3}{9} + \frac{4x^4}{45} - \frac{8x^5}{675} + \frac{16x^6}{14175} - \frac{8x^7}{99225} + O(x^8) \right) \ln(x) \right. \\ \left. + 1 - 12x^2 + \frac{112x^3}{9} - \frac{140x^4}{27} + \frac{808x^5}{675} - \frac{1792x^6}{10125} + \frac{9056x^7}{496125} + O(x^8) \right)$$

Verified OK.

17.4.1 Maple step by step solution

Let's solve

$$y''x + 4y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{4y}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{4y}{x} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$[P_2(x) = 0, P_3(x) = \frac{4}{x}]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 0$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 0$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$y''x + 4y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x \cdot y''$ to series expansion

$$x \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r-1}$$

- Shift index using $k \rightarrow k+1$

$$x \cdot y'' = \sum_{k=-1}^{\infty} a_{k+1} (k+1+r)(k+r) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0 r(-1+r) x^{-1+r} + \left(\sum_{k=0}^{\infty} (a_{k+1} (k+1+r)(k+r) + 4a_k) x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$r(-1+r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \{0, 1\}$$

- Each term in the series must be 0, giving the recursion relation

$$a_{k+1} (k+1+r)(k+r) + 4a_k = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+1} = -\frac{4a_k}{(k+1+r)(k+r)}$$

- Recursion relation for $r = 0$

$$a_{k+1} = -\frac{4a_k}{(k+1)k}$$

- Solution for $r = 0$

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+1} = -\frac{4a_k}{(k+1)k} \right]$$

- Recursion relation for $r = 1$

$$a_{k+1} = -\frac{4a_k}{(k+2)(k+1)}$$

- Solution for $r = 1$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+1}, a_{k+1} = -\frac{4a_k}{(k+2)(k+1)} \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^k \right) + \left(\sum_{k=0}^{\infty} b_k x^{k+1} \right), a_{k+1} = -\frac{4a_k}{(k+1)k}, b_{k+1} = -\frac{4b_k}{(k+2)(k+1)} \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
    <- Bessel successful
<- special function solution successful`

```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 70

```
Order:=8;  
dsolve(x*diff(y(x),x$2)+4*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x \left(1 - 2x + \frac{4}{3}x^2 - \frac{4}{9}x^3 + \frac{4}{45}x^4 - \frac{8}{675}x^5 + \frac{16}{14175}x^6 - \frac{8}{99225}x^7 + O(x^8) \right) \\ + c_2 \left(\ln(x) \left((-4)x + 8x^2 - \frac{16}{3}x^3 + \frac{16}{9}x^4 - \frac{16}{45}x^5 + \frac{32}{675}x^6 - \frac{64}{14175}x^7 + O(x^8) \right) \right. \\ \left. + \left(1 - 12x^2 + \frac{112}{9}x^3 - \frac{140}{27}x^4 + \frac{808}{675}x^5 - \frac{1792}{10125}x^6 + \frac{9056}{496125}x^7 + O(x^8) \right) \right)$$

✓ Solution by Mathematica

Time used: 0.037 (sec). Leaf size: 119

```
AsymptoticDSolveValue[x*y''[x]+4*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(\frac{4}{675}x(8x^5 - 60x^4 + 300x^3 - 900x^2 + 1350x - 675) \log(x) \right. \\ \left. + \frac{-2272x^6 + 15720x^5 - 70500x^4 + 180000x^3 - 202500x^2 + 40500x + 10125}{10125} \right) \\ + c_2 \left(\frac{16x^7}{14175} - \frac{8x^6}{675} + \frac{4x^5}{45} - \frac{4x^4}{9} + \frac{4x^3}{3} - 2x^2 + x \right)$$

17.5 problem 1(e)

17.5.1 Maple step by step solution 1396

Internal problem ID [6044]

Internal file name [OUTPUT/5292_Sunday_June_05_2022_03_29_46_PM_1192623/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 154

Problem number: 1(e).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Repeated root"**

Maple gives the following as the ode type

[_Gegenbauer]

$$(-x^2 + 1)y'' - 2xy' + 2y = 0$$

With the expansion point for the power series method at $x = 1$.

The ode does not have its expansion point at $x = 0$, therefore to simplify the computation of power series expansion, change of variable is made on the independent variable to shift the initial conditions and the expansion point back to zero. The new ode is then solved more easily since the expansion point is now at zero. The solution converted back to the original independent variable. Let

$$t = x - 1$$

The ode is converted to be in terms of the new independent variable t . This results in

$$(-(t + 1)^2 + 1) \left(\frac{d^2}{dt^2} y(t) \right) - 2(t + 1) \left(\frac{d}{dt} y(t) \right) + 2y(t) = 0$$

With its expansion point and initial conditions now at $t = 0$. The transformed ODE is now solved. The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$(-t^2 - 2t) \left(\frac{d^2}{dt^2} y(t) \right) + (-2t - 2) \left(\frac{d}{dt} y(t) \right) + 2y(t) = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$\frac{d^2}{dt^2}y(t) + p(t)\frac{d}{dt}y(t) + q(t)y(t) = 0$$

Where

$$p(t) = \frac{2t + 2}{t(t + 2)}$$

$$q(t) = -\frac{2}{t(t + 2)}$$

Table 213: Table $p(t), q(t)$ singularities.

$p(t) = \frac{2t+2}{t(t+2)}$		$q(t) = -\frac{2}{t(t+2)}$	
singularity	type	singularity	type
$t = -2$	“regular”	$t = -2$	“regular”
$t = 0$	“regular”	$t = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[-2, 0, \infty]$

Irregular singular points : $[\]$

Since $t = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$-\left(\frac{d^2}{dt^2}y(t)\right)t(t + 2) + (-2t - 2)\left(\frac{d}{dt}y(t)\right) + 2y(t) = 0$$

Let the solution be represented as Frobenius power series of the form

$$y(t) = \sum_{n=0}^{\infty} a_n t^{n+r}$$

Then

$$\frac{d}{dt}y(t) = \sum_{n=0}^{\infty} (n + r) a_n t^{n+r-1}$$

$$\frac{d^2}{dt^2}y(t) = \sum_{n=0}^{\infty} (n + r)(n + r - 1) a_n t^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned}
& - \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n t^{n+r-2} \right) t(t+2) \\
& + (-2t-2) \left(\sum_{n=0}^{\infty} (n+r) a_n t^{n+r-1} \right) + 2 \left(\sum_{n=0}^{\infty} a_n t^{n+r} \right) = 0
\end{aligned} \tag{1}$$

Which simplifies to

$$\begin{aligned}
& \sum_{n=0}^{\infty} (-t^{n+r} a_n (n+r)(n+r-1)) + \sum_{n=0}^{\infty} (-2t^{n+r-1} a_n (n+r)(n+r-1)) \\
& + \sum_{n=0}^{\infty} (-2t^{n+r} a_n (n+r)) + \sum_{n=0}^{\infty} (-2(n+r) a_n t^{n+r-1}) + \left(\sum_{n=0}^{\infty} 2a_n t^{n+r} \right) = 0
\end{aligned} \tag{2A}$$

The next step is to make all powers of t be $n+r-1$ in each summation term. Going over each summation term above with power of t in it which is not already t^{n+r-1} and adjusting the power and the corresponding index gives

$$\begin{aligned}
\sum_{n=0}^{\infty} (-t^{n+r} a_n (n+r)(n+r-1)) &= \sum_{n=1}^{\infty} (-a_{n-1} (n+r-1)(n+r-2) t^{n+r-1}) \\
\sum_{n=0}^{\infty} (-2t^{n+r} a_n (n+r)) &= \sum_{n=1}^{\infty} (-2a_{n-1} (n+r-1) t^{n+r-1}) \\
\sum_{n=0}^{\infty} 2a_n t^{n+r} &= \sum_{n=1}^{\infty} 2a_{n-1} t^{n+r-1}
\end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of t are the same and equal to $n+r-1$.

$$\begin{aligned}
& \sum_{n=1}^{\infty} (-a_{n-1} (n+r-1)(n+r-2) t^{n+r-1}) \\
& + \sum_{n=0}^{\infty} (-2t^{n+r-1} a_n (n+r)(n+r-1)) + \sum_{n=1}^{\infty} (-2a_{n-1} (n+r-1) t^{n+r-1}) \\
& + \sum_{n=0}^{\infty} (-2(n+r) a_n t^{n+r-1}) + \left(\sum_{n=1}^{\infty} 2a_{n-1} t^{n+r-1} \right) = 0
\end{aligned} \tag{2B}$$

The indicial equation is obtained from $n = 0$. From Eq (2B) this gives

$$-2t^{n+r-1}a_n(n+r)(n+r-1) - 2(n+r)a_nt^{n+r-1} = 0$$

When $n = 0$ the above becomes

$$-2t^{-1+r}a_0r(-1+r) - 2ra_0t^{-1+r} = 0$$

Or

$$(-2t^{-1+r}r(-1+r) - 2rt^{-1+r})a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$-2t^{-1+r}r^2 = 0$$

Since the above is true for all t then the indicial equation becomes

$$-2r^2 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = 0$$

$$r_2 = 0$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$-2t^{-1+r}r^2 = 0$$

Solving for r gives the roots of the indicial equation as Since the root of the indicial equation is repeated, then we can construct two linearly independent solutions. The first solution has the form

$$y_1(t) = \sum_{n=0}^{\infty} a_n t^{n+r} \tag{1A}$$

Now the second solution y_2 is found using

$$y_2(t) = y_1(t) \ln(t) + \left(\sum_{n=1}^{\infty} b_n t^{n+r} \right) \tag{1B}$$

Then the general solution will be

$$y(t) = c_1 y_1(t) + c_2 y_2(t)$$

In Eq (1B) the sum starts from 1 and not zero. In Eq (1A), a_0 is never zero, and is arbitrary and is typically taken as $a_0 = 1$, and $\{c_1, c_2\}$ are two arbitray constants of

integration which can be found from initial conditions. We start by finding the first solution $y_1(t)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$\begin{aligned} & -a_{n-1}(n+r-1)(n+r-2) - 2a_n(n+r)(n+r-1) \\ & - 2a_{n-1}(n+r-1) - 2a_n(n+r) + 2a_{n-1} = 0 \end{aligned} \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{a_{n-1}(n^2 + 2nr + r^2 - n - r - 2)}{2(n^2 + 2nr + r^2)} \quad (4)$$

Which for the root $r = 0$ becomes

$$a_n = -\frac{a_{n-1}(n^2 - n - 2)}{2n^2} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = \frac{-r^2 - r + 2}{2(r+1)^2}$$

Which for the root $r = 0$ becomes

$$a_1 = 1$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-r^2 - r + 2}{2(r+1)^2}$	1

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{(r+3)r(-1+r)}{4(r+2)(r+1)^2}$$

Which for the root $r = 0$ becomes

$$a_2 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-r^2-r+2}{2(r+1)^2}$	1
a_2	$\frac{(r+3)r(-1+r)}{4(r+2)(r+1)^2}$	0

For $n = 3$, using the above recursive equation gives

$$a_3 = -\frac{(-1+r)r(r+4)}{8(r+3)(r+1)(r+2)}$$

Which for the root $r = 0$ becomes

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-r^2-r+2}{2(r+1)^2}$	1
a_2	$\frac{(r+3)r(-1+r)}{4(r+2)(r+1)^2}$	0
a_3	$-\frac{(-1+r)r(r+4)}{8(r+3)(r+1)(r+2)}$	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{r(-1+r)(r+5)}{16(r+4)(r+1)(r+3)}$$

Which for the root $r = 0$ becomes

$$a_4 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-r^2-r+2}{2(r+1)^2}$	1
a_2	$\frac{(r+3)r(-1+r)}{4(r+2)(r+1)^2}$	0
a_3	$-\frac{(-1+r)r(r+4)}{8(r+3)(r+1)(r+2)}$	0
a_4	$\frac{r(-1+r)(r+5)}{16(r+4)(r+1)(r+3)}$	0

For $n = 5$, using the above recursive equation gives

$$a_5 = -\frac{(-1+r)r(r+6)}{32(r+5)(r+1)(r+4)}$$

Which for the root $r = 0$ becomes

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-r^2-r+2}{2(r+1)^2}$	1
a_2	$\frac{(r+3)r(-1+r)}{4(r+2)(r+1)^2}$	0
a_3	$-\frac{(-1+r)r(r+4)}{8(r+3)(r+1)(r+2)}$	0
a_4	$\frac{r(-1+r)(r+5)}{16(r+4)(r+1)(r+3)}$	0
a_5	$-\frac{(-1+r)r(r+6)}{32(r+5)(r+1)(r+4)}$	0

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{r(-1+r)(r+7)}{64(r+6)(r+1)(r+5)}$$

Which for the root $r = 0$ becomes

$$a_6 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-r^2-r+2}{2(r+1)^2}$	1
a_2	$\frac{(r+3)r(-1+r)}{4(r+2)(r+1)^2}$	0
a_3	$-\frac{(-1+r)r(r+4)}{8(r+3)(r+1)(r+2)}$	0
a_4	$\frac{r(-1+r)(r+5)}{16(r+4)(r+1)(r+3)}$	0
a_5	$-\frac{(-1+r)r(r+6)}{32(r+5)(r+1)(r+4)}$	0
a_6	$\frac{r(-1+r)(r+7)}{64(r+6)(r+1)(r+5)}$	0

For $n = 7$, using the above recursive equation gives

$$a_7 = -\frac{(-1+r)r(r+8)}{128(r+7)(r+1)(r+6)}$$

Which for the root $r = 0$ becomes

$$a_7 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-r^2-r+2}{2(r+1)^2}$	1
a_2	$\frac{(r+3)r(-1+r)}{4(r+2)(r+1)^2}$	0
a_3	$-\frac{(-1+r)r(r+4)}{8(r+3)(r+1)(r+2)}$	0
a_4	$\frac{r(-1+r)(r+5)}{16(r+4)(r+1)(r+3)}$	0
a_5	$-\frac{(-1+r)r(r+6)}{32(r+5)(r+1)(r+4)}$	0
a_6	$\frac{r(-1+r)(r+7)}{64(r+6)(r+1)(r+5)}$	0
a_7	$-\frac{(-1+r)r(r+8)}{128(r+7)(r+1)(r+6)}$	0

Using the above table, then the first solution $y_1(t)$ becomes

$$\begin{aligned} y_1(t) &= a_0 + a_1t + a_2t^2 + a_3t^3 + a_4t^4 + a_5t^5 + a_6t^6 + a_7t^7 + a_8t^8 \dots \\ &= t + 1 + O(t^8) \end{aligned}$$

Now the second solution is found. The second solution is given by

$$y_2(t) = y_1(t) \ln(t) + \left(\sum_{n=1}^{\infty} b_n t^{n+r} \right)$$

Where b_n is found using

$$b_n = \frac{d}{dr} a_{n,r}$$

And the above is then evaluated at $r = 0$. The above table for $a_{n,r}$ is used for this purpose. Computing the derivatives gives the following table

n	$b_{n,r}$	a_n	$b_{n,r} = \frac{d}{dr} a_{n,r}$	$b_n(r = 0)$
b_0	1	1	N/A since b_n starts from 1	N/A
b_1	$\frac{-r^2-r+2}{2(r+1)^2}$	1	$\frac{-r-5}{2(r+1)^3}$	$-\frac{5}{2}$
b_2	$\frac{(r+3)r(-1+r)}{4(r+2)(r+1)^2}$	0	$\frac{r^3+7r^2+7r-3}{2(r+2)^2(r+1)^3}$	$-\frac{3}{8}$
b_3	$-\frac{(-1+r)r(r+4)}{8(r+3)(r+1)(r+2)}$	0	$\frac{3-\frac{75}{8}r^2-\frac{3}{8}r^4-\frac{15}{4}r^3-\frac{9}{2}r}{(r+3)^2(r+1)^2(r+2)^2}$	$\frac{1}{12}$
b_4	$\frac{r(-1+r)(r+5)}{16(r+4)(r+1)(r+3)}$	0	$\frac{r^4+12r^3+38r^2+24r-15}{4(r+4)^2(r+1)^2(r+3)^2}$	$-\frac{5}{192}$
b_5	$-\frac{(-1+r)r(r+6)}{32(r+5)(r+1)(r+4)}$	0	$\frac{-\frac{265}{32}r^2+\frac{15}{4}-\frac{5}{32}r^4-\frac{35}{16}r^3-\frac{25}{4}r}{(r+5)^2(r+1)^2(r+4)^2}$	$\frac{3}{320}$
b_6	$\frac{r(-1+r)(r+7)}{64(r+6)(r+1)(r+5)}$	0	$\frac{\frac{105}{16}r^2-\frac{105}{32}+\frac{3}{32}r^4+\frac{3}{2}r^3+\frac{45}{8}r}{(r+6)^2(r+1)^2(r+5)^2}$	$-\frac{7}{1920}$
b_7	$-\frac{(-1+r)r(r+8)}{128(r+7)(r+1)(r+6)}$	0	$\frac{-\frac{623}{128}r^2-\frac{7}{128}r^4-\frac{63}{64}r^3+\frac{21}{8}-\frac{147}{32}r}{(r+7)^2(r+1)^2(r+6)^2}$	$\frac{1}{672}$

The above table gives all values of b_n needed. Hence the second solution is

$$\begin{aligned} y_2(t) &= y_1(t) \ln(t) + b_0 + b_1 t + b_2 t^2 + b_3 t^3 + b_4 t^4 + b_5 t^5 + b_6 t^6 + b_7 t^7 + b_8 t^8 \dots \\ &= (t + 1 + O(t^8)) \ln(t) - \frac{5t}{2} - \frac{3t^2}{8} + \frac{t^3}{12} - \frac{5t^4}{192} + \frac{3t^5}{320} - \frac{7t^6}{1920} + \frac{t^7}{672} + O(t^8) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(t) &= c_1 y_1(t) + c_2 y_2(t) \\ &= c_1 (t + 1 + O(t^8)) \\ &\quad + c_2 \left((t + 1 + O(t^8)) \ln(t) - \frac{5t}{2} - \frac{3t^2}{8} + \frac{t^3}{12} - \frac{5t^4}{192} + \frac{3t^5}{320} - \frac{7t^6}{1920} + \frac{t^7}{672} + O(t^8) \right) \end{aligned}$$

Hence the final solution is

$$\begin{aligned}
 y(t) &= y_h \\
 &= c_1(t + 1 + O(t^8)) \\
 &\quad + c_2\left((t + 1 + O(t^8)) \ln(t) - \frac{5t}{2} - \frac{3t^2}{8} + \frac{t^3}{12} - \frac{5t^4}{192} + \frac{3t^5}{320} - \frac{7t^6}{1920} + \frac{t^7}{672} + O(t^8)\right)
 \end{aligned}$$

Replacing t in the above with the original independent variable x using $t = x - 1$ results in

$$\begin{aligned}
 y &= c_1(x + O((x - 1)^8)) + c_2\left((x + O((x - 1)^8)) \ln(x - 1) - \frac{5x}{2} + \frac{5}{2} - \frac{3(x - 1)^2}{8}\right. \\
 &\quad \left. + \frac{(x - 1)^3}{12} - \frac{5(x - 1)^4}{192} + \frac{3(x - 1)^5}{320} - \frac{7(x - 1)^6}{1920} + \frac{(x - 1)^7}{672} + O((x - 1)^8)\right)
 \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}
 y &= c_1(x + O((x - 1)^8)) + c_2\left((x + O((x - 1)^8)) \ln(x - 1) - \frac{5x}{2} + \frac{5}{2} - \frac{3(x - 1)^2}{8}\right. \\
 &\quad \left. + \frac{(x - 1)^3}{12} - \frac{5(x - 1)^4}{192} + \frac{3(x - 1)^5}{320} - \frac{7(x - 1)^6}{1920} + \frac{(x - 1)^7}{672} + O((x - 1)^8)\right) \quad (1)
 \end{aligned}$$

Verification of solutions

$$\begin{aligned}
 y &= c_1(x + O((x - 1)^8)) + c_2\left((x + O((x - 1)^8)) \ln(x - 1) - \frac{5x}{2} + \frac{5}{2} - \frac{3(x - 1)^2}{8}\right. \\
 &\quad \left. + \frac{(x - 1)^3}{12} - \frac{5(x - 1)^4}{192} + \frac{3(x - 1)^5}{320} - \frac{7(x - 1)^6}{1920} + \frac{(x - 1)^7}{672} + O((x - 1)^8)\right)
 \end{aligned}$$

Verified OK.

17.5.1 Maple step by step solution

Let's solve

$$(-x^2 + 1)y'' - 2xy' + 2y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{2xy'}{x^2-1} + \frac{2y}{x^2-1}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{2xy'}{x^2-1} - \frac{2y}{x^2-1} = 0$$

- Check to see if x_0 is a regular singular point

- Define functions

$$[P_2(x) = \frac{2x}{x^2-1}, P_3(x) = -\frac{2}{x^2-1}]$$

- $(1+x) \cdot P_2(x)$ is analytic at $x = -1$

$$((1+x) \cdot P_2(x)) \Big|_{x=-1} = 1$$

- $(1+x)^2 \cdot P_3(x)$ is analytic at $x = -1$

$$((1+x)^2 \cdot P_3(x)) \Big|_{x=-1} = 0$$

- $x = -1$ is a regular singular point

Check to see if x_0 is a regular singular point

$$x_0 = -1$$

- Multiply by denominators

$$(x^2 - 1)y'' + 2xy' - 2y = 0$$

- Change variables using $x = u - 1$ so that the regular singular point is at $u = 0$

$$(u^2 - 2u) \left(\frac{d^2}{du^2} y(u) \right) + (2u - 2) \left(\frac{d}{du} y(u) \right) - 2y(u) = 0$$

- Assume series solution for $y(u)$

$$y(u) = \sum_{k=0}^{\infty} a_k u^{k+r}$$

- Rewrite ODE with series expansions

- Convert $u^m \cdot \left(\frac{d}{du} y(u) \right)$ to series expansion for $m = 0..1$

$$u^m \cdot \left(\frac{d}{du} y(u) \right) = \sum_{k=0}^{\infty} a_k (k+r) u^{k+r-1+m}$$

- Shift index using $k \rightarrow k+1-m$

$$u^m \cdot \left(\frac{d}{du} y(u) \right) = \sum_{k=-1+m}^{\infty} a_{k+1-m} (k+1-m+r) u^{k+r}$$

- Convert $u^m \cdot \left(\frac{d^2}{du^2} y(u) \right)$ to series expansion for $m = 1.2$

$$u^m \cdot \left(\frac{d^2}{du^2} y(u) \right) = \sum_{k=0}^{\infty} a_k (k+r) (k+r-1) u^{k+r-2+m}$$

- Shift index using $k \rightarrow k+2-m$

$$u^m \cdot \left(\frac{d^2}{du^2} y(u) \right) = \sum_{k=-2+m}^{\infty} a_{k+2-m} (k+2-m+r) (k+1-m+r) u^{k+r}$$

Rewrite ODE with series expansions

$$-2a_0 r^2 u^{-1+r} + \left(\sum_{k=0}^{\infty} (-2a_{k+1} (k+1+r)^2 + a_k (k+r+2) (k+r-1)) u^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$-2r^2 = 0$$

- Values of r that satisfy the indicial equation

$$r = 0$$

- Each term in the series must be 0, giving the recursion relation

$$-2a_{k+1} (k+1)^2 + a_k (k+2) (k-1) = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+1} = \frac{a_k (k+2)(k-1)}{2(k+1)^2}$$

- Recursion relation for $r = 0$; series terminates at $k = 1$

$$a_{k+1} = \frac{a_k (k+2)(k-1)}{2(k+1)^2}$$

- Apply recursion relation for $k = 0$

$$a_1 = -a_0$$

- Terminating series solution of the ODE for $r = 0$. Use reduction of order to find the second li

$$y(u) = a_0 \cdot (-u + 1)$$

- Revert the change of variables $u = 1 + x$

$$[y = -a_0 x]$$

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacics algorithm  
  A Liouvillian solution exists  
  Reducible group (found an exponential solution)  
  Group is reducible, not completely reducible  
<- Kovacics algorithm successful`
```

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 49

```
Order:=8;  
dsolve((1-x^2)*diff(y(x),x$2)-2*x*diff(y(x),x)+2*y(x)=0,y(x),type='series',x=1);
```

$$y(x) = \left(-\frac{5}{2}(x-1) - \frac{3}{8}(x-1)^2 + \frac{1}{12}(x-1)^3 - \frac{5}{192}(x-1)^4 + \frac{3}{320}(x-1)^5 - \frac{7}{1920}(x-1)^6 + \frac{1}{672}(x-1)^7 + O((x-1)^8) \right) c_2 + (1 + (x-1) + O((x-1)^8)) (c_2 \ln(x-1) + c_1)$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 86

```
AsymptoticDSolveValue[(1-x^2)*y'[x]-2*x*y'[x]+2*y[x]==0,y[x],{x,1,7}]
```

$$y(x) \rightarrow c_1 x + c_2 \left(\frac{1}{672}(x-1)^7 - \frac{7(x-1)^6}{1920} + \frac{3}{320}(x-1)^5 - \frac{5}{192}(x-1)^4 + \frac{1}{12}(x-1)^3 - \frac{3}{8}(x-1)^2 - 2(x-1) + \frac{1-x}{2} + x \log(x-1) \right)$$

17.6 problem 1(f)

17.6.1 Maple step by step solution 1412

Internal problem ID [6045]

Internal file name [OUTPUT/5293_Sunday_June_05_2022_03_29_49_PM_53281409/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 154

Problem number: 1(f).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$(x^2 + x - 2)^2 y'' + 3(x + 2)y' + (x - 1)y = 0$$

With the expansion point for the power series method at $x = -2$.

The ode does not have its expansion point at $x = 0$, therefore to simplify the computation of power series expansion, change of variable is made on the independent variable to shift the initial conditions and the expansion point back to zero. The new ode is then solved more easily since the expansion point is now at zero. The solution converted back to the original independent variable. Let

$$t = x + 2$$

The ode is converted to be in terms of the new independent variable t . This results in

$$\left(\frac{d^2}{dt^2}y(t)\right) t^2(t - 3)^2 + 3t\left(\frac{d}{dt}y(t)\right) + (t - 3)y(t) = 0$$

With its expansion point and initial conditions now at $t = 0$. The transformed ODE is now solved. The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$(t^4 - 6t^3 + 9t^2) \left(\frac{d^2}{dt^2}y(t)\right) + 3t\left(\frac{d}{dt}y(t)\right) + (t - 3)y(t) = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$\frac{d^2}{dt^2}y(t) + p(t)\frac{d}{dt}y(t) + q(t)y(t) = 0$$

Where

$$p(t) = \frac{3}{t(t-3)^2}$$

$$q(t) = \frac{1}{(t-3)t^2}$$

Table 215: Table $p(t), q(t)$ singularities.

$p(t) = \frac{3}{t(t-3)^2}$		$q(t) = \frac{1}{(t-3)t^2}$	
singularity	type	singularity	type
$t = 0$	“regular”	$t = 0$	“regular”
$t = 3$	“irregular”	$t = 3$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0, \infty]$

Irregular singular points : $[3]$

Since $t = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$t^2(t^2 - 6t + 9) \left(\frac{d^2}{dt^2}y(t) \right) + 3t \left(\frac{d}{dt}y(t) \right) + (t - 3)y(t) = 0$$

Let the solution be represented as Frobenius power series of the form

$$y(t) = \sum_{n=0}^{\infty} a_n t^{n+r}$$

Then

$$\frac{d}{dt}y(t) = \sum_{n=0}^{\infty} (n+r) a_n t^{n+r-1}$$

$$\frac{d^2}{dt^2}y(t) = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n t^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned}
 & t^2(t^2 - 6t + 9) \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n t^{n+r-2} \right) \\
 & + 3t \left(\sum_{n=0}^{\infty} (n+r) a_n t^{n+r-1} \right) + (t-3) \left(\sum_{n=0}^{\infty} a_n t^{n+r} \right) = 0
 \end{aligned} \tag{1}$$

Which simplifies to

$$\begin{aligned}
 & \left(\sum_{n=0}^{\infty} t^{n+r+2} a_n (n+r)(n+r-1) \right) + \sum_{n=0}^{\infty} (-6t^{1+n+r} a_n (n+r)(n+r-1)) \\
 & + \left(\sum_{n=0}^{\infty} 9t^{n+r} a_n (n+r)(n+r-1) \right) + \left(\sum_{n=0}^{\infty} 3t^{n+r} a_n (n+r) \right) \\
 & + \left(\sum_{n=0}^{\infty} t^{1+n+r} a_n \right) + \sum_{n=0}^{\infty} (-3a_n t^{n+r}) = 0
 \end{aligned} \tag{2A}$$

The next step is to make all powers of t be $n+r$ in each summation term. Going over each summation term above with power of t in it which is not already t^{n+r} and adjusting the power and the corresponding index gives

$$\begin{aligned}
 \sum_{n=0}^{\infty} t^{n+r+2} a_n (n+r)(n+r-1) &= \sum_{n=2}^{\infty} a_{n-2} (n+r-2)(n-3+r) t^{n+r} \\
 \sum_{n=0}^{\infty} (-6t^{1+n+r} a_n (n+r)(n+r-1)) &= \sum_{n=1}^{\infty} (-6a_{n-1} (n+r-1)(n+r-2) t^{n+r}) \\
 \sum_{n=0}^{\infty} t^{1+n+r} a_n &= \sum_{n=1}^{\infty} a_{n-1} t^{n+r}
 \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers

of t are the same and equal to $n + r$.

$$\begin{aligned} & \left(\sum_{n=2}^{\infty} a_{n-2} (n+r-2) (n-3+r) t^{n+r} \right) \\ & + \sum_{n=1}^{\infty} (-6a_{n-1} (n+r-1) (n+r-2) t^{n+r}) + \left(\sum_{n=0}^{\infty} 9t^{n+r} a_n (n+r) (n+r-1) \right) \quad (2B) \\ & + \left(\sum_{n=0}^{\infty} 3t^{n+r} a_n (n+r) \right) + \left(\sum_{n=1}^{\infty} a_{n-1} t^{n+r} \right) + \sum_{n=0}^{\infty} (-3a_n t^{n+r}) = 0 \end{aligned}$$

The indicial equation is obtained from $n = 0$. From Eq (2B) this gives

$$9t^{n+r} a_n (n+r) (n+r-1) + 3t^{n+r} a_n (n+r) - 3a_n t^{n+r} = 0$$

When $n = 0$ the above becomes

$$9t^r a_0 r (-1+r) + 3t^r a_0 r - 3a_0 t^r = 0$$

Or

$$(9t^r r (-1+r) + 3t^r r - 3t^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(9r^2 - 6r - 3) t^r = 0$$

Since the above is true for all t then the indicial equation becomes

$$9r^2 - 6r - 3 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 1 \\ r_2 &= -\frac{1}{3} \end{aligned}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(9r^2 - 6r - 3) t^r = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = \frac{4}{3}$ is not an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(t) &= t^{r_1} \left(\sum_{n=0}^{\infty} a_n t^n \right) \\ y_2(t) &= t^{r_2} \left(\sum_{n=0}^{\infty} b_n t^n \right) \end{aligned}$$

Or

$$y_1(t) = \sum_{n=0}^{\infty} a_n t^{1+n}$$

$$y_2(t) = \sum_{n=0}^{\infty} b_n t^{n-\frac{1}{3}}$$

We start by finding $y_1(t)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = \frac{6r^2 - 6r - 1}{9r^2 + 12r}$$

For $2 \leq n$ the recursive equation is

$$a_{n-2}(n+r-2)(n-3+r) - 6a_{n-1}(n+r-1)(n+r-2) + 9a_n(n+r)(n+r-1) + 3a_n(n+r) + a_{n-1} - 3a_n = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{n^2 a_{n-2} - 6n^2 a_{n-1} + 2nra_{n-2} - 12nra_{n-1} + r^2 a_{n-2} - 6r^2 a_{n-1} - 5na_{n-2} + 18na_{n-1} - 5ra_{n-2} + 18ra_{n-1}}{3(3n^2 + 6nr + 3r^2 - 2n - 2r - 1)} \quad (4)$$

Which for the root $r = 1$ becomes

$$a_n = \frac{(-a_{n-2} + 6a_{n-1})n^2 + (3a_{n-2} - 6a_{n-1})n - 2a_{n-2} - a_{n-1}}{9n^2 + 12n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 1$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{6r^2-6r-1}{9r^2+12r}$	$-\frac{1}{21}$

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{27r^4 - 3r^3 - 36r^2 + 1}{81r^4 + 378r^3 + 549r^2 + 252r}$$

Which for the root $r = 1$ becomes

$$a_2 = -\frac{11}{1260}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{6r^2-6r-1}{9r^2+12r}$	$-\frac{1}{21}$
a_2	$\frac{27r^4-3r^3-36r^2+1}{81r^4+378r^3+549r^2+252r}$	$-\frac{11}{1260}$

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{108r^6 + 288r^5 - 36r^4 - 462r^3 - 213r^2 + 39r + 11}{27(3r^2 + 16r + 20)r(9r^3 + 42r^2 + 61r + 28)}$$

Which for the root $r = 1$ becomes

$$a_3 = -\frac{53}{29484}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{6r^2-6r-1}{9r^2+12r}$	$-\frac{1}{21}$
a_2	$\frac{27r^4-3r^3-36r^2+1}{81r^4+378r^3+549r^2+252r}$	$-\frac{11}{1260}$
a_3	$\frac{108r^6+288r^5-36r^4-462r^3-213r^2+39r+11}{27(3r^2+16r+20)r(9r^3+42r^2+61r+28)}$	$-\frac{53}{29484}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{405r^8 + 2970r^7 + 6759r^6 + 2142r^5 - 10827r^4 - 12105r^3 - 2121r^2 + 1419r + 265}{81(3r^2 + 16r + 20)r(9r^3 + 42r^2 + 61r + 28)(3r^2 + 22r + 39)}$$

Which for the root $r = 1$ becomes

$$a_4 = -\frac{11093}{28304640}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{6r^2-6r-1}{9r^2+12r}$	$-\frac{1}{21}$
a_2	$\frac{27r^4-3r^3-36r^2+1}{81r^4+378r^3+549r^2+252r}$	$-\frac{11}{1260}$
a_3	$\frac{108r^6+288r^5-36r^4-462r^3-213r^2+39r+11}{27(3r^2+16r+20)r(9r^3+42r^2+61r+28)}$	$-\frac{53}{29484}$
a_4	$\frac{405r^8+2970r^7+6759r^6+2142r^5-10827r^4-12105r^3-2121r^2+1419r+265}{81(3r^2+16r+20)r(9r^3+42r^2+61r+28)(3r^2+22r+39)}$	$-\frac{11093}{28304640}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{1458r^{10} + 20250r^9 + 108297r^8 + 265518r^7 + 217782r^6 - 287388r^5 - 709074r^4 - 427506r^3 + 16353r^2 + 73710r + 11093}{243(3r^2 + 16r + 20)r(9r^3 + 42r^2 + 61r + 28)(3r^2 + 22r + 39)(3r^2 + 28r + 64)}$$

Which for the root $r = 1$ becomes

$$a_5 = -\frac{709507}{8066822400}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{6r^2-6r-1}{9r^2+12r}$	$-\frac{1}{21}$
a_2	$\frac{27r^4-3r^3-36r^2+1}{81r^4+378r^3+549r^2+252r}$	$-\frac{11}{1260}$
a_3	$\frac{108r^6+288r^5-36r^4-462r^3-213r^2+39r+11}{27(3r^2+16r+20)r(9r^3+42r^2+61r+28)}$	$-\frac{53}{29484}$
a_4	$\frac{405r^8+2970r^7+6759r^6+2142r^5-10827r^4-12105r^3-2121r^2+1419r+265}{81(3r^2+16r+20)r(9r^3+42r^2+61r+28)(3r^2+22r+39)}$	$-\frac{11093}{28304640}$
a_5	$\frac{1458r^{10}+20250r^9+108297r^8+265518r^7+217782r^6-287388r^5-709074r^4-427506r^3+16353r^2+73710r+11093}{243(3r^2+16r+20)r(9r^3+42r^2+61r+28)(3r^2+22r+39)(3r^2+28r+64)}$	$-\frac{709507}{8066822400}$

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{5103r^{12} + 113967r^{11} + 1059723r^{10} + 5248125r^9 + 14394024r^8 + 18690912r^7 - 3055158r^6 - 434760r^5 + 1059723r^4 - 113967r^3 + 5103r^2 - 113967r + 1059723}{729(3r^2 + 16r + 20)r(9r^3 + 42r^2 + 61r + 28)(3r^2 + 22r + 39)(3r^2 + 28r + 64)}$$

Which for the root $r = 1$ becomes

$$a_6 = -\frac{5797423}{290405606400}$$

And the table now becomes

n	$a_{n,r}$
a_0	1
a_1	$\frac{6r^2-6r-1}{9r^2+12r}$
a_2	$\frac{27r^4-3r^3-36r^2+1}{81r^4+378r^3+549r^2+252r}$
a_3	$\frac{108r^6+288r^5-36r^4-462r^3-213r^2+39r+11}{27(3r^2+16r+20)r(9r^3+42r^2+61r+28)}$
a_4	$\frac{405r^8+2970r^7+6759r^6+2142r^5-10827r^4-12105r^3-2121r^2+1419r+265}{81(3r^2+16r+20)r(9r^3+42r^2+61r+28)(3r^2+22r+39)}$
a_5	$\frac{1458r^{10}+20250r^9+108297r^8+265518r^7+217782r^6-287388r^5-709074r^4-427506r^3+16353r^2+73710r+11093}{243(3r^2+16r+20)r(9r^3+42r^2+61r+28)(3r^2+22r+39)(3r^2+28r+64)}$
a_6	$\frac{5103r^{12}+113967r^{11}+1059723r^{10}+5248125r^9+14394024r^8+18690912r^7-3055158r^6-43476012r^5-52278174r^4-17968407r^3+7306881r^2}{729(3r^2+16r+20)r(9r^3+42r^2+61r+28)(3r^2+22r+39)(3r^2+28r+64)(3r^2+34r+95)}$

For $n = 7$, using the above recursive equation gives

$$a_7 = \frac{17496r^{14} + 571536r^{13} + 8096760r^{12} + 64902708r^{11} + 320544378r^{10} + 982953738r^9 + 1709577414r^8}{2187(3r^2 + 16r + 20)r(9r^3 + 42r^2 + 61r + 28)}$$

Which for the root $r = 1$ becomes

$$a_7 = -\frac{52991201}{11727918720000}$$

And the table now becomes

n	$a_{n,r}$
a_0	1
a_1	$\frac{6r^2-6r-1}{9r^2+12r}$
a_2	$\frac{27r^4-3r^3-36r^2+1}{81r^4+378r^3+549r^2+252r}$
a_3	$\frac{108r^6+288r^5-36r^4-462r^3-213r^2+39r+11}{27(3r^2+16r+20)r(9r^3+42r^2+61r+28)}$
a_4	$\frac{405r^8+2970r^7+6759r^6+2142r^5-10827r^4-12105r^3-2121r^2+1419r+265}{81(3r^2+16r+20)r(9r^3+42r^2+61r+28)(3r^2+22r+39)}$
a_5	$\frac{1458r^{10}+20250r^9+108297r^8+265518r^7+217782r^6-287388r^5-709074r^4-427506r^3+16353r^2+73710r+11093}{243(3r^2+16r+20)r(9r^3+42r^2+61r+28)(3r^2+22r+39)(3r^2+28r+64)}$
a_6	$\frac{5103r^{12}+113967r^{11}+1059723r^{10}+5248125r^9+14394024r^8+18690912r^7-3055158r^6-43476012r^5-52278174r^4-17968407r^3+7306881r^2}{729(3r^2+16r+20)r(9r^3+42r^2+61r+28)(3r^2+22r+39)(3r^2+28r+64)(3r^2+34r+95)}$
a_7	$\frac{17496r^{14}+571536r^{13}+8096760r^{12}+64902708r^{11}+320544378r^{10}+982953738r^9+1709577414r^8+897903738r^7-2589159015r^6-58493938r^5}{2187(3r^2+16r+20)r(9r^3+42r^2+61r+28)(3r^2+22r+39)(3r^2+28r+64)(3r^2+34r+95)}$

Using the above table, then the solution $y_1(t)$ is

$$y_1(t) = t(a_0 + a_1t + a_2t^2 + a_3t^3 + a_4t^4 + a_5t^5 + a_6t^6 + a_7t^7 + a_8t^8 \dots)$$

$$= t \left(1 - \frac{t}{21} - \frac{11t^2}{1260} - \frac{53t^3}{29484} - \frac{11093t^4}{28304640} - \frac{709507t^5}{8066822400} - \frac{5797423t^6}{290405606400} - \frac{52991201t^7}{11727918720000} + O(t^8) \right)$$

Now the second solution $y_2(t)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$b_1 = \frac{6r^2 - 6r - 1}{9r^2 + 12r}$$

For $2 \leq n$ the recursive equation is

$$b_{n-2}(n+r-2)(n-3+r) - 6b_{n-1}(n+r-1)(n+r-2) + 9b_n(n+r)(n+r-1) + 3b_n(n+r) + b_{n-1} - 3b_n = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = -\frac{n^2b_{n-2} - 6n^2b_{n-1} + 2nrb_{n-2} - 12nrb_{n-1} + r^2b_{n-2} - 6r^2b_{n-1} - 5nb_{n-2} + 18nb_{n-1} - 5rb_{n-2} + 18rb_{n-1}}{3(3n^2 + 6nr + 3r^2 - 2n - 2r - 1)} \quad (4)$$

Which for the root $r = -\frac{1}{3}$ becomes

$$b_n = \frac{(-9b_{n-2} + 54b_{n-1})n^2 + (51b_{n-2} - 198b_{n-1})n - 70b_{n-2} + 159b_{n-1}}{81n^2 - 108n} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = -\frac{1}{3}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{6r^2-6r-1}{9r^2+12r}$	$-\frac{5}{9}$

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{27r^4 - 3r^3 - 36r^2 + 1}{81r^4 + 378r^3 + 549r^2 + 252r}$$

Which for the root $r = -\frac{1}{3}$ becomes

$$b_2 = \frac{23}{324}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{6r^2-6r-1}{9r^2+12r}$	$-\frac{5}{9}$
b_2	$\frac{27r^4-3r^3-36r^2+1}{81r^4+378r^3+549r^2+252r}$	$\frac{23}{324}$

For $n = 3$, using the above recursive equation gives

$$b_3 = \frac{108r^6 + 288r^5 - 36r^4 - 462r^3 - 213r^2 + 39r + 11}{27(3r^2 + 16r + 20)r(9r^3 + 42r^2 + 61r + 28)}$$

Which for the root $r = -\frac{1}{3}$ becomes

$$b_3 = \frac{271}{43740}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{6r^2-6r-1}{9r^2+12r}$	$-\frac{5}{9}$
b_2	$\frac{27r^4-3r^3-36r^2+1}{81r^4+378r^3+549r^2+252r}$	$\frac{23}{324}$
b_3	$\frac{108r^6+288r^5-36r^4-462r^3-213r^2+39r+11}{27(3r^2+16r+20)r(9r^3+42r^2+61r+28)}$	$\frac{271}{43740}$

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{405r^8 + 2970r^7 + 6759r^6 + 2142r^5 - 10827r^4 - 12105r^3 - 2121r^2 + 1419r + 265}{81(3r^2 + 16r + 20)r(9r^3 + 42r^2 + 61r + 28)(3r^2 + 22r + 39)}$$

Which for the root $r = -\frac{1}{3}$ becomes

$$b_4 = \frac{10517}{12597120}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{6r^2-6r-1}{9r^2+12r}$	$-\frac{5}{9}$
b_2	$\frac{27r^4-3r^3-36r^2+1}{81r^4+378r^3+549r^2+252r}$	$\frac{23}{324}$
b_3	$\frac{108r^6+288r^5-36r^4-462r^3-213r^2+39r+11}{27(3r^2+16r+20)r(9r^3+42r^2+61r+28)}$	$\frac{271}{43740}$
b_4	$\frac{405r^8+2970r^7+6759r^6+2142r^5-10827r^4-12105r^3-2121r^2+1419r+265}{81(3r^2+16r+20)r(9r^3+42r^2+61r+28)(3r^2+22r+39)}$	$\frac{10517}{12597120}$

For $n = 5$, using the above recursive equation gives

$$b_5 = \frac{1458r^{10} + 20250r^9 + 108297r^8 + 265518r^7 + 217782r^6 - 287388r^5 - 709074r^4 - 427506r^3 + 16353r^2 + 73710r + 11093}{243(3r^2 + 16r + 20)r(9r^3 + 42r^2 + 61r + 28)(3r^2 + 22r + 39)(3r^2 + 28r + 64)}$$

Which for the root $r = -\frac{1}{3}$ becomes

$$b_5 = \frac{778801}{6235574400}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{6r^2-6r-1}{9r^2+12r}$	$-\frac{5}{9}$
b_2	$\frac{27r^4-3r^3-36r^2+1}{81r^4+378r^3+549r^2+252r}$	$\frac{23}{324}$
b_3	$\frac{108r^6+288r^5-36r^4-462r^3-213r^2+39r+11}{27(3r^2+16r+20)r(9r^3+42r^2+61r+28)}$	$\frac{271}{43740}$
b_4	$\frac{405r^8+2970r^7+6759r^6+2142r^5-10827r^4-12105r^3-2121r^2+1419r+265}{81(3r^2+16r+20)r(9r^3+42r^2+61r+28)(3r^2+22r+39)}$	$\frac{10517}{12597120}$
b_5	$\frac{1458r^{10}+20250r^9+108297r^8+265518r^7+217782r^6-287388r^5-709074r^4-427506r^3+16353r^2+73710r+11093}{243(3r^2+16r+20)r(9r^3+42r^2+61r+28)(3r^2+22r+39)(3r^2+28r+64)}$	$\frac{778801}{6235574400}$

For $n = 6$, using the above recursive equation gives

$$b_6 = \frac{5103r^{12} + 113967r^{11} + 1059723r^{10} + 5248125r^9 + 14394024r^8 + 18690912r^7 - 3055158r^6 - 43476012r^5 + 17968407r^4 + 7306881r^3 + 17968407r^2 - 43476012r + 5248125}{729(3r^2 + 16r + 20)r(9r^3 + 42r^2 + 61r + 28)(3r^2 + 22r + 39)(3r^2 + 28r + 64)(3r^2 + 34r + 95)}$$

Which for the root $r = -\frac{1}{3}$ becomes

$$b_6 = \frac{16965493}{942818849280}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{6r^2-6r-1}{9r^2+12r}$	$-\frac{5}{9}$
b_2	$\frac{27r^4-3r^3-36r^2+1}{81r^4+378r^3+549r^2+252r}$	$\frac{23}{324}$
b_3	$\frac{108r^6+288r^5-36r^4-462r^3-213r^2+39r+11}{27(3r^2+16r+20)r(9r^3+42r^2+61r+28)}$	$\frac{271}{43740}$
b_4	$\frac{405r^8+2970r^7+6759r^6+2142r^5-10827r^4-12105r^3-2121r^2+1419r+265}{81(3r^2+16r+20)r(9r^3+42r^2+61r+28)(3r^2+22r+39)}$	$\frac{10517}{12597120}$
b_5	$\frac{1458r^{10}+20250r^9+108297r^8+265518r^7+217782r^6-287388r^5-709074r^4-427506r^3+16353r^2+73710r+11093}{243(3r^2+16r+20)r(9r^3+42r^2+61r+28)(3r^2+22r+39)(3r^2+28r+64)}$	$\frac{778801}{6235574400}$
b_6	$\frac{5103r^{12}+113967r^{11}+1059723r^{10}+5248125r^9+14394024r^8+18690912r^7-3055158r^6-43476012r^5+17968407r^4+7306881r^3+17968407r^2-43476012r+5248125}{729(3r^2+16r+20)r(9r^3+42r^2+61r+28)(3r^2+22r+39)(3r^2+28r+64)(3r^2+34r+95)}$	$\frac{16965493}{942818849280}$

For $n = 7$, using the above recursive equation gives

$$b_7 = \frac{17496r^{14} + 571536r^{13} + 8096760r^{12} + 64902708r^{11} + 320544378r^{10} + 982953738r^9 + 1709577414r^8}{2187(3r^2 + 16r + 20)r(9r^3 + 42r^2 + 61r + 28)}$$

Which for the root $r = -\frac{1}{3}$ becomes

$$b_7 = \frac{899971067}{458981357990400}$$

And the table now becomes

n	$b_{n,r}$
b_0	1
b_1	$\frac{6r^2-6r-1}{9r^2+12r}$
b_2	$\frac{27r^4-3r^3-36r^2+1}{81r^4+378r^3+549r^2+252r}$
b_3	$\frac{108r^6+288r^5-36r^4-462r^3-213r^2+39r+11}{27(3r^2+16r+20)r(9r^3+42r^2+61r+28)}$
b_4	$\frac{405r^8+2970r^7+6759r^6+2142r^5-10827r^4-12105r^3-2121r^2+1419r+265}{81(3r^2+16r+20)r(9r^3+42r^2+61r+28)(3r^2+22r+39)}$
b_5	$\frac{1458r^{10}+20250r^9+108297r^8+265518r^7+217782r^6-287388r^5-709074r^4-427506r^3+16353r^2+73710r+11093}{243(3r^2+16r+20)r(9r^3+42r^2+61r+28)(3r^2+22r+39)(3r^2+28r+64)}$
b_6	$\frac{5103r^{12}+113967r^{11}+1059723r^{10}+5248125r^9+14394024r^8+18690912r^7-3055158r^6-43476012r^5-52278174r^4-17968407r^3+7306881r^2}{729(3r^2+16r+20)r(9r^3+42r^2+61r+28)(3r^2+22r+39)(3r^2+28r+64)(3r^2+34r+95)}$
b_7	$\frac{17496r^{14}+571536r^{13}+8096760r^{12}+64902708r^{11}+320544378r^{10}+982953738r^9+1709577414r^8+897903738r^7-2589159015r^6-58493938r^5}{2187(3r^2+16r+20)r(9r^3+42r^2+61r+28)(3r^2+22r+39)(3r^2+28r+64)(3r^2+34r+95)}$

Using the above table, then the solution $y_2(t)$ is

$$\begin{aligned} y_2(t) &= t(b_0 + b_1t + b_2t^2 + b_3t^3 + b_4t^4 + b_5t^5 + b_6t^6 + b_7t^7 + b_8t^8 \dots) \\ &= \frac{1 - \frac{5t}{9} + \frac{23t^2}{324} + \frac{271t^3}{43740} + \frac{10517t^4}{12597120} + \frac{778801t^5}{6235574400} + \frac{16965493t^6}{942818849280} + \frac{899971067t^7}{458981357990400} + O(t^8)}{t^{\frac{1}{3}}} \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(t) &= c_1y_1(t) + c_2y_2(t) \\ &= c_1t \left(1 - \frac{t}{21} - \frac{11t^2}{1260} - \frac{53t^3}{29484} - \frac{11093t^4}{28304640} - \frac{709507t^5}{8066822400} - \frac{5797423t^6}{290405606400} \right. \\ &\quad \left. - \frac{52991201t^7}{11727918720000} + O(t^8) \right) \\ &\quad + \frac{c_2 \left(1 - \frac{5t}{9} + \frac{23t^2}{324} + \frac{271t^3}{43740} + \frac{10517t^4}{12597120} + \frac{778801t^5}{6235574400} + \frac{16965493t^6}{942818849280} + \frac{899971067t^7}{458981357990400} + O(t^8) \right)}{t^{\frac{1}{3}}} \end{aligned}$$

Hence the final solution is

$$\begin{aligned}
 y(t) &= y_h \\
 &= c_1 t \left(1 - \frac{t}{21} - \frac{11t^2}{1260} - \frac{53t^3}{29484} - \frac{11093t^4}{28304640} - \frac{709507t^5}{8066822400} - \frac{5797423t^6}{290405606400} \right. \\
 &\quad \left. - \frac{52991201t^7}{11727918720000} + O(t^8) \right) \\
 &\quad + \frac{c_2 \left(1 - \frac{5t}{9} + \frac{23t^2}{324} + \frac{271t^3}{43740} + \frac{10517t^4}{12597120} + \frac{778801t^5}{6235574400} + \frac{16965493t^6}{942818849280} + \frac{899971067t^7}{458981357990400} + O(t^8) \right)}{t^{\frac{1}{3}}}
 \end{aligned}$$

Replacing t in the above with the original independent variable x using $t = x + 2$ results in

$$\begin{aligned}
 y &= c_1 (x + 2) \left(\frac{19}{21} - \frac{x}{21} - \frac{11(x + 2)^2}{1260} - \frac{53(x + 2)^3}{29484} - \frac{11093(x + 2)^4}{28304640} - \frac{709507(x + 2)^5}{8066822400} \right. \\
 &\quad \left. - \frac{5797423(x + 2)^6}{290405606400} - \frac{52991201(x + 2)^7}{11727918720000} + O((x + 2)^8) \right) \\
 &\quad + \frac{c_2 \left(-\frac{1}{9} - \frac{5x}{9} + \frac{23(x+2)^2}{324} + \frac{271(x+2)^3}{43740} + \frac{10517(x+2)^4}{12597120} + \frac{778801(x+2)^5}{6235574400} + \frac{16965493(x+2)^6}{942818849280} + \frac{899971067(x+2)^7}{458981357990400} + O((x + 2)^8) \right)}{(x + 2)^{\frac{1}{3}}}
 \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}
 y &= c_1 (x + 2) \left(\frac{19}{21} - \frac{x}{21} - \frac{11(x + 2)^2}{1260} - \frac{53(x + 2)^3}{29484} - \frac{11093(x + 2)^4}{28304640} - \frac{709507(x + 2)^5}{8066822400} \right. \\
 &\quad \left. - \frac{5797423(x + 2)^6}{290405606400} - \frac{52991201(x + 2)^7}{11727918720000} + O((x + 2)^8) \right) \\
 &\quad + \frac{c_2 \left(-\frac{1}{9} - \frac{5x}{9} + \frac{23(x+2)^2}{324} + \frac{271(x+2)^3}{43740} + \frac{10517(x+2)^4}{12597120} + \frac{778801(x+2)^5}{6235574400} + \frac{16965493(x+2)^6}{942818849280} + \frac{899971067(x+2)^7}{458981357990400} + O((x + 2)^8) \right)}{(x + 2)^{\frac{1}{3}}}
 \end{aligned}$$

Verification of solutions

$$y = c_1(x+2) \left(\frac{19}{21} - \frac{x}{21} - \frac{11(x+2)^2}{1260} - \frac{53(x+2)^3}{29484} - \frac{11093(x+2)^4}{28304640} - \frac{709507(x+2)^5}{8066822400} \right. \\ \left. - \frac{5797423(x+2)^6}{290405606400} - \frac{52991201(x+2)^7}{11727918720000} + O((x+2)^8) \right) \\ + \frac{c_2 \left(-\frac{1}{9} - \frac{5x}{9} + \frac{23(x+2)^2}{324} + \frac{271(x+2)^3}{43740} + \frac{10517(x+2)^4}{12597120} + \frac{778801(x+2)^5}{6235574400} + \frac{16965493(x+2)^6}{942818849280} + \frac{899971067(x+2)^7}{458981357990400} + O((x+2)^8) \right)}{(x+2)^{\frac{1}{3}}}$$

Verified OK.

17.6.1 Maple step by step solution

Let's solve

$$y''(x+2)^2(x-1)^2 + (3x+6)y' + (x-1)y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{3y'}{(x+2)(x-1)^2} - \frac{y}{(x-1)(x+2)^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{3y'}{(x+2)(x-1)^2} + \frac{y}{(x-1)(x+2)^2} = 0$$

- Check to see if x_0 is a regular singular point

- Define functions

$$\left[P_2(x) = \frac{3}{(x+2)(x-1)^2}, P_3(x) = \frac{1}{(x-1)(x+2)^2} \right]$$

- $(x+2) \cdot P_2(x)$ is analytic at $x = -2$

$$\left. ((x+2) \cdot P_2(x)) \right|_{x=-2} = \frac{1}{3}$$

- $(x+2)^2 \cdot P_3(x)$ is analytic at $x = -2$

$$\left. ((x+2)^2 \cdot P_3(x)) \right|_{x=-2} = -\frac{1}{3}$$

- $x = -2$ is a regular singular point

Check to see if x_0 is a regular singular point

$$x_0 = -2$$

- Multiply by denominators

$$y''(x+2)^2(x-1)^2 + (3x+6)y' + (x-1)y = 0$$

- Change variables using $x = u - 2$ so that the regular singular point is at $u = 0$

$$(u^4 - 6u^3 + 9u^2) \left(\frac{d^2}{du^2} y(u) \right) + 3u \left(\frac{d}{du} y(u) \right) + (u-3)y(u) = 0$$

- Assume series solution for $y(u)$

$$y(u) = \sum_{k=0}^{\infty} a_k u^{k+r}$$

- Rewrite ODE with series expansions

- Convert $u^m \cdot y(u)$ to series expansion for $m = 0..1$

$$u^m \cdot y(u) = \sum_{k=0}^{\infty} a_k u^{k+r+m}$$

- Shift index using $k \rightarrow k - m$

$$u^m \cdot y(u) = \sum_{k=m}^{\infty} a_{k-m} u^{k+r}$$

- Convert $u \cdot \left(\frac{d}{du} y(u) \right)$ to series expansion

$$u \cdot \left(\frac{d}{du} y(u) \right) = \sum_{k=0}^{\infty} a_k (k+r) u^{k+r}$$

- Convert $u^m \cdot \left(\frac{d^2}{du^2} y(u) \right)$ to series expansion for $m = 2..4$

$$u^m \cdot \left(\frac{d^2}{du^2} y(u) \right) = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) u^{k+r-2+m}$$

- Shift index using $k \rightarrow k + 2 - m$

$$u^m \cdot \left(\frac{d^2}{du^2} y(u) \right) = \sum_{k=-2+m}^{\infty} a_{k+2-m} (k+2-m+r)(k+1-m+r) u^{k+r}$$

Rewrite ODE with series expansions

$$3a_0(1+3r)(-1+r)u^r + (3a_1(4+3r)r - a_0(6r^2 - 6r - 1))u^{1+r} + \left(\sum_{k=2}^{\infty} (3a_k(3k+3r+1)(k+r) - a_{k-1}(3k+3r+1)(k+r-1) - a_{k-2}(3k+3r+1)(k+r-2)) \right) u^{k+r} = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$3(1+3r)(-1+r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \left\{ 1, -\frac{1}{3} \right\}$$

- Each term must be 0

$$3a_1(4 + 3r)r - a_0(6r^2 - 6r - 1) = 0$$

- Solve for the dependent coefficient(s)

$$a_1 = \frac{a_0(6r^2 - 6r - 1)}{3r(4 + 3r)}$$

- Each term in the series must be 0, giving the recursion relation

$$(9a_k + a_{k-2} - 6a_{k-1})k^2 + (2(9a_k + a_{k-2} - 6a_{k-1})r - 6a_k - 5a_{k-2} + 18a_{k-1})k + (9a_k + a_{k-2} - 6a_{k-1}) = 0$$

- Shift index using $k \rightarrow k + 2$

$$(9a_{k+2} + a_k - 6a_{k+1})(k + 2)^2 + (2(9a_{k+2} + a_k - 6a_{k+1})r - 6a_{k+2} - 5a_k + 18a_{k+1})(k + 2) + (9a_{k+2} + a_k - 6a_{k+1}) = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+2} = -\frac{k^2 a_k - 6k^2 a_{k+1} + 2kra_k - 12kra_{k+1} + r^2 a_k - 6r^2 a_{k+1} - ka_k - 6ka_{k+1} - ra_k - 6ra_{k+1} + a_{k+1}}{3(3k^2 + 6kr + 3r^2 + 10k + 10r + 7)}$$

- Recursion relation for $r = 1$

$$a_{k+2} = -\frac{k^2 a_k - 6k^2 a_{k+1} + ka_k - 18ka_{k+1} - 11a_{k+1}}{3(3k^2 + 16k + 20)}$$

- Solution for $r = 1$

$$\left[y(u) = \sum_{k=0}^{\infty} a_k u^{k+1}, a_{k+2} = -\frac{k^2 a_k - 6k^2 a_{k+1} + ka_k - 18ka_{k+1} - 11a_{k+1}}{3(3k^2 + 16k + 20)}, a_1 = -\frac{a_0}{21} \right]$$

- Revert the change of variables $u = x + 2$

$$\left[y = \sum_{k=0}^{\infty} a_k (x + 2)^{k+1}, a_{k+2} = -\frac{k^2 a_k - 6k^2 a_{k+1} + ka_k - 18ka_{k+1} - 11a_{k+1}}{3(3k^2 + 16k + 20)}, a_1 = -\frac{a_0}{21} \right]$$

- Recursion relation for $r = -\frac{1}{3}$

$$a_{k+2} = -\frac{k^2 a_k - 6k^2 a_{k+1} - \frac{5}{3}ka_k - 2ka_{k+1} + \frac{4}{9}a_k + \frac{7}{3}a_{k+1}}{3(3k^2 + 8k + 4)}$$

- Solution for $r = -\frac{1}{3}$

$$\left[y(u) = \sum_{k=0}^{\infty} a_k u^{k-\frac{1}{3}}, a_{k+2} = -\frac{k^2 a_k - 6k^2 a_{k+1} - \frac{5}{3}ka_k - 2ka_{k+1} + \frac{4}{9}a_k + \frac{7}{3}a_{k+1}}{3(3k^2 + 8k + 4)}, a_1 = -\frac{5a_0}{9} \right]$$

- Revert the change of variables $u = x + 2$

$$\left[y = \sum_{k=0}^{\infty} a_k (x + 2)^{k-\frac{1}{3}}, a_{k+2} = -\frac{k^2 a_k - 6k^2 a_{k+1} - \frac{5}{3}ka_k - 2ka_{k+1} + \frac{4}{9}a_k + \frac{7}{3}a_{k+1}}{3(3k^2 + 8k + 4)}, a_1 = -\frac{5a_0}{9} \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k (x + 2)^{k+1} \right) + \left(\sum_{k=0}^{\infty} b_k (x + 2)^{k-\frac{1}{3}} \right), a_{k+2} = -\frac{k^2 a_k - 6k^2 a_{k+1} + ka_k - 18ka_{k+1} - 11a_{k+1}}{3(3k^2 + 16k + 20)}, a_1 = -\frac{a_0}{21} \right]$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
  -> Bessel
  -> elliptic
  -> Legendre
  -> Kummer
    -> hyper3: Equivalence to 1F1 under a power @ Moebius
  -> hypergeometric
    -> heuristic approach
    -> hyper3: Equivalence to 2F1, 1F1 or 0F1 under a power @ Moebius
  -> Mathieu
    -> Equivalence to the rational form of Mathieu ODE under a power @ Moebius
trying a solution in terms of MeijerG functions
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
<- Heun successful: received ODE is equivalent to the HeunC ODE, case a = 0, e <> 0, c <>
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 57

```
Order:=8;
dsolve((x^2+x-2)^2*diff(y(x),x$2)+3*(x+2)*diff(y(x),x)+(x-1)*y(x)=0,y(x),type='series',x=-2)
```

$$y(x) = \frac{c_1 \left(1 - \frac{5}{9}(x+2) + \frac{23}{324}(x+2)^2 + \frac{271}{43740}(x+2)^3 + \frac{10517}{12597120}(x+2)^4 + \frac{778801}{6235574400}(x+2)^5 + \frac{16965493}{942818849280}(x+2)^6 \right)}{1}$$

✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 148

AsymptoticDSolveValue[(x^2+x-2)^2*y'[x]+3*(x+2)*y'[x]+(x-1)*y[x]==0,y[x],{x,-2,7}]

$$y(x) \rightarrow c_1(x+2) \left(-\frac{52991201(x+2)^7}{11727918720000} - \frac{5797423(x+2)^6}{290405606400} - \frac{709507(x+2)^5}{8066822400} \right. \\ \left. - \frac{11093(x+2)^4}{28304640} - \frac{53(x+2)^3}{29484} - \frac{11(x+2)^2}{1260} + \frac{1}{21}(-x-2) + 1 \right) \\ + \frac{c_2 \left(\frac{899971067(x+2)^7}{458981357990400} + \frac{16965493(x+2)^6}{942818849280} + \frac{778801(x+2)^5}{6235574400} + \frac{10517(x+2)^4}{12597120} + \frac{271(x+2)^3}{43740} + \frac{23}{324}(x+2)^2 - \frac{5(x+2)}{9} + 1 \right)}{\sqrt[3]{x+2}}$$

17.7 problem 1(g)

Internal problem ID [6046]

Internal file name [OUTPUT/5294_Sunday_June_05_2022_03_29_54_PM_91142448/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 154

Problem number: 1(g).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Complex roots"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$x^2y'' + \sin(x)y' + y \cos(x) = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2y'' + \sin(x)y' + y \cos(x) = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{\sin(x)}{x^2}$$

$$q(x) = \frac{\cos(x)}{x^2}$$

Table 217: Table $p(x), q(x)$ singularities.

$p(x) = \frac{\sin(x)}{x^2}$	
singularity	type
$x = 0$	“regular”

$q(x) = \frac{\cos(x)}{x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2 y'' + \sin(x) y' + y \cos(x) = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned}
 & x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\
 & + \sin(x) \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) \cos(x) = 0
 \end{aligned} \tag{1}$$

Expanding $\sin(x)$ as Taylor series around $x = 0$ and keeping only the first 8 terms gives

$$\begin{aligned}\sin(x) &= x - \frac{1}{6}x^3 + \frac{1}{120}x^5 - \frac{1}{5040}x^7 + \frac{1}{362880}x^9 + \dots \\ &= x - \frac{1}{6}x^3 + \frac{1}{120}x^5 - \frac{1}{5040}x^7 + \frac{1}{362880}x^9\end{aligned}$$

Expanding $\cos(x)$ as Taylor series around $x = 0$ and keeping only the first 8 terms gives

$$\begin{aligned}\cos(x) &= \frac{1}{40320}x^8 + 1 - \frac{1}{2}x^2 + \frac{1}{24}x^4 - \frac{1}{720}x^6 + \dots \\ &= \frac{1}{40320}x^8 + 1 - \frac{1}{2}x^2 + \frac{1}{24}x^4 - \frac{1}{720}x^6\end{aligned}$$

Which simplifies to

$$\begin{aligned}&\left(\sum_{n=0}^{\infty} x^{n+r} a_n(n+r)(n+r-1)\right) + \left(\sum_{n=0}^{\infty} \frac{x^{n+r+8} a_n(n+r)}{362880}\right) \\ &+ \sum_{n=0}^{\infty} \left(-\frac{x^{n+r+6} a_n(n+r)}{5040}\right) + \left(\sum_{n=0}^{\infty} \frac{x^{n+r+4} a_n(n+r)}{120}\right) \\ &+ \sum_{n=0}^{\infty} \left(-\frac{x^{n+r+2} a_n(n+r)}{6}\right) + \left(\sum_{n=0}^{\infty} x^{n+r} a_n(n+r)\right) \tag{2A} \\ &+ \left(\sum_{n=0}^{\infty} \frac{x^{n+r+8} a_n}{40320}\right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r}\right) + \sum_{n=0}^{\infty} \left(-\frac{x^{n+r+2} a_n}{2}\right) \\ &+ \left(\sum_{n=0}^{\infty} \frac{x^{n+r+4} a_n}{24}\right) + \sum_{n=0}^{\infty} \left(-\frac{x^{n+r+6} a_n}{720}\right) = 0\end{aligned}$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\begin{aligned}\sum_{n=0}^{\infty} \frac{x^{n+r+8} a_n(n+r)}{362880} &= \sum_{n=8}^{\infty} \frac{a_{n-8}(n-8+r) x^{n+r}}{362880} \\ \sum_{n=0}^{\infty} \left(-\frac{x^{n+r+6} a_n(n+r)}{5040}\right) &= \sum_{n=6}^{\infty} \left(-\frac{a_{n-6}(n-6+r) x^{n+r}}{5040}\right) \\ \sum_{n=0}^{\infty} \frac{x^{n+r+4} a_n(n+r)}{120} &= \sum_{n=4}^{\infty} \frac{a_{n-4}(n-4+r) x^{n+r}}{120}\end{aligned}$$

$$\begin{aligned}
\sum_{n=0}^{\infty} \left(-\frac{x^{n+r+2} a_n (n+r)}{6} \right) &= \sum_{n=2}^{\infty} \left(-\frac{a_{n-2} (n+r-2) x^{n+r}}{6} \right) \\
\sum_{n=0}^{\infty} \frac{x^{n+r+8} a_n}{40320} &= \sum_{n=8}^{\infty} \frac{a_{n-8} x^{n+r}}{40320} \\
\sum_{n=0}^{\infty} \left(-\frac{x^{n+r+2} a_n}{2} \right) &= \sum_{n=2}^{\infty} \left(-\frac{a_{n-2} x^{n+r}}{2} \right) \\
\sum_{n=0}^{\infty} \frac{x^{n+r+4} a_n}{24} &= \sum_{n=4}^{\infty} \frac{a_{n-4} x^{n+r}}{24} \\
\sum_{n=0}^{\infty} \left(-\frac{x^{n+r+6} a_n}{720} \right) &= \sum_{n=6}^{\infty} \left(-\frac{a_{n-6} x^{n+r}}{720} \right)
\end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned}
&\left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=8}^{\infty} \frac{a_{n-8} (n-8+r) x^{n+r}}{362880} \right) \\
&+ \sum_{n=6}^{\infty} \left(-\frac{a_{n-6} (n-6+r) x^{n+r}}{5040} \right) + \left(\sum_{n=4}^{\infty} \frac{a_{n-4} (n-4+r) x^{n+r}}{120} \right) \\
&+ \sum_{n=2}^{\infty} \left(-\frac{a_{n-2} (n+r-2) x^{n+r}}{6} \right) + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) \right) \\
&+ \left(\sum_{n=8}^{\infty} \frac{a_{n-8} x^{n+r}}{40320} \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \sum_{n=2}^{\infty} \left(-\frac{a_{n-2} x^{n+r}}{2} \right) \\
&+ \left(\sum_{n=4}^{\infty} \frac{a_{n-4} x^{n+r}}{24} \right) + \sum_{n=6}^{\infty} \left(-\frac{a_{n-6} x^{n+r}}{720} \right) = 0
\end{aligned} \tag{2B}$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r} a_n (n+r) (n+r-1) + x^{n+r} a_n (n+r) + a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$x^r a_0 r (-1+r) + x^r a_0 r + a_0 x^r = 0$$

Or

$$(x^r r (-1+r) + x^r r + x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(r^2 + 1) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$r^2 + 1 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= i \\ r_2 &= -i \end{aligned}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(r^2 + 1) x^r = 0$$

Solving for r gives the roots of the indicial equation as Since the roots are complex conjugates, then two linearly independent solutions can be constructed using

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^{n+i} \\ y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n-i} \end{aligned}$$

$y_1(x)$ is found first. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

Substituting $n = 2$ in Eq. (2B) gives

$$a_2 = \frac{r + 3}{6r^2 + 24r + 30}$$

Substituting $n = 3$ in Eq. (2B) gives

$$a_3 = 0$$

Substituting $n = 4$ in Eq. (2B) gives

$$a_4 = \frac{-3r^3 - 17r^2 + 5r + 75}{360(r^2 + 4r + 5)(r^2 + 8r + 17)}$$

Substituting $n = 5$ in Eq. (2B) gives

$$a_5 = 0$$

Substituting $n = 6$ in Eq. (2B) gives

$$a_6 = \frac{3r^5 + 15r^4 - 230r^3 - 1818r^2 - 3805r - 2037}{15120(r^2 + 4r + 5)(r^2 + 8r + 17)(r^2 + 12r + 37)}$$

Substituting $n = 7$ in Eq. (2B) gives

$$a_7 = 0$$

For $8 \leq n$ the recursive equation is

$$\begin{aligned} & a_n(n+r)(n+r-1) + \frac{a_{n-8}(n-8+r)}{362880} - \frac{a_{n-6}(n-6+r)}{5040} + \frac{a_{n-4}(n-4+r)}{120} \quad (3) \\ & - \frac{a_{n-2}(n+r-2)}{6} + a_n(n+r) + \frac{a_{n-8}}{40320} + a_n - \frac{a_{n-2}}{2} + \frac{a_{n-4}}{24} - \frac{a_{n-6}}{720} = 0 \end{aligned}$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{na_{n-8} - 72na_{n-6} + 3024na_{n-4} - 60480na_{n-2} + ra_{n-8} - 72ra_{n-6} + 3024ra_{n-4} - 60480ra_{n-2} + a_n}{362880(n^2 + 2nr + r^2 + 1)} \quad (4)$$

Which for the root $r = i$ becomes

$$a_n = -\frac{(a_{n-8} - 72a_{n-6} + 3024a_{n-4} - 60480a_{n-2})(1 + i + n)}{362880n(2i + n)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = i$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{r+3}{6r^2+24r+30}$	$\frac{1}{12} - \frac{i}{24}$
a_3	0	0
a_4	$\frac{-3r^3-17r^2+5r+75}{360(r^2+4r+5)(r^2+8r+17)}$	$\frac{29}{28800} - \frac{67i}{28800}$
a_5	0	0
a_6	$\frac{3r^5+15r^4-230r^3-1818r^2-3805r-2037}{15120(r^2+4r+5)(r^2+8r+17)(r^2+12r+37)}$	$-\frac{893}{14515200} + \frac{17i}{4838400}$
a_7	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned}
y_1(x) &= x^i(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots) \\
&= x^i \left(1 + \left(\frac{1}{12} - \frac{i}{24} \right) x^2 + \left(\frac{29}{28800} - \frac{67i}{28800} \right) x^4 + \left(-\frac{893}{14515200} + \frac{17i}{4838400} \right) x^6 + O(x^8) \right)
\end{aligned}$$

The second solution $y_2(x)$ is found by taking the complex conjugate of $y_1(x)$ which gives

$$\begin{aligned}
y_2(x) &= x^{-i} \left(1 + \left(\frac{1}{12} + \frac{i}{24} \right) x^2 + \left(\frac{29}{28800} + \frac{67i}{28800} \right) x^4 + \left(-\frac{893}{14515200} - \frac{17i}{4838400} \right) x^6 \right. \\
&\quad \left. + O(x^8) \right)
\end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
y_h(x) &= c_1 y_1(x) + c_2 y_2(x) \\
&= c_1 x^i \left(1 + \left(\frac{1}{12} - \frac{i}{24} \right) x^2 + \left(\frac{29}{28800} - \frac{67i}{28800} \right) x^4 + \left(-\frac{893}{14515200} + \frac{17i}{4838400} \right) x^6 \right. \\
&\quad \left. + O(x^8) \right) + c_2 x^{-i} \left(1 + \left(\frac{1}{12} + \frac{i}{24} \right) x^2 + \left(\frac{29}{28800} + \frac{67i}{28800} \right) x^4 \right. \\
&\quad \left. + \left(-\frac{893}{14515200} - \frac{17i}{4838400} \right) x^6 + O(x^8) \right)
\end{aligned}$$

Hence the final solution is

$$\begin{aligned}
 y &= y_h \\
 &= c_1 x^i \left(1 + \left(\frac{1}{12} - \frac{i}{24} \right) x^2 + \left(\frac{29}{28800} - \frac{67i}{28800} \right) x^4 + \left(-\frac{893}{14515200} + \frac{17i}{4838400} \right) x^6 \right. \\
 &\quad \left. + O(x^8) \right) + c_2 x^{-i} \left(1 + \left(\frac{1}{12} + \frac{i}{24} \right) x^2 + \left(\frac{29}{28800} + \frac{67i}{28800} \right) x^4 \right. \\
 &\quad \left. + \left(-\frac{893}{14515200} - \frac{17i}{4838400} \right) x^6 + O(x^8) \right)
 \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}
 y &= c_1 x^i \left(1 + \left(\frac{1}{12} - \frac{i}{24} \right) x^2 + \left(\frac{29}{28800} - \frac{67i}{28800} \right) x^4 + \left(-\frac{893}{14515200} + \frac{17i}{4838400} \right) x^6 \right. \\
 &\quad \left. + O(x^8) \right) + c_2 x^{-i} \left(1 + \left(\frac{1}{12} + \frac{i}{24} \right) x^2 + \left(\frac{29}{28800} + \frac{67i}{28800} \right) x^4 \right. \\
 &\quad \left. + \left(-\frac{893}{14515200} - \frac{17i}{4838400} \right) x^6 + O(x^8) \right)
 \end{aligned}$$

Verification of solutions

$$\begin{aligned}
 y &= c_1 x^i \left(1 + \left(\frac{1}{12} - \frac{i}{24} \right) x^2 + \left(\frac{29}{28800} - \frac{67i}{28800} \right) x^4 + \left(-\frac{893}{14515200} + \frac{17i}{4838400} \right) x^6 \right. \\
 &\quad \left. + O(x^8) \right) + c_2 x^{-i} \left(1 + \left(\frac{1}{12} + \frac{i}{24} \right) x^2 + \left(\frac{29}{28800} + \frac{67i}{28800} \right) x^4 \right. \\
 &\quad \left. + \left(-\frac{893}{14515200} - \frac{17i}{4838400} \right) x^6 + O(x^8) \right)
 \end{aligned}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) * 2F1([a
-> Trying changes of variables to rationalize or make the ODE simpler
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) * 2F1
trying a symmetry of the form [xi=0, eta=F(x)]
trying 2nd order exact linear
trying symmetries linear in x and y(x)
trying to convert to a linear ODE with constant coefficients
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) * 2F1
trying a symmetry of the form [xi=0, eta=F(x)]
trying 2nd order exact linear
trying symmetries linear in x and y(x)
trying to convert to a linear ODE with constant coefficients
-> trying with_periodic_functions in the coefficients
--- Trying Lie symmetry methods, 2nd order ---
`, `-> Computing symmetries using: way = 5
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) * 2F1
trying a symmetry of the form [xi=0, eta=F(x)]
trying 2nd order exact linear
trying symmetries linear in x and y(x)
trying to convert to a linear ODE with constant coefficients
<- unable to find a useful change of variables
trying a symmetry of the form [xi=0, eta=F(x)]
trying differential order: 2; exact nonlinear
trying symmetries linear in x and y(x)
trying to convert to a linear ODE with constant coefficients
trying 2nd order, integrating factor of the form mu(x,y)
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y1425
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) * 2F1
-> Trying changes of variables to rationalize or make the ODE simpler
```

✓ Solution by Maple

Time used: 0.203 (sec). Leaf size: 53

Order:=8;

```
dsolve(x^2*diff(y(x),x$2)+sin(x)*diff(y(x),x)+cos(x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^{-i} \left(1 + \left(\frac{1}{12} + \frac{i}{24} \right) x^2 + \left(\frac{29}{28800} + \frac{67i}{28800} \right) x^4 \right. \\ \left. + \left(-\frac{893}{14515200} - \frac{17i}{4838400} \right) x^6 + O(x^8) \right) + c_2 x^i \left(1 + \left(\frac{1}{12} - \frac{i}{24} \right) x^2 \right. \\ \left. + \left(\frac{29}{28800} - \frac{67i}{28800} \right) x^4 + \left(-\frac{893}{14515200} + \frac{17i}{4838400} \right) x^6 + O(x^8) \right)$$

✓ Solution by Mathematica

Time used: 0.048 (sec). Leaf size: 112

```
AsymptoticDSolveValue[x^2*y''[x]+Sin[x]*y'[x]+Cos[x]*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 x^{-i} \left(\left(-\frac{26459}{59222016000} - \frac{12449i}{7402752000} \right) x^8 - \left(\frac{893}{14515200} + \frac{17i}{4838400} \right) x^6 \right. \\ \left. + \left(\frac{29}{28800} + \frac{67i}{28800} \right) x^4 + \left(\frac{1}{12} + \frac{i}{24} \right) x^2 + 1 \right) \\ + c_2 x^i \left(\left(-\frac{26459}{59222016000} + \frac{12449i}{7402752000} \right) x^8 - \left(\frac{893}{14515200} - \frac{17i}{4838400} \right) x^6 \right. \\ \left. + \left(\frac{29}{28800} - \frac{67i}{28800} \right) x^4 + \left(\frac{1}{12} - \frac{i}{24} \right) x^2 + 1 \right)$$

17.8 problem 2(b)

17.8.1 Maple step by step solution 1438

Internal problem ID [6047]

Internal file name [OUTPUT/5295_Sunday_June_05_2022_03_31_08_PM_14870823/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 154

Problem number: 2(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference is integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$x^2 y'' + xy' + \left(x^2 - \frac{1}{4}\right) y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2 y'' + xy' + \left(x^2 - \frac{1}{4}\right) y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = \frac{4x^2 - 1}{4x^2}$$

Table 218: Table $p(x), q(x)$ singularities.

$p(x) = \frac{1}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = \frac{4x^2-1}{4x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2 y'' + x y' + \left(x^2 - \frac{1}{4}\right) y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned}
 & x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\
 & + x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(x^2 - \frac{1}{4}\right) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0
 \end{aligned} \tag{1}$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) \right) \\ & + \left(\sum_{n=0}^{\infty} x^{n+r+2} a_n \right) + \sum_{n=0}^{\infty} \left(-\frac{a_n x^{n+r}}{4} \right) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} x^{n+r+2} a_n = \sum_{n=2}^{\infty} a_{n-2} x^{n+r}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) \right) \\ & + \left(\sum_{n=2}^{\infty} a_{n-2} x^{n+r} \right) + \sum_{n=0}^{\infty} \left(-\frac{a_n x^{n+r}}{4} \right) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r} a_n (n+r) (n+r-1) + x^{n+r} a_n (n+r) - \frac{a_n x^{n+r}}{4} = 0$$

When $n=0$ the above becomes

$$x^r a_0 r(-1+r) + x^r a_0 r - \frac{a_0 x^r}{4} = 0$$

Or

$$\left(x^r r(-1+r) + x^r r - \frac{x^r}{4} \right) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$\frac{(4r^2 - 1) x^r}{4} = 0$$

Since the above is true for all x then the indicial equation becomes

$$r^2 - \frac{1}{4} = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = \frac{1}{2}$$

$$r_2 = -\frac{1}{2}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$\frac{(4r^2 - 1)x^r}{4} = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = 1$ is an integer, then we can construct two linearly independent solutions

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = C y_1(x) \ln(x) + x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sqrt{x} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = C y_1(x) \ln(x) + \frac{\sum_{n=0}^{\infty} b_n x^n}{\sqrt{x}}$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+\frac{1}{2}}$$

$$y_2(x) = C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n-\frac{1}{2}} \right)$$

Where C above can be zero. We start by finding y_1 . Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

For $2 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) + a_n(n+r) + a_{n-2} - \frac{a_n}{4} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{4a_{n-2}}{4n^2 + 8nr + 4r^2 - 1} \quad (4)$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_n = -\frac{a_{n-2}}{n(n+1)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = \frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = -\frac{4}{4r^2 + 16r + 15}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_2 = -\frac{1}{6}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{4}{4r^2+16r+15}$	$-\frac{1}{6}$

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{4}{4r^2+16r+15}$	$-\frac{1}{6}$
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{16}{(4r^2 + 16r + 15)(4r^2 + 32r + 63)}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_4 = \frac{1}{120}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{4}{4r^2+16r+15}$	$-\frac{1}{6}$
a_3	0	0
a_4	$\frac{16}{(4r^2+16r+15)(4r^2+32r+63)}$	$\frac{1}{120}$

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{4}{4r^2+16r+15}$	$-\frac{1}{6}$
a_3	0	0
a_4	$\frac{16}{(4r^2+16r+15)(4r^2+32r+63)}$	$\frac{1}{120}$
a_5	0	0

For $n = 6$, using the above recursive equation gives

$$a_6 = -\frac{64}{(4r^2 + 16r + 15)(4r^2 + 32r + 63)(4r^2 + 48r + 143)}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_6 = -\frac{1}{5040}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{4}{4r^2+16r+15}$	$-\frac{1}{6}$
a_3	0	0
a_4	$\frac{16}{(4r^2+16r+15)(4r^2+32r+63)}$	$\frac{1}{120}$
a_5	0	0
a_6	$-\frac{64}{(4r^2+16r+15)(4r^2+32r+63)(4r^2+48r+143)}$	$-\frac{1}{5040}$

For $n = 7$, using the above recursive equation gives

$$a_7 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{4}{4r^2+16r+15}$	$-\frac{1}{6}$
a_3	0	0
a_4	$\frac{16}{(4r^2+16r+15)(4r^2+32r+63)}$	$\frac{1}{120}$
a_5	0	0
a_6	$-\frac{64}{(4r^2+16r+15)(4r^2+32r+63)(4r^2+48r+143)}$	$-\frac{1}{5040}$
a_7	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= \sqrt{x}(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots) \\ &= \sqrt{x} \left(1 - \frac{x^2}{6} + \frac{x^4}{120} - \frac{x^6}{5040} + O(x^8) \right) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Let

$$r_1 - r_2 = N$$

Where N is positive integer which is the difference between the two roots. r_1 is taken as the larger root. Hence for this problem we have $N = 1$. Now we need to determine if C is zero or not. This is done by finding $\lim_{r \rightarrow r_2} a_1(r)$. If this limit exists, then $C = 0$, else we need to keep the log term and $C \neq 0$. The above table shows that

$$\begin{aligned} a_N &= a_1 \\ &= 0 \end{aligned}$$

Therefore

$$\begin{aligned} \lim_{r \rightarrow r_2} 0 &= \lim_{r \rightarrow -\frac{1}{2}} 0 \\ &= 0 \end{aligned}$$

The limit is 0. Since the limit exists then the log term is not needed and we can set $C = 0$. Therefore the second solution has the form

$$\begin{aligned} y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+r} \\ &= \sum_{n=0}^{\infty} b_n x^{n-\frac{1}{2}} \end{aligned}$$

Eq (3) derived above is used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq(3) gives

$$b_1 = 0$$

For $2 \leq n$ the recursive equation is

$$b_n(n+r)(n+r-1) + b_n(n+r) + b_{n-2} - \frac{b_n}{4} = 0 \quad (4)$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_n \left(n - \frac{1}{2} \right) \left(n - \frac{3}{2} \right) + b_n \left(n - \frac{1}{2} \right) + b_{n-2} - \frac{b_n}{4} = 0 \quad (4A)$$

Solving for b_n from the recursive equation (4) gives

$$b_n = -\frac{4b_{n-2}}{4n^2 + 8nr + 4r^2 - 1} \quad (5)$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_n = -\frac{4b_{n-2}}{4n^2 - 4n} \quad (6)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = -\frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0

For $n = 2$, using the above recursive equation gives

$$b_2 = -\frac{4}{4r^2 + 16r + 15}$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_2 = -\frac{1}{2}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{4}{4r^2 + 16r + 15}$	$-\frac{1}{2}$

For $n = 3$, using the above recursive equation gives

$$b_3 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{4}{4r^2+16r+15}$	$-\frac{1}{2}$
b_3	0	0

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{16}{(4r^2 + 16r + 15)(4r^2 + 32r + 63)}$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_4 = \frac{1}{24}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{4}{4r^2+16r+15}$	$-\frac{1}{2}$
b_3	0	0
b_4	$\frac{16}{(4r^2+16r+15)(4r^2+32r+63)}$	$\frac{1}{24}$

For $n = 5$, using the above recursive equation gives

$$b_5 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{4}{4r^2+16r+15}$	$-\frac{1}{2}$
b_3	0	0
b_4	$\frac{16}{(4r^2+16r+15)(4r^2+32r+63)}$	$\frac{1}{24}$
b_5	0	0

For $n = 6$, using the above recursive equation gives

$$b_6 = -\frac{64}{(4r^2 + 16r + 15)(4r^2 + 32r + 63)(4r^2 + 48r + 143)}$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_6 = -\frac{1}{720}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{4}{4r^2+16r+15}$	$-\frac{1}{2}$
b_3	0	0
b_4	$\frac{16}{(4r^2+16r+15)(4r^2+32r+63)}$	$\frac{1}{24}$
b_5	0	0
b_6	$-\frac{64}{(4r^2+16r+15)(4r^2+32r+63)(4r^2+48r+143)}$	$-\frac{1}{720}$

For $n = 7$, using the above recursive equation gives

$$b_7 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{4}{4r^2+16r+15}$	$-\frac{1}{2}$
b_3	0	0
b_4	$\frac{16}{(4r^2+16r+15)(4r^2+32r+63)}$	$\frac{1}{24}$
b_5	0	0
b_6	$-\frac{64}{(4r^2+16r+15)(4r^2+32r+63)(4r^2+48r+143)}$	$-\frac{1}{720}$
b_7	0	0

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned} y_2(x) &= \sqrt{x}(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 + b_7x^7 + b_8x^8 \dots) \\ &= \frac{1 - \frac{x^2}{2} + \frac{x^4}{24} - \frac{x^6}{720} + O(x^8)}{\sqrt{x}} \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= c_1\sqrt{x} \left(1 - \frac{x^2}{6} + \frac{x^4}{120} - \frac{x^6}{5040} + O(x^8) \right) + \frac{c_2 \left(1 - \frac{x^2}{2} + \frac{x^4}{24} - \frac{x^6}{720} + O(x^8) \right)}{\sqrt{x}} \end{aligned}$$

Hence the final solution is

$$\begin{aligned} y &= y_h \\ &= c_1\sqrt{x} \left(1 - \frac{x^2}{6} + \frac{x^4}{120} - \frac{x^6}{5040} + O(x^8) \right) + \frac{c_2 \left(1 - \frac{x^2}{2} + \frac{x^4}{24} - \frac{x^6}{720} + O(x^8) \right)}{\sqrt{x}} \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1\sqrt{x} \left(1 - \frac{x^2}{6} + \frac{x^4}{120} - \frac{x^6}{5040} + O(x^8) \right) + \frac{c_2 \left(1 - \frac{x^2}{2} + \frac{x^4}{24} - \frac{x^6}{720} + O(x^8) \right)}{\sqrt{x}} \quad (1)$$

Verification of solutions

$$y = c_1\sqrt{x} \left(1 - \frac{x^2}{6} + \frac{x^4}{120} - \frac{x^6}{5040} + O(x^8) \right) + \frac{c_2 \left(1 - \frac{x^2}{2} + \frac{x^4}{24} - \frac{x^6}{720} + O(x^8) \right)}{\sqrt{x}}$$

Verified OK.

17.8.1 Maple step by step solution

Let's solve

$$x^2y'' + xy' + \left(x^2 - \frac{1}{4}\right)y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{(4x^2-1)y}{4x^2} - \frac{y'}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{y'}{x} + \frac{(4x^2-1)y}{4x^2} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$\left[P_2(x) = \frac{1}{x}, P_3(x) = \frac{4x^2-1}{4x^2} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 1$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = -\frac{1}{4}$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$4x^2y'' + 4xy' + (4x^2 - 1)y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 0..2$

$$x^m \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

- Shift index using $k \rightarrow k - m$

$$x^m \cdot y = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

- Convert $x \cdot y'$ to series expansion

$$x \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r}$$

- Convert $x^2 \cdot y''$ to series expansion

$$x^2 \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(1+2r)(-1+2r)x^r + a_1(3+2r)(1+2r)x^{1+r} + \left(\sum_{k=2}^{\infty} (a_k(2k+2r+1)(2k+2r-1) + 4a_{k-2}) \right) x^{k+r} = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$(1+2r)(-1+2r) = 0$$
- Values of r that satisfy the indicial equation

$$r \in \left\{ -\frac{1}{2}, \frac{1}{2} \right\}$$
- Each term must be 0

$$a_1(3+2r)(1+2r) = 0$$
- Solve for the dependent coefficient(s)

$$a_1 = 0$$
- Each term in the series must be 0, giving the recursion relation

$$a_k(4k^2 + 8kr + 4r^2 - 1) + 4a_{k-2} = 0$$
- Shift index using $k \rightarrow k+2$

$$a_{k+2}(4(k+2)^2 + 8(k+2)r + 4r^2 - 1) + 4a_k = 0$$
- Recursion relation that defines series solution to ODE

$$a_{k+2} = -\frac{4a_k}{4k^2 + 8kr + 4r^2 + 16k + 16r + 15}$$
- Recursion relation for $r = -\frac{1}{2}$

$$a_{k+2} = -\frac{4a_k}{4k^2 + 12k + 8}$$
- Solution for $r = -\frac{1}{2}$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k-\frac{1}{2}}, a_{k+2} = -\frac{4a_k}{4k^2 + 12k + 8}, a_1 = 0 \right]$$
- Recursion relation for $r = \frac{1}{2}$

$$a_{k+2} = -\frac{4a_k}{4k^2 + 20k + 24}$$
- Solution for $r = \frac{1}{2}$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+\frac{1}{2}}, a_{k+2} = -\frac{4a_k}{4k^2 + 20k + 24}, a_1 = 0 \right]$$
- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^{k-\frac{1}{2}} \right) + \left(\sum_{k=0}^{\infty} b_k x^{k+\frac{1}{2}} \right), a_{k+2} = -\frac{4a_k}{4k^2+12k+8}, a_1 = 0, b_{k+2} = -\frac{4b_k}{4k^2+20k+24}, b_1 = 0 \right]$$

Maple trace Kovacic algorithm successful

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Group is reducible or imprimitive
<- Kovacics algorithm successful`

```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 39

```

Order:=8;
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)+(x^2-1/4)*y(x)=0,y(x),type='series',x=0);

```

$$y(x) = \frac{c_1 x \left(1 - \frac{1}{6}x^2 + \frac{1}{120}x^4 - \frac{1}{5040}x^6 + O(x^8) \right) + c_2 \left(1 - \frac{1}{2}x^2 + \frac{1}{24}x^4 - \frac{1}{720}x^6 + O(x^8) \right)}{\sqrt{x}}$$

✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 76

```

AsymptoticDSolveValue[x^2*y''[x]+x*y'[x]+(x^2-1/4)*y[x]==0,y[x],{x,0,7}]

```

$$y(x) \rightarrow c_1 \left(-\frac{x^{11/2}}{720} + \frac{x^{7/2}}{24} - \frac{x^{3/2}}{2} + \frac{1}{\sqrt{x}} \right) + c_2 \left(-\frac{x^{13/2}}{5040} + \frac{x^{9/2}}{120} - \frac{x^{5/2}}{6} + \sqrt{x} \right)$$

17.9 problem 2(c)

17.9.1 Maple step by step solution 1453

Internal problem ID [6048]

Internal file name [OUTPUT/5296_Sunday_June_05_2022_03_31_10_PM_61777425/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 154

Problem number: 2(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$4x^2y'' + (4x^4 - 5x)y' + y(x^2 + 2) = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$4x^2y'' + (4x^4 - 5x)y' + y(x^2 + 2) = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{4x^3 - 5}{4x}$$
$$q(x) = \frac{x^2 + 2}{4x^2}$$

Table 220: Table $p(x), q(x)$ singularities.

$p(x) = \frac{4x^3-5}{4x}$	
singularity	type
$x = 0$	“regular”
$x = \infty$	“regular”
$x = -\infty$	“regular”

$q(x) = \frac{x^2+2}{4x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0, \infty, -\infty]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$4x^2 y'' + (4x^4 - 5x) y' + y(x^2 + 2) = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$4x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) + (4x^4 - 5x) \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) (x^2 + 2) = 0 \quad (1)$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 4x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} 4x^{n+r+3} a_n (n+r) \right) \\ & + \sum_{n=0}^{\infty} (-5x^{n+r} a_n (n+r)) + \left(\sum_{n=0}^{\infty} x^{n+r+2} a_n \right) + \left(\sum_{n=0}^{\infty} 2a_n x^{n+r} \right) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=0}^{\infty} 4x^{n+r+3} a_n (n+r) &= \sum_{n=3}^{\infty} 4a_{n-3} (n-3+r) x^{n+r} \\ \sum_{n=0}^{\infty} x^{n+r+2} a_n &= \sum_{n=2}^{\infty} a_{n-2} x^{n+r} \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 4x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=3}^{\infty} 4a_{n-3} (n-3+r) x^{n+r} \right) \\ & + \sum_{n=0}^{\infty} (-5x^{n+r} a_n (n+r)) + \left(\sum_{n=2}^{\infty} a_{n-2} x^{n+r} \right) + \left(\sum_{n=0}^{\infty} 2a_n x^{n+r} \right) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$4x^{n+r} a_n (n+r) (n+r-1) - 5x^{n+r} a_n (n+r) + 2a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$4x^r a_0 r (-1+r) - 5x^r a_0 r + 2a_0 x^r = 0$$

Or

$$(4x^r r (-1+r) - 5x^r r + 2x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(4r^2 - 9r + 2) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$4r^2 - 9r + 2 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 2 \\ r_2 &= \frac{1}{4} \end{aligned}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(4r^2 - 9r + 2) x^r = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = \frac{7}{4}$ is not an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^{n+2} \\ y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+\frac{1}{4}} \end{aligned}$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

Substituting $n = 2$ in Eq. (2B) gives

$$a_2 = -\frac{1}{r(4r+7)}$$

For $3 \leq n$ the recursive equation is

$$4a_n(n+r)(n+r-1) + 4a_{n-3}(n-3+r) - 5a_n(n+r) + a_{n-2} + 2a_n = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{4na_{n-3} + 4ra_{n-3} - 12a_{n-3} + a_{n-2}}{4n^2 + 8nr + 4r^2 - 9n - 9r + 2} \quad (4)$$

Which for the root $r = 2$ becomes

$$a_n = \frac{-4na_{n-3} + 4a_{n-3} - a_{n-2}}{n(4n + 7)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 2$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r(4r+7)}$	$-\frac{1}{30}$

For $n = 3$, using the above recursive equation gives

$$a_3 = -\frac{4r}{4r^2 + 15r + 11}$$

Which for the root $r = 2$ becomes

$$a_3 = -\frac{8}{57}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r(4r+7)}$	$-\frac{1}{30}$
a_3	$-\frac{4r}{4r^2+15r+11}$	$-\frac{8}{57}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{16r^4 + 120r^3 + 281r^2 + 210r}$$

Which for the root $r = 2$ becomes

$$a_4 = \frac{1}{2760}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r(4r+7)}$	$-\frac{1}{30}$
a_3	$-\frac{4r}{4r^2+15r+11}$	$-\frac{8}{57}$
a_4	$\frac{1}{16r^4+120r^3+281r^2+210r}$	$\frac{1}{2760}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{32r^3 + 120r^2 + 164r + 88}{(4r + 7)r(4r^2 + 15r + 11)(4r^2 + 31r + 57)}$$

Which for the root $r = 2$ becomes

$$a_5 = \frac{64}{12825}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r(4r+7)}$	$-\frac{1}{30}$
a_3	$-\frac{4r}{4r^2+15r+11}$	$-\frac{8}{57}$
a_4	$\frac{1}{16r^4+120r^3+281r^2+210r}$	$\frac{1}{2760}$
a_5	$\frac{32r^3+120r^2+164r+88}{(4r+7)r(4r^2+15r+11)(4r^2+31r+57)}$	$\frac{64}{12825}$

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{256r^6 + 2688r^5 + 10256r^4 + 16848r^3 + 10076r^2 - 15r - 11}{(4r^2 + 15r + 11)r(16r^3 + 120r^2 + 281r + 210)(4r^2 + 39r + 92)}$$

Which for the root $r = 2$ becomes

$$a_6 = \frac{147181}{9753840}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r(4r+7)}$	$-\frac{1}{30}$
a_3	$-\frac{4r}{4r^2+15r+11}$	$-\frac{8}{57}$
a_4	$\frac{1}{16r^4+120r^3+281r^2+210r}$	$\frac{1}{2760}$
a_5	$\frac{32r^3+120r^2+164r+88}{(4r+7)r(4r^2+15r+11)(4r^2+31r+57)}$	$\frac{64}{12825}$
a_6	$\frac{256r^6+2688r^5+10256r^4+16848r^3+10076r^2-15r-11}{(4r^2+15r+11)r(16r^3+120r^2+281r+210)(4r^2+39r+92)}$	$\frac{147181}{9753840}$

For $n = 7$, using the above recursive equation gives

$$a_7 = -\frac{4(48r^5 + 552r^4 + 2567r^3 + 6075r^2 + 7147r + 3168)}{(4r+7)(4r^2+23r+30)r(4r^2+15r+11)(4r^2+31r+57)(4r^2+47r+135)}$$

Which for the root $r = 2$ becomes

$$a_7 = -\frac{4037}{72268875}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r(4r+7)}$	$-\frac{1}{30}$
a_3	$-\frac{4r}{4r^2+15r+11}$	$-\frac{8}{57}$
a_4	$\frac{1}{16r^4+120r^3+281r^2+210r}$	$\frac{1}{2760}$
a_5	$\frac{32r^3+120r^2+164r+88}{(4r+7)r(4r^2+15r+11)(4r^2+31r+57)}$	$\frac{64}{12825}$
a_6	$\frac{256r^6+2688r^5+10256r^4+16848r^3+10076r^2-15r-11}{(4r^2+15r+11)r(16r^3+120r^2+281r+210)(4r^2+39r+92)}$	$\frac{147181}{9753840}$
a_7	$-\frac{4(48r^5+552r^4+2567r^3+6075r^2+7147r+3168)}{(4r+7)(4r^2+23r+30)r(4r^2+15r+11)(4r^2+31r+57)(4r^2+47r+135)}$	$-\frac{4037}{72268875}$

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x^2(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots) \\ &= x^2 \left(1 - \frac{x^2}{30} - \frac{8x^3}{57} + \frac{x^4}{2760} + \frac{64x^5}{12825} + \frac{147181x^6}{9753840} - \frac{4037x^7}{72268875} + O(x^8) \right) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$b_1 = 0$$

Substituting $n = 2$ in Eq. (2B) gives

$$b_2 = -\frac{1}{r(4r+7)}$$

For $3 \leq n$ the recursive equation is

$$4b_n(n+r)(n+r-1) + 4b_{n-3}(n-3+r) - 5b_n(n+r) + b_{n-2} + 2b_n = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = -\frac{4nb_{n-3} + 4rb_{n-3} - 12b_{n-3} + b_{n-2}}{4n^2 + 8nr + 4r^2 - 9n - 9r + 2} \quad (4)$$

Which for the root $r = \frac{1}{4}$ becomes

$$b_n = \frac{-4nb_{n-3} + 11b_{n-3} - b_{n-2}}{n(4n-7)} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = \frac{1}{4}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{1}{r(4r+7)}$	$-\frac{1}{2}$

For $n = 3$, using the above recursive equation gives

$$b_3 = -\frac{4r}{4r^2 + 15r + 11}$$

Which for the root $r = \frac{1}{4}$ becomes

$$b_3 = -\frac{1}{15}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{1}{r(4r+7)}$	$-\frac{1}{2}$
b_3	$-\frac{4r}{4r^2+15r+11}$	$-\frac{1}{15}$

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{1}{16r^4 + 120r^3 + 281r^2 + 210r}$$

Which for the root $r = \frac{1}{4}$ becomes

$$b_4 = \frac{1}{72}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{1}{r(4r+7)}$	$-\frac{1}{2}$
b_3	$-\frac{4r}{4r^2+15r+11}$	$-\frac{1}{15}$
b_4	$\frac{1}{16r^4+120r^3+281r^2+210r}$	$\frac{1}{72}$

For $n = 5$, using the above recursive equation gives

$$b_5 = \frac{32r^3 + 120r^2 + 164r + 88}{(4r + 7)r(4r^2 + 15r + 11)(4r^2 + 31r + 57)}$$

Which for the root $r = \frac{1}{4}$ becomes

$$b_5 = \frac{137}{1950}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{1}{r(4r+7)}$	$-\frac{1}{2}$
b_3	$-\frac{4r}{4r^2+15r+11}$	$-\frac{1}{15}$
b_4	$\frac{1}{16r^4+120r^3+281r^2+210r}$	$\frac{1}{72}$
b_5	$\frac{32r^3+120r^2+164r+88}{(4r+7)r(4r^2+15r+11)(4r^2+31r+57)}$	$\frac{137}{1950}$

For $n = 6$, using the above recursive equation gives

$$b_6 = \frac{256r^6 + 2688r^5 + 10256r^4 + 16848r^3 + 10076r^2 - 15r - 11}{(4r^2 + 15r + 11)r(16r^3 + 120r^2 + 281r + 210)(4r^2 + 39r + 92)}$$

Which for the root $r = \frac{1}{4}$ becomes

$$b_6 = \frac{307}{36720}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{1}{r(4r+7)}$	$-\frac{1}{2}$
b_3	$-\frac{4r}{4r^2+15r+11}$	$-\frac{1}{15}$
b_4	$\frac{1}{16r^4+120r^3+281r^2+210r}$	$\frac{1}{72}$
b_5	$\frac{32r^3+120r^2+164r+88}{(4r+7)r(4r^2+15r+11)(4r^2+31r+57)}$	$\frac{137}{1950}$
b_6	$\frac{256r^6+2688r^5+10256r^4+16848r^3+10076r^2-15r-11}{(4r^2+15r+11)r(16r^3+120r^2+281r+210)(4r^2+39r+92)}$	$\frac{307}{36720}$

For $n = 7$, using the above recursive equation gives

$$b_7 = -\frac{4(48r^5 + 552r^4 + 2567r^3 + 6075r^2 + 7147r + 3168)}{(4r + 7)(4r^2 + 23r + 30)r(4r^2 + 15r + 11)(4r^2 + 31r + 57)(4r^2 + 47r + 135)}$$

Which for the root $r = \frac{1}{4}$ becomes

$$b_7 = -\frac{7169}{3439800}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{1}{r(4r+7)}$	$-\frac{1}{2}$
b_3	$-\frac{4r}{4r^2+15r+11}$	$-\frac{1}{15}$
b_4	$\frac{1}{16r^4+120r^3+281r^2+210r}$	$\frac{1}{72}$
b_5	$\frac{32r^3+120r^2+164r+88}{(4r+7)r(4r^2+15r+11)(4r^2+31r+57)}$	$\frac{137}{1950}$
b_6	$\frac{256r^6+2688r^5+10256r^4+16848r^3+10076r^2-15r-11}{(4r^2+15r+11)r(16r^3+120r^2+281r+210)(4r^2+39r+92)}$	$\frac{307}{36720}$
b_7	$-\frac{4(48r^5+552r^4+2567r^3+6075r^2+7147r+3168)}{(4r+7)(4r^2+23r+30)r(4r^2+15r+11)(4r^2+31r+57)(4r^2+47r+135)}$	$-\frac{7169}{3439800}$

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned} y_2(x) &= x^2(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 + b_7x^7 + b_8x^8 \dots) \\ &= x^{\frac{1}{4}} \left(1 - \frac{x^2}{2} - \frac{x^3}{15} + \frac{x^4}{72} + \frac{137x^5}{1950} + \frac{307x^6}{36720} - \frac{7169x^7}{3439800} + O(x^8) \right) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= c_1x^2 \left(1 - \frac{x^2}{30} - \frac{8x^3}{57} + \frac{x^4}{2760} + \frac{64x^5}{12825} + \frac{147181x^6}{9753840} - \frac{4037x^7}{72268875} + O(x^8) \right) \\ &\quad + c_2x^{\frac{1}{4}} \left(1 - \frac{x^2}{2} - \frac{x^3}{15} + \frac{x^4}{72} + \frac{137x^5}{1950} + \frac{307x^6}{36720} - \frac{7169x^7}{3439800} + O(x^8) \right) \end{aligned}$$

Hence the final solution is

$$\begin{aligned} y &= y_h \\ &= c_1x^2 \left(1 - \frac{x^2}{30} - \frac{8x^3}{57} + \frac{x^4}{2760} + \frac{64x^5}{12825} + \frac{147181x^6}{9753840} - \frac{4037x^7}{72268875} + O(x^8) \right) \\ &\quad + c_2x^{\frac{1}{4}} \left(1 - \frac{x^2}{2} - \frac{x^3}{15} + \frac{x^4}{72} + \frac{137x^5}{1950} + \frac{307x^6}{36720} - \frac{7169x^7}{3439800} + O(x^8) \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 x^2 \left(1 - \frac{x^2}{30} - \frac{8x^3}{57} + \frac{x^4}{2760} + \frac{64x^5}{12825} + \frac{147181x^6}{9753840} - \frac{4037x^7}{72268875} + O(x^8) \right) \\ + c_2 x^{\frac{1}{4}} \left(1 - \frac{x^2}{2} - \frac{x^3}{15} + \frac{x^4}{72} + \frac{137x^5}{1950} + \frac{307x^6}{36720} - \frac{7169x^7}{3439800} + O(x^8) \right) \quad (1)$$

Verification of solutions

$$y = c_1 x^2 \left(1 - \frac{x^2}{30} - \frac{8x^3}{57} + \frac{x^4}{2760} + \frac{64x^5}{12825} + \frac{147181x^6}{9753840} - \frac{4037x^7}{72268875} + O(x^8) \right) \\ + c_2 x^{\frac{1}{4}} \left(1 - \frac{x^2}{2} - \frac{x^3}{15} + \frac{x^4}{72} + \frac{137x^5}{1950} + \frac{307x^6}{36720} - \frac{7169x^7}{3439800} + O(x^8) \right)$$

Verified OK.

17.9.1 Maple step by step solution

Let's solve

$$4x^2 y'' + (4x^4 - 5x) y' + y(x^2 + 2) = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{(x^2+2)y}{4x^2} - \frac{(4x^3-5)y'}{4x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{(4x^3-5)y'}{4x} + \frac{(x^2+2)y}{4x^2} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$\left[P_2(x) = \frac{4x^3-5}{4x}, P_3(x) = \frac{x^2+2}{4x^2} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = -\frac{5}{4}$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = \frac{1}{2}$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$4x^2y'' + x(4x^3 - 5)y' + y(x^2 + 2) = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 0..2$

$$x^m \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

- Shift index using $k \rightarrow k - m$

$$x^m \cdot y = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

- Convert $x^m \cdot y'$ to series expansion for $m = 1..4$

$$x^m \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1+m}$$

- Shift index using $k \rightarrow k + 1 - m$

$$x^m \cdot y' = \sum_{k=-1+m}^{\infty} a_{k+1-m} (k+1-m+r) x^{k+r}$$

- Convert $x^2 \cdot y''$ to series expansion

$$x^2 \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(-1+4r)(-2+r)x^r + a_1(3+4r)(-1+r)x^{1+r} + (a_2(7+4r)r + a_0)x^{2+r} + \left(\sum_{k=3}^{\infty} (a_k(4k+4r)(k+r-1) + a_{k-1}(k+r)(k+r-1) + a_{k-2}(k+r)(k+r-1))x^{k+r} \right)$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$(-1+4r)(-2+r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \left\{ 2, \frac{1}{4} \right\}$$

- The coefficients of each power of x must be 0

$$[a_1(3 + 4r)(-1 + r) = 0, a_2(7 + 4r)r + a_0 = 0]$$

- Solve for the dependent coefficient(s)

$$\left\{ a_1 = 0, a_2 = -\frac{a_0}{r(7+4r)} \right\}$$

- Each term in the series must be 0, giving the recursion relation

$$a_k(4k + 4r - 1)(k + r - 2) + a_{k-2} + 4a_{k-3}(k - 3 + r) = 0$$

- Shift index using $k \rightarrow k + 3$

$$a_{k+3}(4k + 11 + 4r)(k + 1 + r) + a_{k+1} + 4a_k(k + r) = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+3} = -\frac{4ka_k + 4ra_k + a_{k+1}}{(4k+11+4r)(k+1+r)}$$

- Recursion relation for $r = 2$

$$a_{k+3} = -\frac{4ka_k + 8a_k + a_{k+1}}{(4k+19)(k+3)}$$

- Solution for $r = 2$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+2}, a_{k+3} = -\frac{4ka_k + 8a_k + a_{k+1}}{(4k+19)(k+3)}, a_1 = 0, a_2 = -\frac{a_0}{30} \right]$$

- Recursion relation for $r = \frac{1}{4}$

$$a_{k+3} = -\frac{4ka_k + a_k + a_{k+1}}{(4k+12)(k+\frac{5}{4})}$$

- Solution for $r = \frac{1}{4}$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+\frac{1}{4}}, a_{k+3} = -\frac{4ka_k + a_k + a_{k+1}}{(4k+12)(k+\frac{5}{4})}, a_1 = 0, a_2 = -\frac{a_0}{2} \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^{k+2} \right) + \left(\sum_{k=0}^{\infty} b_k x^{k+\frac{1}{4}} \right), a_{k+3} = -\frac{4ka_k + 8a_k + a_{k+1}}{(4k+19)(k+3)}, a_1 = 0, a_2 = -\frac{a_0}{30}, b_{k+3} = -\frac{4kb_k + b_k + a_{k+1}}{(4k+12)(k+\frac{5}{4})} \right]$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
  -> Bessel
  -> elliptic
  -> Legendre
  -> Whittaker
    -> hyper3: Equivalence to 1F1 under a power @ Moebius
  -> hypergeometric
    -> heuristic approach
    -> hyper3: Equivalence to 2F1, 1F1 or 0F1 under a power @ Moebius
  -> Mathieu
    -> Equivalence to the rational form of Mathieu ODE under a power @ Moebius
trying a solution in terms of MeijerG functions
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form  $r_0(x) * Y + r_1(x) * Y$  where  $Y = \exp(\int(r(x), dx)) * 2F1([a$ 
  trying a symmetry of the form [xi=0, eta=F(x)]
  trying differential order: 2; exact nonlinear
  trying symmetries linear in x and y(x)
  trying to convert to a linear ODE with constant coefficients
  trying 2nd order, integrating factor of the form  $\mu(x,y)$ 
  -> Trying a solution in terms of special functions:
    -> Bessel
    -> elliptic
    -> Legendre
    -> Whittaker
      -> hyper3: Equivalence to 1F1 under a power @ Moebius
    -> hypergeometric
      -> heuristic approach
      -> hyper3: Equivalence to 2F1, 1F1 or 0F1 under a power @ Moebius
    -> Mathieu
      -> Equivalence to the rational form of Mathieu ODE under a power @ Moebius
  trying 2nd order exact linear
  trying symmetries linear in x and y(x)
  trying to convert to a linear ODE with constant coefficients
  trying to convert to an ODE of Bessel type
trying to convert to an ODE of Bessel type
-> trying reduction of order to Bessel
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 51

Order:=8;

```
dsolve(4*x^2*dif(y(x),x$2)+(4*x^4-5*x)*dif(y(x),x)+(x^2+2)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^{\frac{1}{4}} \left(1 - \frac{1}{2}x^2 - \frac{1}{15}x^3 + \frac{1}{72}x^4 + \frac{137}{1950}x^5 + \frac{307}{36720}x^6 - \frac{7169}{3439800}x^7 + O(x^8) \right) + c_2 x^2 \left(1 - \frac{1}{30}x^2 - \frac{8}{57}x^3 + \frac{1}{2760}x^4 + \frac{64}{12825}x^5 + \frac{147181}{9753840}x^6 - \frac{4037}{72268875}x^7 + O(x^8) \right)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 106

```
AsymptoticDSolveValue[4*x^2*y'[x]+(4*x^4-5*x)*y'[x]+(x^2+2)*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(-\frac{4037x^7}{72268875} + \frac{147181x^6}{9753840} + \frac{64x^5}{12825} + \frac{x^4}{2760} - \frac{8x^3}{57} - \frac{x^2}{30} + 1 \right) x^2 + c_2 \left(-\frac{7169x^7}{3439800} + \frac{307x^6}{36720} + \frac{137x^5}{1950} + \frac{x^4}{72} - \frac{x^3}{15} - \frac{x^2}{2} + 1 \right) \sqrt[4]{x}$$

17.10 problem 2(d)

Internal problem ID [6049]

Internal file name [OUTPUT/5297_Sunday_June_05_2022_03_31_14_PM_49103670/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 154

Problem number: 2(d).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Complex roots"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$x^2y'' + (-3x^2 + x)y' + e^xy = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2y'' + (-3x^2 + x)y' + e^xy = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{3x-1}{x}$$
$$q(x) = \frac{e^x}{x^2}$$

Table 222: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{3x-1}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = \frac{e^x}{x^2}$	
singularity	type
$x = 0$	“regular”
$x = \infty$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0, \infty]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2 y'' + (-3x^2 + x) y' + e^x y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned} & x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\ & + (-3x^2 + x) \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + e^x \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \end{aligned} \tag{1}$$

Expanding e^x as Taylor series around $x = 0$ and keeping only the first 8 terms gives

$$\begin{aligned} e^x &= 1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{720}x^6 + \frac{1}{5040}x^7 + \frac{1}{40320}x^8 + \dots \\ &= 1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{720}x^6 + \frac{1}{5040}x^7 + \frac{1}{40320}x^8 \end{aligned}$$

Which simplifies to

$$\begin{aligned} &\left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-3x^{1+n+r} a_n (n+r)) \\ &+ \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) \\ &+ \left(\sum_{n=0}^{\infty} x^{1+n+r} a_n \right) + \left(\sum_{n=0}^{\infty} \frac{x^{n+r+2} a_n}{2} \right) + \left(\sum_{n=0}^{\infty} \frac{x^{n+r+3} a_n}{6} \right) \quad (2A) \\ &+ \left(\sum_{n=0}^{\infty} \frac{x^{n+r+4} a_n}{24} \right) + \left(\sum_{n=0}^{\infty} \frac{x^{n+r+5} a_n}{120} \right) + \left(\sum_{n=0}^{\infty} \frac{x^{n+r+6} a_n}{720} \right) \\ &+ \left(\sum_{n=0}^{\infty} \frac{x^{n+r+7} a_n}{5040} \right) + \left(\sum_{n=0}^{\infty} \frac{x^{n+r+8} a_n}{40320} \right) = 0 \end{aligned}$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=0}^{\infty} (-3x^{1+n+r} a_n (n+r)) &= \sum_{n=1}^{\infty} (-3a_{n-1} (n+r-1) x^{n+r}) \\ \sum_{n=0}^{\infty} x^{1+n+r} a_n &= \sum_{n=1}^{\infty} a_{n-1} x^{n+r} \\ \sum_{n=0}^{\infty} \frac{x^{n+r+2} a_n}{2} &= \sum_{n=2}^{\infty} \frac{a_{n-2} x^{n+r}}{2} \\ \sum_{n=0}^{\infty} \frac{x^{n+r+3} a_n}{6} &= \sum_{n=3}^{\infty} \frac{a_{n-3} x^{n+r}}{6} \\ \sum_{n=0}^{\infty} \frac{x^{n+r+4} a_n}{24} &= \sum_{n=4}^{\infty} \frac{a_{n-4} x^{n+r}}{24} \end{aligned}$$

$$\begin{aligned}\sum_{n=0}^{\infty} \frac{x^{n+r+5} a_n}{120} &= \sum_{n=5}^{\infty} \frac{a_{n-5} x^{n+r}}{120} \\ \sum_{n=0}^{\infty} \frac{x^{n+r+6} a_n}{720} &= \sum_{n=6}^{\infty} \frac{a_{n-6} x^{n+r}}{720} \\ \sum_{n=0}^{\infty} \frac{x^{n+r+7} a_n}{5040} &= \sum_{n=7}^{\infty} \frac{a_{n-7} x^{n+r}}{5040} \\ \sum_{n=0}^{\infty} \frac{x^{n+r+8} a_n}{40320} &= \sum_{n=8}^{\infty} \frac{a_{n-8} x^{n+r}}{40320}\end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n + r$.

$$\begin{aligned}&\left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=1}^{\infty} (-3a_{n-1} (n+r-1) x^{n+r}) \\ &+ \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) \\ &+ \left(\sum_{n=1}^{\infty} a_{n-1} x^{n+r} \right) + \left(\sum_{n=2}^{\infty} \frac{a_{n-2} x^{n+r}}{2} \right) + \left(\sum_{n=3}^{\infty} \frac{a_{n-3} x^{n+r}}{6} \right) \\ &+ \left(\sum_{n=4}^{\infty} \frac{a_{n-4} x^{n+r}}{24} \right) + \left(\sum_{n=5}^{\infty} \frac{a_{n-5} x^{n+r}}{120} \right) + \left(\sum_{n=6}^{\infty} \frac{a_{n-6} x^{n+r}}{720} \right) \\ &+ \left(\sum_{n=7}^{\infty} \frac{a_{n-7} x^{n+r}}{5040} \right) + \left(\sum_{n=8}^{\infty} \frac{a_{n-8} x^{n+r}}{40320} \right) = 0\end{aligned}\tag{2B}$$

The indicial equation is obtained from $n = 0$. From Eq (2B) this gives

$$x^{n+r} a_n (n+r) (n+r-1) + x^{n+r} a_n (n+r) + a_n x^{n+r} = 0$$

When $n = 0$ the above becomes

$$x^r a_0 r (-1+r) + x^r a_0 r + a_0 x^r = 0$$

Or

$$(x^r r (-1+r) + x^r r + x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(r^2 + 1) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$r^2 + 1 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= i \\ r_2 &= -i \end{aligned}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(r^2 + 1) x^r = 0$$

Solving for r gives the roots of the indicial equation as Since the roots are complex conjugates, then two linearly independent solutions can be constructed using

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^{n+i} \\ y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n-i} \end{aligned}$$

$y_1(x)$ is found first. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = \frac{3r - 1}{r^2 + 2r + 2}$$

Substituting $n = 2$ in Eq. (2B) gives

$$a_2 = \frac{17r^2 + 4r - 6}{2(r^2 + 2r + 2)(r^2 + 4r + 5)}$$

Substituting $n = 3$ in Eq. (2B) gives

$$a_3 = \frac{-r^4 + 138r^3 + 243r^2 - 45r - 85}{6(r^2 + 2r + 2)(r^2 + 4r + 5)(r^2 + 6r + 10)}$$

Substituting $n = 4$ in Eq. (2B) gives

$$a_4 = \frac{-r^6 - 36r^5 + 1345r^4 + 6100r^3 + 5156r^2 - 3044r - 2260}{24(r^2 + 2r + 2)(r^2 + 4r + 5)(r^2 + 6r + 10)(r^2 + 8r + 17)}$$

Substituting $n = 5$ in Eq. (2B) gives

$$a_5 = \frac{-r^8 - 50r^7 - 1194r^6 + 11665r^5 + 128676r^4 + 315715r^3 + 145319r^2 - 192430r - 97100}{120(r^2 + 2r + 2)(r^2 + 4r + 5)(r^2 + 6r + 10)(r^2 + 8r + 17)(r^2 + 10r + 26)}$$

Substituting $n = 6$ in Eq. (2B) gives

$$a_6 = \frac{-r^{10} - 66r^9 - 2102r^8 - 42690r^7 - 76157r^6 + 2029806r^5 + 11717702r^4 + 20421330r^3 + 3645968r^2 - 192430r - 97100}{720(r^2 + 2r + 2)(r^2 + 4r + 5)(r^2 + 6r + 10)(r^2 + 8r + 17)(r^2 + 10r + 26)(r^2 + 12r + 35)}$$

Substituting $n = 7$ in Eq. (2B) gives

$$a_7 = \frac{-r^{12} - 84r^{11} - 3401r^{10} - 89145r^9 - 1687260r^8 - 12804498r^7 - 15286821r^6 + 243174141r^5 + 112600000r^4 - 192430r - 97100}{5040(r^2 + 2r + 2)(r^2 + 4r + 5)(r^2 + 6r + 10)(r^2 + 8r + 17)(r^2 + 10r + 26)(r^2 + 12r + 35)(r^2 + 14r + 49)}$$

For $8 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) - 3a_{n-1}(n+r-1) + a_n(n+r) + a_n + a_{n-1} + \frac{a_{n-2}}{2} + \frac{a_{n-3}}{6} + \frac{a_{n-4}}{24} + \frac{a_{n-5}}{120} + \frac{a_{n-6}}{720} + \frac{a_{n-7}}{5040} + \frac{a_{n-8}}{40320} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{120960na_{n-1} + 120960ra_{n-1} - a_{n-8} - 8a_{n-7} - 56a_{n-6} - 336a_{n-5} - 1680a_{n-4} - 6720a_{n-3} - 20160a_{n-2}}{40320n^2 + 80640nr + 40320r^2 + 40320} \quad (4)$$

Which for the root $r = i$ becomes

$$a_n = -\frac{40320(4 - 3i - 3n)a_{n-1} + a_{n-8} + 8a_{n-7} + 56a_{n-6} + 336a_{n-5} + 1680a_{n-4} + 6720a_{n-3} + 20160a_{n-2}}{40320n(2i + n)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = i$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$
a_0	1
a_1	$\frac{3r-1}{r^2+2r+2}$
a_2	$\frac{17r^2+4r-6}{2(r^2+2r+2)(r^2+4r+5)}$
a_3	$\frac{-r^4+138r^3+243r^2-45r-85}{6(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)}$
a_4	$\frac{-r^6-36r^5+1345r^4+6100r^3+5156r^2-3044r-2260}{24(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)}$
a_5	$\frac{-r^8-50r^7-1194r^6+11665r^5+128676r^4+315715r^3+145319r^2-192430r-97100}{120(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)(r^2+10r+26)}$
a_6	$\frac{-r^{10}-66r^9-2102r^8-42690r^7-76157r^6+2029806r^5+11717702r^4+20421330r^3+3645968r^2-15274320r-6037400}{720(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)(r^2+10r+26)(r^2+12r+37)}$
a_7	$\frac{-r^{12}-84r^{11}-3401r^{10}-89145r^9-1687260r^8-12804498r^7-15286821r^6+243174141r^5+1126587431r^4+1540608062r^3-207286888r^2-155040}{5040(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)(r^2+10r+26)(r^2+12r+37)(r^2+14r+50)}$

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned}
y_1(x) &= x^i (a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots) \\
&= x^i \left(1 + (1+i)x + \left(\frac{7}{16} + \frac{13i}{16} \right) x^2 + \left(\frac{7}{39} + \frac{395i}{936} \right) x^3 + \left(\frac{2117}{29952} + \frac{5197i}{29952} \right) x^4 \right. \\
&\quad + \left(\frac{5521}{217152} + \frac{642043i}{10857600} \right) x^5 + \left(\frac{782461}{97718400} + \frac{8813057i}{521164800} \right) x^6 \\
&\quad \left. + \left(\frac{1238071931}{580056422400} + \frac{3271304833i}{812078991360} \right) x^7 + O(x^8) \right)
\end{aligned}$$

The second solution $y_2(x)$ is found by taking the complex conjugate of $y_1(x)$ which gives

$$\begin{aligned}
y_2(x) &= x^{-i} \left(1 + (1-i)x + \left(\frac{7}{16} - \frac{13i}{16} \right) x^2 + \left(\frac{7}{39} - \frac{395i}{936} \right) x^3 + \left(\frac{2117}{29952} - \frac{5197i}{29952} \right) x^4 \right. \\
&\quad + \left(\frac{5521}{217152} - \frac{642043i}{10857600} \right) x^5 + \left(\frac{782461}{97718400} - \frac{8813057i}{521164800} \right) x^6 \\
&\quad \left. + \left(\frac{1238071931}{580056422400} - \frac{3271304833i}{812078991360} \right) x^7 + O(x^8) \right)
\end{aligned}$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 y_1(x) + c_2 y_2(x)$$

$$\begin{aligned}
&= c_1 x^i \left(1 + (1+i)x + \left(\frac{7}{16} + \frac{13i}{16} \right) x^2 + \left(\frac{7}{39} + \frac{395i}{936} \right) x^3 + \left(\frac{2117}{29952} + \frac{5197i}{29952} \right) x^4 \right. \\
&\quad + \left(\frac{5521}{217152} + \frac{642043i}{10857600} \right) x^5 + \left(\frac{782461}{97718400} + \frac{8813057i}{521164800} \right) x^6 \\
&\quad \left. + \left(\frac{1238071931}{580056422400} + \frac{3271304833i}{812078991360} \right) x^7 + O(x^8) \right) \\
&+ c_2 x^{-i} \left(1 + (1-i)x + \left(\frac{7}{16} - \frac{13i}{16} \right) x^2 + \left(\frac{7}{39} - \frac{395i}{936} \right) x^3 + \left(\frac{2117}{29952} - \frac{5197i}{29952} \right) x^4 \right. \\
&\quad + \left(\frac{5521}{217152} - \frac{642043i}{10857600} \right) x^5 + \left(\frac{782461}{97718400} - \frac{8813057i}{521164800} \right) x^6 \\
&\quad \left. + \left(\frac{1238071931}{580056422400} - \frac{3271304833i}{812078991360} \right) x^7 + O(x^8) \right)
\end{aligned}$$

Hence the final solution is

$$y = y_h$$

$$\begin{aligned}
&= c_1 x^i \left(1 + (1+i)x + \left(\frac{7}{16} + \frac{13i}{16} \right) x^2 + \left(\frac{7}{39} + \frac{395i}{936} \right) x^3 + \left(\frac{2117}{29952} + \frac{5197i}{29952} \right) x^4 \right. \\
&\quad + \left(\frac{5521}{217152} + \frac{642043i}{10857600} \right) x^5 + \left(\frac{782461}{97718400} + \frac{8813057i}{521164800} \right) x^6 \\
&\quad \left. + \left(\frac{1238071931}{580056422400} + \frac{3271304833i}{812078991360} \right) x^7 + O(x^8) \right) \\
&+ c_2 x^{-i} \left(1 + (1-i)x + \left(\frac{7}{16} - \frac{13i}{16} \right) x^2 + \left(\frac{7}{39} - \frac{395i}{936} \right) x^3 + \left(\frac{2117}{29952} - \frac{5197i}{29952} \right) x^4 \right. \\
&\quad + \left(\frac{5521}{217152} - \frac{642043i}{10857600} \right) x^5 + \left(\frac{782461}{97718400} - \frac{8813057i}{521164800} \right) x^6 \\
&\quad \left. + \left(\frac{1238071931}{580056422400} - \frac{3271304833i}{812078991360} \right) x^7 + O(x^8) \right)
\end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}y = c_1 x^i & \left(1 + (1+i)x + \left(\frac{7}{16} + \frac{13i}{16} \right) x^2 + \left(\frac{7}{39} + \frac{395i}{936} \right) x^3 + \left(\frac{2117}{29952} + \frac{5197i}{29952} \right) x^4 \right. \\ & + \left(\frac{5521}{217152} + \frac{642043i}{10857600} \right) x^5 + \left(\frac{782461}{97718400} + \frac{8813057i}{521164800} \right) x^6 \\ & \left. + \left(\frac{1238071931}{580056422400} + \frac{3271304833i}{812078991360} \right) x^7 + O(x^8) \right) \\ & + c_2 x^{-i} \left(1 + (1-i)x + \left(\frac{7}{16} - \frac{13i}{16} \right) x^2 + \left(\frac{7}{39} - \frac{395i}{936} \right) x^3 + \left(\frac{2117}{29952} - \frac{5197i}{29952} \right) x^4 \right. \\ & + \left(\frac{5521}{217152} - \frac{642043i}{10857600} \right) x^5 + \left(\frac{782461}{97718400} - \frac{8813057i}{521164800} \right) x^6 \\ & \left. + \left(\frac{1238071931}{580056422400} - \frac{3271304833i}{812078991360} \right) x^7 + O(x^8) \right) \quad (1)\end{aligned}$$

Verification of solutions

$$\begin{aligned}y = c_1 x^i & \left(1 + (1+i)x + \left(\frac{7}{16} + \frac{13i}{16} \right) x^2 + \left(\frac{7}{39} + \frac{395i}{936} \right) x^3 + \left(\frac{2117}{29952} + \frac{5197i}{29952} \right) x^4 \right. \\ & + \left(\frac{5521}{217152} + \frac{642043i}{10857600} \right) x^5 + \left(\frac{782461}{97718400} + \frac{8813057i}{521164800} \right) x^6 \\ & \left. + \left(\frac{1238071931}{580056422400} + \frac{3271304833i}{812078991360} \right) x^7 + O(x^8) \right) \\ & + c_2 x^{-i} \left(1 + (1-i)x + \left(\frac{7}{16} - \frac{13i}{16} \right) x^2 + \left(\frac{7}{39} - \frac{395i}{936} \right) x^3 + \left(\frac{2117}{29952} - \frac{5197i}{29952} \right) x^4 \right. \\ & + \left(\frac{5521}{217152} - \frac{642043i}{10857600} \right) x^5 + \left(\frac{782461}{97718400} - \frac{8813057i}{521164800} \right) x^6 \\ & \left. + \left(\frac{1238071931}{580056422400} - \frac{3271304833i}{812078991360} \right) x^7 + O(x^8) \right)\end{aligned}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form  $r_0(x) * Y + r_1(x) * Y$  where  $Y = \exp(\int(r(x), dx)) * 2F1([a$ 
-> Trying changes of variables to rationalize or make the ODE simpler
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
    -> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
    -> trying a solution of the form  $r_0(x) * Y + r_1(x) * Y$  where  $Y = \exp(\int(r(x), dx)) * 2F1$ 
        trying a symmetry of the form [xi=0, eta=F(x)]
        trying 2nd order exact linear
        trying symmetries linear in x and y(x)
        trying to convert to a linear ODE with constant coefficients
<- unable to find a useful change of variables
    trying a symmetry of the form [xi=0, eta=F(x)]
    trying differential order: 2; exact nonlinear
    trying symmetries linear in x and y(x)
    trying to convert to a linear ODE with constant coefficients
    trying 2nd order, integrating factor of the form  $\mu(x,y)$ 
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
    -> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
    -> trying a solution of the form  $r_0(x) * Y + r_1(x) * Y$  where  $Y = \exp(\int(r(x), dx)) * 2F1$ 
    -> Trying changes of variables to rationalize or make the ODE simpler
        trying a symmetry of the form [xi=0, eta=F(x)]
        checking if the LODE is missing y
        -> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
        -> trying a solution of the form  $r_0(x) * Y + r_1(x) * Y$  where  $Y = \exp(\int(r(x), dx)) * 2F1$ 
            trying a symmetry of the form [xi=0, eta=F(x)]
            trying 2nd order exact linear
            trying symmetries linear in x and y(x)
            trying to convert to a linear ODE with constant coefficients
<- unable to find a useful change of variables
    trying a symmetry of the form [xi=0, eta=F(x)]
trying to convert to an ODE of Bessel type
-> trying reduction of order to Riccati
    trying Riccati sub-methods:
        trying Riccati_symmetries
        -> trying a symmetry pattern of the form  $[F(x)*G(y), 0]$ 
        -> trying a symmetry pattern of the form  $[0, F(x)*G(y)]$ 
        -> trying a symmetry pattern of the form  $[F(x), G(x)*y+H(x)]$ 
--- Trying Lie symmetry methods, 2nd order ---
-> Computing symmetries using:  $\text{var} = 2; [0, -]$ 
```

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 85

Order:=8;

dsolve(x^2*diff(y(x),x\$2)+(x-3*x^2)*diff(y(x),x)+exp(x)*y(x)=0,y(x),type='series',x=0);

$$\begin{aligned}
 y(x) = & c_1 x^{-i} \left(1 + (1-i)x + \left(\frac{7}{16} - \frac{13i}{16} \right) x^2 + \left(\frac{7}{39} - \frac{395i}{936} \right) x^3 + \left(\frac{2117}{29952} - \frac{5197i}{29952} \right) x^4 \right. \\
 & + \left(\frac{5521}{217152} - \frac{642043i}{10857600} \right) x^5 + \left(\frac{782461}{97718400} - \frac{8813057i}{521164800} \right) x^6 \\
 & \left. + \left(\frac{1238071931}{580056422400} - \frac{3271304833i}{812078991360} \right) x^7 + O(x^8) \right) \\
 & + c_2 x^i \left(1 + (1+i)x + \left(\frac{7}{16} + \frac{13i}{16} \right) x^2 + \left(\frac{7}{39} + \frac{395i}{936} \right) x^3 + \left(\frac{2117}{29952} + \frac{5197i}{29952} \right) x^4 \right. \\
 & + \left(\frac{5521}{217152} + \frac{642043i}{10857600} \right) x^5 + \left(\frac{782461}{97718400} + \frac{8813057i}{521164800} \right) x^6 \\
 & \left. + \left(\frac{1238071931}{580056422400} + \frac{3271304833i}{812078991360} \right) x^7 + O(x^8) \right)
 \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.043 (sec). Leaf size: 122

AsymptoticDSolveValue[x^2*y'[x]+(x-3*x^2)*y'[x]+Exp[x]*y[x]==0,y[x],{x,0,7}]

$$\begin{aligned}
 y(x) \rightarrow & \left(\frac{1}{97718400} + \frac{11i}{1563494400} \right) c_1 x^i \left((1302761 + 756800i)x^6 \right. \\
 & + (4384656 + 2763936i)x^5 + (12605400 + 8289000i)x^4 \\
 & + (31161600 + 19814400i)x^3 + (66096000 + 33955200i)x^2 \\
 & \left. + (111974400 + 20736000i)x + (66355200 - 45619200i) \right) \\
 & - \left(\frac{11}{1563494400} + \frac{i}{97718400} \right) c_2 x^{-i} \left((756800 + 1302761i)x^6 \right. \\
 & + (2763936 + 4384656i)x^5 + (8289000 + 12605400i)x^4 \\
 & + (19814400 + 31161600i)x^3 + (33955200 + 66096000i)x^2 \\
 & \left. + (20736000 + 111974400i)x - (45619200 - 66355200i) \right)
 \end{aligned}$$

18 Chapter 4. Linear equations with Regular Singular Points. Page 159

18.1 problem 1(a)	1470
18.2 problem 1(b)	1486
18.3 problem 2	1498

18.1 problem 1(a)

18.1.1 Maple step by step solution 1482

Internal problem ID [6050]

Internal file name [OUTPUT/5298_Sunday_June_05_2022_03_32_46_PM_42206577/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 159

Problem number: 1(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

[[_Emden , _Fowler]]

$$3x^2y'' + 5xy' + 3xy = 0$$

With the expansion point for the power series method at $x = 0$.

The ODE is

$$3x^2y'' + 5xy' + 3xy = 0$$

Or

$$x(3y''x + 3y + 5y') = 0$$

For $x \neq 0$ the above simplifies to

$$3y''x + 3y + 5y' = 0$$

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$3x^2y'' + 5xy' + 3xy = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{5}{3x}$$

$$q(x) = \frac{1}{x}$$

Table 223: Table $p(x), q(x)$ singularities.

$p(x) = \frac{5}{3x}$	
singularity	type
$x = 0$	“regular”

$q(x) = \frac{1}{x}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$3x^2y'' + 5xy' + 3xy = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned}
& 3x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\
& + 5x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + 3x \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0
\end{aligned} \tag{1}$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} 3x^{n+r} a_n (n+r)(n+r-1) \right) + \left(\sum_{n=0}^{\infty} 5x^{n+r} a_n (n+r) \right) + \left(\sum_{n=0}^{\infty} 3x^{1+n+r} a_n \right) = 0 \tag{2A}$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} 3x^{1+n+r} a_n = \sum_{n=1}^{\infty} 3a_{n-1} x^{n+r}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\left(\sum_{n=0}^{\infty} 3x^{n+r} a_n (n+r)(n+r-1) \right) + \left(\sum_{n=0}^{\infty} 5x^{n+r} a_n (n+r) \right) + \left(\sum_{n=1}^{\infty} 3a_{n-1} x^{n+r} \right) = 0 \tag{2B}$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$3x^{n+r} a_n (n+r)(n+r-1) + 5x^{n+r} a_n (n+r) = 0$$

When $n=0$ the above becomes

$$3x^r a_0 r(-1+r) + 5x^r a_0 r = 0$$

Or

$$(3x^r r(-1+r) + 5x^r r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$x^r r(2+3r) = 0$$

Since the above is true for all x then the indicial equation becomes

$$3r^2 + 2r = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 0 \\ r_2 &= -\frac{2}{3} \end{aligned}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$x^r r(2 + 3r) = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = \frac{2}{3}$ is not an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^n \\ y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n-\frac{2}{3}} \end{aligned}$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$3a_n(n+r)(n+r-1) + 5a_n(n+r) + 3a_{n-1} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{3a_{n-1}}{3n^2 + 6nr + 3r^2 + 2n + 2r} \quad (4)$$

Which for the root $r = 0$ becomes

$$a_n = -\frac{3a_{n-1}}{n(3n+2)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = -\frac{3}{3r^2 + 8r + 5}$$

Which for the root $r = 0$ becomes

$$a_1 = -\frac{3}{5}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{3}{3r^2+8r+5}$	$-\frac{3}{5}$

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{9}{(3r^2 + 8r + 5)(3r^2 + 14r + 16)}$$

Which for the root $r = 0$ becomes

$$a_2 = \frac{9}{80}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{3}{3r^2+8r+5}$	$-\frac{3}{5}$
a_2	$\frac{9}{(3r^2+8r+5)(3r^2+14r+16)}$	$\frac{9}{80}$

For $n = 3$, using the above recursive equation gives

$$a_3 = -\frac{27}{(3r^2 + 8r + 5)(3r^2 + 14r + 16)(3r^2 + 20r + 33)}$$

Which for the root $r = 0$ becomes

$$a_3 = -\frac{9}{880}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{3}{3r^2+8r+5}$	$-\frac{3}{5}$
a_2	$\frac{9}{(3r^2+8r+5)(3r^2+14r+16)}$	$\frac{9}{80}$
a_3	$-\frac{27}{(3r^2+8r+5)(3r^2+14r+16)(3r^2+20r+33)}$	$-\frac{9}{880}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{81}{(3r^2 + 8r + 5)(3r^2 + 14r + 16)(3r^2 + 20r + 33)(3r^2 + 26r + 56)}$$

Which for the root $r = 0$ becomes

$$a_4 = \frac{27}{49280}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{3}{3r^2+8r+5}$	$-\frac{3}{5}$
a_2	$\frac{9}{(3r^2+8r+5)(3r^2+14r+16)}$	$\frac{9}{80}$
a_3	$-\frac{27}{(3r^2+8r+5)(3r^2+14r+16)(3r^2+20r+33)}$	$-\frac{9}{880}$
a_4	$\frac{81}{(3r^2+8r+5)(3r^2+14r+16)(3r^2+20r+33)(3r^2+26r+56)}$	$\frac{27}{49280}$

For $n = 5$, using the above recursive equation gives

$$a_5 = -\frac{243}{(3r^2 + 8r + 5)(3r^2 + 14r + 16)(3r^2 + 20r + 33)(3r^2 + 26r + 56)(3r^2 + 32r + 85)}$$

Which for the root $r = 0$ becomes

$$a_5 = -\frac{81}{4188800}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{3}{3r^2+8r+5}$	$-\frac{3}{5}$
a_2	$\frac{9}{(3r^2+8r+5)(3r^2+14r+16)}$	$\frac{9}{80}$
a_3	$-\frac{27}{(3r^2+8r+5)(3r^2+14r+16)(3r^2+20r+33)}$	$-\frac{9}{880}$
a_4	$\frac{81}{(3r^2+8r+5)(3r^2+14r+16)(3r^2+20r+33)(3r^2+26r+56)}$	$\frac{27}{49280}$
a_5	$-\frac{243}{(3r^2+8r+5)(3r^2+14r+16)(3r^2+20r+33)(3r^2+26r+56)(3r^2+32r+85)}$	$-\frac{81}{4188800}$

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{729}{(3r^2 + 8r + 5)(3r^2 + 14r + 16)(3r^2 + 20r + 33)(3r^2 + 26r + 56)(3r^2 + 32r + 85)(3r^2 + 38r + 120)}$$

Which for the root $r = 0$ becomes

$$a_6 = \frac{81}{167552000}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{3}{3r^2+8r+5}$	$-\frac{3}{5}$
a_2	$\frac{9}{(3r^2+8r+5)(3r^2+14r+16)}$	$\frac{9}{80}$
a_3	$-\frac{27}{(3r^2+8r+5)(3r^2+14r+16)(3r^2+20r+33)}$	$-\frac{9}{880}$
a_4	$\frac{81}{(3r^2+8r+5)(3r^2+14r+16)(3r^2+20r+33)(3r^2+26r+56)}$	$\frac{27}{49280}$
a_5	$-\frac{243}{(3r^2+8r+5)(3r^2+14r+16)(3r^2+20r+33)(3r^2+26r+56)(3r^2+32r+85)}$	$-\frac{81}{4188800}$
a_6	$\frac{729}{(3r^2+8r+5)(3r^2+14r+16)(3r^2+20r+33)(3r^2+26r+56)(3r^2+32r+85)(3r^2+38r+120)}$	$\frac{81}{167552000}$

For $n = 7$, using the above recursive equation gives

$$a_7 = -\frac{2187}{(3r^2 + 8r + 5)(3r^2 + 14r + 16)(3r^2 + 20r + 33)(3r^2 + 26r + 56)(3r^2 + 32r + 85)(3r^2 + 38r + 120)}$$

Which for the root $r = 0$ becomes

$$a_7 = -\frac{243}{26975872000}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{3}{3r^2+8r+5}$	$-\frac{3}{5}$
a_2	$\frac{9}{(3r^2+8r+5)(3r^2+14r+16)}$	$\frac{9}{80}$
a_3	$-\frac{27}{(3r^2+8r+5)(3r^2+14r+16)(3r^2+20r+33)}$	$-\frac{9}{880}$
a_4	$\frac{81}{(3r^2+8r+5)(3r^2+14r+16)(3r^2+20r+33)(3r^2+26r+56)}$	$\frac{27}{49280}$
a_5	$-\frac{243}{(3r^2+8r+5)(3r^2+14r+16)(3r^2+20r+33)(3r^2+26r+56)(3r^2+32r+85)}$	$-\frac{81}{4188800}$
a_6	$\frac{729}{(3r^2+8r+5)(3r^2+14r+16)(3r^2+20r+33)(3r^2+26r+56)(3r^2+32r+85)(3r^2+38r+120)}$	$\frac{81}{167552000}$
a_7	$-\frac{2187}{(3r^2+8r+5)(3r^2+14r+16)(3r^2+20r+33)(3r^2+26r+56)(3r^2+32r+85)(3r^2+38r+120)(3r^2+44r+161)}$	$-\frac{243}{26975872000}$

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned}
 y_1(x) &= a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots \\
 &= 1 - \frac{3x}{5} + \frac{9x^2}{80} - \frac{9x^3}{880} + \frac{27x^4}{49280} - \frac{81x^5}{4188800} + \frac{81x^6}{167552000} - \frac{243x^7}{26975872000} + O(x^8)
 \end{aligned}$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. For $1 \leq n$ the recursive equation is

$$3b_n(n+r)(n+r-1) + 5b_n(n+r) + 3b_{n-1} = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = -\frac{3b_{n-1}}{3n^2 + 6nr + 3r^2 + 2n + 2r} \quad (4)$$

Which for the root $r = -\frac{2}{3}$ becomes

$$b_n = -\frac{3b_{n-1}}{n(3n-2)} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = -\frac{2}{3}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1

For $n = 1$, using the above recursive equation gives

$$b_1 = -\frac{3}{3r^2 + 8r + 5}$$

Which for the root $r = -\frac{2}{3}$ becomes

$$b_1 = -3$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{3}{3r^2+8r+5}$	-3

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{9}{(3r^2 + 8r + 5)(3r^2 + 14r + 16)}$$

Which for the root $r = -\frac{2}{3}$ becomes

$$b_2 = \frac{9}{8}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{3}{3r^2+8r+5}$	-3
b_2	$\frac{9}{(3r^2+8r+5)(3r^2+14r+16)}$	$\frac{9}{8}$

For $n = 3$, using the above recursive equation gives

$$b_3 = -\frac{27}{(3r^2 + 8r + 5)(3r^2 + 14r + 16)(3r^2 + 20r + 33)}$$

Which for the root $r = -\frac{2}{3}$ becomes

$$b_3 = -\frac{9}{56}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{3}{3r^2+8r+5}$	-3
b_2	$\frac{9}{(3r^2+8r+5)(3r^2+14r+16)}$	$\frac{9}{8}$
b_3	$-\frac{27}{(3r^2+8r+5)(3r^2+14r+16)(3r^2+20r+33)}$	$-\frac{9}{56}$

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{81}{(3r^2 + 8r + 5)(3r^2 + 14r + 16)(3r^2 + 20r + 33)(3r^2 + 26r + 56)}$$

Which for the root $r = -\frac{2}{3}$ becomes

$$b_4 = \frac{27}{2240}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{3}{3r^2+8r+5}$	-3
b_2	$\frac{9}{(3r^2+8r+5)(3r^2+14r+16)}$	$\frac{9}{8}$
b_3	$-\frac{27}{(3r^2+8r+5)(3r^2+14r+16)(3r^2+20r+33)}$	$-\frac{9}{56}$
b_4	$\frac{81}{(3r^2+8r+5)(3r^2+14r+16)(3r^2+20r+33)(3r^2+26r+56)}$	$\frac{27}{2240}$

For $n = 5$, using the above recursive equation gives

$$b_5 = -\frac{243}{(3r^2 + 8r + 5)(3r^2 + 14r + 16)(3r^2 + 20r + 33)(3r^2 + 26r + 56)(3r^2 + 32r + 85)}$$

Which for the root $r = -\frac{2}{3}$ becomes

$$b_5 = -\frac{81}{145600}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{3}{3r^2+8r+5}$	-3
b_2	$\frac{9}{(3r^2+8r+5)(3r^2+14r+16)}$	$\frac{9}{8}$
b_3	$-\frac{27}{(3r^2+8r+5)(3r^2+14r+16)(3r^2+20r+33)}$	$-\frac{9}{56}$
b_4	$\frac{81}{(3r^2+8r+5)(3r^2+14r+16)(3r^2+20r+33)(3r^2+26r+56)}$	$\frac{27}{2240}$
b_5	$-\frac{243}{(3r^2+8r+5)(3r^2+14r+16)(3r^2+20r+33)(3r^2+26r+56)(3r^2+32r+85)}$	$-\frac{81}{145600}$

For $n = 6$, using the above recursive equation gives

$$b_6 = \frac{729}{(3r^2 + 8r + 5)(3r^2 + 14r + 16)(3r^2 + 20r + 33)(3r^2 + 26r + 56)(3r^2 + 32r + 85)(3r^2 + 38r + 120)}$$

Which for the root $r = -\frac{2}{3}$ becomes

$$b_6 = \frac{81}{4659200}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{3}{3r^2+8r+5}$	-3
b_2	$\frac{9}{(3r^2+8r+5)(3r^2+14r+16)}$	$\frac{9}{8}$
b_3	$-\frac{27}{(3r^2+8r+5)(3r^2+14r+16)(3r^2+20r+33)}$	$-\frac{9}{56}$
b_4	$\frac{81}{(3r^2+8r+5)(3r^2+14r+16)(3r^2+20r+33)(3r^2+26r+56)}$	$\frac{27}{2240}$
b_5	$-\frac{243}{(3r^2+8r+5)(3r^2+14r+16)(3r^2+20r+33)(3r^2+26r+56)(3r^2+32r+85)}$	$-\frac{81}{145600}$
b_6	$\frac{729}{(3r^2+8r+5)(3r^2+14r+16)(3r^2+20r+33)(3r^2+26r+56)(3r^2+32r+85)(3r^2+38r+120)}$	$\frac{81}{4659200}$

For $n = 7$, using the above recursive equation gives

$$b_7 = -\frac{2187}{(3r^2 + 8r + 5)(3r^2 + 14r + 16)(3r^2 + 20r + 33)(3r^2 + 26r + 56)(3r^2 + 32r + 85)(3r^2 + 38r + 120)(3r^2 + 44r + 155)}$$

Which for the root $r = -\frac{2}{3}$ becomes

$$b_7 = -\frac{243}{619673600}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{3}{3r^2+8r+5}$	-3
b_2	$\frac{9}{(3r^2+8r+5)(3r^2+14r+16)}$	$\frac{9}{8}$
b_3	$-\frac{27}{(3r^2+8r+5)(3r^2+14r+16)(3r^2+20r+33)}$	$-\frac{9}{56}$
b_4	$\frac{81}{(3r^2+8r+5)(3r^2+14r+16)(3r^2+20r+33)(3r^2+26r+56)}$	$\frac{27}{2240}$
b_5	$-\frac{243}{(3r^2+8r+5)(3r^2+14r+16)(3r^2+20r+33)(3r^2+26r+56)(3r^2+32r+85)}$	$-\frac{81}{145600}$
b_6	$\frac{729}{(3r^2+8r+5)(3r^2+14r+16)(3r^2+20r+33)(3r^2+26r+56)(3r^2+32r+85)(3r^2+38r+120)}$	$\frac{81}{4659200}$
b_7	$-\frac{2187}{(3r^2+8r+5)(3r^2+14r+16)(3r^2+20r+33)(3r^2+26r+56)(3r^2+32r+85)(3r^2+38r+120)(3r^2+44r+161)}$	$-\frac{243}{619673600}$

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned}
 y_2(x) &= 1(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 + b_7x^7 + b_8x^8 \dots) \\
 &= \frac{1 - 3x + \frac{9x^2}{8} - \frac{9x^3}{56} + \frac{27x^4}{2240} - \frac{81x^5}{145600} + \frac{81x^6}{4659200} - \frac{243x^7}{619673600} + O(x^8)}{x^{\frac{2}{3}}}
 \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 y_h(x) &= c_1y_1(x) + c_2y_2(x) \\
 &= c_1 \left(1 - \frac{3x}{5} + \frac{9x^2}{80} - \frac{9x^3}{880} + \frac{27x^4}{49280} - \frac{81x^5}{4188800} + \frac{81x^6}{167552000} - \frac{243x^7}{26975872000} + O(x^8) \right) \\
 &\quad + \frac{c_2 \left(1 - 3x + \frac{9x^2}{8} - \frac{9x^3}{56} + \frac{27x^4}{2240} - \frac{81x^5}{145600} + \frac{81x^6}{4659200} - \frac{243x^7}{619673600} + O(x^8) \right)}{x^{\frac{2}{3}}}
 \end{aligned}$$

Hence the final solution is

$$\begin{aligned}
 y &= y_h \\
 &= c_1 \left(1 - \frac{3x}{5} + \frac{9x^2}{80} - \frac{9x^3}{880} + \frac{27x^4}{49280} - \frac{81x^5}{4188800} + \frac{81x^6}{167552000} - \frac{243x^7}{26975872000} + O(x^8) \right) \\
 &\quad + \frac{c_2 \left(1 - 3x + \frac{9x^2}{8} - \frac{9x^3}{56} + \frac{27x^4}{2240} - \frac{81x^5}{145600} + \frac{81x^6}{4659200} - \frac{243x^7}{619673600} + O(x^8) \right)}{x^{\frac{2}{3}}}
 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \left(1 - \frac{3x}{5} + \frac{9x^2}{80} - \frac{9x^3}{880} + \frac{27x^4}{49280} - \frac{81x^5}{4188800} + \frac{81x^6}{167552000} - \frac{243x^7}{26975872000} + O(x^8) \right) \\ + \frac{c_2 \left(1 - 3x + \frac{9x^2}{8} - \frac{9x^3}{56} + \frac{27x^4}{2240} - \frac{81x^5}{145600} + \frac{81x^6}{4659200} - \frac{243x^7}{619673600} + O(x^8) \right)}{x^{\frac{2}{3}}} \quad (1)$$

Verification of solutions

$$y = c_1 \left(1 - \frac{3x}{5} + \frac{9x^2}{80} - \frac{9x^3}{880} + \frac{27x^4}{49280} - \frac{81x^5}{4188800} + \frac{81x^6}{167552000} - \frac{243x^7}{26975872000} + O(x^8) \right) \\ + \frac{c_2 \left(1 - 3x + \frac{9x^2}{8} - \frac{9x^3}{56} + \frac{27x^4}{2240} - \frac{81x^5}{145600} + \frac{81x^6}{4659200} - \frac{243x^7}{619673600} + O(x^8) \right)}{x^{\frac{2}{3}}}$$

Verified OK.

18.1.1 Maple step by step solution

Let's solve

$$3x^2y'' + 5xy' + 3xy = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{5y'}{3x} - \frac{y}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{5y'}{3x} + \frac{y}{x} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$[P_2(x) = \frac{5}{3x}, P_3(x) = \frac{1}{x}]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = \frac{5}{3}$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 0$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$3y''x + 3y + 5y' = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert y' to series expansion

$$y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1}$$

- Shift index using $k \rightarrow k+1$

$$y' = \sum_{k=-1}^{\infty} a_{k+1} (k+1+r) x^{k+r}$$

- Convert $x \cdot y''$ to series expansion

$$x \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r-1}$$

- Shift index using $k \rightarrow k+1$

$$x \cdot y'' = \sum_{k=-1}^{\infty} a_{k+1} (k+1+r)(k+r) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0 r(2+3r) x^{-1+r} + \left(\sum_{k=0}^{\infty} (a_{k+1} (k+1+r)(3k+5+3r) + 3a_k) x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$r(2+3r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \left\{ 0, -\frac{2}{3} \right\}$$

- Each term in the series must be 0, giving the recursion relation

$$3(k+1+r) \left(k + \frac{5}{3} + r \right) a_{k+1} + 3a_k = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+1} = -\frac{3a_k}{(k+1+r)(3k+5+3r)}$$

- Recursion relation for $r = 0$

$$a_{k+1} = -\frac{3a_k}{(k+1)(3k+5)}$$

- Solution for $r = 0$

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+1} = -\frac{3a_k}{(k+1)(3k+5)} \right]$$

- Recursion relation for $r = -\frac{2}{3}$

$$a_{k+1} = -\frac{3a_k}{(k+\frac{1}{3})(3k+3)}$$

- Solution for $r = -\frac{2}{3}$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k-\frac{2}{3}}, a_{k+1} = -\frac{3a_k}{(k+\frac{1}{3})(3k+3)} \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^k \right) + \left(\sum_{k=0}^{\infty} b_k x^{k-\frac{2}{3}} \right), a_{k+1} = -\frac{3a_k}{(k+1)(3k+5)}, b_{k+1} = -\frac{3b_k}{(k+\frac{1}{3})(3k+3)} \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
    <- Bessel successful
<- special function solution successful`

```

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 52

Order:=8;

dsolve(3*x^2*diff(y(x),x\$2)+5*x*diff(y(x),x)+3*x*y(x)=0,y(x),type='series',x=0);

$$y(x) = \frac{c_1 \left(1 - 3x + \frac{9}{8}x^2 - \frac{9}{56}x^3 + \frac{27}{2240}x^4 - \frac{81}{145600}x^5 + \frac{81}{4659200}x^6 - \frac{243}{619673600}x^7 + O(x^8) \right)}{x^{\frac{2}{3}}} + c_2 \left(1 - \frac{3}{5}x + \frac{9}{80}x^2 - \frac{9}{880}x^3 + \frac{27}{49280}x^4 - \frac{81}{4188800}x^5 + \frac{81}{167552000}x^6 - \frac{243}{26975872000}x^7 + O(x^8) \right)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 111

AsymptoticDSolveValue[3*x^2*y''[x]+5*x*y'[x]+3*x*y[x]==0,y[x],{x,0,7}]

$$y(x) \rightarrow c_1 \left(-\frac{243x^7}{26975872000} + \frac{81x^6}{167552000} - \frac{81x^5}{4188800} + \frac{27x^4}{49280} - \frac{9x^3}{880} + \frac{9x^2}{80} - \frac{3x}{5} + 1 \right) + \frac{c_2 \left(-\frac{243x^7}{619673600} + \frac{81x^6}{4659200} - \frac{81x^5}{145600} + \frac{27x^4}{2240} - \frac{9x^3}{56} + \frac{9x^2}{8} - 3x + 1 \right)}{x^{2/3}}$$

18.2 problem 1(b)

18.2.1 Maple step by step solution 1494

Internal problem ID [6051]

Internal file name [OUTPUT/5299_Sunday_June_05_2022_03_32_48_PM_71658375/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 159

Problem number: 1(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Repeated root"**

Maple gives the following as the ode type

`[_Lienard]`

$$x^2y'' + xy' + yx^2 = 0$$

With the expansion point for the power series method at $x = 0$.

The ODE is

$$x^2y'' + xy' + yx^2 = 0$$

Or

$$x(y''x + y' + xy) = 0$$

For $x \neq 0$ the above simplifies to

$$y''x + y' + xy = 0$$

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2y'' + xy' + yx^2 = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{1}{x}$$

$$q(x) = 1$$

Table 225: Table $p(x), q(x)$ singularities.

$p(x) = \frac{1}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = 1$	
singularity	type

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2 y'' + x y' + y x^2 = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) + x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) x^2 = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) \right) + \left(\sum_{n=0}^{\infty} x^{2+n+r} a_n \right) = 0 \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} x^{2+n+r} a_n = \sum_{n=2}^{\infty} a_{n-2} x^{n+r}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) \right) + \left(\sum_{n=2}^{\infty} a_{n-2} x^{n+r} \right) = 0 \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r} a_n (n+r) (n+r-1) + x^{n+r} a_n (n+r) = 0$$

When $n=0$ the above becomes

$$x^r a_0 r(-1+r) + x^r a_0 r = 0$$

Or

$$(x^r r(-1+r) + x^r r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$x^r r^2 = 0$$

Since the above is true for all x then the indicial equation becomes

$$r^2 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = 0$$

$$r_2 = 0$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$x^r r^2 = 0$$

Solving for r gives the roots of the indicial equation as Since the root of the indicial equation is repeated, then we can construct two linearly independent solutions. The first solution has the form

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+r} \quad (1A)$$

Now the second solution y_2 is found using

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right) \quad (1B)$$

Then the general solution will be

$$y = c_1 y_1(x) + c_2 y_2(x)$$

In Eq (1B) the sum starts from 1 and not zero. In Eq (1A), a_0 is never zero, and is arbitrary and is typically taken as $a_0 = 1$, and $\{c_1, c_2\}$ are two arbitrary constants of integration which can be found from initial conditions. We start by finding the first solution $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

For $2 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) + a_n(n+r) + a_{n-2} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{a_{n-2}}{n^2 + 2nr + r^2} \quad (4)$$

Which for the root $r = 0$ becomes

$$a_n = -\frac{a_{n-2}}{n^2} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = -\frac{1}{(r+2)^2}$$

Which for the root $r = 0$ becomes

$$a_2 = -\frac{1}{4}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{(r+2)^2}$	$-\frac{1}{4}$

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{(r+2)^2}$	$-\frac{1}{4}$
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{(r+2)^2(4+r)^2}$$

Which for the root $r = 0$ becomes

$$a_4 = \frac{1}{64}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{(r+2)^2}$	$-\frac{1}{4}$
a_3	0	0
a_4	$\frac{1}{(r+2)^2(4+r)^2}$	$\frac{1}{64}$

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{(r+2)^2}$	$-\frac{1}{4}$
a_3	0	0
a_4	$\frac{1}{(r+2)^2(4+r)^2}$	$\frac{1}{64}$
a_5	0	0

For $n = 6$, using the above recursive equation gives

$$a_6 = -\frac{1}{(r+2)^2(4+r)^2(r+6)^2}$$

Which for the root $r = 0$ becomes

$$a_6 = -\frac{1}{2304}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{(r+2)^2}$	$-\frac{1}{4}$
a_3	0	0
a_4	$\frac{1}{(r+2)^2(4+r)^2}$	$\frac{1}{64}$
a_5	0	0
a_6	$-\frac{1}{(r+2)^2(4+r)^2(r+6)^2}$	$-\frac{1}{2304}$

For $n = 7$, using the above recursive equation gives

$$a_7 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{(r+2)^2}$	$-\frac{1}{4}$
a_3	0	0
a_4	$\frac{1}{(r+2)^2(4+r)^2}$	$\frac{1}{64}$
a_5	0	0
a_6	$-\frac{1}{(r+2)^2(4+r)^2(r+6)^2}$	$-\frac{1}{2304}$
a_7	0	0

Using the above table, then the first solution $y_1(x)$ becomes

$$\begin{aligned} y_1(x) &= a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots \\ &= 1 - \frac{x^2}{4} + \frac{x^4}{64} - \frac{x^6}{2304} + O(x^8) \end{aligned}$$

Now the second solution is found. The second solution is given by

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right)$$

Where b_n is found using

$$b_n = \frac{d}{dr} a_{n,r}$$

And the above is then evaluated at $r = 0$. The above table for $a_{n,r}$ is used for this purpose. Computing the derivatives gives the following table

n	$b_{n,r}$	a_n	$b_{n,r} = \frac{d}{dr} a_{n,r}$	$b_n(r = 0)$
b_0	1	1	N/A since b_n starts from 1	N/A
b_1	0	0	0	0
b_2	$-\frac{1}{(r+2)^2}$	$-\frac{1}{4}$	$\frac{2}{(r+2)^3}$	$\frac{1}{4}$
b_3	0	0	0	0
b_4	$\frac{1}{(r+2)^2(4+r)^2}$	$\frac{1}{64}$	$\frac{-12-4r}{(r+2)^3(4+r)^3}$	$-\frac{3}{128}$
b_5	0	0	0	0
b_6	$-\frac{1}{(r+2)^2(4+r)^2(r+6)^2}$	$-\frac{1}{2304}$	$\frac{6r^2+48r+88}{(r+2)^3(4+r)^3(r+6)^3}$	$\frac{11}{13824}$
b_7	0	0	0	0

The above table gives all values of b_n needed. Hence the second solution is

$$\begin{aligned} y_2(x) &= y_1(x) \ln(x) + b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 + b_7x^7 + b_8x^8 \dots \\ &= \left(1 - \frac{x^2}{4} + \frac{x^4}{64} - \frac{x^6}{2304} + O(x^8)\right) \ln(x) + \frac{x^2}{4} - \frac{3x^4}{128} + \frac{11x^6}{13824} + O(x^8) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= c_1 \left(1 - \frac{x^2}{4} + \frac{x^4}{64} - \frac{x^6}{2304} + O(x^8)\right) \\ &\quad + c_2 \left(\left(1 - \frac{x^2}{4} + \frac{x^4}{64} - \frac{x^6}{2304} + O(x^8)\right) \ln(x) + \frac{x^2}{4} - \frac{3x^4}{128} + \frac{11x^6}{13824} + O(x^8)\right) \end{aligned}$$

Hence the final solution is

$$\begin{aligned} y &= y_h \\ &= c_1 \left(1 - \frac{x^2}{4} + \frac{x^4}{64} - \frac{x^6}{2304} + O(x^8)\right) \\ &\quad + c_2 \left(\left(1 - \frac{x^2}{4} + \frac{x^4}{64} - \frac{x^6}{2304} + O(x^8)\right) \ln(x) + \frac{x^2}{4} - \frac{3x^4}{128} + \frac{11x^6}{13824} + O(x^8)\right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \left(1 - \frac{x^2}{4} + \frac{x^4}{64} - \frac{x^6}{2304} + O(x^8) \right) + c_2 \left(\left(1 - \frac{x^2}{4} + \frac{x^4}{64} - \frac{x^6}{2304} + O(x^8) \right) \ln(x) + \frac{x^2}{4} - \frac{3x^4}{128} + \frac{11x^6}{13824} + O(x^8) \right) \quad (1)$$

Verification of solutions

$$y = c_1 \left(1 - \frac{x^2}{4} + \frac{x^4}{64} - \frac{x^6}{2304} + O(x^8) \right) + c_2 \left(\left(1 - \frac{x^2}{4} + \frac{x^4}{64} - \frac{x^6}{2304} + O(x^8) \right) \ln(x) + \frac{x^2}{4} - \frac{3x^4}{128} + \frac{11x^6}{13824} + O(x^8) \right)$$

Verified OK.

18.2.1 Maple step by step solution

Let's solve

$$x^2 y'' + xy' + yx^2 = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{y'}{x} - y$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{y'}{x} + y = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$[P_2(x) = \frac{1}{x}, P_3(x) = 1]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 1$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 0$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$y''x + y' + xy = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x \cdot y$ to series expansion

$$x \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+1}$$

- Shift index using $k- > k - 1$

$$x \cdot y = \sum_{k=1}^{\infty} a_{k-1} x^{k+r}$$

- Convert y' to series expansion

$$y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1}$$

- Shift index using $k- > k + 1$

$$y' = \sum_{k=-1}^{\infty} a_{k+1} (k+r+1) x^{k+r}$$

- Convert $x \cdot y''$ to series expansion

$$x \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r-1}$$

- Shift index using $k- > k + 1$

$$x \cdot y'' = \sum_{k=-1}^{\infty} a_{k+1} (k+r+1)(k+r) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0 r^2 x^{-1+r} + a_1 (1+r)^2 x^r + \left(\sum_{k=1}^{\infty} (a_{k+1} (k+r+1)^2 + a_{k-1}) x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$r^2 = 0$$

- Values of r that satisfy the indicial equation

$$r = 0$$

- Each term must be 0

$$a_1(1+r)^2 = 0$$
- Each term in the series must be 0, giving the recursion relation

$$a_{k+1}(k+1)^2 + a_{k-1} = 0$$
- Shift index using $k \rightarrow k+1$

$$a_{k+2}(k+2)^2 + a_k = 0$$
- Recursion relation that defines series solution to ODE

$$a_{k+2} = -\frac{a_k}{(k+2)^2}$$
- Recursion relation for $r = 0$

$$a_{k+2} = -\frac{a_k}{(k+2)^2}$$
- Solution for $r = 0$

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+2} = -\frac{a_k}{(k+2)^2}, a_1 = 0 \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
    <- Bessel successful
<- special function solution successful`

```

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 47

Order:=8;

```
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)+x^2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (c_2 \ln(x) + c_1) \left(1 - \frac{1}{4}x^2 + \frac{1}{64}x^4 - \frac{1}{2304}x^6 + O(x^8) \right) \\ + \left(\frac{1}{4}x^2 - \frac{3}{128}x^4 + \frac{11}{13824}x^6 + O(x^8) \right) c_2$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 81

```
AsymptoticDSolveValue[x^2*y'[x]+x*y'[x]+x^2*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(-\frac{x^6}{2304} + \frac{x^4}{64} - \frac{x^2}{4} + 1 \right) \\ + c_2 \left(\frac{11x^6}{13824} - \frac{3x^4}{128} + \frac{x^2}{4} + \left(-\frac{x^6}{2304} + \frac{x^4}{64} - \frac{x^2}{4} + 1 \right) \log(x) \right)$$

18.3 problem 2

Internal problem ID [6052]

Internal file name [OUTPUT/5300_Sunday_June_05_2022_03_32_50_PM_52482633/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 159

Problem number: 2.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Complex roots"**

Maple gives the following as the ode type

```
[[_Emden , _Fowler]]
```

$$x^2y'' + y'e^xx + y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2y'' + y'e^xx + y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{e^x}{x}$$
$$q(x) = \frac{1}{x^2}$$

Table 227: Table $p(x), q(x)$ singularities.

$p(x) = \frac{e^x}{x}$	
singularity	type
$x = 0$	“regular”
$x = \infty$	“regular”

$q(x) = \frac{1}{x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0, \infty]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2 y'' + y' e^x x + y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) + \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) e^x x + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Expanding $x e^x$ as Taylor series around $x = 0$ and keeping only the first 8 terms gives

$$\begin{aligned} x e^x &= x + x^2 + \frac{1}{2}x^3 + \frac{1}{6}x^4 + \frac{1}{24}x^5 + \frac{1}{120}x^6 + \frac{1}{720}x^7 + \frac{1}{5040}x^8 + \dots \\ &= x + x^2 + \frac{1}{2}x^3 + \frac{1}{6}x^4 + \frac{1}{24}x^5 + \frac{1}{120}x^6 + \frac{1}{720}x^7 + \frac{1}{5040}x^8 \end{aligned}$$

Which simplifies to

$$\begin{aligned}
& \left(\sum_{n=0}^{\infty} x^{n+r} a_n(n+r)(n+r-1) \right) + \left(\sum_{n=0}^{\infty} \frac{x^{n+r+7} a_n(n+r)}{5040} \right) \\
& + \left(\sum_{n=0}^{\infty} \frac{x^{n+r+6} a_n(n+r)}{720} \right) + \left(\sum_{n=0}^{\infty} \frac{x^{n+r+5} a_n(n+r)}{120} \right) \\
& + \left(\sum_{n=0}^{\infty} \frac{x^{n+r+4} a_n(n+r)}{24} \right) + \left(\sum_{n=0}^{\infty} \frac{x^{n+r+3} a_n(n+r)}{6} \right) \\
& + \left(\sum_{n=0}^{\infty} \frac{x^{n+r+2} a_n(n+r)}{2} \right) + \left(\sum_{n=0}^{\infty} x^{1+n+r} a_n(n+r) \right) \\
& + \left(\sum_{n=0}^{\infty} x^{n+r} a_n(n+r) \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0
\end{aligned} \tag{2A}$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\begin{aligned}
\sum_{n=0}^{\infty} \frac{x^{n+r+7} a_n(n+r)}{5040} &= \sum_{n=7}^{\infty} \frac{a_{n-7}(n-7+r) x^{n+r}}{5040} \\
\sum_{n=0}^{\infty} \frac{x^{n+r+6} a_n(n+r)}{720} &= \sum_{n=6}^{\infty} \frac{a_{n-6}(n-6+r) x^{n+r}}{720} \\
\sum_{n=0}^{\infty} \frac{x^{n+r+5} a_n(n+r)}{120} &= \sum_{n=5}^{\infty} \frac{a_{n-5}(n-5+r) x^{n+r}}{120} \\
\sum_{n=0}^{\infty} \frac{x^{n+r+4} a_n(n+r)}{24} &= \sum_{n=4}^{\infty} \frac{a_{n-4}(n-4+r) x^{n+r}}{24} \\
\sum_{n=0}^{\infty} \frac{x^{n+r+3} a_n(n+r)}{6} &= \sum_{n=3}^{\infty} \frac{a_{n-3}(n-3+r) x^{n+r}}{6} \\
\sum_{n=0}^{\infty} \frac{x^{n+r+2} a_n(n+r)}{2} &= \sum_{n=2}^{\infty} \frac{a_{n-2}(n+r-2) x^{n+r}}{2} \\
\sum_{n=0}^{\infty} x^{1+n+r} a_n(n+r) &= \sum_{n=1}^{\infty} a_{n-1}(n+r-1) x^{n+r}
\end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n + r$.

$$\begin{aligned}
& \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=7}^{\infty} \frac{a_{n-7} (n-7+r) x^{n+r}}{5040} \right) \\
& + \left(\sum_{n=6}^{\infty} \frac{a_{n-6} (n-6+r) x^{n+r}}{720} \right) + \left(\sum_{n=5}^{\infty} \frac{a_{n-5} (n-5+r) x^{n+r}}{120} \right) \\
& + \left(\sum_{n=4}^{\infty} \frac{a_{n-4} (n-4+r) x^{n+r}}{24} \right) + \left(\sum_{n=3}^{\infty} \frac{a_{n-3} (n-3+r) x^{n+r}}{6} \right) \\
& + \left(\sum_{n=2}^{\infty} \frac{a_{n-2} (n+r-2) x^{n+r}}{2} \right) + \left(\sum_{n=1}^{\infty} a_{n-1} (n+r-1) x^{n+r} \right) \\
& + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0
\end{aligned} \tag{2B}$$

The indicial equation is obtained from $n = 0$. From Eq (2B) this gives

$$x^{n+r} a_n (n+r) (n+r-1) + x^{n+r} a_n (n+r) + a_n x^{n+r} = 0$$

When $n = 0$ the above becomes

$$x^r a_0 r (-1+r) + x^r a_0 r + a_0 x^r = 0$$

Or

$$(x^r r (-1+r) + x^r r + x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(r^2 + 1) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$r^2 + 1 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = i$$

$$r_2 = -i$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(r^2 + 1) x^r = 0$$

Solving for r gives the roots of the indicial equation as Since the roots are complex conjugates, then two linearly independent solutions can be constructed using

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+i}$$

$$y_2(x) = \sum_{n=0}^{\infty} b_n x^{n-i}$$

$y_1(x)$ is found first. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = -\frac{r}{r^2 + 2r + 2}$$

Substituting $n = 2$ in Eq. (2B) gives

$$a_2 = -\frac{r^3}{2(r^2 + 2r + 2)(r^2 + 4r + 5)}$$

Substituting $n = 3$ in Eq. (2B) gives

$$a_3 = -\frac{r(r^4 - 6r^2 - 9r - 5)}{6(r^2 + 2r + 2)(r^2 + 4r + 5)(r^2 + 6r + 10)}$$

Substituting $n = 4$ in Eq. (2B) gives

$$a_4 = -\frac{r(r^6 - 2r^5 - 43r^4 - 136r^3 - 180r^2 - 112r - 40)}{24(r^2 + 2r + 2)(r^2 + 4r + 5)(r^2 + 6r + 10)(r^2 + 8r + 17)}$$

Substituting $n = 5$ in Eq. (2B) gives

$$a_5 = -\frac{r(r^8 - 10r^7 - 201r^6 - 1035r^5 - 2331r^4 - 2105r^3 + 321r^2 + 1760r + 800)}{120(r^2 + 2r + 2)(r^2 + 4r + 5)(r^2 + 6r + 10)(r^2 + 8r + 17)(r^2 + 10r + 26)}$$

Substituting $n = 6$ in Eq. (2B) gives

$$a_6 = -\frac{r(2+r)(r^9 - 34r^8 - 666r^7 - 3942r^6 - 7855r^5 + 11152r^4 + 77700r^3 + 138084r^2 + 111970r + 413)}{720(r^2 + 2r + 2)(r^2 + 4r + 5)(r^2 + 6r + 10)(r^2 + 8r + 17)(r^2 + 10r + 26)(r^2 + 12r + 37)}$$

For $7 \leq n$ the recursive equation is

$$\begin{aligned}
 & a_n(n+r)(n+r-1) + \frac{a_{n-7}(n-7+r)}{5040} + \frac{a_{n-6}(n-6+r)}{720} \\
 & + \frac{a_{n-5}(n-5+r)}{120} + \frac{a_{n-4}(n-4+r)}{24} + \frac{a_{n-3}(n-3+r)}{6} \\
 & + \frac{a_{n-2}(n+r-2)}{2} + a_{n-1}(n+r-1) + a_n(n+r) + a_n = 0
 \end{aligned} \tag{3}$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{na_{n-7} + 7na_{n-6} + 42na_{n-5} + 210na_{n-4} + 840na_{n-3} + 2520na_{n-2} + 5040na_{n-1} + ra_{n-7} + 7ra_{n-6}}{\dots} \tag{4}$$

Which for the root $r = i$ becomes

$$a_n = \frac{(-a_{n-7} - 7a_{n-6} - 42a_{n-5} - 210a_{n-4} - 840a_{n-3} - 2520a_{n-2} - 5040a_{n-1})n + (7-i)a_{n-7} + (42-7)}{\dots} \tag{5}$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = i$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{r}{r^2+2r+2}$	$-\frac{2}{5} - \frac{i}{5}$
a_2	$-\frac{r^3}{2(r^2+2r+2)(r^2+4r+5)}$	$\frac{3}{80} - \frac{i}{80}$
a_3	$-\frac{r(r^4-6r^2-9r-5)}{6(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)}$	$\frac{67}{9360} + \frac{9i}{1040}$
a_4	$-\frac{r(r^6-2r^5-43r^4-136r^3-180r^2-112r-40)}{24(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)}$	$-\frac{103}{149760} + \frac{229i}{149760}$
a_5	$-\frac{r(r^8-10r^7-201r^6-1035r^5-2331r^4-2105r^3+321r^2+1760r+800)}{120(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)(r^2+10r+26)}$	$-\frac{2831}{7238400} - \frac{607i}{4343040}$
a_6	$-\frac{r(2+r)(r^9-34r^8-666r^7-3942r^6-7855r^5+11152r^4+77700r^3+138084r^2+111970r+41300)}{720(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)(r^2+10r+26)(r^2+12r+37)}$	$-\frac{59077}{1563494400} - \frac{26063i}{260582400}$

For $n = 7$, using the above recursive equation gives

$$a_7 = -\frac{r(r^{12} - 84r^{11} - 2248r^{10} - 19677r^9 - 49342r^8 + 352058r^7 + 3397664r^6 + 13171067r^5 + 29036801r^4 + 5040(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)(r^2+10r+26))}{\dots}$$

Which for the root $r = i$ becomes

$$a_7 = \frac{22952047}{2030197478400} - \frac{8634893i}{580056422400}$$

And the table now becomes

n	$a_{n,r}$
a_0	1
a_1	$-\frac{r}{r^2+2r+2}$
a_2	$-\frac{r^3}{2(r^2+2r+2)(r^2+4r+5)}$
a_3	$-\frac{r(r^4-6r^2-9r-5)}{6(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)}$
a_4	$-\frac{r(r^6-2r^5-43r^4-136r^3-180r^2-112r-40)}{24(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)}$
a_5	$-\frac{r(r^8-10r^7-201r^6-1035r^5-2331r^4-2105r^3+321r^2+1760r+800)}{120(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)(r^2+10r+26)}$
a_6	$-\frac{r(2+r)(r^9-34r^8-666r^7-3942r^6-7855r^5+11152r^4+77700r^3+138084r^2+111970r+41300)}{720(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)(r^2+10r+26)(r^2+12r+37)}$
a_7	$-\frac{r(r^{12}-84r^{11}-2248r^{10}-19677r^9-49342r^8+352058r^7+3397664r^6+13171067r^5+29036801r^4+38565016r^3+30552414r^2+13873510r+1000000)}{5040(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)(r^2+10r+26)(r^2+12r+37)(r^2+14r+50)}$

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned}
 y_1(x) &= x^i(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots) \\
 &= x^i \left(1 + \left(-\frac{2}{5} - \frac{i}{5} \right) x + \left(\frac{3}{80} - \frac{i}{80} \right) x^2 + \left(\frac{67}{9360} + \frac{9i}{1040} \right) x^3 \right. \\
 &\quad \left. + \left(-\frac{103}{149760} + \frac{229i}{149760} \right) x^4 + \left(-\frac{2831}{7238400} - \frac{607i}{4343040} \right) x^5 \right. \\
 &\quad \left. + \left(-\frac{59077}{1563494400} - \frac{26063i}{260582400} \right) x^6 + \left(\frac{22952047}{2030197478400} - \frac{8634893i}{580056422400} \right) x^7 \right. \\
 &\quad \left. + O(x^8) \right)
 \end{aligned}$$

The second solution $y_2(x)$ is found by taking the complex conjugate of $y_1(x)$ which gives

$$\begin{aligned}
 y_2(x) &= x^{-i} \left(1 + \left(-\frac{2}{5} + \frac{i}{5} \right) x + \left(\frac{3}{80} + \frac{i}{80} \right) x^2 + \left(\frac{67}{9360} - \frac{9i}{1040} \right) x^3 \right. \\
 &\quad \left. + \left(-\frac{103}{149760} - \frac{229i}{149760} \right) x^4 + \left(-\frac{2831}{7238400} + \frac{607i}{4343040} \right) x^5 \right. \\
 &\quad \left. + \left(-\frac{59077}{1563494400} + \frac{26063i}{260582400} \right) x^6 + \left(\frac{22952047}{2030197478400} + \frac{8634893i}{580056422400} \right) x^7 \right. \\
 &\quad \left. + O(x^8) \right)
 \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 y_h(x) &= c_1 y_1(x) + c_2 y_2(x) \\
 &= c_1 x^i \left(1 + \left(-\frac{2}{5} - \frac{i}{5} \right) x + \left(\frac{3}{80} - \frac{i}{80} \right) x^2 + \left(\frac{67}{9360} + \frac{9i}{1040} \right) x^3 \right. \\
 &\quad \left. + \left(-\frac{103}{149760} + \frac{229i}{149760} \right) x^4 + \left(-\frac{2831}{7238400} - \frac{607i}{4343040} \right) x^5 \right. \\
 &\quad \left. + \left(-\frac{59077}{1563494400} - \frac{26063i}{260582400} \right) x^6 + \left(\frac{22952047}{2030197478400} - \frac{8634893i}{580056422400} \right) x^7 \right. \\
 &\quad \left. + O(x^8) \right) + c_2 x^{-i} \left(1 + \left(-\frac{2}{5} + \frac{i}{5} \right) x + \left(\frac{3}{80} + \frac{i}{80} \right) x^2 + \left(\frac{67}{9360} - \frac{9i}{1040} \right) x^3 \right. \\
 &\quad \left. + \left(-\frac{103}{149760} - \frac{229i}{149760} \right) x^4 + \left(-\frac{2831}{7238400} + \frac{607i}{4343040} \right) x^5 \right. \\
 &\quad \left. + \left(-\frac{59077}{1563494400} + \frac{26063i}{260582400} \right) x^6 + \left(\frac{22952047}{2030197478400} + \frac{8634893i}{580056422400} \right) x^7 \right. \\
 &\quad \left. + O(x^8) \right)
 \end{aligned}$$

Hence the final solution is

$$\begin{aligned}
 y &= y_h \\
 &= c_1 x^i \left(1 + \left(-\frac{2}{5} - \frac{i}{5} \right) x + \left(\frac{3}{80} - \frac{i}{80} \right) x^2 + \left(\frac{67}{9360} + \frac{9i}{1040} \right) x^3 \right. \\
 &\quad \left. + \left(-\frac{103}{149760} + \frac{229i}{149760} \right) x^4 + \left(-\frac{2831}{7238400} - \frac{607i}{4343040} \right) x^5 \right. \\
 &\quad \left. + \left(-\frac{59077}{1563494400} - \frac{26063i}{260582400} \right) x^6 + \left(\frac{22952047}{2030197478400} - \frac{8634893i}{580056422400} \right) x^7 \right. \\
 &\quad \left. + O(x^8) \right) + c_2 x^{-i} \left(1 + \left(-\frac{2}{5} + \frac{i}{5} \right) x + \left(\frac{3}{80} + \frac{i}{80} \right) x^2 + \left(\frac{67}{9360} - \frac{9i}{1040} \right) x^3 \right. \\
 &\quad \left. + \left(-\frac{103}{149760} - \frac{229i}{149760} \right) x^4 + \left(-\frac{2831}{7238400} + \frac{607i}{4343040} \right) x^5 \right. \\
 &\quad \left. + \left(-\frac{59077}{1563494400} + \frac{26063i}{260582400} \right) x^6 + \left(\frac{22952047}{2030197478400} + \frac{8634893i}{580056422400} \right) x^7 \right. \\
 &\quad \left. + O(x^8) \right)
 \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned} y = c_1 x^i & \left(1 + \left(-\frac{2}{5} - \frac{i}{5} \right) x + \left(\frac{3}{80} - \frac{i}{80} \right) x^2 + \left(\frac{67}{9360} + \frac{9i}{1040} \right) x^3 \right. \\ & + \left(-\frac{103}{149760} + \frac{229i}{149760} \right) x^4 + \left(-\frac{2831}{7238400} - \frac{607i}{4343040} \right) x^5 \\ & + \left(-\frac{59077}{1563494400} - \frac{26063i}{260582400} \right) x^6 + \left(\frac{22952047}{2030197478400} - \frac{8634893i}{580056422400} \right) x^7 \\ & \left. + O(x^8) \right) + c_2 x^{-i} \left(1 + \left(-\frac{2}{5} + \frac{i}{5} \right) x + \left(\frac{3}{80} + \frac{i}{80} \right) x^2 + \left(\frac{67}{9360} - \frac{9i}{1040} \right) x^3 \right. \\ & + \left(-\frac{103}{149760} - \frac{229i}{149760} \right) x^4 + \left(-\frac{2831}{7238400} + \frac{607i}{4343040} \right) x^5 \\ & + \left(-\frac{59077}{1563494400} + \frac{26063i}{260582400} \right) x^6 + \left(\frac{22952047}{2030197478400} + \frac{8634893i}{580056422400} \right) x^7 \\ & \left. + O(x^8) \right) \end{aligned}$$

Verification of solutions

$$\begin{aligned} y = c_1 x^i & \left(1 + \left(-\frac{2}{5} - \frac{i}{5} \right) x + \left(\frac{3}{80} - \frac{i}{80} \right) x^2 + \left(\frac{67}{9360} + \frac{9i}{1040} \right) x^3 \right. \\ & + \left(-\frac{103}{149760} + \frac{229i}{149760} \right) x^4 + \left(-\frac{2831}{7238400} - \frac{607i}{4343040} \right) x^5 \\ & + \left(-\frac{59077}{1563494400} - \frac{26063i}{260582400} \right) x^6 + \left(\frac{22952047}{2030197478400} - \frac{8634893i}{580056422400} \right) x^7 \\ & \left. + O(x^8) \right) + c_2 x^{-i} \left(1 + \left(-\frac{2}{5} + \frac{i}{5} \right) x + \left(\frac{3}{80} + \frac{i}{80} \right) x^2 + \left(\frac{67}{9360} - \frac{9i}{1040} \right) x^3 \right. \\ & + \left(-\frac{103}{149760} - \frac{229i}{149760} \right) x^4 + \left(-\frac{2831}{7238400} + \frac{607i}{4343040} \right) x^5 \\ & + \left(-\frac{59077}{1563494400} + \frac{26063i}{260582400} \right) x^6 + \left(\frac{22952047}{2030197478400} + \frac{8634893i}{580056422400} \right) x^7 \\ & \left. + O(x^8) \right) \end{aligned}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form  $r_0(x) * Y + r_1(x) * Y$  where  $Y = \exp(\int(r(x), dx)) * 2F1([a$ 
-> Trying changes of variables to rationalize or make the ODE simpler
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
    -> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
    -> trying a solution of the form  $r_0(x) * Y + r_1(x) * Y$  where  $Y = \exp(\int(r(x), dx)) * 2F1$ 
        trying a symmetry of the form [xi=0, eta=F(x)]
        trying 2nd order exact linear
        trying symmetries linear in x and y(x)
        trying to convert to a linear ODE with constant coefficients
<- unable to find a useful change of variables
    trying a symmetry of the form [xi=0, eta=F(x)]
    trying differential order: 2; exact nonlinear
    trying symmetries linear in x and y(x)
    trying to convert to a linear ODE with constant coefficients
    trying 2nd order, integrating factor of the form  $\mu(x,y)$ 
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
    -> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
    -> trying a solution of the form  $r_0(x) * Y + r_1(x) * Y$  where  $Y = \exp(\int(r(x), dx)) * 2F1$ 
    -> Trying changes of variables to rationalize or make the ODE simpler
        trying a symmetry of the form [xi=0, eta=F(x)]
        checking if the LODE is missing y
        -> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
        -> trying a solution of the form  $r_0(x) * Y + r_1(x) * Y$  where  $Y = \exp(\int(r(x), dx)) * 2F1$ 
            trying a symmetry of the form [xi=0, eta=F(x)]
            trying 2nd order exact linear
            trying symmetries linear in x and y(x)
            trying to convert to a linear ODE with constant coefficients
<- unable to find a useful change of variables
    trying a symmetry of the form [xi=0, eta=F(x)]
trying to convert to an ODE of Bessel type
-> trying reduction of order to Riccati
    trying Riccati sub-methods:
        trying Riccati_symmetries
        -> trying a symmetry pattern of the form  $[F(x)*G(y), 0]$ 
        -> trying a symmetry pattern of the form  $[0, F(x)*G(y)]$ 
        -> trying a symmetry pattern of the form  $[F(x), G(x)*y+H(x)]$ 
--- Trying Lie symmetry methods, 2nd order ---
-> Computing symmetries using:  $\text{var} = 2; [0, -]$ 
```

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 85

Order:=8;

dsolve(x^2*diff(y(x),x\$2)+x*exp(x)*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);

$$\begin{aligned}
 y(x) = & c_1 x^{-i} \left(1 + \left(-\frac{2}{5} + \frac{i}{5} \right) x + \left(\frac{3}{80} + \frac{i}{80} \right) x^2 + \left(\frac{67}{9360} - \frac{9i}{1040} \right) x^3 \right. \\
 & + \left(-\frac{103}{149760} - \frac{229i}{149760} \right) x^4 + \left(-\frac{2831}{7238400} + \frac{607i}{4343040} \right) x^5 \\
 & + \left(-\frac{59077}{1563494400} + \frac{26063i}{260582400} \right) x^6 + \left(\frac{22952047}{2030197478400} + \frac{8634893i}{580056422400} \right) x^7 \\
 & + O(x^8) \Big) + c_2 x^i \left(1 + \left(-\frac{2}{5} - \frac{i}{5} \right) x + \left(\frac{3}{80} - \frac{i}{80} \right) x^2 + \left(\frac{67}{9360} + \frac{9i}{1040} \right) x^3 \right. \\
 & + \left(-\frac{103}{149760} + \frac{229i}{149760} \right) x^4 + \left(-\frac{2831}{7238400} - \frac{607i}{4343040} \right) x^5 \\
 & + \left(-\frac{59077}{1563494400} - \frac{26063i}{260582400} \right) x^6 + \left(\frac{22952047}{2030197478400} - \frac{8634893i}{580056422400} \right) x^7 \\
 & \left. + O(x^8) \right)
 \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.028 (sec). Leaf size: 122

AsymptoticDSolveValue[x^2*y''[x]+x*Exp[x]*y'[x]+y[x]==0,y[x],{x,0,7}]

$$\begin{aligned}
 y(x) \rightarrow & \left(\frac{11}{1563494400} + \frac{i}{97718400} \right) c_2 x^{-i} \left((4913 + 7070i)x^6 - (8568 - 32328i)x^5 \right. \\
 & - (132840 + 24120i)x^4 - (247680 + 869760i)x^3 + (2540160 - 1918080i)x^2 \\
 & \left. - (4976640 - 35665920i)x + (45619200 - 66355200i) \right) \\
 & - \left(\frac{1}{97718400} + \frac{11i}{1563494400} \right) c_1 x^i \left((7070 + 4913i)x^6 + (32328 - 8568i)x^5 \right. \\
 & - (24120 + 132840i)x^4 - (869760 + 247680i)x^3 - (1918080 - 2540160i)x^2 \\
 & \left. + (35665920 - 4976640i)x - (66355200 - 45619200i) \right)
 \end{aligned}$$

19 Chapter 4. Linear equations with Regular Singular Points. Page 166

19.1	problem 1(i)	1510
19.2	problem 1(ii)	1527
19.3	problem 1(iii)	1548
19.4	problem 3(a)	1561
19.5	problem 3(b)	1574
19.6	problem 3(c)	1591
19.7	problem 3(d)	1606
19.8	problem 3(e)	1623
19.9	problem 3(f)	1640

19.1 problem 1(i)

19.1.1 Maple step by step solution 1522

Internal problem ID [6053]

Internal file name [OUTPUT/5301_Sunday_June_05_2022_03_33_12_PM_94404250/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 166

Problem number: 1(i).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$2x^2y'' + (x^2 + 5x)y' + (x^2 - 2)y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$2x^2y'' + (x^2 + 5x)y' + (x^2 - 2)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{5 + x}{2x}$$
$$q(x) = \frac{x^2 - 2}{2x^2}$$

Table 228: Table $p(x), q(x)$ singularities.

$p(x) = \frac{5+x}{2x}$	
singularity	type
$x = 0$	“regular”

$q(x) = \frac{x^2-2}{2x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$2x^2y'' + (x^2 + 5x)y' + (x^2 - 2)y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$2x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) + (x^2 + 5x) \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (x^2 - 2) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{1+n+r} a_n (n+r) \right) \\ & + \left(\sum_{n=0}^{\infty} 5x^{n+r} a_n (n+r) \right) + \left(\sum_{n=0}^{\infty} x^{n+r+2} a_n \right) + \sum_{n=0}^{\infty} (-2a_n x^{n+r}) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=0}^{\infty} x^{1+n+r} a_n (n+r) &= \sum_{n=1}^{\infty} a_{n-1} (n+r-1) x^{n+r} \\ \sum_{n=0}^{\infty} x^{n+r+2} a_n &= \sum_{n=2}^{\infty} a_{n-2} x^{n+r} \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=1}^{\infty} a_{n-1} (n+r-1) x^{n+r} \right) \\ & + \left(\sum_{n=0}^{\infty} 5x^{n+r} a_n (n+r) \right) + \left(\sum_{n=2}^{\infty} a_{n-2} x^{n+r} \right) + \sum_{n=0}^{\infty} (-2a_n x^{n+r}) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$2x^{n+r} a_n (n+r) (n+r-1) + 5x^{n+r} a_n (n+r) - 2a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$2x^r a_0 r (-1+r) + 5x^r a_0 r - 2a_0 x^r = 0$$

Or

$$(2x^r r (-1+r) + 5x^r r - 2x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(2r^2 + 3r - 2) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$2r^2 + 3r - 2 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = \frac{1}{2}$$

$$r_2 = -2$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(2r^2 + 3r - 2) x^r = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = \frac{5}{2}$ is not an integer, then we can construct two linearly independent solutions

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+\frac{1}{2}}$$

$$y_2(x) = \sum_{n=0}^{\infty} b_n x^{n-2}$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = -\frac{r}{2r^2 + 7r + 3}$$

For $2 \leq n$ the recursive equation is

$$2a_n(n+r)(n+r-1) + a_{n-1}(n+r-1) + 5a_n(n+r) + a_{n-2} - 2a_n = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{na_{n-1} + ra_{n-1} + a_{n-2} - a_{n-1}}{2n^2 + 4nr + 2r^2 + 3n + 3r - 2} \quad (4)$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_n = \frac{-2na_{n-1} - 2a_{n-2} + a_{n-1}}{4n^2 + 10n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = \frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{r}{2r^2+7r+3}$	$-\frac{1}{14}$

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{-r^2 - 6r - 3}{(2r^2 + 7r + 3)(2r^2 + 11r + 12)}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_2 = -\frac{25}{504}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{r}{2r^2+7r+3}$	$-\frac{1}{14}$
a_2	$\frac{-r^2-6r-3}{(2r^2+7r+3)(2r^2+11r+12)}$	$-\frac{25}{504}$

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{3r^3 + 19r^2 + 27r + 6}{(2r^2 + 7r + 3)(2r^2 + 11r + 12)(2r^2 + 15r + 25)}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_3 = \frac{197}{33264}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{r}{2r^2+7r+3}$	$-\frac{1}{14}$
a_2	$\frac{-r^2-6r-3}{(2r^2+7r+3)(2r^2+11r+12)}$	$-\frac{25}{504}$
a_3	$\frac{3r^3+19r^2+27r+6}{(2r^2+7r+3)(2r^2+11r+12)(2r^2+15r+25)}$	$\frac{197}{33264}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{-r^4 - r^3 + 37r^2 + 108r + 57}{(2r^2 + 7r + 3)(2r^2 + 11r + 12)(2r^2 + 15r + 25)(2r^2 + 19r + 42)}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_4 = \frac{1921}{3459456}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{r}{2r^2+7r+3}$	$-\frac{1}{14}$
a_2	$\frac{-r^2-6r-3}{(2r^2+7r+3)(2r^2+11r+12)}$	$-\frac{25}{504}$
a_3	$\frac{3r^3+19r^2+27r+6}{(2r^2+7r+3)(2r^2+11r+12)(2r^2+15r+25)}$	$\frac{197}{33264}$
a_4	$\frac{-r^4-r^3+37r^2+108r+57}{(2r^2+7r+3)(2r^2+11r+12)(2r^2+15r+25)(2r^2+19r+42)}$	$\frac{1921}{3459456}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{-5r^5 - 90r^4 - 574r^3 - 1579r^2 - 1737r - 480}{(2r^2 + 7r + 3)(2r^2 + 11r + 12)(2r^2 + 15r + 25)(2r^2 + 19r + 42)(2r^2 + 23r + 63)}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_5 = -\frac{11653}{103783680}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{r}{2r^2+7r+3}$	$-\frac{1}{14}$
a_2	$\frac{-r^2-6r-3}{(2r^2+7r+3)(2r^2+11r+12)}$	$-\frac{25}{504}$
a_3	$\frac{3r^3+19r^2+27r+6}{(2r^2+7r+3)(2r^2+11r+12)(2r^2+15r+25)}$	$\frac{197}{33264}$
a_4	$\frac{-r^4-r^3+37r^2+108r+57}{(2r^2+7r+3)(2r^2+11r+12)(2r^2+15r+25)(2r^2+19r+42)}$	$\frac{1921}{3459456}$
a_5	$\frac{-5r^5-90r^4-574r^3-1579r^2-1737r-480}{(2r^2+7r+3)(2r^2+11r+12)(2r^2+15r+25)(2r^2+19r+42)(2r^2+23r+63)}$	$-\frac{11653}{103783680}$

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{7r^6 + 140r^5 + 1036r^4 + 3445r^3 + 4703r^2 + 1050r - 1191}{(2r^2 + 7r + 3)(2r^2 + 11r + 12)(2r^2 + 15r + 25)(2r^2 + 19r + 42)(2r^2 + 23r + 63)(2r^2 + 27r + 88)}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_6 = \frac{12923}{21171870720}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{r}{2r^2+7r+3}$	$-\frac{1}{14}$
a_2	$\frac{-r^2-6r-3}{(2r^2+7r+3)(2r^2+11r+12)}$	$-\frac{25}{504}$
a_3	$\frac{3r^3+19r^2+27r+6}{(2r^2+7r+3)(2r^2+11r+12)(2r^2+15r+25)}$	$\frac{197}{33264}$
a_4	$\frac{-r^4-r^3+37r^2+108r+57}{(2r^2+7r+3)(2r^2+11r+12)(2r^2+15r+25)(2r^2+19r+42)}$	$\frac{1921}{3459456}$
a_5	$\frac{-5r^5-90r^4-574r^3-1579r^2-1737r-480}{(2r^2+7r+3)(2r^2+11r+12)(2r^2+15r+25)(2r^2+19r+42)(2r^2+23r+63)}$	$-\frac{11653}{103783680}$
a_6	$\frac{7r^6+140r^5+1036r^4+3445r^3+4703r^2+1050r-1191}{(2r^2+7r+3)(2r^2+11r+12)(2r^2+15r+25)(2r^2+19r+42)(2r^2+23r+63)(2r^2+27r+88)}$	$\frac{12923}{21171870720}$

For $n = 7$, using the above recursive equation gives

$$a_7 = \frac{3r^7 + 133r^6 + 2142r^5 + 16915r^4 + 71246r^3 + 157543r^2 + 160707r + 49386}{(2r^2 + 7r + 3)(2r^2 + 11r + 12)(2r^2 + 15r + 25)(2r^2 + 19r + 42)(2r^2 + 23r + 63)(2r^2 + 27r + 88)}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_7 = \frac{917285}{1126343522304}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{r}{2r^2+7r+3}$	$-\frac{1}{14}$
a_2	$\frac{-r^2-6r-3}{(2r^2+7r+3)(2r^2+11r+12)}$	$-\frac{25}{504}$
a_3	$\frac{3r^3+19r^2+27r+6}{(2r^2+7r+3)(2r^2+11r+12)(2r^2+15r+25)}$	$\frac{197}{33264}$
a_4	$\frac{-r^4-r^3+37r^2+108r+57}{(2r^2+7r+3)(2r^2+11r+12)(2r^2+15r+25)(2r^2+19r+42)}$	$\frac{1921}{3459456}$
a_5	$\frac{-5r^5-90r^4-574r^3-1579r^2-1737r-480}{(2r^2+7r+3)(2r^2+11r+12)(2r^2+15r+25)(2r^2+19r+42)(2r^2+23r+63)}$	$-\frac{11653}{103783680}$
a_6	$\frac{7r^6+140r^5+1036r^4+3445r^3+4703r^2+1050r-1191}{(2r^2+7r+3)(2r^2+11r+12)(2r^2+15r+25)(2r^2+19r+42)(2r^2+23r+63)(2r^2+27r+88)}$	$\frac{12923}{21171870720}$
a_7	$\frac{3r^7+133r^6+2142r^5+16915r^4+71246r^3+157543r^2+160707r+49386}{(2r^2+7r+3)(2r^2+11r+12)(2r^2+15r+25)(2r^2+19r+42)(2r^2+23r+63)(2r^2+27r+88)(2r^2+31r+117)}$	$\frac{917285}{1126343522304}$

Using the above table, then the solution $y_1(x)$ is

$$y_1(x) = \sqrt{x}(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots)$$

$$= \sqrt{x} \left(1 - \frac{x}{14} - \frac{25x^2}{504} + \frac{197x^3}{33264} + \frac{1921x^4}{3459456} - \frac{11653x^5}{103783680} + \frac{12923x^6}{21171870720} + \frac{917285x^7}{1126343522304} + O(x^8) \right)$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$b_1 = -\frac{r}{2r^2 + 7r + 3}$$

For $2 \leq n$ the recursive equation is

$$2b_n(n+r)(n+r-1) + b_{n-1}(n+r-1) + 5b_n(n+r) + b_{n-2} - 2b_n = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = -\frac{nb_{n-1} + rb_{n-1} + b_{n-2} - b_{n-1}}{2n^2 + 4nr + 2r^2 + 3n + 3r - 2} \quad (4)$$

Which for the root $r = -2$ becomes

$$b_n = \frac{-nb_{n-1} - b_{n-2} + 3b_{n-1}}{n(2n-5)} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = -2$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{r}{2r^2+7r+3}$	$-\frac{2}{3}$

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{-r^2 - 6r - 3}{(2r^2 + 7r + 3)(2r^2 + 11r + 12)}$$

Which for the root $r = -2$ becomes

$$b_2 = \frac{5}{6}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{r}{2r^2+7r+3}$	$-\frac{2}{3}$
b_2	$\frac{-r^2-6r-3}{(2r^2+7r+3)(2r^2+11r+12)}$	$\frac{5}{6}$

For $n = 3$, using the above recursive equation gives

$$b_3 = \frac{3r^3 + 19r^2 + 27r + 6}{(2r^2 + 7r + 3)(2r^2 + 11r + 12)(2r^2 + 15r + 25)}$$

Which for the root $r = -2$ becomes

$$b_3 = \frac{2}{9}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{r}{2r^2+7r+3}$	$-\frac{2}{3}$
b_2	$\frac{-r^2-6r-3}{(2r^2+7r+3)(2r^2+11r+12)}$	$\frac{5}{6}$
b_3	$\frac{3r^3+19r^2+27r+6}{(2r^2+7r+3)(2r^2+11r+12)(2r^2+15r+25)}$	$\frac{2}{9}$

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{-r^4 - r^3 + 37r^2 + 108r + 57}{(2r^2 + 7r + 3)(2r^2 + 11r + 12)(2r^2 + 15r + 25)(2r^2 + 19r + 42)}$$

Which for the root $r = -2$ becomes

$$b_4 = -\frac{19}{216}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{r}{2r^2+7r+3}$	$-\frac{2}{3}$
b_2	$\frac{-r^2-6r-3}{(2r^2+7r+3)(2r^2+11r+12)}$	$\frac{5}{6}$
b_3	$\frac{3r^3+19r^2+27r+6}{(2r^2+7r+3)(2r^2+11r+12)(2r^2+15r+25)}$	$\frac{2}{9}$
b_4	$\frac{-r^4-r^3+37r^2+108r+57}{(2r^2+7r+3)(2r^2+11r+12)(2r^2+15r+25)(2r^2+19r+42)}$	$-\frac{19}{216}$

For $n = 5$, using the above recursive equation gives

$$b_5 = \frac{-5r^5 - 90r^4 - 574r^3 - 1579r^2 - 1737r - 480}{(2r^2 + 7r + 3)(2r^2 + 11r + 12)(2r^2 + 15r + 25)(2r^2 + 19r + 42)(2r^2 + 23r + 63)}$$

Which for the root $r = -2$ becomes

$$b_5 = -\frac{1}{540}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{r}{2r^2+7r+3}$	$-\frac{2}{3}$
b_2	$\frac{-r^2-6r-3}{(2r^2+7r+3)(2r^2+11r+12)}$	$\frac{5}{6}$
b_3	$\frac{3r^3+19r^2+27r+6}{(2r^2+7r+3)(2r^2+11r+12)(2r^2+15r+25)}$	$\frac{2}{9}$
b_4	$\frac{-r^4-r^3+37r^2+108r+57}{(2r^2+7r+3)(2r^2+11r+12)(2r^2+15r+25)(2r^2+19r+42)}$	$-\frac{19}{216}$
b_5	$\frac{-5r^5-90r^4-574r^3-1579r^2-1737r-480}{(2r^2+7r+3)(2r^2+11r+12)(2r^2+15r+25)(2r^2+19r+42)(2r^2+23r+63)}$	$-\frac{1}{540}$

For $n = 6$, using the above recursive equation gives

$$b_6 = \frac{7r^6 + 140r^5 + 1036r^4 + 3445r^3 + 4703r^2 + 1050r - 1191}{(2r^2 + 7r + 3)(2r^2 + 11r + 12)(2r^2 + 15r + 25)(2r^2 + 19r + 42)(2r^2 + 23r + 63)(2r^2 + 27r + 88)}$$

Which for the root $r = -2$ becomes

$$b_6 = \frac{101}{45360}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{r}{2r^2+7r+3}$	$-\frac{2}{3}$
b_2	$\frac{-r^2-6r-3}{(2r^2+7r+3)(2r^2+11r+12)}$	$\frac{5}{6}$
b_3	$\frac{3r^3+19r^2+27r+6}{(2r^2+7r+3)(2r^2+11r+12)(2r^2+15r+25)}$	$\frac{2}{9}$
b_4	$\frac{-r^4-r^3+37r^2+108r+57}{(2r^2+7r+3)(2r^2+11r+12)(2r^2+15r+25)(2r^2+19r+42)}$	$-\frac{19}{216}$
b_5	$\frac{-5r^5-90r^4-574r^3-1579r^2-1737r-480}{(2r^2+7r+3)(2r^2+11r+12)(2r^2+15r+25)(2r^2+19r+42)(2r^2+23r+63)}$	$-\frac{1}{540}$
b_6	$\frac{7r^6+140r^5+1036r^4+3445r^3+4703r^2+1050r-1191}{(2r^2+7r+3)(2r^2+11r+12)(2r^2+15r+25)(2r^2+19r+42)(2r^2+23r+63)(2r^2+27r+88)}$	$\frac{101}{45360}$

For $n = 7$, using the above recursive equation gives

$$b_7 = \frac{3r^7 + 133r^6 + 2142r^5 + 16915r^4 + 71246r^3 + 157543r^2 + 160707r + 49386}{(2r^2 + 7r + 3)(2r^2 + 11r + 12)(2r^2 + 15r + 25)(2r^2 + 19r + 42)(2r^2 + 23r + 63)(2r^2 + 27r + 88)}$$

Which for the root $r = -2$ becomes

$$b_7 = -\frac{4}{35721}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{r}{2r^2+7r+3}$	$-\frac{2}{3}$
b_2	$\frac{-r^2-6r-3}{(2r^2+7r+3)(2r^2+11r+12)}$	$\frac{5}{6}$
b_3	$\frac{3r^3+19r^2+27r+6}{(2r^2+7r+3)(2r^2+11r+12)(2r^2+15r+25)}$	$\frac{2}{9}$
b_4	$\frac{-r^4-r^3+37r^2+108r+57}{(2r^2+7r+3)(2r^2+11r+12)(2r^2+15r+25)(2r^2+19r+42)}$	$-\frac{19}{216}$
b_5	$\frac{-5r^5-90r^4-574r^3-1579r^2-1737r-480}{(2r^2+7r+3)(2r^2+11r+12)(2r^2+15r+25)(2r^2+19r+42)(2r^2+23r+63)}$	$-\frac{1}{540}$
b_6	$\frac{7r^6+140r^5+1036r^4+3445r^3+4703r^2+1050r-1191}{(2r^2+7r+3)(2r^2+11r+12)(2r^2+15r+25)(2r^2+19r+42)(2r^2+23r+63)(2r^2+27r+88)}$	$\frac{101}{45360}$
b_7	$\frac{3r^7+133r^6+2142r^5+16915r^4+71246r^3+157543r^2+160707r+49386}{(2r^2+7r+3)(2r^2+11r+12)(2r^2+15r+25)(2r^2+19r+42)(2r^2+23r+63)(2r^2+27r+88)(2r^2+31r+117)}$	$-\frac{4}{35721}$

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned}
y_2(x) &= \sqrt{x}(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 + b_7x^7 + b_8x^8 \dots) \\
&= \frac{1 - \frac{2x}{3} + \frac{5x^2}{6} + \frac{2x^3}{9} - \frac{19x^4}{216} - \frac{x^5}{540} + \frac{101x^6}{45360} - \frac{4x^7}{35721} + O(x^8)}{x^2}
\end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
y_h(x) &= c_1y_1(x) + c_2y_2(x) \\
&= c_1\sqrt{x} \left(1 - \frac{x}{14} - \frac{25x^2}{504} + \frac{197x^3}{33264} + \frac{1921x^4}{3459456} - \frac{11653x^5}{103783680} + \frac{12923x^6}{21171870720} \right. \\
&\quad \left. + \frac{917285x^7}{1126343522304} + O(x^8) \right) \\
&\quad + \frac{c_2 \left(1 - \frac{2x}{3} + \frac{5x^2}{6} + \frac{2x^3}{9} - \frac{19x^4}{216} - \frac{x^5}{540} + \frac{101x^6}{45360} - \frac{4x^7}{35721} + O(x^8) \right)}{x^2}
\end{aligned}$$

Hence the final solution is

$$\begin{aligned}
y &= y_h \\
&= c_1\sqrt{x} \left(1 - \frac{x}{14} - \frac{25x^2}{504} + \frac{197x^3}{33264} + \frac{1921x^4}{3459456} - \frac{11653x^5}{103783680} + \frac{12923x^6}{21171870720} \right. \\
&\quad \left. + \frac{917285x^7}{1126343522304} + O(x^8) \right) \\
&\quad + \frac{c_2 \left(1 - \frac{2x}{3} + \frac{5x^2}{6} + \frac{2x^3}{9} - \frac{19x^4}{216} - \frac{x^5}{540} + \frac{101x^6}{45360} - \frac{4x^7}{35721} + O(x^8) \right)}{x^2}
\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \sqrt{x} \left(1 - \frac{x}{14} - \frac{25x^2}{504} + \frac{197x^3}{33264} + \frac{1921x^4}{3459456} - \frac{11653x^5}{103783680} + \frac{12923x^6}{21171870720} + \frac{917285x^7}{1126343522304} + O(x^8) \right) + \frac{c_2 \left(1 - \frac{2x}{3} + \frac{5x^2}{6} + \frac{2x^3}{9} - \frac{19x^4}{216} - \frac{x^5}{540} + \frac{101x^6}{45360} - \frac{4x^7}{35721} + O(x^8) \right)}{x^2}$$

Verification of solutions

$$y = c_1 \sqrt{x} \left(1 - \frac{x}{14} - \frac{25x^2}{504} + \frac{197x^3}{33264} + \frac{1921x^4}{3459456} - \frac{11653x^5}{103783680} + \frac{12923x^6}{21171870720} + \frac{917285x^7}{1126343522304} + O(x^8) \right) + \frac{c_2 \left(1 - \frac{2x}{3} + \frac{5x^2}{6} + \frac{2x^3}{9} - \frac{19x^4}{216} - \frac{x^5}{540} + \frac{101x^6}{45360} - \frac{4x^7}{35721} + O(x^8) \right)}{x^2}$$

Verified OK.

19.1.1 Maple step by step solution

Let's solve

$$2x^2y'' + (x^2 + 5x)y' + (x^2 - 2)y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{(x^2-2)y}{2x^2} - \frac{(5+x)y'}{2x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{(5+x)y'}{2x} + \frac{(x^2-2)y}{2x^2} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$\left[P_2(x) = \frac{5+x}{2x}, P_3(x) = \frac{x^2-2}{2x^2} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = \frac{5}{2}$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = -1$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$2x^2y'' + x(5+x)y' + (x^2-2)y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 0..2$

$$x^m \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

- Shift index using $k- > k - m$

$$x^m \cdot y = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

- Convert $x^m \cdot y'$ to series expansion for $m = 1..2$

$$x^m \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1+m}$$

- Shift index using $k- > k + 1 - m$

$$x^m \cdot y' = \sum_{k=-1+m}^{\infty} a_{k+1-m} (k+1-m+r) x^{k+r}$$

- Convert $x^2 \cdot y''$ to series expansion

$$x^2 \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(2+r)(-1+2r)x^r + (a_1(3+r)(1+2r) + a_0r)x^{1+r} + \left(\sum_{k=2}^{\infty} (a_k(k+r+2)(2k+2r-1) + a_{k-2}(k+r-2)(k+r-1)) \right) x^{k+r}$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$(2+r)(-1+2r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \left\{-2, \frac{1}{2}\right\}$$

- Each term must be 0

$$a_1(3+r)(1+2r) + a_0r = 0$$

- Solve for the dependent coefficient(s)

$$a_1 = -\frac{a_0r}{2r^2+7r+3}$$

- Each term in the series must be 0, giving the recursion relation

$$2\left(k+r-\frac{1}{2}\right)(k+r+2)a_k + a_{k-1}k + a_{k-1}r + a_{k-2} - a_{k-1} = 0$$

- Shift index using $k- > k+2$

$$2\left(k+\frac{3}{2}+r\right)(k+4+r)a_{k+2} + a_{k+1}(k+2) + a_{k+1}r + a_k - a_{k+1} = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+2} = -\frac{ka_{k+1}+a_{k+1}r+a_k+a_{k+1}}{(2k+3+2r)(k+4+r)}$$

- Recursion relation for $r = -2$

$$a_{k+2} = -\frac{ka_{k+1}+a_k-a_{k+1}}{(2k-1)(k+2)}$$

- Solution for $r = -2$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k-2}, a_{k+2} = -\frac{ka_{k+1}+a_k-a_{k+1}}{(2k-1)(k+2)}, a_1 = -\frac{2a_0}{3} \right]$$

- Recursion relation for $r = \frac{1}{2}$

$$a_{k+2} = -\frac{ka_{k+1}+a_k+\frac{3}{2}a_{k+1}}{(2k+4)\left(k+\frac{9}{2}\right)}$$

- Solution for $r = \frac{1}{2}$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+\frac{1}{2}}, a_{k+2} = -\frac{ka_{k+1}+a_k+\frac{3}{2}a_{k+1}}{(2k+4)\left(k+\frac{9}{2}\right)}, a_1 = -\frac{a_0}{14} \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^{k-2} \right) + \left(\sum_{k=0}^{\infty} b_k x^{k+\frac{1}{2}} \right), a_{k+2} = -\frac{ka_{k+1}+a_k-a_{k+1}}{(2k-1)(k+2)}, a_1 = -\frac{2a_0}{3}, b_{k+2} = -\frac{kb_{k+1}+b_k+\frac{3}{2}b_{k+1}}{(2k+4)\left(k+\frac{9}{2}\right)}, \right]$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
  -> Bessel
  -> elliptic
  -> Legendre
  -> Whittaker
      -> hyper3: Equivalence to 1F1 under a power @ Moebius
      <- hyper3 successful: received ODE is equivalent to the 1F1 ODE
      <- Whittaker successful
<- special function solution successful`
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 55

```
Order:=8;
dsolve(2*x^2*diff(y(x),x$2)+(5*x+x^2)*diff(y(x),x)+(x^2-2)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{c_2 x^{\frac{5}{2}} \left(1 - \frac{1}{14}x - \frac{25}{504}x^2 + \frac{197}{33264}x^3 + \frac{1921}{3459456}x^4 - \frac{11653}{103783680}x^5 + \frac{12923}{21171870720}x^6 + \frac{917285}{1126343522304}x^7 + O(x^8) \right) + c_1}{x^2}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 116

```
AsymptoticDSolveValue[2*x^2*y'[x]+(5*x+x^2)*y'[x]+(x^2-2)*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \sqrt{x} \left(\frac{917285x^7}{1126343522304} + \frac{12923x^6}{21171870720} - \frac{11653x^5}{103783680} + \frac{1921x^4}{3459456} + \frac{197x^3}{33264} - \frac{25x^2}{504} - \frac{x}{14} + 1 \right) + \frac{c_2 \left(-\frac{4x^7}{35721} + \frac{101x^6}{45360} - \frac{x^5}{540} - \frac{19x^4}{216} + \frac{2x^3}{9} + \frac{5x^2}{6} - \frac{2x}{3} + 1 \right)}{x^2}$$

19.2 problem 1(ii)

Internal problem ID [6054]

Internal file name [OUTPUT/5302_Sunday_June_05_2022_03_33_16_PM_20004479/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 166

Problem number: 1(ii).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference is integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$4x^2y'' - 4y'e^xx + 3y \cos(x) = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$4x^2y'' - 4y'e^xx + 3y \cos(x) = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{e^x}{x}$$
$$q(x) = \frac{3 \cos(x)}{4x^2}$$

Table 230: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{e^x}{x}$	
singularity	type
$x = 0$	“regular”
$x = \infty$	“regular”

$q(x) = \frac{3 \cos(x)}{4x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0, \infty]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$4x^2 y'' - 4y' e^x x + 3y \cos(x) = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$4x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) - 4 \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) e^x x + 3 \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) \cos(x) = 0 \quad (1)$$

Expanding $-4x e^x$ as Taylor series around $x = 0$ and keeping only the first 8 terms gives

$$\begin{aligned} -4x e^x &= -4x - 4x^2 - 2x^3 - \frac{2}{3}x^4 - \frac{1}{6}x^5 - \frac{1}{30}x^6 - \frac{1}{180}x^7 - \frac{1}{1260}x^8 + \dots \\ &= -4x - 4x^2 - 2x^3 - \frac{2}{3}x^4 - \frac{1}{6}x^5 - \frac{1}{30}x^6 - \frac{1}{180}x^7 - \frac{1}{1260}x^8 \end{aligned}$$

Expanding $3 \cos(x)$ as Taylor series around $x = 0$ and keeping only the first 8 terms gives

$$\begin{aligned} 3 \cos(x) &= 3 - \frac{3}{2}x^2 + \frac{1}{8}x^4 - \frac{1}{240}x^6 + \frac{1}{13440}x^8 + \dots \\ &= 3 - \frac{3}{2}x^2 + \frac{1}{8}x^4 - \frac{1}{240}x^6 + \frac{1}{13440}x^8 \end{aligned}$$

Which simplifies to

$$\begin{aligned} &\left(\sum_{n=0}^{\infty} 4x^{n+r} a_n(n+r)(n+r-1) \right) + \sum_{n=0}^{\infty} \left(-\frac{x^{n+r+7} a_n(n+r)}{1260} \right) \\ &+ \sum_{n=0}^{\infty} \left(-\frac{x^{n+r+6} a_n(n+r)}{180} \right) + \sum_{n=0}^{\infty} \left(-\frac{x^{n+r+5} a_n(n+r)}{30} \right) \\ &+ \sum_{n=0}^{\infty} \left(-\frac{x^{n+r+4} a_n(n+r)}{6} \right) + \sum_{n=0}^{\infty} \left(-\frac{2x^{n+r+3} a_n(n+r)}{3} \right) \tag{2A} \\ &+ \sum_{n=0}^{\infty} (-2x^{n+r+2} a_n(n+r)) + \sum_{n=0}^{\infty} (-4x^{1+n+r} a_n(n+r)) \\ &+ \sum_{n=0}^{\infty} (-4x^{n+r} a_n(n+r)) + \left(\sum_{n=0}^{\infty} 3a_n x^{n+r} \right) + \sum_{n=0}^{\infty} \left(-\frac{3x^{n+r+2} a_n}{2} \right) \\ &+ \left(\sum_{n=0}^{\infty} \frac{x^{n+r+4} a_n}{8} \right) + \sum_{n=0}^{\infty} \left(-\frac{x^{n+r+6} a_n}{240} \right) + \left(\sum_{n=0}^{\infty} \frac{x^{n+r+8} a_n}{13440} \right) = 0 \end{aligned}$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=0}^{\infty} \left(-\frac{x^{n+r+7} a_n(n+r)}{1260} \right) &= \sum_{n=7}^{\infty} \left(-\frac{a_{n-7}(n-7+r) x^{n+r}}{1260} \right) \\ \sum_{n=0}^{\infty} \left(-\frac{x^{n+r+6} a_n(n+r)}{180} \right) &= \sum_{n=6}^{\infty} \left(-\frac{a_{n-6}(n-6+r) x^{n+r}}{180} \right) \end{aligned}$$

$$\begin{aligned}
\sum_{n=0}^{\infty} \left(-\frac{x^{n+r+5} a_n (n+r)}{30} \right) &= \sum_{n=5}^{\infty} \left(-\frac{a_{n-5} (n-5+r) x^{n+r}}{30} \right) \\
\sum_{n=0}^{\infty} \left(-\frac{x^{n+r+4} a_n (n+r)}{6} \right) &= \sum_{n=4}^{\infty} \left(-\frac{a_{n-4} (n-4+r) x^{n+r}}{6} \right) \\
\sum_{n=0}^{\infty} \left(-\frac{2x^{n+r+3} a_n (n+r)}{3} \right) &= \sum_{n=3}^{\infty} \left(-\frac{2a_{n-3} (n-3+r) x^{n+r}}{3} \right) \\
\sum_{n=0}^{\infty} (-2x^{n+r+2} a_n (n+r)) &= \sum_{n=2}^{\infty} (-2a_{n-2} (n+r-2) x^{n+r}) \\
\sum_{n=0}^{\infty} (-4x^{1+n+r} a_n (n+r)) &= \sum_{n=1}^{\infty} (-4a_{n-1} (n+r-1) x^{n+r}) \\
\sum_{n=0}^{\infty} \left(-\frac{3x^{n+r+2} a_n}{2} \right) &= \sum_{n=2}^{\infty} \left(-\frac{3a_{n-2} x^{n+r}}{2} \right) \\
\sum_{n=0}^{\infty} \frac{x^{n+r+4} a_n}{8} &= \sum_{n=4}^{\infty} \frac{a_{n-4} x^{n+r}}{8} \\
\sum_{n=0}^{\infty} \left(-\frac{x^{n+r+6} a_n}{240} \right) &= \sum_{n=6}^{\infty} \left(-\frac{a_{n-6} x^{n+r}}{240} \right) \\
\sum_{n=0}^{\infty} \frac{x^{n+r+8} a_n}{13440} &= \sum_{n=8}^{\infty} \frac{a_{n-8} x^{n+r}}{13440}
\end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers

of x are the same and equal to $n + r$.

$$\begin{aligned}
& \left(\sum_{n=0}^{\infty} 4x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=7}^{\infty} \left(-\frac{a_{n-7} (n-7+r) x^{n+r}}{1260} \right) \\
& + \sum_{n=6}^{\infty} \left(-\frac{a_{n-6} (n-6+r) x^{n+r}}{180} \right) + \sum_{n=5}^{\infty} \left(-\frac{a_{n-5} (n-5+r) x^{n+r}}{30} \right) \\
& + \sum_{n=4}^{\infty} \left(-\frac{a_{n-4} (n-4+r) x^{n+r}}{6} \right) + \sum_{n=3}^{\infty} \left(-\frac{2a_{n-3} (n-3+r) x^{n+r}}{3} \right) \quad (2B) \\
& + \sum_{n=2}^{\infty} (-2a_{n-2} (n+r-2) x^{n+r}) + \sum_{n=1}^{\infty} (-4a_{n-1} (n+r-1) x^{n+r}) \\
& + \sum_{n=0}^{\infty} (-4x^{n+r} a_n (n+r)) + \left(\sum_{n=0}^{\infty} 3a_n x^{n+r} \right) + \sum_{n=2}^{\infty} \left(-\frac{3a_{n-2} x^{n+r}}{2} \right) \\
& + \left(\sum_{n=4}^{\infty} \frac{a_{n-4} x^{n+r}}{8} \right) + \sum_{n=6}^{\infty} \left(-\frac{a_{n-6} x^{n+r}}{240} \right) + \left(\sum_{n=8}^{\infty} \frac{a_{n-8} x^{n+r}}{13440} \right) = 0
\end{aligned}$$

The indicial equation is obtained from $n = 0$. From Eq (2B) this gives

$$4x^{n+r} a_n (n+r) (n+r-1) - 4x^{n+r} a_n (n+r) + 3a_n x^{n+r} = 0$$

When $n = 0$ the above becomes

$$4x^r a_0 r (-1+r) - 4x^r a_0 r + 3a_0 x^r = 0$$

Or

$$(4x^r r (-1+r) - 4x^r r + 3x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(4r^2 - 8r + 3) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$4r^2 - 8r + 3 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned}
r_1 &= \frac{3}{2} \\
r_2 &= \frac{1}{2}
\end{aligned}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(4r^2 - 8r + 3) x^r = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = 1$ is an integer, then we can construct two linearly independent solutions

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = C y_1(x) \ln(x) + x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = x^{\frac{3}{2}} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = C y_1(x) \ln(x) + \sqrt{x} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+\frac{3}{2}}$$

$$y_2(x) = C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+\frac{1}{2}} \right)$$

Where C above can be zero. We start by finding y_1 . Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = \frac{4r}{4r^2 - 1}$$

Substituting $n = 2$ in Eq. (2B) gives

$$a_2 = \frac{8r^2 + 10r - 1}{16r^3 + 8r^2 - 4r - 2}$$

Substituting $n = 3$ in Eq. (2B) gives

$$a_3 = \frac{16r^4 + 104r^3 + 260r^2 + 154r - 12}{96r^5 + 432r^4 + 528r^3 + 72r^2 - 138r - 45}$$

Substituting $n = 4$ in Eq. (2B) gives

$$a_4 = \frac{128r^6 + 2272r^5 + 14096r^4 + 40832r^3 + 54968r^2 + 23922r - 2097}{3072 \left(r + \frac{1}{2}\right)^2 \left(r + \frac{3}{2}\right) \left(r + \frac{7}{2}\right) \left(r - \frac{1}{2}\right) \left(r + \frac{5}{2}\right)^2}$$

Substituting $n = 5$ in Eq. (2B) gives

$$a_5 = \frac{128r^8 + 5184r^7 + 63840r^6 + 386800r^5 + 1320392r^4 + 2573796r^3 + 2595110r^2 + 925655r - 83940}{30(2r+1)^2(3+2r)(2r+7)(2r-1)(2r+5)^2(4r^2+32r+63)}$$

Substituting $n = 6$ in Eq. (2B) gives

$$a_6 = \frac{2048r^{10} + 166400r^9 + 3645184r^8 + 39318272r^7 + 248963968r^6 + 989572160r^5 + 2499293216r^4 + 3840000000r^3 + 2400000000r^2 + 800000000r - 128000000}{1474560 \left(r + \frac{9}{2}\right)^2 \left(r + \frac{1}{2}\right)^2 \left(r + \frac{3}{2}\right) \left(r + \frac{7}{2}\right)^2 \left(r - \frac{1}{2}\right) \left(r + \frac{5}{2}\right)^2}$$

Substituting $n = 7$ in Eq. (2B) gives

$$a_7 = \frac{2048r^{12} + 316416r^{11} + 11362816r^{10} + 198127872r^9 + 2053881088r^8 + 13811070336r^7 + 62663041472r^6 + 20480000000r^5 + 4608000000r^4 + 640000000r^3 - 128000000r^2 + 12800000r - 1280000}{1260(2r+9)^2(2r+1)^2(3+2r)(2r-1)(2r+5)^2(4r^2+32r+63)}$$

For $8 \leq n$ the recursive equation is

$$\begin{aligned} &4a_n(n+r)(n+r-1) - \frac{a_{n-7}(n-7+r)}{1260} - \frac{a_{n-6}(n-6+r)}{180} \\ &- \frac{a_{n-5}(n-5+r)}{30} - \frac{a_{n-4}(n-4+r)}{6} - \frac{2a_{n-3}(n-3+r)}{3} - 2a_{n-2}(n+r-2) \\ &- 4a_{n-1}(n+r-1) - 4a_n(n+r) + 3a_n - \frac{3a_{n-2}}{2} + \frac{a_{n-4}}{8} - \frac{a_{n-6}}{240} + \frac{a_{n-8}}{13440} = 0 \end{aligned} \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{32na_{n-7} + 224na_{n-6} + 1344na_{n-5} + 6720na_{n-4} + 26880na_{n-3} + 80640na_{n-2} + 161280na_{n-1} + 320000na_{n-8} - 161280n}{161280n(1+n)} \quad (4)$$

Which for the root $r = \frac{3}{2}$ becomes

$$a_n = \frac{32(a_{n-7} + 7a_{n-6} + 42a_{n-5} + 210a_{n-4} + 840a_{n-3} + 2520a_{n-2} + 5040a_{n-1})n - 3a_{n-8} - 176a_{n-7} - 84a_{n-6} - 176a_{n-5} - 176a_{n-4} - 176a_{n-3} - 176a_{n-2} - 176a_{n-1} - 176a_{n-8}}{161280n(1+n)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = \frac{3}{2}$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$
a_0	1
a_1	$\frac{4r}{4r^2-1}$
a_2	$\frac{8r^2+10r-1}{16r^3+8r^2-4r-2}$
a_3	$\frac{16r^4+104r^3+260r^2+154r-12}{96r^5+432r^4+528r^3+72r^2-138r-45}$
a_4	$\frac{128r^6+2272r^5+14096r^4+40832r^3+54968r^2+23922r-2097}{3072(r+\frac{1}{2})^2(r+\frac{3}{2})(r+\frac{7}{2})(r-\frac{1}{2})(r+\frac{5}{2})^2}$
a_5	$\frac{128r^8+5184r^7+63840r^6+386800r^5+1320392r^4+2573796r^3+2595110r^2+925655r-83940}{30(2r+1)^2(3+2r)(2r+7)(2r-1)(2r+5)^2(4r^2+32r+63)}$
a_6	$\frac{2048r^{10}+166400r^9+3645184r^8+39318272r^7+248963968r^6+989572160r^5+2499293216r^4+3862419888r^3+3238012464r^2+1002158970r-100000}{1474560(r+\frac{9}{2})^2(r+\frac{1}{2})^2(r+\frac{3}{2})(r+\frac{7}{2})^2(r-\frac{1}{2})(r+\frac{5}{2})^2(r+\frac{11}{2})}$
a_7	$\frac{2048r^{12}+316416r^{11}+11362816r^{10}+198127872r^9+2053881088r^8+13811070336r^7+62663041472r^6+193775095008r^5+401584712904r^4+100000000000r^3+100000000000r^2+100000000000r-100000000000}{1260(2r+9)^2(2r+1)^2(3+2r)(2r+7)^2(2r-1)(2r+5)^2(2r+11)(4r^2+48r+100)}$

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned}
y_1(x) &= x^{\frac{3}{2}} (a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots) \\
&= x^{\frac{3}{2}} \left(1 + \frac{3x}{4} + \frac{x^2}{2} + \frac{103x^3}{384} + \frac{669x^4}{5120} + \frac{54731x^5}{921600} + \frac{123443x^6}{4838400} + \frac{30273113x^7}{2890137600} + O(x^8) \right)
\end{aligned}$$

Now the second solution $y_2(x)$ is found. Let

$$r_1 - r_2 = N$$

Where N is positive integer which is the difference between the two roots. r_1 is taken as the larger root. Hence for this problem we have $N = 1$. Now we need to determine if C is zero or not. This is done by finding $\lim_{r \rightarrow r_2} a_1(r)$. If this limit exists, then $C = 0$, else we need to keep the log term and $C \neq 0$. The above table shows that

$$\begin{aligned}
a_N &= a_1 \\
&= \frac{4r}{4r^2-1}
\end{aligned}$$

Therefore

$$\begin{aligned}
\lim_{r \rightarrow r_2} \frac{4r}{4r^2-1} &= \lim_{r \rightarrow \frac{1}{2}} \frac{4r}{4r^2-1} \\
&= \text{undefined}
\end{aligned}$$

Since the limit does not exist then the log term is needed. Therefore the second solution has the form

$$y_2(x) = Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right)$$

Therefore

$$\begin{aligned} \frac{d}{dx}y_2(x) &= Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2}(n+r_2)}{x} \right) \\ &= Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} x^{-1+n+r_2} b_n (n+r_2) \right) \\ \frac{d^2}{dx^2}y_2(x) &= Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} \\ &\quad + \sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2}(n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2}(n+r_2)}{x^2} \right) \\ &= Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} + \left(\sum_{n=0}^{\infty} x^{-2+n+r_2} b_n (n+r_2) (-1+n+r_2) \right) \end{aligned}$$

Substituting these back into the given ode $4x^2y'' - 4y'e^x + 3y \cos(x) = 0$ gives

$$\begin{aligned} &4x^2 \left(Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} \right. \\ &\quad \left. + \sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2}(n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2}(n+r_2)}{x^2} \right) \right) \\ &\quad - 4 \left(Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2}(n+r_2)}{x} \right) \right) e^x x \\ &\quad + 3 \left(Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) \right) \cos(x) = 0 \end{aligned}$$

Which can be written as

$$\begin{aligned} &\left((-4e^x y_1'(x) x + 4y_1''(x) x^2 + 3 \cos(x) y_1(x)) \ln(x) \right. \\ &\quad \left. + 4x^2 \left(\frac{2y_1'(x)}{x} - \frac{y_1(x)}{x^2} \right) - 4y_1(x) e^x \right) C - 4 \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2}(n+r_2)}{x} \right) e^x x \\ &\quad + 4x^2 \left(\sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2}(n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2}(n+r_2)}{x^2} \right) \right) \\ &\quad + 3 \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) \cos(x) = 0 \end{aligned} \tag{7}$$

But since $y_1(x)$ is a solution to the ode, then

$$-4e^x y_1'(x) x + 4y_1''(x) x^2 + 3 \cos(x) y_1(x) = 0$$

Eq (7) simplifies to

$$\begin{aligned} & \left(4x^2 \left(\frac{2y_1'(x)}{x} - \frac{y_1(x)}{x^2} \right) - 4y_1(x) e^x \right) C - 4 \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2} (n+r_2)}{x} \right) e^x x \\ & + 4x^2 \left(\sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2} (n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2} (n+r_2)}{x^2} \right) \right) \\ & + 3 \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) \cos(x) = 0 \end{aligned} \quad (8)$$

Substituting $y_1 = \sum_{n=0}^{\infty} a_n x^{n+r_1}$ into the above gives

$$\begin{aligned} & \left(8 \left(\sum_{n=0}^{\infty} x^{-1+n+r_1} a_n (n+r_1) \right) x - 4 \left(\sum_{n=0}^{\infty} a_n x^{n+r_1} \right) (1+e^x) \right) C \\ & - 4 \left(\sum_{n=0}^{\infty} x^{-1+n+r_2} b_n (n+r_2) \right) e^x x \\ & + 4 \left(\sum_{n=0}^{\infty} x^{-2+n+r_2} b_n (n+r_2) (-1+n+r_2) \right) x^2 + 3 \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) \cos(x) = 0 \end{aligned} \quad (9)$$

Since $r_1 = \frac{3}{2}$ and $r_2 = \frac{1}{2}$ then the above becomes

$$\begin{aligned} & \left(8 \left(\sum_{n=0}^{\infty} x^{n+\frac{1}{2}} a_n \left(n + \frac{3}{2} \right) \right) x - 4 \left(\sum_{n=0}^{\infty} a_n x^{n+\frac{3}{2}} \right) (1+e^x) \right) C \\ & - 4 \left(\sum_{n=0}^{\infty} x^{-\frac{1}{2}+n} b_n \left(n + \frac{1}{2} \right) \right) e^x x \\ & + 4 \left(\sum_{n=0}^{\infty} x^{-\frac{3}{2}+n} b_n \left(n + \frac{1}{2} \right) \left(-\frac{1}{2} + n \right) \right) x^2 + 3 \left(\sum_{n=0}^{\infty} b_n x^{n+\frac{1}{2}} \right) \cos(x) = 0 \end{aligned} \quad (10)$$

Expanding $-4C x^{\frac{3}{2}}$ as Taylor series around $x = 0$ and keeping only the first 8 terms gives

$$\begin{aligned} -4C x^{\frac{3}{2}} &= -4C x^{\frac{3}{2}} + \dots \\ &= -4C x^{\frac{3}{2}} \end{aligned}$$

Expanding $-4C x^{\frac{3}{2}} e^x$ as Taylor series around $x = 0$ and keeping only the first 8 terms gives

$$\begin{aligned} -4C x^{\frac{3}{2}} e^x &= -4C x^{\frac{3}{2}} - 4C x^{\frac{5}{2}} - 2C x^{\frac{7}{2}} - \frac{2C x^{\frac{9}{2}}}{3} - \frac{C x^{\frac{11}{2}}}{6} - \frac{C x^{\frac{13}{2}}}{30} - \frac{C x^{\frac{15}{2}}}{180} - \frac{C x^{\frac{17}{2}}}{1260} + \dots \\ &= -4C x^{\frac{3}{2}} - 4C x^{\frac{5}{2}} - 2C x^{\frac{7}{2}} - \frac{2C x^{\frac{9}{2}}}{3} - \frac{C x^{\frac{11}{2}}}{6} - \frac{C x^{\frac{13}{2}}}{30} - \frac{C x^{\frac{15}{2}}}{180} - \frac{C x^{\frac{17}{2}}}{1260} \end{aligned}$$

Expanding $-2\sqrt{x} e^x$ as Taylor series around $x = 0$ and keeping only the first 8 terms gives

$$\begin{aligned} -2\sqrt{x} e^x &= -2\sqrt{x} - 2x^{\frac{3}{2}} - x^{\frac{5}{2}} - \frac{x^{\frac{7}{2}}}{3} - \frac{x^{\frac{9}{2}}}{12} - \frac{x^{\frac{11}{2}}}{60} - \frac{x^{\frac{13}{2}}}{360} - \frac{x^{\frac{15}{2}}}{2520} + \dots \\ &= -2\sqrt{x} - 2x^{\frac{3}{2}} - x^{\frac{5}{2}} - \frac{x^{\frac{7}{2}}}{3} - \frac{x^{\frac{9}{2}}}{12} - \frac{x^{\frac{11}{2}}}{60} - \frac{x^{\frac{13}{2}}}{360} - \frac{x^{\frac{15}{2}}}{2520} \end{aligned}$$

Expanding $3 \cos(x)$ as Taylor series around $x = 0$ and keeping only the first 8 terms gives

$$\begin{aligned} 3 \cos(x) &= 3 - \frac{3}{2}x^2 + \frac{1}{8}x^4 - \frac{1}{240}x^6 + \frac{1}{13440}x^8 + \dots \\ &= 3 - \frac{3}{2}x^2 + \frac{1}{8}x^4 - \frac{1}{240}x^6 + \frac{1}{13440}x^8 \end{aligned}$$

Which simplifies to

$$\begin{aligned}
& \left(\sum_{n=0}^{\infty} (8n+12) C a_n x^{n+\frac{3}{2}} \right) + 2 \left(\sum_{n=0}^{\infty} (-4C x^{n+\frac{3}{2}} a_n) \right) \\
& + \sum_{n=0}^{\infty} (-4C x^{n+\frac{5}{2}} a_n) + \sum_{n=0}^{\infty} (-2C x^{n+\frac{7}{2}} a_n) + \sum_{n=0}^{\infty} \left(-\frac{2C x^{n+\frac{9}{2}} a_n}{3} \right) \\
& + \sum_{n=0}^{\infty} \left(-\frac{C x^{n+\frac{11}{2}} a_n}{6} \right) + \sum_{n=0}^{\infty} \left(-\frac{C x^{n+\frac{13}{2}} a_n}{30} \right) \\
& + \sum_{n=0}^{\infty} \left(-\frac{C x^{n+\frac{15}{2}} a_n}{180} \right) + \sum_{n=0}^{\infty} \left(-\frac{C x^{n+\frac{17}{2}} a_n}{1260} \right) \\
& + \sum_{n=0}^{\infty} \left(-\frac{x^{n+\frac{15}{2}} b_n (2n+1)}{2520} \right) + \sum_{n=0}^{\infty} \left(-\frac{x^{n+\frac{13}{2}} b_n (2n+1)}{360} \right) \\
& + \sum_{n=0}^{\infty} \left(-\frac{x^{n+\frac{11}{2}} b_n (2n+1)}{60} \right) + \sum_{n=0}^{\infty} \left(-\frac{x^{n+\frac{9}{2}} b_n (2n+1)}{12} \right) \\
& + \sum_{n=0}^{\infty} \left(-\frac{x^{n+\frac{7}{2}} b_n (2n+1)}{3} \right) + \left(\sum_{n=0}^{\infty} x^{n+\frac{5}{2}} b_n (-2n-1) \right) \\
& + \left(\sum_{n=0}^{\infty} (-4n-2) b_n x^{n+\frac{3}{2}} \right) + \left(\sum_{n=0}^{\infty} (-4n-2) b_n x^{n+\frac{1}{2}} \right) \\
& + \left(\sum_{n=0}^{\infty} x^{n+\frac{1}{2}} b_n (4n^2-1) \right) + \left(\sum_{n=0}^{\infty} 3b_n x^{n+\frac{1}{2}} \right) + \sum_{n=0}^{\infty} \left(-\frac{3x^{n+\frac{5}{2}} b_n}{2} \right) \\
& + \left(\sum_{n=0}^{\infty} \frac{x^{n+\frac{9}{2}} b_n}{8} \right) + \sum_{n=0}^{\infty} \left(-\frac{x^{n+\frac{13}{2}} b_n}{240} \right) + \left(\sum_{n=0}^{\infty} \frac{x^{n+\frac{17}{2}} b_n}{13440} \right) = 0
\end{aligned} \tag{2A}$$

The next step is to make all powers of x be $n + \frac{1}{2}$ in each summation term. Going over each summation term above with power of x in it which is not already $x^{n+\frac{1}{2}}$ and adjusting the power and the corresponding index gives

$$\begin{aligned}
\sum_{n=0}^{\infty} (8n+12) C a_n x^{n+\frac{3}{2}} &= \sum_{n=1}^{\infty} C a_{n-1} (8n+4) x^{n+\frac{1}{2}} \\
\sum_{n=0}^{\infty} (-4C x^{n+\frac{3}{2}} a_n) &= \sum_{n=1}^{\infty} (-4C a_{n-1} x^{n+\frac{1}{2}}) \\
\sum_{n=0}^{\infty} (-4C x^{n+\frac{5}{2}} a_n) &= \sum_{n=1}^{\infty} (-4C a_{n-1} x^{n+\frac{1}{2}})
\end{aligned}$$

$$\sum_{n=0}^{\infty} \left(-4C x^{n+\frac{5}{2}} a_n \right) = \sum_{n=2}^{\infty} \left(-4C a_{n-2} x^{n+\frac{1}{2}} \right)$$

$$\sum_{n=0}^{\infty} \left(-2C x^{n+\frac{7}{2}} a_n \right) = \sum_{n=3}^{\infty} \left(-2C a_{n-3} x^{n+\frac{1}{2}} \right)$$

$$\sum_{n=0}^{\infty} \left(-\frac{2C x^{n+\frac{9}{2}} a_n}{3} \right) = \sum_{n=4}^{\infty} \left(-\frac{2C a_{n-4} x^{n+\frac{1}{2}}}{3} \right)$$

$$\sum_{n=0}^{\infty} \left(-\frac{C x^{n+\frac{11}{2}} a_n}{6} \right) = \sum_{n=5}^{\infty} \left(-\frac{C a_{n-5} x^{n+\frac{1}{2}}}{6} \right)$$

$$\sum_{n=0}^{\infty} \left(-\frac{C x^{n+\frac{13}{2}} a_n}{30} \right) = \sum_{n=6}^{\infty} \left(-\frac{C a_{n-6} x^{n+\frac{1}{2}}}{30} \right)$$

$$\sum_{n=0}^{\infty} \left(-\frac{C x^{n+\frac{15}{2}} a_n}{180} \right) = \sum_{n=7}^{\infty} \left(-\frac{C a_{n-7} x^{n+\frac{1}{2}}}{180} \right)$$

$$\sum_{n=0}^{\infty} \left(-\frac{C x^{n+\frac{17}{2}} a_n}{1260} \right) = \sum_{n=8}^{\infty} \left(-\frac{C a_{n-8} x^{n+\frac{1}{2}}}{1260} \right)$$

$$\sum_{n=0}^{\infty} \left(-\frac{x^{n+\frac{15}{2}} b_n (2n+1)}{2520} \right) = \sum_{n=7}^{\infty} \left(-\frac{b_{n-7} (2n-13) x^{n+\frac{1}{2}}}{2520} \right)$$

$$\sum_{n=0}^{\infty} \left(-\frac{x^{n+\frac{13}{2}} b_n (2n+1)}{360} \right) = \sum_{n=6}^{\infty} \left(-\frac{b_{n-6} (2n-11) x^{n+\frac{1}{2}}}{360} \right)$$

$$\sum_{n=0}^{\infty} \left(-\frac{x^{n+\frac{11}{2}} b_n (2n+1)}{60} \right) = \sum_{n=5}^{\infty} \left(-\frac{b_{n-5} (2n-9) x^{n+\frac{1}{2}}}{60} \right)$$

$$\sum_{n=0}^{\infty} \left(-\frac{x^{n+\frac{9}{2}} b_n (2n+1)}{12} \right) = \sum_{n=4}^{\infty} \left(-\frac{b_{n-4} (2n-7) x^{n+\frac{1}{2}}}{12} \right)$$

$$\sum_{n=0}^{\infty} \left(-\frac{x^{n+\frac{7}{2}} b_n (2n+1)}{3} \right) = \sum_{n=3}^{\infty} \left(-\frac{b_{n-3} (2n-5) x^{n+\frac{1}{2}}}{3} \right)$$

$$\sum_{n=0}^{\infty} x^{n+\frac{5}{2}} b_n (-2n-1) = \sum_{n=2}^{\infty} b_{n-2} (-2n+3) x^{n+\frac{1}{2}}$$

$$\begin{aligned}
\sum_{n=0}^{\infty} (-4n - 2) b_n x^{n+\frac{3}{2}} &= \sum_{n=1}^{\infty} b_{n-1} (-4n + 2) x^{n+\frac{1}{2}} \\
\sum_{n=0}^{\infty} \left(-\frac{3x^{n+\frac{5}{2}} b_n}{2} \right) &= \sum_{n=2}^{\infty} \left(-\frac{3b_{n-2} x^{n+\frac{1}{2}}}{2} \right) \\
\sum_{n=0}^{\infty} \frac{x^{n+\frac{9}{2}} b_n}{8} &= \sum_{n=4}^{\infty} \frac{b_{n-4} x^{n+\frac{1}{2}}}{8} \\
\sum_{n=0}^{\infty} \left(-\frac{x^{n+\frac{13}{2}} b_n}{240} \right) &= \sum_{n=6}^{\infty} \left(-\frac{b_{n-6} x^{n+\frac{1}{2}}}{240} \right) \\
\sum_{n=0}^{\infty} \frac{x^{n+\frac{17}{2}} b_n}{13440} &= \sum_{n=8}^{\infty} \frac{b_{n-8} x^{n+\frac{1}{2}}}{13440}
\end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers

of x are the same and equal to $n + \frac{1}{2}$.

$$\begin{aligned}
& \left(\sum_{n=1}^{\infty} C a_{n-1} (8n+4) x^{n+\frac{1}{2}} \right) + 2 \left(\sum_{n=1}^{\infty} (-4C a_{n-1} x^{n+\frac{1}{2}}) \right) \\
& + \sum_{n=2}^{\infty} (-4C a_{n-2} x^{n+\frac{1}{2}}) + \sum_{n=3}^{\infty} (-2C a_{n-3} x^{n+\frac{1}{2}}) \\
& + \sum_{n=4}^{\infty} \left(-\frac{2C a_{n-4} x^{n+\frac{1}{2}}}{3} \right) + \sum_{n=5}^{\infty} \left(-\frac{C a_{n-5} x^{n+\frac{1}{2}}}{6} \right) \\
& + \sum_{n=6}^{\infty} \left(-\frac{C a_{n-6} x^{n+\frac{1}{2}}}{30} \right) + \sum_{n=7}^{\infty} \left(-\frac{C a_{n-7} x^{n+\frac{1}{2}}}{180} \right) \\
& + \sum_{n=8}^{\infty} \left(-\frac{C a_{n-8} x^{n+\frac{1}{2}}}{1260} \right) + \sum_{n=7}^{\infty} \left(-\frac{b_{n-7} (2n-13) x^{n+\frac{1}{2}}}{2520} \right) \\
& + \sum_{n=6}^{\infty} \left(-\frac{b_{n-6} (2n-11) x^{n+\frac{1}{2}}}{360} \right) + \sum_{n=5}^{\infty} \left(-\frac{b_{n-5} (2n-9) x^{n+\frac{1}{2}}}{60} \right) \quad (2B) \\
& + \sum_{n=4}^{\infty} \left(-\frac{b_{n-4} (2n-7) x^{n+\frac{1}{2}}}{12} \right) + \sum_{n=3}^{\infty} \left(-\frac{b_{n-3} (2n-5) x^{n+\frac{1}{2}}}{3} \right) \\
& + \left(\sum_{n=2}^{\infty} b_{n-2} (-2n+3) x^{n+\frac{1}{2}} \right) + \left(\sum_{n=1}^{\infty} b_{n-1} (-4n+2) x^{n+\frac{1}{2}} \right) \\
& + \left(\sum_{n=0}^{\infty} (-4n-2) b_n x^{n+\frac{1}{2}} \right) + \left(\sum_{n=0}^{\infty} x^{n+\frac{1}{2}} b_n (4n^2-1) \right) \\
& + \left(\sum_{n=0}^{\infty} 3b_n x^{n+\frac{1}{2}} \right) + \sum_{n=2}^{\infty} \left(-\frac{3b_{n-2} x^{n+\frac{1}{2}}}{2} \right) + \left(\sum_{n=4}^{\infty} \frac{b_{n-4} x^{n+\frac{1}{2}}}{8} \right) \\
& + \sum_{n=6}^{\infty} \left(-\frac{b_{n-6} x^{n+\frac{1}{2}}}{240} \right) + \left(\sum_{n=8}^{\infty} \frac{b_{n-8} x^{n+\frac{1}{2}}}{13440} \right) = 0
\end{aligned}$$

For $n = 0$ in Eq. (2B), we choose arbitrary value for b_0 as $b_0 = 1$. For $n = N$, where $N = 1$ which is the difference between the two roots, we are free to choose $b_1 = 0$. Hence for $n = 1$, Eq (2B) gives

$$4C - 2 = 0$$

Which is solved for C . Solving for C gives

$$C = \frac{1}{2}$$

For $n = 2$, Eq (2B) gives

$$4(-a_0 + 3a_1)C - \frac{5b_0}{2} - 6b_1 + 8b_2 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$8b_2 = 0$$

Solving the above for b_2 gives

$$b_2 = 0$$

For $n = 3$, Eq (2B) gives

$$2(-a_0 - 2a_1 + 10a_2)C - \frac{b_0}{3} - \frac{9b_1}{2} - 10b_2 + 24b_3 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$\frac{13}{6} + 24b_3 = 0$$

Solving the above for b_3 gives

$$b_3 = -\frac{13}{144}$$

For $n = 4$, Eq (2B) gives

$$\frac{2(-a_0 - 3a_1 - 6a_2 + 42a_3)C}{3} + 48b_4 + \frac{b_0}{24} - b_1 - \frac{13b_2}{2} - 14b_3 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$\frac{1715}{576} + 48b_4 = 0$$

Solving the above for b_4 gives

$$b_4 = -\frac{1715}{27648}$$

For $n = 5$, Eq (2B) gives

$$\frac{(-a_0 - 4a_1 - 12a_2 - 24a_3 + 216a_4)C}{6} - \frac{17b_3}{2} - 18b_4 + 80b_5 - \frac{b_0}{60} - \frac{b_1}{8} - \frac{5b_2}{3} = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$\frac{6565}{2304} + 80b_5 = 0$$

Solving the above for b_5 gives

$$b_5 = -\frac{1313}{36864}$$

For $n = 6$, Eq (2B) gives

$$\frac{(-a_0 - 5a_1 - 20a_2 - 60a_3 - 120a_4 + 1320a_5)C}{30} - \frac{7b_2}{24} - \frac{7b_3}{3} - \frac{21b_4}{2} - 22b_5 + 120b_6 - \frac{b_0}{144} - \frac{b_1}{20} = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$\frac{2999423}{1382400} + 120b_6 = 0$$

Solving the above for b_6 gives

$$b_6 = -\frac{2999423}{165888000}$$

For $n = 7$, Eq (2B) gives

$$\frac{(-a_0 - 6a_1 - 30a_2 - 120a_3 - 360a_4 - 720a_5 + 9360a_6)C}{180} - \frac{b_1}{80} - \frac{b_2}{12} - \frac{11b_3}{24} - 3b_4 - \frac{25b_5}{2} - 26b_6 + 168b_7 - \frac{b_0}{2520} = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$\frac{204656267}{145152000} + 168b_7 = 0$$

Solving the above for b_7 gives

$$b_7 = -\frac{204656267}{24385536000}$$

Now that we found all b_n and C , we can calculate the second solution from

$$y_2(x) = Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right)$$

Using the above value found for $C = \frac{1}{2}$ and all b_n , then the second solution becomes

$$y_2(x) = \frac{1}{2} \left(x^{\frac{3}{2}} \left(1 + \frac{3x}{4} + \frac{x^2}{2} + \frac{103x^3}{384} + \frac{669x^4}{5120} + \frac{54731x^5}{921600} + \frac{123443x^6}{4838400} + \frac{30273113x^7}{2890137600} + O(x^8) \right) \right) \ln(x) + \sqrt{x} \left(1 - \frac{13x^3}{144} - \frac{1715x^4}{27648} - \frac{1313x^5}{36864} - \frac{2999423x^6}{165888000} - \frac{204656267x^7}{24385536000} + O(x^8) \right)$$

Therefore the homogeneous solution is

$$\begin{aligned}
 y_h(x) &= c_1 y_1(x) + c_2 y_2(x) \\
 &= c_1 x^{\frac{3}{2}} \left(1 + \frac{3x}{4} + \frac{x^2}{2} + \frac{103x^3}{384} + \frac{669x^4}{5120} + \frac{54731x^5}{921600} + \frac{123443x^6}{4838400} + \frac{30273113x^7}{2890137600} + O(x^8) \right) \\
 &\quad + c_2 \left(\frac{1}{2} \left(x^{\frac{3}{2}} \left(1 + \frac{3x}{4} + \frac{x^2}{2} + \frac{103x^3}{384} + \frac{669x^4}{5120} + \frac{54731x^5}{921600} + \frac{123443x^6}{4838400} + \frac{30273113x^7}{2890137600} + O(x^8) \right) \right) \ln(x) \right. \\
 &\quad \left. + \sqrt{x} \left(1 - \frac{13x^3}{144} - \frac{1715x^4}{27648} - \frac{1313x^5}{36864} - \frac{2999423x^6}{165888000} - \frac{204656267x^7}{24385536000} + O(x^8) \right) \right)
 \end{aligned}$$

Hence the final solution is

$$\begin{aligned}
 y &= y_h \\
 &= c_1 x^{\frac{3}{2}} \left(1 + \frac{3x}{4} + \frac{x^2}{2} + \frac{103x^3}{384} + \frac{669x^4}{5120} + \frac{54731x^5}{921600} + \frac{123443x^6}{4838400} + \frac{30273113x^7}{2890137600} + O(x^8) \right) \\
 &\quad + c_2 \left(\frac{x^{\frac{3}{2}} \left(1 + \frac{3x}{4} + \frac{x^2}{2} + \frac{103x^3}{384} + \frac{669x^4}{5120} + \frac{54731x^5}{921600} + \frac{123443x^6}{4838400} + \frac{30273113x^7}{2890137600} + O(x^8) \right) \ln(x)}{2} \right. \\
 &\quad \left. + \sqrt{x} \left(1 - \frac{13x^3}{144} - \frac{1715x^4}{27648} - \frac{1313x^5}{36864} - \frac{2999423x^6}{165888000} - \frac{204656267x^7}{24385536000} + O(x^8) \right) \right)
 \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}
 y &= c_1 x^{\frac{3}{2}} \left(1 + \frac{3x}{4} + \frac{x^2}{2} + \frac{103x^3}{384} + \frac{669x^4}{5120} + \frac{54731x^5}{921600} + \frac{123443x^6}{4838400} + \frac{30273113x^7}{2890137600} + O(x^8) \right) \\
 &\quad + c_2 \left(\frac{x^{\frac{3}{2}} \left(1 + \frac{3x}{4} + \frac{x^2}{2} + \frac{103x^3}{384} + \frac{669x^4}{5120} + \frac{54731x^5}{921600} + \frac{123443x^6}{4838400} + \frac{30273113x^7}{2890137600} + O(x^8) \right) \ln(x)}{2} \right. \\
 &\quad \left. + \sqrt{x} \left(1 - \frac{13x^3}{144} - \frac{1715x^4}{27648} - \frac{1313x^5}{36864} - \frac{2999423x^6}{165888000} - \frac{204656267x^7}{24385536000} + O(x^8) \right) \right)
 \end{aligned} \tag{1}$$

Verification of solutions

$$y = c_1 x^{\frac{3}{2}} \left(1 + \frac{3x}{4} + \frac{x^2}{2} + \frac{103x^3}{384} + \frac{669x^4}{5120} + \frac{54731x^5}{921600} + \frac{123443x^6}{4838400} + \frac{30273113x^7}{2890137600} + O(x^8) \right) \\ + c_2 \left(\frac{x^{\frac{3}{2}} \left(1 + \frac{3x}{4} + \frac{x^2}{2} + \frac{103x^3}{384} + \frac{669x^4}{5120} + \frac{54731x^5}{921600} + \frac{123443x^6}{4838400} + \frac{30273113x^7}{2890137600} + O(x^8) \right) \ln(x)}{2} \right. \\ \left. + \sqrt{x} \left(1 - \frac{13x^3}{144} - \frac{1715x^4}{27648} - \frac{1313x^5}{36864} - \frac{2999423x^6}{165888000} - \frac{204656267x^7}{24385536000} + O(x^8) \right) \right)$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) * 2F1([a
-> Trying changes of variables to rationalize or make the ODE simpler
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) * 2F1
trying a symmetry of the form [xi=0, eta=F(x)]
trying 2nd order exact linear
trying symmetries linear in x and y(x)
trying to convert to a linear ODE with constant coefficients
-> trying with_periodic_functions in the coefficients
--- Trying Lie symmetry methods, 2nd order ---
`, `-> Computing symmetries using: way = 5
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) * 2F1
trying a symmetry of the form [xi=0, eta=F(x)]
trying 2nd order exact linear
trying symmetries linear in x and y(x)
trying to convert to a linear ODE with constant coefficients
-> trying with_periodic_functions in the coefficients
--- Trying Lie symmetry methods, 2nd order ---
`, `-> Computing symmetries using: way = 5
<- unable to find a useful change of variables
trying a symmetry of the form [xi=0, eta=F(x)]
trying differential order: 2; exact nonlinear
trying symmetries linear in x and y(x)
trying to convert to a linear ODE with constant coefficients
trying 2nd order, integrating factor of the form mu(x,y)
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) * 2F1
-> Trying changes of variables to rationalize or make the ODE simpler
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) *
trying a symmetry of the form [xi=0, eta=F(x)]
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 81

Order:=8;

dsolve(4*x^2*diff(y(x),x\$2)-4*x*exp(x)*diff(y(x),x)+3*cos(x)*y(x)=0,y(x),type='series',x=0);

$$y(x) = \left(x \left(1 + \frac{3}{4}x + \frac{1}{2}x^2 + \frac{103}{384}x^3 + \frac{669}{5120}x^4 + \frac{54731}{921600}x^5 + \frac{123443}{4838400}x^6 + \frac{30273113}{2890137600}x^7 + O(x^8) \right) c_1 + c_2 \left(\ln(x) \left(\frac{1}{2}x + \frac{3}{8}x^2 + \frac{1}{4}x^3 + \frac{103}{768}x^4 + \frac{669}{10240}x^5 + \frac{54731}{1843200}x^6 + \frac{123443}{9676800}x^7 + O(x^8) \right) + \left(1 + x + \frac{3}{4}x^2 + \frac{59}{144}x^3 + \frac{5701}{27648}x^4 + \frac{17519}{184320}x^5 + \frac{6852157}{165888000}x^6 + \frac{417496453}{24385536000}x^7 + O(x^8) \right) \right) \right) \sqrt{x}$$

✓ Solution by Mathematica

Time used: 0.146 (sec). Leaf size: 146

AsymptoticDSolveValue[4*x^2*y''[x]-4*x*Exp[x]*y'[x]+3*Cos[x]*y[x]==0,y[x],{x,0,7}]

$$y(x) \rightarrow c_2 \left(\frac{123443x^{15/2}}{4838400} + \frac{54731x^{13/2}}{921600} + \frac{669x^{11/2}}{5120} + \frac{103x^{9/2}}{384} + \frac{x^{7/2}}{2} + \frac{3x^{5/2}}{4} + x^{3/2} \right) + c_1 \left(\frac{(54731x^5 + 120420x^4 + 247200x^3 + 460800x^2 + 691200x + 921600)x^{3/2} \log(x)}{1843200} + \frac{(192636}{1843200} \right)$$

19.3 problem 1(iii)

Internal problem ID [6055]

Internal file name [OUTPUT/5303_Sunday_June_05_2022_03_33_28_PM_47241310/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 166

Problem number: 1(iii).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Repeated root"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$(-x^2 + 1)x^2y'' + 3(x^2 + x)y' + y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$(-x^4 + x^2)y'' + (3x^2 + 3x)y' + y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{3}{x(x-1)}$$
$$q(x) = -\frac{1}{x^2(x^2-1)}$$

Table 231: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{3}{x(x-1)}$	
singularity	type
$x = 0$	“regular”
$x = 1$	“regular”

$q(x) = -\frac{1}{x^2(x^2-1)}$	
singularity	type
$x = -1$	“regular”
$x = 0$	“regular”
$x = 1$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0, 1, -1, \infty]$

Irregular singular points : $[\]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$-y''x^2(x^2 - 1) + (3x^2 + 3x)y' + y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned} & - \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) x^2(x^2 - 1) \\ & + (3x^2 + 3x) \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \end{aligned} \tag{1}$$

Which simplifies to

$$\begin{aligned} & \sum_{n=0}^{\infty} (-x^{n+r+2} a_n (n+r) (n+r-1)) + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) \\ & + \left(\sum_{n=0}^{\infty} 3x^{1+n+r} a_n (n+r) \right) + \left(\sum_{n=0}^{\infty} 3x^{n+r} a_n (n+r) \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=0}^{\infty} (-x^{n+r+2} a_n (n+r) (n+r-1)) &= \sum_{n=2}^{\infty} (-a_{n-2} (n+r-2) (n-3+r) x^{n+r}) \\ \sum_{n=0}^{\infty} 3x^{1+n+r} a_n (n+r) &= \sum_{n=1}^{\infty} 3a_{n-1} (n+r-1) x^{n+r} \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \sum_{n=2}^{\infty} (-a_{n-2} (n+r-2) (n-3+r) x^{n+r}) + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) \\ & + \left(\sum_{n=1}^{\infty} 3a_{n-1} (n+r-1) x^{n+r} \right) + \left(\sum_{n=0}^{\infty} 3x^{n+r} a_n (n+r) \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r} a_n (n+r) (n+r-1) + 3x^{n+r} a_n (n+r) + a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$x^r a_0 r (-1+r) + 3x^r a_0 r + a_0 x^r = 0$$

Or

$$(x^r r (-1+r) + 3x^r r + x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(r + 1)^2 x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$(r + 1)^2 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = -1$$

$$r_2 = -1$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(r + 1)^2 x^r = 0$$

Solving for r gives the roots of the indicial equation as Since the root of the indicial equation is repeated, then we can construct two linearly independent solutions. The first solution has the form

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+r} \quad (1A)$$

Now the second solution y_2 is found using

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right) \quad (1B)$$

Then the general solution will be

$$y = c_1 y_1(x) + c_2 y_2(x)$$

In Eq (1B) the sum starts from 1 and not zero. In Eq (1A), a_0 is never zero, and is arbitrary and is typically taken as $a_0 = 1$, and $\{c_1, c_2\}$ are two arbitrary constants of integration which can be found from initial conditions. Using the value of the indicial root found earlier, $r = -1$, Eqs (1A,1B) become

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n-1}$$

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n-1} \right)$$

We start by finding the first solution $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = -\frac{3r}{(r+2)^2}$$

For $2 \leq n$ the recursive equation is

$$-a_{n-2}(n+r-2)(n-3+r) + a_n(n+r)(n+r-1) + 3a_{n-1}(n+r-1) + 3a_n(n+r) + a_n = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{n^2 a_{n-2} + 2nra_{n-2} + r^2 a_{n-2} - 5na_{n-2} - 3na_{n-1} - 5ra_{n-2} - 3ra_{n-1} + 6a_{n-2} + 3a_{n-1}}{n^2 + 2nr + r^2 + 2n + 2r + 1} \quad (4)$$

Which for the root $r = -1$ becomes

$$a_n = \frac{n^2 a_{n-2} + (-7a_{n-2} - 3a_{n-1})n + 12a_{n-2} + 6a_{n-1}}{n^2} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = -1$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{3r}{(r+2)^2}$	3

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{r(r^3 + 3r^2 + 9r + 5)}{(r+2)^2(r+3)^2}$$

Which for the root $r = -1$ becomes

$$a_2 = \frac{1}{2}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{3r}{(r+2)^2}$	3
a_2	$\frac{r(r^3+3r^2+9r+5)}{(r+2)^2(r+3)^2}$	$\frac{1}{2}$

For $n = 3$, using the above recursive equation gives

$$a_3 = -\frac{6r(r^4 + 6r^3 + 15r^2 + 16r + 5)}{(r+2)^2(r+3)^2(r+4)^2}$$

Which for the root $r = -1$ becomes

$$a_3 = -\frac{1}{6}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{3r}{(r+2)^2}$	3
a_2	$\frac{r(r^3+3r^2+9r+5)}{(r+2)^2(r+3)^2}$	$\frac{1}{2}$
a_3	$-\frac{6r(r^4+6r^3+15r^2+16r+5)}{(r+2)^2(r+3)^2(r+4)^2}$	$-\frac{1}{6}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{r(r^7 + 14r^6 + 102r^5 + 456r^4 + 1251r^3 + 1980r^2 + 1562r + 430)}{(r+2)^2(r+3)^2(r+4)^2(5+r)^2}$$

Which for the root $r = -1$ becomes

$$a_4 = \frac{1}{16}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{3r}{(r+2)^2}$	3
a_2	$\frac{r(r^3+3r^2+9r+5)}{(r+2)^2(r+3)^2}$	$\frac{1}{2}$
a_3	$-\frac{6r(r^4+6r^3+15r^2+16r+5)}{(r+2)^2(r+3)^2(r+4)^2}$	$-\frac{1}{6}$
a_4	$\frac{r(r^7+14r^6+102r^5+456r^4+1251r^3+1980r^2+1562r+430)}{(r+2)^2(r+3)^2(r+4)^2(5+r)^2}$	$\frac{1}{16}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{-9r^9 - 180r^8 - 1590r^7 - 8064r^6 - 25545r^5 - 51228r^4 - 62136r^3 - 39984r^2 - 9660r}{(r+2)^2(r+3)^2(r+4)^2(5+r)^2(r+6)^2}$$

Which for the root $r = -1$ becomes

$$a_5 = -\frac{43}{1200}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{3r}{(r+2)^2}$	3
a_2	$\frac{r(r^3+3r^2+9r+5)}{(r+2)^2(r+3)^2}$	$\frac{1}{2}$
a_3	$-\frac{6r(r^4+6r^3+15r^2+16r+5)}{(r+2)^2(r+3)^2(r+4)^2}$	$-\frac{1}{6}$
a_4	$\frac{r(r^7+14r^6+102r^5+456r^4+1251r^3+1980r^2+1562r+430)}{(r+2)^2(r+3)^2(r+4)^2(5+r)^2}$	$\frac{1}{16}$
a_5	$\frac{-9r^9-180r^8-1590r^7-8064r^6-25545r^5-51228r^4-62136r^3-39984r^2-9660r}{(r+2)^2(r+3)^2(r+4)^2(5+r)^2(r+6)^2}$	$-\frac{43}{1200}$

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{r(r^{11} + 33r^{10} + 527r^9 + 5313r^8 + 36825r^7 + 180423r^6 + 626549r^5 + 1520715r^4 + 2493694r^3 + 2582}{(r+2)^2(r+3)^2(r+4)^2(5+r)^2(r+6)^2(r+7)^2}$$

Which for the root $r = -1$ becomes

$$a_6 = \frac{161}{7200}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{3r}{(r+2)^2}$	3
a_2	$\frac{r(r^3+3r^2+9r+5)}{(r+2)^2(r+3)^2}$	$\frac{1}{2}$
a_3	$-\frac{6r(r^4+6r^3+15r^2+16r+5)}{(r+2)^2(r+3)^2(r+4)^2}$	$-\frac{1}{6}$
a_4	$\frac{r(r^7+14r^6+102r^5+456r^4+1251r^3+1980r^2+1562r+430)}{(r+2)^2(r+3)^2(r+4)^2(5+r)^2}$	$\frac{1}{16}$
a_5	$\frac{-9r^9-180r^8-1590r^7-8064r^6-25545r^5-51228r^4-62136r^3-39984r^2-9660r}{(r+2)^2(r+3)^2(r+4)^2(5+r)^2(r+6)^2}$	$-\frac{43}{1200}$
a_6	$\frac{r(r^{11}+33r^{10}+527r^9+5313r^8+36825r^7+180423r^6+626549r^5+1520715r^4+2493694r^3+2582664r^2+1473804r+330660)}{(r+2)^2(r+3)^2(r+4)^2(5+r)^2(r+6)^2(r+7)^2}$	$\frac{161}{7200}$

For $n = 7$, using the above recursive equation gives

$$a_7 = -\frac{12r(r^{12} + 42r^{11} + 805r^{10} + 9304r^9 + 72148r^8 + 394846r^7 + 1560105r^6 + 4468272r^5 + 9155771r^4 - (r+2)^2(r+3)^2(r+4)^2(5+r)^2(r+6)^2(r+7)^2)}{(r+2)^2(r+3)^2(r+4)^2(5+r)^2(r+6)^2(r+7)^2}$$

Which for the root $r = -1$ becomes

$$a_7 = -\frac{1837}{117600}$$

And the table now becomes

n	$a_{n,r}$
a_0	1
a_1	$-\frac{3r}{(r+2)^2}$
a_2	$\frac{r(r^3+3r^2+9r+5)}{(r+2)^2(r+3)^2}$
a_3	$-\frac{6r(r^4+6r^3+15r^2+16r+5)}{(r+2)^2(r+3)^2(r+4)^2}$
a_4	$\frac{r(r^7+14r^6+102r^5+456r^4+1251r^3+1980r^2+1562r+430)}{(r+2)^2(r+3)^2(r+4)^2(5+r)^2}$
a_5	$\frac{-9r^9-180r^8-1590r^7-8064r^6-25545r^5-51228r^4-62136r^3-39984r^2-9660r}{(r+2)^2(r+3)^2(r+4)^2(5+r)^2(r+6)^2}$
a_6	$\frac{r(r^{11}+33r^{10}+527r^9+5313r^8+36825r^7+180423r^6+626549r^5+1520715r^4+2493694r^3+2582664r^2+1473804r+330660)}{(r+2)^2(r+3)^2(r+4)^2(5+r)^2(r+6)^2(r+7)^2}$
a_7	$-\frac{12r(r^{12}+42r^{11}+805r^{10}+9304r^9+72148r^8+394846r^7+1560105r^6+4468272r^5+9155771r^4+12971420r^3+11876234r^2+6139136r+128)}{(r+2)^2(r+3)^2(r+4)^2(5+r)^2(r+6)^2(r+7)^2(r+8)^2}$

Using the above table, then the first solution $y_1(x)$ is

$$y_1(x) = \frac{1}{x}(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots)$$

$$= \frac{3x + 1 + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{16} - \frac{43x^5}{1200} + \frac{161x^6}{7200} - \frac{1837x^7}{117600} + O(x^8)}{x}$$

Now the second solution is found. The second solution is given by

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right)$$

Where b_n is found using

$$b_n = \frac{d}{dr} a_{n,r}$$

And the above is then evaluated at $r = -1$. The above table for $a_{n,r}$ is used for this purpose. Computing the derivatives gives the following table

n	$b_{n,r}$
b_0	1
b_1	$-\frac{3r}{(r+2)^2}$
b_2	$\frac{r(r^3+3r^2+9r+5)}{(r+2)^2(r+3)^2}$
b_3	$-\frac{6r(r^4+6r^3+15r^2+16r+5)}{(r+2)^2(r+3)^2(r+4)^2}$
b_4	$\frac{r(r^7+14r^6+102r^5+456r^4+1251r^3+1980r^2+1562r+430)}{(r+2)^2(r+3)^2(r+4)^2(5+r)^2}$
b_5	$\frac{-9r^9-180r^8-1590r^7-8064r^6-25545r^5-51228r^4-62136r^3-39984r^2-9660r}{(r+2)^2(r+3)^2(r+4)^2(5+r)^2(r+6)^2}$
b_6	$\frac{r(r^{11}+33r^{10}+527r^9+5313r^8+36825r^7+180423r^6+626549r^5+1520715r^4+2493694r^3+2582664r^2+1473804r+330660)}{(r+2)^2(r+3)^2(r+4)^2(5+r)^2(r+6)^2(r+7)^2}$
b_7	$-\frac{12r(r^{12}+42r^{11}+805r^{10}+9304r^9+72148r^8+394846r^7+1560105r^6+4468272r^5+9155771r^4+12971420r^3+11876234r^2+6139136r+128400)}{(r+2)^2(r+3)^2(r+4)^2(5+r)^2(r+6)^2(r+7)^2(r+8)^2}$

The above table gives all values of b_n needed. Hence the second solution is

$$y_2(x) = y_1(x) \ln(x) + b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 + b_7x^7 + b_8x^8 \dots$$

$$= \left(\frac{3x + 1 + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{16} - \frac{43x^5}{1200} + \frac{161x^6}{7200} - \frac{1837x^7}{117600} + O(x^8)}{x} \right) \ln(x)$$

$$+ \frac{-9x - \frac{7x^2}{2} + \frac{7x^3}{9} - \frac{25x^4}{96} + \frac{5141x^5}{36000} - \frac{2083x^6}{24000} + \frac{489941x^7}{8232000} + O(x^8)}{x}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 y_h(x) &= c_1 y_1(x) + c_2 y_2(x) \\
 &= \frac{c_1 \left(3x + 1 + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{16} - \frac{43x^5}{1200} + \frac{161x^6}{7200} - \frac{1837x^7}{117600} + O(x^8) \right)}{x} \\
 &\quad + c_2 \left(\frac{\left(3x + 1 + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{16} - \frac{43x^5}{1200} + \frac{161x^6}{7200} - \frac{1837x^7}{117600} + O(x^8) \right) \ln(x)}{x} \right. \\
 &\quad \left. + \frac{-9x - \frac{7x^2}{2} + \frac{7x^3}{9} - \frac{25x^4}{96} + \frac{5141x^5}{36000} - \frac{2083x^6}{24000} + \frac{489941x^7}{8232000} + O(x^8)}{x} \right)
 \end{aligned}$$

Hence the final solution is

$$\begin{aligned}
 y &= y_h \\
 &= \frac{c_1 \left(3x + 1 + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{16} - \frac{43x^5}{1200} + \frac{161x^6}{7200} - \frac{1837x^7}{117600} + O(x^8) \right)}{x} \\
 &\quad + c_2 \left(\frac{\left(3x + 1 + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{16} - \frac{43x^5}{1200} + \frac{161x^6}{7200} - \frac{1837x^7}{117600} + O(x^8) \right) \ln(x)}{x} \right. \\
 &\quad \left. + \frac{-9x - \frac{7x^2}{2} + \frac{7x^3}{9} - \frac{25x^4}{96} + \frac{5141x^5}{36000} - \frac{2083x^6}{24000} + \frac{489941x^7}{8232000} + O(x^8)}{x} \right)
 \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}
 y &= \frac{c_1 \left(3x + 1 + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{16} - \frac{43x^5}{1200} + \frac{161x^6}{7200} - \frac{1837x^7}{117600} + O(x^8) \right)}{x} \\
 &\quad + c_2 \left(\frac{\left(3x + 1 + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{16} - \frac{43x^5}{1200} + \frac{161x^6}{7200} - \frac{1837x^7}{117600} + O(x^8) \right) \ln(x)}{x} \right. \\
 &\quad \left. + \frac{-9x - \frac{7x^2}{2} + \frac{7x^3}{9} - \frac{25x^4}{96} + \frac{5141x^5}{36000} - \frac{2083x^6}{24000} + \frac{489941x^7}{8232000} + O(x^8)}{x} \right) \quad (1)
 \end{aligned}$$

Verification of solutions

$$y = \frac{c_1 \left(3x + 1 + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{16} - \frac{43x^5}{1200} + \frac{161x^6}{7200} - \frac{1837x^7}{117600} + O(x^8) \right)}{x} \\ + c_2 \left(\frac{\left(3x + 1 + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{16} - \frac{43x^5}{1200} + \frac{161x^6}{7200} - \frac{1837x^7}{117600} + O(x^8) \right) \ln(x)}{x} \right. \\ \left. + \frac{-9x - \frac{7x^2}{2} + \frac{7x^3}{9} - \frac{25x^4}{96} + \frac{5141x^5}{36000} - \frac{2083x^6}{24000} + \frac{489941x^7}{8232000} + O(x^8)}{x} \right)$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
  -> Bessel
  -> elliptic
  -> Legendre
  -> Whittaker
    -> hyper3: Equivalence to 1F1 under a power @ Moebius
  -> hypergeometric
    -> heuristic approach
    -> hyper3: Equivalence to 2F1, 1F1 or 0F1 under a power @ Moebius
  -> Mathieu
    -> Equivalence to the rational form of Mathieu ODE under a power @ Moebius
trying a solution in terms of MeijerG functions
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form  $r_0(x) * Y + r_1(x) * Y$  where  $Y = \exp(\int(r(x), dx)) * 2F1([a$ 
  trying a symmetry of the form [xi=0, eta=F(x)]
  trying differential order: 2; exact nonlinear
  trying symmetries linear in x and y(x)
  trying to convert to a linear ODE with constant coefficients
  trying 2nd order, integrating factor of the form  $\mu(x,y)$ 
  -> Trying a solution in terms of special functions:
    -> Bessel
    -> elliptic
    -> Legendre
    -> Whittaker
      -> hyper3: Equivalence to 1F1 under a power @ Moebius
    -> hypergeometric
      -> heuristic approach
      -> hyper3: Equivalence to 2F1, 1F1 or 0F1 under a power @ Moebius
    -> Mathieu
      -> Equivalence to the rational form of Mathieu ODE under a power @ Moebius
  trying 2nd order exact linear
  trying symmetries linear in x and y(x)
  trying to convert to a linear ODE with constant coefficients
  trying to convert to an ODE of Bessel type
trying to convert to an ODE of Bessel type
-> trying reduction of order to Bessel
```


✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 81

Order:=8;

```
dsolve((1-x^2)*x^2*diff(y(x),x$2)+3*(x+x^2)*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$y(x)$

$$= \frac{(c_2 \ln(x) + c_1) \left(1 + 3x + \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{16}x^4 - \frac{43}{1200}x^5 + \frac{161}{7200}x^6 - \frac{1837}{117600}x^7 + O(x^8)\right) + ((-9)x - \frac{7}{2}x^2 + \frac{7}{9}x^3)}{x}$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 84

```
AsymptoticDSolveValue[(1-x^2)*y'[x]+3*(x+x^2)*y'[x]+y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_2 \left(\frac{53x^7}{630} + \frac{5x^6}{24} + \frac{2x^5}{15} - \frac{x^4}{4} - \frac{2x^3}{3} + x \right) + c_1 \left(-\frac{19x^7}{420} - \frac{x^6}{144} + \frac{3x^5}{20} + \frac{5x^4}{24} - \frac{x^2}{2} + 1 \right)$$

19.4 problem 3(a)

19.4.1 Maple step by step solution 1571

Internal problem ID [6056]

Internal file name [OUTPUT/5304_Sunday_June_05_2022_03_33_32_PM_62803089/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 166

Problem number: 3(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Repeated root"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$x^2y'' + 3xy' + (1+x)y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2y'' + 3xy' + (1+x)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{3}{x}$$
$$q(x) = \frac{1+x}{x^2}$$

Table 232: Table $p(x), q(x)$ singularities.

$p(x) = \frac{3}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = \frac{1+x}{x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2 y'' + 3xy' + (1 + x)y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned}
 & x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\
 & + 3x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (1+x) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0
 \end{aligned} \tag{1}$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} 3x^{n+r} a_n (n+r) \right) \\ & + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \left(\sum_{n=0}^{\infty} x^{1+n+r} a_n \right) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} x^{1+n+r} a_n = \sum_{n=1}^{\infty} a_{n-1} x^{n+r}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} 3x^{n+r} a_n (n+r) \right) \\ & + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \left(\sum_{n=1}^{\infty} a_{n-1} x^{n+r} \right) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r} a_n (n+r) (n+r-1) + 3x^{n+r} a_n (n+r) + a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$x^r a_0 r (-1+r) + 3x^r a_0 r + a_0 x^r = 0$$

Or

$$(x^r r (-1+r) + 3x^r r + x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(r+1)^2 x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$(r+1)^2 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= -1 \\ r_2 &= -1 \end{aligned}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(r + 1)^2 x^r = 0$$

Solving for r gives the roots of the indicial equation as Since the root of the indicial equation is repeated, then we can construct two linearly independent solutions. The first solution has the form

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+r} \quad (1A)$$

Now the second solution y_2 is found using

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right) \quad (1B)$$

Then the general solution will be

$$y = c_1 y_1(x) + c_2 y_2(x)$$

In Eq (1B) the sum starts from 1 and not zero. In Eq (1A), a_0 is never zero, and is arbitrary and is typically taken as $a_0 = 1$, and $\{c_1, c_2\}$ are two arbitrary constants of integration which can be found from initial conditions. Using the value of the indicial root found earlier, $r = -1$, Eqs (1A,1B) become

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^{n-1} \\ y_2(x) &= y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n-1} \right) \end{aligned}$$

We start by finding the first solution $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) + 3a_n(n+r) + a_n + a_{n-1} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{a_{n-1}}{n^2 + 2nr + r^2 + 2n + 2r + 1} \quad (4)$$

Which for the root $r = -1$ becomes

$$a_n = -\frac{a_{n-1}}{n^2} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = -1$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = -\frac{1}{(r+2)^2}$$

Which for the root $r = -1$ becomes

$$a_1 = -1$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{(r+2)^2}$	-1

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{(r+2)^2(r+3)^2}$$

Which for the root $r = -1$ becomes

$$a_2 = \frac{1}{4}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{(r+2)^2}$	-1
a_2	$\frac{1}{(r+2)^2(r+3)^2}$	$\frac{1}{4}$

For $n = 3$, using the above recursive equation gives

$$a_3 = -\frac{1}{(r+2)^2 (r+3)^2 (r+4)^2}$$

Which for the root $r = -1$ becomes

$$a_3 = -\frac{1}{36}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{(r+2)^2}$	-1
a_2	$\frac{1}{(r+2)^2(r+3)^2}$	$\frac{1}{4}$
a_3	$-\frac{1}{(r+2)^2(r+3)^2(r+4)^2}$	$-\frac{1}{36}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{(r+2)^2 (r+3)^2 (r+4)^2 (5+r)^2}$$

Which for the root $r = -1$ becomes

$$a_4 = \frac{1}{576}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{(r+2)^2}$	-1
a_2	$\frac{1}{(r+2)^2(r+3)^2}$	$\frac{1}{4}$
a_3	$-\frac{1}{(r+2)^2(r+3)^2(r+4)^2}$	$-\frac{1}{36}$
a_4	$\frac{1}{(r+2)^2(r+3)^2(r+4)^2(5+r)^2}$	$\frac{1}{576}$

For $n = 5$, using the above recursive equation gives

$$a_5 = -\frac{1}{(r+2)^2 (r+3)^2 (r+4)^2 (5+r)^2 (r+6)^2}$$

Which for the root $r = -1$ becomes

$$a_5 = -\frac{1}{14400}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{(r+2)^2}$	-1
a_2	$\frac{1}{(r+2)^2(r+3)^2}$	$\frac{1}{4}$
a_3	$-\frac{1}{(r+2)^2(r+3)^2(r+4)^2}$	$-\frac{1}{36}$
a_4	$\frac{1}{(r+2)^2(r+3)^2(r+4)^2(5+r)^2}$	$\frac{1}{576}$
a_5	$-\frac{1}{(r+2)^2(r+3)^2(r+4)^2(5+r)^2(r+6)^2}$	$-\frac{1}{14400}$

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{1}{(r+2)^2(r+3)^2(r+4)^2(5+r)^2(r+6)^2(r+7)^2}$$

Which for the root $r = -1$ becomes

$$a_6 = \frac{1}{518400}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{(r+2)^2}$	-1
a_2	$\frac{1}{(r+2)^2(r+3)^2}$	$\frac{1}{4}$
a_3	$-\frac{1}{(r+2)^2(r+3)^2(r+4)^2}$	$-\frac{1}{36}$
a_4	$\frac{1}{(r+2)^2(r+3)^2(r+4)^2(5+r)^2}$	$\frac{1}{576}$
a_5	$-\frac{1}{(r+2)^2(r+3)^2(r+4)^2(5+r)^2(r+6)^2}$	$-\frac{1}{14400}$
a_6	$\frac{1}{(r+2)^2(r+3)^2(r+4)^2(5+r)^2(r+6)^2(r+7)^2}$	$\frac{1}{518400}$

For $n = 7$, using the above recursive equation gives

$$a_7 = -\frac{1}{(r+2)^2 (r+3)^2 (r+4)^2 (5+r)^2 (r+6)^2 (r+7)^2 (r+8)^2}$$

Which for the root $r = -1$ becomes

$$a_7 = -\frac{1}{25401600}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{(r+2)^2}$	-1
a_2	$\frac{1}{(r+2)^2(r+3)^2}$	$\frac{1}{4}$
a_3	$-\frac{1}{(r+2)^2(r+3)^2(r+4)^2}$	$-\frac{1}{36}$
a_4	$\frac{1}{(r+2)^2(r+3)^2(r+4)^2(5+r)^2}$	$\frac{1}{576}$
a_5	$-\frac{1}{(r+2)^2(r+3)^2(r+4)^2(5+r)^2(r+6)^2}$	$-\frac{1}{14400}$
a_6	$\frac{1}{(r+2)^2(r+3)^2(r+4)^2(5+r)^2(r+6)^2(r+7)^2}$	$\frac{1}{518400}$
a_7	$-\frac{1}{(r+2)^2(r+3)^2(r+4)^2(5+r)^2(r+6)^2(r+7)^2(r+8)^2}$	$-\frac{1}{25401600}$

Using the above table, then the first solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= \frac{1}{x} (a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots) \\ &= \frac{1 - x + \frac{x^2}{4} - \frac{x^3}{36} + \frac{x^4}{576} - \frac{x^5}{14400} + \frac{x^6}{518400} - \frac{x^7}{25401600} + O(x^8)}{x} \end{aligned}$$

Now the second solution is found. The second solution is given by

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right)$$

Where b_n is found using

$$b_n = \frac{d}{dr} a_{n,r}$$

And the above is then evaluated at $r = -1$. The above table for $a_{n,r}$ is used for this purpose. Computing the derivatives gives the following table

n	$b_{n,r}$	a_n	$b_{n,r} = \frac{d}{dr} a_{n,r}$
b_0	1	1	N/A since b_n starts from 1
b_1	$-\frac{1}{(r+2)^2}$	-1	$\frac{2}{(r+2)^3}$
b_2	$\frac{1}{(r+2)^2(r+3)^2}$	$\frac{1}{4}$	$\frac{-4r-10}{(r+2)^3(r+3)^3}$
b_3	$-\frac{1}{(r+2)^2(r+3)^2(r+4)^2}$	$-\frac{1}{36}$	$\frac{6r^2+36r+52}{(r+2)^3(r+3)^3(r+4)^3}$
b_4	$\frac{1}{(r+2)^2(r+3)^2(r+4)^2(5+r)^2}$	$\frac{1}{576}$	$\frac{-8r^3-84r^2-284r-308}{(r+2)^3(r+3)^3(r+4)^3(5+r)^3}$
b_5	$-\frac{1}{(r+2)^2(r+3)^2(r+4)^2(5+r)^2(r+6)^2}$	$-\frac{1}{14400}$	$\frac{10r^4+160r^3+930r^2+2320r+2088}{(r+2)^3(r+3)^3(r+4)^3(5+r)^3(r+6)^3}$
b_6	$\frac{1}{(r+2)^2(r+3)^2(r+4)^2(5+r)^2(r+6)^2(r+7)^2}$	$\frac{1}{518400}$	$\frac{-12r^5-270r^4-2360r^3-9990r^2-20416r-16056}{(r+2)^3(r+3)^3(r+4)^3(5+r)^3(r+6)^3(r+7)^3}$
b_7	$-\frac{1}{(r+2)^2(r+3)^2(r+4)^2(5+r)^2(r+6)^2(r+7)^2(r+8)^2}$	$-\frac{1}{25401600}$	$\frac{14r^6+420r^5+5110r^4+32200r^3+110544r^2+195440r+138528}{(r+2)^3(r+3)^3(r+4)^3(5+r)^3(r+6)^3(r+7)^3(r+8)^3}$

The above table gives all values of b_n needed. Hence the second solution is

$$\begin{aligned}
y_2(x) &= y_1(x) \ln(x) + b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 + b_7x^7 + b_8x^8 \dots \\
&= \left(1 - x + \frac{x^2}{4} - \frac{x^3}{36} + \frac{x^4}{576} - \frac{x^5}{14400} + \frac{x^6}{518400} - \frac{x^7}{25401600} + O(x^8) \right) \ln(x) \\
&\quad + \frac{2x - \frac{3x^2}{4} + \frac{11x^3}{108} - \frac{25x^4}{3456} + \frac{137x^5}{432000} - \frac{49x^6}{5184000} + \frac{121x^7}{592704000} + O(x^8)}{x}
\end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
y_h(x) &= c_1y_1(x) + c_2y_2(x) \\
&= \frac{c_1 \left(1 - x + \frac{x^2}{4} - \frac{x^3}{36} + \frac{x^4}{576} - \frac{x^5}{14400} + \frac{x^6}{518400} - \frac{x^7}{25401600} + O(x^8) \right)}{x} \\
&\quad + c_2 \left(\frac{\left(1 - x + \frac{x^2}{4} - \frac{x^3}{36} + \frac{x^4}{576} - \frac{x^5}{14400} + \frac{x^6}{518400} - \frac{x^7}{25401600} + O(x^8) \right) \ln(x)}{x} \right. \\
&\quad \left. + \frac{2x - \frac{3x^2}{4} + \frac{11x^3}{108} - \frac{25x^4}{3456} + \frac{137x^5}{432000} - \frac{49x^6}{5184000} + \frac{121x^7}{592704000} + O(x^8)}{x} \right)
\end{aligned}$$

Hence the final solution is

$$\begin{aligned}
 y &= y_h \\
 &= \frac{c_1 \left(1 - x + \frac{x^2}{4} - \frac{x^3}{36} + \frac{x^4}{576} - \frac{x^5}{14400} + \frac{x^6}{518400} - \frac{x^7}{25401600} + O(x^8) \right)}{x} \\
 &\quad + c_2 \left(\frac{\left(1 - x + \frac{x^2}{4} - \frac{x^3}{36} + \frac{x^4}{576} - \frac{x^5}{14400} + \frac{x^6}{518400} - \frac{x^7}{25401600} + O(x^8) \right) \ln(x)}{x} \right. \\
 &\quad \left. + \frac{2x - \frac{3x^2}{4} + \frac{11x^3}{108} - \frac{25x^4}{3456} + \frac{137x^5}{432000} - \frac{49x^6}{5184000} + \frac{121x^7}{592704000} + O(x^8)}{x} \right)
 \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}
 y &= \frac{c_1 \left(1 - x + \frac{x^2}{4} - \frac{x^3}{36} + \frac{x^4}{576} - \frac{x^5}{14400} + \frac{x^6}{518400} - \frac{x^7}{25401600} + O(x^8) \right)}{x} \\
 &\quad + c_2 \left(\frac{\left(1 - x + \frac{x^2}{4} - \frac{x^3}{36} + \frac{x^4}{576} - \frac{x^5}{14400} + \frac{x^6}{518400} - \frac{x^7}{25401600} + O(x^8) \right) \ln(x)}{x} \right. \\
 &\quad \left. + \frac{2x - \frac{3x^2}{4} + \frac{11x^3}{108} - \frac{25x^4}{3456} + \frac{137x^5}{432000} - \frac{49x^6}{5184000} + \frac{121x^7}{592704000} + O(x^8)}{x} \right) \quad (1)
 \end{aligned}$$

Verification of solutions

$$\begin{aligned}
 y &= \frac{c_1 \left(1 - x + \frac{x^2}{4} - \frac{x^3}{36} + \frac{x^4}{576} - \frac{x^5}{14400} + \frac{x^6}{518400} - \frac{x^7}{25401600} + O(x^8) \right)}{x} \\
 &\quad + c_2 \left(\frac{\left(1 - x + \frac{x^2}{4} - \frac{x^3}{36} + \frac{x^4}{576} - \frac{x^5}{14400} + \frac{x^6}{518400} - \frac{x^7}{25401600} + O(x^8) \right) \ln(x)}{x} \right. \\
 &\quad \left. + \frac{2x - \frac{3x^2}{4} + \frac{11x^3}{108} - \frac{25x^4}{3456} + \frac{137x^5}{432000} - \frac{49x^6}{5184000} + \frac{121x^7}{592704000} + O(x^8)}{x} \right)
 \end{aligned}$$

Verified OK.

19.4.1 Maple step by step solution

Let's solve

$$x^2 y'' + 3xy' + (1+x)y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{3y'}{x} - \frac{(1+x)y}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{3y'}{x} + \frac{(1+x)y}{x^2} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$\left[P_2(x) = \frac{3}{x}, P_3(x) = \frac{1+x}{x^2} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 3$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 1$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$x^2 y'' + 3xy' + (1+x)y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 0..1$

$$x^m \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

- Shift index using $k- > k - m$

$$x^m \cdot y = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

- Convert $x \cdot y'$ to series expansion

$$x \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r}$$

- Convert $x^2 \cdot y''$ to series expansion

$$x^2 \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(1+r)^2 x^r + \left(\sum_{k=1}^{\infty} (a_k(k+r+1)^2 + a_{k-1}) x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$(1+r)^2 = 0$$
- Values of r that satisfy the indicial equation

$$r = -1$$
- Each term in the series must be 0, giving the recursion relation

$$a_k(k+r+1)^2 + a_{k-1} = 0$$
- Shift index using $k- > k+1$

$$a_{k+1}(k+2+r)^2 + a_k = 0$$
- Recursion relation that defines series solution to ODE

$$a_{k+1} = -\frac{a_k}{(k+2+r)^2}$$
- Recursion relation for $r = -1$

$$a_{k+1} = -\frac{a_k}{(k+1)^2}$$
- Solution for $r = -1$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k-1}, a_{k+1} = -\frac{a_k}{(k+1)^2} \right]$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
    <- Bessel successful
<- special function solution successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 81

```
Order:=8;
dsolve(x^2*diff(y(x),x$2)+3*x*diff(y(x),x)+(1+x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{(c_2 \ln(x) + c_1) \left(1 - x + \frac{1}{4}x^2 - \frac{1}{36}x^3 + \frac{1}{576}x^4 - \frac{1}{14400}x^5 + \frac{1}{518400}x^6 - \frac{1}{25401600}x^7 + O(x^8)\right) + \left(2x - \frac{3}{4}x^2 + \dots\right)}{x}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 164

```
AsymptoticDSolveValue[x^2*y''[x]+3*x*y'[x]+(1+x)*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow \frac{c_1 \left(-\frac{x^7}{25401600} + \frac{x^6}{518400} - \frac{x^5}{14400} + \frac{x^4}{576} - \frac{x^3}{36} + \frac{x^2}{4} - x + 1\right)}{x} + c_2 \left(\frac{\frac{121x^7}{592704000} - \frac{49x^6}{5184000} + \frac{137x^5}{432000} - \frac{25x^4}{3456} + \frac{11x^3}{108} - \frac{3x^2}{4} + 2x}{x} + \frac{\left(-\frac{x^7}{25401600} + \frac{x^6}{518400} - \frac{x^5}{14400} + \frac{x^4}{576} - \frac{x^3}{36} + \frac{x^2}{4} - x + 1\right) \log(x)}{x} \right)$$

19.5 problem 3(b)

19.5.1 Maple step by step solution 1587

Internal problem ID [6057]

Internal file name [OUTPUT/5305_Sunday_June_05_2022_03_33_35_PM_55245292/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 166

Problem number: 3(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference is integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$x^2y'' + 2x^2y' - 2y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2y'' + 2x^2y' - 2y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = 2$$

$$q(x) = -\frac{2}{x^2}$$

Table 234: Table $p(x), q(x)$ singularities.

$p(x) = 2$	
singularity	type

$q(x) = -\frac{2}{x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2 y'' + 2x^2 y' - 2y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) + 2x^2 \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) - 2 \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r)(n+r-1) \right) + \left(\sum_{n=0}^{\infty} 2x^{1+n+r} a_n (n+r) \right) + \sum_{n=0}^{\infty} (-2a_n x^{n+r}) = 0 \quad (2A)$$

The next step is to make all powers of x be $n + r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} 2x^{1+n+r} a_n(n+r) = \sum_{n=1}^{\infty} 2a_{n-1}(n+r-1)x^{n+r}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n + r$.

$$\left(\sum_{n=0}^{\infty} x^{n+r} a_n(n+r)(n+r-1) \right) + \left(\sum_{n=1}^{\infty} 2a_{n-1}(n+r-1)x^{n+r} \right) + \sum_{n=0}^{\infty} (-2a_n x^{n+r}) = 0 \quad (2B)$$

The indicial equation is obtained from $n = 0$. From Eq (2B) this gives

$$x^{n+r} a_n(n+r)(n+r-1) - 2a_n x^{n+r} = 0$$

When $n = 0$ the above becomes

$$x^r a_0 r(-1+r) - 2a_0 x^r = 0$$

Or

$$(x^r r(-1+r) - 2x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(r^2 - r - 2) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$r^2 - r - 2 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = 2$$

$$r_2 = -1$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(r^2 - r - 2) x^r = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = 3$ is an integer, then we can construct two linearly independent solutions

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = C y_1(x) \ln(x) + x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = x^2 \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = C y_1(x) \ln(x) + \frac{\sum_{n=0}^{\infty} b_n x^n}{x}$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+2}$$

$$y_2(x) = C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n-1} \right)$$

Where C above can be zero. We start by finding y_1 . Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) + 2a_{n-1}(n+r-1) - 2a_n = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{2a_{n-1}(n+r-1)}{n^2 + 2nr + r^2 - n - r - 2} \quad (4)$$

Which for the root $r = 2$ becomes

$$a_n = -\frac{2a_{n-1}(1+n)}{n(n+3)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 2$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = -\frac{2r}{r^2 + r - 2}$$

Which for the root $r = 2$ becomes

$$a_1 = -1$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{2r}{r^2+r-2}$	-1

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{4 + 4r}{r^3 + 4r^2 + r - 6}$$

Which for the root $r = 2$ becomes

$$a_2 = \frac{3}{5}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{2r}{r^2+r-2}$	-1
a_2	$\frac{4+4r}{r^3+4r^2+r-6}$	$\frac{3}{5}$

For $n = 3$, using the above recursive equation gives

$$a_3 = -\frac{8}{r^3 + 6r^2 + 5r - 12}$$

Which for the root $r = 2$ becomes

$$a_3 = -\frac{4}{15}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{2r}{r^2+r-2}$	-1
a_2	$\frac{4+4r}{r^3+4r^2+r-6}$	$\frac{3}{5}$
a_3	$-\frac{8}{r^3+6r^2+5r-12}$	$-\frac{4}{15}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{16}{r^4 + 10r^3 + 27r^2 + 2r - 40}$$

Which for the root $r = 2$ becomes

$$a_4 = \frac{2}{21}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{2r}{r^2+r-2}$	-1
a_2	$\frac{4+4r}{r^3+4r^2+r-6}$	$\frac{3}{5}$
a_3	$-\frac{8}{r^3+6r^2+5r-12}$	$-\frac{4}{15}$
a_4	$\frac{16}{r^4+10r^3+27r^2+2r-40}$	$\frac{2}{21}$

For $n = 5$, using the above recursive equation gives

$$a_5 = -\frac{32}{(r^2 + 9r + 18)(r^3 + 6r^2 + 3r - 10)}$$

Which for the root $r = 2$ becomes

$$a_5 = -\frac{1}{35}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{2r}{r^2+r-2}$	-1
a_2	$\frac{4+4r}{r^3+4r^2+r-6}$	$\frac{3}{5}$
a_3	$-\frac{8}{r^3+6r^2+5r-12}$	$-\frac{4}{15}$
a_4	$\frac{16}{r^4+10r^3+27r^2+2r-40}$	$\frac{2}{21}$
a_5	$-\frac{32}{(r^2+9r+18)(r^3+6r^2+3r-10)}$	$-\frac{1}{35}$

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{64}{(r^2 + 11r + 28)(r^2 + r - 2)(r^2 + 9r + 18)}$$

Which for the root $r = 2$ becomes

$$a_6 = \frac{1}{135}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{2r}{r^2+r-2}$	-1
a_2	$\frac{4+4r}{r^3+4r^2+r-6}$	$\frac{3}{5}$
a_3	$-\frac{8}{r^3+6r^2+5r-12}$	$-\frac{4}{15}$
a_4	$\frac{16}{r^4+10r^3+27r^2+2r-40}$	$\frac{2}{21}$
a_5	$-\frac{32}{(r^2+9r+18)(r^3+6r^2+3r-10)}$	$-\frac{1}{35}$
a_6	$\frac{64}{(r^2+11r+28)(r^2+r-2)(r^2+9r+18)}$	$\frac{1}{135}$

For $n = 7$, using the above recursive equation gives

$$a_7 = -\frac{128}{(r+8)(5+r)(r+3)(2+r)(-1+r)(r+7)(r+4)}$$

Which for the root $r = 2$ becomes

$$a_7 = -\frac{8}{4725}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{2r}{r^2+r-2}$	-1
a_2	$\frac{4+4r}{r^3+4r^2+r-6}$	$\frac{3}{5}$
a_3	$-\frac{8}{r^3+6r^2+5r-12}$	$-\frac{4}{15}$
a_4	$\frac{16}{r^4+10r^3+27r^2+2r-40}$	$\frac{2}{21}$
a_5	$-\frac{32}{(r^2+9r+18)(r^3+6r^2+3r-10)}$	$-\frac{1}{35}$
a_6	$\frac{64}{(r^2+11r+28)(r^2+r-2)(r^2+9r+18)}$	$\frac{1}{135}$
a_7	$-\frac{128}{(r+8)(5+r)(r+3)(2+r)(-1+r)(r+7)(r+4)}$	$-\frac{8}{4725}$

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned}
 y_1(x) &= x^2(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots) \\
 &= x^2\left(1 - x + \frac{3x^2}{5} - \frac{4x^3}{15} + \frac{2x^4}{21} - \frac{x^5}{35} + \frac{x^6}{135} - \frac{8x^7}{4725} + O(x^8)\right)
 \end{aligned}$$

Now the second solution $y_2(x)$ is found. Let

$$r_1 - r_2 = N$$

Where N is positive integer which is the difference between the two roots. r_1 is taken as the larger root. Hence for this problem we have $N = 3$. Now we need to determine if C is zero or not. This is done by finding $\lim_{r \rightarrow r_2} a_3(r)$. If this limit exists, then $C = 0$, else we need to keep the log term and $C \neq 0$. The above table shows that

$$\begin{aligned}
 a_N &= a_3 \\
 &= -\frac{8}{r^3 + 6r^2 + 5r - 12}
 \end{aligned}$$

Therefore

$$\begin{aligned}
 \lim_{r \rightarrow r_2} -\frac{8}{r^3 + 6r^2 + 5r - 12} &= \lim_{r \rightarrow -1} -\frac{8}{r^3 + 6r^2 + 5r - 12} \\
 &= \frac{2}{3}
 \end{aligned}$$

The limit is $\frac{2}{3}$. Since the limit exists then the log term is not needed and we can set $C = 0$. Therefore the second solution has the form

$$\begin{aligned} y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+r} \\ &= \sum_{n=0}^{\infty} b_n x^{n-1} \end{aligned}$$

Eq (3) derived above is used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. For $1 \leq n$ the recursive equation is

$$b_n(n+r)(n+r-1) + 2b_{n-1}(n+r-1) - 2b_n = 0 \quad (4)$$

Which for the root $r = -1$ becomes

$$b_n(n-1)(n-2) + 2b_{n-1}(n-2) - 2b_n = 0 \quad (4A)$$

Solving for b_n from the recursive equation (4) gives

$$b_n = -\frac{2b_{n-1}(n+r-1)}{n^2 + 2nr + r^2 - n - r - 2} \quad (5)$$

Which for the root $r = -1$ becomes

$$b_n = -\frac{2b_{n-1}(n-2)}{n^2 - 3n} \quad (6)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = -1$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1

For $n = 1$, using the above recursive equation gives

$$b_1 = -\frac{2r}{r^2 + r - 2}$$

Which for the root $r = -1$ becomes

$$b_1 = -1$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{2r}{r^2+r-2}$	-1

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{4 + 4r}{(r^2 + r - 2)(r + 3)}$$

Which for the root $r = -1$ becomes

$$b_2 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{2r}{r^2+r-2}$	-1
b_2	$\frac{4+4r}{r^3+4r^2+r-6}$	0

For $n = 3$, using the above recursive equation gives

$$b_3 = -\frac{8}{(r + 4)(r + 3)(-1 + r)}$$

Which for the root $r = -1$ becomes

$$b_3 = \frac{2}{3}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{2r}{r^2+r-2}$	-1
b_2	$\frac{4+4r}{r^3+4r^2+r-6}$	0
b_3	$-\frac{8}{(r+4)(r+3)(-1+r)}$	$\frac{2}{3}$

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{16}{(r+4)(-1+r)(r^2+7r+10)}$$

Which for the root $r = -1$ becomes

$$b_4 = -\frac{2}{3}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{2r}{r^2+r-2}$	-1
b_2	$\frac{4+4r}{r^3+4r^2+r-6}$	0
b_3	$-\frac{8}{(r+4)(r+3)(-1+r)}$	$\frac{2}{3}$
b_4	$\frac{16}{(5+r)(2+r)(r+4)(-1+r)}$	$-\frac{2}{3}$

For $n = 5$, using the above recursive equation gives

$$b_5 = -\frac{32}{(-1+r)(r^2+7r+10)(r^2+9r+18)}$$

Which for the root $r = -1$ becomes

$$b_5 = \frac{2}{5}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{2r}{r^2+r-2}$	-1
b_2	$\frac{4+4r}{r^3+4r^2+r-6}$	0
b_3	$-\frac{8}{(r+4)(r+3)(-1+r)}$	$\frac{2}{3}$
b_4	$\frac{16}{(5+r)(2+r)(r+4)(-1+r)}$	$-\frac{2}{3}$
b_5	$-\frac{32}{(6+r)(r+3)(5+r)(-1+r)(2+r)}$	$\frac{2}{5}$

For $n = 6$, using the above recursive equation gives

$$b_6 = \frac{64}{(r^2 + 9r + 18)(2 + r)(-1 + r)(r^2 + 11r + 28)}$$

Which for the root $r = -1$ becomes

$$b_6 = -\frac{8}{45}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{2r}{r^2+r-2}$	-1
b_2	$\frac{4+4r}{r^3+4r^2+r-6}$	0
b_3	$-\frac{8}{(r+4)(r+3)(-1+r)}$	$\frac{2}{3}$
b_4	$\frac{16}{(5+r)(2+r)(r+4)(-1+r)}$	$-\frac{2}{3}$
b_5	$-\frac{32}{(6+r)(r+3)(5+r)(-1+r)(2+r)}$	$\frac{2}{5}$
b_6	$\frac{64}{(r+7)(r+4)(2+r)(-1+r)(6+r)(r+3)}$	$-\frac{8}{45}$

For $n = 7$, using the above recursive equation gives

$$b_7 = -\frac{128}{(r^2 + 11r + 28)(-1 + r)(2 + r)(r + 3)(r^2 + 13r + 40)}$$

Which for the root $r = -1$ becomes

$$b_7 = \frac{4}{63}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{2r}{r^2+r-2}$	-1
b_2	$\frac{4+4r}{r^3+4r^2+r-6}$	0
b_3	$-\frac{8}{(r+4)(r+3)(-1+r)}$	$\frac{2}{3}$
b_4	$\frac{16}{(5+r)(2+r)(r+4)(-1+r)}$	$-\frac{2}{3}$
b_5	$-\frac{32}{(6+r)(r+3)(5+r)(-1+r)(2+r)}$	$\frac{2}{5}$
b_6	$\frac{64}{(r+7)(r+4)(2+r)(-1+r)(6+r)(r+3)}$	$-\frac{8}{45}$
b_7	$-\frac{128}{(r+8)(5+r)(r+3)(2+r)(-1+r)(r+7)(r+4)}$	$\frac{4}{63}$

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned}
 y_2(x) &= x^2(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 + b_7x^7 + b_8x^8 \dots) \\
 &= \frac{1 - x + \frac{2x^3}{3} - \frac{2x^4}{3} + \frac{2x^5}{5} - \frac{8x^6}{45} + \frac{4x^7}{63} + O(x^8)}{x}
 \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 y_h(x) &= c_1y_1(x) + c_2y_2(x) \\
 &= c_1x^2 \left(1 - x + \frac{3x^2}{5} - \frac{4x^3}{15} + \frac{2x^4}{21} - \frac{x^5}{35} + \frac{x^6}{135} - \frac{8x^7}{4725} + O(x^8) \right) \\
 &\quad + \frac{c_2 \left(1 - x + \frac{2x^3}{3} - \frac{2x^4}{3} + \frac{2x^5}{5} - \frac{8x^6}{45} + \frac{4x^7}{63} + O(x^8) \right)}{x}
 \end{aligned}$$

Hence the final solution is

$$\begin{aligned}
 y &= y_h \\
 &= c_1x^2 \left(1 - x + \frac{3x^2}{5} - \frac{4x^3}{15} + \frac{2x^4}{21} - \frac{x^5}{35} + \frac{x^6}{135} - \frac{8x^7}{4725} + O(x^8) \right) \\
 &\quad + \frac{c_2 \left(1 - x + \frac{2x^3}{3} - \frac{2x^4}{3} + \frac{2x^5}{5} - \frac{8x^6}{45} + \frac{4x^7}{63} + O(x^8) \right)}{x}
 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 x^2 \left(1 - x + \frac{3x^2}{5} - \frac{4x^3}{15} + \frac{2x^4}{21} - \frac{x^5}{35} + \frac{x^6}{135} - \frac{8x^7}{4725} + O(x^8) \right) + \frac{c_2 \left(1 - x + \frac{2x^3}{3} - \frac{2x^4}{3} + \frac{2x^5}{5} - \frac{8x^6}{45} + \frac{4x^7}{63} + O(x^8) \right)}{x} \quad (1)$$

Verification of solutions

$$y = c_1 x^2 \left(1 - x + \frac{3x^2}{5} - \frac{4x^3}{15} + \frac{2x^4}{21} - \frac{x^5}{35} + \frac{x^6}{135} - \frac{8x^7}{4725} + O(x^8) \right) + \frac{c_2 \left(1 - x + \frac{2x^3}{3} - \frac{2x^4}{3} + \frac{2x^5}{5} - \frac{8x^6}{45} + \frac{4x^7}{63} + O(x^8) \right)}{x}$$

Verified OK.

19.5.1 Maple step by step solution

Let's solve

$$x^2 y'' + 2x^2 y' - 2y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -2y' + \frac{2y}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + 2y' - \frac{2y}{x^2} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$[P_2(x) = 2, P_3(x) = -\frac{2}{x^2}]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 0$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = -2$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$x^2 y'' + 2x^2 y' - 2y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^2 \cdot y'$ to series expansion

$$x^2 \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r+1}$$

- Shift index using $k \rightarrow k-1$

$$x^2 \cdot y' = \sum_{k=1}^{\infty} a_{k-1} (k-1+r) x^{k+r}$$

- Convert $x^2 \cdot y''$ to series expansion

$$x^2 \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k-1+r) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(1+r)(-2+r)x^r + \left(\sum_{k=1}^{\infty} (a_k(k+r+1)(k+r-2) + 2a_{k-1}(k-1+r)) x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$(1+r)(-2+r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \{-1, 2\}$$

- Each term in the series must be 0, giving the recursion relation

$$a_k(k+r+1)(k+r-2) + 2a_{k-1}(k-1+r) = 0$$

- Shift index using $k \rightarrow k+1$

$$a_{k+1}(k+2+r)(k-1+r) + 2a_k(k+r) = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+1} = -\frac{2a_k(k+r)}{(k+2+r)(k-1+r)}$$

- Recursion relation for $r = -1$; series terminates at $k = 1$

$$a_{k+1} = -\frac{2a_k(k-1)}{(k+1)(k-2)}$$

- Apply recursion relation for $k = 0$

$$a_1 = -a_0$$

- Terminating series solution of the ODE for $r = -1$. Use reduction of order to find the second

$$y = a_0 \cdot (1 - x)$$

- Recursion relation for $r = 2$

$$a_{k+1} = -\frac{2a_k(k+2)}{(k+4)(k+1)}$$

- Solution for $r = 2$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+2}, a_{k+1} = -\frac{2a_k(k+2)}{(k+4)(k+1)} \right]$$

- Combine solutions and rename parameters

$$\left[y = a_0 \cdot (1 - x) + \left(\sum_{k=0}^{\infty} b_k x^{k+2} \right), b_{k+1} = -\frac{2b_k(k+2)}{(k+4)(k+1)} \right]$$

Maple trace **Kovacic algorithm successful**

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Reducible group (found an exponential solution)
    Reducible group (found another exponential solution)
<- Kovacics algorithm successful`

```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 53

Order:=8;

```
dsolve(x^2*diff(y(x),x$2)+2*x^2*diff(y(x),x)-2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^2 \left(1 - x + \frac{3}{5} x^2 - \frac{4}{15} x^3 + \frac{2}{21} x^4 - \frac{1}{35} x^5 + \frac{1}{135} x^6 - \frac{8}{4725} x^7 + O(x^8) \right) \\ + \frac{c_2 (12 - 12x + 8x^3 - 8x^4 + \frac{24}{5} x^5 - \frac{32}{15} x^6 + \frac{16}{21} x^7 + O(x^8))}{x}$$

✓ Solution by Mathematica

Time used: 0.048 (sec). Leaf size: 87

```
AsymptoticDSolveValue[x^2*y''[x]+2*x^2*y'[x]-2*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(-\frac{8x^5}{45} + \frac{2x^4}{5} - \frac{2x^3}{3} + \frac{2x^2}{3} + \frac{1}{x} - 1 \right) + c_2 \left(\frac{x^8}{135} - \frac{x^7}{35} + \frac{2x^6}{21} - \frac{4x^5}{15} + \frac{3x^4}{5} - x^3 + x^2 \right)$$

19.6 problem 3(c)

19.6.1 Maple step by step solution 1602

Internal problem ID [6058]

Internal file name [OUTPUT/5306_Sunday_June_05_2022_03_33_37_PM_1513034/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 166

Problem number: 3(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference is integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$x^2y'' + 5xy' + (-x^3 + 3)y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2y'' + 5xy' + (-x^3 + 3)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{5}{x}$$
$$q(x) = -\frac{x^3 - 3}{x^2}$$

Table 236: Table $p(x), q(x)$ singularities.

$p(x) = \frac{5}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = -\frac{x^3-3}{x^2}$	
singularity	type
$x = 0$	“regular”
$x = \infty$	“regular”
$x = -\infty$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0, \infty, -\infty]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2 y'' + 5xy' + (-x^3 + 3)y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) + 5x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (-x^3 + 3) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} 5x^{n+r} a_n (n+r) \right) \\ & + \sum_{n=0}^{\infty} (-x^{n+r+3} a_n) + \left(\sum_{n=0}^{\infty} 3a_n x^{n+r} \right) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} (-x^{n+r+3} a_n) = \sum_{n=3}^{\infty} (-a_{n-3} x^{n+r})$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} 5x^{n+r} a_n (n+r) \right) \\ & + \sum_{n=3}^{\infty} (-a_{n-3} x^{n+r}) + \left(\sum_{n=0}^{\infty} 3a_n x^{n+r} \right) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r} a_n (n+r) (n+r-1) + 5x^{n+r} a_n (n+r) + 3a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$x^r a_0 r(-1+r) + 5x^r a_0 r + 3a_0 x^r = 0$$

Or

$$(x^r r(-1+r) + 5x^r r + 3x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(r^2 + 4r + 3) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$r^2 + 4r + 3 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= -1 \\ r_2 &= -3 \end{aligned}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(r^2 + 4r + 3) x^r = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = 2$ is an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= C y_1(x) \ln(x) + x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \frac{\sum_{n=0}^{\infty} a_n x^n}{x} \\ y_2(x) &= C y_1(x) \ln(x) + \frac{\sum_{n=0}^{\infty} b_n x^n}{x^3} \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^{n-1} \\ y_2(x) &= C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n-3} \right) \end{aligned}$$

Where C above can be zero. We start by finding y_1 . Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

Substituting $n = 2$ in Eq. (2B) gives

$$a_2 = 0$$

For $3 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) + 5a_n(n+r) - a_{n-3} + 3a_n = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-3}}{n^2 + 2nr + r^2 + 4n + 4r + 3} \quad (4)$$

Which for the root $r = -1$ becomes

$$a_n = \frac{a_{n-3}}{n(n+2)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = -1$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{1}{r^2 + 10r + 24}$$

Which for the root $r = -1$ becomes

$$a_3 = \frac{1}{15}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	$\frac{1}{r^2+10r+24}$	$\frac{1}{15}$

For $n = 4$, using the above recursive equation gives

$$a_4 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	$\frac{1}{r^2+10r+24}$	$\frac{1}{15}$
a_4	0	0

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	$\frac{1}{r^2+10r+24}$	$\frac{1}{15}$
a_4	0	0
a_5	0	0

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{1}{(r+6)(r+4)(r+9)(r+7)}$$

Which for the root $r = -1$ becomes

$$a_6 = \frac{1}{720}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	$\frac{1}{r^2+10r+24}$	$\frac{1}{15}$
a_4	0	0
a_5	0	0
a_6	$\frac{1}{(r+6)(r+4)(r+9)(r+7)}$	$\frac{1}{720}$

For $n = 7$, using the above recursive equation gives

$$a_7 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	$\frac{1}{r^2+10r+24}$	$\frac{1}{15}$
a_4	0	0
a_5	0	0
a_6	$\frac{1}{(r+6)(r+4)(r+9)(r+7)}$	$\frac{1}{720}$
a_7	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned}
 y_1(x) &= \frac{1}{x} (a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots) \\
 &= \frac{1 + \frac{x^3}{15} + \frac{x^6}{720} + O(x^8)}{x}
 \end{aligned}$$

Now the second solution $y_2(x)$ is found. Let

$$r_1 - r_2 = N$$

Where N is positive integer which is the difference between the two roots. r_1 is taken as the larger root. Hence for this problem we have $N = 2$. Now we need to determine if C is zero or not. This is done by finding $\lim_{r \rightarrow r_2} a_2(r)$. If this limit exists, then $C = 0$, else we need to keep the log term and $C \neq 0$. The above table shows that

$$\begin{aligned} a_N &= a_2 \\ &= 0 \end{aligned}$$

Therefore

$$\begin{aligned} \lim_{r \rightarrow r_2} 0 &= \lim_{r \rightarrow -3} 0 \\ &= 0 \end{aligned}$$

The limit is 0. Since the limit exists then the log term is not needed and we can set $C = 0$. Therefore the second solution has the form

$$\begin{aligned} y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+r} \\ &= \sum_{n=0}^{\infty} b_n x^{n-3} \end{aligned}$$

Eq (3) derived above is used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq(3) gives

$$b_1 = 0$$

Substituting $n = 2$ in Eq(3) gives

$$b_2 = 0$$

For $3 \leq n$ the recursive equation is

$$b_n(n+r)(n+r-1) + 5b_n(n+r) - b_{n-3} + 3b_n = 0 \quad (4)$$

Which for for the root $r = -3$ becomes

$$b_n(n-3)(n-4) + 5b_n(n-3) - b_{n-3} + 3b_n = 0 \quad (4A)$$

Solving for b_n from the recursive equation (4) gives

$$b_n = \frac{b_{n-3}}{n^2 + 2nr + r^2 + 4n + 4r + 3} \quad (5)$$

Which for the root $r = -3$ becomes

$$b_n = \frac{b_{n-3}}{n^2 - 2n} \quad (6)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = -3$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	0	0

For $n = 3$, using the above recursive equation gives

$$b_3 = \frac{1}{r^2 + 10r + 24}$$

Which for the root $r = -3$ becomes

$$b_3 = \frac{1}{3}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	0	0
b_3	$\frac{1}{r^2+10r+24}$	$\frac{1}{3}$

For $n = 4$, using the above recursive equation gives

$$b_4 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	0	0
b_3	$\frac{1}{r^2+10r+24}$	$\frac{1}{3}$
b_4	0	0

For $n = 5$, using the above recursive equation gives

$$b_5 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	0	0
b_3	$\frac{1}{r^2+10r+24}$	$\frac{1}{3}$
b_4	0	0
b_5	0	0

For $n = 6$, using the above recursive equation gives

$$b_6 = \frac{1}{(r^2 + 10r + 24)(r^2 + 16r + 63)}$$

Which for the root $r = -3$ becomes

$$b_6 = \frac{1}{72}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	0	0
b_3	$\frac{1}{r^2+10r+24}$	$\frac{1}{3}$
b_4	0	0
b_5	0	0
b_6	$\frac{1}{(r+6)(r+4)(r+9)(r+7)}$	$\frac{1}{72}$

For $n = 7$, using the above recursive equation gives

$$b_7 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	0	0
b_3	$\frac{1}{r^2+10r+24}$	$\frac{1}{3}$
b_4	0	0
b_5	0	0
b_6	$\frac{1}{(r+6)(r+4)(r+9)(r+7)}$	$\frac{1}{72}$
b_7	0	0

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned} y_2(x) &= \frac{1}{x} (b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 + b_7x^7 + b_8x^8 \dots) \\ &= \frac{1 + \frac{x^3}{3} + \frac{x^6}{72} + O(x^8)}{x^3} \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= \frac{c_1 \left(1 + \frac{x^3}{15} + \frac{x^6}{720} + O(x^8) \right)}{x} + \frac{c_2 \left(1 + \frac{x^3}{3} + \frac{x^6}{72} + O(x^8) \right)}{x^3} \end{aligned}$$

Hence the final solution is

$$\begin{aligned} y &= y_h \\ &= \frac{c_1 \left(1 + \frac{x^3}{15} + \frac{x^6}{720} + O(x^8) \right)}{x} + \frac{c_2 \left(1 + \frac{x^3}{3} + \frac{x^6}{72} + O(x^8) \right)}{x^3} \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1 \left(1 + \frac{x^3}{15} + \frac{x^6}{720} + O(x^8) \right)}{x} + \frac{c_2 \left(1 + \frac{x^3}{3} + \frac{x^6}{72} + O(x^8) \right)}{x^3} \quad (1)$$

Verification of solutions

$$y = \frac{c_1 \left(1 + \frac{x^3}{15} + \frac{x^6}{720} + O(x^8)\right)}{x} + \frac{c_2 \left(1 + \frac{x^3}{3} + \frac{x^6}{72} + O(x^8)\right)}{x^3}$$

Verified OK.

19.6.1 Maple step by step solution

Let's solve

$$x^2 y'' + 5xy' + (-x^3 + 3)y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{5y'}{x} + \frac{(x^3-3)y}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{5y'}{x} - \frac{(x^3-3)y}{x^2} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$\left[P_2(x) = \frac{5}{x}, P_3(x) = -\frac{x^3-3}{x^2} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 5$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 3$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$x^2 y'' + 5xy' + (-x^3 + 3)y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

□ Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 0..3$

$$x^m \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

- Shift index using $k \rightarrow k - m$

$$x^m \cdot y = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

- Convert $x \cdot y'$ to series expansion

$$x \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r}$$

- Convert $x^2 \cdot y''$ to series expansion

$$x^2 \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(3+r)(1+r)x^r + a_1(4+r)(2+r)x^{1+r} + a_2(5+r)(3+r)x^{2+r} + \left(\sum_{k=3}^{\infty} (a_k(k+r+3)(k+r+2)) \right) x^{k+r} = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$(3+r)(1+r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \{-3, -1\}$$

- The coefficients of each power of x must be 0

$$[a_1(4+r)(2+r) = 0, a_2(5+r)(3+r) = 0]$$

- Solve for the dependent coefficient(s)

$$\{a_1 = 0, a_2 = 0\}$$

- Each term in the series must be 0, giving the recursion relation

$$a_k(k+r+3)(k+r+1) - a_{k-3} = 0$$

- Shift index using $k \rightarrow k + 3$

$$a_{k+3}(k+6+r)(k+4+r) - a_k = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+3} = \frac{a_k}{(k+6+r)(k+4+r)}$$

- Recursion relation for $r = -3$

$$a_{k+3} = \frac{a_k}{(k+3)(k+1)}$$

- Solution for $r = -3$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k-3}, a_{k+3} = \frac{a_k}{(k+3)(k+1)}, a_1 = 0, a_2 = 0 \right]$$

- Recursion relation for $r = -1$

$$a_{k+3} = \frac{a_k}{(k+5)(k+3)}$$

- Solution for $r = -1$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k-1}, a_{k+3} = \frac{a_k}{(k+5)(k+3)}, a_1 = 0, a_2 = 0 \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^{k-3} \right) + \left(\sum_{k=0}^{\infty} b_k x^{k-1} \right), a_{k+3} = \frac{a_k}{(k+3)(k+1)}, a_1 = 0, a_2 = 0, b_{k+3} = \frac{b_k}{(k+5)(k+3)}, b_1 = 0, b_2 = 0 \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
        <- Bessel successful
<- special function solution successful`

```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 35

Order:=8;

```
dsolve(x^2*diff(y(x),x$2)+5*x*diff(y(x),x)+(3-x^3)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{c_1 \left(1 + \frac{1}{15}x^3 + \frac{1}{720}x^6 + O(x^8)\right)}{x} + \frac{c_2 \left(-2 - \frac{2}{3}x^3 - \frac{1}{36}x^6 + O(x^8)\right)}{x^3}$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 40

```
AsymptoticDSolveValue[x^2*y''[x]+5*x*y'[x]+(3-3*x^3)*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(\frac{x^3}{8} + \frac{1}{x^3} + 1 \right) + c_2 \left(\frac{x^5}{80} + \frac{x^2}{5} + \frac{1}{x} \right)$$

19.7 problem 3(d)

19.7.1 Maple step by step solution 1619

Internal problem ID [6059]

Internal file name [OUTPUT/5307_Sunday_June_05_2022_03_33_40_PM_38231263/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 166

Problem number: 3(d).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference is integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$x^2 y'' - 2x(1+x)y' + 2(1+x)y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2 y'' + (-2x^2 - 2x)y' + (2 + 2x)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{2(1+x)}{x}$$
$$q(x) = \frac{2+2x}{x^2}$$

Table 238: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{2(1+x)}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = \frac{2+2x}{x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2 y'' + (-2x^2 - 2x) y' + (2 + 2x) y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned} & x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\ & + (-2x^2 - 2x) \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (2+2x) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \end{aligned} \tag{1}$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-2x^{1+n+r} a_n (n+r)) \\ & + \sum_{n=0}^{\infty} (-2x^{n+r} a_n (n+r)) + \left(\sum_{n=0}^{\infty} 2a_n x^{n+r} \right) + \left(\sum_{n=0}^{\infty} 2x^{1+n+r} a_n \right) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=0}^{\infty} (-2x^{1+n+r} a_n (n+r)) &= \sum_{n=1}^{\infty} (-2a_{n-1} (n+r-1) x^{n+r}) \\ \sum_{n=0}^{\infty} 2x^{1+n+r} a_n &= \sum_{n=1}^{\infty} 2a_{n-1} x^{n+r} \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=1}^{\infty} (-2a_{n-1} (n+r-1) x^{n+r}) \\ & + \sum_{n=0}^{\infty} (-2x^{n+r} a_n (n+r)) + \left(\sum_{n=0}^{\infty} 2a_n x^{n+r} \right) + \left(\sum_{n=1}^{\infty} 2a_{n-1} x^{n+r} \right) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r} a_n (n+r) (n+r-1) - 2x^{n+r} a_n (n+r) + 2a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$x^r a_0 r (-1+r) - 2x^r a_0 r + 2a_0 x^r = 0$$

Or

$$(x^r r (-1+r) - 2x^r r + 2x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(r^2 - 3r + 2) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$r^2 - 3r + 2 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = 2$$

$$r_2 = 1$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(r^2 - 3r + 2) x^r = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = 1$ is an integer, then we can construct two linearly independent solutions

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = C y_1(x) \ln(x) + x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = x^2 \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = C y_1(x) \ln(x) + x \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+2}$$

$$y_2(x) = C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{1+n} \right)$$

Where C above can be zero. We start by finding y_1 . Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) - 2a_{n-1}(n+r-1) - 2a_n(n+r) + 2a_n + 2a_{n-1} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{2a_{n-1}}{n+r-1} \quad (4)$$

Which for the root $r = 2$ becomes

$$a_n = \frac{2a_{n-1}}{1+n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 2$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = \frac{2}{r}$$

Which for the root $r = 2$ becomes

$$a_1 = 1$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{2}{r}$	1

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{4}{r(1+r)}$$

Which for the root $r = 2$ becomes

$$a_2 = \frac{2}{3}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{2}{r}$	1
a_2	$\frac{4}{r(1+r)}$	$\frac{2}{3}$

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{8}{r(1+r)(2+r)}$$

Which for the root $r = 2$ becomes

$$a_3 = \frac{1}{3}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{2}{r}$	1
a_2	$\frac{4}{r(1+r)}$	$\frac{2}{3}$
a_3	$\frac{8}{r(1+r)(2+r)}$	$\frac{1}{3}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{16}{r(1+r)(3+r)(2+r)}$$

Which for the root $r = 2$ becomes

$$a_4 = \frac{2}{15}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{2}{r}$	1
a_2	$\frac{4}{r(1+r)}$	$\frac{2}{3}$
a_3	$\frac{8}{r(1+r)(2+r)}$	$\frac{1}{3}$
a_4	$\frac{16}{r(1+r)(3+r)(2+r)}$	$\frac{2}{15}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{32}{r(1+r)(2+r)(3+r)(4+r)}$$

Which for the root $r = 2$ becomes

$$a_5 = \frac{2}{45}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{2}{r}$	1
a_2	$\frac{4}{r(1+r)}$	$\frac{2}{3}$
a_3	$\frac{8}{r(1+r)(2+r)}$	$\frac{1}{3}$
a_4	$\frac{16}{r(1+r)(3+r)(2+r)}$	$\frac{2}{15}$
a_5	$\frac{32}{r(1+r)(2+r)(3+r)(4+r)}$	$\frac{2}{45}$

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{64}{r(1+r)(3+r)(2+r)(4+r)(5+r)}$$

Which for the root $r = 2$ becomes

$$a_6 = \frac{4}{315}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{2}{r}$	1
a_2	$\frac{4}{r(1+r)}$	$\frac{2}{3}$
a_3	$\frac{8}{r(1+r)(2+r)}$	$\frac{1}{3}$
a_4	$\frac{16}{r(1+r)(3+r)(2+r)}$	$\frac{2}{15}$
a_5	$\frac{32}{r(1+r)(2+r)(3+r)(4+r)}$	$\frac{2}{45}$
a_6	$\frac{64}{r(1+r)(3+r)(2+r)(4+r)(5+r)}$	$\frac{4}{315}$

For $n = 7$, using the above recursive equation gives

$$a_7 = \frac{128}{r(1+r)(2+r)(3+r)(4+r)(5+r)(6+r)}$$

Which for the root $r = 2$ becomes

$$a_7 = \frac{1}{315}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{2}{r}$	1
a_2	$\frac{4}{r(1+r)}$	$\frac{2}{3}$
a_3	$\frac{8}{r(1+r)(2+r)}$	$\frac{1}{3}$
a_4	$\frac{16}{r(1+r)(3+r)(2+r)}$	$\frac{2}{15}$
a_5	$\frac{32}{r(1+r)(2+r)(3+r)(4+r)}$	$\frac{2}{45}$
a_6	$\frac{64}{r(1+r)(3+r)(2+r)(4+r)(5+r)}$	$\frac{4}{315}$
a_7	$\frac{128}{r(1+r)(2+r)(3+r)(4+r)(5+r)(6+r)}$	$\frac{1}{315}$

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned}
 y_1(x) &= x^2(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots) \\
 &= x^2\left(1 + x + \frac{2x^2}{3} + \frac{x^3}{3} + \frac{2x^4}{15} + \frac{2x^5}{45} + \frac{4x^6}{315} + \frac{x^7}{315} + O(x^8)\right)
 \end{aligned}$$

Now the second solution $y_2(x)$ is found. Let

$$r_1 - r_2 = N$$

Where N is positive integer which is the difference between the two roots. r_1 is taken as the larger root. Hence for this problem we have $N = 1$. Now we need to determine if C is zero or not. This is done by finding $\lim_{r \rightarrow r_2} a_1(r)$. If this limit exists, then $C = 0$, else we need to keep the log term and $C \neq 0$. The above table shows that

$$\begin{aligned}
 a_N &= a_1 \\
 &= \frac{2}{r}
 \end{aligned}$$

Therefore

$$\begin{aligned}
 \lim_{r \rightarrow r_2} \frac{2}{r} &= \lim_{r \rightarrow 1} \frac{2}{r} \\
 &= 2
 \end{aligned}$$

The limit is 2. Since the limit exists then the log term is not needed and we can set $C = 0$. Therefore the second solution has the form

$$\begin{aligned} y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+r} \\ &= \sum_{n=0}^{\infty} b_n x^{1+n} \end{aligned}$$

Eq (3) derived above is used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. For $1 \leq n$ the recursive equation is

$$b_n(n+r)(n+r-1) - 2b_{n-1}(n+r-1) - 2b_n(n+r) + 2b_n + 2b_{n-1} = 0 \quad (4)$$

Which for the root $r = 1$ becomes

$$b_n(1+n)n - 2b_{n-1}n - 2b_n(1+n) + 2b_n + 2b_{n-1} = 0 \quad (4A)$$

Solving for b_n from the recursive equation (4) gives

$$b_n = \frac{2b_{n-1}}{n+r-1} \quad (5)$$

Which for the root $r = 1$ becomes

$$b_n = \frac{2b_{n-1}}{n} \quad (6)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = 1$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1

For $n = 1$, using the above recursive equation gives

$$b_1 = \frac{2}{r}$$

Which for the root $r = 1$ becomes

$$b_1 = 2$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{2}{r}$	2

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{4}{r(1+r)}$$

Which for the root $r = 1$ becomes

$$b_2 = 2$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{2}{r}$	2
b_2	$\frac{4}{r(1+r)}$	2

For $n = 3$, using the above recursive equation gives

$$b_3 = \frac{8}{r(1+r)(2+r)}$$

Which for the root $r = 1$ becomes

$$b_3 = \frac{4}{3}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{2}{r}$	2
b_2	$\frac{4}{r(1+r)}$	2
b_3	$\frac{8}{r(1+r)(2+r)}$	$\frac{4}{3}$

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{16}{r(1+r)(3+r)(2+r)}$$

Which for the root $r = 1$ becomes

$$b_4 = \frac{2}{3}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{2}{r}$	2
b_2	$\frac{4}{r(1+r)}$	2
b_3	$\frac{8}{r(1+r)(2+r)}$	$\frac{4}{3}$
b_4	$\frac{16}{r(1+r)(3+r)(2+r)}$	$\frac{2}{3}$

For $n = 5$, using the above recursive equation gives

$$b_5 = \frac{32}{r(1+r)(2+r)(3+r)(4+r)}$$

Which for the root $r = 1$ becomes

$$b_5 = \frac{4}{15}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{2}{r}$	2
b_2	$\frac{4}{r(1+r)}$	2
b_3	$\frac{8}{r(1+r)(2+r)}$	$\frac{4}{3}$
b_4	$\frac{16}{r(1+r)(3+r)(2+r)}$	$\frac{2}{3}$
b_5	$\frac{32}{r(1+r)(2+r)(3+r)(4+r)}$	$\frac{4}{15}$

For $n = 6$, using the above recursive equation gives

$$b_6 = \frac{64}{r(1+r)(3+r)(2+r)(4+r)(5+r)}$$

Which for the root $r = 1$ becomes

$$b_6 = \frac{4}{45}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{2}{r}$	2
b_2	$\frac{4}{r(1+r)}$	2
b_3	$\frac{8}{r(1+r)(2+r)}$	$\frac{4}{3}$
b_4	$\frac{16}{r(1+r)(3+r)(2+r)}$	$\frac{2}{3}$
b_5	$\frac{32}{r(1+r)(2+r)(3+r)(4+r)}$	$\frac{4}{15}$
b_6	$\frac{64}{r(1+r)(3+r)(2+r)(4+r)(5+r)}$	$\frac{4}{45}$

For $n = 7$, using the above recursive equation gives

$$b_7 = \frac{128}{r(1+r)(2+r)(3+r)(4+r)(5+r)(6+r)}$$

Which for the root $r = 1$ becomes

$$b_7 = \frac{8}{315}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{2}{r}$	2
b_2	$\frac{4}{r(1+r)}$	2
b_3	$\frac{8}{r(1+r)(2+r)}$	$\frac{4}{3}$
b_4	$\frac{16}{r(1+r)(3+r)(2+r)}$	$\frac{2}{3}$
b_5	$\frac{32}{r(1+r)(2+r)(3+r)(4+r)}$	$\frac{4}{15}$
b_6	$\frac{64}{r(1+r)(3+r)(2+r)(4+r)(5+r)}$	$\frac{4}{45}$
b_7	$\frac{128}{r(1+r)(2+r)(3+r)(4+r)(5+r)(6+r)}$	$\frac{8}{315}$

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned} y_2(x) &= x^2(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 + b_7x^7 + b_8x^8 \dots) \\ &= x \left(1 + 2x + 2x^2 + \frac{4x^3}{3} + \frac{2x^4}{3} + \frac{4x^5}{15} + \frac{4x^6}{45} + \frac{8x^7}{315} + O(x^8) \right) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}y_h(x) &= c_1 y_1(x) + c_2 y_2(x) \\&= c_1 x^2 \left(1 + x + \frac{2x^2}{3} + \frac{x^3}{3} + \frac{2x^4}{15} + \frac{2x^5}{45} + \frac{4x^6}{315} + \frac{x^7}{315} + O(x^8) \right) \\&\quad + c_2 x \left(1 + 2x + 2x^2 + \frac{4x^3}{3} + \frac{2x^4}{3} + \frac{4x^5}{15} + \frac{4x^6}{45} + \frac{8x^7}{315} + O(x^8) \right)\end{aligned}$$

Hence the final solution is

$$\begin{aligned}y &= y_h \\&= c_1 x^2 \left(1 + x + \frac{2x^2}{3} + \frac{x^3}{3} + \frac{2x^4}{15} + \frac{2x^5}{45} + \frac{4x^6}{315} + \frac{x^7}{315} + O(x^8) \right) \\&\quad + c_2 x \left(1 + 2x + 2x^2 + \frac{4x^3}{3} + \frac{2x^4}{3} + \frac{4x^5}{15} + \frac{4x^6}{45} + \frac{8x^7}{315} + O(x^8) \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}y &= c_1 x^2 \left(1 + x + \frac{2x^2}{3} + \frac{x^3}{3} + \frac{2x^4}{15} + \frac{2x^5}{45} + \frac{4x^6}{315} + \frac{x^7}{315} + O(x^8) \right) \\&\quad + c_2 x \left(1 + 2x + 2x^2 + \frac{4x^3}{3} + \frac{2x^4}{3} + \frac{4x^5}{15} + \frac{4x^6}{45} + \frac{8x^7}{315} + O(x^8) \right)\end{aligned} \tag{1}$$

Verification of solutions

$$\begin{aligned}y &= c_1 x^2 \left(1 + x + \frac{2x^2}{3} + \frac{x^3}{3} + \frac{2x^4}{15} + \frac{2x^5}{45} + \frac{4x^6}{315} + \frac{x^7}{315} + O(x^8) \right) \\&\quad + c_2 x \left(1 + 2x + 2x^2 + \frac{4x^3}{3} + \frac{2x^4}{3} + \frac{4x^5}{15} + \frac{4x^6}{45} + \frac{8x^7}{315} + O(x^8) \right)\end{aligned}$$

Verified OK.

19.7.1 Maple step by step solution

Let's solve

$$x^2 y'' + (-2x^2 - 2x) y' + (2 + 2x) y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{2(1+x)y}{x^2} + \frac{2(1+x)y'}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - \frac{2(1+x)y'}{x} + \frac{2(1+x)y}{x^2} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$\left[P_2(x) = -\frac{2(1+x)}{x}, P_3(x) = \frac{2(1+x)}{x^2} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = -2$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 2$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$x^2 y'' - 2x(1+x) y' + (2+2x) y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 0..1$

$$x^m \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

- Shift index using $k \rightarrow k - m$

$$x^m \cdot y = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

- Convert $x^m \cdot y'$ to series expansion for $m = 1..2$

$$x^m \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1+m}$$

- Shift index using $k \rightarrow k + 1 - m$

$$x^m \cdot y' = \sum_{k=-1+m}^{\infty} a_{k+1-m} (k+1-m+r) x^{k+r}$$

- Convert $x^2 \cdot y''$ to series expansion

$$x^2 \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(-1+r)(-2+r)x^r + \left(\sum_{k=1}^{\infty} (a_k(k+r-1)(k+r-2) - 2a_{k-1}(k+r-2)) x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$(-1+r)(-2+r) = 0$$
- Values of r that satisfy the indicial equation

$$r \in \{1, 2\}$$
- Each term in the series must be 0, giving the recursion relation

$$(k+r-2)(a_k(k+r-1) - 2a_{k-1}) = 0$$
- Shift index using $k \rightarrow k + 1$

$$(k+r-1)(a_{k+1}(k+r) - 2a_k) = 0$$
- Recursion relation that defines series solution to ODE

$$a_{k+1} = \frac{2a_k}{k+r}$$
- Recursion relation for $r = 1$

$$a_{k+1} = \frac{2a_k}{k+1}$$
- Solution for $r = 1$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+1}, a_{k+1} = \frac{2a_k}{k+1} \right]$$
- Recursion relation for $r = 2$

$$a_{k+1} = \frac{2a_k}{k+2}$$

- Solution for $r = 2$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+2}, a_{k+1} = \frac{2a_k}{k+2} \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^{k+1} \right) + \left(\sum_{k=0}^{\infty} b_k x^{k+2} \right), a_{k+1} = \frac{2a_k}{k+1}, b_{k+1} = \frac{2b_k}{k+2} \right]$$

Maple trace Kovacic algorithm successful

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Reducible group (found an exponential solution)
    Reducible group (found another exponential solution)
<- Kovacics algorithm successful`

```

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 53

```

Order:=8;
dsolve(x^2*diff(y(x),x$2)-2*x*(x+1)*diff(y(x),x)+2*(x+1)*y(x)=0,y(x),type='series',x=0);

```

$$y(x) = c_1 x^2 \left(1 + x + \frac{2}{3}x^2 + \frac{1}{3}x^3 + \frac{2}{15}x^4 + \frac{2}{45}x^5 + \frac{4}{315}x^6 + \frac{1}{315}x^7 + O(x^8) \right) \\ + c_2 x \left(1 + 2x + 2x^2 + \frac{4}{3}x^3 + \frac{2}{3}x^4 + \frac{4}{15}x^5 + \frac{4}{45}x^6 + \frac{8}{315}x^7 + O(x^8) \right)$$

✓ Solution by Mathematica

Time used: 0.086 (sec). Leaf size: 92

```
AsymptoticDSolveValue[x^2*y''[x]-2*x*(x+1)*y'[x]+2*(1+x)*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(\frac{4x^7}{45} + \frac{4x^6}{15} + \frac{2x^5}{3} + \frac{4x^4}{3} + 2x^3 + 2x^2 + x \right) \\ + c_2 \left(\frac{4x^8}{315} + \frac{2x^7}{45} + \frac{2x^6}{15} + \frac{x^5}{3} + \frac{2x^4}{3} + x^3 + x^2 \right)$$

19.8 problem 3(e)

19.8.1 Maple step by step solution 1636

Internal problem ID [6060]

Internal file name [OUTPUT/5308_Sunday_June_05_2022_03_33_42_PM_72107480/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 166

Problem number: 3(e).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference is integer"**

Maple gives the following as the ode type

[_Bessel]

$$x^2y'' + xy' + (x^2 - 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2y'' + xy' + (x^2 - 1)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = \frac{x^2 - 1}{x^2}$$

Table 240: Table $p(x), q(x)$ singularities.

$p(x) = \frac{1}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = \frac{x^2-1}{x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2 y'' + x y' + (x^2 - 1) y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned}
 & x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\
 & + x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (x^2 - 1) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0
 \end{aligned} \tag{1}$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) \right) \\ & + \left(\sum_{n=0}^{\infty} x^{n+r+2} a_n \right) + \sum_{n=0}^{\infty} (-a_n x^{n+r}) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} x^{n+r+2} a_n = \sum_{n=2}^{\infty} a_{n-2} x^{n+r}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) \right) \\ & + \left(\sum_{n=2}^{\infty} a_{n-2} x^{n+r} \right) + \sum_{n=0}^{\infty} (-a_n x^{n+r}) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r} a_n (n+r) (n+r-1) + x^{n+r} a_n (n+r) - a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$x^r a_0 r(-1+r) + x^r a_0 r - a_0 x^r = 0$$

Or

$$(x^r r(-1+r) + x^r r - x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(r^2 - 1) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$r^2 - 1 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 1 \\ r_2 &= -1 \end{aligned}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(r^2 - 1) x^r = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = 2$ is an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= C y_1(x) \ln(x) + x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= x \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= C y_1(x) \ln(x) + \frac{\sum_{n=0}^{\infty} b_n x^n}{x} \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^{n+1} \\ y_2(x) &= C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n-1} \right) \end{aligned}$$

Where C above can be zero. We start by finding y_1 . Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

For $2 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) + a_n(n+r) + a_{n-2} - a_n = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{a_{n-2}}{n^2 + 2nr + r^2 - 1} \quad (4)$$

Which for the root $r = 1$ becomes

$$a_n = -\frac{a_{n-2}}{n(n+2)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 1$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = -\frac{1}{r^2 + 4r + 3}$$

Which for the root $r = 1$ becomes

$$a_2 = -\frac{1}{8}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2+4r+3}$	$-\frac{1}{8}$

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2+4r+3}$	$-\frac{1}{8}$
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{(r+3)^2(1+r)(5+r)}$$

Which for the root $r = 1$ becomes

$$a_4 = \frac{1}{192}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2+4r+3}$	$-\frac{1}{8}$
a_3	0	0
a_4	$\frac{1}{(r+3)^2(1+r)(5+r)}$	$\frac{1}{192}$

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2+4r+3}$	$-\frac{1}{8}$
a_3	0	0
a_4	$\frac{1}{(r+3)^2(1+r)(5+r)}$	$\frac{1}{192}$
a_5	0	0

For $n = 6$, using the above recursive equation gives

$$a_6 = -\frac{1}{(r+3)^2(1+r)(5+r)^2(r+7)}$$

Which for the root $r = 1$ becomes

$$a_6 = -\frac{1}{9216}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2+4r+3}$	$-\frac{1}{8}$
a_3	0	0
a_4	$\frac{1}{(r+3)^2(1+r)(5+r)}$	$\frac{1}{192}$
a_5	0	0
a_6	$-\frac{1}{(r+3)^2(1+r)(5+r)^2(r+7)}$	$-\frac{1}{9216}$

For $n = 7$, using the above recursive equation gives

$$a_7 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2+4r+3}$	$-\frac{1}{8}$
a_3	0	0
a_4	$\frac{1}{(r+3)^2(1+r)(5+r)}$	$\frac{1}{192}$
a_5	0	0
a_6	$-\frac{1}{(r+3)^2(1+r)(5+r)^2(r+7)}$	$-\frac{1}{9216}$
a_7	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots) \\ &= x\left(1 - \frac{x^2}{8} + \frac{x^4}{192} - \frac{x^6}{9216} + O(x^8)\right) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Let

$$r_1 - r_2 = N$$

Where N is positive integer which is the difference between the two roots. r_1 is taken as the larger root. Hence for this problem we have $N = 2$. Now we need to determine if C is zero or not. This is done by finding $\lim_{r \rightarrow r_2} a_2(r)$. If this limit exists, then $C = 0$, else we need to keep the log term and $C \neq 0$. The above table shows that

$$\begin{aligned} a_N &= a_2 \\ &= -\frac{1}{r^2 + 4r + 3} \end{aligned}$$

Therefore

$$\begin{aligned} \lim_{r \rightarrow r_2} -\frac{1}{r^2 + 4r + 3} &= \lim_{r \rightarrow -1} -\frac{1}{r^2 + 4r + 3} \\ &= \text{undefined} \end{aligned}$$

Since the limit does not exist then the log term is needed. Therefore the second solution has the form

$$y_2(x) = Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right)$$

Therefore

$$\begin{aligned} \frac{d}{dx} y_2(x) &= Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2} (n+r_2)}{x} \right) \\ &= Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} x^{-1+n+r_2} b_n (n+r_2) \right) \\ \frac{d^2}{dx^2} y_2(x) &= Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} \\ &\quad + \sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2} (n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2} (n+r_2)}{x^2} \right) \\ &= Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} + \left(\sum_{n=0}^{\infty} x^{-2+n+r_2} b_n (n+r_2) (-1+n+r_2) \right) \end{aligned}$$

Substituting these back into the given ode $x^2y'' + xy' + (x^2 - 1)y = 0$ gives

$$\begin{aligned}
 & x^2 \left(Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} \right. \\
 & \left. + \sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2} (n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2} (n+r_2)}{x^2} \right) \right) \\
 & + x \left(Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2} (n+r_2)}{x} \right) \right) \\
 & + (x^2 - 1) \left(Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) \right) = 0
 \end{aligned}$$

Which can be written as

$$\begin{aligned}
 & \left((x^2 y_1''(x) + y_1'(x) x + (x^2 - 1) y_1(x)) \ln(x) + x^2 \left(\frac{2y_1'(x)}{x} - \frac{y_1(x)}{x^2} \right) + y_1(x) \right) C \\
 & + x^2 \left(\sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2} (n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2} (n+r_2)}{x^2} \right) \right) \\
 & + x \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2} (n+r_2)}{x} \right) + (x^2 - 1) \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) = 0
 \end{aligned} \tag{7}$$

But since $y_1(x)$ is a solution to the ode, then

$$x^2 y_1''(x) + y_1'(x) x + (x^2 - 1) y_1(x) = 0$$

Eq (7) simplifies to

$$\begin{aligned}
 & \left(x^2 \left(\frac{2y_1'(x)}{x} - \frac{y_1(x)}{x^2} \right) + y_1(x) \right) C \\
 & + x^2 \left(\sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2} (n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2} (n+r_2)}{x^2} \right) \right) \\
 & + x \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2} (n+r_2)}{x} \right) + (x^2 - 1) \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) = 0
 \end{aligned} \tag{8}$$

Substituting $y_1 = \sum_{n=0}^{\infty} a_n x^{n+r_1}$ into the above gives

$$2x \left(\sum_{n=0}^{\infty} x^{-1+n+r_1} a_n (n+r_1) \right) C + \left(\sum_{n=0}^{\infty} x^{-2+n+r_2} b_n (n+r_2) (-1+n+r_2) \right) x^2 \quad (9)$$

$$+ \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) x^2 + \left(\sum_{n=0}^{\infty} x^{-1+n+r_2} b_n (n+r_2) \right) x - \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) = 0$$

Since $r_1 = 1$ and $r_2 = -1$ then the above becomes

$$2x \left(\sum_{n=0}^{\infty} x^n a_n (n+1) \right) C + \left(\sum_{n=0}^{\infty} x^{-3+n} b_n (n-1) (n-2) \right) x^2 \quad (10)$$

$$+ \left(\sum_{n=0}^{\infty} b_n x^{n-1} \right) x^2 + \left(\sum_{n=0}^{\infty} x^{n-2} b_n (n-1) \right) x - \left(\sum_{n=0}^{\infty} b_n x^{n-1} \right) = 0$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} 2C x^{n+1} a_n (n+1) \right) + \left(\sum_{n=0}^{\infty} x^{n-1} b_n (n^2 - 3n + 2) \right) \quad (2A)$$

$$+ \left(\sum_{n=0}^{\infty} x^{n+1} b_n \right) + \left(\sum_{n=0}^{\infty} x^{n-1} b_n (n-1) \right) + \sum_{n=0}^{\infty} (-b_n x^{n-1}) = 0$$

The next step is to make all powers of x be $n-1$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n-1} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} 2C x^{n+1} a_n (n+1) = \sum_{n=2}^{\infty} 2C a_{n-2} (n-1) x^{n-1}$$

$$\sum_{n=0}^{\infty} x^{n+1} b_n = \sum_{n=2}^{\infty} b_{n-2} x^{n-1}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers

of x are the same and equal to $n - 1$.

$$\begin{aligned} & \left(\sum_{n=2}^{\infty} 2Ca_{n-2}(n-1)x^{n-1} \right) + \left(\sum_{n=0}^{\infty} x^{n-1}b_n(n^2 - 3n + 2) \right) \\ & + \left(\sum_{n=2}^{\infty} b_{n-2}x^{n-1} \right) + \left(\sum_{n=0}^{\infty} x^{n-1}b_n(n-1) \right) + \sum_{n=0}^{\infty} (-b_nx^{n-1}) = 0 \end{aligned} \quad (2B)$$

For $n = 0$ in Eq. (2B), we choose arbitrary value for b_0 as $b_0 = 1$. For $n = 1$, Eq (2B) gives

$$-b_1 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$-b_1 = 0$$

Solving the above for b_1 gives

$$b_1 = 0$$

For $n = N$, where $N = 2$ which is the difference between the two roots, we are free to choose $b_2 = 0$. Hence for $n = 2$, Eq (2B) gives

$$2C + 1 = 0$$

Which is solved for C . Solving for C gives

$$C = -\frac{1}{2}$$

For $n = 3$, Eq (2B) gives

$$4Ca_1 + b_1 + 3b_3 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$3b_3 = 0$$

Solving the above for b_3 gives

$$b_3 = 0$$

For $n = 4$, Eq (2B) gives

$$6Ca_2 + b_2 + 8b_4 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$8b_4 + \frac{3}{8} = 0$$

Solving the above for b_4 gives

$$b_4 = -\frac{3}{64}$$

For $n = 5$, Eq (2B) gives

$$8Ca_3 + b_3 + 15b_5 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$15b_5 = 0$$

Solving the above for b_5 gives

$$b_5 = 0$$

For $n = 6$, Eq (2B) gives

$$10Ca_4 + b_4 + 24b_6 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$24b_6 - \frac{7}{96} = 0$$

Solving the above for b_6 gives

$$b_6 = \frac{7}{2304}$$

For $n = 7$, Eq (2B) gives

$$12Ca_5 + b_5 + 35b_7 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$35b_7 = 0$$

Solving the above for b_7 gives

$$b_7 = 0$$

Now that we found all b_n and C , we can calculate the second solution from

$$y_2(x) = Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right)$$

Using the above value found for $C = -\frac{1}{2}$ and all b_n , then the second solution becomes

$$y_2(x) = -\frac{1}{2} \left(x \left(1 - \frac{x^2}{8} + \frac{x^4}{192} - \frac{x^6}{9216} + O(x^8) \right) \right) \ln(x) + \frac{1 - \frac{3x^4}{64} + \frac{7x^6}{2304} + O(x^8)}{x}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 y_h(x) &= c_1 y_1(x) + c_2 y_2(x) \\
 &= c_1 x \left(1 - \frac{x^2}{8} + \frac{x^4}{192} - \frac{x^6}{9216} + O(x^8) \right) \\
 &\quad + c_2 \left(-\frac{1}{2} \left(x \left(1 - \frac{x^2}{8} + \frac{x^4}{192} - \frac{x^6}{9216} + O(x^8) \right) \right) \ln(x) + \frac{1 - \frac{3x^4}{64} + \frac{7x^6}{2304} + O(x^8)}{x} \right)
 \end{aligned}$$

Hence the final solution is

$$\begin{aligned}
 y &= y_h \\
 &= c_1 x \left(1 - \frac{x^2}{8} + \frac{x^4}{192} - \frac{x^6}{9216} + O(x^8) \right) \\
 &\quad + c_2 \left(-\frac{x \left(1 - \frac{x^2}{8} + \frac{x^4}{192} - \frac{x^6}{9216} + O(x^8) \right) \ln(x)}{2} + \frac{1 - \frac{3x^4}{64} + \frac{7x^6}{2304} + O(x^8)}{x} \right)
 \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}
 y &= c_1 x \left(1 - \frac{x^2}{8} + \frac{x^4}{192} - \frac{x^6}{9216} + O(x^8) \right) \\
 &\quad + c_2 \left(-\frac{x \left(1 - \frac{x^2}{8} + \frac{x^4}{192} - \frac{x^6}{9216} + O(x^8) \right) \ln(x)}{2} + \frac{1 - \frac{3x^4}{64} + \frac{7x^6}{2304} + O(x^8)}{x} \right) \quad (1)
 \end{aligned}$$

Verification of solutions

$$\begin{aligned}
 y &= c_1 x \left(1 - \frac{x^2}{8} + \frac{x^4}{192} - \frac{x^6}{9216} + O(x^8) \right) \\
 &\quad + c_2 \left(-\frac{x \left(1 - \frac{x^2}{8} + \frac{x^4}{192} - \frac{x^6}{9216} + O(x^8) \right) \ln(x)}{2} + \frac{1 - \frac{3x^4}{64} + \frac{7x^6}{2304} + O(x^8)}{x} \right)
 \end{aligned}$$

Verified OK.

19.8.1 Maple step by step solution

Let's solve

$$x^2 y'' + xy' + (x^2 - 1)y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{(x^2-1)y}{x^2} - \frac{y'}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{y'}{x} + \frac{(x^2-1)y}{x^2} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$\left[P_2(x) = \frac{1}{x}, P_3(x) = \frac{x^2-1}{x^2} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 1$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = -1$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$x^2 y'' + xy' + (x^2 - 1)y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 0..2$

$$x^m \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

- Shift index using $k \rightarrow k - m$

$$x^m \cdot y = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

- Convert $x \cdot y'$ to series expansion

$$x \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r}$$

- Convert $x^2 \cdot y''$ to series expansion

$$x^2 \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(1+r)(-1+r)x^r + a_1(2+r)r x^{1+r} + \left(\sum_{k=2}^{\infty} (a_k(k+r+1)(k+r-1) + a_{k-2}) x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$(1+r)(-1+r) = 0$$
- Values of r that satisfy the indicial equation

$$r \in \{-1, 1\}$$
- Each term must be 0

$$a_1(2+r)r = 0$$
- Solve for the dependent coefficient(s)

$$a_1 = 0$$
- Each term in the series must be 0, giving the recursion relation

$$a_k(k+r+1)(k+r-1) + a_{k-2} = 0$$
- Shift index using $k \rightarrow k+2$

$$a_{k+2}(k+3+r)(k+r+1) + a_k = 0$$
- Recursion relation that defines series solution to ODE

$$a_{k+2} = -\frac{a_k}{(k+3+r)(k+r+1)}$$
- Recursion relation for $r = -1$

$$a_{k+2} = -\frac{a_k}{(k+2)k}$$
- Solution for $r = -1$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k-1}, a_{k+2} = -\frac{a_k}{(k+2)k}, a_1 = 0 \right]$$

- Recursion relation for $r = 1$

$$a_{k+2} = -\frac{a_k}{(k+4)(k+2)}$$

- Solution for $r = 1$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+1}, a_{k+2} = -\frac{a_k}{(k+4)(k+2)}, a_1 = 0 \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^{k-1} \right) + \left(\sum_{k=0}^{\infty} b_k x^{k+1} \right), a_{k+2} = -\frac{a_k}{(k+2)k}, a_1 = 0, b_{k+2} = -\frac{b_k}{(k+4)(k+2)}, b_1 = 0 \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
    <- Bessel successful
<- special function solution successful`

```

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 53

```

Order:=8;
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)+(x^2-1)*y(x)=0,y(x),type='series',x=0);

```

$$y(x) = \frac{c_1 x^2 \left(1 - \frac{1}{8} x^2 + \frac{1}{192} x^4 - \frac{1}{9216} x^6 + O(x^8) \right) + c_2 (\ln(x) \left(x^2 - \frac{1}{8} x^4 + \frac{1}{192} x^6 + O(x^8) \right) + \left(-2 + \frac{3}{32} x^4 - \frac{7}{1152} x^6 + O(x^8) \right))}{x}$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 75

```
AsymptoticDSolveValue[x^2*y''[x]+x*y'[x]+(x^2-1)*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_2 \left(-\frac{x^7}{9216} + \frac{x^5}{192} - \frac{x^3}{8} + x \right) + c_1 \left(\frac{5x^6 - 90x^4 + 288x^2 + 1152}{1152x} - \frac{1}{384}x(x^4 - 24x^2 + 192) \log(x) \right)$$

19.9 problem 3(f)

19.9.1 Maple step by step solution 1653

Internal problem ID [6061]

Internal file name [OUTPUT/5309_Sunday_June_05_2022_03_33_46_PM_14042674/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 166

Problem number: 3(f).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference is integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries], [_2nd_order , _linear , `
  _with_symmetry_[0,F(x)]`]]
```

$$x^2y'' - 2x^2y' + (4x - 2)y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2y'' - 2x^2y' + (4x - 2)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -2$$
$$q(x) = \frac{4x - 2}{x^2}$$

Table 242: Table $p(x), q(x)$ singularities.

$p(x) = -2$	
singularity	type

$q(x) = \frac{4x-2}{x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2 y'' - 2x^2 y' + (4x - 2)y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned}
 & x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\
 & - 2x^2 \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (4x-2) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0
 \end{aligned} \tag{1}$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-2x^{1+n+r} a_n (n+r)) \\ & + \left(\sum_{n=0}^{\infty} 4x^{1+n+r} a_n \right) + \sum_{n=0}^{\infty} (-2a_n x^{n+r}) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=0}^{\infty} (-2x^{1+n+r} a_n (n+r)) &= \sum_{n=1}^{\infty} (-2a_{n-1} (n+r-1) x^{n+r}) \\ \sum_{n=0}^{\infty} 4x^{1+n+r} a_n &= \sum_{n=1}^{\infty} 4a_{n-1} x^{n+r} \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=1}^{\infty} (-2a_{n-1} (n+r-1) x^{n+r}) \\ & + \left(\sum_{n=1}^{\infty} 4a_{n-1} x^{n+r} \right) + \sum_{n=0}^{\infty} (-2a_n x^{n+r}) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r} a_n (n+r) (n+r-1) - 2a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$x^r a_0 r(-1+r) - 2a_0 x^r = 0$$

Or

$$(x^r r(-1+r) - 2x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(r^2 - r - 2) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$r^2 - r - 2 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 2 \\ r_2 &= -1 \end{aligned}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(r^2 - r - 2) x^r = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = 3$ is an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= C y_1(x) \ln(x) + x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= x^2 \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= C y_1(x) \ln(x) + \frac{\sum_{n=0}^{\infty} b_n x^n}{x} \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^{n+2} \\ y_2(x) &= C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n-1} \right) \end{aligned}$$

Where C above can be zero. We start by finding y_1 . Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) - 2a_{n-1}(n+r-1) + 4a_{n-1} - 2a_n = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{2a_{n-1}(n+r-3)}{n^2 + 2nr + r^2 - n - r - 2} \quad (4)$$

Which for the root $r = 2$ becomes

$$a_n = \frac{2a_{n-1}(n-1)}{n(n+3)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 2$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = \frac{-4 + 2r}{r^2 + r - 2}$$

Which for the root $r = 2$ becomes

$$a_1 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-4+2r}{r^2+r-2}$	0

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{-8 + 4r}{(r+3)r(r+2)}$$

Which for the root $r = 2$ becomes

$$a_2 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-4+2r}{r^2+r-2}$	0
a_2	$\frac{-8+4r}{(r+3)r(r+2)}$	0

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{-16 + 8r}{(r + 3)(r + 2)(r + 4)(r + 1)}$$

Which for the root $r = 2$ becomes

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-4+2r}{r^2+r-2}$	0
a_2	$\frac{-8+4r}{(r+3)r(r+2)}$	0
a_3	$\frac{-16+8r}{(r+3)(r+2)(r+4)(r+1)}$	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{-32 + 16r}{(r + 3)(r + 2)^2(r + 4)(5 + r)}$$

Which for the root $r = 2$ becomes

$$a_4 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-4+2r}{r^2+r-2}$	0
a_2	$\frac{-8+4r}{(r+3)r(r+2)}$	0
a_3	$\frac{-16+8r}{(r+3)(r+2)(r+4)(r+1)}$	0
a_4	$\frac{-32+16r}{(r+3)(r+2)^2(r+4)(5+r)}$	0

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{-64 + 32r}{(r + 3)^2(r + 2)(r + 4)(5 + r)(r + 6)}$$

Which for the root $r = 2$ becomes

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-4+2r}{r^2+r-2}$	0
a_2	$\frac{-8+4r}{(r+3)r(r+2)}$	0
a_3	$\frac{-16+8r}{(r+3)(r+2)(r+4)(r+1)}$	0
a_4	$\frac{-32+16r}{(r+3)(r+2)^2(r+4)(5+r)}$	0
a_5	$\frac{-64+32r}{(r+3)^2(r+2)(r+4)(5+r)(r+6)}$	0

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{-128 + 64r}{(r+3)(r+2)(r+4)^2(5+r)(r+6)(r+7)}$$

Which for the root $r = 2$ becomes

$$a_6 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-4+2r}{r^2+r-2}$	0
a_2	$\frac{-8+4r}{(r+3)r(r+2)}$	0
a_3	$\frac{-16+8r}{(r+3)(r+2)(r+4)(r+1)}$	0
a_4	$\frac{-32+16r}{(r+3)(r+2)^2(r+4)(5+r)}$	0
a_5	$\frac{-64+32r}{(r+3)^2(r+2)(r+4)(5+r)(r+6)}$	0
a_6	$\frac{-128+64r}{(r+3)(r+2)(r+4)^2(5+r)(r+6)(r+7)}$	0

For $n = 7$, using the above recursive equation gives

$$a_7 = \frac{-256 + 128r}{(r+3)(r+2)(r+4)(5+r)^2(r+6)(r+7)(r+8)}$$

Which for the root $r = 2$ becomes

$$a_7 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-4+2r}{r^2+r-2}$	0
a_2	$\frac{-8+4r}{(r+3)r(r+2)}$	0
a_3	$\frac{-16+8r}{(r+3)(r+2)(r+4)(r+1)}$	0
a_4	$\frac{-32+16r}{(r+3)(r+2)^2(r+4)(5+r)}$	0
a_5	$\frac{-64+32r}{(r+3)^2(r+2)(r+4)(5+r)(r+6)}$	0
a_6	$\frac{-128+64r}{(r+3)(r+2)(r+4)^2(5+r)(r+6)(r+7)}$	0
a_7	$\frac{-256+128r}{(r+3)(r+2)(r+4)(5+r)^2(r+6)(r+7)(r+8)}$	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x^2(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots) \\ &= x^2(1 + O(x^8)) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Let

$$r_1 - r_2 = N$$

Where N is positive integer which is the difference between the two roots. r_1 is taken as the larger root. Hence for this problem we have $N = 3$. Now we need to determine if C is zero or not. This is done by finding $\lim_{r \rightarrow r_2} a_3(r)$. If this limit exists, then $C = 0$, else we need to keep the log term and $C \neq 0$. The above table shows that

$$\begin{aligned} a_N &= a_3 \\ &= \frac{-16 + 8r}{(r + 3)(r + 2)(r + 4)(r + 1)} \end{aligned}$$

Therefore

$$\begin{aligned} \lim_{r \rightarrow r_2} \frac{-16 + 8r}{(r + 3)(r + 2)(r + 4)(r + 1)} &= \lim_{r \rightarrow -1} \frac{-16 + 8r}{(r + 3)(r + 2)(r + 4)(r + 1)} \\ &= \text{undefined} \end{aligned}$$

Since the limit does not exist then the log term is needed. Therefore the second solution has the form

$$y_2(x) = Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right)$$

Therefore

$$\begin{aligned}
 \frac{d}{dx}y_2(x) &= Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2}(n+r_2)}{x} \right) \\
 &= Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} x^{-1+n+r_2} b_n (n+r_2) \right) \\
 \frac{d^2}{dx^2}y_2(x) &= Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} \\
 &\quad + \sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2}(n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2}(n+r_2)}{x^2} \right) \\
 &= Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} + \left(\sum_{n=0}^{\infty} x^{-2+n+r_2} b_n (n+r_2) (-1+n+r_2) \right)
 \end{aligned}$$

Substituting these back into the given ode $x^2y'' - 2x^2y' + (4x - 2)y = 0$ gives

$$\begin{aligned}
 &x^2 \left(Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} \right. \\
 &\quad \left. + \sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2}(n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2}(n+r_2)}{x^2} \right) \right) \\
 &\quad - 2x^2 \left(Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2}(n+r_2)}{x} \right) \right) \\
 &\quad + (4x - 2) \left(Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) \right) = 0
 \end{aligned}$$

Which can be written as

$$\begin{aligned}
 &\left((x^2y_1''(x) - 2x^2y_1'(x) + (4x - 2)y_1(x)) \ln(x) + x^2 \left(\frac{2y_1'(x)}{x} - \frac{y_1(x)}{x^2} \right) \right. \\
 &\quad \left. - 2xy_1(x) \right) C + x^2 \left(\sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2}(n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2}(n+r_2)}{x^2} \right) \right) \\
 &\quad - 2x^2 \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2}(n+r_2)}{x} \right) + (4x - 2) \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) = 0
 \end{aligned} \tag{7}$$

But since $y_1(x)$ is a solution to the ode, then

$$x^2 y_1''(x) - 2x^2 y_1'(x) + (4x - 2) y_1(x) = 0$$

Eq (7) simplifies to

$$\begin{aligned} & \left(x^2 \left(\frac{2y_1'(x)}{x} - \frac{y_1(x)}{x^2} \right) - 2xy_1(x) \right) C \\ & + x^2 \left(\sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2} (n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2} (n+r_2)}{x^2} \right) \right) \\ & - 2x^2 \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2} (n+r_2)}{x} \right) + (4x - 2) \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) = 0 \end{aligned} \quad (8)$$

Substituting $y_1 = \sum_{n=0}^{\infty} a_n x^{n+r_1}$ into the above gives

$$\begin{aligned} & \left(2 \left(\sum_{n=0}^{\infty} x^{-1+n+r_1} a_n (n+r_1) \right) x + (-1 - 2x) \left(\sum_{n=0}^{\infty} a_n x^{n+r_1} \right) \right) C \\ & + \left(\sum_{n=0}^{\infty} x^{-2+n+r_2} b_n (n+r_2) (-1+n+r_2) \right) x^2 \\ & - 2x^2 \left(\sum_{n=0}^{\infty} x^{-1+n+r_2} b_n (n+r_2) \right) + 2(2x - 1) \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) = 0 \end{aligned} \quad (9)$$

Since $r_1 = 2$ and $r_2 = -1$ then the above becomes

$$\begin{aligned} & \left(2 \left(\sum_{n=0}^{\infty} x^{1+n} a_n (n+2) \right) x + (-1 - 2x) \left(\sum_{n=0}^{\infty} a_n x^{n+2} \right) \right) C \\ & + \left(\sum_{n=0}^{\infty} x^{-3+n} b_n (n-1) (-2+n) \right) x^2 \\ & - 2x^2 \left(\sum_{n=0}^{\infty} x^{-2+n} b_n (n-1) \right) + 2(2x - 1) \left(\sum_{n=0}^{\infty} b_n x^{n-1} \right) = 0 \end{aligned} \quad (10)$$

Which simplifies to

$$\begin{aligned}
 & \left(\sum_{n=0}^{\infty} 2C x^{n+2} a_n (n+2) \right) + \sum_{n=0}^{\infty} (-C x^{n+2} a_n) \\
 & + \sum_{n=0}^{\infty} (-2C x^{n+3} a_n) + \left(\sum_{n=0}^{\infty} x^{n-1} b_n (n^2 - 3n + 2) \right) \\
 & + \sum_{n=0}^{\infty} (-2x^n b_n (n-1)) + \left(\sum_{n=0}^{\infty} 4b_n x^n \right) + \sum_{n=0}^{\infty} (-2b_n x^{n-1}) = 0
 \end{aligned} \tag{2A}$$

The next step is to make all powers of x be $n - 1$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n-1} and adjusting the power and the corresponding index gives

$$\begin{aligned}
 \sum_{n=0}^{\infty} 2C x^{n+2} a_n (n+2) &= \sum_{n=3}^{\infty} 2C a_{-3+n} (n-1) x^{n-1} \\
 \sum_{n=0}^{\infty} (-C x^{n+2} a_n) &= \sum_{n=3}^{\infty} (-C a_{-3+n} x^{n-1}) \\
 \sum_{n=0}^{\infty} (-2C x^{n+3} a_n) &= \sum_{n=4}^{\infty} (-2C a_{n-4} x^{n-1}) \\
 \sum_{n=0}^{\infty} (-2x^n b_n (n-1)) &= \sum_{n=1}^{\infty} (-2b_{n-1} (-2+n) x^{n-1}) \\
 \sum_{n=0}^{\infty} 4b_n x^n &= \sum_{n=1}^{\infty} 4b_{n-1} x^{n-1}
 \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n - 1$.

$$\begin{aligned}
 & \left(\sum_{n=3}^{\infty} 2C a_{-3+n} (n-1) x^{n-1} \right) + \sum_{n=3}^{\infty} (-C a_{-3+n} x^{n-1}) + \sum_{n=4}^{\infty} (-2C a_{n-4} x^{n-1}) \\
 & + \left(\sum_{n=0}^{\infty} x^{n-1} b_n (n^2 - 3n + 2) \right) + \sum_{n=1}^{\infty} (-2b_{n-1} (-2+n) x^{n-1}) \\
 & + \left(\sum_{n=1}^{\infty} 4b_{n-1} x^{n-1} \right) + \sum_{n=0}^{\infty} (-2b_n x^{n-1}) = 0
 \end{aligned} \tag{2B}$$

For $n = 0$ in Eq. (2B), we choose arbitrary value for b_0 as $b_0 = 1$. For $n = 1$, Eq (2B) gives

$$6b_0 - 2b_1 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$6 - 2b_1 = 0$$

Solving the above for b_1 gives

$$b_1 = 3$$

For $n = 2$, Eq (2B) gives

$$4b_1 - 2b_2 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$12 - 2b_2 = 0$$

Solving the above for b_2 gives

$$b_2 = 6$$

For $n = N$, where $N = 3$ which is the difference between the two roots, we are free to choose $b_3 = 0$. Hence for $n = 3$, Eq (2B) gives

$$3C + 12 = 0$$

Which is solved for C . Solving for C gives

$$C = -4$$

For $n = 4$, Eq (2B) gives

$$(-2a_0 + 5a_1)C + 4b_4 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$8 + 4b_4 = 0$$

Solving the above for b_4 gives

$$b_4 = -2$$

For $n = 5$, Eq (2B) gives

$$(-2a_1 + 7a_2)C - 2b_4 + 10b_5 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$4 + 10b_5 = 0$$

Solving the above for b_5 gives

$$b_5 = -\frac{2}{5}$$

For $n = 6$, Eq (2B) gives

$$(-2a_2 + 9a_3)C - 4b_5 + 18b_6 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$\frac{8}{5} + 18b_6 = 0$$

Solving the above for b_6 gives

$$b_6 = -\frac{4}{45}$$

For $n = 7$, Eq (2B) gives

$$(-2a_3 + 11a_4)C - 6b_6 + 28b_7 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$\frac{8}{15} + 28b_7 = 0$$

Solving the above for b_7 gives

$$b_7 = -\frac{2}{105}$$

Now that we found all b_n and C , we can calculate the second solution from

$$y_2(x) = Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right)$$

Using the above value found for $C = -4$ and all b_n , then the second solution becomes

$$y_2(x) = (-4) (x^2(1 + O(x^8))) \ln(x) + \frac{1 + 3x + 6x^2 - 2x^4 - \frac{2x^5}{5} - \frac{4x^6}{45} - \frac{2x^7}{105} + O(x^8)}{x}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1 y_1(x) + c_2 y_2(x) \\ &= c_1 x^2(1 + O(x^8)) + c_2 \left((-4) (x^2(1 + O(x^8))) \ln(x) \right. \\ &\quad \left. + \frac{1 + 3x + 6x^2 - 2x^4 - \frac{2x^5}{5} - \frac{4x^6}{45} - \frac{2x^7}{105} + O(x^8)}{x} \right) \end{aligned}$$

Hence the final solution is

$$\begin{aligned}
 y &= y_h \\
 &= c_1 x^2 (1 + O(x^8)) \\
 &\quad + c_2 \left(-4x^2 (1 + O(x^8)) \ln(x) + \frac{1 + 3x + 6x^2 - 2x^4 - \frac{2x^5}{5} - \frac{4x^6}{45} - \frac{2x^7}{105} + O(x^8)}{x} \right)
 \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}
 y &= c_1 x^2 (1 + O(x^8)) + c_2 \left(-4x^2 (1 + O(x^8)) \ln(x) \right. \\
 &\quad \left. + \frac{1 + 3x + 6x^2 - 2x^4 - \frac{2x^5}{5} - \frac{4x^6}{45} - \frac{2x^7}{105} + O(x^8)}{x} \right) \quad (1)
 \end{aligned}$$

Verification of solutions

$$\begin{aligned}
 y &= c_1 x^2 (1 + O(x^8)) \\
 &\quad + c_2 \left(-4x^2 (1 + O(x^8)) \ln(x) + \frac{1 + 3x + 6x^2 - 2x^4 - \frac{2x^5}{5} - \frac{4x^6}{45} - \frac{2x^7}{105} + O(x^8)}{x} \right)
 \end{aligned}$$

Verified OK.

19.9.1 Maple step by step solution

Let's solve

$$x^2 y'' - 2x^2 y' + (4x - 2)y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{2(2x-1)y}{x^2} + 2y'$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - 2y' + \frac{2(2x-1)y}{x^2} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point
 - Define functions

$$\left[P_2(x) = -2, P_3(x) = \frac{2(2x-1)}{x^2} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 0$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = -2$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$x^2 y'' - 2x^2 y' + (4x - 2)y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 0..1$

$$x^m \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

- Shift index using $k \rightarrow k - m$

$$x^m \cdot y = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

- Convert $x^2 \cdot y'$ to series expansion

$$x^2 \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r+1}$$

- Shift index using $k \rightarrow k - 1$

$$x^2 \cdot y' = \sum_{k=1}^{\infty} a_{k-1} (k-1+r) x^{k+r}$$

- Convert $x^2 \cdot y''$ to series expansion

$$x^2 \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k-1+r) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(1+r)(-2+r)x^r + \left(\sum_{k=1}^{\infty} (a_k(k+r+1)(k+r-2) - 2a_{k-1}(k-3+r))x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$(1+r)(-2+r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \{-1, 2\}$$

- Each term in the series must be 0, giving the recursion relation

$$a_k(k+r+1)(k+r-2) - 2a_{k-1}(k-3+r) = 0$$

- Shift index using $k \rightarrow k+1$

$$a_{k+1}(k+2+r)(k-1+r) - 2a_k(k+r-2) = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+1} = \frac{2a_k(k+r-2)}{(k+2+r)(k-1+r)}$$

- Recursion relation for $r = -1$; series terminates at $k = 3$

$$a_{k+1} = \frac{2a_k(k-3)}{(k+1)(k-2)}$$

- Series not valid for $r = -1$, division by 0 in the recursion relation at $k = 2$

$$a_{k+1} = \frac{2a_k(k-3)}{(k+1)(k-2)}$$

- Recursion relation for $r = 2$

$$a_{k+1} = \frac{2a_k k}{(k+4)(k+1)}$$

- Solution for $r = 2$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+2}, a_{k+1} = \frac{2a_k k}{(k+4)(k+1)} \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
<- linear_1 successful`

```


✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 55

```
Order:=8;  
dsolve(x^2*diff(y(x),x$2)-2*x^2*diff(y(x),x)+(4*x-2)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^2 (1 + O(x^8)) + \frac{c_2 (\ln(x) ((-48)x^3 + O(x^8)) + (12 + 36x + 72x^2 + 88x^3 - 24x^4 - \frac{24}{5}x^5 - \frac{16}{15}x^6 - \frac{8}{35}x^7 + O(x^8)))}{x}$$

✓ Solution by Mathematica

Time used: 0.1 (sec). Leaf size: 58

```
AsymptoticDSolveValue[x^2*y''[x]-2*x^2*y'[x]+(4*x-2)*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_2 x^2 + c_1 \left(-4x^2 \log(x) - \frac{4x^6 + 18x^5 + 90x^4 - 390x^3 - 270x^2 - 135x - 45}{45x} \right)$$

**20 Chapter 4. Linear equations with Regular
Singular Points. Page 182**

20.1 problem 4 1658

20.1 problem 4

20.1.1 Maple step by step solution 1666

Internal problem ID [6062]

Internal file name [OUTPUT/5310_Sunday_June_05_2022_03_33_49_PM_29452965/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 4. Linear equations with Regular Singular Points. Page 182

Problem number: 4.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second order series method. Ordinary point", "second_order_change_of_variable_on_y_method_2", "second order series method. Taylor series method", "second_order_ode_non_constant_coeff_transformation_on_B"

Maple gives the following as the ode type

[_Gegenbauer]

$$(-x^2 + 1)y'' - 2xy' + 2y = 0$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using

Taylor series gives

$$\begin{aligned}
 y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \cdots \\
 &= y_0 + xy'_0 + \frac{x^2}{2}f|_{x_0, y_0, y'_0} + \frac{x^3}{3!}f'|_{x_0, y_0, y'_0} + \cdots \\
 &= y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0}
 \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \tag{1}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \tag{332}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \tag{333}$$

$$\begin{aligned}
 \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \tag{2}
 \end{aligned}$$

$$\begin{aligned}
 \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \tag{3}
 \end{aligned}$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \tag{4}$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \tag{5}$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \end{aligned}$$

⋮

$$\begin{aligned} F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \tag{6}$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \tag{7}$$

To find $y(x)$ series solution around $x = 0$. Hence

$$F_0 = -\frac{2(-y + xy')}{x^2 - 1}$$

$$\begin{aligned} F_1 &= \frac{dF_0}{dx} \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \\ &= \frac{8(-y + xy') x}{(x^2 - 1)^2} \end{aligned}$$

$$\begin{aligned} F_2 &= \frac{dF_1}{dx} \\ &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} y' + \frac{\partial F_1}{\partial y'} F_1 \\ &= -\frac{8(-y + xy')(5x^2 + 1)}{(x^2 - 1)^3} \end{aligned}$$

$$\begin{aligned} F_3 &= \frac{dF_2}{dx} \\ &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} y' + \frac{\partial F_2}{\partial y'} F_2 \\ &= \frac{240(x^2 + \frac{3}{5})(-y + xy') x}{(x^2 - 1)^4} \end{aligned}$$

$$\begin{aligned} F_4 &= \frac{dF_3}{dx} \\ &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} y' + \frac{\partial F_3}{\partial y'} F_3 \\ &= -\frac{48(-y + xy')(35x^4 + 42x^2 + 3)}{(x^2 - 1)^5} \end{aligned}$$

$$\begin{aligned} F_5 &= \frac{dF_4}{dx} \\ &= \frac{\partial F_4}{\partial x} + \frac{\partial F_4}{\partial y} y' + \frac{\partial F_4}{\partial y'} F_4 \\ &= \frac{13440(-y + xy') x (x^4 + 2x^2 + \frac{3}{7})}{(x^2 - 1)^6} \end{aligned}$$

$$\begin{aligned} F_6 &= \frac{dF_5}{dx} \\ &= \frac{\partial F_5}{\partial x} + \frac{\partial F_5}{\partial y} y' + \frac{\partial F_5}{\partial y'} F_5 \\ &= -\frac{5760(-y + xy')(21x^6 + 63x^4 + 27x^2 + 1)}{(x^2 - 1)^7} \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and $y'(0) = y'(0)$ gives

$$\begin{aligned} F_0 &= -2y(0) \\ F_1 &= 0 \\ F_2 &= -8y(0) \\ F_3 &= 0 \\ F_4 &= -144y(0) \\ F_5 &= 0 \\ F_6 &= -5760y(0) \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$y = \left(1 - x^2 - \frac{1}{3}x^4 - \frac{1}{5}x^6 - \frac{1}{7}x^8\right) y(0) + xy'(0) + O(x^8)$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard power series. The ode is normalized to be

$$(-x^2 + 1)y'' - 2xy' + 2y = 0$$

Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$\begin{aligned} y' &= \sum_{n=1}^{\infty} n a_n x^{n-1} \\ y'' &= \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \end{aligned}$$

Substituting the above back into the ode gives

$$(-x^2 + 1) \left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) - 2x \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) + 2 \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0 \quad (1)$$

Which simplifies to

$$\sum_{n=2}^{\infty} (-x^n a_n n(n-1)) + \left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \sum_{n=1}^{\infty} (-2n a_n x^n) + \left(\sum_{n=0}^{\infty} 2a_n x^n \right) = 0 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\begin{aligned} \sum_{n=2}^{\infty} (-x^n a_n n(n-1)) + \left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n \right) \\ + \sum_{n=1}^{\infty} (-2n a_n x^n) + \left(\sum_{n=0}^{\infty} 2a_n x^n \right) = 0 \end{aligned} \quad (3)$$

$n = 0$ gives

$$2a_2 + 2a_0 = 0$$

$$a_2 = -a_0$$

For $2 \leq n$, the recurrence equation is

$$-n a_n (n-1) + (n+2) a_{n+2} (n+1) - 2n a_n + 2a_n = 0 \quad (4)$$

Solving for a_{n+2} , gives

$$a_{n+2} = \frac{(n-1) a_n}{n+1} \quad (5)$$

For $n = 2$ the recurrence equation gives

$$-4a_2 + 12a_4 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = -\frac{a_0}{3}$$

For $n = 3$ the recurrence equation gives

$$-10a_3 + 20a_5 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = 0$$

For $n = 4$ the recurrence equation gives

$$-18a_4 + 30a_6 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = -\frac{a_0}{5}$$

For $n = 5$ the recurrence equation gives

$$-28a_5 + 42a_7 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = 0$$

For $n = 6$ the recurrence equation gives

$$-40a_6 + 56a_8 = 0$$

Which after substituting the earlier terms found becomes

$$a_8 = -\frac{a_0}{7}$$

For $n = 7$ the recurrence equation gives

$$-54a_7 + 72a_9 = 0$$

Which after substituting the earlier terms found becomes

$$a_9 = 0$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + a_1 x - a_0 x^2 - \frac{1}{3} a_0 x^4 - \frac{1}{5} a_0 x^6 + \dots$$

Collecting terms, the solution becomes

$$y = \left(1 - x^2 - \frac{1}{3} x^4 - \frac{1}{5} x^6 \right) a_0 + a_1 x + O(x^8) \quad (3)$$

At $x = 0$ the solution above becomes

$$y = \left(1 - x^2 - \frac{1}{3} x^4 - \frac{1}{5} x^6 \right) c_1 + c_2 x + O(x^8)$$

Summary

The solution(s) found are the following

$$y = \left(1 - x^2 - \frac{1}{3} x^4 - \frac{1}{5} x^6 - \frac{1}{7} x^8 \right) y(0) + xy'(0) + O(x^8) \quad (1)$$

$$y = \left(1 - x^2 - \frac{1}{3} x^4 - \frac{1}{5} x^6 \right) c_1 + c_2 x + O(x^8) \quad (2)$$

Verification of solutions

$$y = \left(1 - x^2 - \frac{1}{3} x^4 - \frac{1}{5} x^6 - \frac{1}{7} x^8 \right) y(0) + xy'(0) + O(x^8)$$

Verified OK.

$$y = \left(1 - x^2 - \frac{1}{3} x^4 - \frac{1}{5} x^6 \right) c_1 + c_2 x + O(x^8)$$

Verified OK.

20.1.1 Maple step by step solution

Let's solve

$$(-x^2 + 1)y'' - 2xy' + 2y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{2xy'}{x^2-1} + \frac{2y}{x^2-1}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{2xy'}{x^2-1} - \frac{2y}{x^2-1} = 0$$

- Check to see if x_0 is a regular singular point

- Define functions

$$[P_2(x) = \frac{2x}{x^2-1}, P_3(x) = -\frac{2}{x^2-1}]$$

- $(1+x) \cdot P_2(x)$ is analytic at $x = -1$

$$((1+x) \cdot P_2(x)) \Big|_{x=-1} = 1$$

- $(1+x)^2 \cdot P_3(x)$ is analytic at $x = -1$

$$((1+x)^2 \cdot P_3(x)) \Big|_{x=-1} = 0$$

- $x = -1$ is a regular singular point

Check to see if x_0 is a regular singular point

$$x_0 = -1$$

- Multiply by denominators

$$(x^2 - 1)y'' + 2xy' - 2y = 0$$

- Change variables using $x = u - 1$ so that the regular singular point is at $u = 0$

$$(u^2 - 2u) \left(\frac{d^2}{du^2} y(u) \right) + (2u - 2) \left(\frac{d}{du} y(u) \right) - 2y(u) = 0$$

- Assume series solution for $y(u)$

$$y(u) = \sum_{k=0}^{\infty} a_k u^{k+r}$$

- Rewrite ODE with series expansions

- Convert $u^m \cdot \left(\frac{d}{du} y(u) \right)$ to series expansion for $m = 0..1$

$$u^m \cdot \left(\frac{d}{du} y(u) \right) = \sum_{k=0}^{\infty} a_k (k+r) u^{k+r-1+m}$$

- Shift index using $k \rightarrow k+1-m$

$$u^m \cdot \left(\frac{d}{du} y(u) \right) = \sum_{k=-1+m}^{\infty} a_{k+1-m} (k+1-m+r) u^{k+r}$$

- Convert $u^m \cdot \left(\frac{d^2}{du^2} y(u) \right)$ to series expansion for $m = 1.2$

$$u^m \cdot \left(\frac{d^2}{du^2} y(u) \right) = \sum_{k=0}^{\infty} a_k (k+r) (k+r-1) u^{k+r-2+m}$$

- Shift index using $k \rightarrow k+2-m$

$$u^m \cdot \left(\frac{d^2}{du^2} y(u) \right) = \sum_{k=-2+m}^{\infty} a_{k+2-m} (k+2-m+r) (k+1-m+r) u^{k+r}$$

Rewrite ODE with series expansions

$$-2a_0 r^2 u^{-1+r} + \left(\sum_{k=0}^{\infty} (-2a_{k+1} (k+1+r)^2 + a_k (k+r+2) (k+r-1)) u^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$-2r^2 = 0$$

- Values of r that satisfy the indicial equation

$$r = 0$$

- Each term in the series must be 0, giving the recursion relation

$$-2a_{k+1} (k+1)^2 + a_k (k+2) (k-1) = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+1} = \frac{a_k (k+2)(k-1)}{2(k+1)^2}$$

- Recursion relation for $r = 0$; series terminates at $k = 1$

$$a_{k+1} = \frac{a_k (k+2)(k-1)}{2(k+1)^2}$$

- Apply recursion relation for $k = 0$

$$a_1 = -a_0$$

- Terminating series solution of the ODE for $r = 0$. Use reduction of order to find the second li

$$y(u) = a_0 \cdot (-u + 1)$$

- Revert the change of variables $u = 1 + x$

$$[y = -a_0 x]$$

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacics algorithm  
  A Liouvillian solution exists  
  Reducible group (found an exponential solution)  
  Group is reducible, not completely reducible  
<- Kovacics algorithm successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
Order:=8;  
dsolve((1-x^2)*diff(y(x),x$2)-2*x*diff(y(x),x)+2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - x^2 - \frac{1}{3}x^4 - \frac{1}{5}x^6\right) y(0) + D(y)(0)x + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 32

```
AsymptoticDSolveValue[(1-x^2)*y''[x]-2*x*y'[x]+2*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(-\frac{x^6}{5} - \frac{x^4}{3} - x^2 + 1 \right) + c_2 x$$

21 Chapter 5. Existence and uniqueness of solutions to first order equations. Page 190

21.1 problem 1(a)	1670
21.2 problem 1(b)	1685
21.3 problem 1(c)	1700
21.4 problem 1(d)	1720
21.5 problem 1(e)	1733
21.6 problem 2(a)	1748
21.7 problem 3(a)	1752
21.8 problem 3(b)	1756
21.9 problem 4(a)	1760
21.10 problem 4(b)	1764
21.11 problem 4(c)	1768
21.12 problem 4(d)	1772
21.13 problem 5(a)	1776
21.14 problem 5(b)	1782
21.15 problem 5(c)	1788
21.16 problem 6(b)	1792

21.1 problem 1(a)

21.1.1 Solving as separable ode	1670
21.1.2 Solving as linear ode	1672
21.1.3 Solving as homogeneousTypeD2 ode	1673
21.1.4 Solving as first order ode lie symmetry lookup ode	1675
21.1.5 Solving as exact ode	1679
21.1.6 Maple step by step solution	1683

Internal problem ID [6063]

Internal file name [OUTPUT/5311_Sunday_June_05_2022_03_33_51_PM_79278784/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 190

Problem number: 1(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "linear", "separable", "homogeneousTypeD2", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_separable]

$$y' - yx^2 = 0$$

21.1.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= y x^2\end{aligned}$$

Where $f(x) = x^2$ and $g(y) = y$. Integrating both sides gives

$$\begin{aligned}\frac{1}{y} dy &= x^2 dx \\ \int \frac{1}{y} dy &= \int x^2 dx \\ \ln(y) &= \frac{x^3}{3} + c_1 \\ y &= e^{\frac{x^3}{3} + c_1} \\ &= c_1 e^{\frac{x^3}{3}}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{\frac{x^3}{3}} \tag{1}$$




Figure 177: Slope field plot

Verification of solutions

$$y = c_1 e^{\frac{x^3}{3}}$$

Verified OK.

21.1.2 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -x^2$$

$$q(x) = 0$$

Hence the ode is

$$y' - yx^2 = 0$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int -x^2 dx} \\ &= e^{-\frac{x^3}{3}}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}\mu y &= 0 \\ \frac{d}{dx}\left(e^{-\frac{x^3}{3}}y\right) &= 0\end{aligned}$$

Integrating gives

$$e^{-\frac{x^3}{3}}y = c_1$$

Dividing both sides by the integrating factor $\mu = e^{-\frac{x^3}{3}}$ results in

$$y = c_1 e^{\frac{x^3}{3}}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{\frac{x^3}{3}} \tag{1}$$




Figure 178: Slope field plot

Verification of solutions

$$y = c_1 e^{\frac{x^3}{3}}$$

Verified OK.

21.1.3 Solving as homogeneous TypeD2 ode

Using the change of variables $y = u(x)x$ on the above ode results in new ode in $u(x)$

$$u'(x)x + u(x) - u(x)x^3 = 0$$

In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= \frac{u(x^3 - 1)}{x} \end{aligned}$$

Where $f(x) = \frac{x^3-1}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u} du &= \frac{x^3-1}{x} dx \\ \int \frac{1}{u} du &= \int \frac{x^3-1}{x} dx \\ \ln(u) &= \frac{x^3}{3} - \ln(x) + c_2 \\ u &= e^{\frac{x^3}{3} - \ln(x) + c_2} \\ &= c_2 e^{\frac{x^3}{3} - \ln(x)}\end{aligned}$$

Which simplifies to

$$u(x) = \frac{c_2 e^{\frac{x^3}{3}}}{x}$$

Therefore the solution y is

$$\begin{aligned}y &= xu \\ &= c_2 e^{\frac{x^3}{3}}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_2 e^{\frac{x^3}{3}} \tag{1}$$




Figure 179: Slope field plot

Verification of solutions

$$y = c_2 e^{\frac{x^3}{3}}$$

Verified OK.

21.1.4 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = y x^2$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 245: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= e^{\frac{x^3}{3}}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^{\frac{x^3}{3}}} dy \end{aligned}$$

Which results in

$$S = e^{-\frac{x^3}{3}} y$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = y x^2$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= -x^2 e^{-\frac{x^3}{3}} y \\ S_y &= e^{-\frac{x^3}{3}} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by

integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \tag{4}$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$e^{-\frac{x^3}{3}} y = c_1$$



Which simplifies to

$$e^{-\frac{x^3}{3}} y = c_1$$

Which gives

$$y = c_1 e^{\frac{x^3}{3}}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
<p style="text-align: center;">$\frac{dy}{dx} = y x^2$</p> 	<p style="text-align: center;">$R = x$ $S = e^{-\frac{x^3}{3}} y$</p>	<p style="text-align: center;">$\frac{dS}{dR} = 0$</p> 

Summary

The solution(s) found are the following

$$y = c_1 e^{\frac{x^3}{3}} \tag{1}$$




Figure 180: Slope field plot

Verification of solutions

$$y = c_1 e^{\frac{x^3}{3}}$$

Verified OK.

21.1.5 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}\left(\frac{1}{y}\right) dy &= (x^2) dx \\ (-x^2) dx + \left(\frac{1}{y}\right) dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -x^2 \\ N(x, y) &= \frac{1}{y}\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-x^2) \\ &= 0\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{1}{y} \right) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \tag{1}$$

$$\frac{\partial \phi}{\partial y} = N \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -x^2 dx \\ \phi &= -\frac{x^3}{3} + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{1}{y}$. Therefore equation (4) becomes

$$\frac{1}{y} = 0 + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = \frac{1}{y}$$

Integrating the above w.r.t y gives

$$\begin{aligned}\int f'(y) dy &= \int \left(\frac{1}{y} \right) dy \\ f(y) &= \ln(y) + c_1\end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\frac{x^3}{3} + \ln(y) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\frac{x^3}{3} + \ln(y)$$

The solution becomes

$$y = e^{\frac{x^3}{3} + c_1}$$

Summary

The solution(s) found are the following

$$y = e^{\frac{x^3}{3} + c_1} \tag{1}$$




Figure 181: Slope field plot

Verification of solutions

$$y = e^{\frac{x^3}{3} + c_1}$$

Verified OK.

21.1.6 Maple step by step solution

Let's solve

$$y' - yx^2 = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y} = x^2$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y} dx = \int x^2 dx + c_1$$

- Evaluate integral

$$\ln(y) = \frac{x^3}{3} + c_1$$

- Solve for y

$$y = e^{\frac{x^3}{3} + c_1}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)=x^2*y(x),y(x), singsol=all)
```

$$y(x) = c_1 e^{\frac{x^3}{3}}$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 22

```
DSolve[y'[x]==x^2*y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{\frac{x^3}{3}}$$

$$y(x) \rightarrow 0$$

21.2 problem 1(b)

21.2.1 Solving as separable ode	1685
21.2.2 Solving as homogeneousTypeD2 ode	1687
21.2.3 Solving as differentialType ode	1689
21.2.4 Solving as first order ode lie symmetry lookup ode	1690
21.2.5 Solving as exact ode	1694
21.2.6 Maple step by step solution	1698

Internal problem ID [6064]

Internal file name [OUTPUT/5312_Sunday_June_05_2022_03_33_52_PM_12902458/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 190

Problem number: 1(b).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "separable", "differential-Type", "homogeneousTypeD2", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_separable]

$$y'y = x$$

21.2.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{x}{y}\end{aligned}$$

Where $f(x) = x$ and $g(y) = \frac{1}{y}$. Integrating both sides gives

$$\frac{1}{\frac{1}{y}} dy = x dx$$

$$\int \frac{1}{y} dy = \int x dx$$

$$\frac{y^2}{2} = \frac{x^2}{2} + c_1$$

Which results in

$$y = \sqrt{x^2 + 2c_1}$$

$$y = -\sqrt{x^2 + 2c_1}$$

Summary

The solution(s) found are the following

$$y = \sqrt{x^2 + 2c_1} \tag{1}$$

$$y = -\sqrt{x^2 + 2c_1} \tag{2}$$




Figure 182: Slope field plot

Verification of solutions

$$y = \sqrt{x^2 + 2c_1}$$

Verified OK.

$$y = -\sqrt{x^2 + 2c_1}$$

Verified OK.

21.2.2 Solving as homogeneous Type D2 ode

Using the change of variables $y = u(x)x$ on the above ode results in new ode in $u(x)$

$$(u'(x)x + u(x))u(x)x = x$$

In canonical form the ODE is

$$\begin{aligned}u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{u^2 - 1}{ux}\end{aligned}$$

Where $f(x) = -\frac{1}{x}$ and $g(u) = \frac{u^2-1}{u}$. Integrating both sides gives

$$\begin{aligned}\frac{1}{\frac{u^2-1}{u}} du &= -\frac{1}{x} dx \\ \int \frac{1}{\frac{u^2-1}{u}} du &= \int -\frac{1}{x} dx \\ \frac{\ln(u-1)}{2} + \frac{\ln(u+1)}{2} &= -\ln(x) + c_2\end{aligned}$$

The above can be written as

$$\begin{aligned}\left(\frac{1}{2}\right) (\ln(u-1) + \ln(u+1)) &= -\ln(x) + 2c_2 \\ \ln(u-1) + \ln(u+1) &= (2)(-\ln(x) + 2c_2) \\ &= -2\ln(x) + 4c_2\end{aligned}$$

Raising both side to exponential gives

$$e^{\ln(u-1)+\ln(u+1)} = e^{-2\ln(x)+4c_2}$$

Which simplifies to

$$\begin{aligned}u^2 - 1 &= \frac{2c_2}{x^2} \\ &= \frac{c_3}{x^2}\end{aligned}$$

The solution is

$$u(x)^2 - 1 = \frac{c_3}{x^2}$$

Replacing $u(x)$ in the above solution by $\frac{y}{x}$ results in the solution for y in implicit form

$$\begin{aligned}\frac{y^2}{x^2} - 1 &= \frac{c_3}{x^2} \\ \frac{y^2}{x^2} - 1 &= \frac{c_3}{x^2}\end{aligned}$$

Which simplifies to

$$-(x - y)(x + y) = c_3$$

Summary

The solution(s) found are the following

$$-(x - y)(x + y) = c_3 \tag{1}$$




Figure 183: Slope field plot

Verification of solutions

$$-(x - y)(x + y) = c_3$$

Verified OK.

21.2.3 Solving as differential Type ode

Writing the ode as

$$y' = \frac{x}{y} \tag{1}$$

Which becomes

$$(y) dy = (x) dx \tag{2}$$

But the RHS is complete differential because

$$(x) dx = d\left(\frac{x^2}{2}\right)$$

Hence (2) becomes

$$(y) dy = d\left(\frac{x^2}{2}\right)$$

Integrating both sides gives gives these solutions

$$y = \sqrt{x^2 + 2c_1} + c_1$$

$$y = -\sqrt{x^2 + 2c_1} + c_1$$

Summary

The solution(s) found are the following

$$y = \sqrt{x^2 + 2c_1} + c_1 \tag{1}$$

$$y = -\sqrt{x^2 + 2c_1} + c_1 \tag{2}$$




Figure 184: Slope field plot

Verification of solutions

$$y = \sqrt{x^2 + 2c_1} + c_1$$

Verified OK.

$$y = -\sqrt{x^2 + 2c_1} + c_1$$

Verified OK.

21.2.4 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{x}{y}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **separable**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 248: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= \frac{1}{x} \\ \eta(x, y) &= 0\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\eta = 0$ then in this special case

$$R = y$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\xi} dx \\ &= \int \frac{1}{\frac{1}{x}} dx \end{aligned}$$

Which results in

$$S = \frac{x^2}{2}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{x}{y}$$

Evaluating all the partial derivatives gives

$$R_x = 0$$

$$R_y = 1$$

$$S_x = x$$

$$S_y = 0$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = y \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = R$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by

integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \frac{R^2}{2} + c_1 \quad (4)$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{x^2}{2} = \frac{y^2}{2} + c_1$$

Which simplifies to

$$\frac{x^2}{2} = \frac{y^2}{2} + c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{x}{y}$ 	$R = y$ $S = \frac{x^2}{2}$	$\frac{dS}{dR} = R$ 

Summary

The solution(s) found are the following

$$\frac{x^2}{2} = \frac{y^2}{2} + c_1 \quad (1)$$




Figure 185: Slope field plot

Verification of solutions

$$\frac{x^2}{2} = \frac{y^2}{2} + c_1$$

Verified OK.

21.2.5 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}(y) dy &= (x) dx \\ (-x) dx + (y) dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -x \\ N(x, y) &= y\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-x) \\ &= 0\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(y) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\int \frac{\partial \phi}{\partial x} dx = \int M dx$$

$$\int \frac{\partial \phi}{\partial x} dx = \int -x dx$$

$$\phi = -\frac{x^2}{2} + f(y) \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = y$. Therefore equation (4) becomes

$$y = 0 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = y$$

Integrating the above w.r.t y gives

$$\int f'(y) dy = \int (y) dy$$

$$f(y) = \frac{y^2}{2} + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\frac{x^2}{2} + \frac{y^2}{2} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\frac{x^2}{2} + \frac{y^2}{2}$$

Summary

The solution(s) found are the following

$$-\frac{x^2}{2} + \frac{y^2}{2} = c_1 \tag{1}$$




Figure 186: Slope field plot

Verification of solutions

$$-\frac{x^2}{2} + \frac{y^2}{2} = c_1$$

Verified OK.

21.2.6 Maple step by step solution

Let's solve

$$y'y = x$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Integrate both sides with respect to x

$$\int y'y dx = \int x dx + c_1$$

- Evaluate integral

$$\frac{y^2}{2} = \frac{x^2}{2} + c_1$$

- Solve for y

$$\{y = \sqrt{x^2 + 2c_1}, y = -\sqrt{x^2 + 2c_1}\}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
<- Bernoulli successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(y(x)*diff(y(x),x)=x,y(x), singsol=all)
```

$$y(x) = \sqrt{x^2 + c_1}$$
$$y(x) = -\sqrt{x^2 + c_1}$$

✓ Solution by Mathematica

Time used: 0.081 (sec). Leaf size: 35

```
DSolve[y[x]*y'[x]==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sqrt{x^2 + 2c_1}$$

$$y(x) \rightarrow \sqrt{x^2 + 2c_1}$$

21.3 problem 1(c)

21.3.1 Solving as separable ode	1700
21.3.2 Solving as differentialType ode	1705
21.3.3 Solving as first order ode lie symmetry lookup ode	1709
21.3.4 Solving as exact ode	1713
21.3.5 Maple step by step solution	1717

Internal problem ID [6065]

Internal file name [OUTPUT/5313_Sunday_June_05_2022_03_33_53_PM_69528311/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 190

Problem number: 1(c).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exact", "separable", "differential-Type", "first_order_ode_lie_symmetry_lookup"**

Maple gives the following as the ode type

`[_separable]`

$$y' - \frac{x^2 + x}{y - y^2} = 0$$

21.3.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= -\frac{x(1+x)}{y(y-1)}\end{aligned}$$

Where $f(x) = -x(1+x)$ and $g(y) = \frac{1}{y(y-1)}$. Integrating both sides gives

$$\frac{1}{y(y-1)} dy = -x(1+x) dx$$

$$\int \frac{1}{\frac{1}{y(y-1)}} dy = \int -x(1+x) dx$$

$$\frac{1}{3}y^3 - \frac{1}{2}y^2 = -\frac{1}{2}x^2 - \frac{1}{3}x^3 + c_1$$

Which results in

$$y = \frac{\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{2} + \frac{1}{2}$$

$$y = \frac{\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{4} + \frac{1}{2}$$

$$y = \frac{4\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{4} + \frac{1}{2}$$

$$y = \frac{i\sqrt{3}\left(\frac{\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{2} - \frac{1}{2\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}\right)}{2}$$

$$y = \frac{\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{4} + \frac{1}{2}$$

$$y = \frac{i\sqrt{3}\left(\frac{\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{2} - \frac{1}{2\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}\right)}{2}$$

Summary

The solution(s) found are the following

$$y = \frac{\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{2} \quad (1)$$

$$+ \frac{2\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{1} + \frac{1}{2}$$

$$y = \frac{\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{4} \quad (2)$$

$$- \frac{4\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{1} + \frac{1}{2}$$

$$+ i\sqrt{3} \left(\frac{\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{2} - \frac{2\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{2} \right)$$

$$y = \frac{\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{4} \quad (3)$$

$$- \frac{4\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{1} + \frac{1}{2}$$

$$+ i\sqrt{3} \left(\frac{\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{2} - \frac{2\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{2} \right)$$




Figure 187: Slope field plot

Verification of solutions

y

$$= \frac{\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{2} + \frac{1}{2} \frac{2\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{1}$$

Verified OK.

$y =$

$$= \frac{\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{4} + \frac{1}{2} \frac{4\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{1} + i\sqrt{3} \left(\frac{\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{2} - \frac{2\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{2} \right)$$

Verified OK.

$y =$

$$= \frac{\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{4} + \frac{1}{2} \frac{4\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{1} + i\sqrt{3} \left(\frac{\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{2} - \frac{2\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{2} \right)$$

Verified OK.

21.3.2 Solving as differential Type ode

Writing the ode as

$$y' = \frac{x^2 + x}{y - y^2} \quad (1)$$

Which becomes

$$(y^2 - y) dy = (-x(1 + x)) dx \quad (2)$$

But the RHS is complete differential because

$$(-x(1 + x)) dx = d\left(-\frac{1}{2}x^2 - \frac{1}{3}x^3\right)$$

Hence (2) becomes

$$(y^2 - y) dy = d\left(-\frac{1}{2}x^2 - \frac{1}{3}x^3\right)$$

Integrating both sides gives these solutions

$$y = \frac{\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{2} + \frac{\dots}{2}$$

$$y = -\frac{\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{4} - \frac{\dots}{4}$$

$$y = -\frac{\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{4} - \frac{\dots}{4}$$

Summary

The solution(s) found are the following

$$y = \frac{\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{2} + \frac{1}{2} + c_1 \quad (1)$$

$$y = \frac{\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{4} + \frac{1}{2} + i\sqrt{3} \left(\frac{\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{2} - \frac{2\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{2} \right) + c_1 \quad (2)$$

$$y = \frac{\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{4} + \frac{1}{2} + i\sqrt{3} \left(\frac{\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{2} - \frac{2\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{2} \right) + c_1 \quad (3)$$




Figure 188: Slope field plot

Verification of solutions

y

$$= \frac{\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{2} + \frac{1}{2} + c_1$$

Verified OK.

$y =$

$$= \frac{\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{4} + \frac{1}{2} + i\sqrt{3} \left(\frac{\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{2} - \frac{2\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{2} \right) + c_1$$

Verified OK.

$y =$

$$= \frac{\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{4} + \frac{1}{2} + i\sqrt{3} \left(\frac{\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{2} - \frac{2\left(1 - 4x^3 - 6x^2 + 12c_1 + 2\sqrt{4x^6 + 12x^5 - 24c_1x^3 + 9x^4 - 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 + 6c_1}\right)^{\frac{1}{3}}}{2} \right) + c_1$$

Verified OK.

21.3.3 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = -\frac{x(1+x)}{y(y-1)}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **separable**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 251: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= -\frac{1}{x(1+x)} \\ \eta(x, y) &= 0\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\eta = 0$ then in this special case

$$R = y$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\xi} dx \\ &= \int \frac{1}{-\frac{1}{x(1+x)}} dx\end{aligned}$$

Which results in

$$S = -\frac{1}{2}x^2 - \frac{1}{3}x^3$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y}\tag{2}$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -\frac{x(1+x)}{y(y-1)}$$

Evaluating all the partial derivatives gives

$$\begin{aligned}R_x &= 0 \\R_y &= 1 \\S_x &= -x^2 - x \\S_y &= 0\end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = y(y - 1) \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = R(R - 1)$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \frac{1}{3}R^3 - \frac{1}{2}R^2 + c_1 \quad (4)$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$-\frac{1}{2}x^2 - \frac{1}{3}x^3 = \frac{y^3}{3} - \frac{y^2}{2} + c_1$$

Which simplifies to

$$-\frac{1}{2}x^2 - \frac{1}{3}x^3 = \frac{y^3}{3} - \frac{y^2}{2} + c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -\frac{x(1+x)}{y(y-1)}$ 	$R = y$ $S = -\frac{1}{2}x^2 - \frac{1}{3}x^3$	$\frac{dS}{dR} = R(R-1)$ 

Summary

The solution(s) found are the following

$$-\frac{1}{2}x^2 - \frac{1}{3}x^3 = \frac{y^3}{3} - \frac{y^2}{2} + c_1 \quad (1)$$




Figure 189: Slope field plot

Verification of solutions

$$-\frac{1}{2}x^2 - \frac{1}{3}x^3 = \frac{y^3}{3} - \frac{y^2}{2} + c_1$$

Verified OK.

21.3.4 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial\phi}{\partial x} + \frac{\partial\phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}(-y(y-1)) dy &= (x(1+x)) dx \\ (-x(1+x)) dx + (-y(y-1)) dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -x(1+x) \\ N(x, y) &= -y(y-1)\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-x(1+x)) \\ &= 0\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(-y(y-1)) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\int \frac{\partial \phi}{\partial x} dx = \int M dx$$

$$\int \frac{\partial \phi}{\partial x} dx = \int -x(1+x) dx$$

$$\phi = -\frac{1}{2}x^2 - \frac{1}{3}x^3 + f(y) \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = -y(y-1)$. Therefore equation (4) becomes

$$-y(y-1) = 0 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = -y(y-1)$$

Integrating the above w.r.t y gives

$$\int f'(y) dy = \int (-y(y-1)) dy$$

$$f(y) = -\frac{1}{3}y^3 + \frac{1}{2}y^2 + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\frac{1}{2}x^2 - \frac{1}{3}x^3 - \frac{1}{3}y^3 + \frac{1}{2}y^2 + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\frac{1}{2}x^2 - \frac{1}{3}x^3 - \frac{1}{3}y^3 + \frac{1}{2}y^2$$

Summary

The solution(s) found are the following

$$-\frac{x^3}{3} - \frac{y^3}{3} - \frac{x^2}{2} + \frac{y^2}{2} = c_1 \quad (1)$$




Figure 190: Slope field plot

Verification of solutions

$$-\frac{x^3}{3} - \frac{y^3}{3} - \frac{x^2}{2} + \frac{y^2}{2} = c_1$$

Verified OK.

21.3.5 Maple step by step solution

Let's solve

$$y' - \frac{x^2+x}{y-y^2} = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$(y - y^2) y' = x^2 + x$$

- Integrate both sides with respect to x

$$\int (y - y^2) y' dx = \int (x^2 + x) dx + c_1$$

- Evaluate integral

$$-\frac{y^3}{3} + \frac{y^2}{2} = \frac{1}{3}x^3 + \frac{1}{2}x^2 + c_1$$

- Solve for y

$$y = \frac{\left(1 - 4x^3 - 6x^2 - 12c_1 + 2\sqrt{4x^6 + 12x^5 + 24c_1x^3 + 9x^4 + 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 - 6c_1}\right)^{\frac{1}{3}}}{2} + \frac{\left(1 - 4x^3 - 6x^2 - 12c_1 + 2\sqrt{4x^6 + 12x^5 + 24c_1x^3 + 9x^4 + 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 - 6c_1}\right)^{\frac{1}{3}}}{2}$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
<- separable successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 498

```
dsolve(diff(y(x),x)=(x+x^2)/(y(x)-y(x)^2),y(x), singsol=all)
```

$y(x)$

$$= \frac{\left(1 - 4x^3 - 6x^2 - 12c_1 + 2\sqrt{4x^6 + 12x^5 + 24c_1x^3 + 9x^4 + 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 - 6c_1}\right)^{\frac{1}{3}}}{2}$$

$$+ \frac{1}{2 \left(1 - 4x^3 - 6x^2 - 12c_1 + 2\sqrt{4x^6 + 12x^5 + 24c_1x^3 + 9x^4 + 36c_1x^2 - 2x^3 + 36c_1^2 - 3x^2 - 6c_1}\right)^{\frac{1}{3}}}$$

$y(x) =$

$$= \frac{(1 + i\sqrt{3}) \left(-4x^3 - 6x^2 + 2\sqrt{(2x^3 + 3x^2 + 6c_1)(2x^3 + 3x^2 + 6c_1 - 1)} - 12c_1 + 1\right)^{\frac{2}{3}} - i\sqrt{3} - 2(-4x^3 - 6x^2 + 2\sqrt{(2x^3 + 3x^2 + 6c_1)(2x^3 + 3x^2 + 6c_1 - 1)} - 12c_1 + 1)}{4 \left(-4x^3 - 6x^2 + 2\sqrt{(2x^3 + 3x^2 + 6c_1)(2x^3 + 3x^2 + 6c_1 - 1)} - 12c_1 + 1\right)^{\frac{2}{3}} - i\sqrt{3} - 2(-4x^3 - 6x^2 + 2\sqrt{(2x^3 + 3x^2 + 6c_1)(2x^3 + 3x^2 + 6c_1 - 1)} - 12c_1 + 1)}$$

$y(x)$

$$= \frac{(i\sqrt{3} - 1) \left(-4x^3 - 6x^2 + 2\sqrt{(2x^3 + 3x^2 + 6c_1)(2x^3 + 3x^2 + 6c_1 - 1)} - 12c_1 + 1\right)^{\frac{2}{3}} - i\sqrt{3} + 2(-4x^3 - 6x^2 + 2\sqrt{(2x^3 + 3x^2 + 6c_1)(2x^3 + 3x^2 + 6c_1 - 1)} - 12c_1 + 1)}{4 \left(-4x^3 - 6x^2 + 2\sqrt{(2x^3 + 3x^2 + 6c_1)(2x^3 + 3x^2 + 6c_1 - 1)} - 12c_1 + 1\right)^{\frac{2}{3}} - i\sqrt{3} + 2(-4x^3 - 6x^2 + 2\sqrt{(2x^3 + 3x^2 + 6c_1)(2x^3 + 3x^2 + 6c_1 - 1)} - 12c_1 + 1)}$$

✓ Solution by Mathematica

Time used: 4.147 (sec). Leaf size: 346

```
DSolve[y'[x]==(x+x^2)/(y[x]-y[x]^2),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} \left(\sqrt[3]{-4x^3 - 6x^2 + \sqrt{-1 + (-4x^3 - 6x^2 + 1 + 12c_1)^2 + 1 + 12c_1}} \right. \\ \left. + \frac{1}{\sqrt[3]{-4x^3 - 6x^2 + \sqrt{-1 + (-4x^3 - 6x^2 + 1 + 12c_1)^2 + 1 + 12c_1}} + 1} \right)$$
$$y(x) \rightarrow \frac{1}{8} \left(2i(\sqrt{3} + i) \sqrt[3]{-4x^3 - 6x^2 + \sqrt{-1 + (-4x^3 - 6x^2 + 1 + 12c_1)^2 + 1 + 12c_1}} \right. \\ \left. + \frac{-2 - 2i\sqrt{3}}{\sqrt[3]{-4x^3 - 6x^2 + \sqrt{-1 + (-4x^3 - 6x^2 + 1 + 12c_1)^2 + 1 + 12c_1}} + 4} \right)$$
$$y(x) \rightarrow \frac{1}{8} \left(-2(1 + i\sqrt{3}) \sqrt[3]{-4x^3 - 6x^2 + \sqrt{-1 + (-4x^3 - 6x^2 + 1 + 12c_1)^2 + 1 + 12c_1}} \right. \\ \left. + \frac{2i(\sqrt{3} + i)}{\sqrt[3]{-4x^3 - 6x^2 + \sqrt{-1 + (-4x^3 - 6x^2 + 1 + 12c_1)^2 + 1 + 12c_1}} + 4} \right)$$

21.4 problem 1(d)

21.4.1 Solving as separable ode	1720
21.4.2 Solving as first order special form ID 1 ode	1722
21.4.3 Solving as first order ode lie symmetry lookup ode	1723
21.4.4 Solving as exact ode	1727
21.4.5 Maple step by step solution	1731

Internal problem ID [6066]

Internal file name [OUTPUT/5314_Sunday_June_05_2022_03_33_55_PM_6954108/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 190

Problem number: 1(d).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "separable", "first order special form ID 1", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

`[_separable]`

$$y' - \frac{e^{x-y}}{1 + e^x} = 0$$

21.4.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{e^x e^{-y}}{1 + e^x}\end{aligned}$$

Where $f(x) = \frac{e^x}{1+e^x}$ and $g(y) = e^{-y}$. Integrating both sides gives

$$\frac{1}{e^{-y}} dy = \frac{e^x}{1 + e^x} dx$$

$$\int \frac{1}{e^{-y}} dy = \int \frac{e^x}{1 + e^x} dx$$

$$e^y = \ln(1 + e^x) + c_1$$

Which results in

$$y = \ln(\ln(1 + e^x) + c_1)$$

Summary

The solution(s) found are the following

$$y = \ln(\ln(1 + e^x) + c_1) \tag{1}$$




Figure 191: Slope field plot

Verification of solutions

$$y = \ln(\ln(1 + e^x) + c_1)$$

Verified OK.

21.4.2 Solving as first order special form ID 1 ode

Writing the ode as

$$y' = \frac{e^{x-y}}{1 + e^x} \quad (1)$$

And using the substitution $u = e^y$ then

$$u' = y'e^y$$

The above shows that

$$\begin{aligned} y' &= u'(x) e^{-y} \\ &= \frac{u'(x)}{u} \end{aligned}$$

Substituting this in (1) gives

$$\frac{u'(x)}{u} = \frac{e^x}{(1 + e^x) u}$$

The above simplifies to

$$u'(x) = \frac{e^x}{1 + e^x} \quad (2)$$

Now ode (2) is solved for $u(x)$ Integrating both sides gives

$$\begin{aligned} u(x) &= \int \frac{e^x}{1 + e^x} dx \\ &= \ln(1 + e^x) + c_1 \end{aligned}$$

Substituting the solution found for $u(x)$ in $u = e^y$ gives

$$\begin{aligned} y &= \ln(u(x)) \\ &= \ln(\ln(1 + e^x) + c_1) \\ &= \ln(\ln(1 + e^x) + c_1) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \ln(\ln(1 + e^x) + c_1) \quad (1)$$




Figure 192: Slope field plot

Verification of solutions

$$y = \ln(\ln(1 + e^x) + c_1)$$

Verified OK.

21.4.3 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{e^{x-y}}{1 + e^x}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **separable**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 254: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= (1 + e^x)e^{-x} \\ \eta(x, y) &= 0\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\eta = 0$ then in this special case

$$R = y$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\xi} dx \\ &= \int \frac{1}{(1 + e^x) e^{-x}} dx \end{aligned}$$

Which results in

$$S = \ln(1 + e^x)$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{e^{x-y}}{1 + e^x}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 0 \\ R_y &= 1 \\ S_x &= \frac{e^x}{1 + e^x} \\ S_y &= 0 \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = e^y \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = e^R$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = e^R + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\ln(1 + e^x) = e^y + c_1$$



Which simplifies to

$$\ln(1 + e^x) = e^y + c_1$$

Which gives

$$y = \ln(\ln(1 + e^x) - c_1)$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{e^{x-y}}{1+e^x}$ 	$R = y$ $S = \ln(1 + e^x)$	$\frac{dS}{dR} = e^R$ 

Summary

The solution(s) found are the following

$$y = \ln(\ln(1 + e^x) - c_1) \quad (1)$$




Figure 193: Slope field plot

Verification of solutions

$$y = \ln(\ln(1 + e^x) - c_1)$$

Verified OK.

21.4.4 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}(e^y) dy &= \left(\frac{e^x}{1 + e^x} \right) dx \\ \left(-\frac{e^x}{1 + e^x} \right) dx + (e^y) dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -\frac{e^x}{1 + e^x} \\ N(x, y) &= e^y\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(-\frac{e^x}{1 + e^x} \right) \\ &= 0\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(e^y) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \tag{1}$$

$$\frac{\partial \phi}{\partial y} = N \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\frac{e^x}{1 + e^x} dx \\ \phi &= -\ln(1 + e^x) + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = e^y$. Therefore equation (4) becomes

$$e^y = 0 + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = e^y$$

Integrating the above w.r.t y gives

$$\begin{aligned}\int f'(y) dy &= \int (e^y) dy \\ f(y) &= e^y + c_1\end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\ln(1 + e^x) + e^y + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\ln(1 + e^x) + e^y$$

The solution becomes

$$y = \ln(\ln(1 + e^x) + c_1)$$

Summary

The solution(s) found are the following

$$y = \ln(\ln(1 + e^x) + c_1) \tag{1}$$




Figure 194: Slope field plot

Verification of solutions

$$y = \ln(\ln(1 + e^x) + c_1)$$

Verified OK.

21.4.5 Maple step by step solution

Let's solve

$$y' - \frac{e^{x-y}}{1+e^x} = 0$$

- Highest derivative means the order of the ODE is 1

y'

- Separate variables

$$y'e^y = \frac{e^x}{1+e^x}$$

- Integrate both sides with respect to x

$$\int y'e^y dx = \int \frac{e^x}{1+e^x} dx + c_1$$

- Evaluate integral

$$e^y = \ln(1 + e^x) + c_1$$

- Solve for y

$$y = \ln(\ln(1 + e^x) + c_1)$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
<- separable successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)=exp(x-y(x))/(1+exp(x)),y(x), singsol=all)
```

$$y(x) = \ln(\ln(e^x + 1) + c_1)$$

✓ Solution by Mathematica

Time used: 0.465 (sec). Leaf size: 15

```
DSolve[y'[x]==Exp[x-y[x]]/(1+Exp[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \log(\log(e^x + 1) + c_1)$$

21.5 problem 1(e)

21.5.1 Solving as separable ode	1733
21.5.2 Solving as first order ode lie symmetry lookup ode	1735
21.5.3 Solving as exact ode	1739
21.5.4 Solving as riccati ode	1743
21.5.5 Maple step by step solution	1746

Internal problem ID [6067]

Internal file name [OUTPUT/5315_Sunday_June_05_2022_03_33_56_PM_35202324/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 190

Problem number: 1(e).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "riccati", "separable", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_separable]

$$y' - y^2 x^2 = -4x^2$$

21.5.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= x^2(y^2 - 4)\end{aligned}$$

Where $f(x) = x^2$ and $g(y) = y^2 - 4$. Integrating both sides gives

$$\begin{aligned}\frac{1}{y^2 - 4} dy &= x^2 dx \\ \int \frac{1}{y^2 - 4} dy &= \int x^2 dx\end{aligned}$$

$$-\frac{\ln(y+2)}{4} + \frac{\ln(y-2)}{4} = \frac{x^3}{3} + c_1$$

The above can be written as

$$\begin{aligned} \left(-\frac{1}{4}\right) (\ln(y+2) - \ln(y-2)) &= \frac{x^3}{3} + 2c_1 \\ \ln(y+2) - \ln(y-2) &= (-4) \left(\frac{x^3}{3} + 2c_1\right) \\ &= -\frac{4x^3}{3} - 8c_1 \end{aligned}$$

Raising both side to exponential gives

$$e^{\ln(y+2) - \ln(y-2)} = e^{-\frac{4x^3}{3} - 8c_1}$$

Which simplifies to

$$\begin{aligned} \frac{y+2}{y-2} &= -4c_1 e^{-\frac{4x^3}{3}} \\ &= c_2 e^{-\frac{4x^3}{3}} \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{2c_2 e^{-\frac{4x^3}{3}} + 2}{-1 + c_2 e^{-\frac{4x^3}{3}}} \quad (1)$$




Figure 195: Slope field plot

Verification of solutions

$$y = \frac{2c_2 e^{-\frac{4x^3}{3}} + 2}{-1 + c_2 e^{-\frac{4x^3}{3}}}$$

Verified OK.

21.5.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = y^2 x^2 - 4x^2$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **separable**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 257: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= \frac{1}{x^2} \\ \eta(x, y) &= 0\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\eta = 0$ then in this special case

$$R = y$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\xi} dx \\ &= \int \frac{1}{x^2} dx \end{aligned}$$

Which results in

$$S = \frac{x^3}{3}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = y^2x^2 - 4x^2$$

Evaluating all the partial derivatives gives

$$R_x = 0$$

$$R_y = 1$$

$$S_x = x^2$$

$$S_y = 0$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \frac{1}{y^2 - 4} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \frac{1}{R^2 - 4}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by

integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = -\frac{\ln(R+2)}{4} + \frac{\ln(R-2)}{4} + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{x^3}{3} = -\frac{\ln(2+y)}{4} + \frac{\ln(y-2)}{4} + c_1$$



Which simplifies to

$$\frac{x^3}{3} = -\frac{\ln(2+y)}{4} + \frac{\ln(y-2)}{4} + c_1$$

Which gives

$$y = \frac{2 + 2e^{-\frac{4x^3}{3} + 4c_1}}{e^{-\frac{4x^3}{3} + 4c_1} - 1}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = y^2x^2 - 4x^2$ 	$R = y$ $S = \frac{x^3}{3}$	$\frac{dS}{dR} = \frac{1}{R^2-4}$ 

Summary

The solution(s) found are the following

$$y = \frac{2 + 2e^{-\frac{4x^3}{3} + 4c_1}}{e^{-\frac{4x^3}{3} + 4c_1} - 1} \quad (1)$$




Figure 196: Slope field plot

Verification of solutions

$$y = \frac{2 + 2e^{-\frac{4x^3}{3} + 4c_1}}{e^{-\frac{4x^3}{3} + 4c_1} - 1}$$

Verified OK.

21.5.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial\phi}{\partial x} + \frac{\partial\phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}\left(\frac{1}{y^2 - 4}\right) dy &= (x^2) dx \\ (-x^2) dx + \left(\frac{1}{y^2 - 4}\right) dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -x^2 \\ N(x, y) &= \frac{1}{y^2 - 4}\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-x^2) \\ &= 0\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{1}{y^2 - 4} \right) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -x^2 dx \\ \phi &= -\frac{x^3}{3} + f(y)\end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{1}{y^2 - 4}$. Therefore equation (4) becomes

$$\frac{1}{y^2 - 4} = 0 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = \frac{1}{y^2 - 4}$$

Integrating the above w.r.t y gives

$$\int f'(y) dy = \int \left(\frac{1}{y^2 - 4} \right) dy$$
$$f(y) = -\frac{\ln(y+2)}{4} + \frac{\ln(y-2)}{4} + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\frac{x^3}{3} - \frac{\ln(y+2)}{4} + \frac{\ln(y-2)}{4} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\frac{x^3}{3} - \frac{\ln(y+2)}{4} + \frac{\ln(y-2)}{4}$$

The solution becomes

$$y = -\frac{2\left(e^{\frac{4x^3}{3} + 4c_1} + 1\right)}{-1 + e^{\frac{4x^3}{3} + 4c_1}}$$

Summary

The solution(s) found are the following

$$y = -\frac{2\left(e^{\frac{4x^3}{3} + 4c_1} + 1\right)}{-1 + e^{\frac{4x^3}{3} + 4c_1}} \quad (1)$$




Figure 197: Slope field plot

Verification of solutions

$$y = -\frac{2\left(e^{\frac{4x^3}{3}+4c_1} + 1\right)}{-1 + e^{\frac{4x^3}{3}+4c_1}}$$

Verified OK.

21.5.4 Solving as riccati ode

In canonical form the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= y^2 x^2 - 4x^2 \end{aligned}$$

This is a Riccati ODE. Comparing the ODE to solve

$$y' = y^2 x^2 - 4x^2$$

With Riccati ODE standard form

$$y' = f_0(x) + f_1(x)y + f_2(x)y^2$$

Shows that $f_0(x) = -4x^2$, $f_1(x) = 0$ and $f_2(x) = x^2$. Let

$$\begin{aligned} y &= \frac{-u'}{f_2 u} \\ &= \frac{-u'}{x^2 u} \end{aligned} \tag{1}$$

Using the above substitution in the given ODE results (after some simplification) in a second order ODE to solve for $u(x)$ which is

$$f_2 u''(x) - (f_2' + f_1 f_2) u'(x) + f_2^2 f_0 u(x) = 0 \tag{2}$$

But

$$\begin{aligned} f_2' &= 2x \\ f_1 f_2 &= 0 \\ f_2^2 f_0 &= -4x^6 \end{aligned}$$

Substituting the above terms back in equation (2) gives

$$x^2 u''(x) - 2x u'(x) - 4x^6 u(x) = 0$$

Solving the above ODE (this ode solved using Maple, not this program), gives

$$u(x) = c_1 \sinh\left(\frac{2x^3}{3}\right) + c_2 \cosh\left(\frac{2x^3}{3}\right)$$

The above shows that

$$u'(x) = 2x^2 \left(c_1 \cosh\left(\frac{2x^3}{3}\right) + c_2 \sinh\left(\frac{2x^3}{3}\right) \right)$$

Using the above in (1) gives the solution

$$y = -\frac{2 \left(c_1 \cosh\left(\frac{2x^3}{3}\right) + c_2 \sinh\left(\frac{2x^3}{3}\right) \right)}{c_1 \sinh\left(\frac{2x^3}{3}\right) + c_2 \cosh\left(\frac{2x^3}{3}\right)}$$

Dividing both numerator and denominator by c_1 gives, after renaming the constant $\frac{c_2}{c_1} = c_3$ the following solution

$$y = \frac{-2c_3 \cosh\left(\frac{2x^3}{3}\right) - 2 \sinh\left(\frac{2x^3}{3}\right)}{c_3 \sinh\left(\frac{2x^3}{3}\right) + \cosh\left(\frac{2x^3}{3}\right)}$$

Summary

The solution(s) found are the following

$$y = \frac{-2c_3 \cosh\left(\frac{2x^3}{3}\right) - 2 \sinh\left(\frac{2x^3}{3}\right)}{c_3 \sinh\left(\frac{2x^3}{3}\right) + \cosh\left(\frac{2x^3}{3}\right)} \quad (1)$$




Figure 198: Slope field plot

Verification of solutions

$$y = \frac{-2c_3 \cosh\left(\frac{2x^3}{3}\right) - 2 \sinh\left(\frac{2x^3}{3}\right)}{c_3 \sinh\left(\frac{2x^3}{3}\right) + \cosh\left(\frac{2x^3}{3}\right)}$$

Verified OK.

21.5.5 Maple step by step solution

Let's solve

$$y' - y^2x^2 = -4x^2$$

- Highest derivative means the order of the ODE is 1

y'

- Separate variables

$$\frac{y'}{(y-2)(2+y)} = x^2$$

- Integrate both sides with respect to x

$$\int \frac{y'}{(y-2)(2+y)} dx = \int x^2 dx + c_1$$

- Evaluate integral

$$-\frac{\ln(2+y)}{4} + \frac{\ln(y-2)}{4} = \frac{x^3}{3} + c_1$$

- Solve for y

$$y = -\frac{2\left(e^{\frac{4x^3}{3}+4c_1}+1\right)}{-1+e^{\frac{4x^3}{3}+4c_1}}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
<- separable successful`
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 28

```
dsolve(diff(y(x),x)=x^2*y(x)^2-4*x^2,y(x), singsol=all)
```

$$y(x) = \frac{-2 - 2 e^{\frac{4x^3}{3}} c_1}{e^{\frac{4x^3}{3}} c_1 - 1}$$

✓ Solution by Mathematica

Time used: 0.258 (sec). Leaf size: 52

```
DSolve[y'[x]==x^2*y[x]^2-4*x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2 - 2e^{\frac{4x^3}{3} + 4c_1}}{1 + e^{\frac{4x^3}{3} + 4c_1}}$$

$$y(x) \rightarrow -2$$

$$y(x) \rightarrow 2$$

21.6 problem 2(a)

21.6.1 Existence and uniqueness analysis	1748
21.6.2 Solving as quadrature ode	1749
21.6.3 Maple step by step solution	1750

Internal problem ID [6068]

Internal file name [OUTPUT/5316_Sunday_June_05_2022_03_33_58_PM_44743446/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 190

Problem number: 2(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$y' - y^2 = 0$$

With initial conditions

$$[y(x_0) = y_0]$$

21.6.1 Existence and uniqueness analysis

This is non linear first order ODE. In canonical form it is written as

$$\begin{aligned} y' &= f(x, y) \\ &= y^2 \end{aligned}$$

The y domain of $f(x, y)$ when $x = x_0$ is

$$\{-\infty < y < \infty\}$$

But the point $y_0 = y_0$ is not inside this domain. Hence existence and uniqueness theorem does not apply. There could be infinite number of solutions, or one solution or no solution at all.

21.6.2 Solving as quadrature ode

Integrating both sides gives

$$\int \frac{1}{y^2} dy = x + c_1$$
$$-\frac{1}{y} = x + c_1$$

Solving for y gives these solutions

$$y_1 = -\frac{1}{x + c_1}$$

Initial conditions are used to solve for c_1 . Substituting $x = x_0$ and $y = y_0$ in the above solution gives an equation to solve for the constant of integration.

$$y_0 = -\frac{1}{x_0 + c_1}$$

$$c_1 = -\frac{y_0 x_0 + 1}{y_0}$$

Substituting c_1 found above in the general solution gives

$$y = -\frac{1}{x - \frac{y_0 x_0 + 1}{y_0}}$$

Summary

The solution(s) found are the following

$$y = -\frac{1}{x - \frac{y_0 x_0 + 1}{y_0}} \tag{1}$$

Verification of solutions

$$y = -\frac{1}{x - \frac{y_0 x_0 + 1}{y_0}}$$

Verified OK.

21.6.3 Maple step by step solution

Let's solve

$$[y' - y^2 = 0, y(x_0) = y_0]$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y^2} = 1$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y^2} dx = \int 1 dx + c_1$$

- Evaluate integral

$$-\frac{1}{y} = x + c_1$$

- Solve for y

$$y = -\frac{1}{x+c_1}$$

- Use initial condition $y(x_0) = y_0$

$$y_0 = -\frac{1}{x_0+c_1}$$

- Solve for c_1

$$c_1 = -\frac{y_0 x_0 + 1}{y_0}$$

- Substitute $c_1 = -\frac{y_0 x_0 + 1}{y_0}$ into general solution and simplify

$$y = -\frac{y_0}{-1+(x-x_0)y_0}$$

- Solution to the IVP

$$y = -\frac{y_0}{-1+(x-x_0)y_0}$$

Maple trace

```
`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
<- Bernoulli successful`
```

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 18

```
dsolve([diff(y(x),x)=y(x)^2,y(x__0) = y__0],y(x), singsol=all)
```

$$y(x) = -\frac{y_0}{-1 + (x - x_0) y_0}$$

✓ Solution by Mathematica

Time used: 0.028 (sec). Leaf size: 16

```
DSolve[{y'[x]==x^2*y[x],{y[x0]==y0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow y_0 e^{x^2(x-x_0)}$$

21.7 problem 3(a)

21.7.1 Existence and uniqueness analysis	1752
21.7.2 Solving as quadrature ode	1753
21.7.3 Maple step by step solution	1753

Internal problem ID [6069]

Internal file name [OUTPUT/5317_Sunday_June_05_2022_03_33_59_PM_16751227/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 190

Problem number: 3(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$y' - 2\sqrt{y} = 0$$

With initial conditions

$$[y(x_0) = y_0]$$

21.7.1 Existence and uniqueness analysis

This is non linear first order ODE. In canonical form it is written as

$$\begin{aligned}y' &= f(x, y) \\ &= 2\sqrt{y}\end{aligned}$$

The y domain of $f(x, y)$ when $x = x_0$ is

$$\{0 \leq y\}$$

But the point $y_0 = y_0$ is not inside this domain. Hence existence and uniqueness theorem does not apply. There could be infinite number of solutions, or one solution or no solution at all.

21.7.2 Solving as quadrature ode

Integrating both sides gives

$$\int \frac{1}{2\sqrt{y}} dy = \int dx$$
$$\sqrt{y} = x + c_1$$

Initial conditions are used to solve for c_1 . Substituting $x = x_0$ and $y = y_0$ in the above solution gives an equation to solve for the constant of integration.

$$\sqrt{y_0} = x_0 + c_1$$

$$c_1 = -x_0 + \sqrt{y_0}$$

Substituting c_1 found above in the general solution gives

$$\sqrt{y} = x - x_0 + \sqrt{y_0}$$

Solving for y from the above gives

$$y = (2x - 2x_0) \sqrt{y_0} + x^2 - 2xx_0 + x_0^2 + y_0$$

Summary

The solution(s) found are the following

$$y = (2x - 2x_0) \sqrt{y_0} + x^2 - 2xx_0 + x_0^2 + y_0 \quad (1)$$

Verification of solutions

$$y = (2x - 2x_0) \sqrt{y_0} + x^2 - 2xx_0 + x_0^2 + y_0$$

Verified OK.

21.7.3 Maple step by step solution

Let's solve

$$[y' - 2\sqrt{y} = 0, y(x_0) = y_0]$$

- Highest derivative means the order of the ODE is 1
- y'
- Separate variables

$$\frac{y'}{\sqrt{y}} = 2$$

- Integrate both sides with respect to x

$$\int \frac{y'}{\sqrt{y}} dx = \int 2dx + c_1$$

- Evaluate integral

$$2\sqrt{y} = c_1 + 2x$$

- Solve for y

$$y = \frac{1}{4}c_1^2 + c_1x + x^2$$

- Use initial condition $y(x_0) = y_0$

$$y_0 = \frac{1}{4}c_1^2 + c_1x_0 + x_0^2$$

- Solve for c_1

$$c_1 = (2\sqrt{y_0} - 2x_0, -2x_0 - 2\sqrt{y_0})$$

- Substitute $c_1 = (2\sqrt{y_0} - 2x_0, -2x_0 - 2\sqrt{y_0})$ into general solution and simplify

$$y = (2x - 2x_0)\sqrt{y_0} + x^2 - 2xx_0 + x_0^2 + y_0$$

- Solution to the IVP

$$y = (2x - 2x_0)\sqrt{y_0} + x^2 - 2xx_0 + x_0^2 + y_0$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
<- Bernoulli successful`

```

✓ Solution by Maple

Time used: 0.156 (sec). Leaf size: 28

```
dsolve([diff(y(x),x)=2*sqrt(y(x)),y(x__0) = y__0],y(x), singsol=all)
```

$$y(x) = (2x - 2x_0)\sqrt{y_0} + x^2 - 2xx_0 + x_0^2 + y_0$$

✓ Solution by Mathematica

Time used: 0.108 (sec). Leaf size: 33

```
DSolve[{y'[x]==2*Sqrt[y[x]],{y[x0]==y0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow (x - x_0 + \sqrt{y_0})^2$$

$$y(x) \rightarrow (-x + x_0 + \sqrt{y_0})^2$$

21.8 problem 3(b)

21.8.1 Existence and uniqueness analysis	1756
21.8.2 Solving as quadrature ode	1757
21.8.3 Maple step by step solution	1758

Internal problem ID [6070]

Internal file name [OUTPUT/5318_Sunday_June_05_2022_03_34_01_PM_54625177/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 190

Problem number: 3(b).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$y' - 2\sqrt{y} = 0$$

With initial conditions

$$[y(x_0) = 0]$$

21.8.1 Existence and uniqueness analysis

This is non linear first order ODE. In canonical form it is written as

$$\begin{aligned}y' &= f(x, y) \\ &= 2\sqrt{y}\end{aligned}$$

The y domain of $f(x, y)$ when $x = x_0$ is

$$\{0 \leq y\}$$

And the point $y_0 = 0$ is inside this domain. Now we will look at the continuity of

$$\begin{aligned}\frac{\partial f}{\partial y} &= \frac{\partial}{\partial y}(2\sqrt{y}) \\ &= \frac{1}{\sqrt{y}}\end{aligned}$$

The y domain of $\frac{\partial f}{\partial y}$ when $x = x_0$ is

$$\{0 < y\}$$

But the point $y_0 = 0$ is not inside this domain. Hence existence and uniqueness theorem does not apply. Solution exists but no guarantee that unique solution exists.

21.8.2 Solving as quadrature ode

Integrating both sides gives

$$\int \frac{1}{2\sqrt{y}} dy = \int dx$$
$$\sqrt{y} = x + c_1$$

Initial conditions are used to solve for c_1 . Substituting $x = x_0$ and $y = 0$ in the above solution gives an equation to solve for the constant of integration.

$$0 = x_0 + c_1$$

$$c_1 = -x_0$$

Substituting c_1 found above in the general solution gives

$$\sqrt{y} = x - x_0$$

Solving for y from the above gives

$$y = (x - x_0)^2$$

Summary

The solution(s) found are the following

$$y = (x - x_0)^2 \tag{1}$$

Verification of solutions

$$y = (x - x_0)^2$$

Verified OK.

21.8.3 Maple step by step solution

Let's solve

$$[y' - 2\sqrt{y} = 0, y(x_0) = 0]$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{\sqrt{y}} = 2$$

- Integrate both sides with respect to x

$$\int \frac{y'}{\sqrt{y}} dx = \int 2 dx + c_1$$

- Evaluate integral

$$2\sqrt{y} = c_1 + 2x$$

- Solve for y

$$y = \frac{1}{4}c_1^2 + c_1x + x^2$$

- Use initial condition $y(x_0) = 0$

$$0 = \frac{1}{4}c_1^2 + c_1x_0 + x_0^2$$

- Solve for c_1

$$c_1 = (-2x_0, -2x_0)$$

- Substitute $c_1 = (-2x_0, -2x_0)$ into general solution and simplify

$$y = (x - x_0)^2$$

- Solution to the IVP

$$y = (x - x_0)^2$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
<- Bernoulli successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 5

```
dsolve([diff(y(x),x)=2*sqrt(y(x)),y(x__0) = 0],y(x), singsol=all)
```

$$y(x) = 0$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 6

```
DSolve[{y'[x]==2*Sqrt[y[x]],{y[x0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 0$$

21.9 problem 4(a)

21.9.1 Solving as homogeneous ode 1760

Internal problem ID [6071]

Internal file name [OUTPUT/5319_Sunday_June_05_2022_03_34_04_PM_26636281/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 190

Problem number: 4(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

```
[[_homogeneous, `class A`], _rational, [_Abel, `2nd type`, `class A`]]
```

$$y' - \frac{x+y}{x-y} = 0$$

21.9.1 Solving as homogeneous ode

In canonical form, the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= -\frac{x+y}{-x+y} \end{aligned} \tag{1}$$

An ode of the form $y' = \frac{M(x,y)}{N(x,y)}$ is called homogeneous if the functions $M(x, y)$ and $N(x, y)$ are both homogeneous functions and of the same order. Recall that a function $f(x, y)$ is homogeneous of order n if

$$f(t^n x, t^n y) = t^n f(x, y)$$

In this case, it can be seen that both $M = x + y$ and $N = x - y$ are both homogeneous and of the same order $n = 1$. Therefore this is a homogeneous ode. Since this ode is homogeneous, it is converted to separable ODE using the substitution $u = \frac{y}{x}$, or $y = ux$. Hence

$$\frac{dy}{dx} = \frac{du}{dx}x + u$$

Applying the transformation $y = ux$ to the above ODE in (1) gives

$$\frac{du}{dx}x + u = \frac{-u - 1}{u - 1}$$

$$\frac{du}{dx} = \frac{\frac{-u(x)-1}{u(x)-1} - u(x)}{x}$$

Or

$$u'(x) - \frac{\frac{-u(x)-1}{u(x)-1} - u(x)}{x} = 0$$

Or

$$u'(x) xu(x) - u'(x) x + u(x)^2 + 1 = 0$$

Or

$$x(u(x) - 1) u'(x) + u(x)^2 + 1 = 0$$

Which is now solved as separable in $u(x)$. Which is now solved in $u(x)$. In canonical form the ODE is

$$u' = F(x, u)$$

$$= f(x)g(u)$$

$$= -\frac{u^2 + 1}{x(u - 1)}$$

Where $f(x) = -\frac{1}{x}$ and $g(u) = \frac{u^2+1}{u-1}$. Integrating both sides gives

$$\frac{1}{\frac{u^2+1}{u-1}} du = -\frac{1}{x} dx$$

$$\int \frac{1}{\frac{u^2+1}{u-1}} du = \int -\frac{1}{x} dx$$

$$\frac{\ln(u^2 + 1)}{2} - \arctan(u) = -\ln(x) + c_2$$

The solution is

$$\frac{\ln(u(x)^2 + 1)}{2} - \arctan(u(x)) + \ln(x) - c_2 = 0$$

Now u in the above solution is replaced back by y using $u = \frac{y}{x}$ which results in the solution

$$\frac{\ln\left(\frac{y^2}{x^2} + 1\right)}{2} - \arctan\left(\frac{y}{x}\right) + \ln(x) - c_2 = 0$$

Summary

The solution(s) found are the following

$$\frac{\ln\left(\frac{y^2}{x^2} + 1\right)}{2} - \arctan\left(\frac{y}{x}\right) + \ln(x) - c_2 = 0 \quad (1)$$




Figure 199: Slope field plot

Verification of solutions

$$\frac{\ln\left(\frac{y^2}{x^2} + 1\right)}{2} - \arctan\left(\frac{y}{x}\right) + \ln(x) - c_2 = 0$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying homogeneous D  
<- homogeneous successful`
```

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 24

```
dsolve(diff(y(x),x)=(x+y(x))/(x-y(x)),y(x), singsol=all)
```

$$y(x) = \tan \left(\text{RootOf} \left(-2_Z + \ln \left(\sec \left(_Z \right)^2 \right) + 2 \ln (x) + 2c_1 \right) \right) x$$

✓ Solution by Mathematica

Time used: 0.034 (sec). Leaf size: 36

```
DSolve[y'[x]==(x+y[x])/(x-y[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[\frac{1}{2} \log \left(\frac{y(x)^2}{x^2} + 1 \right) - \arctan \left(\frac{y(x)}{x} \right) = -\log(x) + c_1, y(x) \right]$$

21.10 problem 4(b)

21.10.1 Solving as homogeneous ode 1764

Internal problem ID [6072]

Internal file name [OUTPUT/5320_Sunday_June_05_2022_03_34_05_PM_25195445/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 190

Problem number: 4(b).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

```
[[_homogeneous, `class A`], _rational, [_Abel, `2nd type`, `class B`]]
```

$$y' - \frac{y^2}{xy + x^2} = 0$$

21.10.1 Solving as homogeneous ode

In canonical form, the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= \frac{y^2}{x(x+y)} \end{aligned} \tag{1}$$

An ode of the form $y' = \frac{M(x,y)}{N(x,y)}$ is called homogeneous if the functions $M(x, y)$ and $N(x, y)$ are both homogeneous functions and of the same order. Recall that a function $f(x, y)$ is homogeneous of order n if

$$f(t^n x, t^n y) = t^n f(x, y)$$

In this case, it can be seen that both $M = y^2$ and $N = x(x + y)$ are both homogeneous and of the same order $n = 2$. Therefore this is a homogeneous ode. Since this ode is homogeneous, it is converted to separable ODE using the substitution $u = \frac{y}{x}$, or $y = ux$. Hence

$$\frac{dy}{dx} = \frac{du}{dx}x + u$$

Applying the transformation $y = ux$ to the above ODE in (1) gives

$$\begin{aligned}\frac{du}{dx}x + u &= \frac{u^2}{u+1} \\ \frac{du}{dx} &= \frac{\frac{u(x)^2}{u(x)+1} - u(x)}{x}\end{aligned}$$

Or

$$u'(x) - \frac{\frac{u(x)^2}{u(x)+1} - u(x)}{x} = 0$$

Or

$$u'(x) xu(x) + u'(x) x + u(x) = 0$$

Or

$$(u(x) + 1) xu'(x) + u(x) = 0$$

Which is now solved as separable in $u(x)$. Which is now solved in $u(x)$. In canonical form the ODE is

$$\begin{aligned}u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{u}{(u+1)x}\end{aligned}$$

Where $f(x) = -\frac{1}{x}$ and $g(u) = \frac{u}{u+1}$. Integrating both sides gives

$$\begin{aligned}\frac{1}{\frac{u}{u+1}} du &= -\frac{1}{x} dx \\ \int \frac{1}{\frac{u}{u+1}} du &= \int -\frac{1}{x} dx \\ u + \ln(u) &= -\ln(x) + c_2\end{aligned}$$

The solution is

$$u(x) + \ln(u(x)) + \ln(x) - c_2 = 0$$

Now u in the above solution is replaced back by y using $u = \frac{y}{x}$ which results in the solution

$$\frac{y}{x} + \ln\left(\frac{y}{x}\right) + \ln(x) - c_2 = 0$$

Summary

The solution(s) found are the following

$$\frac{y}{x} + \ln\left(\frac{y}{x}\right) + \ln(x) - c_2 = 0 \quad (1)$$




Figure 200: Slope field plot

Verification of solutions

$$\frac{y}{x} + \ln\left(\frac{y}{x}\right) + \ln(x) - c_2 = 0$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying homogeneous D  
<- homogeneous successful`
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 15

```
dsolve(diff(y(x),x)=y(x)^2/(x*y(x)+x^2),y(x), singsol=all)
```

$$y(x) = x \operatorname{LambertW}\left(\frac{e^{-c_1}}{x}\right)$$

✓ Solution by Mathematica

Time used: 2.317 (sec). Leaf size: 21

```
DSolve[y'[x]==y[x]^2/(x*y[x]+x^2),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow xW\left(\frac{e^{c_1}}{x}\right)$$
$$y(x) \rightarrow 0$$

21.11 problem 4(c)

21.11.1 Solving as homogeneous ode 1768

Internal problem ID [6073]

Internal file name [OUTPUT/5321_Sunday_June_05_2022_03_34_07_PM_83140634/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 190

Problem number: 4(c).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

```
[[_homogeneous, `class A`], _rational, _Riccati]
```

$$y' - \frac{x^2 + xy + y^2}{x^2} = 0$$

21.11.1 Solving as homogeneous ode

In canonical form, the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= \frac{x^2 + xy + y^2}{x^2} \end{aligned} \tag{1}$$

An ode of the form $y' = \frac{M(x,y)}{N(x,y)}$ is called homogeneous if the functions $M(x, y)$ and $N(x, y)$ are both homogeneous functions and of the same order. Recall that a function $f(x, y)$ is homogeneous of order n if

$$f(t^n x, t^n y) = t^n f(x, y)$$

In this case, it can be seen that both $M = x^2 + xy + y^2$ and $N = x^2$ are both homogeneous and of the same order $n = 2$. Therefore this is a homogeneous ode. Since this ode is homogeneous, it is converted to separable ODE using the substitution $u = \frac{y}{x}$, or $y = ux$. Hence

$$\frac{dy}{dx} = \frac{du}{dx}x + u$$

Applying the transformation $y = ux$ to the above ODE in (1) gives

$$\begin{aligned}\frac{du}{dx}x + u &= u^2 + u + 1 \\ \frac{du}{dx} &= \frac{u(x)^2 + 1}{x}\end{aligned}$$

Or

$$u'(x) - \frac{u(x)^2 + 1}{x} = 0$$

Or

$$u'(x)x - u(x)^2 - 1 = 0$$

Which is now solved as separable in $u(x)$. Which is now solved in $u(x)$. In canonical form the ODE is

$$\begin{aligned}u' &= F(x, u) \\ &= f(x)g(u) \\ &= \frac{u^2 + 1}{x}\end{aligned}$$

Where $f(x) = \frac{1}{x}$ and $g(u) = u^2 + 1$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u^2 + 1} du &= \frac{1}{x} dx \\ \int \frac{1}{u^2 + 1} du &= \int \frac{1}{x} dx \\ \arctan(u) &= \ln(x) + c_2\end{aligned}$$

The solution is

$$\arctan(u(x)) - \ln(x) - c_2 = 0$$

Now u in the above solution is replaced back by y using $u = \frac{y}{x}$ which results in the solution

$$\arctan\left(\frac{y}{x}\right) - \ln(x) - c_2 = 0$$

Summary

The solution(s) found are the following

$$\arctan\left(\frac{y}{x}\right) - \ln(x) - c_2 = 0 \quad (1)$$




Figure 201: Slope field plot

Verification of solutions

$$\arctan\left(\frac{y}{x}\right) - \ln(x) - c_2 = 0$$

Verified OK.

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying homogeneous D
<- homogeneous successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve(diff(y(x),x)=(x^2+x*y(x)+y(x)^2)/x^2,y(x), singsol=all)
```

$$y(x) = \tan(\ln(x) + c_1) x$$

✓ Solution by Mathematica

Time used: 0.188 (sec). Leaf size: 13

```
DSolve[y'[x]==(x^2+x*y[x]+y[x]^2)/x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x \tan(\log(x) + c_1)$$

21.12 problem 4(d)

21.12.1 Solving as homogeneous ode 1772

Internal problem ID [6074]

Internal file name [OUTPUT/5322_Sunday_June_05_2022_03_34_09_PM_77661497/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 190

Problem number: 4(d).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

```
[[_homogeneous, `class A`], _dAlembert]
```

$$y' - \frac{y + x e^{-\frac{2y}{x}}}{x} = 0$$

21.12.1 Solving as homogeneous ode

In canonical form, the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= \frac{y + x e^{-\frac{2y}{x}}}{x} \end{aligned} \quad (1)$$

An ode of the form $y' = \frac{M(x,y)}{N(x,y)}$ is called homogeneous if the functions $M(x, y)$ and $N(x, y)$ are both homogeneous functions and of the same order. Recall that a function $f(x, y)$ is homogeneous of order n if

$$f(t^n x, t^n y) = t^n f(x, y)$$

In this case, it can be seen that both $M = y + x e^{-\frac{2y}{x}}$ and $N = x$ are both homogeneous and of the same order $n = 1$. Therefore this is a homogeneous ode. Since this ode is homogeneous, it is converted to separable ODE using the substitution $u = \frac{y}{x}$, or $y = ux$.

Hence

$$\frac{dy}{dx} = \frac{du}{dx}x + u$$

Applying the transformation $y = ux$ to the above ODE in (1) gives

$$\begin{aligned}\frac{du}{dx}x + u &= u + e^{-2u} \\ \frac{du}{dx} &= \frac{e^{-2u(x)}}{x}\end{aligned}$$

Or

$$u'(x) - \frac{e^{-2u(x)}}{x} = 0$$

Or

$$u'(x) e^{2u(x)} x - 1 = 0$$

Or

$$u'(x) e^{2u(x)} x - 1 = 0$$

Which is now solved as separable in $u(x)$. Which is now solved in $u(x)$. In canonical form the ODE is

$$\begin{aligned}u' &= F(x, u) \\ &= f(x)g(u) \\ &= \frac{e^{-2u}}{x}\end{aligned}$$

Where $f(x) = \frac{1}{x}$ and $g(u) = e^{-2u}$. Integrating both sides gives

$$\begin{aligned}\frac{1}{e^{-2u}} du &= \frac{1}{x} dx \\ \int \frac{1}{e^{-2u}} du &= \int \frac{1}{x} dx \\ \frac{e^{2u}}{2} &= \ln(x) + c_2\end{aligned}$$

The solution is

$$\frac{e^{2u(x)}}{2} - \ln(x) - c_2 = 0$$

Now u in the above solution is replaced back by y using $u = \frac{y}{x}$ which results in the solution

$$\frac{e^{\frac{2y}{x}}}{2} - \ln(x) - c_2 = 0$$

Summary

The solution(s) found are the following

$$\frac{e^{\frac{2y}{x}}}{2} - \ln(x) - c_2 = 0 \quad (1)$$




Figure 202: Slope field plot

Verification of solutions

$$\frac{e^{\frac{2y}{x}}}{2} - \ln(x) - c_2 = 0$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying homogeneous D  
<- homogeneous successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
dsolve(diff(y(x),x)=(y(x)+x*exp(-2*y(x)/x))/x,y(x), singsol=all)
```

$$y(x) = \frac{(\ln(2) + \ln(\ln(x) + c_1))x}{2}$$

✓ Solution by Mathematica

Time used: 0.412 (sec). Leaf size: 18

```
DSolve[y'[x]==(y[x]+x*Exp[-2*y[x]/x])/x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}x \log(2(\log(x) + c_1))$$

21.13 problem 5(a)

21.13.1 Solving as polynomial ode 1776

Internal problem ID [6075]

Internal file name [OUTPUT/5323_Sunday_June_05_2022_03_34_10_PM_89530852/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 190

Problem number: 5(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

```
[[_homogeneous, `class C`], _rational, [_Abel, `2nd type`, `class A`]]
```

$$y' - \frac{x - y + 2}{-1 + y + x} = 0$$

21.13.1 Solving as polynomial ode

This is ODE of type polynomial. Where the RHS of the ode is ratio of equations of two lines. Writing the ODE in the form

$$y' = \frac{a_1x + b_1y + c_1}{a_2x + b_2y + c_3}$$

Where $a_1 = 1, b_1 = -1, c_1 = 2, a_2 = 1, b_2 = 1, c_2 = -1$. There are now two possible solution methods. The first case is when the two lines $a_1x + b_1y + c_1, a_2x + b_2y + c_3$ are not parallel and the second case is if they are parallel. If they are not parallel, then the transformation $X = x - x_0, Y = y - y_0$ converts the ODE to a homogeneous ODE. The values x_0, y_0 have to be determined. If they are parallel then a transformation $U(x) = a_1x + b_1y$ converts the given ODE in y to a separable ODE in $U(x)$. The first case is when $\frac{a_1}{b_1} \neq \frac{a_2}{b_2}$ and the second case when $\frac{a_1}{b_1} = \frac{a_2}{b_2}$. From the above we see that $\frac{a_1}{b_1} \neq \frac{a_2}{b_2}$. Hence this is case one where lines are not parallel. Using the transformation

$$X = x - x_0$$

$$Y = y - y_0$$

Where the constants x_0, y_0 are obtained by solving the following two linear algebraic equations

$$a_1x_0 + b_1y_0 + c_1 = 0$$

$$a_2x_0 + b_2y_0 + c_2 = 0$$

Substituting the values for $a_1, b_1, c_1, a_2, b_2, c_2$ gives

$$x_0 - y_0 + 2 = 0$$

$$x_0 + y_0 - 1 = 0$$

Solving for x_0, y_0 from the above gives

$$x_0 = -\frac{1}{2}$$

$$y_0 = \frac{3}{2}$$

Therefore the transformation becomes

$$X = x + \frac{1}{2}$$

$$Y = y - \frac{3}{2}$$

Using this transformation in $y' - \frac{x-y+2}{-1+y+x} = 0$ result in

$$\frac{dY}{dX} = \frac{X - Y}{Y + X}$$

This is now a homogeneous ODE which will now be solved for $Y(X)$. In canonical form, the ODE is

$$Y' = F(X, Y)$$
$$= -\frac{-X + Y}{Y + X} \tag{1}$$

An ode of the form $Y' = \frac{M(X,Y)}{N(X,Y)}$ is called homogeneous if the functions $M(X, Y)$ and $N(X, Y)$ are both homogeneous functions and of the same order. Recall that a function $f(X, Y)$ is homogeneous of order n if

$$f(t^n X, t^n Y) = t^n f(X, Y)$$

In this case, it can be seen that both $M = X - Y$ and $N = Y + X$ are both homogeneous and of the same order $n = 1$. Therefore this is a homogeneous ode. Since this ode is

homogeneous, it is converted to separable ODE using the substitution $u = \frac{Y}{X}$, or $Y = uX$. Hence

$$\frac{dY}{dX} = \frac{du}{dX}X + u$$

Applying the transformation $Y = uX$ to the above ODE in (1) gives

$$\begin{aligned} \frac{du}{dX}X + u &= \frac{-u + 1}{u + 1} \\ \frac{du}{dX} &= \frac{\frac{-u(X)+1}{u(X)+1} - u(X)}{X} \end{aligned}$$

Or

$$\frac{d}{dX}u(X) - \frac{\frac{-u(X)+1}{u(X)+1} - u(X)}{X} = 0$$

Or

$$\left(\frac{d}{dX}u(X)\right)Xu(X) + \left(\frac{d}{dX}u(X)\right)X + u(X)^2 + 2u(X) - 1 = 0$$

Or

$$(u(X) + 1)X\left(\frac{d}{dX}u(X)\right) + u(X)^2 + 2u(X) - 1 = 0$$

Which is now solved as separable in $u(X)$. Which is now solved in $u(X)$. In canonical form the ODE is

$$\begin{aligned} u' &= F(X, u) \\ &= f(X)g(u) \\ &= -\frac{u^2 + 2u - 1}{(u + 1)X} \end{aligned}$$

Where $f(X) = -\frac{1}{X}$ and $g(u) = \frac{u^2+2u-1}{u+1}$. Integrating both sides gives

$$\begin{aligned} \frac{1}{\frac{u^2+2u-1}{u+1}} du &= -\frac{1}{X} dX \\ \int \frac{1}{\frac{u^2+2u-1}{u+1}} du &= \int -\frac{1}{X} dX \\ \frac{\ln(u^2 + 2u - 1)}{2} &= -\ln(X) + c_3 \end{aligned}$$

Raising both side to exponential gives

$$\sqrt{u^2 + 2u - 1} = e^{-\ln(X)+c_3}$$

Which simplifies to

$$\sqrt{u^2 + 2u - 1} = \frac{c_4}{X}$$

Which simplifies to

$$\sqrt{u(X)^2 + 2u(X) - 1} = \frac{c_4 e^{c_3}}{X}$$

The solution is

$$\sqrt{u(X)^2 + 2u(X) - 1} = \frac{c_4 e^{c_3}}{X}$$

Now u in the above solution is replaced back by Y using $u = \frac{Y}{X}$ which results in the solution

$$\sqrt{\frac{Y(X)^2}{X^2} + \frac{2Y(X)}{X} - 1} = \frac{c_4 e^{c_3}}{X}$$

The solution is implicit $\sqrt{\frac{Y(X)^2 + 2Y(X)X - X^2}{X^2}} = \frac{c_4 e^{c_3}}{X}$. Replacing $Y = y - y_0$, $X = x - x_0$ gives

$$\sqrt{\frac{-\left(\frac{1}{2} + x\right)^2 + 2\left(y - \frac{3}{2}\right)\left(\frac{1}{2} + x\right) + \left(y - \frac{3}{2}\right)^2}{\left(\frac{1}{2} + x\right)^2}} = \frac{c_4 e^{c_3}}{\frac{1}{2} + x}$$

Summary

The solution(s) found are the following

$$\sqrt{\frac{-\left(\frac{1}{2} + x\right)^2 + 2\left(y - \frac{3}{2}\right)\left(\frac{1}{2} + x\right) + \left(y - \frac{3}{2}\right)^2}{\left(\frac{1}{2} + x\right)^2}} = \frac{c_4 e^{c_3}}{\frac{1}{2} + x} \quad (1)$$




Figure 203: Slope field plot

Verification of solutions

$$\sqrt{\frac{-\left(\frac{1}{2} + x\right)^2 + 2\left(y - \frac{3}{2}\right)\left(\frac{1}{2} + x\right) + \left(y - \frac{3}{2}\right)^2}{\left(\frac{1}{2} + x\right)^2}} = \frac{c_4 e^{c_3}}{\frac{1}{2} + x}$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying homogeneous C  
trying homogeneous types:  
trying homogeneous D  
<- homogeneous successful  
<- homogeneous successful`
```

✓ Solution by Maple

Time used: 0.391 (sec). Leaf size: 33

```
dsolve(diff(y(x),x)=(x-y(x)+2)/(x+y(x)-1),y(x), singsol=all)
```

$$y(x) = \frac{-\sqrt{1 + 8\left(x + \frac{1}{2}\right)^2} c_1^2 + (-2x + 2) c_1}{2c_1}$$

✓ Solution by Mathematica

Time used: 0.154 (sec). Leaf size: 53

```
DSolve[y'[x]==(x-y[x]+2)/(x+y[x]-1),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sqrt{2x^2 + 2x + 1 + c_1} - x + 1$$
$$y(x) \rightarrow \sqrt{2x^2 + 2x + 1 + c_1} - x + 1$$

21.14 problem 5(b)

21.14.1 Solving as polynomial ode 1782

Internal problem ID [6076]

Internal file name [OUTPUT/5324_Sunday_June_05_2022_03_34_13_PM_13344468/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 190

Problem number: 5(b).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

```
[[_homogeneous, `class C`], _rational, [_Abel, `2nd type`, `class A`]]
```

$$y' - \frac{2x + 3y + 1}{x - 2y - 1} = 0$$

21.14.1 Solving as polynomial ode

This is ODE of type polynomial. Where the RHS of the ode is ratio of equations of two lines. Writing the ODE in the form

$$y' = \frac{a_1x + b_1y + c_1}{a_2x + b_2y + c_3}$$

Where $a_1 = 2, b_1 = 3, c_1 = 1, a_2 = 1, b_2 = -2, c_2 = -1$. There are now two possible solution methods. The first case is when the two lines $a_1x + b_1y + c_1, a_2x + b_2y + c_3$ are not parallel and the second case is if they are parallel. If they are not parallel, then the transformation $X = x - x_0, Y = y - y_0$ converts the ODE to a homogeneous ODE. The values x_0, y_0 have to be determined. If they are parallel then a transformation $U(x) = a_1x + b_1y$ converts the given ODE in y to a separable ODE in $U(x)$. The first case is when $\frac{a_1}{b_1} \neq \frac{a_2}{b_2}$ and the second case when $\frac{a_1}{b_1} = \frac{a_2}{b_2}$. From the above we see that $\frac{a_1}{b_1} \neq \frac{a_2}{b_2}$. Hence this is case one where lines are not parallel. Using the transformation

$$X = x - x_0$$

$$Y = y - y_0$$

Where the constants x_0, y_0 are obtained by solving the following two linear algebraic equations

$$a_1x_0 + b_1y_0 + c_1 = 0$$

$$a_2x_0 + b_2y_0 + c_2 = 0$$

Substituting the values for $a_1, b_1, c_1, a_2, b_2, c_2$ gives

$$2x_0 + 3y_0 + 1 = 0$$

$$x_0 - 2y_0 - 1 = 0$$

Solving for x_0, y_0 from the above gives

$$x_0 = \frac{1}{7}$$

$$y_0 = -\frac{3}{7}$$

Therefore the transformation becomes

$$X = x - \frac{1}{7}$$

$$Y = y + \frac{3}{7}$$

Using this transformation in $y' - \frac{2x+3y+1}{x-2y-1} = 0$ result in

$$\frac{dY}{dX} = \frac{2X + 3Y}{X - 2Y}$$

This is now a homogeneous ODE which will now be solved for $Y(X)$. In canonical form, the ODE is

$$\begin{aligned} Y' &= F(X, Y) \\ &= -\frac{2X + 3Y}{-X + 2Y} \end{aligned} \quad (1)$$

An ode of the form $Y' = \frac{M(X,Y)}{N(X,Y)}$ is called homogeneous if the functions $M(X, Y)$ and $N(X, Y)$ are both homogeneous functions and of the same order. Recall that a function $f(X, Y)$ is homogeneous of order n if

$$f(t^n X, t^n Y) = t^n f(X, Y)$$

In this case, it can be seen that both $M = 2X + 3Y$ and $N = X - 2Y$ are both homogeneous and of the same order $n = 1$. Therefore this is a homogeneous ode. Since

this ode is homogeneous, it is converted to separable ODE using the substitution $u = \frac{Y}{X}$, or $Y = uX$. Hence

$$\frac{dY}{dX} = \frac{du}{dX}X + u$$

Applying the transformation $Y = uX$ to the above ODE in (1) gives

$$\begin{aligned} \frac{du}{dX}X + u &= \frac{-3u - 2}{2u - 1} \\ \frac{du}{dX} &= \frac{\frac{-3u(X)-2}{2u(X)-1} - u(X)}{X} \end{aligned}$$

Or

$$\frac{d}{dX}u(X) - \frac{\frac{-3u(X)-2}{2u(X)-1} - u(X)}{X} = 0$$

Or

$$2\left(\frac{d}{dX}u(X)\right)Xu(X) - \left(\frac{d}{dX}u(X)\right)X + 2u(X)^2 + 2u(X) + 2 = 0$$

Or

$$2 + X(2u(X) - 1)\left(\frac{d}{dX}u(X)\right) + 2u(X)^2 + 2u(X) = 0$$

Which is now solved as separable in $u(X)$. Which is now solved in $u(X)$. In canonical form the ODE is

$$\begin{aligned} u' &= F(X, u) \\ &= f(X)g(u) \\ &= -\frac{2(u^2 + u + 1)}{X(2u - 1)} \end{aligned}$$

Where $f(X) = -\frac{2}{X}$ and $g(u) = \frac{u^2+u+1}{2u-1}$. Integrating both sides gives

$$\begin{aligned} \frac{1}{\frac{u^2+u+1}{2u-1}} du &= -\frac{2}{X} dX \\ \int \frac{1}{\frac{u^2+u+1}{2u-1}} du &= \int -\frac{2}{X} dX \\ \ln(u^2 + u + 1) - \frac{4\sqrt{3} \arctan\left(\frac{(2u+1)\sqrt{3}}{3}\right)}{3} &= -2 \ln(X) + c_3 \end{aligned}$$

The solution is

$$\ln(u(X)^2 + u(X) + 1) - \frac{4\sqrt{3} \arctan\left(\frac{(2u(X)+1)\sqrt{3}}{3}\right)}{3} + 2 \ln(X) - c_3 = 0$$

Now u in the above solution is replaced back by Y using $u = \frac{Y}{X}$ which results in the solution

$$\ln \left(\frac{Y(X)^2}{X^2} + \frac{Y(X)}{X} + 1 \right) - \frac{4\sqrt{3} \arctan \left(\frac{\left(\frac{2Y(X)}{X} + 1 \right) \sqrt{3}}{3} \right)}{3} + 2 \ln(X) - c_3 = 0$$

The solution is implicit $\ln \left(\frac{Y(X)^2}{X^2} + \frac{Y(X)}{X} + 1 \right) - \frac{4\sqrt{3} \arctan \left(\frac{(2Y(X)+X)\sqrt{3}}{3X} \right)}{3} + 2 \ln(X) - c_3 = 0$.
Replacing $Y = y - y_0$, $X = x - x_0$ gives

$$\ln \left(\frac{\left(y + \frac{3}{7} \right)^2}{\left(x - \frac{1}{7} \right)^2} + \frac{y + \frac{3}{7}}{x - \frac{1}{7}} + 1 \right) - \frac{4\sqrt{3} \arctan \left(\frac{(2y + \frac{5}{7} + x)\sqrt{3}}{3x - \frac{3}{7}} \right)}{3} + 2 \ln \left(x - \frac{1}{7} \right) - c_3 = 0$$

Summary

The solution(s) found are the following

$$\ln \left(\frac{\left(y + \frac{3}{7} \right)^2}{\left(x - \frac{1}{7} \right)^2} + \frac{y + \frac{3}{7}}{x - \frac{1}{7}} + 1 \right) - \frac{4\sqrt{3} \arctan \left(\frac{(2y + \frac{5}{7} + x)\sqrt{3}}{3x - \frac{3}{7}} \right)}{3} + 2 \ln \left(x - \frac{1}{7} \right) - c_3 = \mathbb{Q}(1)$$




Figure 204: Slope field plot

Verification of solutions

$$\ln \left(\frac{\left(y + \frac{3}{7}\right)^2}{\left(x - \frac{1}{7}\right)^2} + \frac{y + \frac{3}{7}}{x - \frac{1}{7}} + 1 \right) - \frac{4\sqrt{3} \arctan \left(\frac{(2y + \frac{5}{7} + x)\sqrt{3}}{3x - \frac{3}{7}} \right)}{3} + 2 \ln \left(x - \frac{1}{7} \right) - c_3 = 0$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying homogeneous C  
trying homogeneous types:  
trying homogeneous D  
<- homogeneous successful  
<- homogeneous successful`
```

✓ Solution by Maple

Time used: 0.312 (sec). Leaf size: 61

```
dsolve(diff(y(x),x)=(2*x+3*y(x)+1)/(x-2*y(x)-1),y(x), singsol=all)
```

$$y(x) = -\frac{5}{14} - \frac{x}{2} + \frac{\sqrt{3}(7x-1) \tan(\text{RootOf}(-2\sqrt{3} \ln(2) + \sqrt{3} \ln(\sec(_Z)^2(7x-1)^2) + \sqrt{3} \ln(3) + 2\sqrt{3} c_1 - 4_Z))}{14}$$

✓ Solution by Mathematica

Time used: 0.12 (sec). Leaf size: 85

```
DSolve[y'[x]==(2*x+3*y[x]+1)/(x-2*y[x]-1),y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve}\left[32\sqrt{3}\arctan\left(\frac{4y(x)+5x+1}{\sqrt{3}(-2y(x)+x-1)}\right)=3\left(8\log\left(\frac{4(7x^2+7y(x)^2+(7x+5)y(x)+x+1)}{(1-7x)^2}\right)\right.\right. \\ \left.\left.+16\log(7x-1)+7c_1\right),y(x)\right]$$

21.15 problem 5(c)

21.15.1 Solving as polynomial ode 1788

Internal problem ID [6077]

Internal file name [OUTPUT/5325_Sunday_June_05_2022_03_34_16_PM_75701382/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 190

Problem number: 5(c).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

```
[[_homogeneous, `class C`], _rational, [_Abel, `2nd type`, `class A`]]
```

$$y' - \frac{y + x + 1}{2x + 2y - 1} = 0$$

21.15.1 Solving as polynomial ode

This is ODE of type polynomial. Where the RHS of the ode is ratio of equations of two lines. Writing the ODE in the form

$$y' = \frac{a_1x + b_1y + c_1}{a_2x + b_2y + c_3}$$

Where $a_1 = 1, b_1 = 1, c_1 = 1, a_2 = 2, b_2 = 2, c_3 = -1$. There are now two possible solution methods. The first case is when the two lines $a_1x + b_1y + c_1, a_2x + b_2y + c_3$ are not parallel and the second case is if they are parallel. If they are not parallel, then the transformation $X = x - x_0, Y = y - y_0$ converts the ODE to a homogeneous ODE. The values x_0, y_0 have to be determined. If they are parallel then a transformation $U(x) = a_1x + b_1y$ converts the given ODE in y to a separable ODE in $U(x)$. The first case is when $\frac{a_1}{b_1} \neq \frac{a_2}{b_2}$ and the second case when $\frac{a_1}{b_1} = \frac{a_2}{b_2}$. From the above we see that $\frac{a_1}{b_1} = \frac{1}{1} = 1$ and $\frac{a_2}{b_2} = \frac{2}{2} = 1$. Hence this is case two, where the lines are parallel. Let $U(x) = x + y$. Solving for y gives

$$y = -x + U(x)$$

Taking derivative w.r.t x gives

$$y' = -1 + U'(x)$$

Substituting the above into the ODE results in the ODE

$$-1 + U'(x) - \frac{U(x) + 1}{2U(x) - 1} = 0$$

Or

$$-1 + U'(x) + \frac{-U(x) - 1}{2U(x) - 1} = 0$$

Or

$$U'(x) = \frac{3U(x)}{2U(x) - 1}$$

Which is now solved as separable in $U(x)$. In canonical form the ODE is

$$\begin{aligned} U' &= F(x, U) \\ &= f(x)g(U) \\ &= \frac{3U}{2U - 1} \end{aligned}$$

Where $f(x) = 1$ and $g(U) = \frac{3U}{2U-1}$. Integrating both sides gives

$$\begin{aligned} \frac{1}{\frac{3U}{2U-1}} dU &= 1 dx \\ \int \frac{1}{\frac{3U}{2U-1}} dU &= \int 1 dx \\ \frac{2U}{3} - \frac{\ln(U)}{3} &= c_2 + x \end{aligned}$$

The solution is

$$\frac{2U(x)}{3} - \frac{\ln(U(x))}{3} - c_2 - x = 0$$

The solution $\frac{2U(x)}{3} - \frac{\ln(U(x))}{3} - c_2 - x = 0$ is converted to y using $U(x) = x + y$. Which gives

$$-\frac{x}{3} + \frac{2y}{3} - \frac{\ln(x+y)}{3} - c_2 = 0$$

Summary

The solution(s) found are the following

$$-\frac{x}{3} + \frac{2y}{3} - \frac{\ln(x+y)}{3} - c_2 = 0 \quad (1)$$




Figure 205: Slope field plot

Verification of solutions

$$-\frac{x}{3} + \frac{2y}{3} - \frac{\ln(x+y)}{3} - c_2 = 0$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying homogeneous C  
1st order, trying the canonical coordinates of the invariance group  
<- 1st order, canonical coordinates successful  
<- homogeneous successful`
```

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 21

```
dsolve(diff(y(x),x)=(x+y(x)+1)/(2*x+2*y(x)-1),y(x), singsol=all)
```

$$y(x) = -\frac{\text{LambertW}(-2e^{-3x+3c_1})}{2} - x$$

✓ Solution by Mathematica

Time used: 4.2 (sec). Leaf size: 32

```
DSolve[y'[x]==(x+y[x]+1)/(2*x+2*y[x]-1),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x - \frac{1}{2}W(-e^{-3x-1+c_1})$$
$$y(x) \rightarrow -x$$

21.16 problem 6(b)

- 21.16.1 Solving as homogeneousTypeMapleC ode 1792
- 21.16.2 Solving as first order ode lie symmetry calculated ode 1795
- 21.16.3 Solving as riccati ode 1802

Internal problem ID [6078]

Internal file name [OUTPUT/5326_Sunday_June_05_2022_03_34_18_PM_75214191/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 190

Problem number: 6(b).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "riccati", "homogeneousTypeMapleC", "first_order_ode_lie_symmetry_calculated"

Maple gives the following as the ode type

```
[[_homogeneous, `class C`], _rational, _Riccati]
```

$$y' - \frac{(-1 + y + x)^2}{2(x + 2)^2} = 0$$

21.16.1 Solving as homogeneousTypeMapleC ode

Let $Y = y + y_0$ and $X = x + x_0$ then the above is transformed to new ode in $Y(X)$

$$\frac{d}{dX}Y(X) = \frac{(-1 + Y(X) + y_0 + X + x_0)^2}{2(X + x_0 + 2)^2}$$

Solving for possible values of x_0 and y_0 which makes the above ode a homogeneous ode results in

$$x_0 = -2$$

$$y_0 = 3$$

Using these values now it is possible to easily solve for $Y(X)$. The above ode now becomes

$$\frac{d}{dX}Y(X) = \frac{X^2 + 2Y(X)X + Y(X)^2}{2X^2}$$

In canonical form, the ODE is

$$\begin{aligned} Y' &= F(X, Y) \\ &= \frac{X^2 + 2YX + Y^2}{2X^2} \end{aligned} \quad (1)$$

An ode of the form $Y' = \frac{M(X, Y)}{N(X, Y)}$ is called homogeneous if the functions $M(X, Y)$ and $N(X, Y)$ are both homogeneous functions and of the same order. Recall that a function $f(X, Y)$ is homogeneous of order n if

$$f(t^n X, t^n Y) = t^n f(X, Y)$$

In this case, it can be seen that both $M = X^2 + 2YX + Y^2$ and $N = 2X^2$ are both homogeneous and of the same order $n = 2$. Therefore this is a homogeneous ode. Since this ode is homogeneous, it is converted to separable ODE using the substitution $u = \frac{Y}{X}$, or $Y = uX$. Hence

$$\frac{dY}{dX} = \frac{du}{dX}X + u$$

Applying the transformation $Y = uX$ to the above ODE in (1) gives

$$\begin{aligned} \frac{du}{dX}X + u &= \frac{1}{2} + u + \frac{1}{2}u^2 \\ \frac{du}{dX} &= \frac{\frac{1}{2} + \frac{u(X)^2}{2}}{X} \end{aligned}$$

Or

$$\frac{d}{dX}u(X) - \frac{\frac{1}{2} + \frac{u(X)^2}{2}}{X} = 0$$

Or

$$2\left(\frac{d}{dX}u(X)\right)X - u(X)^2 - 1 = 0$$

Which is now solved as separable in $u(X)$. Which is now solved in $u(X)$. In canonical form the ODE is

$$\begin{aligned} u' &= F(X, u) \\ &= f(X)g(u) \\ &= \frac{\frac{u^2}{2} + \frac{1}{2}}{X} \end{aligned}$$

Where $f(X) = \frac{1}{X}$ and $g(u) = \frac{u^2}{2} + \frac{1}{2}$. Integrating both sides gives

$$\begin{aligned}\frac{1}{\frac{u^2}{2} + \frac{1}{2}} du &= \frac{1}{X} dX \\ \int \frac{1}{\frac{u^2}{2} + \frac{1}{2}} du &= \int \frac{1}{X} dX \\ 2 \arctan(u) &= \ln(X) + c_2\end{aligned}$$

The solution is

$$2 \arctan(u(X)) - \ln(X) - c_2 = 0$$

Now u in the above solution is replaced back by Y using $u = \frac{Y}{X}$ which results in the solution

$$2 \arctan\left(\frac{Y(X)}{X}\right) - \ln(X) - c_2 = 0$$

Using the solution for $Y(X)$

$$2 \arctan\left(\frac{Y(X)}{X}\right) - \ln(X) - c_2 = 0$$

And replacing back terms in the above solution using

$$\begin{aligned}Y &= y + y_0 \\ X &= x + x_0\end{aligned}$$

Or

$$\begin{aligned}Y &= 3 + y \\ X &= -2 + x\end{aligned}$$

Then the solution in y becomes

$$2 \arctan\left(\frac{y-3}{x+2}\right) - \ln(x+2) - c_2 = 0$$

Summary

The solution(s) found are the following

$$2 \arctan\left(\frac{y-3}{x+2}\right) - \ln(x+2) - c_2 = 0 \tag{1}$$




Figure 206: Slope field plot

Verification of solutions

$$2 \arctan \left(\frac{y - 3}{x + 2} \right) - \ln(x + 2) - c_2 = 0$$

Verified OK.

21.16.2 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = \frac{(-1 + y + x)^2}{2(x + 2)^2}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \quad (1\text{E})$$

$$\eta = xb_2 + yb_3 + b_1 \quad (2\text{E})$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$\begin{aligned} b_2 + \frac{(-1+y+x)^2(b_3-a_2)}{2(x+2)^2} - \frac{(-1+y+x)^4 a_3}{4(x+2)^4} \\ - \left(\frac{-1+y+x}{(x+2)^2} - \frac{(-1+y+x)^2}{(x+2)^3} \right) (xa_2 + ya_3 + a_1) \\ - \frac{(-1+y+x)(xb_2 + yb_3 + b_1)}{(x+2)^2} = 0 \end{aligned} \quad (5E)$$

Putting the above in normal form gives

$$\frac{2x^4a_2 + x^4a_3 - 2x^4b_3 + 4x^3ya_3 + 4x^3yb_2 - 2x^2y^2a_2 + 2x^2y^2a_3 + 2x^2y^2b_3 + y^4a_3 + 16x^3a_2 - 4x^3a_3 + 4x^3b_1 + 20x^3b_2 + 4x^3b_3 + 4x^2ya_1 - 20x^2ya_2 - 4x^2yb_1 - 16x^2yb_2 + 4xy^2a_1 + 4xy^2a_3 - 8xy^2b_3 + 12y^3a_3 - 12x^2a_1 - 6x^2a_2 - 6x^2a_3 - 12x^2b_1 + 96x^2b_2 - 6x^2b_3 - 8xya_1 - 32xya_2 - 24xya_3 - 16xyb_1 - 16xyb_2 + 8y^2a_1 - 8y^2a_2 - 38y^2a_3 - 8y^2b_3 - 12xa_1 + 32xa_2 + 4xa_3 + 144xb_2 - 8xb_3 - 32ya_1 + 16ya_2 + 28ya_3 - 16yb_1 + 24a_1 - 8a_2 - a_3 + 16b_1 + 64b_2 + 8b_3}{(x+2)^4} = 0$$

Setting the numerator to zero gives

$$\begin{aligned} -2x^4a_2 - x^4a_3 + 2x^4b_3 - 4x^3ya_3 - 4x^3yb_2 + 2x^2y^2a_2 - 2x^2y^2a_3 \\ - 2x^2y^2b_3 - y^4a_3 - 16x^3a_2 + 4x^3a_3 - 4x^3b_1 + 20x^3b_2 + 4x^3b_3 \\ + 4x^2ya_1 - 20x^2ya_2 - 4x^2yb_1 - 16x^2yb_2 + 4xy^2a_1 + 4xy^2a_3 - 8xy^2b_3 \\ + 12y^3a_3 - 12x^2a_1 - 6x^2a_2 - 6x^2a_3 - 12x^2b_1 + 96x^2b_2 - 6x^2b_3 \\ - 8xya_1 - 32xya_2 - 24xya_3 - 16xyb_1 - 16xyb_2 + 8y^2a_1 - 8y^2a_2 \\ - 38y^2a_3 - 8y^2b_3 - 12xa_1 + 32xa_2 + 4xa_3 + 144xb_2 - 8xb_3 - 32ya_1 \\ + 16ya_2 + 28ya_3 - 16yb_1 + 24a_1 - 8a_2 - a_3 + 16b_1 + 64b_2 + 8b_3 = 0 \end{aligned} \quad (6E)$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2\}$$

The above PDE (6E) now becomes

$$\begin{aligned}
& -2a_2v_1^4 + 2a_2v_1^2v_2^2 - a_3v_1^4 - 4a_3v_1^3v_2 - 2a_3v_1^2v_2^2 - a_3v_2^4 - 4b_2v_1^3v_2 \\
& + 2b_3v_1^4 - 2b_3v_1^2v_2^2 + 4a_1v_1^2v_2 + 4a_1v_1v_2^2 - 16a_2v_1^3 - 20a_2v_1^2v_2 + 4a_3v_1^3 \\
& + 4a_3v_1v_2^2 + 12a_3v_2^3 - 4b_1v_1^3 - 4b_1v_1^2v_2 + 20b_2v_1^3 - 16b_2v_1^2v_2 + 4b_3v_1^3 \\
& - 8b_3v_1v_2^2 - 12a_1v_1^2 - 8a_1v_1v_2 + 8a_1v_2^2 - 6a_2v_1^2 - 32a_2v_1v_2 - 8a_2v_2^2 \\
& - 6a_3v_1^2 - 24a_3v_1v_2 - 38a_3v_2^2 - 12b_1v_1^2 - 16b_1v_1v_2 + 96b_2v_1^2 - 16b_2v_1v_2 \\
& - 6b_3v_1^2 - 8b_3v_2^2 - 12a_1v_1 - 32a_1v_2 + 32a_2v_1 + 16a_2v_2 + 4a_3v_1 + 28a_3v_2 \\
& - 16b_1v_2 + 144b_2v_1 - 8b_3v_1 + 24a_1 - 8a_2 - a_3 + 16b_1 + 64b_2 + 8b_3 = 0
\end{aligned} \tag{7E}$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2\}$$

Equation (7E) now becomes

$$\begin{aligned}
& (-2a_2 - a_3 + 2b_3)v_1^4 + (-4a_3 - 4b_2)v_1^3v_2 \\
& + (-16a_2 + 4a_3 - 4b_1 + 20b_2 + 4b_3)v_1^3 + (2a_2 - 2a_3 - 2b_3)v_1^2v_2^2 \\
& + (4a_1 - 20a_2 - 4b_1 - 16b_2)v_1^2v_2 + (-12a_1 - 6a_2 - 6a_3 - 12b_1 + 96b_2 - 6b_3)v_1^2 \\
& + (4a_1 + 4a_3 - 8b_3)v_1v_2^2 + (-8a_1 - 32a_2 - 24a_3 - 16b_1 - 16b_2)v_1v_2 \\
& + (-12a_1 + 32a_2 + 4a_3 + 144b_2 - 8b_3)v_1 - a_3v_2^4 + 12a_3v_2^3 \\
& + (8a_1 - 8a_2 - 38a_3 - 8b_3)v_2^2 + (-32a_1 + 16a_2 + 28a_3 - 16b_1)v_2 \\
& + 24a_1 - 8a_2 - a_3 + 16b_1 + 64b_2 + 8b_3 = 0
\end{aligned} \tag{8E}$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} -a_3 &= 0 \\ 12a_3 &= 0 \\ -4a_3 - 4b_2 &= 0 \\ 4a_1 + 4a_3 - 8b_3 &= 0 \\ -2a_2 - a_3 + 2b_3 &= 0 \\ 2a_2 - 2a_3 - 2b_3 &= 0 \\ -32a_1 + 16a_2 + 28a_3 - 16b_1 &= 0 \\ 4a_1 - 20a_2 - 4b_1 - 16b_2 &= 0 \\ 8a_1 - 8a_2 - 38a_3 - 8b_3 &= 0 \\ -12a_1 + 32a_2 + 4a_3 + 144b_2 - 8b_3 &= 0 \\ -8a_1 - 32a_2 - 24a_3 - 16b_1 - 16b_2 &= 0 \\ -16a_2 + 4a_3 - 4b_1 + 20b_2 + 4b_3 &= 0 \\ -12a_1 - 6a_2 - 6a_3 - 12b_1 + 96b_2 - 6b_3 &= 0 \\ 24a_1 - 8a_2 - a_3 + 16b_1 + 64b_2 + 8b_3 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= 2b_3 \\ a_2 &= b_3 \\ a_3 &= 0 \\ b_1 &= -3b_3 \\ b_2 &= 0 \\ b_3 &= b_3 \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned} \xi &= x + 2 \\ \eta &= -3 + y \end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned}\eta &= \eta - \omega(x, y) \xi \\ &= -3 + y - \left(\frac{(-1 + y + x)^2}{2(x + 2)^2} \right) (x + 2) \\ &= \frac{-x^2 - y^2 - 4x + 6y - 13}{2x + 4} \\ \xi &= 0\end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y} \right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{-x^2 - y^2 - 4x + 6y - 13}{2x + 4}} dy\end{aligned}$$

Which results in

$$S = -2 \arctan \left(\frac{2y - 6}{2x + 4} \right)$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y) S_y}{R_x + \omega(x, y) R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{(-1 + y + x)^2}{2(x + 2)^2}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{2y - 6}{x^2 + y^2 + 4x - 6y + 13} \\ S_y &= \frac{-2x - 4}{x^2 + y^2 + 4x - 6y + 13} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = -\frac{1}{x + 2} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = -\frac{1}{R + 2}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = -\ln(R + 2) + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$-2 \arctan\left(\frac{y - 3}{x + 2}\right) = -\ln(x + 2) + c_1$$



Which simplifies to

$$-2 \arctan\left(\frac{y - 3}{x + 2}\right) = -\ln(x + 2) + c_1$$

Which gives

$$y = -\tan\left(-\frac{\ln(x + 2)}{2} + \frac{c_1}{2}\right)x - 2 \tan\left(-\frac{\ln(x + 2)}{2} + \frac{c_1}{2}\right) + 3$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{(-1+y+x)^2}{2(x+2)^2}$ 	$R = x$ $S = -2 \arctan \left(\frac{-3+y}{x+2} \right)$	$\frac{dS}{dR} = -\frac{1}{R+2}$ 

Summary

The solution(s) found are the following

$$y = -\tan \left(-\frac{\ln(x+2)}{2} + \frac{c_1}{2} \right) x - 2 \tan \left(-\frac{\ln(x+2)}{2} + \frac{c_1}{2} \right) + 3 \quad (1)$$




Figure 207: Slope field plot

Verification of solutions

$$y = -\tan\left(-\frac{\ln(x+2)}{2} + \frac{c_1}{2}\right)x - 2\tan\left(-\frac{\ln(x+2)}{2} + \frac{c_1}{2}\right) + 3$$

Verified OK.

21.16.3 Solving as riccati ode

In canonical form the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= \frac{(-1 + y + x)^2}{2(x+2)^2} \end{aligned}$$

This is a Riccati ODE. Comparing the ODE to solve

$$y' = \frac{x^2}{2(x+2)^2} + \frac{xy}{(x+2)^2} + \frac{y^2}{2(x+2)^2} - \frac{x}{(x+2)^2} - \frac{y}{(x+2)^2} + \frac{1}{2(x+2)^2}$$

With Riccati ODE standard form

$$y' = f_0(x) + f_1(x)y + f_2(x)y^2$$

Shows that $f_0(x) = \frac{(x-1)^2}{2(x+2)^2}$, $f_1(x) = \frac{2x-2}{2(x+2)^2}$ and $f_2(x) = \frac{1}{2(x+2)^2}$. Let

$$\begin{aligned} y &= \frac{-u'}{f_2 u} \\ &= \frac{-u'}{\frac{u}{2(x+2)^2}} \end{aligned} \quad (1)$$

Using the above substitution in the given ODE results (after some simplification) in a second order ODE to solve for $u(x)$ which is

$$f_2 u''(x) - (f_2' + f_1 f_2) u'(x) + f_2^2 f_0 u(x) = 0 \quad (2)$$

But

$$\begin{aligned} f_2' &= -\frac{1}{(x+2)^3} \\ f_1 f_2 &= \frac{2x-2}{4(x+2)^4} \\ f_2^2 f_0 &= \frac{(x-1)^2}{8(x+2)^6} \end{aligned}$$

Substituting the above terms back in equation (2) gives

$$\frac{u''(x)}{2(x+2)^2} - \left(-\frac{1}{(x+2)^3} + \frac{2x-2}{4(x+2)^4} \right) u'(x) + \frac{(x-1)^2 u(x)}{8(x+2)^6} = 0$$

Solving the above ODE (this ode solved using Maple, not this program), gives

$$u(x) = e^{\frac{3}{2x+4}} \left((x+2)^{\frac{i}{2}} c_1 + (x+2)^{-\frac{i}{2}} c_2 \right)$$

The above shows that

$$u'(x) = \frac{\left(-c_2(ix + 2i + 3)(x+2)^{-\frac{i}{2}} + (ix + 2i - 3)c_1(x+2)^{\frac{i}{2}} \right) e^{\frac{3}{2x+4}}}{2(x+2)^2}$$

Using the above in (1) gives the solution

$$y = -\frac{-c_2(ix + 2i + 3)(x+2)^{-\frac{i}{2}} + (ix + 2i - 3)c_1(x+2)^{\frac{i}{2}}}{(x+2)^{\frac{i}{2}} c_1 + (x+2)^{-\frac{i}{2}} c_2}$$

Dividing both numerator and denominator by c_1 gives, after renaming the constant $\frac{c_2}{c_1} = c_3$ the following solution

$$y = \frac{(ix + 2i + 3)(x + 2)^{-\frac{i}{2}} - (ix + 2i - 3)c_3(x + 2)^{\frac{i}{2}}}{(x + 2)^{\frac{i}{2}}c_3 + (x + 2)^{-\frac{i}{2}}}$$

Summary

The solution(s) found are the following

$$y = \frac{(ix + 2i + 3)(x + 2)^{-\frac{i}{2}} - (ix + 2i - 3)c_3(x + 2)^{\frac{i}{2}}}{(x + 2)^{\frac{i}{2}}c_3 + (x + 2)^{-\frac{i}{2}}} \quad (1)$$




Figure 208: Slope field plot

Verification of solutions

$$y = \frac{(ix + 2i + 3)(x + 2)^{-\frac{i}{2}} - (ix + 2i - 3)c_3(x + 2)^{\frac{i}{2}}}{(x + 2)^{\frac{i}{2}}c_3 + (x + 2)^{-\frac{i}{2}}}$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying homogeneous C  
trying homogeneous types:  
trying homogeneous D  
<- homogeneous successful  
<- homogeneous successful`
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 21

```
dsolve(diff(y(x),x)=1/2*((x+y(x)-1)/(x+2))^2,y(x), singsol=all)
```

$$y(x) = 3 + \tan\left(\frac{\ln(x+2)}{2} + \frac{c_1}{2}\right)(x+2)$$

✓ Solution by Mathematica

Time used: 0.411 (sec). Leaf size: 99

```
DSolve[y'[x]==1/2*((x+y[x]-1)/(x+2))^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2^i(x+2)^i x + (2+3i)2^i(x+2)^i - 2ic_1 x - (6+4i)c_1}{i2^i(x+2)^i - 2c_1}$$

$$y(x) \rightarrow ix + (3+2i)$$

$$y(x) \rightarrow ix + (3+2i)$$

22 Chapter 5. Existence and uniqueness of solutions to first order equations. Page 198

22.1 problem 1(a)	1807
22.2 problem 1(b)	1814
22.3 problem 1(c)	1820
22.4 problem 1(d)	1825
22.5 problem 1(e)	1831
22.6 problem 1(f)	1837
22.7 problem 1(g)	1844
22.8 problem 1(h)	1850
22.9 problem 2(a)	1856
22.10 problem 2(b)	1862
22.11 problem 2(c)	1868
22.12 problem 2(d)	1876

22.1 problem 1(a)

22.1.1 Solving as exact ode	1807
22.1.2 Maple step by step solution	1810

Internal problem ID [6079]

Internal file name [OUTPUT/5327_Sunday_June_05_2022_03_34_19_PM_1317435/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 198

Problem number: 1(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exact"**

Maple gives the following as the ode type

```
[[_homogeneous, `class A`], _exact, _rational, _dAlembert]
```

$$2xy + (x^2 + 3y^2) y' = 0$$

22.1.1 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned} (x^2 + 3y^2) dy &= (-2xy) dx \\ (2xy) dx + (x^2 + 3y^2) dy &= 0 \end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= 2xy \\ N(x, y) &= x^2 + 3y^2 \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(2xy) \\ &= 2x \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(x^2 + 3y^2) \\ &= 2x \end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \tag{1}$$

$$\frac{\partial \phi}{\partial y} = N \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int 2xy dx \\ \phi &= yx^2 + f(y)\end{aligned}\tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = x^2 + f'(y)\tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = x^2 + 3y^2$. Therefore equation (4) becomes

$$x^2 + 3y^2 = x^2 + f'(y)\tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 3y^2$$

Integrating the above w.r.t y gives

$$\begin{aligned}\int f'(y) dy &= \int (3y^2) dy \\ f(y) &= y^3 + c_1\end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = yx^2 + y^3 + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = yx^2 + y^3$$

Summary

The solution(s) found are the following

$$y^3 + yx^2 = c_1 \quad (1)$$




Figure 209: Slope field plot

Verification of solutions

$$y^3 + yx^2 = c_1$$

Verified OK.

22.1.2 Maple step by step solution

Let's solve

$$2xy + (x^2 + 3y^2)y' = 0$$

- Highest derivative means the order of the ODE is 1
- y'
- Check if ODE is exact
 - ODE is exact if the lhs is the total derivative of a C^2 function

$$F'(x, y) = 0$$

- Compute derivative of lhs

$$F'(x, y) + \left(\frac{\partial}{\partial y} F(x, y) \right) y' = 0$$

- Evaluate derivatives

$$2x = 2x$$

- Condition met, ODE is exact

- Exact ODE implies solution will be of this form

$$\left[F(x, y) = c_1, M(x, y) = F'(x, y), N(x, y) = \frac{\partial}{\partial y} F(x, y) \right]$$

- Solve for $F(x, y)$ by integrating $M(x, y)$ with respect to x

$$F(x, y) = \int 2xy dx + f_1(y)$$

- Evaluate integral

$$F(x, y) = yx^2 + f_1(y)$$

- Take derivative of $F(x, y)$ with respect to y

$$N(x, y) = \frac{\partial}{\partial y} F(x, y)$$

- Compute derivative

$$x^2 + 3y^2 = x^2 + \frac{d}{dy} f_1(y)$$

- Isolate for $\frac{d}{dy} f_1(y)$

$$\frac{d}{dy} f_1(y) = 3y^2$$

- Solve for $f_1(y)$

$$f_1(y) = y^3$$

- Substitute $f_1(y)$ into equation for $F(x, y)$

$$F(x, y) = yx^2 + y^3$$

- Substitute $F(x, y)$ into the solution of the ODE

$$yx^2 + y^3 = c_1$$

- Solve for y

$$\left\{ \begin{array}{l} y = \frac{\left(108c_1 + 12\sqrt{12x^6 + 81c_1^2}\right)^{\frac{1}{3}}}{6} - \frac{2x^2}{\left(108c_1 + 12\sqrt{12x^6 + 81c_1^2}\right)^{\frac{1}{3}}}, y = -\frac{\left(108c_1 + 12\sqrt{12x^6 + 81c_1^2}\right)^{\frac{1}{3}}}{12} + \frac{x^2}{\left(108c_1 + 12\sqrt{12x^6 + 81c_1^2}\right)^{\frac{1}{3}}} \end{array} \right.$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying homogeneous D
<- homogeneous successful`

```

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 189

```
dsolve(2*x*y(x)+(x^2+3*y(x)^2)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{-12c_1x^2 + \left(108 + 12\sqrt{12c_1^3x^6 + 81}\right)^{\frac{2}{3}}}{6 \left(108 + 12\sqrt{12c_1^3x^6 + 81}\right)^{\frac{1}{3}} \sqrt{c_1}}$$

$$y(x) = -\frac{(1 + i\sqrt{3}) \left(108 + 12\sqrt{12c_1^3x^6 + 81}\right)^{\frac{1}{3}}}{12\sqrt{c_1}} - \frac{x^2(i\sqrt{3} - 1) \sqrt{c_1}}{\left(108 + 12\sqrt{12c_1^3x^6 + 81}\right)^{\frac{1}{3}}}$$

$$y(x) = \frac{(i\sqrt{3} - 1) \left(108 + 12\sqrt{12c_1^3x^6 + 81}\right)^{\frac{1}{3}}}{12\sqrt{c_1}} + \frac{(1 + i\sqrt{3}) x^2 \sqrt{c_1}}{\left(108 + 12\sqrt{12c_1^3x^6 + 81}\right)^{\frac{1}{3}}}$$

✓ Solution by Mathematica

Time used: 27.686 (sec). Leaf size: 442

`DSolve[2*x*y[x]+(x^2+3*y[x]^2)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \frac{-2\sqrt[3]{3}x^2 + \sqrt[3]{2}(\sqrt{12x^6 + 81e^{2c_1}} + 9e^{c_1})^{2/3}}{6^{2/3}\sqrt[3]{\sqrt{12x^6 + 81e^{2c_1}} + 9e^{c_1}}}$$

$$y(x) \rightarrow \frac{i2^{2/3}\sqrt[3]{3}(\sqrt{3} + i)(\sqrt{12x^6 + 81e^{2c_1}} + 9e^{c_1})^{2/3} + 2\sqrt[3]{2}\sqrt[6]{3}(\sqrt{3} + 3i)x^2}{12\sqrt[3]{\sqrt{12x^6 + 81e^{2c_1}} + 9e^{c_1}}}$$

$$y(x) \rightarrow \frac{2^{2/3}\sqrt[3]{3}(-1 - i\sqrt{3})(\sqrt{12x^6 + 81e^{2c_1}} + 9e^{c_1})^{2/3} + 2\sqrt[3]{2}\sqrt[6]{3}(\sqrt{3} - 3i)x^2}{12\sqrt[3]{\sqrt{12x^6 + 81e^{2c_1}} + 9e^{c_1}}}$$

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow \frac{\sqrt[3]{x^6} - x^2}{\sqrt{3}\sqrt[6]{x^6}}$$

$$y(x) \rightarrow \frac{(\sqrt{3} - 3i)x^2 - (\sqrt{3} + 3i)\sqrt[3]{x^6}}{6\sqrt[6]{x^6}}$$

$$y(x) \rightarrow \frac{(\sqrt{3} + 3i)x^2 - (\sqrt{3} - 3i)\sqrt[3]{x^6}}{6\sqrt[6]{x^6}}$$

22.2 problem 1(b)

22.2.1 Solving as exact ode	1814
22.2.2 Maple step by step solution	1818

Internal problem ID [6080]

Internal file name [OUTPUT/5328_Sunday_June_05_2022_03_34_21_PM_20496951/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 198

Problem number: 1(b).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exact"**

Maple gives the following as the ode type

`[_quadrature]`

$$xy + (x + y)y' = -x^2$$

22.2.1 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned} dy &= (-x) dx \\ (x) dx + dy &= 0 \end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= x \\ N(x, y) &= 1 \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(x) \\ &= 0 \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(1) \\ &= 0 \end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \tag{1}$$

$$\frac{\partial \phi}{\partial y} = N \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int x dx \\ \phi &= \frac{x^2}{2} + f(y)\end{aligned}\tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y)\tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = 1$. Therefore equation (4) becomes

$$1 = 0 + f'(y)\tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 1$$

Integrating the above w.r.t y gives

$$\begin{aligned}\int f'(y) dy &= \int (1) dy \\ f(y) &= y + c_1\end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = \frac{x^2}{2} + y + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \frac{x^2}{2} + y$$

The solution becomes

$$y = -\frac{x^2}{2} + c_1$$

Summary

The solution(s) found are the following

$$y = -\frac{x^2}{2} + c_1 \tag{1}$$




Figure 210: Slope field plot

Verification of solutions

$$y = -\frac{x^2}{2} + c_1$$

Verified OK.

22.2.2 Maple step by step solution

Let's solve

$$xy + (x + y)y' = -x^2$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$y' = -x$$

- Integrate both sides with respect to x

$$\int y' dx = \int -x dx + c_1$$

- Evaluate integral

$$y = -\frac{x^2}{2} + c_1$$

- Solve for y

$$y = -\frac{x^2}{2} + c_1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve((x^2+x*y(x))+(x+y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = -x$$
$$y(x) = -\frac{x^2}{2} + c_1$$

✓ Solution by Mathematica

Time used: 0.132 (sec). Leaf size: 53

```
DSolve[(x^2+y[x])+(x+y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x - \sqrt{-\frac{2x^3}{3} + x^2 + c_1}$$

$$y(x) \rightarrow -x + \sqrt{-\frac{2x^3}{3} + x^2 + c_1}$$

22.3 problem 1(c)

22.3.1 Solving as exact ode 1820

22.3.2 Maple step by step solution 1824

Internal problem ID [6081]

Internal file name [OUTPUT/5329_Sunday_June_05_2022_03_34_22_PM_6110514/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 198

Problem number: 1(c).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exact"**

Maple gives the following as the ode type

[_separable]

$$e^y(1+y)y' = -e^x$$

22.3.1 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned} (-e^y(1+y)) dy &= (e^x) dx \\ (-e^x) dx + (-e^y(1+y)) dy &= 0 \end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -e^x \\ N(x, y) &= -e^y(1+y) \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-e^x) \\ &= 0 \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(-e^y(1+y)) \\ &= 0 \end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \tag{1}$$

$$\frac{\partial \phi}{\partial y} = N \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -e^x dx \\ \phi &= -e^x + f(y)\end{aligned}\tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y)\tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = -e^y(1 + y)$. Therefore equation (4) becomes

$$-e^y(1 + y) = 0 + f'(y)\tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = -e^y(1 + y)$$

Integrating the above w.r.t y gives

$$\begin{aligned}\int f'(y) dy &= \int (-e^y(1 + y)) dy \\ f(y) &= -e^y y + c_1\end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -e^x - e^y y + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -e^x - e^y y$$

The solution becomes

$$y = \text{LambertW}(-e^x - c_1)$$

Summary

The solution(s) found are the following

$$y = \text{LambertW}(-e^x - c_1) \tag{1}$$




Figure 211: Slope field plot

Verification of solutions

$$y = \text{LambertW}(-e^x - c_1)$$

Verified OK.

22.3.2 Maple step by step solution

Let's solve

$$e^y(1+y)y' = -e^x$$

- Highest derivative means the order of the ODE is 1

y'

- Integrate both sides with respect to x

$$\int e^y(1+y)y'dx = \int -e^x dx + c_1$$

- Evaluate integral

$$e^y y = -e^x + c_1$$

- Solve for y

$$y = \text{LambertW}(-e^x + c_1)$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
<- separable successful`
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 13

```
dsolve(exp(x)+(exp(y(x))*(y(x)+1))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \text{LambertW}(-c_1 - e^x)$$

✓ Solution by Mathematica

Time used: 60.161 (sec). Leaf size: 14

```
DSolve[Exp[x]+(Exp[y[x]]*(y[x]+1))*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow W(-e^x + c_1)$$

22.4 problem 1(d)

22.4.1 Solving as exact ode 1825

22.4.2 Maple step by step solution 1829

Internal problem ID [6082]

Internal file name [OUTPUT/5330_Sunday_June_05_2022_03_34_24_PM_67663664/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 198

Problem number: 1(d).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exact"**

Maple gives the following as the ode type

[_separable]

$$\cos(x) \cos(y)^2 - \sin(x) \sin(2y) y' = 0$$

22.4.1 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned} \left(\frac{\sin(2y)}{\cos(y)^2} \right) dy &= \left(\frac{\cos(x)}{\sin(x)} \right) dx \\ \left(-\frac{\cos(x)}{\sin(x)} \right) dx + \left(\frac{\sin(2y)}{\cos(y)^2} \right) dy &= 0 \end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -\frac{\cos(x)}{\sin(x)} \\ N(x, y) &= \frac{\sin(2y)}{\cos(y)^2} \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(-\frac{\cos(x)}{\sin(x)} \right) \\ &= 0 \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{\sin(2y)}{\cos(y)^2} \right) \\ &= 0 \end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned} \int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\frac{\cos(x)}{\sin(x)} dx \\ \phi &= -\ln(\sin(x)) + f(y) \end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{\sin(2y)}{\cos(y)^2}$. Therefore equation (4) becomes

$$\frac{\sin(2y)}{\cos(y)^2} = 0 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$\begin{aligned} f'(y) &= \frac{\sin(2y)}{\cos(y)^2} \\ &= 2 \tan(y) \end{aligned}$$

Integrating the above w.r.t y results in

$$\begin{aligned} \int f'(y) dy &= \int (2 \tan(y)) dy \\ f(y) &= -2 \ln(\cos(y)) + c_1 \end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\ln(\sin(x)) - 2\ln(\cos(y)) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\ln(\sin(x)) - 2\ln(\cos(y))$$

Summary

The solution(s) found are the following

$$-\ln(\sin(x)) - 2\ln(\cos(y)) = c_1 \tag{1}$$




Figure 212: Slope field plot

Verification of solutions

$$-\ln(\sin(x)) - 2\ln(\cos(y)) = c_1$$

Verified OK.

22.4.2 Maple step by step solution

Let's solve

$$\cos(x) \cos(y)^2 - \sin(x) \sin(2y) y' = 0$$

- Highest derivative means the order of the ODE is 1

y'

- Integrate both sides with respect to x

$$\int (\cos(x) \cos(y)^2 - \sin(x) \sin(2y) y') dx = \int 0 dx + c_1$$

- Evaluate integral

$$\frac{\sin(x-2y)}{4} + \frac{\sin(2y+x)}{4} + \frac{\sin(x)}{2} = c_1$$

- Solve for y

$$y = \frac{\arccos\left(\frac{2c_1 - \sin(x)}{\sin(x)}\right)}{2}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
<- separable successful`
```

✓ Solution by Maple

Time used: 0.219 (sec). Leaf size: 25

```
dsolve(cos(x)*cos(y(x))^2-sin(x)*sin(2*y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \arccos\left(\frac{1}{\sqrt{c_1 \sin(x)}}\right)$$

$$y(x) = \frac{\pi}{2} + \arcsin\left(\frac{1}{\sqrt{c_1 \sin(x)}}\right)$$

✓ Solution by Mathematica

Time used: 6.536 (sec). Leaf size: 73

```
DSolve[Cos[x]*Cos[y[x]]^2-Sin[x]*Sin[2*y[x]]*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\pi}{2}$$

$$y(x) \rightarrow \frac{\pi}{2}$$

$$y(x) \rightarrow -\arccos\left(-\frac{c_1}{4\sqrt{\sin(x)}}\right)$$

$$y(x) \rightarrow \arccos\left(-\frac{c_1}{4\sqrt{\sin(x)}}\right)$$

$$y(x) \rightarrow -\frac{\pi}{2}$$

$$y(x) \rightarrow \frac{\pi}{2}$$

22.5 problem 1(e)

22.5.1 Solving as exact ode 1831

22.5.2 Maple step by step solution 1835

Internal problem ID [6083]

Internal file name [OUTPUT/5331_Sunday_June_05_2022_03_34_26_PM_63382602/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 198

Problem number: 1(e).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exact"**

Maple gives the following as the ode type

[_separable]

$$y^3 x^2 - x^3 y^2 y' = 0$$

22.5.1 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned} \left(\frac{1}{y}\right) dy &= \left(\frac{1}{x}\right) dx \\ \left(-\frac{1}{x}\right) dx + \left(\frac{1}{y}\right) dy &= 0 \end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -\frac{1}{x} \\ N(x, y) &= \frac{1}{y} \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(-\frac{1}{x}\right) \\ &= 0 \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{1}{y}\right) \\ &= 0 \end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned} \int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\frac{1}{x} dx \\ \phi &= -\ln(x) + f(y) \end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{1}{y}$. Therefore equation (4) becomes

$$\frac{1}{y} = 0 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = \frac{1}{y}$$

Integrating the above w.r.t y gives

$$\begin{aligned} \int f'(y) dy &= \int \left(\frac{1}{y}\right) dy \\ f(y) &= \ln(y) + c_1 \end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\ln(x) + \ln(y) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\ln(x) + \ln(y)$$

The solution becomes

$$y = x e^{c_1}$$

Summary

The solution(s) found are the following

$$y = x e^{c_1} \tag{1}$$




Figure 213: Slope field plot

Verification of solutions

$$y = x e^{c_1}$$

Verified OK.

22.5.2 Maple step by step solution

Let's solve

$$y^3 x^2 - x^3 y^2 y' = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y} = \frac{1}{x}$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y} dx = \int \frac{1}{x} dx + c_1$$

- Evaluate integral

$$\ln(y) = \ln(x) + c_1$$

- Solve for y

$$y = x e^{c_1}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve(x^2*y(x)^3-x^3*y(x)^2*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = c_1 x$$

✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 19

```
DSolve[x^2*y[x]^3-x^3*y[x]^2*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow c_1 x$$

$$y(x) \rightarrow 0$$

22.6 problem 1(f)

22.6.1 Solving as exact ode	1837
22.6.2 Maple step by step solution	1841

Internal problem ID [6084]

Internal file name [OUTPUT/5332_Sunday_June_05_2022_03_34_28_PM_60115467/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 198

Problem number: 1(f).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exact"**

Maple gives the following as the ode type

```
[[_homogeneous, `class A`], _exact, _rational, [_Abel, `2nd  
type`, `class A`]]
```

$$y + (x - y)y' = -x$$

22.6.1 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}(x - y) dy &= (-y - x) dx \\ (x + y) dx + (x - y) dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= x + y \\ N(x, y) &= x - y\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(x + y) \\ &= 1\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(x - y) \\ &= 1\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned} \int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int x + y dx \\ \phi &= \frac{x(2y + x)}{2} + f(y) \end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = x + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = x - y$. Therefore equation (4) becomes

$$x - y = x + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = -y$$

Integrating the above w.r.t y gives

$$\begin{aligned} \int f'(y) dy &= \int (-y) dy \\ f(y) &= -\frac{y^2}{2} + c_1 \end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = \frac{x(2y + x)}{2} - \frac{y^2}{2} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \frac{x(2y + x)}{2} - \frac{y^2}{2}$$

Summary

The solution(s) found are the following

$$\frac{x(2y + x)}{2} - \frac{y^2}{2} = c_1 \tag{1}$$




Figure 214: Slope field plot

Verification of solutions

$$\frac{x(2y + x)}{2} - \frac{y^2}{2} = c_1$$

Verified OK.

22.6.2 Maple step by step solution

Let's solve

$$y + (x - y) y' = -x$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Check if ODE is exact

- ODE is exact if the lhs is the total derivative of a C^2 function

$$F'(x, y) = 0$$

- Compute derivative of lhs

$$F'(x, y) + \left(\frac{\partial}{\partial y} F(x, y) \right) y' = 0$$

- Evaluate derivatives

$$1 = 1$$

- Condition met, ODE is exact

- Exact ODE implies solution will be of this form

$$\left[F(x, y) = c_1, M(x, y) = F'(x, y), N(x, y) = \frac{\partial}{\partial y} F(x, y) \right]$$

- Solve for $F(x, y)$ by integrating $M(x, y)$ with respect to x

$$F(x, y) = \int (x + y) dx + f_1(y)$$

- Evaluate integral

$$F(x, y) = \frac{x^2}{2} + xy + f_1(y)$$

- Take derivative of $F(x, y)$ with respect to y

$$N(x, y) = \frac{\partial}{\partial y} F(x, y)$$

- Compute derivative

$$x - y = x + \frac{d}{dy} f_1(y)$$

- Isolate for $\frac{d}{dy} f_1(y)$

$$\frac{d}{dy} f_1(y) = -y$$

- Solve for $f_1(y)$

$$f_1(y) = -\frac{y^2}{2}$$

- Substitute $f_1(y)$ into equation for $F(x, y)$

$$F(x, y) = \frac{1}{2}x^2 + xy - \frac{1}{2}y^2$$

- Substitute $F(x, y)$ into the solution of the ODE

$$\frac{1}{2}x^2 + xy - \frac{1}{2}y^2 = c_1$$

- Solve for y

$$\{y = x - \sqrt{2x^2 - 2c_1}, y = x + \sqrt{2x^2 - 2c_1}\}$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying homogeneous D
<- homogeneous successful`

```

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 49

```
dsolve((x+y(x))+(x-y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{c_1 x - \sqrt{2x^2 c_1^2 + 1}}{c_1}$$

$$y(x) = \frac{c_1 x + \sqrt{2x^2 c_1^2 + 1}}{c_1}$$

✓ Solution by Mathematica

Time used: 0.449 (sec). Leaf size: 86

```
DSolve[(x+y[x])+(x-y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x - \sqrt{2x^2 + e^{2c_1}}$$

$$y(x) \rightarrow x + \sqrt{2x^2 + e^{2c_1}}$$

$$y(x) \rightarrow x - \sqrt{2}\sqrt{x^2}$$

$$y(x) \rightarrow \sqrt{2}\sqrt{x^2} + x$$

22.7 problem 1(g)

22.7.1 Solving as exact ode 1844

22.7.2 Maple step by step solution 1847

Internal problem ID [6085]

Internal file name [OUTPUT/5333_Sunday_June_05_2022_03_34_29_PM_40997286/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 198

Problem number: 1(g).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exact"**

Maple gives the following as the ode type

[_exact]

$$2 e^{2x} y + 2 \cos(y) x + (e^{2x} - x^2 \sin(y)) y' = 0$$

22.7.1 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned} (e^{2x} - x^2 \sin(y)) dy &= (-2e^{2x}y - 2 \cos(y) x) dx \\ (2e^{2x}y + 2 \cos(y) x) dx + (e^{2x} - x^2 \sin(y)) dy &= 0 \end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= 2e^{2x}y + 2 \cos(y) x \\ N(x, y) &= e^{2x} - x^2 \sin(y) \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} (2e^{2x}y + 2 \cos(y) x) \\ &= 2e^{2x} - 2 \sin(y) x \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} (e^{2x} - x^2 \sin(y)) \\ &= 2e^{2x} - 2 \sin(y) x \end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int 2 e^{2x} y + 2 \cos(y) x dx \\ \phi &= \cos(y) x^2 + e^{2x} y + f(y)\end{aligned}\tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = e^{2x} - x^2 \sin(y) + f'(y)\tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = e^{2x} - x^2 \sin(y)$. Therefore equation (4) becomes

$$e^{2x} - x^2 \sin(y) = e^{2x} - x^2 \sin(y) + f'(y)\tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = \cos(y) x^2 + e^{2x} y + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \cos(y) x^2 + e^{2x} y$$

Summary

The solution(s) found are the following

$$\cos(y) x^2 + e^{2x} y = c_1\tag{1}$$




Figure 215: Slope field plot

Verification of solutions

$$\cos(y) x^2 + e^{2x} y = c_1$$

Verified OK.

22.7.2 Maple step by step solution

Let's solve

$$2e^{2x}y + 2\cos(y)x + (e^{2x} - x^2\sin(y))y' = 0$$

- Highest derivative means the order of the ODE is 1
 y'
- Check if ODE is exact
 - ODE is exact if the lhs is the total derivative of a C^2 function
 $F'(x, y) = 0$
 - Compute derivative of lhs
 $F'(x, y) + \left(\frac{\partial}{\partial y}F(x, y)\right)y' = 0$

- Evaluate derivatives

$$2e^{2x} - 2 \sin(y) x = 2e^{2x} - 2 \sin(y) x$$
- Condition met, ODE is exact
- Exact ODE implies solution will be of this form

$$\left[F(x, y) = c_1, M(x, y) = F'(x, y), N(x, y) = \frac{\partial}{\partial y} F(x, y) \right]$$
- Solve for $F(x, y)$ by integrating $M(x, y)$ with respect to x

$$F(x, y) = \int (2e^{2x}y + 2 \cos(y) x) dx + f_1(y)$$
- Evaluate integral

$$F(x, y) = \cos(y) x^2 + e^{2x}y + f_1(y)$$
- Take derivative of $F(x, y)$ with respect to y

$$N(x, y) = \frac{\partial}{\partial y} F(x, y)$$
- Compute derivative

$$e^{2x} - x^2 \sin(y) = -x^2 \sin(y) + e^{2x} + \frac{d}{dy} f_1(y)$$
- Isolate for $\frac{d}{dy} f_1(y)$

$$\frac{d}{dy} f_1(y) = 0$$
- Solve for $f_1(y)$

$$f_1(y) = 0$$
- Substitute $f_1(y)$ into equation for $F(x, y)$

$$F(x, y) = \cos(y) x^2 + e^{2x}y$$
- Substitute $F(x, y)$ into the solution of the ODE

$$\cos(y) x^2 + e^{2x}y = c_1$$
- Solve for y

$$y = \text{RootOf}(-\cos(_Z) x^2 - _Z e^{2x} + c_1)$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying Chini  
differential order: 1; looking for linear symmetries  
trying exact  
<- exact successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 19

```
dsolve((2*y(x)*exp(2*x)+2*x*cos(y(x)))+(exp(2*x)-x^2*sin(y(x)))*diff(y(x),x)=0,y(x), singsol
```

$$\cos(y(x))x^2 + y(x)e^{2x} + c_1 = 0$$

✓ Solution by Mathematica

Time used: 0.414 (sec). Leaf size: 30

```
DSolve[(2*y[x]*Exp[2*x]+2*x*Cos[y[x]])+(Exp[2*x]-x^2*Sin[y[x]])*y'[x]==0,y[x],x,IncludeSingul
```

$$\text{Solve}\left[2\left(\frac{1}{2}x^2 \cos(y(x)) + \frac{1}{2}e^{2x}y(x)\right) = c_1, y(x)\right]$$

22.8 problem 1(h)

22.8.1 Solving as exact ode 1850

22.8.2 Maple step by step solution 1853

Internal problem ID [6086]

Internal file name [OUTPUT/5334_Sunday_June_05_2022_03_34_33_PM_95864308/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 198

Problem number: 1(h).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exact"**

Maple gives the following as the ode type

[_linear]

$$xy' + y = -3 \ln(x) x^2 - x^2$$

22.8.1 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned} (x) dy &= (-3 \ln(x) x^2 - x^2 - y) dx \\ (3 \ln(x) x^2 + x^2 + y) dx + (x) dy &= 0 \end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= 3 \ln(x) x^2 + x^2 + y \\ N(x, y) &= x \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} (3 \ln(x) x^2 + x^2 + y) \\ &= 1 \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} (x) \\ &= 1 \end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \tag{1}$$

$$\frac{\partial \phi}{\partial y} = N \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int 3 \ln(x) x^2 + x^2 + y dx \\ \phi &= x(\ln(x) x^2 + y) + f(y)\end{aligned}\tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = x + f'(y)\tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = x$. Therefore equation (4) becomes

$$x = x + f'(y)\tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = x(\ln(x) x^2 + y) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = x(\ln(x) x^2 + y)$$

The solution becomes

$$y = -\frac{x^3 \ln(x) - c_1}{x}$$

Summary

The solution(s) found are the following

$$y = -\frac{x^3 \ln(x) - c_1}{x} \quad (1)$$




Figure 216: Slope field plot

Verification of solutions

$$y = -\frac{x^3 \ln(x) - c_1}{x}$$

Verified OK.

22.8.2 Maple step by step solution

Let's solve

$$xy' + y = -3 \ln(x) x^2 - x^2$$

- Highest derivative means the order of the ODE is 1

y'

- Isolate the derivative

$$y' = -\frac{y}{x} - x(3 \ln(x) + 1)$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' + \frac{y}{x} = -x(3 \ln(x) + 1)$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) \left(y' + \frac{y}{x} \right) = -\mu(x) x(3 \ln(x) + 1)$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x) y)$

$$\mu(x) \left(y' + \frac{y}{x} \right) = \mu'(x) y + \mu(x) y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = \frac{\mu(x)}{x}$$

- Solve to find the integrating factor

$$\mu(x) = x$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x) y) \right) dx = \int -\mu(x) x(3 \ln(x) + 1) dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x) y = \int -\mu(x) x(3 \ln(x) + 1) dx + c_1$$

- Solve for y

$$y = \frac{\int -\mu(x)x(3 \ln(x)+1)dx+c_1}{\mu(x)}$$

- Substitute $\mu(x) = x$

$$y = \frac{\int -x^2(3 \ln(x)+1)dx+c_1}{x}$$

- Evaluate the integrals on the rhs

$$y = \frac{-x^3 \ln(x)+c_1}{x}$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve((3*x^2*ln(x)+x^2+y(x))+x*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{-x^3 \ln(x) + c_1}{x}$$

✓ Solution by Mathematica

Time used: 0.035 (sec). Leaf size: 19

```
DSolve[(3*x^2*Log[x]+x^2+y[x])+x*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{-x^3 \log(x) + c_1}{x}$$

22.9 problem 2(a)

22.9.1 Solving as exact ode 1856

22.9.2 Maple step by step solution 1860

Internal problem ID [6087]

Internal file name [OUTPUT/5335_Sunday_June_05_2022_03_34_34_PM_695312/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 198

Problem number: 2(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exact"**

Maple gives the following as the ode type

[_separable]

$$2y^3 + 3xy^2y' = -2$$

22.9.1 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned} \left(-\frac{3y^2}{2(y^3+1)}\right) dy &= \left(\frac{1}{x}\right) dx \\ \left(-\frac{1}{x}\right) dx + \left(-\frac{3y^2}{2(y^3+1)}\right) dy &= 0 \end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -\frac{1}{x} \\ N(x, y) &= -\frac{3y^2}{2(y^3+1)} \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(-\frac{1}{x}\right) \\ &= 0 \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(-\frac{3y^2}{2(y^3+1)}\right) \\ &= 0 \end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned} \int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\frac{1}{x} dx \\ \phi &= -\ln(x) + f(y) \end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = -\frac{3y^2}{2(y^3+1)}$. Therefore equation (4) becomes

$$-\frac{3y^2}{2(y^3+1)} = 0 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = -\frac{3y^2}{2(y^3+1)}$$

Integrating the above w.r.t y gives

$$\begin{aligned} \int f'(y) dy &= \int \left(-\frac{3y^2}{2y^3+2} \right) dy \\ f(y) &= -\frac{\ln(y^3+1)}{2} + c_1 \end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\ln(x) - \frac{\ln(y^3 + 1)}{2} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\ln(x) - \frac{\ln(y^3 + 1)}{2}$$

Summary

The solution(s) found are the following

$$-\ln(x) - \frac{\ln(y^3 + 1)}{2} = c_1 \tag{1}$$




Figure 217: Slope field plot

Verification of solutions

$$-\ln(x) - \frac{\ln(y^3 + 1)}{2} = c_1$$

Verified OK.

22.9.2 Maple step by step solution

Let's solve

$$2y^3 + 3xy^2y' = -2$$

- Highest derivative means the order of the ODE is 1

y'

- Separate variables

$$\frac{y'y^2}{-2y^3-2} = \frac{1}{3x}$$

- Integrate both sides with respect to x

$$\int \frac{y'y^2}{-2y^3-2} dx = \int \frac{1}{3x} dx + c_1$$

- Evaluate integral

$$-\frac{\ln(y^3+1)}{6} = \frac{\ln(x)}{3} + c_1$$

- Solve for y

$$y = \frac{\left(-x e^{3c_1} \left(x^2 (e^{3c_1})^2 - 1\right)\right)^{\frac{1}{3}}}{x e^{3c_1}}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
<- Bernoulli successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 73

```
dsolve((2*y(x)^3+2)+(3*x*y(x)^2)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{((-x^2 + c_1)x)^{\frac{1}{3}}}{x}$$

$$y(x) = -\frac{((-x^2 + c_1)x)^{\frac{1}{3}}(1 + i\sqrt{3})}{2x}$$

$$y(x) = \frac{((-x^2 + c_1)x)^{\frac{1}{3}}(i\sqrt{3} - 1)}{2x}$$

✓ Solution by Mathematica

Time used: 0.281 (sec). Leaf size: 215

```
DSolve[(3*y[x]^3+2)+(3*x*y[x]^2)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\sqrt[3]{-\frac{1}{3}\sqrt[3]{-2x^3 + e^{9c_1}}}}{x}$$

$$y(x) \rightarrow \frac{\sqrt[3]{-2x^3 + e^{9c_1}}}{\sqrt[3]{3}x}$$

$$y(x) \rightarrow \frac{(-1)^{2/3}\sqrt[3]{-2x^3 + e^{9c_1}}}{\sqrt[3]{3}x}$$

$$y(x) \rightarrow \sqrt[3]{-\frac{2}{3}}$$

$$y(x) \rightarrow -\sqrt[3]{\frac{2}{3}}$$

$$y(x) \rightarrow -(-1)^{2/3}\sqrt[3]{\frac{2}{3}}$$

$$y(x) \rightarrow \frac{\sqrt[3]{-\frac{2}{3}x^2}}{(-x^3)^{2/3}}$$

$$y(x) \rightarrow \frac{\sqrt[3]{\frac{2}{3}\sqrt[3]{-x^3}}}{x}$$

$$y(x) \rightarrow \frac{(-1)^{2/3}\sqrt[3]{\frac{2}{3}\sqrt[3]{-x^3}}}{x}$$

22.10 problem 2(b)

22.10.1 Solving as exact ode	1862
22.10.2 Maple step by step solution	1866

Internal problem ID [6088]

Internal file name [OUTPUT/5336_Sunday_June_05_2022_03_34_36_PM_19452203/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 198

Problem number: 2(b).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exact"**

Maple gives the following as the ode type

[_separable]

$$-2y' \sin(y) \sin(x) + \cos(x) \cos(y) = 0$$

22.10.1 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned} \left(\frac{2 \sin(y)}{\cos(y)} \right) dy &= \left(\frac{\cos(x)}{\sin(x)} \right) dx \\ \left(-\frac{\cos(x)}{\sin(x)} \right) dx + \left(\frac{2 \sin(y)}{\cos(y)} \right) dy &= 0 \end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -\frac{\cos(x)}{\sin(x)} \\ N(x, y) &= \frac{2 \sin(y)}{\cos(y)} \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(-\frac{\cos(x)}{\sin(x)} \right) \\ &= 0 \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{2 \sin(y)}{\cos(y)} \right) \\ &= 0 \end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned} \int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\frac{\cos(x)}{\sin(x)} dx \\ \phi &= -\ln(\sin(x)) + f(y) \end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{2 \sin(y)}{\cos(y)}$. Therefore equation (4) becomes

$$\frac{2 \sin(y)}{\cos(y)} = 0 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$\begin{aligned} f'(y) &= \frac{2 \sin(y)}{\cos(y)} \\ &= 2 \tan(y) \end{aligned}$$

Integrating the above w.r.t y results in

$$\begin{aligned} \int f'(y) dy &= \int (2 \tan(y)) dy \\ f(y) &= -2 \ln(\cos(y)) + c_1 \end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\ln(\sin(x)) - 2\ln(\cos(y)) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\ln(\sin(x)) - 2\ln(\cos(y))$$

Summary

The solution(s) found are the following

$$-\ln(\sin(x)) - 2\ln(\cos(y)) = c_1 \tag{1}$$




Figure 218: Slope field plot

Verification of solutions

$$-\ln(\sin(x)) - 2\ln(\cos(y)) = c_1$$

Verified OK.

22.10.2 Maple step by step solution

Let's solve

$$-2y' \sin(y) \sin(x) + \cos(x) \cos(y) = 0$$

- Highest derivative means the order of the ODE is 1

y'

- Separate variables

$$\frac{y' \sin(y)}{\cos(y)} = \frac{\cos(x)}{2 \sin(x)}$$

- Integrate both sides with respect to x

$$\int \frac{y' \sin(y)}{\cos(y)} dx = \int \frac{\cos(x)}{2 \sin(x)} dx + c_1$$

- Evaluate integral

$$-\ln(\cos(y)) = \frac{\ln(\sin(x))}{2} + c_1$$

- Solve for y

$$\left\{ y = \pi - \arccos\left(\frac{\sqrt{\sin(x)e^{-2c_1}}}{\sin(x)}\right), y = \arccos\left(\frac{\sqrt{\sin(x)e^{-2c_1}}}{\sin(x)}\right) \right\}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
<- separable successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve(cos(x)*cos(y(x))-2*sin(x)*sin(y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \arccos\left(\frac{1}{\sqrt{c_1 \sin(x)}}\right)$$

$$y(x) = \frac{\pi}{2} + \arcsin\left(\frac{1}{\sqrt{c_1 \sin(x)}}\right)$$

✓ Solution by Mathematica

Time used: 0.491 (sec). Leaf size: 43

```
DSolve[Cos[x]*cos[y[x]]-(2*Sin[x]*Sin[y[x]])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \text{InverseFunction} \left[\int_1^{\#1} \frac{\sin(K[1])}{\cos(K[1])} dK[1] \& \right] \left[\frac{1}{2} \log(\sin(x)) + c_1 \right]$$

$$y(x) \rightarrow \cos^{(-1)}(0)$$

22.11 problem 2(c)

22.11.1 Solving as exact ode 1868

Internal problem ID [6089]

Internal file name [OUTPUT/5337_Sunday_June_05_2022_03_34_38_PM_97437203/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 198

Problem number: 2(c).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exactWithIntegrationFactor"**

Maple gives the following as the ode type

```
[[_homogeneous, `class G`], _rational, [_Abel, `2nd type`, `class B`]]
```

$$5x^3y^2 + 2y + (3yx^4 + 2x)y' = 0$$

22.11.1 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned} (3y x^4 + 2x) dy &= (-5y^2 x^3 - 2y) dx \\ (5y^2 x^3 + 2y) dx + (3y x^4 + 2x) dy &= 0 \end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= 5y^2 x^3 + 2y \\ N(x, y) &= 3y x^4 + 2x \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} (5y^2 x^3 + 2y) \\ &= 10y x^3 + 2 \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} (3y x^4 + 2x) \\ &= 12y x^3 + 2 \end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= \frac{1}{3y x^4 + 2x} ((10y x^3 + 2) - (12y x^3 + 2)) \\ &= -\frac{2y x^2}{3y x^3 + 2} \end{aligned}$$

Since A depends on y , it can not be used to obtain an integrating factor. We will now try a second method to find an integrating factor. Let

$$\begin{aligned} B &= \frac{1}{M} \left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) \\ &= \frac{1}{5y^2x^3 + 2y} ((12yx^3 + 2) - (10yx^3 + 2)) \\ &= \frac{2x^3}{5yx^3 + 2} \end{aligned}$$

Since B depends on x , it can not be used to obtain an integrating factor. We will now try a third method to find an integrating factor. Let

$$R = \frac{\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y}}{xM - yN}$$

R is now checked to see if it is a function of only $t = xy$. Therefore

$$\begin{aligned} R &= \frac{\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y}}{xM - yN} \\ &= \frac{(12yx^3 + 2) - (10yx^3 + 2)}{x(5y^2x^3 + 2y) - y(3yx^4 + 2x)} \\ &= \frac{1}{yx} \end{aligned}$$

Replacing all powers of terms xy by t gives

$$R = \frac{1}{t}$$

Since R depends on t only, then it can be used to find an integrating factor. Let the integrating factor be μ then

$$\begin{aligned} \mu &= e^{\int R dt} \\ &= e^{\int \left(\frac{1}{t}\right) dt} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^{\ln(t)} \\ &= t \end{aligned}$$

Now t is replaced back with xy giving

$$\mu = xy$$

Multiplying M and N by this integrating factor gives new M and new N which are called \bar{M} and \bar{N} so not to confuse them with the original M and N

$$\begin{aligned}\bar{M} &= \mu M \\ &= xy(5y^2x^3 + 2y) \\ &= 5y^3x^4 + 2y^2x\end{aligned}$$

And

$$\begin{aligned}\bar{N} &= \mu N \\ &= xy(3yx^4 + 2x) \\ &= 3x^5y^2 + 2yx^2\end{aligned}$$

A modified ODE is now obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned}\bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ (5y^3x^4 + 2y^2x) + (3x^5y^2 + 2yx^2) \frac{dy}{dx} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int 5y^3x^4 + 2y^2x dx \\ \phi &= y^2x^2(yx^3 + 1) + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\begin{aligned}\frac{\partial \phi}{\partial y} &= 2yx^2(yx^3 + 1) + x^5y^2 + f'(y) \\ &= 3x^5y^2 + 2yx^2 + f'(y)\end{aligned} \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = 3x^5y^2 + 2yx^2$. Therefore equation (4) becomes

$$3x^5y^2 + 2yx^2 = 3x^5y^2 + 2yx^2 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = y^2x^2(yx^3 + 1) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = y^2x^2(yx^3 + 1)$$

Summary

The solution(s) found are the following

$$y^2x^2(yx^3 + 1) = c_1 \quad (1)$$




Figure 219: Slope field plot

Verification of solutions

$$y^2 x^2 (y x^3 + 1) = c_1$$

Verified OK.

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying homogeneous G
<- homogeneous successful`

```

✓ Solution by Maple

Time used: 0.515 (sec). Leaf size: 350

`dsolve((5*x^3*y(x)^2+2*y(x))+(3*x^4*y(x)+2*x)*diff(y(x),x)=0,y(x), singsol=all)`

$$\begin{aligned}
 y(x) &= \frac{12^{\frac{2}{3}} \left(12^{\frac{1}{3}} c_1^2 + \left((9x^2 + \sqrt{-12c_1^4 + 81x^4}) c_1 \right)^{\frac{2}{3}} \right)^2}{36c_1^2 \left((9x^2 + \sqrt{-12c_1^4 + 81x^4}) c_1 \right)^{\frac{2}{3}}} - 1 \\
 y(x) &= \frac{-\frac{c_1 \left((9x^2 + \sqrt{-12c_1^4 + 81x^4}) c_1 \right)^{\frac{2}{3}}}{3} + \frac{3 \cdot 2^{\frac{1}{3}} \left(x^2 + \frac{\sqrt{-12c_1^4 + 81x^4}}{9} \right) \left(i3^{\frac{1}{6}} - \frac{3^{\frac{2}{3}}}{3} \right) \left((9x^2 + \sqrt{-12c_1^4 + 81x^4}) c_1 \right)^{\frac{1}{3}}}{c_1 \left((9x^2 + \sqrt{-12c_1^4 + 81x^4}) c_1 \right)^{\frac{2}{3}} x^3} - \frac{(i3^{\frac{5}{6}} + 3^{\frac{1}{3}}) 2^{\frac{2}{3}} c_1^3}{6}}{3 \left(\frac{4c_1 \left((9x^2 + \sqrt{-12c_1^4 + 81x^4}) c_1 \right)^{\frac{2}{3}}}{9} + 2^{\frac{1}{3}} \left(x^2 + \frac{\sqrt{-12c_1^4 + 81x^4}}{9} \right) \left(i3^{\frac{1}{6}} + \frac{3^{\frac{2}{3}}}{3} \right) \left((9x^2 + \sqrt{-12c_1^4 + 81x^4}) c_1 \right)^{\frac{1}{3}} \right)} \\
 &= \frac{4 \left((9x^2 + \sqrt{-12c_1^4 + 81x^4}) c_1 \right)^{\frac{2}{3}} x^3 c_1}{4 \left((9x^2 + \sqrt{-12c_1^4 + 81x^4}) c_1 \right)^{\frac{2}{3}} x^3 c_1}
 \end{aligned}$$

✓ Solution by Mathematica

Time used: 49.208 (sec). Leaf size: 400

`DSolve[(5*x^3*y[x]^2+2*y[x])+(3*x^4*y[x]+2*x)*y'[x]==0,y[x],x,IncludeSingularSolutions -> Tr`

$$y(x) \rightarrow \frac{-2x^2 + \frac{2x^4}{\sqrt[3]{\frac{27c_1x^{10}}{2} - x^6 + \frac{3}{2}\sqrt{3}\sqrt{c_1x^{16}}(-4 + 27c_1x^4)}} + 2^{2/3}\sqrt[3]{27c_1x^{10} - 2x^6 + 3\sqrt{3}\sqrt{c_1x^{16}}(-4 + 27c_1x^4)}}}{6x^5}$$

$$y(x) \rightarrow \frac{-4x^2 - \frac{2(1+i\sqrt{3})x^4}{\sqrt[3]{\frac{27c_1x^{10}}{2} - x^6 + \frac{3}{2}\sqrt{3}\sqrt{c_1x^{16}}(-4 + 27c_1x^4)}} + i2^{2/3}(\sqrt{3} + i)\sqrt[3]{27c_1x^{10} - 2x^6 + 3\sqrt{3}\sqrt{c_1x^{16}}(-4 + 27c_1x^4)}}}{12x^5}$$

$$y(x) \rightarrow \frac{4x^2 - \frac{2i(\sqrt{3}+i)x^4}{\sqrt[3]{\frac{27c_1x^{10}}{2} - x^6 + \frac{3}{2}\sqrt{3}\sqrt{c_1x^{16}}(-4 + 27c_1x^4)}} + 2^{2/3}(1 + i\sqrt{3})\sqrt[3]{27c_1x^{10} - 2x^6 + 3\sqrt{3}\sqrt{c_1x^{16}}(-4 + 27c_1x^4)}}}{12x^5}$$

22.12 problem 2(d)

22.12.1 Solving as exact ode	1876
22.12.2 Maple step by step solution	1880

Internal problem ID [6090]

Internal file name [OUTPUT/5338_Sunday_June_05_2022_03_34_41_PM_88081771/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 5. Existence and uniqueness of solutions to first order equations. Page 198

Problem number: 2(d).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**exactWithIntegrationFactor**"

Maple gives the following as the ode type

[_quadrature]

$$e^y + x e^y + x e^y y' = 0$$

22.12.1 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned} (x e^y) dy &= (-e^y - x e^y) dx \\ (e^y + x e^y) dx + (x e^y) dy &= 0 \end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= e^y + x e^y \\ N(x, y) &= x e^y \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(e^y + x e^y) \\ &= e^y(1 + x) \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(x e^y) \\ &= e^y \end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= \frac{e^{-y}}{x} ((e^y + x e^y) - (e^y)) \\ &= 1 \end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int A dx} \\ &= e^{\int 1 dx}\end{aligned}$$

The result of integrating gives

$$\begin{aligned}\mu &= e^x \\ &= e^x\end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} for now so not to confuse them with the original M and N .

$$\begin{aligned}\bar{M} &= \mu M \\ &= e^x(e^y + x e^y) \\ &= (1 + x) e^{x+y}\end{aligned}$$

And

$$\begin{aligned}\bar{N} &= \mu N \\ &= e^x(x e^y) \\ &= x e^{x+y}\end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned}\bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ ((1 + x) e^{x+y}) + (x e^{x+y}) \frac{dy}{dx} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int (1 + x) e^{x+y} dx \\ \phi &= x e^{x+y} + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = x e^{x+y} + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = x e^{x+y}$. Therefore equation (4) becomes

$$x e^{x+y} = x e^{x+y} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = x e^{x+y} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = x e^{x+y}$$

The solution becomes

$$y = -x + \ln\left(\frac{c_1}{x}\right)$$

Summary

The solution(s) found are the following

$$y = -x + \ln\left(\frac{c_1}{x}\right) \quad (1)$$




Figure 220: Slope field plot

Verification of solutions

$$y = -x + \ln\left(\frac{c_1}{x}\right)$$

Verified OK.

22.12.2 Maple step by step solution

Let's solve

$$e^y + x e^y + x e^y y' = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$y' = -\frac{1+x}{x}$$

- Integrate both sides with respect to x

$$\int y' dx = \int -\frac{1+x}{x} dx + c_1$$

- Evaluate integral

$$y = -x - \ln(x) + c_1$$

- Solve for y

$$y = -x - \ln(x) + c_1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 13

```
dsolve((exp(y(x))+x*exp(y(x)))+(x*exp(y(x)))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = -x - \ln(x) + c_1$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 15

```
DSolve[(Exp[y[x]]+x*Exp[y[x]])+(x*Exp[y[x]])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x - \log(x) + c_1$$

23 Chapter 6. Existence and uniqueness of solutions to systems and nth order equations.

Page 238

23.1 problem 1(a)	1883
23.2 problem 1(b)	1902
23.3 problem 1(c)	1906
23.4 problem 1(d)	1912
23.5 problem 1(e)	1920
23.6 problem 1(f)	1928
23.7 problem 2	1950
23.8 problem 3	1957
23.9 problem 5(b)	1972
23.10problem 5(c)	1980

23.1 problem 1(a)

23.1.1 Solving as second order linear constant coeff ode	1883
23.1.2 Solving as second order integrable as is ode	1887
23.1.3 Solving as second order ode missing y ode	1889
23.1.4 Solving as type second_order_integrable_as_is (not using ABC version)	1890
23.1.5 Solving using Kovacic algorithm	1892
23.1.6 Solving as exact linear second order ode ode	1897
23.1.7 Maple step by step solution	1899

Internal problem ID [6091]

Internal file name [OUTPUT/5339_Sunday_June_05_2022_03_34_42_PM_3155877/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 6. Existence and uniqueness of solutions to systems and nth order equations. Page 238

Problem number: 1(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "exact linear second order ode", "second_order_integrable_as_is", "second_order_ode_missing_y", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' + y' = 1$$

23.1.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 1, C = 0, f(x) = 1$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 1, C = 0$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + \lambda e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + \lambda = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 1, C = 0$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{1^2 - (4)(1)(0)} \\ &= -\frac{1}{2} \pm \frac{1}{2} \end{aligned}$$

Hence

$$\lambda_1 = -\frac{1}{2} + \frac{1}{2}$$

$$\lambda_2 = -\frac{1}{2} - \frac{1}{2}$$

Which simplifies to

$$\lambda_1 = 0$$

$$\lambda_2 = -1$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(0)x} + c_2 e^{(-1)x}$$

Or

$$y = c_1 + c_2 e^{-x}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 + c_2 e^{-x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, e^{-x}\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_1 = 1$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = x$$

Therefore the general solution is

$$\begin{aligned}y &= y_h + y_p \\ &= (c_1 + c_2 e^{-x}) + (x)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 + c_2 e^{-x} + x \tag{1}$$




Figure 221: Slope field plot

Verification of solutions

$$y = c_1 + c_2 e^{-x} + x$$

Verified OK.

23.1.2 Solving as second order integrable as is ode

Integrating both sides of the ODE w.r.t x gives

$$\int (y'' + y') dx = \int 1 dx$$
$$y + y' = x + c_1$$

Which is now solved for y .

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = 1$$
$$q(x) = x + c_1$$

Hence the ode is

$$y + y' = x + c_1$$

The integrating factor μ is

$$\mu = e^{\int 1 dx}$$
$$= e^x$$

The ode becomes

$$\frac{d}{dx}(\mu y) = (\mu)(x + c_1)$$
$$\frac{d}{dx}(y e^x) = (e^x)(x + c_1)$$
$$d(y e^x) = (e^x(x + c_1)) dx$$

Integrating gives

$$y e^x = \int e^x(x + c_1) dx$$
$$y e^x = (-1 + x + c_1) e^x + c_2$$

Dividing both sides by the integrating factor $\mu = e^x$ results in

$$y = e^{-x}(-1 + x + c_1) e^x + c_2 e^{-x}$$

which simplifies to

$$y = -1 + x + c_1 + c_2 e^{-x}$$

Summary

The solution(s) found are the following

$$y = -1 + x + c_1 + c_2 e^{-x} \tag{1}$$




Figure 222: Slope field plot

Verification of solutions

$$y = -1 + x + c_1 + c_2 e^{-x}$$

Verified OK.

23.1.3 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p(x) + p'(x) - 1 = 0$$

Which is now solve for $p(x)$ as first order ode. Integrating both sides gives

$$\int \frac{1}{-p+1} dp = \int dx$$
$$-\ln(-p+1) = x + c_1$$

Raising both side to exponential gives

$$\frac{1}{-p+1} = e^{x+c_1}$$

Which simplifies to

$$\frac{1}{-p+1} = c_2 e^x$$

Since $p = y'$ then the new first order ode to solve is

$$y' = -\frac{e^{-x}}{c_2} + 1$$

Integrating both sides gives

$$y = \int \frac{(-1 + c_2 e^x) e^{-x}}{c_2} dx$$
$$= \ln(e^x) + \frac{e^{-x}}{c_2} + c_3$$

Summary

The solution(s) found are the following

$$y = \ln(e^x) + \frac{e^{-x}}{c_2} + c_3 \quad (1)$$




Figure 223: Slope field plot

Verification of solutions

$$y = \ln(e^x) + \frac{e^{-x}}{c_2} + c_3$$

Verified OK.

23.1.4 Solving as type second_order_integrable_as_is (not using ABC version)

Writing the ode as

$$y'' + y' = 1$$

Integrating both sides of the ODE w.r.t x gives

$$\int (y'' + y') dx = \int 1 dx$$

$$y + y' = x + c_1$$

Which is now solved for y .

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$\begin{aligned}p(x) &= 1 \\q(x) &= x + c_1\end{aligned}$$

Hence the ode is

$$y + y' = x + c_1$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int 1 dx} \\ &= e^x\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu)(x + c_1) \\ \frac{d}{dx}(y e^x) &= (e^x)(x + c_1) \\ d(y e^x) &= (e^x(x + c_1)) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}y e^x &= \int e^x(x + c_1) dx \\ y e^x &= (-1 + x + c_1) e^x + c_2\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^x$ results in

$$y = e^{-x}(-1 + x + c_1) e^x + c_2 e^{-x}$$

which simplifies to

$$y = -1 + x + c_1 + c_2 e^{-x}$$

Summary

The solution(s) found are the following

$$y = -1 + x + c_1 + c_2 e^{-x} \tag{1}$$




Figure 224: Slope field plot

Verification of solutions

$$y = -1 + x + c_1 + c_2 e^{-x}$$

Verified OK.

23.1.5 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y' = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 1 \\ C &= 0 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{1}{4} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 1 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \frac{z(x)}{4} \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 274: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0\end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{1}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{x}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{1}{1} dx} \\ &= z_1 e^{-\frac{x}{2}} \\ &= z_1 (e^{-\frac{x}{2}})\end{aligned}$$

Which simplifies to

$$y_1 = e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{1}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-x}}{(y_1)^2} dx \\ &= y_1 (e^x)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-x}) + c_2 (e^{-x}(e^x))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-x} + c_2$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, e^{-x}\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_1 = 1$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^{-x} + c_2) + (x) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-x} + c_2 + x \tag{1}$$




Figure 225: Slope field plot

Verification of solutions

$$y = c_1 e^{-x} + c_2 + x$$

Verified OK.

23.1.6 Solving as exact linear second order ode

An ode of the form

$$p(x) y'' + q(x) y' + r(x) y = s(x)$$

is exact if

$$p''(x) - q'(x) + r(x) = 0 \tag{1}$$

For the given ode we have

$$p(x) = 1$$

$$q(x) = 1$$

$$r(x) = 0$$

$$s(x) = 1$$

Hence

$$p''(x) = 0$$

$$q'(x) = 0$$

Therefore (1) becomes

$$0 - (0) + (0) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(x) y' + (q(x) - p'(x)) y)' = s(x)$$

Integrating gives

$$p(x) y' + (q(x) - p'(x)) y = \int s(x) dx$$

Substituting the above values for p, q, r, s gives

$$y + y' = \int 1 dx$$

We now have a first order ode to solve which is

$$y + y' = x + c_1$$

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = 1$$

$$q(x) = x + c_1$$

Hence the ode is

$$y + y' = x + c_1$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int 1 dx} \\ &= e^x\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu)(x + c_1) \\ \frac{d}{dx}(y e^x) &= (e^x)(x + c_1) \\ d(y e^x) &= (e^x(x + c_1)) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}y e^x &= \int e^x(x + c_1) dx \\ y e^x &= (-1 + x + c_1) e^x + c_2\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^x$ results in

$$y = e^{-x}(-1 + x + c_1) e^x + c_2 e^{-x}$$

which simplifies to

$$y = -1 + x + c_1 + c_2 e^{-x}$$

Summary

The solution(s) found are the following

$$y = -1 + x + c_1 + c_2 e^{-x} \quad (1)$$




Figure 226: Slope field plot

Verification of solutions

$$y = -1 + x + c_1 + c_2 e^{-x}$$

Verified OK.

23.1.7 Maple step by step solution

Let's solve

$$y' + y'' = 1$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + r = 0$$

- Factor the characteristic polynomial
 $r(r + 1) = 0$
- Roots of the characteristic polynomial
 $r = (-1, 0)$
- 1st solution of the homogeneous ODE
 $y_1(x) = e^{-x}$
- 2nd solution of the homogeneous ODE
 $y_2(x) = 1$
- General solution of the ODE
 $y = c_1y_1(x) + c_2y_2(x) + y_p(x)$
- Substitute in solutions of the homogeneous ODE
 $y = c_1e^{-x} + c_2 + y_p(x)$
- Find a particular solution $y_p(x)$ of the ODE
 - Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = 1 \right]$$
 - Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-x} & 1 \\ -e^{-x} & 0 \end{bmatrix}$$
 - Compute Wronskian
 $W(y_1(x), y_2(x)) = e^{-x}$
 - Substitute functions into equation for $y_p(x)$
 $y_p(x) = -e^{-x} \left(\int e^x dx \right) + \int 1 dx$
 - Compute integrals
 $y_p(x) = x - 1$
- Substitute particular solution into general solution to ODE
 $y = c_1e^{-x} + c_2 + x - 1$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
-> Calling odsolve with the ODE`, diff(_b(_a), _a) = -_b(_a)+1, _b(_a)` *** Sublevel 2 ***  
  Methods for first order ODEs:  
    --- Trying classification methods ---  
    trying a quadrature  
    trying 1st order linear  
    <- 1st order linear successful  
  <- high order exact linear fully integrable successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(diff(y(x),x$2)+diff(y(x),x)=1,y(x), singsol=all)
```

$$y(x) = -c_1 e^{-x} + x + c_2$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 18

```
DSolve[y''[x]+y'[x]==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x - c_1 e^{-x} + c_2$$

23.2 problem 1(b)

23.2.1 Solving as second order ode missing y ode 1902

23.2.2 Maple step by step solution 1904

Internal problem ID [6092]

Internal file name [OUTPUT/5340_Sunday_June_05_2022_03_34_43_PM_40990985/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 6. Existence and uniqueness of solutions to systems and nth order equations. Page 238

Problem number: 1(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_ode_missing_y**"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_y]]
```

$$y'' + y'e^x = e^x$$

23.2.1 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x) + e^x p(x) - e^x = 0$$

Which is now solve for $p(x)$ as first order ode. In canonical form the ODE is

$$\begin{aligned} p' &= F(x, p) \\ &= f(x)g(p) \\ &= e^x(-p + 1) \end{aligned}$$

Where $f(x) = e^x$ and $g(p) = -p + 1$. Integrating both sides gives

$$\begin{aligned}\frac{1}{-p+1} dp &= e^x dx \\ \int \frac{1}{-p+1} dp &= \int e^x dx \\ -\ln(p-1) &= e^x + c_1\end{aligned}$$

Raising both side to exponential gives

$$\frac{1}{p-1} = e^{e^x+c_1}$$

Which simplifies to

$$\frac{1}{p-1} = c_2 e^{e^x}$$

Since $p = y'$ then the new first order ode to solve is

$$y' = \frac{(c_2 e^{e^x+c_1} + 1) e^{-e^x-c_1}}{c_2}$$

Integrating both sides gives

$$\begin{aligned}y &= \int \frac{(c_2 e^{e^x+c_1} + 1) e^{-e^x-c_1}}{c_2} dx \\ &= \ln(e^x) - \frac{e^{-c_1} \text{expIntegral}_1(e^x)}{c_2} + c_3\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \ln(e^x) - \frac{e^{-c_1} \text{expIntegral}_1(e^x)}{c_2} + c_3 \quad (1)$$

Verification of solutions

$$y = \ln(e^x) - \frac{e^{-c_1} \text{expIntegral}_1(e^x)}{c_2} + c_3$$

Verified OK.

23.2.2 Maple step by step solution

Let's solve

$$y'' + y'e^x = e^x$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Make substitution $u = y'$ to reduce order of ODE

$$u'(x) + u(x)e^x = e^x$$

- Separate variables

$$\frac{u'(x)}{u(x)-1} = -e^x$$

- Integrate both sides with respect to x

$$\int \frac{u'(x)}{u(x)-1} dx = \int -e^x dx + c_1$$

- Evaluate integral

$$\ln(u(x) - 1) = -e^x + c_1$$

- Solve for $u(x)$

$$u(x) = e^{-e^x+c_1} + 1$$

- Solve 1st ODE for $u(x)$

$$u(x) = e^{-e^x+c_1} + 1$$

- Make substitution $u = y'$

$$y' = e^{-e^x+c_1} + 1$$

- Integrate both sides to solve for y

$$\int y' dx = \int (e^{-e^x+c_1} + 1) dx + c_2$$

- Compute integrals

$$y = x - e^{c_1} \text{Ei}_1(e^x) + c_2$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
-> Calling odsolve with the ODE`, diff(_b(_a), _a) = -exp(_a)*_b(_a)+exp(_a), _b(_a)` ***  
  Methods for first order ODEs:  
    --- Trying classification methods ---  
    trying a quadrature  
    trying 1st order linear  
    <- 1st order linear successful  
  <- high order exact linear fully integrable successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(diff(y(x),x$2)+exp(x)*diff(y(x),x)=exp(x),y(x), singsol=all)
```

$$y(x) = -c_1 \operatorname{ExpIntegral}_1(e^x) + x + c_2$$

✓ Solution by Mathematica

Time used: 0.081 (sec). Leaf size: 18

```
DSolve[y''[x]+Exp[x]*y'[x]==Exp[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \operatorname{ExpIntegralEi}(-e^x) + x + c_2$$

23.3 problem 1(c)

23.3.1 Solving as second order ode missing x ode 1906

23.3.2 Maple step by step solution 1909

Internal problem ID [6093]

Internal file name [OUTPUT/5341_Sunday_June_05_2022_03_34_45_PM_92580117/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 6. Existence and uniqueness of solutions to systems and nth order equations. Page 238

Problem number: 1(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_ode_missing_x**"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x], _Liouville, [_2nd_order, _reducible,
    _mu_x_y1], [_2nd_order, _reducible, _mu_xy]]
```

$$yy'' + 4y'^2 = 0$$

23.3.1 Solving as second order ode missing x ode

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable y an independent variable.

Using

$$y' = p(y)$$

Then

$$\begin{aligned} y'' &= \frac{dp}{dx} \\ &= \frac{dy}{dx} \frac{dp}{dy} \\ &= p \frac{dp}{dy} \end{aligned}$$

Hence the ode becomes

$$yp(y) \left(\frac{d}{dy} p(y) \right) + 4p(y)^2 = 0$$

Which is now solved as first order ode for $p(y)$. In canonical form the ODE is

$$\begin{aligned} p' &= F(y, p) \\ &= f(y)g(p) \\ &= -\frac{4p}{y} \end{aligned}$$

Where $f(y) = -\frac{4}{y}$ and $g(p) = p$. Integrating both sides gives

$$\begin{aligned} \frac{1}{p} dp &= -\frac{4}{y} dy \\ \int \frac{1}{p} dp &= \int -\frac{4}{y} dy \\ \ln(p) &= -4 \ln(y) + c_1 \\ p &= e^{-4 \ln(y) + c_1} \\ &= \frac{c_1}{y^4} \end{aligned}$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = \frac{c_1}{y^4}$$

Integrating both sides gives

$$\begin{aligned} \int \frac{y^4}{c_1} dy &= c_2 + x \\ \frac{y^5}{5c_1} &= c_2 + x \end{aligned}$$

Solving for y gives these solutions

$$y_1 = (5c_1c_2 + 5c_1x)^{\frac{1}{5}}$$

$$y_2 = \left(-\frac{\sqrt{5}}{4} - \frac{1}{4} - \frac{i\sqrt{2}\sqrt{5-\sqrt{5}}}{4} \right) (5c_1c_2 + 5c_1x)^{\frac{1}{5}}$$

$$y_3 = \left(-\frac{\sqrt{5}}{4} - \frac{1}{4} + \frac{i\sqrt{2}\sqrt{5-\sqrt{5}}}{4} \right) (5c_1c_2 + 5c_1x)^{\frac{1}{5}}$$

$$y_4 = \left(\frac{\sqrt{5}}{4} - \frac{1}{4} - \frac{i\sqrt{2}\sqrt{5+\sqrt{5}}}{4} \right) (5c_1c_2 + 5c_1x)^{\frac{1}{5}}$$

$$y_5 = \left(\frac{\sqrt{5}}{4} - \frac{1}{4} + \frac{i\sqrt{2}\sqrt{5+\sqrt{5}}}{4} \right) (5c_1c_2 + 5c_1x)^{\frac{1}{5}}$$

Summary

The solution(s) found are the following

$$y = (5c_1c_2 + 5c_1x)^{\frac{1}{5}} \tag{1}$$

$$y = \left(-\frac{\sqrt{5}}{4} - \frac{1}{4} - \frac{i\sqrt{2}\sqrt{5-\sqrt{5}}}{4} \right) (5c_1c_2 + 5c_1x)^{\frac{1}{5}} \tag{2}$$

$$y = \left(-\frac{\sqrt{5}}{4} - \frac{1}{4} + \frac{i\sqrt{2}\sqrt{5-\sqrt{5}}}{4} \right) (5c_1c_2 + 5c_1x)^{\frac{1}{5}} \tag{3}$$

$$y = \left(\frac{\sqrt{5}}{4} - \frac{1}{4} - \frac{i\sqrt{2}\sqrt{5+\sqrt{5}}}{4} \right) (5c_1c_2 + 5c_1x)^{\frac{1}{5}} \tag{4}$$

$$y = \left(\frac{\sqrt{5}}{4} - \frac{1}{4} + \frac{i\sqrt{2}\sqrt{5+\sqrt{5}}}{4} \right) (5c_1c_2 + 5c_1x)^{\frac{1}{5}} \tag{5}$$

Verification of solutions

$$y = (5c_1c_2 + 5c_1x)^{\frac{1}{5}}$$

Verified OK.

$$y = \left(-\frac{\sqrt{5}}{4} - \frac{1}{4} - \frac{i\sqrt{2}\sqrt{5-\sqrt{5}}}{4} \right) (5c_1c_2 + 5c_1x)^{\frac{1}{5}}$$

Verified OK.

$$y = \left(-\frac{\sqrt{5}}{4} - \frac{1}{4} + \frac{i\sqrt{2}\sqrt{5-\sqrt{5}}}{4} \right) (5c_1c_2 + 5c_1x)^{\frac{1}{5}}$$

Verified OK.

$$y = \left(\frac{\sqrt{5}}{4} - \frac{1}{4} - \frac{i\sqrt{2}\sqrt{5+\sqrt{5}}}{4} \right) (5c_1c_2 + 5c_1x)^{\frac{1}{5}}$$

Verified OK.

$$y = \left(\frac{\sqrt{5}}{4} - \frac{1}{4} + \frac{i\sqrt{2}\sqrt{5+\sqrt{5}}}{4} \right) (5c_1c_2 + 5c_1x)^{\frac{1}{5}}$$

Verified OK.

23.3.2 Maple step by step solution

Let's solve

$$yy'' + 4y'^2 = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Define new dependent variable u

$$u(x) = y'$$

- Compute y''

$$u'(x) = y''$$

- Use chain rule on the lhs

$$y' \left(\frac{d}{dy} u(y) \right) = y''$$

- Substitute in the definition of u

$$u(y) \left(\frac{d}{dy} u(y) \right) = y''$$

- Make substitutions $y' = u(y)$, $y'' = u(y) \left(\frac{d}{dy} u(y) \right)$ to reduce order of ODE

$$y u(y) \left(\frac{d}{dy} u(y) \right) + 4u(y)^2 = 0$$

- Separate variables

$$\frac{\frac{d}{dy} u(y)}{u(y)} = -\frac{4}{y}$$

- Integrate both sides with respect to y

$$\int \frac{\frac{d}{dy} u(y)}{u(y)} dy = \int -\frac{4}{y} dy + c_1$$

- Evaluate integral

$$\ln(u(y)) = -4 \ln(y) + c_1$$

- Solve for $u(y)$

$$u(y) = \frac{e^{c_1}}{y^4}$$

- Solve 1st ODE for $u(y)$

$$u(y) = \frac{e^{c_1}}{y^4}$$

- Revert to original variables with substitution $u(y) = y'$, $y = y$

$$y' = \frac{e^{c_1}}{y^4}$$

- Separate variables

$$y' y^4 = e^{c_1}$$

- Integrate both sides with respect to x

$$\int y' y^4 dx = \int e^{c_1} dx + c_2$$

- Evaluate integral

$$\frac{y^5}{5} = x e^{c_1} + c_2$$

- Solve for y

$$y = (5x e^{c_1} + 5c_2)^{\frac{1}{5}}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying 2nd order Liouville  
<- 2nd_order Liouville successful`
```

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 158

```
dsolve(y(x)*diff(y(x),x$2)+4*diff(y(x),x)^2=0,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = (5c_1x + 5c_2)^{\frac{1}{5}}$$

$$y(x) = -\frac{(i\sqrt{2}\sqrt{5-\sqrt{5}} + \sqrt{5} + 1)(5c_1x + 5c_2)^{\frac{1}{5}}}{4}$$

$$y(x) = \frac{(i\sqrt{2}\sqrt{5-\sqrt{5}} - \sqrt{5} - 1)(5c_1x + 5c_2)^{\frac{1}{5}}}{4}$$

$$y(x) = -\frac{(i\sqrt{2}\sqrt{5+\sqrt{5}} - \sqrt{5} + 1)(5c_1x + 5c_2)^{\frac{1}{5}}}{4}$$

$$y(x) = \frac{(i\sqrt{2}\sqrt{5+\sqrt{5}} + \sqrt{5} - 1)(5c_1x + 5c_2)^{\frac{1}{5}}}{4}$$

✓ Solution by Mathematica

Time used: 0.178 (sec). Leaf size: 20

```
DSolve[y[x]*y'[x]+4*(y'[x])^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2\sqrt[5]{5x - c_1}$$

23.4 problem 1(d)

23.4.1 Solving as second order linear constant coeff ode	1912
23.4.2 Solving as second order ode can be made integrable ode	1914
23.4.3 Solving using Kovacic algorithm	1915
23.4.4 Maple step by step solution	1918

Internal problem ID [6094]

Internal file name [OUTPUT/5342_Sunday_June_05_2022_03_34_47_PM_31515423/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 6. Existence and uniqueness of solutions to systems and nth order equations. Page 238

Problem number: 1(d).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "second_order_ode_can_be_made_integrable"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x]]
```

$$y'' + k^2y = 0$$

23.4.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = k^2$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + k^2 e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$k^2 + \lambda^2 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = k^2$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(k^2)} \\ &= \pm \sqrt{-k^2} \end{aligned}$$

Hence

$$\lambda_1 = +\sqrt{-k^2}$$

$$\lambda_2 = -\sqrt{-k^2}$$

Which simplifies to

$$\lambda_1 = \sqrt{-k^2}$$

$$\lambda_2 = -\sqrt{-k^2}$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(\sqrt{-k^2})x} + c_2 e^{(-\sqrt{-k^2})x}$$

Or

$$y = c_1 e^{\sqrt{-k^2}x} + c_2 e^{-\sqrt{-k^2}x}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{\sqrt{-k^2}x} + c_2 e^{-\sqrt{-k^2}x} \quad (1)$$

Verification of solutions

$$y = c_1 e^{\sqrt{-k^2}x} + c_2 e^{-\sqrt{-k^2}x}$$

Verified OK.

23.4.2 Solving as second order ode can be made integrable ode

Multiplying the ode by y' gives

$$y'y'' + y'k^2y = 0$$

Integrating the above w.r.t x gives

$$\int (y'y'' + y'k^2y) dx = 0$$
$$\frac{y'^2}{2} + \frac{y^2k^2}{2} = c_2$$

Which is now solved for y . Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \sqrt{-y^2k^2 + 2c_1} \quad (1)$$

$$y' = -\sqrt{-y^2k^2 + 2c_1} \quad (2)$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int \frac{1}{\sqrt{-y^2k^2 + 2c_1}} dy = \int dx$$
$$\frac{\arctan\left(\frac{\sqrt{k^2}y}{\sqrt{-y^2k^2+2c_1}}\right)}{\sqrt{k^2}} = c_2 + x$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{1}{\sqrt{-y^2k^2 + 2c_1}} dy = \int dx$$
$$-\frac{\arctan\left(\frac{\sqrt{k^2}y}{\sqrt{-y^2k^2+2c_1}}\right)}{\sqrt{k^2}} = x + c_3$$

Summary

The solution(s) found are the following

$$\frac{\arctan\left(\frac{\sqrt{k^2}y}{\sqrt{-y^2k^2+2c_1}}\right)}{\sqrt{k^2}} = c_2 + x \quad (1)$$

$$-\frac{\arctan\left(\frac{\sqrt{k^2}y}{\sqrt{-y^2k^2+2c_1}}\right)}{\sqrt{k^2}} = x + c_3 \quad (2)$$

Verification of solutions

$$\frac{\arctan\left(\frac{\sqrt{k^2}y}{\sqrt{-y^2k^2+2c_1}}\right)}{\sqrt{k^2}} = c_2 + x$$

Verified OK.

$$-\frac{\arctan\left(\frac{\sqrt{k^2}y}{\sqrt{-y^2k^2+2c_1}}\right)}{\sqrt{k^2}} = x + c_3$$

Verified OK.

23.4.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + k^2y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= k^2 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-k^2}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -k^2 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = (-k^2) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 278: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -k^2$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{\sqrt{-k^2} x}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned} y_1 &= z_1 \\ &= e^{\sqrt{-k^2} x} \end{aligned}$$

Which simplifies to

$$y_1 = e^{\sqrt{-k^2} x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= e^{\sqrt{-k^2} x} \int \frac{1}{e^{2\sqrt{-k^2} x}} dx \\ &= e^{\sqrt{-k^2} x} \left(\frac{\sqrt{-k^2} e^{-2\sqrt{-k^2} x}}{2k^2} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}
y &= c_1 y_1 + c_2 y_2 \\
&= c_1 \left(e^{\sqrt{-k^2} x} \right) + c_2 \left(e^{\sqrt{-k^2} x} \left(\frac{\sqrt{-k^2} e^{-2\sqrt{-k^2} x}}{2k^2} \right) \right)
\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{\sqrt{-k^2} x} + \frac{c_2 \sqrt{-k^2} e^{-\sqrt{-k^2} x}}{2k^2} \quad (1)$$

Verification of solutions

$$y = c_1 e^{\sqrt{-k^2} x} + \frac{c_2 \sqrt{-k^2} e^{-\sqrt{-k^2} x}}{2k^2}$$

Verified OK.

23.4.4 Maple step by step solution

Let's solve

$$y'' + k^2 y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$k^2 + r^2 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4k^2})}{2}$$

- Roots of the characteristic polynomial

$$r = (\sqrt{-k^2}, -\sqrt{-k^2})$$

- 1st solution of the ODE

$$y_1(x) = e^{\sqrt{-k^2} x}$$

- 2nd solution of the ODE

$$y_2(x) = e^{-\sqrt{-k^2} x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = c_1 e^{\sqrt{-k^2} x} + c_2 e^{-\sqrt{-k^2} x}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)+k^2*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 \sin(kx) + c_2 \cos(kx)$$

✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 20

```
DSolve[y''[x]+k^2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos(kx) + c_2 \sin(kx)$$

23.5 problem 1(e)

23.5.1 Solving as second order integrable as is ode	1920
23.5.2 Solving as second order ode missing x ode	1921
23.5.3 Solving as type second_order_integrable_as_is (not using ABC version)	1923
23.5.4 Solving as exact nonlinear second order ode ode	1924
23.5.5 Maple step by step solution	1925

Internal problem ID [6095]

Internal file name [OUTPUT/5343_Sunday_June_05_2022_03_34_48_PM_47713477/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 6. Existence and uniqueness of solutions to systems and nth order equations. Page 238

Problem number: 1(e).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_integrable_as_is", "second_order_ode_missing_x", "exact nonlinear second order ode"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x], [_2nd_order, _exact, _nonlinear],  
  _Lagerstrom, [_2nd_order, _reducible, _mu_x_y1], [_2nd_order,  
  _reducible, _mu_xy]]
```

$$y'' - y'y = 0$$

23.5.1 Solving as second order integrable as is ode

Integrating both sides of the ODE w.r.t x gives

$$\int (y'' - y'y) dx = 0$$
$$-\frac{y^2}{2} + y' = c_1$$

Which is now solved for y . Integrating both sides gives

$$\int \frac{1}{\frac{y^2}{2} + c_1} dy = c_2 + x$$

$$\frac{\sqrt{2} \arctan\left(\frac{y\sqrt{2}}{2\sqrt{c_1}}\right)}{\sqrt{c_1}} = c_2 + x$$

Solving for y gives these solutions

$$y_1 = \tan\left(\frac{\sqrt{c_1}(c_2 + x)\sqrt{2}}{2}\right) \sqrt{c_1} \sqrt{2}$$

Summary

The solution(s) found are the following

$$y = \tan\left(\frac{\sqrt{c_1}(c_2 + x)\sqrt{2}}{2}\right) \sqrt{c_1} \sqrt{2} \quad (1)$$

Verification of solutions

$$y = \tan\left(\frac{\sqrt{c_1}(c_2 + x)\sqrt{2}}{2}\right) \sqrt{c_1} \sqrt{2}$$

Verified OK.

23.5.2 Solving as second order ode missing x ode

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable y an independent variable. Using

$$y' = p(y)$$

Then

$$y'' = \frac{dp}{dx}$$

$$= \frac{dy}{dx} \frac{dp}{dy}$$

$$= p \frac{dp}{dy}$$

Hence the ode becomes

$$p(y) \left(\frac{d}{dy} p(y) \right) - p(y) y = 0$$

Which is now solved as first order ode for $p(y)$. Integrating both sides gives

$$\begin{aligned} p(y) &= \int y \, dy \\ &= \frac{y^2}{2} + c_1 \end{aligned}$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = \frac{y^2}{2} + c_1$$

Integrating both sides gives

$$\begin{aligned} \int \frac{1}{\frac{y^2}{2} + c_1} dy &= c_2 + x \\ \frac{\sqrt{2} \arctan\left(\frac{y\sqrt{2}}{2\sqrt{c_1}}\right)}{\sqrt{c_1}} &= c_2 + x \end{aligned}$$

Solving for y gives these solutions

$$y_1 = \tan\left(\frac{\sqrt{c_1}(c_2 + x)\sqrt{2}}{2}\right) \sqrt{c_1} \sqrt{2}$$

Summary

The solution(s) found are the following

$$y = \tan\left(\frac{\sqrt{c_1}(c_2 + x)\sqrt{2}}{2}\right) \sqrt{c_1} \sqrt{2} \quad (1)$$

Verification of solutions

$$y = \tan\left(\frac{\sqrt{c_1}(c_2 + x)\sqrt{2}}{2}\right) \sqrt{c_1} \sqrt{2}$$

Verified OK.

23.5.3 Solving as type second_order_integrable_as_is (not using ABC version)

Writing the ode as

$$y'' - y'y = 0$$

Integrating both sides of the ODE w.r.t x gives

$$\int (y'' - y'y) dx = 0$$
$$-\frac{y^2}{2} + y' = c_1$$

Which is now solved for y . Integrating both sides gives

$$\int \frac{1}{\frac{y^2}{2} + c_1} dy = c_2 + x$$
$$\frac{\sqrt{2} \arctan\left(\frac{y\sqrt{2}}{2\sqrt{c_1}}\right)}{\sqrt{c_1}} = c_2 + x$$

Solving for y gives these solutions

$$y_1 = \tan\left(\frac{\sqrt{c_1}(c_2 + x)\sqrt{2}}{2}\right) \sqrt{c_1} \sqrt{2}$$

Summary

The solution(s) found are the following

$$y = \tan\left(\frac{\sqrt{c_1}(c_2 + x)\sqrt{2}}{2}\right) \sqrt{c_1} \sqrt{2} \quad (1)$$

Verification of solutions

$$y = \tan\left(\frac{\sqrt{c_1}(c_2 + x)\sqrt{2}}{2}\right) \sqrt{c_1} \sqrt{2}$$

Verified OK.

23.5.4 Solving as exact nonlinear second order ode

An exact non-linear second order ode has the form

$$a_2(x, y, y') y'' + a_1(x, y, y') y' + a_0(x, y, y') = 0$$

Where the following conditions are satisfied

$$\begin{aligned}\frac{\partial a_2}{\partial y} &= \frac{\partial a_1}{\partial y'} \\ \frac{\partial a_2}{\partial x} &= \frac{\partial a_0}{\partial y'} \\ \frac{\partial a_1}{\partial x} &= \frac{\partial a_0}{\partial y}\end{aligned}$$

Looking at the the ode given we see that

$$\begin{aligned}a_2 &= 1 \\ a_1 &= -y \\ a_0 &= 0\end{aligned}$$

Applying the conditions to the above shows this is a nonlinear exact second order ode. Therefore it can be reduced to first order ode given by

$$\begin{aligned}\int a_2 dy' + \int a_1 dy + \int a_0 dx &= c_1 \\ \int 1 dy' + \int -y dy + \int 0 dx &= c_1\end{aligned}$$

Which results in

$$-\frac{y^2}{2} + y' = c_1$$

Which is now solved Integrating both sides gives

$$\begin{aligned}\int \frac{1}{\frac{y^2}{2} + c_1} dy &= c_2 + x \\ \frac{\sqrt{2} \arctan\left(\frac{y\sqrt{2}}{2\sqrt{c_1}}\right)}{\sqrt{c_1}} &= c_2 + x\end{aligned}$$

Solving for y gives these solutions

$$y_1 = \tan\left(\frac{\sqrt{c_1}(c_2 + x)\sqrt{2}}{2}\right) \sqrt{c_1} \sqrt{2}$$

Summary

The solution(s) found are the following

$$y = \tan \left(\frac{\sqrt{c_1} (c_2 + x) \sqrt{2}}{2} \right) \sqrt{c_1} \sqrt{2} \quad (1)$$

Verification of solutions

$$y = \tan \left(\frac{\sqrt{c_1} (c_2 + x) \sqrt{2}}{2} \right) \sqrt{c_1} \sqrt{2}$$

Verified OK.

23.5.5 Maple step by step solution

Let's solve

$$y'' - y'y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Define new dependent variable u

$$u(x) = y'$$

- Compute y''

$$u'(x) = y''$$

- Use chain rule on the lhs

$$y' \left(\frac{d}{dy} u(y) \right) = y''$$

- Substitute in the definition of u

$$u(y) \left(\frac{d}{dy} u(y) \right) = y''$$

- Make substitutions $y' = u(y)$, $y'' = u(y) \left(\frac{d}{dy} u(y) \right)$ to reduce order of ODE

$$u(y) \left(\frac{d}{dy} u(y) \right) - u(y) y = 0$$

- Separate variables

$$\frac{d}{dy} u(y) = y$$

- Integrate both sides with respect to y

$$\int \left(\frac{d}{dy} u(y) \right) dy = \int y dy + c_1$$

- Evaluate integral

$$u(y) = \frac{y^2}{2} + c_1$$
- Solve for $u(y)$

$$u(y) = \frac{y^2}{2} + c_1$$
- Solve 1st ODE for $u(y)$

$$u(y) = \frac{y^2}{2} + c_1$$
- Revert to original variables with substitution $u(y) = y', y = y$

$$y' = \frac{y^2}{2} + c_1$$
- Separate variables

$$\frac{y'}{\frac{y^2}{2} + c_1} = 1$$
- Integrate both sides with respect to x

$$\int \frac{y'}{\frac{y^2}{2} + c_1} dx = \int 1 dx + c_2$$
- Evaluate integral

$$\frac{\sqrt{2} \arctan\left(\frac{y\sqrt{2}}{2\sqrt{c_1}}\right)}{\sqrt{c_1}} = c_2 + x$$
- Solve for y

$$y = \tan\left(\frac{\sqrt{c_1}(c_2+x)\sqrt{2}}{2}\right) \sqrt{c_1} \sqrt{2}$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
-> Calling odsolve with the ODE`, (diff(_b(_a), _a))*_b(_a)-_b(_a)*_a = 0, _b(_a), HINT = []
    symmetry methods on request
`, `1st order, trying reduction of order with given symmetries: `[_a, 2*_b]
```

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 23

```
dsolve(diff(y(x),x$2)=y(x)*diff(y(x),x),y(x), singsol=all)
```

$$y(x) = \frac{\tan\left(\frac{(x+c_2)\sqrt{2}}{2c_1}\right)\sqrt{2}}{c_1}$$

✓ Solution by Mathematica

Time used: 16.739 (sec). Leaf size: 34

```
DSolve[y''[x]==y[x]*y'[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \sqrt{2}\sqrt{c_1} \tan\left(\frac{\sqrt{c_1}(x+c_2)}{\sqrt{2}}\right)$$

23.6 problem 1(f)

- 23.6.1 Solving as second order integrable as is ode 1929
- 23.6.2 Solving as second order ode missing y ode 1930
- 23.6.3 Solving as second order ode non constant coeff transformation
on B ode 1932
- 23.6.4 Solving as type second_order_integrable_as_is (not using ABC
version) 1936
- 23.6.5 Solving using Kovacic algorithm 1938
- 23.6.6 Solving as exact linear second order ode ode 1945
- 23.6.7 Maple step by step solution 1947

Internal problem ID [6096]

Internal file name [OUTPUT/5344_Sunday_June_05_2022_03_34_50_PM_22943806/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 6. Existence and uniqueness of solutions to systems and nth order equations. Page 238

Problem number: 1(f).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "exact linear second order ode", "second_order_integrable_as_is", "second_order_ode_missing_y", "second_order_ode_non_constant_coeff_transformation_on_B"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_y]]
```

$$xy'' - 2y' = x^3$$

23.6.1 Solving as second order integrable as is ode

Integrating both sides of the ODE w.r.t x gives

$$\int (xy'' - 2y') dx = \int x^3 dx$$
$$xy' - 3y = \frac{x^4}{4} + c_1$$

Which is now solved for y .

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -\frac{3}{x}$$
$$q(x) = \frac{x^4 + 4c_1}{4x}$$

Hence the ode is

$$y' - \frac{3y}{x} = \frac{x^4 + 4c_1}{4x}$$

The integrating factor μ is

$$\mu = e^{\int -\frac{3}{x} dx}$$
$$= \frac{1}{x^3}$$

The ode becomes

$$\frac{d}{dx}(\mu y) = (\mu) \left(\frac{x^4 + 4c_1}{4x} \right)$$
$$\frac{d}{dx} \left(\frac{y}{x^3} \right) = \left(\frac{1}{x^3} \right) \left(\frac{x^4 + 4c_1}{4x} \right)$$
$$d \left(\frac{y}{x^3} \right) = \left(\frac{x^4 + 4c_1}{4x^4} \right) dx$$

Integrating gives

$$\frac{y}{x^3} = \int \frac{x^4 + 4c_1}{4x^4} dx$$
$$\frac{y}{x^3} = \frac{x}{4} - \frac{c_1}{3x^3} + c_2$$

Dividing both sides by the integrating factor $\mu = \frac{1}{x^3}$ results in

$$y = x^3 \left(\frac{x}{4} - \frac{c_1}{3x^3} \right) + c_2 x^3$$

which simplifies to

$$y = \frac{1}{4}x^4 - \frac{1}{3}c_1 + c_2x^3$$

Summary

The solution(s) found are the following

$$y = \frac{1}{4}x^4 - \frac{1}{3}c_1 + c_2x^3 \quad (1)$$

Verification of solutions

$$y = \frac{1}{4}x^4 - \frac{1}{3}c_1 + c_2x^3$$

Verified OK.

23.6.2 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x)x - 2p(x) - x^3 = 0$$

Which is now solve for $p(x)$ as first order ode.

Entering Linear first order ODE solver. In canonical form a linear first order is

$$p'(x) + p(x)p(x) = q(x)$$

Where here

$$p(x) = -\frac{2}{x}$$
$$q(x) = x^2$$

Hence the ode is

$$p'(x) - \frac{2p(x)}{x} = x^2$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int -\frac{2}{x} dx} \\ &= \frac{1}{x^2}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu p) &= (\mu)(x^2) \\ \frac{d}{dx}\left(\frac{p}{x^2}\right) &= \left(\frac{1}{x^2}\right)(x^2) \\ d\left(\frac{p}{x^2}\right) &= dx\end{aligned}$$

Integrating gives

$$\begin{aligned}\frac{p}{x^2} &= \int dx \\ \frac{p}{x^2} &= x + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = \frac{1}{x^2}$ results in

$$p(x) = c_1 x^2 + x^3$$

which simplifies to

$$p(x) = x^2(x + c_1)$$

Since $p = y'$ then the new first order ode to solve is

$$y' = x^2(x + c_1)$$

Integrating both sides gives

$$\begin{aligned}y &= \int x^2(x + c_1) dx \\ &= \frac{1}{4}x^4 + \frac{1}{3}c_1 x^3 + c_2\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{1}{4}x^4 + \frac{1}{3}c_1x^3 + c_2 \quad (1)$$

Verification of solutions

$$y = \frac{1}{4}x^4 + \frac{1}{3}c_1x^3 + c_2$$

Verified OK.

23.6.3 Solving as second order ode non constant coeff transformation on B ode

Given an ode of the form

$$Ay'' + By' + Cy = F(x)$$

This method reduces the order ode the ODE by one by applying the transformation

$$y = Bv$$

This results in

$$\begin{aligned} y' &= B'v + v'B \\ y'' &= B''v + B'v' + v''B + v'B' \\ &= v''B + 2v' + B' + B''v \end{aligned}$$

And now the original ode becomes

$$\begin{aligned} A(v''B + 2v'B' + B''v) + B(B'v + v'B) + CBv &= 0 \\ ABv'' + (2AB' + B^2)v' + (AB'' + BB' + CB)v &= 0 \end{aligned} \quad (1)$$

If the term $AB'' + BB' + CB$ is zero, then this method works and can be used to solve

$$ABv'' + (2AB' + B^2)v' = 0$$

By Using $u = v'$ which reduces the order of the above ode to one. The new ode is

$$ABu' + (2AB' + B^2)u = 0$$

The above ode is first order ode which is solved for u . Now a new ode $v' = u$ is solved for v as first order ode. Then the final solution is obtain from $y = Bv$.

This method works only if the term $AB'' + BB' + CB$ is zero. The given ODE shows that

$$\begin{aligned}A &= x \\B &= -2 \\C &= 0 \\F &= x^3\end{aligned}$$

The above shows that for this ode

$$\begin{aligned}AB'' + BB' + CB &= (x)(0) + (-2)(0) + (0)(-2) \\&= 0\end{aligned}$$

Hence the ode in v given in (1) now simplifies to

$$-2xv'' + (4)v' = 0$$

Now by applying $v' = u$ the above becomes

$$-2xu'(x) + 4u(x) = 0$$

Which is now solved for u . In canonical form the ODE is

$$\begin{aligned}u' &= F(x, u) \\&= f(x)g(u) \\&= \frac{2u}{x}\end{aligned}$$

Where $f(x) = \frac{2}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u} du &= \frac{2}{x} dx \\ \int \frac{1}{u} du &= \int \frac{2}{x} dx \\ \ln(u) &= 2 \ln(x) + c_1 \\ u &= e^{2 \ln(x) + c_1} \\ &= c_1 x^2\end{aligned}$$

The ode for v now becomes

$$\begin{aligned}v' &= u \\ &= c_1 x^2\end{aligned}$$

Which is now solved for v . Integrating both sides gives

$$\begin{aligned}v(x) &= \int c_1 x^2 \, dx \\ &= \frac{c_1 x^3}{3} + c_2\end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}y_h(x) &= Bv \\ &= (-2) \left(\frac{c_1 x^3}{3} + c_2 \right) \\ &= -\frac{2c_1 x^3}{3} - 2c_2\end{aligned}$$

And now the particular solution $y_p(x)$ will be found. The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = -2$$

$$y_2 = x^3$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} -2 & x^3 \\ \frac{d}{dx}(-2) & \frac{d}{dx}(x^3) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} -2 & x^3 \\ 0 & 3x^2 \end{vmatrix}$$

Therefore

$$W = (-2)(3x^2) - (x^3)(0)$$

Which simplifies to

$$W = -6x^2$$

Which simplifies to

$$W = -6x^2$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^6}{-6x^3} dx$$

Which simplifies to

$$u_1 = - \int -\frac{x^3}{6} dx$$

Hence

$$u_1 = \frac{x^4}{24}$$

And Eq. (3) becomes

$$u_2 = \int \frac{-2x^3}{-6x^3} dx$$

Which simplifies to

$$u_2 = \int \frac{1}{3} dx$$

Hence

$$u_2 = \frac{x}{3}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{x^4}{4}$$

Hence the complete solution is

$$\begin{aligned}y(x) &= y_h + y_p \\&= \left(-\frac{2c_1x^3}{3} - 2c_2\right) + \left(\frac{x^4}{4}\right) \\&= -\frac{2}{3}c_1x^3 - 2c_2 + \frac{1}{4}x^4\end{aligned}$$

Summary

The solution(s) found are the following

$$y = -\frac{2}{3}c_1x^3 - 2c_2 + \frac{1}{4}x^4 \quad (1)$$

Verification of solutions

$$y = -\frac{2}{3}c_1x^3 - 2c_2 + \frac{1}{4}x^4$$

Verified OK.

23.6.4 Solving as type second_order_integrable_as_is (not using ABC version)

Writing the ode as

$$xy'' - 2y' = x^3$$

Integrating both sides of the ODE w.r.t x gives

$$\begin{aligned}\int (xy'' - 2y') dx &= \int x^3 dx \\xy' - 3y &= \frac{x^4}{4} + c_1\end{aligned}$$

Which is now solved for y .

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -\frac{3}{x}$$
$$q(x) = \frac{x^4 + 4c_1}{4x}$$

Hence the ode is

$$y' - \frac{3y}{x} = \frac{x^4 + 4c_1}{4x}$$

The integrating factor μ is

$$\mu = e^{\int -\frac{3}{x} dx}$$
$$= \frac{1}{x^3}$$

The ode becomes

$$\frac{d}{dx}(\mu y) = (\mu) \left(\frac{x^4 + 4c_1}{4x} \right)$$
$$\frac{d}{dx} \left(\frac{y}{x^3} \right) = \left(\frac{1}{x^3} \right) \left(\frac{x^4 + 4c_1}{4x} \right)$$
$$d \left(\frac{y}{x^3} \right) = \left(\frac{x^4 + 4c_1}{4x^4} \right) dx$$

Integrating gives

$$\frac{y}{x^3} = \int \frac{x^4 + 4c_1}{4x^4} dx$$
$$\frac{y}{x^3} = \frac{x}{4} - \frac{c_1}{3x^3} + c_2$$

Dividing both sides by the integrating factor $\mu = \frac{1}{x^3}$ results in

$$y = x^3 \left(\frac{x}{4} - \frac{c_1}{3x^3} \right) + c_2 x^3$$

which simplifies to

$$y = \frac{1}{4}x^4 - \frac{1}{3}c_1 + c_2x^3$$

Summary

The solution(s) found are the following

$$y = \frac{1}{4}x^4 - \frac{1}{3}c_1 + c_2x^3 \quad (1)$$

Verification of solutions

$$y = \frac{1}{4}x^4 - \frac{1}{3}c_1 + c_2x^3$$

Verified OK.

23.6.5 Solving using Kovacic algorithm

Writing the ode as

$$xy'' - 2y' = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= x \\ B &= -2 \\ C &= 0 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{2}{x^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 2 \\ t &= x^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{2}{x^2}\right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 281: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = \frac{2}{x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = 2$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = 2 \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = -1 \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = \frac{2}{x^2}$$

Since the $\text{gcd}(s, t) = 1$. This gives $b = 2$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = 2 \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = -1 \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{2}{x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	2	-1

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	2	-1

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω .

Trying $\alpha_{\infty}^{-} = -1$ then

$$\begin{aligned} d &= \alpha_{\infty}^{-} - (\alpha_{c_1}^{-}) \\ &= -1 - (-1) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_{\infty}$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^{-}}{x - c_1} \right) + (-)[\sqrt{r}]_{\infty} \\ &= -\frac{1}{x} + (-)(0) \\ &= -\frac{1}{x} \\ &= -\frac{1}{x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r)p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2\left(-\frac{1}{x}\right)(0) + \left(\left(\frac{1}{x^2}\right) + \left(-\frac{1}{x}\right)^2 - \left(\frac{2}{x^2}\right)\right) &= 0 \\ 0 &= 0 \end{aligned}$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= p e^{\int \omega dx} \\ &= e^{\int -\frac{1}{x} dx} \\ &= \frac{1}{x} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\&= z_1 e^{-\int \frac{1}{2} \frac{-2}{x} dx} \\&= z_1 e^{\ln(x)} \\&= z_1(x)\end{aligned}$$

Which simplifies to

$$y_1 = 1$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{-2}{x} dx}}{(y_1)^2} dx \\&= y_1 \int \frac{e^{2\ln(x)}}{(y_1)^2} dx \\&= y_1 \left(\frac{x^3}{3} \right)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\&= c_1(1) + c_2 \left(1 \left(\frac{x^3}{3} \right) \right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$.
 y_h is the solution to

$$xy'' - 2y' = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 + \frac{c_2 x^3}{3}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = 1$$
$$y_2 = \frac{x^3}{3}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} 1 & \frac{x^3}{3} \\ \frac{d}{dx}(1) & \frac{d}{dx}\left(\frac{x^3}{3}\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} 1 & \frac{x^3}{3} \\ 0 & x^2 \end{vmatrix}$$

Therefore

$$W = (1)(x^2) - \left(\frac{x^3}{3}\right)(0)$$

Which simplifies to

$$W = x^2$$

Which simplifies to

$$W = x^2$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^6}{x^3} dx$$

Which simplifies to

$$u_1 = - \int \frac{x^3}{3} dx$$

Hence

$$u_1 = -\frac{x^4}{12}$$

And Eq. (3) becomes

$$u_2 = \int \frac{x^3}{x^3} dx$$

Which simplifies to

$$u_2 = \int 1 dx$$

Hence

$$u_2 = x$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{x^4}{4}$$

Therefore the general solution is

$$\begin{aligned}y &= y_h + y_p \\ &= \left(c_1 + \frac{c_2 x^3}{3} \right) + \left(\frac{x^4}{4} \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 + \frac{1}{3}c_2x^3 + \frac{1}{4}x^4 \quad (1)$$

Verification of solutions

$$y = c_1 + \frac{1}{3}c_2x^3 + \frac{1}{4}x^4$$

Verified OK.

23.6.6 Solving as exact linear second order ode ode

An ode of the form

$$p(x) y'' + q(x) y' + r(x) y = s(x)$$

is exact if

$$p''(x) - q'(x) + r(x) = 0 \quad (1)$$

For the given ode we have

$$\begin{aligned}p(x) &= x \\ q(x) &= -2 \\ r(x) &= 0 \\ s(x) &= x^3\end{aligned}$$

Hence

$$\begin{aligned}p''(x) &= 0 \\ q'(x) &= 0\end{aligned}$$

Therefore (1) becomes

$$0 - (0) + (0) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(x)y' + (q(x) - p'(x))y)' = s(x)$$

Integrating gives

$$p(x)y' + (q(x) - p'(x))y = \int s(x) dx$$

Substituting the above values for p, q, r, s gives

$$xy' - 3y = \int x^3 dx$$

We now have a first order ode to solve which is

$$xy' - 3y = \frac{x^4}{4} + c_1$$

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -\frac{3}{x}$$
$$q(x) = \frac{x^4 + 4c_1}{4x}$$

Hence the ode is

$$y' - \frac{3y}{x} = \frac{x^4 + 4c_1}{4x}$$

The integrating factor μ is

$$\mu = e^{\int -\frac{3}{x} dx}$$
$$= \frac{1}{x^3}$$

The ode becomes

$$\frac{d}{dx}(\mu y) = (\mu) \left(\frac{x^4 + 4c_1}{4x} \right)$$
$$\frac{d}{dx} \left(\frac{y}{x^3} \right) = \left(\frac{1}{x^3} \right) \left(\frac{x^4 + 4c_1}{4x} \right)$$
$$d \left(\frac{y}{x^3} \right) = \left(\frac{x^4 + 4c_1}{4x^4} \right) dx$$

Integrating gives

$$\frac{y}{x^3} = \int \frac{x^4 + 4c_1}{4x^4} dx$$
$$\frac{y}{x^3} = \frac{x}{4} - \frac{c_1}{3x^3} + c_2$$

Dividing both sides by the integrating factor $\mu = \frac{1}{x^3}$ results in

$$y = x^3 \left(\frac{x}{4} - \frac{c_1}{3x^3} \right) + c_2 x^3$$

which simplifies to

$$y = \frac{1}{4}x^4 - \frac{1}{3}c_1 + c_2x^3$$

Summary

The solution(s) found are the following

$$y = \frac{1}{4}x^4 - \frac{1}{3}c_1 + c_2x^3 \quad (1)$$

Verification of solutions

$$y = \frac{1}{4}x^4 - \frac{1}{3}c_1 + c_2x^3$$

Verified OK.

23.6.7 Maple step by step solution

Let's solve

$$y''x - 2y' = x^3$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Make substitution $u = y'$ to reduce order of ODE

$$u'(x)x - 2u(x) = x^3$$

- Isolate the derivative

$$u'(x) = \frac{2u(x)}{x} + x^2$$

- Group terms with $u(x)$ on the lhs of the ODE and the rest on the rhs of the ODE

$$u'(x) - \frac{2u(x)}{x} = x^2$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) \left(u'(x) - \frac{2u(x)}{x} \right) = \mu(x) x^2$$
- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x) u(x))$

$$\mu(x) \left(u'(x) - \frac{2u(x)}{x} \right) = \mu'(x) u(x) + \mu(x) u'(x)$$
- Isolate $\mu'(x)$

$$\mu'(x) = -\frac{2\mu(x)}{x}$$
- Solve to find the integrating factor

$$\mu(x) = \frac{1}{x^2}$$
- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x) u(x)) \right) dx = \int \mu(x) x^2 dx + c_1$$
- Evaluate the integral on the lhs

$$\mu(x) u(x) = \int \mu(x) x^2 dx + c_1$$
- Solve for $u(x)$

$$u(x) = \frac{\int \mu(x) x^2 dx + c_1}{\mu(x)}$$
- Substitute $\mu(x) = \frac{1}{x^2}$

$$u(x) = x^2 \left(\int 1 dx + c_1 \right)$$
- Evaluate the integrals on the rhs

$$u(x) = x^2(x + c_1)$$
- Solve 1st ODE for $u(x)$

$$u(x) = x^2(x + c_1)$$
- Make substitution $u = y'$

$$y' = x^2(x + c_1)$$
- Integrate both sides to solve for y

$$\int y' dx = \int x^2(x + c_1) dx + c_2$$
- Compute integrals

$$y = \frac{1}{4}x^4 + \frac{1}{3}c_1x^3 + c_2$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
-> Calling odsolve with the ODE`, diff(_b(_a), _a) = (_a^3+2*_b(_a))/_a, _b(_a)` *** Suble  
  Methods for first order ODEs:  
  --- Trying classification methods ---  
  trying a quadrature  
  trying 1st order linear  
  <- 1st order linear successful  
<- high order exact linear fully integrable successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(x*diff(y(x),x$2)-2*diff(y(x),x)=x^3,y(x), singsol=all)
```

$$y(x) = \frac{1}{4}x^4 + \frac{1}{3}c_1x^3 + c_2$$

✓ Solution by Mathematica

Time used: 0.031 (sec). Leaf size: 24

```
DSolve[x*y'[x]-2*y'[x]==x^3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^4}{4} + \frac{c_1x^3}{3} + c_2$$

23.7 problem 2

23.7.1 Solving as second order ode missing y ode	1950
23.7.2 Solving as second order ode missing x ode	1952
23.7.3 Maple step by step solution	1954

Internal problem ID [6097]

Internal file name [OUTPUT/5345_Sunday_June_05_2022_03_34_51_PM_97008130/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 6. Existence and uniqueness of solutions to systems and nth order equations. Page 238

Problem number: 2.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_ode_missing_x**", "**second_order_ode_missing_y**"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_xy]]
```

$$y'' - y'^2 = 1$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

23.7.1 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x) - 1 - p(x)^2 = 0$$

Which is now solve for $p(x)$ as first order ode. Integrating both sides gives

$$\int \frac{1}{p^2 + 1} dp = x + c_1$$
$$\arctan(p) = x + c_1$$

Solving for p gives these solutions

$$p_1 = \tan(x + c_1)$$

Initial conditions are used to solve for c_1 . Substituting $x = 0$ and $p = 0$ in the above solution gives an equation to solve for the constant of integration.

$$0 = \tan(c_1)$$

$$c_1 = 0$$

Substituting c_1 found above in the general solution gives

$$p(x) = \tan(x)$$

Since $p = y'$ then the new first order ode to solve is

$$y' = \tan(x)$$

Integrating both sides gives

$$y = \int \tan(x) dx$$
$$= -\ln(\cos(x)) + c_2$$

Initial conditions are used to solve for c_2 . Substituting $x = 0$ and $y = 0$ in the above solution gives an equation to solve for the constant of integration.

$$0 = c_2$$

$$c_2 = 0$$

Substituting c_2 found above in the general solution gives

$$y = -\ln(\cos(x))$$

Initial conditions are used to solve for the constants of integration.

Summary

The solution(s) found are the following

$$y = -\ln(\cos(x)) \quad (1)$$




Figure 227: Solution plot

Verification of solutions

$$y = -\ln(\cos(x))$$

Verified OK.

23.7.2 Solving as second order ode missing x ode

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable y an independent variable. Using

$$y' = p(y)$$

Then

$$\begin{aligned} y'' &= \frac{dp}{dx} \\ &= \frac{dy}{dx} \frac{dp}{dy} \\ &= p \frac{dp}{dy} \end{aligned}$$

Hence the ode becomes

$$p(y) \left(\frac{d}{dy} p(y) \right) - p(y)^2 = 1$$

Which is now solved as first order ode for $p(y)$. Integrating both sides gives

$$\int \frac{p}{p^2 + 1} dp = \int dy$$

$$\frac{\ln(p^2 + 1)}{2} = y + c_1$$

Raising both side to exponential gives

$$\sqrt{p^2 + 1} = e^{y+c_1}$$

Which simplifies to

$$\sqrt{p^2 + 1} = c_2 e^y$$

Unable to solve for constant of integration due to RootOf in solution.

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = \text{RootOf}(_Z^2 - c_2^2 e^{2y} + 1)$$

Integrating both sides gives

$$\int \frac{1}{\text{RootOf}(_Z^2 - c_2^2 e^{2y} + 1)} dy = \int dx$$

$$\int^y \frac{1}{\text{RootOf}(_Z^2 - c_2^2 e^{2-a} + 1)} d_a = x + c_3$$

Unable to solve for constant of integration due to RootOf in solution.

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$\int^y \frac{1}{\text{RootOf}(_Z^2 - c_2^2 e^{2-a} + 1)} d_a = x + c_3 \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$\int^0 \frac{1}{\text{RootOf}(_Z^2 - c_2^2 e^{2-a} + 1)} d_a = c_3 \quad (1A)$$

Taking derivative of the solution gives

$$y' = \text{RootOf} \left(-Z^2 - c_2^2 e^{2 \text{RootOf} \left(- \left(\int^{-Z} \frac{1}{\text{RootOf}(-Z^2 - c_2^2 e^{2-a+1})} d_{-a} \right) + x + c_3 \right)} + 1 \right)$$

substituting $y' = 0$ and $x = 0$ in the above gives

$$0 = \lim_{x \rightarrow 0} \text{RootOf} \left(-Z^2 - c_2^2 e^{2 \text{RootOf} \left(- \left(\int^{-Z} \frac{1}{\text{RootOf}(-Z^2 - c_2^2 e^{2-a+1})} d_{-a} \right) + x + c_3 \right)} + 1 \right) \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_2, c_3\}$. There is no solution for the constants of integrations. This solution is removed.

Verification of solutions N/A

23.7.3 Maple step by step solution

Let's solve

$$\left[y'' - y'^2 = 1, y(0) = 0, y' \Big|_{\{x=0\}} = 0 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Make substitution $u = y'$ to reduce order of ODE

$$u'(x) - u(x)^2 = 1$$

- Separate variables

$$\frac{u'(x)}{u(x)^2+1} = 1$$

- Integrate both sides with respect to x

$$\int \frac{u'(x)}{u(x)^2+1} dx = \int 1 dx + c_1$$

- Evaluate integral

$$\arctan(u(x)) = x + c_1$$

- Solve for $u(x)$

$$u(x) = \tan(x + c_1)$$

- Solve 1st ODE for $u(x)$

$$u(x) = \tan(x + c_1)$$

- Make substitution $u = y'$
 $y' = \tan(x + c_1)$
- Integrate both sides to solve for y
 $\int y' dx = \int \tan(x + c_1) dx + c_2$
- Compute integrals
 $y = \frac{\ln(1 + \tan(x + c_1)^2)}{2} + c_2$
- Check validity of solution $y = \frac{\ln(1 + \tan(x + c_1)^2)}{2} + c_2$
 - Use initial condition $y(0) = 0$
 $0 = \frac{\ln(1 + \tan(c_1)^2)}{2} + c_2$
 - Compute derivative of the solution
 $y' = \tan(x + c_1)$
 - Use the initial condition $y'|_{\{x=0\}} = 0$
 $0 = \tan(c_1)$
 - Solve for c_1 and c_2
 $\{c_1 = 0, c_2 = 0\}$
 - Substitute constant values into general solution and simplify
 $y = \frac{\ln(\sec(x)^2)}{2}$
- Solution to the IVP
 $y = \frac{\ln(\sec(x)^2)}{2}$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying 2nd order Liouville  
trying 2nd order WeierstrassP  
trying 2nd order JacobiSN  
differential order: 2; trying a linearization to 3rd order  
trying 2nd order ODE linearizable_by_differentiation  
trying 2nd order, 2 integrating factors of the form mu(x,y)  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful  
<- 2nd order, 2 integrating factors of the form mu(x,y) successful`
```

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 7

```
dsolve([diff(y(x),x$2)=1+diff(y(x),x)^2,y(0) = 0, D(y)(0) = 0],y(x), singsol=all)
```

$$y(x) = \ln(\sec(x))$$

✓ Solution by Mathematica

Time used: 2.581 (sec). Leaf size: 27

```
DSolve[{y'[x]==1+(y'[x])^2,{y[0]==0,y'[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\log(-\cos(x)) + i\pi$$

$$y(x) \rightarrow -\log(\cos(x))$$

23.8 problem 3

23.8.1 Solving as second order integrable as is ode	1958
23.8.2 Solving as second order ode missing y ode	1961
23.8.3 Solving as second order ode missing x ode	1963
23.8.4 Solving as exact nonlinear second order ode ode	1966
23.8.5 Maple step by step solution	1969

Internal problem ID [6098]

Internal file name [OUTPUT/5346_Sunday_June_05_2022_03_34_53_PM_64136479/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 6. Existence and uniqueness of solutions to systems and nth order equations. Page 238

Problem number: 3.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_integrable_as_is", "second_order_ode_missing_x", "second_order_ode_missing_y", "exact nonlinear second order ode"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_poly_yn]]
```

$$y'' + \frac{1}{2y'^2} = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = -1]$$

23.8.1 Solving as second order integrable as is ode

Integrating both sides of the ODE w.r.t x gives

$$\int 2y''y'^2 dx = \int (-1) dx$$
$$\frac{2y'^3}{3} = -x + c_1$$

Which is now solved for y . Solving the given ode for y' results in 3 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \frac{(-12x + 12c_1)^{\frac{1}{3}}}{2} \quad (1)$$

$$y' = -\frac{(-12x + 12c_1)^{\frac{1}{3}}}{4} + \frac{i\sqrt{3}(-12x + 12c_1)^{\frac{1}{3}}}{4} \quad (2)$$

$$y' = -\frac{(-12x + 12c_1)^{\frac{1}{3}}}{4} - \frac{i\sqrt{3}(-12x + 12c_1)^{\frac{1}{3}}}{4} \quad (3)$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$y = \int \frac{(-12x + 12c_1)^{\frac{1}{3}}}{2} dx$$
$$= \frac{3(-c_1 + x)(-12x + 12c_1)^{\frac{1}{3}}}{8} + c_2$$

Solving equation (2)

Integrating both sides gives

$$y = \int -\frac{(-12x + 12c_1)^{\frac{1}{3}}}{4} + \frac{i\sqrt{3}(-12x + 12c_1)^{\frac{1}{3}}}{4} dx$$
$$= \frac{3(-c_1 + x)(-12x + 12c_1)^{\frac{1}{3}}(i\sqrt{3} - 1)}{16} + c_3$$

Solving equation (3)

Integrating both sides gives

$$y = \int -\frac{(-12x + 12c_1)^{\frac{1}{3}}}{4} - \frac{i\sqrt{3}(-12x + 12c_1)^{\frac{1}{3}}}{4} dx$$
$$= -\frac{3(-c_1 + x)(-12x + 12c_1)^{\frac{1}{3}}(1 + i\sqrt{3})}{16} + c_4$$

Initial conditions are used to solve for the constants of integration.

Looking at the First solution

$$y = \frac{3(-c_1 + x)(-12x + 12c_1)^{\frac{1}{3}}}{8} + c_2 \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 0$ in the above gives

$$1 = -\frac{3c_1^{\frac{4}{3}}12^{\frac{1}{3}}}{8} + c_2 \quad (1A)$$

Taking derivative of the solution gives

$$y' = \frac{3(-12x + 12c_1)^{\frac{1}{3}}}{8} - \frac{3(-c_1 + x)}{2(-12x + 12c_1)^{\frac{2}{3}}}$$

substituting $y' = -1$ and $x = 0$ in the above gives

$$-1 = \frac{c_1^{\frac{1}{3}}12^{\frac{1}{3}}}{2} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. There is no solution for the constants of integrations. This solution is removed.

Looking at the Second solution

$$y = \frac{3(-c_1 + x)(-12x + 12c_1)^{\frac{1}{3}}(i\sqrt{3} - 1)}{16} + c_3 \quad (2)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 0$ in the above gives

$$1 = -\frac{3 \cdot 2^{\frac{2}{3}} \left(i3^{\frac{5}{6}} - 3^{\frac{1}{3}} \right) c_1^{\frac{4}{3}}}{16} + c_3 \quad (1A)$$

Taking derivative of the solution gives

$$y' = \frac{3(-12x + 12c_1)^{\frac{1}{3}}(i\sqrt{3} - 1)}{16} - \frac{3(-c_1 + x)(i\sqrt{3} - 1)}{4(-12x + 12c_1)^{\frac{2}{3}}}$$

substituting $y' = -1$ and $x = 0$ in the above gives

$$-1 = \frac{2^{\frac{2}{3}} c_1^{\frac{1}{3}} \left(i3^{\frac{5}{6}} - 3^{\frac{1}{3}} \right)}{4} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_3\}$. Solving for the constants gives

$$c_1 = -\frac{2}{3}$$

$$c_3 = \frac{3}{2}$$

Substituting these values back in above solution results in

$$y = \frac{i(-12x - 8)^{\frac{1}{3}} \sqrt{3}}{8} - \frac{(-12x - 8)^{\frac{1}{3}}}{8} + \frac{3i(-12x - 8)^{\frac{1}{3}} \sqrt{3} x}{16} - \frac{3(-12x - 8)^{\frac{1}{3}} x}{16} + \frac{3}{2}$$

Looking at the Third solution

$$y = -\frac{3(-c_1 + x)(-12x + 12c_1)^{\frac{1}{3}}(1 + i\sqrt{3})}{16} + c_4 \quad (3)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 0$ in the above gives

$$1 = \frac{3\left(i3^{\frac{5}{6}} + 3^{\frac{1}{3}}\right)2^{\frac{2}{3}}c_1^{\frac{4}{3}}}{16} + c_4 \quad (1A)$$

Taking derivative of the solution gives

$$y' = -\frac{3(-12x + 12c_1)^{\frac{1}{3}}(1 + i\sqrt{3})}{16} + \frac{3(-c_1 + x)(1 + i\sqrt{3})}{4(-12x + 12c_1)^{\frac{2}{3}}}$$

substituting $y' = -1$ and $x = 0$ in the above gives

$$-1 = -\frac{2^{\frac{2}{3}} c_1^{\frac{1}{3}} \left(i3^{\frac{5}{6}} + 3^{\frac{1}{3}} \right)}{4} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_4\}$. There is no solution for the constants of integrations. This solution is removed.

Summary

The solution(s) found are the following

$$y = \frac{3}{2} + \frac{3(i\sqrt{3} - 1) \left(\frac{2}{3} + x\right) (-12x - 8)^{\frac{1}{3}}}{16} \quad (1)$$

Verification of solutions

$$y = \frac{3}{2} + \frac{3(i\sqrt{3} - 1) \left(\frac{2}{3} + x\right) (-12x - 8)^{\frac{1}{3}}}{16}$$

Verified OK.

23.8.2 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$2p'(x)p(x)^2 + 1 = 0$$

Which is now solve for $p(x)$ as first order ode. Integrating both sides gives

$$\begin{aligned} \int -2p^2 dp &= x + c_1 \\ -\frac{2p^3}{3} &= x + c_1 \end{aligned}$$

Solving for p gives these solutions

$$\begin{aligned} p_1 &= \frac{(-12x - 12c_1)^{\frac{1}{3}}}{2} \\ p_2 &= -\frac{(-12x - 12c_1)^{\frac{1}{3}}}{4} - \frac{i\sqrt{3}(-12x - 12c_1)^{\frac{1}{3}}}{4} \\ p_3 &= -\frac{(-12x - 12c_1)^{\frac{1}{3}}}{4} + \frac{i\sqrt{3}(-12x - 12c_1)^{\frac{1}{3}}}{4} \end{aligned}$$

Initial conditions are used to solve for c_1 . Substituting $x = 0$ and $p = -1$ in the above solution gives an equation to solve for the constant of integration.

$$-1 = \frac{i(-12c_1)^{\frac{1}{3}}\sqrt{3}}{4} - \frac{(-12c_1)^{\frac{1}{3}}}{4}$$

$$c_1 = \frac{2}{3}$$

Substituting c_1 found above in the general solution gives

$$p(x) = \frac{i(-12x - 8)^{\frac{1}{3}} \sqrt{3}}{4} - \frac{(-12x - 8)^{\frac{1}{3}}}{4}$$

Initial conditions are used to solve for c_1 . Substituting $x = 0$ and $p = -1$ in the above solution gives an equation to solve for the constant of integration.

$$-1 = -\frac{i(-12c_1)^{\frac{1}{3}} \sqrt{3}}{4} - \frac{(-12c_1)^{\frac{1}{3}}}{4}$$

Warning: Unable to solve for constant of integration. Initial conditions are used to solve for c_1 . Substituting $x = 0$ and $p = -1$ in the above solution gives an equation to solve for the constant of integration.

$$-1 = \frac{(-12c_1)^{\frac{1}{3}}}{2}$$

Warning: Unable to solve for constant of integration. Since $p = y'$ then the new first order ode to solve is

$$y' = \frac{i(-12x - 8)^{\frac{1}{3}} \sqrt{3}}{4} - \frac{(-12x - 8)^{\frac{1}{3}}}{4}$$

Integrating both sides gives

$$\begin{aligned} y &= \int \frac{i(-12x - 8)^{\frac{1}{3}} \sqrt{3}}{4} - \frac{(-12x - 8)^{\frac{1}{3}}}{4} dx \\ &= \frac{(2 + 3x)(-12x - 8)^{\frac{1}{3}}(i\sqrt{3} - 1)}{16} + c_2 \end{aligned}$$

Initial conditions are used to solve for c_2 . Substituting $x = 0$ and $y = 1$ in the above solution gives an equation to solve for the constant of integration.

$$1 = -\frac{1}{2} + c_2$$

$$c_2 = \frac{3}{2}$$

Substituting c_2 found above in the general solution gives

$$y = \frac{i(-12x - 8)^{\frac{1}{3}} \sqrt{3}}{8} - \frac{(-12x - 8)^{\frac{1}{3}}}{8} + \frac{3i(-12x - 8)^{\frac{1}{3}} \sqrt{3} x}{16} - \frac{3(-12x - 8)^{\frac{1}{3}} x}{16} + \frac{3}{2}$$

Initial conditions are used to solve for the constants of integration.

Summary

The solution(s) found are the following

$$y = \frac{i(-12x - 8)^{\frac{1}{3}} \sqrt{3}}{8} - \frac{(-12x - 8)^{\frac{1}{3}}}{8} + \frac{3i(-12x - 8)^{\frac{1}{3}} \sqrt{3} x}{16} - \frac{3(-12x - 8)^{\frac{1}{3}} x}{16} + \left(\frac{3}{2}\right)$$

Verification of solutions

$$y = \frac{i(-12x - 8)^{\frac{1}{3}} \sqrt{3}}{8} - \frac{(-12x - 8)^{\frac{1}{3}}}{8} + \frac{3i(-12x - 8)^{\frac{1}{3}} \sqrt{3} x}{16} - \frac{3(-12x - 8)^{\frac{1}{3}} x}{16} + \frac{3}{2}$$

Verified OK.

23.8.3 Solving as second order ode missing x ode

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable y an independent variable.

Using

$$y' = p(y)$$

Then

$$\begin{aligned} y'' &= \frac{dp}{dx} \\ &= \frac{dy}{dx} \frac{dp}{dy} \\ &= p \frac{dp}{dy} \end{aligned}$$

Hence the ode becomes

$$2p(y)^3 \left(\frac{d}{dy} p(y) \right) = -1$$

Which is now solved as first order ode for $p(y)$. Integrating both sides gives

$$\begin{aligned} \int -2p^3 dp &= y + c_1 \\ -\frac{p^4}{2} &= y + c_1 \end{aligned}$$

Solving for p gives these solutions

$$\begin{aligned} p_1 &= (-2c_1 - 2y)^{\frac{1}{4}} \\ p_2 &= -i(-2c_1 - 2y)^{\frac{1}{4}} \\ p_3 &= i(-2c_1 - 2y)^{\frac{1}{4}} \\ p_4 &= -(-2c_1 - 2y)^{\frac{1}{4}} \end{aligned}$$

Initial conditions are used to solve for c_1 . Substituting $y = 1$ and $p = -1$ in the above solution gives an equation to solve for the constant of integration.

$$-1 = -(-2c_1 - 2)^{\frac{1}{4}}$$

$$c_1 = -\frac{3}{2}$$

Substituting c_1 found above in the general solution gives

$$p(y) = -(3 - 2y)^{\frac{1}{4}}$$

Initial conditions are used to solve for c_1 . Substituting $y = 1$ and $p = -1$ in the above solution gives an equation to solve for the constant of integration.

$$-1 = i(-2c_1 - 2)^{\frac{1}{4}}$$

Warning: Unable to solve for constant of integration. Initial conditions are used to solve for c_1 . Substituting $y = 1$ and $p = -1$ in the above solution gives an equation to solve for the constant of integration.

$$-1 = -i(-2c_1 - 2)^{\frac{1}{4}}$$

Warning: Unable to solve for constant of integration. Initial conditions are used to solve for c_1 . Substituting $y = 1$ and $p = -1$ in the above solution gives an equation to solve for the constant of integration.

$$-1 = (-2c_1 - 2)^{\frac{1}{4}}$$

Warning: Unable to solve for constant of integration. For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = -(3 - 2y)^{\frac{1}{4}}$$

Integrating both sides gives

$$\int -\frac{1}{(3 - 2y)^{\frac{1}{4}}} dy = \int dx$$

$$\frac{2(3 - 2y)^{\frac{3}{4}}}{3} = c_2 + x$$

Initial conditions are used to solve for c_2 . Substituting $x = 0$ and $y = 1$ in the above solution gives an equation to solve for the constant of integration.

$$\frac{2}{3} = c_2$$

$$c_2 = \frac{2}{3}$$

Substituting c_2 found above in the general solution gives

$$\frac{2(3 - 2y)^{\frac{3}{4}}}{3} = \frac{2}{3} + x$$

Solving for y from the above gives

$$y = \frac{3}{2} + \frac{(-3x - 2) \left(\frac{3x}{2} + 1\right)^{\frac{1}{3}}}{4}$$

Initial conditions are used to solve for the constants of integration.

Summary

The solution(s) found are the following

$$y = \frac{3}{2} + \frac{(-3x - 2) \left(\frac{3x}{2} + 1\right)^{\frac{1}{3}}}{4} \quad (1)$$




Figure 228: Solution plot

Verification of solutions

$$y = \frac{3}{2} + \frac{(-3x - 2) \left(\frac{3x}{2} + 1\right)^{\frac{1}{3}}}{4}$$

Verified OK.

23.8.4 Solving as exact nonlinear second order ode

An exact non-linear second order ode has the form

$$a_2(x, y, y') y'' + a_1(x, y, y') y' + a_0(x, y, y') = 0$$

Where the following conditions are satisfied

$$\begin{aligned} \frac{\partial a_2}{\partial y} &= \frac{\partial a_1}{\partial y'} \\ \frac{\partial a_2}{\partial x} &= \frac{\partial a_0}{\partial y'} \\ \frac{\partial a_1}{\partial x} &= \frac{\partial a_0}{\partial y} \end{aligned}$$

Looking at the the ode given we see that

$$\begin{aligned} a_2 &= 2y'^2 \\ a_1 &= 0 \\ a_0 &= 1 \end{aligned}$$

Applying the conditions to the above shows this is a nonlinear exact second order ode. Therefore it can be reduced to first order ode given by

$$\begin{aligned} \int a_2 dy' + \int a_1 dy + \int a_0 dx &= c_1 \\ \int 2y'^2 dy' + \int 0 dy + \int 1 dx &= c_1 \end{aligned}$$

Which results in

$$\frac{2y'^3}{3} + x = c_1$$

Which is now solved Solving the given ode for y' results in 3 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \frac{(-12x + 12c_1)^{\frac{1}{3}}}{2} \tag{1}$$

$$y' = -\frac{(-12x + 12c_1)^{\frac{1}{3}}}{4} + \frac{i\sqrt{3}(-12x + 12c_1)^{\frac{1}{3}}}{4} \tag{2}$$

$$y' = -\frac{(-12x + 12c_1)^{\frac{1}{3}}}{4} - \frac{i\sqrt{3}(-12x + 12c_1)^{\frac{1}{3}}}{4} \tag{3}$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\begin{aligned}y &= \int \frac{(-12x + 12c_1)^{\frac{1}{3}}}{2} dx \\&= \frac{3(-c_1 + x)(-12x + 12c_1)^{\frac{1}{3}}}{8} + c_2\end{aligned}$$

Solving equation (2)

Integrating both sides gives

$$\begin{aligned}y &= \int -\frac{(-12x + 12c_1)^{\frac{1}{3}}}{4} + \frac{i\sqrt{3}(-12x + 12c_1)^{\frac{1}{3}}}{4} dx \\&= \frac{3(-c_1 + x)(-12x + 12c_1)^{\frac{1}{3}}(i\sqrt{3} - 1)}{16} + c_3\end{aligned}$$

Solving equation (3)

Integrating both sides gives

$$\begin{aligned}y &= \int -\frac{(-12x + 12c_1)^{\frac{1}{3}}}{4} - \frac{i\sqrt{3}(-12x + 12c_1)^{\frac{1}{3}}}{4} dx \\&= -\frac{3(-c_1 + x)(-12x + 12c_1)^{\frac{1}{3}}(1 + i\sqrt{3})}{16} + c_4\end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the First solution

$$y = \frac{3(-c_1 + x)(-12x + 12c_1)^{\frac{1}{3}}}{8} + c_2 \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 0$ in the above gives

$$1 = -\frac{3c_1^{\frac{4}{3}}12^{\frac{1}{3}}}{8} + c_2 \quad (1A)$$

Taking derivative of the solution gives

$$y' = \frac{3(-12x + 12c_1)^{\frac{1}{3}}}{8} - \frac{3(-c_1 + x)}{2(-12x + 12c_1)^{\frac{2}{3}}}$$

substituting $y' = -1$ and $x = 0$ in the above gives

$$-1 = \frac{c_1^{\frac{1}{3}} 12^{\frac{1}{3}}}{2} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. There is no solution for the constants of integrations. This solution is removed.

Looking at the Second solution

$$y = \frac{3(-c_1 + x)(-12x + 12c_1)^{\frac{1}{3}}(i\sqrt{3} - 1)}{16} + c_3 \quad (2)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 0$ in the above gives

$$1 = -\frac{3 \cdot 2^{\frac{2}{3}} \left(i3^{\frac{5}{6}} - 3^{\frac{1}{3}} \right) c_1^{\frac{4}{3}}}{16} + c_3 \quad (1A)$$

Taking derivative of the solution gives

$$y' = \frac{3(-12x + 12c_1)^{\frac{1}{3}}(i\sqrt{3} - 1)}{16} - \frac{3(-c_1 + x)(i\sqrt{3} - 1)}{4(-12x + 12c_1)^{\frac{2}{3}}}$$

substituting $y' = -1$ and $x = 0$ in the above gives

$$-1 = \frac{2^{\frac{2}{3}} c_1^{\frac{1}{3}} \left(i3^{\frac{5}{6}} - 3^{\frac{1}{3}} \right)}{4} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_3\}$. Solving for the constants gives

$$c_1 = -\frac{2}{3}$$

$$c_3 = \frac{3}{2}$$

Substituting these values back in above solution results in

$$y = \frac{i(-12x - 8)^{\frac{1}{3}}\sqrt{3}}{8} - \frac{(-12x - 8)^{\frac{1}{3}}}{8} + \frac{3i(-12x - 8)^{\frac{1}{3}}\sqrt{3}x}{16} - \frac{3(-12x - 8)^{\frac{1}{3}}x}{16} + \frac{3}{2}$$

Looking at the Third solution

$$y = -\frac{3(-c_1 + x)(-12x + 12c_1)^{\frac{1}{3}}(1 + i\sqrt{3})}{16} + c_4 \quad (3)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 0$ in the above gives

$$1 = \frac{3\left(i3^{\frac{5}{6}} + 3^{\frac{1}{3}}\right)2^{\frac{2}{3}}c_1^{\frac{4}{3}}}{16} + c_4 \quad (1A)$$

Taking derivative of the solution gives

$$y' = -\frac{3(-12x + 12c_1)^{\frac{1}{3}}(1 + i\sqrt{3})}{16} + \frac{3(-c_1 + x)(1 + i\sqrt{3})}{4(-12x + 12c_1)^{\frac{2}{3}}}$$

substituting $y' = -1$ and $x = 0$ in the above gives

$$-1 = -\frac{2^{\frac{2}{3}}c_1^{\frac{1}{3}}\left(i3^{\frac{5}{6}} + 3^{\frac{1}{3}}\right)}{4} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_4\}$. There is no solution for the constants of integrations. This solution is removed.

Summary

The solution(s) found are the following

$$y = \frac{3}{2} + \frac{3(i\sqrt{3} - 1)\left(\frac{2}{3} + x\right)(-12x - 8)^{\frac{1}{3}}}{16} \quad (1)$$

Verification of solutions

$$y = \frac{3}{2} + \frac{3(i\sqrt{3} - 1)\left(\frac{2}{3} + x\right)(-12x - 8)^{\frac{1}{3}}}{16}$$

Verified OK.

23.8.5 Maple step by step solution

Let's solve

$$\left[2y''y'^2 = -1, y(0) = 1, y'|_{\{x=0\}} = -1\right]$$

- Highest derivative means the order of the ODE is 2

y''

- Make substitution $u = y'$ to reduce order of ODE

$$2u'(x)u(x)^2 = -1$$

- Integrate both sides with respect to x

$$\int 2u'(x) u(x)^2 dx = \int (-1) dx + c_1$$
- Evaluate integral

$$\frac{2u(x)^3}{3} = -x + c_1$$
- Solve for $u(x)$

$$u(x) = \frac{(-12x+12c_1)^{\frac{1}{3}}}{2}$$
- Solve 1st ODE for $u(x)$

$$u(x) = \frac{(-12x+12c_1)^{\frac{1}{3}}}{2}$$
- Make substitution $u = y'$

$$y' = \frac{(-12x+12c_1)^{\frac{1}{3}}}{2}$$
- Integrate both sides to solve for y

$$\int y' dx = \int \frac{(-12x+12c_1)^{\frac{1}{3}}}{2} dx + c_2$$
- Compute integrals

$$y = \frac{3(-c_1+x)(-12x+12c_1)^{\frac{1}{3}}}{8} + c_2$$
- Check validity of solution $y = \frac{3(-c_1+x)(-12x+12c_1)^{\frac{1}{3}}}{8} + c_2$
 - Use initial condition $y(0) = 1$

$$1 = -\frac{3c_1^{\frac{4}{3}}12^{\frac{1}{3}}}{8} + c_2$$
 - Compute derivative of the solution

$$y' = \frac{3(-12x+12c_1)^{\frac{1}{3}}}{8} - \frac{3(-c_1+x)}{2(-12x+12c_1)^{\frac{2}{3}}}$$
 - Use the initial condition $y' \Big|_{\{x=0\}} = -1$

$$-1 = \frac{c_1^{\frac{1}{3}}12^{\frac{1}{3}}}{2}$$
 - Solve for c_1 and c_2
 - The solution does not satisfy the initial conditions

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
-> Calling odsolve with the ODE`, diff(_b(_a), _a) = -(1/2)/_b(_a)^2, _b(_a), HINT = [[1, 0]
    symmetry methods on request
`, `1st order, trying reduction of order with given symmetries:`[1, 0], [_a, 1/3*_b]
```

✓ Solution by Maple

Time used: 0.469 (sec). Leaf size: 26

```
dsolve([diff(y(x),x$2)=-1/(2*diff(y(x),x)^2),y(0) = 1, D(y)(0) = -1],y(x), singsol=all)
```

$$y(x) = \frac{3(x + \frac{2}{3})(-12x - 8)^{\frac{1}{3}}(i\sqrt{3} - 1)}{16} + \frac{3}{2}$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 27

```
DSolve[{y'[x]==-1/(2*(y'[x])^2),{y[0]==1,y'[0]==-1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{8}(12 - (-2)^{2/3}(-3x - 2)^{4/3})$$

23.9 problem 5(b)

23.9.1 Solving as second order ode can be made integrable ode 1972

23.9.2 Solving as second order ode missing x ode 1974

Internal problem ID [6099]

Internal file name [OUTPUT/5347_Sunday_June_05_2022_03_34_57_PM_92223807/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 6. Existence and uniqueness of solutions to systems and nth order equations. Page 238

Problem number: 5(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_ode_missing_x", "second_order_ode_can_be_made_integrable"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x], [_2nd_order , _reducible , _mu_x_y1]]
```

$$y'' + \sin(y) = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = \beta]$$

23.9.1 Solving as second order ode can be made integrable ode

Multiplying the ode by y' gives

$$y'y'' + y'\sin(y) = 0$$

Integrating the above w.r.t x gives

$$\int (y'y'' + y'\sin(y)) dx = 0$$
$$\frac{y'^2}{2} - \cos(y) = c_2$$

Which is now solved for y . Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \sqrt{2 \cos (y) + 2c_1} \quad (1)$$

$$y' = -\sqrt{2 \cos (y) + 2c_1} \quad (2)$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int \frac{1}{\sqrt{2 \cos (y) + 2c_1}} dy = \int dx$$

$$\frac{2\sqrt{\frac{\cos(y)+c_1}{c_1+1}} \operatorname{InverseJacobiAM}\left(\frac{y}{2}, \frac{2}{\sqrt{2+2c_1}}\right)}{\sqrt{2 \cos (y) + 2c_1}} = c_2 + x$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{1}{\sqrt{2 \cos (y) + 2c_1}} dy = \int dx$$

$$-\frac{2\sqrt{\frac{\cos(y)+c_1}{c_1+1}} \operatorname{InverseJacobiAM}\left(\frac{y}{2}, \frac{2}{\sqrt{2+2c_1}}\right)}{\sqrt{2 \cos (y) + 2c_1}} = x + c_3$$

Initial conditions are used to solve for the constants of integration.

Looking at the First solution

$$\frac{2\sqrt{\frac{\cos(y)+c_1}{c_1+1}} \operatorname{InverseJacobiAM}\left(\frac{y}{2}, \frac{2}{\sqrt{2+2c_1}}\right)}{\sqrt{2 \cos (y) + 2c_1}} = c_2 + x \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_2 \quad (1A)$$

Taking derivative of the solution gives

Expression too large to display

substituting $y' = \beta$ and $x = 0$ in the above gives

$$\text{Expression too large to display} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Warning, unable to solve for constants of integrations.

Looking at the Second solution

$$-\frac{2\sqrt{\frac{\cos(y)+c_1}{c_1+1}} \operatorname{InverseJacobiAM}\left(\frac{y}{2}, \frac{2}{\sqrt{2+2c_1}}\right)}{\sqrt{2\cos(y)+2c_1}} = x + c_3 \quad (2)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_3 \quad (1A)$$

Taking derivative of the solution gives

$$\text{Expression too large to display}$$

substituting $y' = \beta$ and $x = 0$ in the above gives

$$\text{Expression too large to display} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_3\}$. Warning, unable to solve for constants of integrations.

Verification of solutions N/A

23.9.2 Solving as second order ode missing x ode

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable y an independent variable. Using

$$y' = p(y)$$

Then

$$\begin{aligned} y'' &= \frac{dp}{dx} \\ &= \frac{dy}{dx} \frac{dp}{dy} \\ &= p \frac{dp}{dy} \end{aligned}$$

Hence the ode becomes

$$p(y) \left(\frac{d}{dy} p(y) \right) = -\sin(y)$$

Which is now solved as first order ode for $p(y)$. In canonical form the ODE is

$$\begin{aligned} p' &= F(y, p) \\ &= f(y)g(p) \\ &= -\frac{\sin(y)}{p} \end{aligned}$$

Where $f(y) = -\sin(y)$ and $g(p) = \frac{1}{p}$. Integrating both sides gives

$$\begin{aligned} \frac{1}{p} dp &= -\sin(y) dy \\ \int \frac{1}{p} dp &= \int -\sin(y) dy \\ \frac{p^2}{2} &= \cos(y) + c_1 \end{aligned}$$

The solution is

$$\frac{p(y)^2}{2} - \cos(y) - c_1 = 0$$

Initial conditions are used to solve for c_1 . Substituting $y = 0$ and $p = \beta$ in the above solution gives an equation to solve for the constant of integration.

$$\frac{\beta^2}{2} - 1 - c_1 = 0$$

$$c_1 = -1 + \frac{\beta^2}{2}$$

Substituting c_1 found above in the general solution gives

$$\frac{p^2}{2} - \cos(y) + 1 - \frac{\beta^2}{2} = 0$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$\frac{y'^2}{2} - \cos(y) + 1 - \frac{\beta^2}{2} = 0$$

Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \sqrt{\beta^2 + 2 \cos(y) - 2} \quad (1)$$

$$y' = -\sqrt{\beta^2 + 2 \cos(y) - 2} \quad (2)$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int \frac{1}{\sqrt{\beta^2 + 2 \cos(y) - 2}} dy = \int dx$$

$$\frac{2\sqrt{\frac{\beta^2 + 2 \cos(y) - 2}{\beta^2}} \operatorname{InverseJacobiAM}\left(\frac{y}{2}, \frac{2}{\beta}\right)}{\operatorname{csgn}(\beta)^2 \sqrt{\beta^2 + 2 \cos(y) - 2}} = c_2 + x$$

Simplifying the solution $\frac{2\sqrt{\frac{\beta^2 + 2 \cos(y) - 2}{\beta^2}} \operatorname{InverseJacobiAM}\left(\frac{y}{2}, \frac{2}{\beta}\right)}{\operatorname{csgn}(\beta)^2 \sqrt{\beta^2 + 2 \cos(y) - 2}} = c_2 + x$ to $\frac{2\sqrt{\frac{\beta^2 + 2 \cos(y) - 2}{\beta^2}} \operatorname{InverseJacobiAM}\left(\frac{y}{2}, \frac{2}{\beta}\right)}{\sqrt{\beta^2 + 2 \cos(y) - 2}} = c_2 + x$ Initial conditions are used to solve for c_2 . Substituting $x = 0$ and $y = 0$ in the above solution gives an equation to solve for the constant of integration.

$$0 = c_2$$

$$c_2 = 0$$

Substituting c_2 found above in the general solution gives

$$\frac{2\sqrt{\frac{\beta^2 + 2 \cos(y) - 2}{\beta^2}} \operatorname{InverseJacobiAM}\left(\frac{y}{2}, \frac{2}{\beta}\right)}{\sqrt{\beta^2 + 2 \cos(y) - 2}} = x$$

The above simplifies to

$$-x\sqrt{\beta^2 + 2 \cos(y) - 2} + 2\sqrt{\frac{\beta^2 + 2 \cos(y) - 2}{\beta^2}} \operatorname{InverseJacobiAM}\left(\frac{y}{2}, \frac{2}{\beta}\right) = 0$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{1}{\sqrt{\beta^2 + 2 \cos(y) - 2}} dy = \int dx$$

$$-\frac{2\sqrt{\frac{\beta^2 + 2 \cos(y) - 2}{\beta^2}} \operatorname{InverseJacobiAM}\left(\frac{y}{2}, \frac{2}{\beta}\right)}{\operatorname{csgn}(\beta)^2 \sqrt{\beta^2 + 2 \cos(y) - 2}} = x + c_3$$

Simplifying the solution $-\frac{2\sqrt{\frac{\beta^2+2\cos(y)-2}{\beta^2}} \operatorname{InverseJacobiAM}\left(\frac{y}{2}, \frac{2}{\beta}\right)}{\operatorname{csgn}(\beta)^2 \sqrt{\beta^2+2\cos(y)-2}} = x+c_3$ to $-\frac{2\sqrt{\frac{\beta^2+2\cos(y)-2}{\beta^2}} \operatorname{InverseJacobiAM}\left(\frac{y}{2}, \frac{2}{\beta}\right)}{\sqrt{\beta^2+2\cos(y)-2}} = x+c_3$ Initial conditions are used to solve for c_3 . Substituting $x=0$ and $y=0$ in the above solution gives an equation to solve for the constant of integration.

$$0 = c_3$$

$$c_3 = 0$$

Substituting c_3 found above in the general solution gives

$$-\frac{2\sqrt{\frac{\beta^2+2\cos(y)-2}{\beta^2}} \operatorname{InverseJacobiAM}\left(\frac{y}{2}, \frac{2}{\beta}\right)}{\sqrt{\beta^2+2\cos(y)-2}} = x$$

The above simplifies to

$$-x\sqrt{\beta^2+2\cos(y)-2} - 2\sqrt{\frac{\beta^2+2\cos(y)-2}{\beta^2}} \operatorname{InverseJacobiAM}\left(\frac{y}{2}, \frac{2}{\beta}\right) = 0$$

Initial conditions are used to solve for the constants of integration.

Summary

The solution(s) found are the following

$$-x\sqrt{\beta^2+2\cos(y)-2} + 2\sqrt{\frac{\beta^2+2\cos(y)-2}{\beta^2}} \operatorname{InverseJacobiAM}\left(\frac{y}{2}, \frac{2}{\beta}\right) = 0 \quad (1)$$

$$-x\sqrt{\beta^2+2\cos(y)-2} - 2\sqrt{\frac{\beta^2+2\cos(y)-2}{\beta^2}} \operatorname{InverseJacobiAM}\left(\frac{y}{2}, \frac{2}{\beta}\right) = 0 \quad (2)$$

Verification of solutions

$$-x\sqrt{\beta^2+2\cos(y)-2} + 2\sqrt{\frac{\beta^2+2\cos(y)-2}{\beta^2}} \operatorname{InverseJacobiAM}\left(\frac{y}{2}, \frac{2}{\beta}\right) = 0$$

Verified OK.

$$-x\sqrt{\beta^2+2\cos(y)-2} - 2\sqrt{\frac{\beta^2+2\cos(y)-2}{\beta^2}} \operatorname{InverseJacobiAM}\left(\frac{y}{2}, \frac{2}{\beta}\right) = 0$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
`, `-> Computing symmetries using: way = exp_sym
-> Calling odsolve with the ODE`, (diff(_b(_a), _a))*_b(_a)+sin(_a) = 0, _b(_a)` *** Suble
  Methods for first order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  trying 1st order linear
  trying Bernoulli
  <- Bernoulli successful
<- differential order: 2; canonical coordinates successful
<- differential order 2; missing variables successful`
```

✓ Solution by Maple

Time used: 1.062 (sec). Leaf size: 53

```
dsolve([diff(y(x),x$2)+sin(y(x))=0,y(0) = 0, D(y)(0) = beta],y(x), singsol=all)
```

$$y(x) = \text{RootOf} \left(- \left(\int_0^{-z} \frac{1}{\sqrt{2 \cos(_a) + \beta^2 - 2}} d_a \right) + x \right)$$
$$y(x) = \text{RootOf} \left(\int_0^{-z} \frac{1}{\sqrt{2 \cos(_a) + \beta^2 - 2}} d_a + x \right)$$

✓ Solution by Mathematica

Time used: 0.621 (sec). Leaf size: 19

```
DSolve[{y''[x]+Sin[y[x]]==0,{y[0]==0,y'[0]==\[Beta]}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2 \operatorname{JacobiAmplitude}\left(\frac{x\beta}{2}, \frac{4}{\beta^2}\right)$$

23.10 problem 5(c)

23.10.1 Solving as second order ode can be made integrable ode 1980

23.10.2 Solving as second order ode missing x ode 1982

Internal problem ID [6100]

Internal file name [OUTPUT/5348_Sunday_June_05_2022_03_35_01_PM_69825552/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 6. Existence and uniqueness of solutions to systems and nth order equations. Page 238

Problem number: 5(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_ode_missing_x", "second_order_ode_can_be_made_integrable"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x], [_2nd_order , _reducible , _mu_x_y1]]
```

$$y'' + \sin(y) = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 2]$$

23.10.1 Solving as second order ode can be made integrable ode

Multiplying the ode by y' gives

$$y'y'' + y'\sin(y) = 0$$

Integrating the above w.r.t x gives

$$\int (y'y'' + y'\sin(y)) dx = 0$$
$$\frac{y'^2}{2} - \cos(y) = c_2$$

Which is now solved for y . Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \sqrt{2 \cos (y) + 2c_1} \quad (1)$$

$$y' = -\sqrt{2 \cos (y) + 2c_1} \quad (2)$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int \frac{1}{\sqrt{2 \cos (y) + 2c_1}} dy = \int dx$$

$$\frac{2\sqrt{\frac{\cos(y)+c_1}{c_1+1}} \operatorname{InverseJacobiAM}\left(\frac{y}{2}, \frac{2}{\sqrt{2+2c_1}}\right)}{\sqrt{2 \cos (y) + 2c_1}} = c_2 + x$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{1}{\sqrt{2 \cos (y) + 2c_1}} dy = \int dx$$

$$-\frac{2\sqrt{\frac{\cos(y)+c_1}{c_1+1}} \operatorname{InverseJacobiAM}\left(\frac{y}{2}, \frac{2}{\sqrt{2+2c_1}}\right)}{\sqrt{2 \cos (y) + 2c_1}} = x + c_3$$

Initial conditions are used to solve for the constants of integration.

Looking at the First solution

$$\frac{2\sqrt{\frac{\cos(y)+c_1}{c_1+1}} \operatorname{InverseJacobiAM}\left(\frac{y}{2}, \frac{2}{\sqrt{2+2c_1}}\right)}{\sqrt{2 \cos (y) + 2c_1}} = c_2 + x \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_2 \quad (1A)$$

Taking derivative of the solution gives

Expression too large to display

substituting $y' = 2$ and $x = 0$ in the above gives

$$\text{Expression too large to display} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. There is no solution for the constants of integrations. This solution is removed.

Looking at the Second solution

$$-\frac{2\sqrt{\frac{\cos(y)+c_1}{c_1+1}} \operatorname{InverseJacobiAM}\left(\frac{y}{2}, \frac{2}{\sqrt{2+2c_1}}\right)}{\sqrt{2\cos(y)+2c_1}} = x + c_3 \quad (2)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_3 \quad (1A)$$

Taking derivative of the solution gives

$$\text{Expression too large to display}$$

substituting $y' = 2$ and $x = 0$ in the above gives

$$\text{Expression too large to display} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_3\}$. There is no solution for the constants of integrations. This solution is removed.

Verification of solutions N/A

23.10.2 Solving as second order ode missing x ode

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable y an independent variable. Using

$$y' = p(y)$$

Then

$$\begin{aligned} y'' &= \frac{dp}{dx} \\ &= \frac{dy}{dx} \frac{dp}{dy} \\ &= p \frac{dp}{dy} \end{aligned}$$

Hence the ode becomes

$$p(y) \left(\frac{d}{dy} p(y) \right) = -\sin(y)$$

Which is now solved as first order ode for $p(y)$. In canonical form the ODE is

$$\begin{aligned} p' &= F(y, p) \\ &= f(y)g(p) \\ &= -\frac{\sin(y)}{p} \end{aligned}$$

Where $f(y) = -\sin(y)$ and $g(p) = \frac{1}{p}$. Integrating both sides gives

$$\begin{aligned} \frac{1}{p} dp &= -\sin(y) dy \\ \int \frac{1}{p} dp &= \int -\sin(y) dy \\ \frac{p^2}{2} &= \cos(y) + c_1 \end{aligned}$$

The solution is

$$\frac{p(y)^2}{2} - \cos(y) - c_1 = 0$$

Initial conditions are used to solve for c_1 . Substituting $y = 0$ and $p = 2$ in the above solution gives an equation to solve for the constant of integration.

$$1 - c_1 = 0$$

$$c_1 = 1$$

Substituting c_1 found above in the general solution gives

$$\frac{p^2}{2} - \cos(y) - 1 = 0$$

Solving for $p(y)$ from the above gives

$$p(y) = \sqrt{2} \sqrt{\cos(y) + 1}$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = \sqrt{2} \sqrt{\cos(y) + 1}$$

Integrating both sides gives

$$\int \frac{\sqrt{2}}{2\sqrt{\cos(y) + 1}} dy = \int dx$$

$$\int^y \frac{\sqrt{2}}{2\sqrt{\cos(_a) + 1}} d_a = c_2 + x$$

Initial conditions are used to solve for c_2 . Substituting $x = 0$ and $y = 0$ in the above solution gives an equation to solve for the constant of integration.

$$\int^0 \frac{\sqrt{2}}{2\sqrt{\cos(_a) + 1}} d_a = c_2$$

$$c_2 = \frac{\left(\int^0 \sec\left(\frac{_a}{2}\right) d_a\right)}{2}$$

Substituting c_2 found above in the general solution gives

$$\int^y \frac{\sqrt{2}}{2\sqrt{\cos(_a) + 1}} d_a = \frac{\left(\int^0 \sec\left(\frac{_a}{2}\right) d_a\right)}{2} + x$$

Simplifying the solution $\frac{\left(\int^y \operatorname{csgn}\left(\cos\left(\frac{_a}{2}\right)\right) \sec\left(\frac{_a}{2}\right) d_a\right)}{2} = \frac{\left(\int^0 \sec\left(\frac{_a}{2}\right) d_a\right)}{2} + x$ to $\frac{\left(\int^y \sec\left(\frac{_a}{2}\right) d_a\right)}{2} = \frac{\left(\int^0 \sec\left(\frac{_a}{2}\right) d_a\right)}{2} + x$ Initial conditions are used to solve for the constants of integration.

Summary

The solution(s) found are the following

$$\frac{\left(\int^y \sec\left(\frac{_a}{2}\right) d_a\right)}{2} = \frac{\left(\int^0 \sec\left(\frac{_a}{2}\right) d_a\right)}{2} + x \quad (1)$$

Verification of solutions

$$\frac{\left(\int^y \sec\left(\frac{_a}{2}\right) d_a\right)}{2} = \frac{\left(\int^0 \sec\left(\frac{_a}{2}\right) d_a\right)}{2} + x$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
`, `-> Computing symmetries using: way = exp_sym
-> Calling odsolve with the ODE`, (diff(_b(_a), _a))*_b(_a)+sin(_a) = 0, _b(_a)` *** Suble
  Methods for first order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  trying 1st order linear
  trying Bernoulli
  <- Bernoulli successful
<- differential order: 2; canonical coordinates successful
<- differential order 2; missing variables successful`
```

✓ Solution by Maple

Time used: 1.296 (sec). Leaf size: 23

```
dsolve([diff(y(x),x$2)+sin(y(x))=0,y(0) = 0, D(y)(0) = 2],y(x), singsol=all)
```

$$y(x) = \text{RootOf} \left(- \left(\int_0^{-Z} \sec \left(\frac{a}{2} \right) \text{csgn} \left(\cos \left(\frac{a}{2} \right) \right) d_a \right) + 2x \right)$$

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[{y'[x]+Sin[y[x]]==0,{y[0]==0,y'[0]==2}},y[x],x,IncludeSingularSolutions -> True]
```

```
{}
```

24 Chapter 6. Existence and uniqueness of solutions to systems and nth order equations.

Page 250

24.1 problem 3	1987
24.2 problem 4	1995
24.3 problem 5	2004

24.1 problem 3

24.1.1 Solution using Matrix exponential method 1987

24.1.2 Solution using explicit Eigenvalue and Eigenvector method . . . 1988

Internal problem ID [6101]

Internal file name [OUTPUT/5349_Sunday_June_05_2022_03_35_04_PM_96258689/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 6. Existence and uniqueness of solutions to systems and nth order equations. Page 250

Problem number: 3.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**system of linear ODEs**"

Solve

$$y_1'(x) = y_1(x)$$

$$y_2'(x) = y_1(x) + y_2(x)$$

With initial conditions

$$[y_1(0) = 1, y_2(0) = 2]$$

24.1.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(x) = A\vec{x}(x)$$

Or

$$\begin{bmatrix} y_1'(x) \\ y_2'(x) \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} y_1(x) \\ y_2(x) \end{bmatrix}$$

For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} e^x & 0 \\ x e^x & e^x \end{bmatrix}$$

Therefore the homogeneous solution is

$$\begin{aligned}\vec{x}_h(x) &= e^{Ax} \vec{x}_0 \\ &= \begin{bmatrix} e^x & 0 \\ x e^x & e^x \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix} \\ &= \begin{bmatrix} e^x \\ x e^x + 2 e^x \end{bmatrix} \\ &= \begin{bmatrix} e^x \\ e^x(x + 2) \end{bmatrix}\end{aligned}$$

Since no forcing function is given, then the final solution is $\vec{x}_h(x)$ above.

24.1.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(x) = A \vec{x}(x)$$

Or

$$\begin{bmatrix} y_1'(x) \\ y_2'(x) \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} y_1(x) \\ y_2(x) \end{bmatrix}$$

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det \left(\begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) = 0$$

Therefore

$$\det \left(\begin{bmatrix} 1 - \lambda & 0 \\ 1 & 1 - \lambda \end{bmatrix} \right) = 0$$

Since the matrix A is triangular matrix, then the determinant is the product of the elements along the diagonal. Therefore the above becomes

$$(1 - \lambda)(1 - \lambda) = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = 1$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
1	1	real eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = 1$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} - (1) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} 0 & 0 & 0 \\ 1 & 0 & 0 \end{array} \right]$$

Since the current pivot $A(1, 1)$ is zero, then the current pivot row is replaced with a row with a non-zero pivot. Swapping row 1 and row 2 gives

$$\left[\begin{array}{cc|c} 1 & 0 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = 0\}$

Hence the solution is

$$\begin{bmatrix} 0 \\ t \end{bmatrix} = \begin{bmatrix} 0 \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} 0 \\ t \end{bmatrix} = t \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
1	2	1	Yes	$\begin{bmatrix} 0 \\ 1 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. eigenvalue 1 is real and repeated eigenvalue of multiplicity 2. There are two possible cases that can happen. This is illustrated in this diagram




Figure 229: Possible case for repeated λ of multiplicity 2

This eigenvalue has algebraic multiplicity of 2, and geometric multiplicity 1, therefore this is defective eigenvalue. The defect is 1. This falls into case 2 shown above. We need to generate the missing additional generalized eigenvector \vec{v}_2 by solving

$$(A - \lambda I) \vec{v}_2 = \vec{v}_1$$

Where \vec{v}_1 is the normal (rank 1) eigenvector found above. Hence we need to solve

$$\left(\begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} - (1) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

Solving for \vec{v}_2 gives

$$\vec{v}_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

We have found two generalized eigenvectors for eigenvalue 1. Therefore the two basis solution associated with this eigenvalue are

$$\begin{aligned}\vec{x}_1(x) &= \vec{v}_1 e^{\lambda t} \\ &= \begin{bmatrix} 0 \\ 1 \end{bmatrix} e^x \\ &= \begin{bmatrix} 0 \\ e^x \end{bmatrix}\end{aligned}$$

And

$$\begin{aligned}\vec{x}_2(x) &= (\vec{v}_1 x + \vec{v}_2) e^{\lambda t} \\ &= \left(\begin{bmatrix} 0 \\ 1 \end{bmatrix} t + \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right) e^x \\ &= \begin{bmatrix} e^x \\ e^x(1+x) \end{bmatrix}\end{aligned}$$

Therefore the final solution is

$$\vec{x}_h(x) = c_1 \vec{x}_1(x) + c_2 \vec{x}_2(x)$$

Which is written as

$$\begin{bmatrix} y_1(x) \\ y_2(x) \end{bmatrix} = c_1 \begin{bmatrix} 0 \\ e^x \end{bmatrix} + c_2 \begin{bmatrix} e^x \\ e^x(1+x) \end{bmatrix}$$

Which becomes

$$\begin{bmatrix} y_1(x) \\ y_2(x) \end{bmatrix} = \begin{bmatrix} c_2 e^x \\ e^x(c_2 x + c_1 + c_2) \end{bmatrix}$$

Since initial conditions are given, the solution above needs to be updated by solving for the constants of integrations using the given initial conditions

$$\begin{bmatrix} y_1(0) = 1 \\ y_2(0) = 2 \end{bmatrix} \tag{1}$$

Substituting initial conditions into the above solution at $x = 0$ gives

$$\begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} c_2 \\ c_1 + c_2 \end{bmatrix}$$

Solving for the constants of integrations gives

$$\begin{bmatrix} c_1 = 1 \\ c_2 = 1 \end{bmatrix}$$

Substituting these constants back in original solution in Eq. (1) gives

$$\begin{bmatrix} y_1(x) \\ y_2(x) \end{bmatrix} = \begin{bmatrix} e^x \\ e^x(x + 2) \end{bmatrix}$$

The following is the phase plot of the system.






Figure 230: Phase plot

The following are plots of each solution.



✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 16

```
dsolve([diff(y__1(x),x) = y__1(x), diff(y__2(x),x) = y__1(x)+y__2(x), y__1(0) = 1, y__2(0) = 2])
```

$$y_1(x) = e^x$$

$$y_2(x) = (x + 2)e^x$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 18

```
DSolve[{y1'[x]==y1[x],y2'[x]==y1[x]+y2[x]},{y1[0]==1,y2[0]==2},{y1[x],y2[x]},x,IncludeSingularSolutions->False]
```

$$y1(x) \rightarrow e^x$$

$$y2(x) \rightarrow e^x(x + 2)$$

24.2 problem 4

24.2.1 Solution using Matrix exponential method 1995

24.2.2 Solution using explicit Eigenvalue and Eigenvector method . . . 1996

Internal problem ID [6102]

Internal file name [OUTPUT/5350_Sunday_June_05_2022_03_35_05_PM_35774872/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 6. Existence and uniqueness of solutions to systems and nth order equations. Page 250

Problem number: 4.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**system of linear ODEs**"

Solve

$$\begin{aligned}y_1'(x) &= y_2(x) \\y_2'(x) &= 6y_1(x) + y_2(x)\end{aligned}$$

With initial conditions

$$[y_1(0) = 1, y_2(0) = -1]$$

24.2.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(x) = A\vec{x}(x)$$

Or

$$\begin{bmatrix} y_1'(x) \\ y_2'(x) \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 6 & 1 \end{bmatrix} \begin{bmatrix} y_1(x) \\ y_2(x) \end{bmatrix}$$

For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} \frac{(2e^{5x}+3)e^{-2x}}{5} & \frac{(e^{5x}-1)e^{-2x}}{5} \\ \frac{6(e^{5x}-1)e^{-2x}}{5} & \frac{(3e^{5x}+2)e^{-2x}}{5} \end{bmatrix}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 \vec{x}_h(x) &= e^{Ax} \vec{x}_0 \\
 &= \begin{bmatrix} \frac{(2e^{5x}+3)e^{-2x}}{5} & \frac{(e^{5x}-1)e^{-2x}}{5} \\ \frac{6(e^{5x}-1)e^{-2x}}{5} & \frac{(3e^{5x}+2)e^{-2x}}{5} \end{bmatrix} \begin{bmatrix} 1 \\ -1 \end{bmatrix} \\
 &= \begin{bmatrix} \frac{(2e^{5x}+3)e^{-2x}}{5} - \frac{(e^{5x}-1)e^{-2x}}{5} \\ \frac{6(e^{5x}-1)e^{-2x}}{5} - \frac{(3e^{5x}+2)e^{-2x}}{5} \end{bmatrix} \\
 &= \begin{bmatrix} \frac{(e^{5x}+4)e^{-2x}}{5} \\ \frac{(3e^{5x}-8)e^{-2x}}{5} \end{bmatrix}
 \end{aligned}$$

Since no forcing function is given, then the final solution is $\vec{x}_h(x)$ above.

24.2.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(x) = A\vec{x}(x)$$

Or

$$\begin{bmatrix} y_1'(x) \\ y_2'(x) \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 6 & 1 \end{bmatrix} \begin{bmatrix} y_1(x) \\ y_2(x) \end{bmatrix}$$

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det\left(\begin{bmatrix} 0 & 1 \\ 6 & 1 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}\right) = 0$$

Therefore

$$\det\left(\begin{bmatrix} -\lambda & 1 \\ 6 & 1 - \lambda \end{bmatrix}\right) = 0$$

Which gives the characteristic equation

$$\lambda^2 - \lambda - 6 = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = -2$$

$$\lambda_2 = 3$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
-2	1	real eigenvalue
3	1	real eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = -2$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 0 & 1 \\ 6 & 1 \end{bmatrix} - (-2) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 1 \\ 6 & 3 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} 2 & 1 & 0 \\ 6 & 3 & 0 \end{array} \right]$$

$$R_2 = R_2 - 3R_1 \implies \left[\begin{array}{cc|c} 2 & 1 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} 2 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = -\frac{t}{2}\}$

Hence the solution is

$$\begin{bmatrix} -\frac{t}{2} \\ t \end{bmatrix} = \begin{bmatrix} -\frac{t}{2} \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} -\frac{t}{2} \\ t \end{bmatrix} = t \begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix} = \begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix}$$

Which is normalized to

$$\begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix} = \begin{bmatrix} -1 \\ 2 \end{bmatrix}$$

Considering the eigenvalue $\lambda_2 = 3$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 0 & 1 \\ 6 & 1 \end{bmatrix} - (3) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} -3 & 1 \\ 6 & -2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} -3 & 1 & 0 \\ 6 & -2 & 0 \end{array} \right]$$

$$R_2 = R_2 + 2R_1 \implies \left[\begin{array}{cc|c} -3 & 1 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} -3 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = \frac{t}{3}\}$

Hence the solution is

$$\begin{bmatrix} \frac{t}{3} \\ t \end{bmatrix} = \begin{bmatrix} \frac{t}{3} \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} \frac{t}{3} \\ t \end{bmatrix} = t \begin{bmatrix} \frac{1}{3} \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} \frac{1}{3} \\ 1 \end{bmatrix} = \begin{bmatrix} \frac{1}{3} \\ 1 \end{bmatrix}$$

Which is normalized to

$$\begin{bmatrix} \frac{1}{3} \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 3 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
-2	1	1	No	$\begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix}$
3	1	1	No	$\begin{bmatrix} \frac{1}{3} \\ 1 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. Since eigenvalue -2 is real and distinct then the corresponding eigenvector solution is

$$\begin{aligned}\vec{x}_1(x) &= \vec{v}_1 e^{-2x} \\ &= \begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix} e^{-2x}\end{aligned}$$

Since eigenvalue 3 is real and distinct then the corresponding eigenvector solution is

$$\begin{aligned}\vec{x}_2(x) &= \vec{v}_2 e^{3x} \\ &= \begin{bmatrix} \frac{1}{3} \\ 1 \end{bmatrix} e^{3x}\end{aligned}$$

Therefore the final solution is

$$\vec{x}_h(x) = c_1 \vec{x}_1(x) + c_2 \vec{x}_2(x)$$

Which is written as

$$\begin{bmatrix} y_1(x) \\ y_2(x) \end{bmatrix} = c_1 \begin{bmatrix} -\frac{e^{-2x}}{2} \\ e^{-2x} \end{bmatrix} + c_2 \begin{bmatrix} \frac{e^{3x}}{3} \\ e^{3x} \end{bmatrix}$$

Which becomes

$$\begin{bmatrix} y_1(x) \\ y_2(x) \end{bmatrix} = \begin{bmatrix} \frac{(2c_2 e^{5x} - 3c_1) e^{-2x}}{6} \\ (c_2 e^{5x} + c_1) e^{-2x} \end{bmatrix}$$

Since initial conditions are given, the solution above needs to be updated by solving for the constants of integrations using the given initial conditions

$$\begin{bmatrix} y_1(0) = 1 \\ y_2(0) = -1 \end{bmatrix} \quad (1)$$

Substituting initial conditions into the above solution at $x = 0$ gives

$$\begin{bmatrix} 1 \\ -1 \end{bmatrix} = \begin{bmatrix} \frac{c_2}{3} - \frac{c_1}{2} \\ c_2 + c_1 \end{bmatrix}$$

Solving for the constants of integrations gives

$$\begin{bmatrix} c_1 = -\frac{8}{5} \\ c_2 = \frac{3}{5} \end{bmatrix}$$

Substituting these constants back in original solution in Eq. (1) gives

$$\begin{bmatrix} y_1(x) \\ y_2(x) \end{bmatrix} = \begin{bmatrix} \frac{\left(\frac{6e^{5x}}{5} + \frac{24}{5}\right)e^{-2x}}{6} \\ \left(\frac{3e^{5x}}{5} - \frac{8}{5}\right)e^{-2x} \end{bmatrix}$$

The following is the phase plot of the system.






Figure 231: Phase plot

The following are plots of each solution.



✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 34

```
dsolve([diff(y__1(x),x) = y__2(x), diff(y__2(x),x) = 6*y__1(x)+y__2(x), y__1(0) = 1, y__2(0)
```

$$y_1(x) = \frac{4e^{-2x}}{5} + \frac{e^{3x}}{5}$$
$$y_2(x) = -\frac{8e^{-2x}}{5} + \frac{3e^{3x}}{5}$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 42

```
DSolve[{y1'[x]==y2[x],y2'[x]==6*y1[x]+y2[x]},{y1[0]==1,y2[0]==-1},{y1[x],y2[x]},x,IncludeSin
```

$$y_1(x) \rightarrow \frac{1}{5}e^{-2x}(e^{5x} + 4)$$
$$y_2(x) \rightarrow \frac{1}{5}e^{-2x}(3e^{5x} - 8)$$

24.3 problem 5

24.3.1 Solution using Matrix exponential method 2004

24.3.2 Solution using explicit Eigenvalue and Eigenvector method . . . 2006

Internal problem ID [6103]

Internal file name [OUTPUT/5351_Sunday_June_05_2022_03_35_06_PM_78469615/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 6. Existence and uniqueness of solutions to systems and nth order equations. Page 250

Problem number: 5.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**system of linear ODEs**"

Solve

$$\begin{aligned}y_1'(x) &= y_1(x) + y_2(x) \\y_2'(x) &= y_1(x) + y_2(x) + e^{3x}\end{aligned}$$

With initial conditions

$$[y_1(0) = 0, y_2(0) = 0]$$

24.3.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(x) = A\vec{x}(x) + \vec{G}(x)$$

Or

$$\begin{bmatrix} y_1'(x) \\ y_2'(x) \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} y_1(x) \\ y_2(x) \end{bmatrix} + \begin{bmatrix} 0 \\ e^{3x} \end{bmatrix}$$

Since the system is nonhomogeneous, then the solution is given by

$$\vec{x}(x) = \vec{x}_h(x) + \vec{x}_p(x)$$

Where $\vec{x}_h(x)$ is the homogeneous solution to $\vec{x}'(x) = A\vec{x}(x)$ and $\vec{x}_p(x)$ is a particular solution to $\vec{x}'(x) = A\vec{x}(x) + \vec{G}(x)$. The particular solution will be found using variation of parameters method applied to the fundamental matrix. For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} \frac{1}{2} + \frac{e^{2x}}{2} & \frac{e^{2x}}{2} - \frac{1}{2} \\ \frac{e^{2x}}{2} - \frac{1}{2} & \frac{1}{2} + \frac{e^{2x}}{2} \end{bmatrix}$$

Therefore the homogeneous solution is

$$\begin{aligned} \vec{x}_h(x) &= e^{Ax} \vec{x}_0 \\ &= \begin{bmatrix} \frac{1}{2} + \frac{e^{2x}}{2} & \frac{e^{2x}}{2} - \frac{1}{2} \\ \frac{e^{2x}}{2} - \frac{1}{2} & \frac{1}{2} + \frac{e^{2x}}{2} \end{bmatrix} \begin{bmatrix} 0 \\ 0 \end{bmatrix} \\ &= \begin{bmatrix} 0 \\ 0 \end{bmatrix} \end{aligned}$$

The particular solution given by

$$\vec{x}_p(x) = e^{Ax} \int e^{-Ax} \vec{G}(x) dx$$

But

$$\begin{aligned} e^{-Ax} &= (e^{Ax})^{-1} \\ &= \begin{bmatrix} \frac{1}{2} + \frac{e^{-2x}}{2} & -\frac{1}{2} + \frac{e^{-2x}}{2} \\ -\frac{1}{2} + \frac{e^{-2x}}{2} & \frac{1}{2} + \frac{e^{-2x}}{2} \end{bmatrix} \end{aligned}$$

Hence

$$\begin{aligned} \vec{x}_p(x) &= \begin{bmatrix} \frac{1}{2} + \frac{e^{2x}}{2} & \frac{e^{2x}}{2} - \frac{1}{2} \\ \frac{e^{2x}}{2} - \frac{1}{2} & \frac{1}{2} + \frac{e^{2x}}{2} \end{bmatrix} \int \begin{bmatrix} \frac{1}{2} + \frac{e^{-2x}}{2} & -\frac{1}{2} + \frac{e^{-2x}}{2} \\ -\frac{1}{2} + \frac{e^{-2x}}{2} & \frac{1}{2} + \frac{e^{-2x}}{2} \end{bmatrix} \begin{bmatrix} 0 \\ e^{3x} \end{bmatrix} dx \\ &= \begin{bmatrix} \frac{1}{2} + \frac{e^{2x}}{2} & \frac{e^{2x}}{2} - \frac{1}{2} \\ \frac{e^{2x}}{2} - \frac{1}{2} & \frac{1}{2} + \frac{e^{2x}}{2} \end{bmatrix} \begin{bmatrix} -\frac{e^{3x}}{6} + \frac{e^x}{2} \\ \frac{e^{3x}}{6} + \frac{e^x}{2} \end{bmatrix} \\ &= \begin{bmatrix} \frac{e^{3x}}{3} \\ \frac{2e^{3x}}{3} \end{bmatrix} \end{aligned}$$

Hence the complete solution is

$$\begin{aligned}\vec{x}(x) &= \vec{x}_h(x) + \vec{x}_p(x) \\ &= \begin{bmatrix} \frac{e^{3x}}{3} \\ \frac{2e^{3x}}{3} \end{bmatrix}\end{aligned}$$

24.3.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(x) = A\vec{x}(x) + \vec{G}(x)$$

Or

$$\begin{bmatrix} y_1'(x) \\ y_2'(x) \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} y_1(x) \\ y_2(x) \end{bmatrix} + \begin{bmatrix} 0 \\ e^{3x} \end{bmatrix}$$

Since the system is nonhomogeneous, then the solution is given by

$$\vec{x}(x) = \vec{x}_h(x) + \vec{x}_p(x)$$

Where $\vec{x}_h(x)$ is the homogeneous solution to $\vec{x}'(x) = A\vec{x}(x)$ and $\vec{x}_p(x)$ is a particular solution to $\vec{x}'(x) = A\vec{x}(x) + \vec{G}(x)$. The particular solution will be found using variation of parameters method applied to the fundamental matrix.

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det\left(\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}\right) = 0$$

Therefore

$$\det\left(\begin{bmatrix} 1 - \lambda & 1 \\ 1 & 1 - \lambda \end{bmatrix}\right) = 0$$

Which gives the characteristic equation

$$\lambda^2 - 2\lambda = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = 0$$

$$\lambda_2 = 2$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
0	1	real eigenvalue
2	1	real eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = 0$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} - (0) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} 1 & 1 & 0 \\ 1 & 1 & 0 \end{array} \right]$$

$$R_2 = R_2 - R_1 \implies \left[\begin{array}{cc|c} 1 & 1 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = -t\}$

Hence the solution is

$$\begin{bmatrix} -t \\ t \end{bmatrix} = \begin{bmatrix} -t \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} -t \\ t \end{bmatrix} = t \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} -t \\ t \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

Considering the eigenvalue $\lambda_2 = 2$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} - (2) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$
$$\begin{bmatrix} -1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} -1 & 1 & 0 \\ 1 & -1 & 0 \end{array} \right]$$

$$R_2 = R_2 + R_1 \implies \left[\begin{array}{cc|c} -1 & 1 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} -1 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = t\}$

Hence the solution is

$$\begin{bmatrix} t \\ t \end{bmatrix} = \begin{bmatrix} t \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} t \\ t \end{bmatrix} = t \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} t \\ t \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
0	1	1	No	$\begin{bmatrix} -1 \\ 1 \end{bmatrix}$
2	1	1	No	$\begin{bmatrix} 1 \\ 1 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. Since eigenvalue 0 is real and distinct then the corresponding eigenvector solution is

$$\begin{aligned} \vec{x}_1(x) &= \vec{v}_1 e^0 \\ &= \begin{bmatrix} -1 \\ 1 \end{bmatrix} e^0 \end{aligned}$$

Since eigenvalue 2 is real and distinct then the corresponding eigenvector solution is

$$\begin{aligned}\vec{x}_2(x) &= \vec{v}_2 e^{2x} \\ &= \begin{bmatrix} 1 \\ 1 \end{bmatrix} e^{2x}\end{aligned}$$

Therefore the homogeneous solution is

$$\vec{x}_h(x) = c_1 \vec{x}_1(x) + c_2 \vec{x}_2(x)$$

Which is written as

$$\begin{bmatrix} y_1(x) \\ y_2(x) \end{bmatrix} = c_1 \begin{bmatrix} -1 \\ 1 \end{bmatrix} + c_2 \begin{bmatrix} e^{2x} \\ e^{2x} \end{bmatrix}$$

Now that we found homogeneous solution above, we need to find a particular solution $\vec{x}_p(x)$. We will use Variation of parameters. The fundamental matrix is

$$\Phi = \begin{bmatrix} \vec{x}_1 & \vec{x}_2 & \cdots \end{bmatrix}$$

Where \vec{x}_i are the solution basis found above. Therefore the fundamental matrix is

$$\Phi(x) = \begin{bmatrix} -1 & e^{2x} \\ 1 & e^{2x} \end{bmatrix}$$

The particular solution is then given by

$$\vec{x}_p(x) = \Phi \int \Phi^{-1} \vec{G}(x) dx$$

But

$$\Phi^{-1} = \begin{bmatrix} -\frac{1}{2} & \frac{1}{2} \\ \frac{e^{-2x}}{2} & \frac{e^{-2x}}{2} \end{bmatrix}$$

Hence

$$\begin{aligned}
 \vec{x}_p(x) &= \begin{bmatrix} -1 & e^{2x} \\ 1 & e^{2x} \end{bmatrix} \int \begin{bmatrix} -\frac{1}{2} & \frac{1}{2} \\ \frac{e^{-2x}}{2} & \frac{e^{-2x}}{2} \end{bmatrix} \begin{bmatrix} 0 \\ e^{3x} \end{bmatrix} dx \\
 &= \begin{bmatrix} -1 & e^{2x} \\ 1 & e^{2x} \end{bmatrix} \int \begin{bmatrix} \frac{e^{3x}}{2} \\ \frac{e^x}{2} \end{bmatrix} dx \\
 &= \begin{bmatrix} -1 & e^{2x} \\ 1 & e^{2x} \end{bmatrix} \begin{bmatrix} \frac{e^{3x}}{6} \\ \frac{e^x}{2} \end{bmatrix} \\
 &= \begin{bmatrix} \frac{e^{3x}}{3} \\ \frac{2e^{3x}}{3} \end{bmatrix}
 \end{aligned}$$

Now that we found particular solution, the final solution is

$$\begin{aligned}
 \vec{x}(x) &= \vec{x}_h(x) + \vec{x}_p(x) \\
 \begin{bmatrix} y_1(x) \\ y_2(x) \end{bmatrix} &= \begin{bmatrix} -c_1 \\ c_1 \end{bmatrix} + \begin{bmatrix} c_2 e^{2x} \\ c_2 e^{2x} \end{bmatrix} + \begin{bmatrix} \frac{e^{3x}}{3} \\ \frac{2e^{3x}}{3} \end{bmatrix}
 \end{aligned}$$

Which becomes

$$\begin{bmatrix} y_1(x) \\ y_2(x) \end{bmatrix} = \begin{bmatrix} -c_1 + c_2 e^{2x} + \frac{e^{3x}}{3} \\ c_1 + c_2 e^{2x} + \frac{2e^{3x}}{3} \end{bmatrix}$$

Since initial conditions are given, the solution above needs to be updated by solving for the constants of integrations using the given initial conditions

$$\begin{bmatrix} y_1(0) = 0 \\ y_2(0) = 0 \end{bmatrix} \tag{1}$$

Substituting initial conditions into the above solution at $x = 0$ gives


$$\begin{bmatrix} 0 \\ 0 \end{bmatrix} = \begin{bmatrix} -c_1 + c_2 + \frac{1}{3} \\ c_1 + c_2 + \frac{2}{3} \end{bmatrix}$$

Solving for the constants of integrations gives



$$\begin{bmatrix} c_1 = -\frac{1}{6} \\ c_2 = -\frac{1}{2} \end{bmatrix}$$

Substituting these constants back in original solution in Eq. (1) gives

$$\begin{bmatrix} y_1(x) \\ y_2(x) \end{bmatrix} = \begin{bmatrix} \frac{1}{6} - \frac{e^{2x}}{2} + \frac{e^{3x}}{3} \\ -\frac{1}{6} - \frac{e^{2x}}{2} + \frac{2e^{3x}}{3} \end{bmatrix}$$



The following are plots of each solution.



✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 36

```
dsolve([diff(y__1(x),x) = y__1(x)+y__2(x), diff(y__2(x),x) = y__1(x)+y__2(x)+exp(3*x), y__1(
```

$$y_1(x) = -\frac{e^{2x}}{2} + \frac{e^{3x}}{3} + \frac{1}{6}$$
$$y_2(x) = -\frac{e^{2x}}{2} + \frac{2e^{3x}}{3} - \frac{1}{6}$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 46

```
DSolve[{y1'[x]==y1[x]+y2[x],y2'[x]==y1[x]+y2[x]+Exp[3*x]},{y1[0]==0,y2[0]==0},{y1[x],y2[x]},
```

$$y_1(x) \rightarrow \frac{1}{6}(e^x - 1)^2(2e^x + 1)$$
$$y_2(x) \rightarrow \frac{1}{6}(-3e^{2x} + 4e^{3x} - 1)$$

25 Chapter 6. Existence and uniqueness of solutions to systems and nth order equations.

Page 254

25.1 problem 2 2015

25.1 problem 2

Internal problem ID [6104]

Internal file name [OUTPUT/5352_Sunday_June_05_2022_03_35_08_PM_97644046/index.tex]

Book: An introduction to Ordinary Differential Equations. Earl A. Coddington. Dover. NY 1961

Section: Chapter 6. Existence and uniqueness of solutions to systems and nth order equations. Page 254

Problem number: 2.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**system of linear ODEs**" Unable to solve or complete t

Solve

$$y_1'(x) = 3y_1(x) + xy_3(x)$$

$$y_2'(x) = y_2(x) + x^3y_3(x)$$

$$y_3'(x) = 2xy_2(x) - y_2(x) + e^x y_3(x)$$

Does not currently support non autonomous system of first order linear differential equations. The following is the phase plot

X Solution by Maple

```
dsolve([diff(y__1(x),x)=3*y__1(x)+x*y__3(x),diff(y__2(x),x)=y__2(x)+x^3*y__3(x),diff(y__3(x)
```

No solution found

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[{y1'[x]==3*y1[x]+x*y3[x],y2'[x]==y2[x]+x^3*y3[x],y3'[x]==2*x*y1[x]-y2[x]+Exp[x]*y3[x]
```

Not solved