


A Solution Manual For

Applied Differential equations, Newby Curle.

Van Nostrand Reinhold. 1972



Nasser M. Abbasi

December 13, 2025

Compiled on December 13, 2025 at 7:55am

Contents

1	Lookup tables for all problems in current book	3
2	Book Solved Problems	5

Chapter

1

Lookup tables for all problems in current book

Local contents

1.1	Examples, page 35	4
-----	-----------------------------	---

1.1 Examples, page 35

Table 1.1: Lookup table for all problems in current section

ID	problem	ODE	Solved?	Maple	Mma	Sympy
4086	1	$y = y' + \frac{y'^2}{2}$	✓	✓	✓	✓
4087	2	$(-y'x + y)^2 = 1 + y'^2$	✓	✓	✓	✗
4088	3	$-x + y = y'^2 \left(1 - \frac{2y'}{3}\right)$	✓	✓	✓	✗
4089	4	$x^2 y' = (-1 + y)x + (-1 + y)^2$	✓	✓	✓	✓

Chapter 2

Book Solved Problems

Local contents

2.1	Examples, page 35	6
-----	-----------------------------	---

2.1 Examples, page 35

Local contents

2.1.1	Problem 1	7
2.1.2	Problem 2	13
2.1.3	Problem 3	19
2.1.4	Problem 4	24

2.1.1 Problem 1

Local contents

2.1.1.1	Solved using first_order_ode_dAlembert	7
2.1.1.2	Solved using first_order_ode_parametric method	9
2.1.1.3	✓ Maple	10
2.1.1.4	✓ Mathematica	12
2.1.1.5	✓ Sympy	12

Internal problem ID [4086]

Book : Applied Differential equations, Newby Curle. Van Nostrand Reinhold. 1972

Section : Examples, page 35

Problem number : 1

Date solved : Saturday, December 06, 2025 at 04:17:25 PM

CAS classification : [_quadrature]

0.098 (sec) 2.1.1.1 Solved using first_order_ode_dAlembert

Entering first
order ode
dAlembert solver

$$y = y' + \frac{y'^2}{2}$$

Let $p = y'$ the ode becomes

$$y = p + \frac{1}{2}p^2$$

Solving for y from the above results in

$$y = p + \frac{1}{2}p^2 \quad (1)$$

This has the form

$$y = xf(p) + g(p) \quad (*)$$

Where f, g are functions of $p = y'(x)$. The above ode is dAlembert ode which is now solved.

Taking derivative of (*) w.r.t. x gives

$$\begin{aligned} p &= f + (xf' + g')\frac{dp}{dx} \\ p - f &= (xf' + g')\frac{dp}{dx} \end{aligned} \quad (2)$$

Comparing the form $y = xf + g$ to (1A) shows that

$$\begin{aligned} f &= 0 \\ g &= p + \frac{1}{2}p^2 \end{aligned}$$

Hence (2) becomes

$$p = (1 + p)p'(x) \quad (2A)$$

The singular solution is found by setting $\frac{dp}{dx} = 0$ in the above which gives

$$p = 0$$

Solving the above for p results in

$$p_1 = 0$$

Substituting these in (1A) and keeping singular solution that verifies the ode gives

$$y = 0$$

The general solution is found when $\frac{dp}{dx} \neq 0$. From eq. (2A). This results in

$$p'(x) = \frac{p(x)}{1 + p(x)} \quad (3)$$

This ODE is now solved for $p(x)$. No inversion is needed.

Integrating gives

$$\begin{aligned} \int \frac{1+p}{p} dp &= dx \\ p + \ln(p) &= x + c_1 \end{aligned}$$

Singular solutions are found by solving

$$\frac{p}{1+p} = 0$$

for $p(x)$. This is because we had to divide by this in the above step. This gives the following singular solution(s), which also have to satisfy the given ODE.

$$p(x) = 0$$

Substituting the above solution for p in (2A) gives

$$y = e^{-\text{LambertW}(e^{x+c_1})+x+c_1} + \frac{e^{-2\text{LambertW}(e^{x+c_1})+2x+2c_1}}{2}$$

$$y = 0$$

Simplifying the above gives

$$y = 0$$

$$y = \frac{\text{LambertW}(e^{x+c_1})(2 + \text{LambertW}(e^{x+c_1}))}{2}$$

$$y = 0$$

Summary of solutions found

$$y = 0$$

$$y = \frac{\text{LambertW}(e^{x+c_1})(2 + \text{LambertW}(e^{x+c_1}))}{2}$$

0.178 (sec) **2.1.1.2 Solved using first_order_ode_parametric method**

Entering first
order ode
parametric solver

$$y = y' + \frac{y'^2}{2}$$

Let y' be a parameter λ . The ode becomes

$$y - \lambda - \frac{1}{2}\lambda^2 = 0$$

Isolating y gives

$$\begin{aligned} y &= \lambda + \frac{1}{2}\lambda^2 \\ &= \lambda + \frac{1}{2}\lambda^2 \\ &= F(x, \lambda) \end{aligned}$$

Now we generate an ode in $x(\lambda)$ using

$$\begin{aligned} \frac{d}{d\lambda}x(\lambda) &= \frac{\frac{\partial F}{\partial \lambda}}{\lambda - \frac{\partial F}{\partial x}} \\ &= \frac{1 + \lambda}{\lambda} \\ &= \frac{1 + \lambda}{\lambda} \end{aligned}$$

Which is now solved for x .

Entering first
order ode
quadrature solver

Since the ode has the form $\frac{d}{d\lambda}x(\lambda) = f(\lambda)$, then we only need to integrate $f(\lambda)$.

$$\int dx = \int \frac{1+\lambda}{\lambda} d\lambda$$

$$x(\lambda) = \lambda + \ln(\lambda) + c_1$$

Now that we found solution x we have two equations with parameter λ . They are

$$y = \lambda + \frac{1}{2}\lambda^2$$

$$x = \lambda + \ln(\lambda) + c_1$$

Eliminating λ gives the solution for y . Simplifying the above gives

$$y = \frac{\text{LambertW}(e^{x-c_1})(\text{LambertW}(e^{x-c_1}) + 2)}{2}$$

$$y = \frac{\text{LambertW}(e^{x-c_1})(\text{LambertW}(e^{x-c_1}) + 2)}{2}$$

Summary of solutions found

$$y = \frac{\text{LambertW}(e^{x-c_1})(\text{LambertW}(e^{x-c_1}) + 2)}{2}$$

2.1.1.3 ✓ **Maple.** Time used: 0.014 (sec). Leaf size: 106

```
ode:=y(x) = diff(y(x),x)+1/2*diff(y(x),x)^2;
dsolve(ode,y(x), singsol=all);
```

$$y = \frac{\text{LambertW}(-\sqrt{2}e^{-1+x-c_1})(\text{LambertW}(-\sqrt{2}e^{-1+x-c_1}) + 2)}{2}$$

$$y = \frac{e^{2\text{RootOf}(-Z-2x+2e^{-Z}-2+2c_1-\ln(2)+\ln(e^{-Z}(e^{-Z}-2)^2))}}{2}$$

$$- e^{\text{RootOf}(-Z-2x+2e^{-Z}-2+2c_1-\ln(2)+\ln(e^{-Z}(e^{-Z}-2)^2))}$$

Maple trace

Methods for first order ODEs:

```
-> Solving 1st order ODE of high degree, 1st attempt
trying 1st order WeierstrassP solution for high degree ODE
trying 1st order WeierstrassPPrime solution for high degree ODE
```

```

trying 1st order JacobiSN solution for high degree ODE
trying 1st order ODE linearizable_by_differentiation
trying differential order: 1; missing variables
<- differential order: 1; missing x successful

```

Maple step by step

Let's solve

$$y(x) = \frac{d}{dx}y(x) + \frac{\left(\frac{d}{dx}y(x)\right)^2}{2}$$

- Highest derivative means the order of the ODE is 1

$$\frac{d}{dx}y(x)$$

- Solve for the highest derivative

$$\left[\frac{d}{dx}y(x) = -1 - \sqrt{1 + 2y(x)}, \frac{d}{dx}y(x) = -1 + \sqrt{1 + 2y(x)} \right]$$

- Solve the equation $\frac{d}{dx}y(x) = -1 - \sqrt{1 + 2y(x)}$

- Separate variables

$$\frac{\frac{d}{dx}y(x)}{-1 - \sqrt{1 + 2y(x)}} = 1$$

- Integrate both sides with respect to x

$$\int \frac{\frac{d}{dx}y(x)}{-1 - \sqrt{1 + 2y(x)}} dx = \int 1 dx + C_1$$

- Evaluate integral

$$\frac{\ln(y(x))}{2} - \sqrt{1 + 2y(x)} - \frac{\ln(-1 + \sqrt{1 + 2y(x)})}{2} + \frac{\ln(1 + \sqrt{1 + 2y(x)})}{2} = x + C_1$$

- Solve the equation $\frac{d}{dx}y(x) = -1 + \sqrt{1 + 2y(x)}$

- Separate variables

$$\frac{\frac{d}{dx}y(x)}{-1 + \sqrt{1 + 2y(x)}} = 1$$

- Integrate both sides with respect to x

$$\int \frac{\frac{d}{dx}y(x)}{-1 + \sqrt{1 + 2y(x)}} dx = \int 1 dx + C_1$$

- Evaluate integral

$$\frac{\ln(y(x))}{2} + \sqrt{1 + 2y(x)} + \frac{\ln(-1 + \sqrt{1 + 2y(x)})}{2} - \frac{\ln(1 + \sqrt{1 + 2y(x)})}{2} = x + C_1$$

- Set of solutions

$$\left\{ \frac{\ln(y(x))}{2} - \sqrt{1 + 2y(x)} - \frac{\ln(-1 + \sqrt{1 + 2y(x)})}{2} + \frac{\ln(1 + \sqrt{1 + 2y(x)})}{2} = x + C_1, \frac{\ln(y(x))}{2} + \sqrt{1 + 2y(x)} + \frac{\ln(-1 + \sqrt{1 + 2y(x)})}{2} - \frac{\ln(1 + \sqrt{1 + 2y(x)})}{2} = x + C_1 \right\}$$

2.1.1.4 ✓ **Mathematica.** Time used: 11.291 (sec). Leaf size: 66

```
ode=y[x]==D[y[x],x]+1/2*(D[y[x],x])^2;
ic={};
DSolve[{ode,ic},y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow \frac{1}{2} W(-e^{x-1-c_1}) (2 + W(-e^{x-1-c_1}))$$

$$y(x) \rightarrow \frac{1}{2} W(e^{x-1+c_1}) (2 + W(e^{x-1+c_1}))$$

$$y(x) \rightarrow 0$$

2.1.1.5 ✓ **Sympy.** Time used: 0.540 (sec). Leaf size: 51

```
from sympy import *
x = symbols("x")
y = Function("y")
ode = Eq(y(x) - Derivative(y(x), x)**2/2 - Derivative(y(x), x), 0)
ics = {}
dsolve(ode, func=y(x), ics=ics)
```

$$\left[x + \sqrt{2y(x) + 1} - \log(\sqrt{2y(x) + 1} + 1) = C_1, \quad x \right. \\ \left. - \sqrt{2y(x) + 1} - \log(\sqrt{2y(x) + 1} - 1) = C_1 \right]$$

2.1.2 Problem 2

Local contents

2.1.2.1	Solved using first_order_ode_clairaut	13
2.1.2.2	✓ Maple	16
2.1.2.3	✓ Mathematica	17
2.1.2.4	✗ Sympy	18

Internal problem ID [4087]

Book : Applied Differential equations, Newby Curle. Van Nostrand Reinhold. 1972

Section : Examples, page 35

Problem number : 2

Date solved : Saturday, December 06, 2025 at 04:17:26 PM

CAS classification :

[[_1st_order, _with_linear_symmetries], _rational, _Clairaut]

0.088 (sec) **2.1.2.1 Solved using first_order_ode_clairaut**

Entering first
order ode clairaut
solver

$$(y - xy')^2 = 1 + y'^2$$

This is Clairaut ODE. It has the form

$$y = xy' + g(y')$$

Where g is function of $y'(x)$. Let $p = y'$ the ode becomes

$$(-xp + y)^2 = p^2 + 1$$

Solving for y from the above results in

$$y = xp + \sqrt{p^2 + 1} \tag{1A}$$

$$y = xp - \sqrt{p^2 + 1} \tag{2A}$$

Each of the above ode's is a Clairaut ode which is now solved.

Solving ode 1A Writing the equation (1A) as

$$y = xp + g(p)$$

We now write $g \equiv g(p)$ to make notation simpler but we should always remember that g is function of p which in turn is function of x . Hence the above becomes

$$y = xp + g \tag{1}$$

Where

$$g = \sqrt{p^2 + 1}$$

Taking derivative of (1) w.r.t. x gives

$$\begin{aligned} p &= \frac{d}{dx}(xp + g) \\ p &= \left(p + x \frac{dp}{dx}\right) + \left(g' \frac{dp}{dx}\right) \\ p &= p + (x + g') \frac{dp}{dx} \\ 0 &= (x + g') \frac{dp}{dx} \end{aligned}$$

Where g' is derivative of $g(p)$ w.r.t. p .

The general solution is given by

$$\begin{aligned} \frac{dp}{dx} &= 0 \\ p &= c_1 \end{aligned}$$

Substituting this in (1) gives the general solution as

$$y = c_1 x + \sqrt{c_1^2 + 1}$$

The singular solution is found from solving for p from

$$x + g'(p) = 0$$

And substituting the result back in (1). Since we found above that $g = \sqrt{p^2 + 1}$, then the above equation becomes

$$\begin{aligned} x + g'(p) &= x + \frac{p}{\sqrt{p^2 + 1}} \\ &= 0 \end{aligned}$$

Solving the above for p results in

$$p_1 = y = (-x^2 + 1) \sqrt{-\frac{1}{x^2 - 1}}$$

Substituting the above back in (1) results in

$$y = (-x^2 + 1) \sqrt{-\frac{1}{x^2 - 1}}$$

Solving ode 2A Writing the equation (1A) as

$$y = xp + g(p)$$

We now write $g \equiv g(p)$ to make notation simpler but we should always remember that g is function of p which in turn is function of x . Hence the above becomes

$$y = xp + g \tag{1}$$

Where

$$g = -\sqrt{p^2 + 1}$$

Taking derivative of (1) w.r.t. x gives

$$\begin{aligned} p &= \frac{d}{dx}(xp + g) \\ p &= \left(p + x \frac{dp}{dx}\right) + \left(g' \frac{dp}{dx}\right) \\ p &= p + (x + g') \frac{dp}{dx} \\ 0 &= (x + g') \frac{dp}{dx} \end{aligned}$$

Where g' is derivative of $g(p)$ w.r.t. p .

The general solution is given by

$$\begin{aligned} \frac{dp}{dx} &= 0 \\ p &= c_1 \end{aligned}$$

Substituting this in (1) gives the general solution as

$$y = c_2x - \sqrt{c_2^2 + 1}$$

The singular solution is found from solving for p from

$$x + g'(p) = 0$$

And substituting the result back in (1). Since we found above that $g = -\sqrt{p^2 + 1}$, then the above equation becomes

$$\begin{aligned} x + g'(p) &= x - \frac{p}{\sqrt{p^2 + 1}} \\ &= 0 \end{aligned}$$

Solving the above for p results in

$$p_1 = y = \sqrt{-\frac{1}{x^2 - 1}} (x^2 - 1)$$

Substituting the above back in (1) results in

$$y = \sqrt{-\frac{1}{x^2 - 1}} (x^2 - 1)$$

Summary of solutions found

$$y = \sqrt{-\frac{1}{x^2 - 1}} (x^2 - 1)$$

$$y = (-x^2 + 1) \sqrt{-\frac{1}{x^2 - 1}}$$

$$y = c_1 x + \sqrt{c_1^2 + 1}$$

$$y = c_2 x - \sqrt{c_2^2 + 1}$$

2.1.2.2 ✓ Maple. Time used: 0.042 (sec). Leaf size: 57

```
ode:=(-diff(y(x),x)*x+y(x))^2 = 1+diff(y(x),x)^2;
dsolve(ode,y(x), singsol=all);
```

$$y = \sqrt{-x^2 + 1}$$

$$y = -\sqrt{-x^2 + 1}$$

$$y = c_1 x - \sqrt{c_1^2 + 1}$$

$$y = c_1 x + \sqrt{c_1^2 + 1}$$

Maple trace

Methods for first order ODEs:

*** Sublevel 2 ***

Methods for first order ODEs:

-> Solving 1st order ODE of high degree, 1st attempt

trying 1st order WeierstrassP solution for high degree ODE

trying 1st order WeierstrassPPrime solution for high degree ODE

trying 1st order JacobiSN solution for high degree ODE

```

trying 1st order ODE linearizable_by_differentiation
trying differential order: 1; missing variables
trying dAlembert
<- dAlembert successful
<- dAlembert successful

```

Maple step by step

Let's solve

$$\left(y(x) - x\left(\frac{d}{dx}y(x)\right)\right)^2 = 1 + \left(\frac{d}{dx}y(x)\right)^2$$

- Highest derivative means the order of the ODE is 1

$$\frac{d}{dx}y(x)$$

- Solve for the highest derivative

$$\left[\frac{d}{dx}y(x) = \frac{y(x)x - \sqrt{y(x)^2 + x^2 - 1}}{x^2 - 1}, \frac{d}{dx}y(x) = \frac{y(x)x + \sqrt{y(x)^2 + x^2 - 1}}{x^2 - 1}\right]$$

- Solve the equation $\frac{d}{dx}y(x) = \frac{y(x)x - \sqrt{y(x)^2 + x^2 - 1}}{x^2 - 1}$

- Solve the equation $\frac{d}{dx}y(x) = \frac{y(x)x + \sqrt{y(x)^2 + x^2 - 1}}{x^2 - 1}$

- Set of solutions

$$\{workingODE, workingODE\}$$

2.1.2.3 ✓ Mathematica. Time used: 0.076 (sec). Leaf size: 73

```

ode=(y[x]-x*D[y[x],x])^2==1+(D[y[x],x])^2;
ic={};
DSolve[{ode,ic},y[x],x,IncludeSingularSolutions->True]

```

$$y(x) \rightarrow c_1 x - \sqrt{1 + c_1^2}$$

$$y(x) \rightarrow c_1 x + \sqrt{1 + c_1^2}$$

$$y(x) \rightarrow -\sqrt{1 - x^2}$$

$$y(x) \rightarrow \sqrt{1 - x^2}$$

2.1.2.4 Sympy

```
from sympy import *
x = symbols("x")
y = Function("y")
ode = Eq((-x*Derivative(y(x), x) + y(x))**2 - Derivative(y(x), x)**2 - 1,0)
ics = {}
dsolve(ode,func=y(x),ics=ics)
```

Timed Out

2.1.3 Problem 3

Local contents

2.1.3.1	Solved using first_order_ode_dAlembert	19
2.1.3.2	Solved using first_order_ode_parametric method	21
2.1.3.3	✓ Maple	22
2.1.3.4	✓ Mathematica	23
2.1.3.5	✗ Sympy	23

Internal problem ID [4088]

Book : Applied Differential equations, Newby Curle. Van Nostrand Reinhold. 1972

Section : Examples, page 35

Problem number : 3

Date solved : Saturday, December 06, 2025 at 04:17:27 PM

CAS classification : [[_homogeneous, 'class C'], _dAlembert]

0.056 (sec) 2.1.3.1 Solved using first_order_ode_dAlembert

Entering first
order ode
dAlembert solver

$$y - x = y'^2 \left(1 - \frac{2y'}{3} \right)$$

Let $p = y'$ the ode becomes

$$y - x = p^2 \left(1 - \frac{2p}{3} \right)$$

Solving for y from the above results in

$$y = p^2 - \frac{2}{3}p^3 + x \quad (1)$$

This has the form

$$y = xf(p) + g(p) \quad (*)$$

Where f, g are functions of $p = y'(x)$. The above ode is dAlembert ode which is now solved.

Taking derivative of (*) w.r.t. x gives

$$\begin{aligned} p &= f + (xf' + g') \frac{dp}{dx} \\ p - f &= (xf' + g') \frac{dp}{dx} \end{aligned} \quad (2)$$

Comparing the form $y = xf + g$ to (1A) shows that

$$\begin{aligned} f &= 1 \\ g &= p^2 - \frac{2}{3}p^3 \end{aligned}$$

Hence (2) becomes

$$p - 1 = (-2p^2 + 2p) p'(x) \quad (2A)$$

The singular solution is found by setting $\frac{dp}{dx} = 0$ in the above which gives

$$p - 1 = 0$$

Solving the above for p results in

$$p_1 = 1$$

Substituting these in (1A) and keeping singular solution that verifies the ode gives

$$y = x + \frac{1}{3}$$

The general solution is found when $\frac{dp}{dx} \neq 0$. From eq. (2A). This results in

$$p'(x) = \frac{p(x) - 1}{-2p(x)^2 + 2p(x)} \quad (3)$$

This ODE is now solved for $p(x)$. No inversion is needed.

Integrating gives

$$\begin{aligned} \int -2p dp &= dx \\ -p^2 &= x + c_1 \end{aligned}$$

Substituting the above solution for p in (2A) gives

$$y = -c_1 - \frac{2(-x - c_1)^{3/2}}{3}$$

Simplifying the above gives

$$\begin{aligned} y &= x + \frac{1}{3} \\ y &= \frac{(2x + 2c_1) \sqrt{-x - c_1}}{3} - c_1 \end{aligned}$$

Summary of solutions found

$$y = x + \frac{1}{3}$$

$$y = \frac{(2x + 2c_1)\sqrt{-x - c_1}}{3} - c_1$$

0.086 (sec) **2.1.3.2 Solved using first_order_ode_parametric method**

Entering first
order ode
parametric solver

$$y - x = y'^2 \left(1 - \frac{2y'}{3} \right)$$

Let y' be a parameter λ . The ode becomes

$$y - x - \lambda^2 \left(1 - \frac{2\lambda}{3} \right) = 0$$

Isolating y gives

$$\begin{aligned} y &= x + \lambda^2 - \frac{2}{3}\lambda^3 \\ &= x + \lambda^2 - \frac{2}{3}\lambda^3 \\ &= F(x, \lambda) \end{aligned}$$

Now we generate an ode in $x(\lambda)$ using

$$\begin{aligned} \frac{d}{d\lambda} x(\lambda) &= \frac{\frac{\partial F}{\partial \lambda}}{\lambda - \frac{\partial F}{\partial x}} \\ &= \frac{-2\lambda^2 + 2\lambda}{\lambda - 1} \\ &= -2\lambda \end{aligned}$$

Which is now solved for x .

Entering first
order ode
quadrature solver

Since the ode has the form $\frac{d}{d\lambda} x(\lambda) = f(\lambda)$, then we only need to integrate $f(\lambda)$.

$$\begin{aligned} \int dx &= \int -2\lambda d\lambda \\ x(\lambda) &= -\lambda^2 + c_1 \end{aligned}$$

Now that we found solution x we have two equations with parameter λ . They are

$$\begin{aligned} y &= x + \lambda^2 - \frac{2}{3}\lambda^3 \\ x &= -\lambda^2 + c_1 \end{aligned}$$

Eliminating λ gives the solution for y . Simplifying the above gives

$$y = c_1 - \frac{2\sqrt{(-x + c_1)^3}}{3}$$

$$y = c_1 + \frac{2\sqrt{(-x + c_1)^3}}{3}$$

Summary of solutions found

$$y = c_1 - \frac{2\sqrt{(-x + c_1)^3}}{3}$$

$$y = c_1 + \frac{2\sqrt{(-x + c_1)^3}}{3}$$

2.1.3.3 ✓ Maple. Time used: 0.024 (sec). Leaf size: 49

```
ode:=-x+y(x) = diff(y(x),x)^2*(1-2/3*diff(y(x),x));
dsolve(ode,y(x), singsol=all);
```

$$y = x + \frac{1}{3}$$

$$y = \frac{(2x - 2c_1)\sqrt{c_1 - x}}{3} + c_1$$

$$y = \frac{(-2x + 2c_1)\sqrt{c_1 - x}}{3} + c_1$$

Maple trace

Methods for first order ODEs:

*** Sublevel 2 ***

Methods for first order ODEs:

-> Solving 1st order ODE of high degree, 1st attempt

trying 1st order WeierstrassP solution for high degree ODE

trying 1st order WeierstrassPPrime solution for high degree ODE

trying 1st order JacobiSN solution for high degree ODE

trying 1st order ODE linearizable_by_differentiation

trying differential order: 1; missing variables

trying dAlembert

<- dAlembert successful

Maple step by step

Let's solve

$$y(x) - x = \left(\frac{d}{dx}y(x)\right)^2 \left(1 - \frac{2\frac{d}{dx}y(x)}{3}\right)$$

- Highest derivative means the order of the ODE is 1

$$\frac{d}{dx}y(x)$$

- Solve for the highest derivative

$$\left[\frac{d}{dx}y(x) = \frac{\left(1-6y(x)+6x+2\sqrt{-3y(x)+3x+9y(x)^2-18y(x)x+9x^2}\right)^{1/3}}{2} + \frac{1}{2\left(1-6y(x)+6x+2\sqrt{-3y(x)+3x+9y(x)^2-18y(x)x+9x^2}\right)}$$

- Solve the equation $\frac{d}{dx}y(x) = \frac{\left(1-6y(x)+6x+2\sqrt{-3y(x)+3x+9y(x)^2-18y(x)x+9x^2}\right)^{1/3}}{2} + \frac{1}{2\left(1-6y(x)+6x+2\sqrt{-3y(x)+3x+9y(x)^2-18y(x)x+9x^2}\right)}$

- Solve the equation $\frac{d}{dx}y(x) = -\frac{\left(1-6y(x)+6x+2\sqrt{-3y(x)+3x+9y(x)^2-18y(x)x+9x^2}\right)^{1/3}}{4} - \frac{1}{4\left(1-6y(x)+6x+2\sqrt{-3y(x)+3x+9y(x)^2-18y(x)x+9x^2}\right)}$

- Solve the equation $\frac{d}{dx}y(x) = -\frac{\left(1-6y(x)+6x+2\sqrt{-3y(x)+3x+9y(x)^2-18y(x)x+9x^2}\right)^{1/3}}{4} - \frac{1}{4\left(1-6y(x)+6x+2\sqrt{-3y(x)+3x+9y(x)^2-18y(x)x+9x^2}\right)}$

- Set of solutions

$$\{workingODE, workingODE, workingODE\}$$

2.1.3.4 ✓ Mathematica. Time used: 160.121 (sec). Leaf size: 14234

```
ode=y[x]-x==D[y[x],x]^2*(1-2/3*D[y[x],x]);
ic={};
DSolve[{ode,ic},y[x],x,IncludeSingularSolutions->True]
```

Too large to display

2.1.3.5 ✗ Sympy

```
from sympy import *
x = symbols("x")
y = Function("y")
ode = Eq(-x - (1 - 2*Derivative(y(x), x)/3)*Derivative(y(x), x)**2 + y(x), 0)
ics = {}
dsolve(ode, func=y(x), ics=ics)
```

Timed Out

2.1.4 Problem 4

Local contents

2.1.4.1	Solved using first_order_ode_homog_type_maple_C	24
2.1.4.2	Solved using first_order_ode_LIE	28
2.1.4.3	Solved using first_order_ode_riccati	33
2.1.4.4	Solved using first_order_ode_riccati_by_guessing_particular_solution	40
2.1.4.5	✓ Maple	42
2.1.4.6	✓ Mathematica	43
2.1.4.7	✓ Sympy	43

Internal problem ID [4089]

Book : Applied Differential equations, Newby Curle. Van Nostrand Reinhold. 1972

Section : Examples, page 35

Problem number : 4

Date solved : Saturday, December 06, 2025 at 04:17:28 PM

CAS classification : [[_homogeneous, 'class C'], _rational, _Riccati]

0.235 (sec) 2.1.4.1 Solved using first_order_ode_homog_type_maple_C

Entering first
order ode homog
type maple C
solver

$$y'x^2 = x(y-1) + (y-1)^2$$

Let $Y = y - y_0$ and $X = x - x_0$ then the above is transformed to new ode in $Y(X)$

$$\frac{d}{dX}Y(X) = \frac{(Y(X) + y_0 - 1)(Y(X) + y_0 + X + x_0 - 1)}{(X + x_0)^2}$$

Solving for possible values of x_0 and y_0 which makes the above ode a homogeneous ode results in

$$x_0 = 0$$

$$y_0 = 1$$

Using these values now it is possible to easily solve for $Y(X)$. The above ode now becomes

$$\frac{d}{dX}Y(X) = \frac{Y(X)X + Y(X)^2}{X^2}$$

In canonical form, the ODE is

$$\begin{aligned} Y' &= F(X, Y) \\ &= \frac{Y(X + Y)}{X^2} \end{aligned} \quad (1)$$

An ode of the form $Y' = \frac{M(X, Y)}{N(X, Y)}$ is called homogeneous if the functions $M(X, Y)$ and $N(X, Y)$ are both homogeneous functions and of the same order. Recall that a function $f(X, Y)$ is homogeneous of order n if

$$f(t^n X, t^n Y) = t^n f(X, Y)$$

In this case, it can be seen that both $M = Y(X + Y)$ and $N = X^2$ are both homogeneous and of the same order $n = 2$. Therefore this is a homogeneous ode. Since this ode is homogeneous, it is converted to separable ODE using the substitution $u = \frac{Y}{X}$, or $Y = uX$. Hence

$$\frac{dY}{dX} = \frac{du}{dX}X + u$$

Applying the transformation $Y = uX$ to the above ODE in (1) gives

$$\begin{aligned} \frac{du}{dX}X + u &= u^2 + u \\ \frac{du}{dX} &= \frac{u(X)^2}{X} \end{aligned}$$

Or

$$\frac{d}{dX}u(X) - \frac{u(X)^2}{X} = 0$$

Or

$$-u(X)^2 + \left(\frac{d}{dX}u(X) \right) X = 0$$

Which is now solved as separable in $u(X)$.

The ode

$$\frac{d}{dX}u(X) = \frac{u(X)^2}{X} \quad (2.1)$$

is separable as it can be written as

$$\begin{aligned} \frac{d}{dX}u(X) &= \frac{u(X)^2}{X} \\ &= f(X)g(u) \end{aligned}$$

Where

$$\begin{aligned} f(X) &= \frac{1}{X} \\ g(u) &= u^2 \end{aligned}$$

Integrating gives

$$\int \frac{1}{g(u)} du = \int f(X) dX$$

$$\int \frac{1}{u^2} du = \int \frac{1}{X} dX$$

$$-\frac{1}{u(X)} = \ln(X) + c_1$$

We now need to find the singular solutions, these are found by finding for what values $g(u)$ is zero, since we had to divide by this above. Solving $g(u) = 0$ or

$$u^2 = 0$$

for $u(X)$ gives

$$u(X) = 0$$

Now we go over each such singular solution and check if it verifies the ode itself and any initial conditions given. If it does not then the singular solution will not be used.

Therefore the solutions found are

$$-\frac{1}{u(X)} = \ln(X) + c_1$$

$$u(X) = 0$$

Converting $-\frac{1}{u(X)} = \ln(X) + c_1$ back to $Y(X)$ gives

$$-\frac{X}{Y(X)} = \ln(X) + c_1$$

Converting $u(X) = 0$ back to $Y(X)$ gives

$$Y(X) = 0$$

Using the solution for $Y(X)$

$$-\frac{X}{Y(X)} = \ln(X) + c_1 \tag{A}$$

And replacing back terms in the above solution using

$$\begin{aligned} Y &= y + y_0 \\ X &= x_0 + x \end{aligned}$$

Or

$$\begin{aligned} Y &= y + 1 \\ X &= x \end{aligned}$$

Then the solution in y becomes using EQ (A)

$$-\frac{x}{y-1} = \ln(x) + c_1$$

Using the solution for $Y(X)$

$$Y(X) = 0 \tag{A}$$

And replacing back terms in the above solution using

$$\begin{aligned} Y &= y + y_0 \\ X &= x_0 + x \end{aligned}$$

Or


$$\begin{aligned} Y &= y + 1 \\ X &= x \end{aligned}$$

Then the solution in y becomes using EQ (A)

$$y - 1 = 0$$

Solving for y gives

$$\begin{aligned} y &= 1 \\ y &= \frac{\ln(x) + c_1 - x}{\ln(x) + c_1} \end{aligned}$$

Figure 2.1: Slope field $y'x^2 = x(y-1) + (y-1)^2$ Summary of solutions found

$$y = 1$$

$$y = \frac{\ln(x) + c_1 - x}{\ln(x) + c_1}$$

0.911 (sec) **2.1.4.2 Solved using first_order_ode_LIE**

Entering first
order ode LIE
solver

$$y'x^2 = x(y-1) + (y-1)^2$$

Writing the ode as

$$y' = \frac{yx + y^2 - x - 2y + 1}{x^2}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \quad (1\text{E})$$

$$\eta = xb_2 + yb_3 + b_1 \quad (2\text{E})$$

Equation (7E) now becomes

$$\begin{aligned} & -2b_2v_1^3v_2 + (-b_1 + 2b_2 - b_3)v_1^3 + (-b_3 + a_2)v_1^2v_2^2 + (a_1 - 2a_2 + a_3 - 2b_1)v_1^2v_2 \\ & + (-a_1 + a_2 - a_3 + 2b_1 + b_3)v_1^2 + (2a_1 + 2a_3)v_1v_2^2 + (-4a_1 - 4a_3)v_1v_2 \\ & + (2a_1 + 2a_3)v_1 - a_3v_2^4 + 4a_3v_2^3 - 6a_3v_2^2 + 4a_3v_2 - a_3 = 0 \end{aligned} \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} -6a_3 &= 0 \\ -a_3 &= 0 \\ 4a_3 &= 0 \\ -2b_2 &= 0 \\ -4a_1 - 4a_3 &= 0 \\ 2a_1 + 2a_3 &= 0 \\ -b_3 + a_2 &= 0 \\ -b_1 + 2b_2 - b_3 &= 0 \\ a_1 - 2a_2 + a_3 - 2b_1 &= 0 \\ -a_1 + a_2 - a_3 + 2b_1 + b_3 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= 0 \\ a_2 &= b_3 \\ a_3 &= 0 \\ b_1 &= -b_3 \\ b_2 &= 0 \\ b_3 &= b_3 \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned} \xi &= x \\ \eta &= y - 1 \end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned} \eta &= \eta - \omega(x, y) \xi \\ &= y - 1 - \left(\frac{yx + y^2 - x - 2y + 1}{x^2} \right) (x) \\ &= -\frac{(y-1)^2}{x} \\ \xi &= 0 \end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{-\frac{(y-1)^2}{x}} dy \end{aligned}$$

Which results in

$$S = \frac{x}{y-1}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{yx + y^2 - x - 2y + 1}{x^2}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{1}{y-1} \\ S_y &= -\frac{x}{(y-1)^2} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = -\frac{1}{R} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = -\frac{1}{R}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S .



Since the ode has the form $\frac{d}{dR}S(R) = f(R)$, then we only need to integrate $f(R)$.

$$\begin{aligned} \int dS &= \int -\frac{1}{R} dR \\ S(R) &= -\ln(R) + c_2 \end{aligned}$$

To complete the solution, we just need to transform the above back to x, y coordinates. This results in

$$\frac{x}{y-1} = -\ln(x) + c_2$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{yx+y^2-x-2y+1}{x^2}$ 	$\begin{aligned} R &= x \\ S &= \frac{x}{y-1} \end{aligned}$	$\frac{dS}{dR} = -\frac{1}{R}$ 

Solving for y gives

$$y = \frac{\ln(x) - c_2 - x}{\ln(x) - c_2}$$




Figure 2.2: Slope field $y'x^2 = x(y-1) + (y-1)^2$

Summary of solutions found

$$y = \frac{\ln(x) - c_2 - x}{\ln(x) - c_2}$$

0.273 (sec) **2.1.4.3 Solved using first_order_ode_riccati**

Entering first
order ode riccati
solver

$$y'x^2 = x(y-1) + (y-1)^2$$

In canonical form the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= \frac{yx + y^2 - x - 2y + 1}{x^2} \end{aligned}$$

This is a Riccati ODE. Comparing the ODE to solve

$$y' = \frac{y}{x} + \frac{y^2}{x^2} - \frac{1}{x} - \frac{2y}{x^2} + \frac{1}{x^2}$$

With Riccati ODE standard form

$$y' = f_0(x) + f_1(x)y + f_2(x)y^2$$

Shows that $f_0(x) = \frac{1-x}{x^2}$, $f_1(x) = \frac{x-2}{x^2}$ and $f_2(x) = \frac{1}{x^2}$. Let

$$\begin{aligned} y &= \frac{-u'}{f_2 u} \\ &= \frac{-u'}{\frac{u}{x^2}} \end{aligned} \quad (1)$$

Using the above substitution in the given ODE results (after some simplification) in a second order ODE to solve for $u(x)$ which is

$$f_2 u''(x) - (f_2' + f_1 f_2) u'(x) + f_2^2 f_0 u(x) = 0 \quad (2)$$

But

$$\begin{aligned} f_2' &= -\frac{2}{x^3} \\ f_1 f_2 &= \frac{x-2}{x^4} \\ f_2^2 f_0 &= \frac{1-x}{x^6} \end{aligned}$$

Substituting the above terms back in equation (2) gives

$$\frac{u''(x)}{x^2} - \left(-\frac{2}{x^3} + \frac{x-2}{x^4} \right) u'(x) + \frac{(1-x)u(x)}{x^6} = 0$$

Entering kovacic
solver

Writing the ode as

$$\frac{d^2 u}{dx^2} + \frac{(2+x) \left(\frac{du}{dx} \right)}{x^4} + \frac{(1-x)u}{x^6} = 0 \quad (1)$$

$$A \frac{d^2 u}{dx^2} + B \frac{du}{dx} + C u = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= \frac{1}{x^2} \\ B &= \frac{2+x}{x^4} \\ C &= \frac{1-x}{x^6} \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = u e^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{4x^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 4x^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(-\frac{1}{4x^2}\right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then u is found using the inverse transformation

$$u = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 2.5: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = -\frac{1}{4x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = -\frac{1}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{1}{2} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{2} \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = -\frac{1}{4x^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = -\frac{1}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{1}{2} \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{2} \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = -\frac{1}{4x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	$\frac{1}{2}$	$\frac{1}{2}$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	$\frac{1}{2}$	$\frac{1}{2}$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = \frac{1}{2}$ then

$$\begin{aligned} d &= \alpha_\infty^- - (\alpha_{c_1}^-) \\ &= \frac{1}{2} - \left(\frac{1}{2}\right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c) [\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty) [\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= \left((-) [\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{x - c_1} \right) + (-) [\sqrt{r}]_\infty \\ &= \frac{1}{2x} + (-) (0) \\ &= \frac{1}{2x} \\ &= \frac{1}{2x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2\left(\frac{1}{2x}\right)(0) + \left(\left(-\frac{1}{2x^2}\right) + \left(\frac{1}{2x}\right)^2 - \left(-\frac{1}{4x^2}\right)\right) &= 0 \\ 0 &= 0 \end{aligned}$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= pe^{\int \omega dx} \\ &= e^{\int \frac{1}{2x} dx} \\ &= \sqrt{x} \end{aligned}$$

The first solution to the original ode in u is found from

$$\begin{aligned} u_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{\frac{2+x}{x^4}}{\frac{1}{x^2}} dx} \\ &= z_1 e^{-\frac{\ln(x)}{2} + \frac{1}{x}} \\ &= z_1 \left(\frac{e^{\frac{1}{x}}}{\sqrt{x}} \right) \end{aligned}$$

Which simplifies to

$$u_1 = e^{\frac{1}{x}}$$

The second solution u_2 to the original ode is found using reduction of order

$$u_2 = u_1 \int \frac{e^{\int -\frac{B}{A} dx}}{u_1^2} dx$$

Substituting gives

$$\begin{aligned} u_2 &= u_1 \int \frac{e^{\int -\frac{\frac{2+x}{x^4}}{\frac{1}{x^2}} dx}}{(u_1)^2} dx \\ &= u_1 \int \frac{e^{-\ln(x) + \frac{2}{x}}}{(u_1)^2} dx \\ &= u_1 \left(-e^{-\ln(\frac{1}{x}) - \ln(x)} \ln\left(\frac{1}{x}\right) \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} u &= c_1 u_1 + c_2 u_2 \\ &= c_1 \left(e^{\frac{1}{x}} \right) + c_2 \left(e^{\frac{1}{x}} \left(-e^{-\ln(\frac{1}{x}) - \ln(x)} \ln \left(\frac{1}{x} \right) \right) \right) \end{aligned}$$

Taking derivative gives

$$u'(x) = -\frac{c_1 e^{\frac{1}{x}}}{x^2} + \frac{c_2 e^{\frac{1}{x}} \ln \left(\frac{1}{x} \right)}{x^2} + \frac{c_2 e^{\frac{1}{x}}}{x} \quad (4)$$

Substituting equations (3,4) into (1) results in


$$\begin{aligned} y &= \frac{-u'}{f_2 u} \\ y &= \frac{-u'}{\frac{u}{x^2}} \\ y &= \frac{-c_2 x - \ln \left(\frac{1}{x} \right) c_2 + c_1}{-\ln \left(\frac{1}{x} \right) c_2 + c_1} \end{aligned}$$

Doing change of constants, the above solution becomes

$$y = -\frac{\left(-\frac{e^{\frac{1}{x}}}{x^2} + \frac{c_3 e^{\frac{1}{x}} \ln \left(\frac{1}{x} \right)}{x^2} + \frac{c_3 e^{\frac{1}{x}}}{x} \right) x^2}{e^{\frac{1}{x}} - c_3 e^{\frac{1}{x}} \ln \left(\frac{1}{x} \right)}$$

Simplifying the above gives

$$y = \frac{\ln \left(\frac{1}{x} \right) c_3 + c_3 x - 1}{\ln \left(\frac{1}{x} \right) c_3 - 1}$$

Figure 2.3: Slope field $y'x^2 = x(y-1) + (y-1)^2$ Summary of solutions found

$$y = \frac{\ln\left(\frac{1}{x}\right) c_3 + c_3 x - 1}{\ln\left(\frac{1}{x}\right) c_3 - 1}$$

2.1.4.4 Solved using**first_order_ode_riccati_by_guessing_particular_solution**

0.320 (sec)

Entering first
order ode riccati
guess solver

$$y'x^2 = x(y-1) + (y-1)^2$$

This is a Riccati ODE. Comparing the above ODE to solve with the Riccati standard form

$$y' = f_0(x) + f_1(x)y + f_2(x)y^2$$

Shows that

$$\begin{aligned} f_0(x) &= \frac{1-x}{x^2} \\ f_1(x) &= \frac{x-2}{x^2} \\ f_2(x) &= \frac{1}{x^2} \end{aligned}$$

Using trial and error, the following particular solution was found

$$y_p = 1$$

Since a particular solution is known, then the general solution is given by

$$y = y_p + \frac{\phi(x)}{c_1 - \int \phi(x) f_2 dx}$$

Where

$$\phi(x) = e^{\int 2f_2 y_p + f_1 dx}$$

Evaluating the above gives the general solution as

$$y = 1 + \frac{x}{c_1 - \ln(x)}$$





Figure 2.4: Slope field $y'x^2 = x(y-1) + (y-1)^2$

Figure 2.5: Slope field $y'x^2 = x(y-1) + (y-1)^2$ Summary of solutions found

$$y = 1 + \frac{x}{c_1 - \ln(x)}$$

2.1.4.5 ✓ Maple. Time used: 0.005 (sec). Leaf size: 15

```
ode:=x^2*diff(y(x),x) = (-1+y(x))*x+(-1+y(x))^2;
dsolve(ode,y(x), singsol=all);
```

$$y = 1 - \frac{x}{\ln(x) + c_1}$$

Maple trace

```
Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying homogeneous C
trying homogeneous types:
trying homogeneous D
```

```
<- homogeneous successful
<- homogeneous successful
```

Maple step by step

Let's solve

$$x^2 \left(\frac{d}{dx} y(x) \right) = x(y(x) - 1) + (y(x) - 1)^2$$

- Highest derivative means the order of the ODE is 1

$$\frac{d}{dx} y(x)$$

- Solve for the highest derivative

$$\frac{d}{dx} y(x) = \frac{x(y(x)-1) + (y(x)-1)^2}{x^2}$$

2.1.4.6 ✓ **Mathematica.** Time used: 0.13 (sec). Leaf size: 23

```
ode=x^2*D[y[x],x]==x*(y[x]-1)+(y[x]-1)^2;
ic={};
DSolve[{ode,ic},y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow 1 + \frac{x}{-\log(x) + c_1}$$

$$y(x) \rightarrow 1$$

2.1.4.7 ✓ **Sympy.** Time used: 0.202 (sec). Leaf size: 8

```
from sympy import *
x = symbols("x")
y = Function("y")
ode = Eq(x**2*Derivative(y(x), x) - x*(y(x) - 1) - (y(x) - 1)**2, 0)
ics = {}
dsolve(ode, func=y(x), ics=ics)
```

$$y(x) = 1 - 8x^3$$