


A Solution Manual For

**Differential Equations: Theory,
Technique, and Practice by George
Simmons, Steven Krantz. McGraw-Hill
NY. 2007. 1st Edition.**



Nasser M. Abbasi

May 15, 2024

Contents

1	Chapter 1. What is a differential equation. Section 1.2 THE NATURE OF SOLUTIONS. Page 9	4
2	Chapter 1. What is a differential equation. Section 1.3 SEPARABLE EQUATIONS. Page 12	234
3	Chapter 1. What is a differential equation. Section 1.4 First Order Linear Equations. Page 15	317
4	Chapter 1. What is a differential equation. Section 1.5. Exact Equations. Page 20	506
5	Chapter 1. What is a differential equation. Section 1.7. Homogeneous Equations. Page 28	693
6	Chapter 1. What is a differential equation. Section 1.8. Integrating Factors. Page 32	851
7	Chapter 1. What is a differential equation. Section 1.9. Reduction of Order. Page 38	932
8	Chapter 1. What is a differential equation. Problems for Review and Discovery. Page 53	1030
9	Chapter 2. Second-Order Linear Equations. Section 2.1. Linear Equations with Constant Coefficients. Page 62	1281
10	Chapter 2. Second-Order Linear Equations. Section 2.2. THE METHOD OF UNDETERMINED COEFFICIENTS. Page 67	1638
11	Chapter 2. Second-Order Linear Equations. Section 2.3. THE METHOD OF VARIATION OF PARAMETERS. Page 71	1822
12	Chapter 2. Second-Order Linear Equations. Section 2.4. THE USE OF A KNOWN SOLUTION TO FIND ANOTHER. Page 74	2117
13	Chapter 2. Section 2.7. HIGHER ORDER LINEAR EQUATIONS, COUPLED HARMONIC OSCILLATORS Page 98	2168

14 Chapter 2. Problems for Review and Discovery. Drill excercises. Page 105	2295
15 Chapter 2. Problems for Review and Discovery. Challenge excercises. Page 105	2663
16 Chapter 2. Problems for Review and Discovery. Problems for Dis- cussion and Exploration. Page 105	2700
17 Chapter 4. Power Series Solutions and Special Functions. Section 4.2. Series Solutions of First-Order Differential Equations Page 1622731	
18 Chapter 4. Power Series Solutions and Special Functions. Section 4.3. Second-Order Linear Equations: Ordinary Points. Page 169	2961
19 Chapter 4. Power Series Solutions and Special Functions. Section 4.4. REGULAR SINGULAR POINTS. Page 175	3118
20 Chapter 4. Power Series Solutions and Special Functions. Section 4.5. More on Regular Singular Points. Page 183	3357
21 Chapter 4. Power Series Solutions and Special Functions. Section 4.6. Gauss's Hypergeometric Equation. Page 187	3488
22 Chapter 4. Power Series Solutions and Special Functions. Problems for review and discover. (A) Drill Exercises . Page 194	3592
23 Chapter 4. Power Series Solutions and Special Functions. Problems for review and discover. (B) Challenge Problems . Page 194	3806
24 Chapter 7. Laplace Transforms. Section 7.5 The Unit Step and Impulse Functions. Page 303	3853
25 Chapter 7. Laplace Transforms. Section 7.5 Problesm for review and discovery. Section A, Drill exercises. Page 309	3887
26 Chapter 7. Laplace Transforms. Section 7.5 Problesm for review and discovery. Section B, Challenge Problems. Page 310	3935
27 Chapter 10. Systems of First-Order Equations. Section 10.2 Linear Systems. Page 380	3944

28	Chapter 10. Systems of First-Order Equations. Section 10.3 Homogeneous Linear Systems with Constant Coefficients. Page 387	3999
29	Chapter 10. Systems of First-Order Equations. Section A. Drill exercises. Page 400	4086
30	Chapter 10. Systems of First-Order Equations. Section B. Challenge Problems. Page 401	4224

1 Chapter 1. What is a differential equation.

Section 1.2 THE NATURE OF SOLUTIONS.

Page 9

1.1	problem 1(a)	5
1.2	problem 1(b)	8
1.3	problem 1(c)	23
1.4	problem 1(d)	35
1.5	problem 1(e)	38
1.6	problem 1(f)	48
1.7	problem 1(h)	58
1.8	problem 1(i)	60
1.9	problem 1(j)	70
1.10	problem 1(k)	84
1.11	problem 1(L)	99
1.12	problem 1(m)	112
1.13	problem 1(n)	126
1.14	problem 1(o)	139
1.15	problem 2(a)	142
1.16	problem 2(b)	145
1.17	problem 2(c)	148
1.18	problem 2(d)	151
1.19	problem 2(e)	154
1.20	problem 2(f)	157
1.21	problem 2(g)	160
1.22	problem 2(h)	163
1.23	problem 2(i)	166
1.24	problem 2(j)	169
1.25	problem 3(a)	172
1.26	problem 3(b)	176
1.27	problem 3(c)	180
1.28	problem 3(d)	184
1.29	problem 3(e)	188
1.30	problem 3(f)	192
1.31	problem 4	196
1.32	problem 5	209
1.33	problem 6	217
1.34	problem 7	229

1.1 problem 1(a)

1.1.1 Solving as quadrature ode	5
1.1.2 Maple step by step solution	6

Internal problem ID [6105]

Internal file name [OUTPUT/5353_Sunday_June_05_2022_03_35_10_PM_5110884/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.2 THE NATURE OF SOLUTIONS. Page 9

Problem number: 1(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

`[_quadrature]`

$$y' = 2x$$

1.1.1 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned} y &= \int 2x \, dx \\ &= x^2 + c_1 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = x^2 + c_1 \tag{1}$$




Figure 1: Slope field plot

Verification of solutions

$$y = x^2 + c_1$$

Verified OK.

1.1.2 Maple step by step solution

Let's solve

$$y' = 2x$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Integrate both sides with respect to x

$$\int y' dx = \int 2x dx + c_1$$

- Evaluate integral

$$y = x^2 + c_1$$

- Solve for y

$$y = x^2 + c_1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
dsolve(diff(y(x),x)=2*x,y(x), singsol=all)
```

$$y(x) = x^2 + c_1$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 11

```
DSolve[y'[x]==2*x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^2 + c_1$$

1.2 problem 1(b)

1.2.1	Solving as separable ode	8
1.2.2	Solving as linear ode	10
1.2.3	Solving as homogeneousTypeD2 ode	11
1.2.4	Solving as first order ode lie symmetry lookup ode	13
1.2.5	Solving as exact ode	17
1.2.6	Maple step by step solution	21

Internal problem ID [6106]

Internal file name [OUTPUT/5354_Sunday_June_05_2022_03_35_11_PM_70934078/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.2 THE NATURE OF SOLUTIONS. Page 9

Problem number: 1(b).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "linear", "separable", "homogeneousTypeD2", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_separable]

$$xy' - 2y = 0$$

1.2.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{2y}{x}\end{aligned}$$

Where $f(x) = \frac{2}{x}$ and $g(y) = y$. Integrating both sides gives

$$\begin{aligned}\frac{1}{y} dy &= \frac{2}{x} dx \\ \int \frac{1}{y} dy &= \int \frac{2}{x} dx \\ \ln(y) &= 2 \ln(x) + c_1 \\ y &= e^{2 \ln(x) + c_1} \\ &= c_1 x^2\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 x^2 \tag{1}$$




Figure 2: Slope field plot

Verification of solutions

$$y = c_1 x^2$$

Verified OK.

1.2.2 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -\frac{2}{x}$$
$$q(x) = 0$$

Hence the ode is

$$y' - \frac{2y}{x} = 0$$

The integrating factor μ is

$$\mu = e^{\int -\frac{2}{x} dx}$$
$$= \frac{1}{x^2}$$

The ode becomes

$$\frac{d}{dx} \mu y = 0$$
$$\frac{d}{dx} \left(\frac{y}{x^2} \right) = 0$$

Integrating gives

$$\frac{y}{x^2} = c_1$$

Dividing both sides by the integrating factor $\mu = \frac{1}{x^2}$ results in

$$y = c_1 x^2$$

Summary

The solution(s) found are the following

$$y = c_1 x^2 \tag{1}$$




Figure 3: Slope field plot

Verification of solutions

$$y = c_1 x^2$$

Verified OK.

1.2.3 Solving as homogeneousTypeD2 ode

Using the change of variables $y = u(x)x$ on the above ode results in new ode in $u(x)$

$$x(u'(x)x + u(x)) - 2u(x)x = 0$$

In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= \frac{u}{x} \end{aligned}$$

Where $f(x) = \frac{1}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u} du &= \frac{1}{x} dx \\ \int \frac{1}{u} du &= \int \frac{1}{x} dx \\ \ln(u) &= \ln(x) + c_2 \\ u &= e^{\ln(x)+c_2} \\ &= c_2 x\end{aligned}$$

Therefore the solution y is

$$\begin{aligned}y &= ux \\ &= c_2 x^2\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_2 x^2 \tag{1}$$




Figure 4: Slope field plot

Verification of solutions

$$y = c_2 x^2$$

Verified OK.

1.2.4 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{2y}{x}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 2: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= x^2\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{x^2} dy \end{aligned}$$

Which results in

$$S = \frac{y}{x^2}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{2y}{x}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= -\frac{2y}{x^3} \\ S_y &= \frac{1}{x^2} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \tag{4}$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{y}{x^2} = c_1$$



Which simplifies to

$$\frac{y}{x^2} = c_1$$

Which gives

$$y = c_1 x^2$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{2y}{x}$ 	$R = x$ $S = \frac{y}{x^2}$	$\frac{dS}{dR} = 0$ 

Summary

The solution(s) found are the following

$$y = c_1 x^2 \tag{1}$$




Figure 5: Slope field plot

Verification of solutions

$$y = c_1 x^2$$

Verified OK.

1.2.5 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}\left(\frac{1}{2y}\right) dy &= \left(\frac{1}{x}\right) dx \\ \left(-\frac{1}{x}\right) dx + \left(\frac{1}{2y}\right) dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -\frac{1}{x} \\ N(x, y) &= \frac{1}{2y}\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(-\frac{1}{x}\right) \\ &= 0\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{1}{2y} \right) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \tag{1}$$

$$\frac{\partial \phi}{\partial y} = N \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\frac{1}{x} dx \\ \phi &= -\ln(x) + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{1}{2y}$. Therefore equation (4) becomes

$$\frac{1}{2y} = 0 + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = \frac{1}{2y}$$

Integrating the above w.r.t y gives

$$\begin{aligned}\int f'(y) dy &= \int \left(\frac{1}{2y} \right) dy \\ f(y) &= \frac{\ln(y)}{2} + c_1\end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\ln(x) + \frac{\ln(y)}{2} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\ln(x) + \frac{\ln(y)}{2}$$

The solution becomes

$$y = e^{2c_1} x^2$$

Summary

The solution(s) found are the following

$$y = e^{2c_1} x^2 \tag{1}$$




Figure 6: Slope field plot

Verification of solutions

$$y = e^{2c_1} x^2$$

Verified OK.

1.2.6 Maple step by step solution

Let's solve

$$xy' - 2y = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y} = \frac{2}{x}$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y} dx = \int \frac{2}{x} dx + c_1$$

- Evaluate integral

$$\ln(y) = 2 \ln(x) + c_1$$

- Solve for y

$$y = x^2 e^{c_1}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
dsolve(x*diff(y(x),x)=2*y(x),y(x), singsol=all)
```

$$y(x) = c_1 x^2$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 16

```
DSolve[x*y'[x]==2*y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 x^2$$

$$y(x) \rightarrow 0$$

1.3 problem 1(c)

1.3.1	Solving as separable ode	23
1.3.2	Solving as first order ode lie symmetry lookup ode	25
1.3.3	Solving as exact ode	29
1.3.4	Maple step by step solution	33

Internal problem ID [6107]

Internal file name [OUTPUT/5355_Sunday_June_05_2022_03_35_12_PM_30046570/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.2 THE NATURE OF SOLUTIONS. Page 9

Problem number: 1(c).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "separable", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_separable]

$$y'y = e^{2x}$$

1.3.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{e^{2x}}{y}\end{aligned}$$

Where $f(x) = e^{2x}$ and $g(y) = \frac{1}{y}$. Integrating both sides gives

$$\frac{1}{y} dy = e^{2x} dx$$

$$\int \frac{1}{y} dy = \int e^{2x} dx$$

$$\frac{y^2}{2} = \frac{e^{2x}}{2} + c_1$$

Which results in

$$y = \sqrt{e^{2x} + 2c_1}$$

$$y = -\sqrt{e^{2x} + 2c_1}$$

Summary

The solution(s) found are the following

$$y = \sqrt{e^{2x} + 2c_1} \tag{1}$$

$$y = -\sqrt{e^{2x} + 2c_1} \tag{2}$$




Figure 7: Slope field plot

Verification of solutions

$$y = \sqrt{e^{2x} + 2c_1}$$

Verified OK.

$$y = -\sqrt{e^{2x} + 2c_1}$$

Verified OK.

1.3.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{e^{2x}}{y}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **separable**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 5: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= e^{-2x} \\ \eta(x, y) &= 0\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\eta = 0$ then in this special case

$$R = y$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\xi} dx \\ &= \int \frac{1}{e^{-2x}} dx \end{aligned}$$

Which results in

$$S = \frac{e^{2x}}{2}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{e^{2x}}{y}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 0 \\ R_y &= 1 \\ S_x &= e^{2x} \\ S_y &= 0 \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = y \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = R$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \frac{R^2}{2} + c_1 \quad (4)$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{e^{2x}}{2} = \frac{y^2}{2} + c_1$$

Which simplifies to

$$\frac{e^{2x}}{2} = \frac{y^2}{2} + c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{e^{2x}}{y}$ 	$R = y$ $S = \frac{e^{2x}}{2}$	$\frac{dS}{dR} = R$ 

Summary

The solution(s) found are the following

$$\frac{e^{2x}}{2} = \frac{y^2}{2} + c_1 \quad (1)$$




Figure 8: Slope field plot

Verification of solutions

$$\frac{e^{2x}}{2} = \frac{y^2}{2} + c_1$$

Verified OK.

1.3.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}(y) dy &= (e^{2x}) dx \\ (-e^{2x}) dx + (y) dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -e^{2x} \\ N(x, y) &= y\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-e^{2x}) \\ &= 0\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(y) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\int \frac{\partial \phi}{\partial x} dx = \int M dx$$

$$\int \frac{\partial \phi}{\partial x} dx = \int -e^{2x} dx$$

$$\phi = -\frac{e^{2x}}{2} + f(y) \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = y$. Therefore equation (4) becomes

$$y = 0 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = y$$

Integrating the above w.r.t y gives

$$\int f'(y) dy = \int (y) dy$$

$$f(y) = \frac{y^2}{2} + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\frac{e^{2x}}{2} + \frac{y^2}{2} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\frac{e^{2x}}{2} + \frac{y^2}{2}$$

Summary

The solution(s) found are the following

$$-\frac{e^{2x}}{2} + \frac{y^2}{2} = c_1 \tag{1}$$




Figure 9: Slope field plot

Verification of solutions

$$-\frac{e^{2x}}{2} + \frac{y^2}{2} = c_1$$

Verified OK.

1.3.4 Maple step by step solution

Let's solve

$$y'y = e^{2x}$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Integrate both sides with respect to x

$$\int y'y dx = \int e^{2x} dx + c_1$$

- Evaluate integral

$$\frac{y^2}{2} = \frac{e^{2x}}{2} + c_1$$

- Solve for y

$$\{y = \sqrt{e^{2x} + 2c_1}, y = -\sqrt{e^{2x} + 2c_1}\}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
<- Bernoulli successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 25

```
dsolve(y(x)*diff(y(x),x)=exp(2*x),y(x), singsol=all)
```

$$y(x) = \sqrt{e^{2x} + c_1}$$
$$y(x) = -\sqrt{e^{2x} + c_1}$$

✓ Solution by Mathematica

Time used: 0.658 (sec). Leaf size: 39

```
DSolve[y[x]*y'[x]==Exp[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sqrt{e^{2x} + 2c_1}$$

$$y(x) \rightarrow \sqrt{e^{2x} + 2c_1}$$

1.4 problem 1(d)

1.4.1 Solving as quadrature ode	35
1.4.2 Maple step by step solution	36

Internal problem ID [6108]

Internal file name [OUTPUT/5356_Sunday_June_05_2022_03_35_14_PM_47845420/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.2 THE NATURE OF SOLUTIONS. Page 9

Problem number: 1(d).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$y' - yk = 0$$

1.4.1 Solving as quadrature ode

Integrating both sides gives

$$\int \frac{1}{yk} dy = \int dx$$
$$\frac{\ln(y)}{k} = x + c_1$$

Raising both side to exponential gives

$$e^{\frac{\ln(y)}{k}} = e^{x+c_1}$$

Which simplifies to

$$y^{\frac{1}{k}} = c_2 e^x$$

Summary

The solution(s) found are the following

$$y = (c_2 e^x)^k \tag{1}$$

Verification of solutions

$$y = (c_2 e^x)^k$$

Verified OK.

1.4.2 Maple step by step solution

Let's solve

$$y' - yk = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y} = k$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y} dx = \int k dx + c_1$$

- Evaluate integral

$$\ln(y) = kx + c_1$$

- Solve for y

$$y = e^{kx+c_1}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 10

```
dsolve(diff(y(x),x)=k*y(x),y(x), singsol=all)
```

$$y(x) = c_1 e^{kx}$$

✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 18

```
DSolve[y'[x]==k*y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{kx}$$

$$y(x) \rightarrow 0$$

1.5 problem 1(e)

1.5.1	Solving as second order linear constant coeff ode	38
1.5.2	Solving as second order ode can be made integrable ode	40
1.5.3	Solving using Kovacic algorithm	42
1.5.4	Maple step by step solution	46

Internal problem ID [6109]

Internal file name [OUTPUT/5357_Sunday_June_05_2022_03_35_15_PM_52805837/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.2 THE NATURE OF SOLUTIONS. Page 9

Problem number: 1(e).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "second_order_ode_can_be_made_integrable"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x]]
```

$$y'' + 4y = 0$$

1.5.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 4$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 4 e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 4 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 4$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(4)} \\ &= \pm 2i \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +2i \\ \lambda_2 &= -2i \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= 2i \\ \lambda_2 &= -2i \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 2$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (c_1 \cos(2x) + c_2 \sin(2x))$$

Or

$$y = c_1 \cos(2x) + c_2 \sin(2x)$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(2x) + c_2 \sin(2x) \tag{1}$$




Figure 10: Slope field plot

Verification of solutions

$$y = c_1 \cos(2x) + c_2 \sin(2x)$$

Verified OK.

1.5.2 Solving as second order ode can be made integrable ode

Multiplying the ode by y' gives

$$y'y'' + 4y'y = 0$$

Integrating the above w.r.t x gives

$$\int (y'y'' + 4y'y) dx = 0$$

$$\frac{y'^2}{2} + 2y^2 = c_2$$

Which is now solved for y . Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \sqrt{-4y^2 + 2c_1} \quad (1)$$

$$y' = -\sqrt{-4y^2 + 2c_1} \quad (2)$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int \frac{1}{\sqrt{-4y^2 + 2c_1}} dy = \int dx$$
$$\frac{\arctan\left(\frac{2y}{\sqrt{-4y^2 + 2c_1}}\right)}{2} = c_2 + x$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{1}{\sqrt{-4y^2 + 2c_1}} dy = \int dx$$
$$-\frac{\arctan\left(\frac{2y}{\sqrt{-4y^2 + 2c_1}}\right)}{2} = x + c_3$$

Summary

The solution(s) found are the following

$$\frac{\arctan\left(\frac{2y}{\sqrt{-4y^2 + 2c_1}}\right)}{2} = c_2 + x \quad (1)$$

$$-\frac{\arctan\left(\frac{2y}{\sqrt{-4y^2 + 2c_1}}\right)}{2} = x + c_3 \quad (2)$$




Figure 11: Slope field plot

Verification of solutions

$$\frac{\arctan\left(\frac{2y}{\sqrt{-4y^2+2c_1}}\right)}{2} = c_2 + x$$

Verified OK.

$$-\frac{\arctan\left(\frac{2y}{\sqrt{-4y^2+2c_1}}\right)}{2} = x + c_3$$

Verified OK.

1.5.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 4y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= 1 \\B &= 0 \\C &= 4\end{aligned}\tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x)\tag{4}$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\&= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned}\tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-4}{1}\tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= -4 \\t &= 1\end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -4z(x)\tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 9: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
O(\infty) &= \deg(t) - \deg(s) \\
&= 0 - 0 \\
&= 0
\end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -4$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(2x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}
y_1 &= z_1 \\
&= \cos(2x)
\end{aligned}$$

Which simplifies to

$$y_1 = \cos(2x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(2x) \int \frac{1}{\cos(2x)^2} dx \\ &= \cos(2x) \left(\frac{\tan(2x)}{2} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1(\cos(2x)) + c_2 \left(\cos(2x) \left(\frac{\tan(2x)}{2} \right) \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(2x) + \frac{c_2 \sin(2x)}{2} \tag{1}$$




Figure 12: Slope field plot

Verification of solutions

$$y = c_1 \cos(2x) + \frac{c_2 \sin(2x)}{2}$$

Verified OK.

1.5.4 Maple step by step solution

Let's solve

$$y'' + 4y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 + 4 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-16})}{2}$$

- Roots of the characteristic polynomial

$$r = (-2I, 2I)$$

- 1st solution of the ODE
 $y_1(x) = \cos(2x)$
- 2nd solution of the ODE
 $y_2(x) = \sin(2x)$
- General solution of the ODE
 $y = c_1 y_1(x) + c_2 y_2(x)$
- Substitute in solutions
 $y = c_1 \cos(2x) + c_2 \sin(2x)$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)+4*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 \sin(2x) + c_2 \cos(2x)$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 20

```
DSolve[y''[x]+4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos(2x) + c_2 \sin(2x)$$

1.6 problem 1(f)

1.6.1	Solving as second order linear constant coeff ode	48
1.6.2	Solving as second order ode can be made integrable ode	50
1.6.3	Solving using Kovacic algorithm	52
1.6.4	Maple step by step solution	56

Internal problem ID [6110]

Internal file name [OUTPUT/5358_Sunday_June_05_2022_03_35_16_PM_82047167/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.2 THE NATURE OF SOLUTIONS. Page 9

Problem number: 1(f).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "second_order_ode_can_be_made_integrable"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x]]
```

$$y'' - 4y = 0$$

1.6.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = -4$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 4 e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 4 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = -4$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(-4)} \\ &= \pm 2 \end{aligned}$$

Hence

$$\lambda_1 = +2$$

$$\lambda_2 = -2$$

Which simplifies to

$$\lambda_1 = 2$$

$$\lambda_2 = -2$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(2)x} + c_2 e^{(-2)x}$$

Or

$$y = c_1 e^{2x} + c_2 e^{-2x}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{2x} + c_2 e^{-2x} \tag{1}$$




Figure 13: Slope field plot

Verification of solutions

$$y = c_1 e^{2x} + c_2 e^{-2x}$$

Verified OK.

1.6.2 Solving as second order ode can be made integrable ode

Multiplying the ode by y' gives

$$y' y'' - 4y' y = 0$$

Integrating the above w.r.t x gives

$$\int (y' y'' - 4y' y) dx = 0$$

$$\frac{y'^2}{2} - 2y^2 = c_2$$

Which is now solved for y . Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \sqrt{4y^2 + 2c_1} \tag{1}$$

$$y' = -\sqrt{4y^2 + 2c_1} \tag{2}$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int \frac{1}{\sqrt{4y^2 + 2c_1}} dy = \int dx$$
$$\frac{\ln(y\sqrt{4} + \sqrt{4y^2 + 2c_1})\sqrt{4}}{4} = c_2 + x$$

Raising both side to exponential gives

$$e^{\frac{\ln(y\sqrt{4} + \sqrt{4y^2 + 2c_1})\sqrt{4}}{4}} = e^{c_2 + x}$$

Which simplifies to

$$\sqrt{2y + \sqrt{4y^2 + 2c_1}} = c_3 e^x$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{1}{\sqrt{4y^2 + 2c_1}} dy = \int dx$$
$$-\frac{\ln(y\sqrt{4} + \sqrt{4y^2 + 2c_1})\sqrt{4}}{4} = x + c_4$$

Raising both side to exponential gives

$$e^{-\frac{\ln(y\sqrt{4} + \sqrt{4y^2 + 2c_1})\sqrt{4}}{4}} = e^{x + c_4}$$

Which simplifies to

$$\frac{1}{\sqrt{2y + \sqrt{4y^2 + 2c_1}}} = c_5 e^x$$

Summary

The solution(s) found are the following

$$y = \frac{(e^{4x} c_3^4 - 2c_1) e^{-2x}}{4c_3^2} \quad (1)$$

$$y = -\frac{(2c_1 c_5^4 e^{4x} - 1) e^{-2x}}{4c_5^2} \quad (2)$$




Figure 14: Slope field plot

Verification of solutions

$$y = \frac{(e^{4x}c_3^4 - 2c_1) e^{-2x}}{4c_3^2}$$

Verified OK.

$$y = -\frac{(2c_1c_5^4e^{4x} - 1) e^{-2x}}{4c_5^2}$$

Verified OK.

1.6.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 4y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= 1 \\B &= 0 \\C &= -4\end{aligned}\tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x)\tag{4}$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\&= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned}\tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{4}{1}\tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= 4 \\t &= 1\end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 4z(x)\tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 11: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
\mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
&= 0 - 0 \\
&= 0
\end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 4$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-2x}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}
y_1 &= z_1 \\
&= e^{-2x}
\end{aligned}$$

Which simplifies to

$$y_1 = e^{-2x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= e^{-2x} \int \frac{1}{e^{-4x}} dx \\ &= e^{-2x} \left(\frac{e^{4x}}{4} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-2x}) + c_2 \left(e^{-2x} \left(\frac{e^{4x}}{4} \right) \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-2x} + \frac{c_2 e^{2x}}{4} \tag{1}$$




Figure 15: Slope field plot

Verification of solutions

$$y = c_1 e^{-2x} + \frac{c_2 e^{2x}}{4}$$

Verified OK.

1.6.4 Maple step by step solution

Let's solve

$$y'' - 4y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 - 4 = 0$$

- Factor the characteristic polynomial

$$(r - 2)(r + 2) = 0$$

- Roots of the characteristic polynomial

- $r = (-2, 2)$
- 1st solution of the ODE
 $y_1(x) = e^{-2x}$
 - 2nd solution of the ODE
 $y_2(x) = e^{2x}$
 - General solution of the ODE
 $y = c_1y_1(x) + c_2y_2(x)$
 - Substitute in solutions
 $y = c_1e^{-2x} + c_2e^{2x}$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)-4*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{2x}c_1 + c_2e^{-2x}$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 22

```
DSolve[y''[x]-4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2x}(c_1e^{4x} + c_2)$$

1.7 problem 1(h)

Internal problem ID [6111]

Internal file name [OUTPUT/5359_Sunday_June_05_2022_03_35_17_PM_19549517/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.2 THE NATURE OF SOLUTIONS. Page 9

Problem number: 1(h).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**unknown**"

Maple gives the following as the ode type

`[_rational]`

Unable to solve or complete the solution.

$$xy' + y - y'\sqrt{1 - yx^2} = 0$$

Unable to determine ODE type.

Maple trace

```
`Methods for first order ODEs:
--- Trying classification methods ---
trying homogeneous types:
differential order: 1; looking for linear symmetries
trying exact
Looking for potential symmetries
trying an equivalence to an Abel ODE
trying 1st order ODE linearizable_by_differentiation
--- Trying Lie symmetry methods, 1st order ---
`, `-> Computing symmetries using: way = 2
`, `-> Computing symmetries using: way = 3
`, `-> Computing symmetries using: way = 4
`, `-> Computing symmetries using: way = 5
trying symmetry patterns for 1st order ODEs
-> trying a symmetry pattern of the form [F(x)*G(y), 0]
-> trying a symmetry pattern of the form [0, F(x)*G(y)]
-> trying symmetry patterns of the forms [F(x),G(y)] and [G(y),F(x)]
-> trying a symmetry pattern of the form [F(x),G(x)]
-> trying a symmetry pattern of the form [F(y),G(y)]
-> trying a symmetry pattern of the form [F(x)+G(y), 0]
-> trying a symmetry pattern of the form [0, F(x)+G(y)]
-> trying a symmetry pattern of the form [F(x),G(x)*y+H(x)]
-> trying a symmetry pattern of conformal type`
```

X Solution by Maple

```
dsolve(x*diff(y(x),x)+y(x)=diff(y(x),x)*sqrt(1-x^2*y(x)),y(x), singsol=all)
```

No solution found

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[x*y'[x]+y[x]==y[x]*Sqrt[1-x^2*y[x]],y[x],x,IncludeSingularSolutions -> True]
```

Not solved

1.8 problem 1(i)

1.8.1	Solving as homogeneousTypeD2 ode	60
1.8.2	Solving as exact ode	62
1.8.3	Solving as riccati ode	67

Internal problem ID [6112]

Internal file name [OUTPUT/5360_Sunday_June_05_2022_03_35_19_PM_88813315/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.2 THE NATURE OF SOLUTIONS. Page 9

Problem number: 1(i).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**riccati**", "**exactByInspection**", "**homogeneousTypeD2**"

Maple gives the following as the ode type

```
[[_homogeneous, `class D`], _rational, _Riccati]
```

$$-y + xy' - y^2 = x^2$$

1.8.1 Solving as homogeneousTypeD2 ode

Using the change of variables $y = u(x)x$ on the above ode results in new ode in $u(x)$

$$-u(x)x + x(u'(x)x + u(x)) - u(x)^2 x^2 = x^2$$

Integrating both sides gives

$$\int \frac{1}{u^2 + 1} du = c_2 + x$$
$$\arctan(u) = c_2 + x$$

Solving for u gives these solutions

$$u_1 = \tan(c_2 + x)$$

Therefore the solution y is

$$\begin{aligned}y &= xu \\ &= x \tan(c_2 + x)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = x \tan(c_2 + x) \tag{1}$$




Figure 16: Slope field plot

Verification of solutions

$$y = x \tan(c_2 + x)$$

Verified OK.

1.8.2 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial\phi}{\partial x} + \frac{\partial\phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} (x) dy &= (x^2 + y^2 + y) dx \\ (-x^2 - y^2 - y) dx + (x) dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -x^2 - y^2 - y \\ N(x, y) &= x \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-x^2 - y^2 - y) \\ &= -2y - 1\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(x) \\ &= 1\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. By inspection $\frac{1}{x^2+y^2}$ is an integrating factor. Therefore by multiplying $M = -y^2 - x^2 - y$ and $N = x$ by this integrating factor the ode becomes exact. The new M, N are

$$\begin{aligned}M &= \frac{-y^2 - x^2 - y}{x^2 + y^2} \\ N &= \frac{x}{x^2 + y^2}\end{aligned}$$

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned} \left(\frac{x}{x^2 + y^2} \right) dy &= \left(-\frac{-x^2 - y^2 - y}{x^2 + y^2} \right) dx \\ \left(\frac{-x^2 - y^2 - y}{x^2 + y^2} \right) dx + \left(\frac{x}{x^2 + y^2} \right) dy &= 0 \end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= \frac{-x^2 - y^2 - y}{x^2 + y^2} \\ N(x, y) &= \frac{x}{x^2 + y^2} \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(\frac{-x^2 - y^2 - y}{x^2 + y^2} \right) \\ &= \frac{-x^2 + y^2}{(x^2 + y^2)^2} \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{x}{x^2 + y^2} \right) \\ &= \frac{-x^2 + y^2}{(x^2 + y^2)^2} \end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned} \int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int \frac{-x^2 - y^2 - y}{x^2 + y^2} dx \\ \phi &= -x - \arctan\left(\frac{x}{y}\right) + f(y) \end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\begin{aligned} \frac{\partial \phi}{\partial y} &= \frac{x}{y^2 \left(\frac{x^2}{y^2} + 1\right)} + f'(y) \\ &= \frac{x}{x^2 + y^2} + f'(y) \end{aligned} \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{x}{x^2 + y^2}$. Therefore equation (4) becomes

$$\frac{x}{x^2 + y^2} = \frac{x}{x^2 + y^2} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = -x - \arctan\left(\frac{x}{y}\right) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -x - \arctan\left(\frac{x}{y}\right)$$

The solution becomes

$$y = -\frac{x}{\tan(x + c_1)}$$

Summary

The solution(s) found are the following

$$y = -\frac{x}{\tan(x + c_1)} \tag{1}$$




Figure 17: Slope field plot

Verification of solutions

$$y = -\frac{x}{\tan(x + c_1)}$$

Verified OK.

1.8.3 Solving as riccati ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= \frac{x^2 + y^2 + y}{x}\end{aligned}$$

This is a Riccati ODE. Comparing the ODE to solve

$$y' = x + \frac{y^2}{x} + \frac{y}{x}$$

With Riccati ODE standard form

$$y' = f_0(x) + f_1(x)y + f_2(x)y^2$$

Shows that $f_0(x) = x$, $f_1(x) = \frac{1}{x}$ and $f_2(x) = \frac{1}{x}$. Let

$$\begin{aligned}y &= \frac{-u'}{f_2 u} \\ &= \frac{-u'}{\frac{u}{x}}\end{aligned}\tag{1}$$

Using the above substitution in the given ODE results (after some simplification) in a second order ODE to solve for $u(x)$ which is

$$f_2 u''(x) - (f_2' + f_1 f_2) u'(x) + f_2^2 f_0 u(x) = 0\tag{2}$$

But

$$\begin{aligned}f_2' &= -\frac{1}{x^2} \\ f_1 f_2 &= \frac{1}{x^2} \\ f_2^2 f_0 &= \frac{1}{x}\end{aligned}$$

Substituting the above terms back in equation (2) gives

$$\frac{u''(x)}{x} + \frac{u(x)}{x} = 0$$

Solving the above ODE (this ode solved using Maple, not this program), gives

$$u(x) = \sin(x) c_1 + c_2 \cos(x)$$

The above shows that

$$u'(x) = \cos(x) c_1 - c_2 \sin(x)$$

Using the above in (1) gives the solution

$$y = -\frac{(\cos(x) c_1 - c_2 \sin(x)) x}{\sin(x) c_1 + c_2 \cos(x)}$$

Dividing both numerator and denominator by c_1 gives, after renaming the constant $\frac{c_2}{c_1} = c_3$ the following solution

$$y = \frac{x(-c_3 \cos(x) + \sin(x))}{c_3 \sin(x) + \cos(x)}$$

Summary

The solution(s) found are the following

$$y = \frac{x(-c_3 \cos(x) + \sin(x))}{c_3 \sin(x) + \cos(x)} \tag{1}$$




Figure 18: Slope field plot

Verification of solutions

$$y = \frac{x(-c_3 \cos(x) + \sin(x))}{c_3 \sin(x) + \cos(x)}$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying homogeneous D  
<- homogeneous successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 10

```
dsolve(x*diff(y(x),x)=y(x)+x^2+y(x)^2,y(x), singsol=all)
```

$$y(x) = \tan(x + c_1) x$$

✓ Solution by Mathematica

Time used: 0.184 (sec). Leaf size: 12

```
DSolve[x*y'[x]==y[x]+x^2+y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x \tan(x + c_1)$$

1.9 problem 1(j)

1.9.1	Solving as homogeneousTypeD2 ode	70
1.9.2	Solving as first order ode lie symmetry calculated ode	72
1.9.3	Solving as exact ode	77

Internal problem ID [6113]

Internal file name [OUTPUT/5361_Sunday_June_05_2022_03_35_20_PM_40589996/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.2 THE NATURE OF SOLUTIONS. Page 9

Problem number: 1(j).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**homogeneousTypeD2**", "**exactWithIntegrationFactor**", "**first_order_ode_lie_symmetry_calculated**"

Maple gives the following as the ode type

```
[[_homogeneous, `class A`], _rational, _dAlembert]
```

$$y' - \frac{xy}{x^2 + y^2} = 0$$

1.9.1 Solving as homogeneousTypeD2 ode

Using the change of variables $y = u(x)x$ on the above ode results in new ode in $u(x)$

$$u'(x)x + u(x) - \frac{x^2 u(x)}{x^2 + u(x)^2 x^2} = 0$$

In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{u^3}{x(u^2 + 1)} \end{aligned}$$

Where $f(x) = -\frac{1}{x}$ and $g(u) = \frac{u^3}{u^2+1}$. Integrating both sides gives

$$\begin{aligned}\frac{1}{\frac{u^3}{u^2+1}} du &= -\frac{1}{x} dx \\ \int \frac{1}{\frac{u^3}{u^2+1}} du &= \int -\frac{1}{x} dx \\ \ln(u) - \frac{1}{2u^2} &= -\ln(x) + c_2\end{aligned}$$

The solution is

$$\ln(u(x)) - \frac{1}{2u(x)^2} + \ln(x) - c_2 = 0$$

Replacing $u(x)$ in the above solution by $\frac{y}{x}$ results in the solution for y in implicit form

$$\begin{aligned}\ln\left(\frac{y}{x}\right) - \frac{x^2}{2y^2} + \ln(x) - c_2 &= 0 \\ \ln\left(\frac{y}{x}\right) - \frac{x^2}{2y^2} + \ln(x) - c_2 &= 0\end{aligned}$$

Summary

The solution(s) found are the following

$$\ln\left(\frac{y}{x}\right) - \frac{x^2}{2y^2} + \ln(x) - c_2 = 0 \tag{1}$$




Figure 19: Slope field plot

Verification of solutions

$$\ln\left(\frac{y}{x}\right) - \frac{x^2}{2y^2} + \ln(x) - c_2 = 0$$

Verified OK.

1.9.2 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = \frac{xy}{x^2 + y^2}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \quad (1\text{E})$$

$$\eta = xb_2 + yb_3 + b_1 \quad (2\text{E})$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$b_2 + \frac{xy(b_3 - a_2)}{x^2 + y^2} - \frac{x^2 y^2 a_3}{(x^2 + y^2)^2} - \left(\frac{y}{x^2 + y^2} - \frac{2x^2 y}{(x^2 + y^2)^2} \right) (xa_2 + ya_3 + a_1) \quad (5E)$$

$$- \left(\frac{x}{x^2 + y^2} - \frac{2xy^2}{(x^2 + y^2)^2} \right) (xb_2 + yb_3 + b_1) = 0$$

Putting the above in normal form gives

$$\frac{-3x^2 y^2 b_2 + 2x y^3 a_2 - 2x y^3 b_3 + y^4 a_3 - y^4 b_2 + x^3 b_1 - x^2 y a_1 - x y^2 b_1 + y^3 a_1}{(x^2 + y^2)^2} = 0$$

Setting the numerator to zero gives

$$3x^2 y^2 b_2 - 2x y^3 a_2 + 2x y^3 b_3 - y^4 a_3 + y^4 b_2 - x^3 b_1 + x^2 y a_1 + x y^2 b_1 - y^3 a_1 = 0 \quad (6E)$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2\}$$

The above PDE (6E) now becomes

$$-2a_2 v_1 v_2^3 - a_3 v_2^4 + 3b_2 v_1^2 v_2^2 + b_2 v_2^4 + 2b_3 v_1 v_2^3 + a_1 v_1^2 v_2 - a_1 v_2^3 - b_1 v_1^3 + b_1 v_1 v_2^2 = 0 \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2\}$$

Equation (7E) now becomes

$$-b_1 v_1^3 + 3b_2 v_1^2 v_2^2 + a_1 v_1^2 v_2 + (-2a_2 + 2b_3) v_1 v_2^3 + b_1 v_1 v_2^2 + (-a_3 + b_2) v_2^4 - a_1 v_2^3 = 0 \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned}
 a_1 &= 0 \\
 b_1 &= 0 \\
 -a_1 &= 0 \\
 -b_1 &= 0 \\
 3b_2 &= 0 \\
 -2a_2 + 2b_3 &= 0 \\
 -a_3 + b_2 &= 0
 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned}
 a_1 &= 0 \\
 a_2 &= b_3 \\
 a_3 &= 0 \\
 b_1 &= 0 \\
 b_2 &= 0 \\
 b_3 &= b_3
 \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned}
 \xi &= x \\
 \eta &= y
 \end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned}
 \eta &= \eta - \omega(x, y) \xi \\
 &= y - \left(\frac{xy}{x^2 + y^2} \right) (x) \\
 &= \frac{y^3}{x^2 + y^2} \\
 \xi &= 0
 \end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{y^3}{x^2+y^2}} dy \end{aligned}$$

Which results in

$$S = \ln(y) - \frac{x^2}{2y^2}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{xy}{x^2 + y^2}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= -\frac{x}{y^2} \\ S_y &= \frac{x^2 + y^2}{y^3} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \tag{4}$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{2y^2 \ln(y) - x^2}{2y^2} = c_1$$



Which simplifies to

$$\frac{2y^2 \ln(y) - x^2}{2y^2} = c_1$$

Which gives

$$y = e^{\frac{\text{LambertW}(x^2 e^{-2c_1})}{2} + c_1}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{xy}{x^2 + y^2}$ 	$R = x$ $S = \frac{2 \ln(y) y^2 - x^2}{2y^2}$	$\frac{dS}{dR} = 0$ 

Summary

The solution(s) found are the following

$$y = e^{\frac{\text{LambertW}(x^2 e^{-2c_1})}{2} + c_1} \quad (1)$$




Figure 20: Slope field plot

Verification of solutions

$$y = e^{\frac{\text{LambertW}(x^2 e^{-2c_1})}{2} + c_1}$$

Verified OK.

1.9.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} (x^2 + y^2) dy &= (xy) dx \\ (-xy) dx + (x^2 + y^2) dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -xy \\ N(x, y) &= x^2 + y^2 \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-xy) \\ &= -x \end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(x^2 + y^2) \\ &= 2x\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned}A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= \frac{1}{x^2 + y^2} ((-x) - (2x)) \\ &= -\frac{3x}{x^2 + y^2}\end{aligned}$$

Since A depends on y , it can not be used to obtain an integrating factor. We will now try a second method to find an integrating factor. Let

$$\begin{aligned}B &= \frac{1}{M} \left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) \\ &= -\frac{1}{yx} ((2x) - (-x)) \\ &= -\frac{3}{y}\end{aligned}$$

Since B does not depend on x , it can be used to obtain an integrating factor. Let the integrating factor be μ . Then

$$\begin{aligned}\mu &= e^{\int B \, dy} \\ &= e^{\int -\frac{3}{y} \, dy}\end{aligned}$$

The result of integrating gives

$$\begin{aligned}\mu &= e^{-3 \ln(y)} \\ &= \frac{1}{y^3}\end{aligned}$$

M and N are now multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} so not to confuse them with the original M and N .

$$\begin{aligned}\bar{M} &= \mu M \\ &= \frac{1}{y^3}(-xy) \\ &= -\frac{x}{y^2}\end{aligned}$$

And

$$\begin{aligned}\bar{N} &= \mu N \\ &= \frac{1}{y^3}(x^2 + y^2) \\ &= \frac{x^2 + y^2}{y^3}\end{aligned}$$

So now a modified ODE is obtained from the original ODE which will be exact and can be solved using the standard method. The modified ODE is

$$\begin{aligned}\bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ \left(-\frac{x}{y^2}\right) + \left(\frac{x^2 + y^2}{y^3}\right) \frac{dy}{dx} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\frac{x}{y^2} dx \\ \phi &= -\frac{x^2}{2y^2} + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = \frac{x^2}{y^3} + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{x^2 + y^2}{y^3}$. Therefore equation (4) becomes

$$\frac{x^2 + y^2}{y^3} = \frac{x^2}{y^3} + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = \frac{1}{y}$$

Integrating the above w.r.t y gives

$$\int f'(y) dy = \int \left(\frac{1}{y}\right) dy$$
$$f(y) = \ln(y) + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = \ln(y) - \frac{x^2}{2y^2} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \ln(y) - \frac{x^2}{2y^2}$$

The solution becomes

$$y = e^{\frac{\text{LambertW}(x^2 e^{-2c_1})}{2} + c_1}$$

Summary

The solution(s) found are the following

$$y = e^{\frac{\text{LambertW}(x^2 e^{-2c_1})}{2} + c_1} \quad (1)$$




Figure 21: Slope field plot

Verification of solutions

$$y = e^{\frac{\text{LambertW}(x^2 e^{-2c_1})}{2}} + c_1$$

Verified OK.

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying homogeneous D
<- homogeneous successful`

```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 16

```
dsolve(diff(y(x),x)=(x*y(x))/(x^2+y(x)^2),y(x), singsol=all)
```

$$y(x) = \sqrt{\frac{1}{\text{LambertW}(c_1 x^2)}} x$$

✓ Solution by Mathematica

Time used: 7.664 (sec). Leaf size: 49

```
DSolve[y'[x]==(x*y[x])/(x^2+y[x]^2),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{x}{\sqrt{W(e^{-2c_1 x^2})}}$$
$$y(x) \rightarrow \frac{x}{\sqrt{W(e^{-2c_1 x^2})}}$$
$$y(x) \rightarrow 0$$

1.10 problem 1(k)

1.10.1 Solving as homogeneousTypeD2 ode	84
1.10.2 Solving as first order ode lie symmetry lookup ode	86
1.10.3 Solving as bernoulli ode	90
1.10.4 Solving as exact ode	94

Internal problem ID [6114]

Internal file name [OUTPUT/5362_Sunday_June_05_2022_03_35_22_PM_9672469/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.2 THE NATURE OF SOLUTIONS. Page 9

Problem number: 1(k).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**bernoulli**", "**homogeneousTypeD2**", "**exactWithIntegrationFactor**", "**first_order_ode_lie_symmetry_lookup**"

Maple gives the following as the ode type

```
[[_homogeneous, `class A`], _rational, _Bernoulli]
```

$$2xyy' - y^2 = x^2$$

1.10.1 Solving as homogeneousTypeD2 ode

Using the change of variables $y = u(x)x$ on the above ode results in new ode in $u(x)$

$$2x^2u(x)(u'(x)x + u(x)) - u(x)^2x^2 = x^2$$

In canonical form the ODE is

$$\begin{aligned}u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{u^2 - 1}{2ux}\end{aligned}$$

Where $f(x) = -\frac{1}{2x}$ and $g(u) = \frac{u^2-1}{u}$. Integrating both sides gives

$$\begin{aligned}\frac{1}{\frac{u^2-1}{u}} du &= -\frac{1}{2x} dx \\ \int \frac{1}{\frac{u^2-1}{u}} du &= \int -\frac{1}{2x} dx \\ \frac{\ln(u-1)}{2} + \frac{\ln(u+1)}{2} &= -\frac{\ln(x)}{2} + c_2\end{aligned}$$

The above can be written as

$$\begin{aligned}\left(\frac{1}{2}\right) (\ln(u-1) + \ln(u+1)) &= -\frac{\ln(x)}{2} + 2c_2 \\ \ln(u-1) + \ln(u+1) &= (2) \left(-\frac{\ln(x)}{2} + 2c_2\right) \\ &= -\ln(x) + 4c_2\end{aligned}$$

Raising both side to exponential gives

$$e^{\ln(u-1)+\ln(u+1)} = e^{-\ln(x)+2c_2}$$

Which simplifies to

$$\begin{aligned}u^2 - 1 &= \frac{2c_2}{x} \\ &= \frac{c_3}{x}\end{aligned}$$

The solution is

$$u(x)^2 - 1 = \frac{c_3}{x}$$

Replacing $u(x)$ in the above solution by $\frac{y}{x}$ results in the solution for y in implicit form

$$\begin{aligned}\frac{y^2}{x^2} - 1 &= \frac{c_3}{x} \\ \frac{y^2}{x^2} - 1 &= \frac{c_3}{x}\end{aligned}$$

Summary

The solution(s) found are the following

$$\frac{y^2}{x^2} - 1 = \frac{c_3}{x} \tag{1}$$




Figure 22: Slope field plot

Verification of solutions

$$\frac{y^2}{x^2} - 1 = \frac{c_3}{x}$$

Verified OK.

1.10.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{x^2 + y^2}{2xy}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \tag{A}$$

The type of this ode is known. It is of type **Bernoulli**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 13: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= \frac{x}{y}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{x}{y}} dy \end{aligned}$$

Which results in

$$S = \frac{y^2}{2x}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{x^2 + y^2}{2xy}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= -\frac{y^2}{2x^2} \\ S_y &= \frac{y}{x} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \frac{1}{2} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \frac{1}{2}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \frac{R}{2} + c_1 \quad (4)$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{y^2}{2x} = \frac{x}{2} + c_1$$

Which simplifies to

$$\frac{y^2}{2x} = \frac{x}{2} + c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{x^2+y^2}{2xy}$ 	$R = x$ $S = \frac{y^2}{2x}$	$\frac{dS}{dR} = \frac{1}{2}$ 

Summary

The solution(s) found are the following

$$\frac{y^2}{2x} = \frac{x}{2} + c_1 \quad (1)$$




Figure 23: Slope field plot

Verification of solutions

$$\frac{y^2}{2x} = \frac{x}{2} + c_1$$

Verified OK.

1.10.3 Solving as bernoulli ode

In canonical form, the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= \frac{x^2 + y^2}{2xy} \end{aligned}$$

This is a Bernoulli ODE.

$$y' = \frac{1}{2x}y + \frac{x}{2} \frac{1}{y} \tag{1}$$

The standard Bernoulli ODE has the form

$$y' = f_0(x)y + f_1(x)y^n \tag{2}$$

The first step is to divide the above equation by y^n which gives

$$\frac{y'}{y^n} = f_0(x)y^{1-n} + f_1(x) \tag{3}$$

The next step is use the substitution $w = y^{1-n}$ in equation (3) which generates a new ODE in $w(x)$ which will be linear and can be easily solved using an integrating factor. Backsubstitution then gives the solution $y(x)$ which is what we want.

This method is now applied to the ODE at hand. Comparing the ODE (1) With (2) Shows that

$$\begin{aligned} f_0(x) &= \frac{1}{2x} \\ f_1(x) &= \frac{x}{2} \\ n &= -1 \end{aligned}$$

Dividing both sides of ODE (1) by $y^n = \frac{1}{y}$ gives

$$y'y = \frac{y^2}{2x} + \frac{x}{2} \quad (4)$$

Let

$$\begin{aligned} w &= y^{1-n} \\ &= y^2 \end{aligned} \quad (5)$$

Taking derivative of equation (5) w.r.t x gives

$$w' = 2yy' \quad (6)$$

Substituting equations (5) and (6) into equation (4) gives

$$\begin{aligned} \frac{w'(x)}{2} &= \frac{w(x)}{2x} + \frac{x}{2} \\ w' &= \frac{w}{x} + x \end{aligned} \quad (7)$$

The above now is a linear ODE in $w(x)$ which is now solved.

Entering Linear first order ODE solver. In canonical form a linear first order is

$$w'(x) + p(x)w(x) = q(x)$$

Where here

$$\begin{aligned} p(x) &= -\frac{1}{x} \\ q(x) &= x \end{aligned}$$

Hence the ode is

$$w'(x) - \frac{w(x)}{x} = x$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int -\frac{1}{x} dx} \\ &= \frac{1}{x}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu w) &= (\mu)(x) \\ \frac{d}{dx}\left(\frac{w}{x}\right) &= \left(\frac{1}{x}\right)(x) \\ d\left(\frac{w}{x}\right) &= dx\end{aligned}$$

Integrating gives

$$\begin{aligned}\frac{w}{x} &= \int dx \\ \frac{w}{x} &= x + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = \frac{1}{x}$ results in

$$w(x) = c_1 x + x^2$$

which simplifies to

$$w(x) = x(x + c_1)$$

Replacing w in the above by y^2 using equation (5) gives the final solution.

$$y^2 = x(x + c_1)$$

Solving for y gives

$$\begin{aligned}y(x) &= \sqrt{x(x + c_1)} \\ y(x) &= -\sqrt{x(x + c_1)}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \sqrt{x(x + c_1)} \quad (1)$$

$$y = -\sqrt{x(x + c_1)} \quad (2)$$




Figure 24: Slope field plot

Verification of solutions

$$y = \sqrt{x(x + c_1)}$$

Verified OK.

$$y = -\sqrt{x(x + c_1)}$$

Verified OK.

1.10.4 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial\phi}{\partial x} + \frac{\partial\phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} (2xy) dy &= (x^2 + y^2) dx \\ (-x^2 - y^2) dx + (2xy) dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -x^2 - y^2 \\ N(x, y) &= 2xy \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-x^2 - y^2) \\ &= -2y\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(2xy) \\ &= 2y\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned}A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= \frac{1}{2yx} ((-2y) - (2y)) \\ &= -\frac{2}{x}\end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int A dx} \\ &= e^{\int -\frac{2}{x} dx}\end{aligned}$$

The result of integrating gives

$$\begin{aligned}\mu &= e^{-2 \ln(x)} \\ &= \frac{1}{x^2}\end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} for now so not to confuse them with the original M and N .

$$\begin{aligned}\bar{M} &= \mu M \\ &= \frac{1}{x^2}(-x^2 - y^2) \\ &= \frac{-x^2 - y^2}{x^2}\end{aligned}$$

And

$$\begin{aligned}\bar{N} &= \mu N \\ &= \frac{1}{x^2}(2xy) \\ &= \frac{2y}{x}\end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned}\bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ \left(\frac{-x^2 - y^2}{x^2} \right) + \left(\frac{2y}{x} \right) \frac{dy}{dx} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \quad (1)$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int \frac{-x^2 - y^2}{x^2} dx \\ \phi &= -x + \frac{y^2}{x} + f(y)\end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = \frac{2y}{x} + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{2y}{x}$. Therefore equation (4) becomes

$$\frac{2y}{x} = \frac{2y}{x} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = -x + \frac{y^2}{x} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -x + \frac{y^2}{x}$$

Summary

The solution(s) found are the following

$$-x + \frac{y^2}{x} = c_1 \tag{1}$$




Figure 25: Slope field plot

Verification of solutions

$$-x + \frac{y^2}{x} = c_1$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
<- Bernoulli successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(2*x*y(x)*diff(y(x),x)=x^2+y(x)^2,y(x), singsol=all)
```

$$y(x) = \sqrt{(x + c_1)x}$$
$$y(x) = -\sqrt{(x + c_1)x}$$

✓ Solution by Mathematica

Time used: 0.182 (sec). Leaf size: 38

```
DSolve[2*x*y[x]*y'[x]==x^2+y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sqrt{x}\sqrt{x + c_1}$$
$$y(x) \rightarrow \sqrt{x}\sqrt{x + c_1}$$

1.11 problem 1(L)

Internal problem ID [6115]

Internal file name [OUTPUT/5363_Sunday_June_05_2022_03_35_24_PM_66181348/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.2 THE NATURE OF SOLUTIONS. Page 9

Problem number: 1(L).

ODE order: 1.

ODE degree: 2.

The type(s) of ODE detected by this program : "first_order_ode_lie_symmetry_calculated"

Maple gives the following as the ode type

```
[[_homogeneous, `class G`], _rational]
```

$$xy' + y - x^4y'^2 = 0$$

Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \frac{1 + \sqrt{1 + 4yx^2}}{2x^3} \quad (1)$$

$$y' = -\frac{-1 + \sqrt{1 + 4yx^2}}{2x^3} \quad (2)$$

Now each one of the above ODE is solved.

Solving equation (1)

Writing the ode as

$$y' = \frac{1 + \sqrt{4yx^2 + 1}}{2x^3}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2\xi_y - \omega_x\xi - \omega_y\eta = 0 \quad (\text{A})$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \quad (1E)$$

$$\eta = xb_2 + yb_3 + b_1 \quad (2E)$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$\begin{aligned} b_2 + \frac{(1 + \sqrt{4yx^2 + 1})(b_3 - a_2)}{2x^3} - \frac{(1 + \sqrt{4yx^2 + 1})^2 a_3}{4x^6} \\ - \left(-\frac{3(1 + \sqrt{4yx^2 + 1})}{2x^4} + \frac{2y}{x^2\sqrt{4yx^2 + 1}} \right) (xa_2 + ya_3 + a_1) \\ - \frac{xb_2 + yb_3 + b_1}{x\sqrt{4yx^2 + 1}} = 0 \end{aligned} \quad (5E)$$

Putting the above in normal form gives

$$\begin{aligned} \frac{-4b_2x^6\sqrt{4yx^2 + 1} + 4x^6b_2 - 8x^5ya_2 - 4x^5yb_3 - 16x^4y^2a_3 + 4x^5b_1 - 16x^4ya_1 - 4\sqrt{4yx^2 + 1}x^3a_2 - 2\sqrt{4yx^2 + 1}x^3a_3}{x^6\sqrt{4yx^2 + 1}} \\ = 0 \end{aligned}$$

Setting the numerator to zero gives

$$\begin{aligned} 4b_2x^6\sqrt{4yx^2 + 1} - 4x^6b_2 + 8x^5ya_2 + 4x^5yb_3 + 16x^4y^2a_3 \\ - 4x^5b_1 + 16x^4ya_1 + 4\sqrt{4yx^2 + 1}x^3a_2 + 2\sqrt{4yx^2 + 1}x^3b_3 \\ + 6\sqrt{4yx^2 + 1}x^2ya_3 - (4yx^2 + 1)^{\frac{3}{2}}a_3 + 6\sqrt{4yx^2 + 1}x^2a_1 \\ + 4x^3a_2 + 2x^3b_3 - 2x^2ya_3 + 6x^2a_1 - a_3\sqrt{4yx^2 + 1} - 2a_3 = 0 \end{aligned} \quad (6E)$$

Simplifying the above gives

$$\begin{aligned} 4b_2x^6\sqrt{4yx^2 + 1} - 4x^6b_2 - 8x^5ya_2 - 4x^5yb_3 - 8x^4y^2a_3 + 4(4yx^2 + 1)x^3a_2 \\ + 2(4yx^2 + 1)x^3b_3 + 6(4yx^2 + 1)x^2ya_3 - 4x^5b_1 - 8x^4ya_1 + 6(4yx^2 + 1)x^2a_1 \\ + 4\sqrt{4yx^2 + 1}x^3a_2 + 2\sqrt{4yx^2 + 1}x^3b_3 + 6\sqrt{4yx^2 + 1}x^2ya_3 \\ - (4yx^2 + 1)^{\frac{3}{2}}a_3 + 6\sqrt{4yx^2 + 1}x^2a_1 - 2(4yx^2 + 1)a_3 - a_3\sqrt{4yx^2 + 1} = 0 \end{aligned} \quad (6E)$$

Since the PDE has radicals, simplifying gives

$$\begin{aligned}
& 4b_2x^6\sqrt{4yx^2+1} - 4x^6b_2 + 8x^5ya_2 + 4x^5yb_3 + 16x^4y^2a_3 - 4x^5b_1 + 16x^4ya_1 \\
& + 4\sqrt{4yx^2+1}x^3a_2 + 2\sqrt{4yx^2+1}x^3b_3 + 2\sqrt{4yx^2+1}x^2ya_3 + 4x^3a_2 \\
& + 2x^3b_3 + 6\sqrt{4yx^2+1}x^2a_1 - 2x^2ya_3 + 6x^2a_1 - 2a_3\sqrt{4yx^2+1} - 2a_3 = 0
\end{aligned}$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y, \sqrt{4yx^2+1}\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2, \sqrt{4yx^2+1} = v_3\}$$

The above PDE (6E) now becomes

$$\begin{aligned}
& 4b_2v_1^6v_3 + 8v_1^5v_2a_2 + 16v_1^4v_2^2a_3 - 4v_1^6b_2 + 4v_1^5v_2b_3 + 16v_1^4v_2a_1 \\
& - 4v_1^5b_1 + 4v_3v_1^3a_2 + 2v_3v_1^2v_2a_3 + 2v_3v_1^3b_3 + 6v_3v_1^2a_1 \\
& + 4v_1^3a_2 - 2v_1^2v_2a_3 + 2v_1^3b_3 + 6v_1^2a_1 - 2a_3v_3 - 2a_3 = 0
\end{aligned} \tag{7E}$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2, v_3\}$$

Equation (7E) now becomes

$$\begin{aligned}
& 4b_2v_1^6v_3 - 4v_1^6b_2 + (8a_2 + 4b_3)v_1^5v_2 - 4v_1^5b_1 + 16v_1^4v_2^2a_3 \\
& + 16v_1^4v_2a_1 + (4a_2 + 2b_3)v_1^3v_3 + (4a_2 + 2b_3)v_1^3 + 2v_3v_1^2v_2a_3 \\
& - 2v_1^2v_2a_3 + 6v_3v_1^2a_1 + 6v_1^2a_1 - 2a_3v_3 - 2a_3 = 0
\end{aligned} \tag{8E}$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned}
 6a_1 &= 0 \\
 16a_1 &= 0 \\
 -2a_3 &= 0 \\
 2a_3 &= 0 \\
 16a_3 &= 0 \\
 -4b_1 &= 0 \\
 -4b_2 &= 0 \\
 4b_2 &= 0 \\
 4a_2 + 2b_3 &= 0 \\
 8a_2 + 4b_3 &= 0
 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned}
 a_1 &= 0 \\
 a_2 &= a_2 \\
 a_3 &= 0 \\
 b_1 &= 0 \\
 b_2 &= 0 \\
 b_3 &= -2a_2
 \end{aligned}$$

Substituting the above solution in the ansatz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned}
 \xi &= x \\
 \eta &= -2y
 \end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned}
 \eta &= \eta - \omega(x, y) \xi \\
 &= -2y - \left(\frac{1 + \sqrt{4y x^2 + 1}}{2x^3} \right) (x) \\
 &= \frac{-4y x^2 - \sqrt{4y x^2 + 1} - 1}{2x^2} \\
 \xi &= 0
 \end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{-4yx^2 - \sqrt{4yx^2 + 1} - 1}{2x^2}} dy \end{aligned}$$

Which results in

$$S = -\frac{\ln(y)}{2} - \operatorname{arctanh}\left(\sqrt{4yx^2 + 1}\right)$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{1 + \sqrt{4yx^2 + 1}}{2x^3}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{1}{x\sqrt{4yx^2 + 1}} \\ S_y &= \frac{-1 + \frac{1}{\sqrt{4yx^2 + 1}}}{2y} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$-\frac{\ln(y)}{2} - \operatorname{arctanh}\left(\sqrt{1+4yx^2}\right) = c_1$$

Which simplifies to

$$-\frac{\ln(y)}{2} - \operatorname{arctanh}\left(\sqrt{1+4yx^2}\right) = c_1$$

Summary

The solution(s) found are the following

$$-\frac{\ln(y)}{2} - \operatorname{arctanh}\left(\sqrt{1+4yx^2}\right) = c_1 \quad (1)$$

Verification of solutions

$$-\frac{\ln(y)}{2} - \operatorname{arctanh}\left(\sqrt{1+4yx^2}\right) = c_1$$

Verified OK.

Solving equation (2)

Writing the ode as

$$y' = -\frac{-1 + \sqrt{4yx^2 + 1}}{2x^3}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (A)$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \quad (1E)$$

$$\eta = xb_2 + yb_3 + b_1 \quad (2E)$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$b_2 - \frac{(-1 + \sqrt{4yx^2 + 1})(b_3 - a_2)}{2x^3} - \frac{(-1 + \sqrt{4yx^2 + 1})^2 a_3}{4x^6} - \left(-\frac{2y}{x^2 \sqrt{4yx^2 + 1}} + \frac{-\frac{3}{2} + \frac{3\sqrt{4yx^2 + 1}}{2}}{x^4} \right) (xa_2 + ya_3 + a_1) + \frac{xb_2 + yb_3 + b_1}{x\sqrt{4yx^2 + 1}} = 0 \quad (5E)$$

Putting the above in normal form gives

$$\frac{-4b_2x^6\sqrt{4yx^2 + 1} - 4x^6b_2 + 8x^5ya_2 + 4x^5yb_3 + 16x^4y^2a_3 - 4x^5b_1 + 16x^4ya_1 - 4\sqrt{4yx^2 + 1}x^3a_2 - 2\sqrt{4yx^2 + 1}x^3a_3}{x^6} = 0$$

Setting the numerator to zero gives

$$4b_2x^6\sqrt{4yx^2 + 1} + 4x^6b_2 - 8x^5ya_2 - 4x^5yb_3 - 16x^4y^2a_3 + 4x^5b_1 - 16x^4ya_1 + 4\sqrt{4yx^2 + 1}x^3a_2 + 2\sqrt{4yx^2 + 1}x^3b_3 + 6\sqrt{4yx^2 + 1}x^2ya_3 - (4yx^2 + 1)^{\frac{3}{2}}a_3 + 6\sqrt{4yx^2 + 1}x^2a_1 - 4x^3a_2 - 2x^3b_3 + 2x^2ya_3 - 6x^2a_1 - a_3\sqrt{4yx^2 + 1} + 2a_3 = 0 \quad (6E)$$

Simplifying the above gives

$$4b_2x^6\sqrt{4yx^2 + 1} + 4x^6b_2 + 8x^5ya_2 + 4x^5yb_3 + 8x^4y^2a_3 - 4(4yx^2 + 1)x^3a_2 - 2(4yx^2 + 1)x^3b_3 - 6(4yx^2 + 1)x^2ya_3 + 4x^5b_1 + 8x^4ya_1 - 6(4yx^2 + 1)x^2a_1 + 4\sqrt{4yx^2 + 1}x^3a_2 + 2\sqrt{4yx^2 + 1}x^3b_3 + 6\sqrt{4yx^2 + 1}x^2ya_3 - (4yx^2 + 1)^{\frac{3}{2}}a_3 + 6\sqrt{4yx^2 + 1}x^2a_1 + 2(4yx^2 + 1)a_3 - a_3\sqrt{4yx^2 + 1} = 0 \quad (6E)$$

Since the PDE has radicals, simplifying gives

$$\begin{aligned}
& 4b_2x^6\sqrt{4yx^2+1} + 4x^6b_2 - 8x^5ya_2 - 4x^5yb_3 - 16x^4y^2a_3 + 4x^5b_1 - 16x^4ya_1 \\
& + 4\sqrt{4yx^2+1}x^3a_2 + 2\sqrt{4yx^2+1}x^3b_3 + 2\sqrt{4yx^2+1}x^2ya_3 - 4x^3a_2 \\
& - 2x^3b_3 + 6\sqrt{4yx^2+1}x^2a_1 + 2x^2ya_3 - 6x^2a_1 - 2a_3\sqrt{4yx^2+1} + 2a_3 = 0
\end{aligned}$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y, \sqrt{4yx^2+1}\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2, \sqrt{4yx^2+1} = v_3\}$$

The above PDE (6E) now becomes

$$\begin{aligned}
& 4b_2v_1^6v_3 - 8v_1^5v_2a_2 - 16v_1^4v_2^2a_3 + 4v_1^6b_2 - 4v_1^5v_2b_3 - 16v_1^4v_2a_1 \\
& + 4v_1^5b_1 + 4v_3v_1^3a_2 + 2v_3v_1^2v_2a_3 + 2v_3v_1^3b_3 + 6v_3v_1^2a_1 \\
& - 4v_1^3a_2 + 2v_1^2v_2a_3 - 2v_1^3b_3 - 6v_1^2a_1 - 2a_3v_3 + 2a_3 = 0
\end{aligned} \tag{7E}$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2, v_3\}$$

Equation (7E) now becomes

$$\begin{aligned}
& 4b_2v_1^6v_3 + 4v_1^6b_2 + (-8a_2 - 4b_3)v_1^5v_2 + 4v_1^5b_1 - 16v_1^4v_2^2a_3 \\
& - 16v_1^4v_2a_1 + (4a_2 + 2b_3)v_1^3v_3 + (-4a_2 - 2b_3)v_1^3 \\
& + 2v_3v_1^2v_2a_3 + 2v_1^2v_2a_3 + 6v_3v_1^2a_1 - 6v_1^2a_1 - 2a_3v_3 + 2a_3 = 0
\end{aligned} \tag{8E}$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned}
 -16a_1 &= 0 \\
 -6a_1 &= 0 \\
 6a_1 &= 0 \\
 -16a_3 &= 0 \\
 -2a_3 &= 0 \\
 2a_3 &= 0 \\
 4b_1 &= 0 \\
 4b_2 &= 0 \\
 -8a_2 - 4b_3 &= 0 \\
 -4a_2 - 2b_3 &= 0 \\
 4a_2 + 2b_3 &= 0
 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned}
 a_1 &= 0 \\
 a_2 &= a_2 \\
 a_3 &= 0 \\
 b_1 &= 0 \\
 b_2 &= 0 \\
 b_3 &= -2a_2
 \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned}
 \xi &= x \\
 \eta &= -2y
 \end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned}
 \eta &= \eta - \omega(x, y) \xi \\
 &= -2y - \left(-\frac{-1 + \sqrt{4y x^2 + 1}}{2x^3} \right) (x) \\
 &= \frac{-4y x^2 + \sqrt{4y x^2 + 1} - 1}{2x^2} \\
 \xi &= 0
 \end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{-4yx^2 + \sqrt{4yx^2 + 1} - 1}{2x^2}} dy \end{aligned}$$

Which results in

$$S = -\frac{\ln(y)}{2} + \operatorname{arctanh}\left(\sqrt{4yx^2 + 1}\right)$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -\frac{-1 + \sqrt{4yx^2 + 1}}{2x^3}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= -\frac{1}{x\sqrt{4yx^2 + 1}} \\ S_y &= \frac{-\frac{1}{\sqrt{4yx^2 + 1}} - 1}{2y} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$-\frac{\ln(y)}{2} + \operatorname{arctanh}\left(\sqrt{1+4yx^2}\right) = c_1$$

Which simplifies to

$$-\frac{\ln(y)}{2} + \operatorname{arctanh}\left(\sqrt{1+4yx^2}\right) = c_1$$

Summary

The solution(s) found are the following

$$-\frac{\ln(y)}{2} + \operatorname{arctanh}\left(\sqrt{1+4yx^2}\right) = c_1 \quad (1)$$

Verification of solutions

$$-\frac{\ln(y)}{2} + \operatorname{arctanh}\left(\sqrt{1+4yx^2}\right) = c_1$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:
  *** Sublevel 2 ***
  Methods for first order ODEs:
  -> Solving 1st order ODE of high degree, 1st attempt
  trying 1st order WeierstrassP solution for high degree ODE
  trying 1st order WeierstrassPPrime solution for high degree ODE
  trying 1st order JacobiSN solution for high degree ODE
  trying 1st order ODE linearizable_by_differentiation
  trying differential order: 1; missing variables
  trying dAlembert
  trying simple symmetries for implicit equations
  Successful isolation of dy/dx: 2 solutions were found. Trying to solve each resulting ODE
    *** Sublevel 3 ***
    Methods for first order ODEs:
    --- Trying classification methods ---
    trying homogeneous types:
    trying homogeneous G
    <- homogeneous successful
    -----
  * Tackling next ODE.
    *** Sublevel 3 ***
    Methods for first order ODEs:
    --- Trying classification methods ---
    trying homogeneous types:
    trying homogeneous G
    <- homogeneous successful`
```

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 81

```
dsolve(y(x)+x*diff(y(x),x)=x^4*(diff(y(x),x))^2,y(x), singsol=all)
```

$$\begin{aligned}y(x) &= -\frac{1}{4x^2} \\y(x) &= \frac{-c_1 i - x}{c_1^2 x} \\y(x) &= \frac{c_1 i - x}{x c_1^2} \\y(x) &= \frac{c_1 i - x}{x c_1^2} \\y(x) &= \frac{-c_1 i - x}{c_1^2 x}\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.5 (sec). Leaf size: 123

```
DSolve[y[x]+x*y'[x]==x^4*(y'[x])^2,y[x],x,IncludeSingularSolutions -> True]
```

$$\begin{aligned}&\text{Solve} \left[\frac{x\sqrt{4x^2y(x)+1}\operatorname{arctanh}\left(\sqrt{4x^2y(x)+1}\right)}{\sqrt{4x^4y(x)+x^2}} - \frac{1}{2}\log(y(x)) = c_1, y(x) \right] \\&\text{Solve} \left[\frac{x\sqrt{4x^2y(x)+1}\operatorname{arctanh}\left(\sqrt{4x^2y(x)+1}\right)}{\sqrt{4x^4y(x)+x^2}} - \frac{1}{2}\log(y(x)) = c_1, y(x) \right] \\&y(x) \rightarrow 0\end{aligned}$$

1.12 problem 1(m)

1.12.1 Solving as homogeneousTypeD2 ode	112
1.12.2 Solving as first order ode lie symmetry calculated ode	114
1.12.3 Solving as exact ode	119

Internal problem ID [6116]

Internal file name [OUTPUT/5364_Sunday_June_05_2022_03_35_28_PM_30647876/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.2 THE NATURE OF SOLUTIONS. Page 9

Problem number: 1(m).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exactByInspection", "homogeneousTypeD2", "first_order_ode_lie_symmetry_calculated"

Maple gives the following as the ode type

```
[[_homogeneous, `class A`], _rational, [_Abel, `2nd type`, `class B`]]
```

$$y' - \frac{y^2}{xy - x^2} = 0$$

1.12.1 Solving as homogeneousTypeD2 ode

Using the change of variables $y = u(x)x$ on the above ode results in new ode in $u(x)$

$$u'(x)x + u(x) - \frac{u(x)^2 x^2}{x^2 u(x) - x^2} = 0$$

In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= \frac{u}{x(u-1)} \end{aligned}$$

Where $f(x) = \frac{1}{x}$ and $g(u) = \frac{u}{u-1}$. Integrating both sides gives

$$\frac{1}{\frac{u}{u-1}} du = \frac{1}{x} dx$$

$$\int \frac{1}{\frac{u}{u-1}} du = \int \frac{1}{x} dx$$

$$u - \ln(u) = \ln(x) + c_2$$

The solution is

$$u(x) - \ln(u(x)) - \ln(x) - c_2 = 0$$

Replacing $u(x)$ in the above solution by $\frac{y}{x}$ results in the solution for y in implicit form

$$\frac{y}{x} - \ln\left(\frac{y}{x}\right) - \ln(x) - c_2 = 0$$

$$\frac{y}{x} - \ln\left(\frac{y}{x}\right) - \ln(x) - c_2 = 0$$

Summary

The solution(s) found are the following

$$\frac{y}{x} - \ln\left(\frac{y}{x}\right) - \ln(x) - c_2 = 0 \tag{1}$$




Figure 26: Slope field plot

Verification of solutions

$$\frac{y}{x} - \ln\left(\frac{y}{x}\right) - \ln(x) - c_2 = 0$$

Verified OK.

1.12.2 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = \frac{y^2}{x(-x+y)}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \quad (\text{1E})$$

$$\eta = xb_2 + yb_3 + b_1 \quad (\text{2E})$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$b_2 + \frac{y^2(b_3 - a_2)}{x(-x+y)} - \frac{y^4 a_3}{x^2(-x+y)^2}$$
$$- \left(-\frac{y^2}{x^2(-x+y)} + \frac{y^2}{x(-x+y)^2} \right) (xa_2 + ya_3 + a_1) \quad (\text{5E})$$
$$- \left(\frac{2y}{x(-x+y)} - \frac{y^2}{x(-x+y)^2} \right) (xb_2 + yb_3 + b_1) = 0$$

Putting the above in normal form gives

$$\frac{x^4 b_2 - x^2 y^2 a_2 + x^2 y^2 b_3 - 2x y^3 a_3 + 2x^2 y b_1 - 2x y^2 a_1 - x y^2 b_1 + y^3 a_1}{x^2 (x - y)^2} = 0$$

Setting the numerator to zero gives

$$x^4b_2 - x^2y^2a_2 + x^2y^2b_3 - 2xy^3a_3 + 2x^2yb_1 - 2xy^2a_1 - xy^2b_1 + y^3a_1 = 0 \quad (6E)$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2\}$$

The above PDE (6E) now becomes

$$-a_2v_1^2v_2^2 - 2a_3v_1v_2^3 + b_2v_1^4 + b_3v_1^2v_2^2 - 2a_1v_1v_2^2 + a_1v_2^3 + 2b_1v_1^2v_2 - b_1v_1v_2^2 = 0 \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2\}$$

Equation (7E) now becomes

$$b_2v_1^4 + (b_3 - a_2)v_1^2v_2^2 + 2b_1v_1^2v_2 - 2a_3v_1v_2^3 + (-2a_1 - b_1)v_1v_2^2 + a_1v_2^3 = 0 \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} a_1 &= 0 \\ b_2 &= 0 \\ -2a_3 &= 0 \\ 2b_1 &= 0 \\ -2a_1 - b_1 &= 0 \\ b_3 - a_2 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= 0 \\ a_2 &= b_3 \\ a_3 &= 0 \\ b_1 &= 0 \\ b_2 &= 0 \\ b_3 &= b_3 \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned}\xi &= x \\ \eta &= y\end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned}\eta &= \eta - \omega(x, y) \xi \\ &= y - \left(\frac{y^2}{x(-x + y)} \right) (x) \\ &= \frac{yx}{x - y} \\ \xi &= 0\end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y} \right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{yx}{x-y}} dy\end{aligned}$$

Which results in

$$S = \ln(y) - \frac{y}{x}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y) S_y}{R_x + \omega(x, y) R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{y^2}{x(-x+y)}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{y}{x^2} \\ S_y &= \frac{x-y}{xy} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \tag{2A}$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \tag{4}$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{x \ln(y) - y}{x} = c_1$$



Which simplifies to

$$\frac{x \ln(y) - y}{x} = c_1$$

Which gives

$$y = e^{-\text{LambertW}\left(-\frac{e^{c_1}}{x}\right) + c_1}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{y^2}{x(-x+y)}$ 	$R = x$ $S = \frac{\ln(y) x - y}{x}$	$\frac{dS}{dR} = 0$ 

Summary

The solution(s) found are the following

$$y = e^{-\text{LambertW}\left(-\frac{e^{c_1}}{x}\right) + c_1} \tag{1}$$




Figure 27: Slope field plot

Verification of solutions

$$y = e^{-\text{LambertW}\left(-\frac{e^{c_1}}{x}\right) + c_1}$$

Verified OK.

1.12.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned}(x(-x + y)) dy &= (y^2) dx \\ (-y^2) dx + (x(-x + y)) dy &= 0\end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -y^2 \\ N(x, y) &= x(-x + y)\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-y^2) \\ &= -2y\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(x(-x + y)) \\ &= -2x + y\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. By inspection $\frac{1}{x^2y}$ is an integrating factor. Therefore by multiplying $M = -y^2$ and $N = x(-x + y)$ by this integrating factor the ode becomes exact. The new M, N are

$$M = -\frac{y}{x^2}$$

$$N = \frac{-x + y}{xy}$$

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\frac{\partial \phi}{\partial x} = M$$

$$\frac{\partial \phi}{\partial y} = N$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} \left(\frac{-x+y}{xy}\right) dy &= \left(\frac{y}{x^2}\right) dx \\ \left(-\frac{y}{x^2}\right) dx + \left(\frac{-x+y}{xy}\right) dy &= 0 \end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -\frac{y}{x^2} \\ N(x, y) &= \frac{-x+y}{xy} \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(-\frac{y}{x^2}\right) \\ &= -\frac{1}{x^2} \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{-x+y}{xy}\right) \\ &= -\frac{1}{x^2} \end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \tag{1}$$

$$\frac{\partial \phi}{\partial y} = N \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned} \int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\frac{y}{x^2} dx \\ \phi &= \frac{y}{x} + f(y) \end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial\phi}{\partial y} = \frac{1}{x} + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial\phi}{\partial y} = \frac{-x+y}{xy}$. Therefore equation (4) becomes

$$\frac{-x+y}{xy} = \frac{1}{x} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = -\frac{1}{y}$$

Integrating the above w.r.t y gives

$$\int f'(y) dy = \int \left(-\frac{1}{y}\right) dy$$

$$f(y) = -\ln(y) + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = \frac{y}{x} - \ln(y) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \frac{y}{x} - \ln(y)$$

The solution becomes

$$y = e^{-\text{LambertW}\left(-\frac{e^{-c_1}}{x}\right) - c_1}$$

Summary

The solution(s) found are the following

$$y = e^{-\text{LambertW}\left(-\frac{e^{-c_1}}{x}\right) - c_1} \quad (1)$$




Figure 28: Slope field plot

Verification of solutions

$$y = e^{-\text{LambertW}\left(-\frac{e^{-c_1}}{x}\right) - c_1}$$

Verified OK.

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying homogeneous D
<- homogeneous successful`

```

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 17

```
dsolve(diff(y(x),x)=y(x)^2/(x*y(x)-x^2),y(x), singsol=all)
```

$$y(x) = -x \operatorname{LambertW}\left(-\frac{e^{-c_1}}{x}\right)$$

✓ Solution by Mathematica

Time used: 2.335 (sec). Leaf size: 25

```
DSolve[y'[x]==y[x]^2/(x*y[x]-x^2),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -xW\left(-\frac{e^{-c_1}}{x}\right)$$
$$y(x) \rightarrow 0$$

1.13 problem 1(n)

- 1.13.1 Solving as first order ode lie symmetry calculated ode 126
- 1.13.2 Solving as exact ode 132

Internal problem ID [6117]

Internal file name [OUTPUT/5365_Sunday_June_05_2022_03_35_29_PM_26351614/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.2 THE NATURE OF SOLUTIONS. Page 9

Problem number: 1(n).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exactWithIntegrationFactor", "first_order_ode_lie_symmetry_calculated"**

Maple gives the following as the ode type

```
[[_1st_order , _with_linear_symmetries]]
```

$$(y \cos(y) - \sin(y) + x)y' - y = 0$$

1.13.1 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = \frac{y}{y \cos(y) - \sin(y) + x}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \tag{A}$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \tag{1E}$$

$$\eta = xb_2 + yb_3 + b_1 \tag{2E}$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$\begin{aligned} b_2 + \frac{y(b_3 - a_2)}{y \cos(y) - \sin(y) + x} \\ - \frac{y^2 a_3}{(y \cos(y) - \sin(y) + x)^2} + \frac{y(xa_2 + ya_3 + a_1)}{(y \cos(y) - \sin(y) + x)^2} \\ - \left(\frac{1}{y \cos(y) - \sin(y) + x} + \frac{y^2 \sin(y)}{(y \cos(y) - \sin(y) + x)^2} \right) (xb_2 + yb_3 + b_1) = 0 \end{aligned} \quad (5E)$$

Putting the above in normal form gives

$$\frac{\cos(y)^2 y^2 b_2 - \sin(y) x y^2 b_2 - \sin(y) y^3 b_3 - 2 \cos(y) \sin(y) y b_2 + \cos(y) x y b_2 - \cos(y) y^2 a_2 - \sin(y) y^2 b_1}{(y \cos(y) - \sin(y) + x)^2} = 0$$

Setting the numerator to zero gives

$$\begin{aligned} \cos(y)^2 y^2 b_2 - \sin(y) x y^2 b_2 - \sin(y) y^3 b_3 - 2 \cos(y) \sin(y) y b_2 \\ + \cos(y) x y b_2 - \cos(y) y^2 a_2 - \sin(y) y^2 b_1 - \cos(y) y b_1 \\ + \sin(y)^2 b_2 - \sin(y) x b_2 + \sin(y) y a_2 + \sin(y) b_1 - x b_1 + y a_1 = 0 \end{aligned} \quad (6E)$$

Simplifying the above gives

$$\begin{aligned} -x b_1 + y a_1 + \frac{y^2 b_2}{2} + \frac{b_2}{2} + \frac{y^2 b_2 \cos(2y)}{2} - \sin(y) x y^2 b_2 - \sin(y) y^3 b_3 \\ - y b_2 \sin(2y) + \cos(y) x y b_2 - \cos(y) y^2 a_2 - \sin(y) y^2 b_1 \\ - \cos(y) y b_1 - \frac{b_2 \cos(2y)}{2} - \sin(y) x b_2 + \sin(y) y a_2 + \sin(y) b_1 = 0 \end{aligned} \quad (6E)$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y, \cos(y), \cos(2y), \sin(y), \sin(2y)\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2, \cos(y) = v_3, \cos(2y) = v_4, \sin(y) = v_5, \sin(2y) = v_6\}$$

The above PDE (6E) now becomes

$$\begin{aligned} -v_1 b_1 + v_2 a_1 + \frac{1}{2} v_2^2 b_2 + \frac{1}{2} b_2 + \frac{1}{2} v_2^2 b_2 v_4 - v_5 v_1 v_2^2 b_2 - v_5 v_2^3 b_3 - v_2 b_2 v_6 \\ + v_3 v_1 v_2 b_2 - v_3 v_2^2 a_2 - v_5 v_2^2 b_1 - v_3 v_2 b_1 - \frac{1}{2} b_2 v_4 - v_5 v_1 b_2 + v_5 v_2 a_2 + v_5 b_1 = 0 \end{aligned} \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2, v_3, v_4, v_5, v_6\}$$

Equation (7E) now becomes

$$\begin{aligned} -v_1 b_1 + v_2 a_1 + \frac{1}{2} v_2^2 b_2 + \frac{1}{2} b_2 + \frac{1}{2} v_2^2 b_2 v_4 - v_5 v_1 v_2^2 b_2 - v_5 v_2^3 b_3 - v_2 b_2 v_6 \\ + v_3 v_1 v_2 b_2 - v_3 v_2^2 a_2 - v_5 v_2^2 b_1 - v_3 v_2 b_1 - \frac{1}{2} b_2 v_4 - v_5 v_1 b_2 + v_5 v_2 a_2 + v_5 b_1 = 0 \end{aligned} \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} a_1 &= 0 \\ a_2 &= 0 \\ b_1 &= 0 \\ b_2 &= 0 \\ -a_2 &= 0 \\ -b_1 &= 0 \\ -b_2 &= 0 \\ -\frac{b_2}{2} &= 0 \\ \frac{b_2}{2} &= 0 \\ -b_3 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$a_1 = 0$$

$$a_2 = 0$$

$$a_3 = a_3$$

$$b_1 = 0$$

$$b_2 = 0$$

$$b_3 = 0$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\xi = y$$

$$\eta = 0$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned} \eta &= \eta - \omega(x, y) \xi \\ &= 0 - \left(\frac{y}{y \cos(y) - \sin(y) + x} \right) (y) \\ &= - \frac{y^2}{y \cos(y) - \sin(y) + x} \\ \xi &= 0 \end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y} \right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{-\frac{y^2}{y \cos(y) - \sin(y) + x}} dy \end{aligned}$$

Which results in

$$S = -\frac{\sin(y)}{y} + \frac{x}{y}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{y}{y \cos(y) - \sin(y) + x}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{1}{y} \\ S_y &= \frac{-y \cos(y) + \sin(y) - x}{y^2} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \quad (4)$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{-\sin(y) + x}{y} = c_1$$

Which simplifies to

$$\frac{-\sin(y) + x}{y} = c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{y}{y \cos(y) - \sin(y) + x}$ 	$R = x$ $S = \frac{-\sin(y) + x}{y}$	$\frac{dS}{dR} = 0$ 

Summary

The solution(s) found are the following

$$\frac{-\sin(y) + x}{y} = c_1 \tag{1}$$




Figure 29: Slope field plot

Verification of solutions

$$\frac{-\sin(y) + x}{y} = c_1$$

Verified OK.

1.13.2 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned}(y \cos(y) - \sin(y) + x) dy &= (y) dx \\ (-y) dx + (y \cos(y) - \sin(y) + x) dy &= 0\end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -y \\ N(x, y) &= y \cos(y) - \sin(y) + x\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-y) \\ &= -1\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(y \cos(y) - \sin(y) + x) \\ &= 1\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= \frac{1}{y \cos(y) - \sin(y) + x} ((-1) - (1)) \\ &= -\frac{2}{y \cos(y) - \sin(y) + x} \end{aligned}$$

Since A depends on y , it can not be used to obtain an integrating factor. We will now try a second method to find an integrating factor. Let

$$\begin{aligned} B &= \frac{1}{M} \left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) \\ &= -\frac{1}{y} ((1) - (-1)) \\ &= -\frac{2}{y} \end{aligned}$$

Since B does not depend on x , it can be used to obtain an integrating factor. Let the integrating factor be μ . Then

$$\begin{aligned} \mu &= e^{\int B \, dy} \\ &= e^{\int -\frac{2}{y} \, dy} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^{-2 \ln(y)} \\ &= \frac{1}{y^2} \end{aligned}$$

M and N are now multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} so not to confuse them with the original M and N .

$$\begin{aligned} \bar{M} &= \mu M \\ &= \frac{1}{y^2} (-y) \\ &= -\frac{1}{y} \end{aligned}$$

And

$$\begin{aligned}\bar{N} &= \mu N \\ &= \frac{1}{y^2}(y \cos(y) - \sin(y) + x) \\ &= \frac{y \cos(y) - \sin(y) + x}{y^2}\end{aligned}$$

So now a modified ODE is obtained from the original ODE which will be exact and can be solved using the standard method. The modified ODE is

$$\begin{aligned}\bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ \left(-\frac{1}{y}\right) + \left(\frac{y \cos(y) - \sin(y) + x}{y^2}\right) \frac{dy}{dx} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \quad (1)$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\frac{1}{y} dx \\ \phi &= -\frac{x}{y} + f(y)\end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = \frac{x}{y^2} + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{y \cos(y) - \sin(y) + x}{y^2}$. Therefore equation (4) becomes

$$\frac{y \cos(y) - \sin(y) + x}{y^2} = \frac{x}{y^2} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = \frac{y \cos(y) - \sin(y)}{y^2}$$

Integrating the above w.r.t y gives

$$\int f'(y) dy = \int \left(\frac{y \cos(y) - \sin(y)}{y^2} \right) dy$$

$$f(y) = \frac{\sin(y)}{y} + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\frac{x}{y} + \frac{\sin(y)}{y} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\frac{x}{y} + \frac{\sin(y)}{y}$$

Summary

The solution(s) found are the following

$$-\frac{x}{y} + \frac{\sin(y)}{y} = c_1 \tag{1}$$




Figure 30: Slope field plot

Verification of solutions

$$-\frac{x}{y} + \frac{\sin(y)}{y} = c_1$$

Verified OK.

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
<- 1st order linear successful
<- inverse linear successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
dsolve((y(x)*cos(y(x))-sin(y(x))+x)*diff(y(x),x)=y(x),y(x), singsol=all)
```

$$x - c_1 y(x) - \sin(y(x)) = 0$$

✓ Solution by Mathematica

Time used: 0.245 (sec). Leaf size: 14

```
DSolve[(y[x]*Cos[y[x]]-Sin[y[x]]+x)*y'[x]==y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve}[x = \sin(y(x)) + c_1 y(x), y(x)]$$

1.14 problem 1(o)

1.14.1 Solving as quadrature ode	139
1.14.2 Maple step by step solution	140

Internal problem ID [6118]

Internal file name [OUTPUT/5366_Sunday_June_05_2022_03_35_32_PM_75099697/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.2 THE NATURE OF SOLUTIONS. Page 9

Problem number: 1(o).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

`[_quadrature]`

$$y^2 + y'y^2 = -1$$

1.14.1 Solving as quadrature ode

Integrating both sides gives

$$\int -\frac{y^2}{y^2 + 1} dy = \int dx$$
$$\int -\frac{-a^2}{-a^2 + 1} d_a = x + c_1$$

Summary

The solution(s) found are the following

$$\int -\frac{-a^2}{-a^2 + 1} d_a = x + c_1 \tag{1}$$




Figure 31: Slope field plot

Verification of solutions

$$\int^y -\frac{a^2}{-a^2 + 1} da = x + c_1$$

Verified OK.

1.14.2 Maple step by step solution

Let's solve

$$y^2 + y'y^2 = -1$$

- Highest derivative means the order of the ODE is 1

y'

- Separate variables

$$\frac{y'y^2}{-1-y^2} = 1$$

- Integrate both sides with respect to x

$$\int \frac{y'y^2}{-1-y^2} dx = \int 1 dx + c_1$$

- Evaluate integral

$$-y + \arctan(y) = x + c_1$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
<- separable successful`

```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 14

```
dsolve(1+y(x)^2+y(x)^2*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \tan(\text{RootOf}(-_Z + x + c_1 + \tan(_Z)))$$

✓ Solution by Mathematica

Time used: 0.186 (sec). Leaf size: 35

```
DSolve[1+y[x]^2+y[x]^2*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\begin{aligned}
 y(x) &\rightarrow \text{InverseFunction}[\#1 - \arctan(\#1)\&][-x + c_1] \\
 y(x) &\rightarrow -i \\
 y(x) &\rightarrow i
 \end{aligned}$$

1.15 problem 2(a)

1.15.1 Solving as quadrature ode	142
1.15.2 Maple step by step solution	143

Internal problem ID [6119]

Internal file name [OUTPUT/5367_Sunday_June_05_2022_03_35_33_PM_12539902/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.2 THE NATURE OF SOLUTIONS. Page 9

Problem number: 2(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

`[_quadrature]`

$$y' = e^{3x} - x$$

1.15.1 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned} y &= \int e^{3x} - x \, dx \\ &= -\frac{x^2}{2} + \frac{e^{3x}}{3} + c_1 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = -\frac{x^2}{2} + \frac{e^{3x}}{3} + c_1 \tag{1}$$




Figure 32: Slope field plot

Verification of solutions

$$y = -\frac{x^2}{2} + \frac{e^{3x}}{3} + c_1$$

Verified OK.

1.15.2 Maple step by step solution

Let's solve

$$y' = e^{3x} - x$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Integrate both sides with respect to x

$$\int y' dx = \int (e^{3x} - x) dx + c_1$$

- Evaluate integral

$$y = -\frac{x^2}{2} + \frac{e^{3x}}{3} + c_1$$

- Solve for y

$$y = -\frac{x^2}{2} + \frac{e^{3x}}{3} + c_1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x)=exp(3*x)-x,y(x), singsol=all)
```

$$y(x) = -\frac{x^2}{2} + \frac{e^{3x}}{3} + c_1$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 24

```
DSolve[y'[x]==Exp[3*x]-x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{x^2}{2} + \frac{e^{3x}}{3} + c_1$$

1.16 problem 2(b)

1.16.1 Solving as quadrature ode	145
1.16.2 Maple step by step solution	146

Internal problem ID [6120]

Internal file name [OUTPUT/5368_Sunday_June_05_2022_03_35_34_PM_99531044/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.2 THE NATURE OF SOLUTIONS. Page 9

Problem number: 2(b).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

`[_quadrature]`

$$y' = x e^{x^2}$$

1.16.1 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned} y &= \int x e^{x^2} dx \\ &= \frac{e^{x^2}}{2} + c_1 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{e^{x^2}}{2} + c_1 \tag{1}$$




Figure 33: Slope field plot

Verification of solutions

$$y = \frac{e^{x^2}}{2} + c_1$$

Verified OK.

1.16.2 Maple step by step solution

Let's solve

$$y' = x e^{x^2}$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Integrate both sides with respect to x

$$\int y' dx = \int x e^{x^2} dx + c_1$$

- Evaluate integral

$$y = \frac{e^{x^2}}{2} + c_1$$

- Solve for y

$$y = \frac{e^{x^2}}{2} + c_1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)=x*exp(x^2),y(x), singsol=all)
```

$$y(x) = \frac{e^{x^2}}{2} + c_1$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 17

```
DSolve[y'[x]==x*Exp[x^2],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{x^2}}{2} + c_1$$

1.17 problem 2(c)

1.17.1 Solving as quadrature ode	148
1.17.2 Maple step by step solution	149

Internal problem ID [6121]

Internal file name [OUTPUT/5369_Sunday_June_05_2022_03_35_35_PM_66222368/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.2 THE NATURE OF SOLUTIONS. Page 9

Problem number: 2(c).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

`[_quadrature]`

$$(1 + x)y' = x$$

1.17.1 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned}y &= \int \frac{x}{1+x} dx \\ &= x - \ln(1+x) + c_1\end{aligned}$$

Summary

The solution(s) found are the following

$$y = x - \ln(1+x) + c_1 \tag{1}$$




Figure 34: Slope field plot

Verification of solutions

$$y = x - \ln(1 + x) + c_1$$

Verified OK.

1.17.2 Maple step by step solution

Let's solve

$$(1 + x)y' = x$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$y' = \frac{x}{1+x}$$

- Integrate both sides with respect to x

$$\int y'dx = \int \frac{x}{1+x} dx + c_1$$

- Evaluate integral

$$y = x - \ln(1 + x) + c_1$$

- Solve for y

$$y = x - \ln(1 + x) + c_1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve((1+x)*diff(y(x),x)=x,y(x), singsol=all)
```

$$y(x) = x - \ln(x + 1) + c_1$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 15

```
DSolve[(1+x)*y'[x]==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x - \log(x + 1) + c_1$$

1.18 problem 2(d)

1.18.1 Solving as quadrature ode	151
1.18.2 Maple step by step solution	152

Internal problem ID [6122]

Internal file name [OUTPUT/5370_Sunday_June_05_2022_03_35_36_PM_57886836/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.2 THE NATURE OF SOLUTIONS. Page 9

Problem number: 2(d).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

`[_quadrature]`

$$(x^2 + 1) y' = x$$

1.18.1 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned} y &= \int \frac{x}{x^2 + 1} dx \\ &= \frac{\ln(x^2 + 1)}{2} + c_1 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{\ln(x^2 + 1)}{2} + c_1 \tag{1}$$




Figure 35: Slope field plot

Verification of solutions

$$y = \frac{\ln(x^2 + 1)}{2} + c_1$$

Verified OK.

1.18.2 Maple step by step solution

Let's solve

$$(x^2 + 1)y' = x$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$y' = \frac{x}{x^2+1}$$

- Integrate both sides with respect to x

$$\int y' dx = \int \frac{x}{x^2+1} dx + c_1$$

- Evaluate integral

$$y = \frac{\ln(x^2+1)}{2} + c_1$$

- Solve for y

$$y = \frac{\ln(x^2+1)}{2} + c_1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 14

```
dsolve((1+x^2)*diff(y(x),x)=x,y(x), singsol=all)
```

$$y(x) = \frac{\ln(x^2 + 1)}{2} + c_1$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 18

```
DSolve[(1+x^2)*y'[x]==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} \log(x^2 + 1) + c_1$$

1.19 problem 2(e)

1.19.1 Solving as quadrature ode	154
1.19.2 Maple step by step solution	155

Internal problem ID [6123]

Internal file name [OUTPUT/5371_Sunday_June_05_2022_03_35_37_PM_74157158/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.2 THE NATURE OF SOLUTIONS. Page 9

Problem number: 2(e).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

`[_quadrature]`

$$(x^2 + 1) y' = \arctan(x)$$

1.19.1 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned} y &= \int \frac{\arctan(x)}{x^2 + 1} dx \\ &= \frac{\arctan(x)^2}{2} + c_1 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{\arctan(x)^2}{2} + c_1 \tag{1}$$




Figure 36: Slope field plot

Verification of solutions

$$y = \frac{\arctan(x)^2}{2} + c_1$$

Verified OK.

1.19.2 Maple step by step solution

Let's solve

$$(x^2 + 1)y' = \arctan(x)$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$y' = \frac{\arctan(x)}{x^2+1}$$

- Integrate both sides with respect to x

$$\int y'dx = \int \frac{\arctan(x)}{x^2+1} dx + c_1$$

- Evaluate integral

$$y = \frac{\arctan(x)^2}{2} + c_1$$

- Solve for y

$$y = \frac{\arctan(x)^2}{2} + c_1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve((1+x^2)*diff(y(x),x)=arctan(x),y(x), singsol=all)
```

$$y(x) = \frac{\arctan(x)^2}{2} + c_1$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 16

```
DSolve[(1+x^2)*y'[x]==ArcTan[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\arctan(x)^2}{2} + c_1$$

1.20 problem 2(f)

1.20.1 Solving as quadrature ode	157
1.20.2 Maple step by step solution	158

Internal problem ID [6124]

Internal file name [OUTPUT/5372_Sunday_June_05_2022_03_35_38_PM_11524109/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.2 THE NATURE OF SOLUTIONS. Page 9

Problem number: 2(f).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

`[_quadrature]`

$$xy' = 1$$

1.20.1 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned}y &= \int \frac{1}{x} dx \\ &= \ln(x) + c_1\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \ln(x) + c_1 \tag{1}$$




Figure 37: Slope field plot

Verification of solutions

$$y = \ln(x) + c_1$$

Verified OK.

1.20.2 Maple step by step solution

Let's solve

$$xy' = 1$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$y' = \frac{1}{x}$$

- Integrate both sides with respect to x

$$\int y' dx = \int \frac{1}{x} dx + c_1$$

- Evaluate integral

- $y = \ln(x) + c_1$
 • Solve for y
 $y = \ln(x) + c_1$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
<- quadrature successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 8

```
dsolve(x*diff(y(x),x)=1,y(x), singsol=all)
```

$$y(x) = \ln(x) + c_1$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 10

```
DSolve[x*y'[x]==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \log(x) + c_1$$

1.21 problem 2(g)

1.21.1 Solving as quadrature ode	160
1.21.2 Maple step by step solution	161

Internal problem ID [6125]

Internal file name [OUTPUT/5373_Sunday_June_05_2022_03_35_39_PM_21493951/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.2 THE NATURE OF SOLUTIONS. Page 9

Problem number: 2(g).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

`[_quadrature]`

$$y' = \arcsin(x)$$

1.21.1 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned} y &= \int \arcsin(x) \, dx \\ &= x \arcsin(x) + \sqrt{-x^2 + 1} + c_1 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = x \arcsin(x) + \sqrt{-x^2 + 1} + c_1 \tag{1}$$




Figure 38: Slope field plot

Verification of solutions

$$y = x \arcsin(x) + \sqrt{-x^2 + 1} + c_1$$

Verified OK.

1.21.2 Maple step by step solution

Let's solve

$$y' = \arcsin(x)$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Integrate both sides with respect to x

$$\int y' dx = \int \arcsin(x) dx + c_1$$

- Evaluate integral

$$y = x \arcsin(x) + \sqrt{-x^2 + 1} + c_1$$

- Solve for y

$$y = x \arcsin(x) + \sqrt{-x^2 + 1} + c_1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(diff(y(x),x)=arcsin(x),y(x), singsol=all)
```

$$y(x) = x \arcsin(x) + \sqrt{-x^2 + 1} + c_1$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 23

```
DSolve[y'[x]==ArcSin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x \arcsin(x) + \sqrt{1 - x^2} + c_1$$

1.22 problem 2(h)

1.22.1 Solving as quadrature ode	163
1.22.2 Maple step by step solution	164

Internal problem ID [6126]

Internal file name [OUTPUT/5374_Sunday_June_05_2022_03_35_40_PM_19514843/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.2 THE NATURE OF SOLUTIONS. Page 9

Problem number: 2(h).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

`[_quadrature]`

$$\sin(x) y' = 1$$

1.22.1 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned} y &= \int \frac{1}{\sin(x)} dx \\ &= \ln(\csc(x) - \cot(x)) + c_1 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \ln(\csc(x) - \cot(x)) + c_1 \tag{1}$$




Figure 39: Slope field plot

Verification of solutions

$$y = \ln(\csc(x) - \cot(x)) + c_1$$

Verified OK.

1.22.2 Maple step by step solution

Let's solve

$$\sin(x) y' = 1$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$y' = \frac{1}{\sin(x)}$$

- Integrate both sides with respect to x

$$\int y' dx = \int \frac{1}{\sin(x)} dx + c_1$$

- Evaluate integral

$$y = \ln(\csc(x) - \cot(x)) + c_1$$

- Solve for y

$$y = \ln(\csc(x) - \cot(x)) + c_1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(sin(x)*diff(y(x),x)=1,y(x), singsol=all)
```

$$y(x) = -\ln(\csc(x) + \cot(x)) + c_1$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 13

```
DSolve[Sin[x]*y'[x]==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\operatorname{arctanh}(\cos(x)) + c_1$$

1.23 problem 2(i)

1.23.1 Solving as quadrature ode	166
1.23.2 Maple step by step solution	167

Internal problem ID [6127]

Internal file name [OUTPUT/5375_Sunday_June_05_2022_03_35_41_PM_50895992/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.2 THE NATURE OF SOLUTIONS. Page 9

Problem number: 2(i).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"quadrature"**

Maple gives the following as the ode type

`[_quadrature]`

$$(x^3 + 1) y' = x$$

1.23.1 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned} y &= \int \frac{x}{x^3 + 1} dx \\ &= \frac{\ln(x^2 - x + 1)}{6} + \frac{\sqrt{3} \arctan\left(\frac{(2x-1)\sqrt{3}}{3}\right)}{3} - \frac{\ln(1+x)}{3} + c_1 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{\ln(x^2 - x + 1)}{6} + \frac{\sqrt{3} \arctan\left(\frac{(2x-1)\sqrt{3}}{3}\right)}{3} - \frac{\ln(1+x)}{3} + c_1 \quad (1)$$




Figure 40: Slope field plot

Verification of solutions

$$y = \frac{\ln(x^2 - x + 1)}{6} + \frac{\sqrt{3} \arctan\left(\frac{(2x-1)\sqrt{3}}{3}\right)}{3} - \frac{\ln(1+x)}{3} + c_1$$

Verified OK.

1.23.2 Maple step by step solution

Let's solve

$$(x^3 + 1)y' = x$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$y' = \frac{x}{x^3+1}$$

- Integrate both sides with respect to x

$$\int y' dx = \int \frac{x}{x^3+1} dx + c_1$$

- Evaluate integral

$$y = \frac{\ln(x^2-x+1)}{6} + \frac{\sqrt{3} \arctan\left(\frac{(2x-1)\sqrt{3}}{3}\right)}{3} - \frac{\ln(1+x)}{3} + c_1$$

- Solve for y

$$y = \frac{\ln(x^2-x+1)}{6} + \frac{\sqrt{3} \arctan\left(\frac{(2x-1)\sqrt{3}}{3}\right)}{3} - \frac{\ln(1+x)}{3} + c_1$$

Maple trace

```
`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 39

```
dsolve((1+x^3)*diff(y(x),x)=x,y(x), singsol=all)
```

$$y(x) = \frac{\ln(x^2 - x + 1)}{6} + \frac{\sqrt{3} \arctan\left(\frac{(2x-1)\sqrt{3}}{3}\right)}{3} - \frac{\ln(x + 1)}{3} + c_1$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 48

```
DSolve[(1+x^3)*y'[x]==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{6} \left(2\sqrt{3} \arctan\left(\frac{2x-1}{\sqrt{3}}\right) + \log(x^2 - x + 1) - 2\log(x + 1) + 6c_1 \right)$$

1.24 problem 2(j)

1.24.1 Solving as quadrature ode	169
1.24.2 Maple step by step solution	170

Internal problem ID [6128]

Internal file name [OUTPUT/5376_Sunday_June_05_2022_03_35_43_PM_51844411/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.2 THE NATURE OF SOLUTIONS. Page 9

Problem number: 2(j).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[**_quadrature**]

$$(x^2 - 3x + 2) y' = x$$

1.24.1 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned} y &= \int \frac{x}{x^2 - 3x + 2} dx \\ &= -\ln(x - 1) + 2 \ln(-2 + x) + c_1 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = -\ln(x - 1) + 2 \ln(-2 + x) + c_1 \tag{1}$$




Figure 41: Slope field plot

Verification of solutions

$$y = -\ln(x-1) + 2\ln(-2+x) + c_1$$

Verified OK.

1.24.2 Maple step by step solution

Let's solve

$$(x^2 - 3x + 2)y' = x$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$y' = \frac{x}{x^2 - 3x + 2}$$

- Integrate both sides with respect to x

$$\int y' dx = \int \frac{x}{x^2 - 3x + 2} dx + c_1$$

- Evaluate integral

$$y = -\ln(x - 1) + 2\ln(-2 + x) + c_1$$

- Solve for y

$$y = -\ln(x - 1) + 2\ln(-2 + x) + c_1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve((x^2-3*x+2)*diff(y(x),x)=x,y(x), singsol=all)
```

$$y(x) = 2\ln(-2 + x) - \ln(x - 1) + c_1$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 24

```
DSolve[(x^2-3*x+2)*y'[x]==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\log(1 - x) + 2\log(2 - x) + c_1$$

1.25 problem 3(a)

1.25.1 Existence and uniqueness analysis	172
1.25.2 Solving as quadrature ode	173
1.25.3 Maple step by step solution	174

Internal problem ID [6129]

Internal file name [OUTPUT/5377_Sunday_June_05_2022_03_35_44_PM_61552955/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.2 THE NATURE OF SOLUTIONS. Page 9

Problem number: 3(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$y' = x e^x$$

With initial conditions

$$[y(1) = 3]$$

1.25.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = 0$$

$$q(x) = x e^x$$

Hence the ode is

$$y' = x e^x$$

The domain of $p(x) = 0$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 1$ is inside this domain. The domain of $q(x) = x e^x$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 1$ is also inside this domain. Hence solution exists and is unique.

1.25.2 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned} y &= \int x e^x dx \\ &= (x - 1) e^x + c_1 \end{aligned}$$

Initial conditions are used to solve for c_1 . Substituting $x = 1$ and $y = 3$ in the above solution gives an equation to solve for the constant of integration.

$$3 = c_1$$

$$c_1 = 3$$


Substituting c_1 found above in the general solution gives

$$y = x e^x - e^x + 3$$


Summary

The solution(s) found are the following

$$y = x e^x - e^x + 3 \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = x e^x - e^x + 3$$

Verified OK.

1.25.3 Maple step by step solution

Let's solve

$$[y' = x e^x, y(1) = 3]$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Integrate both sides with respect to x

$$\int y' dx = \int x e^x dx + c_1$$

- Evaluate integral

$$y = (x - 1) e^x + c_1$$

- Solve for y

$$y = x e^x - e^x + c_1$$

- Use initial condition $y(1) = 3$

$$3 = c_1$$

- Solve for c_1
 $c_1 = 3$
- Substitute $c_1 = 3$ into general solution and simplify
 $y = 3 + (x - 1)e^x$
- Solution to the IVP
 $y = 3 + (x - 1)e^x$

Maple trace

```
`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 12

```
dsolve([diff(y(x),x)=x*exp(x),y(1) = 3],y(x), singsol=all)
```

$$y(x) = (x - 1)e^x + 3$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 14

```
DSolve[{y'[x]==x*Exp[x],{y[1]==3}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x(x - 1) + 3$$

1.26 problem 3(b)

1.26.1 Existence and uniqueness analysis	176
1.26.2 Solving as quadrature ode	177
1.26.3 Maple step by step solution	178

Internal problem ID [6130]

Internal file name [OUTPUT/5378_Sunday_June_05_2022_03_35_45_PM_64565389/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.2 THE NATURE OF SOLUTIONS. Page 9

Problem number: 3(b).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$y' = 2 \cos(x) \sin(x)$$

With initial conditions

$$[y(0) = 1]$$

1.26.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = 0$$

$$q(x) = \sin(2x)$$

Hence the ode is

$$y' = \sin(2x)$$

The domain of $p(x) = 0$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = \sin(2x)$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

1.26.2 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned} y &= \int 2 \cos(x) \sin(x) \, dx \\ &= \sin(x)^2 + c_1 \end{aligned}$$

Initial conditions are used to solve for c_1 . Substituting $x = 0$ and $y = 1$ in the above solution gives an equation to solve for the constant of integration.

$$1 = c_1$$

$$c_1 = 1$$


Substituting c_1 found above in the general solution gives

$$y = 1 + \sin(x)^2$$


Summary

The solution(s) found are the following

$$y = 1 + \sin(x)^2 \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = 1 + \sin(x)^2$$

Verified OK.

1.26.3 Maple step by step solution

Let's solve

$$[y' = 2 \cos(x) \sin(x), y(0) = 1]$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Integrate both sides with respect to x

$$\int y' dx = \int 2 \cos(x) \sin(x) dx + c_1$$

- Evaluate integral

$$y = \sin(x)^2 + c_1$$

- Solve for y

$$y = \sin(x)^2 + c_1$$

- Use initial condition $y(0) = 1$

$$1 = c_1$$

- Solve for c_1
 $c_1 = 1$
- Substitute $c_1 = 1$ into general solution and simplify
 $y = -\cos(x)^2 + 2$
- Solution to the IVP
 $y = -\cos(x)^2 + 2$

Maple trace

```
`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 12

```
dsolve([diff(y(x),x)=2*sin(x)*cos(x),y(0) = 1],y(x), singsol=all)
```

$$y(x) = -\frac{\cos(2x)}{2} + \frac{3}{2}$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 17

```
DSolve[{y'[x]==2*Sin[x]*Cos[x],{y[0]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}(3 - \cos(2x))$$

1.27 problem 3(c)

1.27.1 Existence and uniqueness analysis	180
1.27.2 Solving as quadrature ode	181
1.27.3 Maple step by step solution	182

Internal problem ID [6131]

Internal file name [OUTPUT/5379_Sunday_June_05_2022_03_35_46_PM_85888940/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.2 THE NATURE OF SOLUTIONS. Page 9

Problem number: 3(c).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$y' = \ln(x)$$

With initial conditions

$$[y(e) = 0]$$

1.27.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = 0$$

$$q(x) = \ln(x)$$

Hence the ode is

$$y' = \ln(x)$$

The domain of $p(x) = 0$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = e$ is inside this domain. The domain of $q(x) = \ln(x)$ is

$$\{0 < x\}$$

And the point $x_0 = e$ is also inside this domain. Hence solution exists and is unique.

1.27.2 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned} y &= \int \ln(x) \, dx \\ &= \ln(x) x - x + c_1 \end{aligned}$$

Initial conditions are used to solve for c_1 . Substituting $x = e$ and $y = 0$ in the above solution gives an equation to solve for the constant of integration.

$$0 = c_1$$

$$c_1 = 0$$


Substituting c_1 found above in the general solution gives

$$y = \ln(x) x - x$$


Summary

The solution(s) found are the following

$$y = \ln(x) x - x \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \ln(x)x - x$$

Verified OK.

1.27.3 Maple step by step solution

Let's solve

$$[y' = \ln(x), y(e) = 0]$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Integrate both sides with respect to x

$$\int y' dx = \int \ln(x) dx + c_1$$

- Evaluate integral

$$y = \ln(x)x - x + c_1$$

- Solve for y

$$y = \ln(x)x - x + c_1$$

- Use initial condition $y(e) = 0$

$$0 = c_1$$

- Solve for c_1
 $c_1 = 0$
- Substitute $c_1 = 0$ into general solution and simplify
 $y = x(\ln(x) - 1)$
- Solution to the IVP
 $y = x(\ln(x) - 1)$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
<- quadrature successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 10

```
dsolve([diff(y(x),x)=ln(x),y(exp(1)) = 0],y(x), singsol=all)
```

$$y(x) = x(\ln(x) - 1)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 11

```
DSolve[{y'[x]==Log[x],{y[Exp[1]]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x(\log(x) - 1)$$

1.28 problem 3(d)

1.28.1 Existence and uniqueness analysis	184
1.28.2 Solving as quadrature ode	185
1.28.3 Maple step by step solution	186

Internal problem ID [6132]

Internal file name [OUTPUT/5380_Sunday_June_05_2022_03_35_47_PM_40340673/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.2 THE NATURE OF SOLUTIONS. Page 9

Problem number: 3(d).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$(x^2 - 1) y' = 1$$

With initial conditions

$$[y(2) = 0]$$

1.28.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = 0$$
$$q(x) = \frac{1}{x^2 - 1}$$

Hence the ode is

$$y' = \frac{1}{x^2 - 1}$$

The domain of $p(x) = 0$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 2$ is inside this domain. The domain of $q(x) = \frac{1}{x^2-1}$ is

$$\{-\infty \leq x < -1, -1 < x < 1, 1 < x \leq \infty\}$$

And the point $x_0 = 2$ is also inside this domain. Hence solution exists and is unique.

1.28.2 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned} y &= \int \frac{1}{x^2 - 1} dx \\ &= -\operatorname{arctanh}(x) + c_1 \end{aligned}$$

Initial conditions are used to solve for c_1 . Substituting $x = 2$ and $y = 0$ in the above solution gives an equation to solve for the constant of integration.

$$0 = -\operatorname{arccoth}(2) + \frac{i\pi}{2} + c_1$$

$$c_1 = \operatorname{arccoth}(2) - \frac{i\pi}{2}$$

Substituting c_1 found above in the general solution gives

$$y = -\operatorname{arctanh}(x) + \operatorname{arccoth}(2) - \frac{i\pi}{2}$$

Summary

The solution(s) found are the following

$$y = -\operatorname{arctanh}(x) + \operatorname{arccoth}(2) - \frac{i\pi}{2} \quad (1)$$

Verification of solutions

$$y = -\operatorname{arctanh}(x) + \operatorname{arccoth}(2) - \frac{i\pi}{2}$$

Verified OK.

1.28.3 Maple step by step solution

Let's solve

$$[(x^2 - 1) y' = 1, y(2) = 0]$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$y' = \frac{1}{x^2-1}$$

- Integrate both sides with respect to x

$$\int y' dx = \int \frac{1}{x^2-1} dx + c_1$$

- Evaluate integral

$$y = -\operatorname{arctanh}(x) + c_1$$

- Solve for y

$$y = -\operatorname{arctanh}(x) + c_1$$

- Use initial condition $y(2) = 0$

$$0 = -\operatorname{arctanh}\left(\frac{1}{2}\right) + \frac{I\pi}{2} + c_1$$

- Solve for c_1

$$c_1 = \operatorname{arctanh}\left(\frac{1}{2}\right) - \frac{I\pi}{2}$$

- Substitute $c_1 = \operatorname{arctanh}\left(\frac{1}{2}\right) - \frac{I\pi}{2}$ into general solution and simplify

$$y = -\operatorname{arctanh}(x) + \operatorname{arctanh}\left(\frac{1}{2}\right) - \frac{I\pi}{2}$$

- Solution to the IVP

$$y = -\operatorname{arctanh}(x) + \operatorname{arctanh}\left(\frac{1}{2}\right) - \frac{I\pi}{2}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 15

```
dsolve([(x^2-1)*diff(y(x),x)=1,y(2) = 0],y(x), singsol=all)
```

$$y(x) = -\operatorname{arctanh}(x) + \operatorname{arctanh}\left(\frac{1}{2}\right) - \frac{i\pi}{2}$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 27

```
DSolve[{(x^2-1)*y'[x]==1,{y[2]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}(\log(3-3x) - \log(x+1) - i\pi)$$

1.29 problem 3(e)

1.29.1 Existence and uniqueness analysis	188
1.29.2 Solving as quadrature ode	189
1.29.3 Maple step by step solution	190

Internal problem ID [6133]

Internal file name [OUTPUT/5381_Sunday_June_05_2022_03_35_49_PM_75045527/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.2 THE NATURE OF SOLUTIONS. Page 9

Problem number: 3(e).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$x(x^2 - 4)y' = 1$$

With initial conditions

$$[y(1) = 0]$$

1.29.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = 0$$
$$q(x) = \frac{1}{x^3 - 4x}$$

Hence the ode is

$$y' = \frac{1}{x^3 - 4x}$$

The domain of $p(x) = 0$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 1$ is inside this domain. The domain of $q(x) = \frac{1}{x^3 - 4x}$ is

$$\{-\infty \leq x < -2, -2 < x < 0, 0 < x < 2, 2 < x \leq \infty\}$$

And the point $x_0 = 1$ is also inside this domain. Hence solution exists and is unique.

1.29.2 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned} y &= \int \frac{1}{x(x^2 - 4)} dx \\ &= -\frac{\ln(x)}{4} + \frac{\ln(x+2)}{8} + \frac{\ln(-2+x)}{8} + c_1 \end{aligned}$$

Initial conditions are used to solve for c_1 . Substituting $x = 1$ and $y = 0$ in the above solution gives an equation to solve for the constant of integration.

$$0 = \frac{\ln(3)}{8} + \frac{i\pi}{8} + c_1$$

$$c_1 = -\frac{\ln(3)}{8} - \frac{i\pi}{8}$$

Substituting c_1 found above in the general solution gives

$$y = -\frac{\ln(x)}{4} + \frac{\ln(x+2)}{8} + \frac{\ln(-2+x)}{8} - \frac{\ln(3)}{8} - \frac{i\pi}{8}$$

Summary

The solution(s) found are the following

$$y = -\frac{\ln(x)}{4} + \frac{\ln(x+2)}{8} + \frac{\ln(-2+x)}{8} - \frac{\ln(3)}{8} - \frac{i\pi}{8} \quad (1)$$

Verification of solutions

$$y = -\frac{\ln(x)}{4} + \frac{\ln(x+2)}{8} + \frac{\ln(-2+x)}{8} - \frac{\ln(3)}{8} - \frac{i\pi}{8}$$

Verified OK.

1.29.3 Maple step by step solution

Let's solve

$$[x(x^2 - 4) y' = 1, y(1) = 0]$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$y' = \frac{1}{x(x^2-4)}$$

- Integrate both sides with respect to x

$$\int y' dx = \int \frac{1}{x(x^2-4)} dx + c_1$$

- Evaluate integral

$$y = -\frac{\ln(x)}{4} + \frac{\ln(x+2)}{8} + \frac{\ln(-2+x)}{8} + c_1$$

- Solve for y

$$y = -\frac{\ln(x)}{4} + \frac{\ln(x+2)}{8} + \frac{\ln(-2+x)}{8} + c_1$$

- Use initial condition $y(1) = 0$

$$0 = \frac{\ln(3)}{8} + \frac{I\pi}{8} + c_1$$

- Solve for c_1

$$c_1 = -\frac{\ln(3)}{8} - \frac{I\pi}{8}$$

- Substitute $c_1 = -\frac{\ln(3)}{8} - \frac{I\pi}{8}$ into general solution and simplify

$$y = -\frac{\ln(x)}{4} + \frac{\ln(x+2)}{8} + \frac{\ln(-2+x)}{8} - \frac{\ln(3)}{8} - \frac{I\pi}{8}$$

- Solution to the IVP

$$y = -\frac{\ln(x)}{4} + \frac{\ln(x+2)}{8} + \frac{\ln(-2+x)}{8} - \frac{\ln(3)}{8} - \frac{I\pi}{8}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 29

```
dsolve([x*(x^2-4)*diff(y(x),x)=1,y(1) = 0],y(x), singsol=all)
```

$$y(x) = \frac{\ln(x+2)}{8} + \frac{\ln(-2+x)}{8} - \frac{\ln(x)}{4} - \frac{\ln(3)}{8} - \frac{i\pi}{8}$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 26

```
DSolve[{x*(x^2-4)*y'[x]==1,{y[1]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{8} \left(\log \left(\frac{1}{3} (4 - x^2) \right) - 2 \log(x) \right)$$

1.30 problem 3(f)

1.30.1 Existence and uniqueness analysis	192
1.30.2 Solving as quadrature ode	193
1.30.3 Maple step by step solution	194

Internal problem ID [6134]

Internal file name [OUTPUT/5382_Sunday_June_05_2022_03_35_50_PM_41186660/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.2 THE NATURE OF SOLUTIONS. Page 9

Problem number: 3(f).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$(1 + x)(x^2 + 1)y' = 2x^2 + x$$

With initial conditions

$$[y(0) = 1]$$

1.30.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = 0$$

$$q(x) = \frac{2x^2 + x}{(1 + x)(x^2 + 1)}$$

Hence the ode is

$$y' = \frac{2x^2 + x}{(1+x)(x^2+1)}$$

The domain of $p(x) = 0$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = \frac{2x^2+x}{(1+x)(x^2+1)}$ is

$$\{x < -1 \vee -1 < x\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

1.30.2 Solving as quadrature ode

Integrating both sides gives

$$\begin{aligned} y &= \int \frac{x(1+2x)}{(1+x)(x^2+1)} dx \\ &= \frac{\ln(1+x)}{2} + \frac{3 \ln(x^2+1)}{4} - \frac{\arctan(x)}{2} + c_1 \end{aligned}$$

Initial conditions are used to solve for c_1 . Substituting $x = 0$ and $y = 1$ in the above solution gives an equation to solve for the constant of integration.

$$1 = c_1$$

$$c_1 = 1$$


Substituting c_1 found above in the general solution gives

$$y = \frac{\ln(1+x)}{2} + \frac{3 \ln(x^2+1)}{4} - \frac{\arctan(x)}{2} + 1$$


Summary

The solution(s) found are the following

$$y = \frac{\ln(1+x)}{2} + \frac{3 \ln(x^2+1)}{4} - \frac{\arctan(x)}{2} + 1 \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{\ln(1+x)}{2} + \frac{3 \ln(x^2+1)}{4} - \frac{\arctan(x)}{2} + 1$$

Verified OK.

1.30.3 Maple step by step solution

Let's solve

$$[(1+x)(x^2+1)y' = 2x^2+x, y(0) = 1]$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$y' = \frac{2x^2+x}{(1+x)(x^2+1)}$$

- Integrate both sides with respect to x

$$\int y' dx = \int \frac{2x^2+x}{(1+x)(x^2+1)} dx + c_1$$

- Evaluate integral

$$y = \frac{\ln(1+x)}{2} + \frac{3 \ln(x^2+1)}{4} - \frac{\arctan(x)}{2} + c_1$$

- Solve for y

$$y = \frac{\ln(1+x)}{2} + \frac{3\ln(x^2+1)}{4} - \frac{\arctan(x)}{2} + c_1$$

- Use initial condition $y(0) = 1$

$$1 = c_1$$

- Solve for c_1

$$c_1 = 1$$

- Substitute $c_1 = 1$ into general solution and simplify

$$y = \frac{\ln(1+x)}{2} + \frac{3\ln(x^2+1)}{4} - \frac{\arctan(x)}{2} + 1$$

- Solution to the IVP

$$y = \frac{\ln(1+x)}{2} + \frac{3\ln(x^2+1)}{4} - \frac{\arctan(x)}{2} + 1$$

Maple trace

```
`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 24

```
dsolve([(x+1)*(x^2+1)*diff(y(x),x)=2*x^2+x,y(0) = 1],y(x), singsol=all)
```

$$y(x) = \frac{3\ln(x^2+1)}{4} - \frac{\arctan(x)}{2} + \frac{\ln(x+1)}{2} + 1$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 29

```
DSolve[{(x+1)*(x^2+1)*y'[x]==2*x^2+x,{y[0]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{4}(-2\arctan(x) + 3\log(x^2+1) + 2\log(x+1) + 4)$$

1.31 problem 4

1.31.1 Solving as linear ode	196
1.31.2 Solving as first order ode lie symmetry lookup ode	198
1.31.3 Solving as exact ode	202
1.31.4 Maple step by step solution	206

Internal problem ID [6135]

Internal file name [OUTPUT/5383_Sunday_June_05_2022_03_35_51_PM_88971099/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.2 THE NATURE OF SOLUTIONS. Page 9

Problem number: 4.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**linear**", "**exactWithIntegrationFactor**", "**first_order_ode_lie_symmetry_lookup**"

Maple gives the following as the ode type

`[_linear]`

$$-2xy + y' = 1$$

1.31.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -2x$$

$$q(x) = 1$$

Hence the ode is

$$-2xy + y' = 1$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int -2x dx} \\ &= e^{-x^2}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= \mu \\ \frac{d}{dx}(e^{-x^2} y) &= e^{-x^2} \\ d(e^{-x^2} y) &= e^{-x^2} dx\end{aligned}$$

Integrating gives

$$\begin{aligned}e^{-x^2} y &= \int e^{-x^2} dx \\ e^{-x^2} y &= \frac{\sqrt{\pi} \operatorname{erf}(x)}{2} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{-x^2}$ results in

$$y = \frac{e^{x^2} \sqrt{\pi} \operatorname{erf}(x)}{2} + c_1 e^{x^2}$$

which simplifies to

$$y = e^{x^2} \left(\frac{\sqrt{\pi} \operatorname{erf}(x)}{2} + c_1 \right)$$

Summary

The solution(s) found are the following

$$y = e^{x^2} \left(\frac{\sqrt{\pi} \operatorname{erf}(x)}{2} + c_1 \right) \tag{1}$$




Figure 46: Slope field plot

Verification of solutions

$$y = e^{x^2} \left(\frac{\sqrt{\pi} \operatorname{erf}(x)}{2} + c_1 \right)$$

Verified OK.

1.31.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$\begin{aligned} y' &= 2xy + 1 \\ y' &= \omega(x, y) \end{aligned}$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 32: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= e^{x^2}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^{x^2}} dy \end{aligned}$$

Which results in

$$S = e^{-x^2} y$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = 2xy + 1$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= -2x e^{-x^2} y \\ S_y &= e^{-x^2} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = e^{-x^2} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = e^{-R^2}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by

integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \frac{\sqrt{\pi} \operatorname{erf}(R)}{2} + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$e^{-x^2} y = \frac{\sqrt{\pi} \operatorname{erf}(x)}{2} + c_1$$



Which simplifies to

$$e^{-x^2} y = \frac{\sqrt{\pi} \operatorname{erf}(x)}{2} + c_1$$

Which gives

$$y = \frac{e^{x^2} (\sqrt{\pi} \operatorname{erf}(x) + 2c_1)}{2}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = 2xy + 1$ 	$R = x$ $S = e^{-x^2} y$	$\frac{dS}{dR} = e^{-R^2}$ 

Summary

The solution(s) found are the following

$$y = \frac{e^{x^2} (\sqrt{\pi} \operatorname{erf}(x) + 2c_1)}{2} \quad (1)$$




Figure 47: Slope field plot

Verification of solutions

$$y = \frac{e^{x^2} (\sqrt{\pi} \operatorname{erf}(x) + 2c_1)}{2}$$

Verified OK.

1.31.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned}dy &= (2xy + 1) dx \\ (-2xy - 1) dx + dy &= 0\end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -2xy - 1 \\ N(x, y) &= 1\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-2xy - 1) \\ &= -2x\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(1) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= 1((-2x) - (0)) \\ &= -2x \end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int A \, dx} \\ &= e^{\int -2x \, dx} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^{-x^2} \\ &= e^{-x^2} \end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \overline{M} and \overline{N} for now so not to confuse them with the original M and N .

$$\begin{aligned} \overline{M} &= \mu M \\ &= e^{-x^2}(-2xy - 1) \\ &= (-2xy - 1)e^{-x^2} \end{aligned}$$

And

$$\begin{aligned} \overline{N} &= \mu N \\ &= e^{-x^2}(1) \\ &= e^{-x^2} \end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned} \overline{M} + \overline{N} \frac{dy}{dx} &= 0 \\ ((-2xy - 1)e^{-x^2}) + (e^{-x^2}) \frac{dy}{dx} &= 0 \end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \overline{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \overline{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \overline{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int (-2xy - 1) e^{-x^2} dx \\ \phi &= -\frac{\sqrt{\pi} \operatorname{erf}(x)}{2} + e^{-x^2} y + f(y)\end{aligned}\quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = e^{-x^2} + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = e^{-x^2}$. Therefore equation (4) becomes

$$e^{-x^2} = e^{-x^2} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = -\frac{\sqrt{\pi} \operatorname{erf}(x)}{2} + e^{-x^2} y + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\frac{\sqrt{\pi} \operatorname{erf}(x)}{2} + e^{-x^2} y$$

The solution becomes

$$y = \frac{e^{x^2} (\sqrt{\pi} \operatorname{erf}(x) + 2c_1)}{2}$$

Summary

The solution(s) found are the following

$$y = \frac{e^{x^2} (\sqrt{\pi} \operatorname{erf}(x) + 2c_1)}{2} \quad (1)$$




Figure 48: Slope field plot

Verification of solutions

$$y = \frac{e^{x^2} (\sqrt{\pi} \operatorname{erf}(x) + 2c_1)}{2}$$

Verified OK.

1.31.4 Maple step by step solution

Let's solve

$$-2xy + y' = 1$$

- Highest derivative means the order of the ODE is 1
- y'
- Isolate the derivative

$$y' = 1 + 2xy$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$-2xy + y' = 1$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) (-2xy + y') = \mu(x)$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x)y)$

$$\mu(x) (-2xy + y') = \mu'(x)y + \mu(x)y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = -2\mu(x)x$$

- Solve to find the integrating factor

$$\mu(x) = e^{-x^2}$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x)y) \right) dx = \int \mu(x) dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x)y = \int \mu(x) dx + c_1$$

- Solve for y

$$y = \frac{\int \mu(x) dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = e^{-x^2}$

$$y = \frac{\int e^{-x^2} dx + c_1}{e^{-x^2}}$$

- Evaluate the integrals on the rhs

$$y = \frac{\frac{\sqrt{\pi} \operatorname{erf}(x)}{2} + c_1}{e^{-x^2}}$$

- Simplify

$$y = \frac{e^{x^2} (\sqrt{\pi} \operatorname{erf}(x) + 2c_1)}{2}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve(diff(y(x),x)=2*x*y(x)+1,y(x), singsol=all)
```

$$y(x) = \frac{(\sqrt{\pi} \operatorname{erf}(x) + 2c_1) e^{x^2}}{2}$$

✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 27

```
DSolve[y'[x]==2*x*y[x]+1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} e^{x^2} (\sqrt{\pi} \operatorname{erf}(x) + 2c_1)$$

1.32 problem 5

1.32.1 Solving as second order linear constant coeff ode	209
1.32.2 Solving using Kovacic algorithm	211
1.32.3 Maple step by step solution	215

Internal problem ID [6136]

Internal file name [OUTPUT/5384_Sunday_June_05_2022_03_35_53_PM_83939676/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.2 THE NATURE OF SOLUTIONS. Page 9

Problem number: 5.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' - 5y' + 4y = 0$$

1.32.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -5, C = 4$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 5\lambda e^{\lambda x} + 4e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 5\lambda + 4 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -5, C = 4$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{5}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-5^2 - (4)(1)(4)} \\ &= \frac{5}{2} \pm \frac{3}{2}\end{aligned}$$

Hence

$$\lambda_1 = \frac{5}{2} + \frac{3}{2}$$

$$\lambda_2 = \frac{5}{2} - \frac{3}{2}$$

Which simplifies to

$$\lambda_1 = 4$$

$$\lambda_2 = 1$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(4)x} + c_2 e^{(1)x}$$

Or

$$y = c_1 e^{4x} + c_2 e^x$$

Summary

The solution(s) found are the following

$$y = c_1 e^{4x} + c_2 e^x \tag{1}$$




Figure 49: Slope field plot

Verification of solutions

$$y = c_1 e^{4x} + c_2 e^x$$

Verified OK.

1.32.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 5y' + 4y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -5 \\ C &= 4 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{9}{4} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 9 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \frac{9z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 35: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{9}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{3x}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-5}{1} dx} \\ &= z_1 e^{\frac{5x}{2}} \\ &= z_1 \left(e^{\frac{5x}{2}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = e^x$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-5}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{5x}}{(y_1)^2} dx \\ &= y_1 \left(\frac{e^{3x}}{3} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1(e^x) + c_2\left(e^x\left(\frac{e^{3x}}{3}\right)\right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^x + \frac{c_2 e^{4x}}{3} \quad (1)$$




Figure 50: Slope field plot

Verification of solutions

$$y = c_1 e^x + \frac{c_2 e^{4x}}{3}$$

Verified OK.

1.32.3 Maple step by step solution

Let's solve

$$y'' - 5y' + 4y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 - 5r + 4 = 0$$

- Factor the characteristic polynomial

$$(r - 1)(r - 4) = 0$$

- Roots of the characteristic polynomial

$$r = (1, 4)$$

- 1st solution of the ODE

$$y_1(x) = e^x$$

- 2nd solution of the ODE

$$y_2(x) = e^{4x}$$

- General solution of the ODE

$$y = c_1y_1(x) + c_2y_2(x)$$

- Substitute in solutions

$$y = c_1e^x + c_2e^{4x}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 15

```
dsolve(diff(y(x),x$2)-5*diff(y(x),x)+4*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{4x}c_1 + e^x c_2$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 20

```
DSolve[y''[x]-5*y'[x]+4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x (c_2 e^{3x} + c_1)$$

1.33 problem 6

- 1.33.1 Solving as first order ode lie symmetry calculated ode 217
- 1.33.2 Solving as exact ode 223

Internal problem ID [6137]

Internal file name [OUTPUT/5385_Sunday_June_05_2022_03_35_54_PM_77836984/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.2 THE NATURE OF SOLUTIONS. Page 9

Problem number: 6.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exactWithIntegrationFactor", "first_order_ode_lie_symmetry_calculated"**

Maple gives the following as the ode type

```
[[_homogeneous, `class G`], _rational, [_Abel, `2nd type`, `class B`]]
```

$$y' - \frac{2xy^2}{1 - yx^2} = 0$$

1.33.1 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = -\frac{2y^2x}{yx^2 - 1}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2\xi_y - \omega_x\xi - \omega_y\eta = 0 \tag{A}$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \tag{1E}$$

$$\eta = xb_2 + yb_3 + b_1 \tag{2E}$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$\begin{aligned} b_2 - \frac{2y^2x(b_3 - a_2)}{yx^2 - 1} - \frac{4y^4x^2a_3}{(yx^2 - 1)^2} - \left(-\frac{2y^2}{yx^2 - 1} + \frac{4y^3x^2}{(yx^2 - 1)^2} \right) (xa_2 + ya_3 + a_1) \\ - \left(-\frac{4xy}{yx^2 - 1} + \frac{2y^2x^3}{(yx^2 - 1)^2} \right) (xb_2 + yb_3 + b_1) = 0 \end{aligned} \quad (5E)$$

Putting the above in normal form gives

$$\frac{3x^4y^2b_2 - 6y^4x^2a_3 + 2x^3y^2b_1 - 2x^2y^3a_1 - 6x^2yb_2 - 4xy^2a_2 - 2xy^2b_3 - 2y^3a_3 - 4xyb_1 - 2y^2a_1 + b_2}{(yx^2 - 1)^2} = 0$$

Setting the numerator to zero gives

$$\begin{aligned} 3x^4y^2b_2 - 6y^4x^2a_3 + 2x^3y^2b_1 - 2x^2y^3a_1 - 6x^2yb_2 \\ - 4xy^2a_2 - 2xy^2b_3 - 2y^3a_3 - 4xyb_1 - 2y^2a_1 + b_2 = 0 \end{aligned} \quad (6E)$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2\}$$

The above PDE (6E) now becomes

$$\begin{aligned} -6a_3v_1^2v_2^4 + 3b_2v_1^4v_2^2 - 2a_1v_1^2v_2^3 + 2b_1v_1^3v_2^2 - 4a_2v_1v_2^2 \\ - 2a_3v_2^3 - 6b_2v_1^2v_2 - 2b_3v_1v_2^2 - 2a_1v_2^2 - 4b_1v_1v_2 + b_2 = 0 \end{aligned} \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2\}$$

Equation (7E) now becomes

$$3b_2v_1^4v_2^2 + 2b_1v_1^3v_2^2 - 6a_3v_1^2v_2^4 - 2a_1v_1^2v_2^3 - 6b_2v_1^2v_2 + (-4a_2 - 2b_3)v_1v_2^2 - 4b_1v_1v_2 - 2a_3v_2^3 - 2a_1v_2^2 + b_2 = 0 \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} b_2 &= 0 \\ -2a_1 &= 0 \\ -6a_3 &= 0 \\ -2a_3 &= 0 \\ -4b_1 &= 0 \\ 2b_1 &= 0 \\ -6b_2 &= 0 \\ 3b_2 &= 0 \\ -4a_2 - 2b_3 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= 0 \\ a_2 &= a_2 \\ a_3 &= 0 \\ b_1 &= 0 \\ b_2 &= 0 \\ b_3 &= -2a_2 \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned} \xi &= x \\ \eta &= -2y \end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned} \eta &= \eta - \omega(x, y) \xi \\ &= -2y - \left(-\frac{2y^2x}{yx^2 - 1} \right) (x) \\ &= \frac{2y}{yx^2 - 1} \\ \xi &= 0 \end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{2y}{y x^2 - 1}} dy \end{aligned}$$

Which results in

$$S = \frac{y x^2}{2} - \frac{\ln(y)}{2}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -\frac{2y^2x}{y x^2 - 1}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= xy \\ S_y &= \frac{y x^2 - 1}{2y} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \tag{2A}$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \tag{4}$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{yx^2}{2} - \frac{\ln(y)}{2} = c_1$$



Which simplifies to

$$\frac{yx^2}{2} - \frac{\ln(y)}{2} = c_1$$

Which gives

$$y = e^{-\text{LambertW}(-x^2e^{-2c_1}) - 2c_1}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -\frac{2y^2x}{yx^2-1}$ 	$R = x$ $S = \frac{yx^2}{2} - \frac{\ln(y)}{2}$	$\frac{dS}{dR} = 0$ 

Summary

The solution(s) found are the following

$$y = e^{-\text{LambertW}(-x^2e^{-2c_1})-2c_1} \tag{1}$$




Figure 51: Slope field plot

Verification of solutions

$$y = e^{-\text{LambertW}(-x^2 e^{-2c_1}) - 2c_1}$$

Verified OK.

1.33.2 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}(y x^2 - 1) dy &= (-2y^2 x) dx \\ (2y^2 x) dx + (y x^2 - 1) dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= 2y^2 x \\ N(x, y) &= y x^2 - 1\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(2y^2 x) \\ &= 4xy\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(y x^2 - 1) \\ &= 2xy\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= \frac{1}{y x^2 - 1} ((4xy) - (2xy)) \\ &= \frac{2xy}{y x^2 - 1} \end{aligned}$$

Since A depends on y , it can not be used to obtain an integrating factor. We will now try a second method to find an integrating factor. Let

$$\begin{aligned} B &= \frac{1}{M} \left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) \\ &= \frac{1}{2y^2x} ((2xy) - (4xy)) \\ &= -\frac{1}{y} \end{aligned}$$

Since B does not depend on x , it can be used to obtain an integrating factor. Let the integrating factor be μ . Then

$$\begin{aligned} \mu &= e^{\int B \, dy} \\ &= e^{\int -\frac{1}{y} \, dy} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^{-\ln(y)} \\ &= \frac{1}{y} \end{aligned}$$

M and N are now multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} so not to confuse them with the original M and N .

$$\begin{aligned} \bar{M} &= \mu M \\ &= \frac{1}{y} (2y^2x) \\ &= 2xy \end{aligned}$$

And

$$\begin{aligned} \bar{N} &= \mu N \\ &= \frac{1}{y} (y x^2 - 1) \\ &= \frac{y x^2 - 1}{y} \end{aligned}$$

So now a modified ODE is obtained from the original ODE which will be exact and can be solved using the standard method. The modified ODE is

$$\begin{aligned}\bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ (2xy) + \left(\frac{y x^2 - 1}{y} \right) \frac{dy}{dx} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \quad (1)$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int 2xy dx \\ \phi &= y x^2 + f(y)\end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = x^2 + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{y x^2 - 1}{y}$. Therefore equation (4) becomes

$$\frac{y x^2 - 1}{y} = x^2 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = -\frac{1}{y}$$

Integrating the above w.r.t y gives

$$\begin{aligned}\int f'(y) dy &= \int \left(-\frac{1}{y} \right) dy \\ f(y) &= -\ln(y) + c_1\end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = y x^2 - \ln(y) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = y x^2 - \ln(y)$$

The solution becomes

$$y = e^{-\text{LambertW}(-e^{-c_1} x^2) - c_1}$$

Summary

The solution(s) found are the following

$$y = e^{-\text{LambertW}(-e^{-c_1} x^2) - c_1} \tag{1}$$




Figure 52: Slope field plot

Verification of solutions

$$y = e^{-\text{LambertW}(-e^{-c_1} x^2) - c_1}$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying homogeneous G  
<- homogeneous successful`
```

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 19

```
dsolve(diff(y(x),x)=2*x*y(x)^2/(1-x^2*y(x)),y(x), singsol=all)
```

$$y(x) = -\frac{\text{LambertW}(-x^2 e^{-2c_1})}{x^2}$$

✓ Solution by Mathematica

Time used: 4.457 (sec). Leaf size: 27

```
DSolve[y'[x]==2*x*y[x]^2/(1-x^2*y[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{W(-e^{-1+c_1} x^2)}{x^2}$$
$$y(x) \rightarrow 0$$

1.34 problem 7

1.34.1 Maple step by step solution 230

Internal problem ID [6138]

Internal file name [OUTPUT/5386_Sunday_June_05_2022_03_35_56_PM_8521084/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.2 THE NATURE OF SOLUTIONS. Page 9

Problem number: 7.

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_3rd_order , _missing_x]]
```

$$2y''' + y'' - 5y' + 2y = 0$$

The characteristic equation is

$$2\lambda^3 + \lambda^2 - 5\lambda + 2 = 0$$

The roots of the above equation are

$$\lambda_1 = 1$$

$$\lambda_2 = \frac{1}{2}$$

$$\lambda_3 = -2$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 e^{-2x} + c_2 e^x + c_3 e^{\frac{x}{2}}$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = e^{-2x}$$

$$y_2 = e^x$$

$$y_3 = e^{\frac{x}{2}}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-2x} + c_2 e^x + c_3 e^{\frac{x}{2}} \quad (1)$$

Verification of solutions

$$y = c_1 e^{-2x} + c_2 e^x + c_3 e^{\frac{x}{2}}$$

Verified OK.

1.34.1 Maple step by step solution

Let's solve

$$2y''' + y'' - 5y' + 2y = 0$$

- Highest derivative means the order of the ODE is 3

$$y'''$$

- Isolate 3rd derivative

$$y''' = -\frac{y''}{2} + \frac{5y'}{2} - y$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y''' + \frac{y''}{2} - \frac{5y'}{2} + y = 0$$

- Convert linear ODE into a system of first order ODEs

- Define new variable $y_1(x)$

$$y_1(x) = y$$

- Define new variable $y_2(x)$

$$y_2(x) = y'$$

- Define new variable $y_3(x)$

$$y_3(x) = y''$$

- Isolate for $y_3'(x)$ using original ODE

$$y_3'(x) = -\frac{y_3(x)}{2} + \frac{5y_2(x)}{2} - y_1(x)$$

Convert linear ODE into a system of first order ODEs

$$\left[y_2(x) = y_1'(x), y_3(x) = y_2'(x), y_3'(x) = -\frac{y_3(x)}{2} + \frac{5y_2(x)}{2} - y_1(x) \right]$$

- Define vector

$$\vec{y}(x) = \begin{bmatrix} y_1(x) \\ y_2(x) \\ y_3(x) \end{bmatrix}$$

- System to solve

$$\vec{y}'(x) = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ -1 & \frac{5}{2} & -\frac{1}{2} \end{bmatrix} \cdot \vec{y}(x)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ -1 & \frac{5}{2} & -\frac{1}{2} \end{bmatrix}$$

- Rewrite the system as

$$\vec{y}'(x) = A \cdot \vec{y}(x)$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[-2, \begin{bmatrix} \frac{1}{4} \\ -\frac{1}{2} \\ 1 \end{bmatrix} \right], \left[\frac{1}{2}, \begin{bmatrix} 4 \\ 2 \\ 1 \end{bmatrix} \right], \left[1, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[-2, \begin{bmatrix} \frac{1}{4} \\ -\frac{1}{2} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_1 = e^{-2x} \cdot \begin{bmatrix} \frac{1}{4} \\ -\frac{1}{2} \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[\frac{1}{2}, \begin{bmatrix} 4 \\ 2 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_2 = e^{\frac{x}{2}} \cdot \begin{bmatrix} 4 \\ 2 \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[1, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_3 = e^x \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{y} = c_1 \vec{y}_1 + c_2 \vec{y}_2 + c_3 \vec{y}_3$$

- Substitute solutions into the general solution

$$\vec{y} = c_1 e^{-2x} \cdot \begin{bmatrix} \frac{1}{4} \\ -\frac{1}{2} \\ 1 \end{bmatrix} + c_2 e^{\frac{x}{2}} \cdot \begin{bmatrix} 4 \\ 2 \\ 1 \end{bmatrix} + c_3 e^x \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y = \frac{(4c_3 e^{3x} + 16c_2 e^{\frac{5x}{2}} + c_1) e^{-2x}}{4}$$

Maple trace

```
`Methods for third order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(2*diff(y(x),x$3)+diff(y(x),x$2)-5*diff(y(x),x)+2*y(x)=0,y(x), singsol=all)
```

$$y(x) = \left(c_3 e^{3x} + c_1 e^{\frac{5x}{2}} + c_2 \right) e^{-2x}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 30

```
DSolve[2*y'''[x]+y''[x]-5*y'[x]+2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{x/2} + c_2 e^{-2x} + c_3 e^x$$

2 Chapter 1. What is a differential equation.

Section 1.3 SEPARABLE EQUATIONS. Page 12

2.1	problem 1(a)	235
2.2	problem 1(b)	239
2.3	problem 1(c)	242
2.4	problem 1(d)	245
2.5	problem 1(e)	249
2.6	problem 1(f)	253
2.7	problem 1(g)	257
2.8	problem 1(h)	261
2.9	problem 1(i)	264
2.10	problem 1(j)	268
2.11	problem 2(a)	272
2.12	problem 2(b)	277
2.13	problem 2(c)	281
2.14	problem 2(d)	287
2.15	problem 2(e)	292
2.16	problem 2(e)	297
2.17	problem 3	302
2.18	problem 4	308

2.1 problem 1(a)

2.1.1 Solving as separable ode	235
2.1.2 Maple step by step solution	237

Internal problem ID [6139]

Internal file name [OUTPUT/5387_Sunday_June_05_2022_03_35_57_PM_84684605/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.3 SEPARABLE EQUATIONS.

Page 12

Problem number: 1(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

`[_separable]`

$$y'x^5 + y^5 = 0$$

2.1.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= -\frac{y^5}{x^5}\end{aligned}$$

Where $f(x) = -\frac{1}{x^5}$ and $g(y) = y^5$. Integrating both sides gives

$$\begin{aligned}\frac{1}{y^5} dy &= -\frac{1}{x^5} dx \\ \int \frac{1}{y^5} dy &= \int -\frac{1}{x^5} dx \\ -\frac{1}{4y^4} &= \frac{1}{4x^4} + c_1\end{aligned}$$

The solution is

$$-\frac{1}{4y^4} - \frac{1}{4x^4} - c_1 = 0$$

Summary

The solution(s) found are the following

$$-\frac{1}{4y^4} - \frac{1}{4x^4} - c_1 = 0 \tag{1}$$




Figure 53: Slope field plot

Verification of solutions

$$-\frac{1}{4y^4} - \frac{1}{4x^4} - c_1 = 0$$

Verified OK.

2.1.2 Maple step by step solution

Let's solve

$$y'x^5 + y^5 = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y^5} = -\frac{1}{x^5}$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y^5} dx = \int -\frac{1}{x^5} dx + c_1$$

- Evaluate integral

$$-\frac{1}{4y^4} = \frac{1}{4x^4} + c_1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
<- Bernoulli successful`
```

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 67

```
dsolve(x^5*diff(y(x),x)+y(x)^5=0,y(x), singsol=all)
```

$$y(x) = \frac{x}{(c_1x^4 - 1)^{\frac{1}{4}}}$$

$$y(x) = -\frac{x}{(c_1x^4 - 1)^{\frac{1}{4}}}$$

$$y(x) = \frac{x}{\sqrt{-\sqrt{c_1x^4 - 1}}}$$

$$y(x) = -\frac{x}{\sqrt{-\sqrt{c_1x^4 - 1}}}$$

✓ Solution by Mathematica

Time used: 0.489 (sec). Leaf size: 145

```
DSolve[x^5*y'[x]+y[x]^5==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{x}{\sqrt[4]{-1-4c_1x^4}}$$

$$y(x) \rightarrow -\frac{ix}{\sqrt[4]{-1-4c_1x^4}}$$

$$y(x) \rightarrow \frac{ix}{\sqrt[4]{-1-4c_1x^4}}$$

$$y(x) \rightarrow \frac{x}{\sqrt[4]{-1-4c_1x^4}}$$

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow -\frac{(1+i)x}{\sqrt{2}}$$

$$y(x) \rightarrow -\frac{(1-i)x}{\sqrt{2}}$$

$$y(x) \rightarrow \frac{(1-i)x}{\sqrt{2}}$$

$$y(x) \rightarrow \frac{(1+i)x}{\sqrt{2}}$$

2.2 problem 1(b)

2.2.1 Solving as separable ode	239
2.2.2 Maple step by step solution	240

Internal problem ID [6140]

Internal file name [OUTPUT/5388_Sunday_June_05_2022_03_35_59_PM_76582465/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.3 SEPARABLE EQUATIONS.

Page 12

Problem number: 1(b).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

`[_separable]`

$$y' - 4xy = 0$$

2.2.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= 4xy\end{aligned}$$

Where $f(x) = 4x$ and $g(y) = y$. Integrating both sides gives

$$\begin{aligned}\frac{1}{y} dy &= 4x dx \\ \int \frac{1}{y} dy &= \int 4x dx \\ \ln(y) &= 2x^2 + c_1 \\ y &= e^{2x^2+c_1} \\ &= c_1 e^{2x^2}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{2x^2} \quad (1)$$




Figure 54: Slope field plot

Verification of solutions

$$y = c_1 e^{2x^2}$$

Verified OK.

2.2.2 Maple step by step solution

Let's solve

$$y' - 4xy = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y} = 4x$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y} dx = \int 4x dx + c_1$$

- Evaluate integral

$$\ln(y) = 2x^2 + c_1$$

- Solve for y

$$y = e^{2x^2 + c_1}$$

Maple trace

```
`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)=4*x*y(x),y(x), singsol=all)
```

$$y(x) = c_1 e^{2x^2}$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 20

```
DSolve[y'[x]==4*x*y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{2x^2}$$

$$y(x) \rightarrow 0$$

2.3 problem 1(c)

2.3.1 Solving as separable ode	242
2.3.2 Maple step by step solution	243

Internal problem ID [6141]

Internal file name [OUTPUT/5389_Sunday_June_05_2022_03_36_01_PM_92610766/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.3 SEPARABLE EQUATIONS.

Page 12

Problem number: 1(c).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

`[_separable]`

$$y' + y \tan(x) = 0$$

2.3.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= -y \tan(x)\end{aligned}$$

Where $f(x) = -\tan(x)$ and $g(y) = y$. Integrating both sides gives

$$\begin{aligned}\frac{1}{y} dy &= -\tan(x) dx \\ \int \frac{1}{y} dy &= \int -\tan(x) dx \\ \ln(y) &= \ln(\cos(x)) + c_1 \\ y &= e^{\ln(\cos(x)) + c_1} \\ &= \cos(x) c_1\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \cos(x) c_1 \quad (1)$$




Figure 55: Slope field plot

Verification of solutions

$$y = \cos(x) c_1$$

Verified OK.

2.3.2 Maple step by step solution

Let's solve

$$y' + y \tan(x) = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y} = -\tan(x)$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y} dx = \int -\tan(x) dx + c_1$$
- Evaluate integral

$$\ln(y) = \ln(\cos(x)) + c_1$$
- Solve for y

$$y = \cos(x) e^{c_1}$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 8

```
dsolve(diff(y(x),x)+y(x)*tan(x)=0,y(x), singsol=all)
```

$$y(x) = \cos(x) c_1$$

✓ Solution by Mathematica

Time used: 0.037 (sec). Leaf size: 15

```
DSolve[y'[x]+y[x]*Tan[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos(x)$$

$$y(x) \rightarrow 0$$

2.4 problem 1(d)

2.4.1 Solving as separable ode	245
2.4.2 Maple step by step solution	247

Internal problem ID [6142]

Internal file name [OUTPUT/5390_Sunday_June_05_2022_03_36_02_PM_91456469/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.3 SEPARABLE EQUATIONS.

Page 12

Problem number: 1(d).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

`[_separable]`

$$(x^2 + 1) y' + y^2 = -1$$

2.4.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{-y^2 - 1}{x^2 + 1} \end{aligned}$$

Where $f(x) = \frac{1}{x^2+1}$ and $g(y) = -y^2 - 1$. Integrating both sides gives

$$\begin{aligned} \frac{1}{-y^2 - 1} dy &= \frac{1}{x^2 + 1} dx \\ \int \frac{1}{-y^2 - 1} dy &= \int \frac{1}{x^2 + 1} dx \\ -\arctan(y) &= \arctan(x) + c_1 \end{aligned}$$

Which results in

$$y = -\tan(\arctan(x) + c_1)$$

Summary

The solution(s) found are the following

$$y = -\tan(\arctan(x) + c_1) \tag{1}$$




Figure 56: Slope field plot

Verification of solutions

$$y = -\tan(\arctan(x) + c_1)$$

Verified OK.

2.4.2 Maple step by step solution

Let's solve

$$(x^2 + 1) y' + y^2 = -1$$

- Highest derivative means the order of the ODE is 1

y'

- Separate variables

$$\frac{y'}{-1-y^2} = \frac{1}{x^2+1}$$

- Integrate both sides with respect to x

$$\int \frac{y'}{-1-y^2} dx = \int \frac{1}{x^2+1} dx + c_1$$

- Evaluate integral

$$-\arctan(y) = \arctan(x) + c_1$$

- Solve for y

$$y = -\tan(\arctan(x) + c_1)$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
<- separable successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve((1+x^2)*diff(y(x),x)+1+y(x)^2=0,y(x), singsol=all)
```

$$y(x) = -\tan(\arctan(x) + c_1)$$

✓ Solution by Mathematica

Time used: 0.233 (sec). Leaf size: 29

```
DSolve[(1+x^2)*y'[x]+1+y[x]^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\tan(\arctan(x) - c_1)$$

$$y(x) \rightarrow -i$$

$$y(x) \rightarrow i$$

2.5 problem 1(e)

2.5.1 Solving as separable ode	249
2.5.2 Maple step by step solution	251

Internal problem ID [6143]

Internal file name [OUTPUT/5391_Sunday_June_05_2022_03_36_04_PM_1499382/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.3 SEPARABLE EQUATIONS.

Page 12

Problem number: 1(e).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

`[_separable]`

$$y \ln(y) - xy' = 0$$

2.5.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{y \ln(y)}{x} \end{aligned}$$

Where $f(x) = \frac{1}{x}$ and $g(y) = \ln(y)y$. Integrating both sides gives

$$\begin{aligned} \frac{1}{\ln(y)y} dy &= \frac{1}{x} dx \\ \int \frac{1}{\ln(y)y} dy &= \int \frac{1}{x} dx \\ \ln(\ln(y)) &= \ln(x) + c_1 \end{aligned}$$

Raising both side to exponential gives

$$\ln(y) = e^{\ln(x)+c_1}$$

Which simplifies to

$$\ln(y) = c_2x$$

Summary

The solution(s) found are the following

$$y = e^{c_2x e^{c_1}} \quad (1)$$




Figure 57: Slope field plot

Verification of solutions

$$y = e^{c_2x e^{c_1}}$$

Verified OK.

2.5.2 Maple step by step solution

Let's solve

$$y \ln(y) - xy' = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y \ln(y)} = \frac{1}{x}$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y \ln(y)} dx = \int \frac{1}{x} dx + c_1$$

- Evaluate integral

$$\ln(\ln(y)) = \ln(x) + c_1$$

- Solve for y

$$y = e^{x e^{c_1}}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
<- separable successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 8

```
dsolve(y(x)*ln(y(x))-x*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = e^{c_1 x}$$

✓ Solution by Mathematica

Time used: 0.172 (sec). Leaf size: 18

```
DSolve[y[x]*Log[y[x]]-x*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{e^{c_1 x}}$$

$$y(x) \rightarrow 1$$

2.6 problem 1(f)

2.6.1 Solving as separable ode	253
2.6.2 Maple step by step solution	255

Internal problem ID [6144]

Internal file name [OUTPUT/5392_Sunday_June_05_2022_03_36_06_PM_57599098/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.3 SEPARABLE EQUATIONS.

Page 12

Problem number: 1(f).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

`[_separable]`

$$xy' - (-4x^2 + 1) \tan(y) = 0$$

2.6.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= -\frac{\tan(y)(4x^2 - 1)}{x}\end{aligned}$$

Where $f(x) = -\frac{4x^2-1}{x}$ and $g(y) = \tan(y)$. Integrating both sides gives

$$\begin{aligned}\frac{1}{\tan(y)} dy &= -\frac{4x^2 - 1}{x} dx \\ \int \frac{1}{\tan(y)} dy &= \int -\frac{4x^2 - 1}{x} dx \\ \ln(\sin(y)) &= -2x^2 + \ln(x) + c_1\end{aligned}$$

Raising both side to exponential gives

$$\sin(y) = e^{-2x^2 + \ln(x) + c_1}$$

Which simplifies to

$$\sin(y) = c_2 e^{-2x^2 + \ln(x)}$$

Summary

The solution(s) found are the following

$$y = \arcsin\left(c_2 e^{-2x^2 + c_1} x\right) \quad (1)$$




Figure 58: Slope field plot

Verification of solutions

$$y = \arcsin\left(c_2 e^{-2x^2 + c_1} x\right)$$

Verified OK.

2.6.2 Maple step by step solution

Let's solve

$$xy' - (-4x^2 + 1) \tan(y) = 0$$

- Highest derivative means the order of the ODE is 1

y'

- Separate variables

$$\frac{y'}{\tan(y)} = \frac{-4x^2+1}{x}$$

- Integrate both sides with respect to x

$$\int \frac{y'}{\tan(y)} dx = \int \frac{-4x^2+1}{x} dx + c_1$$

- Evaluate integral

$$\ln(\sin(y)) = -2x^2 + \ln(x) + c_1$$

- Solve for y

$$y = \arcsin\left(\frac{x}{e^{2x^2-c_1}}\right)$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
<- separable successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 16

```
dsolve(x*diff(y(x),x)=(1-4*x^2)*tan(y(x)),y(x), singsol=all)
```

$$y(x) = \arcsin\left(\frac{x e^{-2x^2}}{c_1}\right)$$

✓ Solution by Mathematica

Time used: 53.453 (sec). Leaf size: 23

```
DSolve[x*y'[x]==(1-4*x^2)*Tan[y[x]],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \arcsin\left(xe^{-2x^2+c_1}\right)$$

$$y(x) \rightarrow 0$$

2.7 problem 1(g)

2.7.1 Solving as separable ode	257
2.7.2 Maple step by step solution	259

Internal problem ID [6145]

Internal file name [OUTPUT/5393_Sunday_June_05_2022_03_36_07_PM_45941527/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.3 SEPARABLE EQUATIONS.

Page 12

Problem number: 1(g).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

`[_separable]`

$$y' \sin(y) = x^2$$

2.7.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{x^2}{\sin(y)}\end{aligned}$$

Where $f(x) = x^2$ and $g(y) = \frac{1}{\sin(y)}$. Integrating both sides gives

$$\begin{aligned}\frac{1}{\frac{1}{\sin(y)}} dy &= x^2 dx \\ \int \frac{1}{\frac{1}{\sin(y)}} dy &= \int x^2 dx \\ -\cos(y) &= \frac{x^3}{3} + c_1\end{aligned}$$

Which results in

$$y = \pi - \arccos\left(\frac{x^3}{3} + c_1\right)$$

Summary

The solution(s) found are the following

$$y = \pi - \arccos\left(\frac{x^3}{3} + c_1\right) \tag{1}$$




Figure 59: Slope field plot

Verification of solutions

$$y = \pi - \arccos\left(\frac{x^3}{3} + c_1\right)$$

Verified OK.

2.7.2 Maple step by step solution

Let's solve

$$y' \sin(y) = x^2$$

- Highest derivative means the order of the ODE is 1

y'

- Integrate both sides with respect to x

$$\int y' \sin(y) dx = \int x^2 dx + c_1$$

- Evaluate integral

$$-\cos(y) = \frac{x^3}{3} + c_1$$

- Solve for y

$$y = \pi - \arccos\left(\frac{x^3}{3} + c_1\right)$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
<- separable successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve(diff(y(x),x)*sin(y(x))=x^2,y(x), singsol=all)
```

$$y(x) = \frac{\pi}{2} + \arcsin\left(\frac{x^3}{3} + c_1\right)$$

✓ Solution by Mathematica

Time used: 0.499 (sec). Leaf size: 37

```
DSolve[y'[x]*Sin[y[x]]==x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\arccos\left(-\frac{x^3}{3} - c_1\right)$$

$$y(x) \rightarrow \arccos\left(-\frac{x^3}{3} - c_1\right)$$

2.8 problem 1(h)

2.8.1 Solving as separable ode	261
2.8.2 Maple step by step solution	262

Internal problem ID [6146]

Internal file name [OUTPUT/5394_Sunday_June_05_2022_03_36_09_PM_7071013/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.3 SEPARABLE EQUATIONS.

Page 12

Problem number: 1(h).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

`[_separable]`

$$y' - y \tan(x) = 0$$

2.8.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= y \tan(x)\end{aligned}$$

Where $f(x) = \tan(x)$ and $g(y) = y$. Integrating both sides gives

$$\begin{aligned}\frac{1}{y} dy &= \tan(x) dx \\ \int \frac{1}{y} dy &= \int \tan(x) dx \\ \ln(y) &= -\ln(\cos(x)) + c_1 \\ y &= e^{-\ln(\cos(x))+c_1} \\ &= \frac{c_1}{\cos(x)}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1}{\cos(x)} \quad (1)$$




Figure 60: Slope field plot

Verification of solutions

$$y = \frac{c_1}{\cos(x)}$$

Verified OK.

2.8.2 Maple step by step solution

Let's solve

$$y' - y \tan(x) = 0$$

- Highest derivative means the order of the ODE is 1
- y'
- Separate variables

$$\frac{y'}{y} = \tan(x)$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y} dx = \int \tan(x) dx + c_1$$

- Evaluate integral

$$\ln(y) = -\ln(\cos(x)) + c_1$$

- Solve for y

$$y = \frac{e^{c_1}}{\cos(x)}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 8

```
dsolve(diff(y(x),x)-y(x)*tan(x)=0,y(x), singsol=all)
```

$$y(x) = \sec(x) c_1$$

✓ Solution by Mathematica

Time used: 0.034 (sec). Leaf size: 15

```
DSolve[y'[x]-y[x]*Tan[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \sec(x)$$

$$y(x) \rightarrow 0$$

2.9 problem 1(i)

2.9.1 Solving as separable ode	264
2.9.2 Maple step by step solution	266

Internal problem ID [6147]

Internal file name [OUTPUT/5395_Sunday_June_05_2022_03_36_10_PM_98519124/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.3 SEPARABLE EQUATIONS.

Page 12

Problem number: 1(i).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

`[_separable]`

$$xyy' - y = -1$$

2.9.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{y-1}{xy}\end{aligned}$$

Where $f(x) = \frac{1}{x}$ and $g(y) = \frac{y-1}{y}$. Integrating both sides gives

$$\frac{1}{\frac{y-1}{y}} dy = \frac{1}{x} dx$$

$$\int \frac{1}{\frac{y-1}{y}} dy = \int \frac{1}{x} dx$$

$$y + \ln(y-1) = \ln(x) + c_1$$

Which results in

$$y = \text{LambertW}(x e^{-1+c_1}) + 1$$

Since c_1 is constant, then exponential powers of this constant are constants also, and these can be simplified to just c_1 in the above solution. Which simplifies to

$$y = \text{LambertW}(x e^{-1+c_1}) + 1$$

gives

$$y = \text{LambertW}(x e^{-1}c_1) + 1$$

Summary

The solution(s) found are the following

$$y = \text{LambertW}(x e^{-1}c_1) + 1 \tag{1}$$




Figure 61: Slope field plot

Verification of solutions

$$y = \text{LambertW}(x e^{-1}c_1) + 1$$

Verified OK.

2.9.2 Maple step by step solution

Let's solve

$$xyy' - y = -1$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'y}{y-1} = \frac{1}{x}$$

- Integrate both sides with respect to x

$$\int \frac{y'y}{y-1} dx = \int \frac{1}{x} dx + c_1$$

- Evaluate integral

$$y + \ln(y - 1) = \ln(x) + c_1$$

- Solve for y

$$y = \text{LambertW}(x e^{-1+c_1}) + 1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
<- separable successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 12

```
dsolve(x*y(x)*diff(y(x),x)=y(x)-1,y(x), singsol=all)
```

$$y(x) = \text{LambertW}(x c_1 e^{-1}) + 1$$

✓ Solution by Mathematica

Time used: 3.215 (sec). Leaf size: 21

```
DSolve[x*y[x]*y'[x]==y[x]-1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 1 + W(e^{-1+c_1x})$$

$$y(x) \rightarrow 1$$

2.10 problem 1(j)

2.10.1 Solving as separable ode	268
2.10.2 Maple step by step solution	270

Internal problem ID [6148]

Internal file name [OUTPUT/5396_Sunday_June_05_2022_03_36_12_PM_71582804/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.3 SEPARABLE EQUATIONS.

Page 12

Problem number: 1(j).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

`[_separable]`

$$xy^2 - x^2y' = 0$$

2.10.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{y^2}{x}\end{aligned}$$

Where $f(x) = \frac{1}{x}$ and $g(y) = y^2$. Integrating both sides gives

$$\begin{aligned}\frac{1}{y^2} dy &= \frac{1}{x} dx \\ \int \frac{1}{y^2} dy &= \int \frac{1}{x} dx \\ -\frac{1}{y} &= \ln(x) + c_1\end{aligned}$$

Which results in

$$y = -\frac{1}{\ln(x) + c_1}$$

Summary

The solution(s) found are the following

$$y = -\frac{1}{\ln(x) + c_1} \tag{1}$$




Figure 62: Slope field plot

Verification of solutions

$$y = -\frac{1}{\ln(x) + c_1}$$

Verified OK.

2.10.2 Maple step by step solution

Let's solve

$$xy^2 - x^2y' = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y^2} = \frac{1}{x}$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y^2} dx = \int \frac{1}{x} dx + c_1$$

- Evaluate integral

$$-\frac{1}{y} = \ln(x) + c_1$$

- Solve for y

$$y = -\frac{1}{\ln(x) + c_1}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
<- Bernoulli successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 12

```
dsolve(x*y(x)^2-diff(y(x),x)*x^2=0,y(x), singsol=all)
```

$$y(x) = \frac{1}{-\ln(x) + c_1}$$

✓ Solution by Mathematica

Time used: 0.107 (sec). Leaf size: 19

```
DSolve[x*y[x]^2-y'[x]*x^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{\log(x) + c_1}$$
$$y(x) \rightarrow 0$$

2.11 problem 2(a)

2.11.1 Existence and uniqueness analysis	272
2.11.2 Solving as separable ode	273
2.11.3 Maple step by step solution	275

Internal problem ID [6149]

Internal file name [OUTPUT/5397_Sunday_June_05_2022_03_36_14_PM_99221309/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.3 SEPARABLE EQUATIONS.

Page 12

Problem number: 2(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "separable", "differential-Type", "homogeneousTypeMapleC", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

`[_separable]`

$$y'y = 1 + x$$

With initial conditions

$$[y(1) = 3]$$

2.11.1 Existence and uniqueness analysis

This is non linear first order ODE. In canonical form it is written as

$$\begin{aligned} y' &= f(x, y) \\ &= \frac{1 + x}{y} \end{aligned}$$

The x domain of $f(x, y)$ when $y = 3$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 1$ is inside this domain. The y domain of $f(x, y)$ when $x = 1$ is

$$\{y < 0 \vee 0 < y\}$$

And the point $y_0 = 3$ is inside this domain. Now we will look at the continuity of

$$\begin{aligned}\frac{\partial f}{\partial y} &= \frac{\partial}{\partial y} \left(\frac{1+x}{y} \right) \\ &= -\frac{1+x}{y^2}\end{aligned}$$

The x domain of $\frac{\partial f}{\partial y}$ when $y = 3$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 1$ is inside this domain. The y domain of $\frac{\partial f}{\partial y}$ when $x = 1$ is

$$\{y < 0 \vee 0 < y\}$$

And the point $y_0 = 3$ is inside this domain. Therefore solution exists and is unique.

2.11.2 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{1+x}{y}\end{aligned}$$

Where $f(x) = 1 + x$ and $g(y) = \frac{1}{y}$. Integrating both sides gives

$$\begin{aligned}\frac{1}{\frac{1}{y}} dy &= 1 + x dx \\ \int \frac{1}{\frac{1}{y}} dy &= \int 1 + x dx \\ \frac{y^2}{2} &= \frac{1}{2}x^2 + x + c_1\end{aligned}$$

Which results in

$$y = \sqrt{x^2 + 2c_1 + 2x}$$

$$y = -\sqrt{x^2 + 2c_1 + 2x}$$

Initial conditions are used to solve for c_1 . Substituting $x = 1$ and $y = 3$ in the above solution gives an equation to solve for the constant of integration.

$$3 = -\sqrt{3 + 2c_1}$$

Warning: Unable to solve for constant of integration. Initial conditions are used to solve for c_1 . Substituting $x = 1$ and $y = 3$ in the above solution gives an equation to solve for the constant of integration.

$$3 = \sqrt{3 + 2c_1}$$

$$c_1 = 3$$


Substituting c_1 found above in the general solution gives

$$y = \sqrt{x^2 + 2x + 6}$$


Summary

The solution(s) found are the following

$$y = \sqrt{x^2 + 2x + 6} \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \sqrt{x^2 + 2x + 6}$$

Verified OK.

2.11.3 Maple step by step solution

Let's solve

$$[y'y = 1 + x, y(1) = 3]$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Integrate both sides with respect to x

$$\int y'y dx = \int (1 + x) dx + c_1$$

- Evaluate integral

$$\frac{y^2}{2} = \frac{1}{2}x^2 + x + c_1$$

- Solve for y

$$\{y = \sqrt{x^2 + 2c_1 + 2x}, y = -\sqrt{x^2 + 2c_1 + 2x}\}$$

- Use initial condition $y(1) = 3$

$$3 = \sqrt{3 + 2c_1}$$

- Solve for c_1

$$c_1 = 3$$

- Substitute $c_1 = 3$ into general solution and simplify

$$y = \sqrt{x^2 + 2x + 6}$$

- Use initial condition $y(1) = 3$

$$3 = -\sqrt{3 + 2c_1}$$

- Solution does not satisfy initial condition

- Solution to the IVP

$$y = \sqrt{x^2 + 2x + 6}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
<- Bernoulli successful`
```

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 14

```
dsolve([diff(y(x),x)*y(x)=x+1,y(1) = 3],y(x), singsol=all)
```

$$y(x) = \sqrt{x^2 + 2x + 6}$$

✓ Solution by Mathematica

Time used: 0.097 (sec). Leaf size: 17

```
DSolve[{y'[x]*y[x]==x+1,{y[1]==3}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \sqrt{x^2 + 2x + 6}$$

2.12 problem 2(b)

2.12.1 Existence and uniqueness analysis	277
2.12.2 Solving as separable ode	278
2.12.3 Maple step by step solution	279

Internal problem ID [6150]

Internal file name [OUTPUT/5398_Sunday_June_05_2022_03_36_15_PM_15070873/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.3 SEPARABLE EQUATIONS.

Page 12

Problem number: 2(b).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "linear", "separable", "homogeneousTypeD2", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_separable]

$$x^2y' - y = 0$$

With initial conditions

$$[y(1) = 0]$$

2.12.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -\frac{1}{x^2}$$
$$q(x) = 0$$

Hence the ode is

$$y' - \frac{y}{x^2} = 0$$

The domain of $p(x) = -\frac{1}{x^2}$ is

$$\{x < 0 \vee 0 < x\}$$

And the point $x_0 = 1$ is inside this domain. Hence solution exists and is unique.

2.12.2 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{y}{x^2}\end{aligned}$$


Where $f(x) = \frac{1}{x^2}$ and $g(y) = y$. Since unique solution exists and $g(y)$ evaluated at $y_0 = 0$ is zero, then the solution is

$$\begin{aligned}y &= y_0 \\ &= 0\end{aligned}$$


Summary

The solution(s) found are the following

$$y = 0 \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = 0$$

Verified OK.

2.12.3 Maple step by step solution

Let's solve

$$[x^2y' - y = 0, y(1) = 0]$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y} = \frac{1}{x^2}$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y} dx = \int \frac{1}{x^2} dx + c_1$$

- Evaluate integral

$$\ln(y) = c_1 - \frac{1}{x}$$

- Solve for y

$$y = e^{\frac{c_1 x - 1}{x}}$$

- Use initial condition $y(1) = 0$

$$0 = e^{-1+c_1}$$

- Solution does not satisfy initial condition

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 5

```
dsolve([diff(y(x),x)*x^2=y(x),y(1) = 0],y(x), singsol=all)
```

$$y(x) = 0$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 6

```
DSolve[{y'[x]*x^2==y[x],{y[1]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 0$$

2.13 problem 2(c)

2.13.1 Existence and uniqueness analysis	281
2.13.2 Solving as separable ode	282
2.13.3 Maple step by step solution	284

Internal problem ID [6151]

Internal file name [OUTPUT/5399_Sunday_June_05_2022_03_36_17_PM_95195842/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.3 SEPARABLE EQUATIONS.

Page 12

Problem number: 2(c).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "separable", "differential-Type", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_separable]

$$\frac{y'}{x^2 + 1} - \frac{x}{y} = 0$$

With initial conditions

$$[y(1) = 3]$$

2.13.1 Existence and uniqueness analysis

This is non linear first order ODE. In canonical form it is written as

$$\begin{aligned} y' &= f(x, y) \\ &= \frac{x(x^2 + 1)}{y} \end{aligned}$$

The x domain of $f(x, y)$ when $y = 3$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 1$ is inside this domain. The y domain of $f(x, y)$ when $x = 1$ is

$$\{y < 0 \vee 0 < y\}$$

And the point $y_0 = 3$ is inside this domain. Now we will look at the continuity of

$$\begin{aligned}\frac{\partial f}{\partial y} &= \frac{\partial}{\partial y} \left(\frac{x(x^2 + 1)}{y} \right) \\ &= -\frac{x(x^2 + 1)}{y^2}\end{aligned}$$

The x domain of $\frac{\partial f}{\partial y}$ when $y = 3$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 1$ is inside this domain. The y domain of $\frac{\partial f}{\partial y}$ when $x = 1$ is

$$\{y < 0 \vee 0 < y\}$$

And the point $y_0 = 3$ is inside this domain. Therefore solution exists and is unique.

2.13.2 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{x(x^2 + 1)}{y}\end{aligned}$$

Where $f(x) = x(x^2 + 1)$ and $g(y) = \frac{1}{y}$. Integrating both sides gives

$$\begin{aligned}\frac{1}{\frac{1}{y}} dy &= x(x^2 + 1) dx \\ \int \frac{1}{\frac{1}{y}} dy &= \int x(x^2 + 1) dx \\ \frac{y^2}{2} &= \frac{(x^2 + 1)^2}{4} + c_1\end{aligned}$$

Which results in

$$y = \frac{\sqrt{2x^4 + 4x^2 + 8c_1 + 2}}{2}$$
$$y = -\frac{\sqrt{2x^4 + 4x^2 + 8c_1 + 2}}{2}$$

Initial conditions are used to solve for c_1 . Substituting $x = 1$ and $y = 3$ in the above solution gives an equation to solve for the constant of integration.

$$3 = -\sqrt{2 + 2c_1}$$

Warning: Unable to solve for constant of integration. Initial conditions are used to solve for c_1 . Substituting $x = 1$ and $y = 3$ in the above solution gives an equation to solve for the constant of integration.

$$3 = \sqrt{2 + 2c_1}$$

$$c_1 = \frac{7}{2}$$


Substituting c_1 found above in the general solution gives

$$y = \frac{\sqrt{2x^4 + 4x^2 + 30}}{2}$$


Summary

The solution(s) found are the following

$$y = \frac{\sqrt{2x^4 + 4x^2 + 30}}{2} \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{\sqrt{2x^4 + 4x^2 + 30}}{2}$$

Verified OK.

2.13.3 Maple step by step solution

Let's solve

$$\left[\frac{y'}{x^2+1} - \frac{x}{y} = 0, y(1) = 3 \right]$$

- Highest derivative means the order of the ODE is 1

y'

- Separate variables

$$y'y = x(x^2 + 1)$$

- Integrate both sides with respect to x

$$\int y'y dx = \int x(x^2 + 1) dx + c_1$$

- Evaluate integral

$$\frac{y^2}{2} = \frac{(x^2+1)^2}{4} + c_1$$

- Solve for y

$$\left\{ y = -\frac{\sqrt{2x^4+4x^2+8c_1+2}}{2}, y = \frac{\sqrt{2x^4+4x^2+8c_1+2}}{2} \right\}$$

- Use initial condition $y(1) = 3$

$$3 = -\frac{\sqrt{8+8c_1}}{2}$$

- Solution does not satisfy initial condition

- Use initial condition $y(1) = 3$

$$3 = \frac{\sqrt{8+8c_1}}{2}$$

- Solve for c_1

$$c_1 = \frac{7}{2}$$

- Substitute $c_1 = \frac{7}{2}$ into general solution and simplify

$$y = \frac{\sqrt{2x^4+4x^2+30}}{2}$$

- Solution to the IVP

$$y = \frac{\sqrt{2x^4+4x^2+30}}{2}$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
<- Bernoulli successful`

```

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 20

```
dsolve([diff(y(x),x)/(1+x^2)=x/y(x),y(1) = 3],y(x), singsol=all)
```

$$y(x) = \frac{\sqrt{2x^4 + 4x^2 + 30}}{2}$$

✓ Solution by Mathematica

Time used: 0.104 (sec). Leaf size: 25

```
DSolve[{y'[x]/(1+x^2)==x/y[x],{y[1]==3}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\sqrt{x^4 + 2x^2 + 15}}{\sqrt{2}}$$

2.14 problem 2(d)

2.14.1 Existence and uniqueness analysis	287
2.14.2 Solving as separable ode	288
2.14.3 Maple step by step solution	290

Internal problem ID [6152]

Internal file name [OUTPUT/5400_Sunday_June_05_2022_03_36_18_PM_50028180/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.3 SEPARABLE EQUATIONS.

Page 12

Problem number: 2(d).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "separable", "differential-Type", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_separable]

$$y'y^2 = x + 2$$

With initial conditions

$$[y(0) = 4]$$

2.14.1 Existence and uniqueness analysis

This is non linear first order ODE. In canonical form it is written as

$$\begin{aligned}y' &= f(x, y) \\ &= \frac{x + 2}{y^2}\end{aligned}$$

The x domain of $f(x, y)$ when $y = 4$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The y domain of $f(x, y)$ when $x = 0$ is

$$\{y < 0 \vee 0 < y\}$$

And the point $y_0 = 4$ is inside this domain. Now we will look at the continuity of

$$\begin{aligned}\frac{\partial f}{\partial y} &= \frac{\partial}{\partial y} \left(\frac{x+2}{y^2} \right) \\ &= -\frac{2(x+2)}{y^3}\end{aligned}$$

The x domain of $\frac{\partial f}{\partial y}$ when $y = 4$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The y domain of $\frac{\partial f}{\partial y}$ when $x = 0$ is

$$\{y < 0 \vee 0 < y\}$$

And the point $y_0 = 4$ is inside this domain. Therefore solution exists and is unique.

2.14.2 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{x+2}{y^2}\end{aligned}$$

Where $f(x) = x + 2$ and $g(y) = \frac{1}{y^2}$. Integrating both sides gives

$$\begin{aligned}\frac{1}{y^2} dy &= x + 2 dx \\ \int \frac{1}{y^2} dy &= \int x + 2 dx \\ \frac{y^3}{3} &= \frac{1}{2}x^2 + 2x + c_1\end{aligned}$$

Which results in

$$y = \frac{(12x^2 + 24c_1 + 48x)^{\frac{1}{3}}}{2}$$
$$y = -\frac{(12x^2 + 24c_1 + 48x)^{\frac{1}{3}}}{4} + \frac{i\sqrt{3}(12x^2 + 24c_1 + 48x)^{\frac{1}{3}}}{4}$$
$$y = -\frac{(12x^2 + 24c_1 + 48x)^{\frac{1}{3}}}{4} - \frac{i\sqrt{3}(12x^2 + 24c_1 + 48x)^{\frac{1}{3}}}{4}$$

Initial conditions are used to solve for c_1 . Substituting $x = 0$ and $y = 4$ in the above solution gives an equation to solve for the constant of integration.

$$4 = -\frac{ic_1^{\frac{1}{3}}24^{\frac{1}{3}}\sqrt{3}}{4} - \frac{c_1^{\frac{1}{3}}24^{\frac{1}{3}}}{4}$$

Warning: Unable to solve for constant of integration. Initial conditions are used to solve for c_1 . Substituting $x = 0$ and $y = 4$ in the above solution gives an equation to solve for the constant of integration.

$$4 = \frac{ic_1^{\frac{1}{3}}24^{\frac{1}{3}}\sqrt{3}}{4} - \frac{c_1^{\frac{1}{3}}24^{\frac{1}{3}}}{4}$$

Warning: Unable to solve for constant of integration. Initial conditions are used to solve for c_1 . Substituting $x = 0$ and $y = 4$ in the above solution gives an equation to solve for the constant of integration.

$$4 = \frac{c_1^{\frac{1}{3}}24^{\frac{1}{3}}}{2}$$

$$c_1 = \frac{64}{3}$$


Substituting c_1 found above in the general solution gives

$$y = \frac{(12x^2 + 48x + 512)^{\frac{1}{3}}}{2}$$


Summary

The solution(s) found are the following

$$y = \frac{(12x^2 + 48x + 512)^{\frac{1}{3}}}{2} \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{(12x^2 + 48x + 512)^{\frac{1}{3}}}{2}$$

Verified OK.

2.14.3 Maple step by step solution

Let's solve

$$[y'y^2 = x + 2, y(0) = 4]$$

- Highest derivative means the order of the ODE is 1

y'

- Integrate both sides with respect to x

$$\int y'y^2 dx = \int (x + 2) dx + c_1$$

- Evaluate integral

$$\frac{y^3}{3} = \frac{1}{2}x^2 + 2x + c_1$$

- Solve for y

$$y = \frac{(12x^2 + 24c_1 + 48x)^{\frac{1}{3}}}{2}$$

- Use initial condition $y(0) = 4$

$$4 = \frac{c_1^{\frac{1}{3}} 24^{\frac{1}{3}}}{2}$$

- Solve for c_1

$$c_1 = \frac{64}{3}$$

- Substitute $c_1 = \frac{64}{3}$ into general solution and simplify

$$y = \frac{(12x^2 + 48x + 512)^{\frac{1}{3}}}{2}$$

- Solution to the IVP

$$y = \frac{(12x^2 + 48x + 512)^{\frac{1}{3}}}{2}$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
<- Bernoulli successful`

```

✓ Solution by Maple

Time used: 0.141 (sec). Leaf size: 18

```
dsolve([y(x)^2*diff(y(x),x)=x+2,y(0) = 4],y(x), singsol=all)
```

$$y(x) = \frac{(12x^2 + 48x + 512)^{\frac{1}{3}}}{2}$$

✓ Solution by Mathematica

Time used: 0.211 (sec). Leaf size: 21

```
DSolve[{y[x]^2*y'[x]==x+2,{y[0]==4}],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \sqrt[3]{\frac{3x^2}{2} + 6x + 64}$$

2.15 problem 2(e)

2.15.1 Existence and uniqueness analysis	292
2.15.2 Solving as separable ode	293
2.15.3 Maple step by step solution	295

Internal problem ID [6153]

Internal file name [OUTPUT/5401_Sunday_June_05_2022_03_36_20_PM_57043492/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.3 SEPARABLE EQUATIONS.

Page 12

Problem number: 2(e).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "riccati", "separable",
"first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_separable]

$$y' - y^2x^2 = 0$$

With initial conditions

$$[y(-1) = 2]$$

2.15.1 Existence and uniqueness analysis

This is non linear first order ODE. In canonical form it is written as

$$\begin{aligned}y' &= f(x, y) \\ &= y^2x^2\end{aligned}$$

The x domain of $f(x, y)$ when $y = 2$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = -1$ is inside this domain. The y domain of $f(x, y)$ when $x = -1$ is

$$\{-\infty < y < \infty\}$$

And the point $y_0 = 2$ is inside this domain. Now we will look at the continuity of

$$\begin{aligned}\frac{\partial f}{\partial y} &= \frac{\partial}{\partial y}(y^2 x^2) \\ &= 2y x^2\end{aligned}$$

The x domain of $\frac{\partial f}{\partial y}$ when $y = 2$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = -1$ is inside this domain. The y domain of $\frac{\partial f}{\partial y}$ when $x = -1$ is

$$\{-\infty < y < \infty\}$$

And the point $y_0 = 2$ is inside this domain. Therefore solution exists and is unique.

2.15.2 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= y^2 x^2\end{aligned}$$

Where $f(x) = x^2$ and $g(y) = y^2$. Integrating both sides gives

$$\begin{aligned}\frac{1}{y^2} dy &= x^2 dx \\ \int \frac{1}{y^2} dy &= \int x^2 dx \\ -\frac{1}{y} &= \frac{x^3}{3} + c_1\end{aligned}$$

Which results in

$$y = -\frac{3}{x^3 + 3c_1}$$

Initial conditions are used to solve for c_1 . Substituting $x = -1$ and $y = 2$ in the above solution gives an equation to solve for the constant of integration.

$$2 = -\frac{3}{3c_1 - 1}$$

$$c_1 = -\frac{1}{6}$$


Substituting c_1 found above in the general solution gives

$$y = -\frac{6}{2x^3 - 1}$$


Summary

The solution(s) found are the following

$$y = -\frac{6}{2x^3 - 1} \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = -\frac{6}{2x^3 - 1}$$

Verified OK.

2.15.3 Maple step by step solution

Let's solve

$$[y' - y^2x^2 = 0, y(-1) = 2]$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y^2} = x^2$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y^2} dx = \int x^2 dx + c_1$$

- Evaluate integral

$$-\frac{1}{y} = \frac{x^3}{3} + c_1$$

- Solve for y

$$y = -\frac{3}{x^3 + 3c_1}$$

- Use initial condition $y(-1) = 2$

$$2 = -\frac{3}{3c_1 - 1}$$

- Solve for c_1

$$c_1 = -\frac{1}{6}$$

- Substitute $c_1 = -\frac{1}{6}$ into general solution and simplify

$$y = -\frac{6}{2x^3 - 1}$$

- Solution to the IVP

$$y = -\frac{6}{2x^3 - 1}$$

Maple trace

```
`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
<- Bernoulli successful`
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 15

```
dsolve([diff(y(x),x)=x^2*y(x)^2,y(-1) = 2],y(x), singsol=all)
```

$$y(x) = -\frac{6}{2x^3 - 1}$$

✓ Solution by Mathematica

Time used: 0.106 (sec). Leaf size: 16

```
DSolve[{y'[x]==x^2*y[x]^2,{y[-1]==2}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{6}{1 - 2x^3}$$

2.16 problem 2(e)

2.16.1 Existence and uniqueness analysis	297
2.16.2 Solving as separable ode	298
2.16.3 Maple step by step solution	300

Internal problem ID [6154]

Internal file name [OUTPUT/5402_Sunday_June_05_2022_03_36_21_PM_69555050/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.3 SEPARABLE EQUATIONS.

Page 12

Problem number: 2(e).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "separable", "differential-Type", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

`[_separable]`

$$y'(1 + y) = -x^2 + 1$$

With initial conditions

$$[y(-1) = -2]$$

2.16.1 Existence and uniqueness analysis

This is non linear first order ODE. In canonical form it is written as

$$\begin{aligned} y' &= f(x, y) \\ &= -\frac{x^2 - 1}{1 + y} \end{aligned}$$

The x domain of $f(x, y)$ when $y = -2$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = -1$ is inside this domain. The y domain of $f(x, y)$ when $x = -1$ is

$$\{-\infty < y < \infty\}$$

And the point $y_0 = -2$ is inside this domain. Now we will look at the continuity of

$$\begin{aligned}\frac{\partial f}{\partial y} &= \frac{\partial}{\partial y} \left(-\frac{x^2 - 1}{1 + y} \right) \\ &= \frac{x^2 - 1}{(1 + y)^2}\end{aligned}$$

The x domain of $\frac{\partial f}{\partial y}$ when $y = -2$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = -1$ is inside this domain. The y domain of $\frac{\partial f}{\partial y}$ when $x = -1$ is

$$\{-\infty < y < \infty\}$$

And the point $y_0 = -2$ is inside this domain. Therefore solution exists and is unique.

2.16.2 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{-x^2 + 1}{1 + y}\end{aligned}$$

Where $f(x) = -x^2 + 1$ and $g(y) = \frac{1}{1+y}$. Integrating both sides gives

$$\begin{aligned}\frac{1}{1+y} dy &= -x^2 + 1 dx \\ \int \frac{1}{1+y} dy &= \int -x^2 + 1 dx \\ y + \frac{1}{2}y^2 &= -\frac{1}{3}x^3 + x + c_1\end{aligned}$$

Which results in

$$y = -1 + \frac{\sqrt{-6x^3 + 18c_1 + 18x + 9}}{3}$$

$$y = -1 - \frac{\sqrt{-6x^3 + 18c_1 + 18x + 9}}{3}$$

Initial conditions are used to solve for c_1 . Substituting $x = -1$ and $y = -2$ in the above solution gives an equation to solve for the constant of integration.

$$-2 = -1 - \frac{\sqrt{-3 + 18c_1}}{3}$$

$$c_1 = \frac{2}{3}$$

Substituting c_1 found above in the general solution gives

$$y = -1 - \frac{\sqrt{-6x^3 + 18x + 21}}{3}$$

Initial conditions are used to solve for c_1 . Substituting $x = -1$ and $y = -2$ in the above solution gives an equation to solve for the constant of integration.


$$-2 = -1 + \frac{\sqrt{-3 + 18c_1}}{3}$$

Summary


The solution(s) found are the following

Warning: Unable to solve for constant of integration.

$$y = -1 - \frac{\sqrt{-6x^3 + 18x + 21}}{3}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = -1 - \frac{\sqrt{-6x^3 + 18x + 21}}{3}$$

Verified OK.

2.16.3 Maple step by step solution

Let's solve

$$[y'(1 + y) = -x^2 + 1, y(-1) = -2]$$

- Highest derivative means the order of the ODE is 1

y'

- Integrate both sides with respect to x

$$\int y'(1 + y) dx = \int (-x^2 + 1) dx + c_1$$

- Evaluate integral

$$\frac{y^2}{2} + y = -\frac{1}{3}x^3 + x + c_1$$

- Solve for y

$$\left\{ y = -1 - \frac{\sqrt{-6x^3 + 18c_1 + 18x + 9}}{3}, y = -1 + \frac{\sqrt{-6x^3 + 18c_1 + 18x + 9}}{3} \right\}$$

- Use initial condition $y(-1) = -2$

$$-2 = -1 - \frac{\sqrt{-3 + 18c_1}}{3}$$

- Solve for c_1

$$c_1 = \frac{2}{3}$$

- Substitute $c_1 = \frac{2}{3}$ into general solution and simplify

$$y = -1 - \frac{\sqrt{-6x^3 + 18x + 21}}{3}$$

- Use initial condition $y(-1) = -2$

$$-2 = -1 + \frac{\sqrt{-3 + 18c_1}}{3}$$

- Solution does not satisfy initial condition

- Solution to the IVP

$$y = -1 - \frac{\sqrt{-6x^3 + 18x + 21}}{3}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
<- separable successful`
```

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 20

```
dsolve([diff(y(x),x)*(1+y(x))=1-x^2,y(-1) = -2],y(x), singsol=all)
```

$$y(x) = -1 - \frac{\sqrt{-6x^3 + 18x + 21}}{3}$$

✓ Solution by Mathematica

Time used: 0.151 (sec). Leaf size: 28

```
DSolve[{y'[x]*(1+y[x])==1-x^2,{y[-1]==-2}],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\sqrt{-2x^3 + 6x + 7}}{\sqrt{3}} - 1$$

2.17 problem 3

2.17.1 Solving as second order integrable as is ode	302
2.17.2 Solving as second order ode missing y ode	303
2.17.3 Solving as exact nonlinear second order ode ode	304
2.17.4 Maple step by step solution	305

Internal problem ID [6155]

Internal file name [OUTPUT/5403_Sunday_June_05_2022_03_36_23_PM_21409840/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.3 SEPARABLE EQUATIONS.

Page 12

Problem number: 3.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_integrable_as_is", "second_order_ode_missing_y", "exact nonlinear second order ode"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_y]]
```

$$0 = -\frac{y''}{y'} + x^2$$

2.17.1 Solving as second order integrable as is ode

Integrating both sides of the ODE w.r.t x gives

$$\int -\frac{y''}{y'} dx = \int -x^2 dx$$
$$-\ln(y') = -\frac{x^3}{3} + c_1$$

Which is now solved for y . Integrating both sides gives

$$y = \int e^{\frac{x^3}{3} - c_1} dx$$
$$= \frac{e^{-c_1} 3^{\frac{1}{3}} (-1)^{\frac{2}{3}} \left(\frac{2x\sqrt{3}(-1)^{\frac{1}{3}}\pi}{3\Gamma(\frac{2}{3})(-x^3)^{\frac{1}{3}}} - \frac{x(-1)^{\frac{1}{3}}\Gamma(\frac{1}{3}, -\frac{x^3}{3})}{(-x^3)^{\frac{1}{3}}} \right)}{3} + c_2$$

Summary

The solution(s) found are the following

$$y = -\frac{e^{-c_1} 3^{\frac{1}{3}} (-1)^{\frac{2}{3}} \left(\frac{2x\sqrt{3}(-1)^{\frac{1}{3}}\pi}{3\Gamma(\frac{2}{3})(-x^3)^{\frac{1}{3}}} - \frac{x(-1)^{\frac{1}{3}}\Gamma(\frac{1}{3}, -\frac{x^3}{3})}{(-x^3)^{\frac{1}{3}}} \right)}{3} + c_2 \quad (1)$$

Verification of solutions

$$y = -\frac{e^{-c_1} 3^{\frac{1}{3}} (-1)^{\frac{2}{3}} \left(\frac{2x\sqrt{3}(-1)^{\frac{1}{3}}\pi}{3\Gamma(\frac{2}{3})(-x^3)^{\frac{1}{3}}} - \frac{x(-1)^{\frac{1}{3}}\Gamma(\frac{1}{3}, -\frac{x^3}{3})}{(-x^3)^{\frac{1}{3}}} \right)}{3} + c_2$$

Verified OK.

2.17.2 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$-\frac{p'(x)}{p(x)} + x^2 = 0$$

Which is now solve for $p(x)$ as first order ode. In canonical form the ODE is

$$\begin{aligned} p' &= F(x, p) \\ &= f(x)g(p) \\ &= x^2 p \end{aligned}$$

Where $f(x) = x^2$ and $g(p) = p$. Integrating both sides gives

$$\begin{aligned} \frac{1}{p} dp &= x^2 dx \\ \int \frac{1}{p} dp &= \int x^2 dx \\ \ln(p) &= \frac{x^3}{3} + c_1 \\ p &= e^{\frac{x^3}{3} + c_1} \\ &= c_1 e^{\frac{x^3}{3}} \end{aligned}$$

Since $p = y'$ then the new first order ode to solve is

$$y' = c_1 e^{\frac{x^3}{3}}$$

Integrating both sides gives

$$\begin{aligned} y &= \int c_1 e^{\frac{x^3}{3}} dx \\ &= -\frac{c_1 \mathfrak{I}^{\frac{1}{3}}(-1)^{\frac{2}{3}} \left(\frac{2x\sqrt{3}(-1)^{\frac{1}{3}}\pi}{3\Gamma(\frac{2}{3})(-x^3)^{\frac{1}{3}}} - \frac{x(-1)^{\frac{1}{3}}\Gamma(\frac{1}{3}, -\frac{x^3}{3})}{(-x^3)^{\frac{1}{3}}} \right)}{3} + c_2 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = -\frac{c_1 \mathfrak{I}^{\frac{1}{3}}(-1)^{\frac{2}{3}} \left(\frac{2x\sqrt{3}(-1)^{\frac{1}{3}}\pi}{3\Gamma(\frac{2}{3})(-x^3)^{\frac{1}{3}}} - \frac{x(-1)^{\frac{1}{3}}\Gamma(\frac{1}{3}, -\frac{x^3}{3})}{(-x^3)^{\frac{1}{3}}} \right)}{3} + c_2 \quad (1)$$

Verification of solutions

$$y = -\frac{c_1 \mathfrak{I}^{\frac{1}{3}}(-1)^{\frac{2}{3}} \left(\frac{2x\sqrt{3}(-1)^{\frac{1}{3}}\pi}{3\Gamma(\frac{2}{3})(-x^3)^{\frac{1}{3}}} - \frac{x(-1)^{\frac{1}{3}}\Gamma(\frac{1}{3}, -\frac{x^3}{3})}{(-x^3)^{\frac{1}{3}}} \right)}{3} + c_2$$

Verified OK.

2.17.3 Solving as exact nonlinear second order ode

An exact non-linear second order ode has the form

$$a_2(x, y, y') y'' + a_1(x, y, y') y' + a_0(x, y, y') = 0$$

Where the following conditions are satisfied

$$\begin{aligned} \frac{\partial a_2}{\partial y} &= \frac{\partial a_1}{\partial y'} \\ \frac{\partial a_2}{\partial x} &= \frac{\partial a_0}{\partial y'} \\ \frac{\partial a_1}{\partial x} &= \frac{\partial a_0}{\partial y} \end{aligned}$$

Looking at the the ode given we see that

$$\begin{aligned} a_2 &= -\frac{1}{y'} \\ a_1 &= 0 \\ a_0 &= x^2 \end{aligned}$$

Applying the conditions to the above shows this is a nonlinear exact second order ode. Therefore it can be reduced to first order ode given by

$$\int a_2 dy' + \int a_1 dy + \int a_0 dx = c_1$$

$$\int -\frac{1}{y'} dy' + \int 0 dy + \int x^2 dx = c_1$$

Which results in

$$\frac{x^3}{3} - \ln(y') = c_1$$

Which is now solved Integrating both sides gives

$$y = \int e^{\frac{x^3}{3} - c_1} dx$$

$$= -\frac{e^{-c_1} 3^{\frac{1}{3}} (-1)^{\frac{2}{3}} \left(\frac{2x\sqrt{3}(-1)^{\frac{1}{3}}\pi}{3\Gamma(\frac{2}{3})(-x^3)^{\frac{1}{3}}} - \frac{x(-1)^{\frac{1}{3}}\Gamma(\frac{1}{3}, -\frac{x^3}{3})}{(-x^3)^{\frac{1}{3}}} \right)}{3} + c_2$$

Summary

The solution(s) found are the following

$$y = -\frac{e^{-c_1} 3^{\frac{1}{3}} (-1)^{\frac{2}{3}} \left(\frac{2x\sqrt{3}(-1)^{\frac{1}{3}}\pi}{3\Gamma(\frac{2}{3})(-x^3)^{\frac{1}{3}}} - \frac{x(-1)^{\frac{1}{3}}\Gamma(\frac{1}{3}, -\frac{x^3}{3})}{(-x^3)^{\frac{1}{3}}} \right)}{3} + c_2 \quad (1)$$

Verification of solutions

$$y = -\frac{e^{-c_1} 3^{\frac{1}{3}} (-1)^{\frac{2}{3}} \left(\frac{2x\sqrt{3}(-1)^{\frac{1}{3}}\pi}{3\Gamma(\frac{2}{3})(-x^3)^{\frac{1}{3}}} - \frac{x(-1)^{\frac{1}{3}}\Gamma(\frac{1}{3}, -\frac{x^3}{3})}{(-x^3)^{\frac{1}{3}}} \right)}{3} + c_2$$

Verified OK.

2.17.4 Maple step by step solution

Let's solve

$$-\frac{y''}{y'} = -x^2$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = x^2 y'$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$-x^2 y' + y'' = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^k$$

- Rewrite ODE with series expansions

- Convert $x^2 \cdot y'$ to series expansion

$$x^2 \cdot y' = \sum_{k=0}^{\infty} a_k k x^{k+1}$$

- Shift index using $k- > k-1$

$$x^2 \cdot y' = \sum_{k=1}^{\infty} a_{k-1} (k-1) x^k$$

- Convert y'' to series expansion

$$y'' = \sum_{k=2}^{\infty} a_k k(k-1) x^{k-2}$$

- Shift index using $k- > k+2$

$$y'' = \sum_{k=0}^{\infty} a_{k+2} (k+2)(k+1) x^k$$

Rewrite ODE with series expansions

$$2a_2 + \left(\sum_{k=1}^{\infty} (a_{k+2} (k+2)(k+1) - a_{k-1} (k-1)) x^k \right) = 0$$

- Each term must be 0

$$2a_2 = 0$$

- Each term in the series must be 0, giving the recursion relation

$$(k^2 + 3k + 2) a_{k+2} - a_{k-1} (k-1) = 0$$

- Shift index using $k- > k+1$

$$((k+1)^2 + 3k + 5) a_{k+3} - a_k k = 0$$

- Recursion relation that defines the series solution to the ODE

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+3} = \frac{a_k k}{k^2 + 5k + 6}, 2a_2 = 0 \right]$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
checking if the LODE is missing y  
<- LODE missing y successful`
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 25

```
dsolve(diff(y(x),x$2)/diff(y(x),x)=x^2,y(x), singsol=all)
```

$$y(x) = -3\Gamma\left(\frac{1}{3}, -\frac{x^3}{3}\right) c_2 \Gamma\left(\frac{2}{3}\right) + 2\sqrt{3}\pi c_2 + c_1$$

✓ Solution by Mathematica

Time used: 0.042 (sec). Leaf size: 39

```
DSolve[y''[x]/y'[x]==x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_1(-x^3)^{2/3} \Gamma\left(\frac{1}{3}, -\frac{x^3}{3}\right)}{3^{2/3}x^2} + c_2$$

2.18 problem 4

2.18.1 Solving as second order integrable as is ode	308
2.18.2 Solving as second order ode missing y ode	310
2.18.3 Solving as exact nonlinear second order ode ode	312
2.18.4 Maple step by step solution	313

Internal problem ID [6156]

Internal file name [OUTPUT/5404_Sunday_June_05_2022_03_36_24_PM_12653789/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.3 SEPARABLE EQUATIONS.

Page 12

Problem number: 4.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second_order_integrable_as_is", "second_order_ode_missing_y", "exact nonlinear second order ode"**

Maple gives the following as the ode type

```
[[_2nd_order, _missing_y], [_2nd_order, _exact, _nonlinear], [_2nd_order, _reducible, _mu_y_y1], [_2nd_order, _reducible, _mu_poly_yn]]
```

$$y''y' = x(1 + x)$$

2.18.1 Solving as second order integrable as is ode

Integrating both sides of the ODE w.r.t x gives

$$\int y''y' dx = \int (x^2 + x) dx$$
$$\frac{y'^2}{2} = \frac{1}{3}x^3 + \frac{1}{2}x^2 + c_1$$

Which is now solved for y . Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \frac{\sqrt{6x^3 + 9x^2 + 18c_1}}{3} \quad (1)$$

$$y' = -\frac{\sqrt{6x^3 + 9x^2 + 18c_1}}{3} \quad (2)$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\begin{aligned} y &= \int \frac{\sqrt{6x^3 + 9x^2 + 18c_1}}{3} dx \\ &= \int \frac{\sqrt{6x^3 + 9x^2 + 18c_1}}{3} dx + c_2 \end{aligned}$$

Solving equation (2)

Integrating both sides gives

$$\begin{aligned} y &= \int -\frac{\sqrt{6x^3 + 9x^2 + 18c_1}}{3} dx \\ &= \int -\frac{\sqrt{6x^3 + 9x^2 + 18c_1}}{3} dx + c_3 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \int \frac{\sqrt{6x^3 + 9x^2 + 18c_1}}{3} dx + c_2 \quad (1)$$

$$y = \int -\frac{\sqrt{6x^3 + 9x^2 + 18c_1}}{3} dx + c_3 \quad (2)$$

Verification of solutions

$$y = \int \frac{\sqrt{6x^3 + 9x^2 + 18c_1}}{3} dx + c_2$$

Verified OK.

$$y = \int -\frac{\sqrt{6x^3 + 9x^2 + 18c_1}}{3} dx + c_3$$

Verified OK.

2.18.2 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p(x)p'(x) - x^2 - x = 0$$

Which is now solve for $p(x)$ as first order ode. In canonical form the ODE is

$$\begin{aligned} p' &= F(x, p) \\ &= f(x)g(p) \\ &= \frac{x(1+x)}{p} \end{aligned}$$

Where $f(x) = x(1+x)$ and $g(p) = \frac{1}{p}$. Integrating both sides gives

$$\begin{aligned} \frac{1}{p} dp &= x(1+x) dx \\ \int \frac{1}{p} dp &= \int x(1+x) dx \\ \frac{p^2}{2} &= \frac{1}{3}x^3 + \frac{1}{2}x^2 + c_1 \end{aligned}$$

The solution is

$$\frac{p(x)^2}{2} - \frac{x^3}{3} - \frac{x^2}{2} - c_1 = 0$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$\frac{y'^2}{2} - \frac{x^3}{3} - \frac{x^2}{2} - c_1 = 0$$

Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \frac{\sqrt{6x^3 + 9x^2 + 18c_1}}{3} \quad (1)$$

$$y' = -\frac{\sqrt{6x^3 + 9x^2 + 18c_1}}{3} \quad (2)$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\begin{aligned} y &= \int \frac{\sqrt{6x^3 + 9x^2 + 18c_1}}{3} dx \\ &= \int \frac{\sqrt{6x^3 + 9x^2 + 18c_1}}{3} dx + c_2 \end{aligned}$$

Solving equation (2)

Integrating both sides gives

$$\begin{aligned} y &= \int -\frac{\sqrt{6x^3 + 9x^2 + 18c_1}}{3} dx \\ &= \int -\frac{\sqrt{6x^3 + 9x^2 + 18c_1}}{3} dx + c_3 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \int \frac{\sqrt{6x^3 + 9x^2 + 18c_1}}{3} dx + c_2 \quad (1)$$

$$y = \int -\frac{\sqrt{6x^3 + 9x^2 + 18c_1}}{3} dx + c_3 \quad (2)$$

Verification of solutions

$$y = \int \frac{\sqrt{6x^3 + 9x^2 + 18c_1}}{3} dx + c_2$$

Verified OK.

$$y = \int -\frac{\sqrt{6x^3 + 9x^2 + 18c_1}}{3} dx + c_3$$

Verified OK.

2.18.3 Solving as exact nonlinear second order ode

An exact non-linear second order ode has the form

$$a_2(x, y, y') y'' + a_1(x, y, y') y' + a_0(x, y, y') = 0$$

Where the following conditions are satisfied

$$\begin{aligned}\frac{\partial a_2}{\partial y} &= \frac{\partial a_1}{\partial y'} \\ \frac{\partial a_2}{\partial x} &= \frac{\partial a_0}{\partial y'} \\ \frac{\partial a_1}{\partial x} &= \frac{\partial a_0}{\partial y}\end{aligned}$$

Looking at the the ode given we see that

$$\begin{aligned}a_2 &= y' \\ a_1 &= 0 \\ a_0 &= -x^2 - x\end{aligned}$$

Applying the conditions to the above shows this is a nonlinear exact second order ode. Therefore it can be reduced to first order ode given by

$$\begin{aligned}\int a_2 dy' + \int a_1 dy + \int a_0 dx &= c_1 \\ \int y' dy' + \int 0 dy + \int -x^2 - x dx &= c_1\end{aligned}$$

Which results in

$$-\frac{x^3}{3} - \frac{x^2}{2} + \frac{y'^2}{2} = c_1$$

Which is now solved Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \frac{\sqrt{6x^3 + 9x^2 + 18c_1}}{3} \quad (1)$$

$$y' = -\frac{\sqrt{6x^3 + 9x^2 + 18c_1}}{3} \quad (2)$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\begin{aligned}y &= \int \frac{\sqrt{6x^3 + 9x^2 + 18c_1}}{3} dx \\ &= \int \frac{\sqrt{6x^3 + 9x^2 + 18c_1}}{3} dx + c_2\end{aligned}$$

Solving equation (2)

Integrating both sides gives

$$\begin{aligned}y &= \int -\frac{\sqrt{6x^3 + 9x^2 + 18c_1}}{3} dx \\ &= \int -\frac{\sqrt{6x^3 + 9x^2 + 18c_1}}{3} dx + c_3\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \int \frac{\sqrt{6x^3 + 9x^2 + 18c_1}}{3} dx + c_2 \quad (1)$$

$$y = \int -\frac{\sqrt{6x^3 + 9x^2 + 18c_1}}{3} dx + c_3 \quad (2)$$

Verification of solutions

$$y = \int \frac{\sqrt{6x^3 + 9x^2 + 18c_1}}{3} dx + c_2$$

Verified OK.

$$y = \int -\frac{\sqrt{6x^3 + 9x^2 + 18c_1}}{3} dx + c_3$$

Verified OK.

2.18.4 Maple step by step solution

Let's solve

$$y'y'' = x^2 + x$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Make substitution $u = y'$ to reduce order of ODE

$$u(x) u'(x) = x^2 + x$$

- Integrate both sides with respect to x

$$\int u(x) u'(x) dx = \int (x^2 + x) dx + c_1$$

- Evaluate integral

$$\frac{u(x)^2}{2} = \frac{1}{3}x^3 + \frac{1}{2}x^2 + c_1$$

- Solve for $u(x)$

$$\left\{ u(x) = -\frac{\sqrt{6x^3+9x^2+18c_1}}{3}, u(x) = \frac{\sqrt{6x^3+9x^2+18c_1}}{3} \right\}$$

- Solve 1st ODE for $u(x)$

$$u(x) = -\frac{\sqrt{6x^3+9x^2+18c_1}}{3}$$

- Make substitution $u = y'$

$$y' = -\frac{\sqrt{6x^3+9x^2+18c_1}}{3}$$

- Integrate both sides to solve for y

$$\int y' dx = \int -\frac{\sqrt{6x^3+9x^2+18c_1}}{3} dx + c_2$$

- Compute integrals

$$12Ic_1\sqrt{3} \left(\frac{\left(\frac{-1-12c_1+2\sqrt{36c_1^2+6c_1}}{2} \right)^{\frac{1}{3}}}{2} - \frac{1}{2\left(\frac{-1-12c_1+2\sqrt{36c_1^2+6c_1}}{2} \right)^{\frac{1}{3}}} \right) \sqrt{\left(\begin{array}{l} -1 \\ x+ \end{array} \right)}$$

$$y = -\frac{2x\sqrt{6x^3+9x^2+18c_1}}{15} - \frac{\sqrt{6x^3+9x^2+18c_1}}{15} - \dots$$

- Solve 2nd ODE for $u(x)$

$$u(x) = \frac{\sqrt{6x^3+9x^2+18c_1}}{3}$$

- Make substitution $u = y'$

$$y' = \frac{\sqrt{6x^3+9x^2+18c_1}}{3}$$

- Integrate both sides to solve for y

$$\int y' dx = \int \frac{\sqrt{6x^3+9x^2+18c_1}}{3} dx + c_2$$

- Compute integrals

$$12Ic_1\sqrt{3} \left(\frac{\left(\frac{-1-12c_1+2\sqrt{36c_1^2+6c_1}}{2} \right)^{\frac{1}{3}}}{2 \left(-1-12c_1+2\sqrt{36c_1^2+6c_1} \right)^{\frac{1}{3}}} - \frac{1}{2 \left(-1-12c_1+2\sqrt{36c_1^2+6c_1} \right)^{\frac{1}{3}}} \right) \sqrt{-1-x+\dots}$$

$$y = \frac{2x\sqrt{6x^3+9x^2+18c_1}}{15} + \frac{\sqrt{6x^3+9x^2+18c_1}}{15} + \dots$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
`, `-> Computing symmetries using: way = exp_sym
-> Calling odsolve with the ODE`, diff(_b(_a), _a) = _a*(_a+1)/_b(_a), _b(_a)
  Methods for first order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  trying 1st order linear
  trying Bernoulli
  <- Bernoulli successful
<- differential order: 2; canonical coordinates successful
<- differential order 2; missing variables successful`
```

*** Subleve

✓ Solution by Maple

Time used: 0.094 (sec). Leaf size: 51

```
dsolve(diff(y(x),x$2)*diff(y(x),x)=x*(1+x),y(x), singsol=all)
```

$$y(x) = -\frac{\left(\int \sqrt{6x^3 + 9x^2 + 9c_1} dx\right)}{3} + c_2$$
$$y(x) = \frac{\left(\int \sqrt{6x^3 + 9x^2 + 9c_1} dx\right)}{3} + c_2$$

✓ Solution by Mathematica

Time used: 61.466 (sec). Leaf size: 12885

```
DSolve[y''[x]*y'[x]==x*(1+x),y[x],x,IncludeSingularSolutions -> True]
```

Too large to display

3 Chapter 1. What is a differential equation.
Section 1.4 First Order Linear Equations. Page
15

3.1	problem 1(a)	318
3.2	problem 1(b)	322
3.3	problem 1(c)	326
3.4	problem 1(d)	331
3.5	problem 1(e)	335
3.6	problem 1(f)	339
3.7	problem 1(g)	343
3.8	problem 1(h)	347
3.9	problem 1(i)	352
3.10	problem 1(j)	356
3.11	problem 2(a)	361
3.12	problem 2(b)	365
3.13	problem 2(c)	370
3.14	problem 2(d)	373
3.15	problem 2(e)	378
3.16	problem 2(f)	383
3.17	problem 3(a)	388
3.18	problem 3(b)	402
3.19	problem 3(c)	416
3.20	problem 3(d)	433
3.21	problem 4(a)	449
3.22	problem 4(b)	455
3.23	problem 4(c)	467
3.24	problem 6	478
3.25	problem 7	493

3.1 problem 1(a)

3.1.1 Solving as linear ode	318
3.1.2 Maple step by step solution	320

Internal problem ID [6157]

Internal file name [OUTPUT/5405_Sunday_June_05_2022_03_36_26_PM_99668702/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.4 First Order Linear Equations. Page 15

Problem number: 1(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

`[_separable]`

$$y' - xy = 0$$

3.1.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -x$$

$$q(x) = 0$$

Hence the ode is

$$y' - xy = 0$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int -x dx} \\ &= e^{-\frac{x^2}{2}}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}\mu y &= 0 \\ \frac{d}{dx}\left(e^{-\frac{x^2}{2}}y\right) &= 0\end{aligned}$$

Integrating gives

$$e^{-\frac{x^2}{2}}y = c_1$$

Dividing both sides by the integrating factor $\mu = e^{-\frac{x^2}{2}}$ results in

$$y = c_1 e^{\frac{x^2}{2}}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{\frac{x^2}{2}} \tag{1}$$




Figure 69: Slope field plot

Verification of solutions

$$y = c_1 e^{\frac{x^2}{2}}$$

Verified OK.

3.1.2 Maple step by step solution

Let's solve

$$y' - xy = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y} = x$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y} dx = \int x dx + c_1$$

- Evaluate integral

$$\ln(y) = \frac{x^2}{2} + c_1$$

- Solve for y

$$y = e^{\frac{x^2}{2} + c_1}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)-x*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{\frac{x^2}{2}} c_1$$

✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 22

```
DSolve[y'[x]-x*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{\frac{x^2}{2}}$$

$$y(x) \rightarrow 0$$

3.2 problem 1(b)

3.2.1 Solving as linear ode	322
3.2.2 Maple step by step solution	324

Internal problem ID [6158]

Internal file name [OUTPUT/5406_Sunday_June_05_2022_03_36_27_PM_50883306/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.4 First Order Linear Equations. Page 15

Problem number: 1(b).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

`[_separable]`

$$y' + xy = x$$

3.2.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = x$$

$$q(x) = x$$

Hence the ode is

$$y' + xy = x$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int x dx} \\ &= e^{\frac{x^2}{2}}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu)(x) \\ \frac{d}{dx}\left(e^{\frac{x^2}{2}} y\right) &= \left(e^{\frac{x^2}{2}}\right)(x) \\ d\left(e^{\frac{x^2}{2}} y\right) &= \left(e^{\frac{x^2}{2}} x\right) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}e^{\frac{x^2}{2}} y &= \int e^{\frac{x^2}{2}} x dx \\ e^{\frac{x^2}{2}} y &= e^{\frac{x^2}{2}} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{\frac{x^2}{2}}$ results in

$$y = e^{-\frac{x^2}{2}} e^{\frac{x^2}{2}} + c_1 e^{-\frac{x^2}{2}}$$

which simplifies to

$$y = 1 + c_1 e^{-\frac{x^2}{2}}$$

Summary

The solution(s) found are the following

$$y = 1 + c_1 e^{-\frac{x^2}{2}} \tag{1}$$




Figure 70: Slope field plot

Verification of solutions

$$y = 1 + c_1 e^{-\frac{x^2}{2}}$$

Verified OK.

3.2.2 Maple step by step solution

Let's solve

$$y' + xy = x$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y-1} = -x$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y-1} dx = \int -x dx + c_1$$

- Evaluate integral

$$\ln(y - 1) = -\frac{x^2}{2} + c_1$$

- Solve for y

$$y = e^{-\frac{x^2}{2} + c_1} + 1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(diff(y(x),x)+x*y(x)=x,y(x), singsol=all)
```

$$y(x) = 1 + e^{-\frac{x^2}{2}} c_1$$

✓ Solution by Mathematica

Time used: 0.056 (sec). Leaf size: 24

```
DSolve[y'[x]+x*y[x]==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 1 + c_1 e^{-\frac{x^2}{2}}$$

$$y(x) \rightarrow 1$$

3.3 problem 1(c)

3.3.1 Solving as linear ode	326
3.3.2 Maple step by step solution	328

Internal problem ID [6159]

Internal file name [OUTPUT/5407_Sunday_June_05_2022_03_36_28_PM_85180385/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.4 First Order Linear Equations. Page 15

Problem number: 1(c).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

`[_linear]`

$$y + y' = \frac{1}{e^{2x} + 1}$$

3.3.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = 1$$
$$q(x) = \frac{1}{e^{2x} + 1}$$

Hence the ode is

$$y + y' = \frac{1}{e^{2x} + 1}$$

The integrating factor μ is

$$\mu = e^{\int 1 dx}$$
$$= e^x$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu) \left(\frac{1}{e^{2x} + 1} \right) \\ \frac{d}{dx}(y e^x) &= (e^x) \left(\frac{1}{e^{2x} + 1} \right) \\ d(y e^x) &= \left(\frac{e^x}{e^{2x} + 1} \right) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}y e^x &= \int \frac{e^x}{e^{2x} + 1} dx \\ y e^x &= \arctan(e^x) + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^x$ results in

$$y = e^{-x} \arctan(e^x) + c_1 e^{-x}$$

which simplifies to

$$y = e^{-x}(\arctan(e^x) + c_1)$$

Summary

The solution(s) found are the following

$$y = e^{-x}(\arctan(e^x) + c_1) \tag{1}$$




Figure 71: Slope field plot

Verification of solutions

$$y = e^{-x}(\arctan(e^x) + c_1)$$

Verified OK.

3.3.2 Maple step by step solution

Let's solve

$$y + y' = \frac{1}{e^{2x}+1}$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = -y + \frac{1}{e^{2x}+1}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y + y' = \frac{1}{e^{2x}+1}$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x)(y + y') = \frac{\mu(x)}{e^{2x+1}}$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x)y)$

$$\mu(x)(y + y') = \mu'(x)y + \mu(x)y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = \mu(x)$$

- Solve to find the integrating factor

$$\mu(x) = e^x$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x)y) \right) dx = \int \frac{\mu(x)}{e^{2x+1}} dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x)y = \int \frac{\mu(x)}{e^{2x+1}} dx + c_1$$

- Solve for y

$$y = \frac{\int \frac{\mu(x)}{e^{2x+1}} dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = e^x$

$$y = \frac{\int \frac{e^x}{e^{2x+1}} dx + c_1}{e^x}$$

- Evaluate the integrals on the rhs

$$y = \frac{\arctan(e^x) + c_1}{e^x}$$

- Simplify

$$y = e^{-x}(\arctan(e^x) + c_1)$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 14

```
dsolve(diff(y(x),x)+y(x)=1/(1+exp(2*x)),y(x), singsol=all)
```

$$y(x) = (\arctan(e^x) + c_1) e^{-x}$$

✓ Solution by Mathematica

Time used: 0.061 (sec). Leaf size: 18

```
DSolve[y'[x]+y[x]==1/(1+Exp[2*x]),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x}(\arctan(e^x) + c_1)$$

3.4 problem 1(d)

3.4.1 Solving as linear ode	331
3.4.2 Maple step by step solution	333

Internal problem ID [6160]

Internal file name [OUTPUT/5408_Sunday_June_05_2022_03_36_30_PM_7315470/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.4 First Order Linear Equations. Page 15

Problem number: 1(d).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

```
[[_linear, `class A`]]
```

$$y + y' = 2x e^{-x} + x^2$$

3.4.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = 1$$

$$q(x) = x(2e^{-x} + x)$$

Hence the ode is

$$y + y' = x(2e^{-x} + x)$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int 1 dx} \\ &= e^x\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu) (x(2e^{-x} + x)) \\ \frac{d}{dx}(y e^x) &= (e^x) (x(2e^{-x} + x)) \\ d(y e^x) &= (x(x e^x + 2)) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}y e^x &= \int x(x e^x + 2) dx \\ y e^x &= x^2 e^x - 2x e^x + 2e^x + x^2 + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^x$ results in

$$y = e^{-x}(x^2 e^x - 2x e^x + 2e^x + x^2) + c_1 e^{-x}$$

which simplifies to

$$y = (x^2 + c_1) e^{-x} + x^2 - 2x + 2$$

Summary

The solution(s) found are the following

$$y = (x^2 + c_1) e^{-x} + x^2 - 2x + 2 \tag{1}$$




Figure 72: Slope field plot

Verification of solutions

$$y = (x^2 + c_1) e^{-x} + x^2 - 2x + 2$$

Verified OK.

3.4.2 Maple step by step solution

Let's solve

$$y + y' = 2x e^{-x} + x^2$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = -y + 2x e^{-x} + x^2$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y + y' = 2x e^{-x} + x^2$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) (y + y') = \mu(x) (2x e^{-x} + x^2)$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x) y)$

$$\mu(x) (y + y') = \mu'(x) y + \mu(x) y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = \mu(x)$$

- Solve to find the integrating factor

$$\mu(x) = e^x$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x) y) \right) dx = \int \mu(x) (2x e^{-x} + x^2) dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x) y = \int \mu(x) (2x e^{-x} + x^2) dx + c_1$$

- Solve for y

$$y = \frac{\int \mu(x) (2x e^{-x} + x^2) dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = e^x$

$$y = \frac{\int e^x (2x e^{-x} + x^2) dx + c_1}{e^x}$$

- Evaluate the integrals on the rhs

$$y = \frac{x^2 e^x - 2x e^x + 2e^x + x^2 + c_1}{e^x}$$

- Simplify

$$y = (x^2 + c_1) e^{-x} + x^2 - 2x + 2$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 22

```
dsolve(diff(y(x),x)+y(x)=2*x*exp(-x)+x^2,y(x), singsol=all)
```

$$y(x) = (x^2 + c_1) e^{-x} + x^2 - 2x + 2$$

✓ Solution by Mathematica

Time used: 0.069 (sec). Leaf size: 29

```
DSolve[y'[x]+y[x]==2*x*Exp[-x]+x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x}(x^2 + e^x(x^2 - 2x + 2) + c_1)$$

3.5 problem 1(e)

3.5.1 Solving as linear ode	335
3.5.2 Maple step by step solution	337

Internal problem ID [6161]

Internal file name [OUTPUT/5409_Sunday_June_05_2022_03_36_31_PM_35181252/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.4 First Order Linear Equations. Page 15

Problem number: 1(e).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

[_linear]

$$-xy' + 2y = x^3$$

3.5.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -\frac{2}{x}$$
$$q(x) = -x^2$$

Hence the ode is

$$y' - \frac{2y}{x} = -x^2$$

The integrating factor μ is

$$\mu = e^{\int -\frac{2}{x} dx}$$
$$= \frac{1}{x^2}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu) (-x^2) \\ \frac{d}{dx}\left(\frac{y}{x^2}\right) &= \left(\frac{1}{x^2}\right) (-x^2) \\ d\left(\frac{y}{x^2}\right) &= -1 dx\end{aligned}$$

Integrating gives

$$\begin{aligned}\frac{y}{x^2} &= \int -1 dx \\ \frac{y}{x^2} &= -x + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = \frac{1}{x^2}$ results in

$$y = c_1 x^2 - x^3$$

which simplifies to

$$y = x^2(-x + c_1)$$

Summary

The solution(s) found are the following

$$y = x^2(-x + c_1) \tag{1}$$




Figure 73: Slope field plot

Verification of solutions

$$y = x^2(-x + c_1)$$

Verified OK.

3.5.2 Maple step by step solution

Let's solve

$$-xy' + 2y = x^3$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = \frac{2y}{x} - x^2$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' - \frac{2y}{x} = -x^2$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) \left(y' - \frac{2y}{x} \right) = -\mu(x) x^2$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x)y)$

$$\mu(x) \left(y' - \frac{2y}{x} \right) = \mu'(x)y + \mu(x)y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = -\frac{2\mu(x)}{x}$$

- Solve to find the integrating factor

$$\mu(x) = \frac{1}{x^2}$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x)y) \right) dx = \int -\mu(x)x^2 dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x)y = \int -\mu(x)x^2 dx + c_1$$

- Solve for y

$$y = \frac{\int -\mu(x)x^2 dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = \frac{1}{x^2}$

$$y = x^2 \left(\int (-1) dx + c_1 \right)$$

- Evaluate the integrals on the rhs

$$y = x^2(-x + c_1)$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(2*y(x)-x^3=x*diff(y(x),x),y(x), singsol=all)
```

$$y(x) = (-x + c_1) x^2$$

✓ Solution by Mathematica

Time used: 0.029 (sec). Leaf size: 15

```
DSolve[2*y[x]-x^3==x*y'[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^2(-x + c_1)$$

3.6 problem 1(f)

3.6.1 Solving as linear ode	339
3.6.2 Maple step by step solution	341

Internal problem ID [6162]

Internal file name [OUTPUT/5410_Sunday_June_05_2022_03_36_32_PM_80153340/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.4 First Order Linear Equations. Page 15

Problem number: 1(f).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

`[_separable]`

$$y' + 2xy = 0$$

3.6.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = 2x$$

$$q(x) = 0$$

Hence the ode is

$$y' + 2xy = 0$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int 2x dx} \\ &= e^{x^2}\end{aligned}$$

The ode becomes

$$\frac{d}{dx}\mu y = 0$$
$$\frac{d}{dx}(e^{x^2}y) = 0$$

Integrating gives

$$e^{x^2}y = c_1$$

Dividing both sides by the integrating factor $\mu = e^{x^2}$ results in

$$y = c_1 e^{-x^2}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-x^2} \tag{1}$$




Figure 74: Slope field plot

Verification of solutions

$$y = c_1 e^{-x^2}$$

Verified OK.

3.6.2 Maple step by step solution

Let's solve

$$y' + 2xy = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y} = -2x$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y} dx = \int -2x dx + c_1$$

- Evaluate integral

$$\ln(y) = -x^2 + c_1$$

- Solve for y

$$y = e^{-x^2+c_1}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)+2*x*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{-x^2} c_1$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 20

```
DSolve[y'[x]+2*x*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-x^2}$$

$$y(x) \rightarrow 0$$

3.7 problem 1(g)

3.7.1 Solving as linear ode	343
3.7.2 Maple step by step solution	345

Internal problem ID [6163]

Internal file name [OUTPUT/5411_Sunday_June_05_2022_03_36_33_PM_54552225/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.4 First Order Linear Equations.

Page 15

Problem number: 1(g).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

[_linear]

$$xy' - 3y = x^4$$

3.7.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -\frac{3}{x}$$
$$q(x) = x^3$$

Hence the ode is

$$y' - \frac{3y}{x} = x^3$$

The integrating factor μ is

$$\mu = e^{\int -\frac{3}{x} dx}$$
$$= \frac{1}{x^3}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu)(x^3) \\ \frac{d}{dx}\left(\frac{y}{x^3}\right) &= \left(\frac{1}{x^3}\right)(x^3) \\ d\left(\frac{y}{x^3}\right) &= dx\end{aligned}$$

Integrating gives

$$\begin{aligned}\frac{y}{x^3} &= \int dx \\ \frac{y}{x^3} &= x + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = \frac{1}{x^3}$ results in

$$y = c_1 x^3 + x^4$$

which simplifies to

$$y = x^3(x + c_1)$$

Summary

The solution(s) found are the following

$$y = x^3(x + c_1) \tag{1}$$




Figure 75: Slope field plot

Verification of solutions

$$y = x^3(x + c_1)$$

Verified OK.

3.7.2 Maple step by step solution

Let's solve

$$xy' - 3y = x^4$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = \frac{3y}{x} + x^3$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' - \frac{3y}{x} = x^3$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) \left(y' - \frac{3y}{x} \right) = \mu(x) x^3$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x)y)$

$$\mu(x) \left(y' - \frac{3y}{x} \right) = \mu'(x)y + \mu(x)y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = -\frac{3\mu(x)}{x}$$

- Solve to find the integrating factor

$$\mu(x) = \frac{1}{x^3}$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x)y) \right) dx = \int \mu(x) x^3 dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x)y = \int \mu(x) x^3 dx + c_1$$

- Solve for y

$$y = \frac{\int \mu(x) x^3 dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = \frac{1}{x^3}$

$$y = x^3 \left(\int 1 dx + c_1 \right)$$

- Evaluate the integrals on the rhs

$$y = x^3(x + c_1)$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve(x*diff(y(x),x)-3*y(x)=x^4,y(x), singsol=all)
```

$$y(x) = (x + c_1) x^3$$

✓ Solution by Mathematica

Time used: 0.029 (sec). Leaf size: 13

```
DSolve[x*y'[x]-3*y[x]==x^4,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^3(x + c_1)$$

3.8 problem 1(h)

3.8.1 Solving as linear ode	347
3.8.2 Maple step by step solution	349

Internal problem ID [6164]

Internal file name [OUTPUT/5412_Sunday_June_05_2022_03_36_35_PM_43760870/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.4 First Order Linear Equations. Page 15

Problem number: 1(h).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

`[_linear]`

$$(x^2 + 1) y' + 2xy = \cot(x)$$

3.8.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = \frac{2x}{x^2 + 1}$$
$$q(x) = \frac{\cot(x)}{x^2 + 1}$$

Hence the ode is

$$y' + \frac{2xy}{x^2 + 1} = \frac{\cot(x)}{x^2 + 1}$$

The integrating factor μ is

$$\mu = e^{\int \frac{2x}{x^2+1} dx}$$
$$= x^2 + 1$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu) \left(\frac{\cot(x)}{x^2 + 1} \right) \\ \frac{d}{dx}((x^2 + 1)y) &= (x^2 + 1) \left(\frac{\cot(x)}{x^2 + 1} \right) \\ d((x^2 + 1)y) &= \cot(x) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}(x^2 + 1)y &= \int \cot(x) dx \\ (x^2 + 1)y &= \ln(\sin(x)) + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = x^2 + 1$ results in

$$y = \frac{\ln(\sin(x))}{x^2 + 1} + \frac{c_1}{x^2 + 1}$$

which simplifies to

$$y = \frac{\ln(\sin(x)) + c_1}{x^2 + 1}$$

Summary

The solution(s) found are the following

$$y = \frac{\ln(\sin(x)) + c_1}{x^2 + 1} \tag{1}$$




Figure 76: Slope field plot

Verification of solutions

$$y = \frac{\ln(\sin(x)) + c_1}{x^2 + 1}$$

Verified OK.

3.8.2 Maple step by step solution

Let's solve

$$(x^2 + 1)y' + 2xy = \cot(x)$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = -\frac{2xy}{x^2+1} + \frac{\cot(x)}{x^2+1}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' + \frac{2xy}{x^2+1} = \frac{\cot(x)}{x^2+1}$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) \left(y' + \frac{2xy}{x^2+1} \right) = \frac{\mu(x) \cot(x)}{x^2+1}$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x) y)$

$$\mu(x) \left(y' + \frac{2xy}{x^2+1} \right) = \mu'(x) y + \mu(x) y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = \frac{2\mu(x)x}{x^2+1}$$

- Solve to find the integrating factor

$$\mu(x) = x^2 + 1$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x) y) \right) dx = \int \frac{\mu(x) \cot(x)}{x^2+1} dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x) y = \int \frac{\mu(x) \cot(x)}{x^2+1} dx + c_1$$

- Solve for y

$$y = \frac{\int \frac{\mu(x) \cot(x)}{x^2+1} dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = x^2 + 1$

$$y = \frac{\int \cot(x) dx + c_1}{x^2+1}$$

- Evaluate the integrals on the rhs

$$y = \frac{\ln(\sin(x)) + c_1}{x^2+1}$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve((1+x^2)*diff(y(x),x)+2*x*y(x)=cot(x),y(x), singsol=all)
```

$$y(x) = \frac{\ln(\sin(x)) + c_1}{x^2 + 1}$$

✓ Solution by Mathematica

Time used: 0.06 (sec). Leaf size: 19

```
DSolve[(1+x^2)*y'[x]+2*x*y[x]==Cot[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\log(\sin(x)) + c_1}{x^2 + 1}$$

3.9 problem 1(i)

3.9.1 Solving as linear ode	352
3.9.2 Maple step by step solution	354

Internal problem ID [6165]

Internal file name [OUTPUT/5413_Sunday_June_05_2022_03_36_36_PM_27167940/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.4 First Order Linear Equations. Page 15

Problem number: 1(i).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

`[_linear]`

$$y' + y \cot(x) = 2x \csc(x)$$

3.9.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = \cot(x)$$

$$q(x) = 2x \csc(x)$$

Hence the ode is

$$y' + y \cot(x) = 2x \csc(x)$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int \cot(x) dx} \\ &= \sin(x) \end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu) (2x \csc(x)) \\ \frac{d}{dx}(y \sin(x)) &= (\sin(x)) (2x \csc(x)) \\ d(y \sin(x)) &= (2x) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}y \sin(x) &= \int 2x dx \\ y \sin(x) &= x^2 + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = \sin(x)$ results in

$$y = \csc(x) x^2 + c_1 \csc(x)$$

which simplifies to

$$y = \csc(x) (x^2 + c_1)$$

Summary

The solution(s) found are the following

$$y = \csc(x) (x^2 + c_1) \tag{1}$$




Figure 77: Slope field plot

Verification of solutions

$$y = \csc(x) (x^2 + c_1)$$

Verified OK.

3.9.2 Maple step by step solution

Let's solve

$$y' + y \cot(x) = 2x \csc(x)$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = -y \cot(x) + 2x \csc(x)$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' + y \cot(x) = 2x \csc(x)$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) (y' + y \cot(x)) = 2\mu(x) x \csc(x)$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x) y)$

$$\mu(x) (y' + y \cot(x)) = \mu'(x) y + \mu(x) y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = \mu(x) \cot(x)$$

- Solve to find the integrating factor

$$\mu(x) = \sin(x)$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x) y) \right) dx = \int 2\mu(x) x \csc(x) dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x) y = \int 2\mu(x) x \csc(x) dx + c_1$$

- Solve for y

$$y = \frac{\int 2\mu(x) x \csc(x) dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = \sin(x)$

$$y = \frac{\int 2x \csc(x) \sin(x) dx + c_1}{\sin(x)}$$

- Evaluate the integrals on the rhs

$$y = \frac{x^2 + c_1}{\sin(x)}$$

- Simplify

$$y = \csc(x) (x^2 + c_1)$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)+y(x)*cot(x)=2*x*csc(x),y(x), singsol=all)
```

$$y(x) = \csc(x) (x^2 + c_1)$$

✓ Solution by Mathematica

Time used: 0.048 (sec). Leaf size: 14

```
DSolve[y'[x]+y[x]*Cot[x]==2*x*Csc[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow (x^2 + c_1) \csc(x)$$

3.10 problem 1(j)

3.10.1 Solving as linear ode	356
3.10.2 Maple step by step solution	358

Internal problem ID [6166]

Internal file name [OUTPUT/5414_Sunday_June_05_2022_03_36_37_PM_36928189/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.4 First Order Linear Equations. Page 15

Problem number: 1(j).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

`[_linear]`

$$y + xy \cot(x) + xy' = x$$

3.10.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -\frac{-x \cot(x) - 1}{x}$$

$$q(x) = 1$$

Hence the ode is

$$y' - \frac{(-x \cot(x) - 1)y}{x} = 1$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int -\frac{-x \cot(x) - 1}{x} dx} \\ &= e^{\ln(\sin(x)) + \ln(x)} \end{aligned}$$

Which simplifies to

$$\mu = \sin(x) x$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= \mu \\ \frac{d}{dx}(\sin(x) xy) &= \sin(x) x \\ d(\sin(x) xy) &= \sin(x) x dx\end{aligned}$$

Integrating gives

$$\begin{aligned}\sin(x) xy &= \int \sin(x) x dx \\ \sin(x) xy &= \sin(x) - \cos(x) x + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = \sin(x) x$ results in

$$y = \frac{\csc(x) (\sin(x) - \cos(x) x)}{x} + \frac{c_1 \csc(x)}{x}$$

which simplifies to

$$y = \frac{1 - x \cot(x) + c_1 \csc(x)}{x}$$

Summary

The solution(s) found are the following

$$y = \frac{1 - x \cot(x) + c_1 \csc(x)}{x} \tag{1}$$




Figure 78: Slope field plot

Verification of solutions

$$y = \frac{1 - x \cot(x) + c_1 \csc(x)}{x}$$

Verified OK.

3.10.2 Maple step by step solution

Let's solve

$$y + xy \cot(x) + xy' = x$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = 1 - \frac{(x \cot(x) + 1)y}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' + \frac{(x \cot(x) + 1)y}{x} = 1$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) \left(y' + \frac{(x \cot(x) + 1)y}{x} \right) = \mu(x)$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x)y)$

$$\mu(x) \left(y' + \frac{(x \cot(x) + 1)y}{x} \right) = \mu'(x)y + \mu(x)y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = \frac{\mu(x)(x \cot(x) + 1)}{x}$$

- Solve to find the integrating factor

$$\mu(x) = \sin(x)x$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x)y) \right) dx = \int \mu(x) dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x)y = \int \mu(x) dx + c_1$$

- Solve for y

$$y = \frac{\int \mu(x) dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = \sin(x)x$

$$y = \frac{\int \sin(x)x dx + c_1}{\sin(x)x}$$

- Evaluate the integrals on the rhs

$$y = \frac{\sin(x) - \cos(x)x + c_1}{\sin(x)x}$$

- Simplify

$$y = \frac{1 - x \cot(x) + c_1 \csc(x)}{x}$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(y(x)-x+x*y(x)*cot(x)+x*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{-\cot(x)x + 1 + \csc(x)c_1}{x}$$

✓ Solution by Mathematica

Time used: 0.072 (sec). Leaf size: 21

```
DSolve[y[x]-x+x*y[x]*Cot[x]+x*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{-x \cot(x) + c_1 \csc(x) + 1}{x}$$

3.11 problem 2(a)

3.11.1 Existence and uniqueness analysis	361
3.11.2 Solving as linear ode	362
3.11.3 Maple step by step solution	363

Internal problem ID [6167]

Internal file name [OUTPUT/5415_Sunday_June_05_2022_03_36_39_PM_42878629/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.4 First Order Linear Equations. Page 15

Problem number: 2(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "linear", "separable", "homogeneousTypeD2", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_separable]

$$y' - xy = 0$$

With initial conditions

$$[y(1) = 3]$$

3.11.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -x$$

$$q(x) = 0$$

Hence the ode is

$$y' - xy = 0$$

The domain of $p(x) = -x$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 1$ is inside this domain. Hence solution exists and is unique.

3.11.2 Solving as linear ode

Entering Linear first order ODE solver. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int -x dx} \\ &= e^{-\frac{x^2}{2}}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}\mu y &= 0 \\ \frac{d}{dx}\left(e^{-\frac{x^2}{2}}y\right) &= 0\end{aligned}$$

Integrating gives

$$e^{-\frac{x^2}{2}}y = c_1$$

Dividing both sides by the integrating factor $\mu = e^{-\frac{x^2}{2}}$ results in

$$y = c_1 e^{\frac{x^2}{2}}$$

Initial conditions are used to solve for c_1 . Substituting $x = 1$ and $y = 3$ in the above solution gives an equation to solve for the constant of integration.

$$3 = c_1 e^{\frac{1}{2}}$$

$$c_1 = 3 e^{-\frac{1}{2}}$$


Substituting c_1 found above in the general solution gives

$$y = 3 e^{\frac{(x-1)(1+x)}{2}}$$


Summary

The solution(s) found are the following

$$y = 3e^{\frac{(x-1)(1+x)}{2}} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = 3e^{\frac{(x-1)(1+x)}{2}}$$

Verified OK.

3.11.3 Maple step by step solution

Let's solve

$$[y' - xy = 0, y(1) = 3]$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y} = x$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y} dx = \int x dx + c_1$$

- Evaluate integral
 $\ln(y) = \frac{x^2}{2} + c_1$
- Solve for y
 $y = e^{\frac{x^2}{2} + c_1}$
- Use initial condition $y(1) = 3$
 $3 = e^{\frac{1}{2} + c_1}$
- Solve for c_1
 $c_1 = -\frac{1}{2} + \ln(3)$
- Substitute $c_1 = -\frac{1}{2} + \ln(3)$ into general solution and simplify
 $y = 3e^{\frac{(x-1)(1+x)}{2}}$
- Solution to the IVP
 $y = 3e^{\frac{(x-1)(1+x)}{2}}$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 15

```
dsolve([diff(y(x),x)-x*y(x)=0,y(1) = 3],y(x), singsol=all)
```

$$y(x) = 3e^{\frac{(x-1)(x+1)}{2}}$$

✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 18

```
DSolve[{y'[x]-x*y[x]==0,{y[1]==3}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 3e^{\frac{1}{2}(x^2-1)}$$

3.12 problem 2(b)

3.12.1 Existence and uniqueness analysis	365
3.12.2 Solving as linear ode	366
3.12.3 Maple step by step solution	368

Internal problem ID [6168]

Internal file name [OUTPUT/5416_Sunday_June_05_2022_03_36_41_PM_87143776/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.4 First Order Linear Equations. Page 15

Problem number: 2(b).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "linear", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

`[_linear]`

$$-2xy + y' = 6x e^{x^2}$$

With initial conditions

$$[y(1) = 1]$$

3.12.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -2x$$

$$q(x) = 6x e^{x^2}$$

Hence the ode is

$$-2xy + y' = 6x e^{x^2}$$

The domain of $p(x) = -2x$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 1$ is inside this domain. The domain of $q(x) = 6x e^{x^2}$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 1$ is also inside this domain. Hence solution exists and is unique.

3.12.2 Solving as linear ode

Entering Linear first order ODE solver. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int -2x dx} \\ &= e^{-x^2}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu) (6x e^{x^2}) \\ \frac{d}{dx}(e^{-x^2} y) &= (e^{-x^2}) (6x e^{x^2}) \\ d(e^{-x^2} y) &= (6x) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}e^{-x^2} y &= \int 6x dx \\ e^{-x^2} y &= 3x^2 + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{-x^2}$ results in

$$y = 3x^2 e^{x^2} + c_1 e^{x^2}$$

which simplifies to

$$y = e^{x^2} (3x^2 + c_1)$$

Initial conditions are used to solve for c_1 . Substituting $x = 1$ and $y = 1$ in the above solution gives an equation to solve for the constant of integration.

$$1 = e(c_1 + 3)$$

$$c_1 = -(3e - 1)e^{-1}$$


Substituting c_1 found above in the general solution gives

$$y = 3x^2e^{x^2} + e^{(x-1)(1+x)} - 3e^{x^2}$$


Summary

The solution(s) found are the following

$$y = 3x^2e^{x^2} + e^{(x-1)(1+x)} - 3e^{x^2} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = 3x^2e^{x^2} + e^{(x-1)(1+x)} - 3e^{x^2}$$

Verified OK.

3.12.3 Maple step by step solution

Let's solve

$$\left[-2xy + y' = 6x e^{x^2}, y(1) = 1\right]$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = 2xy + 6x e^{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$-2xy + y' = 6x e^{x^2}$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) (-2xy + y') = 6\mu(x) x e^{x^2}$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x) y)$

$$\mu(x) (-2xy + y') = \mu'(x) y + \mu(x) y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = -2\mu(x) x$$

- Solve to find the integrating factor

$$\mu(x) = e^{-x^2}$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x) y)\right) dx = \int 6\mu(x) x e^{x^2} dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x) y = \int 6\mu(x) x e^{x^2} dx + c_1$$

- Solve for y

$$y = \frac{\int 6\mu(x) x e^{x^2} dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = e^{-x^2}$

$$y = \frac{\int 6x e^{-x^2} e^{x^2} dx + c_1}{e^{-x^2}}$$

- Evaluate the integrals on the rhs

$$y = \frac{3x^2 + c_1}{e^{-x^2}}$$

- Simplify

$$y = e^{x^2}(3x^2 + c_1)$$

- Use initial condition $y(1) = 1$

$$1 = e(c_1 + 3)$$

- Solve for c_1

$$c_1 = -\frac{3e-1}{e}$$

- Substitute $c_1 = -\frac{3e-1}{e}$ into general solution and simplify

$$y = (-3 + 3x^2 + e^{-1})e^{x^2}$$

- Solution to the IVP

$$y = (-3 + 3x^2 + e^{-1})e^{x^2}$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 18

```
dsolve([diff(y(x),x)-2*x*y(x)=6*x*exp(x^2),y(1) = 1],y(x), singsol=all)
```

$$y(x) = (3x^2 - 3 + e^{-1})e^{x^2}$$

✓ Solution by Mathematica

Time used: 0.058 (sec). Leaf size: 23

```
DSolve[{y'[x]-2*x*y[x]==6*x*Exp[x^2],{y[1]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{x^2-1}(3e(x^2 - 1) + 1)$$

3.13 problem 2(c)

3.13.1 Existence and uniqueness analysis	370
3.13.2 Solving as linear ode	371

Internal problem ID [6169]

Internal file name [OUTPUT/5417_Sunday_June_05_2022_03_36_43_PM_66708007/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.4 First Order Linear Equations. Page 15

Problem number: 2(c).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "linear", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_linear]

Unable to solve or complete the solution.

$$x \ln(x) y' + y = 3x^3$$

With initial conditions

$$[y(1) = 0]$$

3.13.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = \frac{1}{x \ln(x)}$$
$$q(x) = \frac{3x^2}{\ln(x)}$$

Hence the ode is

$$y' + \frac{y}{x \ln(x)} = \frac{3x^2}{\ln(x)}$$

The domain of $p(x) = \frac{1}{x \ln(x)}$ is

$$\{0 < x < 1, 1 < x \leq \infty\}$$

But the point $x_0 = 1$ is not inside this domain. Hence existence and uniqueness theorem does not apply. There could be infinite number of solutions, or one solution or no solution at all.

3.13.2 Solving as linear ode

Entering Linear first order ODE solver. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int \frac{1}{x \ln(x)} dx} \\ &= \ln(x)\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu) \left(\frac{3x^2}{\ln(x)} \right) \\ \frac{d}{dx}(\ln(x) y) &= (\ln(x)) \left(\frac{3x^2}{\ln(x)} \right) \\ d(\ln(x) y) &= (3x^2) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}\ln(x) y &= \int 3x^2 dx \\ \ln(x) y &= x^3 + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = \ln(x)$ results in

$$y = \frac{x^3}{\ln(x)} + \frac{c_1}{\ln(x)}$$

which simplifies to

$$y = \frac{x^3 + c_1}{\ln(x)}$$

Initial conditions are used to solve for c_1 . Substituting $x = 1$ and $y = 0$ in the above solution gives an equation to solve for the constant of integration. Warning: Unable to solve for constant of integration.

Verification of solutions N/A

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✗ Solution by Maple

```
dsolve([(x*ln(x))*diff(y(x),x)+y(x)=3*x^3,y(1) = 0],y(x), singsol=all)
```

No solution found

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[{(x*Log[x])*y'[x]+y[x]==3*x^3,{y[1]==0}},y[x],x,IncludeSingularSolutions -> True]
```

Not solved

3.14 problem 2(d)

3.14.1 Existence and uniqueness analysis	373
3.14.2 Solving as linear ode	374
3.14.3 Maple step by step solution	376

Internal problem ID [6170]

Internal file name [OUTPUT/5418_Sunday_June_05_2022_03_36_44_PM_70578026/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.4 First Order Linear Equations. Page 15

Problem number: 2(d).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "linear", "homogeneousTypeD2", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_linear]

$$y' - \frac{y}{x} = x^2$$

With initial conditions

$$[y(1) = 3]$$

3.14.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -\frac{1}{x}$$
$$q(x) = x^2$$

Hence the ode is

$$y' - \frac{y}{x} = x^2$$

The domain of $p(x) = -\frac{1}{x}$ is

$$\{x < 0 \vee 0 < x\}$$

And the point $x_0 = 1$ is inside this domain. The domain of $q(x) = x^2$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 1$ is also inside this domain. Hence solution exists and is unique.

3.14.2 Solving as linear ode

Entering Linear first order ODE solver. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int -\frac{1}{x} dx} \\ &= \frac{1}{x}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu)(x^2) \\ \frac{d}{dx}\left(\frac{y}{x}\right) &= \left(\frac{1}{x}\right)(x^2) \\ d\left(\frac{y}{x}\right) &= x dx\end{aligned}$$

Integrating gives

$$\begin{aligned}\frac{y}{x} &= \int x dx \\ \frac{y}{x} &= \frac{x^2}{2} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = \frac{1}{x}$ results in

$$y = \frac{1}{2}x^3 + c_1x$$

Initial conditions are used to solve for c_1 . Substituting $x = 1$ and $y = 3$ in the above solution gives an equation to solve for the constant of integration.

$$3 = \frac{1}{2} + c_1$$

$$c_1 = \frac{5}{2}$$


Substituting c_1 found above in the general solution gives

$$y = \frac{1}{2}x^3 + \frac{5}{2}x$$


Summary

The solution(s) found are the following

$$y = \frac{1}{2}x^3 + \frac{5}{2}x \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{1}{2}x^3 + \frac{5}{2}x$$

Verified OK.

3.14.3 Maple step by step solution

Let's solve

$$\left[y' - \frac{y}{x} = x^2, y(1) = 3 \right]$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = \frac{y}{x} + x^2$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' - \frac{y}{x} = x^2$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) \left(y' - \frac{y}{x} \right) = \mu(x) x^2$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x) y)$

$$\mu(x) \left(y' - \frac{y}{x} \right) = \mu'(x) y + \mu(x) y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = -\frac{\mu(x)}{x}$$

- Solve to find the integrating factor

$$\mu(x) = \frac{1}{x}$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x) y) \right) dx = \int \mu(x) x^2 dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x) y = \int \mu(x) x^2 dx + c_1$$

- Solve for y

$$y = \frac{\int \mu(x) x^2 dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = \frac{1}{x}$

$$y = x \left(\int x dx + c_1 \right)$$

- Evaluate the integrals on the rhs

$$y = x \left(\frac{x^2}{2} + c_1 \right)$$

- Simplify

$$y = \frac{x(x^2+2c_1)}{2}$$

- Use initial condition $y(1) = 3$

$$3 = \frac{1}{2} + c_1$$

- Solve for c_1

$$c_1 = \frac{5}{2}$$

- Substitute $c_1 = \frac{5}{2}$ into general solution and simplify

$$y = \frac{x(x^2+5)}{2}$$

- Solution to the IVP

$$y = \frac{x(x^2+5)}{2}$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve([diff(y(x),x)-y(x)/x=x^2,y(1) = 3],y(x), singsol=all)
```

$$y(x) = \frac{(x^2 + 5)x}{2}$$

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 15

```
DSolve[{y'[x]-y[x]/x==x^2,{y[1]==3}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}x(x^2 + 5)$$

3.15 problem 2(e)

3.15.1 Existence and uniqueness analysis	378
3.15.2 Solving as linear ode	379
3.15.3 Maple step by step solution	381

Internal problem ID [6171]

Internal file name [OUTPUT/5419_Sunday_June_05_2022_03_36_46_PM_67387346/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.4 First Order Linear Equations. Page 15

Problem number: 2(e).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "linear", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

```
[[_linear, `class A`]]
```

$$4y + y' = e^{-x}$$

With initial conditions

$$[y(0) = 0]$$

3.15.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = 4$$

$$q(x) = e^{-x}$$

Hence the ode is

$$4y + y' = e^{-x}$$

The domain of $p(x) = 4$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = e^{-x}$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

3.15.2 Solving as linear ode

Entering Linear first order ODE solver. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int 4dx} \\ &= e^{4x}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu)(e^{-x}) \\ \frac{d}{dx}(e^{4x}y) &= (e^{4x})(e^{-x}) \\ d(e^{4x}y) &= e^{3x} dx\end{aligned}$$

Integrating gives

$$\begin{aligned}e^{4x}y &= \int e^{3x} dx \\ e^{4x}y &= \frac{e^{3x}}{3} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{4x}$ results in

$$y = \frac{e^{-4x}e^{3x}}{3} + c_1e^{-4x}$$

which simplifies to

$$y = \frac{e^{-x}}{3} + c_1e^{-4x}$$

Initial conditions are used to solve for c_1 . Substituting $x = 0$ and $y = 0$ in the above solution gives an equation to solve for the constant of integration.

$$0 = \frac{1}{3} + c_1$$

$$c_1 = -\frac{1}{3}$$


Substituting c_1 found above in the general solution gives

$$y = \frac{e^{-x}}{3} - \frac{e^{-4x}}{3}$$


Summary

The solution(s) found are the following

$$y = \frac{e^{-x}}{3} - \frac{e^{-4x}}{3} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{e^{-x}}{3} - \frac{e^{-4x}}{3}$$

Verified OK.

3.15.3 Maple step by step solution

Let's solve

$$[4y + y' = e^{-x}, y(0) = 0]$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = -4y + e^{-x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$4y + y' = e^{-x}$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) (4y + y') = \mu(x) e^{-x}$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x) y)$

$$\mu(x) (4y + y') = \mu'(x) y + \mu(x) y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = 4\mu(x)$$

- Solve to find the integrating factor

$$\mu(x) = e^{4x}$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x) y) \right) dx = \int \mu(x) e^{-x} dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x) y = \int \mu(x) e^{-x} dx + c_1$$

- Solve for y

$$y = \frac{\int \mu(x) e^{-x} dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = e^{4x}$

$$y = \frac{\int e^{-x} e^{4x} dx + c_1}{e^{4x}}$$

- Evaluate the integrals on the rhs

$$y = \frac{\frac{e^{3x}}{3} + c_1}{e^{4x}}$$

- Simplify

$$y = \frac{(e^{3x} + 3c_1)e^{-4x}}{3}$$

- Use initial condition $y(0) = 0$
 $0 = \frac{1}{3} + c_1$
- Solve for c_1
 $c_1 = -\frac{1}{3}$
- Substitute $c_1 = -\frac{1}{3}$ into general solution and simplify
 $y = \frac{(e^{3x}-1)e^{-4x}}{3}$
- Solution to the IVP
 $y = \frac{(e^{3x}-1)e^{-4x}}{3}$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 16

```
dsolve([diff(y(x),x)+4*y(x)=exp(-x),y(0) = 0],y(x), singsol=all)
```

$$y(x) = \frac{(e^{3x} - 1)e^{-4x}}{3}$$

✓ Solution by Mathematica

Time used: 0.056 (sec). Leaf size: 21

```
DSolve[{y'[x]+4*y[x]==Exp[-x],{y[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{3}e^{-4x}(e^{3x} - 1)$$

3.16 problem 2(f)

3.16.1 Existence and uniqueness analysis	383
3.16.2 Solving as linear ode	384
3.16.3 Maple step by step solution	385

Internal problem ID [6172]

Internal file name [OUTPUT/5420_Sunday_June_05_2022_03_36_48_PM_64201154/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.4 First Order Linear Equations. Page 15

Problem number: 2(f).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "linear", "separable", "differentialType", "homogeneousTypeMapleC", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_separable]

$$x^2y' + xy = 2x$$

With initial conditions

$$[y(1) = 1]$$

3.16.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = \frac{1}{x}$$
$$q(x) = \frac{2}{x}$$

Hence the ode is

$$y' + \frac{y}{x} = \frac{2}{x}$$

The domain of $p(x) = \frac{1}{x}$ is

$$\{x < 0 \vee 0 < x\}$$

And the point $x_0 = 1$ is inside this domain. The domain of $q(x) = \frac{2}{x}$ is

$$\{x < 0 \vee 0 < x\}$$

And the point $x_0 = 1$ is also inside this domain. Hence solution exists and is unique.

3.16.2 Solving as linear ode

Entering Linear first order ODE solver. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int \frac{1}{x} dx} \\ &= x\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu) \left(\frac{2}{x}\right) \\ \frac{d}{dx}(xy) &= (x) \left(\frac{2}{x}\right) \\ d(xy) &= 2 dx\end{aligned}$$

Integrating gives

$$\begin{aligned}xy &= \int 2 dx \\ xy &= 2x + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = x$ results in

$$y = 2 + \frac{c_1}{x}$$

Initial conditions are used to solve for c_1 . Substituting $x = 1$ and $y = 1$ in the above solution gives an equation to solve for the constant of integration.

$$1 = 2 + c_1$$

$$c_1 = -1$$


Substituting c_1 found above in the general solution gives

$$y = \frac{2x - 1}{x}$$


Summary

The solution(s) found are the following

$$y = \frac{2x - 1}{x} \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{2x - 1}{x}$$

Verified OK.

3.16.3 Maple step by step solution

Let's solve

$$[x^2y' + xy = 2x, y(1) = 1]$$

- Highest derivative means the order of the ODE is 1

y'

- Separate variables

$$\frac{y'}{y-2} = -\frac{1}{x}$$
- Integrate both sides with respect to x

$$\int \frac{y'}{y-2} dx = \int -\frac{1}{x} dx + c_1$$
- Evaluate integral

$$\ln(y-2) = -\ln(x) + c_1$$
- Solve for y

$$y = \frac{e^{c_1+2x}}{x}$$
- Use initial condition $y(1) = 1$

$$1 = e^{c_1} + 2$$
- Solve for c_1

$$c_1 = \ln(-1)$$
- Substitute $c_1 = \ln(-1)$ into general solution and simplify

$$y = \frac{2x-1}{x}$$
- Solution to the IVP

$$y = \frac{2x-1}{x}$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 13

```
dsolve([x^2*diff(y(x),x)+x*y(x)=2*x,y(1) = 1],y(x), singsol=all)
```

$$y(x) = \frac{2x-1}{x}$$

✓ Solution by Mathematica

Time used: 0.039 (sec). Leaf size: 12

```
DSolve[{x^2*y'[x]+x*y[x]==2*x,{y[1]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2 - \frac{1}{x}$$

3.17 problem 3(a)

3.17.1 Solving as first order ode lie symmetry lookup ode	388
3.17.2 Solving as bernoulli ode	392
3.17.3 Solving as exact ode	396

Internal problem ID [6173]

Internal file name [OUTPUT/5421_Sunday_June_05_2022_03_36_49_PM_92467487/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.4 First Order Linear Equations. Page 15

Problem number: 3(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "bernoulli", "exactWithIntegrationFactor", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

```
[[_homogeneous, `class G`], _rational, _Bernoulli]
```

$xy' + y - y^3x^4 = 0$

3.17.1 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{y(y^2x^4 - 1)}{x}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2\xi_y - \omega_x\xi - \omega_y\eta = 0 \tag{A}$$

The type of this ode is known. It is of type Bernoulli. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 71: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= y^3x^2\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{y^3 x^2} dy \end{aligned}$$

Which results in

$$S = -\frac{1}{2y^2 x^2}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{y(y^2 x^4 - 1)}{x}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{1}{y^2 x^3} \\ S_y &= \frac{1}{y^3 x^2} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = x \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = R$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \frac{R^2}{2} + c_1 \quad (4)$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$-\frac{1}{2y^2x^2} = \frac{x^2}{2} + c_1$$

Which simplifies to

$$-\frac{1}{2y^2x^2} = \frac{x^2}{2} + c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{y(y^2x^4-1)}{x}$ 	$R = x$ $S = -\frac{1}{2y^2x^2}$	$\frac{dS}{dR} = R$ 

Summary

The solution(s) found are the following

$$-\frac{1}{2y^2x^2} = \frac{x^2}{2} + c_1 \quad (1)$$




Figure 84: Slope field plot

Verification of solutions

$$-\frac{1}{2y^2x^2} = \frac{x^2}{2} + c_1$$

Verified OK.

3.17.2 Solving as bernoulli ode

In canonical form, the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= \frac{y(y^2x^4 - 1)}{x} \end{aligned}$$

This is a Bernoulli ODE.

$$y' = -\frac{1}{x}y + x^3y^3 \quad (1)$$

The standard Bernoulli ODE has the form

$$y' = f_0(x)y + f_1(x)y^n \quad (2)$$

The first step is to divide the above equation by y^n which gives

$$\frac{y'}{y^n} = f_0(x)y^{1-n} + f_1(x) \quad (3)$$

The next step is use the substitution $w = y^{1-n}$ in equation (3) which generates a new ODE in $w(x)$ which will be linear and can be easily solved using an integrating factor. Backsubstitution then gives the solution $y(x)$ which is what we want.

This method is now applied to the ODE at hand. Comparing the ODE (1) With (2) Shows that

$$\begin{aligned} f_0(x) &= -\frac{1}{x} \\ f_1(x) &= x^3 \\ n &= 3 \end{aligned}$$

Dividing both sides of ODE (1) by $y^n = y^3$ gives

$$y' \frac{1}{y^3} = -\frac{1}{y^2 x} + x^3 \quad (4)$$

Let

$$\begin{aligned} w &= y^{1-n} \\ &= \frac{1}{y^2} \end{aligned} \quad (5)$$

Taking derivative of equation (5) w.r.t x gives

$$w' = -\frac{2}{y^3} y' \quad (6)$$

Substituting equations (5) and (6) into equation (4) gives

$$\begin{aligned} -\frac{w'(x)}{2} &= -\frac{w(x)}{x} + x^3 \\ w' &= \frac{2w}{x} - 2x^3 \end{aligned} \quad (7)$$

The above now is a linear ODE in $w(x)$ which is now solved.

Entering Linear first order ODE solver. In canonical form a linear first order is

$$w'(x) + p(x)w(x) = q(x)$$

Where here

$$\begin{aligned} p(x) &= -\frac{2}{x} \\ q(x) &= -2x^3 \end{aligned}$$

Hence the ode is

$$w'(x) - \frac{2w(x)}{x} = -2x^3$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int -\frac{2}{x} dx} \\ &= \frac{1}{x^2}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu w) &= (\mu)(-2x^3) \\ \frac{d}{dx}\left(\frac{w}{x^2}\right) &= \left(\frac{1}{x^2}\right)(-2x^3) \\ d\left(\frac{w}{x^2}\right) &= (-2x) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}\frac{w}{x^2} &= \int -2x dx \\ \frac{w}{x^2} &= -x^2 + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = \frac{1}{x^2}$ results in

$$w(x) = -x^4 + c_1x^2$$

Replacing w in the above by $\frac{1}{y^2}$ using equation (5) gives the final solution.

$$\frac{1}{y^2} = -x^4 + c_1x^2$$

Solving for y gives

$$\begin{aligned}y(x) &= \frac{1}{\sqrt{-x^2 + c_1} x} \\ y(x) &= -\frac{1}{\sqrt{-x^2 + c_1} x}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{1}{\sqrt{-x^2 + c_1 x}} \quad (1)$$

$$y = -\frac{1}{\sqrt{-x^2 + c_1 x}} \quad (2)$$




Figure 85: Slope field plot

Verification of solutions

$$y = \frac{1}{\sqrt{-x^2 + c_1 x}}$$

Verified OK.

$$y = -\frac{1}{\sqrt{-x^2 + c_1 x}}$$

Verified OK.

3.17.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial\phi}{\partial x} + \frac{\partial\phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} (x) dy &= (y^3 x^4 - y) dx \\ (-y^3 x^4 + y) dx + (x) dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -y^3 x^4 + y \\ N(x, y) &= x \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-y^3x^4 + y) \\ &= -3y^2x^4 + 1\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(x) \\ &= 1\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned}A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= \frac{1}{x} ((-3y^2x^4 + 1) - (1)) \\ &= -3y^2x^3\end{aligned}$$

Since A depends on y , it can not be used to obtain an integrating factor. We will now try a second method to find an integrating factor. Let

$$\begin{aligned}B &= \frac{1}{M} \left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) \\ &= \frac{1}{-y^3x^4 + y} ((1) - (-3y^2x^4 + 1)) \\ &= -\frac{3yx^4}{y^2x^4 - 1}\end{aligned}$$

Since B depends on x , it can not be used to obtain an integrating factor. We will now try a third method to find an integrating factor. Let

$$R = \frac{\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y}}{xM - yN}$$

R is now checked to see if it is a function of only $t = xy$. Therefore

$$\begin{aligned} R &= \frac{\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y}}{xM - yN} \\ &= \frac{(1) - (-3y^2x^4 + 1)}{x(-y^3x^4 + y) - y(x)} \\ &= -\frac{3}{yx} \end{aligned}$$

Replacing all powers of terms xy by t gives

$$R = -\frac{3}{t}$$

Since R depends on t only, then it can be used to find an integrating factor. Let the integrating factor be μ then

$$\begin{aligned} \mu &= e^{\int R dt} \\ &= e^{\int (-\frac{3}{t}) dt} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^{-3\ln(t)} \\ &= \frac{1}{t^3} \end{aligned}$$

Now t is replaced back with xy giving

$$\mu = \frac{1}{x^3y^3}$$

Multiplying M and N by this integrating factor gives new M and new N which are called \bar{M} and \bar{N} so not to confuse them with the original M and N

$$\begin{aligned} \bar{M} &= \mu M \\ &= \frac{1}{x^3y^3}(-y^3x^4 + y) \\ &= \frac{-y^2x^4 + 1}{y^2x^3} \end{aligned}$$

And

$$\begin{aligned} \bar{N} &= \mu N \\ &= \frac{1}{x^3y^3}(x) \\ &= \frac{1}{y^3x^2} \end{aligned}$$

A modified ODE is now obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned} \overline{M} + \overline{N} \frac{dy}{dx} &= 0 \\ \left(\frac{-y^2x^4 + 1}{y^2x^3} \right) + \left(\frac{1}{y^3x^2} \right) \frac{dy}{dx} &= 0 \end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \overline{M} \quad (1)$$

$$\frac{\partial \phi}{\partial y} = \overline{N} \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned} \int \frac{\partial \phi}{\partial x} dx &= \int \overline{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int \frac{-y^2x^4 + 1}{y^2x^3} dx \\ \phi &= \frac{-y^2x^4 - 1}{2y^2x^2} + f(y) \end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\begin{aligned} \frac{\partial \phi}{\partial y} &= -\frac{x^2}{y} - \frac{-y^2x^4 - 1}{y^3x^2} + f'(y) \\ &= \frac{1}{y^3x^2} + f'(y) \end{aligned} \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{1}{y^3x^2}$. Therefore equation (4) becomes

$$\frac{1}{y^3x^2} = \frac{1}{y^3x^2} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = \frac{-y^2x^4 - 1}{2y^2x^2} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \frac{-y^2x^4 - 1}{2y^2x^2}$$

Summary

The solution(s) found are the following

$$\frac{-y^2x^4 - 1}{2y^2x^2} = c_1 \tag{1}$$




Figure 86: Slope field plot

Verification of solutions

$$\frac{-y^2x^4 - 1}{2y^2x^2} = c_1$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
<- Bernoulli successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 34

```
dsolve(x*diff(y(x),x)+y(x)=x^4*y(x)^3,y(x), singsol=all)
```

$$y(x) = \frac{1}{\sqrt{-x^2 + c_1 x}}$$
$$y(x) = -\frac{1}{\sqrt{-x^2 + c_1 x}}$$

✓ Solution by Mathematica

Time used: 0.414 (sec). Leaf size: 48

```
DSolve[x*y'[x]+y[x]==x^4*y[x]^3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{\sqrt{-x^4 + c_1 x^2}}$$
$$y(x) \rightarrow \frac{1}{\sqrt{-x^4 + c_1 x^2}}$$
$$y(x) \rightarrow 0$$

3.18 problem 3(b)

3.18.1 Solving as first order ode lie symmetry lookup ode	402
3.18.2 Solving as bernoulli ode	406
3.18.3 Solving as exact ode	410

Internal problem ID [6174]

Internal file name [OUTPUT/5422_Sunday_June_05_2022_03_36_51_PM_7313356/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.4 First Order Linear Equations. Page 15

Problem number: 3(b).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "bernoulli", "exactWithIntegrationFactor", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_Bernoulli]

$$xy^2y' + y^3 = \cos(x)x$$

3.18.1 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{-y^3 + \cos(x)x}{xy^2}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2\xi_y - \omega_x\xi - \omega_y\eta = 0 \tag{A}$$

The type of this ode is known. It is of type Bernoulli. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 73: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= \frac{1}{y^2x^3}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{1}{y^2 x^3}} dy \end{aligned}$$

Which results in

$$S = \frac{y^3 x^3}{3}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{-y^3 + \cos(x) x}{x y^2}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= y^3 x^2 \\ S_y &= y^2 x^3 \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \cos(x) x^3 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \cos(R) R^3$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \sin(R) R^3 + 3 \cos(R) R^2 - 6 \cos(R) - 6 \sin(R) R + c_1 \quad (4)$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{x^3 y^3}{3} = \sin(x) x^3 + 3 \cos(x) x^2 - 6 \cos(x) - 6 \sin(x) x + c_1$$

Which simplifies to

$$\frac{x^3 y^3}{3} = \sin(x) x^3 + 3 \cos(x) x^2 - 6 \cos(x) - 6 \sin(x) x + c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{-y^3 + \cos(x)x}{x y^2}$ 	$R = x$ $S = \frac{y^3 x^3}{3}$	$\frac{dS}{dR} = \cos(R) R^3$ 

Summary

The solution(s) found are the following

$$\frac{x^3 y^3}{3} = \sin(x) x^3 + 3 \cos(x) x^2 - 6 \cos(x) - 6 \sin(x) x + c_1 \quad (1)$$




Figure 87: Slope field plot

Verification of solutions

$$\frac{x^3 y^3}{3} = \sin(x) x^3 + 3 \cos(x) x^2 - 6 \cos(x) - 6 \sin(x) x + c_1$$

Verified OK.

3.18.2 Solving as bernoulli ode

In canonical form, the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= \frac{-y^3 + \cos(x) x}{x y^2} \end{aligned}$$

This is a Bernoulli ODE.

$$y' = -\frac{1}{x}y + \cos(x) \frac{1}{y^2} \quad (1)$$

The standard Bernoulli ODE has the form

$$y' = f_0(x)y + f_1(x)y^n \quad (2)$$

The first step is to divide the above equation by y^n which gives

$$\frac{y'}{y^n} = f_0(x)y^{1-n} + f_1(x) \quad (3)$$

The next step is use the substitution $w = y^{1-n}$ in equation (3) which generates a new ODE in $w(x)$ which will be linear and can be easily solved using an integrating factor. Backsubstitution then gives the solution $y(x)$ which is what we want.

This method is now applied to the ODE at hand. Comparing the ODE (1) With (2) Shows that

$$\begin{aligned}f_0(x) &= -\frac{1}{x} \\f_1(x) &= \cos(x) \\n &= -2\end{aligned}$$

Dividing both sides of ODE (1) by $y^n = \frac{1}{y^2}$ gives

$$y'y^2 = -\frac{y^3}{x} + \cos(x) \quad (4)$$

Let

$$\begin{aligned}w &= y^{1-n} \\&= y^3\end{aligned} \quad (5)$$

Taking derivative of equation (5) w.r.t x gives

$$w' = 3y^2y' \quad (6)$$

Substituting equations (5) and (6) into equation (4) gives

$$\begin{aligned}\frac{w'(x)}{3} &= -\frac{w(x)}{x} + \cos(x) \\w' &= -\frac{3w}{x} + 3\cos(x)\end{aligned} \quad (7)$$

The above now is a linear ODE in $w(x)$ which is now solved.

Entering Linear first order ODE solver. In canonical form a linear first order is

$$w'(x) + p(x)w(x) = q(x)$$

Where here

$$\begin{aligned}p(x) &= \frac{3}{x} \\q(x) &= 3\cos(x)\end{aligned}$$

Hence the ode is

$$w'(x) + \frac{3w(x)}{x} = 3 \cos(x)$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int \frac{3}{x} dx} \\ &= x^3\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu w) &= (\mu) (3 \cos(x)) \\ \frac{d}{dx}(x^3 w) &= (x^3) (3 \cos(x)) \\ d(x^3 w) &= (3 \cos(x) x^3) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}x^3 w &= \int 3 \cos(x) x^3 dx \\ x^3 w &= 3 \sin(x) x^3 + 9 \cos(x) x^2 - 18 \cos(x) - 18 \sin(x) x + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = x^3$ results in

$$w(x) = \frac{3 \sin(x) x^3 + 9 \cos(x) x^2 - 18 \cos(x) - 18 \sin(x) x + c_1}{x^3}$$

which simplifies to

$$w(x) = \frac{9(x^2 - 2) \cos(x) + 3(x^3 - 6x) \sin(x) + c_1}{x^3}$$

Replacing w in the above by y^3 using equation (5) gives the final solution.

$$y^3 = \frac{9(x^2 - 2) \cos(x) + 3(x^3 - 6x) \sin(x) + c_1}{x^3}$$

Solving for y gives

$$\begin{aligned}y(x) &= \frac{(9(x^2 - 2) \cos(x) + 3(x^3 - 6x) \sin(x) + c_1)^{\frac{1}{3}}}{x} \\ y(x) &= \frac{(9(x^2 - 2) \cos(x) + 3(x^3 - 6x) \sin(x) + c_1)^{\frac{1}{3}} (i\sqrt{3} - 1)}{2x} \\ y(x) &= -\frac{(9(x^2 - 2) \cos(x) + 3(x^3 - 6x) \sin(x) + c_1)^{\frac{1}{3}} (1 + i\sqrt{3})}{2x}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{(9(x^2 - 2) \cos(x) + 3(x^3 - 6x) \sin(x) + c_1)^{\frac{1}{3}}}{x} \quad (1)$$

$$y = \frac{(9(x^2 - 2) \cos(x) + 3(x^3 - 6x) \sin(x) + c_1)^{\frac{1}{3}} (i\sqrt{3} - 1)}{2x} \quad (2)$$

$$y = -\frac{(9(x^2 - 2) \cos(x) + 3(x^3 - 6x) \sin(x) + c_1)^{\frac{1}{3}} (1 + i\sqrt{3})}{2x} \quad (3)$$




Figure 88: Slope field plot

Verification of solutions

$$y = \frac{(9(x^2 - 2) \cos(x) + 3(x^3 - 6x) \sin(x) + c_1)^{\frac{1}{3}}}{x}$$

Verified OK.

$$y = \frac{(9(x^2 - 2) \cos(x) + 3(x^3 - 6x) \sin(x) + c_1)^{\frac{1}{3}} (i\sqrt{3} - 1)}{2x}$$

Verified OK.

$$y = -\frac{(9(x^2 - 2) \cos(x) + 3(x^3 - 6x) \sin(x) + c_1)^{\frac{1}{3}} (1 + i\sqrt{3})}{2x}$$

Verified OK.

3.18.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition

$\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned} (y^2 x) dy &= (-y^3 + \cos(x) x) dx \\ (y^3 - \cos(x) x) dx + (y^2 x) dy &= 0 \end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= y^3 - \cos(x) x \\ N(x, y) &= y^2 x \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} (y^3 - \cos(x) x) \\ &= 3y^2 \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} (y^2 x) \\ &= y^2 \end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= \frac{1}{y^2 x} ((3y^2) - (y^2)) \\ &= \frac{2}{x} \end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int A dx} \\ &= e^{\int \frac{2}{x} dx}\end{aligned}$$

The result of integrating gives

$$\begin{aligned}\mu &= e^{2\ln(x)} \\ &= x^2\end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \overline{M} and \overline{N} for now so not to confuse them with the original M and N .

$$\begin{aligned}\overline{M} &= \mu M \\ &= x^2(y^3 - \cos(x)x) \\ &= (y^3 - \cos(x)x)x^2\end{aligned}$$

And

$$\begin{aligned}\overline{N} &= \mu N \\ &= x^2(y^2x) \\ &= y^2x^3\end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned}\overline{M} + \overline{N} \frac{dy}{dx} &= 0 \\ ((y^3 - \cos(x)x)x^2) + (y^2x^3) \frac{dy}{dx} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \overline{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \overline{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \overline{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int (y^3 - \cos(x)x)x^2 dx \\ \phi &= \frac{(-3x^3 + 18x)\sin(x)}{3} + \frac{y^3x^3}{3} - 3\cos(x)x^2 + 6\cos(x) + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = y^2 x^3 + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = y^2 x^3$. Therefore equation (4) becomes

$$y^2 x^3 = y^2 x^3 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = \frac{(-3x^3 + 18x) \sin(x)}{3} + \frac{y^3 x^3}{3} - 3 \cos(x) x^2 + 6 \cos(x) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \frac{(-3x^3 + 18x) \sin(x)}{3} + \frac{y^3 x^3}{3} - 3 \cos(x) x^2 + 6 \cos(x)$$

Summary

The solution(s) found are the following

$$\frac{(-3x^3 + 18x) \sin(x)}{3} + \frac{x^3 y^3}{3} - 3 \cos(x) x^2 + 6 \cos(x) = c_1 \quad (1)$$




Figure 89: Slope field plot

Verification of solutions

$$\frac{(-3x^3 + 18x) \sin(x)}{3} + \frac{x^3 y^3}{3} - 3 \cos(x) x^2 + 6 \cos(x) = c_1$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
<- Bernoulli successful`
```

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 112

```
dsolve(x*y(x)^2*diff(y(x),x)+y(x)^3=x*cos(x),y(x), singsol=all)
```

$$y(x) = \frac{(9(x^2 - 2) \cos(x) + 3(x^3 - 6x) \sin(x) + c_1)^{\frac{1}{3}}}{x}$$
$$y(x) = -\frac{(9(x^2 - 2) \cos(x) + 3(x^3 - 6x) \sin(x) + c_1)^{\frac{1}{3}}(1 + i\sqrt{3})}{2x}$$
$$y(x) = \frac{(9(x^2 - 2) \cos(x) + 3(x^3 - 6x) \sin(x) + c_1)^{\frac{1}{3}}(i\sqrt{3} - 1)}{2x}$$

✓ Solution by Mathematica

Time used: 0.513 (sec). Leaf size: 114

```
DSolve[x*y[x]^2*y'[x]+y[x]^3==x*Cos[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\sqrt[3]{3x(x^2 - 6) \sin(x) + 9(x^2 - 2) \cos(x) + c_1}}{x}$$
$$y(x) \rightarrow -\frac{\sqrt[3]{-1} \sqrt[3]{3x(x^2 - 6) \sin(x) + 9(x^2 - 2) \cos(x) + c_1}}{x}$$
$$y(x) \rightarrow \frac{(-1)^{2/3} \sqrt[3]{3x(x^2 - 6) \sin(x) + 9(x^2 - 2) \cos(x) + c_1}}{x}$$

3.19 problem 3(c)

3.19.1 Solving as first order ode lie symmetry lookup ode	416
3.19.2 Solving as bernoulli ode	420
3.19.3 Solving as exact ode	424
3.19.4 Solving as riccati ode	429

Internal problem ID [6175]

Internal file name [OUTPUT/5423_Sunday_June_05_2022_03_36_55_PM_77365129/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.4 First Order Linear Equations. Page 15

Problem number: 3(c).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**riccati**", "**bernoulli**", "**exactWithIntegrationFactor**", "**first_order_ode_lie_symmetry_lookup**"

Maple gives the following as the ode type

```
[[_homogeneous, `class G`], _rational, _Bernoulli]
```

$$xy' + y - xy^2 = 0$$

3.19.1 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{y(xy - 1)}{x}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \tag{A}$$

The type of this ode is known. It is of type **Bernoulli**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 75: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= y^2x\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{y^2 x} dy \end{aligned}$$

Which results in

$$S = -\frac{1}{yx}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{y(xy - 1)}{x}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{1}{y x^2} \\ S_y &= \frac{1}{y^2 x} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \frac{1}{x} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \frac{1}{R}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \ln(R) + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$-\frac{1}{xy} = \ln(x) + c_1$$



Which simplifies to

$$-\frac{1}{xy} = \ln(x) + c_1$$

Which gives

$$y = -\frac{1}{x(\ln(x) + c_1)}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{y(xy-1)}{x}$ 	$R = x$ $S = -\frac{1}{yx}$	$\frac{dS}{dR} = \frac{1}{R}$ 

Summary

The solution(s) found are the following

$$y = -\frac{1}{x(\ln(x) + c_1)} \quad (1)$$




Figure 90: Slope field plot

Verification of solutions

$$y = -\frac{1}{x(\ln(x) + c_1)}$$

Verified OK.

3.19.2 Solving as bernoulli ode

In canonical form, the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= \frac{y(xy - 1)}{x} \end{aligned}$$

This is a Bernoulli ODE.

$$y' = -\frac{1}{x}y + y^2 \quad (1)$$

The standard Bernoulli ODE has the form

$$y' = f_0(x)y + f_1(x)y^n \quad (2)$$

The first step is to divide the above equation by y^n which gives

$$\frac{y'}{y^n} = f_0(x)y^{1-n} + f_1(x) \quad (3)$$

The next step is use the substitution $w = y^{1-n}$ in equation (3) which generates a new ODE in $w(x)$ which will be linear and can be easily solved using an integrating factor. Backsubstitution then gives the solution $y(x)$ which is what we want.

This method is now applied to the ODE at hand. Comparing the ODE (1) With (2) Shows that

$$\begin{aligned} f_0(x) &= -\frac{1}{x} \\ f_1(x) &= 1 \\ n &= 2 \end{aligned}$$

Dividing both sides of ODE (1) by $y^n = y^2$ gives

$$y' \frac{1}{y^2} = -\frac{1}{yx} + 1 \quad (4)$$

Let

$$\begin{aligned} w &= y^{1-n} \\ &= \frac{1}{y} \end{aligned} \quad (5)$$

Taking derivative of equation (5) w.r.t x gives

$$w' = -\frac{1}{y^2}y' \quad (6)$$

Substituting equations (5) and (6) into equation (4) gives

$$\begin{aligned} -w'(x) &= -\frac{w(x)}{x} + 1 \\ w' &= \frac{w}{x} - 1 \end{aligned} \quad (7)$$

The above now is a linear ODE in $w(x)$ which is now solved.

Entering Linear first order ODE solver. In canonical form a linear first order is

$$w'(x) + p(x)w(x) = q(x)$$

Where here

$$p(x) = -\frac{1}{x}$$
$$q(x) = -1$$

Hence the ode is

$$w'(x) - \frac{w(x)}{x} = -1$$

The integrating factor μ is

$$\mu = e^{\int -\frac{1}{x} dx}$$
$$= \frac{1}{x}$$

The ode becomes

$$\frac{d}{dx}(\mu w) = (\mu)(-1)$$
$$\frac{d}{dx}\left(\frac{w}{x}\right) = \left(\frac{1}{x}\right)(-1)$$
$$d\left(\frac{w}{x}\right) = \left(-\frac{1}{x}\right) dx$$

Integrating gives

$$\frac{w}{x} = \int -\frac{1}{x} dx$$
$$\frac{w}{x} = -\ln(x) + c_1$$

Dividing both sides by the integrating factor $\mu = \frac{1}{x}$ results in

$$w(x) = -\ln(x)x + c_1x$$

which simplifies to

$$w(x) = x(-\ln(x) + c_1)$$

Replacing w in the above by $\frac{1}{y}$ using equation (5) gives the final solution.

$$\frac{1}{y} = x(-\ln(x) + c_1)$$

Or

$$y = \frac{1}{x(-\ln(x) + c_1)}$$

Summary

The solution(s) found are the following

$$y = \frac{1}{x(-\ln(x) + c_1)} \tag{1}$$




Figure 91: Slope field plot

Verification of solutions

$$y = \frac{1}{x(-\ln(x) + c_1)}$$

Verified OK.

3.19.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} (x) dy &= (y^2 x - y) dx \\ (-y^2 x + y) dx + (x) dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -y^2 x + y \\ N(x, y) &= x \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-y^2x + y) \\ &= -2xy + 1\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(x) \\ &= 1\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned}A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= \frac{1}{x} ((-2xy + 1) - (1)) \\ &= -2y\end{aligned}$$

Since A depends on y , it can not be used to obtain an integrating factor. We will now try a second method to find an integrating factor. Let

$$\begin{aligned}B &= \frac{1}{M} \left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) \\ &= \frac{1}{-y^2x + y} ((1) - (-2xy + 1)) \\ &= -\frac{2x}{xy - 1}\end{aligned}$$

Since B depends on x , it can not be used to obtain an integrating factor. We will now try a third method to find an integrating factor. Let

$$R = \frac{\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y}}{xM - yN}$$

R is now checked to see if it is a function of only $t = xy$. Therefore

$$\begin{aligned} R &= \frac{\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y}}{xM - yN} \\ &= \frac{(1) - (-2xy + 1)}{x(-y^2x + y) - y(x)} \\ &= -\frac{2}{yx} \end{aligned}$$

Replacing all powers of terms xy by t gives

$$R = -\frac{2}{t}$$

Since R depends on t only, then it can be used to find an integrating factor. Let the integrating factor be μ then

$$\begin{aligned} \mu &= e^{\int R dt} \\ &= e^{\int (-\frac{2}{t}) dt} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^{-2\ln(t)} \\ &= \frac{1}{t^2} \end{aligned}$$

Now t is replaced back with xy giving

$$\mu = \frac{1}{y^2x^2}$$

Multiplying M and N by this integrating factor gives new M and new N which are called \bar{M} and \bar{N} so not to confuse them with the original M and N

$$\begin{aligned} \bar{M} &= \mu M \\ &= \frac{1}{y^2x^2}(-y^2x + y) \\ &= \frac{-xy + 1}{yx^2} \end{aligned}$$

And

$$\begin{aligned} \bar{N} &= \mu N \\ &= \frac{1}{y^2x^2}(x) \\ &= \frac{1}{y^2x} \end{aligned}$$

A modified ODE is now obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned} \overline{M} + \overline{N} \frac{dy}{dx} &= 0 \\ \left(\frac{-xy + 1}{y x^2} \right) + \left(\frac{1}{y^2 x} \right) \frac{dy}{dx} &= 0 \end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \overline{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \overline{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned} \int \frac{\partial \phi}{\partial x} dx &= \int \overline{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int \frac{-xy + 1}{y x^2} dx \\ \phi &= -\ln(x) - \frac{1}{yx} + f(y) \end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = \frac{1}{y^2 x} + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{1}{y^2 x}$. Therefore equation (4) becomes

$$\frac{1}{y^2 x} = \frac{1}{y^2 x} + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = -\ln(x) - \frac{1}{yx} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\ln(x) - \frac{1}{yx}$$

The solution becomes

$$y = -\frac{1}{x(\ln(x) + c_1)}$$

Summary

The solution(s) found are the following

$$y = -\frac{1}{x(\ln(x) + c_1)} \tag{1}$$




Figure 92: Slope field plot

Verification of solutions

$$y = -\frac{1}{x(\ln(x) + c_1)}$$

Verified OK.

3.19.4 Solving as riccati ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= \frac{y(xy - 1)}{x}\end{aligned}$$

This is a Riccati ODE. Comparing the ODE to solve

$$y' = y^2 - \frac{y}{x}$$

With Riccati ODE standard form

$$y' = f_0(x) + f_1(x)y + f_2(x)y^2$$

Shows that $f_0(x) = 0$, $f_1(x) = -\frac{1}{x}$ and $f_2(x) = 1$. Let

$$\begin{aligned}y &= \frac{-u'}{f_2 u} \\ &= \frac{-u'}{u}\end{aligned}\tag{1}$$

Using the above substitution in the given ODE results (after some simplification) in a second order ODE to solve for $u(x)$ which is

$$f_2 u''(x) - (f_2' + f_1 f_2) u'(x) + f_2^2 f_0 u(x) = 0\tag{2}$$

But

$$\begin{aligned}f_2' &= 0 \\ f_1 f_2 &= -\frac{1}{x} \\ f_2^2 f_0 &= 0\end{aligned}$$

Substituting the above terms back in equation (2) gives

$$u''(x) + \frac{u'(x)}{x} = 0$$

Solving the above ODE (this ode solved using Maple, not this program), gives

$$u(x) = c_1 + c_2 \ln(x)$$

The above shows that

$$u'(x) = \frac{c_2}{x}$$

Using the above in (1) gives the solution

$$y = -\frac{c_2}{x(c_1 + c_2 \ln(x))}$$

Dividing both numerator and denominator by c_1 gives, after renaming the constant $\frac{c_2}{c_1} = c_3$ the following solution

$$y = -\frac{1}{x(c_3 + \ln(x))}$$

Summary

The solution(s) found are the following

$$y = -\frac{1}{x(c_3 + \ln(x))} \tag{1}$$




Figure 93: Slope field plot

Verification of solutions

$$y = -\frac{1}{x(c_3 + \ln(x))}$$

Verified OK.

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
<- Bernoulli successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 16

```
dsolve(x*diff(y(x),x)+y(x)=x*y(x)^2,y(x), singsol=all)
```

$$y(x) = \frac{1}{(-\ln(x) + c_1)x}$$

✓ Solution by Mathematica

Time used: 0.128 (sec). Leaf size: 22

```
DSolve[x*y'[x]+y[x]==x*y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{-x \log(x) + c_1 x}$$
$$y(x) \rightarrow 0$$

3.20 problem 3(d)

3.20.1 Solving as separable ode	433
3.20.2 Solving as first order ode lie symmetry lookup ode	435
3.20.3 Solving as bernoulli ode	439
3.20.4 Solving as exact ode	443
3.20.5 Maple step by step solution	447

Internal problem ID [6176]

Internal file name [OUTPUT/5424_Sunday_June_05_2022_03_36_57_PM_49217883/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.4 First Order Linear Equations. Page 15

Problem number: 3(d).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "bernoulli", "separable", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_separable]

$$y' + xy - xy^4 = 0$$

3.20.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= x(y^4 - y)\end{aligned}$$

Where $f(x) = x$ and $g(y) = y^4 - y$. Integrating both sides gives

$$\frac{1}{y^4 - y} dy = x dx$$

$$\int \frac{1}{y^4 - y} dy = \int x dx$$

$$\frac{\ln(y-1)}{3} - \ln(y) + \frac{\ln(y^2 + y + 1)}{3} = \frac{x^2}{2} + c_1$$

Raising both side to exponential gives

$$e^{\frac{\ln(y-1)}{3} - \ln(y) + \frac{\ln(y^2 + y + 1)}{3}} = e^{\frac{x^2}{2} + c_1}$$

Which simplifies to

$$\frac{(y-1)^{\frac{1}{3}} (y^2 + y + 1)^{\frac{1}{3}}}{y} = c_2 e^{\frac{x^2}{2}}$$

The solution is

$$\frac{(y-1)^{\frac{1}{3}} (y^2 + y + 1)^{\frac{1}{3}}}{y} = c_2 e^{\frac{x^2}{2}}$$

Summary

The solution(s) found are the following

$$\frac{(y-1)^{\frac{1}{3}} (y^2 + y + 1)^{\frac{1}{3}}}{y} = c_2 e^{\frac{x^2}{2}} \quad (1)$$




Figure 94: Slope field plot

Verification of solutions

$$\frac{(y-1)^{\frac{1}{3}}(y^2+y+1)^{\frac{1}{3}}}{y} = c_2 e^{\frac{x^2}{2}}$$

Verified OK.

3.20.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$\begin{aligned}y' &= y^4 x - xy \\y' &= \omega(x, y)\end{aligned}$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **separable**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 77: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= \frac{1}{x} \\ \eta(x, y) &= 0\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\eta = 0$ then in this special case

$$R = y$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\xi} dx \\ &= \int \frac{1}{\frac{1}{x}} dx \end{aligned}$$

Which results in

$$S = \frac{x^2}{2}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = y^4 x - xy$$

Evaluating all the partial derivatives gives

$$R_x = 0$$

$$R_y = 1$$

$$S_x = x$$

$$S_y = 0$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \frac{1}{y^4 - y} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \frac{1}{R^4 - R}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by

integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \frac{\ln(R-1)}{3} - \ln(R) + \frac{\ln(R^2 + R + 1)}{3} + c_1 \quad (4)$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{x^2}{2} = \frac{\ln(y-1)}{3} - \ln(y) + \frac{\ln(y^2 + y + 1)}{3} + c_1$$

Which simplifies to

$$\frac{x^2}{2} = \frac{\ln(y-1)}{3} - \ln(y) + \frac{\ln(y^2 + y + 1)}{3} + c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = y^4 x - xy$ 	$R = y$ $S = \frac{x^2}{2}$	$\frac{dS}{dR} = \frac{1}{R^4 - R}$ 

Summary

The solution(s) found are the following

$$\frac{x^2}{2} = \frac{\ln(y-1)}{3} - \ln(y) + \frac{\ln(y^2 + y + 1)}{3} + c_1 \quad (1)$$




Figure 95: Slope field plot

Verification of solutions

$$\frac{x^2}{2} = \frac{\ln(y-1)}{3} - \ln(y) + \frac{\ln(y^2+y+1)}{3} + c_1$$

Verified OK.

3.20.3 Solving as bernoulli ode

In canonical form, the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= y^4x - xy \end{aligned}$$

This is a Bernoulli ODE.

$$y' = -xy + xy^4 \tag{1}$$

The standard Bernoulli ODE has the form

$$y' = f_0(x)y + f_1(x)y^n \tag{2}$$

The first step is to divide the above equation by y^n which gives

$$\frac{y'}{y^n} = f_0(x)y^{1-n} + f_1(x) \tag{3}$$

The next step is use the substitution $w = y^{1-n}$ in equation (3) which generates a new ODE in $w(x)$ which will be linear and can be easily solved using an integrating factor. Backsubstitution then gives the solution $y(x)$ which is what we want.

This method is now applied to the ODE at hand. Comparing the ODE (1) With (2) Shows that

$$\begin{aligned}f_0(x) &= -x \\f_1(x) &= x \\n &= 4\end{aligned}$$

Dividing both sides of ODE (1) by $y^n = y^4$ gives

$$y' \frac{1}{y^4} = -\frac{x}{y^3} + x \quad (4)$$

Let

$$\begin{aligned}w &= y^{1-n} \\&= \frac{1}{y^3}\end{aligned} \quad (5)$$

Taking derivative of equation (5) w.r.t x gives

$$w' = -\frac{3}{y^4}y' \quad (6)$$

Substituting equations (5) and (6) into equation (4) gives

$$\begin{aligned}-\frac{w'(x)}{3} &= -w(x)x + x \\w' &= 3xw - 3x\end{aligned} \quad (7)$$

The above now is a linear ODE in $w(x)$ which is now solved.

Entering Linear first order ODE solver. In canonical form a linear first order is

$$w'(x) + p(x)w(x) = q(x)$$

Where here

$$\begin{aligned}p(x) &= -3x \\q(x) &= -3x\end{aligned}$$

Hence the ode is

$$w'(x) - 3w(x)x = -3x$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int -3x dx} \\ &= e^{-\frac{3x^2}{2}}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu w) &= (\mu)(-3x) \\ \frac{d}{dx}\left(e^{-\frac{3x^2}{2}} w\right) &= \left(e^{-\frac{3x^2}{2}}\right)(-3x) \\ d\left(e^{-\frac{3x^2}{2}} w\right) &= \left(-3x e^{-\frac{3x^2}{2}}\right) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}e^{-\frac{3x^2}{2}} w &= \int -3x e^{-\frac{3x^2}{2}} dx \\ e^{-\frac{3x^2}{2}} w &= e^{-\frac{3x^2}{2}} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{-\frac{3x^2}{2}}$ results in

$$w(x) = e^{\frac{3x^2}{2}} e^{-\frac{3x^2}{2}} + c_1 e^{\frac{3x^2}{2}}$$

which simplifies to

$$w(x) = 1 + c_1 e^{\frac{3x^2}{2}}$$

Replacing w in the above by $\frac{1}{y^3}$ using equation (5) gives the final solution.

$$\frac{1}{y^3} = 1 + c_1 e^{\frac{3x^2}{2}}$$

Solving for y gives

$$\begin{aligned}y(x) &= \frac{1}{\left(1 + c_1 e^{\frac{3x^2}{2}}\right)^{\frac{1}{3}}} \\ y(x) &= \frac{i\sqrt{3} - 1}{2\left(1 + c_1 e^{\frac{3x^2}{2}}\right)^{\frac{1}{3}}} \\ y(x) &= -\frac{1 + i\sqrt{3}}{2\left(1 + c_1 e^{\frac{3x^2}{2}}\right)^{\frac{1}{3}}}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{1}{\left(1 + c_1 e^{\frac{3x^2}{2}}\right)^{\frac{1}{3}}} \quad (1)$$

$$y = \frac{i\sqrt{3} - 1}{2\left(1 + c_1 e^{\frac{3x^2}{2}}\right)^{\frac{1}{3}}} \quad (2)$$

$$y = -\frac{1 + i\sqrt{3}}{2\left(1 + c_1 e^{\frac{3x^2}{2}}\right)^{\frac{1}{3}}} \quad (3)$$




Figure 96: Slope field plot

Verification of solutions

$$y = \frac{1}{\left(1 + c_1 e^{\frac{3x^2}{2}}\right)^{\frac{1}{3}}}$$

Verified OK.

$$y = \frac{i\sqrt{3} - 1}{2 \left(1 + c_1 e^{\frac{3x^2}{2}}\right)^{\frac{1}{3}}}$$

Verified OK.

$$y = -\frac{1 + i\sqrt{3}}{2 \left(1 + c_1 e^{\frac{3x^2}{2}}\right)^{\frac{1}{3}}}$$

Verified OK.

3.20.4 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned} & \left(\frac{1}{y^4 - y} \right) dy = (x) dx \\ (-x) dx + \left(\frac{1}{y^4 - y} \right) dy &= 0 \end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -x \\ N(x, y) &= \frac{1}{y^4 - y} \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-x) \\ &= 0 \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{1}{y^4 - y} \right) \\ &= 0 \end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -x dx \\ \phi &= -\frac{x^2}{2} + f(y)\end{aligned}\quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y)\quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{1}{y^4 - y}$. Therefore equation (4) becomes

$$\frac{1}{y^4 - y} = 0 + f'(y)\quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$\begin{aligned}f'(y) &= \frac{1}{y(y^3 - 1)} \\ &= \frac{1}{y^4 - y}\end{aligned}$$

Integrating the above w.r.t y results in

$$\begin{aligned}\int f'(y) dy &= \int \left(\frac{1}{y^4 - y} \right) dy \\ f(y) &= \frac{\ln(y - 1)}{3} - \ln(y) + \frac{\ln(y^2 + y + 1)}{3} + c_1\end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\frac{x^2}{2} + \frac{\ln(y - 1)}{3} - \ln(y) + \frac{\ln(y^2 + y + 1)}{3} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\frac{x^2}{2} + \frac{\ln(y-1)}{3} - \ln(y) + \frac{\ln(y^2 + y + 1)}{3}$$

Summary

The solution(s) found are the following

$$-\frac{x^2}{2} + \frac{\ln(y-1)}{3} - \ln(y) + \frac{\ln(y^2 + y + 1)}{3} = c_1 \quad (1)$$




Figure 97: Slope field plot

Verification of solutions

$$-\frac{x^2}{2} + \frac{\ln(y-1)}{3} - \ln(y) + \frac{\ln(y^2 + y + 1)}{3} = c_1$$

Verified OK.

3.20.5 Maple step by step solution

Let's solve

$$y' + xy - xy^4 = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y(y-1)(y^2+y+1)} = x$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y(y-1)(y^2+y+1)} dx = \int x dx + c_1$$

- Evaluate integral

$$\frac{\ln(y-1)}{3} - \ln(y) + \frac{\ln(y^2+y+1)}{3} = \frac{x^2}{2} + c_1$$

- Solve for y

$$y = \frac{\left(-\left(e^{\frac{3x^2}{2}+3c_1-1}\right)^2\right)^{\frac{1}{3}}}{e^{\frac{3x^2}{2}+3c_1-1}}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
<- Bernoulli successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 66

```
dsolve(diff(y(x),x)+x*y(x)=x*y(x)^4,y(x), singsol=all)
```

$$y(x) = \frac{1}{\left(e^{\frac{3x^2}{2}} c_1 + 1\right)^{\frac{1}{3}}}$$

$$y(x) = -\frac{1 + i\sqrt{3}}{2 \left(e^{\frac{3x^2}{2}} c_1 + 1\right)^{\frac{1}{3}}}$$

$$y(x) = \frac{i\sqrt{3} - 1}{2 \left(e^{\frac{3x^2}{2}} c_1 + 1\right)^{\frac{1}{3}}}$$

✓ Solution by Mathematica

Time used: 1.97 (sec). Leaf size: 116

```
DSolve[y'[x]+x*y[x]==x*y[x]^4,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{\sqrt[3]{1 + e^{\frac{3x^2}{2}} + 3c_1}}$$

$$y(x) \rightarrow -\frac{\sqrt[3]{-1}}{\sqrt[3]{1 + e^{\frac{3x^2}{2}} + 3c_1}}$$

$$y(x) \rightarrow \frac{(-1)^{2/3}}{\sqrt[3]{1 + e^{\frac{3x^2}{2}} + 3c_1}}$$

$$y(x) \rightarrow 0$$

$$y(x) \rightarrow 1$$

$$y(x) \rightarrow -\sqrt[3]{-1}$$

$$y(x) \rightarrow (-1)^{2/3}$$

3.21 problem 4(a)

3.21.1 Solving as exact ode	449
3.21.2 Maple step by step solution	453

Internal problem ID [6177]

Internal file name [OUTPUT/5425_Sunday_June_05_2022_03_36_59_PM_41933859/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.4 First Order Linear Equations. Page 15

Problem number: 4(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact"

Maple gives the following as the ode type

```
[_exact , [_1st_order , ` _with_symmetry_ [F(x)*G(y) , 0] `]]
```

$(e^y - 2xy)y' - y^2 = 0$

3.21.1 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned}(e^y - 2xy) dy &= (y^2) dx \\ (-y^2) dx + (e^y - 2xy) dy &= 0\end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -y^2 \\ N(x, y) &= e^y - 2xy\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-y^2) \\ &= -2y\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(e^y - 2xy) \\ &= -2y\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned} \int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -y^2 dx \\ \phi &= -y^2x + f(y) \end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = -2xy + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = e^y - 2xy$. Therefore equation (4) becomes

$$e^y - 2xy = -2xy + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = e^y$$

Integrating the above w.r.t y gives

$$\begin{aligned} \int f'(y) dy &= \int (e^y) dy \\ f(y) &= e^y + c_1 \end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -y^2x + e^y + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -y^2x + e^y$$

Summary

The solution(s) found are the following

$$-xy^2 + e^y = c_1 \tag{1}$$




Figure 98: Slope field plot

Verification of solutions

$$-xy^2 + e^y = c_1$$

Verified OK.

3.21.2 Maple step by step solution

Let's solve

$$(e^y - 2xy) y' - y^2 = 0$$

- Highest derivative means the order of the ODE is 1
 y'
- Check if ODE is exact
 - ODE is exact if the lhs is the total derivative of a C^2 function
 $F'(x, y) = 0$
 - Compute derivative of lhs
 $F'(x, y) + \left(\frac{\partial}{\partial y} F(x, y)\right) y' = 0$
 - Evaluate derivatives
 $-2y = -2y$
 - Condition met, ODE is exact
- Exact ODE implies solution will be of this form
 $\left[F(x, y) = c_1, M(x, y) = F'(x, y), N(x, y) = \frac{\partial}{\partial y} F(x, y) \right]$
- Solve for $F(x, y)$ by integrating $M(x, y)$ with respect to x
 $F(x, y) = \int -y^2 dx + f_1(y)$
- Evaluate integral
 $F(x, y) = -y^2 x + f_1(y)$
- Take derivative of $F(x, y)$ with respect to y
 $N(x, y) = \frac{\partial}{\partial y} F(x, y)$
- Compute derivative
 $e^y - 2xy = -2xy + \frac{d}{dy} f_1(y)$
- Isolate for $\frac{d}{dy} f_1(y)$
 $\frac{d}{dy} f_1(y) = e^y$
- Solve for $f_1(y)$
 $f_1(y) = e^y$
- Substitute $f_1(y)$ into equation for $F(x, y)$

$$F(x, y) = -y^2x + e^y$$

- Substitute $F(x, y)$ into the solution of the ODE

$$-y^2x + e^y = c_1$$

- Solve for y

$$y = \text{RootOf}(-Z^2x - e^{-Z} + c_1)$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
<- 1st order linear successful
<- inverse linear successful`

```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 23

```
dsolve((exp(y(x))-2*x*y(x))*diff(y(x),x)=y(x)^2,y(x), singsol=all)
```

$$\frac{xy(x)^2 - e^{y(x)} - c_1}{y(x)^2} = 0$$

✓ Solution by Mathematica

Time used: 0.236 (sec). Leaf size: 22

```
DSolve[(Exp[y[x]]-2*x*y[x])*y'[x]==y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve}\left[x = \frac{e^{y(x)}}{y(x)^2} + \frac{c_1}{y(x)^2}, y(x)\right]$$

3.22 problem 4(b)

3.22.1 Solving as first order ode lie symmetry calculated ode	455
3.22.2 Solving as exact ode	460

Internal problem ID [6178]

Internal file name [OUTPUT/5426_Sunday_June_05_2022_03_37_00_PM_28995455/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.4 First Order Linear Equations. Page 15

Problem number: 4(b).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exactWithIntegrationFactor", "first_order_ode_lie_symmetry_calculated"**

Maple gives the following as the ode type

```
[[_1st_order , _with_linear_symmetries]]
```

$-xy' + y - y'y^2e^y = 0$

3.22.1 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = \frac{y}{y^2e^y + x}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2\xi_y - \omega_x\xi - \omega_y\eta = 0 \tag{A}$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \tag{1E}$$

$$\eta = xb_2 + yb_3 + b_1 \tag{2E}$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$b_2 + \frac{y(b_3 - a_2)}{y^2 e^y + x} - \frac{y^2 a_3}{(y^2 e^y + x)^2} + \frac{y(xa_2 + ya_3 + a_1)}{(y^2 e^y + x)^2} \quad (5E)$$

$$- \left(\frac{1}{y^2 e^y + x} - \frac{y(2e^y y + y^2 e^y)}{(y^2 e^y + x)^2} \right) (xb_2 + yb_3 + b_1) = 0$$

Putting the above in normal form gives

$$\frac{e^{2y} y^4 b_2 + e^y x y^3 b_2 + e^y y^4 b_3 + 3e^y x y^2 b_2 - e^y y^3 a_2 + e^y y^3 b_1 + 2e^y y^3 b_3 + e^y y^2 b_1 - xb_1 + ya_1}{(y^2 e^y + x)^2}$$

$$= 0$$

Setting the numerator to zero gives

$$e^{2y} y^4 b_2 + e^y x y^3 b_2 + e^y y^4 b_3 + 3e^y x y^2 b_2 - e^y y^3 a_2 \quad (6E)$$

$$+ e^y y^3 b_1 + 2e^y y^3 b_3 + e^y y^2 b_1 - xb_1 + ya_1 = 0$$

Simplifying the above gives

$$e^{2y} y^4 b_2 + e^y x y^3 b_2 + e^y y^4 b_3 + 3e^y x y^2 b_2 - e^y y^3 a_2 \quad (6E)$$

$$+ e^y y^3 b_1 + 2e^y y^3 b_3 + e^y y^2 b_1 - xb_1 + ya_1 = 0$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y, e^y, e^{2y}\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2, e^y = v_3, e^{2y} = v_4\}$$

The above PDE (6E) now becomes

$$v_3 v_1 v_2^3 b_2 + v_4 v_2^4 b_2 + v_3 v_2^4 b_3 - v_3 v_2^3 a_2 + v_3 v_2^3 b_1 \quad (7E)$$

$$+ 3v_3 v_1 v_2^2 b_2 + 2v_3 v_2^3 b_3 + v_3 v_2^2 b_1 + v_2 a_1 - v_1 b_1 = 0$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2, v_3, v_4\}$$

Equation (7E) now becomes

$$v_3 v_1 v_2^3 b_2 + 3v_3 v_1 v_2^2 b_2 - v_1 b_1 + v_3 v_2^4 b_3 + v_4 v_2^4 b_2 + (-a_2 + b_1 + 2b_3) v_2^3 v_3 + v_3 v_2^2 b_1 + v_2 a_1 = 0 \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} a_1 &= 0 \\ b_1 &= 0 \\ b_2 &= 0 \\ b_3 &= 0 \\ -b_1 &= 0 \\ 3b_2 &= 0 \\ -a_2 + b_1 + 2b_3 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= 0 \\ a_2 &= 0 \\ a_3 &= a_3 \\ b_1 &= 0 \\ b_2 &= 0 \\ b_3 &= 0 \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned} \xi &= y \\ \eta &= 0 \end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned} \eta &= \eta - \omega(x, y) \xi \\ &= 0 - \left(\frac{y}{y^2 e^y + x} \right) (y) \\ &= -\frac{y^2}{y^2 e^y + x} \\ \xi &= 0 \end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{-\frac{y^2}{y^2 e^y + x}} dy \end{aligned}$$

Which results in

$$S = \frac{x}{y} - e^y$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{y}{y^2 e^y + x}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{1}{y} \\ S_y &= \frac{-y^2 e^y - x}{y^2} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \tag{2A}$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \tag{4}$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{-e^y y + x}{y} = c_1$$

Which simplifies to

$$\frac{-e^y y + x}{y} = c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{y}{y^2 e^y + x}$ 	$R = x$ $S = \frac{-e^y y + x}{y}$	$\frac{dS}{dR} = 0$ 

Summary

The solution(s) found are the following

$$\frac{-e^y y + x}{y} = c_1 \quad (1)$$




Figure 99: Slope field plot

Verification of solutions

$$\frac{-e^y y + x}{y} = c_1$$

Verified OK.

3.22.2 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (A)$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} (-y^2 e^y - x) dy &= (-y) dx \\ (y) dx + (-y^2 e^y - x) dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= y \\ N(x, y) &= -y^2 e^y - x \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(y) \\ &= 1 \end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(-y^2e^y - x) \\ &= -1\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned}A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= -\frac{1}{y^2e^y + x} ((1) - (-1)) \\ &= -\frac{2}{y^2e^y + x}\end{aligned}$$

Since A depends on y , it can not be used to obtain an integrating factor. We will now try a second method to find an integrating factor. Let

$$\begin{aligned}B &= \frac{1}{M} \left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) \\ &= \frac{1}{y} ((-1) - (1)) \\ &= -\frac{2}{y}\end{aligned}$$

Since B does not depend on x , it can be used to obtain an integrating factor. Let the integrating factor be μ . Then

$$\begin{aligned}\mu &= e^{\int B \, dy} \\ &= e^{\int -\frac{2}{y} \, dy}\end{aligned}$$

The result of integrating gives

$$\begin{aligned}\mu &= e^{-2\ln(y)} \\ &= \frac{1}{y^2}\end{aligned}$$

M and N are now multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} so not to confuse them with the original M and N .

$$\begin{aligned}\bar{M} &= \mu M \\ &= \frac{1}{y^2}(y) \\ &= \frac{1}{y}\end{aligned}$$

And

$$\begin{aligned}\bar{N} &= \mu N \\ &= \frac{1}{y^2}(-y^2 e^y - x) \\ &= \frac{-y^2 e^y - x}{y^2}\end{aligned}$$

So now a modified ODE is obtained from the original ODE which will be exact and can be solved using the standard method. The modified ODE is

$$\begin{aligned}\bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ \left(\frac{1}{y}\right) + \left(\frac{-y^2 e^y - x}{y^2}\right) \frac{dy}{dx} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int \frac{1}{y} dx \\ \phi &= \frac{x}{y} + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = -\frac{x}{y^2} + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{-y^2 e^y - x}{y^2}$. Therefore equation (4) becomes

$$\frac{-y^2 e^y - x}{y^2} = -\frac{x}{y^2} + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = -e^y$$

Integrating the above w.r.t y gives

$$\int f'(y) dy = \int (-e^y) dy$$

$$f(y) = -e^y + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = \frac{x}{y} - e^y + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \frac{x}{y} - e^y$$

Summary

The solution(s) found are the following

$$\frac{x}{y} - e^y = c_1 \tag{1}$$




Figure 100: Slope field plot

Verification of solutions

$$\frac{x}{y} - e^y = c_1$$

Verified OK.

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
<- 1st order linear successful
<- inverse linear successful`

```

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 17

```
dsolve(y(x)-x*diff(y(x),x)=diff(y(x),x)*y(x)^2*exp(y(x)),y(x), singsol=all)
```

$$-y(x) e^{y(x)} - c_1 y(x) + x = 0$$

✓ Solution by Mathematica

Time used: 0.203 (sec). Leaf size: 18

```
DSolve[y[x]-x*y'[x]==y'[x]*y[x]^2*Exp[y[x]],y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve}[x = e^{y(x)}y(x) + c_1y(x), y(x)]$$

3.23 problem 4(c)

3.23.1 Solving as first order ode lie symmetry calculated ode 467

Internal problem ID [6179]

Internal file name [OUTPUT/5427_Sunday_June_05_2022_03_37_03_PM_5587567/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.4 First Order Linear Equations. Page 15

Problem number: 4(c).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**first_order_ode_lie_symmetry_calculated**"

Maple gives the following as the ode type

```
[[_1st_order, `_with_symmetry_[F(x)*G(y),0]`], [_Abel, `_2nd
type`, `_class C`]]
```

$$xy' - x^3(y - 1)y' = -2$$

3.23.1 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = \frac{2}{x(yx^2 - x^2 - 1)}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 3 to use as anstaz gives

$$\xi = x^3 a_7 + y x^2 a_8 + y^2 x a_9 + y^3 a_{10} + x^2 a_4 + x y a_5 + y^2 a_6 + x a_2 + y a_3 + a_1 \quad (1\text{E})$$

$$\eta = x^3 b_7 + y x^2 b_8 + y^2 x b_9 + y^3 b_{10} + x^2 b_4 + x y b_5 + y^2 b_6 + x b_2 + y b_3 + b_1 \quad (2\text{E})$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, a_4, a_5, a_6, a_7, a_8, a_9, a_{10}, b_1, b_2, b_3, b_4, b_5, b_6, b_7, b_8, b_9, b_{10}\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$\begin{aligned} & 3x^2b_7 + 2xyb_8 + y^2b_9 + 2xb_4 + yb_5 + b_2 \tag{5E} \\ & + \frac{-6x^2a_7 + 2x^2b_8 - 4xya_8 + 4xyb_9 - 2y^2a_9 + 6y^2b_{10} - 4xa_4 + 2xb_5 - 2ya_5 + 4yb_6 - 2a_2 + 2b_3}{x(yx^2 - x^2 - 1)} \\ & - \frac{4(x^2a_8 + 2xya_9 + 3y^2a_{10} + xa_5 + 2ya_6 + a_3)}{x^2(yx^2 - x^2 - 1)^2} \\ & - \left(-\frac{2}{x^2(yx^2 - x^2 - 1)} - \frac{2(2xy - 2x)}{x(yx^2 - x^2 - 1)^2} \right) (x^3a_7 + yx^2a_8 \\ & + y^2xa_9 + y^3a_{10} + x^2a_4 + xya_5 + y^2a_6 + xa_2 + ya_3 + a_1) \\ & + \frac{2x(x^3b_7 + yx^2b_8 + y^2xb_9 + y^3b_{10} + x^2b_4 + xyb_5 + y^2b_6 + xb_2 + yb_3 + b_1)}{(yx^2 - x^2 - 1)^2} = 0 \end{aligned}$$

Putting the above in normal form gives

$$\begin{aligned} & \frac{2x^3yb_8 + y^2b_9x^2 - 2a_1 - 4a_3 + yb_5x^2 + 2x^4ya_4 + 4x^3y^2a_5 + 6x^3y^2b_6 - 4x^3ya_5 - 4x^3yb_6 - 4xyb_6 + 6x^2y^3a_6}{1} \\ & = 0 \end{aligned}$$

Setting the numerator to zero gives

$$\begin{aligned} & 2x^3yb_8 + y^2b_9x^2 - 2a_1 - 4a_3 + yb_5x^2 + 2x^4ya_4 + 4x^3y^2a_5 \\ & + 6x^3y^2b_6 - 4x^3ya_5 - 4x^3yb_6 - 4xyb_6 + 6x^2y^3a_6 - 6x^2y^2a_6 + x^6b_2 \\ & + 4x^4b_2 - 4x^3a_2 - 2x^3b_3 - 2xb_3 - 6x^2a_1 + 2x^3b_1 + 2x^3b_4 + 2x^7y^2b_4 \\ & - 4x^7yb_4 - 4x^5yb_4 + x^6y^3b_5 - 2x^6y^2b_5 + x^6yb_5 - 2x^4y^2b_5 + 6x^4yb_5 \\ & - 4x^2a_8 - 12y^2a_{10} + 3x^8b_7 + 8x^6b_7 - 2x^5b_8 - 2x^3b_8 - 4xa_5 - 8ya_6 \\ & + 2x^7b_4 + 6x^5b_4 - 2x^4a_4 - 2x^4b_5 - 2x^2b_5 + b_2x^2 + x^6y^2b_2 - 2x^6yb_2 \\ & - 2x^4yb_2 + 4x^3ya_2 + 4x^3yb_3 + 6x^2y^2a_3 + 6x^2ya_1 - 8xya_9 + 3x^4b_7 \\ & + 3x^8y^2b_7 - 6x^8yb_7 - 6x^6yb_7 + 2yx^2a_8 + 2x^7y^3b_8 - 4x^7y^2b_8 \\ & + 2x^7yb_8 - 4x^5y^2b_8 + 8x^5yb_8 + x^6y^4b_9 - 2x^6y^3b_9 + x^6y^2b_9 \\ & - 2x^4y^3b_9 + 8x^4y^2b_9 + 2x^4y^2a_8 + 4x^3y^3a_9 + 8x^3y^3b_{10} - 2x^4ya_8 \\ & - 4x^4yb_9 - 4x^3y^2a_9 - 6x^3y^2b_{10} - 4x^2yb_9 - 6xy^2b_{10} + 6x^2y^4a_{10} \\ & - 6x^2y^3a_{10} - 6x^2ya_3 - 2ya_3 + 2x^2a_4 - 2y^2a_6 + 4x^3a_7 - 2y^3a_{10} = 0 \end{aligned} \tag{6E}$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2\}$$

The above PDE (6E) now becomes

$$\begin{aligned}
& 4v_1^3v_2b_3 + 6v_1^2v_2^2a_3 + 6v_1^2v_2a_1 - 8v_1v_2a_9 + 3v_1^8v_2^2b_7 - 6v_1^8v_2b_7 \\
& - 6v_1^6v_2b_7 + 2v_2v_1^2a_8 + 2v_1^7v_2^3b_8 - 4v_1^7v_2^2b_8 + 2v_1^7v_2b_8 - 4v_1^5v_2^2b_8 \\
& + 8v_1^5v_2b_8 + v_1^6v_2^4b_9 - 2v_1^6v_2^3b_9 + v_1^6v_2^2b_9 - 2v_1^4v_2^3b_9 + 8v_1^4v_2^2b_9 \\
& + 2v_1^4v_2^2a_8 + 4v_1^3v_2^3a_9 + 8v_1^3v_2^3b_{10} - 2v_1^4v_2a_8 - 4v_1^4v_2b_9 - 4v_1^3v_2^2a_9 \\
& - 6v_1^3v_2^2b_{10} - 4v_1^2v_2b_9 - 6v_1v_2^2b_{10} + 6v_1^2v_2^4a_{10} - 6v_1^2v_2^3a_{10} - 6v_1^2v_2a_3 \\
& - 2a_1 - 4a_3 + 2v_1^3v_2b_8 + v_2^2b_9v_1^2 + v_2b_5v_1^2 + 2v_1^4v_2a_4 + 4v_1^3v_2^2a_5 \\
& + 6v_1^3v_2^2b_6 - 4v_1^3v_2a_5 - 4v_1^3v_2b_6 - 4v_1v_2b_6 + 6v_1^2v_2^3a_6 - 6v_1^2v_2^2a_6 \\
& + 2v_1^7v_2^2b_4 - 4v_1^7v_2b_4 - 4v_1^5v_2b_4 + v_1^6v_2^3b_5 - 2v_1^6v_2^2b_5 + v_1^6v_2b_5 \\
& - 2v_1^4v_2^2b_5 + 6v_1^4v_2b_5 + v_1^6v_2^2b_2 - 2v_1^6v_2b_2 - 2v_1^4v_2b_2 + 4v_1^3v_2a_2 \\
& + v_1^6b_2 + 4v_1^4b_2 - 4v_1^3a_2 - 2v_1^3b_3 - 2v_1b_3 - 6v_1^2a_1 + 2v_1^3b_1 \\
& + 2v_1^3b_4 - 4v_1^2a_8 - 12v_2^2a_{10} + 3v_1^8b_7 + 8v_1^6b_7 - 2v_1^5b_8 - 2v_1^3b_8 \\
& - 4v_1a_5 - 8v_2a_6 + 2v_1^7b_4 + 6v_1^5b_4 - 2v_1^4a_4 - 2v_1^4b_5 - 2v_1^2b_5 \\
& + b_2v_1^2 + 3v_1^4b_7 - 2v_2a_3 + 2v_1^2a_4 - 2v_2^2a_6 + 4v_1^3a_7 - 2v_2^3a_{10} = 0
\end{aligned} \tag{7E}$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2\}$$

Equation (7E) now becomes

$$\begin{aligned}
& 3v_1^8v_2^2b_7 - 6v_1^8v_2b_7 + 2v_1^7v_2^3b_8 - 4v_1^5v_2^2b_8 + v_1^6v_2^4b_9 - 2v_1^4v_2^3b_9 - 6v_1v_2^2b_{10} \\
& + 6v_1^2v_2^4a_{10} - 2a_1 - 4a_3 + (8b_9 + 2a_8 - 2b_5)v_2^2v_1^4 + (4a_9 + 8b_{10})v_2^3v_1^3 \\
& + (-2a_8 - 4b_9 + 2a_4 + 6b_5 - 2b_2)v_2v_1^4 + (-4a_9 - 6b_{10} + 4a_5 + 6b_6)v_2^2v_1^3 \\
& + (-6a_{10} + 6a_6)v_2^3v_1^2 + (4b_3 + 2b_8 - 4a_5 - 4b_6 + 4a_2)v_2v_1^3 \\
& + (6a_3 + b_9 - 6a_6)v_2^2v_1^2 + (6a_1 + 2a_8 - 4b_9 - 6a_3 + b_5)v_2v_1^2 \\
& + (-8a_9 - 4b_6)v_2v_1 + (-6b_7 + b_5 - 2b_2)v_2v_1^6 \\
& + (-4b_8 + 2b_4)v_2^2v_1^7 + (2b_8 - 4b_4)v_2v_1^7 + (8b_8 - 4b_4)v_2v_1^5 \\
& + (-2b_9 + b_5)v_2^3v_1^6 + (b_9 - 2b_5 + b_2)v_2^2v_1^6 + (-4a_5 - 2b_3)v_1 \\
& + (-2a_3 - 8a_6)v_2 + (-4a_2 - 2b_3 + 2b_1 + 2b_4 - 2b_8 + 4a_7)v_1^3 \\
& + (-6a_1 - 4a_8 - 2b_5 + b_2 + 2a_4)v_1^2 \\
& + (-12a_{10} - 2a_6)v_2^2 + (b_2 + 8b_7)v_1^6 + (-2b_8 + 6b_4)v_1^5 \\
& + (4b_2 - 2a_4 - 2b_5 + 3b_7)v_1^4 + 3v_1^8b_7 + 2v_1^7b_4 - 2v_2^3a_{10} = 0
\end{aligned} \tag{8E}$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned}
 b_9 &= 0 \\
 -2a_{10} &= 0 \\
 6a_{10} &= 0 \\
 2b_4 &= 0 \\
 -6b_7 &= 0 \\
 3b_7 &= 0 \\
 -4b_8 &= 0 \\
 2b_8 &= 0 \\
 -2b_9 &= 0 \\
 -6b_{10} &= 0 \\
 -2a_1 - 4a_3 &= 0 \\
 -2a_3 - 8a_6 &= 0 \\
 -4a_5 - 2b_3 &= 0 \\
 -8a_9 - 4b_6 &= 0 \\
 4a_9 + 8b_{10} &= 0 \\
 -12a_{10} - 2a_6 &= 0 \\
 -6a_{10} + 6a_6 &= 0 \\
 b_2 + 8b_7 &= 0 \\
 -4b_8 + 2b_4 &= 0 \\
 -2b_8 + 6b_4 &= 0 \\
 2b_8 - 4b_4 &= 0 \\
 8b_8 - 4b_4 &= 0 \\
 -2b_9 + b_5 &= 0 \\
 6a_3 + b_9 - 6a_6 &= 0 \\
 -6b_7 + b_5 - 2b_2 &= 0 \\
 b_9 - 2b_5 + b_2 &= 0 \\
 8b_9 + 2a_8 - 2b_5 &= 0 \\
 -4a_9 - 6b_{10} + 4a_5 + 6b_6 &= 0 \\
 4b_2 - 2a_4 - 2b_5 + 3b_7 &= 0 \\
 -6a_1 - 4a_8 - 2b_5 + b_2 + 2a_4 &= 0 \\
 6a_1 + 2a_8 - 4b_9 - 6a_3 + b_5 &= 0 \\
 -2a_8 - 4b_9 + 2a_4 + 6b_5 - 2b_2 &= 0 \\
 4b_3 + 2b_8 - 4a_5 - 4b_6 + 4a_2 &= 0 \\
 -4a_2 - 2b_3 + 2b_1 + 2b_4 - 2b_8 + 4a_7 &= 0
 \end{aligned}$$

Solving the above equations for the unknowns gives

$$a_1 = 0$$

$$a_2 = 0$$

$$a_3 = 0$$

$$a_4 = 0$$

$$a_5 = 0$$

$$a_6 = 0$$

$$a_7 = -\frac{b_1}{2}$$

$$a_8 = 0$$

$$a_9 = 0$$

$$a_{10} = 0$$

$$b_1 = b_1$$

$$b_2 = 0$$

$$b_3 = 0$$

$$b_4 = 0$$

$$b_5 = 0$$

$$b_6 = 0$$

$$b_7 = 0$$

$$b_8 = 0$$

$$b_9 = 0$$

$$b_{10} = 0$$

Substituting the above solution in the ansatz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\xi = -\frac{x^3}{2}$$

$$\eta = 1$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned}\eta &= \eta - \omega(x, y) \xi \\ &= 1 - \left(\frac{2}{x(yx^2 - x^2 - 1)} \right) \left(-\frac{x^3}{2} \right) \\ &= \frac{yx^2 - 1}{yx^2 - x^2 - 1} \\ \xi &= 0\end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y} \right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{yx^2 - 1}{yx^2 - x^2 - 1}} dy\end{aligned}$$

Which results in

$$S = y - \ln(yx^2 - 1)$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{2}{x(yx^2 - x^2 - 1)}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= -\frac{2xy}{yx^2 - 1} \\ S_y &= 1 - \frac{x^2}{yx^2 - 1} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = -\frac{2}{x} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = -\frac{2}{R}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = -2 \ln(R) + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$y - \ln(yx^2 - 1) = -2 \ln(x) + c_1$$



Which simplifies to

$$y - \ln(yx^2 - 1) = -2 \ln(x) + c_1$$

Which gives

$$y = -\frac{\text{LambertW}\left(-e^{\frac{2 \ln(x)x^2 - c_1 x^2 + 1}{x^2}}\right) x^2 - 1}{x^2}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{2}{x(yx^2 - x^2 - 1)}$ 	$R = x$ $S = y - \ln(yx^2 - 1)$	$\frac{dS}{dR} = -\frac{2}{R}$ 

Summary

The solution(s) found are the following

$$y = -\frac{\text{LambertW}\left(-e^{\frac{2 \ln(x)x^2 - c_1 x^2 + 1}{x^2}}\right) x^2 - 1}{x^2} \quad (1)$$




Figure 101: Slope field plot

Verification of solutions

$$y = -\frac{\text{LambertW}\left(-e^{\frac{2\ln(x)x^2 - c_1x^2 + 1}{x^2}}\right)x^2 - 1}{x^2}$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying Chini  
differential order: 1; looking for linear symmetries  
trying exact  
trying Abel  
<- Abel AIR successful: ODE belongs to the 1F1 1-parameter class`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 22

```
dsolve(x*diff(y(x),x)+2=x^3*(y(x)-1)*diff(y(x),x),y(x), singsol=all)
```

$$y(x) = \frac{-\text{LambertW}\left(c_1 e^{\frac{1}{x^2}}\right) x^2 + 1}{x^2}$$

✓ Solution by Mathematica

Time used: 0.405 (sec). Leaf size: 33

```
DSolve[x*y'[x]+2==x^3*(y[x]-1)*y'[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{x^2} - W\left(e^{\frac{1}{x^2} + \frac{1}{2}(-2-9\sqrt[3]{-2}c_1)}\right)$$

3.24 problem 6

3.24.1 Solving as separable ode	478
3.24.2 Solving as linear ode	480
3.24.3 Solving as homogeneousTypeD2 ode	481
3.24.4 Solving as first order ode lie symmetry lookup ode	483
3.24.5 Solving as exact ode	487
3.24.6 Maple step by step solution	491

Internal problem ID [6180]

Internal file name [OUTPUT/5428_Sunday_June_05_2022_03_37_04_PM_41613258/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.4 First Order Linear Equations. Page 15

Problem number: 6.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "linear", "separable", "homogeneousTypeD2", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_separable]

$$xy' - 2yx^2 - \ln(x)y = 0$$

3.24.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{y(2x^2 + \ln(x))}{x}\end{aligned}$$

Where $f(x) = \frac{2x^2 + \ln(x)}{x}$ and $g(y) = y$. Integrating both sides gives

$$\begin{aligned}\frac{1}{y} dy &= \frac{2x^2 + \ln(x)}{x} dx \\ \int \frac{1}{y} dy &= \int \frac{2x^2 + \ln(x)}{x} dx \\ \ln(y) &= x^2 + \frac{\ln(x)^2}{2} + c_1 \\ y &= e^{x^2 + \frac{\ln(x)^2}{2} + c_1} \\ &= c_1 e^{x^2 + \frac{\ln(x)^2}{2}}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{x^2 + \frac{\ln(x)^2}{2}} \quad (1)$$




Figure 102: Slope field plot

Verification of solutions

$$y = c_1 e^{x^2 + \frac{\ln(x)^2}{2}}$$

Verified OK.

3.24.2 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -\frac{2x^2 + \ln(x)}{x}$$

$$q(x) = 0$$

Hence the ode is

$$y' - \frac{y(2x^2 + \ln(x))}{x} = 0$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int -\frac{2x^2 + \ln(x)}{x} dx} \\ &= e^{-x^2 - \frac{\ln(x)^2}{2}}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}\mu y &= 0 \\ \frac{d}{dx}\left(e^{-x^2 - \frac{\ln(x)^2}{2}}y\right) &= 0\end{aligned}$$

Integrating gives

$$e^{-x^2 - \frac{\ln(x)^2}{2}}y = c_1$$

Dividing both sides by the integrating factor $\mu = e^{-x^2 - \frac{\ln(x)^2}{2}}$ results in

$$y = c_1 e^{x^2 + \frac{\ln(x)^2}{2}}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{x^2 + \frac{\ln(x)^2}{2}} \quad (1)$$




Figure 103: Slope field plot

Verification of solutions

$$y = c_1 e^{x^2 + \frac{\ln(x)^2}{2}}$$

Verified OK.

3.24.3 Solving as homogeneousTypeD2 ode

Using the change of variables $y = u(x)x$ on the above ode results in new ode in $u(x)$

$$x(u'(x)x + u(x)) - 2u(x)x^3 - \ln(x)u(x)x = 0$$

In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= \frac{u(2x^2 + \ln(x) - 1)}{x} \end{aligned}$$

Where $f(x) = \frac{2x^2 + \ln(x) - 1}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned} \frac{1}{u} du &= \frac{2x^2 + \ln(x) - 1}{x} dx \\ \int \frac{1}{u} du &= \int \frac{2x^2 + \ln(x) - 1}{x} dx \\ \ln(u) &= x^2 + \frac{\ln(x)^2}{2} - \ln(x) + c_2 \\ u &= e^{x^2 + \frac{\ln(x)^2}{2} - \ln(x) + c_2} \\ &= c_2 e^{x^2 + \frac{\ln(x)^2}{2} - \ln(x)} \end{aligned}$$

Therefore the solution y is

$$\begin{aligned} y &= xu \\ &= xc_2 e^{x^2 + \frac{\ln(x)^2}{2} - \ln(x)} \end{aligned}$$

Summary

The solution(s) found are the following

$$y = xc_2 e^{x^2 + \frac{\ln(x)^2}{2} - \ln(x)} \quad (1)$$




Figure 104: Slope field plot

Verification of solutions

$$y = xc_2e^{x^2 + \frac{\ln(x)^2}{2} - \ln(x)}$$

Verified OK.

3.24.4 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{y(2x^2 + \ln(x))}{x}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2\xi_y - \omega_x\xi - \omega_y\eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 81: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= e^{x^2 + \frac{\ln(x)^2}{2}}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^{x^2 + \frac{\ln(x)^2}{2}}} dy \end{aligned}$$

Which results in

$$S = e^{-x^2 - \frac{\ln(x)^2}{2}} y$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{y(2x^2 + \ln(x))}{x}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= -\frac{(2x^2 + \ln(x)) e^{-x^2 - \frac{\ln(x)^2}{2}} y}{x} \\ S_y &= e^{-x^2 - \frac{\ln(x)^2}{2}} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \tag{4}$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$e^{-x^2 - \frac{\ln(x)^2}{2}} y = c_1$$



Which simplifies to

$$e^{-x^2 - \frac{\ln(x)^2}{2}} y = c_1$$

Which gives

$$y = c_1 e^{x^2 + \frac{\ln(x)^2}{2}}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{y(2x^2 + \ln(x))}{x}$ 	$R = x$ $S = e^{-x^2 - \frac{\ln(x)^2}{2}} y$	$\frac{dS}{dR} = 0$ 

Summary

The solution(s) found are the following

$$y = c_1 e^{x^2 + \frac{\ln(x)^2}{2}} \tag{1}$$




Figure 105: Slope field plot

Verification of solutions

$$y = c_1 e^{x^2 + \frac{\ln(x)^2}{2}}$$

Verified OK.

3.24.5 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned}\left(\frac{1}{y}\right) dy &= \left(\frac{2x^2 + \ln(x)}{x}\right) dx \\ \left(-\frac{2x^2 + \ln(x)}{x}\right) dx + \left(\frac{1}{y}\right) dy &= 0\end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -\frac{2x^2 + \ln(x)}{x} \\ N(x, y) &= \frac{1}{y}\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(-\frac{2x^2 + \ln(x)}{x}\right) \\ &= 0\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{1}{y} \right) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\frac{2x^2 + \ln(x)}{x} dx \\ \phi &= -x^2 - \frac{\ln(x)^2}{2} + f(y)\end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{1}{y}$. Therefore equation (4) becomes

$$\frac{1}{y} = 0 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = \frac{1}{y}$$

Integrating the above w.r.t y gives

$$\int f'(y) dy = \int \left(\frac{1}{y}\right) dy$$

$$f(y) = \ln(y) + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -x^2 - \frac{\ln(x)^2}{2} + \ln(y) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -x^2 - \frac{\ln(x)^2}{2} + \ln(y)$$

The solution becomes

$$y = e^{x^2 + \frac{\ln(x)^2}{2} + c_1}$$

Summary

The solution(s) found are the following

$$y = e^{x^2 + \frac{\ln(x)^2}{2} + c_1} \tag{1}$$




Figure 106: Slope field plot

Verification of solutions

$$y = e^{x^2 + \frac{\ln(x)^2}{2} + c_1}$$

Verified OK.

3.24.6 Maple step by step solution

Let's solve

$$xy' - 2yx^2 - \ln(x)y = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y} = \frac{2x^2 + \ln(x)}{x}$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y} dx = \int \frac{2x^2 + \ln(x)}{x} dx + c_1$$

- Evaluate integral

$$\ln(y) = x^2 + \frac{\ln(x)^2}{2} + c_1$$

- Solve for y

$$y = e^{x^2 + \frac{\ln(x)^2}{2} + c_1}$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(x*diff(y(x),x)=2*x^2*y(x)+y(x)*ln(x),y(x), singsol=all)
```

$$y(x) = c_1 e^{\frac{\ln(x)^2}{2} + x^2}$$

✓ Solution by Mathematica

Time used: 0.038 (sec). Leaf size: 27

```
DSolve[x*y'[x]==2*x^2*y[x]+y[x]*Log[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{x^2 + \frac{\log^2(x)}{2}}$$

$$y(x) \rightarrow 0$$

3.25 problem 7

3.25.1 Solving as linear ode	493
3.25.2 Solving as first order ode lie symmetry lookup ode	495
3.25.3 Solving as exact ode	499
3.25.4 Maple step by step solution	503

Internal problem ID [6181]

Internal file name [OUTPUT/5429_Sunday_June_05_2022_03_37_05_PM_32380097/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.4 First Order Linear Equations. Page 15

Problem number: 7.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**linear**", "**exactWithIntegrationFactor**", "**first_order_ode_lie_symmetry_lookup**"

Maple gives the following as the ode type

`[_linear]`

$$y' \sin(2x) - 2y = 2 \cos(x)$$

3.25.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -2 \csc(2x)$$

$$q(x) = \csc(x)$$

Hence the ode is

$$y' - 2 \csc(2x)y = \csc(x)$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int -2 \csc(2x) dx} \\ &= \frac{1}{\csc(2x) - \cot(2x)}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu) (\csc(x)) \\ \frac{d}{dx} \left(\frac{y}{\csc(2x) - \cot(2x)} \right) &= \left(\frac{1}{\csc(2x) - \cot(2x)} \right) (\csc(x)) \\ d \left(\frac{y}{\csc(2x) - \cot(2x)} \right) &= (\csc(x) \cot(x)) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}\frac{y}{\csc(2x) - \cot(2x)} &= \int \csc(x) \cot(x) dx \\ \frac{y}{\csc(2x) - \cot(2x)} &= -\csc(x) + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = \frac{1}{\csc(2x) - \cot(2x)}$ results in

$$y = -(\csc(2x) - \cot(2x)) \csc(x) + c_1(\csc(2x) - \cot(2x))$$

which simplifies to

$$y = c_1 \tan(x) - \sec(x)$$

Summary

The solution(s) found are the following

$$y = c_1 \tan(x) - \sec(x) \tag{1}$$




Figure 107: Slope field plot

Verification of solutions

$$y = c_1 \tan(x) - \sec(x)$$

Verified OK.

3.25.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{2y + 2 \cos(x)}{\sin(2x)}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 84: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned} \xi(x, y) &= 0 \\ \eta(x, y) &= \frac{1}{\csc(2x) + \cot(2x)} \end{aligned} \tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{1}{\csc(2x) + \cot(2x)}} dy \end{aligned}$$

Which results in

$$S = (\csc(2x) + \cot(2x)) y$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{2y + 2 \cos(x)}{\sin(2x)}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= -2y(1 + \cos(2x)) \csc(2x)^2 \\ S_y &= \csc(2x) + \cot(2x) \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \csc(x) \cot(x) \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \csc(R) \cot(R)$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = -\csc(R) + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$(\csc(2x) + \cot(2x))y = -\csc(x) + c_1$$



Which simplifies to

$$y \cot(x) - c_1 + \csc(x) = 0$$

Which gives

$$y = -\frac{\csc(x) - c_1}{\cot(x)}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{2y+2\cos(x)}{\sin(2x)}$ 	$R = x$ $S = (\csc(2x) + \cot(2x))y$	$\frac{dS}{dR} = \csc(R) \cot(R)$ 

Summary

The solution(s) found are the following

$$y = -\frac{\csc(x) - c_1}{\cot(x)} \quad (1)$$




Figure 108: Slope field plot

Verification of solutions

$$y = -\frac{\csc(x) - c_1}{\cot(x)}$$

Verified OK.

3.25.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}(\sin(2x)) dy &= (2y + 2 \cos(x)) dx \\ (-2y - 2 \cos(x)) dx + (\sin(2x)) dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -2y - 2 \cos(x) \\ N(x, y) &= \sin(2x)\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-2y - 2 \cos(x)) \\ &= -2\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(\sin(2x)) \\ &= 2 \cos(2x)\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= \csc(2x) ((-2) - (2 \cos(2x))) \\ &= -2 \csc(2x) - 2 \cot(2x) \end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int A dx} \\ &= e^{\int -2 \csc(2x) - 2 \cot(2x) dx} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^{\frac{\ln(\cot(2x)^2 + 1)}{2} + \ln(\csc(2x) + \cot(2x))} \\ &= \text{csgn}(\csc(2x)) \csc(2x) (\csc(2x) + \cot(2x)) \end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \overline{M} and \overline{N} for now so not to confuse them with the original M and N .

$$\begin{aligned} \overline{M} &= \mu M \\ &= \text{csgn}(\csc(2x)) \csc(2x) (\csc(2x) + \cot(2x)) (-2y - 2 \cos(x)) \\ &= (-y - \cos(x)) \text{csgn}(\csc(2x)) \csc(x)^2 \end{aligned}$$

And

$$\begin{aligned} \overline{N} &= \mu N \\ &= \text{csgn}(\csc(2x)) \csc(2x) (\csc(2x) + \cot(2x)) (\sin(2x)) \\ &= \text{csgn}(\csc(2x)) (\csc(2x) + \cot(2x)) \end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned} \overline{M} + \overline{N} \frac{dy}{dx} &= 0 \\ ((-y - \cos(x)) \text{csgn}(\csc(2x)) \csc(x)^2) + (\text{csgn}(\csc(2x)) (\csc(2x) + \cot(2x))) \frac{dy}{dx} &= 0 \end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \overline{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \overline{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\int \frac{\partial \phi}{\partial x} dx = \int \overline{M} dx$$

$$\int \frac{\partial \phi}{\partial x} dx = \int (-y - \cos(x)) \operatorname{csgn}(\csc(2x)) \csc(x)^2 dx$$

$$\phi = \operatorname{csgn}(\csc(2x)) (y \cot(x) + \csc(x)) + f(y) \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = \operatorname{csgn}(\csc(2x)) \cot(x) + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \operatorname{csgn}(\csc(2x)) (\csc(2x) + \cot(2x))$. Therefore equation (4) becomes

$$\operatorname{csgn}(\csc(2x)) (\csc(2x) + \cot(2x)) = \operatorname{csgn}(\csc(2x)) \cot(x) + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = \operatorname{csgn}(\csc(2x)) (y \cot(x) + \csc(x)) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \operatorname{csgn}(\csc(2x)) (y \cot(x) + \csc(x))$$

The solution becomes

$$y = -\frac{\csc(x) \operatorname{csgn}(\csc(2x)) - c_1}{\operatorname{csgn}(\csc(2x)) \cot(x)}$$

Summary

The solution(s) found are the following

Simplifying the solution $y = -\frac{\csc(x) \operatorname{csgn}(\csc(2x)) - c_1}{\operatorname{csgn}(\csc(2x)) \cot(x)}$ to $y = -\frac{\csc(x) - c_1}{\cot(x)}$

$y =$




Figure 109: Slope field plot

Verification of solutions

$$y = -\frac{\csc(x) - c_1}{\cot(x)}$$

Verified OK.

3.25.4 Maple step by step solution

Let's solve

$$y' \sin(2x) - 2y = 2 \cos(x)$$

- Highest derivative means the order of the ODE is 1

y'

- Isolate the derivative

$$y' = \frac{2y}{\sin(2x)} + \frac{2\cos(x)}{\sin(2x)}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' - \frac{2y}{\sin(2x)} = \frac{2\cos(x)}{\sin(2x)}$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) \left(y' - \frac{2y}{\sin(2x)} \right) = \frac{2\mu(x)\cos(x)}{\sin(2x)}$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x)y)$

$$\mu(x) \left(y' - \frac{2y}{\sin(2x)} \right) = \mu'(x)y + \mu(x)y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = -\frac{2\mu(x)}{\sin(2x)}$$

- Solve to find the integrating factor

$$\mu(x) = \frac{1}{\tan(x)}$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x)y) \right) dx = \int \frac{2\mu(x)\cos(x)}{\sin(2x)} dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x)y = \int \frac{2\mu(x)\cos(x)}{\sin(2x)} dx + c_1$$

- Solve for y

$$y = \frac{\int \frac{2\mu(x)\cos(x)}{\sin(2x)} dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = \frac{1}{\tan(x)}$

$$y = \tan(x) \left(\int \frac{2\cos(x)}{\tan(x)\sin(2x)} dx + c_1 \right)$$

- Evaluate the integrals on the rhs

$$y = \tan(x) (-\csc(x) + c_1)$$

- Simplify

$$y = c_1 \tan(x) - \sec(x)$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(diff(y(x),x)*sin(2*x)=2*y(x)+2*cos(x),y(x), singsol=all)
```

$$y(x) = \tan(x) c_1 - \sec(x)$$

✓ Solution by Mathematica

Time used: 0.04 (sec). Leaf size: 15

```
DSolve[y'[x]*Sin[2*x]==2*y[x]+2*Cos[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \sec(x)(-1 + c_1 \sin(x))$$

4 Chapter 1. What is a differential equation.

Section 1.5. Exact Equations. Page 20

4.1	problem 1	507
4.2	problem 2	519
4.3	problem 3	522
4.4	problem 4	531
4.5	problem 5	534
4.6	problem 6	550
4.7	problem 7	562
4.8	problem 8	568
4.9	problem 9	574
4.10	problem 10	592
4.11	problem 11	599
4.12	problem 12	608
4.13	problem 13	614
4.14	problem 14	623
4.15	problem 15	637
4.16	problem 16	646
4.17	problem 17	652
4.18	problem 18	658
4.19	problem 19	664
4.20	problem 20	671
4.21	problem 21	685

4.1 problem 1

4.1.1	Solving as first order ode lie symmetry calculated ode	507
4.1.2	Solving as exact ode	512
4.1.3	Maple step by step solution	516

Internal problem ID [6182]

Internal file name [OUTPUT/5430_Sunday_June_05_2022_03_37_07_PM_5121642/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.5. Exact Equations. Page 20

Problem number: 1.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "first_order_ode_lie_symmetry_calculated"

Maple gives the following as the ode type

```
[[_homogeneous, `class G`], _exact, _rational, [_Abel, `2nd
  type`, `class B`]]
```

$$\left(x + \frac{2}{y}\right) y' + y = 0$$

4.1.1 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = -\frac{y^2}{xy + 2}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \tag{A}$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \tag{1E}$$

$$\eta = xb_2 + yb_3 + b_1 \tag{2E}$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$b_2 - \frac{y^2(b_3 - a_2)}{xy + 2} - \frac{y^4 a_3}{(xy + 2)^2} - \frac{y^3(xa_2 + ya_3 + a_1)}{(xy + 2)^2} - \left(-\frac{2y}{xy + 2} + \frac{y^2 x}{(xy + 2)^2} \right) (xb_2 + yb_3 + b_1) = 0 \quad (5E)$$

Putting the above in normal form gives

$$\frac{2x^2 y^2 b_2 - 2y^4 a_3 + x y^2 b_1 - y^3 a_1 + 8xyb_2 + 2y^2 a_2 + 2y^2 b_3 + 4yb_1 + 4b_2}{(xy + 2)^2} = 0$$

Setting the numerator to zero gives

$$2x^2 y^2 b_2 - 2y^4 a_3 + x y^2 b_1 - y^3 a_1 + 8xyb_2 + 2y^2 a_2 + 2y^2 b_3 + 4yb_1 + 4b_2 = 0 \quad (6E)$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2\}$$

The above PDE (6E) now becomes

$$-2a_3 v_2^4 + 2b_2 v_1^2 v_2^2 - a_1 v_2^3 + b_1 v_1 v_2^2 + 2a_2 v_2^2 + 8b_2 v_1 v_2 + 2b_3 v_2^2 + 4b_1 v_2 + 4b_2 = 0 \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2\}$$

Equation (7E) now becomes

$$2b_2 v_1^2 v_2^2 + b_1 v_1 v_2^2 + 8b_2 v_1 v_2 - 2a_3 v_2^4 - a_1 v_2^3 + (2a_2 + 2b_3) v_2^2 + 4b_1 v_2 + 4b_2 = 0 \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned}b_1 &= 0 \\-a_1 &= 0 \\-2a_3 &= 0 \\4b_1 &= 0 \\2b_2 &= 0 \\4b_2 &= 0 \\8b_2 &= 0 \\2a_2 + 2b_3 &= 0\end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned}a_1 &= 0 \\a_2 &= -b_3 \\a_3 &= 0 \\b_1 &= 0 \\b_2 &= 0 \\b_3 &= b_3\end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned}\xi &= -x \\ \eta &= y\end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned}\eta &= \eta - \omega(x, y) \xi \\ &= y - \left(-\frac{y^2}{xy + 2} \right) (-x) \\ &= \frac{2y}{xy + 2} \\ \xi &= 0\end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{2y}{xy+2}} dy \end{aligned}$$

Which results in

$$S = \frac{xy}{2} + \ln(y)$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -\frac{y^2}{xy + 2}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{y}{2} \\ S_y &= \frac{x}{2} + \frac{1}{y} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \tag{4}$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{xy}{2} + \ln(y) = c_1$$



Which simplifies to

$$\frac{xy}{2} + \ln(y) = c_1$$

Which gives

$$y = e^{-\text{LambertW}\left(\frac{x e^{c_1}}{2}\right) + c_1}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -\frac{y^2}{xy+2}$ 	$R = x$ $S = \frac{xy}{2} + \ln(y)$	$\frac{dS}{dR} = 0$ 

Summary

The solution(s) found are the following

$$y = e^{-\text{LambertW}\left(\frac{x e^{c_1}}{2}\right) + c_1} \quad (1)$$




Figure 110: Slope field plot

Verification of solutions

$$y = e^{-\text{LambertW}\left(\frac{x e^{c_1}}{2}\right) + c_1}$$

Verified OK.

4.1.2 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} \left(x + \frac{2}{y}\right) dy &= (-y) dx \\ (y) dx + \left(x + \frac{2}{y}\right) dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= y \\ N(x, y) &= x + \frac{2}{y} \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(y) \\ &= 1 \end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(x + \frac{2}{y} \right) \\ &= 1\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int y dx \\ \phi &= xy + f(y)\end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = x + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = x + \frac{2}{y}$. Therefore equation (4) becomes

$$x + \frac{2}{y} = x + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = \frac{2}{y}$$

Integrating the above w.r.t y gives

$$\begin{aligned}\int f'(y) dy &= \int \left(\frac{2}{y} \right) dy \\ f(y) &= 2 \ln(y) + c_1\end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = xy + 2 \ln(y) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = xy + 2 \ln(y)$$

The solution becomes

$$y = e^{-\text{LambertW}\left(\frac{x e^{\frac{c_1}{2}}}{2}\right) + \frac{c_1}{2}}$$

Summary

The solution(s) found are the following

$$y = e^{-\text{LambertW}\left(\frac{x e^{\frac{c_1}{2}}}{2}\right) + \frac{c_1}{2}} \tag{1}$$




Figure 111: Slope field plot

Verification of solutions

$$y = e^{-\text{LambertW}\left(\frac{x e^{\frac{c_1}{2}}}{2}\right) + \frac{c_1}{2}}$$

Verified OK.

4.1.3 Maple step by step solution

Let's solve

$$\left(x + \frac{2}{y}\right) y' + y = 0$$

- Highest derivative means the order of the ODE is 1
 y'
- Check if ODE is exact
 - ODE is exact if the lhs is the total derivative of a C^2 function
 $F'(x, y) = 0$
 - Compute derivative of lhs
 $F'(x, y) + \left(\frac{\partial}{\partial y} F(x, y)\right) y' = 0$
 - Evaluate derivatives
 $1 = 1$
 - Condition met, ODE is exact
- Exact ODE implies solution will be of this form
 $\left[F(x, y) = c_1, M(x, y) = F'(x, y), N(x, y) = \frac{\partial}{\partial y} F(x, y)\right]$
- Solve for $F(x, y)$ by integrating $M(x, y)$ with respect to x
 $F(x, y) = \int y dx + f_1(y)$
- Evaluate integral
 $F(x, y) = xy + f_1(y)$
- Take derivative of $F(x, y)$ with respect to y
 $N(x, y) = \frac{\partial}{\partial y} F(x, y)$
- Compute derivative
 $x + \frac{2}{y} = x + \frac{d}{dy} f_1(y)$

- Isolate for $\frac{d}{dy}f_1(y)$

$$\frac{d}{dy}f_1(y) = \frac{2}{y}$$
- Solve for $f_1(y)$

$$f_1(y) = 2 \ln(y)$$
- Substitute $f_1(y)$ into equation for $F(x, y)$

$$F(x, y) = xy + 2 \ln(y)$$
- Substitute $F(x, y)$ into the solution of the ODE

$$xy + 2 \ln(y) = c_1$$
- Solve for y

$$y = e^{-\text{LambertW}\left(\frac{x e^{\frac{c_1}{2}}}{2}\right) + \frac{c_1}{2}}$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
<- 1st order linear successful
<- inverse linear successful`

```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 17

```
dsolve((x+2/y(x))*diff(y(x),x)+y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{2 \text{LambertW}\left(\frac{x e^{\frac{c_1}{2}}}{2}\right)}{x}$$

✓ Solution by Mathematica

Time used: 17.046 (sec). Leaf size: 58

```
DSolve[(x+2/y[x])*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2W\left(-\frac{1}{2}\sqrt{e^{c_1}x^2}\right)}{x}$$

$$y(x) \rightarrow \frac{2W\left(\frac{1}{2}\sqrt{e^{c_1}x^2}\right)}{x}$$

$$y(x) \rightarrow 0$$

4.2 problem 2

Internal problem ID [6183]

Internal file name [OUTPUT/5431_Sunday_June_05_2022_03_37_08_PM_33515694/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.5. Exact Equations. Page 20

Problem number: 2.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**unknown**"

Maple gives the following as the ode type

[`x=_G(y,y')`]

Unable to solve or complete the solution.

$$\sin(x) \tan(y) + \cos(x) \sec(x)^2 yy' = -1$$

Unable to determine ODE type.

Maple trace

```
`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying Chini
differential order: 1; looking for linear symmetries
trying exact
Looking for potential symmetries
trying inverse_Riccati
trying an equivalence to an Abel ODE
differential order: 1; trying a linearization to 2nd order
--- trying a change of variables {x -> y(x), y(x) -> x}
differential order: 1; trying a linearization to 2nd order
trying 1st order ODE linearizable_by_differentiation
--- Trying Lie symmetry methods, 1st order ---
`, `-> Computing symmetries using: way = 3
`, `-> Computing symmetries using: way = 4
`, `-> Computing symmetries using: way = 5
trying symmetry patterns for 1st order ODEs
-> trying a symmetry pattern of the form [F(x)*G(y), 0]
-> trying a symmetry pattern of the form [0, F(x)*G(y)]
-> trying symmetry patterns of the forms [F(x),G(y)] and [G(y),F(x)]
`, `-> Computing symmetries using: way = HINT
-> Calling odsolve with the ODE`, diff(y(x), x)-2*y(x)*(tan(x)*sin(2*x)-1)/sin(2*x), y(x)
  Methods for first order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  trying 1st order linear
  <- 1st order linear successful
-> Calling odsolve with the ODE`, diff(y(x), x)-y(x)*(2*cos(x)*tan(x)-sin(x))/cos(x), y(x)
  Methods for first order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  trying 1st order linear
  <- 1st order linear successful
`, `-> Computing symmetries using: way = HINT
-> Calling odsolve with the ODE`, diff(y(x), x)+y(x)/x, y(x)`      *** Sublevel 2 ***
  Methods for first order ODEs: 520
  --- Trying classification methods ---
  trying a quadrature
  trying 1st order linear
```

X Solution by Maple

```
dsolve((sin(x)*tan(y(x))+1)+(cos(x)*sec(x)^2*y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

No solution found

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[(Sin[x]*Tan[y[x]]+1)+(Cos[x]*Sec[x]^2*y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions
```

Not solved

4.3 problem 3

4.3.1 Solving as differentialType ode	522
4.3.2 Solving as exact ode	524
4.3.3 Maple step by step solution	527

Internal problem ID [6184]

Internal file name [OUTPUT/5432_Sunday_June_05_2022_03_37_56_PM_80988887/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.5. Exact Equations. Page 20

Problem number: 3.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "differentialType"

Maple gives the following as the ode type

```
[_exact, _rational]
```

$$y + (x + y^3) y' = x^3$$

4.3.1 Solving as differentialType ode

Writing the ode as

$$y' = \frac{-y + x^3}{x + y^3} \tag{1}$$

Which becomes

$$(y^3) dy = (-x) dy + (x^3 - y) dx \tag{2}$$

But the RHS is complete differential because

$$(-x) dy + (x^3 - y) dx = d\left(\frac{1}{4}x^4 - xy\right)$$

Hence (2) becomes

$$(y^3) dy = d\left(\frac{1}{4}x^4 - xy\right)$$

Integrating both sides gives gives the solution as

$$\frac{y^4}{4} = \frac{x^4}{4} - xy + c_1$$

Summary

The solution(s) found are the following

$$\frac{y^4}{4} = \frac{x^4}{4} - xy + c_1 \tag{1}$$




Figure 112: Slope field plot

Verification of solutions

$$\frac{y^4}{4} = \frac{x^4}{4} - xy + c_1$$

Verified OK.

4.3.2 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} (y^3 + x) dy &= (x^3 - y) dx \\ (-x^3 + y) dx + (y^3 + x) dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -x^3 + y \\ N(x, y) &= y^3 + x \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-x^3 + y) \\ &= 1\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(y^3 + x) \\ &= 1\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \tag{1}$$

$$\frac{\partial \phi}{\partial y} = N \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -x^3 + y dx \\ \phi &= -\frac{1}{4}x^4 + xy + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = x + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = y^3 + x$. Therefore equation (4) becomes

$$y^3 + x = x + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = y^3$$

Integrating the above w.r.t y gives

$$\int f'(y) dy = \int (y^3) dy$$

$$f(y) = \frac{y^4}{4} + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\frac{1}{4}x^4 + xy + \frac{1}{4}y^4 + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\frac{1}{4}x^4 + xy + \frac{1}{4}y^4$$

Summary

The solution(s) found are the following

$$\frac{y^4}{4} - \frac{x^4}{4} + xy = c_1 \tag{1}$$




Figure 113: Slope field plot

Verification of solutions

$$\frac{y^4}{4} - \frac{x^4}{4} + xy = c_1$$

Verified OK.

4.3.3 Maple step by step solution

Let's solve

$$y + (x + y^3) y' = x^3$$

- Highest derivative means the order of the ODE is 1
 y'
- Check if ODE is exact
 - ODE is exact if the lhs is the total derivative of a C^2 function
 $F'(x, y) = 0$
 - Compute derivative of lhs
 $F'(x, y) + \left(\frac{\partial}{\partial y} F(x, y)\right) y' = 0$

- Evaluate derivatives
 $1 = 1$
- Condition met, ODE is exact
- Exact ODE implies solution will be of this form

$$\left[F(x, y) = c_1, M(x, y) = F'(x, y), N(x, y) = \frac{\partial}{\partial y} F(x, y) \right]$$
- Solve for $F(x, y)$ by integrating $M(x, y)$ with respect to x

$$F(x, y) = \int (-x^3 + y) dx + f_1(y)$$
- Evaluate integral

$$F(x, y) = -\frac{x^4}{4} + xy + f_1(y)$$
- Take derivative of $F(x, y)$ with respect to y

$$N(x, y) = \frac{\partial}{\partial y} F(x, y)$$
- Compute derivative

$$y^3 + x = x + \frac{d}{dy} f_1(y)$$
- Isolate for $\frac{d}{dy} f_1(y)$

$$\frac{d}{dy} f_1(y) = y^3$$
- Solve for $f_1(y)$

$$f_1(y) = \frac{y^4}{4}$$
- Substitute $f_1(y)$ into equation for $F(x, y)$

$$F(x, y) = -\frac{1}{4}x^4 + xy + \frac{1}{4}y^4$$
- Substitute $F(x, y)$ into the solution of the ODE

$$-\frac{1}{4}x^4 + xy + \frac{1}{4}y^4 = c_1$$
- Solve for y

$$y = \text{RootOf}(_Z^4 - x^4 + 4x_Z - 4c_1)$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying Chini  
differential order: 1; looking for linear symmetries  
trying exact  
<- exact successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve((y(x)-x^3)+(x+y(x)^3)*diff(y(x),x)=0,y(x), singsol=all)
```

$$-\frac{x^4}{4} + xy(x) + \frac{y(x)^4}{4} + c_1 = 0$$

✓ Solution by Mathematica

Time used: 60.218 (sec). Leaf size: 1210

`DSolve[(y[x]-x^3)+(x+y[x]^3)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \sqrt{\sqrt[3]{9x^2 + \sqrt{3}\sqrt{27x^4 + (x^4 + 4c_1)^3}} - \frac{\sqrt[3]{3}(x^4+4c_1)}{\sqrt[3]{9x^2 + \sqrt{3}\sqrt{27x^4 + (x^4 + 4c_1)^3}}} + \sqrt{\sqrt[3]{9x^2 + \sqrt{3}\sqrt{27x^4 + (x^4 + 4c_1)^3}}}$$

$$y(x) \rightarrow \sqrt{\sqrt[3]{9x^2 + \sqrt{3}\sqrt{27x^4 + (x^4 + 4c_1)^3}} - \frac{6\sqrt{2}x}{\sqrt[3]{9x^2 + \sqrt{3}\sqrt{27x^4 + (x^4 + 4c_1)^3}}} - \sqrt[3]{9x^2 + \sqrt{3}\sqrt{27x^4 + (x^4 + 4c_1)^3}}}$$

$$y(x) \rightarrow \sqrt{\sqrt[3]{9x^2 + \sqrt{3}\sqrt{27x^4 + (x^4 + 4c_1)^3}} - \frac{\sqrt[3]{3}(x^4+4c_1)}{\sqrt[3]{9x^2 + \sqrt{3}\sqrt{27x^4 + (x^4 + 4c_1)^3}}} - \sqrt{\sqrt[3]{9x^2 + \sqrt{3}\sqrt{27x^4 + (x^4 + 4c_1)^3}}}$$

$$y(x) \rightarrow \sqrt{\sqrt[3]{9x^2 + \sqrt{3}\sqrt{27x^4 + (x^4 + 4c_1)^3}} - \frac{\sqrt[3]{3}(x^4+4c_1)}{\sqrt[3]{9x^2 + \sqrt{3}\sqrt{27x^4 + (x^4 + 4c_1)^3}}} + \sqrt{\sqrt[3]{9x^2 + \sqrt{3}\sqrt{27x^4 + (x^4 + 4c_1)^3}}}$$

4.4 problem 4

Internal problem ID [6185]

Internal file name [OUTPUT/5433_Sunday_June_05_2022_03_37_57_PM_4773145/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.5. Exact Equations. Page 20

Problem number: 4.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**unknown**"

Maple gives the following as the ode type

```
[_rational, [_Abel, `2nd type`, `class B`]]
```

Unable to solve or complete the solution.

$$2y^2 - (4 - 2y + 4xy)y' = 4x - 5$$

Unable to determine ODE type.

Maple trace

```
`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying Chini
differential order: 1; looking for linear symmetries
trying exact
trying Abel
Looking for potential symmetries
Looking for potential symmetries
Looking for potential symmetries
trying inverse_Riccati
trying an equivalence to an Abel ODE
differential order: 1; trying a linearization to 2nd order
--- trying a change of variables {x -> y(x), y(x) -> x}
differential order: 1; trying a linearization to 2nd order
trying 1st order ODE linearizable_by_differentiation
--- Trying Lie symmetry methods, 1st order ---
`, `-> Computing symmetries using: way = 3
`, `-> Computing symmetries using: way = 4
`, `-> Computing symmetries using: way = 2
trying symmetry patterns for 1st order ODEs
-> trying a symmetry pattern of the form [F(x)*G(y), 0]
-> trying a symmetry pattern of the form [0, F(x)*G(y)]
-> trying symmetry patterns of the forms [F(x),G(y)] and [G(y),F(x)]
-> trying a symmetry pattern of the form [F(x),G(x)]
-> trying a symmetry pattern of the form [F(y),G(y)]
-> trying a symmetry pattern of the form [F(x)+G(y), 0]
-> trying a symmetry pattern of the form [0, F(x)+G(y)]
-> trying a symmetry pattern of the form [F(x),G(x)*y+H(x)]
-> trying a symmetry pattern of conformal type`
```

X Solution by Maple

```
dsolve((2*y(x)^2-4*x+5)=(4-2*y(x)+4*x*y(x))*diff(y(x),x),y(x), singsol=all)
```

No solution found

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[(2*y[x]^2-4*x+5)==(4-2*y[x]+4*x*y[x])*y'[x],y[x],x,IncludeSingularSolutions -> True]
```

Not solved

4.5 problem 5

4.5.1	Solving as separable ode	534
4.5.2	Solving as linear ode	536
4.5.3	Solving as homogeneousTypeD2 ode	537
4.5.4	Solving as differentialType ode	539
4.5.5	Solving as first order ode lie symmetry lookup ode	540
4.5.6	Solving as exact ode	544
4.5.7	Maple step by step solution	548

Internal problem ID [6186]

Internal file name [OUTPUT/5434_Sunday_June_05_2022_03_38_01_PM_47822472/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.5. Exact Equations. Page 20

Problem number: 5.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "linear", "separable", "differentialType", "homogeneousTypeD2", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

`[_separable]`

$$y + y \cos(xy) + (x + x \cos(xy)) y' = 0$$

4.5.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= f(x)g(y) \\ &= -\frac{y}{x} \end{aligned}$$

Where $f(x) = -\frac{1}{x}$ and $g(y) = y$. Integrating both sides gives

$$\begin{aligned}\frac{1}{y} dy &= -\frac{1}{x} dx \\ \int \frac{1}{y} dy &= \int -\frac{1}{x} dx \\ \ln(y) &= -\ln(x) + c_1 \\ y &= e^{-\ln(x)+c_1} \\ &= \frac{c_1}{x}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1}{x} \tag{1}$$




Figure 114: Slope field plot

Verification of solutions

$$y = \frac{c_1}{x}$$

Verified OK.

4.5.2 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = \frac{1}{x}$$
$$q(x) = 0$$

Hence the ode is

$$y' + \frac{y}{x} = 0$$

The integrating factor μ is

$$\mu = e^{\int \frac{1}{x} dx}$$
$$= x$$

The ode becomes

$$\frac{d}{dx} \mu y = 0$$
$$\frac{d}{dx} (xy) = 0$$

Integrating gives

$$xy = c_1$$

Dividing both sides by the integrating factor $\mu = x$ results in

$$y = \frac{c_1}{x}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1}{x} \tag{1}$$




Figure 115: Slope field plot

Verification of solutions

$$y = \frac{c_1}{x}$$

Verified OK.

4.5.3 Solving as homogeneousTypeD2 ode

Using the change of variables $y = u(x)x$ on the above ode results in new ode in $u(x)$

$$u(x)x + u(x)x \cos(x^2u(x)) + (x + x \cos(x^2u(x)))(u'(x)x + u(x)) = 0$$

In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{2u}{x} \end{aligned}$$

Where $f(x) = -\frac{2}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u} du &= -\frac{2}{x} dx \\ \int \frac{1}{u} du &= \int -\frac{2}{x} dx \\ \ln(u) &= -2 \ln(x) + c_2 \\ u &= e^{-2 \ln(x) + c_2} \\ &= \frac{c_2}{x^2}\end{aligned}$$

Therefore the solution y is

$$\begin{aligned}y &= ux \\ &= \frac{c_2}{x}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{c_2}{x} \tag{1}$$




Figure 116: Slope field plot

Verification of solutions

$$y = \frac{c_2}{x}$$

Verified OK.

4.5.4 Solving as differentialType ode

Writing the ode as

$$y' = \frac{-y - y \cos(xy)}{x + x \cos(xy)} \quad (1)$$

Which becomes

$$0 = (-x) dy + (-y) dx \quad (2)$$

But the RHS is complete differential because

$$(-x) dy + (-y) dx = d(-xy)$$

Hence (2) becomes

$$0 = d(-xy)$$

Integrating both sides gives gives these solutions

$$y = \frac{c_1}{x} + c_1$$

Summary

The solution(s) found are the following

$$y = \frac{c_1}{x} + c_1 \quad (1)$$




Figure 117: Slope field plot

Verification of solutions

$$y = \frac{c_1}{x} + c_2$$

Verified OK.

4.5.5 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = -\frac{y}{x}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 89: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= \frac{1}{x}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{1}{x}} dy \end{aligned}$$

Which results in

$$S = xy$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -\frac{y}{x}$$

Evaluating all the partial derivatives gives

$$R_x = 1$$

$$R_y = 0$$

$$S_x = y$$

$$S_y = x$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by

integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \tag{4}$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$xy = c_1$$



Which simplifies to

$$xy = c_1$$

Which gives

$$y = \frac{c_1}{x}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -\frac{y}{x}$ 	$R = x$ $S = xy$	$\frac{dS}{dR} = 0$ 

Summary

The solution(s) found are the following

$$y = \frac{c_1}{x} \tag{1}$$




Figure 118: Slope field plot

Verification of solutions

$$y = \frac{c_1}{x}$$

Verified OK.

4.5.6 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}\left(-\frac{1}{y}\right) dy &= \left(\frac{1}{x}\right) dx \\ \left(-\frac{1}{x}\right) dx + \left(-\frac{1}{y}\right) dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -\frac{1}{x} \\ N(x, y) &= -\frac{1}{y}\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(-\frac{1}{x}\right) \\ &= 0\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(-\frac{1}{y} \right) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \tag{1}$$

$$\frac{\partial \phi}{\partial y} = N \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\frac{1}{x} dx \\ \phi &= -\ln(x) + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = -\frac{1}{y}$. Therefore equation (4) becomes

$$-\frac{1}{y} = 0 + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = -\frac{1}{y}$$

Integrating the above w.r.t y gives

$$\begin{aligned}\int f'(y) dy &= \int \left(-\frac{1}{y} \right) dy \\ f(y) &= -\ln(y) + c_1\end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\ln(x) - \ln(y) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\ln(x) - \ln(y)$$

The solution becomes

$$y = \frac{e^{-c_1}}{x}$$

Summary

The solution(s) found are the following

$$y = \frac{e^{-c_1}}{x} \tag{1}$$




Figure 119: Slope field plot

Verification of solutions

$$y = \frac{e^{-c_1}}{x}$$

Verified OK.

4.5.7 Maple step by step solution

Let's solve

$$y + y \cos(xy) + (x + x \cos(xy)) y' = 0$$

- Highest derivative means the order of the ODE is 1

y'

- Integrate both sides with respect to x

$$\int (y + y \cos(xy) + (x + x \cos(xy)) y') dx = \int 0 dx + c_1$$

- Evaluate integral

$$xy + \sin(xy) = c_1$$

Maple trace

```
`Classification methods on request
Methods to be used are: [exact]
-----
* Tackling ODE using method: exact
--- Trying classification methods ---
trying exact
<- exact successful`
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 17

```
dsolve((y(x)+y(x)*cos(x*y(x)))+(x+x*cos(x*y(x)))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{\pi}{x}$$
$$y(x) = \frac{c_1}{x}$$

✓ Solution by Mathematica

Time used: 0.033 (sec). Leaf size: 49

```
DSolve[(y[x]+y[x]*Cos[x*y[x]])+(x+x*Cos[x*y[x]])*y'[x]==0,y[x],x,IncludeSingularSolutions ->
```

$$y(x) \rightarrow -\frac{\pi}{x}$$

$$y(x) \rightarrow \frac{\pi}{x}$$

$$y(x) \rightarrow \frac{C_1}{x}$$

$$y(x) \rightarrow -\frac{\pi}{x}$$

$$y(x) \rightarrow \frac{\pi}{x}$$

4.6 problem 6

4.6.1	Solving as separable ode	550
4.6.2	Solving as first order ode lie symmetry lookup ode	552
4.6.3	Solving as exact ode	556
4.6.4	Maple step by step solution	560

Internal problem ID [6187]

Internal file name [OUTPUT/5435_Sunday_June_05_2022_03_38_03_PM_36685332/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.5. Exact Equations. Page 20

Problem number: 6.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exact", "separable", "first_order_ode_lie_symmetry_lookup"**

Maple gives the following as the ode type

`[_separable]`

$$\cos(x) \cos(y)^2 + 2 \sin(x) \sin(y) \cos(y) y' = 0$$

4.6.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= f(x)g(y) \\ &= -\frac{\cos(x) \cot(y)}{2 \sin(x)} \end{aligned}$$

Where $f(x) = -\frac{\cos(x)}{2 \sin(x)}$ and $g(y) = \cot(y)$. Integrating both sides gives

$$\begin{aligned} \frac{1}{\cot(y)} dy &= -\frac{\cos(x)}{2 \sin(x)} dx \\ \int \frac{1}{\cot(y)} dy &= \int -\frac{\cos(x)}{2 \sin(x)} dx \end{aligned}$$

$$-\ln(\cos(y)) = -\frac{\ln(\sin(x))}{2} + c_1$$

Raising both side to exponential gives

$$\frac{1}{\cos(y)} = e^{-\frac{\ln(\sin(x))}{2} + c_1}$$

Which simplifies to

$$\sec(y) = \frac{c_2}{\sqrt{\sin(x)}}$$

Which simplifies to

$$y = \operatorname{arcsec}\left(\frac{c_2 e^{c_1}}{\sqrt{\sin(x)}}\right)$$

Summary

The solution(s) found are the following

$$y = \operatorname{arcsec}\left(\frac{c_2 e^{c_1}}{\sqrt{\sin(x)}}\right) \tag{1}$$




Figure 120: Slope field plot

Verification of solutions

$$y = \operatorname{arcsec} \left(\frac{c_2 e^{c_1}}{\sqrt{\sin(x)}} \right)$$

Verified OK.

4.6.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = -\frac{\cos(x) \cos(y)}{2 \sin(x) \sin(y)}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **separable**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 92: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= -\frac{2 \sin(x)}{\cos(x)} \\ \eta(x, y) &= 0\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\eta = 0$ then in this special case

$$R = y$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\xi} dx \\ &= \int \frac{1}{-\frac{2 \sin(x)}{\cos(x)}} dx \end{aligned}$$

Which results in

$$S = -\frac{\ln(\sin(x))}{2}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -\frac{\cos(x) \cos(y)}{2 \sin(x) \sin(y)}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 0 \\ R_y &= 1 \\ S_x &= -\frac{\cot(x)}{2} \\ S_y &= 0 \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \tan(y) \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \tan(R)$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = -\ln(\cos(R)) + c_1 \quad (4)$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$-\frac{\ln(\sin(x))}{2} = -\ln(\cos(y)) + c_1$$

Which simplifies to

$$-\frac{\ln(\sin(x))}{2} = -\ln(\cos(y)) + c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -\frac{\cos(x)\cos(y)}{2\sin(x)\sin(y)}$ 	$R = y$ $S = -\frac{\ln(\sin(x))}{2}$	$\frac{dS}{dR} = \tan(R)$ 

Summary

The solution(s) found are the following

$$-\frac{\ln(\sin(x))}{2} = -\ln(\cos(y)) + c_1 \quad (1)$$




Figure 121: Slope field plot

Verification of solutions

$$-\frac{\ln(\sin(x))}{2} = -\ln(\cos(y)) + c_1$$

Verified OK.

4.6.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial\phi}{\partial x} + \frac{\partial\phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}\left(-\frac{2 \sin (y)}{\cos (y)}\right) dy &= \left(\frac{\cos (x)}{\sin (x)}\right) dx \\ \left(-\frac{\cos (x)}{\sin (x)}\right) dx + \left(-\frac{2 \sin (y)}{\cos (y)}\right) dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -\frac{\cos (x)}{\sin (x)} \\ N(x, y) &= -\frac{2 \sin (y)}{\cos (y)}\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(-\frac{\cos (x)}{\sin (x)}\right) \\ &= 0\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(-\frac{2 \sin(y)}{\cos(y)} \right) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\frac{\cos(x)}{\sin(x)} dx \\ \phi &= -\ln(\sin(x)) + f(y)\end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = -\frac{2 \sin(y)}{\cos(y)}$. Therefore equation (4) becomes

$$-\frac{2 \sin(y)}{\cos(y)} = 0 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$\begin{aligned}f'(y) &= -\frac{2 \sin(y)}{\cos(y)} \\ &= -2 \tan(y)\end{aligned}$$

Integrating the above w.r.t y results in

$$\int f'(y) dy = \int (-2 \tan(y)) dy$$

$$f(y) = 2 \ln(\cos(y)) + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\ln(\sin(x)) + 2 \ln(\cos(y)) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\ln(\sin(x)) + 2 \ln(\cos(y))$$

Summary

The solution(s) found are the following

$$-\ln(\sin(x)) + 2 \ln(\cos(y)) = c_1 \tag{1}$$




Figure 122: Slope field plot

Verification of solutions

$$-\ln(\sin(x)) + 2\ln(\cos(y)) = c_1$$

Verified OK.

4.6.4 Maple step by step solution

Let's solve

$$\cos(x)\cos(y)^2 + 2\sin(x)\sin(y)\cos(y)y' = 0$$

- Highest derivative means the order of the ODE is 1

y'

- Separate variables

$$\frac{y' \sin(y)}{\cos(y)} = -\frac{\cos(x)}{2\sin(x)}$$

- Integrate both sides with respect to x

$$\int \frac{y' \sin(y)}{\cos(y)} dx = \int -\frac{\cos(x)}{2\sin(x)} dx + c_1$$

- Evaluate integral

$$-\ln(\cos(y)) = -\frac{\ln(\sin(x))}{2} + c_1$$

- Solve for y

$$\left\{ y = \pi - \arccos\left(\frac{\sqrt{e^{2c_1}\sin(x)}}{e^{2c_1}}\right), y = \arccos\left(\frac{\sqrt{e^{2c_1}\sin(x)}}{e^{2c_1}}\right) \right\}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
<- separable successful`
```

✓ Solution by Maple

Time used: 0.25 (sec). Leaf size: 31

```
dsolve((cos(x)*cos(y(x))^2)+(2*sin(x)*sin(y(x))*cos(y(x)))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{\pi}{2}$$

$$y(x) = \arccos\left(\sqrt{c_1 \sin(x)}\right)$$

$$y(x) = \frac{\pi}{2} + \arcsin\left(\sqrt{c_1 \sin(x)}\right)$$

✓ Solution by Mathematica

Time used: 5.453 (sec). Leaf size: 73

```
DSolve[(Cos[x]*Cos[y[x]]^2)+(2*Sin[x]*Sin[y[x]]*Cos[y[x]])*y'[x]==0,y[x],x,IncludeSingularSo
```

$$y(x) \rightarrow -\frac{\pi}{2}$$

$$y(x) \rightarrow \frac{\pi}{2}$$

$$y(x) \rightarrow -\arccos\left(-\frac{1}{4}c_1\sqrt{\sin(x)}\right)$$

$$y(x) \rightarrow \arccos\left(-\frac{1}{4}c_1\sqrt{\sin(x)}\right)$$

$$y(x) \rightarrow -\frac{\pi}{2}$$

$$y(x) \rightarrow \frac{\pi}{2}$$

4.7 problem 7

4.7.1 Solving as exact ode	562
4.7.2 Maple step by step solution	565

Internal problem ID [6188]

Internal file name [OUTPUT/5436_Sunday_June_05_2022_03_38_05_PM_34313907/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.5. Exact Equations. Page 20

Problem number: 7.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exact"**

Maple gives the following as the ode type

[_exact]

$$(\sin(x) \sin(y) - x e^y) y' - e^y - \cos(x) \cos(y) = 0$$

4.7.1 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned} (\sin(x) \sin(y) - x e^y) dy &= (e^y + \cos(x) \cos(y)) dx \\ (-\cos(x) \cos(y) - e^y) dx &+ (\sin(x) \sin(y) - x e^y) dy = 0 \end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -\cos(x) \cos(y) - e^y \\ N(x, y) &= \sin(x) \sin(y) - x e^y \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} (-\cos(x) \cos(y) - e^y) \\ &= \cos(x) \sin(y) - e^y \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} (\sin(x) \sin(y) - x e^y) \\ &= \cos(x) \sin(y) - e^y \end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\cos(x) \cos(y) - e^y dx \\ \phi &= -\sin(x) \cos(y) - x e^y + f(y)\end{aligned}\quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = \sin(x) \sin(y) - x e^y + f'(y)\quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \sin(x) \sin(y) - x e^y$. Therefore equation (4) becomes

$$\sin(x) \sin(y) - x e^y = \sin(x) \sin(y) - x e^y + f'(y)\quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = -\sin(x) \cos(y) - x e^y + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\sin(x) \cos(y) - x e^y$$

Summary

The solution(s) found are the following

$$-\sin(x) \cos(y) - x e^y = c_1\quad (1)$$




Figure 123: Slope field plot

Verification of solutions

$$-\sin(x) \cos(y) - x e^y = c_1$$

Verified OK.

4.7.2 Maple step by step solution

Let's solve

$$(\sin(x) \sin(y) - x e^y) y' - e^y - \cos(x) \cos(y) = 0$$

- Highest derivative means the order of the ODE is 1
 y'
- Check if ODE is exact
 - ODE is exact if the lhs is the total derivative of a C^2 function
 $F'(x, y) = 0$
 - Compute derivative of lhs
 $F'(x, y) + \left(\frac{\partial}{\partial y} F(x, y)\right) y' = 0$

- Evaluate derivatives
 $\cos(x) \sin(y) - e^y = \cos(x) \sin(y) - e^y$
- Condition met, ODE is exact
- Exact ODE implies solution will be of this form

$$\left[F(x, y) = c_1, M(x, y) = F'(x, y), N(x, y) = \frac{\partial}{\partial y} F(x, y) \right]$$
- Solve for $F(x, y)$ by integrating $M(x, y)$ with respect to x

$$F(x, y) = \int (-\cos(x) \cos(y) - e^y) dx + f_1(y)$$
- Evaluate integral

$$F(x, y) = -\sin(x) \cos(y) - x e^y + f_1(y)$$
- Take derivative of $F(x, y)$ with respect to y

$$N(x, y) = \frac{\partial}{\partial y} F(x, y)$$
- Compute derivative

$$\sin(x) \sin(y) - x e^y = \sin(x) \sin(y) - x e^y + \frac{d}{dy} f_1(y)$$
- Isolate for $\frac{d}{dy} f_1(y)$

$$\frac{d}{dy} f_1(y) = 0$$
- Solve for $f_1(y)$

$$f_1(y) = 0$$
- Substitute $f_1(y)$ into equation for $F(x, y)$

$$F(x, y) = -\sin(x) \cos(y) - x e^y$$
- Substitute $F(x, y)$ into the solution of the ODE

$$-\sin(x) \cos(y) - x e^y = c_1$$
- Solve for y

$$y = \text{RootOf}\left(-Z - \ln\left(-\frac{\sin(x) \cos(-Z) + c_1}{x}\right)\right)$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying Chini  
differential order: 1; looking for linear symmetries  
trying exact  
<- exact successful`
```

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 16

```
dsolve((sin(x)*sin(y(x))-x*exp(y(x)))*diff(y(x),x)=exp(y(x))+cos(x)*cos(y(x))),y(x), singsol=
```

$$c_1 + \sin(x) \cos(y(x)) + x e^{y(x)} = 0$$

✓ Solution by Mathematica

Time used: 0.594 (sec). Leaf size: 21

```
DSolve[(Sin[x]*Sin[y[x]]-x*Exp[y[x]])*y'[x]==Exp[y[x]]+Cos[x]*Cos[y[x]],y[x],x,IncludeSingul
```

$$\text{Solve}[2(xe^{y(x)} + \sin(x) \cos(y(x))) = c_1, y(x)]$$

4.8 problem 8

4.8.1 Solving as exact ode	568
4.8.2 Maple step by step solution	572

Internal problem ID [6189]

Internal file name [OUTPUT/5437_Sunday_June_05_2022_03_38_09_PM_22462015/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.5. Exact Equations. Page 20

Problem number: 8.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exact"**

Maple gives the following as the ode type

`[_separable]`

$$-\frac{\sin\left(\frac{x}{y}\right)}{y} + \frac{x \sin\left(\frac{x}{y}\right) y'}{y^2} = 0$$

4.8.1 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}\left(\frac{1}{y}\right) dy &= \left(\frac{1}{x}\right) dx \\ \left(-\frac{1}{x}\right) dx + \left(\frac{1}{y}\right) dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -\frac{1}{x} \\ N(x, y) &= \frac{1}{y}\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(-\frac{1}{x}\right) \\ &= 0\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{1}{y} \right) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\frac{1}{x} dx \\ \phi &= -\ln(x) + f(y)\end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{1}{y}$. Therefore equation (4) becomes

$$\frac{1}{y} = 0 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = \frac{1}{y}$$

Integrating the above w.r.t y gives

$$\begin{aligned}\int f'(y) dy &= \int \left(\frac{1}{y} \right) dy \\ f(y) &= \ln(y) + c_1\end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\ln(x) + \ln(y) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\ln(x) + \ln(y)$$

The solution becomes

$$y = x e^{c_1}$$

Summary

The solution(s) found are the following

$$y = x e^{c_1} \tag{1}$$




Figure 124: Slope field plot

Verification of solutions

$$y = x e^{c_1}$$

Verified OK.

4.8.2 Maple step by step solution

Let's solve

$$-\frac{\sin\left(\frac{x}{y}\right)}{y} + \frac{x \sin\left(\frac{x}{y}\right)y'}{y^2} = 0$$

- Highest derivative means the order of the ODE is 1

y'

- Integrate both sides with respect to x

$$\int \left(-\frac{\sin\left(\frac{x}{y}\right)}{y} + \frac{x \sin\left(\frac{x}{y}\right)y'}{y^2} \right) dx = \int 0 dx + c_1$$

- Evaluate integral

$$\cos\left(\frac{x}{y}\right) = c_1$$

- Solve for y

$$y = \frac{x}{\arccos(c_1)}$$

Maple trace

```
`Classification methods on request
```

```
Methods to be used are: [exact]
```

```
-----
```

```
* Tackling ODE using method: exact
```

```
--- Trying classification methods ---
```

```
trying exact
```

```
<- exact successful`
```

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 13

```
dsolve(-1/y(x)*sin(x/y(x))+(x/y(x)^2*sin(x/y(x)))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{x}{\pi - c_1}$$

✓ Solution by Mathematica

Time used: 0.032 (sec). Leaf size: 19

```
DSolve[-1/y[x]*Sin[x/y[x]]+(x/y[x]^2*Sin[x/y[x]])*y'[x]==0,y[x],x,IncludeSingularSolutions
```

$$y(x) \rightarrow c_1 x$$

$$y(x) \rightarrow \text{ComplexInfinity}$$

$$y(x) \rightarrow \text{ComplexInfinity}$$

4.9 problem 9

4.9.1	Solving as separable ode	574
4.9.2	Solving as linear ode	576
4.9.3	Solving as homogeneousTypeD2 ode	578
4.9.4	Solving as homogeneousTypeMapleC ode	579
4.9.5	Solving as first order ode lie symmetry lookup ode	582
4.9.6	Solving as exact ode	586
4.9.7	Maple step by step solution	590

Internal problem ID [6190]

Internal file name [OUTPUT/5438_Sunday_June_05_2022_03_38_11_PM_5955005/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.5. Exact Equations. Page 20

Problem number: 9.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "linear", "separable", "homogeneousTypeD2", "homogeneousTypeMapleC", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_separable]

$$y + (1 - x)y' = -1$$

4.9.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{1 + y}{x - 1}\end{aligned}$$

Where $f(x) = \frac{1}{x-1}$ and $g(y) = 1 + y$. Integrating both sides gives

$$\begin{aligned}\frac{1}{1+y} dy &= \frac{1}{x-1} dx \\ \int \frac{1}{1+y} dy &= \int \frac{1}{x-1} dx \\ \ln(1+y) &= \ln(x-1) + c_1\end{aligned}$$

Raising both side to exponential gives

$$1 + y = e^{\ln(x-1)+c_1}$$

Which simplifies to

$$1 + y = c_2(x - 1)$$

Summary

The solution(s) found are the following

$$y = c_2 e^{\ln(x-1)+c_1} - 1 \tag{1}$$




Figure 125: Slope field plot

Verification of solutions

$$y = c_2 e^{\ln(x-1)+c_1} - 1$$

Verified OK.

4.9.2 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -\frac{1}{x-1}$$
$$q(x) = \frac{1}{x-1}$$

Hence the ode is

$$y' - \frac{y}{x-1} = \frac{1}{x-1}$$

The integrating factor μ is

$$\mu = e^{\int -\frac{1}{x-1} dx}$$
$$= \frac{1}{x-1}$$

The ode becomes

$$\frac{d}{dx}(\mu y) = (\mu) \left(\frac{1}{x-1} \right)$$
$$\frac{d}{dx} \left(\frac{y}{x-1} \right) = \left(\frac{1}{x-1} \right) \left(\frac{1}{x-1} \right)$$
$$d \left(\frac{y}{x-1} \right) = \frac{1}{(x-1)^2} dx$$

Integrating gives

$$\frac{y}{x-1} = \int \frac{1}{(x-1)^2} dx$$
$$\frac{y}{x-1} = -\frac{1}{x-1} + c_1$$

Dividing both sides by the integrating factor $\mu = \frac{1}{x-1}$ results in

$$y = -1 + c_1(x - 1)$$

Summary

The solution(s) found are the following

$$y = -1 + c_1(x - 1) \tag{1}$$




Figure 126: Slope field plot

Verification of solutions

$$y = -1 + c_1(x - 1)$$

Verified OK.

4.9.3 Solving as homogeneous Type D2 ode

Using the change of variables $y = u(x)x$ on the above ode results in new ode in $u(x)$

$$u(x)x + (1-x)(u'(x)x + u(x)) = -1$$

In canonical form the ODE is

$$\begin{aligned}u' &= F(x, u) \\ &= f(x)g(u) \\ &= \frac{u+1}{x(x-1)}\end{aligned}$$

Where $f(x) = \frac{1}{x(x-1)}$ and $g(u) = u+1$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u+1} du &= \frac{1}{x(x-1)} dx \\ \int \frac{1}{u+1} du &= \int \frac{1}{x(x-1)} dx \\ \ln(u+1) &= \ln(x-1) - \ln(x) + c_2\end{aligned}$$

Raising both side to exponential gives

$$u+1 = e^{\ln(x-1) - \ln(x) + c_2}$$

Which simplifies to

$$u+1 = c_3 e^{\ln(x-1) - \ln(x)}$$

Which simplifies to

$$u(x) = c_3 \left(e^{c_2} - \frac{e^{c_2}}{x} \right) - 1$$

Therefore the solution y is

$$\begin{aligned}y &= ux \\ &= x \left(c_3 \left(e^{c_2} - \frac{e^{c_2}}{x} \right) - 1 \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = x \left(c_3 \left(e^{c_2} - \frac{e^{c_2}}{x} \right) - 1 \right) \quad (1)$$




Figure 127: Slope field plot

Verification of solutions

$$y = x \left(c_3 \left(e^{c_2} - \frac{e^{c_2}}{x} \right) - 1 \right)$$

Verified OK.

4.9.4 Solving as homogeneousTypeMapleC ode

Let $Y = y + y_0$ and $X = x + x_0$ then the above is transformed to new ode in $Y(X)$

$$\frac{d}{dX}Y(X) = \frac{1 + Y(X) + y_0}{X + x_0 - 1}$$

Solving for possible values of x_0 and y_0 which makes the above ode a homogeneous ode results in

$$\begin{aligned} x_0 &= 1 \\ y_0 &= -1 \end{aligned}$$

Using these values now it is possible to easily solve for $Y(X)$. The above ode now becomes

$$\frac{d}{dX}Y(X) = \frac{Y(X)}{X}$$

In canonical form, the ODE is

$$\begin{aligned} Y' &= F(X, Y) \\ &= \frac{Y}{X} \end{aligned} \tag{1}$$

An ode of the form $Y' = \frac{M(X, Y)}{N(X, Y)}$ is called homogeneous if the functions $M(X, Y)$ and $N(X, Y)$ are both homogeneous functions and of the same order. Recall that a function $f(X, Y)$ is homogeneous of order n if

$$f(t^n X, t^n Y) = t^n f(X, Y)$$

In this case, it can be seen that both $M = Y$ and $N = X$ are both homogeneous and of the same order $n = 1$. Therefore this is a homogeneous ode. Since this ode is homogeneous, it is converted to separable ODE using the substitution $u = \frac{Y}{X}$, or $Y = uX$. Hence

$$\frac{dY}{dX} = \frac{du}{dX}X + u$$

Applying the transformation $Y = uX$ to the above ODE in (1) gives

$$\begin{aligned} \frac{du}{dX}X + u &= u \\ \frac{du}{dX} &= 0 \end{aligned}$$

Or

$$\frac{d}{dX}u(X) = 0$$

Which is now solved as separable in $u(X)$. Which is now solved in $u(X)$. Integrating both sides gives

$$\begin{aligned} u(X) &= \int 0 \, dX \\ &= c_2 \end{aligned}$$

Now u in the above solution is replaced back by Y using $u = \frac{Y}{X}$ which results in the solution

$$Y(X) = Xc_2$$

Using the solution for $Y(X)$

$$Y(X) = Xc_2$$

And replacing back terms in the above solution using

$$Y = y + y_0$$

$$X = x + x_0$$

Or

$$Y = y - 1$$

$$X = 1 + x$$

Then the solution in y becomes

$$1 + y = c_2(x - 1)$$

Summary

The solution(s) found are the following

$$1 + y = c_2(x - 1) \tag{1}$$




Figure 128: Slope field plot

Verification of solutions

$$1 + y = c_2(x - 1)$$

Verified OK.

4.9.5 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{1+y}{x-1}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 97: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= x - 1\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{x-1} dy\end{aligned}$$

Which results in

$$S = \frac{y}{x-1}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y}\tag{2}$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{1+y}{x-1}$$

Evaluating all the partial derivatives gives

$$\begin{aligned}R_x &= 1 \\R_y &= 0 \\S_x &= -\frac{y}{(x-1)^2} \\S_y &= \frac{1}{x-1}\end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \frac{1}{(x-1)^2} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \frac{1}{(R-1)^2}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = -\frac{1}{R-1} + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{y}{x-1} = -\frac{1}{x-1} + c_1$$



Which simplifies to

$$\frac{-c_1x + c_1 + y + 1}{x-1} = 0$$

Which gives

$$y = c_1x - c_1 - 1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{1+y}{x-1}$ 	$R = x$ $S = \frac{y}{x-1}$	$\frac{dS}{dR} = \frac{1}{(R-1)^2}$ 

Summary

The solution(s) found are the following

$$y = c_1 x - c_1 - 1 \tag{1}$$




Figure 129: Slope field plot

Verification of solutions

$$y = c_1x - c_1 - 1$$

Verified OK.

4.9.6 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial\phi}{\partial x} + \frac{\partial\phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned}\left(\frac{1}{1+y}\right) dy &= \left(\frac{1}{x-1}\right) dx \\ \left(-\frac{1}{x-1}\right) dx + \left(\frac{1}{1+y}\right) dy &= 0\end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -\frac{1}{x-1} \\ N(x, y) &= \frac{1}{1+y}\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(-\frac{1}{x-1}\right) \\ &= 0\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{1}{1+y} \right) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\frac{1}{x-1} dx \\ \phi &= -\ln(x-1) + f(y)\end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{1}{1+y}$. Therefore equation (4) becomes

$$\frac{1}{1+y} = 0 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = \frac{1}{1+y}$$

Integrating the above w.r.t y gives

$$\begin{aligned}\int f'(y) dy &= \int \left(\frac{1}{1+y} \right) dy \\ f(y) &= \ln(1+y) + c_1\end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\ln(x-1) + \ln(1+y) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\ln(x-1) + \ln(1+y)$$

The solution becomes

$$y = x e^{c_1} - e^{c_1} - 1$$

Summary

The solution(s) found are the following

$$y = x e^{c_1} - e^{c_1} - 1 \tag{1}$$




Figure 130: Slope field plot

Verification of solutions

$$y = x e^{c_1} - e^{c_1} - 1$$

Verified OK.

4.9.7 Maple step by step solution

Let's solve

$$y + (1 - x)y' = -1$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{-1-y} = \frac{1}{1-x}$$

- Integrate both sides with respect to x

$$\int \frac{y'}{-1-y} dx = \int \frac{1}{1-x} dx + c_1$$

- Evaluate integral

$$-\ln(-1 - y) = -\ln(1 - x) + c_1$$

- Solve for y

$$y = -\frac{e^{c_1-x+1}}{e^{c_1}}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve((1+y(x))+(1-x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = -1 + c_1(x - 1)$$

✓ Solution by Mathematica

Time used: 0.029 (sec). Leaf size: 18

```
DSolve[(1+y[x])+(1-x)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -1 + c_1(x - 1)$$

$$y(x) \rightarrow -1$$

4.10 problem 10

4.10.1 Solving as exact ode	592
4.10.2 Maple step by step solution	595

Internal problem ID [6191]

Internal file name [OUTPUT/5439_Sunday_June_05_2022_03_38_12_PM_82103197/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.5. Exact Equations. Page 20

Problem number: 10.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exact"**

Maple gives the following as the ode type

```
[_exact , [_1st_order , ` _with_symmetry_ [F(x) , G(x)*y+H(x)] `]]
```

$$2y^3x + y \cos(x) + (3y^2x^2 + \sin(x)) y' = 0$$

4.10.1 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned} (3y^2x^2 + \sin(x)) dy &= (-2xy^3 - y \cos(x)) dx \\ (2xy^3 + y \cos(x)) dx + (3y^2x^2 + \sin(x)) dy &= 0 \end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= 2xy^3 + y \cos(x) \\ N(x, y) &= 3y^2x^2 + \sin(x) \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} (2xy^3 + y \cos(x)) \\ &= 6y^2x + \cos(x) \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} (3y^2x^2 + \sin(x)) \\ &= 6y^2x + \cos(x) \end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int 2x y^3 + y \cos(x) dx \\ \phi &= y(y^2 x^2 + \sin(x)) + f(y)\end{aligned}\quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 3y^2 x^2 + \sin(x) + f'(y)\quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = 3y^2 x^2 + \sin(x)$. Therefore equation (4) becomes

$$3y^2 x^2 + \sin(x) = 3y^2 x^2 + \sin(x) + f'(y)\quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = y(y^2 x^2 + \sin(x)) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = y(y^2 x^2 + \sin(x))$$

Summary

The solution(s) found are the following

$$y(y^2 x^2 + \sin(x)) = c_1\quad (1)$$




Figure 131: Slope field plot

Verification of solutions

$$y(y^2 x^2 + \sin(x)) = c_1$$

Verified OK.

4.10.2 Maple step by step solution

Let's solve

$$2y^3 x + y \cos(x) + (3y^2 x^2 + \sin(x)) y' = 0$$

- Highest derivative means the order of the ODE is 1
 y'
- Check if ODE is exact
 - ODE is exact if the lhs is the total derivative of a C^2 function
 $F'(x, y) = 0$
 - Compute derivative of lhs
 $F'(x, y) + \left(\frac{\partial}{\partial y} F(x, y)\right) y' = 0$

- Evaluate derivatives

$$6y^2x + \cos(x) = 6y^2x + \cos(x)$$

- Condition met, ODE is exact

- Exact ODE implies solution will be of this form

$$\left[F(x, y) = c_1, M(x, y) = F'(x, y), N(x, y) = \frac{\partial}{\partial y} F(x, y) \right]$$

- Solve for $F(x, y)$ by integrating $M(x, y)$ with respect to x

$$F(x, y) = \int (2xy^3 + y \cos(x)) dx + f_1(y)$$

- Evaluate integral

$$F(x, y) = y \sin(x) + y^3x^2 + f_1(y)$$

- Take derivative of $F(x, y)$ with respect to y

$$N(x, y) = \frac{\partial}{\partial y} F(x, y)$$

- Compute derivative

$$3y^2x^2 + \sin(x) = \sin(x) + 3y^2x^2 + \frac{d}{dy} f_1(y)$$

- Isolate for $\frac{d}{dy} f_1(y)$

$$\frac{d}{dy} f_1(y) = 0$$

- Solve for $f_1(y)$

$$f_1(y) = 0$$

- Substitute $f_1(y)$ into equation for $F(x, y)$

$$F(x, y) = y \sin(x) + y^3x^2$$

- Substitute $F(x, y)$ into the solution of the ODE

$$y \sin(x) + y^3x^2 = c_1$$

- Solve for y

$$\left\{ y = \frac{\left(12\sqrt{3} \sqrt{27x^2c_1^2 + 4 \sin(x)^3} + 108c_1x\right)^{\frac{1}{3}}}{6x} - \frac{2 \sin(x)}{x \left(12\sqrt{3} \sqrt{27x^2c_1^2 + 4 \sin(x)^3} + 108c_1x\right)^{\frac{1}{3}}}, y = -\frac{\left(12\sqrt{3} \sqrt{27x^2c_1^2 + 4 \sin(x)^3}\right)^{\frac{1}{3}}}{12x} \right.$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying Chini  
differential order: 1; looking for linear symmetries  
trying exact  
<- exact successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 302

```
dsolve((2*x*y(x)^3+y(x)*cos(x))+(3*x^2*y(x)^2+sin(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{\left(12\sqrt{3}\sqrt{27x^2c_1^2 + 4\sin(x)^3} - 108c_1x\right)^{\frac{2}{3}} - 12\sin(x)}{6x\left(12\sqrt{3}\sqrt{27x^2c_1^2 + 4\sin(x)^3} - 108c_1x\right)^{\frac{1}{3}}}$$

$$y(x) = \frac{i\sqrt{3}\left(12\sqrt{3}\sqrt{27x^2c_1^2 + 4\sin(x)^3} - 108c_1x\right)^{\frac{2}{3}} + 12i\sin(x)\sqrt{3} + \left(12\sqrt{3}\sqrt{27x^2c_1^2 + 4\sin(x)^3} - 108c_1x\right)^{\frac{1}{3}}}{12x\left(12\sqrt{3}\sqrt{27x^2c_1^2 + 4\sin(x)^3} - 108c_1x\right)^{\frac{1}{3}}}$$

$$y(x) = \frac{i\sqrt{3}\left(12\sqrt{3}\sqrt{27x^2c_1^2 + 4\sin(x)^3} - 108c_1x\right)^{\frac{2}{3}} + 12i\sin(x)\sqrt{3} - \left(12\sqrt{3}\sqrt{27x^2c_1^2 + 4\sin(x)^3} - 108c_1x\right)^{\frac{1}{3}}}{12x\left(12\sqrt{3}\sqrt{27x^2c_1^2 + 4\sin(x)^3} - 108c_1x\right)^{\frac{1}{3}}}$$

✓ Solution by Mathematica

Time used: 34.571 (sec). Leaf size: 339

`DSolve[(2*x*y[x]^3+y[x]*Cos[x])+(3*x^2*y[x]^2+Sin[x])*y'[x]==0,y[x],x,IncludeSingularSolutio`

$$y(x) \rightarrow \frac{\sqrt[3]{9c_1x^4 + \sqrt{12x^6 \sin^3(x) + 81c_1^2x^8}}}{\sqrt[3]{23^{2/3}x^2}} - \frac{\sqrt[3]{\frac{2}{3}} \sin(x)}{\sqrt[3]{9c_1x^4 + \sqrt{12x^6 \sin^3(x) + 81c_1^2x^8}}}$$

$$y(x) \rightarrow \frac{(1 + i\sqrt{3}) \sin(x)}{2^{2/3} \sqrt[3]{27c_1x^4 + 3\sqrt{12x^6 \sin^3(x) + 81c_1^2x^8}}}$$

$$- \frac{(1 - i\sqrt{3}) \sqrt[3]{27c_1x^4 + \sqrt{108x^6 \sin^3(x) + 729c_1^2x^8}}}{6\sqrt[3]{2}x^2}$$

$$y(x) \rightarrow \frac{(1 - i\sqrt{3}) \sin(x)}{2^{2/3} \sqrt[3]{27c_1x^4 + 3\sqrt{12x^6 \sin^3(x) + 81c_1^2x^8}}}$$

$$- \frac{(1 + i\sqrt{3}) \sqrt[3]{27c_1x^4 + \sqrt{108x^6 \sin^3(x) + 729c_1^2x^8}}}{6\sqrt[3]{2}x^2}$$

4.11 problem 11

4.11.1 Solving as exact ode	599
4.11.2 Solving as riccati ode	603
4.11.3 Maple step by step solution	605

Internal problem ID [6192]

Internal file name [OUTPUT/5440_Sunday_June_05_2022_03_38_16_PM_44427014/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.5. Exact Equations. Page 20

Problem number: 11.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "riccati"

Maple gives the following as the ode type

[_exact, _rational, _Riccati]

$$\frac{y}{1-y^2x^2} + \frac{xy'}{1-y^2x^2} = 1$$

4.11.1 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned}\left(\frac{x}{-y^2x^2 + 1}\right) dy &= \left(-\frac{y}{-y^2x^2 + 1} + 1\right) dx \\ \left(\frac{y}{-y^2x^2 + 1} - 1\right) dx &+ \left(\frac{x}{-y^2x^2 + 1}\right) dy = 0\end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= \frac{y}{-y^2x^2 + 1} - 1 \\ N(x, y) &= \frac{x}{-y^2x^2 + 1}\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(\frac{y}{-y^2x^2 + 1} - 1 \right) \\ &= \frac{y^2x^2 + 1}{(y^2x^2 - 1)^2}\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{x}{-y^2x^2 + 1} \right) \\ &= \frac{y^2x^2 + 1}{(y^2x^2 - 1)^2}\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int \frac{y}{-y^2x^2 + 1} - 1 dx \\ \phi &= -x + \frac{\ln(xy + 1)}{2} - \frac{\ln(xy - 1)}{2} + f(y)\end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\begin{aligned}\frac{\partial \phi}{\partial y} &= \frac{x}{2xy + 2} - \frac{x}{2(xy - 1)} + f'(y) \\ &= -\frac{x}{y^2x^2 - 1} + f'(y)\end{aligned} \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{x}{-y^2x^2 + 1}$. Therefore equation (4) becomes

$$\frac{x}{-y^2x^2 + 1} = -\frac{x}{y^2x^2 - 1} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = -x + \frac{\ln(xy + 1)}{2} - \frac{\ln(xy - 1)}{2} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -x + \frac{\ln(xy + 1)}{2} - \frac{\ln(xy - 1)}{2}$$

The solution becomes

$$y = -\frac{e^{-2c_1-2x} + 1}{x(e^{-2c_1-2x} - 1)}$$

Summary

The solution(s) found are the following

$$y = -\frac{e^{-2c_1-2x} + 1}{x(e^{-2c_1-2x} - 1)} \quad (1)$$




Figure 132: Slope field plot

Verification of solutions

$$y = -\frac{e^{-2c_1-2x} + 1}{x(e^{-2c_1-2x} - 1)}$$

Verified OK.

4.11.2 Solving as riccati ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= -\frac{y^2 x^2 + y - 1}{x}\end{aligned}$$

This is a Riccati ODE. Comparing the ODE to solve

$$y' = -y^2 x - \frac{y}{x} + \frac{1}{x}$$

With Riccati ODE standard form

$$y' = f_0(x) + f_1(x)y + f_2(x)y^2$$

Shows that $f_0(x) = \frac{1}{x}$, $f_1(x) = -\frac{1}{x}$ and $f_2(x) = -x$. Let

$$\begin{aligned}y &= \frac{-u'}{f_2 u} \\ &= \frac{-u'}{-x u}\end{aligned}\tag{1}$$

Using the above substitution in the given ODE results (after some simplification) in a second order ODE to solve for $u(x)$ which is

$$f_2 u''(x) - (f_2' + f_1 f_2) u'(x) + f_2^2 f_0 u(x) = 0\tag{2}$$

But

$$\begin{aligned}f_2' &= -1 \\ f_1 f_2 &= 1 \\ f_2^2 f_0 &= x\end{aligned}$$

Substituting the above terms back in equation (2) gives

$$-x u''(x) + x u(x) = 0$$

Solving the above ODE (this ode solved using Maple, not this program), gives

$$u(x) = c_1 e^x + c_2 e^{-x}$$

The above shows that

$$u'(x) = c_1 e^x - c_2 e^{-x}$$

Using the above in (1) gives the solution

$$y = \frac{c_1 e^x - c_2 e^{-x}}{x (c_1 e^x + c_2 e^{-x})}$$

Dividing both numerator and denominator by c_1 gives, after renaming the constant $\frac{c_2}{c_1} = c_3$ the following solution

$$y = \frac{-1 + e^{2x} c_3}{x (e^{2x} c_3 + 1)}$$

Summary

The solution(s) found are the following

$$y = \frac{-1 + e^{2x} c_3}{x (e^{2x} c_3 + 1)} \tag{1}$$




Figure 133: Slope field plot

Verification of solutions

$$y = \frac{-1 + e^{2x} c_3}{x (e^{2x} c_3 + 1)}$$

Verified OK.

4.11.3 Maple step by step solution

Let's solve

$$\frac{y}{1-y^2x^2} + \frac{xy'}{1-y^2x^2} = 1$$

- Highest derivative means the order of the ODE is 1
- Check if ODE is exact
 - ODE is exact if the lhs is the total derivative of a C^2 function
 - Compute derivative of lhs

$$F'(x, y) + \left(\frac{\partial}{\partial y} F(x, y) \right) y' = 0$$

- Evaluate derivatives

$$\frac{1}{-y^2x^2+1} + \frac{2y^2x^2}{(-y^2x^2+1)^2} = \frac{1}{-y^2x^2+1} + \frac{2y^2x^2}{(-y^2x^2+1)^2}$$

- Condition met, ODE is exact

- Exact ODE implies solution will be of this form

$$\left[F(x, y) = c_1, M(x, y) = F'(x, y), N(x, y) = \frac{\partial}{\partial y} F(x, y) \right]$$

- Solve for $F(x, y)$ by integrating $M(x, y)$ with respect to x

$$F(x, y) = \int \left(\frac{y}{-y^2x^2+1} - 1 \right) dx + f_1(y)$$

- Evaluate integral

$$F(x, y) = -x + \frac{\ln(xy+1)}{2} - \frac{\ln(xy-1)}{2} + f_1(y)$$

- Take derivative of $F(x, y)$ with respect to y

$$N(x, y) = \frac{\partial}{\partial y} F(x, y)$$

- Compute derivative

$$\frac{x}{-y^2x^2+1} = \frac{x}{2(xy+1)} - \frac{x}{2(xy-1)} + \frac{d}{dy} f_1(y)$$

- Isolate for $\frac{d}{dy} f_1(y)$

$$\frac{d}{dy} f_1(y) = \frac{x}{-y^2x^2+1} - \frac{x}{2(xy+1)} + \frac{x}{2(xy-1)}$$

- Solve for $f_1(y)$

$$f_1(y) = 0$$

- Substitute $f_1(y)$ into equation for $F(x, y)$

$$F(x, y) = -x + \frac{\ln(xy+1)}{2} - \frac{\ln(xy-1)}{2}$$

- Substitute $F(x, y)$ into the solution of the ODE

$$-x + \frac{\ln(xy+1)}{2} - \frac{\ln(xy-1)}{2} = c_1$$

- Solve for y

$$y = -\frac{e^{-2c_1-2x}+1}{x(e^{-2c_1-2x}-1)}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying Chini  
differential order: 1; looking for linear symmetries  
trying exact  
<- exact successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
dsolve(y(x)/(1-x^2*y(x)^2)+x/(1-x^2*y(x)^2)*diff(y(x),x)=1,y(x), singsol=all)
```

$$y(x) = \frac{e^{2x} + c_1}{x(e^{2x} - c_1)}$$

✓ Solution by Mathematica

Time used: 0.16 (sec). Leaf size: 18

```
DSolve[y[x]/(1-x^2*y[x]^2)+x/(1-x^2*y[x]^2)*y'[x]==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\tanh(x + ic_1)}{x}$$

4.12 problem 12

4.12.1 Solving as exact ode	608
4.12.2 Maple step by step solution	611

Internal problem ID [6193]

Internal file name [OUTPUT/5441_Sunday_June_05_2022_03_38_19_PM_43906110/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.5. Exact Equations. Page 20

Problem number: 12.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exact"**

Maple gives the following as the ode type

[_exact]

$$2xy^4 + \sin(y) + (4y^3x^2 + \cos(y)x)y' = 0$$

4.12.1 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned} (4y^3 x^2 + \cos(y) x) dy &= (-2y^4 x - \sin(y)) dx \\ (2y^4 x + \sin(y)) dx + (4y^3 x^2 + \cos(y) x) dy &= 0 \end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= 2y^4 x + \sin(y) \\ N(x, y) &= 4y^3 x^2 + \cos(y) x \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} (2y^4 x + \sin(y)) \\ &= 8x y^3 + \cos(y) \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} (4y^3 x^2 + \cos(y) x) \\ &= 8x y^3 + \cos(y) \end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \tag{1}$$

$$\frac{\partial \phi}{\partial y} = N \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int 2y^4 x + \sin(y) dx \\ \phi &= x(y^4 x + \sin(y)) + f(y)\end{aligned}\quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = x(4y^3 + \cos(y)) + f'(y)\quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = 4y^3 x^2 + \cos(y) x$. Therefore equation (4) becomes

$$4y^3 x^2 + \cos(y) x = 4y^3 x^2 + \cos(y) x + f'(y)\quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = x(y^4 x + \sin(y)) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = x(y^4 x + \sin(y))$$

Summary

The solution(s) found are the following

$$x(xy^4 + \sin(y)) = c_1\quad (1)$$




Figure 134: Slope field plot

Verification of solutions

$$x(xy^4 + \sin(y)) = c_1$$

Verified OK.

4.12.2 Maple step by step solution

Let's solve

$$2xy^4 + \sin(y) + (4y^3x^2 + \cos(y)x)y' = 0$$

- Highest derivative means the order of the ODE is 1
 y'
- Check if ODE is exact
 - ODE is exact if the lhs is the total derivative of a C^2 function
 $F'(x, y) = 0$
 - Compute derivative of lhs
 $F'(x, y) + \left(\frac{\partial}{\partial y}F(x, y)\right)y' = 0$

- Evaluate derivatives

$$8x y^3 + \cos(y) = 8x y^3 + \cos(y)$$
- Condition met, ODE is exact
- Exact ODE implies solution will be of this form

$$\left[F(x, y) = c_1, M(x, y) = F'(x, y), N(x, y) = \frac{\partial}{\partial y} F(x, y) \right]$$
- Solve for $F(x, y)$ by integrating $M(x, y)$ with respect to x

$$F(x, y) = \int (2y^4 x + \sin(y)) dx + f_1(y)$$
- Evaluate integral

$$F(x, y) = y^4 x^2 + \sin(y) x + f_1(y)$$
- Take derivative of $F(x, y)$ with respect to y

$$N(x, y) = \frac{\partial}{\partial y} F(x, y)$$
- Compute derivative

$$4y^3 x^2 + \cos(y) x = 4y^3 x^2 + \cos(y) x + \frac{d}{dy} f_1(y)$$
- Isolate for $\frac{d}{dy} f_1(y)$

$$\frac{d}{dy} f_1(y) = 0$$
- Solve for $f_1(y)$

$$f_1(y) = 0$$
- Substitute $f_1(y)$ into equation for $F(x, y)$

$$F(x, y) = y^4 x^2 + \sin(y) x$$
- Substitute $F(x, y)$ into the solution of the ODE

$$y^4 x^2 + \sin(y) x = c_1$$
- Solve for y

$$y = \text{RootOf}(-x^2_Z^4 - \sin(_Z) x + c_1)$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying Chini  
differential order: 1; looking for linear symmetries  
trying exact  
<- exact successful`
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 18

```
dsolve((2*x*y(x)^4+sin(y(x)))+(4*x^2*y(x)^3+x*cos(y(x)))*diff(y(x),x)=0,y(x), singsol=all)
```

$$x^2y(x)^4 + x \sin(y(x)) + c_1 = 0$$

✓ Solution by Mathematica

Time used: 0.227 (sec). Leaf size: 20

```
DSolve[(2*x*y[x]^4+Sin[y[x]])+(4*x^2*y[x]^3+x*Cos[y[x]])*y'[x]==0,y[x],x,IncludeSingularSolu
```

$$\text{Solve}[x^2y(x)^4 + x \sin(y(x)) = c_1, y(x)]$$

4.13 problem 13

4.13.1 Solving as exact ode	614
4.13.2 Solving as riccati ode	618
4.13.3 Maple step by step solution	620

Internal problem ID [6194]

Internal file name [OUTPUT/5442_Sunday_June_05_2022_03_38_24_PM_56095593/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.5. Exact Equations. Page 20

Problem number: 13.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "riccati"

Maple gives the following as the ode type

[_exact, _rational, _Riccati]

$$\frac{xy' + y}{1 - y^2x^2} = -x$$

4.13.1 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned}\left(\frac{x}{-y^2x^2 + 1}\right) dy &= \left(-\frac{y}{-y^2x^2 + 1} - x\right) dx \\ \left(\frac{y}{-y^2x^2 + 1} + x\right) dx &+ \left(\frac{x}{-y^2x^2 + 1}\right) dy = 0\end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= \frac{y}{-y^2x^2 + 1} + x \\ N(x, y) &= \frac{x}{-y^2x^2 + 1}\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(\frac{y}{-y^2x^2 + 1} + x\right) \\ &= \frac{y^2x^2 + 1}{(y^2x^2 - 1)^2}\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{x}{-y^2x^2 + 1} \right) \\ &= \frac{y^2x^2 + 1}{(y^2x^2 - 1)^2}\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int \frac{y}{-y^2x^2 + 1} + x dx \\ \phi &= \frac{x^2}{2} + \frac{\ln(xy + 1)}{2} - \frac{\ln(xy - 1)}{2} + f(y)\end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\begin{aligned}\frac{\partial \phi}{\partial y} &= \frac{x}{2xy + 2} - \frac{x}{2(xy - 1)} + f'(y) \\ &= -\frac{x}{y^2x^2 - 1} + f'(y)\end{aligned} \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{x}{-y^2x^2 + 1}$. Therefore equation (4) becomes

$$\frac{x}{-y^2x^2 + 1} = -\frac{x}{y^2x^2 - 1} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = \frac{x^2}{2} + \frac{\ln(xy + 1)}{2} - \frac{\ln(xy - 1)}{2} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \frac{x^2}{2} + \frac{\ln(xy + 1)}{2} - \frac{\ln(xy - 1)}{2}$$

The solution becomes

$$y = -\frac{e^{x^2-2c_1} + 1}{x(e^{x^2-2c_1} - 1)}$$

Summary

The solution(s) found are the following

$$y = -\frac{e^{x^2-2c_1} + 1}{x(e^{x^2-2c_1} - 1)} \quad (1)$$




Figure 135: Slope field plot

Verification of solutions

$$y = -\frac{e^{x^2-2c_1} + 1}{x(e^{x^2-2c_1} - 1)}$$

Verified OK.

4.13.2 Solving as riccati ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= \frac{y^2 x^3 - x - y}{x}\end{aligned}$$

This is a Riccati ODE. Comparing the ODE to solve

$$y' = y^2 x^2 - 1 - \frac{y}{x}$$

With Riccati ODE standard form

$$y' = f_0(x) + f_1(x)y + f_2(x)y^2$$

Shows that $f_0(x) = -1$, $f_1(x) = -\frac{1}{x}$ and $f_2(x) = x^2$. Let

$$\begin{aligned}y &= \frac{-u'}{f_2 u} \\ &= \frac{-u'}{x^2 u}\end{aligned}\tag{1}$$

Using the above substitution in the given ODE results (after some simplification) in a second order ODE to solve for $u(x)$ which is

$$f_2 u''(x) - (f_2' + f_1 f_2) u'(x) + f_2^2 f_0 u(x) = 0\tag{2}$$

But

$$\begin{aligned}f_2' &= 2x \\ f_1 f_2 &= -x \\ f_2^2 f_0 &= -x^4\end{aligned}$$

Substituting the above terms back in equation (2) gives

$$x^2 u''(x) - x u'(x) - x^4 u(x) = 0$$

Solving the above ODE (this ode solved using Maple, not this program), gives

$$u(x) = c_1 \sinh\left(\frac{x^2}{2}\right) + c_2 \cosh\left(\frac{x^2}{2}\right)$$

The above shows that

$$u'(x) = x\left(c_1 \cosh\left(\frac{x^2}{2}\right) + c_2 \sinh\left(\frac{x^2}{2}\right)\right)$$

Using the above in (1) gives the solution

$$y = -\frac{c_1 \cosh\left(\frac{x^2}{2}\right) + c_2 \sinh\left(\frac{x^2}{2}\right)}{x\left(c_1 \sinh\left(\frac{x^2}{2}\right) + c_2 \cosh\left(\frac{x^2}{2}\right)\right)}$$

Dividing both numerator and denominator by c_1 gives, after renaming the constant $\frac{c_2}{c_1} = c_3$ the following solution

$$y = \frac{-c_3 \cosh\left(\frac{x^2}{2}\right) - \sinh\left(\frac{x^2}{2}\right)}{x\left(c_3 \sinh\left(\frac{x^2}{2}\right) + \cosh\left(\frac{x^2}{2}\right)\right)}$$

Summary

The solution(s) found are the following

$$y = \frac{-c_3 \cosh\left(\frac{x^2}{2}\right) - \sinh\left(\frac{x^2}{2}\right)}{x\left(c_3 \sinh\left(\frac{x^2}{2}\right) + \cosh\left(\frac{x^2}{2}\right)\right)} \quad (1)$$




Figure 136: Slope field plot

Verification of solutions

$$y = \frac{-c_3 \cosh\left(\frac{x^2}{2}\right) - \sinh\left(\frac{x^2}{2}\right)}{x \left(c_3 \sinh\left(\frac{x^2}{2}\right) + \cosh\left(\frac{x^2}{2}\right)\right)}$$

Verified OK.

4.13.3 Maple step by step solution

Let's solve

$$\frac{xy' + y}{1 - y^2 x^2} = -x$$

- Highest derivative means the order of the ODE is 1
- Check if ODE is exact
 - ODE is exact if the lhs is the total derivative of a C^2 function
 - Compute derivative of lhs

$$F'(x, y) + \left(\frac{\partial}{\partial y} F(x, y) \right) y' = 0$$

- Evaluate derivatives

$$\frac{1}{-y^2x^2+1} + \frac{2y^2x^2}{(-y^2x^2+1)^2} = \frac{1}{-y^2x^2+1} + \frac{2y^2x^2}{(-y^2x^2+1)^2}$$

- Condition met, ODE is exact

- Exact ODE implies solution will be of this form

$$\left[F(x, y) = c_1, M(x, y) = F'(x, y), N(x, y) = \frac{\partial}{\partial y} F(x, y) \right]$$

- Solve for $F(x, y)$ by integrating $M(x, y)$ with respect to x

$$F(x, y) = \int \left(\frac{y}{-y^2x^2+1} + x \right) dx + f_1(y)$$

- Evaluate integral

$$F(x, y) = \frac{x^2}{2} + \frac{\ln(xy+1)}{2} - \frac{\ln(xy-1)}{2} + f_1(y)$$

- Take derivative of $F(x, y)$ with respect to y

$$N(x, y) = \frac{\partial}{\partial y} F(x, y)$$

- Compute derivative

$$\frac{x}{-y^2x^2+1} = \frac{x}{2(xy+1)} - \frac{x}{2(xy-1)} + \frac{d}{dy} f_1(y)$$

- Isolate for $\frac{d}{dy} f_1(y)$

$$\frac{d}{dy} f_1(y) = \frac{x}{-y^2x^2+1} - \frac{x}{2(xy+1)} + \frac{x}{2(xy-1)}$$

- Solve for $f_1(y)$

$$f_1(y) = 0$$

- Substitute $f_1(y)$ into equation for $F(x, y)$

$$F(x, y) = \frac{x^2}{2} + \frac{\ln(xy+1)}{2} - \frac{\ln(xy-1)}{2}$$

- Substitute $F(x, y)$ into the solution of the ODE

$$\frac{x^2}{2} + \frac{\ln(xy+1)}{2} - \frac{\ln(xy-1)}{2} = c_1$$

- Solve for y

$$y = -\frac{e^{x^2-2c_1}+1}{x(e^{x^2-2c_1}-1)}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying Chini  
<- Chini successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 19

```
dsolve((y(x)+x*diff(y(x),x))/(1-x^2*y(x)^2)+x=0,y(x), singsol=all)
```

$$y(x) = \frac{i \tan\left(\frac{ix^2}{2} + c_1\right)}{x}$$

✓ Solution by Mathematica

Time used: 0.142 (sec). Leaf size: 25

```
DSolve[(y[x]+x*y'[x])/(1-x^2*y[x]^2)+x==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\tanh\left(\frac{1}{2}(x^2 - 2ic_1)\right)}{x}$$

4.14 problem 14

4.14.1 Solving as first order ode lie symmetry calculated ode	623
4.14.2 Solving as exact ode	631
4.14.3 Maple step by step solution	635

Internal problem ID [6195]

Internal file name [OUTPUT/5443_Sunday_June_05_2022_03_38_25_PM_23417214/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.5. Exact Equations. Page 20

Problem number: 14.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "first_order_ode_lie_symmetry_calculated"

Maple gives the following as the ode type

```
[_exact, [_1st_order, `_with_symmetry_[F(x),G(y)]`]]
```

$$2x(1 + \sqrt{x^2 - y}) - \sqrt{x^2 - y}y' = 0$$

4.14.1 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = \frac{2x(1 + \sqrt{x^2 - y})}{\sqrt{x^2 - y}}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2\xi_y - \omega_x\xi - \omega_y\eta = 0 \quad (\text{A})$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 2 to use as anstaz gives

$$\xi = x^2a_4 + xya_5 + y^2a_6 + xa_2 + ya_3 + a_1 \quad (1\text{E})$$

$$\eta = x^2b_4 + xyb_5 + y^2b_6 + xb_2 + yb_3 + b_1 \quad (2\text{E})$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, a_4, a_5, a_6, b_1, b_2, b_3, b_4, b_5, b_6\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$\begin{aligned} & 2xb_4 + yb_5 + b_2 + \frac{2x(1 + \sqrt{x^2 - y})(-2xa_4 + xb_5 - ya_5 + 2yb_6 - a_2 + b_3)}{\sqrt{x^2 - y}} \\ & - \frac{4x^2(1 + \sqrt{x^2 - y})^2(xa_5 + 2ya_6 + a_3)}{x^2 - y} - \left(\frac{2 + 2\sqrt{x^2 - y}}{\sqrt{x^2 - y}} + \frac{2x^2}{x^2 - y} \right. \\ & \left. - \frac{2x^2(1 + \sqrt{x^2 - y})}{(x^2 - y)^{\frac{3}{2}}} \right) (x^2a_4 + xya_5 + y^2a_6 + xa_2 + ya_3 + a_1) \\ & - \left(-\frac{x}{x^2 - y} + \frac{x(1 + \sqrt{x^2 - y})}{(x^2 - y)^{\frac{3}{2}}} \right) (x^2b_4 + xyb_5 + y^2b_6 + xb_2 + yb_3 + b_1) = 0 \end{aligned} \quad (5E)$$

Putting the above in normal form gives

Expression too large to display

Setting the numerator to zero gives

Expression too large to display (6E)

Simplifying the above gives

Expression too large to display (6E)

Since the PDE has radicals, simplifying gives

$$\begin{aligned}
& -2\sqrt{x^2 - y}x^4a_1 - 2\sqrt{x^2 - y}y^4a_6 - 8x^6\sqrt{x^2 - y}ya_6 \\
& + 4x^5\sqrt{x^2 - y}yb_6 - 8x^4\sqrt{x^2 - y}ya_6 + 4\sqrt{x^2 - y}x^3ya_5 \\
& - 6x^2\sqrt{x^2 - y}y^2a_4 + 8\sqrt{x^2 - y}x^2y^2a_6 - 4x\sqrt{x^2 - y}y^3a_5 \\
& + 6\sqrt{x^2 - y}x^4ya_3 + 8\sqrt{x^2 - y}x^3ya_2 - 4\sqrt{x^2 - y}x^3yb_3 \\
& + 4\sqrt{x^2 - y}x^2ya_1 - 2\sqrt{x^2 - y}x^2yb_2 + 2\sqrt{x^2 - y}xy^2b_3 \\
& + 4\sqrt{x^2 - y}x^5ya_5 + 14\sqrt{x^2 - y}x^4y^2a_6 + 12\sqrt{x^2 - y}x^4ya_4 \\
& - 3\sqrt{x^2 - y}x^4yb_5 + 4\sqrt{x^2 - y}x^3y^2a_5 - 8\sqrt{x^2 - y}x^3y^2b_6 \\
& - 4\sqrt{x^2 - y}x^2y^3a_6 - 4\sqrt{x^2 - y}x^3yb_4 + 4\sqrt{x^2 - y}xy^3b_6 - 8x^6a_3 \\
& + 2x^5b_3 - 2y^3a_3 - 2y^2a_1 + 4\sqrt{x^2 - y}x^2ya_3 - 4x\sqrt{x^2 - y}y^2a_2 \\
& + 2x\sqrt{x^2 - y}y^2b_4 - 2x^5a_2 - x^4b_2 - x^3b_1 + \sqrt{x^2 - y}x^4b_2 - 4x^6a_4 \\
& - x^5b_4 - 8x^7a_5 + 2x^6b_5 - 2y^4a_6 - 4\sqrt{x^2 - y}x^5a_2 + \sqrt{x^2 - y}y^3b_5 \\
& + 14x^5ya_5 + 32x^4y^2a_6 + 10x^4ya_4 - 5x^4yb_5 - 2x^3y^2a_5 - 4xy^2a_2 \\
& + 2x^2ya_1 + x^2yb_2 + 3xy^2b_3 + xyb_1 - 9x^3y^2b_6 - 14x^2y^3a_6 \\
& + x^3yb_4 + 3x^2y^2b_5 + 5xy^3b_6 - 6\sqrt{x^2 - y}x^6a_4 + 2\sqrt{x^2 - y}x^5b_4 \\
& - 16x^6ya_6 + 4x^5yb_6 - 6x^2y^2a_4 - 4xy^3a_5 - 4x^7\sqrt{x^2 - y}a_5 \\
& + 2x^6\sqrt{x^2 - y}b_5 - 4x^5\sqrt{x^2 - y}a_5 + 6x^3ya_2 - 5x^3yb_3 - 6x^2y^2a_3 \\
& + 16x^4ya_3 - 4x^6\sqrt{x^2 - y}a_3 + 2x^5\sqrt{x^2 - y}b_3 - 4x^4\sqrt{x^2 - y}a_3 \\
& - 2\sqrt{x^2 - y}y^3a_3 - 2\sqrt{x^2 - y}y^2a_1 + \sqrt{x^2 - y}y^2b_2 = 0
\end{aligned}$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y, \sqrt{x^2 - y}\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2, \sqrt{x^2 - y} = v_3\}$$

The above PDE (6E) now becomes

$$\begin{aligned}
& -4v_1^7v_3a_5 - 8v_1^6v_3v_2a_6 - 4v_1^6v_3a_3 - 6v_3v_1^6a_4 - 8v_1^7a_5 + 4v_3v_1^5v_2a_5 \\
& - 16v_1^6v_2a_6 + 14v_3v_1^4v_2^2a_6 + 2v_1^6v_3b_5 + 4v_1^5v_3v_2b_6 - 4v_3v_1^5a_2 \\
& - 8v_1^6a_3 + 6v_3v_1^4v_2a_3 - 4v_1^6a_4 + 12v_3v_1^4v_2a_4 + 14v_1^5v_2a_5 - 4v_1^5v_3a_5 \\
& + 4v_3v_1^3v_2^2a_5 + 32v_1^4v_2^2a_6 - 8v_1^4v_3v_2a_6 - 4v_3v_1^2v_2^3a_6 + 2v_1^5v_3b_3 \\
& + 2v_3v_1^5b_4 + 2v_1^6b_5 - 3v_3v_1^4v_2b_5 + 4v_1^5v_2b_6 - 8v_3v_1^3v_2^2b_6 - 2v_3v_1^4a_1 \\
& - 2v_1^5a_2 + 8v_3v_1^3v_2a_2 + 16v_1^4v_2a_3 - 4v_1^4v_3a_3 + 10v_1^4v_2a_4 - 6v_1^2v_3v_2^2a_4 \\
& - 2v_1^3v_2^2a_5 + 4v_3v_1^3v_2a_5 - 4v_1v_3v_2^3a_5 - 14v_1^2v_2^3a_6 + 8v_3v_1^2v_2^2a_6 \\
& - 2v_3v_1^4a_6 + v_3v_1^4b_2 + 2v_1^5b_3 - 4v_3v_1^3v_2b_3 - v_1^5b_4 - 4v_3v_1^3v_2b_4 \\
& - 5v_1^4v_2b_5 - 9v_1^3v_2^2b_6 + 4v_3v_1v_2^3b_6 + 4v_3v_1^2v_2a_1 + 6v_1^3v_2a_2 - 4v_1v_3v_2^2a_2 \\
& - 6v_1^2v_2^2a_3 + 4v_3v_1^2v_2a_3 - 2v_3v_2^3a_3 - 6v_1^2v_2^2a_4 - 4v_1v_2^3a_5 - 2v_1^4a_6 \\
& - v_1^4b_2 - 2v_3v_1^2v_2b_2 - 5v_1^3v_2b_3 + 2v_3v_1v_2^2b_3 + v_1^3v_2b_4 + 2v_1v_3v_2^2b_4 \\
& + 3v_1^2v_2^2b_5 + v_3v_2^3b_5 + 5v_1v_2^3b_6 + 2v_1^2v_2a_1 - 2v_3v_2^2a_1 - 4v_1v_2^2a_2 \\
& - 2v_2^3a_3 - v_1^3b_1 + v_1^2v_2b_2 + v_3v_2^2b_2 + 3v_1v_2^2b_3 - 2v_2^2a_1 + v_1v_2b_1 = 0
\end{aligned} \tag{7E}$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2, v_3\}$$

Equation (7E) now becomes

$$\begin{aligned}
& 14v_3v_1^4v_2^2a_6 - 4v_3v_1^2v_2^3a_6 + (-4a_3 - 6a_4 + 2b_5)v_1^6v_3 + (14a_5 + 4b_6)v_1^5v_2 \\
& + (-4a_2 - 4a_5 + 2b_3 + 2b_4)v_1^5v_3 + (16a_3 + 10a_4 - 5b_5)v_1^4v_2 \\
& + (-2a_1 - 4a_3 + b_2)v_1^4v_3 + (-2a_5 - 9b_6)v_1^3v_2^2 + (6a_2 - 5b_3 + b_4)v_1^3v_2 \\
& - 8v_1^6v_3v_2a_6 + (4a_5 + 4b_6)v_1^5v_2v_3 + (6a_3 + 12a_4 - 8a_6 - 3b_5)v_1^4v_2v_3 \\
& + (4a_5 - 8b_6)v_1^3v_2^2v_3 + (8a_2 + 4a_5 - 4b_3 - 4b_4)v_1^3v_2v_3 \\
& + (-6a_4 + 8a_6)v_1^2v_2^2v_3 + (4a_1 + 4a_3 - 2b_2)v_1^2v_2v_3 \\
& + (-4a_5 + 4b_6)v_1v_2^3v_3 + (-4a_2 + 2b_3 + 2b_4)v_1v_2^2v_3 \\
& + (-6a_3 - 6a_4 + 3b_5)v_1^2v_2^2 + (2a_1 + b_2)v_1^2v_2 + (-4a_5 + 5b_6)v_1v_2^3 \\
& + (-4a_2 + 3b_3)v_1v_2^2 + (-2a_3 + b_5)v_2^3v_3 + (-2a_1 + b_2)v_2^2v_3 \\
& - 2v_3v_2^4a_6 + 32v_1^4v_2^2a_6 + v_1v_2b_1 - 14v_1^2v_2^3a_6 - 16v_1^6v_2a_6 \\
& - 4v_1^7v_3a_5 + (-8a_3 - 4a_4 + 2b_5)v_1^6 + (-2a_2 + 2b_3 - b_4)v_1^5 \\
& - 2v_2^3a_3 - 2v_2^2a_1 - v_1^4b_2 - v_1^3b_1 - 8v_1^7a_5 - 2v_2^4a_6 = 0
\end{aligned} \tag{8E}$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} b_1 &= 0 \\ -2a_1 &= 0 \\ -2a_3 &= 0 \\ -8a_5 &= 0 \\ -4a_5 &= 0 \\ -16a_6 &= 0 \\ -14a_6 &= 0 \\ -8a_6 &= 0 \\ -4a_6 &= 0 \\ -2a_6 &= 0 \\ 14a_6 &= 0 \\ 32a_6 &= 0 \\ -b_1 &= 0 \\ -b_2 &= 0 \\ -2a_1 + b_2 &= 0 \\ 2a_1 + b_2 &= 0 \\ -4a_2 + 3b_3 &= 0 \\ -2a_3 + b_5 &= 0 \\ -6a_4 + 8a_6 &= 0 \\ -4a_5 + 4b_6 &= 0 \\ -4a_5 + 5b_6 &= 0 \\ -2a_5 - 9b_6 &= 0 \\ 4a_5 - 8b_6 &= 0 \\ 4a_5 + 4b_6 &= 0 \\ 14a_5 + 4b_6 &= 0 \\ -2a_1 - 4a_3 + b_2 &= 0 \\ 4a_1 + 4a_3 - 2b_2 &= 0 \\ -4a_2 + 2b_3 + 2b_4 &= 0 \\ -2a_2 + 2b_3 - b_4 &= 0 \\ 6a_2 - 5b_3 + b_4 &= 0 \\ -8a_3 - 4a_4 + 2b_5 &= 0 \\ -6a_3 - 6a_4 + 3b_5 &= 0 \\ -4a_3 - 6a_4 + 2b_5 &= 0 \\ 16a_3 + 10a_4 - 5b_5 &= 0 \\ -4a_2 - 4a_5 + 2b_3 + 2b_4 &= 0 \\ 8a_2 + 4a_5 - 4b_3 - 4b_4 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned}
 a_1 &= 0 \\
 a_2 &= \frac{3b_4}{2} \\
 a_3 &= 0 \\
 a_4 &= 0 \\
 a_5 &= 0 \\
 a_6 &= 0 \\
 b_1 &= 0 \\
 b_2 &= 0 \\
 b_3 &= 2b_4 \\
 b_4 &= b_4 \\
 b_5 &= 0 \\
 b_6 &= 0
 \end{aligned}$$

Substituting the above solution in the ansatz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned}
 \xi &= \frac{3x}{2} \\
 \eta &= x^2 + 2y
 \end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned}
 \eta &= \eta - \omega(x, y) \xi \\
 &= x^2 + 2y - \left(\frac{2x(1 + \sqrt{x^2 - y})}{\sqrt{x^2 - y}} \right) \left(\frac{3x}{2} \right) \\
 &= \frac{-2x^2\sqrt{x^2 - y} + 2\sqrt{x^2 - y}y - 3x^2}{\sqrt{x^2 - y}} \\
 \xi &= 0
 \end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{-2x^2\sqrt{x^2-y}+2\sqrt{x^2-y}y-3x^2}{\sqrt{x^2-y}}} dy \end{aligned}$$

Which results in

$$S = \frac{\ln\left(2(x^2 - y)^{\frac{3}{2}} + 3x^2\right)}{3}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{2x(1 + \sqrt{x^2 - y})}{\sqrt{x^2 - y}}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{2x(x^2 + \sqrt{x^2 - y} - y)}{\sqrt{x^2 - y} ((2x^2 - 2y)\sqrt{x^2 - y} + 3x^2)} \\ S_y &= -\frac{\sqrt{x^2 - y}}{(2x^2 - 2y)\sqrt{x^2 - y} + 3x^2} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \tag{4}$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{\ln((2x^2 - 2y)\sqrt{x^2 - y} + 3x^2)}{3} = c_1$$

Which simplifies to

$$\frac{\ln((2x^2 - 2y)\sqrt{x^2 - y} + 3x^2)}{3} = c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{2x(1 + \sqrt{x^2 - y})}{\sqrt{x^2 - y}}$ 	$R = x$ $S = \frac{\ln((2x^2 - 2y)\sqrt{x^2 - y} + 3x^2)}{3}$	$\frac{dS}{dR} = 0$ 

Summary

The solution(s) found are the following

$$\frac{\ln((2x^2 - 2y)\sqrt{x^2 - y} + 3x^2)}{3} = c_1 \quad (1)$$




Figure 137: Slope field plot

Verification of solutions

$$\frac{\ln((2x^2 - 2y)\sqrt{x^2 - y} + 3x^2)}{3} = c_1$$

Verified OK.

4.14.2 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (A)$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} & \left(-\sqrt{x^2 - y} \right) dy = \left(-2x \left(1 + \sqrt{x^2 - y} \right) \right) dx \\ \left(2x \left(1 + \sqrt{x^2 - y} \right) \right) dx + \left(-\sqrt{x^2 - y} \right) dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= 2x \left(1 + \sqrt{x^2 - y} \right) \\ N(x, y) &= -\sqrt{x^2 - y} \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(2x \left(1 + \sqrt{x^2 - y} \right) \right) \\ &= -\frac{x}{\sqrt{x^2 - y}} \end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(-\sqrt{x^2 - y} \right) \\ &= -\frac{x}{\sqrt{x^2 - y}}\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \tag{1}$$

$$\frac{\partial \phi}{\partial y} = N \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int 2x(1 + \sqrt{x^2 - y}) dx \\ \phi &= \frac{(2x^2 - 2y)\sqrt{x^2 - y}}{3} + x^2 + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\begin{aligned}\frac{\partial \phi}{\partial y} &= -\frac{2\sqrt{x^2 - y}}{3} - \frac{2x^2 - 2y}{6\sqrt{x^2 - y}} + f'(y) \\ &= -\sqrt{x^2 - y} + f'(y)\end{aligned} \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = -\sqrt{x^2 - y}$. Therefore equation (4) becomes

$$-\sqrt{x^2 - y} = -\sqrt{x^2 - y} + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = \frac{(2x^2 - 2y) \sqrt{x^2 - y}}{3} + x^2 + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \frac{(2x^2 - 2y) \sqrt{x^2 - y}}{3} + x^2$$

Summary

The solution(s) found are the following

$$\frac{(2x^2 - 2y) \sqrt{x^2 - y}}{3} + x^2 = c_1 \quad (1)$$




Figure 138: Slope field plot

Verification of solutions

$$\frac{(2x^2 - 2y) \sqrt{x^2 - y}}{3} + x^2 = c_1$$

Verified OK.

4.14.3 Maple step by step solution

Let's solve

$$2x(1 + \sqrt{x^2 - y}) - \sqrt{x^2 - y} y' = 0$$

- Highest derivative means the order of the ODE is 1

y'

- Check if ODE is exact

- ODE is exact if the lhs is the total derivative of a C^2 function

$$F'(x, y) = 0$$

- Compute derivative of lhs

$$F'(x, y) + \left(\frac{\partial}{\partial y} F(x, y) \right) y' = 0$$

- Evaluate derivatives

$$-\frac{x}{\sqrt{x^2 - y}} = -\frac{x}{\sqrt{x^2 - y}}$$

- Condition met, ODE is exact

- Exact ODE implies solution will be of this form

$$\left[F(x, y) = c_1, M(x, y) = F'(x, y), N(x, y) = \frac{\partial}{\partial y} F(x, y) \right]$$

- Solve for $F(x, y)$ by integrating $M(x, y)$ with respect to x

$$F(x, y) = \int 2x(1 + \sqrt{x^2 - y}) dx + f_1(y)$$

- Evaluate integral

$$F(x, y) = \frac{2(x^2 - y)^{\frac{3}{2}}}{3} + x^2 + f_1(y)$$

- Take derivative of $F(x, y)$ with respect to y

$$N(x, y) = \frac{\partial}{\partial y} F(x, y)$$

- Compute derivative

$$-\sqrt{x^2 - y} = -\sqrt{x^2 - y} + \frac{d}{dy} f_1(y)$$

- Isolate for $\frac{d}{dy} f_1(y)$

$$\frac{d}{dy} f_1(y) = 0$$

- Solve for $f_1(y)$

$$f_1(y) = 0$$

- Substitute $f_1(y)$ into equation for $F(x, y)$

$$F(x, y) = \frac{2(x^2 - y)^{\frac{3}{2}}}{3} + x^2$$

- Substitute $F(x, y)$ into the solution of the ODE

$$\frac{2(x^2 - y)^{\frac{3}{2}}}{3} + x^2 = c_1$$

- Solve for y

$$\left\{ y = x^2 - \left(-\frac{(-12x^2 + 12c_1)^{\frac{1}{3}}}{4} - \frac{I\sqrt{3}(-12x^2 + 12c_1)^{\frac{1}{3}}}{4} \right)^2, y = x^2 - \left(-\frac{(-12x^2 + 12c_1)^{\frac{1}{3}}}{4} + \frac{I\sqrt{3}(-12x^2 + 12c_1)^{\frac{1}{3}}}{4} \right)^2 \right.$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying homogeneous types:
differential order: 1; looking for linear symmetries
trying exact
<- exact successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 30

```
dsolve((2*x*(1+sqrt(x^2-y(x))))=sqrt(x^2-y(x))*diff(y(x),x),y(x), singsol=all)
```

$$\frac{(2x^2 - 2y(x)) \sqrt{x^2 - y(x)}}{3} + x^2 + c_1 = 0$$

✓ Solution by Mathematica

Time used: 0.898 (sec). Leaf size: 121

```
DSolve[2*x*(1+Sqrt[x^2-y[x]])==Sqrt[x^2-y[x]]*y'[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^2 + \left(\frac{3}{2}\right)^{2/3} \sqrt[3]{-(x^2 + c_1)^2}$$

$$y(x) \rightarrow x^2 - \frac{\sqrt[6]{3}(\sqrt{3} - 3i) \sqrt[3]{-(x^2 + c_1)^2}}{2 \cdot 2^{2/3}}$$

$$y(x) \rightarrow x^2 - \frac{\sqrt[6]{3}(\sqrt{3} + 3i) \sqrt[3]{-(x^2 + c_1)^2}}{2 \cdot 2^{2/3}}$$

4.15 problem 15

4.15.1 Solving as separable ode	637
4.15.2 Solving as first order ode lie symmetry lookup ode	639
4.15.3 Solving as exact ode	640
4.15.4 Maple step by step solution	644

Internal problem ID [6196]

Internal file name [OUTPUT/5444_Sunday_June_05_2022_03_38_28_PM_66698641/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.5. Exact Equations. Page 20

Problem number: 15.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exact", "separable", "first_order_ode_lie_symmetry_lookup"**

Maple gives the following as the ode type

`[_separable]`

$$x \ln(y) + xy + (\ln(x)y + xy)y' = 0$$

4.15.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= -\frac{x(\ln(y) + y)}{y(x + \ln(x))}\end{aligned}$$

Where $f(x) = -\frac{x}{x + \ln(x)}$ and $g(y) = \frac{\ln(y) + y}{y}$. Integrating both sides gives

$$\begin{aligned}\frac{1}{\frac{\ln(y) + y}{y}} dy &= -\frac{x}{x + \ln(x)} dx \\ \int \frac{1}{\frac{\ln(y) + y}{y}} dy &= \int -\frac{x}{x + \ln(x)} dx\end{aligned}$$

$$\int^y \frac{-a}{\ln(-a) + -a} d-a = \int -\frac{x}{x + \ln(x)} dx + c_1$$

Which results in

$$\int^y \frac{-a}{\ln(-a) + -a} d-a = \int -\frac{x}{x + \ln(x)} dx + c_1$$

The solution is

$$\int^y \frac{-a}{\ln(-a) + -a} d-a - \left(\int -\frac{x}{x + \ln(x)} dx \right) - c_1 = 0$$

Summary

The solution(s) found are the following

$$\int^y \frac{-a}{\ln(-a) + -a} d-a - \left(\int -\frac{x}{x + \ln(x)} dx \right) - c_1 = 0 \tag{1}$$




Figure 139: Slope field plot

Verification of solutions

$$\int^y \frac{-a}{\ln(-a) + -a} d-a - \left(\int -\frac{x}{x + \ln(x)} dx \right) - c_1 = 0$$

Verified OK.

4.15.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = -\frac{x(\ln(y) + y)}{y(x + \ln(x))}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **separable**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 105: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= -\frac{x + \ln(x)}{x} \\ \eta(x, y) &= 0\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\eta = 0$ then in this special case

$$R = y$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\xi} dx \\ &= \int \frac{1}{-\frac{x + \ln(x)}{x}} dx\end{aligned}$$

Which results in

$$S = \int -\frac{x}{x + \ln(x)} dx$$

4.15.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0\tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} \left(-\frac{y}{\ln(y) + y} \right) dy &= \left(\frac{x}{x + \ln(x)} \right) dx \\ \left(-\frac{x}{x + \ln(x)} \right) dx + \left(-\frac{y}{\ln(y) + y} \right) dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -\frac{x}{x + \ln(x)} \\ N(x, y) &= -\frac{y}{\ln(y) + y} \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(-\frac{x}{x + \ln(x)} \right) \\ &= 0 \end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(-\frac{y}{\ln(y) + y} \right) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \tag{1}$$

$$\frac{\partial \phi}{\partial y} = N \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\frac{x}{x + \ln(x)} dx \\ \phi &= \int -\frac{-a}{-a + \ln(-a)} d_{-a} + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = -\frac{y}{\ln(y)+y}$. Therefore equation (4) becomes

$$-\frac{y}{\ln(y) + y} = 0 + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = -\frac{y}{\ln(y) + y}$$

Integrating the above w.r.t y gives

$$\int f'(y) dy = \int \left(-\frac{y}{\ln(y) + y} \right) dy$$

$$f(y) = \int_0^y -\frac{-a}{-a + \ln(-a)} d_{-a} + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = \int^x -\frac{-a}{-a + \ln(-a)} d_{-a} + \int_0^y -\frac{-a}{-a + \ln(-a)} d_{-a} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \int^x -\frac{-a}{-a + \ln(-a)} d_{-a} + \int_0^y -\frac{-a}{-a + \ln(-a)} d_{-a}$$

Summary

The solution(s) found are the following

$$\int^x -\frac{-a}{-a + \ln(-a)} d_{-a} + \int_0^y -\frac{-a}{-a + \ln(-a)} d_{-a} = c_1 \quad (1)$$




Figure 140: Slope field plot

Verification of solutions

$$\int^x -\frac{-a}{-a + \ln(-a)} d_{-a} + \int_0^y -\frac{-a}{-a + \ln(-a)} d_{-a} = c_1$$

Verified OK.

4.15.4 Maple step by step solution

Let's solve

$$x \ln(y) + xy + (\ln(x) y + xy) y' = 0$$

- Highest derivative means the order of the ODE is 1

y'

- Separate variables

$$\frac{yy'}{\ln(y)+y} = -\frac{x}{x+\ln(x)}$$

- Integrate both sides with respect to x

$$\int \frac{yy'}{\ln(y)+y} dx = \int -\frac{x}{x+\ln(x)} dx + c_1$$

- Cannot compute integral

$$\int \frac{yy'}{\ln(y)+y} dx = \int -\frac{x}{x+\ln(x)} dx + c_1$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
<- separable successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 28

```
dsolve((x*ln(y(x))+x*y(x))+(y(x)*ln(x)+x*y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$\int \frac{x}{\ln(x) + x} dx + \int^{y(x)} \frac{-a}{-a + \ln(-a)} d_a + c_1 = 0$$

✓ Solution by Mathematica

Time used: 36.692 (sec). Leaf size: 54

```
DSolve[(x*Log[y[x]]+x*y[x])+(y[x]*Log[x]+x*y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions ->
```

$$y(x) \rightarrow \text{InverseFunction} \left[\int_1^{\#1} \frac{K[1]}{K[1] + \log(K[1])} dK[1] \& \right] \left[\int_1^x -\frac{K[2]}{K[2] + \log(K[2])} dK[2] + c_1 \right]$$

$$y(x) \rightarrow W(1)$$

4.16 problem 16

4.16.1 Solving as exact ode	646
4.16.2 Maple step by step solution	649

Internal problem ID [6197]

Internal file name [OUTPUT/5445_Sunday_June_05_2022_03_38_30_PM_69843768/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.5. Exact Equations. Page 20

Problem number: 16.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exact"**

Maple gives the following as the ode type

[_exact]

$$e^{y^2} - \csc(y) \csc(x)^2 + (2xy e^{y^2} - \csc(y) \cot(y) \cot(x)) y' = 0$$

4.16.1 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned} & \left(2y e^{y^2} x - \csc(y) \cot(y) \cot(x) \right) dy = \left(-e^{y^2} + \csc(y) \csc(x)^2 \right) dx \\ & \left(e^{y^2} - \csc(y) \csc(x)^2 \right) dx + \left(2y e^{y^2} x - \csc(y) \cot(y) \cot(x) \right) dy = 0 \end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= e^{y^2} - \csc(y) \csc(x)^2 \\ N(x, y) &= 2y e^{y^2} x - \csc(y) \cot(y) \cot(x) \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(e^{y^2} - \csc(y) \csc(x)^2 \right) \\ &= 2y e^{y^2} + \csc(x)^2 \csc(y) \cot(y) \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(2y e^{y^2} x - \csc(y) \cot(y) \cot(x) \right) \\ &= 2y e^{y^2} + \csc(x)^2 \csc(y) \cot(y) \end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int e^{y^2} - \csc(y) \csc(x)^2 dx \\ \phi &= \csc(y) \cot(x) + e^{y^2} x + f(y)\end{aligned}\quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 2y e^{y^2} x - \csc(y) \cot(y) \cot(x) + f'(y)\quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = 2y e^{y^2} x - \csc(y) \cot(y) \cot(x)$. Therefore equation (4) becomes

$$2y e^{y^2} x - \csc(y) \cot(y) \cot(x) = 2y e^{y^2} x - \csc(y) \cot(y) \cot(x) + f'(y)\quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = \csc(y) \cot(x) + e^{y^2} x + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \csc(y) \cot(x) + e^{y^2} x$$

Summary

The solution(s) found are the following

$$\csc(y) \cot(x) + e^{y^2} x = c_1\quad (1)$$




Figure 141: Slope field plot

Verification of solutions

$$\csc(y) \cot(x) + e^{y^2} x = c_1$$

Verified OK.

4.16.2 Maple step by step solution

Let's solve

$$e^{y^2} - \csc(y) \csc(x)^2 + (2xy e^{y^2} - \csc(y) \cot(y) \cot(x)) y' = 0$$

- Highest derivative means the order of the ODE is 1
 y'
- Check if ODE is exact
 - ODE is exact if the lhs is the total derivative of a C^2 function
 $F'(x, y) = 0$
 - Compute derivative of lhs
 $F'(x, y) + \left(\frac{\partial}{\partial y} F(x, y)\right) y' = 0$

- Evaluate derivatives

$$2y e^{y^2} + \csc(x)^2 \csc(y) \cot(y) = 2y e^{y^2} - \csc(y) \cot(y) (-1 - \cot(x)^2)$$

- Simplify

$$2y e^{y^2} + \csc(x)^2 \csc(y) \cot(y) = 2y e^{y^2} + \csc(x)^2 \csc(y) \cot(y)$$

- Condition met, ODE is exact

- Exact ODE implies solution will be of this form

$$\left[F(x, y) = c_1, M(x, y) = F'(x, y), N(x, y) = \frac{\partial}{\partial y} F(x, y) \right]$$

- Solve for $F(x, y)$ by integrating $M(x, y)$ with respect to x

$$F(x, y) = \int \left(e^{y^2} - \csc(y) \csc(x)^2 \right) dx + f_1(y)$$

- Evaluate integral

$$F(x, y) = \csc(y) \cot(x) + e^{y^2} x + f_1(y)$$

- Take derivative of $F(x, y)$ with respect to y

$$N(x, y) = \frac{\partial}{\partial y} F(x, y)$$

- Compute derivative

$$2y e^{y^2} x - \csc(y) \cot(y) \cot(x) = -\csc(y) \cot(y) \cot(x) + 2y e^{y^2} x + \frac{d}{dy} f_1(y)$$

- Isolate for $\frac{d}{dy} f_1(y)$

$$\frac{d}{dy} f_1(y) = 0$$

- Solve for $f_1(y)$

$$f_1(y) = 0$$

- Substitute $f_1(y)$ into equation for $F(x, y)$

$$F(x, y) = \csc(y) \cot(x) + e^{y^2} x$$

- Substitute $F(x, y)$ into the solution of the ODE

$$\csc(y) \cot(x) + e^{y^2} x = c_1$$

- Solve for y

$$y = \text{RootOf} \left(-Z^2 - \ln \left(\frac{c_1 \tan(x) \sin(-Z) - 1}{x \sin(-Z) \tan(x)} \right) \right)$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying Chini  
differential order: 1; looking for linear symmetries  
trying exact  
<- exact successful`
```

✓ Solution by Maple

Time used: 0.141 (sec). Leaf size: 18

```
dsolve((exp(y(x)^2)-csc(y(x))*csc(x)^2)+(2*x*y(x)*exp(y(x)^2)-csc(y(x))*cot(y(x))*cot(x))*di
```

$$\csc(y(x)) \cot(x) + x e^{y(x)^2} + c_1 = 0$$

✓ Solution by Mathematica

Time used: 175.525 (sec). Leaf size: 23

```
DSolve[(Exp[y[x]^2]-Csc[y[x]]*Csc[x]^2)+(2*x*y[x]*Exp[y[x]^2]-Csc[y[x]]*Cot[y[x]]*Cot[x])*y'
```

$$\text{Solve}\left[-2xe^{y(x)^2} - 2\cot(x)\csc(y(x)) = c_1, y(x)\right]$$

4.17 problem 17

4.17.1 Solving as exact ode	652
4.17.2 Maple step by step solution	656

Internal problem ID [6198]

Internal file name [OUTPUT/5446_Sunday_June_05_2022_03_38_44_PM_7799592/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.5. Exact Equations. Page 20

Problem number: 17.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exact"**

Maple gives the following as the ode type

```
[_exact , _Bernoulli]
```

$$y^2 \sin(2x) - 2y \cos(x)^2 y' = -1$$

4.17.1 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned} (-2 \cos(x)^2 y) dy &= (-1 - y^2 \sin(2x)) dx \\ (y^2 \sin(2x) + 1) dx &+ (-2 \cos(x)^2 y) dy = 0 \end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= y^2 \sin(2x) + 1 \\ N(x, y) &= -2 \cos(x)^2 y \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} (y^2 \sin(2x) + 1) \\ &= 2 \sin(2x) y \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} (-2 \cos(x)^2 y) \\ &= 2 \sin(2x) y \end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int y^2 \sin(2x) + 1 dx \\ \phi &= x - \frac{y^2 \cos(2x)}{2} + f(y)\end{aligned}\tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = -y \cos(2x) + f'(y)\tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = -2 \cos(x)^2 y$. Therefore equation (4) becomes

$$-2 \cos(x)^2 y = -y \cos(2x) + f'(y)\tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$\begin{aligned}f'(y) &= -2 \cos(x)^2 y + y \cos(2x) \\ &= -y\end{aligned}$$

Integrating the above w.r.t y results in

$$\begin{aligned}\int f'(y) dy &= \int (-y) dy \\ f(y) &= -\frac{y^2}{2} + c_1\end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = x - \frac{y^2 \cos(2x)}{2} - \frac{y^2}{2} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = x - \frac{y^2 \cos(2x)}{2} - \frac{y^2}{2}$$

Summary

The solution(s) found are the following

$$x - \frac{\cos(2x) y^2}{2} - \frac{y^2}{2} = c_1 \quad (1)$$




Figure 142: Slope field plot

Verification of solutions

$$x - \frac{\cos(2x) y^2}{2} - \frac{y^2}{2} = c_1$$

Verified OK.

4.17.2 Maple step by step solution

Let's solve

$$y^2 \sin(2x) - 2y \cos(x)^2 y' = -1$$

- Highest derivative means the order of the ODE is 1
 y'
- Check if ODE is exact
 - ODE is exact if the lhs is the total derivative of a C^2 function
 $F'(x, y) = 0$
 - Compute derivative of lhs
 $F'(x, y) + \left(\frac{\partial}{\partial y} F(x, y)\right) y' = 0$
 - Evaluate derivatives
 $2 \sin(2x) y = 4 \cos(x) \sin(x) y$
 - Simplify
 $2 \sin(2x) y = 2 \sin(2x) y$
 - Condition met, ODE is exact
- Exact ODE implies solution will be of this form
 $\left[F(x, y) = c_1, M(x, y) = F'(x, y), N(x, y) = \frac{\partial}{\partial y} F(x, y) \right]$
- Solve for $F(x, y)$ by integrating $M(x, y)$ with respect to x
 $F(x, y) = \int (y^2 \sin(2x) + 1) dx + f_1(y)$
- Evaluate integral
 $F(x, y) = x - \frac{y^2 \cos(2x)}{2} + f_1(y)$
- Take derivative of $F(x, y)$ with respect to y
 $N(x, y) = \frac{\partial}{\partial y} F(x, y)$
- Compute derivative
 $-2 \cos(x)^2 y = -y \cos(2x) + \frac{d}{dy} f_1(y)$
- Isolate for $\frac{d}{dy} f_1(y)$
 $\frac{d}{dy} f_1(y) = -2 \cos(x)^2 y + y \cos(2x)$
- Solve for $f_1(y)$

$$f_1(y) = -\frac{y^2(2\cos(x)^2 - \cos(2x))}{2}$$

- Substitute $f_1(y)$ into equation for $F(x, y)$

$$F(x, y) = x - \frac{y^2 \cos(2x)}{2} - \frac{y^2(2\cos(x)^2 - \cos(2x))}{2}$$

- Substitute $F(x, y)$ into the solution of the ODE

$$x - \frac{y^2 \cos(2x)}{2} - \frac{y^2(2\cos(x)^2 - \cos(2x))}{2} = c_1$$

- Solve for y

$$\left\{ y = \frac{\sqrt{-c_1+x}}{\cos(x)}, y = -\frac{\sqrt{-c_1+x}}{\cos(x)} \right\}$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
<- Bernoulli successful`

```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 24

```
dsolve((1+y(x)^2*sin(2*x))-(2*y(x)*cos(x)^2)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \sec(x) \sqrt{x + c_1}$$

$$y(x) = -\sec(x) \sqrt{x + c_1}$$

✓ Solution by Mathematica

Time used: 0.294 (sec). Leaf size: 32

```
DSolve[(1+y[x]^2*Sin[2*x])-(2*y[x]*Cos[x]^2)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sqrt{x + c_1} \sec(x)$$

$$y(x) \rightarrow \sqrt{x + c_1} \sec(x)$$

4.18 problem 18

4.18.1 Solving as exact ode	658
4.18.2 Maple step by step solution	662

Internal problem ID [6199]

Internal file name [OUTPUT/5447_Sunday_June_05_2022_03_38_47_PM_64144589/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.5. Exact Equations. Page 20

Problem number: 18.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exact"**

Maple gives the following as the ode type

`[_separable]`

$$\frac{x}{(x^2 + y^2)^{\frac{3}{2}}} + \frac{yy'}{(x^2 + y^2)^{\frac{3}{2}}} = 0$$

4.18.1 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}(-y) dy &= (x) dx \\ (-x) dx + (-y) dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -x \\ N(x, y) &= -y\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-x) \\ &= 0\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(-y) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\int \frac{\partial \phi}{\partial x} dx = \int M dx$$

$$\int \frac{\partial \phi}{\partial x} dx = \int -x dx$$

$$\phi = -\frac{x^2}{2} + f(y) \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = -y$. Therefore equation (4) becomes

$$-y = 0 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = -y$$

Integrating the above w.r.t y gives

$$\int f'(y) dy = \int (-y) dy$$

$$f(y) = -\frac{y^2}{2} + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\frac{x^2}{2} - \frac{y^2}{2} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\frac{x^2}{2} - \frac{y^2}{2}$$

Summary

The solution(s) found are the following

$$-\frac{x^2}{2} - \frac{y^2}{2} = c_1 \tag{1}$$




Figure 143: Slope field plot

Verification of solutions

$$-\frac{x^2}{2} - \frac{y^2}{2} = c_1$$

Verified OK.

4.18.2 Maple step by step solution

Let's solve

$$\frac{x}{(x^2+y^2)^{\frac{3}{2}}} + \frac{yy'}{(x^2+y^2)^{\frac{3}{2}}} = 0$$

- Highest derivative means the order of the ODE is 1

y'

- Integrate both sides with respect to x

$$\int \left(\frac{x}{(x^2+y^2)^{\frac{3}{2}}} + \frac{yy'}{(x^2+y^2)^{\frac{3}{2}}} \right) dx = \int 0 dx + c_1$$

- Evaluate integral

$$-\frac{1}{\sqrt{x^2+y^2}} = c_1$$

- Solve for y

$$\left\{ y = \frac{\sqrt{-x^2c_1^2+1}}{c_1}, y = -\frac{\sqrt{-x^2c_1^2+1}}{c_1} \right\}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
<- Bernoulli successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 27

```
dsolve((x/(x^2+y(x)^2)^(3/2))+y(x)/(x^2+y(x)^2)^(3/2))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \sqrt{-x^2 + c_1}$$
$$y(x) = -\sqrt{-x^2 + c_1}$$

✓ Solution by Mathematica

Time used: 0.086 (sec). Leaf size: 39

```
DSolve[(x/(x^2+y[x]^2)^(3/2))+y[x]/(x^2+y[x]^2)^(3/2))*y'[x]==0,y[x],x,IncludeSingularSolut
```

$$y(x) \rightarrow -\sqrt{-x^2 + 2c_1}$$

$$y(x) \rightarrow \sqrt{-x^2 + 2c_1}$$

4.19 problem 19

4.19.1 Solving as exact ode	664
4.19.2 Maple step by step solution	668

Internal problem ID [6200]

Internal file name [OUTPUT/5448_Sunday_June_05_2022_03_38_48_PM_80823751/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.5. Exact Equations. Page 20

Problem number: 19.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exact"**

Maple gives the following as the ode type

```
[_exact, [_1st_order, ` _with_symmetry_[F(x)*G(y),0] `]]
```

$$3x^2(1 + \ln(y)) + \left(\frac{x^3}{y} - 2y\right) y' = 0$$

4.19.1 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned}\left(\frac{x^3}{y} - 2y\right) dy &= (-3x^2(1 + \ln(y))) dx \\ (3x^2(1 + \ln(y))) dx + \left(\frac{x^3}{y} - 2y\right) dy &= 0\end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= 3x^2(1 + \ln(y)) \\ N(x, y) &= \frac{x^3}{y} - 2y\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} (3x^2(1 + \ln(y))) \\ &= \frac{3x^2}{y}\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{x^3}{y} - 2y \right) \\ &= \frac{3x^2}{y}\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int 3x^2(1 + \ln(y)) dx \\ \phi &= x^3(1 + \ln(y)) + f(y)\end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = \frac{x^3}{y} + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{x^3}{y} - 2y$. Therefore equation (4) becomes

$$\frac{x^3}{y} - 2y = \frac{x^3}{y} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = -2y$$

Integrating the above w.r.t y gives

$$\begin{aligned}\int f'(y) dy &= \int (-2y) dy \\ f(y) &= -y^2 + c_1\end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = x^3(1 + \ln(y)) - y^2 + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = x^3(1 + \ln(y)) - y^2$$

The solution becomes

$$y = e^{-\frac{x^3 \operatorname{LambertW}\left(-2e^{-\frac{-2x^3+2c_1}{x^3}}\right) + 2x^3 - 2c_1}{2x^3}}$$

Summary

The solution(s) found are the following

$$y = e^{-\frac{x^3 \operatorname{LambertW}\left(-2e^{-\frac{-2x^3+2c_1}{x^3}}\right) + 2x^3 - 2c_1}{2x^3}} \quad (1)$$




Figure 144: Slope field plot

Verification of solutions

$$y = e^{-\frac{x^3 \operatorname{LambertW}\left(-\frac{2e^{-2x^3+2c_1}}{x^3}\right) + 2x^3 - 2c_1}{2x^3}}$$

Verified OK.

4.19.2 Maple step by step solution

Let's solve

$$3x^2(1 + \ln(y)) + \left(\frac{x^3}{y} - 2y\right) y' = 0$$

- Highest derivative means the order of the ODE is 1
- Check if ODE is exact
 - ODE is exact if the lhs is the total derivative of a C^2 function
 - Compute derivative of lhs
 - Evaluate derivatives
 - Condition met, ODE is exact
- Exact ODE implies solution will be of this form
- Solve for $F(x, y)$ by integrating $M(x, y)$ with respect to x
- Evaluate integral
- Take derivative of $F(x, y)$ with respect to y
- Compute derivative

- Isolate for $\frac{d}{dy}f_1(y)$

$$\frac{d}{dy}f_1(y) = -2y$$
- Solve for $f_1(y)$

$$f_1(y) = -y^2$$
- Substitute $f_1(y)$ into equation for $F(x, y)$

$$F(x, y) = x^3(1 + \ln(y)) - y^2$$
- Substitute $F(x, y)$ into the solution of the ODE

$$x^3(1 + \ln(y)) - y^2 = c_1$$
- Solve for y

$$y = e^{-\frac{x^3 \text{LambertW}\left(-\frac{2e^{-\frac{x^3+c_1}{x^3}}}{x^3}\right) + 2x^3 - 2c_1}{2x^3}}$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying Chini
differential order: 1; looking for linear symmetries
trying exact
<- exact successful`

```

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 43

```
dsolve((3*x^2*(1+ln(y(x))))+(x^3/y(x)-2*y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = e^{-\frac{x^3 \text{LambertW}\left(-\frac{2e^{-\frac{-2x^3-2c_1}{x^3}}}{x^3}\right) + 2x^3 + 2c_1}{2x^3}}$$

✓ Solution by Mathematica

Time used: 60.17 (sec). Leaf size: 79

```
DSolve[(3*x^2*(1+Log[y[x]]))+(x^3/y[x]-2*y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> T
```

$$y(x) \rightarrow -\frac{ix^{3/2} \sqrt{W\left(-\frac{2e^{-2+\frac{2c_1}{x^3}}}{x^3}\right)}}{\sqrt{2}}$$
$$y(x) \rightarrow \frac{ix^{3/2} \sqrt{W\left(-\frac{2e^{-2+\frac{2c_1}{x^3}}}{x^3}\right)}}{\sqrt{2}}$$

4.20 problem 20

4.20.1 Solving as first order ode lie symmetry calculated ode	671
4.20.2 Solving as exact ode	677
4.20.3 Maple step by step solution	681

Internal problem ID [6201]

Internal file name [OUTPUT/5449_Sunday_June_05_2022_03_38_52_PM_17513810/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.5. Exact Equations. Page 20

Problem number: 20.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "first_order_ode_lie_symmetry_calculated"

Maple gives the following as the ode type

```
[[_1st_order , _with_linear_symmetries], _exact , _rational]
```

$$\frac{-xy' + y}{(x + y)^2} + y' = 1$$

4.20.1 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = \frac{x^2 + 2xy + y^2 - y}{x^2 + 2xy + y^2 - x}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \tag{A}$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \tag{1E}$$

$$\eta = xb_2 + yb_3 + b_1 \tag{2E}$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$\begin{aligned} b_2 + \frac{(x^2 + 2xy + y^2 - y)(b_3 - a_2)}{x^2 + 2xy + y^2 - x} - \frac{(x^2 + 2xy + y^2 - y)^2 a_3}{(x^2 + 2xy + y^2 - x)^2} \\ - \left(\frac{2y + 2x}{x^2 + 2xy + y^2 - x} - \frac{(x^2 + 2xy + y^2 - y)(2x + 2y - 1)}{(x^2 + 2xy + y^2 - x)^2} \right) (xa_2 \\ + ya_3 + a_1) - \left(\frac{2x + 2y - 1}{x^2 + 2xy + y^2 - x} \right. \\ \left. - \frac{(x^2 + 2xy + y^2 - y)(2y + 2x)}{(x^2 + 2xy + y^2 - x)^2} \right) (xb_2 + yb_3 + b_1) = 0 \end{aligned} \quad (5E)$$

Putting the above in normal form gives

$$\frac{-x^4a_2 + x^4a_3 - x^4b_2 - x^4b_3 + 4x^3ya_2 + 4x^3ya_3 - 4x^3yb_2 - 4x^3yb_3 + 6x^2y^2a_2 + 6x^2y^2a_3 - 6x^2y^2b_2 - 6x^2y^2b_3 - 4xy^3a_2 - 4xy^3a_3 + 4xy^3b_2 + 4xy^3b_3 - y^4a_2 - y^4a_3 + y^4b_2 + y^4b_3 + 2x^3a_2 + x^3b_2 - x^3b_3 + x^2ya_2 + 3x^2ya_3 - 2x^2yb_2 + 2xy^2a_3 - 3xy^2b_2 - xy^2b_3 + y^3a_2 - y^3a_3 - 2y^3b_3 + x^2a_1 + 3x^2b_1 - 2xya_1 + 2xyb_1 - 3y^2a_1 - y^2b_1 - xb_1 + ya_1}{1} = 0$$

Setting the numerator to zero gives

$$\begin{aligned} -x^4a_2 - x^4a_3 + x^4b_2 + x^4b_3 - 4x^3ya_2 - 4x^3ya_3 + 4x^3yb_2 + 4x^3yb_3 \\ - 6x^2y^2a_2 - 6x^2y^2a_3 + 6x^2y^2b_2 + 6x^2y^2b_3 - 4xy^3a_2 - 4xy^3a_3 + 4xy^3b_2 \\ + 4xy^3b_3 - y^4a_2 - y^4a_3 + y^4b_2 + y^4b_3 + 2x^3a_2 + x^3b_2 - x^3b_3 + x^2ya_2 \\ + 3x^2ya_3 - 2x^2yb_2 + 2xy^2a_3 - 3xy^2b_2 - xy^2b_3 + y^3a_2 - y^3a_3 - 2y^3b_3 \\ + x^2a_1 + 3x^2b_1 - 2xya_1 + 2xyb_1 - 3y^2a_1 - y^2b_1 - xb_1 + ya_1 = 0 \end{aligned} \quad (6E)$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2\}$$

The above PDE (6E) now becomes

$$\begin{aligned}
& -a_2v_1^4 - 4a_2v_1^3v_2 - 6a_2v_1^2v_2^2 - 4a_2v_1v_2^3 - a_2v_2^4 - a_3v_1^4 - 4a_3v_1^3v_2 - 6a_3v_1^2v_2^2 \\
& - 4a_3v_1v_2^3 - a_3v_2^4 + b_2v_1^4 + 4b_2v_1^3v_2 + 6b_2v_1^2v_2^2 + 4b_2v_1v_2^3 + b_2v_2^4 + b_3v_1^4 \\
& + 4b_3v_1^3v_2 + 6b_3v_1^2v_2^2 + 4b_3v_1v_2^3 + b_3v_2^4 + 2a_2v_1^3 + a_2v_1^2v_2 + a_2v_2^3 + 3a_3v_1^2v_2 \\
& + 2a_3v_1v_2^2 - a_3v_2^3 + b_2v_1^3 - 2b_2v_1^2v_2 - 3b_2v_1v_2^2 - b_3v_1^3 - b_3v_1v_2^2 - 2b_3v_2^3 \\
& + a_1v_1^2 - 2a_1v_1v_2 - 3a_1v_2^2 + 3b_1v_1^2 + 2b_1v_1v_2 - b_1v_2^2 + a_1v_2 - b_1v_1 = 0
\end{aligned} \tag{7E}$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2\}$$

Equation (7E) now becomes

$$\begin{aligned}
& (-a_2 - a_3 + b_2 + b_3)v_1^4 + (-4a_2 - 4a_3 + 4b_2 + 4b_3)v_1^3v_2 + (2a_2 + b_2 - b_3)v_1^3 \\
& + (-6a_2 - 6a_3 + 6b_2 + 6b_3)v_1^2v_2^2 + (a_2 + 3a_3 - 2b_2)v_1^2v_2 + (a_1 + 3b_1)v_1^2 \\
& + (-4a_2 - 4a_3 + 4b_2 + 4b_3)v_1v_2^3 + (2a_3 - 3b_2 - b_3)v_1v_2^2 + (-2a_1 + 2b_1)v_1v_2 \\
& - b_1v_1 + (-a_2 - a_3 + b_2 + b_3)v_2^4 + (a_2 - a_3 - 2b_3)v_2^3 + (-3a_1 - b_1)v_2^2 + a_1v_2 = 0
\end{aligned} \tag{8E}$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned}
a_1 &= 0 \\
-b_1 &= 0 \\
-3a_1 - b_1 &= 0 \\
-2a_1 + 2b_1 &= 0 \\
a_1 + 3b_1 &= 0 \\
a_2 - a_3 - 2b_3 &= 0 \\
a_2 + 3a_3 - 2b_2 &= 0 \\
2a_2 + b_2 - b_3 &= 0 \\
2a_3 - 3b_2 - b_3 &= 0 \\
-6a_2 - 6a_3 + 6b_2 + 6b_3 &= 0 \\
-4a_2 - 4a_3 + 4b_2 + 4b_3 &= 0 \\
-a_2 - a_3 + b_2 + b_3 &= 0
\end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= 0 \\ a_2 &= b_3 \\ a_3 &= -b_3 \\ b_1 &= 0 \\ b_2 &= -b_3 \\ b_3 &= b_3 \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned} \xi &= x - y \\ \eta &= -x + y \end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned} \eta &= \eta - \omega(x, y) \xi \\ &= -x + y - \left(\frac{x^2 + 2xy + y^2 - y}{x^2 + 2xy + y^2 - x} \right) (x - y) \\ &= \frac{-2x^3 - 2yx^2 + 2y^2x + 2y^3 + x^2 - y^2}{x^2 + 2xy + y^2 - x} \\ \xi &= 0 \end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y} \right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{-2x^3 - 2yx^2 + 2y^2x + 2y^3 + x^2 - y^2}{x^2 + 2xy + y^2 - x}} dy \end{aligned}$$

Which results in

$$S = -\frac{\ln(x+y)}{2} + \frac{\ln(2x+2y-1)}{2} + \frac{\ln(-x+y)}{2}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{x^2 + 2xy + y^2 - y}{x^2 + 2xy + y^2 - x}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= -\frac{1}{2y+2x} + \frac{1}{2x+2y-1} + \frac{1}{-2y+2x} \\ S_y &= -\frac{1}{2y+2x} + \frac{1}{2x+2y-1} - \frac{1}{-2y+2x} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \quad (4)$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$-\frac{\ln(x+y)}{2} + \frac{\ln(2x+2y-1)}{2} + \frac{\ln(-x+y)}{2} = c_1$$

Which simplifies to

$$-\frac{\ln(x+y)}{2} + \frac{\ln(2x+2y-1)}{2} + \frac{\ln(-x+y)}{2} = c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{x^2+2xy+y^2-y}{x^2+2xy+y^2-x}$ 	$R = x$ $S = -\frac{\ln(x+y)}{2} + \frac{\ln(2x)}{2}$	$\frac{dS}{dR} = 0$ 

Summary

The solution(s) found are the following

$$-\frac{\ln(x+y)}{2} + \frac{\ln(2x+2y-1)}{2} + \frac{\ln(-x+y)}{2} = c_1 \quad (1)$$




Figure 145: Slope field plot

Verification of solutions

$$-\frac{\ln(x+y)}{2} + \frac{\ln(2x+2y-1)}{2} + \frac{\ln(-x+y)}{2} = c_1$$

Verified OK.

4.20.2 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial\phi}{\partial x} + \frac{\partial\phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned}\left(-\frac{x}{(x+y)^2} + 1\right) dy &= \left(-\frac{y}{(x+y)^2} + 1\right) dx \\ \left(\frac{y}{(x+y)^2} - 1\right) dx &+ \left(-\frac{x}{(x+y)^2} + 1\right) dy = 0\end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= \frac{y}{(x+y)^2} - 1 \\ N(x, y) &= -\frac{x}{(x+y)^2} + 1\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(\frac{y}{(x+y)^2} - 1 \right) \\ &= \frac{x-y}{(x+y)^3}\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(-\frac{x}{(x+y)^2} + 1 \right) \\ &= \frac{x-y}{(x+y)^3}\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int \frac{y}{(x+y)^2} - 1 dx \\ \phi &= -x - \frac{y}{x+y} + f(y)\end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\begin{aligned}\frac{\partial \phi}{\partial y} &= -\frac{1}{x+y} + \frac{y}{(x+y)^2} + f'(y) \\ &= -\frac{x}{(x+y)^2} + f'(y)\end{aligned} \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = -\frac{x}{(x+y)^2} + 1$. Therefore equation (4) becomes

$$-\frac{x}{(x+y)^2} + 1 = -\frac{x}{(x+y)^2} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 1$$

Integrating the above w.r.t y gives

$$\int f'(y) dy = \int (1) dy$$
$$f(y) = y + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -x - \frac{y}{x+y} + y + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -x - \frac{y}{x+y} + y$$

Summary

The solution(s) found are the following

$$-x - \frac{y}{x+y} + y = c_1 \tag{1}$$




Figure 146: Slope field plot

Verification of solutions

$$-x - \frac{y}{x+y} + y = c_1$$

Verified OK.

4.20.3 Maple step by step solution

Let's solve

$$\frac{-xy'+y}{(x+y)^2} + y' = 1$$

- Highest derivative means the order of the ODE is 1
- y'
- Check if ODE is exact
- ODE is exact if the lhs is the total derivative of a C^2 function

$$F'(x, y) = 0$$

- Compute derivative of lhs

$$F'(x, y) + \left(\frac{\partial}{\partial y} F(x, y)\right) y' = 0$$

- Evaluate derivatives

$$\frac{1}{(x+y)^2} - \frac{2y}{(x+y)^3} = -\frac{1}{(x+y)^2} + \frac{2x}{(x+y)^3}$$

- Simplify

$$\frac{x-y}{(x+y)^3} = \frac{x-y}{(x+y)^3}$$

- Condition met, ODE is exact

- Exact ODE implies solution will be of this form

$$\left[F(x, y) = c_1, M(x, y) = F'(x, y), N(x, y) = \frac{\partial}{\partial y} F(x, y) \right]$$

- Solve for $F(x, y)$ by integrating $M(x, y)$ with respect to x

$$F(x, y) = \int \left(\frac{y}{(x+y)^2} - 1 \right) dx + f_1(y)$$

- Evaluate integral

$$F(x, y) = -x - \frac{y}{x+y} + f_1(y)$$

- Take derivative of $F(x, y)$ with respect to y

$$N(x, y) = \frac{\partial}{\partial y} F(x, y)$$

- Compute derivative

$$-\frac{x}{(x+y)^2} + 1 = -\frac{1}{x+y} + \frac{y}{(x+y)^2} + \frac{d}{dy}f_1(y)$$

- Isolate for $\frac{d}{dy}f_1(y)$

$$\frac{d}{dy}f_1(y) = -\frac{x}{(x+y)^2} + 1 + \frac{1}{x+y} - \frac{y}{(x+y)^2}$$

- Solve for $f_1(y)$

$$f_1(y) = y$$

- Substitute $f_1(y)$ into equation for $F(x, y)$

$$F(x, y) = -x - \frac{y}{x+y} + y$$

- Substitute $F(x, y)$ into the solution of the ODE

$$-x - \frac{y}{x+y} + y = c_1$$

- Solve for y

$$\left\{ y = \frac{1}{2} + \frac{c_1}{2} - \frac{\sqrt{c_1^2 + 4c_1x + 4x^2 + 2c_1 + 1}}{2}, y = \frac{1}{2} + \frac{c_1}{2} + \frac{\sqrt{c_1^2 + 4c_1x + 4x^2 + 2c_1 + 1}}{2} \right\}$$

Maple trace

```
`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying Chini
differential order: 1; looking for linear symmetries
differential order: 1; found: 1 linear symmetries. Trying reduction of order
1st order, trying the canonical coordinates of the invariance group
  -> Calling odsolve with the ODE`, diff(y(x), x) = (x-y(x))/(y(x)-x), y(x)`
      Methods for first order ODEs:
          --- Trying classification methods ---
              trying a quadrature
              trying 1st order linear
              <- 1st order linear successful
          <- 1st order, canonical coordinates successful`
```

*** Suble

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 61

```
dsolve((y(x)-x*diff(y(x),x))/(x+y(x))^2+diff(y(x),x)=1,y(x), singsol=all)
```

$$y(x) = \frac{c_1}{4} + \frac{1}{4} - \frac{\sqrt{c_1^2 + (8x + 2)c_1 + 16\left(x - \frac{1}{4}\right)^2}}{4}$$
$$y(x) = \frac{c_1}{4} + \frac{1}{4} + \frac{\sqrt{c_1^2 + (8x + 2)c_1 + 16\left(x - \frac{1}{4}\right)^2}}{4}$$

✓ Solution by Mathematica

Time used: 0.468 (sec). Leaf size: 76

```
DSolve[(y[x]-x*y'[x])/(x+y[x])^2+y'[x]==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} \left(-\sqrt{4x^2 + 4c_1x + (1 + c_1)^2} + 1 + c_1 \right)$$

$$y(x) \rightarrow \frac{1}{2} \left(\sqrt{4x^2 + 4c_1x + (1 + c_1)^2} + 1 + c_1 \right)$$

$$y(x) \rightarrow -x$$

4.21 problem 21

- 4.21.1 Solving as exact ode 685
4.21.2 Maple step by step solution 689

Internal problem ID [6202]

Internal file name [OUTPUT/5450_Sunday_June_05_2022_03_38_54_PM_42825175/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.5. Exact Equations. Page 20

Problem number: 21.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exact"**

Maple gives the following as the ode type

```
[[_homogeneous, `class A`], _exact, _rational, _dAlembert]
```

$$\frac{4y^2 - 2x^2}{4xy^2 - x^3} + \frac{(8y^2 - x^2)y'}{4y^3 - yx^2} = 0$$

4.21.1 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned}\left(\frac{-x^2 + 8y^2}{-y x^2 + 4y^3}\right) dy &= \left(-\frac{-2x^2 + 4y^2}{-x^3 + 4y^2 x}\right) dx \\ \left(\frac{-2x^2 + 4y^2}{-x^3 + 4y^2 x}\right) dx &+ \left(\frac{-x^2 + 8y^2}{-y x^2 + 4y^3}\right) dy = 0\end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= \frac{-2x^2 + 4y^2}{-x^3 + 4y^2 x} \\ N(x, y) &= \frac{-x^2 + 8y^2}{-y x^2 + 4y^3}\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(\frac{-2x^2 + 4y^2}{-x^3 + 4y^2 x}\right) \\ &= \frac{8yx}{(x^2 - 4y^2)^2}\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{-x^2 + 8y^2}{-yx^2 + 4y^3} \right) \\ &= \frac{8yx}{(x^2 - 4y^2)^2}\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int \frac{-2x^2 + 4y^2}{-x^3 + 4y^2x} dx \\ \phi &= \frac{\ln(2y + x)}{2} + \ln(x) + \frac{\ln(x - 2y)}{2} + f(y)\end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\begin{aligned}\frac{\partial \phi}{\partial y} &= \frac{1}{2y + x} - \frac{1}{x - 2y} + f'(y) \\ &= -\frac{4y}{x^2 - 4y^2} + f'(y)\end{aligned} \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{-x^2 + 8y^2}{-yx^2 + 4y^3}$. Therefore equation (4) becomes

$$\frac{-x^2 + 8y^2}{-yx^2 + 4y^3} = -\frac{4y}{x^2 - 4y^2} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = \frac{1}{y}$$

Integrating the above w.r.t y gives

$$\int f'(y) dy = \int \left(\frac{1}{y}\right) dy$$

$$f(y) = \ln(y) + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = \frac{\ln(2y+x)}{2} + \ln(x) + \frac{\ln(x-2y)}{2} + \ln(y) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \frac{\ln(2y+x)}{2} + \ln(x) + \frac{\ln(x-2y)}{2} + \ln(y)$$

Summary

The solution(s) found are the following

$$\frac{\ln(2y+x)}{2} + \ln(x) + \frac{\ln(x-2y)}{2} + \ln(y) = c_1 \tag{1}$$




Figure 147: Slope field plot

Verification of solutions

$$\frac{\ln(2y+x)}{2} + \ln(x) + \frac{\ln(x-2y)}{2} + \ln(y) = c_1$$

Verified OK.

4.21.2 Maple step by step solution

Let's solve

$$\frac{4y^2-2x^2}{4xy^2-x^3} + \frac{(8y^2-x^2)y'}{4y^3-yx^2} = 0$$

- Highest derivative means the order of the ODE is 1

y'

- Check if ODE is exact

- ODE is exact if the lhs is the total derivative of a C^2 function

$$F'(x, y) = 0$$

- Compute derivative of lhs

$$F'(x, y) + \left(\frac{\partial}{\partial y} F(x, y)\right) y' = 0$$

- Evaluate derivatives

$$\frac{8y}{-x^3+4y^2x} - \frac{8(-2x^2+4y^2)xy}{(-x^3+4y^2x)^2} = -\frac{2x}{-yx^2+4y^3} + \frac{2(-x^2+8y^2)xy}{(-yx^2+4y^3)^2}$$

- Simplify

$$\frac{8yx}{(x^2-4y^2)^2} = \frac{8yx}{(x^2-4y^2)^2}$$

- Condition met, ODE is exact

- Exact ODE implies solution will be of this form

$$\left[F(x, y) = c_1, M(x, y) = F'(x, y), N(x, y) = \frac{\partial}{\partial y} F(x, y) \right]$$

- Solve for $F(x, y)$ by integrating $M(x, y)$ with respect to x

$$F(x, y) = \int \frac{-2x^2+4y^2}{-x^3+4y^2x} dx + f_1(y)$$

- Evaluate integral

$$F(x, y) = \frac{\ln(2y+x)}{2} + \ln(x) + \frac{\ln(x-2y)}{2} + f_1(y)$$

- Take derivative of $F(x, y)$ with respect to y

$$N(x, y) = \frac{\partial}{\partial y} F(x, y)$$

- Compute derivative

$$\frac{-x^2+8y^2}{-yx^2+4y^3} = \frac{1}{2y+x} - \frac{1}{x-2y} + \frac{d}{dy}f_1(y)$$

- Isolate for $\frac{d}{dy}f_1(y)$

$$\frac{d}{dy}f_1(y) = \frac{-x^2+8y^2}{-yx^2+4y^3} - \frac{1}{2y+x} + \frac{1}{x-2y}$$

- Solve for $f_1(y)$

$$f_1(y) = -\frac{\ln(x-2y)}{2} + \frac{\ln(-x+2y)}{2} + \ln(y)$$

- Substitute $f_1(y)$ into equation for $F(x, y)$

$$F(x, y) = \frac{\ln(2y+x)}{2} + \ln(x) + \frac{\ln(-x+2y)}{2} + \ln(y)$$

- Substitute $F(x, y)$ into the solution of the ODE

$$\frac{\ln(2y+x)}{2} + \ln(x) + \frac{\ln(-x+2y)}{2} + \ln(y) = c_1$$

- Solve for y

$$y = -\frac{x\text{RootOf}\left(2_Z^6x^5+4(e^{c_1})^2_Z^6-5x^4_Z^4+4x^3_Z^2-x^2\right)^2-1}{2\text{RootOf}\left(2_Z^6x^5+4(e^{c_1})^2_Z^6-5x^4_Z^4+4x^3_Z^2-x^2\right)^2}$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying homogeneous D
<- homogeneous successful`

```

✓ Solution by Maple

Time used: 0.703 (sec). Leaf size: 157

```
dsolve(( (4*y(x)^2-2*x^2)/(4*x*y(x)^2-x^3))+( (8*y(x)^2-x^2)/(4*y(x)^3-x^2*y(x)) )*diff(y(x)
```

$$y(x) = -\frac{\sqrt{2} \sqrt{c_1 x (c_1^3 x^3 - \sqrt{c_1^6 x^6 + 16})}}{4c_1^2 x}$$

$$y(x) = \frac{\sqrt{2} \sqrt{c_1 x (c_1^3 x^3 - \sqrt{c_1^6 x^6 + 16})}}{4c_1^2 x}$$

$$y(x) = -\frac{\sqrt{2} \sqrt{c_1 x (c_1^3 x^3 + \sqrt{c_1^6 x^6 + 16})}}{4c_1^2 x}$$

$$y(x) = \frac{\sqrt{2} \sqrt{c_1 x (c_1^3 x^3 + \sqrt{c_1^6 x^6 + 16})}}{4c_1^2 x}$$

✓ Solution by Mathematica

Time used: 12.331 (sec). Leaf size: 297

`DSolve[((4*y[x]^2-2*x^2)/(4*x*y[x]^2-x^3))+((8*y[x]^2-x^2)/(4*y[x]^3-x^2*y[x]))*y'[x]==0,`

$$y(x) \rightarrow -\frac{\sqrt{x^2 - \frac{\sqrt{x^6 - 16e^{2c_1}}}{x}}}{2\sqrt{2}}$$

$$y(x) \rightarrow \frac{\sqrt{x^2 - \frac{\sqrt{x^6 - 16e^{2c_1}}}{x}}}{2\sqrt{2}}$$

$$y(x) \rightarrow -\frac{\sqrt{\frac{x^3 + \sqrt{x^6 - 16e^{2c_1}}}{x}}}{2\sqrt{2}}$$

$$y(x) \rightarrow \frac{\sqrt{\frac{x^3 + \sqrt{x^6 - 16e^{2c_1}}}{x}}}{2\sqrt{2}}$$

$$y(x) \rightarrow -\frac{\sqrt{x^2 - \frac{\sqrt{x^6}}{x}}}{2\sqrt{2}}$$

$$y(x) \rightarrow \frac{\sqrt{x^2 - \frac{\sqrt{x^6}}{x}}}{2\sqrt{2}}$$

$$y(x) \rightarrow -\frac{\sqrt{\frac{\sqrt{x^6} + x^3}{x}}}{2\sqrt{2}}$$

$$y(x) \rightarrow \frac{\sqrt{\frac{\sqrt{x^6} + x^3}{x}}}{2\sqrt{2}}$$

5 Chapter 1. What is a differential equation.

Section 1.7. Homogeneous Equations. Page 28

5.1	problem 1(a)	694
5.2	problem 1(b)	698
5.3	problem 1(c)	702
5.4	problem 1(d)	706
5.5	problem 1(e)	710
5.6	problem 1(f)	714
5.7	problem 1(g)	720
5.8	problem 1(h)	725
5.9	problem 1(i)	729
5.10	problem 1(j)	733
5.11	problem 4(a)	737
5.12	problem 4(b)	748
5.13	problem 4(c)	756
5.14	problem 4(d)	767
5.15	problem 4(e)	778
5.16	problem 5(a)	795
5.17	problem 5(b)	809
5.18	problem 5(c)	822
5.19	problem 7(a)	835
5.20	problem 7(b)	839
5.21	problem 7(c)	843
5.22	problem 7(d)	847

5.1 problem 1(a)

5.1.1 Solving as homogeneous ode 694

Internal problem ID [6203]

Internal file name [OUTPUT/5451_Sunday_June_05_2022_03_38_58_PM_29077920/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.7. Homogeneous Equations. Page 28

Problem number: 1(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

```
[[_homogeneous, `class A`], _rational, _Bernoulli]
```

$$-2y^2 + xy' = -x^2$$

5.1.1 Solving as homogeneous ode

In canonical form, the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= \frac{-x^2 + 2y^2}{xy} \end{aligned} \tag{1}$$

An ode of the form $y' = \frac{M(x,y)}{N(x,y)}$ is called homogeneous if the functions $M(x, y)$ and $N(x, y)$ are both homogeneous functions and of the same order. Recall that a function $f(x, y)$ is homogeneous of order n if

$$f(t^n x, t^n y) = t^n f(x, y)$$

In this case, it can be seen that both $M = -x^2 + 2y^2$ and $N = xy$ are both homogeneous and of the same order $n = 2$. Therefore this is a homogeneous ode. Since this ode is homogeneous, it is converted to separable ODE using the substitution $u = \frac{y}{x}$, or $y = ux$. Hence

$$\frac{dy}{dx} = \frac{du}{dx}x + u$$

Applying the transformation $y = ux$ to the above ODE in (1) gives

$$\frac{du}{dx}x + u = -\frac{1}{u} + 2u$$

$$\frac{du}{dx} = \frac{-\frac{1}{u(x)} + u(x)}{x}$$

Or

$$u'(x) - \frac{-\frac{1}{u(x)} + u(x)}{x} = 0$$

Or

$$u'(x)u(x)x - u(x)^2 + 1 = 0$$

Which is now solved as separable in $u(x)$. Which is now solved in $u(x)$. In canonical form the ODE is

$$u' = F(x, u)$$

$$= f(x)g(u)$$

$$= \frac{u^2 - 1}{ux}$$

Where $f(x) = \frac{1}{x}$ and $g(u) = \frac{u^2-1}{u}$. Integrating both sides gives

$$\frac{1}{\frac{u^2-1}{u}} du = \frac{1}{x} dx$$

$$\int \frac{1}{\frac{u^2-1}{u}} du = \int \frac{1}{x} dx$$

$$\frac{\ln(u-1)}{2} + \frac{\ln(u+1)}{2} = \ln(x) + c_2$$

The above can be written as

$$\left(\frac{1}{2}\right) (\ln(u-1) + \ln(u+1)) = \ln(x) + 2c_2$$

$$\ln(u-1) + \ln(u+1) = (2) (\ln(x) + 2c_2)$$

$$= 2\ln(x) + 4c_2$$

Raising both side to exponential gives

$$e^{\ln(u-1)+\ln(u+1)} = e^{2\ln(x)+2c_2}$$

Which simplifies to

$$\begin{aligned}u^2 - 1 &= 2c_2x^2 \\ &= c_3x^2\end{aligned}$$

The solution is

$$u(x)^2 - 1 = c_3x^2$$

Now u in the above solution is replaced back by y using $u = \frac{y}{x}$ which results in the solution

$$\frac{y^2}{x^2} - 1 = c_3x^2$$

Summary

The solution(s) found are the following

$$\frac{y^2}{x^2} - 1 = c_3x^2 \quad (1)$$




Figure 148: Slope field plot

Verification of solutions

$$\frac{y^2}{x^2} - 1 = c_3x^2$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
<- Bernoulli successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 30

```
dsolve((x^2-2*y(x)^2)+(x*y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \sqrt{c_1 x^2 + 1} x$$
$$y(x) = -\sqrt{c_1 x^2 + 1} x$$

✓ Solution by Mathematica

Time used: 0.451 (sec). Leaf size: 39

```
DSolve[(x^2-2*y[x]^2)+(x*y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sqrt{x^2 + c_1 x^4}$$
$$y(x) \rightarrow \sqrt{x^2 + c_1 x^4}$$

5.2 problem 1(b)

5.2.1 Solving as homogeneous ode 698

Internal problem ID [6204]

Internal file name [OUTPUT/5452_Sunday_June_05_2022_03_39_00_PM_22095681/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.7. Homogeneous Equations. Page 28

Problem number: 1(b).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

```
[[_homogeneous, `class A`], _rational, _Bernoulli]
```

$$x^2y' - 3xy - 2y^2 = 0$$

5.2.1 Solving as homogeneous ode

In canonical form, the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= \frac{y(2y + 3x)}{x^2}\end{aligned}\tag{1}$$

An ode of the form $y' = \frac{M(x,y)}{N(x,y)}$ is called homogeneous if the functions $M(x, y)$ and $N(x, y)$ are both homogeneous functions and of the same order. Recall that a function $f(x, y)$ is homogeneous of order n if

$$f(t^n x, t^n y) = t^n f(x, y)$$

In this case, it can be seen that both $M = y(2y + 3x)$ and $N = x^2$ are both homogeneous and of the same order $n = 2$. Therefore this is a homogeneous ode. Since this ode is homogeneous, it is converted to separable ODE using the substitution $u = \frac{y}{x}$, or $y = ux$.

Hence

$$\frac{dy}{dx} = \frac{du}{dx}x + u$$

Applying the transformation $y = ux$ to the above ODE in (1) gives

$$\frac{du}{dx}x + u = 2u^2 + 3u$$

$$\frac{du}{dx} = \frac{2u(x)^2 + 2u(x)}{x}$$

Or

$$u'(x) - \frac{2u(x)^2 + 2u(x)}{x} = 0$$

Or

$$u'(x)x - 2u(x)^2 - 2u(x) = 0$$

Which is now solved as separable in $u(x)$. Which is now solved in $u(x)$. In canonical form the ODE is

$$u' = F(x, u)$$

$$= f(x)g(u)$$

$$= \frac{2u(u+1)}{x}$$

Where $f(x) = \frac{2}{x}$ and $g(u) = u(u+1)$. Integrating both sides gives

$$\frac{1}{u(u+1)} du = \frac{2}{x} dx$$

$$\int \frac{1}{u(u+1)} du = \int \frac{2}{x} dx$$

$$-\ln(u+1) + \ln(u) = 2\ln(x) + c_2$$

Raising both side to exponential gives

$$e^{-\ln(u+1)+\ln(u)} = e^{2\ln(x)+c_2}$$

Which simplifies to

$$\frac{u}{u+1} = c_3x^2$$

Now u in the above solution is replaced back by y using $u = \frac{y}{x}$ which results in the solution

$$y = -\frac{x^3c_3}{c_3x^2 - 1}$$

Summary

The solution(s) found are the following

$$y = -\frac{x^3 c_3}{c_3 x^2 - 1} \quad (1)$$




Figure 149: Slope field plot

Verification of solutions

$$y = -\frac{x^3 c_3}{c_3 x^2 - 1}$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
<- Bernoulli successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(x^2*diff(y(x),x)-3*x*y(x)-2*y(x)^2=0,y(x), singsol=all)
```

$$y(x) = \frac{x^3}{-x^2 + c_1}$$

✓ Solution by Mathematica

Time used: 0.141 (sec). Leaf size: 25

```
DSolve[x^2*y'[x]-3*x*y[x]-2*y[x]^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{x^3}{x^2 - c_1}$$
$$y(x) \rightarrow 0$$

5.3 problem 1(c)

5.3.1 Solving as homogeneous ode 702

Internal problem ID [6205]

Internal file name [OUTPUT/5453_Sunday_June_05_2022_03_39_02_PM_43511898/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.7. Homogeneous Equations. Page 28

Problem number: 1(c).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

```
[[_homogeneous , `class A`], _dAlembert]
```

$$x^2 y' - 3(x^2 + y^2) \arctan\left(\frac{y}{x}\right) - xy = 0$$

5.3.1 Solving as homogeneous ode

In canonical form, the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= \frac{3 \arctan\left(\frac{y}{x}\right) x^2 + 3 \arctan\left(\frac{y}{x}\right) y^2 + xy}{x^2} \end{aligned} \quad (1)$$

An ode of the form $y' = \frac{M(x,y)}{N(x,y)}$ is called homogeneous if the functions $M(x, y)$ and $N(x, y)$ are both homogeneous functions and of the same order. Recall that a function $f(x, y)$ is homogeneous of order n if

$$f(t^n x, t^n y) = t^n f(x, y)$$

In this case, it can be seen that both $M = 3 \arctan\left(\frac{y}{x}\right) x^2 + 3 \arctan\left(\frac{y}{x}\right) y^2 + xy$ and $N = x^2$ are both homogeneous and of the same order $n = 2$. Therefore this is a homogeneous ode. Since this ode is homogeneous, it is converted to separable ODE using the substitution $u = \frac{y}{x}$, or $y = ux$. Hence

$$\frac{dy}{dx} = \frac{du}{dx}x + u$$

Applying the transformation $y = ux$ to the above ODE in (1) gives

$$\frac{du}{dx}x + u = 3 \arctan(u) + 3 \arctan(u) u^2 + u$$

$$\frac{du}{dx} = \frac{3 \arctan(u(x)) + 3 \arctan(u(x)) u(x)^2}{x}$$

Or

$$u'(x) - \frac{3 \arctan(u(x)) + 3 \arctan(u(x)) u(x)^2}{x} = 0$$

Or

$$-3 \arctan(u(x)) u(x)^2 + u'(x) x - 3 \arctan(u(x)) = 0$$

Which is now solved as separable in $u(x)$. Which is now solved in $u(x)$. In canonical form the ODE is

$$u' = F(x, u)$$

$$= f(x)g(u)$$

$$= \frac{3 \arctan(u) (u^2 + 1)}{x}$$

Where $f(x) = \frac{3}{x}$ and $g(u) = \arctan(u) (u^2 + 1)$. Integrating both sides gives

$$\frac{1}{\arctan(u) (u^2 + 1)} du = \frac{3}{x} dx$$

$$\int \frac{1}{\arctan(u) (u^2 + 1)} du = \int \frac{3}{x} dx$$

$$\ln(\arctan(u)) = 3 \ln(x) + c_2$$

Raising both side to exponential gives

$$\arctan(u) = e^{3 \ln(x) + c_2}$$

Which simplifies to

$$\arctan(u) = c_3 x^3$$

Now u in the above solution is replaced back by y using $u = \frac{y}{x}$ which results in the solution

$$y = x \tan(c_3 e^{c_2} x^3)$$

Summary

The solution(s) found are the following

$$y = x \tan(c_3 e^{c_2} x^3) \tag{1}$$




Figure 150: Slope field plot

Verification of solutions

$$y = x \tan(c_3 e^{c_2 x^3})$$

Verified OK.

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying homogeneous D
<- homogeneous successful`

```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 12

```
dsolve(x^2*diff(y(x),x)=3*(x^2+y(x)^2)*arctan(y(x)/x)+x*y(x),y(x), singsol=all)
```

$$y(x) = \tan(c_1 x^3) x$$

✓ Solution by Mathematica

Time used: 5.758 (sec). Leaf size: 30

```
DSolve[x^2*y'[x]==3*(x^2+y[x]^2)*ArcTan[x,y[x]]+x*y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x \tan(x^3(\cosh(3c_1) - \sinh(3c_1)))$$
$$y(x) \rightarrow 0$$

5.4 problem 1(d)

5.4.1 Solving as homogeneous ode 706

Internal problem ID [6206]

Internal file name [OUTPUT/5454_Sunday_June_05_2022_03_39_05_PM_14435377/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.7. Homogeneous Equations. Page 28

Problem number: 1(d).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

```
[[_homogeneous , `class A`], _dAlembert]
```

$$x \sin\left(\frac{y}{x}\right) y' - y \sin\left(\frac{y}{x}\right) = x$$

5.4.1 Solving as homogeneous ode

In canonical form, the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= \frac{y \sin\left(\frac{y}{x}\right) + x}{x \sin\left(\frac{y}{x}\right)} \end{aligned} \quad (1)$$

An ode of the form $y' = \frac{M(x,y)}{N(x,y)}$ is called homogeneous if the functions $M(x, y)$ and $N(x, y)$ are both homogeneous functions and of the same order. Recall that a function $f(x, y)$ is homogeneous of order n if

$$f(t^n x, t^n y) = t^n f(x, y)$$

In this case, it can be seen that both $M = y \sin\left(\frac{y}{x}\right) + x$ and $N = x \sin\left(\frac{y}{x}\right)$ are both homogeneous and of the same order $n = 1$. Therefore this is a homogeneous ode. Since this ode is homogeneous, it is converted to separable ODE using the substitution $u = \frac{y}{x}$, or $y = ux$. Hence

$$\frac{dy}{dx} = \frac{du}{dx}x + u$$

Applying the transformation $y = ux$ to the above ODE in (1) gives

$$\begin{aligned}\frac{du}{dx}x + u &= u + \frac{1}{\sin(u)} \\ \frac{du}{dx} &= \frac{1}{\sin(u(x))x}\end{aligned}$$

Or

$$u'(x) - \frac{1}{\sin(u(x))x} = 0$$

Or

$$u'(x) \sin(u(x))x - 1 = 0$$

Which is now solved as separable in $u(x)$. Which is now solved in $u(x)$. In canonical form the ODE is

$$\begin{aligned}u' &= F(x, u) \\ &= f(x)g(u) \\ &= \frac{1}{\sin(u)x}\end{aligned}$$

Where $f(x) = \frac{1}{x}$ and $g(u) = \frac{1}{\sin(u)}$. Integrating both sides gives

$$\begin{aligned}\frac{1}{\frac{1}{\sin(u)}} du &= \frac{1}{x} dx \\ \int \frac{1}{\frac{1}{\sin(u)}} du &= \int \frac{1}{x} dx \\ -\cos(u) &= \ln(x) + c_2\end{aligned}$$

The solution is

$$-\cos(u(x)) - \ln(x) - c_2 = 0$$

Now u in the above solution is replaced back by y using $u = \frac{y}{x}$ which results in the solution

$$-\cos\left(\frac{y}{x}\right) - \ln(x) - c_2 = 0$$

Summary

The solution(s) found are the following

$$-\cos\left(\frac{y}{x}\right) - \ln(x) - c_2 = 0 \tag{1}$$




Figure 151: Slope field plot

Verification of solutions

$$-\cos\left(\frac{y}{x}\right) - \ln(x) - c_2 = 0$$

Verified OK.

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying homogeneous D
<- homogeneous successful`

```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 15

```
dsolve(x*sin(y(x)/x)*diff(y(x),x)=y(x)*sin(y(x)/x)+x,y(x), singsol=all)
```

$$y(x) = \left(\frac{\pi}{2} + \arcsin(\ln(x) + c_1) \right) x$$

✓ Solution by Mathematica

Time used: 0.461 (sec). Leaf size: 34

```
DSolve[x*Sin[y[x]/x]*y'[x]==y[x]*Sin[y[x]/x]+x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x \arccos(-\log(x) - c_1)$$

$$y(x) \rightarrow x \arccos(-\log(x) - c_1)$$

5.5 problem 1(e)

5.5.1 Solving as homogeneous ode 710

Internal problem ID [6207]

Internal file name [OUTPUT/5455_Sunday_June_05_2022_03_39_07_PM_87586332/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.7. Homogeneous Equations. Page 28

Problem number: 1(e).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

```
[[_homogeneous , `class A`], _dAlembert]
```

$$xy' - y - 2xe^{-\frac{y}{x}} = 0$$

5.5.1 Solving as homogeneous ode

In canonical form, the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= \frac{y + 2xe^{-\frac{y}{x}}}{x} \end{aligned} \quad (1)$$

An ode of the form $y' = \frac{M(x,y)}{N(x,y)}$ is called homogeneous if the functions $M(x, y)$ and $N(x, y)$ are both homogeneous functions and of the same order. Recall that a function $f(x, y)$ is homogeneous of order n if

$$f(t^n x, t^n y) = t^n f(x, y)$$

In this case, it can be seen that both $M = y + 2xe^{-\frac{y}{x}}$ and $N = x$ are both homogeneous and of the same order $n = 1$. Therefore this is a homogeneous ode. Since this ode is homogeneous, it is converted to separable ODE using the substitution $u = \frac{y}{x}$, or $y = ux$. Hence

$$\frac{dy}{dx} = \frac{du}{dx}x + u$$

Applying the transformation $y = ux$ to the above ODE in (1) gives

$$\frac{du}{dx}x + u = u + 2e^{-u}$$

$$\frac{du}{dx} = \frac{2e^{-u(x)}}{x}$$

Or

$$u'(x) - \frac{2e^{-u(x)}}{x} = 0$$

Or

$$u'(x)e^{u(x)}x - 2 = 0$$

Which is now solved as separable in $u(x)$. Which is now solved in $u(x)$. In canonical form the ODE is

$$u' = F(x, u)$$

$$= f(x)g(u)$$

$$= \frac{2e^{-u}}{x}$$

Where $f(x) = \frac{2}{x}$ and $g(u) = e^{-u}$. Integrating both sides gives

$$\frac{1}{e^{-u}} du = \frac{2}{x} dx$$

$$\int \frac{1}{e^{-u}} du = \int \frac{2}{x} dx$$

$$e^u = 2 \ln(x) + c_2$$

The solution is

$$e^{u(x)} - 2 \ln(x) - c_2 = 0$$

Now u in the above solution is replaced back by y using $u = \frac{y}{x}$ which results in the solution

$$e^{\frac{y}{x}} - 2 \ln(x) - c_2 = 0$$

Summary

The solution(s) found are the following

$$e^{\frac{y}{x}} - 2 \ln(x) - c_2 = 0 \tag{1}$$




Figure 152: Slope field plot

Verification of solutions

$$e^{\frac{y}{x}} - 2 \ln(x) - c_2 = 0$$

Verified OK.

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying homogeneous D
<- homogeneous successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 14

```
dsolve(x*diff(y(x),x)=y(x)+2*x*exp(-y(x)/x),y(x), singsol=all)
```

$$y(x) = (\ln(2) + \ln(\ln(x) + c_1))x$$

✓ Solution by Mathematica

Time used: 0.42 (sec). Leaf size: 15

```
DSolve[x*y'[x]==y[x]+2*x*Exp[-y[x]/x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x \log(2 \log(x) + c_1)$$

5.6 problem 1(f)

5.6.1 Solving as homogeneous ode	714
5.6.2 Maple step by step solution	717

Internal problem ID [6208]

Internal file name [OUTPUT/5456_Sunday_June_05_2022_03_39_10_PM_3528254/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.7. Homogeneous Equations. Page 28

Problem number: 1(f).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

```
[[_homogeneous, `class A`], _exact, _rational, [_Abel, `2nd
type`, `class A`]]
```

$$-y - (x + y)y' = -x$$

5.6.1 Solving as homogeneous ode

In canonical form, the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= -\frac{-x + y}{x + y}\end{aligned}\tag{1}$$

An ode of the form $y' = \frac{M(x,y)}{N(x,y)}$ is called homogeneous if the functions $M(x, y)$ and $N(x, y)$ are both homogeneous functions and of the same order. Recall that a function $f(x, y)$ is homogeneous of order n if

$$f(t^n x, t^n y) = t^n f(x, y)$$

In this case, it can be seen that both $M = x - y$ and $N = x + y$ are both homogeneous and of the same order $n = 1$. Therefore this is a homogeneous ode. Since this ode is

homogeneous, it is converted to separable ODE using the substitution $u = \frac{y}{x}$, or $y = ux$. Hence

$$\frac{dy}{dx} = \frac{du}{dx}x + u$$

Applying the transformation $y = ux$ to the above ODE in (1) gives

$$\begin{aligned} \frac{du}{dx}x + u &= \frac{-u + 1}{u + 1} \\ \frac{du}{dx} &= \frac{\frac{-u(x)+1}{u(x)+1} - u(x)}{x} \end{aligned}$$

Or

$$u'(x) - \frac{\frac{-u(x)+1}{u(x)+1} - u(x)}{x} = 0$$

Or

$$u'(x) xu(x) + u'(x) x + u(x)^2 + 2u(x) - 1 = 0$$

Or

$$x(u(x) + 1) u'(x) + u(x)^2 + 2u(x) - 1 = 0$$

Which is now solved as separable in $u(x)$. Which is now solved in $u(x)$. In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{u^2 + 2u - 1}{x(u + 1)} \end{aligned}$$

Where $f(x) = -\frac{1}{x}$ and $g(u) = \frac{u^2+2u-1}{u+1}$. Integrating both sides gives

$$\begin{aligned} \frac{1}{\frac{u^2+2u-1}{u+1}} du &= -\frac{1}{x} dx \\ \int \frac{1}{\frac{u^2+2u-1}{u+1}} du &= \int -\frac{1}{x} dx \\ \frac{\ln(u^2 + 2u - 1)}{2} &= -\ln(x) + c_2 \end{aligned}$$

Raising both side to exponential gives

$$\sqrt{u^2 + 2u - 1} = e^{-\ln(x)+c_2}$$

Which simplifies to

$$\sqrt{u^2 + 2u - 1} = \frac{c_3}{x}$$

Which simplifies to

$$\sqrt{u(x)^2 + 2u(x) - 1} = \frac{c_3 e^{c_2}}{x}$$

The solution is

$$\sqrt{u(x)^2 + 2u(x) - 1} = \frac{c_3 e^{c_2}}{x}$$

Now u in the above solution is replaced back by y using $u = \frac{y}{x}$ which results in the solution

$$\sqrt{\frac{y^2}{x^2} + \frac{2y}{x} - 1} = \frac{c_3 e^{c_2}}{x}$$

Summary

The solution(s) found are the following

$$\sqrt{\frac{y^2 + 2xy - x^2}{x^2}} = \frac{c_3 e^{c_2}}{x} \quad (1)$$




Figure 153: Slope field plot

Verification of solutions

$$\sqrt{\frac{y^2 + 2xy - x^2}{x^2}} = \frac{c_3 e^{c_2}}{x}$$

Verified OK.

5.6.2 Maple step by step solution

Let's solve

$$-y - (x + y) y' = -x$$

- Highest derivative means the order of the ODE is 1
- Check if ODE is exact
 - ODE is exact if the lhs is the total derivative of a C^2 function
 - $F'(x, y) = 0$
 - Compute derivative of lhs
 - $F'(x, y) + \left(\frac{\partial}{\partial y} F(x, y)\right) y' = 0$
 - Evaluate derivatives
 - $-1 = -1$
 - Condition met, ODE is exact
- Exact ODE implies solution will be of this form
- $\left[F(x, y) = c_1, M(x, y) = F'(x, y), N(x, y) = \frac{\partial}{\partial y} F(x, y) \right]$
- Solve for $F(x, y)$ by integrating $M(x, y)$ with respect to x
- $F(x, y) = \int (x - y) dx + f_1(y)$
- Evaluate integral
- $F(x, y) = \frac{x^2}{2} - xy + f_1(y)$
- Take derivative of $F(x, y)$ with respect to y
- $N(x, y) = \frac{\partial}{\partial y} F(x, y)$
- Compute derivative
- $-y - x = -x + \frac{d}{dy} f_1(y)$
- Isolate for $\frac{d}{dy} f_1(y)$

$$\frac{d}{dy} f_1(y) = -y$$

- Solve for $f_1(y)$

$$f_1(y) = -\frac{y^2}{2}$$

- Substitute $f_1(y)$ into equation for $F(x, y)$

$$F(x, y) = \frac{1}{2}x^2 - xy - \frac{1}{2}y^2$$

- Substitute $F(x, y)$ into the solution of the ODE

$$\frac{1}{2}x^2 - xy - \frac{1}{2}y^2 = c_1$$

- Solve for y

$$\{y = -x - \sqrt{2x^2 - 2c_1}, y = -x + \sqrt{2x^2 - 2c_1}\}$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying homogeneous D
<- homogeneous successful`

```

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 51

```
dsolve((x-y(x))-(x+y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{-c_1x - \sqrt{2x^2c_1^2 + 1}}{c_1}$$

$$y(x) = \frac{-c_1x + \sqrt{2x^2c_1^2 + 1}}{c_1}$$

✓ Solution by Mathematica

Time used: 0.493 (sec). Leaf size: 94

```
DSolve[(x-y[x])-(x+y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x - \sqrt{2x^2 + e^{2c_1}}$$

$$y(x) \rightarrow -x + \sqrt{2x^2 + e^{2c_1}}$$

$$y(x) \rightarrow -\sqrt{2}\sqrt{x^2} - x$$

$$y(x) \rightarrow \sqrt{2}\sqrt{x^2} - x$$

5.7 problem 1(g)

5.7.1 Solving as homogeneous ode	720
5.7.2 Maple step by step solution	722

Internal problem ID [6209]

Internal file name [OUTPUT/5457_Sunday_June_05_2022_03_39_12_PM_9353929/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.7. Homogeneous Equations. Page 28

Problem number: 1(g).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

`[_linear]`

$$xy' + 6y = 2x$$

5.7.1 Solving as homogeneous ode

In canonical form, the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= -\frac{2(-x + 3y)}{x} \end{aligned} \tag{1}$$

An ode of the form $y' = \frac{M(x,y)}{N(x,y)}$ is called homogeneous if the functions $M(x, y)$ and $N(x, y)$ are both homogeneous functions and of the same order. Recall that a function $f(x, y)$ is homogeneous of order n if

$$f(t^n x, t^n y) = t^n f(x, y)$$

In this case, it can be seen that both $M = 2x - 6y$ and $N = x$ are both homogeneous and of the same order $n = 1$. Therefore this is a homogeneous ode. Since this ode is homogeneous, it is converted to separable ODE using the substitution $u = \frac{y}{x}$, or $y = ux$. Hence

$$\frac{dy}{dx} = \frac{du}{dx}x + u$$

Applying the transformation $y = ux$ to the above ODE in (1) gives

$$\begin{aligned}\frac{du}{dx}x + u &= 2 - 6u \\ \frac{du}{dx} &= \frac{2 - 7u(x)}{x}\end{aligned}$$

Or

$$u'(x) - \frac{2 - 7u(x)}{x} = 0$$

Or

$$u'(x)x + 7u(x) - 2 = 0$$

Which is now solved as separable in $u(x)$. Which is now solved in $u(x)$. In canonical form the ODE is

$$\begin{aligned}u' &= F(x, u) \\ &= f(x)g(u) \\ &= \frac{2 - 7u}{x}\end{aligned}$$

Where $f(x) = \frac{1}{x}$ and $g(u) = 2 - 7u$. Integrating both sides gives

$$\begin{aligned}\frac{1}{2 - 7u} du &= \frac{1}{x} dx \\ \int \frac{1}{2 - 7u} du &= \int \frac{1}{x} dx \\ -\frac{\ln(2 - 7u)}{7} &= \ln(x) + c_2\end{aligned}$$

Raising both side to exponential gives

$$\frac{1}{(2 - 7u)^{\frac{1}{7}}} = e^{\ln(x) + c_2}$$

Which simplifies to

$$\frac{1}{(2 - 7u)^{\frac{1}{7}}} = c_3x$$

Now u in the above solution is replaced back by y using $u = \frac{y}{x}$ which results in the solution

$$y = \frac{(2c_3^7x^7e^{7c_2} - 1)e^{-7c_2}}{7x^6c_3^7}$$

Summary

The solution(s) found are the following

$$y = \frac{(2c_3^7 x^7 e^{7c_2} - 1) e^{-7c_2}}{7x^6 c_3^7} \quad (1)$$




Figure 154: Slope field plot

Verification of solutions

$$y = \frac{(2c_3^7 x^7 e^{7c_2} - 1) e^{-7c_2}}{7x^6 c_3^7}$$

Verified OK.

5.7.2 Maple step by step solution

Let's solve

$$xy' + 6y = 2x$$

- Highest derivative means the order of the ODE is 1
- y'
- Isolate the derivative

$$y' = 2 - \frac{6y}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' + \frac{6y}{x} = 2$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) \left(y' + \frac{6y}{x} \right) = 2\mu(x)$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x)y)$

$$\mu(x) \left(y' + \frac{6y}{x} \right) = \mu'(x)y + \mu(x)y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = \frac{6\mu(x)}{x}$$

- Solve to find the integrating factor

$$\mu(x) = x^6$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x)y) \right) dx = \int 2\mu(x) dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x)y = \int 2\mu(x) dx + c_1$$

- Solve for y

$$y = \frac{\int 2\mu(x)dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = x^6$

$$y = \frac{\int 2x^6 dx + c_1}{x^6}$$

- Evaluate the integrals on the rhs

$$y = \frac{\frac{2x^7}{7} + c_1}{x^6}$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```


✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 13

```
dsolve(x*diff(y(x),x)=2*x-6*y(x),y(x), singsol=all)
```

$$y(x) = \frac{2x}{7} + \frac{c_1}{x^6}$$

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 17

```
DSolve[x*y'[x]==2*x-6*y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2x}{7} + \frac{c_1}{x^6}$$

5.8 problem 1(h)

5.8.1 Solving as homogeneous ode 725

Internal problem ID [6210]

Internal file name [OUTPUT/5458_Sunday_June_05_2022_03_39_14_PM_33678347/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.7. Homogeneous Equations. Page 28

Problem number: 1(h).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

```
[[_homogeneous, `class A`], _rational, _dAlembert]
```

$$xy' - \sqrt{x^2 + y^2} = 0$$

5.8.1 Solving as homogeneous ode

In canonical form, the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= \frac{\sqrt{x^2 + y^2}}{x} \end{aligned} \tag{1}$$

An ode of the form $y' = \frac{M(x,y)}{N(x,y)}$ is called homogeneous if the functions $M(x, y)$ and $N(x, y)$ are both homogeneous functions and of the same order. Recall that a function $f(x, y)$ is homogeneous of order n if

$$f(t^n x, t^n y) = t^n f(x, y)$$

In this case, it can be seen that both $M = \sqrt{x^2 + y^2}$ and $N = x$ are both homogeneous and of the same order $n = 1$. Therefore this is a homogeneous ode. Since this ode is homogeneous, it is converted to separable ODE using the substitution $u = \frac{y}{x}$, or $y = ux$. Hence

$$\frac{dy}{dx} = \frac{du}{dx}x + u$$

Applying the transformation $y = ux$ to the above ODE in (1) gives

$$\begin{aligned}\frac{du}{dx}x + u &= \sqrt{u^2 + 1} \\ \frac{du}{dx} &= \frac{\sqrt{u(x)^2 + 1} - u(x)}{x}\end{aligned}$$

Or

$$u'(x) - \frac{\sqrt{u(x)^2 + 1} - u(x)}{x} = 0$$

Or

$$u'(x)x - \sqrt{u(x)^2 + 1} + u(x) = 0$$

Which is now solved as separable in $u(x)$. Which is now solved in $u(x)$. In canonical form the ODE is

$$\begin{aligned}u' &= F(x, u) \\ &= f(x)g(u) \\ &= \frac{\sqrt{u^2 + 1} - u}{x}\end{aligned}$$

Where $f(x) = \frac{1}{x}$ and $g(u) = \sqrt{u^2 + 1} - u$. Integrating both sides gives

$$\begin{aligned}\frac{1}{\sqrt{u^2 + 1} - u} du &= \frac{1}{x} dx \\ \int \frac{1}{\sqrt{u^2 + 1} - u} du &= \int \frac{1}{x} dx \\ \frac{u^2}{2} + \frac{\sqrt{u^2 + 1}u}{2} + \frac{\operatorname{arcsinh}(u)}{2} &= \ln(x) + c_2\end{aligned}$$

The solution is

$$\frac{u(x)^2}{2} + \frac{\sqrt{u(x)^2 + 1}u(x)}{2} + \frac{\operatorname{arcsinh}(u(x))}{2} - \ln(x) - c_2 = 0$$

Now u in the above solution is replaced back by y using $u = \frac{y}{x}$ which results in the solution

$$\frac{y^2}{2x^2} + \frac{\sqrt{\frac{y^2}{x^2} + 1}y}{2x} + \frac{\operatorname{arcsinh}\left(\frac{y}{x}\right)}{2} - \ln(x) - c_2 = 0$$

Summary

The solution(s) found are the following

$$\frac{\sqrt{\frac{x^2+y^2}{x^2}} yx + \operatorname{arcsinh}\left(\frac{y}{x}\right) x^2 + y^2 - 2x^2(\ln(x) + c_2)}{2x^2} = 0 \quad (1)$$




Figure 155: Slope field plot

Verification of solutions

$$\frac{\sqrt{\frac{x^2+y^2}{x^2}} yx + \operatorname{arcsinh}\left(\frac{y}{x}\right) x^2 + y^2 - 2x^2(\ln(x) + c_2)}{2x^2} = 0$$

Verified OK. {0 < x}

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying homogeneous types:  
trying homogeneous G  
1st order, trying the canonical coordinates of the invariance group  
<- 1st order, canonical coordinates successful  
<- homogeneous successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 51

```
dsolve(x*diff(y(x),x)=sqrt(x^2+y(x)^2),y(x), singsol=all)
```

$$\frac{y(x)^2 + \sqrt{x^2 + y(x)^2} y(x) + x^2 \left(\ln \left(y(x) + \sqrt{x^2 + y(x)^2} \right) - c_1 - 3 \ln(x) \right)}{x^2} = 0$$

✓ Solution by Mathematica

Time used: 0.296 (sec). Leaf size: 66

```
DSolve[x*y'[x]==Sqrt[x^2+y[x]^2],y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[\frac{1}{2} \left(\frac{y(x) \left(\sqrt{\frac{y(x)^2}{x^2} + 1} + \frac{y(x)}{x} \right)}{x} - \log \left(\sqrt{\frac{y(x)^2}{x^2} + 1} - \frac{y(x)}{x} \right) \right) = \log(x) + c_1, y(x) \right]$$

5.9 problem 1(i)

5.9.1 Solving as homogeneous ode 729

Internal problem ID [6211]

Internal file name [OUTPUT/5459_Sunday_June_05_2022_03_39_17_PM_43009656/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.7. Homogeneous Equations. Page 28

Problem number: 1(i).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

```
[[_homogeneous, `class A`], _rational, _Bernoulli]
```

$$x^2y' - 2xy - y^2 = 0$$

5.9.1 Solving as homogeneous ode

In canonical form, the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= \frac{y(y + 2x)}{x^2}\end{aligned}\tag{1}$$

An ode of the form $y' = \frac{M(x,y)}{N(x,y)}$ is called homogeneous if the functions $M(x, y)$ and $N(x, y)$ are both homogeneous functions and of the same order. Recall that a function $f(x, y)$ is homogeneous of order n if

$$f(t^n x, t^n y) = t^n f(x, y)$$

In this case, it can be seen that both $M = y(y + 2x)$ and $N = x^2$ are both homogeneous and of the same order $n = 2$. Therefore this is a homogeneous ode. Since this ode is homogeneous, it is converted to separable ODE using the substitution $u = \frac{y}{x}$, or $y = ux$.

Hence

$$\frac{dy}{dx} = \frac{du}{dx}x + u$$

Applying the transformation $y = ux$ to the above ODE in (1) gives

$$\begin{aligned}\frac{du}{dx}x + u &= u^2 + 2u \\ \frac{du}{dx} &= \frac{u(x)^2 + u(x)}{x}\end{aligned}$$

Or

$$u'(x) - \frac{u(x)^2 + u(x)}{x} = 0$$

Or

$$u'(x)x - u(x)^2 - u(x) = 0$$

Which is now solved as separable in $u(x)$. Which is now solved in $u(x)$. In canonical form the ODE is

$$\begin{aligned}u' &= F(x, u) \\ &= f(x)g(u) \\ &= \frac{u(u+1)}{x}\end{aligned}$$

Where $f(x) = \frac{1}{x}$ and $g(u) = u(u+1)$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u(u+1)} du &= \frac{1}{x} dx \\ \int \frac{1}{u(u+1)} du &= \int \frac{1}{x} dx \\ -\ln(u+1) + \ln(u) &= \ln(x) + c_2\end{aligned}$$

Raising both side to exponential gives

$$e^{-\ln(u+1)+\ln(u)} = e^{\ln(x)+c_2}$$

Which simplifies to

$$\frac{u}{u+1} = c_3x$$

Now u in the above solution is replaced back by y using $u = \frac{y}{x}$ which results in the solution

$$y = -\frac{x^2c_3}{c_3x - 1}$$

Summary

The solution(s) found are the following

$$y = -\frac{x^2 c_3}{c_3 x - 1} \quad (1)$$




Figure 156: Slope field plot

Verification of solutions

$$y = -\frac{x^2 c_3}{c_3 x - 1}$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
<- Bernoulli successful`
```


✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(x^2*diff(y(x),x)=y(x)^2+2*x*y(x),y(x), singsol=all)
```

$$y(x) = \frac{x^2}{-x + c_1}$$

✓ Solution by Mathematica

Time used: 0.141 (sec). Leaf size: 23

```
DSolve[x^2*y'[x]==y[x]^2+2*x*y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{x^2}{x - c_1}$$
$$y(x) \rightarrow 0$$

5.10 problem 1(j)

5.10.1 Solving as homogeneous ode 733

Internal problem ID [6212]

Internal file name [OUTPUT/5460_Sunday_June_05_2022_03_39_20_PM_57557365/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.7. Homogeneous Equations. Page 28

Problem number: 1(j).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

```
[[_homogeneous, `class A`], _rational, _Bernoulli]
```

$$y^3 - xy^2y' = -x^3$$

5.10.1 Solving as homogeneous ode

In canonical form, the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= \frac{x^3 + y^3}{xy^2} \end{aligned} \tag{1}$$

An ode of the form $y' = \frac{M(x,y)}{N(x,y)}$ is called homogeneous if the functions $M(x, y)$ and $N(x, y)$ are both homogeneous functions and of the same order. Recall that a function $f(x, y)$ is homogeneous of order n if

$$f(t^n x, t^n y) = t^n f(x, y)$$

In this case, it can be seen that both $M = x^3 + y^3$ and $N = y^2x$ are both homogeneous and of the same order $n = 3$. Therefore this is a homogeneous ode. Since this ode is homogeneous, it is converted to separable ODE using the substitution $u = \frac{y}{x}$, or $y = ux$. Hence

$$\frac{dy}{dx} = \frac{du}{dx}x + u$$

Applying the transformation $y = ux$ to the above ODE in (1) gives

$$\begin{aligned}\frac{du}{dx}x + u &= \frac{1}{u^2} + u \\ \frac{du}{dx} &= \frac{1}{u(x)^2 x}\end{aligned}$$

Or

$$u'(x) - \frac{1}{u(x)^2 x} = 0$$

Or

$$u'(x) u(x)^2 x - 1 = 0$$

Which is now solved as separable in $u(x)$. Which is now solved in $u(x)$. In canonical form the ODE is

$$\begin{aligned}u' &= F(x, u) \\ &= f(x)g(u) \\ &= \frac{1}{u^2 x}\end{aligned}$$

Where $f(x) = \frac{1}{x}$ and $g(u) = \frac{1}{u^2}$. Integrating both sides gives

$$\begin{aligned}\frac{1}{\frac{1}{u^2}} du &= \frac{1}{x} dx \\ \int \frac{1}{\frac{1}{u^2}} du &= \int \frac{1}{x} dx \\ \frac{u^3}{3} &= \ln(x) + c_2\end{aligned}$$

The solution is

$$\frac{u(x)^3}{3} - \ln(x) - c_2 = 0$$

Now u in the above solution is replaced back by y using $u = \frac{y}{x}$ which results in the solution

$$\frac{y^3}{3x^3} - \ln(x) - c_2 = 0$$

Summary

The solution(s) found are the following

$$\frac{y^3}{3x^3} - \ln(x) - c_2 = 0 \tag{1}$$




Figure 157: Slope field plot

Verification of solutions

$$\frac{y^3}{3x^3} - \ln(x) - c_2 = 0$$

Verified OK.

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
<- Bernoulli successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 58

```
dsolve((x^3+y(x)^3)-(x*y(x)^2)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = (c_1 + 3 \ln(x))^{\frac{1}{3}} x$$
$$y(x) = -\frac{(c_1 + 3 \ln(x))^{\frac{1}{3}} (1 + i\sqrt{3}) x}{2}$$
$$y(x) = \frac{(c_1 + 3 \ln(x))^{\frac{1}{3}} (i\sqrt{3} - 1) x}{2}$$

✓ Solution by Mathematica

Time used: 0.197 (sec). Leaf size: 63

```
DSolve[(x^3+y[x]^3)-(x*y[x]^2)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x \sqrt[3]{3 \log(x) + c_1}$$
$$y(x) \rightarrow -\sqrt[3]{-1} x \sqrt[3]{3 \log(x) + c_1}$$
$$y(x) \rightarrow (-1)^{2/3} x \sqrt[3]{3 \log(x) + c_1}$$

5.11 problem 4(a)

5.11.1 Solving as homogeneousTypeMapleC ode 737

5.11.2 Solving as first order ode lie symmetry calculated ode 740

Internal problem ID [6213]

Internal file name [OUTPUT/5461_Sunday_June_05_2022_03_39_22_PM_1042896/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.7. Homogeneous Equations. Page 28

Problem number: 4(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"homogeneousTypeMapleC", "first_order_ode_lie_symmetry_calculated"**

Maple gives the following as the ode type

```
[[_homogeneous, `class C`], _rational, [_Abel, `2nd type`, `class A`]]
```

$$y' - \frac{x + y + 4}{x - y - 6} = 0$$

5.11.1 Solving as homogeneousTypeMapleC ode

Let $Y = y + y_0$ and $X = x + x_0$ then the above is transformed to new ode in $Y(X)$

$$\frac{d}{dX}Y(X) = -\frac{X + x_0 + Y(X) + y_0 + 4}{-X - x_0 + Y(X) + y_0 + 6}$$

Solving for possible values of x_0 and y_0 which makes the above ode a homogeneous ode results in

$$x_0 = 1$$

$$y_0 = -5$$

Using these values now it is possible to easily solve for $Y(X)$. The above ode now becomes

$$\frac{d}{dX}Y(X) = -\frac{X + Y(X)}{-X + Y(X)}$$

In canonical form, the ODE is

$$\begin{aligned} Y' &= F(X, Y) \\ &= -\frac{X + Y}{-X + Y} \end{aligned} \quad (1)$$

An ode of the form $Y' = \frac{M(X, Y)}{N(X, Y)}$ is called homogeneous if the functions $M(X, Y)$ and $N(X, Y)$ are both homogeneous functions and of the same order. Recall that a function $f(X, Y)$ is homogeneous of order n if

$$f(t^n X, t^n Y) = t^n f(X, Y)$$

In this case, it can be seen that both $M = X + Y$ and $N = X - Y$ are both homogeneous and of the same order $n = 1$. Therefore this is a homogeneous ode. Since this ode is homogeneous, it is converted to separable ODE using the substitution $u = \frac{Y}{X}$, or $Y = uX$. Hence

$$\frac{dY}{dX} = \frac{du}{dX}X + u$$

Applying the transformation $Y = uX$ to the above ODE in (1) gives

$$\begin{aligned} \frac{du}{dX}X + u &= \frac{-u - 1}{u - 1} \\ \frac{du}{dX} &= \frac{\frac{-u(X)-1}{u(X)-1} - u(X)}{X} \end{aligned}$$

Or

$$\frac{d}{dX}u(X) - \frac{\frac{-u(X)-1}{u(X)-1} - u(X)}{X} = 0$$

Or

$$\left(\frac{d}{dX}u(X)\right)Xu(X) - \left(\frac{d}{dX}u(X)\right)X + u(X)^2 + 1 = 0$$

Or

$$X(u(X) - 1) \left(\frac{d}{dX}u(X)\right) + u(X)^2 + 1 = 0$$

Which is now solved as separable in $u(X)$. Which is now solved in $u(X)$. In canonical form the ODE is

$$\begin{aligned} u' &= F(X, u) \\ &= f(X)g(u) \\ &= -\frac{u^2 + 1}{X(u - 1)} \end{aligned}$$

Where $f(X) = -\frac{1}{X}$ and $g(u) = \frac{u^2+1}{u-1}$. Integrating both sides gives

$$\frac{1}{\frac{u^2+1}{u-1}} du = -\frac{1}{X} dX$$

$$\int \frac{1}{\frac{u^2+1}{u-1}} du = \int -\frac{1}{X} dX$$

$$\frac{\ln(u^2 + 1)}{2} - \arctan(u) = -\ln(X) + c_2$$

The solution is

$$\frac{\ln(u(X)^2 + 1)}{2} - \arctan(u(X)) + \ln(X) - c_2 = 0$$

Now u in the above solution is replaced back by Y using $u = \frac{Y}{X}$ which results in the solution

$$\frac{\ln\left(\frac{Y(X)^2}{X^2} + 1\right)}{2} - \arctan\left(\frac{Y(X)}{X}\right) + \ln(X) - c_2 = 0$$

Using the solution for $Y(X)$

$$\frac{\ln\left(\frac{Y(X)^2}{X^2} + 1\right)}{2} - \arctan\left(\frac{Y(X)}{X}\right) + \ln(X) - c_2 = 0$$

And replacing back terms in the above solution using

$$Y = y + y_0$$

$$X = x + x_0$$

Or

$$Y = y - 5$$

$$X = 1 + x$$

Then the solution in y becomes

$$\frac{\ln\left(\frac{(y+5)^2}{(x-1)^2} + 1\right)}{2} - \arctan\left(\frac{y+5}{x-1}\right) + \ln(x-1) - c_2 = 0$$

Summary

The solution(s) found are the following

$$\frac{\ln\left(\frac{(y+5)^2}{(x-1)^2} + 1\right)}{2} - \arctan\left(\frac{y+5}{x-1}\right) + \ln(x-1) - c_2 = 0 \quad (1)$$




Figure 158: Slope field plot

Verification of solutions

$$\frac{\ln\left(\frac{(y+5)^2}{(x-1)^2} + 1\right)}{2} - \arctan\left(\frac{y+5}{x-1}\right) + \ln(x-1) - c_2 = 0$$

Verified OK.

5.11.2 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = -\frac{x+y+4}{-x+y+6}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \quad (\text{1E})$$

$$\eta = xb_2 + yb_3 + b_1 \quad (\text{2E})$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$\begin{aligned} b_2 - \frac{(x+y+4)(b_3 - a_2)}{-x+y+6} - \frac{(x+y+4)^2 a_3}{(-x+y+6)^2} \\ - \left(-\frac{1}{-x+y+6} - \frac{x+y+4}{(-x+y+6)^2} \right) (xa_2 + ya_3 + a_1) \\ - \left(-\frac{1}{-x+y+6} + \frac{x+y+4}{(-x+y+6)^2} \right) (xb_2 + yb_3 + b_1) = 0 \end{aligned} \quad (\text{5E})$$

Putting the above in normal form gives

$$\frac{x^2 a_2 + x^2 a_3 + x^2 b_2 - x^2 b_3 - 2xy a_2 + 2xy a_3 + 2xy b_2 + 2xy b_3 - y^2 a_2 - y^2 a_3 - y^2 b_2 + y^2 b_3 - 12xa_2 + 8xa_3 - 12xb_2 + 8xb_3 - 10ya_2 + 10ya_3 - 10yb_2 + 10yb_3 - 10a_1 + 10a_2 - 10a_3 - 10b_1 + 10b_2 - 10b_3}{(x - y + 6)^3} = 0$$

Setting the numerator to zero gives

$$\begin{aligned} -x^2 a_2 - x^2 a_3 - x^2 b_2 + x^2 b_3 + 2xy a_2 - 2xy a_3 - 2xy b_2 - 2xy b_3 + y^2 a_2 \\ + y^2 a_3 + y^2 b_2 - y^2 b_3 + 12xa_2 - 8xa_3 - 2xb_1 - 10xb_2 - 2xb_3 + 2ya_1 + 10ya_2 \\ + 2ya_3 + 12yb_2 - 8yb_3 + 10a_1 + 24a_2 - 16a_3 + 2b_1 + 36b_2 - 24b_3 = 0 \end{aligned} \quad (\text{6E})$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2\}$$

The above PDE (6E) now becomes

$$\begin{aligned} & -a_2v_1^2 + 2a_2v_1v_2 + a_2v_2^2 - a_3v_1^2 - 2a_3v_1v_2 + a_3v_2^2 - b_2v_1^2 - 2b_2v_1v_2 + b_2v_2^2 \\ & + b_3v_1^2 - 2b_3v_1v_2 - b_3v_2^2 + 2a_1v_2 + 12a_2v_1 + 10a_2v_2 - 8a_3v_1 + 2a_3v_2 - 2b_1v_1 \\ & - 10b_2v_1 + 12b_2v_2 - 2b_3v_1 - 8b_3v_2 + 10a_1 + 24a_2 - 16a_3 + 2b_1 + 36b_2 - 24b_3 = 0 \end{aligned} \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2\}$$

Equation (7E) now becomes

$$\begin{aligned} & (-a_2 - a_3 - b_2 + b_3)v_1^2 + (2a_2 - 2a_3 - 2b_2 - 2b_3)v_1v_2 \\ & + (12a_2 - 8a_3 - 2b_1 - 10b_2 - 2b_3)v_1 + (a_2 + a_3 + b_2 - b_3)v_2^2 \\ & + (2a_1 + 10a_2 + 2a_3 + 12b_2 - 8b_3)v_2 + 10a_1 + 24a_2 - 16a_3 + 2b_1 + 36b_2 - 24b_3 = 0 \end{aligned} \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} -a_2 - a_3 - b_2 + b_3 &= 0 \\ a_2 + a_3 + b_2 - b_3 &= 0 \\ 2a_2 - 2a_3 - 2b_2 - 2b_3 &= 0 \\ 2a_1 + 10a_2 + 2a_3 + 12b_2 - 8b_3 &= 0 \\ 12a_2 - 8a_3 - 2b_1 - 10b_2 - 2b_3 &= 0 \\ 10a_1 + 24a_2 - 16a_3 + 2b_1 + 36b_2 - 24b_3 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= -5b_2 - b_3 \\ a_2 &= b_3 \\ a_3 &= -b_2 \\ b_1 &= -b_2 + 5b_3 \\ b_2 &= b_2 \\ b_3 &= b_3 \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned}\xi &= x - 1 \\ \eta &= y + 5\end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned}\eta &= \eta - \omega(x, y) \xi \\ &= y + 5 - \left(-\frac{x + y + 4}{-x + y + 6} \right) (x - 1) \\ &= \frac{-x^2 - y^2 + 2x - 10y - 26}{x - y - 6} \\ \xi &= 0\end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y} \right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{-x^2 - y^2 + 2x - 10y - 26}{x - y - 6}} dy\end{aligned}$$

Which results in

$$S = \frac{\ln(x^2 + y^2 - 2x + 10y + 26)}{2} + \frac{2(1 - x) \arctan\left(\frac{2y+10}{2x-2}\right)}{2x - 2}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y) S_y}{R_x + \omega(x, y) R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -\frac{x + y + 4}{-x + y + 6}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{x + y + 4}{x^2 + y^2 - 2x + 10y + 26} \\ S_y &= \frac{-x + y + 6}{x^2 + y^2 - 2x + 10y + 26} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \tag{2A}$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \tag{4}$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{\ln(y^2 + x^2 + 10y - 2x + 26)}{2} - \arctan\left(\frac{y + 5}{x - 1}\right) = c_1$$

Which simplifies to

$$\frac{\ln(y^2 + x^2 + 10y - 2x + 26)}{2} - \arctan\left(\frac{y + 5}{x - 1}\right) = c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -\frac{x+y+4}{-x+y+6}$ 	$R = x$ $S = \frac{\ln(x^2 + y^2 - 2x + 1)}{2}$	$\frac{dS}{dR} = 0$ 

Summary

The solution(s) found are the following

$$\frac{\ln(y^2 + x^2 + 10y - 2x + 26)}{2} - \arctan\left(\frac{y + 5}{x - 1}\right) = c_1 \quad (1)$$




Figure 159: Slope field plot

Verification of solutions

$$\frac{\ln(y^2 + x^2 + 10y - 2x + 26)}{2} - \arctan\left(\frac{y + 5}{x - 1}\right) = c_1$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying homogeneous C  
trying homogeneous types:  
trying homogeneous D  
<- homogeneous successful  
<- homogeneous successful`
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 31

```
dsolve(diff(y(x),x)=(x+y(x)+4)/(x-y(x)-6),y(x), singsol=all)
```

$$y(x) = -5 - \tan(\text{RootOf}(2_Z + \ln(\sec(_Z)^2) + 2\ln(x-1) + 2c_1))(x-1)$$

✓ Solution by Mathematica

Time used: 0.063 (sec). Leaf size: 58

```
DSolve[y'[x]==(x+y[x]+4)/(x-y[x]-6),y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve}\left[2 \arctan\left(\frac{y(x) + x + 4}{y(x) - x + 6}\right) + \log\left(\frac{x^2 + y(x)^2 + 10y(x) - 2x + 26}{2(x-1)^2}\right) + 2 \log(x-1) + c_1 = 0, y(x)\right]$$

5.12 problem 4(b)

5.12.1 Solving as first order ode lie symmetry calculated ode 748

Internal problem ID [6214]

Internal file name [OUTPUT/5462_Sunday_June_05_2022_03_39_25_PM_7414833/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.7. Homogeneous Equations. Page 28

Problem number: 4(b).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**first_order_ode_lie_symmetry_calculated**"

Maple gives the following as the ode type

```
[[_homogeneous, `class C`], _rational, [_Abel, `2nd type`, `class A`]]
```

$$y' - \frac{x + y + 4}{x + y - 6} = 0$$

5.12.1 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = \frac{x + y + 4}{x + y - 6}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \quad (\text{1E})$$

$$\eta = xb_2 + yb_3 + b_1 \quad (\text{2E})$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$\begin{aligned} b_2 + \frac{(x+y+4)(b_3-a_2)}{x+y-6} - \frac{(x+y+4)^2 a_3}{(x+y-6)^2} \\ - \left(\frac{1}{x+y-6} - \frac{x+y+4}{(x+y-6)^2} \right) (xa_2 + ya_3 + a_1) \\ - \left(\frac{1}{x+y-6} - \frac{x+y+4}{(x+y-6)^2} \right) (xb_2 + yb_3 + b_1) = 0 \end{aligned} \quad (5E)$$

Putting the above in normal form gives

$$\frac{x^2 a_2 + x^2 a_3 - x^2 b_2 - x^2 b_3 + 2xy a_2 + 2xy a_3 - 2xy b_2 - 2xy b_3 + y^2 a_2 + y^2 a_3 - y^2 b_2 - y^2 b_3 - 12xa_2 + 8xa_3 - 12yb_2 + 8yb_3}{(x+y-6)^2} = 0$$

Setting the numerator to zero gives

$$\begin{aligned} -x^2 a_2 - x^2 a_3 + x^2 b_2 + x^2 b_3 - 2xy a_2 - 2xy a_3 + 2xy b_2 + 2xy b_3 - y^2 a_2 \\ - y^2 a_3 + y^2 b_2 + y^2 b_3 + 12xa_2 - 8xa_3 - 2xb_2 - 2xb_3 + 2ya_2 + 2ya_3 \\ - 12yb_2 + 8yb_3 + 10a_1 + 24a_2 - 16a_3 + 10b_1 + 36b_2 - 24b_3 = 0 \end{aligned} \quad (6E)$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2\}$$

The above PDE (6E) now becomes

$$\begin{aligned} -a_2 v_1^2 - 2a_2 v_1 v_2 - a_2 v_2^2 - a_3 v_1^2 - 2a_3 v_1 v_2 - a_3 v_2^2 + b_2 v_1^2 + 2b_2 v_1 v_2 \\ + b_2 v_2^2 + b_3 v_1^2 + 2b_3 v_1 v_2 + b_3 v_2^2 + 12a_2 v_1 + 2a_2 v_2 - 8a_3 v_1 + 2a_3 v_2 - 2b_2 v_1 \\ - 12b_2 v_2 - 2b_3 v_1 + 8b_3 v_2 + 10a_1 + 24a_2 - 16a_3 + 10b_1 + 36b_2 - 24b_3 = 0 \end{aligned} \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2\}$$

Equation (7E) now becomes

$$\begin{aligned} &(-a_2 - a_3 + b_2 + b_3) v_1^2 + (-2a_2 - 2a_3 + 2b_2 + 2b_3) v_1 v_2 \\ &+ (12a_2 - 8a_3 - 2b_2 - 2b_3) v_1 + (-a_2 - a_3 + b_2 + b_3) v_2^2 \\ &+ (2a_2 + 2a_3 - 12b_2 + 8b_3) v_2 + 10a_1 + 24a_2 - 16a_3 + 10b_1 + 36b_2 - 24b_3 = 0 \end{aligned} \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} -2a_2 - 2a_3 + 2b_2 + 2b_3 &= 0 \\ -a_2 - a_3 + b_2 + b_3 &= 0 \\ 2a_2 + 2a_3 - 12b_2 + 8b_3 &= 0 \\ 12a_2 - 8a_3 - 2b_2 - 2b_3 &= 0 \\ 10a_1 + 24a_2 - 16a_3 + 10b_1 + 36b_2 - 24b_3 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= -2b_3 - b_1 \\ a_2 &= b_3 \\ a_3 &= b_3 \\ b_1 &= b_1 \\ b_2 &= b_3 \\ b_3 &= b_3 \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned} \xi &= -1 \\ \eta &= 1 \end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned} \eta &= \eta - \omega(x, y) \xi \\ &= 1 - \left(\frac{x + y + 4}{x + y - 6} \right) (-1) \\ &= \frac{2x + 2y - 2}{x + y - 6} \\ \xi &= 0 \end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{2x+2y-2}{x+y-6}} dy \end{aligned}$$

Which results in

$$S = \frac{y}{2} - \frac{5 \ln(x+y-1)}{2}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{x+y+4}{x+y-6}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= -\frac{5}{2x+2y-2} \\ S_y &= \frac{x+y-6}{2x+2y-2} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \frac{1}{2} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \frac{1}{2}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \frac{R}{2} + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{y}{2} - \frac{5 \ln(-1 + y + x)}{2} = \frac{x}{2} + c_1$$



Which simplifies to

$$\frac{y}{2} - \frac{5 \ln(-1 + y + x)}{2} = \frac{x}{2} + c_1$$

Which gives

$$y = -5 \operatorname{LambertW} \left(-\frac{e^{-\frac{2x}{5} - \frac{2c_1}{5} + \frac{1}{5}}}{5} \right) - x + 1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{x+y+4}{x+y-6}$ 	$R = x$ $S = \frac{y}{2} - \frac{5 \ln(x+y-1)}{2}$	$\frac{dS}{dR} = \frac{1}{2}$ 

Summary

The solution(s) found are the following

$$y = -5 \text{LambertW} \left(-\frac{e^{-\frac{2x}{5} - \frac{2c_1}{5} + \frac{1}{5}}}{5} \right) - x + 1 \quad (1)$$




Figure 160: Slope field plot

Verification of solutions

$$y = -5 \operatorname{LambertW} \left(-\frac{e^{-\frac{2x}{5} - \frac{2c_1}{5} + \frac{1}{5}}}{5} \right) - x + 1$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying homogeneous C  
1st order, trying the canonical coordinates of the invariance group  
<- 1st order, canonical coordinates successful  
<- homogeneous successful`
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 21

```
dsolve(diff(y(x),x)=(x+y(x)+4)/(x+y(x)-6),y(x), singsol=all)
```

$$y(x) = -x - 5 \operatorname{LambertW}\left(-\frac{c_1 e^{-\frac{2x}{5} + \frac{1}{5}}}{5}\right) + 1$$

✓ Solution by Mathematica

Time used: 4.043 (sec). Leaf size: 35

```
DSolve[y'[x]==(x+y[x]+4)/(x+y[x]-6),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -5W\left(-e^{-\frac{2x}{5}-1+c_1}\right) - x + 1$$
$$y(x) \rightarrow 1 - x$$

5.13 problem 4(c)

5.13.1 Solving as homogeneousTypeMapleC ode 756

5.13.2 Solving as first order ode lie symmetry calculated ode 759

Internal problem ID [6215]

Internal file name [OUTPUT/5463_Sunday_June_05_2022_03_39_28_PM_94865919/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.7. Homogeneous Equations. Page 28

Problem number: 4(c).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"homogeneousTypeMapleC", "first_order_ode_lie_symmetry_calculated"**

Maple gives the following as the ode type

```
[[_homogeneous, `class C`], _rational, [_Abel, `2nd type`, `class A`]]
```

$$-2y + (y - 1)y' = -2x$$

5.13.1 Solving as homogeneousTypeMapleC ode

Let $Y = y + y_0$ and $X = x + x_0$ then the above is transformed to new ode in $Y(X)$

$$\frac{d}{dX}Y(X) = \frac{-2X - 2x_0 + 2Y(X) + 2y_0}{Y(X) + y_0 - 1}$$

Solving for possible values of x_0 and y_0 which makes the above ode a homogeneous ode results in

$$x_0 = 1$$

$$y_0 = 1$$

Using these values now it is possible to easily solve for $Y(X)$. The above ode now becomes

$$\frac{d}{dX}Y(X) = \frac{-2X + 2Y(X)}{Y(X)}$$

In canonical form, the ODE is

$$\begin{aligned} Y' &= F(X, Y) \\ &= \frac{-2X + 2Y}{Y} \end{aligned} \quad (1)$$

An ode of the form $Y' = \frac{M(X, Y)}{N(X, Y)}$ is called homogeneous if the functions $M(X, Y)$ and $N(X, Y)$ are both homogeneous functions and of the same order. Recall that a function $f(X, Y)$ is homogeneous of order n if

$$f(t^n X, t^n Y) = t^n f(X, Y)$$

In this case, it can be seen that both $M = -2X + 2Y$ and $N = Y$ are both homogeneous and of the same order $n = 1$. Therefore this is a homogeneous ode. Since this ode is homogeneous, it is converted to separable ODE using the substitution $u = \frac{Y}{X}$, or $Y = uX$. Hence

$$\frac{dY}{dX} = \frac{du}{dX}X + u$$

Applying the transformation $Y = uX$ to the above ODE in (1) gives

$$\begin{aligned} \frac{du}{dX}X + u &= -\frac{2}{u} + 2 \\ \frac{du}{dX} &= \frac{-\frac{2}{u(X)} + 2 - u(X)}{X} \end{aligned}$$

Or

$$\frac{d}{dX}u(X) - \frac{-\frac{2}{u(X)} + 2 - u(X)}{X} = 0$$

Or

$$\left(\frac{d}{dX}u(X) \right) u(X) X + u(X)^2 - 2u(X) + 2 = 0$$

Which is now solved as separable in $u(X)$. Which is now solved in $u(X)$. In canonical form the ODE is

$$\begin{aligned} u' &= F(X, u) \\ &= f(X)g(u) \\ &= -\frac{u^2 - 2u + 2}{uX} \end{aligned}$$

Where $f(X) = -\frac{1}{X}$ and $g(u) = \frac{u^2-2u+2}{u}$. Integrating both sides gives

$$\frac{1}{\frac{u^2-2u+2}{u}} du = -\frac{1}{X} dX$$

$$\int \frac{1}{\frac{u^2-2u+2}{u}} du = \int -\frac{1}{X} dX$$

$$\frac{\ln(u^2 - 2u + 2)}{2} + \arctan(u - 1) = -\ln(X) + c_2$$

The solution is

$$\frac{\ln(u(X)^2 - 2u(X) + 2)}{2} + \arctan(u(X) - 1) + \ln(X) - c_2 = 0$$

Now u in the above solution is replaced back by Y using $u = \frac{Y}{X}$ which results in the solution

$$\frac{\ln\left(\frac{Y(X)^2}{X^2} - \frac{2Y(X)}{X} + 2\right)}{2} - \arctan\left(-\frac{Y(X)}{X} + 1\right) + \ln(X) - c_2 = 0$$

Using the solution for $Y(X)$

$$\frac{\ln\left(\frac{Y(X)^2}{X^2} - \frac{2Y(X)}{X} + 2\right)}{2} - \arctan\left(-\frac{Y(X)}{X} + 1\right) + \ln(X) - c_2 = 0$$

And replacing back terms in the above solution using

$$Y = y + y_0$$

$$X = x + x_0$$

Or

$$Y = 1 + y$$

$$X = 1 + x$$

Then the solution in y becomes

$$\frac{\ln\left(\frac{(y-1)^2}{(x-1)^2} - \frac{2(y-1)}{x-1} + 2\right)}{2} + \arctan\left(\frac{y-1}{x-1} - 1\right) + \ln(x-1) - c_2 = 0$$

Summary

The solution(s) found are the following

$$\frac{\ln\left(\frac{(y-1)^2}{(x-1)^2} - \frac{2(y-1)}{x-1} + 2\right)}{2} + \arctan\left(\frac{y-1}{x-1} - 1\right) + \ln(x-1) - c_2 = 0 \quad (1)$$




Figure 161: Slope field plot

Verification of solutions

$$\frac{\ln\left(\frac{(y-1)^2}{(x-1)^2} - \frac{2(y-1)}{x-1} + 2\right)}{2} + \arctan\left(\frac{y-1}{x-1} - 1\right) + \ln(x-1) - c_2 = 0$$

Verified OK.

5.13.2 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = \frac{-2x + 2y}{y - 1}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \quad (\text{1E})$$

$$\eta = xb_2 + yb_3 + b_1 \quad (\text{2E})$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$\begin{aligned} b_2 + \frac{2(-x+y)(b_3-a_2)}{y-1} - \frac{4(-x+y)^2 a_3}{(y-1)^2} + \frac{2xa_2 + 2ya_3 + 2a_1}{y-1} \\ - \left(\frac{2}{y-1} - \frac{2(-x+y)}{(y-1)^2} \right) (xb_2 + yb_3 + b_1) = 0 \end{aligned} \quad (\text{5E})$$

Putting the above in normal form gives

$$\frac{4x^2a_3 + 2x^2b_2 - 4xya_2 - 8xya_3 + 4xyb_3 + 2y^2a_2 + 2y^2a_3 - y^2b_2 - 2y^2b_3 + 4xa_2 + 2xb_1 - 2xb_2 - 2xb_3 - 2a_1 + 2b_1 + b_2}{(y-1)^2} = 0$$

Setting the numerator to zero gives

$$\begin{aligned} -4x^2a_3 - 2x^2b_2 + 4xya_2 + 8xya_3 - 4xyb_3 - 2y^2a_2 - 2y^2a_3 + y^2b_2 + 2y^2b_3 \\ - 4xa_2 - 2xb_1 + 2xb_2 + 2xb_3 + 2ya_1 + 2ya_2 - 2ya_3 - 2yb_2 - 2a_1 + 2b_1 + b_2 = 0 \end{aligned} \quad (\text{6E})$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2\}$$

The above PDE (6E) now becomes

$$\begin{aligned}
 &4a_2v_1v_2 - 2a_2v_2^2 - 4a_3v_1^2 + 8a_3v_1v_2 - 2a_3v_2^2 - 2b_2v_1^2 + b_2v_2^2 - 4b_3v_1v_2 + 2b_3v_2^2 \\
 &+ 2a_1v_2 - 4a_2v_1 + 2a_2v_2 - 2a_3v_2 - 2b_1v_1 + 2b_2v_1 - 2b_2v_2 + 2b_3v_1 - 2a_1 + 2b_1 + b_2 \\
 &= 0
 \end{aligned} \tag{7E}$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2\}$$

Equation (7E) now becomes

$$\begin{aligned}
 &(-4a_3 - 2b_2)v_1^2 + (4a_2 + 8a_3 - 4b_3)v_1v_2 + (-4a_2 - 2b_1 + 2b_2 + 2b_3)v_1 \\
 &+ (-2a_2 - 2a_3 + b_2 + 2b_3)v_2^2 + (2a_1 + 2a_2 - 2a_3 - 2b_2)v_2 - 2a_1 + 2b_1 + b_2 = 0
 \end{aligned} \tag{8E}$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned}
 -4a_3 - 2b_2 &= 0 \\
 -2a_1 + 2b_1 + b_2 &= 0 \\
 4a_2 + 8a_3 - 4b_3 &= 0 \\
 2a_1 + 2a_2 - 2a_3 - 2b_2 &= 0 \\
 -4a_2 - 2b_1 + 2b_2 + 2b_3 &= 0 \\
 -2a_2 - 2a_3 + b_2 + 2b_3 &= 0
 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned}
 a_1 &= a_3 - b_3 \\
 a_2 &= -2a_3 + b_3 \\
 a_3 &= a_3 \\
 b_1 &= 2a_3 - b_3 \\
 b_2 &= -2a_3 \\
 b_3 &= b_3
 \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned}
 \xi &= x - 1 \\
 \eta &= y - 1
 \end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned}\eta &= \eta - \omega(x, y) \xi \\ &= y - 1 - \left(\frac{-2x + 2y}{y - 1} \right) (x - 1) \\ &= \frac{2x^2 - 2xy + y^2 - 2x + 1}{y - 1} \\ \xi &= 0\end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y} \right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{2x^2 - 2xy + y^2 - 2x + 1}{y - 1}} dy\end{aligned}$$

Which results in

$$S = \frac{\ln(2x^2 - 2xy + y^2 - 2x + 1)}{2} + \frac{2(x - 1) \arctan\left(\frac{-2x + 2y}{2x - 2}\right)}{2x - 2}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y) S_y}{R_x + \omega(x, y) R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{-2x + 2y}{y - 1}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{-2y + 2x}{2x^2 + (-2y - 2)x + y^2 + 1} \\ S_y &= \frac{y - 1}{2x^2 + (-2y - 2)x + y^2 + 1} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \quad (4)$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{\ln(2x^2 + (-2y - 2)x + y^2 + 1)}{2} - \arctan\left(\frac{x - y}{x - 1}\right) = c_1$$

Which simplifies to

$$\frac{\ln(2x^2 + (-2y - 2)x + y^2 + 1)}{2} - \arctan\left(\frac{x - y}{x - 1}\right) = c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{-2x+2y}{y-1}$ 	$R = x$ $S = \frac{\ln(2x^2 + (-2y - 2)x + y^2 + 1)}{2}$	$\frac{dS}{dR} = 0$ 

Summary

The solution(s) found are the following

$$\frac{\ln(2x^2 + (-2y - 2)x + y^2 + 1)}{2} - \arctan\left(\frac{x - y}{x - 1}\right) = c_1 \quad (1)$$




Figure 162: Slope field plot

Verification of solutions

$$\frac{\ln(2x^2 + (-2y - 2)x + y^2 + 1)}{2} - \arctan\left(\frac{x - y}{x - 1}\right) = c_1$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying homogeneous C  
trying homogeneous types:  
trying homogeneous D  
<- homogeneous successful  
<- homogeneous successful`
```

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 31

```
dsolve((2*x-2*y(x))+(y(x)-1)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = -\tan(\text{RootOf}(-2_Z + \ln(\sec(_Z)^2) + 2\ln(x-1) + 2c_1))(x-1) + x$$

✓ Solution by Mathematica

Time used: 0.057 (sec). Leaf size: 60

```
DSolve[(2*x-2*y[x])+(y[x]-1)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve}\left[2\arctan\left(\frac{y(x)-2x+1}{y(x)-1}\right) + \log\left(\frac{2x^2-2xy(x)+y(x)^2-2x+1}{2(x-1)^2}\right) + 2\log(x-1) + c_1 = 0, y(x)\right]$$

5.14 problem 4(d)

5.14.1 Solving as homogeneousTypeMapleC ode 767

5.14.2 Solving as first order ode lie symmetry calculated ode 771

Internal problem ID [6216]

Internal file name [OUTPUT/5464_Sunday_June_05_2022_03_39_50_PM_24835712/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.7. Homogeneous Equations. Page 28

Problem number: 4(d).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"homogeneousTypeMapleC", "first_order_ode_lie_symmetry_calculated"**

Maple gives the following as the ode type

```
[[_homogeneous, `class C`], _rational, [_Abel, `2nd type`, `class A`]]
```

$$y' - \frac{-1 + y + x}{x + 4y + 2} = 0$$

5.14.1 Solving as homogeneousTypeMapleC ode

Let $Y = y + y_0$ and $X = x + x_0$ then the above is transformed to new ode in $Y(X)$

$$\frac{d}{dX}Y(X) = \frac{-1 + Y(X) + y_0 + X + x_0}{X + x_0 + 4Y(X) + 4y_0 + 2}$$

Solving for possible values of x_0 and y_0 which makes the above ode a homogeneous ode results in

$$x_0 = 2$$

$$y_0 = -1$$

Using these values now it is possible to easily solve for $Y(X)$. The above ode now becomes

$$\frac{d}{dX}Y(X) = \frac{Y(X) + X}{X + 4Y(X)}$$

In canonical form, the ODE is

$$\begin{aligned} Y' &= F(X, Y) \\ &= \frac{Y + X}{X + 4Y} \end{aligned} \quad (1)$$

An ode of the form $Y' = \frac{M(X, Y)}{N(X, Y)}$ is called homogeneous if the functions $M(X, Y)$ and $N(X, Y)$ are both homogeneous functions and of the same order. Recall that a function $f(X, Y)$ is homogeneous of order n if

$$f(t^n X, t^n Y) = t^n f(X, Y)$$

In this case, it can be seen that both $M = Y + X$ and $N = X + 4Y$ are both homogeneous and of the same order $n = 1$. Therefore this is a homogeneous ode. Since this ode is homogeneous, it is converted to separable ODE using the substitution $u = \frac{Y}{X}$, or $Y = uX$. Hence

$$\frac{dY}{dX} = \frac{du}{dX}X + u$$

Applying the transformation $Y = uX$ to the above ODE in (1) gives

$$\begin{aligned} \frac{du}{dX}X + u &= \frac{u + 1}{4u + 1} \\ \frac{du}{dX} &= \frac{\frac{u(X)+1}{4u(X)+1} - u(X)}{X} \end{aligned}$$

Or

$$\frac{d}{dX}u(X) - \frac{\frac{u(X)+1}{4u(X)+1} - u(X)}{X} = 0$$

Or

$$4 \left(\frac{d}{dX}u(X) \right) Xu(X) + \left(\frac{d}{dX}u(X) \right) X + 4u(X)^2 - 1 = 0$$

Or

$$-1 + X(4u(X) + 1) \left(\frac{d}{dX}u(X) \right) + 4u(X)^2 = 0$$

Which is now solved as separable in $u(X)$. Which is now solved in $u(X)$. In canonical form the ODE is

$$\begin{aligned} u' &= F(X, u) \\ &= f(X)g(u) \\ &= -\frac{4u^2 - 1}{X(4u + 1)} \end{aligned}$$

Where $f(X) = -\frac{1}{X}$ and $g(u) = \frac{4u^2-1}{4u+1}$. Integrating both sides gives

$$\begin{aligned}\frac{1}{\frac{4u^2-1}{4u+1}} du &= -\frac{1}{X} dX \\ \int \frac{1}{\frac{4u^2-1}{4u+1}} du &= \int -\frac{1}{X} dX \\ \frac{3 \ln(2u-1)}{4} + \frac{\ln(2u+1)}{4} &= -\ln(X) + c_2\end{aligned}$$

The above can be written as

$$\begin{aligned}\frac{3 \ln(2u-1) + \ln(2u+1)}{4} &= -\ln(X) + c_2 \\ 3 \ln(2u-1) + \ln(2u+1) &= (4)(-\ln(X) + c_2) \\ &= -4 \ln(X) + 4c_2\end{aligned}$$

Raising both side to exponential gives

$$e^{3 \ln(2u-1) + \ln(2u+1)} = e^{-4 \ln(X) + 4c_2}$$

Which simplifies to

$$\begin{aligned}(2u-1)^3 (2u+1) &= \frac{4c_2}{X^4} \\ &= \frac{c_3}{X^4}\end{aligned}$$

Which simplifies to

$$u(X) = \frac{\text{RootOf}(-Z^4 - 2_Z Z^3 - \frac{c_3 e^{4c_2}}{X^4} + 2_Z Z - 1)}{2}$$

Now u in the above solution is replaced back by Y using $u = \frac{Y}{X}$ which results in the solution

$$Y(X) = \frac{X \text{RootOf}(-Z^4 X^4 - 2_Z Z^3 X^4 - c_3 e^{4c_2} + 2_Z Z X^4 - X^4)}{2}$$

Using the solution for $Y(X)$

$$Y(X) = \frac{X \text{RootOf}(-Z^4 X^4 - 2_Z Z^3 X^4 - c_3 e^{4c_2} + 2_Z Z X^4 - X^4)}{2}$$

And replacing back terms in the above solution using

$$Y = y + y_0$$

$$X = x + x_0$$

Or

$$Y = y - 1$$

$$X = x + 2$$

Then the solution in y becomes

$$1 + y = \frac{(-2 + x) \text{RootOf}((x^4 - 8x^3 + 24x^2 - 32x + 16)_Z^4 + (-2x^4 + 16x^3 - 48x^2 + 64x - 32)_Z^3 + \dots)}{2}$$

Summary

The solution(s) found are the following

$$1 + y = \frac{(-2 + x) \text{RootOf}((x^4 - 8x^3 + 24x^2 - 32x + 16)_Z^4 + (-2x^4 + 16x^3 - 48x^2 + 64x - 32)_Z^3 + (2x^4 - \dots)}{2} \quad (1)$$




Figure 163: Slope field plot

Verification of solutions

$1 + y$

$$= \frac{(-2 + x) \text{RootOf}((x^4 - 8x^3 + 24x^2 - 32x + 16)Z^4 + (-2x^4 + 16x^3 - 48x^2 + 64x - 32)Z^3 + (2x^4 - 16x^3 + 48x^2 - 64x + 32)Z^2 + (-2x^4 + 16x^3 - 48x^2 + 64x - 32)Z + (2x^4 - 16x^3 + 48x^2 - 64x + 32))}{2}$$

Verified OK.

5.14.2 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = \frac{x + y - 1}{x + 4y + 2}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \quad (\text{1E})$$

$$\eta = xb_2 + yb_3 + b_1 \quad (\text{2E})$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$b_2 + \frac{(x + y - 1)(b_3 - a_2)}{x + 4y + 2} - \frac{(x + y - 1)^2 a_3}{(x + 4y + 2)^2}$$

$$- \left(\frac{1}{x + 4y + 2} - \frac{x + y - 1}{(x + 4y + 2)^2} \right) (xa_2 + ya_3 + a_1)$$

$$- \left(\frac{1}{x + 4y + 2} - \frac{4(x + y - 1)}{(x + 4y + 2)^2} \right) (xb_2 + yb_3 + b_1) = 0 \quad (\text{5E})$$

Putting the above in normal form gives

$$\frac{x^2 a_2 + x^2 a_3 - 4x^2 b_2 - x^2 b_3 + 8xy a_2 + 2xy a_3 - 8xy b_2 - 8xy b_3 + 4y^2 a_2 + 4y^2 a_3 - 16y^2 b_2 - 4y^2 b_3 + 4xa_1 - 4yb_1}{(x + 4y + 2)^3} = 0$$

Setting the numerator to zero gives

$$\begin{aligned}
 & -x^2a_2 - x^2a_3 + 4x^2b_2 + x^2b_3 - 8xya_2 - 2xya_3 + 8xyb_2 + 8xyb_3 - 4y^2a_2 \\
 & - 4y^2a_3 + 16y^2b_2 + 4y^2b_3 - 4xa_2 + 2xa_3 + 3xb_1 - 2xb_2 + xb_3 - 3ya_1 \\
 & + 2ya_2 - ya_3 + 16yb_2 - 8yb_3 - 3a_1 + 2a_2 - a_3 - 6b_1 + 4b_2 - 2b_3 = 0
 \end{aligned} \tag{6E}$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2\}$$

The above PDE (6E) now becomes

$$\begin{aligned}
 & -a_2v_1^2 - 8a_2v_1v_2 - 4a_2v_2^2 - a_3v_1^2 - 2a_3v_1v_2 - 4a_3v_2^2 + 4b_2v_1^2 + 8b_2v_1v_2 + 16b_2v_2^2 \\
 & + b_3v_1^2 + 8b_3v_1v_2 + 4b_3v_2^2 - 3a_1v_2 - 4a_2v_1 + 2a_2v_2 + 2a_3v_1 - a_3v_2 + 3b_1v_1 \\
 & - 2b_2v_1 + 16b_2v_2 + b_3v_1 - 8b_3v_2 - 3a_1 + 2a_2 - a_3 - 6b_1 + 4b_2 - 2b_3 = 0
 \end{aligned} \tag{7E}$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2\}$$

Equation (7E) now becomes

$$\begin{aligned}
 & (-a_2 - a_3 + 4b_2 + b_3) v_1^2 + (-8a_2 - 2a_3 + 8b_2 + 8b_3) v_1v_2 \\
 & + (-4a_2 + 2a_3 + 3b_1 - 2b_2 + b_3) v_1 + (-4a_2 - 4a_3 + 16b_2 + 4b_3) v_2^2 \\
 & + (-3a_1 + 2a_2 - a_3 + 16b_2 - 8b_3) v_2 - 3a_1 + 2a_2 - a_3 - 6b_1 + 4b_2 - 2b_3 = 0
 \end{aligned} \tag{8E}$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned}
 & -8a_2 - 2a_3 + 8b_2 + 8b_3 = 0 \\
 & -4a_2 - 4a_3 + 16b_2 + 4b_3 = 0 \\
 & -a_2 - a_3 + 4b_2 + b_3 = 0 \\
 & -3a_1 + 2a_2 - a_3 + 16b_2 - 8b_3 = 0 \\
 & -4a_2 + 2a_3 + 3b_1 - 2b_2 + b_3 = 0 \\
 & -3a_1 + 2a_2 - a_3 - 6b_1 + 4b_2 - 2b_3 = 0
 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= 4b_2 - 2b_3 \\ a_2 &= b_3 \\ a_3 &= 4b_2 \\ b_1 &= -2b_2 + b_3 \\ b_2 &= b_2 \\ b_3 &= b_3 \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned} \xi &= -2 + x \\ \eta &= 1 + y \end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned} \eta &= \eta - \omega(x, y) \xi \\ &= 1 + y - \left(\frac{x + y - 1}{x + 4y + 2} \right) (-2 + x) \\ &= \frac{-x^2 + 4y^2 + 4x + 8y}{x + 4y + 2} \\ \xi &= 0 \end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y} \right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{-x^2 + 4y^2 + 4x + 8y}{x + 4y + 2}} dy \end{aligned}$$

Which results in

$$S = \frac{3 \ln(2y + 4 - x)}{4} + \frac{\ln(2y + x)}{4}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{x + y - 1}{x + 4y + 2}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{x + y - 1}{(2y + x)(x - 4 - 2y)} \\ S_y &= \frac{-x - 4y - 2}{(2y + x)(x - 4 - 2y)} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \quad (4)$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{3 \ln(2y + 4 - x)}{4} + \frac{\ln(2y + x)}{4} = c_1$$

Which simplifies to

$$\frac{3 \ln(2y + 4 - x)}{4} + \frac{\ln(2y + x)}{4} = c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{x+y-1}{x+4y+2}$ 	$R = x$ $S = \frac{3 \ln(2y + 4 - x)}{4} + \frac{\ln(2y + x)}{4}$	$\frac{dS}{dR} = 0$ 

Summary

The solution(s) found are the following

$$\frac{3 \ln(2y + 4 - x)}{4} + \frac{\ln(2y + x)}{4} = c_1 \tag{1}$$




Figure 164: Slope field plot

Verification of solutions

$$\frac{3 \ln(2y + 4 - x)}{4} + \frac{\ln(2y + x)}{4} = c_1$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying homogeneous C  
trying homogeneous types:  
trying homogeneous D  
<- homogeneous successful  
<- homogeneous successful`
```

✓ Solution by Maple

Time used: 4.563 (sec). Leaf size: 139

```
dsolve(diff(y(x),x)=(x+y(x)-1)/(x+4*y(x)+2),y(x), singsol=all)
```

$$y(x) = \frac{(x-4) \operatorname{RootOf}(_Z^{16} + (2c_1x^4 - 16c_1x^3 + 48c_1x^2 - 64c_1x + 32c_1)_Z^4 - c_1x^4 + 8c_1x^3 - 24c_1x^2 + 32c_1x - 16c_1), _Z^4 - c_1x^4 + 8c_1x^3 - 24c_1x^2 + 32c_1x - 16c_1)}{2 \operatorname{RootOf}(_Z^{16} + (2c_1x^4 - 16c_1x^3 + 48c_1x^2 - 64c_1x + 32c_1)_Z^4 - c_1x^4 + 8c_1x^3 - 24c_1x^2 + 32c_1x - 16c_1), _Z^4 - c_1x^4 + 8c_1x^3 - 24c_1x^2 + 32c_1x - 16c_1)}$$

✓ Solution by Mathematica

Time used: 60.343 (sec). Leaf size: 8141

```
DSolve[y'[x]==(x+y[x]-1)/(x+4*y[x]+2),y[x],x,IncludeSingularSolutions -> True]
```

Too large to display

5.15 problem 4(e)

5.15.1 Solving as linear ode	778
5.15.2 Solving as homogeneousTypeMapleC ode	780
5.15.3 Solving as first order ode lie symmetry lookup ode	784
5.15.4 Solving as exact ode	788
5.15.5 Maple step by step solution	793

Internal problem ID [6217]

Internal file name [OUTPUT/5465_Sunday_June_05_2022_03_39_56_PM_96534507/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.7. Homogeneous Equations. Page 28

Problem number: 4(e).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**linear**", "**homogeneousTypeMapleC**", "**exactWithIntegrationFactor**", "**first_order_ode_lie_symmetry_lookup**"

Maple gives the following as the ode type

[_linear]

$$3y - 4(1 + x)y' = 1 - 2x$$

5.15.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -\frac{3}{4x + 4}$$
$$q(x) = \frac{2x - 1}{4x + 4}$$

Hence the ode is

$$y' - \frac{3y}{4x + 4} = \frac{2x - 1}{4x + 4}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int -\frac{3}{4x+4} dx} \\ &= \frac{1}{(1+x)^{\frac{3}{4}}}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu) \left(\frac{2x-1}{4x+4} \right) \\ \frac{d}{dx} \left(\frac{y}{(1+x)^{\frac{3}{4}}} \right) &= \left(\frac{1}{(1+x)^{\frac{3}{4}}} \right) \left(\frac{2x-1}{4x+4} \right) \\ d \left(\frac{y}{(1+x)^{\frac{3}{4}}} \right) &= \left(\frac{2x-1}{4(1+x)^{\frac{7}{4}}} \right) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}\frac{y}{(1+x)^{\frac{3}{4}}} &= \int \frac{2x-1}{4(1+x)^{\frac{7}{4}}} dx \\ \frac{y}{(1+x)^{\frac{3}{4}}} &= \frac{2x+3}{(1+x)^{\frac{3}{4}}} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = \frac{1}{(1+x)^{\frac{3}{4}}}$ results in

$$y = 2x + 3 + c_1(1+x)^{\frac{3}{4}}$$

Summary

The solution(s) found are the following

$$y = 2x + 3 + c_1(1+x)^{\frac{3}{4}} \tag{1}$$




Figure 165: Slope field plot

Verification of solutions

$$y = 2x + 3 + c_1(1 + x)^{\frac{3}{4}}$$

Verified OK.

5.15.2 Solving as homogeneousTypeMapleC ode

Let $Y = y + y_0$ and $X = x + x_0$ then the above is transformed to new ode in $Y(X)$

$$\frac{d}{dX}Y(X) = \frac{2X + 2x_0 + 3Y(X) + 3y_0 - 1}{4 + 4X + 4x_0}$$

Solving for possible values of x_0 and y_0 which makes the above ode a homogeneous ode results in

$$x_0 = -1$$

$$y_0 = 1$$

Using these values now it is possible to easily solve for $Y(X)$. The above ode now becomes

$$\frac{d}{dX}Y(X) = \frac{2X + 3Y(X)}{4X}$$

In canonical form, the ODE is

$$\begin{aligned} Y' &= F(X, Y) \\ &= \frac{2X + 3Y}{4X} \end{aligned} \tag{1}$$

An ode of the form $Y' = \frac{M(X, Y)}{N(X, Y)}$ is called homogeneous if the functions $M(X, Y)$ and $N(X, Y)$ are both homogeneous functions and of the same order. Recall that a function $f(X, Y)$ is homogeneous of order n if

$$f(t^n X, t^n Y) = t^n f(X, Y)$$

In this case, it can be seen that both $M = 2X + 3Y$ and $N = 4X$ are both homogeneous and of the same order $n = 1$. Therefore this is a homogeneous ode. Since this ode is homogeneous, it is converted to separable ODE using the substitution $u = \frac{Y}{X}$, or $Y = uX$. Hence

$$\frac{dY}{dX} = \frac{du}{dX}X + u$$

Applying the transformation $Y = uX$ to the above ODE in (1) gives

$$\begin{aligned} \frac{du}{dX}X + u &= \frac{1}{2} + \frac{3u}{4} \\ \frac{du}{dX} &= \frac{\frac{1}{2} - \frac{u(X)}{4}}{X} \end{aligned}$$

Or

$$\frac{d}{dX}u(X) - \frac{\frac{1}{2} - \frac{u(X)}{4}}{X} = 0$$

Or

$$4\left(\frac{d}{dX}u(X)\right)X + u(X) - 2 = 0$$

Which is now solved as separable in $u(X)$. Which is now solved in $u(X)$. In canonical form the ODE is

$$\begin{aligned} u' &= F(X, u) \\ &= f(X)g(u) \\ &= \frac{-\frac{u}{4} + \frac{1}{2}}{X} \end{aligned}$$

Where $f(X) = \frac{1}{X}$ and $g(u) = -\frac{u}{4} + \frac{1}{2}$. Integrating both sides gives

$$\begin{aligned}\frac{1}{-\frac{u}{4} + \frac{1}{2}} du &= \frac{1}{X} dX \\ \int \frac{1}{-\frac{u}{4} + \frac{1}{2}} du &= \int \frac{1}{X} dX \\ -4 \ln(u - 2) &= \ln(X) + c_2\end{aligned}$$

Raising both side to exponential gives

$$\frac{1}{(u - 2)^4} = e^{\ln(X) + c_2}$$

Which simplifies to

$$\frac{1}{(u - 2)^4} = c_3 X$$

Which simplifies to

$$\frac{1}{(u(X) - 2)^4} = c_3 X e^{c_2}$$

The solution is

$$\frac{1}{(u(X) - 2)^4} = c_3 X e^{c_2}$$

Now u in the above solution is replaced back by Y using $u = \frac{Y}{X}$ which results in the solution

$$\frac{1}{\left(\frac{Y(X)}{X} - 2\right)^4} = c_3 X e^{c_2}$$

Using the solution for $Y(X)$

$$\frac{X^4}{(Y(X) - 2X)^4} = c_3 X e^{c_2}$$

And replacing back terms in the above solution using

$$\begin{aligned}Y &= y + y_0 \\ X &= x + x_0\end{aligned}$$

Or

$$\begin{aligned} Y &= 1 + y \\ X &= x - 1 \end{aligned}$$

Then the solution in y becomes

$$\frac{(1+x)^4}{(y-3-2x)^4} = c_3(1+x)e^{c_2}$$

Summary

The solution(s) found are the following

$$\frac{(1+x)^4}{(y-3-2x)^4} = c_3(1+x)e^{c_2} \quad (1)$$




Figure 166: Slope field plot

Verification of solutions

$$\frac{(1+x)^4}{(y-3-2x)^4} = c_3(1+x)e^{c_2}$$

Verified OK.

5.15.3 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{2x + 3y - 1}{4x + 4}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 116: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= (1 + x)^{\frac{3}{4}}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{(1 + x)^{\frac{3}{4}}} dy\end{aligned}$$

Which results in

$$S = \frac{y}{(1 + x)^{\frac{3}{4}}}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y}\tag{2}$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{2x + 3y - 1}{4x + 4}$$

Evaluating all the partial derivatives gives

$$\begin{aligned}R_x &= 1 \\R_y &= 0 \\S_x &= -\frac{3y}{4(1+x)^{\frac{7}{4}}} \\S_y &= \frac{1}{(1+x)^{\frac{3}{4}}}\end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \frac{2x - 1}{4(1+x)^{\frac{7}{4}}} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \frac{2R - 1}{4(1+R)^{\frac{7}{4}}}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \frac{2R + 3}{(1+R)^{\frac{3}{4}}} + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{y}{(1+x)^{\frac{3}{4}}} = \frac{2x + 3}{(1+x)^{\frac{3}{4}}} + c_1$$



Which simplifies to

$$\frac{y}{(1+x)^{\frac{3}{4}}} = \frac{2x + 3}{(1+x)^{\frac{3}{4}}} + c_1$$

Which gives

$$y = 2x + 3 + c_1(1+x)^{\frac{3}{4}}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{2x+3y-1}{4x+4}$ 	$R = x$ $S = \frac{y}{(1+x)^{\frac{3}{4}}}$	$\frac{dS}{dR} = \frac{2R-1}{4(1+R)^{\frac{7}{4}}}$ 

Summary

The solution(s) found are the following

$$y = 2x + 3 + c_1(1+x)^{\frac{3}{4}} \quad (1)$$




Figure 167: Slope field plot

Verification of solutions

$$y = 2x + 3 + c_1(1 + x)^{\frac{3}{4}}$$

Verified OK.

5.15.4 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}(-4 - 4x) dy &= (-2x - 3y + 1) dx \\ (2x + 3y - 1) dx + (-4 - 4x) dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= 2x + 3y - 1 \\ N(x, y) &= -4 - 4x\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(2x + 3y - 1) \\ &= 3\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(-4 - 4x) \\ &= -4\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= \frac{1}{-4 - 4x} ((3) - (-4)) \\ &= -\frac{7}{4x + 4} \end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int A dx} \\ &= e^{\int -\frac{7}{4x+4} dx} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^{-\frac{7 \ln(1+x)}{4}} \\ &= \frac{1}{(1+x)^{\frac{7}{4}}} \end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} for now so not to confuse them with the original M and N .

$$\begin{aligned} \bar{M} &= \mu M \\ &= \frac{1}{(1+x)^{\frac{7}{4}}} (2x + 3y - 1) \\ &= \frac{2x + 3y - 1}{(1+x)^{\frac{7}{4}}} \end{aligned}$$

And

$$\begin{aligned} \bar{N} &= \mu N \\ &= \frac{1}{(1+x)^{\frac{7}{4}}} (-4 - 4x) \\ &= -\frac{4}{(1+x)^{\frac{3}{4}}} \end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned} \bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ \left(\frac{2x + 3y - 1}{(1+x)^{\frac{7}{4}}} \right) + \left(-\frac{4}{(1+x)^{\frac{3}{4}}} \right) \frac{dy}{dx} &= 0 \end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \quad (1)$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned} \int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int \frac{2x + 3y - 1}{(1+x)^{\frac{7}{4}}} dx \\ \phi &= \frac{8x + 12 - 4y}{(1+x)^{\frac{3}{4}}} + f(y) \end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = -\frac{4}{(1+x)^{\frac{3}{4}}} + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = -\frac{4}{(1+x)^{\frac{3}{4}}}$. Therefore equation (4) becomes

$$-\frac{4}{(1+x)^{\frac{3}{4}}} = -\frac{4}{(1+x)^{\frac{3}{4}}} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = \frac{8x + 12 - 4y}{(1+x)^{\frac{3}{4}}} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \frac{8x + 12 - 4y}{(1+x)^{\frac{3}{4}}}$$

The solution becomes

$$y = -\frac{c_1(1+x)^{\frac{3}{4}}}{4} + 2x + 3$$

Summary

The solution(s) found are the following

$$y = -\frac{c_1(1+x)^{\frac{3}{4}}}{4} + 2x + 3 \quad (1)$$




Figure 168: Slope field plot

Verification of solutions

$$y = -\frac{c_1(1+x)^{\frac{3}{4}}}{4} + 2x + 3$$

Verified OK.

5.15.5 Maple step by step solution

Let's solve

$$3y - 4(1+x)y' = 1 - 2x$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = \frac{3y}{4(1+x)} + \frac{2x-1}{4(1+x)}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' - \frac{3y}{4(1+x)} = \frac{2x-1}{4(1+x)}$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) \left(y' - \frac{3y}{4(1+x)} \right) = \frac{\mu(x)(2x-1)}{4(1+x)}$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x)y)$

$$\mu(x) \left(y' - \frac{3y}{4(1+x)} \right) = \mu'(x)y + \mu(x)y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = -\frac{3\mu(x)}{4(1+x)}$$

- Solve to find the integrating factor

$$\mu(x) = \frac{1}{(1+x)^{\frac{3}{4}}}$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x)y) \right) dx = \int \frac{\mu(x)(2x-1)}{4(1+x)} dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x)y = \int \frac{\mu(x)(2x-1)}{4(1+x)} dx + c_1$$

- Solve for y

$$y = \frac{\int \frac{\mu(x)(2x-1)}{4(1+x)} dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = \frac{1}{(1+x)^{\frac{3}{4}}}$

$$y = (1+x)^{\frac{3}{4}} \left(\int \frac{2x-1}{4(1+x)^{\frac{7}{4}}} dx + c_1 \right)$$

- Evaluate the integrals on the rhs

$$y = (1+x)^{\frac{3}{4}} \left(\frac{2x+3}{(1+x)^{\frac{3}{4}}} + c_1 \right)$$

- Simplify

$$y = 2x + 3 + c_1(1+x)^{\frac{3}{4}}$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 16

```
dsolve((2*x+3*y(x)-1)-4*(x+1)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = (x+1)^{\frac{3}{4}} c_1 + 2x + 3$$

✓ Solution by Mathematica

Time used: 0.065 (sec). Leaf size: 20

```
DSolve[(2*x+3*y[x]-1)-4*(x+1)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2x + c_1(x+1)^{3/4} + 3$$

5.16 problem 5(a)

5.16.1 Solving as first order ode lie symmetry lookup ode	795
5.16.2 Solving as bernoulli ode	799
5.16.3 Solving as exact ode	803

Internal problem ID [6218]

Internal file name [OUTPUT/5466_Sunday_June_05_2022_03_39_57_PM_28256205/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.7. Homogeneous Equations. Page 28

Problem number: 5(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "bernoulli", "exactWithIntegrationFactor", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

```
[[_homogeneous, `class G`], _rational, _Bernoulli]
```

$$y' - \frac{-xy^2 + 1}{2yx^2} = 0$$

5.16.1 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = -\frac{y^2x - 1}{2yx^2}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2\xi_y - \omega_x\xi - \omega_y\eta = 0 \tag{A}$$

The type of this ode is known. It is of type Bernoulli. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 119: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= \frac{1}{yx}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{1}{yx}} dy \end{aligned}$$

Which results in

$$S = \frac{y^2 x}{2}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -\frac{y^2 x - 1}{2y x^2}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{y^2}{2} \\ S_y &= xy \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \frac{1}{2x} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \frac{1}{2R}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \frac{\ln(R)}{2} + c_1 \quad (4)$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{xy^2}{2} = \frac{\ln(x)}{2} + c_1$$

Which simplifies to

$$\frac{xy^2}{2} = \frac{\ln(x)}{2} + c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -\frac{y^2x-1}{2yx^2}$ 	$R = x$ $S = \frac{y^2x}{2}$	$\frac{dS}{dR} = \frac{1}{2R}$ 

Summary

The solution(s) found are the following

$$\frac{xy^2}{2} = \frac{\ln(x)}{2} + c_1 \quad (1)$$




Figure 169: Slope field plot

Verification of solutions

$$\frac{xy^2}{2} = \frac{\ln(x)}{2} + c_1$$

Verified OK.

5.16.2 Solving as bernoulli ode

In canonical form, the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= -\frac{y^2x - 1}{2yx^2} \end{aligned}$$

This is a Bernoulli ODE.

$$y' = -\frac{1}{2x}y + \frac{1}{2x^2} \frac{1}{y} \quad (1)$$

The standard Bernoulli ODE has the form

$$y' = f_0(x)y + f_1(x)y^n \quad (2)$$

The first step is to divide the above equation by y^n which gives

$$\frac{y'}{y^n} = f_0(x)y^{1-n} + f_1(x) \quad (3)$$

The next step is use the substitution $w = y^{1-n}$ in equation (3) which generates a new ODE in $w(x)$ which will be linear and can be easily solved using an integrating factor. Backsubstitution then gives the solution $y(x)$ which is what we want.

This method is now applied to the ODE at hand. Comparing the ODE (1) With (2) Shows that

$$\begin{aligned}f_0(x) &= -\frac{1}{2x} \\f_1(x) &= \frac{1}{2x^2} \\n &= -1\end{aligned}$$

Dividing both sides of ODE (1) by $y^n = \frac{1}{y}$ gives

$$y'y = -\frac{y^2}{2x} + \frac{1}{2x^2} \quad (4)$$

Let

$$\begin{aligned}w &= y^{1-n} \\&= y^2\end{aligned} \quad (5)$$

Taking derivative of equation (5) w.r.t x gives

$$w' = 2yy' \quad (6)$$

Substituting equations (5) and (6) into equation (4) gives

$$\begin{aligned}\frac{w'(x)}{2} &= -\frac{w(x)}{2x} + \frac{1}{2x^2} \\w' &= -\frac{w}{x} + \frac{1}{x^2}\end{aligned} \quad (7)$$

The above now is a linear ODE in $w(x)$ which is now solved.

Entering Linear first order ODE solver. In canonical form a linear first order is

$$w'(x) + p(x)w(x) = q(x)$$

Where here

$$\begin{aligned}p(x) &= \frac{1}{x} \\q(x) &= \frac{1}{x^2}\end{aligned}$$

Hence the ode is

$$w'(x) + \frac{w(x)}{x} = \frac{1}{x^2}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int \frac{1}{x} dx} \\ &= x\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu w) &= (\mu) \left(\frac{1}{x^2} \right) \\ \frac{d}{dx}(xw) &= (x) \left(\frac{1}{x^2} \right) \\ d(xw) &= \frac{1}{x} dx\end{aligned}$$

Integrating gives

$$\begin{aligned}xw &= \int \frac{1}{x} dx \\ xw &= \ln(x) + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = x$ results in

$$w(x) = \frac{\ln(x)}{x} + \frac{c_1}{x}$$

which simplifies to

$$w(x) = \frac{\ln(x) + c_1}{x}$$

Replacing w in the above by y^2 using equation (5) gives the final solution.

$$y^2 = \frac{\ln(x) + c_1}{x}$$

Solving for y gives

$$\begin{aligned}y(x) &= \frac{\sqrt{x(\ln(x) + c_1)}}{x} \\ y(x) &= -\frac{\sqrt{x(\ln(x) + c_1)}}{x}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{\sqrt{x (\ln(x) + c_1)}}{x} \quad (1)$$

$$y = -\frac{\sqrt{x (\ln(x) + c_1)}}{x} \quad (2)$$




Figure 170: Slope field plot

Verification of solutions

$$y = \frac{\sqrt{x (\ln(x) + c_1)}}{x}$$

Verified OK.

$$y = -\frac{\sqrt{x (\ln(x) + c_1)}}{x}$$

Verified OK.

5.16.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} dy &= \left(\frac{-y^2 x + 1}{2y x^2} \right) dx \\ \left(-\frac{-y^2 x + 1}{2y x^2} \right) dx + dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$M(x, y) = -\frac{-y^2x + 1}{2yx^2}$$
$$N(x, y) = 1$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\frac{\partial M}{\partial y} = \frac{\partial}{\partial y} \left(-\frac{-y^2x + 1}{2yx^2} \right)$$
$$= \frac{y^2x + 1}{2y^2x^2}$$

And

$$\frac{\partial N}{\partial x} = \frac{\partial}{\partial x}(1)$$
$$= 0$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$A = \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right)$$
$$= 1 \left(\left(\frac{1}{x} + \frac{-y^2x + 1}{2y^2x^2} \right) - (0) \right)$$
$$= \frac{y^2x + 1}{2y^2x^2}$$

Since A depends on y , it can not be used to obtain an integrating factor. We will now try a second method to find an integrating factor. Let

$$B = \frac{1}{M} \left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right)$$
$$= \frac{2yx^2}{y^2x - 1} \left((0) - \left(\frac{1}{x} + \frac{-y^2x + 1}{2y^2x^2} \right) \right)$$
$$= \frac{-y^2x - 1}{y(y^2x - 1)}$$

Since B depends on x , it can not be used to obtain an integrating factor. We will now try a third method to find an integrating factor. Let

$$R = \frac{\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y}}{xM - yN}$$

R is now checked to see if it is a function of only $t = xy$. Therefore

$$\begin{aligned} R &= \frac{\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y}}{xM - yN} \\ &= \frac{(0) - \left(\frac{1}{x} + \frac{-y^2x+1}{2y^2x^2}\right)}{x\left(-\frac{-y^2x+1}{2yx^2}\right) - y(1)} \\ &= \frac{1}{yx} \end{aligned}$$

Replacing all powers of terms xy by t gives

$$R = \frac{1}{t}$$

Since R depends on t only, then it can be used to find an integrating factor. Let the integrating factor be μ then

$$\begin{aligned} \mu &= e^{\int R dt} \\ &= e^{\int \left(\frac{1}{t}\right) dt} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^{\ln(t)} \\ &= t \end{aligned}$$

Now t is replaced back with xy giving

$$\mu = xy$$

Multiplying M and N by this integrating factor gives new M and new N which are called \overline{M} and \overline{N} so not to confuse them with the original M and N

$$\begin{aligned} \overline{M} &= \mu M \\ &= xy \left(-\frac{-y^2x+1}{2yx^2} \right) \\ &= \frac{y^2x-1}{2x} \end{aligned}$$

And

$$\begin{aligned}\bar{N} &= \mu N \\ &= xy(1) \\ &= xy\end{aligned}$$

A modified ODE is now obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned}\bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ \left(\frac{y^2x - 1}{2x} \right) + (xy) \frac{dy}{dx} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int \frac{y^2x - 1}{2x} dx \\ \phi &= \frac{y^2x}{2} - \frac{\ln(x)}{2} + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = xy + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = xy$. Therefore equation (4) becomes

$$xy = xy + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = \frac{y^2 x}{2} - \frac{\ln(x)}{2} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \frac{y^2 x}{2} - \frac{\ln(x)}{2}$$

Summary

The solution(s) found are the following

$$\frac{xy^2}{2} - \frac{\ln(x)}{2} = c_1 \tag{1}$$




Figure 171: Slope field plot

Verification of solutions

$$\frac{xy^2}{2} - \frac{\ln(x)}{2} = c_1$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
<- Bernoulli successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 32

```
dsolve(diff(y(x),x)=(1-x*y(x)^2)/(2*x^2*y(x)),y(x), singsol=all)
```

$$y(x) = \frac{\sqrt{x(\ln(x) + c_1)}}{x}$$
$$y(x) = -\frac{\sqrt{x(\ln(x) + c_1)}}{x}$$

✓ Solution by Mathematica

Time used: 0.184 (sec). Leaf size: 40

```
DSolve[y'[x]==(1-x*y[x]^2)/(2*x^2*y[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\sqrt{\log(x) + c_1}}{\sqrt{x}}$$
$$y(x) \rightarrow \frac{\sqrt{\log(x) + c_1}}{\sqrt{x}}$$

5.17 problem 5(b)

5.17.1 Solving as first order ode lie symmetry lookup ode	809
5.17.2 Solving as bernoulli ode	813
5.17.3 Solving as exact ode	817

Internal problem ID [6219]

Internal file name [OUTPUT/5467_Sunday_June_05_2022_03_40_00_PM_44970004/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.7. Homogeneous Equations. Page 28

Problem number: 5(b).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "bernoulli", "exactWithIntegrationFactor", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

```
[[_homogeneous, `class G`], _rational, _Bernoulli]
```

$$y' - \frac{2 + 3xy^2}{4yx^2} = 0$$

5.17.1 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{3y^2x + 2}{4yx^2}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2\xi_y - \omega_x\xi - \omega_y\eta = 0 \tag{A}$$

The type of this ode is known. It is of type Bernoulli. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 121: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= \frac{x^{\frac{3}{2}}}{y}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{x^{\frac{3}{2}}}{y}} dy \end{aligned}$$

Which results in

$$S = \frac{y^2}{2x^{\frac{3}{2}}}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{3y^2x + 2}{4yx^2}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= -\frac{3y^2}{4x^{\frac{5}{2}}} \\ S_y &= \frac{y}{x^{\frac{3}{2}}} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \frac{1}{2x^{\frac{7}{2}}} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \frac{1}{2R^{\frac{7}{2}}}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = -\frac{1}{5R^{\frac{5}{2}}} + c_1 \quad (4)$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{y^2}{2x^{\frac{3}{2}}} = -\frac{1}{5x^{\frac{5}{2}}} + c_1$$

Which simplifies to

$$\frac{y^2}{2x^{\frac{3}{2}}} = -\frac{1}{5x^{\frac{5}{2}}} + c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{3y^2x+2}{4yx^2}$ 	$R = x$ $S = \frac{y^2}{2x^{\frac{3}{2}}}$	$\frac{dS}{dR} = \frac{1}{2R^{\frac{7}{2}}}$ 

Summary

The solution(s) found are the following

$$\frac{y^2}{2x^{\frac{3}{2}}} = -\frac{1}{5x^{\frac{5}{2}}} + c_1 \quad (1)$$




Figure 172: Slope field plot

Verification of solutions

$$\frac{y^2}{2x^{\frac{3}{2}}} = -\frac{1}{5x^{\frac{5}{2}}} + c_1$$

Verified OK.

5.17.2 Solving as bernoulli ode

In canonical form, the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= \frac{3y^2x + 2}{4y x^2} \end{aligned}$$

This is a Bernoulli ODE.

$$y' = \frac{3}{4x}y + \frac{1}{2x^2} \frac{1}{y} \tag{1}$$

The standard Bernoulli ODE has the form

$$y' = f_0(x)y + f_1(x)y^n \tag{2}$$

The first step is to divide the above equation by y^n which gives

$$\frac{y'}{y^n} = f_0(x)y^{1-n} + f_1(x) \tag{3}$$

The next step is use the substitution $w = y^{1-n}$ in equation (3) which generates a new ODE in $w(x)$ which will be linear and can be easily solved using an integrating factor. Backsubstitution then gives the solution $y(x)$ which is what we want.

This method is now applied to the ODE at hand. Comparing the ODE (1) With (2) Shows that

$$\begin{aligned}f_0(x) &= \frac{3}{4x} \\f_1(x) &= \frac{1}{2x^2} \\n &= -1\end{aligned}$$

Dividing both sides of ODE (1) by $y^n = \frac{1}{y}$ gives

$$y'y = \frac{3y^2}{4x} + \frac{1}{2x^2} \quad (4)$$

Let

$$\begin{aligned}w &= y^{1-n} \\&= y^2\end{aligned} \quad (5)$$

Taking derivative of equation (5) w.r.t x gives

$$w' = 2yy' \quad (6)$$

Substituting equations (5) and (6) into equation (4) gives

$$\begin{aligned}\frac{w'(x)}{2} &= \frac{3w(x)}{4x} + \frac{1}{2x^2} \\w' &= \frac{3w}{2x} + \frac{1}{x^2}\end{aligned} \quad (7)$$

The above now is a linear ODE in $w(x)$ which is now solved.

Entering Linear first order ODE solver. In canonical form a linear first order is

$$w'(x) + p(x)w(x) = q(x)$$

Where here

$$\begin{aligned}p(x) &= -\frac{3}{2x} \\q(x) &= \frac{1}{x^2}\end{aligned}$$

Hence the ode is

$$w'(x) - \frac{3w(x)}{2x} = \frac{1}{x^2}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int -\frac{3}{2x} dx} \\ &= \frac{1}{x^{\frac{3}{2}}}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu w) &= (\mu) \left(\frac{1}{x^2} \right) \\ \frac{d}{dx} \left(\frac{w}{x^{\frac{3}{2}}} \right) &= \left(\frac{1}{x^{\frac{3}{2}}} \right) \left(\frac{1}{x^2} \right) \\ d \left(\frac{w}{x^{\frac{3}{2}}} \right) &= \frac{1}{x^{\frac{7}{2}}} dx\end{aligned}$$

Integrating gives

$$\begin{aligned}\frac{w}{x^{\frac{3}{2}}} &= \int \frac{1}{x^{\frac{7}{2}}} dx \\ \frac{w}{x^{\frac{3}{2}}} &= -\frac{2}{5x^{\frac{5}{2}}} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = \frac{1}{x^{\frac{3}{2}}}$ results in

$$w(x) = -\frac{2}{5x} + c_1 x^{\frac{3}{2}}$$

Replacing w in the above by y^2 using equation (5) gives the final solution.

$$y^2 = -\frac{2}{5x} + c_1 x^{\frac{3}{2}}$$

Solving for y gives

$$\begin{aligned}y(x) &= \frac{\sqrt{5} \sqrt{5x^{\frac{7}{2}} c_1 - 2x}}{5x} \\ y(x) &= -\frac{\sqrt{5} \sqrt{5x^{\frac{7}{2}} c_1 - 2x}}{5x}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{\sqrt{5} \sqrt{5x^{\frac{7}{2}}c_1 - 2x}}{5x} \quad (1)$$

$$y = -\frac{\sqrt{5} \sqrt{5x^{\frac{7}{2}}c_1 - 2x}}{5x} \quad (2)$$




Figure 173: Slope field plot

Verification of solutions

$$y = \frac{\sqrt{5} \sqrt{5x^{\frac{7}{2}}c_1 - 2x}}{5x}$$

Verified OK.

$$y = -\frac{\sqrt{5} \sqrt{5x^{\frac{7}{2}}c_1 - 2x}}{5x}$$

Verified OK.

5.17.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} (4y x^2) dy &= (3y^2 x + 2) dx \\ (-3y^2 x - 2) dx + (4y x^2) dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -3y^2 x - 2 \\ N(x, y) &= 4y x^2 \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-3y^2x - 2) \\ &= -6xy\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(4yx^2) \\ &= 8xy\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned}A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= \frac{1}{4yx^2} ((-6xy) - (8xy)) \\ &= -\frac{7}{2x}\end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int A dx} \\ &= e^{\int -\frac{7}{2x} dx}\end{aligned}$$

The result of integrating gives

$$\begin{aligned}\mu &= e^{-\frac{7 \ln(x)}{2}} \\ &= \frac{1}{x^{\frac{7}{2}}}\end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} for now so not to confuse them with the original M and N .

$$\begin{aligned}\bar{M} &= \mu M \\ &= \frac{1}{x^{\frac{7}{2}}}(-3y^2x - 2) \\ &= \frac{-3y^2x - 2}{x^{\frac{7}{2}}}\end{aligned}$$

And

$$\begin{aligned}\bar{N} &= \mu N \\ &= \frac{1}{x^{\frac{7}{2}}}(4y x^2) \\ &= \frac{4y}{x^{\frac{3}{2}}}\end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned}\bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ \left(\frac{-3y^2x - 2}{x^{\frac{7}{2}}} \right) + \left(\frac{4y}{x^{\frac{3}{2}}} \right) \frac{dy}{dx} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int \frac{-3y^2x - 2}{x^{\frac{7}{2}}} dx \\ \phi &= \frac{2y^2x + \frac{4}{5}}{x^{\frac{5}{2}}} + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = \frac{4y}{x^{\frac{3}{2}}} + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{4y}{x^{\frac{3}{2}}}$. Therefore equation (4) becomes

$$\frac{4y}{x^{\frac{3}{2}}} = \frac{4y}{x^{\frac{3}{2}}} + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = \frac{2y^2x + \frac{4}{5}}{x^{\frac{5}{2}}} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \frac{2y^2x + \frac{4}{5}}{x^{\frac{5}{2}}}$$

Summary

The solution(s) found are the following

$$\frac{2xy^2 + \frac{4}{5}}{x^{\frac{5}{2}}} = c_1 \tag{1}$$




Figure 174: Slope field plot

Verification of solutions

$$\frac{2xy^2 + \frac{4}{5}}{x^{\frac{5}{2}}} = c_1$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
<- Bernoulli successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 47

```
dsolve(diff(y(x),x)=(2+3*x*y(x)^2)/(4*x^2*y(x)),y(x), singsol=all)
```

$$y(x) = -\frac{\sqrt{5} \sqrt{5x^{\frac{7}{2}}c_1 - 2x}}{5x}$$
$$y(x) = \frac{\sqrt{5} \sqrt{5x^{\frac{7}{2}}c_1 - 2x}}{5x}$$

✓ Solution by Mathematica

Time used: 3.667 (sec). Leaf size: 51

```
DSolve[y'[x]==(2+3*x*y[x]^2)/(4*x^2*y[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sqrt{-\frac{2}{5x} + c_1x^{3/2}}$$
$$y(x) \rightarrow \sqrt{-\frac{2}{5x} + c_1x^{3/2}}$$

5.18 problem 5(c)

5.18.1 Solving as first order ode lie symmetry calculated ode 822

5.18.2 Solving as exact ode 828

Internal problem ID [6220]

Internal file name [OUTPUT/5468_Sunday_June_05_2022_03_40_04_PM_85038005/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.7. Homogeneous Equations. Page 28

Problem number: 5(c).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exactWithIntegrationFactor", "first_order_ode_lie_symmetry_calculated"**

Maple gives the following as the ode type

```
[[_homogeneous, `class G`], _rational, [_Abel, `2nd type`, `class B`]]
```

$$y' - \frac{-xy^2 + y}{x + yx^2} = 0$$

5.18.1 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = -\frac{y(xy - 1)}{x(xy + 1)}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \tag{A}$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \tag{1E}$$

$$\eta = xb_2 + yb_3 + b_1 \tag{2E}$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$\begin{aligned} b_2 - \frac{y(xy-1)(b_3-a_2)}{x(xy+1)} - \frac{y^2(xy-1)^2 a_3}{x^2(xy+1)^2} \\ - \left(-\frac{y^2}{x(xy+1)} + \frac{y(xy-1)}{x^2(xy+1)} + \frac{y^2(xy-1)}{x(xy+1)^2} \right) (xa_2 + ya_3 + a_1) \\ - \left(-\frac{xy-1}{x(xy+1)} - \frac{y}{xy+1} + \frac{y(xy-1)}{(xy+1)^2} \right) (xb_2 + yb_3 + b_1) = 0 \end{aligned} \quad (5E)$$

Putting the above in normal form gives

$$\frac{2x^4y^2b_2 - 2x^2y^4a_3 + x^3y^2b_1 - x^2y^3a_1 + 4x^3yb_2 + 2x^2y^2a_2 + 2x^2y^2b_3 + 4xy^3a_3 + 2x^2yb_1 + 2xy^2a_1 - xb_1 - ya_1}{x^2(xy+1)^2} = 0$$

Setting the numerator to zero gives

$$\begin{aligned} 2x^4y^2b_2 - 2x^2y^4a_3 + x^3y^2b_1 - x^2y^3a_1 + 4x^3yb_2 + 2x^2y^2a_2 \\ + 2x^2y^2b_3 + 4xy^3a_3 + 2x^2yb_1 + 2xy^2a_1 - xb_1 + ya_1 = 0 \end{aligned} \quad (6E)$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2\}$$

The above PDE (6E) now becomes

$$\begin{aligned} -2a_3v_1^2v_2^4 + 2b_2v_1^4v_2^2 - a_1v_1^2v_2^3 + b_1v_1^3v_2^2 + 2a_2v_1^2v_2^2 + 4a_3v_1v_2^3 \\ + 4b_2v_1^3v_2 + 2b_3v_1^2v_2^2 + 2a_1v_1v_2^2 + 2b_1v_1^2v_2 + a_1v_2 - b_1v_1 = 0 \end{aligned} \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2\}$$

Equation (7E) now becomes

$$2b_2v_1^4v_2^2 + b_1v_1^3v_2^2 + 4b_2v_1^3v_2 - 2a_3v_1^2v_2^4 - a_1v_1^2v_2^3 + (2a_2 + 2b_3)v_1^2v_2^2 + 2b_1v_1^2v_2 + 4a_3v_1v_2^3 + 2a_1v_1v_2^2 - b_1v_1 + a_1v_2 = 0 \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} a_1 &= 0 \\ b_1 &= 0 \\ -a_1 &= 0 \\ 2a_1 &= 0 \\ -2a_3 &= 0 \\ 4a_3 &= 0 \\ -b_1 &= 0 \\ 2b_1 &= 0 \\ 2b_2 &= 0 \\ 4b_2 &= 0 \\ 2a_2 + 2b_3 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= 0 \\ a_2 &= -b_3 \\ a_3 &= 0 \\ b_1 &= 0 \\ b_2 &= 0 \\ b_3 &= b_3 \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned} \xi &= -x \\ \eta &= y \end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned}\eta &= \eta - \omega(x, y) \xi \\ &= y - \left(-\frac{y(xy - 1)}{x(xy + 1)} \right) (-x) \\ &= \frac{2y}{xy + 1} \\ \xi &= 0\end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y} \right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{2y}{xy+1}} dy\end{aligned}$$

Which results in

$$S = \frac{xy}{2} + \frac{\ln(y)}{2}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -\frac{y(xy - 1)}{x(xy + 1)}$$

Evaluating all the partial derivatives gives

$$\begin{aligned}R_x &= 1 \\R_y &= 0 \\S_x &= \frac{y}{2} \\S_y &= \frac{xy + 1}{2y}\end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \frac{1}{2x} \tag{2A}$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \frac{1}{2R}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \frac{\ln(R)}{2} + c_1 \tag{4}$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{xy}{2} + \frac{\ln(y)}{2} = \frac{\ln(x)}{2} + c_1$$



Which simplifies to

$$\frac{xy}{2} + \frac{\ln(y)}{2} = \frac{\ln(x)}{2} + c_1$$

Which gives

$$y = \frac{\text{LambertW}(e^{2c_1}x^2)}{x}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -\frac{y(xy-1)}{x(xy+1)}$ 	$R = x$ $S = \frac{xy}{2} + \frac{\ln(y)}{2}$	$\frac{dS}{dR} = \frac{1}{2R}$ 

Summary

The solution(s) found are the following

$$y = \frac{\text{LambertW}(e^{2c_1} x^2)}{x} \tag{1}$$




Figure 175: Slope field plot

Verification of solutions

$$y = \frac{\text{LambertW}(e^{2c_1 x^2})}{x}$$

Verified OK.

5.18.2 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned}dy &= \left(\frac{-y^2x + y}{yx^2 + x} \right) dx \\ \left(-\frac{-y^2x + y}{yx^2 + x} \right) dx + dy &= 0\end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -\frac{-y^2x + y}{yx^2 + x} \\ N(x, y) &= 1\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(-\frac{-y^2x + y}{yx^2 + x} \right) \\ &= \frac{y^2x^2 + 2xy - 1}{x(xy + 1)^2}\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(1) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned}A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= 1 \left(\left(-\frac{-2xy+1}{yx^2+x} + \frac{(-y^2x+y)x^2}{(yx^2+x)^2} \right) - (0) \right) \\ &= \frac{y^2x^2+2xy-1}{x(xy+1)^2}\end{aligned}$$

Since A depends on y , it can not be used to obtain an integrating factor. We will now try a second method to find an integrating factor. Let

$$\begin{aligned}B &= \frac{1}{M} \left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) \\ &= \frac{x(xy+1)}{y(xy-1)} \left((0) - \left(-\frac{-2xy+1}{yx^2+x} + \frac{(-y^2x+y)x^2}{(yx^2+x)^2} \right) \right) \\ &= \frac{-y^2x^2-2xy+1}{y^3x^2-y}\end{aligned}$$

Since B depends on x , it can not be used to obtain an integrating factor. We will now try a third method to find an integrating factor. Let

$$R = \frac{\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y}}{xM - yN}$$

R is now checked to see if it is a function of only $t = xy$. Therefore

$$\begin{aligned}R &= \frac{\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y}}{xM - yN} \\ &= \frac{(0) - \left(-\frac{-2xy+1}{yx^2+x} + \frac{(-y^2x+y)x^2}{(yx^2+x)^2} \right)}{x \left(-\frac{-y^2x+y}{yx^2+x} \right) - y(1)} \\ &= \frac{y^2x^2+2xy-1}{2xy(xy+1)}\end{aligned}$$

Replacing all powers of terms xy by t gives

$$R = \frac{t^2 + 2t - 1}{2t(t+1)}$$

Since R depends on t only, then it can be used to find an integrating factor. Let the integrating factor be μ then

$$\begin{aligned}\mu &= e^{\int R dt} \\ &= e^{\int \left(\frac{t^2+2t-1}{2t(t+1)}\right) dt}\end{aligned}$$

The result of integrating gives

$$\begin{aligned}\mu &= e^{\frac{t}{2} + \ln(t+1) - \frac{\ln(t)}{2}} \\ &= \frac{(t+1)e^{\frac{t}{2}}}{\sqrt{t}}\end{aligned}$$

Now t is replaced back with xy giving

$$\mu = \frac{(xy+1)e^{\frac{xy}{2}}}{\sqrt{xy}}$$

Multiplying M and N by this integrating factor gives new \bar{M} and new \bar{N} which are called \bar{M} and \bar{N} so not to confuse them with the original M and N

$$\begin{aligned}\bar{M} &= \mu M \\ &= \frac{(xy+1)e^{\frac{xy}{2}}}{\sqrt{xy}} \left(-\frac{y^2x+y}{yx^2+x} \right) \\ &= \frac{e^{\frac{xy}{2}}y(xy-1)}{\sqrt{xy}x}\end{aligned}$$

And

$$\begin{aligned}\bar{N} &= \mu N \\ &= \frac{(xy+1)e^{\frac{xy}{2}}}{\sqrt{xy}}(1) \\ &= \frac{(xy+1)e^{\frac{xy}{2}}}{\sqrt{xy}}\end{aligned}$$

A modified ODE is now obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned}\bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ \left(\frac{e^{\frac{xy}{2}}y(xy-1)}{\sqrt{xy}x} \right) + \left(\frac{(xy+1)e^{\frac{xy}{2}}}{\sqrt{xy}} \right) \frac{dy}{dx} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \quad (1)$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned} \int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int \frac{e^{\frac{xy}{2}} y(xy - 1)}{\sqrt{xy} x} dx \\ \phi &= \frac{2e^{\frac{xy}{2}} y}{\sqrt{xy}} + f(y) \end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\begin{aligned} \frac{\partial \phi}{\partial y} &= \frac{x e^{\frac{xy}{2}} y}{\sqrt{xy}} + \frac{2e^{\frac{xy}{2}}}{\sqrt{xy}} - \frac{e^{\frac{xy}{2}} yx}{(xy)^{\frac{3}{2}}} + f'(y) \\ &= \frac{(xy + 1)e^{\frac{xy}{2}}}{\sqrt{xy}} + f'(y) \end{aligned} \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{(xy+1)e^{\frac{xy}{2}}}{\sqrt{xy}}$. Therefore equation (4) becomes

$$\frac{(xy + 1)e^{\frac{xy}{2}}}{\sqrt{xy}} = \frac{(xy + 1)e^{\frac{xy}{2}}}{\sqrt{xy}} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = \frac{2e^{\frac{xy}{2}} y}{\sqrt{xy}} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \frac{2 e^{\frac{xy}{2}} y}{\sqrt{xy}}$$

The solution becomes

$$y = \frac{c_1^2 x e^{-\text{LambertW}\left(\frac{x^2 c_1^2}{4}\right)}}{4}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1^2 x e^{-\text{LambertW}\left(\frac{x^2 c_1^2}{4}\right)}}{4} \quad (1)$$




Figure 176: Slope field plot

Verification of solutions

$$y = \frac{c_1^2 x e^{-\text{LambertW}\left(\frac{x^2 c_1^2}{4}\right)}}{4}$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying homogeneous G  
<- homogeneous successful`
```

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 17

```
dsolve(diff(y(x),x)=(y(x)-x*y(x)^2)/(x+x^2*y(x)),y(x), singsol=all)
```

$$y(x) = \frac{\text{LambertW}(x^2 e^{-2c_1})}{x}$$

✓ Solution by Mathematica

Time used: 60.444 (sec). Leaf size: 31

```
DSolve[y'[x]==(y[x]-x*y[x]^2)/(x+x^2*y[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{W\left(e^{\frac{1}{2}(-2-9\sqrt[3]{-2c_1})}x^2\right)}{x}$$

5.19 problem 7(a)

5.19.1 Solving as homogeneous ode 835

Internal problem ID [6221]

Internal file name [OUTPUT/5469_Sunday_June_05_2022_03_40_06_PM_28975778/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.7. Homogeneous Equations. Page 28

Problem number: 7(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

```
[[_homogeneous , `class A`], _dAlembert]
```

$$y' - \sin\left(\frac{y}{x}\right) + \cos\left(\frac{y}{x}\right) = 0$$

5.19.1 Solving as homogeneous ode

In canonical form, the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= \sin\left(\frac{y}{x}\right) - \cos\left(\frac{y}{x}\right) \end{aligned} \quad (1)$$

An ode of the form $y' = \frac{M(x,y)}{N(x,y)}$ is called homogeneous if the functions $M(x, y)$ and $N(x, y)$ are both homogeneous functions and of the same order. Recall that a function $f(x, y)$ is homogeneous of order n if

$$f(t^n x, t^n y) = t^n f(x, y)$$

In this case, it can be seen that both $M = \sin\left(\frac{y}{x}\right) - \cos\left(\frac{y}{x}\right)$ and $N = 1$ are both homogeneous and of the same order $n = 0$. Therefore this is a homogeneous ode. Since this ode is homogeneous, it is converted to separable ODE using the substitution $u = \frac{y}{x}$, or $y = ux$. Hence

$$\frac{dy}{dx} = \frac{du}{dx}x + u$$

Applying the transformation $y = ux$ to the above ODE in (1) gives

$$\begin{aligned} \frac{du}{dx}x + u &= \sin(u) - \cos(u) \\ \frac{du}{dx} &= \frac{\sin(u(x)) - \cos(u(x)) - u(x)}{x} \end{aligned}$$

Or

$$u'(x) - \frac{\sin(u(x)) - \cos(u(x)) - u(x)}{x} = 0$$

Or

$$u'(x)x - \sin(u(x)) + \cos(u(x)) + u(x) = 0$$

Which is now solved as separable in $u(x)$. Which is now solved in $u(x)$. In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= \frac{\sin(u) - \cos(u) - u}{x} \end{aligned}$$

Where $f(x) = \frac{1}{x}$ and $g(u) = \sin(u) - \cos(u) - u$. Integrating both sides gives

$$\begin{aligned} \frac{1}{\sin(u) - \cos(u) - u} du &= \frac{1}{x} dx \\ \int \frac{1}{\sin(u) - \cos(u) - u} du &= \int \frac{1}{x} dx \\ \int^u \frac{1}{\sin(_a) - \cos(_a) - _a} d_a &= \ln(x) + c_2 \end{aligned}$$

Which results in

$$\int^u \frac{1}{\sin(_a) - \cos(_a) - _a} d_a = \ln(x) + c_2$$

The solution is

$$\int^{u(x)} \frac{1}{\sin(_a) - \cos(_a) - _a} d_a - \ln(x) - c_2 = 0$$

Now u in the above solution is replaced back by y using $u = \frac{y}{x}$ which results in the solution

$$\int^{\frac{y}{x}} \frac{1}{\sin(_a) - \cos(_a) - _a} d_a - \ln(x) - c_2 = 0$$

Summary

The solution(s) found are the following

$$\int^{\frac{y}{x}} \frac{1}{\sin(_a) - \cos(_a) - _a} d_a - \ln(x) - c_2 = 0 \quad (1)$$




Figure 177: Slope field plot

Verification of solutions

$$\int^{\frac{y}{x}} \frac{1}{\sin(_a) - \cos(_a) - _a} d_a - \ln(x) - c_2 = 0$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying homogeneous D  
<- homogeneous successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 25

```
dsolve(diff(y(x),x)=sin(y(x)/x)-cos(y(x)/x),y(x), singsol=all)
```

$$y(x) = \text{RootOf} \left(\int^{-Z} \frac{1}{-\sin(_a) + \cos(_a) + _a} d_a + \ln(x) + c_1 \right) x$$

✓ Solution by Mathematica

Time used: 0.367 (sec). Leaf size: 36

```
DSolve[y'[x]==Sin[y[x]/x]-Cos[y[x]/x],y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[\int_1^{\frac{y(x)}{x}} \frac{1}{\cos(K[1]) + K[1] - \sin(K[1])} dK[1] = -\log(x) + c_1, y(x) \right]$$

5.20 problem 7(b)

5.20.1 Solving as homogeneous ode 839

Internal problem ID [6222]

Internal file name [OUTPUT/5470_Sunday_June_05_2022_03_40_08_PM_58372649/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.7. Homogeneous Equations. Page 28

Problem number: 7(b).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

```
[[_homogeneous , `class A`], _dAlembert]
```

$$e^{\frac{x}{y}} - \frac{y'y}{x} = 0$$

5.20.1 Solving as homogeneous ode

In canonical form, the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= \frac{x e^{\frac{x}{y}}}{y} \end{aligned} \tag{1}$$

An ode of the form $y' = \frac{M(x,y)}{N(x,y)}$ is called homogeneous if the functions $M(x, y)$ and $N(x, y)$ are both homogeneous functions and of the same order. Recall that a function $f(x, y)$ is homogeneous of order n if

$$f(t^n x, t^n y) = t^n f(x, y)$$

In this case, it can be seen that both $M = x e^{\frac{x}{y}}$ and $N = y$ are both homogeneous and of the same order $n = 1$. Therefore this is a homogeneous ode. Since this ode is homogeneous, it is converted to separable ODE using the substitution $u = \frac{y}{x}$, or $y = ux$. Hence

$$\frac{dy}{dx} = \frac{du}{dx}x + u$$

Applying the transformation $y = ux$ to the above ODE in (1) gives

$$\frac{du}{dx}x + u = \frac{e^{\frac{1}{u}}}{u}$$

$$\frac{du}{dx} = \frac{\frac{e^{\frac{1}{u(x)}}}{u(x)} - u(x)}{x}$$

Or

$$u'(x) - \frac{\frac{e^{\frac{1}{u(x)}}}{u(x)} - u(x)}{x} = 0$$

Or

$$u'(x)u(x)x + u(x)^2 - e^{\frac{1}{u(x)}} = 0$$

Which is now solved as separable in $u(x)$. Which is now solved in $u(x)$. In canonical form the ODE is

$$u' = F(x, u)$$

$$= f(x)g(u)$$

$$= -\frac{u^2 - e^{\frac{1}{u}}}{ux}$$

Where $f(x) = -\frac{1}{x}$ and $g(u) = \frac{u^2 - e^{\frac{1}{u}}}{u}$. Integrating both sides gives

$$\frac{1}{\frac{u^2 - e^{\frac{1}{u}}}{u}} du = -\frac{1}{x} dx$$

$$\int \frac{1}{\frac{u^2 - e^{\frac{1}{u}}}{u}} du = \int -\frac{1}{x} dx$$

$$\int^u \frac{-a}{-a^2 - e^{-\frac{1}{a}}} d_a = -\ln(x) + c_2$$

Which results in

$$\int^u \frac{-a}{-a^2 - e^{-\frac{1}{a}}} d_a = -\ln(x) + c_2$$

The solution is

$$\int^{u(x)} \frac{-a}{-a^2 - e^{-\frac{1}{a}}} d_a + \ln(x) - c_2 = 0$$

Now u in the above solution is replaced back by y using $u = \frac{y}{x}$ which results in the solution

$$\int^{\frac{y}{x}} \frac{-a}{-a^2 - e^{-\frac{1}{a}}} d_a + \ln(x) - c_2 = 0$$

Summary

The solution(s) found are the following

$$-\left(\int^{\frac{y}{x}} \frac{-a}{-a^2 + e^{-\frac{1}{a}}} d_a\right) + \ln(x) - c_2 = 0 \quad (1)$$




Figure 178: Slope field plot

Verification of solutions

$$-\left(\int^{\frac{y}{x}} \frac{-a}{-a^2 + e^{-\frac{1}{a}}} d_a\right) + \ln(x) - c_2 = 0$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying homogeneous D  
<- homogeneous successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 31

```
dsolve(exp(x/y(x))-y(x)/x*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \text{RootOf} \left(- \left(\int^{-Z} \frac{-a}{-a^2 + e^{-\frac{1}{a}}} d_a \right) + \ln(x) + c_1 \right) x$$

✓ Solution by Mathematica

Time used: 0.192 (sec). Leaf size: 41

```
DSolve[Exp[x/y[x]]-y[x]/x*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[\int_1^{\frac{y(x)}{x}} \frac{K[1]}{K[1]^2 - e^{\frac{1}{K[1]}}} dK[1] = -\log(x) + c_1, y(x) \right]$$

5.21 problem 7(c)

5.21.1 Solving as homogeneous ode 843

Internal problem ID [6223]

Internal file name [OUTPUT/5471_Sunday_June_05_2022_03_40_11_PM_7650062/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.7. Homogeneous Equations.

Page 28

Problem number: 7(c).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

```
[[_homogeneous, `class A`], _dAlembert]
```

$$y' - \frac{x^2 - xy}{y^2 \cos\left(\frac{x}{y}\right)} = 0$$

5.21.1 Solving as homogeneous ode

In canonical form, the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= -\frac{x(-x + y)}{y^2 \cos\left(\frac{x}{y}\right)} \end{aligned} \tag{1}$$

An ode of the form $y' = \frac{M(x,y)}{N(x,y)}$ is called homogeneous if the functions $M(x, y)$ and $N(x, y)$ are both homogeneous functions and of the same order. Recall that a function $f(x, y)$ is homogeneous of order n if

$$f(t^n x, t^n y) = t^n f(x, y)$$

In this case, it can be seen that both $M = x(x - y)$ and $N = \cos\left(\frac{x}{y}\right) y^2$ are both homogeneous and of the same order $n = 2$. Therefore this is a homogeneous ode. Since

this ode is homogeneous, it is converted to separable ODE using the substitution $u = \frac{y}{x}$, or $y = ux$. Hence

$$\frac{dy}{dx} = \frac{du}{dx}x + u$$

Applying the transformation $y = ux$ to the above ODE in (1) gives

$$\begin{aligned} \frac{du}{dx}x + u &= \frac{1}{u^2 \cos\left(\frac{1}{u}\right)} - \frac{1}{u \cos\left(\frac{1}{u}\right)} \\ \frac{du}{dx} &= \frac{\frac{1}{u(x)^2 \cos\left(\frac{1}{u(x)}\right)} - \frac{1}{u(x) \cos\left(\frac{1}{u(x)}\right)} - u(x)}{x} \end{aligned}$$

Or

$$u'(x) - \frac{\frac{1}{u(x)^2 \cos\left(\frac{1}{u(x)}\right)} - \frac{1}{u(x) \cos\left(\frac{1}{u(x)}\right)} - u(x)}{x} = 0$$

Or

$$u'(x) u(x)^2 \cos\left(\frac{1}{u(x)}\right) x + u(x)^3 \cos\left(\frac{1}{u(x)}\right) + u(x) - 1 = 0$$

Or

$$u(x)^2 (xu'(x) + u(x)) \cos\left(\frac{1}{u(x)}\right) + u(x) - 1 = 0$$

Which is now solved as separable in $u(x)$. Which is now solved in $u(x)$. In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{(u-1) \sec\left(\frac{1}{u}\right) + u^3}{x u^2} \end{aligned}$$

Where $f(x) = -\frac{1}{x}$ and $g(u) = \frac{(u-1) \sec\left(\frac{1}{u}\right) + u^3}{u^2}$. Integrating both sides gives

$$\begin{aligned} \frac{1}{\frac{(u-1) \sec\left(\frac{1}{u}\right) + u^3}{u^2}} du &= -\frac{1}{x} dx \\ \int \frac{1}{\frac{(u-1) \sec\left(\frac{1}{u}\right) + u^3}{u^2}} du &= \int -\frac{1}{x} dx \\ \int \frac{-a^2}{\left(-a-1\right) \sec\left(\frac{1}{-a}\right) + -a^3} d_{-a} &= -\ln(x) + c_2 \end{aligned}$$

Which results in

$$\int^u \frac{-a^2}{(-a-1)\sec\left(\frac{1}{-a}\right) + -a^3} d_{-a} = -\ln(x) + c_2$$

The solution is

$$\int^{u(x)} \frac{-a^2}{(-a-1)\sec\left(\frac{1}{-a}\right) + -a^3} d_{-a} + \ln(x) - c_2 = 0$$

Now u in the above solution is replaced back by y using $u = \frac{y}{x}$ which results in the solution

$$\int^{\frac{y}{x}} \frac{-a^2}{(-a-1)\sec\left(\frac{1}{-a}\right) + -a^3} d_{-a} + \ln(x) - c_2 = 0$$

Summary

The solution(s) found are the following

$$\int^{\frac{y}{x}} \frac{-a^2 \cos\left(\frac{1}{-a}\right)}{-a^3 \cos\left(\frac{1}{-a}\right) + -a - 1} d_{-a} + \ln(x) - c_2 = 0 \quad (1)$$




Figure 179: Slope field plot

Verification of solutions

$$\int \frac{\frac{y}{x} - a^2 \cos\left(\frac{1}{-a}\right)}{-a^3 \cos\left(\frac{1}{-a}\right) + -a - 1} d_{-a} + \ln(x) - c_2 = 0$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying homogeneous D  
<- homogeneous successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 36

```
dsolve(diff(y(x),x)=(x^2-x*y(x))/(y(x)^2*cos(x/y(x))),y(x), singsol=all)
```

$$y(x) = \text{RootOf} \left(\int^{-z} \frac{-a^2 \cos\left(\frac{1}{-a}\right)}{-a^3 \cos\left(\frac{1}{-a}\right) + -a - 1} d_{-a} + \ln(x) + c_1 \right) x$$

✓ Solution by Mathematica

Time used: 1.114 (sec). Leaf size: 49

```
DSolve[y'[x]==(x^2-x*y[x])/(y[x]^2*Cos[x/y[x]]),y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[\int_1^{\frac{y(x)}{x}} \frac{\cos\left(\frac{1}{K[1]}\right) K[1]^2}{\cos\left(\frac{1}{K[1]}\right) K[1]^3 + K[1] - 1} dK[1] = -\log(x) + c_1, y(x) \right]$$

5.22 problem 7(d)

5.22.1 Solving as homogeneous ode 847

Internal problem ID [6224]

Internal file name [OUTPUT/5472_Sunday_June_05_2022_03_40_15_PM_31766225/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.7. Homogeneous Equations.

Page 28

Problem number: 7(d).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program :

Maple gives the following as the ode type

```
[[_homogeneous , `class A`], _dAlembert]
```

$$y' - \frac{y \tan\left(\frac{y}{x}\right)}{x} = 0$$

5.22.1 Solving as homogeneous ode

In canonical form, the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= \frac{y \tan\left(\frac{y}{x}\right)}{x} \end{aligned} \tag{1}$$

An ode of the form $y' = \frac{M(x,y)}{N(x,y)}$ is called homogeneous if the functions $M(x, y)$ and $N(x, y)$ are both homogeneous functions and of the same order. Recall that a function $f(x, y)$ is homogeneous of order n if

$$f(t^n x, t^n y) = t^n f(x, y)$$

In this case, it can be seen that both $M = y \tan\left(\frac{y}{x}\right)$ and $N = x$ are both homogeneous and of the same order $n = 1$. Therefore this is a homogeneous ode. Since this ode is homogeneous, it is converted to separable ODE using the substitution $u = \frac{y}{x}$, or $y = ux$.

Hence

$$\frac{dy}{dx} = \frac{du}{dx}x + u$$

Applying the transformation $y = ux$ to the above ODE in (1) gives

$$\frac{du}{dx}x + u = u \tan(u)$$

$$\frac{du}{dx} = \frac{u(x) \tan(u(x)) - u(x)}{x}$$

Or

$$u'(x) - \frac{u(x) \tan(u(x)) - u(x)}{x} = 0$$

Or

$$u'(x)x - u(x) \tan(u(x)) + u(x) = 0$$

Which is now solved as separable in $u(x)$. Which is now solved in $u(x)$. In canonical form the ODE is

$$u' = F(x, u)$$

$$= f(x)g(u)$$

$$= \frac{u(\tan(u) - 1)}{x}$$

Where $f(x) = \frac{1}{x}$ and $g(u) = u(\tan(u) - 1)$. Integrating both sides gives

$$\frac{1}{u(\tan(u) - 1)} du = \frac{1}{x} dx$$

$$\int \frac{1}{u(\tan(u) - 1)} du = \int \frac{1}{x} dx$$

$$\int^u \frac{1}{_a(\tan(_a) - 1)} d_a = \ln(x) + c_2$$

Which results in

$$\int^u \frac{1}{_a(\tan(_a) - 1)} d_a = \ln(x) + c_2$$

The solution is

$$\int^{u(x)} \frac{1}{_a(\tan(_a) - 1)} d_a - \ln(x) - c_2 = 0$$

Now u in the above solution is replaced back by y using $u = \frac{y}{x}$ which results in the solution

$$\int^{\frac{y}{x}} \frac{1}{_a(\tan(_a) - 1)} d_a - \ln(x) - c_2 = 0$$

Summary

The solution(s) found are the following

$$\int^{\frac{y}{x}} \frac{1}{-a(\tan(-a) - 1)} d_{-a} - \ln(x) - c_2 = 0 \quad (1)$$




Figure 180: Slope field plot

Verification of solutions

$$\int^{\frac{y}{x}} \frac{1}{-a(\tan(-a) - 1)} d_{-a} - \ln(x) - c_2 = 0$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying homogeneous D  
<- homogeneous successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve(diff(y(x),x)=y(x)/x*tan(y(x)/x),y(x), singsol=all)
```

$$y(x) = \text{RootOf} \left(\ln(x) + c_1 - \left(\int^{-Z} \frac{1}{-a(-1 + \tan(-a))} d_{-a} \right) \right) x$$

✓ Solution by Mathematica

Time used: 1.796 (sec). Leaf size: 33

```
DSolve[y'[x]==y[x]/x*Tan[y[x]/x],y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[\int_1^{\frac{y(x)}{x}} \frac{1}{K[1](\tan(K[1]) - 1)} dK[1] = \log(x) + c_1, y(x) \right]$$

6 Chapter 1. What is a differential equation.

Section 1.8. Integrating Factors. Page 32

6.1	problem 1(a)	852
6.2	problem 1(b)	859
6.3	problem 1(c)	865
6.4	problem 1(d)	872
6.5	problem 1(e)	878
6.6	problem 1(f)	884
6.7	problem 1(g)	892
6.8	problem 1(h)	898
6.9	problem 1(i)	904
6.10	problem 1(j)	910
6.11	problem 1(k)	917
6.12	problem 4	923

6.1 problem 1(a)

6.1.1 Solving as exact ode 852

Internal problem ID [6225]

Internal file name [OUTPUT/5473_Sunday_June_05_2022_03_40_17_PM_80284096/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.8. Integrating Factors. Page 32

Problem number: 1(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exactWithIntegrationFactor"**

Maple gives the following as the ode type

```
[[_homogeneous, `class A`], _rational, _dAlembert]
```

$$(3x^2 - y^2)y' - 2xy = 0$$

6.1.1 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned} (3x^2 - y^2) dy &= (2xy) dx \\ (-2xy) dx + (3x^2 - y^2) dy &= 0 \end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -2xy \\ N(x, y) &= 3x^2 - y^2 \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-2xy) \\ &= -2x \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(3x^2 - y^2) \\ &= 6x \end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= \frac{1}{3x^2 - y^2} ((-2x) - (6x)) \\ &= -\frac{8x}{3x^2 - y^2} \end{aligned}$$

Since A depends on y , it can not be used to obtain an integrating factor. We will now try a second method to find an integrating factor. Let

$$\begin{aligned} B &= \frac{1}{M} \left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) \\ &= -\frac{1}{2yx} ((6x) - (-2x)) \\ &= -\frac{4}{y} \end{aligned}$$

Since B does not depend on x , it can be used to obtain an integrating factor. Let the integrating factor be μ . Then

$$\begin{aligned} \mu &= e^{\int B \, dy} \\ &= e^{\int -\frac{4}{y} \, dy} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^{-4 \ln(y)} \\ &= \frac{1}{y^4} \end{aligned}$$

M and N are now multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} so not to confuse them with the original M and N .

$$\begin{aligned} \bar{M} &= \mu M \\ &= \frac{1}{y^4} (-2xy) \\ &= -\frac{2x}{y^3} \end{aligned}$$

And

$$\begin{aligned} \bar{N} &= \mu N \\ &= \frac{1}{y^4} (3x^2 - y^2) \\ &= \frac{3x^2 - y^2}{y^4} \end{aligned}$$

So now a modified ODE is obtained from the original ODE which will be exact and can be solved using the standard method. The modified ODE is

$$\begin{aligned} \bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ \left(-\frac{2x}{y^3} \right) + \left(\frac{3x^2 - y^2}{y^4} \right) \frac{dy}{dx} &= 0 \end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \quad (1)$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned} \int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\frac{2x}{y^3} dx \\ \phi &= -\frac{x^2}{y^3} + f(y) \end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = \frac{3x^2}{y^4} + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{3x^2 - y^2}{y^4}$. Therefore equation (4) becomes

$$\frac{3x^2 - y^2}{y^4} = \frac{3x^2}{y^4} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = -\frac{1}{y^2}$$

Integrating the above w.r.t y gives

$$\begin{aligned} \int f'(y) dy &= \int \left(-\frac{1}{y^2}\right) dy \\ f(y) &= \frac{1}{y} + c_1 \end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\frac{x^2}{y^3} + \frac{1}{y} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\frac{x^2}{y^3} + \frac{1}{y}$$

Summary

The solution(s) found are the following

$$-\frac{x^2}{y^3} + \frac{1}{y} = c_1 \tag{1}$$




Figure 181: Slope field plot

Verification of solutions

$$-\frac{x^2}{y^3} + \frac{1}{y} = c_1$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying homogeneous D  
<- homogeneous successful`
```

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 317

```
dsolve((3*x^2-y(x)^2)*diff(y(x),x)-2*x*y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{1 + \frac{\left(12\sqrt{3}x\sqrt{27x^2c_1^2 - 4c_1 - 108x^2c_1^2 + 8}\right)^{\frac{1}{3}}}{2} + \frac{2}{\left(12\sqrt{3}x\sqrt{27x^2c_1^2 - 4c_1 - 108x^2c_1^2 + 8}\right)^{\frac{1}{3}}}}{3c_1}$$
$$y(x) = \frac{(1 + i\sqrt{3}) \left(12\sqrt{3}x\sqrt{27x^2c_1^2 - 4c_1 - 108x^2c_1^2 + 8}\right)^{\frac{2}{3}} - 4i\sqrt{3} - 4 \left(12\sqrt{3}x\sqrt{27x^2c_1^2 - 4c_1 - 108x^2c_1^2 + 8}\right)^{\frac{1}{3}}}{12 \left(12\sqrt{3}x\sqrt{27x^2c_1^2 - 4c_1 - 108x^2c_1^2 + 8}\right)^{\frac{1}{3}} c_1}$$
$$y(x) = \frac{(i\sqrt{3} - 1) \left(12\sqrt{3}x\sqrt{27x^2c_1^2 - 4c_1 - 108x^2c_1^2 + 8}\right)^{\frac{2}{3}} - 4i\sqrt{3} + 4 \left(12\sqrt{3}x\sqrt{27x^2c_1^2 - 4c_1 - 108x^2c_1^2 + 8}\right)^{\frac{1}{3}}}{12 \left(12\sqrt{3}x\sqrt{27x^2c_1^2 - 4c_1 - 108x^2c_1^2 + 8}\right)^{\frac{1}{3}} c_1}$$

✓ Solution by Mathematica

Time used: 60.184 (sec). Leaf size: 458

`DSolve[(3*x^2-y[x]^2)*y'[x]-2*x*y[x]==0,y[x],x,IncludeSingularSolutions -> True]`

$$\begin{aligned}
 y(x) &\rightarrow \frac{1}{3} \left(\frac{\sqrt[3]{27e^{c_1}x^2 + 3\sqrt{81e^{2c_1}x^4 - 12e^{4c_1}x^2} - 2e^{3c_1}}}{\sqrt[3]{2}} \right. \\
 &\quad \left. + \frac{\sqrt[3]{2}e^{2c_1}}{\sqrt[3]{27e^{c_1}x^2 + 3\sqrt{81e^{2c_1}x^4 - 12e^{4c_1}x^2} - 2e^{3c_1}}} - e^{c_1} \right) \\
 y(x) &\rightarrow \frac{i(\sqrt{3} + i) \sqrt[3]{27e^{c_1}x^2 + 3\sqrt{81e^{2c_1}x^4 - 12e^{4c_1}x^2} - 2e^{3c_1}}}{6\sqrt[3]{2}} \\
 &\quad - \frac{i(\sqrt{3} - i) e^{2c_1}}{3 \cdot 2^{2/3} \sqrt[3]{27e^{c_1}x^2 + 3\sqrt{81e^{2c_1}x^4 - 12e^{4c_1}x^2} - 2e^{3c_1}}} - \frac{e^{c_1}}{3} \\
 y(x) &\rightarrow - \frac{i(\sqrt{3} - i) \sqrt[3]{27e^{c_1}x^2 + 3\sqrt{81e^{2c_1}x^4 - 12e^{4c_1}x^2} - 2e^{3c_1}}}{6\sqrt[3]{2}} \\
 &\quad + \frac{i(\sqrt{3} + i) e^{2c_1}}{3 \cdot 2^{2/3} \sqrt[3]{27e^{c_1}x^2 + 3\sqrt{81e^{2c_1}x^4 - 12e^{4c_1}x^2} - 2e^{3c_1}}} - \frac{e^{c_1}}{3}
 \end{aligned}$$

6.2 problem 1(b)

6.2.1 Solving as exact ode 859

Internal problem ID [6226]

Internal file name [OUTPUT/5474_Sunday_June_05_2022_03_40_20_PM_72852053/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.8. Integrating Factors. Page 32

Problem number: 1(b).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**exactWithIntegrationFactor**"

Maple gives the following as the ode type

```
[_rational, [_1st_order, `_with_symmetry_[F(x),G(x)]`], [_Abel,
`2nd type`, `class B`]]
```

$$xy + (x^2 - xy)y' = 1$$

6.2.1 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned}(x^2 - xy) dy &= (-xy + 1) dx \\ (xy - 1) dx + (x^2 - xy) dy &= 0\end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= xy - 1 \\ N(x, y) &= x^2 - xy\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(xy - 1) \\ &= x\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(x^2 - xy) \\ &= 2x - y\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= \frac{1}{x(x-y)} ((x) - (2x-y)) \\ &= -\frac{1}{x} \end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int A \, dx} \\ &= e^{\int -\frac{1}{x} \, dx} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^{-\ln(x)} \\ &= \frac{1}{x} \end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} for now so not to confuse them with the original M and N .

$$\begin{aligned} \bar{M} &= \mu M \\ &= \frac{1}{x}(xy - 1) \\ &= \frac{xy - 1}{x} \end{aligned}$$

And

$$\begin{aligned} \bar{N} &= \mu N \\ &= \frac{1}{x}(x^2 - xy) \\ &= x - y \end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned} \bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ \left(\frac{xy - 1}{x} \right) + (x - y) \frac{dy}{dx} &= 0 \end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \overline{M} \quad (1)$$

$$\frac{\partial \phi}{\partial y} = \overline{N} \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned} \int \frac{\partial \phi}{\partial x} dx &= \int \overline{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int \frac{xy - 1}{x} dx \\ \phi &= xy - \ln(x) + f(y) \end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = x + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = x - y$. Therefore equation (4) becomes

$$x - y = x + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = -y$$

Integrating the above w.r.t y gives

$$\begin{aligned} \int f'(y) dy &= \int (-y) dy \\ f(y) &= -\frac{y^2}{2} + c_1 \end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = xy - \ln(x) - \frac{y^2}{2} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = xy - \ln(x) - \frac{y^2}{2}$$

Summary

The solution(s) found are the following

$$xy - \ln(x) - \frac{y^2}{2} = c_1 \quad (1)$$




Figure 182: Slope field plot

Verification of solutions

$$xy - \ln(x) - \frac{y^2}{2} = c_1$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying Chini  
differential order: 1; looking for linear symmetries  
trying exact  
<- exact successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 39

```
dsolve((x*y(x)-1)+(x^2-x*y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = x - \sqrt{x^2 - 2 \ln(x) + 2c_1}$$
$$y(x) = x + \sqrt{x^2 - 2 \ln(x) + 2c_1}$$

✓ Solution by Mathematica

Time used: 0.393 (sec). Leaf size: 68

```
DSolve[(x*y[x]-1)+(x^2-x*y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x + \sqrt{-\frac{1}{x} \sqrt{-x(x^2 - 2 \log(x) + c_1)}}$$
$$y(x) \rightarrow x + x \left(-\frac{1}{x}\right)^{3/2} \sqrt{-x(x^2 - 2 \log(x) + c_1)}$$

6.3 problem 1(c)

6.3.1 Solving as exact ode 865

Internal problem ID [6227]

Internal file name [OUTPUT/5475_Sunday_June_05_2022_03_40_22_PM_75831642/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.8. Integrating Factors. Page 32

Problem number: 1(c).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exactWithIntegrationFactor"**

Maple gives the following as the ode type

```
[[_homogeneous , `class G`], _rational]
```

$$xy' + y + 3x^3y^4y' = 0$$

6.3.1 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned} (3y^4 x^3 + x) dy &= (-y) dx \\ (y) dx + (3y^4 x^3 + x) dy &= 0 \end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= y \\ N(x, y) &= 3y^4 x^3 + x \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(y) \\ &= 1 \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(3y^4 x^3 + x) \\ &= 9y^4 x^2 + 1 \end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= \frac{1}{3y^4 x^3 + x} ((1) - (9y^4 x^2 + 1)) \\ &= -\frac{9y^4 x}{3y^4 x^2 + 1} \end{aligned}$$

Since A depends on y , it can not be used to obtain an integrating factor. We will now try a second method to find an integrating factor. Let

$$\begin{aligned} B &= \frac{1}{M} \left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) \\ &= \frac{1}{y} ((9y^4x^2 + 1) - (1)) \\ &= 9y^3x^2 \end{aligned}$$

Since B depends on x , it can not be used to obtain an integrating factor. We will now try a third method to find an integrating factor. Let

$$R = \frac{\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y}}{xM - yN}$$

R is now checked to see if it is a function of only $t = xy$. Therefore

$$\begin{aligned} R &= \frac{\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y}}{xM - yN} \\ &= \frac{(9y^4x^2 + 1) - (1)}{x(y) - y(3y^4x^3 + x)} \\ &= -\frac{3}{yx} \end{aligned}$$

Replacing all powers of terms xy by t gives

$$R = -\frac{3}{t}$$

Since R depends on t only, then it can be used to find an integrating factor. Let the integrating factor be μ then

$$\begin{aligned} \mu &= e^{\int R dt} \\ &= e^{\int (-\frac{3}{t}) dt} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^{-3 \ln(t)} \\ &= \frac{1}{t^3} \end{aligned}$$

Now t is replaced back with xy giving

$$\mu = \frac{1}{x^3y^3}$$

Multiplying M and N by this integrating factor gives new M and new N which are called \bar{M} and \bar{N} so not to confuse them with the original M and N

$$\begin{aligned}\bar{M} &= \mu M \\ &= \frac{1}{x^3 y^3}(y) \\ &= \frac{1}{y^2 x^3}\end{aligned}$$

And

$$\begin{aligned}\bar{N} &= \mu N \\ &= \frac{1}{x^3 y^3}(3y^4 x^3 + x) \\ &= \frac{3y^4 x^2 + 1}{x^2 y^3}\end{aligned}$$

A modified ODE is now obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned}\bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ \left(\frac{1}{y^2 x^3} \right) + \left(\frac{3y^4 x^2 + 1}{x^2 y^3} \right) \frac{dy}{dx} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int \frac{1}{y^2 x^3} dx \\ \phi &= -\frac{1}{2y^2 x^2} + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = \frac{1}{y^3 x^2} + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{3y^4x^2+1}{x^2y^3}$. Therefore equation (4) becomes

$$\frac{3y^4x^2+1}{x^2y^3} = \frac{1}{y^3x^2} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 3y$$

Integrating the above w.r.t y gives

$$\int f'(y) dy = \int (3y) dy$$
$$f(y) = \frac{3y^2}{2} + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\frac{1}{2y^2x^2} + \frac{3y^2}{2} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\frac{1}{2y^2x^2} + \frac{3y^2}{2}$$

Summary

The solution(s) found are the following

$$-\frac{1}{2y^2x^2} + \frac{3y^2}{2} = c_1 \quad (1)$$




Figure 183: Slope field plot

Verification of solutions

$$-\frac{1}{2y^2x^2} + \frac{3y^2}{2} = c_1$$

Verified OK.

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying homogeneous G
<- homogeneous successful`

```

✓ Solution by Maple

Time used: 0.156 (sec). Leaf size: 133

```
dsolve(x*diff(y(x),x)+y(x)+3*x^3*y(x)^4*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = -\frac{\sqrt{6} \sqrt{xc_1 \left(x - \sqrt{12c_1^2 + x^2} \right)}}{6xc_1}$$

$$y(x) = \frac{\sqrt{6} \sqrt{xc_1 \left(x - \sqrt{12c_1^2 + x^2} \right)}}{6xc_1}$$

$$y(x) = -\frac{\sqrt{6} \sqrt{xc_1 \left(x + \sqrt{12c_1^2 + x^2} \right)}}{6xc_1}$$

$$y(x) = \frac{\sqrt{6} \sqrt{xc_1 \left(x + \sqrt{12c_1^2 + x^2} \right)}}{6xc_1}$$

✓ Solution by Mathematica

Time used: 9.711 (sec). Leaf size: 166

```
DSolve[x*y'[x]+y[x]+3*x^3*y[x]^4*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\sqrt{c_1 - \frac{\sqrt{x^2(3+c_1^2x^2)}}{x^2}}}{\sqrt{3}}$$

$$y(x) \rightarrow \frac{\sqrt{c_1 - \frac{\sqrt{x^2(3+c_1^2x^2)}}{x^2}}}{\sqrt{3}}$$

$$y(x) \rightarrow -\frac{\sqrt{\frac{\sqrt{x^2(3+c_1^2x^2)}}{x^2} + c_1}}{\sqrt{3}}$$

$$y(x) \rightarrow \frac{\sqrt{\frac{\sqrt{x^2(3+c_1^2x^2)}}{x^2} + c_1}}{\sqrt{3}}$$

$$y(x) \rightarrow 0$$

6.4 problem 1(d)

6.4.1 Solving as exact ode 872

Internal problem ID [6228]

Internal file name [OUTPUT/5476_Sunday_June_05_2022_03_40_25_PM_67230230/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.8. Integrating Factors. Page 32

Problem number: 1(d).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exactWithIntegrationFactor"**

Maple gives the following as the ode type

```
[[_1st_order , `_with_symmetry_[F(x)*G(y),0]`]]
```

$$(e^x \cot(y) + 2y \csc(y)) y' = -e^x$$

6.4.1 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned} (e^x \cot(y) + 2y \csc(y)) dy &= (-e^x) dx \\ (e^x) dx + (e^x \cot(y) + 2y \csc(y)) dy &= 0 \end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= e^x \\ N(x, y) &= e^x \cot(y) + 2y \csc(y) \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(e^x) \\ &= 0 \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(e^x \cot(y) + 2y \csc(y)) \\ &= e^x \cot(y) \end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= \frac{\sin(y)}{e^x \cos(y) + 2y} ((0) - (e^x \cot(y))) \\ &= -\frac{e^x \cos(y)}{e^x \cos(y) + 2y} \end{aligned}$$

Since A depends on y , it can not be used to obtain an integrating factor. We will now try a second method to find an integrating factor. Let

$$\begin{aligned} B &= \frac{1}{M} \left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) \\ &= e^{-x} ((e^x \cot(y)) - (0)) \\ &= \cot(y) \end{aligned}$$

Since B does not depend on x , it can be used to obtain an integrating factor. Let the integrating factor be μ . Then

$$\begin{aligned} \mu &= e^{\int B \, dy} \\ &= e^{\int \cot(y) \, dy} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^{\ln(\sin(y))} \\ &= \sin(y) \end{aligned}$$

M and N are now multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} so not to confuse them with the original M and N .

$$\begin{aligned} \bar{M} &= \mu M \\ &= \sin(y) (e^x) \\ &= e^x \sin(y) \end{aligned}$$

And

$$\begin{aligned} \bar{N} &= \mu N \\ &= \sin(y) (e^x \cot(y) + 2y \csc(y)) \\ &= e^x \cos(y) + 2y \end{aligned}$$

So now a modified ODE is obtained from the original ODE which will be exact and can be solved using the standard method. The modified ODE is

$$\begin{aligned} \overline{M} + \overline{N} \frac{dy}{dx} &= 0 \\ (e^x \sin(y)) + (e^x \cos(y) + 2y) \frac{dy}{dx} &= 0 \end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \overline{M} \quad (1)$$

$$\frac{\partial \phi}{\partial y} = \overline{N} \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned} \int \frac{\partial \phi}{\partial x} dx &= \int \overline{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int e^x \sin(y) dx \\ \phi &= e^x \sin(y) + f(y) \end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = e^x \cos(y) + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = e^x \cos(y) + 2y$. Therefore equation (4) becomes

$$e^x \cos(y) + 2y = e^x \cos(y) + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 2y$$

Integrating the above w.r.t y gives

$$\begin{aligned} \int f'(y) dy &= \int (2y) dy \\ f(y) &= y^2 + c_1 \end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = e^x \sin(y) + y^2 + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = e^x \sin(y) + y^2$$

Summary

The solution(s) found are the following

$$e^x \sin(y) + y^2 = c_1 \tag{1}$$




Figure 184: Slope field plot

Verification of solutions

$$e^x \sin(y) + y^2 = c_1$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying Chini  
differential order: 1; looking for linear symmetries  
trying exact  
<- exact successful`
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 15

```
dsolve(exp(x)+(exp(x)*cot(y(x))+2*y(x)*csc(y(x)))*diff(y(x),x)=0,y(x), singsol=all)
```

$$e^x \sin(y(x)) + y(x)^2 + c_1 = 0$$

✓ Solution by Mathematica

Time used: 0.322 (sec). Leaf size: 18

```
DSolve[Exp[x]+(Exp[x]*Cot[y[x]]+2*y[x]*Csc[y[x]])*y'[x]==0,y[x],x,IncludeSingularSolutions -
```

$$\text{Solve}[y(x)^2 + e^x \sin(y(x)) = c_1, y(x)]$$

6.5 problem 1(e)

6.5.1 Solving as exact ode	878
6.5.2 Maple step by step solution	882

Internal problem ID [6229]

Internal file name [OUTPUT/5477_Sunday_June_05_2022_03_40_39_PM_65128346/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.8. Integrating Factors. Page 32

Problem number: 1(e).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exact"**

Maple gives the following as the ode type

`[_separable]`

$$(x + 2) \sin(y) + y' \cos(y) x = 0$$

6.5.1 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned}\left(-\frac{\cos(y)}{\sin(y)}\right) dy &= \left(\frac{x+2}{x}\right) dx \\ \left(-\frac{x+2}{x}\right) dx + \left(-\frac{\cos(y)}{\sin(y)}\right) dy &= 0\end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -\frac{x+2}{x} \\ N(x, y) &= -\frac{\cos(y)}{\sin(y)}\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(-\frac{x+2}{x}\right) \\ &= 0\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(-\frac{\cos(y)}{\sin(y)} \right) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\frac{x+2}{x} dx \\ \phi &= -x - 2 \ln(x) + f(y)\end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = -\frac{\cos(y)}{\sin(y)}$. Therefore equation (4) becomes

$$-\frac{\cos(y)}{\sin(y)} = 0 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$\begin{aligned}f'(y) &= -\frac{\cos(y)}{\sin(y)} \\ &= -\cot(y)\end{aligned}$$

Integrating the above w.r.t y results in

$$\int f'(y) dy = \int (-\cot(y)) dy$$

$$f(y) = -\ln(\sin(y)) + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -x - 2\ln(x) - \ln(\sin(y)) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -x - 2\ln(x) - \ln(\sin(y))$$

Summary

The solution(s) found are the following

$$-x - 2\ln(x) - \ln(\sin(y)) = c_1 \tag{1}$$




Figure 185: Slope field plot

Verification of solutions

$$-x - 2 \ln(x) - \ln(\sin(y)) = c_1$$

Verified OK.

6.5.2 Maple step by step solution

Let's solve

$$(x + 2) \sin(y) + y' \cos(y) x = 0$$

- Highest derivative means the order of the ODE is 1

y'

- Separate variables

$$\frac{y' \cos(y)}{\sin(y)} = -\frac{x+2}{x}$$

- Integrate both sides with respect to x

$$\int \frac{y' \cos(y)}{\sin(y)} dx = \int -\frac{x+2}{x} dx + c_1$$

- Evaluate integral

$$\ln(\sin(y)) = -x - 2 \ln(x) + c_1$$

- Solve for y

$$y = \arcsin\left(\frac{e^{-x+c_1}}{x^2}\right)$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
<- separable successful`
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 16

```
dsolve((x+2)*sin(y(x))+x*cos(y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \arcsin\left(\frac{e^{-x}}{c_1 x^2}\right)$$

✓ Solution by Mathematica

Time used: 51.335 (sec). Leaf size: 23

```
DSolve[(x+2)*Sin[y[x]]+x*Cos[y[x]]*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \arcsin\left(\frac{e^{-x+c_1}}{x^2}\right)$$

$$y(x) \rightarrow 0$$

6.6 problem 1(f)

6.6.1 Solving as exact ode 884

Internal problem ID [6230]

Internal file name [OUTPUT/5478_Sunday_June_05_2022_03_40_42_PM_72967215/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.8. Integrating Factors. Page 32

Problem number: 1(f).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exactWithIntegrationFactor"**

Maple gives the following as the ode type

```
[[_homogeneous , `class G`], _rational]
```

$$y + (x - 2y^3x^2)y' = 0$$

6.6.1 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned} (-2y^3x^2 + x) dy &= (-y) dx \\ (y) dx + (-2y^3x^2 + x) dy &= 0 \end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= y \\ N(x, y) &= -2y^3x^2 + x \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(y) \\ &= 1 \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(-2y^3x^2 + x) \\ &= -4xy^3 + 1 \end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= \frac{1}{-2y^3x^2 + x} ((1) - (-4xy^3 + 1)) \\ &= -\frac{4y^3}{2xy^3 - 1} \end{aligned}$$

Since A depends on y , it can not be used to obtain an integrating factor. We will now try a second method to find an integrating factor. Let

$$\begin{aligned} B &= \frac{1}{M} \left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) \\ &= \frac{1}{y} ((-4xy^3 + 1) - (1)) \\ &= -4y^2x \end{aligned}$$

Since B depends on x , it can not be used to obtain an integrating factor. We will now try a third method to find an integrating factor. Let

$$R = \frac{\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y}}{xM - yN}$$

R is now checked to see if it is a function of only $t = xy$. Therefore

$$\begin{aligned} R &= \frac{\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y}}{xM - yN} \\ &= \frac{(-4xy^3 + 1) - (1)}{x(y) - y(-2y^3x^2 + x)} \\ &= -\frac{2}{yx} \end{aligned}$$

Replacing all powers of terms xy by t gives

$$R = -\frac{2}{t}$$

Since R depends on t only, then it can be used to find an integrating factor. Let the integrating factor be μ then

$$\begin{aligned} \mu &= e^{\int R dt} \\ &= e^{\int (-\frac{2}{t}) dt} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^{-2 \ln(t)} \\ &= \frac{1}{t^2} \end{aligned}$$

Now t is replaced back with xy giving

$$\mu = \frac{1}{y^2x^2}$$

Multiplying M and N by this integrating factor gives new M and new N which are called \bar{M} and \bar{N} so not to confuse them with the original M and N

$$\begin{aligned}\bar{M} &= \mu M \\ &= \frac{1}{y^2 x^2}(y) \\ &= \frac{1}{y x^2}\end{aligned}$$

And

$$\begin{aligned}\bar{N} &= \mu N \\ &= \frac{1}{y^2 x^2}(-2y^3 x^2 + x) \\ &= \frac{-2x y^3 + 1}{y^2 x}\end{aligned}$$

A modified ODE is now obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned}\bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ \left(\frac{1}{y x^2} \right) + \left(\frac{-2x y^3 + 1}{y^2 x} \right) \frac{dy}{dx} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int \frac{1}{y x^2} dx \\ \phi &= -\frac{1}{y x} + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = \frac{1}{y^2 x} + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{-2xy^3+1}{y^2x}$. Therefore equation (4) becomes

$$\frac{-2xy^3+1}{y^2x} = \frac{1}{y^2x} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = -2y$$

Integrating the above w.r.t y gives

$$\int f'(y) dy = \int (-2y) dy$$
$$f(y) = -y^2 + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\frac{1}{yx} - y^2 + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\frac{1}{yx} - y^2$$

Summary

The solution(s) found are the following

$$-\frac{1}{xy} - y^2 = c_1 \quad (1)$$




Figure 186: Slope field plot

Verification of solutions

$$-\frac{1}{xy} - y^2 = c_1$$

Verified OK.

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying homogeneous G
<- homogeneous successful`

```

✓ Solution by Maple

Time used: 0.125 (sec). Leaf size: 320

`dsolve(y(x)+(x-2*x^2*y(x)^3)*diff(y(x),x)=0,y(x), singsol=all)`

$$y(x) = \frac{12^{\frac{1}{3}} \left(12^{\frac{1}{3}} c_1 x^2 + \left(-9x^2 \left(-\frac{\sqrt{3} \sqrt{27c_1^3 - 4x^2}}{9c_1} + c_1 \right) c_1^2 \right)^{\frac{2}{3}} \right)}{6c_1 x \left(-9x^2 \left(-\frac{\sqrt{3} \sqrt{27c_1^3 - 4x^2}}{9c_1} + c_1 \right) c_1^2 \right)^{\frac{1}{3}}}$$

$$y(x) = \frac{2^{\frac{2}{3}} \left((-i\sqrt{3} - 1) \left(-9x^2 \left(-\frac{\sqrt{3} \sqrt{27c_1^3 - 4x^2}}{9c_1} + c_1 \right) c_1^2 \right)^{\frac{2}{3}} + 2^{\frac{2}{3}} x^2 c_1 \left(i3^{\frac{5}{6}} - 3^{\frac{1}{3}} \right) \right) 3^{\frac{1}{3}}}{12 \left(-9x^2 \left(-\frac{\sqrt{3} \sqrt{27c_1^3 - 4x^2}}{9c_1} + c_1 \right) c_1^2 \right)^{\frac{1}{3}} x c_1}$$

$$y(x) = \frac{2^{\frac{2}{3}} \left((1 - i\sqrt{3}) \left(-9x^2 \left(-\frac{\sqrt{3} \sqrt{27c_1^3 - 4x^2}}{9c_1} + c_1 \right) c_1^2 \right)^{\frac{2}{3}} + 2^{\frac{2}{3}} \left(i3^{\frac{5}{6}} + 3^{\frac{1}{3}} \right) x^2 c_1 \right) 3^{\frac{1}{3}}}{12 \left(-9x^2 \left(-\frac{\sqrt{3} \sqrt{27c_1^3 - 4x^2}}{9c_1} + c_1 \right) c_1^2 \right)^{\frac{1}{3}} x c_1}$$

✓ Solution by Mathematica

Time used: 28.221 (sec). Leaf size: 327

`DSolve[y[x]+(x-2*x^2*y[x]^3)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \frac{2\sqrt[3]{3}c_1x^2 + \sqrt[3]{2}(-9x^2 + \sqrt{81x^4 - 12c_1^3x^6})^{2/3}}{6^{2/3}x^3\sqrt[3]{-9x^2 + \sqrt{81x^4 - 12c_1^3x^6}}}$$

$$y(x) \rightarrow \frac{i\sqrt[3]{3}(\sqrt{3} + i)(-18x^2 + 2\sqrt{81x^4 - 12c_1^3x^6})^{2/3} - 2\sqrt[3]{2}\sqrt[6]{3}(\sqrt{3} + 3i)c_1x^2}{12x^3\sqrt[3]{-9x^2 + \sqrt{81x^4 - 12c_1^3x^6}}}$$

$$y(x) \rightarrow \frac{\sqrt[3]{3}(-1 - i\sqrt{3})(-18x^2 + 2\sqrt{81x^4 - 12c_1^3x^6})^{2/3} - 2\sqrt[3]{2}\sqrt[6]{3}(\sqrt{3} - 3i)c_1x^2}{12x^3\sqrt[3]{-9x^2 + \sqrt{81x^4 - 12c_1^3x^6}}}$$

$$y(x) \rightarrow 0$$

6.7 problem 1(g)

6.7.1 Solving as exact ode 892

Internal problem ID [6231]

Internal file name [OUTPUT/5479_Sunday_June_05_2022_03_40_47_PM_13544762/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.8. Integrating Factors. Page 32

Problem number: 1(g).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exactWithIntegrationFactor"**

Maple gives the following as the ode type

```
[[_homogeneous, `class G`], _rational, _Bernoulli]
```

$$3y^2 + 2xyy' = -x$$

6.7.1 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned} (2xy) dy &= (-3y^2 - x) dx \\ (3y^2 + x) dx + (2xy) dy &= 0 \end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= 3y^2 + x \\ N(x, y) &= 2xy \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} (3y^2 + x) \\ &= 6y \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} (2xy) \\ &= 2y \end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= \frac{1}{2yx} ((6y) - (2y)) \\ &= \frac{2}{x} \end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int A dx} \\ &= e^{\int \frac{2}{x} dx}\end{aligned}$$

The result of integrating gives

$$\begin{aligned}\mu &= e^{2\ln(x)} \\ &= x^2\end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} for now so not to confuse them with the original M and N .

$$\begin{aligned}\bar{M} &= \mu M \\ &= x^2(3y^2 + x) \\ &= x^2(3y^2 + x)\end{aligned}$$

And

$$\begin{aligned}\bar{N} &= \mu N \\ &= x^2(2xy) \\ &= 2y x^3\end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned}\bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ (x^2(3y^2 + x)) + (2y x^3) \frac{dy}{dx} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int x^2(3y^2 + x) dx \\ \phi &= y^2 x^3 + \frac{1}{4} x^4 + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 2y x^3 + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = 2y x^3$. Therefore equation (4) becomes

$$2y x^3 = 2y x^3 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = y^2 x^3 + \frac{1}{4} x^4 + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = y^2 x^3 + \frac{1}{4} x^4$$

Summary

The solution(s) found are the following

$$x^3 y^2 + \frac{x^4}{4} = c_1 \quad (1)$$




Figure 187: Slope field plot

Verification of solutions

$$x^3y^2 + \frac{x^4}{4} = c_1$$

Verified OK.

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
<- Bernoulli successful`

```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 41

```
dsolve((x+3*y(x)^2)+(2*x*y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = -\frac{\sqrt{-x(x^4 - 4c_1)}}{2x^2}$$
$$y(x) = \frac{\sqrt{-x(x^4 - 4c_1)}}{2x^2}$$

✓ Solution by Mathematica

Time used: 0.206 (sec). Leaf size: 55

```
DSolve[(x+3*y[x]^2)+(2*x*y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\sqrt{-x^4 + 4c_1}}{2x^{3/2}}$$
$$y(x) \rightarrow \frac{\sqrt{-x^4 + 4c_1}}{2x^{3/2}}$$

6.8 problem 1(h)

6.8.1 Solving as exact ode 898

Internal problem ID [6232]

Internal file name [OUTPUT/5480_Sunday_June_05_2022_03_40_49_PM_66946067/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.8. Integrating Factors. Page 32

Problem number: 1(h).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**exactWithIntegrationFactor**"

Maple gives the following as the ode type

```
[[_1st_order , `_with_symmetry_[F(x)*G(y),0]`]]
```

$$y + (2x - e^y y) y' = 0$$

6.8.1 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned} (2x - e^y y) dy &= (-y) dx \\ (y) dx + (2x - e^y y) dy &= 0 \end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= y \\ N(x, y) &= 2x - e^y y \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(y) \\ &= 1 \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(2x - e^y y) \\ &= 2 \end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= \frac{1}{2x - e^y y} ((1) - (2)) \\ &= \frac{1}{e^y y - 2x} \end{aligned}$$

Since A depends on y , it can not be used to obtain an integrating factor. We will now try a second method to find an integrating factor. Let

$$\begin{aligned} B &= \frac{1}{M} \left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) \\ &= \frac{1}{y} ((2) - (1)) \\ &= \frac{1}{y} \end{aligned}$$

Since B does not depend on x , it can be used to obtain an integrating factor. Let the integrating factor be μ . Then

$$\begin{aligned} \mu &= e^{\int B \, dy} \\ &= e^{\int \frac{1}{y} \, dy} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^{\ln(y)} \\ &= y \end{aligned}$$

M and N are now multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} so not to confuse them with the original M and N .

$$\begin{aligned} \bar{M} &= \mu M \\ &= y(y) \\ &= y^2 \end{aligned}$$

And

$$\begin{aligned} \bar{N} &= \mu N \\ &= y(2x - e^y y) \\ &= -y^2 e^y + 2xy \end{aligned}$$

So now a modified ODE is obtained from the original ODE which will be exact and can be solved using the standard method. The modified ODE is

$$\begin{aligned} \bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ (y^2) + (-y^2 e^y + 2xy) \frac{dy}{dx} &= 0 \end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \quad (1)$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned} \int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int y^2 dx \\ \phi &= y^2 x + f(y) \end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 2xy + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = -y^2 e^y + 2xy$. Therefore equation (4) becomes

$$-y^2 e^y + 2xy = 2xy + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = -y^2 e^y$$

Integrating the above w.r.t y gives

$$\begin{aligned} \int f'(y) dy &= \int (-y^2 e^y) dy \\ f(y) &= -(y^2 - 2y + 2) e^y + c_1 \end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = y^2 x - (y^2 - 2y + 2) e^y + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = y^2 x - (y^2 - 2y + 2) e^y$$

Summary

The solution(s) found are the following

$$xy^2 - (y^2 - 2y + 2) e^y = c_1 \tag{1}$$




Figure 188: Slope field plot

Verification of solutions

$$xy^2 - (y^2 - 2y + 2) e^y = c_1$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
<- 1st order linear successful  
<- inverse linear successful`
```

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 34

```
dsolve(y(x)+(2*x-y(x)*exp(y(x)))*diff(y(x),x)=0,y(x), singsol=all)
```

$$\frac{(-y(x)^2 + 2y(x) - 2) e^{y(x)} + xy(x)^2 - c_1}{y(x)^2} = 0$$

✓ Solution by Mathematica

Time used: 0.259 (sec). Leaf size: 32

```
DSolve[y[x]+(2*x-y[x]*Exp[y[x]])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[x = \frac{e^{y(x)}(y(x)^2 - 2y(x) + 2)}{y(x)^2} + \frac{c_1}{y(x)^2}, y(x) \right]$$

6.9 problem 1(i)

6.9.1 Solving as exact ode 904

Internal problem ID [6233]

Internal file name [OUTPUT/5481_Sunday_June_05_2022_03_40_52_PM_86106885/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.8. Integrating Factors. Page 32

Problem number: 1(i).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**exactWithIntegrationFactor**"

Maple gives the following as the ode type

[`y=_G(x,y)´]

$$y \ln(y) - 2xy + (x + y)y' = 0$$

6.9.1 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned} (x + y) dy &= (-\ln(y)y + 2xy) dx \\ (\ln(y)y - 2xy) dx + (x + y) dy &= 0 \end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= \ln(y)y - 2xy \\ N(x, y) &= x + y \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(\ln(y)y - 2xy) \\ &= 1 + \ln(y) - 2x \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(x + y) \\ &= 1 \end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= \frac{1}{x + y} ((1 + \ln(y) - 2x) - (1)) \\ &= \frac{\ln(y) - 2x}{x + y} \end{aligned}$$

Since A depends on y , it can not be used to obtain an integrating factor. We will now try a second method to find an integrating factor. Let

$$\begin{aligned} B &= \frac{1}{M} \left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) \\ &= \frac{1}{y(\ln(y) - 2x)} ((1) - (1 + \ln(y) - 2x)) \\ &= -\frac{1}{y} \end{aligned}$$

Since B does not depend on x , it can be used to obtain an integrating factor. Let the integrating factor be μ . Then

$$\begin{aligned} \mu &= e^{\int B \, dy} \\ &= e^{\int -\frac{1}{y} \, dy} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^{-\ln(y)} \\ &= \frac{1}{y} \end{aligned}$$

M and N are now multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} so not to confuse them with the original M and N .

$$\begin{aligned} \bar{M} &= \mu M \\ &= \frac{1}{y} (\ln(y) y - 2xy) \\ &= \ln(y) - 2x \end{aligned}$$

And

$$\begin{aligned} \bar{N} &= \mu N \\ &= \frac{1}{y} (x + y) \\ &= \frac{x + y}{y} \end{aligned}$$

So now a modified ODE is obtained from the original ODE which will be exact and can be solved using the standard method. The modified ODE is

$$\begin{aligned} \bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ (\ln(y) - 2x) + \left(\frac{x + y}{y} \right) \frac{dy}{dx} &= 0 \end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \overline{M} \quad (1)$$

$$\frac{\partial \phi}{\partial y} = \overline{N} \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned} \int \frac{\partial \phi}{\partial x} dx &= \int \overline{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int \ln(y) - 2x dx \\ \phi &= x(\ln(y) - x) + f(y) \end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = \frac{x}{y} + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{x+y}{y}$. Therefore equation (4) becomes

$$\frac{x+y}{y} = \frac{x}{y} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 1$$

Integrating the above w.r.t y gives

$$\begin{aligned} \int f'(y) dy &= \int (1) dy \\ f(y) &= y + c_1 \end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = x(\ln(y) - x) + y + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = x(\ln(y) - x) + y$$

The solution becomes

$$y = e^{-\frac{x \operatorname{LambertW}\left(\frac{e^{\frac{x^2+c_1}{x}}}{x}\right) - x^2 - c_1}{x}}$$

Summary

The solution(s) found are the following

$$y = e^{-\frac{x \operatorname{LambertW}\left(\frac{e^{\frac{x^2+c_1}{x}}}{x}\right) - x^2 - c_1}{x}} \quad (1)$$




Figure 189: Slope field plot

Verification of solutions

$$y = e^{-\frac{x \operatorname{LambertW}\left(\frac{e^{\frac{x^2+c_1}{x}}}{x}\right) - x^2 - c_1}{x}}$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying Chini  
differential order: 1; looking for linear symmetries  
trying exact  
<- exact successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 36

```
dsolve((y(x)*ln(y(x))-2*x*y(x))+(x+y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = e^{\frac{x^2 - x \operatorname{LambertW}\left(\frac{e^{-\frac{x^2 - c_1}{x}}}{x}\right) - c_1}{x}}$$

✓ Solution by Mathematica

Time used: 1.073 (sec). Leaf size: 22

```
DSolve[(y[x]*Log[y[x]]-2*x*y[x])+(x+y[x])*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow xW\left(\frac{e^{x+\frac{c_1}{x}}}{x}\right)$$

6.10 problem 1(j)

6.10.1 Solving as exact ode 910

Internal problem ID [6234]

Internal file name [OUTPUT/5482_Sunday_June_05_2022_03_40_55_PM_15432646/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.8. Integrating Factors. Page 32

Problem number: 1(j).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exactWithIntegrationFactor"**

Maple gives the following as the ode type

```
[_rational, [_Abel, `2nd type`, `class B`]]
```

$$y^2 + xy + (x^2 + xy + 1) y' = -1$$

6.10.1 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned} (x^2 + xy + 1) dy &= (-xy - y^2 - 1) dx \\ (xy + y^2 + 1) dx + (x^2 + xy + 1) dy &= 0 \end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= xy + y^2 + 1 \\ N(x, y) &= x^2 + xy + 1 \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} (xy + y^2 + 1) \\ &= 2y + x \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} (x^2 + xy + 1) \\ &= y + 2x \end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= \frac{1}{x^2 + xy + 1} ((2y + x) - (y + 2x)) \\ &= \frac{-x + y}{x^2 + xy + 1} \end{aligned}$$

Since A depends on y , it can not be used to obtain an integrating factor. We will now try a second method to find an integrating factor. Let

$$\begin{aligned} B &= \frac{1}{M} \left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) \\ &= \frac{1}{xy + y^2 + 1} ((y + 2x) - (2y + x)) \\ &= \frac{x - y}{xy + y^2 + 1} \end{aligned}$$

Since B depends on x , it can not be used to obtain an integrating factor. We will now try a third method to find an integrating factor. Let

$$R = \frac{\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y}}{xM - yN}$$

R is now checked to see if it is a function of only $t = xy$. Therefore

$$\begin{aligned} R &= \frac{\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y}}{xM - yN} \\ &= \frac{(y + 2x) - (2y + x)}{x(xy + y^2 + 1) - y(x^2 + xy + 1)} \\ &= 1 \end{aligned}$$

Replacing all powers of terms xy by t gives

$$R = 1$$

Since R depends on t only, then it can be used to find an integrating factor. Let the integrating factor be μ then

$$\begin{aligned} \mu &= e^{\int R dt} \\ &= e^{\int (1) dt} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^t \\ &= e^t \end{aligned}$$

Now t is replaced back with xy giving

$$\mu = e^{xy}$$

Multiplying M and N by this integrating factor gives new \bar{M} and new \bar{N} which are called \bar{M} and \bar{N} so not to confuse them with the original M and N

$$\begin{aligned}\bar{M} &= \mu M \\ &= e^{xy}(xy + y^2 + 1) \\ &= (xy + y^2 + 1) e^{xy}\end{aligned}$$

And

$$\begin{aligned}\bar{N} &= \mu N \\ &= e^{xy}(x^2 + xy + 1) \\ &= (x^2 + xy + 1) e^{xy}\end{aligned}$$

A modified ODE is now obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned}\bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ ((xy + y^2 + 1) e^{xy}) + ((x^2 + xy + 1) e^{xy}) \frac{dy}{dx} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int (xy + y^2 + 1) e^{xy} dx \\ \phi &= (x + y) e^{xy} + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\begin{aligned}\frac{\partial \phi}{\partial y} &= e^{xy} + (x + y) x e^{xy} + f'(y) \\ &= (x^2 + xy + 1) e^{xy} + f'(y)\end{aligned} \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = (x^2 + xy + 1)e^{xy}$. Therefore equation (4) becomes

$$(x^2 + xy + 1)e^{xy} = (x^2 + xy + 1)e^{xy} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = (x + y)e^{xy} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = (x + y)e^{xy}$$

The solution becomes

$$y = \frac{-x^2 + \text{LambertW}(c_1 x e^{x^2})}{x}$$

Summary

The solution(s) found are the following

$$y = \frac{-x^2 + \text{LambertW}(c_1 x e^{x^2})}{x} \quad (1)$$




Figure 190: Slope field plot

Verification of solutions

$$y = \frac{-x^2 + \text{LambertW}\left(c_1 x e^{x^2}\right)}{x}$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying Chini
differential order: 1; looking for linear symmetries
trying exact
trying Abel
<- Abel successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve((y(x)^2+x*y(x)+1)+(x^2+x*y(x)+1)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{-x^2 + \text{LambertW}(-2xc_1e^{(x-1)(x+1)})}{x}$$

✓ Solution by Mathematica

Time used: 6.606 (sec). Leaf size: 56

```
DSolve[(y[x]^2+x*y[x]+1)+(x^2+x*y[x]+1)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x + \frac{W\left(x\left(-e^{x^2-1+c_1}\right)\right)}{x}$$
$$y(x) \rightarrow -x$$
$$y(x) \rightarrow \frac{W\left(-e^{x^2-1}x\right)}{x} - x$$

6.11 problem 1(k)

6.11.1 Solving as exact ode 917

Internal problem ID [6235]

Internal file name [OUTPUT/5483_Sunday_June_05_2022_03_40_58_PM_37883864/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.8. Integrating Factors. Page 32

Problem number: 1(k).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**exactWithIntegrationFactor**"

Maple gives the following as the ode type

`[_rational, _Bernoulli]`

$$y^3x + 3y'y^2 = -x^3$$

6.11.1 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned} (3y^2) dy &= (-xy^3 - x^3) dx \\ (xy^3 + x^3) dx + (3y^2) dy &= 0 \end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= xy^3 + x^3 \\ N(x, y) &= 3y^2 \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(xy^3 + x^3) \\ &= 3y^2x \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(3y^2) \\ &= 0 \end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= \frac{1}{3y^2} ((3y^2x) - (0)) \\ &= x \end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int A dx} \\ &= e^{\int x dx}\end{aligned}$$

The result of integrating gives

$$\begin{aligned}\mu &= e^{\frac{x^2}{2}} \\ &= e^{\frac{x^2}{2}}\end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} for now so not to confuse them with the original M and N .

$$\begin{aligned}\bar{M} &= \mu M \\ &= e^{\frac{x^2}{2}} (x y^3 + x^3) \\ &= x(y^3 + x^2) e^{\frac{x^2}{2}}\end{aligned}$$

And

$$\begin{aligned}\bar{N} &= \mu N \\ &= e^{\frac{x^2}{2}} (3y^2) \\ &= 3y^2 e^{\frac{x^2}{2}}\end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned}\bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ \left(x(y^3 + x^2) e^{\frac{x^2}{2}} \right) + \left(3y^2 e^{\frac{x^2}{2}} \right) \frac{dy}{dx} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \overline{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int x(y^3 + x^2) e^{\frac{x^2}{2}} dx \\ \phi &= (y^3 + x^2 - 2) e^{\frac{x^2}{2}} + f(y)\end{aligned}\quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 3y^2 e^{\frac{x^2}{2}} + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = 3y^2 e^{\frac{x^2}{2}}$. Therefore equation (4) becomes

$$3y^2 e^{\frac{x^2}{2}} = 3y^2 e^{\frac{x^2}{2}} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = (y^3 + x^2 - 2) e^{\frac{x^2}{2}} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = (y^3 + x^2 - 2) e^{\frac{x^2}{2}}$$

Summary

The solution(s) found are the following

$$(y^3 + x^2 - 2) e^{\frac{x^2}{2}} = c_1 \quad (1)$$




Figure 191: Slope field plot

Verification of solutions

$$(y^3 + x^2 - 2) e^{\frac{x^2}{2}} = c_1$$

Verified OK.

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
<- Bernoulli successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 81

```
dsolve((x^3+x*y(x)^3)+(3*y(x)^2)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \left(e^{-\frac{x^2}{2}} c_1 - x^2 + 2 \right)^{\frac{1}{3}}$$
$$y(x) = -\frac{\left(e^{-\frac{x^2}{2}} c_1 - x^2 + 2 \right)^{\frac{1}{3}} (1 + i\sqrt{3})}{2}$$
$$y(x) = \frac{\left(e^{-\frac{x^2}{2}} c_1 - x^2 + 2 \right)^{\frac{1}{3}} (i\sqrt{3} - 1)}{2}$$

✓ Solution by Mathematica

Time used: 10.689 (sec). Leaf size: 95

```
DSolve[(x^3+x*y[x]^3)+(3*y[x]^2)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \sqrt[3]{-x^2 + c_1 e^{-\frac{x^2}{2}} + 2}$$
$$y(x) \rightarrow -\sqrt[3]{-1} \sqrt[3]{-x^2 + c_1 e^{-\frac{x^2}{2}} + 2}$$
$$y(x) \rightarrow (-1)^{2/3} \sqrt[3]{-x^2 + c_1 e^{-\frac{x^2}{2}} + 2}$$

6.12 problem 4

6.12.1 Solving as first order ode lie symmetry calculated ode 923

Internal problem ID [6236]

Internal file name [OUTPUT/5484_Sunday_June_05_2022_03_41_00_PM_64600096/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.8. Integrating Factors. Page 32

Problem number: 4.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**first_order_ode_lie_symmetry_calculated**"

Maple gives the following as the ode type

```
[[_homogeneous, `class G`]]
```

$$y' - \frac{2y}{x} - \frac{x^3}{y} - x \tan\left(\frac{y}{x^2}\right) = 0$$

6.12.1 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = \frac{x^2 \tan\left(\frac{y}{x^2}\right) y + x^4 + 2y^2}{xy}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \quad (\text{1E})$$

$$\eta = xb_2 + yb_3 + b_1 \quad (\text{2E})$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$\begin{aligned} b_2 + \frac{(x^2 \tan(\frac{y}{x^2}) y + x^4 + 2y^2)(b_3 - a_2)}{xy} - \frac{(x^2 \tan(\frac{y}{x^2}) y + x^4 + 2y^2)^2 a_3}{x^2 y^2} \\ - \left(\frac{2x \tan(\frac{y}{x^2}) y - \frac{2y^2(1 + \tan(\frac{y}{x^2})^2)}{x} + 4x^3}{xy} \right. \\ \left. - \frac{x^2 \tan(\frac{y}{x^2}) y + x^4 + 2y^2}{x^2 y} \right) (xa_2 + ya_3 + a_1) \\ - \left(\frac{(1 + \tan(\frac{y}{x^2})^2) y + x^2 \tan(\frac{y}{x^2}) + 4y}{xy} \right. \\ \left. - \frac{x^2 \tan(\frac{y}{x^2}) y + x^4 + 2y^2}{y^2 x} \right) (xb_2 + yb_3 + b_1) = 0 \end{aligned} \quad (5E)$$

Putting the above in normal form gives

$$\begin{aligned} - \frac{\tan(\frac{y}{x^2})^2 x^4 y^2 a_3 + 2 \tan(\frac{y}{x^2}) x^6 y a_3 + x^8 a_3 + \tan(\frac{y}{x^2})^2 x^2 y^2 b_2 - 2 \tan(\frac{y}{x^2})^2 x y^3 a_2 + \tan(\frac{y}{x^2})^2 x y^3 b_3 - 2}{x^2 y^2} \\ = 0 \end{aligned}$$

Setting the numerator to zero gives

$$\begin{aligned} - \tan(\frac{y}{x^2})^2 x^4 y^2 a_3 - 2 \tan(\frac{y}{x^2}) x^6 y a_3 - x^8 a_3 - \tan(\frac{y}{x^2})^2 x^2 y^2 b_2 \\ + 2 \tan(\frac{y}{x^2})^2 x y^3 a_2 - \tan(\frac{y}{x^2})^2 x y^3 b_3 + 2 \tan(\frac{y}{x^2})^2 y^4 a_3 \\ - 2 \tan(\frac{y}{x^2}) x^3 y^2 a_2 + \tan(\frac{y}{x^2}) x^3 y^2 b_3 - 5 \tan(\frac{y}{x^2}) x^2 y^3 a_3 \\ + x^6 b_2 - 4x^5 y a_2 + 2x^5 y b_3 - 7x^4 y^2 a_3 - \tan(\frac{y}{x^2})^2 x y^2 b_1 \\ + 2 \tan(\frac{y}{x^2})^2 y^3 a_1 - \tan(\frac{y}{x^2}) x^2 y^2 a_1 + x^5 b_1 - 3x^4 y a_1 \\ - 2b_2 y^2 x^2 + 2x y^3 a_2 - x y^3 b_3 - 3x y^2 b_1 + 4y^3 a_1 = 0 \end{aligned} \quad (6E)$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\left\{x, y, \tan\left(\frac{y}{x^2}\right)\right\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\left\{x = v_1, y = v_2, \tan\left(\frac{y}{x^2}\right) = v_3\right\}$$

The above PDE (6E) now becomes

$$\begin{aligned} & -v_1^8 a_3 - 2v_3 v_1^6 v_2 a_3 - v_3^2 v_1^4 v_2^2 a_3 - 4v_1^5 v_2 a_2 - 2v_3 v_1^3 v_2^2 a_2 + 2v_3^2 v_1 v_2^3 a_2 \\ & - 7v_1^4 v_2^2 a_3 - 5v_3 v_1^2 v_2^3 a_3 + 2v_3^2 v_2^4 a_3 + v_1^6 b_2 - v_3^2 v_1^2 v_2^2 b_2 + 2v_1^5 v_2 b_3 \\ & + v_3 v_1^3 v_2^2 b_3 - v_3^2 v_1 v_2^3 b_3 - 3v_1^4 v_2 a_1 - v_3 v_1^2 v_2^2 a_1 + 2v_3^2 v_2^3 a_1 + v_1^5 b_1 \\ & - v_3^2 v_1 v_2^2 b_1 + 2v_1 v_2^3 a_2 - 2b_2 v_2^2 v_1^2 - v_1 v_2^3 b_3 + 4v_2^3 a_1 - 3v_1 v_2^2 b_1 = 0 \end{aligned} \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2, v_3\}$$

Equation (7E) now becomes

$$\begin{aligned} & -v_1^8 a_3 - 2v_3 v_1^6 v_2 a_3 + v_1^6 b_2 + (-4a_2 + 2b_3) v_1^5 v_2 + v_1^5 b_1 - v_3^2 v_1^4 v_2^2 a_3 \\ & - 7v_1^4 v_2^2 a_3 - 3v_1^4 v_2 a_1 + (-2a_2 + b_3) v_1^3 v_2^2 v_3 - 5v_3 v_1^2 v_2^3 a_3 - v_3^2 v_1^2 v_2^2 b_2 \\ & - v_3 v_1^2 v_2^2 a_1 - 2b_2 v_2^2 v_1^2 + (2a_2 - b_3) v_1 v_2^3 v_3^2 + (2a_2 - b_3) v_1 v_2^3 \\ & - v_3^2 v_1 v_2^2 b_1 - 3v_1 v_2^2 b_1 + 2v_3^2 v_2^4 a_3 + 2v_3^2 v_2^3 a_1 + 4v_2^3 a_1 = 0 \end{aligned} \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned}b_1 &= 0 \\b_2 &= 0 \\-3a_1 &= 0 \\-a_1 &= 0 \\2a_1 &= 0 \\4a_1 &= 0 \\-7a_3 &= 0 \\-5a_3 &= 0 \\-2a_3 &= 0 \\-a_3 &= 0 \\2a_3 &= 0 \\-3b_1 &= 0 \\-b_1 &= 0 \\-2b_2 &= 0 \\-b_2 &= 0 \\-4a_2 + 2b_3 &= 0 \\-2a_2 + b_3 &= 0 \\2a_2 - b_3 &= 0\end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned}a_1 &= 0 \\a_2 &= a_2 \\a_3 &= 0 \\b_1 &= 0 \\b_2 &= 0 \\b_3 &= 2a_2\end{aligned}$$

Substituting the above solution in the ansatz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned}\xi &= x \\ \eta &= 2y\end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned}\eta &= \eta - \omega(x, y) \xi \\ &= 2y - \left(\frac{x^2 \tan\left(\frac{y}{x^2}\right) y + x^4 + 2y^2}{xy} \right) (x) \\ &= \frac{-x^2 \tan\left(\frac{y}{x^2}\right) y - x^4}{y} \\ \xi &= 0\end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{-x^2 \tan\left(\frac{y}{x^2}\right) y - x^4}{y}} dy\end{aligned}$$

Which results in

$$S = -\ln \left(\cos\left(\frac{y}{x^2}\right) + \frac{\sin\left(\frac{y}{x^2}\right) y}{x^2} \right)$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y) S_y}{R_x + \omega(x, y) R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{x^2 \tan\left(\frac{y}{x^2}\right) y + x^4 + 2y^2}{xy}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{2y^2 \cos\left(\frac{y}{x^2}\right)}{x^3 \left(\cos\left(\frac{y}{x^2}\right) x^2 + \sin\left(\frac{y}{x^2}\right) y\right)} \\ S_y &= -\frac{\cos\left(\frac{y}{x^2}\right) y}{x^2 \left(\cos\left(\frac{y}{x^2}\right) x^2 + \sin\left(\frac{y}{x^2}\right) y\right)} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = -\frac{1}{x} \tag{2A}$$



We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = -\frac{1}{R}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = -\ln(R) + c_1 \tag{4}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{x^2 \tan\left(\frac{y}{x^2}\right) y + x^4 + 2y^2}{xy}$ 	$R = x$ $S = -\ln\left(\cos\left(\frac{y}{x^2}\right) x^2\right) + \dots$	$\frac{dS}{dR} = -\frac{1}{R}$ 

Summary

The solution(s) found are the following

$$-\ln\left(\cos\left(\frac{y}{x^2}\right) x^2 + \sin\left(\frac{y}{x^2}\right) y\right) + 2\ln(x) = -\ln(x) + c_1 \quad (1)$$




Figure 192: Slope field plot

Verification of solutions

$$-\ln\left(\cos\left(\frac{y}{x^2}\right)x^2 + \sin\left(\frac{y}{x^2}\right)y\right) + 2\ln(x) = -\ln(x) + c_1$$

Verified OK.

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying homogeneous G
<- homogeneous successful`

```

✓ Solution by Maple

Time used: 1.157 (sec). Leaf size: 176

```
dsolve(diff(y(x),x)=2*y(x)/x+x^3/y(x)+x*tan(y(x)/x^2),y(x), singsol=all)
```

$$y(x) = \frac{x^2 \left(\csc \left(\text{RootOf} \left(2c_1^2 \sin(_Z)^2 _Z^2 - 2c_1^2 \cos(_Z)^2 - 4c_1 \sin(_Z) x _Z + 2x^2 \right) \right) x - \cot \left(\text{RootOf} \left(2c_1^2 \sin(_Z)^2 _Z^2 - 2c_1^2 \cos(_Z)^2 - 4c_1 \sin(_Z) x _Z + 2x^2 \right) \right) c_1}{c_1}$$

$$y(x) = \frac{\left(\cot \left(\text{RootOf} \left(2c_1^2 \sin(_Z)^2 _Z^2 - 2c_1^2 \cos(_Z)^2 - 4c_1 \sin(_Z) x _Z + 2x^2 \right) \right) c_1 + \csc \left(\text{RootOf} \left(2c_1^2 \sin(_Z)^2 _Z^2 - 2c_1^2 \cos(_Z)^2 - 4c_1 \sin(_Z) x _Z + 2x^2 \right) \right) c_1}{c_1}$$

✓ Solution by Mathematica

Time used: 1.103 (sec). Leaf size: 36

```
DSolve[y'[x]==2*y[x]/x+x^3/y[x]+x*Tan[y[x]/x^2],y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[3 \log(x) - \log \left(y(x) \sin \left(\frac{y(x)}{x^2} \right) + x^2 \cos \left(\frac{y(x)}{x^2} \right) \right) = c_1, y(x) \right]$$

7 Chapter 1. What is a differential equation.

Section 1.9. Reduction of Order. Page 38

7.1	problem 1(a)	933
7.2	problem 1(b)	941
7.3	problem 1(c)	944
7.4	problem 1(d)	953
7.5	problem 1(e)	957
7.6	problem 1(f)	963
7.7	problem 1(g)	967
7.8	problem 2(a)	986
7.9	problem 2(b)	998
7.10	problem 2(c)	1006
7.11	problem 3(a)	1020
7.12	problem 3(b)	1025

7.1 problem 1(a)

7.1.1	Solving as second order integrable as is ode	933
7.1.2	Solving as second order ode missing x ode	934
7.1.3	Solving as type second_order_integrable_as_is (not using ABC version)	936
7.1.4	Solving as exact nonlinear second order ode ode	937
7.1.5	Maple step by step solution	938

Internal problem ID [6237]

Internal file name [OUTPUT/5485_Sunday_June_05_2022_03_41_08_PM_41183519/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.9. Reduction of Order. Page 38

Problem number: 1(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_integrable_as_is", "second_order_ode_missing_x", "exact nonlinear second order ode"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x], [_2nd_order, _exact, _nonlinear],  
  _Liouville, [_2nd_order, _reducible, _mu_x_y1], [_2nd_order,  
  _reducible, _mu_xy]]
```

$$yy'' + y'^2 = 0$$

7.1.1 Solving as second order integrable as is ode

Integrating both sides of the ODE w.r.t x gives

$$\int (yy'' + y'^2) dx = 0$$
$$y'y = c_1$$

Which is now solved for y . Integrating both sides gives

$$\int \frac{y}{c_1} dy = c_2 + x$$
$$\frac{y^2}{2c_1} = c_2 + x$$

Solving for y gives these solutions

$$y_1 = \sqrt{2c_1c_2 + 2c_1x}$$
$$y_2 = -\sqrt{2c_1c_2 + 2c_1x}$$

Summary

The solution(s) found are the following

$$y = \sqrt{2c_1c_2 + 2c_1x} \quad (1)$$

$$y = -\sqrt{2c_1c_2 + 2c_1x} \quad (2)$$

Verification of solutions

$$y = \sqrt{2c_1c_2 + 2c_1x}$$

Verified OK.

$$y = -\sqrt{2c_1c_2 + 2c_1x}$$

Verified OK.

7.1.2 Solving as second order ode missing x ode

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable y an independent variable. Using

$$y' = p(y)$$

Then

$$y'' = \frac{dp}{dx}$$
$$= \frac{dy}{dx} \frac{dp}{dy}$$
$$= p \frac{dp}{dy}$$

Hence the ode becomes

$$yp(y) \left(\frac{d}{dy} p(y) \right) + p(y)^2 = 0$$

Which is now solved as first order ode for $p(y)$. In canonical form the ODE is

$$\begin{aligned} p' &= F(y, p) \\ &= f(y)g(p) \\ &= -\frac{p}{y} \end{aligned}$$

Where $f(y) = -\frac{1}{y}$ and $g(p) = p$. Integrating both sides gives

$$\begin{aligned} \frac{1}{p} dp &= -\frac{1}{y} dy \\ \int \frac{1}{p} dp &= \int -\frac{1}{y} dy \\ \ln(p) &= -\ln(y) + c_1 \\ p &= e^{-\ln(y)+c_1} \\ &= \frac{c_1}{y} \end{aligned}$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = \frac{c_1}{y}$$

Integrating both sides gives

$$\begin{aligned} \int \frac{y}{c_1} dy &= c_2 + x \\ \frac{y^2}{2c_1} &= c_2 + x \end{aligned}$$

Solving for y gives these solutions

$$\begin{aligned} y_1 &= \sqrt{2c_1c_2 + 2c_1x} \\ y_2 &= -\sqrt{2c_1c_2 + 2c_1x} \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \sqrt{2c_1c_2 + 2c_1x} \tag{1}$$

$$y = -\sqrt{2c_1c_2 + 2c_1x} \tag{2}$$

Verification of solutions

$$y = \sqrt{2c_1c_2 + 2c_1x}$$

Verified OK.

$$y = -\sqrt{2c_1c_2 + 2c_1x}$$

Verified OK.

7.1.3 Solving as type second_order_integrable_as_is (not using ABC version)

Writing the ode as

$$yy'' + y'^2 = 0$$

Integrating both sides of the ODE w.r.t x gives

$$\int (yy'' + y'^2) dx = 0$$

$$y'y = c_1$$

Which is now solved for y . Integrating both sides gives

$$\int \frac{y}{c_1} dy = c_2 + x$$

$$\frac{y^2}{2c_1} = c_2 + x$$

Solving for y gives these solutions

$$y_1 = \sqrt{2c_1c_2 + 2c_1x}$$

$$y_2 = -\sqrt{2c_1c_2 + 2c_1x}$$

Summary

The solution(s) found are the following

$$y = \sqrt{2c_1c_2 + 2c_1x} \tag{1}$$

$$y = -\sqrt{2c_1c_2 + 2c_1x} \tag{2}$$

Verification of solutions

$$y = \sqrt{2c_1c_2 + 2c_1x}$$

Verified OK.

$$y = -\sqrt{2c_1c_2 + 2c_1x}$$

Verified OK.

7.1.4 Solving as exact nonlinear second order ode

An exact non-linear second order ode has the form

$$a_2(x, y, y') y'' + a_1(x, y, y') y' + a_0(x, y, y') = 0$$

Where the following conditions are satisfied

$$\begin{aligned}\frac{\partial a_2}{\partial y} &= \frac{\partial a_1}{\partial y'} \\ \frac{\partial a_2}{\partial x} &= \frac{\partial a_0}{\partial y'} \\ \frac{\partial a_1}{\partial x} &= \frac{\partial a_0}{\partial y}\end{aligned}$$

Looking at the the ode given we see that

$$\begin{aligned}a_2 &= y \\ a_1 &= y' \\ a_0 &= 0\end{aligned}$$

Applying the conditions to the above shows this is a nonlinear exact second order ode. Therefore it can be reduced to first order ode given by

$$\begin{aligned}\int a_2 dy' + \int a_1 dy + \int a_0 dx &= c_1 \\ \int y dy' + \int y' dy + \int 0 dx &= c_1\end{aligned}$$

Which results in

$$2y'y = c_1$$

Which is now solved Integrating both sides gives

$$\begin{aligned}\int \frac{2y}{c_1} dy &= c_2 + x \\ \frac{y^2}{c_1} &= c_2 + x\end{aligned}$$

Solving for y gives these solutions

$$\begin{aligned}y_1 &= \sqrt{c_1 c_2 + c_1 x} \\ y_2 &= -\sqrt{c_1 c_2 + c_1 x}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \sqrt{c_1 c_2 + c_1 x} \quad (1)$$

$$y = -\sqrt{c_1 c_2 + c_1 x} \quad (2)$$

Verification of solutions

$$y = \sqrt{c_1 c_2 + c_1 x}$$

Verified OK.

$$y = -\sqrt{c_1 c_2 + c_1 x}$$

Verified OK.

7.1.5 Maple step by step solution

Let's solve

$$yy'' + y'^2 = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Define new dependent variable u

$$u(x) = y'$$

- Compute y''

$$u'(x) = y''$$

- Use chain rule on the lhs

$$y' \left(\frac{d}{dy} u(y) \right) = y''$$

- Substitute in the definition of u

$$u(y) \left(\frac{d}{dy} u(y) \right) = y''$$

- Make substitutions $y' = u(y)$, $y'' = u(y) \left(\frac{d}{dy} u(y) \right)$ to reduce order of ODE

$$yu(y) \left(\frac{d}{dy} u(y) \right) + u(y)^2 = 0$$

- Separate variables

$$\frac{\frac{d}{dy} u(y)}{u(y)} = -\frac{1}{y}$$

- Integrate both sides with respect to y

$$\int \frac{\frac{d}{dy}u(y)}{u(y)} dy = \int -\frac{1}{y} dy + c_1$$

- Evaluate integral

$$\ln(u(y)) = -\ln(y) + c_1$$

- Solve for $u(y)$

$$u(y) = \frac{e^{c_1}}{y}$$

- Solve 1st ODE for $u(y)$

$$u(y) = \frac{e^{c_1}}{y}$$

- Revert to original variables with substitution $u(y) = y', y = y$

$$y' = \frac{e^{c_1}}{y}$$

- Separate variables

$$y'y = e^{c_1}$$

- Integrate both sides with respect to x

$$\int y'y dx = \int e^{c_1} dx + c_2$$

- Evaluate integral

$$\frac{y^2}{2} = x e^{c_1} + c_2$$

- Solve for y

$$\{y = \sqrt{2x e^{c_1} + 2c_2}, y = -\sqrt{2x e^{c_1} + 2c_2}\}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
<- 2nd_order Liouville successful`

```


✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 33

```
dsolve(y(x)*diff(y(x),x$2)+(diff(y(x),x))^2=0,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = \sqrt{2c_1x + 2c_2}$$

$$y(x) = -\sqrt{2c_1x + 2c_2}$$

✓ Solution by Mathematica

Time used: 0.166 (sec). Leaf size: 20

```
DSolve[y[x]*y'[x]+(y'[x])^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2\sqrt{2x - c_1}$$

7.2 problem 1(b)

Internal problem ID [6238]

Internal file name [OUTPUT/5486_Sunday_June_05_2022_03_41_11_PM_32735869/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.9. Reduction of Order. Page 38

Problem number: 1(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**unknown**"

Maple gives the following as the ode type


[NONE]

Unable to solve or complete the solution.

$$xyy'' - y' - y'^3 = 0$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
--- trying a change of variables {x -> y(x), y(x) -> x}
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
-> trying 2nd order, dynamical_symmetries, fully reducible to Abel through one integrating f
    --- trying a change of variables {x -> y(x), y(x) -> x} and re-entering methods for dynam
    -> trying 2nd order, dynamical_symmetries, fully reducible to Abel through one integratin
trying 2nd order, integrating factors of the form mu(x,y)/(y)^n, only the singular cases
trying symmetries linear in x and y(x)
trying differential order: 2; exact nonlinear
trying 2nd order, integrating factor of the form mu(y)
trying 2nd order, integrating factor of the form mu(x,y)
trying 2nd order, integrating factor of the form mu(x,y)/(y)^n, only the general case
trying 2nd order, integrating factor of the form mu(y,y)
trying differential order: 2; mu polynomial in y
trying 2nd order, integrating factor of the form mu(x,y)
differential order: 2; looking for linear symmetries
-> trying 2nd order, the S-function method
    -> trying a change of variables {x -> y(x), y(x) -> x} and re-entering methods for the S-
    -> trying 2nd order, the S-function method
    -> trying 2nd order, No Point Symmetries Class V
        --- trying a change of variables {x -> y(x), y(x) -> x} and re-entering methods for dy
        -> trying 2nd order, No Point Symmetries Class V
    -> trying 2nd order, No Point Symmetries Class V
        --- trying a change of variables {x -> y(x), y(x) -> x} and re-entering methods for dy
        -> trying 2nd order, No Point Symmetries Class V
    -> trying 2nd order, No Point Symmetries Class V
        --- trying a change of variables {x -> y(x), y(x) -> x} and re-entering methods for dy
        -> trying 2nd order, No Point Symmetries Class V
trying 2nd order, integrating factor of the form mu(x,y)/(y)^n, only the general case
-> trying 2nd order, dynamical_symmetries, only a reduction of order through one integrating
    --- trying a change of variables {x -> y(x), y(x) -> x} and re-entering methods for dynam
    -> trying 2nd order, dynamical_symmetries, only a reduction of order through one integrat
--- Trying Lie symmetry methods, 2nd order ---
`, `-> Computing symmetries using: way = 3
`, `-> Computing symmetries using: way = 5
`, `-> Computing symmetries using: way = formal`
```

 Solution by Maple

```
dsolve(x*y(x)*diff(y(x),x$2)=diff(y(x),x)+(diff(y(x),x))^3,y(x), singsol=all)
```

No solution found

 Solution by Mathematica

Time used: 1.417 (sec). Leaf size: 103

```
DSolve[x*y'[x]==y'[x]+(y'[x])^3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 - ie^{-c_1} \sqrt{-1 + e^{2c_1} x^2}$$

$$y(x) \rightarrow ie^{-c_1} \sqrt{-1 + e^{2c_1} x^2} + c_2$$

$$y(x) \rightarrow c_2 - i\sqrt{x^2}$$

$$y(x) \rightarrow i\sqrt{x^2} + c_2$$

7.3 problem 1(c)

7.3.1	Solving as second order linear constant coeff ode	944
7.3.2	Solving as second order ode can be made integrable ode	946
7.3.3	Solving using Kovacic algorithm	947
7.3.4	Maple step by step solution	951

Internal problem ID [6239]

Internal file name [OUTPUT/5487_Sunday_June_05_2022_03_41_13_PM_65395190/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.9. Reduction of Order. Page 38

Problem number: 1(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "second_order_ode_can_be_made_integrable"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x]]
```

$$y'' - k^2y = 0$$

7.3.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = -k^2$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - k^2 e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$-k^2 + \lambda^2 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = -k^2$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(-k^2)} \\ &= \pm \sqrt{k^2} \end{aligned}$$

Hence

$$\lambda_1 = +\sqrt{k^2}$$

$$\lambda_2 = -\sqrt{k^2}$$

Which simplifies to

$$\lambda_1 = \sqrt{k^2}$$

$$\lambda_2 = -\sqrt{k^2}$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(\sqrt{k^2})x} + c_2 e^{(-\sqrt{k^2})x}$$

Or

$$y = c_1 e^{\sqrt{k^2}x} + c_2 e^{-\sqrt{k^2}x}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{\sqrt{k^2}x} + c_2 e^{-\sqrt{k^2}x} \quad (1)$$

Verification of solutions

$$y = c_1 e^{\sqrt{k^2}x} + c_2 e^{-\sqrt{k^2}x}$$

Verified OK.

7.3.2 Solving as second order ode can be made integrable ode

Multiplying the ode by y' gives

$$y'y'' - y'k^2y = 0$$

Integrating the above w.r.t x gives

$$\int (y'y'' - y'k^2y) dx = 0$$

$$\frac{y'^2}{2} - \frac{y^2k^2}{2} = c_2$$

Which is now solved for y . Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \sqrt{y^2k^2 + 2c_1} \quad (1)$$

$$y' = -\sqrt{y^2k^2 + 2c_1} \quad (2)$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int \frac{1}{\sqrt{y^2k^2 + 2c_1}} dy = \int dx$$

$$\frac{\ln\left(\frac{k^2y}{\sqrt{k^2} + \sqrt{y^2k^2 + 2c_1}}\right)}{\sqrt{k^2}} = c_2 + x$$

Raising both side to exponential gives

$$e^{\frac{\ln\left(\frac{k^2y}{\sqrt{k^2} + \sqrt{y^2k^2 + 2c_1}}\right)}{\sqrt{k^2}}} = e^{c_2 + x}$$

Which simplifies to

$$\left(k \operatorname{csgn}(k) y + \sqrt{y^2k^2 + 2c_1}\right)^{\frac{1}{\sqrt{k^2}}} = c_3 e^x$$

Simplifying the solution $y = -\frac{\operatorname{csgn}(k)\left(2(c_3e^x)^{-\operatorname{csgn}(k)k}c_1 - (c_3e^x)^{\operatorname{csgn}(k)k}\right)}{2k}$ to $y = -\frac{2(c_3e^x)^{-k}c_1 - (c_3e^x)^k}{2k}$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{1}{\sqrt{y^2 k^2 + 2c_1}} dy = \int dx$$

$$-\frac{\ln\left(\frac{k^2 y}{\sqrt{k^2}} + \sqrt{y^2 k^2 + 2c_1}\right)}{\sqrt{k^2}} = x + c_4$$

Raising both side to exponential gives

$$e^{-\frac{\ln\left(\frac{k^2 y}{\sqrt{k^2}} + \sqrt{y^2 k^2 + 2c_1}\right)}{\sqrt{k^2}}} = e^{x+c_4}$$

Which simplifies to

$$\left(k \operatorname{csgn}(k) y + \sqrt{y^2 k^2 + 2c_1}\right)^{-\frac{\operatorname{csgn}(k)}{k}} = c_5 e^x$$

Simplifying the solution $y = \frac{\operatorname{csgn}(k)\left((c_5 e^x)^{-\operatorname{csgn}(k)k} - 2(c_5 e^x)^{\operatorname{csgn}(k)k} c_1\right)}{2k}$ to $y = \frac{(c_5 e^x)^{-k} - 2(c_5 e^x)^k c_1}{2k}$

Summary

The solution(s) found are the following

$$y = -\frac{2(c_3 e^x)^{-k} c_1 - (c_3 e^x)^k}{2k} \quad (1)$$

$$y = \frac{(c_5 e^x)^{-k} - 2(c_5 e^x)^k c_1}{2k} \quad (2)$$

Verification of solutions

$$y = -\frac{2(c_3 e^x)^{-k} c_1 - (c_3 e^x)^k}{2k}$$

Verified OK.

$$y = \frac{(c_5 e^x)^{-k} - 2(c_5 e^x)^k c_1}{2k}$$

Verified OK.

7.3.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' - k^2 y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= 1 \\B &= 0 \\C &= -k^2\end{aligned}\tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x)\tag{4}$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\&= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned}\tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{k^2}{1}\tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= k^2 \\t &= 1\end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = (k^2) z(x)\tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 125: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = k^2$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{\sqrt{k^2}x}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}
 y_1 &= z_1 \\
 &= e^{\sqrt{k^2}x}
 \end{aligned}$$

Which simplifies to

$$y_1 = e^{\sqrt{k^2} x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= e^{\sqrt{k^2} x} \int \frac{1}{e^{2\sqrt{k^2} x}} dx \\ &= e^{\sqrt{k^2} x} \left(-\frac{\operatorname{csgn}(k) e^{-2kx \operatorname{csgn}(k)}}{2k} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(e^{\sqrt{k^2} x} \right) + c_2 \left(e^{\sqrt{k^2} x} \left(-\frac{\operatorname{csgn}(k) e^{-2kx \operatorname{csgn}(k)}}{2k} \right) \right) \end{aligned}$$

Simplifying the solution $y = c_1 e^{\sqrt{k^2} x} - \frac{c_2 \operatorname{csgn}(k) e^{-kx \operatorname{csgn}(k)}}{2k}$ to $y = c_1 e^{\sqrt{k^2} x} - \frac{c_2 e^{-kx}}{2k}$

Summary

The solution(s) found are the following

$$y = c_1 e^{\sqrt{k^2} x} - \frac{c_2 e^{-kx}}{2k} \quad (1)$$

Verification of solutions

$$y = c_1 e^{\sqrt{k^2} x} - \frac{c_2 e^{-kx}}{2k}$$

Verified OK.

7.3.4 Maple step by step solution

Let's solve

$$y'' - k^2y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$-k^2 + r^2 = 0$$

- Factor the characteristic polynomial

$$-(k - r)(k + r) = 0$$

- Roots of the characteristic polynomial

$$r = (k, -k)$$

- 1st solution of the ODE

$$y_1(x) = e^{kx}$$

- 2nd solution of the ODE

$$y_2(x) = e^{-kx}$$

- General solution of the ODE

$$y = c_1y_1(x) + c_2y_2(x)$$

- Substitute in solutions

$$y = c_1e^{kx} + c_2e^{-kx}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(diff(y(x),x$2)-k^2*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{-kx} + c_2 e^{kx}$$

✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 23

```
DSolve[y''[x]-k^2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{kx} + c_2 e^{-kx}$$

7.4 problem 1(d)

7.4.1 Solving as second order ode missing y ode 953

Internal problem ID [6240]

Internal file name [OUTPUT/5488_Sunday_June_05_2022_03_41_15_PM_99919298/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.9. Reduction of Order. Page 38

Problem number: 1(d).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_ode_missing_y"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_y]]
```

$$x^2y'' - 2xy' - y'^2 = 0$$

7.4.1 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$x^2p'(x) + (-2x - p(x))p(x) = 0$$

Which is now solve for $p(x)$ as first order ode. Using the change of variables $p(x) = u(x) x$ on the above ode results in new ode in $u(x)$

$$x^2(u'(x) x + u(x)) + (-2x - u(x) x) u(x) x = 0$$

In canonical form the ODE is

$$\begin{aligned}u' &= F(x, u) \\ &= f(x)g(u) \\ &= \frac{u(u+1)}{x}\end{aligned}$$

Where $f(x) = \frac{1}{x}$ and $g(u) = u(u+1)$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u(u+1)} du &= \frac{1}{x} dx \\ \int \frac{1}{u(u+1)} du &= \int \frac{1}{x} dx \\ -\ln(u+1) + \ln(u) &= \ln(x) + c_2\end{aligned}$$

Raising both side to exponential gives

$$e^{-\ln(u+1)+\ln(u)} = e^{\ln(x)+c_2}$$

Which simplifies to

$$\frac{u}{u+1} = c_3x$$

Therefore the solution $p(x)$ is

$$\begin{aligned}p(x) &= xu \\ &= -\frac{x^2c_3}{c_3x-1}\end{aligned}$$

Since $p = y'$ then the new first order ode to solve is

$$y' = -\frac{x^2c_3}{c_3x-1}$$

Integrating both sides gives

$$\begin{aligned}y &= \int -\frac{x^2c_3}{c_3x-1} dx \\ &= -\frac{x^2}{2} - \frac{x}{c_3} - \frac{\ln(c_3x-1)}{c_3^2} + c_4\end{aligned}$$

Summary

The solution(s) found are the following

$$y = -\frac{x^2}{2} - \frac{x}{c_3} - \frac{\ln(c_3x - 1)}{c_3^2} + c_4 \quad (1)$$

Verification of solutions

$$y = -\frac{x^2}{2} - \frac{x}{c_3} - \frac{\ln(c_3x - 1)}{c_3^2} + c_4$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
-> Calling odsolve with the ODE`, diff(_b(_a), _a) = _b(_a)*(_b(_a)+2*_a)/_a^2, _b(_a), HINT
symmetry methods on request
`, `1st order, trying reduction of order with given symmetries: `[ _a, _b]
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 26

```
dsolve(x^2*diff(y(x),x$2)=2*x*diff(y(x),x)+(diff(y(x),x))^2,y(x), singsol=all)
```

$$y(x) = -\frac{x^2}{2} - c_1x - c_1^2 \ln(x - c_1) + c_2$$

✓ Solution by Mathematica

Time used: 0.435 (sec). Leaf size: 41

```
DSolve[x^2*y'[x]==2*x*y'[x]+(y'[x])^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{x^2}{2} - c_1 x - c_1^2 \log(x - c_1) + \frac{3c_1^2}{2} + c_2$$

7.5 problem 1(e)

7.5.1 Solving as second order ode missing x ode	957
7.5.2 Maple step by step solution	960

Internal problem ID [6241]

Internal file name [OUTPUT/5489_Sunday_June_05_2022_03_41_18_PM_49576723/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.9. Reduction of Order. Page 38

Problem number: 1(e).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_ode_missing_x**"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]]
```

$$2yy'' - y'^2 = 1$$

7.5.1 Solving as second order ode missing x ode

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable y an independent variable. Using

$$y' = p(y)$$

Then

$$\begin{aligned}y'' &= \frac{dp}{dx} \\ &= \frac{dy}{dx} \frac{dp}{dy} \\ &= p \frac{dp}{dy}\end{aligned}$$

Hence the ode becomes

$$2yp(y) \left(\frac{d}{dy} p(y) \right) - p(y)^2 = 1$$

Which is now solved as first order ode for $p(y)$. In canonical form the ODE is

$$\begin{aligned} p' &= F(y, p) \\ &= f(y)g(p) \\ &= \frac{p^2 + 1}{2yp} \end{aligned}$$

Where $f(y) = \frac{1}{2y}$ and $g(p) = \frac{p^2+1}{p}$. Integrating both sides gives

$$\begin{aligned} \frac{1}{\frac{p^2+1}{p}} dp &= \frac{1}{2y} dy \\ \int \frac{1}{\frac{p^2+1}{p}} dp &= \int \frac{1}{2y} dy \\ \frac{\ln(p^2 + 1)}{2} &= \frac{\ln(y)}{2} + c_1 \end{aligned}$$

Raising both side to exponential gives

$$\sqrt{p^2 + 1} = e^{\frac{\ln(y)}{2} + c_1}$$

Which simplifies to

$$\sqrt{p^2 + 1} = c_2 \sqrt{y}$$

Which simplifies to

$$\sqrt{p(y)^2 + 1} = c_2 \sqrt{y} e^{c_1}$$

The solution is

$$\sqrt{p(y)^2 + 1} = c_2 \sqrt{y} e^{c_1}$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$\sqrt{1 + y'^2} = c_2 \sqrt{y} e^{c_1}$$

Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \sqrt{-1 + c_2^2 y e^{2c_1}} \tag{1}$$

$$y' = -\sqrt{-1 + c_2^2 y e^{2c_1}} \tag{2}$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int \frac{1}{\sqrt{-1 + c_2^2 y e^{2c_1}}} dy = \int dx$$
$$\frac{2\sqrt{-1 + c_2^2 y e^{2c_1}} e^{-2c_1}}{c_2^2} = x + c_3$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{1}{\sqrt{-1 + c_2^2 y e^{2c_1}}} dy = \int dx$$
$$-\frac{2\sqrt{-1 + c_2^2 y e^{2c_1}} e^{-2c_1}}{c_2^2} = x + c_4$$

Summary

The solution(s) found are the following

$$y = \frac{(c_3^2 e^{4c_1} c_2^4 + 2c_3 e^{4c_1} c_2^4 x + e^{4c_1} c_2^4 x^2 + 4) e^{-2c_1}}{4c_2^2} \quad (1)$$

$$y = \frac{(c_4^2 e^{4c_1} c_2^4 + 2c_4 e^{4c_1} c_2^4 x + e^{4c_1} c_2^4 x^2 + 4) e^{-2c_1}}{4c_2^2} \quad (2)$$

Verification of solutions

$$y = \frac{(c_3^2 e^{4c_1} c_2^4 + 2c_3 e^{4c_1} c_2^4 x + e^{4c_1} c_2^4 x^2 + 4) e^{-2c_1}}{4c_2^2}$$

Verified OK.

$$y = \frac{(c_4^2 e^{4c_1} c_2^4 + 2c_4 e^{4c_1} c_2^4 x + e^{4c_1} c_2^4 x^2 + 4) e^{-2c_1}}{4c_2^2}$$

Verified OK.

7.5.2 Maple step by step solution

Let's solve

$$2yy'' - y'^2 = 1$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Define new dependent variable u

$$u(x) = y'$$

- Compute y''

$$u'(x) = y''$$

- Use chain rule on the lhs

$$y' \left(\frac{d}{dy} u(y) \right) = y''$$

- Substitute in the definition of u

$$u(y) \left(\frac{d}{dy} u(y) \right) = y''$$

- Make substitutions $y' = u(y)$, $y'' = u(y) \left(\frac{d}{dy} u(y) \right)$ to reduce order of ODE

$$2yu(y) \left(\frac{d}{dy} u(y) \right) - u(y)^2 = 1$$

- Separate variables

$$\frac{\left(\frac{d}{dy} u(y) \right) u(y)}{u(y)^2 + 1} = \frac{1}{2y}$$

- Integrate both sides with respect to y

$$\int \frac{\left(\frac{d}{dy} u(y) \right) u(y)}{u(y)^2 + 1} dy = \int \frac{1}{2y} dy + c_1$$

- Evaluate integral

$$\frac{\ln(u(y)^2 + 1)}{2} = \frac{\ln(y)}{2} + c_1$$

- Solve for $u(y)$

$$\left\{ u(y) = \frac{\sqrt{-e^{-2c_1}(e^{-2c_1} - y)}}{e^{-2c_1}}, u(y) = -\frac{\sqrt{-e^{-2c_1}(e^{-2c_1} - y)}}{e^{-2c_1}} \right\}$$

- Solve 1st ODE for $u(y)$

$$u(y) = \frac{\sqrt{-e^{-2c_1}(e^{-2c_1} - y)}}{e^{-2c_1}}$$

- Revert to original variables with substitution $u(y) = y'$, $y = y$

$$y' = \frac{\sqrt{-e^{-2c_1}(e^{-2c_1}-y)}}{e^{-2c_1}}$$

- Separate variables

$$\frac{y'}{\sqrt{-e^{-2c_1}(e^{-2c_1}-y)}} = \frac{1}{e^{-2c_1}}$$

- Integrate both sides with respect to x

$$\int \frac{y'}{\sqrt{-e^{-2c_1}(e^{-2c_1}-y)}} dx = \int \frac{1}{e^{-2c_1}} dx + c_2$$

- Evaluate integral

$$\frac{2\sqrt{e^{-2c_1}y-(e^{-2c_1})^2}}{e^{-2c_1}} = \frac{x}{e^{-2c_1}} + c_2$$

- Solve for y

$$y = \frac{c_2^2(e^{-2c_1})^2 + 2c_2e^{-2c_1}x + 4(e^{-2c_1})^2 + x^2}{4e^{-2c_1}}$$

- Solve 2nd ODE for $u(y)$

$$u(y) = -\frac{\sqrt{-e^{-2c_1}(e^{-2c_1}-y)}}{e^{-2c_1}}$$

- Revert to original variables with substitution $u(y) = y', y = y$

$$y' = -\frac{\sqrt{-e^{-2c_1}(e^{-2c_1}-y)}}{e^{-2c_1}}$$

- Separate variables

$$\frac{y'}{\sqrt{-e^{-2c_1}(e^{-2c_1}-y)}} = -\frac{1}{e^{-2c_1}}$$

- Integrate both sides with respect to x

$$\int \frac{y'}{\sqrt{-e^{-2c_1}(e^{-2c_1}-y)}} dx = \int -\frac{1}{e^{-2c_1}} dx + c_2$$

- Evaluate integral

$$\frac{2\sqrt{e^{-2c_1}y-(e^{-2c_1})^2}}{e^{-2c_1}} = -\frac{x}{e^{-2c_1}} + c_2$$

- Solve for y

$$y = \frac{c_2^2(e^{-2c_1})^2 - 2c_2e^{-2c_1}x + 4(e^{-2c_1})^2 + x^2}{4e^{-2c_1}}$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
-> Calling odsolve with the ODE`, diff(diff(diff(y(x), x), x), x), y(x)` *** Sublevel 2 **
  Methods for third order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  <- quadrature successful
<- 2nd order ODE linearizable_by_differentiation successful`
```

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 22

```
dsolve(2*y(x)*diff(y(x),x$2)=1+(diff(y(x),x))^2,y(x), singsol=all)
```

$$y(x) = \frac{(c_1^2 + 1)x^2}{4c_2} + c_1x + c_2$$

✓ Solution by Mathematica

Time used: 0.032 (sec). Leaf size: 34

```
DSolve[2*y[x]*y'[x]==1+(y'[x])^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{(1 + c_1^2)x^2}{4c_2} + c_1x + c_2$$
$$y(x) \rightarrow \text{Indeterminate}$$

7.6 problem 1(f)

7.6.1 Solving as second order ode missing x ode	963
7.6.2 Maple step by step solution	965

Internal problem ID [6242]

Internal file name [OUTPUT/5490_Sunday_June_05_2022_03_41_21_PM_68794890/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.9. Reduction of Order. Page 38

Problem number: 1(f).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_ode_missing_x**"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x], _Liouville, [_2nd_order, _reducible,
    _mu_x_y1], [_2nd_order, _reducible, _mu_xy]]
```

$$yy'' - y'^2 = 0$$

7.6.1 Solving as second order ode missing x ode

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable y an independent variable. Using

$$y' = p(y)$$

Then

$$\begin{aligned} y'' &= \frac{dp}{dx} \\ &= \frac{dy}{dx} \frac{dp}{dy} \\ &= p \frac{dp}{dy} \end{aligned}$$

Hence the ode becomes

$$yp(y) \left(\frac{d}{dy} p(y) \right) - p(y)^2 = 0$$

Which is now solved as first order ode for $p(y)$. In canonical form the ODE is

$$\begin{aligned} p' &= F(y, p) \\ &= f(y)g(p) \\ &= \frac{p}{y} \end{aligned}$$

Where $f(y) = \frac{1}{y}$ and $g(p) = p$. Integrating both sides gives

$$\begin{aligned} \frac{1}{p} dp &= \frac{1}{y} dy \\ \int \frac{1}{p} dp &= \int \frac{1}{y} dy \\ \ln(p) &= \ln(y) + c_1 \\ p &= e^{\ln(y)+c_1} \\ &= c_1 y \end{aligned}$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = yc_1$$

Integrating both sides gives

$$\begin{aligned} \int \frac{1}{c_1 y} dy &= \int dx \\ \frac{\ln(y)}{c_1} &= c_2 + x \end{aligned}$$

Raising both side to exponential gives

$$e^{\frac{\ln(y)}{c_1}} = e^{c_2+x}$$

Which simplifies to

$$y^{\frac{1}{c_1}} = c_3 e^x$$

Summary

The solution(s) found are the following

$$y = (c_3 e^x)^{c_1} \tag{1}$$

Verification of solutions

$$y = (c_3 e^x)^{c_1}$$

Verified OK.

7.6.2 Maple step by step solution

Let's solve

$$yy'' - y'^2 = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Define new dependent variable u

$$u(x) = y'$$

- Compute y''

$$u'(x) = y''$$

- Use chain rule on the lhs

$$y' \left(\frac{d}{dy} u(y) \right) = y''$$

- Substitute in the definition of u

$$u(y) \left(\frac{d}{dy} u(y) \right) = y''$$

- Make substitutions $y' = u(y)$, $y'' = u(y) \left(\frac{d}{dy} u(y) \right)$ to reduce order of ODE

$$yu(y) \left(\frac{d}{dy} u(y) \right) - u(y)^2 = 0$$

- Separate variables

$$\frac{\frac{d}{dy} u(y)}{u(y)} = \frac{1}{y}$$

- Integrate both sides with respect to y

$$\int \frac{\frac{d}{dy} u(y)}{u(y)} dy = \int \frac{1}{y} dy + c_1$$

- Evaluate integral

$$\ln(u(y)) = \ln(y) + c_1$$

- Solve for $u(y)$

$$u(y) = y e^{c_1}$$

- Solve 1st ODE for $u(y)$

$$u(y) = y e^{c_1}$$

- Revert to original variables with substitution $u(y) = y'$, $y = y$

$$y' = y e^{c_1}$$

- Separate variables

$$\frac{y'}{y} = e^{c_1}$$
- Integrate both sides with respect to x

$$\int \frac{y'}{y} dx = \int e^{c_1} dx + c_2$$
- Evaluate integral

$$\ln(y) = x e^{c_1} + c_2$$
- Solve for y

$$y = e^{x e^{c_1} + c_2}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
<- 2nd_order Liouville successful`

```

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 14

```
dsolve(y(x)*diff(y(x),x$2)-(diff(y(x),x))^2=0,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = e^{c_1 x} c_2$$

✓ Solution by Mathematica

Time used: 0.101 (sec). Leaf size: 14

```
DSolve[y[x]*y'[x]-(y'[x])^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 e^{c_1 x}$$

7.7 problem 1(g)

7.7.1	Solving as second order integrable as is ode	968
7.7.2	Solving as second order ode missing y ode	968
7.7.3	Solving as second order ode non constant coeff transformation on B ode	970
7.7.4	Solving as type second_order_integrable_as_is (not using ABC version)	974
7.7.5	Solving using Kovacic algorithm	975
7.7.6	Solving as exact linear second order ode ode	982
7.7.7	Maple step by step solution	983

Internal problem ID [6243]

Internal file name [OUTPUT/5491_Sunday_June_05_2022_03_41_23_PM_37012367/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.9. Reduction of Order. Page 38

Problem number: 1(g).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "exact linear second order ode", "second_order_integrable_as_is", "second_order_ode_missing_y", "second_order_ode_non_constant_coeff_transformation_on_B"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_y]]
```

$$xy'' + y' = 4x$$

7.7.1 Solving as second order integrable as is ode

Integrating both sides of the ODE w.r.t x gives

$$\int (xy'' + y') dx = \int 4x dx$$
$$xy' = 2x^2 + c_1$$

Which is now solved for y . Integrating both sides gives

$$y = \int \frac{2x^2 + c_1}{x} dx$$
$$= x^2 + c_1 \ln(x) + c_2$$

Summary

The solution(s) found are the following

$$y = x^2 + c_1 \ln(x) + c_2 \quad (1)$$

Verification of solutions

$$y = x^2 + c_1 \ln(x) + c_2$$

Verified OK.

7.7.2 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x)x + p(x) - 4x = 0$$

Which is now solve for $p(x)$ as first order ode.

Entering Linear first order ODE solver. In canonical form a linear first order is

$$p'(x) + p(x)p(x) = q(x)$$

Where here

$$p(x) = \frac{1}{x}$$
$$q(x) = 4$$

Hence the ode is

$$p'(x) + \frac{p(x)}{x} = 4$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int \frac{1}{x} dx} \\ &= x\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu p) &= (\mu) (4) \\ \frac{d}{dx}(xp) &= (x) (4) \\ d(xp) &= (4x) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}xp &= \int 4x dx \\ xp &= 2x^2 + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = x$ results in

$$p(x) = 2x + \frac{c_1}{x}$$

Since $p = y'$ then the new first order ode to solve is

$$y' = 2x + \frac{c_1}{x}$$

Integrating both sides gives

$$\begin{aligned}y &= \int \frac{2x^2 + c_1}{x} dx \\ &= x^2 + c_1 \ln(x) + c_2\end{aligned}$$

Summary

The solution(s) found are the following

$$y = x^2 + c_1 \ln(x) + c_2 \tag{1}$$

Verification of solutions

$$y = x^2 + c_1 \ln(x) + c_2$$

Verified OK.

7.7.3 Solving as second order ode non constant coeff transformation on B ode

Given an ode of the form

$$Ay'' + By' + Cy = F(x)$$

This method reduces the order ode the ODE by one by applying the transformation

$$y = Bv$$

This results in

$$\begin{aligned} y' &= B'v + v'B \\ y'' &= B''v + B'v' + v''B + v'B' \\ &= v''B + 2v' + B' + B''v \end{aligned}$$

And now the original ode becomes

$$\begin{aligned} A(v''B + 2v'B' + B''v) + B(B'v + v'B) + CBv &= 0 \\ ABv'' + (2AB' + B^2)v' + (AB'' + BB' + CB)v &= 0 \end{aligned} \quad (1)$$

If the term $AB'' + BB' + CB$ is zero, then this method works and can be used to solve

$$ABv'' + (2AB' + B^2)v' = 0$$

By Using $u = v'$ which reduces the order of the above ode to one. The new ode is

$$ABu' + (2AB' + B^2)u = 0$$

The above ode is first order ode which is solved for u . Now a new ode $v' = u$ is solved for v as first order ode. Then the final solution is obtain from $y = Bv$.

This method works only if the term $AB'' + BB' + CB$ is zero. The given ODE shows that

$$\begin{aligned} A &= x \\ B &= 1 \\ C &= 0 \\ F &= 4x \end{aligned}$$

The above shows that for this ode

$$\begin{aligned} AB'' + BB' + CB &= (x)(0) + (1)(0) + (0)(1) \\ &= 0 \end{aligned}$$

Hence the ode in v given in (1) now simplifies to

$$xv'' + (1)v' = 0$$

Now by applying $v' = u$ the above becomes

$$xu'(x) + u(x) = 0$$

Which is now solved for u . In canonical form the ODE is

$$\begin{aligned}u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{u}{x}\end{aligned}$$

Where $f(x) = -\frac{1}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u} du &= -\frac{1}{x} dx \\ \int \frac{1}{u} du &= \int -\frac{1}{x} dx \\ \ln(u) &= -\ln(x) + c_1 \\ u &= e^{-\ln(x)+c_1} \\ &= \frac{c_1}{x}\end{aligned}$$

The ode for v now becomes

$$\begin{aligned}v' &= u \\ &= \frac{c_1}{x}\end{aligned}$$

Which is now solved for v . Integrating both sides gives

$$\begin{aligned}v(x) &= \int \frac{c_1}{x} dx \\ &= c_1 \ln(x) + c_2\end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}y_h(x) &= Bv \\ &= (1)(c_1 \ln(x) + c_2) \\ &= c_1 \ln(x) + c_2\end{aligned}$$

And now the particular solution $y_p(x)$ will be found. The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1y_1 + u_2y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned} y_1 &= 1 \\ y_2 &= \ln(x) \end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} 1 & \ln(x) \\ \frac{d}{dx}(1) & \frac{d}{dx}(\ln(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} 1 & \ln(x) \\ 0 & \frac{1}{x} \end{vmatrix}$$

Therefore

$$W = (1) \left(\frac{1}{x} \right) - (\ln(x))(0)$$

Which simplifies to

$$W = \frac{1}{x}$$

Which simplifies to

$$W = \frac{1}{x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{4 \ln(x) x}{1} dx$$

Which simplifies to

$$u_1 = - \int 4 \ln(x) x dx$$

Hence

$$u_1 = -2 \ln(x) x^2 + x^2$$

And Eq. (3) becomes

$$u_2 = \int \frac{4x}{1} dx$$

Which simplifies to

$$u_2 = \int 4x dx$$

Hence

$$u_2 = 2x^2$$

Which simplifies to

$$u_1 = x^2(1 - 2 \ln(x))$$

$$u_2 = 2x^2$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = x^2(1 - 2 \ln(x)) + 2 \ln(x) x^2$$

Which simplifies to

$$y_p(x) = x^2$$

Hence the complete solution is

$$\begin{aligned}y(x) &= y_h + y_p \\ &= (c_1 \ln(x) + c_2) + (x^2) \\ &= x^2 + c_1 \ln(x) + c_2\end{aligned}$$

Summary

The solution(s) found are the following

$$y = x^2 + c_1 \ln(x) + c_2 \quad (1)$$

Verification of solutions

$$y = x^2 + c_1 \ln(x) + c_2$$

Verified OK.

7.7.4 Solving as type second_order_integrable_as_is (not using ABC version)

Writing the ode as

$$xy'' + y' = 4x$$

Integrating both sides of the ODE w.r.t x gives

$$\begin{aligned}\int (xy'' + y') dx &= \int 4x dx \\ xy' &= 2x^2 + c_1\end{aligned}$$

Which is now solved for y . Integrating both sides gives

$$\begin{aligned}y &= \int \frac{2x^2 + c_1}{x} dx \\ &= x^2 + c_1 \ln(x) + c_2\end{aligned}$$

Summary

The solution(s) found are the following

$$y = x^2 + c_1 \ln(x) + c_2 \quad (1)$$

Verification of solutions

$$y = x^2 + c_1 \ln(x) + c_2$$

Verified OK.

7.7.5 Solving using Kovacic algorithm

Writing the ode as

$$xy'' + y' = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= x \\ B &= 1 \\ C &= 0 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{4x^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 4x^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(-\frac{1}{4x^2}\right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 129: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = -\frac{1}{4x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = -\frac{1}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{1}{2} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{2} \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ , which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = -\frac{1}{4x^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = -\frac{1}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{1}{2} \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{2} \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = -\frac{1}{4x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	$\frac{1}{2}$	$\frac{1}{2}$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	$\frac{1}{2}$	$\frac{1}{2}$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω .

Trying $\alpha_{\infty}^{-} = \frac{1}{2}$ then

$$\begin{aligned} d &= \alpha_{\infty}^{-} - (\alpha_{c_1}^{-}) \\ &= \frac{1}{2} - \left(\frac{1}{2}\right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x-c} \right) + s(\infty)[\sqrt{r}]_{\infty}$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^{-}}{x-c_1} \right) + (-)[\sqrt{r}]_{\infty} \\ &= \frac{1}{2x} + (-)(0) \\ &= \frac{1}{2x} \\ &= \frac{1}{2x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r)p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2\left(\frac{1}{2x}\right)(0) + \left(\left(-\frac{1}{2x^2}\right) + \left(\frac{1}{2x}\right)^2 - \left(-\frac{1}{4x^2}\right) \right) &= 0 \\ 0 &= 0 \end{aligned}$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= p e^{\int \omega dx} \\ &= e^{\int \frac{1}{2x} dx} \\ &= \sqrt{x} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\&= z_1 e^{-\int \frac{1}{2} \frac{1}{x} dx} \\&= z_1 e^{-\frac{\ln(x)}{2}} \\&= z_1 \left(\frac{1}{\sqrt{x}} \right)\end{aligned}$$

Which simplifies to

$$y_1 = 1$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{1}{x} dx}}{(y_1)^2} dx \\&= y_1 \int \frac{e^{-\ln(x)}}{(y_1)^2} dx \\&= y_1 (\ln(x))\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\&= c_1 (1) + c_2 (1(\ln(x)))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$xy'' + y' = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 + c_2 \ln(x)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned} y_1 &= 1 \\ y_2 &= \ln(x) \end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} 1 & \ln(x) \\ \frac{d}{dx}(1) & \frac{d}{dx}(\ln(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} 1 & \ln(x) \\ 0 & \frac{1}{x} \end{vmatrix}$$

Therefore

$$W = (1) \left(\frac{1}{x} \right) - (\ln(x)) (0)$$

Which simplifies to

$$W = \frac{1}{x}$$

Which simplifies to

$$W = \frac{1}{x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{4 \ln(x) x}{1} dx$$

Which simplifies to

$$u_1 = - \int 4 \ln(x) x dx$$

Hence

$$u_1 = -2 \ln(x) x^2 + x^2$$

And Eq. (3) becomes

$$u_2 = \int \frac{4x}{1} dx$$

Which simplifies to

$$u_2 = \int 4x dx$$

Hence

$$u_2 = 2x^2$$

Which simplifies to

$$u_1 = x^2(1 - 2 \ln(x))$$

$$u_2 = 2x^2$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = x^2(1 - 2 \ln(x)) + 2 \ln(x) x^2$$

Which simplifies to

$$y_p(x) = x^2$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 + c_2 \ln(x)) + (x^2) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 + c_2 \ln(x) + x^2 \quad (1)$$

Verification of solutions

$$y = c_1 + c_2 \ln(x) + x^2$$

Verified OK.

7.7.6 Solving as exact linear second order ode ode

An ode of the form

$$p(x) y'' + q(x) y' + r(x) y = s(x)$$

is exact if

$$p''(x) - q'(x) + r(x) = 0 \quad (1)$$

For the given ode we have

$$\begin{aligned} p(x) &= x \\ q(x) &= 1 \\ r(x) &= 0 \\ s(x) &= 4x \end{aligned}$$

Hence

$$\begin{aligned} p''(x) &= 0 \\ q'(x) &= 0 \end{aligned}$$

Therefore (1) becomes

$$0 - (0) + (0) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(x)y' + (q(x) - p'(x))y)' = s(x)$$

Integrating gives

$$p(x)y' + (q(x) - p'(x))y = \int s(x) dx$$

Substituting the above values for p, q, r, s gives

$$xy' = \int 4x dx$$

We now have a first order ode to solve which is

$$xy' = 2x^2 + c_1$$

Integrating both sides gives

$$\begin{aligned} y &= \int \frac{2x^2 + c_1}{x} dx \\ &= x^2 + c_1 \ln(x) + c_2 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = x^2 + c_1 \ln(x) + c_2 \tag{1}$$

Verification of solutions

$$y = x^2 + c_1 \ln(x) + c_2$$

Verified OK.

7.7.7 Maple step by step solution

Let's solve

$$y''x + y' = 4x$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Make substitution $u = y'$ to reduce order of ODE

$$u'(x)x + u(x) = 4x$$

- Isolate the derivative

$$u'(x) = 4 - \frac{u(x)}{x}$$

- Group terms with $u(x)$ on the lhs of the ODE and the rest on the rhs of the ODE

$$u'(x) + \frac{u(x)}{x} = 4$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) \left(u'(x) + \frac{u(x)}{x} \right) = 4\mu(x)$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x) u(x))$

$$\mu(x) \left(u'(x) + \frac{u(x)}{x} \right) = \mu'(x) u(x) + \mu(x) u'(x)$$

- Isolate $\mu'(x)$

$$\mu'(x) = \frac{\mu(x)}{x}$$

- Solve to find the integrating factor

$$\mu(x) = x$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x) u(x)) \right) dx = \int 4\mu(x) dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x) u(x) = \int 4\mu(x) dx + c_1$$

- Solve for $u(x)$

$$u(x) = \frac{\int 4\mu(x) dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = x$

$$u(x) = \frac{\int 4x dx + c_1}{x}$$

- Evaluate the integrals on the rhs

$$u(x) = \frac{2x^2 + c_1}{x}$$

- Solve 1st ODE for $u(x)$

$$u(x) = \frac{2x^2 + c_1}{x}$$

- Make substitution $u = y'$

$$y' = \frac{2x^2 + c_1}{x}$$

- Integrate both sides to solve for y

$$\int y' dx = \int \frac{2x^2 + c_1}{x} dx + c_2$$

- Compute integrals

$$y = x^2 + c_1 \ln(x) + c_2$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
-> Calling odsolve with the ODE`, diff(_b(_a), _a) = -(_b(_a)-4*_a)/_a, _b(_a)`
  Methods for first order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  trying 1st order linear
  <- 1st order linear successful
<- high order exact linear fully integrable successful`

```

*** Sublev

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(x*diff(y(x),x$2)+diff(y(x),x)=4*x,y(x), singsol=all)
```

$$y(x) = x^2 + c_1 \ln(x) + c_2$$

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 16

```
DSolve[x*y'[x]+y'[x]==4*x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^2 + c_1 \log(x) + c_2$$

7.8 problem 2(a)

7.8.1	Solving as second order integrable as is ode	986
7.8.2	Solving as second order ode missing y ode	990
7.8.3	Solving as type second_order_integrable_as_is (not using ABC version)	993

Internal problem ID [6244]

Internal file name [OUTPUT/5492_Sunday_June_05_2022_03_41_26_PM_64351548/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.9. Reduction of Order. Page 38

Problem number: 2(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_integrable_as_is", "second_order_ode_missing_y"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_y], [_2nd_order, _exact, _nonlinear], [_2nd_order, _reducible, _mu_poly_yn]]
```

$$(x^2 + 2y') y'' + 2xy' = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 0]$$

7.8.1 Solving as second order integrable as is ode

Integrating both sides of the ODE w.r.t x gives

$$\int ((x^2 + 2y') y'' + 2xy') dx = 0$$
$$x^2 y' + y'^2 = c_1$$

Which is now solved for y . Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = -\frac{x^2}{2} + \frac{\sqrt{x^4 + 4c_1}}{2} \quad (1)$$

$$y' = -\frac{x^2}{2} - \frac{\sqrt{x^4 + 4c_1}}{2} \quad (2)$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\begin{aligned} y &= \int -\frac{x^2}{2} + \frac{\sqrt{x^4 + 4c_1}}{2} dx \\ &= \frac{x\sqrt{x^4 + 4c_1}}{6} + \frac{c_1\sqrt{2} \sqrt{4 - \frac{2ix^2}{\sqrt{c_1}}} \sqrt{4 + \frac{2ix^2}{\sqrt{c_1}}} \text{EllipticF}\left(\frac{x\sqrt{2}\sqrt{\frac{i}{\sqrt{c_1}}}}{2}, i\right)}{3\sqrt{\frac{i}{\sqrt{c_1}}} \sqrt{x^4 + 4c_1}} - \frac{x^3}{6} + c_2 \end{aligned}$$

Solving equation (2)

Integrating both sides gives

$$\begin{aligned} y &= \int -\frac{x^2}{2} - \frac{\sqrt{x^4 + 4c_1}}{2} dx \\ &= -\frac{x^3}{6} - \frac{x\sqrt{x^4 + 4c_1}}{6} - \frac{c_1\sqrt{2} \sqrt{4 - \frac{2ix^2}{\sqrt{c_1}}} \sqrt{4 + \frac{2ix^2}{\sqrt{c_1}}} \text{EllipticF}\left(\frac{x\sqrt{2}\sqrt{\frac{i}{\sqrt{c_1}}}}{2}, i\right)}{3\sqrt{\frac{i}{\sqrt{c_1}}} \sqrt{x^4 + 4c_1}} + c_3 \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the First solution

$$y = \frac{x\sqrt{x^4 + 4c_1}}{6} + \frac{c_1\sqrt{2} \sqrt{4 - \frac{2ix^2}{\sqrt{c_1}}} \sqrt{4 + \frac{2ix^2}{\sqrt{c_1}}} \text{EllipticF}\left(\frac{x\sqrt{2}\sqrt{\frac{i}{\sqrt{c_1}}}}{2}, i\right)}{3\sqrt{\frac{i}{\sqrt{c_1}}} \sqrt{x^4 + 4c_1}} - \frac{x^3}{6} + c_2 \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 0$ in the above gives

$$1 = c_2 \quad (1A)$$

Taking derivative of the solution gives

$$y' = \frac{\sqrt{x^4 + 4c_1}}{6} + \frac{x^4}{3\sqrt{x^4 + 4c_1}} - \frac{2i\sqrt{c_1}\sqrt{2}\sqrt{4 + \frac{2ix^2}{\sqrt{c_1}}}\text{EllipticF}\left(\frac{x\sqrt{2}\sqrt{\frac{i}{\sqrt{c_1}}}, i\right)x}{3\sqrt{\frac{i}{\sqrt{c_1}}}\sqrt{4 - \frac{2ix^2}{\sqrt{c_1}}}\sqrt{x^4 + 4c_1}} + \frac{2i\sqrt{c_1}\sqrt{2}\sqrt{4 - \frac{2ix^2}{\sqrt{c_1}}}\text{E}}{3\sqrt{\frac{i}{\sqrt{c_1}}}\sqrt{4 + \frac{2ix^2}{\sqrt{c_1}}}\sqrt{x^4 + 4c_1}}$$

substituting $y' = 0$ and $x = 0$ in the above gives

$$0 = \sqrt{c_1} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 0$$

$$c_2 = 1$$

Substituting these values back in above solution results in

$$y = \lim_{c_1 \rightarrow 0} \left(-\frac{-\sqrt{\frac{i}{\sqrt{c_1}}}x^5 + x^3\sqrt{\frac{i}{\sqrt{c_1}}}\sqrt{x^4 + 4c_1} - 4c_1\sqrt{2}\sqrt{\frac{-ix^2 + 2\sqrt{c_1}}{\sqrt{c_1}}}\sqrt{\frac{ix^2 + 2\sqrt{c_1}}{\sqrt{c_1}}}\text{EllipticF}\left(\frac{x\sqrt{2}\sqrt{\frac{i}{\sqrt{c_1}}}, i\right) - 4c_1}{6\sqrt{\frac{i}{\sqrt{c_1}}}\sqrt{x^4 + 4c_1}} \right)$$

Looking at the Second solution

$$y = -\frac{x^3}{6} - \frac{x\sqrt{x^4 + 4c_1}}{6} - \frac{c_1\sqrt{2}\sqrt{4 - \frac{2ix^2}{\sqrt{c_1}}}\sqrt{4 + \frac{2ix^2}{\sqrt{c_1}}}\text{EllipticF}\left(\frac{x\sqrt{2}\sqrt{\frac{i}{\sqrt{c_1}}}, i\right)}{3\sqrt{\frac{i}{\sqrt{c_1}}}\sqrt{x^4 + 4c_1}} + c_3 \quad (2)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 0$ in the above gives

$$1 = c_3 \quad (1A)$$

Taking derivative of the solution gives

$$y' = -\frac{x^2}{2} - \frac{\sqrt{x^4 + 4c_1}}{6} - \frac{x^4}{3\sqrt{x^4 + 4c_1}} + \frac{2i\sqrt{c_1}\sqrt{2}\sqrt{4 + \frac{2ix^2}{\sqrt{c_1}}}\text{EllipticF}\left(\frac{x\sqrt{2}\sqrt{\frac{i}{\sqrt{c_1}}}, i\right)x}{3\sqrt{\frac{i}{\sqrt{c_1}}}\sqrt{4 - \frac{2ix^2}{\sqrt{c_1}}}\sqrt{x^4 + 4c_1}} - \frac{2i\sqrt{c_1}\sqrt{2}\sqrt{4 - \frac{2ix^2}{\sqrt{c_1}}}\text{E}}{3\sqrt{\frac{i}{\sqrt{c_1}}}\sqrt{4 + \frac{2ix^2}{\sqrt{c_1}}}\sqrt{x^4 + 4c_1}}$$

substituting $y' = 0$ and $x = 0$ in the above gives

$$0 = -\sqrt{c_1} \quad (2A)$$

Equations {1A,2A} are now solved for {c₁, c₃}. Solving for the constants gives

$$\begin{aligned} c_1 &= 0 \\ c_3 &= 1 \end{aligned}$$

Substituting these values back in above solution results in

$$y = \lim_{c_1 \rightarrow 0} \left(- \frac{\sqrt{\frac{i}{\sqrt{c_1}}} x^5 + x^3 \sqrt{\frac{i}{\sqrt{c_1}}} \sqrt{x^4 + 4c_1} + 4c_1 \sqrt{2} \sqrt{\frac{-ix^2 + 2\sqrt{c_1}}{\sqrt{c_1}}} \sqrt{\frac{ix^2 + 2\sqrt{c_1}}{\sqrt{c_1}}} \text{EllipticF} \left(\frac{x\sqrt{2}\sqrt{\frac{i}{\sqrt{c_1}}}}{2}, i \right) + 4\sqrt{\frac{i}{\sqrt{c_1}}} \sqrt{x^4 + 4c_1}}{6\sqrt{\frac{i}{\sqrt{c_1}}} \sqrt{x^4 + 4c_1}} \right)$$

Summary

The solution(s) found are the following

$$y = \left(\lim_{c_1 \rightarrow 0} \left(- \frac{4 \left((1+i) \text{EllipticF} \left(\frac{\left(\frac{1}{2} - \frac{i}{2}\right)x}{\frac{1}{c_1^{\frac{1}{4}}}}, i \right) c_1^{\frac{5}{4}} \sqrt{\frac{ix^2 + 2\sqrt{c_1}}{\sqrt{c_1}}} \sqrt{\frac{-ix^2 - 2\sqrt{c_1}}{\sqrt{c_1}}} + \left(-\frac{x^3}{4} + \frac{3}{2}\right) \sqrt{x^4 + 4c_1} + \frac{x^5}{4} + c_1 x \right)}{\sqrt{x^4 + 4c_1}} \right) \right) \quad (1)$$

$$y = - \frac{6 \left(\lim_{c_1 \rightarrow 0} \frac{4(1+i) \text{EllipticF} \left(\frac{\left(\frac{1}{2} - \frac{i}{2}\right)x}{\frac{1}{c_1^{\frac{1}{4}}}}, i \right) c_1^{\frac{5}{4}} \sqrt{\frac{ix^2 + 2\sqrt{c_1}}{\sqrt{c_1}}} \sqrt{\frac{-ix^2 - 2\sqrt{c_1}}{\sqrt{c_1}}} + 4 \left(\frac{x^3}{4} - \frac{3}{2}\right) \sqrt{x^4 + 4c_1} + x^5 + 4c_1 x}{\sqrt{x^4 + 4c_1}} \right)}{6} \quad (2)$$

Verification of solutions

$$y = - \frac{6 \left(\lim_{c_1 \rightarrow 0} \left(- \frac{4 \left((1+i) \text{EllipticF} \left(\frac{\left(\frac{1}{2} - \frac{i}{2}\right)x}{\frac{1}{c_1^{\frac{1}{4}}}}, i \right) c_1^{\frac{5}{4}} \sqrt{\frac{ix^2 + 2\sqrt{c_1}}{\sqrt{c_1}}} \sqrt{\frac{-ix^2 - 2\sqrt{c_1}}{\sqrt{c_1}}} + \left(-\frac{x^3}{4} + \frac{3}{2}\right) \sqrt{x^4 + 4c_1} + \frac{x^5}{4} + c_1 x \right)}{\sqrt{x^4 + 4c_1}} \right) \right)}{6}$$

Verified OK.

$$y = - \frac{6 \left(\lim_{c_1 \rightarrow 0} \frac{4(1+i) \text{EllipticF} \left(\frac{\left(\frac{1}{2} - \frac{i}{2}\right)x}{\frac{1}{c_1^{\frac{1}{4}}}}, i \right) c_1^{\frac{5}{4}} \sqrt{\frac{ix^2 + 2\sqrt{c_1}}{\sqrt{c_1}}} \sqrt{\frac{-ix^2 - 2\sqrt{c_1}}{\sqrt{c_1}}} + 4 \left(\frac{x^3}{4} - \frac{3}{2}\right) \sqrt{x^4 + 4c_1} + x^5 + 4c_1 x}{\sqrt{x^4 + 4c_1}} \right)}{6}$$

Verified OK.

7.8.2 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$(x^2 + 2p(x)) p'(x) + 2xp(x) = 0$$

Which is now solve for $p(x)$ as first order ode. Writing the ode as

$$p'(x) = -\frac{2xp(x)}{x^2 + 2p(x)} \quad (1)$$

Which becomes

$$(2p) dp = (-x^2) dp + (-2xp) dx \quad (2)$$

But the RHS is complete differential because

$$(-x^2) dp + (-2xp) dx = d(-x^2p)$$

Hence (2) becomes

$$(2p) dp = d(-x^2p)$$

Integrating both sides gives gives these solutions

$$p(x) = -\frac{x^2}{2} + \frac{\sqrt{x^4 + 4c_1}}{2} + c_1$$
$$p(x) = -\frac{x^2}{2} - \frac{\sqrt{x^4 + 4c_1}}{2} + c_1$$

Initial conditions are used to solve for c_1 . Substituting $x = 0$ and $p = 0$ in the above solution gives an equation to solve for the constant of integration.

$$0 = -\sqrt{c_1} + c_1$$

$$c_1 = 0$$

Substituting c_1 found above in the general solution gives

$$p(x) = -\frac{x^2}{2} - \frac{\sqrt{x^4}}{2}$$

Initial conditions are used to solve for c_1 . Substituting $x = 0$ and $p = 0$ in the above solution gives an equation to solve for the constant of integration.

$$0 = \sqrt{c_1} + c_1$$

$$c_1 = 0$$

Substituting c_1 found above in the general solution gives

$$p(x) = -\frac{x^2}{2} + \frac{\sqrt{x^4}}{2}$$

For solution (1) found earlier, since $p = y'$ then the new first order ode to solve is

$$y' = -\frac{x^2}{2} + \frac{\sqrt{x^4}}{2}$$

Integrating both sides gives

$$\begin{aligned} y &= \int -\frac{x^2}{2} + \frac{\sqrt{x^4}}{2} dx \\ &= \frac{\sqrt{x^4}x}{6} - \frac{x^3}{6} + c_2 \end{aligned}$$

Initial conditions are used to solve for c_2 . Substituting $x = 0$ and $y = 1$ in the above solution gives an equation to solve for the constant of integration.

$$1 = c_2$$

$$c_2 = 1$$

Substituting c_2 found above in the general solution gives

$$y = \frac{\sqrt{x^4}x}{6} - \frac{x^3}{6} + 1$$

Since $p = y'$ then the new first order ode to solve is

$$y' = -\frac{x^2}{2} - \frac{\sqrt{x^4}}{2}$$

Integrating both sides gives

$$\begin{aligned}y &= \int -\frac{x^2}{2} - \frac{\sqrt{x^4}}{2} dx \\ &= -\frac{x^3}{6} - \frac{\sqrt{x^4} x}{6} + c_3\end{aligned}$$

Initial conditions are used to solve for c_3 . Substituting $x = 0$ and $y = 1$ in the above solution gives an equation to solve for the constant of integration.

$$1 = c_3$$

$$c_3 = 1$$

Substituting c_3 found above in the general solution gives

$$y = -\frac{x^3}{6} - \frac{\sqrt{x^4} x}{6} + 1$$

Initial conditions are used to solve for the constants of integration.

Summary

The solution(s) found are the following

$$y = \frac{\sqrt{x^4} x}{6} - \frac{x^3}{6} + 1 \tag{1}$$

$$y = -\frac{x^3}{6} - \frac{\sqrt{x^4} x}{6} + 1 \tag{2}$$




Figure 193: Solution plot

Verification of solutions

$$y = \frac{\sqrt{x^4} x}{6} - \frac{x^3}{6} + 1$$

Verified OK.

$$y = -\frac{x^3}{6} - \frac{\sqrt{x^4} x}{6} + 1$$

Verified OK.

7.8.3 Solving as type second_order_integrable_as_is (not using ABC version)

Writing the ode as

$$(x^2 + 2y') y'' + 2xy' = 0$$

Integrating both sides of the ODE w.r.t x gives

$$\int ((x^2 + 2y') y'' + 2xy') dx = 0$$

$$x^2 y' + y'^2 = c_1$$

Which is now solved for y . Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = -\frac{x^2}{2} + \frac{\sqrt{x^4 + 4c_1}}{2} \quad (1)$$

$$y' = -\frac{x^2}{2} - \frac{\sqrt{x^4 + 4c_1}}{2} \quad (2)$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$y = \int -\frac{x^2}{2} + \frac{\sqrt{x^4 + 4c_1}}{2} dx$$

$$= \frac{x\sqrt{x^4 + 4c_1}}{6} + \frac{c_1\sqrt{2} \sqrt{4 - \frac{2ix^2}{\sqrt{c_1}}} \sqrt{4 + \frac{2ix^2}{\sqrt{c_1}}} \text{EllipticF}\left(\frac{x\sqrt{2}\sqrt{\frac{i}{\sqrt{c_1}}}}{2}, i\right)}{3\sqrt{\frac{i}{\sqrt{c_1}}}\sqrt{x^4 + 4c_1}} - \frac{x^3}{6} + c_2$$

Solving equation (2)

Integrating both sides gives

$$y = \int -\frac{x^2}{2} - \frac{\sqrt{x^4 + 4c_1}}{2} dx$$

$$= -\frac{x^3}{6} - \frac{x\sqrt{x^4 + 4c_1}}{6} - \frac{c_1\sqrt{2} \sqrt{4 - \frac{2ix^2}{\sqrt{c_1}}} \sqrt{4 + \frac{2ix^2}{\sqrt{c_1}}} \text{EllipticF}\left(\frac{x\sqrt{2}\sqrt{\frac{i}{\sqrt{c_1}}}}{2}, i\right)}{3\sqrt{\frac{i}{\sqrt{c_1}}} \sqrt{x^4 + 4c_1}} + c_3$$

Initial conditions are used to solve for the constants of integration.

Looking at the First solution

$$y = \frac{x\sqrt{x^4 + 4c_1}}{6} + \frac{c_1\sqrt{2} \sqrt{4 - \frac{2ix^2}{\sqrt{c_1}}} \sqrt{4 + \frac{2ix^2}{\sqrt{c_1}}} \text{EllipticF}\left(\frac{x\sqrt{2}\sqrt{\frac{i}{\sqrt{c_1}}}}{2}, i\right)}{3\sqrt{\frac{i}{\sqrt{c_1}}} \sqrt{x^4 + 4c_1}} - \frac{x^3}{6} + c_2 \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 0$ in the above gives

$$1 = c_2 \quad (1A)$$

Taking derivative of the solution gives

$$y' = \frac{\sqrt{x^4 + 4c_1}}{6} + \frac{x^4}{3\sqrt{x^4 + 4c_1}} - \frac{2i\sqrt{c_1} \sqrt{2} \sqrt{4 + \frac{2ix^2}{\sqrt{c_1}}} \text{EllipticF}\left(\frac{x\sqrt{2}\sqrt{\frac{i}{\sqrt{c_1}}}}{2}, i\right) x}{3\sqrt{\frac{i}{\sqrt{c_1}}} \sqrt{4 - \frac{2ix^2}{\sqrt{c_1}}} \sqrt{x^4 + 4c_1}} + \frac{2i\sqrt{c_1} \sqrt{2} \sqrt{4 - \frac{2ix^2}{\sqrt{c_1}}}}{3\sqrt{\frac{i}{\sqrt{c_1}}} \sqrt{4 + \frac{2ix^2}{\sqrt{c_1}}}}$$

substituting $y' = 0$ and $x = 0$ in the above gives

$$0 = \sqrt{c_1} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 0$$

$$c_2 = 1$$

Substituting these values back in above solution results in

$$y = \lim_{c_1 \rightarrow 0} \left(\frac{-\sqrt{\frac{i}{\sqrt{c_1}}} x^5 + x^3 \sqrt{\frac{i}{\sqrt{c_1}}} \sqrt{x^4 + 4c_1} - 4c_1\sqrt{2} \sqrt{\frac{-ix^2 + 2\sqrt{c_1}}{\sqrt{c_1}}} \sqrt{\frac{ix^2 + 2\sqrt{c_1}}{\sqrt{c_1}}} \text{EllipticF}\left(\frac{x\sqrt{2}\sqrt{\frac{i}{\sqrt{c_1}}}}{2}, i\right) - 4c_3}{6\sqrt{\frac{i}{\sqrt{c_1}}} \sqrt{x^4 + 4c_1}} \right)$$

Looking at the Second solution

$$y = -\frac{x^3}{6} - \frac{x\sqrt{x^4 + 4c_1}}{6} - \frac{c_1\sqrt{2}\sqrt{4 - \frac{2ix^2}{\sqrt{c_1}}}\sqrt{4 + \frac{2ix^2}{\sqrt{c_1}}}\text{EllipticF}\left(\frac{x\sqrt{2}\sqrt{\frac{i}{\sqrt{c_1}}}}{2}, i\right)}{3\sqrt{\frac{i}{\sqrt{c_1}}}\sqrt{x^4 + 4c_1}} + c_3 \quad (2)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 0$ in the above gives

$$1 = c_3 \quad (1A)$$

Taking derivative of the solution gives

$$y' = -\frac{x^2}{2} - \frac{\sqrt{x^4 + 4c_1}}{6} - \frac{x^4}{3\sqrt{x^4 + 4c_1}} + \frac{2i\sqrt{c_1}\sqrt{2}\sqrt{4 + \frac{2ix^2}{\sqrt{c_1}}}\text{EllipticF}\left(\frac{x\sqrt{2}\sqrt{\frac{i}{\sqrt{c_1}}}}{2}, i\right)x}{3\sqrt{\frac{i}{\sqrt{c_1}}}\sqrt{4 - \frac{2ix^2}{\sqrt{c_1}}}\sqrt{x^4 + 4c_1}} - \frac{2i\sqrt{c_1}\sqrt{2}\sqrt{4 - \frac{2ix^2}{\sqrt{c_1}}}\sqrt{x^4 + 4c_1}}{3\sqrt{\frac{i}{\sqrt{c_1}}}\sqrt{x^4 + 4c_1}}$$

substituting $y' = 0$ and $x = 0$ in the above gives

$$0 = -\sqrt{c_1} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_3\}$. Solving for the constants gives

$$c_1 = 0$$

$$c_3 = 1$$

Substituting these values back in above solution results in

$$y = \lim_{c_1 \rightarrow 0} \left(-\frac{\sqrt{\frac{i}{\sqrt{c_1}}}x^5 + x^3\sqrt{\frac{i}{\sqrt{c_1}}}\sqrt{x^4 + 4c_1} + 4c_1\sqrt{2}\sqrt{\frac{-ix^2 + 2\sqrt{c_1}}{\sqrt{c_1}}}\sqrt{\frac{ix^2 + 2\sqrt{c_1}}{\sqrt{c_1}}}\text{EllipticF}\left(\frac{x\sqrt{2}\sqrt{\frac{i}{\sqrt{c_1}}}}{2}, i\right) + 4\sqrt{c_1}}{6\sqrt{\frac{i}{\sqrt{c_1}}}\sqrt{x^4 + 4c_1}} \right)$$

Summary

The solution(s) found are the following

$$y = \frac{\lim_{c_1 \rightarrow 0} \left(\frac{4 \left((1+i) \operatorname{EllipticF} \left(\frac{\left(\frac{1}{2} - \frac{i}{2}\right)x}{c_1^{\frac{1}{4}}}, i \right) c_1^{\frac{5}{4}} \sqrt{\frac{ix^2+2\sqrt{c_1}}{\sqrt{c_1}}} \sqrt{-\frac{ix^2-2\sqrt{c_1}}{\sqrt{c_1}}} + \left(-\frac{x^3}{4} + \frac{3}{2}\right) \sqrt{x^4+4c_1} + \frac{x^5}{4} + c_1x \right)}{\sqrt{x^4+4c_1}} \right)}{6} \quad (1)$$
$$y = - \frac{\lim_{c_1 \rightarrow 0} \left(\frac{4(1+i) \operatorname{EllipticF} \left(\frac{\left(\frac{1}{2} - \frac{i}{2}\right)x}{c_1^{\frac{1}{4}}}, i \right) c_1^{\frac{5}{4}} \sqrt{\frac{ix^2+2\sqrt{c_1}}{\sqrt{c_1}}} \sqrt{-\frac{ix^2-2\sqrt{c_1}}{\sqrt{c_1}}} + 4\left(\frac{x^3}{4} - \frac{3}{2}\right) \sqrt{x^4+4c_1} + x^5 + 4c_1x}{\sqrt{x^4+4c_1}} \right)}{6} \quad (2)$$

Verification of solutions

$$y = - \frac{\lim_{c_1 \rightarrow 0} \left(\frac{4 \left((1+i) \operatorname{EllipticF} \left(\frac{\left(\frac{1}{2} - \frac{i}{2}\right)x}{c_1^{\frac{1}{4}}}, i \right) c_1^{\frac{5}{4}} \sqrt{\frac{ix^2+2\sqrt{c_1}}{\sqrt{c_1}}} \sqrt{-\frac{ix^2-2\sqrt{c_1}}{\sqrt{c_1}}} + \left(-\frac{x^3}{4} + \frac{3}{2}\right) \sqrt{x^4+4c_1} + \frac{x^5}{4} + c_1x \right)}{\sqrt{x^4+4c_1}} \right)}{6}$$

Verified OK.

$$y = - \frac{\lim_{c_1 \rightarrow 0} \left(\frac{4(1+i) \operatorname{EllipticF} \left(\frac{\left(\frac{1}{2} - \frac{i}{2}\right)x}{c_1^{\frac{1}{4}}}, i \right) c_1^{\frac{5}{4}} \sqrt{\frac{ix^2+2\sqrt{c_1}}{\sqrt{c_1}}} \sqrt{-\frac{ix^2-2\sqrt{c_1}}{\sqrt{c_1}}} + 4\left(\frac{x^3}{4} - \frac{3}{2}\right) \sqrt{x^4+4c_1} + x^5 + 4c_1x}{\sqrt{x^4+4c_1}} \right)}{6}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
-> Calling odsolve with the ODE`, diff(_b(_a), _a) = -2*_b(_a)*_a/(_a^2+2*_b(_a)), _b(_a), H
symmetry methods on request
`, `1st order, trying reduction of order with given symmetries: `[a, 2*_b]
```

✓ Solution by Maple

Time used: 0.079 (sec). Leaf size: 15

```
dsolve([(x^2+2*diff(y(x),x))*diff(y(x),x$2)+2*x*diff(y(x),x)=0,y(0) = 1, D(y)(0) = 0],y(x),
```

$$y(x) = 1$$
$$y(x) = -\frac{x^3}{3} + 1$$

✓ Solution by Mathematica

Time used: 0.252 (sec). Leaf size: 32

```
DSolve[{(x^2+2*y'[x])*y''[x]+2*x*y'[x]==0,{y[0]==1,y'[0]==0}},y[x],x,IncludeSingularSolution
```

$y(x) \rightarrow$ Indeterminate

$$y(x) \rightarrow 0 \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, \frac{1}{4}, \frac{5}{4}, \operatorname{ComplexInfinity}\right) - \frac{x^3}{6} + 1$$

7.9 problem 2(b)

7.9.1 Solving as second order ode missing x ode	998
7.9.2 Maple step by step solution	1002

Internal problem ID [6245]

Internal file name [OUTPUT/5493_Sunday_June_05_2022_03_41_28_PM_68194147/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.9. Reduction of Order. Page 38

Problem number: 2(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_ode_missing_x"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x], [_2nd_order,
    _with_potential_symmetries], [_2nd_order, _reducible, _mu_xy
]]
```

$$yy'' - y'y^2 - y'^2 = 0$$

With initial conditions

$$\left[y(0) = -\frac{1}{2}, y'(0) = 1 \right]$$

7.9.1 Solving as second order ode missing x ode

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable y an independent variable. Using

$$y' = p(y)$$

Then

$$\begin{aligned}y'' &= \frac{dp}{dx} \\ &= \frac{dy}{dx} \frac{dp}{dy} \\ &= p \frac{dp}{dy}\end{aligned}$$

Hence the ode becomes

$$yp(y) \left(\frac{d}{dy} p(y) \right) + (-p(y) - y^2) p(y) = 0$$

Which is now solved as first order ode for $p(y)$.

Entering Linear first order ODE solver. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int -\frac{1}{y} dy} \\ &= \frac{1}{y}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dy}(\mu p) &= (\mu)(y) \\ \frac{d}{dy} \left(\frac{p}{y} \right) &= \left(\frac{1}{y} \right) (y) \\ d \left(\frac{p}{y} \right) &= dy\end{aligned}$$

Integrating gives

$$\begin{aligned}\frac{p}{y} &= \int dy \\ \frac{p}{y} &= y + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = \frac{1}{y}$ results in

$$p(y) = c_1 y + y^2$$

which simplifies to

$$p(y) = y(y + c_1)$$

Initial conditions are used to solve for c_1 . Substituting $y = -\frac{1}{2}$ and $p = 1$ in the above solution gives an equation to solve for the constant of integration.

$$1 = \frac{1}{4} - \frac{c_1}{2}$$

$$c_1 = -\frac{3}{2}$$

Substituting c_1 found above in the general solution gives

$$p(y) = \frac{y(2y - 3)}{2}$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = \frac{y(2y - 3)}{2}$$

Integrating both sides gives

$$\int \frac{2}{y(2y - 3)} dy = \int dx$$

$$-\frac{2 \ln(y)}{3} + \frac{2 \ln(2y - 3)}{3} = c_2 + x$$

The above can be written as

$$\left(-\frac{2}{3}\right) (\ln(y) - \ln(2y - 3)) = c_2 + x$$

$$\ln(y) - \ln(2y - 3) = \left(-\frac{3}{2}\right) (c_2 + x)$$

$$= -\frac{3c_2}{2} - \frac{3x}{2}$$

Raising both side to exponential gives

$$e^{\ln(y) - \ln(2y - 3)} = -\frac{3c_2 e^{-\frac{3x}{2}}}{2}$$

Which simplifies to

$$\frac{y}{2y - 3} = c_3 e^{-\frac{3x}{2}}$$

Initial conditions are used to solve for c_3 . Substituting $x = 0$ and $y = -\frac{1}{2}$ in the above solution gives an equation to solve for the constant of integration.

$$-\frac{1}{2} = \frac{3c_3}{-1 + 2c_3}$$

$$c_3 = \frac{1}{8}$$

Substituting c_3 found above in the general solution gives

$$y = \frac{3e^{-\frac{3x}{2}}}{2e^{-\frac{3x}{2}} - 8}$$

Initial conditions are used to solve for the constants of integration.

Summary

The solution(s) found are the following

$$y = \frac{3e^{-\frac{3x}{2}}}{2e^{-\frac{3x}{2}} - 8} \quad (1)$$




Figure 194: Solution plot

Verification of solutions

$$y = \frac{3e^{-\frac{3x}{2}}}{2e^{-\frac{3x}{2}} - 8}$$

Verified OK.

7.9.2 Maple step by step solution

Let's solve

$$\left[yy'' + (-y' - y^2) y' = 0, y(0) = -\frac{1}{2}, y' \Big|_{\{x=0\}} = 1 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Define new dependent variable u

$$u(x) = y'$$

- Compute y''

$$u'(x) = y''$$

- Use chain rule on the lhs

$$y' \left(\frac{d}{dy} u(y) \right) = y''$$

- Substitute in the definition of u

$$u(y) \left(\frac{d}{dy} u(y) \right) = y''$$

- Make substitutions $y' = u(y)$, $y'' = u(y) \left(\frac{d}{dy} u(y) \right)$ to reduce order of ODE

$$yu(y) \left(\frac{d}{dy} u(y) \right) + (-u(y) - y^2) u(y) = 0$$

- Isolate the derivative

$$\frac{d}{dy} u(y) = \frac{u(y)}{y} + y$$

- Group terms with $u(y)$ on the lhs of the ODE and the rest on the rhs of the ODE

$$\frac{d}{dy} u(y) - \frac{u(y)}{y} = y$$

- The ODE is linear; multiply by an integrating factor $\mu(y)$

$$\mu(y) \left(\frac{d}{dy} u(y) - \frac{u(y)}{y} \right) = \mu(y) y$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dy} (\mu(y) u(y))$

$$\mu(y) \left(\frac{d}{dy} u(y) - \frac{u(y)}{y} \right) = \left(\frac{d}{dy} \mu(y) \right) u(y) + \mu(y) \left(\frac{d}{dy} u(y) \right)$$

- Isolate $\frac{d}{dy} \mu(y)$

$$\frac{d}{dy} \mu(y) = -\frac{\mu(y)}{y}$$

- Solve to find the integrating factor

$$\mu(y) = \frac{1}{y}$$

- Integrate both sides with respect to y

$$\int \left(\frac{d}{dy}(\mu(y) u(y)) \right) dy = \int \mu(y) y dy + c_1$$

- Evaluate the integral on the lhs

$$\mu(y) u(y) = \int \mu(y) y dy + c_1$$

- Solve for $u(y)$

$$u(y) = \frac{\int \mu(y) y dy + c_1}{\mu(y)}$$

- Substitute $\mu(y) = \frac{1}{y}$

$$u(y) = y \left(\int 1 dy + c_1 \right)$$

- Evaluate the integrals on the rhs

$$u(y) = y(y + c_1)$$

- Solve 1st ODE for $u(y)$

$$u(y) = y(y + c_1)$$

- Revert to original variables with substitution $u(y) = y', y = y$

$$y' = y(y + c_1)$$

- Separate variables

$$\frac{y'}{y(y+c_1)} = 1$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y(y+c_1)} dx = \int 1 dx + c_2$$

- Evaluate integral

$$\frac{\ln(y)}{c_1} - \frac{\ln(y+c_1)}{c_1} = c_2 + x$$

- Solve for y

$$y = -\frac{c_1 e^{c_2 c_1 + c_1 x}}{-1 + e^{c_2 c_1 + c_1 x}}$$

- Check validity of solution $y = -\frac{c_1 e^{c_2 c_1 + c_1 x}}{-1 + e^{c_2 c_1 + c_1 x}}$

- Use initial condition $y(0) = -\frac{1}{2}$

$$-\frac{1}{2} = -\frac{c_1 e^{c_2 c_1}}{-1 + e^{c_2 c_1}}$$

- Compute derivative of the solution

$$y' = \frac{c_1^2 (e^{c_2 c_1 + c_1 x})^2}{(-1 + e^{c_2 c_1 + c_1 x})^2} - \frac{c_1^2 e^{c_2 c_1 + c_1 x}}{-1 + e^{c_2 c_1 + c_1 x}}$$

- Use the initial condition $y'|_{\{x=0\}} = 1$

$$1 = \frac{c_1^2(e^{c_2 c_1})^2}{(-1+e^{c_2 c_1})^2} - \frac{c_1^2 e^{c_2 c_1}}{-1+e^{c_2 c_1}}$$

- Solve for c_1 and c_2

$$\left\{ c_1 = -\frac{3}{2}, c_2 = \frac{4 \ln(2)}{3} \right\}$$

- Substitute constant values into general solution and simplify

$$y = \frac{3e^{-\frac{3x}{2}}}{2e^{-\frac{3x}{2}} - 8}$$

- Solution to the IVP

$$y = \frac{3e^{-\frac{3x}{2}}}{2e^{-\frac{3x}{2}} - 8}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
<- differential order: 2; canonical coordinates successful
<- differential order 2; missing variables successful`

```

✓ Solution by Maple

Time used: 0.203 (sec). Leaf size: 16

```
dsolve([y(x)*diff(y(x),x$2)=y(x)^2*diff(y(x),x)+(diff(y(x),x))^2,y(0) = -1/2, D(y)(0) = 1],y
```

$$y(x) = -\frac{3}{8e^{\frac{3x}{2}} - 2}$$

✓ Solution by Mathematica

Time used: 1.982 (sec). Leaf size: 20

```
DSolve[{y[x]*y'[x]==y[x]^2*y'[x]+(y'[x])^2,{y[0]==-1/2,y'[0]==1}},y[x],x,IncludeSingularSol
```

$$y(x) \rightarrow \frac{3}{2 - 8e^{3x/2}}$$

7.10 problem 2(c)

7.10.1 Solving as second order integrable as is ode	1007
7.10.2 Solving as second order ode missing x ode	1009
7.10.3 Solving as type second_order_integrable_as_is (not using ABC version)	1011
7.10.4 Solving as exact nonlinear second order ode ode	1013
7.10.5 Maple step by step solution	1016

Internal problem ID [6246]

Internal file name [OUTPUT/5494_Sunday_June_05_2022_03_41_29_PM_83387578/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.9. Reduction of Order. Page 38

Problem number: 2(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_integrable_as_is", "second_order_ode_missing_x", "exact nonlinear second order ode"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x], [_2nd_order, _exact, _nonlinear], [  
_2nd_order, _reducible, _mu_xy]]
```

$$y'' - y'e^y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 2]$$

7.10.1 Solving as second order integrable as is ode

Integrating both sides of the ODE w.r.t x gives

$$\int (y'' - y'e^y) dx = 0$$
$$-e^y + y' = c_1$$

Which is now solved for y . Integrating both sides gives

$$\int \frac{1}{e^y + c_1} dy = \int dx$$
$$\frac{\ln(e^y)}{c_1} - \frac{\ln(e^y + c_1)}{c_1} = c_2 + x$$

The above can be written as

$$\left(\frac{1}{c_1}\right) (\ln(e^y) - \ln(e^y + c_1)) = c_2 + x$$
$$\ln(e^y) - \ln(e^y + c_1) = (c_1)(c_2 + x)$$
$$= c_1(c_2 + x)$$

Raising both side to exponential gives

$$e^{\ln(e^y) - \ln(e^y + c_1)} = c_1 c_2 e^{c_1 x}$$

Which simplifies to

$$\frac{e^y}{e^y + c_1} = c_3 e^{c_1 x}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = \ln\left(-\frac{c_3 c_1}{-1 + c_3 e^{c_1 x}}\right) + c_1 x \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = \ln\left(-\frac{c_3 c_1}{-1 + c_3}\right) \quad (1A)$$

Taking derivative of the solution gives

$$y' = -\frac{c_3 e^{c_1 x} c_1}{-1 + c_3 e^{c_1 x}} + c_1$$

substituting $y' = 2$ and $x = 0$ in the above gives

$$2 = -\frac{c_1}{-1 + c_3} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_3\}$. Solving for the constants gives

$$c_1 = 1$$
$$c_3 = \frac{1}{2}$$

Substituting these values back in above solution results in

$$y = \ln\left(-\frac{1}{e^x - 2}\right) + x$$

Summary

The solution(s) found are the following

$$y = \ln\left(-\frac{1}{e^x - 2}\right) + x \quad (1)$$




Figure 195: Solution plot

Verification of solutions

$$y = \ln\left(-\frac{1}{e^x - 2}\right) + x$$

Verified OK.

7.10.2 Solving as second order ode missing x ode

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable y an independent variable. Using

$$y' = p(y)$$

Then

$$\begin{aligned}y'' &= \frac{dp}{dx} \\ &= \frac{dy}{dx} \frac{dp}{dy} \\ &= p \frac{dp}{dy}\end{aligned}$$

Hence the ode becomes

$$p(y) \left(\frac{d}{dy} p(y) \right) - p(y) e^y = 0$$

Which is now solved as first order ode for $p(y)$. Integrating both sides gives

$$\begin{aligned}p(y) &= \int e^y dy \\ &= e^y + c_1\end{aligned}$$

Initial conditions are used to solve for c_1 . Substituting $y = 0$ and $p = 2$ in the above solution gives an equation to solve for the constant of integration.

$$2 = c_1 + 1$$

$$c_1 = 1$$

Substituting c_1 found above in the general solution gives

$$p(y) = 1 + e^y$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = 1 + e^y$$

Integrating both sides gives

$$\int \frac{1}{1+e^y} dy = \int dx$$
$$-\ln(1+e^y) + \ln(e^y) = c_2 + x$$

Raising both side to exponential gives

$$e^{-\ln(1+e^y)+\ln(e^y)} = e^{c_2+x}$$

Which simplifies to

$$\frac{e^y}{1+e^y} = c_3 e^x$$

Initial conditions are used to solve for c_3 . Substituting $x = 0$ and $y = 0$ in the above solution gives an equation to solve for the constant of integration.

$$0 = \ln\left(-\frac{c_3}{-1+c_3}\right)$$

$$c_3 = \frac{1}{2}$$

Substituting c_3 found above in the general solution gives

$$y = \ln\left(-\frac{1}{e^x-2}\right) + x$$

Initial conditions are used to solve for the constants of integration.

Summary

The solution(s) found are the following

$$y = \ln\left(-\frac{1}{e^x-2}\right) + x \tag{1}$$




Figure 196: Solution plot

Verification of solutions

$$y = \ln\left(-\frac{1}{e^x - 2}\right) + x$$

Verified OK.

7.10.3 Solving as type second_order_integrable_as_is (not using ABC version)

Writing the ode as

$$y'' - y'e^y = 0$$

Integrating both sides of the ODE w.r.t x gives

$$\int (y'' - y'e^y) dx = 0$$

$$-e^y + y' = c_1$$

Which is now solved for y . Integrating both sides gives

$$\int \frac{1}{e^y + c_1} dy = \int dx$$

$$\frac{\ln(e^y)}{c_1} - \frac{\ln(e^y + c_1)}{c_1} = c_2 + x$$

The above can be written as

$$\begin{aligned}\left(\frac{1}{c_1}\right)(\ln(e^y) - \ln(e^y + c_1)) &= c_2 + x \\ \ln(e^y) - \ln(e^y + c_1) &= (c_1)(c_2 + x) \\ &= c_1(c_2 + x)\end{aligned}$$

Raising both side to exponential gives

$$e^{\ln(e^y) - \ln(e^y + c_1)} = c_1 c_2 e^{c_1 x}$$

Which simplifies to

$$\frac{e^y}{e^y + c_1} = c_3 e^{c_1 x}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = \ln\left(-\frac{c_3 c_1}{-1 + c_3 e^{c_1 x}}\right) + c_1 x \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = \ln\left(-\frac{c_3 c_1}{-1 + c_3}\right) \quad (1A)$$

Taking derivative of the solution gives

$$y' = -\frac{c_3 e^{c_1 x} c_1}{-1 + c_3 e^{c_1 x}} + c_1$$

substituting $y' = 2$ and $x = 0$ in the above gives

$$2 = -\frac{c_1}{-1 + c_3} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_3\}$. Solving for the constants gives

$$\begin{aligned}c_1 &= 1 \\ c_3 &= \frac{1}{2}\end{aligned}$$

Substituting these values back in above solution results in

$$y = \ln \left(-\frac{1}{e^x - 2} \right) + x$$

Summary

The solution(s) found are the following

$$y = \ln \left(-\frac{1}{e^x - 2} \right) + x \tag{1}$$




Figure 197: Solution plot

Verification of solutions

$$y = \ln \left(-\frac{1}{e^x - 2} \right) + x$$

Verified OK.

7.10.4 Solving as exact nonlinear second order ode

An exact non-linear second order ode has the form

$$a_2(x, y, y') y'' + a_1(x, y, y') y' + a_0(x, y, y') = 0$$

Where the following conditions are satisfied

$$\begin{aligned}\frac{\partial a_2}{\partial y} &= \frac{\partial a_1}{\partial y'} \\ \frac{\partial a_2}{\partial x} &= \frac{\partial a_0}{\partial y'} \\ \frac{\partial a_1}{\partial x} &= \frac{\partial a_0}{\partial y}\end{aligned}$$

Looking at the the ode given we see that

$$\begin{aligned}a_2 &= 1 \\ a_1 &= -e^y \\ a_0 &= 0\end{aligned}$$

Applying the conditions to the above shows this is a nonlinear exact second order ode. Therefore it can be reduced to first order ode given by

$$\begin{aligned}\int a_2 dy' + \int a_1 dy + \int a_0 dx &= c_1 \\ \int 1 dy' + \int -e^y dy + \int 0 dx &= c_1\end{aligned}$$

Which results in

$$-e^y + y' = c_1$$

Which is now solved Integrating both sides gives

$$\begin{aligned}\int \frac{1}{e^y + c_1} dy &= \int dx \\ \frac{\ln(e^y)}{c_1} - \frac{\ln(e^y + c_1)}{c_1} &= c_2 + x\end{aligned}$$

The above can be written as

$$\begin{aligned}\left(\frac{1}{c_1}\right) (\ln(e^y) - \ln(e^y + c_1)) &= c_2 + x \\ \ln(e^y) - \ln(e^y + c_1) &= (c_1)(c_2 + x) \\ &= c_1(c_2 + x)\end{aligned}$$

Raising both side to exponential gives

$$e^{\ln(e^y) - \ln(e^y + c_1)} = c_1 c_2 e^{c_1 x}$$

Which simplifies to

$$\frac{e^y}{e^y + c_1} = c_3 e^{c_1 x}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = \ln \left(-\frac{c_3 c_1}{-1 + c_3 e^{c_1 x}} \right) + c_1 x \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = \ln \left(-\frac{c_3 c_1}{-1 + c_3} \right) \quad (1A)$$

Taking derivative of the solution gives

$$y' = -\frac{c_3 e^{c_1 x} c_1}{-1 + c_3 e^{c_1 x}} + c_1$$

substituting $y' = 2$ and $x = 0$ in the above gives

$$2 = -\frac{c_1}{-1 + c_3} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_3\}$. Solving for the constants gives

$$\begin{aligned} c_1 &= 1 \\ c_3 &= \frac{1}{2} \end{aligned}$$

Substituting these values back in above solution results in

$$y = \ln \left(-\frac{1}{e^x - 2} \right) + x$$

Summary

The solution(s) found are the following

$$y = \ln \left(-\frac{1}{e^x - 2} \right) + x \quad (1)$$




Figure 198: Solution plot

Verification of solutions

$$y = \ln \left(-\frac{1}{e^x - 2} \right) + x$$

Verified OK.

7.10.5 Maple step by step solution

Let's solve

$$\left[y'' - y'e^y = 0, y(0) = 0, y'|_{\{x=0\}} = 2 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Define new dependent variable u

$$u(x) = y'$$

- Compute y''

$$u'(x) = y''$$

- Use chain rule on the lhs

$$y' \left(\frac{d}{dy} u(y) \right) = y''$$

- Substitute in the definition of u

$$u(y) \left(\frac{d}{dy} u(y) \right) = y''$$
- Make substitutions $y' = u(y)$, $y'' = u(y) \left(\frac{d}{dy} u(y) \right)$ to reduce order of ODE

$$u(y) \left(\frac{d}{dy} u(y) \right) - u(y) e^y = 0$$
- Separate variables

$$\frac{d}{dy} u(y) = e^y$$
- Integrate both sides with respect to y

$$\int \left(\frac{d}{dy} u(y) \right) dy = \int e^y dy + c_1$$
- Evaluate integral

$$u(y) = e^y + c_1$$
- Solve for $u(y)$

$$u(y) = e^y + c_1$$
- Solve 1st ODE for $u(y)$

$$u(y) = e^y + c_1$$
- Revert to original variables with substitution $u(y) = y'$, $y = y$

$$y' = e^y + c_1$$
- Separate variables

$$\frac{y'}{e^y + c_1} = 1$$
- Integrate both sides with respect to x

$$\int \frac{y'}{e^y + c_1} dx = \int 1 dx + c_2$$
- Evaluate integral

$$\frac{\ln(e^y)}{c_1} - \frac{\ln(e^y + c_1)}{c_1} = c_2 + x$$
- Solve for y

$$y = c_2 c_1 + c_1 x + \ln \left(-\frac{c_1}{-1 + e^{c_2 c_1 + c_1 x}} \right)$$
- Check validity of solution $y = c_2 c_1 + c_1 x + \ln \left(-\frac{c_1}{-1 + e^{c_2 c_1 + c_1 x}} \right)$
 - Use initial condition $y(0) = 0$

$$0 = c_2 c_1 + \ln \left(-\frac{c_1}{-1 + e^{c_2 c_1}} \right)$$

- Compute derivative of the solution

$$y' = -\frac{c_1 e^{c_2 c_1 + c_1 x}}{-1 + e^{c_2 c_1 + c_1 x}} + c_1$$
- Use the initial condition $y'|_{\{x=0\}} = 2$

$$2 = -\frac{c_1 e^{c_2 c_1}}{-1 + e^{c_2 c_1}} + c_1$$
- Solve for c_1 and c_2

$$\{c_1 = 1, c_2 = -\ln(2)\}$$
- Substitute constant values into general solution and simplify

$$y = \ln\left(-\frac{1}{e^x - 2}\right) + x$$
- Solution to the IVP

$$y = \ln\left(-\frac{1}{e^x - 2}\right) + x$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
-> Calling odsolve with the ODE`, (diff(_b(_a), _a))*_b(_a)-exp(_a)*_b(_a) = 0, _b(_a), HINT
symmetry methods on request
`, `1st order, trying reduction of order with given symmetries:`[1, _b]

```

✓ Solution by Maple

Time used: 0.735 (sec). Leaf size: 21

```
dsolve([diff(y(x),x$2)=diff(y(x),x)*exp(y(x)),y(0) = 0, D(y)(0) = 2],y(x), singsol=all)
```

$$y(x) = x + \ln\left(-\frac{1}{e^x - 2}\right)$$

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[{y'[x]==y[x]*Exp[y[x]},{y[0]==0,y'[0]==2}},y[x],x,IncludeSingularSolutions -> True]
```

```
{}
```


7.11 problem 3(a)

- 7.11.1 Solving as second order ode missing y ode 1020
- 7.11.2 Solving as second order ode missing x ode 1021
- 7.11.3 Maple step by step solution 1023

Internal problem ID [6247]

Internal file name [OUTPUT/5495_Sunday_June_05_2022_03_41_33_PM_45489118/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.9. Reduction of Order. Page 38

Problem number: 3(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_ode_missing_x", "second_order_ode_missing_y"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_xy]]
```

$$y'' - y'^2 = 1$$

7.11.1 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x) - 1 - p(x)^2 = 0$$

Which is now solve for $p(x)$ as first order ode. Integrating both sides gives

$$\int \frac{1}{p^2 + 1} dp = x + c_1$$
$$\arctan(p) = x + c_1$$

Solving for p gives these solutions

$$p_1 = \tan(x + c_1)$$

Since $p = y'$ then the new first order ode to solve is

$$y' = \tan(x + c_1)$$

Integrating both sides gives

$$\begin{aligned} y &= \int \tan(x + c_1) \, dx \\ &= \frac{\ln(1 + \tan(x + c_1)^2)}{2} + c_2 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{\ln(1 + \tan(x + c_1)^2)}{2} + c_2 \quad (1)$$

Verification of solutions

$$y = \frac{\ln(1 + \tan(x + c_1)^2)}{2} + c_2$$

Verified OK.

7.11.2 Solving as second order ode missing x ode

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable y an independent variable. Using

$$y' = p(y)$$

Then

$$\begin{aligned} y'' &= \frac{dp}{dx} \\ &= \frac{dy}{dx} \frac{dp}{dy} \\ &= p \frac{dp}{dy} \end{aligned}$$

Hence the ode becomes

$$p(y) \left(\frac{d}{dy} p(y) \right) - p(y)^2 = 1$$

Which is now solved as first order ode for $p(y)$. Integrating both sides gives

$$\int \frac{p}{p^2 + 1} dp = \int dy$$

$$\frac{\ln(p^2 + 1)}{2} = y + c_1$$

Raising both side to exponential gives

$$\sqrt{p^2 + 1} = e^{y+c_1}$$

Which simplifies to

$$\sqrt{p^2 + 1} = c_2 e^y$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = \text{RootOf}(_Z^2 - c_2^2 e^{2y} + 1)$$

Integrating both sides gives

$$\int \frac{1}{\text{RootOf}(_Z^2 - c_2^2 e^{2y} + 1)} dy = \int dx$$

$$\int^y \frac{1}{\text{RootOf}(_Z^2 - c_2^2 e^{2-a} + 1)} d_a = x + c_3$$

Summary

The solution(s) found are the following

$$\int^y \frac{1}{\text{RootOf}(_Z^2 - c_2^2 e^{2-a} + 1)} d_a = x + c_3 \quad (1)$$

Verification of solutions

$$\int^y \frac{1}{\text{RootOf}(_Z^2 - c_2^2 e^{2-a} + 1)} d_a = x + c_3$$

Verified OK.

7.11.3 Maple step by step solution

Let's solve

$$y'' - y'^2 = 1$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Make substitution $u = y'$ to reduce order of ODE

$$u'(x) - u(x)^2 = 1$$

- Separate variables

$$\frac{u'(x)}{u(x)^2+1} = 1$$

- Integrate both sides with respect to x

$$\int \frac{u'(x)}{u(x)^2+1} dx = \int 1 dx + c_1$$

- Evaluate integral

$$\arctan(u(x)) = x + c_1$$

- Solve for $u(x)$

$$u(x) = \tan(x + c_1)$$

- Solve 1st ODE for $u(x)$

$$u(x) = \tan(x + c_1)$$

- Make substitution $u = y'$

$$y' = \tan(x + c_1)$$

- Integrate both sides to solve for y

$$\int y' dx = \int \tan(x + c_1) dx + c_2$$

- Compute integrals

$$y = \frac{\ln(1+\tan(x+c_1)^2)}{2} + c_2$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying 2nd order Liouville  
trying 2nd order WeierstrassP  
trying 2nd order JacobiSN  
differential order: 2; trying a linearization to 3rd order  
trying 2nd order ODE linearizable_by_differentiation  
trying 2nd order, 2 integrating factors of the form mu(x,y)  
<- 2nd order, 2 integrating factors of the form mu(x,y) successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)=1+(diff(y(x),x))^2,y(x), singsol=all)
```

$$y(x) = -\ln(-\cos(x)c_2 + c_1 \sin(x))$$

✓ Solution by Mathematica

Time used: 1.938 (sec). Leaf size: 16

```
DSolve[y''[x]==1+(y'[x])^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 - \log(\cos(x + c_1))$$

7.12 problem 3(b)

- 7.12.1 Solving as second order ode missing y ode 1025
- 7.12.2 Solving as second order ode missing x ode 1026
- 7.12.3 Maple step by step solution 1028

Internal problem ID [6248]

Internal file name [OUTPUT/5496_Sunday_June_05_2022_03_41_34_PM_27172925/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Section 1.9. Reduction of Order. Page 38

Problem number: 3(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_ode_missing_x", "second_order_ode_missing_y"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_xy]]
```

$$y'' + y'^2 = 1$$

7.12.1 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x) + p(x)^2 - 1 = 0$$

Which is now solve for $p(x)$ as first order ode. Integrating both sides gives

$$\int \frac{1}{-p^2 + 1} dp = x + c_1$$
$$\operatorname{arctanh}(p) = x + c_1$$

Solving for p gives these solutions

$$p_1 = \tanh(x + c_1)$$

Since $p = y'$ then the new first order ode to solve is

$$y' = \tanh(x + c_1)$$

Integrating both sides gives

$$\begin{aligned} y &= \int \tanh(x + c_1) \, dx \\ &= \ln(\cosh(x + c_1)) + c_2 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \ln(\cosh(x + c_1)) + c_2 \tag{1}$$

Verification of solutions

$$y = \ln(\cosh(x + c_1)) + c_2$$

Verified OK.

7.12.2 Solving as second order ode missing x ode

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable y an independent variable. Using

$$y' = p(y)$$

Then

$$\begin{aligned} y'' &= \frac{dp}{dx} \\ &= \frac{dy}{dx} \frac{dp}{dy} \\ &= p \frac{dp}{dy} \end{aligned}$$

Hence the ode becomes

$$p(y) \left(\frac{d}{dy} p(y) \right) + p(y)^2 = 1$$

Which is now solved as first order ode for $p(y)$. Integrating both sides gives

$$\int -\frac{p}{p^2 - 1} dp = \int dy$$

$$-\frac{\ln(p - 1)}{2} - \frac{\ln(p + 1)}{2} = y + c_1$$

The above can be written as

$$\left(-\frac{1}{2}\right) (\ln(p - 1) + \ln(p + 1)) = y + c_1$$

$$\ln(p - 1) + \ln(p + 1) = (-2)(y + c_1)$$

$$= -2c_1 - 2y$$

Raising both side to exponential gives

$$e^{\ln(p-1)+\ln(p+1)} = -2c_1 e^{-2y}$$

Which simplifies to

$$p^2 - 1 = c_2 e^{-2y}$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = \text{RootOf}(_Z^2 - c_2 e^{-2y} - 1)$$

Integrating both sides gives

$$\int \frac{1}{\text{RootOf}(_Z^2 - c_2 e^{-2y} - 1)} dy = \int dx$$

$$\int^y \frac{1}{\text{RootOf}(_Z^2 - c_2 e^{-2_a} - 1)} d_a = x + c_3$$

Summary

The solution(s) found are the following

$$\int^y \frac{1}{\text{RootOf}(_Z^2 - c_2 e^{-2_a} - 1)} d_a = x + c_3 \quad (1)$$

Verification of solutions

$$\int^y \frac{1}{\text{RootOf}(_Z^2 - c_2 e^{-2_a} - 1)} d_a = x + c_3$$

Verified OK.

7.12.3 Maple step by step solution

Let's solve

$$y'' + y'^2 = 1$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Make substitution $u = y'$ to reduce order of ODE

$$u'(x) + u(x)^2 = 1$$

- Separate variables

$$\frac{u'(x)}{-u(x)^2+1} = 1$$

- Integrate both sides with respect to x

$$\int \frac{u'(x)}{-u(x)^2+1} dx = \int 1 dx + c_1$$

- Evaluate integral

$$\operatorname{arctanh}(u(x)) = x + c_1$$

- Solve for $u(x)$

$$u(x) = \tanh(x + c_1)$$

- Solve 1st ODE for $u(x)$

$$u(x) = \tanh(x + c_1)$$

- Make substitution $u = y'$

$$y' = \tanh(x + c_1)$$

- Integrate both sides to solve for y

$$\int y' dx = \int \tanh(x + c_1) dx + c_2$$

- Compute integrals

$$y = \ln(\cosh(x + c_1)) + c_2$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying 2nd order Liouville  
trying 2nd order WeierstrassP  
trying 2nd order JacobiSN  
differential order: 2; trying a linearization to 3rd order  
trying 2nd order ODE linearizable_by_differentiation  
trying 2nd order, 2 integrating factors of the form mu(x,y)  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful  
<- 2nd order, 2 integrating factors of the form mu(x,y) successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 21

```
dsolve(diff(y(x),x$2)+(diff(y(x),x))^2=1,y(x), singsol=all)
```

$$y(x) = x - \ln(2) + \ln(e^{-2x}c_1 - c_2)$$

✓ Solution by Mathematica

Time used: 0.388 (sec). Leaf size: 46

```
DSolve[y''[x]+(y'[x])^2==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\log(e^x) + \log(e^{2x} + e^{2c_1}) + c_2$$

$$y(x) \rightarrow -\log(e^x) + \log(e^{2x}) + c_2$$

8 Chapter 1. What is a differential equation.

Problems for Review and Discovery. Page 53

8.1	problem 1(a)	1031
8.2	problem 1(b)	1046
8.3	problem 1(c)	1060
8.4	problem 1(d)	1075
8.5	problem 1(e)	1087
8.6	problem 1(f)	1097
8.7	problem 1(g)	1111
8.8	problem 1(h)	1126
8.9	problem 2(a)	1138
8.10	problem 2(b)	1153
8.11	problem 2(c)	1167
8.12	problem 2(d)	1183
8.13	problem 2(e)	1195
8.14	problem 2(f)	1208
8.15	problem 2(g)	1218
8.16	problem 2(h)	1231
8.17	problem 4(a)	1244
8.18	problem 4(b)	1248
8.19	problem 4(c)	1254
8.20	problem 4(d)	1259

8.1 problem 1(a)

8.1.1	Solving as linear ode	1031
8.1.2	Solving as homogeneousTypeD2 ode	1033
8.1.3	Solving as differentialType ode	1035
8.1.4	Solving as first order ode lie symmetry lookup ode	1036
8.1.5	Solving as exact ode	1040
8.1.6	Maple step by step solution	1044

Internal problem ID [6249]

Internal file name [OUTPUT/5497_Sunday_June_05_2022_03_41_36_PM_62340482/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Problems for Review and Discovery. Page 53

Problem number: 1(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "linear", "differentialType", "homogeneousTypeD2", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

`[_linear]`

$$xy' + y = x$$

8.1.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = \frac{1}{x}$$
$$q(x) = 1$$

Hence the ode is

$$y' + \frac{y}{x} = 1$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int \frac{1}{x} dx} \\ &= x\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= \mu \\ \frac{d}{dx}(xy) &= x \\ d(xy) &= x dx\end{aligned}$$

Integrating gives

$$\begin{aligned}xy &= \int x dx \\ xy &= \frac{x^2}{2} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = x$ results in

$$y = \frac{x}{2} + \frac{c_1}{x}$$

Summary

The solution(s) found are the following

$$y = \frac{x}{2} + \frac{c_1}{x} \tag{1}$$




Figure 199: Slope field plot

Verification of solutions

$$y = \frac{x}{2} + \frac{c_1}{x}$$

Verified OK.

8.1.2 Solving as homogeneousTypeD2 ode

Using the change of variables $y = u(x)x$ on the above ode results in new ode in $u(x)$

$$x(u'(x)x + u(x)) + u(x)x = x$$

In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= \frac{-2u + 1}{x} \end{aligned}$$

Where $f(x) = \frac{1}{x}$ and $g(u) = -2u + 1$. Integrating both sides gives

$$\frac{1}{-2u + 1} du = \frac{1}{x} dx$$

$$\int \frac{1}{-2u+1} du = \int \frac{1}{x} dx$$

$$-\frac{\ln(-2u+1)}{2} = \ln(x) + c_2$$

Raising both side to exponential gives

$$\frac{1}{\sqrt{-2u+1}} = e^{\ln(x)+c_2}$$

Which simplifies to

$$\frac{1}{\sqrt{-2u+1}} = c_3 x$$

Therefore the solution y is

$$y = ux$$

$$= \frac{(e^{2c_2} c_3^2 x^2 - 1) e^{-2c_2}}{2x c_3^2}$$

Summary

The solution(s) found are the following

$$y = \frac{(e^{2c_2} c_3^2 x^2 - 1) e^{-2c_2}}{2x c_3^2} \quad (1)$$




Figure 200: Slope field plot

Verification of solutions

$$y = \frac{(e^{2c_2} c_3^2 x^2 - 1) e^{-2c_2}}{2x c_3^2}$$

Verified OK.

8.1.3 Solving as differential Type ode

Writing the ode as

$$y' = \frac{x - y}{x} \quad (1)$$

Which becomes

$$0 = (-x) dy + (x - y) dx \quad (2)$$

But the RHS is complete differential because

$$(-x) dy + (x - y) dx = d\left(\frac{1}{2}x^2 - xy\right)$$

Hence (2) becomes

$$0 = d\left(\frac{1}{2}x^2 - xy\right)$$

Integrating both sides gives gives these solutions

$$y = \frac{x^2 + 2c_1}{2x} + c_1$$

Summary

The solution(s) found are the following

$$y = \frac{x^2 + 2c_1}{2x} + c_1 \quad (1)$$




Figure 201: Slope field plot

Verification of solutions

$$y = \frac{x^2 + 2c_1}{2x} + c_1$$

Verified OK.

8.1.4 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = -\frac{-x + y}{x}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 135: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int(n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= \frac{1}{x}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{1}{x}} dy \end{aligned}$$

Which results in

$$S = xy$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -\frac{-x + y}{x}$$

Evaluating all the partial derivatives gives

$$R_x = 1$$

$$R_y = 0$$

$$S_x = y$$

$$S_y = x$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = x \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = R$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by

integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \frac{R^2}{2} + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$xy = \frac{x^2}{2} + c_1$$



Which simplifies to

$$xy = \frac{x^2}{2} + c_1$$

Which gives

$$y = \frac{x^2 + 2c_1}{2x}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -\frac{-x+y}{x}$ 	$R = x$ $S = xy$	$\frac{dS}{dR} = R$ 

Summary

The solution(s) found are the following

$$y = \frac{x^2 + 2c_1}{2x} \quad (1)$$




Figure 202: Slope field plot

Verification of solutions

$$y = \frac{x^2 + 2c_1}{2x}$$

Verified OK.

8.1.5 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial\phi}{\partial x} + \frac{\partial\phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}(x) dy &= (x - y) dx \\ (-x + y) dx + (x) dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -x + y \\ N(x, y) &= x\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-x + y) \\ &= 1\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(x) \\ &= 1\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\int \frac{\partial \phi}{\partial x} dx = \int M dx$$

$$\int \frac{\partial \phi}{\partial x} dx = \int -x + y dx$$

$$\phi = -\frac{x(x - 2y)}{2} + f(y) \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = x + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = x$. Therefore equation (4) becomes

$$x = x + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = -\frac{x(x - 2y)}{2} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\frac{x(x - 2y)}{2}$$

The solution becomes

$$y = \frac{x^2 + 2c_1}{2x}$$

Summary

The solution(s) found are the following

$$y = \frac{x^2 + 2c_1}{2x} \tag{1}$$




Figure 203: Slope field plot

Verification of solutions

$$y = \frac{x^2 + 2c_1}{2x}$$

Verified OK.

8.1.6 Maple step by step solution

Let's solve

$$xy' + y = x$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = 1 - \frac{y}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' + \frac{y}{x} = 1$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) \left(y' + \frac{y}{x} \right) = \mu(x)$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x)y)$

$$\mu(x) \left(y' + \frac{y}{x} \right) = \mu'(x)y + \mu(x)y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = \frac{\mu(x)}{x}$$

- Solve to find the integrating factor

$$\mu(x) = x$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x)y) \right) dx = \int \mu(x) dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x)y = \int \mu(x) dx + c_1$$

- Solve for y

$$y = \frac{\int \mu(x) dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = x$

$$y = \frac{\int x dx + c_1}{x}$$

- Evaluate the integrals on the rhs

$$y = \frac{\frac{x^2}{2} + c_1}{x}$$

- Simplify

$$y = \frac{x^2 + 2c_1}{2x}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(x*diff(y(x),x)+y(x)=x,y(x), singsol=all)
```

$$y(x) = \frac{x}{2} + \frac{c_1}{x}$$

✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 17

```
DSolve[x*y'[x]+y[x]==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x}{2} + \frac{c_1}{x}$$

8.2 problem 1(b)

8.2.1	Solving as linear ode	1046
8.2.2	Solving as first order ode lie symmetry lookup ode	1048
8.2.3	Solving as exact ode	1052
8.2.4	Maple step by step solution	1057

Internal problem ID [6250]

Internal file name [OUTPUT/5498_Sunday_June_05_2022_03_41_38_PM_88803413/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Problems for Review and Discovery. Page 53

Problem number: 1(b).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "linear", "exactWithIntegrationFactor", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

`[_linear]`

$$x^2y' + y = x^2$$

8.2.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = \frac{1}{x^2}$$

$$q(x) = 1$$

Hence the ode is

$$y' + \frac{y}{x^2} = 1$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int \frac{1}{x^2} dx} \\ &= e^{-\frac{1}{x}}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= \mu \\ \frac{d}{dx}\left(e^{-\frac{1}{x}} y\right) &= e^{-\frac{1}{x}} \\ d\left(e^{-\frac{1}{x}} y\right) &= e^{-\frac{1}{x}} dx\end{aligned}$$

Integrating gives

$$\begin{aligned}e^{-\frac{1}{x}} y &= \int e^{-\frac{1}{x}} dx \\ e^{-\frac{1}{x}} y &= e^{-\frac{1}{x}} x - \text{expIntegral}_1\left(\frac{1}{x}\right) + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{-\frac{1}{x}}$ results in

$$y = e^{\frac{1}{x}} \left(e^{-\frac{1}{x}} x - \text{expIntegral}_1\left(\frac{1}{x}\right) \right) + c_1 e^{\frac{1}{x}}$$

which simplifies to

$$y = x - \text{expIntegral}_1\left(\frac{1}{x}\right) e^{\frac{1}{x}} + c_1 e^{\frac{1}{x}}$$

Summary

The solution(s) found are the following

$$y = x - \text{expIntegral}_1\left(\frac{1}{x}\right) e^{\frac{1}{x}} + c_1 e^{\frac{1}{x}} \tag{1}$$




Figure 204: Slope field plot

Verification of solutions

$$y = x - \expIntegral_1 \left(\frac{1}{x} \right) e^{\frac{1}{x}} + c_1 e^{\frac{1}{x}}$$

Verified OK.

8.2.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = -\frac{-x^2 + y}{x^2}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 138: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= e^{\frac{1}{x}}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right)S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^{\frac{1}{x}}} dy \end{aligned}$$

Which results in

$$S = e^{-\frac{1}{x}} y$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -\frac{-x^2 + y}{x^2}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{e^{-\frac{1}{x}} y}{x^2} \\ S_y &= e^{-\frac{1}{x}} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = e^{-\frac{1}{x}} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = e^{-\frac{1}{R}}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = R e^{-\frac{1}{R}} - \text{expIntegral}_1\left(\frac{1}{R}\right) + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$e^{-\frac{1}{x}}y = e^{-\frac{1}{x}}x - \text{expIntegral}_1\left(\frac{1}{x}\right) + c_1$$



Which simplifies to

$$(-x + y) e^{-\frac{1}{x}} - c_1 + \text{expIntegral}_1\left(\frac{1}{x}\right) = 0$$

Which gives

$$y = \left(e^{-\frac{1}{x}}x - \text{expIntegral}_1\left(\frac{1}{x}\right) + c_1 \right) e^{\frac{1}{x}}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -\frac{-x^2+y}{x^2}$ 	$R = x$ $S = e^{-\frac{1}{x}}y$	$\frac{dS}{dR} = e^{-\frac{1}{R}}$ 

Summary

The solution(s) found are the following

$$y = \left(e^{-\frac{1}{x}} x - \text{expIntegral}_1\left(\frac{1}{x}\right) + c_1 \right) e^{\frac{1}{x}} \quad (1)$$




Figure 205: Slope field plot

Verification of solutions

$$y = \left(e^{-\frac{1}{x}} x - \text{expIntegral}_1\left(\frac{1}{x}\right) + c_1 \right) e^{\frac{1}{x}}$$

Verified OK.

8.2.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (A)$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the

ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial\phi}{\partial x} + \frac{\partial\phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} (x^2) dy &= (x^2 - y) dx \\ (-x^2 + y) dx + (x^2) dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -x^2 + y \\ N(x, y) &= x^2 \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-x^2 + y) \\ &= 1\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(x^2) \\ &= 2x\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned}A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= \frac{1}{x^2} ((1) - (2x)) \\ &= \frac{1 - 2x}{x^2}\end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int A dx} \\ &= e^{\int \frac{1-2x}{x^2} dx}\end{aligned}$$

The result of integrating gives

$$\begin{aligned}\mu &= e^{-2\ln(x) - \frac{1}{x}} \\ &= \frac{e^{-\frac{1}{x}}}{x^2}\end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} for now so not to confuse them with the original M and N .

$$\begin{aligned}\bar{M} &= \mu M \\ &= \frac{e^{-\frac{1}{x}}}{x^2} (-x^2 + y) \\ &= -\frac{e^{-\frac{1}{x}}(x^2 - y)}{x^2}\end{aligned}$$

And

$$\begin{aligned}\bar{N} &= \mu N \\ &= \frac{e^{-\frac{1}{x}}}{x^2} (x^2) \\ &= e^{-\frac{1}{x}}\end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned}\bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ \left(-\frac{e^{-\frac{1}{x}}(x^2 - y)}{x^2} \right) + \left(e^{-\frac{1}{x}} \right) \frac{dy}{dx} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\frac{e^{-\frac{1}{x}}(x^2 - y)}{x^2} dx \\ \phi &= (-x + y) e^{-\frac{1}{x}} + \text{expIntegral}_1 \left(\frac{1}{x} \right) + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = e^{-\frac{1}{x}} + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = e^{-\frac{1}{x}}$. Therefore equation (4) becomes

$$e^{-\frac{1}{x}} = e^{-\frac{1}{x}} + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = (-x + y) e^{-\frac{1}{x}} + \expIntegral_1\left(\frac{1}{x}\right) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = (-x + y) e^{-\frac{1}{x}} + \expIntegral_1\left(\frac{1}{x}\right)$$

The solution becomes

$$y = \left(e^{-\frac{1}{x}} x - \expIntegral_1\left(\frac{1}{x}\right) + c_1 \right) e^{\frac{1}{x}}$$

Summary

The solution(s) found are the following

$$y = \left(e^{-\frac{1}{x}} x - \expIntegral_1\left(\frac{1}{x}\right) + c_1 \right) e^{\frac{1}{x}} \quad (1)$$




Figure 206: Slope field plot

Verification of solutions

$$y = \left(e^{-\frac{1}{x}} x - \text{expIntegral}_1\left(\frac{1}{x}\right) + c_1 \right) e^{\frac{1}{x}}$$

Verified OK.

8.2.4 Maple step by step solution

Let's solve

$$x^2 y' + y = x^2$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = 1 - \frac{y}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' + \frac{y}{x^2} = 1$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) \left(y' + \frac{y}{x^2} \right) = \mu(x)$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x)y)$

$$\mu(x) \left(y' + \frac{y}{x^2} \right) = \mu'(x)y + \mu(x)y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = \frac{\mu(x)}{x^2}$$

- Solve to find the integrating factor

$$\mu(x) = e^{-\frac{1}{x}}$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x)y) \right) dx = \int \mu(x) dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x)y = \int \mu(x) dx + c_1$$

- Solve for y

$$y = \frac{\int \mu(x) dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = e^{-\frac{1}{x}}$

$$y = \frac{\int e^{-\frac{1}{x}} dx + c_1}{e^{-\frac{1}{x}}}$$

- Evaluate the integrals on the rhs

$$y = \frac{e^{-\frac{1}{x}} x - \text{Ei}_1\left(\frac{1}{x}\right) + c_1}{e^{-\frac{1}{x}}}$$

- Simplify

$$y = x - \text{Ei}_1\left(\frac{1}{x}\right) e^{\frac{1}{x}} + c_1 e^{\frac{1}{x}}$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(x^2*diff(y(x),x)+y(x)=x^2,y(x), singsol=all)
```

$$y(x) = x - \operatorname{expIntegral}_1\left(\frac{1}{x}\right) e^{\frac{1}{x}} + e^{\frac{1}{x}} c_1$$

✓ Solution by Mathematica

Time used: 0.036 (sec). Leaf size: 27

```
DSolve[x^2*y'[x]+y[x]==x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{\frac{1}{x}} \operatorname{ExpIntegralEi}\left(-\frac{1}{x}\right) + x + c_1 e^{\frac{1}{x}}$$

8.3 problem 1(c)

8.3.1	Solving as separable ode	1060
8.3.2	Solving as linear ode	1062
8.3.3	Solving as homogeneousTypeD2 ode	1063
8.3.4	Solving as first order ode lie symmetry lookup ode	1065
8.3.5	Solving as exact ode	1069
8.3.6	Maple step by step solution	1073

Internal problem ID [6251]

Internal file name [OUTPUT/5499_Sunday_June_05_2022_03_41_39_PM_75934341/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Problems for Review and Discovery. Page 53

Problem number: 1(c).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "linear", "separable", "homogeneousTypeD2", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_separable]

$$x^2y' - y = 0$$

8.3.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{y}{x^2}\end{aligned}$$

Where $f(x) = \frac{1}{x^2}$ and $g(y) = y$. Integrating both sides gives

$$\begin{aligned}\frac{1}{y} dy &= \frac{1}{x^2} dx \\ \int \frac{1}{y} dy &= \int \frac{1}{x^2} dx \\ \ln(y) &= c_1 - \frac{1}{x} \\ y &= e^{c_1 - \frac{1}{x}} \\ &= c_1 e^{-\frac{1}{x}}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-\frac{1}{x}} \tag{1}$$




Figure 207: Slope field plot

Verification of solutions

$$y = c_1 e^{-\frac{1}{x}}$$

Verified OK.

8.3.2 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -\frac{1}{x^2}$$
$$q(x) = 0$$

Hence the ode is

$$y' - \frac{y}{x^2} = 0$$

The integrating factor μ is

$$\mu = e^{\int -\frac{1}{x^2} dx}$$
$$= e^{\frac{1}{x}}$$

The ode becomes

$$\frac{d}{dx} \mu y = 0$$
$$\frac{d}{dx} \left(e^{\frac{1}{x}} y \right) = 0$$

Integrating gives

$$e^{\frac{1}{x}} y = c_1$$

Dividing both sides by the integrating factor $\mu = e^{\frac{1}{x}}$ results in

$$y = c_1 e^{-\frac{1}{x}}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-\frac{1}{x}} \tag{1}$$




Figure 208: Slope field plot

Verification of solutions

$$y = c_1 e^{-\frac{1}{x}}$$

Verified OK.

8.3.3 Solving as homogeneousTypeD2 ode

Using the change of variables $y = u(x)x$ on the above ode results in new ode in $u(x)$

$$x^2(u'(x)x + u(x)) - u(x)x = 0$$

In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{u(x-1)}{x^2} \end{aligned}$$

Where $f(x) = -\frac{x-1}{x^2}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u} du &= -\frac{x-1}{x^2} dx \\ \int \frac{1}{u} du &= \int -\frac{x-1}{x^2} dx \\ \ln(u) &= -\ln(x) - \frac{1}{x} + c_2 \\ u &= e^{-\ln(x) - \frac{1}{x} + c_2} \\ &= c_2 e^{-\ln(x) - \frac{1}{x}}\end{aligned}$$

Which simplifies to

$$u(x) = \frac{c_2 e^{-\frac{1}{x}}}{x}$$

Therefore the solution y is

$$\begin{aligned}y &= ux \\ &= c_2 e^{-\frac{1}{x}}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_2 e^{-\frac{1}{x}} \tag{1}$$




Figure 209: Slope field plot

Verification of solutions

$$y = c_2 e^{-\frac{1}{x}}$$

Verified OK.

8.3.4 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{y}{x^2}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 141: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= e^{-\frac{1}{x}}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right)S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^{-\frac{1}{x}}} dy \end{aligned}$$

Which results in

$$S = e^{\frac{1}{x}} y$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{y}{x^2}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= -\frac{e^{\frac{1}{x}} y}{x^2} \\ S_y &= e^{\frac{1}{x}} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \tag{4}$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$e^{\frac{1}{x}} y = c_1$$



Which simplifies to

$$e^{\frac{1}{x}} y = c_1$$

Which gives

$$y = c_1 e^{-\frac{1}{x}}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{y}{x^2}$ 	$R = x$ $S = e^{\frac{1}{x}} y$	$\frac{dS}{dR} = 0$ 

Summary

The solution(s) found are the following

$$y = c_1 e^{-\frac{1}{x}} \tag{1}$$




Figure 210: Slope field plot

Verification of solutions

$$y = c_1 e^{-\frac{1}{x}}$$

Verified OK.

8.3.5 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}\left(\frac{1}{y}\right) dy &= \left(\frac{1}{x^2}\right) dx \\ \left(-\frac{1}{x^2}\right) dx + \left(\frac{1}{y}\right) dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -\frac{1}{x^2} \\ N(x, y) &= \frac{1}{y}\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(-\frac{1}{x^2}\right) \\ &= 0\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{1}{y} \right) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \tag{1}$$

$$\frac{\partial \phi}{\partial y} = N \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\frac{1}{x^2} dx \\ \phi &= \frac{1}{x} + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{1}{y}$. Therefore equation (4) becomes

$$\frac{1}{y} = 0 + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = \frac{1}{y}$$

Integrating the above w.r.t y gives

$$\begin{aligned}\int f'(y) dy &= \int \left(\frac{1}{y} \right) dy \\ f(y) &= \ln(y) + c_1\end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = \frac{1}{x} + \ln(y) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \frac{1}{x} + \ln(y)$$

The solution becomes

$$y = e^{\frac{c_1 x - 1}{x}}$$

Summary

The solution(s) found are the following

$$y = e^{\frac{c_1 x - 1}{x}} \tag{1}$$




Figure 211: Slope field plot

Verification of solutions

$$y = e^{\frac{c_1 x - 1}{x}}$$

Verified OK.

8.3.6 Maple step by step solution

Let's solve

$$x^2 y' - y = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y} = \frac{1}{x^2}$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y} dx = \int \frac{1}{x^2} dx + c_1$$

- Evaluate integral

$$\ln(y) = c_1 - \frac{1}{x}$$

- Solve for y

$$y = e^{\frac{c_1 x - 1}{x}}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(x^2*diff(y(x),x)=y(x),y(x), singsol=all)
```

$$y(x) = c_1 e^{-\frac{1}{x}}$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 20

```
DSolve[x^2*y'[x]==y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-1/x}$$

$$y(x) \rightarrow 0$$

8.4 problem 1(d)

8.4.1	Solving as separable ode	1075
8.4.2	Solving as first order ode lie symmetry lookup ode	1077
8.4.3	Solving as exact ode	1081
8.4.4	Maple step by step solution	1085

Internal problem ID [6252]

Internal file name [OUTPUT/5500_Sunday_June_05_2022_03_41_40_PM_74561563/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Problems for Review and Discovery. Page 53

Problem number: 1(d).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exact", "separable", "first_order_ode_lie_symmetry_lookup"**

Maple gives the following as the ode type

`[_separable]`

$$y' \sec(x) - \sec(y) = 0$$

8.4.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{\sec(y)}{\sec(x)} \end{aligned}$$

Where $f(x) = \frac{1}{\sec(x)}$ and $g(y) = \sec(y)$. Integrating both sides gives

$$\frac{1}{\sec(y)} dy = \frac{1}{\sec(x)} dx$$

$$\int \frac{1}{\sec(y)} dy = \int \frac{1}{\sec(x)} dx$$

$$\sin(y) = \sin(x) + c_1$$

Which results in

$$y = \arcsin(\sin(x) + c_1)$$

Summary

The solution(s) found are the following

$$y = \arcsin(\sin(x) + c_1) \tag{1}$$




Figure 212: Slope field plot

Verification of solutions

$$y = \arcsin(\sin(x) + c_1)$$

Verified OK.

8.4.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{\sec(y)}{\sec(x)}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **separable**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 144: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= \sec(x) \\ \eta(x, y) &= 0\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\eta = 0$ then in this special case

$$R = y$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\xi} dx \\ &= \int \frac{1}{\sec(x)} dx\end{aligned}$$

Which results in

$$S = \sin(x)$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y}\tag{2}$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{\sec(y)}{\sec(x)}$$

Evaluating all the partial derivatives gives

$$\begin{aligned}R_x &= 0 \\R_y &= 1 \\S_x &= \cos(x) \\S_y &= 0\end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \cos(y) \tag{2A}$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \cos(R)$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \sin(R) + c_1 \tag{4}$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\sin(x) = \sin(y) + c_1$$



Which simplifies to

$$\sin(x) = \sin(y) + c_1$$

Which gives

$$y = -\arcsin(-\sin(x) + c_1)$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{\sec(y)}{\sec(x)}$ 	$R = y$ $S = \sin(x)$	$\frac{dS}{dR} = \cos(R)$ 

Summary

The solution(s) found are the following

$$y = -\arcsin(-\sin(x) + c_1) \tag{1}$$




Figure 213: Slope field plot

Verification of solutions

$$y = -\arcsin(-\sin(x) + c_1)$$

Verified OK.

8.4.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned}\left(\frac{1}{\sec(y)}\right) dy &= \left(\frac{1}{\sec(x)}\right) dx \\ \left(-\frac{1}{\sec(x)}\right) dx + \left(\frac{1}{\sec(y)}\right) dy &= 0\end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -\frac{1}{\sec(x)} \\ N(x, y) &= \frac{1}{\sec(y)}\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(-\frac{1}{\sec(x)}\right) \\ &= 0\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{1}{\sec(y)} \right) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \tag{1}$$

$$\frac{\partial \phi}{\partial y} = N \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\frac{1}{\sec(x)} dx \\ \phi &= -\sin(x) + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{1}{\sec(y)}$. Therefore equation (4) becomes

$$\frac{1}{\sec(y)} = 0 + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$\begin{aligned}f'(y) &= \frac{1}{\sec(y)} \\ &= \cos(y)\end{aligned}$$

Integrating the above w.r.t y results in

$$\int f'(y) dy = \int (\cos(y)) dy$$
$$f(y) = \sin(y) + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\sin(x) + \sin(y) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\sin(x) + \sin(y)$$

Summary

The solution(s) found are the following

$$-\sin(x) + \sin(y) = c_1 \tag{1}$$




Figure 214: Slope field plot

Verification of solutions

$$-\sin(x) + \sin(y) = c_1$$

Verified OK.

8.4.4 Maple step by step solution

Let's solve

$$y' \sec(x) - \sec(y) = 0$$

- Highest derivative means the order of the ODE is 1

y'

- Separate variables

$$\frac{y'}{\sec(y)} = \frac{1}{\sec(x)}$$

- Integrate both sides with respect to x

$$\int \frac{y'}{\sec(y)} dx = \int \frac{1}{\sec(x)} dx + c_1$$

- Evaluate integral

$$\sin(y) = \sin(x) + c_1$$

- Solve for y

$$y = \arcsin(\sin(x) + c_1)$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
<- separable successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 9

```
dsolve(sec(x)*diff(y(x),x)=sec(y(x)),y(x), singsol=all)
```

$$y(x) = \arcsin(\sin(x) + c_1)$$

✓ Solution by Mathematica

Time used: 0.335 (sec). Leaf size: 11

```
DSolve[Sec[x]*y'[x]==Sec[y[x]],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \arcsin(\sin(x) + c_1)$$

8.5 problem 1(e)

- 8.5.1 Solving as homogeneousTypeD2 ode 1087
- 8.5.2 Solving as first order ode lie symmetry calculated ode 1089

Internal problem ID [6253]

Internal file name [OUTPUT/5501_Sunday_June_05_2022_03_41_42_PM_4078066/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Problems for Review and Discovery. Page 53

Problem number: 1(e).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**homogeneousTypeD2**", "**first_order_ode_lie_symmetry_calculated**"

Maple gives the following as the ode type

```
[[_homogeneous, `class A`], _rational, _dAlembert]
```

$$y' - \frac{x^2 + y^2}{x^2 - y^2} = 0$$

8.5.1 Solving as homogeneousTypeD2 ode

Using the change of variables $y = u(x)x$ on the above ode results in new ode in $u(x)$

$$u'(x)x + u(x) - \frac{x^2 + u(x)^2 x^2}{x^2 - u(x)^2 x^2} = 0$$

In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{u^3 + u^2 - u + 1}{x(u^2 - 1)} \end{aligned}$$

Where $f(x) = -\frac{1}{x}$ and $g(u) = \frac{u^3+u^2-u+1}{u^2-1}$. Integrating both sides gives

$$\begin{aligned}\frac{1}{\frac{u^3+u^2-u+1}{u^2-1}} du &= -\frac{1}{x} dx \\ \int \frac{1}{\frac{u^3+u^2-u+1}{u^2-1}} du &= \int -\frac{1}{x} dx \\ \int^u \frac{u^2-1}{u^3+u^2-u+1} du &= -\ln(x) + c_2\end{aligned}$$

Which results in

$$\int^u \frac{u^2-1}{u^3+u^2-u+1} du = -\ln(x) + c_2$$

The solution is

$$\int^{u(x)} \frac{u^2-1}{u^3+u^2-u+1} du + \ln(x) - c_2 = 0$$

Replacing $u(x)$ in the above solution by $\frac{y}{x}$ results in the solution for y in implicit form

$$\begin{aligned}\int^{\frac{y}{x}} \frac{u^2-1}{u^3+u^2-u+1} du + \ln(x) - c_2 &= 0 \\ \int^{\frac{y}{x}} \frac{u^2-1}{u^3+u^2-u+1} du + \ln(x) - c_2 &= 0\end{aligned}$$

Summary

The solution(s) found are the following

$$\int^{\frac{y}{x}} \frac{u^2-1}{u^3+u^2-u+1} du + \ln(x) - c_2 = 0 \quad (1)$$




Figure 215: Slope field plot

Verification of solutions

$$\int^{\frac{y}{x}} \frac{-a^2 - 1}{-a^3 + -a^2 - -a + 1} da + \ln(x) - c_2 = 0$$

Verified OK.

8.5.2 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = -\frac{x^2 + y^2}{-x^2 + y^2}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \tag{A}$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \tag{1E}$$

$$\eta = xb_2 + yb_3 + b_1 \tag{2E}$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$\begin{aligned} b_2 - \frac{(x^2 + y^2)(b_3 - a_2)}{-x^2 + y^2} - \frac{(x^2 + y^2)^2 a_3}{(-x^2 + y^2)^2} \\ - \left(-\frac{2x}{-x^2 + y^2} - \frac{2(x^2 + y^2)x}{(-x^2 + y^2)^2} \right) (xa_2 + ya_3 + a_1) \\ - \left(-\frac{2y}{-x^2 + y^2} + \frac{2(x^2 + y^2)y}{(-x^2 + y^2)^2} \right) (xb_2 + yb_3 + b_1) = 0 \end{aligned} \quad (5E)$$

Putting the above in normal form gives

$$\frac{x^4 a_2 + x^4 a_3 - x^4 b_2 - x^4 b_3 + 4x^3 y b_2 - 4x^2 y^2 a_2 + 2x^2 y^2 a_3 + 2x^2 y^2 b_2 + 4x^2 y^2 b_3 - 4x y^3 a_3 - y^4 a_2 + y^4 a_3}{(x^2 - y^2)^2} = 0$$

Setting the numerator to zero gives

$$\begin{aligned} -x^4 a_2 - x^4 a_3 + x^4 b_2 + x^4 b_3 - 4x^3 y b_2 + 4x^2 y^2 a_2 - 2x^2 y^2 a_3 - 2x^2 y^2 b_2 \\ - 4x^2 y^2 b_3 + 4x y^3 a_3 + y^4 a_2 - y^4 a_3 + y^4 b_2 - y^4 b_3 - 4x^2 y b_1 + 4x y^2 a_1 = 0 \end{aligned} \quad (6E)$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2\}$$

The above PDE (6E) now becomes

$$\begin{aligned} -a_2 v_1^4 + 4a_2 v_1^2 v_2^2 + a_2 v_2^4 - a_3 v_1^4 - 2a_3 v_1^2 v_2^2 + 4a_3 v_1 v_2^3 - a_3 v_2^4 + b_2 v_1^4 - 4b_2 v_1^3 v_2 \\ - 2b_2 v_1^2 v_2^2 + b_2 v_2^4 + b_3 v_1^4 - 4b_3 v_1^2 v_2^2 - b_3 v_2^4 + 4a_1 v_1 v_2^2 - 4b_1 v_1^2 v_2 = 0 \end{aligned} \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2\}$$

Equation (7E) now becomes

$$\begin{aligned} &(-a_2 - a_3 + b_2 + b_3)v_1^4 - 4b_2v_1^3v_2 + (4a_2 - 2a_3 - 2b_2 - 4b_3)v_1^2v_2^2 \\ &- 4b_1v_1^2v_2 + 4a_3v_1v_2^3 + 4a_1v_1v_2^2 + (a_2 - a_3 + b_2 - b_3)v_2^4 = 0 \end{aligned} \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} 4a_1 &= 0 \\ 4a_3 &= 0 \\ -4b_1 &= 0 \\ -4b_2 &= 0 \\ -a_2 - a_3 + b_2 + b_3 &= 0 \\ a_2 - a_3 + b_2 - b_3 &= 0 \\ 4a_2 - 2a_3 - 2b_2 - 4b_3 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= 0 \\ a_2 &= b_3 \\ a_3 &= 0 \\ b_1 &= 0 \\ b_2 &= 0 \\ b_3 &= b_3 \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned} \xi &= x \\ \eta &= y \end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Therefore

$$\begin{aligned} \frac{dy}{dx} &= \frac{\eta}{\xi} \\ &= \frac{y}{x} \\ &= \frac{y}{x} \end{aligned}$$

This is easily solved to give

$$y = c_1 x$$

Where now the coordinate R is taken as the constant of integration. Hence

$$R = \frac{y}{x}$$

And S is found from

$$\begin{aligned} dS &= \frac{dx}{\xi} \\ &= \frac{dx}{x} \end{aligned}$$

Integrating gives

$$\begin{aligned} S &= \int \frac{dx}{x} \\ &= \ln(x) \end{aligned}$$

Where the constant of integration is set to zero as we just need one solution. Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -\frac{x^2 + y^2}{-x^2 + y^2}$$

Evaluating all the partial derivatives gives

$$\begin{aligned}R_x &= -\frac{y}{x^2} \\R_y &= \frac{1}{x} \\S_x &= \frac{1}{x} \\S_y &= 0\end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \frac{x^3 - y^2x}{x^3 - yx^2 + y^2x + y^3} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \frac{-R^2 + 1}{R^3 + R^2 - R + 1}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \int -\frac{R^2 - 1}{R^3 + R^2 - R + 1} dR + c_1 \quad (4)$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\ln(x) = \int^{\frac{y}{x}} -\frac{-a^2 - 1}{-a^3 + -a^2 - -a + 1} d-a + c_1$$

Which simplifies to

$$\ln(x) = \int^{\frac{y}{x}} -\frac{-a^2 - 1}{-a^3 + -a^2 - -a + 1} d-a + c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -\frac{x^2+y^2}{-x^2+y^2}$ 	$R = \frac{y}{x}$ $S = \ln(x)$	$\frac{dS}{dR} = \frac{-R^2+1}{R^3+R^2-R+1}$ 

Summary

The solution(s) found are the following

$$\ln(x) = \int^{\frac{y}{x}} \frac{-a^2 - 1}{-a^3 + a^2 - a + 1} da + c_1 \tag{1}$$




Figure 216: Slope field plot

Verification of solutions

$$\ln(x) = \int^x \frac{-a^2 - 1}{-a^3 + -a^2 - a + 1} da + c_1$$

Verified OK.

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying homogeneous D
<- homogeneous successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 34

```
dsolve(diff(y(x),x)=(x^2+y(x)^2)/(x^2-y(x)^2),y(x), singsol=all)
```

$$y(x) = \text{RootOf} \left(\int^{-Z} \frac{-a^2 - 1}{-a^3 + -a^2 - -a + 1} d-a + \ln(x) + c_1 \right) x$$

✓ Solution by Mathematica

Time used: 0.115 (sec). Leaf size: 67

```
DSolve[y'[x]==(x^2+y[x]^2)/(x^2-y[x]^2),y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[\text{RootSum} \left[\#1^3 + \#1^2 - \#1 \right. \right. \\ \left. \left. + 1 \&, \frac{\#1 \log \left(\frac{y(x)}{x} - \#1 \right) - \log \left(\frac{y(x)}{x} - \#1 \right)}{3\#1 - 1} \& \right] = -\log(x) + c_1, y(x) \right]$$

8.6 problem 1(f)

- 8.6.1 Solving as homogeneousTypeD2 ode 1097
- 8.6.2 Solving as first order ode lie symmetry calculated ode 1099
- 8.6.3 Solving as exact ode 1104

Internal problem ID [6254]

Internal file name [OUTPUT/5502_Sunday_June_05_2022_03_41_43_PM_36474534/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Problems for Review and Discovery. Page 53

Problem number: 1(f).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exactByInspection", "homogeneousTypeD2", "first_order_ode_lie_symmetry_calculated"

Maple gives the following as the ode type

```
[[_homogeneous, `class A`], _rational, [_Abel, `2nd type`, `class A`]]
```

$$y' - \frac{2y + x}{2x - y} = 0$$

8.6.1 Solving as homogeneousTypeD2 ode

Using the change of variables $y = u(x)x$ on the above ode results in new ode in $u(x)$

$$u'(x)x + u(x) - \frac{2u(x)x + x}{2x - u(x)x} = 0$$

In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{u^2 + 1}{x(u - 2)} \end{aligned}$$

Where $f(x) = -\frac{1}{x}$ and $g(u) = \frac{u^2+1}{u-2}$. Integrating both sides gives

$$\frac{1}{\frac{u^2+1}{u-2}} du = -\frac{1}{x} dx$$
$$\int \frac{1}{\frac{u^2+1}{u-2}} du = \int -\frac{1}{x} dx$$

$$\frac{\ln(u^2 + 1)}{2} - 2 \arctan(u) = -\ln(x) + c_2$$

The solution is

$$\frac{\ln(u(x)^2 + 1)}{2} - 2 \arctan(u(x)) + \ln(x) - c_2 = 0$$

Replacing $u(x)$ in the above solution by $\frac{y}{x}$ results in the solution for y in implicit form

$$\frac{\ln\left(\frac{y^2}{x^2} + 1\right)}{2} - 2 \arctan\left(\frac{y}{x}\right) + \ln(x) - c_2 = 0$$
$$\frac{\ln\left(\frac{y^2}{x^2} + 1\right)}{2} - 2 \arctan\left(\frac{y}{x}\right) + \ln(x) - c_2 = 0$$

Summary

The solution(s) found are the following

$$\frac{\ln\left(\frac{y^2}{x^2} + 1\right)}{2} - 2 \arctan\left(\frac{y}{x}\right) + \ln(x) - c_2 = 0 \quad (1)$$




Figure 217: Slope field plot

Verification of solutions

$$\frac{\ln\left(\frac{y^2}{x^2} + 1\right)}{2} - 2 \arctan\left(\frac{y}{x}\right) + \ln(x) - c_2 = 0$$

Verified OK.

8.6.2 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = -\frac{2y + x}{-2x + y}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \quad (1\text{E})$$

$$\eta = xb_2 + yb_3 + b_1 \quad (2\text{E})$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$\begin{aligned} b_2 - \frac{(2y+x)(b_3-a_2)}{-2x+y} - \frac{(2y+x)^2 a_3}{(-2x+y)^2} \\ - \left(-\frac{1}{-2x+y} - \frac{2(2y+x)}{(-2x+y)^2} \right) (xa_2 + ya_3 + a_1) \\ - \left(-\frac{2}{-2x+y} + \frac{2y+x}{(-2x+y)^2} \right) (xb_2 + yb_3 + b_1) = 0 \end{aligned} \quad (5E)$$

Putting the above in normal form gives

$$\frac{2x^2a_2 + x^2a_3 + x^2b_2 - 2x^2b_3 - 2xya_2 + 4xya_3 + 4xyb_2 + 2xyb_3 - 2y^2a_2 - y^2a_3 - y^2b_2 + 2y^2b_3 + 5xb_1 - 5ya_1}{(2x-y)^2} = 0$$

Setting the numerator to zero gives

$$\begin{aligned} -2x^2a_2 - x^2a_3 - x^2b_2 + 2x^2b_3 + 2xya_2 - 4xya_3 - 4xyb_2 \\ - 2xyb_3 + 2y^2a_2 + y^2a_3 + y^2b_2 - 2y^2b_3 - 5xb_1 + 5ya_1 = 0 \end{aligned} \quad (6E)$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2\}$$

The above PDE (6E) now becomes

$$\begin{aligned} -2a_2v_1^2 + 2a_2v_1v_2 + 2a_2v_2^2 - a_3v_1^2 - 4a_3v_1v_2 + a_3v_2^2 - b_2v_1^2 \\ - 4b_2v_1v_2 + b_2v_2^2 + 2b_3v_1^2 - 2b_3v_1v_2 - 2b_3v_2^2 + 5a_1v_2 - 5b_1v_1 = 0 \end{aligned} \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2\}$$

Equation (7E) now becomes

$$\begin{aligned} (-2a_2 - a_3 - b_2 + 2b_3)v_1^2 + (2a_2 - 4a_3 - 4b_2 - 2b_3)v_1v_2 \\ - 5b_1v_1 + (2a_2 + a_3 + b_2 - 2b_3)v_2^2 + 5a_1v_2 = 0 \end{aligned} \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} 5a_1 &= 0 \\ -5b_1 &= 0 \\ -2a_2 - a_3 - b_2 + 2b_3 &= 0 \\ 2a_2 - 4a_3 - 4b_2 - 2b_3 &= 0 \\ 2a_2 + a_3 + b_2 - 2b_3 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= 0 \\ a_2 &= b_3 \\ a_3 &= -b_2 \\ b_1 &= 0 \\ b_2 &= b_2 \\ b_3 &= b_3 \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned} \xi &= x \\ \eta &= y \end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned} \eta &= \eta - \omega(x, y)\xi \\ &= y - \left(-\frac{2y+x}{-2x+y} \right) (x) \\ &= \frac{-x^2 - y^2}{2x - y} \\ \xi &= 0 \end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{-x^2-y^2}{2x-y}} dy \end{aligned}$$

Which results in

$$S = \frac{\ln(x^2 + y^2)}{2} - 2 \arctan\left(\frac{y}{x}\right)$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -\frac{2y + x}{-2x + y}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{2y + x}{x^2 + y^2} \\ S_y &= \frac{-2x + y}{x^2 + y^2} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \tag{2A}$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \tag{4}$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{\ln(x^2 + y^2)}{2} - 2 \arctan\left(\frac{y}{x}\right) = c_1$$

Which simplifies to

$$\frac{\ln(x^2 + y^2)}{2} - 2 \arctan\left(\frac{y}{x}\right) = c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -\frac{2y+x}{-2x+y}$ 	$R = x$ $S = \frac{\ln(x^2 + y^2)}{2} - 2 \arctan\left(\frac{y}{x}\right)$	$\frac{dS}{dR} = 0$ 

Summary

The solution(s) found are the following

$$\frac{\ln(x^2 + y^2)}{2} - 2 \arctan\left(\frac{y}{x}\right) = c_1 \quad (1)$$




Figure 218: Slope field plot

Verification of solutions

$$\frac{\ln(x^2 + y^2)}{2} - 2 \arctan\left(\frac{y}{x}\right) = c_1$$

Verified OK.

8.6.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (A)$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} (-2x + y) dy &= (-2y - x) dx \\ (2y + x) dx + (-2x + y) dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= 2y + x \\ N(x, y) &= -2x + y \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(2y + x) \\ &= 2 \end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(-2x + y) \\ &= -2\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. By inspection $\frac{1}{x^2+y^2}$ is an integrating factor. Therefore by multiplying $M = 2y + x$ and $N = -2x + y$ by this integrating factor the ode becomes exact. The new M, N are

$$\begin{aligned}M &= \frac{2y + x}{x^2 + y^2} \\ N &= \frac{-2x + y}{x^2 + y^2}\end{aligned}$$

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might

or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned} \left(\frac{-2x + y}{x^2 + y^2}\right) dy &= \left(-\frac{2y + x}{x^2 + y^2}\right) dx \\ \left(\frac{2y + x}{x^2 + y^2}\right) dx + \left(\frac{-2x + y}{x^2 + y^2}\right) dy &= 0 \end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= \frac{2y + x}{x^2 + y^2} \\ N(x, y) &= \frac{-2x + y}{x^2 + y^2} \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(\frac{2y + x}{x^2 + y^2}\right) \\ &= \frac{2x^2 - 2xy - 2y^2}{(x^2 + y^2)^2} \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{-2x + y}{x^2 + y^2}\right) \\ &= \frac{2x^2 - 2xy - 2y^2}{(x^2 + y^2)^2} \end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int \frac{2y + x}{x^2 + y^2} dx \\ \phi &= \frac{\ln(x^2 + y^2)}{2} + 2 \arctan\left(\frac{x}{y}\right) + f(y)\end{aligned}\quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\begin{aligned}\frac{\partial \phi}{\partial y} &= \frac{y}{x^2 + y^2} - \frac{2x}{y^2 \left(\frac{x^2}{y^2} + 1\right)} + f'(y) \\ &= \frac{-2x + y}{x^2 + y^2} + f'(y)\end{aligned}\quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{-2x + y}{x^2 + y^2}$. Therefore equation (4) becomes

$$\frac{-2x + y}{x^2 + y^2} = \frac{-2x + y}{x^2 + y^2} + f'(y)\quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = \frac{\ln(x^2 + y^2)}{2} + 2 \arctan\left(\frac{x}{y}\right) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \frac{\ln(x^2 + y^2)}{2} + 2 \arctan\left(\frac{x}{y}\right)$$

Summary

The solution(s) found are the following

$$\frac{\ln(x^2 + y^2)}{2} + 2 \arctan\left(\frac{x}{y}\right) = c_1 \quad (1)$$




Figure 219: Slope field plot

Verification of solutions

$$\frac{\ln(x^2 + y^2)}{2} + 2 \arctan\left(\frac{x}{y}\right) = c_1$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying homogeneous D  
<- homogeneous successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 24

```
dsolve(diff(y(x),x)=(x+2*y(x))/(2*x-y(x)),y(x), singsol=all)
```

$$y(x) = \tan \left(\text{RootOf} \left(-4_Z + \ln \left(\sec \left(_Z \right)^2 \right) + 2 \ln (x) + 2c_1 \right) \right) x$$

✓ Solution by Mathematica

Time used: 0.06 (sec). Leaf size: 36

```
DSolve[y'[x]==(x+2*y[x])/(2*x-y[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[\frac{1}{2} \log \left(\frac{y(x)^2}{x^2} + 1 \right) - 2 \arctan \left(\frac{y(x)}{x} \right) = -\log(x) + c_1, y(x) \right]$$

8.7 problem 1(g)

8.7.1	Solving as separable ode	1111
8.7.2	Solving as linear ode	1113
8.7.3	Solving as homogeneousTypeD2 ode	1114
8.7.4	Solving as first order ode lie symmetry lookup ode	1116
8.7.5	Solving as exact ode	1120
8.7.6	Maple step by step solution	1124

Internal problem ID [6255]

Internal file name [OUTPUT/5503_Sunday_June_05_2022_03_41_46_PM_59101253/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Problems for Review and Discovery. Page 53

Problem number: 1(g).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "linear", "separable", "homogeneousTypeD2", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_separable]

$$x^2y' + 2xy = 0$$

8.7.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= -\frac{2y}{x}\end{aligned}$$

Where $f(x) = -\frac{2}{x}$ and $g(y) = y$. Integrating both sides gives

$$\begin{aligned}\frac{1}{y} dy &= -\frac{2}{x} dx \\ \int \frac{1}{y} dy &= \int -\frac{2}{x} dx \\ \ln(y) &= -2 \ln(x) + c_1 \\ y &= e^{-2 \ln(x) + c_1} \\ &= \frac{c_1}{x^2}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1}{x^2} \tag{1}$$




Figure 220: Slope field plot

Verification of solutions

$$y = \frac{c_1}{x^2}$$

Verified OK.

8.7.2 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = \frac{2}{x}$$
$$q(x) = 0$$

Hence the ode is

$$y' + \frac{2y}{x} = 0$$

The integrating factor μ is

$$\mu = e^{\int \frac{2}{x} dx}$$
$$= x^2$$

The ode becomes

$$\frac{d}{dx} \mu y = 0$$
$$\frac{d}{dx} (y x^2) = 0$$

Integrating gives

$$y x^2 = c_1$$

Dividing both sides by the integrating factor $\mu = x^2$ results in

$$y = \frac{c_1}{x^2}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1}{x^2} \tag{1}$$




Figure 221: Slope field plot

Verification of solutions

$$y = \frac{c_1}{x^2}$$

Verified OK.

8.7.3 Solving as homogeneousTypeD2 ode

Using the change of variables $y = u(x)x$ on the above ode results in new ode in $u(x)$

$$x^2(u'(x)x + u(x)) + 2x^2u(x) = 0$$

In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{3u}{x} \end{aligned}$$

Where $f(x) = -\frac{3}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u} du &= -\frac{3}{x} dx \\ \int \frac{1}{u} du &= \int -\frac{3}{x} dx \\ \ln(u) &= -3 \ln(x) + c_2 \\ u &= e^{-3 \ln(x) + c_2} \\ &= \frac{c_2}{x^3}\end{aligned}$$

Therefore the solution y is

$$\begin{aligned}y &= xu \\ &= \frac{c_2}{x^2}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{c_2}{x^2} \tag{1}$$




Figure 222: Slope field plot

Verification of solutions

$$y = \frac{c_2}{x^2}$$

Verified OK.

8.7.4 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = -\frac{2y}{x}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 147: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= \frac{1}{x^2}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{1}{x^2}} dy \end{aligned}$$

Which results in

$$S = y x^2$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -\frac{2y}{x}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= 2xy \\ S_y &= x^2 \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by

integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \tag{4}$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$yx^2 = c_1$$



Which simplifies to

$$yx^2 = c_1$$

Which gives

$$y = \frac{c_1}{x^2}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -\frac{2y}{x}$ 	$R = x$ $S = yx^2$	$\frac{dS}{dR} = 0$ 

Summary

The solution(s) found are the following

$$y = \frac{c_1}{x^2} \tag{1}$$




Figure 223: Slope field plot

Verification of solutions

$$y = \frac{c_1}{x^2}$$

Verified OK.

8.7.5 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}\left(-\frac{1}{2y}\right) dy &= \left(\frac{1}{x}\right) dx \\ \left(-\frac{1}{x}\right) dx + \left(-\frac{1}{2y}\right) dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -\frac{1}{x} \\ N(x, y) &= -\frac{1}{2y}\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(-\frac{1}{x}\right) \\ &= 0\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(-\frac{1}{2y} \right) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \tag{1}$$

$$\frac{\partial \phi}{\partial y} = N \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\frac{1}{x} dx \\ \phi &= -\ln(x) + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = -\frac{1}{2y}$. Therefore equation (4) becomes

$$-\frac{1}{2y} = 0 + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = -\frac{1}{2y}$$

Integrating the above w.r.t y gives

$$\begin{aligned}\int f'(y) dy &= \int \left(-\frac{1}{2y} \right) dy \\ f(y) &= -\frac{\ln(y)}{2} + c_1\end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\ln(x) - \frac{\ln(y)}{2} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\ln(x) - \frac{\ln(y)}{2}$$

The solution becomes

$$y = \frac{e^{-2c_1}}{x^2}$$

Summary

The solution(s) found are the following

$$y = \frac{e^{-2c_1}}{x^2} \tag{1}$$




Figure 224: Slope field plot

Verification of solutions

$$y = \frac{e^{-2c_1}}{x^2}$$

Verified OK.

8.7.6 Maple step by step solution

Let's solve

$$x^2y' + 2xy = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Integrate both sides with respect to x

$$\int (x^2y' + 2xy) dx = \int 0 dx + c_1$$

- Evaluate integral

$$yx^2 = c_1$$

- Solve for y

$$y = \frac{c_1}{x^2}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
dsolve(2*x*y(x)+x^2*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{c_1}{x^2}$$

✓ Solution by Mathematica

Time used: 0.037 (sec). Leaf size: 16

```
DSolve[2*x*y[x]+x^2*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_1}{x^2}$$

$$y(x) \rightarrow 0$$

8.8 problem 1(h)

8.8.1	Solving as separable ode	1126
8.8.2	Solving as first order ode lie symmetry lookup ode	1128
8.8.3	Solving as exact ode	1132
8.8.4	Maple step by step solution	1136

Internal problem ID [6256]

Internal file name [OUTPUT/5504_Sunday_June_05_2022_03_41_47_PM_95497302/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Problems for Review and Discovery. Page 53

Problem number: 1(h).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "separable", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_separable]

$$-\sin(x)\sin(y) + \cos(x)\cos(y)y' = 0$$

8.8.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{\sin(x)\tan(y)}{\cos(x)}\end{aligned}$$

Where $f(x) = \frac{\sin(x)}{\cos(x)}$ and $g(y) = \tan(y)$. Integrating both sides gives

$$\frac{1}{\tan(y)} dy = \frac{\sin(x)}{\cos(x)} dx$$

$$\int \frac{1}{\tan(y)} dy = \int \frac{\sin(x)}{\cos(x)} dx$$

$$\ln(\sin(y)) = -\ln(\cos(x)) + c_1$$

Raising both side to exponential gives

$$\sin(y) = e^{-\ln(\cos(x))+c_1}$$

Which simplifies to

$$\sin(y) = \frac{c_2}{\cos(x)}$$

Summary

The solution(s) found are the following

$$y = \arcsin\left(\frac{c_2 e^{c_1}}{\cos(x)}\right) \tag{1}$$




Figure 225: Slope field plot

Verification of solutions

$$y = \arcsin\left(\frac{c_2 e^{c_1}}{\cos(x)}\right)$$

Verified OK.

8.8.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{\sin(x) \sin(y)}{\cos(x) \cos(y)}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **separable**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 150: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= \frac{\cos(x)}{\sin(x)} \\ \eta(x, y) &= 0\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\eta = 0$ then in this special case

$$R = y$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\xi} dx \\ &= \int \frac{1}{\frac{\cos(x)}{\sin(x)}} dx\end{aligned}$$

Which results in

$$S = -\ln(\cos(x))$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y}\tag{2}$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{\sin(x) \sin(y)}{\cos(x) \cos(y)}$$

Evaluating all the partial derivatives gives

$$\begin{aligned}R_x &= 0 \\R_y &= 1 \\S_x &= \tan(x) \\S_y &= 0\end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \cot(y) \tag{2A}$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \cot(R)$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \ln(\sin(R)) + c_1 \tag{4}$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$-\ln(\cos(x)) = \ln(\sin(y)) + c_1$$



Which simplifies to

$$-\ln(\cos(x)) = \ln(\sin(y)) + c_1$$

Which gives

$$y = \arcsin\left(\frac{e^{-c_1}}{\cos(x)}\right)$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{\sin(x) \sin(y)}{\cos(x) \cos(y)}$ 	$R = y$ $S = -\ln(\cos(x))$	$\frac{dS}{dR} = \cot(R)$ 

Summary

The solution(s) found are the following

$$y = \arcsin\left(\frac{e^{-c_1}}{\cos(x)}\right) \tag{1}$$




Figure 226: Slope field plot

Verification of solutions

$$y = \arcsin\left(\frac{e^{-c_1}}{\cos(x)}\right)$$

Verified OK.

8.8.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial\phi}{\partial x} + \frac{\partial\phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}\left(\frac{\cos(y)}{\sin(y)}\right) dy &= \left(\frac{\sin(x)}{\cos(x)}\right) dx \\ \left(-\frac{\sin(x)}{\cos(x)}\right) dx + \left(\frac{\cos(y)}{\sin(y)}\right) dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -\frac{\sin(x)}{\cos(x)} \\ N(x, y) &= \frac{\cos(y)}{\sin(y)}\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(-\frac{\sin(x)}{\cos(x)}\right) \\ &= 0\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{\cos(y)}{\sin(y)} \right) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\frac{\sin(x)}{\cos(x)} dx \\ \phi &= \ln(\cos(x)) + f(y)\end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{\cos(y)}{\sin(y)}$. Therefore equation (4) becomes

$$\frac{\cos(y)}{\sin(y)} = 0 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$\begin{aligned}f'(y) &= \frac{\cos(y)}{\sin(y)} \\ &= \cot(y)\end{aligned}$$

Integrating the above w.r.t y results in

$$\int f'(y) dy = \int (\cot(y)) dy$$
$$f(y) = \ln(\sin(y)) + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = \ln(\cos(x)) + \ln(\sin(y)) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \ln(\cos(x)) + \ln(\sin(y))$$

Summary

The solution(s) found are the following

$$\ln(\cos(x)) + \ln(\sin(y)) = c_1 \tag{1}$$




Figure 227: Slope field plot

Verification of solutions

$$\ln(\cos(x)) + \ln(\sin(y)) = c_1$$

Verified OK.

8.8.4 Maple step by step solution

Let's solve

$$-\sin(x)\sin(y) + \cos(x)\cos(y)y' = 0$$

- Highest derivative means the order of the ODE is 1

y'

- Integrate both sides with respect to x

$$\int (-\sin(x)\sin(y) + \cos(x)\cos(y)y') dx = \int 0 dx + c_1$$

- Evaluate integral

$$\cos(x)\sin(y) = c_1$$

- Solve for y

$$y = \arcsin\left(\frac{c_1}{\cos(x)}\right)$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
<- separable successful`
```

✓ Solution by Maple

Time used: 0.14 (sec). Leaf size: 9

```
dsolve(-sin(x)*sin(y(x))+cos(x)*cos(y(x))*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \arcsin(\sec(x) c_1)$$

✓ Solution by Mathematica

Time used: 3.473 (sec). Leaf size: 19

```
DSolve[-Sin[x]*Sin[y[x]]+Cos[x]*Cos[y[x]]*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \arcsin\left(\frac{1}{2}c_1 \sec(x)\right)$$

$$y(x) \rightarrow 0$$

8.9 problem 2(a)

8.9.1	Existence and uniqueness analysis	1139
8.9.2	Solving as linear ode	1139
8.9.3	Solving as homogeneousTypeD2 ode	1141
8.9.4	Solving as first order ode lie symmetry lookup ode	1142
8.9.5	Solving as exact ode	1146
8.9.6	Maple step by step solution	1151

Internal problem ID [6257]

Internal file name [OUTPUT/5505_Sunday_June_05_2022_03_41_49_PM_16109693/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Problems for Review and Discovery. Page 53

Problem number: 2(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"linear"**, **"homogeneousTypeD2"**, **"exactWithIntegrationFactor"**, **"first_order_ode_lie_symmetry_lookup"**

Maple gives the following as the ode type

`[_linear]`

$$-y + xy' = 2x$$

With initial conditions

$$[y(1) = 0]$$

8.9.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -\frac{1}{x}$$

$$q(x) = 2$$

Hence the ode is

$$y' - \frac{y}{x} = 2$$

The domain of $p(x) = -\frac{1}{x}$ is

$$\{x < 0 \vee 0 < x\}$$

And the point $x_0 = 1$ is inside this domain. The domain of $q(x) = 2$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 1$ is also inside this domain. Hence solution exists and is unique.

8.9.2 Solving as linear ode

Entering Linear first order ODE solver. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int -\frac{1}{x} dx} \\ &= \frac{1}{x}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu)(2) \\ \frac{d}{dx}\left(\frac{y}{x}\right) &= \left(\frac{1}{x}\right)(2) \\ d\left(\frac{y}{x}\right) &= \left(\frac{2}{x}\right) dx\end{aligned}$$

Integrating gives

$$\frac{y}{x} = \int \frac{2}{x} dx$$
$$\frac{y}{x} = 2 \ln(x) + c_1$$

Dividing both sides by the integrating factor $\mu = \frac{1}{x}$ results in

$$y = 2 \ln(x) x + c_1 x$$

which simplifies to

$$y = x(2 \ln(x) + c_1)$$

Initial conditions are used to solve for c_1 . Substituting $x = 1$ and $y = 0$ in the above solution gives an equation to solve for the constant of integration.

$$0 = c_1$$

$$c_1 = 0$$


Substituting c_1 found above in the general solution gives

$$y = 2 \ln(x) x$$


Summary

The solution(s) found are the following

$$y = 2 \ln(x) x \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = 2 \ln(x) x$$

Verified OK.

8.9.3 Solving as homogeneous Type D2 ode

Using the change of variables $y = u(x) x$ on the above ode results in new ode in $u(x)$

$$-u(x) x + x(u'(x) x + u(x)) = 2x$$

Integrating both sides gives

$$\begin{aligned} u(x) &= \int \frac{2}{x} dx \\ &= 2 \ln(x) + c_2 \end{aligned}$$

Therefore the solution y is

$$\begin{aligned} y &= ux \\ &= x(2 \ln(x) + c_2) \end{aligned}$$

Initial conditions are used to solve for c_2 . Substituting $x = 1$ and $y = 0$ in the above solution gives an equation to solve for the constant of integration.

$$0 = c_2$$

$$c_2 = 0$$


Substituting c_2 found above in the general solution gives

$$y = 2 \ln(x) x$$


Summary

The solution(s) found are the following

$$y = 2 \ln(x) x \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = 2 \ln(x) x$$

Verified OK.

8.9.4 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{2x + y}{x}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \tag{A}$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 153: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= x\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{x} dy \end{aligned}$$

Which results in

$$S = \frac{y}{x}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{2x + y}{x}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= -\frac{y}{x^2} \\ S_y &= \frac{1}{x} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \frac{2}{x} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \frac{2}{R}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = 2 \ln(R) + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{y}{x} = 2 \ln(x) + c_1$$



Which simplifies to

$$\frac{y}{x} = 2 \ln(x) + c_1$$

Which gives

$$y = x(2 \ln(x) + c_1)$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{2x+y}{x}$ 	$R = x$ $S = \frac{y}{x}$	$\frac{dS}{dR} = \frac{2}{R}$ 

Initial conditions are used to solve for c_1 . Substituting $x = 1$ and $y = 0$ in the above solution gives an equation to solve for the constant of integration.

$$0 = c_1$$

$$c_1 = 0$$


Substituting c_1 found above in the general solution gives

$$y = 2 \ln(x) x$$


Summary

The solution(s) found are the following

$$y = 2 \ln(x) x \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = 2 \ln(x) x$$

Verified OK.

8.9.5 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} (x) dy &= (2x + y) dx \\ (-2x - y) dx + (x) dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -2x - y \\ N(x, y) &= x \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-2x - y) \\ &= -1 \end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(x) \\ &= 1\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned}A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= \frac{1}{x} ((-1) - (1)) \\ &= -\frac{2}{x}\end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int A \, dx} \\ &= e^{\int -\frac{2}{x} \, dx}\end{aligned}$$

The result of integrating gives

$$\begin{aligned}\mu &= e^{-2 \ln(x)} \\ &= \frac{1}{x^2}\end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} for now so not to confuse them with the original M and N .

$$\begin{aligned}\bar{M} &= \mu M \\ &= \frac{1}{x^2}(-2x - y) \\ &= \frac{-2x - y}{x^2}\end{aligned}$$

And

$$\begin{aligned}\bar{N} &= \mu N \\ &= \frac{1}{x^2}(x) \\ &= \frac{1}{x}\end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned}\overline{M} + \overline{N} \frac{dy}{dx} &= 0 \\ \left(\frac{-2x - y}{x^2} \right) + \left(\frac{1}{x} \right) \frac{dy}{dx} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \overline{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \overline{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \overline{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int \frac{-2x - y}{x^2} dx \\ \phi &= -2 \ln(x) + \frac{y}{x} + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = \frac{1}{x} + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{1}{x}$. Therefore equation (4) becomes

$$\frac{1}{x} = \frac{1}{x} + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = -2 \ln(x) + \frac{y}{x} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -2 \ln(x) + \frac{y}{x}$$

The solution becomes

$$y = x(2 \ln(x) + c_1)$$

Initial conditions are used to solve for c_1 . Substituting $x = 1$ and $y = 0$ in the above solution gives an equation to solve for the constant of integration.

$$0 = c_1$$

$$c_1 = 0$$


Substituting c_1 found above in the general solution gives

$$y = 2 \ln(x) x$$


Summary

The solution(s) found are the following

$$y = 2 \ln(x) x \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = 2 \ln(x) x$$

Verified OK.

8.9.6 Maple step by step solution

Let's solve

$$[-y + xy' = 2x, y(1) = 0]$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = 2 + \frac{y}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' - \frac{y}{x} = 2$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) \left(y' - \frac{y}{x} \right) = 2\mu(x)$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x)y)$

$$\mu(x) \left(y' - \frac{y}{x} \right) = \mu'(x)y + \mu(x)y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = -\frac{\mu(x)}{x}$$

- Solve to find the integrating factor

$$\mu(x) = \frac{1}{x}$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x)y) \right) dx = \int 2\mu(x) dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x)y = \int 2\mu(x) dx + c_1$$

- Solve for y

$$y = \frac{\int 2\mu(x)dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = \frac{1}{x}$

$$y = x \left(\int \frac{2}{x} dx + c_1 \right)$$

- Evaluate the integrals on the rhs
 $y = x(2 \ln(x) + c_1)$
- Use initial condition $y(1) = 0$
 $0 = c_1$
- Solve for c_1
 $c_1 = 0$
- Substitute $c_1 = 0$ into general solution and simplify
 $y = 2 \ln(x) x$
- Solution to the IVP
 $y = 2 \ln(x) x$

Maple trace

```
`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
dsolve([x*diff(y(x),x)-y(x)=2*x,y(1) = 0],y(x), singsol=all)
```

$$y(x) = 2 \ln(x) x$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 10

```
DSolve[{x*y'[x]-y[x]==2*x,{y[1]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2x \log(x)$$

8.10 problem 2(b)

8.10.1 Existence and uniqueness analysis	1153
8.10.2 Solving as linear ode	1154
8.10.3 Solving as first order ode lie symmetry lookup ode	1156
8.10.4 Solving as exact ode	1160
8.10.5 Maple step by step solution	1165

Internal problem ID [6258]

Internal file name [OUTPUT/5506_Sunday_June_05_2022_03_41_50_PM_37330925/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Problems for Review and Discovery. Page 53

Problem number: 2(b).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"linear"**, **"exactWithIntegrationFactor"**, **"first_order_ode_lie_symmetry_lookup"**

Maple gives the following as the ode type

[_linear]

$$x^2y' - 2y = 3x^2$$

With initial conditions

$$[y(1) = 2]$$

8.10.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -\frac{2}{x^2}$$

$$q(x) = 3$$

Hence the ode is

$$y' - \frac{2y}{x^2} = 3$$

The domain of $p(x) = -\frac{2}{x^2}$ is

$$\{x < 0 \vee 0 < x\}$$

And the point $x_0 = 1$ is inside this domain. The domain of $q(x) = 3$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 1$ is also inside this domain. Hence solution exists and is unique.

8.10.2 Solving as linear ode

Entering Linear first order ODE solver. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int -\frac{2}{x^2} dx} \\ &= e^{\frac{2}{x}}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu)(3) \\ \frac{d}{dx}\left(e^{\frac{2}{x}}y\right) &= \left(e^{\frac{2}{x}}\right)(3) \\ d\left(e^{\frac{2}{x}}y\right) &= \left(3e^{\frac{2}{x}}\right) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}e^{\frac{2}{x}}y &= \int 3e^{\frac{2}{x}} dx \\ e^{\frac{2}{x}}y &= 3x e^{\frac{2}{x}} + 6 \expIntegral_1\left(-\frac{2}{x}\right) + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{\frac{2}{x}}$ results in

$$y = e^{-\frac{2}{x}}\left(3x e^{\frac{2}{x}} + 6 \expIntegral_1\left(-\frac{2}{x}\right)\right) + c_1 e^{-\frac{2}{x}}$$

which simplifies to

$$y = 3x + 6 \expIntegral_1\left(-\frac{2}{x}\right) e^{-\frac{2}{x}} + c_1 e^{-\frac{2}{x}}$$

Initial conditions are used to solve for c_1 . Substituting $x = 1$ and $y = 2$ in the above solution gives an equation to solve for the constant of integration.

$$2 = 3 + 6 \expIntegral_1(-2) e^{-2} + c_1 e^{-2}$$

$$c_1 = -(6 \expIntegral_1(-2) e^{-2} + 1) e^2$$


Substituting c_1 found above in the general solution gives

$$y = -6 e^{-\frac{2}{x}} \expIntegral_1(-2) + 6 \expIntegral_1\left(-\frac{2}{x}\right) e^{-\frac{2}{x}} - e^{\frac{2x-2}{x}} + 3x$$


Summary

The solution(s) found are the following

$$y = -6 e^{-\frac{2}{x}} \expIntegral_1(-2) + 6 \expIntegral_1\left(-\frac{2}{x}\right) e^{-\frac{2}{x}} - e^{\frac{2x-2}{x}} + 3x \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = -6 e^{-\frac{2}{x}} \expIntegral_1(-2) + 6 \expIntegral_1\left(-\frac{2}{x}\right) e^{-\frac{2}{x}} - e^{\frac{2x-2}{x}} + 3x$$

Verified OK.

8.10.3 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{3x^2 + 2y}{x^2}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 156: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= e^{-\frac{2}{x}}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^{-\frac{2}{x}}} dy\end{aligned}$$

Which results in

$$S = e^{\frac{2}{x}} y$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y}\tag{2}$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{3x^2 + 2y}{x^2}$$

Evaluating all the partial derivatives gives

$$\begin{aligned}R_x &= 1 \\R_y &= 0 \\S_x &= -\frac{2 e^{\frac{2}{x}} y}{x^2} \\S_y &= e^{\frac{2}{x}}\end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 3 e^{\frac{2}{x}} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 3 e^{\frac{2}{R}}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = 3R e^{\frac{2}{R}} + 6 \expIntegral_1 \left(-\frac{2}{R} \right) + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$e^{\frac{2}{x}} y = 3x e^{\frac{2}{x}} + 6 \expIntegral_1 \left(-\frac{2}{x} \right) + c_1$$



Which simplifies to

$$-6 \expIntegral_1 \left(-\frac{2}{x} \right) + (-3x + y) e^{\frac{2}{x}} - c_1 = 0$$

Which gives

$$y = \left(3x e^{\frac{2}{x}} + 6 \expIntegral_1 \left(-\frac{2}{x} \right) + c_1 \right) e^{-\frac{2}{x}}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{3x^2 + 2y}{x^2}$ 	$R = x$ $S = e^{\frac{2}{x}} y$	$\frac{dS}{dR} = 3 e^{\frac{2}{R}}$ 

Initial conditions are used to solve for c_1 . Substituting $x = 1$ and $y = 2$ in the above solution gives an equation to solve for the constant of integration.

$$2 = 3 + 6 \exp\text{Integral}_1(-2) e^{-2} + c_1 e^{-2}$$

$$c_1 = -(6 \exp\text{Integral}_1(-2) e^{-2} + 1) e^2$$


Substituting c_1 found above in the general solution gives

$$y = -6 e^{-\frac{2}{x}} \exp\text{Integral}_1(-2) + 6 \exp\text{Integral}_1\left(-\frac{2}{x}\right) e^{-\frac{2}{x}} - e^{\frac{2x-2}{x}} + 3x$$


Summary

The solution(s) found are the following

$$y = -6 e^{-\frac{2}{x}} \exp\text{Integral}_1(-2) + 6 \exp\text{Integral}_1\left(-\frac{2}{x}\right) e^{-\frac{2}{x}} - e^{\frac{2x-2}{x}} + 3x \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = -6 e^{-\frac{2}{x}} \expIntegral_1(-2) + 6 \expIntegral_1\left(-\frac{2}{x}\right) e^{-\frac{2}{x}} - e^{\frac{2x-2}{x}} + 3x$$

Verified OK.

8.10.4 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned} (x^2) dy &= (3x^2 + 2y) dx \\ (-3x^2 - 2y) dx + (x^2) dy &= 0 \end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -3x^2 - 2y \\ N(x, y) &= x^2 \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} (-3x^2 - 2y) \\ &= -2 \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} (x^2) \\ &= 2x \end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= \frac{1}{x^2} ((-2) - (2x)) \\ &= \frac{-2 - 2x}{x^2} \end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int A \, dx} \\ &= e^{\int \frac{-2-2x}{x^2} \, dx}\end{aligned}$$

The result of integrating gives

$$\begin{aligned}\mu &= e^{-2\ln(x) + \frac{2}{x}} \\ &= \frac{e^{\frac{2}{x}}}{x^2}\end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} for now so not to confuse them with the original M and N .

$$\begin{aligned}\bar{M} &= \mu M \\ &= \frac{e^{\frac{2}{x}}}{x^2}(-3x^2 - 2y) \\ &= \frac{(-3x^2 - 2y) e^{\frac{2}{x}}}{x^2}\end{aligned}$$

And

$$\begin{aligned}\bar{N} &= \mu N \\ &= \frac{e^{\frac{2}{x}}}{x^2}(x^2) \\ &= e^{\frac{2}{x}}\end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned}\bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ \left(\frac{(-3x^2 - 2y) e^{\frac{2}{x}}}{x^2} \right) + \left(e^{\frac{2}{x}} \right) \frac{dy}{dx} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\int \frac{\partial \phi}{\partial x} dx = \int \bar{M} dx$$

$$\int \frac{\partial \phi}{\partial x} dx = \int \frac{(-3x^2 - 2y) e^{\frac{2}{x}}}{x^2} dx$$

$$\phi = -6 \expIntegral_1 \left(-\frac{2}{x} \right) + (-3x + y) e^{\frac{2}{x}} + f(y) \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = e^{\frac{2}{x}} + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = e^{\frac{2}{x}}$. Therefore equation (4) becomes

$$e^{\frac{2}{x}} = e^{\frac{2}{x}} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = -6 \expIntegral_1 \left(-\frac{2}{x} \right) + (-3x + y) e^{\frac{2}{x}} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -6 \expIntegral_1 \left(-\frac{2}{x} \right) + (-3x + y) e^{\frac{2}{x}}$$

The solution becomes

$$y = \left(3x e^{\frac{2}{x}} + 6 \expIntegral_1 \left(-\frac{2}{x} \right) + c_1 \right) e^{-\frac{2}{x}}$$

Initial conditions are used to solve for c_1 . Substituting $x = 1$ and $y = 2$ in the above solution gives an equation to solve for the constant of integration.

$$2 = 3 + 6 \expIntegral_1(-2) e^{-2} + c_1 e^{-2}$$

$$c_1 = -(6 \expIntegral_1(-2) e^{-2} + 1) e^2$$


Substituting c_1 found above in the general solution gives

$$y = -6 e^{-\frac{2}{x}} \expIntegral_1(-2) + 6 \expIntegral_1\left(-\frac{2}{x}\right) e^{-\frac{2}{x}} - e^{\frac{2x-2}{x}} + 3x$$


Summary

The solution(s) found are the following

$$y = -6 e^{-\frac{2}{x}} \expIntegral_1(-2) + 6 \expIntegral_1\left(-\frac{2}{x}\right) e^{-\frac{2}{x}} - e^{\frac{2x-2}{x}} + 3x \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = -6 e^{-\frac{2}{x}} \expIntegral_1(-2) + 6 \expIntegral_1\left(-\frac{2}{x}\right) e^{-\frac{2}{x}} - e^{\frac{2x-2}{x}} + 3x$$

Verified OK.

8.10.5 Maple step by step solution

Let's solve

$$[x^2y' - 2y = 3x^2, y(1) = 2]$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = 3 + \frac{2y}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' - \frac{2y}{x^2} = 3$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) \left(y' - \frac{2y}{x^2} \right) = 3\mu(x)$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x)y)$

$$\mu(x) \left(y' - \frac{2y}{x^2} \right) = \mu'(x)y + \mu(x)y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = -\frac{2\mu(x)}{x^2}$$

- Solve to find the integrating factor

$$\mu(x) = e^{\frac{2}{x}}$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x)y) \right) dx = \int 3\mu(x) dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x)y = \int 3\mu(x) dx + c_1$$

- Solve for y

$$y = \frac{\int 3\mu(x)dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = e^{\frac{2}{x}}$

$$y = \frac{\int 3e^{\frac{2}{x}}dx + c_1}{e^{\frac{2}{x}}}$$

- Evaluate the integrals on the rhs

$$y = \frac{3xe^{\frac{2}{x}} + 6\text{Ei}_1\left(-\frac{2}{x}\right) + c_1}{e^{\frac{2}{x}}}$$

- Simplify

$$y = 3x + 6 \operatorname{Ei}_1\left(-\frac{2}{x}\right) e^{-\frac{2}{x}} + c_1 e^{-\frac{2}{x}}$$

- Use initial condition $y(1) = 2$

$$2 = 3 + 6 \operatorname{Ei}_1(-2) e^{-2} + c_1 e^{-2}$$

- Solve for c_1

$$c_1 = -\frac{6 \operatorname{Ei}_1(-2) e^{-2} + 1}{e^{-2}}$$

- Substitute $c_1 = -\frac{6 \operatorname{Ei}_1(-2) e^{-2} + 1}{e^{-2}}$ into general solution and simplify

$$y = -6 e^{-\frac{2}{x}} \operatorname{Ei}_1(-2) + 6 \operatorname{Ei}_1\left(-\frac{2}{x}\right) e^{-\frac{2}{x}} - e^{\frac{2x-2}{x}} + 3x$$

- Solution to the IVP

$$y = -6 e^{-\frac{2}{x}} \operatorname{Ei}_1(-2) + 6 \operatorname{Ei}_1\left(-\frac{2}{x}\right) e^{-\frac{2}{x}} - e^{\frac{2x-2}{x}} + 3x$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.218 (sec). Leaf size: 39

```
dsolve([x^2*diff(y(x),x)-2*y(x)=3*x^2,y(1) = 2],y(x), singsol=all)
```

$$y(x) = 3x - e^{2-\frac{2}{x}} + 6 \operatorname{expIntegral}_1\left(-\frac{2}{x}\right) e^{-\frac{2}{x}} - 6 \operatorname{expIntegral}_1(-2) e^{-\frac{2}{x}}$$

✓ Solution by Mathematica

Time used: 0.039 (sec). Leaf size: 41

```
DSolve[{x^2*y'[x]-2*y[x]==3*x^2,{y[1]==2}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2/x} \left(-6 \operatorname{ExpIntegralEi}\left(\frac{2}{x}\right) + 6 \operatorname{ExpIntegralEi}(2) + 3e^{2/x}x - e^2 \right)$$

8.11 problem 2(c)

8.11.1 Existence and uniqueness analysis	1168
8.11.2 Solving as separable ode	1168
8.11.3 Solving as differentialType ode	1170
8.11.4 Solving as first order ode lie symmetry lookup ode	1174
8.11.5 Solving as exact ode	1178
8.11.6 Maple step by step solution	1181

Internal problem ID [6259]

Internal file name [OUTPUT/5507_Sunday_June_05_2022_03_41_59_PM_92155662/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Problems for Review and Discovery. Page 53

Problem number: 2(c).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "separable", "differential-Type", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

`[_separable]`

$$y'y^2 = x$$

With initial conditions

$$[y(-1) = 3]$$

8.11.1 Existence and uniqueness analysis

This is non linear first order ODE. In canonical form it is written as

$$\begin{aligned}y' &= f(x, y) \\ &= \frac{x}{y^2}\end{aligned}$$

The x domain of $f(x, y)$ when $y = 3$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = -1$ is inside this domain. The y domain of $f(x, y)$ when $x = -1$ is

$$\{y < 0 \vee 0 < y\}$$

And the point $y_0 = 3$ is inside this domain. Now we will look at the continuity of

$$\begin{aligned}\frac{\partial f}{\partial y} &= \frac{\partial}{\partial y} \left(\frac{x}{y^2} \right) \\ &= -\frac{2x}{y^3}\end{aligned}$$

The x domain of $\frac{\partial f}{\partial y}$ when $y = 3$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = -1$ is inside this domain. The y domain of $\frac{\partial f}{\partial y}$ when $x = -1$ is

$$\{y < 0 \vee 0 < y\}$$

And the point $y_0 = 3$ is inside this domain. Therefore solution exists and is unique.

8.11.2 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{x}{y^2}\end{aligned}$$

Where $f(x) = x$ and $g(y) = \frac{1}{y^2}$. Integrating both sides gives

$$\begin{aligned}\frac{1}{y^2} dy &= x dx \\ \int \frac{1}{y^2} dy &= \int x dx \\ \frac{y^3}{3} &= \frac{x^2}{2} + c_1\end{aligned}$$

Which results in

$$\begin{aligned}y &= \frac{(12x^2 + 24c_1)^{\frac{1}{3}}}{2} \\ y &= -\frac{(12x^2 + 24c_1)^{\frac{1}{3}}}{4} + \frac{i\sqrt{3}(12x^2 + 24c_1)^{\frac{1}{3}}}{4} \\ y &= -\frac{(12x^2 + 24c_1)^{\frac{1}{3}}}{4} - \frac{i\sqrt{3}(12x^2 + 24c_1)^{\frac{1}{3}}}{4}\end{aligned}$$

Initial conditions are used to solve for c_1 . Substituting $x = -1$ and $y = 3$ in the above solution gives an equation to solve for the constant of integration.

$$3 = -\frac{i(12 + 24c_1)^{\frac{1}{3}} \sqrt{3}}{4} - \frac{(12 + 24c_1)^{\frac{1}{3}}}{4}$$

Warning: Unable to solve for constant of integration. Initial conditions are used to solve for c_1 . Substituting $x = -1$ and $y = 3$ in the above solution gives an equation to solve for the constant of integration.

$$3 = \frac{i(12 + 24c_1)^{\frac{1}{3}} \sqrt{3}}{4} - \frac{(12 + 24c_1)^{\frac{1}{3}}}{4}$$

Warning: Unable to solve for constant of integration. Initial conditions are used to solve for c_1 . Substituting $x = -1$ and $y = 3$ in the above solution gives an equation to solve for the constant of integration.

$$3 = \frac{(12 + 24c_1)^{\frac{1}{3}}}{2}$$

$$c_1 = \frac{17}{2}$$


Substituting c_1 found above in the general solution gives

$$y = \frac{(12x^2 + 204)^{\frac{1}{3}}}{2}$$


Summary

The solution(s) found are the following

$$y = \frac{(12x^2 + 204)^{\frac{1}{3}}}{2} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{(12x^2 + 204)^{\frac{1}{3}}}{2}$$

Verified OK.

8.11.3 Solving as differential Type ode

Writing the ode as

$$y' = \frac{x}{y^2} \quad (1)$$

Which becomes

$$(y^2) dy = (x) dx \quad (2)$$

But the RHS is complete differential because

$$(x) dx = d\left(\frac{x^2}{2}\right)$$

Hence (2) becomes

$$(y^2) dy = d\left(\frac{y^3}{3}\right)$$

Integrating both sides gives gives these solutions

$$y = \frac{(12x^2 + 24c_1)^{\frac{1}{3}}}{2} + c_1$$

$$y = -\frac{(12x^2 + 24c_1)^{\frac{1}{3}}}{4} + \frac{i\sqrt{3}(12x^2 + 24c_1)^{\frac{1}{3}}}{4} + c_1$$

$$y = -\frac{(12x^2 + 24c_1)^{\frac{1}{3}}}{4} - \frac{i\sqrt{3}(12x^2 + 24c_1)^{\frac{1}{3}}}{4} + c_1$$

Initial conditions are used to solve for c_1 . Substituting $x = -1$ and $y = 3$ in the above solution gives an equation to solve for the constant of integration.

$$3 = -\frac{(12 + 24c_1)^{\frac{1}{3}}}{4} - \frac{i(12 + 24c_1)^{\frac{1}{3}} \sqrt{3}}{4} + c_1$$

$$c_1 = \frac{i\sqrt{3} \left((42 + 2\sqrt{457})^{\frac{1}{3}} - \frac{-2i\sqrt{3}-2}{(42+2\sqrt{457})^{\frac{1}{3}}} \right)}{4} + \frac{(42 + 2\sqrt{457})^{\frac{1}{3}}}{4} - \frac{-2i\sqrt{3} - 2}{4(42 + 2\sqrt{457})^{\frac{1}{3}}} + 3$$

Substituting c_1 found above in the general solution gives

$$y = \frac{-4(42 + 2\sqrt{457})^{\frac{1}{3}} - i\sqrt{3} \left(i\sqrt{3} (42 + 2\sqrt{457})^{\frac{2}{3}} + 2x^2(42 + 2\sqrt{457})^{\frac{1}{3}} + 4i\sqrt{3} + (42 + 2\sqrt{457})^{\frac{2}{3}} + 12 \right)}{4}$$

Initial conditions are used to solve for c_1 . Substituting $x = -1$ and $y = 3$ in the above solution gives an equation to solve for the constant of integration.

$$3 = -\frac{(12 + 24c_1)^{\frac{1}{3}}}{4} + \frac{i(12 + 24c_1)^{\frac{1}{3}} \sqrt{3}}{4} + c_1$$

$$c_1 = -\frac{i\sqrt{3} \left((42 + 2\sqrt{457})^{\frac{1}{3}} - \frac{2i\sqrt{3}-2}{(42+2\sqrt{457})^{\frac{1}{3}}} \right)}{4} + \frac{(42 + 2\sqrt{457})^{\frac{1}{3}}}{4} - \frac{2i\sqrt{3} - 2}{4(42 + 2\sqrt{457})^{\frac{1}{3}}} + 3$$

Substituting c_1 found above in the general solution gives

$$y = \frac{-i\sqrt{3} (42 + 2\sqrt{457})^{\frac{7}{9}} + i\sqrt{3} \left(-6i\sqrt{3} (42 + 2\sqrt{457})^{\frac{2}{3}} + 12x^2(42 + 2\sqrt{457})^{\frac{1}{3}} - 24i\sqrt{3} + 6(42 + 2\sqrt{457})^{\frac{1}{3}} \right)}{2(42 + 2\sqrt{457})^{\frac{4}{9}}}$$

Initial conditions are used to solve for c_1 . Substituting $x = -1$ and $y = 3$ in the above solution gives an equation to solve for the constant of integration.

$$3 = \frac{(12 + 24c_1)^{\frac{1}{3}}}{2} + c_1$$

$$c_1 = -\frac{(42 + 2\sqrt{457})^{\frac{1}{3}}}{2} + \frac{2}{(42 + 2\sqrt{457})^{\frac{1}{3}}} + 3$$

Substituting c_1 found above in the general solution gives

$$y = \frac{-(42 + 2\sqrt{457})^{\frac{7}{9}} + \left(12x^2(42 + 2\sqrt{457})^{\frac{1}{3}} - 12(42 + 2\sqrt{457})^{\frac{2}{3}} + 72(42 + 2\sqrt{457})^{\frac{1}{3}} + 48 \right)^{\frac{1}{3}} (42 + 2\sqrt{457})^{\frac{1}{3}}}{2(42 + 2\sqrt{457})^{\frac{4}{9}}}$$


Summary

The solution(s) found are the following


$$y = \frac{-(42 + 2\sqrt{457})^{\frac{7}{9}} + \left(12x^2(42 + 2\sqrt{457})^{\frac{1}{3}} - 12(42 + 2\sqrt{457})^{\frac{2}{3}} + 72(42 + 2\sqrt{457})^{\frac{1}{3}} + 48 \right)^{\frac{1}{3}} (42 + 2\sqrt{457})^{\frac{1}{3}}}{2(42 + 2\sqrt{457})^{\frac{4}{9}}} \quad (1)$$

$$y = \frac{-i\sqrt{3} (42 + 2\sqrt{457})^{\frac{7}{9}} + i\sqrt{3} \left(-6i\sqrt{3} (42 + 2\sqrt{457})^{\frac{2}{3}} + 12x^2(42 + 2\sqrt{457})^{\frac{1}{3}} - 24i\sqrt{3} + 6(42 + 2\sqrt{457})^{\frac{1}{3}} \right)}{2(42 + 2\sqrt{457})^{\frac{4}{9}}} \quad (2)$$

$$y = \frac{-4(42 + 2\sqrt{457})^{\frac{1}{9}} - i\sqrt{3} \left(i\sqrt{3} (42 + 2\sqrt{457})^{\frac{2}{3}} + 2x^2(42 + 2\sqrt{457})^{\frac{1}{3}} + 4i\sqrt{3} + (42 + 2\sqrt{457})^{\frac{2}{3}} + 12(42 + 2\sqrt{457})^{\frac{1}{3}} \right)}{2(42 + 2\sqrt{457})^{\frac{4}{9}}} \quad (3)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

y

$$= \frac{-(42 + 2\sqrt{457})^{\frac{7}{9}} + \left(12x^2(42 + 2\sqrt{457})^{\frac{1}{3}} - 12(42 + 2\sqrt{457})^{\frac{2}{3}} + 72(42 + 2\sqrt{457})^{\frac{1}{3}} + 48\right)^{\frac{1}{3}} (42 + 2\sqrt{457})}{2(42 + 2\sqrt{457})^{\frac{4}{9}}}$$

Verified OK. {positive}

y

$$= \frac{-i\sqrt{3}(42 + 2\sqrt{457})^{\frac{7}{9}} + i\sqrt{3}\left(-6i\sqrt{3}(42 + 2\sqrt{457})^{\frac{2}{3}} + 12x^2(42 + 2\sqrt{457})^{\frac{1}{3}} - 24i\sqrt{3} + 6(42 + 2\sqrt{457})\right)}{2(42 + 2\sqrt{457})^{\frac{4}{9}}}$$

Verified OK. {positive}

y

$$= \frac{-4(42 + 2\sqrt{457})^{\frac{1}{9}} - i\sqrt{3}\left(i\sqrt{3}(42 + 2\sqrt{457})^{\frac{2}{3}} + 2x^2(42 + 2\sqrt{457})^{\frac{1}{3}} + 4i\sqrt{3} + (42 + 2\sqrt{457})^{\frac{2}{3}} + 12(42 + 2\sqrt{457})\right)}{2(42 + 2\sqrt{457})^{\frac{4}{9}}}$$

Verified OK. {positive}

8.11.4 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{x}{y^2}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **separable**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 159: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= \frac{1}{x} \\ \eta(x, y) &= 0\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\eta = 0$ then in this special case

$$R = y$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\xi} dx \\ &= \int \frac{1}{\frac{1}{x}} dx\end{aligned}$$

Which results in

$$S = \frac{x^2}{2}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y}\tag{2}$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{x}{y^2}$$

Evaluating all the partial derivatives gives

$$R_x = 0$$

$$R_y = 1$$

$$S_x = x$$

$$S_y = 0$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = y^2 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = R^2$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \frac{R^3}{3} + c_1 \quad (4)$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{x^2}{2} = \frac{y^3}{3} + c_1$$

Which simplifies to

$$\frac{x^2}{2} = \frac{y^3}{3} + c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{x}{y^2}$ 	$R = y$ $S = \frac{x^2}{2}$	$\frac{dS}{dR} = R^2$ 

Initial conditions are used to solve for c_1 . Substituting $x = -1$ and $y = 3$ in the above solution gives an equation to solve for the constant of integration.

$$\frac{1}{2} = 9 + c_1$$

$$c_1 = -\frac{17}{2}$$

Substituting c_1 found above in the general solution gives

$$\frac{x^2}{2} = \frac{y^3}{3} - \frac{17}{2}$$

Summary

The solution(s) found are the following

$$\frac{x^2}{2} = \frac{y^3}{3} - \frac{17}{2} \tag{1}$$

Verification of solutions

$$\frac{x^2}{2} = \frac{y^3}{3} - \frac{17}{2}$$

Verified OK.

8.11.5 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial\phi}{\partial x} + \frac{\partial\phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} (y^2) dy &= (x) dx \\ (-x) dx + (y^2) dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -x \\ N(x, y) &= y^2 \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-x) \\ &= 0\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(y^2) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \tag{1}$$

$$\frac{\partial \phi}{\partial y} = N \tag{2}$$

Integrating (1) w.r.t. x gives

$$\int \frac{\partial \phi}{\partial x} dx = \int M dx$$

$$\int \frac{\partial \phi}{\partial x} dx = \int -x dx$$

$$\phi = -\frac{x^2}{2} + f(y) \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = y^2$. Therefore equation (4) becomes

$$y^2 = 0 + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = y^2$$

Integrating the above w.r.t y gives

$$\int f'(y) dy = \int (y^2) dy$$

$$f(y) = \frac{y^3}{3} + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = \frac{y^3}{3} - \frac{x^2}{2} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \frac{y^3}{3} - \frac{x^2}{2}$$

Initial conditions are used to solve for c_1 . Substituting $x = -1$ and $y = 3$ in the above solution gives an equation to solve for the constant of integration.

$$\frac{17}{2} = c_1$$

$$c_1 = \frac{17}{2}$$

Substituting c_1 found above in the general solution gives

$$\frac{y^3}{3} - \frac{x^2}{2} = \frac{17}{2}$$

Summary

The solution(s) found are the following

$$\frac{y^3}{3} - \frac{x^2}{2} = \frac{17}{2} \tag{1}$$

Verification of solutions

$$\frac{y^3}{3} - \frac{x^2}{2} = \frac{17}{2}$$

Verified OK.

8.11.6 Maple step by step solution

Let's solve

$$[y'y^2 = x, y(-1) = 3]$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Integrate both sides with respect to x

$$\int y'y^2 dx = \int x dx + c_1$$

- Evaluate integral

$$\frac{y^3}{3} = \frac{x^2}{2} + c_1$$

- Solve for y

$$y = \frac{(12x^2 + 24c_1)^{\frac{1}{3}}}{2}$$

- Use initial condition $y(-1) = 3$

$$3 = \frac{(12 + 24c_1)^{\frac{1}{3}}}{2}$$

- Solve for c_1

$$c_1 = \frac{17}{2}$$

- Substitute $c_1 = \frac{17}{2}$ into general solution and simplify

$$y = \frac{(12x^2 + 204)^{\frac{1}{3}}}{2}$$

- Solution to the IVP

$$y = \frac{(12x^2 + 204)^{\frac{1}{3}}}{2}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
<- Bernoulli successful`
```

✓ Solution by Maple

Time used: 0.094 (sec). Leaf size: 15

```
dsolve([y(x)^2*diff(y(x),x)=x,y(-1) = 3],y(x), singsol=all)
```

$$y(x) = \frac{(12x^2 + 204)^{\frac{1}{3}}}{2}$$

✓ Solution by Mathematica

Time used: 0.191 (sec). Leaf size: 22

```
DSolve[{y[x]^2*y'[x]==x,{y[-1]==3}],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \sqrt[3]{\frac{3}{2}\sqrt[3]{x^2 + 17}}$$

8.12 problem 2(d)

8.12.1 Existence and uniqueness analysis	1183
8.12.2 Solving as separable ode	1184
8.12.3 Solving as first order ode lie symmetry lookup ode	1186
8.12.4 Solving as exact ode	1190

Internal problem ID [6260]

Internal file name [OUTPUT/5508_Sunday_June_05_2022_03_42_00_PM_51902994/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Problems for Review and Discovery. Page 53

Problem number: 2(d).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exact", "separable", "first_order_ode_lie_symmetry_lookup"**

Maple gives the following as the ode type

`[_separable]`

$$y' \csc(x) - \csc(y) = 0$$

With initial conditions

$$\left[y\left(\frac{\pi}{2}\right) = 1 \right]$$

8.12.1 Existence and uniqueness analysis

This is non linear first order ODE. In canonical form it is written as

$$y' = f(x, y) = \frac{\csc(y)}{\csc(x)}$$

The x domain of $f(x, y)$ when $y = 1$ is

$$\{x < \pi_{Z411} \vee \pi_{Z411} < x\}$$

And the point $x_0 = \frac{\pi}{2}$ is inside this domain. The y domain of $f(x, y)$ when $x = \frac{\pi}{2}$ is

$$\{y < \pi_{-Z412} \vee \pi_{-Z412} < y\}$$

And the point $y_0 = 1$ is inside this domain. Now we will look at the continuity of

$$\begin{aligned} \frac{\partial f}{\partial y} &= \frac{\partial}{\partial y} \left(\frac{\csc(y)}{\csc(x)} \right) \\ &= -\frac{\csc(y) \cot(y)}{\csc(x)} \end{aligned}$$

The x domain of $\frac{\partial f}{\partial y}$ when $y = 1$ is

$$\{x < \pi_{-Z411} \vee \pi_{-Z411} < x\}$$

And the point $x_0 = \frac{\pi}{2}$ is inside this domain. The y domain of $\frac{\partial f}{\partial y}$ when $x = \frac{\pi}{2}$ is

$$\{y < \pi_{-Z412} \vee \pi_{-Z412} < y\}$$

And the point $y_0 = 1$ is inside this domain. Therefore solution exists and is unique.

8.12.2 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{\csc(y)}{\csc(x)} \end{aligned}$$

Where $f(x) = \frac{1}{\csc(x)}$ and $g(y) = \csc(y)$. Integrating both sides gives

$$\begin{aligned} \frac{1}{\csc(y)} dy &= \frac{1}{\csc(x)} dx \\ \int \frac{1}{\csc(y)} dy &= \int \frac{1}{\csc(x)} dx \\ -\cos(y) &= -\cos(x) + c_1 \end{aligned}$$

Which results in

$$y = \pi - \arccos(-\cos(x) + c_1)$$

Initial conditions are used to solve for c_1 . Substituting $x = \frac{\pi}{2}$ and $y = 1$ in the above solution gives an equation to solve for the constant of integration.

$$1 = \frac{\pi}{2} + \arcsin(c_1)$$

$$c_1 = -\cos(1)$$


Substituting c_1 found above in the general solution gives

$$y = \frac{\pi}{2} - \arcsin(\cos(x) + \cos(1))$$


Summary

The solution(s) found are the following

$$y = \frac{\pi}{2} - \arcsin(\cos(x) + \cos(1)) \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{\pi}{2} - \arcsin(\cos(x) + \cos(1))$$

Verified OK.

8.12.3 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{\text{csc}(y)}{\text{csc}(x)}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **separable**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 162: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= \csc(x) \\ \eta(x, y) &= 0\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\eta = 0$ then in this special case

$$R = y$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\xi} dx \\ &= \int \frac{1}{\csc(x)} dx\end{aligned}$$

Which results in

$$S = -\cos(x)$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y}\tag{2}$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{\csc(y)}{\csc(x)}$$

Evaluating all the partial derivatives gives

$$\begin{aligned}R_x &= 0 \\R_y &= 1 \\S_x &= \sin(x) \\S_y &= 0\end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \sin(y) \tag{2A}$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \sin(R)$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = -\cos(R) + c_1 \tag{4}$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$-\cos(x) = -\cos(y) + c_1$$



Which simplifies to

$$-\cos(x) = -\cos(y) + c_1$$

Which gives

$$y = \arccos(\cos(x) + c_1)$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{\csc(y)}{\csc(x)}$ 	$R = y$ $S = -\cos(x)$	$\frac{dS}{dR} = \sin(R)$ 

Initial conditions are used to solve for c_1 . Substituting $x = \frac{\pi}{2}$ and $y = 1$ in the above solution gives an equation to solve for the constant of integration.

$$1 = \frac{\pi}{2} - \arcsin(c_1)$$

$$c_1 = \cos(1)$$


Substituting c_1 found above in the general solution gives

$$y = \frac{\pi}{2} - \arcsin(\cos(x) + \cos(1))$$


Summary

The solution(s) found are the following

$$y = \frac{\pi}{2} - \arcsin(\cos(x) + \cos(1)) \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{\pi}{2} - \arcsin(\cos(x) + \cos(1))$$

Verified OK.

8.12.4 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned} \left(\frac{1}{\csc(y)}\right) dy &= \left(\frac{1}{\csc(x)}\right) dx \\ \left(-\frac{1}{\csc(x)}\right) dx + \left(\frac{1}{\csc(y)}\right) dy &= 0 \end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -\frac{1}{\csc(x)} \\ N(x, y) &= \frac{1}{\csc(y)} \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(-\frac{1}{\csc(x)}\right) \\ &= 0 \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{1}{\csc(y)}\right) \\ &= 0 \end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned} \int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\frac{1}{\csc(x)} dx \\ \phi &= \cos(x) + f(y) \end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{1}{\csc(y)}$. Therefore equation (4) becomes

$$\frac{1}{\csc(y)} = 0 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$\begin{aligned} f'(y) &= \frac{1}{\csc(y)} \\ &= \sin(y) \end{aligned}$$

Integrating the above w.r.t y results in

$$\begin{aligned} \int f'(y) dy &= \int (\sin(y)) dy \\ f(y) &= -\cos(y) + c_1 \end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = \cos(x) - \cos(y) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \cos(x) - \cos(y)$$

Initial conditions are used to solve for c_1 . Substituting $x = \frac{\pi}{2}$ and $y = 1$ in the above solution gives an equation to solve for the constant of integration.

$$-\cos(1) = c_1$$

$$c_1 = -\cos(1)$$

Substituting c_1 found above in the general solution gives

$$\cos(x) - \cos(y) = -\cos(1)$$


Solving for y from the above gives

$$y = \arccos(\cos(x) + \cos(1))$$


Summary

The solution(s) found are the following

$$y = \arccos(\cos(x) + \cos(1)) \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \arccos(\cos(x) + \cos(1))$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
<- separable successful`
```

✓ Solution by Maple

Time used: 0.172 (sec). Leaf size: 10

```
dsolve([csc(x)*diff(y(x),x)=csc(y(x)),y(1/2*Pi) = 1],y(x), singsol=all)
```

$$y(x) = \arccos(\cos(x) + \cos(1))$$

✓ Solution by Mathematica

Time used: 0.398 (sec). Leaf size: 11

```
DSolve[{Csc[x]*y'[x]==Csc[y[x]],{y[Pi/2]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \arccos(\cos(x) + \cos(1))$$

8.13 problem 2(e)

8.13.1 Existence and uniqueness analysis	1195
8.13.2 Solving as homogeneousTypeD2 ode	1196
8.13.3 Solving as first order ode lie symmetry calculated ode	1197
8.13.4 Solving as exact ode	1202

Internal problem ID [6261]

Internal file name [OUTPUT/5509_Sunday_June_05_2022_03_42_02_PM_76246877/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Problems for Review and Discovery. Page 53

Problem number: 2(e).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exactByInspection", "homogeneousTypeD2", "first_order_ode_lie_symmetry_calculated"**

Maple gives the following as the ode type

```
[[_homogeneous, `class A`], _rational, [_Abel, `2nd type`, `class A`]]
```

$$y' - \frac{x+y}{x-y} = 0$$

With initial conditions

$$[y(1) = 1]$$

8.13.1 Existence and uniqueness analysis

This is non linear first order ODE. In canonical form it is written as

$$\begin{aligned} y' &= f(x, y) \\ &= -\frac{x+y}{-x+y} \end{aligned}$$

The x domain of $f(x, y)$ when $y = 1$ is

$$\{x < 1 \vee 1 < x\}$$

But the point $x_0 = 1$ is not inside this domain. Hence existence and uniqueness theorem does not apply. There could be infinite number of solutions, or one solution or no solution at all.

8.13.2 Solving as homogeneous Type D2 ode

Using the change of variables $y = u(x)x$ on the above ode results in new ode in $u(x)$

$$u'(x)x + u(x) - \frac{x + u(x)x}{x - u(x)x} = 0$$

In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{u^2 + 1}{x(u - 1)} \end{aligned}$$

Where $f(x) = -\frac{1}{x}$ and $g(u) = \frac{u^2+1}{u-1}$. Integrating both sides gives

$$\begin{aligned} \frac{1}{\frac{u^2+1}{u-1}} du &= -\frac{1}{x} dx \\ \int \frac{1}{\frac{u^2+1}{u-1}} du &= \int -\frac{1}{x} dx \\ \frac{\ln(u^2 + 1)}{2} - \arctan(u) &= -\ln(x) + c_2 \end{aligned}$$

The solution is

$$\frac{\ln(u(x)^2 + 1)}{2} - \arctan(u(x)) + \ln(x) - c_2 = 0$$

Replacing $u(x)$ in the above solution by $\frac{y}{x}$ results in the solution for y in implicit form

$$\begin{aligned} \frac{\ln\left(\frac{y^2}{x^2} + 1\right)}{2} - \arctan\left(\frac{y}{x}\right) + \ln(x) - c_2 &= 0 \\ \frac{\ln\left(\frac{y^2}{x^2} + 1\right)}{2} - \arctan\left(\frac{y}{x}\right) + \ln(x) - c_2 &= 0 \end{aligned}$$

Substituting initial conditions and solving for c_2 gives $c_2 = \frac{\ln(2)}{2} - \frac{\pi}{4}$. Hence the solution

Summary

The solution(s) found are the following
becomes

$$\frac{\ln\left(\frac{y^2}{x^2} + 1\right)}{2} - \arctan\left(\frac{y}{x}\right) + \ln(x) - \frac{\ln(2)}{2} + \frac{\pi}{4} = 0 \quad (1)$$

Verification of solutions

$$\frac{\ln\left(\frac{y^2}{x^2} + 1\right)}{2} - \arctan\left(\frac{y}{x}\right) + \ln(x) - \frac{\ln(2)}{2} + \frac{\pi}{4} = 0$$

Verified OK.

8.13.3 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = -\frac{x+y}{-x+y}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (A)$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \quad (1E)$$

$$\eta = xb_2 + yb_3 + b_1 \quad (2E)$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$b_2 - \frac{(x+y)(b_3 - a_2)}{-x+y} - \frac{(x+y)^2 a_3}{(-x+y)^2}$$

$$- \left(-\frac{1}{-x+y} - \frac{x+y}{(-x+y)^2} \right) (xa_2 + ya_3 + a_1) \quad (5E)$$

$$- \left(-\frac{1}{-x+y} + \frac{x+y}{(-x+y)^2} \right) (xb_2 + yb_3 + b_1) = 0$$

Putting the above in normal form gives

$$\frac{x^2a_2 + x^2a_3 + x^2b_2 - x^2b_3 - 2xya_2 + 2xya_3 + 2xyb_2 + 2xyb_3 - y^2a_2 - y^2a_3 - y^2b_2 + y^2b_3 + 2xb_1 - 2ya_1}{(x - y)^2} = 0$$

Setting the numerator to zero gives

$$\begin{aligned} & -x^2a_2 - x^2a_3 - x^2b_2 + x^2b_3 + 2xya_2 - 2xya_3 - 2xyb_2 \\ & - 2xyb_3 + y^2a_2 + y^2a_3 + y^2b_2 - y^2b_3 - 2xb_1 + 2ya_1 = 0 \end{aligned} \quad (6E)$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2\}$$

The above PDE (6E) now becomes

$$\begin{aligned} & -a_2v_1^2 + 2a_2v_1v_2 + a_2v_2^2 - a_3v_1^2 - 2a_3v_1v_2 + a_3v_2^2 - b_2v_1^2 \\ & - 2b_2v_1v_2 + b_2v_2^2 + b_3v_1^2 - 2b_3v_1v_2 - b_3v_2^2 + 2a_1v_2 - 2b_1v_1 = 0 \end{aligned} \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2\}$$

Equation (7E) now becomes

$$\begin{aligned} & (-a_2 - a_3 - b_2 + b_3)v_1^2 + (2a_2 - 2a_3 - 2b_2 - 2b_3)v_1v_2 \\ & - 2b_1v_1 + (a_2 + a_3 + b_2 - b_3)v_2^2 + 2a_1v_2 = 0 \end{aligned} \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} 2a_1 &= 0 \\ -2b_1 &= 0 \\ -a_2 - a_3 - b_2 + b_3 &= 0 \\ a_2 + a_3 + b_2 - b_3 &= 0 \\ 2a_2 - 2a_3 - 2b_2 - 2b_3 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= 0 \\ a_2 &= b_3 \\ a_3 &= -b_2 \\ b_1 &= 0 \\ b_2 &= b_2 \\ b_3 &= b_3 \end{aligned}$$

Substituting the above solution in the ansatz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned} \xi &= x \\ \eta &= y \end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned} \eta &= \eta - \omega(x, y) \xi \\ &= y - \left(-\frac{x+y}{-x+y} \right) (x) \\ &= \frac{-x^2 - y^2}{x-y} \\ \xi &= 0 \end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y} \right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{-x^2 - y^2}{x-y}} dy \end{aligned}$$

Which results in

$$S = \frac{\ln(x^2 + y^2)}{2} - \arctan\left(\frac{y}{x}\right)$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -\frac{x + y}{-x + y}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{x + y}{x^2 + y^2} \\ S_y &= \frac{-x + y}{x^2 + y^2} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \quad (4)$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{\ln(x^2 + y^2)}{2} - \arctan\left(\frac{y}{x}\right) = c_1$$

Which simplifies to

$$\frac{\ln(x^2 + y^2)}{2} - \arctan\left(\frac{y}{x}\right) = c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -\frac{x+y}{-x+y}$ 	$R = x$ $S = \frac{\ln(x^2 + y^2)}{2} - \arctan\left(\frac{y}{x}\right)$	$\frac{dS}{dR} = 0$ 

Initial conditions are used to solve for c_1 . Substituting $x = 1$ and $y = 1$ in the above solution gives an equation to solve for the constant of integration.

$$\frac{\ln(2)}{2} - \frac{\pi}{4} = c_1$$

$$c_1 = \frac{\ln(2)}{2} - \frac{\pi}{4}$$

Substituting c_1 found above in the general solution gives

$$\frac{\ln(x^2 + y^2)}{2} - \arctan\left(\frac{y}{x}\right) = \frac{\ln(2)}{2} - \frac{\pi}{4}$$

Summary

The solution(s) found are the following

$$\frac{\ln(x^2 + y^2)}{2} - \arctan\left(\frac{y}{x}\right) = \frac{\ln(2)}{2} - \frac{\pi}{4} \quad (1)$$

Verification of solutions

$$\frac{\ln(x^2 + y^2)}{2} - \arctan\left(\frac{y}{x}\right) = \frac{\ln(2)}{2} - \frac{\pi}{4}$$

Verified OK.

8.13.4 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial\phi}{\partial x} + \frac{\partial\phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} (-x + y) dy &= (-y - x) dx \\ (x + y) dx + (-x + y) dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= x + y \\N(x, y) &= -x + y\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(x + y) \\&= 1\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(-x + y) \\&= -1\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. By inspection $\frac{1}{x^2+y^2}$ is an integrating factor. Therefore by multiplying $M = x + y$ and $N = -x + y$ by this integrating factor the ode becomes exact. The new M, N are

$$\begin{aligned}M &= \frac{x + y}{x^2 + y^2} \\N &= \frac{-x + y}{x^2 + y^2}\end{aligned}$$

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned}\left(\frac{-x+y}{x^2+y^2}\right) dy &= \left(-\frac{x+y}{x^2+y^2}\right) dx \\ \left(\frac{x+y}{x^2+y^2}\right) dx + \left(\frac{-x+y}{x^2+y^2}\right) dy &= 0\end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= \frac{x+y}{x^2+y^2} \\ N(x, y) &= \frac{-x+y}{x^2+y^2}\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(\frac{x+y}{x^2+y^2}\right) \\ &= \frac{x^2 - 2xy - y^2}{(x^2+y^2)^2}\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{-x+y}{x^2+y^2} \right) \\ &= \frac{x^2 - 2xy - y^2}{(x^2+y^2)^2}\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int \frac{x+y}{x^2+y^2} dx \\ \phi &= \frac{\ln(x^2+y^2)}{2} + \arctan\left(\frac{x}{y}\right) + f(y)\end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\begin{aligned}\frac{\partial \phi}{\partial y} &= \frac{y}{x^2+y^2} - \frac{x}{y^2 \left(\frac{x^2}{y^2} + 1 \right)} + f'(y) \\ &= \frac{-x+y}{x^2+y^2} + f'(y)\end{aligned} \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{-x+y}{x^2+y^2}$. Therefore equation (4) becomes

$$\frac{-x+y}{x^2+y^2} = \frac{-x+y}{x^2+y^2} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = \frac{\ln(x^2 + y^2)}{2} + \arctan\left(\frac{x}{y}\right) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \frac{\ln(x^2 + y^2)}{2} + \arctan\left(\frac{x}{y}\right)$$

Initial conditions are used to solve for c_1 . Substituting $x = 1$ and $y = 1$ in the above solution gives an equation to solve for the constant of integration.

$$\frac{\ln(2)}{2} + \frac{\pi}{4} = c_1$$

$$c_1 = \frac{\ln(2)}{2} + \frac{\pi}{4}$$

Substituting c_1 found above in the general solution gives

$$\frac{\ln(x^2 + y^2)}{2} + \arctan\left(\frac{x}{y}\right) = \frac{\ln(2)}{2} + \frac{\pi}{4}$$

Summary

The solution(s) found are the following

$$\frac{\ln(x^2 + y^2)}{2} + \arctan\left(\frac{x}{y}\right) = \frac{\ln(2)}{2} + \frac{\pi}{4} \quad (1)$$

Verification of solutions

$$\frac{\ln(x^2 + y^2)}{2} + \arctan\left(\frac{x}{y}\right) = \frac{\ln(2)}{2} + \frac{\pi}{4}$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying homogeneous D  
<- homogeneous successful`
```

✓ Solution by Maple

Time used: 0.234 (sec). Leaf size: 30

```
dsolve([diff(y(x),x)=(x+y(x))/(x-y(x)),y(1) = 1],y(x), singsol=all)
```

$$y(x) = \tan(\text{RootOf}(4_Z - 2 \ln(\sec(Z)^2) - 4 \ln(x) + 2 \ln(2) - \pi)) x$$

✓ Solution by Mathematica

Time used: 0.044 (sec). Leaf size: 46

```
DSolve[{y'[x]==(x+y[x])/(x-y[x]),{y[1]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve}\left[\frac{1}{2} \log\left(\frac{y(x)^2}{x^2} + 1\right) - \arctan\left(\frac{y(x)}{x}\right) = \frac{1}{4}(2 \log(2) - \pi) - \log(x), y(x)\right]$$

8.14 problem 2(f)

8.14.1 Solving as homogeneousTypeD2 ode 1208

8.14.2 Solving as first order ode lie symmetry calculated ode 1210

Internal problem ID [6262]

Internal file name [OUTPUT/5510_Sunday_June_05_2022_03_42_05_PM_82967625/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Problems for Review and Discovery. Page 53

Problem number: 2(f).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**homogeneousTypeD2**", "**first_order_ode_lie_symmetry_calculated**"

Maple gives the following as the ode type

```
[[_homogeneous, `class A`], _rational, _dAlembert]
```

$$y' - \frac{x^2 + 2y^2}{x^2 - 2y^2} = 0$$

8.14.1 Solving as homogeneousTypeD2 ode

Using the change of variables $y = u(x)x$ on the above ode results in new ode in $u(x)$

$$u'(x)x + u(x) - \frac{x^2 + 2u(x)^2 x^2}{x^2 - 2u(x)^2 x^2} = 0$$

In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{2u^3 + 2u^2 - u + 1}{x(2u^2 - 1)} \end{aligned}$$

Where $f(x) = -\frac{1}{x}$ and $g(u) = \frac{2u^3+2u^2-u+1}{2u^2-1}$. Integrating both sides gives

$$\begin{aligned} \frac{1}{\frac{2u^3+2u^2-u+1}{2u^2-1}} du &= -\frac{1}{x} dx \\ \int \frac{1}{\frac{2u^3+2u^2-u+1}{2u^2-1}} du &= \int -\frac{1}{x} dx \\ \int^u \frac{2u^2-1}{2u^3+2u^2-u+1} du &= -\ln(x) + c_2 \end{aligned}$$

Which results in

$$\int^u \frac{2u^2-1}{2u^3+2u^2-u+1} du = -\ln(x) + c_2$$

The solution is

$$\int^{u(x)} \frac{2u^2-1}{2u^3+2u^2-u+1} du + \ln(x) - c_2 = 0$$

Replacing $u(x)$ in the above solution by $\frac{y}{x}$ results in the solution for y in implicit form

$$\begin{aligned} \int^{\frac{y}{x}} \frac{2u^2-1}{2u^3+2u^2-u+1} du + \ln(x) - c_2 &= 0 \\ \int^{\frac{y}{x}} \frac{2u^2-1}{2u^3+2u^2-u+1} du + \ln(x) - c_2 &= 0 \end{aligned}$$

Summary

The solution(s) found are the following

$$\int^{\frac{y}{x}} \frac{2u^2-1}{2u^3+2u^2-u+1} du + \ln(x) - c_2 = 0 \quad (1)$$




Figure 240: Slope field plot

Verification of solutions

$$\int^{\frac{y}{x}} \frac{2a^2 - 1}{2a^3 + 2a^2 - a + 1} da + \ln(x) - c_2 = 0$$

Verified OK.

8.14.2 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = -\frac{x^2 + 2y^2}{-x^2 + 2y^2}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \tag{A}$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \tag{1E}$$

$$\eta = xb_2 + yb_3 + b_1 \tag{2E}$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$\begin{aligned} b_2 - \frac{(x^2 + 2y^2)(b_3 - a_2)}{-x^2 + 2y^2} - \frac{(x^2 + 2y^2)^2 a_3}{(-x^2 + 2y^2)^2} \\ - \left(-\frac{2x}{-x^2 + 2y^2} - \frac{2(x^2 + 2y^2)x}{(-x^2 + 2y^2)^2} \right) (xa_2 + ya_3 + a_1) \\ - \left(-\frac{4y}{-x^2 + 2y^2} + \frac{4(x^2 + 2y^2)y}{(-x^2 + 2y^2)^2} \right) (xb_2 + yb_3 + b_1) = 0 \end{aligned} \quad (5E)$$

Putting the above in normal form gives

$$\frac{x^4 a_2 + x^4 a_3 - x^4 b_2 - x^4 b_3 + 8x^3 y b_2 - 8x^2 y^2 a_2 + 4x^2 y^2 a_3 + 4x^2 y^2 b_2 + 8x^2 y^2 b_3 - 8x y^3 a_3 - 4y^4 a_2 + 4y^4 a_3}{(x^2 - 2y^2)^2} = 0$$

Setting the numerator to zero gives

$$\begin{aligned} -x^4 a_2 - x^4 a_3 + x^4 b_2 + x^4 b_3 - 8x^3 y b_2 + 8x^2 y^2 a_2 - 4x^2 y^2 a_3 - 4x^2 y^2 b_2 \\ - 8x^2 y^2 b_3 + 8x y^3 a_3 + 4y^4 a_2 - 4y^4 a_3 + 4y^4 b_2 - 4y^4 b_3 - 8x^2 y b_1 + 8x y^2 a_1 = 0 \end{aligned} \quad (6E)$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2\}$$

The above PDE (6E) now becomes

$$\begin{aligned} -a_2 v_1^4 + 8a_2 v_1^2 v_2^2 + 4a_2 v_2^4 - a_3 v_1^4 - 4a_3 v_1^2 v_2^2 + 8a_3 v_1 v_2^3 - 4a_3 v_2^4 + b_2 v_1^4 \\ - 8b_2 v_1^3 v_2 - 4b_2 v_1^2 v_2^2 + 4b_2 v_2^4 + b_3 v_1^4 - 8b_3 v_1^2 v_2^2 - 4b_3 v_2^4 + 8a_1 v_1 v_2^2 - 8b_1 v_1^2 v_2 = 0 \end{aligned} \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2\}$$

Equation (7E) now becomes

$$\begin{aligned} &(-a_2 - a_3 + b_2 + b_3)v_1^4 - 8b_2v_1^3v_2 + (8a_2 - 4a_3 - 4b_2 - 8b_3)v_1^2v_2^2 \\ &- 8b_1v_1^2v_2 + 8a_3v_1v_2^3 + 8a_1v_1v_2^2 + (4a_2 - 4a_3 + 4b_2 - 4b_3)v_2^4 = 0 \end{aligned} \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} 8a_1 &= 0 \\ 8a_3 &= 0 \\ -8b_1 &= 0 \\ -8b_2 &= 0 \\ -a_2 - a_3 + b_2 + b_3 &= 0 \\ 4a_2 - 4a_3 + 4b_2 - 4b_3 &= 0 \\ 8a_2 - 4a_3 - 4b_2 - 8b_3 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= 0 \\ a_2 &= b_3 \\ a_3 &= 0 \\ b_1 &= 0 \\ b_2 &= 0 \\ b_3 &= b_3 \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned} \xi &= x \\ \eta &= y \end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Therefore

$$\begin{aligned} \frac{dy}{dx} &= \frac{\eta}{\xi} \\ &= \frac{y}{x} \\ &= \frac{y}{x} \end{aligned}$$

This is easily solved to give

$$y = c_1 x$$

Where now the coordinate R is taken as the constant of integration. Hence

$$R = \frac{y}{x}$$

And S is found from

$$\begin{aligned} dS &= \frac{dx}{\xi} \\ &= \frac{dx}{x} \end{aligned}$$

Integrating gives

$$\begin{aligned} S &= \int \frac{dx}{x} \\ &= \ln(x) \end{aligned}$$

Where the constant of integration is set to zero as we just need one solution. Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -\frac{x^2 + 2y^2}{-x^2 + 2y^2}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= -\frac{y}{x^2} \\ R_y &= \frac{1}{x} \\ S_x &= \frac{1}{x} \\ S_y &= 0 \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \frac{x(x^2 - 2y^2)}{x^3 - yx^2 + 2y^2x + 2y^3} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \frac{-2R^2 + 1}{2R^3 + 2R^2 - R + 1}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \int -\frac{2R^2 - 1}{2R^3 + 2R^2 - R + 1} dR + c_1 \quad (4)$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\ln(x) = \int^{\frac{y}{x}} -\frac{2a^2 - 1}{2a^3 + 2a^2 - a + 1} da + c_1$$

Which simplifies to

$$\ln(x) = \int^{\frac{y}{x}} -\frac{2a^2 - 1}{2a^3 + 2a^2 - a + 1} da + c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -\frac{x^2+2y^2}{-x^2+2y^2}$ 	$R = \frac{y}{x}$ $S = \ln(x)$	$\frac{dS}{dR} = \frac{-2R^2+1}{2R^3+2R^2-R+1}$ 

Summary

The solution(s) found are the following

$$\ln(x) = \int^{\frac{y}{x}} -\frac{2a^2 - 1}{2a^3 + 2a^2 - a + 1} da + c_1 \tag{1}$$




Figure 241: Slope field plot

Verification of solutions

$$\ln(x) = \int^{\frac{y}{x}} -\frac{2a^2 - 1}{2a^3 + 2a^2 - a + 1} da + c_1$$

Verified OK.

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying homogeneous D
<- homogeneous successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 40

```
dsolve(diff(y(x),x)=(x^2+2*y(x)^2)/(x^2-2*y(x)^2),y(x), singsol=all)
```

$$y(x) = \text{RootOf} \left(\int^{-Z} \frac{2_a^2 - 1}{2_a^3 + 2_a^2 - _a + 1} d_a + \ln(x) + c_1 \right) x$$

✓ Solution by Mathematica

Time used: 0.106 (sec). Leaf size: 80

```
DSolve[y'[x]==(x^2+2*y[x]^2)/(x^2-2*y[x]^2),y[x],x,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[\text{RootSum} \left[2\#1^3 + 2\#1^2 - \#1 \right. \right. \\ \left. \left. + 1 \&, \frac{2\#1^2 \log\left(\frac{y(x)}{x} - \#1\right) - \log\left(\frac{y(x)}{x} - \#1\right)}{6\#1^2 + 4\#1 - 1} \& \right] = -\log(x) + c_1, y(x) \right]$$

8.15 problem 2(g)

8.15.1 Existence and uniqueness analysis	1218
8.15.2 Solving as separable ode	1219
8.15.3 Solving as first order ode lie symmetry lookup ode	1221
8.15.4 Solving as exact ode	1225
8.15.5 Maple step by step solution	1229

Internal problem ID [6263]

Internal file name [OUTPUT/5511_Sunday_June_05_2022_03_42_16_PM_21153099/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Problems for Review and Discovery. Page 53

Problem number: 2(g).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "separable", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_separable]

$$2 \cos (y) x - x^2 \sin (y) y' = 0$$

With initial conditions

$$[y(1) = 1]$$

8.15.1 Existence and uniqueness analysis

This is non linear first order ODE. In canonical form it is written as

$$\begin{aligned} y' &= f(x, y) \\ &= \frac{2 \cos (y)}{x \sin (y)} \end{aligned}$$

The x domain of $f(x, y)$ when $y = 1$ is

$$\{x < 0 \vee 0 < x\}$$

And the point $x_0 = 1$ is inside this domain. The y domain of $f(x, y)$ when $x = 1$ is

$$\{y < \pi \vee \pi < y\}$$

And the point $y_0 = 1$ is inside this domain. Now we will look at the continuity of

$$\begin{aligned} \frac{\partial f}{\partial y} &= \frac{\partial}{\partial y} \left(\frac{2 \cos(y)}{x \sin(y)} \right) \\ &= -\frac{2}{x} - \frac{2 \cos(y)^2}{x \sin(y)^2} \end{aligned}$$

The x domain of $\frac{\partial f}{\partial y}$ when $y = 1$ is

$$\{x < 0 \vee 0 < x\}$$

And the point $x_0 = 1$ is inside this domain. The y domain of $\frac{\partial f}{\partial y}$ when $x = 1$ is

$$\{y < \pi \vee \pi < y\}$$

And the point $y_0 = 1$ is inside this domain. Therefore solution exists and is unique.

8.15.2 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{2 \cot(y)}{x} \end{aligned}$$

Where $f(x) = \frac{2}{x}$ and $g(y) = \cot(y)$. Integrating both sides gives

$$\begin{aligned} \frac{1}{\cot(y)} dy &= \frac{2}{x} dx \\ \int \frac{1}{\cot(y)} dy &= \int \frac{2}{x} dx \\ -\ln(\cos(y)) &= 2 \ln(x) + c_1 \end{aligned}$$

Raising both side to exponential gives

$$\frac{1}{\cos(y)} = e^{2 \ln(x) + c_1}$$

Which simplifies to

$$\sec(y) = c_2 x^2$$

Initial conditions are used to solve for c_1 . Substituting $x = 1$ and $y = 1$ in the above solution gives an equation to solve for the constant of integration.

$$1 = \frac{\pi}{2} - \arcsin\left(\frac{e^{-c_1}}{c_2}\right)$$

$$c_1 = -\ln(\cos(1) c_2)$$


Substituting c_1 found above in the general solution gives

$$y = \frac{\pi}{2} - \arcsin\left(\frac{\cos(1)}{x^2}\right)$$


Summary

The solution(s) found are the following

$$y = \frac{\pi}{2} - \arcsin\left(\frac{\cos(1)}{x^2}\right) \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{\pi}{2} - \arcsin\left(\frac{\cos(1)}{x^2}\right)$$

Verified OK.

8.15.3 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{2 \cos(y)}{x \sin(y)}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **separable**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 164: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= \frac{x}{2} \\ \eta(x, y) &= 0\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\eta = 0$ then in this special case

$$R = y$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\xi} dx \\ &= \int \frac{1}{\frac{x}{2}} dx\end{aligned}$$

Which results in

$$S = 2 \ln(x)$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y}\tag{2}$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{2 \cos(y)}{x \sin(y)}$$

Evaluating all the partial derivatives gives

$$R_x = 0$$

$$R_y = 1$$

$$S_x = \frac{2}{x}$$

$$S_y = 0$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \tan(y) \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \tan(R)$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = -\ln(\cos(R)) + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$2 \ln(x) = -\ln(\cos(y)) + c_1$$



Which simplifies to

$$2 \ln(x) = -\ln(\cos(y)) + c_1$$

Which gives

$$y = \arccos\left(\frac{e^{c_1}}{x^2}\right)$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{2 \cos(y)}{x \sin(y)}$ 	$R = y$ $S = 2 \ln(x)$	$\frac{dS}{dR} = \tan(R)$ 

Initial conditions are used to solve for c_1 . Substituting $x = 1$ and $y = 1$ in the above solution gives an equation to solve for the constant of integration.

$$1 = \frac{\pi}{2} - \arcsin(e^{c_1})$$

$$c_1 = \ln(\cos(1))$$


Substituting c_1 found above in the general solution gives

$$y = \frac{\pi}{2} - \arcsin\left(\frac{\cos(1)}{x^2}\right)$$


Summary

The solution(s) found are the following

$$y = \frac{\pi}{2} - \arcsin\left(\frac{\cos(1)}{x^2}\right) \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{\pi}{2} - \arcsin\left(\frac{\cos(1)}{x^2}\right)$$

Verified OK.

8.15.4 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial\phi}{\partial x} + \frac{\partial\phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned} \left(\frac{\sin(y)}{2 \cos(y)} \right) dy &= \left(\frac{1}{x} \right) dx \\ \left(-\frac{1}{x} \right) dx + \left(\frac{\sin(y)}{2 \cos(y)} \right) dy &= 0 \end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -\frac{1}{x} \\ N(x, y) &= \frac{\sin(y)}{2 \cos(y)} \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(-\frac{1}{x} \right) \\ &= 0 \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{\sin(y)}{2 \cos(y)} \right) \\ &= 0 \end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned} \int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\frac{1}{x} dx \\ \phi &= -\ln(x) + f(y) \end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{\sin(y)}{2 \cos(y)}$. Therefore equation (4) becomes

$$\frac{\sin(y)}{2 \cos(y)} = 0 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$\begin{aligned} f'(y) &= \frac{\sin(y)}{2 \cos(y)} \\ &= \frac{\tan(y)}{2} \end{aligned}$$

Integrating the above w.r.t y results in

$$\begin{aligned} \int f'(y) dy &= \int \left(\frac{\tan(y)}{2} \right) dy \\ f(y) &= -\frac{\ln(\cos(y))}{2} + c_1 \end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\ln(x) - \frac{\ln(\cos(y))}{2} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\ln(x) - \frac{\ln(\cos(y))}{2}$$

Initial conditions are used to solve for c_1 . Substituting $x = 1$ and $y = 1$ in the above solution gives an equation to solve for the constant of integration.

$$-\frac{\ln(\cos(1))}{2} = c_1$$

$$c_1 = -\frac{\ln(\cos(1))}{2}$$

Substituting c_1 found above in the general solution gives

$$-\ln(x) - \frac{\ln(\cos(y))}{2} = -\frac{\ln(\cos(1))}{2}$$


Solving for y from the above gives

$$y = \arccos\left(\frac{\cos(1)}{x^2}\right)$$


Summary

The solution(s) found are the following

$$y = \arccos\left(\frac{\cos(1)}{x^2}\right) \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \arccos\left(\frac{\cos(1)}{x^2}\right)$$

Verified OK.

8.15.5 Maple step by step solution

Let's solve

$$[2 \cos(y) x - x^2 \sin(y) y' = 0, y(1) = 1]$$

- Highest derivative means the order of the ODE is 1

y'

- Integrate both sides with respect to x

$$\int (2 \cos(y) x - x^2 \sin(y) y') dx = \int 0 dx + c_1$$

- Evaluate integral

$$\cos(y) x^2 = c_1$$

- Solve for y

$$y = \arccos\left(\frac{c_1}{x^2}\right)$$

- Use initial condition $y(1) = 1$

$$1 = \arccos(c_1)$$

- Solve for c_1
 $c_1 = \cos(1)$
- Substitute $c_1 = \cos(1)$ into general solution and simplify
 $y = \arccos\left(\frac{\cos(1)}{x^2}\right)$
- Solution to the IVP
 $y = \arccos\left(\frac{\cos(1)}{x^2}\right)$

Maple trace

```
`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
<- separable successful`
```

✓ Solution by Maple

Time used: 0.64 (sec). Leaf size: 11

```
dsolve([2*x*cos(y(x))-x^2*sin(y(x))*diff(y(x),x)=0,y(1) = 1],y(x), singsol=all)
```

$$y(x) = \arccos\left(\frac{\cos(1)}{x^2}\right)$$

✓ Solution by Mathematica

Time used: 29.379 (sec). Leaf size: 12

```
DSolve[{2*x*Cos[y[x]]-x^2*Sin[y[x]]*y'[x]==0,{y[1]==1}},y[x],x,IncludeSingularSolutions -> T
```

$$y(x) \rightarrow \arccos\left(\frac{\cos(1)}{x^2}\right)$$

8.16 problem 2(h)

8.16.1 Existence and uniqueness analysis	1232
8.16.2 Solving as separable ode	1232
8.16.3 Solving as linear ode	1233
8.16.4 Solving as homogeneousTypeD2 ode	1234
8.16.5 Solving as first order ode lie symmetry lookup ode	1235
8.16.6 Solving as exact ode	1239
8.16.7 Maple step by step solution	1242

Internal problem ID [6264]

Internal file name [OUTPUT/5512_Sunday_June_05_2022_03_42_20_PM_46598502/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Problems for Review and Discovery. Page 53

Problem number: 2(h).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "linear", "separable", "homogeneousTypeD2", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_separable]

$$\frac{1}{y} - \frac{xy'}{y^2} = 0$$

With initial conditions

$$[y(0) = 2]$$

8.16.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -\frac{1}{x}$$
$$q(x) = 0$$

Hence the ode is

$$y' - \frac{y}{x} = 0$$

The domain of $p(x) = -\frac{1}{x}$ is

$$\{x < 0 \vee 0 < x\}$$

But the point $x_0 = 0$ is not inside this domain. Hence existence and uniqueness theorem does not apply. There could be infinite number of solutions, or one solution or no solution at all.

8.16.2 Solving as separable ode

In canonical form the ODE is

$$y' = F(x, y)$$
$$= f(x)g(y)$$
$$= \frac{y}{x}$$

Where $f(x) = \frac{1}{x}$ and $g(y) = y$. Integrating both sides gives

$$\frac{1}{y} dy = \frac{1}{x} dx$$
$$\int \frac{1}{y} dy = \int \frac{1}{x} dx$$
$$\ln(y) = \ln(x) + c_1$$
$$y = e^{\ln(x)+c_1}$$
$$= c_1 x$$

Initial conditions are used to solve for c_1 . Substituting $x = 0$ and $y = 2$ in the above solution gives an equation to solve for the constant of integration.

$$2 = 0$$

Summary

This shows that no solution exist. The solution(s) found are the following

$$y = c_1x$$

Verification of solutions

$$y = c_1x$$

Warning, solution could not be verified

8.16.3 Solving as linear ode

Entering Linear first order ODE solver. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int -\frac{1}{x} dx} \\ &= \frac{1}{x}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}\mu y &= 0 \\ \frac{d}{dx}\left(\frac{y}{x}\right) &= 0\end{aligned}$$

Integrating gives

$$\frac{y}{x} = c_1$$

Dividing both sides by the integrating factor $\mu = \frac{1}{x}$ results in

$$y = c_1x$$

Initial conditions are used to solve for c_1 . Substituting $x = 0$ and $y = 2$ in the above solution gives an equation to solve for the constant of integration.

$$2 = 0$$

Summary

This shows that no solution exist. The solution(s) found are the following

$$y = c_1x$$

Verification of solutions

$$y = c_1x$$

Warning, solution could not be verified

8.16.4 Solving as homogeneous Type D2 ode

Using the change of variables $y = u(x)x$ on the above ode results in new ode in $u(x)$

$$\frac{1}{u(x)x} - \frac{u'(x)x + u(x)}{xu(x)^2} = 0$$

Integrating both sides gives

$$\begin{aligned} u(x) &= \int 0 \, dx \\ &= c_2 \end{aligned}$$

Therefore the solution y is

$$\begin{aligned} y &= xu \\ &= c_2x \end{aligned}$$

Initial conditions are used to solve for c_2 . Substituting $x = 0$ and $y = 2$ in the above solution gives an equation to solve for the constant of integration.

$$2 = 0$$

Summary

This shows that no solution exist. The solution(s) found are the following

$$y = c_2x$$

Verification of solutions

$$y = c_2x$$

Warning, solution could not be verified

8.16.5 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{y}{x}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 167: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= x\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{x} dy\end{aligned}$$

Which results in

$$S = \frac{y}{x}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y}\tag{2}$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{y}{x}$$

Evaluating all the partial derivatives gives

$$\begin{aligned}R_x &= 1 \\ R_y &= 0 \\ S_x &= -\frac{y}{x^2} \\ S_y &= \frac{1}{x}\end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{y}{x} = c_1$$



Which simplifies to

$$\frac{y}{x} = c_1$$

Which gives

$$y = c_1 x$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{y}{x}$ 	$R = x$ $S = \frac{y}{x}$	$\frac{dS}{dR} = 0$ 

Initial conditions are used to solve for c_1 . Substituting $x = 0$ and $y = 2$ in the above solution gives an equation to solve for the constant of integration.

$$2 = 0$$

Summary

This shows that no solution exist. The solution(s) found are the following

$$y = c_1x$$

Verification of solutions

$$y = c_1x$$

Warning, solution could not be verified

8.16.6 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial\phi}{\partial x} + \frac{\partial\phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} \left(\frac{1}{y}\right) dy &= \left(\frac{1}{x}\right) dx \\ \left(-\frac{1}{x}\right) dx + \left(\frac{1}{y}\right) dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$M(x, y) = -\frac{1}{x}$$
$$N(x, y) = \frac{1}{y}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\frac{\partial M}{\partial y} = \frac{\partial}{\partial y} \left(-\frac{1}{x} \right)$$
$$= 0$$

And

$$\frac{\partial N}{\partial x} = \frac{\partial}{\partial x} \left(\frac{1}{y} \right)$$
$$= 0$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \tag{1}$$

$$\frac{\partial \phi}{\partial y} = N \tag{2}$$

Integrating (1) w.r.t. x gives

$$\int \frac{\partial \phi}{\partial x} dx = \int M dx$$
$$\int \frac{\partial \phi}{\partial x} dx = \int -\frac{1}{x} dx$$
$$\phi = -\ln(x) + f(y) \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{1}{y}$. Therefore equation (4) becomes

$$\frac{1}{y} = 0 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = \frac{1}{y}$$

Integrating the above w.r.t y gives

$$\int f'(y) dy = \int \left(\frac{1}{y}\right) dy$$
$$f(y) = \ln(y) + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\ln(x) + \ln(y) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\ln(x) + \ln(y)$$

The solution becomes

$$y = x e^{c_1}$$

Initial conditions are used to solve for c_1 . Substituting $x = 0$ and $y = 2$ in the above solution gives an equation to solve for the constant of integration.

$$2 = 0$$

Summary

This shows that no solution exist. The solution(s) found are the following

$$y = x e^{c_1}$$

Verification of solutions

$$y = x e^{c_1}$$

Warning, solution could not be verified

8.16.7 Maple step by step solution

Let's solve

$$\left[\frac{1}{y} - \frac{xy'}{y^2} = 0, y(0) = 2 \right]$$

- Highest derivative means the order of the ODE is 1

y'

- Integrate both sides with respect to x

$$\int \left(\frac{1}{y} - \frac{xy'}{y^2} \right) dx = \int 0 dx + c_1$$

- Evaluate integral

$$\frac{x}{y} = c_1$$

- Solve for y

$$y = \frac{x}{c_1}$$

- Use initial condition $y(0) = 2$

$$2 = 0$$

- Solution does not satisfy initial condition

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✗ Solution by Maple

```
dsolve([1/y(x)-x/y(x)^2*diff(y(x),x)=0,y(0) = 2],y(x), singsol=all)
```

No solution found

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[{1/y[x]-x/y[x]^2*y'[x]==0,{y[0]==2}},y[x],x,IncludeSingularSolutions -> True]
```

```
{}
```


8.17 problem 4(a)

8.17.1 Solving as second order ode missing x ode 1244

8.17.2 Maple step by step solution 1246

Internal problem ID [6265]

Internal file name [OUTPUT/5513_Sunday_June_05_2022_03_42_21_PM_66243305/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Problems for Review and Discovery.

Page 53

Problem number: 4(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_ode_missing_x**"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x], _Liouville, [_2nd_order, _reducible,
    _mu_x_y1], [_2nd_order, _reducible, _mu_xy]]
```

$$yy'' - y'^2 = 0$$

8.17.1 Solving as second order ode missing x ode

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable y an independent variable.

Using

$$y' = p(y)$$

Then

$$\begin{aligned} y'' &= \frac{dp}{dx} \\ &= \frac{dy}{dx} \frac{dp}{dy} \\ &= p \frac{dp}{dy} \end{aligned}$$

Hence the ode becomes

$$yp(y) \left(\frac{d}{dy} p(y) \right) - p(y)^2 = 0$$

Which is now solved as first order ode for $p(y)$. In canonical form the ODE is

$$\begin{aligned} p' &= F(y, p) \\ &= f(y)g(p) \\ &= \frac{p}{y} \end{aligned}$$

Where $f(y) = \frac{1}{y}$ and $g(p) = p$. Integrating both sides gives

$$\begin{aligned} \frac{1}{p} dp &= \frac{1}{y} dy \\ \int \frac{1}{p} dp &= \int \frac{1}{y} dy \\ \ln(p) &= \ln(y) + c_1 \\ p &= e^{\ln(y)+c_1} \\ &= c_1 y \end{aligned}$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = yc_1$$

Integrating both sides gives

$$\begin{aligned} \int \frac{1}{c_1 y} dy &= \int dx \\ \frac{\ln(y)}{c_1} &= c_2 + x \end{aligned}$$

Raising both side to exponential gives

$$e^{\frac{\ln(y)}{c_1}} = e^{c_2+x}$$

Which simplifies to

$$y^{\frac{1}{c_1}} = c_3 e^x$$

Summary

The solution(s) found are the following

$$y = (c_3 e^x)^{c_1} \tag{1}$$

Verification of solutions

$$y = (c_3 e^x)^{c_1}$$

Verified OK.

8.17.2 Maple step by step solution

Let's solve

$$yy'' - y'^2 = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Define new dependent variable u

$$u(x) = y'$$

- Compute y''

$$u'(x) = y''$$

- Use chain rule on the lhs

$$y' \left(\frac{d}{dy} u(y) \right) = y''$$

- Substitute in the definition of u

$$u(y) \left(\frac{d}{dy} u(y) \right) = y''$$

- Make substitutions $y' = u(y)$, $y'' = u(y) \left(\frac{d}{dy} u(y) \right)$ to reduce order of ODE

$$yu(y) \left(\frac{d}{dy} u(y) \right) - u(y)^2 = 0$$

- Separate variables

$$\frac{\frac{d}{dy} u(y)}{u(y)} = \frac{1}{y}$$

- Integrate both sides with respect to y

$$\int \frac{\frac{d}{dy} u(y)}{u(y)} dy = \int \frac{1}{y} dy + c_1$$

- Evaluate integral

$$\ln(u(y)) = \ln(y) + c_1$$

- Solve for $u(y)$

$$u(y) = y e^{c_1}$$

- Solve 1st ODE for $u(y)$

$$u(y) = y e^{c_1}$$

- Revert to original variables with substitution $u(y) = y'$, $y = y$

$$y' = y e^{c_1}$$

- Separate variables

$$\frac{y'}{y} = e^{c_1}$$
- Integrate both sides with respect to x

$$\int \frac{y'}{y} dx = \int e^{c_1} dx + c_2$$
- Evaluate integral

$$\ln(y) = x e^{c_1} + c_2$$
- Solve for y

$$y = e^{x e^{c_1} + c_2}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
<- 2nd_order Liouville successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 14

```
dsolve(y(x)*diff(y(x),x$2)-(diff(y(x),x))^2=0,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = e^{c_1 x} c_2$$

✓ Solution by Mathematica

Time used: 0.113 (sec). Leaf size: 14

```
DSolve[y[x]*y'[x]-(y'[x])^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 e^{c_1 x}$$

8.18 problem 4(b)

8.18.1 Solving as second order ode missing y ode 1248

8.18.2 Maple step by step solution 1250

Internal problem ID [6266]

Internal file name [OUTPUT/5514_Sunday_June_05_2022_03_42_24_PM_11812817/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Problems for Review and Discovery. Page 53

Problem number: 4(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_ode_missing_y**"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_y], [_2nd_order, _reducible, _mu_y_y1]]
```

$$xy'' - y' + 2y'^3 = 0$$

8.18.1 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x)x + (-1 + 2p(x)^2)p(x) = 0$$

Which is now solve for $p(x)$ as first order ode. In canonical form the ODE is

$$\begin{aligned} p' &= F(x, p) \\ &= f(x)g(p) \\ &= -\frac{2p^3 - p}{x} \end{aligned}$$

Where $f(x) = -\frac{1}{x}$ and $g(p) = 2p^3 - p$. Integrating both sides gives

$$\begin{aligned}\frac{1}{2p^3 - p} dp &= -\frac{1}{x} dx \\ \int \frac{1}{2p^3 - p} dp &= \int -\frac{1}{x} dx \\ \frac{\ln(2p^2 - 1)}{2} - \ln(p) &= -\ln(x) + c_1\end{aligned}$$

Raising both side to exponential gives

$$e^{\frac{\ln(2p^2 - 1)}{2} - \ln(p)} = e^{-\ln(x) + c_1}$$

Which simplifies to

$$\frac{\sqrt{2p^2 - 1}}{p} = \frac{c_2}{x}$$

The solution is

$$\frac{\sqrt{-1 + 2p(x)^2}}{p(x)} = \frac{c_2}{x}$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$\frac{\sqrt{-1 + 2y'^2}}{y'} = \frac{c_2}{x}$$

Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = -\frac{x}{\sqrt{-c_2^2 + 2x^2}} \tag{1}$$

$$y' = \frac{x}{\sqrt{-c_2^2 + 2x^2}} \tag{2}$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\begin{aligned}y &= \int -\frac{x}{\sqrt{-c_2^2 + 2x^2}} dx \\ &= -\frac{\sqrt{-c_2^2 + 2x^2}}{2} + c_3\end{aligned}$$

Solving equation (2)

Integrating both sides gives

$$\begin{aligned}y &= \int \frac{x}{\sqrt{-c_2^2 + 2x^2}} dx \\ &= \frac{\sqrt{-c_2^2 + 2x^2}}{2} + c_4\end{aligned}$$

Summary

The solution(s) found are the following

$$y = -\frac{\sqrt{-c_2^2 + 2x^2}}{2} + c_3 \quad (1)$$

$$y = \frac{\sqrt{-c_2^2 + 2x^2}}{2} + c_4 \quad (2)$$

Verification of solutions

$$y = -\frac{\sqrt{-c_2^2 + 2x^2}}{2} + c_3$$

Verified OK.

$$y = \frac{\sqrt{-c_2^2 + 2x^2}}{2} + c_4$$

Verified OK.

8.18.2 Maple step by step solution

Let's solve

$$y''x + (-1 + 2y'^2)y' = 0$$

- Highest derivative means the order of the ODE is 2

y''

- Make substitution $u = y'$ to reduce order of ODE

$$u'(x)x + (-1 + 2u(x)^2)u(x) = 0$$

- Separate variables

$$\frac{u'(x)}{(-1+2u(x)^2)u(x)} = -\frac{1}{x}$$

- Integrate both sides with respect to x

$$\int \frac{u'(x)}{(-1+2u(x)^2)u(x)} dx = \int -\frac{1}{x} dx + c_1$$

- Evaluate integral

$$\frac{\ln(-1+2u(x)^2)}{2} - \ln(u(x)) = -\ln(x) + c_1$$

- Solve for $u(x)$

$$\left\{ u(x) = \frac{x}{\sqrt{2x^2-(e^{c_1})^2}}, u(x) = -\frac{x}{\sqrt{2x^2-(e^{c_1})^2}} \right\}$$

- Solve 1st ODE for $u(x)$

$$u(x) = \frac{x}{\sqrt{2x^2-(e^{c_1})^2}}$$

- Make substitution $u = y'$

$$y' = \frac{x}{\sqrt{2x^2-(e^{c_1})^2}}$$

- Integrate both sides to solve for y

$$\int y' dx = \int \frac{x}{\sqrt{2x^2-(e^{c_1})^2}} dx + c_2$$

- Compute integrals

$$y = \frac{\sqrt{2x^2-(e^{c_1})^2}}{2} + c_2$$

- Solve 2nd ODE for $u(x)$

$$u(x) = -\frac{x}{\sqrt{2x^2-(e^{c_1})^2}}$$

- Make substitution $u = y'$

$$y' = -\frac{x}{\sqrt{2x^2-(e^{c_1})^2}}$$

- Integrate both sides to solve for y

$$\int y' dx = \int -\frac{x}{\sqrt{2x^2-(e^{c_1})^2}} dx + c_2$$

- Compute integrals

$$y = -\frac{\sqrt{2x^2-(e^{c_1})^2}}{2} + c_2$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
--- trying a change of variables {x -> y(x), y(x) -> x}
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
-> Calling odsolve with the ODE`, diff(_b(_a), _a) = -_b(_a)*(2*_b(_a)^2-1)/_a, _b(_a), HINT
symmetry methods on request
`, `1st order, trying reduction of order with given symmetries: `[_a, 0]
```

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 37

```
dsolve(x*diff(y(x),x$2)=diff(y(x),x)-2*(diff(y(x),x))^3,y(x), singsol=all)
```

$$y(x) = \frac{\sqrt{2x^2 - c_1}}{2} + c_2$$
$$y(x) = -\frac{\sqrt{2x^2 - c_1}}{2} + c_2$$

✓ Solution by Mathematica

Time used: 0.628 (sec). Leaf size: 96

```
DSolve[x*y'[x]==y'[x]-2*(y'[x])^3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 - \frac{1}{2}\sqrt{2x^2 + e^{2c_1}}$$

$$y(x) \rightarrow \frac{1}{2}\sqrt{2x^2 + e^{2c_1}} + c_2$$

$$y(x) \rightarrow -\frac{\sqrt{x^2}}{\sqrt{2}} + c_2$$

$$y(x) \rightarrow \frac{\sqrt{x^2}}{\sqrt{2}} + c_2$$

8.19 problem 4(c)

8.19.1 Solving as second order ode missing x ode 1254

8.19.2 Maple step by step solution 1256

Internal problem ID [6267]

Internal file name [OUTPUT/5515_Sunday_June_05_2022_03_42_26_PM_82411712/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Problems for Review and Discovery. Page 53

Problem number: 4(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_ode_missing_x**"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1],  
 [_2nd_order, _reducible, _mu_xy]]
```

$$yy'' + y' = 0$$

8.19.1 Solving as second order ode missing x ode

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable y an independent variable. Using

$$y' = p(y)$$

Then

$$\begin{aligned} y'' &= \frac{dp}{dx} \\ &= \frac{dy}{dx} \frac{dp}{dy} \\ &= p \frac{dp}{dy} \end{aligned}$$

Hence the ode becomes

$$yp(y) \left(\frac{d}{dy} p(y) \right) + p(y) = 0$$

Which is now solved as first order ode for $p(y)$. Integrating both sides gives

$$\begin{aligned} p(y) &= \int -\frac{1}{y} dy \\ &= -\ln(y) + c_1 \end{aligned}$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = -\ln(y) + c_1$$

Integrating both sides gives

$$\begin{aligned} \int \frac{1}{-\ln(y) + c_1} dy &= \int dx \\ e^{c_1 \expIntegral_1(-\ln(y) + c_1)} &= c_2 + x \end{aligned}$$

Raising both side to exponential gives

$$e^{e^{c_1 \expIntegral_1(-\ln(y) + c_1)}} = e^{c_2 + x}$$

Which simplifies to

$$e^{e^{c_1 \expIntegral_1(-\ln(y) + c_1)}} = c_3 e^x$$

Summary

The solution(s) found are the following

$$y = e^{\text{RootOf}(-\expIntegral_1(-Z+c_1)e^{c_1} + \ln(c_3) + x)} \quad (1)$$

Verification of solutions

$$y = e^{\text{RootOf}(-\expIntegral_1(-Z+c_1)e^{c_1} + \ln(c_3) + x)}$$

Verified OK.

8.19.2 Maple step by step solution

Let's solve

$$yy'' + y' = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Define new dependent variable u

$$u(x) = y'$$

- Compute y''

$$u'(x) = y''$$

- Use chain rule on the lhs

$$y' \left(\frac{d}{dy} u(y) \right) = y''$$

- Substitute in the definition of u

$$u(y) \left(\frac{d}{dy} u(y) \right) = y''$$

- Make substitutions $y' = u(y)$, $y'' = u(y) \left(\frac{d}{dy} u(y) \right)$ to reduce order of ODE

$$yu(y) \left(\frac{d}{dy} u(y) \right) + u(y) = 0$$

- Separate variables

$$\frac{d}{dy} u(y) = -\frac{1}{y}$$

- Integrate both sides with respect to y

$$\int \left(\frac{d}{dy} u(y) \right) dy = \int -\frac{1}{y} dy + c_1$$

- Evaluate integral

$$u(y) = -\ln(y) + c_1$$

- Solve for $u(y)$

$$u(y) = -\ln(y) + c_1$$

- Solve 1st ODE for $u(y)$

$$u(y) = -\ln(y) + c_1$$

- Revert to original variables with substitution $u(y) = y'$, $y = y$

$$y' = -\ln(y) + c_1$$

- Separate variables

$$\frac{y'}{-\ln(y)+c_1} = 1$$

- Integrate both sides with respect to x

$$\int \frac{y'}{-\ln(y)+c_1} dx = \int 1 dx + c_2$$

- Evaluate integral

$$\text{Ei}_1(-\ln(y) + c_1) e^{c_1} = c_2 + x$$

- Solve for y

$$\{e^{\text{RootOf}(-\text{Ei}_1(-Z+c_1)e^{c_1}+c_2+x)}\}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
-> Calling odsolve with the ODE`, (diff(_b(_a), _a))*_b(_a)+_b(_a)/_a = 0, _b(_a), HINT = [
symmetry methods on request
`, `1st order, trying reduction of order with given symmetries: `[_a, 0]

```

✓ Solution by Maple

Time used: 0.125 (sec). Leaf size: 24

```
dsolve(y(x)*diff(y(x),x$2)+diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = e^{\text{RootOf}(-e^{c_1} \text{expIntegral}_1(-_Z+c_1)+x+c_2)}$$

✓ Solution by Mathematica

Time used: 0.248 (sec). Leaf size: 80

```
DSolve[y[x]*y'[x]+y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \text{InverseFunction}[-e^{c_1} \text{ExpIntegralEi}(\log(\#1) - c_1)\&][x + c_2]$$

$$y(x) \rightarrow \text{InverseFunction}[-e^{-c_1} \text{ExpIntegralEi}(\log(\#1) - -c_1)\&][x + c_2]$$

$$y(x) \rightarrow \text{InverseFunction}[-e^{c_1} \text{ExpIntegralEi}(\log(\#1) - c_1)\&][x + c_2]$$

8.20 problem 4(d)

8.20.1 Solving as second order integrable as is ode	1260
8.20.2 Solving as second order ode missing y ode	1261
8.20.3 Solving as second order ode non constant coeff transformation on B ode	1263
8.20.4 Solving as type second_order_integrable_as_is (not using ABC version)	1267
8.20.5 Solving using Kovacic algorithm	1269
8.20.6 Solving as exact linear second order ode ode	1276
8.20.7 Maple step by step solution	1278

Internal problem ID [6268]

Internal file name [OUTPUT/5516_Sunday_June_05_2022_03_42_29_PM_92411154/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 1. What is a differential equation. Problems for Review and Discovery.
Page 53

Problem number: 4(d).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "exact linear second order ode", "second_order_integrable_as_is", "second_order_ode_missing_y", "second_order_ode_non_constant_coeff_transformation_on_B"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_y]]
```

$$xy'' - 3y' = 5x$$

8.20.1 Solving as second order integrable as is ode

Integrating both sides of the ODE w.r.t x gives

$$\int (xy'' - 3y') dx = \int 5x dx$$
$$xy' - 4y = \frac{5x^2}{2} + c_1$$

Which is now solved for y .

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -\frac{4}{x}$$
$$q(x) = \frac{5x^2 + 2c_1}{2x}$$

Hence the ode is

$$y' - \frac{4y}{x} = \frac{5x^2 + 2c_1}{2x}$$

The integrating factor μ is

$$\mu = e^{\int -\frac{4}{x} dx}$$
$$= \frac{1}{x^4}$$

The ode becomes

$$\frac{d}{dx}(\mu y) = (\mu) \left(\frac{5x^2 + 2c_1}{2x} \right)$$
$$\frac{d}{dx} \left(\frac{y}{x^4} \right) = \left(\frac{1}{x^4} \right) \left(\frac{5x^2 + 2c_1}{2x} \right)$$
$$d \left(\frac{y}{x^4} \right) = \left(\frac{5x^2 + 2c_1}{2x^5} \right) dx$$

Integrating gives

$$\frac{y}{x^4} = \int \frac{5x^2 + 2c_1}{2x^5} dx$$
$$\frac{y}{x^4} = -\frac{c_1}{4x^4} - \frac{5}{4x^2} + c_2$$

Dividing both sides by the integrating factor $\mu = \frac{1}{x^4}$ results in

$$y = x^4 \left(-\frac{c_1}{4x^4} - \frac{5}{4x^2} \right) + c_2 x^4$$

which simplifies to

$$y = -\frac{5}{4}x^2 - \frac{1}{4}c_1 + c_2x^4$$

Summary

The solution(s) found are the following

$$y = -\frac{5}{4}x^2 - \frac{1}{4}c_1 + c_2x^4 \quad (1)$$

Verification of solutions

$$y = -\frac{5}{4}x^2 - \frac{1}{4}c_1 + c_2x^4$$

Verified OK.

8.20.2 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x)x - 3p(x) - 5x = 0$$

Which is now solve for $p(x)$ as first order ode.

Entering Linear first order ODE solver. In canonical form a linear first order is

$$p'(x) + p(x)p(x) = q(x)$$

Where here

$$p(x) = -\frac{3}{x}$$
$$q(x) = 5$$

Hence the ode is

$$p'(x) - \frac{3p(x)}{x} = 5$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int -\frac{3}{x} dx} \\ &= \frac{1}{x^3}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu p) &= (\mu)(5) \\ \frac{d}{dx}\left(\frac{p}{x^3}\right) &= \left(\frac{1}{x^3}\right)(5) \\ d\left(\frac{p}{x^3}\right) &= \left(\frac{5}{x^3}\right) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}\frac{p}{x^3} &= \int \frac{5}{x^3} dx \\ \frac{p}{x^3} &= -\frac{5}{2x^2} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = \frac{1}{x^3}$ results in

$$p(x) = -\frac{5}{2}x + c_1x^3$$

Since $p = y'$ then the new first order ode to solve is

$$y' = -\frac{5}{2}x + c_1x^3$$

Integrating both sides gives

$$\begin{aligned}y &= \int -\frac{5}{2}x + c_1x^3 dx \\ &= \frac{(2c_1x^2 - 5)^2}{16c_1} + c_2\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{(2c_1x^2 - 5)^2}{16c_1} + c_2 \quad (1)$$

Verification of solutions

$$y = \frac{(2c_1x^2 - 5)^2}{16c_1} + c_2$$

Verified OK.

8.20.3 Solving as second order ode non constant coeff transformation on B ode

Given an ode of the form

$$Ay'' + By' + Cy = F(x)$$

This method reduces the order ode the ODE by one by applying the transformation

$$y = Bv$$

This results in

$$\begin{aligned}y' &= B'v + v'B \\y'' &= B''v + B'v' + v''B + v'B' \\ &= v''B + 2v' + B' + B''v\end{aligned}$$

And now the original ode becomes

$$\begin{aligned}A(v''B + 2v'B' + B''v) + B(B'v + v'B) + CBv &= 0 \\ABv'' + (2AB' + B^2)v' + (AB'' + BB' + CB)v &= 0\end{aligned}\tag{1}$$

If the term $AB'' + BB' + CB$ is zero, then this method works and can be used to solve

$$ABv'' + (2AB' + B^2)v' = 0$$

By Using $u = v'$ which reduces the order of the above ode to one. The new ode is

$$ABu' + (2AB' + B^2)u = 0$$

The above ode is first order ode which is solved for u . Now a new ode $v' = u$ is solved for v as first order ode. Then the final solution is obtain from $y = Bv$.

This method works only if the term $AB'' + BB' + CB$ is zero. The given ODE shows that

$$\begin{aligned}A &= x \\B &= -3 \\C &= 0 \\F &= 5x\end{aligned}$$

The above shows that for this ode

$$\begin{aligned} AB'' + BB' + CB &= (x)(0) + (-3)(0) + (0)(-3) \\ &= 0 \end{aligned}$$

Hence the ode in v given in (1) now simplifies to

$$-3xv'' + (9)v' = 0$$

Now by applying $v' = u$ the above becomes

$$-3xu'(x) + 9u(x) = 0$$

Which is now solved for u . In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= \frac{3u}{x} \end{aligned}$$

Where $f(x) = \frac{3}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned} \frac{1}{u} du &= \frac{3}{x} dx \\ \int \frac{1}{u} du &= \int \frac{3}{x} dx \\ \ln(u) &= 3 \ln(x) + c_1 \\ u &= e^{3 \ln(x) + c_1} \\ &= c_1 x^3 \end{aligned}$$

The ode for v now becomes

$$\begin{aligned} v' &= u \\ &= c_1 x^3 \end{aligned}$$

Which is now solved for v . Integrating both sides gives

$$\begin{aligned} v(x) &= \int c_1 x^3 dx \\ &= \frac{c_1 x^4}{4} + c_2 \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}y_h(x) &= Bv \\ &= (-3) \left(\frac{c_1 x^4}{4} + c_2 \right) \\ &= -\frac{3c_1 x^4}{4} - 3c_2\end{aligned}$$

And now the particular solution $y_p(x)$ will be found. The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = -3$$

$$y_2 = x^4$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} -3 & x^4 \\ \frac{d}{dx}(-3) & \frac{d}{dx}(x^4) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} -3 & x^4 \\ 0 & 4x^3 \end{vmatrix}$$

Therefore

$$W = (-3)(4x^3) - (x^4)(0)$$

Which simplifies to

$$W = -12x^3$$

Which simplifies to

$$W = -12x^3$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{5x^5}{-12x^4} dx$$

Which simplifies to

$$u_1 = - \int -\frac{5x}{12} dx$$

Hence

$$u_1 = \frac{5x^2}{24}$$

And Eq. (3) becomes

$$u_2 = \int \frac{-15x}{-12x^4} dx$$

Which simplifies to

$$u_2 = \int \frac{5}{4x^3} dx$$

Hence

$$u_2 = -\frac{5}{8x^2}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{5x^2}{4}$$

Hence the complete solution is

$$\begin{aligned}y(x) &= y_h + y_p \\&= \left(-\frac{3c_1x^4}{4} - 3c_2\right) + \left(-\frac{5x^2}{4}\right) \\&= -\frac{3}{4}c_1x^4 - 3c_2 - \frac{5}{4}x^2\end{aligned}$$

Summary

The solution(s) found are the following

$$y = -\frac{3}{4}c_1x^4 - 3c_2 - \frac{5}{4}x^2 \quad (1)$$

Verification of solutions

$$y = -\frac{3}{4}c_1x^4 - 3c_2 - \frac{5}{4}x^2$$

Verified OK.

8.20.4 Solving as type `second_order_integrable_as_is` (not using ABC version)

Writing the ode as

$$xy'' - 3y' = 5x$$

Integrating both sides of the ODE w.r.t x gives

$$\begin{aligned}\int (xy'' - 3y') dx &= \int 5x dx \\xy' - 4y &= \frac{5x^2}{2} + c_1\end{aligned}$$

Which is now solved for y .

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$\begin{aligned}p(x) &= -\frac{4}{x} \\q(x) &= \frac{5x^2 + 2c_1}{2x}\end{aligned}$$

Hence the ode is

$$y' - \frac{4y}{x} = \frac{5x^2 + 2c_1}{2x}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int -\frac{4}{x} dx} \\ &= \frac{1}{x^4}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu) \left(\frac{5x^2 + 2c_1}{2x} \right) \\ \frac{d}{dx} \left(\frac{y}{x^4} \right) &= \left(\frac{1}{x^4} \right) \left(\frac{5x^2 + 2c_1}{2x} \right) \\ d \left(\frac{y}{x^4} \right) &= \left(\frac{5x^2 + 2c_1}{2x^5} \right) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}\frac{y}{x^4} &= \int \frac{5x^2 + 2c_1}{2x^5} dx \\ \frac{y}{x^4} &= -\frac{c_1}{4x^4} - \frac{5}{4x^2} + c_2\end{aligned}$$

Dividing both sides by the integrating factor $\mu = \frac{1}{x^4}$ results in

$$y = x^4 \left(-\frac{c_1}{4x^4} - \frac{5}{4x^2} \right) + c_2 x^4$$

which simplifies to

$$y = -\frac{5}{4}x^2 - \frac{1}{4}c_1 + c_2x^4$$

Summary

The solution(s) found are the following

$$y = -\frac{5}{4}x^2 - \frac{1}{4}c_1 + c_2x^4 \quad (1)$$

Verification of solutions

$$y = -\frac{5}{4}x^2 - \frac{1}{4}c_1 + c_2x^4$$

Verified OK.

8.20.5 Solving using Kovacic algorithm

Writing the ode as

$$xy'' - 3y' = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= x \\ B &= -3 \\ C &= 0 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{15}{4x^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 15 \\ t &= 4x^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{15}{4x^2} \right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 173: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = \frac{15}{4x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = \frac{15}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{5}{2} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = -\frac{3}{2} \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = \frac{15}{4x^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = \frac{15}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{5}{2} \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = -\frac{3}{2} \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{15}{4x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	$\frac{5}{2}$	$-\frac{3}{2}$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	$\frac{5}{2}$	$-\frac{3}{2}$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω .

Trying $\alpha_{\infty}^{-} = -\frac{3}{2}$ then

$$\begin{aligned} d &= \alpha_{\infty}^{-} - (\alpha_{c_1}^{-}) \\ &= -\frac{3}{2} - \left(-\frac{3}{2}\right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_{\infty}$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^{-}}{x - c_1} \right) + (-)[\sqrt{r}]_{\infty} \\ &= -\frac{3}{2x} + (-)(0) \\ &= -\frac{3}{2x} \\ &= -\frac{3}{2x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2\left(-\frac{3}{2x}\right)(0) + \left(\left(\frac{3}{2x^2}\right) + \left(-\frac{3}{2x}\right)^2 - \left(\frac{15}{4x^2}\right)\right) &= 0 \\ 0 &= 0 \end{aligned}$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= p e^{\int \omega dx} \\ &= e^{\int -\frac{3}{2x} dx} \\ &= \frac{1}{x^{\frac{3}{2}}} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\&= z_1 e^{-\int \frac{1}{2} \frac{-3}{x} dx} \\&= z_1 e^{\frac{3 \ln(x)}{2}} \\&= z_1 \left(x^{\frac{3}{2}}\right)\end{aligned}$$

Which simplifies to

$$y_1 = 1$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{-3}{x} dx}}{(y_1)^2} dx \\&= y_1 \int \frac{e^{3 \ln(x)}}{(y_1)^2} dx \\&= y_1 \left(\frac{x^4}{4}\right)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\&= c_1(1) + c_2 \left(1 \left(\frac{x^4}{4}\right)\right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$.
 y_h is the solution to

$$xy'' - 3y' = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 + \frac{c_2 x^4}{4}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = 1$$

$$y_2 = \frac{x^4}{4}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} 1 & \frac{x^4}{4} \\ \frac{d}{dx}(1) & \frac{d}{dx}\left(\frac{x^4}{4}\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} 1 & \frac{x^4}{4} \\ 0 & x^3 \end{vmatrix}$$

Therefore

$$W = (1)(x^3) - \left(\frac{x^4}{4}\right) (0)$$

Which simplifies to

$$W = x^3$$

Which simplifies to

$$W = x^3$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{5x^5}{x^4} dx$$

Which simplifies to

$$u_1 = - \int \frac{5x}{4} dx$$

Hence

$$u_1 = -\frac{5x^2}{8}$$

And Eq. (3) becomes

$$u_2 = \int \frac{5x}{x^4} dx$$

Which simplifies to

$$u_2 = \int \frac{5}{x^3} dx$$

Hence

$$u_2 = -\frac{5}{2x^2}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{5x^2}{4}$$

Therefore the general solution is

$$\begin{aligned}y &= y_h + y_p \\ &= \left(c_1 + \frac{c_2 x^4}{4} \right) + \left(-\frac{5x^2}{4} \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 + \frac{1}{4}c_2x^4 - \frac{5}{4}x^2 \quad (1)$$

Verification of solutions

$$y = c_1 + \frac{1}{4}c_2x^4 - \frac{5}{4}x^2$$

Verified OK.

8.20.6 Solving as exact linear second order ode

An ode of the form

$$p(x) y'' + q(x) y' + r(x) y = s(x)$$

is exact if

$$p''(x) - q'(x) + r(x) = 0 \quad (1)$$

For the given ode we have

$$\begin{aligned}p(x) &= x \\ q(x) &= -3 \\ r(x) &= 0 \\ s(x) &= 5x\end{aligned}$$

Hence

$$\begin{aligned}p''(x) &= 0 \\ q'(x) &= 0\end{aligned}$$

Therefore (1) becomes

$$0 - (0) + (0) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(x)y' + (q(x) - p'(x))y)' = s(x)$$

Integrating gives

$$p(x)y' + (q(x) - p'(x))y = \int s(x) dx$$

Substituting the above values for p, q, r, s gives

$$xy' - 4y = \int 5x dx$$

We now have a first order ode to solve which is

$$xy' - 4y = \frac{5x^2}{2} + c_1$$

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -\frac{4}{x}$$
$$q(x) = \frac{5x^2 + 2c_1}{2x}$$

Hence the ode is

$$y' - \frac{4y}{x} = \frac{5x^2 + 2c_1}{2x}$$

The integrating factor μ is

$$\mu = e^{\int -\frac{4}{x} dx}$$
$$= \frac{1}{x^4}$$

The ode becomes

$$\frac{d}{dx}(\mu y) = (\mu) \left(\frac{5x^2 + 2c_1}{2x} \right)$$
$$\frac{d}{dx} \left(\frac{y}{x^4} \right) = \left(\frac{1}{x^4} \right) \left(\frac{5x^2 + 2c_1}{2x} \right)$$
$$d \left(\frac{y}{x^4} \right) = \left(\frac{5x^2 + 2c_1}{2x^5} \right) dx$$

Integrating gives

$$\frac{y}{x^4} = \int \frac{5x^2 + 2c_1}{2x^5} dx$$
$$\frac{y}{x^4} = -\frac{c_1}{4x^4} - \frac{5}{4x^2} + c_2$$

Dividing both sides by the integrating factor $\mu = \frac{1}{x^4}$ results in

$$y = x^4 \left(-\frac{c_1}{4x^4} - \frac{5}{4x^2} \right) + c_2 x^4$$

which simplifies to

$$y = -\frac{5}{4}x^2 - \frac{1}{4}c_1 + c_2x^4$$

Summary

The solution(s) found are the following

$$y = -\frac{5}{4}x^2 - \frac{1}{4}c_1 + c_2x^4 \quad (1)$$

Verification of solutions

$$y = -\frac{5}{4}x^2 - \frac{1}{4}c_1 + c_2x^4$$

Verified OK.

8.20.7 Maple step by step solution

Let's solve

$$y''x - 3y' = 5x$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Make substitution $u = y'$ to reduce order of ODE

$$u'(x)x - 3u(x) = 5x$$

- Isolate the derivative

$$u'(x) = 5 + \frac{3u(x)}{x}$$

- Group terms with $u(x)$ on the lhs of the ODE and the rest on the rhs of the ODE

$$u'(x) - \frac{3u(x)}{x} = 5$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) \left(u'(x) - \frac{3u(x)}{x} \right) = 5\mu(x)$$
- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x) u(x))$

$$\mu(x) \left(u'(x) - \frac{3u(x)}{x} \right) = \mu'(x) u(x) + \mu(x) u'(x)$$
- Isolate $\mu'(x)$

$$\mu'(x) = -\frac{3\mu(x)}{x}$$
- Solve to find the integrating factor

$$\mu(x) = \frac{1}{x^3}$$
- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x) u(x)) \right) dx = \int 5\mu(x) dx + c_1$$
- Evaluate the integral on the lhs

$$\mu(x) u(x) = \int 5\mu(x) dx + c_1$$
- Solve for $u(x)$

$$u(x) = \frac{\int 5\mu(x) dx + c_1}{\mu(x)}$$
- Substitute $\mu(x) = \frac{1}{x^3}$

$$u(x) = x^3 \left(\int \frac{5}{x^3} dx + c_1 \right)$$
- Evaluate the integrals on the rhs

$$u(x) = x^3 \left(-\frac{5}{2x^2} + c_1 \right)$$
- Simplify

$$u(x) = -\frac{5}{2}x + c_1x^3$$
- Solve 1st ODE for $u(x)$

$$u(x) = -\frac{5}{2}x + c_1x^3$$
- Make substitution $u = y'$

$$y' = -\frac{5}{2}x + c_1x^3$$
- Integrate both sides to solve for y

$$\int y' dx = \int \left(-\frac{5}{2}x + c_1x^3 \right) dx + c_2$$
- Compute integrals

$$y = \frac{(2c_1x^2 - 5)^2}{16c_1} + c_2$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
-> Calling odsolve with the ODE`, diff(_b(_a), _a) = (3*_b(_a)+5*_a)/_a, _b(_a)` *** Suble  
  Methods for first order ODEs:  
  --- Trying classification methods ---  
  trying a quadrature  
  trying 1st order linear  
  <- 1st order linear successful  
<- high order exact linear fully integrable successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(x*diff(y(x),x$2)-3*diff(y(x),x)=5*x,y(x), singsol=all)
```

$$y(x) = \frac{(2c_1x^2 - 5)^2}{16c_1} + c_2$$

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 24

```
DSolve[x*y'[x]-3*y'[x]==5*x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_1x^4}{4} - \frac{5x^2}{4} + c_2$$

9 Chapter 2. Second-Order Linear Equations.
Section 2.1. Linear Equations with Constant
Coefficients. Page 62

9.1	problem 1(a)	1282
9.2	problem 1(b)	1290
9.3	problem 1(c)	1299
9.4	problem 1(d)	1309
9.5	problem 1(e)	1317
9.6	problem 1(f)	1326
9.7	problem 1(g)	1334
9.8	problem 1(h)	1343
9.9	problem 1(i)	1352
9.10	problem 1(j)	1362
9.11	problem 1(k)	1370
9.12	problem 1(l)	1379
9.13	problem 1(m)	1387
9.14	problem 1(n)	1397
9.15	problem 1(o)	1406
9.16	problem 1(p)	1414
9.17	problem 1(q)	1422
9.18	problem 1(r)	1430
9.19	problem 2(a)	1438
9.20	problem 2(b)	1448
9.21	problem 2(c)	1458
9.22	problem 2(d)	1470
9.23	problem 2(e)	1480
9.24	problem 2(f)	1490
9.25	problem 5(a)	1500
9.26	problem 5(b)	1517
9.27	problem 5(c)	1533
9.28	problem 5(d)	1547
9.29	problem 5(e)	1557
9.30	problem 5(f)	1573
9.31	problem 5(g)	1587
9.32	problem 5(h)	1605
9.33	problem 5(i)	1622

9.1 problem 1(a)

- 9.1.1 Solving as second order linear constant coeff ode 1282
- 9.1.2 Solving using Kovacic algorithm 1284
- 9.1.3 Maple step by step solution 1288

Internal problem ID [6269]

Internal file name [OUTPUT/5517_Sunday_June_05_2022_03_42_31_PM_66854464/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.1. Linear Equations with Constant Coefficients. Page 62

Problem number: 1(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' + y' - 6y = 0$$

9.1.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 1, C = -6$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + \lambda e^{\lambda x} - 6 e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + \lambda - 6 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 1, C = -6$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{-1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{1^2 - (4)(1)(-6)} \\ &= -\frac{1}{2} \pm \frac{5}{2}\end{aligned}$$

Hence

$$\lambda_1 = -\frac{1}{2} + \frac{5}{2}$$

$$\lambda_2 = -\frac{1}{2} - \frac{5}{2}$$

Which simplifies to

$$\lambda_1 = 2$$

$$\lambda_2 = -3$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(2)x} + c_2 e^{(-3)x}$$

Or

$$y = c_1 e^{2x} + c_2 e^{-3x}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{2x} + c_2 e^{-3x} \tag{1}$$




Figure 245: Slope field plot

Verification of solutions

$$y = c_1 e^{2x} + c_2 e^{-3x}$$

Verified OK.

9.1.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y' - 6y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 1 \\ C &= -6 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{25}{4} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 25 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \frac{25z(x)}{4} \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 175: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{25}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{5x}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{1}{1} dx} \\ &= z_1 e^{-\frac{x}{2}} \\ &= z_1 \left(e^{-\frac{x}{2}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-3x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{1}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-x}}{(y_1)^2} dx \\ &= y_1 \left(\frac{e^{5x}}{5} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-3x}) + c_2 \left(e^{-3x} \left(\frac{e^{5x}}{5} \right) \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-3x} + \frac{c_2 e^{2x}}{5} \quad (1)$$




Figure 246: Slope field plot

Verification of solutions

$$y = c_1 e^{-3x} + \frac{c_2 e^{2x}}{5}$$

Verified OK.

9.1.3 Maple step by step solution

Let's solve

$$y'' + y' - 6y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 + r - 6 = 0$$

- Factor the characteristic polynomial

$$(r + 3)(r - 2) = 0$$

- Roots of the characteristic polynomial

$$r = (-3, 2)$$

- 1st solution of the ODE

$$y_1(x) = e^{-3x}$$

- 2nd solution of the ODE

$$y_2(x) = e^{2x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = c_1 e^{-3x} + c_2 e^{2x}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)+diff(y(x),x)-6*y(x)=0,y(x), singsol=all)
```

$$y(x) = (e^{5x}c_1 + c_2) e^{-3x}$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 22

```
DSolve[y''[x]+y'[x]-6*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-3x}(c_2e^{5x} + c_1)$$

9.2 problem 1(b)

9.2.1	Solving as second order linear constant coeff ode	1290
9.2.2	Solving as linear second order ode solved by an integrating factor ode	1292
9.2.3	Solving using Kovacic algorithm	1293
9.2.4	Maple step by step solution	1297

Internal problem ID [6270]

Internal file name [OUTPUT/5518_Sunday_June_05_2022_03_42_32_PM_57703630/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.1. Linear Equations with Constant Coefficients. Page 62

Problem number: 1(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "linear_second_order_ode_solved_by_an_integrating_factor"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' + 2y' + y = 0$$

9.2.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 2, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 2\lambda e^{\lambda x} + e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 2\lambda + 1 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 2, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-2}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{(2)^2 - (4)(1)(1)} \\ &= -1 \end{aligned}$$

Hence this is the case of a double root $\lambda_{1,2} = 1$. Therefore the solution is

$$y = c_1 e^{-x} + c_2 x e^{-x} \quad (1)$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-x} + x e^{-x} c_2 \quad (1)$$




Figure 247: Slope field plot

Verification of solutions

$$y = c_1 e^{-x} + x e^{-x} c_2$$

Verified OK.

9.2.2 Solving as linear second order ode solved by an integrating factor ode

The ode satisfies this form

$$y'' + p(x)y' + \frac{(p(x)^2 + p'(x))y}{2} = f(x)$$

Where $p(x) = 2$. Therefore, there is an integrating factor given by

$$\begin{aligned}M(x) &= e^{\frac{1}{2} \int p dx} \\ &= e^{\int 2 dx} \\ &= e^x\end{aligned}$$

Multiplying both sides of the ODE by the integrating factor $M(x)$ makes the left side of the ODE a complete differential

$$\begin{aligned}(M(x)y)'' &= 0 \\ (ye^x)'' &= 0\end{aligned}$$

Integrating once gives

$$(ye^x)' = c_1$$

Integrating again gives

$$ye^x = c_1x + c_2$$

Hence the solution is

$$y = \frac{c_1x + c_2}{e^x}$$

Or

$$y = c_1x e^{-x} + c_2e^{-x}$$

Summary

The solution(s) found are the following

$$y = c_1x e^{-x} + c_2e^{-x} \tag{1}$$




Figure 248: Slope field plot

Verification of solutions

$$y = c_1 x e^{-x} + c_2 e^{-x}$$

Verified OK.

9.2.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 2y' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 2 \\ C &= 1 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} = \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \tag{6}$$

Comparing the above to (5) shows that

$$s = 0$$

$$t = 1$$

Therefore eq. (4) becomes

$$z''(x) = 0 \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 177: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty\end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{2}{1} dx} \\ &= z_1 e^{-x} \\ &= z_1 (e^{-x})\end{aligned}$$

Which simplifies to

$$y_1 = e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{2}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-2x}}{(y_1)^2} dx \\ &= y_1(x)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1(e^{-x}) + c_2(e^{-x}(x))\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-x} + x e^{-x} c_2 \tag{1}$$




Figure 249: Slope field plot

Verification of solutions

$$y = c_1 e^{-x} + x e^{-x} c_2$$

Verified OK.

9.2.4 Maple step by step solution

Let's solve

$$y'' + 2y' + y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 + 2r + 1 = 0$$

- Factor the characteristic polynomial

$$(r + 1)^2 = 0$$

- Root of the characteristic polynomial

$$r = -1$$

- 1st solution of the ODE

$$y_1(x) = e^{-x}$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = x e^{-x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = c_1 e^{-x} + x e^{-x} c_2$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 14

```
dsolve(diff(y(x),x$2)+2*diff(y(x),x)+y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{-x}(c_2x + c_1)$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 18

```
DSolve[y''[x]+2*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x}(c_2x + c_1)$$

9.3 problem 1(c)

9.3.1	Solving as second order linear constant coeff ode	1299
9.3.2	Solving as second order ode can be made integrable ode	1301
9.3.3	Solving using Kovacic algorithm	1303
9.3.4	Maple step by step solution	1307

Internal problem ID [6271]

Internal file name [OUTPUT/5519_Sunday_June_05_2022_03_42_33_PM_3420271/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.1. Linear Equations with Constant Coefficients. Page 62

Problem number: 1(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "second_order_ode_can_be_made_integrable"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x]]
```

$$y'' + 8y = 0$$

9.3.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 8$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 8e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 8 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 8$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(8)} \\ &= \pm 2i\sqrt{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +2i\sqrt{2} \\ \lambda_2 &= -2i\sqrt{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= 2i\sqrt{2} \\ \lambda_2 &= -2i\sqrt{2} \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 2\sqrt{2}$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (c_1 \cos(2x\sqrt{2}) + c_2 \sin(2x\sqrt{2}))$$

Or

$$y = c_1 \cos(2x\sqrt{2}) + c_2 \sin(2x\sqrt{2})$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(2x\sqrt{2}) + c_2 \sin(2x\sqrt{2}) \quad (1)$$




Figure 250: Slope field plot

Verification of solutions

$$y = c_1 \cos(2x\sqrt{2}) + c_2 \sin(2x\sqrt{2})$$

Verified OK.

9.3.2 Solving as second order ode can be made integrable ode

Multiplying the ode by y' gives

$$y'y'' + 8y'y = 0$$

Integrating the above w.r.t x gives

$$\int (y'y'' + 8y'y) dx = 0$$

$$\frac{y'^2}{2} + 4y^2 = c_2$$

Which is now solved for y . Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \sqrt{-8y^2 + 2c_1} \tag{1}$$

$$y' = -\sqrt{-8y^2 + 2c_1} \tag{2}$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int \frac{1}{\sqrt{-8y^2 + 2c_1}} dy = \int dx$$
$$\frac{\sqrt{2} \arctan\left(\frac{2\sqrt{2}y}{\sqrt{-8y^2 + 2c_1}}\right)}{4} = c_2 + x$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{1}{\sqrt{-8y^2 + 2c_1}} dy = \int dx$$
$$-\frac{\sqrt{2} \arctan\left(\frac{2\sqrt{2}y}{\sqrt{-8y^2 + 2c_1}}\right)}{4} = x + c_3$$

Summary

The solution(s) found are the following

$$\frac{\sqrt{2} \arctan\left(\frac{2\sqrt{2}y}{\sqrt{-8y^2 + 2c_1}}\right)}{4} = c_2 + x \quad (1)$$

$$-\frac{\sqrt{2} \arctan\left(\frac{2\sqrt{2}y}{\sqrt{-8y^2 + 2c_1}}\right)}{4} = x + c_3 \quad (2)$$




Figure 251: Slope field plot

Verification of solutions

$$\frac{\sqrt{2} \arctan\left(\frac{2\sqrt{2}y}{\sqrt{-8y^2+2c_1}}\right)}{4} = c_2 + x$$

Verified OK.

$$-\frac{\sqrt{2} \arctan\left(\frac{2\sqrt{2}y}{\sqrt{-8y^2+2c_1}}\right)}{4} = x + c_3$$

Verified OK.

9.3.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 8y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= 1 \\B &= 0 \\C &= 8\end{aligned}\tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x)\tag{4}$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\&= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned}\tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-8}{1}\tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= -8 \\t &= 1\end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -8z(x)\tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 179: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -8$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(2x\sqrt{2})$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}
 y_1 &= z_1 \\
 &= \cos(2x\sqrt{2})
 \end{aligned}$$

Which simplifies to

$$y_1 = \cos(2x\sqrt{2})$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(2x\sqrt{2}) \int \frac{1}{\cos(2x\sqrt{2})^2} dx \\ &= \cos(2x\sqrt{2}) \left(\frac{\sqrt{2} \tan(2x\sqrt{2})}{4} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(\cos(2x\sqrt{2}) \right) + c_2 \left(\cos(2x\sqrt{2}) \left(\frac{\sqrt{2} \tan(2x\sqrt{2})}{4} \right) \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(2x\sqrt{2}) + \frac{c_2 \sqrt{2} \sin(2x\sqrt{2})}{4} \quad (1)$$




Figure 252: Slope field plot

Verification of solutions

$$y = c_1 \cos(2x\sqrt{2}) + \frac{c_2\sqrt{2} \sin(2x\sqrt{2})}{4}$$

Verified OK.

9.3.4 Maple step by step solution

Let's solve

$$y'' + 8y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 + 8 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-32})}{2}$$

- Roots of the characteristic polynomial

$$r = (-2i\sqrt{2}, 2i\sqrt{2})$$

- 1st solution of the ODE

$$y_1(x) = \cos(2x\sqrt{2})$$

- 2nd solution of the ODE

$$y_2(x) = \sin(2x\sqrt{2})$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = c_1 \cos(2x\sqrt{2}) + c_2 \sin(2x\sqrt{2})$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(diff(y(x),x$2)+8*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 \sin(2\sqrt{2}x) + c_2 \cos(2\sqrt{2}x)$$

✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 30

```
DSolve[y''[x]+8*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos(2\sqrt{2}x) + c_2 \sin(2\sqrt{2}x)$$

9.4 problem 1(d)

- 9.4.1 Solving as second order linear constant coeff ode 1309
- 9.4.2 Solving using Kovacic algorithm 1311
- 9.4.3 Maple step by step solution 1315

Internal problem ID [6272]

Internal file name [OUTPUT/5520_Sunday_June_05_2022_03_42_34_PM_15542383/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.1. Linear Equations with Constant Coefficients. Page 62

Problem number: 1(d).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$2y'' - 4y' + 4y = 0$$

9.4.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 2, B = -4, C = 4$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$2\lambda^2 e^{\lambda x} - 4\lambda e^{\lambda x} + 4e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$2\lambda^2 - 4\lambda + 4 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 2, B = -4, C = 4$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{4}{(2)(2)} \pm \frac{1}{(2)(2)} \sqrt{-4^2 - (4)(2)(4)} \\ &= 1 \pm i\end{aligned}$$

Hence

$$\lambda_1 = 1 + i$$

$$\lambda_2 = 1 - i$$

Which simplifies to

$$\lambda_1 = 1 + i$$

$$\lambda_2 = 1 - i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 1$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^x(\cos(x) c_1 + c_2 \sin(x))$$

Summary

The solution(s) found are the following

$$y = e^x(\cos(x) c_1 + c_2 \sin(x)) \quad (1)$$




Figure 253: Slope field plot

Verification of solutions

$$y = e^x (\cos(x) c_1 + c_2 \sin(x))$$

Verified OK.

9.4.2 Solving using Kovacic algorithm

Writing the ode as

$$2y'' - 4y' + 4y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 2 \\ B &= -4 \\ C &= 4 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 181: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0\end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-4}{2} dx} \\ &= z_1 e^x \\ &= z_1(e^x)\end{aligned}$$

Which simplifies to

$$y_1 = \cos(x) e^x$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{-4}{2} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{2x}}{(y_1)^2} dx \\ &= y_1(\tan(x))\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1(\cos(x) e^x) + c_2(\cos(x) e^x (\tan(x)))\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(x) e^x + c_2 \sin(x) e^x \quad (1)$$




Figure 254: Slope field plot

Verification of solutions

$$y = c_1 \cos(x) e^x + c_2 \sin(x) e^x$$

Verified OK.

9.4.3 Maple step by step solution

Let's solve

$$2y'' - 4y' + 4y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = 2y' - 2y$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - 2y' + 2y = 0$$

- Characteristic polynomial of ODE

$$r^2 - 2r + 2 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{2 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (1 - I, 1 + I)$$

- 1st solution of the ODE

$$y_1(x) = \cos(x) e^x$$

- 2nd solution of the ODE

$$y_2(x) = \sin(x) e^x$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = c_1 \cos(x) e^x + c_2 \sin(x) e^x$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```


✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 16

```
dsolve(2*diff(y(x),x$2)-4*diff(y(x),x)+4*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^x(c_1 \sin(x) + \cos(x) c_2)$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 20

```
DSolve[2*y''[x]-4*y'[x]+4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x(c_2 \cos(x) + c_1 \sin(x))$$

9.5 problem 1(e)

9.5.1	Solving as second order linear constant coeff ode	1317
9.5.2	Solving as linear second order ode solved by an integrating factor ode	1319
9.5.3	Solving using Kovacic algorithm	1320
9.5.4	Maple step by step solution	1324

Internal problem ID [6273]

Internal file name [OUTPUT/5521_Sunday_June_05_2022_03_42_36_PM_76880388/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.1. Linear Equations with Constant Coefficients. Page 62

Problem number: 1(e).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "linear_second_order_ode_solved_by_an_integrating_factor"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' - 4y' + 4y = 0$$

9.5.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -4, C = 4$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 4\lambda e^{\lambda x} + 4e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 4\lambda + 4 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -4, C = 4$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{4}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{(-4)^2 - (4)(1)(4)} \\ &= 2 \end{aligned}$$

Hence this is the case of a double root $\lambda_{1,2} = -2$. Therefore the solution is

$$y = c_1 e^{2x} + c_2 e^{2x} x \tag{1}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{2x} + x e^{2x} c_2 \tag{1}$$




Figure 255: Slope field plot

Verification of solutions

$$y = c_1 e^{2x} + x e^{2x} c_2$$

Verified OK.

9.5.2 Solving as linear second order ode solved by an integrating factor ode

The ode satisfies this form

$$y'' + p(x)y' + \frac{(p(x)^2 + p'(x))y}{2} = f(x)$$

Where $p(x) = -4$. Therefore, there is an integrating factor given by

$$\begin{aligned}M(x) &= e^{\frac{1}{2} \int p dx} \\ &= e^{\int -4 dx} \\ &= e^{-2x}\end{aligned}$$

Multiplying both sides of the ODE by the integrating factor $M(x)$ makes the left side of the ODE a complete differential

$$\begin{aligned}(M(x)y)'' &= 0 \\ (ye^{-2x})'' &= 0\end{aligned}$$

Integrating once gives

$$(ye^{-2x})' = c_1$$

Integrating again gives

$$(ye^{-2x}) = c_1x + c_2$$

Hence the solution is

$$y = \frac{c_1x + c_2}{e^{-2x}}$$

Or

$$y = c_1x e^{2x} + c_2e^{2x}$$

Summary

The solution(s) found are the following

$$y = c_1x e^{2x} + c_2e^{2x} \tag{1}$$




Figure 256: Slope field plot

Verification of solutions

$$y = c_1 x e^{2x} + c_2 e^{2x}$$

Verified OK.

9.5.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 4y' + 4y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -4 \\ C &= 4 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 0 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 0 \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 183: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-4}{1} dx} \\ &= z_1 e^{2x} \\ &= z_1 (e^{2x}) \end{aligned}$$

Which simplifies to

$$y_1 = e^{2x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-4}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{4x}}{(y_1)^2} dx \\ &= y_1(x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{2x}) + c_2 (e^{2x}(x))\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{2x} + x e^{2x} c_2 \tag{1}$$




Figure 257: Slope field plot

Verification of solutions

$$y = c_1 e^{2x} + x e^{2x} c_2$$

Verified OK.

9.5.4 Maple step by step solution

Let's solve

$$y'' - 4y' + 4y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 - 4r + 4 = 0$$

- Factor the characteristic polynomial

$$(r - 2)^2 = 0$$

- Root of the characteristic polynomial

$$r = 2$$

- 1st solution of the ODE

$$y_1(x) = e^{2x}$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = e^{2x}x$$

- General solution of the ODE

$$y = c_1y_1(x) + c_2y_2(x)$$

- Substitute in solutions

$$y = c_1e^{2x} + xe^{2x}c_2$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(diff(y(x),x$2)-4*diff(y(x),x)+4*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{2x}(c_2x + c_1)$$

✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 18

```
DSolve[y''[x]-4*y'[x]+4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{2x}(c_2x + c_1)$$

9.6 problem 1(f)

- 9.6.1 Solving as second order linear constant coeff ode 1326
- 9.6.2 Solving using Kovacic algorithm 1328
- 9.6.3 Maple step by step solution 1332

Internal problem ID [6274]

Internal file name [OUTPUT/5522_Sunday_June_05_2022_03_42_37_PM_46056091/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.1. Linear Equations with Constant Coefficients. Page 62

Problem number: 1(f).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' - 9y' + 20y = 0$$

9.6.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -9, C = 20$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 9\lambda e^{\lambda x} + 20 e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 9\lambda + 20 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -9, C = 20$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{9}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-9^2 - (4)(1)(20)} \\ &= \frac{9}{2} \pm \frac{1}{2}\end{aligned}$$

Hence

$$\lambda_1 = \frac{9}{2} + \frac{1}{2}$$

$$\lambda_2 = \frac{9}{2} - \frac{1}{2}$$

Which simplifies to

$$\lambda_1 = 5$$

$$\lambda_2 = 4$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(5)x} + c_2 e^{(4)x}$$

Or

$$y = c_1 e^{5x} + c_2 e^{4x}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{5x} + c_2 e^{4x} \tag{1}$$




Figure 258: Slope field plot

Verification of solutions

$$y = c_1 e^{5x} + c_2 e^{4x}$$

Verified OK.

9.6.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 9y' + 20y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = -9 \tag{3}$$

$$C = 20$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{1}{4} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 1 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \frac{z(x)}{4} \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 185: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0\end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{1}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{x}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-9}{1} dx} \\ &= z_1 e^{\frac{9x}{2}} \\ &= z_1 \left(e^{\frac{9x}{2}} \right)\end{aligned}$$

Which simplifies to

$$y_1 = e^{4x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{-9}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{9x}}{(y_1)^2} dx \\ &= y_1(e^x)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{4x}) + c_2 (e^{4x}(e^x))\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{4x} + c_2 e^{5x} \tag{1}$$




Figure 259: Slope field plot

Verification of solutions

$$y = c_1 e^{4x} + c_2 e^{5x}$$

Verified OK.

9.6.3 Maple step by step solution

Let's solve

$$y'' - 9y' + 20y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 - 9r + 20 = 0$$

- Factor the characteristic polynomial

$$(r - 4)(r - 5) = 0$$

- Roots of the characteristic polynomial

$$r = (4, 5)$$

- 1st solution of the ODE

$$y_1(x) = e^{4x}$$

- 2nd solution of the ODE

$$y_2(x) = e^{5x}$$

- General solution of the ODE

$$y = c_1y_1(x) + c_2y_2(x)$$

- Substitute in solutions

$$y = c_1e^{4x} + c_2e^{5x}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)-9*diff(y(x),x)+20*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{4x}c_1 + c_2e^{5x}$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 20

```
DSolve[y''[x]-9*y'[x]+20*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{4x}(c_2e^x + c_1)$$

9.7 problem 1(g)

9.7.1 Solving as second order linear constant coeff ode	1334
9.7.2 Solving using Kovacic algorithm	1336
9.7.3 Maple step by step solution	1340

Internal problem ID [6275]

Internal file name [OUTPUT/5523_Sunday_June_05_2022_03_42_38_PM_47340201/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.1. Linear Equations with Constant Coefficients. Page 62

Problem number: 1(g).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$2y'' + 2y' + 3y = 0$$

9.7.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 2, B = 2, C = 3$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$2\lambda^2 e^{\lambda x} + 2\lambda e^{\lambda x} + 3e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$2\lambda^2 + 2\lambda + 3 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 2, B = 2, C = 3$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{-2}{(2)(2)} \pm \frac{1}{(2)(2)} \sqrt{2^2 - (4)(2)(3)} \\ &= -\frac{1}{2} \pm \frac{i\sqrt{5}}{2}\end{aligned}$$

Hence

$$\begin{aligned}\lambda_1 &= -\frac{1}{2} + \frac{i\sqrt{5}}{2} \\ \lambda_2 &= -\frac{1}{2} - \frac{i\sqrt{5}}{2}\end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= -\frac{1}{2} + \frac{i\sqrt{5}}{2} \\ \lambda_2 &= -\frac{1}{2} - \frac{i\sqrt{5}}{2}\end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = -\frac{1}{2}$ and $\beta = \frac{\sqrt{5}}{2}$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{x\sqrt{5}}{2} \right) + c_2 \sin \left(\frac{x\sqrt{5}}{2} \right) \right)$$

Summary

The solution(s) found are the following

$$y = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{x\sqrt{5}}{2} \right) + c_2 \sin \left(\frac{x\sqrt{5}}{2} \right) \right) \quad (1)$$




Figure 260: Slope field plot

Verification of solutions

$$y = e^{-\frac{ix}{2}} \left(c_1 \cos \left(\frac{x\sqrt{5}}{2} \right) + c_2 \sin \left(\frac{x\sqrt{5}}{2} \right) \right)$$

Verified OK.

9.7.2 Solving using Kovacic algorithm

Writing the ode as

$$2y'' + 2y' + 3y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 2 \\ B &= 2 \\ C &= 3 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-5}{4} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -5 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -\frac{5z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 187: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0\end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -\frac{5}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos\left(\frac{x\sqrt{5}}{2}\right)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{2}{2} dx} \\ &= z_1 e^{-\frac{x}{2}} \\ &= z_1 \left(e^{-\frac{x}{2}}\right)\end{aligned}$$

Which simplifies to

$$y_1 = e^{-\frac{x}{2}} \cos\left(\frac{x\sqrt{5}}{2}\right)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{2}{2} dx}}{(y_1)^2} dx \\&= y_1 \int \frac{e^{-x}}{(y_1)^2} dx \\&= y_1 \left(\frac{2\sqrt{5} \tan\left(\frac{x\sqrt{5}}{2}\right)}{5} \right)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\&= c_1 \left(e^{-\frac{x}{2}} \cos\left(\frac{x\sqrt{5}}{2}\right) \right) + c_2 \left(e^{-\frac{x}{2}} \cos\left(\frac{x\sqrt{5}}{2}\right) \left(\frac{2\sqrt{5} \tan\left(\frac{x\sqrt{5}}{2}\right)}{5} \right) \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-\frac{x}{2}} \cos\left(\frac{x\sqrt{5}}{2}\right) + \frac{2c_2 e^{-\frac{x}{2}} \sqrt{5} \sin\left(\frac{x\sqrt{5}}{2}\right)}{5} \quad (1)$$




Figure 261: Slope field plot

Verification of solutions

$$y = c_1 e^{-\frac{x}{2}} \cos\left(\frac{x\sqrt{5}}{2}\right) + \frac{2c_2 e^{-\frac{x}{2}} \sqrt{5} \sin\left(\frac{x\sqrt{5}}{2}\right)}{5}$$

Verified OK.

9.7.3 Maple step by step solution

Let's solve

$$2y'' + 2y' + 3y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -y' - \frac{3y}{2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + y' + \frac{3y}{2} = 0$$

- Characteristic polynomial of ODE

$$r^2 + r + \frac{3}{2} = 0$$
- Use quadratic formula to solve for r

$$r = \frac{(-1) \pm (\sqrt{-5})}{2}$$
- Roots of the characteristic polynomial

$$r = \left(-\frac{1}{2} - \frac{i\sqrt{5}}{2}, -\frac{1}{2} + \frac{i\sqrt{5}}{2} \right)$$
- 1st solution of the ODE

$$y_1(x) = e^{-\frac{x}{2}} \cos\left(\frac{x\sqrt{5}}{2}\right)$$
- 2nd solution of the ODE

$$y_2(x) = e^{-\frac{x}{2}} \sin\left(\frac{x\sqrt{5}}{2}\right)$$
- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$
- Substitute in solutions

$$y = c_1 e^{-\frac{x}{2}} \cos\left(\frac{x\sqrt{5}}{2}\right) + c_2 e^{-\frac{x}{2}} \sin\left(\frac{x\sqrt{5}}{2}\right)$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 28

```
dsolve(2*diff(y(x),x$2)+2*diff(y(x),x)+3*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{-\frac{x}{2}} \left(c_1 \sin\left(\frac{\sqrt{5}x}{2}\right) + c_2 \cos\left(\frac{\sqrt{5}x}{2}\right) \right)$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 42

```
DSolve[2*y''[x]+2*y'[x]+3*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x/2} \left(c_2 \cos \left(\frac{\sqrt{5}x}{2} \right) + c_1 \sin \left(\frac{\sqrt{5}x}{2} \right) \right)$$

9.8 problem 1(h)

9.8.1	Solving as second order linear constant coeff ode	1343
9.8.2	Solving as linear second order ode solved by an integrating factor ode	1345
9.8.3	Solving using Kovacic algorithm	1346
9.8.4	Maple step by step solution	1350

Internal problem ID [6276]

Internal file name [OUTPUT/5524_Sunday_June_05_2022_03_42_39_PM_51360054/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.1. Linear Equations with Constant Coefficients. Page 62

Problem number: 1(h).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "linear_second_order_ode_solved_by_an_integrating_factor"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$4y'' - 12y' + 9y = 0$$

9.8.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 4, B = -12, C = 9$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$4\lambda^2 e^{\lambda x} - 12\lambda e^{\lambda x} + 9 e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$4\lambda^2 - 12\lambda + 9 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 4, B = -12, C = 9$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{12}{(2)(4)} \pm \frac{1}{(2)(4)} \sqrt{(-12)^2 - (4)(4)(9)} \\ &= \frac{3}{2} \end{aligned}$$

Hence this is the case of a double root $\lambda_{1,2} = -\frac{3}{2}$. Therefore the solution is

$$y = c_1 e^{\frac{3x}{2}} + c_2 x e^{\frac{3x}{2}} \quad (1)$$

Summary

The solution(s) found are the following

$$y = c_1 e^{\frac{3x}{2}} + c_2 x e^{\frac{3x}{2}} \quad (1)$$




Figure 262: Slope field plot

Verification of solutions

$$y = c_1 e^{\frac{3x}{2}} + c_2 x e^{\frac{3x}{2}}$$

Verified OK.

9.8.2 Solving as linear second order ode solved by an integrating factor ode

The ode satisfies this form

$$y'' + p(x)y' + \frac{(p(x)^2 + p'(x))y}{2} = f(x)$$

Where $p(x) = -3$. Therefore, there is an integrating factor given by

$$\begin{aligned}M(x) &= e^{\frac{1}{2} \int p dx} \\ &= e^{\int -3 dx} \\ &= e^{-\frac{3x}{2}}\end{aligned}$$

Multiplying both sides of the ODE by the integrating factor $M(x)$ makes the left side of the ODE a complete differential

$$\begin{aligned}(M(x)y)'' &= 0 \\ \left(e^{-\frac{3x}{2}}y\right)'' &= 0\end{aligned}$$

Integrating once gives

$$\left(e^{-\frac{3x}{2}}y\right)' = c_1$$

Integrating again gives

$$\left(e^{-\frac{3x}{2}}y\right) = c_1x + c_2$$

Hence the solution is

$$y = \frac{c_1x + c_2}{e^{-\frac{3x}{2}}}$$

Or

$$y = c_1x e^{\frac{3x}{2}} + c_2 e^{\frac{3x}{2}}$$

Summary

The solution(s) found are the following

$$y = c_1x e^{\frac{3x}{2}} + c_2 e^{\frac{3x}{2}} \quad (1)$$




Figure 263: Slope field plot

Verification of solutions

$$y = c_1 x e^{\frac{3x}{2}} + c_2 e^{\frac{3x}{2}}$$

Verified OK.

9.8.3 Solving using Kovacic algorithm

Writing the ode as

$$4y'' - 12y' + 9y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 4 \\ B &= -12 \\ C &= 9 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} = \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \tag{6}$$

Comparing the above to (5) shows that

$$s = 0$$

$$t = 1$$

Therefore eq. (4) becomes

$$z''(x) = 0 \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 189: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty\end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-12}{4} dx} \\ &= z_1 e^{\frac{3x}{2}} \\ &= z_1 \left(e^{\frac{3x}{2}} \right)\end{aligned}$$

Which simplifies to

$$y_1 = e^{\frac{3x}{2}}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{-12}{4} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{3x}}{(y_1)^2} dx \\ &= y_1(x)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(e^{\frac{3x}{2}} \right) + c_2 \left(e^{\frac{3x}{2}} (x) \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{\frac{3x}{2}} + c_2 x e^{\frac{3x}{2}} \quad (1)$$




Figure 264: Slope field plot

Verification of solutions

$$y = c_1 e^{\frac{3x}{2}} + c_2 x e^{\frac{3x}{2}}$$

Verified OK.

9.8.4 Maple step by step solution

Let's solve

$$4y'' - 12y' + 9y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = 3y' - \frac{9y}{4}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - 3y' + \frac{9y}{4} = 0$$

- Characteristic polynomial of ODE

$$r^2 - 3r + \frac{9}{4} = 0$$

- Factor the characteristic polynomial

$$\frac{(2r-3)^2}{4} = 0$$

- Root of the characteristic polynomial

$$r = \frac{3}{2}$$

- 1st solution of the ODE

$$y_1(x) = e^{\frac{3x}{2}}$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = x e^{\frac{3x}{2}}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = c_1 e^{\frac{3x}{2}} + c_2 x e^{\frac{3x}{2}}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 14

```
dsolve(4*diff(y(x),x$2)-12*diff(y(x),x)+9*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{\frac{3x}{2}}(c_2x + c_1)$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 20

```
DSolve[4*y''[x]-12*y'[x]+9*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{3x/2}(c_2x + c_1)$$

9.9 problem 1(i)

9.9.1	Solving as second order linear constant coeff ode	1352
9.9.2	Solving as second order ode can be made integrable ode	1354
9.9.3	Solving using Kovacic algorithm	1356
9.9.4	Maple step by step solution	1360

Internal problem ID [6277]

Internal file name [OUTPUT/5525_Sunday_June_05_2022_03_42_40_PM_28163000/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.1. Linear Equations with Constant Coefficients. Page 62

Problem number: 1(i).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "second_order_ode_can_be_made_integrable"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x]]
```

$$y'' + y = 0$$

9.9.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +i \\ \lambda_2 &= -i \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= i \\ \lambda_2 &= -i \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (\cos(x) c_1 + c_2 \sin(x))$$

Or

$$y = \cos(x) c_1 + c_2 \sin(x)$$

Summary

The solution(s) found are the following

$$y = \cos(x) c_1 + c_2 \sin(x) \tag{1}$$




Figure 265: Slope field plot

Verification of solutions

$$y = \cos(x) c_1 + c_2 \sin(x)$$

Verified OK.

9.9.2 Solving as second order ode can be made integrable ode

Multiplying the ode by y' gives

$$y' y'' + y' y = 0$$

Integrating the above w.r.t x gives

$$\int (y' y'' + y' y) dx = 0$$

$$\frac{y'^2}{2} + \frac{y^2}{2} = c_2$$

Which is now solved for y . Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \sqrt{-y^2 + 2c_1} \tag{1}$$

$$y' = -\sqrt{-y^2 + 2c_1} \tag{2}$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int \frac{1}{\sqrt{-y^2 + 2c_1}} dy = \int dx$$
$$\arctan\left(\frac{y}{\sqrt{-y^2 + 2c_1}}\right) = c_2 + x$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{1}{\sqrt{-y^2 + 2c_1}} dy = \int dx$$
$$-\arctan\left(\frac{y}{\sqrt{-y^2 + 2c_1}}\right) = x + c_3$$

Summary

The solution(s) found are the following

$$\arctan\left(\frac{y}{\sqrt{-y^2 + 2c_1}}\right) = c_2 + x \tag{1}$$

$$-\arctan\left(\frac{y}{\sqrt{-y^2 + 2c_1}}\right) = x + c_3 \tag{2}$$




Figure 266: Slope field plot

Verification of solutions

$$\arctan\left(\frac{y}{\sqrt{-y^2 + 2c_1}}\right) = c_2 + x$$

Verified OK.

$$-\arctan\left(\frac{y}{\sqrt{-y^2 + 2c_1}}\right) = x + c_3$$

Verified OK.

9.9.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= 1 \\B &= 0 \\C &= 1\end{aligned}\tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x)\tag{4}$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\&= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned}\tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1}\tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= -1 \\t &= 1\end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x)\tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 191: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}
 y_1 &= z_1 \\
 &= \cos(x)
 \end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x)) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1(\cos(x)) + c_2(\cos(x) (\tan(x))) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \cos(x) c_1 + c_2 \sin(x) \tag{1}$$




Figure 267: Slope field plot

Verification of solutions

$$y = \cos(x) c_1 + c_2 \sin(x)$$

Verified OK.

9.9.4 Maple step by step solution

Let's solve

$$y'' + y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

- $r = (-I, I)$
- 1st solution of the ODE
 $y_1(x) = \cos(x)$
- 2nd solution of the ODE
 $y_2(x) = \sin(x)$
- General solution of the ODE
 $y = c_1 y_1(x) + c_2 y_2(x)$
- Substitute in solutions
 $y = \cos(x) c_1 + c_2 \sin(x)$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(diff(y(x),x$2)+y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 \sin(x) + \cos(x) c_2$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 16

```
DSolve[y''[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos(x) + c_2 \sin(x)$$

9.10 problem 1(j)

- 9.10.1 Solving as second order linear constant coeff ode 1362
- 9.10.2 Solving using Kovacic algorithm 1364
- 9.10.3 Maple step by step solution 1368

Internal problem ID [6278]

Internal file name [OUTPUT/5526_Sunday_June_05_2022_03_42_41_PM_11821967/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.1. Linear Equations with Constant Coefficients. Page 62

Problem number: 1(j).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' - 6y' + 25y = 0$$

9.10.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -6, C = 25$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 6\lambda e^{\lambda x} + 25 e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 6\lambda + 25 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -6, C = 25$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{6}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-6^2 - (4)(1)(25)} \\ &= 3 \pm 4i\end{aligned}$$

Hence

$$\lambda_1 = 3 + 4i$$

$$\lambda_2 = 3 - 4i$$

Which simplifies to

$$\lambda_1 = 3 + 4i$$

$$\lambda_2 = 3 - 4i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 3$ and $\beta = 4$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^{3x}(c_1 \cos(4x) + c_2 \sin(4x))$$

Summary

The solution(s) found are the following

$$y = e^{3x}(c_1 \cos(4x) + c_2 \sin(4x)) \quad (1)$$




Figure 268: Slope field plot

Verification of solutions

$$y = e^{3x}(c_1 \cos(4x) + c_2 \sin(4x))$$

Verified OK.

9.10.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 6y' + 25y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -6 \\ C &= 25 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} = \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-16}{1} \tag{6}$$

Comparing the above to (5) shows that

$$s = -16$$

$$t = 1$$

Therefore eq. (4) becomes

$$z''(x) = -16z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 193: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -16$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(4x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-6}{1} dx} \\ &= z_1 e^{3x} \\ &= z_1 (e^{3x}) \end{aligned}$$

Which simplifies to

$$y_1 = e^{3x} \cos(4x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-6}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{6x}}{(y_1)^2} dx \\ &= y_1 \left(\frac{\tan(4x)}{4} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{3x} \cos(4x)) + c_2 \left(e^{3x} \cos(4x) \left(\frac{\tan(4x)}{4} \right) \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{3x} \cos(4x) + \frac{c_2 e^{3x} \sin(4x)}{4} \quad (1)$$




Figure 269: Slope field plot

Verification of solutions

$$y = c_1 e^{3x} \cos(4x) + \frac{c_2 e^{3x} \sin(4x)}{4}$$

Verified OK.

9.10.3 Maple step by step solution

Let's solve

$$y'' - 6y' + 25y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 - 6r + 25 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{6 \pm (\sqrt{-64})}{2}$$

- Roots of the characteristic polynomial

$$r = (3 - 4I, 3 + 4I)$$

- 1st solution of the ODE

$$y_1(x) = e^{3x} \cos(4x)$$

- 2nd solution of the ODE

$$y_2(x) = e^{3x} \sin(4x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = c_1 e^{3x} \cos(4x) + c_2 e^{3x} \sin(4x)$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 22

```
dsolve(diff(y(x),x$2)-6*diff(y(x),x)+25*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{3x}(c_1 \sin(4x) + c_2 \cos(4x))$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 26

```
DSolve[y''[x]-6*y'[x]+25*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{3x}(c_2 \cos(4x) + c_1 \sin(4x))$$

9.11 problem 1(k)

9.11.1 Solving as second order linear constant coeff ode	1370
9.11.2 Solving as linear second order ode solved by an integrating factor ode	1372
9.11.3 Solving using Kovacic algorithm	1373
9.11.4 Maple step by step solution	1377

Internal problem ID [6279]

Internal file name [OUTPUT/5527_Sunday_June_05_2022_03_42_43_PM_67433330/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.1. Linear Equations with Constant Coefficients. Page 62

Problem number: 1(k).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "linear_second_order_ode_solved_by_an_integrating_factor"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$4y'' + 20y' + 25y = 0$$

9.11.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 4, B = 20, C = 25$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$4\lambda^2 e^{\lambda x} + 20\lambda e^{\lambda x} + 25 e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$4\lambda^2 + 20\lambda + 25 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 4, B = 20, C = 25$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-20}{(2)(4)} \pm \frac{1}{(2)(4)} \sqrt{(20)^2 - (4)(4)(25)} \\ &= -\frac{5}{2} \end{aligned}$$

Hence this is the case of a double root $\lambda_{1,2} = \frac{5}{2}$. Therefore the solution is

$$y = c_1 e^{-\frac{5x}{2}} + c_2 x e^{-\frac{5x}{2}} \quad (1)$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-\frac{5x}{2}} + c_2 x e^{-\frac{5x}{2}} \quad (1)$$




Figure 270: Slope field plot

Verification of solutions

$$y = c_1 e^{-\frac{5x}{2}} + c_2 x e^{-\frac{5x}{2}}$$

Verified OK.

9.11.2 Solving as linear second order ode solved by an integrating factor ode

The ode satisfies this form

$$y'' + p(x)y' + \frac{(p(x)^2 + p'(x))y}{2} = f(x)$$

Where $p(x) = 5$. Therefore, there is an integrating factor given by

$$\begin{aligned}M(x) &= e^{\frac{1}{2} \int p dx} \\ &= e^{\int 5 dx} \\ &= e^{\frac{5x}{2}}\end{aligned}$$

Multiplying both sides of the ODE by the integrating factor $M(x)$ makes the left side of the ODE a complete differential

$$\begin{aligned}(M(x)y)'' &= 0 \\ \left(e^{\frac{5x}{2}}y\right)'' &= 0\end{aligned}$$

Integrating once gives

$$\left(e^{\frac{5x}{2}}y\right)' = c_1$$

Integrating again gives

$$\left(e^{\frac{5x}{2}}y\right) = c_1x + c_2$$

Hence the solution is

$$y = \frac{c_1x + c_2}{e^{\frac{5x}{2}}}$$

Or

$$y = c_1x e^{-\frac{5x}{2}} + e^{-\frac{5x}{2}}c_2$$

Summary

The solution(s) found are the following

$$y = c_1x e^{-\frac{5x}{2}} + e^{-\frac{5x}{2}}c_2 \quad (1)$$




Figure 271: Slope field plot

Verification of solutions

$$y = c_1 x e^{-\frac{5x}{2}} + e^{-\frac{5x}{2}} c_2$$

Verified OK.

9.11.3 Solving using Kovacic algorithm

Writing the ode as

$$4y'' + 20y' + 25y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 4 \\ B &= 20 \\ C &= 25 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} = \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \tag{6}$$

Comparing the above to (5) shows that

$$s = 0$$

$$t = 1$$

Therefore eq. (4) becomes

$$z''(x) = 0 \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 195: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{20}{4} dx} \\ &= z_1 e^{-\frac{5x}{2}} \\ &= z_1 \left(e^{-\frac{5x}{2}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-\frac{5x}{2}}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{20}{4} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-5x}}{(y_1)^2} dx \\ &= y_1(x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(e^{-\frac{5x}{2}} \right) + c_2 \left(e^{-\frac{5x}{2}} (x) \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-\frac{5x}{2}} + c_2 x e^{-\frac{5x}{2}} \quad (1)$$




Figure 272: Slope field plot

Verification of solutions

$$y = c_1 e^{-\frac{5x}{2}} + c_2 x e^{-\frac{5x}{2}}$$

Verified OK.

9.11.4 Maple step by step solution

Let's solve

$$4y'' + 20y' + 25y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -5y' - \frac{25y}{4}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + 5y' + \frac{25y}{4} = 0$$

- Characteristic polynomial of ODE

$$r^2 + 5r + \frac{25}{4} = 0$$

- Factor the characteristic polynomial

$$\frac{(2r+5)^2}{4} = 0$$

- Root of the characteristic polynomial

$$r = -\frac{5}{2}$$

- 1st solution of the ODE

$$y_1(x) = e^{-\frac{5x}{2}}$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = x e^{-\frac{5x}{2}}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = c_1 e^{-\frac{5x}{2}} + c_2 x e^{-\frac{5x}{2}}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 14

```
dsolve(4*diff(y(x),x$2)+20*diff(y(x),x)+25*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{-\frac{5x}{2}}(c_2x + c_1)$$

✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 20

```
DSolve[4*y''[x]+20*y'[x]+25*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-5x/2}(c_2x + c_1)$$

9.12 problem 1(1)

9.12.1 Solving as second order linear constant coeff ode	1379
9.12.2 Solving using Kovacic algorithm	1381
9.12.3 Maple step by step solution	1385

Internal problem ID [6280]

Internal file name [OUTPUT/5528_Sunday_June_05_2022_03_42_44_PM_3209467/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.1. Linear Equations with Constant Coefficients. Page 62

Problem number: 1(1).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' + 2y' + 3y = 0$$

9.12.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 2, C = 3$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 2\lambda e^{\lambda x} + 3e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 2\lambda + 3 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 2, C = 3$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{-2}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{2^2 - (4)(1)(3)} \\ &= -1 \pm i\sqrt{2}\end{aligned}$$

Hence

$$\begin{aligned}\lambda_1 &= -1 + i\sqrt{2} \\ \lambda_2 &= -1 - i\sqrt{2}\end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= -1 + i\sqrt{2} \\ \lambda_2 &= -1 - i\sqrt{2}\end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = -1$ and $\beta = \sqrt{2}$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^{-x}(c_1 \cos(x\sqrt{2}) + c_2 \sin(x\sqrt{2}))$$

Summary

The solution(s) found are the following

$$y = e^{-x}(c_1 \cos(x\sqrt{2}) + c_2 \sin(x\sqrt{2})) \quad (1)$$




Figure 273: Slope field plot

Verification of solutions

$$y = e^{-x} \left(c_1 \cos(x\sqrt{2}) + c_2 \sin(x\sqrt{2}) \right)$$

Verified OK.

9.12.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 2y' + 3y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 2 \tag{3}$$

$$C = 3$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-2}{1} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -2 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -2z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 197: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -2$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x\sqrt{2})$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{2}{1} dx} \\ &= z_1 e^{-x} \\ &= z_1 (e^{-x}) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-x} \cos(x\sqrt{2})$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{2}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-2x}}{(y_1)^2} dx \\ &= y_1 \left(\frac{\sqrt{2} \tan(x\sqrt{2})}{2} \right) \end{aligned}$$

Therefore the solution is

$$y = c_1 y_1 + c_2 y_2$$

$$= c_1 \left(e^{-x} \cos(x\sqrt{2}) \right) + c_2 \left(e^{-x} \cos(x\sqrt{2}) \left(\frac{\sqrt{2} \tan(x\sqrt{2})}{2} \right) \right)$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-x} \cos(x\sqrt{2}) + \frac{c_2 e^{-x} \sqrt{2} \sin(x\sqrt{2})}{2} \quad (1)$$




Figure 274: Slope field plot

Verification of solutions

$$y = c_1 e^{-x} \cos(x\sqrt{2}) + \frac{c_2 e^{-x} \sqrt{2} \sin(x\sqrt{2})}{2}$$

Verified OK.

9.12.3 Maple step by step solution

Let's solve

$$y'' + 2y' + 3y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 + 2r + 3 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-2) \pm (\sqrt{-8})}{2}$$

- Roots of the characteristic polynomial

$$r = (-1 - I\sqrt{2}, -1 + I\sqrt{2})$$

- 1st solution of the ODE

$$y_1(x) = e^{-x} \cos(x\sqrt{2})$$

- 2nd solution of the ODE

$$y_2(x) = e^{-x} \sin(x\sqrt{2})$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = c_1 e^{-x} \cos(x\sqrt{2}) + c_2 e^{-x} \sin(x\sqrt{2})$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 26

```
dsolve(diff(y(x),x$2)+2*diff(y(x),x)+3*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{-x} \left(c_1 \sin(\sqrt{2}x) + c_2 \cos(\sqrt{2}x) \right)$$

✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 34

```
DSolve[y''[x]+2*y'[x]+3*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x} \left(c_2 \cos(\sqrt{2}x) + c_1 \sin(\sqrt{2}x) \right)$$

9.13 problem 1(m)

9.13.1 Solving as second order linear constant coeff ode	1387
9.13.2 Solving as second order ode can be made integrable ode	1389
9.13.3 Solving using Kovacic algorithm	1391
9.13.4 Maple step by step solution	1395

Internal problem ID [6281]

Internal file name [OUTPUT/5529_Sunday_June_05_2022_03_42_45_PM_74314027/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.1. Linear Equations with Constant Coefficients. Page 62

Problem number: 1(m).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "second_order_ode_can_be_made_integrable"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x]]
```

$$y'' - 4y = 0$$

9.13.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = -4$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 4 e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 4 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = -4$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(-4)} \\ &= \pm 2 \end{aligned}$$

Hence

$$\lambda_1 = +2$$

$$\lambda_2 = -2$$

Which simplifies to

$$\lambda_1 = 2$$

$$\lambda_2 = -2$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(2)x} + c_2 e^{(-2)x}$$

Or

$$y = c_1 e^{2x} + c_2 e^{-2x}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{2x} + c_2 e^{-2x} \tag{1}$$




Figure 275: Slope field plot

Verification of solutions

$$y = c_1 e^{2x} + c_2 e^{-2x}$$

Verified OK.

9.13.2 Solving as second order ode can be made integrable ode

Multiplying the ode by y' gives

$$y' y'' - 4y' y = 0$$

Integrating the above w.r.t x gives

$$\int (y' y'' - 4y' y) dx = 0$$

$$\frac{y'^2}{2} - 2y^2 = c_2$$

Which is now solved for y . Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \sqrt{4y^2 + 2c_1} \tag{1}$$

$$y' = -\sqrt{4y^2 + 2c_1} \tag{2}$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int \frac{1}{\sqrt{4y^2 + 2c_1}} dy = \int dx$$
$$\frac{\ln(y\sqrt{4} + \sqrt{4y^2 + 2c_1})\sqrt{4}}{4} = c_2 + x$$

Raising both side to exponential gives

$$e^{\frac{\ln(y\sqrt{4} + \sqrt{4y^2 + 2c_1})\sqrt{4}}{4}} = e^{c_2 + x}$$

Which simplifies to

$$\sqrt{2y + \sqrt{4y^2 + 2c_1}} = c_3 e^x$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{1}{\sqrt{4y^2 + 2c_1}} dy = \int dx$$
$$-\frac{\ln(y\sqrt{4} + \sqrt{4y^2 + 2c_1})\sqrt{4}}{4} = x + c_4$$

Raising both side to exponential gives

$$e^{-\frac{\ln(y\sqrt{4} + \sqrt{4y^2 + 2c_1})\sqrt{4}}{4}} = e^{x + c_4}$$

Which simplifies to

$$\frac{1}{\sqrt{2y + \sqrt{4y^2 + 2c_1}}} = c_5 e^x$$

Summary

The solution(s) found are the following

$$y = \frac{(e^{4x} c_3^4 - 2c_1) e^{-2x}}{4c_3^2} \quad (1)$$

$$y = -\frac{(2c_1 c_5^4 e^{4x} - 1) e^{-2x}}{4c_5^2} \quad (2)$$




Figure 276: Slope field plot

Verification of solutions

$$y = \frac{(e^{4x}c_3^4 - 2c_1)e^{-2x}}{4c_3^2}$$

Verified OK.

$$y = -\frac{(2c_1c_5^4e^{4x} - 1)e^{-2x}}{4c_5^2}$$

Verified OK.

9.13.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 4y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= 1 \\B &= 0 \\C &= -4\end{aligned}\tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x)\tag{4}$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\&= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned}\tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{4}{1}\tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= 4 \\t &= 1\end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 4z(x)\tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 199: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 4$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-2x}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}
 y_1 &= z_1 \\
 &= e^{-2x}
 \end{aligned}$$

Which simplifies to

$$y_1 = e^{-2x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= e^{-2x} \int \frac{1}{e^{-4x}} dx \\ &= e^{-2x} \left(\frac{e^{4x}}{4} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-2x}) + c_2 \left(e^{-2x} \left(\frac{e^{4x}}{4} \right) \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-2x} + \frac{c_2 e^{2x}}{4} \tag{1}$$




Figure 277: Slope field plot

Verification of solutions

$$y = c_1 e^{-2x} + \frac{c_2 e^{2x}}{4}$$

Verified OK.

9.13.4 Maple step by step solution

Let's solve

$$y'' - 4y = 0$$

- Highest derivative means the order of the ODE is 2
- $$y''$$
- Characteristic polynomial of ODE
- $$r^2 - 4 = 0$$
- Factor the characteristic polynomial
- $$(r - 2)(r + 2) = 0$$
- Roots of the characteristic polynomial

- $r = (-2, 2)$
- 1st solution of the ODE
 $y_1(x) = e^{-2x}$
- 2nd solution of the ODE
 $y_2(x) = e^{2x}$
- General solution of the ODE
 $y = c_1 y_1(x) + c_2 y_2(x)$
- Substitute in solutions
 $y = c_1 e^{-2x} + c_2 e^{2x}$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)=4*y(x),y(x), singsol=all)
```

$$y(x) = e^{2x} c_1 + c_2 e^{-2x}$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 22

```
DSolve[y''[x]==4*y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2x} (c_1 e^{4x} + c_2)$$

9.14 problem 1(n)

- 9.14.1 Solving as second order linear constant coeff ode 1397
- 9.14.2 Solving using Kovacic algorithm 1399
- 9.14.3 Maple step by step solution 1403

Internal problem ID [6282]

Internal file name [OUTPUT/5530_Sunday_June_05_2022_03_42_46_PM_92651194/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.1. Linear Equations with Constant Coefficients. Page 62

Problem number: 1(n).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$4y'' - 8y' + 7y = 0$$

9.14.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 4, B = -8, C = 7$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$4\lambda^2 e^{\lambda x} - 8\lambda e^{\lambda x} + 7e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$4\lambda^2 - 8\lambda + 7 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 4, B = -8, C = 7$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{8}{(2)(4)} \pm \frac{1}{(2)(4)} \sqrt{-8^2 - (4)(4)(7)} \\ &= 1 \pm \frac{i\sqrt{3}}{2}\end{aligned}$$

Hence

$$\begin{aligned}\lambda_1 &= 1 + \frac{i\sqrt{3}}{2} \\ \lambda_2 &= 1 - \frac{i\sqrt{3}}{2}\end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= \frac{i\sqrt{3}}{2} + 1 \\ \lambda_2 &= 1 - \frac{i\sqrt{3}}{2}\end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 1$ and $\beta = \frac{\sqrt{3}}{2}$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^x \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

Summary

The solution(s) found are the following

$$y = e^x \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right) \quad (1)$$




Figure 278: Slope field plot

Verification of solutions

$$y = e^x \left(c_1 \cos \left(\frac{\sqrt{3} x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3} x}{2} \right) \right)$$

Verified OK.

9.14.2 Solving using Kovacic algorithm

Writing the ode as

$$4y'' - 8y' + 7y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 4 \\ B &= -8 \\ C &= 7 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-3}{4} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -3 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -\frac{3z(x)}{4} \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 201: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0\end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -\frac{3}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos\left(\frac{\sqrt{3}x}{2}\right)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-8}{4} dx} \\ &= z_1 e^x \\ &= z_1 (e^x)\end{aligned}$$

Which simplifies to

$$y_1 = e^x \cos\left(\frac{\sqrt{3}x}{2}\right)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{8}{4} dx}}{(y_1)^2} dx \\&= y_1 \int \frac{e^{2x}}{(y_1)^2} dx \\&= y_1 \left(\frac{2\sqrt{3} \tan\left(\frac{\sqrt{3}x}{2}\right)}{3} \right)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\&= c_1 \left(e^x \cos\left(\frac{\sqrt{3}x}{2}\right) \right) + c_2 \left(e^x \cos\left(\frac{\sqrt{3}x}{2}\right) \left(\frac{2\sqrt{3} \tan\left(\frac{\sqrt{3}x}{2}\right)}{3} \right) \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^x \cos\left(\frac{\sqrt{3}x}{2}\right) + \frac{2c_2 e^x \sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3} \quad (1)$$




Figure 279: Slope field plot

Verification of solutions

$$y = c_1 e^x \cos\left(\frac{\sqrt{3}x}{2}\right) + \frac{2c_2 e^x \sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3}$$

Verified OK.

9.14.3 Maple step by step solution

Let's solve

$$4y'' - 8y' + 7y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = 2y' - \frac{7y}{4}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - 2y' + \frac{7y}{4} = 0$$

- Characteristic polynomial of ODE

$$r^2 - 2r + \frac{7}{4} = 0$$
- Use quadratic formula to solve for r

$$r = \frac{2 \pm (\sqrt{-3})}{2}$$
- Roots of the characteristic polynomial

$$r = \left(1 - \frac{i\sqrt{3}}{2}, \frac{i\sqrt{3}}{2} + 1\right)$$
- 1st solution of the ODE

$$y_1(x) = e^x \cos\left(\frac{\sqrt{3}x}{2}\right)$$
- 2nd solution of the ODE

$$y_2(x) = e^x \sin\left(\frac{\sqrt{3}x}{2}\right)$$
- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$
- Substitute in solutions

$$y = c_1 e^x \cos\left(\frac{\sqrt{3}x}{2}\right) + c_2 e^x \sin\left(\frac{\sqrt{3}x}{2}\right)$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(4*diff(y(x),x$2)-8*diff(y(x),x)+7*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^x \left(c_1 \sin\left(\frac{\sqrt{3}x}{2}\right) + c_2 \cos\left(\frac{\sqrt{3}x}{2}\right) \right)$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 38

```
DSolve[4*y''[x]-8*y'[x]+7*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x \left(c_2 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_1 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

9.15 problem 1(o)

- 9.15.1 Solving as second order linear constant coeff ode 1406
- 9.15.2 Solving using Kovacic algorithm 1408
- 9.15.3 Maple step by step solution 1412

Internal problem ID [6283]

Internal file name [OUTPUT/5531_Sunday_June_05_2022_03_42_47_PM_58331744/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.1. Linear Equations with Constant Coefficients. Page 62

Problem number: 1(o).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$2y'' + y' - y = 0$$

9.15.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 2, B = 1, C = -1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$2\lambda^2 e^{\lambda x} + \lambda e^{\lambda x} - e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$2\lambda^2 + \lambda - 1 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 2, B = 1, C = -1$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{-1}{(2)(2)} \pm \frac{1}{(2)(2)} \sqrt{1^2 - (4)(2)(-1)} \\ &= -\frac{1}{4} \pm \frac{3}{4}\end{aligned}$$

Hence

$$\lambda_1 = -\frac{1}{4} + \frac{3}{4}$$

$$\lambda_2 = -\frac{1}{4} - \frac{3}{4}$$

Which simplifies to

$$\lambda_1 = \frac{1}{2}$$

$$\lambda_2 = -1$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(\frac{1}{2})x} + c_2 e^{(-1)x}$$

Or

$$y = c_1 e^{\frac{x}{2}} + c_2 e^{-x}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{\frac{x}{2}} + c_2 e^{-x} \tag{1}$$




Figure 280: Slope field plot

Verification of solutions

$$y = c_1 e^{\frac{x}{2}} + c_2 e^{-x}$$

Verified OK.

9.15.2 Solving using Kovacic algorithm

Writing the ode as

$$2y'' + y' - y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 2 \\ B &= 1 \\ C &= -1 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{9}{16} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 9 \\ t &= 16 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \frac{9z(x)}{16} \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 203: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{9}{16}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{3x}{4}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{1}{2} dx} \\ &= z_1 e^{-\frac{x}{4}} \\ &= z_1 \left(e^{-\frac{x}{4}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{1}{2} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-\frac{x}{2}}}{(y_1)^2} dx \\ &= y_1 \left(\frac{2 e^{\frac{3x}{2}}}{3} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-x}) + c_2 \left(e^{-x} \left(\frac{2e^{\frac{3x}{2}}}{3} \right) \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-x} + \frac{2c_2 e^{\frac{x}{2}}}{3} \quad (1)$$




Figure 281: Slope field plot

Verification of solutions

$$y = c_1 e^{-x} + \frac{2c_2 e^{\frac{x}{2}}}{3}$$

Verified OK.

9.15.3 Maple step by step solution

Let's solve

$$2y'' + y' - y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{y'}{2} + \frac{y}{2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{y'}{2} - \frac{y}{2} = 0$$

- Characteristic polynomial of ODE

$$r^2 + \frac{1}{2}r - \frac{1}{2} = 0$$

- Factor the characteristic polynomial

$$\frac{(r+1)(2r-1)}{2} = 0$$

- Roots of the characteristic polynomial

$$r = \left(-1, \frac{1}{2}\right)$$

- 1st solution of the ODE

$$y_1(x) = e^{-x}$$

- 2nd solution of the ODE

$$y_2(x) = e^{\frac{x}{2}}$$

- General solution of the ODE

$$y = c_1y_1(x) + c_2y_2(x)$$

- Substitute in solutions

$$y = c_1e^{-x} + c_2e^{\frac{x}{2}}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(2*diff(y(x),x$2)+diff(y(x),x)-y(x)=0,y(x), singsol=all)
```

$$y(x) = \left(c_1 e^{\frac{3x}{2}} + c_2 \right) e^{-x}$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 24

```
DSolve[2*y'[x]+y'[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x} (c_1 e^{3x/2} + c_2)$$

9.16 problem 1(p)

- 9.16.1 Solving as second order linear constant coeff ode 1414
- 9.16.2 Solving using Kovacic algorithm 1416
- 9.16.3 Maple step by step solution 1420

Internal problem ID [6284]

Internal file name [OUTPUT/5532_Sunday_June_05_2022_03_42_49_PM_74376436/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.1. Linear Equations with Constant Coefficients. Page 62

Problem number: 1(p).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' + 4y' + 5y = 0$$

9.16.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 4, C = 5$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 4\lambda e^{\lambda x} + 5e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 4\lambda + 5 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 4, C = 5$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{-4}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{4^2 - (4)(1)(5)} \\ &= -2 \pm i\end{aligned}$$

Hence

$$\begin{aligned}\lambda_1 &= -2 + i \\ \lambda_2 &= -2 - i\end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= -2 + i \\ \lambda_2 &= -2 - i\end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = -2$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^{-2x}(\cos(x) c_1 + c_2 \sin(x))$$

Summary

The solution(s) found are the following

$$y = e^{-2x}(\cos(x) c_1 + c_2 \sin(x)) \tag{1}$$




Figure 282: Slope field plot

Verification of solutions

$$y = e^{-2x}(\cos(x) c_1 + c_2 \sin(x))$$

Verified OK.

9.16.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 4y' + 5y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 4 \\ C &= 5 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 205: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0\end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{4}{1} dx} \\ &= z_1 e^{-2x} \\ &= z_1 (e^{-2x})\end{aligned}$$

Which simplifies to

$$y_1 = e^{-2x} \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{4}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-4x}}{(y_1)^2} dx \\ &= y_1 (\tan(x))\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-2x} \cos(x)) + c_2 (e^{-2x} \cos(x) (\tan(x)))\end{aligned}$$

Summary

The solution(s) found are the following

$$y = e^{-2x} \cos(x) c_1 + e^{-2x} \sin(x) c_2 \quad (1)$$




Figure 283: Slope field plot

Verification of solutions

$$y = e^{-2x} \cos(x) c_1 + e^{-2x} \sin(x) c_2$$

Verified OK.

9.16.3 Maple step by step solution

Let's solve

$$y'' + 4y' + 5y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 + 4r + 5 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-4) \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (-2 - I, -2 + I)$$

- 1st solution of the ODE

$$y_1(x) = e^{-2x} \cos(x)$$

- 2nd solution of the ODE

$$y_2(x) = \sin(x) e^{-2x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = e^{-2x} \cos(x) c_1 + e^{-2x} \sin(x) c_2$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(diff(y(x),x$2)+4*diff(y(x),x)+5*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{-2x}(c_1 \sin(x) + \cos(x) c_2)$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 22

```
DSolve[y''[x]+4*y'[x]+5*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2x}(c_2 \cos(x) + c_1 \sin(x))$$

9.17 problem 1(q)

- 9.17.1 Solving as second order linear constant coeff ode 1422
- 9.17.2 Solving using Kovacic algorithm 1424
- 9.17.3 Maple step by step solution 1428

Internal problem ID [6285]

Internal file name [OUTPUT/5533_Sunday_June_05_2022_03_42_50_PM_37612645/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.1. Linear Equations with Constant Coefficients. Page 62

Problem number: 1(q).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' + 4y' + 5y = 0$$

9.17.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 4, C = 5$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 4\lambda e^{\lambda x} + 5e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 4\lambda + 5 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 4, C = 5$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{-4}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{4^2 - (4)(1)(5)} \\ &= -2 \pm i\end{aligned}$$

Hence

$$\begin{aligned}\lambda_1 &= -2 + i \\ \lambda_2 &= -2 - i\end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= -2 + i \\ \lambda_2 &= -2 - i\end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = -2$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^{-2x}(\cos(x) c_1 + c_2 \sin(x))$$

Summary

The solution(s) found are the following

$$y = e^{-2x}(\cos(x) c_1 + c_2 \sin(x)) \tag{1}$$




Figure 284: Slope field plot

Verification of solutions

$$y = e^{-2x}(\cos(x) c_1 + c_2 \sin(x))$$

Verified OK.

9.17.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 4y' + 5y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 4 \\ C &= 5 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} = \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \tag{6}$$

Comparing the above to (5) shows that

$$s = -1$$

$$t = 1$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 207: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0\end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{4}{1} dx} \\ &= z_1 e^{-2x} \\ &= z_1 (e^{-2x})\end{aligned}$$

Which simplifies to

$$y_1 = e^{-2x} \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{4}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-4x}}{(y_1)^2} dx \\ &= y_1 (\tan(x))\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-2x} \cos(x)) + c_2 (e^{-2x} \cos(x) (\tan(x)))\end{aligned}$$

Summary

The solution(s) found are the following

$$y = e^{-2x} \cos(x) c_1 + e^{-2x} \sin(x) c_2 \quad (1)$$




Figure 285: Slope field plot

Verification of solutions

$$y = e^{-2x} \cos(x) c_1 + e^{-2x} \sin(x) c_2$$

Verified OK.

9.17.3 Maple step by step solution

Let's solve

$$y'' + 4y' + 5y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 + 4r + 5 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-4) \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (-2 - I, -2 + I)$$

- 1st solution of the ODE

$$y_1(x) = e^{-2x} \cos(x)$$

- 2nd solution of the ODE

$$y_2(x) = \sin(x) e^{-2x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = e^{-2x} \cos(x) c_1 + e^{-2x} \sin(x) c_2$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(diff(y(x),x$2)+4*diff(y(x),x)+5*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{-2x}(c_1 \sin(x) + \cos(x) c_2)$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 22

```
DSolve[y''[x]+4*y'[x]+5*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-2x}(c_2 \cos(x) + c_1 \sin(x))$$

9.18 problem 1(r)

- 9.18.1 Solving as second order linear constant coeff ode 1430
- 9.18.2 Solving using Kovacic algorithm 1432
- 9.18.3 Maple step by step solution 1436

Internal problem ID [6286]

Internal file name [OUTPUT/5534_Sunday_June_05_2022_03_42_51_PM_2680923/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.1. Linear Equations with Constant Coefficients. Page 62

Problem number: 1(r).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' + 4y' - 5y = 0$$

9.18.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 4, C = -5$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 4\lambda e^{\lambda x} - 5e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 4\lambda - 5 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 4, C = -5$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{-4}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{4^2 - (4)(1)(-5)} \\ &= -2 \pm 3\end{aligned}$$

Hence

$$\lambda_1 = -2 + 3$$

$$\lambda_2 = -2 - 3$$

Which simplifies to

$$\lambda_1 = 1$$

$$\lambda_2 = -5$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(1)x} + c_2 e^{(-5)x}$$

Or

$$y = c_1 e^x + c_2 e^{-5x}$$

Summary

The solution(s) found are the following

$$y = c_1 e^x + c_2 e^{-5x} \tag{1}$$




Figure 286: Slope field plot

Verification of solutions

$$y = c_1 e^x + c_2 e^{-5x}$$

Verified OK.

9.18.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 4y' - 5y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 4 \\ C &= -5 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{9}{1} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 9 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 9z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 209: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0\end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 9$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-3x}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{4}{1} dx} \\ &= z_1 e^{-2x} \\ &= z_1 (e^{-2x})\end{aligned}$$

Which simplifies to

$$y_1 = e^{-5x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{4}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-4x}}{(y_1)^2} dx \\ &= y_1 \left(\frac{e^{6x}}{6} \right)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-5x}) + c_2 \left(e^{-5x} \left(\frac{e^{6x}}{6} \right) \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-5x} + \frac{c_2 e^x}{6} \quad (1)$$




Figure 287: Slope field plot

Verification of solutions

$$y = c_1 e^{-5x} + \frac{c_2 e^x}{6}$$

Verified OK.

9.18.3 Maple step by step solution

Let's solve

$$y'' + 4y' - 5y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 + 4r - 5 = 0$$

- Factor the characteristic polynomial

$$(r + 5)(r - 1) = 0$$

- Roots of the characteristic polynomial

$$r = (-5, 1)$$

- 1st solution of the ODE

$$y_1(x) = e^{-5x}$$

- 2nd solution of the ODE

$$y_2(x) = e^x$$

- General solution of the ODE

$$y = c_1y_1(x) + c_2y_2(x)$$

- Substitute in solutions

$$y = c_1e^{-5x} + c_2e^x$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)+4*diff(y(x),x)-5*y(x)=0,y(x), singsol=all)
```

$$y(x) = (e^{6x}c_2 + c_1) e^{-5x}$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 20

```
DSolve[y''[x]+4*y'[x]-5*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-5x} + c_2 e^x$$

9.19 problem 2(a)

9.19.1 Existence and uniqueness analysis	1438
9.19.2 Solving as second order linear constant coeff ode	1439
9.19.3 Solving using Kovacic algorithm	1441
9.19.4 Maple step by step solution	1446

Internal problem ID [6287]

Internal file name [OUTPUT/5535_Sunday_June_05_2022_03_42_52_PM_50697687/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.1. Linear Equations with Constant Coefficients. Page 62

Problem number: 2(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' - 5y' + 6y = 0$$

With initial conditions

$$[y(1) = e^2, y'(1) = 3e^2]$$

9.19.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = -5$$

$$q(x) = 6$$

$$F = 0$$

Hence the ode is

$$y'' - 5y' + 6y = 0$$

The domain of $p(x) = -5$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 1$ is inside this domain. The domain of $q(x) = 6$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 1$ is also inside this domain. Hence solution exists and is unique.

9.19.2 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -5, C = 6$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 5\lambda e^{\lambda x} + 6 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 5\lambda + 6 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -5, C = 6$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{5}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-5^2 - (4)(1)(6)} \\ &= \frac{5}{2} \pm \frac{1}{2} \end{aligned}$$

Hence

$$\lambda_1 = \frac{5}{2} + \frac{1}{2}$$

$$\lambda_2 = \frac{5}{2} - \frac{1}{2}$$

Which simplifies to

$$\lambda_1 = 3$$

$$\lambda_2 = 2$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(3)x} + c_2 e^{(2)x}$$

Or

$$y = e^{3x} c_1 + c_2 e^{2x}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = e^{3x} c_1 + c_2 e^{2x} \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = e^2$ and $x = 1$ in the above gives

$$e^2 = e^3 c_1 + c_2 e^2 \tag{1A}$$

Taking derivative of the solution gives

$$y' = 3 e^{3x} c_1 + 2 c_2 e^{2x}$$

substituting $y' = 3 e^2$ and $x = 1$ in the above gives

$$3 e^2 = 3 e^3 c_1 + 2 c_2 e^2 \tag{2A}$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = e^{-1}$$

$$c_2 = 0$$

Substituting these values back in above solution results in

$$y = e^{3x} e^{-1}$$


Which simplifies to

$$y = e^{3x-1}$$


Summary

The solution(s) found are the following

$$y = e^{3x-1} \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = e^{3x-1}$$

Verified OK.

9.19.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 5y' + 6y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = -5 \tag{3}$$

$$C = 6$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{1}{4} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 1 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \frac{z(x)}{4} \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 211: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{1}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{x}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}
 y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\
 &= z_1 e^{-\int \frac{1}{2} \frac{-5}{1} dx} \\
 &= z_1 e^{\frac{5x}{2}} \\
 &= z_1 \left(e^{\frac{5x}{2}} \right)
 \end{aligned}$$

Which simplifies to

$$y_1 = e^{2x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{5}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{5x}}{(y_1)^2} dx \\ &= y_1(e^x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1(e^{2x}) + c_2(e^{2x}(e^x)) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 e^{2x} + c_2 e^{3x} \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = e^2$ and $x = 1$ in the above gives

$$e^2 = c_1 e^2 + c_2 e^3 \tag{1A}$$

Taking derivative of the solution gives

$$y' = 2c_1 e^{2x} + 3c_2 e^{3x}$$

substituting $y' = 3e^2$ and $x = 1$ in the above gives

$$3e^2 = 2c_1 e^2 + 3c_2 e^3 \tag{2A}$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$\begin{aligned}c_1 &= 0 \\c_2 &= e^{-1}\end{aligned}$$

Substituting these values back in above solution results in

$$y = e^{3x}e^{-1}$$


Which simplifies to

$$y = e^{3x-1}$$


Summary

The solution(s) found are the following

$$y = e^{3x-1} \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = e^{3x-1}$$

Verified OK.

9.19.4 Maple step by step solution

Let's solve

$$\left[y'' - 5y' + 6y = 0, y(1) = e^2, y' \Big|_{\{x=1\}} = 3e^2 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 - 5r + 6 = 0$$

- Factor the characteristic polynomial

$$(r - 2)(r - 3) = 0$$

- Roots of the characteristic polynomial

$$r = (2, 3)$$

- 1st solution of the ODE

$$y_1(x) = e^{2x}$$

- 2nd solution of the ODE

$$y_2(x) = e^{3x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = c_1 e^{2x} + c_2 e^{3x}$$

- Check validity of solution $y = c_1 e^{2x} + c_2 e^{3x}$

- Use initial condition $y(1) = e^2$

$$e^2 = c_1 e^2 + c_2 e^3$$

- Compute derivative of the solution

$$y' = 2c_1 e^{2x} + 3c_2 e^{3x}$$

- Use the initial condition $y' \Big|_{\{x=1\}} = 3e^2$

$$3e^2 = 2c_1 e^2 + 3c_2 e^3$$

- Solve for c_1 and c_2

$$\left\{ c_1 = 0, c_2 = \frac{e^2}{e^3} \right\}$$

- Substitute constant values into general solution and simplify

$$y = e^{3x-1}$$

- Solution to the IVP

$$y = e^{3x-1}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 11

```
dsolve([diff(y(x),x$2)-5*diff(y(x),x)+6*y(x)=0,y(1) = exp(2), D(y)(1) = 3*exp(2)],y(x), sing
```

$$y(x) = e^{3x-1}$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 12

```
DSolve[{y'[x]-5*y'[x]+6*y[x]==0,{y[1]==Exp[2],y'[1]==3*Exp[2]}},y[x],x,IncludeSingularSolut
```

$$y(x) \rightarrow e^{3x-1}$$

9.20 problem 2(b)

9.20.1 Existence and uniqueness analysis	1448
9.20.2 Solving as second order linear constant coeff ode	1449
9.20.3 Solving using Kovacic algorithm	1451
9.20.4 Maple step by step solution	1455

Internal problem ID [6288]

Internal file name [OUTPUT/5536_Sunday_June_05_2022_03_42_54_PM_84101560/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.1. Linear Equations with Constant Coefficients. Page 62

Problem number: 2(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' - 6y' + 5y = 0$$

With initial conditions

$$[y(0) = 3, y'(0) = 11]$$

9.20.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = -6$$

$$q(x) = 5$$

$$F = 0$$

Hence the ode is

$$y'' - 6y' + 5y = 0$$

The domain of $p(x) = -6$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = 5$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

9.20.2 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -6, C = 5$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 6\lambda e^{\lambda x} + 5 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 6\lambda + 5 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -6, C = 5$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{6}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-6^2 - (4)(1)(5)} \\ &= 3 \pm 2 \end{aligned}$$

Hence

$$\lambda_1 = 3 + 2$$

$$\lambda_2 = 3 - 2$$

Which simplifies to

$$\lambda_1 = 5$$

$$\lambda_2 = 1$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(5)x} + c_2 e^{(1)x}$$

Or

$$y = c_1 e^{5x} + c_2 e^x$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 e^{5x} + c_2 e^x \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 3$ and $x = 0$ in the above gives

$$3 = c_1 + c_2 \tag{1A}$$

Taking derivative of the solution gives

$$y' = 5c_1 e^{5x} + c_2 e^x$$

substituting $y' = 11$ and $x = 0$ in the above gives

$$11 = 5c_1 + c_2 \tag{2A}$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 2$$

$$c_2 = 1$$


Substituting these values back in above solution results in

$$y = 2e^{5x} + e^x$$


Summary

The solution(s) found are the following

$$y = 2e^{5x} + e^x \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = 2e^{5x} + e^x$$

Verified OK.

9.20.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 6y' + 5y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = -6 \quad (3)$$

$$C = 5$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} = \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{4}{1} \tag{6}$$

Comparing the above to (5) shows that

$$s = 4$$

$$t = 1$$

Therefore eq. (4) becomes

$$z''(x) = 4z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 213: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 4$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-2x}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-6}{1} dx} \\ &= z_1 e^{3x} \\ &= z_1 (e^{3x}) \end{aligned}$$

Which simplifies to

$$y_1 = e^x$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-6}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{6x}}{(y_1)^2} dx \\ &= y_1 \left(\frac{e^{4x}}{4} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1(e^x) + c_2\left(e^x\left(\frac{e^{4x}}{4}\right)\right)\end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 e^x + \frac{c_2 e^{5x}}{4} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 3$ and $x = 0$ in the above gives

$$3 = c_1 + \frac{c_2}{4} \quad (1A)$$

Taking derivative of the solution gives

$$y' = c_1 e^x + \frac{5c_2 e^{5x}}{4}$$

substituting $y' = 11$ and $x = 0$ in the above gives

$$11 = c_1 + \frac{5c_2}{4} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 1$$

$$c_2 = 8$$


Substituting these values back in above solution results in

$$y = 2e^{5x} + e^x$$


Summary

The solution(s) found are the following

$$y = 2e^{5x} + e^x \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = 2e^{5x} + e^x$$

Verified OK.

9.20.4 Maple step by step solution

Let's solve

$$\left[y'' - 6y' + 5y = 0, y(0) = 3, y' \Big|_{\{x=0\}} = 11 \right]$$

- Highest derivative means the order of the ODE is 2
- y''
- Characteristic polynomial of ODE

$$r^2 - 6r + 5 = 0$$
- Factor the characteristic polynomial

$$(r - 1)(r - 5) = 0$$
- Roots of the characteristic polynomial

$$r = (1, 5)$$
- 1st solution of the ODE

$$y_1(x) = e^x$$

- 2nd solution of the ODE

$$y_2(x) = e^{5x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = c_1 e^x + c_2 e^{5x}$$

- Check validity of solution $y = c_1 e^x + c_2 e^{5x}$

- Use initial condition $y(0) = 3$

$$3 = c_1 + c_2$$

- Compute derivative of the solution

$$y' = c_1 e^x + 5c_2 e^{5x}$$

- Use the initial condition $y' \Big|_{\{x=0\}} = 11$

$$11 = c_1 + 5c_2$$

- Solve for c_1 and c_2

$$\{c_1 = 1, c_2 = 2\}$$

- Substitute constant values into general solution and simplify

$$y = 2e^{5x} + e^x$$

- Solution to the IVP

$$y = 2e^{5x} + e^x$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 13

```
dsolve([diff(y(x),x$2)-6*diff(y(x),x)+5*y(x)=0,y(0) = 3, D(y)(0) = 11],y(x), singsol=all)
```

$$y(x) = 2e^{5x} + e^x$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 18

```
DSolve[{y'[x]-5*y'[x]+6*y[x]==0,{y[0]==3,y'[0]==11}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{2x}(5e^x - 2)$$

9.21 problem 2(c)

- 9.21.1 Existence and uniqueness analysis 1459
- 9.21.2 Solving as second order linear constant coeff ode 1459
- 9.21.3 Solving as linear second order ode solved by an integrating factor
ode 1461
- 9.21.4 Solving using Kovacic algorithm 1463
- 9.21.5 Maple step by step solution 1467

Internal problem ID [6289]

Internal file name [OUTPUT/5537_Sunday_June_05_2022_03_42_55_PM_27619051/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.1. Linear Equations with Constant Coefficients. Page 62

Problem number: 2(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "linear_second_order_ode_solved_by_an_integrating_factor"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x]]
```

$$y'' - 6y' + 9y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 5]$$

9.21.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = -6$$

$$q(x) = 9$$

$$F = 0$$

Hence the ode is

$$y'' - 6y' + 9y = 0$$

The domain of $p(x) = -6$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = 9$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

9.21.2 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -6, C = 9$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 6\lambda e^{\lambda x} + 9e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 6\lambda + 9 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -6, C = 9$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{6}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{(-6)^2 - (4)(1)(9)} \\ &= 3\end{aligned}$$

Hence this is the case of a double root $\lambda_{1,2} = -3$. Therefore the solution is

$$y = c_1 e^{3x} + c_2 x e^{3x} \quad (1)$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = e^{3x} c_1 + c_2 x e^{3x} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_1 \quad (1A)$$

Taking derivative of the solution gives

$$y' = 3e^{3x} c_1 + c_2 e^{3x} + 3c_2 x e^{3x}$$

substituting $y' = 5$ and $x = 0$ in the above gives

$$5 = 3c_1 + c_2 \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 0$$

$$c_2 = 5$$


Substituting these values back in above solution results in

$$y = 5 e^{3x} x$$


Summary

The solution(s) found are the following

$$y = 5 e^{3x} x \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = 5e^{3x}x$$

Verified OK.

9.21.3 Solving as linear second order ode solved by an integrating factor ode

The ode satisfies this form

$$y'' + p(x)y' + \frac{(p(x)^2 + p'(x))y}{2} = f(x)$$

Where $p(x) = -6$. Therefore, there is an integrating factor given by

$$\begin{aligned} M(x) &= e^{\frac{1}{2} \int p dx} \\ &= e^{\int -6 dx} \\ &= e^{-3x} \end{aligned}$$

Multiplying both sides of the ODE by the integrating factor $M(x)$ makes the left side of the ODE a complete differential

$$\begin{aligned} (M(x)y)'' &= 0 \\ (e^{-3x}y)'' &= 0 \end{aligned}$$

Integrating once gives

$$(e^{-3x}y)' = c_1$$

Integrating again gives

$$(e^{-3x}y) = c_1x + c_2$$

Hence the solution is

$$y = \frac{c_1x + c_2}{e^{-3x}}$$

Or

$$y = c_1x e^{3x} + c_2e^{3x}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1x e^{3x} + c_2e^{3x} \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_2 \tag{1A}$$

Taking derivative of the solution gives

$$y' = e^{3x}c_1 + 3c_1x e^{3x} + 3c_2e^{3x}$$

substituting $y' = 5$ and $x = 0$ in the above gives

$$5 = c_1 + 3c_2 \tag{2A}$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 5$$

$$c_2 = 0$$


Substituting these values back in above solution results in

$$y = 5 e^{3x}x$$


Summary

The solution(s) found are the following

$$y = 5 e^{3x}x \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = 5 e^{3x} x$$

Verified OK.

9.21.4 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 6y' + 9y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = -6 \tag{3}$$

$$C = 9$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} \tag{5}$$

$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \tag{6}$$

Comparing the above to (5) shows that

$$s = 0$$

$$t = 1$$

Therefore eq. (4) becomes

$$z''(x) = 0 \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 215: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty\end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-6}{1} dx} \\ &= z_1 e^{3x} \\ &= z_1 (e^{3x})\end{aligned}$$

Which simplifies to

$$y_1 = e^{3x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{-6}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{6x}}{(y_1)^2} dx \\ &= y_1(x)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{3x}) + c_2 (e^{3x}(x))\end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = e^{3x} c_1 + c_2 x e^{3x} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = c_1 \quad (1A)$$

Taking derivative of the solution gives

$$y' = 3e^{3x} c_1 + c_2 e^{3x} + 3c_2 x e^{3x}$$

substituting $y' = 5$ and $x = 0$ in the above gives

$$5 = 3c_1 + c_2 \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 0$$

$$c_2 = 5$$


Substituting these values back in above solution results in

$$y = 5 e^{3x} x$$


Summary

The solution(s) found are the following

$$y = 5 e^{3x} x \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = 5e^{3x}$$

Verified OK.

9.21.5 Maple step by step solution

Let's solve

$$\left[y'' - 6y' + 9y = 0, y(0) = 0, y'|_{\{x=0\}} = 5 \right]$$

- Highest derivative means the order of the ODE is 2
 y''
- Characteristic polynomial of ODE
 $r^2 - 6r + 9 = 0$
- Factor the characteristic polynomial
 $(r - 3)^2 = 0$
- Root of the characteristic polynomial
 $r = 3$
- 1st solution of the ODE

$$y_1(x) = e^{3x}$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = e^{3x}x$$

- General solution of the ODE

$$y = c_1y_1(x) + c_2y_2(x)$$

- Substitute in solutions

$$y = e^{3x}c_1 + c_2x e^{3x}$$

- Check validity of solution $y = e^{3x}c_1 + c_2xe^{3x}$

- Use initial condition $y(0) = 0$

$$0 = c_1$$

- Compute derivative of the solution

$$y' = 3e^{3x}c_1 + c_2e^{3x} + 3c_2xe^{3x}$$

- Use the initial condition $y' \Big|_{\{x=0\}} = 5$

$$5 = 3c_1 + c_2$$

- Solve for c_1 and c_2

$$\{c_1 = 0, c_2 = 5\}$$

- Substitute constant values into general solution and simplify

$$y = 5e^{3x}x$$

- Solution to the IVP

$$y = 5e^{3x}x$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve([diff(y(x),x$2)-6*diff(y(x),x)+9*y(x)=0,y(0) = 0, D(y)(0) = 5],y(x), singsol=all)
```

$$y(x) = 5x e^{3x}$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 13

```
DSolve[{y''[x]-6*y'[x]+9*y[x]==0,{y[0]==0,y'[0]==5}},y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow 5e^{3x}x$$

9.22 problem 2(d)

9.22.1 Existence and uniqueness analysis	1470
9.22.2 Solving as second order linear constant coeff ode	1471
9.22.3 Solving using Kovacic algorithm	1473
9.22.4 Maple step by step solution	1477

Internal problem ID [6290]

Internal file name [OUTPUT/5538_Sunday_June_05_2022_03_42_57_PM_17264032/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.1. Linear Equations with Constant Coefficients. Page 62

Problem number: 2(d).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' + 4y' + 5y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 0]$$

9.22.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = 4$$

$$q(x) = 5$$

$$F = 0$$

Hence the ode is

$$y'' + 4y' + 5y = 0$$

The domain of $p(x) = 4$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = 5$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

9.22.2 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 4, C = 5$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 4\lambda e^{\lambda x} + 5e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 4\lambda + 5 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 4, C = 5$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-4}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{4^2 - (4)(1)(5)} \\ &= -2 \pm i \end{aligned}$$

Hence

$$\lambda_1 = -2 + i$$

$$\lambda_2 = -2 - i$$

Which simplifies to

$$\lambda_1 = -2 + i$$

$$\lambda_2 = -2 - i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = -2$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^{-2x}(\cos(x) c_1 + c_2 \sin(x))$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = e^{-2x}(\cos(x) c_1 + c_2 \sin(x)) \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 0$ in the above gives

$$1 = c_1 \tag{1A}$$

Taking derivative of the solution gives

$$y' = -2 e^{-2x}(\cos(x) c_1 + c_2 \sin(x)) + e^{-2x}(-\sin(x) c_1 + c_2 \cos(x))$$

substituting $y' = 0$ and $x = 0$ in the above gives

$$0 = -2c_1 + c_2 \tag{2A}$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 1$$

$$c_2 = 2$$


Substituting these values back in above solution results in

$$y = e^{-2x}(2 \sin(x) + \cos(x))$$


Summary

The solution(s) found are the following

$$y = e^{-2x}(2 \sin(x) + \cos(x)) \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = e^{-2x}(2 \sin(x) + \cos(x))$$

Verified OK.

9.22.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 4y' + 5y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 4 \quad (3)$$

$$C = 5$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} = \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \tag{6}$$

Comparing the above to (5) shows that

$$s = -1$$

$$t = 1$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 217: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0\end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{4}{1} dx} \\ &= z_1 e^{-2x} \\ &= z_1 (e^{-2x})\end{aligned}$$

Which simplifies to

$$y_1 = e^{-2x} \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{4}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-4x}}{(y_1)^2} dx \\ &= y_1 (\tan(x))\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-2x} \cos(x)) + c_2 (e^{-2x} \cos(x) (\tan(x)))\end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = e^{-2x} \cos(x) c_1 + e^{-2x} \sin(x) c_2 \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 0$ in the above gives

$$1 = c_1 \quad (1A)$$

Taking derivative of the solution gives

$$y' = -2 e^{-2x} \cos(x) c_1 - e^{-2x} \sin(x) c_1 - 2 e^{-2x} \sin(x) c_2 + e^{-2x} \cos(x) c_2$$

substituting $y' = 0$ and $x = 0$ in the above gives

$$0 = -2c_1 + c_2 \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 1$$

$$c_2 = 2$$

Substituting these values back in above solution results in

$$y = e^{-2x} \cos(x) + 2 \sin(x) e^{-2x}$$


Which simplifies to

$$y = e^{-2x} (2 \sin(x) + \cos(x))$$


Summary

The solution(s) found are the following

$$y = e^{-2x} (2 \sin(x) + \cos(x)) \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = e^{-2x}(2 \sin(x) + \cos(x))$$

Verified OK.

9.22.4 Maple step by step solution

Let's solve

$$\left[y'' + 4y' + 5y = 0, y(0) = 1, y'|_{\{x=0\}} = 0 \right]$$

- Highest derivative means the order of the ODE is 2
- y''
- Characteristic polynomial of ODE
- $r^2 + 4r + 5 = 0$
- Use quadratic formula to solve for r
- $r = \frac{(-4) \pm (\sqrt{-4})}{2}$
- Roots of the characteristic polynomial
- $r = (-2 - I, -2 + I)$
- 1st solution of the ODE

$$y_1(x) = e^{-2x} \cos(x)$$

- 2nd solution of the ODE

$$y_2(x) = \sin(x) e^{-2x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = e^{-2x} \cos(x) c_1 + e^{-2x} \sin(x) c_2$$

- Check validity of solution $y = e^{-2x} \cos(x) c_1 + e^{-2x} \sin(x) c_2$

- Use initial condition $y(0) = 1$

$$1 = c_1$$

- Compute derivative of the solution

$$y' = -2 e^{-2x} \cos(x) c_1 - e^{-2x} \sin(x) c_1 - 2 e^{-2x} \sin(x) c_2 + e^{-2x} \cos(x) c_2$$

- Use the initial condition $y' \Big|_{\{x=0\}} = 0$

$$0 = -2c_1 + c_2$$

- Solve for c_1 and c_2

$$\{c_1 = 1, c_2 = 2\}$$

- Substitute constant values into general solution and simplify

$$y = e^{-2x}(2 \sin(x) + \cos(x))$$

- Solution to the IVP

$$y = e^{-2x}(2 \sin(x) + \cos(x))$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 16

```
dsolve([diff(y(x),x$2)+4*diff(y(x),x)+5*y(x)=0,y(0) = 1, D(y)(0) = 0],y(x), singsol=all)
```

$$y(x) = e^{-2x}(2 \sin(x) + \cos(x))$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 18

```
DSolve[{y''[x]+4*y'[x]+5*y[x]==0,{y[0]==1,y'[0]==0}},y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow e^{-2x}(2 \sin(x) + \cos(x))$$

9.23 problem 2(e)

9.23.1 Existence and uniqueness analysis	1480
9.23.2 Solving as second order linear constant coeff ode	1481
9.23.3 Solving using Kovacic algorithm	1483
9.23.4 Maple step by step solution	1487

Internal problem ID [6291]

Internal file name [OUTPUT/5539_Sunday_June_05_2022_03_42_58_PM_81195951/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.1. Linear Equations with Constant Coefficients. Page 62

Problem number: 2(e).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' + 4y' + 2y = 0$$

With initial conditions

$$\left[y(0) = -1, y'(0) = 2 + 3\sqrt{2} \right]$$

9.23.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = 4$$

$$q(x) = 2$$

$$F = 0$$

Hence the ode is

$$y'' + 4y' + 2y = 0$$

The domain of $p(x) = 4$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = 2$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

9.23.2 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 4, C = 2$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 4\lambda e^{\lambda x} + 2e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 4\lambda + 2 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 4, C = 2$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-4}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{4^2 - (4)(1)(2)} \\ &= -2 \pm \sqrt{2} \end{aligned}$$

Hence

$$\lambda_1 = -2 + \sqrt{2}$$

$$\lambda_2 = -2 - \sqrt{2}$$

Which simplifies to

$$\lambda_1 = -2 + \sqrt{2}$$

$$\lambda_2 = -2 - \sqrt{2}$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(-2+\sqrt{2})x} + c_2 e^{(-2-\sqrt{2})x}$$

Or

$$y = c_1 e^{(-2+\sqrt{2})x} + c_2 e^{(-2-\sqrt{2})x}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 e^{(-2+\sqrt{2})x} + c_2 e^{(-2-\sqrt{2})x} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = -1$ and $x = 0$ in the above gives

$$-1 = c_1 + c_2 \quad (1A)$$

Taking derivative of the solution gives

$$y' = c_1 (-2 + \sqrt{2}) e^{(-2+\sqrt{2})x} + c_2 (-2 - \sqrt{2}) e^{(-2-\sqrt{2})x}$$

substituting $y' = 2 + 3\sqrt{2}$ and $x = 0$ in the above gives

$$2 + 3\sqrt{2} = (c_1 - c_2) \sqrt{2} - 2c_1 - 2c_2 \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 1$$

$$c_2 = -2$$


Substituting these values back in above solution results in

$$y = e^{(-2+\sqrt{2})x} - 2e^{(-2-\sqrt{2})x}$$


Summary

The solution(s) found are the following

$$y = e^{(-2+\sqrt{2})x} - 2e^{-(2+\sqrt{2})x} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = e^{(-2+\sqrt{2})x} - 2e^{-(2+\sqrt{2})x}$$

Verified OK.

9.23.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 4y' + 2y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 4 \quad (3)$$

$$C = 2$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} = \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{2}{1} \tag{6}$$

Comparing the above to (5) shows that

$$s = 2$$

$$t = 1$$

Therefore eq. (4) becomes

$$z''(x) = 2z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 219: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 2$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-x\sqrt{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{4}{1} dx} \\ &= z_1 e^{-2x} \\ &= z_1 (e^{-2x}) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-(2+\sqrt{2})x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{4}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-4x}}{(y_1)^2} dx \\ &= y_1 \left(\frac{\sqrt{2} e^{2x\sqrt{2}}}{4} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}
 y &= c_1 y_1 + c_2 y_2 \\
 &= c_1 \left(e^{-(2+\sqrt{2})x} \right) + c_2 \left(e^{-(2+\sqrt{2})x} \left(\frac{\sqrt{2} e^{2x\sqrt{2}}}{4} \right) \right)
 \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 e^{-(2+\sqrt{2})x} + \frac{c_2 e^{(-2+\sqrt{2})x} \sqrt{2}}{4} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = -1$ and $x = 0$ in the above gives

$$-1 = c_1 + \frac{c_2 \sqrt{2}}{4} \quad (1A)$$

Taking derivative of the solution gives

$$y' = c_1 (-2 - \sqrt{2}) e^{-(2+\sqrt{2})x} + \frac{c_2 (-2 + \sqrt{2}) e^{(-2+\sqrt{2})x} \sqrt{2}}{4}$$

substituting $y' = 2 + 3\sqrt{2}$ and $x = 0$ in the above gives

$$2 + 3\sqrt{2} = \frac{(-2c_1 - c_2) \sqrt{2}}{2} - 2c_1 + \frac{c_2}{2} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$\begin{aligned}
 c_1 &= -2 \\
 c_2 &= 2\sqrt{2}
 \end{aligned}$$


Substituting these values back in above solution results in

$$y = e^{(-2+\sqrt{2})x} - 2e^{-(2+\sqrt{2})x}$$


Summary

The solution(s) found are the following

$$y = e^{(-2+\sqrt{2})x} - 2e^{-(2+\sqrt{2})x} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = e^{(-2+\sqrt{2})x} - 2e^{-(2+\sqrt{2})x}$$

Verified OK.

9.23.4 Maple step by step solution

Let's solve

$$\left[y'' + 4y' + 2y = 0, y(0) = -1, y'|_{\{x=0\}} = 2 + 3\sqrt{2} \right]$$

- Highest derivative means the order of the ODE is 2

y''

- Characteristic polynomial of ODE

$$r^2 + 4r + 2 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-4) \pm (\sqrt{8})}{2}$$

- Roots of the characteristic polynomial

$$r = (-2 - \sqrt{2}, -2 + \sqrt{2})$$

- 1st solution of the ODE

$$y_1(x) = e^{(-2-\sqrt{2})x}$$

- 2nd solution of the ODE

$$y_2(x) = e^{(-2+\sqrt{2})x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = c_1 e^{(-2-\sqrt{2})x} + c_2 e^{(-2+\sqrt{2})x}$$

- Check validity of solution $y = c_1 e^{(-2-\sqrt{2})x} + c_2 e^{(-2+\sqrt{2})x}$

- Use initial condition $y(0) = -1$

$$-1 = c_1 + c_2$$

- Compute derivative of the solution

$$y' = c_1(-2 - \sqrt{2}) e^{(-2-\sqrt{2})x} + c_2(-2 + \sqrt{2}) e^{(-2+\sqrt{2})x}$$

- Use the initial condition $y'|_{\{x=0\}} = 2 + 3\sqrt{2}$

$$2 + 3\sqrt{2} = (-2 - \sqrt{2}) c_1 + (-2 + \sqrt{2}) c_2$$

- Solve for c_1 and c_2

$$\{c_1 = -2, c_2 = 1\}$$

- Substitute constant values into general solution and simplify

$$y = e^{(-2+\sqrt{2})x} - 2e^{(-2-\sqrt{2})x}$$

- Solution to the IVP

$$y = e^{(-2+\sqrt{2})x} - 2e^{(-2-\sqrt{2})x}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 24

```
dsolve([diff(y(x),x$2)+4*diff(y(x),x)+2*y(x)=0,y(0) = -1, D(y)(0) = 2+3*sqrt(2)],y(x), sings
```

$$y(x) = e^{(-2+\sqrt{2})x} - 2e^{-(2+\sqrt{2})x}$$

✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 30

```
DSolve[{y'[x]+4*y'[x]+2*y[x]==0,{y[0]==-1,y'[0]==2+3*Sqrt[2]}},y[x],x,IncludeSingularSoluti
```

$$y(x) \rightarrow e^{-((2+\sqrt{2})x)} \left(e^{2\sqrt{2}x} - 2 \right)$$

9.24 problem 2(f)

9.24.1 Existence and uniqueness analysis	1490
9.24.2 Solving as second order linear constant coeff ode	1491
9.24.3 Solving using Kovacic algorithm	1493
9.24.4 Maple step by step solution	1498

Internal problem ID [6292]

Internal file name [OUTPUT/5540_Sunday_June_05_2022_03_43_00_PM_87683680/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.1. Linear Equations with Constant Coefficients. Page 62

Problem number: 2(f).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' + 8y' - 9y = 0$$

With initial conditions

$$[y(1) = 2, y'(1) = 0]$$

9.24.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = 8$$

$$q(x) = -9$$

$$F = 0$$

Hence the ode is

$$y'' + 8y' - 9y = 0$$

The domain of $p(x) = 8$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 1$ is inside this domain. The domain of $q(x) = -9$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 1$ is also inside this domain. Hence solution exists and is unique.

9.24.2 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 8, C = -9$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 8\lambda e^{\lambda x} - 9e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 8\lambda - 9 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 8, C = -9$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-8}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{8^2 - (4)(1)(-9)} \\ &= -4 \pm 5 \end{aligned}$$

Hence

$$\lambda_1 = -4 + 5$$

$$\lambda_2 = -4 - 5$$

Which simplifies to

$$\lambda_1 = 1$$

$$\lambda_2 = -9$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(1)x} + c_2 e^{(-9)x}$$

Or

$$y = c_1 e^x + c_2 e^{-9x}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 e^x + c_2 e^{-9x} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 2$ and $x = 1$ in the above gives

$$2 = (c_1 e^{10} + c_2) e^{-9} \quad (1A)$$

Taking derivative of the solution gives

$$y' = c_1 e^x - 9c_2 e^{-9x}$$

substituting $y' = 0$ and $x = 1$ in the above gives

$$0 = (c_1 e^{10} - 9c_2) e^{-9} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = \frac{9 e^{-1}}{5}$$

$$c_2 = \frac{e^9}{5}$$


Substituting these values back in above solution results in

$$y = \frac{9 e^x e^{-1}}{5} + \frac{e^{-9x} e^9}{5}$$


Summary

The solution(s) found are the following

$$y = \frac{9e^xe^{-1}}{5} + \frac{e^{-9x}e^9}{5} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{9e^xe^{-1}}{5} + \frac{e^{-9x}e^9}{5}$$

Verified OK.

9.24.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 8y' - 9y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 8 \\ C &= -9 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{25}{1} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 25 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 25z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 221: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 25$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-5x}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}
 y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\
 &= z_1 e^{-\int \frac{1}{2} \frac{8}{1} dx} \\
 &= z_1 e^{-4x} \\
 &= z_1 (e^{-4x})
 \end{aligned}$$

Which simplifies to

$$y_1 = e^{-9x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{8}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-8x}}{(y_1)^2} dx \\ &= y_1 \left(\frac{e^{10x}}{10} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-9x}) + c_2 \left(e^{-9x} \left(\frac{e^{10x}}{10} \right) \right) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 e^{-9x} + \frac{c_2 e^x}{10} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 2$ and $x = 1$ in the above gives

$$2 = \frac{(c_2 e^{10} + 10c_1) e^{-9}}{10} \quad (1A)$$

Taking derivative of the solution gives

$$y' = -9c_1 e^{-9x} + \frac{c_2 e^x}{10}$$

substituting $y' = 0$ and $x = 1$ in the above gives

$$0 = \frac{(c_2 e^{10} - 90c_1) e^{-9}}{10} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = \frac{e^9}{5}$$

$$c_2 = 18e^{-1}$$


Substituting these values back in above solution results in

$$y = \frac{9e^x e^{-1}}{5} + \frac{e^{-9x} e^9}{5}$$


Summary

The solution(s) found are the following

$$y = \frac{9e^x e^{-1}}{5} + \frac{e^{-9x} e^9}{5} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{9e^x e^{-1}}{5} + \frac{e^{-9x} e^9}{5}$$

Verified OK.

9.24.4 Maple step by step solution

Let's solve

$$\left[y'' + 8y' - 9y = 0, y(1) = 2, y' \Big|_{\{x=1\}} = 0 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 + 8r - 9 = 0$$

- Factor the characteristic polynomial

$$(r + 9)(r - 1) = 0$$

- Roots of the characteristic polynomial

$$r = (-9, 1)$$

- 1st solution of the ODE

$$y_1(x) = e^{-9x}$$

- 2nd solution of the ODE

$$y_2(x) = e^x$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = c_1 e^{-9x} + c_2 e^x$$

- Check validity of solution $y = c_1 e^{-9x} + c_2 e^x$

- Use initial condition $y(1) = 2$

$$2 = c_1 e^{-9} + c_2 e$$

- Compute derivative of the solution

$$y' = -9c_1 e^{-9x} + c_2 e^x$$

- Use the initial condition $y' \Big|_{\{x=1\}} = 0$

$$0 = -9c_1 e^{-9} + c_2 e$$

- Solve for c_1 and c_2

$$\left\{ c_1 = \frac{1}{5e^{-9}}, c_2 = \frac{9}{5e} \right\}$$

- Substitute constant values into general solution and simplify

$$y = \frac{9e^{x-1}}{5} + \frac{e^{-9x+9}}{5}$$

- Solution to the IVP

$$y = \frac{9e^{x-1}}{5} + \frac{e^{-9x+9}}{5}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 19

```
dsolve([diff(y(x),x$2)+8*diff(y(x),x)-9*y(x)=0,y(1) = 2, D(y)(1) = 0],y(x), singsol=all)
```

$$y(x) = \frac{9e^{x-1}}{5} + \frac{e^{9-9x}}{5}$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 26

```
DSolve[{y''[x]+8*y'[x]-9*y[x]==0,{y[1]==2,y'[1]==0}},y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow \frac{1}{5}e^{9-9x} + \frac{9e^{x-1}}{5}$$

9.25 problem 5(a)

9.25.1 Solving as second order euler ode ode	1500
9.25.2 Solving as second order change of variable on x method 2 ode .	1502
9.25.3 Solving as second order change of variable on x method 1 ode .	1505
9.25.4 Solving as second order change of variable on y method 2 ode .	1507
9.25.5 Solving using Kovacic algorithm	1509
9.25.6 Maple step by step solution	1514

Internal problem ID [6293]

Internal file name [OUTPUT/5541_Sunday_June_05_2022_03_43_01_PM_56111531/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.1. Linear Equations with Constant Coefficients. Page 62

Problem number: 5(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_euler_ode", "second_order_change_of_variable_on_x_method_1", "second_order_change_of_variable_on_x_method_2", "second_order_change_of_variable_on_y_method_2"

Maple gives the following as the ode type

```
[[_Emden, _Fowler]]
```

$$x^2y'' + 3xy' + 10y = 0$$

9.25.1 Solving as second order euler ode ode

This is Euler second order ODE. Let the solution be $y = x^r$, then $y' = rx^{r-1}$ and $y'' = r(r-1)x^{r-2}$. Substituting these back into the given ODE gives

$$x^2(r(r-1))x^{r-2} + 3rx^{r-1} + 10x^r = 0$$

Simplifying gives

$$r(r-1)x^r + 3rx^r + 10x^r = 0$$

Since $x^r \neq 0$ then dividing throughout by x^r gives

$$r(r - 1) + 3r + 10 = 0$$

Or

$$r^2 + 2r + 10 = 0 \quad (1)$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = -1 - 3i$$

$$r_2 = -1 + 3i$$

The roots are complex conjugate of each others. Let the roots be

$$r_1 = \alpha + i\beta$$

$$r_2 = \alpha - i\beta$$

Where in this case $\alpha = -1$ and $\beta = -3$. Hence the solution becomes

$$\begin{aligned} y &= c_1 x^{r_1} + c_2 x^{r_2} \\ &= c_1 x^{\alpha+i\beta} + c_2 x^{\alpha-i\beta} \\ &= x^\alpha (c_1 x^{i\beta} + c_2 x^{-i\beta}) \\ &= x^\alpha (c_1 e^{\ln(x^{i\beta})} + c_2 e^{\ln(x^{-i\beta})}) \\ &= x^\alpha (c_1 e^{i(\beta \ln x)} + c_2 e^{-i(\beta \ln x)}) \end{aligned}$$

Using the values for $\alpha = -1, \beta = -3$, the above becomes

$$y = x^{-1} (c_1 e^{-3i \ln(x)} + c_2 e^{3i \ln(x)})$$

Using Euler relation, the expression $c_1 e^{iA} + c_2 e^{-iA}$ is transformed to $c_1 \cos A + c_1 \sin A$ where the constants are free to change. Applying this to the above result gives

$$y = \frac{1}{x} (c_1 \cos(3 \ln(x)) + c_2 \sin(3 \ln(x)))$$

Summary

The solution(s) found are the following

$$y = \frac{c_1 \cos(3 \ln(x)) + c_2 \sin(3 \ln(x))}{x} \quad (1)$$

Verification of solutions

$$y = \frac{c_1 \cos(3 \ln(x)) + c_2 \sin(3 \ln(x))}{x}$$

Verified OK.

9.25.2 Solving as second order change of variable on x method 2 ode

In normal form the ode

$$x^2 y'' + 3xy' + 10y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{3}{x}$$
$$q(x) = \frac{10}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x)\tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned} \tau &= \int e^{-(\int p(x)dx)} dx \\ &= \int e^{-(\int \frac{3}{x} dx)} dx \\ &= \int e^{-3\ln(x)} dx \\ &= \int \frac{1}{x^3} dx \\ &= -\frac{1}{2x^2} \end{aligned} \quad (6)$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned}q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\ &= \frac{\frac{10}{x^2}}{\frac{1}{x^6}} \\ &= 10x^4\end{aligned}\tag{7}$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned}\frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) + 10x^4y(\tau) &= 0\end{aligned}$$

But in terms of τ

$$10x^4 = \frac{5}{2\tau^2}$$

Hence the above ode becomes

$$\frac{d^2}{d\tau^2}y(\tau) + \frac{5y(\tau)}{2\tau^2} = 0$$

The above ode is now solved for $y(\tau)$. The ode can be written as

$$2\left(\frac{d^2}{d\tau^2}y(\tau)\right)\tau^2 + 5y(\tau) = 0$$

Which shows it is a Euler ODE. This is Euler second order ODE. Let the solution be $y(\tau) = \tau^r$, then $y' = r\tau^{r-1}$ and $y'' = r(r-1)\tau^{r-2}$. Substituting these back into the given ODE gives

$$2\tau^2(r(r-1))\tau^{r-2} + 0r\tau^{r-1} + 5\tau^r = 0$$

Simplifying gives

$$2r(r-1)\tau^r + 0\tau^r + 5\tau^r = 0$$

Since $\tau^r \neq 0$ then dividing throughout by τ^r gives

$$2r(r-1) + 0 + 5 = 0$$

Or

$$2r^2 - 2r + 5 = 0\tag{1}$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = \frac{1}{2} - \frac{3i}{2}$$

$$r_2 = \frac{1}{2} + \frac{3i}{2}$$

The roots are complex conjugate of each others. Let the roots be

$$r_1 = \alpha + i\beta$$

$$r_2 = \alpha - i\beta$$

Where in this case $\alpha = \frac{1}{2}$ and $\beta = -\frac{3}{2}$. Hence the solution becomes

$$y(\tau) = c_1\tau^{r_1} + c_2\tau^{r_2}$$

$$= c_1\tau^{\alpha+i\beta} + c_2\tau^{\alpha-i\beta}$$

$$= \tau^\alpha (c_1\tau^{i\beta} + c_2\tau^{-i\beta})$$

$$= \tau^\alpha (c_1e^{\ln(\tau^{i\beta})} + c_2e^{\ln(\tau^{-i\beta})})$$

$$= \tau^\alpha (c_1e^{i(\beta \ln \tau)} + c_2e^{-i(\beta \ln \tau)})$$

Using the values for $\alpha = \frac{1}{2}$, $\beta = -\frac{3}{2}$, the above becomes

$$y(\tau) = \tau^{\frac{1}{2}} \left(c_1 e^{-\frac{3i \ln(\tau)}{2}} + c_2 e^{\frac{3i \ln(\tau)}{2}} \right)$$

Using Euler relation, the expression $c_1e^{iA} + c_2e^{-iA}$ is transformed to $c_1 \cos A + c_1 \sin A$ where the constants are free to change. Applying this to the above result gives

$$y(\tau) = \sqrt{\tau} \left(c_1 \cos \left(\frac{3 \ln(\tau)}{2} \right) + c_2 \sin \left(\frac{3 \ln(\tau)}{2} \right) \right)$$

The above solution is now transformed back to y using (6) which results in

$$y = \frac{\left(c_1 \cos \left(-\frac{3 \ln(2)}{2} + \frac{3 \ln\left(-\frac{1}{x^2}\right)}{2} \right) + c_2 \sin \left(-\frac{3 \ln(2)}{2} + \frac{3 \ln\left(-\frac{1}{x^2}\right)}{2} \right) \right) \sqrt{2} \sqrt{-\frac{1}{x^2}}}{2}$$

Summary

The solution(s) found are the following

$$y = \frac{\left(c_1 \cos \left(-\frac{3 \ln(2)}{2} + \frac{3 \ln\left(-\frac{1}{x^2}\right)}{2} \right) + c_2 \sin \left(-\frac{3 \ln(2)}{2} + \frac{3 \ln\left(-\frac{1}{x^2}\right)}{2} \right) \right) \sqrt{2} \sqrt{-\frac{1}{x^2}}}{2} \quad (1)$$

Verification of solutions

$$y = \frac{\left(c_1 \cos \left(-\frac{3 \ln(2)}{2} + \frac{3 \ln\left(-\frac{1}{x^2}\right)}{2} \right) + c_2 \sin \left(-\frac{3 \ln(2)}{2} + \frac{3 \ln\left(-\frac{1}{x^2}\right)}{2} \right) \right) \sqrt{2} \sqrt{-\frac{1}{x^2}}}{2}$$

Verified OK.

9.25.3 Solving as second order change of variable on x method 1 ode

In normal form the ode

$$x^2 y'' + 3xy' + 10y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{3}{x}$$
$$q(x) = \frac{10}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) results

$$\frac{d^2}{d\tau^2}y(\tau) + p_1 \left(\frac{d}{d\tau}y(\tau) \right) + q_1 y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $q_1 = c^2$ where c is some constant. Therefore from (5)

$$\tau' = \frac{1}{c} \sqrt{q}$$
$$= \frac{\sqrt{10} \sqrt{\frac{1}{x^2}}}{c} \quad (6)$$
$$\tau'' = -\frac{\sqrt{10}}{c \sqrt{\frac{1}{x^2}} x^3}$$

Substituting the above into (4) results in

$$\begin{aligned}
 p_1(\tau) &= \frac{\tau''(x) + p(x) \tau'(x)}{\tau'(x)^2} \\
 &= \frac{-\frac{\sqrt{10}}{c\sqrt{\frac{1}{x^2}}x^3} + \frac{3}{x}\frac{\sqrt{10}\sqrt{\frac{1}{x^2}}}{c}}{\left(\frac{\sqrt{10}\sqrt{\frac{1}{x^2}}}{c}\right)^2} \\
 &= \frac{c\sqrt{10}}{5}
 \end{aligned}$$

Therefore ode (3) now becomes

$$\begin{aligned}
 y(\tau)'' + p_1 y(\tau)' + q_1 y(\tau) &= 0 \\
 \frac{d^2}{d\tau^2} y(\tau) + \frac{c\sqrt{10}}{5} \left(\frac{d}{d\tau} y(\tau)\right) + c^2 y(\tau) &= 0
 \end{aligned} \tag{7}$$

The above ode is now solved for $y(\tau)$. Since the ode is now constant coefficients, it can be easily solved to give

$$y(\tau) = e^{-\frac{\sqrt{10}c\tau}{10}} \left(c_1 \cos\left(\frac{3\sqrt{10}c\tau}{10}\right) + c_2 \sin\left(\frac{3\sqrt{10}c\tau}{10}\right) \right)$$

Now from (6)

$$\begin{aligned}
 \tau &= \int \frac{1}{c} \sqrt{q} dx \\
 &= \frac{\int \sqrt{10} \sqrt{\frac{1}{x^2}} dx}{c} \\
 &= \frac{\sqrt{10} \sqrt{\frac{1}{x^2}} x \ln(x)}{c}
 \end{aligned}$$

Substituting the above into the solution obtained gives

$$y = \frac{c_1 \cos(3 \ln(x)) + c_2 \sin(3 \ln(x))}{x}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1 \cos(3 \ln(x)) + c_2 \sin(3 \ln(x))}{x} \tag{1}$$

Verification of solutions

$$y = \frac{c_1 \cos(3 \ln(x)) + c_2 \sin(3 \ln(x))}{x}$$

Verified OK.

9.25.4 Solving as second order change of variable on y method 2 ode

In normal form the ode

$$x^2 y'' + 3xy' + 10y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{3}{x}$$
$$q(x) = \frac{10}{x^2}$$

Applying change of variables on the dependent variable $y = v(x)x^n$ to (2) gives the following ode where the dependent variable is $v(x)$ and not y .

$$v''(x) + \left(\frac{2n}{x} + p\right)v'(x) + \left(\frac{n(n-1)}{x^2} + \frac{np}{x} + q\right)v(x) = 0 \quad (3)$$

Let the coefficient of $v(x)$ above be zero. Hence

$$\frac{n(n-1)}{x^2} + \frac{np}{x} + q = 0 \quad (4)$$

Substituting the earlier values found for $p(x)$ and $q(x)$ into (4) gives

$$\frac{n(n-1)}{x^2} + \frac{3n}{x^2} + \frac{10}{x^2} = 0 \quad (5)$$

Solving (5) for n gives

$$n = -1 + 3i \quad (6)$$

Substituting this value in (3) gives

$$v''(x) + \left(\frac{-2 + 6i}{x} + \frac{3}{x}\right)v'(x) = 0$$
$$v''(x) + \frac{(1 + 6i)v'(x)}{x} = 0 \quad (7)$$

Using the substitution

$$u(x) = v'(x)$$

Then (7) becomes

$$u'(x) + \frac{(1 + 6i)u(x)}{x} = 0 \quad (8)$$

The above is now solved for $u(x)$. In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= \frac{(-1 - 6i)u}{x} \end{aligned}$$

Where $f(x) = \frac{-1-6i}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned} \frac{1}{u} du &= \frac{-1 - 6i}{x} dx \\ \int \frac{1}{u} du &= \int \frac{-1 - 6i}{x} dx \\ \ln(u) &= (-1 - 6i) \ln(x) + c_1 \\ u &= e^{(-1-6i)\ln(x)+c_1} \\ &= c_1 e^{(-1-6i)\ln(x)} \end{aligned}$$

Which simplifies to

$$u(x) = \frac{c_1 x^{-6i}}{x}$$

Now that $u(x)$ is known, then

$$\begin{aligned} v'(x) &= u(x) \\ v(x) &= \int u(x) dx + c_2 \\ &= \frac{ic_1 x^{-6i}}{6} + c_2 \end{aligned}$$

Hence

$$\begin{aligned} y &= v(x) x^n \\ &= \left(\frac{ic_1 x^{-6i}}{6} + c_2 \right) x^{-1+3i} \\ &= c_2 x^{-1+3i} + \frac{ic_1 x^{-1-3i}}{6} \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \left(\frac{ic_1 x^{-6i}}{6} + c_2 \right) x^{-1+3i} \quad (1)$$

Verification of solutions

$$y = \left(\frac{ic_1 x^{-6i}}{6} + c_2 \right) x^{-1+3i}$$

Verified OK.

9.25.5 Solving using Kovacic algorithm

Writing the ode as

$$x^2 y'' + 3xy' + 10y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= x^2 \\ B &= 3x \\ C &= 10 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-37}{4x^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -37 \\ t &= 4x^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(-\frac{37}{4x^2} \right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 223: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole

larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = -\frac{37}{4x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = -\frac{37}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{1}{2} + 3i \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{2} - 3i \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = -\frac{37}{4x^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = -\frac{37}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{1}{2} + 3i \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{2} - 3i \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = -\frac{37}{4x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	$\frac{1}{2} + 3i$	$\frac{1}{2} - 3i$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	$\frac{1}{2} + 3i$	$\frac{1}{2} - 3i$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = \frac{1}{2} - 3i$ then

$$\begin{aligned} d &= \alpha_\infty^- - (\alpha_{c_1}^-) \\ &= \frac{1}{2} - 3i - \left(\frac{1}{2} - 3i \right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{x - c_1} \right) + (-)[\sqrt{r}]_\infty \\ &= \frac{\frac{1}{2} - 3i}{x} + (-)(0) \\ &= \frac{\frac{1}{2} - 3i}{x} \\ &= \frac{\frac{1}{2} - 3i}{x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$(0) + 2\left(\frac{\frac{1}{2} - 3i}{x}\right)(0) + \left(\left(\frac{-\frac{1}{2} + 3i}{x^2}\right) + \left(\frac{\frac{1}{2} - 3i}{x}\right)^2 - \left(-\frac{37}{4x^2}\right)\right) = 0$$

$$0 = 0$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= pe^{\int \omega dx} \\ &= e^{\int \frac{\frac{1}{2} - 3i}{x} dx} \\ &= x^{\frac{1}{2} - 3i} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{3x}{x^2} dx} \\ &= z_1 e^{-\frac{3 \ln(x)}{2}} \\ &= z_1 \left(\frac{1}{x^{\frac{3}{2}}}\right) \end{aligned}$$

Which simplifies to

$$y_1 = x^{-1-3i}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{3x}{x^2} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-3 \ln(x)}}{(y_1)^2} dx \\ &= y_1 \left(-\frac{ix^{6i}}{6}\right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (x^{-1-3i}) + c_2 \left(x^{-1-3i} \left(-\frac{i x^{6i}}{6} \right) \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 x^{-1-3i} - \frac{i c_2 x^{-1+3i}}{6} \quad (1)$$

Verification of solutions

$$y = c_1 x^{-1-3i} - \frac{i c_2 x^{-1+3i}}{6}$$

Verified OK.

9.25.6 Maple step by step solution

Let's solve

$$x^2 y'' + 3xy' + 10y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{3y'}{x} - \frac{10y}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{3y'}{x} + \frac{10y}{x^2} = 0$$

- Multiply by denominators of the ODE

$$x^2 y'' + 3xy' + 10y = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x , using the chain rule

$$y' = \left(\frac{d}{dt} y(t) \right) t'(x)$$

- Compute derivative

$$y' = \frac{d}{dt} y(t)$$

- Calculate the 2nd derivative of y with respect to x , using the chain rule

$$y'' = \left(\frac{d^2}{dt^2} y(t) \right) t'(x)^2 + t''(x) \left(\frac{d}{dt} y(t) \right)$$

- Compute derivative

$$y'' = \frac{\frac{d^2}{dt^2} y(t)}{x^2} - \frac{\frac{d}{dt} y(t)}{x^2}$$

Substitute the change of variables back into the ODE

$$x^2 \left(\frac{\frac{d^2}{dt^2} y(t)}{x^2} - \frac{\frac{d}{dt} y(t)}{x^2} \right) + 3 \frac{d}{dt} y(t) + 10y(t) = 0$$

- Simplify

$$\frac{d^2}{dt^2} y(t) + 2 \frac{d}{dt} y(t) + 10y(t) = 0$$

- Characteristic polynomial of ODE

$$r^2 + 2r + 10 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-2) \pm (\sqrt{-36})}{2}$$

- Roots of the characteristic polynomial

$$r = (-1 - 3I, -1 + 3I)$$

- 1st solution of the ODE

$$y_1(t) = e^{-t} \cos(3t)$$

- 2nd solution of the ODE

$$y_2(t) = e^{-t} \sin(3t)$$

- General solution of the ODE

$$y(t) = c_1 y_1(t) + c_2 y_2(t)$$

- Substitute in solutions

$$y(t) = c_1 e^{-t} \cos(3t) + c_2 e^{-t} \sin(3t)$$

- Change variables back using $t = \ln(x)$

$$y = \frac{c_1 \cos(3 \ln(x))}{x} + \frac{c_2 \sin(3 \ln(x))}{x}$$

- Simplify

$$y = \frac{c_1 \cos(3 \ln(x))}{x} + \frac{c_2 \sin(3 \ln(x))}{x}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
<- LODE of Euler type successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(x^2*diff(y(x),x$2)+3*x*diff(y(x),x)+10*y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{c_1 \sin(3 \ln(x)) + c_2 \cos(3 \ln(x))}{x}$$

✓ Solution by Mathematica

Time used: 0.029 (sec). Leaf size: 26

```
DSolve[x^2*y'[x]+3*x*y'[x]+10*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_2 \cos(3 \log(x)) + c_1 \sin(3 \log(x))}{x}$$

9.26 problem 5(b)

9.26.1 Solving as second order euler ode ode	1517
9.26.2 Solving as second order change of variable on x method 2 ode .	1518
9.26.3 Solving as second order change of variable on x method 1 ode .	1521
9.26.4 Solving as second order change of variable on y method 2 ode .	1523
9.26.5 Solving using Kovacic algorithm	1525
9.26.6 Maple step by step solution	1530

Internal problem ID [6294]

Internal file name [OUTPUT/5542_Sunday_June_05_2022_03_43_02_PM_19522388/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.1. Linear Equations with Constant Coefficients. Page 62

Problem number: 5(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_euler_ode", "second_order_change_of_variable_on_x_method_1", "second_order_change_of_variable_on_x_method_2", "second_order_change_of_variable_on_y_method_2"

Maple gives the following as the ode type

```
[[_Emden, _Fowler]]
```

$$2x^2y'' + 10xy' + 8y = 0$$

9.26.1 Solving as second order euler ode ode

This is Euler second order ODE. Let the solution be $y = x^r$, then $y' = rx^{r-1}$ and $y'' = r(r-1)x^{r-2}$. Substituting these back into the given ODE gives

$$2x^2(r(r-1))x^{r-2} + 10rxr^{r-1} + 8x^r = 0$$

Simplifying gives

$$2r(r-1)x^r + 10rx^r + 8x^r = 0$$

Since $x^r \neq 0$ then dividing throughout by x^r gives

$$2r(r - 1) + 10r + 8 = 0$$

Or

$$2r^2 + 8r + 8 = 0 \quad (1)$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = -2$$

$$r_2 = -2$$

Since the roots are equal, then the general solution is

$$y = c_1 y_1 + c_2 y_2$$

Where $y_1 = x^r$ and $y_2 = x^r \ln(x)$. Hence

$$y = \frac{c_1}{x^2} + \frac{c_2 \ln(x)}{x^2}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1}{x^2} + \frac{c_2 \ln(x)}{x^2} \quad (1)$$

Verification of solutions

$$y = \frac{c_1}{x^2} + \frac{c_2 \ln(x)}{x^2}$$

Verified OK.

9.26.2 Solving as second order change of variable on x method 2 ode

In normal form the ode

$$2x^2 y'' + 10xy' + 8y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{5}{x}$$
$$q(x) = \frac{4}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x)\tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned} \tau &= \int e^{-(\int p(x)dx)} dx \\ &= \int e^{-(\int \frac{5}{x} dx)} dx \\ &= \int e^{-5\ln(x)} dx \\ &= \int \frac{1}{x^5} dx \\ &= -\frac{1}{4x^4} \end{aligned} \quad (6)$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned} q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\ &= \frac{\frac{4}{x^2}}{\frac{1}{x^{10}}} \\ &= 4x^8 \end{aligned} \quad (7)$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned} \frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) + 4x^8y(\tau) &= 0 \end{aligned}$$

But in terms of τ

$$4x^8 = \frac{1}{4\tau^2}$$

Hence the above ode becomes

$$\frac{d^2}{d\tau^2}y(\tau) + \frac{y(\tau)}{4\tau^2} = 0$$

The above ode is now solved for $y(\tau)$. The ode can be written as

$$4\left(\frac{d^2}{d\tau^2}y(\tau)\right)\tau^2 + y(\tau) = 0$$

Which shows it is a Euler ODE. This is Euler second order ODE. Let the solution be $y(\tau) = \tau^r$, then $y' = r\tau^{r-1}$ and $y'' = r(r-1)\tau^{r-2}$. Substituting these back into the given ODE gives

$$4\tau^2(r(r-1))\tau^{r-2} + 0r\tau^{r-1} + \tau^r = 0$$

Simplifying gives

$$4r(r-1)\tau^r + 0\tau^r + \tau^r = 0$$

Since $\tau^r \neq 0$ then dividing throughout by τ^r gives

$$4r(r-1) + 0 + 1 = 0$$

Or

$$4r^2 - 4r + 1 = 0 \tag{1}$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = \frac{1}{2}$$
$$r_2 = \frac{1}{2}$$

Since the roots are equal, then the general solution is

$$y(\tau) = c_1y_1 + c_2y_2$$

Where $y_1 = \tau^r$ and $y_2 = \tau^r \ln(\tau)$. Hence

$$y(\tau) = c_1\sqrt{\tau} + c_2\sqrt{\tau} \ln(\tau)$$

The above solution is now transformed back to y using (6) which results in

$$y = \frac{(-2c_2 \ln(2) + c_2 \ln(-\frac{1}{x^4}) + c_1) \sqrt{-\frac{1}{x^4}}}{2}$$

Summary

The solution(s) found are the following

$$y = \frac{(-2c_2 \ln(2) + c_2 \ln(-\frac{1}{x^4}) + c_1) \sqrt{-\frac{1}{x^4}}}{2} \quad (1)$$

Verification of solutions

$$y = \frac{(-2c_2 \ln(2) + c_2 \ln(-\frac{1}{x^4}) + c_1) \sqrt{-\frac{1}{x^4}}}{2}$$

Verified OK.

9.26.3 Solving as second order change of variable on x method 1 ode

In normal form the ode

$$2x^2y'' + 10xy' + 8y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{5}{x}$$
$$q(x) = \frac{4}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) results

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $q_1 = c^2$ where c is some constant. Therefore from (5)

$$\tau' = \frac{1}{c}\sqrt{q}$$
$$= \frac{2\sqrt{\frac{1}{x^2}}}{c} \quad (6)$$
$$\tau'' = -\frac{2}{c\sqrt{\frac{1}{x^2}}x^3}$$

Substituting the above into (4) results in

$$\begin{aligned}
 p_1(\tau) &= \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \\
 &= \frac{-\frac{2}{c\sqrt{\frac{1}{x^2}}x^3} + \frac{5}{x}\frac{2\sqrt{\frac{1}{x^2}}}{c}}{\left(\frac{2\sqrt{\frac{1}{x^2}}}{c}\right)^2} \\
 &= 2c
 \end{aligned}$$

Therefore ode (3) now becomes

$$\begin{aligned}
 y(\tau)'' + p_1y(\tau)' + q_1y(\tau) &= 0 \\
 \frac{d^2}{d\tau^2}y(\tau) + 2c\left(\frac{d}{d\tau}y(\tau)\right) + c^2y(\tau) &= 0 \tag{7}
 \end{aligned}$$

The above ode is now solved for $y(\tau)$. Since the ode is now constant coefficients, it can be easily solved to give

$$y(\tau) = e^{-c\tau} c_1$$

Now from (6)

$$\begin{aligned}
 \tau &= \int \frac{1}{c}\sqrt{q} dx \\
 &= \frac{\int 2\sqrt{\frac{1}{x^2}} dx}{c} \\
 &= \frac{2\sqrt{\frac{1}{x^2}} x \ln(x)}{c}
 \end{aligned}$$

Substituting the above into the solution obtained gives

$$y = \frac{c_1}{x^2}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1}{x^2} \tag{1}$$

Verification of solutions

$$y = \frac{c_1}{x^2}$$

Verified OK.

9.26.4 Solving as second order change of variable on y method 2 ode

In normal form the ode

$$2x^2y'' + 10xy' + 8y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{5}{x}$$
$$q(x) = \frac{4}{x^2}$$

Applying change of variables on the dependent variable $y = v(x)x^n$ to (2) gives the following ode where the dependent variables is $v(x)$ and not y .

$$v''(x) + \left(\frac{2n}{x} + p\right)v'(x) + \left(\frac{n(n-1)}{x^2} + \frac{np}{x} + q\right)v(x) = 0 \quad (3)$$

Let the coefficient of $v(x)$ above be zero. Hence

$$\frac{n(n-1)}{x^2} + \frac{np}{x} + q = 0 \quad (4)$$

Substituting the earlier values found for $p(x)$ and $q(x)$ into (4) gives

$$\frac{n(n-1)}{x^2} + \frac{5n}{x^2} + \frac{4}{x^2} = 0 \quad (5)$$

Solving (5) for n gives

$$n = -2 \quad (6)$$

Substituting this value in (3) gives

$$v''(x) + \frac{v'(x)}{x} = 0$$
$$v''(x) + \frac{v'(x)}{x} = 0 \quad (7)$$

Using the substitution

$$u(x) = v'(x)$$

Then (7) becomes

$$u'(x) + \frac{u(x)}{x} = 0 \quad (8)$$

The above is now solved for $u(x)$. In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{u}{x} \end{aligned}$$

Where $f(x) = -\frac{1}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned} \frac{1}{u} du &= -\frac{1}{x} dx \\ \int \frac{1}{u} du &= \int -\frac{1}{x} dx \\ \ln(u) &= -\ln(x) + c_1 \\ u &= e^{-\ln(x)+c_1} \\ &= \frac{c_1}{x} \end{aligned}$$

Now that $u(x)$ is known, then

$$\begin{aligned} v'(x) &= u(x) \\ v(x) &= \int u(x) dx + c_2 \\ &= c_1 \ln(x) + c_2 \end{aligned}$$

Hence

$$\begin{aligned} y &= v(x) x^n \\ &= \frac{c_1 \ln(x) + c_2}{x^2} \\ &= \frac{c_1 \ln(x) + c_2}{x^2} \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1 \ln(x) + c_2}{x^2} \quad (1)$$

Verification of solutions

$$y = \frac{c_1 \ln(x) + c_2}{x^2}$$

Verified OK.

9.26.5 Solving using Kovacic algorithm

Writing the ode as

$$2x^2y'' + 10xy' + 8y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 2x^2 \\ B &= 10x \\ C &= 8 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{4x^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 4x^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(-\frac{1}{4x^2}\right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 225: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = -\frac{1}{4x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = -\frac{1}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{1}{2} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{2} \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = -\frac{1}{4x^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = -\frac{1}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{1}{2} \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{2} \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = -\frac{1}{4x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	$\frac{1}{2}$	$\frac{1}{2}$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	$\frac{1}{2}$	$\frac{1}{2}$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to

determine possible non negative integer d from these using

$$d = \alpha_{\infty}^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_{∞}^{\pm} . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_{\infty}^{-} = \frac{1}{2}$ then

$$\begin{aligned} d &= \alpha_{\infty}^{-} - (\alpha_{c_1}^{-}) \\ &= \frac{1}{2} - \left(\frac{1}{2}\right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_{\infty}$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^{-}}{x - c_1} \right) + (-)[\sqrt{r}]_{\infty} \\ &= \frac{1}{2x} + (-)(0) \\ &= \frac{1}{2x} \\ &= \frac{1}{2x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2\left(\frac{1}{2x}\right)(0) + \left(\left(-\frac{1}{2x^2}\right) + \left(\frac{1}{2x}\right)^2 - \left(-\frac{1}{4x^2}\right) \right) &= 0 \\ 0 &= 0 \end{aligned}$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned}z_1(x) &= pe^{\int \omega dx} \\ &= e^{\int \frac{1}{2x} dx} \\ &= \sqrt{x}\end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{10x}{2x^2} dx} \\ &= z_1 e^{-\frac{5 \ln(x)}{2}} \\ &= z_1 \left(\frac{1}{x^{\frac{5}{2}}} \right)\end{aligned}$$

Which simplifies to

$$y_1 = \frac{1}{x^2}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{10x}{2x^2} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-5 \ln(x)}}{(y_1)^2} dx \\ &= y_1 (\ln(x))\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(\frac{1}{x^2} \right) + c_2 \left(\frac{1}{x^2} (\ln(x)) \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1}{x^2} + \frac{c_2 \ln(x)}{x^2} \quad (1)$$

Verification of solutions

$$y = \frac{c_1}{x^2} + \frac{c_2 \ln(x)}{x^2}$$

Verified OK.

9.26.6 Maple step by step solution

Let's solve

$$2x^2y'' + 10xy' + 8y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{5y'}{x} - \frac{4y}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{5y'}{x} + \frac{4y}{x^2} = 0$$

- Multiply by denominators of the ODE

$$x^2y'' + 5xy' + 4y = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x , using the chain rule

$$y' = \left(\frac{d}{dt}y(t)\right) t'(x)$$

- Compute derivative

$$y' = \frac{\frac{d}{dt}y(t)}{x}$$

- Calculate the 2nd derivative of y with respect to x , using the chain rule

$$y'' = \left(\frac{d^2}{dt^2}y(t)\right) t'(x)^2 + t''(x) \left(\frac{d}{dt}y(t)\right)$$

- Compute derivative

$$y'' = \frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2}$$

Substitute the change of variables back into the ODE

$$x^2 \left(\frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2} \right) + 5 \frac{d}{dt}y(t) + 4y(t) = 0$$

- Simplify

$$\frac{d^2}{dt^2}y(t) + 4 \frac{d}{dt}y(t) + 4y(t) = 0$$

- Characteristic polynomial of ODE

$$r^2 + 4r + 4 = 0$$

- Factor the characteristic polynomial

$$(r + 2)^2 = 0$$

- Root of the characteristic polynomial

$$r = -2$$

- 1st solution of the ODE

$$y_1(t) = e^{-2t}$$

- Repeated root, multiply $y_1(t)$ by t to ensure linear independence

$$y_2(t) = t e^{-2t}$$

- General solution of the ODE

$$y(t) = c_1 y_1(t) + c_2 y_2(t)$$

- Substitute in solutions

$$y(t) = c_1 e^{-2t} + c_2 t e^{-2t}$$

- Change variables back using $t = \ln(x)$

$$y = \frac{c_1}{x^2} + \frac{c_2 \ln(x)}{x^2}$$

- Simplify

$$y = \frac{c_1}{x^2} + \frac{c_2 \ln(x)}{x^2}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
<- LODE of Euler type successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(2*x^2*diff(y(x),x$2)+10*x*diff(y(x),x)+8*y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{c_2 \ln(x) + c_1}{x^2}$$

✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 18

```
DSolve[2*x^2*y''[x]+10*x*y'[x]+8*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2c_2 \log(x) + c_1}{x^2}$$

9.27 problem 5(c)

9.27.1 Solving as second order euler ode ode	1533
9.27.2 Solving as second order change of variable on x method 2 ode .	1534
9.27.3 Solving as second order change of variable on y method 2 ode .	1537
9.27.4 Solving using Kovacic algorithm	1539
9.27.5 Maple step by step solution	1544

Internal problem ID [6295]

Internal file name [OUTPUT/5543_Sunday_June_05_2022_03_43_03_PM_17006049/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.1. Linear Equations with Constant Coefficients. Page 62

Problem number: 5(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_euler_ode", "second_order_change_of_variable_on_x_method_2", "second_order_change_of_variable_on_y_method_2"

Maple gives the following as the ode type

```
[[_Emden , _Fowler]]
```

$$x^2y'' + 2xy' - 12y = 0$$

9.27.1 Solving as second order euler ode ode

This is Euler second order ODE. Let the solution be $y = x^r$, then $y' = rx^{r-1}$ and $y'' = r(r-1)x^{r-2}$. Substituting these back into the given ODE gives

$$x^2(r(r-1))x^{r-2} + 2rxr^{r-1} - 12x^r = 0$$

Simplifying gives

$$r(r-1)x^r + 2rx^r - 12x^r = 0$$

Since $x^r \neq 0$ then dividing throughout by x^r gives

$$r(r-1) + 2r - 12 = 0$$

Or

$$r^2 + r - 12 = 0 \quad (1)$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = -4$$

$$r_2 = 3$$

Since the roots are real and distinct, then the general solution is

$$y = c_1 y_1 + c_2 y_2$$

Where $y_1 = x^{r_1}$ and $y_2 = x^{r_2}$. Hence

$$y = \frac{c_1}{x^4} + c_2 x^3$$

Summary

The solution(s) found are the following

$$y = \frac{c_1}{x^4} + c_2 x^3 \quad (1)$$

Verification of solutions

$$y = \frac{c_1}{x^4} + c_2 x^3$$

Verified OK.

9.27.2 Solving as second order change of variable on x method 2 ode

In normal form the ode

$$x^2 y'' + 2xy' - 12y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{2}{x}$$
$$q(x) = -\frac{12}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x)\tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned} \tau &= \int e^{-(\int p(x)dx)} dx \\ &= \int e^{-(\int \frac{2}{x} dx)} dx \\ &= \int e^{-2\ln(x)} dx \\ &= \int \frac{1}{x^2} dx \\ &= -\frac{1}{x} \end{aligned} \quad (6)$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned} q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\ &= \frac{-\frac{12}{x^2}}{\frac{1}{x^4}} \\ &= -12x^2 \end{aligned} \quad (7)$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned} \frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) - 12x^2y(\tau) &= 0 \end{aligned}$$

But in terms of τ

$$-12x^2 = -\frac{12}{\tau^2}$$

Hence the above ode becomes

$$\frac{d^2}{d\tau^2}y(\tau) - \frac{12y(\tau)}{\tau^2} = 0$$

The above ode is now solved for $y(\tau)$. The ode can be written as

$$\left(\frac{d^2}{d\tau^2}y(\tau)\right)\tau^2 - 12y(\tau) = 0$$

Which shows it is a Euler ODE. This is Euler second order ODE. Let the solution be $y(\tau) = \tau^r$, then $y' = r\tau^{r-1}$ and $y'' = r(r-1)\tau^{r-2}$. Substituting these back into the given ODE gives

$$\tau^2(r(r-1))\tau^{r-2} + 0r\tau^{r-1} - 12\tau^r = 0$$

Simplifying gives

$$r(r-1)\tau^r + 0\tau^r - 12\tau^r = 0$$

Since $\tau^r \neq 0$ then dividing throughout by τ^r gives

$$r(r-1) + 0 - 12 = 0$$

Or

$$r^2 - r - 12 = 0 \tag{1}$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = -3$$

$$r_2 = 4$$

Since the roots are real and distinct, then the general solution is

$$y(\tau) = c_1y_1 + c_2y_2$$

Where $y_1 = \tau^{r_1}$ and $y_2 = \tau^{r_2}$. Hence

$$y(\tau) = \frac{c_1}{\tau^3} + c_2\tau^4$$

The above solution is now transformed back to y using (6) which results in

$$y = \frac{-c_1x^7 + c_2}{x^4}$$

Summary

The solution(s) found are the following

$$y = \frac{-c_1 x^7 + c_2}{x^4} \quad (1)$$

Verification of solutions

$$y = \frac{-c_1 x^7 + c_2}{x^4}$$

Verified OK.

9.27.3 Solving as second order change of variable on y method 2 ode

In normal form the ode

$$x^2 y'' + 2xy' - 12y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{2}{x}$$
$$q(x) = -\frac{12}{x^2}$$

Applying change of variables on the dependent variable $y = v(x)x^n$ to (2) gives the following ode where the dependent variables is $v(x)$ and not y .

$$v''(x) + \left(\frac{2n}{x} + p\right)v'(x) + \left(\frac{n(n-1)}{x^2} + \frac{np}{x} + q\right)v(x) = 0 \quad (3)$$

Let the coefficient of $v(x)$ above be zero. Hence

$$\frac{n(n-1)}{x^2} + \frac{np}{x} + q = 0 \quad (4)$$

Substituting the earlier values found for $p(x)$ and $q(x)$ into (4) gives

$$\frac{n(n-1)}{x^2} + \frac{2n}{x^2} - \frac{12}{x^2} = 0 \quad (5)$$

Solving (5) for n gives

$$n = 3 \quad (6)$$

Substituting this value in (3) gives

$$\begin{aligned}v''(x) + \frac{8v'(x)}{x} &= 0 \\v''(x) + \frac{8v'(x)}{x} &= 0\end{aligned}\tag{7}$$

Using the substitution

$$u(x) = v'(x)$$

Then (7) becomes

$$u'(x) + \frac{8u(x)}{x} = 0\tag{8}$$

The above is now solved for $u(x)$. In canonical form the ODE is

$$\begin{aligned}u' &= F(x, u) \\&= f(x)g(u) \\&= -\frac{8u}{x}\end{aligned}$$

Where $f(x) = -\frac{8}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u} du &= -\frac{8}{x} dx \\ \int \frac{1}{u} du &= \int -\frac{8}{x} dx \\ \ln(u) &= -8 \ln(x) + c_1 \\ u &= e^{-8 \ln(x) + c_1} \\ &= \frac{c_1}{x^8}\end{aligned}$$

Now that $u(x)$ is known, then

$$\begin{aligned}v'(x) &= u(x) \\ v(x) &= \int u(x) dx + c_2 \\ &= -\frac{c_1}{7x^7} + c_2\end{aligned}$$

Hence

$$\begin{aligned}y &= v(x) x^n \\&= \left(-\frac{c_1}{7x^7} + c_2\right) x^3 \\&= \frac{7c_2x^7 - c_1}{7x^4}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \left(-\frac{c_1}{7x^7} + c_2\right) x^3 \quad (1)$$

Verification of solutions

$$y = \left(-\frac{c_1}{7x^7} + c_2\right) x^3$$

Verified OK.

9.27.4 Solving using Kovacic algorithm

Writing the ode as

$$x^2y'' + 2xy' - 12y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= x^2 \\B &= 2x \\C &= -12\end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\&= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{12}{x^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 12 \\ t &= x^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{12}{x^2}\right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 227: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = \frac{12}{x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = 12$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = 4 \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = -3 \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = \frac{12}{x^2}$$

Since the $\text{gcd}(s, t) = 1$. This gives $b = 12$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = 4 \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = -3 \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{12}{x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	4	-3

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	4	-3

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = -3$ then

$$\begin{aligned} d &= \alpha_\infty^- - (\alpha_{c_1}^-) \\ &= -3 - (-3) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{x - c_1} \right) + (-)[\sqrt{r}]_\infty \\ &= -\frac{3}{x} + (-)(0) \\ &= -\frac{3}{x} \\ &= -\frac{3}{x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$(0) + 2\left(-\frac{3}{x}\right)(0) + \left(\left(\frac{3}{x^2}\right) + \left(-\frac{3}{x}\right)^2 - \left(\frac{12}{x^2}\right)\right) = 0$$
$$0 = 0$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$z_1(x) = pe^{\int \omega dx}$$
$$= e^{\int -\frac{3}{x} dx}$$
$$= \frac{1}{x^3}$$

The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$
$$= z_1 e^{-\int \frac{1}{2} \frac{2x}{x^2} dx}$$
$$= z_1 e^{-\ln(x)}$$
$$= z_1 \left(\frac{1}{x}\right)$$

Which simplifies to

$$y_1 = \frac{1}{x^4}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$y_2 = y_1 \int \frac{e^{\int -\frac{2x}{x^2} dx}}{(y_1)^2} dx$$
$$= y_1 \int \frac{e^{-2\ln(x)}}{(y_1)^2} dx$$
$$= y_1 \left(\frac{x^7}{7}\right)$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(\frac{1}{x^4} \right) + c_2 \left(\frac{1}{x^4} \left(\frac{x^7}{7} \right) \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1}{x^4} + \frac{c_2 x^3}{7} \quad (1)$$

Verification of solutions

$$y = \frac{c_1}{x^4} + \frac{c_2 x^3}{7}$$

Verified OK.

9.27.5 Maple step by step solution

Let's solve

$$x^2 y'' + 2xy' - 12y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{2y'}{x} + \frac{12y}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{2y'}{x} - \frac{12y}{x^2} = 0$$

- Multiply by denominators of the ODE

$$x^2 y'' + 2xy' - 12y = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x , using the chain rule

$$y' = \left(\frac{d}{dt} y(t) \right) t'(x)$$

- Compute derivative

$$y' = \frac{\frac{d}{dt}y(t)}{x}$$

- Calculate the 2nd derivative of y with respect to x , using the chain rule

$$y'' = \left(\frac{d^2}{dt^2}y(t) \right) t'(x)^2 + t''(x) \left(\frac{d}{dt}y(t) \right)$$

- Compute derivative

$$y'' = \frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2}$$

Substitute the change of variables back into the ODE

$$x^2 \left(\frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2} \right) + 2 \frac{d}{dt}y(t) - 12y(t) = 0$$

- Simplify

$$\frac{d^2}{dt^2}y(t) + \frac{d}{dt}y(t) - 12y(t) = 0$$

- Characteristic polynomial of ODE

$$r^2 + r - 12 = 0$$

- Factor the characteristic polynomial

$$(r + 4)(r - 3) = 0$$

- Roots of the characteristic polynomial

$$r = (-4, 3)$$

- 1st solution of the ODE

$$y_1(t) = e^{-4t}$$

- 2nd solution of the ODE

$$y_2(t) = e^{3t}$$

- General solution of the ODE

$$y(t) = c_1 y_1(t) + c_2 y_2(t)$$

- Substitute in solutions

$$y(t) = c_1 e^{-4t} + c_2 e^{3t}$$

- Change variables back using $t = \ln(x)$

$$y = \frac{c_1}{x^4} + c_2 x^3$$

- Simplify

$$y = \frac{c_1}{x^4} + c_2 x^3$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
<- LODE of Euler type successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 15

```
dsolve(x^2*diff(y(x),x$2)+2*x*diff(y(x),x)-12*y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{c_1 x^7 + c_2}{x^4}$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 18

```
DSolve[x^2*y''[x]+2*x*y'[x]-12*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_2 x^7 + c_1}{x^4}$$

9.28 problem 5(d)

- 9.28.1 Solving as second order euler ode ode 1547
- 9.28.2 Solving using Kovacic algorithm 1548
- 9.28.3 Maple step by step solution 1553

Internal problem ID [6296]

Internal file name [OUTPUT/5544_Sunday_June_05_2022_03_43_04_PM_95902784/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.1. Linear Equations with Constant Coefficients. Page 62

Problem number: 5(d).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_euler_ode"

Maple gives the following as the ode type

```
[[_Emden , _Fowler]]
```

$$4x^2y'' - 3y = 0$$

9.28.1 Solving as second order euler ode ode

This is Euler second order ODE. Let the solution be $y = x^r$, then $y' = rx^{r-1}$ and $y'' = r(r-1)x^{r-2}$. Substituting these back into the given ODE gives

$$4x^2(r(r-1))x^{r-2} + 0rx^{r-1} - 3x^r = 0$$

Simplifying gives

$$4r(r-1)x^r + 0x^r - 3x^r = 0$$

Since $x^r \neq 0$ then dividing throughout by x^r gives

$$4r(r-1) + 0 - 3 = 0$$

Or

$$4r^2 - 4r - 3 = 0 \tag{1}$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = -\frac{1}{2}$$

$$r_2 = \frac{3}{2}$$

Since the roots are real and distinct, then the general solution is

$$y = c_1 y_1 + c_2 y_2$$

Where $y_1 = x^{r_1}$ and $y_2 = x^{r_2}$. Hence

$$y = \frac{c_1}{\sqrt{x}} + c_2 x^{\frac{3}{2}}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1}{\sqrt{x}} + c_2 x^{\frac{3}{2}} \quad (1)$$

Verification of solutions

$$y = \frac{c_1}{\sqrt{x}} + c_2 x^{\frac{3}{2}}$$

Verified OK.

9.28.2 Solving using Kovacic algorithm

Writing the ode as

$$4x^2 y'' - 3y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$A = 4x^2$$

$$B = 0$$

$$C = -3 \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{3}{4x^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 3 \\ t &= 4x^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{3}{4x^2} \right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 229: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = \frac{3}{4x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = \frac{3}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{3}{2} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = -\frac{1}{2} \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = \frac{3}{4x^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = \frac{3}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{3}{2} \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = -\frac{1}{2} \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{3}{4x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	$\frac{3}{2}$	$-\frac{1}{2}$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	$\frac{3}{2}$	$-\frac{1}{2}$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = -\frac{1}{2}$ then

$$\begin{aligned} d &= \alpha_\infty^- - (\alpha_{c_1}^-) \\ &= -\frac{1}{2} - \left(-\frac{1}{2}\right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{x - c_1} \right) + (-)[\sqrt{r}]_\infty \\ &= -\frac{1}{2x} + (-)(0) \\ &= -\frac{1}{2x} \\ &= -\frac{1}{2x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$(0) + 2\left(-\frac{1}{2x}\right)(0) + \left(\left(\frac{1}{2x^2}\right) + \left(-\frac{1}{2x}\right)^2 - \left(\frac{3}{4x^2}\right)\right) = 0$$

$$0 = 0$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= pe^{\int \omega dx} \\ &= e^{\int -\frac{1}{2x} dx} \\ &= \frac{1}{\sqrt{x}} \end{aligned}$$

The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned} y_1 &= z_1 \\ &= \frac{1}{\sqrt{x}} \end{aligned}$$

Which simplifies to

$$y_1 = \frac{1}{\sqrt{x}}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned}y_2 &= y_1 \int \frac{1}{y_1^2} dx \\&= \frac{1}{\sqrt{x}} \int \frac{1}{x} dx \\&= \frac{1}{\sqrt{x}} \left(\frac{x^2}{2} \right)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\&= c_1 \left(\frac{1}{\sqrt{x}} \right) + c_2 \left(\frac{1}{\sqrt{x}} \left(\frac{x^2}{2} \right) \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1}{\sqrt{x}} + \frac{c_2 x^{\frac{3}{2}}}{2} \quad (1)$$

Verification of solutions

$$y = \frac{c_1}{\sqrt{x}} + \frac{c_2 x^{\frac{3}{2}}}{2}$$

Verified OK.

9.28.3 Maple step by step solution

Let's solve

$$4x^2 y'' - 3y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = \frac{3y}{4x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - \frac{3y}{4x^2} = 0$$

- Multiply by denominators of the ODE

$$4x^2 y'' - 3y = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x , using the chain rule

$$y' = \left(\frac{d}{dt} y(t) \right) t'(x)$$

- Compute derivative

$$y' = \frac{\frac{d}{dt} y(t)}{x}$$

- Calculate the 2nd derivative of y with respect to x , using the chain rule

$$y'' = \left(\frac{d^2}{dt^2} y(t) \right) t'(x)^2 + t''(x) \left(\frac{d}{dt} y(t) \right)$$

- Compute derivative

$$y'' = \frac{\frac{d^2}{dt^2} y(t)}{x^2} - \frac{\frac{d}{dt} y(t)}{x^2}$$

Substitute the change of variables back into the ODE

$$4x^2 \left(\frac{\frac{d^2}{dt^2} y(t)}{x^2} - \frac{\frac{d}{dt} y(t)}{x^2} \right) - 3y(t) = 0$$

- Simplify

$$4 \frac{d^2}{dt^2} y(t) - 4 \frac{d}{dt} y(t) - 3y(t) = 0$$

- Isolate 2nd derivative

$$\frac{d^2}{dt^2} y(t) = \frac{d}{dt} y(t) + \frac{3y(t)}{4}$$

- Group terms with $y(t)$ on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$\frac{d^2}{dt^2} y(t) - \frac{d}{dt} y(t) - \frac{3y(t)}{4} = 0$$

- Characteristic polynomial of ODE

$$r^2 - r - \frac{3}{4} = 0$$

- Factor the characteristic polynomial

$$\frac{(2r+1)(2r-3)}{4} = 0$$

- Roots of the characteristic polynomial

$$r = \left(-\frac{1}{2}, \frac{3}{2} \right)$$

- 1st solution of the ODE
 $y_1(t) = e^{-\frac{t}{2}}$
- 2nd solution of the ODE
 $y_2(t) = e^{\frac{3t}{2}}$
- General solution of the ODE
 $y(t) = c_1 y_1(t) + c_2 y_2(t)$
- Substitute in solutions
 $y(t) = c_1 e^{-\frac{t}{2}} + c_2 e^{\frac{3t}{2}}$
- Change variables back using $t = \ln(x)$
 $y = \frac{c_1}{\sqrt{x}} + c_2 x^{\frac{3}{2}}$
- Simplify
 $y = \frac{c_1}{\sqrt{x}} + c_2 x^{\frac{3}{2}}$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
<- LODE of Euler type successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(4*x^2*diff(y(x),x$2)-3*y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{c_2 x^2 + c_1}{\sqrt{x}}$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 20

```
DSolve[4*x^2*y'[x]-3*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_2 x^2 + c_1}{\sqrt{x}}$$

9.29 problem 5(e)

9.29.1 Solving as second order euler ode ode	1557
9.29.2 Solving as second order change of variable on x method 2 ode .	1558
9.29.3 Solving as second order change of variable on x method 1 ode .	1561
9.29.4 Solving as second order change of variable on y method 2 ode .	1563
9.29.5 Solving using Kovacic algorithm	1565
9.29.6 Maple step by step solution	1570

Internal problem ID [6297]

Internal file name [OUTPUT/5545_Sunday_June_05_2022_03_43_06_PM_80595365/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.1. Linear Equations with Constant Coefficients. Page 62

Problem number: 5(e).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_euler_ode", "second_order_change_of_variable_on_x_method_1", "second_order_change_of_variable_on_x_method_2", "second_order_change_of_variable_on_y_method_2"

Maple gives the following as the ode type

```
[[_Emden, _Fowler], [_2nd_order, _linear, `_with_symmetry_[0,F(x)]`]]
```

$$x^2y'' - 3xy' + 4y = 0$$

9.29.1 Solving as second order euler ode ode

This is Euler second order ODE. Let the solution be $y = x^r$, then $y' = rx^{r-1}$ and $y'' = r(r-1)x^{r-2}$. Substituting these back into the given ODE gives

$$x^2(r(r-1))x^{r-2} - 3rxr^{r-1} + 4x^r = 0$$

Simplifying gives

$$r(r-1)x^r - 3rx^r + 4x^r = 0$$

Since $x^r \neq 0$ then dividing throughout by x^r gives

$$r(r - 1) - 3r + 4 = 0$$

Or

$$r^2 - 4r + 4 = 0 \quad (1)$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = 2$$

$$r_2 = 2$$

Since the roots are equal, then the general solution is

$$y = c_1 y_1 + c_2 y_2$$

Where $y_1 = x^r$ and $y_2 = x^r \ln(x)$. Hence

$$y = c_1 x^2 + c_2 x^2 \ln(x)$$

Summary

The solution(s) found are the following

$$y = c_1 x^2 + c_2 x^2 \ln(x) \quad (1)$$

Verification of solutions

$$y = c_1 x^2 + c_2 x^2 \ln(x)$$

Verified OK.

9.29.2 Solving as second order change of variable on x method 2 ode

In normal form the ode

$$x^2 y'' - 3xy' + 4y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = -\frac{3}{x}$$
$$q(x) = \frac{4}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x)\tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned} \tau &= \int e^{-(\int p(x)dx)} dx \\ &= \int e^{-(\int -\frac{3}{x}dx)} dx \\ &= \int e^{3\ln(x)} dx \\ &= \int x^3 dx \\ &= \frac{x^4}{4} \end{aligned} \quad (6)$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned} q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\ &= \frac{4}{x^6} \\ &= \frac{4}{x^8} \end{aligned} \quad (7)$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned} \frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) + \frac{4y(\tau)}{x^8} &= 0 \end{aligned}$$

But in terms of τ

$$\frac{4}{x^8} = \frac{1}{4\tau^2}$$

Hence the above ode becomes

$$\frac{d^2}{d\tau^2}y(\tau) + \frac{y(\tau)}{4\tau^2} = 0$$

The above ode is now solved for $y(\tau)$. The ode can be written as

$$4\left(\frac{d^2}{d\tau^2}y(\tau)\right)\tau^2 + y(\tau) = 0$$

Which shows it is a Euler ODE. This is Euler second order ODE. Let the solution be $y(\tau) = \tau^r$, then $y' = r\tau^{r-1}$ and $y'' = r(r-1)\tau^{r-2}$. Substituting these back into the given ODE gives

$$4\tau^2(r(r-1))\tau^{r-2} + 0r\tau^{r-1} + \tau^r = 0$$

Simplifying gives

$$4r(r-1)\tau^r + 0\tau^r + \tau^r = 0$$

Since $\tau^r \neq 0$ then dividing throughout by τ^r gives

$$4r(r-1) + 0 + 1 = 0$$

Or

$$4r^2 - 4r + 1 = 0 \tag{1}$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = \frac{1}{2}$$
$$r_2 = \frac{1}{2}$$

Since the roots are equal, then the general solution is

$$y(\tau) = c_1y_1 + c_2y_2$$

Where $y_1 = \tau^r$ and $y_2 = \tau^r \ln(\tau)$. Hence

$$y(\tau) = c_1\sqrt{\tau} + c_2\sqrt{\tau} \ln(\tau)$$

The above solution is now transformed back to y using (6) which results in

$$y = \frac{(-2c_2 \ln(2) + c_2 \ln(x^4) + c_1) \sqrt{x^4}}{2}$$

Summary

The solution(s) found are the following

$$y = \frac{(-2c_2 \ln(2) + c_2 \ln(x^4) + c_1) \sqrt{x^4}}{2} \quad (1)$$

Verification of solutions

$$y = \frac{(-2c_2 \ln(2) + c_2 \ln(x^4) + c_1) \sqrt{x^4}}{2}$$

Verified OK.

9.29.3 Solving as second order change of variable on x method 1 ode

In normal form the ode

$$x^2 y'' - 3xy' + 4y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = -\frac{3}{x}$$
$$q(x) = \frac{4}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) results

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $q_1 = c^2$ where c is some constant. Therefore from (5)

$$\tau' = \frac{1}{c}\sqrt{q}$$
$$= \frac{2\sqrt{\frac{1}{x^2}}}{c} \quad (6)$$
$$\tau'' = -\frac{2}{c\sqrt{\frac{1}{x^2}}x^3}$$

Substituting the above into (4) results in

$$\begin{aligned}
 p_1(\tau) &= \frac{\tau''(x) + p(x) \tau'(x)}{\tau'(x)^2} \\
 &= \frac{-\frac{2}{c\sqrt{\frac{1}{x^2}}x^3} - \frac{3}{x} \frac{2\sqrt{\frac{1}{x^2}}}{c}}{\left(\frac{2\sqrt{\frac{1}{x^2}}}{c}\right)^2} \\
 &= -2c
 \end{aligned}$$

Therefore ode (3) now becomes

$$\begin{aligned}
 y(\tau)'' + p_1 y(\tau)' + q_1 y(\tau) &= 0 \\
 \frac{d^2}{d\tau^2} y(\tau) - 2c \left(\frac{d}{d\tau} y(\tau) \right) + c^2 y(\tau) &= 0 \tag{7}
 \end{aligned}$$

The above ode is now solved for $y(\tau)$. Since the ode is now constant coefficients, it can be easily solved to give

$$y(\tau) = e^{c\tau} c_1$$

Now from (6)

$$\begin{aligned}
 \tau &= \int \frac{1}{c} \sqrt{q} dx \\
 &= \frac{\int 2\sqrt{\frac{1}{x^2}} dx}{c} \\
 &= \frac{2\sqrt{\frac{1}{x^2}} x \ln(x)}{c}
 \end{aligned}$$

Substituting the above into the solution obtained gives

$$y = c_1 x^2$$

Summary

The solution(s) found are the following

$$y = c_1 x^2 \tag{1}$$

Verification of solutions

$$y = c_1 x^2$$

Verified OK.

9.29.4 Solving as second order change of variable on y method 2 ode

In normal form the ode

$$x^2y'' - 3xy' + 4y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = -\frac{3}{x}$$
$$q(x) = \frac{4}{x^2}$$

Applying change of variables on the dependent variable $y = v(x)x^n$ to (2) gives the following ode where the dependent variables is $v(x)$ and not y .

$$v''(x) + \left(\frac{2n}{x} + p\right)v'(x) + \left(\frac{n(n-1)}{x^2} + \frac{np}{x} + q\right)v(x) = 0 \quad (3)$$

Let the coefficient of $v(x)$ above be zero. Hence

$$\frac{n(n-1)}{x^2} + \frac{np}{x} + q = 0 \quad (4)$$

Substituting the earlier values found for $p(x)$ and $q(x)$ into (4) gives

$$\frac{n(n-1)}{x^2} - \frac{3n}{x^2} + \frac{4}{x^2} = 0 \quad (5)$$

Solving (5) for n gives

$$n = 2 \quad (6)$$

Substituting this value in (3) gives

$$v''(x) + \frac{v'(x)}{x} = 0$$
$$v''(x) + \frac{v'(x)}{x} = 0 \quad (7)$$

Using the substitution

$$u(x) = v'(x)$$

Then (7) becomes

$$u'(x) + \frac{u(x)}{x} = 0 \quad (8)$$

The above is now solved for $u(x)$. In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{u}{x} \end{aligned}$$

Where $f(x) = -\frac{1}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned} \frac{1}{u} du &= -\frac{1}{x} dx \\ \int \frac{1}{u} du &= \int -\frac{1}{x} dx \\ \ln(u) &= -\ln(x) + c_1 \\ u &= e^{-\ln(x)+c_1} \\ &= \frac{c_1}{x} \end{aligned}$$

Now that $u(x)$ is known, then

$$\begin{aligned} v'(x) &= u(x) \\ v(x) &= \int u(x) dx + c_2 \\ &= c_1 \ln(x) + c_2 \end{aligned}$$

Hence

$$\begin{aligned} y &= v(x) x^n \\ &= (c_1 \ln(x) + c_2) x^2 \\ &= (c_1 \ln(x) + c_2) x^2 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = (c_1 \ln(x) + c_2) x^2 \quad (1)$$

Verification of solutions

$$y = (c_1 \ln(x) + c_2) x^2$$

Verified OK.

9.29.5 Solving using Kovacic algorithm

Writing the ode as

$$x^2y'' - 3xy' + 4y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= x^2 \\ B &= -3x \\ C &= 4 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{4x^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 4x^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(-\frac{1}{4x^2}\right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 231: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = -\frac{1}{4x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = -\frac{1}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{1}{2} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{2} \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ , which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = -\frac{1}{4x^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = -\frac{1}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{1}{2} \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{2} \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = -\frac{1}{4x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	$\frac{1}{2}$	$\frac{1}{2}$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	$\frac{1}{2}$	$\frac{1}{2}$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω .

Trying $\alpha_{\infty}^{-} = \frac{1}{2}$ then

$$\begin{aligned} d &= \alpha_{\infty}^{-} - (\alpha_{c_1}^{-}) \\ &= \frac{1}{2} - \left(\frac{1}{2}\right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_{\infty}$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^{-}}{x - c_1} \right) + (-)[\sqrt{r}]_{\infty} \\ &= \frac{1}{2x} + (-)(0) \\ &= \frac{1}{2x} \\ &= \frac{1}{2x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2\left(\frac{1}{2x}\right)(0) + \left(\left(-\frac{1}{2x^2}\right) + \left(\frac{1}{2x}\right)^2 - \left(-\frac{1}{4x^2}\right) \right) &= 0 \\ 0 &= 0 \end{aligned}$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= p e^{\int \omega dx} \\ &= e^{\int \frac{1}{2x} dx} \\ &= \sqrt{x} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\&= z_1 e^{-\int \frac{1}{2} \frac{-3x}{x^2} dx} \\&= z_1 e^{\frac{3 \ln(x)}{2}} \\&= z_1 \left(x^{\frac{3}{2}} \right)\end{aligned}$$

Which simplifies to

$$y_1 = x^2$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{-3x}{x^2} dx}}{(y_1)^2} dx \\&= y_1 \int \frac{e^{3 \ln(x)}}{(y_1)^2} dx \\&= y_1 (\ln(x))\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\&= c_1 (x^2) + c_2 (x^2 (\ln(x)))\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 x^2 + c_2 x^2 \ln(x) \tag{1}$$

Verification of solutions

$$y = c_1 x^2 + c_2 x^2 \ln(x)$$

Verified OK.

9.29.6 Maple step by step solution

Let's solve

$$x^2y'' - 3xy' + 4y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = \frac{3y'}{x} - \frac{4y}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - \frac{3y'}{x} + \frac{4y}{x^2} = 0$$

- Multiply by denominators of the ODE

$$x^2y'' - 3xy' + 4y = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x , using the chain rule

$$y' = \left(\frac{d}{dt}y(t)\right) t'(x)$$

- Compute derivative

$$y' = \frac{\frac{d}{dt}y(t)}{x}$$

- Calculate the 2nd derivative of y with respect to x , using the chain rule

$$y'' = \left(\frac{d^2}{dt^2}y(t)\right) t'(x)^2 + t''(x) \left(\frac{d}{dt}y(t)\right)$$

- Compute derivative

$$y'' = \frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2}$$

Substitute the change of variables back into the ODE

$$x^2 \left(\frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2} \right) - 3 \frac{d}{dt}y(t) + 4y(t) = 0$$

- Simplify

$$\frac{d^2}{dt^2}y(t) - 4 \frac{d}{dt}y(t) + 4y(t) = 0$$

- Characteristic polynomial of ODE

$$r^2 - 4r + 4 = 0$$

- Factor the characteristic polynomial
 $(r - 2)^2 = 0$
- Root of the characteristic polynomial
 $r = 2$
- 1st solution of the ODE
 $y_1(t) = e^{2t}$
- Repeated root, multiply $y_1(t)$ by t to ensure linear independence
 $y_2(t) = t e^{2t}$
- General solution of the ODE
 $y(t) = c_1 y_1(t) + c_2 y_2(t)$
- Substitute in solutions
 $y(t) = c_1 e^{2t} + c_2 t e^{2t}$
- Change variables back using $t = \ln(x)$
 $y = c_1 x^2 + c_2 x^2 \ln(x)$
- Simplify
 $y = x^2(c_1 + c_2 \ln(x))$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
<- LODE of Euler type successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(x^2*diff(y(x),x$2)-3*x*diff(y(x),x)+4*y(x)=0,y(x), singsol=all)
```

$$y(x) = x^2(c_2 \ln(x) + c_1)$$

✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 18

```
DSolve[x^2*y'[x]-3*x*y'[x]+4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^2(2c_2 \log(x) + c_1)$$

9.30 problem 5(f)

9.30.1 Solving as second order euler ode ode	1573
9.30.2 Solving as second order change of variable on x method 2 ode .	1574
9.30.3 Solving as second order change of variable on y method 2 ode .	1577
9.30.4 Solving using Kovacic algorithm	1579
9.30.5 Maple step by step solution	1584

Internal problem ID [6298]

Internal file name [OUTPUT/5546_Sunday_June_05_2022_03_43_07_PM_12118679/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.1. Linear Equations with Constant Coefficients. Page 62

Problem number: 5(f).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_euler_ode", "second_order_change_of_variable_on_x_method_2", "second_order_change_of_variable_on_y_method_2"

Maple gives the following as the ode type

```
[[_Emden, _Fowler], [_2nd_order, _linear, `_with_symmetry_[0,F(x)]`]]
```

$$x^2y'' + 2xy' - 6y = 0$$

9.30.1 Solving as second order euler ode ode

This is Euler second order ODE. Let the solution be $y = x^r$, then $y' = rx^{r-1}$ and $y'' = r(r-1)x^{r-2}$. Substituting these back into the given ODE gives

$$x^2(r(r-1))x^{r-2} + 2rxr^{r-1} - 6x^r = 0$$

Simplifying gives

$$r(r-1)x^r + 2rx^r - 6x^r = 0$$

Since $x^r \neq 0$ then dividing throughout by x^r gives

$$r(r-1) + 2r - 6 = 0$$

Or

$$r^2 + r - 6 = 0 \quad (1)$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = -3$$

$$r_2 = 2$$

Since the roots are real and distinct, then the general solution is

$$y = c_1 y_1 + c_2 y_2$$

Where $y_1 = x^{r_1}$ and $y_2 = x^{r_2}$. Hence

$$y = \frac{c_1}{x^3} + c_2 x^2$$

Summary

The solution(s) found are the following

$$y = \frac{c_1}{x^3} + c_2 x^2 \quad (1)$$

Verification of solutions

$$y = \frac{c_1}{x^3} + c_2 x^2$$

Verified OK.

9.30.2 Solving as second order change of variable on x method 2 ode

In normal form the ode

$$x^2 y'' + 2xy' - 6y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{2}{x}$$
$$q(x) = -\frac{6}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x)\tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned} \tau &= \int e^{-(\int p(x)dx)} dx \\ &= \int e^{-(\int \frac{2}{x} dx)} dx \\ &= \int e^{-2\ln(x)} dx \\ &= \int \frac{1}{x^2} dx \\ &= -\frac{1}{x} \end{aligned} \quad (6)$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned} q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\ &= \frac{-\frac{6}{x^2}}{\frac{1}{x^4}} \\ &= -6x^2 \end{aligned} \quad (7)$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned} \frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) - 6x^2y(\tau) &= 0 \end{aligned}$$

But in terms of τ

$$-6x^2 = -\frac{6}{\tau^2}$$

Hence the above ode becomes

$$\frac{d^2}{d\tau^2}y(\tau) - \frac{6y(\tau)}{\tau^2} = 0$$

The above ode is now solved for $y(\tau)$. The ode can be written as

$$\left(\frac{d^2}{d\tau^2}y(\tau)\right)\tau^2 - 6y(\tau) = 0$$

Which shows it is a Euler ODE. This is Euler second order ODE. Let the solution be $y(\tau) = \tau^r$, then $y' = r\tau^{r-1}$ and $y'' = r(r-1)\tau^{r-2}$. Substituting these back into the given ODE gives

$$\tau^2(r(r-1))\tau^{r-2} + 0r\tau^{r-1} - 6\tau^r = 0$$

Simplifying gives

$$r(r-1)\tau^r + 0\tau^r - 6\tau^r = 0$$

Since $\tau^r \neq 0$ then dividing throughout by τ^r gives

$$r(r-1) + 0 - 6 = 0$$

Or

$$r^2 - r - 6 = 0 \tag{1}$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = -2$$

$$r_2 = 3$$

Since the roots are real and distinct, then the general solution is

$$y(\tau) = c_1y_1 + c_2y_2$$

Where $y_1 = \tau^{r_1}$ and $y_2 = \tau^{r_2}$. Hence

$$y(\tau) = \frac{c_1}{\tau^2} + c_2\tau^3$$

The above solution is now transformed back to y using (6) which results in

$$y = \frac{c_1x^5 - c_2}{x^3}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1 x^5 - c_2}{x^3} \quad (1)$$

Verification of solutions

$$y = \frac{c_1 x^5 - c_2}{x^3}$$

Verified OK.

9.30.3 Solving as second order change of variable on y method 2 ode

In normal form the ode

$$x^2 y'' + 2xy' - 6y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{2}{x}$$
$$q(x) = -\frac{6}{x^2}$$

Applying change of variables on the dependent variable $y = v(x)x^n$ to (2) gives the following ode where the dependent variables is $v(x)$ and not y .

$$v''(x) + \left(\frac{2n}{x} + p\right)v'(x) + \left(\frac{n(n-1)}{x^2} + \frac{np}{x} + q\right)v(x) = 0 \quad (3)$$

Let the coefficient of $v(x)$ above be zero. Hence

$$\frac{n(n-1)}{x^2} + \frac{np}{x} + q = 0 \quad (4)$$

Substituting the earlier values found for $p(x)$ and $q(x)$ into (4) gives

$$\frac{n(n-1)}{x^2} + \frac{2n}{x^2} - \frac{6}{x^2} = 0 \quad (5)$$

Solving (5) for n gives

$$n = 2 \quad (6)$$

Substituting this value in (3) gives

$$\begin{aligned}v''(x) + \frac{6v'(x)}{x} &= 0 \\v''(x) + \frac{6v'(x)}{x} &= 0\end{aligned}\tag{7}$$

Using the substitution

$$u(x) = v'(x)$$

Then (7) becomes

$$u'(x) + \frac{6u(x)}{x} = 0\tag{8}$$

The above is now solved for $u(x)$. In canonical form the ODE is

$$\begin{aligned}u' &= F(x, u) \\&= f(x)g(u) \\&= -\frac{6u}{x}\end{aligned}$$

Where $f(x) = -\frac{6}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u} du &= -\frac{6}{x} dx \\ \int \frac{1}{u} du &= \int -\frac{6}{x} dx \\ \ln(u) &= -6 \ln(x) + c_1 \\ u &= e^{-6 \ln(x) + c_1} \\ &= \frac{c_1}{x^6}\end{aligned}$$

Now that $u(x)$ is known, then

$$\begin{aligned}v'(x) &= u(x) \\ v(x) &= \int u(x) dx + c_2 \\ &= -\frac{c_1}{5x^5} + c_2\end{aligned}$$

Hence

$$\begin{aligned}y &= v(x) x^n \\ &= \left(-\frac{c_1}{5x^5} + c_2\right) x^2 \\ &= \frac{5c_2x^5 - c_1}{5x^3}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \left(-\frac{c_1}{5x^5} + c_2\right) x^2 \quad (1)$$

Verification of solutions

$$y = \left(-\frac{c_1}{5x^5} + c_2\right) x^2$$

Verified OK.

9.30.4 Solving using Kovacic algorithm

Writing the ode as

$$x^2y'' + 2xy' - 6y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= x^2 \\ B &= 2x \\ C &= -6\end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{6}{x^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 6 \\ t &= x^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{6}{x^2}\right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 233: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = \frac{6}{x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = 6$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = 3 \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = -2 \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = \frac{6}{x^2}$$

Since the $\text{gcd}(s, t) = 1$. This gives $b = 6$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = 3 \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = -2 \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{6}{x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	3	-2

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	3	-2

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = -2$ then

$$\begin{aligned} d &= \alpha_\infty^- - (\alpha_{c_1}^-) \\ &= -2 - (-2) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{x - c_1} \right) + (-)[\sqrt{r}]_\infty \\ &= -\frac{2}{x} + (-)(0) \\ &= -\frac{2}{x} \\ &= -\frac{2}{x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$(0) + 2\left(-\frac{2}{x}\right)(0) + \left(\left(\frac{2}{x^2}\right) + \left(-\frac{2}{x}\right)^2 - \left(\frac{6}{x^2}\right)\right) = 0$$
$$0 = 0$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$z_1(x) = pe^{\int \omega dx}$$
$$= e^{\int -\frac{2}{x} dx}$$
$$= \frac{1}{x^2}$$

The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$
$$= z_1 e^{-\int \frac{1}{2} \frac{2x}{x^2} dx}$$
$$= z_1 e^{-\ln(x)}$$
$$= z_1 \left(\frac{1}{x}\right)$$

Which simplifies to

$$y_1 = \frac{1}{x^3}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$y_2 = y_1 \int \frac{e^{\int -\frac{2x}{x^2} dx}}{(y_1)^2} dx$$
$$= y_1 \int \frac{e^{-2\ln(x)}}{(y_1)^2} dx$$
$$= y_1 \left(\frac{x^5}{5}\right)$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(\frac{1}{x^3} \right) + c_2 \left(\frac{1}{x^3} \left(\frac{x^5}{5} \right) \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1}{x^3} + \frac{c_2 x^2}{5} \quad (1)$$

Verification of solutions

$$y = \frac{c_1}{x^3} + \frac{c_2 x^2}{5}$$

Verified OK.

9.30.5 Maple step by step solution

Let's solve

$$x^2 y'' + 2xy' - 6y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{2y'}{x} + \frac{6y}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{2y'}{x} - \frac{6y}{x^2} = 0$$

- Multiply by denominators of the ODE

$$x^2 y'' + 2xy' - 6y = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x , using the chain rule

$$y' = \left(\frac{d}{dt} y(t) \right) t'(x)$$

- Compute derivative

$$y' = \frac{\frac{d}{dt}y(t)}{x}$$

- Calculate the 2nd derivative of y with respect to x , using the chain rule

$$y'' = \left(\frac{d^2}{dt^2}y(t) \right) t'(x)^2 + t''(x) \left(\frac{d}{dt}y(t) \right)$$

- Compute derivative

$$y'' = \frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2}$$

Substitute the change of variables back into the ODE

$$x^2 \left(\frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2} \right) + 2 \frac{d}{dt}y(t) - 6y(t) = 0$$

- Simplify

$$\frac{d^2}{dt^2}y(t) + \frac{d}{dt}y(t) - 6y(t) = 0$$

- Characteristic polynomial of ODE

$$r^2 + r - 6 = 0$$

- Factor the characteristic polynomial

$$(r + 3)(r - 2) = 0$$

- Roots of the characteristic polynomial

$$r = (-3, 2)$$

- 1st solution of the ODE

$$y_1(t) = e^{-3t}$$

- 2nd solution of the ODE

$$y_2(t) = e^{2t}$$

- General solution of the ODE

$$y(t) = c_1 y_1(t) + c_2 y_2(t)$$

- Substitute in solutions

$$y(t) = c_1 e^{-3t} + c_2 e^{2t}$$

- Change variables back using $t = \ln(x)$

$$y = \frac{c_1}{x^3} + c_2 x^2$$

- Simplify

$$y = \frac{c_1}{x^3} + c_2 x^2$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
<- LODE of Euler type successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(x^2*diff(y(x),x$2)+2*x*diff(y(x),x)-6*y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{c_1 x^5 + c_2}{x^3}$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 18

```
DSolve[x^2*y'[x]+2*x*y'[x]-6*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_2 x^5 + c_1}{x^3}$$

9.31 problem 5(g)

9.31.1 Solving as second order euler ode ode	1587
9.31.2 Solving as second order change of variable on x method 2 ode .	1589
9.31.3 Solving as second order change of variable on x method 1 ode .	1592
9.31.4 Solving as second order change of variable on y method 2 ode .	1594
9.31.5 Solving using Kovacic algorithm	1597
9.31.6 Maple step by step solution	1602

Internal problem ID [6299]

Internal file name [OUTPUT/5547_Sunday_June_05_2022_03_43_08_PM_87551141/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.1. Linear Equations with Constant Coefficients. Page 62

Problem number: 5(g).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_euler_ode", "second_order_change_of_variable_on_x_method_1", "second_order_change_of_variable_on_x_method_2", "second_order_change_of_variable_on_y_method_2"

Maple gives the following as the ode type

```
[[_Emden, _Fowler]]
```

$$x^2y'' + 2xy' + 3y = 0$$

9.31.1 Solving as second order euler ode ode

This is Euler second order ODE. Let the solution be $y = x^r$, then $y' = rx^{r-1}$ and $y'' = r(r-1)x^{r-2}$. Substituting these back into the given ODE gives

$$x^2(r(r-1))x^{r-2} + 2rxr^{r-1} + 3x^r = 0$$

Simplifying gives

$$r(r-1)x^r + 2rx^r + 3x^r = 0$$

Since $x^r \neq 0$ then dividing throughout by x^r gives

$$r(r - 1) + 2r + 3 = 0$$

Or

$$r^2 + r + 3 = 0 \quad (1)$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = -\frac{1}{2} - \frac{i\sqrt{11}}{2}$$

$$r_2 = -\frac{1}{2} + \frac{i\sqrt{11}}{2}$$

The roots are complex conjugate of each others. Let the roots be

$$r_1 = \alpha + i\beta$$

$$r_2 = \alpha - i\beta$$

Where in this case $\alpha = -\frac{1}{2}$ and $\beta = -\frac{\sqrt{11}}{2}$. Hence the solution becomes

$$y = c_1 x^{r_1} + c_2 x^{r_2}$$

$$= c_1 x^{\alpha+i\beta} + c_2 x^{\alpha-i\beta}$$

$$= x^\alpha (c_1 x^{i\beta} + c_2 x^{-i\beta})$$

$$= x^\alpha (c_1 e^{\ln(x^{i\beta})} + c_2 e^{\ln(x^{-i\beta})})$$

$$= x^\alpha (c_1 e^{i(\beta \ln x)} + c_2 e^{-i(\beta \ln x)})$$

Using the values for $\alpha = -\frac{1}{2}$, $\beta = -\frac{\sqrt{11}}{2}$, the above becomes

$$y = x^{-\frac{1}{2}} \left(c_1 e^{-\frac{i\sqrt{11} \ln(x)}{2}} + c_2 e^{\frac{i\sqrt{11} \ln(x)}{2}} \right)$$

Using Euler relation, the expression $c_1 e^{iA} + c_2 e^{-iA}$ is transformed to $c_1 \cos A + c_1 \sin A$ where the constants are free to change. Applying this to the above result gives

$$y = \frac{1}{\sqrt{x}} \left(c_1 \cos \left(\frac{\sqrt{11} \ln(x)}{2} \right) + c_2 \sin \left(\frac{\sqrt{11} \ln(x)}{2} \right) \right)$$

Summary

The solution(s) found are the following

$$y = \frac{c_1 \cos \left(\frac{\sqrt{11} \ln(x)}{2} \right) + c_2 \sin \left(\frac{\sqrt{11} \ln(x)}{2} \right)}{\sqrt{x}} \quad (1)$$

Verification of solutions

$$y = \frac{c_1 \cos\left(\frac{\sqrt{11} \ln(x)}{2}\right) + c_2 \sin\left(\frac{\sqrt{11} \ln(x)}{2}\right)}{\sqrt{x}}$$

Verified OK.

9.31.2 Solving as second order change of variable on x method 2 ode

In normal form the ode

$$x^2 y'' + 2xy' + 3y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{2}{x}$$
$$q(x) = \frac{3}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x)\tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned}
 \tau &= \int e^{-(\int p(x)dx)} dx \\
 &= \int e^{-(\int \frac{2}{x} dx)} dx \\
 &= \int e^{-2\ln(x)} dx \\
 &= \int \frac{1}{x^2} dx \\
 &= -\frac{1}{x}
 \end{aligned} \tag{6}$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned}
 q_1(\tau) &= \frac{q(x)}{\tau' (x)^2} \\
 &= \frac{\frac{3}{x^2}}{\frac{1}{x^4}} \\
 &= 3x^2
 \end{aligned} \tag{7}$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned}
 \frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) &= 0 \\
 \frac{d^2}{d\tau^2}y(\tau) + 3x^2y(\tau) &= 0
 \end{aligned}$$

But in terms of τ

$$3x^2 = \frac{3}{\tau^2}$$

Hence the above ode becomes

$$\frac{d^2}{d\tau^2}y(\tau) + \frac{3y(\tau)}{\tau^2} = 0$$

The above ode is now solved for $y(\tau)$. The ode can be written as

$$\left(\frac{d^2}{d\tau^2}y(\tau)\right)\tau^2 + 3y(\tau) = 0$$

Which shows it is a Euler ODE. This is Euler second order ODE. Let the solution be $y(\tau) = \tau^r$, then $y' = r\tau^{r-1}$ and $y'' = r(r-1)\tau^{r-2}$. Substituting these back into the given ODE gives

$$\tau^2(r(r-1))\tau^{r-2} + 0r\tau^{r-1} + 3\tau^r = 0$$

Simplifying gives

$$r(r-1)\tau^r + 0\tau^r + 3\tau^r = 0$$

Since $\tau^r \neq 0$ then dividing throughout by τ^r gives

$$r(r-1) + 0 + 3 = 0$$

Or

$$r^2 - r + 3 = 0 \quad (1)$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = \frac{1}{2} - \frac{i\sqrt{11}}{2}$$
$$r_2 = \frac{1}{2} + \frac{i\sqrt{11}}{2}$$

The roots are complex conjugate of each others. Let the roots be

$$r_1 = \alpha + i\beta$$
$$r_2 = \alpha - i\beta$$

Where in this case $\alpha = \frac{1}{2}$ and $\beta = -\frac{\sqrt{11}}{2}$. Hence the solution becomes

$$y(\tau) = c_1\tau^{r_1} + c_2\tau^{r_2}$$
$$= c_1\tau^{\alpha+i\beta} + c_2\tau^{\alpha-i\beta}$$
$$= \tau^\alpha(c_1\tau^{i\beta} + c_2\tau^{-i\beta})$$
$$= \tau^\alpha(c_1e^{\ln(\tau^{i\beta})} + c_2e^{\ln(\tau^{-i\beta})})$$
$$= \tau^\alpha(c_1e^{i(\beta \ln \tau)} + c_2e^{-i(\beta \ln \tau)})$$

Using the values for $\alpha = \frac{1}{2}$, $\beta = -\frac{\sqrt{11}}{2}$, the above becomes

$$y(\tau) = \tau^{\frac{1}{2}} \left(c_1 e^{-\frac{i\sqrt{11} \ln(\tau)}{2}} + c_2 e^{\frac{i\sqrt{11} \ln(\tau)}{2}} \right)$$

Using Euler relation, the expression $c_1e^{iA} + c_2e^{-iA}$ is transformed to $c_1 \cos A + c_1 \sin A$ where the constants are free to change. Applying this to the above result gives

$$y(\tau) = \sqrt{\tau} \left(c_1 \cos \left(\frac{\sqrt{11} \ln(\tau)}{2} \right) + c_2 \sin \left(\frac{\sqrt{11} \ln(\tau)}{2} \right) \right)$$

The above solution is now transformed back to y using (6) which results in

$$y = \sqrt{-\frac{1}{x}} \left(c_1 \cos \left(\frac{\sqrt{11} \ln \left(-\frac{1}{x} \right)}{2} \right) + c_2 \sin \left(\frac{\sqrt{11} \ln \left(-\frac{1}{x} \right)}{2} \right) \right)$$

Summary

The solution(s) found are the following

$$y = \sqrt{-\frac{1}{x}} \left(c_1 \cos \left(\frac{\sqrt{11} \ln \left(-\frac{1}{x} \right)}{2} \right) + c_2 \sin \left(\frac{\sqrt{11} \ln \left(-\frac{1}{x} \right)}{2} \right) \right) \quad (1)$$

Verification of solutions

$$y = \sqrt{-\frac{1}{x}} \left(c_1 \cos \left(\frac{\sqrt{11} \ln \left(-\frac{1}{x} \right)}{2} \right) + c_2 \sin \left(\frac{\sqrt{11} \ln \left(-\frac{1}{x} \right)}{2} \right) \right)$$

Verified OK.

9.31.3 Solving as second order change of variable on x method 1 ode

In normal form the ode

$$x^2 y'' + 2xy' + 3y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{2}{x}$$

$$q(x) = \frac{3}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) results

$$\frac{d^2}{d\tau^2} y(\tau) + p_1 \left(\frac{d}{d\tau} y(\tau) \right) + q_1 y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $q_1 = c^2$ where c is some constant. Therefore from (5)

$$\begin{aligned}\tau' &= \frac{1}{c}\sqrt{q} \\ &= \frac{\sqrt{3}\sqrt{\frac{1}{x^2}}}{c} \\ \tau'' &= -\frac{\sqrt{3}}{c\sqrt{\frac{1}{x^2}}x^3}\end{aligned}\tag{6}$$

Substituting the above into (4) results in

$$\begin{aligned}p_1(\tau) &= \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \\ &= \frac{-\frac{\sqrt{3}}{c\sqrt{\frac{1}{x^2}}x^3} + \frac{2}{x}\frac{\sqrt{3}\sqrt{\frac{1}{x^2}}}{c}}{\left(\frac{\sqrt{3}\sqrt{\frac{1}{x^2}}}{c}\right)^2} \\ &= \frac{c\sqrt{3}}{3}\end{aligned}$$

Therefore ode (3) now becomes

$$\begin{aligned}y(\tau)'' + p_1y(\tau)' + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) + \frac{c\sqrt{3}}{3}\left(\frac{d}{d\tau}y(\tau)\right) + c^2y(\tau) &= 0\end{aligned}\tag{7}$$

The above ode is now solved for $y(\tau)$. Since the ode is now constant coefficients, it can be easily solved to give

$$y(\tau) = e^{-\frac{\sqrt{3}c\tau}{6}} \left(c_1 \cos\left(\frac{c\sqrt{33}\tau}{6}\right) + c_2 \sin\left(\frac{c\sqrt{33}\tau}{6}\right) \right)$$

Now from (6)

$$\begin{aligned}\tau &= \int \frac{1}{c}\sqrt{q} dx \\ &= \frac{\int \sqrt{3}\sqrt{\frac{1}{x^2}} dx}{c} \\ &= \frac{\sqrt{3}\sqrt{\frac{1}{x^2}}x \ln(x)}{c}\end{aligned}$$

Substituting the above into the solution obtained gives

$$y = \frac{c_1 \cos\left(\frac{\sqrt{11} \ln(x)}{2}\right) + c_2 \sin\left(\frac{\sqrt{11} \ln(x)}{2}\right)}{\sqrt{x}}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1 \cos\left(\frac{\sqrt{11} \ln(x)}{2}\right) + c_2 \sin\left(\frac{\sqrt{11} \ln(x)}{2}\right)}{\sqrt{x}} \quad (1)$$

Verification of solutions

$$y = \frac{c_1 \cos\left(\frac{\sqrt{11} \ln(x)}{2}\right) + c_2 \sin\left(\frac{\sqrt{11} \ln(x)}{2}\right)}{\sqrt{x}}$$

Verified OK.

9.31.4 Solving as second order change of variable on y method 2 ode

In normal form the ode

$$x^2 y'' + 2xy' + 3y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{2}{x}$$

$$q(x) = \frac{3}{x^2}$$

Applying change of variables on the dependent variable $y = v(x)x^n$ to (2) gives the following ode where the dependent variables is $v(x)$ and not y .

$$v''(x) + \left(\frac{2n}{x} + p\right)v'(x) + \left(\frac{n(n-1)}{x^2} + \frac{np}{x} + q\right)v(x) = 0 \quad (3)$$

Let the coefficient of $v(x)$ above be zero. Hence

$$\frac{n(n-1)}{x^2} + \frac{np}{x} + q = 0 \quad (4)$$

Substituting the earlier values found for $p(x)$ and $q(x)$ into (4) gives

$$\frac{n(n-1)}{x^2} + \frac{2n}{x^2} + \frac{3}{x^2} = 0 \quad (5)$$

Solving (5) for n gives

$$n = -\frac{1}{2} + \frac{i\sqrt{11}}{2} \quad (6)$$

Substituting this value in (3) gives

$$\begin{aligned} v''(x) + \left(\frac{-1 + i\sqrt{11}}{x} + \frac{2}{x} \right) v'(x) &= 0 \\ v''(x) + \frac{(i\sqrt{11} + 1)v'(x)}{x} &= 0 \end{aligned} \quad (7)$$

Using the substitution

$$u(x) = v'(x)$$

Then (7) becomes

$$u'(x) + \frac{(i\sqrt{11} + 1)u(x)}{x} = 0 \quad (8)$$

The above is now solved for $u(x)$. In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= \frac{(-1 - i\sqrt{11})u}{x} \end{aligned}$$

Where $f(x) = \frac{-1 - i\sqrt{11}}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned} \frac{1}{u} du &= \frac{-1 - i\sqrt{11}}{x} dx \\ \int \frac{1}{u} du &= \int \frac{-1 - i\sqrt{11}}{x} dx \\ \ln(u) &= (-1 - i\sqrt{11}) \ln(x) + c_1 \\ u &= e^{(-1 - i\sqrt{11}) \ln(x) + c_1} \\ &= c_1 e^{(-1 - i\sqrt{11}) \ln(x)} \end{aligned}$$

Which simplifies to

$$u(x) = \frac{c_1 x^{-i\sqrt{11}}}{x}$$

Now that $u(x)$ is known, then

$$\begin{aligned}v'(x) &= u(x) \\v(x) &= \int u(x) dx + c_2 \\&= \frac{i\sqrt{11} c_1 x^{-i\sqrt{11}}}{11} + c_2\end{aligned}$$

Hence

$$\begin{aligned}y &= v(x) x^n \\&= \left(\frac{i\sqrt{11} c_1 x^{-i\sqrt{11}}}{11} + c_2 \right) x^{-\frac{1}{2} + \frac{i\sqrt{11}}{2}} \\&= \frac{x^{-\frac{1}{2} - \frac{i\sqrt{11}}{2}} \left(i\sqrt{11} c_1 + 11c_2 x^{i\sqrt{11}} \right)}{11}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \left(\frac{i\sqrt{11} c_1 x^{-i\sqrt{11}}}{11} + c_2 \right) x^{-\frac{1}{2} + \frac{i\sqrt{11}}{2}} \quad (1)$$

Verification of solutions

$$y = \left(\frac{i\sqrt{11} c_1 x^{-i\sqrt{11}}}{11} + c_2 \right) x^{-\frac{1}{2} + \frac{i\sqrt{11}}{2}}$$

Verified OK.

9.31.5 Solving using Kovacic algorithm

Writing the ode as

$$x^2y'' + 2xy' + 3y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= x^2 \\ B &= 2x \\ C &= 3 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-3}{x^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -3 \\ t &= x^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(-\frac{3}{x^2}\right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 235: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = -\frac{3}{x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = -3$. Hence

$$[\sqrt{r}]_c = 0$$

$$\alpha_c^+ = \frac{1}{2} + \sqrt{1 + 4b} = \frac{1}{2} + \frac{i\sqrt{11}}{2}$$

$$\alpha_c^- = \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{2} - \frac{i\sqrt{11}}{2}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = -\frac{3}{x^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = -3$. Hence

$$[\sqrt{r}]_\infty = 0$$

$$\alpha_\infty^+ = \frac{1}{2} + \sqrt{1 + 4b} = \frac{1}{2} + \frac{i\sqrt{11}}{2}$$

$$\alpha_\infty^- = \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{2} - \frac{i\sqrt{11}}{2}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = -\frac{3}{x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	$\frac{1}{2} + \frac{i\sqrt{11}}{2}$	$\frac{1}{2} - \frac{i\sqrt{11}}{2}$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	$\frac{1}{2} + \frac{i\sqrt{11}}{2}$	$\frac{1}{2} - \frac{i\sqrt{11}}{2}$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω .

Trying $\alpha_{\infty}^{-} = \frac{1}{2} - \frac{i\sqrt{11}}{2}$ then

$$\begin{aligned} d &= \alpha_{\infty}^{-} - (\alpha_{c_1}^{-}) \\ &= \frac{1}{2} - \frac{i\sqrt{11}}{2} - \left(\frac{1}{2} - \frac{i\sqrt{11}}{2} \right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_{\infty}$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^{-}}{x - c_1} \right) + (-)[\sqrt{r}]_{\infty} \\ &= \frac{\frac{1}{2} - \frac{i\sqrt{11}}{2}}{x} + (-)(0) \\ &= \frac{\frac{1}{2} - \frac{i\sqrt{11}}{2}}{x} \\ &= \frac{1 - i\sqrt{11}}{2x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r)p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2 \left(\frac{\frac{1}{2} - \frac{i\sqrt{11}}{2}}{x} \right) (0) + \left(\left(-\frac{\frac{1}{2} - \frac{i\sqrt{11}}{2}}{x^2} \right) + \left(\frac{\frac{1}{2} - \frac{i\sqrt{11}}{2}}{x} \right)^2 - \left(-\frac{3}{x^2} \right) \right) &= 0 \\ 0 &= 0 \end{aligned}$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= p e^{\int \omega dx} \\ &= e^{\int \frac{\frac{1}{2} - \frac{i\sqrt{11}}{2}}{x} dx} \\ &= x^{\frac{1}{2} - \frac{i\sqrt{11}}{2}} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\&= z_1 e^{-\int \frac{1}{2} \frac{2x}{x^2} dx} \\&= z_1 e^{-\ln(x)} \\&= z_1 \left(\frac{1}{x} \right)\end{aligned}$$

Which simplifies to

$$y_1 = x^{-\frac{1}{2} - \frac{i\sqrt{11}}{2}}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{2x}{x^2} dx}}{(y_1)^2} dx \\&= y_1 \int \frac{e^{-2\ln(x)}}{(y_1)^2} dx \\&= y_1 \left(-\frac{ix^{i\sqrt{11}}\sqrt{11}}{11} \right)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\&= c_1 \left(x^{-\frac{1}{2} - \frac{i\sqrt{11}}{2}} \right) + c_2 \left(x^{-\frac{1}{2} - \frac{i\sqrt{11}}{2}} \left(-\frac{ix^{i\sqrt{11}}\sqrt{11}}{11} \right) \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 x^{-\frac{1}{2} - \frac{i\sqrt{11}}{2}} - \frac{ic_2 \sqrt{11} x^{-\frac{1}{2} + \frac{i\sqrt{11}}{2}}}{11} \quad (1)$$

Verification of solutions

$$y = c_1 x^{-\frac{1}{2} - \frac{i\sqrt{11}}{2}} - \frac{ic_2 \sqrt{11} x^{-\frac{1}{2} + \frac{i\sqrt{11}}{2}}}{11}$$

Verified OK.

9.31.6 Maple step by step solution

Let's solve

$$x^2 y'' + 2xy' + 3y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{2y'}{x} - \frac{3y}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{2y'}{x} + \frac{3y}{x^2} = 0$$

- Multiply by denominators of the ODE

$$x^2 y'' + 2xy' + 3y = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x , using the chain rule

$$y' = \left(\frac{d}{dt}y(t)\right) t'(x)$$

- Compute derivative

$$y' = \frac{\frac{d}{dt}y(t)}{x}$$

- Calculate the 2nd derivative of y with respect to x , using the chain rule

$$y'' = \left(\frac{d^2}{dt^2}y(t)\right) t'(x)^2 + t''(x) \left(\frac{d}{dt}y(t)\right)$$

- Compute derivative

$$y'' = \frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2}$$

Substitute the change of variables back into the ODE

$$x^2 \left(\frac{d^2 y(t)}{dt^2} - \frac{d}{dt} \frac{y(t)}{x^2} \right) + 2 \frac{d}{dt} y(t) + 3y(t) = 0$$

- Simplify

$$\frac{d^2}{dt^2} y(t) + \frac{d}{dt} y(t) + 3y(t) = 0$$

- Characteristic polynomial of ODE

$$r^2 + r + 3 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-1) \pm (\sqrt{-11})}{2}$$

- Roots of the characteristic polynomial

$$r = \left(-\frac{1}{2} - \frac{i\sqrt{11}}{2}, -\frac{1}{2} + \frac{i\sqrt{11}}{2} \right)$$

- 1st solution of the ODE

$$y_1(t) = e^{-\frac{t}{2}} \cos\left(\frac{\sqrt{11}t}{2}\right)$$

- 2nd solution of the ODE

$$y_2(t) = e^{-\frac{t}{2}} \sin\left(\frac{\sqrt{11}t}{2}\right)$$

- General solution of the ODE

$$y(t) = c_1 y_1(t) + c_2 y_2(t)$$

- Substitute in solutions

$$y(t) = c_1 e^{-\frac{t}{2}} \cos\left(\frac{\sqrt{11}t}{2}\right) + c_2 e^{-\frac{t}{2}} \sin\left(\frac{\sqrt{11}t}{2}\right)$$

- Change variables back using $t = \ln(x)$

$$y = \frac{c_1 \cos\left(\frac{\sqrt{11} \ln(x)}{2}\right)}{\sqrt{x}} + \frac{c_2 \sin\left(\frac{\sqrt{11} \ln(x)}{2}\right)}{\sqrt{x}}$$

- Simplify

$$y = \frac{c_1 \cos\left(\frac{\sqrt{11} \ln(x)}{2}\right)}{\sqrt{x}} + \frac{c_2 \sin\left(\frac{\sqrt{11} \ln(x)}{2}\right)}{\sqrt{x}}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
<- LODE of Euler type successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 29

```
dsolve(x^2*diff(y(x),x$2)+2*x*diff(y(x),x)+3*y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{c_1 \sin\left(\frac{\sqrt{11} \ln(x)}{2}\right) + c_2 \cos\left(\frac{\sqrt{11} \ln(x)}{2}\right)}{\sqrt{x}}$$

✓ Solution by Mathematica

Time used: 0.031 (sec). Leaf size: 42

```
DSolve[x^2*y''[x]+2*x*y'[x]+3*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_2 \cos\left(\frac{1}{2}\sqrt{11} \log(x)\right) + c_1 \sin\left(\frac{1}{2}\sqrt{11} \log(x)\right)}{\sqrt{x}}$$

9.32 problem 5(h)

9.32.1 Solving as second order euler ode ode	1605
9.32.2 Solving as second order change of variable on x method 2 ode .	1606
9.32.3 Solving as second order change of variable on x method 1 ode .	1609
9.32.4 Solving as second order change of variable on y method 2 ode .	1611
9.32.5 Solving using Kovacic algorithm	1613
9.32.6 Maple step by step solution	1619

Internal problem ID [6300]

Internal file name [OUTPUT/5548_Sunday_June_05_2022_03_43_09_PM_86161255/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.1. Linear Equations with Constant Coefficients. Page 62

Problem number: 5(h).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_euler_ode", "second_order_change_of_variable_on_x_method_1", "second_order_change_of_variable_on_x_method_2", "second_order_change_of_variable_on_y_method_2"

Maple gives the following as the ode type

```
[[_Emden, _Fowler], [_2nd_order, _linear, `_with_symmetry_[0,F(x)]`]]
```

$$x^2y'' + xy' - 2y = 0$$

9.32.1 Solving as second order euler ode ode

This is Euler second order ODE. Let the solution be $y = x^r$, then $y' = rx^{r-1}$ and $y'' = r(r-1)x^{r-2}$. Substituting these back into the given ODE gives

$$x^2(r(r-1))x^{r-2} + rx^{r-1} - 2x^r = 0$$

Simplifying gives

$$r(r-1)x^r + rx^r - 2x^r = 0$$

Since $x^r \neq 0$ then dividing throughout by x^r gives

$$r(r - 1) + r - 2 = 0$$

Or

$$r^2 - 2 = 0 \quad (1)$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = \sqrt{2}$$
$$r_2 = -\sqrt{2}$$

Since the roots are real and distinct, then the general solution is

$$y = c_1 y_1 + c_2 y_2$$

Where $y_1 = x^{r_1}$ and $y_2 = x^{r_2}$. Hence

$$y = c_1 x^{\sqrt{2}} + c_2 x^{-\sqrt{2}}$$

Summary

The solution(s) found are the following

$$y = c_1 x^{\sqrt{2}} + c_2 x^{-\sqrt{2}} \quad (1)$$

Verification of solutions

$$y = c_1 x^{\sqrt{2}} + c_2 x^{-\sqrt{2}}$$

Verified OK.

9.32.2 Solving as second order change of variable on x method 2 ode

In normal form the ode

$$x^2 y'' + xy' - 2y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = -\frac{2}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x)\tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned} \tau &= \int e^{-(\int p(x)dx)} dx \\ &= \int e^{-(\int \frac{1}{x}dx)} dx \\ &= \int e^{-\ln(x)} dx \\ &= \int \frac{1}{x} dx \\ &= \ln(x) \end{aligned} \quad (6)$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned} q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\ &= \frac{-\frac{2}{x^2}}{\frac{1}{x^2}} \\ &= -2 \end{aligned} \quad (7)$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned} \frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) - 2y(\tau) &= 0 \end{aligned}$$

The above ode is now solved for $y(\tau)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(\tau) + By'(\tau) + Cy(\tau) = 0$$

Where in the above $A = 1, B = 0, C = -2$. Let the solution be $y(\tau) = e^{\lambda\tau}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda\tau} - 2e^{\lambda\tau} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda\tau}$ gives

$$\lambda^2 - 2 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = -2$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(-2)} \\ &= \pm\sqrt{2} \end{aligned}$$

Hence

$$\lambda_1 = +\sqrt{2}$$

$$\lambda_2 = -\sqrt{2}$$

Which simplifies to

$$\lambda_1 = \sqrt{2}$$

$$\lambda_2 = -\sqrt{2}$$

Since roots are real and distinct, then the solution is

$$y(\tau) = c_1 e^{\lambda_1 \tau} + c_2 e^{\lambda_2 \tau}$$

$$y(\tau) = c_1 e^{(\sqrt{2})\tau} + c_2 e^{(-\sqrt{2})\tau}$$

Or

$$y(\tau) = c_1 e^{\sqrt{2}\tau} + c_2 e^{-\sqrt{2}\tau}$$

The above solution is now transformed back to y using (6) which results in

$$y = c_1 x^{\sqrt{2}} + c_2 x^{-\sqrt{2}}$$

Summary

The solution(s) found are the following

$$y = c_1 x^{\sqrt{2}} + c_2 x^{-\sqrt{2}} \quad (1)$$

Verification of solutions

$$y = c_1 x^{\sqrt{2}} + c_2 x^{-\sqrt{2}}$$

Verified OK.

9.32.3 Solving as second order change of variable on x method 1 ode

In normal form the ode

$$x^2 y'' + xy' - 2y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = -\frac{2}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) results

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $q_1 = c^2$ where c is some constant. Therefore from (5)

$$\begin{aligned}\tau' &= \frac{1}{c}\sqrt{q} \\ &= \frac{\sqrt{-\frac{2}{x^2}}}{c} \\ \tau'' &= \frac{2}{c\sqrt{-\frac{2}{x^2}}x^3}\end{aligned}\tag{6}$$

Substituting the above into (4) results in

$$\begin{aligned}p_1(\tau) &= \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \\ &= \frac{\frac{2}{c\sqrt{-\frac{2}{x^2}}x^3} + \frac{1}{x}\frac{\sqrt{-\frac{2}{x^2}}}{c}}{\left(\frac{\sqrt{-\frac{2}{x^2}}}{c}\right)^2} \\ &= 0\end{aligned}$$

Therefore ode (3) now becomes

$$\begin{aligned}y(\tau)'' + p_1y(\tau)' + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) + c^2y(\tau) &= 0\end{aligned}\tag{7}$$

The above ode is now solved for $y(\tau)$. Since the ode is now constant coefficients, it can be easily solved to give

$$y(\tau) = c_1 \cos(c\tau) + c_2 \sin(c\tau)$$

Now from (6)

$$\begin{aligned}\tau &= \int \frac{1}{c}\sqrt{q} dx \\ &= \frac{\int \sqrt{-\frac{2}{x^2}} dx}{c} \\ &= \frac{\sqrt{-\frac{2}{x^2}} x \ln(x)}{c}\end{aligned}$$

Substituting the above into the solution obtained gives

$$y = c_1 \cosh\left(\sqrt{2} \ln(x)\right) + ic_2 \sinh\left(\sqrt{2} \ln(x)\right)$$

Summary

The solution(s) found are the following

$$y = c_1 \cosh \left(\sqrt{2} \ln (x) \right) + ic_2 \sinh \left(\sqrt{2} \ln (x) \right) \quad (1)$$

Verification of solutions

$$y = c_1 \cosh \left(\sqrt{2} \ln (x) \right) + ic_2 \sinh \left(\sqrt{2} \ln (x) \right)$$

Verified OK.

9.32.4 Solving as second order change of variable on y method 2 ode

In normal form the ode

$$x^2 y'' + xy' - 2y = 0 \quad (1)$$

Becomes

$$y'' + p(x) y' + q(x) y = 0 \quad (2)$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = -\frac{2}{x^2}$$

Applying change of variables on the dependent variable $y = v(x) x^n$ to (2) gives the following ode where the dependent variables is $v(x)$ and not y .

$$v''(x) + \left(\frac{2n}{x} + p \right) v'(x) + \left(\frac{n(n-1)}{x^2} + \frac{np}{x} + q \right) v(x) = 0 \quad (3)$$

Let the coefficient of $v(x)$ above be zero. Hence

$$\frac{n(n-1)}{x^2} + \frac{np}{x} + q = 0 \quad (4)$$

Substituting the earlier values found for $p(x)$ and $q(x)$ into (4) gives

$$\frac{n(n-1)}{x^2} + \frac{n}{x^2} - \frac{2}{x^2} = 0 \quad (5)$$

Solving (5) for n gives

$$n = \sqrt{2} \quad (6)$$

Substituting this value in (3) gives

$$\begin{aligned}v''(x) + \left(\frac{2\sqrt{2}}{x} + \frac{1}{x}\right)v'(x) &= 0 \\v''(x) + \frac{(1 + 2\sqrt{2})v'(x)}{x} &= 0\end{aligned}\tag{7}$$

Using the substitution

$$u(x) = v'(x)$$

Then (7) becomes

$$u'(x) + \frac{(1 + 2\sqrt{2})u(x)}{x} = 0\tag{8}$$

The above is now solved for $u(x)$. In canonical form the ODE is

$$\begin{aligned}u' &= F(x, u) \\&= f(x)g(u) \\&= \frac{(-1 - 2\sqrt{2})u}{x}\end{aligned}$$

Where $f(x) = \frac{-1-2\sqrt{2}}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u} du &= \frac{-1 - 2\sqrt{2}}{x} dx \\ \int \frac{1}{u} du &= \int \frac{-1 - 2\sqrt{2}}{x} dx \\ \ln(u) &= (-1 - 2\sqrt{2}) \ln(x) + c_1 \\ u &= e^{(-1-2\sqrt{2}) \ln(x) + c_1} \\ &= c_1 e^{(-1-2\sqrt{2}) \ln(x)}\end{aligned}$$

Which simplifies to

$$u(x) = \frac{c_1 x^{-2\sqrt{2}}}{x}$$

Now that $u(x)$ is known, then

$$\begin{aligned}v'(x) &= u(x) \\v(x) &= \int u(x) dx + c_2 \\&= -\frac{\sqrt{2} c_1 x^{-2\sqrt{2}}}{4} + c_2\end{aligned}$$

Hence

$$\begin{aligned}y &= v(x) x^n \\&= \left(-\frac{\sqrt{2} c_1 x^{-2\sqrt{2}}}{4} + c_2 \right) x^{\sqrt{2}} \\&= x^{\sqrt{2}} c_2 - \frac{x^{-\sqrt{2}} c_1 \sqrt{2}}{4}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \left(-\frac{\sqrt{2} c_1 x^{-2\sqrt{2}}}{4} + c_2 \right) x^{\sqrt{2}} \quad (1)$$

Verification of solutions

$$y = \left(-\frac{\sqrt{2} c_1 x^{-2\sqrt{2}}}{4} + c_2 \right) x^{\sqrt{2}}$$

Verified OK.

9.32.5 Solving using Kovacic algorithm

Writing the ode as

$$x^2 y'' + xy' - 2y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= x^2 \\B &= x \\C &= -2\end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{7}{4x^2} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 7 \\ t &= 4x^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{7}{4x^2} \right) z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 237: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 2 - 0 \\
 &= 2
 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = \frac{7}{4x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition

of r given above. Therefore $b = \frac{7}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{1}{2} + \sqrt{2} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{2} - \sqrt{2} \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ , which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = \frac{7}{4x^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = \frac{7}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{1}{2} + \sqrt{2} \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{2} - \sqrt{2} \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{7}{4x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	$\frac{1}{2} + \sqrt{2}$	$\frac{1}{2} - \sqrt{2}$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	$\frac{1}{2} + \sqrt{2}$	$\frac{1}{2} - \sqrt{2}$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω .

Trying $\alpha_{\infty}^{-} = \frac{1}{2} - \sqrt{2}$ then

$$\begin{aligned} d &= \alpha_{\infty}^{-} - (\alpha_{c_1}^{-}) \\ &= \frac{1}{2} - \sqrt{2} - \left(\frac{1}{2} - \sqrt{2}\right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_{\infty}$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^{-}}{x - c_1} \right) + (-)[\sqrt{r}]_{\infty} \\ &= \frac{\frac{1}{2} - \sqrt{2}}{x} + (-)(0) \\ &= \frac{\frac{1}{2} - \sqrt{2}}{x} \\ &= \frac{1 - 2\sqrt{2}}{2x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2 \left(\frac{\frac{1}{2} - \sqrt{2}}{x} \right) (0) + \left(\left(-\frac{\frac{1}{2} - \sqrt{2}}{x^2} \right) + \left(\frac{\frac{1}{2} - \sqrt{2}}{x} \right)^2 - \left(\frac{7}{4x^2} \right) \right) &= 0 \\ 0 &= 0 \end{aligned}$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= p e^{\int \omega dx} \\ &= e^{\int \frac{\frac{1}{2} - \sqrt{2}}{x} dx} \\ &= x^{\frac{1}{2} - \sqrt{2}} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\&= z_1 e^{-\int \frac{1}{2} \frac{x}{x^2} dx} \\&= z_1 e^{-\frac{\ln(x)}{2}} \\&= z_1 \left(\frac{1}{\sqrt{x}} \right)\end{aligned}$$

Which simplifies to

$$y_1 = x^{-\sqrt{2}}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{x}{x^2} dx}}{(y_1)^2} dx \\&= y_1 \int \frac{e^{-\ln(x)}}{(y_1)^2} dx \\&= y_1 \left(\frac{\sqrt{2} x^{2\sqrt{2}}}{4} \right)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\&= c_1 \left(x^{-\sqrt{2}} \right) + c_2 \left(x^{-\sqrt{2}} \left(\frac{\sqrt{2} x^{2\sqrt{2}}}{4} \right) \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = x^{-\sqrt{2}} c_1 + \frac{c_2 \sqrt{2} x^{\sqrt{2}}}{4} \tag{1}$$

Verification of solutions

$$y = x^{-\sqrt{2}}c_1 + \frac{c_2\sqrt{2}x^{\sqrt{2}}}{4}$$

Verified OK.

9.32.6 Maple step by step solution

Let's solve

$$x^2y'' + xy' - 2y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{y'}{x} + \frac{2y}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{y'}{x} - \frac{2y}{x^2} = 0$$

- Multiply by denominators of the ODE

$$x^2y'' + xy' - 2y = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x , using the chain rule

$$y' = \left(\frac{d}{dt}y(t)\right) t'(x)$$

- Compute derivative

$$y' = \frac{\frac{d}{dt}y(t)}{x}$$

- Calculate the 2nd derivative of y with respect to x , using the chain rule

$$y'' = \left(\frac{d^2}{dt^2}y(t)\right) t'(x)^2 + t''(x) \left(\frac{d}{dt}y(t)\right)$$

- Compute derivative

$$y'' = \frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2}$$

Substitute the change of variables back into the ODE

$$x^2 \left(\frac{d^2 y(t)}{dt^2} - \frac{d y(t)}{dt} \right) + \frac{d y(t)}{dt} - 2y(t) = 0$$

- Simplify

$$\frac{d^2 y(t)}{dt^2} - 2y(t) = 0$$

- Characteristic polynomial of ODE

$$r^2 - 2 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{8})}{2}$$

- Roots of the characteristic polynomial

$$r = (\sqrt{2}, -\sqrt{2})$$

- 1st solution of the ODE

$$y_1(t) = e^{\sqrt{2}t}$$

- 2nd solution of the ODE

$$y_2(t) = e^{-\sqrt{2}t}$$

- General solution of the ODE

$$y(t) = c_1 y_1(t) + c_2 y_2(t)$$

- Substitute in solutions

$$y(t) = c_1 e^{\sqrt{2}t} + c_2 e^{-\sqrt{2}t}$$

- Change variables back using $t = \ln(x)$

$$y = c_1 e^{\sqrt{2} \ln(x)} + c_2 e^{-\sqrt{2} \ln(x)}$$

- Simplify

$$y = c_1 x^{\sqrt{2}} + c_2 x^{-\sqrt{2}}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
<- LODE of Euler type successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 21

```
dsolve(x^2*diff(y(x),x)+x*diff(y(x),x)-2*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1x^{\sqrt{2}} + c_2x^{-\sqrt{2}}$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 28

```
DSolve[x^2*y'[x]+x*y'[x]-2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1x^{-\sqrt{2}} + c_2x^{\sqrt{2}}$$

9.33 problem 5(i)

9.33.1 Solving as second order euler ode ode	1622
9.33.2 Solving as second order change of variable on x method 2 ode .	1623
9.33.3 Solving as second order change of variable on x method 1 ode .	1626
9.33.4 Solving as second order change of variable on y method 2 ode .	1628
9.33.5 Solving using Kovacic algorithm	1630
9.33.6 Maple step by step solution	1635

Internal problem ID [6301]

Internal file name [OUTPUT/5549_Sunday_June_05_2022_03_43_11_PM_13904494/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.1. Linear Equations with Constant Coefficients. Page 62

Problem number: 5(i).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_euler_ode", "second_order_change_of_variable_on_x_method_1", "second_order_change_of_variable_on_x_method_2", "second_order_change_of_variable_on_y_method_2"

Maple gives the following as the ode type

```
[[_Emden, _Fowler], [_2nd_order, _linear, `_with_symmetry_[0,F(x)]`]]
```

$$x^2y'' + xy' - 16y = 0$$

9.33.1 Solving as second order euler ode ode

This is Euler second order ODE. Let the solution be $y = x^r$, then $y' = rx^{r-1}$ and $y'' = r(r-1)x^{r-2}$. Substituting these back into the given ODE gives

$$x^2(r(r-1))x^{r-2} + xrx^{r-1} - 16x^r = 0$$

Simplifying gives

$$r(r-1)x^r + rx^r - 16x^r = 0$$

Since $x^r \neq 0$ then dividing throughout by x^r gives

$$r(r - 1) + r - 16 = 0$$

Or

$$r^2 - 16 = 0 \quad (1)$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = -4$$

$$r_2 = 4$$

Since the roots are real and distinct, then the general solution is

$$y = c_1 y_1 + c_2 y_2$$

Where $y_1 = x^{r_1}$ and $y_2 = x^{r_2}$. Hence

$$y = \frac{c_1}{x^4} + c_2 x^4$$

Summary

The solution(s) found are the following

$$y = \frac{c_1}{x^4} + c_2 x^4 \quad (1)$$

Verification of solutions

$$y = \frac{c_1}{x^4} + c_2 x^4$$

Verified OK.

9.33.2 Solving as second order change of variable on x method 2 ode

In normal form the ode

$$x^2 y'' + xy' - 16y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = -\frac{16}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x)\tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned} \tau &= \int e^{-(\int p(x)dx)} dx \\ &= \int e^{-(\int \frac{1}{x}dx)} dx \\ &= \int e^{-\ln(x)} dx \\ &= \int \frac{1}{x} dx \\ &= \ln(x) \end{aligned} \quad (6)$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned} q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\ &= \frac{-16}{x^2} \\ &= \frac{1}{x^2} \\ &= -16 \end{aligned} \quad (7)$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned} \frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) - 16y(\tau) &= 0 \end{aligned}$$

The above ode is now solved for $y(\tau)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(\tau) + By'(\tau) + Cy(\tau) = 0$$

Where in the above $A = 1, B = 0, C = -16$. Let the solution be $y(\tau) = e^{\lambda\tau}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda\tau} - 16 e^{\lambda\tau} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda\tau}$ gives

$$\lambda^2 - 16 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = -16$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(-16)} \\ &= \pm 4 \end{aligned}$$

Hence

$$\lambda_1 = +4$$

$$\lambda_2 = -4$$

Which simplifies to

$$\lambda_1 = 4$$

$$\lambda_2 = -4$$

Since roots are real and distinct, then the solution is

$$y(\tau) = c_1 e^{\lambda_1 \tau} + c_2 e^{\lambda_2 \tau}$$

$$y(\tau) = c_1 e^{(4)\tau} + c_2 e^{(-4)\tau}$$

Or

$$y(\tau) = c_1 e^{4\tau} + c_2 e^{-4\tau}$$

The above solution is now transformed back to y using (6) which results in

$$y = \frac{c_1 x^8 + c_2}{x^4}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1 x^8 + c_2}{x^4} \quad (1)$$

Verification of solutions

$$y = \frac{c_1 x^8 + c_2}{x^4}$$

Verified OK.

9.33.3 Solving as second order change of variable on x method 1 ode

In normal form the ode

$$x^2 y'' + x y' - 16y = 0 \quad (1)$$

Becomes

$$y'' + p(x) y' + q(x) y = 0 \quad (2)$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = -\frac{16}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) results

$$\frac{d^2}{d\tau^2} y(\tau) + p_1 \left(\frac{d}{d\tau} y(\tau) \right) + q_1 y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x) \tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $q_1 = c^2$ where c is some constant. Therefore from (5)

$$\begin{aligned}\tau' &= \frac{1}{c}\sqrt{q} \\ &= \frac{4\sqrt{-\frac{1}{x^2}}}{c} \\ \tau'' &= \frac{4}{c\sqrt{-\frac{1}{x^2}}x^3}\end{aligned}\tag{6}$$

Substituting the above into (4) results in

$$\begin{aligned}p_1(\tau) &= \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \\ &= \frac{\frac{4}{c\sqrt{-\frac{1}{x^2}}x^3} + \frac{1}{x}\frac{4\sqrt{-\frac{1}{x^2}}}{c}}{\left(\frac{4\sqrt{-\frac{1}{x^2}}}{c}\right)^2} \\ &= 0\end{aligned}$$

Therefore ode (3) now becomes

$$\begin{aligned}y(\tau)'' + p_1y(\tau)' + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) + c^2y(\tau) &= 0\end{aligned}\tag{7}$$

The above ode is now solved for $y(\tau)$. Since the ode is now constant coefficients, it can be easily solved to give

$$y(\tau) = c_1 \cos(c\tau) + c_2 \sin(c\tau)$$

Now from (6)

$$\begin{aligned}\tau &= \int \frac{1}{c}\sqrt{q} dx \\ &= \frac{\int 4\sqrt{-\frac{1}{x^2}} dx}{c} \\ &= \frac{4\sqrt{-\frac{1}{x^2}}x \ln(x)}{c}\end{aligned}$$

Substituting the above into the solution obtained gives

$$y = c_1 \cosh(4 \ln(x)) + ic_2 \sinh(4 \ln(x))$$

Summary

The solution(s) found are the following

$$y = c_1 \cosh(4 \ln(x)) + ic_2 \sinh(4 \ln(x)) \quad (1)$$

Verification of solutions

$$y = c_1 \cosh(4 \ln(x)) + ic_2 \sinh(4 \ln(x))$$

Verified OK.

9.33.4 Solving as second order change of variable on y method 2 ode

In normal form the ode

$$x^2 y'' + xy' - 16y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = -\frac{16}{x^2}$$

Applying change of variables on the dependent variable $y = v(x)x^n$ to (2) gives the following ode where the dependent variables is $v(x)$ and not y .

$$v''(x) + \left(\frac{2n}{x} + p\right)v'(x) + \left(\frac{n(n-1)}{x^2} + \frac{np}{x} + q\right)v(x) = 0 \quad (3)$$

Let the coefficient of $v(x)$ above be zero. Hence

$$\frac{n(n-1)}{x^2} + \frac{np}{x} + q = 0 \quad (4)$$

Substituting the earlier values found for $p(x)$ and $q(x)$ into (4) gives

$$\frac{n(n-1)}{x^2} + \frac{n}{x^2} - \frac{16}{x^2} = 0 \quad (5)$$

Solving (5) for n gives

$$n = 4 \quad (6)$$

Substituting this value in (3) gives

$$\begin{aligned}v''(x) + \frac{9v'(x)}{x} &= 0 \\v''(x) + \frac{9v'(x)}{x} &= 0\end{aligned}\tag{7}$$

Using the substitution

$$u(x) = v'(x)$$

Then (7) becomes

$$u'(x) + \frac{9u(x)}{x} = 0\tag{8}$$

The above is now solved for $u(x)$. In canonical form the ODE is

$$\begin{aligned}u' &= F(x, u) \\&= f(x)g(u) \\&= -\frac{9u}{x}\end{aligned}$$

Where $f(x) = -\frac{9}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u} du &= -\frac{9}{x} dx \\ \int \frac{1}{u} du &= \int -\frac{9}{x} dx \\ \ln(u) &= -9 \ln(x) + c_1 \\ u &= e^{-9 \ln(x) + c_1} \\ &= \frac{c_1}{x^9}\end{aligned}$$

Now that $u(x)$ is known, then

$$\begin{aligned}v'(x) &= u(x) \\ v(x) &= \int u(x) dx + c_2 \\ &= -\frac{c_1}{8x^8} + c_2\end{aligned}$$

Hence

$$\begin{aligned}y &= v(x) x^n \\ &= \left(-\frac{c_1}{8x^8} + c_2\right) x^4 \\ &= \frac{8c_2x^8 - c_1}{8x^4}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \left(-\frac{c_1}{8x^8} + c_2\right) x^4 \quad (1)$$

Verification of solutions

$$y = \left(-\frac{c_1}{8x^8} + c_2\right) x^4$$

Verified OK.

9.33.5 Solving using Kovacic algorithm

Writing the ode as

$$x^2y'' + xy' - 16y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= x^2 \\ B &= x \\ C &= -16\end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{63}{4x^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 63 \\ t &= 4x^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{63}{4x^2} \right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 239: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = \frac{63}{4x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = \frac{63}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{9}{2} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = -\frac{7}{2} \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = \frac{63}{4x^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = \frac{63}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{9}{2} \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = -\frac{7}{2} \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{63}{4x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	$\frac{9}{2}$	$-\frac{7}{2}$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	$\frac{9}{2}$	$-\frac{7}{2}$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = -\frac{7}{2}$ then

$$\begin{aligned} d &= \alpha_\infty^- - (\alpha_{c_1}^-) \\ &= -\frac{7}{2} - \left(-\frac{7}{2}\right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c) [\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty) [\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= \left((-) [\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{x - c_1} \right) + (-) [\sqrt{r}]_\infty \\ &= -\frac{7}{2x} + (-) (0) \\ &= -\frac{7}{2x} \\ &= -\frac{7}{2x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \tag{1A}$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$(0) + 2\left(-\frac{7}{2x}\right)(0) + \left(\left(\frac{7}{2x^2}\right) + \left(-\frac{7}{2x}\right)^2 - \left(\frac{63}{4x^2}\right)\right) = 0$$
$$0 = 0$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$z_1(x) = pe^{\int \omega dx}$$
$$= e^{\int -\frac{7}{2x} dx}$$
$$= \frac{1}{x^{\frac{7}{2}}}$$

The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$
$$= z_1 e^{-\int \frac{1}{2} \frac{x}{x^2} dx}$$
$$= z_1 e^{-\frac{\ln(x)}{2}}$$
$$= z_1 \left(\frac{1}{\sqrt{x}}\right)$$

Which simplifies to

$$y_1 = \frac{1}{x^4}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$y_2 = y_1 \int \frac{e^{\int -\frac{x}{x^2} dx}}{(y_1)^2} dx$$
$$= y_1 \int \frac{e^{-\ln(x)}}{(y_1)^2} dx$$
$$= y_1 \left(\frac{x^8}{8}\right)$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(\frac{1}{x^4} \right) + c_2 \left(\frac{1}{x^4} \left(\frac{x^8}{8} \right) \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1}{x^4} + \frac{c_2 x^4}{8} \quad (1)$$

Verification of solutions

$$y = \frac{c_1}{x^4} + \frac{c_2 x^4}{8}$$

Verified OK.

9.33.6 Maple step by step solution

Let's solve

$$x^2 y'' + x y' - 16y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{y'}{x} + \frac{16y}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{y'}{x} - \frac{16y}{x^2} = 0$$

- Multiply by denominators of the ODE

$$x^2 y'' + x y' - 16y = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x , using the chain rule

$$y' = \left(\frac{d}{dt} y(t) \right) t'(x)$$

- Compute derivative

$$y' = \frac{\frac{d}{dt}y(t)}{x}$$

- Calculate the 2nd derivative of y with respect to x , using the chain rule

$$y'' = \left(\frac{d^2}{dt^2}y(t) \right) t'(x)^2 + t''(x) \left(\frac{d}{dt}y(t) \right)$$

- Compute derivative

$$y'' = \frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2}$$

Substitute the change of variables back into the ODE

$$x^2 \left(\frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2} \right) + \frac{d}{dt}y(t) - 16y(t) = 0$$

- Simplify

$$\frac{d^2}{dt^2}y(t) - 16y(t) = 0$$

- Characteristic polynomial of ODE

$$r^2 - 16 = 0$$

- Factor the characteristic polynomial

$$(r - 4)(r + 4) = 0$$

- Roots of the characteristic polynomial

$$r = (-4, 4)$$

- 1st solution of the ODE

$$y_1(t) = e^{-4t}$$

- 2nd solution of the ODE

$$y_2(t) = e^{4t}$$

- General solution of the ODE

$$y(t) = c_1y_1(t) + c_2y_2(t)$$

- Substitute in solutions

$$y(t) = c_1e^{-4t} + c_2e^{4t}$$

- Change variables back using $t = \ln(x)$

$$y = \frac{c_1}{x^4} + c_2x^4$$

- Simplify

$$y = \frac{c_1}{x^4} + c_2x^4$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
<- LODE of Euler type successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)-16*y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{c_1 x^8 + c_2}{x^4}$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 18

```
DSolve[x^2*y'[x]+x*y'[x]-16*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_2 x^8 + c_1}{x^4}$$

**10 Chapter 2. Second-Order Linear Equations.
Section 2.2. THE METHOD OF
UNDETERMINED COEFFICIENTS. Page 67**

10.1	problem 1(a)	1639
10.2	problem 1(b)	1650
10.3	problem 1(c)	1661
10.4	problem 1(d)	1673
10.5	problem 1(e)	1684
10.6	problem 1(f)	1695
10.7	problem 1(g)	1705
10.8	problem 1(h)	1716
10.9	problem 1(i)	1736
10.10	problem 1(j)	1748
10.11	problem 1(k)	1759
10.12	problem 3(a)	1779
10.13	problem 3(b)	1790
10.14	problem 4(a)	1801
10.15	problem 4(b)	1812

10.1 problem 1(a)

- 10.1.1 Solving as second order linear constant coeff ode 1639
- 10.1.2 Solving using Kovacic algorithm 1642
- 10.1.3 Maple step by step solution 1647

Internal problem ID [6302]

Internal file name [OUTPUT/5550_Sunday_June_05_2022_03_43_12_PM_7678405/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.2. THE METHOD OF UNDETERMINED COEFFICIENTS. Page 67

Problem number: 1(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' + 3y' - 10y = 6e^{4x}$$

10.1.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 3, C = -10, f(x) = 6e^{4x}$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 3y' - 10y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 3, C = -10$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 3\lambda e^{\lambda x} - 10 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 3\lambda - 10 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 3, C = -10$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-3}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{3^2 - (4)(1)(-10)} \\ &= -\frac{3}{2} \pm \frac{7}{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= -\frac{3}{2} + \frac{7}{2} \\ \lambda_2 &= -\frac{3}{2} - \frac{7}{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= 2 \\ \lambda_2 &= -5 \end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned} y &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ y &= c_1 e^{(2)x} + c_2 e^{(-5)x} \end{aligned}$$

Or

$$y = c_1 e^{2x} + c_2 e^{-5x}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^{2x} + c_2 e^{-5x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$6 e^{4x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{4x}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^{-5x}, e^{2x}\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^{4x}$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$18A_1 e^{4x} = 6 e^{4x}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{3} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{e^{4x}}{3}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^{2x} + c_2 e^{-5x}) + \left(\frac{e^{4x}}{3} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{2x} + c_2 e^{-5x} + \frac{e^{4x}}{3} \quad (1)$$




Figure 301: Slope field plot

Verification of solutions

$$y = c_1 e^{2x} + c_2 e^{-5x} + \frac{e^{4x}}{3}$$

Verified OK.

10.1.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 3y' - 10y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 3 \\ C &= -10 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{49}{4} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 49 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \frac{49z(x)}{4} \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 241: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{49}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{7x}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{3}{1} dx} \\ &= z_1 e^{-\frac{3x}{2}} \\ &= z_1 \left(e^{-\frac{3x}{2}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-5x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{3}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-3x}}{(y_1)^2} dx \\ &= y_1 \left(\frac{e^{7x}}{7} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-5x}) + c_2 \left(e^{-5x} \left(\frac{e^{7x}}{7} \right) \right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 3y' - 10y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-5x} + \frac{c_2 e^{2x}}{7}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$6e^{4x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{4x}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ \frac{e^{2x}}{7}, e^{-5x} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^{4x}$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$18A_1e^{4x} = 6e^{4x}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{3} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{e^{4x}}{3}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1e^{-5x} + \frac{c_2e^{2x}}{7} \right) + \left(\frac{e^{4x}}{3} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1e^{-5x} + \frac{c_2e^{2x}}{7} + \frac{e^{4x}}{3} \quad (1)$$




Figure 302: Slope field plot

Verification of solutions

$$y = c_1 e^{-5x} + \frac{c_2 e^{2x}}{7} + \frac{e^{4x}}{3}$$

Verified OK.

10.1.3 Maple step by step solution

Let's solve

$$y'' + 3y' - 10y = 6e^{4x}$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 3r - 10 = 0$$

- Factor the characteristic polynomial

$$(r + 5)(r - 2) = 0$$

- Roots of the characteristic polynomial

$$r = (-5, 2)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-5x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{2x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{-5x} + c_2 e^{2x} + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = 6 e^{4x} \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-5x} & e^{2x} \\ -5e^{-5x} & 2e^{2x} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 7 e^{-3x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = \frac{6(e^{7x} (\int e^{2x} dx) - (\int e^{9x} dx)) e^{-5x}}{7}$$

- Compute integrals

$$y_p(x) = \frac{e^{4x}}{3}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{-5x} + c_2 e^{2x} + \frac{e^{4x}}{3}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 25

```
dsolve(diff(y(x),x$2)+3*diff(y(x),x)-10*y(x)=6*exp(4*x),y(x), singsol=all)
```

$$y(x) = \frac{(e^{9x} + 3e^{7x}c_1 + 3c_2)e^{-5x}}{3}$$

✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 31

```
DSolve[y''[x]+3*y'[x]-10*y[x]==6*Exp[4*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{4x}}{3} + c_1 e^{-5x} + c_2 e^{2x}$$

10.2 problem 1(b)

10.2.1 Solving as second order linear constant coeff ode	1650
10.2.2 Solving using Kovacic algorithm	1653
10.2.3 Maple step by step solution	1658

Internal problem ID [6303]

Internal file name [OUTPUT/5551_Sunday_June_05_2022_03_43_14_PM_75364838/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.2. THE METHOD OF UN-DETERMINED COEFFICIENTS. Page 67

Problem number: 1(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + 4y = 3 \sin(x)$$

10.2.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 4, f(x) = 3 \sin(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 4y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 4$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 4 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 4 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 4$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(4)} \\ &= \pm 2i \end{aligned}$$

Hence

$$\lambda_1 = +2i$$

$$\lambda_2 = -2i$$

Which simplifies to

$$\lambda_1 = 2i$$

$$\lambda_2 = -2i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 2$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (c_1 \cos(2x) + c_2 \sin(2x))$$

Or

$$y = c_1 \cos(2x) + c_2 \sin(2x)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \cos(2x) + c_2 \sin(2x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$3 \sin(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(2x), \sin(2x)\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(x) + A_2 \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$3A_1 \cos(x) + 3A_2 \sin(x) = 3 \sin(x)$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 0, A_2 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \sin(x)$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(2x) + c_2 \sin(2x)) + (\sin(x)) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(2x) + c_2 \sin(2x) + \sin(x) \quad (1)$$




Figure 303: Slope field plot

Verification of solutions

$$y = c_1 \cos(2x) + c_2 \sin(2x) + \sin(x)$$

Verified OK.

10.2.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 4y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 0 \quad (3)$$

$$C = 4$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-4}{1} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -4 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -4z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 243: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -4$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(2x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned} y_1 &= z_1 \\ &= \cos(2x) \end{aligned}$$

Which simplifies to

$$y_1 = \cos(2x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(2x) \int \frac{1}{\cos(2x)^2} dx \\ &= \cos(2x) \left(\frac{\tan(2x)}{2} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(2x)) + c_2 \left(\cos(2x) \left(\frac{\tan(2x)}{2} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 4y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 \cos(2x) + \frac{c_2 \sin(2x)}{2}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$3 \sin (x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos (x), \sin (x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ \frac{\sin (2x)}{2}, \cos (2x) \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos (x) + A_2 \sin (x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$3A_1 \cos (x) + 3A_2 \sin (x) = 3 \sin (x)$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 0, A_2 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \sin (x)$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 \cos (2x) + \frac{c_2 \sin (2x)}{2} \right) + (\sin (x)) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \cos (2x) + \frac{c_2 \sin (2x)}{2} + \sin (x) \quad (1)$$




Figure 304: Slope field plot

Verification of solutions

$$y = c_1 \cos(2x) + \frac{c_2 \sin(2x)}{2} + \sin(x)$$

Verified OK.

10.2.3 Maple step by step solution

Let's solve

$$y'' + 4y = 3 \sin(x)$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 4 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-16})}{2}$$

- Roots of the characteristic polynomial

$$r = (-2I, 2I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(2x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(2x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 \cos(2x) + c_2 \sin(2x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = 3 \sin(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(2x) & \sin(2x) \\ -2 \sin(2x) & 2 \cos(2x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 2$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -3 \cos(2x) \left(\int \cos(x) \sin(x)^2 dx \right) + \frac{3 \sin(2x) \left(\int (\sin(3x) - \sin(x)) dx \right)}{4}$$

- Compute integrals

$$y_p(x) = \sin(x)$$

- Substitute particular solution into general solution to ODE

$$y = c_1 \cos(2x) + c_2 \sin(2x) + \sin(x)$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 19

```
dsolve(diff(y(x),x$2)+4*y(x)=3*sin(x),y(x), singsol=all)
```

$$y(x) = \sin(2x) c_2 + \cos(2x) c_1 + \sin(x)$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 22

```
DSolve[y''[x]+4*y[x]==3*Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \sin(x) + c_1 \cos(2x) + c_2 \sin(2x)$$

10.3 problem 1(c)

10.3.1 Solving as second order linear constant coeff ode	1661
10.3.2 Solving as linear second order ode solved by an integrating factor ode	1664
10.3.3 Solving using Kovacic algorithm	1666
10.3.4 Maple step by step solution	1671

Internal problem ID [6304]

Internal file name [OUTPUT/5552_Sunday_June_05_2022_03_43_16_PM_66884/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.2. THE METHOD OF UNDETERMINED COEFFICIENTS. Page 67

Problem number: 1(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "linear_second_order_ode_solved_by_an_integrating_factor"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' + 10y' + 25y = 14e^{-5x}$$

10.3.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 10, C = 25, f(x) = 14e^{-5x}$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 10y' + 25y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 10, C = 25$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 10\lambda e^{\lambda x} + 25 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 10\lambda + 25 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 10, C = 25$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-10}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{(10)^2 - (4)(1)(25)} \\ &= -5 \end{aligned}$$

Hence this is the case of a double root $\lambda_{1,2} = 5$. Therefore the solution is

$$y = c_1 e^{-5x} + c_2 x e^{-5x} \quad (1)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^{-5x} + c_2 x e^{-5x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$14 e^{-5x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$\{e^{-5x}\}$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{x e^{-5x}, e^{-5x}\}$$

Since e^{-5x} is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x e^{-5x}\}]$$

Since $x e^{-5x}$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x^2 e^{-5x}\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x^2 e^{-5x}$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$2A_1 e^{-5x} = 14 e^{-5x}$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 7]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = 7x^2 e^{-5x}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^{-5x} + c_2 x e^{-5x}) + (7x^2 e^{-5x}) \end{aligned}$$

Which simplifies to

$$y = e^{-5x}(c_2 x + c_1) + 7x^2 e^{-5x}$$

Summary

The solution(s) found are the following

$$y = e^{-5x}(c_2 x + c_1) + 7x^2 e^{-5x} \quad (1)$$




Figure 305: Slope field plot

Verification of solutions

$$y = e^{-5x}(c_2x + c_1) + 7x^2e^{-5x}$$

Verified OK.

10.3.2 Solving as linear second order ode solved by an integrating factor ode

The ode satisfies this form

$$y'' + p(x)y' + \frac{(p(x))^2 + p'(x)}{2}y = f(x)$$

Where $p(x) = 10$. Therefore, there is an integrating factor given by

$$\begin{aligned} M(x) &= e^{\frac{1}{2} \int p dx} \\ &= e^{\int 10 dx} \\ &= e^{5x} \end{aligned}$$

Multiplying both sides of the ODE by the integrating factor $M(x)$ makes the left side of the ODE a complete differential

$$(M(x)y)'' = 14e^{5x}e^{-5x}$$

$$(e^{5x}y)'' = 14e^{5x}e^{-5x}$$

Integrating once gives

$$(e^{5x}y)' = 14x + c_1$$

Integrating again gives

$$(e^{5x}y) = x(c_1 + 7x) + c_2$$

Hence the solution is

$$y = \frac{x(c_1 + 7x) + c_2}{e^{5x}}$$

Or

$$y = c_1x e^{-5x} + 7x^2 e^{-5x} + c_2 e^{-5x}$$

Summary

The solution(s) found are the following

$$y = c_1x e^{-5x} + 7x^2 e^{-5x} + c_2 e^{-5x} \tag{1}$$




Figure 306: Slope field plot

Verification of solutions

$$y = c_1 x e^{-5x} + 7x^2 e^{-5x} + c_2 e^{-5x}$$

Verified OK.

10.3.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 10y' + 25y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 10 \\ C &= 25 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 0 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 0 \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 245: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{10}{1} dx} \\ &= z_1 e^{-5x} \\ &= z_1 (e^{-5x}) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-5x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{10}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-10x}}{(y_1)^2} dx \\ &= y_1(x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-5x}) + c_2 (e^{-5x}(x)) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 10y' + 25y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-5x} + c_2 x e^{-5x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$14 e^{-5x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[{\{e^{-5x}\}}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{x e^{-5x}, e^{-5x}\}$$

Since e^{-5x} is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[{\{x e^{-5x}\}}]$$

Since $x e^{-5x}$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[{\{x^2 e^{-5x}\}}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x^2 e^{-5x}$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$2A_1 e^{-5x} = 14 e^{-5x}$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 7]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = 7x^2 e^{-5x}$$

Therefore the general solution is

$$\begin{aligned}y &= y_h + y_p \\ &= (c_1 e^{-5x} + c_2 x e^{-5x}) + (7x^2 e^{-5x})\end{aligned}$$

Which simplifies to

$$y = e^{-5x}(c_2 x + c_1) + 7x^2 e^{-5x}$$

Summary

The solution(s) found are the following

$$y = e^{-5x}(c_2 x + c_1) + 7x^2 e^{-5x} \quad (1)$$




Figure 307: Slope field plot

Verification of solutions

$$y = e^{-5x}(c_2 x + c_1) + 7x^2 e^{-5x}$$

Verified OK.

10.3.4 Maple step by step solution

Let's solve

$$y'' + 10y' + 25y = 14e^{-5x}$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 10r + 25 = 0$$

- Factor the characteristic polynomial

$$(r + 5)^2 = 0$$

- Root of the characteristic polynomial

$$r = -5$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-5x}$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = x e^{-5x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{-5x} + c_2 x e^{-5x} + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = 14e^{-5x} \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-5x} & x e^{-5x} \\ -5e^{-5x} & e^{-5x} - 5x e^{-5x} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = e^{-10x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -14e^{-5x} \left(\int x dx - \left(\int 1 dx \right) x \right)$$

- Compute integrals

$$y_p(x) = 7x^2e^{-5x}$$

- Substitute particular solution into general solution to ODE

$$y = c_2x e^{-5x} + 7x^2e^{-5x} + c_1e^{-5x}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 19

```
dsolve(diff(y(x),x$2)+10*diff(y(x),x)+25*y(x)=14*exp(-5*x),y(x), singsol=all)
```

$$y(x) = e^{-5x}(c_1x + 7x^2 + c_2)$$

✓ Solution by Mathematica

Time used: 0.039 (sec). Leaf size: 23

```
DSolve[y''[x]+10*y'[x]+25*y[x]==14*Exp[-5*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-5x}(7x^2 + c_2x + c_1)$$

10.4 problem 1(d)

10.4.1 Solving as second order linear constant coeff ode	1673
10.4.2 Solving using Kovacic algorithm	1676
10.4.3 Maple step by step solution	1681

Internal problem ID [6305]

Internal file name [OUTPUT/5553_Sunday_June_05_2022_03_43_18_PM_32183444/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.2. THE METHOD OF UNDETERMINED COEFFICIENTS. Page 67

Problem number: 1(d).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - 2y' + 5y = 25x^2 + 12$$

10.4.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = -2, C = 5, f(x) = 25x^2 + 12$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 2y' + 5y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -2, C = 5$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 2\lambda e^{\lambda x} + 5 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 2\lambda + 5 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -2, C = 5$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{2}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-2^2 - (4)(1)(5)} \\ &= 1 \pm 2i \end{aligned}$$

Hence

$$\lambda_1 = 1 + 2i$$

$$\lambda_2 = 1 - 2i$$

Which simplifies to

$$\lambda_1 = 1 + 2i$$

$$\lambda_2 = 1 - 2i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 1$ and $\beta = 2$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^x (c_1 \cos(2x) + c_2 \sin(2x))$$

Therefore the homogeneous solution y_h is

$$y_h = e^x (c_1 \cos(2x) + c_2 \sin(2x))$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x^2 + 1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x, x^2\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^x \cos(2x), e^x \sin(2x)\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_3x^2 + A_2x + A_1$$

The unknowns $\{A_1, A_2, A_3\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$5A_3x^2 + 5A_2x - 4xA_3 + 5A_1 - 2A_2 + 2A_3 = 25x^2 + 12$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 2, A_2 = 4, A_3 = 5]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = 5x^2 + 4x + 2$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (e^x(c_1 \cos(2x) + c_2 \sin(2x))) + (5x^2 + 4x + 2) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = e^x(c_1 \cos(2x) + c_2 \sin(2x)) + 5x^2 + 4x + 2 \quad (1)$$




Figure 308: Slope field plot

Verification of solutions

$$y = e^x(c_1 \cos(2x) + c_2 \sin(2x)) + 5x^2 + 4x + 2$$

Verified OK.

10.4.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 2y' + 5y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -2 \\ C &= 5 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} = \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-4}{1} \tag{6}$$

Comparing the above to (5) shows that

$$s = -4$$

$$t = 1$$

Therefore eq. (4) becomes

$$z''(x) = -4z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 247: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -4$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(2x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-2}{1} dx} \\ &= z_1 e^x \\ &= z_1 (e^x) \end{aligned}$$

Which simplifies to

$$y_1 = e^x \cos(2x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-2}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{2x}}{(y_1)^2} dx \\ &= y_1 \left(\frac{\tan(2x)}{2} \right) \end{aligned}$$

Therefore the solution is

$$y = c_1 y_1 + c_2 y_2$$

$$= c_1 (e^x \cos(2x)) + c_2 \left(e^x \cos(2x) \left(\frac{\tan(2x)}{2} \right) \right)$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 2y' + 5y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = \cos(2x) e^x c_1 + \frac{\sin(2x) e^x c_2}{2}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x^2 + 1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x, x^2\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^x \cos(2x), \frac{e^x \sin(2x)}{2} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_3 x^2 + A_2 x + A_1$$

The unknowns $\{A_1, A_2, A_3\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$5A_3x^2 + 5A_2x - 4xA_3 + 5A_1 - 2A_2 + 2A_3 = 25x^2 + 12$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 2, A_2 = 4, A_3 = 5]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = 5x^2 + 4x + 2$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(\cos(2x) e^x c_1 + \frac{\sin(2x) e^x c_2}{2} \right) + (5x^2 + 4x + 2) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \cos(2x) e^x c_1 + \frac{\sin(2x) e^x c_2}{2} + 5x^2 + 4x + 2 \quad (1)$$




Figure 309: Slope field plot

Verification of solutions

$$y = \cos(2x) e^x c_1 + \frac{\sin(2x) e^x c_2}{2} + 5x^2 + 4x + 2$$

Verified OK.

10.4.3 Maple step by step solution

Let's solve

$$y'' - 2y' + 5y = 25x^2 + 12$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - 2r + 5 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{2 \pm (\sqrt{-16})}{2}$$

- Roots of the characteristic polynomial

$$r = (1 - 2I, 1 + 2I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^x \cos(2x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^x \sin(2x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = \cos(2x) e^x c_1 + \sin(2x) e^x c_2 + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = 25x^2 + 12 \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^x \cos(2x) & e^x \sin(2x) \\ e^x \cos(2x) - 2e^x \sin(2x) & e^x \sin(2x) + 2e^x \cos(2x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 2e^{2x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{e^x(\cos(2x)(\int e^{-x} \sin(2x)(25x^2+12)dx) - \sin(2x)(\int e^{-x} \cos(2x)(25x^2+12)dx))}{2}$$

- Compute integrals

$$y_p(x) = 5x^2 + 4x + 2$$

- Substitute particular solution into general solution to ODE

$$y = \sin(2x) e^x c_2 + \cos(2x) e^x c_1 + 5x^2 + 4x + 2$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 30

```
dsolve(diff(y(x),x$2)-2*diff(y(x),x)+5*y(x)=25*x^2+12,y(x), singsol=all)
```

$$y(x) = e^x \sin(2x) c_2 + e^x \cos(2x) c_1 + 5x^2 + 4x + 2$$

✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 35

```
DSolve[y''[x]-2*y'[x]+5*y[x]==25*x^2+12,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 5x^2 + 4x + c_2 e^x \cos(2x) + c_1 e^x \sin(2x) + 2$$

10.5 problem 1(e)

10.5.1 Solving as second order linear constant coeff ode	1684
10.5.2 Solving using Kovacic algorithm	1687
10.5.3 Maple step by step solution	1692

Internal problem ID [6306]

Internal file name [OUTPUT/5554_Sunday_June_05_2022_03_43_20_PM_43368877/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.2. THE METHOD OF UN-DETERMINED COEFFICIENTS. Page 67

Problem number: 1(e).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - y' - 6y = 20e^{-2x}$$

10.5.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = -1, C = -6, f(x) = 20e^{-2x}$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - y' - 6y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -1, C = -6$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - \lambda e^{\lambda x} - 6 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - \lambda - 6 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -1, C = -6$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-1^2 - (4)(1)(-6)} \\ &= \frac{1}{2} \pm \frac{5}{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= \frac{1}{2} + \frac{5}{2} \\ \lambda_2 &= \frac{1}{2} - \frac{5}{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= 3 \\ \lambda_2 &= -2 \end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned} y &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ y &= c_1 e^{(3)x} + c_2 e^{(-2)x} \end{aligned}$$

Or

$$y = e^{3x} c_1 + c_2 e^{-2x}$$

Therefore the homogeneous solution y_h is

$$y_h = e^{3x} c_1 + c_2 e^{-2x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$20 e^{-2x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{-2x}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^{-2x}, e^{3x}\}$$

Since e^{-2x} is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x e^{-2x}\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x e^{-2x}$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-5A_1 e^{-2x} = 20 e^{-2x}$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = -4]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -4x e^{-2x}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (e^{3x} c_1 + c_2 e^{-2x}) + (-4x e^{-2x}) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = e^{3x} c_1 + c_2 e^{-2x} - 4x e^{-2x} \quad (1)$$




Figure 310: Slope field plot

Verification of solutions

$$y = e^{3x} c_1 + c_2 e^{-2x} - 4x e^{-2x}$$

Verified OK.

10.5.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' - y' - 6y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = -1 \quad (3)$$

$$C = -6$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{25}{4} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 25 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \frac{25z(x)}{4} \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 249: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{25}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{5x}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}
 y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\
 &= z_1 e^{-\int \frac{1}{2} \frac{-1}{1} dx} \\
 &= z_1 e^{\frac{x}{2}} \\
 &= z_1 \left(e^{\frac{x}{2}} \right)
 \end{aligned}$$

Which simplifies to

$$y_1 = e^{-2x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-1}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^x}{(y_1)^2} dx \\ &= y_1 \left(\frac{e^{5x}}{5} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-2x}) + c_2 \left(e^{-2x} \left(\frac{e^{5x}}{5} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - y' - 6y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-2x} + \frac{c_2 e^{3x}}{5}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$20 e^{-2x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{-2x}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ \frac{e^{3x}}{5}, e^{-2x} \right\}$$

Since e^{-2x} is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x e^{-2x}\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x e^{-2x}$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-5A_1 e^{-2x} = 20 e^{-2x}$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = -4]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -4x e^{-2x}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 e^{-2x} + \frac{c_2 e^{3x}}{5} \right) + (-4x e^{-2x}) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-2x} + \frac{c_2 e^{3x}}{5} - 4x e^{-2x} \quad (1)$$




Figure 311: Slope field plot

Verification of solutions

$$y = c_1 e^{-2x} + \frac{c_2 e^{3x}}{5} - 4x e^{-2x}$$

Verified OK.

10.5.3 Maple step by step solution

Let's solve

$$y'' - y' - 6y = 20e^{-2x}$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - r - 6 = 0$$

- Factor the characteristic polynomial

$$(r + 2)(r - 3) = 0$$

- Roots of the characteristic polynomial

$$r = (-2, 3)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-2x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{3x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{-2x} + c_2 e^{3x} + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = 20 e^{-2x} \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-2x} & e^{3x} \\ -2e^{-2x} & 3e^{3x} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 5 e^x$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -4(-e^{5x}(\int e^{-5x} dx) + \int 1 dx) e^{-2x}$$

- Compute integrals

$$y_p(x) = -\frac{4(5x+1)e^{-2x}}{5}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{-2x} + c_2 e^{3x} - \frac{4(5x+1)e^{-2x}}{5}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 20

```
dsolve(diff(y(x),x$2)-diff(y(x),x)-6*y(x)=20*exp(-2*x),y(x), singsol=all)
```

$$y(x) = (c_2 e^{5x} + c_1 - 4x) e^{-2x}$$

✓ Solution by Mathematica

Time used: 0.038 (sec). Leaf size: 32

```
DSolve[y''[x]-y'[x]-6*y[x]==20*Exp[-2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{5} e^{-2x} (-20x + 5c_2 e^{5x} - 4 + 5c_1)$$

10.6 problem 1(f)

10.6.1 Solving as second order linear constant coeff ode	1695
10.6.2 Solving using Kovacic algorithm	1698
10.6.3 Maple step by step solution	1703

Internal problem ID [6307]

Internal file name [OUTPUT/5555_Sunday_June_05_2022_03_43_21_PM_40034600/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.2. THE METHOD OF UN-DETERMINED COEFFICIENTS. Page 67

Problem number: 1(f).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' - 3y' + 2y = 14 \sin(2x) - 18 \cos(2x)$$

10.6.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = -3, C = 2, f(x) = 14 \sin(2x) - 18 \cos(2x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 3y' + 2y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -3, C = 2$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 3\lambda e^{\lambda x} + 2e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 3\lambda + 2 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -3, C = 2$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{3}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-3^2 - (4)(1)(2)} \\ &= \frac{3}{2} \pm \frac{1}{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= \frac{3}{2} + \frac{1}{2} \\ \lambda_2 &= \frac{3}{2} - \frac{1}{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= 2 \\ \lambda_2 &= 1 \end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned} y &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ y &= c_1 e^{(2)x} + c_2 e^{(1)x} \end{aligned}$$

Or

$$y = c_1 e^{2x} + c_2 e^x$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^{2x} + c_2 e^x$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$14 \sin(2x) - 18 \cos(2x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(2x), \sin(2x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^x, e^{2x}\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(2x) + A_2 \sin(2x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-2A_1 \cos(2x) - 2A_2 \sin(2x) + 6A_1 \sin(2x) - 6A_2 \cos(2x) = 14 \sin(2x) - 18 \cos(2x)$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 3, A_2 = 2]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = 3 \cos(2x) + 2 \sin(2x)$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^{2x} + c_2 e^x) + (3 \cos(2x) + 2 \sin(2x)) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{2x} + c_2 e^x + 3 \cos(2x) + 2 \sin(2x) \quad (1)$$




Figure 312: Slope field plot

Verification of solutions

$$y = c_1 e^{2x} + c_2 e^x + 3 \cos(2x) + 2 \sin(2x)$$

Verified OK.

10.6.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 3y' + 2y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -3 \\ C &= 2 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{1}{4} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 1 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \frac{z(x)}{4} \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 251: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0\end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{1}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{x}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-3}{1} dx} \\ &= z_1 e^{\frac{3x}{2}} \\ &= z_1 \left(e^{\frac{3x}{2}} \right)\end{aligned}$$

Which simplifies to

$$y_1 = e^x$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{-3}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{3x}}{(y_1)^2} dx \\ &= y_1(e^x)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1(e^x) + c_2(e^x(e^x))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 3y' + 2y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^x + c_2 e^{2x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$14 \sin(2x) - 18 \cos(2x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(2x), \sin(2x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^x, e^{2x}\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(2x) + A_2 \sin(2x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-2A_1 \cos(2x) - 2A_2 \sin(2x) + 6A_1 \sin(2x) - 6A_2 \cos(2x) = 14 \sin(2x) - 18 \cos(2x)$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 3, A_2 = 2]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = 3 \cos(2x) + 2 \sin(2x)$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^x + c_2 e^{2x}) + (3 \cos(2x) + 2 \sin(2x)) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^x + c_2 e^{2x} + 3 \cos(2x) + 2 \sin(2x) \quad (1)$$




Figure 313: Slope field plot

Verification of solutions

$$y = c_1 e^x + c_2 e^{2x} + 3 \cos(2x) + 2 \sin(2x)$$

Verified OK.

10.6.3 Maple step by step solution

Let's solve

$$y'' - 3y' + 2y = 14 \sin(2x) - 18 \cos(2x)$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - 3r + 2 = 0$$

- Factor the characteristic polynomial

$$(r - 1)(r - 2) = 0$$

- Roots of the characteristic polynomial

$$r = (1, 2)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^x$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{2x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^x + c_2 e^{2x} + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = 14 \sin(2x) - 18 \cos(2x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^x & e^{2x} \\ e^x & 2e^{2x} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = e^{3x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = 2e^x \left(\int (-7 \sin(2x) + 9 \cos(2x)) e^{-x} dx \right) - 2e^{2x} \left(\int (-7 \sin(2x) + 9 \cos(2x)) e^{-2x} dx \right)$$

- Compute integrals

$$y_p(x) = 3 \cos(2x) + 2 \sin(2x)$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^x + c_2 e^{2x} + 3 \cos(2x) + 2 \sin(2x)$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
<- double symmetry of the form [xi=0, eta=F(x)] successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve(diff(y(x),x$2)-3*diff(y(x),x)+2*y(x)=14*sin(2*x)-18*cos(2*x),y(x), singsol=all)
```

$$y(x) = e^{2x} c_1 + e^x c_2 + 2 \sin(2x) + 3 \cos(2x)$$

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 31

```
DSolve[y''[x]-3*y'[x]+2*y[x]==14*Sin[2*x]-18*Cos[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2 \sin(2x) + 3 \cos(2x) + e^x (c_2 e^x + c_1)$$

10.7 problem 1(g)

10.7.1 Solving as second order linear constant coeff ode	1705
10.7.2 Solving using Kovacic algorithm	1708
10.7.3 Maple step by step solution	1713

Internal problem ID [6308]

Internal file name [OUTPUT/5556_Sunday_June_05_2022_03_43_23_PM_88038474/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.2. THE METHOD OF UN-DETERMINED COEFFICIENTS. Page 67

Problem number: 1(g).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + y = 2 \cos(x)$$

10.7.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 1, f(x) = 2 \cos(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\lambda_1 = +i$$

$$\lambda_2 = -i$$

Which simplifies to

$$\lambda_1 = i$$

$$\lambda_2 = -i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (\cos(x) c_1 + c_2 \sin(x))$$

Or

$$y = \cos(x) c_1 + c_2 \sin(x)$$

Therefore the homogeneous solution y_h is

$$y_h = \cos(x) c_1 + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$2 \cos(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since $\cos(x)$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{\cos(x) x, \sin(x) x\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 \cos(x) x + A_2 \sin(x) x$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-2A_1 \sin(x) + 2A_2 \cos(x) = 2 \cos(x)$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 0, A_2 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \sin(x) x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (\cos(x) c_1 + c_2 \sin(x)) + (\sin(x) x) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \cos(x) c_1 + c_2 \sin(x) + \sin(x) x \quad (1)$$




Figure 314: Slope field plot

Verification of solutions

$$y = \cos(x) c_1 + c_2 \sin(x) + \sin(x) x$$

Verified OK.

10.7.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 0 \quad (3)$$

$$C = 1$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 253: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}
 y_1 &= z_1 \\
 &= \cos(x)
 \end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x)) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(x)) + c_2 (\cos(x) (\tan(x))) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = \cos(x) c_1 + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$2 \cos(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since $\cos(x)$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{\cos(x)x, \sin(x)x\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 \cos(x)x + A_2 \sin(x)x$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-2A_1 \sin(x) + 2A_2 \cos(x) = 2 \cos(x)$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 0, A_2 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \sin(x)x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (\cos(x)c_1 + c_2 \sin(x)) + (\sin(x)x) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \cos(x)c_1 + c_2 \sin(x) + \sin(x)x \quad (1)$$




Figure 315: Slope field plot

Verification of solutions

$$y = \cos(x) c_1 + c_2 \sin(x) + \sin(x) x$$

Verified OK.

10.7.3 Maple step by step solution

Let's solve

$$y'' + y = 2 \cos(x)$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (-I, I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = \cos(x) c_1 + c_2 \sin(x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = 2 \cos(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\cos(x) \left(\int \sin(2x) dx \right) + 2 \sin(x) \left(\int \cos(x)^2 dx \right)$$

- Compute integrals

$$y_p(x) = \frac{\cos(x)}{2} + \sin(x) x$$

- Substitute particular solution into general solution to ODE

$$y = \cos(x) c_1 + c_2 \sin(x) + \frac{\cos(x)}{2} + \sin(x) x$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
dsolve(diff(y(x),x$2)+y(x)=2*cos(x),y(x), singsol=all)
```

$$y(x) = (x + c_2) \sin(x) + \cos(x) c_1$$

✓ Solution by Mathematica

Time used: 0.036 (sec). Leaf size: 20

```
DSolve[y''[x]+y[x]==2*Cos[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow (1 + c_1) \cos(x) + (x + c_2) \sin(x)$$

10.8 problem 1(h)

10.8.1 Solving as second order linear constant coeff ode	1716
10.8.2 Solving as second order integrable as is ode	1720
10.8.3 Solving as second order ode missing y ode	1722
10.8.4 Solving as type second_order_integrable_as_is (not using ABC version)	1724
10.8.5 Solving using Kovacic algorithm	1726
10.8.6 Solving as exact linear second order ode ode	1730
10.8.7 Maple step by step solution	1733

Internal problem ID [6309]

Internal file name [OUTPUT/5557_Sunday_June_05_2022_03_43_25_PM_97123339/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.2. THE METHOD OF UN-DETERMINED COEFFICIENTS. Page 67

Problem number: 1(h).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "exact linear second order ode", "second_order_integrable_as_is", "second_order_ode_missing_y", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_y]]
```

$$y'' - 2y' = 12x - 10$$

10.8.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = -2, C = 0, f(x) = 12x - 10$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 2y' = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -2, C = 0$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 2\lambda e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 2\lambda = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -2, C = 0$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{2}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-2^2 - (4)(1)(0)} \\ &= 1 \pm 1 \end{aligned}$$

Hence

$$\lambda_1 = 1 + 1$$

$$\lambda_2 = 1 - 1$$

Which simplifies to

$$\lambda_1 = 2$$

$$\lambda_2 = 0$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(2)x} + c_2 e^{(0)x}$$

Or

$$y = c_1 e^{2x} + c_2$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^{2x} + c_2$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$1 + x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, e^{2x}\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x, x^2\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_2 x^2 + A_1 x$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-4xA_2 - 2A_1 + 2A_2 = 12x - 10$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 2, A_2 = -3]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -3x^2 + 2x$$

Therefore the general solution is

$$\begin{aligned}y &= y_h + y_p \\ &= (c_1 e^{2x} + c_2) + (-3x^2 + 2x)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{2x} + c_2 - 3x^2 + 2x \tag{1}$$




Figure 316: Slope field plot

Verification of solutions

$$y = c_1 e^{2x} + c_2 - 3x^2 + 2x$$

Verified OK.

10.8.2 Solving as second order integrable as is ode

Integrating both sides of the ODE w.r.t x gives

$$\int (y'' - 2y') dx = \int (12x - 10) dx$$
$$y' - 2y = 6x^2 - 10x + c_1$$

Which is now solved for y .

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -2$$
$$q(x) = 6x^2 + c_1 - 10x$$

Hence the ode is

$$y' - 2y = 6x^2 + c_1 - 10x$$

The integrating factor μ is

$$\mu = e^{\int (-2) dx}$$
$$= e^{-2x}$$

The ode becomes

$$\frac{d}{dx}(\mu y) = (\mu)(6x^2 + c_1 - 10x)$$
$$\frac{d}{dx}(y e^{-2x}) = (e^{-2x})(6x^2 + c_1 - 10x)$$
$$d(y e^{-2x}) = ((6x^2 + c_1 - 10x) e^{-2x}) dx$$

Integrating gives

$$y e^{-2x} = \int (6x^2 + c_1 - 10x) e^{-2x} dx$$
$$y e^{-2x} = -\frac{(6x^2 + c_1 - 4x - 2) e^{-2x}}{2} + c_2$$

Dividing both sides by the integrating factor $\mu = e^{-2x}$ results in

$$y = -\frac{e^{2x}(6x^2 + c_1 - 4x - 2) e^{-2x}}{2} + c_2 e^{2x}$$

which simplifies to

$$y = -3x^2 - \frac{c_1}{2} + 2x + 1 + c_2e^{2x}$$

Summary

The solution(s) found are the following

$$y = -3x^2 - \frac{c_1}{2} + 2x + 1 + c_2e^{2x} \tag{1}$$




Figure 317: Slope field plot

Verification of solutions

$$y = -3x^2 - \frac{c_1}{2} + 2x + 1 + c_2e^{2x}$$

Verified OK.

10.8.3 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x) - 2p(x) - 12x + 10 = 0$$

Which is now solve for $p(x)$ as first order ode.

Entering Linear first order ODE solver. In canonical form a linear first order is

$$p'(x) + p(x)p(x) = q(x)$$

Where here

$$\begin{aligned} p(x) &= -2 \\ q(x) &= 12x - 10 \end{aligned}$$

Hence the ode is

$$p'(x) - 2p(x) = 12x - 10$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int (-2) dx} \\ &= e^{-2x} \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx}(\mu p) &= (\mu)(12x - 10) \\ \frac{d}{dx}(e^{-2x} p) &= (e^{-2x})(12x - 10) \\ d(e^{-2x} p) &= ((12x - 10)e^{-2x}) dx \end{aligned}$$

Integrating gives

$$\begin{aligned} e^{-2x} p &= \int (12x - 10) e^{-2x} dx \\ e^{-2x} p &= -2 e^{-2x} (3x - 1) + c_1 \end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{-2x}$ results in

$$p(x) = -2 e^{2x} e^{-2x} (3x - 1) + c_1 e^{2x}$$

which simplifies to

$$p(x) = -6x + 2 + c_1 e^{2x}$$

Since $p = y'$ then the new first order ode to solve is

$$y' = -6x + 2 + c_1 e^{2x}$$

Integrating both sides gives

$$\begin{aligned} y &= \int -6x + 2 + c_1 e^{2x} \, dx \\ &= 2x + \frac{c_1 e^{2x}}{2} - 3x^2 + c_2 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = 2x + \frac{c_1 e^{2x}}{2} - 3x^2 + c_2 \tag{1}$$




Figure 318: Slope field plot

Verification of solutions

$$y = 2x + \frac{c_1 e^{2x}}{2} - 3x^2 + c_2$$

Verified OK.

10.8.4 Solving as type second_order_integrable_as_is (not using ABC version)

Writing the ode as

$$y'' - 2y' = 12x - 10$$

Integrating both sides of the ODE w.r.t x gives

$$\int (y'' - 2y') dx = \int (12x - 10) dx$$
$$y' - 2y = 6x^2 - 10x + c_1$$

Which is now solved for y .

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -2$$
$$q(x) = 6x^2 + c_1 - 10x$$

Hence the ode is

$$y' - 2y = 6x^2 + c_1 - 10x$$

The integrating factor μ is

$$\mu = e^{\int (-2) dx}$$
$$= e^{-2x}$$

The ode becomes

$$\frac{d}{dx}(\mu y) = (\mu) (6x^2 + c_1 - 10x)$$
$$\frac{d}{dx}(y e^{-2x}) = (e^{-2x}) (6x^2 + c_1 - 10x)$$
$$d(y e^{-2x}) = ((6x^2 + c_1 - 10x) e^{-2x}) dx$$

Integrating gives

$$y e^{-2x} = \int (6x^2 + c_1 - 10x) e^{-2x} dx$$
$$y e^{-2x} = -\frac{(6x^2 + c_1 - 4x - 2) e^{-2x}}{2} + c_2$$

Dividing both sides by the integrating factor $\mu = e^{-2x}$ results in

$$y = -\frac{e^{2x}(6x^2 + c_1 - 4x - 2) e^{-2x}}{2} + c_2 e^{2x}$$

which simplifies to

$$y = -3x^2 - \frac{c_1}{2} + 2x + 1 + c_2 e^{2x}$$

Summary

The solution(s) found are the following

$$y = -3x^2 - \frac{c_1}{2} + 2x + 1 + c_2 e^{2x} \quad (1)$$




Figure 319: Slope field plot

Verification of solutions

$$y = -3x^2 - \frac{c_1}{2} + 2x + 1 + c_2 e^{2x}$$

Verified OK.

10.8.5 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 2y' = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -2 \\ C &= 0 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{1}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 255: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-x}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-2}{1} dx} \end{aligned}$$

$$\begin{aligned}
 &= z_1 e^x \\
 &= z_1 (e^x)
 \end{aligned}$$

Which simplifies to

$$y_1 = 1$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}
 y_2 &= y_1 \int \frac{e^{\int -\frac{-2}{1} dx}}{(y_1)^2} dx \\
 &= y_1 \int \frac{e^{2x}}{(y_1)^2} dx \\
 &= y_1 \left(\frac{e^{2x}}{2} \right)
 \end{aligned}$$

Therefore the solution is

$$\begin{aligned}
 y &= c_1 y_1 + c_2 y_2 \\
 &= c_1 (1) + c_2 \left(1 \left(\frac{e^{2x}}{2} \right) \right)
 \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 2y' = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 + \frac{c_2 e^{2x}}{2}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$1 + x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ 1, \frac{e^{2x}}{2} \right\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x, x^2\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_2x^2 + A_1x$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-4xA_2 - 2A_1 + 2A_2 = 12x - 10$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 2, A_2 = -3]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -3x^2 + 2x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 + \frac{c_2 e^{2x}}{2} \right) + (-3x^2 + 2x) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 + \frac{c_2 e^{2x}}{2} - 3x^2 + 2x \quad (1)$$




Figure 320: Slope field plot

Verification of solutions

$$y = c_1 + \frac{c_2 e^{2x}}{2} - 3x^2 + 2x$$

Verified OK.

10.8.6 Solving as exact linear second order ode ode

An ode of the form

$$p(x)y'' + q(x)y' + r(x)y = s(x)$$

is exact if

$$p''(x) - q'(x) + r(x) = 0 \quad (1)$$

For the given ode we have

$$\begin{aligned}p(x) &= 1 \\q(x) &= -2 \\r(x) &= 0 \\s(x) &= 12x - 10\end{aligned}$$

Hence

$$\begin{aligned}p''(x) &= 0 \\q'(x) &= 0\end{aligned}$$

Therefore (1) becomes

$$0 - (0) + (0) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(x)y' + (q(x) - p'(x))y)' = s(x)$$

Integrating gives

$$p(x)y' + (q(x) - p'(x))y = \int s(x) dx$$

Substituting the above values for p, q, r, s gives

$$y' - 2y = \int 12x - 10 dx$$

We now have a first order ode to solve which is

$$y' - 2y = 6x^2 + c_1 - 10x$$

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$\begin{aligned}p(x) &= -2 \\q(x) &= 6x^2 + c_1 - 10x\end{aligned}$$

Hence the ode is

$$y' - 2y = 6x^2 + c_1 - 10x$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int (-2)dx} \\ &= e^{-2x}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu)(6x^2 + c_1 - 10x) \\ \frac{d}{dx}(y e^{-2x}) &= (e^{-2x})(6x^2 + c_1 - 10x) \\ d(y e^{-2x}) &= ((6x^2 + c_1 - 10x) e^{-2x}) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}y e^{-2x} &= \int (6x^2 + c_1 - 10x) e^{-2x} dx \\ y e^{-2x} &= -\frac{(6x^2 + c_1 - 4x - 2) e^{-2x}}{2} + c_2\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{-2x}$ results in

$$y = -\frac{e^{2x}(6x^2 + c_1 - 4x - 2) e^{-2x}}{2} + c_2 e^{2x}$$

which simplifies to

$$y = -3x^2 - \frac{c_1}{2} + 2x + 1 + c_2 e^{2x}$$

Summary

The solution(s) found are the following

$$y = -3x^2 - \frac{c_1}{2} + 2x + 1 + c_2 e^{2x} \quad (1)$$




Figure 321: Slope field plot

Verification of solutions

$$y = -3x^2 - \frac{c_1}{2} + 2x + 1 + c_2e^{2x}$$

Verified OK.

10.8.7 Maple step by step solution

Let's solve

$$y'' - 2y' = 12x - 10$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - 2r = 0$$

- Factor the characteristic polynomial

$$r(r - 2) = 0$$

- Roots of the characteristic polynomial

$$r = (0, 2)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = 1$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{2x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 + c_2 e^{2x} + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x), y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x), y_2(x))} dx \right), f(x) = 12x - 10 \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} 1 & e^{2x} \\ 0 & 2e^{2x} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 2e^{2x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\left(\int (6x - 5) dx \right) + e^{2x} \left(\int (6x - 5) e^{-2x} dx \right)$$

- Compute integrals

$$y_p(x) = -3x^2 + 2x + 1$$

- Substitute particular solution into general solution to ODE

$$y = c_1 + c_2 e^{2x} - 3x^2 + 2x + 1$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
-> Calling odsolve with the ODE`, diff(_b(_a), _a) = 2*_b(_a)+12*_a-10, _b(_a)`  
  Methods for first order ODEs:  
    --- Trying classification methods ---  
    trying a quadrature  
    trying 1st order linear  
    <- 1st order linear successful  
  <- high order exact linear fully integrable successful`
```

*** Sublev

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(diff(y(x),x$2)-2*diff(y(x),x)=12*x-10,y(x), singsol=all)
```

$$y(x) = \frac{e^{2x}c_1}{2} - 3x^2 + 2x + c_2$$

✓ Solution by Mathematica

Time used: 0.103 (sec). Leaf size: 27

```
DSolve[y''[x]-2*y'[x]==12*x-10,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -3x^2 + 2x + \frac{1}{2}c_1e^{2x} + c_2$$

10.9 problem 1(i)

10.9.1 Solving as second order linear constant coeff ode	1736
10.9.2 Solving as linear second order ode solved by an integrating factor ode	1739
10.9.3 Solving using Kovacic algorithm	1741
10.9.4 Maple step by step solution	1746

Internal problem ID [6310]

Internal file name [OUTPUT/5558_Sunday_June_05_2022_03_43_26_PM_74812156/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.2. THE METHOD OF UNDETERMINED COEFFICIENTS. Page 67

Problem number: 1(i).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "linear_second_order_ode_solved_by_an_integrating_factor"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - 2y' + y = 6e^x$$

10.9.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = -2, C = 1, f(x) = 6e^x$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 2y' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -2, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 2\lambda e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 2\lambda + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -2, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{2}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{(-2)^2 - (4)(1)(1)} \\ &= 1 \end{aligned}$$

Hence this is the case of a double root $\lambda_{1,2} = -1$. Therefore the solution is

$$y = c_1 e^x + c_2 x e^x \quad (1)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^x + c_2 x e^x$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$6 e^x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{x e^x, e^x\}$$

Since e^x is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x e^x\}]$$

Since $x e^x$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x^2 e^x\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x^2 e^x$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$2A_1 e^x = 6 e^x$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 3]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = 3x^2 e^x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^x + c_2 x e^x) + (3x^2 e^x) \end{aligned}$$

Which simplifies to

$$y = e^x(c_2 x + c_1) + 3x^2 e^x$$

Summary

The solution(s) found are the following

$$y = e^x(c_2 x + c_1) + 3x^2 e^x \quad (1)$$




Figure 322: Slope field plot

Verification of solutions

$$y = e^x(c_2x + c_1) + 3x^2e^x$$

Verified OK.

10.9.2 Solving as linear second order ode solved by an integrating factor ode

The ode satisfies this form

$$y'' + p(x)y' + \frac{(p(x))^2 + p'(x)}{2}y = f(x)$$

Where $p(x) = -2$. Therefore, there is an integrating factor given by

$$\begin{aligned} M(x) &= e^{\frac{1}{2} \int p dx} \\ &= e^{\int -2 dx} \\ &= e^{-x} \end{aligned}$$

Multiplying both sides of the ODE by the integrating factor $M(x)$ makes the left side of the ODE a complete differential

$$(M(x)y)'' = 6e^{-x}e^x$$

$$(e^{-x}y)'' = 6e^{-x}e^x$$

Integrating once gives

$$(e^{-x}y)' = 6x + c_1$$

Integrating again gives

$$(e^{-x}y) = x(3x + c_1) + c_2$$

Hence the solution is

$$y = \frac{x(3x + c_1) + c_2}{e^{-x}}$$

Or

$$y = c_1x e^x + 3x^2e^x + c_2e^x$$

Summary

The solution(s) found are the following

$$y = c_1x e^x + 3x^2e^x + c_2e^x \tag{1}$$




Figure 323: Slope field plot

Verification of solutions

$$y = c_1 x e^x + 3x^2 e^x + c_2 e^x$$

Verified OK.

10.9.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 2y' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -2 \\ C &= 1 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 0 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 0 \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 257: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-2}{1} dx} \\ &= z_1 e^x \\ &= z_1 (e^x) \end{aligned}$$

Which simplifies to

$$y_1 = e^x$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-2}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{2x}}{(y_1)^2} dx \\ &= y_1(x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^x) + c_2 (e^x(x)) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 2y' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^x + c_2 x e^x$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$6 e^x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[e^x]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{x e^x, e^x\}$$

Since e^x is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[x e^x]$$

Since $x e^x$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[x^2 e^x]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x^2 e^x$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$2A_1 e^x = 6 e^x$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 3]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = 3x^2 e^x$$

Therefore the general solution is

$$\begin{aligned}y &= y_h + y_p \\ &= (c_1 e^x + c_2 x e^x) + (3x^2 e^x)\end{aligned}$$

Which simplifies to

$$y = e^x(c_2 x + c_1) + 3x^2 e^x$$

Summary

The solution(s) found are the following

$$y = e^x(c_2 x + c_1) + 3x^2 e^x \quad (1)$$




Figure 324: Slope field plot

Verification of solutions

$$y = e^x(c_2 x + c_1) + 3x^2 e^x$$

Verified OK.

10.9.4 Maple step by step solution

Let's solve

$$y'' - 2y' + y = 6e^x$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - 2r + 1 = 0$$

- Factor the characteristic polynomial

$$(r - 1)^2 = 0$$

- Root of the characteristic polynomial

$$r = 1$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^x$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = x e^x$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^x + c_2 x e^x + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = 6e^x \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^x & x e^x \\ e^x & x e^x + e^x \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = e^{2x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -6e^x \left(\int x dx - \left(\int 1 dx \right) x \right)$$

- Compute integrals

$$y_p(x) = 3x^2e^x$$

- Substitute particular solution into general solution to ODE

$$y = c_2x e^x + 3x^2e^x + c_1e^x$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)-2*diff(y(x),x)+y(x)=6*exp(x),y(x), singsol=all)
```

$$y(x) = e^x(c_1x + 3x^2 + c_2)$$

✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 21

```
DSolve[y''[x]-2*y'[x]+y[x]==6*Exp[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x(3x^2 + c_2x + c_1)$$

10.10 problem 1(j)

10.10.1 Solving as second order linear constant coeff ode	1748
10.10.2 Solving using Kovacic algorithm	1751
10.10.3 Maple step by step solution	1756

Internal problem ID [6311]

Internal file name [OUTPUT/5559_Sunday_June_05_2022_03_43_28_PM_75454561/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.2. THE METHOD OF UN-DETERMINED COEFFICIENTS. Page 67

Problem number: 1(j).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' - 2y' + 2y = \sin(x) e^x$$

10.10.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = -2, C = 2, f(x) = \sin(x) e^x$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 2y' + 2y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -2, C = 2$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 2\lambda e^{\lambda x} + 2 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 2\lambda + 2 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -2, C = 2$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{2}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-2^2 - (4)(1)(2)} \\ &= 1 \pm i \end{aligned}$$

Hence

$$\lambda_1 = 1 + i$$

$$\lambda_2 = 1 - i$$

Which simplifies to

$$\lambda_1 = 1 + i$$

$$\lambda_2 = 1 - i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 1$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^x (\cos(x) c_1 + c_2 \sin(x))$$

Therefore the homogeneous solution y_h is

$$y_h = e^x (\cos(x) c_1 + c_2 \sin(x))$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x) e^x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x) e^x, \sin(x) e^x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x) e^x, \sin(x) e^x\}$$

Since $\cos(x) e^x$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{\cos(x) e^x x, \sin(x) e^x x\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 \cos(x) e^x x + A_2 \sin(x) e^x x$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-2A_1 \sin(x) e^x + 2A_2 \cos(x) e^x = \sin(x) e^x$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{2}, A_2 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{\cos(x) e^x x}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (e^x (\cos(x) c_1 + c_2 \sin(x))) + \left(-\frac{\cos(x) e^x x}{2} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = e^x (\cos(x) c_1 + c_2 \sin(x)) - \frac{\cos(x) e^x x}{2} \quad (1)$$




Figure 325: Slope field plot

Verification of solutions

$$y = e^x (\cos(x) c_1 + c_2 \sin(x)) - \frac{\cos(x) e^x x}{2}$$

Verified OK.

10.10.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 2y' + 2y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= 1 \\B &= -2 \\C &= 2\end{aligned}\tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x)\tag{4}$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\&= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned}\tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1}\tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= -1 \\t &= 1\end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x)\tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 259: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}
 y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\
 &= z_1 e^{-\int \frac{1}{2} \frac{-2}{1} dx} \\
 &= z_1 e^x \\
 &= z_1(e^x)
 \end{aligned}$$

Which simplifies to

$$y_1 = \cos(x) e^x$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-2}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{2x}}{(y_1)^2} dx \\ &= y_1(\tan(x)) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1(\cos(x) e^x) + c_2(\cos(x) e^x(\tan(x))) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 2y' + 2y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 \cos(x) e^x + c_2 \sin(x) e^x$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x) e^x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x) e^x, \sin(x) e^x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x) e^x, \sin(x) e^x\}$$

Since $\cos(x) e^x$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{\cos(x) e^x x, \sin(x) e^x x\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 \cos(x) e^x x + A_2 \sin(x) e^x x$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-2A_1 \sin(x) e^x + 2A_2 \cos(x) e^x = \sin(x) e^x$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{2}, A_2 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{\cos(x) e^x x}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(x) e^x + c_2 \sin(x) e^x) + \left(-\frac{\cos(x) e^x x}{2} \right) \end{aligned}$$

Which simplifies to

$$y = e^x (\cos(x) c_1 + c_2 \sin(x)) - \frac{\cos(x) e^x x}{2}$$

Summary

The solution(s) found are the following

$$y = e^x (\cos(x) c_1 + c_2 \sin(x)) - \frac{\cos(x) e^x x}{2} \quad (1)$$




Figure 326: Slope field plot

Verification of solutions

$$y = e^x (\cos(x) c_1 + c_2 \sin(x)) - \frac{\cos(x) e^x x}{2}$$

Verified OK.

10.10.3 Maple step by step solution

Let's solve

$$y'' - 2y' + 2y = \sin(x) e^x$$

- Highest derivative means the order of the ODE is 2
- y''
- Characteristic polynomial of homogeneous ODE

$$r^2 - 2r + 2 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{2 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (1 - I, 1 + I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x) e^x$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(x) e^x$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 \cos(x) e^x + c_2 \sin(x) e^x + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \sin(x) e^x \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) e^x & \sin(x) e^x \\ -\sin(x) e^x + \cos(x) e^x & \cos(x) e^x + \sin(x) e^x \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = e^{2x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = \frac{e^x \left(\sin(x) \left(\int \sin(2x) dx \right) - 2 \cos(x) \left(\int \sin(x)^2 dx \right) \right)}{2}$$

- Compute integrals

$$y_p(x) = \frac{(\sin(x) - 2 \cos(x)x) e^x}{4}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 \cos(x) e^x + c_2 \sin(x) e^x + \frac{(\sin(x) - 2 \cos(x)x) e^x}{4}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve(diff(y(x),x$2)-2*diff(y(x),x)+2*y(x)=exp(x)*sin(x),y(x), singsol=all)
```

$$y(x) = -\frac{((-2c_1 + x) \cos(x) + (-2c_2 - 1) \sin(x)) e^x}{2}$$

✓ Solution by Mathematica

Time used: 0.052 (sec). Leaf size: 28

```
DSolve[y''[x]-2*y'[x]+2*y[x]==Exp[x]*Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{2}e^x((x - 2c_2) \cos(x) - 2c_1 \sin(x))$$

10.11 problem 1(k)

10.11.1 Solving as second order linear constant coeff ode	1759
10.11.2 Solving as second order integrable as is ode	1763
10.11.3 Solving as second order ode missing y ode	1765
10.11.4 Solving as type second_order_integrable_as_is (not using ABC version)	1767
10.11.5 Solving using Kovacic algorithm	1769
10.11.6 Solving as exact linear second order ode ode	1773
10.11.7 Maple step by step solution	1776

Internal problem ID [6312]

Internal file name [OUTPUT/5560_Sunday_June_05_2022_03_43_29_PM_43068580/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.2. THE METHOD OF UNDETERMINED COEFFICIENTS. Page 67

Problem number: 1(k).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "exact linear second order ode", "second_order_integrable_as_is", "second_order_ode_missing_y", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_y]]
```

$$y'' + y' = 10x^4 + 2$$

10.11.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 1, C = 0, f(x) = 10x^4 + 2$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 1, C = 0$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + \lambda e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + \lambda = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 1, C = 0$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{1^2 - (4)(1)(0)} \\ &= -\frac{1}{2} \pm \frac{1}{2} \end{aligned}$$

Hence

$$\lambda_1 = -\frac{1}{2} + \frac{1}{2}$$

$$\lambda_2 = -\frac{1}{2} - \frac{1}{2}$$

Which simplifies to

$$\lambda_1 = 0$$

$$\lambda_2 = -1$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(0)x} + c_2 e^{(-1)x}$$

Or

$$y = c_1 + c_2 e^{-x}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 + c_2 e^{-x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x^4 + 1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x, x^2, x^3, x^4\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, e^{-x}\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x, x^2, x^3, x^4, x^5\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_5 x^5 + A_4 x^4 + A_3 x^3 + A_2 x^2 + A_1 x$$

The unknowns $\{A_1, A_2, A_3, A_4, A_5\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$5x^4 A_5 + 4x^3 A_4 + 20x^3 A_5 + 3x^2 A_3 + 12x^2 A_4 + 2x A_2 + 6x A_3 + A_1 + 2A_2 = 10x^4 + 2$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 242, A_2 = -120, A_3 = 40, A_4 = -10, A_5 = 2]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = 2x^5 - 10x^4 + 40x^3 - 120x^2 + 242x$$

Therefore the general solution is

$$\begin{aligned}y &= y_h + y_p \\ &= (c_1 + c_2 e^{-x}) + (2x^5 - 10x^4 + 40x^3 - 120x^2 + 242x)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 + c_2 e^{-x} + 2x^5 - 10x^4 + 40x^3 - 120x^2 + 242x \quad (1)$$




Figure 327: Slope field plot

Verification of solutions

$$y = c_1 + c_2 e^{-x} + 2x^5 - 10x^4 + 40x^3 - 120x^2 + 242x$$

Verified OK.

10.11.2 Solving as second order integrable as is ode

Integrating both sides of the ODE w.r.t x gives

$$\int (y'' + y') dx = \int (10x^4 + 2) dx$$
$$y + y' = 2x^5 + 2x + c_1$$

Which is now solved for y .

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = 1$$
$$q(x) = 2x^5 + c_1 + 2x$$

Hence the ode is

$$y + y' = 2x^5 + c_1 + 2x$$

The integrating factor μ is

$$\mu = e^{\int 1 dx}$$
$$= e^x$$

The ode becomes

$$\frac{d}{dx}(\mu y) = (\mu) (2x^5 + c_1 + 2x)$$
$$\frac{d}{dx}(y e^x) = (e^x) (2x^5 + c_1 + 2x)$$
$$d(y e^x) = ((2x^5 + c_1 + 2x) e^x) dx$$

Integrating gives

$$y e^x = \int (2x^5 + c_1 + 2x) e^x dx$$
$$y e^x = (2x^5 - 10x^4 + 40x^3 - 120x^2 + c_1 + 242x - 242) e^x + c_2$$

Dividing both sides by the integrating factor $\mu = e^x$ results in

$$y = e^{-x} (2x^5 - 10x^4 + 40x^3 - 120x^2 + c_1 + 242x - 242) e^x + c_2 e^{-x}$$

which simplifies to

$$y = 2x^5 - 10x^4 + 40x^3 - 120x^2 + c_1 + 242x - 242 + c_2e^{-x}$$

Summary

The solution(s) found are the following

$$y = 2x^5 - 10x^4 + 40x^3 - 120x^2 + c_1 + 242x - 242 + c_2e^{-x} \quad (1)$$




Figure 328: Slope field plot

Verification of solutions

$$y = 2x^5 - 10x^4 + 40x^3 - 120x^2 + c_1 + 242x - 242 + c_2e^{-x}$$

Verified OK.

10.11.3 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p(x) + p'(x) - 10x^4 - 2 = 0$$

Which is now solve for $p(x)$ as first order ode.

Entering Linear first order ODE solver. In canonical form a linear first order is

$$p'(x) + p(x) = 10x^4 + 2$$

Where here

$$p(x) = 1$$

$$q(x) = 10x^4 + 2$$

Hence the ode is

$$p(x) + p'(x) = 10x^4 + 2$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int 1 dx} \\ &= e^x\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu p) &= (\mu) (10x^4 + 2) \\ \frac{d}{dx}(e^x p) &= (e^x) (10x^4 + 2) \\ d(e^x p) &= ((10x^4 + 2) e^x) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}e^x p &= \int (10x^4 + 2) e^x dx \\ e^x p &= 2 e^x (5x^4 - 20x^3 + 60x^2 - 120x + 121) + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^x$ results in

$$p(x) = 2 e^{-x} e^x (5x^4 - 20x^3 + 60x^2 - 120x + 121) + c_1 e^{-x}$$

which simplifies to

$$p(x) = 10x^4 - 40x^3 + 120x^2 - 240x + 242 + c_1 e^{-x}$$

Since $p = y'$ then the new first order ode to solve is

$$y' = 10x^4 - 40x^3 + 120x^2 - 240x + 242 + c_1 e^{-x}$$

Integrating both sides gives

$$\begin{aligned} y &= \int 10x^4 - 40x^3 + 120x^2 - 240x + 242 + c_1 e^{-x} dx \\ &= 242x - c_1 e^{-x} - 120x^2 + 40x^3 - 10x^4 + 2x^5 + c_2 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = 242x - c_1 e^{-x} - 120x^2 + 40x^3 - 10x^4 + 2x^5 + c_2 \quad (1)$$




Figure 329: Slope field plot

Verification of solutions

$$y = 242x - c_1 e^{-x} - 120x^2 + 40x^3 - 10x^4 + 2x^5 + c_2$$

Verified OK.

10.11.4 Solving as type second_order_integrable_as_is (not using ABC version)

Writing the ode as

$$y'' + y' = 10x^4 + 2$$

Integrating both sides of the ODE w.r.t x gives

$$\int (y'' + y') dx = \int (10x^4 + 2) dx$$
$$y + y' = 2x^5 + 2x + c_1$$

Which is now solved for y .

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = 1$$
$$q(x) = 2x^5 + c_1 + 2x$$

Hence the ode is

$$y + y' = 2x^5 + c_1 + 2x$$

The integrating factor μ is

$$\mu = e^{\int 1 dx}$$
$$= e^x$$

The ode becomes

$$\frac{d}{dx}(\mu y) = (\mu) (2x^5 + c_1 + 2x)$$
$$\frac{d}{dx}(y e^x) = (e^x) (2x^5 + c_1 + 2x)$$
$$d(y e^x) = ((2x^5 + c_1 + 2x) e^x) dx$$

Integrating gives

$$y e^x = \int (2x^5 + c_1 + 2x) e^x dx$$
$$y e^x = (2x^5 - 10x^4 + 40x^3 - 120x^2 + c_1 + 242x - 242) e^x + c_2$$

Dividing both sides by the integrating factor $\mu = e^x$ results in

$$y = e^{-x} (2x^5 - 10x^4 + 40x^3 - 120x^2 + c_1 + 242x - 242) e^x + c_2 e^{-x}$$

which simplifies to

$$y = 2x^5 - 10x^4 + 40x^3 - 120x^2 + c_1 + 242x - 242 + c_2 e^{-x}$$

Summary

The solution(s) found are the following

$$y = 2x^5 - 10x^4 + 40x^3 - 120x^2 + c_1 + 242x - 242 + c_2 e^{-x} \quad (1)$$




Figure 330: Slope field plot

Verification of solutions

$$y = 2x^5 - 10x^4 + 40x^3 - 120x^2 + c_1 + 242x - 242 + c_2 e^{-x}$$

Verified OK.

10.11.5 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y' = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 1 \quad (3)$$

$$C = 0$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$r = \frac{s}{t} \quad (5)$$
$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{1}{4} \quad (6)$$

Comparing the above to (5) shows that

$$s = 1$$

$$t = 4$$

Therefore eq. (4) becomes

$$z''(x) = \frac{z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 261: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{1}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{x}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{1}{1} dx} \end{aligned}$$

$$\begin{aligned}
&= z_1 e^{-\frac{x}{2}} \\
&= z_1 (e^{-\frac{x}{2}})
\end{aligned}$$

Which simplifies to

$$y_1 = e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}
y_2 &= y_1 \int \frac{e^{\int -\frac{1}{1} dx}}{(y_1)^2} dx \\
&= y_1 \int \frac{e^{-x}}{(y_1)^2} dx \\
&= y_1 (e^x)
\end{aligned}$$

Therefore the solution is

$$\begin{aligned}
y &= c_1 y_1 + c_2 y_2 \\
&= c_1 (e^{-x}) + c_2 (e^{-x}(e^x))
\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-x} + c_2$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x^4 + 1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x, x^2, x^3, x^4\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, e^{-x}\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x, x^2, x^3, x^4, x^5\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_5x^5 + A_4x^4 + A_3x^3 + A_2x^2 + A_1x$$

The unknowns $\{A_1, A_2, A_3, A_4, A_5\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$5x^4A_5 + 4x^3A_4 + 20x^3A_5 + 3x^2A_3 + 12x^2A_4 + 2xA_2 + 6xA_3 + A_1 + 2A_2 = 10x^4 + 2$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 242, A_2 = -120, A_3 = 40, A_4 = -10, A_5 = 2]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = 2x^5 - 10x^4 + 40x^3 - 120x^2 + 242x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1e^{-x} + c_2) + (2x^5 - 10x^4 + 40x^3 - 120x^2 + 242x) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-x} + c_2 + 2x^5 - 10x^4 + 40x^3 - 120x^2 + 242x \quad (1)$$




Figure 331: Slope field plot

Verification of solutions

$$y = c_1 e^{-x} + c_2 + 2x^5 - 10x^4 + 40x^3 - 120x^2 + 242x$$

Verified OK.

10.11.6 Solving as exact linear second order ode ode

An ode of the form

$$p(x)y'' + q(x)y' + r(x)y = s(x)$$

is exact if

$$p''(x) - q'(x) + r(x) = 0 \quad (1)$$

For the given ode we have

$$\begin{aligned}p(x) &= 1 \\q(x) &= 1 \\r(x) &= 0 \\s(x) &= 10x^4 + 2\end{aligned}$$

Hence

$$\begin{aligned}p''(x) &= 0 \\q'(x) &= 0\end{aligned}$$

Therefore (1) becomes

$$0 - (0) + (0) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(x)y' + (q(x) - p'(x))y)' = s(x)$$

Integrating gives

$$p(x)y' + (q(x) - p'(x))y = \int s(x) dx$$

Substituting the above values for p, q, r, s gives

$$y + y' = \int 10x^4 + 2 dx$$

We now have a first order ode to solve which is

$$y + y' = 2x^5 + c_1 + 2x$$

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$\begin{aligned}p(x) &= 1 \\q(x) &= 2x^5 + c_1 + 2x\end{aligned}$$

Hence the ode is

$$y + y' = 2x^5 + c_1 + 2x$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int 1 dx} \\ &= e^x\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu)(2x^5 + c_1 + 2x) \\ \frac{d}{dx}(y e^x) &= (e^x)(2x^5 + c_1 + 2x) \\ d(y e^x) &= ((2x^5 + c_1 + 2x) e^x) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}y e^x &= \int (2x^5 + c_1 + 2x) e^x dx \\ y e^x &= (2x^5 - 10x^4 + 40x^3 - 120x^2 + c_1 + 242x - 242) e^x + c_2\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^x$ results in

$$y = e^{-x} (2x^5 - 10x^4 + 40x^3 - 120x^2 + c_1 + 242x - 242) e^x + c_2 e^{-x}$$

which simplifies to

$$y = 2x^5 - 10x^4 + 40x^3 - 120x^2 + c_1 + 242x - 242 + c_2 e^{-x}$$

Summary

The solution(s) found are the following

$$y = 2x^5 - 10x^4 + 40x^3 - 120x^2 + c_1 + 242x - 242 + c_2 e^{-x} \quad (1)$$




Figure 332: Slope field plot

Verification of solutions

$$y = 2x^5 - 10x^4 + 40x^3 - 120x^2 + c_1 + 242x - 242 + c_2e^{-x}$$

Verified OK.

10.11.7 Maple step by step solution

Let's solve

$$y' + y'' = 10x^4 + 2$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + r = 0$$

- Factor the characteristic polynomial

$$r(r + 1) = 0$$

- Roots of the characteristic polynomial

$$r = (-1, 0)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = 1$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{-x} + c_2 + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = 10x^4 + 2 \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-x} & 1 \\ -e^{-x} & 0 \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = e^{-x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -2e^{-x} \left(\int (5x^4 + 1) e^x dx \right) + 2 \left(\int (5x^4 + 1) dx \right)$$

- Compute integrals

$$y_p(x) = 2x^5 - 10x^4 + 40x^3 - 120x^2 + 242x - 242$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{-x} + c_2 + 2x^5 - 10x^4 + 40x^3 - 120x^2 + 242x - 242$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
-> Calling odsolve with the ODE`, diff(_b(_a), _a) = 10*_a^4-_b(_a)+2, _b(_a)  
Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful  
<- high order exact linear fully integrable successful`
```

*** Subleve

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 36

```
dsolve(diff(y(x),x$2)+diff(y(x),x)=10*x^4+2,y(x), singsol=all)
```

$$y(x) = -c_1 e^{-x} - 120x^2 + 40x^3 - 10x^4 + 2x^5 + 242x + c_2$$

✓ Solution by Mathematica

Time used: 0.106 (sec). Leaf size: 40

```
DSolve[y''[x]+y'[x]==10*x^4+2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2x^5 - 10x^4 + 40x^3 - 120x^2 + 242x - c_1 e^{-x} + c_2$$

10.12 problem 3(a)

10.12.1 Solving as second order linear constant coeff ode	1779
10.12.2 Solving using Kovacic algorithm	1783
10.12.3 Maple step by step solution	1788

Internal problem ID [6313]

Internal file name [OUTPUT/5561_Sunday_June_05_2022_03_43_31_PM_45233317/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.2. THE METHOD OF UNDETERMINED COEFFICIENTS. Page 67

Problem number: 3(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + 4y = 4 \cos(2x) + 6 \cos(x) + 8x^2 - 4x$$

10.12.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 4, f(x) = 4 \cos(2x) + 6 \cos(x) + 8x^2 - 4x$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 4y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 4$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 4 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 4 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 4$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(4)} \\ &= \pm 2i \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +2i \\ \lambda_2 &= -2i \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= 2i \\ \lambda_2 &= -2i \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 2$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (c_1 \cos(2x) + c_2 \sin(2x))$$

Or

$$y = c_1 \cos(2x) + c_2 \sin(2x)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \cos(2x) + c_2 \sin(2x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$4 \cos(2x) + 6 \cos(x) + 8x^2 - 4x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}, \{\cos(2x), \sin(2x)\}, \{1, x, x^2\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(2x), \sin(2x)\}$$

Since $\cos(2x)$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{\cos(x), \sin(x)\}, \{x \cos(2x), x \sin(2x)\}, \{1, x, x^2\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 \cos(x) + A_2 \sin(x) + A_3 x \cos(2x) + A_4 x \sin(2x) + A_5 + A_6 x + A_7 x^2$$

The unknowns $\{A_1, A_2, A_3, A_4, A_5, A_6, A_7\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$\begin{aligned} 3A_1 \cos(x) + 3A_2 \sin(x) - 4A_3 \sin(2x) + 4A_4 \cos(2x) + 2A_7 + 4A_5 + 4A_6 x + 4A_7 x^2 \\ = 4 \cos(2x) + 6 \cos(x) + 8x^2 - 4x \end{aligned}$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 2, A_2 = 0, A_3 = 0, A_4 = 1, A_5 = -1, A_6 = -1, A_7 = 2]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = 2 \cos(x) - 1 + x \sin(2x) - x + 2x^2$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(2x) + c_2 \sin(2x)) + (2 \cos(x) - 1 + x \sin(2x) - x + 2x^2) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(2x) + c_2 \sin(2x) + 2 \cos(x) - 1 + x \sin(2x) - x + 2x^2 \quad (1)$$




Figure 333: Slope field plot

Verification of solutions

$$y = c_1 \cos(2x) + c_2 \sin(2x) + 2 \cos(x) - 1 + x \sin(2x) - x + 2x^2$$

Verified OK.

10.12.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 4y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 4 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-4}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -4 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -4z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 263: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -4$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(2x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}y_1 &= z_1 \\ &= \cos(2x)\end{aligned}$$

Which simplifies to

$$y_1 = \cos(2x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned}y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(2x) \int \frac{1}{\cos(2x)^2} dx \\ &= \cos(2x) \left(\frac{\tan(2x)}{2} \right)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1(\cos(2x)) + c_2 \left(\cos(2x) \left(\frac{\tan(2x)}{2} \right) \right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$.
 y_h is the solution to

$$y'' + 4y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 \cos(2x) + \frac{c_2 \sin(2x)}{2}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$8 \cos(x)^2 - 4 + 6 \cos(x) + 8x^2 - 4x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}, \{\cos(2x), \sin(2x)\}, \{1, x, x^2\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ \frac{\sin(2x)}{2}, \cos(2x) \right\}$$

Since $\cos(2x)$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{\cos(x), \sin(x)\}, \{x \cos(2x), x \sin(2x)\}, \{1, x, x^2\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 \cos(x) + A_2 \sin(x) + A_3 x \cos(2x) + A_4 x \sin(2x) + A_5 + A_6 x + A_7 x^2$$

The unknowns $\{A_1, A_2, A_3, A_4, A_5, A_6, A_7\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$\begin{aligned} 3A_1 \cos(x) + 3A_2 \sin(x) - 4A_3 \sin(2x) + 4A_4 \cos(2x) + 2A_7 + 4A_5 + 4A_6 x + 4A_7 x^2 \\ = 4 \cos(2x) + 6 \cos(x) + 8x^2 - 4x \end{aligned}$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 2, A_2 = 0, A_3 = 0, A_4 = 1, A_5 = -1, A_6 = -1, A_7 = 2]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = 2 \cos(x) - 1 + x \sin(2x) - x + 2x^2$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 \cos(2x) + \frac{c_2 \sin(2x)}{2} \right) + (2 \cos(x) - 1 + x \sin(2x) - x + 2x^2) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(2x) + \frac{c_2 \sin(2x)}{2} + 2 \cos(x) - 1 + x \sin(2x) - x + 2x^2 \quad (1)$$




Figure 334: Slope field plot

Verification of solutions

$$y = c_1 \cos(2x) + \frac{c_2 \sin(2x)}{2} + 2 \cos(x) - 1 + x \sin(2x) - x + 2x^2$$

Verified OK.

10.12.3 Maple step by step solution

Let's solve

$$y'' + 4y = 4 \cos(2x) + 6 \cos(x) + 8x^2 - 4x$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 4 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-16})}{2}$$

- Roots of the characteristic polynomial

$$r = (-2i, 2i)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(2x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(2x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 \cos(2x) + c_2 \sin(2x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right) \right], f(x) = 4 \cos(2x) + 6 \cos(x) +$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(2x) & \sin(2x) \\ -2 \sin(2x) & 2 \cos(2x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 2$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\cos(2x) \left(\int \sin(2x) (2 \cos(2x) + 3 \cos(x) + 4x^2 - 2x) dx \right) + \sin(2x) \left(\int \cos(2x) (-2 + \dots) dx \right)$$

- Compute integrals

$$y_p(x) = 2 \cos(x) \sin(x) x + \frac{\cos(x)^2}{2} + 2x^2 + 2 \cos(x) - x - \frac{5}{4}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 \cos(2x) + c_2 \sin(2x) + 2 \cos(x) \sin(x) x + \frac{\cos(x)^2}{2} + 2x^2 + 2 \cos(x) - x - \frac{5}{4}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 37

```
dsolve(diff(y(x), x$2)+4*y(x)=4*cos(2*x)+6*cos(x)+8*x^2-4*x,y(x), singsol=all)
```

$$y(x) = -1 + \frac{(4c_1 + 1) \cos(2x)}{4} + (x + c_2) \sin(2x) + 2x^2 - x + 2 \cos(x)$$

✓ Solution by Mathematica

Time used: 0.515 (sec). Leaf size: 43

```
DSolve[y''[x]+4*y[x]==4*Cos[2*x]+6*Cos[x]+8*x^2-4*x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2x^2 - x + x \sin(2x) + 2 \cos(x) + \left(\frac{1}{2} + c_1 \right) \cos(2x) + c_2 \sin(2x) - 1$$

10.13 problem 3(b)

10.13.1 Solving as second order linear constant coeff ode	1790
10.13.2 Solving using Kovacic algorithm	1794
10.13.3 Maple step by step solution	1799

Internal problem ID [6314]

Internal file name [OUTPUT/5562_Sunday_June_05_2022_03_43_33_PM_29618251/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.2. THE METHOD OF UNDETERMINED COEFFICIENTS. Page 67

Problem number: 3(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + 9y = 2 \sin(3x) + 4 \sin(x) - 26e^{-2x} + 27x^3$$

10.13.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 9, f(x) = 2 \sin(3x) + 4 \sin(x) - 26e^{-2x} + 27x^3$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 9y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 9$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 9 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 9 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 9$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(9)} \\ &= \pm 3i \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +3i \\ \lambda_2 &= -3i \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= 3i \\ \lambda_2 &= -3i \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 3$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (c_1 \cos(3x) + c_2 \sin(3x))$$

Or

$$y = c_1 \cos(3x) + c_2 \sin(3x)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \cos(3x) + c_2 \sin(3x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$2 \sin(3x) + 4 \sin(x) - 26 e^{-2x} + 27x^3$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$\{e^{-2x}, \{\cos(x), \sin(x)\}, \{\cos(3x), \sin(3x)\}, \{1, x, x^2, x^3\}\}$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(3x), \sin(3x)\}$$

Since $\cos(3x)$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$\{e^{-2x}, \{\cos(x), \sin(x)\}, \{x \cos(3x), x \sin(3x)\}, \{1, x, x^2, x^3\}\}$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 e^{-2x} + A_2 \cos(x) + A_3 \sin(x) + A_4 x \cos(3x) + A_5 x \sin(3x) + A_6 + A_7 x + A_8 x^2 + A_9 x^3$$

The unknowns $\{A_1, A_2, A_3, A_4, A_5, A_6, A_7, A_8, A_9\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$13A_1 e^{-2x} + 8A_2 \cos(x) + 8A_3 \sin(x) - 6A_4 \sin(3x) + 6A_5 \cos(3x) + 2A_8 + 6A_9 x + 9A_6 + 9A_7 x + 9A_8 x^2 + 9A_9 x^3 = 2 \sin(3x) + 4 \sin(x) - 26 e^{-2x} + 27x^3$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -2, A_2 = 0, A_3 = \frac{1}{2}, A_4 = -\frac{1}{3}, A_5 = 0, A_6 = 0, A_7 = -2, A_8 = 0, A_9 = 3 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -2e^{-2x} + \frac{\sin(x)}{2} - \frac{x \cos(3x)}{3} - 2x + 3x^3$$

Therefore the general solution is

$$y = y_h + y_p$$

$$= (c_1 \cos(3x) + c_2 \sin(3x)) + \left(-2e^{-2x} + \frac{\sin(x)}{2} - \frac{x \cos(3x)}{3} - 2x + 3x^3 \right)$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(3x) + c_2 \sin(3x) - 2e^{-2x} + \frac{\sin(x)}{2} - \frac{x \cos(3x)}{3} - 2x + 3x^3 \quad (1)$$




Figure 335: Slope field plot

Verification of solutions

$$y = c_1 \cos(3x) + c_2 \sin(3x) - 2e^{-2x} + \frac{\sin(x)}{2} - \frac{x \cos(3x)}{3} - 2x + 3x^3$$

Verified OK.

10.13.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 9y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 9 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-9}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -9 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -9z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 265: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -9$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(3x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}y_1 &= z_1 \\ &= \cos(3x)\end{aligned}$$

Which simplifies to

$$y_1 = \cos(3x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned}y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(3x) \int \frac{1}{\cos(3x)^2} dx \\ &= \cos(3x) \left(\frac{\tan(3x)}{3} \right)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1(\cos(3x)) + c_2 \left(\cos(3x) \left(\frac{\tan(3x)}{3} \right) \right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$.
 y_h is the solution to

$$y'' + 9y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 \cos(3x) + \frac{c_2 \sin(3x)}{3}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$2 \sin(3x) + 4 \sin(x) - 26 e^{-2x} + 27x^3$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{-2x}\}, \{\cos(x), \sin(x)\}, \{\cos(3x), \sin(3x)\}, \{1, x, x^2, x^3\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ \frac{\sin(3x)}{3}, \cos(3x) \right\}$$

Since $\cos(3x)$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{e^{-2x}\}, \{\cos(x), \sin(x)\}, \{x \cos(3x), x \sin(3x)\}, \{1, x, x^2, x^3\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 e^{-2x} + A_2 \cos(x) + A_3 \sin(x) + A_4 x \cos(3x) + A_5 x \sin(3x) + A_6 + A_7 x + A_8 x^2 + A_9 x^3$$

The unknowns $\{A_1, A_2, A_3, A_4, A_5, A_6, A_7, A_8, A_9\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$13A_1 e^{-2x} + 8A_2 \cos(x) + 8A_3 \sin(x) - 6A_4 \sin(3x) + 6A_5 \cos(3x) + 2A_8 + 6A_9 x + 9A_6 + 9A_7 x + 9A_8 x^2 + 9A_9 x^3 = 2 \sin(3x) + 4 \sin(x) - 26 e^{-2x} + 27x^3$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -2, A_2 = 0, A_3 = \frac{1}{2}, A_4 = -\frac{1}{3}, A_5 = 0, A_6 = 0, A_7 = -2, A_8 = 0, A_9 = 3 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -2e^{-2x} + \frac{\sin(x)}{2} - \frac{x \cos(3x)}{3} - 2x + 3x^3$$

Therefore the general solution is

$$y = y_h + y_p$$

$$= \left(c_1 \cos(3x) + \frac{c_2 \sin(3x)}{3} \right) + \left(-2e^{-2x} + \frac{\sin(x)}{2} - \frac{x \cos(3x)}{3} - 2x + 3x^3 \right)$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(3x) + \frac{c_2 \sin(3x)}{3} - 2e^{-2x} + \frac{\sin(x)}{2} - \frac{x \cos(3x)}{3} - 2x + 3x^3 \quad (1)$$




Figure 336: Slope field plot

Verification of solutions

$$y = c_1 \cos(3x) + \frac{c_2 \sin(3x)}{3} - 2e^{-2x} + \frac{\sin(x)}{2} - \frac{x \cos(3x)}{3} - 2x + 3x^3$$

Verified OK.

10.13.3 Maple step by step solution

Let's solve

$$y'' + 9y = 2 \sin(3x) + 4 \sin(x) - 26e^{-2x} + 27x^3$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 9 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-36})}{2}$$

- Roots of the characteristic polynomial

$$r = (-3I, 3I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(3x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(3x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 \cos(3x) + c_2 \sin(3x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right) \right], f(x) = 2 \sin(3x) + 4 \sin(x) -$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(3x) & \sin(3x) \\ -3 \sin(3x) & 3 \cos(3x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 3$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = \frac{\cos(3x) \left(\int \sin(3x) (-2 \sin(3x) - 4 \sin(x) + 26 e^{-2x} - 27x^3) dx \right)}{3} - \frac{\sin(3x) \left(\int \cos(3x) (-2 \sin(3x) - 4 \sin(x) + 26 e^{-2x} - 27x^3) dx \right)}{3}$$

- Compute integrals

$$y_p(x) = 2 \cos(x)^2 \sin(x) - \frac{4 \cos(x)^3 x}{3} + 3x^3 + \cos(x) x - 2 e^{-2x} - 2x$$

- Substitute particular solution into general solution to ODE

$$y = c_1 \cos(3x) + c_2 \sin(3x) + 2 \cos(x)^2 \sin(x) - \frac{4 \cos(x)^3 x}{3} + 3x^3 + \cos(x) x - 2 e^{-2x} - 2x$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 47

```
dsolve(diff(y(x), x$2)+9*y(x)=2*sin(3*x)+4*sin(x)-26*exp(-2*x)+27*x^3,y(x), singsol=all)
```

$$y(x) = \frac{(-2x + 6c_1) \cos(3x)}{6} + \frac{(6c_2 + 3) \sin(3x)}{6} + 3x^3 - 2x + \frac{\sin(x)}{2} - 2e^{-2x}$$

✓ Solution by Mathematica

Time used: 2.215 (sec). Leaf size: 55

```
DSolve[y''[x]+9*y[x]==2*Sin[3*x]+4*Sin[x]-26*Exp[-2*x]+27*x^3,y[x],x,IncludeSingularSolution->True]
```

$$y(x) \rightarrow 3x^3 - 2x - 2e^{-2x} + \frac{\sin(x)}{2} + \frac{1}{18} \sin(3x) + \left(-\frac{x}{3} + c_1 \right) \cos(3x) + c_2 \sin(3x)$$

10.14 problem 4(a)

10.14.1 Solving as second order linear constant coeff ode	1801
10.14.2 Solving using Kovacic algorithm	1804
10.14.3 Maple step by step solution	1809

Internal problem ID [6315]

Internal file name [OUTPUT/5563_Sunday_June_05_2022_03_43_35_PM_96652659/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.2. THE METHOD OF UN-DETERMINED COEFFICIENTS. Page 67

Problem number: 4(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - 3y = e^{2x}$$

10.14.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = -3, f(x) = e^{2x}$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 3y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = -3$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 3 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 3 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = -3$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(-3)} \\ &= \pm \sqrt{3} \end{aligned}$$

Hence

$$\lambda_1 = +\sqrt{3}$$

$$\lambda_2 = -\sqrt{3}$$

Which simplifies to

$$\lambda_1 = \sqrt{3}$$

$$\lambda_2 = -\sqrt{3}$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(\sqrt{3})x} + c_2 e^{(-\sqrt{3})x}$$

Or

$$y = c_1 e^{\sqrt{3}x} + c_2 e^{-\sqrt{3}x}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^{\sqrt{3}x} + c_2 e^{-\sqrt{3}x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$e^{2x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[e^{2x}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^{\sqrt{3}x}, e^{-\sqrt{3}x}\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^{2x}$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_1 e^{2x} = e^{2x}$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = e^{2x}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^{\sqrt{3}x} + c_2 e^{-\sqrt{3}x}) + (e^{2x}) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{\sqrt{3}x} + c_2 e^{-\sqrt{3}x} + e^{2x} \tag{1}$$




Figure 337: Slope field plot

Verification of solutions

$$y = c_1 e^{\sqrt{3}x} + c_2 e^{-\sqrt{3}x} + e^{2x}$$

Verified OK.

10.14.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 3y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= -3 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} = \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{3}{1} \tag{6}$$

Comparing the above to (5) shows that

$$s = 3$$

$$t = 1$$

Therefore eq. (4) becomes

$$z''(x) = 3z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 267: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 3$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\sqrt{3}x}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned} y_1 &= z_1 \\ &= e^{-\sqrt{3}x} \end{aligned}$$

Which simplifies to

$$y_1 = e^{-\sqrt{3}x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= e^{-\sqrt{3}x} \int \frac{1}{e^{-2\sqrt{3}x}} dx \\ &= e^{-\sqrt{3}x} \left(\frac{e^{2\sqrt{3}x} \sqrt{3}}{6} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(e^{-\sqrt{3}x} \right) + c_2 \left(e^{-\sqrt{3}x} \left(\frac{e^{2\sqrt{3}x} \sqrt{3}}{6} \right) \right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 3y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-\sqrt{3}x} + \frac{c_2 \sqrt{3} e^{\sqrt{3}x}}{6}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$e^{2x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{2x}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ \frac{\sqrt{3} e^{\sqrt{3}x}}{6}, e^{-\sqrt{3}x} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^{2x}$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_1 e^{2x} = e^{2x}$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = e^{2x}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 e^{-\sqrt{3}x} + \frac{c_2 \sqrt{3} e^{\sqrt{3}x}}{6} \right) + (e^{2x}) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-\sqrt{3}x} + \frac{c_2 \sqrt{3} e^{\sqrt{3}x}}{6} + e^{2x} \quad (1)$$




Figure 338: Slope field plot

Verification of solutions

$$y = c_1 e^{-\sqrt{3}x} + \frac{c_2 \sqrt{3} e^{\sqrt{3}x}}{6} + e^{2x}$$

Verified OK.

10.14.3 Maple step by step solution

Let's solve

$$y'' - 3y = e^{2x}$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - 3 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{12})}{2}$$

- Roots of the characteristic polynomial
 $r = (\sqrt{3}, -\sqrt{3})$
- 1st solution of the homogeneous ODE
 $y_1(x) = e^{\sqrt{3}x}$
- 2nd solution of the homogeneous ODE
 $y_2(x) = e^{-\sqrt{3}x}$
- General solution of the ODE
 $y = c_1y_1(x) + c_2y_2(x) + y_p(x)$
- Substitute in solutions of the homogeneous ODE
 $y = c_1e^{\sqrt{3}x} + c_2e^{-\sqrt{3}x} + y_p(x)$
- Find a particular solution $y_p(x)$ of the ODE
 - Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = e^{2x} \right]$$
 - Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{\sqrt{3}x} & e^{-\sqrt{3}x} \\ \sqrt{3}e^{\sqrt{3}x} & -\sqrt{3}e^{-\sqrt{3}x} \end{bmatrix}$$
 - Compute Wronskian
 $W(y_1(x), y_2(x)) = -2\sqrt{3}$
 - Substitute functions into equation for $y_p(x)$

$$y_p(x) = \frac{\sqrt{3} \left(e^{\sqrt{3}x} \left(\int e^{-x(-2+\sqrt{3})} dx \right) - e^{-\sqrt{3}x} \left(\int e^{x(2+\sqrt{3})} dx \right) \right)}{6}$$
 - Compute integrals
 $y_p(x) = e^{2x}$
- Substitute particular solution into general solution to ODE
 $y = c_1e^{\sqrt{3}x} + c_2e^{-\sqrt{3}x} + e^{2x}$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(diff(y(x),x$2)-3*y(x)=exp(2*x),y(x), singsol=all)
```

$$y(x) = e^{\sqrt{3}x}c_2 + e^{-\sqrt{3}x}c_1 + e^{2x}$$

✓ Solution by Mathematica

Time used: 0.103 (sec). Leaf size: 36

```
DSolve[y''[x]-3*y[x]==Exp[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{2x} + c_1 e^{\sqrt{3}x} + c_2 e^{-\sqrt{3}x}$$

10.15 problem 4(b)

10.15.1 Maple step by step solution 1816

Internal problem ID [6316]

Internal file name [OUTPUT/5564_Sunday_June_05_2022_03_43_37_PM_76520915/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.2. THE METHOD OF UN-DETERMINED COEFFICIENTS. Page 67

Problem number: 4(b).

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_3rd_order , _missing_y]]
```

$$y''' + y' = \sin(x)$$

This is higher order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE And y_p is a particular solution to the nonhomogeneous ODE. y_h is the solution to

$$y''' + y' = 0$$

The characteristic equation is

$$\lambda^3 + \lambda = 0$$

The roots of the above equation are

$$\lambda_1 = 0$$

$$\lambda_2 = i$$

$$\lambda_3 = -i$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 + e^{-ix}c_2 + e^{ix}c_3$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = 1$$

$$y_2 = e^{-ix}$$

$$y_3 = e^{ix}$$

Now the particular solution to the given ODE is found

$$y''' + y' = \sin(x)$$

Let the particular solution be

$$y_p = U_1y_1 + U_2y_2 + U_3y_3$$

Where y_i are the basis solutions found above for the homogeneous solution y_h and $U_i(x)$ are functions to be determined as follows

$$U_i = (-1)^{n-i} \int \frac{F(x)W_i(x)}{aW(x)} dx$$

Where $W(x)$ is the Wronskian and $W_i(x)$ is the Wronskian that results after deleting the last row and the i -th column of the determinant and n is the order of the ODE or equivalently, the number of basis solutions, and a is the coefficient of the leading derivative in the ODE, and $F(x)$ is the RHS of the ODE. Therefore, the first step is to find the Wronskian $W(x)$. This is given by

$$W(x) = \begin{vmatrix} y_1 & y_2 & y_3 \\ y_1' & y_2' & y_3' \\ y_1'' & y_2'' & y_3'' \end{vmatrix}$$

Substituting the fundamental set of solutions y_i found above in the Wronskian gives

$$W = \begin{bmatrix} 1 & e^{-ix} & e^{ix} \\ 0 & -ie^{-ix} & ie^{ix} \\ 0 & -e^{-ix} & -e^{ix} \end{bmatrix}$$

$$|W| = 2ie^{-ix}e^{ix}$$

The determinant simplifies to

$$|W| = 2i$$

Now we determine W_i for each U_i .

$$\begin{aligned} W_1(x) &= \det \begin{bmatrix} e^{-ix} & e^{ix} \\ -ie^{-ix} & ie^{ix} \end{bmatrix} \\ &= 2i \end{aligned}$$

$$\begin{aligned} W_2(x) &= \det \begin{bmatrix} 1 & e^{ix} \\ 0 & ie^{ix} \end{bmatrix} \\ &= ie^{ix} \end{aligned}$$

$$\begin{aligned} W_3(x) &= \det \begin{bmatrix} 1 & e^{-ix} \\ 0 & -ie^{-ix} \end{bmatrix} \\ &= -ie^{-ix} \end{aligned}$$

Now we are ready to evaluate each $U_i(x)$.

$$\begin{aligned} U_1 &= (-1)^{3-1} \int \frac{F(x)W_1(x)}{aW(x)} dx \\ &= (-1)^2 \int \frac{(\sin(x))(2i)}{(1)(2i)} dx \\ &= \int \frac{2i \sin(x)}{2i} dx \\ &= \int (\sin(x)) dx \\ &= -\cos(x) \end{aligned}$$

$$\begin{aligned} U_2 &= (-1)^{3-2} \int \frac{F(x)W_2(x)}{aW(x)} dx \\ &= (-1)^1 \int \frac{(\sin(x))(ie^{ix})}{(1)(2i)} dx \\ &= - \int \frac{i \sin(x) e^{ix}}{2i} dx \\ &= - \int \left(\frac{\sin(x) e^{ix}}{2} \right) dx \\ &= - \left(\int \frac{\sin(x) e^{ix}}{2} dx \right) \end{aligned}$$

$$\begin{aligned}
U_3 &= (-1)^{3-3} \int \frac{F(x)W_3(x)}{aW(x)} dx \\
&= (-1)^0 \int \frac{(\sin(x))(-ie^{-ix})}{(1)(2i)} dx \\
&= \int \frac{-i \sin(x) e^{-ix}}{2i} dx \\
&= \int \left(-\frac{\sin(x) e^{-ix}}{2} \right) dx \\
&= \int -\frac{\sin(x) e^{-ix}}{2} dx
\end{aligned}$$

Now that all the U_i functions have been determined, the particular solution is found from

$$y_p = U_1 y_1 + U_2 y_2 + U_3 y_3$$

Hence

$$\begin{aligned}
y_p &= (-\cos(x)) \\
&\quad + \left(-\left(\int \frac{\sin(x) e^{ix}}{2} dx \right) \right) (e^{-ix}) \\
&\quad + \left(\int -\frac{\sin(x) e^{-ix}}{2} dx \right) (e^{ix})
\end{aligned}$$

Therefore the particular solution is

$$y_p = -\cos(x) - \frac{(\int \sin(x) e^{ix} dx) e^{-ix}}{2} - \frac{(\int \sin(x) e^{-ix} dx) e^{ix}}{2}$$

Which simplifies to

$$y_p = -\left(\int \sin(x)^2 dx \right) \sin(x) - \frac{\cos(x) (\int \sin(2x) dx + 2)}{2}$$

Therefore the general solution is

$$\begin{aligned}
y &= y_h + y_p \\
&= (c_1 + e^{-ix} c_2 + e^{ix} c_3) + \left(-\left(\int \sin(x)^2 dx \right) \sin(x) - \frac{\cos(x) (\int \sin(2x) dx + 2)}{2} \right)
\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 + e^{-ix} c_2 + e^{ix} c_3 - \left(\int \sin(x)^2 dx \right) \sin(x) - \frac{\cos(x) (\int \sin(2x) dx + 2)}{2} \quad (1)$$

Verification of solutions

$$y = c_1 + e^{-ix} c_2 + e^{ix} c_3 - \left(\int \sin(x)^2 dx \right) \sin(x) - \frac{\cos(x) (\int \sin(2x) dx + 2)}{2}$$

Verified OK.

10.15.1 Maple step by step solution

Let's solve

$$y''' + y' = \sin(x)$$

- Highest derivative means the order of the ODE is 3

$$y'''$$

- Convert linear ODE into a system of first order ODEs

- Define new variable $y_1(x)$

$$y_1(x) = y$$

- Define new variable $y_2(x)$

$$y_2(x) = y'$$

- Define new variable $y_3(x)$

$$y_3(x) = y''$$

- Isolate for $y_3'(x)$ using original ODE

$$y_3'(x) = \sin(x) - y_2(x)$$

Convert linear ODE into a system of first order ODEs

$$[y_2(x) = y_1'(x), y_3(x) = y_2'(x), y_3'(x) = \sin(x) - y_2(x)]$$

- Define vector

$$\vec{y}(x) = \begin{bmatrix} y_1(x) \\ y_2(x) \\ y_3(x) \end{bmatrix}$$

- System to solve

$$\vec{y}'(x) = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{bmatrix} \cdot \vec{y}(x) + \begin{bmatrix} 0 \\ 0 \\ \sin(x) \end{bmatrix}$$

- Define the forcing function

$$\vec{f}(x) = \begin{bmatrix} 0 \\ 0 \\ \sin(x) \end{bmatrix}$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{bmatrix}$$

- Rewrite the system as

$$\vec{y}'(x) = A \cdot \vec{y}(x) + \vec{f}$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[0, \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \right], \left[-I, \begin{bmatrix} -1 \\ I \\ 1 \end{bmatrix} \right], \left[I, \begin{bmatrix} -1 \\ -I \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[0, \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$\left[-I, \begin{bmatrix} -1 \\ I \\ 1 \end{bmatrix} \right]$$

- Solution from eigenpair

$$e^{-Ix} \cdot \begin{bmatrix} -1 \\ I \\ 1 \end{bmatrix}$$

- Use Euler identity to write solution in terms of sin and cos

$$(\cos(x) - I \sin(x)) \cdot \begin{bmatrix} -1 \\ I \\ 1 \end{bmatrix}$$

- Simplify expression

$$\begin{bmatrix} I \sin(x) - \cos(x) \\ I(\cos(x) - I \sin(x)) \\ \cos(x) - I \sin(x) \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\left[\vec{y}_2(x) = \begin{bmatrix} -\cos(x) \\ \sin(x) \\ \cos(x) \end{bmatrix}, \vec{y}_3(x) = \begin{bmatrix} \sin(x) \\ \cos(x) \\ -\sin(x) \end{bmatrix} \right]$$

- General solution of the system of ODEs can be written in terms of the particular solution $\vec{y}_p(x)$

$$\vec{y}(x) = c_1 \vec{y}_1 + c_2 \vec{y}_2(x) + c_3 \vec{y}_3(x) + \vec{y}_p(x)$$

- Fundamental matrix

- Let $\phi(x)$ be the matrix whose columns are the independent solutions of the homogeneous system

$$\phi(x) = \begin{bmatrix} 1 & -\cos(x) & \sin(x) \\ 0 & \sin(x) & \cos(x) \\ 0 & \cos(x) & -\sin(x) \end{bmatrix}$$

- The fundamental matrix, $\Phi(x)$ is a normalized version of $\phi(x)$ satisfying $\Phi(0) = I$ where I is the identity matrix

$$\Phi(x) = \phi(x) \cdot \frac{1}{\phi(0)}$$

- Substitute the value of $\phi(x)$ and $\phi(0)$

$$\Phi(x) = \begin{bmatrix} 1 & -\cos(x) & \sin(x) \\ 0 & \sin(x) & \cos(x) \\ 0 & \cos(x) & -\sin(x) \end{bmatrix} \cdot \frac{1}{\begin{bmatrix} 1 & -1 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}}$$

- Evaluate and simplify to get the fundamental matrix

$$\Phi(x) = \begin{bmatrix} 1 & \sin(x) & 1 - \cos(x) \\ 0 & \cos(x) & \sin(x) \\ 0 & -\sin(x) & \cos(x) \end{bmatrix}$$

- Find a particular solution of the system of ODEs using variation of parameters

- Let the particular solution be the fundamental matrix multiplied by $\vec{v}(x)$ and solve for $\vec{v}(x)$

$$\vec{y}_p(x) = \Phi(x) \cdot \vec{v}(x)$$

- Take the derivative of the particular solution

$$\vec{y}'_p(x) = \Phi'(x) \cdot \vec{v}(x) + \Phi(x) \cdot \vec{v}'(x)$$

- Substitute particular solution and its derivative into the system of ODEs

$$\Phi'(x) \cdot \vec{v}(x) + \Phi(x) \cdot \vec{v}'(x) = A \cdot \Phi(x) \cdot \vec{v}(x) + \vec{f}(x)$$

- The fundamental matrix has columns that are solutions to the homogeneous system so its derivative is $A \cdot \Phi(x)$

$$A \cdot \Phi(x) \cdot \vec{v}(x) + \Phi(x) \cdot \vec{v}'(x) = A \cdot \Phi(x) \cdot \vec{v}(x) + \vec{f}(x)$$

- Cancel like terms

$$\Phi(x) \cdot \vec{v}'(x) = \vec{f}(x)$$

- Multiply by the inverse of the fundamental matrix

$$\vec{v}'(x) = \frac{1}{\Phi(x)} \cdot \vec{f}(x)$$

- Integrate to solve for $\vec{v}(x)$

$$\vec{v}(x) = \int_0^x \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds$$

- Plug $\vec{v}(x)$ into the equation for the particular solution

$$\vec{y}_p(x) = \Phi(x) \cdot \left(\int_0^x \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds \right)$$

- Plug in the fundamental matrix and the forcing function and compute

$$\vec{y}_p(x) = \begin{bmatrix} 1 - \frac{\sin(x)x}{2} - \cos(x) \\ \frac{\sin(x)}{2} - \frac{\cos(x)x}{2} \\ \frac{\sin(x)x}{2} \end{bmatrix}$$

- Plug particular solution back into general solution

$$\vec{y}(x) = c_1 \vec{y}_1 + c_2 \vec{y}_2(x) + c_3 \vec{y}_3(x) + \begin{bmatrix} 1 - \frac{\sin(x)x}{2} - \cos(x) \\ \frac{\sin(x)}{2} - \frac{\cos(x)x}{2} \\ \frac{\sin(x)x}{2} \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y = 1 + (-1 - c_2) \cos(x) + \frac{(2c_3 - x)\sin(x)}{2} + c_1$$

Maple trace

```

`Methods for third order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 3; linear nonhomogeneous with symmetry [0,1]
-> Calling odsolve with the ODE`, diff(diff(_b(_a), _a), _a) = -_b(_a)+sin(_a), _b(_a)`
Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
    <- solving first the homogeneous part of the ODE successful
<- differential order: 3; linear nonhomogeneous with symmetry [0,1] successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve(diff(y(x),x$3)+diff(y(x),x)=sin(x),y(x), singsol=all)
```

$$y(x) = (-1 - c_2) \cos(x) + \frac{(2c_1 - x) \sin(x)}{2} + c_3$$

✓ Solution by Mathematica

Time used: 0.118 (sec). Leaf size: 31

```
DSolve[y'''[x]+y'[x]==Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{2}(1 + 2c_2) \cos(x) + \left(-\frac{x}{2} + c_1\right) \sin(x) + c_3$$

11 Chapter 2. Second-Order Linear Equations.
Section 2.3. THE METHOD OF VARIATION
OF PARAMETERS. Page 71

11.1 problem 1(a)	1823
11.2 problem 1(b)	1836
11.3 problem 1(c)	1851
11.4 problem 1(d)	1862
11.5 problem 1(e)	1875
11.6 problem 1(f)	1886
11.7 problem 2(a)	1898
11.8 problem 2(b)	1911
11.9 problem 2(c)	1924
11.10 problem 2(d)	1938
11.11 problem 2(e)	1949
11.12 problem 2(f)	1962
11.13 problem 2(g)	1975
11.14 problem 3	1988
11.15 problem 4	2000
11.16 problem 5(a)	2011
11.17 problem 5(b)	2030
11.18 problem 5(c)	2046
11.19 problem 5(d)	2066
11.20 problem 5(e)	2081

11.1 problem 1(a)

- 11.1.1 Solving as second order linear constant coeff ode 1823
- 11.1.2 Solving using Kovacic algorithm 1828
- 11.1.3 Maple step by step solution 1834

Internal problem ID [6317]

Internal file name [OUTPUT/5565_Sunday_June_05_2022_03_43_39_PM_26639439/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.3. THE METHOD OF VARIATION OF PARAMETERS. Page 71

Problem number: 1(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + 4y = \tan(2x)$$

11.1.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 4, f(x) = \tan(2x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 4y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 4$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 4 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 4 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 4$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(4)} \\ &= \pm 2i \end{aligned}$$

Hence

$$\lambda_1 = +2i$$

$$\lambda_2 = -2i$$

Which simplifies to

$$\lambda_1 = 2i$$

$$\lambda_2 = -2i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 2$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (c_1 \cos(2x) + c_2 \sin(2x))$$

Or

$$y = c_1 \cos(2x) + c_2 \sin(2x)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \cos(2x) + c_2 \sin(2x)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \cos(2x)$$

$$y_2 = \sin(2x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(2x) & \sin(2x) \\ \frac{d}{dx}(\cos(2x)) & \frac{d}{dx}(\sin(2x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(2x) & \sin(2x) \\ -2 \sin(2x) & 2 \cos(2x) \end{vmatrix}$$

Therefore

$$W = (\cos(2x))(2 \cos(2x)) - (\sin(2x))(-2 \sin(2x))$$

Which simplifies to

$$W = 2 \cos (2x)^2 + 2 \sin (2x)^2$$

Which simplifies to

$$W = 2$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\sin (2x) \tan (2x)}{2} dx$$

Which simplifies to

$$u_1 = - \int \frac{\sin (2x) \tan (2x)}{2} dx$$

Hence

$$u_1 = \frac{\sin (2x)}{4} - \frac{\ln (\sec (2x) + \tan (2x))}{4}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\cos (2x) \tan (2x)}{2} dx$$

Which simplifies to

$$u_2 = \int \frac{\sin (2x)}{2} dx$$

Hence

$$u_2 = -\frac{\cos (2x)}{4}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(\frac{\sin (2x)}{4} - \frac{\ln (\sec (2x) + \tan (2x))}{4} \right) \cos (2x) - \frac{\cos (2x) \sin (2x)}{4}$$

Which simplifies to

$$y_p(x) = -\frac{\cos (2x) \ln (\sec (2x) + \tan (2x))}{4}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(2x) + c_2 \sin(2x)) + \left(-\frac{\cos(2x) \ln(\sec(2x) + \tan(2x))}{4} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(2x) + c_2 \sin(2x) - \frac{\cos(2x) \ln(\sec(2x) + \tan(2x))}{4} \quad (1)$$




Figure 339: Slope field plot

Verification of solutions

$$y = c_1 \cos(2x) + c_2 \sin(2x) - \frac{\cos(2x) \ln(\sec(2x) + \tan(2x))}{4}$$

Verified OK.

11.1.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 4y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 4 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-4}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -4 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -4z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 270: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -4$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(2x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}y_1 &= z_1 \\ &= \cos(2x)\end{aligned}$$

Which simplifies to

$$y_1 = \cos(2x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned}y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(2x) \int \frac{1}{\cos(2x)^2} dx \\ &= \cos(2x) \left(\frac{\tan(2x)}{2} \right)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1(\cos(2x)) + c_2 \left(\cos(2x) \left(\frac{\tan(2x)}{2} \right) \right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$.
 y_h is the solution to

$$y'' + 4y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 \cos(2x) + \frac{c_2 \sin(2x)}{2}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \cos(2x)$$

$$y_2 = \frac{\sin(2x)}{2}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(2x) & \frac{\sin(2x)}{2} \\ \frac{d}{dx}(\cos(2x)) & \frac{d}{dx}\left(\frac{\sin(2x)}{2}\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(2x) & \frac{\sin(2x)}{2} \\ -2 \sin(2x) & \cos(2x) \end{vmatrix}$$

Therefore

$$W = (\cos(2x))(\cos(2x)) - \left(\frac{\sin(2x)}{2}\right)(-2\sin(2x))$$

Which simplifies to

$$W = \cos(2x)^2 + \sin(2x)^2$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\frac{\sin(2x)\tan(2x)}{2}}{1} dx$$

Which simplifies to

$$u_1 = - \int \frac{\sin(2x)\tan(2x)}{2} dx$$

Hence

$$u_1 = \frac{\sin(2x)}{4} - \frac{\ln(\sec(2x) + \tan(2x))}{4}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\cos(2x)\tan(2x)}{1} dx$$

Which simplifies to

$$u_2 = \int \sin(2x) dx$$

Hence

$$u_2 = -\frac{\cos(2x)}{2}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(\frac{\sin(2x)}{4} - \frac{\ln(\sec(2x) + \tan(2x))}{4}\right) \cos(2x) - \frac{\cos(2x)\sin(2x)}{4}$$

Which simplifies to

$$y_p(x) = -\frac{\cos(2x) \ln(\sec(2x) + \tan(2x))}{4}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 \cos(2x) + \frac{c_2 \sin(2x)}{2} \right) + \left(-\frac{\cos(2x) \ln(\sec(2x) + \tan(2x))}{4} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(2x) + \frac{c_2 \sin(2x)}{2} - \frac{\cos(2x) \ln(\sec(2x) + \tan(2x))}{4} \quad (1)$$




Figure 340: Slope field plot

Verification of solutions

$$y = c_1 \cos(2x) + \frac{c_2 \sin(2x)}{2} - \frac{\cos(2x) \ln(\sec(2x) + \tan(2x))}{4}$$

Verified OK.

11.1.3 Maple step by step solution

Let's solve

$$y'' + 4y = \tan(2x)$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 4 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-16})}{2}$$

- Roots of the characteristic polynomial

$$r = (-2i, 2i)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(2x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(2x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 \cos(2x) + c_2 \sin(2x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \tan(2x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(2x) & \sin(2x) \\ -2\sin(2x) & 2\cos(2x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 2$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{\cos(2x)(\int \sin(2x) \tan(2x) dx)}{2} + \frac{\sin(2x)(\int \sin(2x) dx)}{2}$$

- Compute integrals

$$y_p(x) = -\frac{\cos(2x) \ln(\sec(2x) + \tan(2x))}{4}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 \cos(2x) + c_2 \sin(2x) - \frac{\cos(2x) \ln(\sec(2x) + \tan(2x))}{4}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 33

```
dsolve(diff(y(x),x$2)+4*y(x)=tan(2*x),y(x), singsol=all)
```

$$y(x) = \sin(2x) c_2 + \cos(2x) c_1 - \frac{\cos(2x) \ln(\sec(2x) + \tan(2x))}{4}$$

✓ Solution by Mathematica

Time used: 0.048 (sec). Leaf size: 40

```
DSolve[y''[x]+4*y[x]==Tan[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{4} \cos(2x) \operatorname{arctanh}(\sin(2x)) + c_1 \cos(2x) + \frac{1}{4} (-1 + 4c_2) \sin(2x)$$

11.2 problem 1(b)

11.2.1 Solving as second order linear constant coeff ode	1836
11.2.2 Solving as linear second order ode solved by an integrating factor ode	1840
11.2.3 Solving using Kovacic algorithm	1842
11.2.4 Maple step by step solution	1848

Internal problem ID [6318]

Internal file name [OUTPUT/5566_Sunday_June_05_2022_03_43_40_PM_41678587/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.3. THE METHOD OF VARIATION OF PARAMETERS. Page 71

Problem number: 1(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "linear_second_order_ode_solved_by_an_integrating_factor"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + 2y' + y = e^{-x} \ln(x)$$

11.2.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 2, C = 1, f(x) = e^{-x} \ln(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 2y' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 2, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 2\lambda e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 2\lambda + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 2, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-2}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{(2)^2 - (4)(1)(1)} \\ &= -1 \end{aligned}$$

Hence this is the case of a double root $\lambda_{1,2} = 1$. Therefore the solution is

$$y = c_1 e^{-x} + c_2 x e^{-x} \quad (1)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^{-x} + x e^{-x} c_2$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned} y_1 &= e^{-x} \\ y_2 &= x e^{-x} \end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^{-x} & x e^{-x} \\ \frac{d}{dx}(e^{-x}) & \frac{d}{dx}(x e^{-x}) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^{-x} & x e^{-x} \\ -e^{-x} & e^{-x} - x e^{-x} \end{vmatrix}$$

Therefore

$$W = (e^{-x})(e^{-x} - x e^{-x}) - (x e^{-x})(-e^{-x})$$

Which simplifies to

$$W = e^{-2x}$$

Which simplifies to

$$W = e^{-2x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x e^{-2x} \ln(x)}{e^{-2x}} dx$$

Which simplifies to

$$u_1 = - \int \ln(x) x dx$$

Hence

$$u_1 = -\frac{\ln(x) x^2}{2} + \frac{x^2}{4}$$

And Eq. (3) becomes

$$u_2 = \int \frac{e^{-2x} \ln(x)}{e^{-2x}} dx$$

Which simplifies to

$$u_2 = \int \ln(x) dx$$

Hence

$$u_2 = \ln(x) x - x$$

Which simplifies to

$$u_1 = -\frac{x^2(-1 + 2 \ln(x))}{4}$$

$$u_2 = x(\ln(x) - 1)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{x^2(-1 + 2 \ln(x)) e^{-x}}{4} + x^2(\ln(x) - 1) e^{-x}$$

Which simplifies to

$$y_p(x) = \frac{x^2 e^{-x}(-3 + 2 \ln(x))}{4}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^{-x} + x e^{-x} c_2) + \left(\frac{x^2 e^{-x}(-3 + 2 \ln(x))}{4} \right) \end{aligned}$$

Which simplifies to

$$y = e^{-x}(c_2 x + c_1) + \frac{x^2 e^{-x}(-3 + 2 \ln(x))}{4}$$

Summary

The solution(s) found are the following

$$y = e^{-x}(c_2x + c_1) + \frac{x^2e^{-x}(-3 + 2 \ln(x))}{4} \quad (1)$$




Figure 341: Slope field plot

Verification of solutions

$$y = e^{-x}(c_2x + c_1) + \frac{x^2e^{-x}(-3 + 2 \ln(x))}{4}$$

Verified OK.

11.2.2 Solving as linear second order ode solved by an integrating factor ode

The ode satisfies this form

$$y'' + p(x)y' + \frac{(p(x))^2 + p'(x)}{2}y = f(x)$$

Where $p(x) = 2$. Therefore, there is an integrating factor given by

$$\begin{aligned}M(x) &= e^{\frac{1}{2} \int p \, dx} \\ &= e^{\int 2 \, dx} \\ &= e^x\end{aligned}$$

Multiplying both sides of the ODE by the integrating factor $M(x)$ makes the left side of the ODE a complete differential

$$\begin{aligned}(M(x)y)'' &= e^x e^{-x} \ln(x) \\ (y e^x)'' &= e^x e^{-x} \ln(x)\end{aligned}$$

Integrating once gives

$$(y e^x)' = x(\ln(x) - 1) + c_1$$

Integrating again gives

$$(y e^x) = \frac{x(2 \ln(x) x + 4c_1 - 3x)}{4} + c_2$$

Hence the solution is

$$y = \frac{\frac{x(2 \ln(x)x + 4c_1 - 3x)}{4} + c_2}{e^x}$$

Or

$$y = \frac{x^2 e^{-x} \ln(x)}{2} + c_1 x e^{-x} - \frac{3x^2 e^{-x}}{4} + c_2 e^{-x}$$

Summary

The solution(s) found are the following

$$y = \frac{x^2 e^{-x} \ln(x)}{2} + c_1 x e^{-x} - \frac{3x^2 e^{-x}}{4} + c_2 e^{-x} \quad (1)$$




Figure 342: Slope field plot

Verification of solutions

$$y = \frac{x^2 e^{-x} \ln(x)}{2} + c_1 x e^{-x} - \frac{3x^2 e^{-x}}{4} + c_2 e^{-x}$$

Verified OK.

11.2.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 2y' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 2 \\ C &= 1 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} \tag{5}$$

$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \tag{6}$$

Comparing the above to (5) shows that

$$s = 0$$

$$t = 1$$

Therefore eq. (4) becomes

$$z''(x) = 0 \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 272: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty\end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{2}{1} dx} \\ &= z_1 e^{-x} \\ &= z_1 (e^{-x})\end{aligned}$$

Which simplifies to

$$y_1 = e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{2}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-2x}}{(y_1)^2} dx \\ &= y_1(x)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-x}) + c_2 (e^{-x}(x))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 2y' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-x} + x e^{-x} c_2$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned}y_1 &= e^{-x} \\ y_2 &= x e^{-x}\end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^{-x} & x e^{-x} \\ \frac{d}{dx}(e^{-x}) & \frac{d}{dx}(x e^{-x}) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^{-x} & x e^{-x} \\ -e^{-x} & e^{-x} - x e^{-x} \end{vmatrix}$$

Therefore

$$W = (e^{-x})(e^{-x} - x e^{-x}) - (x e^{-x})(-e^{-x})$$

Which simplifies to

$$W = e^{-2x}$$

Which simplifies to

$$W = e^{-2x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x e^{-2x} \ln(x)}{e^{-2x}} dx$$

Which simplifies to

$$u_1 = - \int \ln(x) x dx$$

Hence

$$u_1 = -\frac{\ln(x) x^2}{2} + \frac{x^2}{4}$$

And Eq. (3) becomes

$$u_2 = \int \frac{e^{-2x} \ln(x)}{e^{-2x}} dx$$

Which simplifies to

$$u_2 = \int \ln(x) dx$$

Hence

$$u_2 = \ln(x)x - x$$

Which simplifies to

$$u_1 = -\frac{x^2(-1 + 2 \ln(x))}{4}$$
$$u_2 = x(\ln(x) - 1)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{x^2(-1 + 2 \ln(x)) e^{-x}}{4} + x^2(\ln(x) - 1) e^{-x}$$

Which simplifies to

$$y_p(x) = \frac{x^2 e^{-x}(-3 + 2 \ln(x))}{4}$$

Therefore the general solution is

$$y = y_h + y_p$$
$$= (c_1 e^{-x} + x e^{-x} c_2) + \left(\frac{x^2 e^{-x}(-3 + 2 \ln(x))}{4} \right)$$

Which simplifies to

$$y = e^{-x}(c_2 x + c_1) + \frac{x^2 e^{-x}(-3 + 2 \ln(x))}{4}$$

Summary

The solution(s) found are the following

$$y = e^{-x}(c_2 x + c_1) + \frac{x^2 e^{-x}(-3 + 2 \ln(x))}{4} \quad (1)$$




Figure 343: Slope field plot

Verification of solutions

$$y = e^{-x}(c_2x + c_1) + \frac{x^2e^{-x}(-3 + 2 \ln(x))}{4}$$

Verified OK.

11.2.4 Maple step by step solution

Let's solve

$$y'' + 2y' + y = e^{-x} \ln(x)$$

- Highest derivative means the order of the ODE is 2
- $$y''$$
- Characteristic polynomial of homogeneous ODE
- $$r^2 + 2r + 1 = 0$$
- Factor the characteristic polynomial
- $$(r + 1)^2 = 0$$
- Root of the characteristic polynomial

$$r = -1$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-x}$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = x e^{-x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{-x} + x e^{-x} c_2 + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = e^{-x} \ln(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-x} & x e^{-x} \\ -e^{-x} & e^{-x} - x e^{-x} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = e^{-2x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = e^{-x} \left(- \left(\int \ln(x) x dx \right) + \left(\int \ln(x) dx \right) x \right)$$

- Compute integrals

$$y_p(x) = \frac{x^2 e^{-x} (-3 + 2 \ln(x))}{4}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{-x} + x e^{-x} c_2 + \frac{x^2 e^{-x} (-3 + 2 \ln(x))}{4}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 30

```
dsolve(diff(y(x),x$2)+2*diff(y(x),x)+y(x)=exp(-x)*ln(x),y(x), singsol=all)
```

$$y(x) = \frac{e^{-x}(2 \ln(x) x^2 + 4c_1 x - 3x^2 + 4c_2)}{4}$$

✓ Solution by Mathematica

Time used: 0.037 (sec). Leaf size: 36

```
DSolve[y''[x]+2*y'[x]+y[x]==Exp[-x]*Log[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{4}e^{-x}(-3x^2 + 2x^2 \log(x) + 4c_2x + 4c_1)$$

11.3 problem 1(c)

- 11.3.1 Solving as second order linear constant coeff ode 1851
- 11.3.2 Solving using Kovacic algorithm 1854
- 11.3.3 Maple step by step solution 1859

Internal problem ID [6319]

Internal file name [OUTPUT/5567_Sunday_June_05_2022_03_43_42_PM_84385027/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.3. THE METHOD OF VARIATION OF PARAMETERS. Page 71

Problem number: 1(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' - 2y' - 3y = 64x e^{-x}$$

11.3.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = -2, C = -3, f(x) = 64x e^{-x}$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 2y' - 3y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -2, C = -3$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 2\lambda e^{\lambda x} - 3e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 2\lambda - 3 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -2, C = -3$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{2}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-2^2 - (4)(1)(-3)} \\ &= 1 \pm 2 \end{aligned}$$

Hence

$$\lambda_1 = 1 + 2$$

$$\lambda_2 = 1 - 2$$

Which simplifies to

$$\lambda_1 = 3$$

$$\lambda_2 = -1$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(3)x} + c_2 e^{(-1)x}$$

Or

$$y = e^{3x} c_1 + c_2 e^{-x}$$

Therefore the homogeneous solution y_h is

$$y_h = e^{3x} c_1 + c_2 e^{-x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$64x e^{-x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{x e^{-x}, e^{-x}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^{-x}, e^{3x}\}$$

Since e^{-x} is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x e^{-x}, x^2 e^{-x}\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x e^{-x} + A_2 x^2 e^{-x}$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-4A_1 e^{-x} + 2A_2 e^{-x} - 8A_2 x e^{-x} = 64x e^{-x}$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = -4, A_2 = -8]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -4x e^{-x} - 8x^2 e^{-x}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (e^{3x} c_1 + c_2 e^{-x}) + (-4x e^{-x} - 8x^2 e^{-x}) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = e^{3x}c_1 + c_2e^{-x} - 4xe^{-x} - 8x^2e^{-x} \quad (1)$$




Figure 344: Slope field plot

Verification of solutions

$$y = e^{3x}c_1 + c_2e^{-x} - 4xe^{-x} - 8x^2e^{-x}$$

Verified OK.

11.3.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 2y' - 3y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = -2 \quad (3)$$

$$C = -3$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{4}{1} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 4 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 4z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 274: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 4$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-2x}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}
 y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\
 &= z_1 e^{-\int \frac{1}{2} \frac{-2}{1} dx} \\
 &= z_1 e^x \\
 &= z_1 (e^x)
 \end{aligned}$$

Which simplifies to

$$y_1 = e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{2}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{2x}}{(y_1)^2} dx \\ &= y_1 \left(\frac{e^{4x}}{4} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-x}) + c_2 \left(e^{-x} \left(\frac{e^{4x}}{4} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 2y' - 3y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-x} + \frac{c_2 e^{3x}}{4}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$64x e^{-x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{x e^{-x}, e^{-x}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ \frac{e^{3x}}{4}, e^{-x} \right\}$$

Since e^{-x} is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x e^{-x}, x^2 e^{-x}\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x e^{-x} + A_2 x^2 e^{-x}$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-4A_1 e^{-x} + 2A_2 e^{-x} - 8A_2 x e^{-x} = 64x e^{-x}$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = -4, A_2 = -8]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -4x e^{-x} - 8x^2 e^{-x}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 e^{-x} + \frac{c_2 e^{3x}}{4} \right) + (-4x e^{-x} - 8x^2 e^{-x}) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-x} + \frac{c_2 e^{3x}}{4} - 4x e^{-x} - 8x^2 e^{-x} \quad (1)$$




Figure 345: Slope field plot

Verification of solutions

$$y = c_1 e^{-x} + \frac{c_2 e^{3x}}{4} - 4x e^{-x} - 8x^2 e^{-x}$$

Verified OK.

11.3.3 Maple step by step solution

Let's solve

$$y'' - 2y' - 3y = 64x e^{-x}$$

- Highest derivative means the order of the ODE is 2
- y''
- Characteristic polynomial of homogeneous ODE

$$r^2 - 2r - 3 = 0$$

- Factor the characteristic polynomial

$$(r + 1)(r - 3) = 0$$

- Roots of the characteristic polynomial

$$r = (-1, 3)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{3x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{-x} + c_2 e^{3x} + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x), y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x), y_2(x))} dx \right), f(x) = 64x e^{-x} \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-x} & e^{3x} \\ -e^{-x} & 3e^{3x} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 4e^{2x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -16e^{-x} \left(\int x dx \right) + 16e^{3x} \left(\int e^{-4x} x dx \right)$$

- Compute integrals

$$y_p(x) = e^{-x}(-8x^2 - 4x - 1)$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{-x} + c_2 e^{3x} + e^{-x}(-8x^2 - 4x - 1)$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 26

```
dsolve(diff(y(x),x$2)-2*diff(y(x),x)-3*y(x)=64*x*exp(-x),y(x), singsol=all)
```

$$y(x) = (-8x^2 + c_1 - 4x) e^{-x} + c_2 e^{3x}$$

✓ Solution by Mathematica

Time used: 0.049 (sec). Leaf size: 31

```
DSolve[y''[x]-2*y'[x]-3*y[x]==64*x*Exp[-x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x}(-8x^2 - 4x + c_2 e^{4x} - 1 + c_1)$$

11.4 problem 1(d)

- 11.4.1 Solving as second order linear constant coeff ode 1862
- 11.4.2 Solving using Kovacic algorithm 1867
- 11.4.3 Maple step by step solution 1873

Internal problem ID [6320]

Internal file name [OUTPUT/5568_Sunday_June_05_2022_03_43_43_PM_22734416/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.3. THE METHOD OF VARIATION OF PARAMETERS. Page 71

Problem number: 1(d).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + 2y' + 5y = e^{-x} \sec(2x)$$

11.4.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 2, C = 5, f(x) = e^{-x} \sec(2x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 2y' + 5y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 2, C = 5$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 2\lambda e^{\lambda x} + 5 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 2\lambda + 5 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 2, C = 5$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-2}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{2^2 - (4)(1)(5)} \\ &= -1 \pm 2i \end{aligned}$$

Hence

$$\lambda_1 = -1 + 2i$$

$$\lambda_2 = -1 - 2i$$

Which simplifies to

$$\lambda_1 = -1 + 2i$$

$$\lambda_2 = -1 - 2i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = -1$ and $\beta = 2$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^{-x} (c_1 \cos(2x) + c_2 \sin(2x))$$

Therefore the homogeneous solution y_h is

$$y_h = e^{-x} (c_1 \cos(2x) + c_2 \sin(2x))$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = e^{-x} \cos(2x)$$

$$y_2 = e^{-x} \sin(2x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^{-x} \cos(2x) & e^{-x} \sin(2x) \\ \frac{d}{dx}(e^{-x} \cos(2x)) & \frac{d}{dx}(e^{-x} \sin(2x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^{-x} \cos(2x) & e^{-x} \sin(2x) \\ -e^{-x} \cos(2x) - 2e^{-x} \sin(2x) & -e^{-x} \sin(2x) + 2e^{-x} \cos(2x) \end{vmatrix}$$

Therefore

$$W = (e^{-x} \cos(2x)) (-e^{-x} \sin(2x) + 2e^{-x} \cos(2x)) - (e^{-x} \sin(2x)) (-e^{-x} \cos(2x) - 2e^{-x} \sin(2x))$$

Which simplifies to

$$W = 2e^{-2x} \cos(2x)^2 + 2e^{-2x} \sin(2x)^2$$

Which simplifies to

$$W = 2 e^{-2x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{e^{-2x} \sin(2x) \sec(2x)}{2 e^{-2x}} dx$$

Which simplifies to

$$u_1 = - \int \frac{\tan(2x)}{2} dx$$

Hence

$$u_1 = - \frac{\ln(1 + \tan(2x)^2)}{8}$$

And Eq. (3) becomes

$$u_2 = \int \frac{e^{-2x} \cos(2x) \sec(2x)}{2 e^{-2x}} dx$$

Which simplifies to

$$u_2 = \int \frac{1}{2} dx$$

Hence

$$u_2 = \frac{x}{2}$$

Which simplifies to

$$u_1 = - \frac{\ln(\sec(2x)^2)}{8}$$

$$u_2 = \frac{x}{2}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = - \frac{\ln(\sec(2x)^2) e^{-x} \cos(2x)}{8} + \frac{x e^{-x} \sin(2x)}{2}$$

Which simplifies to

$$y_p(x) = -\frac{e^{-x}(\ln(\sec(2x)^2)\cos(2x) - 4x\sin(2x))}{8}$$

Therefore the general solution is

$$y = y_h + y_p = (e^{-x}(c_1 \cos(2x) + c_2 \sin(2x))) + \left(-\frac{e^{-x}(\ln(\sec(2x)^2)\cos(2x) - 4x\sin(2x))}{8}\right)$$

Summary

The solution(s) found are the following

$$y = e^{-x}(c_1 \cos(2x) + c_2 \sin(2x)) - \frac{e^{-x}(\ln(\sec(2x)^2)\cos(2x) - 4x\sin(2x))}{8} \quad (1)$$




Figure 346: Slope field plot

Verification of solutions

$$y = e^{-x}(c_1 \cos(2x) + c_2 \sin(2x)) - \frac{e^{-x}(\ln(\sec(2x)^2)\cos(2x) - 4x\sin(2x))}{8}$$

Verified OK.

11.4.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 2y' + 5y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 2 \\ C &= 5 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-4}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -4 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -4z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 276: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -4$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(2x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{2}{1} dx} \end{aligned}$$

$$\begin{aligned}
&= z_1 e^{-x} \\
&= z_1 (e^{-x})
\end{aligned}$$

Which simplifies to

$$y_1 = e^{-x} \cos(2x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}
y_2 &= y_1 \int \frac{e^{\int -\frac{2}{1} dx}}{(y_1)^2} dx \\
&= y_1 \int \frac{e^{-2x}}{(y_1)^2} dx \\
&= y_1 \left(\frac{\tan(2x)}{2} \right)
\end{aligned}$$

Therefore the solution is

$$\begin{aligned}
y &= c_1 y_1 + c_2 y_2 \\
&= c_1 (e^{-x} \cos(2x)) + c_2 \left(e^{-x} \cos(2x) \left(\frac{\tan(2x)}{2} \right) \right)
\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 2y' + 5y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = e^{-x} \cos(2x) c_1 + \frac{e^{-x} \sin(2x) c_2}{2}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = e^{-x} \cos(2x)$$

$$y_2 = \frac{e^{-x} \sin(2x)}{2}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^{-x} \cos(2x) & \frac{e^{-x} \sin(2x)}{2} \\ \frac{d}{dx}(e^{-x} \cos(2x)) & \frac{d}{dx}\left(\frac{e^{-x} \sin(2x)}{2}\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^{-x} \cos(2x) & \frac{e^{-x} \sin(2x)}{2} \\ -e^{-x} \cos(2x) - 2e^{-x} \sin(2x) & -\frac{e^{-x} \sin(2x)}{2} + e^{-x} \cos(2x) \end{vmatrix}$$

Therefore

$$W = (e^{-x} \cos(2x)) \left(-\frac{e^{-x} \sin(2x)}{2} + e^{-x} \cos(2x) \right) - \left(\frac{e^{-x} \sin(2x)}{2} \right) (-e^{-x} \cos(2x) - 2e^{-x} \sin(2x))$$

Which simplifies to

$$W = e^{-2x} \cos(2x)^2 + e^{-2x} \sin(2x)^2$$

Which simplifies to

$$W = e^{-2x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\frac{e^{-2x} \sin(2x) \sec(2x)}{2}}{e^{-2x}} dx$$

Which simplifies to

$$u_1 = - \int \frac{\tan(2x)}{2} dx$$

Hence

$$u_1 = - \frac{\ln(1 + \tan(2x)^2)}{8}$$

And Eq. (3) becomes

$$u_2 = \int \frac{e^{-2x} \cos(2x) \sec(2x)}{e^{-2x}} dx$$

Which simplifies to

$$u_2 = \int 1 dx$$

Hence

$$u_2 = x$$

Which simplifies to

$$u_1 = - \frac{\ln(\sec(2x)^2)}{8}$$

$$u_2 = x$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = - \frac{\ln(\sec(2x)^2) e^{-x} \cos(2x)}{8} + \frac{x e^{-x} \sin(2x)}{2}$$

Which simplifies to

$$y_p(x) = -\frac{e^{-x}(\ln(\sec(2x)^2)\cos(2x) - 4x\sin(2x))}{8}$$

Therefore the general solution is

$$y = y_h + y_p = \left(e^{-x}\cos(2x)c_1 + \frac{e^{-x}\sin(2x)c_2}{2} \right) + \left(-\frac{e^{-x}(\ln(\sec(2x)^2)\cos(2x) - 4x\sin(2x))}{8} \right)$$

Summary

The solution(s) found are the following

$$y = e^{-x}\cos(2x)c_1 + \frac{e^{-x}\sin(2x)c_2}{2} - \frac{e^{-x}(\ln(\sec(2x)^2)\cos(2x) - 4x\sin(2x))}{8} \quad (1)$$




Figure 347: Slope field plot

Verification of solutions

$$y = e^{-x}\cos(2x)c_1 + \frac{e^{-x}\sin(2x)c_2}{2} - \frac{e^{-x}(\ln(\sec(2x)^2)\cos(2x) - 4x\sin(2x))}{8}$$

Verified OK.

11.4.3 Maple step by step solution

Let's solve

$$y'' + 2y' + 5y = e^{-x} \sec(2x)$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 2r + 5 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-2) \pm (\sqrt{-16})}{2}$$

- Roots of the characteristic polynomial

$$r = (-1 - 2I, -1 + 2I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-x} \cos(2x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{-x} \sin(2x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = e^{-x} \cos(2x) c_1 + e^{-x} \sin(2x) c_2 + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = e^{-x} \sec(2x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-x} \cos(2x) & e^{-x} \sin(2x) \\ -e^{-x} \cos(2x) - 2e^{-x} \sin(2x) & -e^{-x} \sin(2x) + 2e^{-x} \cos(2x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 2e^{-2x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{e^{-x}(\cos(2x)(\int \tan(2x)dx) - \sin(2x)(\int 1dx))}{2}$$

- Compute integrals

$$y_p(x) = -\frac{e^{-x}(\ln(\sec(2x)^2)\cos(2x) - 4x\sin(2x))}{8}$$

- Substitute particular solution into general solution to ODE

$$y = e^{-x} \cos(2x) c_1 + e^{-x} \sin(2x) c_2 - \frac{e^{-x}(\ln(\sec(2x)^2)\cos(2x) - 4x\sin(2x))}{8}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 39

```
dsolve(diff(y(x),x$2)+2*diff(y(x),x)+5*y(x)=exp(-x)*sec(2*x),y(x), singsol=all)
```

$$y(x) = \frac{\left(\frac{\cos(2x)\ln(\cos(2x))}{2} + 2\cos(2x)c_1 + \sin(2x)(2c_2 + x)\right)e^{-x}}{2}$$

✓ Solution by Mathematica

Time used: 0.068 (sec). Leaf size: 42

```
DSolve[y''[x]+2*y'[x]+5*y[x]==Exp[-x]*Sec[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{4}e^{-x}(2(x + 2c_1)\sin(2x) + \cos(2x)(\log(\cos(2x)) + 4c_2))$$

11.5 problem 1(e)

- 11.5.1 Solving as second order linear constant coeff ode 1875
- 11.5.2 Solving using Kovacic algorithm 1878
- 11.5.3 Maple step by step solution 1883

Internal problem ID [6321]

Internal file name [OUTPUT/5569_Sunday_June_05_2022_03_43_45_PM_50315069/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.3. THE METHOD OF VARIATION OF PARAMETERS. Page 71

Problem number: 1(e).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$2y'' + 3y' + y = e^{-3x}$$

11.5.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 2, B = 3, C = 1, f(x) = e^{-3x}$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$2y'' + 3y' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 2, B = 3, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$2\lambda^2 e^{\lambda x} + 3\lambda e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$2\lambda^2 + 3\lambda + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 2, B = 3, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-3}{(2)(2)} \pm \frac{1}{(2)(2)} \sqrt{3^2 - (4)(2)(1)} \\ &= -\frac{3}{4} \pm \frac{1}{4} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= -\frac{3}{4} + \frac{1}{4} \\ \lambda_2 &= -\frac{3}{4} - \frac{1}{4} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= -\frac{1}{2} \\ \lambda_2 &= -1 \end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned} y &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ y &= c_1 e^{(-\frac{1}{2})x} + c_2 e^{(-1)x} \end{aligned}$$

Or

$$y = c_1 e^{-\frac{x}{2}} + c_2 e^{-x}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^{-\frac{x}{2}} + c_2 e^{-x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$e^{-3x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{-3x}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^{-x}, e^{-\frac{x}{2}}\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^{-3x}$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$10A_1 e^{-3x} = e^{-3x}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{10} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{e^{-3x}}{10}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^{-\frac{x}{2}} + c_2 e^{-x}) + \left(\frac{e^{-3x}}{10} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-\frac{x}{2}} + c_2 e^{-x} + \frac{e^{-3x}}{10} \quad (1)$$




Figure 348: Slope field plot

Verification of solutions

$$y = c_1 e^{-\frac{x}{2}} + c_2 e^{-x} + \frac{e^{-3x}}{10}$$

Verified OK.

11.5.2 Solving using Kovacic algorithm

Writing the ode as

$$2y'' + 3y' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 2 \\ B &= 3 \\ C &= 1 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{1}{16} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 1 \\ t &= 16 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \frac{z(x)}{16} \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 278: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{1}{16}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{x}{4}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{3}{2} dx} \\ &= z_1 e^{-\frac{3x}{4}} \\ &= z_1 \left(e^{-\frac{3x}{4}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{3}{2} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-\frac{3x}{2}}}{(y_1)^2} dx \\ &= y_1 \left(2 e^{\frac{x}{2}} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-x}) + c_2 (e^{-x} (2 e^{\frac{x}{2}}))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$2y'' + 3y' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-x} + 2c_2 e^{-\frac{x}{2}}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$e^{-3x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{-3x}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{2e^{-\frac{x}{2}}, e^{-x}\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^{-3x}$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$10A_1 e^{-3x} = e^{-3x}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{10} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{e^{-3x}}{10}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^{-x} + 2c_2 e^{-\frac{x}{2}}) + \left(\frac{e^{-3x}}{10} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-x} + 2c_2 e^{-\frac{x}{2}} + \frac{e^{-3x}}{10} \quad (1)$$




Figure 349: Slope field plot

Verification of solutions

$$y = c_1 e^{-x} + 2c_2 e^{-\frac{x}{2}} + \frac{e^{-3x}}{10}$$

Verified OK.

11.5.3 Maple step by step solution

Let's solve

$$2y'' + 3y' + y = e^{-3x}$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{3y'}{2} - \frac{y}{2} + \frac{e^{-3x}}{2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{3y'}{2} + \frac{y}{2} = \frac{e^{-3x}}{2}$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + \frac{3}{2}r + \frac{1}{2} = 0$$

- Factor the characteristic polynomial

$$\frac{(r+1)(2r+1)}{2} = 0$$

- Roots of the characteristic polynomial

$$r = \left(-1, -\frac{1}{2}\right)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{-\frac{x}{2}}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{-x} + c_2 e^{-\frac{x}{2}} + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \frac{e^{-3x}}{2} \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-x} & e^{-\frac{x}{2}} \\ -e^{-x} & -\frac{e^{-\frac{x}{2}}}{2} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = \frac{e^{-\frac{3x}{2}}}{2}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -e^{-x} \left(\int e^{-2x} dx \right) + e^{-\frac{x}{2}} \left(\int e^{-\frac{5x}{2}} dx \right)$$

- Compute integrals

$$y_p(x) = \frac{e^{-3x}}{10}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{-x} + c_2 e^{-\frac{x}{2}} + \frac{e^{-3x}}{10}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
<- double symmetry of the form [xi=0, eta=F(x)] successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
dsolve(2*diff(y(x),x$2)+3*diff(y(x),x)+y(x)=exp(-3*x),y(x), singsol=all)
```

$$y(x) = -2c_1 e^{-x} + \frac{e^{-3x}}{10} + e^{-\frac{x}{2}} c_2$$

✓ Solution by Mathematica

Time used: 0.11 (sec). Leaf size: 45

```
DSolve[y''[x]+3*y'[x]+y[x]==Exp[-3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-3x} + c_1 e^{-\frac{1}{2}(3+\sqrt{5})x} + c_2 e^{\frac{1}{2}(\sqrt{5}-3)x}$$

11.6 problem 1(f)

- 11.6.1 Solving as second order linear constant coeff ode 1886
- 11.6.2 Solving using Kovacic algorithm 1890
- 11.6.3 Maple step by step solution 1895

Internal problem ID [6322]

Internal file name [OUTPUT/5570_Sunday_June_05_2022_03_43_47_PM_29822636/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.3. THE METHOD OF VARIATION OF PARAMETERS. Page 71

Problem number: 1(f).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' - 3y' + 2y = \frac{1}{1 + e^{-x}}$$

11.6.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = -3, C = 2, f(x) = \frac{1}{1+e^{-x}}$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 3y' + 2y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -3, C = 2$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 3\lambda e^{\lambda x} + 2e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 3\lambda + 2 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -3, C = 2$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{3}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-3^2 - (4)(1)(2)} \\ &= \frac{3}{2} \pm \frac{1}{2} \end{aligned}$$

Hence

$$\lambda_1 = \frac{3}{2} + \frac{1}{2}$$

$$\lambda_2 = \frac{3}{2} - \frac{1}{2}$$

Which simplifies to

$$\lambda_1 = 2$$

$$\lambda_2 = 1$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(2)x} + c_2 e^{(1)x}$$

Or

$$y = c_1 e^{2x} + c_2 e^x$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^{2x} + c_2 e^x$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = e^{2x}$$

$$y_2 = e^x$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^{2x} & e^x \\ \frac{d}{dx}(e^{2x}) & \frac{d}{dx}(e^x) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^{2x} & e^x \\ 2e^{2x} & e^x \end{vmatrix}$$

Therefore

$$W = (e^{2x})(e^x) - (e^x)(2e^{2x})$$

Which simplifies to

$$W = -e^{2x} e^x$$

Which simplifies to

$$W = -e^{3x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{e^x}{\frac{1+e^{-x}}{-e^{3x}}} dx$$

Which simplifies to

$$u_1 = - \int -\frac{e^{-2x}}{1+e^{-x}} dx$$

Hence

$$u_1 = \ln(1+e^x) - e^{-x} - \ln(e^x)$$

And Eq. (3) becomes

$$u_2 = \int \frac{e^{2x}}{\frac{1+e^{-x}}{-e^{3x}}} dx$$

Which simplifies to

$$u_2 = \int -\frac{e^{-x}}{1+e^{-x}} dx$$

Hence

$$u_2 = \ln(1+e^{-x})$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = (\ln(1+e^x) - e^{-x} - \ln(e^x)) e^{2x} + e^x \ln(1+e^{-x})$$

Which simplifies to

$$y_p(x) = e^x (\ln(1+e^x) e^x - e^x \ln(e^x) + \ln(1+e^{-x}) - 1)$$

Therefore the general solution is

$$\begin{aligned}
 y &= y_h + y_p \\
 &= (c_1 e^{2x} + c_2 e^x) + (e^x (\ln(1 + e^x) e^x - e^x \ln(e^x) + \ln(1 + e^{-x}) - 1))
 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{2x} + c_2 e^x + e^x (\ln(1 + e^x) e^x - e^x \ln(e^x) + \ln(1 + e^{-x}) - 1) \quad (1)$$

Verification of solutions

$$y = c_1 e^{2x} + c_2 e^x + e^x (\ln(1 + e^x) e^x - e^x \ln(e^x) + \ln(1 + e^{-x}) - 1)$$

Verified OK.

11.6.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 3y' + 2y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned}
 A &= 1 \\
 B &= -3 \\
 C &= 2
 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned}
 r &= \frac{s}{t} \\
 &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}
 \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{1}{4} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 1 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \frac{z(x)}{4} \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 280: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{1}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{x}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-3}{1} dx} \\ &= z_1 e^{\frac{3x}{2}} \\ &= z_1 \left(e^{\frac{3x}{2}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = e^x$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-3}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{3x}}{(y_1)^2} dx \\ &= y_1(e^x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1(e^x) + c_2(e^x(e^x)) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 3y' + 2y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1e^x + c_2e^{2x}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1y_1 + u_2y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = e^x$$

$$y_2 = e^{2x}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^x & e^{2x} \\ \frac{d}{dx}(e^x) & \frac{d}{dx}(e^{2x}) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^x & e^{2x} \\ e^x & 2e^{2x} \end{vmatrix}$$

Therefore

$$W = (e^x)(2e^{2x}) - (e^{2x})(e^x)$$

Which simplifies to

$$W = e^{2x}e^x$$

Which simplifies to

$$W = e^{3x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{e^{2x}}{\frac{1+e^{-x}}{e^{3x}}} dx$$

Which simplifies to

$$u_1 = - \int \frac{e^{-x}}{1+e^{-x}} dx$$

Hence

$$u_1 = \ln(1+e^{-x})$$

And Eq. (3) becomes

$$u_2 = \int \frac{e^x}{\frac{1+e^{-x}}{e^{3x}}} dx$$

Which simplifies to

$$u_2 = \int \frac{e^{-2x}}{1+e^{-x}} dx$$

Hence

$$u_2 = \ln(1+e^x) - e^{-x} - \ln(e^x)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = (\ln(1+e^x) - e^{-x} - \ln(e^x))e^{2x} + e^x \ln(1+e^{-x})$$

Which simplifies to

$$y_p(x) = e^x (\ln(1 + e^x) e^x - e^x \ln(e^x) + \ln(1 + e^{-x}) - 1)$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^x + c_2 e^{2x}) + (e^x (\ln(1 + e^x) e^x - e^x \ln(e^x) + \ln(1 + e^{-x}) - 1)) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^x + c_2 e^{2x} + e^x (\ln(1 + e^x) e^x - e^x \ln(e^x) + \ln(1 + e^{-x}) - 1) \quad (1)$$

Verification of solutions

$$y = c_1 e^x + c_2 e^{2x} + e^x (\ln(1 + e^x) e^x - e^x \ln(e^x) + \ln(1 + e^{-x}) - 1)$$

Verified OK.

11.6.3 Maple step by step solution

Let's solve

$$y'' - 3y' + 2y = \frac{1}{1+e^{-x}}$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - 3r + 2 = 0$$

- Factor the characteristic polynomial

$$(r - 1)(r - 2) = 0$$

- Roots of the characteristic polynomial

$$r = (1, 2)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^x$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{2x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$
- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^x + c_2 e^{2x} + y_p(x)$$
- Find a particular solution $y_p(x)$ of the ODE
 - Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \frac{1}{1+e^{-x}} \right]$$
 - Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^x & e^{2x} \\ e^x & 2e^{2x} \end{bmatrix}$$
 - Compute Wronskian

$$W(y_1(x), y_2(x)) = e^{3x}$$
 - Substitute functions into equation for $y_p(x)$

$$y_p(x) = -e^x \left(\int \frac{e^{-x}}{1+e^{-x}} dx \right) + e^{2x} \left(\int \frac{e^{-2x}}{1+e^{-x}} dx \right)$$
 - Compute integrals

$$y_p(x) = e^x (\ln(1 + e^x) e^x - e^x \ln(e^x) + \ln(1 + e^{-x}) - 1)$$
- Substitute particular solution into general solution to ODE

$$y = c_1 e^x + c_2 e^{2x} + e^x (\ln(1 + e^x) e^x - e^x \ln(e^x) + \ln(1 + e^{-x}) - 1)$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
<- double symmetry of the form [xi=0, eta=F(x)] successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
dsolve(diff(y(x),x$2)-3*diff(y(x),x)+2*y(x)=1/(1+exp(-x)),y(x), singsol=all)
```

$$y(x) = (\ln(e^x + 1)(e^x + 1) + (-e^x - 1)\ln(e^x) + e^x c_1 + c_2 - 1)e^x$$

✓ Solution by Mathematica

Time used: 0.086 (sec). Leaf size: 34

```
DSolve[y''[x]-3*y'[x]+2*y[x]==1/(1+Exp[-x]),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x(2(e^x + 1) \operatorname{arctanh}(2e^x + 1) + c_2 e^x - 1 + c_1)$$

11.7 problem 2(a)

11.7.1 Solving as second order linear constant coeff ode	1898
11.7.2 Solving using Kovacic algorithm	1902
11.7.3 Maple step by step solution	1908

Internal problem ID [6323]

Internal file name [OUTPUT/5571_Sunday_June_05_2022_03_43_48_PM_87300387/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.3. THE METHOD OF VARIATION OF PARAMETERS. Page 71

Problem number: 2(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + y = \sec(x)$$

11.7.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 1, f(x) = \sec(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\lambda_1 = +i$$

$$\lambda_2 = -i$$

Which simplifies to

$$\lambda_1 = i$$

$$\lambda_2 = -i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (\cos(x) c_1 + c_2 \sin(x))$$

Or

$$y = \cos(x) c_1 + c_2 \sin(x)$$

Therefore the homogeneous solution y_h is

$$y_h = \cos(x) c_1 + c_2 \sin(x)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \cos(x)$$

$$y_2 = \sin(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ \frac{d}{dx}(\cos(x)) & \frac{d}{dx}(\sin(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{vmatrix}$$

Therefore

$$W = (\cos(x))(\cos(x)) - (\sin(x))(-\sin(x))$$

Which simplifies to

$$W = \cos(x)^2 + \sin(x)^2$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\sin(x) \sec(x)}{1} dx$$

Which simplifies to

$$u_1 = - \int \tan(x) dx$$

Hence

$$u_1 = \ln(\cos(x))$$

And Eq. (3) becomes

$$u_2 = \int \frac{\sec(x) \cos(x)}{1} dx$$

Which simplifies to

$$u_2 = \int 1 dx$$

Hence

$$u_2 = x$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \ln(\cos(x)) \cos(x) + \sin(x) x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (\cos(x) c_1 + c_2 \sin(x)) + (\ln(\cos(x)) \cos(x) + \sin(x) x) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \cos(x) c_1 + c_2 \sin(x) + \ln(\cos(x)) \cos(x) + \sin(x) x \quad (1)$$




Figure 350: Slope field plot

Verification of solutions

$$y = \cos(x) c_1 + c_2 \sin(x) + \ln(\cos(x)) \cos(x) + \sin(x) x$$

Verified OK.

11.7.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 0 \quad (3)$$

$$C = 1$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 282: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned} y_1 &= z_1 \\ &= \cos(x) \end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x)) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1(\cos(x)) + c_2(\cos(x) (\tan(x))) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = \cos(x) c_1 + c_2 \sin(x)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of

parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \cos(x)$$

$$y_2 = \sin(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ \frac{d}{dx}(\cos(x)) & \frac{d}{dx}(\sin(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{vmatrix}$$

Therefore

$$W = (\cos(x))(\cos(x)) - (\sin(x))(-\sin(x))$$

Which simplifies to

$$W = \cos(x)^2 + \sin(x)^2$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\sin(x) \sec(x)}{1} dx$$

Which simplifies to

$$u_1 = - \int \tan(x) dx$$

Hence

$$u_1 = \ln(\cos(x))$$

And Eq. (3) becomes

$$u_2 = \int \frac{\sec(x) \cos(x)}{1} dx$$

Which simplifies to

$$u_2 = \int 1 dx$$

Hence

$$u_2 = x$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \ln(\cos(x)) \cos(x) + \sin(x) x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (\cos(x) c_1 + c_2 \sin(x)) + (\ln(\cos(x)) \cos(x) + \sin(x) x) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \cos(x) c_1 + c_2 \sin(x) + \ln(\cos(x)) \cos(x) + \sin(x) x \quad (1)$$




Figure 351: Slope field plot

Verification of solutions

$$y = \cos(x) c_1 + c_2 \sin(x) + \ln(\cos(x)) \cos(x) + \sin(x) x$$

Verified OK.

11.7.3 Maple step by step solution

Let's solve

$$y'' + y = \sec(x)$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (-I, I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = \cos(x) c_1 + c_2 \sin(x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \sec(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\cos(x) \left(\int \tan(x) dx \right) + \sin(x) \left(\int 1 dx \right)$$

- Compute integrals

$$y_p(x) = \ln(\cos(x)) \cos(x) + \sin(x) x$$

- Substitute particular solution into general solution to ODE

$$y = \cos(x) c_1 + c_2 \sin(x) + \ln(\cos(x)) \cos(x) + \sin(x) x$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 22

```
dsolve(diff(y(x),x$2)+y(x)=sec(x),y(x), singsol=all)
```

$$y(x) = -\ln(\sec(x)) \cos(x) + \cos(x) c_1 + (x + c_2) \sin(x)$$

✓ Solution by Mathematica

Time used: 0.035 (sec). Leaf size: 22

```
DSolve[y''[x]+y[x]==Sec[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow (x + c_2) \sin(x) + \cos(x)(\log(\cos(x)) + c_1)$$

11.8 problem 2(b)

- 11.8.1 Solving as second order linear constant coeff ode 1911
- 11.8.2 Solving using Kovacic algorithm 1916
- 11.8.3 Maple step by step solution 1922

Internal problem ID [6324]

Internal file name [OUTPUT/5572_Sunday_June_05_2022_03_43_50_PM_43269960/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.3. THE METHOD OF VARIATION OF PARAMETERS. Page 71

Problem number: 2(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + y = \cot(x)^2$$

11.8.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 1, f(x) = \frac{1}{\tan(x)^2}$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +i \\ \lambda_2 &= -i \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= i \\ \lambda_2 &= -i \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (\cos(x) c_1 + c_2 \sin(x))$$

Or

$$y = \cos(x) c_1 + c_2 \sin(x)$$

Therefore the homogeneous solution y_h is

$$y_h = \cos(x) c_1 + c_2 \sin(x)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \cos(x)$$

$$y_2 = \sin(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ \frac{d}{dx}(\cos(x)) & \frac{d}{dx}(\sin(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{vmatrix}$$

Therefore

$$W = (\cos(x))(\cos(x)) - (\sin(x))(-\sin(x))$$

Which simplifies to

$$W = \cos(x)^2 + \sin(x)^2$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\frac{\sin(x)}{\tan(x)^2}}{1} dx$$

Which simplifies to

$$u_1 = - \int \cos(x) \cot(x) dx$$

Hence

$$u_1 = -\cos(x) - \ln(\csc(x) - \cot(x))$$

And Eq. (3) becomes

$$u_2 = \int \frac{\frac{\cos(x)}{\tan(x)^2}}{1} dx$$

Which simplifies to

$$u_2 = \int \cos(x) \cot(x)^2 dx$$

Hence

$$u_2 = -\frac{\cos(x)^4}{\sin(x)} - (2 + \cos(x)^2) \sin(x)$$

Which simplifies to

$$u_1 = -\cos(x) - \ln(\csc(x) - \cot(x))$$

$$u_2 = \cos(x) \cot(x) - 2 \csc(x)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = (-\cos(x) - \ln(\csc(x) - \cot(x)))\cos(x) + (\cos(x)\cot(x) - 2\csc(x))\sin(x)$$

Which simplifies to

$$y_p(x) = -\cos(x)\ln(\csc(x) - \cot(x)) - 2$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (\cos(x)c_1 + c_2\sin(x)) + (-\cos(x)\ln(\csc(x) - \cot(x)) - 2) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \cos(x)c_1 + c_2\sin(x) - \cos(x)\ln(\csc(x) - \cot(x)) - 2 \quad (1)$$




Figure 352: Slope field plot

Verification of solutions

$$y = \cos(x)c_1 + c_2\sin(x) - \cos(x)\ln(\csc(x) - \cot(x)) - 2$$

Verified OK.

11.8.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 1 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 284: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}y_1 &= z_1 \\ &= \cos(x)\end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned}y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x))\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(x)) + c_2 (\cos(x) (\tan(x)))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = \cos(x) c_1 + c_2 \sin(x)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \cos(x)$$

$$y_2 = \sin(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ \frac{d}{dx}(\cos(x)) & \frac{d}{dx}(\sin(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{vmatrix}$$

Therefore

$$W = (\cos(x))(\cos(x)) - (\sin(x))(-\sin(x))$$

Which simplifies to

$$W = \cos(x)^2 + \sin(x)^2$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\sin(x) \cot(x)^2}{1} dx$$

Which simplifies to

$$u_1 = - \int \cos(x) \cot(x) dx$$

Hence

$$u_1 = - \cos(x) - \ln(\csc(x) - \cot(x))$$

And Eq. (3) becomes

$$u_2 = \int \frac{\cos(x) \cot(x)^2}{1} dx$$

Which simplifies to

$$u_2 = \int \cos(x) \cot(x)^2 dx$$

Hence

$$u_2 = - \frac{\cos(x)^4}{\sin(x)} - (2 + \cos(x)^2) \sin(x)$$

Which simplifies to

$$u_1 = - \cos(x) - \ln(\csc(x) - \cot(x))$$

$$u_2 = \cos(x) \cot(x) - 2 \csc(x)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = (- \cos(x) - \ln(\csc(x) - \cot(x))) \cos(x) + (\cos(x) \cot(x) - 2 \csc(x)) \sin(x)$$

Which simplifies to

$$y_p(x) = - \cos(x) \ln(\csc(x) - \cot(x)) - 2$$

Therefore the general solution is

$$\begin{aligned}y &= y_h + y_p \\ &= (\cos(x) c_1 + c_2 \sin(x)) + (-\cos(x) \ln(\csc(x) - \cot(x)) - 2)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \cos(x) c_1 + c_2 \sin(x) - \cos(x) \ln(\csc(x) - \cot(x)) - 2 \quad (1)$$




Figure 353: Slope field plot

Verification of solutions

$$y = \cos(x) c_1 + c_2 \sin(x) - \cos(x) \ln(\csc(x) - \cot(x)) - 2$$

Verified OK.

11.8.3 Maple step by step solution

Let's solve

$$y'' + y = \frac{1}{\tan(x)^2}$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (-I, I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = \cos(x) c_1 + c_2 \sin(x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \frac{1}{\tan(x)^2} \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\cos(x) \left(\int \cos(x) \cot(x) dx \right) + \sin(x) \left(\int \cos(x) \cot(x)^2 dx \right)$$

- Compute integrals

$$y_p(x) = -\cos(x) \ln(\csc(x) - \cot(x)) - 2$$

- Substitute particular solution into general solution to ODE

$$y = \cos(x) c_1 + c_2 \sin(x) - \cos(x) \ln(\csc(x) - \cot(x)) - 2$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 26

```
dsolve(diff(y(x), x$2)+y(x)=cot(x)^2, y(x), singsol=all)
```

$$y(x) = \sin(x) c_2 + \cos(x) c_1 - 2 - \cos(x) \ln(\csc(x) - \cot(x))$$

✓ Solution by Mathematica

Time used: 0.075 (sec). Leaf size: 34

```
DSolve[y''[x]+y[x]==Cot[x]^2, y[x], x, IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 \sin(x) + \cos(x) \left(-\log\left(\sin\left(\frac{x}{2}\right)\right) + \log\left(\cos\left(\frac{x}{2}\right)\right) + c_1 \right) - 2$$

11.9 problem 2(c)

11.9.1 Solving as second order linear constant coeff ode	1924
11.9.2 Solving using Kovacic algorithm	1929
11.9.3 Maple step by step solution	1935

Internal problem ID [6325]

Internal file name [OUTPUT/5573_Sunday_June_05_2022_03_43_51_PM_1270245/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.3. THE METHOD OF VARIATION OF PARAMETERS. Page 71

Problem number: 2(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + y = \cot(2x)$$

11.9.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 1, f(x) = \cot(2x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\lambda_1 = +i$$

$$\lambda_2 = -i$$

Which simplifies to

$$\lambda_1 = i$$

$$\lambda_2 = -i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (\cos(x) c_1 + c_2 \sin(x))$$

Or

$$y = \cos(x) c_1 + c_2 \sin(x)$$

Therefore the homogeneous solution y_h is

$$y_h = \cos(x) c_1 + c_2 \sin(x)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \cos(x)$$

$$y_2 = \sin(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ \frac{d}{dx}(\cos(x)) & \frac{d}{dx}(\sin(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{vmatrix}$$

Therefore

$$W = (\cos(x))(\cos(x)) - (\sin(x))(-\sin(x))$$

Which simplifies to

$$W = \cos(x)^2 + \sin(x)^2$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\sin(x) \cot(2x)}{1} dx$$

Which simplifies to

$$u_1 = - \int \left(\cos(x) - \frac{\sec(x)}{2} \right) dx$$

Hence

$$u_1 = -\sin(x) + \frac{\ln(\sec(x) + \tan(x))}{2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\cos(x) \cot(2x)}{1} dx$$

Which simplifies to

$$u_2 = \int \left(\cos(x) \cot(x) - \frac{\csc(x)}{2} \right) dx$$

Hence

$$u_2 = \cos(x) + \ln(\csc(x) - \cot(x)) + \frac{\ln(\csc(x) + \cot(x))}{2}$$

Therefore the particular solution, from equation (1) is

$$\begin{aligned} y_p(x) = & \left(-\sin(x) + \frac{\ln(\sec(x) + \tan(x))}{2} \right) \cos(x) \\ & + \left(\cos(x) + \ln(\csc(x) - \cot(x)) + \frac{\ln(\csc(x) + \cot(x))}{2} \right) \sin(x) \end{aligned}$$

Which simplifies to

$$y_p(x) = \frac{\cos(x) \ln(\sec(x) + \tan(x))}{2} + \sin(x) \ln(\csc(x) - \cot(x)) + \frac{\sin(x) \ln(\csc(x) + \cot(x))}{2}$$

Therefore the general solution is

$$y = y_h + y_p$$

$$= (\cos(x) c_1 + c_2 \sin(x)) + \left(\frac{\cos(x) \ln(\sec(x) + \tan(x))}{2} + \sin(x) \ln(\csc(x) - \cot(x)) + \frac{\sin(x) \ln(\csc(x) + \cot(x))}{2} \right)$$

Summary

The solution(s) found are the following

$$y = \cos(x) c_1 + c_2 \sin(x) + \frac{\cos(x) \ln(\sec(x) + \tan(x))}{2} + \sin(x) \ln(\csc(x) - \cot(x)) + \frac{\sin(x) \ln(\csc(x) + \cot(x))}{2} \quad (1)$$




Figure 354: Slope field plot

Verification of solutions

$$y = \cos(x) c_1 + c_2 \sin(x) + \frac{\cos(x) \ln(\sec(x) + \tan(x))}{2} \\ + \sin(x) \ln(\csc(x) - \cot(x)) + \frac{\sin(x) \ln(\csc(x) + \cot(x))}{2}$$

Verified OK.

11.9.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1 \\ B = 0 \\ C = 1 \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} \\ = \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \tag{6}$$

Comparing the above to (5) shows that

$$s = -1 \\ t = 1$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 286: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned} y_1 &= z_1 \\ &= \cos(x) \end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x)) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(x)) + c_2 (\cos(x) (\tan(x))) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = \cos(x) c_1 + c_2 \sin(x)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \cos(x)$$

$$y_2 = \sin(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ \frac{d}{dx}(\cos(x)) & \frac{d}{dx}(\sin(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{vmatrix}$$

Therefore

$$W = (\cos(x))(\cos(x)) - (\sin(x))(-\sin(x))$$

Which simplifies to

$$W = \cos(x)^2 + \sin(x)^2$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\sin(x) \cot(2x)}{1} dx$$

Which simplifies to

$$u_1 = - \int \left(\cos(x) - \frac{\sec(x)}{2} \right) dx$$

Hence

$$u_1 = -\sin(x) + \frac{\ln(\sec(x) + \tan(x))}{2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\cos(x) \cot(2x)}{1} dx$$

Which simplifies to

$$u_2 = \int \left(\cos(x) \cot(x) - \frac{\csc(x)}{2} \right) dx$$

Hence

$$u_2 = \cos(x) + \ln(\csc(x) - \cot(x)) + \frac{\ln(\csc(x) + \cot(x))}{2}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(-\sin(x) + \frac{\ln(\sec(x) + \tan(x))}{2} \right) \cos(x) \\ + \left(\cos(x) + \ln(\csc(x) - \cot(x)) + \frac{\ln(\csc(x) + \cot(x))}{2} \right) \sin(x)$$

Which simplifies to

$$y_p(x) = \frac{\cos(x) \ln(\sec(x) + \tan(x))}{2} + \sin(x) \ln(\csc(x) - \cot(x)) + \frac{\sin(x) \ln(\csc(x) + \cot(x))}{2}$$

Therefore the general solution is

$$y = y_h + y_p = (\cos(x) c_1 + c_2 \sin(x)) + \left(\frac{\cos(x) \ln(\sec(x) + \tan(x))}{2} + \sin(x) \ln(\csc(x) - \cot(x)) + \frac{\sin(x) \ln(\csc(x) + \cot(x))}{2} \right)$$

Summary

The solution(s) found are the following

$$y = \cos(x) c_1 + c_2 \sin(x) + \frac{\cos(x) \ln(\sec(x) + \tan(x))}{2} + \sin(x) \ln(\csc(x) - \cot(x)) + \frac{\sin(x) \ln(\csc(x) + \cot(x))}{2} \quad (1)$$




Figure 355: Slope field plot

Verification of solutions

$$y = \cos(x) c_1 + c_2 \sin(x) + \frac{\cos(x) \ln(\sec(x) + \tan(x))}{2} \\ + \sin(x) \ln(\csc(x) - \cot(x)) + \frac{\sin(x) \ln(\csc(x) + \cot(x))}{2}$$

Verified OK.

11.9.3 Maple step by step solution

Let's solve

$$y'' + y = \cot(2x)$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (-I, I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = \cos(x) c_1 + c_2 \sin(x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \cot(2x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix}$$

- o Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$

- o Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{\cos(x)(\int(2\cos(x)-\sec(x))dx)}{2} + \frac{\sin(x)(\int(2\cos(x)\cot(x)-\csc(x))dx)}{2}$$

- o Compute integrals

$$y_p(x) = \frac{\cos(x)\ln(\sec(x)+\tan(x))}{2} + \sin(x)\ln(\csc(x) - \cot(x)) + \frac{\sin(x)\ln(\csc(x)+\cot(x))}{2}$$

- Substitute particular solution into general solution to ODE

$$y = \cos(x)c_1 + c_2\sin(x) + \frac{\cos(x)\ln(\sec(x)+\tan(x))}{2} + \sin(x)\ln(\csc(x) - \cot(x)) + \frac{\sin(x)\ln(\csc(x)+\cot(x))}{2}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 35

```
dsolve(diff(y(x), x$2)+y(x)=cot(2*x), y(x), singsol=all)
```

$$y(x) = \sin(x)c_2 + \cos(x)c_1 + \frac{\sin(x)\ln(\csc(x) - \cot(x))}{2} + \frac{\cos(x)\ln(\sec(x) + \tan(x))}{2}$$

✓ Solution by Mathematica

Time used: 0.082 (sec). Leaf size: 46

```
DSolve[y''[x]+y[x]==Cot[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} \left(\cos(x) \operatorname{arctanh}(\sin(x)) + 2c_1 \cos(x) \right. \\ \left. + \sin(x) \left(\log \left(\sin \left(\frac{x}{2} \right) \right) - \log \left(\cos \left(\frac{x}{2} \right) \right) + 2c_2 \right) \right)$$

11.10 problem 2(d)

11.10.1 Solving as second order linear constant coeff ode	1938
11.10.2 Solving using Kovacic algorithm	1942
11.10.3 Maple step by step solution	1946

Internal problem ID [6326]

Internal file name [OUTPUT/5574_Sunday_June_05_2022_03_43_53_PM_39095857/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.3. THE METHOD OF VARIATION OF PARAMETERS. Page 71

Problem number: 2(d).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + y = \cos(x)x$$

11.10.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 1, f(x) = \cos(x)x$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\lambda_1 = +i$$

$$\lambda_2 = -i$$

Which simplifies to

$$\lambda_1 = i$$

$$\lambda_2 = -i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (\cos(x) c_1 + c_2 \sin(x))$$

Or

$$y = \cos(x) c_1 + c_2 \sin(x)$$

Therefore the homogeneous solution y_h is

$$y_h = \cos(x) c_1 + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\cos(x) x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x) x, \sin(x) x, \cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since $\cos(x)$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x^2 \sin(x), \cos(x) x, \cos(x) x^2, \sin(x) x\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x^2 \sin(x) + A_2 \cos(x) x + A_3 \cos(x) x^2 + A_4 \sin(x) x$$

The unknowns $\{A_1, A_2, A_3, A_4\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$\begin{aligned} 2A_1 \sin(x) + 4A_1 x \cos(x) - 2A_2 \sin(x) - 4A_3 \sin(x) x + 2A_3 \cos(x) + 2A_4 \cos(x) \\ = \cos(x) x \end{aligned}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{4}, A_2 = \frac{1}{4}, A_3 = 0, A_4 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{x^2 \sin(x)}{4} + \frac{\cos(x) x}{4}$$

Therefore the general solution is

$$\begin{aligned}y &= y_h + y_p \\ &= (\cos(x) c_1 + c_2 \sin(x)) + \left(\frac{x^2 \sin(x)}{4} + \frac{\cos(x) x}{4} \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \cos(x) c_1 + c_2 \sin(x) + \frac{x^2 \sin(x)}{4} + \frac{\cos(x) x}{4} \quad (1)$$




Figure 356: Slope field plot

Verification of solutions

$$y = \cos(x) c_1 + c_2 \sin(x) + \frac{x^2 \sin(x)}{4} + \frac{\cos(x) x}{4}$$

Verified OK.

11.10.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 1 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 288: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}y_1 &= z_1 \\ &= \cos(x)\end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned}y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x))\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(x)) + c_2 (\cos(x) (\tan(x)))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = \cos(x) c_1 + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\cos(x) x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x) x, \sin(x) x, \cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since $\cos(x)$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x^2 \sin(x), \cos(x) x, \cos(x) x^2, \sin(x) x\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x^2 \sin(x) + A_2 \cos(x) x + A_3 \cos(x) x^2 + A_4 \sin(x) x$$

The unknowns $\{A_1, A_2, A_3, A_4\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$\begin{aligned} 2A_1 \sin(x) + 4A_1 x \cos(x) - 2A_2 \sin(x) - 4A_3 \sin(x) x + 2A_3 \cos(x) + 2A_4 \cos(x) \\ = \cos(x) x \end{aligned}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{4}, A_2 = \frac{1}{4}, A_3 = 0, A_4 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{x^2 \sin(x)}{4} + \frac{\cos(x) x}{4}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (\cos(x) c_1 + c_2 \sin(x)) + \left(\frac{x^2 \sin(x)}{4} + \frac{\cos(x) x}{4} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \cos(x) c_1 + c_2 \sin(x) + \frac{x^2 \sin(x)}{4} + \frac{\cos(x) x}{4} \quad (1)$$




Figure 357: Slope field plot

Verification of solutions

$$y = \cos(x) c_1 + c_2 \sin(x) + \frac{x^2 \sin(x)}{4} + \frac{\cos(x) x}{4}$$

Verified OK.

11.10.3 Maple step by step solution

Let's solve

$$y'' + y = \cos(x) x$$

- Highest derivative means the order of the ODE is 2
- y''
- Characteristic polynomial of homogeneous ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (-I, I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = \cos(x) c_1 + c_2 \sin(x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x), y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x), y_2(x))} dx \right), f(x) = \cos(x)x \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{\cos(x) \left(\int x \sin(2x) dx \right)}{2} + \sin(x) \left(\int \cos(x)^2 x dx \right)$$

- Compute integrals

$$y_p(x) = \frac{x^2 \sin(x)}{4} - \frac{\sin(x)}{4} + \frac{\cos(x)x}{4}$$

- Substitute particular solution into general solution to ODE

$$y = \cos(x) c_1 + c_2 \sin(x) + \frac{x^2 \sin(x)}{4} - \frac{\sin(x)}{4} + \frac{\cos(x)x}{4}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(diff(y(x),x$2)+y(x)=x*cos(x),y(x), singsol=all)
```

$$y(x) = \frac{(x^2 + 4c_2 - 1) \sin(x)}{4} + \frac{\cos(x)(4c_1 + x)}{4}$$

✓ Solution by Mathematica

Time used: 0.053 (sec). Leaf size: 34

```
DSolve[y''[x]+y[x]==x*Cos[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{8}((2x^2 - 1 + 8c_2) \sin(x) + 2(x + 4c_1) \cos(x))$$

11.11 problem 2(e)

11.11.1 Solving as second order linear constant coeff ode	1949
11.11.2 Solving using Kovacic algorithm	1954
11.11.3 Maple step by step solution	1959

Internal problem ID [6327]

Internal file name [OUTPUT/5575_Sunday_June_05_2022_03_43_54_PM_39686071/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.3. THE METHOD OF VARIATION OF PARAMETERS. Page 71

Problem number: 2(e).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + y = \tan(x)$$

11.11.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 1, f(x) = \tan(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\lambda_1 = +i$$

$$\lambda_2 = -i$$

Which simplifies to

$$\lambda_1 = i$$

$$\lambda_2 = -i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (\cos(x) c_1 + c_2 \sin(x))$$

Or

$$y = \cos(x) c_1 + c_2 \sin(x)$$

Therefore the homogeneous solution y_h is

$$y_h = \cos(x) c_1 + c_2 \sin(x)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \cos(x)$$

$$y_2 = \sin(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ \frac{d}{dx}(\cos(x)) & \frac{d}{dx}(\sin(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{vmatrix}$$

Therefore

$$W = (\cos(x))(\cos(x)) - (\sin(x))(-\sin(x))$$

Which simplifies to

$$W = \cos(x)^2 + \sin(x)^2$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\sin(x) \tan(x)}{1} dx$$

Which simplifies to

$$u_1 = - \int \sin(x) \tan(x) dx$$

Hence

$$u_1 = \sin(x) - \ln(\sec(x) + \tan(x))$$

And Eq. (3) becomes

$$u_2 = \int \frac{\tan(x) \cos(x)}{1} dx$$

Which simplifies to

$$u_2 = \int \sin(x) dx$$

Hence

$$u_2 = -\cos(x)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = (\sin(x) - \ln(\sec(x) + \tan(x))) \cos(x) - \cos(x) \sin(x)$$

Which simplifies to

$$y_p(x) = -\cos(x) \ln(\sec(x) + \tan(x))$$

Therefore the general solution is

$$\begin{aligned}y &= y_h + y_p \\ &= (\cos(x) c_1 + c_2 \sin(x)) + (-\cos(x) \ln(\sec(x) + \tan(x)))\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \cos(x) c_1 + c_2 \sin(x) - \cos(x) \ln(\sec(x) + \tan(x)) \quad (1)$$




Figure 358: Slope field plot

Verification of solutions

$$y = \cos(x) c_1 + c_2 \sin(x) - \cos(x) \ln(\sec(x) + \tan(x))$$

Verified OK.

11.11.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 1 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 290: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}y_1 &= z_1 \\ &= \cos(x)\end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned}y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x))\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(x)) + c_2 (\cos(x) (\tan(x)))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = \cos(x) c_1 + c_2 \sin(x)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \cos(x)$$

$$y_2 = \sin(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ \frac{d}{dx}(\cos(x)) & \frac{d}{dx}(\sin(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{vmatrix}$$

Therefore

$$W = (\cos(x))(\cos(x)) - (\sin(x))(-\sin(x))$$

Which simplifies to

$$W = \cos(x)^2 + \sin(x)^2$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\sin(x) \tan(x)}{1} dx$$

Which simplifies to

$$u_1 = - \int \sin(x) \tan(x) dx$$

Hence

$$u_1 = \sin(x) - \ln(\sec(x) + \tan(x))$$

And Eq. (3) becomes

$$u_2 = \int \frac{\tan(x) \cos(x)}{1} dx$$

Which simplifies to

$$u_2 = \int \sin(x) dx$$

Hence

$$u_2 = -\cos(x)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = (\sin(x) - \ln(\sec(x) + \tan(x))) \cos(x) - \cos(x) \sin(x)$$

Which simplifies to

$$y_p(x) = -\cos(x) \ln(\sec(x) + \tan(x))$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (\cos(x) c_1 + c_2 \sin(x)) + (-\cos(x) \ln(\sec(x) + \tan(x))) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \cos(x) c_1 + c_2 \sin(x) - \cos(x) \ln(\sec(x) + \tan(x)) \quad (1)$$




Figure 359: Slope field plot

Verification of solutions

$$y = \cos(x) c_1 + c_2 \sin(x) - \cos(x) \ln(\sec(x) + \tan(x))$$

Verified OK.

11.11.3 Maple step by step solution

Let's solve

$$y'' + y = \tan(x)$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$
- Roots of the characteristic polynomial

$$r = (-I, I)$$
- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x)$$
- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(x)$$
- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$
- Substitute in solutions of the homogeneous ODE

$$y = \cos(x) c_1 + c_2 \sin(x) + y_p(x)$$
- Find a particular solution $y_p(x)$ of the ODE
 - Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x), y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x), y_2(x))} dx \right), f(x) = \tan(x) \right]$$
 - Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix}$$
 - Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$
 - Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\cos(x) \left(\int \sin(x) \tan(x) dx \right) + \sin(x) \left(\int \sin(x) dx \right)$$
 - Compute integrals

$$y_p(x) = -\cos(x) \ln(\sec(x) + \tan(x))$$
- Substitute particular solution into general solution to ODE

$$y = \cos(x) c_1 + c_2 \sin(x) - \cos(x) \ln(\sec(x) + \tan(x))$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(diff(y(x),x$2)+y(x)=tan(x),y(x), singsol=all)
```

$$y(x) = \sin(x) c_2 + \cos(x) c_1 - \cos(x) \ln(\sec(x) + \tan(x))$$

✓ Solution by Mathematica

Time used: 0.038 (sec). Leaf size: 23

```
DSolve[y''[x]+y[x]==Tan[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \cos(x)(-\operatorname{arctanh}(\sin(x))) + c_1 \cos(x) + c_2 \sin(x)$$

11.12 problem 2(f)

11.12.1 Solving as second order linear constant coeff ode	1962
11.12.2 Solving using Kovacic algorithm	1967
11.12.3 Maple step by step solution	1972

Internal problem ID [6328]

Internal file name [OUTPUT/5576_Sunday_June_05_2022_03_43_56_PM_78572068/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.3. THE METHOD OF VARIATION OF PARAMETERS. Page 71

Problem number: 2(f).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + y = \sec(x) \tan(x)$$

11.12.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 1, f(x) = \sec(x) \tan(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\lambda_1 = +i$$

$$\lambda_2 = -i$$

Which simplifies to

$$\lambda_1 = i$$

$$\lambda_2 = -i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (\cos(x) c_1 + c_2 \sin(x))$$

Or

$$y = \cos(x) c_1 + c_2 \sin(x)$$

Therefore the homogeneous solution y_h is

$$y_h = \cos(x) c_1 + c_2 \sin(x)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \cos(x)$$

$$y_2 = \sin(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ \frac{d}{dx}(\cos(x)) & \frac{d}{dx}(\sin(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{vmatrix}$$

Therefore

$$W = (\cos(x))(\cos(x)) - (\sin(x))(-\sin(x))$$

Which simplifies to

$$W = \cos(x)^2 + \sin(x)^2$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\sec(x) \sin(x) \tan(x)}{1} dx$$

Which simplifies to

$$u_1 = - \int \tan(x)^2 dx$$

Hence

$$u_1 = - \tan(x) + x$$

And Eq. (3) becomes

$$u_2 = \int \frac{\tan(x) \cos(x) \sec(x)}{1} dx$$

Which simplifies to

$$u_2 = \int \tan(x) dx$$

Hence

$$u_2 = - \ln(\cos(x))$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = (- \tan(x) + x) \cos(x) - \ln(\cos(x)) \sin(x)$$

Which simplifies to

$$y_p(x) = - \sin(x) + \cos(x) x - \ln(\cos(x)) \sin(x)$$

Therefore the general solution is

$$\begin{aligned}y &= y_h + y_p \\ &= (\cos(x) c_1 + c_2 \sin(x)) + (-\sin(x) + \cos(x) x - \ln(\cos(x)) \sin(x))\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \cos(x) c_1 + c_2 \sin(x) - \sin(x) + \cos(x) x - \ln(\cos(x)) \sin(x) \quad (1)$$




Figure 360: Slope field plot

Verification of solutions

$$y = \cos(x) c_1 + c_2 \sin(x) - \sin(x) + \cos(x) x - \ln(\cos(x)) \sin(x)$$

Verified OK.

11.12.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 1 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 292: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}y_1 &= z_1 \\ &= \cos(x)\end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned}y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x))\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(x)) + c_2 (\cos(x) (\tan(x)))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = \cos(x) c_1 + c_2 \sin(x)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \cos(x)$$

$$y_2 = \sin(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ \frac{d}{dx}(\cos(x)) & \frac{d}{dx}(\sin(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{vmatrix}$$

Therefore

$$W = (\cos(x))(\cos(x)) - (\sin(x))(-\sin(x))$$

Which simplifies to

$$W = \cos(x)^2 + \sin(x)^2$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\sec(x) \sin(x) \tan(x)}{1} dx$$

Which simplifies to

$$u_1 = - \int \tan(x)^2 dx$$

Hence

$$u_1 = - \tan(x) + x$$

And Eq. (3) becomes

$$u_2 = \int \frac{\tan(x) \cos(x) \sec(x)}{1} dx$$

Which simplifies to

$$u_2 = \int \tan(x) dx$$

Hence

$$u_2 = - \ln(\cos(x))$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = (- \tan(x) + x) \cos(x) - \ln(\cos(x)) \sin(x)$$

Which simplifies to

$$y_p(x) = - \sin(x) + \cos(x) x - \ln(\cos(x)) \sin(x)$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (\cos(x) c_1 + c_2 \sin(x)) + (- \sin(x) + \cos(x) x - \ln(\cos(x)) \sin(x)) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \cos(x) c_1 + c_2 \sin(x) - \sin(x) + \cos(x) x - \ln(\cos(x)) \sin(x) \quad (1)$$




Figure 361: Slope field plot

Verification of solutions

$$y = \cos(x) c_1 + c_2 \sin(x) - \sin(x) + \cos(x) x - \ln(\cos(x)) \sin(x)$$

Verified OK.

11.12.3 Maple step by step solution

Let's solve

$$y'' + y = \sec(x) \tan(x)$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$
- Roots of the characteristic polynomial

$$r = (-I, I)$$
- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x)$$
- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(x)$$
- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$
- Substitute in solutions of the homogeneous ODE

$$y = \cos(x) c_1 + c_2 \sin(x) + y_p(x)$$
- Find a particular solution $y_p(x)$ of the ODE
 - Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \sec(x) \tan(x) \right]$$
 - Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix}$$
 - Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$
 - Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\cos(x) \left(\int \tan(x)^2 dx \right) + \sin(x) \left(\int \tan(x) dx \right)$$
 - Compute integrals

$$y_p(x) = -\sin(x) + \cos(x) x - \ln(\cos(x)) \sin(x)$$
- Substitute particular solution into general solution to ODE

$$y = \cos(x) c_1 + c_2 \sin(x) - \sin(x) + \cos(x) x - \ln(\cos(x)) \sin(x)$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(diff(y(x),x$2)+y(x)=sec(x)*tan(x),y(x), singsol=all)
```

$$y(x) = \ln(\sec(x)) \sin(x) + (c_2 - 1) \sin(x) + \cos(x)(x + c_1)$$

✓ Solution by Mathematica

Time used: 0.04 (sec). Leaf size: 29

```
DSolve[y''[x]+y[x]==Sec[x]*Tan[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \cos(x) \arctan(\tan(x)) + c_1 \cos(x) + \sin(x)(-\log(\cos(x)) - 1 + c_2)$$

11.13 problem 2(g)

11.13.1 Solving as second order linear constant coeff ode	1975
11.13.2 Solving using Kovacic algorithm	1979
11.13.3 Maple step by step solution	1985

Internal problem ID [6329]

Internal file name [OUTPUT/5577_Sunday_June_05_2022_03_43_57_PM_17656438/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.3. THE METHOD OF VARIATION OF PARAMETERS. Page 71

Problem number: 2(g).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + y = \sec(x) \csc(x)$$

11.13.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 1, f(x) = \sec(x) \csc(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\lambda_1 = +i$$

$$\lambda_2 = -i$$

Which simplifies to

$$\lambda_1 = i$$

$$\lambda_2 = -i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (\cos(x) c_1 + c_2 \sin(x))$$

Or

$$y = \cos(x) c_1 + c_2 \sin(x)$$

Therefore the homogeneous solution y_h is

$$y_h = \cos(x) c_1 + c_2 \sin(x)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \cos(x)$$

$$y_2 = \sin(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ \frac{d}{dx}(\cos(x)) & \frac{d}{dx}(\sin(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{vmatrix}$$

Therefore

$$W = (\cos(x))(\cos(x)) - (\sin(x))(-\sin(x))$$

Which simplifies to

$$W = \cos(x)^2 + \sin(x)^2$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\sin(x) \sec(x) \csc(x)}{1} dx$$

Which simplifies to

$$u_1 = - \int \sec(x) dx$$

Hence

$$u_1 = - \ln(\sec(x) + \tan(x))$$

And Eq. (3) becomes

$$u_2 = \int \frac{\cos(x) \sec(x) \csc(x)}{1} dx$$

Which simplifies to

$$u_2 = \int \csc(x) dx$$

Hence

$$u_2 = - \ln(\csc(x) + \cot(x))$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = - \cos(x) \ln(\sec(x) + \tan(x)) - \sin(x) \ln(\csc(x) + \cot(x))$$

Therefore the general solution is

$$y = y_h + y_p$$

$$= (\cos(x) c_1 + c_2 \sin(x)) + (- \cos(x) \ln(\sec(x) + \tan(x)) - \sin(x) \ln(\csc(x) + \cot(x)))$$

Summary

The solution(s) found are the following

$$y = \cos(x) c_1 + c_2 \sin(x) - \cos(x) \ln(\sec(x) + \tan(x)) - \sin(x) \ln(\csc(x) + \cot(x))$$




Figure 362: Slope field plot

Verification of solutions

$$y = \cos(x) c_1 + c_2 \sin(x) - \cos(x) \ln(\sec(x) + \tan(x)) - \sin(x) \ln(\csc(x) + \cot(x))$$

Verified OK.

11.13.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 0 \tag{3}$$

$$C = 1$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 294: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}
 y_1 &= z_1 \\
 &= \cos(x)
 \end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x)) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1(\cos(x)) + c_2(\cos(x) (\tan(x))) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = \cos(x) c_1 + c_2 \sin(x)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of

parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \cos(x)$$

$$y_2 = \sin(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ \frac{d}{dx}(\cos(x)) & \frac{d}{dx}(\sin(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{vmatrix}$$

Therefore

$$W = (\cos(x))(\cos(x)) - (\sin(x))(-\sin(x))$$

Which simplifies to

$$W = \cos(x)^2 + \sin(x)^2$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\sin(x) \sec(x) \csc(x)}{1} dx$$

Which simplifies to

$$u_1 = - \int \sec(x) dx$$

Hence

$$u_1 = - \ln(\sec(x) + \tan(x))$$

And Eq. (3) becomes

$$u_2 = \int \frac{\cos(x) \sec(x) \csc(x)}{1} dx$$

Which simplifies to

$$u_2 = \int \csc(x) dx$$

Hence

$$u_2 = - \ln(\csc(x) + \cot(x))$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = - \cos(x) \ln(\sec(x) + \tan(x)) - \sin(x) \ln(\csc(x) + \cot(x))$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (\cos(x) c_1 + c_2 \sin(x)) + (- \cos(x) \ln(\sec(x) + \tan(x)) - \sin(x) \ln(\csc(x) + \cot(x))) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \cos(x) c_1 + c_2 \sin(x) - \cos(x) \ln(\sec(x) + \tan(x)) - \sin(x) \ln(\csc(x) + \cot(x))$$




Figure 363: Slope field plot

Verification of solutions

$$y = \cos(x) c_1 + c_2 \sin(x) - \cos(x) \ln(\sec(x) + \tan(x)) - \sin(x) \ln(\csc(x) + \cot(x))$$

Verified OK.

11.13.3 Maple step by step solution

Let's solve

$$y'' + y = \sec(x) \csc(x)$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (-I, I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = \cos(x) c_1 + c_2 \sin(x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \sec(x) \csc(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\cos(x) \left(\int \sec(x) dx \right) + \sin(x) \left(\int \csc(x) dx \right)$$

- Compute integrals

$$y_p(x) = -\cos(x) \ln(\sec(x) + \tan(x)) - \sin(x) \ln(\csc(x) + \cot(x))$$

- Substitute particular solution into general solution to ODE

$$y = \cos(x) c_1 + c_2 \sin(x) - \cos(x) \ln(\sec(x) + \tan(x)) - \sin(x) \ln(\csc(x) + \cot(x))$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
dsolve(diff(y(x),x$2)+y(x)=sec(x)*csc(x),y(x), singsol=all)
```

$$y(x) = \sin(x) c_2 + \cos(x) c_1 + \sin(x) \ln(\csc(x) - \cot(x)) - \cos(x) \ln(\sec(x) + \tan(x))$$

✓ Solution by Mathematica

Time used: 0.033 (sec). Leaf size: 30

```
DSolve[y''[x]+y[x]==Sec[x]*Csc[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sin(x)\operatorname{arctanh}(\cos(x)) + c_1 \cos(x) + c_2 \sin(x) + \cos(x) \left(-\operatorname{coth}^{-1}(\sin(x))\right)$$

11.14 problem 3

- 11.14.1 Solving as second order linear constant coeff ode 1988
- 11.14.2 Solving as linear second order ode solved by an integrating factor
ode 1991
- 11.14.3 Solving using Kovacic algorithm 1993
- 11.14.4 Maple step by step solution 1998

Internal problem ID [6330]

Internal file name [OUTPUT/5578_Sunday_June_05_2022_03_43_59_PM_97172001/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.3. THE METHOD OF VARIATION OF PARAMETERS. Page 71

Problem number: 3.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "linear_second_order_ode_solved_by_an_integrating_factor"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - 2y' + y = 2x$$

11.14.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = -2, C = 1, f(x) = 2x$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 2y' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -2, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 2\lambda e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 2\lambda + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -2, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{2}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{(-2)^2 - (4)(1)(1)} \\ &= 1 \end{aligned}$$

Hence this is the case of a double root $\lambda_{1,2} = -1$. Therefore the solution is

$$y = c_1 e^x + c_2 x e^x \quad (1)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^x + c_2 x e^x$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{x e^x, e^x\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_2x + A_1$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_2x + A_1 - 2A_2 = 2x$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 4, A_2 = 2]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = 2x + 4$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1e^x + c_2xe^x) + (2x + 4) \end{aligned}$$

Which simplifies to

$$y = e^x(c_2x + c_1) + 2x + 4$$

Summary

The solution(s) found are the following

$$y = e^x(c_2x + c_1) + 2x + 4 \tag{1}$$




Figure 364: Slope field plot

Verification of solutions

$$y = e^x(c_2x + c_1) + 2x + 4$$

Verified OK.

11.14.2 Solving as linear second order ode solved by an integrating factor ode

The ode satisfies this form

$$y'' + p(x)y' + \frac{(p(x)^2 + p'(x))y}{2} = f(x)$$

Where $p(x) = -2$. Therefore, there is an integrating factor given by

$$\begin{aligned} M(x) &= e^{\frac{1}{2} \int p dx} \\ &= e^{\int -2 dx} \\ &= e^{-x} \end{aligned}$$

Multiplying both sides of the ODE by the integrating factor $M(x)$ makes the left side of the ODE a complete differential

$$(M(x)y)'' = 2x e^{-x}$$

$$(e^{-x}y)'' = 2x e^{-x}$$

Integrating once gives

$$(e^{-x}y)' = -2(1+x)e^{-x} + c_1$$

Integrating again gives

$$(e^{-x}y) = (2x+4)e^{-x} + c_1x + c_2$$

Hence the solution is

$$y = \frac{(2x+4)e^{-x} + c_1x + c_2}{e^{-x}}$$

Or

$$y = c_1x e^x + c_2e^x + 2x + 4$$

Summary

The solution(s) found are the following

$$y = c_1x e^x + c_2e^x + 2x + 4 \tag{1}$$




Figure 365: Slope field plot

Verification of solutions

$$y = c_1 x e^x + c_2 e^x + 2x + 4$$

Verified OK.

11.14.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 2y' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -2 \\ C &= 1 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} = \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \tag{6}$$

Comparing the above to (5) shows that

$$s = 0$$

$$t = 1$$

Therefore eq. (4) becomes

$$z''(x) = 0 \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 296: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty\end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-2}{1} dx} \\ &= z_1 e^x \\ &= z_1(e^x)\end{aligned}$$

Which simplifies to

$$y_1 = e^x$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{-2}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{2x}}{(y_1)^2} dx \\ &= y_1(x)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^x) + c_2 (e^x(x))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 2y' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^x + c_2 x e^x$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{x e^x, e^x\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_2 x + A_1$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_2 x + A_1 - 2A_2 = 2x$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 4, A_2 = 2]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = 2x + 4$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^x + c_2 x e^x) + (2x + 4) \end{aligned}$$

Which simplifies to

$$y = e^x(c_2 x + c_1) + 2x + 4$$

Summary

The solution(s) found are the following

$$y = e^x(c_2 x + c_1) + 2x + 4 \tag{1}$$




Figure 366: Slope field plot

Verification of solutions

$$y = e^x(c_2x + c_1) + 2x + 4$$

Verified OK.

11.14.4 Maple step by step solution

Let's solve

$$y'' - 2y' + y = 2x$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - 2r + 1 = 0$$

- Factor the characteristic polynomial

$$(r - 1)^2 = 0$$

- Root of the characteristic polynomial

$$r = 1$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^x$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = x e^x$$

- General solution of the ODE

$$y = c_1y_1(x) + c_2y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1e^x + c_2xe^x + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = 2x \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^x & x e^x \\ e^x & x e^x + e^x \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = e^{2x}$$
- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -2e^x \left(\int x^2 e^{-x} dx - \left(\int x e^{-x} dx \right) x \right)$$
- Compute integrals

$$y_p(x) = 2x + 4$$
- Substitute particular solution into general solution to ODE

$$y = c_2 x e^x + c_1 e^x + 2x + 4$$

Maple trace

```

Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)-2*diff(y(x),x)+y(x)=2*x,y(x), singsol=all)
```

$$y(x) = (c_1 x + c_2) e^x + 2x + 4$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 26

```
DSolve[y''[x]+2*y'[x]+y[x]==2*x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x}(2e^x(x - 2) + c_2 x + c_1)$$

11.15 problem 4

11.15.1 Solving as second order linear constant coeff ode	2000
11.15.2 Solving using Kovacic algorithm	2003
11.15.3 Maple step by step solution	2008

Internal problem ID [6331]

Internal file name [OUTPUT/5579_Sunday_June_05_2022_03_44_00_PM_17667703/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.3. THE METHOD OF VARIATION OF PARAMETERS. Page 71

Problem number: 4.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"kovacic", "second_order_linear_constant_coeff"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - y' - 6y = e^{-x}$$

11.15.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = -1, C = -6, f(x) = e^{-x}$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - y' - 6y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -1, C = -6$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - \lambda e^{\lambda x} - 6 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - \lambda - 6 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -1, C = -6$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-1^2 - (4)(1)(-6)} \\ &= \frac{1}{2} \pm \frac{5}{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= \frac{1}{2} + \frac{5}{2} \\ \lambda_2 &= \frac{1}{2} - \frac{5}{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= 3 \\ \lambda_2 &= -2 \end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned} y &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ y &= c_1 e^{(3)x} + c_2 e^{(-2)x} \end{aligned}$$

Or

$$y = e^{3x} c_1 + c_2 e^{-2x}$$

Therefore the homogeneous solution y_h is

$$y_h = e^{3x} c_1 + c_2 e^{-2x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$e^{-x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{-x}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^{-2x}, e^{3x}\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^{-x}$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-4A_1 e^{-x} = e^{-x}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{4} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{e^{-x}}{4}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (e^{3x} c_1 + c_2 e^{-2x}) + \left(-\frac{e^{-x}}{4} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = e^{3x} c_1 + c_2 e^{-2x} - \frac{e^{-x}}{4} \quad (1)$$




Figure 367: Slope field plot

Verification of solutions

$$y = e^{3x}c_1 + c_2e^{-2x} - \frac{e^{-x}}{4}$$

Verified OK.

11.15.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' - y' - 6y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -1 \\ C &= -6 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{25}{4} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 25 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \frac{25z(x)}{4} \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 298: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{25}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{5x}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-1}{1} dx} \\ &= z_1 e^{\frac{x}{2}} \\ &= z_1 \left(e^{\frac{x}{2}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-2x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-1}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^x}{(y_1)^2} dx \\ &= y_1 \left(\frac{e^{5x}}{5} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-2x}) + c_2 \left(e^{-2x} \left(\frac{e^{5x}}{5} \right) \right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - y' - 6y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-2x} + \frac{c_2 e^{3x}}{5}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$e^{-x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{-x}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ \frac{e^{3x}}{5}, e^{-2x} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^{-x}$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-4A_1e^{-x} = e^{-x}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{4} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{e^{-x}}{4}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1e^{-2x} + \frac{c_2e^{3x}}{5} \right) + \left(-\frac{e^{-x}}{4} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1e^{-2x} + \frac{c_2e^{3x}}{5} - \frac{e^{-x}}{4} \quad (1)$$




Figure 368: Slope field plot

Verification of solutions

$$y = c_1 e^{-2x} + \frac{c_2 e^{3x}}{5} - \frac{e^{-x}}{4}$$

Verified OK.

11.15.3 Maple step by step solution

Let's solve

$$y'' - y' - 6y = e^{-x}$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - r - 6 = 0$$

- Factor the characteristic polynomial

$$(r + 2)(r - 3) = 0$$

- Roots of the characteristic polynomial

$$r = (-2, 3)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-2x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{3x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{-2x} + c_2 e^{3x} + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = e^{-x} \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-2x} & e^{3x} \\ -2e^{-2x} & 3e^{3x} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 5e^x$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = \frac{(e^{5x}(\int e^{-4x} dx) - (\int e^x dx)e^{-2x})}{5}$$

- Compute integrals

$$y_p(x) = -\frac{e^{-x}}{4}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{-2x} + c_2 e^{3x} - \frac{e^{-x}}{4}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(diff(y(x),x$2)-diff(y(x),x)-6*y(x)=exp(-x),y(x), singsol=all)
```

$$y(x) = -\frac{(-4c_2e^{5x} + e^x - 4c_1)e^{-2x}}{4}$$

✓ Solution by Mathematica

Time used: 0.037 (sec). Leaf size: 31

```
DSolve[y''[x]-y'[x]-6*y[x]==Exp[-x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{e^{-x}}{4} + c_1e^{-2x} + c_2e^{3x}$$

11.16 problem 5(a)

- 11.16.1 Solving as second order change of variable on y method 2 ode . 2011
- 11.16.2 Solving as second order ode non constant coeff transformation
on B ode 2016
- 11.16.3 Solving using Kovacic algorithm 2021

Internal problem ID [6332]

Internal file name [OUTPUT/5580_Sunday_June_05_2022_03_44_02_PM_71792521/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.3. THE METHOD OF VARIATION OF PARAMETERS. Page 71

Problem number: 5(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_change_of_variable_on_y_method_2", "second_order_ode_non_constant_coeff_transformation_on_B"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$(x^2 - 1)y'' - 2xy' + 2y = (x^2 - 1)^2$$

11.16.1 Solving as second order change of variable on y method 2 ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = x^2 - 1$, $B = -2x$, $C = 2$, $f(x) = x^4 - 2x^2 + 1$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$(x^2 - 1)y'' - 2xy' + 2y = 0$$

In normal form the ode

$$(x^2 - 1) y'' - 2xy' + 2y = 0 \quad (1)$$

Becomes

$$y'' + p(x) y' + q(x) y = 0 \quad (2)$$

Where

$$p(x) = -\frac{2x}{x^2 - 1}$$
$$q(x) = \frac{2}{x^2 - 1}$$

Applying change of variables on the dependent variable $y = v(x) x^n$ to (2) gives the following ode where the dependent variable is $v(x)$ and not y .

$$v''(x) + \left(\frac{2n}{x} + p\right) v'(x) + \left(\frac{n(n-1)}{x^2} + \frac{np}{x} + q\right) v(x) = 0 \quad (3)$$

Let the coefficient of $v(x)$ above be zero. Hence

$$\frac{n(n-1)}{x^2} + \frac{np}{x} + q = 0 \quad (4)$$

Substituting the earlier values found for $p(x)$ and $q(x)$ into (4) gives

$$\frac{n(n-1)}{x^2} - \frac{2n}{x^2 - 1} + \frac{2}{x^2 - 1} = 0 \quad (5)$$

Solving (5) for n gives

$$n = 1 \quad (6)$$

Substituting this value in (3) gives

$$v''(x) + \left(\frac{2}{x} - \frac{2x}{x^2 - 1}\right) v'(x) = 0$$
$$v''(x) - \frac{2v'(x)}{x^3 - x} = 0 \quad (7)$$

Using the substitution

$$u(x) = v'(x)$$

Then (7) becomes

$$u'(x) - \frac{2u(x)}{x^3 - x} = 0 \quad (8)$$

The above is now solved for $u(x)$. In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= \frac{2u}{x(x^2 - 1)} \end{aligned}$$

Where $f(x) = \frac{2}{x(x^2-1)}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned} \frac{1}{u} du &= \frac{2}{x(x^2 - 1)} dx \\ \int \frac{1}{u} du &= \int \frac{2}{x(x^2 - 1)} dx \\ \ln(u) &= \ln(1+x) + \ln(x-1) - 2\ln(x) + c_1 \\ u &= e^{\ln(1+x)+\ln(x-1)-2\ln(x)+c_1} \\ &= c_1 e^{\ln(1+x)+\ln(x-1)-2\ln(x)} \end{aligned}$$

Which simplifies to

$$u(x) = c_1 \left(1 - \frac{1}{x^2} \right)$$

Now that $u(x)$ is known, then

$$\begin{aligned} v'(x) &= u(x) \\ v(x) &= \int u(x) dx + c_2 \\ &= c_1 \left(x + \frac{1}{x} \right) + c_2 \end{aligned}$$

Hence

$$\begin{aligned} y &= v(x) x^n \\ &= \left(c_1 \left(x + \frac{1}{x} \right) + c_2 \right) x \\ &= c_1 x^2 + c_2 x + c_1 \end{aligned}$$

Now the particular solution to this ODE is found

$$(x^2 - 1) y'' - 2xy' + 2y = x^4 - 2x^2 + 1$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = x$$

$$y_2 = x^2 + 1$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} x & x^2 + 1 \\ \frac{d}{dx}(x) & \frac{d}{dx}(x^2 + 1) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} x & x^2 + 1 \\ 1 & 2x \end{vmatrix}$$

Therefore

$$W = (x)(2x) - (x^2 + 1)(1)$$

Which simplifies to

$$W = x^2 - 1$$

Which simplifies to

$$W = x^2 - 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{(x^2 + 1)(x^4 - 2x^2 + 1)}{(x^2 - 1)^2} dx$$

Which simplifies to

$$u_1 = - \int (x^2 + 1) dx$$

Hence

$$u_1 = -\frac{1}{3}x^3 - x$$

And Eq. (3) becomes

$$u_2 = \int \frac{x(x^4 - 2x^2 + 1)}{(x^2 - 1)^2} dx$$

Which simplifies to

$$u_2 = \int x dx$$

Hence

$$u_2 = \frac{x^2}{2}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(-\frac{1}{3}x^3 - x\right)x + \frac{(x^2 + 1)x^2}{2}$$

Which simplifies to

$$y_p(x) = \frac{1}{6}x^4 - \frac{1}{2}x^2$$

Therefore the general solution is

$$\begin{aligned}y &= y_h + y_p \\&= \left(\left(c_1 \left(x + \frac{1}{x} \right) + c_2 \right) x \right) + \left(\frac{1}{6}x^4 - \frac{1}{2}x^2 \right) \\&= \frac{x^4}{6} - \frac{x^2}{2} + \left(c_1 \left(x + \frac{1}{x} \right) + c_2 \right) x\end{aligned}$$

Which simplifies to

$$y = \frac{1}{6}x^4 - \frac{1}{2}x^2 + c_1x^2 + c_2x + c_1$$

Summary

The solution(s) found are the following

$$y = \frac{1}{6}x^4 - \frac{1}{2}x^2 + c_1x^2 + c_2x + c_1 \quad (1)$$

Verification of solutions

$$y = \frac{1}{6}x^4 - \frac{1}{2}x^2 + c_1x^2 + c_2x + c_1$$

Verified OK.

11.16.2 Solving as second order ode non constant coeff transformation on B ode

Given an ode of the form

$$Ay'' + By' + Cy = F(x)$$

This method reduces the order ode the ODE by one by applying the transformation

$$y = Bv$$

This results in

$$\begin{aligned}y' &= B'v + v'B \\y'' &= B''v + B'v' + v''B + v'B' \\&= v''B + 2v' + B' + B''v\end{aligned}$$

And now the original ode becomes

$$\begin{aligned} A(v''B + 2v'B' + B''v) + B(B'v + v'B) + CBv &= 0 \\ ABv'' + (2AB' + B^2)v' + (AB'' + BB' + CB)v &= 0 \end{aligned} \quad (1)$$

If the term $AB'' + BB' + CB$ is zero, then this method works and can be used to solve

$$ABv'' + (2AB' + B^2)v' = 0$$

By Using $u = v'$ which reduces the order of the above ode to one. The new ode is

$$ABu' + (2AB' + B^2)u = 0$$

The above ode is first order ode which is solved for u . Now a new ode $v' = u$ is solved for v as first order ode. Then the final solution is obtain from $y = Bv$.

This method works only if the term $AB'' + BB' + CB$ is zero. The given ODE shows that

$$\begin{aligned} A &= x^2 - 1 \\ B &= -2x \\ C &= 2 \\ F &= x^4 - 2x^2 + 1 \end{aligned}$$

The above shows that for this ode

$$\begin{aligned} AB'' + BB' + CB &= (x^2 - 1)(0) + (-2x)(-2) + (2)(-2x) \\ &= 0 \end{aligned}$$

Hence the ode in v given in (1) now simplifies to

$$-2x^3 + 2xv'' + (4)v' = 0$$

Now by applying $v' = u$ the above becomes

$$(-2x^3 + 2x)u'(x) + 4u(x) = 0$$

Which is now solved for u . In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= \frac{2u}{x(x^2 - 1)} \end{aligned}$$

Where $f(x) = \frac{2}{x(x^2-1)}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u} du &= \frac{2}{x(x^2-1)} dx \\ \int \frac{1}{u} du &= \int \frac{2}{x(x^2-1)} dx \\ \ln(u) &= \ln(1+x) + \ln(x-1) - 2\ln(x) + c_1 \\ u &= e^{\ln(1+x)+\ln(x-1)-2\ln(x)+c_1} \\ &= c_1 e^{\ln(1+x)+\ln(x-1)-2\ln(x)}\end{aligned}$$

Which simplifies to

$$u(x) = c_1 \left(1 - \frac{1}{x^2}\right)$$

The ode for v now becomes

$$\begin{aligned}v' &= u \\ &= c_1 \left(1 - \frac{1}{x^2}\right)\end{aligned}$$

Which is now solved for v . Integrating both sides gives

$$\begin{aligned}v(x) &= \int \frac{c_1(x^2-1)}{x^2} dx \\ &= c_1 \left(x + \frac{1}{x}\right) + c_2\end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}y_h(x) &= Bv \\ &= (-2x) \left(c_1 \left(x + \frac{1}{x}\right) + c_2\right) \\ &= -2c_1x^2 - 2c_2x - 2c_1\end{aligned}$$

And now the particular solution $y_p(x)$ will be found. The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1y_1 + u_2y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned}y_1 &= x \\y_2 &= -2x^2 - 2\end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} x & -2x^2 - 2 \\ \frac{d}{dx}(x) & \frac{d}{dx}(-2x^2 - 2) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} x & -2x^2 - 2 \\ 1 & -4x \end{vmatrix}$$

Therefore

$$W = (x)(-4x) - (-2x^2 - 2) \quad (1)$$

Which simplifies to

$$W = -2x^2 + 2$$

Which simplifies to

$$W = -2x^2 + 2$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{(-2x^2 - 2)(x^4 - 2x^2 + 1)}{(x^2 - 1)(-2x^2 + 2)} dx$$

Which simplifies to

$$u_1 = - \int (x^2 + 1) dx$$

Hence

$$u_1 = -\frac{1}{3}x^3 - x$$

And Eq. (3) becomes

$$u_2 = \int \frac{x(x^4 - 2x^2 + 1)}{(x^2 - 1)(-2x^2 + 2)} dx$$

Which simplifies to

$$u_2 = \int -\frac{x}{2} dx$$

Hence

$$u_2 = -\frac{x^2}{4}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(-\frac{1}{3}x^3 - x\right) x - \frac{x^2(-2x^2 - 2)}{4}$$

Which simplifies to

$$y_p(x) = \frac{1}{6}x^4 - \frac{1}{2}x^2$$

Hence the complete solution is

$$\begin{aligned} y(x) &= y_h + y_p \\ &= (-2c_1x^2 - 2c_2x - 2c_1) + \left(\frac{1}{6}x^4 - \frac{1}{2}x^2\right) \\ &= \frac{x^4}{6} + \frac{(-1 - 4c_1)x^2}{2} - 2c_2x - 2c_1 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{x^4}{6} + \frac{(-1 - 4c_1)x^2}{2} - 2c_2x - 2c_1 \quad (1)$$

Verification of solutions

$$y = \frac{x^4}{6} + \frac{(-1 - 4c_1)x^2}{2} - 2c_2x - 2c_1$$

Verified OK.

11.16.3 Solving using Kovacic algorithm

Writing the ode as

$$(x^2 - 1)y'' - 2xy' + 2y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= x^2 - 1 \\ B &= -2x \\ C &= 2 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{3}{(x^2 - 1)^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 3 \\ t &= (x^2 - 1)^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{3}{(x^2 - 1)^2} \right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 300: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 4 - 0 \\ &= 4 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = (x^2 - 1)^2$. There is a pole at $x = 1$ of order 2. There is a pole at $x = -1$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 4 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 4 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = \frac{3}{4(x-1)^2} - \frac{3}{4(x-1)} + \frac{3}{4(1+x)} + \frac{3}{4(1+x)^2}$$

For the pole at $x = 1$ let b be the coefficient of $\frac{1}{(x-1)^2}$ in the partial fractions decomposition of r given above. Therefore $b = \frac{3}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1+4b} = \frac{3}{2} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1+4b} = -\frac{1}{2} \end{aligned}$$

For the pole at $x = -1$ let b be the coefficient of $\frac{1}{(1+x)^2}$ in the partial fractions decomposition of r given above. Therefore $b = \frac{3}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1+4b} = \frac{3}{2} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1+4b} = -\frac{1}{2} \end{aligned}$$

Since the order of r at ∞ is $4 > 2$ then

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= 0 \\ \alpha_\infty^- &= 1 \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{3}{(x^2 - 1)^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
1	2	0	$\frac{3}{2}$	$-\frac{1}{2}$
-1	2	0	$\frac{3}{2}$	$-\frac{1}{2}$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
4	0	0	1

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^- = 1$ then

$$\begin{aligned} d &= \alpha_\infty^- - (\alpha_{c_1}^- + \alpha_{c_2}^+) \\ &= 1 - (1) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{x - c_1} \right) + \left((+)[\sqrt{r}]_{c_2} + \frac{\alpha_{c_2}^+}{x - c_2} \right) + (-)[\sqrt{r}]_\infty \\ &= -\frac{1}{2(x-1)} + \frac{3}{2(1+x)} + (-)(0) \\ &= -\frac{1}{2(x-1)} + \frac{3}{2(1+x)} \\ &= \frac{-2+x}{x^2-1} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$(0) + 2 \left(-\frac{1}{2(x-1)} + \frac{3}{2(1+x)} \right) (0) + \left(\left(\frac{1}{2(x-1)^2} - \frac{3}{2(1+x)^2} \right) + \left(-\frac{1}{2(x-1)} + \frac{3}{2(1+x)} \right)^2 - \left(\frac{1}{2(x-1)} + \frac{3}{2(1+x)} \right) \right) (0) = 0$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= p e^{\int \omega dx} \\ &= e^{\int \left(-\frac{1}{2(x-1)} + \frac{3}{2(1+x)} \right) dx} \\ &= \frac{(1+x)^{\frac{3}{2}}}{\sqrt{x-1}} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-2x}{x^2-1} dx} \\ &= z_1 e^{\frac{\ln(x-1)}{2} + \frac{\ln(1+x)}{2}} \\ &= z_1 \left(\sqrt{x-1} \sqrt{1+x} \right) \end{aligned}$$

Which simplifies to

$$y_1 = (1+x)^2$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-2x}{x^2-1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{\ln(x-1) + \ln(1+x)}}{(y_1)^2} dx \\ &= y_1 \left(-\frac{x}{(1+x)^2} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 ((1+x)^2) + c_2 \left((1+x)^2 \left(-\frac{x}{(1+x)^2} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$(x^2 - 1) y'' - 2xy' + 2y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1(1 + x)^2 - c_2x$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1y_1 + u_2y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = (1 + x)^2$$

$$y_2 = -x$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} (1 + x)^2 & -x \\ \frac{d}{dx}((1 + x)^2) & \frac{d}{dx}(-x) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} (1+x)^2 & -x \\ 2+2x & -1 \end{vmatrix}$$

Therefore

$$W = ((1+x)^2)(-1) - (-x)(2+2x)$$

Which simplifies to

$$W = x^2 - 1$$

Which simplifies to

$$W = x^2 - 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{-x(x^4 - 2x^2 + 1)}{(x^2 - 1)^2} dx$$

Which simplifies to

$$u_1 = - \int -x dx$$

Hence

$$u_1 = \frac{x^2}{2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{(1+x)^2(x^4 - 2x^2 + 1)}{(x^2 - 1)^2} dx$$

Which simplifies to

$$u_2 = \int (1+x)^2 dx$$

Hence

$$u_2 = \frac{(1+x)^3}{3}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{x^2(1+x)^2}{2} - \frac{(1+x)^3 x}{3}$$

Which simplifies to

$$y_p(x) = \frac{x(x^3 - 3x - 2)}{6}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1(1+x)^2 - c_2x) + \left(\frac{x(x^3 - 3x - 2)}{6} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1(1+x)^2 - c_2x + \frac{x(x^3 - 3x - 2)}{6} \quad (1)$$

Verification of solutions

$$y = c_1(1+x)^2 - c_2x + \frac{x(x^3 - 3x - 2)}{6}$$

Verified OK.

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Reducible group (found an exponential solution)
<- Kovacics algorithm successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 20

```
dsolve((x^2-1)*diff(y(x),x$2)-2*x*diff(y(x),x)+2*y(x)=(x^2-1)^2,y(x), singsol=all)
```

$$y(x) = c_2x + c_1x^2 + c_1 + \frac{1}{2} + \frac{1}{6}x^4$$

✓ Solution by Mathematica

Time used: 0.375 (sec). Leaf size: 111

```
DSolve[(x^2-1)*y'[x]-2*x*y'[x]+2*y[x]==(x^2-1)^2,y[x],x,IncludeSingularSolutions -> True]
```

$y(x)$

$$\rightarrow \frac{-x^6 + 4x^4 - 2x^3 + 3x^2 \left(-1 + 2c_1 \sqrt{-(x^2 - 1)^2} \right) + 2x \left(-6c_1 \sqrt{-(x^2 - 1)^2} + 3c_2 \sqrt{-(x^2 - 1)^2 + 1} \right)}{6 - 6x^2}$$

11.17 problem 5(b)

11.17.1 Solving as second order change of variable on y method 2 ode . 2030

11.17.2 Solving using Kovacic algorithm 2035

Internal problem ID [6333]

Internal file name [OUTPUT/5581_Sunday_June_05_2022_03_44_04_PM_12485339/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.3. THE METHOD OF VARIATION OF PARAMETERS. Page 71

Problem number: 5(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_change_of_variable_on_y_method_2"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$(x^2 + x)y'' + (-x^2 + 2)y' - (x + 2)y = x(1 + x)^2$$

11.17.1 Solving as second order change of variable on y method 2 ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = x^2 + x$, $B = -x^2 + 2$, $C = -x - 2$, $f(x) = x(1 + x)^2$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$(x^2 + x)y'' + (-x^2 + 2)y' + (-x - 2)y = 0$$

In normal form the ode

$$(x^2 + x) y'' + (-x^2 + 2) y' + (-x - 2) y = 0 \quad (1)$$

Becomes

$$y'' + p(x) y' + q(x) y = 0 \quad (2)$$

Where

$$p(x) = \frac{-x^2 + 2}{x(1+x)}$$
$$q(x) = \frac{-x - 2}{x(1+x)}$$

Applying change of variables on the dependent variable $y = v(x) x^n$ to (2) gives the following ode where the dependent variables is $v(x)$ and not y .

$$v''(x) + \left(\frac{2n}{x} + p\right) v'(x) + \left(\frac{n(n-1)}{x^2} + \frac{np}{x} + q\right) v(x) = 0 \quad (3)$$

Let the coefficient of $v(x)$ above be zero. Hence

$$\frac{n(n-1)}{x^2} + \frac{np}{x} + q = 0 \quad (4)$$

Substituting the earlier values found for $p(x)$ and $q(x)$ into (4) gives

$$\frac{n(n-1)}{x^2} + \frac{n(-x^2 + 2)}{x^2(1+x)} + \frac{-x - 2}{x(1+x)} = 0 \quad (5)$$

Solving (5) for n gives

$$n = -1 \quad (6)$$

Substituting this value in (3) gives

$$v''(x) + \left(-\frac{2}{x} + \frac{-x^2 + 2}{x(1+x)}\right) v'(x) = 0$$
$$v''(x) + \frac{(-x - 2) v'(x)}{1+x} = 0 \quad (7)$$

Using the substitution

$$u(x) = v'(x)$$

Then (7) becomes

$$u'(x) + \frac{(-x-2)u(x)}{1+x} = 0 \quad (8)$$

The above is now solved for $u(x)$. In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= \frac{u(x+2)}{1+x} \end{aligned}$$

Where $f(x) = \frac{x+2}{1+x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned} \frac{1}{u} du &= \frac{x+2}{1+x} dx \\ \int \frac{1}{u} du &= \int \frac{x+2}{1+x} dx \\ \ln(u) &= x + \ln(1+x) + c_1 \\ u &= e^{x+\ln(1+x)+c_1} \\ &= c_1 e^{x+\ln(1+x)} \end{aligned}$$

Now that $u(x)$ is known, then

$$\begin{aligned} v'(x) &= u(x) \\ v(x) &= \int u(x) dx + c_2 \\ &= \frac{xc_1 e^{x+\ln(1+x)}}{1+x} + c_2 \end{aligned}$$

Hence

$$\begin{aligned} y &= v(x) x^n \\ &= \frac{\frac{xc_1 e^{x+\ln(1+x)}}{1+x} + c_2}{x} \\ &= \frac{c_1 x e^x + c_2}{x} \end{aligned}$$

Now the particular solution to this ODE is found

$$(x^2 + x)y'' + (-x^2 + 2)y' + (-x - 2)y = x(1+x)^2$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \frac{1}{x}$$

$$y_2 = \frac{x e^x}{1+x} + \frac{e^x}{1+x}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \frac{1}{x} & \frac{x e^x}{1+x} + \frac{e^x}{1+x} \\ \frac{d}{dx} \left(\frac{1}{x} \right) & \frac{d}{dx} \left(\frac{x e^x}{1+x} + \frac{e^x}{1+x} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \frac{1}{x} & \frac{x e^x}{1+x} + \frac{e^x}{1+x} \\ -\frac{1}{x^2} & \frac{2 e^x}{1+x} - \frac{x e^x}{(1+x)^2} + \frac{x e^x}{1+x} - \frac{e^x}{(1+x)^2} \end{vmatrix}$$

Therefore

$$W = \left(\frac{1}{x} \right) \left(\frac{2 e^x}{1+x} - \frac{x e^x}{(1+x)^2} + \frac{x e^x}{1+x} - \frac{e^x}{(1+x)^2} \right) - \left(\frac{x e^x}{1+x} + \frac{e^x}{1+x} \right) \left(-\frac{1}{x^2} \right)$$

Which simplifies to

$$W = \frac{(1+x)e^x}{x^2}$$

Which simplifies to

$$W = \frac{(1+x)e^x}{x^2}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\left(\frac{xe^x}{1+x} + \frac{e^x}{1+x}\right) x(1+x)^2}{\frac{(x^2+x)(1+x)e^x}{x^2}} dx$$

Which simplifies to

$$u_1 = - \int x^2 dx$$

Hence

$$u_1 = -\frac{x^3}{3}$$

And Eq. (3) becomes

$$u_2 = \int \frac{(1+x)^2}{\frac{(x^2+x)(1+x)e^x}{x^2}} dx$$

Which simplifies to

$$u_2 = \int x e^{-x} dx$$

Hence

$$u_2 = -(1+x)e^{-x}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{x^2}{3} - (1+x)e^{-x} \left(\frac{xe^x}{1+x} + \frac{e^x}{1+x} \right)$$

Which simplifies to

$$y_p(x) = -\frac{1}{3}x^2 - 1 - x$$

Therefore the general solution is

$$\begin{aligned}
 y &= y_h + y_p \\
 &= \left(\frac{\frac{xc_1 e^{x+\ln(1+x)}}{1+x} + c_2}{x} \right) + \left(-\frac{1}{3}x^2 - 1 - x \right) \\
 &= -\frac{x^2}{3} - 1 - x + \frac{\frac{xc_1 e^{x+\ln(1+x)}}{1+x} + c_2}{x}
 \end{aligned}$$

Which simplifies to

$$y = \frac{3c_1 x e^x - x^3 - 3x^2 + 3c_2 - 3x}{3x}$$

Summary

The solution(s) found are the following

$$y = \frac{3c_1 x e^x - x^3 - 3x^2 + 3c_2 - 3x}{3x} \quad (1)$$

Verification of solutions

$$y = \frac{3c_1 x e^x - x^3 - 3x^2 + 3c_2 - 3x}{3x}$$

Verified OK.

11.17.2 Solving using Kovacic algorithm

Writing the ode as

$$(x^2 + x) y'' + (-x^2 + 2) y' + (-x - 2) y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned}
 A &= x^2 + x \\
 B &= -x^2 + 2 \\
 C &= -x - 2
 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{x^2 + 4x + 6}{4(1+x)^2} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= x^2 + 4x + 6 \\ t &= 4(1+x)^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{x^2 + 4x + 6}{4(1+x)^2} \right) z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 301: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 2 \\ &= 0 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4(1+x)^2$. There is a pole at $x = -1$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Therefore

$$L = [1, 2]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = \frac{1}{4} + \frac{1}{2+2x} + \frac{3}{4(1+x)^2}$$

For the pole at $x = -1$ let b be the coefficient of $\frac{1}{(1+x)^2}$ in the partial fractions decomposition of r given above. Therefore $b = \frac{3}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1+4b} = \frac{3}{2} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1+4b} = -\frac{1}{2} \end{aligned}$$

Since the order of r at ∞ is $O_r(\infty) = 0$ then

$$v = \frac{-O_r(\infty)}{2} = \frac{0}{2} = 0$$

$[\sqrt{r}]_\infty$ is the sum of terms involving x^i for $0 \leq i \leq v$ in the Laurent series for \sqrt{r} at ∞ . Therefore

$$\begin{aligned} [\sqrt{r}]_\infty &= \sum_{i=0}^v a_i x^i \\ &= \sum_{i=0}^0 a_i x^i \end{aligned} \tag{8}$$

Let a be the coefficient of $x^v = x^0$ in the above sum. The Laurent series of \sqrt{r} at ∞ is

$$\sqrt{r} \approx \frac{1}{2} + \frac{1}{2x} - \frac{1}{x^3} + \frac{11}{4x^4} - \frac{21}{4x^5} + \frac{15}{2x^6} - \frac{6}{x^7} - \frac{117}{16x^8} + \dots \quad (9)$$

Comparing Eq. (9) with Eq. (8) shows that

$$a = \frac{1}{2}$$

From Eq. (9) the sum up to $v = 0$ gives

$$\begin{aligned} [\sqrt{r}]_{\infty} &= \sum_{i=0}^0 a_i x^i \\ &= \frac{1}{2} \end{aligned} \quad (10)$$

Now we need to find b , where b be the coefficient of $x^{v-1} = x^{-1} = \frac{1}{x}$ in r minus the coefficient of same term but in $([\sqrt{r}]_{\infty})^2$ where $[\sqrt{r}]_{\infty}$ was found above in Eq (10). Hence

$$([\sqrt{r}]_{\infty})^2 = \frac{1}{4}$$

This shows that the coefficient of $\frac{1}{x}$ in the above is 0. Now we need to find the coefficient of $\frac{1}{x}$ in r . How this is done depends on if $v = 0$ or not. Since $v = 0$ then starting from $r = \frac{s}{t}$ and doing long division in the form

$$r = Q + \frac{R}{t}$$

Where Q is the quotient and R is the remainder. Then the coefficient of $\frac{1}{x}$ in r will be the coefficient in R of the term in x of degree of t minus one, divided by the leading coefficient in t . Doing long division gives

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{x^2 + 4x + 6}{4x^2 + 8x + 4} \\ &= Q + \frac{R}{4x^2 + 8x + 4} \\ &= \left(\frac{1}{4}\right) + \left(\frac{2x + 5}{4x^2 + 8x + 4}\right) \\ &= \frac{1}{4} + \frac{2x + 5}{4x^2 + 8x + 4} \end{aligned}$$

Since the degree of t is 2, then we see that the coefficient of the term x in the remainder R is 2. Dividing this by leading coefficient in t which is 4 gives $\frac{1}{2}$. Now b can be found.

$$\begin{aligned} b &= \left(\frac{1}{2}\right) - (0) \\ &= \frac{1}{2} \end{aligned}$$

Hence

$$\begin{aligned} [\sqrt{r}]_{\infty} &= \frac{1}{2} \\ \alpha_{\infty}^{+} &= \frac{1}{2} \left(\frac{b}{a} - v \right) = \frac{1}{2} \left(\frac{\frac{1}{2}}{\frac{1}{2}} - 0 \right) = \frac{1}{2} \\ \alpha_{\infty}^{-} &= \frac{1}{2} \left(-\frac{b}{a} - v \right) = \frac{1}{2} \left(-\frac{\frac{1}{2}}{\frac{1}{2}} - 0 \right) = -\frac{1}{2} \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{x^2 + 4x + 6}{4(1+x)^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^{+}	α_c^{-}
-1	2	0	$\frac{3}{2}$	$-\frac{1}{2}$

Order of r at ∞	$[\sqrt{r}]_{\infty}$	α_{∞}^{+}	α_{∞}^{-}
0	$\frac{1}{2}$	$\frac{1}{2}$	$-\frac{1}{2}$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^{\pm} have been determined for all the poles in the set Γ and $[\sqrt{r}]_{\infty}$ and its associated α_{∞}^{\pm} have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_{\infty}^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_{∞}^{\pm} . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_{\infty}^{-} = -\frac{1}{2}$ then

$$\begin{aligned} d &= \alpha_{\infty}^{-} - (\alpha_{c_1}^{-}) \\ &= -\frac{1}{2} - \left(-\frac{1}{2}\right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_\infty$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{x - c_1} \right) + (-)[\sqrt{r}]_\infty \\ &= -\frac{1}{2(1+x)} + (-) \left(\frac{1}{2} \right) \\ &= -\frac{1}{2(1+x)} - \frac{1}{2} \\ &= -\frac{x+2}{2(1+x)} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2 \left(-\frac{1}{2(1+x)} - \frac{1}{2} \right) (0) + \left(\left(\frac{1}{2(1+x)^2} \right) + \left(-\frac{1}{2(1+x)} - \frac{1}{2} \right)^2 - \left(\frac{x^2 + 4x + 6}{4(1+x)^2} \right) \right) &= 0 \\ 0 &= 0 \end{aligned}$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= p e^{\int \omega dx} \\ &= e^{\int \left(-\frac{1}{2(1+x)} - \frac{1}{2} \right) dx} \\ &= \frac{e^{-\frac{x}{2}}}{\sqrt{1+x}} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\&= z_1 e^{-\int \frac{1}{2} \frac{-x^2+2}{x^2+x} dx} \\&= z_1 e^{\frac{x}{2} + \frac{\ln(1+x)}{2} - \ln(x)} \\&= z_1 \left(\frac{\sqrt{1+x} e^{\frac{x}{2}}}{x} \right)\end{aligned}$$

Which simplifies to

$$y_1 = \frac{1}{x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{-x^2+2}{x^2+x} dx}}{(y_1)^2} dx \\&= y_1 \int \frac{e^{x+\ln(1+x)-2\ln(x)}}{(y_1)^2} dx \\&= y_1(x e^x)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\&= c_1 \left(\frac{1}{x} \right) + c_2 \left(\frac{1}{x} (x e^x) \right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$(x^2 + x)y'' + (-x^2 + 2)y' + (-x - 2)y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = \frac{c_1}{x} + c_2e^x$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1y_1 + u_2y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \frac{1}{x}$$

$$y_2 = e^x$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \frac{1}{x} & e^x \\ \frac{d}{dx}\left(\frac{1}{x}\right) & \frac{d}{dx}(e^x) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \frac{1}{x} & e^x \\ -\frac{1}{x^2} & e^x \end{vmatrix}$$

Therefore

$$W = \left(\frac{1}{x}\right) (e^x) - (e^x) \left(-\frac{1}{x^2}\right)$$

Which simplifies to

$$W = \frac{(1+x)e^x}{x^2}$$

Which simplifies to

$$W = \frac{(1+x)e^x}{x^2}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{e^x x (1+x)^2}{\frac{(x^2+x)(1+x)e^x}{x^2}} dx$$

Which simplifies to

$$u_1 = - \int x^2 dx$$

Hence

$$u_1 = -\frac{x^3}{3}$$

And Eq. (3) becomes

$$u_2 = \int \frac{(1+x)^2}{\frac{(x^2+x)(1+x)e^x}{x^2}} dx$$

Which simplifies to

$$u_2 = \int x e^{-x} dx$$

Hence

$$u_2 = -(1+x)e^{-x}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{x^2}{3} - (1+x)e^{-x}e^x$$

Which simplifies to

$$y_p(x) = -\frac{1}{3}x^2 - 1 - x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(\frac{c_1}{x} + c_2e^x\right) + \left(-\frac{1}{3}x^2 - 1 - x\right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1}{x} + c_2e^x - \frac{x^2}{3} - 1 - x \quad (1)$$

Verification of solutions

$$y = \frac{c_1}{x} + c_2e^x - \frac{x^2}{3} - 1 - x$$

Verified OK.

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Reducible group (found an exponential solution)
    Reducible group (found another exponential solution)
<- Kovacics algorithm successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 32

```
dsolve((x^2+x)*diff(y(x),x$2)+(2-x^2)*diff(y(x),x)-(2+x)*y(x)=x*(x+1)^2,y(x), singsol=all)
```

$$y(x) = \frac{3x e^x c_1 - x^3 - 3x^2 + 3c_2 - 3x}{3x}$$

✓ Solution by Mathematica

Time used: 0.143 (sec). Leaf size: 45

```
DSolve[(x^2+x)*y'[x]+(2-x^2)*y'[x]-(2+x)*y[x]==x*(x+1)^2,y[x],x,IncludeSingularSolutions ->
```

$$y(x) \rightarrow -\frac{x^2}{3} - x + \sqrt{2}c_2 e^{x+\frac{1}{2}} + \frac{c_1}{\sqrt{2}ex} - 1$$

11.18 problem 5(c)

- 11.18.1 Solving as second order change of variable on y method 2 ode . 2046
- 11.18.2 Solving as second order ode non constant coeff transformation
on B ode 2051
- 11.18.3 Solving using Kovacic algorithm 2056

Internal problem ID [6334]

Internal file name [OUTPUT/5582_Sunday_June_05_2022_03_44_07_PM_42183739/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.3. THE METHOD OF VARIATION OF PARAMETERS. Page 71

Problem number: 5(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_change_of_variable_on_y_method_2", "second_order_ode_non_constant_coeff_transformation_on_B"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$(1 - x)y'' + xy' - y = (1 - x)^2$$

11.18.1 Solving as second order change of variable on y method 2 ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1 - x$, $B = x$, $C = -1$, $f(x) = (x - 1)^2$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$(1 - x)y'' + xy' - y = 0$$

In normal form the ode

$$(1 - x) y'' + x y' - y = 0 \quad (1)$$

Becomes

$$y'' + p(x) y' + q(x) y = 0 \quad (2)$$

Where

$$p(x) = -\frac{x}{x-1}$$
$$q(x) = \frac{1}{x-1}$$

Applying change of variables on the dependent variable $y = v(x) x^n$ to (2) gives the following ode where the dependent variables is $v(x)$ and not y .

$$v''(x) + \left(\frac{2n}{x} + p\right) v'(x) + \left(\frac{n(n-1)}{x^2} + \frac{np}{x} + q\right) v(x) = 0 \quad (3)$$

Let the coefficient of $v(x)$ above be zero. Hence

$$\frac{n(n-1)}{x^2} + \frac{np}{x} + q = 0 \quad (4)$$

Substituting the earlier values found for $p(x)$ and $q(x)$ into (4) gives

$$\frac{n(n-1)}{x^2} - \frac{n}{x-1} + \frac{1}{x-1} = 0 \quad (5)$$

Solving (5) for n gives

$$n = 1 \quad (6)$$

Substituting this value in (3) gives

$$v''(x) + \left(\frac{2}{x} - \frac{x}{x-1}\right) v'(x) = 0$$
$$v''(x) + \left(\frac{2}{x} - \frac{x}{x-1}\right) v'(x) = 0 \quad (7)$$

Using the substitution

$$u(x) = v'(x)$$

Then (7) becomes

$$u'(x) + \left(\frac{2}{x} - \frac{x}{x-1} \right) u(x) = 0 \quad (8)$$

The above is now solved for $u(x)$. In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= \frac{u(x^2 - 2x + 2)}{x(x-1)} \end{aligned}$$

Where $f(x) = \frac{x^2 - 2x + 2}{x(x-1)}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned} \frac{1}{u} du &= \frac{x^2 - 2x + 2}{x(x-1)} dx \\ \int \frac{1}{u} du &= \int \frac{x^2 - 2x + 2}{x(x-1)} dx \\ \ln(u) &= x + \ln(x-1) - 2 \ln(x) + c_1 \\ u &= e^{x + \ln(x-1) - 2 \ln(x) + c_1} \\ &= c_1 e^{x + \ln(x-1) - 2 \ln(x)} \end{aligned}$$

Which simplifies to

$$u(x) = c_1 \left(\frac{e^x}{x} - \frac{e^x}{x^2} \right)$$

Now that $u(x)$ is known, then

$$\begin{aligned} v'(x) &= u(x) \\ v(x) &= \int u(x) dx + c_2 \\ &= \frac{e^x c_1}{x} + c_2 \end{aligned}$$

Hence

$$\begin{aligned} y &= v(x) x^n \\ &= \left(\frac{e^x c_1}{x} + c_2 \right) x \\ &= c_1 e^x + c_2 x \end{aligned}$$

Now the particular solution to this ODE is found

$$(1 - x)y'' + xy' - y = (x - 1)^2$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1y_1 + u_2y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = x$$

$$y_2 = e^x$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} x & e^x \\ \frac{d}{dx}(x) & \frac{d}{dx}(e^x) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} x & e^x \\ 1 & e^x \end{vmatrix}$$

Therefore

$$W = (x)(e^x) - (e^x)(1)$$

Which simplifies to

$$W = x e^x - e^x$$

Which simplifies to

$$W = (x - 1) e^x$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{(x - 1)^2 e^x}{(1 - x)(x - 1) e^x} dx$$

Which simplifies to

$$u_1 = - \int (-1) dx$$

Hence

$$u_1 = x$$

And Eq. (3) becomes

$$u_2 = \int \frac{x(x - 1)^2}{(1 - x)(x - 1) e^x} dx$$

Which simplifies to

$$u_2 = \int -x e^{-x} dx$$

Hence

$$u_2 = (1 + x) e^{-x}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = x^2 + (1 + x) e^{-x} e^x$$

Which simplifies to

$$y_p(x) = x^2 + x + 1$$

Therefore the general solution is

$$\begin{aligned}
 y &= y_h + y_p \\
 &= \left(\left(\frac{e^x c_1}{x} + c_2 \right) x \right) + (x^2 + x + 1) \\
 &= x^2 + x + 1 + \left(\frac{e^x c_1}{x} + c_2 \right) x
 \end{aligned}$$

Which simplifies to

$$y = c_1 e^x + c_2 x + x^2 + x + 1$$

Summary

The solution(s) found are the following

$$y = c_1 e^x + c_2 x + x^2 + x + 1 \quad (1)$$

Verification of solutions

$$y = c_1 e^x + c_2 x + x^2 + x + 1$$

Verified OK.

11.18.2 Solving as second order ode non constant coeff transformation on B ode

Given an ode of the form

$$Ay'' + By' + Cy = F(x)$$

This method reduces the order ode the ODE by one by applying the transformation

$$y = Bv$$

This results in

$$\begin{aligned}
 y' &= B'v + v'B \\
 y'' &= B''v + B'v' + v''B + v'B' \\
 &= v''B + 2v' + B' + B''v
 \end{aligned}$$

And now the original ode becomes

$$\begin{aligned}
 A(v''B + 2v'B' + B''v) + B(B'v + v'B) + CBv &= 0 \\
 ABv'' + (2AB' + B^2)v' + (AB'' + BB' + CB)v &= 0
 \end{aligned} \quad (1)$$

If the term $AB'' + BB' + CB$ is zero, then this method works and can be used to solve

$$ABv'' + (2AB' + B^2)v' = 0$$

By Using $u = v'$ which reduces the order of the above ode to one. The new ode is

$$ABu' + (2AB' + B^2)u = 0$$

The above ode is first order ode which is solved for u . Now a new ode $v' = u$ is solved for v as first order ode. Then the final solution is obtain from $y = Bv$.

This method works only if the term $AB'' + BB' + CB$ is zero. The given ODE shows that

$$\begin{aligned}A &= 1 - x \\B &= x \\C &= -1 \\F &= (x - 1)^2\end{aligned}$$

The above shows that for this ode

$$\begin{aligned}AB'' + BB' + CB &= (1 - x)(0) + (x)(1) + (-1)(x) \\&= 0\end{aligned}$$

Hence the ode in v given in (1) now simplifies to

$$-x(x - 1)v'' + (x^2 - 2x + 2)v' = 0$$

Now by applying $v' = u$ the above becomes

$$(-x^2 + x)u'(x) + (x^2 - 2x + 2)u(x) = 0$$

Which is now solved for u . In canonical form the ODE is

$$\begin{aligned}u' &= F(x, u) \\&= f(x)g(u) \\&= \frac{u(x^2 - 2x + 2)}{x(x - 1)}\end{aligned}$$

Where $f(x) = \frac{x^2-2x+2}{x(x-1)}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u} du &= \frac{x^2 - 2x + 2}{x(x-1)} dx \\ \int \frac{1}{u} du &= \int \frac{x^2 - 2x + 2}{x(x-1)} dx \\ \ln(u) &= x + \ln(x-1) - 2\ln(x) + c_1 \\ u &= e^{x+\ln(x-1)-2\ln(x)+c_1} \\ &= c_1 e^{x+\ln(x-1)-2\ln(x)}\end{aligned}$$

Which simplifies to

$$u(x) = c_1 \left(\frac{e^x}{x} - \frac{e^x}{x^2} \right)$$

The ode for v now becomes

$$\begin{aligned}v' &= u \\ &= c_1 \left(\frac{e^x}{x} - \frac{e^x}{x^2} \right)\end{aligned}$$

Which is now solved for v . Integrating both sides gives

$$\begin{aligned}v(x) &= \int \frac{c_1(x-1)e^x}{x^2} dx \\ &= \frac{e^x c_1}{x} + c_2\end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}y_h(x) &= Bv \\ &= (x) \left(\frac{e^x c_1}{x} + c_2 \right) \\ &= c_1 e^x + c_2 x\end{aligned}$$

And now the particular solution $y_p(x)$ will be found. The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = x$$

$$y_2 = e^x$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} x & e^x \\ \frac{d}{dx}(x) & \frac{d}{dx}(e^x) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} x & e^x \\ 1 & e^x \end{vmatrix}$$

Therefore

$$W = (x)(e^x) - (e^x)(1)$$

Which simplifies to

$$W = x e^x - e^x$$

Which simplifies to

$$W = (x - 1) e^x$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{(x - 1)^2 e^x}{(1 - x)(x - 1) e^x} dx$$

Which simplifies to

$$u_1 = - \int (-1) dx$$

Hence

$$u_1 = x$$

And Eq. (3) becomes

$$u_2 = \int \frac{x(x-1)^2}{(1-x)(x-1)e^x} dx$$

Which simplifies to

$$u_2 = \int -x e^{-x} dx$$

Hence

$$u_2 = (1+x)e^{-x}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = x^2 + (1+x)e^{-x}e^x$$

Which simplifies to

$$y_p(x) = x^2 + x + 1$$

Hence the complete solution is

$$\begin{aligned} y(x) &= y_h + y_p \\ &= (c_1e^x + c_2x) + (x^2 + x + 1) \\ &= c_1e^x + c_2x + x^2 + x + 1 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1e^x + c_2x + x^2 + x + 1 \tag{1}$$

Verification of solutions

$$y = c_1e^x + c_2x + x^2 + x + 1$$

Verified OK.

11.18.3 Solving using Kovacic algorithm

Writing the ode as

$$(1 - x)y'' + xy' - y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 - x \\ B &= x \\ C &= -1 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{x^2 - 4x + 6}{4(x - 1)^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= x^2 - 4x + 6 \\ t &= 4(x - 1)^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{x^2 - 4x + 6}{4(x - 1)^2} \right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 302: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 2 \\ &= 0 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4(x - 1)^2$. There is a pole at $x = 1$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Therefore

$$L = [1, 2]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = \frac{1}{4} + \frac{3}{4(x-1)^2} - \frac{1}{2(x-1)}$$

For the pole at $x = 1$ let b be the coefficient of $\frac{1}{(x-1)^2}$ in the partial fractions decomposition of r given above. Therefore $b = \frac{3}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{3}{2} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = -\frac{1}{2} \end{aligned}$$

Since the order of r at ∞ is $O_r(\infty) = 0$ then

$$v = \frac{-O_r(\infty)}{2} = \frac{0}{2} = 0$$

$[\sqrt{r}]_\infty$ is the sum of terms involving x^i for $0 \leq i \leq v$ in the Laurent series for \sqrt{r} at ∞ . Therefore

$$\begin{aligned} [\sqrt{r}]_\infty &= \sum_{i=0}^v a_i x^i \\ &= \sum_{i=0}^0 a_i x^i \end{aligned} \tag{8}$$

Let a be the coefficient of $x^v = x^0$ in the above sum. The Laurent series of \sqrt{r} at ∞ is

$$\sqrt{r} \approx \frac{1}{2} - \frac{1}{2x} + \frac{1}{x^3} + \frac{11}{4x^4} + \frac{21}{4x^5} + \frac{15}{2x^6} + \frac{6}{x^7} - \frac{117}{16x^8} + \dots \tag{9}$$

Comparing Eq. (9) with Eq. (8) shows that

$$a = \frac{1}{2}$$

From Eq. (9) the sum up to $v = 0$ gives

$$\begin{aligned} [\sqrt{r}]_\infty &= \sum_{i=0}^0 a_i x^i \\ &= \frac{1}{2} \end{aligned} \tag{10}$$

Now we need to find b , where b be the coefficient of $x^{v-1} = x^{-1} = \frac{1}{x}$ in r minus the coefficient of same term but in $([\sqrt{r}]_\infty)^2$ where $[\sqrt{r}]_\infty$ was found above in Eq (10). Hence

$$([\sqrt{r}]_\infty)^2 = \frac{1}{4}$$

This shows that the coefficient of $\frac{1}{x}$ in the above is 0. Now we need to find the coefficient of $\frac{1}{x}$ in r . How this is done depends on if $v = 0$ or not. Since $v = 0$ then starting from $r = \frac{s}{t}$ and doing long division in the form

$$r = Q + \frac{R}{t}$$

Where Q is the quotient and R is the remainder. Then the coefficient of $\frac{1}{x}$ in r will be the coefficient in R of the term in x of degree of t minus one, divided by the leading coefficient in t . Doing long division gives

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{x^2 - 4x + 6}{4x^2 - 8x + 4} \\ &= Q + \frac{R}{4x^2 - 8x + 4} \\ &= \left(\frac{1}{4}\right) + \left(\frac{-2x + 5}{4x^2 - 8x + 4}\right) \\ &= \frac{1}{4} + \frac{-2x + 5}{4x^2 - 8x + 4} \end{aligned}$$

Since the degree of t is 2, then we see that the coefficient of the term x in the remainder R is -2 . Dividing this by leading coefficient in t which is 4 gives $-\frac{1}{2}$. Now b can be found.

$$\begin{aligned} b &= \left(-\frac{1}{2}\right) - (0) \\ &= -\frac{1}{2} \end{aligned}$$

Hence

$$\begin{aligned} [\sqrt{r}]_{\infty} &= \frac{1}{2} \\ \alpha_{\infty}^+ &= \frac{1}{2} \left(\frac{b}{a} - v\right) = \frac{1}{2} \left(\frac{-\frac{1}{2}}{\frac{1}{2}} - 0\right) = -\frac{1}{2} \\ \alpha_{\infty}^- &= \frac{1}{2} \left(-\frac{b}{a} - v\right) = \frac{1}{2} \left(-\frac{-\frac{1}{2}}{\frac{1}{2}} - 0\right) = \frac{1}{2} \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{x^2 - 4x + 6}{4(x - 1)^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
1	2	0	$\frac{3}{2}$	$-\frac{1}{2}$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
0	$\frac{1}{2}$	$-\frac{1}{2}$	$\frac{1}{2}$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^+ = -\frac{1}{2}$ then

$$\begin{aligned} d &= \alpha_\infty^+ - (\alpha_{c_1}^-) \\ &= -\frac{1}{2} - \left(-\frac{1}{2}\right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c) [\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty) [\sqrt{r}]_\infty$$

Substituting the above values in the above results in

$$\begin{aligned} \omega &= \left((-) [\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{x - c_1} \right) + (+) [\sqrt{r}]_\infty \\ &= -\frac{1}{2(x-1)} + \left(\frac{1}{2}\right) \\ &= -\frac{1}{2(x-1)} + \frac{1}{2} \\ &= \frac{-2+x}{2x-2} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \tag{1A}$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$(0) + 2\left(-\frac{1}{2(x-1)} + \frac{1}{2}\right) (0) + \left(\left(\frac{1}{2(x-1)^2}\right) + \left(-\frac{1}{2(x-1)} + \frac{1}{2}\right)^2 - \left(\frac{x^2 - 4x + 6}{4(x-1)^2}\right)\right) = 0$$
$$0 = 0$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= pe^{\int \omega dx} \\ &= e^{\int \left(-\frac{1}{2(x-1)} + \frac{1}{2}\right) dx} \\ &= \frac{e^{\frac{x}{2}}}{\sqrt{x-1}} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{x}{1-x} dx} \\ &= z_1 e^{\frac{x}{2} + \frac{\ln(x-1)}{2}} \\ &= z_1 (\sqrt{x-1} e^{\frac{x}{2}}) \end{aligned}$$

Which simplifies to

$$y_1 = e^x$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{x}{1-x} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{x + \ln(x-1)}}{(y_1)^2} dx \\ &= y_1 (-x e^{-x}) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1(e^x) + c_2(e^x(-x e^{-x}))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$(1 - x)y'' + xy' - y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^x - c_2 x$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = e^x$$

$$y_2 = -x$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^x & -x \\ \frac{d}{dx}(e^x) & \frac{d}{dx}(-x) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^x & -x \\ e^x & -1 \end{vmatrix}$$

Therefore

$$W = (e^x)(-1) - (-x)(e^x)$$

Which simplifies to

$$W = x e^x - e^x$$

Which simplifies to

$$W = (x - 1) e^x$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{-x(x-1)^2}{(1-x)(x-1)e^x} dx$$

Which simplifies to

$$u_1 = - \int x e^{-x} dx$$

Hence

$$u_1 = (1+x) e^{-x}$$

And Eq. (3) becomes

$$u_2 = \int \frac{(x-1)^2 e^x}{(1-x)(x-1)e^x} dx$$

Which simplifies to

$$u_2 = \int (-1) dx$$

Hence

$$u_2 = -x$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = x^2 + (1 + x) e^{-x} e^x$$

Which simplifies to

$$y_p(x) = x^2 + x + 1$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^x - c_2 x) + (x^2 + x + 1) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^x - c_2 x + x^2 + x + 1 \tag{1}$$

Verification of solutions

$$y = c_1 e^x - c_2 x + x^2 + x + 1$$

Verified OK.

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Reducible group (found an exponential solution)
    Reducible group (found another exponential solution)
<- Kovacics algorithm successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve((1-x)*diff(y(x),x$2)+x*diff(y(x),x)-y(x)=(1-x)^2,y(x), singsol=all)
```

$$y(x) = c_2x + e^x c_1 + x^2 + 1$$

✓ Solution by Mathematica

Time used: 0.087 (sec). Leaf size: 22

```
DSolve[(1-x)*y''[x]+x*y'[x]-y[x]==(1-x)^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^2 + x - c_2x + c_1e^x + 1$$

11.19 problem 5(d)

- 11.19.1 Solving as second order ode non constant coeff transformation on B ode 2066
- 11.19.2 Solving using Kovacic algorithm 2071

Internal problem ID [6335]

Internal file name [OUTPUT/5583_Sunday_June_05_2022_03_44_09_PM_29873215/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.3. THE METHOD OF VARIATION OF PARAMETERS. Page 71

Problem number: 5(d).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_ode_non_constant_coeff_transformation_on_B"

Maple gives the following as the ode type

```
[[_2nd_order, _with_linear_symmetries]]
```

$$xy'' - (1 + x)y' + y = x^2e^{2x}$$

11.19.1 Solving as second order ode non constant coeff transformation on B ode

Given an ode of the form

$$Ay'' + By' + Cy = F(x)$$

This method reduces the order ode the ODE by one by applying the transformation

$$y = Bv$$

This results in

$$\begin{aligned}y' &= B'v + v'B \\y'' &= B''v + B'v' + v''B + v'B' \\ &= v''B + 2v' + B' + B''v\end{aligned}$$

And now the original ode becomes

$$\begin{aligned} A(v''B + 2v'B' + B''v) + B(B'v + v'B) + CBv &= 0 \\ ABv'' + (2AB' + B^2)v' + (AB'' + BB' + CB)v &= 0 \end{aligned} \quad (1)$$

If the term $AB'' + BB' + CB$ is zero, then this method works and can be used to solve

$$ABv'' + (2AB' + B^2)v' = 0$$

By Using $u = v'$ which reduces the order of the above ode to one. The new ode is

$$ABu' + (2AB' + B^2)u = 0$$

The above ode is first order ode which is solved for u . Now a new ode $v' = u$ is solved for v as first order ode. Then the final solution is obtain from $y = Bv$.

This method works only if the term $AB'' + BB' + CB$ is zero. The given ODE shows that

$$\begin{aligned} A &= x \\ B &= -1 - x \\ C &= 1 \\ F &= x^2e^{2x} \end{aligned}$$

The above shows that for this ode

$$\begin{aligned} AB'' + BB' + CB &= (x)(0) + (-1 - x)(-1) + (1)(-1 - x) \\ &= 0 \end{aligned}$$

Hence the ode in v given in (1) now simplifies to

$$-x(1 + x)v'' + (x^2 + 1)v' = 0$$

Now by applying $v' = u$ the above becomes

$$-x(1 + x)u'(x) + (x^2 + 1)u(x) = 0$$

Which is now solved for u . In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= \frac{(x^2 + 1)u}{x(1 + x)} \end{aligned}$$

Where $f(x) = \frac{x^2+1}{x(1+x)}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u} du &= \frac{x^2 + 1}{x(1+x)} dx \\ \int \frac{1}{u} du &= \int \frac{x^2 + 1}{x(1+x)} dx \\ \ln(u) &= x - 2 \ln(1+x) + \ln(x) + c_1 \\ u &= e^{x-2 \ln(1+x)+\ln(x)+c_1} \\ &= c_1 e^{x-2 \ln(1+x)+\ln(x)}\end{aligned}$$

Which simplifies to

$$u(x) = \frac{c_1 x e^x}{(1+x)^2}$$

The ode for v now becomes

$$\begin{aligned}v' &= u \\ &= \frac{c_1 x e^x}{(1+x)^2}\end{aligned}$$

Which is now solved for v . Integrating both sides gives

$$\begin{aligned}v(x) &= \int \frac{c_1 x e^x}{(1+x)^2} dx \\ &= \frac{c_1 e^x}{1+x} + c_2\end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}y_h(x) &= Bv \\ &= (-1-x) \left(\frac{c_1 e^x}{1+x} + c_2 \right) \\ &= -c_1 e^x - (1+x) c_2\end{aligned}$$

And now the particular solution $y_p(x)$ will be found. The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = -1 - x$$

$$y_2 = e^x$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} -1 - x & e^x \\ \frac{d}{dx}(-1 - x) & \frac{d}{dx}(e^x) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} -1 - x & e^x \\ -1 & e^x \end{vmatrix}$$

Therefore

$$W = (-1 - x)(e^x) - (e^x)(-1)$$

Which simplifies to

$$W = -x e^x$$

Which simplifies to

$$W = -x e^x$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{e^x x^2 e^{2x}}{-x^2 e^x} dx$$

Which simplifies to

$$u_1 = - \int -e^{2x} dx$$

Hence

$$u_1 = \frac{e^{2x}}{2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{(-1-x)x^2 e^{2x}}{-x^2 e^x} dx$$

Which simplifies to

$$u_2 = \int e^x(1+x) dx$$

Hence

$$u_2 = x e^x$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{e^{2x}(-1-x)}{2} + e^{2x}x$$

Which simplifies to

$$y_p(x) = \frac{(x-1)e^{2x}}{2}$$

Hence the complete solution is

$$\begin{aligned} y(x) &= y_h + y_p \\ &= (-c_1 e^x - (1+x)c_2) + \left(\frac{(x-1)e^{2x}}{2} \right) \\ &= -c_1 e^x - (1+x)c_2 + \frac{(x-1)e^{2x}}{2} \end{aligned}$$

Summary

The solution(s) found are the following

$$y = -c_1 e^x - (1+x)c_2 + \frac{(x-1)e^{2x}}{2} \tag{1}$$

Verification of solutions

$$y = -c_1 e^x - (1+x)c_2 + \frac{(x-1)e^{2x}}{2}$$

Verified OK.

11.19.2 Solving using Kovacic algorithm

Writing the ode as

$$xy'' + (-1-x)y' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= x \\ B &= -1-x \\ C &= 1 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{x^2 - 2x + 3}{4x^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= x^2 - 2x + 3 \\ t &= 4x^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{x^2 - 2x + 3}{4x^2} \right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 303: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 2 \\ &= 0 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Therefore

$$L = [1, 2]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = \frac{1}{4} - \frac{1}{2x} + \frac{3}{4x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = \frac{3}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{3}{2} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = -\frac{1}{2} \end{aligned}$$

Since the order of r at ∞ is $O_r(\infty) = 0$ then

$$v = \frac{-O_r(\infty)}{2} = \frac{0}{2} = 0$$

$[\sqrt{r}]_\infty$ is the sum of terms involving x^i for $0 \leq i \leq v$ in the Laurent series for \sqrt{r} at ∞ . Therefore

$$\begin{aligned} [\sqrt{r}]_\infty &= \sum_{i=0}^v a_i x^i \\ &= \sum_{i=0}^0 a_i x^i \end{aligned} \tag{8}$$

Let a be the coefficient of $x^v = x^0$ in the above sum. The Laurent series of \sqrt{r} at ∞ is

$$\sqrt{r} \approx \frac{1}{2} - \frac{1}{2x} + \frac{1}{2x^2} + \frac{1}{2x^3} + \frac{1}{4x^4} - \frac{1}{4x^5} - \frac{3}{4x^6} - \frac{3}{4x^7} + \dots \tag{9}$$

Comparing Eq. (9) with Eq. (8) shows that

$$a = \frac{1}{2}$$

From Eq. (9) the sum up to $v = 0$ gives

$$\begin{aligned} [\sqrt{r}]_\infty &= \sum_{i=0}^0 a_i x^i \\ &= \frac{1}{2} \end{aligned} \tag{10}$$

Now we need to find b , where b be the coefficient of $x^{v-1} = x^{-1} = \frac{1}{x}$ in r minus the coefficient of same term but in $([\sqrt{r}]_\infty)^2$ where $[\sqrt{r}]_\infty$ was found above in Eq (10). Hence

$$([\sqrt{r}]_\infty)^2 = \frac{1}{4}$$

This shows that the coefficient of $\frac{1}{x}$ in the above is 0. Now we need to find the coefficient of $\frac{1}{x}$ in r . How this is done depends on if $v = 0$ or not. Since $v = 0$ then starting from $r = \frac{s}{t}$ and doing long division in the form

$$r = Q + \frac{R}{t}$$

Where Q is the quotient and R is the remainder. Then the coefficient of $\frac{1}{x}$ in r will be the coefficient in R of the term in x of degree of t minus one, divided by the leading coefficient in t . Doing long division gives

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{x^2 - 2x + 3}{4x^2} \\ &= Q + \frac{R}{4x^2} \\ &= \left(\frac{1}{4}\right) + \left(\frac{-2x + 3}{4x^2}\right) \\ &= \frac{1}{4} + \frac{-2x + 3}{4x^2} \end{aligned}$$

Since the degree of t is 2, then we see that the coefficient of the term x in the remainder R is -2 . Dividing this by leading coefficient in t which is 4 gives $-\frac{1}{2}$. Now b can be found.

$$\begin{aligned} b &= \left(-\frac{1}{2}\right) - (0) \\ &= -\frac{1}{2} \end{aligned}$$

Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= \frac{1}{2} \\ \alpha_\infty^+ &= \frac{1}{2} \left(\frac{b}{a} - v\right) = \frac{1}{2} \left(\frac{-\frac{1}{2}}{\frac{1}{2}} - 0\right) = -\frac{1}{2} \\ \alpha_\infty^- &= \frac{1}{2} \left(-\frac{b}{a} - v\right) = \frac{1}{2} \left(-\frac{-\frac{1}{2}}{\frac{1}{2}} - 0\right) = \frac{1}{2} \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{x^2 - 2x + 3}{4x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	$\frac{3}{2}$	$-\frac{1}{2}$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
0	$\frac{1}{2}$	$-\frac{1}{2}$	$\frac{1}{2}$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^+ = -\frac{1}{2}$ then

$$\begin{aligned} d &= \alpha_\infty^+ - (\alpha_{c_1}^-) \\ &= -\frac{1}{2} - \left(-\frac{1}{2}\right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_\infty$$

Substituting the above values in the above results in

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{x - c_1} \right) + (+)[\sqrt{r}]_\infty \\ &= -\frac{1}{2x} + \left(\frac{1}{2}\right) \\ &= \frac{1}{2} - \frac{1}{2x} \\ &= \frac{x - 1}{2x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$(0) + 2\left(\frac{1}{2} - \frac{1}{2x}\right) (0) + \left(\left(\frac{1}{2x^2}\right) + \left(\frac{1}{2} - \frac{1}{2x}\right)^2 - \left(\frac{x^2 - 2x + 3}{4x^2}\right)\right) = 0$$

$$0 = 0$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= p e^{\int \omega dx} \\ &= e^{\int \left(\frac{1}{2} - \frac{1}{2x}\right) dx} \\ &= \frac{e^{\frac{x}{2}}}{\sqrt{x}} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-1-x}{x} dx} \\ &= z_1 e^{\frac{x}{2} + \frac{\ln(x)}{2}} \\ &= z_1 (\sqrt{x} e^{\frac{x}{2}}) \end{aligned}$$

Which simplifies to

$$y_1 = e^x$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{1-x}{x} dx}}{(y_1)^2} dx \\&= y_1 \int \frac{e^{x+\ln(x)}}{(y_1)^2} dx \\&= y_1 (-(1+x)e^{-x})\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\&= c_1(e^x) + c_2(e^x(-(1+x)e^{-x}))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$xy'' + (-1-x)y' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^x + (-1-x)c_2$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned}y_1 &= e^x \\y_2 &= -1-x\end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^x & -1-x \\ \frac{d}{dx}(e^x) & \frac{d}{dx}(-1-x) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^x & -1-x \\ e^x & -1 \end{vmatrix}$$

Therefore

$$W = (e^x)(-1) - (-1-x)(e^x)$$

Which simplifies to

$$W = x e^x$$

Which simplifies to

$$W = x e^x$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{(-1-x)x^2 e^{2x}}{x^2 e^x} dx$$

Which simplifies to

$$u_1 = - \int -e^x(1+x) dx$$

Hence

$$u_1 = x e^x$$

And Eq. (3) becomes

$$u_2 = \int \frac{e^x x^2 e^{2x}}{x^2 e^x} dx$$

Which simplifies to

$$u_2 = \int e^{2x} dx$$

Hence

$$u_2 = \frac{e^{2x}}{2}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{e^{2x}(-1-x)}{2} + e^{2x}x$$

Which simplifies to

$$y_p(x) = \frac{(x-1)e^{2x}}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^x + (-1-x)c_2) + \left(\frac{(x-1)e^{2x}}{2} \right) \end{aligned}$$

Which simplifies to

$$y = c_1 e^x - (1+x)c_2 + \frac{(x-1)e^{2x}}{2}$$

Summary

The solution(s) found are the following

$$y = c_1 e^x - (1+x)c_2 + \frac{(x-1)e^{2x}}{2} \tag{1}$$

Verification of solutions

$$y = c_1 e^x - (1+x)c_2 + \frac{(x-1)e^{2x}}{2}$$

Verified OK.

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Reducible group (found an exponential solution)
    Reducible group (found another exponential solution)
<- Kovacics algorithm successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 23

```
dsolve(x*diff(y(x),x$2)-(1+x)*diff(y(x),x)+y(x)=x^2*exp(2*x),y(x), singsol=all)
```

$$y(x) = (x + 1)c_2 + e^x c_1 + \frac{(x - 1)e^{2x}}{2}$$

✓ Solution by Mathematica

Time used: 0.066 (sec). Leaf size: 31

```
DSolve[x*y''[x]-(1+x)*y'[x]+y[x]==x^2*Exp[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}e^{2x}(x - 1) + c_1 e^x - c_2(x + 1)$$

11.20 problem 5(e)

11.20.1 Solving as second order euler ode	2082
11.20.2 Solving as linear second order ode solved by an integrating factor ode	2085
11.20.3 Solving as second order change of variable on x method 2 ode	2086
11.20.4 Solving as second order change of variable on x method 1 ode	2092
11.20.5 Solving as second order change of variable on y method 1 ode	2096
11.20.6 Solving as second order change of variable on y method 2 ode	2101
11.20.7 Solving as second order ode non constant coeff transformation on B ode	2106
11.20.8 Solving using Kovacic algorithm	2110

Internal problem ID [6336]

Internal file name [OUTPUT/5584_Sunday_June_05_2022_03_44_11_PM_30279075/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.3. THE METHOD OF VARIATION OF PARAMETERS. Page 71

Problem number: 5(e).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_euler_ode", "second_order_change_of_variable_on_x_method_1", "second_order_change_of_variable_on_x_method_2", "second_order_change_of_variable_on_y_method_1", "second_order_change_of_variable_on_y_method_2", "linear_second_order_ode_solved_by_an_integrating_factor", "second_order_ode_non_constant_coeff_transformation_on_B"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$x^2 y'' - 2xy' + 2y = x e^{-x}$$

11.20.1 Solving as second order euler ode ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = x^2$, $B = -2x$, $C = 2$, $f(x) = x e^{-x}$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$x^2y'' - 2xy' + 2y = 0$$

This is Euler second order ODE. Let the solution be $y = x^r$, then $y' = rx^{r-1}$ and $y'' = r(r-1)x^{r-2}$. Substituting these back into the given ODE gives

$$x^2(r(r-1))x^{r-2} - 2xr x^{r-1} + 2x^r = 0$$

Simplifying gives

$$r(r-1)x^r - 2rx^r + 2x^r = 0$$

Since $x^r \neq 0$ then dividing throughout by x^r gives

$$r(r-1) - 2r + 2 = 0$$

Or

$$r^2 - 3r + 2 = 0 \tag{1}$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = 1$$

$$r_2 = 2$$

Since the roots are real and distinct, then the general solution is

$$y = c_1y_1 + c_2y_2$$

Where $y_1 = x^{r_1}$ and $y_2 = x^{r_2}$. Hence

$$y = c_2x^2 + c_1x$$

Next, we find the particular solution to the ODE

$$x^2y'' - 2xy' + 2y = x e^{-x}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1y_1 + u_2y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = x$$

$$y_2 = x^2$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} x & x^2 \\ \frac{d}{dx}(x) & \frac{d}{dx}(x^2) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} x & x^2 \\ 1 & 2x \end{vmatrix}$$

Therefore

$$W = (x)(2x) - (x^2)(1)$$

Which simplifies to

$$W = x^2$$

Which simplifies to

$$W = x^2$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^3 e^{-x}}{x^4} dx$$

Which simplifies to

$$u_1 = - \int \frac{e^{-x}}{x} dx$$

Hence

$$u_1 = \text{expIntegral}_1(x)$$

And Eq. (3) becomes

$$u_2 = \int \frac{x^2 e^{-x}}{x^4} dx$$

Which simplifies to

$$u_2 = \int \frac{e^{-x}}{x^2} dx$$

Hence

$$u_2 = -\frac{e^{-x}}{x} + \text{expIntegral}_1(x)$$

Which simplifies to

$$u_1 = \text{expIntegral}_1(x)$$
$$u_2 = \frac{\text{expIntegral}_1(x) x - e^{-x}}{x}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \text{expIntegral}_1(x) x + (\text{expIntegral}_1(x) x - e^{-x}) x$$

Which simplifies to

$$y_p(x) = x(-e^{-x} + (1 + x) \text{expIntegral}_1(x))$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (-e^{-x} + (1+x) \exp \int_1^x (x) dx + c_2 x + c_1) x \end{aligned}$$

Summary

The solution(s) found are the following

$$y = (-e^{-x} + (1+x) \exp \int_1^x (x) dx + c_2 x + c_1) x \quad (1)$$

Verification of solutions

$$y = (-e^{-x} + (1+x) \exp \int_1^x (x) dx + c_2 x + c_1) x$$

Verified OK.

11.20.2 Solving as linear second order ode solved by an integrating factor ode

The ode satisfies this form

$$y'' + p(x)y' + \frac{(p(x)^2 + p'(x))y}{2} = f(x)$$

Where $p(x) = -\frac{2}{x}$. Therefore, there is an integrating factor given by

$$\begin{aligned} M(x) &= e^{\frac{1}{2} \int p dx} \\ &= e^{\int -\frac{2}{x} dx} \\ &= \frac{1}{x} \end{aligned}$$

Multiplying both sides of the ODE by the integrating factor $M(x)$ makes the left side of the ODE a complete differential

$$\begin{aligned} (M(x)y)'' &= \frac{e^{-x}}{x^2} \\ \left(\frac{y}{x}\right)'' &= \frac{e^{-x}}{x^2} \end{aligned}$$

Integrating once gives

$$\left(\frac{y}{x}\right)' = \frac{\exp \int_1^x (x) dx - e^{-x}}{x} + c_1$$

Integrating again gives

$$\left(\frac{y}{x}\right) = \exp\int_1(x) x + \exp\int_1(x) - e^{-x} + c_1x + c_2$$

Hence the solution is

$$y = \frac{\exp\int_1(x) x + \exp\int_1(x) - e^{-x} + c_1x + c_2}{\frac{1}{x}}$$

Or

$$y = c_1x^2 + x^2 \exp\int_1(x) + c_2x + x \exp\int_1(x) - x e^{-x}$$

Summary

The solution(s) found are the following

$$y = c_1x^2 + x^2 \exp\int_1(x) + c_2x + x \exp\int_1(x) - x e^{-x} \quad (1)$$

Verification of solutions

$$y = c_1x^2 + x^2 \exp\int_1(x) + c_2x + x \exp\int_1(x) - x e^{-x}$$

Verified OK.

11.20.3 Solving as second order change of variable on x method 2 ode

This is second order non-homogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$.
 y_h is the solution to

$$x^2y'' - 2xy' + 2y = 0$$

In normal form the ode

$$x^2y'' - 2xy' + 2y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = -\frac{2}{x}$$
$$q(x) = \frac{2}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x)\tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned} \tau &= \int e^{-(\int p(x)dx)} dx \\ &= \int e^{-(\int -\frac{2}{x}dx)} dx \\ &= \int e^{2\ln(x)} dx \\ &= \int x^2 dx \\ &= \frac{x^3}{3} \end{aligned} \quad (6)$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned} q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\ &= \frac{\frac{2}{x^2}}{x^4} \\ &= \frac{2}{x^6} \end{aligned} \quad (7)$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned}\frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) + \frac{2y(\tau)}{x^6} &= 0\end{aligned}$$

But in terms of τ

$$\frac{2}{x^6} = \frac{2}{9\tau^2}$$

Hence the above ode becomes

$$\frac{d^2}{d\tau^2}y(\tau) + \frac{2y(\tau)}{9\tau^2} = 0$$

The above ode is now solved for $y(\tau)$. The ode can be written as

$$9\left(\frac{d^2}{d\tau^2}y(\tau)\right)\tau^2 + 2y(\tau) = 0$$

Which shows it is a Euler ODE. This is Euler second order ODE. Let the solution be $y(\tau) = \tau^r$, then $y' = r\tau^{r-1}$ and $y'' = r(r-1)\tau^{r-2}$. Substituting these back into the given ODE gives

$$9\tau^2(r(r-1))\tau^{r-2} + 0r\tau^{r-1} + 2\tau^r = 0$$

Simplifying gives

$$9r(r-1)\tau^r + 0\tau^r + 2\tau^r = 0$$

Since $\tau^r \neq 0$ then dividing throughout by τ^r gives

$$9r(r-1) + 0 + 2 = 0$$

Or

$$9r^2 - 9r + 2 = 0 \tag{1}$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$\begin{aligned}r_1 &= \frac{1}{3} \\ r_2 &= \frac{2}{3}\end{aligned}$$

Since the roots are real and distinct, then the general solution is

$$y(\tau) = c_1y_1 + c_2y_2$$

Where $y_1 = \tau^{r_1}$ and $y_2 = \tau^{r_2}$. Hence

$$y(\tau) = c_1\tau^{\frac{1}{3}} + c_2\tau^{\frac{2}{3}}$$

The above solution is now transformed back to y using (6) which results in

$$y = \frac{c_1 3^{\frac{2}{3}} (x^3)^{\frac{1}{3}}}{3} + \frac{c_2 3^{\frac{1}{3}} (x^3)^{\frac{2}{3}}}{3}$$

Therefore the homogeneous solution y_h is

$$y_h = \frac{c_1 3^{\frac{2}{3}} (x^3)^{\frac{1}{3}}}{3} + \frac{c_2 3^{\frac{1}{3}} (x^3)^{\frac{2}{3}}}{3}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = (x^3)^{\frac{1}{3}}$$

$$y_2 = (x^3)^{\frac{2}{3}}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} (x^3)^{\frac{1}{3}} & (x^3)^{\frac{2}{3}} \\ \frac{d}{dx} \left((x^3)^{\frac{1}{3}} \right) & \frac{d}{dx} \left((x^3)^{\frac{2}{3}} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} (x^3)^{\frac{1}{3}} & (x^3)^{\frac{2}{3}} \\ \frac{x^2}{(x^3)^{\frac{2}{3}}} & \frac{2x^2}{(x^3)^{\frac{1}{3}}} \end{vmatrix}$$

Therefore

$$W = \left((x^3)^{\frac{1}{3}} \right) \left(\frac{2x^2}{(x^3)^{\frac{1}{3}}} \right) - \left((x^3)^{\frac{2}{3}} \right) \left(\frac{x^2}{(x^3)^{\frac{2}{3}}} \right)$$

Which simplifies to

$$W = x^2$$

Which simplifies to

$$W = x^2$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{(x^3)^{\frac{2}{3}} x e^{-x}}{x^4} dx$$

Which simplifies to

$$u_1 = - \int \frac{(x^3)^{\frac{2}{3}} e^{-x}}{x^3} dx$$

Hence

$$u_1 = \frac{(x^3)^{\frac{2}{3}} \text{expIntegral}_1(x)}{x^2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{(x^3)^{\frac{1}{3}} x e^{-x}}{x^4} dx$$

Which simplifies to

$$u_2 = \int \frac{(x^3)^{\frac{1}{3}} e^{-x}}{x^3} dx$$

Hence

$$u_2 = -\frac{(x^3)^{\frac{1}{3}}(-x \operatorname{expIntegral}_1(x) + e^{-x})}{x^2}$$

Which simplifies to

$$u_1 = \frac{(x^3)^{\frac{2}{3}} \operatorname{expIntegral}_1(x)}{x^2}$$
$$u_2 = \frac{(x^3)^{\frac{1}{3}}(x \operatorname{expIntegral}_1(x) - e^{-x})}{x^2}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = x \operatorname{expIntegral}_1(x) + (x \operatorname{expIntegral}_1(x) - e^{-x})x$$

Which simplifies to

$$y_p(x) = x(-e^{-x} + (1+x) \operatorname{expIntegral}_1(x))$$

Therefore the general solution is

$$y = y_h + y_p$$
$$= \left(\frac{c_1 3^{\frac{2}{3}}(x^3)^{\frac{1}{3}}}{3} + \frac{c_2 3^{\frac{1}{3}}(x^3)^{\frac{2}{3}}}{3} \right) + (x(-e^{-x} + (1+x) \operatorname{expIntegral}_1(x)))$$

Summary

The solution(s) found are the following

$$y = \frac{c_1 3^{\frac{2}{3}}(x^3)^{\frac{1}{3}}}{3} + \frac{c_2 3^{\frac{1}{3}}(x^3)^{\frac{2}{3}}}{3} + x(-e^{-x} + (1+x) \operatorname{expIntegral}_1(x)) \quad (1)$$

Verification of solutions

$$y = \frac{c_1 3^{\frac{2}{3}}(x^3)^{\frac{1}{3}}}{3} + \frac{c_2 3^{\frac{1}{3}}(x^3)^{\frac{2}{3}}}{3} + x(-e^{-x} + (1+x) \operatorname{expIntegral}_1(x))$$

Verified OK.

11.20.4 Solving as second order change of variable on x method 1 ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = x^2$, $B = -2x$, $C = 2$, $f(x) = xe^{-x}$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$x^2y'' - 2xy' + 2y = 0$$

In normal form the ode

$$x^2y'' - 2xy' + 2y = 0 \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \tag{2}$$

Where

$$p(x) = -\frac{2}{x}$$

$$q(x) = \frac{2}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) results

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \tag{3}$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \tag{4}$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \tag{5}$$

Let $q_1 = c^2$ where c is some constant. Therefore from (5)

$$\begin{aligned} \tau' &= \frac{1}{c}\sqrt{q} \\ &= \frac{\sqrt{2}\sqrt{\frac{1}{x^2}}}{c} \\ \tau'' &= -\frac{\sqrt{2}}{c\sqrt{\frac{1}{x^2}}x^3} \end{aligned} \tag{6}$$

Substituting the above into (4) results in

$$\begin{aligned}
 p_1(\tau) &= \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \\
 &= \frac{-\frac{\sqrt{2}}{c\sqrt{\frac{1}{x^2}}x^3} - \frac{2}{x}\frac{\sqrt{2}\sqrt{\frac{1}{x^2}}}{c}}{\left(\frac{\sqrt{2}\sqrt{\frac{1}{x^2}}}{c}\right)^2} \\
 &= -\frac{3c\sqrt{2}}{2}
 \end{aligned}$$

Therefore ode (3) now becomes

$$\begin{aligned}
 y(\tau)'' + p_1y(\tau)' + q_1y(\tau) &= 0 \\
 \frac{d^2}{d\tau^2}y(\tau) - \frac{3c\sqrt{2}}{2}\left(\frac{d}{d\tau}y(\tau)\right) + c^2y(\tau) &= 0 \tag{7}
 \end{aligned}$$

The above ode is now solved for $y(\tau)$. Since the ode is now constant coefficients, it can be easily solved to give

$$y(\tau) = e^{\frac{3\sqrt{2}c\tau}{4}} \left(c_1 \cosh\left(\frac{\sqrt{2}c\tau}{4}\right) + ic_2 \sinh\left(\frac{\sqrt{2}c\tau}{4}\right) \right)$$

Now from (6)

$$\begin{aligned}
 \tau &= \int \frac{1}{c}\sqrt{q} dx \\
 &= \frac{\int \sqrt{2}\sqrt{\frac{1}{x^2}} dx}{c} \\
 &= \frac{\sqrt{2}\sqrt{\frac{1}{x^2}} x \ln(x)}{c}
 \end{aligned}$$

Substituting the above into the solution obtained gives

$$y = x^{\frac{3}{2}} \left(c_1 \cosh\left(\frac{\ln(x)}{2}\right) + ic_2 \sinh\left(\frac{\ln(x)}{2}\right) \right)$$

Now the particular solution to this ODE is found

$$x^2y'' - 2xy' + 2y = xe^{-x}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of

parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = (x^3)^{\frac{1}{3}}$$

$$y_2 = (x^3)^{\frac{2}{3}}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} (x^3)^{\frac{1}{3}} & (x^3)^{\frac{2}{3}} \\ \frac{d}{dx} \left((x^3)^{\frac{1}{3}} \right) & \frac{d}{dx} \left((x^3)^{\frac{2}{3}} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} (x^3)^{\frac{1}{3}} & (x^3)^{\frac{2}{3}} \\ \frac{x^2}{(x^3)^{\frac{2}{3}}} & \frac{2x^2}{(x^3)^{\frac{1}{3}}} \end{vmatrix}$$

Therefore

$$W = \left((x^3)^{\frac{1}{3}} \right) \left(\frac{2x^2}{(x^3)^{\frac{1}{3}}} \right) - \left((x^3)^{\frac{2}{3}} \right) \left(\frac{x^2}{(x^3)^{\frac{2}{3}}} \right)$$

Which simplifies to

$$W = x^2$$

Which simplifies to

$$W = x^2$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{(x^3)^{\frac{2}{3}} x e^{-x}}{x^4} dx$$

Which simplifies to

$$u_1 = - \int \frac{(x^3)^{\frac{2}{3}} e^{-x}}{x^3} dx$$

Hence

$$u_1 = \frac{(x^3)^{\frac{2}{3}} \text{expIntegral}_1(x)}{x^2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{(x^3)^{\frac{1}{3}} x e^{-x}}{x^4} dx$$

Which simplifies to

$$u_2 = \int \frac{(x^3)^{\frac{1}{3}} e^{-x}}{x^3} dx$$

Hence

$$u_2 = - \frac{(x^3)^{\frac{1}{3}} (-x \text{expIntegral}_1(x) + e^{-x})}{x^2}$$

Which simplifies to

$$u_1 = \frac{(x^3)^{\frac{2}{3}} \text{expIntegral}_1(x)}{x^2}$$
$$u_2 = \frac{(x^3)^{\frac{1}{3}} (x \text{expIntegral}_1(x) - e^{-x})}{x^2}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = x \text{expIntegral}_1(x) + (x \text{expIntegral}_1(x) - e^{-x}) x$$

Which simplifies to

$$y_p(x) = x(-e^{-x} + (1+x) \exp \text{Integral}_1(x))$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(x^{\frac{3}{2}} \left(c_1 \cosh \left(\frac{\ln(x)}{2} \right) + ic_2 \sinh \left(\frac{\ln(x)}{2} \right) \right) \right) + (x(-e^{-x} + (1+x) \exp \text{Integral}_1(x))) \\ &= x(-e^{-x} + (1+x) \exp \text{Integral}_1(x)) + x^{\frac{3}{2}} \left(c_1 \cosh \left(\frac{\ln(x)}{2} \right) + ic_2 \sinh \left(\frac{\ln(x)}{2} \right) \right) \end{aligned}$$

Which simplifies to

$$y = x(-e^{-x} + (1+x) \exp \text{Integral}_1(x)) + x^{\frac{3}{2}} \left(c_1 \cosh \left(\frac{\ln(x)}{2} \right) + ic_2 \sinh \left(\frac{\ln(x)}{2} \right) \right)$$

Summary

The solution(s) found are the following

$$y = x(-e^{-x} + (1+x) \exp \text{Integral}_1(x)) + x^{\frac{3}{2}} \left(c_1 \cosh \left(\frac{\ln(x)}{2} \right) + ic_2 \sinh \left(\frac{\ln(x)}{2} \right) \right) \quad (1)$$

Verification of solutions

$$y = x(-e^{-x} + (1+x) \exp \text{Integral}_1(x)) + x^{\frac{3}{2}} \left(c_1 \cosh \left(\frac{\ln(x)}{2} \right) + ic_2 \sinh \left(\frac{\ln(x)}{2} \right) \right)$$

Verified OK.

11.20.5 Solving as second order change of variable on y method 1 ode

This is second order non-homogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$.
 y_h is the solution to

$$x^2y'' - 2xy' + 2y = 0$$

In normal form the given ode is written as

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = -\frac{2}{x}$$
$$q(x) = \frac{2}{x^2}$$

Calculating the Liouville ode invariant Q given by

$$\begin{aligned} Q &= q - \frac{p'}{2} - \frac{p^2}{4} \\ &= \frac{2}{x^2} - \frac{\left(-\frac{2}{x}\right)'}{2} - \frac{\left(-\frac{2}{x}\right)^2}{4} \\ &= \frac{2}{x^2} - \frac{\left(\frac{2}{x^2}\right)}{2} - \frac{\left(\frac{4}{x^2}\right)}{4} \\ &= \frac{2}{x^2} - \left(\frac{1}{x^2}\right) - \frac{1}{x^2} \\ &= 0 \end{aligned}$$

Since the Liouville ode invariant does not depend on the independent variable x then the transformation

$$y = v(x)z(x) \quad (3)$$

is used to change the original ode to a constant coefficients ode in v . In (3) the term $z(x)$ is given by

$$\begin{aligned} z(x) &= e^{-\left(\int \frac{p(x)}{2} dx\right)} \\ &= e^{-\int \frac{-2}{2} dx} \\ &= x \end{aligned} \quad (5)$$

Hence (3) becomes

$$y = v(x)x \quad (4)$$

Applying this change of variable to the original ode results in

$$x^2v''(x) = e^{-x}$$

Which is now solved for $v(x)$ Simplifying the ode gives

$$v''(x) = \frac{e^{-x}}{x^2}$$

Integrating once gives

$$v'(x) = -\frac{e^{-x}}{x} + \expIntegral_1(x) + c_1$$

Integrating again gives

$$v(x) = x \expIntegral_1(x) - e^{-x} + \expIntegral_1(x) + c_1x + c_2$$

Now that $v(x)$ is known, then

$$\begin{aligned} y &= v(x) z(x) \\ &= (x \expIntegral_1(x) + \expIntegral_1(x) - e^{-x} + c_1x + c_2) (z(x)) \end{aligned} \quad (7)$$

But from (5)

$$z(x) = x$$

Hence (7) becomes

$$y = (x \expIntegral_1(x) + \expIntegral_1(x) - e^{-x} + c_1x + c_2) x$$

Therefore the homogeneous solution y_h is

$$y_h = (x \expIntegral_1(x) + \expIntegral_1(x) - e^{-x} + c_1x + c_2) x$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1y_1 + u_2y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = (x^3)^{\frac{1}{3}}$$

$$y_2 = (x^3)^{\frac{2}{3}}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} (x^3)^{\frac{1}{3}} & (x^3)^{\frac{2}{3}} \\ \frac{d}{dx} \left((x^3)^{\frac{1}{3}} \right) & \frac{d}{dx} \left((x^3)^{\frac{2}{3}} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} (x^3)^{\frac{1}{3}} & (x^3)^{\frac{2}{3}} \\ \frac{x^2}{(x^3)^{\frac{2}{3}}} & \frac{2x^2}{(x^3)^{\frac{1}{3}}} \end{vmatrix}$$

Therefore

$$W = \left((x^3)^{\frac{1}{3}} \right) \left(\frac{2x^2}{(x^3)^{\frac{1}{3}}} \right) - \left((x^3)^{\frac{2}{3}} \right) \left(\frac{x^2}{(x^3)^{\frac{2}{3}}} \right)$$

Which simplifies to

$$W = x^2$$

Which simplifies to

$$W = x^2$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{(x^3)^{\frac{2}{3}} x e^{-x}}{x^4} dx$$

Which simplifies to

$$u_1 = - \int \frac{(x^3)^{\frac{2}{3}} e^{-x}}{x^3} dx$$

Hence

$$u_1 = \frac{(x^3)^{\frac{2}{3}} \exp \text{Integral}_1(x)}{x^2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{(x^3)^{\frac{1}{3}} x e^{-x}}{x^4} dx$$

Which simplifies to

$$u_2 = \int \frac{(x^3)^{\frac{1}{3}} e^{-x}}{x^3} dx$$

Hence

$$u_2 = -\frac{(x^3)^{\frac{1}{3}} (-x \exp \text{Integral}_1(x) + e^{-x})}{x^2}$$

Which simplifies to

$$u_1 = \frac{(x^3)^{\frac{2}{3}} \exp \text{Integral}_1(x)}{x^2}$$
$$u_2 = \frac{(x^3)^{\frac{1}{3}} (x \exp \text{Integral}_1(x) - e^{-x})}{x^2}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = x \exp \text{Integral}_1(x) + (x \exp \text{Integral}_1(x) - e^{-x}) x$$

Which simplifies to

$$y_p(x) = x(-e^{-x} + (1+x) \exp \text{Integral}_1(x))$$

Therefore the general solution is

$$y = y_h + y_p$$
$$= ((x \exp \text{Integral}_1(x) + \exp \text{Integral}_1(x) - e^{-x} + c_1 x + c_2) x) + (x(-e^{-x} + (1+x) \exp \text{Integral}_1(x)))$$

Which simplifies to

$$y = (-e^{-x} + (1+x) \exp \text{Integral}_1(x) + c_1x + c_2) x + x(-e^{-x} + (1+x) \exp \text{Integral}_1(x))$$

Summary

The solution(s) found are the following

$$y = (-e^{-x} + (1+x) \exp \text{Integral}_1(x) + c_1x + c_2) x + x(-e^{-x} + (1+x) \exp \text{Integral}_1(x)) \quad (1)$$

Verification of solutions

$$y = (-e^{-x} + (1+x) \exp \text{Integral}_1(x) + c_1x + c_2) x + x(-e^{-x} + (1+x) \exp \text{Integral}_1(x))$$

Verified OK.

11.20.6 Solving as second order change of variable on y method 2 ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = x^2$, $B = -2x$, $C = 2$, $f(x) = x e^{-x}$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$x^2y'' - 2xy' + 2y = 0$$

In normal form the ode

$$x^2y'' - 2xy' + 2y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = -\frac{2}{x}$$
$$q(x) = \frac{2}{x^2}$$

Applying change of variables on the dependent variable $y = v(x)x^n$ to (2) gives the following ode where the dependent variables is $v(x)$ and not y .

$$v''(x) + \left(\frac{2n}{x} + p\right)v'(x) + \left(\frac{n(n-1)}{x^2} + \frac{np}{x} + q\right)v(x) = 0 \quad (3)$$

Let the coefficient of $v(x)$ above be zero. Hence

$$\frac{n(n-1)}{x^2} + \frac{np}{x} + q = 0 \quad (4)$$

Substituting the earlier values found for $p(x)$ and $q(x)$ into (4) gives

$$\frac{n(n-1)}{x^2} - \frac{2n}{x^2} + \frac{2}{x^2} = 0 \quad (5)$$

Solving (5) for n gives

$$n = 2 \quad (6)$$

Substituting this value in (3) gives

$$\begin{aligned} v''(x) + \frac{2v'(x)}{x} &= 0 \\ v''(x) + \frac{2v'(x)}{x} &= 0 \end{aligned} \quad (7)$$

Using the substitution

$$u(x) = v'(x)$$

Then (7) becomes

$$u'(x) + \frac{2u(x)}{x} = 0 \quad (8)$$

The above is now solved for $u(x)$. In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{2u}{x} \end{aligned}$$

Where $f(x) = -\frac{2}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u} du &= -\frac{2}{x} dx \\ \int \frac{1}{u} du &= \int -\frac{2}{x} dx \\ \ln(u) &= -2 \ln(x) + c_1 \\ u &= e^{-2 \ln(x) + c_1} \\ &= \frac{c_1}{x^2}\end{aligned}$$

Now that $u(x)$ is known, then

$$\begin{aligned}v'(x) &= u(x) \\ v(x) &= \int u(x) dx + c_2 \\ &= -\frac{c_1}{x} + c_2\end{aligned}$$

Hence

$$\begin{aligned}y &= v(x) x^n \\ &= \left(-\frac{c_1}{x} + c_2\right) x^2 \\ &= (c_2 x - c_1) x\end{aligned}$$

Now the particular solution to this ODE is found

$$x^2 y'' - 2xy' + 2y = x e^{-x}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned}y_1 &= x \\ y_2 &= x^2\end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} x & x^2 \\ \frac{d}{dx}(x) & \frac{d}{dx}(x^2) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} x & x^2 \\ 1 & 2x \end{vmatrix}$$

Therefore

$$W = (x)(2x) - (x^2)(1)$$

Which simplifies to

$$W = x^2$$

Which simplifies to

$$W = x^2$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^3 e^{-x}}{x^4} dx$$

Which simplifies to

$$u_1 = - \int \frac{e^{-x}}{x} dx$$

Hence

$$u_1 = \exp \text{Integral}_1(x)$$

And Eq. (3) becomes

$$u_2 = \int \frac{x^2 e^{-x}}{x^4} dx$$

Which simplifies to

$$u_2 = \int \frac{e^{-x}}{x^2} dx$$

Hence

$$u_2 = -\frac{e^{-x}}{x} + \text{expIntegral}_1(x)$$

Which simplifies to

$$u_1 = \text{expIntegral}_1(x)$$
$$u_2 = \frac{x \text{expIntegral}_1(x) - e^{-x}}{x}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = x \text{expIntegral}_1(x) + (x \text{expIntegral}_1(x) - e^{-x}) x$$

Which simplifies to

$$y_p(x) = x(-e^{-x} + (1+x) \text{expIntegral}_1(x))$$

Therefore the general solution is

$$y = y_h + y_p$$
$$= \left(\left(-\frac{c_1}{x} + c_2 \right) x^2 \right) + (x(-e^{-x} + (1+x) \text{expIntegral}_1(x)))$$
$$= x(-e^{-x} + (1+x) \text{expIntegral}_1(x)) + \left(-\frac{c_1}{x} + c_2 \right) x^2$$

Which simplifies to

$$y = x(-e^{-x} + (1+x) \text{expIntegral}_1(x)) + c_2 x - c_1$$

Summary

The solution(s) found are the following

$$y = x(-e^{-x} + (1+x) \exp \int_1^x (x) + c_2x - c_1) \quad (1)$$

Verification of solutions

$$y = x(-e^{-x} + (1+x) \exp \int_1^x (x) + c_2x - c_1)$$

Verified OK.

11.20.7 Solving as second order ode non constant coeff transformation on B ode

Given an ode of the form

$$Ay'' + By' + Cy = F(x)$$

This method reduces the order ode the ODE by one by applying the transformation

$$y = Bv$$

This results in

$$\begin{aligned} y' &= B'v + v'B \\ y'' &= B''v + B'v' + v''B + v'B' \\ &= v''B + 2v' + B' + B''v \end{aligned}$$

And now the original ode becomes

$$\begin{aligned} A(v''B + 2v'B' + B''v) + B(B'v + v'B) + CBv &= 0 \\ ABv'' + (2AB' + B^2)v' + (AB'' + BB' + CB)v &= 0 \end{aligned} \quad (1)$$

If the term $AB'' + BB' + CB$ is zero, then this method works and can be used to solve

$$ABv'' + (2AB' + B^2)v' = 0$$

By Using $u = v'$ which reduces the order of the above ode to one. The new ode is

$$ABu' + (2AB' + B^2)u = 0$$

The above ode is first order ode which is solved for u . Now a new ode $v' = u$ is solved for v as first order ode. Then the final solution is obtain from $y = Bv$.

This method works only if the term $AB'' + BB' + CB$ is zero. The given ODE shows that

$$\begin{aligned}A &= x^2 \\B &= -2x \\C &= 2 \\F &= x e^{-x}\end{aligned}$$

The above shows that for this ode

$$\begin{aligned}AB'' + BB' + CB &= (x^2)(0) + (-2x)(-2) + (2)(-2x) \\&= 0\end{aligned}$$

Hence the ode in v given in (1) now simplifies to

$$-2x^3v'' + (0)v' = 0$$

Now by applying $v' = u$ the above becomes

$$-2x^3u'(x) = 0$$

Which is now solved for u . Integrating both sides gives

$$\begin{aligned}u(x) &= \int 0 \, dx \\&= c_1\end{aligned}$$

The ode for v now becomes

$$\begin{aligned}v' &= u \\&= c_1\end{aligned}$$

Which is now solved for v . Integrating both sides gives

$$\begin{aligned}v(x) &= \int c_1 \, dx \\&= c_1x + c_2\end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}y_h(x) &= Bv \\&= (-2x)(c_1x + c_2) \\&= -2x(c_1x + c_2)\end{aligned}$$

And now the particular solution $y_p(x)$ will be found. The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1y_1 + u_2y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = x$$

$$y_2 = x^2$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} x & x^2 \\ \frac{d}{dx}(x) & \frac{d}{dx}(x^2) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} x & x^2 \\ 1 & 2x \end{vmatrix}$$

Therefore

$$W = (x)(2x) - (x^2)(1)$$

Which simplifies to

$$W = x^2$$

Which simplifies to

$$W = x^2$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^3 e^{-x}}{x^4} dx$$

Which simplifies to

$$u_1 = - \int \frac{e^{-x}}{x} dx$$

Hence

$$u_1 = \text{expIntegral}_1(x)$$

And Eq. (3) becomes

$$u_2 = \int \frac{x^2 e^{-x}}{x^4} dx$$

Which simplifies to

$$u_2 = \int \frac{e^{-x}}{x^2} dx$$

Hence

$$u_2 = -\frac{e^{-x}}{x} + \text{expIntegral}_1(x)$$

Which simplifies to

$$u_1 = \text{expIntegral}_1(x)$$
$$u_2 = \frac{x \text{expIntegral}_1(x) - e^{-x}}{x}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = x \text{expIntegral}_1(x) + (x \text{expIntegral}_1(x) - e^{-x}) x$$

Which simplifies to

$$y_p(x) = x(-e^{-x} + (1 + x) \text{expIntegral}_1(x))$$

Hence the complete solution is

$$\begin{aligned}
 y(x) &= y_h + y_p \\
 &= (-2x(c_1x + c_2)) + (x(-e^{-x} + (1+x) \exp\text{Integral}_1(x))) \\
 &= -x(-x \exp\text{Integral}_1(x) + 2c_1x + e^{-x} - \exp\text{Integral}_1(x) + 2c_2)
 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = -x(-x \exp\text{Integral}_1(x) + 2c_1x + e^{-x} - \exp\text{Integral}_1(x) + 2c_2) \quad (1)$$

Verification of solutions

$$y = -x(-x \exp\text{Integral}_1(x) + 2c_1x + e^{-x} - \exp\text{Integral}_1(x) + 2c_2)$$

Verified OK.

11.20.8 Solving using Kovacic algorithm

Writing the ode as

$$x^2y'' - 2xy' + 2y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned}
 A &= x^2 \\
 B &= -2x \\
 C &= 2
 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned}
 r &= \frac{s}{t} \\
 &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}
 \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 0 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 0 \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 304: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-2x}{x^2} dx} \\ &= z_1 e^{\ln(x)} \\ &= z_1(x) \end{aligned}$$

Which simplifies to

$$y_1 = x$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-2x}{x^2} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{2\ln(x)}}{(y_1)^2} dx \\ &= y_1(x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1(x) + c_2(x(x)) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$x^2y'' - 2xy' + 2y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_2x^2 + c_1x$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1y_1 + u_2y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = x$$

$$y_2 = x^2$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} x & x^2 \\ \frac{d}{dx}(x) & \frac{d}{dx}(x^2) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} x & x^2 \\ 1 & 2x \end{vmatrix}$$

Therefore

$$W = (x)(2x) - (x^2) \quad (1)$$

Which simplifies to

$$W = x^2$$

Which simplifies to

$$W = x^2$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^3 e^{-x}}{x^4} dx$$

Which simplifies to

$$u_1 = - \int \frac{e^{-x}}{x} dx$$

Hence

$$u_1 = \text{expIntegral}_1(x)$$

And Eq. (3) becomes

$$u_2 = \int \frac{x^2 e^{-x}}{x^4} dx$$

Which simplifies to

$$u_2 = \int \frac{e^{-x}}{x^2} dx$$

Hence

$$u_2 = -\frac{e^{-x}}{x} + \text{expIntegral}_1(x)$$

Which simplifies to

$$u_1 = \exp\text{Integral}_1(x)$$
$$u_2 = \frac{x \exp\text{Integral}_1(x) - e^{-x}}{x}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = x \exp\text{Integral}_1(x) + (x \exp\text{Integral}_1(x) - e^{-x}) x$$

Which simplifies to

$$y_p(x) = x(-e^{-x} + (1 + x) \exp\text{Integral}_1(x))$$

Therefore the general solution is

$$y = y_h + y_p$$
$$= (c_2x^2 + c_1x) + (x(-e^{-x} + (1 + x) \exp\text{Integral}_1(x)))$$

Which simplifies to

$$y = x(c_2x + c_1) + x(-e^{-x} + (1 + x) \exp\text{Integral}_1(x))$$

Summary

The solution(s) found are the following

$$y = x(c_2x + c_1) + x(-e^{-x} + (1 + x) \exp\text{Integral}_1(x)) \quad (1)$$

Verification of solutions

$$y = x(c_2x + c_1) + x(-e^{-x} + (1 + x) \exp\text{Integral}_1(x))$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    checking if the LODE is of Euler type  
    <- LODE of Euler type successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 24

```
dsolve(x^2*diff(y(x),x$2)-2*x*diff(y(x),x)+2*y(x)=x*exp(-x),y(x), singsol=all)
```

$$y(x) = (-e^{-x} + \text{expIntegral}_1(x)(x+1) + c_2x + c_1)x$$

✓ Solution by Mathematica

Time used: 0.073 (sec). Leaf size: 30

```
DSolve[x^2*y'[x]-2*x*y'[x]+2*y[x]==x*Exp[-x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x(-(x+1)\text{ExpIntegralEi}(-x) - e^{-x} + c_2x + c_1)$$

12 Chapter 2. Second-Order Linear Equations.
Section 2.4. THE USE OF A KNOWN
SOLUTION TO FIND ANOTHER. Page 74

12.1	problem 1(a)	2118
12.2	problem 1(b)	2122
12.3	problem 2	2126
12.4	problem 3	2130
12.5	problem 4	2135
12.6	problem 5	2140
12.7	problem 6(a)	2145
12.8	problem 6(b)	2150
12.9	problem 6(c)	2154
12.10	problem 7	2159
12.11	problem 8	2163

12.1 problem 1(a)

12.1.1 Maple step by step solution 2120

Internal problem ID [6337]

Internal file name [OUTPUT/5585_Sunday_June_05_2022_03_44_13_PM_9921427/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.4. THE USE OF A KNOWN SOLUTION TO FIND ANOTHER. Page 74

Problem number: 1(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "reduction_of_order", "second_order_linear_constant_coeff", "second_order_ode_can_be_made_integrable"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x]]
```

$$y'' + y = 0$$

Given that one solution of the ode is

$$y_1 = \sin(x)$$

Given one basis solution $y_1(x)$, then the second basis solution is given by

$$y_2(x) = y_1 \left(\int \frac{e^{-\int p dx}}{y_1^2} dx \right)$$

Where $p(x)$ is the coefficient of y' when the ode is written in the normal form

$$y'' + p(x)y' + q(x)y = f(x)$$

Looking at the ode to solve shows that

$$p(x) = 0$$

Therefore

$$y_2(x) = \sin(x) \left(\int \frac{e^{-(\int 0 dx)}}{\sin(x)^2} dx \right)$$

$$y_2(x) = \sin(x) \int \frac{1}{\sin(x)^2} dx$$

$$y_2(x) = \sin(x) \left(\int \csc(x)^2 dx \right)$$

$$y_2(x) = -\sin(x) \cot(x)$$

Hence the solution is

$$\begin{aligned} y &= c_1 y_1(x) + c_2 y_2(x) \\ &= \sin(x) c_1 - c_2 \sin(x) \cot(x) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \sin(x) c_1 - c_2 \sin(x) \cot(x) \tag{1}$$




Figure 369: Slope field plot

Verification of solutions

$$y = \sin(x) c_1 - c_2 \sin(x) \cot(x)$$

Verified OK.

12.1.1 Maple step by step solution

Let's solve

$$y'' + y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (-I, I)$$

- 1st solution of the ODE

$$y_1(x) = \cos(x)$$

- 2nd solution of the ODE

$$y_2(x) = \sin(x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = \cos(x) c_1 + c_2 \sin(x)$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve([diff(y(x),x$2)+y(x)=0,sin(x)],singsol=all)
```

$$y(x) = c_1 \sin(x) + \cos(x) c_2$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 16

```
DSolve[y''[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos(x) + c_2 \sin(x)$$

12.2 problem 1(b)

12.2.1 Maple step by step solution 2124

Internal problem ID [6338]

Internal file name [OUTPUT/5586_Sunday_June_05_2022_03_44_14_PM_46831161/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.4. THE USE OF A KNOWN SOLUTION TO FIND ANOTHER. Page 74

Problem number: 1(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "reduction_of_order", "second_order_linear_constant_coeff", "second_order_ode_can_be_made_integrable"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x]]
```

$$y'' - y = 0$$

Given that one solution of the ode is

$$y_1 = e^x$$

Given one basis solution $y_1(x)$, then the second basis solution is given by

$$y_2(x) = y_1 \left(\int \frac{e^{-\int p dx}}{y_1^2} dx \right)$$

Where $p(x)$ is the coefficient of y' when the ode is written in the normal form

$$y'' + p(x)y' + q(x)y = f(x)$$

Looking at the ode to solve shows that

$$p(x) = 0$$

Therefore

$$y_2(x) = e^x \left(\int e^{-(\int 0 dx)} e^{-2x} dx \right)$$

$$y_2(x) = e^x \int \frac{1}{e^{2x}}, dx$$

$$y_2(x) = e^x \left(\int e^{-2x} dx \right)$$

$$y_2(x) = -\frac{e^x e^{-2x}}{2}$$

Hence the solution is

$$\begin{aligned} y &= c_1 y_1(x) + c_2 y_2(x) \\ &= c_1 e^x - \frac{c_2 e^x e^{-2x}}{2} \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^x - \frac{c_2 e^x e^{-2x}}{2} \tag{1}$$




Figure 370: Slope field plot

Verification of solutions

$$y = c_1 e^x - \frac{c_2 e^x e^{-2x}}{2}$$

Verified OK.

12.2.1 Maple step by step solution

Let's solve

$$y'' - y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 - 1 = 0$$

- Factor the characteristic polynomial

$$(r - 1)(r + 1) = 0$$

- Roots of the characteristic polynomial

$$r = (-1, 1)$$

- 1st solution of the ODE

$$y_1(x) = e^{-x}$$

- 2nd solution of the ODE

$$y_2(x) = e^x$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = c_1 e^{-x} + c_2 e^x$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
dsolve([diff(y(x),x$2)-y(x)=0,exp(x)],singsol=all)
```

$$y(x) = c_1 e^{-x} + e^x c_2$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 20

```
DSolve[y''[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^x + c_2 e^{-x}$$

12.3 problem 2

12.3.1 Maple step by step solution 2127

Internal problem ID [6339]

Internal file name [OUTPUT/5587_Sunday_June_05_2022_03_44_15_PM_23721771/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.4. THE USE OF A KNOWN SOLUTION TO FIND ANOTHER. Page 74

Problem number: 2.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "reduction_of_order", "exact linear second order ode", "second_order_integrable_as_is", "second_order_ode_missing_y", "second_order_ode_non_constant_coeff_transformation_on_B"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_y]]
```

$$xy'' + 3y' = 0$$

Given that one solution of the ode is

$$y_1 = 1$$

Given one basis solution $y_1(x)$, then the second basis solution is given by

$$y_2(x) = y_1 \left(\int \frac{e^{-\int p dx}}{y_1^2} dx \right)$$

Where $p(x)$ is the coefficient of y' when the ode is written in the normal form

$$y'' + p(x)y' + q(x)y = f(x)$$

Looking at the ode to solve shows that

$$p(x) = \frac{3}{x}$$

Therefore

$$y_2(x) = \int e^{-(\int \frac{3}{x} dx)} dx$$

$$y_2(x) = 1 \int \frac{1}{x^3}, dx$$

$$y_2(x) = \int \frac{1}{x^3} dx$$

$$y_2(x) = -\frac{1}{2x^2}$$

Hence the solution is

$$\begin{aligned} y &= c_1 y_1(x) + c_2 y_2(x) \\ &= c_1 - \frac{c_2}{2x^2} \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 - \frac{c_2}{2x^2} \tag{1}$$

Verification of solutions

$$y = c_1 - \frac{c_2}{2x^2}$$

Verified OK.

12.3.1 Maple step by step solution

Let's solve

$$y''x + 3y' = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{3y'}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{3y'}{x} = 0$$

- Multiply by denominators of the ODE

$$y''x + 3y' = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x , using the chain rule

$$y' = \left(\frac{d}{dt}y(t)\right) t'(x)$$

- Compute derivative

$$y' = \frac{\frac{d}{dt}y(t)}{x}$$

- Calculate the 2nd derivative of y with respect to x , using the chain rule

$$y'' = \left(\frac{d^2}{dt^2}y(t)\right) t'(x)^2 + t''(x) \left(\frac{d}{dt}y(t)\right)$$

- Compute derivative

$$y'' = \frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2}$$

Substitute the change of variables back into the ODE

$$\left(\frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2}\right) x + \frac{3\left(\frac{d}{dt}y(t)\right)}{x} = 0$$

- Simplify

$$\frac{\frac{d^2}{dt^2}y(t) + 2\frac{d}{dt}y(t)}{x} = 0$$

- Isolate 2nd derivative

$$\frac{d^2}{dt^2}y(t) = -2\frac{d}{dt}y(t)$$

- Group terms with $y(t)$ on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$\frac{d^2}{dt^2}y(t) + 2\frac{d}{dt}y(t) = 0$$

- Characteristic polynomial of ODE

$$r^2 + 2r = 0$$

- Factor the characteristic polynomial

$$r(r + 2) = 0$$

- Roots of the characteristic polynomial

$$r = (-2, 0)$$

- 1st solution of the ODE

$$y_1(t) = e^{-2t}$$

- 2nd solution of the ODE
 $y_2(t) = 1$
- General solution of the ODE
 $y(t) = c_1 y_1(t) + c_2 y_2(t)$
- Substitute in solutions
 $y(t) = c_1 e^{-2t} + c_2$
- Change variables back using $t = \ln(x)$
 $y = \frac{c_1}{x^2} + c_2$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
<- LODE of Euler type successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve([x*diff(y(x),x$2)+3*diff(y(x),x)=0,1],singsol=all)
```

$$y(x) = c_1 + \frac{c_2}{x^2}$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 17

```
DSolve[x*y''[x]+3*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 - \frac{c_1}{2x^2}$$

12.4 problem 3

12.4.1 Maple step by step solution 2132

Internal problem ID [6340]

Internal file name [OUTPUT/5588_Sunday_June_05_2022_03_44_16_PM_95367239/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.4. THE USE OF A KNOWN SOLUTION TO FIND ANOTHER. Page 74

Problem number: 3.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "reduction_of_order", "second_order_euler_ode", "second_order_change_of_variable_on_x_method_1", "second_order_change_of_variable_on_x_method_2", "second_order_change_of_variable_on_y_method_2"

Maple gives the following as the ode type

```
[[_Emden, _Fowler], [_2nd_order, _linear, `_with_symmetry_[0,F(x)]`]]
```

$$x^2y'' + xy' - 4y = 0$$

Given that one solution of the ode is

$$y_1 = x^2$$

Given one basis solution $y_1(x)$, then the second basis solution is given by

$$y_2(x) = y_1 \left(\int \frac{e^{-\int p dx}}{y_1^2} dx \right)$$

Where $p(x)$ is the coefficient of y' when the ode is written in the normal form

$$y'' + p(x)y' + q(x)y = f(x)$$

Looking at the ode to solve shows that

$$p(x) = \frac{1}{x}$$

Therefore

$$y_2(x) = x^2 \left(\int \frac{e^{-\left(\int \frac{1}{x} dx\right)}}{x^4} dx \right)$$

$$y_2(x) = x^2 \int \frac{\frac{1}{x}}{x^4} dx$$

$$y_2(x) = x^2 \left(\int \frac{1}{x^5} dx \right)$$

$$y_2(x) = -\frac{1}{4x^2}$$

Hence the solution is

$$\begin{aligned} y &= c_1 y_1(x) + c_2 y_2(x) \\ &= c_1 x^2 - \frac{c_2}{4x^2} \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 x^2 - \frac{c_2}{4x^2} \tag{1}$$

Verification of solutions

$$y = c_1 x^2 - \frac{c_2}{4x^2}$$

Verified OK.

12.4.1 Maple step by step solution

Let's solve

$$x^2 y'' + xy' - 4y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{y'}{x} + \frac{4y}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{y'}{x} - \frac{4y}{x^2} = 0$$

- Multiply by denominators of the ODE

$$x^2 y'' + xy' - 4y = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x , using the chain rule

$$y' = \left(\frac{d}{dt}y(t)\right) t'(x)$$

- Compute derivative

$$y' = \frac{\frac{d}{dt}y(t)}{x}$$

- Calculate the 2nd derivative of y with respect to x , using the chain rule

$$y'' = \left(\frac{d^2}{dt^2}y(t)\right) t'(x)^2 + t''(x) \left(\frac{d}{dt}y(t)\right)$$

- Compute derivative

$$y'' = \frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2}$$

Substitute the change of variables back into the ODE

$$x^2 \left(\frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2}\right) + \frac{d}{dt}y(t) - 4y(t) = 0$$

- Simplify

$$\frac{d^2}{dt^2}y(t) - 4y(t) = 0$$

- Characteristic polynomial of ODE

$$r^2 - 4 = 0$$

- Factor the characteristic polynomial
 $(r - 2)(r + 2) = 0$
- Roots of the characteristic polynomial
 $r = (-2, 2)$
- 1st solution of the ODE
 $y_1(t) = e^{-2t}$
- 2nd solution of the ODE
 $y_2(t) = e^{2t}$
- General solution of the ODE
 $y(t) = c_1 y_1(t) + c_2 y_2(t)$
- Substitute in solutions
 $y(t) = c_1 e^{-2t} + c_2 e^{2t}$
- Change variables back using $t = \ln(x)$
 $y = \frac{c_1}{x^2} + c_2 x^2$
- Simplify
 $y = \frac{c_1}{x^2} + c_2 x^2$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
<- LODE of Euler type successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
dsolve([x^2*diff(y(x),x$2)+x*diff(y(x),x)-4*y(x)=0,x^2],singsol=all)
```

$$y(x) = \frac{c_1 x^4 + c_2}{x^2}$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 18

```
DSolve[x^2*y''[x]+x*y'[x]-4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_2 x^4 + c_1}{x^2}$$

12.5 problem 4

12.5.1 Maple step by step solution 2136

Internal problem ID [6341]

Internal file name [OUTPUT/5589_Sunday_June_05_2022_03_44_17_PM_96299379/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.4. THE USE OF A KNOWN SOLUTION TO FIND ANOTHER. Page 74

Problem number: 4.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**reduction_of_order**", "**second_order_change_of_variable_on_y_method_2**", "**second_order_ode_non_constant_coeff_transformation_on_B**"

Maple gives the following as the ode type

[_Gegenbauer]

$$(-x^2 + 1)y'' - 2xy' + 2y = 0$$

Given that one solution of the ode is

$$y_1 = x$$

Given one basis solution $y_1(x)$, then the second basis solution is given by

$$y_2(x) = y_1 \left(\int \frac{e^{-\int p dx}}{y_1^2} dx \right)$$

Where $p(x)$ is the coefficient of y' when the ode is written in the normal form

$$y'' + p(x)y' + q(x)y = f(x)$$

Looking at the ode to solve shows that

$$p(x) = -\frac{2x}{-x^2 + 1}$$

Therefore

$$y_2(x) = x \left(\int \frac{e^{-\left(\int -\frac{2x}{x^2+1} dx\right)}}{x^2} dx \right)$$

$$y_2(x) = x \int \frac{e^{-\ln(x-1)-\ln(1+x)}}{x^2} dx$$

$$y_2(x) = x \left(\int \frac{1}{x^2(x^2-1)} dx \right)$$

$$y_2(x) = x \left(-\frac{\ln(1+x)}{2} + \frac{\ln(x-1)}{2} + \frac{1}{x} \right)$$

Hence the solution is

$$\begin{aligned} y &= c_1 y_1(x) + c_2 y_2(x) \\ &= c_1 x + c_2 x \left(-\frac{\ln(1+x)}{2} + \frac{\ln(x-1)}{2} + \frac{1}{x} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 x + c_2 x \left(-\frac{\ln(1+x)}{2} + \frac{\ln(x-1)}{2} + \frac{1}{x} \right) \quad (1)$$

Verification of solutions

$$y = c_1 x + c_2 x \left(-\frac{\ln(1+x)}{2} + \frac{\ln(x-1)}{2} + \frac{1}{x} \right)$$

Verified OK.

12.5.1 Maple step by step solution

Let's solve

$$(-x^2 + 1)y'' - 2xy' + 2y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{2xy'}{x^2-1} + \frac{2y}{x^2-1}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{2xy'}{x^2-1} - \frac{2y}{x^2-1} = 0$$

- Check to see if x_0 is a regular singular point

- Define functions

$$[P_2(x) = \frac{2x}{x^2-1}, P_3(x) = -\frac{2}{x^2-1}]$$

- $(1+x) \cdot P_2(x)$ is analytic at $x = -1$

$$((1+x) \cdot P_2(x)) \Big|_{x=-1} = 1$$

- $(1+x)^2 \cdot P_3(x)$ is analytic at $x = -1$

$$((1+x)^2 \cdot P_3(x)) \Big|_{x=-1} = 0$$

- $x = -1$ is a regular singular point

Check to see if x_0 is a regular singular point

$$x_0 = -1$$

- Multiply by denominators

$$(x^2 - 1)y'' + 2xy' - 2y = 0$$

- Change variables using $x = u - 1$ so that the regular singular point is at $u = 0$

$$(u^2 - 2u) \left(\frac{d^2}{du^2} y(u) \right) + (2u - 2) \left(\frac{d}{du} y(u) \right) - 2y(u) = 0$$

- Assume series solution for $y(u)$

$$y(u) = \sum_{k=0}^{\infty} a_k u^{k+r}$$

- Rewrite ODE with series expansions

- Convert $u^m \cdot \left(\frac{d}{du} y(u) \right)$ to series expansion for $m = 0..1$

$$u^m \cdot \left(\frac{d}{du} y(u) \right) = \sum_{k=0}^{\infty} a_k (k+r) u^{k+r-1+m}$$

- Shift index using $k \rightarrow k + 1 - m$

$$u^m \cdot \left(\frac{d}{du} y(u) \right) = \sum_{k=-1+m}^{\infty} a_{k+1-m} (k+1-m+r) u^{k+r}$$

- Convert $u^m \cdot \left(\frac{d^2}{du^2} y(u) \right)$ to series expansion for $m = 1..2$

$$u^m \cdot \left(\frac{d^2}{du^2} y(u) \right) = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) u^{k+r-2+m}$$

- Shift index using $k \rightarrow k + 2 - m$

$$u^m \cdot \left(\frac{d^2}{du^2} y(u) \right) = \sum_{k=-2+m}^{\infty} a_{k+2-m} (k+2-m+r) (k+1-m+r) u^{k+r}$$

Rewrite ODE with series expansions

$$-2a_0 r^2 u^{-1+r} + \left(\sum_{k=0}^{\infty} (-2a_{k+1} (k+1+r)^2 + a_k (k+r+2) (k+r-1)) u^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$-2r^2 = 0$$

- Values of r that satisfy the indicial equation

$$r = 0$$

- Each term in the series must be 0, giving the recursion relation

$$-2a_{k+1} (k+1)^2 + a_k (k+2) (k-1) = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+1} = \frac{a_k (k+2)(k-1)}{2(k+1)^2}$$

- Recursion relation for $r = 0$; series terminates at $k = 1$

$$a_{k+1} = \frac{a_k (k+2)(k-1)}{2(k+1)^2}$$

- Apply recursion relation for $k = 0$

$$a_1 = -a_0$$

- Terminating series solution of the ODE for $r = 0$. Use reduction of order to find the second li

$$y(u) = a_0 \cdot (-u + 1)$$

- Revert the change of variables $u = 1 + x$

$$[y = -a_0 x]$$

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacics algorithm  
  A Liouvillian solution exists  
  Reducible group (found an exponential solution)  
  Group is reducible, not completely reducible  
<- Kovacics algorithm successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve([(1-x^2)*diff(y(x),x$2)-2*x*diff(y(x),x)+2*y(x)=0,x],singsol=all)
```

$$y(x) = -\frac{c_2 \ln(x+1)x}{2} + \frac{c_2 \ln(x-1)x}{2} + c_1x + c_2$$

✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 33

```
DSolve[(1-x^2)*y'[x]-2*x*y'[x]+2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1x - \frac{1}{2}c_2(x \log(1-x) - x \log(x+1) + 2)$$

12.6 problem 5

12.6.1 Maple step by step solution 2141

Internal problem ID [6342]

Internal file name [OUTPUT/5590_Sunday_June_05_2022_03_44_19_PM_36590290/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.4. THE USE OF A KNOWN SOLUTION TO FIND ANOTHER. Page 74

Problem number: 5.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**reduction_of_order**", "**second_order_bessel_ode**", "**second_order_change_of_variable_on_y_method_1**"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$x^2 y'' + xy' + \left(x^2 - \frac{1}{4}\right) y = 0$$

Given that one solution of the ode is

$$y_1 = \frac{\sin(x)}{\sqrt{x}}$$

Given one basis solution $y_1(x)$, then the second basis solution is given by

$$y_2(x) = y_1 \left(\int \frac{e^{-\int p dx}}{y_1^2} dx \right)$$

Where $p(x)$ is the coefficient of y' when the ode is written in the normal form

$$y'' + p(x)y' + q(x)y = f(x)$$

Looking at the ode to solve shows that

$$p(x) = \frac{1}{x}$$

Therefore

$$y_2(x) = \frac{\sin(x) \left(\int \frac{e^{-\left(\int \frac{1}{x} dx\right)x}}{\sin(x)^2} dx \right)}{\sqrt{x}}$$

$$y_2(x) = \frac{\sin(x)}{\sqrt{x}} \int \frac{\frac{1}{x}}{\sin(x)^2} dx$$

$$y_2(x) = \frac{\sin(x) \left(\int \csc(x)^2 dx \right)}{\sqrt{x}}$$

$$y_2(x) = -\frac{\sin(x) \cot(x)}{\sqrt{x}}$$

Hence the solution is

$$\begin{aligned} y &= c_1 y_1(x) + c_2 y_2(x) \\ &= \frac{\sin(x) c_1}{\sqrt{x}} - \frac{c_2 \sin(x) \cot(x)}{\sqrt{x}} \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{\sin(x) c_1}{\sqrt{x}} - \frac{c_2 \sin(x) \cot(x)}{\sqrt{x}} \tag{1}$$

Verification of solutions

$$y = \frac{\sin(x) c_1}{\sqrt{x}} - \frac{c_2 \sin(x) \cot(x)}{\sqrt{x}}$$

Verified OK.

12.6.1 Maple step by step solution

Let's solve

$$x^2 y'' + xy' + \left(x^2 - \frac{1}{4}\right) y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{(4x^2-1)y}{4x^2} - \frac{y'}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{y'}{x} + \frac{(4x^2-1)y}{4x^2} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$\left[P_2(x) = \frac{1}{x}, P_3(x) = \frac{4x^2-1}{4x^2} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 1$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = -\frac{1}{4}$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$4x^2y'' + 4xy' + (4x^2 - 1)y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 0..2$

$$x^m \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

- Shift index using $k \rightarrow k - m$

$$x^m \cdot y = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

- Convert $x \cdot y'$ to series expansion

$$x \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r}$$

- Convert $x^2 \cdot y''$ to series expansion

$$x^2 \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(1+2r)(-1+2r)x^r + a_1(3+2r)(1+2r)x^{1+r} + \left(\sum_{k=2}^{\infty} (a_k(2k+2r+1)(2k+2r-1) + 4a_{k-2}) \right) x^{k+r}$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$(1+2r)(-1+2r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \left\{ -\frac{1}{2}, \frac{1}{2} \right\}$$

- Each term must be 0

$$a_1(3+2r)(1+2r) = 0$$

- Solve for the dependent coefficient(s)

$$a_1 = 0$$

- Each term in the series must be 0, giving the recursion relation

$$a_k(4k^2 + 8kr + 4r^2 - 1) + 4a_{k-2} = 0$$

- Shift index using $k \rightarrow k+2$

$$a_{k+2}(4(k+2)^2 + 8(k+2)r + 4r^2 - 1) + 4a_k = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+2} = -\frac{4a_k}{4k^2 + 8kr + 4r^2 + 16k + 16r + 15}$$

- Recursion relation for $r = -\frac{1}{2}$

$$a_{k+2} = -\frac{4a_k}{4k^2 + 12k + 8}$$

- Solution for $r = -\frac{1}{2}$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k-\frac{1}{2}}, a_{k+2} = -\frac{4a_k}{4k^2 + 12k + 8}, a_1 = 0 \right]$$

- Recursion relation for $r = \frac{1}{2}$

$$a_{k+2} = -\frac{4a_k}{4k^2 + 20k + 24}$$

- Solution for $r = \frac{1}{2}$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+\frac{1}{2}}, a_{k+2} = -\frac{4a_k}{4k^2 + 20k + 24}, a_1 = 0 \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^{k-\frac{1}{2}} \right) + \left(\sum_{k=0}^{\infty} b_k x^{k+\frac{1}{2}} \right), a_{k+2} = -\frac{4a_k}{4k^2 + 12k + 8}, a_1 = 0, b_{k+2} = -\frac{4b_k}{4k^2 + 20k + 24}, b_1 = 0 \right]$$

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacics algorithm  
  A Liouvillian solution exists  
  Group is reducible or imprimitive  
<- Kovacics algorithm successful`
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 17

```
dsolve([x^2*diff(y(x),x$2)+x*diff(y(x),x)+(x^2-1/4)*y(x)=0,x^(-1/2)*sin(x)],singsol=all)
```

$$y(x) = \frac{c_1 \sin(x) + \cos(x) c_2}{\sqrt{x}}$$

✓ Solution by Mathematica

Time used: 0.044 (sec). Leaf size: 39

```
DSolve[x^2*y''[x]+x*y'[x]+(x^2-1/4)*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{-ix}(2c_1 - ic_2 e^{2ix})}{2\sqrt{x}}$$

12.7 problem 6(a)

12.7.1 Maple step by step solution 2146

Internal problem ID [6343]

Internal file name [OUTPUT/5591_Sunday_June_05_2022_03_44_20_PM_15281440/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.4. THE USE OF A KNOWN SOLUTION TO FIND ANOTHER. Page 74

Problem number: 6(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**reduction_of_order**", "**second_order_change_of_variable_on_y_method_2**", "**second_order_ode_non_constant_coeff_transformation_on_B**"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - \frac{xy'}{x-1} + \frac{y}{x-1} = 0$$

Given that one solution of the ode is

$$y_1 = x$$

Given one basis solution $y_1(x)$, then the second basis solution is given by

$$y_2(x) = y_1 \left(\int \frac{e^{-\int p dx}}{y_1^2} dx \right)$$

Where $p(x)$ is the coefficient of y' when the ode is written in the normal form

$$y'' + p(x)y' + q(x)y = f(x)$$

Looking at the ode to solve shows that

$$p(x) = -\frac{x}{x-1}$$

Therefore

$$y_2(x) = x \left(\int \frac{e^{-\left(\int -\frac{x}{x-1} dx\right)}}{x^2} dx \right)$$

$$y_2(x) = x \int \frac{e^{x+\ln(x-1)}}{x^2} dx$$

$$y_2(x) = x \left(\int \frac{e^x(x-1)}{x^2} dx \right)$$

$$y_2(x) = e^x$$

Hence the solution is

$$\begin{aligned} y &= c_1 y_1(x) + c_2 y_2(x) \\ &= c_1 x + c_2 e^x \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 x + c_2 e^x \tag{1}$$

Verification of solutions

$$y = c_1 x + c_2 e^x$$

Verified OK.

12.7.1 Maple step by step solution

Let's solve

$$y'' - \frac{xy'}{x-1} + \frac{y}{x-1} = 0$$

- Highest derivative means the order of the ODE is 2

y''

- Check to see if $x_0 = 1$ is a regular singular point

- Define functions

$$\left[P_2(x) = -\frac{x}{x-1}, P_3(x) = \frac{1}{x-1} \right]$$

- $(x-1) \cdot P_2(x)$ is analytic at $x = 1$

$$\left. ((x-1) \cdot P_2(x)) \right|_{x=1} = -1$$

- $(x - 1)^2 \cdot P_3(x)$ is analytic at $x = 1$

$$\left. ((x - 1)^2 \cdot P_3(x)) \right|_{x=1} = 0$$

- $x = 1$ is a regular singular point

Check to see if $x_0 = 1$ is a regular singular point

$$x_0 = 1$$

- Multiply by denominators

$$(x - 1)y'' - xy' + y = 0$$

- Change variables using $x = u + 1$ so that the regular singular point is at $u = 0$

$$u \left(\frac{d^2}{du^2} y(u) \right) + (-u - 1) \left(\frac{d}{du} y(u) \right) + y(u) = 0$$

- Assume series solution for $y(u)$

$$y(u) = \sum_{k=0}^{\infty} a_k u^{k+r}$$

- Rewrite ODE with series expansions

- Convert $u^m \cdot \left(\frac{d}{du} y(u) \right)$ to series expansion for $m = 0..1$

$$u^m \cdot \left(\frac{d}{du} y(u) \right) = \sum_{k=0}^{\infty} a_k (k + r) u^{k+r-1+m}$$

- Shift index using $k \rightarrow k + 1 - m$

$$u^m \cdot \left(\frac{d}{du} y(u) \right) = \sum_{k=-1+m}^{\infty} a_{k+1-m} (k + 1 - m + r) u^{k+r}$$

- Convert $u \cdot \left(\frac{d^2}{du^2} y(u) \right)$ to series expansion

$$u \cdot \left(\frac{d^2}{du^2} y(u) \right) = \sum_{k=0}^{\infty} a_k (k + r) (k + r - 1) u^{k+r-1}$$

- Shift index using $k \rightarrow k + 1$

$$u \cdot \left(\frac{d^2}{du^2} y(u) \right) = \sum_{k=-1}^{\infty} a_{k+1} (k + 1 + r) (k + r) u^{k+r}$$

Rewrite ODE with series expansions

$$a_0 r(-2 + r) u^{-1+r} + \left(\sum_{k=0}^{\infty} (a_{k+1} (k + 1 + r) (k + r - 1) - a_k (k + r - 1)) u^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$r(-2 + r) = 0$$

- Values of r that satisfy the indicial equation
 $r \in \{0, 2\}$
- Each term in the series must be 0, giving the recursion relation
 $(k + r - 1)(a_{k+1}(k + 1 + r) - a_k) = 0$
- Recursion relation that defines series solution to ODE
 $a_{k+1} = \frac{a_k}{k+1+r}$
- Recursion relation for $r = 0$
 $a_{k+1} = \frac{a_k}{k+1}$
- Solution for $r = 0$
 $\left[y(u) = \sum_{k=0}^{\infty} a_k u^k, a_{k+1} = \frac{a_k}{k+1} \right]$
- Revert the change of variables $u = x - 1$
 $\left[y = \sum_{k=0}^{\infty} a_k (x - 1)^k, a_{k+1} = \frac{a_k}{k+1} \right]$
- Recursion relation for $r = 2$
 $a_{k+1} = \frac{a_k}{k+3}$
- Solution for $r = 2$
 $\left[y(u) = \sum_{k=0}^{\infty} a_k u^{k+2}, a_{k+1} = \frac{a_k}{k+3} \right]$
- Revert the change of variables $u = x - 1$
 $\left[y = \sum_{k=0}^{\infty} a_k (x - 1)^{k+2}, a_{k+1} = \frac{a_k}{k+3} \right]$
- Combine solutions and rename parameters
 $\left[y = \left(\sum_{k=0}^{\infty} a_k (x - 1)^k \right) + \left(\sum_{k=0}^{\infty} b_k (x - 1)^{k+2} \right), a_{k+1} = \frac{a_k}{k+1}, b_{k+1} = \frac{b_k}{k+3} \right]$

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacics algorithm  
  A Liouvillian solution exists  
  Reducible group (found an exponential solution)  
  Reducible group (found another exponential solution)  
<- Kovacics algorithm successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 12

```
dsolve([diff(y(x),x$2)-x/(x-1)*diff(y(x),x)+1/(x-1)*y(x)=0,x],singsol=all)
```

$$y(x) = c_1x + e^x c_2$$

✓ Solution by Mathematica

Time used: 0.061 (sec). Leaf size: 17

```
DSolve[y''[x]-x/(x-1)*y'[x]+1/(x-1)*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1e^x - c_2x$$

12.8 problem 6(b)

12.8.1 Maple step by step solution 2151

Internal problem ID [6344]

Internal file name [OUTPUT/5592_Sunday_June_05_2022_03_44_21_PM_71927023/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.4. THE USE OF A KNOWN SOLUTION TO FIND ANOTHER. Page 74

Problem number: 6(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**reduction_of_order**", "**second_order_euler_ode**", "**second_order_change_of_variable_on_x_method_2**", "**second_order_change_of_variable_on_y_method_2**", "**second_order_ode_non_constant_coeff_transformation_on_B**"

Maple gives the following as the ode type

[[_Emden , _Fowler]]

$$x^2y'' + 2xy' - 2y = 0$$

Given that one solution of the ode is

$$y_1 = x$$

Given one basis solution $y_1(x)$, then the second basis solution is given by

$$y_2(x) = y_1 \left(\int \frac{e^{-\int p dx}}{y_1^2} dx \right)$$

Where $p(x)$ is the coefficient of y' when the ode is written in the normal form

$$y'' + p(x)y' + q(x)y = f(x)$$

Looking at the ode to solve shows that

$$p(x) = \frac{2}{x}$$

Therefore

$$y_2(x) = x \left(\int \frac{e^{-\left(\int \frac{2}{x} dx\right)}}{x^2} dx \right)$$

$$y_2(x) = x \int \frac{1}{x^2} dx$$

$$y_2(x) = x \left(\int \frac{1}{x^2} dx \right)$$

$$y_2(x) = -\frac{1}{3x^2}$$

Hence the solution is

$$\begin{aligned} y &= c_1 y_1(x) + c_2 y_2(x) \\ &= c_1 x - \frac{c_2}{3x^2} \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 x - \frac{c_2}{3x^2} \tag{1}$$

Verification of solutions

$$y = c_1 x - \frac{c_2}{3x^2}$$

Verified OK.

12.8.1 Maple step by step solution

Let's solve

$$x^2 y'' + 2xy' - 2y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{2y'}{x} + \frac{2y}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{2y'}{x} - \frac{2y}{x^2} = 0$$

- Multiply by denominators of the ODE

$$x^2 y'' + 2xy' - 2y = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x , using the chain rule

$$y' = \left(\frac{d}{dt} y(t) \right) t'(x)$$

- Compute derivative

$$y' = \frac{\frac{d}{dt} y(t)}{x}$$

- Calculate the 2nd derivative of y with respect to x , using the chain rule

$$y'' = \left(\frac{d^2}{dt^2} y(t) \right) t'(x)^2 + t''(x) \left(\frac{d}{dt} y(t) \right)$$

- Compute derivative

$$y'' = \frac{\frac{d^2}{dt^2} y(t)}{x^2} - \frac{\frac{d}{dt} y(t)}{x^2}$$

Substitute the change of variables back into the ODE

$$x^2 \left(\frac{\frac{d^2}{dt^2} y(t)}{x^2} - \frac{\frac{d}{dt} y(t)}{x^2} \right) + 2 \frac{d}{dt} y(t) - 2y(t) = 0$$

- Simplify

$$\frac{d^2}{dt^2} y(t) + \frac{d}{dt} y(t) - 2y(t) = 0$$

- Characteristic polynomial of ODE

$$r^2 + r - 2 = 0$$

- Factor the characteristic polynomial

$$(r + 2)(r - 1) = 0$$

- Roots of the characteristic polynomial

$$r = (-2, 1)$$

- 1st solution of the ODE

$$y_1(t) = e^{-2t}$$

- 2nd solution of the ODE

$$y_2(t) = e^t$$

- General solution of the ODE

$$y(t) = c_1 y_1(t) + c_2 y_2(t)$$

- Substitute in solutions

$$y(t) = c_1 e^{-2t} + c_2 e^t$$

- Change variables back using $t = \ln(x)$

$$y = \frac{c_1}{x^2} + c_2 x$$

- Simplify

$$y = \frac{c_1}{x^2} + c_2 x$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
<- LODE of Euler type successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve([x^2*diff(y(x),x$2)+2*x*diff(y(x),x)-2*y(x)=0,x],singsol=all)
```

$$y(x) = \frac{c_2 x^3 + c_1}{x^2}$$

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 16

```
DSolve[x^2*y''[x]+2*x*y'[x]-2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_1}{x^2} + c_2 x$$

12.9 problem 6(c)

12.9.1 Maple step by step solution 2155

Internal problem ID [6345]

Internal file name [OUTPUT/5593_Sunday_June_05_2022_03_44_22_PM_15722899/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.4. THE USE OF A KNOWN SOLUTION TO FIND ANOTHER. Page 74

Problem number: 6(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**reduction_of_order**", "**second_order_change_of_variable_on_y_method_1**", "**second_order_change_of_variable_on_y_method_2**"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$x^2 y'' - x(x+2)y' + (x+2)y = 0$$

Given that one solution of the ode is

$$y_1 = x$$

Given one basis solution $y_1(x)$, then the second basis solution is given by

$$y_2(x) = y_1 \left(\int \frac{e^{-\int p dx}}{y_1^2} dx \right)$$

Where $p(x)$ is the coefficient of y' when the ode is written in the normal form

$$y'' + p(x)y' + q(x)y = f(x)$$

Looking at the ode to solve shows that

$$p(x) = \frac{-x^2 - 2x}{x^2}$$

Therefore

$$y_2(x) = x \left(\int \frac{e^{-\left(\int \frac{-x^2-2x}{x^2} dx\right)}}{x^2} dx \right)$$

$$y_2(x) = x \int \frac{e^{x+2\ln(x)}}{x^2} dx$$

$$y_2(x) = \left(\int e^x dx \right) x$$

$$y_2(x) = x e^x$$

Hence the solution is

$$\begin{aligned} y &= c_1 y_1(x) + c_2 y_2(x) \\ &= c_1 x + c_2 x e^x \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 x + c_2 x e^x \tag{1}$$

Verification of solutions

$$y = c_1 x + c_2 x e^x$$

Verified OK.

12.9.1 Maple step by step solution

Let's solve

$$x^2 y'' + (-x^2 - 2x) y' + (x + 2) y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{(x+2)y}{x^2} + \frac{(x+2)y'}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - \frac{(x+2)y'}{x} + \frac{(x+2)y}{x^2} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$[P_2(x) = -\frac{x+2}{x}, P_3(x) = \frac{x+2}{x^2}]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = -2$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 2$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$x^2 y'' - x(x+2)y' + (x+2)y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 0..1$

$$x^m \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

- Shift index using $k \rightarrow k - m$

$$x^m \cdot y = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

- Convert $x^m \cdot y'$ to series expansion for $m = 1..2$

$$x^m \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1+m}$$

- Shift index using $k \rightarrow k + 1 - m$

$$x^m \cdot y' = \sum_{k=-1+m}^{\infty} a_{k+1-m} (k+1-m+r) x^{k+r}$$

- Convert $x^2 \cdot y''$ to series expansion

$$x^2 \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(-1+r)(-2+r)x^r + \left(\sum_{k=1}^{\infty} (a_k(k+r-1)(k+r-2) - a_{k-1}(k+r-2))x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation
 $(-1+r)(-2+r) = 0$
- Values of r that satisfy the indicial equation
 $r \in \{1, 2\}$
- Each term in the series must be 0, giving the recursion relation
 $(k+r-2)(a_k(k+r-1) - a_{k-1}) = 0$
- Shift index using $k \rightarrow k+1$
 $(k+r-1)(a_{k+1}(k+r) - a_k) = 0$
- Recursion relation that defines series solution to ODE
 $a_{k+1} = \frac{a_k}{k+r}$
- Recursion relation for $r = 1$
 $a_{k+1} = \frac{a_k}{k+1}$
- Solution for $r = 1$
 $\left[y = \sum_{k=0}^{\infty} a_k x^{k+1}, a_{k+1} = \frac{a_k}{k+1} \right]$
- Recursion relation for $r = 2$
 $a_{k+1} = \frac{a_k}{k+2}$
- Solution for $r = 2$
 $\left[y = \sum_{k=0}^{\infty} a_k x^{k+2}, a_{k+1} = \frac{a_k}{k+2} \right]$
- Combine solutions and rename parameters
 $\left[y = \left(\sum_{k=0}^{\infty} a_k x^{k+1} \right) + \left(\sum_{k=0}^{\infty} b_k x^{k+2} \right), a_{k+1} = \frac{a_k}{k+1}, b_{k+1} = \frac{b_k}{k+2} \right]$

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacics algorithm  
  A Liouvillian solution exists  
  Reducible group (found an exponential solution)  
  Reducible group (found another exponential solution)  
<- Kovacics algorithm successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 12

```
dsolve([x^2*diff(y(x),x$2)-x*(x+2)*diff(y(x),x)+(x+2)*y(x)=0,x],singsol=all)
```

$$y(x) = x(e^x c_2 + c_1)$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 16

```
DSolve[x^2*y'[x]-x*(x+2)*y'[x]+(x+2)*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x(c_2 e^x + c_1)$$

12.10 problem 7

12.10.1 Solving as second order change of variable on y method 2 ode . 2159

Internal problem ID [6346]

Internal file name [OUTPUT/5594_Sunday_June_05_2022_03_44_24_PM_57895568/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.4. THE USE OF A KNOWN SOLUTION TO FIND ANOTHER. Page 74

Problem number: 7.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_change_of_variable_on_y_method_2**"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - xf(x)y' + f(x)y = 0$$

12.10.1 Solving as second order change of variable on y method 2 ode

In normal form the ode

$$y'' - xf(x)y' + f(x)y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = -f(x)x$$

$$q(x) = f(x)$$

Applying change of variables on the dependent variable $y = v(x)x^n$ to (2) gives the following ode where the dependent variables is $v(x)$ and not y .

$$v''(x) + \left(\frac{2n}{x} + p\right)v'(x) + \left(\frac{n(n-1)}{x^2} + \frac{np}{x} + q\right)v(x) = 0 \quad (3)$$

Let the coefficient of $v(x)$ above be zero. Hence

$$\frac{n(n-1)}{x^2} + \frac{np}{x} + q = 0 \quad (4)$$

Substituting the earlier values found for $p(x)$ and $q(x)$ into (4) gives

$$\frac{n(n-1)}{x^2} - nf(x) + f(x) = 0 \quad (5)$$

Solving (5) for n gives

$$n = 1 \quad (6)$$

Substituting this value in (3) gives

$$\begin{aligned} v''(x) + \left(\frac{2}{x} - f(x)x\right)v'(x) &= 0 \\ v''(x) + \left(\frac{2}{x} - f(x)x\right)v'(x) &= 0 \end{aligned} \quad (7)$$

Using the substitution

$$u(x) = v'(x)$$

Then (7) becomes

$$u'(x) + \left(\frac{2}{x} - f(x)x\right)u(x) = 0 \quad (8)$$

The above is now solved for $u(x)$. In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= \frac{u(f(x)x^2 - 2)}{x} \end{aligned}$$

Where $f(x) = \frac{f(x)x^2 - 2}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned} \frac{1}{u} du &= \frac{f(x)x^2 - 2}{x} dx \\ \int \frac{1}{u} du &= \int \frac{f(x)x^2 - 2}{x} dx \\ \ln(u) &= \int \frac{f(x)x^2 - 2}{x} dx + c_1 \\ u &= e^{\int \frac{f(x)x^2 - 2}{x} dx + c_1} \\ &= c_1 e^{\int \frac{f(x)x^2 - 2}{x} dx} \end{aligned}$$

Now that $u(x)$ is known, then

$$\begin{aligned}v'(x) &= u(x) \\v(x) &= \int u(x) dx + c_2 \\&= \int c_1 e^{\int \frac{f(x)x^2-2}{x} dx} dx + c_2\end{aligned}$$

Hence

$$\begin{aligned}y &= v(x) x^n \\&= \left(\int c_1 e^{\int \frac{f(x)x^2-2}{x} dx} dx + c_2 \right) x \\&= \left(c_1 \left(\int e^{\int \frac{f(x)x^2-2}{x} dx} dx \right) + c_2 \right) x\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \left(\int c_1 e^{\int \frac{f(x)x^2-2}{x} dx} dx + c_2 \right) x \quad (1)$$

Verification of solutions

$$y = \left(\int c_1 e^{\int \frac{f(x)x^2-2}{x} dx} dx + c_2 \right) x$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) * 2F1([a
-> Trying changes of variables to rationalize or make the ODE simpler
<- unable to find a useful change of variables
    trying a symmetry of the form [xi=0, eta=F(x)]
    trying 2nd order exact linear
    trying symmetries linear in x and y(x)
<- linear symmetries successful`
```

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 27

```
dsolve(diff(y(x),x$2)-x*f(x)*diff(y(x),x)+f(x)*y(x)=0,y(x), singsol=all)
```

$$y(x) = x \left(\left(\int e^{\int \frac{-2+f(x)x^2}{x} dx} dx \right) c_1 + c_2 \right)$$

✓ Solution by Mathematica

Time used: 0.287 (sec). Leaf size: 44

```
DSolve[y''[x]-x*f[x]*y'[x]+f[x]*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x \left(c_2 \int_1^x \frac{\exp \left(- \int_1^{K[2]} -f(K[1])K[1]dK[1] \right)}{K[2]^2} dK[2] + c_1 \right)$$

12.11 problem 8

12.11.1 Maple step by step solution 2164

Internal problem ID [6347]

Internal file name [OUTPUT/5595_Sunday_June_05_2022_03_44_26_PM_43339838/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Second-Order Linear Equations. Section 2.4. THE USE OF A KNOWN SOLUTION TO FIND ANOTHER. Page 74

Problem number: 8.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**reduction_of_order**"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$xy'' - (1 + 2x)y' + (1 + x)y = 0$$

Given that one solution of the ode is

$$y_1 = e^x$$

Given one basis solution $y_1(x)$, then the second basis solution is given by

$$y_2(x) = y_1 \left(\int \frac{e^{-\int p dx}}{y_1^2} dx \right)$$

Where $p(x)$ is the coefficient of y' when the ode is written in the normal form

$$y'' + p(x)y' + q(x)y = f(x)$$

Looking at the ode to solve shows that

$$p(x) = \frac{-1 - 2x}{x}$$

Therefore

$$y_2(x) = e^x \left(\int e^{-\left(\int \frac{-1-2x}{x} dx\right)} e^{-2x} dx \right)$$

$$y_2(x) = e^x \int \frac{e^{2x+\ln(x)}}{e^{2x}}, dx$$

$$y_2(x) = e^x \left(\int x dx \right)$$

$$y_2(x) = \frac{x^2 e^x}{2}$$

Hence the solution is

$$\begin{aligned} y &= c_1 y_1(x) + c_2 y_2(x) \\ &= c_1 e^x + \frac{c_2 x^2 e^x}{2} \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^x + \frac{c_2 x^2 e^x}{2} \tag{1}$$

Verification of solutions

$$y = c_1 e^x + \frac{c_2 x^2 e^x}{2}$$

Verified OK.

12.11.1 Maple step by step solution

Let's solve

$$y'' x + (-1 - 2x) y' + (1 + x) y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{(1+x)y}{x} + \frac{(1+2x)y'}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - \frac{(1+2x)y'}{x} + \frac{(1+x)y}{x} = 0$$

□ Check to see if $x_0 = 0$ is a regular singular point

○ Define functions

$$[P_2(x) = -\frac{1+2x}{x}, P_3(x) = \frac{1+x}{x}]$$

○ $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = -1$$

○ $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 0$$

○ $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

• Multiply by denominators

$$y''x + (-1 - 2x)y' + (1 + x)y = 0$$

• Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

□ Rewrite ODE with series expansions

○ Convert $x^m \cdot y$ to series expansion for $m = 0..1$

$$x^m \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

○ Shift index using $k- > k - m$

$$x^m \cdot y = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

○ Convert $x^m \cdot y'$ to series expansion for $m = 0..1$

$$x^m \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1+m}$$

○ Shift index using $k- > k + 1 - m$

$$x^m \cdot y' = \sum_{k=-1+m}^{\infty} a_{k+1-m} (k+1-m+r) x^{k+r}$$

○ Convert $x \cdot y''$ to series expansion

$$x \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r-1}$$

- Shift index using $k \rightarrow k + 1$

$$x \cdot y'' = \sum_{k=-1}^{\infty} a_{k+1}(k+1+r)(k+r)x^{k+r}$$

Rewrite ODE with series expansions

$$a_0r(-2+r)x^{-1+r} + (a_1(1+r)(-1+r) - a_0(-1+2r))x^r + \left(\sum_{k=1}^{\infty} (a_{k+1}(k+1+r)(k+r-1) \right.$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$r(-2+r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \{0, 2\}$$

- Each term must be 0

$$a_1(1+r)(-1+r) - a_0(-1+2r) = 0$$

- Each term in the series must be 0, giving the recursion relation

$$a_{k+1}(k+1+r)(k+r-1) + a_k(-2k-2r+1) + a_{k-1} = 0$$

- Shift index using $k \rightarrow k + 1$

$$a_{k+2}(k+2+r)(k+r) + a_{k+1}(-2k-1-2r) + a_k = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+2} = \frac{2ka_{k+1} + 2ra_{k+1} - a_k + a_{k+1}}{(k+2+r)(k+r)}$$

- Recursion relation for $r = 0$

$$a_{k+2} = \frac{2ka_{k+1} - a_k + a_{k+1}}{(k+2)k}$$

- Series not valid for $r = 0$, division by 0 in the recursion relation at $k = 0$

$$a_{k+2} = \frac{2ka_{k+1} - a_k + a_{k+1}}{(k+2)k}$$

- Recursion relation for $r = 2$

$$a_{k+2} = \frac{2ka_{k+1} - a_k + 5a_{k+1}}{(k+4)(k+2)}$$

- Solution for $r = 2$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+2}, a_{k+2} = \frac{2ka_{k+1} - a_k + 5a_{k+1}}{(k+4)(k+2)}, 3a_1 - 3a_0 = 0 \right]$$

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacics algorithm  
  A Liouvillian solution exists  
  Reducible group (found an exponential solution)  
<- Kovacics algorithm successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve([x*diff(y(x),x$2)-(2*x+1)*diff(y(x),x)+(x+1)*y(x)=0,exp(x)],singsol=all)
```

$$y(x) = e^x (c_2 x^2 + c_1)$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 23

```
DSolve[x*y''[x]-(2*x+1)*y'[x]+(x+1)*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} e^x (c_2 x^2 + 2c_1)$$

13 Chapter 2. Section 2.7. HIGHER ORDER LINEAR EQUATIONS, COUPLED HARMONIC OSCILLATORS Page 98

13.1 problem 1	2169
13.2 problem 2	2174
13.3 problem 3	2179
13.4 problem 4	2184
13.5 problem 5	2189
13.6 problem 6	2191
13.7 problem 7	2193
13.8 problem 8	2199
13.9 problem 9	2205
13.10problem 10	2207
13.11problem 11	2209
13.12problem 12	2216
13.13problem 13	2223
13.14problem 14	2228
13.15problem 15	2230
13.16problem 16(a)	2238
13.17problem 16(b)	2244
13.18problem 17	2248
13.19problem 18	2256
13.20problem 19(a)	2266
13.21problem 19(b)	2273
13.22problem 19(c)	2280
13.23problem 20	2287

13.1 problem 1

13.1.1 Maple step by step solution 2170

Internal problem ID [6348]

Internal file name [OUTPUT/5596_Sunday_June_05_2022_03_44_27_PM_11838903/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Section 2.7. HIGHER ORDER LINEAR EQUATIONS, COUPLED HARMONIC OSCILLATORS Page 98

Problem number: 1.

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_3rd_order , _missing_x]]
```

$$y''' - 3y'' + 2y' = 0$$

The characteristic equation is

$$\lambda^3 - 3\lambda^2 + 2\lambda = 0$$

The roots of the above equation are

$$\lambda_1 = 0$$

$$\lambda_2 = 2$$

$$\lambda_3 = 1$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 + c_2e^x + e^{2x}c_3$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = 1$$

$$y_2 = e^x$$

$$y_3 = e^{2x}$$

Summary

The solution(s) found are the following

$$y = c_1 + c_2 e^x + e^{2x} c_3 \quad (1)$$

Verification of solutions

$$y = c_1 + c_2 e^x + e^{2x} c_3$$

Verified OK.

13.1.1 Maple step by step solution

Let's solve

$$y''' - 3y'' + 2y' = 0$$

- Highest derivative means the order of the ODE is 3
 y'''
- Convert linear ODE into a system of first order ODEs
 - Define new variable $y_1(x)$
 $y_1(x) = y$
 - Define new variable $y_2(x)$
 $y_2(x) = y'$
 - Define new variable $y_3(x)$
 $y_3(x) = y''$
 - Isolate for $y_3'(x)$ using original ODE
 $y_3'(x) = 3y_3(x) - 2y_2(x)$
Convert linear ODE into a system of first order ODEs
 $[y_2(x) = y_1'(x), y_3(x) = y_2'(x), y_3'(x) = 3y_3(x) - 2y_2(x)]$
- Define vector
$$\vec{y}(x) = \begin{bmatrix} y_1(x) \\ y_2(x) \\ y_3(x) \end{bmatrix}$$
- System to solve

$$\vec{y}'(x) = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & -2 & 3 \end{bmatrix} \cdot \vec{y}(x)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & -2 & 3 \end{bmatrix}$$

- Rewrite the system as

$$\vec{y}'(x) = A \cdot \vec{y}(x)$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[0, \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \right], \left[1, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right], \left[2, \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[0, \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

- Consider eigenpair

$$\left[1, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_2 = e^x \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[2, \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_3 = e^{2x} \cdot \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{y} = c_1 \vec{y}_1 + c_2 \vec{y}_2 + c_3 \vec{y}_3$$

- Substitute solutions into the general solution

$$\vec{y} = c_2 e^x \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + e^{2x} c_3 \cdot \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} + \begin{bmatrix} c_1 \\ 0 \\ 0 \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y = c_2 e^x + \frac{e^{2x} c_3}{4} + c_1$$

Maple trace

```

`Methods for third order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve(diff(y(x),x$3)-3*diff(y(x),x$2)+2*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = c_1 + c_2 e^{2x} + c_3 e^x$$

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 25

```
DSolve[y'''[x]-3*y''[x]+2*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^x + \frac{1}{2} c_2 e^{2x} + c_3$$

13.2 problem 2

13.2.1 Maple step by step solution 2175

Internal problem ID [6349]

Internal file name [OUTPUT/5597_Sunday_June_05_2022_03_44_29_PM_35554333/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Section 2.7. HIGHER ORDER LINEAR EQUATIONS, COUPLED HARMONIC OSCILLATORS Page 98

Problem number: 2.

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_3rd_order , _missing_x]]
```

$$y''' - 3y'' + 4y' - 2y = 0$$

The characteristic equation is

$$\lambda^3 - 3\lambda^2 + 4\lambda - 2 = 0$$

The roots of the above equation are

$$\lambda_1 = 1$$

$$\lambda_2 = 1 - i$$

$$\lambda_3 = 1 + i$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 e^x + e^{(1+i)x} c_2 + e^{(1-i)x} c_3$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = e^x$$

$$y_2 = e^{(1+i)x}$$

$$y_3 = e^{(1-i)x}$$

Summary

The solution(s) found are the following

$$y = c_1 e^x + e^{(1+i)x} c_2 + e^{(1-i)x} c_3 \quad (1)$$

Verification of solutions

$$y = c_1 e^x + e^{(1+i)x} c_2 + e^{(1-i)x} c_3$$

Verified OK.

13.2.1 Maple step by step solution

Let's solve

$$y''' - 3y'' + 4y' - 2y = 0$$

- Highest derivative means the order of the ODE is 3

$$y'''$$

- Convert linear ODE into a system of first order ODEs

- Define new variable $y_1(x)$

$$y_1(x) = y$$

- Define new variable $y_2(x)$

$$y_2(x) = y'$$

- Define new variable $y_3(x)$

$$y_3(x) = y''$$

- Isolate for $y_3'(x)$ using original ODE

$$y_3'(x) = 3y_3(x) - 4y_2(x) + 2y_1(x)$$

Convert linear ODE into a system of first order ODEs

$$[y_2(x) = y_1'(x), y_3(x) = y_2'(x), y_3'(x) = 3y_3(x) - 4y_2(x) + 2y_1(x)]$$

- Define vector

$$\vec{y}(x) = \begin{bmatrix} y_1(x) \\ y_2(x) \\ y_3(x) \end{bmatrix}$$

- System to solve

$$\vec{y}'(x) = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 2 & -4 & 3 \end{bmatrix} \cdot \vec{y}(x)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 2 & -4 & 3 \end{bmatrix}$$

- Rewrite the system as

$$\vec{y}'(x) = A \cdot \vec{y}(x)$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[1, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right], \left[1 - \mathbf{I}, \begin{bmatrix} \frac{1}{2} \\ \frac{1}{2} + \frac{\mathbf{I}}{2} \\ 1 \end{bmatrix} \right], \left[1 + \mathbf{I}, \begin{bmatrix} -\frac{\mathbf{I}}{2} \\ \frac{1}{2} - \frac{\mathbf{I}}{2} \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[1, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_1 = e^x \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$\left[1 - \mathbf{I}, \begin{bmatrix} \frac{\mathbf{I}}{2} \\ \frac{1}{2} + \frac{\mathbf{I}}{2} \\ 1 \end{bmatrix} \right]$$

- Solution from eigenpair

$$e^{(1-i)x} \cdot \begin{bmatrix} \frac{1}{2} \\ \frac{1}{2} + \frac{i}{2} \\ 1 \end{bmatrix}$$

- Use Euler identity to write solution in terms of sin and cos

$$e^x \cdot (\cos(x) - i \sin(x)) \cdot \begin{bmatrix} \frac{1}{2} \\ \frac{1}{2} + \frac{i}{2} \\ 1 \end{bmatrix}$$

- Simplify expression

$$e^x \cdot \begin{bmatrix} \frac{1}{2}(\cos(x) - i \sin(x)) \\ (\frac{1}{2} + \frac{i}{2})(\cos(x) - i \sin(x)) \\ \cos(x) - i \sin(x) \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\left[\vec{y}_2(x) = e^x \cdot \begin{bmatrix} \frac{\sin(x)}{2} \\ \frac{\sin(x)}{2} + \frac{\cos(x)}{2} \\ \cos(x) \end{bmatrix}, \vec{y}_3(x) = e^x \cdot \begin{bmatrix} \frac{\cos(x)}{2} \\ \frac{\cos(x)}{2} - \frac{\sin(x)}{2} \\ -\sin(x) \end{bmatrix} \right]$$

- General solution to the system of ODEs

$$\vec{y} = c_1 \vec{y}_1 + c_2 \vec{y}_2(x) + c_3 \vec{y}_3(x)$$

- Substitute solutions into the general solution

$$\vec{y} = c_1 e^x \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + c_2 e^x \cdot \begin{bmatrix} \frac{\sin(x)}{2} \\ \frac{\sin(x)}{2} + \frac{\cos(x)}{2} \\ \cos(x) \end{bmatrix} + c_3 e^x \cdot \begin{bmatrix} \frac{\cos(x)}{2} \\ \frac{\cos(x)}{2} - \frac{\sin(x)}{2} \\ -\sin(x) \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y = \frac{e^x(c_2 \sin(x) + c_3 \cos(x) + 2c_1)}{2}$$

Maple trace

```
`Methods for third order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$3)-3*diff(y(x),x$2)+4*diff(y(x),x)-2*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^x(c_1 + \sin(x)c_2 + c_3 \cos(x))$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 22

```
DSolve[y'''[x]-3*y''[x]+4*y'[x]-2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x(c_2 \cos(x) + c_1 \sin(x) + c_3)$$

13.3 problem 3

13.3.1 Maple step by step solution 2180

Internal problem ID [6350]

Internal file name [OUTPUT/5598_Sunday_June_05_2022_03_44_31_PM_84736631/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Section 2.7. HIGHER ORDER LINEAR EQUATIONS, COUPLED HARMONIC OSCILLATORS Page 98

Problem number: 3.

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_3rd_order , _missing_x]]
```

$$y''' - y = 0$$

The characteristic equation is

$$\lambda^3 - 1 = 0$$

The roots of the above equation are

$$\lambda_1 = 1$$

$$\lambda_2 = -\frac{1}{2} - \frac{i\sqrt{3}}{2}$$

$$\lambda_3 = -\frac{1}{2} + \frac{i\sqrt{3}}{2}$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 e^x + e^{\left(-\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)x} c_2 + e^{\left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)x} c_3$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = e^x$$

$$y_2 = e^{\left(-\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)x}$$

$$y_3 = e^{\left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)x}$$

Summary

The solution(s) found are the following

$$y = c_1 e^x + e^{\left(-\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)x} c_2 + e^{\left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)x} c_3 \quad (1)$$

Verification of solutions

$$y = c_1 e^x + e^{\left(-\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)x} c_2 + e^{\left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)x} c_3$$

Verified OK.

13.3.1 Maple step by step solution

Let's solve

$$y''' - y = 0$$

- Highest derivative means the order of the ODE is 3

$$y'''$$

- Convert linear ODE into a system of first order ODEs

- Define new variable $y_1(x)$

$$y_1(x) = y$$

- Define new variable $y_2(x)$

$$y_2(x) = y'$$

- Define new variable $y_3(x)$

$$y_3(x) = y''$$

- Isolate for $y_3'(x)$ using original ODE

$$y_3'(x) = y_1(x)$$

Convert linear ODE into a system of first order ODEs

$$[y_2(x) = y_1'(x), y_3(x) = y_2'(x), y_3'(x) = y_1(x)]$$

- Define vector

$$\vec{y}(x) = \begin{bmatrix} y_1(x) \\ y_2(x) \\ y_3(x) \end{bmatrix}$$

- System to solve

$$\vec{y}'(x) = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix} \cdot \vec{y}(x)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}$$

- Rewrite the system as

$$\vec{y}'(x) = A \cdot \vec{y}(x)$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\begin{array}{c} \left[\begin{array}{c} 1 \\ 1 \\ 1 \end{array} \right] \\ \left[\begin{array}{c} 1 \\ 1 \\ 1 \end{array} \right] \end{array} \right], \left[-\frac{1}{2} - \frac{i\sqrt{3}}{2}, \begin{array}{c} \frac{1}{\left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)^2} \\ \frac{1}{-\frac{1}{2} - \frac{i\sqrt{3}}{2}} \\ 1 \end{array} \right], \left[-\frac{1}{2} + \frac{i\sqrt{3}}{2}, \begin{array}{c} \frac{1}{\left(-\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)^2} \\ \frac{1}{-\frac{1}{2} + \frac{i\sqrt{3}}{2}} \\ 1 \end{array} \right] \right]$$

- Consider eigenpair

$$\left[\begin{array}{c} \left[\begin{array}{c} 1 \\ 1 \\ 1 \end{array} \right] \\ 1 \end{array} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_1 = e^x \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$\left[-\frac{1}{2} - \frac{i\sqrt{3}}{2}, \begin{array}{c} \frac{1}{\left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)^2} \\ \frac{1}{-\frac{1}{2} - \frac{i\sqrt{3}}{2}} \\ 1 \end{array} \right]$$

- Solution from eigenpair

$$e^{\left(-\frac{1}{2}-\frac{i\sqrt{3}}{2}\right)x} \cdot \begin{bmatrix} \frac{1}{\left(-\frac{1}{2}-\frac{i\sqrt{3}}{2}\right)^2} \\ \frac{1}{-\frac{1}{2}-\frac{i\sqrt{3}}{2}} \\ 1 \end{bmatrix}$$

- Use Euler identity to write solution in terms of sin and cos

$$e^{-\frac{x}{2}} \cdot \left(\cos\left(\frac{\sqrt{3}x}{2}\right) - i \sin\left(\frac{\sqrt{3}x}{2}\right) \right) \cdot \begin{bmatrix} \frac{1}{\left(-\frac{1}{2}-\frac{i\sqrt{3}}{2}\right)^2} \\ \frac{1}{-\frac{1}{2}-\frac{i\sqrt{3}}{2}} \\ 1 \end{bmatrix}$$

- Simplify expression

$$e^{-\frac{x}{2}} \cdot \begin{bmatrix} \frac{\cos\left(\frac{\sqrt{3}x}{2}\right) - i \sin\left(\frac{\sqrt{3}x}{2}\right)}{\left(-\frac{1}{2}-\frac{i\sqrt{3}}{2}\right)^2} \\ \frac{\cos\left(\frac{\sqrt{3}x}{2}\right) - i \sin\left(\frac{\sqrt{3}x}{2}\right)}{-\frac{1}{2}-\frac{i\sqrt{3}}{2}} \\ \cos\left(\frac{\sqrt{3}x}{2}\right) - i \sin\left(\frac{\sqrt{3}x}{2}\right) \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\vec{y}_2(x) = e^{-\frac{x}{2}} \cdot \begin{bmatrix} -\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)}{2} - \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} \\ -\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} \\ \cos\left(\frac{\sqrt{3}x}{2}\right) \end{bmatrix}, \vec{y}_3(x) = e^{-\frac{x}{2}} \cdot \begin{bmatrix} -\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)}{2} \\ \frac{\cos\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)}{2} \\ -\sin\left(\frac{\sqrt{3}x}{2}\right) \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{y} = c_1 \vec{y}_1 + c_2 \vec{y}_2(x) + c_3 \vec{y}_3(x)$$

- Substitute solutions into the general solution

$$\vec{y} = c_1 e^x \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + c_2 e^{-\frac{x}{2}} \cdot \begin{bmatrix} -\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)}{2} - \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} \\ -\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} \\ \cos\left(\frac{\sqrt{3}x}{2}\right) \end{bmatrix} + c_3 e^{-\frac{x}{2}} \cdot \begin{bmatrix} -\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)}{2} \\ \frac{\cos\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)}{2} \\ -\sin\left(\frac{\sqrt{3}x}{2}\right) \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y = -\frac{e^{-\frac{x}{2}}(c_3\sqrt{3}+c_2)\cos\left(\frac{\sqrt{3}x}{2}\right)}{2} - \frac{e^{-\frac{x}{2}}(\sqrt{3}c_2-c_3)\sin\left(\frac{\sqrt{3}x}{2}\right)}{2} + c_1e^x$$

Maple trace

```

`Methods for third order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 35

```
dsolve(diff(y(x),x$3)-y(x)=0,y(x), singsol=all)
```

$$y(x) = e^x c_1 + c_2 e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) + c_3 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 52

```
DSolve[y'''[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x/2} \left(c_1 e^{3x/2} + c_2 \cos\left(\frac{\sqrt{3}x}{2}\right) + c_3 \sin\left(\frac{\sqrt{3}x}{2}\right) \right)$$

13.4 problem 4

13.4.1 Maple step by step solution 2185

Internal problem ID [6351]

Internal file name [OUTPUT/5599_Sunday_June_05_2022_03_44_32_PM_83856143/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Section 2.7. HIGHER ORDER LINEAR EQUATIONS, COUPLED HARMONIC OSCILLATORS Page 98

Problem number: 4.

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_3rd_order , _missing_x]]
```

$$y''' + y = 0$$

The characteristic equation is

$$\lambda^3 + 1 = 0$$

The roots of the above equation are

$$\begin{aligned}\lambda_1 &= -1 \\ \lambda_2 &= \frac{1}{2} - \frac{i\sqrt{3}}{2} \\ \lambda_3 &= \frac{1}{2} + \frac{i\sqrt{3}}{2}\end{aligned}$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 e^{-x} + e^{\left(\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)x} c_2 + e^{\left(\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)x} c_3$$

The fundamental set of solutions for the homogeneous solution are the following

$$\begin{aligned}y_1 &= e^{-x} \\ y_2 &= e^{\left(\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)x} \\ y_3 &= e^{\left(\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)x}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-x} + e^{\left(\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)x} c_2 + e^{\left(\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)x} c_3 \quad (1)$$

Verification of solutions

$$y = c_1 e^{-x} + e^{\left(\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)x} c_2 + e^{\left(\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)x} c_3$$

Verified OK.

13.4.1 Maple step by step solution

Let's solve

$$y''' + y = 0$$

- Highest derivative means the order of the ODE is 3

$$y'''$$

- Convert linear ODE into a system of first order ODEs

- Define new variable $y_1(x)$

$$y_1(x) = y$$

- Define new variable $y_2(x)$

$$y_2(x) = y'$$

- Define new variable $y_3(x)$

$$y_3(x) = y''$$

- Isolate for $y_3'(x)$ using original ODE

$$y_3'(x) = -y_1(x)$$

Convert linear ODE into a system of first order ODEs

$$[y_2(x) = y_1'(x), y_3(x) = y_2'(x), y_3'(x) = -y_1(x)]$$

- Define vector

$$\vec{y}(x) = \begin{bmatrix} y_1(x) \\ y_2(x) \\ y_3(x) \end{bmatrix}$$

- System to solve

$$\vec{y}'(x) = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ -1 & 0 & 0 \end{bmatrix} \cdot \vec{y}(x)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ -1 & 0 & 0 \end{bmatrix}$$

- Rewrite the system as

$$\vec{y}'(x) = A \cdot \vec{y}(x)$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[-1, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right], \left[\frac{1}{2} - \frac{i\sqrt{3}}{2}, \begin{bmatrix} \frac{1}{\left(\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)^2} \\ \frac{1}{\frac{1}{2} - \frac{i\sqrt{3}}{2}} \\ 1 \end{bmatrix} \right], \left[\frac{1}{2} + \frac{i\sqrt{3}}{2}, \begin{bmatrix} \frac{1}{\left(\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)^2} \\ \frac{1}{\frac{1}{2} + \frac{i\sqrt{3}}{2}} \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[-1, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_1 = e^{-x} \cdot \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$\left[\frac{1}{2} - \frac{i\sqrt{3}}{2}, \begin{bmatrix} \frac{1}{\left(\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)^2} \\ \frac{1}{\frac{1}{2} - \frac{i\sqrt{3}}{2}} \\ 1 \end{bmatrix} \right]$$

- Solution from eigenpair

$$e^{\left(\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)x} \cdot \begin{bmatrix} \frac{1}{\left(\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)^2} \\ \frac{1}{\frac{1}{2} - \frac{i\sqrt{3}}{2}} \\ 1 \end{bmatrix}$$

- Use Euler identity to write solution in terms of sin and cos

$$e^{\frac{x}{2}} \cdot \left(\cos\left(\frac{\sqrt{3}x}{2}\right) - i \sin\left(\frac{\sqrt{3}x}{2}\right) \right) \cdot \begin{bmatrix} \frac{1}{\left(\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)^2} \\ \frac{1}{\frac{1}{2} - \frac{i\sqrt{3}}{2}} \\ 1 \end{bmatrix}$$

- Simplify expression

$$e^{\frac{x}{2}} \cdot \begin{bmatrix} \frac{\cos\left(\frac{\sqrt{3}x}{2}\right) - i \sin\left(\frac{\sqrt{3}x}{2}\right)}{\left(\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)^2} \\ \frac{\cos\left(\frac{\sqrt{3}x}{2}\right) - i \sin\left(\frac{\sqrt{3}x}{2}\right)}{\frac{1}{2} - \frac{i\sqrt{3}}{2}} \\ \cos\left(\frac{\sqrt{3}x}{2}\right) - i \sin\left(\frac{\sqrt{3}x}{2}\right) \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\begin{bmatrix} \vec{y}_2(x) = e^{\frac{x}{2}} \cdot \begin{bmatrix} -\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} \\ \frac{\cos\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} \\ \cos\left(\frac{\sqrt{3}x}{2}\right) \end{bmatrix}, \vec{y}_3(x) = e^{\frac{x}{2}} \cdot \begin{bmatrix} \frac{\cos\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)}{2} \\ \frac{\cos\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} - \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)}{2} \\ -\sin\left(\frac{\sqrt{3}x}{2}\right) \end{bmatrix} \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{y} = c_1 \vec{y}_1 + c_2 \vec{y}_2(x) + c_3 \vec{y}_3(x)$$

- Substitute solutions into the general solution

$$\vec{y} = c_1 e^{-x} \cdot \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} + c_2 e^{\frac{x}{2}} \cdot \begin{bmatrix} -\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} \\ \frac{\cos\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} \\ \cos\left(\frac{\sqrt{3}x}{2}\right) \end{bmatrix} + c_3 e^{\frac{x}{2}} \cdot \begin{bmatrix} \frac{\cos\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)}{2} \\ \frac{\cos\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} - \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)}{2} \\ -\sin\left(\frac{\sqrt{3}x}{2}\right) \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y = e^{-x} \left(-\frac{e^{\frac{3x}{2}} (-c_3\sqrt{3}+c_2) \cos\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{e^{\frac{3x}{2}} (\sqrt{3}c_2+c_3) \sin\left(\frac{\sqrt{3}x}{2}\right)}{2} + c_1 \right)$$

Maple trace

```
`Methods for third order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 37

```
dsolve(diff(y(x),x$3)+y(x)=0,y(x), singsol=all)
```

$$y(x) = \left(c_2 e^{\frac{3x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) + c_3 e^{\frac{3x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + c_1 \right) e^{-x}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 56

```
DSolve[y'''[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x} \left(c_3 e^{3x/2} \cos\left(\frac{\sqrt{3}x}{2}\right) + c_2 e^{3x/2} \sin\left(\frac{\sqrt{3}x}{2}\right) + c_1 \right)$$

13.5 problem 5

Internal problem ID [6352]

Internal file name [OUTPUT/5600_Sunday_June_05_2022_03_44_33_PM_8107633/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Section 2.7. HIGHER ORDER LINEAR EQUATIONS, COUPLED HARMONIC OSCILLATORS Page 98

Problem number: 5.

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_3rd_order , _missing_x]]
```

$$y''' + 3y'' + 3y' + y = 0$$

The characteristic equation is

$$\lambda^3 + 3\lambda^2 + 3\lambda + 1 = 0$$

The roots of the above equation are

$$\lambda_1 = -1$$

$$\lambda_2 = -1$$

$$\lambda_3 = -1$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 e^{-x} + x e^{-x} c_2 + c_3 x^2 e^{-x}$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = e^{-x}$$

$$y_2 = x e^{-x}$$

$$y_3 = x^2 e^{-x}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-x} + x e^{-x} c_2 + c_3 x^2 e^{-x} \quad (1)$$

Verification of solutions

$$y = c_1 e^{-x} + x e^{-x} c_2 + c_3 x^2 e^{-x}$$

Verified OK.

Maple trace

```
`Methods for third order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(diff(y(x),x$3)+3*diff(y(x),x$2)+3*diff(y(x),x)+y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{-x} (c_3 x^2 + c_2 x + c_1)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 23

```
DSolve[y'''[x]+3*y''[x]+3*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x} (x(c_3 x + c_2) + c_1)$$

13.6 problem 6

Internal problem ID [6353]

Internal file name [OUTPUT/5601_Sunday_June_05_2022_03_44_35_PM_64782230/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Section 2.7. HIGHER ORDER LINEAR EQUATIONS, COUPLED HARMONIC OSCILLATORS Page 98

Problem number: 6.

ODE order: 4.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_high_order , _missing_x]]
```

$$y'''' + 4y''' + 6y'' + 4y' + y = 0$$

The characteristic equation is

$$\lambda^4 + 4\lambda^3 + 6\lambda^2 + 4\lambda + 1 = 0$$

The roots of the above equation are

$$\lambda_1 = -1$$

$$\lambda_2 = -1$$

$$\lambda_3 = -1$$

$$\lambda_4 = -1$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 e^{-x} + x e^{-x} c_2 + c_3 x^2 e^{-x} + x^3 e^{-x} c_4$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = e^{-x}$$

$$y_2 = x e^{-x}$$

$$y_3 = x^2 e^{-x}$$

$$y_4 = x^3 e^{-x}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-x} + x e^{-x} c_2 + c_3 x^2 e^{-x} + x^3 e^{-x} c_4 \quad (1)$$

Verification of solutions

$$y = c_1 e^{-x} + x e^{-x} c_2 + c_3 x^2 e^{-x} + x^3 e^{-x} c_4$$

Verified OK.

Maple trace

```
`Methods for high order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```
dsolve(diff(y(x),x$4)+4*diff(y(x),x$3)+6*diff(y(x),x$2)+4*diff(y(x),x)+y(x)=0,y(x), singsol=
```

$$y(x) = e^{-x} (c_4 x^3 + c_3 x^2 + c_2 x + c_1)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 28

```
DSolve[y''''[x]+4*y'''[x]+6*y''[x]+4*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x} (x(x(c_4 x + c_3) + c_2) + c_1)$$

13.7 problem 7

13.7.1 Maple step by step solution 2194

Internal problem ID [6354]

Internal file name [OUTPUT/5602_Sunday_June_05_2022_03_44_36_PM_20432115/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Section 2.7. HIGHER ORDER LINEAR EQUATIONS, COUPLED HARMONIC OSCILLATORS Page 98

Problem number: 7.

ODE order: 4.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_high_order , _missing_x]]
```

$$y'''' - y = 0$$

The characteristic equation is

$$\lambda^4 - 1 = 0$$

The roots of the above equation are

$$\lambda_1 = 1$$

$$\lambda_2 = -1$$

$$\lambda_3 = i$$

$$\lambda_4 = -i$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 e^{-x} + c_2 e^x + e^{-ix} c_3 + e^{ix} c_4$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = e^{-x}$$

$$y_2 = e^x$$

$$y_3 = e^{-ix}$$

$$y_4 = e^{ix}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-x} + c_2 e^x + e^{-ix} c_3 + e^{ix} c_4 \quad (1)$$

Verification of solutions

$$y = c_1 e^{-x} + c_2 e^x + e^{-ix} c_3 + e^{ix} c_4$$

Verified OK.

13.7.1 Maple step by step solution

Let's solve

$$y'''' - y = 0$$

- Highest derivative means the order of the ODE is 4

$$y''''$$

- Convert linear ODE into a system of first order ODEs

- Define new variable $y_1(x)$

$$y_1(x) = y$$

- Define new variable $y_2(x)$

$$y_2(x) = y'$$

- Define new variable $y_3(x)$

$$y_3(x) = y''$$

- Define new variable $y_4(x)$

$$y_4(x) = y'''$$

- Isolate for $y_4'(x)$ using original ODE

$$y_4'(x) = y_1(x)$$

Convert linear ODE into a system of first order ODEs

$$[y_2(x) = y_1'(x), y_3(x) = y_2'(x), y_4(x) = y_3'(x), y_4'(x) = y_1(x)]$$

- Define vector

$$\vec{y}(x) = \begin{bmatrix} y_1(x) \\ y_2(x) \\ y_3(x) \\ y_4(x) \end{bmatrix}$$

- System to solve

$$\vec{y}'(x) = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \end{bmatrix} \cdot \vec{y}(x)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \end{bmatrix}$$

- Rewrite the system as

$$\vec{y}'(x) = A \cdot \vec{y}(x)$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[-1, \begin{bmatrix} -1 \\ 1 \\ -1 \\ 1 \end{bmatrix} \right] \right], \left[\left[1, \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} \right] \right], \left[\left[-I, \begin{bmatrix} -I \\ -1 \\ I \\ 1 \end{bmatrix} \right] \right], \left[\left[I, \begin{bmatrix} I \\ -1 \\ -I \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[\left[-1, \begin{bmatrix} -1 \\ 1 \\ -1 \\ 1 \end{bmatrix} \right] \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_1 = e^{-x} \cdot \begin{bmatrix} -1 \\ 1 \\ -1 \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[1, \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_2 = e^x \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$\left[-I, \begin{bmatrix} -I \\ -1 \\ I \\ 1 \end{bmatrix} \right]$$

- Solution from eigenpair

$$e^{-Ix} \cdot \begin{bmatrix} -I \\ -1 \\ I \\ 1 \end{bmatrix}$$

- Use Euler identity to write solution in terms of sin and cos

$$(\cos(x) - I \sin(x)) \cdot \begin{bmatrix} -I \\ -1 \\ I \\ 1 \end{bmatrix}$$

- Simplify expression

$$\begin{bmatrix} -I(\cos(x) - I \sin(x)) \\ -\cos(x) + I \sin(x) \\ I(\cos(x) - I \sin(x)) \\ \cos(x) - I \sin(x) \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\vec{y}_3(x) = \begin{bmatrix} -\sin(x) \\ -\cos(x) \\ \sin(x) \\ \cos(x) \end{bmatrix}, \vec{y}_4(x) = \begin{bmatrix} -\cos(x) \\ \sin(x) \\ \cos(x) \\ -\sin(x) \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{y} = c_1 \vec{y}_1 + c_2 \vec{y}_2 + c_3 \vec{y}_3(x) + c_4 \vec{y}_4(x)$$

- Substitute solutions into the general solution

$$\vec{y} = c_1 e^{-x} \cdot \begin{bmatrix} -1 \\ 1 \\ -1 \\ 1 \end{bmatrix} + c_2 e^x \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} + \begin{bmatrix} -c_3 \sin(x) - c_4 \cos(x) \\ -c_3 \cos(x) + c_4 \sin(x) \\ c_3 \sin(x) + c_4 \cos(x) \\ c_3 \cos(x) - c_4 \sin(x) \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y = -c_1 e^{-x} + c_2 e^x - c_4 \cos(x) - c_3 \sin(x)$$

Maple trace

```
`Methods for high order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 23

```
dsolve(diff(y(x),x$4)-y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{-x} + e^x c_2 + c_3 \sin(x) + c_4 \cos(x)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 30

```
DSolve[y''''[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^x + c_3 e^{-x} + c_2 \cos(x) + c_4 \sin(x)$$

13.8 problem 8

13.8.1 Maple step by step solution 2200

Internal problem ID [6355]

Internal file name [OUTPUT/5603_Sunday_June_05_2022_03_44_37_PM_61774592/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Section 2.7. HIGHER ORDER LINEAR EQUATIONS, COUPLED HARMONIC OSCILLATORS Page 98

Problem number: 8.

ODE order: 4.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_high_order , _missing_x]]
```

$$y'''' + 5y'' + 4y = 0$$

The characteristic equation is

$$\lambda^4 + 5\lambda^2 + 4 = 0$$

The roots of the above equation are

$$\lambda_1 = 2i$$

$$\lambda_2 = -2i$$

$$\lambda_3 = i$$

$$\lambda_4 = -i$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 e^{-ix} + e^{-2ix} c_2 + e^{2ix} c_3 + e^{ix} c_4$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = e^{-ix}$$

$$y_2 = e^{-2ix}$$

$$y_3 = e^{2ix}$$

$$y_4 = e^{ix}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-ix} + e^{-2ix} c_2 + e^{2ix} c_3 + e^{ix} c_4 \quad (1)$$

Verification of solutions

$$y = c_1 e^{-ix} + e^{-2ix} c_2 + e^{2ix} c_3 + e^{ix} c_4$$

Verified OK.

13.8.1 Maple step by step solution

Let's solve

$$y'''' + 5y'' + 4y = 0$$

- Highest derivative means the order of the ODE is 4

$$y''''$$

- Convert linear ODE into a system of first order ODEs

- Define new variable $y_1(x)$

$$y_1(x) = y$$

- Define new variable $y_2(x)$

$$y_2(x) = y'$$

- Define new variable $y_3(x)$

$$y_3(x) = y''$$

- Define new variable $y_4(x)$

$$y_4(x) = y'''$$

- Isolate for $y_4'(x)$ using original ODE

$$y_4'(x) = -5y_3(x) - 4y_1(x)$$

Convert linear ODE into a system of first order ODEs

$$[y_2(x) = y_1'(x), y_3(x) = y_2'(x), y_4(x) = y_3'(x), y_4'(x) = -5y_3(x) - 4y_1(x)]$$

- Define vector

$$\vec{y}(x) = \begin{bmatrix} y_1(x) \\ y_2(x) \\ y_3(x) \\ y_4(x) \end{bmatrix}$$

- System to solve

$$\vec{y}'(x) = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -4 & 0 & -5 & 0 \end{bmatrix} \cdot \vec{y}(x)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -4 & 0 & -5 & 0 \end{bmatrix}$$

- Rewrite the system as

$$\vec{y}'(x) = A \cdot \vec{y}(x)$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[-2I, \begin{bmatrix} -\frac{1}{8} \\ -\frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right], \left[-I, \begin{bmatrix} -1 \\ -1 \\ 1 \\ 1 \end{bmatrix} \right], \left[I, \begin{bmatrix} 1 \\ -1 \\ -1 \\ 1 \end{bmatrix} \right], \left[2I, \begin{bmatrix} \frac{1}{8} \\ -\frac{1}{4} \\ -\frac{1}{2} \\ 1 \end{bmatrix} \right] \right]$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$\left[-2I, \begin{bmatrix} -\frac{1}{8} \\ -\frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right]$$

- Solution from eigenpair

$$e^{-2Ix} \cdot \begin{bmatrix} -\frac{I}{8} \\ -\frac{1}{4} \\ \frac{I}{2} \\ 1 \end{bmatrix}$$

- Use Euler identity to write solution in terms of sin and cos

$$(\cos(2x) - I \sin(2x)) \cdot \begin{bmatrix} -\frac{I}{8} \\ -\frac{1}{4} \\ \frac{I}{2} \\ 1 \end{bmatrix}$$

- Simplify expression

$$\begin{bmatrix} -\frac{I}{8}(\cos(2x) - I \sin(2x)) \\ -\frac{\cos(2x)}{4} + \frac{I \sin(2x)}{4} \\ \frac{I}{2}(\cos(2x) - I \sin(2x)) \\ \cos(2x) - I \sin(2x) \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\vec{y}_1(x) = \begin{bmatrix} -\frac{\sin(2x)}{8} \\ -\frac{\cos(2x)}{4} \\ \frac{\sin(2x)}{2} \\ \cos(2x) \end{bmatrix}, \vec{y}_2(x) = \begin{bmatrix} -\frac{\cos(2x)}{8} \\ \frac{\sin(2x)}{4} \\ \frac{\cos(2x)}{2} \\ -\sin(2x) \end{bmatrix}$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$-I, \begin{bmatrix} -I \\ -1 \\ I \\ 1 \end{bmatrix}$$

- Solution from eigenpair

$$e^{-Ix} \cdot \begin{bmatrix} -I \\ -1 \\ I \\ 1 \end{bmatrix}$$

- Use Euler identity to write solution in terms of sin and cos

$$(\cos(x) - I \sin(x)) \cdot \begin{bmatrix} -I \\ -1 \\ I \\ 1 \end{bmatrix}$$

- Simplify expression

$$\begin{bmatrix} -I(\cos(x) - I \sin(x)) \\ -\cos(x) + I \sin(x) \\ I(\cos(x) - I \sin(x)) \\ \cos(x) - I \sin(x) \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\vec{y}_3(x) = \begin{bmatrix} -\sin(x) \\ -\cos(x) \\ \sin(x) \\ \cos(x) \end{bmatrix}, \vec{y}_4(x) = \begin{bmatrix} -\cos(x) \\ \sin(x) \\ \cos(x) \\ -\sin(x) \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{y} = c_1 \vec{y}_1(x) + c_2 \vec{y}_2(x) + c_3 \vec{y}_3(x) + c_4 \vec{y}_4(x)$$

- Substitute solutions into the general solution

$$\vec{y} = \begin{bmatrix} -c_4 \cos(x) - c_3 \sin(x) - \frac{c_2 \cos(2x)}{8} - \frac{\sin(2x)c_1}{8} \\ -c_3 \cos(x) + c_4 \sin(x) + \frac{c_2 \sin(2x)}{4} - \frac{c_1 \cos(2x)}{4} \\ c_3 \sin(x) + c_4 \cos(x) + \frac{c_2 \cos(2x)}{2} + \frac{\sin(2x)c_1}{2} \\ c_3 \cos(x) - c_4 \sin(x) - c_2 \sin(2x) + c_1 \cos(2x) \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y = -c_4 \cos(x) - c_3 \sin(x) - \frac{c_2 \cos(2x)}{8} - \frac{\sin(2x)c_1}{8}$$

Maple trace

```
`Methods for high order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
dsolve(diff(y(x),x$4)+5*diff(y(x),x$2)+4*y(x)=0,y(x), singsol=all)
```

$$y(x) = 2c_2 \cos(x)^2 + (2c_1 \sin(x) + c_4) \cos(x) + c_3 \sin(x) - c_2$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 30

```
DSolve[y''''[x]+5*y'''[x]+4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos(2x) + c_4 \sin(x) + \cos(x)(2c_2 \sin(x) + c_3)$$

13.9 problem 9

Internal problem ID [6356]

Internal file name [OUTPUT/5604_Sunday_June_05_2022_03_44_38_PM_42445937/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Section 2.7. HIGHER ORDER LINEAR EQUATIONS, COUPLED HARMONIC OSCILLATORS Page 98

Problem number: 9.

ODE order: 4.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_high_order , _missing_x]]
```

$$y'''' - 2a^2y'' + ya^4 = 0$$

The characteristic equation is

$$a^4 - 2a^2\lambda^2 + \lambda^4 = 0$$

The roots of the above equation are

$$\lambda_1 = a$$

$$\lambda_2 = a$$

$$\lambda_3 = -a$$

$$\lambda_4 = -a$$

Therefore the homogeneous solution is

$$y_h(x) = c_1e^{ax} + c_2xe^{ax} + e^{-ax}c_3 + xe^{-ax}c_4$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = e^{ax}$$

$$y_2 = xe^{ax}$$

$$y_3 = e^{-ax}$$

$$y_4 = xe^{-ax}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{ax} + c_2 x e^{ax} + e^{-ax} c_3 + x e^{-ax} c_4 \quad (1)$$

Verification of solutions

$$y = c_1 e^{ax} + c_2 x e^{ax} + e^{-ax} c_3 + x e^{-ax} c_4$$

Verified OK.

Maple trace

```
`Methods for high order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 26

```
dsolve(diff(y(x),x$4)-2*a^2*diff(y(x),x$2)+a^4*y(x)=0,y(x), singsol=all)
```

$$y(x) = (c_4 x + c_3) e^{-ax} + e^{ax} (c_2 x + c_1)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 38

```
DSolve[y''''[x]-2*a^2*y''[x]+a^4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-ax} (c_3 e^{2ax} + x(c_4 e^{2ax} + c_2) + c_1)$$

13.10 problem 10

Internal problem ID [6357]

Internal file name [OUTPUT/5605_Sunday_June_05_2022_03_44_40_PM_24673305/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Section 2.7. HIGHER ORDER LINEAR EQUATIONS, COUPLED HARMONIC OSCILLATORS Page 98

Problem number: 10.

ODE order: 4.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_high_order , _missing_x]]
```

$$y'''' + 2a^2y'' + ya^4 = 0$$

The characteristic equation is

$$a^4 + 2a^2\lambda^2 + \lambda^4 = 0$$

The roots of the above equation are

$$\lambda_1 = ia$$

$$\lambda_2 = -ia$$

$$\lambda_3 = ia$$

$$\lambda_4 = -ia$$

Therefore the homogeneous solution is

$$y_h(x) = e^{-iax}c_1 + xe^{-iax}c_2 + e^{iax}c_3 + xe^{iax}c_4$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = e^{-iax}$$

$$y_2 = xe^{-iax}$$

$$y_3 = e^{iax}$$

$$y_4 = xe^{iax}$$

Summary

The solution(s) found are the following

$$y = e^{-iax} c_1 + x e^{-iax} c_2 + e^{iax} c_3 + x e^{iax} c_4 \quad (1)$$

Verification of solutions

$$y = e^{-iax} c_1 + x e^{-iax} c_2 + e^{iax} c_3 + x e^{iax} c_4$$

Verified OK.

Maple trace

```
`Methods for high order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 25

```
dsolve(diff(y(x),x$4)+2*a^2*diff(y(x),x$2)+a^4*y(x)=0,y(x), singsol=all)
```

$$y(x) = (c_4 x + c_2) \cos(ax) + \sin(ax) (c_3 x + c_1)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 30

```
DSolve[y''''[x]+2*a^2*y''[x]+a^4*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow (c_2 x + c_1) \cos(ax) + (c_4 x + c_3) \sin(ax)$$

13.11 problem 11

13.11.1 Maple step by step solution 2210

Internal problem ID [6358]

Internal file name [OUTPUT/5606_Sunday_June_05_2022_03_44_41_PM_46376091/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Section 2.7. HIGHER ORDER LINEAR EQUATIONS, COUPLED HARMONIC OSCILLATORS Page 98

Problem number: 11.

ODE order: 4.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_high_order , _missing_x]]
```

$$y'''' + 2y''' + 2y'' + 2y' + y = 0$$

The characteristic equation is

$$\lambda^4 + 2\lambda^3 + 2\lambda^2 + 2\lambda + 1 = 0$$

The roots of the above equation are

$$\lambda_1 = i$$

$$\lambda_2 = -i$$

$$\lambda_3 = -1$$

$$\lambda_4 = -1$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 e^{-x} + x e^{-x} c_2 + e^{-ix} c_3 + e^{ix} c_4$$

The fundamental set of solutions for the homogeneous solution are the following

$$\begin{aligned}y_1 &= e^{-x} \\y_2 &= x e^{-x} \\y_3 &= e^{-ix} \\y_4 &= e^{ix}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-x} + x e^{-x} c_2 + e^{-ix} c_3 + e^{ix} c_4 \quad (1)$$

Verification of solutions

$$y = c_1 e^{-x} + x e^{-x} c_2 + e^{-ix} c_3 + e^{ix} c_4$$

Verified OK.

13.11.1 Maple step by step solution

Let's solve

$$y'''' + 2y''' + 2y'' + 2y' + y = 0$$

- Highest derivative means the order of the ODE is 4
 y''''
 - Convert linear ODE into a system of first order ODEs
 - Define new variable $y_1(x)$
 $y_1(x) = y$
 - Define new variable $y_2(x)$
 $y_2(x) = y'$
 - Define new variable $y_3(x)$
 $y_3(x) = y''$
 - Define new variable $y_4(x)$
 $y_4(x) = y'''$
 - Isolate for $y_4'(x)$ using original ODE
 $y_4'(x) = -2y_4(x) - 2y_3(x) - 2y_2(x) - y_1(x)$
- Convert linear ODE into a system of first order ODEs

$$[y_2(x) = y_1'(x), y_3(x) = y_2'(x), y_4(x) = y_3'(x), y_4'(x) = -2y_4(x) - 2y_3(x) - 2y_2(x) - y_1(x)]$$

- Define vector

$$\vec{y}(x) = \begin{bmatrix} y_1(x) \\ y_2(x) \\ y_3(x) \\ y_4(x) \end{bmatrix}$$

- System to solve

$$\vec{y}'(x) = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & -2 & -2 & -2 \end{bmatrix} \cdot \vec{y}(x)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & -2 & -2 & -2 \end{bmatrix}$$

- Rewrite the system as

$$\vec{y}'(x) = A \cdot \vec{y}(x)$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[\begin{bmatrix} -1 \\ 1 \\ -1 \\ 1 \end{bmatrix} \right], \left[\begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} \right], \left[\begin{bmatrix} -I \\ -1 \\ I \\ 1 \end{bmatrix} \right], \left[\begin{bmatrix} I \\ -1 \\ -I \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair, with eigenvalue of algebraic multiplicity 2

$$\left[-1, \begin{bmatrix} -1 \\ 1 \\ -1 \\ 1 \end{bmatrix} \right]$$

- First solution from eigenvalue -1

$$\vec{y}_1(x) = e^{-x} \cdot \begin{bmatrix} -1 \\ 1 \\ -1 \\ 1 \end{bmatrix}$$

- Form of the 2nd homogeneous solution where \vec{p} is to be solved for, $\lambda = -1$ is the eigenvalue, a

$$\vec{y}_2(x) = e^{\lambda x} (x\vec{v} + \vec{p})$$

- Note that the x multiplying \vec{v} makes this solution linearly independent to the 1st solution obt

- Substitute $\vec{y}_2(x)$ into the homogeneous system

$$\lambda e^{\lambda x} (x\vec{v} + \vec{p}) + e^{\lambda x} \vec{v} = (e^{\lambda x} A) \cdot (x\vec{v} + \vec{p})$$

- Use the fact that \vec{v} is an eigenvector of A

$$\lambda e^{\lambda x} (x\vec{v} + \vec{p}) + e^{\lambda x} \vec{v} = e^{\lambda x} (\lambda x\vec{v} + A \cdot \vec{p})$$

- Simplify equation

$$\lambda \vec{p} + \vec{v} = A \cdot \vec{p}$$

- Make use of the identity matrix I

$$(\lambda \cdot I) \cdot \vec{p} + \vec{v} = A \cdot \vec{p}$$

- Condition \vec{p} must meet for $\vec{y}_2(x)$ to be a solution to the homogeneous system

$$(A - \lambda \cdot I) \cdot \vec{p} = \vec{v}$$

- Choose \vec{p} to use in the second solution to the homogeneous system from eigenvalue -1

$$\left(\begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & -2 & -2 & -2 \end{bmatrix} - (-1) \cdot \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \right) \cdot \vec{p} = \begin{bmatrix} -1 \\ 1 \\ -1 \\ 1 \end{bmatrix}$$

- Choice of \vec{p}

$$\vec{p} = \begin{bmatrix} -1 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

- Second solution from eigenvalue -1

$$\vec{y}_2(x) = e^{-x} \cdot \left(x \cdot \begin{bmatrix} -1 \\ 1 \\ -1 \\ 1 \end{bmatrix} + \begin{bmatrix} -1 \\ 0 \\ 0 \\ 0 \end{bmatrix} \right)$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$\left[-I, \begin{bmatrix} -I \\ -1 \\ I \\ 1 \end{bmatrix} \right]$$

- Solution from eigenpair

$$e^{-Ix} \cdot \begin{bmatrix} -I \\ -1 \\ I \\ 1 \end{bmatrix}$$

- Use Euler identity to write solution in terms of sin and cos

$$(\cos(x) - I \sin(x)) \cdot \begin{bmatrix} -I \\ -1 \\ I \\ 1 \end{bmatrix}$$

- Simplify expression

$$\begin{bmatrix} -I(\cos(x) - I \sin(x)) \\ -\cos(x) + I \sin(x) \\ I(\cos(x) - I \sin(x)) \\ \cos(x) - I \sin(x) \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\vec{y}_3(x) = \begin{bmatrix} -\sin(x) \\ -\cos(x) \\ \sin(x) \\ \cos(x) \end{bmatrix}, \vec{y}_4(x) = \begin{bmatrix} -\cos(x) \\ \sin(x) \\ \cos(x) \\ -\sin(x) \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{y} = c_1 \vec{y}_1(x) + c_2 \vec{y}_2(x) + c_3 \vec{y}_3(x) + c_4 \vec{y}_4(x)$$

- Substitute solutions into the general solution

$$\vec{y} = c_1 e^{-x} \cdot \begin{bmatrix} -1 \\ 1 \\ -1 \\ 1 \end{bmatrix} + c_2 e^{-x} \cdot \left(x \cdot \begin{bmatrix} -1 \\ 1 \\ -1 \\ 1 \end{bmatrix} + \begin{bmatrix} -1 \\ 0 \\ 0 \\ 0 \end{bmatrix} \right) + \begin{bmatrix} -c_3 \sin(x) - c_4 \cos(x) \\ -c_3 \cos(x) + c_4 \sin(x) \\ c_3 \sin(x) + c_4 \cos(x) \\ c_3 \cos(x) - c_4 \sin(x) \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y = ((-1 - x) c_2 - c_1) e^{-x} - c_3 \sin(x) - c_4 \cos(x)$$

Maple trace

```

`Methods for high order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(diff(y(x),x$4)+2*diff(y(x),x$3)+2*diff(y(x),x$2)+2*diff(y(x),x)+y(x)=0,y(x), singsol=
```

$$y(x) = e^{-x}(c_2x + c_1) + c_3 \sin(x) + c_4 \cos(x)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 34

```
DSolve[y''''[x]+2*y'''[x]+2*y''[x]+2*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x}(c_4x + c_1e^x \cos(x) + c_2e^x \sin(x) + c_3)$$

13.12 problem 12

13.12.1 Maple step by step solution 2217

Internal problem ID [6359]

Internal file name [OUTPUT/5607_Sunday_June_05_2022_03_44_42_PM_46403322/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Section 2.7. HIGHER ORDER LINEAR EQUATIONS, COUPLED HARMONIC OSCILLATORS Page 98

Problem number: 12.

ODE order: 4.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_high_order , _missing_x]]
```

$$y'''' + 2y''' - 2y'' - 6y' + 5y = 0$$

The characteristic equation is

$$\lambda^4 + 2\lambda^3 - 2\lambda^2 - 6\lambda + 5 = 0$$

The roots of the above equation are

$$\lambda_1 = -2 - i$$

$$\lambda_2 = -2 + i$$

$$\lambda_3 = 1$$

$$\lambda_4 = 1$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 e^x + c_2 x e^x + e^{(-2+i)x} c_3 + e^{(-2-i)x} c_4$$

The fundamental set of solutions for the homogeneous solution are the following

$$\begin{aligned}y_1 &= e^x \\y_2 &= x e^x \\y_3 &= e^{(-2+i)x} \\y_4 &= e^{(-2-i)x}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^x + c_2 x e^x + e^{(-2+i)x} c_3 + e^{(-2-i)x} c_4 \quad (1)$$

Verification of solutions

$$y = c_1 e^x + c_2 x e^x + e^{(-2+i)x} c_3 + e^{(-2-i)x} c_4$$

Verified OK.

13.12.1 Maple step by step solution

Let's solve

$$y'''' + 2y''' - 2y'' - 6y' + 5y = 0$$

- Highest derivative means the order of the ODE is 4

$$y''''$$

- Convert linear ODE into a system of first order ODEs

- Define new variable $y_1(x)$

$$y_1(x) = y$$

- Define new variable $y_2(x)$

$$y_2(x) = y'$$

- Define new variable $y_3(x)$

$$y_3(x) = y''$$

- Define new variable $y_4(x)$

$$y_4(x) = y'''$$

- Isolate for $y_4'(x)$ using original ODE

$$y_4'(x) = -2y_4(x) + 2y_3(x) + 6y_2(x) - 5y_1(x)$$

Convert linear ODE into a system of first order ODEs

$$[y_2(x) = y_1'(x), y_3(x) = y_2'(x), y_4(x) = y_3'(x), y_4'(x) = -2y_4(x) + 2y_3(x) + 6y_2(x) - 5y_1(x)]$$

- Define vector

$$\vec{y}(x) = \begin{bmatrix} y_1(x) \\ y_2(x) \\ y_3(x) \\ y_4(x) \end{bmatrix}$$

- System to solve

$$\vec{y}'(x) = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -5 & 6 & 2 & -2 \end{bmatrix} \cdot \vec{y}(x)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -5 & 6 & 2 & -2 \end{bmatrix}$$

- Rewrite the system as

$$\vec{y}'(x) = A \cdot \vec{y}(x)$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[\begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} \right], \left[\begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} \right], \left[-2 - I, \begin{bmatrix} -\frac{2}{125} + \frac{11I}{125} \\ \frac{3}{25} - \frac{4I}{25} \\ -\frac{2}{5} + \frac{I}{5} \\ 1 \end{bmatrix} \right], \left[-2 + I, \begin{bmatrix} -\frac{2}{125} - \frac{11I}{125} \\ \frac{3}{25} + \frac{4I}{25} \\ -\frac{2}{5} - \frac{I}{5} \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair, with eigenvalue of algebraic multiplicity 2

$$\left[1, \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} \right]$$

- First solution from eigenvalue 1

$$\vec{y}_1(x) = e^x \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}$$

- Form of the 2nd homogeneous solution where \vec{p} is to be solved for, $\lambda = 1$ is the eigenvalue, and

$$\vec{y}_2(x) = e^{\lambda x} (x\vec{v} + \vec{p})$$

- Note that the x multiplying \vec{v} makes this solution linearly independent to the 1st solution obtained

- Substitute $\vec{y}_2(x)$ into the homogeneous system

$$\lambda e^{\lambda x} (x\vec{v} + \vec{p}) + e^{\lambda x} \vec{v} = (e^{\lambda x} A) \cdot (x\vec{v} + \vec{p})$$

- Use the fact that \vec{v} is an eigenvector of A

$$\lambda e^{\lambda x} (x\vec{v} + \vec{p}) + e^{\lambda x} \vec{v} = e^{\lambda x} (\lambda x\vec{v} + A \cdot \vec{p})$$

- Simplify equation

$$\lambda \vec{p} + \vec{v} = A \cdot \vec{p}$$

- Make use of the identity matrix I

$$(\lambda \cdot I) \cdot \vec{p} + \vec{v} = A \cdot \vec{p}$$

- Condition \vec{p} must meet for $\vec{y}_2(x)$ to be a solution to the homogeneous system

$$(A - \lambda \cdot I) \cdot \vec{p} = \vec{v}$$

- Choose \vec{p} to use in the second solution to the homogeneous system from eigenvalue 1

$$\left(\begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -5 & 6 & 2 & -2 \end{bmatrix} - 1 \cdot \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \right) \cdot \vec{p} = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}$$

- Choice of \vec{p}

$$\vec{p} = \begin{bmatrix} -1 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

- Second solution from eigenvalue 1

$$\vec{y}_2(x) = e^x \cdot \left(x \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} + \begin{bmatrix} -1 \\ 0 \\ 0 \\ 0 \end{bmatrix} \right)$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$\left[-2 - I, \begin{bmatrix} -\frac{2}{125} + \frac{11I}{125} \\ \frac{3}{25} - \frac{4I}{25} \\ -\frac{2}{5} + \frac{I}{5} \\ 1 \end{bmatrix} \right]$$

- Solution from eigenpair

$$e^{(-2-I)x} \cdot \begin{bmatrix} -\frac{2}{125} + \frac{11I}{125} \\ \frac{3}{25} - \frac{4I}{25} \\ -\frac{2}{5} + \frac{I}{5} \\ 1 \end{bmatrix}$$

- Use Euler identity to write solution in terms of sin and cos

$$e^{-2x} \cdot (\cos(x) - I \sin(x)) \cdot \begin{bmatrix} -\frac{2}{125} + \frac{11I}{125} \\ \frac{3}{25} - \frac{4I}{25} \\ -\frac{2}{5} + \frac{I}{5} \\ 1 \end{bmatrix}$$

- Simplify expression

$$e^{-2x} \cdot \begin{bmatrix} \left(-\frac{2}{125} + \frac{11I}{125}\right) (\cos(x) - I \sin(x)) \\ \left(\frac{3}{25} - \frac{4I}{25}\right) (\cos(x) - I \sin(x)) \\ \left(-\frac{2}{5} + \frac{I}{5}\right) (\cos(x) - I \sin(x)) \\ \cos(x) - I \sin(x) \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\vec{y}_3(x) = e^{-2x} \cdot \begin{bmatrix} -\frac{2 \cos(x)}{125} + \frac{11 \sin(x)}{125} \\ \frac{3 \cos(x)}{25} - \frac{4 \sin(x)}{25} \\ -\frac{2 \cos(x)}{5} + \frac{\sin(x)}{5} \\ \cos(x) \end{bmatrix}, \vec{y}_4(x) = e^{-2x} \cdot \begin{bmatrix} \frac{2 \sin(x)}{125} + \frac{11 \cos(x)}{125} \\ -\frac{3 \sin(x)}{25} - \frac{4 \cos(x)}{25} \\ \frac{2 \sin(x)}{5} + \frac{\cos(x)}{5} \\ -\sin(x) \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{y} = c_1 \vec{y}_1(x) + c_2 \vec{y}_2(x) + c_3 \vec{y}_3(x) + c_4 \vec{y}_4(x)$$

- Substitute solutions into the general solution

$$\vec{y} = c_1 e^x \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} + c_2 e^x \cdot \left(x \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} + \begin{bmatrix} -1 \\ 0 \\ 0 \\ 0 \end{bmatrix} \right) + e^{-2x} c_3 \cdot \begin{bmatrix} -\frac{2 \cos(x)}{125} + \frac{11 \sin(x)}{125} \\ \frac{3 \cos(x)}{25} - \frac{4 \sin(x)}{25} \\ -\frac{2 \cos(x)}{5} + \frac{\sin(x)}{5} \\ \cos(x) \end{bmatrix} + e^{-2x} c_4 \cdot \begin{bmatrix} \frac{2 \sin(x)}{125} + \frac{11 \cos(x)}{125} \\ -\frac{3 \sin(x)}{25} - \frac{4 \cos(x)}{25} \\ \frac{2 \sin(x)}{5} + \frac{\cos(x)}{5} \\ -\sin(x) \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y = \left((c_2(x-1) + c_1) e^{3x} + \frac{(-2c_3 + 11c_4) \cos(x)}{125} + \frac{11(c_3 + \frac{2c_4}{11}) \sin(x)}{125} \right) e^{-2x}$$

Maple trace

```

`Methods for high order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 28

```
dsolve(diff(y(x),x$4)+2*diff(y(x),x$3)-2*diff(y(x),x$2)-6*diff(y(x),x)+5*y(x))=0,y(x), singso
```

$$y(x) = ((c_2x + c_1) e^{3x} + c_3 \sin(x) + c_4 \cos(x)) e^{-2x}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 35

```
DSolve[y''''[x]+2*y'''[x]-2*y''[x]-6*y'[x]+5*y[x]==0,y[x],x,IncludeSingularSolutions -> True
```

$$y(x) \rightarrow e^{-2x} (e^{3x} (c_4x + c_3) + c_2 \cos(x) + c_1 \sin(x))$$

13.13 problem 13

13.13.1 Maple step by step solution 2224

Internal problem ID [6360]

Internal file name [OUTPUT/5608_Sunday_June_05_2022_03_44_44_PM_58446449/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Section 2.7. HIGHER ORDER LINEAR EQUATIONS, COUPLED HARMONIC OSCILLATORS Page 98

Problem number: 13.

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_3rd_order , _missing_x]]
```

$$y''' - 6y'' + 11y' - 6y = 0$$

The characteristic equation is

$$\lambda^3 - 6\lambda^2 + 11\lambda - 6 = 0$$

The roots of the above equation are

$$\lambda_1 = 1$$

$$\lambda_2 = 2$$

$$\lambda_3 = 3$$

Therefore the homogeneous solution is

$$y_h(x) = c_1e^x + c_2e^{2x} + c_3e^{3x}$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = e^x$$

$$y_2 = e^{2x}$$

$$y_3 = e^{3x}$$

Summary

The solution(s) found are the following

$$y = c_1e^x + c_2e^{2x} + c_3e^{3x} \quad (1)$$

Verification of solutions

$$y = c_1e^x + c_2e^{2x} + c_3e^{3x}$$

Verified OK.

13.13.1 Maple step by step solution

Let's solve

$$y''' - 6y'' + 11y' - 6y = 0$$

- Highest derivative means the order of the ODE is 3

$$y'''$$

- Convert linear ODE into a system of first order ODEs

- Define new variable $y_1(x)$

$$y_1(x) = y$$

- Define new variable $y_2(x)$

$$y_2(x) = y'$$

- Define new variable $y_3(x)$

$$y_3(x) = y''$$

- Isolate for $y_3'(x)$ using original ODE

$$y_3'(x) = 6y_3(x) - 11y_2(x) + 6y_1(x)$$

Convert linear ODE into a system of first order ODEs

$$[y_2(x) = y_1'(x), y_3(x) = y_2'(x), y_3'(x) = 6y_3(x) - 11y_2(x) + 6y_1(x)]$$

- Define vector

$$\vec{y}(x) = \begin{bmatrix} y_1(x) \\ y_2(x) \\ y_3(x) \end{bmatrix}$$

- System to solve

$$\vec{y}'(x) = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 6 & -11 & 6 \end{bmatrix} \cdot \vec{y}(x)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 6 & -11 & 6 \end{bmatrix}$$

- Rewrite the system as

$$\vec{y}'(x) = A \cdot \vec{y}(x)$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[1, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right], \left[2, \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right], \left[3, \begin{bmatrix} \frac{1}{9} \\ \frac{1}{3} \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[1, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_1 = e^x \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[2, \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_2 = e^{2x} \cdot \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[3, \begin{bmatrix} \frac{1}{9} \\ \frac{1}{3} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_3 = e^{3x} \cdot \begin{bmatrix} \frac{1}{9} \\ \frac{1}{3} \\ 1 \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{y} = c_1 \vec{y}_1 + c_2 \vec{y}_2 + c_3 \vec{y}_3$$

- Substitute solutions into the general solution

$$\vec{y} = c_1 e^x \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + c_2 e^{2x} \cdot \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} + c_3 e^{3x} \cdot \begin{bmatrix} \frac{1}{9} \\ \frac{1}{3} \\ 1 \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y = c_1 e^x + \frac{c_2 e^{2x}}{4} + \frac{c_3 e^{3x}}{9}$$

Maple trace

```

`Methods for third order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(diff(y(x),x$3)-6*diff(y(x),x$2)+11*diff(y(x),x)-6*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{2x}c_1 + c_2e^{3x} + c_3e^x$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 25

```
DSolve[y'''[x]-6*y''[x]+11*y'[x]-6*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^x(e^x(c_3e^x + c_2) + c_1)$$

13.14 problem 14

Internal problem ID [6361]

Internal file name [OUTPUT/5609_Sunday_June_05_2022_03_44_45_PM_71354160/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Section 2.7. HIGHER ORDER LINEAR EQUATIONS, COUPLED HARMONIC OSCILLATORS Page 98

Problem number: 14.

ODE order: 4.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_high_order , _missing_x]]
```

$$y'''' + y''' - 3y'' - 5y' - 2y = 0$$

The characteristic equation is

$$\lambda^4 + \lambda^3 - 3\lambda^2 - 5\lambda - 2 = 0$$

The roots of the above equation are

$$\lambda_1 = 2$$

$$\lambda_2 = -1$$

$$\lambda_3 = -1$$

$$\lambda_4 = -1$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 e^{-x} + x e^{-x} c_2 + c_3 x^2 e^{-x} + e^{2x} c_4$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = e^{-x}$$

$$y_2 = x e^{-x}$$

$$y_3 = x^2 e^{-x}$$

$$y_4 = e^{2x}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-x} + x e^{-x} c_2 + c_3 x^2 e^{-x} + e^{2x} c_4 \quad (1)$$

Verification of solutions

$$y = c_1 e^{-x} + x e^{-x} c_2 + c_3 x^2 e^{-x} + e^{2x} c_4$$

Verified OK.

Maple trace

```
`Methods for high order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 26

```
dsolve(diff(y(x),x$4)+diff(y(x),x$3)-3*diff(y(x),x$2)-5*diff(y(x),x)-2*y(x)=0,y(x), singsol=
```

$$y(x) = (c_4 x^2 + c_3 x + c_2) e^{-x} + e^{2x} c_1$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 32

```
DSolve[y''''[x]+y'''[x]-3*y''[x]-5*y'[x]-2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x} (c_3 x^2 + c_2 x + c_4 e^{3x} + c_1)$$

13.15 problem 15

13.15.1 Maple step by step solution 2231

Internal problem ID [6362]

Internal file name [OUTPUT/5610_Sunday_June_05_2022_03_44_46_PM_33953325/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Section 2.7. HIGHER ORDER LINEAR EQUATIONS, COUPLED HARMONIC OSCILLATORS Page 98

Problem number: 15.

ODE order: 5.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_high_order , _missing_x]]
```

$$y^{(5)} - 6y'''' - 8y''' + 48y'' + 16y' - 96y = 0$$

The characteristic equation is

$$\lambda^5 - 6\lambda^4 - 8\lambda^3 + 48\lambda^2 + 16\lambda - 96 = 0$$

The roots of the above equation are

$$\lambda_1 = 6$$

$$\lambda_2 = 2$$

$$\lambda_3 = 2$$

$$\lambda_4 = -2$$

$$\lambda_5 = -2$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 e^{-2x} + c_2 x e^{-2x} + e^{2x} c_3 + x e^{2x} c_4 + e^{6x} c_5$$

The fundamental set of solutions for the homogeneous solution are the following

$$\begin{aligned}y_1 &= e^{-2x} \\y_2 &= x e^{-2x} \\y_3 &= e^{2x} \\y_4 &= e^{2x}x \\y_5 &= e^{6x}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-2x} + c_2 x e^{-2x} + e^{2x} c_3 + x e^{2x} c_4 + e^{6x} c_5 \quad (1)$$

Verification of solutions

$$y = c_1 e^{-2x} + c_2 x e^{-2x} + e^{2x} c_3 + x e^{2x} c_4 + e^{6x} c_5$$

Verified OK.

13.15.1 Maple step by step solution

Let's solve

$$y^{(5)} - 6y'''' - 8y''' + 48y'' + 16y' - 96y = 0$$

- Highest derivative means the order of the ODE is 5
- Convert linear ODE into a system of first order ODEs
 - Define new variable $y_1(x)$
$$y_1(x) = y$$
 - Define new variable $y_2(x)$
$$y_2(x) = y'$$
 - Define new variable $y_3(x)$
$$y_3(x) = y''$$
 - Define new variable $y_4(x)$
$$y_4(x) = y'''$$
 - Define new variable $y_5(x)$
$$y_5(x) = y''''$$

- Isolate for $y_5'(x)$ using original ODE

$$y_5'(x) = 6y_5(x) + 8y_4(x) - 48y_3(x) - 16y_2(x) + 96y_1(x)$$

Convert linear ODE into a system of first order ODEs

$$[y_2(x) = y_1'(x), y_3(x) = y_2'(x), y_4(x) = y_3'(x), y_5(x) = y_4'(x), y_5'(x) = 6y_5(x) + 8y_4(x) - 48y_3(x)]$$

- Define vector

$$\vec{y}(x) = \begin{bmatrix} y_1(x) \\ y_2(x) \\ y_3(x) \\ y_4(x) \\ y_5(x) \end{bmatrix}$$

- System to solve

$$\vec{y}'(x) = \begin{bmatrix} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 96 & -16 & -48 & 8 & 6 \end{bmatrix} \cdot \vec{y}(x)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 96 & -16 & -48 & 8 & 6 \end{bmatrix}$$

- Rewrite the system as

$$\vec{y}'(x) = A \cdot \vec{y}(x)$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[\begin{array}{c} -2, \\ \left[\begin{array}{c} \frac{1}{16} \\ -\frac{1}{8} \\ \frac{1}{4} \\ -\frac{1}{2} \\ 1 \end{array} \right] \end{array} \right], \left[\begin{array}{c} -2, \\ \left[\begin{array}{c} 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{array} \right] \end{array} \right], \left[\begin{array}{c} 2, \\ \left[\begin{array}{c} \frac{1}{16} \\ \frac{1}{8} \\ \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{array} \right] \end{array} \right], \left[\begin{array}{c} 2, \\ \left[\begin{array}{c} 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{array} \right] \end{array} \right], \left[\begin{array}{c} 6, \\ \left[\begin{array}{c} \frac{1}{1296} \\ \frac{1}{216} \\ \frac{1}{36} \\ \frac{1}{6} \\ 1 \end{array} \right] \end{array} \right] \right]$$

- Consider eigenpair, with eigenvalue of algebraic multiplicity 2

$$\left[\begin{array}{c} -2, \\ \left[\begin{array}{c} \frac{1}{16} \\ -\frac{1}{8} \\ \frac{1}{4} \\ -\frac{1}{2} \\ 1 \end{array} \right] \end{array} \right]$$

- First solution from eigenvalue -2

$$\vec{y}_1(x) = e^{-2x} \cdot \left[\begin{array}{c} \frac{1}{16} \\ -\frac{1}{8} \\ \frac{1}{4} \\ -\frac{1}{2} \\ 1 \end{array} \right]$$

- Form of the 2nd homogeneous solution where \vec{p} is to be solved for, $\lambda = -2$ is the eigenvalue, a

$$\vec{y}_2(x) = e^{\lambda x} (x\vec{v} + \vec{p})$$

- Note that the x multiplying \vec{v} makes this solution linearly independent to the 1st solution obt

- Substitute $\vec{y}_2(x)$ into the homogeneous system

$$\lambda e^{\lambda x} (x\vec{v} + \vec{p}) + e^{\lambda x} \vec{v} = (e^{\lambda x} A) \cdot (x\vec{v} + \vec{p})$$

- Use the fact that \vec{v} is an eigenvector of A

$$\lambda e^{\lambda x} (x\vec{v} + \vec{p}) + e^{\lambda x} \vec{v} = e^{\lambda x} (\lambda x\vec{v} + A \cdot \vec{p})$$

- Simplify equation

$$\lambda \vec{p} + \vec{v} = A \cdot \vec{p}$$

- Make use of the identity matrix I

$$(\lambda \cdot I) \cdot \vec{p} + \vec{v} = A \cdot \vec{p}$$

- Condition \vec{p} must meet for $\vec{y}_2(x)$ to be a solution to the homogeneous system

$$(A - \lambda \cdot I) \cdot \vec{p} = \vec{v}$$

- Choose \vec{p} to use in the second solution to the homogeneous system from eigenvalue -2

$$\left(\begin{bmatrix} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 96 & -16 & -48 & 8 & 6 \end{bmatrix} - (-2) \cdot \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix} \right) \cdot \vec{p} = \begin{bmatrix} \frac{1}{16} \\ -\frac{1}{8} \\ \frac{1}{4} \\ -\frac{1}{2} \\ 1 \end{bmatrix}$$

- Choice of \vec{p}

$$\vec{p} = \begin{bmatrix} \frac{1}{32} \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

- Second solution from eigenvalue -2

$$\vec{y}_2(x) = e^{-2x} \cdot \left(x \cdot \begin{bmatrix} \frac{1}{16} \\ -\frac{1}{8} \\ \frac{1}{4} \\ -\frac{1}{2} \\ 1 \end{bmatrix} + \begin{bmatrix} \frac{1}{32} \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} \right)$$

- Consider eigenpair, with eigenvalue of algebraic multiplicity 2

$$\left[2, \begin{bmatrix} \frac{1}{16} \\ \frac{1}{8} \\ \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right]$$

- First solution from eigenvalue 2

$$\vec{y}_3(x) = e^{2x} \cdot \begin{bmatrix} \frac{1}{16} \\ \frac{1}{8} \\ \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix}$$

- Form of the 2nd homogeneous solution where \vec{p} is to be solved for, $\lambda = 2$ is the eigenvalue, and

$$\vec{y}_4(x) = e^{\lambda x} (x\vec{v} + \vec{p})$$

- Note that the x multiplying \vec{v} makes this solution linearly independent to the 1st solution obtained

- Substitute $\vec{y}_4(x)$ into the homogeneous system

$$\lambda e^{\lambda x} (x\vec{v} + \vec{p}) + e^{\lambda x} \vec{v} = (e^{\lambda x} A) \cdot (x\vec{v} + \vec{p})$$

- Use the fact that \vec{v} is an eigenvector of A

$$\lambda e^{\lambda x} (x\vec{v} + \vec{p}) + e^{\lambda x} \vec{v} = e^{\lambda x} (\lambda x\vec{v} + A \cdot \vec{p})$$

- Simplify equation

$$\lambda \vec{p} + \vec{v} = A \cdot \vec{p}$$

- Make use of the identity matrix I

$$(\lambda \cdot I) \cdot \vec{p} + \vec{v} = A \cdot \vec{p}$$

- Condition \vec{p} must meet for $\vec{y}_4(x)$ to be a solution to the homogeneous system

$$(A - \lambda \cdot I) \cdot \vec{p} = \vec{v}$$

- Choose \vec{p} to use in the second solution to the homogeneous system from eigenvalue 2

$$\left(\begin{bmatrix} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 96 & -16 & -48 & 8 & 6 \end{bmatrix} - 2 \cdot \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix} \right) \cdot \vec{p} = \begin{bmatrix} \frac{1}{16} \\ \frac{1}{8} \\ \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix}$$

- Choice of \vec{p}

$$\vec{p} = \begin{bmatrix} -\frac{1}{32} \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

- Second solution from eigenvalue 2

$$\vec{y}_4(x) = e^{2x} \cdot \left(x \cdot \begin{bmatrix} \frac{1}{16} \\ \frac{1}{8} \\ \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} + \begin{bmatrix} -\frac{1}{32} \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} \right)$$

- Consider eigenpair

$$\left[6, \begin{bmatrix} \frac{1}{1296} \\ \frac{1}{216} \\ \frac{1}{36} \\ \frac{1}{6} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_5 = e^{6x} \cdot \begin{bmatrix} \frac{1}{1296} \\ \frac{1}{216} \\ \frac{1}{36} \\ \frac{1}{6} \\ 1 \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{y} = c_1 \vec{y}_1(x) + c_2 \vec{y}_2(x) + c_3 \vec{y}_3(x) + c_4 \vec{y}_4(x) + c_5 \vec{y}_5$$

- Substitute solutions into the general solution

$$\vec{y} = c_1 e^{-2x} \cdot \begin{bmatrix} \frac{1}{16} \\ -\frac{1}{8} \\ \frac{1}{4} \\ -\frac{1}{2} \\ 1 \end{bmatrix} + c_2 e^{-2x} \cdot \left(x \cdot \begin{bmatrix} \frac{1}{16} \\ -\frac{1}{8} \\ \frac{1}{4} \\ -\frac{1}{2} \\ 1 \end{bmatrix} + \begin{bmatrix} \frac{1}{32} \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} \right) + e^{2x} c_3 \cdot \begin{bmatrix} \frac{1}{16} \\ \frac{1}{8} \\ \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} + e^{2x} c_4 \cdot \left(x \cdot \begin{bmatrix} \frac{1}{16} \\ \frac{1}{8} \\ \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right)$$

- First component of the vector is the solution to the ODE

$$y = \frac{(c_2(1+2x)+2c_1)e^{-2x}}{32} + \frac{((2x-1)c_4+2c_3)e^{2x}}{32} + \frac{e^{6x}c_5}{1296}$$

Maple trace

```

`Methods for high order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 31

```
dsolve(diff(y(x),x$5)-6*diff(y(x),x$4)-8*diff(y(x),x$3)+48*diff(y(x),x$2)+16*diff(y(x),x)-96*y(x),x)-96*y(x),x)
```

$$y(x) = (c_5 x + c_4) e^{-2x} + (c_3 x + c_2) e^{2x} + c_1 e^{6x}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 43

```
DSolve[y''''[x]-6*y''''[x]-8*y''''[x]+48*y''[x]+16*y'[x]-96*y[x]==0,y[x],x,IncludeSingularSolutions->True]
```

$$y(x) \rightarrow e^{-2x} (c_2 x + c_3 e^{4x} + c_4 e^{4x} x + c_5 e^{8x} + c_1)$$

13.16 problem 16(a)

13.16.1 Maple step by step solution 2239

Internal problem ID [6363]

Internal file name [OUTPUT/5611_Sunday_June_05_2022_03_44_47_PM_74242486/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Section 2.7. HIGHER ORDER LINEAR EQUATIONS, COUPLED HARMONIC OSCILLATORS Page 98

Problem number: 16(a).

ODE order: 4.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_high_order , _quadrature]]
```

$$y'''' = 0$$

The characteristic equation is

$$\lambda^4 = 0$$

The roots of the above equation are

$$\lambda_1 = 0$$

$$\lambda_2 = 0$$

$$\lambda_3 = 0$$

$$\lambda_4 = 0$$

Therefore the homogeneous solution is

$$y_h(x) = c_4x^3 + c_3x^2 + c_2x + c_1$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = 1$$

$$y_2 = x$$

$$y_3 = x^2$$

$$y_4 = x^3$$

Summary

The solution(s) found are the following

$$y = c_4x^3 + c_3x^2 + c_2x + c_1 \quad (1)$$

Verification of solutions

$$y = c_4x^3 + c_3x^2 + c_2x + c_1$$

Verified OK.

13.16.1 Maple step by step solution

Let's solve

$$y'''' = 0$$

- Highest derivative means the order of the ODE is 4
 y''''
- Convert linear ODE into a system of first order ODEs
 - Define new variable $y_1(x)$
 $y_1(x) = y$
 - Define new variable $y_2(x)$
 $y_2(x) = y'$
 - Define new variable $y_3(x)$
 $y_3(x) = y''$
 - Define new variable $y_4(x)$
 $y_4(x) = y'''$
 - Isolate for $y_4'(x)$ using original ODE
 $y_4'(x) = 0$
Convert linear ODE into a system of first order ODEs
 $[y_2(x) = y_1'(x), y_3(x) = y_2'(x), y_4(x) = y_3'(x), y_4'(x) = 0]$
- Define vector

$$\vec{y}(x) = \begin{bmatrix} y_1(x) \\ y_2(x) \\ y_3(x) \\ y_4(x) \end{bmatrix}$$

- System to solve

$$\vec{y}'(x) = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \cdot \vec{y}(x)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

- Rewrite the system as

$$\vec{y}'(x) = A \cdot \vec{y}(x)$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[0, \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} \right], \left[0, \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} \right], \left[0, \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} \right], \left[0, \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[0, \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

- Consider eigenpair

$$\left[0, \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_2 = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

- Consider eigenpair

$$\left[0, \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_3 = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

- Consider eigenpair

$$\begin{bmatrix} 0, \\ \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} \end{bmatrix}$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_4 = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{y} = c_1 \vec{y}_1 + c_2 \vec{y}_2 + c_3 \vec{y}_3 + c_4 \vec{y}_4$$

- Substitute solutions into the general solution

$$\vec{y} = \begin{bmatrix} c_1 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y = c_1$$

Maple trace

```
`Methods for high order ODEs:
--- Trying classification methods ---
trying a quadrature
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(diff(y(x),x$4)=0,y(x), singsol=all)
```

$$y(x) = \frac{1}{6}c_1x^3 + \frac{1}{2}c_2x^2 + c_3x + c_4$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 22

```
DSolve[y''''[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x(x(c_4x + c_3) + c_2) + c_1$$

13.17 problem 16(b)

Internal problem ID [6364]

Internal file name [OUTPUT/5612_Sunday_June_05_2022_03_44_49_PM_1015288/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Section 2.7. HIGHER ORDER LINEAR EQUATIONS, COUPLED HARMONIC OSCILLATORS Page 98

Problem number: 16(b).

ODE order: 4.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_high_order , _quadrature]]
```

$$y'''' = \sin(x) + 24$$

This is higher order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE And y_p is a particular solution to the nonhomogeneous ODE. y_h is the solution to

$$y'''' = 0$$

The characteristic equation is

$$\lambda^4 = 0$$

The roots of the above equation are

$$\lambda_1 = 0$$

$$\lambda_2 = 0$$

$$\lambda_3 = 0$$

$$\lambda_4 = 0$$

Therefore the homogeneous solution is

$$y_h(x) = c_4x^3 + c_3x^2 + c_2x + c_1$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = 1$$

$$y_2 = x$$

$$y_3 = x^2$$

$$y_4 = x^3$$

Now the particular solution to the given ODE is found

$$y'''' = \sin(x) + 24$$

The particular solution is found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x) + 24$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1\}, \{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, x, x^2, x^3\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x\}, \{\cos(x), \sin(x)\}]$$

Since x is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x^2\}, \{\cos(x), \sin(x)\}]$$

Since x^2 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x^3\}, \{\cos(x), \sin(x)\}]$$

Since x^3 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x^4\}, \{\cos(x), \sin(x)\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1x^4 + A_2 \cos(x) + A_3 \sin(x)$$

The unknowns $\{A_1, A_2, A_3\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$24A_1 + A_2 \cos(x) + A_3 \sin(x) = \sin(x) + 24$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 1, A_2 = 0, A_3 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = x^4 + \sin(x)$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_4x^3 + c_3x^2 + c_2x + c_1) + (x^4 + \sin(x)) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_4x^3 + c_3x^2 + c_2x + c_1 + x^4 + \sin(x) \quad (1)$$

Verification of solutions

$$y = c_4x^3 + c_3x^2 + c_2x + c_1 + x^4 + \sin(x)$$

Verified OK.

Maple trace

```
`Methods for high order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(diff(y(x),x$4)=sin(x)+24,y(x), singsol=all)
```

$$y(x) = \frac{c_1 x^3}{6} + x^4 + \frac{c_2 x^2}{2} + \sin(x) + c_3 x + c_4$$

✓ Solution by Mathematica

Time used: 0.033 (sec). Leaf size: 29

```
DSolve[y''''[x]==Sin[x]+24,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^4 + c_4 x^3 + c_3 x^2 + \sin(x) + c_2 x + c_1$$

13.18 problem 17

13.18.1 Maple step by step solution 2250

Internal problem ID [6365]

Internal file name [OUTPUT/5613_Sunday_June_05_2022_03_44_50_PM_84082901/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Section 2.7. HIGHER ORDER LINEAR EQUATIONS, COUPLED HARMONIC OSCILLATORS Page 98

Problem number: 17.

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_3rd_order , _missing_y]]
```

$$y''' - 3y'' + 2y' = 10 + 42e^{3x}$$

This is higher order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE And y_p is a particular solution to the nonhomogeneous ODE. y_h is the solution to

$$y''' - 3y'' + 2y' = 0$$

The characteristic equation is

$$\lambda^3 - 3\lambda^2 + 2\lambda = 0$$

The roots of the above equation are

$$\lambda_1 = 0$$

$$\lambda_2 = 2$$

$$\lambda_3 = 1$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 + c_2 e^x + e^{2x} c_3$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = 1$$

$$y_2 = e^x$$

$$y_3 = e^{2x}$$

Now the particular solution to the given ODE is found

$$y''' - 3y'' + 2y' = 10 + 42 e^{3x}$$

The particular solution is found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$10 + 42 e^{3x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1\}, \{e^{3x}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, e^x, e^{2x}\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x\}, \{e^{3x}\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x + A_2 e^{3x}$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$6A_2 e^{3x} + 2A_1 = 10 + 42 e^{3x}$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 5, A_2 = 7]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = 5x + 7e^{3x}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 + c_2e^x + e^{2x}c_3) + (5x + 7e^{3x}) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 + c_2e^x + e^{2x}c_3 + 5x + 7e^{3x} \quad (1)$$

Verification of solutions

$$y = c_1 + c_2e^x + e^{2x}c_3 + 5x + 7e^{3x}$$

Verified OK.

13.18.1 Maple step by step solution

Let's solve

$$y''' - 3y'' + 2y' = 10 + 42e^{3x}$$

- Highest derivative means the order of the ODE is 3

y'''

- Convert linear ODE into a system of first order ODEs

- Define new variable $y_1(x)$

$$y_1(x) = y$$

- Define new variable $y_2(x)$

$$y_2(x) = y'$$

- Define new variable $y_3(x)$

$$y_3(x) = y''$$

- Isolate for $y_3'(x)$ using original ODE

$$y_3'(x) = 10 + 42 e^{3x} + 3y_3(x) - 2y_2(x)$$

Convert linear ODE into a system of first order ODEs

$$[y_2(x) = y_1'(x), y_3(x) = y_2'(x), y_3'(x) = 10 + 42 e^{3x} + 3y_3(x) - 2y_2(x)]$$

- Define vector

$$\vec{y}(x) = \begin{bmatrix} y_1(x) \\ y_2(x) \\ y_3(x) \end{bmatrix}$$

- System to solve

$$\vec{y}'(x) = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & -2 & 3 \end{bmatrix} \cdot \vec{y}(x) + \begin{bmatrix} 0 \\ 0 \\ 10 + 42 e^{3x} \end{bmatrix}$$

- Define the forcing function

$$\vec{f}(x) = \begin{bmatrix} 0 \\ 0 \\ 10 + 42 e^{3x} \end{bmatrix}$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & -2 & 3 \end{bmatrix}$$

- Rewrite the system as

$$\vec{y}'(x) = A \cdot \vec{y}(x) + \vec{f}$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[0, \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \right], \left[1, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right], \left[2, \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[0, \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

- Consider eigenpair

$$\left[1, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_2 = e^x \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[2, \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_3 = e^{2x} \cdot \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix}$$

- General solution of the system of ODEs can be written in terms of the particular solution $\vec{y}_p(x)$

$$\vec{y}(x) = c_1 \vec{y}_1 + c_2 \vec{y}_2 + c_3 \vec{y}_3 + \vec{y}_p(x)$$
- Fundamental matrix
 - Let $\phi(x)$ be the matrix whose columns are the independent solutions of the homogeneous system

$$\phi(x) = \begin{bmatrix} 1 & e^x & \frac{e^{2x}}{4} \\ 0 & e^x & \frac{e^{2x}}{2} \\ 0 & e^x & e^{2x} \end{bmatrix}$$

- The fundamental matrix, $\Phi(x)$ is a normalized version of $\phi(x)$ satisfying $\Phi(0) = I$ where I is the identity matrix

$$\Phi(x) = \phi(x) \cdot \frac{1}{\phi(0)}$$

- Substitute the value of $\phi(x)$ and $\phi(0)$

$$\Phi(x) = \begin{bmatrix} 1 & e^x & \frac{e^{2x}}{4} \\ 0 & e^x & \frac{e^{2x}}{2} \\ 0 & e^x & e^{2x} \end{bmatrix} \cdot \frac{1}{\begin{bmatrix} 1 & 1 & \frac{1}{4} \\ 0 & 1 & \frac{1}{2} \\ 0 & 1 & 1 \end{bmatrix}}$$

- Evaluate and simplify to get the fundamental matrix

$$\Phi(x) = \begin{bmatrix} 1 & -\frac{3}{2} + 2e^x - \frac{e^{2x}}{2} & \frac{1}{2} - e^x + \frac{e^{2x}}{2} \\ 0 & 2e^x - e^{2x} & -e^x + e^{2x} \\ 0 & 2e^x - 2e^{2x} & -e^x + 2e^{2x} \end{bmatrix}$$

- Find a particular solution of the system of ODEs using variation of parameters

- Let the particular solution be the fundamental matrix multiplied by $\vec{v}(x)$ and solve for $\vec{v}(x)$

$$\vec{y}_p(x) = \Phi(x) \cdot \vec{v}(x)$$

- Take the derivative of the particular solution

$$\vec{y}'_p(x) = \Phi'(x) \cdot \vec{v}(x) + \Phi(x) \cdot \vec{v}'(x)$$

- Substitute particular solution and its derivative into the system of ODEs

$$\Phi'(x) \cdot \vec{v}(x) + \Phi(x) \cdot \vec{v}'(x) = A \cdot \Phi(x) \cdot \vec{v}(x) + \vec{f}(x)$$

- The fundamental matrix has columns that are solutions to the homogeneous system so its derivative is $A \cdot \Phi(x)$

$$A \cdot \Phi(x) \cdot \vec{v}(x) + \Phi(x) \cdot \vec{v}'(x) = A \cdot \Phi(x) \cdot \vec{v}(x) + \vec{f}(x)$$

- Cancel like terms

$$\Phi(x) \cdot \vec{v}'(x) = \vec{f}(x)$$

- Multiply by the inverse of the fundamental matrix

$$\vec{v}'(x) = \frac{1}{\Phi(x)} \cdot \vec{f}(x)$$

- Integrate to solve for $\vec{v}(x)$

$$\vec{v}(x) = \int_0^x \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds$$

- Plug $\vec{v}(x)$ into the equation for the particular solution

$$\vec{y}_p(x) = \Phi(x) \cdot \left(\int_0^x \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds \right)$$

- Plug in the fundamental matrix and the forcing function and compute

$$\vec{y}_p(x) = \begin{bmatrix} 5x + 7e^{3x} + 11e^x + \frac{1}{2} - \frac{37e^{2x}}{2} \\ 21e^{3x} + 5 + 11e^x - 37e^{2x} \\ 63e^{3x} + 11e^x - 74e^{2x} \end{bmatrix}$$

- Plug particular solution back into general solution

$$\vec{y}(x) = c_1 \vec{y}_1 + c_2 \vec{y}_2 + c_3 \vec{y}_3 + \begin{bmatrix} 5x + 7e^{3x} + 11e^x + \frac{1}{2} - \frac{37e^{2x}}{2} \\ 21e^{3x} + 5 + 11e^x - 37e^{2x} \\ 63e^{3x} + 11e^x - 74e^{2x} \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y = \frac{1}{2} + \frac{(c_3 - 74)e^{2x}}{4} + 7e^{3x} + (11 + c_2)e^x + 5x + c_1$$

Maple trace

```

`Methods for third order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 3; linear nonhomogeneous with symmetry [0,1]
-> Calling odsolve with the ODE`, diff(diff(_b(_a), _a), _a) = 3*(diff(_b(_a), _a))-2*_b(_a)
Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
<- double symmetry of the form [xi=0, eta=F(x)] successful
<- differential order: 3; linear nonhomogeneous with symmetry [0,1] successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 26

```
dsolve(diff(y(x),x$3)-3*diff(y(x),x$2)+2*diff(y(x),x)=10+42*exp(3*x),y(x), singsol=all)
```

$$y(x) = \frac{e^{2x}c_1}{2} + e^x c_2 + 7e^{3x} + 5x + c_3$$

✓ Solution by Mathematica

Time used: 0.102 (sec). Leaf size: 35

```
DSolve[y'''[x]-3*y''[x]+2*y'[x]==10+42*Exp[3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 5x + 7e^{3x} + c_1e^x + \frac{1}{2}c_2e^{2x} + c_3$$

13.19 problem 18

13.19.1 Maple step by step solution 2260

Internal problem ID [6366]

Internal file name [OUTPUT/5614_Sunday_June_05_2022_03_44_52_PM_11593060/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Section 2.7. HIGHER ORDER LINEAR EQUATIONS, COUPLED HARMONIC OSCILLATORS Page 98

Problem number: 18.

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_3rd_order , _missing_x]]
```

$$y''' - y' = 1$$

With initial conditions

$$[y(0) = 4, y'(0) = 4, y''(0) = 4]$$

This is higher order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE And y_p is a particular solution to the nonhomogeneous ODE. y_h is the solution to

$$y''' - y' = 0$$

The characteristic equation is

$$\lambda^3 - \lambda = 0$$

The roots of the above equation are

$$\lambda_1 = 0$$

$$\lambda_2 = 1$$

$$\lambda_3 = -1$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 e^{-x} + c_2 + c_3 e^x$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = e^{-x}$$

$$y_2 = 1$$

$$y_3 = e^x$$

Now the particular solution to the given ODE is found

$$y''' - y' = 1$$

The particular solution is found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, e^x, e^{-x}\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-A_1 = 1$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = -1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^{-x} + c_2 + c_3 e^x) + (-x) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 e^{-x} + c_2 + c_3 e^x - x \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 4$ and $x = 0$ in the above gives

$$4 = c_1 + c_2 + c_3 \tag{1A}$$

Taking derivative of the solution gives

$$y' = -c_1 e^{-x} + c_3 e^x - 1$$

substituting $y' = 4$ and $x = 0$ in the above gives

$$4 = -c_1 + c_3 - 1 \tag{2A}$$

Taking two derivatives of the solution gives

$$y'' = c_1 e^{-x} + c_3 e^x$$

substituting $y'' = 4$ and $x = 0$ in the above gives

$$4 = c_1 + c_3 \tag{3A}$$

Equations {1A,2A,3A} are now solved for $\{c_1, c_2, c_3\}$. Solving for the constants gives

$$c_1 = -\frac{1}{2}$$

$$c_2 = 0$$

$$c_3 = \frac{9}{2}$$

Substituting these values back in above solution results in

$$y = -\frac{e^{-x}}{2} + \frac{9e^x}{2} - x$$

Summary

The solution(s) found are the following

$$y = -\frac{e^{-x}}{2} + \frac{9e^x}{2} - x \tag{1}$$




Figure 371: Solution plot

Verification of solutions

$$y = -\frac{e^{-x}}{2} + \frac{9e^x}{2} - x$$

Verified OK.

13.19.1 Maple step by step solution

Let's solve

$$\left[y''' - y' = 1, y(0) = 4, y'|_{\{x=0\}} = 4, y''|_{\{x=0\}} = 4 \right]$$

- Highest derivative means the order of the ODE is 3
 y'''

□ Convert linear ODE into a system of first order ODEs

- Define new variable $y_1(x)$

$$y_1(x) = y$$

- Define new variable $y_2(x)$

$$y_2(x) = y'$$

- Define new variable $y_3(x)$

$$y_3(x) = y''$$

- Isolate for $y'_3(x)$ using original ODE

$$y'_3(x) = 1 + y_2(x)$$

Convert linear ODE into a system of first order ODEs

$$[y_2(x) = y'_1(x), y_3(x) = y'_2(x), y'_3(x) = 1 + y_2(x)]$$

- Define vector

$$\vec{y}(x) = \begin{bmatrix} y_1(x) \\ y_2(x) \\ y_3(x) \end{bmatrix}$$

- System to solve

$$\vec{y}'(x) = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \cdot \vec{y}(x) + \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

- Define the forcing function

$$\vec{f}(x) = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$$

- Rewrite the system as

$$\vec{y}'(x) = A \cdot \vec{y}(x) + \vec{f}$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[-1, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right], \left[0, \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \right], \left[1, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[-1, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_1 = e^{-x} \cdot \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[0, \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_2 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

- Consider eigenpair

$$\left[1, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_3 = e^x \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

- General solution of the system of ODEs can be written in terms of the particular solution $\vec{y}_p(x)$

$$\vec{y}(x) = c_1 \vec{y}_1 + c_2 \vec{y}_2 + c_3 \vec{y}_3 + \vec{y}_p(x)$$

- Fundamental matrix

- Let $\phi(x)$ be the matrix whose columns are the independent solutions of the homogeneous system

$$\phi(x) = \begin{bmatrix} e^{-x} & 1 & e^x \\ -e^{-x} & 0 & e^x \\ e^{-x} & 0 & e^x \end{bmatrix}$$

- The fundamental matrix, $\Phi(x)$ is a normalized version of $\phi(x)$ satisfying $\Phi(0) = I$ where I is the identity matrix

$$\Phi(x) = \phi(x) \cdot \frac{1}{\phi(0)}$$

- Substitute the value of $\phi(x)$ and $\phi(0)$

$$\Phi(x) = \begin{bmatrix} e^{-x} & 1 & e^x \\ -e^{-x} & 0 & e^x \\ e^{-x} & 0 & e^x \end{bmatrix} \cdot \frac{1}{\begin{bmatrix} 1 & 1 & 1 \\ -1 & 0 & 1 \\ 1 & 0 & 1 \end{bmatrix}}$$

- Evaluate and simplify to get the fundamental matrix

$$\Phi(x) = \begin{bmatrix} 1 & -\frac{e^{-x}}{2} + \frac{e^x}{2} & \frac{e^{-x}}{2} - 1 + \frac{e^x}{2} \\ 0 & \frac{e^{-x}}{2} + \frac{e^x}{2} & -\frac{e^{-x}}{2} + \frac{e^x}{2} \\ 0 & -\frac{e^{-x}}{2} + \frac{e^x}{2} & \frac{e^{-x}}{2} + \frac{e^x}{2} \end{bmatrix}$$

- Find a particular solution of the system of ODEs using variation of parameters

- Let the particular solution be the fundamental matrix multiplied by $\vec{v}(x)$ and solve for $\vec{v}(x)$

$$\vec{y}_p(x) = \Phi(x) \cdot \vec{v}(x)$$

- Take the derivative of the particular solution

$$\vec{y}'_p(x) = \Phi'(x) \cdot \vec{v}(x) + \Phi(x) \cdot \vec{v}'(x)$$

- Substitute particular solution and its derivative into the system of ODEs

$$\Phi'(x) \cdot \vec{v}(x) + \Phi(x) \cdot \vec{v}'(x) = A \cdot \Phi(x) \cdot \vec{v}(x) + \vec{f}(x)$$

- The fundamental matrix has columns that are solutions to the homogeneous system so its derivative is

$$A \cdot \Phi(x) \cdot \vec{v}(x) + \Phi(x) \cdot \vec{v}'(x) = A \cdot \Phi(x) \cdot \vec{v}(x) + \vec{f}(x)$$

- Cancel like terms

$$\Phi(x) \cdot \vec{v}'(x) = \vec{f}(x)$$

- Multiply by the inverse of the fundamental matrix

$$\vec{v}'(x) = \frac{1}{\Phi(x)} \cdot \vec{f}(x)$$

- Integrate to solve for $\vec{v}(x)$

$$\vec{v}(x) = \int_0^x \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds$$

- Plug $\vec{v}(x)$ into the equation for the particular solution

$$\vec{y}_p(x) = \Phi(x) \cdot \left(\int_0^x \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds \right)$$

- Plug in the fundamental matrix and the forcing function and compute

$$\vec{y}_p(x) = \begin{bmatrix} -x - \frac{e^{-x}}{2} + \frac{e^x}{2} \\ \frac{e^{-x}}{2} - 1 + \frac{e^x}{2} \\ -\frac{e^{-x}}{2} + \frac{e^x}{2} \end{bmatrix}$$

- Plug particular solution back into general solution

$$\vec{y}(x) = c_1 \vec{y}_1 + c_2 \vec{y}_2 + c_3 \vec{y}_3 + \begin{bmatrix} -x - \frac{e^{-x}}{2} + \frac{e^x}{2} \\ \frac{e^{-x}}{2} - 1 + \frac{e^x}{2} \\ -\frac{e^{-x}}{2} + \frac{e^x}{2} \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y = c_1 e^{-x} + c_3 e^x - x - \frac{e^{-x}}{2} + \frac{e^x}{2} + c_2$$

- Use the initial condition $y(0) = 4$

$$4 = c_1 + c_2 + c_3$$

- Calculate the 1st derivative of the solution

$$y' = -c_1 e^{-x} + c_3 e^x + \frac{e^{-x}}{2} - 1 + \frac{e^x}{2}$$

- Use the initial condition $y' \Big|_{\{x=0\}} = 4$

$$4 = -c_1 + c_3$$

- Calculate the 2nd derivative of the solution

$$y'' = c_1 e^{-x} + c_3 e^x - \frac{e^{-x}}{2} + \frac{e^x}{2}$$

- Use the initial condition $y'' \Big|_{\{x=0\}} = 4$

$$4 = c_1 + c_3$$

- Solve for the unknown coefficients

$$\{c_1 = 0, c_2 = 0, c_3 = 4\}$$

- Solution to the IVP

$$y = -\frac{e^{-x}}{2} + \frac{9e^x}{2} - x$$

Maple trace

```

`Methods for third order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 3; linear nonhomogeneous with symmetry [0,1]
-> Calling odsolve with the ODE`, diff(diff(_b(_a), _a), _a) = _b(_a)+1, _b(_a)`
Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
    <- solving first the homogeneous part of the ODE successful
<- differential order: 3; linear nonhomogeneous with symmetry [0,1] successful`

```

*** Suble

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 18

```
dsolve([diff(y(x),x$3)-diff(y(x),x)=1,y(0) = 4, D(y)(0) = 4, (D@@2)(y)(0) = 4],y(x), singsol
```

$$y(x) = -\frac{e^{-x}}{2} + \frac{9e^x}{2} - x$$

✓ Solution by Mathematica

Time used: 0.029 (sec). Leaf size: 25

```
DSolve[{y'''[x]-y'[x]==1,{y[0]==4,y'[0]==4,y''[0]==4}},y[x],x,IncludeSingularSolutions -> Tr
```

$$y(x) \rightarrow -x - \frac{e^{-x}}{2} + \frac{9e^x}{2}$$

13.20 problem 19(a)

13.20.1 Maple step by step solution 2268

Internal problem ID [6367]

Internal file name [OUTPUT/5615_Sunday_June_05_2022_03_44_54_PM_29400279/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Section 2.7. HIGHER ORDER LINEAR EQUATIONS, COUPLED HARMONIC OSCILLATORS Page 98

Problem number: 19(a).

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_ODE_non_constant_coefficients_of_type_Euler**"

Maple gives the following as the ode type

```
[[_3rd_order , _missing_y]]
```

$$x^3 y''' + 3x^2 y'' = 0$$

This is Euler ODE of higher order. Let $y = x^\lambda$. Hence

$$\begin{aligned}y' &= \lambda x^{\lambda-1} \\y'' &= \lambda(\lambda-1) x^{\lambda-2} \\y''' &= \lambda(\lambda-1)(\lambda-2) x^{\lambda-3}\end{aligned}$$

Substituting these back into

$$x^3 y''' + 3x^2 y'' = 0$$

gives

$$3x^2 \lambda(\lambda-1) x^{\lambda-2} + x^3 \lambda(\lambda-1)(\lambda-2) x^{\lambda-3} = 0$$

Which simplifies to

$$3\lambda(\lambda-1) x^\lambda + \lambda(\lambda-1)(\lambda-2) x^\lambda = 0$$

And since $x^\lambda \neq 0$ then dividing through by x^λ , the above becomes

$$3\lambda(\lambda - 1) + \lambda(\lambda - 1)(\lambda - 2) = 0$$

Simplifying gives the characteristic equation as

$$\lambda^3 - \lambda = 0$$

Solving the above gives the following roots

$$\lambda_1 = 0$$

$$\lambda_2 = 1$$

$$\lambda_3 = -1$$

This table summarises the result

root	multiplicity	type of root
-1	1	real root
0	1	real root
1	1	real root

The solution is generated by going over the above table. For each real root λ of multiplicity one generates a $c_1 x^\lambda$ basis solution. Each real root of multiplicity two, generates $c_1 x^\lambda$ and $c_2 x^\lambda \ln(x)$ basis solutions. Each real root of multiplicity three, generates $c_1 x^\lambda$ and $c_2 x^\lambda \ln(x)$ and $c_3 x^\lambda \ln(x)^2$ basis solutions, and so on. Each complex root $\alpha \pm i\beta$ of multiplicity one generates $x^\alpha (c_1 \cos(\beta \ln(x)) + c_2 \sin(\beta \ln(x)))$ basis solutions. And each complex root $\alpha \pm i\beta$ of multiplicity two generates $\ln(x) x^\alpha (c_1 \cos(\beta \ln(x)) + c_2 \sin(\beta \ln(x)))$ basis solutions. And each complex root $\alpha \pm i\beta$ of multiplicity three generates $\ln(x)^2 x^\alpha (c_1 \cos(\beta \ln(x)) + c_2 \sin(\beta \ln(x)))$ basis solutions. And so on. Using the above show that the solution is

$$y = \frac{c_1}{x} + c_2 + c_3 x$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = \frac{1}{x}$$

$$y_2 = 1$$

$$y_3 = x$$

Summary

The solution(s) found are the following

$$y = \frac{c_1}{x} + c_2 + c_3 x \tag{1}$$

Verification of solutions

$$y = \frac{c_1}{x} + c_2 + c_3x$$

Verified OK.

13.20.1 Maple step by step solution

Let's solve

$$x^3y''' + 3x^2y'' = 0$$

- Highest derivative means the order of the ODE is 3

$$y'''$$

- Isolate 3rd derivative

$$y''' = -\frac{3y''}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y''' + \frac{3y''}{x} = 0$$

- Multiply by denominators of the ODE

$$y'''x + 3y'' = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x , using the chain rule

$$y' = \left(\frac{d}{dt}y(t)\right) t'(x)$$

- Compute derivative

$$y' = \frac{\frac{d}{dt}y(t)}{x}$$

- Calculate the 2nd derivative of y with respect to x , using the chain rule

$$y'' = \left(\frac{d^2}{dt^2}y(t)\right) t'(x)^2 + t''(x) \left(\frac{d}{dt}y(t)\right)$$

- Compute derivative

$$y'' = \frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2}$$

- Calculate the 3rd derivative of y with respect to x , using the chain rule

$$y''' = \left(\frac{d^3}{dt^3}y(t)\right) t'(x)^3 + 3t'(x)t''(x) \left(\frac{d^2}{dt^2}y(t)\right) + t'''(x) \left(\frac{d}{dt}y(t)\right)$$

- Compute derivative

$$y''' = \frac{\frac{d^3}{dt^3}y(t)}{x^3} - \frac{3\left(\frac{d^2}{dt^2}y(t)\right)}{x^3} + \frac{2\left(\frac{d}{dt}y(t)\right)}{x^3}$$

Substitute the change of variables back into the ODE

$$\left(\frac{\frac{d^3}{dt^3}y(t)}{x^3} - \frac{3\left(\frac{d^2}{dt^2}y(t)\right)}{x^3} + \frac{2\left(\frac{d}{dt}y(t)\right)}{x^3}\right)x + \frac{3\left(\frac{d^2}{dt^2}y(t)\right)}{x^2} - \frac{3\left(\frac{d}{dt}y(t)\right)}{x^2} = 0$$

- Simplify

$$\frac{\frac{d^3}{dt^3}y(t) - \frac{d}{dt}y(t)}{x^2} = 0$$

- Isolate 3rd derivative

$$\frac{d^3}{dt^3}y(t) = \frac{d}{dt}y(t)$$

- Group terms with $y(t)$ on the lhs of the ODE and the rest on the rhs of the ODE; ODE is lin

$$\frac{d^3}{dt^3}y(t) - \frac{d}{dt}y(t) = 0$$

- Convert linear ODE into a system of first order ODEs

- Define new variable $y_1(t)$

$$y_1(t) = y(t)$$

- Define new variable $y_2(t)$

$$y_2(t) = \frac{d}{dt}y(t)$$

- Define new variable $y_3(t)$

$$y_3(t) = \frac{d^2}{dt^2}y(t)$$

- Isolate for $\frac{d}{dt}y_3(t)$ using original ODE

$$\frac{d}{dt}y_3(t) = y_2(t)$$

Convert linear ODE into a system of first order ODEs

$$\left[y_2(t) = \frac{d}{dt}y_1(t), y_3(t) = \frac{d}{dt}y_2(t), \frac{d}{dt}y_3(t) = y_2(t)\right]$$

- Define vector

$$\vec{y}(t) = \begin{bmatrix} y_1(t) \\ y_2(t) \\ y_3(t) \end{bmatrix}$$

- System to solve

$$\frac{d}{dt} \vec{y}(t) = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \cdot \vec{y}(t)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$$

- Rewrite the system as

$$\frac{d}{dt} \vec{y}(t) = A \cdot \vec{y}(t)$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[-1, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right], \left[0, \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \right], \left[1, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[-1, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_1 = e^{-t} \cdot \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[0, \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_2 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

- Consider eigenpair

$$\left[1, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_3 = e^t \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{y} = c_1 \vec{y}_1 + c_2 \vec{y}_2 + c_3 \vec{y}_3$$

- Substitute solutions into the general solution

$$\vec{y} = c_1 e^{-t} \cdot \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} + c_3 e^t \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + \begin{bmatrix} c_2 \\ 0 \\ 0 \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y(t) = c_1 e^{-t} + c_3 e^t + c_2$$

- Change variables back using $t = \ln(x)$

$$y = \frac{c_1}{x} + c_2 + c_3 x$$

Maple trace

```

`Methods for third order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
<- LODE of Euler type successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 14

```
dsolve(x^3*diff(y(x),x$3)+3*x^2*diff(y(x),x$2)=0,y(x), singsol=all)
```

$$y(x) = c_1 + \frac{c_2}{x} + c_3x$$

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 21

```
DSolve[x^3*y'''[x]+3*x^2*y''[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_1}{2x} + c_3x + c_2$$

13.21 problem 19(b)

13.21.1 Maple step by step solution 2275

Internal problem ID [6368]

Internal file name [OUTPUT/5616_Sunday_June_05_2022_03_44_56_PM_29921558/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Section 2.7. HIGHER ORDER LINEAR EQUATIONS, COUPLED HARMONIC OSCILLATORS Page 98

Problem number: 19(b).

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_ODE_non_constant_coefficients_of_type_Euler**"

Maple gives the following as the ode type

```
[[_3rd_order , _exact , _linear , _homogeneous]]
```

$$x^3 y''' + x^2 y'' - 2xy' + 2y = 0$$

This is Euler ODE of higher order. Let $y = x^\lambda$. Hence

$$y' = \lambda x^{\lambda-1}$$

$$y'' = \lambda(\lambda - 1) x^{\lambda-2}$$

$$y''' = \lambda(\lambda - 1)(\lambda - 2) x^{\lambda-3}$$

Substituting these back into

$$x^3 y''' + x^2 y'' - 2xy' + 2y = 0$$

gives

$$-2x\lambda x^{\lambda-1} + x^2\lambda(\lambda - 1) x^{\lambda-2} + x^3\lambda(\lambda - 1)(\lambda - 2) x^{\lambda-3} + 2x^\lambda = 0$$

Which simplifies to

$$-2\lambda x^\lambda + \lambda(\lambda - 1) x^\lambda + \lambda(\lambda - 1)(\lambda - 2) x^\lambda + 2x^\lambda = 0$$

And since $x^\lambda \neq 0$ then dividing through by x^λ , the above becomes

$$-2\lambda + \lambda(\lambda - 1) + \lambda(\lambda - 1)(\lambda - 2) + 2 = 0$$

Simplifying gives the characteristic equation as

$$\lambda^3 - 2\lambda^2 - \lambda + 2 = 0$$

Solving the above gives the following roots

$$\lambda_1 = 1$$

$$\lambda_2 = 2$$

$$\lambda_3 = -1$$

This table summarises the result

root	multiplicity	type of root
-1	1	real root
1	1	real root
2	1	real root

The solution is generated by going over the above table. For each real root λ of multiplicity one generates a c_1x^λ basis solution. Each real root of multiplicity two, generates c_1x^λ and $c_2x^\lambda \ln(x)$ basis solutions. Each real root of multiplicity three, generates c_1x^λ and $c_2x^\lambda \ln(x)$ and $c_3x^\lambda \ln(x)^2$ basis solutions, and so on. Each complex root $\alpha \pm i\beta$ of multiplicity one generates $x^\alpha(c_1 \cos(\beta \ln(x)) + c_2 \sin(\beta \ln(x)))$ basis solutions. And each complex root $\alpha \pm i\beta$ of multiplicity two generates $\ln(x) x^\alpha(c_1 \cos(\beta \ln(x)) + c_2 \sin(\beta \ln(x)))$ basis solutions. And each complex root $\alpha \pm i\beta$ of multiplicity three generates $\ln(x)^2 x^\alpha(c_1 \cos(\beta \ln(x)) + c_2 \sin(\beta \ln(x)))$ basis solutions. And so on. Using the above show that the solution is

$$y = \frac{c_1}{x} + c_2x + c_3x^2$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = \frac{1}{x}$$

$$y_2 = x$$

$$y_3 = x^2$$

Summary

The solution(s) found are the following

$$y = \frac{c_1}{x} + c_2x + c_3x^2 \tag{1}$$

Verification of solutions

$$y = \frac{c_1}{x} + c_2x + c_3x^2$$

Verified OK.

13.21.1 Maple step by step solution

Let's solve

$$x^3y''' + x^2y'' - 2xy' + 2y = 0$$

- Highest derivative means the order of the ODE is 3

$$y'''$$

- Isolate 3rd derivative

$$y''' = -\frac{2y}{x^3} - \frac{y'x - 2y'}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y''' + \frac{y''}{x} - \frac{2y'}{x^2} + \frac{2y}{x^3} = 0$$

- Multiply by denominators of the ODE

$$x^3y''' + x^2y'' - 2xy' + 2y = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x , using the chain rule

$$y' = \left(\frac{d}{dt}y(t)\right) t'(x)$$

- Compute derivative

$$y' = \frac{\frac{d}{dt}y(t)}{x}$$

- Calculate the 2nd derivative of y with respect to x , using the chain rule

$$y'' = \left(\frac{d^2}{dt^2}y(t)\right) t'(x)^2 + t''(x) \left(\frac{d}{dt}y(t)\right)$$

- Compute derivative

$$y'' = \frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2}$$

- Calculate the 3rd derivative of y with respect to x , using the chain rule

$$y''' = \left(\frac{d^3}{dt^3}y(t)\right) t'(x)^3 + 3t'(x)t''(x) \left(\frac{d^2}{dt^2}y(t)\right) + t'''(x) \left(\frac{d}{dt}y(t)\right)$$

- Compute derivative

$$y''' = \frac{\frac{d^3}{dt^3}y(t)}{x^3} - \frac{3\left(\frac{d^2}{dt^2}y(t)\right)}{x^3} + \frac{2\left(\frac{d}{dt}y(t)\right)}{x^3}$$

Substitute the change of variables back into the ODE

$$x^3 \left(\frac{\frac{d^3}{dt^3}y(t)}{x^3} - \frac{3\left(\frac{d^2}{dt^2}y(t)\right)}{x^3} + \frac{2\left(\frac{d}{dt}y(t)\right)}{x^3} \right) + x^2 \left(\frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2} \right) - 2\frac{d}{dt}y(t) + 2y(t) = 0$$

- Simplify

$$\frac{d^3}{dt^3}y(t) - 2\frac{d^2}{dt^2}y(t) - \frac{d}{dt}y(t) + 2y(t) = 0$$

- Convert linear ODE into a system of first order ODEs

- Define new variable $y_1(t)$

$$y_1(t) = y(t)$$

- Define new variable $y_2(t)$

$$y_2(t) = \frac{d}{dt}y(t)$$

- Define new variable $y_3(t)$

$$y_3(t) = \frac{d^2}{dt^2}y(t)$$

- Isolate for $\frac{d}{dt}y_3(t)$ using original ODE

$$\frac{d}{dt}y_3(t) = 2y_3(t) + y_2(t) - 2y_1(t)$$

Convert linear ODE into a system of first order ODEs

$$\left[y_2(t) = \frac{d}{dt}y_1(t), y_3(t) = \frac{d}{dt}y_2(t), \frac{d}{dt}y_3(t) = 2y_3(t) + y_2(t) - 2y_1(t) \right]$$

- Define vector

$$\vec{y}(t) = \begin{bmatrix} y_1(t) \\ y_2(t) \\ y_3(t) \end{bmatrix}$$

- System to solve

$$\frac{d}{dt}\vec{y}(t) = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ -2 & 1 & 2 \end{bmatrix} \cdot \vec{y}(t)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ -2 & 1 & 2 \end{bmatrix}$$

- Rewrite the system as

$$\frac{d}{dt} \vec{y}(t) = A \cdot \vec{y}(t)$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[-1, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right], \left[1, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right], \left[2, \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[-1, \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_1 = e^{-t} \cdot \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[1, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_2 = e^t \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[2, \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_3 = e^{2t} \cdot \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{y} = c_1 \vec{y}_1 + c_2 \vec{y}_2 + c_3 \vec{y}_3$$

- Substitute solutions into the general solution

$$\vec{y} = c_1 e^{-t} \cdot \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} + c_2 e^t \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + c_3 e^{2t} \cdot \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y(t) = c_1 e^{-t} + c_2 e^t + \frac{c_3 e^{2t}}{4}$$

- Change variables back using $t = \ln(x)$

$$y = \frac{c_1}{x} + c_2 x + \frac{c_3 x^2}{4}$$

Maple trace

```

`Methods for third order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
<- LODE of Euler type successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 20

```
dsolve(x^3*diff(y(x),x$3)+x^2*diff(y(x),x$2)-2*x*diff(y(x),x)+2*y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{c_1 x^3 + c_3 x^2 + c_2}{x}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 22

```
DSolve[x^3*y'''[x]+x^2*y''[x]-2*x*y'[x]+2*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_3 x^2 + c_2 x + \frac{c_1}{x}$$

13.22 problem 19(c)

13.22.1 Maple step by step solution 2282

Internal problem ID [6369]

Internal file name [OUTPUT/5617_Sunday_June_05_2022_03_44_57_PM_49950363/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Section 2.7. HIGHER ORDER LINEAR EQUATIONS, COUPLED HARMONIC OSCILLATORS Page 98

Problem number: 19(c).

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : **"higher_order_ODE_non_constant_coefficients_of_type_Euler"**

Maple gives the following as the ode type

```
[[_3rd_order , _with_linear_symmetries]]
```

$$x^3y''' + 2x^2y'' + xy' - y = 0$$

This is Euler ODE of higher order. Let $y = x^\lambda$. Hence

$$y' = \lambda x^{\lambda-1}$$

$$y'' = \lambda(\lambda - 1) x^{\lambda-2}$$

$$y''' = \lambda(\lambda - 1)(\lambda - 2) x^{\lambda-3}$$

Substituting these back into

$$x^3y''' + 2x^2y'' + xy' - y = 0$$

gives

$$x\lambda x^{\lambda-1} + 2x^2\lambda(\lambda - 1) x^{\lambda-2} + x^3\lambda(\lambda - 1)(\lambda - 2) x^{\lambda-3} - x^\lambda = 0$$

Which simplifies to

$$\lambda x^\lambda + 2\lambda(\lambda - 1) x^\lambda + \lambda(\lambda - 1)(\lambda - 2) x^\lambda - x^\lambda = 0$$

And since $x^\lambda \neq 0$ then dividing through by x^λ , the above becomes

$$\lambda + 2\lambda(\lambda - 1) + \lambda(\lambda - 1)(\lambda - 2) - 1 = 0$$

Simplifying gives the characteristic equation as

$$\lambda^3 - \lambda^2 + \lambda - 1 = 0$$

Solving the above gives the following roots

$$\lambda_1 = 1$$

$$\lambda_2 = i$$

$$\lambda_3 = -i$$

This table summarises the result

root	multiplicity	type of root
1	1	real root
$\pm 1i$	1	complex conjugate root

The solution is generated by going over the above table. For each real root λ of multiplicity one generates a c_1x^λ basis solution. Each real root of multiplicity two, generates c_1x^λ and $c_2x^\lambda \ln(x)$ basis solutions. Each real root of multiplicity three, generates c_1x^λ and $c_2x^\lambda \ln(x)$ and $c_3x^\lambda \ln(x)^2$ basis solutions, and so on. Each complex root $\alpha \pm i\beta$ of multiplicity one generates $x^\alpha(c_1 \cos(\beta \ln(x)) + c_2 \sin(\beta \ln(x)))$ basis solutions. And each complex root $\alpha \pm i\beta$ of multiplicity two generates $\ln(x) x^\alpha(c_1 \cos(\beta \ln(x)) + c_2 \sin(\beta \ln(x)))$ basis solutions. And each complex root $\alpha \pm i\beta$ of multiplicity three generates $\ln(x)^2 x^\alpha(c_1 \cos(\beta \ln(x)) + c_2 \sin(\beta \ln(x)))$ basis solutions. And so on. Using the above show that the solution is

$$y = c_1x + c_2 \cos(\ln(x)) + c_3 \sin(\ln(x))$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = x$$

$$y_2 = \cos(\ln(x))$$

$$y_3 = \sin(\ln(x))$$

Summary

The solution(s) found are the following

$$y = c_1x + c_2 \cos(\ln(x)) + c_3 \sin(\ln(x)) \quad (1)$$

Verification of solutions

$$y = c_1 x + c_2 \cos(\ln(x)) + c_3 \sin(\ln(x))$$

Verified OK.

13.22.1 Maple step by step solution

Let's solve

$$x^3 y''' + 2x^2 y'' + xy' - y = 0$$

- Highest derivative means the order of the ODE is 3

$$y'''$$

- Isolate 3rd derivative

$$y''' = \frac{y}{x^3} - \frac{2y''x + y'}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y''' + \frac{2y''}{x} + \frac{y'}{x^2} - \frac{y}{x^3} = 0$$

- Multiply by denominators of the ODE

$$x^3 y''' + 2x^2 y'' + xy' - y = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x , using the chain rule

$$y' = \left(\frac{d}{dt}y(t)\right) t'(x)$$

- Compute derivative

$$y' = \frac{\frac{d}{dt}y(t)}{x}$$

- Calculate the 2nd derivative of y with respect to x , using the chain rule

$$y'' = \left(\frac{d^2}{dt^2}y(t)\right) t'(x)^2 + t''(x) \left(\frac{d}{dt}y(t)\right)$$

- Compute derivative

$$y'' = \frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2}$$

- Calculate the 3rd derivative of y with respect to x , using the chain rule

$$y''' = \left(\frac{d^3}{dt^3}y(t)\right) t'(x)^3 + 3t'(x)t''(x) \left(\frac{d^2}{dt^2}y(t)\right) + t'''(x) \left(\frac{d}{dt}y(t)\right)$$

- Compute derivative

$$y''' = \frac{\frac{d^3}{dt^3}y(t)}{x^3} - \frac{3\left(\frac{d^2}{dt^2}y(t)\right)}{x^3} + \frac{2\left(\frac{d}{dt}y(t)\right)}{x^3}$$

Substitute the change of variables back into the ODE

$$x^3 \left(\frac{\frac{d^3}{dt^3}y(t)}{x^3} - \frac{3\left(\frac{d^2}{dt^2}y(t)\right)}{x^3} + \frac{2\left(\frac{d}{dt}y(t)\right)}{x^3} \right) + 2x^2 \left(\frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2} \right) + \frac{d}{dt}y(t) - y(t) = 0$$

- Simplify

$$\frac{d^3}{dt^3}y(t) - \frac{d^2}{dt^2}y(t) + \frac{d}{dt}y(t) - y(t) = 0$$

- Convert linear ODE into a system of first order ODEs

- Define new variable $y_1(t)$

$$y_1(t) = y(t)$$

- Define new variable $y_2(t)$

$$y_2(t) = \frac{d}{dt}y(t)$$

- Define new variable $y_3(t)$

$$y_3(t) = \frac{d^2}{dt^2}y(t)$$

- Isolate for $\frac{d}{dt}y_3(t)$ using original ODE

$$\frac{d}{dt}y_3(t) = y_3(t) - y_2(t) + y_1(t)$$

Convert linear ODE into a system of first order ODEs

$$\left[\frac{d}{dt}y_1(t) = y_2(t), \frac{d}{dt}y_2(t) = y_3(t), \frac{d}{dt}y_3(t) = y_3(t) - y_2(t) + y_1(t) \right]$$

- Define vector

$$\vec{y}(t) = \begin{bmatrix} y_1(t) \\ y_2(t) \\ y_3(t) \end{bmatrix}$$

- System to solve

$$\frac{d}{dt}\vec{y}(t) = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & -1 & 1 \end{bmatrix} \cdot \vec{y}(t)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & -1 & 1 \end{bmatrix}$$

- Rewrite the system as

$$\frac{d}{dt}\vec{y}(t) = A \cdot \vec{y}(t)$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[1, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right], \left[-I, \begin{bmatrix} -1 \\ I \\ 1 \end{bmatrix} \right], \left[I, \begin{bmatrix} -1 \\ -I \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[1, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_1 = e^t \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$\left[-I, \begin{bmatrix} -1 \\ I \\ 1 \end{bmatrix} \right]$$

- Solution from eigenpair

$$e^{-It} \cdot \begin{bmatrix} -1 \\ I \\ 1 \end{bmatrix}$$

- Use Euler identity to write solution in terms of sin and cos

$$(\cos(t) - I \sin(t)) \cdot \begin{bmatrix} -1 \\ I \\ 1 \end{bmatrix}$$

- Simplify expression

$$\begin{bmatrix} -\cos(t) + I \sin(t) \\ I(\cos(t) - I \sin(t)) \\ \cos(t) - I \sin(t) \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\left[\vec{y}_2(t) = \begin{bmatrix} -\cos(t) \\ \sin(t) \\ \cos(t) \end{bmatrix}, \vec{y}_3(t) = \begin{bmatrix} \sin(t) \\ \cos(t) \\ -\sin(t) \end{bmatrix} \right]$$

- General solution to the system of ODEs

$$\vec{y} = c_1 \vec{y}_1 + c_2 \vec{y}_2(t) + c_3 \vec{y}_3(t)$$

- Substitute solutions into the general solution

$$\vec{y} = c_1 e^t \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + \begin{bmatrix} -c_2 \cos(t) + c_3 \sin(t) \\ c_2 \sin(t) + c_3 \cos(t) \\ c_2 \cos(t) - c_3 \sin(t) \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y(t) = c_1 e^t + c_3 \sin(t) - c_2 \cos(t)$$

- Change variables back using $t = \ln(x)$

$$y = c_1 x + c_3 \sin(\ln(x)) - c_2 \cos(\ln(x))$$

Maple trace

```

`Methods for third order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
<- LODE of Euler type successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 18

```
dsolve(x^3*diff(y(x),x$3)+2*x^2*diff(y(x),x$2)+x*diff(y(x),x)-y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 x + \sin(\ln(x)) c_2 + c_3 \cos(\ln(x))$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 22

```
DSolve[x^3*y'''[x]+2*x^2*y''[x]+x*y'[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_3 x + c_1 \cos(\log(x)) + c_2 \sin(\log(x))$$

13.23 problem 20

13.23.1 Maple step by step solution 2289

Internal problem ID [6370]

Internal file name [OUTPUT/5618_Sunday_June_05_2022_03_44_58_PM_22166040/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Section 2.7. HIGHER ORDER LINEAR EQUATIONS, COUPLED HARMONIC OSCILLATORS Page 98

Problem number: 20.

ODE order: 4.

ODE degree: 1.

The type(s) of ODE detected by this program : **"higher_order_missing_y"**

Maple gives the following as the ode type

[[_high_order , _missing_y]]

$$x^3 y'''' + 8x^2 y''' + 8xy'' - 8y' = 0$$

Since y is missing from the ode then we can use the substitution $y' = v(x)$ to reduce the order by one. The ODE becomes

$$x^3 v'''(x) + 8x^2 v''(x) + 8v'(x)x - 8v(x) = 0$$

This is Euler ODE of higher order. Let $v(x) = x^\lambda$. Hence

$$\begin{aligned}v'(x) &= \lambda x^{\lambda-1} \\v''(x) &= \lambda(\lambda-1) x^{\lambda-2} \\v'''(x) &= \lambda(\lambda-1)(\lambda-2) x^{\lambda-3}\end{aligned}$$

Substituting these back into

$$x^3 v'''(x) + 8x^2 v''(x) + 8v'(x)x - 8v(x) = 0$$

gives

$$8x\lambda x^{\lambda-1} + 8x^2\lambda(\lambda-1) x^{\lambda-2} + x^3\lambda(\lambda-1)(\lambda-2) x^{\lambda-3} - 8x^\lambda = 0$$

Which simplifies to

$$8\lambda x^\lambda + 8\lambda(\lambda - 1)x^\lambda + \lambda(\lambda - 1)(\lambda - 2)x^\lambda - 8x^\lambda = 0$$

And since $x^\lambda \neq 0$ then dividing through by x^λ , the above becomes

$$8\lambda + 8\lambda(\lambda - 1) + \lambda(\lambda - 1)(\lambda - 2) - 8 = 0$$

Simplifying gives the characteristic equation as

$$\lambda^3 + 5\lambda^2 + 2\lambda - 8 = 0$$

Solving the above gives the following roots

$$\lambda_1 = 1$$

$$\lambda_2 = -4$$

$$\lambda_3 = -2$$

This table summarises the result

root	multiplicity	type of root
-2	1	real root
1	1	real root
-4	1	real root

The solution is generated by going over the above table. For each real root λ of multiplicity one generates a c_1x^λ basis solution. Each real root of multiplicity two, generates c_1x^λ and $c_2x^\lambda \ln(x)$ basis solutions. Each real root of multiplicity three, generates c_1x^λ and $c_2x^\lambda \ln(x)$ and $c_3x^\lambda \ln(x)^2$ basis solutions, and so on. Each complex root $\alpha \pm i\beta$ of multiplicity one generates $x^\alpha(c_1 \cos(\beta \ln(x)) + c_2 \sin(\beta \ln(x)))$ basis solutions. And each complex root $\alpha \pm i\beta$ of multiplicity two generates $\ln(x) x^\alpha(c_1 \cos(\beta \ln(x)) + c_2 \sin(\beta \ln(x)))$ basis solutions. And each complex root $\alpha \pm i\beta$ of multiplicity three generates $\ln(x)^2 x^\alpha(c_1 \cos(\beta \ln(x)) + c_2 \sin(\beta \ln(x)))$ basis solutions. And so on. Using the above show that the solution is

$$v(x) = \frac{c_1}{x^2} + c_2x + \frac{c_3}{x^4}$$

The fundamental set of solutions for the homogeneous solution are the following

$$v_1 = \frac{1}{x^2}$$

$$v_2 = x$$

$$v_3 = \frac{1}{x^4}$$

But since $y' = v(x)$ then we now need to solve the ode $y' = \frac{c_1}{x^2} + c_2x + \frac{c_3}{x^4}$. Integrating both sides gives

$$y = \int \frac{c_2x^5 + c_1x^2 + c_3}{x^4} dx$$

$$= \frac{c_2x^2}{2} - \frac{c_3}{3x^3} - \frac{c_1}{x} + c_1$$

Summary

The solution(s) found are the following

$$y = \frac{c_2x^2}{2} - \frac{c_3}{3x^3} - \frac{c_1}{x} + c_1 \quad (1)$$

Verification of solutions

$$y = \frac{c_2x^2}{2} - \frac{c_3}{3x^3} - \frac{c_1}{x} + c_1$$

Verified OK.

13.23.1 Maple step by step solution

Let's solve

$$x^3y'''' + 8x^2y''' + 8y''x - 8y' = 0$$

- Highest derivative means the order of the ODE is 4

$$y''''$$

- Isolate 4th derivative

$$y'''' = -\frac{8(x^2y''' + y''x - y')}{x^3}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'''' + \frac{8y'''}{x} + \frac{8y''}{x^2} - \frac{8y'}{x^3} = 0$$

- Multiply by denominators of the ODE

$$x^3y'''' + 8x^2y''' + 8y''x - 8y' = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x , using the chain rule

$$y' = \left(\frac{d}{dt}y(t)\right) t'(x)$$

- Compute derivative

$$y' = \frac{\frac{d}{dt}y(t)}{x}$$

- Calculate the 2nd derivative of y with respect to x , using the chain rule

$$y'' = \left(\frac{d^2}{dt^2}y(t) \right) t'(x)^2 + t''(x) \left(\frac{d}{dt}y(t) \right)$$

- Compute derivative

$$y'' = \frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2}$$

- Calculate the 3rd derivative of y with respect to x , using the chain rule

$$y''' = \left(\frac{d^3}{dt^3}y(t) \right) t'(x)^3 + 3t''(x) t'(x) \left(\frac{d^2}{dt^2}y(t) \right) + t'''(x) \left(\frac{d}{dt}y(t) \right)$$

- Compute derivative

$$y''' = \frac{\frac{d^3}{dt^3}y(t)}{x^3} - \frac{3\left(\frac{d^2}{dt^2}y(t)\right)}{x^3} + \frac{2\left(\frac{d}{dt}y(t)\right)}{x^3}$$

- Calculate the 4th derivative of y with respect to x , using the chain rule

$$y'''' = \left(\frac{d^4}{dt^4}y(t) \right) t'(x)^4 + 3t''(x)^2 t'(x) \left(\frac{d^3}{dt^3}y(t) \right) + 3t''(x)^2 \left(\frac{d^2}{dt^2}y(t) \right) + 3\left(t'''(x) \left(\frac{d^2}{dt^2}y(t) \right) \right) + \left(\frac{d^3}{dt^3}y(t) \right)$$

- Compute derivative

$$y'''' = \frac{\frac{d^4}{dt^4}y(t)}{x^4} - \frac{3\left(\frac{d^3}{dt^3}y(t)\right)}{x^4} + \frac{5\left(\frac{d^2}{dt^2}y(t)\right)}{x^4} + \frac{3\left(\frac{2\left(\frac{d^2}{dt^2}y(t)\right)}{x^3} - \frac{\frac{d^3}{dt^3}y(t)}{x^3}\right)}{x} - \frac{6\left(\frac{d}{dt}y(t)\right)}{x^4}$$

Substitute the change of variables back into the ODE

$$x^3 \left(\frac{\frac{d^4}{dt^4}y(t)}{x^4} - \frac{3\left(\frac{d^3}{dt^3}y(t)\right)}{x^4} + \frac{5\left(\frac{d^2}{dt^2}y(t)\right)}{x^4} + \frac{3\left(\frac{2\left(\frac{d^2}{dt^2}y(t)\right)}{x^3} - \frac{\frac{d^3}{dt^3}y(t)}{x^3}\right)}{x} - \frac{6\left(\frac{d}{dt}y(t)\right)}{x^4} \right) + 8x^2 \left(\frac{\frac{d^3}{dt^3}y(t)}{x^3} - \frac{3\left(\frac{d^2}{dt^2}y(t)\right)}{x^3} \right)$$

- Simplify

$$\frac{\frac{d^4}{dt^4}y(t) + 2\frac{d^3}{dt^3}y(t) - 5\frac{d^2}{dt^2}y(t) - 6\frac{d}{dt}y(t)}{x} = 0$$

- Isolate 4th derivative

$$\frac{d^4}{dt^4}y(t) = -2\frac{d^3}{dt^3}y(t) + 5\frac{d^2}{dt^2}y(t) + 6\frac{d}{dt}y(t)$$

- Group terms with y(t) on the lhs of the ODE and the rest on the rhs of the ODE; ODE is lin

$$\frac{d^4}{dt^4}y(t) + 2\frac{d^3}{dt^3}y(t) - 5\frac{d^2}{dt^2}y(t) - 6\frac{d}{dt}y(t) = 0$$

- Convert linear ODE into a system of first order ODEs

- Define new variable y₁(t)

$$y_1(t) = y(t)$$

- Define new variable $y_2(t)$

$$y_2(t) = \frac{d}{dt}y(t)$$

- Define new variable $y_3(t)$

$$y_3(t) = \frac{d^2}{dt^2}y(t)$$

- Define new variable $y_4(t)$

$$y_4(t) = \frac{d^3}{dt^3}y(t)$$

- Isolate for $\frac{d}{dt}y_4(t)$ using original ODE

$$\frac{d}{dt}y_4(t) = -2y_4(t) + 5y_3(t) + 6y_2(t)$$

Convert linear ODE into a system of first order ODEs

$$\left[y_2(t) = \frac{d}{dt}y_1(t), y_3(t) = \frac{d}{dt}y_2(t), y_4(t) = \frac{d}{dt}y_3(t), \frac{d}{dt}y_4(t) = -2y_4(t) + 5y_3(t) + 6y_2(t) \right]$$

- Define vector

$$\vec{y}(t) = \begin{bmatrix} y_1(t) \\ y_2(t) \\ y_3(t) \\ y_4(t) \end{bmatrix}$$

- System to solve

$$\frac{d}{dt}\vec{y}(t) = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 6 & 5 & -2 \end{bmatrix} \cdot \vec{y}(t)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 6 & 5 & -2 \end{bmatrix}$$

- Rewrite the system as

$$\frac{d}{dt}\vec{y}(t) = A \cdot \vec{y}(t)$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[\begin{array}{c} -3, \\ \left[\begin{array}{c} -\frac{1}{27} \\ \frac{1}{9} \\ -\frac{1}{3} \\ 1 \end{array} \right] \end{array} \right], \left[-1, \left[\begin{array}{c} -1 \\ 1 \\ -1 \\ 1 \end{array} \right] \right], \left[0, \left[\begin{array}{c} 1 \\ 0 \\ 0 \\ 0 \end{array} \right] \right], \left[2, \left[\begin{array}{c} \frac{1}{8} \\ \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{array} \right] \right] \right]$$

- Consider eigenpair

$$\left[\begin{array}{c} -3, \\ \left[\begin{array}{c} -\frac{1}{27} \\ \frac{1}{9} \\ -\frac{1}{3} \\ 1 \end{array} \right] \end{array} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_1 = e^{-3t} \cdot \left[\begin{array}{c} -\frac{1}{27} \\ \frac{1}{9} \\ -\frac{1}{3} \\ 1 \end{array} \right]$$

- Consider eigenpair

$$\left[\begin{array}{c} -1, \\ \left[\begin{array}{c} -1 \\ 1 \\ -1 \\ 1 \end{array} \right] \end{array} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_2 = e^{-t} \cdot \left[\begin{array}{c} -1 \\ 1 \\ -1 \\ 1 \end{array} \right]$$

- Consider eigenpair

$$\left[0, \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_3 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

- Consider eigenpair

$$\left[2, \begin{bmatrix} \frac{1}{8} \\ \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_4 = e^{2t} \cdot \begin{bmatrix} \frac{1}{8} \\ \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{y} = c_1 \vec{y}_1 + c_2 \vec{y}_2 + c_3 \vec{y}_3 + c_4 \vec{y}_4$$

- Substitute solutions into the general solution

$$\vec{y} = c_1 e^{-3t} \cdot \begin{bmatrix} -\frac{1}{27} \\ \frac{1}{9} \\ -\frac{1}{3} \\ 1 \end{bmatrix} + c_2 e^{-t} \cdot \begin{bmatrix} -1 \\ 1 \\ -1 \\ 1 \end{bmatrix} + c_4 e^{2t} \cdot \begin{bmatrix} \frac{1}{8} \\ \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} + \begin{bmatrix} c_3 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y(t) = -\frac{\left(27c_2e^{2t} - 27c_3e^{3t} - \frac{27c_4e^{5t}}{8} + c_1\right)e^{-3t}}{27}$$

- Change variables back using $t = \ln(x)$

$$y = -\frac{27c_2x^2 - 27c_3x^3 - \frac{27}{8}c_4x^5 + c_1}{27x^3}$$

- Simplify

$$y = \frac{c_4x^2}{8} + c_3 - \frac{c_2}{x} - \frac{c_1}{27x^3}$$

Maple trace

```

`Methods for high order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
<- LODE of Euler type successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(x^3*diff(y(x),x$4)+8*x^2*diff(y(x),x$3)+8*x*diff(y(x),x$2)-8*diff(y(x),x)=0,y(x), sin
```

$$y(x) = c_1 + c_2x^2 + \frac{c_3}{x} + \frac{c_4}{x^3}$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 33

```
DSolve[x^3*y''''[x]+8*x^2*y'''[x]+8*x*y''[x]-8*y'[x]==0,y[x],x,IncludeSingularSolutions -> T
```

$$y(x) \rightarrow -\frac{c_1}{3x^3} + \frac{c_3x^2}{2} - \frac{c_2}{x} + c_4$$

14 Chapter 2. Problems for Review and Discovery. Drill excercises. Page 105

14.1 problem 1(a)	2296
14.2 problem 1(b)	2304
14.3 problem 1(c)	2313
14.4 problem 1(d)	2322
14.5 problem 1(e)	2331
14.6 problem 1(f)	2342
14.7 problem 1(g)	2353
14.8 problem 1(h)	2365
14.9 problem 2(a)	2376
14.10problem 2(b)	2389
14.11problem 2(c)	2404
14.12problem 2(d)	2418
14.13problem 2(e)	2433
14.14problem 2(f)	2447
14.15problem 2(g)	2460
14.16problem 2(h)	2481
14.17problem 3(a)	2509
14.18problem 3(b)	2519
14.19problem 3(c)	2530
14.20problem 3(d)	2541
14.21problem 3(e)	2553
14.22problem 3(f)	2568
14.23problem 3(g)	2584
14.24problem 3(h)	2618
14.25problem 4(a)	2632
14.26problem 4(b)	2639
14.27problem 4(c)	2646
14.28problem 4(d)	2654

14.1 problem 1(a)

- 14.1.1 Solving as second order linear constant coeff ode 2296
- 14.1.2 Solving using Kovacic algorithm 2298
- 14.1.3 Maple step by step solution 2302

Internal problem ID [6371]

Internal file name [OUTPUT/5619_Sunday_June_05_2022_03_44_59_PM_18712078/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Problems for Review and Discovery. Drill excercises. Page 105

Problem number: 1(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"kovacic", "second_order_linear_constant_coeff"**

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' - 3y' + y = 0$$

14.1.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -3, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 3\lambda e^{\lambda x} + e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 3\lambda + 1 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -3, C = 1$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{3}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-3^2 - (4)(1)(1)} \\ &= \frac{3}{2} \pm \frac{\sqrt{5}}{2}\end{aligned}$$

Hence

$$\lambda_1 = \frac{3}{2} + \frac{\sqrt{5}}{2}$$

$$\lambda_2 = \frac{3}{2} - \frac{\sqrt{5}}{2}$$

Which simplifies to

$$\lambda_1 = \frac{3}{2} + \frac{\sqrt{5}}{2}$$

$$\lambda_2 = \frac{3}{2} - \frac{\sqrt{5}}{2}$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{\left(\frac{3}{2} + \frac{\sqrt{5}}{2}\right)x} + c_2 e^{\left(\frac{3}{2} - \frac{\sqrt{5}}{2}\right)x}$$

Or

$$y = c_1 e^{\left(\frac{3}{2} + \frac{\sqrt{5}}{2}\right)x} + c_2 e^{\left(\frac{3}{2} - \frac{\sqrt{5}}{2}\right)x}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{\left(\frac{3}{2} + \frac{\sqrt{5}}{2}\right)x} + c_2 e^{\left(\frac{3}{2} - \frac{\sqrt{5}}{2}\right)x} \quad (1)$$




Figure 372: Slope field plot

Verification of solutions

$$y = c_1 e^{\left(\frac{3}{2} + \frac{\sqrt{5}}{2}\right)x} + c_2 e^{\left(\frac{3}{2} - \frac{\sqrt{5}}{2}\right)x}$$

Verified OK.

14.1.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 3y' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -3 \\ C &= 1 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{5}{4} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 5 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \frac{5z(x)}{4} \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 332: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{5}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{x\sqrt{5}}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-3}{1} dx} \\ &= z_1 e^{\frac{3x}{2}} \\ &= z_1 \left(e^{\frac{3x}{2}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-\frac{(\sqrt{5}-3)x}{2}}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-3}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{3x}}{(y_1)^2} dx \\ &= y_1 \left(\frac{\sqrt{5} e^{x\sqrt{5}}}{5} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}
 y &= c_1 y_1 + c_2 y_2 \\
 &= c_1 \left(e^{-\frac{(\sqrt{5}-3)x}{2}} \right) + c_2 \left(e^{-\frac{(\sqrt{5}-3)x}{2}} \left(\frac{\sqrt{5} e^{x\sqrt{5}}}{5} \right) \right)
 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-\frac{(\sqrt{5}-3)x}{2}} + \frac{c_2 \sqrt{5} e^{\frac{(3+\sqrt{5})x}{2}}}{5} \tag{1}$$




Figure 373: Slope field plot

Verification of solutions

$$y = c_1 e^{-\frac{(\sqrt{5}-3)x}{2}} + \frac{c_2 \sqrt{5} e^{\frac{(3+\sqrt{5})x}{2}}}{5}$$

Verified OK.

14.1.3 Maple step by step solution

Let's solve

$$y'' - 3y' + y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 - 3r + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{3 \pm (\sqrt{5})}{2}$$

- Roots of the characteristic polynomial

$$r = \left(\frac{3}{2} - \frac{\sqrt{5}}{2}, \frac{3}{2} + \frac{\sqrt{5}}{2} \right)$$

- 1st solution of the ODE

$$y_1(x) = e^{\left(\frac{3}{2} - \frac{\sqrt{5}}{2}\right)x}$$

- 2nd solution of the ODE

$$y_2(x) = e^{\left(\frac{3}{2} + \frac{\sqrt{5}}{2}\right)x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = c_1 e^{\left(\frac{3}{2} - \frac{\sqrt{5}}{2}\right)x} + c_2 e^{\left(\frac{3}{2} + \frac{\sqrt{5}}{2}\right)x}$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve(diff(y(x),x$2)-3*diff(y(x),x)+y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{\frac{(\sqrt{5}+3)x}{2}} + c_2 e^{-\frac{(\sqrt{5}-3)x}{2}}$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 35

```
DSolve[y''[x]-3*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-\frac{1}{2}(\sqrt{5}-3)x} (c_2 e^{\sqrt{5}x} + c_1)$$

14.2 problem 1(b)

14.2.1 Solving as second order linear constant coeff ode	2304
14.2.2 Solving using Kovacic algorithm	2306
14.2.3 Maple step by step solution	2310

Internal problem ID [6372]

Internal file name [OUTPUT/5620_Sunday_June_05_2022_03_45_01_PM_40855532/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Problems for Review and Discovery. Drill excercises. Page 105

Problem number: 1(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"kovacic", "second_order_linear_constant_coeff"**

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' + y' + y = 0$$

14.2.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 1, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + \lambda e^{\lambda x} + e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + \lambda + 1 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 1, C = 1$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{-1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{1^2 - (4)(1)(1)} \\ &= -\frac{1}{2} \pm \frac{i\sqrt{3}}{2}\end{aligned}$$

Hence

$$\begin{aligned}\lambda_1 &= -\frac{1}{2} + \frac{i\sqrt{3}}{2} \\ \lambda_2 &= -\frac{1}{2} - \frac{i\sqrt{3}}{2}\end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= -\frac{1}{2} + \frac{i\sqrt{3}}{2} \\ \lambda_2 &= -\frac{1}{2} - \frac{i\sqrt{3}}{2}\end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = -\frac{1}{2}$ and $\beta = \frac{\sqrt{3}}{2}$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

Summary

The solution(s) found are the following

$$y = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right) \quad (1)$$




Figure 374: Slope field plot

Verification of solutions

$$y = e^{-\frac{\sqrt{3}x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

Verified OK.

14.2.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 1 \tag{3}$$

$$C = 1$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-3}{4} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -3 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -\frac{3z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 334: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0\end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -\frac{3}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos\left(\frac{\sqrt{3}x}{2}\right)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{1}{1} dx} \\ &= z_1 e^{-\frac{x}{2}} \\ &= z_1 \left(e^{-\frac{x}{2}}\right)\end{aligned}$$

Which simplifies to

$$y_1 = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{1}{x} dx}}{(y_1)^2} dx \\&= y_1 \int \frac{e^{-x}}{(y_1)^2} dx \\&= y_1 \left(\frac{2\sqrt{3} \tan\left(\frac{\sqrt{3}x}{2}\right)}{3} \right)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\&= c_1 \left(e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) \right) + c_2 \left(e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) \left(\frac{2\sqrt{3} \tan\left(\frac{\sqrt{3}x}{2}\right)}{3} \right) \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) c_1 + \frac{2c_2 e^{-\frac{x}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3} \quad (1)$$




Figure 375: Slope field plot

Verification of solutions

$$y = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) c_1 + \frac{2c_2 e^{-\frac{x}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3}$$

Verified OK.

14.2.3 Maple step by step solution

Let's solve

$$y'' + y' + y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 + r + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-1) \pm (\sqrt{-3})}{2}$$

- Roots of the characteristic polynomial

$$r = \left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}, -\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)$$
- 1st solution of the ODE

$$y_1(x) = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)$$
- 2nd solution of the ODE

$$y_2(x) = e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)$$
- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$
- Substitute in solutions

$$y = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) c_1 + e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) c_2$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 28

```
dsolve(diff(y(x),x$2)+diff(y(x),x)+y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{-\frac{x}{2}} \left(c_1 \sin\left(\frac{\sqrt{3}x}{2}\right) + c_2 \cos\left(\frac{\sqrt{3}x}{2}\right) \right)$$

✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 42

```
DSolve[y''[x]+y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x/2} \left(c_2 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_1 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

14.3 problem 1(c)

14.3.1 Solving as second order linear constant coeff ode	2313
14.3.2 Solving as linear second order ode solved by an integrating factor ode	2315
14.3.3 Solving using Kovacic algorithm	2316
14.3.4 Maple step by step solution	2320

Internal problem ID [6373]

Internal file name [OUTPUT/5621_Sunday_June_05_2022_03_45_02_PM_44670485/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Problems for Review and Discovery. Drill excercises. Page 105

Problem number: 1(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "linear_second_order_ode_solved_by_an_integrating_factor"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' + 6y' + 9y = 0$$

14.3.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 6, C = 9$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 6\lambda e^{\lambda x} + 9 e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 6\lambda + 9 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 6, C = 9$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-6}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{(6)^2 - (4)(1)(9)} \\ &= -3 \end{aligned}$$

Hence this is the case of a double root $\lambda_{1,2} = 3$. Therefore the solution is

$$y = c_1 e^{-3x} + c_2 x e^{-3x} \tag{1}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-3x} + x e^{-3x} c_2 \tag{1}$$




Figure 376: Slope field plot

Verification of solutions

$$y = c_1 e^{-3x} + x e^{-3x} c_2$$

Verified OK.

14.3.2 Solving as linear second order ode solved by an integrating factor ode

The ode satisfies this form

$$y'' + p(x)y' + \frac{(p(x)^2 + p'(x))y}{2} = f(x)$$

Where $p(x) = 6$. Therefore, there is an integrating factor given by

$$\begin{aligned}M(x) &= e^{\frac{1}{2} \int p dx} \\ &= e^{\int 6 dx} \\ &= e^{3x}\end{aligned}$$

Multiplying both sides of the ODE by the integrating factor $M(x)$ makes the left side of the ODE a complete differential

$$\begin{aligned}(M(x)y)'' &= 0 \\ (e^{3x}y)'' &= 0\end{aligned}$$

Integrating once gives

$$(e^{3x}y)' = c_1$$

Integrating again gives

$$(e^{3x}y) = c_1x + c_2$$

Hence the solution is

$$y = \frac{c_1x + c_2}{e^{3x}}$$

Or

$$y = c_1x e^{-3x} + c_2e^{-3x}$$

Summary

The solution(s) found are the following

$$y = c_1x e^{-3x} + c_2e^{-3x} \tag{1}$$




Figure 377: Slope field plot

Verification of solutions

$$y = c_1 x e^{-3x} + c_2 e^{-3x}$$

Verified OK.

14.3.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 6y' + 9y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 6 \\ C &= 9 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 0 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 0 \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 336: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty\end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{6}{1} dx} \\ &= z_1 e^{-3x} \\ &= z_1 (e^{-3x})\end{aligned}$$

Which simplifies to

$$y_1 = e^{-3x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{6}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-6x}}{(y_1)^2} dx \\ &= y_1(x)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-3x}) + c_2 (e^{-3x}(x))\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-3x} + x e^{-3x} c_2 \quad (1)$$




Figure 378: Slope field plot

Verification of solutions

$$y = c_1 e^{-3x} + x e^{-3x} c_2$$

Verified OK.

14.3.4 Maple step by step solution

Let's solve

$$y'' + 6y' + 9y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 + 6r + 9 = 0$$

- Factor the characteristic polynomial

$$(r + 3)^2 = 0$$

- Root of the characteristic polynomial

$$r = -3$$

- 1st solution of the ODE

$$y_1(x) = e^{-3x}$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = x e^{-3x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = c_1 e^{-3x} + x e^{-3x} c_2$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(diff(y(x),x$2)+6*diff(y(x),x)+9*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{-3x}(c_2x + c_1)$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 18

```
DSolve[y''[x]+6*y'[x]+9*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-3x}(c_2x + c_1)$$

14.4 problem 1(d)

- 14.4.1 Solving as second order linear constant coeff ode 2322
- 14.4.2 Solving using Kovacic algorithm 2324
- 14.4.3 Maple step by step solution 2328

Internal problem ID [6374]

Internal file name [OUTPUT/5622_Sunday_June_05_2022_03_45_03_PM_54715595/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Problems for Review and Discovery. Drill excercises. Page 105

Problem number: 1(d).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"kovacic", "second_order_linear_constant_coeff"**

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' - y' + 6y = 0$$

14.4.1 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -1, C = 6$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - \lambda e^{\lambda x} + 6 e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - \lambda + 6 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -1, C = 6$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-1^2 - (4)(1)(6)} \\ &= \frac{1}{2} \pm \frac{i\sqrt{23}}{2}\end{aligned}$$

Hence

$$\begin{aligned}\lambda_1 &= \frac{1}{2} + \frac{i\sqrt{23}}{2} \\ \lambda_2 &= \frac{1}{2} - \frac{i\sqrt{23}}{2}\end{aligned}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= \frac{1}{2} + \frac{i\sqrt{23}}{2} \\ \lambda_2 &= \frac{1}{2} - \frac{i\sqrt{23}}{2}\end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = \frac{1}{2}$ and $\beta = \frac{\sqrt{23}}{2}$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^{\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{23} x}{2} \right) + c_2 \sin \left(\frac{\sqrt{23} x}{2} \right) \right)$$

Summary

The solution(s) found are the following

$$y = e^{\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{23} x}{2} \right) + c_2 \sin \left(\frac{\sqrt{23} x}{2} \right) \right) \quad (1)$$




Figure 379: Slope field plot

Verification of solutions

$$y = e^{\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{23} x}{2} \right) + c_2 \sin \left(\frac{\sqrt{23} x}{2} \right) \right)$$

Verified OK.

14.4.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' - y' + 6y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -1 \\ C &= 6 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-23}{4} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -23 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -\frac{23z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 338: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0\end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -\frac{23}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos\left(\frac{\sqrt{23}x}{2}\right)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-1}{1} dx} \\ &= z_1 e^{\frac{x}{2}} \\ &= z_1 \left(e^{\frac{x}{2}}\right)\end{aligned}$$

Which simplifies to

$$y_1 = e^{\frac{x}{2}} \cos\left(\frac{\sqrt{23}x}{2}\right)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{-1}{1} dx}}{(y_1)^2} dx \\&= y_1 \int \frac{e^x}{(y_1)^2} dx \\&= y_1 \left(\frac{2\sqrt{23} \tan\left(\frac{\sqrt{23}x}{2}\right)}{23} \right)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\&= c_1 \left(e^{\frac{x}{2}} \cos\left(\frac{\sqrt{23}x}{2}\right) \right) + c_2 \left(e^{\frac{x}{2}} \cos\left(\frac{\sqrt{23}x}{2}\right) \left(\frac{2\sqrt{23} \tan\left(\frac{\sqrt{23}x}{2}\right)}{23} \right) \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{\frac{x}{2}} \cos\left(\frac{\sqrt{23}x}{2}\right) + \frac{2c_2 e^{\frac{x}{2}} \sqrt{23} \sin\left(\frac{\sqrt{23}x}{2}\right)}{23} \quad (1)$$




Figure 380: Slope field plot

Verification of solutions

$$y = c_1 e^{\frac{x}{2}} \cos\left(\frac{\sqrt{23}x}{2}\right) + \frac{2c_2 e^{\frac{x}{2}} \sqrt{23} \sin\left(\frac{\sqrt{23}x}{2}\right)}{23}$$

Verified OK.

14.4.3 Maple step by step solution

Let's solve

$$y'' - y' + 6y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 - r + 6 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{1 \pm (\sqrt{-23})}{2}$$

- Roots of the characteristic polynomial

$$r = \left(\frac{1}{2} - \frac{i\sqrt{23}}{2}, \frac{1}{2} + \frac{i\sqrt{23}}{2} \right)$$

- 1st solution of the ODE

$$y_1(x) = e^{\frac{x}{2}} \cos\left(\frac{\sqrt{23}x}{2}\right)$$

- 2nd solution of the ODE

$$y_2(x) = e^{\frac{x}{2}} \sin\left(\frac{\sqrt{23}x}{2}\right)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = c_1 e^{\frac{x}{2}} \cos\left(\frac{\sqrt{23}x}{2}\right) + c_2 e^{\frac{x}{2}} \sin\left(\frac{\sqrt{23}x}{2}\right)$$

Maple trace

```

Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 28

```
dsolve(diff(y(x),x$2)-diff(y(x),x)+6*y(x)=0,y(x), singsol=all)
```

$$y(x) = e^{\frac{x}{2}} \left(c_1 \sin\left(\frac{\sqrt{23}x}{2}\right) + c_2 \cos\left(\frac{\sqrt{23}x}{2}\right) \right)$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 42

```
DSolve[y''[x]-y'[x]+6*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{x/2} \left(c_2 \cos \left(\frac{\sqrt{23}x}{2} \right) + c_1 \sin \left(\frac{\sqrt{23}x}{2} \right) \right)$$

14.5 problem 1(e)

14.5.1 Solving as second order linear constant coeff ode	2331
14.5.2 Solving using Kovacic algorithm	2334
14.5.3 Maple step by step solution	2339

Internal problem ID [6375]

Internal file name [OUTPUT/5623_Sunday_June_05_2022_03_45_04_PM_72793540/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Problems for Review and Discovery. Drill excercises. Page 105

Problem number: 1(e).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"kovacic", "second_order_linear_constant_coeff"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - 2y' - 5y = x$$

14.5.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = -2, C = -5, f(x) = x$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 2y' - 5y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -2, C = -5$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 2\lambda e^{\lambda x} - 5 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 2\lambda - 5 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -2, C = -5$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{2}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-2^2 - (4)(1)(-5)} \\ &= 1 \pm \sqrt{6} \end{aligned}$$

Hence

$$\lambda_1 = 1 + \sqrt{6}$$

$$\lambda_2 = 1 - \sqrt{6}$$

Which simplifies to

$$\lambda_1 = 1 + \sqrt{6}$$

$$\lambda_2 = 1 - \sqrt{6}$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(1+\sqrt{6})x} + c_2 e^{(1-\sqrt{6})x}$$

Or

$$y = c_1 e^{(1+\sqrt{6})x} + c_2 e^{(1-\sqrt{6})x}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^{(1+\sqrt{6})x} + c_2 e^{(1-\sqrt{6})x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{(1-\sqrt{6})x}, e^{(1+\sqrt{6})x} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_2x + A_1$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-5A_2x - 5A_1 - 2A_2 = x$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{2}{25}, A_2 = -\frac{1}{5} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{x}{5} + \frac{2}{25}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 e^{(1+\sqrt{6})x} + c_2 e^{(1-\sqrt{6})x} \right) + \left(-\frac{x}{5} + \frac{2}{25} \right) \end{aligned}$$

Which simplifies to

$$y = c_1 e^{(1+\sqrt{6})x} + c_2 e^{-(1+\sqrt{6})x} - \frac{x}{5} + \frac{2}{25}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{(1+\sqrt{6})x} + c_2 e^{-(1+\sqrt{6})x} - \frac{x}{5} + \frac{2}{25} \quad (1)$$




Figure 381: Slope field plot

Verification of solutions

$$y = c_1 e^{(1+\sqrt{6})x} + c_2 e^{-(1+\sqrt{6})x} - \frac{x}{5} + \frac{2}{25}$$

Verified OK.

14.5.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 2y' - 5y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= 1 \\B &= -2 \\C &= -5\end{aligned}\tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x)\tag{4}$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\&= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned}\tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{6}{1}\tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= 6 \\t &= 1\end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 6z(x)\tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 340: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 6$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-x\sqrt{6}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}
 y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\
 &= z_1 e^{-\int \frac{1}{2} \frac{-2}{1} dx} \\
 &= z_1 e^x \\
 &= z_1 (e^x)
 \end{aligned}$$

Which simplifies to

$$y_1 = e^{-(-1+\sqrt{6})x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-2}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{2x}}{(y_1)^2} dx \\ &= y_1 \left(\frac{\sqrt{6} e^{2x\sqrt{6}}}{12} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(e^{-(-1+\sqrt{6})x} \right) + c_2 \left(e^{-(-1+\sqrt{6})x} \left(\frac{\sqrt{6} e^{2x\sqrt{6}}}{12} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 2y' - 5y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-(-1+\sqrt{6})x} + \frac{c_2 \sqrt{6} e^{(1+\sqrt{6})x}}{12}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ \frac{\sqrt{6} e^{(1+\sqrt{6})x}}{12}, e^{-(-1+\sqrt{6})x} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_2x + A_1$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-5A_2x - 5A_1 - 2A_2 = x$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{2}{25}, A_2 = -\frac{1}{5} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{x}{5} + \frac{2}{25}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 e^{-(-1+\sqrt{6})x} + \frac{c_2 \sqrt{6} e^{(1+\sqrt{6})x}}{12} \right) + \left(-\frac{x}{5} + \frac{2}{25} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-(1+\sqrt{6})x} + \frac{c_2 \sqrt{6} e^{(1+\sqrt{6})x}}{12} - \frac{x}{5} + \frac{2}{25} \quad (1)$$




Figure 382: Slope field plot

Verification of solutions

$$y = c_1 e^{-(1+\sqrt{6})x} + \frac{c_2 \sqrt{6} e^{(1+\sqrt{6})x}}{12} - \frac{x}{5} + \frac{2}{25}$$

Verified OK.

14.5.3 Maple step by step solution

Let's solve

$$y'' - 2y' - 5y = x$$

- Highest derivative means the order of the ODE is 2

y''

- Characteristic polynomial of homogeneous ODE

$$r^2 - 2r - 5 = 0$$
- Use quadratic formula to solve for r

$$r = \frac{2 \pm (\sqrt{24})}{2}$$
- Roots of the characteristic polynomial

$$r = (1 - \sqrt{6}, 1 + \sqrt{6})$$
- 1st solution of the homogeneous ODE

$$y_1(x) = e^{(1-\sqrt{6})x}$$
- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{(1+\sqrt{6})x}$$
- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$
- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{(1-\sqrt{6})x} + c_2 e^{(1+\sqrt{6})x} + y_p(x)$$
- Find a particular solution $y_p(x)$ of the ODE
 - Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = x \right]$$
 - Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{(1-\sqrt{6})x} & e^{(1+\sqrt{6})x} \\ (1 - \sqrt{6}) e^{(1-\sqrt{6})x} & (1 + \sqrt{6}) e^{(1+\sqrt{6})x} \end{bmatrix}$$
 - Compute Wronskian

$$W(y_1(x), y_2(x)) = 2\sqrt{6} e^{2x}$$
 - Substitute functions into equation for $y_p(x)$

$$y_p(x) = \frac{\sqrt{6} \left(-e^{-(1+\sqrt{6})x} \left(\int x e^{-(1+\sqrt{6})x} dx \right) + e^{(1+\sqrt{6})x} \left(\int x e^{-(1+\sqrt{6})x} dx \right) \right)}{12}$$
 - Compute integrals

$$y_p(x) = -\frac{x}{5} + \frac{2}{25}$$
- Substitute particular solution into general solution to ODE

$$y = c_1 e^{(1-\sqrt{6})x} + c_2 e^{(1+\sqrt{6})x} - \frac{x}{5} + \frac{2}{25}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 30

```
dsolve(diff(y(x),x$2)-2*diff(y(x),x)-5*y(x)=x,y(x), singsol=all)
```

$$y(x) = e^{(1+\sqrt{6})x} c_2 + e^{-(1+\sqrt{6})x} c_1 - \frac{x}{5} + \frac{2}{25}$$

✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 43

```
DSolve[y''[x]-2*y'[x]-5*y[x]==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{x}{5} + c_1 e^{x-\sqrt{6}x} + c_2 e^{(1+\sqrt{6})x} + \frac{2}{25}$$

14.6 problem 1(f)

14.6.1 Solving as second order linear constant coeff ode	2342
14.6.2 Solving using Kovacic algorithm	2345
14.6.3 Maple step by step solution	2350

Internal problem ID [6376]

Internal file name [OUTPUT/5624_Sunday_June_05_2022_03_45_06_PM_61507785/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Problems for Review and Discovery. Drill excercises. Page 105

Problem number: 1(f).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' + y = e^x$$

14.6.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 1, f(x) = e^x$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\lambda_1 = +i$$

$$\lambda_2 = -i$$

Which simplifies to

$$\lambda_1 = i$$

$$\lambda_2 = -i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (\cos(x) c_1 + c_2 \sin(x))$$

Or

$$y = \cos(x) c_1 + c_2 \sin(x)$$

Therefore the homogeneous solution y_h is

$$y_h = \cos(x) c_1 + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$e^x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^x$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$2A_1 e^x = e^x$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{2} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{e^x}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (\cos(x) c_1 + c_2 \sin(x)) + \left(\frac{e^x}{2} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \cos(x) c_1 + c_2 \sin(x) + \frac{e^x}{2} \quad (1)$$




Figure 383: Slope field plot

Verification of solutions

$$y = \cos(x) c_1 + c_2 \sin(x) + \frac{e^x}{2}$$

Verified OK.

14.6.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= 1 \\B &= 0 \\C &= 1\end{aligned}\tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x)\tag{4}$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\&= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned}\tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1}\tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= -1 \\t &= 1\end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x)\tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 342: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}
 y_1 &= z_1 \\
 &= \cos(x)
 \end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x)) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(x)) + c_2 (\cos(x) (\tan(x))) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = \cos(x) c_1 + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$e^x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^x$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$2A_1 e^x = e^x$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{2} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{e^x}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (\cos(x) c_1 + c_2 \sin(x)) + \left(\frac{e^x}{2} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \cos(x) c_1 + c_2 \sin(x) + \frac{e^x}{2} \quad (1)$$




Figure 384: Slope field plot

Verification of solutions

$$y = \cos(x) c_1 + c_2 \sin(x) + \frac{e^x}{2}$$

Verified OK.

14.6.3 Maple step by step solution

Let's solve

$$y'' + y = e^x$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (-I, I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = \cos(x) c_1 + c_2 \sin(x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = e^x \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\cos(x) \left(\int \sin(x) e^x dx \right) + \sin(x) \left(\int \cos(x) e^x dx \right)$$

- Compute integrals

$$y_p(x) = \frac{e^x}{2}$$

- Substitute particular solution into general solution to ODE

$$y = \cos(x) c_1 + c_2 \sin(x) + \frac{e^x}{2}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 17

```
dsolve(diff(y(x),x$2)+y(x)=exp(x),y(x), singsol=all)
```

$$y(x) = \sin(x) c_2 + \cos(x) c_1 + \frac{e^x}{2}$$

✓ Solution by Mathematica

Time used: 0.058 (sec). Leaf size: 23

```
DSolve[y''[x]+y[x]==Exp[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^x}{2} + c_1 \cos(x) + c_2 \sin(x)$$

14.7 problem 1(g)

14.7.1 Solving as second order linear constant coeff ode	2353
14.7.2 Solving using Kovacic algorithm	2357
14.7.3 Maple step by step solution	2362

Internal problem ID [6377]

Internal file name [OUTPUT/5625_Sunday_June_05_2022_03_45_07_PM_38430624/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Problems for Review and Discovery. Drill excercises. Page 105

Problem number: 1(g).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + y' + y = \sin(x)$$

14.7.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 1, C = 1, f(x) = \sin(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 1, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + \lambda e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + \lambda + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 1, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{1^2 - (4)(1)(1)} \\ &= -\frac{1}{2} \pm \frac{i\sqrt{3}}{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= -\frac{1}{2} + \frac{i\sqrt{3}}{2} \\ \lambda_2 &= -\frac{1}{2} - \frac{i\sqrt{3}}{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= -\frac{1}{2} + \frac{i\sqrt{3}}{2} \\ \lambda_2 &= -\frac{1}{2} - \frac{i\sqrt{3}}{2} \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = -\frac{1}{2}$ and $\beta = \frac{\sqrt{3}}{2}$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

Therefore the homogeneous solution y_h is

$$y_h = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3}x}{2} \right), e^{-\frac{x}{2}} \sin \left(\frac{\sqrt{3}x}{2} \right) \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(x) + A_2 \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-A_1 \sin(x) + A_2 \cos(x) = \sin(x)$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = -1, A_2 = 0]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\cos(x)$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right) \right) + (-\cos(x)) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right) - \cos(x) \quad (1)$$




Figure 385: Slope field plot

Verification of solutions

$$y = e^{-\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{3}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{3}x}{2} \right) \right) - \cos(x)$$

Verified OK.

14.7.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 1 \\ C &= 1 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-3}{4} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -3 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -\frac{3z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 344: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -\frac{3}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos\left(\frac{\sqrt{3}x}{2}\right)$$

Using the above, the solution for the original ode can now be found. The first solution

to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\&= z_1 e^{-\int \frac{1}{2} \frac{1}{1} dx} \\&= z_1 e^{-\frac{x}{2}} \\&= z_1 \left(e^{-\frac{x}{2}} \right)\end{aligned}$$

Which simplifies to

$$y_1 = e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3}x}{2} \right)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{1}{1} dx}}{(y_1)^2} dx \\&= y_1 \int \frac{e^{-x}}{(y_1)^2} dx \\&= y_1 \left(\frac{2\sqrt{3} \tan \left(\frac{\sqrt{3}x}{2} \right)}{3} \right)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\&= c_1 \left(e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3}x}{2} \right) \right) + c_2 \left(e^{-\frac{x}{2}} \cos \left(\frac{\sqrt{3}x}{2} \right) \left(\frac{2\sqrt{3} \tan \left(\frac{\sqrt{3}x}{2} \right)}{3} \right) \right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) c_1 + \frac{2c_2 e^{-\frac{x}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right), \frac{2 e^{-\frac{x}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(x) + A_2 \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-A_1 \sin(x) + A_2 \cos(x) = \sin(x)$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = -1, A_2 = 0]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\cos(x)$$

Therefore the general solution is

$$y = y_h + y_p$$

$$= \left(e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) c_1 + \frac{2c_2 e^{-\frac{x}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3} \right) + (-\cos(x))$$

Summary

The solution(s) found are the following

$$y = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) c_1 + \frac{2c_2 e^{-\frac{x}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3} - \cos(x) \quad (1)$$




Figure 386: Slope field plot

Verification of solutions

$$y = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) c_1 + \frac{2c_2 e^{-\frac{x}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{3} - \cos(x)$$

Verified OK.

14.7.3 Maple step by step solution

Let's solve

$$y'' + y' + y = \sin(x)$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + r + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-1) \pm (\sqrt{-3})}{2}$$

- Roots of the characteristic polynomial

$$r = \left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}, -\frac{1}{2} + \frac{i\sqrt{3}}{2} \right)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) c_1 + e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) c_2 + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \sin(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) & e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) \\ -\frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{2} - \frac{e^{-\frac{x}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{2} & -\frac{e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) \sqrt{3}}{2} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = \frac{\sqrt{3}e^{-x}}{2}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{2\sqrt{3}e^{-\frac{x}{2}} \left(\cos\left(\frac{\sqrt{3}x}{2}\right) \left(\int e^{\frac{x}{2}} \sin(x) \sin\left(\frac{\sqrt{3}x}{2}\right) dx \right) - \sin\left(\frac{\sqrt{3}x}{2}\right) \left(\int e^{\frac{x}{2}} \sin(x) \cos\left(\frac{\sqrt{3}x}{2}\right) dx \right) \right)}{3}$$

- Compute integrals

$$y_p(x) = -\cos(x)$$

- Substitute particular solution into general solution to ODE

$$y = e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) c_2 + e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) c_1 - \cos(x)$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 35

```
dsolve(diff(y(x),x$2)+diff(y(x),x)+y(x)=sin(x),y(x), singsol=all)
```

$$y(x) = c_2 e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) + e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) c_1 - \cos(x)$$

✓ Solution by Mathematica

Time used: 1.371 (sec). Leaf size: 53

```
DSolve[y''[x]+y'[x]+y[x]==Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x/2} \left(-e^{x/2} \cos(x) + c_2 \cos\left(\frac{\sqrt{3}x}{2}\right) + c_1 \sin\left(\frac{\sqrt{3}x}{2}\right) \right)$$

14.8 problem 1(h)

14.8.1 Solving as second order linear constant coeff ode	2365
14.8.2 Solving using Kovacic algorithm	2368
14.8.3 Maple step by step solution	2373

Internal problem ID [6378]

Internal file name [OUTPUT/5626_Sunday_June_05_2022_03_45_09_PM_98016615/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Problems for Review and Discovery. Drill excercises. Page 105

Problem number: 1(h).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"kovacic", "second_order_linear_constant_coeff"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - y = e^{3x}$$

14.8.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = -1, f(x) = e^{3x}$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = -1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = -1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(-1)} \\ &= \pm 1 \end{aligned}$$

Hence

$$\lambda_1 = +1$$

$$\lambda_2 = -1$$

Which simplifies to

$$\lambda_1 = 1$$

$$\lambda_2 = -1$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(1)x} + c_2 e^{(-1)x}$$

Or

$$y = c_1 e^x + c_2 e^{-x}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^x + c_2 e^{-x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$e^{3x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{3x}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^x, e^{-x}\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^{3x}$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$8A_1 e^{3x} = e^{3x}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{8} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{e^{3x}}{8}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^x + c_2 e^{-x}) + \left(\frac{e^{3x}}{8} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^x + c_2 e^{-x} + \frac{e^{3x}}{8} \quad (1)$$




Figure 387: Slope field plot

Verification of solutions

$$y = c_1 e^x + c_2 e^{-x} + \frac{e^{3x}}{8}$$

Verified OK.

14.8.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' - y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 0 \tag{3}$$

$$C = -1$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} = \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{1}{1} \tag{6}$$

Comparing the above to (5) shows that

$$s = 1 \\ t = 1$$

Therefore eq. (4) becomes

$$z''(x) = z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 346: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-x}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned} y_1 &= z_1 \\ &= e^{-x} \end{aligned}$$

Which simplifies to

$$y_1 = e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= e^{-x} \int \frac{1}{e^{-2x}} dx \\ &= e^{-x} \left(\frac{e^{2x}}{2} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-x}) + c_2 \left(e^{-x} \left(\frac{e^{2x}}{2} \right) \right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-x} + \frac{c_2 e^x}{2}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$e^{3x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{3x}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ \frac{e^x}{2}, e^{-x} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^{3x}$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$8A_1e^{3x} = e^{3x}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{8} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{e^{3x}}{8}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1e^{-x} + \frac{c_2e^x}{2} \right) + \left(\frac{e^{3x}}{8} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1e^{-x} + \frac{c_2e^x}{2} + \frac{e^{3x}}{8} \quad (1)$$




Figure 388: Slope field plot

Verification of solutions

$$y = c_1 e^{-x} + \frac{c_2 e^x}{2} + \frac{e^{3x}}{8}$$

Verified OK.

14.8.3 Maple step by step solution

Let's solve

$$y'' - y = e^{3x}$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - 1 = 0$$

- Factor the characteristic polynomial

$$(r - 1)(r + 1) = 0$$

- Roots of the characteristic polynomial

$$r = (-1, 1)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^x$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{-x} + c_2 e^x + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = e^{3x} \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-x} & e^x \\ -e^{-x} & e^x \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 2$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{e^{-x}(\int e^{4x} dx)}{2} + \frac{e^x(\int e^{2x} dx)}{2}$$

- Compute integrals

$$y_p(x) = \frac{e^{3x}}{8}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{-x} + c_2 e^x + \frac{e^{3x}}{8}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 21

```
dsolve(diff(y(x),x$2)-y(x)=exp(3*x),y(x), singsol=all)
```

$$y(x) = c_2 e^{-x} + e^x c_1 + \frac{e^{3x}}{8}$$

✓ Solution by Mathematica

Time used: 0.029 (sec). Leaf size: 29

```
DSolve[y''[x]-y[x]==Exp[3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{3x}}{8} + c_1 e^x + c_2 e^{-x}$$

14.9 problem 2(a)

14.9.1 Existence and uniqueness analysis	2376
14.9.2 Solving as second order linear constant coeff ode	2377
14.9.3 Solving as second order ode can be made integrable ode	2379
14.9.4 Solving using Kovacic algorithm	2382
14.9.5 Maple step by step solution	2386

Internal problem ID [6379]

Internal file name [OUTPUT/5627_Sunday_June_05_2022_03_45_11_PM_75680248/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Problems for Review and Discovery. Drill excercises. Page 105

Problem number: 2(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "second_order_ode_can_be_made_integrable"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' + 9y = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 2]$$

14.9.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = 0$$

$$q(x) = 9$$

$$F = 0$$

Hence the ode is

$$y'' + 9y = 0$$

The domain of $p(x) = 0$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = 9$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

14.9.2 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 9$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 9e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 9 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 9$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(9)} \\ &= \pm 3i \end{aligned}$$

Hence

$$\lambda_1 = +3i$$

$$\lambda_2 = -3i$$

Which simplifies to

$$\lambda_1 = 3i$$

$$\lambda_2 = -3i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 3$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0(c_1 \cos(3x) + c_2 \sin(3x))$$

Or

$$y = c_1 \cos(3x) + c_2 \sin(3x)$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 \cos(3x) + c_2 \sin(3x) \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 0$ in the above gives

$$1 = c_1 \tag{1A}$$

Taking derivative of the solution gives

$$y' = -3c_1 \sin(3x) + 3c_2 \cos(3x)$$

substituting $y' = 2$ and $x = 0$ in the above gives

$$2 = 3c_2 \tag{2A}$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 1$$

$$c_2 = \frac{2}{3}$$


Substituting these values back in above solution results in

$$y = \frac{2 \sin(3x)}{3} + \cos(3x)$$


Summary

The solution(s) found are the following

$$y = \frac{2 \sin(3x)}{3} + \cos(3x) \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{2 \sin(3x)}{3} + \cos(3x)$$

Verified OK.

14.9.3 Solving as second order ode can be made integrable ode

Multiplying the ode by y' gives

$$y'y'' + 9y'y' = 0$$

Integrating the above w.r.t x gives

$$\int (y'y'' + 9y'y') dx = 0$$

$$\frac{y'^2}{2} + \frac{9y^2}{2} = c_2$$

Which is now solved for y . Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \sqrt{-9y^2 + 2c_1} \quad (1)$$

$$y' = -\sqrt{-9y^2 + 2c_1} \quad (2)$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int \frac{1}{\sqrt{-9y^2 + 2c_1}} dy = \int dx$$

$$\frac{\arctan\left(\frac{3y}{\sqrt{-9y^2 + 2c_1}}\right)}{3} = c_2 + x$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{1}{\sqrt{-9y^2 + 2c_1}} dy = \int dx$$

$$-\frac{\arctan\left(\frac{3y}{\sqrt{-9y^2 + 2c_1}}\right)}{3} = x + c_3$$

Initial conditions are used to solve for the constants of integration.

Looking at the First solution

$$\frac{\arctan\left(\frac{3y}{\sqrt{-9y^2 + 2c_1}}\right)}{3} = c_2 + x \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 0$ in the above gives

$$\frac{\arctan\left(\frac{3}{\sqrt{-9+2c_1}}\right)}{3} = c_2 \quad (1A)$$

Taking derivative of the solution gives

$$y' = \frac{(3 \tan(3c_2 + 3x))^2 + 3}{3} \sqrt{2} \sqrt{\frac{c_1}{\tan(3c_2 + 3x)^2 + 1}} - \frac{\tan(3c_2 + 3x)^2 \sqrt{2} c_1 (3 \tan(3c_2 + 3x))^2 + 3}{3 \sqrt{\frac{c_1}{\tan(3c_2 + 3x)^2 + 1}} (\tan(3c_2 + 3x)^2 + 1)^2}$$

substituting $y' = 2$ and $x = 0$ in the above gives

$$2 = \frac{\cos(3c_2)^2 c_1 \sqrt{2}}{\sqrt{c_1 \cos(3c_2)^2}} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = \frac{13}{2}$$

$$c_2 = \frac{\arctan\left(\frac{3}{2}\right)}{3}$$

Substituting these values back in above solution results in

$$\frac{\arctan\left(\frac{3y}{\sqrt{-9y^2+13}}\right)}{3} = \frac{\arctan\left(\frac{3}{2}\right)}{3} + x$$

Looking at the Second solution

$$-\frac{\arctan\left(\frac{3y}{\sqrt{-9y^2+2c_1}}\right)}{3} = x + c_3 \quad (2)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 0$ in the above gives

$$-\frac{\arctan\left(\frac{3}{\sqrt{-9+2c_1}}\right)}{3} = c_3 \quad (1A)$$

Taking derivative of the solution gives

$$y' = -\frac{(3 \tan(3x + 3c_3))^2 + 3}{3} \sqrt{2} \sqrt{\frac{c_1}{\tan(3x + 3c_3)^2 + 1}} + \frac{\tan(3x + 3c_3)^2 \sqrt{2} c_1 (3 \tan(3x + 3c_3))^2 + 3}{3 \sqrt{\frac{c_1}{\tan(3x + 3c_3)^2 + 1}} (\tan(3x + 3c_3)^2 + 1)^2}$$

substituting $y' = 2$ and $x = 0$ in the above gives

$$2 = -\frac{\cos(3c_3)^2 c_1 \sqrt{2}}{\sqrt{c_1 \cos(3c_3)^2}} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_3\}$. There is no solution for the constants of integrations. This solution is removed.

Summary

The solution(s) found are the following

$$\frac{\arctan\left(\frac{3y}{\sqrt{-9y^2+13}}\right)}{3} = \frac{\arctan\left(\frac{3}{2}\right)}{3} + x \quad (1)$$

Verification of solutions

$$\frac{\arctan\left(\frac{3y}{\sqrt{-9y^2+13}}\right)}{3} = \frac{\arctan\left(\frac{3}{2}\right)}{3} + x$$

Verified OK.

14.9.4 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 9y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 9 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-9}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -9 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -9z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 348: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -9$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(3x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned} y_1 &= z_1 \\ &= \cos(3x) \end{aligned}$$

Which simplifies to

$$y_1 = \cos(3x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(3x) \int \frac{1}{\cos(3x)^2} dx \\ &= \cos(3x) \left(\frac{\tan(3x)}{3} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(3x)) + c_2 \left(\cos(3x) \left(\frac{\tan(3x)}{3} \right) \right) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 \cos(3x) + \frac{c_2 \sin(3x)}{3} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 0$ in the above gives

$$1 = c_1 \quad (1A)$$

Taking derivative of the solution gives

$$y' = -3c_1 \sin(3x) + c_2 \cos(3x)$$

substituting $y' = 2$ and $x = 0$ in the above gives

$$2 = c_2 \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 1$$

$$c_2 = 2$$


Substituting these values back in above solution results in

$$y = \frac{2 \sin(3x)}{3} + \cos(3x)$$


Summary

The solution(s) found are the following

$$y = \frac{2 \sin(3x)}{3} + \cos(3x) \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{2 \sin(3x)}{3} + \cos(3x)$$

Verified OK.

14.9.5 Maple step by step solution

Let's solve

$$\left[y'' + 9y = 0, y(0) = 1, y' \Big|_{\{x=0\}} = 2 \right]$$

- Highest derivative means the order of the ODE is 2
- y''
- Characteristic polynomial of ODE
- $r^2 + 9 = 0$
- Use quadratic formula to solve for r
- $r = \frac{0 \pm (\sqrt{-36})}{2}$
- Roots of the characteristic polynomial
- $r = (-3I, 3I)$
- 1st solution of the ODE

$$y_1(x) = \cos(3x)$$

- 2nd solution of the ODE

$$y_2(x) = \sin(3x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x)$$

- Substitute in solutions

$$y = c_1 \cos(3x) + c_2 \sin(3x)$$

- Check validity of solution $y = c_1 \cos(3x) + c_2 \sin(3x)$

- Use initial condition $y(0) = 1$

$$1 = c_1$$

- Compute derivative of the solution

$$y' = -3c_1 \sin(3x) + 3c_2 \cos(3x)$$

- Use the initial condition $y' \Big|_{\{x=0\}} = 2$

$$2 = 3c_2$$

- Solve for c_1 and c_2

$$\left\{ c_1 = 1, c_2 = \frac{2}{3} \right\}$$

- Substitute constant values into general solution and simplify

$$y = \frac{2\sin(3x)}{3} + \cos(3x)$$

- Solution to the IVP

$$y = \frac{2\sin(3x)}{3} + \cos(3x)$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```


✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 15

```
dsolve([diff(y(x),x$2)+9*y(x)=0,y(0) = 1, D(y)(0) = 2],y(x), singsol=all)
```

$$y(x) = \frac{2 \sin(3x)}{3} + \cos(3x)$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 18

```
DSolve[{y'[x]+9*y[x]==0,{y[0]==1,y'[0]==2}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2}{3} \sin(3x) + \cos(3x)$$

14.10 problem 2(b)

14.10.1 Existence and uniqueness analysis	2389
14.10.2 Solving as second order linear constant coeff ode	2390
14.10.3 Solving using Kovacic algorithm	2395
14.10.4 Maple step by step solution	2401

Internal problem ID [6380]

Internal file name [OUTPUT/5628_Sunday_June_05_2022_03_45_12_PM_67246059/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Problems for Review and Discovery. Drill exercises. Page 105

Problem number: 2(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"kovacic", "second_order_linear_constant_coeff"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - y' + 4y = x$$

With initial conditions

$$[y(1) = 2, y'(1) = 1]$$

14.10.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = -1$$

$$q(x) = 4$$

$$F = x$$

Hence the ode is

$$y'' - y' + 4y = x$$

The domain of $p(x) = -1$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 1$ is inside this domain. The domain of $q(x) = 4$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 1$ is also inside this domain. The domain of $F = x$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 1$ is also inside this domain. Hence solution exists and is unique.

14.10.2 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = -1, C = 4, f(x) = x$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - y' + 4y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -1, C = 4$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - \lambda e^{\lambda x} + 4 e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - \lambda + 4 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -1, C = 4$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-1^2 - (4)(1)(4)} \\ &= \frac{1}{2} \pm \frac{i\sqrt{15}}{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= \frac{1}{2} + \frac{i\sqrt{15}}{2} \\ \lambda_2 &= \frac{1}{2} - \frac{i\sqrt{15}}{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= \frac{1}{2} + \frac{i\sqrt{15}}{2} \\ \lambda_2 &= \frac{1}{2} - \frac{i\sqrt{15}}{2} \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = \frac{1}{2}$ and $\beta = \frac{\sqrt{15}}{2}$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^{\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{15} x}{2} \right) + c_2 \sin \left(\frac{\sqrt{15} x}{2} \right) \right)$$

Therefore the homogeneous solution y_h is

$$y_h = e^{\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{15} x}{2} \right) + c_2 \sin \left(\frac{\sqrt{15} x}{2} \right) \right)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right), e^{\frac{x}{2}} \sin\left(\frac{\sqrt{15}x}{2}\right) \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_2x + A_1$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$4A_2x + 4A_1 - A_2 = x$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{16}, A_2 = \frac{1}{4} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{x}{4} + \frac{1}{16}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(e^{\frac{x}{2}} \left(c_1 \cos\left(\frac{\sqrt{15}x}{2}\right) + c_2 \sin\left(\frac{\sqrt{15}x}{2}\right) \right) \right) + \left(\frac{x}{4} + \frac{1}{16} \right) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = e^{\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{15} x}{2} \right) + c_2 \sin \left(\frac{\sqrt{15} x}{2} \right) \right) + \frac{x}{4} + \frac{1}{16} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 2$ and $x = 1$ in the above gives

$$2 = e^{\frac{1}{2}} \cos \left(\frac{\sqrt{15}}{2} \right) c_1 + e^{\frac{1}{2}} \sin \left(\frac{\sqrt{15}}{2} \right) c_2 + \frac{5}{16} \quad (1A)$$

Taking derivative of the solution gives

$$y' = \frac{e^{\frac{x}{2}} \left(c_1 \cos \left(\frac{\sqrt{15} x}{2} \right) + c_2 \sin \left(\frac{\sqrt{15} x}{2} \right) \right)}{2} + e^{\frac{x}{2}} \left(-\frac{c_1 \sqrt{15} \sin \left(\frac{\sqrt{15} x}{2} \right)}{2} + \frac{c_2 \sqrt{15} \cos \left(\frac{\sqrt{15} x}{2} \right)}{2} \right) + \frac{1}{4}$$

substituting $y' = 1$ and $x = 1$ in the above gives

$$1 = \frac{e^{\frac{1}{2}} (\sqrt{15} c_2 + c_1) \cos \left(\frac{\sqrt{15}}{2} \right)}{2} + \frac{1}{4} - \frac{e^{\frac{1}{2}} (\sqrt{15} c_1 - c_2) \sin \left(\frac{\sqrt{15}}{2} \right)}{2} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = \frac{e^{-\frac{1}{2}} \sqrt{15} \left(9\sqrt{15} \cos \left(\frac{\sqrt{15}}{2} \right) + \sin \left(\frac{\sqrt{15}}{2} \right) \right)}{80}$$

$$c_2 = \frac{e^{-\frac{1}{2}} \sqrt{15} \left(9\sqrt{15} \sin \left(\frac{\sqrt{15}}{2} \right) - \cos \left(\frac{\sqrt{15}}{2} \right) \right)}{80}$$

Substituting these values back in above solution results in

$$y = \frac{1}{16} + \frac{x}{4} + \frac{27 \sin \left(\frac{\sqrt{15} x}{2} \right) e^{\frac{x}{2} - \frac{1}{2}} \sin \left(\frac{\sqrt{15}}{2} \right)}{16} - \frac{\sin \left(\frac{\sqrt{15} x}{2} \right) e^{\frac{x}{2} - \frac{1}{2}} \sqrt{15} \cos \left(\frac{\sqrt{15}}{2} \right)}{80} + \frac{27 \cos \left(\frac{\sqrt{15} x}{2} \right) \cos \left(\frac{\sqrt{15}}{2} \right)}{16}$$


Which simplifies to

$$y = \frac{\left(\left(\sqrt{15} \sin \left(\frac{\sqrt{15}}{2} \right) + 135 \cos \left(\frac{\sqrt{15}}{2} \right) \right) \cos \left(\frac{\sqrt{15} x}{2} \right) - \sin \left(\frac{\sqrt{15} x}{2} \right) \left(\sqrt{15} \cos \left(\frac{\sqrt{15}}{2} \right) - 135 \sin \left(\frac{\sqrt{15}}{2} \right) \right) \right) e^{\frac{x}{2} - \frac{1}{2}}}{80} + \frac{x}{4} + \frac{1}{16}$$


Summary

The solution(s) found are the following

$$y = \frac{\left(\left(\sqrt{15} \sin\left(\frac{\sqrt{15}}{2}\right) + 135 \cos\left(\frac{\sqrt{15}}{2}\right) \right) \cos\left(\frac{\sqrt{15}x}{2}\right) - \sin\left(\frac{\sqrt{15}x}{2}\right) \left(\sqrt{15} \cos\left(\frac{\sqrt{15}}{2}\right) - 135 \sin\left(\frac{\sqrt{15}}{2}\right) \right) \right) e^{\frac{x}{2} - \frac{1}{16}}}{80} + \frac{x}{4} + \frac{1}{16} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{\left(\left(\sqrt{15} \sin\left(\frac{\sqrt{15}}{2}\right) + 135 \cos\left(\frac{\sqrt{15}}{2}\right) \right) \cos\left(\frac{\sqrt{15}x}{2}\right) - \sin\left(\frac{\sqrt{15}x}{2}\right) \left(\sqrt{15} \cos\left(\frac{\sqrt{15}}{2}\right) - 135 \sin\left(\frac{\sqrt{15}}{2}\right) \right) \right) e^{\frac{x}{2} - \frac{1}{16}}}{80} + \frac{x}{4} + \frac{1}{16}$$

Verified OK.

14.10.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' - y' + 4y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = -1 \quad (3)$$

$$C = 4$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$r = \frac{s}{t} \quad (5)$$
$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-15}{4} \quad (6)$$

Comparing the above to (5) shows that

$$s = -15$$

$$t = 4$$

Therefore eq. (4) becomes

$$z''(x) = -\frac{15z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 350: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -\frac{15}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos\left(\frac{\sqrt{15}x}{2}\right)$$

Using the above, the solution for the original ode can now be found. The first solution

to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\&= z_1 e^{-\int \frac{1}{2} \frac{-1}{1} dx} \\&= z_1 e^{\frac{x}{2}} \\&= z_1 \left(e^{\frac{x}{2}} \right)\end{aligned}$$

Which simplifies to

$$y_1 = e^{\frac{x}{2}} \cos \left(\frac{\sqrt{15} x}{2} \right)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{-1}{1} dx}}{(y_1)^2} dx \\&= y_1 \int \frac{e^x}{(y_1)^2} dx \\&= y_1 \left(\frac{2\sqrt{15} \tan \left(\frac{\sqrt{15} x}{2} \right)}{15} \right)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\&= c_1 \left(e^{\frac{x}{2}} \cos \left(\frac{\sqrt{15} x}{2} \right) \right) + c_2 \left(e^{\frac{x}{2}} \cos \left(\frac{\sqrt{15} x}{2} \right) \left(\frac{2\sqrt{15} \tan \left(\frac{\sqrt{15} x}{2} \right)}{15} \right) \right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - y' + 4y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right) + \frac{2c_2 e^{\frac{x}{2}} \sqrt{15} \sin\left(\frac{\sqrt{15}x}{2}\right)}{15}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right), \frac{2 e^{\frac{x}{2}} \sqrt{15} \sin\left(\frac{\sqrt{15}x}{2}\right)}{15} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_2 x + A_1$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$4A_2 x + 4A_1 - A_2 = x$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{16}, A_2 = \frac{1}{4} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{x}{4} + \frac{1}{16}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 e^{\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right) + \frac{2c_2 e^{\frac{x}{2}} \sqrt{15} \sin\left(\frac{\sqrt{15}x}{2}\right)}{15} \right) + \left(\frac{x}{4} + \frac{1}{16} \right) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 e^{\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right) + \frac{2c_2 e^{\frac{x}{2}} \sqrt{15} \sin\left(\frac{\sqrt{15}x}{2}\right)}{15} + \frac{x}{4} + \frac{1}{16} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 2$ and $x = 1$ in the above gives

$$2 = e^{\frac{1}{2}} \cos\left(\frac{\sqrt{15}}{2}\right) c_1 + \frac{2c_2 e^{\frac{1}{2}} \sqrt{15} \sin\left(\frac{\sqrt{15}}{2}\right)}{15} + \frac{5}{16} \quad (1A)$$

Taking derivative of the solution gives

$$y' = \frac{c_1 e^{\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right)}{2} - \frac{c_1 e^{\frac{x}{2}} \sqrt{15} \sin\left(\frac{\sqrt{15}x}{2}\right)}{2} + \frac{c_2 e^{\frac{x}{2}} \sqrt{15} \sin\left(\frac{\sqrt{15}x}{2}\right)}{15} + c_2 e^{\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right) + \frac{1}{4}$$

substituting $y' = 1$ and $x = 1$ in the above gives

$$1 = \frac{e^{\frac{1}{2}}(c_1 + 2c_2) \cos\left(\frac{\sqrt{15}}{2}\right)}{2} + \frac{1}{4} - \frac{(c_1 - \frac{2c_2}{15}) \sqrt{15} e^{\frac{1}{2}} \sin\left(\frac{\sqrt{15}}{2}\right)}{2} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$\begin{aligned} c_1 &= \frac{\left(\sqrt{15} \sin\left(\frac{\sqrt{15}}{2}\right) + 135 \cos\left(\frac{\sqrt{15}}{2}\right)\right) e^{-\frac{1}{2}}}{80} \\ c_2 &= \frac{3\left(9\sqrt{15} \sin\left(\frac{\sqrt{15}}{2}\right) - \cos\left(\frac{\sqrt{15}}{2}\right)\right) e^{-\frac{1}{2}}}{32} \end{aligned}$$

Substituting these values back in above solution results in

$$y = \frac{1}{16} + \frac{x}{4} + \frac{27 \sin\left(\frac{\sqrt{15}x}{2}\right) e^{\frac{x}{2} - \frac{1}{2}} \sin\left(\frac{\sqrt{15}}{2}\right) - \sin\left(\frac{\sqrt{15}x}{2}\right) e^{\frac{x}{2} - \frac{1}{2}} \sqrt{15} \cos\left(\frac{\sqrt{15}}{2}\right) + 27 \cos\left(\frac{\sqrt{15}x}{2}\right) \cos\left(\frac{\sqrt{15}}{2}\right)}{16} + \frac{27 \cos\left(\frac{\sqrt{15}x}{2}\right) \cos\left(\frac{\sqrt{15}}{2}\right)}{16}$$


Which simplifies to

$$y = \frac{\left(\left(\sqrt{15} \sin\left(\frac{\sqrt{15}}{2}\right) + 135 \cos\left(\frac{\sqrt{15}}{2}\right)\right) \cos\left(\frac{\sqrt{15}x}{2}\right) - \sin\left(\frac{\sqrt{15}x}{2}\right) \left(\sqrt{15} \cos\left(\frac{\sqrt{15}}{2}\right) - 135 \sin\left(\frac{\sqrt{15}}{2}\right)\right)\right) e^{\frac{x}{2} - \frac{1}{2}}}{80} + \frac{x}{4} + \frac{1}{16}$$


Summary

The solution(s) found are the following

$$y = \frac{\left(\left(\sqrt{15} \sin\left(\frac{\sqrt{15}}{2}\right) + 135 \cos\left(\frac{\sqrt{15}}{2}\right)\right) \cos\left(\frac{\sqrt{15}x}{2}\right) - \sin\left(\frac{\sqrt{15}x}{2}\right) \left(\sqrt{15} \cos\left(\frac{\sqrt{15}}{2}\right) - 135 \sin\left(\frac{\sqrt{15}}{2}\right)\right)\right) e^{\frac{x}{2} - \frac{1}{2}}}{80} + \frac{x}{4} + \frac{1}{16} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$\begin{aligned} & y \\ = & \frac{\left(\left(\sqrt{15} \sin\left(\frac{\sqrt{15}}{2}\right) + 135 \cos\left(\frac{\sqrt{15}}{2}\right) \right) \cos\left(\frac{\sqrt{15}x}{2}\right) - \sin\left(\frac{\sqrt{15}x}{2}\right) \left(\sqrt{15} \cos\left(\frac{\sqrt{15}}{2}\right) - 135 \sin\left(\frac{\sqrt{15}}{2}\right) \right) \right) e^{\frac{x}{2} - \frac{1}{2}}}{80} \\ & + \frac{x}{4} + \frac{1}{16} \end{aligned}$$

Verified OK.

14.10.4 Maple step by step solution

Let's solve

$$\left[y'' - y' + 4y = x, y(1) = 2, y'|_{\{x=1\}} = 1 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - r + 4 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{1 \pm (\sqrt{-15})}{2}$$

- Roots of the characteristic polynomial

$$r = \left(\frac{1}{2} - \frac{i\sqrt{15}}{2}, \frac{1}{2} + \frac{i\sqrt{15}}{2} \right)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{\frac{x}{2}} \sin\left(\frac{\sqrt{15}x}{2}\right)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right) + e^{\frac{x}{2}} c_2 \sin\left(\frac{\sqrt{15}x}{2}\right) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = x \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right) & e^{\frac{x}{2}} \sin\left(\frac{\sqrt{15}x}{2}\right) \\ \frac{e^{\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right)}{2} - \frac{e^{\frac{x}{2}} \sqrt{15} \sin\left(\frac{\sqrt{15}x}{2}\right)}{2} & \frac{e^{\frac{x}{2}} \sin\left(\frac{\sqrt{15}x}{2}\right)}{2} + \frac{e^{\frac{x}{2}} \sqrt{15} \cos\left(\frac{\sqrt{15}x}{2}\right)}{2} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = \frac{\sqrt{15}e^x}{2}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{2e^{\frac{x}{2}}\sqrt{15}\left(\cos\left(\frac{\sqrt{15}x}{2}\right)\left(\int xe^{-\frac{x}{2}}\sin\left(\frac{\sqrt{15}x}{2}\right)dx\right) - \sin\left(\frac{\sqrt{15}x}{2}\right)\left(\int xe^{-\frac{x}{2}}\cos\left(\frac{\sqrt{15}x}{2}\right)dx\right)\right)}{15}$$

- Compute integrals

$$y_p(x) = \frac{x}{4} + \frac{1}{16}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right) + e^{\frac{x}{2}} c_2 \sin\left(\frac{\sqrt{15}x}{2}\right) + \frac{x}{4} + \frac{1}{16}$$

- Check validity of solution $y = c_1 e^{\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right) + e^{\frac{x}{2}} c_2 \sin\left(\frac{\sqrt{15}x}{2}\right) + \frac{x}{4} + \frac{1}{16}$

- Use initial condition $y(1) = 2$

$$2 = e^{\frac{1}{2}} \cos\left(\frac{\sqrt{15}}{2}\right) c_1 + e^{\frac{1}{2}} \sin\left(\frac{\sqrt{15}}{2}\right) c_2 + \frac{5}{16}$$

- Compute derivative of the solution

$$y' = \frac{c_1 e^{\frac{x}{2}} \cos\left(\frac{\sqrt{15}x}{2}\right)}{2} - \frac{c_1 e^{\frac{x}{2}} \sqrt{15} \sin\left(\frac{\sqrt{15}x}{2}\right)}{2} + \frac{e^{\frac{x}{2}} c_2 \sin\left(\frac{\sqrt{15}x}{2}\right)}{2} + \frac{e^{\frac{x}{2}} c_2 \sqrt{15} \cos\left(\frac{\sqrt{15}x}{2}\right)}{2} + \frac{1}{4}$$

- Use the initial condition $y'|_{\{x=1\}} = 1$

$$1 = \frac{1}{4} + \frac{e^{\frac{1}{2}} \cos\left(\frac{\sqrt{15}}{2}\right) c_1}{2} - \frac{e^{\frac{1}{2}} c_1 \sqrt{15} \sin\left(\frac{\sqrt{15}}{2}\right)}{2} + \frac{e^{\frac{1}{2}} \sin\left(\frac{\sqrt{15}}{2}\right) c_2}{2} + \frac{c_2 e^{\frac{1}{2}} \sqrt{15} \cos\left(\frac{\sqrt{15}}{2}\right)}{2}$$

- Solve for c_1 and c_2

$$\left\{ c_1 = \frac{(9\sqrt{15} \cos\left(\frac{\sqrt{15}}{2}\right) + \sin\left(\frac{\sqrt{15}}{2}\right))\sqrt{15}}{80e^{\frac{1}{2}} \left(\sin\left(\frac{\sqrt{15}}{2}\right)^2 + \cos\left(\frac{\sqrt{15}}{2}\right)^2\right)}, c_2 = \frac{\sqrt{15} \left(9\sqrt{15} \sin\left(\frac{\sqrt{15}}{2}\right) - \cos\left(\frac{\sqrt{15}}{2}\right)\right)}{80e^{\frac{1}{2}} \left(\sin\left(\frac{\sqrt{15}}{2}\right)^2 + \cos\left(\frac{\sqrt{15}}{2}\right)^2\right)} \right\}$$

- Substitute constant values into general solution and simplify

$$y = \frac{\left(\left(\sqrt{15} \sin\left(\frac{\sqrt{15}}{2}\right) + 135 \cos\left(\frac{\sqrt{15}}{2}\right)\right) \cos\left(\frac{\sqrt{15}x}{2}\right) - \sin\left(\frac{\sqrt{15}x}{2}\right) \left(\sqrt{15} \cos\left(\frac{\sqrt{15}}{2}\right) - 135 \sin\left(\frac{\sqrt{15}}{2}\right)\right)\right) e^{\frac{x}{2}} - \frac{1}{2}}{80} + \frac{x}{4} + \frac{1}{16}$$

- Solution to the IVP

$$y = \frac{\left(\left(\sqrt{15} \sin\left(\frac{\sqrt{15}}{2}\right) + 135 \cos\left(\frac{\sqrt{15}}{2}\right)\right) \cos\left(\frac{\sqrt{15}x}{2}\right) - \sin\left(\frac{\sqrt{15}x}{2}\right) \left(\sqrt{15} \cos\left(\frac{\sqrt{15}}{2}\right) - 135 \sin\left(\frac{\sqrt{15}}{2}\right)\right)\right) e^{\frac{x}{2} - \frac{1}{2}}}{80} + \frac{x}{4} + \frac{1}{16}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.344 (sec). Leaf size: 73

```
dsolve([diff(y(x),x$2)-diff(y(x),x)+4*y(x)=x,y(1) = 2, D(y)(1) = 1],y(x), singsol=all)
```

$$y(x) = \frac{\left(\left(\sqrt{15} \sin\left(\frac{\sqrt{15}}{2}\right) + 135 \cos\left(\frac{\sqrt{15}}{2}\right)\right) \cos\left(\frac{\sqrt{15}x}{2}\right) - \sin\left(\frac{\sqrt{15}x}{2}\right) \left(\sqrt{15} \cos\left(\frac{\sqrt{15}}{2}\right) - 135 \sin\left(\frac{\sqrt{15}}{2}\right)\right)\right) e^{\frac{x}{2} - \frac{1}{2}}}{80} + \frac{x}{4} + \frac{1}{16}$$

✓ Solution by Mathematica

Time used: 0.036 (sec). Leaf size: 67

```
DSolve[{y'[x]-y'[x]+4*y[x]==x,{y[1]==2,y'[1]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{80} \left(20x - \sqrt{15} e^{\frac{x-1}{2}} \sin\left(\frac{1}{2}\sqrt{15}(x-1)\right) + 135 e^{\frac{x-1}{2}} \cos\left(\frac{1}{2}\sqrt{15}(x-1)\right) + 5 \right)$$

14.11 problem 2(c)

14.11.1 Existence and uniqueness analysis	2404
14.11.2 Solving as second order linear constant coeff ode	2405
14.11.3 Solving using Kovacic algorithm	2409
14.11.4 Maple step by step solution	2415

Internal problem ID [6381]

Internal file name [OUTPUT/5629_Sunday_June_05_2022_03_45_15_PM_77622315/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Problems for Review and Discovery. Drill exercises. Page 105

Problem number: 2(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"kovacic", "second_order_linear_constant_coeff"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' + 2y' + 5y = e^x$$

With initial conditions

$$[y(0) = -1, y'(0) = 1]$$

14.11.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = 2$$

$$q(x) = 5$$

$$F = e^x$$

Hence the ode is

$$y'' + 2y' + 5y = e^x$$

The domain of $p(x) = 2$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = 5$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. The domain of $F = e^x$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

14.11.2 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 2, C = 5, f(x) = e^x$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 2y' + 5y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 2, C = 5$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 2\lambda e^{\lambda x} + 5 e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 2\lambda + 5 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 2, C = 5$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-2}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{2^2 - (4)(1)(5)} \\ &= -1 \pm 2i \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= -1 + 2i \\ \lambda_2 &= -1 - 2i \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= -1 + 2i \\ \lambda_2 &= -1 - 2i \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = -1$ and $\beta = 2$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^{-x} (c_1 \cos(2x) + c_2 \sin(2x))$$

Therefore the homogeneous solution y_h is

$$y_h = e^{-x} (c_1 \cos(2x) + c_2 \sin(2x))$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$e^x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^{-x} \cos(2x), e^{-x} \sin(2x)\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^x$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$8A_1 e^x = e^x$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{8} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{e^x}{8}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (e^{-x}(c_1 \cos(2x) + c_2 \sin(2x))) + \left(\frac{e^x}{8}\right) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = e^{-x}(c_1 \cos(2x) + c_2 \sin(2x)) + \frac{e^x}{8} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = -1$ and $x = 0$ in the above gives

$$-1 = c_1 + \frac{1}{8} \quad (1A)$$

Taking derivative of the solution gives

$$y' = -e^{-x}(c_1 \cos(2x) + c_2 \sin(2x)) + e^{-x}(-2 \sin(2x) c_1 + 2c_2 \cos(2x)) + \frac{e^x}{8}$$

substituting $y' = 1$ and $x = 0$ in the above gives

$$1 = -c_1 + \frac{1}{8} + 2c_2 \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = -\frac{9}{8}$$

$$c_2 = -\frac{1}{8}$$

Substituting these values back in above solution results in

$$y = -\frac{9 e^{-x} \cos(2x)}{8} - \frac{e^{-x} \sin(2x)}{8} + \frac{e^x}{8}$$


Which simplifies to

$$y = \frac{(-9 \cos(2x) - \sin(2x)) e^{-x}}{8} + \frac{e^x}{8}$$


Summary

The solution(s) found are the following

$$y = \frac{(-9 \cos(2x) - \sin(2x)) e^{-x}}{8} + \frac{e^x}{8} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{(-9 \cos(2x) - \sin(2x)) e^{-x}}{8} + \frac{e^x}{8}$$

Verified OK.

14.11.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 2y' + 5y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 2 \tag{3}$$

$$C = 5$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} \tag{5}$$

$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-4}{1} \tag{6}$$

Comparing the above to (5) shows that

$$s = -4$$

$$t = 1$$

Therefore eq. (4) becomes

$$z''(x) = -4z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 352: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -4$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(2x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{2}{1} dx} \\ &= z_1 e^{-x} \\ &= z_1 (e^{-x}) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-x} \cos(2x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{2}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-2x}}{(y_1)^2} dx \\ &= y_1 \left(\frac{\tan(2x)}{2} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-x} \cos(2x)) + c_2 \left(e^{-x} \cos(2x) \left(\frac{\tan(2x)}{2} \right) \right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 2y' + 5y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-x} \cos(2x) + \frac{e^{-x} c_2 \sin(2x)}{2}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$e^x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{-x} \cos(2x), \frac{e^{-x} \sin(2x)}{2} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^x$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$8A_1e^x = e^x$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{8} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{e^x}{8}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 e^{-x} \cos(2x) + \frac{e^{-x} c_2 \sin(2x)}{2} \right) + \left(\frac{e^x}{8} \right) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 e^{-x} \cos(2x) + \frac{e^{-x} c_2 \sin(2x)}{2} + \frac{e^x}{8} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = -1$ and $x = 0$ in the above gives

$$-1 = c_1 + \frac{1}{8} \quad (1A)$$

Taking derivative of the solution gives

$$y' = -c_1 e^{-x} \cos(2x) - 2c_1 e^{-x} \sin(2x) - \frac{e^{-x} c_2 \sin(2x)}{2} + e^{-x} c_2 \cos(2x) + \frac{e^x}{8}$$

substituting $y' = 1$ and $x = 0$ in the above gives

$$1 = -c_1 + \frac{1}{8} + c_2 \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = -\frac{9}{8}$$
$$c_2 = -\frac{1}{4}$$

Substituting these values back in above solution results in

$$y = -\frac{9 e^{-x} \cos(2x)}{8} - \frac{e^{-x} \sin(2x)}{8} + \frac{e^x}{8}$$


Which simplifies to

$$y = \frac{(-9 \cos(2x) - \sin(2x)) e^{-x}}{8} + \frac{e^x}{8}$$


Summary

The solution(s) found are the following

$$y = \frac{(-9 \cos(2x) - \sin(2x)) e^{-x}}{8} + \frac{e^x}{8} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{(-9 \cos(2x) - \sin(2x)) e^{-x}}{8} + \frac{e^x}{8}$$

Verified OK.

14.11.4 Maple step by step solution

Let's solve

$$\left[y'' + 2y' + 5y = e^x, y(0) = -1, y' \Big|_{\{x=0\}} = 1 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 2r + 5 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-2) \pm (\sqrt{-16})}{2}$$

- Roots of the characteristic polynomial

$$r = (-1 - 2I, -1 + 2I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-x} \cos(2x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{-x} \sin(2x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{-x} \cos(2x) + e^{-x} c_2 \sin(2x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = e^x \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-x} \cos(2x) & e^{-x} \sin(2x) \\ -e^{-x} \cos(2x) - 2e^{-x} \sin(2x) & -e^{-x} \sin(2x) + 2e^{-x} \cos(2x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 2e^{-2x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{e^{-x}(\cos(2x)(\int e^{2x} \sin(2x) dx) - \sin(2x)(\int e^{2x} \cos(2x) dx))}{2}$$

- Compute integrals

$$y_p(x) = \frac{e^x}{8}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{-x} \cos(2x) + e^{-x} c_2 \sin(2x) + \frac{e^x}{8}$$

- Check validity of solution $y = c_1 e^{-x} \cos(2x) + e^{-x} c_2 \sin(2x) + \frac{e^x}{8}$

- Use initial condition $y(0) = -1$

$$-1 = c_1 + \frac{1}{8}$$

- Compute derivative of the solution

$$y' = -c_1 e^{-x} \cos(2x) - 2c_1 e^{-x} \sin(2x) - e^{-x} c_2 \sin(2x) + 2e^{-x} c_2 \cos(2x) + \frac{e^x}{8}$$

- Use the initial condition $y' \Big|_{\{x=0\}} = 1$

$$1 = -c_1 + \frac{1}{8} + 2c_2$$

- Solve for c_1 and c_2

$$\left\{ c_1 = -\frac{9}{8}, c_2 = -\frac{1}{8} \right\}$$

- Substitute constant values into general solution and simplify

$$y = \frac{(-9 \cos(2x) - \sin(2x))e^{-x}}{8} + \frac{e^x}{8}$$

- Solution to the IVP

$$y = \frac{(-9 \cos(2x) - \sin(2x))e^{-x}}{8} + \frac{e^x}{8}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 29

```
dsolve([diff(y(x),x$2)+2*diff(y(x),x)+5*y(x)=exp(x),y(0) = -1, D(y)(0) = 1],y(x), singsol=all)
```

$$y(x) = \frac{(-\sin(2x) - 9\cos(2x))e^{-x}}{8} + \frac{e^x}{8}$$

✓ Solution by Mathematica

Time used: 0.108 (sec). Leaf size: 32

```
DSolve[{y'[x]+2*y'[x]+5*y[x]==Exp[x],{y[0]==-1,y'[0]==1}},y[x],x,IncludeSingularSolutions->False]
```

$$y(x) \rightarrow \frac{1}{8}e^{-x}(e^{2x} - \sin(2x) - 9\cos(2x))$$

14.12 problem 2(d)

14.12.1 Existence and uniqueness analysis	2418
14.12.2 Solving as second order linear constant coeff ode	2419
14.12.3 Solving using Kovacic algorithm	2424
14.12.4 Maple step by step solution	2430

Internal problem ID [6382]

Internal file name [OUTPUT/5630_Sunday_June_05_2022_03_45_17_PM_47741063/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Problems for Review and Discovery. Drill exercises. Page 105

Problem number: 2(d).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + 3y' + 4y = \sin(x)$$

With initial conditions

$$\left[y\left(\frac{\pi}{2}\right) = 1, y'\left(\frac{\pi}{2}\right) = -1 \right]$$

14.12.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = 3$$

$$q(x) = 4$$

$$F = \sin(x)$$

Hence the ode is

$$y'' + 3y' + 4y = \sin(x)$$

The domain of $p(x) = 3$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = \frac{\pi}{2}$ is inside this domain. The domain of $q(x) = 4$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = \frac{\pi}{2}$ is also inside this domain. The domain of $F = \sin(x)$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = \frac{\pi}{2}$ is also inside this domain. Hence solution exists and is unique.

14.12.2 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 3, C = 4, f(x) = \sin(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 3y' + 4y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 3, C = 4$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 3\lambda e^{\lambda x} + 4e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 3\lambda + 4 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 3, C = 4$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-3}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{3^2 - (4)(1)(4)} \\ &= -\frac{3}{2} \pm \frac{i\sqrt{7}}{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= -\frac{3}{2} + \frac{i\sqrt{7}}{2} \\ \lambda_2 &= -\frac{3}{2} - \frac{i\sqrt{7}}{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= -\frac{3}{2} + \frac{i\sqrt{7}}{2} \\ \lambda_2 &= -\frac{3}{2} - \frac{i\sqrt{7}}{2} \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = -\frac{3}{2}$ and $\beta = \frac{\sqrt{7}}{2}$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^{-\frac{3x}{2}} \left(c_1 \cos \left(\frac{\sqrt{7}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{7}x}{2} \right) \right)$$

Therefore the homogeneous solution y_h is

$$y_h = e^{-\frac{3x}{2}} \left(c_1 \cos \left(\frac{\sqrt{7}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{7}x}{2} \right) \right)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{-\frac{3x}{2}} \cos\left(\frac{\sqrt{7}x}{2}\right), e^{-\frac{3x}{2}} \sin\left(\frac{\sqrt{7}x}{2}\right) \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(x) + A_2 \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$3A_1 \cos(x) + 3A_2 \sin(x) - 3A_1 \sin(x) + 3A_2 \cos(x) = \sin(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{6}, A_2 = \frac{1}{6} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{\cos(x)}{6} + \frac{\sin(x)}{6}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(e^{-\frac{3x}{2}} \left(c_1 \cos\left(\frac{\sqrt{7}x}{2}\right) + c_2 \sin\left(\frac{\sqrt{7}x}{2}\right) \right) \right) + \left(-\frac{\cos(x)}{6} + \frac{\sin(x)}{6} \right) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = e^{-\frac{3x}{2}} \left(c_1 \cos \left(\frac{\sqrt{7}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{7}x}{2} \right) \right) - \frac{\cos(x)}{6} + \frac{\sin(x)}{6} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = \frac{\pi}{2}$ in the above gives

$$1 = e^{-\frac{3\pi}{4}} \cos \left(\frac{\sqrt{7}\pi}{4} \right) c_1 + e^{-\frac{3\pi}{4}} \sin \left(\frac{\sqrt{7}\pi}{4} \right) c_2 + \frac{1}{6} \quad (1A)$$

Taking derivative of the solution gives

$$y' = -\frac{3e^{-\frac{3x}{2}} \left(c_1 \cos \left(\frac{\sqrt{7}x}{2} \right) + c_2 \sin \left(\frac{\sqrt{7}x}{2} \right) \right)}{2} + e^{-\frac{3x}{2}} \left(-\frac{c_1 \sqrt{7} \sin \left(\frac{\sqrt{7}x}{2} \right)}{2} + \frac{c_2 \sqrt{7} \cos \left(\frac{\sqrt{7}x}{2} \right)}{2} \right) + \frac{\sin(x)}{6}$$

substituting $y' = -1$ and $x = \frac{\pi}{2}$ in the above gives

$$-1 = -\frac{3 \left(-\frac{\sqrt{7}c_2}{3} + c_1 \right) e^{-\frac{3\pi}{4}} \cos \left(\frac{\sqrt{7}\pi}{4} \right)}{2} + \frac{1}{6} - \frac{e^{-\frac{3\pi}{4}} (\sqrt{7}c_1 + 3c_2) \sin \left(\frac{\sqrt{7}\pi}{4} \right)}{2} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = \frac{e^{\frac{3\pi}{4}} \sqrt{7} \left(5\sqrt{7} \cos \left(\frac{\sqrt{7}\pi}{4} \right) - \sin \left(\frac{\sqrt{7}\pi}{4} \right) \right)}{42}$$

$$c_2 = \frac{e^{\frac{3\pi}{4}} \sqrt{7} \left(5\sqrt{7} \sin \left(\frac{\sqrt{7}\pi}{4} \right) + \cos \left(\frac{\sqrt{7}\pi}{4} \right) \right)}{42}$$

Substituting these values back in above solution results in

$$y = \frac{\sin(x)}{6} - \frac{\cos(x)}{6} + \frac{5 \cos \left(\frac{\sqrt{7}x}{2} \right) \cos \left(\frac{\sqrt{7}\pi}{4} \right) e^{-\frac{3x}{2} + \frac{3\pi}{4}}}{6} - \frac{\cos \left(\frac{\sqrt{7}x}{2} \right) \sqrt{7} \sin \left(\frac{\sqrt{7}\pi}{4} \right) e^{-\frac{3x}{2} + \frac{3\pi}{4}}}{42} + \frac{5 \sin \left(\frac{\sqrt{7}x}{2} \right) \sin \left(\frac{\sqrt{7}\pi}{4} \right) e^{-\frac{3x}{2} + \frac{3\pi}{4}}}{42}$$


Which simplifies to

$$y = \frac{\left(\left(\sqrt{7} \sin \left(\frac{\sqrt{7}x}{2} \right) + 35 \cos \left(\frac{\sqrt{7}x}{2} \right) \right) \cos \left(\frac{\sqrt{7}\pi}{4} \right) - \sin \left(\frac{\sqrt{7}\pi}{4} \right) \left(\sqrt{7} \cos \left(\frac{\sqrt{7}x}{2} \right) - 35 \sin \left(\frac{\sqrt{7}x}{2} \right) \right) \right) e^{-\frac{3x}{2} + \frac{3\pi}{4}}}{42} - \frac{\cos(x)}{6} + \frac{\sin(x)}{6}$$


Summary

The solution(s) found are the following

$$y = \frac{\left(\left(\sqrt{7} \sin\left(\frac{\sqrt{7}x}{2}\right) + 35 \cos\left(\frac{\sqrt{7}x}{2}\right) \right) \cos\left(\frac{\sqrt{7}\pi}{4}\right) - \sin\left(\frac{\sqrt{7}\pi}{4}\right) \left(\sqrt{7} \cos\left(\frac{\sqrt{7}x}{2}\right) - 35 \sin\left(\frac{\sqrt{7}x}{2}\right) \right) \right) e^{-\frac{3x}{2} + \frac{3\pi}{4}}}{42} - \frac{\cos(x)}{6} + \frac{\sin(x)}{6} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{\left(\left(\sqrt{7} \sin\left(\frac{\sqrt{7}x}{2}\right) + 35 \cos\left(\frac{\sqrt{7}x}{2}\right) \right) \cos\left(\frac{\sqrt{7}\pi}{4}\right) - \sin\left(\frac{\sqrt{7}\pi}{4}\right) \left(\sqrt{7} \cos\left(\frac{\sqrt{7}x}{2}\right) - 35 \sin\left(\frac{\sqrt{7}x}{2}\right) \right) \right) e^{-\frac{3x}{2} + \frac{3\pi}{4}}}{42} - \frac{\cos(x)}{6} + \frac{\sin(x)}{6}$$

Verified OK.

14.12.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 3y' + 4y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 3 \end{aligned} \quad (3)$$

$$C = 4$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-7}{4} \quad (6)$$

Comparing the above to (5) shows that

$$s = -7$$

$$t = 4$$

Therefore eq. (4) becomes

$$z''(x) = -\frac{7z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 354: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -\frac{7}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos\left(\frac{\sqrt{7}x}{2}\right)$$

Using the above, the solution for the original ode can now be found. The first solution

to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\&= z_1 e^{-\int \frac{1}{2} \frac{3}{1} dx} \\&= z_1 e^{-\frac{3x}{2}} \\&= z_1 \left(e^{-\frac{3x}{2}} \right)\end{aligned}$$

Which simplifies to

$$y_1 = e^{-\frac{3x}{2}} \cos \left(\frac{\sqrt{7} x}{2} \right)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{3}{1} dx}}{(y_1)^2} dx \\&= y_1 \int \frac{e^{-3x}}{(y_1)^2} dx \\&= y_1 \left(\frac{2\sqrt{7} \tan \left(\frac{\sqrt{7} x}{2} \right)}{7} \right)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\&= c_1 \left(e^{-\frac{3x}{2}} \cos \left(\frac{\sqrt{7} x}{2} \right) \right) + c_2 \left(e^{-\frac{3x}{2}} \cos \left(\frac{\sqrt{7} x}{2} \right) \left(\frac{2\sqrt{7} \tan \left(\frac{\sqrt{7} x}{2} \right)}{7} \right) \right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 3y' + 4y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-\frac{3x}{2}} \cos\left(\frac{\sqrt{7}x}{2}\right) + \frac{2c_2 e^{-\frac{3x}{2}} \sqrt{7} \sin\left(\frac{\sqrt{7}x}{2}\right)}{7}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{-\frac{3x}{2}} \cos\left(\frac{\sqrt{7}x}{2}\right), \frac{2 e^{-\frac{3x}{2}} \sqrt{7} \sin\left(\frac{\sqrt{7}x}{2}\right)}{7} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(x) + A_2 \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$3A_1 \cos(x) + 3A_2 \sin(x) - 3A_1 \sin(x) + 3A_2 \cos(x) = \sin(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{6}, A_2 = \frac{1}{6} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{\cos(x)}{6} + \frac{\sin(x)}{6}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 e^{-\frac{3x}{2}} \cos\left(\frac{\sqrt{7}x}{2}\right) + \frac{2c_2 e^{-\frac{3x}{2}} \sqrt{7} \sin\left(\frac{\sqrt{7}x}{2}\right)}{7} \right) + \left(-\frac{\cos(x)}{6} + \frac{\sin(x)}{6} \right) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 e^{-\frac{3x}{2}} \cos\left(\frac{\sqrt{7}x}{2}\right) + \frac{2c_2 e^{-\frac{3x}{2}} \sqrt{7} \sin\left(\frac{\sqrt{7}x}{2}\right)}{7} - \frac{\cos(x)}{6} + \frac{\sin(x)}{6} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = \frac{\pi}{2}$ in the above gives

$$1 = e^{-\frac{3\pi}{4}} \cos\left(\frac{\sqrt{7}\pi}{4}\right) c_1 + \frac{2c_2 e^{-\frac{3\pi}{4}} \sqrt{7} \sin\left(\frac{\sqrt{7}\pi}{4}\right)}{7} + \frac{1}{6} \quad (1A)$$

Taking derivative of the solution gives

$$y' = -\frac{3c_1 e^{-\frac{3x}{2}} \cos\left(\frac{\sqrt{7}x}{2}\right)}{2} - \frac{c_1 e^{-\frac{3x}{2}} \sqrt{7} \sin\left(\frac{\sqrt{7}x}{2}\right)}{2} - \frac{3c_2 e^{-\frac{3x}{2}} \sqrt{7} \sin\left(\frac{\sqrt{7}x}{2}\right)}{7} + c_2 e^{-\frac{3x}{2}} \cos\left(\frac{\sqrt{7}x}{2}\right) + \frac{\sin(x)}{6}$$

substituting $y' = -1$ and $x = \frac{\pi}{2}$ in the above gives

$$-1 = -\frac{3(c_1 - \frac{2c_2}{3}) e^{-\frac{3\pi}{4}} \cos\left(\frac{\sqrt{7}\pi}{4}\right)}{2} + \frac{1}{6} - \frac{\sqrt{7}(c_1 + \frac{6c_2}{7}) e^{-\frac{3\pi}{4}} \sin\left(\frac{\sqrt{7}\pi}{4}\right)}{2} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$\begin{aligned} c_1 &= -\frac{\left(\sqrt{7} \sin\left(\frac{\sqrt{7}\pi}{4}\right) - 35 \cos\left(\frac{\sqrt{7}\pi}{4}\right)\right) e^{\frac{3\pi}{4}}}{42} \\ c_2 &= \frac{\left(5\sqrt{7} \sin\left(\frac{\sqrt{7}\pi}{4}\right) + \cos\left(\frac{\sqrt{7}\pi}{4}\right)\right) e^{\frac{3\pi}{4}}}{12} \end{aligned}$$

Substituting these values back in above solution results in

$$y = \frac{\sin(x)}{6} - \frac{\cos(x)}{6} + \frac{5 \cos\left(\frac{\sqrt{7}x}{2}\right) \cos\left(\frac{\sqrt{7}\pi}{4}\right) e^{-\frac{3x}{2} + \frac{3\pi}{4}}}{6} - \frac{\cos\left(\frac{\sqrt{7}x}{2}\right) \sqrt{7} \sin\left(\frac{\sqrt{7}\pi}{4}\right) e^{-\frac{3x}{2} + \frac{3\pi}{4}}}{42} + \frac{5 \sin\left(\frac{\sqrt{7}x}{2}\right)}{6}$$


Which simplifies to

$$\begin{aligned} y &= \frac{\left(\left(\sqrt{7} \sin\left(\frac{\sqrt{7}x}{2}\right) + 35 \cos\left(\frac{\sqrt{7}x}{2}\right)\right) \cos\left(\frac{\sqrt{7}\pi}{4}\right) - \sin\left(\frac{\sqrt{7}\pi}{4}\right) \left(\sqrt{7} \cos\left(\frac{\sqrt{7}x}{2}\right) - 35 \sin\left(\frac{\sqrt{7}x}{2}\right)\right)\right) e^{-\frac{3x}{2} + \frac{3\pi}{4}}}{42} \\ &\quad - \frac{\cos(x)}{6} + \frac{\sin(x)}{6} \end{aligned}$$


Summary

The solution(s) found are the following

$$\begin{aligned} y &= \frac{\left(\left(\sqrt{7} \sin\left(\frac{\sqrt{7}x}{2}\right) + 35 \cos\left(\frac{\sqrt{7}x}{2}\right)\right) \cos\left(\frac{\sqrt{7}\pi}{4}\right) - \sin\left(\frac{\sqrt{7}\pi}{4}\right) \left(\sqrt{7} \cos\left(\frac{\sqrt{7}x}{2}\right) - 35 \sin\left(\frac{\sqrt{7}x}{2}\right)\right)\right) e^{-\frac{3x}{2} + \frac{3\pi}{4}}}{42} \quad (1) \\ &\quad - \frac{\cos(x)}{6} + \frac{\sin(x)}{6} \end{aligned}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$\begin{aligned} & y \\ &= \frac{\left(\left(\sqrt{7} \sin\left(\frac{\sqrt{7}x}{2}\right) + 35 \cos\left(\frac{\sqrt{7}x}{2}\right) \right) \cos\left(\frac{\sqrt{7}\pi}{4}\right) - \sin\left(\frac{\sqrt{7}\pi}{4}\right) \left(\sqrt{7} \cos\left(\frac{\sqrt{7}x}{2}\right) - 35 \sin\left(\frac{\sqrt{7}x}{2}\right) \right) \right) e^{-\frac{3x}{2} + \frac{3\pi}{4}}}{42} \\ &= -\frac{\cos(x)}{6} + \frac{\sin(x)}{6} \end{aligned}$$

Verified OK.

14.12.4 Maple step by step solution

Let's solve

$$\left[y'' + 3y' + 4y = \sin(x), y\left(\frac{\pi}{2}\right) = 1, y'\Big|_{\{x=\frac{\pi}{2}\}} = -1 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 3r + 4 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-3) \pm (\sqrt{-7})}{2}$$

- Roots of the characteristic polynomial

$$r = \left(-\frac{3}{2} - \frac{i\sqrt{7}}{2}, -\frac{3}{2} + \frac{i\sqrt{7}}{2} \right)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-\frac{3x}{2}} \cos\left(\frac{\sqrt{7}x}{2}\right)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{-\frac{3x}{2}} \sin\left(\frac{\sqrt{7}x}{2}\right)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{-\frac{3x}{2}} \cos\left(\frac{\sqrt{7}x}{2}\right) + e^{-\frac{3x}{2}} c_2 \sin\left(\frac{\sqrt{7}x}{2}\right) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \sin(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-\frac{3x}{2}} \cos\left(\frac{\sqrt{7}x}{2}\right) & e^{-\frac{3x}{2}} \sin\left(\frac{\sqrt{7}x}{2}\right) \\ -\frac{3e^{-\frac{3x}{2}} \cos\left(\frac{\sqrt{7}x}{2}\right)}{2} - \frac{e^{-\frac{3x}{2}} \sqrt{7} \sin\left(\frac{\sqrt{7}x}{2}\right)}{2} & -\frac{3e^{-\frac{3x}{2}} \sin\left(\frac{\sqrt{7}x}{2}\right)}{2} + \frac{e^{-\frac{3x}{2}} \sqrt{7} \cos\left(\frac{\sqrt{7}x}{2}\right)}{2} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = \frac{\sqrt{7}e^{-3x}}{2}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{2e^{-\frac{3x}{2}} \sqrt{7} \left(\cos\left(\frac{\sqrt{7}x}{2}\right) \left(\int \sin(x) e^{\frac{3x}{2}} \sin\left(\frac{\sqrt{7}x}{2}\right) dx \right) - \sin\left(\frac{\sqrt{7}x}{2}\right) \left(\int \sin(x) e^{\frac{3x}{2}} \cos\left(\frac{\sqrt{7}x}{2}\right) dx \right) \right)}{7}$$

- Compute integrals

$$y_p(x) = -\frac{\cos(x)}{6} + \frac{\sin(x)}{6}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{-\frac{3x}{2}} \cos\left(\frac{\sqrt{7}x}{2}\right) + e^{-\frac{3x}{2}} c_2 \sin\left(\frac{\sqrt{7}x}{2}\right) - \frac{\cos(x)}{6} + \frac{\sin(x)}{6}$$

- Check validity of solution $y = c_1 e^{-\frac{3x}{2}} \cos\left(\frac{\sqrt{7}x}{2}\right) + e^{-\frac{3x}{2}} c_2 \sin\left(\frac{\sqrt{7}x}{2}\right) - \frac{\cos(x)}{6} + \frac{\sin(x)}{6}$

- Use initial condition $y\left(\frac{\pi}{2}\right) = 1$

$$1 = e^{-\frac{3\pi}{4}} \cos\left(\frac{\sqrt{7}\pi}{4}\right) c_1 + e^{-\frac{3\pi}{4}} \sin\left(\frac{\sqrt{7}\pi}{4}\right) c_2 + \frac{1}{6}$$

- Compute derivative of the solution

$$y' = -\frac{3c_1 e^{-\frac{3x}{2}} \cos\left(\frac{\sqrt{7}x}{2}\right)}{2} - \frac{c_1 e^{-\frac{3x}{2}} \sqrt{7} \sin\left(\frac{\sqrt{7}x}{2}\right)}{2} - \frac{3e^{-\frac{3x}{2}} c_2 \sin\left(\frac{\sqrt{7}x}{2}\right)}{2} + \frac{e^{-\frac{3x}{2}} c_2 \sqrt{7} \cos\left(\frac{\sqrt{7}x}{2}\right)}{2} + \frac{\sin(x)}{6} + \frac{\cos(x)}{6}$$

- Use the initial condition $y' \Big|_{\{x=\frac{\pi}{2}\}} = -1$

$$-1 = \frac{1}{6} - \frac{c_1 e^{-\frac{3\pi}{4}} \sqrt{7} \sin\left(\frac{\sqrt{7}\pi}{4}\right)}{2} - \frac{3e^{-\frac{3\pi}{4}} \cos\left(\frac{\sqrt{7}\pi}{4}\right) c_1}{2} + \frac{e^{-\frac{3\pi}{4}} c_2 \sqrt{7} \cos\left(\frac{\sqrt{7}\pi}{4}\right)}{2} - \frac{3e^{-\frac{3\pi}{4}} \sin\left(\frac{\sqrt{7}\pi}{4}\right) c_2}{2}$$

- Solve for c_1 and c_2

$$\left\{ c_1 = \frac{\left(5\sqrt{7} \cos\left(\frac{\sqrt{7}\pi}{4}\right) - \sin\left(\frac{\sqrt{7}\pi}{4}\right)\right) \sqrt{7}}{42e^{-\frac{3\pi}{4}} \left(\cos\left(\frac{\sqrt{7}\pi}{4}\right)^2 + \sin\left(\frac{\sqrt{7}\pi}{4}\right)^2\right)}, c_2 = \frac{\sqrt{7} \left(5\sqrt{7} \sin\left(\frac{\sqrt{7}\pi}{4}\right) + \cos\left(\frac{\sqrt{7}\pi}{4}\right)\right)}{42e^{-\frac{3\pi}{4}} \left(\cos\left(\frac{\sqrt{7}\pi}{4}\right)^2 + \sin\left(\frac{\sqrt{7}\pi}{4}\right)^2\right)} \right\}$$

- Substitute constant values into general solution and simplify

$$y = \frac{\left(\left(\sqrt{7} \sin\left(\frac{\sqrt{7}x}{2}\right) + 35 \cos\left(\frac{\sqrt{7}x}{2}\right)\right) \cos\left(\frac{\sqrt{7}\pi}{4}\right) - \sin\left(\frac{\sqrt{7}\pi}{4}\right) \left(\sqrt{7} \cos\left(\frac{\sqrt{7}x}{2}\right) - 35 \sin\left(\frac{\sqrt{7}x}{2}\right)\right)\right) e^{-\frac{3x}{2} + \frac{3\pi}{4}}}{42} - \frac{\cos(x)}{6} + \frac{\sin(x)}{6}$$

- Solution to the IVP

$$y = \frac{\left(\left(\sqrt{7} \sin\left(\frac{\sqrt{7}x}{2}\right) + 35 \cos\left(\frac{\sqrt{7}x}{2}\right)\right) \cos\left(\frac{\sqrt{7}\pi}{4}\right) - \sin\left(\frac{\sqrt{7}\pi}{4}\right) \left(\sqrt{7} \cos\left(\frac{\sqrt{7}x}{2}\right) - 35 \sin\left(\frac{\sqrt{7}x}{2}\right)\right)\right) e^{-\frac{3x}{2} + \frac{3\pi}{4}}}{42} - \frac{\cos(x)}{6} + \frac{\sin(x)}{6}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.547 (sec). Leaf size: 93

```
dsolve([diff(y(x),x$2)+3*diff(y(x),x)+4*y(x)=sin(x),y(1/2*Pi) = 1, D(y)(1/2*Pi) = -1],y(x),
```

$$y(x) = \frac{\left(\left(\sqrt{7} \sin\left(\frac{\sqrt{7}x}{2}\right) + 35 \cos\left(\frac{\sqrt{7}x}{2}\right)\right) \cos\left(\frac{\sqrt{7}\pi}{4}\right) - \sin\left(\frac{\sqrt{7}\pi}{4}\right) \left(\sqrt{7} \cos\left(\frac{\sqrt{7}x}{2}\right) - 35 \sin\left(\frac{\sqrt{7}x}{2}\right)\right)\right) e^{-\frac{3x}{2} + \frac{3\pi}{4}}}{42} - \frac{\cos(x)}{6} + \frac{\sin(x)}{6}$$

✓ Solution by Mathematica

Time used: 1.46 (sec). Leaf size: 79

```
DSolve[{y''[x]+3*y'[x]+4*y[x]==Sin[x],{y[Pi/2]==1,y'[Pi/2]==-1}},y[x],x,IncludeSingularSolut
```

$$y(x) \rightarrow \frac{1}{42} \left(-\sqrt{7} e^{\frac{3}{4}(\pi-2x)} \sin\left(\frac{1}{4}\sqrt{7}(\pi-2x)\right) + 7 \sin(x) + 35 e^{\frac{3}{4}(\pi-2x)} \cos\left(\frac{1}{4}\sqrt{7}(\pi-2x)\right) - 7 \cos(x) \right)$$

14.13 problem 2(e)

14.13.1 Existence and uniqueness analysis	2433
14.13.2 Solving as second order linear constant coeff ode	2434
14.13.3 Solving using Kovacic algorithm	2438
14.13.4 Maple step by step solution	2444

Internal problem ID [6383]

Internal file name [OUTPUT/5631_Sunday_June_05_2022_03_45_20_PM_81601001/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Problems for Review and Discovery. Drill exercises. Page 105

Problem number: 2(e).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"kovacic", "second_order_linear_constant_coeff"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' + y = e^{-x}$$

With initial conditions

$$[y(2) = 0, y'(2) = -2]$$

14.13.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = 0$$

$$q(x) = 1$$

$$F = e^{-x}$$

Hence the ode is

$$y'' + y = e^{-x}$$

The domain of $p(x) = 0$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 2$ is inside this domain. The domain of $q(x) = 1$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 2$ is also inside this domain. The domain of $F = e^{-x}$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 2$ is also inside this domain. Hence solution exists and is unique.

14.13.2 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 1, f(x) = e^{-x}$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +i \\ \lambda_2 &= -i \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= i \\ \lambda_2 &= -i \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (\cos(x) c_1 + c_2 \sin(x))$$

Or

$$y = \cos(x) c_1 + c_2 \sin(x)$$

Therefore the homogeneous solution y_h is

$$y_h = \cos(x) c_1 + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$e^{-x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{-x}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^{-x}$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$2A_1 e^{-x} = e^{-x}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{2} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{e^{-x}}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (\cos(x) c_1 + c_2 \sin(x)) + \left(\frac{e^{-x}}{2} \right) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = \cos(x) c_1 + c_2 \sin(x) + \frac{e^{-x}}{2} \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 2$ in the above gives

$$0 = \cos(2) c_1 + c_2 \sin(2) + \frac{e^{-2}}{2} \quad (1A)$$

Taking derivative of the solution gives

$$y' = -\sin(x) c_1 + c_2 \cos(x) - \frac{e^{-x}}{2}$$

substituting $y' = -2$ and $x = 2$ in the above gives

$$-2 = -\sin(2) c_1 + c_2 \cos(2) - \frac{e^{-2}}{2} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = -\frac{e^{-2} \cos(2)}{2} - \frac{\sin(2) e^{-2}}{2} + 2 \sin(2)$$

$$c_2 = \frac{e^{-2} \cos(2)}{2} - \frac{\sin(2) e^{-2}}{2} - 2 \cos(2)$$

Substituting these values back in above solution results in

$$y = -\frac{\cos(x) e^{-2} \cos(2)}{2} - \frac{\cos(x) \sin(2) e^{-2}}{2} + 2 \cos(x) \sin(2) + \frac{\sin(x) e^{-2} \cos(2)}{2} - \frac{\sin(x) \sin(2) e^{-2}}{2}$$


Which simplifies to

$$y = \frac{e^{-x}}{2} + \frac{((\sin(x) - \cos(x)) \cos(2) - \cos(x) \sin(2) - \sin(x) \sin(2)) e^{-2}}{2} - 2 \sin(x) \cos(2) + 2 \cos(x) \sin(2)$$


Summary

The solution(s) found are the following

$$y = \frac{e^{-x}}{2} + \frac{((\sin(x) - \cos(x)) \cos(2) - \cos(x) \sin(2) - \sin(x) \sin(2)) e^{-2}}{2} - 2 \sin(x) \cos(2) + 2 \cos(x) \sin(2) \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{e^{-x}}{2} + \frac{((\sin(x) - \cos(x)) \cos(2) - \cos(x) \sin(2) - \sin(x) \sin(2)) e^{-2} - 2 \sin(x) \cos(2) + 2 \cos(x) \sin(2))}{2}$$

Verified OK.

14.13.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 0 \tag{3}$$

$$C = 1$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} \tag{5}$$

$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \tag{6}$$

Comparing the above to (5) shows that

$$s = -1$$

$$t = 1$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 356: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0\end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}y_1 &= z_1 \\ &= \cos(x)\end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned}y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x))\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1(\cos(x)) + c_2(\cos(x)(\tan(x)))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = \cos(x) c_1 + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$e^{-x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{-x}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^{-x}$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$2A_1 e^{-x} = e^{-x}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{2} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{e^{-x}}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (\cos(x) c_1 + c_2 \sin(x)) + \left(\frac{e^{-x}}{2} \right) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = \cos(x) c_1 + c_2 \sin(x) + \frac{e^{-x}}{2} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 2$ in the above gives

$$0 = \cos(2) c_1 + c_2 \sin(2) + \frac{e^{-2}}{2} \quad (1A)$$

Taking derivative of the solution gives

$$y' = -\sin(x) c_1 + c_2 \cos(x) - \frac{e^{-x}}{2}$$

substituting $y' = -2$ and $x = 2$ in the above gives

$$-2 = -\sin(2) c_1 + c_2 \cos(2) - \frac{e^{-2}}{2} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$\begin{aligned} c_1 &= -\frac{e^{-2} \cos(2)}{2} - \frac{\sin(2) e^{-2}}{2} + 2 \sin(2) \\ c_2 &= \frac{e^{-2} \cos(2)}{2} - \frac{\sin(2) e^{-2}}{2} - 2 \cos(2) \end{aligned}$$

Substituting these values back in above solution results in

$$y = -\frac{\cos(x) e^{-2} \cos(2)}{2} - \frac{\cos(x) \sin(2) e^{-2}}{2} + 2 \cos(x) \sin(2) + \frac{\sin(x) e^{-2} \cos(2)}{2} - \frac{\sin(x) \sin(2) e^{-2}}{2}$$


Which simplifies to

$$y = \frac{e^{-x}}{2} + \frac{((\sin(x) - \cos(x)) \cos(2) - \cos(x) \sin(2) - \sin(x) \sin(2)) e^{-2}}{-2 \sin(x) \cos(2) + 2 \cos(x) \sin(2)}$$


Summary

The solution(s) found are the following

$$y = \frac{e^{-x}}{2} + \frac{((\sin(x) - \cos(x)) \cos(2) - \cos(x) \sin(2) - \sin(x) \sin(2)) e^{-2}}{-2 \sin(x) \cos(2) + 2 \cos(x) \sin(2)} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{e^{-x}}{2} + \frac{((\sin(x) - \cos(x)) \cos(2) - \cos(x) \sin(2) - \sin(x) \sin(2)) e^{-2}}{-2 \sin(x) \cos(2) + 2 \cos(x) \sin(2)}$$

Verified OK.

14.13.4 Maple step by step solution

Let's solve

$$\left[y'' + y = e^{-x}, y(2) = 0, y'|_{\{x=2\}} = -2 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (-I, I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = \cos(x) c_1 + c_2 \sin(x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = e^{-x} \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\cos(x) \left(\int e^{-x} \sin(x) dx \right) + \sin(x) \left(\int e^{-x} \cos(x) dx \right)$$

- Compute integrals

$$y_p(x) = \frac{e^{-x}}{2}$$

- Substitute particular solution into general solution to ODE

$$y = \cos(x) c_1 + c_2 \sin(x) + \frac{e^{-x}}{2}$$

- Check validity of solution $y = \cos(x) c_1 + c_2 \sin(x) + \frac{e^{-x}}{2}$

- Use initial condition $y(2) = 0$

$$0 = \cos(2) c_1 + c_2 \sin(2) + \frac{e^{-2}}{2}$$

- Compute derivative of the solution

$$y' = -\sin(x) c_1 + c_2 \cos(x) - \frac{e^{-x}}{2}$$

- Use the initial condition $y' \Big|_{\{x=2\}} = -2$

$$-2 = -\sin(2) c_1 + c_2 \cos(2) - \frac{e^{-2}}{2}$$

- Solve for c_1 and c_2

$$\left\{ c_1 = -\frac{e^{-2} \cos(2) + \sin(2) e^{-2} - 4 \sin(2)}{2(\cos(2)^2 + \sin(2)^2)}, c_2 = \frac{e^{-2} \cos(2) - \sin(2) e^{-2} - 4 \cos(2)}{2(\cos(2)^2 + \sin(2)^2)} \right\}$$

- Substitute constant values into general solution and simplify

$$y = \frac{e^{-x}}{2} + \frac{((\sin(x) - \cos(x)) \cos(2) - \cos(x) \sin(2) - \sin(x) \sin(2)) e^{-2}}{2} - 2 \sin(x) \cos(2) + 2 \cos(x) \sin(2)$$

- Solution to the IVP

$$y = \frac{e^{-x}}{2} + \frac{((\sin(x) - \cos(x)) \cos(2) - \cos(x) \sin(2) - \sin(x) \sin(2)) e^{-2}}{2} - 2 \sin(x) \cos(2) + 2 \cos(x) \sin(2)$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.234 (sec). Leaf size: 50

```
dsolve([diff(y(x),x$2)+y(x)=exp(-x),y(2) = 0, D(y)(2) = -2],y(x), singsol=all)
```

$$y(x) = \frac{e^{-x}}{2} + \frac{((- \cos(x) + \sin(x)) \cos(2) - \sin(2) \cos(x) - \sin(2) \sin(x)) e^{-2}}{-2 \cos(2) \sin(x) + 2 \sin(2) \cos(x)}$$

✓ Solution by Mathematica

Time used: 0.062 (sec). Leaf size: 48

```
DSolve[{y'[x]+y[x]==Exp[-x],{y[2]==0,y'[2]==-2}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} e^{-x-2} ((4e^2 - 1) e^x \sin(2 - x) - e^x \cos(2 - x) + e^2)$$

14.14 problem 2(f)

14.14.1 Solving as second order linear constant coeff ode	2447
14.14.2 Solving using Kovacic algorithm	2451
14.14.3 Maple step by step solution	2457

Internal problem ID [6384]

Internal file name [OUTPUT/5632_Sunday_June_05_2022_03_45_22_PM_8971909/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Problems for Review and Discovery. Drill excercises. Page 105

Problem number: 2(f).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' - y = \cos(x)$$

With initial conditions

$$[y(0) = 3, y'(2) = 2]$$

14.14.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = -1, f(x) = \cos(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = -1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = -1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(-1)} \\ &= \pm 1 \end{aligned}$$

Hence

$$\lambda_1 = +1$$

$$\lambda_2 = -1$$

Which simplifies to

$$\lambda_1 = 1$$

$$\lambda_2 = -1$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(1)x} + c_2 e^{(-1)x}$$

Or

$$y = c_1 e^x + c_2 e^{-x}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^x + c_2 e^{-x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\cos(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^x, e^{-x}\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(x) + A_2 \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-2A_1 \cos(x) - 2A_2 \sin(x) = \cos(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{2}, A_2 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{\cos(x)}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^x + c_2 e^{-x}) + \left(-\frac{\cos(x)}{2} \right) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 e^x + c_2 e^{-x} - \frac{\cos(x)}{2} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 3$ and $x = 0$ in the above gives

$$3 = c_1 + c_2 - \frac{1}{2} \quad (1A)$$

Taking derivative of the solution gives

$$y' = c_1 e^x - c_2 e^{-x} + \frac{\sin(x)}{2}$$

substituting $y' = 2$ and $x = 2$ in the above gives

$$2 = c_1 e^2 - c_2 e^{-2} + \frac{\sin(2)}{2} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = -\frac{\sin(2)e^2 - 4e^2 - 7}{2(e^4 + 1)}$$

$$c_2 = \frac{(7e^2 + \sin(2) - 4)e^2}{2e^4 + 2}$$

Substituting these values back in above solution results in

$$y = \frac{-\cos(x)e^8 - 2\cos(x)e^4 + \sin(2)e^{2-x} + \sin(2)e^{-x+6} - \sin(2)e^{x+6} - \sin(2)e^{x+2} - \cos(x) + 7e^{8-x}}{2e^8 + 4e^4 + 2}$$

Which simplifies to

$$y = \frac{(\sin(2) - 4)e^{2-x} + (\sin(2) - 4)e^{-x+6} + 7e^{4-x} + 7e^{8-x} + (4 - \sin(2))e^{x+2} + (4 - \sin(2))e^{x+6} + 7e^{x+6}}{2e^8 + 4e^4 + 2}$$

Summary

The solution(s) found are the following

$$y = \frac{(\sin(2) - 4)e^{2-x} + (\sin(2) - 4)e^{-x+6} + 7e^{4-x} + 7e^{8-x} + (4 - \sin(2))e^{x+2} + (4 - \sin(2))e^{x+6} + 7e^{x+6}}{2e^8 + 4e^4 + 2} \quad (1)$$




Figure 399: Solution plot

Verification of solutions

y

$$= \frac{(\sin(2) - 4)e^{2-x} + (\sin(2) - 4)e^{-x+6} + 7e^{4-x} + 7e^{8-x} + (4 - \sin(2))e^{x+2} + (4 - \sin(2))e^{x+6} + 7e^{x+8}}{2e^8 + 4e^4 + 2}$$

Verified OK.

14.14.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' - y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 0$$

$$C = -1$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} \tag{5}$$

$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{1}{1} \tag{6}$$

Comparing the above to (5) shows that

$$s = 1$$

$$t = 1$$

Therefore eq. (4) becomes

$$z''(x) = z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 358: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-x}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned} y_1 &= z_1 \\ &= e^{-x} \end{aligned}$$

Which simplifies to

$$y_1 = e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= e^{-x} \int \frac{1}{e^{-2x}} dx \\ &= e^{-x} \left(\frac{e^{2x}}{2} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-x}) + c_2 \left(e^{-x} \left(\frac{e^{2x}}{2} \right) \right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-x} + \frac{c_2 e^x}{2}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\cos(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ \frac{e^x}{2}, e^{-x} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(x) + A_2 \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-2A_1 \cos(x) - 2A_2 \sin(x) = \cos(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{2}, A_2 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{\cos(x)}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 e^{-x} + \frac{c_2 e^x}{2} \right) + \left(-\frac{\cos(x)}{2} \right) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 e^{-x} + \frac{c_2 e^x}{2} - \frac{\cos(x)}{2} \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 3$ and $x = 0$ in the above gives

$$3 = c_1 + \frac{c_2}{2} - \frac{1}{2} \tag{1A}$$

Taking derivative of the solution gives

$$y' = -c_1 e^{-x} + \frac{c_2 e^x}{2} + \frac{\sin(x)}{2}$$

substituting $y' = 2$ and $x = 2$ in the above gives

$$2 = -c_1 e^{-2} + \frac{c_2 e^2}{2} + \frac{\sin(2)}{2} \tag{2A}$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = \frac{(7e^2 + \sin(2) - 4)e^2}{2e^4 + 2}$$

$$c_2 = -\frac{\sin(2)e^2 - 4e^2 - 7}{e^4 + 1}$$

Substituting these values back in above solution results in

$$y = \frac{-\cos(x)e^8 - 2\cos(x)e^4 + \sin(2)e^{2-x} + \sin(2)e^{-x+6} - \sin(2)e^{x+6} - \sin(2)e^{x+2} - \cos(x) + 7e^{8-x}}{2e^8 + 4e^4 + 2}$$

Which simplifies to

$$y = \frac{(\sin(2) - 4)e^{2-x} + (\sin(2) - 4)e^{-x+6} + 7e^{4-x} + 7e^{8-x} + (4 - \sin(2))e^{x+2} + (4 - \sin(2))e^{x+6} + 7e^{x+8}}{2e^8 + 4e^4 + 2}$$

Summary

The solution(s) found are the following

$$y = \frac{(\sin(2) - 4)e^{2-x} + (\sin(2) - 4)e^{-x+6} + 7e^{4-x} + 7e^{8-x} + (4 - \sin(2))e^{x+2} + (4 - \sin(2))e^{x+6} + 7e^{x+8}}{2e^8 + 4e^4 + 2} \quad (1)$$




Figure 400: Solution plot

Verification of solutions

$$y = \frac{(\sin(2) - 4)e^{2-x} + (\sin(2) - 4)e^{-x+6} + 7e^{4-x} + 7e^{8-x} + (4 - \sin(2))e^{x+2} + (4 - \sin(2))e^{x+6} + 7e^{x+8}}{2e^8 + 4e^4 + 2}$$

Verified OK.

14.14.3 Maple step by step solution

Let's solve

$$\left[y'' - y = \cos(x), y(0) = 3, y' \Big|_{\{x=2\}} = 2 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - 1 = 0$$

- Factor the characteristic polynomial

$$(r - 1)(r + 1) = 0$$

- Roots of the characteristic polynomial

$$r = (-1, 1)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^x$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{-x} + c_2 e^x + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \cos(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-x} & e^x \\ -e^{-x} & e^x \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 2$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{e^{-x}(\int \cos(x)e^x dx)}{2} + \frac{e^x(\int e^{-x} \cos(x) dx)}{2}$$

- Compute integrals

$$y_p(x) = -\frac{\cos(x)}{2}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{-x} + c_2 e^x - \frac{\cos(x)}{2}$$

- Check validity of solution $y = c_1 e^{-x} + c_2 e^x - \frac{\cos(x)}{2}$

- Use initial condition $y(0) = 3$

$$3 = c_1 + c_2 - \frac{1}{2}$$

- Compute derivative of the solution

$$y' = -c_1 e^{-x} + c_2 e^x + \frac{\sin(x)}{2}$$

- Use the initial condition $y' \Big|_{\{x=2\}} = 2$

$$2 = -c_1 e^{-2} + c_2 e^2 + \frac{\sin(2)}{2}$$

- Solve for c_1 and c_2

$$\left\{ c_1 = \frac{7e^2 + \sin(2) - 4}{2(e^2 + e^{-2})}, c_2 = \frac{7e^{-2} - \sin(2) + 4}{2(e^2 + e^{-2})} \right\}$$

- Substitute constant values into general solution and simplify

$$y = \frac{(\sin(2)-4)e^{2-x} + 7e^{4-x} + (4-\sin(2))e^{x+2} + (-e^4-1)\cos(x) + 7e^x}{2e^4+2}$$

- Solution to the IVP

$$y = \frac{(\sin(2)-4)e^{2-x} + 7e^{4-x} + (4-\sin(2))e^{x+2} + (-e^4-1)\cos(x) + 7e^x}{2e^4+2}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.406 (sec). Leaf size: 54

```
dsolve([diff(y(x),x$2)-y(x)=cos(x),y(0) = 3, D(y)(2) = 2],y(x), singsol=all)
```

$$y(x) = \frac{(\sin(2) - 4)e^{2-x} + 7e^{4-x} + (-\sin(2) + 4)e^{x+2} + (-e^4 - 1)\cos(x) + 7e^x}{2e^4 + 2}$$

✓ Solution by Mathematica

Time used: 0.038 (sec). Leaf size: 67

```
DSolve[{y'[x]-y[x]==Cos[x],{y[0]==3,y'[2]==2}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{-x}(7e^{2x} - e^{2x+2}(\sin(2) - 4) + (1 + e^4)(-e^x)\cos(x) + 7e^4 + e^2(\sin(2) - 4))}{2(1 + e^4)}$$

14.15 problem 2(g)

14.15.1 Existence and uniqueness analysis	2461
14.15.2 Solving as second order ode quadrature ode	2461
14.15.3 Solving as second order linear constant coeff ode	2463
14.15.4 Solving as second order integrable as is ode	2467
14.15.5 Solving as second order ode missing y ode	2468
14.15.6 Solving using Kovacic algorithm	2470
14.15.7 Solving as exact linear second order ode ode	2476
14.15.8 Maple step by step solution	2478

Internal problem ID [6385]

Internal file name [OUTPUT/5633_Sunday_June_05_2022_03_45_24_PM_52149414/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Problems for Review and Discovery. Drill excercises. Page 105

Problem number: 2(g).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "exact linear second order ode", "second_order_integrable_as_is", "second_order_ode_missing_y", "second_order_ode_quadrature", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _quadrature]]
```

$$y'' = \tan(x)$$

With initial conditions

$$[y(1) = 1, y'(1) = -1]$$

14.15.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$\begin{aligned}p(x) &= 0 \\q(x) &= 0 \\F &= \tan(x)\end{aligned}$$

Hence the ode is

$$y'' = \tan(x)$$

The domain of $p(x) = 0$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 1$ is inside this domain. The domain of $F = \tan(x)$ is

$$\left\{ x < \frac{1}{2}\pi + \pi \vee \frac{1}{2}\pi + \pi < x \right\}$$

And the point $x_0 = 1$ is also inside this domain. Hence solution exists and is unique.

14.15.2 Solving as second order ode quadrature ode

Integrating once gives

$$y' = -\ln(\cos(x)) + c_1$$

Integrating again gives

$$y = -\frac{ix^2}{2} - x \ln(\cos(x)) + x \ln(e^{2ix} + 1) - \frac{i \operatorname{polylog}(2, -e^{2ix})}{2} + c_1x + c_2$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = -\frac{ix^2}{2} - x \ln(\cos(x)) + x \ln(e^{2ix} + 1) - \frac{i \operatorname{polylog}(2, -e^{2ix})}{2} + c_1x + c_2 \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 1$ in the above gives

$$1 = -\frac{i}{2} - \ln(\cos(1)) + \ln(e^{2i} + 1) - \frac{i \operatorname{dilog}(e^{2i} + 1)}{2} + c_1 + c_2 \quad (1A)$$

Taking derivative of the solution gives

$$y' = -ix - \ln(\cos(x)) + \frac{x \sin(x)}{\cos(x)} + \frac{2ix e^{2ix}}{e^{2ix} + 1} + c_1$$

substituting $y' = -1$ and $x = 1$ in the above gives

$$-1 = \frac{(i - \ln(\cos(1)) + \tan(1) + c_1) e^{2i} - i + c_1 + \tan(1) - \ln(\cos(1))}{e^{2i} + 1} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = \frac{e^{2i} \ln(\cos(1)) - ie^{2i} - e^{2i} \tan(1) - 1 + i - e^{2i} + \ln(\cos(1)) - \tan(1)}{e^{2i} + 1}$$

$$c_2 = -\frac{-ie^{2i} \operatorname{dilog}(e^{2i} + 1) + 2 \ln(e^{2i} + 1) e^{2i} - 3ie^{2i} - 2e^{2i} \tan(1) - i \operatorname{dilog}(e^{2i} + 1) - 4e^{2i} + 2 \ln(e^{2i} + 1)}{2(e^{2i} + 1)}$$

Substituting these values back in above solution results in

$$y = \frac{4 - 2e^{2i}x - 2x + 2 \tan(1) - i + 4e^{2i} - 2 \ln(e^{2i} + 1) + 2ix + ie^{2i} \operatorname{dilog}(e^{2i} + 1) + 3ie^{2i} + 2e^{2i} \tan(1)}{2(e^{2i} + 1)}$$

Summary

The solution(s) found are the following

$$y = \frac{(-ie^{2i} - i) \operatorname{polylog}(2, -e^{2ix}) + 2x(e^{2i} + 1) \ln(e^{2ix} + 1) + (ie^{2i} + i) \operatorname{dilog}(e^{2i} + 1) + (-2e^{2i} - 2) \ln(e^{2i} + 1)}{2(e^{2i} + 1)} \quad (1)$$

Verification of solutions

$$y = \frac{(-ie^{2i} - i) \operatorname{polylog}(2, -e^{2ix}) + 2x(e^{2i} + 1) \ln(e^{2ix} + 1) + (ie^{2i} + i) \operatorname{dilog}(e^{2i} + 1) + (-2e^{2i} - 2) \ln(e^{2i} + 1)}{2(e^{2i} + 1)}$$

Verified OK.

14.15.3 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 0, f(x) = \tan(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 0$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} = 0 \tag{1}$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 = 0 \tag{2}$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 0$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{(0)^2 - (4)(1)(0)} \\ &= 0 \end{aligned}$$

Hence this is the case of a double root $\lambda_{1,2} = 0$. Therefore the solution is

$$y = c_1 1 + c_2 x \tag{1}$$

Therefore the homogeneous solution y_h is

$$y_h = c_2 x + c_1$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1y_1 + u_2y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned} y_1 &= 1 \\ y_2 &= x \end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} 1 & x \\ \frac{d}{dx}(1) & \frac{d}{dx}(x) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} 1 & x \\ 0 & 1 \end{vmatrix}$$

Therefore

$$W = (1)(1) - (x)(0)$$

Which simplifies to

$$W = 1$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x \tan(x)}{1} dx$$

Which simplifies to

$$u_1 = - \int x \tan(x) dx$$

Hence

$$u_1 = -\frac{i\pi^2}{24} - \frac{ix^2}{2} + x \ln(e^{2ix} + 1) - \frac{i \operatorname{polylog}(2, -e^{2ix})}{2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\tan(x)}{1} dx$$

Which simplifies to

$$u_2 = \int \tan(x) dx$$

Hence

$$u_2 = -\ln(\cos(x))$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{i\pi^2}{24} - \frac{ix^2}{2} + x \ln(e^{2ix} + 1) - \frac{i \operatorname{polylog}(2, -e^{2ix})}{2} - x \ln(\cos(x))$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_2x + c_1) + \left(-\frac{i\pi^2}{24} - \frac{ix^2}{2} + x \ln(e^{2ix} + 1) - \frac{i \operatorname{polylog}(2, -e^{2ix})}{2} - x \ln(\cos(x)) \right) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_2x + c_1 - \frac{i\pi^2}{24} - \frac{ix^2}{2} + x \ln(e^{2ix} + 1) - \frac{i \operatorname{polylog}(2, -e^{2ix})}{2} - x \ln(\cos(x)) \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 1$ in the above gives

$$1 = c_2 + c_1 - \frac{i\pi^2}{24} - \frac{i}{2} + \ln(e^{2i} + 1) - \frac{i \operatorname{dilog}(e^{2i} + 1)}{2} - \ln(\cos(1)) \quad (1A)$$

Taking derivative of the solution gives

$$y' = c_2 - ix + \frac{2ix e^{2ix}}{e^{2ix} + 1} - \ln(\cos(x)) + \frac{x \sin(x)}{\cos(x)}$$

substituting $y' = -1$ and $x = 1$ in the above gives

$$-1 = \frac{(i - \ln(\cos(1)) + \tan(1) + c_2) e^{2i} - i + c_2 + \tan(1) - \ln(\cos(1))}{e^{2i} + 1} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = -\frac{-i\pi^2 e^{2i} - 12ie^{2i} \operatorname{dilog}(e^{2i} + 1) - i\pi^2 + 24 \ln(e^{2i} + 1) e^{2i} - 36ie^{2i} - 24 e^{2i} \tan(1) - 12i \operatorname{dilog}(e^{2i} + 1)}{24(e^{2i} + 1)}$$

$$c_2 = \frac{e^{2i} \ln(\cos(1)) - ie^{2i} - e^{2i} \tan(1) - 1 + i - e^{2i} + \ln(\cos(1)) - \tan(1)}{e^{2i} + 1}$$

Substituting these values back in above solution results in

$$y = \frac{4 - 2e^{2i}x - 2x + 2 \tan(1) - i + 4e^{2i} - 2 \ln(e^{2i} + 1) + 2ix + ie^{2i} \operatorname{dilog}(e^{2i} + 1) + 3ie^{2i} + 2e^{2i} \tan(1)}{24(e^{2i} + 1)}$$

Summary

The solution(s) found are the following

$$y = \frac{(-ie^{2i} - i) \operatorname{polylog}(2, -e^{2ix}) + 2x(e^{2i} + 1) \ln(e^{2ix} + 1) + (ie^{2i} + i) \operatorname{dilog}(e^{2i} + 1) + (-2e^{2i} - 2) \ln(e^{2i} + 1)}{24(e^{2i} + 1)} \quad (1)$$

Verification of solutions

$$y = \frac{(-ie^{2i} - i) \operatorname{polylog}(2, -e^{2ix}) + 2x(e^{2i} + 1) \ln(e^{2ix} + 1) + (ie^{2i} + i) \operatorname{dilog}(e^{2i} + 1) + (-2e^{2i} - 2) \ln(e^{2i} + 1)}{24(e^{2i} + 1)}$$

Verified OK.

14.15.4 Solving as second order integrable as is ode

Integrating both sides of the ODE w.r.t x gives

$$\int y' dx = \int \tan(x) dx$$

$$y' = -\ln(\cos(x)) + c_1$$

Which is now solved for y . Integrating both sides gives

$$y = \int -\ln(\cos(x)) + c_1 dx$$

$$= -\frac{ix^2}{2} - x \ln(\cos(x)) + x \ln(e^{2ix} + 1) - \frac{i \operatorname{polylog}(2, -e^{2ix})}{2} + c_1 x + c_2$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = -\frac{ix^2}{2} - x \ln(\cos(x)) + x \ln(e^{2ix} + 1) - \frac{i \operatorname{polylog}(2, -e^{2ix})}{2} + c_1 x + c_2 \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 1$ in the above gives

$$1 = -\frac{i}{2} - \ln(\cos(1)) + \ln(e^{2i} + 1) - \frac{i \operatorname{dilog}(e^{2i} + 1)}{2} + c_1 + c_2 \quad (1A)$$

Taking derivative of the solution gives

$$y' = -ix - \ln(\cos(x)) + \frac{x \sin(x)}{\cos(x)} + \frac{2ix e^{2ix}}{e^{2ix} + 1} + c_1$$

substituting $y' = -1$ and $x = 1$ in the above gives

$$-1 = \frac{(i - \ln(\cos(1)) + \tan(1) + c_1) e^{2i} - i + c_1 + \tan(1) - \ln(\cos(1))}{e^{2i} + 1} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = \frac{e^{2i} \ln(\cos(1)) - ie^{2i} - e^{2i} \tan(1) - 1 + i - e^{2i} + \ln(\cos(1)) - \tan(1)}{e^{2i} + 1}$$

$$c_2 = -\frac{-ie^{2i} \operatorname{dilog}(e^{2i} + 1) + 2 \ln(e^{2i} + 1) e^{2i} - 3ie^{2i} - 2e^{2i} \tan(1) - i \operatorname{dilog}(e^{2i} + 1) - 4e^{2i} + 2 \ln(e^{2i} + 1)}{2(e^{2i} + 1)}$$

Substituting these values back in above solution results in

$$y = \frac{4 - 2e^{2i}x - 2x + 2 \tan(1) - i + 4e^{2i} - 2 \ln(e^{2i} + 1) + 2ix + ie^{2i} \operatorname{dilog}(e^{2i} + 1) + 3ie^{2i} + 2e^{2i} \tan(1)}{2(e^{2i} + 1)}$$

Summary

The solution(s) found are the following

$$y = \frac{(-ie^{2i} - i) \operatorname{polylog}(2, -e^{2ix}) + 2x(e^{2i} + 1) \ln(e^{2ix} + 1) + (ie^{2i} + i) \operatorname{dilog}(e^{2i} + 1) + (-2e^{2i} - 2) \ln(e^{2i} - 2)}{(1)}$$

Verification of solutions

$$y = \frac{(-ie^{2i} - i) \operatorname{polylog}(2, -e^{2ix}) + 2x(e^{2i} + 1) \ln(e^{2ix} + 1) + (ie^{2i} + i) \operatorname{dilog}(e^{2i} + 1) + (-2e^{2i} - 2) \ln(e^{2i} - 2)}{(1)}$$

Verified OK.

14.15.5 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x) - \tan(x) = 0$$

Which is now solve for $p(x)$ as first order ode. Integrating both sides gives

$$\begin{aligned} p(x) &= \int \tan(x) \, dx \\ &= -\ln(\cos(x)) + c_1 \end{aligned}$$

Initial conditions are used to solve for c_1 . Substituting $x = 1$ and $p = -1$ in the above solution gives an equation to solve for the constant of integration.

$$-1 = -\ln(\cos(1)) + c_1$$

$$c_1 = \ln(\cos(1)) - 1$$

Substituting c_1 found above in the general solution gives

$$p(x) = -\ln(\cos(x)) + \ln(\cos(1)) - 1$$

Since $p = y'$ then the new first order ode to solve is

$$y' = -\ln(\cos(x)) + \ln(\cos(1)) - 1$$

Integrating both sides gives

$$\begin{aligned} y &= \int -\ln(\cos(x)) + \ln(\cos(1)) - 1 \, dx \\ &= -x - \frac{ix^2}{2} - x \ln(\cos(x)) + x \ln(e^{2ix} + 1) - \frac{i \operatorname{polylog}(2, -e^{2ix})}{2} + \ln(\cos(1))x + c_2 \end{aligned}$$

Initial conditions are used to solve for c_2 . Substituting $x = 1$ and $y = 1$ in the above solution gives an equation to solve for the constant of integration.

$$1 = -1 - \frac{i}{2} + \ln(e^{2i} + 1) - \frac{i \operatorname{dilog}(e^{2i} + 1)}{2} + c_2$$

$$c_2 = 2 + \frac{i}{2} - \ln(e^{2i} + 1) + \frac{i \operatorname{dilog}(e^{2i} + 1)}{2}$$

Substituting c_2 found above in the general solution gives

$$y = -x - \frac{ix^2}{2} - x \ln(\cos(x)) + x \ln(e^{2ix} + 1) - \frac{i \operatorname{polylog}(2, -e^{2ix})}{2} + \ln(\cos(1))x + 2 + \frac{i}{2} - \ln(e^{2i} + 1) + \frac{i \operatorname{dilog}(e^{2i} + 1)}{2}$$

Initial conditions are used to solve for the constants of integration.

Summary

The solution(s) found are the following

$$\begin{aligned} y &= -x - \frac{ix^2}{2} - x \ln(\cos(x)) + x \ln(e^{2ix} + 1) - \frac{i \operatorname{polylog}(2, -e^{2ix})}{2} \\ &\quad + \ln(\cos(1))x + 2 + \frac{i}{2} - \ln(e^{2i} + 1) + \frac{i \operatorname{dilog}(e^{2i} + 1)}{2} \end{aligned} \tag{1}$$

Verification of solutions

$$\begin{aligned} y &= -x - \frac{ix^2}{2} - x \ln(\cos(x)) + x \ln(e^{2ix} + 1) - \frac{i \operatorname{polylog}(2, -e^{2ix})}{2} \\ &\quad + \ln(\cos(1))x + 2 + \frac{i}{2} - \ln(e^{2i} + 1) + \frac{i \operatorname{dilog}(e^{2i} + 1)}{2} \end{aligned}$$

Verified OK.

14.15.6 Solving using Kovacic algorithm

Writing the ode as

$$y'' = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 0 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 0 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 0 \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 360: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}y_1 &= z_1 \\ &= 1\end{aligned}$$

Which simplifies to

$$y_1 = 1$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned}y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= 1 \int \frac{1}{1} dx \\ &= 1(x)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1(1) + c_2(1(x))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_2 x + c_1$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned} y_1 &= 1 \\ y_2 &= x \end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} 1 & x \\ \frac{d}{dx}(1) & \frac{d}{dx}(x) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} 1 & x \\ 0 & 1 \end{vmatrix}$$

Therefore

$$W = (1)(1) - (x)(0)$$

Which simplifies to

$$W = 1$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x \tan(x)}{1} dx$$

Which simplifies to

$$u_1 = - \int x \tan(x) dx$$

Hence

$$u_1 = -\frac{i\pi^2}{24} - \frac{ix^2}{2} + x \ln(e^{2ix} + 1) - \frac{i \operatorname{polylog}(2, -e^{2ix})}{2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\tan(x)}{1} dx$$

Which simplifies to

$$u_2 = \int \tan(x) dx$$

Hence

$$u_2 = -\ln(\cos(x))$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{i\pi^2}{24} - \frac{ix^2}{2} + x \ln(e^{2ix} + 1) - \frac{i \operatorname{polylog}(2, -e^{2ix})}{2} - x \ln(\cos(x))$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_2x + c_1) + \left(-\frac{i\pi^2}{24} - \frac{ix^2}{2} + x \ln(e^{2ix} + 1) - \frac{i \operatorname{polylog}(2, -e^{2ix})}{2} - x \ln(\cos(x)) \right) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_2x + c_1 - \frac{i\pi^2}{24} - \frac{ix^2}{2} + x \ln(e^{2ix} + 1) - \frac{i \operatorname{polylog}(2, -e^{2ix})}{2} - x \ln(\cos(x)) \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 1$ in the above gives

$$1 = c_2 + c_1 - \frac{i\pi^2}{24} - \frac{i}{2} + \ln(e^{2i} + 1) - \frac{i \operatorname{dilog}(e^{2i} + 1)}{2} - \ln(\cos(1)) \quad (1A)$$

Taking derivative of the solution gives

$$y' = c_2 - ix + \frac{2ix e^{2ix}}{e^{2ix} + 1} - \ln(\cos(x)) + \frac{x \sin(x)}{\cos(x)}$$

substituting $y' = -1$ and $x = 1$ in the above gives

$$-1 = \frac{(i - \ln(\cos(1)) + \tan(1) + c_2) e^{2i} - i + c_2 + \tan(1) - \ln(\cos(1))}{e^{2i} + 1} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = -\frac{-i\pi^2 e^{2i} - 12ie^{2i} \operatorname{dilog}(e^{2i} + 1) - i\pi^2 + 24 \ln(e^{2i} + 1) e^{2i} - 36ie^{2i} - 24 e^{2i} \tan(1) - 12i \operatorname{dilog}(e^{2i} + 1)}{24(e^{2i} + 1)}$$

$$c_2 = \frac{e^{2i} \ln(\cos(1)) - ie^{2i} - e^{2i} \tan(1) - 1 + i - e^{2i} + \ln(\cos(1)) - \tan(1)}{e^{2i} + 1}$$

Substituting these values back in above solution results in

$$y = \frac{4 - 2e^{2i}x - 2x + 2 \tan(1) - i + 4e^{2i} - 2 \ln(e^{2i} + 1) + 2ix + ie^{2i} \operatorname{dilog}(e^{2i} + 1) + 3ie^{2i} + 2e^{2i} \tan(1)}{24(e^{2i} + 1)}$$

Summary

The solution(s) found are the following

$$y = \frac{(-ie^{2i} - i) \operatorname{polylog}(2, -e^{2ix}) + 2x(e^{2i} + 1) \ln(e^{2ix} + 1) + (ie^{2i} + i) \operatorname{dilog}(e^{2i} + 1) + (-2e^{2i} - 2) \ln(e^{2i} + 1)}{24(e^{2i} + 1)} \quad (1)$$

Verification of solutions

$$y = \frac{(-ie^{2i} - i) \operatorname{polylog}(2, -e^{2ix}) + 2x(e^{2i} + 1) \ln(e^{2ix} + 1) + (ie^{2i} + i) \operatorname{dilog}(e^{2i} + 1) + (-2e^{2i} - 2) \ln(e^{2i} + 1)}{24(e^{2i} + 1)}$$

Verified OK.

14.15.7 Solving as exact linear second order ode

An ode of the form

$$p(x)y'' + q(x)y' + r(x)y = s(x)$$

is exact if

$$p''(x) - q'(x) + r(x) = 0 \tag{1}$$

For the given ode we have

$$p(x) = 1$$

$$q(x) = 0$$

$$r(x) = 0$$

$$s(x) = \tan(x)$$

Hence

$$p''(x) = 0$$

$$q'(x) = 0$$

Therefore (1) becomes

$$0 - (0) + (0) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(x)y' + (q(x) - p'(x))y)' = s(x)$$

Integrating gives

$$p(x)y' + (q(x) - p'(x))y = \int s(x) dx$$

Substituting the above values for p, q, r, s gives

$$y' = \int \tan(x) dx$$

We now have a first order ode to solve which is

$$y' = -\ln(\cos(x)) + c_1$$

Integrating both sides gives

$$y = \int -\ln(\cos(x)) + c_1 \, dx$$

$$= -\frac{ix^2}{2} - x \ln(\cos(x)) + x \ln(e^{2ix} + 1) - \frac{i \operatorname{polylog}(2, -e^{2ix})}{2} + c_1 x + c_2$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = -\frac{ix^2}{2} - x \ln(\cos(x)) + x \ln(e^{2ix} + 1) - \frac{i \operatorname{polylog}(2, -e^{2ix})}{2} + c_1 x + c_2 \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 1$ in the above gives

$$1 = -\frac{i}{2} - \ln(\cos(1)) + \ln(e^{2i} + 1) - \frac{i \operatorname{dilog}(e^{2i} + 1)}{2} + c_1 + c_2 \quad (1A)$$

Taking derivative of the solution gives

$$y' = -ix - \ln(\cos(x)) + \frac{x \sin(x)}{\cos(x)} + \frac{2ix e^{2ix}}{e^{2ix} + 1} + c_1$$

substituting $y' = -1$ and $x = 1$ in the above gives

$$-1 = \frac{(i - \ln(\cos(1)) + \tan(1) + c_1) e^{2i} - i + c_1 + \tan(1) - \ln(\cos(1))}{e^{2i} + 1} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = \frac{e^{2i} \ln(\cos(1)) - ie^{2i} - e^{2i} \tan(1) - 1 + i - e^{2i} + \ln(\cos(1)) - \tan(1)}{e^{2i} + 1}$$

$$c_2 = -\frac{-ie^{2i} \operatorname{dilog}(e^{2i} + 1) + 2 \ln(e^{2i} + 1) e^{2i} - 3ie^{2i} - 2e^{2i} \tan(1) - i \operatorname{dilog}(e^{2i} + 1) - 4e^{2i} + 2 \ln(e^{2i} + 1)}{2(e^{2i} + 1)}$$

Substituting these values back in above solution results in

$$y = \frac{4 - 2e^{2i}x - 2x + 2 \tan(1) - i + 4e^{2i} - 2 \ln(e^{2i} + 1) + 2ix + ie^{2i} \operatorname{dilog}(e^{2i} + 1) + 3ie^{2i} + 2e^{2i} \tan(1)}{2(e^{2i} + 1)}$$

Summary

The solution(s) found are the following

$$y = \frac{(-ie^{2i} - i) \operatorname{polylog}(2, -e^{2ix}) + 2x(e^{2i} + 1) \ln(e^{2ix} + 1) + (ie^{2i} + i) \operatorname{dilog}(e^{2i} + 1) + (-2e^{2i} - 2) \ln(e^{2i} + 1)}{2(e^{2i} + 1)} \quad (1)$$

Verification of solutions

$$y = \frac{(-ie^{2i} - i) \operatorname{polylog}(2, -e^{2ix}) + 2x(e^{2i} + 1) \ln(e^{2ix} + 1) + (ie^{2i} + i) \operatorname{dilog}(e^{2i} + 1) + (-2e^{2i} - 2) \ln(e^{2i} - 2)}{1}$$

Verified OK.

14.15.8 Maple step by step solution

Let's solve

$$\left[y'' = \tan(x), y(1) = 1, y' \Big|_{\{x=1\}} = -1 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{0})}{2}$$

- Roots of the characteristic polynomial

$$r = 0$$

- 1st solution of the homogeneous ODE

$$y_1(x) = 1$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = x$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 + c_2 x + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x), y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x), y_2(x))} dx \right), f(x) = \tan(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} 1 & x \\ 0 & 1 \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\left(\int x \tan(x) dx\right) + x\left(\int \tan(x) dx\right)$$

- Compute integrals

$$y_p(x) = -\frac{Ix^2}{2} - x \ln(\cos(x)) + x \ln(e^{2Ix} + 1) - \frac{I \text{polylog}(2, -e^{2Ix})}{2}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 + c_2x - \frac{Ix^2}{2} - x \ln(\cos(x)) + x \ln(e^{2Ix} + 1) - \frac{I \text{polylog}(2, -e^{2Ix})}{2}$$

- Check validity of solution $y = c_1 + c_2x - \frac{Ix^2}{2} - x \ln(\cos(x)) + x \ln(e^{2Ix} + 1) - \frac{I \text{polylog}(2, -e^{2Ix})}{2}$

- Use initial condition $y(1) = 1$

$$1 = c_1 + c_2 - \frac{I}{2} - \ln(\cos(1)) + \ln(e^{2I} + 1) - \frac{I \text{polylog}(2, -e^{2I})}{2}$$

- Compute derivative of the solution

$$y' = c_2 - Ix + \frac{2Ix e^{2Ix}}{e^{2Ix} + 1} - \ln(\cos(x)) + \frac{x \sin(x)}{\cos(x)}$$

- Use the initial condition $y' \Big|_{\{x=1\}} = -1$

$$-1 = c_2 - I + \frac{2I e^{2I}}{e^{2I} + 1} - \ln(\cos(1)) + \frac{\sin(1)}{\cos(1)}$$

- Solve for c_1 and c_2

$$\left\{ c_1 = -\frac{-I \text{polylog}(2, -e^{2I}) \cos(1) e^{2I} + 2 \ln(e^{2I} + 1) e^{2I} \cos(1) - 3 I e^{2I} \cos(1) - I \text{polylog}(2, -e^{2I}) \cos(1) - 2 e^{2I} \sin(1) - 4 e^{2I} \cos(1) + 2}{2 \cos(1) (e^{2I} + 1)} \right.$$

- Substitute constant values into general solution and simplify

$$y = \frac{(-I e^{2I} - I) \text{polylog}(2, -e^{2Ix}) + 2x (e^{2I} + 1) \ln(e^{2Ix} + 1) + (I e^{2I} + I) \text{polylog}(2, -e^{2I}) + (-2 e^{2I} - 2) \ln(e^{2I} + 1) + (2 \ln(\cos(1)))x - 2x \ln(\cos(1))}{2 e^{2I}}$$

- Solution to the IVP

$$y = \frac{(-I e^{2I} - I) \text{polylog}(2, -e^{2Ix}) + 2x (e^{2I} + 1) \ln(e^{2Ix} + 1) + (I e^{2I} + I) \text{polylog}(2, -e^{2I}) + (-2 e^{2I} - 2) \ln(e^{2I} + 1) + (2 \ln(\cos(1)))x - 2x \ln(\cos(1))}{2 e^{2I}}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
<- quadrature successful`
```

✓ Solution by Maple

Time used: 1.984 (sec). Leaf size: 135

```
dsolve([diff(y(x),x$2)=tan(x),y(1) = 1, D(y)(1) = -1],y(x), singsol=all)
```

$$y(x) = \frac{(-ie^{2i} - i) \operatorname{polylog}(2, -e^{2ix}) + 2x(e^{2i} + 1) \ln(e^{2ix} + 1) + (ie^{2i} + i) \operatorname{polylog}(2, -e^{2i}) + (-2e^{2i} - 2) \ln(e^{2i} + 1)}{2}$$

✓ Solution by Mathematica

Time used: 0.038 (sec). Leaf size: 86

```
DSolve[{y'[x]==Tan[x],{y[1]==1,y'[1]==-1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}(-i \operatorname{PolyLog}(2, -e^{2ix}) + i \operatorname{PolyLog}(2, -e^{2i}) - ix^2 - 2x + 2x \log(1 + e^{2ix}) - 2x \log(\cos(x)) + 2x \log(\cos(1)) + (4 + i) - 2 \log(1 + e^{2i}))$$

14.16 problem 2(h)

14.16.1 Existence and uniqueness analysis	2482
14.16.2 Solving as second order linear constant coeff ode	2482
14.16.3 Solving as second order integrable as is ode	2487
14.16.4 Solving as second order ode missing y ode	2490
14.16.5 Solving as type second_order_integrable_as_is (not using ABC version)	2492
14.16.6 Solving using Kovacic algorithm	2495
14.16.7 Solving as exact linear second order ode ode	2502
14.16.8 Maple step by step solution	2505

Internal problem ID [6386]

Internal file name [OUTPUT/5634_Sunday_June_05_2022_03_45_26_PM_96787947/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Problems for Review and Discovery. Drill excercises. Page 105

Problem number: 2(h).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "exact linear second order ode", "second_order_integrable_as_is", "second_order_ode_missing_y", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_y]]
```

$$y'' - 2y' = \ln(x)$$

With initial conditions

$$[y(1) = e, y'(1) = e^{-1}]$$

14.16.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$\begin{aligned}p(x) &= -2 \\q(x) &= 0 \\F &= \ln(x)\end{aligned}$$

Hence the ode is

$$y'' - 2y' = \ln(x)$$

The domain of $p(x) = -2$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 1$ is inside this domain. The domain of $F = \ln(x)$ is

$$\{0 < x\}$$

And the point $x_0 = 1$ is also inside this domain. Hence solution exists and is unique.

14.16.2 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = -2, C = 0, f(x) = \ln(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 2y' = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -2, C = 0$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 2\lambda e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 2\lambda = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -2, C = 0$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{2}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-2^2 - (4)(1)(0)} \\ &= 1 \pm 1 \end{aligned}$$

Hence

$$\lambda_1 = 1 + 1$$

$$\lambda_2 = 1 - 1$$

Which simplifies to

$$\lambda_1 = 2$$

$$\lambda_2 = 0$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(2)x} + c_2 e^{(0)x}$$

Or

$$y = c_1 e^{2x} + c_2$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^{2x} + c_2$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of

parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = e^{2x}$$

$$y_2 = 1$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^{2x} & 1 \\ \frac{d}{dx}(e^{2x}) & \frac{d}{dx}(1) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^{2x} & 1 \\ 2e^{2x} & 0 \end{vmatrix}$$

Therefore

$$W = (e^{2x})(0) - (1)(2e^{2x})$$

Which simplifies to

$$W = -2e^{2x}$$

Which simplifies to

$$W = -2e^{2x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\ln(x)}{-2e^{2x}} dx$$

Which simplifies to

$$u_1 = - \int -\frac{\ln(x) e^{-2x}}{2} dx$$

Hence

$$u_1 = -\frac{\ln(x) e^{-2x}}{4} - \frac{\text{expIntegral}_1(2x)}{4}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\ln(x) e^{2x}}{-2e^{2x}} dx$$

Which simplifies to

$$u_2 = \int -\frac{\ln(x)}{2} dx$$

Hence

$$u_2 = -\frac{\ln(x) x}{2} + \frac{x}{2}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(-\frac{\ln(x) e^{-2x}}{4} - \frac{\text{expIntegral}_1(2x)}{4} \right) e^{2x} - \frac{\ln(x) x}{2} + \frac{x}{2}$$

Which simplifies to

$$y_p(x) = -\frac{e^{2x} \text{expIntegral}_1(2x)}{4} + \frac{(-1 - 2x) \ln(x)}{4} + \frac{x}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^{2x} + c_2) + \left(-\frac{e^{2x} \text{expIntegral}_1(2x)}{4} + \frac{(-1 - 2x) \ln(x)}{4} + \frac{x}{2} \right) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 e^{2x} + c_2 - \frac{e^{2x} \operatorname{expIntegral}_1(2x)}{4} + \frac{(-1 - 2x) \ln(x)}{4} + \frac{x}{2} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = e$ and $x = 1$ in the above gives

$$e = c_1 e^2 + c_2 - \frac{e^2 \operatorname{expIntegral}_1(2)}{4} + \frac{1}{2} \quad (1A)$$

Taking derivative of the solution gives

$$y' = 2c_1 e^{2x} - \frac{e^{2x} \operatorname{expIntegral}_1(2x)}{2} + \frac{e^{2x} e^{-2x}}{4x} - \frac{\ln(x)}{2} + \frac{-1 - 2x}{4x} + \frac{1}{2}$$

substituting $y' = e^{-1}$ and $x = 1$ in the above gives

$$e^{-1} = e^2 \left(2c_1 - \frac{\operatorname{expIntegral}_1(2)}{2} \right) \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = \frac{\operatorname{expIntegral}_1(2)}{4} + \frac{e^{-3}}{2}$$

$$c_2 = e - \frac{1}{2} - \frac{e^{-1}}{2}$$


Substituting these values back in above solution results in

$$y = \frac{e^{2x} \operatorname{expIntegral}_1(2)}{4} + \frac{e^{2x-3}}{2} + e - \frac{1}{2} - \frac{e^{-1}}{2} - \frac{e^{2x} \operatorname{expIntegral}_1(2x)}{4} - \frac{\ln(x)x}{2} - \frac{\ln(x)}{4} + \frac{x}{2}$$


Summary

The solution(s) found are the following

$$y = \frac{e^{2x} \operatorname{expIntegral}_1(2)}{4} + \frac{e^{2x-3}}{2} + e - \frac{1}{2} - \frac{e^{-1}}{2} - \frac{e^{2x} \operatorname{expIntegral}_1(2x)}{4} - \frac{\ln(x)x}{2} - \frac{\ln(x)}{4} + \frac{x}{2} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{e^{2x} \operatorname{ExpIntegralE}_1(2)}{4} + \frac{e^{2x-3}}{2} + e^{-\frac{1}{2}} - \frac{e^{-1}}{2} - \frac{e^{2x} \operatorname{ExpIntegralE}_1(2x)}{4} - \frac{\ln(x)x}{2} - \frac{\ln(x)}{4} + \frac{x}{2}$$

Verified OK.

14.16.3 Solving as second order integrable as is ode

Integrating both sides of the ODE w.r.t x gives

$$\int (y'' - 2y') dx = \int \ln(x) dx$$

$$y' - 2y = \ln(x)x - x + c_1$$

Which is now solved for y .

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -2$$

$$q(x) = \ln(x)x - x + c_1$$

Hence the ode is

$$y' - 2y = \ln(x) x - x + c_1$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int (-2) dx} \\ &= e^{-2x}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu) (\ln(x) x - x + c_1) \\ \frac{d}{dx}(y e^{-2x}) &= (e^{-2x}) (\ln(x) x - x + c_1) \\ d(y e^{-2x}) &= ((\ln(x) x - x + c_1) e^{-2x}) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}y e^{-2x} &= \int (\ln(x) x - x + c_1) e^{-2x} dx \\ y e^{-2x} &= \left(-\frac{1}{4} - \frac{x}{2}\right) e^{-2x} \ln(x) - \frac{c_1 e^{-2x}}{2} + \frac{x e^{-2x}}{2} - \frac{\text{expIntegral}_1(2x)}{4} + c_2\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{-2x}$ results in

$$y = e^{2x} \left(\left(-\frac{1}{4} - \frac{x}{2}\right) e^{-2x} \ln(x) - \frac{c_1 e^{-2x}}{2} + \frac{x e^{-2x}}{2} - \frac{\text{expIntegral}_1(2x)}{4} \right) + c_2 e^{2x}$$

which simplifies to

$$y = -\frac{e^{2x} \text{expIntegral}_1(2x)}{4} + c_2 e^{2x} + \frac{(-1 - 2x) \ln(x)}{4} + \frac{x}{2} - \frac{c_1}{2}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = -\frac{e^{2x} \text{expIntegral}_1(2x)}{4} + c_2 e^{2x} + \frac{(-1 - 2x) \ln(x)}{4} + \frac{x}{2} - \frac{c_1}{2} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = e$ and $x = 1$ in the above gives

$$e = -\frac{e^2 \text{expIntegral}_1(2)}{4} + c_2 e^2 + \frac{1}{2} - \frac{c_1}{2} \quad (1A)$$

Taking derivative of the solution gives

$$y' = -\frac{e^{2x} \operatorname{expIntegral}_1(2x)}{2} + \frac{e^{2x}e^{-2x}}{4x} + 2c_2e^{2x} - \frac{\ln(x)}{2} + \frac{-1-2x}{4x} + \frac{1}{2}$$

substituting $y' = e^{-1}$ and $x = 1$ in the above gives

$$e^{-1} = e^2 \left(-\frac{\operatorname{expIntegral}_1(2)}{2} + 2c_2 \right) \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = -2e + 1 + e^{-1}$$

$$c_2 = \frac{\operatorname{expIntegral}_1(2)}{4} + \frac{e^{-3}}{2}$$


Substituting these values back in above solution results in

$$y = \frac{e^{2x} \operatorname{expIntegral}_1(2)}{4} + \frac{e^{2x-3}}{2} + e - \frac{1}{2} - \frac{e^{-1}}{2} - \frac{e^{2x} \operatorname{expIntegral}_1(2x)}{4} - \frac{\ln(x)x}{2} - \frac{\ln(x)}{4} + \frac{x}{2}$$


Summary

The solution(s) found are the following

$$y = \frac{e^{2x} \operatorname{expIntegral}_1(2)}{4} + \frac{e^{2x-3}}{2} + e - \frac{1}{2} - \frac{e^{-1}}{2} - \frac{e^{2x} \operatorname{expIntegral}_1(2x)}{4} - \frac{\ln(x)x}{2} - \frac{\ln(x)}{4} + \frac{x}{2} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{e^{2x} \operatorname{expIntegral}_1(2)}{4} + \frac{e^{2x-3}}{2} + e - \frac{1}{2} - \frac{e^{-1}}{2} \\ - \frac{e^{2x} \operatorname{expIntegral}_1(2x)}{4} - \frac{\ln(x)x}{2} - \frac{\ln(x)}{4} + \frac{x}{2}$$

Verified OK.

14.16.4 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x) - 2p(x) - \ln(x) = 0$$

Which is now solve for $p(x)$ as first order ode.

Entering Linear first order ODE solver. The integrating factor μ is

$$\mu = e^{\int (-2) dx} \\ = e^{-2x}$$

The ode becomes

$$\frac{d}{dx}(\mu p) = (\mu) (\ln(x)) \\ \frac{d}{dx}(e^{-2x} p) = (e^{-2x}) (\ln(x)) \\ d(e^{-2x} p) = (\ln(x) e^{-2x}) dx$$

Integrating gives

$$e^{-2x} p = \int \ln(x) e^{-2x} dx \\ e^{-2x} p = -\frac{\ln(x) e^{-2x}}{2} - \frac{\operatorname{expIntegral}_1(2x)}{2} + c_1$$

Dividing both sides by the integrating factor $\mu = e^{-2x}$ results in

$$p(x) = e^{2x} \left(-\frac{\ln(x) e^{-2x}}{2} - \frac{\text{expIntegral}_1(2x)}{2} \right) + c_1 e^{2x}$$

which simplifies to

$$p(x) = -\frac{\ln(x)}{2} - \frac{e^{2x} \text{expIntegral}_1(2x)}{2} + c_1 e^{2x}$$

Initial conditions are used to solve for c_1 . Substituting $x = 1$ and $p = e^{-1}$ in the above solution gives an equation to solve for the constant of integration.

$$e^{-1} = -\frac{e^2 \text{expIntegral}_1(2)}{2} + c_1 e^2$$

$$c_1 = \frac{(e^2 \text{expIntegral}_1(2) + 2 e^{-1}) e^{-2}}{2}$$

Substituting c_1 found above in the general solution gives

$$p(x) = \frac{e^{2x} \text{expIntegral}_1(2)}{2} + e^{2x-3} - \frac{e^{2x} \text{expIntegral}_1(2x)}{2} - \frac{\ln(x)}{2}$$

Since $p = y'$ then the new first order ode to solve is

$$y' = \frac{e^{2x} \text{expIntegral}_1(2)}{2} + e^{2x-3} - \frac{e^{2x} \text{expIntegral}_1(2x)}{2} - \frac{\ln(x)}{2}$$

Integrating both sides gives

$$\begin{aligned} y &= \int \left(\frac{e^{2x} \text{expIntegral}_1(2)}{2} + e^{2x-3} - \frac{e^{2x} \text{expIntegral}_1(2x)}{2} - \frac{\ln(x)}{2} \right) dx \\ &= \int \left(\frac{e^{2x} \text{expIntegral}_1(2)}{2} + e^{2x-3} - \frac{e^{2x} \text{expIntegral}_1(2x)}{2} - \frac{\ln(x)}{2} \right) dx + c_2 \end{aligned}$$

Initial conditions are used to solve for c_2 . Substituting $x = 1$ and $y = e$ in the above solution gives an equation to solve for the constant of integration.

$$e = \int^1 \left(\frac{e^{2-a} \text{expIntegral}_1(2)}{2} + e^{2-a-3} - \frac{e^{2-a} \text{expIntegral}_1(2-a)}{2} - \frac{\ln(a)}{2} \right) d_a + c_2$$

$$c_2 = -\frac{\left(\int^1 (-e^{2-a} \text{expIntegral}_1(2-a) + e^{2-a} \text{expIntegral}_1(2) + 2 e^{2-a-3} - \ln(a)) d_a \right)}{2} + e$$

Substituting c_2 found above in the general solution gives

$$y = \int \left(\frac{e^{2x} \operatorname{ExpIntegralE}_1(2)}{2} + e^{2x-3} - \frac{e^{2x} \operatorname{ExpIntegralE}_1(2x)}{2} - \frac{\ln(x)}{2} \right) dx - \frac{\left(\int^1 (-e^{2-a} \operatorname{ExpIntegralE}_1(2-a) \right.$$

Initial conditions are used to solve for the constants of integration.

Summary

The solution(s) found are the following

$$y = \int \left(\frac{e^{2x} \operatorname{ExpIntegralE}_1(2)}{2} + e^{2x-3} - \frac{e^{2x} \operatorname{ExpIntegralE}_1(2x)}{2} - \frac{\ln(x)}{2} \right) dx - \frac{\left(\int^1 (-e^{2-a} \operatorname{ExpIntegralE}_1(2-a) + e^{2-a} \operatorname{ExpIntegralE}_1(2) + 2e^{2-a-3} - \ln(-a)) d_a \right)}{2} + e \quad (1)$$

Verification of solutions

$$y = \int \left(\frac{e^{2x} \operatorname{ExpIntegralE}_1(2)}{2} + e^{2x-3} - \frac{e^{2x} \operatorname{ExpIntegralE}_1(2x)}{2} - \frac{\ln(x)}{2} \right) dx - \frac{\left(\int^1 (-e^{2-a} \operatorname{ExpIntegralE}_1(2-a) + e^{2-a} \operatorname{ExpIntegralE}_1(2) + 2e^{2-a-3} - \ln(-a)) d_a \right)}{2} + e$$

Verified OK.

14.16.5 Solving as type second_order_integrable_as_is (not using ABC version)

Writing the ode as

$$y'' - 2y' = \ln(x)$$

Integrating both sides of the ODE w.r.t x gives

$$\int (y'' - 2y') dx = \int \ln(x) dx$$

$$y' - 2y = \ln(x)x - x + c_1$$

Which is now solved for y .

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$\begin{aligned}p(x) &= -2 \\q(x) &= \ln(x) x - x + c_1\end{aligned}$$

Hence the ode is

$$y' - 2y = \ln(x) x - x + c_1$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int (-2) dx} \\ &= e^{-2x}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu) (\ln(x) x - x + c_1) \\ \frac{d}{dx}(y e^{-2x}) &= (e^{-2x}) (\ln(x) x - x + c_1) \\ d(y e^{-2x}) &= ((\ln(x) x - x + c_1) e^{-2x}) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}y e^{-2x} &= \int (\ln(x) x - x + c_1) e^{-2x} dx \\ y e^{-2x} &= \left(-\frac{1}{4} - \frac{x}{2}\right) e^{-2x} \ln(x) - \frac{c_1 e^{-2x}}{2} + \frac{x e^{-2x}}{2} - \frac{\text{expIntegral}_1(2x)}{4} + c_2\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{-2x}$ results in

$$y = e^{2x} \left(\left(-\frac{1}{4} - \frac{x}{2}\right) e^{-2x} \ln(x) - \frac{c_1 e^{-2x}}{2} + \frac{x e^{-2x}}{2} - \frac{\text{expIntegral}_1(2x)}{4} \right) + c_2 e^{2x}$$

which simplifies to

$$y = -\frac{e^{2x} \text{expIntegral}_1(2x)}{4} + c_2 e^{2x} + \frac{(-1 - 2x) \ln(x)}{4} + \frac{x}{2} - \frac{c_1}{2}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = -\frac{e^{2x} \text{expIntegral}_1(2x)}{4} + c_2 e^{2x} + \frac{(-1 - 2x) \ln(x)}{4} + \frac{x}{2} - \frac{c_1}{2} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = e$ and $x = 1$ in the above gives

$$e = -\frac{e^2 \exp\text{Integral}_1(2)}{4} + c_2 e^2 + \frac{1}{2} - \frac{c_1}{2} \quad (1A)$$

Taking derivative of the solution gives

$$y' = -\frac{e^{2x} \exp\text{Integral}_1(2x)}{2} + \frac{e^{2x} e^{-2x}}{4x} + 2c_2 e^{2x} - \frac{\ln(x)}{2} + \frac{-1 - 2x}{4x} + \frac{1}{2}$$

substituting $y' = e^{-1}$ and $x = 1$ in the above gives

$$e^{-1} = e^2 \left(-\frac{\exp\text{Integral}_1(2)}{2} + 2c_2 \right) \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = -2e + 1 + e^{-1}$$

$$c_2 = \frac{\exp\text{Integral}_1(2)}{4} + \frac{e^{-3}}{2}$$


Substituting these values back in above solution results in

$$y = \frac{e^{2x} \exp\text{Integral}_1(2)}{4} + \frac{e^{2x-3}}{2} + e - \frac{1}{2} - \frac{e^{-1}}{2} - \frac{e^{2x} \exp\text{Integral}_1(2x)}{4} - \frac{\ln(x)x}{2} - \frac{\ln(x)}{4} + \frac{x}{2}$$


Summary

The solution(s) found are the following

$$y = \frac{e^{2x} \exp\text{Integral}_1(2)}{4} + \frac{e^{2x-3}}{2} + e - \frac{1}{2} - \frac{e^{-1}}{2} - \frac{e^{2x} \exp\text{Integral}_1(2x)}{4} - \frac{\ln(x)x}{2} - \frac{\ln(x)}{4} + \frac{x}{2} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{e^{2x} \operatorname{ExpIntegralE}_1(2)}{4} + \frac{e^{2x-3}}{2} + e - \frac{1}{2} - \frac{e^{-1}}{2} - \frac{e^{2x} \operatorname{ExpIntegralE}_1(2x)}{4} - \frac{\ln(x)x}{2} - \frac{\ln(x)}{4} + \frac{x}{2}$$

Verified OK.

14.16.6 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 2y' = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = -2 \tag{3}$$

$$C = 0$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} = \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{1}{1} \tag{6}$$

Comparing the above to (5) shows that

$$s = 1$$

$$t = 1$$

Therefore eq. (4) becomes

$$z''(x) = z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 362: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-x}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-2}{1} dx} \\ &= z_1 e^x \\ &= z_1 (e^x) \end{aligned}$$

Which simplifies to

$$y_1 = 1$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-2}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{2x}}{(y_1)^2} dx \\ &= y_1 \left(\frac{e^{2x}}{2} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1y_1 + c_2y_2 \\ &= c_1(1) + c_2\left(1\left(\frac{e^{2x}}{2}\right)\right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 2y' = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 + \frac{c_2e^{2x}}{2}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1y_1 + u_2y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned}y_1 &= 1 \\ y_2 &= \frac{e^{2x}}{2}\end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} 1 & \frac{e^{2x}}{2} \\ \frac{d}{dx}(1) & \frac{d}{dx}\left(\frac{e^{2x}}{2}\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} 1 & \frac{e^{2x}}{2} \\ 0 & e^{2x} \end{vmatrix}$$

Therefore

$$W = (1)(e^{2x}) - \left(\frac{e^{2x}}{2}\right)(0)$$

Which simplifies to

$$W = e^{2x}$$

Which simplifies to

$$W = e^{2x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\ln(x)e^{2x}}{\frac{e^{2x}}{2}} dx$$

Which simplifies to

$$u_1 = - \int \frac{\ln(x)}{2} dx$$

Hence

$$u_1 = -\frac{\ln(x)x}{2} + \frac{x}{2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\ln(x)}{e^{2x}} dx$$

Which simplifies to

$$u_2 = \int \ln(x) e^{-2x} dx$$

Hence

$$u_2 = -\frac{\ln(x) e^{-2x}}{2} - \frac{\text{expIntegral}_1(2x)}{2}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{\ln(x)x}{2} + \frac{x}{2} + \frac{e^{2x} \left(-\frac{\ln(x)e^{-2x}}{2} - \frac{\text{expIntegral}_1(2x)}{2} \right)}{2}$$

Which simplifies to

$$y_p(x) = -\frac{e^{2x} \text{expIntegral}_1(2x)}{4} + \frac{(-1 - 2x) \ln(x)}{4} + \frac{x}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 + \frac{c_2 e^{2x}}{2} \right) + \left(-\frac{e^{2x} \text{expIntegral}_1(2x)}{4} + \frac{(-1 - 2x) \ln(x)}{4} + \frac{x}{2} \right) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 + \frac{c_2 e^{2x}}{2} - \frac{e^{2x} \text{expIntegral}_1(2x)}{4} + \frac{(-1 - 2x) \ln(x)}{4} + \frac{x}{2} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = e$ and $x = 1$ in the above gives

$$e = c_1 + \frac{c_2 e^2}{2} - \frac{e^2 \text{expIntegral}_1(2)}{4} + \frac{1}{2} \quad (1A)$$

Taking derivative of the solution gives

$$y' = c_2 e^{2x} - \frac{e^{2x} \text{expIntegral}_1(2x)}{2} + \frac{e^{2x} e^{-2x}}{4x} - \frac{\ln(x)}{2} + \frac{-1 - 2x}{4x} + \frac{1}{2}$$

substituting $y' = e^{-1}$ and $x = 1$ in the above gives

$$e^{-1} = e^2 \left(c_2 - \frac{\text{expIntegral}_1(2)}{2} \right) \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = e - \frac{1}{2} - \frac{e^{-1}}{2}$$

$$c_2 = \frac{\text{expIntegral}_1(2)}{2} + e^{-3}$$


Substituting these values back in above solution results in

$$y = \frac{e^{2x} \text{expIntegral}_1(2)}{4} + \frac{e^{2x-3}}{2} + e - \frac{1}{2} - \frac{e^{-1}}{2} - \frac{e^{2x} \text{expIntegral}_1(2x)}{4} - \frac{\ln(x)x}{2} - \frac{\ln(x)}{4} + \frac{x}{2}$$


Summary

The solution(s) found are the following

$$y = \frac{e^{2x} \text{expIntegral}_1(2)}{4} + \frac{e^{2x-3}}{2} + e - \frac{1}{2} - \frac{e^{-1}}{2} - \frac{e^{2x} \text{expIntegral}_1(2x)}{4} - \frac{\ln(x)x}{2} - \frac{\ln(x)}{4} + \frac{x}{2} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{e^{2x} \exp \int_1^x (2) dx}{4} + \frac{e^{2x-3}}{2} + e^{-\frac{1}{2}} - \frac{e^{-1}}{2} - \frac{e^{2x} \exp \int_1^x (2x) dx}{4} - \frac{\ln(x)x}{2} - \frac{\ln(x)}{4} + \frac{x}{2}$$

Verified OK.

14.16.7 Solving as exact linear second order ode

An ode of the form

$$p(x) y'' + q(x) y' + r(x) y = s(x)$$

is exact if

$$p''(x) - q'(x) + r(x) = 0 \tag{1}$$

For the given ode we have

$$\begin{aligned} p(x) &= 1 \\ q(x) &= -2 \\ r(x) &= 0 \\ s(x) &= \ln(x) \end{aligned}$$

Hence

$$\begin{aligned} p''(x) &= 0 \\ q'(x) &= 0 \end{aligned}$$

Therefore (1) becomes

$$0 - (0) + (0) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(x) y' + (q(x) - p'(x)) y)' = s(x)$$

Integrating gives

$$p(x) y' + (q(x) - p'(x)) y = \int s(x) dx$$

Substituting the above values for p, q, r, s gives

$$y' - 2y = \int \ln(x) dx$$

We now have a first order ode to solve which is

$$y' - 2y = \ln(x) x - x + c_1$$

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$\begin{aligned} p(x) &= -2 \\ q(x) &= \ln(x) x - x + c_1 \end{aligned}$$

Hence the ode is

$$y' - 2y = \ln(x) x - x + c_1$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int (-2) dx} \\ &= e^{-2x} \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dx}(\mu y) &= (\mu) (\ln(x) x - x + c_1) \\ \frac{d}{dx}(y e^{-2x}) &= (e^{-2x}) (\ln(x) x - x + c_1) \\ d(y e^{-2x}) &= ((\ln(x) x - x + c_1) e^{-2x}) dx \end{aligned}$$

Integrating gives

$$\begin{aligned} y e^{-2x} &= \int (\ln(x) x - x + c_1) e^{-2x} dx \\ y e^{-2x} &= \left(-\frac{1}{4} - \frac{x}{2}\right) e^{-2x} \ln(x) - \frac{c_1 e^{-2x}}{2} + \frac{x e^{-2x}}{2} - \frac{\text{expIntegral}_1(2x)}{4} + c_2 \end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{-2x}$ results in

$$y = e^{2x} \left(\left(-\frac{1}{4} - \frac{x}{2}\right) e^{-2x} \ln(x) - \frac{c_1 e^{-2x}}{2} + \frac{x e^{-2x}}{2} - \frac{\text{expIntegral}_1(2x)}{4} \right) + c_2 e^{2x}$$

which simplifies to

$$y = -\frac{e^{2x} \operatorname{expIntegral}_1(2x)}{4} + c_2 e^{2x} + \frac{(-1 - 2x) \ln(x)}{4} + \frac{x}{2} - \frac{c_1}{2}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = -\frac{e^{2x} \operatorname{expIntegral}_1(2x)}{4} + c_2 e^{2x} + \frac{(-1 - 2x) \ln(x)}{4} + \frac{x}{2} - \frac{c_1}{2} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = e$ and $x = 1$ in the above gives

$$e = -\frac{e^2 \operatorname{expIntegral}_1(2)}{4} + c_2 e^2 + \frac{1}{2} - \frac{c_1}{2} \quad (1A)$$

Taking derivative of the solution gives

$$y' = -\frac{e^{2x} \operatorname{expIntegral}_1(2x)}{2} + \frac{e^{2x} e^{-2x}}{4x} + 2c_2 e^{2x} - \frac{\ln(x)}{2} + \frac{-1 - 2x}{4x} + \frac{1}{2}$$

substituting $y' = e^{-1}$ and $x = 1$ in the above gives

$$e^{-1} = e^2 \left(-\frac{\operatorname{expIntegral}_1(2)}{2} + 2c_2 \right) \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = -2e + 1 + e^{-1}$$

$$c_2 = \frac{\operatorname{expIntegral}_1(2)}{4} + \frac{e^{-3}}{2}$$


Substituting these values back in above solution results in

$$y = \frac{e^{2x} \operatorname{expIntegral}_1(2)}{4} + \frac{e^{2x-3}}{2} + e - \frac{1}{2} - \frac{e^{-1}}{2} - \frac{e^{2x} \operatorname{expIntegral}_1(2x)}{4} - \frac{\ln(x)x}{2} - \frac{\ln(x)}{4} + \frac{x}{2}$$


Summary

The solution(s) found are the following

$$y = \frac{e^{2x} \operatorname{expIntegral}_1(2)}{4} + \frac{e^{2x-3}}{2} + e - \frac{1}{2} - \frac{e^{-1}}{2} - \frac{e^{2x} \operatorname{expIntegral}_1(2x)}{4} - \frac{\ln(x)x}{2} - \frac{\ln(x)}{4} + \frac{x}{2} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{e^{2x} \operatorname{expIntegral}_1(2)}{4} + \frac{e^{2x-3}}{2} + e - \frac{1}{2} - \frac{e^{-1}}{2} - \frac{e^{2x} \operatorname{expIntegral}_1(2x)}{4} - \frac{\ln(x)x}{2} - \frac{\ln(x)}{4} + \frac{x}{2}$$

Verified OK.

14.16.8 Maple step by step solution

Let's solve

$$\left[y'' - 2y' = \ln(x), y(1) = e, y' \Big|_{\{x=1\}} = \frac{1}{e} \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - 2r = 0$$

- Factor the characteristic polynomial

$$r(r - 2) = 0$$

- Roots of the characteristic polynomial

$$r = (0, 2)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = 1$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{2x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 + c_2 e^{2x} + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \ln(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} 1 & e^{2x} \\ 0 & 2e^{2x} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 2e^{2x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{\int \ln(x) dx}{2} + \frac{e^{2x} \int \ln(x) e^{-2x} dx}{2}$$

- Compute integrals

$$y_p(x) = -\frac{e^{2x} \text{Ei}_1(2x)}{4} + \frac{(-1-2x) \ln(x)}{4} + \frac{x}{2}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 + c_2 e^{2x} - \frac{e^{2x} \text{Ei}_1(2x)}{4} + \frac{(-1-2x) \ln(x)}{4} + \frac{x}{2}$$

- Check validity of solution $y = c_1 + c_2 e^{2x} - \frac{e^{2x} \text{Ei}_1(2x)}{4} + \frac{(-1-2x) \ln(x)}{4} + \frac{x}{2}$

- Use initial condition $y(1) = e$

$$e = c_1 + c_2 e^2 - \frac{e^2 \text{Ei}_1(2)}{4} + \frac{1}{2}$$

- Compute derivative of the solution

$$y' = -\frac{e^{2x} \text{Ei}_1(2x)}{2} + \frac{e^{2x} e^{-2x}}{4x} + 2c_2 e^{2x} - \frac{\ln(x)}{2} + \frac{-1-2x}{4x} + \frac{1}{2}$$

- Use the initial condition $y' \Big|_{\{x=1\}} = \frac{1}{e}$

$$\frac{1}{e} = -\frac{e^2 \text{Ei}_1(2)}{2} + \frac{e^2 e^{-2}}{4} + 2c_2 e^2 - \frac{1}{4}$$

- Solve for c_1 and c_2

$$\left\{ c_1 = \frac{e^2 e^{-2} e + 8(e)^2 - 5e^{-4}}{8e}, c_2 = -\frac{e^2 e^{-2} e - 2e^2 e \text{Ei}_1(2) - e^{-4}}{8e^2 e} \right\}$$

- Substitute constant values into general solution and simplify

$$y = \frac{e^{2x-3}}{2} - \frac{e^{2x} \text{Ei}_1(2x)}{4} + \frac{e^{2x} \text{Ei}_1(2)}{4} + \frac{(-1-2x) \ln(x)}{4} + \frac{x}{2} - \frac{e^{-1}}{2} + e - \frac{1}{2}$$

- Solution to the IVP

$$y = \frac{e^{2x-3}}{2} - \frac{e^{2x} \text{Ei}_1(2x)}{4} + \frac{e^{2x} \text{Ei}_1(2)}{4} + \frac{(-1-2x) \ln(x)}{4} + \frac{x}{2} - \frac{e^{-1}}{2} + e - \frac{1}{2}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
-> Calling odsolve with the ODE`, diff(_b(_a), _a) = 2*_b(_a)+ln(_a), _b(_a)`
  Methods for first order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  trying 1st order linear
  <- 1st order linear successful
<- high order exact linear fully integrable successful`

```

*** Sublevel

✓ Solution by Maple

Time used: 0.171 (sec). Leaf size: 42

```
dsolve([diff(y(x),x$2)-2*diff(y(x),x)=ln(x),y(1) = exp(1), D(y)(1) = 1/exp(1)],y(x), singsol
```

$$y(x) = \frac{\left(\int_1^x (e^{2-z1} \exp \text{Integral}_1(2) - e^{2-z1} \exp \text{Integral}_1(2-z1) + 2e^{2-z1-3} - \ln(z1)) d_z1 \right)}{2} + e$$

✓ Solution by Mathematica

Time used: 0.101 (sec). Leaf size: 66

```
DSolve[{y'[x]-2*y'[x]==Log[x],{y[1]==Exp[1],y'[1]==1/Exp[1]}},y[x],x,IncludeSingularSolutio
```

$$y(x) \rightarrow \frac{1}{4} \left(e^{2x} \text{ExpIntegralEi}(-2x) - \text{ExpIntegralEi}(-2)e^{2x} + 2x + 2e^{2x-3} - 2x \log(x) - \log(-x) + i\pi + 4e - \frac{2}{e} - 2 \right)$$

14.17 problem 3(a)

14.17.1 Solving as second order linear constant coeff ode	2509
14.17.2 Solving using Kovacic algorithm	2512
14.17.3 Maple step by step solution	2517

Internal problem ID [6387]

Internal file name [OUTPUT/5635_Sunday_June_05_2022_03_45_29_PM_41144827/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Problems for Review and Discovery. Drill excercises. Page 105

Problem number: 3(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' + 3y' + 2y = 2x - 1$$

14.17.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 3, C = 2, f(x) = 2x - 1$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 3y' + 2y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 3, C = 2$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 3\lambda e^{\lambda x} + 2e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 3\lambda + 2 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 3, C = 2$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-3}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{3^2 - (4)(1)(2)} \\ &= -\frac{3}{2} \pm \frac{1}{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= -\frac{3}{2} + \frac{1}{2} \\ \lambda_2 &= -\frac{3}{2} - \frac{1}{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= -1 \\ \lambda_2 &= -2 \end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned} y &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ y &= c_1 e^{(-1)x} + c_2 e^{(-2)x} \end{aligned}$$

Or

$$y = c_1 e^{-x} + c_2 e^{-2x}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^{-x} + c_2 e^{-2x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$1 + x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^{-2x}, e^{-x}\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_2x + A_1$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$2A_2x + 2A_1 + 3A_2 = 2x - 1$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = -2, A_2 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -2 + x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1e^{-x} + c_2e^{-2x}) + (-2 + x) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1e^{-x} + c_2e^{-2x} - 2 + x \tag{1}$$




Figure 406: Slope field plot

Verification of solutions

$$y = c_1 e^{-x} + c_2 e^{-2x} - 2 + x$$

Verified OK.

14.17.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 3y' + 2y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 3 \tag{3}$$

$$C = 2$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{1}{4} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 1 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \frac{z(x)}{4} \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 364: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0\end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{1}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{x}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{3}{1} dx} \\ &= z_1 e^{-\frac{3x}{2}} \\ &= z_1 \left(e^{-\frac{3x}{2}} \right)\end{aligned}$$

Which simplifies to

$$y_1 = e^{-2x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{3}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-3x}}{(y_1)^2} dx \\ &= y_1(e^x)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-2x}) + c_2 (e^{-2x}(e^x))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 3y' + 2y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-2x} + c_2 e^{-x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$1 + x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^{-2x}, e^{-x}\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_2 x + A_1$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$2A_2 x + 2A_1 + 3A_2 = 2x - 1$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = -2, A_2 = 1]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -2 + x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^{-2x} + c_2 e^{-x}) + (-2 + x) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-2x} + c_2 e^{-x} - 2 + x \tag{1}$$




Figure 407: Slope field plot

Verification of solutions

$$y = c_1 e^{-2x} + c_2 e^{-x} - 2 + x$$

Verified OK.

14.17.3 Maple step by step solution

Let's solve

$$y'' + 3y' + 2y = 2x - 1$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 3r + 2 = 0$$

- Factor the characteristic polynomial

$$(r + 2)(r + 1) = 0$$

- Roots of the characteristic polynomial

$$r = (-2, -1)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-2x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{-x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{-2x} + c_2 e^{-x} + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = 2x - 1 \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-2x} & e^{-x} \\ -2e^{-2x} & -e^{-x} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = e^{-3x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -e^{-2x} \left(\int (2x - 1) e^{2x} dx \right) + e^{-x} \left(\int (2x - 1) e^x dx \right)$$

- Compute integrals

$$y_p(x) = -2 + x$$

- Substitute particular solution into general solution to ODE

$$y = c_1e^{-2x} + c_2e^{-x} - 2 + x$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
<- double symmetry of the form [xi=0, eta=F(x)] successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve(diff(y(x),x$2)+3*diff(y(x),x)+2*y(x)=2*x-1,y(x), singsol=all)
```

$$y(x) = -e^{-2x}c_1 + c_2e^{-x} + x - 2$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 24

```
DSolve[y''[x]+3*y'[x]+2*y[x]==2*x-1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x + c_1e^{-2x} + c_2e^{-x} - 2$$

14.18 problem 3(b)

14.18.1 Solving as second order linear constant coeff ode	2519
14.18.2 Solving using Kovacic algorithm	2522
14.18.3 Maple step by step solution	2527

Internal problem ID [6388]

Internal file name [OUTPUT/5636_Sunday_June_05_2022_03_45_31_PM_14675524/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Problems for Review and Discovery. Drill excercises. Page 105

Problem number: 3(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - 3y' + 2y = e^{-x}$$

14.18.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = -3, C = 2, f(x) = e^{-x}$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 3y' + 2y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -3, C = 2$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 3\lambda e^{\lambda x} + 2e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 3\lambda + 2 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -3, C = 2$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{3}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-3^2 - (4)(1)(2)} \\ &= \frac{3}{2} \pm \frac{1}{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= \frac{3}{2} + \frac{1}{2} \\ \lambda_2 &= \frac{3}{2} - \frac{1}{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= 2 \\ \lambda_2 &= 1 \end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned} y &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ y &= c_1 e^{(2)x} + c_2 e^{(1)x} \end{aligned}$$

Or

$$y = c_1 e^{2x} + c_2 e^x$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^{2x} + c_2 e^x$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$e^{-x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{-x}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^x, e^{2x}\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^{-x}$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$6A_1 e^{-x} = e^{-x}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{6} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{e^{-x}}{6}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^{2x} + c_2 e^x) + \left(\frac{e^{-x}}{6} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{2x} + c_2 e^x + \frac{e^{-x}}{6} \quad (1)$$




Figure 408: Slope field plot

Verification of solutions

$$y = c_1 e^{2x} + c_2 e^x + \frac{e^{-x}}{6}$$

Verified OK.

14.18.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 3y' + 2y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -3 \\ C &= 2 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{1}{4} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 1 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \frac{z(x)}{4} \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 366: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0\end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{1}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{x}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-3}{1} dx} \\ &= z_1 e^{\frac{3x}{2}} \\ &= z_1 \left(e^{\frac{3x}{2}} \right)\end{aligned}$$

Which simplifies to

$$y_1 = e^x$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{-3}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{3x}}{(y_1)^2} dx \\ &= y_1(e^x)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1(e^x) + c_2(e^x(e^x))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 3y' + 2y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^x + c_2 e^{2x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$e^{-x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{-x}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^x, e^{2x}\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^{-x}$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$6A_1 e^{-x} = e^{-x}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{6} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{e^{-x}}{6}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^x + c_2 e^{2x}) + \left(\frac{e^{-x}}{6} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^x + c_2 e^{2x} + \frac{e^{-x}}{6} \tag{1}$$




Figure 409: Slope field plot

Verification of solutions

$$y = c_1 e^x + c_2 e^{2x} + \frac{e^{-x}}{6}$$

Verified OK.

14.18.3 Maple step by step solution

Let's solve

$$y'' - 3y' + 2y = e^{-x}$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - 3r + 2 = 0$$

- Factor the characteristic polynomial

$$(r - 1)(r - 2) = 0$$

- Roots of the characteristic polynomial

$$r = (1, 2)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^x$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{2x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^x + c_2 e^{2x} + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = e^{-x} \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^x & e^{2x} \\ e^x & 2e^{2x} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = e^{3x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -e^x \left(\int e^{-2x} dx \right) + e^{2x} \left(\int e^{-3x} dx \right)$$

- Compute integrals

$$y_p(x) = \frac{e^{-x}}{6}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^x + c_2 e^{2x} + \frac{e^{-x}}{6}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
<- double symmetry of the form [xi=0, eta=F(x)] successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(diff(y(x),x$2)-3*diff(y(x),x)+2*y(x)=exp(-x),y(x), singsol=all)
```

$$y(x) = e^{2x} c_1 + \frac{e^{-x}}{6} + e^x c_2$$

✓ Solution by Mathematica

Time used: 0.04 (sec). Leaf size: 29

```
DSolve[y''[x]-3*y'[x]+2*y[x]==Exp[-x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{-x}}{6} + c_1 e^x + c_2 e^{2x}$$

14.19 problem 3(c)

14.19.1 Solving as second order linear constant coeff ode	2530
14.19.2 Solving using Kovacic algorithm	2533
14.19.3 Maple step by step solution	2538

Internal problem ID [6389]

Internal file name [OUTPUT/5637_Sunday_June_05_2022_03_45_32_PM_77945810/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Problems for Review and Discovery. Drill excercises. Page 105

Problem number: 3(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' - y' - 2y = \cos(x)$$

14.19.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = -1, C = -2, f(x) = \cos(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - y' - 2y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -1, C = -2$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - \lambda e^{\lambda x} - 2e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - \lambda - 2 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -1, C = -2$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{1}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-1^2 - (4)(1)(-2)} \\ &= \frac{1}{2} \pm \frac{3}{2} \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= \frac{1}{2} + \frac{3}{2} \\ \lambda_2 &= \frac{1}{2} - \frac{3}{2} \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= 2 \\ \lambda_2 &= -1 \end{aligned}$$

Since roots are real and distinct, then the solution is

$$\begin{aligned} y &= c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x} \\ y &= c_1 e^{(2)x} + c_2 e^{(-1)x} \end{aligned}$$

Or

$$y = c_1 e^{2x} + c_2 e^{-x}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^{2x} + c_2 e^{-x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\cos(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^{-x}, e^{2x}\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(x) + A_2 \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-3A_1 \cos(x) - 3A_2 \sin(x) + A_1 \sin(x) - A_2 \cos(x) = \cos(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{3}{10}, A_2 = -\frac{1}{10} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{3 \cos(x)}{10} - \frac{\sin(x)}{10}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^{2x} + c_2 e^{-x}) + \left(-\frac{3 \cos(x)}{10} - \frac{\sin(x)}{10} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{2x} + c_2 e^{-x} - \frac{3 \cos(x)}{10} - \frac{\sin(x)}{10} \quad (1)$$




Figure 410: Slope field plot

Verification of solutions

$$y = c_1 e^{2x} + c_2 e^{-x} - \frac{3 \cos(x)}{10} - \frac{\sin(x)}{10}$$

Verified OK.

14.19.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' - y' - 2y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -1 \\ C &= -2 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{9}{4} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 9 \\ t &= 4 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \frac{9z(x)}{4} \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 368: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = \frac{9}{4}$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-\frac{3x}{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-1}{1} dx} \\ &= z_1 e^{\frac{x}{2}} \\ &= z_1 \left(e^{\frac{x}{2}} \right) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{-1}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^x}{(y_1)^2} dx \\ &= y_1 \left(\frac{e^{3x}}{3} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-x}) + c_2 \left(e^{-x} \left(\frac{e^{3x}}{3} \right) \right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - y' - 2y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-x} + \frac{c_2 e^{2x}}{3}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\cos(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x), \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ \frac{e^{2x}}{3}, e^{-x} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(x) + A_2 \sin(x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-3A_1 \cos(x) - 3A_2 \sin(x) + A_1 \sin(x) - A_2 \cos(x) = \cos(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{3}{10}, A_2 = -\frac{1}{10} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{3 \cos(x)}{10} - \frac{\sin(x)}{10}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 e^{-x} + \frac{c_2 e^{2x}}{3} \right) + \left(-\frac{3 \cos(x)}{10} - \frac{\sin(x)}{10} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-x} + \frac{c_2 e^{2x}}{3} - \frac{3 \cos(x)}{10} - \frac{\sin(x)}{10} \quad (1)$$




Figure 411: Slope field plot

Verification of solutions

$$y = c_1 e^{-x} + \frac{c_2 e^{2x}}{3} - \frac{3 \cos(x)}{10} - \frac{\sin(x)}{10}$$

Verified OK.

14.19.3 Maple step by step solution

Let's solve

$$y'' - y' - 2y = \cos(x)$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - r - 2 = 0$$

- Factor the characteristic polynomial

$$(r + 1)(r - 2) = 0$$

- Roots of the characteristic polynomial

$$r = (-1, 2)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{2x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{-x} + c_2 e^{2x} + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \cos(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-x} & e^{2x} \\ -e^{-x} & 2e^{2x} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 3e^x$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{e^{-x} \left(\int \cos(x) e^x dx \right)}{3} + \frac{e^{2x} \left(\int e^{-2x} \cos(x) dx \right)}{3}$$

- Compute integrals

$$y_p(x) = -\frac{3 \cos(x)}{10} - \frac{\sin(x)}{10}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{-x} + c_2 e^{2x} - \frac{3 \cos(x)}{10} - \frac{\sin(x)}{10}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve(diff(y(x),x$2)-diff(y(x),x)-2*y(x)=cos(x),y(x), singsol=all)
```

$$y(x) = c_2 e^{2x} + c_1 e^{-x} - \frac{3 \cos(x)}{10} - \frac{\sin(x)}{10}$$

✓ Solution by Mathematica

Time used: 0.056 (sec). Leaf size: 34

```
DSolve[y''[x]-y'[x]-2*y[x]==Cos[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\sin(x)}{10} - \frac{3 \cos(x)}{10} + c_1 e^{-x} + c_2 e^{2x}$$

14.20 problem 3(d)

14.20.1 Solving as second order linear constant coeff ode	2541
14.20.2 Solving using Kovacic algorithm	2545
14.20.3 Maple step by step solution	2550

Internal problem ID [6390]

Internal file name [OUTPUT/5638_Sunday_June_05_2022_03_45_34_PM_63092397/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Problems for Review and Discovery. Drill excercises. Page 105

Problem number: 3(d).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + 2y' - y = \sin(x) e^x x$$

14.20.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 2, C = -1, f(x) = \sin(x) e^x x$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 2y' - y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 2, C = -1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 2\lambda e^{\lambda x} - e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 2\lambda - 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 2, C = -1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-2}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{2^2 - (4)(1)(-1)} \\ &= -1 \pm \sqrt{2} \end{aligned}$$

Hence

$$\lambda_1 = -1 + \sqrt{2}$$

$$\lambda_2 = -1 - \sqrt{2}$$

Which simplifies to

$$\lambda_1 = \sqrt{2} - 1$$

$$\lambda_2 = -1 - \sqrt{2}$$

Since roots are real and distinct, then the solution is

$$y = c_1 e^{\lambda_1 x} + c_2 e^{\lambda_2 x}$$

$$y = c_1 e^{(\sqrt{2}-1)x} + c_2 e^{(-1-\sqrt{2})x}$$

Or

$$y = c_1 e^{(\sqrt{2}-1)x} + c_2 e^{(-1-\sqrt{2})x}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^{(\sqrt{2}-1)x} + c_2 e^{(-1-\sqrt{2})x}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x) e^x x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x) e^x, \sin(x) e^x, \cos(x) e^x x, \sin(x) e^x x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{(-1-\sqrt{2})x}, e^{(\sqrt{2}-1)x} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(x) e^x + A_2 \sin(x) e^x + A_3 \cos(x) e^x x + A_4 \sin(x) e^x x$$

The unknowns $\{A_1, A_2, A_3, A_4\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$\begin{aligned} & -4A_1 \sin(x) e^x + 4A_2 \cos(x) e^x - 4A_3 \sin(x) e^x x - 2A_3 \sin(x) e^x \\ & + 4A_3 \cos(x) e^x + 4A_4 \cos(x) e^x x + 2A_4 \cos(x) e^x + 4A_4 \sin(x) e^x \\ & + A_1 \cos(x) e^x + A_2 \sin(x) e^x + A_3 \cos(x) e^x x + A_4 \sin(x) e^x x = \sin(x) e^x x \end{aligned}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{62}{289}, A_2 = \frac{44}{289}, A_3 = -\frac{4}{17}, A_4 = \frac{1}{17} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{62 \cos(x) e^x}{289} + \frac{44 \sin(x) e^x}{289} - \frac{4 \cos(x) e^x x}{17} + \frac{\sin(x) e^x x}{17}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 e^{(\sqrt{2}-1)x} + c_2 e^{(-1-\sqrt{2})x} \right) \\ &\quad + \left(\frac{62 \cos(x) e^x}{289} + \frac{44 \sin(x) e^x}{289} - \frac{4 \cos(x) e^x x}{17} + \frac{\sin(x) e^x x}{17} \right) \end{aligned}$$

Which simplifies to

$$y = c_1 e^{(\sqrt{2}-1)x} + c_2 e^{-(1+\sqrt{2})x} + \frac{62 \cos(x) e^x}{289} + \frac{44 \sin(x) e^x}{289} - \frac{4 \cos(x) e^x x}{17} + \frac{\sin(x) e^x x}{17}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{(\sqrt{2}-1)x} + c_2 e^{-(1+\sqrt{2})x} + \frac{62 \cos(x) e^x}{289} + \frac{44 \sin(x) e^x}{289} - \frac{4 \cos(x) e^x x}{17} + \frac{\sin(x) e^x x}{17} \quad (1)$$




Figure 412: Slope field plot

Verification of solutions

$$y = c_1 e^{(\sqrt{2}-1)x} + c_2 e^{-(1+\sqrt{2})x} + \frac{62 \cos(x) e^x}{289} + \frac{44 \sin(x) e^x}{289} - \frac{4 \cos(x) e^x x}{17} + \frac{\sin(x) e^x x}{17}$$

Verified OK.

14.20.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 2y' - y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 2 \\ C &= -1 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{2}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 2 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 2z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 370: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 2$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = e^{-x\sqrt{2}}$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{2}{1} dx} \end{aligned}$$

$$\begin{aligned}
&= z_1 e^{-x} \\
&= z_1 (e^{-x})
\end{aligned}$$

Which simplifies to

$$y_1 = e^{-(1+\sqrt{2})x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}
y_2 &= y_1 \int \frac{e^{\int -\frac{2}{1} dx}}{(y_1)^2} dx \\
&= y_1 \int \frac{e^{-2x}}{(y_1)^2} dx \\
&= y_1 \left(\frac{\sqrt{2} e^{2x\sqrt{2}}}{4} \right)
\end{aligned}$$

Therefore the solution is

$$\begin{aligned}
y &= c_1 y_1 + c_2 y_2 \\
&= c_1 \left(e^{-(1+\sqrt{2})x} \right) + c_2 \left(e^{-(1+\sqrt{2})x} \left(\frac{\sqrt{2} e^{2x\sqrt{2}}}{4} \right) \right)
\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 2y' - y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-(1+\sqrt{2})x} + \frac{c_2 \sqrt{2} e^{(\sqrt{2}-1)x}}{4}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(x) e^x x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(x) e^x, \sin(x) e^x, \cos(x) e^x x, \sin(x) e^x x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ \frac{\sqrt{2} e^{(\sqrt{2}-1)x}}{4}, e^{-(1+\sqrt{2})x} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(x) e^x + A_2 \sin(x) e^x + A_3 \cos(x) e^x x + A_4 \sin(x) e^x x$$

The unknowns $\{A_1, A_2, A_3, A_4\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$\begin{aligned} & -4A_1 \sin(x) e^x + 4A_2 \cos(x) e^x - 4A_3 \sin(x) e^x x - 2A_3 \sin(x) e^x \\ & + 4A_3 \cos(x) e^x + 4A_4 \cos(x) e^x x + 2A_4 \cos(x) e^x + 4A_4 \sin(x) e^x \\ & + A_1 \cos(x) e^x + A_2 \sin(x) e^x + A_3 \cos(x) e^x x + A_4 \sin(x) e^x x = \sin(x) e^x x \end{aligned}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{62}{289}, A_2 = \frac{44}{289}, A_3 = -\frac{4}{17}, A_4 = \frac{1}{17} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{62 \cos(x) e^x}{289} + \frac{44 \sin(x) e^x}{289} - \frac{4 \cos(x) e^x x}{17} + \frac{\sin(x) e^x x}{17}$$

Therefore the general solution is

$$\begin{aligned}
 y &= y_h + y_p \\
 &= \left(c_1 e^{-(1+\sqrt{2})x} + \frac{c_2 \sqrt{2} e^{(\sqrt{2}-1)x}}{4} \right) \\
 &\quad + \left(\frac{62 \cos(x) e^x}{289} + \frac{44 \sin(x) e^x}{289} - \frac{4 \cos(x) e^x x}{17} + \frac{\sin(x) e^x x}{17} \right)
 \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}
 y &= c_1 e^{-(1+\sqrt{2})x} + \frac{c_2 \sqrt{2} e^{(\sqrt{2}-1)x}}{4} + \frac{62 \cos(x) e^x}{289} \\
 &\quad + \frac{44 \sin(x) e^x}{289} - \frac{4 \cos(x) e^x x}{17} + \frac{\sin(x) e^x x}{17}
 \end{aligned} \tag{1}$$




Figure 413: Slope field plot

Verification of solutions

$$y = c_1 e^{-(1+\sqrt{2})x} + \frac{c_2 \sqrt{2} e^{(\sqrt{2}-1)x}}{4} + \frac{62 \cos(x) e^x}{289} + \frac{44 \sin(x) e^x}{289} - \frac{4 \cos(x) e^x x}{17} + \frac{\sin(x) e^x x}{17}$$

Verified OK.

14.20.3 Maple step by step solution

Let's solve

$$y'' + 2y' - y = \sin(x) e^x x$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 2r - 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-2) \pm (\sqrt{8})}{2}$$

- Roots of the characteristic polynomial

$$r = (-1 - \sqrt{2}, \sqrt{2} - 1)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{(-1-\sqrt{2})x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{(\sqrt{2}-1)x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{(-1-\sqrt{2})x} + c_2 e^{(\sqrt{2}-1)x} + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \sin(x) e^x x \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{(-1-\sqrt{2})x} & e^{(\sqrt{2}-1)x} \\ (-1-\sqrt{2})e^{(-1-\sqrt{2})x} & (\sqrt{2}-1)e^{(\sqrt{2}-1)x} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 2\sqrt{2}e^{-2x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{\sqrt{2} \left(e^{-(1+\sqrt{2})x} \left(\int x e^{(2+\sqrt{2})x} \sin(x) dx \right) - e^{(\sqrt{2}-1)x} \left(\int x e^{-(2+\sqrt{2})x} \sin(x) dx \right) \right)}{4}$$

- Compute integrals

$$y_p(x) = \frac{e^x(17 \sin(x)x - 68 \cos(x)x + 44 \sin(x) + 62 \cos(x))}{289}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{(-1-\sqrt{2})x} + c_2 e^{(\sqrt{2}-1)x} + \frac{e^x(17 \sin(x)x - 68 \cos(x)x + 44 \sin(x) + 62 \cos(x))}{289}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 44

```
dsolve(diff(y(x),x$2)+2*diff(y(x),x)-y(x)=x*exp(x)*sin(x),y(x), singsol=all)
```

$$y(x) = e^{-(1+\sqrt{2})x} c_1 + e^{(\sqrt{2}-1)x} c_2 - \frac{4e^x \left(\left(x - \frac{31}{34} \right) \cos(x) - \frac{\sin(x) \left(x + \frac{44}{17} \right)}{4} \right)}{17}$$

✓ Solution by Mathematica

Time used: 0.04 (sec). Leaf size: 59

```
DSolve[y''[x]+2*y'[x]-y[x]==x*Exp[x]*Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-((1+\sqrt{2})x)} + c_2 e^{(\sqrt{2}-1)x} + \frac{1}{289} e^x ((17x + 44) \sin(x) + (62 - 68x) \cos(x))$$

14.21 problem 3(e)

14.21.1 Solving as second order linear constant coeff ode	2553
14.21.2 Solving using Kovacic algorithm	2558
14.21.3 Maple step by step solution	2565

Internal problem ID [6391]

Internal file name [OUTPUT/5639_Sunday_June_05_2022_03_45_36_PM_95764535/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Problems for Review and Discovery. Drill excercises. Page 105

Problem number: 3(e).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"kovacic", "second_order_linear_constant_coeff"**

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + 9y = \sec(2x)$$

14.21.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 9, f(x) = \sec(2x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 9y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 9$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 9 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 9 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 9$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(9)} \\ &= \pm 3i \end{aligned}$$

Hence

$$\lambda_1 = +3i$$

$$\lambda_2 = -3i$$

Which simplifies to

$$\lambda_1 = 3i$$

$$\lambda_2 = -3i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 3$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (c_1 \cos(3x) + c_2 \sin(3x))$$

Or

$$y = c_1 \cos(3x) + c_2 \sin(3x)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \cos(3x) + c_2 \sin(3x)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \cos(3x)$$

$$y_2 = \sin(3x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(3x) & \sin(3x) \\ \frac{d}{dx}(\cos(3x)) & \frac{d}{dx}(\sin(3x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(3x) & \sin(3x) \\ -3 \sin(3x) & 3 \cos(3x) \end{vmatrix}$$

Therefore

$$W = (\cos(3x))(3 \cos(3x)) - (\sin(3x))(-3 \sin(3x))$$

Which simplifies to

$$W = 3 \cos (3x)^2 + 3 \sin (3x)^2$$

Which simplifies to

$$W = 3$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\sin (3x) \sec (2x)}{3} dx$$

Which simplifies to

$$u_1 = - \int \frac{\sin (3x) \sec (2x)}{3} dx$$

Hence

$$u_1 = - \frac{\sqrt{2} \operatorname{arctanh} (\sqrt{2} \cos (x))}{6} + \frac{2 \cos (x)}{3}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\cos (3x) \sec (2x)}{3} dx$$

Which simplifies to

$$u_2 = \int \frac{\cos (3x) \sec (2x)}{3} dx$$

Hence

$$u_2 = - \frac{\sqrt{2} \operatorname{arctanh} (\sin (x) \sqrt{2})}{6} + \frac{2 \sin (x)}{3}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(- \frac{\sqrt{2} \operatorname{arctanh} (\sqrt{2} \cos (x))}{6} + \frac{2 \cos (x)}{3} \right) \cos (3x) \\ + \left(- \frac{\sqrt{2} \operatorname{arctanh} (\sin (x) \sqrt{2})}{6} + \frac{2 \sin (x)}{3} \right) \sin (3x)$$

Which simplifies to

$$y_p(x) = -\frac{2}{3} + \frac{(-4 \cos(x)^3 + 3 \cos(x)) \sqrt{2} \operatorname{arctanh}(\sqrt{2} \cos(x))}{6} \\ + \frac{\sin(x) (-4 \cos(x)^2 + 1) \sqrt{2} \operatorname{arctanh}(\sin(x) \sqrt{2})}{6} + \frac{4 \cos(x)^2}{3}$$

Therefore the general solution is

$$y = y_h + y_p$$

$$= (c_1 \cos(3x) + c_2 \sin(3x)) + \left(-\frac{2}{3} + \frac{(-4 \cos(x)^3 + 3 \cos(x)) \sqrt{2} \operatorname{arctanh}(\sqrt{2} \cos(x))}{6} \right. \\ \left. + \frac{\sin(x) (-4 \cos(x)^2 + 1) \sqrt{2} \operatorname{arctanh}(\sin(x) \sqrt{2})}{6} + \frac{4 \cos(x)^2}{3} \right)$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(3x) + c_2 \sin(3x) - \frac{2}{3} + \frac{(-4 \cos(x)^3 + 3 \cos(x)) \sqrt{2} \operatorname{arctanh}(\sqrt{2} \cos(x))}{6} \\ + \frac{\sin(x) (-4 \cos(x)^2 + 1) \sqrt{2} \operatorname{arctanh}(\sin(x) \sqrt{2})}{6} + \frac{4 \cos(x)^2}{3} \quad (1)$$




Figure 414: Slope field plot

Verification of solutions

$$y = c_1 \cos(3x) + c_2 \sin(3x) - \frac{2}{3} + \frac{(-4 \cos(x)^3 + 3 \cos(x)) \sqrt{2} \operatorname{arctanh}(\sqrt{2} \cos(x))}{6} + \frac{\sin(x) (-4 \cos(x)^2 + 1) \sqrt{2} \operatorname{arctanh}(\sin(x) \sqrt{2})}{6} + \frac{4 \cos(x)^2}{3}$$

Verified OK.

14.21.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 9y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 9 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-9}{1} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -9 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -9z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 372: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -9$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(3x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}
 y_1 &= z_1 \\
 &= \cos(3x)
 \end{aligned}$$

Which simplifies to

$$y_1 = \cos(3x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(3x) \int \frac{1}{\cos(3x)^2} dx \\ &= \cos(3x) \left(\frac{\tan(3x)}{3} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(3x)) + c_2 \left(\cos(3x) \left(\frac{\tan(3x)}{3} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 9y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 \cos(3x) + \frac{c_2 \sin(3x)}{3}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of

parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \cos(3x)$$

$$y_2 = \frac{\sin(3x)}{3}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(3x) & \frac{\sin(3x)}{3} \\ \frac{d}{dx}(\cos(3x)) & \frac{d}{dx}\left(\frac{\sin(3x)}{3}\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(3x) & \frac{\sin(3x)}{3} \\ -3 \sin(3x) & \cos(3x) \end{vmatrix}$$

Therefore

$$W = (\cos(3x))(\cos(3x)) - \left(\frac{\sin(3x)}{3}\right)(-3 \sin(3x))$$

Which simplifies to

$$W = \cos(3x)^2 + \sin(3x)^2$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\frac{\sin(3x) \sec(2x)}{3}}{1} dx$$

Which simplifies to

$$u_1 = - \int \frac{\sin(3x) \sec(2x)}{3} dx$$

Hence

$$u_1 = - \frac{\sqrt{2} \operatorname{arctanh}(\sqrt{2} \cos(x))}{6} + \frac{2 \cos(x)}{3}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\cos(3x) \sec(2x)}{1} dx$$

Which simplifies to

$$u_2 = \int \cos(3x) \sec(2x) dx$$

Hence

$$u_2 = - \frac{\sqrt{2} \operatorname{arctanh}(\sin(x) \sqrt{2})}{2} + 2 \sin(x)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(- \frac{\sqrt{2} \operatorname{arctanh}(\sqrt{2} \cos(x))}{6} + \frac{2 \cos(x)}{3} \right) \cos(3x) \\ + \frac{\left(- \frac{\sqrt{2} \operatorname{arctanh}(\sin(x) \sqrt{2})}{2} + 2 \sin(x) \right) \sin(3x)}{3}$$

Which simplifies to

$$y_p(x) = -\frac{2}{3} + \frac{(-4 \cos(x)^3 + 3 \cos(x)) \sqrt{2} \operatorname{arctanh}(\sqrt{2} \cos(x))}{6} \\ + \frac{\sin(x) (-4 \cos(x)^2 + 1) \sqrt{2} \operatorname{arctanh}(\sin(x) \sqrt{2})}{6} + \frac{4 \cos(x)^2}{3}$$

Therefore the general solution is

$$y = y_h + y_p \\ = \left(c_1 \cos(3x) + \frac{c_2 \sin(3x)}{3} \right) \\ + \left(-\frac{2}{3} + \frac{(-4 \cos(x)^3 + 3 \cos(x)) \sqrt{2} \operatorname{arctanh}(\sqrt{2} \cos(x))}{6} \right. \\ \left. + \frac{\sin(x) (-4 \cos(x)^2 + 1) \sqrt{2} \operatorname{arctanh}(\sin(x) \sqrt{2})}{6} + \frac{4 \cos(x)^2}{3} \right)$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(3x) + \frac{c_2 \sin(3x)}{3} - \frac{2}{3} + \frac{(-4 \cos(x)^3 + 3 \cos(x)) \sqrt{2} \operatorname{arctanh}(\sqrt{2} \cos(x))}{6} \\ + \frac{\sin(x) (-4 \cos(x)^2 + 1) \sqrt{2} \operatorname{arctanh}(\sin(x) \sqrt{2})}{6} + \frac{4 \cos(x)^2}{3} \quad (1)$$




Figure 415: Slope field plot

Verification of solutions

$$y = c_1 \cos(3x) + \frac{c_2 \sin(3x)}{3} - \frac{2}{3} + \frac{(-4 \cos(x)^3 + 3 \cos(x)) \sqrt{2} \operatorname{arctanh}(\sqrt{2} \cos(x))}{6} + \frac{\sin(x) (-4 \cos(x)^2 + 1) \sqrt{2} \operatorname{arctanh}(\sin(x) \sqrt{2})}{6} + \frac{4 \cos(x)^2}{3}$$

Verified OK.

14.21.3 Maple step by step solution

Let's solve

$$y'' + 9y = \sec(2x)$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 9 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-36})}{2}$$

- Roots of the characteristic polynomial

$$r = (-3I, 3I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(3x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(3x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 \cos(3x) + c_2 \sin(3x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x), y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x), y_2(x))} dx \right), f(x) = \sec(2x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(3x) & \sin(3x) \\ -3 \sin(3x) & 3 \cos(3x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 3$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{\cos(3x) \left(\int \sin(3x) \sec(2x) dx \right)}{3} + \frac{\sin(3x) \left(\int \cos(3x) \sec(2x) dx \right)}{3}$$

- Compute integrals

$$y_p(x) = -\frac{2}{3} + \frac{(-4 \cos(x)^3 + 3 \cos(x)) \sqrt{2} \operatorname{arctanh}(\sqrt{2} \cos(x))}{6} + \frac{\sin(x) (-4 \cos(x)^2 + 1) \sqrt{2} \operatorname{arctanh}(\sin(x) \sqrt{2})}{6} + \frac{4 \cos(x)^2}{3}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 \cos(3x) + c_2 \sin(3x) - \frac{2}{3} + \frac{(-4 \cos(x)^3 + 3 \cos(x)) \sqrt{2} \operatorname{arctanh}(\sqrt{2} \cos(x))}{6} + \frac{\sin(x) (-4 \cos(x)^2 + 1) \sqrt{2} \operatorname{arctanh}(\sin(x) \sqrt{2})}{6}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 81

```
dsolve(diff(y(x),x$2)+9*y(x)=sec(2*x),y(x), singsol=all)
```

$$y(x) = -\frac{2}{3} + \frac{(-4 \cos(x)^3 + 3 \cos(x)) \sqrt{2} \operatorname{arctanh}(\cos(x) \sqrt{2})}{6} \\ + \frac{\sin(x) (1 - 4 \cos(x)^2) \sqrt{2} \operatorname{arctanh}(\sin(x) \sqrt{2})}{6} + 4c_1 \cos(x)^3 \\ + \frac{4(3 \sin(x) c_2 + 1) \cos(x)^2}{3} - 3 \cos(x) c_1 - \sin(x) c_2$$

✓ Solution by Mathematica

Time used: 0.25 (sec). Leaf size: 102

```
DSolve[y''[x]+9*y[x]==Sec[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{6} \left(-\sqrt{2} \sin(3x) \operatorname{arctanh}(\sqrt{2} \sin(x)) - \sqrt{2} \cos(3x) \operatorname{arctanh}(\sqrt{2} - \tan\left(\frac{x}{2}\right)) \right. \\ \left. - \sqrt{2} \cos(3x) \operatorname{arctanh}\left(\tan\left(\frac{x}{2}\right) + \sqrt{2}\right) + 4 \cos(2x) + 6c_1 \cos(3x) + 6c_2 \sin(3x) \right)$$

14.22 problem 3(f)

14.22.1 Solving as second order linear constant coeff ode	2568
14.22.2 Solving as linear second order ode solved by an integrating factor ode	2573
14.22.3 Solving using Kovacic algorithm	2575
14.22.4 Maple step by step solution	2581

Internal problem ID [6392]

Internal file name [OUTPUT/5640_Sunday_June_05_2022_03_45_38_PM_40351328/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Problems for Review and Discovery. Drill excercises. Page 105

Problem number: 3(f).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "linear_second_order_ode_solved_by_an_integrating_factor"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + 4y' + 4y = \ln(x) x$$

14.22.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 4, C = 4, f(x) = \ln(x) x$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 4y' + 4y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 4, C = 4$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 4\lambda e^{\lambda x} + 4e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 4\lambda + 4 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 4, C = 4$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-4}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{(4)^2 - (4)(1)(4)} \\ &= -2 \end{aligned}$$

Hence this is the case of a double root $\lambda_{1,2} = 2$. Therefore the solution is

$$y = c_1 e^{-2x} + c_2 x e^{-2x} \quad (1)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^{-2x} + c_2 x e^{-2x}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned} y_1 &= e^{-2x} \\ y_2 &= x e^{-2x} \end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^{-2x} & x e^{-2x} \\ \frac{d}{dx}(e^{-2x}) & \frac{d}{dx}(x e^{-2x}) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^{-2x} & x e^{-2x} \\ -2e^{-2x} & e^{-2x} - 2x e^{-2x} \end{vmatrix}$$

Therefore

$$W = (e^{-2x})(e^{-2x} - 2x e^{-2x}) - (x e^{-2x})(-2e^{-2x})$$

Which simplifies to

$$W = e^{-4x}$$

Which simplifies to

$$W = e^{-4x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^2 e^{-2x} \ln(x)}{e^{-4x}} dx$$

Which simplifies to

$$u_1 = - \int x^2 \ln(x) e^{2x} dx$$

Hence

$$u_1 = -\left(\frac{1}{4} - \frac{1}{2}x + \frac{1}{2}x^2\right) e^{2x} \ln(x) + \frac{e^{2x}x}{4} - \frac{3e^{2x}}{8} - \frac{\text{expIntegral}_1(-2x)}{4}$$

And Eq. (3) becomes

$$u_2 = \int \frac{e^{-2x} \ln(x) x}{e^{-4x}} dx$$

Which simplifies to

$$u_2 = \int x \ln(x) e^{2x} dx$$

Hence

$$u_2 = \left(-\frac{1}{4} + \frac{x}{2}\right) e^{2x} \ln(x) - \frac{\text{expIntegral}_1(-2x)}{4} - \frac{e^{2x}}{4}$$

Which simplifies to

$$u_1 = -\frac{\text{expIntegral}_1(-2x)}{4} + \frac{((-4x^2 + 4x - 2) \ln(x) + 2x - 3) e^{2x}}{8}$$

$$u_2 = -\frac{\text{expIntegral}_1(-2x)}{4} + \frac{(-1 + (2x - 1) \ln(x)) e^{2x}}{4}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(-\frac{\text{expIntegral}_1(-2x)}{4} + \frac{((-4x^2 + 4x - 2) \ln(x) + 2x - 3) e^{2x}}{8}\right) e^{-2x}$$

$$+ \left(-\frac{\text{expIntegral}_1(-2x)}{4} + \frac{(-1 + (2x - 1) \ln(x)) e^{2x}}{4}\right) x e^{-2x}$$

Which simplifies to

$$y_p(x) = -\frac{e^{-2x}(1+x) \text{expIntegral}_1(-2x)}{4} - \frac{3}{8} + \frac{(2x-2) \ln(x)}{8}$$

Therefore the general solution is

$$y = y_h + y_p$$

$$= (c_1 e^{-2x} + c_2 x e^{-2x}) + \left(-\frac{e^{-2x}(1+x) \text{expIntegral}_1(-2x)}{4} - \frac{3}{8} + \frac{(2x-2) \ln(x)}{8}\right)$$

Which simplifies to

$$y = e^{-2x}(c_2x + c_1) - \frac{e^{-2x}(1+x) \operatorname{expIntegral}_1(-2x)}{4} - \frac{3}{8} + \frac{(2x-2) \ln(x)}{8}$$

Summary

The solution(s) found are the following

$$y = e^{-2x}(c_2x + c_1) - \frac{e^{-2x}(1+x) \operatorname{expIntegral}_1(-2x)}{4} - \frac{3}{8} + \frac{(2x-2) \ln(x)}{8} \quad (1)$$




Figure 416: Slope field plot

Verification of solutions

$$y = e^{-2x}(c_2x + c_1) - \frac{e^{-2x}(1+x) \operatorname{expIntegral}_1(-2x)}{4} - \frac{3}{8} + \frac{(2x-2) \ln(x)}{8}$$

Verified OK.

14.22.2 Solving as linear second order ode solved by an integrating factor ode

The ode satisfies this form

$$y'' + p(x)y' + \frac{(p(x)^2 + p'(x))y}{2} = f(x)$$

Where $p(x) = 4$. Therefore, there is an integrating factor given by

$$\begin{aligned} M(x) &= e^{\frac{1}{2} \int p dx} \\ &= e^{\int 4 dx} \\ &= e^{2x} \end{aligned}$$

Multiplying both sides of the ODE by the integrating factor $M(x)$ makes the left side of the ODE a complete differential

$$\begin{aligned} (M(x)y)'' &= x \ln(x) e^{2x} \\ (e^{2x}y)'' &= x \ln(x) e^{2x} \end{aligned}$$

Integrating once gives

$$(e^{2x}y)' = -\frac{\exp\text{Integral}_1(-2x)}{4} + \frac{(-1 + (2x - 1) \ln(x)) e^{2x}}{4} + c_1$$

Integrating again gives

$$(e^{2x}y) = \frac{(-2 - 2x) \exp\text{Integral}_1(-2x)}{8} + \frac{(-3 + (2x - 2) \ln(x)) e^{2x}}{8} + c_1x + c_2$$

Hence the solution is

$$y = \frac{\frac{(-2-2x) \exp\text{Integral}_1(-2x)}{8} + \frac{(-3+(2x-2) \ln(x)) e^{2x}}{8} + c_1x + c_2}{e^{2x}}$$

Or

$$\begin{aligned} y &= \frac{x \ln(x)}{4} - \frac{\ln(x)}{4} + c_1x e^{-2x} - \frac{x e^{-2x} \exp\text{Integral}_1(-2x)}{4} \\ &\quad + c_2 e^{-2x} - \frac{e^{-2x} \exp\text{Integral}_1(-2x)}{4} - \frac{3}{8} \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{x \ln(x)}{4} - \frac{\ln(x)}{4} + c_1 x e^{-2x} - \frac{x e^{-2x} \operatorname{expIntegral}_1(-2x)}{4} + c_2 e^{-2x} - \frac{e^{-2x} \operatorname{expIntegral}_1(-2x)}{4} - \frac{3}{8} \quad (1)$$




Figure 417: Slope field plot

Verification of solutions

$$y = \frac{x \ln(x)}{4} - \frac{\ln(x)}{4} + c_1 x e^{-2x} - \frac{x e^{-2x} \operatorname{expIntegral}_1(-2x)}{4} + c_2 e^{-2x} - \frac{e^{-2x} \operatorname{expIntegral}_1(-2x)}{4} - \frac{3}{8}$$

Verified OK.

14.22.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 4y' + 4y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 4 \\ C &= 4 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 0 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 0 \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 374: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{4}{1} dx} \end{aligned}$$

$$\begin{aligned}
&= z_1 e^{-2x} \\
&= z_1 (e^{-2x})
\end{aligned}$$

Which simplifies to

$$y_1 = e^{-2x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}
y_2 &= y_1 \int \frac{e^{\int -\frac{4}{1} dx}}{(y_1)^2} dx \\
&= y_1 \int \frac{e^{-4x}}{(y_1)^2} dx \\
&= y_1(x)
\end{aligned}$$

Therefore the solution is

$$\begin{aligned}
y &= c_1 y_1 + c_2 y_2 \\
&= c_1 (e^{-2x}) + c_2 (e^{-2x}(x))
\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 4y' + 4y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-2x} + c_2 x e^{-2x}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = e^{-2x}$$

$$y_2 = x e^{-2x}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^{-2x} & x e^{-2x} \\ \frac{d}{dx}(e^{-2x}) & \frac{d}{dx}(x e^{-2x}) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^{-2x} & x e^{-2x} \\ -2 e^{-2x} & e^{-2x} - 2x e^{-2x} \end{vmatrix}$$

Therefore

$$W = (e^{-2x})(e^{-2x} - 2x e^{-2x}) - (x e^{-2x})(-2 e^{-2x})$$

Which simplifies to

$$W = e^{-4x}$$

Which simplifies to

$$W = e^{-4x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^2 e^{-2x} \ln(x)}{e^{-4x}} dx$$

Which simplifies to

$$u_1 = - \int x^2 \ln(x) e^{2x} dx$$

Hence

$$u_1 = - \left(\frac{1}{4} - \frac{1}{2}x + \frac{1}{2}x^2 \right) e^{2x} \ln(x) + \frac{e^{2x}x}{4} - \frac{3e^{2x}}{8} - \frac{\text{expIntegral}_1(-2x)}{4}$$

And Eq. (3) becomes

$$u_2 = \int \frac{e^{-2x} \ln(x) x}{e^{-4x}} dx$$

Which simplifies to

$$u_2 = \int x \ln(x) e^{2x} dx$$

Hence

$$u_2 = \left(-\frac{1}{4} + \frac{x}{2} \right) e^{2x} \ln(x) - \frac{\text{expIntegral}_1(-2x)}{4} - \frac{e^{2x}}{4}$$

Which simplifies to

$$u_1 = -\frac{\text{expIntegral}_1(-2x)}{4} + \frac{((-4x^2 + 4x - 2) \ln(x) + 2x - 3) e^{2x}}{8}$$

$$u_2 = -\frac{\text{expIntegral}_1(-2x)}{4} + \frac{(-1 + (2x - 1) \ln(x)) e^{2x}}{4}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(-\frac{\text{expIntegral}_1(-2x)}{4} + \frac{((-4x^2 + 4x - 2) \ln(x) + 2x - 3) e^{2x}}{8} \right) e^{-2x}$$

$$+ \left(-\frac{\text{expIntegral}_1(-2x)}{4} + \frac{(-1 + (2x - 1) \ln(x)) e^{2x}}{4} \right) x e^{-2x}$$

Which simplifies to

$$y_p(x) = -\frac{e^{-2x}(1+x) \expIntegral_1(-2x)}{4} - \frac{3}{8} + \frac{(2x-2) \ln(x)}{8}$$

Therefore the general solution is

$$y = y_h + y_p$$

$$= (c_1 e^{-2x} + c_2 x e^{-2x}) + \left(-\frac{e^{-2x}(1+x) \expIntegral_1(-2x)}{4} - \frac{3}{8} + \frac{(2x-2) \ln(x)}{8} \right)$$

Which simplifies to

$$y = e^{-2x}(c_2 x + c_1) - \frac{e^{-2x}(1+x) \expIntegral_1(-2x)}{4} - \frac{3}{8} + \frac{(2x-2) \ln(x)}{8}$$

Summary

The solution(s) found are the following

$$y = e^{-2x}(c_2 x + c_1) - \frac{e^{-2x}(1+x) \expIntegral_1(-2x)}{4} - \frac{3}{8} + \frac{(2x-2) \ln(x)}{8} \quad (1)$$




Figure 418: Slope field plot

Verification of solutions

$$y = e^{-2x}(c_2x + c_1) - \frac{e^{-2x}(1+x)\text{expIntegral}_1(-2x)}{4} - \frac{3}{8} + \frac{(2x-2)\ln(x)}{8}$$

Verified OK.

14.22.4 Maple step by step solution

Let's solve

$$y'' + 4y' + 4y = x \ln(x)$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 4r + 4 = 0$$

- Factor the characteristic polynomial

$$(r + 2)^2 = 0$$

- Root of the characteristic polynomial

$$r = -2$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-2x}$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = x e^{-2x}$$

- General solution of the ODE

$$y = c_1y_1(x) + c_2y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1e^{-2x} + c_2xe^{-2x} + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = x \ln(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-2x} & x e^{-2x} \\ -2 e^{-2x} & e^{-2x} - 2x e^{-2x} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = e^{-4x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = e^{-2x} \left(- \int x^2 \ln(x) e^{2x} dx \right) + \left(\int x \ln(x) e^{2x} dx \right) x$$

- Compute integrals

$$y_p(x) = -\frac{e^{-2x}(1+x)\text{Ei}_1(-2x)}{4} - \frac{3}{8} + \frac{(2x-2)\ln(x)}{8}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{-2x} + c_2 x e^{-2x} - \frac{e^{-2x}(1+x)\text{Ei}_1(-2x)}{4} - \frac{3}{8} + \frac{(2x-2)\ln(x)}{8}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 37

```
dsolve(diff(y(x),x$2)+4*diff(y(x),x)+4*y(x)=x*ln(x),y(x), singsol=all)
```

$$y(x) = -\frac{3}{8} - \frac{e^{-2x}(x+1)\text{expIntegral}_1(-2x)}{4} + (c_1 x + c_2) e^{-2x} + \frac{\ln(x)(x-1)}{4}$$

✓ Solution by Mathematica

Time used: 0.09 (sec). Leaf size: 52

```
DSolve[y''[x]+4*y'[x]+4*y[x]==x*Log[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{8}e^{-2x}(2(x+1)\text{ExpIntegralEi}(2x) - 3e^{2x} + 2e^{2x}(x-1)\log(x) + 8c_2x + 8c_1)$$

14.23 problem 3(g)

14.23.1 Solving as second order euler ode ode	2585
14.23.2 Solving as second order change of variable on x method 2 ode .	2588
14.23.3 Solving as second order change of variable on x method 1 ode .	2594
14.23.4 Solving as second order change of variable on y method 2 ode .	2599
14.23.5 Solving as second order integrable as is ode	2604
14.23.6 Solving as type second_order_integrable_as_is (not using ABC version)	2605
14.23.7 Solving using Kovacic algorithm	2607
14.23.8 Solving as exact linear second order ode ode	2614

Internal problem ID [6393]

Internal file name [OUTPUT/5641_Sunday_June_05_2022_03_45_40_PM_37710944/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Problems for Review and Discovery. Drill excercises. Page 105

Problem number: 3(g).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_euler_ode", "exact linear second order ode", "second_order_integrable_as_is", "second_order_change_of_variable_on_x_method_1", "second_order_change_of_variable_on_x_method_2", "second_order_change_of_variable_on_y_method_2"

Maple gives the following as the ode type

```
[[_2nd_order, _exact, _linear, _nonhomogeneous]]
```

$$x^2y'' + 3xy' + y = \frac{2}{x}$$

14.23.1 Solving as second order euler ode ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = x^2, B = 3x, C = 1, f(x) = \frac{2}{x}$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$x^2y'' + 3xy' + y = 0$$

This is Euler second order ODE. Let the solution be $y = x^r$, then $y' = rx^{r-1}$ and $y'' = r(r-1)x^{r-2}$. Substituting these back into the given ODE gives

$$x^2(r(r-1))x^{r-2} + 3rx^{r-1} + x^r = 0$$

Simplifying gives

$$r(r-1)x^r + 3rx^r + x^r = 0$$

Since $x^r \neq 0$ then dividing throughout by x^r gives

$$r(r-1) + 3r + 1 = 0$$

Or

$$r^2 + 2r + 1 = 0 \tag{1}$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = -1$$

$$r_2 = -1$$

Since the roots are equal, then the general solution is

$$y = c_1y_1 + c_2y_2$$

Where $y_1 = x^r$ and $y_2 = x^r \ln(x)$. Hence

$$y = \frac{c_1}{x} + \frac{c_2 \ln(x)}{x}$$

Next, we find the particular solution to the ODE

$$x^2y'' + 3xy' + y = \frac{2}{x}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1y_1 + u_2y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \frac{1}{x}$$

$$y_2 = \frac{\ln(x)}{x}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \frac{1}{x} & \frac{\ln(x)}{x} \\ \frac{d}{dx} \left(\frac{1}{x} \right) & \frac{d}{dx} \left(\frac{\ln(x)}{x} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \frac{1}{x} & \frac{\ln(x)}{x} \\ -\frac{1}{x^2} & -\frac{\ln(x)}{x^2} + \frac{1}{x^2} \end{vmatrix}$$

Therefore

$$W = \left(\frac{1}{x}\right) \left(-\frac{\ln(x)}{x^2} + \frac{1}{x^2}\right) - \left(\frac{\ln(x)}{x}\right) \left(-\frac{1}{x^2}\right)$$

Which simplifies to

$$W = \frac{1}{x^3}$$

Which simplifies to

$$W = \frac{1}{x^3}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\frac{2\ln(x)}{x^2}}{\frac{1}{x}} dx$$

Which simplifies to

$$u_1 = - \int \frac{2\ln(x)}{x} dx$$

Hence

$$u_1 = -\ln(x)^2$$

And Eq. (3) becomes

$$u_2 = \int \frac{\frac{2}{x^2}}{\frac{1}{x}} dx$$

Which simplifies to

$$u_2 = \int \frac{2}{x} dx$$

Hence

$$u_2 = 2\ln(x)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{\ln(x)^2}{x}$$

Therefore the general solution is

$$\begin{aligned}y &= y_h + y_p \\ &= \frac{\ln(x)^2 + c_1 + c_2 \ln(x)}{x}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{\ln(x)^2 + c_1 + c_2 \ln(x)}{x} \quad (1)$$

Verification of solutions

$$y = \frac{\ln(x)^2 + c_1 + c_2 \ln(x)}{x}$$

Verified OK.

14.23.2 Solving as second order change of variable on x method 2 ode

This is second order non-homogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$.
 y_h is the solution to

$$x^2 y'' + 3xy' + y = 0$$

In normal form the ode

$$x^2 y'' + 3xy' + y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$\begin{aligned}p(x) &= \frac{3}{x} \\ q(x) &= \frac{1}{x^2}\end{aligned}$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x)\tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned} \tau &= \int e^{-(\int p(x)dx)} dx \\ &= \int e^{-(\int \frac{3}{x} dx)} dx \\ &= \int e^{-3\ln(x)} dx \\ &= \int \frac{1}{x^3} dx \\ &= -\frac{1}{2x^2} \end{aligned} \quad (6)$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned} q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\ &= \frac{\frac{1}{x^2}}{\frac{1}{x^6}} \\ &= x^4 \end{aligned} \quad (7)$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned} \frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) + x^4y(\tau) &= 0 \end{aligned}$$

But in terms of τ

$$x^4 = \frac{1}{4\tau^2}$$

Hence the above ode becomes

$$\frac{d^2}{d\tau^2}y(\tau) + \frac{y(\tau)}{4\tau^2} = 0$$

The above ode is now solved for $y(\tau)$. The ode can be written as

$$4\left(\frac{d^2}{d\tau^2}y(\tau)\right)\tau^2 + y(\tau) = 0$$

Which shows it is a Euler ODE. This is Euler second order ODE. Let the solution be $y(\tau) = \tau^r$, then $y' = r\tau^{r-1}$ and $y'' = r(r-1)\tau^{r-2}$. Substituting these back into the given ODE gives

$$4\tau^2(r(r-1))\tau^{r-2} + 0r\tau^{r-1} + \tau^r = 0$$

Simplifying gives

$$4r(r-1)\tau^r + 0\tau^r + \tau^r = 0$$

Since $\tau^r \neq 0$ then dividing throughout by τ^r gives

$$4r(r-1) + 0 + 1 = 0$$

Or

$$4r^2 - 4r + 1 = 0 \tag{1}$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = \frac{1}{2}$$
$$r_2 = \frac{1}{2}$$

Since the roots are equal, then the general solution is

$$y(\tau) = c_1y_1 + c_2y_2$$

Where $y_1 = \tau^r$ and $y_2 = \tau^r \ln(\tau)$. Hence

$$y(\tau) = c_1\sqrt{\tau} + c_2\sqrt{\tau} \ln(\tau)$$

The above solution is now transformed back to y using (6) which results in

$$y = \frac{\sqrt{2} \sqrt{-\frac{1}{x^2}} (c_1 - c_2 \ln(2) + c_2 \ln(-\frac{1}{x^2}))}{2}$$

Therefore the homogeneous solution y_h is

$$y_h = \frac{\sqrt{2} \sqrt{-\frac{1}{x^2}} (c_1 - c_2 \ln(2) + c_2 \ln(-\frac{1}{x^2}))}{2}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \sqrt{-\frac{1}{x^2}}$$

$$y_2 = -\frac{\sqrt{2} \sqrt{-\frac{1}{x^2}} \ln(2)}{2} + \frac{\sqrt{2} \sqrt{-\frac{1}{x^2}} \ln(-\frac{1}{x^2})}{2}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \sqrt{-\frac{1}{x^2}} & -\frac{\sqrt{2} \sqrt{-\frac{1}{x^2}} \ln(2)}{2} + \frac{\sqrt{2} \sqrt{-\frac{1}{x^2}} \ln(-\frac{1}{x^2})}{2} \\ \frac{d}{dx} \left(\sqrt{-\frac{1}{x^2}} \right) & \frac{d}{dx} \left(-\frac{\sqrt{2} \sqrt{-\frac{1}{x^2}} \ln(2)}{2} + \frac{\sqrt{2} \sqrt{-\frac{1}{x^2}} \ln(-\frac{1}{x^2})}{2} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \sqrt{-\frac{1}{x^2}} & -\frac{\sqrt{2}\sqrt{-\frac{1}{x^2}}\ln(2)}{2} + \frac{\sqrt{2}\sqrt{-\frac{1}{x^2}}\ln(-\frac{1}{x^2})}{2} \\ \frac{1}{\sqrt{-\frac{1}{x^2}}x^3} & -\frac{\sqrt{2}\ln(2)}{2\sqrt{-\frac{1}{x^2}}x^3} + \frac{\sqrt{2}\ln(-\frac{1}{x^2})}{2\sqrt{-\frac{1}{x^2}}x^3} - \frac{\sqrt{2}\sqrt{-\frac{1}{x^2}}}{x} \end{vmatrix}$$

Therefore

$$W = \left(\sqrt{-\frac{1}{x^2}}\right) \left(-\frac{\sqrt{2}\ln(2)}{2\sqrt{-\frac{1}{x^2}}x^3} + \frac{\sqrt{2}\ln(-\frac{1}{x^2})}{2\sqrt{-\frac{1}{x^2}}x^3} - \frac{\sqrt{2}\sqrt{-\frac{1}{x^2}}}{x}\right) - \left(-\frac{\sqrt{2}\sqrt{-\frac{1}{x^2}}\ln(2)}{2} + \frac{\sqrt{2}\sqrt{-\frac{1}{x^2}}\ln(-\frac{1}{x^2})}{2}\right) \left(\frac{1}{\sqrt{-\frac{1}{x^2}}x^3}\right)$$

Which simplifies to

$$W = \frac{\sqrt{2}}{x^3}$$

Which simplifies to

$$W = \frac{\sqrt{2}}{x^3}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{-\sqrt{2}\sqrt{-\frac{1}{x^2}}\ln(2) + \sqrt{2}\sqrt{-\frac{1}{x^2}}\ln(-\frac{1}{x^2})}{\frac{x}{\frac{\sqrt{2}}{x}}} dx$$

Which simplifies to

$$u_1 = - \int \left(-\ln(2) + \ln\left(-\frac{1}{x^2}\right)\right) \sqrt{-\frac{1}{x^2}} dx$$

Hence

$$u_1 = x\sqrt{-\frac{1}{x^2}}\ln(2)\ln(x) + \frac{x\sqrt{-\frac{1}{x^2}}\ln(-\frac{1}{x^2})^2}{4}$$

And Eq. (3) becomes

$$u_2 = \int \frac{2\sqrt{-\frac{1}{x^2}}}{\frac{\sqrt{2}}{x}} dx$$

Which simplifies to

$$u_2 = \int \sqrt{2} \sqrt{-\frac{1}{x^2}} dx$$

Hence

$$u_2 = \sqrt{2} \sqrt{-\frac{1}{x^2}} x \ln(x)$$

Which simplifies to

$$u_1 = \sqrt{-\frac{1}{x^2}} x \left(\ln(2) \ln(x) + \frac{\ln\left(-\frac{1}{x^2}\right)^2}{4} \right)$$

$$u_2 = \sqrt{2} \sqrt{-\frac{1}{x^2}} x \ln(x)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{\ln(2) \ln(x) + \frac{\ln\left(-\frac{1}{x^2}\right)^2}{4}}{x} + \sqrt{2} \sqrt{-\frac{1}{x^2}} x \ln(x) \left(-\frac{\sqrt{2} \sqrt{-\frac{1}{x^2}} \ln(2)}{2} + \frac{\sqrt{2} \sqrt{-\frac{1}{x^2}} \ln\left(-\frac{1}{x^2}\right)}{2} \right)$$

Which simplifies to

$$y_p(x) = -\frac{\ln\left(-\frac{1}{x^2}\right) (4 \ln(x) + \ln\left(-\frac{1}{x^2}\right))}{4x}$$

Therefore the general solution is

$$y = y_h + y_p$$

$$= \left(\frac{\sqrt{2} \sqrt{-\frac{1}{x^2}} (c_1 - c_2 \ln(2) + c_2 \ln\left(-\frac{1}{x^2}\right))}{2} \right) + \left(-\frac{\ln\left(-\frac{1}{x^2}\right) (4 \ln(x) + \ln\left(-\frac{1}{x^2}\right))}{4x} \right)$$

Summary

The solution(s) found are the following

$$y = \frac{\sqrt{2} \sqrt{-\frac{1}{x^2}} (c_1 - c_2 \ln(2) + c_2 \ln(-\frac{1}{x^2}))}{2} - \frac{\ln(-\frac{1}{x^2}) (4 \ln(x) + \ln(-\frac{1}{x^2}))}{4x} \quad (1)$$

Verification of solutions

$$y = \frac{\sqrt{2} \sqrt{-\frac{1}{x^2}} (c_1 - c_2 \ln(2) + c_2 \ln(-\frac{1}{x^2}))}{2} - \frac{\ln(-\frac{1}{x^2}) (4 \ln(x) + \ln(-\frac{1}{x^2}))}{4x}$$

Verified OK.

14.23.3 Solving as second order change of variable on x method 1 ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = x^2$, $B = 3x$, $C = 1$, $f(x) = \frac{2}{x}$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$x^2 y'' + 3xy' + y = 0$$

In normal form the ode

$$x^2 y'' + 3xy' + y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{3}{x}$$
$$q(x) = \frac{1}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) results

$$\frac{d^2}{d\tau^2} y(\tau) + p_1 \left(\frac{d}{d\tau} y(\tau) \right) + q_1 y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x) \tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $q_1 = c^2$ where c is some constant. Therefore from (5)

$$\begin{aligned} \tau' &= \frac{1}{c} \sqrt{q} \\ &= \frac{\sqrt{\frac{1}{x^2}}}{c} \\ \tau'' &= -\frac{1}{c \sqrt{\frac{1}{x^2}} x^3} \end{aligned} \quad (6)$$

Substituting the above into (4) results in

$$\begin{aligned} p_1(\tau) &= \frac{\tau''(x) + p(x) \tau'(x)}{\tau'(x)^2} \\ &= \frac{-\frac{1}{c \sqrt{\frac{1}{x^2}} x^3} + \frac{3}{x} \frac{\sqrt{\frac{1}{x^2}}}{c}}{\left(\frac{\sqrt{\frac{1}{x^2}}}{c}\right)^2} \\ &= 2c \end{aligned}$$

Therefore ode (3) now becomes

$$\begin{aligned} y(\tau)'' + p_1 y(\tau)' + q_1 y(\tau) &= 0 \\ \frac{d^2}{d\tau^2} y(\tau) + 2c \left(\frac{d}{d\tau} y(\tau) \right) + c^2 y(\tau) &= 0 \end{aligned} \quad (7)$$

The above ode is now solved for $y(\tau)$. Since the ode is now constant coefficients, it can be easily solved to give

$$y(\tau) = e^{-c\tau} c_1$$

Now from (6)

$$\begin{aligned} \tau &= \int \frac{1}{c} \sqrt{q} dx \\ &= \frac{\int \sqrt{\frac{1}{x^2}} dx}{c} \\ &= \frac{\sqrt{\frac{1}{x^2}} x \ln(x)}{c} \end{aligned}$$

Substituting the above into the solution obtained gives

$$y = \frac{c_1}{x}$$

Now the particular solution to this ODE is found

$$x^2 y'' + 3xy' + y = \frac{2}{x}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \sqrt{-\frac{1}{x^2}}$$

$$y_2 = -\frac{\sqrt{2} \sqrt{-\frac{1}{x^2}} \ln(2)}{2} + \frac{\sqrt{2} \sqrt{-\frac{1}{x^2}} \ln\left(-\frac{1}{x^2}\right)}{2}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = -\int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \sqrt{-\frac{1}{x^2}} & -\frac{\sqrt{2} \sqrt{-\frac{1}{x^2}} \ln(2)}{2} + \frac{\sqrt{2} \sqrt{-\frac{1}{x^2}} \ln\left(-\frac{1}{x^2}\right)}{2} \\ \frac{d}{dx} \left(\sqrt{-\frac{1}{x^2}} \right) & \frac{d}{dx} \left(-\frac{\sqrt{2} \sqrt{-\frac{1}{x^2}} \ln(2)}{2} + \frac{\sqrt{2} \sqrt{-\frac{1}{x^2}} \ln\left(-\frac{1}{x^2}\right)}{2} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \sqrt{-\frac{1}{x^2}} & -\frac{\sqrt{2}\sqrt{-\frac{1}{x^2}}\ln(2)}{2} + \frac{\sqrt{2}\sqrt{-\frac{1}{x^2}}\ln(-\frac{1}{x^2})}{2} \\ \frac{1}{\sqrt{-\frac{1}{x^2}}x^3} & -\frac{\sqrt{2}\ln(2)}{2\sqrt{-\frac{1}{x^2}}x^3} + \frac{\sqrt{2}\ln(-\frac{1}{x^2})}{2\sqrt{-\frac{1}{x^2}}x^3} - \frac{\sqrt{2}\sqrt{-\frac{1}{x^2}}}{x} \end{vmatrix}$$

Therefore

$$W = \left(\sqrt{-\frac{1}{x^2}}\right) \left(-\frac{\sqrt{2}\ln(2)}{2\sqrt{-\frac{1}{x^2}}x^3} + \frac{\sqrt{2}\ln(-\frac{1}{x^2})}{2\sqrt{-\frac{1}{x^2}}x^3} - \frac{\sqrt{2}\sqrt{-\frac{1}{x^2}}}{x}\right) - \left(-\frac{\sqrt{2}\sqrt{-\frac{1}{x^2}}\ln(2)}{2} + \frac{\sqrt{2}\sqrt{-\frac{1}{x^2}}\ln(-\frac{1}{x^2})}{2}\right) \left(\frac{1}{\sqrt{-\frac{1}{x^2}}x^3}\right)$$

Which simplifies to

$$W = \frac{\sqrt{2}}{x^3}$$

Which simplifies to

$$W = \frac{\sqrt{2}}{x^3}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{-\sqrt{2}\sqrt{-\frac{1}{x^2}}\ln(2) + \sqrt{2}\sqrt{-\frac{1}{x^2}}\ln(-\frac{1}{x^2})}{\frac{x}{\frac{\sqrt{2}}{x}}} dx$$

Which simplifies to

$$u_1 = - \int \left(-\ln(2) + \ln\left(-\frac{1}{x^2}\right)\right) \sqrt{-\frac{1}{x^2}} dx$$

Hence

$$u_1 = x\sqrt{-\frac{1}{x^2}}\ln(2)\ln(x) + \frac{x\sqrt{-\frac{1}{x^2}}\ln(-\frac{1}{x^2})^2}{4}$$

And Eq. (3) becomes

$$u_2 = \int \frac{2\sqrt{-\frac{1}{x^2}}}{\frac{\sqrt{2}}{x}} dx$$

Which simplifies to

$$u_2 = \int \sqrt{2} \sqrt{-\frac{1}{x^2}} dx$$

Hence

$$u_2 = \sqrt{2} \sqrt{-\frac{1}{x^2}} x \ln(x)$$

Which simplifies to

$$u_1 = \sqrt{-\frac{1}{x^2}} x \left(\ln(2) \ln(x) + \frac{\ln\left(-\frac{1}{x^2}\right)^2}{4} \right)$$

$$u_2 = \sqrt{2} \sqrt{-\frac{1}{x^2}} x \ln(x)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{\ln(2) \ln(x) + \frac{\ln\left(-\frac{1}{x^2}\right)^2}{4}}{x} + \sqrt{2} \sqrt{-\frac{1}{x^2}} x \ln(x) \left(-\frac{\sqrt{2} \sqrt{-\frac{1}{x^2}} \ln(2)}{2} + \frac{\sqrt{2} \sqrt{-\frac{1}{x^2}} \ln\left(-\frac{1}{x^2}\right)}{2} \right)$$

Which simplifies to

$$y_p(x) = -\frac{\ln\left(-\frac{1}{x^2}\right) (4 \ln(x) + \ln\left(-\frac{1}{x^2}\right))}{4x}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(\frac{c_1}{x}\right) + \left(-\frac{\ln\left(-\frac{1}{x^2}\right) (4 \ln(x) + \ln\left(-\frac{1}{x^2}\right))}{4x}\right) \\ &= -\frac{\ln\left(-\frac{1}{x^2}\right) (4 \ln(x) + \ln\left(-\frac{1}{x^2}\right))}{4x} + \frac{c_1}{x} \end{aligned}$$

Which simplifies to

$$y = \frac{-\frac{\ln\left(-\frac{1}{x^2}\right)\left(4\ln(x)+\ln\left(-\frac{1}{x^2}\right)\right)}{4} + c_1}{x}$$

Summary

The solution(s) found are the following

$$y = \frac{-\frac{\ln\left(-\frac{1}{x^2}\right)\left(4\ln(x)+\ln\left(-\frac{1}{x^2}\right)\right)}{4} + c_1}{x} \quad (1)$$

Verification of solutions

$$y = \frac{-\frac{\ln\left(-\frac{1}{x^2}\right)\left(4\ln(x)+\ln\left(-\frac{1}{x^2}\right)\right)}{4} + c_1}{x}$$

Verified OK.

14.23.4 Solving as second order change of variable on y method 2 ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = x^2$, $B = 3x$, $C = 1$, $f(x) = \frac{2}{x}$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$x^2y'' + 3xy' + y = 0$$

In normal form the ode

$$x^2y'' + 3xy' + y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{3}{x}$$
$$q(x) = \frac{1}{x^2}$$

Applying change of variables on the dependent variable $y = v(x)x^n$ to (2) gives the following ode where the dependent variables is $v(x)$ and not y .

$$v''(x) + \left(\frac{2n}{x} + p\right)v'(x) + \left(\frac{n(n-1)}{x^2} + \frac{np}{x} + q\right)v(x) = 0 \quad (3)$$

Let the coefficient of $v(x)$ above be zero. Hence

$$\frac{n(n-1)}{x^2} + \frac{np}{x} + q = 0 \quad (4)$$

Substituting the earlier values found for $p(x)$ and $q(x)$ into (4) gives

$$\frac{n(n-1)}{x^2} + \frac{3n}{x^2} + \frac{1}{x^2} = 0 \quad (5)$$

Solving (5) for n gives

$$n = -1 \quad (6)$$

Substituting this value in (3) gives

$$v''(x) + \frac{v'(x)}{x} = 0$$
$$v''(x) + \frac{v'(x)}{x} = 0 \quad (7)$$

Using the substitution

$$u(x) = v'(x)$$

Then (7) becomes

$$u'(x) + \frac{u(x)}{x} = 0 \quad (8)$$

The above is now solved for $u(x)$. In canonical form the ODE is

$$\begin{aligned}u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{u}{x}\end{aligned}$$

Where $f(x) = -\frac{1}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u} du &= -\frac{1}{x} dx \\ \int \frac{1}{u} du &= \int -\frac{1}{x} dx \\ \ln(u) &= -\ln(x) + c_1 \\ u &= e^{-\ln(x)+c_1} \\ &= \frac{c_1}{x}\end{aligned}$$

Now that $u(x)$ is known, then

$$\begin{aligned}v'(x) &= u(x) \\ v(x) &= \int u(x) dx + c_2 \\ &= c_1 \ln(x) + c_2\end{aligned}$$

Hence

$$\begin{aligned}y &= v(x) x^n \\ &= \frac{c_1 \ln(x) + c_2}{x} \\ &= \frac{c_1 \ln(x) + c_2}{x}\end{aligned}$$

Now the particular solution to this ODE is found

$$x^2 y'' + 3xy' + y = \frac{2}{x}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \frac{1}{x}$$

$$y_2 = \frac{\ln(x)}{x}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \frac{1}{x} & \frac{\ln(x)}{x} \\ \frac{d}{dx} \left(\frac{1}{x} \right) & \frac{d}{dx} \left(\frac{\ln(x)}{x} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \frac{1}{x} & \frac{\ln(x)}{x} \\ -\frac{1}{x^2} & -\frac{\ln(x)}{x^2} + \frac{1}{x^2} \end{vmatrix}$$

Therefore

$$W = \left(\frac{1}{x} \right) \left(-\frac{\ln(x)}{x^2} + \frac{1}{x^2} \right) - \left(\frac{\ln(x)}{x} \right) \left(-\frac{1}{x^2} \right)$$

Which simplifies to

$$W = \frac{1}{x^3}$$

Which simplifies to

$$W = \frac{1}{x^3}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\frac{2 \ln(x)}{x^2}}{\frac{1}{x}} dx$$

Which simplifies to

$$u_1 = - \int \frac{2 \ln(x)}{x} dx$$

Hence

$$u_1 = - \ln(x)^2$$

And Eq. (3) becomes

$$u_2 = \int \frac{\frac{2}{x^2}}{\frac{1}{x}} dx$$

Which simplifies to

$$u_2 = \int \frac{2}{x} dx$$

Hence

$$u_2 = 2 \ln(x)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{\ln(x)^2}{x}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(\frac{c_1 \ln(x) + c_2}{x} \right) + \left(\frac{\ln(x)^2}{x} \right) \\ &= \frac{\ln(x)^2}{x} + \frac{c_1 \ln(x) + c_2}{x} \end{aligned}$$

Which simplifies to

$$y = \frac{\ln(x)^2 + c_1 \ln(x) + c_2}{x}$$

Summary

The solution(s) found are the following

$$y = \frac{\ln(x)^2 + c_1 \ln(x) + c_2}{x} \quad (1)$$

Verification of solutions

$$y = \frac{\ln(x)^2 + c_1 \ln(x) + c_2}{x}$$

Verified OK.

14.23.5 Solving as second order integrable as is ode

Integrating both sides of the ODE w.r.t x gives

$$\int (x^2 y'' + 3xy' + y) dx = \int \frac{2}{x} dx$$
$$x^2 y' + xy = 2 \ln(x) + c_1$$

Which is now solved for y .

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = \frac{1}{x}$$
$$q(x) = \frac{2 \ln(x) + c_1}{x^2}$$

Hence the ode is

$$y' + \frac{y}{x} = \frac{2 \ln(x) + c_1}{x^2}$$

The integrating factor μ is

$$\mu = e^{\int \frac{1}{x} dx}$$
$$= x$$

The ode becomes

$$\frac{d}{dx}(\mu y) = (\mu) \left(\frac{2 \ln(x) + c_1}{x^2} \right)$$
$$\frac{d}{dx}(xy) = (x) \left(\frac{2 \ln(x) + c_1}{x^2} \right)$$
$$d(xy) = \left(\frac{2 \ln(x) + c_1}{x} \right) dx$$

Integrating gives

$$xy = \int \frac{2 \ln(x) + c_1}{x} dx$$
$$xy = \ln(x)^2 + c_1 \ln(x) + c_2$$

Dividing both sides by the integrating factor $\mu = x$ results in

$$y = \frac{\ln(x)^2 + c_1 \ln(x)}{x} + \frac{c_2}{x}$$

which simplifies to

$$y = \frac{\ln(x)^2 + c_1 \ln(x) + c_2}{x}$$

Summary

The solution(s) found are the following

$$y = \frac{\ln(x)^2 + c_1 \ln(x) + c_2}{x} \quad (1)$$

Verification of solutions

$$y = \frac{\ln(x)^2 + c_1 \ln(x) + c_2}{x}$$

Verified OK.

14.23.6 Solving as type second_order_integrable_as_is (not using ABC version)

Writing the ode as

$$x^2 y'' + 3xy' + y = \frac{2}{x}$$

Integrating both sides of the ODE w.r.t x gives

$$\int (x^2 y'' + 3xy' + y) dx = \int \frac{2}{x} dx$$
$$x^2 y' + xy = 2 \ln(x) + c_1$$

Which is now solved for y .

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = \frac{1}{x}$$
$$q(x) = \frac{2 \ln(x) + c_1}{x^2}$$

Hence the ode is

$$y' + \frac{y}{x} = \frac{2 \ln(x) + c_1}{x^2}$$

The integrating factor μ is

$$\mu = e^{\int \frac{1}{x} dx}$$
$$= x$$

The ode becomes

$$\frac{d}{dx}(\mu y) = (\mu) \left(\frac{2 \ln(x) + c_1}{x^2} \right)$$
$$\frac{d}{dx}(xy) = (x) \left(\frac{2 \ln(x) + c_1}{x^2} \right)$$
$$d(xy) = \left(\frac{2 \ln(x) + c_1}{x} \right) dx$$

Integrating gives

$$xy = \int \frac{2 \ln(x) + c_1}{x} dx$$
$$xy = \ln(x)^2 + c_1 \ln(x) + c_2$$

Dividing both sides by the integrating factor $\mu = x$ results in

$$y = \frac{\ln(x)^2 + c_1 \ln(x)}{x} + \frac{c_2}{x}$$

which simplifies to

$$y = \frac{\ln(x)^2 + c_1 \ln(x) + c_2}{x}$$

Summary

The solution(s) found are the following

$$y = \frac{\ln(x)^2 + c_1 \ln(x) + c_2}{x} \tag{1}$$

Verification of solutions

$$y = \frac{\ln(x)^2 + c_1 \ln(x) + c_2}{x}$$

Verified OK.

14.23.7 Solving using Kovacic algorithm

Writing the ode as

$$x^2y'' + 3xy' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= x^2 \\ B &= 3x \\ C &= 1 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{4x^2} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 4x^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(-\frac{1}{4x^2}\right) z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 376: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4x^2$. There is a pole at $x = 0$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = -\frac{1}{4x^2}$$

For the pole at $x = 0$ let b be the coefficient of $\frac{1}{x^2}$ in the partial fractions decomposition of r given above. Therefore $b = -\frac{1}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{1}{2} \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{2} \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ , which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = -\frac{1}{4x^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = -\frac{1}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{1}{2} \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{2} \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = -\frac{1}{4x^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
0	2	0	$\frac{1}{2}$	$\frac{1}{2}$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	$\frac{1}{2}$	$\frac{1}{2}$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω .

Trying $\alpha_{\infty}^{-} = \frac{1}{2}$ then

$$\begin{aligned} d &= \alpha_{\infty}^{-} - (\alpha_{c_1}^{-}) \\ &= \frac{1}{2} - \left(\frac{1}{2}\right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_{\infty}$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^{-}}{x - c_1} \right) + (-)[\sqrt{r}]_{\infty} \\ &= \frac{1}{2x} + (-)(0) \\ &= \frac{1}{2x} \\ &= \frac{1}{2x} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r)p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2\left(\frac{1}{2x}\right)(0) + \left(\left(-\frac{1}{2x^2}\right) + \left(\frac{1}{2x}\right)^2 - \left(-\frac{1}{4x^2}\right) \right) &= 0 \\ 0 &= 0 \end{aligned}$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= pe^{\int \omega dx} \\ &= e^{\int \frac{1}{2x} dx} \\ &= \sqrt{x} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\&= z_1 e^{-\int \frac{1}{2} \frac{3x}{x^2} dx} \\&= z_1 e^{-\frac{3 \ln(x)}{2}} \\&= z_1 \left(\frac{1}{x^{\frac{3}{2}}} \right)\end{aligned}$$

Which simplifies to

$$y_1 = \frac{1}{x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{3x}{x^2} dx}}{(y_1)^2} dx \\&= y_1 \int \frac{e^{-3 \ln(x)}}{(y_1)^2} dx \\&= y_1 (\ln(x))\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\&= c_1 \left(\frac{1}{x} \right) + c_2 \left(\frac{1}{x} (\ln(x)) \right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$x^2 y'' + 3xy' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = \frac{c_1}{x} + \frac{c_2 \ln(x)}{x}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \frac{1}{x}$$

$$y_2 = \frac{\ln(x)}{x}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \frac{1}{x} & \frac{\ln(x)}{x} \\ \frac{d}{dx} \left(\frac{1}{x} \right) & \frac{d}{dx} \left(\frac{\ln(x)}{x} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \frac{1}{x} & \frac{\ln(x)}{x} \\ -\frac{1}{x^2} & -\frac{\ln(x)}{x^2} + \frac{1}{x^2} \end{vmatrix}$$

Therefore

$$W = \left(\frac{1}{x}\right) \left(-\frac{\ln(x)}{x^2} + \frac{1}{x^2}\right) - \left(\frac{\ln(x)}{x}\right) \left(-\frac{1}{x^2}\right)$$

Which simplifies to

$$W = \frac{1}{x^3}$$

Which simplifies to

$$W = \frac{1}{x^3}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\frac{2\ln(x)}{x^2}}{\frac{1}{x}} dx$$

Which simplifies to

$$u_1 = - \int \frac{2\ln(x)}{x} dx$$

Hence

$$u_1 = -\ln(x)^2$$

And Eq. (3) becomes

$$u_2 = \int \frac{\frac{2}{x^2}}{\frac{1}{x}} dx$$

Which simplifies to

$$u_2 = \int \frac{2}{x} dx$$

Hence

$$u_2 = 2\ln(x)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{\ln(x)^2}{x}$$

Therefore the general solution is

$$\begin{aligned}y &= y_h + y_p \\ &= \left(\frac{c_1}{x} + \frac{c_2 \ln(x)}{x} \right) + \left(\frac{\ln(x)^2}{x} \right)\end{aligned}$$

Which simplifies to

$$y = \frac{c_1 + c_2 \ln(x)}{x} + \frac{\ln(x)^2}{x}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1 + c_2 \ln(x)}{x} + \frac{\ln(x)^2}{x} \quad (1)$$

Verification of solutions

$$y = \frac{c_1 + c_2 \ln(x)}{x} + \frac{\ln(x)^2}{x}$$

Verified OK.

14.23.8 Solving as exact linear second order ode

An ode of the form

$$p(x) y'' + q(x) y' + r(x) y = s(x)$$

is exact if

$$p''(x) - q'(x) + r(x) = 0 \quad (1)$$

For the given ode we have

$$\begin{aligned}p(x) &= x^2 \\ q(x) &= 3x \\ r(x) &= 1 \\ s(x) &= \frac{2}{x}\end{aligned}$$

Hence

$$\begin{aligned}p''(x) &= 2 \\ q'(x) &= 3\end{aligned}$$

Therefore (1) becomes

$$2 - (3) + (1) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(x)y' + (q(x) - p'(x))y)' = s(x)$$

Integrating gives

$$p(x)y' + (q(x) - p'(x))y = \int s(x) dx$$

Substituting the above values for p, q, r, s gives

$$x^2y' + xy = \int \frac{2}{x} dx$$

We now have a first order ode to solve which is

$$x^2y' + xy = 2 \ln(x) + c_1$$

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = \frac{1}{x}$$
$$q(x) = \frac{2 \ln(x) + c_1}{x^2}$$

Hence the ode is

$$y' + \frac{y}{x} = \frac{2 \ln(x) + c_1}{x^2}$$

The integrating factor μ is

$$\mu = e^{\int \frac{1}{x} dx}$$
$$= x$$

The ode becomes

$$\frac{d}{dx}(\mu y) = (\mu) \left(\frac{2 \ln(x) + c_1}{x^2} \right)$$
$$\frac{d}{dx}(xy) = (x) \left(\frac{2 \ln(x) + c_1}{x^2} \right)$$
$$d(xy) = \left(\frac{2 \ln(x) + c_1}{x} \right) dx$$

Integrating gives

$$xy = \int \frac{2 \ln(x) + c_1}{x} dx$$
$$xy = \ln(x)^2 + c_1 \ln(x) + c_2$$

Dividing both sides by the integrating factor $\mu = x$ results in

$$y = \frac{\ln(x)^2 + c_1 \ln(x)}{x} + \frac{c_2}{x}$$

which simplifies to

$$y = \frac{\ln(x)^2 + c_1 \ln(x) + c_2}{x}$$

Summary

The solution(s) found are the following

$$y = \frac{\ln(x)^2 + c_1 \ln(x) + c_2}{x} \quad (1)$$

Verification of solutions

$$y = \frac{\ln(x)^2 + c_1 \ln(x) + c_2}{x}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
<- high order exact linear fully integrable successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 18

```
dsolve(x^2*diff(y(x),x$2)+3*x*diff(y(x),x)+y(x)=2/x,y(x), singsol=all)
```

$$y(x) = \frac{c_2 + \ln(x)^2 + c_1 \ln(x)}{x}$$

✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 21

```
DSolve[x^2*y''[x]+3*x*y'[x]+y[x]==2/x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\log^2(x) + c_2 \log(x) + c_1}{x}$$

14.24 problem 3(h)

14.24.1 Solving as second order linear constant coeff ode	2618
14.24.2 Solving using Kovacic algorithm	2623
14.24.3 Maple step by step solution	2629

Internal problem ID [6394]

Internal file name [OUTPUT/5642_Sunday_June_05_2022_03_45_41_PM_39558801/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Problems for Review and Discovery. Drill excercises. Page 105

Problem number: 3(h).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + 4y = \tan(x)^2$$

14.24.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 4, f(x) = \tan(x)^2$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 4y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 4$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 4 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 4 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 4$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(4)} \\ &= \pm 2i \end{aligned}$$

Hence

$$\lambda_1 = +2i$$

$$\lambda_2 = -2i$$

Which simplifies to

$$\lambda_1 = 2i$$

$$\lambda_2 = -2i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 2$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (c_1 \cos(2x) + c_2 \sin(2x))$$

Or

$$y = c_1 \cos(2x) + c_2 \sin(2x)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \cos(2x) + c_2 \sin(2x)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \cos(2x)$$

$$y_2 = \sin(2x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(2x) & \sin(2x) \\ \frac{d}{dx}(\cos(2x)) & \frac{d}{dx}(\sin(2x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(2x) & \sin(2x) \\ -2 \sin(2x) & 2 \cos(2x) \end{vmatrix}$$

Therefore

$$W = (\cos(2x))(2 \cos(2x)) - (\sin(2x))(-2 \sin(2x))$$

Which simplifies to

$$W = 2 \cos (2x)^2 + 2 \sin (2x)^2$$

Which simplifies to

$$W = 2$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\sin (2x) \tan (x)^2}{2} dx$$

Which simplifies to

$$u_1 = - \int \sin (x)^2 \tan (x) dx$$

Hence

$$u_1 = \frac{\sin (x)^2}{2} + \ln (\cos (x))$$

And Eq. (3) becomes

$$u_2 = \int \frac{\tan (x)^2 \cos (2x)}{2} dx$$

Which simplifies to

$$u_2 = \int \left(\sin (x)^2 - \frac{\tan (x)^2}{2} \right) dx$$

Hence

$$u_2 = -\frac{\cos (x) \sin (x)}{2} + x - \frac{\tan (x)}{2}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(\frac{\sin (x)^2}{2} + \ln (\cos (x)) \right) \cos (2x) + \left(-\frac{\cos (x) \sin (x)}{2} + x - \frac{\tan (x)}{2} \right) \sin (2x)$$

Which simplifies to

$$y_p(x) = (2 \cos(x)^2 - 1) \ln(\cos(x)) + 2 \cos(x) \sin(x) x - \frac{3 \sin(x)^2}{2}$$

Therefore the general solution is

$$y = y_h + y_p$$

$$= (c_1 \cos(2x) + c_2 \sin(2x)) + \left((2 \cos(x)^2 - 1) \ln(\cos(x)) + 2 \cos(x) \sin(x) x - \frac{3 \sin(x)^2}{2} \right)$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(2x) + c_2 \sin(2x) + (2 \cos(x)^2 - 1) \ln(\cos(x)) + 2 \cos(x) \sin(x) x - \frac{3 \sin(x)^2}{2} \quad (1)$$




Figure 419: Slope field plot

Verification of solutions

$$y = c_1 \cos(2x) + c_2 \sin(2x) + (2 \cos(x)^2 - 1) \ln(\cos(x)) + 2 \cos(x) \sin(x) x - \frac{3 \sin(x)^2}{2}$$

Verified OK.

14.24.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 4y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 4 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-4}{1} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -4 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -4z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 377: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -4$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(2x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned} y_1 &= z_1 \\ &= \cos(2x) \end{aligned}$$

Which simplifies to

$$y_1 = \cos(2x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(2x) \int \frac{1}{\cos(2x)^2} dx \\ &= \cos(2x) \left(\frac{\tan(2x)}{2} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(2x)) + c_2 \left(\cos(2x) \left(\frac{\tan(2x)}{2} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 4y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 \cos(2x) + \frac{c_2 \sin(2x)}{2}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \cos(2x)$$

$$y_2 = \frac{\sin(2x)}{2}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(2x) & \frac{\sin(2x)}{2} \\ \frac{d}{dx}(\cos(2x)) & \frac{d}{dx}\left(\frac{\sin(2x)}{2}\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(2x) & \frac{\sin(2x)}{2} \\ -2 \sin(2x) & \cos(2x) \end{vmatrix}$$

Therefore

$$W = (\cos(2x))(\cos(2x)) - \left(\frac{\sin(2x)}{2}\right)(-2 \sin(2x))$$

Which simplifies to

$$W = \cos(2x)^2 + \sin(2x)^2$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\frac{\sin(2x) \tan(x)^2}{2}}{1} dx$$

Which simplifies to

$$u_1 = - \int \sin(x)^2 \tan(x) dx$$

Hence

$$u_1 = \frac{\sin(x)^2}{2} + \ln(\cos(x))$$

And Eq. (3) becomes

$$u_2 = \int \frac{\tan(x)^2 \cos(2x)}{1} dx$$

Which simplifies to

$$u_2 = \int (2 \sin(x)^2 - \tan(x)^2) dx$$

Hence

$$u_2 = -\cos(x) \sin(x) - \tan(x) + 2x$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(\frac{\sin(x)^2}{2} + \ln(\cos(x)) \right) \cos(2x) + \frac{(-\cos(x)\sin(x) - \tan(x) + 2x)\sin(2x)}{2}$$

Which simplifies to

$$y_p(x) = (2\cos(x)^2 - 1)\ln(\cos(x)) + 2\cos(x)\sin(x)x - \frac{3\sin(x)^2}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 \cos(2x) + \frac{c_2 \sin(2x)}{2} \right) \\ &\quad + \left((2\cos(x)^2 - 1)\ln(\cos(x)) + 2\cos(x)\sin(x)x - \frac{3\sin(x)^2}{2} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned} y &= c_1 \cos(2x) + \frac{c_2 \sin(2x)}{2} + (2\cos(x)^2 - 1)\ln(\cos(x)) \\ &\quad + 2\cos(x)\sin(x)x - \frac{3\sin(x)^2}{2} \end{aligned} \tag{1}$$




Figure 420: Slope field plot

Verification of solutions

$$y = c_1 \cos(2x) + \frac{c_2 \sin(2x)}{2} + (2 \cos(x)^2 - 1) \ln(\cos(x)) + 2 \cos(x) \sin(x) x - \frac{3 \sin(x)^2}{2}$$

Verified OK.

14.24.3 Maple step by step solution

Let's solve

$$y'' + 4y = \tan(x)^2$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 4 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-16})}{2}$$

- Roots of the characteristic polynomial

$$r = (-2I, 2I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(2x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(2x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 \cos(2x) + c_2 \sin(2x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \tan(x)^2 \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(2x) & \sin(2x) \\ -2\sin(2x) & 2\cos(2x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 2$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\cos(2x) \left(\int \sin(x)^2 \tan(x) dx \right) + \frac{\sin(2x) \left(\int (2\sin(x)^2 - \tan(x)^2) dx \right)}{2}$$

- Compute integrals

$$y_p(x) = (2\cos(x)^2 - 1) \ln(\cos(x)) + 2\cos(x)\sin(x)x - \frac{3\sin(x)^2}{2}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 \cos(2x) + c_2 \sin(2x) + (2\cos(x)^2 - 1) \ln(\cos(x)) + 2\cos(x)\sin(x)x - \frac{3\sin(x)^2}{2}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 42

```
dsolve(diff(y(x),x$2)+4*y(x)=tan(x)^2,y(x), singsol=all)
```

$$y(x) = (2 \cos(x)^2 - 1) \ln(\cos(x)) + 2c_1 \cos(x)^2 + 2(x + c_2) \sin(x) \cos(x) - \frac{3 \sin(x)^2}{2} - c_1$$

✓ Solution by Mathematica

Time used: 0.091 (sec). Leaf size: 32

```
DSolve[y''[x]+4*y[x]==Tan[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow (x + c_2) \sin(2x) + \cos(2x) \left(\log(\cos(x)) + \frac{1}{4} + c_1 \right) - \frac{3}{4}$$

14.25 problem 4(a)

14.25.1 Maple step by step solution 2636

Internal problem ID [6395]

Internal file name [OUTPUT/5643_Sunday_June_05_2022_03_45_43_PM_45346907/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Problems for Review and Discovery. Drill exercises. Page 105

Problem number: 4(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"reduction_of_order", "second_order_linear_constant_coeff"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - y = 3e^{2x}$$

Given that one solution of the ode is

$$y_1 = e^{2x}$$

This is second order nonhomogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = -1, f(x) = 3e^{2x}$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the inhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - y = 0$$

Given one basis solution $y_1(x)$, then the second basis solution is given by

$$y_2(x) = y_1 \left(\int \frac{e^{-\int p dx}}{y_1^2} dx \right)$$

Where $p(x)$ is the coefficient of y' when the ode is written in the normal form

$$y'' + p(x)y' + q(x)y = f(x)$$

Looking at the ode to solve shows that

$$p(x) = 0$$

Therefore

$$y_2(x) = e^{2x} \left(\int e^{-(\int 0 dx)} e^{-4x} dx \right)$$

$$y_2(x) = e^{2x} \int \frac{1}{e^{4x}} dx$$

$$y_2(x) = e^{2x} \left(\int e^{-4x} dx \right)$$

$$y_2(x) = -\frac{e^{2x} e^{-4x}}{4}$$

Hence the solution is

$$\begin{aligned} y &= c_1 y_1(x) + c_2 y_2(x) \\ &= c_1 e^{2x} - \frac{c_2 e^{2x} e^{-4x}}{4} \end{aligned}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^{2x} - \frac{c_2 e^{2x} e^{-4x}}{4}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned} y_1 &= e^{2x} \\ y_2 &= -\frac{e^{2x} e^{-4x}}{4} \end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^{2x} & -\frac{e^{2x}e^{-4x}}{4} \\ \frac{d}{dx}(e^{2x}) & \frac{d}{dx}\left(-\frac{e^{2x}e^{-4x}}{4}\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^{2x} & -\frac{e^{2x}e^{-4x}}{4} \\ 2e^{2x} & \frac{e^{2x}e^{-4x}}{2} \end{vmatrix}$$

Therefore

$$W = (e^{2x}) \left(\frac{e^{2x}e^{-4x}}{2} \right) - \left(-\frac{e^{2x}e^{-4x}}{4} \right) (2e^{2x})$$

Which simplifies to

$$W = e^{4x}e^{-4x}$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{-\frac{3e^{4x}e^{-4x}}{4}}{1} dx$$

Which simplifies to

$$u_1 = - \int -\frac{3}{4} dx$$

Hence

$$u_1 = \frac{3x}{4}$$

And Eq. (3) becomes

$$u_2 = \int \frac{3e^{4x}}{1} dx$$

Which simplifies to

$$u_2 = \int 3e^{4x} dx$$

Hence

$$u_2 = \frac{3e^{4x}}{4}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{3e^{2x}x}{4} - \frac{3e^{4x}e^{2x}e^{-4x}}{16}$$

Which simplifies to

$$y_p(x) = \frac{3(4x - 1)e^{2x}}{16}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1e^{2x} - \frac{c_2e^{2x}e^{-4x}}{4} \right) + \left(\frac{3(4x - 1)e^{2x}}{16} \right) \end{aligned}$$

Which simplifies to

$$y = c_1e^{2x} - \frac{c_2e^{-2x}}{4} + \frac{3(4x - 1)e^{2x}}{16}$$

Summary

The solution(s) found are the following

$$y = c_1e^{2x} - \frac{c_2e^{-2x}}{4} + \frac{3(4x - 1)e^{2x}}{16} \quad (1)$$




Figure 421: Slope field plot

Verification of solutions

$$y = c_1 e^{2x} - \frac{c_2 e^{-2x}}{4} + \frac{3(4x - 1) e^{2x}}{16}$$

Verified OK.

14.25.1 Maple step by step solution

Let's solve

$$y'' - y = 3e^{2x}$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - 1 = 0$$

- Factor the characteristic polynomial

$$(r - 1)(r + 1) = 0$$

- Roots of the characteristic polynomial

$$r = (-1, 1)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^x$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{-x} + c_2 e^x + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = 3 e^{2x} \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-x} & e^x \\ -e^{-x} & e^x \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 2$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{3 e^{-x} (\int e^{3x} dx)}{2} + \frac{3 e^x (\int e^x dx)}{2}$$

- Compute integrals

$$y_p(x) = e^{2x}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{-x} + c_2 e^x + e^{2x}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve([diff(y(x),x$2)-y(x)=3*exp(2*x),exp(2*x)],singsol=all)
```

$$y(x) = c_2 e^{-x} + e^x c_1 + e^{2x}$$

✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 25

```
DSolve[y''[x]-y[x]==3*Exp[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{2x} + c_1 e^x + c_2 e^{-x}$$

14.26 problem 4(b)

14.26.1 Maple step by step solution 2643

Internal problem ID [6396]

Internal file name [OUTPUT/5644_Sunday_June_05_2022_03_45_45_PM_62158437/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Problems for Review and Discovery. Drill exercises. Page 105

Problem number: 4(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"reduction_of_order", "second_order_linear_constant_coeff"**

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + y = -8 \sin(3x)$$

Given that one solution of the ode is

$$y_1 = \sin(3x)$$

This is second order nonhomogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 1, f(x) = -8 \sin(3x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the inhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

Given one basis solution $y_1(x)$, then the second basis solution is given by

$$y_2(x) = y_1 \left(\int \frac{e^{-\int p dx}}{y_1^2} dx \right)$$

Where $p(x)$ is the coefficient of y' when the ode is written in the normal form

$$y'' + p(x)y' + q(x)y = f(x)$$

Looking at the ode to solve shows that

$$p(x) = 0$$

Therefore

$$y_2(x) = \sin(3x) \left(\int \frac{e^{-(\int 0 dx)}}{\sin(3x)^2} dx \right)$$

$$y_2(x) = \sin(3x) \int \frac{1}{\sin(3x)^2} dx$$

$$y_2(x) = \sin(3x) \left(\int \csc(3x)^2 dx \right)$$

$$y_2(x) = -\frac{\sin(3x) \cot(3x)}{3}$$

Hence the solution is

$$\begin{aligned} y &= c_1 y_1(x) + c_2 y_2(x) \\ &= c_1 \sin(3x) - \frac{c_2 \sin(3x) \cot(3x)}{3} \end{aligned}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \sin(3x) - \frac{c_2 \sin(3x) \cot(3x)}{3}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned} y_1 &= \sin(3x) \\ y_2 &= -\frac{\sin(3x) \cot(3x)}{3} \end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \sin(3x) & -\frac{\sin(3x)\cot(3x)}{3} \\ \frac{d}{dx}(\sin(3x)) & \frac{d}{dx}\left(-\frac{\sin(3x)\cot(3x)}{3}\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \sin(3x) & -\frac{\sin(3x)\cot(3x)}{3} \\ 3\cos(3x) & -\cos(3x)\cot(3x) - \frac{\sin(3x)(-3-3\cot(3x)^2)}{3} \end{vmatrix}$$

Therefore

$$W = (\sin(3x)) \left(-\cos(3x)\cot(3x) - \frac{\sin(3x)(-3-3\cot(3x)^2)}{3} \right) - \left(-\frac{\sin(3x)\cot(3x)}{3} \right) (3\cos(3x))$$

Which simplifies to

$$W = \sin(3x)^2 \cot(3x)^2 + \sin(3x)^2$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\frac{8\sin(3x)^2 \cot(3x)}{3}}{1} dx$$

Which simplifies to

$$u_1 = - \int \frac{4 \sin(6x)}{3} dx$$

Hence

$$u_1 = \frac{2 \cos(6x)}{9}$$

And Eq. (3) becomes

$$u_2 = \int \frac{-8 \sin(3x)^2}{1} dx$$

Which simplifies to

$$u_2 = \int (-4 + 4 \cos(6x)) dx$$

Hence

$$u_2 = -4x + \frac{2 \sin(6x)}{3}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{2 \cos(6x) \sin(3x)}{9} - \frac{\left(-4x + \frac{2 \sin(6x)}{3}\right) \sin(3x) \cot(3x)}{3}$$

Which simplifies to

$$y_p(x) = -\frac{2 \sin(3x)}{9} + \frac{4x \cos(3x)}{3}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 \sin(3x) - \frac{c_2 \sin(3x) \cot(3x)}{3} \right) + \left(-\frac{2 \sin(3x)}{9} + \frac{4x \cos(3x)}{3} \right) \end{aligned}$$

Which simplifies to

$$y = c_1 \sin(3x) - \frac{c_2 \cos(3x)}{3} - \frac{2 \sin(3x)}{9} + \frac{4x \cos(3x)}{3}$$

Summary

The solution(s) found are the following

$$y = c_1 \sin(3x) - \frac{c_2 \cos(3x)}{3} - \frac{2 \sin(3x)}{9} + \frac{4x \cos(3x)}{3} \quad (1)$$




Figure 422: Slope field plot

Verification of solutions

$$y = c_1 \sin(3x) - \frac{c_2 \cos(3x)}{3} - \frac{2 \sin(3x)}{9} + \frac{4x \cos(3x)}{3}$$

Verified OK.

14.26.1 Maple step by step solution

Let's solve

$$y'' + y = -8 \sin(3x)$$

- Highest derivative means the order of the ODE is 2
- y''
- Characteristic polynomial of homogeneous ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (-I, I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = \cos(x) c_1 + c_2 \sin(x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x), y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x), y_2(x))} dx \right), f(x) = -8 \sin(3x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -4 \cos(x) \left(\int (-\cos(2x) + \cos(4x)) dx \right) + 4 \sin(x) \left(\int (-\sin(4x) - \sin(2x)) dx \right)$$

- Compute integrals

$$y_p(x) = \sin(3x)$$

- Substitute particular solution into general solution to ODE

$$y = \cos(x) c_1 + c_2 \sin(x) + \sin(3x)$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve([diff(y(x),x$2)+y(x)=-8*sin(3*x),sin(3*x)],singsol=all)
```

$$y(x) = \sin(x) c_2 + \cos(x) c_1 + \sin(3x)$$

✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 20

```
DSolve[y''[x]+y[x]==-8*Sin[3*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \sin(3x) + c_1 \cos(x) + c_2 \sin(x)$$

14.27 problem 4(c)

14.27.1 Maple step by step solution 2651

Internal problem ID [6397]

Internal file name [OUTPUT/5645_Sunday_June_05_2022_03_45_47_PM_58516173/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Problems for Review and Discovery. Drill exercises. Page 105

Problem number: 4(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"reduction_of_order", "second_order_linear_constant_coeff"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' + y' + y = x^2 + 2x + 2$$

Given that one solution of the ode is

$$y_1 = x^2$$

This is second order nonhomogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 1, C = 1, f(x) = x^2 + 2x + 2$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the inhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' + y = 0$$

Given one basis solution $y_1(x)$, then the second basis solution is given by

$$y_2(x) = y_1 \left(\int \frac{e^{-(\int p dx)}}{y_1^2} dx \right)$$

Where $p(x)$ is the coefficient of y' when the ode is written in the normal form

$$y'' + p(x)y' + q(x)y = f(x)$$

Looking at the ode to solve shows that

$$p(x) = 1$$

Therefore

$$y_2(x) = x^2 \left(\int \frac{e^{-(\int 1 dx)}}{x^4} dx \right)$$

$$y_2(x) = x^2 \int \frac{e^{-x}}{x^4} dx$$

$$y_2(x) = x^2 \left(\int \frac{e^{-x}}{x^4} dx \right)$$

$$y_2(x) = x^2 \left(-\frac{e^{-x}}{3x^3} + \frac{e^{-x}}{6x^2} - \frac{e^{-x}}{6x} + \frac{\text{expIntegral}_1(x)}{6} \right)$$

Hence the solution is

$$\begin{aligned} y &= c_1 y_1(x) + c_2 y_2(x) \\ &= c_1 x^2 + c_2 x^2 \left(-\frac{e^{-x}}{3x^3} + \frac{e^{-x}}{6x^2} - \frac{e^{-x}}{6x} + \frac{\text{expIntegral}_1(x)}{6} \right) \end{aligned}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 x^2 + c_2 x^2 \left(-\frac{e^{-x}}{3x^3} + \frac{e^{-x}}{6x^2} - \frac{e^{-x}}{6x} + \frac{\text{expIntegral}_1(x)}{6} \right)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = x^2$$

$$y_2 = x^2 \left(-\frac{e^{-x}}{3x^3} + \frac{e^{-x}}{6x^2} - \frac{e^{-x}}{6x} + \frac{\text{expIntegral}_1(x)}{6} \right)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} x^2 & x^2 \left(-\frac{e^{-x}}{3x^3} + \frac{e^{-x}}{6x^2} - \frac{e^{-x}}{6x} + \frac{\text{expIntegral}_1(x)}{6} \right) \\ \frac{d}{dx}(x^2) & \frac{d}{dx} \left(x^2 \left(-\frac{e^{-x}}{3x^3} + \frac{e^{-x}}{6x^2} - \frac{e^{-x}}{6x} + \frac{\text{expIntegral}_1(x)}{6} \right) \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} x^2 & x^2 \left(-\frac{e^{-x}}{3x^3} + \frac{e^{-x}}{6x^2} - \frac{e^{-x}}{6x} + \frac{\text{expIntegral}_1(x)}{6} \right) \\ 2x & 2x \left(-\frac{e^{-x}}{3x^3} + \frac{e^{-x}}{6x^2} - \frac{e^{-x}}{6x} + \frac{\text{expIntegral}_1(x)}{6} \right) + \frac{e^{-x}}{x^2} \end{vmatrix}$$

Therefore

$$W = (x^2) \left(2x \left(-\frac{e^{-x}}{3x^3} + \frac{e^{-x}}{6x^2} - \frac{e^{-x}}{6x} + \frac{\text{expIntegral}_1(x)}{6} \right) + \frac{e^{-x}}{x^2} \right) - \left(x^2 \left(-\frac{e^{-x}}{3x^3} + \frac{e^{-x}}{6x^2} - \frac{e^{-x}}{6x} + \frac{\text{expIntegral}_1(x)}{6} \right) \right) (2x)$$

Which simplifies to

$$W = e^{-x}$$

Which simplifies to

$$W = e^{-x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x^2 \left(-\frac{e^{-x}}{3x^3} + \frac{e^{-x}}{6x^2} - \frac{e^{-x}}{6x} + \frac{\exp\text{Integral}_1(x)}{6} \right) (x^2 + 2x + 2)}{e^{-x}} dx$$

Which simplifies to

$$u_1 = - \int \frac{(x^2 + 2x + 2) (e^x \exp\text{Integral}_1(x) x^3 - x^2 + x - 2)}{6x} dx$$

Hence

$$u_1 = - \left(\int_0^x \frac{(\alpha^2 + 2\alpha + 2) (e^\alpha \exp\text{Integral}_1(\alpha) \alpha^3 - \alpha^2 + \alpha - 2)}{6\alpha} d\alpha \right)$$

And Eq. (3) becomes

$$u_2 = \int \frac{(x^2 + 2x + 2) x^2}{e^{-x}} dx$$

Which simplifies to

$$u_2 = \int x^2 (x^2 + 2x + 2) e^x dx$$

Hence

$$u_2 = (x^4 - 2x^3 + 8x^2 - 16x + 16) e^x$$

Which simplifies to

$$u_1 = - \frac{\left(\int_0^x \frac{(\alpha^2 + 2\alpha + 2) (e^\alpha \exp\text{Integral}_1(\alpha) \alpha^3 - \alpha^2 + \alpha - 2)}{\alpha} d\alpha \right)}{6}$$

$$u_2 = (x^4 - 2x^3 + 8x^2 - 16x + 16) e^x$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\frac{\left(\int_0^x \frac{(\alpha^2+2\alpha+2)(e^\alpha \exp\text{Integral}_1(\alpha)\alpha^3-\alpha^2+\alpha-2)}{\alpha} d\alpha\right) x^2}{6} + (x^4 - 2x^3 + 8x^2 - 16x + 16) e^x x^2 \left(-\frac{e^{-x}}{3x^3} + \frac{e^{-x}}{6x^2} - \frac{e^{-x}}{6x} + \frac{\exp\text{Integral}_1(x)}{6}\right)$$

Which simplifies to

$$y_p(x) = \frac{-\left(\int_0^x \frac{(\alpha^2+2\alpha+2)(e^\alpha \exp\text{Integral}_1(\alpha)\alpha^3-\alpha^2+\alpha-2)}{\alpha} d\alpha\right) x^3 + (x^4 - 2x^3 + 8x^2 - 16x + 16) (e^x \exp\text{Integral}_1(x) x^3 - 6x)}{6x}$$

Therefore the general solution is

$$y = y_h + y_p = \left(c_1 x^2 + c_2 x^2 \left(-\frac{e^{-x}}{3x^3} + \frac{e^{-x}}{6x^2} - \frac{e^{-x}}{6x} + \frac{\exp\text{Integral}_1(x)}{6}\right)\right) + \left(-\frac{\left(\int_0^x \frac{(\alpha^2+2\alpha+2)(e^\alpha \exp\text{Integral}_1(\alpha)\alpha^3-\alpha^2+\alpha-2)}{\alpha} d\alpha\right) x^3 + (x^4 - 2x^3 + 8x^2 - 16x + 16) (e^x \exp\text{Integral}_1(x) x^3 - 6x)}{6x}\right)$$

Which simplifies to

$$y = \frac{-c_2(x^2 - x + 2) e^{-x} + 6\left(\frac{c_2 \exp\text{Integral}_1(x)}{6} + c_1\right) x^3}{6x} + \frac{-\left(\int_0^x \frac{(\alpha^2+2\alpha+2)(e^\alpha \exp\text{Integral}_1(\alpha)\alpha^3-\alpha^2+\alpha-2)}{\alpha} d\alpha\right) x^3 + (x^4 - 2x^3 + 8x^2 - 16x + 16) (e^x \exp\text{Integral}_1(x) x^3 - 6x)}{6x}$$

Summary

The solution(s) found are the following

$$y = \frac{-c_2(x^2 - x + 2) e^{-x} + 6\left(\frac{c_2 \exp\text{Integral}_1(x)}{6} + c_1\right) x^3}{6x} + \frac{-\left(\int_0^x \frac{(\alpha^2+2\alpha+2)(e^\alpha \exp\text{Integral}_1(\alpha)\alpha^3-\alpha^2+\alpha-2)}{\alpha} d\alpha\right) x^3 + (x^4 - 2x^3 + 8x^2 - 16x + 16) (e^x \exp\text{Integral}_1(x) x^3 - 6x)}{6x} \tag{1}$$




Figure 423: Slope field plot

Verification of solutions

$$y = \frac{-c_2(x^2 - x + 2)e^{-x} + 6\left(\frac{c_2 \exp\text{Integral}_1(x)}{6} + c_1\right)x^3}{6x} + \frac{-\left(\int_0^x \frac{(\alpha^2 + 2\alpha + 2)(e^\alpha \exp\text{Integral}_1(\alpha)\alpha^3 - \alpha^2 + \alpha - 2)}{\alpha} d\alpha\right)x^3 + (x^4 - 2x^3 + 8x^2 - 16x + 16)(e^x \exp\text{Integral}_1(x)x)}{6x}$$

Verified OK.

14.27.1 Maple step by step solution

Let's solve

$$y'' + y' + y = x^2 + 2x + 2$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + r + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-1) \pm (\sqrt{-3})}{2}$$

- Roots of the characteristic polynomial

$$r = \left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}, -\frac{1}{2} + \frac{i\sqrt{3}}{2} \right)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) c_1 + e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) c_2 + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = x^2 + 2x + 2 \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) & e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) \\ -\frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{2} - \frac{e^{-\frac{x}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{2} & -\frac{e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) \sqrt{3}}{2} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = \frac{\sqrt{3}e^{-x}}{2}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{2\sqrt{3}e^{-\frac{x}{2}} \left(\cos\left(\frac{\sqrt{3}x}{2}\right) \left(\int e^{\frac{x}{2}} (x^2+2x+2) \sin\left(\frac{\sqrt{3}x}{2}\right) dx \right) - \sin\left(\frac{\sqrt{3}x}{2}\right) \left(\int e^{\frac{x}{2}} (x^2+2x+2) \cos\left(\frac{\sqrt{3}x}{2}\right) dx \right) \right)}{3}$$

- Compute integrals

$$y_p(x) = x^2$$

- Substitute particular solution into general solution to ODE

$$y = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) c_1 + e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) c_2 + x^2$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 34

```
dsolve([diff(y(x),x$2)+diff(y(x),x)+y(x)=x^2+2*x+2,x^2],singsol=all)
```

$$y(x) = c_2 e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) + e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) c_1 + x^2$$

✓ Solution by Mathematica

Time used: 0.024 (sec). Leaf size: 51

```
DSolve[y''[x]+y'[x]+y[x]==x^2+2*x+2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x^2 + c_2 e^{-x/2} \cos\left(\frac{\sqrt{3}x}{2}\right) + c_1 e^{-x/2} \sin\left(\frac{\sqrt{3}x}{2}\right)$$

14.28 problem 4(d)

14.28.1 Maple step by step solution 2660

Internal problem ID [6398]

Internal file name [OUTPUT/5646_Sunday_June_05_2022_03_45_51_PM_29806910/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Problems for Review and Discovery. Drill exercises. Page 105

Problem number: 4(d).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "reduction_of_order", "exact linear second order ode", "second_order_integrable_as_is", "second_order_ode_missing_y", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_y]]
```

$$y'' + y' = \frac{x-1}{x}$$

Given that one solution of the ode is

$$y_1 = \ln(x)$$

This is second order nonhomogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 1, C = 0, f(x) = 1 - \frac{1}{x}$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the inhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y' = 0$$

Given one basis solution $y_1(x)$, then the second basis solution is given by

$$y_2(x) = y_1 \left(\int \frac{e^{-(\int p dx)}}{y_1^2} dx \right)$$

Where $p(x)$ is the coefficient of y' when the ode is written in the normal form

$$y'' + p(x)y' + q(x)y = f(x)$$

Looking at the ode to solve shows that

$$p(x) = 1$$

Therefore

$$y_2(x) = \ln(x) \left(\int \frac{e^{-(\int 1 dx)}}{\ln(x)^2} dx \right)$$

$$y_2(x) = \ln(x) \int \frac{e^{-x}}{\ln(x)^2} dx$$

$$y_2(x) = \ln(x) \left(\int \frac{e^{-x}}{\ln(x)^2} dx \right)$$

$$y_2(x) = \ln(x) \left(-\frac{x e^{-x}}{\ln(x)} + \int -\frac{e^{-x}(x-1)}{\ln(x)} dx \right)$$

Hence the solution is

$$\begin{aligned} y &= c_1 y_1(x) + c_2 y_2(x) \\ &= c_1 \ln(x) + c_2 \ln(x) \left(-\frac{x e^{-x}}{\ln(x)} + \int -\frac{e^{-x}(x-1)}{\ln(x)} dx \right) \end{aligned}$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \ln(x) + c_2 \ln(x) \left(-\frac{x e^{-x}}{\ln(x)} + \int -\frac{e^{-x}(x-1)}{\ln(x)} dx \right)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of

parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \ln(x)$$

$$y_2 = \ln(x) \left(-\frac{x e^{-x}}{\ln(x)} + \int -\frac{e^{-x}(x-1)}{\ln(x)} dx \right)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \ln(x) & \ln(x) \left(-\frac{x e^{-x}}{\ln(x)} + \int -\frac{e^{-x}(x-1)}{\ln(x)} dx \right) \\ \frac{d}{dx}(\ln(x)) & \frac{d}{dx} \left(\ln(x) \left(-\frac{x e^{-x}}{\ln(x)} + \int -\frac{e^{-x}(x-1)}{\ln(x)} dx \right) \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \ln(x) & \ln(x) \left(-\frac{x e^{-x}}{\ln(x)} + \int -\frac{e^{-x}(x-1)}{\ln(x)} dx \right) \\ \frac{1}{x} & \frac{-\frac{x e^{-x}}{\ln(x)} + \int -\frac{e^{-x}(x-1)}{\ln(x)} dx}{x} + \ln(x) \left(-\frac{e^{-x}}{\ln(x)} + \frac{x e^{-x}}{\ln(x)} + \frac{e^{-x}}{\ln(x)^2} - \frac{e^{-x}(x-1)}{\ln(x)} \right) \end{vmatrix}$$

Therefore

$$W = (\ln(x)) \left(\frac{-\frac{x e^{-x}}{\ln(x)} + \int -\frac{e^{-x}(x-1)}{\ln(x)} dx}{x} + \ln(x) \left(-\frac{e^{-x}}{\ln(x)} + \frac{x e^{-x}}{\ln(x)} + \frac{e^{-x}}{\ln(x)^2} - \frac{e^{-x}(x-1)}{\ln(x)} \right) \right) - \left(\ln(x) \left(-\frac{x e^{-x}}{\ln(x)} + \int -\frac{e^{-x}(x-1)}{\ln(x)} dx \right) \right) \left(\frac{1}{x} \right)$$

Which simplifies to

$$W = e^{-x}$$

Which simplifies to

$$W = e^{-x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\ln(x) \left(-\frac{x e^{-x}}{\ln(x)} + \int -\frac{e^{-x}(x-1)}{\ln(x)} dx \right) \left(1 - \frac{1}{x} \right)}{e^{-x}} dx$$

Which simplifies to

$$u_1 = - \int - \frac{(x-1) \left(\left(\int \frac{e^{-x}(x-1)}{\ln(x)} dx \right) e^x \ln(x) + x \right)}{x} dx$$

Hence

$$u_1 = - \left(\int_0^x - \frac{(\alpha-1) \left(\left(\int \frac{e^{-\alpha}(\alpha-1)}{\ln(\alpha)} d\alpha \right) e^\alpha \ln(\alpha) + \alpha \right)}{\alpha} d\alpha \right)$$

And Eq. (3) becomes

$$u_2 = \int \frac{\ln(x) \left(1 - \frac{1}{x} \right)}{e^{-x}} dx$$

Which simplifies to

$$u_2 = \int \frac{\ln(x) (x-1) e^x}{x} dx$$

Hence

$$u_2 = \int_0^x \frac{\ln(\alpha)(\alpha-1)e^\alpha}{\alpha} d\alpha$$

Which simplifies to

$$u_1 = \int_0^x \frac{(\alpha-1) \left(\left(\int \frac{e^{-\alpha(\alpha-1)}}{\ln(\alpha)} d\alpha \right) e^\alpha \ln(\alpha) + \alpha \right)}{\alpha} d\alpha$$

$$u_2 = \int_0^x \frac{\ln(\alpha)(\alpha-1)e^\alpha}{\alpha} d\alpha$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(\int_0^x \frac{(\alpha-1) \left(\left(\int \frac{e^{-\alpha(\alpha-1)}}{\ln(\alpha)} d\alpha \right) e^\alpha \ln(\alpha) + \alpha \right)}{\alpha} d\alpha \right) \ln(x)$$

$$+ \left(\int_0^x \frac{\ln(\alpha)(\alpha-1)e^\alpha}{\alpha} d\alpha \right) \ln(x) \left(-\frac{x e^{-x}}{\ln(x)} + \int -\frac{e^{-x}(x-1)}{\ln(x)} dx \right)$$

Which simplifies to

$$y_p(x) = -e^{-x} \left(\int_0^x \frac{\ln(\alpha)(\alpha-1)e^\alpha}{\alpha} d\alpha \right) x$$

$$- \ln(x) \left(\int_0^x \frac{\ln(\alpha)(\alpha-1)e^\alpha}{\alpha} d\alpha \right) \left(\int \frac{e^{-x}(x-1)}{\ln(x)} dx \right)$$

$$+ \left(\int_0^x \frac{(\alpha-1) \left(\left(\int \frac{e^{-\alpha(\alpha-1)}}{\ln(\alpha)} d\alpha \right) e^\alpha \ln(\alpha) + \alpha \right)}{\alpha} d\alpha \right) \ln(x)$$

Therefore the general solution is

$$y = y_h + y_p$$

$$= \left(c_1 \ln(x) + c_2 \ln(x) \left(-\frac{x e^{-x}}{\ln(x)} + \int -\frac{e^{-x}(x-1)}{\ln(x)} dx \right) \right) + \left(-e^{-x} \left(\int_0^x \frac{\ln(\alpha)(\alpha-1)e^\alpha}{\alpha} d\alpha \right) x - \ln(x) \left(\int_0^x \frac{\ln(\alpha)(\alpha-1)e^\alpha}{\alpha} d\alpha \right) \right)$$

Which simplifies to

$$y = -x e^{-x} c_2 - \ln(x) \left(\int \frac{e^{-x}(x-1)}{\ln(x)} dx \right) c_2 + c_1 \ln(x) - e^{-x} \left(\int_0^x \frac{\ln(\alpha)(\alpha-1)e^\alpha}{\alpha} d\alpha \right) x - \ln(x) \left(\int_0^x \frac{\ln(\alpha)(\alpha-1)e^\alpha}{\alpha} d\alpha \right)$$

Summary

The solution(s) found are the following

$$\begin{aligned} y = & -x e^{-x} c_2 - \ln(x) \left(\int \frac{e^{-x}(x-1)}{\ln(x)} dx \right) c_2 \\ & + c_1 \ln(x) - e^{-x} \left(\int_0^x \frac{\ln(\alpha)(\alpha-1)e^\alpha}{\alpha} d\alpha \right) x \\ & - \ln(x) \left(\int_0^x \frac{\ln(\alpha)(\alpha-1)e^\alpha}{\alpha} d\alpha \right) \left(\int \frac{e^{-x}(x-1)}{\ln(x)} dx \right) \\ & + \left(\int_0^x \frac{(\alpha-1) \left(\left(\int \frac{e^{-\alpha}(\alpha-1)}{\ln(\alpha)} d\alpha \right) e^\alpha \ln(\alpha) + \alpha \right)}{\alpha} d\alpha \right) \ln(x) \end{aligned} \quad (1)$$




Figure 424: Slope field plot

Verification of solutions

$$y = -x e^{-x} c_2 - \ln(x) \left(\int \frac{e^{-x}(x-1)}{\ln(x)} dx \right) c_2 + c_1 \ln(x) - e^{-x} \left(\int_0^x \frac{\ln(\alpha)(\alpha-1)e^\alpha}{\alpha} d\alpha \right) x \\ - \ln(x) \left(\int_0^x \frac{\ln(\alpha)(\alpha-1)e^\alpha}{\alpha} d\alpha \right) \left(\int \frac{e^{-x}(x-1)}{\ln(x)} dx \right) \\ + \left(\int_0^x \frac{(\alpha-1) \left(\int \frac{e^{-\alpha}(\alpha-1)}{\ln(\alpha)} d\alpha \right) e^\alpha \ln(\alpha) + \alpha}{\alpha} d\alpha \right) \ln(x)$$

Verified OK.

14.28.1 Maple step by step solution

Let's solve

$$y' + y'' = \frac{x-1}{x}$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + r = 0$$

- Factor the characteristic polynomial

$$r(r+1) = 0$$

- Roots of the characteristic polynomial

$$r = (-1, 0)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = 1$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{-x} + c_2 + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \frac{x-1}{x} \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-x} & 1 \\ -e^{-x} & 0 \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = e^{-x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -e^{-x} \left(\int \frac{e^x(x-1)}{x} dx \right) + \int \frac{x-1}{x} dx$$

- Compute integrals

$$y_p(x) = -e^{-x} \text{Ei}_1(-x) - 1 - \ln(x) + x$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{-x} + c_2 - e^{-x} \text{Ei}_1(-x) - 1 - \ln(x) + x$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
-> Calling odsolve with the ODE`, diff(_b(_a), _a) = -(_b(_a)*_a-_a+1)/_a, _b(_a)` *** Sub
  Methods for first order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  trying 1st order linear
  <- 1st order linear successful
<- high order exact linear fully integrable successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 26

```
dsolve([diff(y(x),x$2)+diff(y(x),x)=(x-1)/x,ln(x)],singsol=all)
```

$$y(x) = \int (1 + e^{-x} \text{expIntegral}_1(-x) + c_1 e^{-x}) dx + c_2$$

✓ Solution by Mathematica

Time used: 0.12 (sec). Leaf size: 30

```
DSolve[y''[x]+y'[x]==(x-1)/x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{-x} \text{ExpIntegralEi}(x) + x - \log(x) - c_1 e^{-x} + c_2$$

**15 Chapter 2. Problems for Review and Discovery.
Challenge excercises. Page 105**

15.1 problem 3 2664
15.2 problem 4 2689

15.1 problem 3

15.1.1 Existence and uniqueness analysis	2665
15.1.2 Solving as second order euler ode ode	2665
15.1.3 Solving as linear second order ode solved by an integrating factor ode	2667
15.1.4 Solving as second order change of variable on x method 2 ode .	2668
15.1.5 Solving as second order change of variable on x method 1 ode .	2671
15.1.6 Solving as second order change of variable on y method 1 ode .	2674
15.1.7 Solving as second order change of variable on y method 2 ode .	2676
15.1.8 Solving as second order ode non constant coeff transformation on B ode	2679
15.1.9 Solving using Kovacic algorithm	2681
15.1.10 Maple step by step solution	2685

Internal problem ID [6399]

Internal file name [OUTPUT/5647_Sunday_June_05_2022_03_45_55_PM_3219856/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Problems for Review and Discovery. Challenge excercises. Page 105

Problem number: 3.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_euler_ode", "second_order_change_of_variable_on_x_method_1", "second_order_change_of_variable_on_x_method_2", "second_order_change_of_variable_on_y_method_1", "second_order_change_of_variable_on_y_method_2", "linear_second_order_ode_solved_by_an_integrating_factor", "second_order_ode_non_constant_coeff_transformation_on_B"

Maple gives the following as the ode type

```
[[_Emden, _Fowler], [_2nd_order, _linear, ` _with_symmetry_[0,F(x)]`]]
```

$$x^2y'' - 2xy' + 2y = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

15.1.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$\begin{aligned} p(x) &= -\frac{2}{x} \\ q(x) &= \frac{2}{x^2} \\ F &= 0 \end{aligned}$$

Hence the ode is

$$y'' - \frac{2y'}{x} + \frac{2y}{x^2} = 0$$

The domain of $p(x) = -\frac{2}{x}$ is

$$\{x < 0 \vee 0 < x\}$$

But the point $x_0 = 0$ is not inside this domain. Hence existence and uniqueness theorem does not apply. There could be infinite number of solutions, or one solution or no solution at all.

15.1.2 Solving as second order euler ode ode

This is Euler second order ODE. Let the solution be $y = x^r$, then $y' = rx^{r-1}$ and $y'' = r(r-1)x^{r-2}$. Substituting these back into the given ODE gives

$$x^2(r(r-1))x^{r-2} - 2rxr^{r-1} + 2x^r = 0$$

Simplifying gives

$$r(r-1)x^r - 2rx^r + 2x^r = 0$$

Since $x^r \neq 0$ then dividing throughout by x^r gives

$$r(r-1) - 2r + 2 = 0$$

Or

$$r^2 - 3r + 2 = 0 \tag{1}$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = 1$$

$$r_2 = 2$$

Since the roots are real and distinct, then the general solution is

$$y = c_1y_1 + c_2y_2$$

Where $y_1 = x^{r_1}$ and $y_2 = x^{r_2}$. Hence

$$y = c_2x^2 + c_1x$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_2x^2 + c_1x \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = 0 \tag{1A}$$

Taking derivative of the solution gives

$$y' = 2c_2x + c_1$$

substituting $y' = 0$ and $x = 0$ in the above gives

$$0 = c_1 \tag{2A}$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 0$$

Substituting these values back in above solution results in

$$y = c_2x^2$$

Summary

The solution(s) found are the following

$$y = c_2x^2 \tag{1}$$

Verification of solutions

$$y = c_2x^2$$

Verified OK.

15.1.3 Solving as linear second order ode solved by an integrating factor ode

The ode satisfies this form

$$y'' + p(x)y' + \frac{(p(x)^2 + p'(x))y}{2} = f(x)$$

Where $p(x) = -\frac{2}{x}$. Therefore, there is an integrating factor given by

$$\begin{aligned}M(x) &= e^{\frac{1}{2} \int p dx} \\ &= e^{\int -\frac{2}{x} dx} \\ &= \frac{1}{x}\end{aligned}$$

Multiplying both sides of the ODE by the integrating factor $M(x)$ makes the left side of the ODE a complete differential

$$\begin{aligned}(M(x)y)'' &= 0 \\ \left(\frac{y}{x}\right)'' &= 0\end{aligned}$$

Integrating once gives

$$\left(\frac{y}{x}\right)' = c_1$$

Integrating again gives

$$\left(\frac{y}{x}\right) = c_1x + c_2$$

Hence the solution is

$$y = \frac{c_1x + c_2}{\frac{1}{x}}$$

Or

$$y = c_1x^2 + c_2x$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1x^2 + c_2x \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = 0 \quad (1A)$$

Taking derivative of the solution gives

$$y' = 2c_1x + c_2$$

substituting $y' = 0$ and $x = 0$ in the above gives

$$0 = c_2 \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_2 = 0$$

Substituting these values back in above solution results in

$$y = c_1x^2$$

Summary

The solution(s) found are the following

$$y = c_1x^2 \quad (1)$$

Verification of solutions

$$y = c_1x^2$$

Verified OK.

15.1.4 Solving as second order change of variable on x method 2 ode

In normal form the ode

$$x^2y'' - 2xy' + 2y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = -\frac{2}{x}$$
$$q(x) = \frac{2}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x)\tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned} \tau &= \int e^{-(\int p(x)dx)} dx \\ &= \int e^{-(\int -\frac{2}{x} dx)} dx \\ &= \int e^{2\ln(x)} dx \\ &= \int x^2 dx \\ &= \frac{x^3}{3} \end{aligned} \quad (6)$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned} q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\ &= \frac{\frac{2}{x^2}}{x^4} \\ &= \frac{2}{x^6} \end{aligned} \quad (7)$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned} \frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) + \frac{2y(\tau)}{x^6} &= 0 \end{aligned}$$

But in terms of τ

$$\frac{2}{x^6} = \frac{2}{9\tau^2}$$

Hence the above ode becomes

$$\frac{d^2}{d\tau^2}y(\tau) + \frac{2y(\tau)}{9\tau^2} = 0$$

The above ode is now solved for $y(\tau)$. The ode can be written as

$$9\left(\frac{d^2}{d\tau^2}y(\tau)\right)\tau^2 + 2y(\tau) = 0$$

Which shows it is a Euler ODE. This is Euler second order ODE. Let the solution be $y(\tau) = \tau^r$, then $y' = r\tau^{r-1}$ and $y'' = r(r-1)\tau^{r-2}$. Substituting these back into the given ODE gives

$$9\tau^2(r(r-1))\tau^{r-2} + 0r\tau^{r-1} + 2\tau^r = 0$$

Simplifying gives

$$9r(r-1)\tau^r + 0\tau^r + 2\tau^r = 0$$

Since $\tau^r \neq 0$ then dividing throughout by τ^r gives

$$9r(r-1) + 0 + 2 = 0$$

Or

$$9r^2 - 9r + 2 = 0 \tag{1}$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = \frac{1}{3}$$
$$r_2 = \frac{2}{3}$$

Since the roots are real and distinct, then the general solution is

$$y(\tau) = c_1y_1 + c_2y_2$$

Where $y_1 = \tau^{r_1}$ and $y_2 = \tau^{r_2}$. Hence

$$y(\tau) = c_1\tau^{\frac{1}{3}} + c_2\tau^{\frac{2}{3}}$$

The above solution is now transformed back to y using (6) which results in

$$y = \frac{c_1 3^{\frac{2}{3}}(x^3)^{\frac{1}{3}}}{3} + \frac{c_2 3^{\frac{1}{3}}(x^3)^{\frac{2}{3}}}{3}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = \frac{c_1 3^{\frac{2}{3}} (x^3)^{\frac{1}{3}}}{3} + \frac{c_2 3^{\frac{1}{3}} (x^3)^{\frac{2}{3}}}{3} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = 0 \quad (1A)$$

Taking derivative of the solution gives

$$y' = \frac{c_1 3^{\frac{2}{3}} x^2}{3 (x^3)^{\frac{2}{3}}} + \frac{2c_2 3^{\frac{1}{3}} x^2}{3 (x^3)^{\frac{1}{3}}}$$

substituting $y' = 0$ and $x = 0$ in the above gives

$$0 = \frac{3^{\frac{1}{3}} \left(\lim_{x \rightarrow 0} \frac{x^2 (2c_2 (x^3)^{\frac{1}{3}} + c_1 3^{\frac{1}{3}})}{(x^3)^{\frac{2}{3}}} \right)}{3} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Warning, unable to solve for constants of integrations.

Verification of solutions N/A

15.1.5 Solving as second order change of variable on x method 1 ode

In normal form the ode

$$x^2 y'' - 2xy' + 2y = 0 \quad (1)$$

Becomes

$$y'' + p(x) y' + q(x) y = 0 \quad (2)$$

Where

$$p(x) = -\frac{2}{x}$$
$$q(x) = \frac{2}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) results

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $q_1 = c^2$ where c is some constant. Therefore from (5)

$$\begin{aligned} \tau' &= \frac{1}{c}\sqrt{q} \\ &= \frac{\sqrt{2}\sqrt{\frac{1}{x^2}}}{c} \\ \tau'' &= -\frac{\sqrt{2}}{c\sqrt{\frac{1}{x^2}}x^3} \end{aligned} \quad (6)$$

Substituting the above into (4) results in

$$\begin{aligned} p_1(\tau) &= \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \\ &= \frac{-\frac{\sqrt{2}}{c\sqrt{\frac{1}{x^2}}x^3} - \frac{2}{x}\frac{\sqrt{2}\sqrt{\frac{1}{x^2}}}{c}}{\left(\frac{\sqrt{2}\sqrt{\frac{1}{x^2}}}{c}\right)^2} \\ &= -\frac{3c\sqrt{2}}{2} \end{aligned}$$

Therefore ode (3) now becomes

$$\begin{aligned} y(\tau)'' + p_1y(\tau)' + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) - \frac{3c\sqrt{2}}{2}\left(\frac{d}{d\tau}y(\tau)\right) + c^2y(\tau) &= 0 \end{aligned} \quad (7)$$

The above ode is now solved for $y(\tau)$. Since the ode is now constant coefficients, it can be easily solved to give

$$y(\tau) = e^{\frac{3\sqrt{2}c\tau}{4}} \left(c_1 \cosh\left(\frac{\sqrt{2}c\tau}{4}\right) + ic_2 \sinh\left(\frac{\sqrt{2}c\tau}{4}\right) \right)$$

Now from (6)

$$\begin{aligned}\tau &= \int \frac{1}{c} \sqrt{q} dx \\ &= \frac{\int \sqrt{2} \sqrt{\frac{1}{x^2}} dx}{c} \\ &= \frac{\sqrt{2} \sqrt{\frac{1}{x^2}} x \ln(x)}{c}\end{aligned}$$

Substituting the above into the solution obtained gives

$$y = x^{\frac{3}{2}} \left(c_1 \cosh \left(\frac{\ln(x)}{2} \right) + ic_2 \sinh \left(\frac{\ln(x)}{2} \right) \right)$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = x^{\frac{3}{2}} \left(c_1 \cosh \left(\frac{\ln(x)}{2} \right) + ic_2 \sinh \left(\frac{\ln(x)}{2} \right) \right) \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = 0 \quad (1A)$$

Taking derivative of the solution gives

$$y' = \frac{3\sqrt{x} \left(c_1 \cosh \left(\frac{\ln(x)}{2} \right) + ic_2 \sinh \left(\frac{\ln(x)}{2} \right) \right)}{2} + x^{\frac{3}{2}} \left(\frac{c_1 \sinh \left(\frac{\ln(x)}{2} \right)}{2x} + \frac{ic_2 \cosh \left(\frac{\ln(x)}{2} \right)}{2x} \right)$$

substituting $y' = 0$ and $x = 0$ in the above gives

$$0 = \frac{c_1}{2} - \frac{ic_2}{2} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = ic_2$$

Substituting these values back in above solution results in

$$y = ic_2 x^{\frac{3}{2}} \cosh \left(\frac{\ln(x)}{2} \right) + ix^{\frac{3}{2}} \sinh \left(\frac{\ln(x)}{2} \right) c_2$$

Summary

The solution(s) found are the following

$$y = i \left(\cosh \left(\frac{\ln(x)}{2} \right) + \sinh \left(\frac{\ln(x)}{2} \right) \right) c_2 x^{\frac{3}{2}} \quad (1)$$

Verification of solutions

$$y = i \left(\cosh \left(\frac{\ln(x)}{2} \right) + \sinh \left(\frac{\ln(x)}{2} \right) \right) c_2 x^{\frac{3}{2}}$$

Verified OK.

15.1.6 Solving as second order change of variable on y method 1 ode

In normal form the given ode is written as

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = -\frac{2}{x}$$
$$q(x) = \frac{2}{x^2}$$

Calculating the Liouville ode invariant Q given by

$$\begin{aligned} Q &= q - \frac{p'}{2} - \frac{p^2}{4} \\ &= \frac{2}{x^2} - \frac{\left(-\frac{2}{x}\right)'}{2} - \frac{\left(-\frac{2}{x}\right)^2}{4} \\ &= \frac{2}{x^2} - \frac{\left(\frac{2}{x^2}\right)}{2} - \frac{\left(\frac{4}{x^2}\right)}{4} \\ &= \frac{2}{x^2} - \left(\frac{1}{x^2}\right) - \frac{1}{x^2} \\ &= 0 \end{aligned}$$

Since the Liouville ode invariant does not depend on the independent variable x then the transformation

$$y = v(x)z(x) \quad (3)$$

is used to change the original ode to a constant coefficients ode in v . In (3) the term $z(x)$ is given by

$$\begin{aligned} z(x) &= e^{-\left(\int \frac{p(x)}{2} dx\right)} \\ &= e^{-\int \frac{-2}{2x}} \\ &= x \end{aligned} \tag{5}$$

Hence (3) becomes

$$y = v(x) x \tag{4}$$

Applying this change of variable to the original ode results in

$$x^3 v''(x) = 0$$

Which is now solved for $v(x)$ Integrating twice gives the solution

$$v(x) = c_1 x + c_2$$

Now that $v(x)$ is known, then

$$\begin{aligned} y &= v(x) z(x) \\ &= (c_1 x + c_2) (z(x)) \end{aligned} \tag{7}$$

But from (5)

$$z(x) = x$$

Hence (7) becomes

$$y = x(c_1 x + c_2)$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = x(c_1 x + c_2) \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = 0 \tag{1A}$$

Taking derivative of the solution gives

$$y' = 2c_1x + c_2$$

substituting $y' = 0$ and $x = 0$ in the above gives

$$0 = c_2 \tag{2A}$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_2 = 0$$

Substituting these values back in above solution results in

$$y = c_1x^2$$

Summary

The solution(s) found are the following

$$y = c_1x^2 \tag{1}$$

Verification of solutions

$$y = c_1x^2$$

Verified OK.

15.1.7 Solving as second order change of variable on y method 2 ode

In normal form the ode

$$x^2y'' - 2xy' + 2y = 0 \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \tag{2}$$

Where

$$p(x) = -\frac{2}{x}$$
$$q(x) = \frac{2}{x^2}$$

Applying change of variables on the dependent variable $y = v(x)x^n$ to (2) gives the following ode where the dependent variables is $v(x)$ and not y .

$$v''(x) + \left(\frac{2n}{x} + p\right)v'(x) + \left(\frac{n(n-1)}{x^2} + \frac{np}{x} + q\right)v(x) = 0 \tag{3}$$

Let the coefficient of $v(x)$ above be zero. Hence

$$\frac{n(n-1)}{x^2} + \frac{np}{x} + q = 0 \quad (4)$$

Substituting the earlier values found for $p(x)$ and $q(x)$ into (4) gives

$$\frac{n(n-1)}{x^2} - \frac{2n}{x^2} + \frac{2}{x^2} = 0 \quad (5)$$

Solving (5) for n gives

$$n = 2 \quad (6)$$

Substituting this value in (3) gives

$$\begin{aligned} v''(x) + \frac{2v'(x)}{x} &= 0 \\ v''(x) + \frac{2v'(x)}{x} &= 0 \end{aligned} \quad (7)$$

Using the substitution

$$u(x) = v'(x)$$

Then (7) becomes

$$u'(x) + \frac{2u(x)}{x} = 0 \quad (8)$$

The above is now solved for $u(x)$. In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{2u}{x} \end{aligned}$$

Where $f(x) = -\frac{2}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned} \frac{1}{u} du &= -\frac{2}{x} dx \\ \int \frac{1}{u} du &= \int -\frac{2}{x} dx \\ \ln(u) &= -2 \ln(x) + c_1 \\ u &= e^{-2 \ln(x) + c_1} \\ &= \frac{c_1}{x^2} \end{aligned}$$

Now that $u(x)$ is known, then

$$\begin{aligned}v'(x) &= u(x) \\v(x) &= \int u(x) dx + c_2 \\&= -\frac{c_1}{x} + c_2\end{aligned}$$

Hence

$$\begin{aligned}y &= v(x) x^n \\&= \left(-\frac{c_1}{x} + c_2\right) x^2 \\&= (c_2 x - c_1) x\end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = \left(-\frac{c_1}{x} + c_2\right) x^2 \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = 0 \quad (1A)$$

Taking derivative of the solution gives

$$y' = c_1 + 2\left(-\frac{c_1}{x} + c_2\right) x$$

substituting $y' = 0$ and $x = 0$ in the above gives

$$0 = -c_1 \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 0$$

Substituting these values back in above solution results in

$$y = c_2 x^2$$

Summary

The solution(s) found are the following

$$y = c_2 x^2 \quad (1)$$

Verification of solutions

$$y = c_2 x^2$$

Verified OK.

15.1.8 Solving as second order ode non constant coeff transformation on B ode

Given an ode of the form

$$Ay'' + By' + Cy = F(x)$$

This method reduces the order ode the ODE by one by applying the transformation

$$y = Bv$$

This results in

$$\begin{aligned} y' &= B'v + v'B \\ y'' &= B''v + B'v' + v''B + v'B' \\ &= v''B + 2v' + B' + B''v \end{aligned}$$

And now the original ode becomes

$$\begin{aligned} A(v''B + 2v'B' + B''v) + B(B'v + v'B) + CBv &= 0 \\ ABv'' + (2AB' + B^2)v' + (AB'' + BB' + CB)v &= 0 \end{aligned} \quad (1)$$

If the term $AB'' + BB' + CB$ is zero, then this method works and can be used to solve

$$ABv'' + (2AB' + B^2)v' = 0$$

By Using $u = v'$ which reduces the order of the above ode to one. The new ode is

$$ABu' + (2AB' + B^2)u = 0$$

The above ode is first order ode which is solved for u . Now a new ode $v' = u$ is solved for v as first order ode. Then the final solution is obtain from $y = Bv$.

This method works only if the term $AB'' + BB' + CB$ is zero. The given ODE shows that

$$A = x^2$$

$$B = -2x$$

$$C = 2$$

$$F = 0$$

The above shows that for this ode

$$\begin{aligned} AB'' + BB' + CB &= (x^2)(0) + (-2x)(-2) + (2)(-2x) \\ &= 0 \end{aligned}$$

Hence the ode in v given in (1) now simplifies to

$$-2x^3v'' + (0)v' = 0$$

Now by applying $v' = u$ the above becomes

$$-2x^3u'(x) = 0$$

Which is now solved for u . Integrating both sides gives

$$\begin{aligned} u(x) &= \int 0 \, dx \\ &= c_1 \end{aligned}$$

The ode for v now becomes

$$\begin{aligned} v' &= u \\ &= c_1 \end{aligned}$$

Which is now solved for v . Integrating both sides gives

$$\begin{aligned} v(x) &= \int c_1 \, dx \\ &= c_1x + c_2 \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y(x) &= Bv \\ &= (-2x)(c_1x + c_2) \\ &= -2x(c_1x + c_2) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = -2x(c_1x + c_2) \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = 0 \quad (1A)$$

Taking derivative of the solution gives

$$y' = -4c_1x - 2c_2$$

substituting $y' = 0$ and $x = 0$ in the above gives

$$0 = -2c_2 \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_2 = 0$$

Substituting these values back in above solution results in

$$y = -2c_1x^2$$

Summary

The solution(s) found are the following

$$y = -2c_1x^2 \quad (1)$$

Verification of solutions

$$y = -2c_1x^2$$

Verified OK.

15.1.9 Solving using Kovacic algorithm

Writing the ode as

$$x^2y'' - 2xy' + 2y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= x^2 \\B &= -2x \\C &= 2\end{aligned}\tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x)\tag{4}$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\&= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned}\tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1}\tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= 0 \\t &= 1\end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 0\tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 383: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - -\infty \\
 &= \infty
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}
 y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\
 &= z_1 e^{-\int \frac{1}{2} \frac{-2x}{x^2} dx} \\
 &= z_1 e^{\ln(x)} \\
 &= z_1(x)
 \end{aligned}$$

Which simplifies to

$$y_1 = x$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{2x}{x^2} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{2\ln(x)}}{(y_1)^2} dx \\ &= y_1(x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1(x) + c_2(x(x)) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_2 x^2 + c_1 x \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$0 = 0 \tag{1A}$$

Taking derivative of the solution gives

$$y' = 2c_2 x + c_1$$

substituting $y' = 0$ and $x = 0$ in the above gives

$$0 = c_1 \tag{2A}$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 0$$

Substituting these values back in above solution results in

$$y = c_2 x^2$$

Summary

The solution(s) found are the following

$$y = c_2 x^2 \tag{1}$$

Verification of solutions

$$y = c_2 x^2$$

Verified OK.

15.1.10 Maple step by step solution

Let's solve

$$\left[x^2 y'' - 2xy' + 2y = 0, y(0) = 0, y'|_{\{x=0\}} = 0 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = \frac{2y'}{x} - \frac{2y}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - \frac{2y'}{x} + \frac{2y}{x^2} = 0$$

- Multiply by denominators of the ODE

$$x^2 y'' - 2xy' + 2y = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x , using the chain rule

$$y' = \left(\frac{d}{dt} y(t) \right) t'(x)$$

- Compute derivative

$$y' = \frac{\frac{d}{dt}y(t)}{x}$$

- Calculate the 2nd derivative of y with respect to x , using the chain rule

$$y'' = \left(\frac{d^2}{dt^2}y(t) \right) t'(x)^2 + t''(x) \left(\frac{d}{dt}y(t) \right)$$

- Compute derivative

$$y'' = \frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2}$$

Substitute the change of variables back into the ODE

$$x^2 \left(\frac{\frac{d^2}{dt^2}y(t)}{x^2} - \frac{\frac{d}{dt}y(t)}{x^2} \right) - 2 \frac{d}{dt}y(t) + 2y(t) = 0$$

- Simplify

$$\frac{d^2}{dt^2}y(t) - 3 \frac{d}{dt}y(t) + 2y(t) = 0$$

- Characteristic polynomial of ODE

$$r^2 - 3r + 2 = 0$$

- Factor the characteristic polynomial

$$(r - 1)(r - 2) = 0$$

- Roots of the characteristic polynomial

$$r = (1, 2)$$

- 1st solution of the ODE

$$y_1(t) = e^t$$

- 2nd solution of the ODE

$$y_2(t) = e^{2t}$$

- General solution of the ODE

$$y(t) = c_1 y_1(t) + c_2 y_2(t)$$

- Substitute in solutions

$$y(t) = c_1 e^t + c_2 e^{2t}$$

- Change variables back using $t = \ln(x)$

$$y = c_2 x^2 + c_1 x$$

- Simplify

$$y = x(c_2 x + c_1)$$

- Check validity of solution $y = x(c_2x + c_1)$
 - Use initial condition $y(0) = 0$

$$0 = 0$$
 - Compute derivative of the solution

$$y' = 2c_2x + c_1$$
 - Use the initial condition $y' \Big|_{\{x=0\}} = 0$

$$0 = c_1$$
 - Solve for c_1 and c_2

$$\{c_1 = 0, c_2 = c_2\}$$
 - Substitute constant values into general solution and simplify

$$y = c_2x^2$$
- Solution to the IVP

$$y = c_2x^2$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
<- LODE of Euler type successful`

```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 9

```

dsolve([x^2*diff(y(x),x$2)-2*x*diff(y(x),x)+2*y(x)=0,y(0) = 0, D(y)(0) = 0],y(x), singsol=all)

```

$$y(x) = c_1x^2$$

✓ Solution by Mathematica

Time used: 0.077 (sec). Leaf size: 11

```
DSolve[{x^2*y'[x]-2*x*y'[x]+2*y[x]==0,{y[0]==0,y'[0]==0}},y[x],x,IncludeSingularSolutions
```

$$y(x) \rightarrow c_2 x^2$$

15.2 problem 4

15.2.1 Solving as second order linear constant coeff ode	2689
15.2.2 Solving using Kovacic algorithm	2692
15.2.3 Maple step by step solution	2697

Internal problem ID [6400]

Internal file name [OUTPUT/5648_Sunday_June_05_2022_03_45_56_PM_80774343/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Problems for Review and Discovery. Challenge excercises. Page 105

Problem number: 4.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + 9y = -3 \cos(2x)$$

15.2.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 9, f(x) = -3 \cos(2x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 9y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 9$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 9 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 9 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 9$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(9)} \\ &= \pm 3i \end{aligned}$$

Hence

$$\lambda_1 = +3i$$

$$\lambda_2 = -3i$$

Which simplifies to

$$\lambda_1 = 3i$$

$$\lambda_2 = -3i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 3$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (c_1 \cos(3x) + c_2 \sin(3x))$$

Or

$$y = c_1 \cos(3x) + c_2 \sin(3x)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \cos(3x) + c_2 \sin(3x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$-3 \cos(2x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(2x), \sin(2x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(3x), \sin(3x)\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(2x) + A_2 \sin(2x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$5A_1 \cos(2x) + 5A_2 \sin(2x) = -3 \cos(2x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{3}{5}, A_2 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{3 \cos(2x)}{5}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(3x) + c_2 \sin(3x)) + \left(-\frac{3 \cos(2x)}{5} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(3x) + c_2 \sin(3x) - \frac{3 \cos(2x)}{5} \quad (1)$$




Figure 425: Slope field plot

Verification of solutions

$$y = c_1 \cos(3x) + c_2 \sin(3x) - \frac{3 \cos(2x)}{5}$$

Verified OK.

15.2.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 9y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= 1 \\B &= 0 \\C &= 9\end{aligned}\tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x)\tag{4}$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\&= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned}\tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-9}{1}\tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= -9 \\t &= 1\end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -9z(x)\tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 385: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -9$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(3x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned} y_1 &= z_1 \\ &= \cos(3x) \end{aligned}$$

Which simplifies to

$$y_1 = \cos(3x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(3x) \int \frac{1}{\cos(3x)^2} dx \\ &= \cos(3x) \left(\frac{\tan(3x)}{3} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(3x)) + c_2 \left(\cos(3x) \left(\frac{\tan(3x)}{3} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$.
 y_h is the solution to

$$y'' + 9y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 \cos(3x) + \frac{c_2 \sin(3x)}{3}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$-3 \cos(2x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(2x), \sin(2x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ \frac{\sin(3x)}{3}, \cos(3x) \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(2x) + A_2 \sin(2x)$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$5A_1 \cos(2x) + 5A_2 \sin(2x) = -3 \cos(2x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{3}{5}, A_2 = 0 \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{3 \cos(2x)}{5}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(c_1 \cos(3x) + \frac{c_2 \sin(3x)}{3} \right) + \left(-\frac{3 \cos(2x)}{5} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(3x) + \frac{c_2 \sin(3x)}{3} - \frac{3 \cos(2x)}{5} \quad (1)$$




Figure 426: Slope field plot

Verification of solutions

$$y = c_1 \cos(3x) + \frac{c_2 \sin(3x)}{3} - \frac{3 \cos(2x)}{5}$$

Verified OK.

15.2.3 Maple step by step solution

Let's solve

$$y'' + 9y = -3 \cos(2x)$$

- Highest derivative means the order of the ODE is 2
- y''
- Characteristic polynomial of homogeneous ODE

$$r^2 + 9 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-36})}{2}$$

- Roots of the characteristic polynomial

$$r = (-3I, 3I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(3x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(3x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 \cos(3x) + c_2 \sin(3x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x), y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x), y_2(x))} dx \right), f(x) = -3 \cos(2x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(3x) & \sin(3x) \\ -3 \sin(3x) & 3 \cos(3x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 3$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{\cos(3x)(\int (-\sin(5x) - \sin(x)) dx)}{2} + \frac{\sin(3x)(\int (-\cos(x) - \cos(5x)) dx)}{2}$$

- Compute integrals

$$y_p(x) = -\frac{3 \cos(2x)}{5}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 \cos(3x) + c_2 \sin(3x) - \frac{3 \cos(2x)}{5}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(diff(y(x),x$2)+9*y(x)=-3*cos(2*x),y(x), singsol=all)
```

$$y(x) = \sin(3x) c_2 + \cos(3x) c_1 - \frac{3 \cos(2x)}{5}$$

✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 28

```
DSolve[y''[x]+9*y[x]==-3*Cos[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{3}{5} \cos(2x) + c_1 \cos(3x) + c_2 \sin(3x)$$

**16 Chapter 2. Problems for Review and Discovery.
Problems for Discussion and Exploration. Page
105**

16.1 problem 1	2701
16.2 problem 2	2714
16.3 problem 4	2725

16.1 problem 1

16.1.1 Solving as linear ode	2701
16.1.2 Solving as first order ode lie symmetry lookup ode	2703
16.1.3 Solving as exact ode	2707
16.1.4 Maple step by step solution	2711

Internal problem ID [6401]

Internal file name [OUTPUT/5649_Sunday_June_05_2022_03_45_58_PM_11655965/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Problems for Review and Discovery. Problems for Discussion and Exploration. Page 105

Problem number: 1.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"linear", "exactWithIntegrationFactor", "first_order_ode_lie_symmetry_lookup"**

Maple gives the following as the ode type

```
[[_linear, `class A`]]
```

$$y + y' = \cos(x)$$

16.1.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = 1$$

$$q(x) = \cos(x)$$

Hence the ode is

$$y + y' = \cos(x)$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int 1 dx} \\ &= e^x\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu) (\cos(x)) \\ \frac{d}{dx}(y e^x) &= (e^x) (\cos(x)) \\ d(y e^x) &= (\cos(x) e^x) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}y e^x &= \int \cos(x) e^x dx \\ y e^x &= \frac{\cos(x) e^x}{2} + \frac{\sin(x) e^x}{2} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^x$ results in

$$y = e^{-x} \left(\frac{\cos(x) e^x}{2} + \frac{\sin(x) e^x}{2} \right) + c_1 e^{-x}$$

which simplifies to

$$y = \frac{\sin(x)}{2} + \frac{\cos(x)}{2} + c_1 e^{-x}$$

Summary

The solution(s) found are the following

$$y = \frac{\sin(x)}{2} + \frac{\cos(x)}{2} + c_1 e^{-x} \tag{1}$$




Figure 427: Slope field plot

Verification of solutions

$$y = \frac{\sin(x)}{2} + \frac{\cos(x)}{2} + c_1 e^{-x}$$

Verified OK.

16.1.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = -y + \cos(x)$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 387: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= e^{-x}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^{-x}} dy \end{aligned}$$

Which results in

$$S = y e^x$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -y + \cos(x)$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= y e^x \\ S_y &= e^x \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \cos(x) e^x \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \cos(R) e^R$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by

integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 + \frac{e^R(\cos(R) + \sin(R))}{2} \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$e^x y = \frac{e^x(\sin(x) + \cos(x))}{2} + c_1$$



Which simplifies to

$$e^x y = \frac{e^x(\sin(x) + \cos(x))}{2} + c_1$$

Which gives

$$y = \frac{e^{-x}(\cos(x) e^x + \sin(x) e^x + 2c_1)}{2}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -y + \cos(x)$ 	$R = x$ $S = y e^x$	$\frac{dS}{dR} = \cos(R) e^R$ 

Summary

The solution(s) found are the following

$$y = \frac{e^{-x}(\cos(x) e^x + \sin(x) e^x + 2c_1)}{2} \quad (1)$$




Figure 428: Slope field plot

Verification of solutions

$$y = \frac{e^{-x}(\cos(x)e^x + \sin(x)e^x + 2c_1)}{2}$$

Verified OK.

16.1.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial\phi}{\partial x} + \frac{\partial\phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}dy &= (-y + \cos(x)) dx \\ (y - \cos(x)) dx + dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= y - \cos(x) \\ N(x, y) &= 1\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(y - \cos(x)) \\ &= 1\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(1) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= 1((1) - (0)) \\ &= 1 \end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int A dx} \\ &= e^{\int 1 dx} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^x \\ &= e^x \end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} for now so not to confuse them with the original M and N .

$$\begin{aligned} \bar{M} &= \mu M \\ &= e^x(y - \cos(x)) \\ &= (y - \cos(x)) e^x \end{aligned}$$

And

$$\begin{aligned} \bar{N} &= \mu N \\ &= e^x(1) \\ &= e^x \end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned} \bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ ((y - \cos(x)) e^x) + (e^x) \frac{dy}{dx} &= 0 \end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int (y - \cos(x)) e^x dx \\ \phi &= -\frac{e^x(-2y + \cos(x) + \sin(x))}{2} + f(y)\end{aligned}\quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = e^x + f'(y)\quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = e^x$. Therefore equation (4) becomes

$$e^x = e^x + f'(y)\quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = -\frac{e^x(-2y + \cos(x) + \sin(x))}{2} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\frac{e^x(-2y + \cos(x) + \sin(x))}{2}$$

The solution becomes

$$y = \frac{e^{-x}(\cos(x) e^x + \sin(x) e^x + 2c_1)}{2}$$

Summary

The solution(s) found are the following

$$y = \frac{e^{-x}(\cos(x) e^x + \sin(x) e^x + 2c_1)}{2} \quad (1)$$




Figure 429: Slope field plot

Verification of solutions

$$y = \frac{e^{-x}(\cos(x) e^x + \sin(x) e^x + 2c_1)}{2}$$

Verified OK.

16.1.4 Maple step by step solution

Let's solve

$$y + y' = \cos(x)$$

- Highest derivative means the order of the ODE is 1
- y'
- Isolate the derivative

$$y' = -y + \cos(x)$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y + y' = \cos(x)$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x)(y + y') = \mu(x)\cos(x)$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x)y)$

$$\mu(x)(y + y') = \mu'(x)y + \mu(x)y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = \mu(x)$$

- Solve to find the integrating factor

$$\mu(x) = e^x$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x)y) \right) dx = \int \mu(x)\cos(x) dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x)y = \int \mu(x)\cos(x) dx + c_1$$

- Solve for y

$$y = \frac{\int \mu(x)\cos(x)dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = e^x$

$$y = \frac{\int \cos(x)e^x dx + c_1}{e^x}$$

- Evaluate the integrals on the rhs

$$y = \frac{\frac{\cos(x)e^x}{2} + \frac{\sin(x)e^x}{2} + c_1}{e^x}$$

- Simplify

$$y = \frac{\sin(x)}{2} + \frac{\cos(x)}{2} + c_1e^{-x}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(diff(y(x),x)+y(x)=cos(x),y(x), singsol=all)
```

$$y(x) = \frac{\cos(x)}{2} + \frac{\sin(x)}{2} + c_1 e^{-x}$$

✓ Solution by Mathematica

Time used: 0.036 (sec). Leaf size: 23

```
DSolve[y'[x]+y[x]==Cos[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2}(\sin(x) + \cos(x) + 2c_1 e^{-x})$$

16.2 problem 2

16.2.1 Existence and uniqueness analysis	2714
16.2.2 Solving as second order linear constant coeff ode	2715
16.2.3 Solving as second order ode can be made integrable ode	2717
16.2.4 Solving using Kovacic algorithm	2719
16.2.5 Maple step by step solution	2723

Internal problem ID [6402]

Internal file name [OUTPUT/5650_Sunday_June_05_2022_03_46_00_PM_10350138/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Problems for Review and Discovery. Problems for Discussion and Exploration. Page 105

Problem number: 2.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "second_order_ode_can_be_made_integrable"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' + 3y = 0$$

With initial conditions

$$[y(0) = -1]$$

16.2.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = 0$$

$$q(x) = 3$$

$$F = 0$$

Hence the ode is

$$y'' + 3y = 0$$

The domain of $p(x) = 0$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = 3$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

16.2.2 Solving as second order linear constant coeff ode

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 3$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 3e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 3 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 3$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(3)} \\ &= \pm i\sqrt{3} \end{aligned}$$

Hence

$$\lambda_1 = +i\sqrt{3}$$

$$\lambda_2 = -i\sqrt{3}$$

Which simplifies to

$$\begin{aligned}\lambda_1 &= i\sqrt{3} \\ \lambda_2 &= -i\sqrt{3}\end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = \sqrt{3}$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0(c_1 \cos(\sqrt{3}x) + c_2 \sin(\sqrt{3}x))$$

Or

$$y = c_1 \cos(\sqrt{3}x) + c_2 \sin(\sqrt{3}x)$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 \cos(\sqrt{3}x) + c_2 \sin(\sqrt{3}x) \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = -1$ and $x = 0$ in the above gives

$$-1 = c_1 \tag{1A}$$

Equations {1A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = -1$$

Substituting these values back in above solution results in

$$y = -\cos(\sqrt{3}x) + c_2 \sin(\sqrt{3}x)$$

Summary

The solution(s) found are the following

$$y = -\cos(\sqrt{3}x) + c_2 \sin(\sqrt{3}x) \quad (1)$$

Verification of solutions

$$y = -\cos(\sqrt{3}x) + c_2 \sin(\sqrt{3}x)$$

Verified OK.

16.2.3 Solving as second order ode can be made integrable ode

Multiplying the ode by y' gives

$$y'y'' + 3y'y = 0$$

Integrating the above w.r.t x gives

$$\int (y'y'' + 3y'y) dx = 0$$
$$\frac{y'^2}{2} + \frac{3y^2}{2} = c_2$$

Which is now solved for y . Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \sqrt{-3y^2 + 2c_1} \quad (1)$$

$$y' = -\sqrt{-3y^2 + 2c_1} \quad (2)$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int \frac{1}{\sqrt{-3y^2 + 2c_1}} dy = \int dx$$
$$\frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}y}{\sqrt{-3y^2 + 2c_1}}\right)}{3} = c_2 + x$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{1}{\sqrt{-3y^2 + 2c_1}} dy = \int dx$$

$$-\frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}y}{\sqrt{-3y^2 + 2c_1}}\right)}{3} = x + c_3$$

Initial conditions are used to solve for the constants of integration.

Looking at the First solution

$$\frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}y}{\sqrt{-3y^2 + 2c_1}}\right)}{3} = c_2 + x \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = -1$ and $x = 0$ in the above gives

$$-\frac{\arctan\left(\frac{\sqrt{3}}{\sqrt{-3+2c_1}}\right) \sqrt{3}}{3} = c_2 \quad (1A)$$

Equations {1A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_2 = -\frac{\arctan\left(\frac{\sqrt{3}}{\sqrt{-3+2c_1}}\right) \sqrt{3}}{3}$$

Substituting these values back in above solution results in

$$\frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}y}{\sqrt{-3y^2 + 2c_1}}\right)}{3} = -\frac{\arctan\left(\frac{\sqrt{3}}{\sqrt{-3+2c_1}}\right) \sqrt{3}}{3} + x$$

Looking at the Second solution

$$-\frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}y}{\sqrt{-3y^2 + 2c_1}}\right)}{3} = x + c_3 \quad (2)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = -1$ and $x = 0$ in the above gives

$$\frac{\arctan\left(\frac{\sqrt{3}}{\sqrt{-3+2c_1}}\right) \sqrt{3}}{3} = c_3 \quad (1A)$$

Equations {1A} are now solved for $\{c_1, c_3\}$. Solving for the constants gives

$$c_3 = \frac{\arctan\left(\frac{\sqrt{3}}{\sqrt{-3+2c_1}}\right) \sqrt{3}}{3}$$

Substituting these values back in above solution results in

$$-\frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}y}{\sqrt{-3y^2+2c_1}}\right)}{3} = x + \frac{\arctan\left(\frac{\sqrt{3}}{\sqrt{-3+2c_1}}\right) \sqrt{3}}{3}$$

Summary

The solution(s) found are the following

$$\frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}y}{\sqrt{-3y^2+2c_1}}\right)}{3} = -\frac{\arctan\left(\frac{\sqrt{3}}{\sqrt{-3+2c_1}}\right) \sqrt{3}}{3} + x \quad (1)$$

$$-\frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}y}{\sqrt{-3y^2+2c_1}}\right)}{3} = x + \frac{\arctan\left(\frac{\sqrt{3}}{\sqrt{-3+2c_1}}\right) \sqrt{3}}{3} \quad (2)$$

Verification of solutions

$$\frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}y}{\sqrt{-3y^2+2c_1}}\right)}{3} = -\frac{\arctan\left(\frac{\sqrt{3}}{\sqrt{-3+2c_1}}\right) \sqrt{3}}{3} + x$$

Verified OK.

$$-\frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}y}{\sqrt{-3y^2+2c_1}}\right)}{3} = x + \frac{\arctan\left(\frac{\sqrt{3}}{\sqrt{-3+2c_1}}\right) \sqrt{3}}{3}$$

Verified OK.

16.2.4 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 3y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 3 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-3}{1} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -3 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -3z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 390: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -3$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(\sqrt{3}x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}
 y_1 &= z_1 \\
 &= \cos(\sqrt{3}x)
 \end{aligned}$$

Which simplifies to

$$y_1 = \cos(\sqrt{3}x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(\sqrt{3}x) \int \frac{1}{\cos^2(\sqrt{3}x)} dx \\ &= \cos(\sqrt{3}x) \left(\frac{\sqrt{3} \tan(\sqrt{3}x)}{3} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 \left(\cos(\sqrt{3}x) \right) + c_2 \left(\cos(\sqrt{3}x) \left(\frac{\sqrt{3} \tan(\sqrt{3}x)}{3} \right) \right) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = c_1 \cos(\sqrt{3}x) + \frac{c_2 \sqrt{3} \sin(\sqrt{3}x)}{3} \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = -1$ and $x = 0$ in the above gives

$$-1 = c_1 \quad (1A)$$

Equations {1A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = -1$$

Substituting these values back in above solution results in

$$y = -\cos(\sqrt{3}x) + \frac{c_2\sqrt{3}\sin(\sqrt{3}x)}{3}$$

Summary

The solution(s) found are the following

$$y = -\cos(\sqrt{3}x) + \frac{c_2\sqrt{3}\sin(\sqrt{3}x)}{3} \quad (1)$$

Verification of solutions

$$y = -\cos(\sqrt{3}x) + \frac{c_2\sqrt{3}\sin(\sqrt{3}x)}{3}$$

Verified OK.

16.2.5 Maple step by step solution

Let's solve

$$[y'' + 3y = 0, y(0) = -1]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 + 3 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-12})}{2}$$

- Roots of the characteristic polynomial

$$r = (-i\sqrt{3}, i\sqrt{3})$$

- 1st solution of the ODE

$$y_1(x) = \cos(\sqrt{3}x)$$

- 2nd solution of the ODE

$$y_2(x) = \sin(\sqrt{3}x)$$

- General solution of the ODE

$$y = c_1y_1(x) + c_2y_2(x)$$

- Substitute in solutions

$$y = c_1 \cos(\sqrt{3}x) + c_2 \sin(\sqrt{3}x)$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 21

```
dsolve([diff(y(x),x$2)=-3*y(x),y(0) = -1],y(x), singsol=all)
```

$$y(x) = c_1 \sin(\sqrt{3}x) - \cos(\sqrt{3}x)$$

✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 27

```
DSolve[{y''[x]==-3*y[x],{y[0]==-1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\cos(\sqrt{3}x) + c_2 \sin(\sqrt{3}x)$$

16.3 problem 4

16.3.1 Solving as second order ode can be made integrable ode 2725

16.3.2 Solving as second order ode missing x ode 2727

Internal problem ID [6403]

Internal file name [OUTPUT/5651_Sunday_June_05_2022_03_46_01_PM_55027395/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 2. Problems for Review and Discovery. Problems for Discussion and Exploration. Page 105

Problem number: 4.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_ode_missing_x", "second_order_ode_can_be_made_integrable"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x], [_2nd_order , _reducible , _mu_x_y1]]
```

$$y'' + \sin(y) = 0$$

16.3.1 Solving as second order ode can be made integrable ode

Multiplying the ode by y' gives

$$y'y'' + y' \sin(y) = 0$$

Integrating the above w.r.t x gives

$$\int (y'y'' + y' \sin(y)) dx = 0$$
$$\frac{y'^2}{2} - \cos(y) = c_2$$

Which is now solved for y . Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \sqrt{2 \cos(y) + 2c_1} \tag{1}$$

$$y' = -\sqrt{2 \cos(y) + 2c_1} \tag{2}$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int \frac{1}{\sqrt{2 \cos (y) + 2c_1}} dy = \int dx$$

$$\frac{2\sqrt{\frac{\cos(y)+c_1}{c_1+1}} \operatorname{InverseJacobiAM}\left(\frac{y}{2}, \frac{2}{\sqrt{2+2c_1}}\right)}{\sqrt{2 \cos (y) + 2c_1}} = c_2 + x$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{1}{\sqrt{2 \cos (y) + 2c_1}} dy = \int dx$$

$$-\frac{2\sqrt{\frac{\cos(y)+c_1}{c_1+1}} \operatorname{InverseJacobiAM}\left(\frac{y}{2}, \frac{2}{\sqrt{2+2c_1}}\right)}{\sqrt{2 \cos (y) + 2c_1}} = x + c_3$$

Summary

The solution(s) found are the following

$$\frac{2\sqrt{\frac{\cos(y)+c_1}{c_1+1}} \operatorname{InverseJacobiAM}\left(\frac{y}{2}, \frac{2}{\sqrt{2+2c_1}}\right)}{\sqrt{2 \cos (y) + 2c_1}} = c_2 + x \quad (1)$$

$$-\frac{2\sqrt{\frac{\cos(y)+c_1}{c_1+1}} \operatorname{InverseJacobiAM}\left(\frac{y}{2}, \frac{2}{\sqrt{2+2c_1}}\right)}{\sqrt{2 \cos (y) + 2c_1}} = x + c_3 \quad (2)$$

Verification of solutions

$$\frac{2\sqrt{\frac{\cos(y)+c_1}{c_1+1}} \operatorname{InverseJacobiAM}\left(\frac{y}{2}, \frac{2}{\sqrt{2+2c_1}}\right)}{\sqrt{2 \cos (y) + 2c_1}} = c_2 + x$$

Verified OK.

$$-\frac{2\sqrt{\frac{\cos(y)+c_1}{c_1+1}} \operatorname{InverseJacobiAM}\left(\frac{y}{2}, \frac{2}{\sqrt{2+2c_1}}\right)}{\sqrt{2 \cos (y) + 2c_1}} = x + c_3$$

Verified OK.

16.3.2 Solving as second order ode missing x ode

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable y an independent variable. Using

$$y' = p(y)$$

Then

$$\begin{aligned}y'' &= \frac{dp}{dx} \\ &= \frac{dy}{dx} \frac{dp}{dy} \\ &= p \frac{dp}{dy}\end{aligned}$$

Hence the ode becomes

$$p(y) \left(\frac{d}{dy} p(y) \right) = -\sin(y)$$

Which is now solved as first order ode for $p(y)$. In canonical form the ODE is

$$\begin{aligned}p' &= F(y, p) \\ &= f(y)g(p) \\ &= -\frac{\sin(y)}{p}\end{aligned}$$

Where $f(y) = -\sin(y)$ and $g(p) = \frac{1}{p}$. Integrating both sides gives

$$\begin{aligned}\frac{1}{p} dp &= -\sin(y) dy \\ \int \frac{1}{p} dp &= \int -\sin(y) dy \\ \frac{p^2}{2} &= \cos(y) + c_1\end{aligned}$$

The solution is

$$\frac{p(y)^2}{2} - \cos(y) - c_1 = 0$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$\frac{y'^2}{2} - \cos(y) - c_1 = 0$$

Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \sqrt{2 \cos(y) + 2c_1} \quad (1)$$

$$y' = -\sqrt{2 \cos(y) + 2c_1} \quad (2)$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int \frac{1}{\sqrt{2 \cos(y) + 2c_1}} dy = \int dx$$

$$\frac{2\sqrt{\frac{\cos(y)+c_1}{c_1+1}} \operatorname{InverseJacobiAM}\left(\frac{y}{2}, \frac{2}{\sqrt{2+2c_1}}\right)}{\sqrt{2 \cos(y) + 2c_1}} = c_2 + x$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{1}{\sqrt{2 \cos(y) + 2c_1}} dy = \int dx$$

$$-\frac{2\sqrt{\frac{\cos(y)+c_1}{c_1+1}} \operatorname{InverseJacobiAM}\left(\frac{y}{2}, \frac{2}{\sqrt{2+2c_1}}\right)}{\sqrt{2 \cos(y) + 2c_1}} = x + c_3$$

Summary

The solution(s) found are the following

$$\frac{2\sqrt{\frac{\cos(y)+c_1}{c_1+1}} \operatorname{InverseJacobiAM}\left(\frac{y}{2}, \frac{2}{\sqrt{2+2c_1}}\right)}{\sqrt{2 \cos(y) + 2c_1}} = c_2 + x \quad (1)$$

$$-\frac{2\sqrt{\frac{\cos(y)+c_1}{c_1+1}} \operatorname{InverseJacobiAM}\left(\frac{y}{2}, \frac{2}{\sqrt{2+2c_1}}\right)}{\sqrt{2 \cos(y) + 2c_1}} = x + c_3 \quad (2)$$

Verification of solutions

$$\frac{2\sqrt{\frac{\cos(y)+c_1}{c_1+1}} \operatorname{InverseJacobiAM}\left(\frac{y}{2}, \frac{2}{\sqrt{2+2c_1}}\right)}{\sqrt{2\cos(y)+2c_1}} = c_2 + x$$

Verified OK.

$$-\frac{2\sqrt{\frac{\cos(y)+c_1}{c_1+1}} \operatorname{InverseJacobiAM}\left(\frac{y}{2}, \frac{2}{\sqrt{2+2c_1}}\right)}{\sqrt{2\cos(y)+2c_1}} = x + c_3$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
`, `-> Computing symmetries using: way = exp_sym
-> Calling odsolve with the ODE`, (diff(_b(_a), _a))*_b(_a)+sin(_a) = 0, _b(_a)` *** Suble
  Methods for first order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  trying 1st order linear
  trying Bernoulli
  <- Bernoulli successful
  <- differential order: 2; canonical coordinates successful
  <- differential order 2; missing variables successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 47

```
dsolve(diff(y(x),x$2)+sin(y(x))=0,y(x), singsol=all)
```

$$\int^{y(x)} \frac{1}{\sqrt{2 \cos(_a) + c_1}} d_a - x - c_2 = 0$$
$$-\left(\int^{y(x)} \frac{1}{\sqrt{2 \cos(_a) + c_1}} d_a \right) - x - c_2 = 0$$

✓ Solution by Mathematica

Time used: 3.86 (sec). Leaf size: 69

```
DSolve[y''[x]+Sin[y[x]]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -2 \text{JacobiAmplitude} \left(\frac{1}{2} \sqrt{(c_1 + 2)(x + c_2)^2}, \frac{4}{c_1 + 2} \right)$$

$$y(x) \rightarrow 2 \text{JacobiAmplitude} \left(\frac{1}{2} \sqrt{(c_1 + 2)(x + c_2)^2}, \frac{4}{c_1 + 2} \right)$$

17 Chapter 4. Power Series Solutions and Special Functions. Section 4.2. Series Solutions of First-Order Differential Equations Page 162

17.1 problem 1(a) solving using series	2732
17.2 problem 1(a) solving directly	2742
17.3 problem 1(b) solving using series	2757
17.4 problem 1(b) solving directly	2769
17.5 problem 1(c) solving using series	2772
17.6 problem 1(c) solving directly	2784
17.7 problem 1(d) solving using series	2787
17.8 problem 1(d) solving directly	2797
17.9 problem 1(e) solving using series	2800
17.10 problem 1(e) solving directly	2810
17.11 problem 1(f) solving using series	2813
17.12 problem 1(f) solving directly	2825
17.13 problem 2(a) solving using series	2838
17.14 problem 2(a) solving directly	2844
17.15 problem 2(b) solving using series	2858
17.16 problem 2(b) solving directly	2861
17.17 problem 2(c) solving using series	2876
17.18 problem 2(c) solving directly	2883
17.19 problem 2(d) solving using series	2897
17.20 problem 3	2911
17.21 problem 4	2923
17.22 problem 5 solved using series	2935
17.23 problem 5 solved directly	2947

17.1 problem 1(a) solving using series

17.1.1 Solving as series ode	2732
17.1.2 Maple step by step solution	2740

Internal problem ID [6404]

Internal file name [OUTPUT/5652_Sunday_June_05_2022_03_46_04_PM_97963013/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.2. Series Solutions of First-Order Differential Equations Page 162

Problem number: 1(a) solving using series.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"first order ode series method. Ordinary point", "first order ode series method. Taylor series method"**

Maple gives the following as the ode type

`[_separable]`

$$y' - 2xy = 0$$

With the expansion point for the power series method at $x = 0$.

17.1.1 Solving as series ode

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving first order ode. Let

$$y' = f(x, y)$$

Where $f(x, y)$ is analytic at expansion point x_0 . We can always shift to $x_0 = 0$ if x_0 is not zero. So from now we assume $x_0 = 0$. Assume also that $y(x_0) = y_0$. Using Taylor

series

$$\begin{aligned}
 y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \dots \\
 &= y_0 + xf + \frac{x^2}{2} \left. \frac{df}{dx} \right|_{x_0, y_0} + \frac{x^3}{3!} \left. \frac{d^2f}{dx^2} \right|_{x_0, y_0} + \dots \\
 &= y_0 + \sum_{n=0}^{\infty} \frac{x^{n+1}}{(n+1)!} \left. \frac{d^n f}{dx^n} \right|_{x_0, y_0}
 \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} f \tag{1}$$

$$\begin{aligned}
 \frac{d^2f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) f
 \end{aligned} \tag{2}$$

$$\begin{aligned}
 \frac{d^3f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2f}{dx^2} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{d^2f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2f}{dx^2} \right) f
 \end{aligned} \tag{3}$$

⋮

And so on. Hence if we name $F_0 = f(x, y)$ then the above can be written as

$$F_0 = f(x, y) \tag{4}$$

$$\begin{aligned}
 F_n &= \frac{d}{dx}(F_{n-1}) \\
 &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) F_0
 \end{aligned} \tag{5}$$

For example, for $n = 1$ we see that

$$\begin{aligned}
 F_1 &= \frac{d}{dx}(F_0) \\
 &= \frac{\partial}{\partial x} F_0 + \left(\frac{\partial F_0}{\partial y} \right) F_0 \\
 &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} f
 \end{aligned}$$

Which is (1). And when $n = 2$

$$\begin{aligned} F_2 &= \frac{d}{dx}(F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) F_0 \\ &= \frac{\partial}{\partial x} \left(\frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} f \right) + \frac{\partial}{\partial y} \left(\frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} f \right) f \\ &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) f \end{aligned}$$

Which is (2) and so on. Therefore (4,5) can be used from now on along with

$$y(x) = y_0 + \sum_{n=0}^{\infty} \frac{x^{n+1}}{(n+1)!} F_n|_{x_0, y_0} \quad (6)$$

Hence

$$\begin{aligned}F_0 &= 2xy \\F_1 &= \frac{dF_0}{dx} \\&= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} F_0 \\&= (4x^2 + 2) y \\F_2 &= \frac{dF_1}{dx} \\&= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} F_1 \\&= (8x^3 + 12x) y \\F_3 &= \frac{dF_2}{dx} \\&= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} F_2 \\&= 4y(4x^4 + 12x^2 + 3) \\F_4 &= \frac{dF_3}{dx} \\&= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} F_3 \\&= 32y \left(x^4 + 5x^2 + \frac{15}{4} \right) x \\F_5 &= \frac{dF_4}{dx} \\&= \frac{\partial F_4}{\partial x} + \frac{\partial F_4}{\partial y} F_4 \\&= (64x^6 + 480x^4 + 720x^2 + 120) y \\F_6 &= \frac{dF_5}{dx} \\&= \frac{\partial F_5}{\partial x} + \frac{\partial F_5}{\partial y} F_5 \\&= (128x^7 + 1344x^5 + 3360x^3 + 1680x) y\end{aligned}$$

And so on. Evaluating all the above at initial conditions $x(0) = 0$ and $y(0) = y(0)$ gives

$$\begin{aligned} F_0 &= 0 \\ F_1 &= 2y(0) \\ F_2 &= 0 \\ F_3 &= 12y(0) \\ F_4 &= 0 \\ F_5 &= 120y(0) \\ F_6 &= 0 \end{aligned}$$

Substituting all the above in (6) and simplifying gives the solution as

$$y = \left(1 + x^2 + \frac{1}{2}x^4 + \frac{1}{6}x^6 \right) y(0) + O(x^8)$$

Since $x = 0$ is also an ordinary point, then standard power series can also be used. Writing the ODE as

$$\begin{aligned} y' + q(x)y &= p(x) \\ y' - 2xy &= 0 \end{aligned}$$

Where

$$\begin{aligned} q(x) &= -2x \\ p(x) &= 0 \end{aligned}$$

Next, the type of the expansion point $x = 0$ is determined. This point can be an ordinary point, a regular singular point (also called removable singularity), or irregular singular point (also called non-removable singularity or essential singularity). When $x = 0$ is an ordinary point, then the standard power series is used. If the point is a regular singular point, Frobenius series is used instead. Irregular singular point requires more advanced methods (asymptotic methods) and is not supported now. Hopefully this will be added in the future. $x = 0$ is called an ordinary point $q(x)$ has a Taylor series expansion around the point $x = 0$. $x = 0$ is called a regular singular point if $q(x)$ is not analytic at $x = 0$ but $xq(x)$ has Taylor series expansion. And finally, $x = 0$ is an irregular singular point if the point is not ordinary and not regular singular. This is the most complicated case. Now the expansion point $x = 0$ is checked to see if it is an ordinary point or not. Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$y' = \sum_{n=1}^{\infty} n a_n x^{n-1}$$

Substituting the above back into the ode gives

$$\left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) - 2x \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) + \sum_{n=0}^{\infty} (-2x^{1+n} a_n) = 0 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=1}^{\infty} n a_n x^{n-1} &= \sum_{n=0}^{\infty} (1+n) a_{1+n} x^n \\ \sum_{n=0}^{\infty} (-2x^{1+n} a_n) &= \sum_{n=1}^{\infty} (-2a_{n-1} x^n) \end{aligned}$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} (1+n) a_{1+n} x^n \right) + \sum_{n=1}^{\infty} (-2a_{n-1} x^n) = 0 \quad (3)$$

For $1 \leq n$, the recurrence equation is

$$(1+n) a_{1+n} - 2a_{n-1} = 0 \quad (4)$$

Solving for a_{1+n} , gives

$$a_{1+n} = \frac{2a_{n-1}}{1+n} \quad (5)$$

For $n = 1$ the recurrence equation gives

$$2a_2 - 2a_0 = 0$$

Which after substituting the earlier terms found becomes

$$a_2 = a_0$$

For $n = 2$ the recurrence equation gives

$$3a_3 - 2a_1 = 0$$

Which after substituting the earlier terms found becomes

$$a_3 = 0$$

For $n = 3$ the recurrence equation gives

$$4a_4 - 2a_2 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = \frac{a_0}{2}$$

For $n = 4$ the recurrence equation gives

$$5a_5 - 2a_3 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = 0$$

For $n = 5$ the recurrence equation gives

$$6a_6 - 2a_4 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = \frac{a_0}{6}$$

For $n = 6$ the recurrence equation gives

$$7a_7 - 2a_5 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = 0$$

For $n = 7$ the recurrence equation gives

$$8a_8 - 2a_6 = 0$$

Which after substituting the earlier terms found becomes

$$a_8 = \frac{a_0}{24}$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + a_0 x^2 + \frac{1}{2} a_0 x^4 + \frac{1}{6} a_0 x^6 + \dots$$

Collecting terms, the solution becomes

$$y = \left(1 + x^2 + \frac{1}{2} x^4 + \frac{1}{6} x^6 \right) a_0 + O(x^8) \quad (3)$$

Summary

The solution(s) found are the following

$$y = \left(1 + x^2 + \frac{1}{2} x^4 + \frac{1}{6} x^6 \right) y(0) + O(x^8) \quad (1)$$

$$y = \left(1 + x^2 + \frac{1}{2} x^4 + \frac{1}{6} x^6 \right) c_1 + O(x^8) \quad (2)$$




Figure 430: Slope field plot

Verification of solutions

$$y = \left(1 + x^2 + \frac{1}{2}x^4 + \frac{1}{6}x^6\right) y(0) + O(x^8)$$

Verified OK.

$$y = \left(1 + x^2 + \frac{1}{2}x^4 + \frac{1}{6}x^6\right) c_1 + O(x^8)$$

Verified OK.

17.1.2 Maple step by step solution

Let's solve

$$y' - 2xy = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y} = 2x$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y} dx = \int 2x dx + c_1$$

- Evaluate integral

$$\ln(y) = x^2 + c_1$$

- Solve for y

$$y = e^{x^2+c_1}$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 24

```

Order:=8;
dsolve(diff(y(x),x)=2*x*y(x),y(x),type='series',x=0);

```

$$y(x) = \left(1 + x^2 + \frac{1}{2}x^4 + \frac{1}{6}x^6\right) y(0) + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 25

```

AsymptoticDSolveValue[y'[x]==2*x*y[x],y[x],{x,0,7}]

```

$$y(x) \rightarrow c_1 \left(\frac{x^6}{6} + \frac{x^4}{2} + x^2 + 1 \right)$$

17.2 problem 1(a) solving directly

17.2.1 Solving as separable ode	2742
17.2.2 Solving as linear ode	2744
17.2.3 Solving as homogeneousTypeD2 ode	2745
17.2.4 Solving as first order ode lie symmetry lookup ode	2747
17.2.5 Solving as exact ode	2751
17.2.6 Maple step by step solution	2755

Internal problem ID [6405]

Internal file name [OUTPUT/5653_Sunday_June_05_2022_03_46_06_PM_96428372/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.2. Series Solutions of First-Order Differential Equations Page 162

Problem number: 1(a) solving directly.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "linear", "separable", "homogeneousTypeD2", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_separable]

$$y' - 2xy = 0$$

17.2.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= 2xy\end{aligned}$$

Where $f(x) = 2x$ and $g(y) = y$. Integrating both sides gives

$$\begin{aligned}\frac{1}{y} dy &= 2x dx \\ \int \frac{1}{y} dy &= \int 2x dx \\ \ln(y) &= x^2 + c_1 \\ y &= e^{x^2 + c_1} \\ &= c_1 e^{x^2}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{x^2} \tag{1}$$




Figure 431: Slope field plot

Verification of solutions

$$y = c_1 e^{x^2}$$

Verified OK.

17.2.2 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -2x$$

$$q(x) = 0$$

Hence the ode is

$$y' - 2xy = 0$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int -2x dx} \\ &= e^{-x^2}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}\mu y &= 0 \\ \frac{d}{dx}(e^{-x^2}y) &= 0\end{aligned}$$

Integrating gives

$$e^{-x^2}y = c_1$$

Dividing both sides by the integrating factor $\mu = e^{-x^2}$ results in

$$y = c_1 e^{x^2}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{x^2} \tag{1}$$




Figure 432: Slope field plot

Verification of solutions

$$y = c_1 e^{x^2}$$

Verified OK.

17.2.3 Solving as homogeneousTypeD2 ode

Using the change of variables $y = u(x)x$ on the above ode results in new ode in $u(x)$

$$u'(x)x + u(x) - 2x^2u(x) = 0$$

In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= \frac{u(2x^2 - 1)}{x} \end{aligned}$$

Where $f(x) = \frac{2x^2-1}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u} du &= \frac{2x^2-1}{x} dx \\ \int \frac{1}{u} du &= \int \frac{2x^2-1}{x} dx \\ \ln(u) &= x^2 - \ln(x) + c_2 \\ u &= e^{x^2 - \ln(x) + c_2} \\ &= c_2 e^{x^2 - \ln(x)}\end{aligned}$$

Which simplifies to

$$u(x) = \frac{c_2 e^{x^2}}{x}$$

Therefore the solution y is

$$\begin{aligned}y &= xu \\ &= c_2 e^{x^2}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_2 e^{x^2} \tag{1}$$




Figure 433: Slope field plot

Verification of solutions

$$y = c_2 e^{x^2}$$

Verified OK.

17.2.4 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$\begin{aligned} y' &= 2xy \\ y' &= \omega(x, y) \end{aligned}$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 393: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= e^{x^2}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right)S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^{x^2}} dy \end{aligned}$$

Which results in

$$S = e^{-x^2} y$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = 2xy$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= -2x e^{-x^2} y \\ S_y &= e^{-x^2} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by

integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \tag{4}$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$e^{-x^2} y = c_1$$



Which simplifies to

$$e^{-x^2} y = c_1$$

Which gives

$$y = c_1 e^{x^2}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = 2xy$ 	$R = x$ $S = e^{-x^2} y$	$\frac{dS}{dR} = 0$ 

Summary

The solution(s) found are the following

$$y = c_1 e^{x^2} \tag{1}$$




Figure 434: Slope field plot

Verification of solutions

$$y = c_1 e^{x^2}$$

Verified OK.

17.2.5 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}\left(\frac{1}{2y}\right) dy &= (x) dx \\ (-x) dx + \left(\frac{1}{2y}\right) dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -x \\ N(x, y) &= \frac{1}{2y}\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-x) \\ &= 0\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{1}{2y} \right) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \tag{1}$$

$$\frac{\partial \phi}{\partial y} = N \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -x dx \\ \phi &= -\frac{x^2}{2} + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{1}{2y}$. Therefore equation (4) becomes

$$\frac{1}{2y} = 0 + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = \frac{1}{2y}$$

Integrating the above w.r.t y gives

$$\int f'(y) dy = \int \left(\frac{1}{2y} \right) dy$$
$$f(y) = \frac{\ln(y)}{2} + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\frac{x^2}{2} + \frac{\ln(y)}{2} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\frac{x^2}{2} + \frac{\ln(y)}{2}$$

The solution becomes

$$y = e^{x^2+2c_1}$$

Summary

The solution(s) found are the following

$$y = e^{x^2+2c_1} \tag{1}$$




Figure 435: Slope field plot

Verification of solutions

$$y = e^{x^2+2c_1}$$

Verified OK.

17.2.6 Maple step by step solution

Let's solve

$$y' - 2xy = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y} = 2x$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y} dx = \int 2x dx + c_1$$

- Evaluate integral

- $\ln(y) = x^2 + c_1$
Solve for y
 $y = e^{x^2+c_1}$

Maple trace

```
`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 10

```
dsolve(diff(y(x),x)=2*x*y(x),y(x), singsol=all)
```

$$y(x) = e^{x^2} c_1$$

✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 18

```
DSolve[y'[x]==2*x*y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{x^2}$$

$$y(x) \rightarrow 0$$

17.3 problem 1(b) solving using series

17.3.1 Solving as series ode	2757
17.3.2 Maple step by step solution	2767

Internal problem ID [6406]

Internal file name [OUTPUT/5654_Sunday_June_05_2022_03_46_07_PM_96437827/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.2. Series Solutions of First-Order Differential Equations Page 162

Problem number: 1(b) solving using series.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"first order ode series method. Ordinary point", "first order ode series method. Taylor series method"**

Maple gives the following as the ode type

[_quadrature]

$$y + y' = 1$$

With the expansion point for the power series method at $x = 0$.

17.3.1 Solving as series ode

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving first order ode. Let

$$y' = f(x, y)$$

Where $f(x, y)$ is analytic at expansion point x_0 . We can always shift to $x_0 = 0$ if x_0 is not zero. So from now we assume $x_0 = 0$. Assume also that $y(x_0) = y_0$. Using Taylor

series

$$\begin{aligned}
 y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \dots \\
 &= y_0 + xf + \frac{x^2}{2} \left. \frac{df}{dx} \right|_{x_0, y_0} + \frac{x^3}{3!} \left. \frac{d^2f}{dx^2} \right|_{x_0, y_0} + \dots \\
 &= y_0 + \sum_{n=0}^{\infty} \frac{x^{n+1}}{(n+1)!} \left. \frac{d^n f}{dx^n} \right|_{x_0, y_0}
 \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} f \tag{1}$$

$$\begin{aligned}
 \frac{d^2f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) f
 \end{aligned} \tag{2}$$

$$\begin{aligned}
 \frac{d^3f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2f}{dx^2} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{d^2f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2f}{dx^2} \right) f
 \end{aligned} \tag{3}$$

⋮

And so on. Hence if we name $F_0 = f(x, y)$ then the above can be written as

$$F_0 = f(x, y) \tag{4}$$

$$\begin{aligned}
 F_n &= \frac{d}{dx}(F_{n-1}) \\
 &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) F_0
 \end{aligned} \tag{5}$$

For example, for $n = 1$ we see that

$$\begin{aligned}
 F_1 &= \frac{d}{dx}(F_0) \\
 &= \frac{\partial}{\partial x} F_0 + \left(\frac{\partial F_0}{\partial y} \right) F_0 \\
 &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} f
 \end{aligned}$$

Which is (1). And when $n = 2$

$$\begin{aligned} F_2 &= \frac{d}{dx}(F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) F_0 \\ &= \frac{\partial}{\partial x} \left(\frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} f \right) + \frac{\partial}{\partial y} \left(\frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} f \right) f \\ &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) f \end{aligned}$$

Which is (2) and so on. Therefore (4,5) can be used from now on along with

$$y(x) = y_0 + \sum_{n=0}^{\infty} \frac{x^{n+1}}{(n+1)!} F_n|_{x_0, y_0} \quad (6)$$

Hence

$$\begin{aligned}F_0 &= 1 - y \\F_1 &= \frac{dF_0}{dx} \\&= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} F_0 \\&= y - 1 \\F_2 &= \frac{dF_1}{dx} \\&= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} F_1 \\&= 1 - y \\F_3 &= \frac{dF_2}{dx} \\&= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} F_2 \\&= y - 1 \\F_4 &= \frac{dF_3}{dx} \\&= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} F_3 \\&= 1 - y \\F_5 &= \frac{dF_4}{dx} \\&= \frac{\partial F_4}{\partial x} + \frac{\partial F_4}{\partial y} F_4 \\&= y - 1 \\F_6 &= \frac{dF_5}{dx} \\&= \frac{\partial F_5}{\partial x} + \frac{\partial F_5}{\partial y} F_5 \\&= 1 - y\end{aligned}$$

And so on. Evaluating all the above at initial conditions $x(0) = 0$ and $y(0) = y(0)$ gives

$$F_0 = 1 - y(0)$$

$$F_1 = y(0) - 1$$

$$F_2 = 1 - y(0)$$

$$F_3 = y(0) - 1$$

$$F_4 = 1 - y(0)$$

$$F_5 = y(0) - 1$$

$$F_6 = 1 - y(0)$$

Substituting all the above in (6) and simplifying gives the solution as

$$y = \left(1 - x + \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{24}x^4 - \frac{1}{120}x^5 + \frac{1}{720}x^6 - \frac{1}{5040}x^7 \right) y(0) \\ + x - \frac{x^2}{2} + \frac{x^3}{6} - \frac{x^4}{24} + \frac{x^5}{120} - \frac{x^6}{720} + \frac{x^7}{5040} + O(x^8)$$

Since $x = 0$ is also an ordinary point, then standard power series can also be used. Writing the ODE as

$$y' + q(x)y = p(x) \\ y + y' = 1$$

Where

$$q(x) = 1 \\ p(x) = 1$$

Next, the type of the expansion point $x = 0$ is determined. This point can be an ordinary point, a regular singular point (also called removable singularity), or irregular singular point (also called non-removable singularity or essential singularity). When $x = 0$ is an ordinary point, then the standard power series is used. If the point is a regular singular point, Frobenius series is used instead. Irregular singular point requires more advanced methods (asymptotic methods) and is not supported now. Hopefully this will be added in the future. $x = 0$ is called an ordinary point $q(x)$ has a Taylor series expansion around the point $x = 0$. $x = 0$ is called a regular singular point if $q(x)$ is not analytic at $x = 0$ but $xq(x)$ has Taylor series expansion. And finally, $x = 0$ is an irregular singular point if the point is not ordinary and not regular singular. This

is the most complicated case. Now the expansion point $x = 0$ is checked to see if it is an ordinary point or not. Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$y' = \sum_{n=1}^{\infty} n a_n x^{n-1}$$

Substituting the above back into the ode gives

$$\left(\sum_{n=0}^{\infty} a_n x^n \right) + \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) = 1 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} a_n x^n \right) + \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) = 1 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} (n+1) a_{n+1} x^n \right) + \left(\sum_{n=0}^{\infty} a_n x^n \right) = 1 \quad (3)$$

For $0 \leq n$, the recurrence equation is

$$((n+1) a_{n+1} + a_n) x^n = 1 \quad (4)$$

For $n = 0$ the recurrence equation gives

$$\begin{aligned} (a_1 + a_0) 1 &= 1 \\ a_1 + a_0 &= 1 \end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_1 = 1 - a_0$$

For $n = 1$ the recurrence equation gives

$$\begin{aligned}(2a_2 + a_1)x &= 0 \\ 2a_2 + a_1 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_2 = -\frac{1}{2} + \frac{a_0}{2}$$

For $n = 2$ the recurrence equation gives

$$\begin{aligned}(3a_3 + a_2)x^2 &= 0 \\ 3a_3 + a_2 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_3 = \frac{1}{6} - \frac{a_0}{6}$$

For $n = 3$ the recurrence equation gives

$$\begin{aligned}(4a_4 + a_3)x^3 &= 0 \\ 4a_4 + a_3 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_4 = -\frac{1}{24} + \frac{a_0}{24}$$

For $n = 4$ the recurrence equation gives

$$\begin{aligned}(5a_5 + a_4)x^4 &= 0 \\ 5a_5 + a_4 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_5 = \frac{1}{120} - \frac{a_0}{120}$$

For $n = 5$ the recurrence equation gives

$$\begin{aligned}(6a_6 + a_5)x^5 &= 0 \\ 6a_6 + a_5 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_6 = -\frac{1}{720} + \frac{a_0}{720}$$

For $n = 6$ the recurrence equation gives

$$\begin{aligned}(7a_7 + a_6)x^6 &= 0 \\ 7a_7 + a_6 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_7 = \frac{1}{5040} - \frac{a_0}{5040}$$

For $n = 7$ the recurrence equation gives

$$\begin{aligned}(8a_8 + a_7)x^7 &= 0 \\ 8a_8 + a_7 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_8 = -\frac{1}{40320} + \frac{a_0}{40320}$$

And so on. Therefore the solution is

$$\begin{aligned}y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots\end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + (1 - a_0)x + \left(-\frac{1}{2} + \frac{a_0}{2}\right)x^2 + \left(\frac{1}{6} - \frac{a_0}{6}\right)x^3 + \left(-\frac{1}{24} + \frac{a_0}{24}\right)x^4 \\ + \left(\frac{1}{120} - \frac{a_0}{120}\right)x^5 + \left(-\frac{1}{720} + \frac{a_0}{720}\right)x^6 + \left(\frac{1}{5040} - \frac{a_0}{5040}\right)x^7 + \dots$$

Collecting terms, the solution becomes

$$y = \left(1 - x + \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{24}x^4 - \frac{1}{120}x^5 + \frac{1}{720}x^6 - \frac{1}{5040}x^7\right)a_0 \quad (3) \\ + x - \frac{x^2}{2} + \frac{x^3}{6} - \frac{x^4}{24} + \frac{x^5}{120} - \frac{x^6}{720} + \frac{x^7}{5040} + O(x^8)$$

Summary

The solution(s) found are the following

$$y = \left(1 - x + \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{24}x^4 - \frac{1}{120}x^5 + \frac{1}{720}x^6 - \frac{1}{5040}x^7\right)y(0) \quad (1) \\ + x - \frac{x^2}{2} + \frac{x^3}{6} - \frac{x^4}{24} + \frac{x^5}{120} - \frac{x^6}{720} + \frac{x^7}{5040} + O(x^8)$$

$$y = \left(1 - x + \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{24}x^4 - \frac{1}{120}x^5 + \frac{1}{720}x^6 - \frac{1}{5040}x^7\right)c_1 \quad (2) \\ + x - \frac{x^2}{2} + \frac{x^3}{6} - \frac{x^4}{24} + \frac{x^5}{120} - \frac{x^6}{720} + \frac{x^7}{5040} + O(x^8)$$




Figure 436: Slope field plot

Verification of solutions

$$y = \left(1 - x + \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{24}x^4 - \frac{1}{120}x^5 + \frac{1}{720}x^6 - \frac{1}{5040}x^7 \right) y(0) + x - \frac{x^2}{2} + \frac{x^3}{6} - \frac{x^4}{24} + \frac{x^5}{120} - \frac{x^6}{720} + \frac{x^7}{5040} + O(x^8)$$

Verified OK.

$$y = \left(1 - x + \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{24}x^4 - \frac{1}{120}x^5 + \frac{1}{720}x^6 - \frac{1}{5040}x^7 \right) c_1 + x - \frac{x^2}{2} + \frac{x^3}{6} - \frac{x^4}{24} + \frac{x^5}{120} - \frac{x^6}{720} + \frac{x^7}{5040} + O(x^8)$$

Verified OK.

17.3.2 Maple step by step solution

Let's solve

$$y + y' = 1$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{1-y} = 1$$

- Integrate both sides with respect to x

$$\int \frac{y'}{1-y} dx = \int 1 dx + c_1$$

- Evaluate integral

$$-\ln(1-y) = x + c_1$$

- Solve for y

$$y = -e^{-c_1-x} + 1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 60

```
Order:=8;  
dsolve(diff(y(x),x)+y(x)=1,y(x),type='series',x=0);
```

$$y(x) = \left(1 - x + \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{24}x^4 - \frac{1}{120}x^5 + \frac{1}{720}x^6 - \frac{1}{5040}x^7\right) y(0) \\ + x - \frac{x^2}{2} + \frac{x^3}{6} - \frac{x^4}{24} + \frac{x^5}{120} - \frac{x^6}{720} + \frac{x^7}{5040} + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 97

```
AsymptoticDSolveValue[y'[x]+y[x]==1,y[x],{x,0,7}]
```

$$y(x) \rightarrow \frac{x^7}{5040} - \frac{x^6}{720} + \frac{x^5}{120} - \frac{x^4}{24} + \frac{x^3}{6} - \frac{x^2}{2} + c_1 \left(-\frac{x^7}{5040} + \frac{x^6}{720} - \frac{x^5}{120} + \frac{x^4}{24} - \frac{x^3}{6} + \frac{x^2}{2} - x + 1 \right) + x$$

17.4 problem 1(b) solving directly

17.4.1 Solving as quadrature ode	2769
17.4.2 Maple step by step solution	2770

Internal problem ID [6407]

Internal file name [OUTPUT/5655_Sunday_June_05_2022_03_46_09_PM_61613456/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.2. Series Solutions of First-Order Differential Equations Page 162

Problem number: 1(b) solving directly.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

`[_quadrature]`

$$y + y' = 1$$

17.4.1 Solving as quadrature ode

Integrating both sides gives

$$\int \frac{1}{1-y} dy = \int dx$$
$$-\ln(1-y) = x + c_1$$

Raising both side to exponential gives

$$\frac{1}{1-y} = e^{x+c_1}$$

Which simplifies to

$$\frac{1}{1-y} = c_2 e^x$$

Summary

The solution(s) found are the following

$$y = -\frac{e^{-x}}{c_2} + 1 \quad (1)$$




Figure 437: Slope field plot

Verification of solutions

$$y = -\frac{e^{-x}}{c_2} + 1$$

Verified OK.

17.4.2 Maple step by step solution

Let's solve

$$y + y' = 1$$

- Highest derivative means the order of the ODE is 1
- y'
- Separate variables

$$\frac{y'}{1-y} = 1$$

- Integrate both sides with respect to x

$$\int \frac{y'}{1-y} dx = \int 1 dx + c_1$$

- Evaluate integral

$$-\ln(1-y) = x + c_1$$

- Solve for y

$$y = -e^{-c_1-x} + 1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(y(x),x)+y(x)=1,y(x), singsol=all)
```

$$y(x) = 1 + c_1 e^{-x}$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 20

```
DSolve[y'[x]+y[x]==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 1 + c_1 e^{-x}$$

$$y(x) \rightarrow 1$$

17.5 problem 1(c) solving using series

17.5.1 Solving as series ode	2772
17.5.2 Maple step by step solution	2782

Internal problem ID [6408]

Internal file name [OUTPUT/5656_Sunday_June_05_2022_03_46_10_PM_64285777/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.2. Series Solutions of First-Order Differential Equations Page 162

Problem number: 1(c) solving using series.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"first order ode series method. Ordinary point", "first order ode series method. Taylor series method"**

Maple gives the following as the ode type

`[_quadrature]`

$$y' - y = 2$$

With the expansion point for the power series method at $x = 0$.

17.5.1 Solving as series ode

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving first order ode. Let

$$y' = f(x, y)$$

Where $f(x, y)$ is analytic at expansion point x_0 . We can always shift to $x_0 = 0$ if x_0 is not zero. So from now we assume $x_0 = 0$. Assume also that $y(x_0) = y_0$. Using Taylor

series

$$\begin{aligned}
 y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \dots \\
 &= y_0 + xf + \frac{x^2}{2} \left. \frac{df}{dx} \right|_{x_0, y_0} + \frac{x^3}{3!} \left. \frac{d^2f}{dx^2} \right|_{x_0, y_0} + \dots \\
 &= y_0 + \sum_{n=0}^{\infty} \frac{x^{n+1}}{(n+1)!} \left. \frac{d^n f}{dx^n} \right|_{x_0, y_0}
 \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} f \tag{1}$$

$$\begin{aligned}
 \frac{d^2f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) f
 \end{aligned} \tag{2}$$

$$\begin{aligned}
 \frac{d^3f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2f}{dx^2} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{d^2f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2f}{dx^2} \right) f
 \end{aligned} \tag{3}$$

⋮

And so on. Hence if we name $F_0 = f(x, y)$ then the above can be written as

$$F_0 = f(x, y) \tag{4}$$

$$\begin{aligned}
 F_n &= \frac{d}{dx}(F_{n-1}) \\
 &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) F_0
 \end{aligned} \tag{5}$$

For example, for $n = 1$ we see that

$$\begin{aligned}
 F_1 &= \frac{d}{dx}(F_0) \\
 &= \frac{\partial}{\partial x} F_0 + \left(\frac{\partial F_0}{\partial y} \right) F_0 \\
 &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} f
 \end{aligned}$$

Which is (1). And when $n = 2$

$$\begin{aligned} F_2 &= \frac{d}{dx}(F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) F_0 \\ &= \frac{\partial}{\partial x} \left(\frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} f \right) + \frac{\partial}{\partial y} \left(\frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} f \right) f \\ &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) f \end{aligned}$$

Which is (2) and so on. Therefore (4,5) can be used from now on along with

$$y(x) = y_0 + \sum_{n=0}^{\infty} \frac{x^{n+1}}{(n+1)!} F_n|_{x_0, y_0} \quad (6)$$

Hence

$$\begin{aligned}F_0 &= 2 + y \\F_1 &= \frac{dF_0}{dx} \\&= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} F_0 \\&= 2 + y \\F_2 &= \frac{dF_1}{dx} \\&= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} F_1 \\&= 2 + y \\F_3 &= \frac{dF_2}{dx} \\&= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} F_2 \\&= 2 + y \\F_4 &= \frac{dF_3}{dx} \\&= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} F_3 \\&= 2 + y \\F_5 &= \frac{dF_4}{dx} \\&= \frac{\partial F_4}{\partial x} + \frac{\partial F_4}{\partial y} F_4 \\&= 2 + y \\F_6 &= \frac{dF_5}{dx} \\&= \frac{\partial F_5}{\partial x} + \frac{\partial F_5}{\partial y} F_5 \\&= 2 + y\end{aligned}$$

And so on. Evaluating all the above at initial conditions $x(0) = 0$ and $y(0) = y(0)$ gives

$$F_0 = y(0) + 2$$

$$F_1 = y(0) + 2$$

$$F_2 = y(0) + 2$$

$$F_3 = y(0) + 2$$

$$F_4 = y(0) + 2$$

$$F_5 = y(0) + 2$$

$$F_6 = y(0) + 2$$

Substituting all the above in (6) and simplifying gives the solution as

$$y = \left(1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{720}x^6 + \frac{1}{5040}x^7 \right) y(0) \\ + 2x + x^2 + \frac{x^3}{3} + \frac{x^4}{12} + \frac{x^5}{60} + \frac{x^6}{360} + \frac{x^7}{2520} + O(x^8)$$

Since $x = 0$ is also an ordinary point, then standard power series can also be used. Writing the ODE as

$$y' + q(x)y = p(x) \\ y' - y = 2$$

Where

$$q(x) = -1 \\ p(x) = 2$$

Next, the type of the expansion point $x = 0$ is determined. This point can be an ordinary point, a regular singular point (also called removable singularity), or irregular singular point (also called non-removable singularity or essential singularity). When $x = 0$ is an ordinary point, then the standard power series is used. If the point is a regular singular point, Frobenius series is used instead. Irregular singular point requires more advanced methods (asymptotic methods) and is not supported now. Hopefully this will be added in the future. $x = 0$ is called an ordinary point $q(x)$ has a Taylor series expansion around the point $x = 0$. $x = 0$ is called a regular singular point if $q(x)$ is not not analytic at $x = 0$ but $xq(x)$ has Taylor series expansion. And finally, $x = 0$ is an irregular singular point if the point is not ordinary and not regular singular. This

is the most complicated case. Now the expansion point $x = 0$ is checked to see if it is an ordinary point or not. Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$y' = \sum_{n=1}^{\infty} n a_n x^{n-1}$$

Substituting the above back into the ode gives

$$\left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) - \left(\sum_{n=0}^{\infty} a_n x^n \right) = 2 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) + \sum_{n=0}^{\infty} (-a_n x^n) = 2 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} (n+1) a_{n+1} x^n \right) + \sum_{n=0}^{\infty} (-a_n x^n) = 2 \quad (3)$$

For $0 \leq n$, the recurrence equation is

$$((n+1) a_{n+1} - a_n) x^n = 2 \quad (4)$$

For $n = 0$ the recurrence equation gives

$$\begin{aligned} (a_1 - a_0) 1 &= 2 \\ a_1 - a_0 &= 2 \end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_1 = 2 + a_0$$

For $n = 1$ the recurrence equation gives

$$\begin{aligned}(2a_2 - a_1)x &= 0 \\ 2a_2 - a_1 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_2 = 1 + \frac{a_0}{2}$$

For $n = 2$ the recurrence equation gives

$$\begin{aligned}(3a_3 - a_2)x^2 &= 0 \\ 3a_3 - a_2 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_3 = \frac{1}{3} + \frac{a_0}{6}$$

For $n = 3$ the recurrence equation gives

$$\begin{aligned}(4a_4 - a_3)x^3 &= 0 \\ 4a_4 - a_3 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_4 = \frac{1}{12} + \frac{a_0}{24}$$

For $n = 4$ the recurrence equation gives

$$\begin{aligned}(5a_5 - a_4)x^4 &= 0 \\ 5a_5 - a_4 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_5 = \frac{1}{60} + \frac{a_0}{120}$$

For $n = 5$ the recurrence equation gives

$$\begin{aligned}(6a_6 - a_5)x^5 &= 0 \\ 6a_6 - a_5 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_6 = \frac{1}{360} + \frac{a_0}{720}$$

For $n = 6$ the recurrence equation gives

$$\begin{aligned}(7a_7 - a_6)x^6 &= 0 \\ 7a_7 - a_6 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_7 = \frac{1}{2520} + \frac{a_0}{5040}$$

For $n = 7$ the recurrence equation gives

$$\begin{aligned}(8a_8 - a_7)x^7 &= 0 \\ 8a_8 - a_7 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_8 = \frac{1}{20160} + \frac{a_0}{40320}$$

And so on. Therefore the solution is

$$\begin{aligned}y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots\end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + (2 + a_0)x + \left(1 + \frac{a_0}{2}\right)x^2 + \left(\frac{1}{3} + \frac{a_0}{6}\right)x^3 + \left(\frac{1}{12} + \frac{a_0}{24}\right)x^4 \\ + \left(\frac{1}{60} + \frac{a_0}{120}\right)x^5 + \left(\frac{1}{360} + \frac{a_0}{720}\right)x^6 + \left(\frac{1}{2520} + \frac{a_0}{5040}\right)x^7 + \dots$$

Collecting terms, the solution becomes

$$y = \left(1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{720}x^6 + \frac{1}{5040}x^7\right)a_0 \quad (3) \\ + 2x + x^2 + \frac{x^3}{3} + \frac{x^4}{12} + \frac{x^5}{60} + \frac{x^6}{360} + \frac{x^7}{2520} + O(x^8)$$

Summary

The solution(s) found are the following

$$y = \left(1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{720}x^6 + \frac{1}{5040}x^7\right)y(0) \quad (1) \\ + 2x + x^2 + \frac{x^3}{3} + \frac{x^4}{12} + \frac{x^5}{60} + \frac{x^6}{360} + \frac{x^7}{2520} + O(x^8)$$

$$y = \left(1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{720}x^6 + \frac{1}{5040}x^7\right)c_1 \quad (2) \\ + 2x + x^2 + \frac{x^3}{3} + \frac{x^4}{12} + \frac{x^5}{60} + \frac{x^6}{360} + \frac{x^7}{2520} + O(x^8)$$




Figure 438: Slope field plot

Verification of solutions

$$y = \left(1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{720}x^6 + \frac{1}{5040}x^7 \right) y(0) + 2x + x^2 + \frac{x^3}{3} + \frac{x^4}{12} + \frac{x^5}{60} + \frac{x^6}{360} + \frac{x^7}{2520} + O(x^8)$$

Verified OK.

$$y = \left(1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{720}x^6 + \frac{1}{5040}x^7 \right) c_1 + 2x + x^2 + \frac{x^3}{3} + \frac{x^4}{12} + \frac{x^5}{60} + \frac{x^6}{360} + \frac{x^7}{2520} + O(x^8)$$

Verified OK.

17.5.2 Maple step by step solution

Let's solve

$$y' - y = 2$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{2+y} = 1$$

- Integrate both sides with respect to x

$$\int \frac{y'}{2+y} dx = \int 1 dx + c_1$$

- Evaluate integral

$$\ln(2 + y) = x + c_1$$

- Solve for y

$$y = e^{x+c_1} - 2$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 58

```
Order:=8;  
dsolve(diff(y(x),x)-y(x)=2,y(x),type='series',x=0);
```

$$y(x) = \left(1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{720}x^6 + \frac{1}{5040}x^7\right) y(0) \\ + 2x + x^2 + \frac{x^3}{3} + \frac{x^4}{12} + \frac{x^5}{60} + \frac{x^6}{360} + \frac{x^7}{2520} + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 93

```
AsymptoticDSolveValue[y'[x]-y[x]==2,y[x],{x,0,7}]
```

$$y(x) \rightarrow \frac{x^7}{2520} + \frac{x^6}{360} + \frac{x^5}{60} + \frac{x^4}{12} + \frac{x^3}{3} + x^2 + c_1 \left(\frac{x^7}{5040} + \frac{x^6}{720} + \frac{x^5}{120} + \frac{x^4}{24} + \frac{x^3}{6} + \frac{x^2}{2} + x + 1 \right) + 2x$$

17.6 problem 1(c) solving directly

17.6.1 Solving as quadrature ode	2784
17.6.2 Maple step by step solution	2785

Internal problem ID [6409]

Internal file name [OUTPUT/5657_Sunday_June_05_2022_03_46_12_PM_90120699/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.2. Series Solutions of First-Order Differential Equations Page 162

Problem number: 1(c) solving directly.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$y' - y = 2$$

17.6.1 Solving as quadrature ode

Integrating both sides gives

$$\int \frac{1}{y+2} dy = \int dx$$
$$\ln(y+2) = x + c_1$$

Raising both side to exponential gives

$$y + 2 = e^{x+c_1}$$

Which simplifies to

$$y + 2 = c_2 e^x$$

Summary

The solution(s) found are the following

$$y = c_2 e^x - 2 \tag{1}$$




Figure 439: Slope field plot

Verification of solutions

$$y = c_2 e^x - 2$$

Verified OK.

17.6.2 Maple step by step solution

Let's solve

$$y' - y = 2$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{2+y} = 1$$

- Integrate both sides with respect to x

$$\int \frac{y'}{2+y} dx = \int 1 dx + c_1$$

- Evaluate integral

- $\ln(2 + y) = x + c_1$
Solve for y
 $y = e^{x+c_1} - 2$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 10

```
dsolve(diff(y(x),x)-y(x)=2,y(x), singsol=all)
```

$$y(x) = -2 + e^x c_1$$

✓ Solution by Mathematica

Time used: 0.035 (sec). Leaf size: 18

```
DSolve[y'[x]-y[x]==2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -2 + c_1 e^x$$

$$y(x) \rightarrow -2$$

17.7 problem 1(d) solving using series

17.7.1 Solving as series ode	2787
17.7.2 Maple step by step solution	2795

Internal problem ID [6410]

Internal file name [OUTPUT/5658_Sunday_June_05_2022_03_46_13_PM_14284799/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.2. Series Solutions of First-Order Differential Equations Page 162

Problem number: 1(d) solving using series.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"first order ode series method. Ordinary point", "first order ode series method. Taylor series method"**

Maple gives the following as the ode type

`[_quadrature]`

$$y + y' = 0$$

With the expansion point for the power series method at $x = 0$.

17.7.1 Solving as series ode

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving first order ode. Let

$$y' = f(x, y)$$

Where $f(x, y)$ is analytic at expansion point x_0 . We can always shift to $x_0 = 0$ if x_0 is not zero. So from now we assume $x_0 = 0$. Assume also that $y(x_0) = y_0$. Using Taylor

series

$$\begin{aligned}
 y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \dots \\
 &= y_0 + xf + \frac{x^2}{2} \left. \frac{df}{dx} \right|_{x_0, y_0} + \frac{x^3}{3!} \left. \frac{d^2f}{dx^2} \right|_{x_0, y_0} + \dots \\
 &= y_0 + \sum_{n=0}^{\infty} \frac{x^{n+1}}{(n+1)!} \left. \frac{d^n f}{dx^n} \right|_{x_0, y_0}
 \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} f \tag{1}$$

$$\begin{aligned}
 \frac{d^2f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) f
 \end{aligned} \tag{2}$$

$$\begin{aligned}
 \frac{d^3f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2f}{dx^2} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{d^2f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2f}{dx^2} \right) f
 \end{aligned} \tag{3}$$

⋮

And so on. Hence if we name $F_0 = f(x, y)$ then the above can be written as

$$F_0 = f(x, y) \tag{4}$$

$$\begin{aligned}
 F_n &= \frac{d}{dx}(F_{n-1}) \\
 &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) F_0
 \end{aligned} \tag{5}$$

For example, for $n = 1$ we see that

$$\begin{aligned}
 F_1 &= \frac{d}{dx}(F_0) \\
 &= \frac{\partial}{\partial x} F_0 + \left(\frac{\partial F_0}{\partial y} \right) F_0 \\
 &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} f
 \end{aligned}$$

Which is (1). And when $n = 2$

$$\begin{aligned} F_2 &= \frac{d}{dx}(F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) F_0 \\ &= \frac{\partial}{\partial x} \left(\frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} f \right) + \frac{\partial}{\partial y} \left(\frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} f \right) f \\ &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) f \end{aligned}$$

Which is (2) and so on. Therefore (4,5) can be used from now on along with

$$y(x) = y_0 + \sum_{n=0}^{\infty} \frac{x^{n+1}}{(n+1)!} F_n|_{x_0, y_0} \quad (6)$$

Hence

$$\begin{aligned}F_0 &= -y \\F_1 &= \frac{dF_0}{dx} \\&= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} F_0 \\&= y \\F_2 &= \frac{dF_1}{dx} \\&= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} F_1 \\&= -y \\F_3 &= \frac{dF_2}{dx} \\&= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} F_2 \\&= y \\F_4 &= \frac{dF_3}{dx} \\&= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} F_3 \\&= -y \\F_5 &= \frac{dF_4}{dx} \\&= \frac{\partial F_4}{\partial x} + \frac{\partial F_4}{\partial y} F_4 \\&= y \\F_6 &= \frac{dF_5}{dx} \\&= \frac{\partial F_5}{\partial x} + \frac{\partial F_5}{\partial y} F_5 \\&= -y\end{aligned}$$

And so on. Evaluating all the above at initial conditions $x(0) = 0$ and $y(0) = y(0)$ gives

$$F_0 = -y(0)$$

$$F_1 = y(0)$$

$$F_2 = -y(0)$$

$$F_3 = y(0)$$

$$F_4 = -y(0)$$

$$F_5 = y(0)$$

$$F_6 = -y(0)$$

Substituting all the above in (6) and simplifying gives the solution as

$$y = \left(1 - x + \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{24}x^4 - \frac{1}{120}x^5 + \frac{1}{720}x^6 - \frac{1}{5040}x^7 \right) y(0) + O(x^8)$$

Since $x = 0$ is also an ordinary point, then standard power series can also be used. Writing the ODE as

$$y' + q(x)y = p(x)$$

$$y + y' = 0$$

Where

$$q(x) = 1$$

$$p(x) = 0$$

Next, the type of the expansion point $x = 0$ is determined. This point can be an ordinary point, a regular singular point (also called removable singularity), or irregular singular point (also called non-removable singularity or essential singularity). When $x = 0$ is an ordinary point, then the standard power series is used. If the point is a regular singular point, Frobenius series is used instead. Irregular singular point requires more advanced methods (asymptotic methods) and is not supported now. Hopefully this will be added in the future. $x = 0$ is called an ordinary point $q(x)$ has a Taylor series expansion around the point $x = 0$. $x = 0$ is called a regular singular point if $q(x)$ is not analytic at $x = 0$ but $xq(x)$ has Taylor series expansion. And finally, $x = 0$ is an irregular singular point if the point is not ordinary and not regular singular. This is the most complicated case. Now the expansion point $x = 0$ is checked to see if it is an ordinary point or not. Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$y' = \sum_{n=1}^{\infty} n a_n x^{n-1}$$

Substituting the above back into the ode gives

$$\left(\sum_{n=0}^{\infty} a_n x^n \right) + \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} a_n x^n \right) + \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) = 0 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} (n+1) a_{n+1} x^n \right) + \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0 \quad (3)$$

For $0 \leq n$, the recurrence equation is

$$(n+1) a_{n+1} + a_n = 0 \quad (4)$$

Solving for a_{n+1} , gives

$$a_{n+1} = -\frac{a_n}{n+1} \quad (5)$$

For $n = 0$ the recurrence equation gives

$$a_1 + a_0 = 0$$

Which after substituting the earlier terms found becomes

$$a_1 = -a_0$$

For $n = 1$ the recurrence equation gives

$$2a_2 + a_1 = 0$$

Which after substituting the earlier terms found becomes

$$a_2 = \frac{a_0}{2}$$

For $n = 2$ the recurrence equation gives

$$3a_3 + a_2 = 0$$

Which after substituting the earlier terms found becomes

$$a_3 = -\frac{a_0}{6}$$

For $n = 3$ the recurrence equation gives

$$4a_4 + a_3 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = \frac{a_0}{24}$$

For $n = 4$ the recurrence equation gives

$$5a_5 + a_4 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = -\frac{a_0}{120}$$

For $n = 5$ the recurrence equation gives

$$6a_6 + a_5 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = \frac{a_0}{720}$$

For $n = 6$ the recurrence equation gives

$$7a_7 + a_6 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = -\frac{a_0}{5040}$$

For $n = 7$ the recurrence equation gives

$$8a_8 + a_7 = 0$$

Which after substituting the earlier terms found becomes

$$a_8 = \frac{a_0}{40320}$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 - a_0 x + \frac{1}{2} a_0 x^2 - \frac{1}{6} a_0 x^3 + \frac{1}{24} a_0 x^4 - \frac{1}{120} a_0 x^5 + \frac{1}{720} a_0 x^6 - \frac{1}{5040} a_0 x^7 + \dots$$

Collecting terms, the solution becomes

$$y = \left(1 - x + \frac{1}{2} x^2 - \frac{1}{6} x^3 + \frac{1}{24} x^4 - \frac{1}{120} x^5 + \frac{1}{720} x^6 - \frac{1}{5040} x^7 \right) a_0 + O(x^8) \quad (3)$$

Summary

The solution(s) found are the following

$$y = \left(1 - x + \frac{1}{2} x^2 - \frac{1}{6} x^3 + \frac{1}{24} x^4 - \frac{1}{120} x^5 + \frac{1}{720} x^6 - \frac{1}{5040} x^7 \right) y(0) + O(x^8) \quad (1)$$

$$y = \left(1 - x + \frac{1}{2} x^2 - \frac{1}{6} x^3 + \frac{1}{24} x^4 - \frac{1}{120} x^5 + \frac{1}{720} x^6 - \frac{1}{5040} x^7 \right) c_1 + O(x^8) \quad (2)$$




Figure 440: Slope field plot

Verification of solutions

$$y = \left(1 - x + \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{24}x^4 - \frac{1}{120}x^5 + \frac{1}{720}x^6 - \frac{1}{5040}x^7\right) y(0) + O(x^8)$$

Verified OK.

$$y = \left(1 - x + \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{24}x^4 - \frac{1}{120}x^5 + \frac{1}{720}x^6 - \frac{1}{5040}x^7\right) c_1 + O(x^8)$$

Verified OK.

17.7.2 Maple step by step solution

Let's solve

$$y + y' = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y} = -1$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y} dx = \int (-1) dx + c_1$$

- Evaluate integral

$$\ln(y) = -x + c_1$$

- Solve for y

$$y = e^{-x+c_1}$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 46

```

Order:=8;
dsolve(diff(y(x),x)+y(x)=0,y(x),type='series',x=0);

```

$$y(x) = \left(1 - x + \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{24}x^4 - \frac{1}{120}x^5 + \frac{1}{720}x^6 - \frac{1}{5040}x^7\right) y(0) + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 53

```

AsymptoticDSolveValue[y'[x]+y[x]==0,y[x],{x,0,7}]

```

$$y(x) \rightarrow c_1 \left(-\frac{x^7}{5040} + \frac{x^6}{720} - \frac{x^5}{120} + \frac{x^4}{24} - \frac{x^3}{6} + \frac{x^2}{2} - x + 1 \right)$$

17.8 problem 1(d) solving directly

17.8.1 Solving as quadrature ode 2797

17.8.2 Maple step by step solution 2798

Internal problem ID [6411]

Internal file name [OUTPUT/5659_Sunday_June_05_2022_03_46_15_PM_41740201/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.2. Series Solutions of First-Order Differential Equations Page 162

Problem number: 1(d) solving directly.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[**_quadrature**]

$$y + y' = 0$$

17.8.1 Solving as quadrature ode

Integrating both sides gives

$$\int -\frac{1}{y} dy = \int dx$$
$$-\ln(y) = x + c_1$$

Raising both side to exponential gives

$$\frac{1}{y} = e^{x+c_1}$$

Which simplifies to

$$\frac{1}{y} = c_2 e^x$$

Summary

The solution(s) found are the following

$$y = \frac{e^{-x}}{c_2} \quad (1)$$




Figure 441: Slope field plot

Verification of solutions

$$y = \frac{e^{-x}}{c_2}$$

Verified OK.

17.8.2 Maple step by step solution

Let's solve

$$y + y' = 0$$

- Highest derivative means the order of the ODE is 1
- y'
- Separate variables

$$\frac{y'}{y} = -1$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y} dx = \int (-1) dx + c_1$$

- Evaluate integral

$$\ln(y) = -x + c_1$$

- Solve for y

$$y = e^{-x+c_1}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 10

```
dsolve(diff(y(x),x)+y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 e^{-x}$$

✓ Solution by Mathematica

Time used: 0.021 (sec). Leaf size: 18

```
DSolve[y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-x}$$

$$y(x) \rightarrow 0$$

17.9 problem 1(e) solving using series

17.9.1 Solving as series ode	2800
17.9.2 Maple step by step solution	2808

Internal problem ID [6412]

Internal file name [OUTPUT/5660_Sunday_June_05_2022_03_46_16_PM_73875935/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.2. Series Solutions of First-Order Differential Equations Page 162

Problem number: 1(e) solving using series.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"first order ode series method. Ordinary point", "first order ode series method. Taylor series method"**

Maple gives the following as the ode type

`[_quadrature]`

$$y' - y = 0$$

With the expansion point for the power series method at $x = 0$.

17.9.1 Solving as series ode

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving first order ode. Let

$$y' = f(x, y)$$

Where $f(x, y)$ is analytic at expansion point x_0 . We can always shift to $x_0 = 0$ if x_0 is not zero. So from now we assume $x_0 = 0$. Assume also that $y(x_0) = y_0$. Using Taylor

series

$$\begin{aligned}
 y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \dots \\
 &= y_0 + xf + \frac{x^2}{2} \left. \frac{df}{dx} \right|_{x_0, y_0} + \frac{x^3}{3!} \left. \frac{d^2f}{dx^2} \right|_{x_0, y_0} + \dots \\
 &= y_0 + \sum_{n=0}^{\infty} \frac{x^{n+1}}{(n+1)!} \left. \frac{d^n f}{dx^n} \right|_{x_0, y_0}
 \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} f \tag{1}$$

$$\begin{aligned}
 \frac{d^2f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) f
 \end{aligned} \tag{2}$$

$$\begin{aligned}
 \frac{d^3f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2f}{dx^2} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{d^2f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2f}{dx^2} \right) f
 \end{aligned} \tag{3}$$

⋮

And so on. Hence if we name $F_0 = f(x, y)$ then the above can be written as

$$F_0 = f(x, y) \tag{4}$$

$$\begin{aligned}
 F_n &= \frac{d}{dx}(F_{n-1}) \\
 &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) F_0
 \end{aligned} \tag{5}$$

For example, for $n = 1$ we see that

$$\begin{aligned}
 F_1 &= \frac{d}{dx}(F_0) \\
 &= \frac{\partial}{\partial x} F_0 + \left(\frac{\partial F_0}{\partial y} \right) F_0 \\
 &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} f
 \end{aligned}$$

Which is (1). And when $n = 2$

$$\begin{aligned} F_2 &= \frac{d}{dx}(F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) F_0 \\ &= \frac{\partial}{\partial x} \left(\frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} f \right) + \frac{\partial}{\partial y} \left(\frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} f \right) f \\ &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) f \end{aligned}$$

Which is (2) and so on. Therefore (4,5) can be used from now on along with

$$y(x) = y_0 + \sum_{n=0}^{\infty} \frac{x^{n+1}}{(n+1)!} F_n|_{x_0, y_0} \quad (6)$$

Hence

$$\begin{aligned}F_0 &= y \\F_1 &= \frac{dF_0}{dx} \\&= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} F_0 \\&= y \\F_2 &= \frac{dF_1}{dx} \\&= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} F_1 \\&= y \\F_3 &= \frac{dF_2}{dx} \\&= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} F_2 \\&= y \\F_4 &= \frac{dF_3}{dx} \\&= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} F_3 \\&= y \\F_5 &= \frac{dF_4}{dx} \\&= \frac{\partial F_4}{\partial x} + \frac{\partial F_4}{\partial y} F_4 \\&= y \\F_6 &= \frac{dF_5}{dx} \\&= \frac{\partial F_5}{\partial x} + \frac{\partial F_5}{\partial y} F_5 \\&= y\end{aligned}$$

And so on. Evaluating all the above at initial conditions $x(0) = 0$ and $y(0) = y(0)$ gives

$$F_0 = y(0)$$

$$F_1 = y(0)$$

$$F_2 = y(0)$$

$$F_3 = y(0)$$

$$F_4 = y(0)$$

$$F_5 = y(0)$$

$$F_6 = y(0)$$

Substituting all the above in (6) and simplifying gives the solution as

$$y = \left(1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{720}x^6 + \frac{1}{5040}x^7\right) y(0) + O(x^8)$$

Since $x = 0$ is also an ordinary point, then standard power series can also be used. Writing the ODE as

$$y' + q(x)y = p(x)$$

$$y' - y = 0$$

Where

$$q(x) = -1$$

$$p(x) = 0$$

Next, the type of the expansion point $x = 0$ is determined. This point can be an ordinary point, a regular singular point (also called removable singularity), or irregular singular point (also called non-removable singularity or essential singularity). When $x = 0$ is an ordinary point, then the standard power series is used. If the point is a regular singular point, Frobenius series is used instead. Irregular singular point requires more advanced methods (asymptotic methods) and is not supported now. Hopefully this will be added in the future. $x = 0$ is called an ordinary point $q(x)$ has a Taylor series expansion around the point $x = 0$. $x = 0$ is called a regular singular point if $q(x)$ is not not analytic at $x = 0$ but $xq(x)$ has Taylor series expansion. And finally, $x = 0$ is an irregular singular point if the point is not ordinary and not regular singular. This is the most complicated case. Now the expansion point $x = 0$ is checked to see if it is an ordinary point or not. Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$y' = \sum_{n=1}^{\infty} n a_n x^{n-1}$$

Substituting the above back into the ode gives

$$\left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) - \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) + \sum_{n=0}^{\infty} (-a_n x^n) = 0 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} (n+1) a_{n+1} x^n \right) + \sum_{n=0}^{\infty} (-a_n x^n) = 0 \quad (3)$$

For $0 \leq n$, the recurrence equation is

$$(n+1) a_{n+1} - a_n = 0 \quad (4)$$

Solving for a_{n+1} , gives

$$a_{n+1} = \frac{a_n}{n+1} \quad (5)$$

For $n = 0$ the recurrence equation gives

$$a_1 - a_0 = 0$$

Which after substituting the earlier terms found becomes

$$a_1 = a_0$$

For $n = 1$ the recurrence equation gives

$$2a_2 - a_1 = 0$$

Which after substituting the earlier terms found becomes

$$a_2 = \frac{a_0}{2}$$

For $n = 2$ the recurrence equation gives

$$3a_3 - a_2 = 0$$

Which after substituting the earlier terms found becomes

$$a_3 = \frac{a_0}{6}$$

For $n = 3$ the recurrence equation gives

$$4a_4 - a_3 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = \frac{a_0}{24}$$

For $n = 4$ the recurrence equation gives

$$5a_5 - a_4 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = \frac{a_0}{120}$$

For $n = 5$ the recurrence equation gives

$$6a_6 - a_5 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = \frac{a_0}{720}$$

For $n = 6$ the recurrence equation gives

$$7a_7 - a_6 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = \frac{a_0}{5040}$$

For $n = 7$ the recurrence equation gives

$$8a_8 - a_7 = 0$$

Which after substituting the earlier terms found becomes

$$a_8 = \frac{a_0}{40320}$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + a_0 x + \frac{1}{2} a_0 x^2 + \frac{1}{6} a_0 x^3 + \frac{1}{24} a_0 x^4 + \frac{1}{120} a_0 x^5 + \frac{1}{720} a_0 x^6 + \frac{1}{5040} a_0 x^7 + \dots$$

Collecting terms, the solution becomes

$$y = \left(1 + x + \frac{1}{2} x^2 + \frac{1}{6} x^3 + \frac{1}{24} x^4 + \frac{1}{120} x^5 + \frac{1}{720} x^6 + \frac{1}{5040} x^7 \right) a_0 + O(x^8) \quad (3)$$

Summary

The solution(s) found are the following

$$y = \left(1 + x + \frac{1}{2} x^2 + \frac{1}{6} x^3 + \frac{1}{24} x^4 + \frac{1}{120} x^5 + \frac{1}{720} x^6 + \frac{1}{5040} x^7 \right) y(0) + O(x^8) \quad (1)$$

$$y = \left(1 + x + \frac{1}{2} x^2 + \frac{1}{6} x^3 + \frac{1}{24} x^4 + \frac{1}{120} x^5 + \frac{1}{720} x^6 + \frac{1}{5040} x^7 \right) c_1 + O(x^8) \quad (2)$$




Figure 442: Slope field plot

Verification of solutions

$$y = \left(1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{720}x^6 + \frac{1}{5040}x^7\right) y(0) + O(x^8)$$

Verified OK.

$$y = \left(1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{720}x^6 + \frac{1}{5040}x^7\right) c_1 + O(x^8)$$

Verified OK.

17.9.2 Maple step by step solution

Let's solve

$$y' - y = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y} = 1$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y} dx = \int 1 dx + c_1$$
- Evaluate integral

$$\ln(y) = x + c_1$$
- Solve for y

$$y = e^{x+c_1}$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 44

```

Order:=8;
dsolve(diff(y(x),x)-y(x)=0,y(x),type='series',x=0);

```

$$y(x) = \left(1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{720}x^6 + \frac{1}{5040}x^7\right) y(0) + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 51

```

AsymptoticDSolveValue[y'[x]-y[x]==0,y[x],{x,0,7}]

```

$$y(x) \rightarrow c_1 \left(\frac{x^7}{5040} + \frac{x^6}{720} + \frac{x^5}{120} + \frac{x^4}{24} + \frac{x^3}{6} + \frac{x^2}{2} + x + 1 \right)$$

17.10 problem 1(e) solving directly

17.10.1 Solving as quadrature ode 2810

17.10.2 Maple step by step solution 2811

Internal problem ID [6413]

Internal file name [OUTPUT/5661_Sunday_June_05_2022_03_46_18_PM_50855044/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.2. Series Solutions of First-Order Differential Equations Page 162

Problem number: 1(e) solving directly.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

`[_quadrature]`

$$y' - y = 0$$

17.10.1 Solving as quadrature ode

Integrating both sides gives

$$\int \frac{1}{y} dy = x + c_1$$

$$\ln(y) = x + c_1$$

$$y = e^{x+c_1}$$

$$y = c_1 e^x$$

Summary

The solution(s) found are the following

$$y = c_1 e^x \tag{1}$$




Figure 443: Slope field plot

Verification of solutions

$$y = c_1 e^x$$

Verified OK.

17.10.2 Maple step by step solution

Let's solve

$$y' - y = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y} = 1$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y} dx = \int 1 dx + c_1$$

- Evaluate integral

- $\ln(y) = x + c_1$
Solve for y
 $y = e^{x+c_1}$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 8

```
dsolve(diff(y(x),x)-y(x)=0,y(x), singsol=all)
```

$$y(x) = e^x c_1$$

✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 16

```
DSolve[y'[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^x$$

$$y(x) \rightarrow 0$$

17.11 problem 1(f) solving using series

17.11.1 Solving as series ode	2813
17.11.2 Maple step by step solution	2823

Internal problem ID [6414]

Internal file name [OUTPUT/5662_Sunday_June_05_2022_03_46_19_PM_71072901/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.2. Series Solutions of First-Order Differential Equations Page 162

Problem number: 1(f) solving using series.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"first order ode series method. Ordinary point", "first order ode series method. Taylor series method"**

Maple gives the following as the ode type

```
[[_linear, `class A`]]
```

$$y' - y = x^2$$

With the expansion point for the power series method at $x = 0$.

17.11.1 Solving as series ode

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving first order ode. Let

$$y' = f(x, y)$$

Where $f(x, y)$ is analytic at expansion point x_0 . We can always shift to $x_0 = 0$ if x_0 is not zero. So from now we assume $x_0 = 0$. Assume also that $y(x_0) = y_0$. Using Taylor

series

$$\begin{aligned}
 y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \dots \\
 &= y_0 + xf + \frac{x^2}{2} \left. \frac{df}{dx} \right|_{x_0, y_0} + \frac{x^3}{3!} \left. \frac{d^2f}{dx^2} \right|_{x_0, y_0} + \dots \\
 &= y_0 + \sum_{n=0}^{\infty} \frac{x^{n+1}}{(n+1)!} \left. \frac{d^n f}{dx^n} \right|_{x_0, y_0}
 \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} f \tag{1}$$

$$\begin{aligned}
 \frac{d^2f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) f
 \end{aligned} \tag{2}$$

$$\begin{aligned}
 \frac{d^3f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2f}{dx^2} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{d^2f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2f}{dx^2} \right) f
 \end{aligned} \tag{3}$$

⋮

And so on. Hence if we name $F_0 = f(x, y)$ then the above can be written as

$$F_0 = f(x, y) \tag{4}$$

$$\begin{aligned}
 F_n &= \frac{d}{dx}(F_{n-1}) \\
 &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) F_0
 \end{aligned} \tag{5}$$

For example, for $n = 1$ we see that

$$\begin{aligned}
 F_1 &= \frac{d}{dx}(F_0) \\
 &= \frac{\partial}{\partial x} F_0 + \left(\frac{\partial F_0}{\partial y} \right) F_0 \\
 &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} f
 \end{aligned}$$

Which is (1). And when $n = 2$

$$\begin{aligned} F_2 &= \frac{d}{dx}(F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) F_0 \\ &= \frac{\partial}{\partial x} \left(\frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} f \right) + \frac{\partial}{\partial y} \left(\frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} f \right) f \\ &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) f \end{aligned}$$

Which is (2) and so on. Therefore (4,5) can be used from now on along with

$$y(x) = y_0 + \sum_{n=0}^{\infty} \frac{x^{n+1}}{(n+1)!} F_n|_{x_0, y_0} \quad (6)$$

Hence

$$F_0 = x^2 + y$$

$$\begin{aligned} F_1 &= \frac{dF_0}{dx} \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} F_0 \\ &= 2x + x^2 + y \end{aligned}$$

$$\begin{aligned} F_2 &= \frac{dF_1}{dx} \\ &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} F_1 \\ &= 2 + 2x + x^2 + y \end{aligned}$$

$$\begin{aligned} F_3 &= \frac{dF_2}{dx} \\ &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} F_2 \\ &= 2 + 2x + x^2 + y \end{aligned}$$

$$\begin{aligned} F_4 &= \frac{dF_3}{dx} \\ &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} F_3 \\ &= 2 + 2x + x^2 + y \end{aligned}$$

$$\begin{aligned} F_5 &= \frac{dF_4}{dx} \\ &= \frac{\partial F_4}{\partial x} + \frac{\partial F_4}{\partial y} F_4 \\ &= 2 + 2x + x^2 + y \end{aligned}$$

$$\begin{aligned} F_6 &= \frac{dF_5}{dx} \\ &= \frac{\partial F_5}{\partial x} + \frac{\partial F_5}{\partial y} F_5 \\ &= 2 + 2x + x^2 + y \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x(0) = 0$ and $y(0) = y(0)$ gives

$$\begin{aligned} F_0 &= y(0) \\ F_1 &= y(0) \\ F_2 &= y(0) + 2 \\ F_3 &= y(0) + 2 \\ F_4 &= y(0) + 2 \\ F_5 &= y(0) + 2 \\ F_6 &= y(0) + 2 \end{aligned}$$

Substituting all the above in (6) and simplifying gives the solution as

$$\begin{aligned} y &= \left(1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{720}x^6 + \frac{1}{5040}x^7 \right) y(0) \\ &\quad + \frac{x^3}{3} + \frac{x^4}{12} + \frac{x^5}{60} + \frac{x^6}{360} + \frac{x^7}{2520} + O(x^8) \end{aligned}$$

Since $x = 0$ is also an ordinary point, then standard power series can also be used. Writing the ODE as

$$\begin{aligned} y' + q(x)y &= p(x) \\ y' - y &= x^2 \end{aligned}$$

Where

$$\begin{aligned} q(x) &= -1 \\ p(x) &= x^2 \end{aligned}$$

Next, the type of the expansion point $x = 0$ is determined. This point can be an ordinary point, a regular singular point (also called removable singularity), or irregular singular point (also called non-removable singularity or essential singularity). When $x = 0$ is an ordinary point, then the standard power series is used. If the point is a regular singular point, Frobenius series is used instead. Irregular singular point requires more advanced methods (asymptotic methods) and is not supported now. Hopefully this will be added in the future. $x = 0$ is called an ordinary point $q(x)$ has a Taylor series expansion around the point $x = 0$. $x = 0$ is called a regular singular point if $q(x)$ is not not analytic at $x = 0$ but $xq(x)$ has Taylor series expansion. And finally, $x = 0$ is an irregular singular point if the point is not ordinary and not regular singular. This

is the most complicated case. Now the expansion point $x = 0$ is checked to see if it is an ordinary point or not. Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$y' = \sum_{n=1}^{\infty} n a_n x^{n-1}$$

Substituting the above back into the ode gives

$$\left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) - \left(\sum_{n=0}^{\infty} a_n x^n \right) = x^2 \quad (1)$$

Expanding x^2 as Taylor series around $x = 0$ and keeping only the first 8 terms gives

$$\begin{aligned} x^2 &= x^2 + \dots \\ &= x^2 \end{aligned}$$

Hence the ODE in Eq (1) becomes

$$\left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) - \left(\sum_{n=0}^{\infty} a_n x^n \right) = x^2 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) + \sum_{n=0}^{\infty} (-a_n x^n) = x^2 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} (n+1) a_{n+1} x^n \right) + \sum_{n=0}^{\infty} (-a_n x^n) = x^2 \quad (3)$$

For $0 \leq n$, the recurrence equation is

$$((n + 1) a_{n+1} - a_n) x^n = x^2 \quad (4)$$

For $n = 0$ the recurrence equation gives

$$\begin{aligned} (a_1 - a_0) 1 &= 0 \\ a_1 - a_0 &= 0 \end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_1 = a_0$$

For $n = 1$ the recurrence equation gives

$$\begin{aligned} (2a_2 - a_1) x &= 0 \\ 2a_2 - a_1 &= 0 \end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_2 = \frac{a_0}{2}$$

For $n = 2$ the recurrence equation gives

$$\begin{aligned} (3a_3 - a_2) x^2 &= x^2 \\ 3a_3 - a_2 &= 1 \end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_3 = \frac{1}{3} + \frac{a_0}{6}$$

For $n = 3$ the recurrence equation gives

$$\begin{aligned} (4a_4 - a_3) x^3 &= 0 \\ 4a_4 - a_3 &= 0 \end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_4 = \frac{1}{12} + \frac{a_0}{24}$$

For $n = 4$ the recurrence equation gives

$$\begin{aligned}(5a_5 - a_4)x^4 &= 0 \\ 5a_5 - a_4 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_5 = \frac{1}{60} + \frac{a_0}{120}$$

For $n = 5$ the recurrence equation gives

$$\begin{aligned}(6a_6 - a_5)x^5 &= 0 \\ 6a_6 - a_5 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_6 = \frac{1}{360} + \frac{a_0}{720}$$

For $n = 6$ the recurrence equation gives

$$\begin{aligned}(7a_7 - a_6)x^6 &= 0 \\ 7a_7 - a_6 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_7 = \frac{1}{2520} + \frac{a_0}{5040}$$

For $n = 7$ the recurrence equation gives

$$\begin{aligned}(8a_8 - a_7)x^7 &= 0 \\ 8a_8 - a_7 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_8 = \frac{1}{20160} + \frac{a_0}{40320}$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$\begin{aligned} y &= a_0 + a_0 x + \frac{a_0 x^2}{2} + \left(\frac{1}{3} + \frac{a_0}{6}\right) x^3 + \left(\frac{1}{12} + \frac{a_0}{24}\right) x^4 \\ &\quad + \left(\frac{1}{60} + \frac{a_0}{120}\right) x^5 + \left(\frac{1}{360} + \frac{a_0}{720}\right) x^6 + \left(\frac{1}{2520} + \frac{a_0}{5040}\right) x^7 + \dots \end{aligned}$$

Collecting terms, the solution becomes

$$\begin{aligned} y &= \left(1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{720}x^6 + \frac{1}{5040}x^7\right) a_0 \\ &\quad + \frac{x^3}{3} + \frac{x^4}{12} + \frac{x^5}{60} + \frac{x^6}{360} + \frac{x^7}{2520} + O(x^8) \end{aligned} \quad (3)$$

Summary

The solution(s) found are the following

$$\begin{aligned} y &= \left(1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{720}x^6 + \frac{1}{5040}x^7\right) y(0) \\ &\quad + \frac{x^3}{3} + \frac{x^4}{12} + \frac{x^5}{60} + \frac{x^6}{360} + \frac{x^7}{2520} + O(x^8) \end{aligned} \quad (1)$$

$$\begin{aligned} y &= \left(1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{720}x^6 + \frac{1}{5040}x^7\right) c_1 \\ &\quad + \frac{x^3}{3} + \frac{x^4}{12} + \frac{x^5}{60} + \frac{x^6}{360} + \frac{x^7}{2520} + O(x^8) \end{aligned} \quad (2)$$




Figure 444: Slope field plot

Verification of solutions

$$y = \left(1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{720}x^6 + \frac{1}{5040}x^7 \right) y(0) + \frac{x^3}{3} + \frac{x^4}{12} + \frac{x^5}{60} + \frac{x^6}{360} + \frac{x^7}{2520} + O(x^8)$$

Verified OK.

$$y = \left(1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{720}x^6 + \frac{1}{5040}x^7 \right) c_1 + \frac{x^3}{3} + \frac{x^4}{12} + \frac{x^5}{60} + \frac{x^6}{360} + \frac{x^7}{2520} + O(x^8)$$

Verified OK.

17.11.2 Maple step by step solution

Let's solve

$$y' - y = x^2$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = x^2 + y$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' - y = x^2$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) (y' - y) = \mu(x) x^2$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x) y)$

$$\mu(x) (y' - y) = \mu'(x) y + \mu(x) y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = -\mu(x)$$

- Solve to find the integrating factor

$$\mu(x) = e^{-x}$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x) y) \right) dx = \int \mu(x) x^2 dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x) y = \int \mu(x) x^2 dx + c_1$$

- Solve for y

$$y = \frac{\int \mu(x) x^2 dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = e^{-x}$

$$y = \frac{\int x^2 e^{-x} dx + c_1}{e^{-x}}$$

- Evaluate the integrals on the rhs

$$y = \frac{-(x^2 + 2x + 2)e^{-x} + c_1}{e^{-x}}$$

- Simplify

$$y = c_1 e^x - x^2 - 2x - 2$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 54

```
Order:=8;  
dsolve(diff(y(x),x)-y(x)=x^2,y(x),type='series',x=0);
```

$$y(x) = \left(1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{720}x^6 + \frac{1}{5040}x^7\right) y(0) \\ + \frac{x^3}{3} + \frac{x^4}{12} + \frac{x^5}{60} + \frac{x^6}{360} + \frac{x^7}{2520} + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 87

```
AsymptoticDSolveValue[y'[x]-y[x]==x^2,y[x],{x,0,7}]
```

$$y(x) \rightarrow \frac{x^7}{2520} + \frac{x^6}{360} + \frac{x^5}{60} + \frac{x^4}{12} + \frac{x^3}{3} + c_1 \left(\frac{x^7}{5040} + \frac{x^6}{720} + \frac{x^5}{120} + \frac{x^4}{24} + \frac{x^3}{6} + \frac{x^2}{2} + x + 1 \right)$$

17.12 problem 1(f) solving directly

17.12.1 Solving as linear ode	2825
17.12.2 Solving as first order ode lie symmetry lookup ode	2827
17.12.3 Solving as exact ode	2831
17.12.4 Maple step by step solution	2835

Internal problem ID [6415]

Internal file name [OUTPUT/5663_Sunday_June_05_2022_03_46_21_PM_30766124/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.2. Series Solutions of First-Order Differential Equations Page 162

Problem number: 1(f) solving directly.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"linear", "exactWithIntegrationFactor", "first_order_ode_lie_symmetry_lookup"**

Maple gives the following as the ode type

```
[[_linear, `class A`]]
```

$$y' - y = x^2$$

17.12.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -1$$

$$q(x) = x^2$$

Hence the ode is

$$y' - y = x^2$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int (-1) dx} \\ &= e^{-x}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu)(x^2) \\ \frac{d}{dx}(e^{-x}y) &= (e^{-x})(x^2) \\ d(e^{-x}y) &= (x^2 e^{-x}) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}e^{-x}y &= \int x^2 e^{-x} dx \\ e^{-x}y &= -(x^2 + 2x + 2) e^{-x} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{-x}$ results in

$$y = -e^x(x^2 + 2x + 2) e^{-x} + c_1 e^x$$

which simplifies to

$$y = c_1 e^x - x^2 - 2x - 2$$

Summary

The solution(s) found are the following

$$y = c_1 e^x - x^2 - 2x - 2 \tag{1}$$




Figure 445: Slope field plot

Verification of solutions

$$y = c_1 e^x - x^2 - 2x - 2$$

Verified OK.

17.12.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = x^2 + y$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 405: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= e^x\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^x} dy \end{aligned}$$

Which results in

$$S = e^{-x}y$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = x^2 + y$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= -e^{-x}y \\ S_y &= e^{-x} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = x^2 e^{-x} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = R^2 e^{-R}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by

integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = -(R^2 + 2R + 2) e^{-R} + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$y e^{-x} = -(x^2 + 2x + 2) e^{-x} + c_1$$



Which simplifies to

$$(2 + 2x + x^2 + y) e^{-x} - c_1 = 0$$

Which gives

$$y = -(x^2 e^{-x} + 2x e^{-x} + 2 e^{-x} - c_1) e^x$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = x^2 + y$ 	$R = x$ $S = e^{-x} y$	$\frac{dS}{dR} = R^2 e^{-R}$ 

Summary

The solution(s) found are the following

$$y = -(x^2 e^{-x} + 2x e^{-x} + 2 e^{-x} - c_1) e^x \quad (1)$$




Figure 446: Slope field plot

Verification of solutions

$$y = -(x^2 e^{-x} + 2x e^{-x} + 2 e^{-x} - c_1) e^x$$

Verified OK.

17.12.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}dy &= (x^2 + y) dx \\ (-x^2 - y) dx + dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -x^2 - y \\ N(x, y) &= 1\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-x^2 - y) \\ &= -1\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(1) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= 1((-1) - (0)) \\ &= -1 \end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int A dx} \\ &= e^{\int -1 dx} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^{-x} \\ &= e^{-x} \end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} for now so not to confuse them with the original M and N .

$$\begin{aligned} \bar{M} &= \mu M \\ &= e^{-x}(-x^2 - y) \\ &= -e^{-x}(x^2 + y) \end{aligned}$$

And

$$\begin{aligned} \bar{N} &= \mu N \\ &= e^{-x}(1) \\ &= e^{-x} \end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned} \bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ (-e^{-x}(x^2 + y)) + (e^{-x}) \frac{dy}{dx} &= 0 \end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -e^{-x}(x^2 + y) dx \\ \phi &= (x^2 + 2x + y + 2) e^{-x} + f(y)\end{aligned}\quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = e^{-x} + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = e^{-x}$. Therefore equation (4) becomes

$$e^{-x} = e^{-x} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = (x^2 + 2x + y + 2) e^{-x} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = (x^2 + 2x + y + 2) e^{-x}$$

The solution becomes

$$y = -(x^2 e^{-x} + 2x e^{-x} + 2 e^{-x} - c_1) e^x$$

Summary

The solution(s) found are the following

$$y = -(x^2 e^{-x} + 2x e^{-x} + 2 e^{-x} - c_1) e^x \quad (1)$$




Figure 447: Slope field plot

Verification of solutions

$$y = -(x^2 e^{-x} + 2x e^{-x} + 2 e^{-x} - c_1) e^x$$

Verified OK.

17.12.4 Maple step by step solution

Let's solve

$$y' - y = x^2$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = x^2 + y$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' - y = x^2$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) (y' - y) = \mu(x) x^2$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x) y)$

$$\mu(x) (y' - y) = \mu'(x) y + \mu(x) y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = -\mu(x)$$

- Solve to find the integrating factor

$$\mu(x) = e^{-x}$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x) y) \right) dx = \int \mu(x) x^2 dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x) y = \int \mu(x) x^2 dx + c_1$$

- Solve for y

$$y = \frac{\int \mu(x) x^2 dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = e^{-x}$

$$y = \frac{\int x^2 e^{-x} dx + c_1}{e^{-x}}$$

- Evaluate the integrals on the rhs

$$y = \frac{-(x^2 + 2x + 2)e^{-x} + c_1}{e^{-x}}$$

- Simplify

$$y = c_1 e^x - x^2 - 2x - 2$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(diff(y(x),x)-y(x)=x^2,y(x), singsol=all)
```

$$y(x) = -x^2 - 2x - 2 + e^x c_1$$

✓ Solution by Mathematica

Time used: 0.046 (sec). Leaf size: 21

```
DSolve[y'[x]-y[x]==x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x^2 - 2x + c_1 e^x - 2$$

17.13 problem 2(a) solving using series

17.13.1 Solving as series ode	2838
17.13.2 Maple step by step solution	2842

Internal problem ID [6416]

Internal file name [OUTPUT/5664_Sunday_June_05_2022_03_46_22_PM_92286864/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.2. Series Solutions of First-Order Differential Equations Page 162

Problem number: 2(a) solving using series.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"first order ode series method. Regular singular point"**

Maple gives the following as the ode type

[_separable]

$$-y + xy' = 0$$

With the expansion point for the power series method at $x = 0$.

17.13.1 Solving as series ode

Writing the ODE as

$$y' + q(x)y = p(x)$$
$$y' - \frac{y}{x} = 0$$

Where

$$q(x) = -\frac{1}{x}$$
$$p(x) = 0$$

Next, the type of the expansion point $x = 0$ is determined. This point can be an ordinary point, a regular singular point (also called removable singularity), or irregular

singular point (also called non-removable singularity or essential singularity). When $x = 0$ is an ordinary point, then the standard power series is used. If the point is a regular singular point, Frobenius series is used instead. Irregular singular point requires more advanced methods (asymptotic methods) and is not supported now. Hopefully this will be added in the future. $x = 0$ is called an ordinary point if $q(x)$ has a Taylor series expansion around the point $x = 0$. $x = 0$ is called a regular singular point if $q(x)$ is not analytic at $x = 0$ but $xq(x)$ has Taylor series expansion. And finally, $x = 0$ is an irregular singular point if the point is not ordinary and not regular singular. This is the most complicated case. Now the expansion point $x = 0$ is checked to see if it is an ordinary point or not.

Since $x = 0$ is not an ordinary point, we now check to see if it is a regular singular point. $xq(x) = -1$ has a Taylor series around $x = 0$. Since $x = 0$ is regular singular point, then Frobenius power series is used. Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

Substituting the above back into the ode gives

$$\left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) - \frac{\sum_{n=0}^{\infty} a_n x^{n+r}}{x} = 0 \quad (1)$$

Hence the ODE in Eq (1) becomes

$$\left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) - \frac{\sum_{n=0}^{\infty} a_n x^{n+r}}{x} = 0 \quad (1)$$

Expanding the second term in (1) gives

$$\left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + -1 \cdot \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \frac{1}{x} \cdot \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \sum_{n=0}^{\infty} (-x^{n+r-1} a_n) = 0 \quad (2A)$$

The next step is to make all powers of x be $n + r - 1$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r-1} and adjusting the power and the corresponding index gives Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n + r - 1$.

$$\left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \sum_{n=0}^{\infty} (-x^{n+r-1} a_n) = 0 \quad (2B)$$

The indicial equation is obtained from $n = 0$. From Eq (2) this gives

$$(n+r) a_n x^{n+r-1} - x^{n+r-1} a_n = 0$$

When $n = 0$ the above becomes

$$r a_0 x^{-1+r} - x^{-1+r} a_0 = 0$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(-1+r) x^{-1+r} = 0$$

Since the above is true for all x then the indicial equation simplifies to

$$-1+r = 0$$

Solving for r gives the root of the indicial equation as

$$r = 1$$

We start by finding y_h . Replacing $r = 1$ found above results in

$$\left(\sum_{n=0}^{\infty} (n+1) a_n x^n \right) + \sum_{n=0}^{\infty} (-x^n a_n) = 0$$

From the above we see that there is no recurrence relation since there is only one summation term. Therefore all a_n terms are zero except for a_0 . Hence

$$y_h = a_0 x^r$$

Therefore the homogeneous solution is

$$y_h(x) = a_0(x + O(x^8))$$

At $x = 0$ the solution above becomes

$$y = c_1(x + O(x^8))$$

Summary

The solution(s) found are the following

$$y = c_1(x + O(x^8)) \tag{1}$$




Figure 448: Slope field plot

Verification of solutions

$$y = c_1(x + O(x^8))$$

Verified OK.

17.13.2 Maple step by step solution

Let's solve

$$y' - \frac{y}{x} = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y} = \frac{1}{x}$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y} dx = \int \frac{1}{x} dx + c_1$$

- Evaluate integral

$$\ln(y) = \ln(x) + c_1$$

- Solve for y

$$y = x e^{c_1}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 13

```
Order:=8;  
dsolve(x*diff(y(x),x)=y(x),y(x),type='series',x=0);
```

$$y(x) = c_1 x + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 7

```
AsymptoticDSolveValue[x*y'[x]==y[x],y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 x$$

17.14 problem 2(a) solving directly

17.14.1 Solving as separable ode	2844
17.14.2 Solving as linear ode	2846
17.14.3 Solving as homogeneousTypeD2 ode	2847
17.14.4 Solving as first order ode lie symmetry lookup ode	2848
17.14.5 Solving as exact ode	2852
17.14.6 Maple step by step solution	2856

Internal problem ID [6417]

Internal file name [OUTPUT/5665_Sunday_June_05_2022_03_46_25_PM_59981860/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.2. Series Solutions of First-Order Differential Equations Page 162

Problem number: 2(a) solving directly.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "linear", "separable", "homogeneousTypeD2", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

`[_separable]`

$$-y + xy' = 0$$

17.14.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{y}{x}\end{aligned}$$

Where $f(x) = \frac{1}{x}$ and $g(y) = y$. Integrating both sides gives

$$\begin{aligned}\frac{1}{y} dy &= \frac{1}{x} dx \\ \int \frac{1}{y} dy &= \int \frac{1}{x} dx \\ \ln(y) &= \ln(x) + c_1 \\ y &= e^{\ln(x)+c_1} \\ &= c_1 x\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 x \tag{1}$$




Figure 449: Slope field plot

Verification of solutions

$$y = c_1 x$$

Verified OK.

17.14.2 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -\frac{1}{x}$$
$$q(x) = 0$$

Hence the ode is

$$y' - \frac{y}{x} = 0$$

The integrating factor μ is

$$\mu = e^{\int -\frac{1}{x} dx}$$
$$= \frac{1}{x}$$

The ode becomes

$$\frac{d}{dx} \mu y = 0$$
$$\frac{d}{dx} \left(\frac{y}{x} \right) = 0$$

Integrating gives

$$\frac{y}{x} = c_1$$

Dividing both sides by the integrating factor $\mu = \frac{1}{x}$ results in

$$y = c_1 x$$

Summary

The solution(s) found are the following

$$y = c_1 x \tag{1}$$




Figure 450: Slope field plot

Verification of solutions

$$y = c_1 x$$

Verified OK.

17.14.3 Solving as homogeneous Type D2 ode

Using the change of variables $y = u(x)x$ on the above ode results in new ode in $u(x)$

$$-u(x)x + x(u'(x)x + u(x)) = 0$$

Integrating both sides gives

$$\begin{aligned} u(x) &= \int 0 \, dx \\ &= c_2 \end{aligned}$$

Therefore the solution y is

$$\begin{aligned} y &= ux \\ &= c_2 x \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_2x \tag{1}$$




Figure 451: Slope field plot

Verification of solutions

$$y = c_2x$$

Verified OK.

17.14.4 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{y}{x}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2\xi_y - \omega_x\xi - \omega_y\eta = 0 \tag{A}$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 409: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int(n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= x\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{x} dy \end{aligned}$$

Which results in

$$S = \frac{y}{x}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{y}{x}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= -\frac{y}{x^2} \\ S_y &= \frac{1}{x} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \tag{4}$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{y}{x} = c_1$$



Which simplifies to

$$\frac{y}{x} = c_1$$

Which gives

$$y = c_1 x$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
<div style="text-align: center;"> $\frac{dy}{dx} = \frac{y}{x}$ </div> 	$R = x$ $S = \frac{y}{x}$	<div style="text-align: center;"> $\frac{dS}{dR} = 0$ </div> 

Summary

The solution(s) found are the following

$$y = c_1 x \tag{1}$$




Figure 452: Slope field plot

Verification of solutions

$$y = c_1 x$$

Verified OK.

17.14.5 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}\left(\frac{1}{y}\right) dy &= \left(\frac{1}{x}\right) dx \\ \left(-\frac{1}{x}\right) dx + \left(\frac{1}{y}\right) dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -\frac{1}{x} \\ N(x, y) &= \frac{1}{y}\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(-\frac{1}{x}\right) \\ &= 0\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{1}{y} \right) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\frac{1}{x} dx \\ \phi &= -\ln(x) + f(y)\end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{1}{y}$. Therefore equation (4) becomes

$$\frac{1}{y} = 0 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = \frac{1}{y}$$

Integrating the above w.r.t y gives

$$\begin{aligned}\int f'(y) dy &= \int \left(\frac{1}{y} \right) dy \\ f(y) &= \ln(y) + c_1\end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\ln(x) + \ln(y) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\ln(x) + \ln(y)$$

The solution becomes

$$y = x e^{c_1}$$

Summary

The solution(s) found are the following

$$y = x e^{c_1} \tag{1}$$




Figure 453: Slope field plot

Verification of solutions

$$y = x e^{c_1}$$

Verified OK.

17.14.6 Maple step by step solution

Let's solve

$$-y + xy' = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y} = \frac{1}{x}$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y} dx = \int \frac{1}{x} dx + c_1$$

- Evaluate integral

$$\ln(y) = \ln(x) + c_1$$

- Solve for y

$$y = x e^{c_1}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 7

```
dsolve(x*diff(y(x),x)=y(x),y(x), singsol=all)
```

$$y(x) = c_1 x$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 14

```
DSolve[x*y'[x]==y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 x$$

$$y(x) \rightarrow 0$$

17.15 problem 2(b) solving using series

17.15.1 Solving as series ode	2858
17.15.2 Maple step by step solution	2859

Internal problem ID [6418]

Internal file name [OUTPUT/5666_Sunday_June_05_2022_03_46_26_PM_70853944/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.2. Series Solutions of First-Order Differential Equations Page 162

Problem number: 2(b) solving using series.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"first order ode series method. Irregular singular point"**

Maple gives the following as the ode type

`[_separable]`

Unable to solve or complete the solution.

$$x^2y' - y = 0$$

With the expansion point for the power series method at $x = 0$.

17.15.1 Solving as series ode

Writing the ODE as

$$y' + q(x)y = p(x)$$
$$y' - \frac{y}{x^2} = 0$$

Where

$$q(x) = -\frac{1}{x^2}$$
$$p(x) = 0$$

Next, the type of the expansion point $x = 0$ is determined. This point can be an ordinary point, a regular singular point (also called removable singularity), or irregular singular point (also called non-removable singularity or essential singularity). When $x = 0$ is an ordinary point, then the standard power series is used. If the point is a regular singular point, Frobenius series is used instead. Irregular singular point requires more advanced methods (asymptotic methods) and is not supported now. Hopefully this will be added in the future. $x = 0$ is called an ordinary point if $q(x)$ has a Taylor series expansion around the point $x = 0$. $x = 0$ is called a regular singular point if $q(x)$ is not analytic at $x = 0$ but $xq(x)$ has Taylor series expansion. And finally, $x = 0$ is an irregular singular point if the point is not ordinary and not regular singular. This is the most complicated case. Now the expansion point $x = 0$ is checked to see if it is an ordinary point or not.

Since $x = 0$ is not an ordinary point, we now check to see if it is a regular singular point.

$$xq(x) = -\frac{1}{x}$$

does not have a Taylor series around $x = 0$.

Unable to solve since $x = 0$ is not regular singular point. Terminating.

Verification of solutions N/A

17.15.2 Maple step by step solution

Let's solve

$$x^2y' - y = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y} = \frac{1}{x^2}$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y} dx = \int \frac{1}{x^2} dx + c_1$$

- Evaluate integral

$$\ln(y) = c_1 - \frac{1}{x}$$

- Solve for y

$$y = e^{\frac{c_1 x - 1}{x}}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

Solution by Maple

```
Order:=8;  
dsolve(x^2*diff(y(x),x)=y(x),y(x),type='series',x=0);
```

No solution found

Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 13

```
AsymptoticDSolveValue[x^2*y'[x]==y[x],y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 e^{-1/x}$$

17.16 problem 2(b) solving directly

17.16.1 Solving as separable ode	2861
17.16.2 Solving as linear ode	2863
17.16.3 Solving as homogeneousTypeD2 ode	2864
17.16.4 Solving as first order ode lie symmetry lookup ode	2866
17.16.5 Solving as exact ode	2870
17.16.6 Maple step by step solution	2874

Internal problem ID [6419]

Internal file name [OUTPUT/5667_Sunday_June_05_2022_03_46_27_PM_99396460/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.2. Series Solutions of First-Order Differential Equations Page 162

Problem number: 2(b) solving directly.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "linear", "separable", "homogeneousTypeD2", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_separable]

$$x^2y' - y = 0$$

17.16.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{y}{x^2}\end{aligned}$$

Where $f(x) = \frac{1}{x^2}$ and $g(y) = y$. Integrating both sides gives

$$\begin{aligned}\frac{1}{y} dy &= \frac{1}{x^2} dx \\ \int \frac{1}{y} dy &= \int \frac{1}{x^2} dx \\ \ln(y) &= c_1 - \frac{1}{x} \\ y &= e^{c_1 - \frac{1}{x}} \\ &= c_1 e^{-\frac{1}{x}}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-\frac{1}{x}} \tag{1}$$




Figure 454: Slope field plot

Verification of solutions

$$y = c_1 e^{-\frac{1}{x}}$$

Verified OK.

17.16.2 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -\frac{1}{x^2}$$
$$q(x) = 0$$

Hence the ode is

$$y' - \frac{y}{x^2} = 0$$

The integrating factor μ is

$$\mu = e^{\int -\frac{1}{x^2} dx}$$
$$= e^{\frac{1}{x}}$$

The ode becomes

$$\frac{d}{dx} \mu y = 0$$
$$\frac{d}{dx} \left(e^{\frac{1}{x}} y \right) = 0$$

Integrating gives

$$e^{\frac{1}{x}} y = c_1$$

Dividing both sides by the integrating factor $\mu = e^{\frac{1}{x}}$ results in

$$y = c_1 e^{-\frac{1}{x}}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-\frac{1}{x}} \tag{1}$$




Figure 455: Slope field plot

Verification of solutions

$$y = c_1 e^{-\frac{1}{x}}$$

Verified OK.

17.16.3 Solving as homogeneousTypeD2 ode

Using the change of variables $y = u(x)x$ on the above ode results in new ode in $u(x)$

$$x^2(u'(x)x + u(x)) - u(x)x = 0$$

In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{u(x-1)}{x^2} \end{aligned}$$

Where $f(x) = -\frac{x-1}{x^2}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u} du &= -\frac{x-1}{x^2} dx \\ \int \frac{1}{u} du &= \int -\frac{x-1}{x^2} dx \\ \ln(u) &= -\ln(x) - \frac{1}{x} + c_2 \\ u &= e^{-\ln(x) - \frac{1}{x} + c_2} \\ &= c_2 e^{-\ln(x) - \frac{1}{x}}\end{aligned}$$

Which simplifies to

$$u(x) = \frac{c_2 e^{-\frac{1}{x}}}{x}$$

Therefore the solution y is

$$\begin{aligned}y &= ux \\ &= c_2 e^{-\frac{1}{x}}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_2 e^{-\frac{1}{x}} \tag{1}$$




Figure 456: Slope field plot

Verification of solutions

$$y = c_2 e^{-\frac{1}{x}}$$

Verified OK.

17.16.4 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{y}{x^2}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 413: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= e^{-\frac{1}{x}}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right)S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^{-\frac{1}{x}}} dy \end{aligned}$$

Which results in

$$S = e^{\frac{1}{x}} y$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{y}{x^2}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= -\frac{e^{\frac{1}{x}} y}{x^2} \\ S_y &= e^{\frac{1}{x}} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \tag{4}$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$e^{\frac{1}{x}} y = c_1$$



Which simplifies to

$$e^{\frac{1}{x}} y = c_1$$

Which gives

$$y = c_1 e^{-\frac{1}{x}}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{y}{x^2}$ 	$R = x$ $S = e^{\frac{1}{x}} y$	$\frac{dS}{dR} = 0$ 

Summary

The solution(s) found are the following

$$y = c_1 e^{-\frac{1}{x}} \tag{1}$$




Figure 457: Slope field plot

Verification of solutions

$$y = c_1 e^{-\frac{1}{x}}$$

Verified OK.

17.16.5 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}\left(\frac{1}{y}\right) dy &= \left(\frac{1}{x^2}\right) dx \\ \left(-\frac{1}{x^2}\right) dx + \left(\frac{1}{y}\right) dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -\frac{1}{x^2} \\ N(x, y) &= \frac{1}{y}\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(-\frac{1}{x^2}\right) \\ &= 0\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{1}{y} \right) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \tag{1}$$

$$\frac{\partial \phi}{\partial y} = N \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\frac{1}{x^2} dx \\ \phi &= \frac{1}{x} + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{1}{y}$. Therefore equation (4) becomes

$$\frac{1}{y} = 0 + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = \frac{1}{y}$$

Integrating the above w.r.t y gives

$$\begin{aligned}\int f'(y) dy &= \int \left(\frac{1}{y} \right) dy \\ f(y) &= \ln(y) + c_1\end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = \frac{1}{x} + \ln(y) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \frac{1}{x} + \ln(y)$$

The solution becomes

$$y = e^{\frac{c_1 x - 1}{x}}$$

Summary

The solution(s) found are the following

$$y = e^{\frac{c_1 x - 1}{x}} \tag{1}$$




Figure 458: Slope field plot

Verification of solutions

$$y = e^{\frac{c_1 x - 1}{x}}$$

Verified OK.

17.16.6 Maple step by step solution

Let's solve

$$x^2 y' - y = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y} = \frac{1}{x^2}$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y} dx = \int \frac{1}{x^2} dx + c_1$$

- Evaluate integral

$$\ln(y) = c_1 - \frac{1}{x}$$

- Solve for y

$$y = e^{\frac{c_1 x - 1}{x}}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(x^2*diff(y(x),x)=y(x),y(x), singsol=all)
```

$$y(x) = c_1 e^{-\frac{1}{x}}$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 20

```
DSolve[x^2*y'[x]==y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 e^{-1/x}$$

$$y(x) \rightarrow 0$$

17.17 problem 2(c) solving using series

17.17.1 Solving as series ode	2876
17.17.2 Maple step by step solution	2881

Internal problem ID [6420]

Internal file name [OUTPUT/5668_Sunday_June_05_2022_03_46_29_PM_78088919/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.2. Series Solutions of First-Order Differential Equations Page 162

Problem number: 2(c) solving using series.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"first order ode series method. Regular singular point"**

Maple gives the following as the ode type

[_linear]

$$y' - \frac{y}{x} = x^2$$

With the expansion point for the power series method at $x = 0$.

17.17.1 Solving as series ode

Writing the ODE as

$$y' + q(x)y = p(x)$$
$$y' - \frac{y}{x} = x^2$$

Where

$$q(x) = -\frac{1}{x}$$
$$p(x) = x^2$$

Next, the type of the expansion point $x = 0$ is determined. This point can be an ordinary point, a regular singular point (also called removable singularity), or irregular

singular point (also called non-removable singularity or essential singularity). When $x = 0$ is an ordinary point, then the standard power series is used. If the point is a regular singular point, Frobenius series is used instead. Irregular singular point requires more advanced methods (asymptotic methods) and is not supported now. Hopefully this will be added in the future. $x = 0$ is called an ordinary point if $q(x)$ has a Taylor series expansion around the point $x = 0$. $x = 0$ is called a regular singular point if $q(x)$ is not analytic at $x = 0$ but $xq(x)$ has Taylor series expansion. And finally, $x = 0$ is an irregular singular point if the point is not ordinary and not regular singular. This is the most complicated case. Now the expansion point $x = 0$ is checked to see if it is an ordinary point or not.

Since $x = 0$ is not an ordinary point, we now check to see if it is a regular singular point. $xq(x) = -1$ has a Taylor series around $x = 0$. Since $x = 0$ is regular singular point, then Frobenius power series is used. Since this is an inhomogeneous, then let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ode $y' - \frac{y}{x} = 0$, and y_p is a particular solution to the inhomogeneous ode. First, we solve for y_h . Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

Substituting the above back into the ode gives

$$\left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) - \frac{\sum_{n=0}^{\infty} a_n x^{n+r}}{x} = 0 \quad (1)$$

Hence the ODE in Eq (1) becomes

$$\left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) - \frac{\sum_{n=0}^{\infty} a_n x^{n+r}}{x} = 0 \quad (1)$$

Expanding the second term in (1) gives

$$\left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + -1 \cdot \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \frac{1}{x} \cdot \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \sum_{n=0}^{\infty} (-x^{n+r-1} a_n) = 0 \quad (2A)$$

The next step is to make all powers of x be $n+r-1$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r-1} and adjusting the power and the corresponding index gives Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r-1$.

$$\left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \sum_{n=0}^{\infty} (-x^{n+r-1} a_n) = 0 \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2) this gives

$$(n+r) a_n x^{n+r-1} - x^{n+r-1} a_n = 0$$

When $n=0$ the above becomes

$$r a_0 x^{-1+r} - x^{-1+r} a_0 = 0$$

The corresponding balance equation is found by replacing r by m and a by c to avoid confusing terms between particular solution and the homogeneous solution. Hence the balance equation is

$$(x^{-1+m} m - x^{-1+m}) c_0 = x^2$$

This equation will used later to find the particular solution.

Since $a_0 \neq 0$ then the indicial equation becomes

$$(-1+r) x^{-1+r} = 0$$

Since the above is true for all x then the indicial equation simplifies to

$$-1+r = 0$$

Solving for r gives the root of the indicial equation as

$$r = 1$$

We start by finding y_h . From the above we see that there is no recurrence relation since there is only one summation term. Therefore all a_n terms are zero except for a_0 . Hence

$$y_h = a_0 x^r$$

Therefore the homogeneous solution is

$$y_h(x) = a_0(x + O(x^8))$$

Now we determine the particular solution y_p by solving the balance equation

$$(x^{-1+m}m - x^{-1+m}) c_0 = x^2$$

For c_0 and x . This results in

$$\begin{aligned} c_0 &= \frac{1}{2} \\ m &= 3 \end{aligned}$$

The particular solution is therefore

$$\begin{aligned} y_p &= \sum_{n=0}^{\infty} c_n x^{n+m} \\ &= \sum_{n=0}^{\infty} c_n x^{n+3} \end{aligned}$$

Where in the above $c_0 = \frac{1}{2}$. The remaining c_n values are found using the same recurrence relation used to find the homogeneous solution but using c_0 in place of a_0 and using $m = 3$ in place of the root of the indicial equation used to find the homogeneous solution. The following are the values of a_n found in terms of the indicial root r . These will be now used to find c_n by replacing $a_0 = \frac{1}{2}$ and $r = 3$. The following table gives the a_n values found and the corresponding c_n values which will be used to find the particular solution

n	a_n	c_n
0	$a_0 = 1$	$c_0 = \frac{1}{2}$

The particular solution is now found using

$$\begin{aligned} y_p &= x^m \sum_{n=0}^{\infty} c_n x^n \\ &= x^3 \sum_{n=0}^{\infty} c_n x^n \end{aligned}$$

Using the values found above for c_n into the above sum gives

$$y_p = x^3 \left(\frac{1}{2} \right)$$

At $x = 0$ the solution above becomes

$$y = \frac{x^3}{2} + O(x^8) + c_1(x + O(x^8))$$

Summary

The solution(s) found are the following

$$y = \frac{x^3}{2} + O(x^8) + c_1(x + O(x^8)) \quad (1)$$




Figure 459: Slope field plot

Verification of solutions

$$y = \frac{x^3}{2} + O(x^8) + c_1(x + O(x^8))$$

Verified OK.

17.17.2 Maple step by step solution

Let's solve

$$y' - \frac{y}{x} = x^2$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = \frac{y}{x} + x^2$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' - \frac{y}{x} = x^2$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) \left(y' - \frac{y}{x} \right) = \mu(x) x^2$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x) y)$

$$\mu(x) \left(y' - \frac{y}{x} \right) = \mu'(x) y + \mu(x) y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = -\frac{\mu(x)}{x}$$

- Solve to find the integrating factor

$$\mu(x) = \frac{1}{x}$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x) y) \right) dx = \int \mu(x) x^2 dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x) y = \int \mu(x) x^2 dx + c_1$$

- Solve for y

$$y = \frac{\int \mu(x) x^2 dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = \frac{1}{x}$

$$y = x \left(\int x dx + c_1 \right)$$

- Evaluate the integrals on the rhs

$$y = x \left(\frac{x^2}{2} + c_1 \right)$$

- Simplify

$$y = \frac{x(x^2 + 2c_1)}{2}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 24

```
Order:=8;  
dsolve(diff(y(x),x)-(1/x)*y(x)=x^2,y(x),type='series',x=0);
```

$$y(x) = c_1x(1 + O(x^8)) + x^3\left(\frac{1}{2} + O(x^5)\right)$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 15

```
AsymptoticDSolveValue[y'[x]-1/x*y[x]==x^2,y[x],{x,0,7}]
```

$$y(x) \rightarrow \frac{x^3}{2} + c_1x$$

17.18 problem 2(c) solving directly

17.18.1 Solving as linear ode	2883
17.18.2 Solving as homogeneousTypeD2 ode	2885
17.18.3 Solving as first order ode lie symmetry lookup ode	2886
17.18.4 Solving as exact ode	2890
17.18.5 Maple step by step solution	2895

Internal problem ID [6421]

Internal file name [OUTPUT/5669_Sunday_June_05_2022_03_46_30_PM_49881333/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.2. Series Solutions of First-Order Differential Equations Page 162

Problem number: 2(c) solving directly.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"linear"**, **"homogeneousTypeD2"**, **"exactWithIntegrationFactor"**, **"first_order_ode_lie_symmetry_lookup"**

Maple gives the following as the ode type

[_linear]

$$y' - \frac{y}{x} = x^2$$

17.18.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = -\frac{1}{x}$$
$$q(x) = x^2$$

Hence the ode is

$$y' - \frac{y}{x} = x^2$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int -\frac{1}{x} dx} \\ &= \frac{1}{x}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu)(x^2) \\ \frac{d}{dx}\left(\frac{y}{x}\right) &= \left(\frac{1}{x}\right)(x^2) \\ d\left(\frac{y}{x}\right) &= x dx\end{aligned}$$

Integrating gives

$$\begin{aligned}\frac{y}{x} &= \int x dx \\ \frac{y}{x} &= \frac{x^2}{2} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = \frac{1}{x}$ results in

$$y = \frac{1}{2}x^3 + c_1x$$

Summary

The solution(s) found are the following

$$y = \frac{1}{2}x^3 + c_1x \tag{1}$$




Figure 460: Slope field plot

Verification of solutions

$$y = \frac{1}{2}x^3 + c_1x$$

Verified OK.

17.18.2 Solving as homogeneousTypeD2 ode

Using the change of variables $y = u(x)x$ on the above ode results in new ode in $u(x)$

$$u'(x)x = x^2$$

Integrating both sides gives

$$\begin{aligned} u(x) &= \int x \, dx \\ &= \frac{x^2}{2} + c_2 \end{aligned}$$

Therefore the solution y is

$$\begin{aligned} y &= xu \\ &= x\left(\frac{x^2}{2} + c_2\right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = x \left(\frac{x^2}{2} + c_2 \right) \quad (1)$$




Figure 461: Slope field plot

Verification of solutions

$$y = x \left(\frac{x^2}{2} + c_2 \right)$$

Verified OK.

17.18.3 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{x^3 + y}{x}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 417: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= x\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{x} dy \end{aligned}$$

Which results in

$$S = \frac{y}{x}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{x^3 + y}{x}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= -\frac{y}{x^2} \\ S_y &= \frac{1}{x} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = x \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = R$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \frac{R^2}{2} + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{y}{x} = \frac{x^2}{2} + c_1$$



Which simplifies to

$$\frac{y}{x} = \frac{x^2}{2} + c_1$$

Which gives

$$y = \frac{x(x^2 + 2c_1)}{2}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{x^3 + y}{x}$ 	$R = x$ $S = \frac{y}{x}$	$\frac{dS}{dR} = R$ 

Summary

The solution(s) found are the following

$$y = \frac{x(x^2 + 2c_1)}{2} \quad (1)$$




Figure 462: Slope field plot

Verification of solutions

$$y = \frac{x(x^2 + 2c_1)}{2}$$

Verified OK.

17.18.4 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (A)$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} dy &= \left(\frac{y}{x} + x^2 \right) dx \\ \left(-\frac{y}{x} - x^2 \right) dx + dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -\frac{y}{x} - x^2 \\ N(x, y) &= 1 \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(-\frac{y}{x} - x^2 \right) \\ &= -\frac{1}{x} \end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(1) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned}A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= 1 \left(\left(-\frac{1}{x} \right) - (0) \right) \\ &= -\frac{1}{x}\end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int A dx} \\ &= e^{\int -\frac{1}{x} dx}\end{aligned}$$

The result of integrating gives

$$\begin{aligned}\mu &= e^{-\ln(x)} \\ &= \frac{1}{x}\end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} for now so not to confuse them with the original M and N .

$$\begin{aligned}\bar{M} &= \mu M \\ &= \frac{1}{x} \left(-\frac{y}{x} - x^2 \right) \\ &= \frac{-x^3 - y}{x^2}\end{aligned}$$

And

$$\begin{aligned}\bar{N} &= \mu N \\ &= \frac{1}{x}(1) \\ &= \frac{1}{x}\end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned} \overline{M} + \overline{N} \frac{dy}{dx} &= 0 \\ \left(\frac{-x^3 - y}{x^2} \right) + \left(\frac{1}{x} \right) \frac{dy}{dx} &= 0 \end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \overline{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \overline{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned} \int \frac{\partial \phi}{\partial x} dx &= \int \overline{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int \frac{-x^3 - y}{x^2} dx \\ \phi &= \frac{-x^3 + 2y}{2x} + f(y) \end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = \frac{1}{x} + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{1}{x}$. Therefore equation (4) becomes

$$\frac{1}{x} = \frac{1}{x} + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = \frac{-x^3 + 2y}{2x} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \frac{-x^3 + 2y}{2x}$$

The solution becomes

$$y = \frac{x(x^2 + 2c_1)}{2}$$

Summary

The solution(s) found are the following

$$y = \frac{x(x^2 + 2c_1)}{2} \tag{1}$$




Figure 463: Slope field plot

Verification of solutions

$$y = \frac{x(x^2 + 2c_1)}{2}$$

Verified OK.

17.18.5 Maple step by step solution

Let's solve

$$y' - \frac{y}{x} = x^2$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = \frac{y}{x} + x^2$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' - \frac{y}{x} = x^2$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) \left(y' - \frac{y}{x} \right) = \mu(x) x^2$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x) y)$

$$\mu(x) \left(y' - \frac{y}{x} \right) = \mu'(x) y + \mu(x) y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = -\frac{\mu(x)}{x}$$

- Solve to find the integrating factor

$$\mu(x) = \frac{1}{x}$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x) y) \right) dx = \int \mu(x) x^2 dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x) y = \int \mu(x) x^2 dx + c_1$$

- Solve for y

$$y = \frac{\int \mu(x) x^2 dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = \frac{1}{x}$

$$y = x \left(\int x dx + c_1 \right)$$

- Evaluate the integrals on the rhs

$$y = x \left(\frac{x^2}{2} + c_1 \right)$$

- Simplify

$$y = \frac{x(x^2 + 2c_1)}{2}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 14

```
dsolve(diff(y(x),x)-(1/x)*y(x)=x^2,y(x), singsol=all)
```

$$y(x) = \frac{x(x^2 + 2c_1)}{2}$$

✓ Solution by Mathematica

Time used: 0.025 (sec). Leaf size: 17

```
DSolve[y'[x]-1/x*y[x]==x^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^3}{2} + c_1 x$$

17.19 problem 2(d) solving using series

17.19.1 Solving as linear ode	2897
17.19.2 Solving as differentialType ode	2899
17.19.3 Solving as first order ode lie symmetry lookup ode	2901
17.19.4 Solving as exact ode	2905
17.19.5 Maple step by step solution	2909

Internal problem ID [6422]

Internal file name [OUTPUT/5670_Sunday_June_05_2022_03_46_32_PM_70367049/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.2. Series Solutions of First-Order Differential Equations Page 162

Problem number: 2(d) solving using series.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exact", "linear", "differentialType", "first_order_ode_lie_symmetry_lookup"**

Maple gives the following as the ode type

`[_linear]`

$$y' + \frac{y}{x} = x$$

17.19.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = \frac{1}{x}$$

$$q(x) = x$$

Hence the ode is

$$y' + \frac{y}{x} = x$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int \frac{1}{x} dx} \\ &= x\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu)(x) \\ \frac{d}{dx}(xy) &= (x)(x) \\ d(xy) &= x^2 dx\end{aligned}$$

Integrating gives

$$\begin{aligned}xy &= \int x^2 dx \\ xy &= \frac{x^3}{3} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = x$ results in

$$y = \frac{x^2}{3} + \frac{c_1}{x}$$

Summary

The solution(s) found are the following

$$y = \frac{x^2}{3} + \frac{c_1}{x} \tag{1}$$




Figure 464: Slope field plot

Verification of solutions

$$y = \frac{x^2}{3} + \frac{c_1}{x}$$

Verified OK.

17.19.2 Solving as differentialType ode

Writing the ode as

$$y' = -\frac{y}{x} + x \tag{1}$$

Which becomes

$$0 = (-x) dy + (x^2 - y) dx \tag{2}$$

But the RHS is complete differential because

$$(-x) dy + (x^2 - y) dx = d\left(\frac{1}{3}x^3 - xy\right)$$

Hence (2) becomes

$$0 = d\left(\frac{1}{3}x^3 - xy\right)$$

Integrating both sides gives gives these solutions

$$y = \frac{x^3 + 3c_1}{3x} + c_1$$

Summary

The solution(s) found are the following

$$y = \frac{x^3 + 3c_1}{3x} + c_1 \tag{1}$$




Figure 465: Slope field plot

Verification of solutions

$$y = \frac{x^3 + 3c_1}{3x} + c_1$$

Verified OK.

17.19.3 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = -\frac{-x^2 + y}{x}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 420: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= \frac{1}{x}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{1}{x}} dy\end{aligned}$$

Which results in

$$S = xy$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y}\tag{2}$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -\frac{-x^2 + y}{x}$$

Evaluating all the partial derivatives gives

$$R_x = 1$$

$$R_y = 0$$

$$S_x = y$$

$$S_y = x$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = x^2 \tag{2A}$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = R^2$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \frac{R^3}{3} + c_1 \tag{4}$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$xy = \frac{x^3}{3} + c_1$$



Which simplifies to

$$xy = \frac{x^3}{3} + c_1$$

Which gives

$$y = \frac{x^3 + 3c_1}{3x}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -\frac{-x^2+y}{x}$ 	$R = x$ $S = xy$	$\frac{dS}{dR} = R^2$ 

Summary

The solution(s) found are the following

$$y = \frac{x^3 + 3c_1}{3x} \tag{1}$$




Figure 466: Slope field plot

Verification of solutions

$$y = \frac{x^3 + 3c_1}{3x}$$

Verified OK.

17.19.4 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}(x) dy &= (x^2 - y) dx \\ (-x^2 + y) dx + (x) dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -x^2 + y \\ N(x, y) &= x\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-x^2 + y) \\ &= 1\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(x) \\ &= 1\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\int \frac{\partial \phi}{\partial x} dx = \int M dx$$

$$\int \frac{\partial \phi}{\partial x} dx = \int -x^2 + y dx$$

$$\phi = -\frac{1}{3}x^3 + xy + f(y) \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = x + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = x$. Therefore equation (4) becomes

$$x = x + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = -\frac{1}{3}x^3 + xy + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\frac{1}{3}x^3 + xy$$

The solution becomes

$$y = \frac{x^3 + 3c_1}{3x}$$

Summary

The solution(s) found are the following

$$y = \frac{x^3 + 3c_1}{3x} \tag{1}$$




Figure 467: Slope field plot

Verification of solutions

$$y = \frac{x^3 + 3c_1}{3x}$$

Verified OK.

17.19.5 Maple step by step solution

Let's solve

$$y' + \frac{y}{x} = x$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = -\frac{y}{x} + x$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' + \frac{y}{x} = x$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) \left(y' + \frac{y}{x} \right) = \mu(x) x$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x) y)$

$$\mu(x) \left(y' + \frac{y}{x} \right) = \mu'(x) y + \mu(x) y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = \frac{\mu(x)}{x}$$

- Solve to find the integrating factor

$$\mu(x) = x$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x) y) \right) dx = \int \mu(x) x dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x) y = \int \mu(x) x dx + c_1$$

- Solve for y

$$y = \frac{\int \mu(x) x dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = x$

$$y = \frac{\int x^2 dx + c_1}{x}$$

- Evaluate the integrals on the rhs

$$y = \frac{\frac{x^3}{3} + c_1}{x}$$

- Simplify

$$y = \frac{x^3 + 3c_1}{3x}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve(diff(y(x),x)+(1/x)*y(x)=x,y(x), singsol=all)
```

$$y(x) = \frac{x^3 + 3c_1}{3x}$$

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 19

```
DSolve[y'[x]+1/x*y[x]==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^2}{3} + \frac{c_1}{x}$$

17.20 problem 3

17.20.1 Solving as series ode	2911
17.20.2 Maple step by step solution	2921

Internal problem ID [6423]

Internal file name [OUTPUT/5671_Sunday_June_05_2022_03_46_33_PM_79704206/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.2. Series Solutions of First-Order Differential Equations Page 162

Problem number: 3.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"first order ode series method. Ordinary point", "first order ode series method. Taylor series method"**

Maple gives the following as the ode type

[_quadrature]

$$y' = \frac{1}{\sqrt{-x^2 + 1}}$$

With the expansion point for the power series method at $x = 0$.

17.20.1 Solving as series ode

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving first order ode. Let

$$y' = f(x, y)$$

Where $f(x, y)$ is analytic at expansion point x_0 . We can always shift to $x_0 = 0$ if x_0 is not zero. So from now we assume $x_0 = 0$. Assume also that $y(x_0) = y_0$. Using Taylor

series

$$\begin{aligned}
 y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \dots \\
 &= y_0 + xf + \frac{x^2}{2} \left. \frac{df}{dx} \right|_{x_0, y_0} + \frac{x^3}{3!} \left. \frac{d^2f}{dx^2} \right|_{x_0, y_0} + \dots \\
 &= y_0 + \sum_{n=0}^{\infty} \frac{x^{n+1}}{(n+1)!} \left. \frac{d^n f}{dx^n} \right|_{x_0, y_0}
 \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} f \tag{1}$$

$$\begin{aligned}
 \frac{d^2f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) f
 \end{aligned} \tag{2}$$

$$\begin{aligned}
 \frac{d^3f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2f}{dx^2} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{d^2f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2f}{dx^2} \right) f
 \end{aligned} \tag{3}$$

⋮

And so on. Hence if we name $F_0 = f(x, y)$ then the above can be written as

$$F_0 = f(x, y) \tag{4}$$

$$\begin{aligned}
 F_n &= \frac{d}{dx}(F_{n-1}) \\
 &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) F_0
 \end{aligned} \tag{5}$$

For example, for $n = 1$ we see that

$$\begin{aligned}
 F_1 &= \frac{d}{dx}(F_0) \\
 &= \frac{\partial}{\partial x} F_0 + \left(\frac{\partial F_0}{\partial y} \right) F_0 \\
 &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} f
 \end{aligned}$$

Which is (1). And when $n = 2$

$$\begin{aligned} F_2 &= \frac{d}{dx}(F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) F_0 \\ &= \frac{\partial}{\partial x} \left(\frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} f \right) + \frac{\partial}{\partial y} \left(\frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} f \right) f \\ &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) f \end{aligned}$$

Which is (2) and so on. Therefore (4,5) can be used from now on along with

$$y(x) = y_0 + \sum_{n=0}^{\infty} \frac{x^{n+1}}{(n+1)!} F_n|_{x_0, y_0} \quad (6)$$

Hence

$$\begin{aligned}F_0 &= \frac{1}{\sqrt{-x^2 + 1}} \\F_1 &= \frac{dF_0}{dx} \\&= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} F_0 \\&= \frac{x}{(-x^2 + 1)^{\frac{3}{2}}} \\F_2 &= \frac{dF_1}{dx} \\&= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} F_1 \\&= \frac{2x^2 + 1}{(-x^2 + 1)^{\frac{5}{2}}} \\F_3 &= \frac{dF_2}{dx} \\&= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} F_2 \\&= \frac{6x^3 + 9x}{(-x^2 + 1)^{\frac{7}{2}}} \\F_4 &= \frac{dF_3}{dx} \\&= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} F_3 \\&= \frac{24x^4 + 72x^2 + 9}{(-x^2 + 1)^{\frac{9}{2}}} \\F_5 &= \frac{dF_4}{dx} \\&= \frac{\partial F_4}{\partial x} + \frac{\partial F_4}{\partial y} F_4 \\&= \frac{120x(x^4 + 5x^2 + \frac{15}{8})}{(-x^2 + 1)^{\frac{11}{2}}} \\F_6 &= \frac{dF_5}{dx} \\&= \frac{\partial F_5}{\partial x} + \frac{\partial F_5}{\partial y} F_5 \\&= \frac{720x^6 + 5400x^4 + 4050x^2 + 225}{(-x^2 + 1)^{\frac{13}{2}}}\end{aligned}$$

And so on. Evaluating all the above at initial conditions $x(0) = 0$ and $y(0) = y(0)$ gives

$$\begin{aligned} F_0 &= 1 \\ F_1 &= 0 \\ F_2 &= 1 \\ F_3 &= 0 \\ F_4 &= 9 \\ F_5 &= 0 \\ F_6 &= 225 \end{aligned}$$

Substituting all the above in (6) and simplifying gives the solution as

$$y = y(0) + x + \frac{x^3}{6} + \frac{3x^5}{40} + \frac{5x^7}{112} + O(x^8)$$

Since $x = 0$ is also an ordinary point, then standard power series can also be used. Writing the ODE as

$$\begin{aligned} y' + q(x)y &= p(x) \\ y' &= \frac{1}{\sqrt{-x^2 + 1}} \end{aligned}$$

Where

$$\begin{aligned} q(x) &= 0 \\ p(x) &= \frac{1}{\sqrt{-x^2 + 1}} \end{aligned}$$

Next, the type of the expansion point $x = 0$ is determined. This point can be an ordinary point, a regular singular point (also called removable singularity), or irregular singular point (also called non-removable singularity or essential singularity). When $x = 0$ is an ordinary point, then the standard power series is used. If the point is a regular singular point, Frobenius series is used instead. Irregular singular point requires more advanced methods (asymptotic methods) and is not supported now. Hopefully this will be added in the future. $x = 0$ is called an ordinary point $q(x)$ has a Taylor series expansion around the point $x = 0$. $x = 0$ is called a regular singular point if $q(x)$ is not analytic at $x = 0$ but $xq(x)$ has Taylor series expansion. And finally, $x = 0$ is an irregular singular point if the point is not ordinary and not regular singular. This is the most complicated case. Now the expansion point $x = 0$ is checked to see if it is an ordinary point or not. Now the ode is normalized by writing it as

$$y' \sqrt{-x^2 + 1} = 1$$

Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$y' = \sum_{n=1}^{\infty} n a_n x^{n-1}$$

Substituting the above back into the ode gives

$$\left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) \sqrt{-x^2 + 1} = 1 \quad (1)$$

Expanding $\sqrt{-x^2 + 1}$ as Taylor series around $x = 0$ and keeping only the first 8 terms gives

$$\begin{aligned} \sqrt{-x^2 + 1} &= 1 - \frac{1}{2}x^2 - \frac{1}{8}x^4 - \frac{1}{16}x^6 - \frac{5}{128}x^8 + \dots \\ &= 1 - \frac{1}{2}x^2 - \frac{1}{8}x^4 - \frac{1}{16}x^6 - \frac{5}{128}x^8 \end{aligned}$$

Hence the ODE in Eq (1) becomes

$$\left(1 - \frac{1}{2}x^2 - \frac{1}{8}x^4 - \frac{1}{16}x^6 - \frac{5}{128}x^8 \right) \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) = 1 \quad (1)$$

Expanding the first term in (1) gives

$$1 \cdot \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) - \frac{x^2}{2} \cdot \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) - \frac{x^4}{8} \cdot \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) - \frac{x^6}{16} \cdot \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) - \frac{5x^8}{128} \cdot \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) \quad (1)$$

Which simplifies to

$$\begin{aligned} &\left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) + \sum_{n=1}^{\infty} \left(-\frac{n x^{1+n} a_n}{2} \right) + \sum_{n=1}^{\infty} \left(-\frac{n x^{n+3} a_n}{8} \right) \\ &+ \sum_{n=1}^{\infty} \left(-\frac{n x^{n+5} a_n}{16} \right) + \sum_{n=1}^{\infty} \left(-\frac{5n x^{n+7} a_n}{128} \right) = 1 \end{aligned} \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the

power and the corresponding index gives

$$\begin{aligned} \sum_{n=1}^{\infty} n a_n x^{n-1} &= \sum_{n=0}^{\infty} (1+n) a_{1+n} x^n \\ \sum_{n=1}^{\infty} \left(-\frac{n x^{1+n} a_n}{2} \right) &= \sum_{n=2}^{\infty} \left(-\frac{(n-1) a_{n-1} x^n}{2} \right) \\ \sum_{n=1}^{\infty} \left(-\frac{n x^{n+3} a_n}{8} \right) &= \sum_{n=4}^{\infty} \left(-\frac{(n-3) a_{n-3} x^n}{8} \right) \\ \sum_{n=1}^{\infty} \left(-\frac{n x^{n+5} a_n}{16} \right) &= \sum_{n=6}^{\infty} \left(-\frac{(n-5) a_{n-5} x^n}{16} \right) \\ \sum_{n=1}^{\infty} \left(-\frac{5n x^{n+7} a_n}{128} \right) &= \sum_{n=8}^{\infty} \left(-\frac{5(n-7) a_{n-7} x^n}{128} \right) \end{aligned}$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\begin{aligned} &\left(\sum_{n=0}^{\infty} (1+n) a_{1+n} x^n \right) + \sum_{n=2}^{\infty} \left(-\frac{(n-1) a_{n-1} x^n}{2} \right) + \sum_{n=4}^{\infty} \left(-\frac{(n-3) a_{n-3} x^n}{8} \right) \\ &+ \sum_{n=6}^{\infty} \left(-\frac{(n-5) a_{n-5} x^n}{16} \right) + \sum_{n=8}^{\infty} \left(-\frac{5(n-7) a_{n-7} x^n}{128} \right) = 1 \end{aligned} \quad (3)$$

$n = 0$ gives

$$\begin{aligned} (a_1) 1 &= 1 \\ a_1 &= 1 \end{aligned}$$

Or

$$a_1 = 1$$

$n = 2$ gives

$$3a_3 - \frac{a_1}{2} = 0$$

Which after substituting earlier equations, simplifies to

$$3a_3 - \frac{1}{2} = 0$$

Or

$$a_3 = \frac{1}{6}$$

$n = 3$ gives

$$4a_4 - a_2 = 0$$

Which after substituting earlier equations, simplifies to

$$4a_4 = 0$$

Or

$$a_4 = 0$$

$n = 4$ gives

$$5a_5 - \frac{3a_3}{2} - \frac{a_1}{8} = 0$$

Which after substituting earlier equations, simplifies to

$$5a_5 - \frac{3}{8} = 0$$

Or

$$a_5 = \frac{3}{40}$$

$n = 5$ gives

$$6a_6 - 2a_4 - \frac{a_2}{4} = 0$$

Which after substituting earlier equations, simplifies to

$$6a_6 = 0$$

Or

$$a_6 = 0$$

$n = 6$ gives

$$7a_7 - \frac{5a_5}{2} - \frac{3a_3}{8} - \frac{a_1}{16} = 0$$

Which after substituting earlier equations, simplifies to

$$7a_7 - \frac{5}{16} = 0$$

Or

$$a_7 = \frac{5}{112}$$

$n = 7$ gives

$$8a_8 - 3a_6 - \frac{a_4}{2} - \frac{a_2}{8} = 0$$

Which after substituting earlier equations, simplifies to

$$8a_8 = 0$$

Or

$$a_8 = 0$$

For $8 \leq n$, the recurrence equation is

$$\left((1+n)a_{1+n} - \frac{(n-1)a_{n-1}}{2} - \frac{(n-3)a_{n-3}}{8} - \frac{(n-5)a_{n-5}}{16} - \frac{5(n-7)a_{n-7}}{128} \right) x^n = 1 \quad (4)$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + x + \frac{1}{6}x^3 + \frac{3}{40}x^5 + \frac{5}{112}x^7 + \dots$$

Collecting terms, the solution becomes

$$y = a_0 + x + \frac{x^3}{6} + \frac{3x^5}{40} + \frac{5x^7}{112} + O(x^8) \quad (3)$$

Summary

The solution(s) found are the following

$$y = y(0) + x + \frac{x^3}{6} + \frac{3x^5}{40} + \frac{5x^7}{112} + O(x^8) \quad (1)$$

$$y = c_1 + x + \frac{x^3}{6} + \frac{3x^5}{40} + \frac{5x^7}{112} + O(x^8) \quad (2)$$




Figure 468: Slope field plot

Verification of solutions

$$y = y(0) + x + \frac{x^3}{6} + \frac{3x^5}{40} + \frac{5x^7}{112} + O(x^8)$$

Verified OK.

$$y = c_1 + x + \frac{x^3}{6} + \frac{3x^5}{40} + \frac{5x^7}{112} + O(x^8)$$

Verified OK.

17.20.2 Maple step by step solution

Let's solve

$$y' \sqrt{-x^2 + 1} = 1$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$y' = \frac{1}{\sqrt{-x^2+1}}$$

- Integrate both sides with respect to x

$$\int y' dx = \int \frac{1}{\sqrt{-x^2+1}} dx + c_1$$

- Evaluate integral

$$y = \arcsin(x) + c_1$$

- Solve for y

$$y = \arcsin(x) + c_1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
<- quadrature successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
Order:=8;  
dsolve(diff(y(x),x)=(1-x^2)^(-1/2),y(x),type='series',x=0);
```

$$y(x) = y(0) + x + \frac{x^3}{6} + \frac{3x^5}{40} + \frac{5x^7}{112} + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 28

```
AsymptoticDSolveValue[y'[x]==(1-x^2)^(-1/2),y[x],{x,0,7}]
```

$$y(x) \rightarrow \frac{5x^7}{112} + \frac{3x^5}{40} + \frac{x^3}{6} + x + c_1$$

17.21 problem 4

17.21.1 Solving as series ode	2923
17.21.2 Maple step by step solution	2933

Internal problem ID [6424]

Internal file name [OUTPUT/5672_Sunday_June_05_2022_03_46_35_PM_19500384/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.2. Series Solutions of First-Order Differential Equations Page 162

Problem number: 4.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"first order ode series method. Ordinary point", "first order ode series method. Taylor series method"**

Maple gives the following as the ode type

`[_quadrature]`

$$y' - y = 1$$

With the expansion point for the power series method at $x = 0$.

17.21.1 Solving as series ode

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving first order ode. Let

$$y' = f(x, y)$$

Where $f(x, y)$ is analytic at expansion point x_0 . We can always shift to $x_0 = 0$ if x_0 is not zero. So from now we assume $x_0 = 0$. Assume also that $y(x_0) = y_0$. Using Taylor

series

$$\begin{aligned}
 y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \dots \\
 &= y_0 + xf + \frac{x^2}{2} \left. \frac{df}{dx} \right|_{x_0, y_0} + \frac{x^3}{3!} \left. \frac{d^2f}{dx^2} \right|_{x_0, y_0} + \dots \\
 &= y_0 + \sum_{n=0}^{\infty} \frac{x^{n+1}}{(n+1)!} \left. \frac{d^n f}{dx^n} \right|_{x_0, y_0}
 \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} f \tag{1}$$

$$\begin{aligned}
 \frac{d^2f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) f
 \end{aligned} \tag{2}$$

$$\begin{aligned}
 \frac{d^3f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2f}{dx^2} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{d^2f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2f}{dx^2} \right) f
 \end{aligned} \tag{3}$$

⋮

And so on. Hence if we name $F_0 = f(x, y)$ then the above can be written as

$$F_0 = f(x, y) \tag{4}$$

$$\begin{aligned}
 F_n &= \frac{d}{dx}(F_{n-1}) \\
 &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) F_0
 \end{aligned} \tag{5}$$

For example, for $n = 1$ we see that

$$\begin{aligned}
 F_1 &= \frac{d}{dx}(F_0) \\
 &= \frac{\partial}{\partial x} F_0 + \left(\frac{\partial F_0}{\partial y} \right) F_0 \\
 &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} f
 \end{aligned}$$

Which is (1). And when $n = 2$

$$\begin{aligned} F_2 &= \frac{d}{dx}(F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) F_0 \\ &= \frac{\partial}{\partial x} \left(\frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} f \right) + \frac{\partial}{\partial y} \left(\frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} f \right) f \\ &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) f \end{aligned}$$

Which is (2) and so on. Therefore (4,5) can be used from now on along with

$$y(x) = y_0 + \sum_{n=0}^{\infty} \frac{x^{n+1}}{(n+1)!} F_n|_{x_0, y_0} \quad (6)$$

Hence

$$\begin{aligned}F_0 &= 1 + y \\F_1 &= \frac{dF_0}{dx} \\&= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} F_0 \\&= 1 + y \\F_2 &= \frac{dF_1}{dx} \\&= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} F_1 \\&= 1 + y \\F_3 &= \frac{dF_2}{dx} \\&= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} F_2 \\&= 1 + y \\F_4 &= \frac{dF_3}{dx} \\&= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} F_3 \\&= 1 + y \\F_5 &= \frac{dF_4}{dx} \\&= \frac{\partial F_4}{\partial x} + \frac{\partial F_4}{\partial y} F_4 \\&= 1 + y \\F_6 &= \frac{dF_5}{dx} \\&= \frac{\partial F_5}{\partial x} + \frac{\partial F_5}{\partial y} F_5 \\&= 1 + y\end{aligned}$$

And so on. Evaluating all the above at initial conditions $x(0) = 0$ and $y(0) = y(0)$ gives

$$F_0 = 1 + y(0)$$

$$F_1 = 1 + y(0)$$

$$F_2 = 1 + y(0)$$

$$F_3 = 1 + y(0)$$

$$F_4 = 1 + y(0)$$

$$F_5 = 1 + y(0)$$

$$F_6 = 1 + y(0)$$

Substituting all the above in (6) and simplifying gives the solution as

$$y = \left(1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{720}x^6 + \frac{1}{5040}x^7 \right) y(0) \\ + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \frac{x^5}{120} + \frac{x^6}{720} + \frac{x^7}{5040} + O(x^8)$$

Since $x = 0$ is also an ordinary point, then standard power series can also be used. Writing the ODE as

$$y' + q(x)y = p(x) \\ y' - y = 1$$

Where

$$q(x) = -1 \\ p(x) = 1$$

Next, the type of the expansion point $x = 0$ is determined. This point can be an ordinary point, a regular singular point (also called removable singularity), or irregular singular point (also called non-removable singularity or essential singularity). When $x = 0$ is an ordinary point, then the standard power series is used. If the point is a regular singular point, Frobenius series is used instead. Irregular singular point requires more advanced methods (asymptotic methods) and is not supported now. Hopefully this will be added in the future. $x = 0$ is called an ordinary point $q(x)$ has a Taylor series expansion around the point $x = 0$. $x = 0$ is called a regular singular point if $q(x)$ is not analytic at $x = 0$ but $xq(x)$ has Taylor series expansion. And finally, $x = 0$ is an irregular singular point if the point is not ordinary and not regular singular. This

is the most complicated case. Now the expansion point $x = 0$ is checked to see if it is an ordinary point or not. Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$y' = \sum_{n=1}^{\infty} n a_n x^{n-1}$$

Substituting the above back into the ode gives

$$\left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) - \left(\sum_{n=0}^{\infty} a_n x^n \right) = 1 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) + \sum_{n=0}^{\infty} (-a_n x^n) = 1 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} (n+1) a_{n+1} x^n \right) + \sum_{n=0}^{\infty} (-a_n x^n) = 1 \quad (3)$$

For $0 \leq n$, the recurrence equation is

$$((n+1) a_{n+1} - a_n) x^n = 1 \quad (4)$$

For $n = 0$ the recurrence equation gives

$$\begin{aligned} (a_1 - a_0) 1 &= 1 \\ a_1 - a_0 &= 1 \end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_1 = 1 + a_0$$

For $n = 1$ the recurrence equation gives

$$\begin{aligned}(2a_2 - a_1)x &= 0 \\ 2a_2 - a_1 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_2 = \frac{1}{2} + \frac{a_0}{2}$$

For $n = 2$ the recurrence equation gives

$$\begin{aligned}(3a_3 - a_2)x^2 &= 0 \\ 3a_3 - a_2 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_3 = \frac{1}{6} + \frac{a_0}{6}$$

For $n = 3$ the recurrence equation gives

$$\begin{aligned}(4a_4 - a_3)x^3 &= 0 \\ 4a_4 - a_3 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_4 = \frac{1}{24} + \frac{a_0}{24}$$

For $n = 4$ the recurrence equation gives

$$\begin{aligned}(5a_5 - a_4)x^4 &= 0 \\ 5a_5 - a_4 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_5 = \frac{1}{120} + \frac{a_0}{120}$$

For $n = 5$ the recurrence equation gives

$$\begin{aligned}(6a_6 - a_5)x^5 &= 0 \\ 6a_6 - a_5 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_6 = \frac{1}{720} + \frac{a_0}{720}$$

For $n = 6$ the recurrence equation gives

$$\begin{aligned}(7a_7 - a_6)x^6 &= 0 \\ 7a_7 - a_6 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_7 = \frac{1}{5040} + \frac{a_0}{5040}$$

For $n = 7$ the recurrence equation gives

$$\begin{aligned}(8a_8 - a_7)x^7 &= 0 \\ 8a_8 - a_7 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_8 = \frac{1}{40320} + \frac{a_0}{40320}$$

And so on. Therefore the solution is

$$\begin{aligned}y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots\end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + (1 + a_0)x + \left(\frac{1}{2} + \frac{a_0}{2}\right)x^2 + \left(\frac{1}{6} + \frac{a_0}{6}\right)x^3 + \left(\frac{1}{24} + \frac{a_0}{24}\right)x^4 \\ + \left(\frac{1}{120} + \frac{a_0}{120}\right)x^5 + \left(\frac{1}{720} + \frac{a_0}{720}\right)x^6 + \left(\frac{1}{5040} + \frac{a_0}{5040}\right)x^7 + \dots$$

Collecting terms, the solution becomes

$$y = \left(1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{720}x^6 + \frac{1}{5040}x^7\right) a_0 \quad (3) \\ + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \frac{x^5}{120} + \frac{x^6}{720} + \frac{x^7}{5040} + O(x^8)$$

Summary

The solution(s) found are the following

$$y = \left(1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{720}x^6 + \frac{1}{5040}x^7\right) y(0) \quad (1) \\ + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \frac{x^5}{120} + \frac{x^6}{720} + \frac{x^7}{5040} + O(x^8)$$

$$y = \left(1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{720}x^6 + \frac{1}{5040}x^7\right) c_1 \quad (2) \\ + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \frac{x^5}{120} + \frac{x^6}{720} + \frac{x^7}{5040} + O(x^8)$$




Figure 469: Slope field plot

Verification of solutions

$$y = \left(1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{720}x^6 + \frac{1}{5040}x^7 \right) y(0) \\ + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \frac{x^5}{120} + \frac{x^6}{720} + \frac{x^7}{5040} + O(x^8)$$

Verified OK.

$$y = \left(1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{720}x^6 + \frac{1}{5040}x^7 \right) c_1 \\ + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \frac{x^5}{120} + \frac{x^6}{720} + \frac{x^7}{5040} + O(x^8)$$

Verified OK.

17.21.2 Maple step by step solution

Let's solve

$$y' - y = 1$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{1+y} = 1$$

- Integrate both sides with respect to x

$$\int \frac{y'}{1+y} dx = \int 1 dx + c_1$$

- Evaluate integral

$$\ln(1 + y) = x + c_1$$

- Solve for y

$$y = e^{x+c_1} - 1$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 58

```
Order:=8;  
dsolve(diff(y(x),x)=1+y(x),y(x),type='series',x=0);
```

$$y(x) = \left(1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{720}x^6 + \frac{1}{5040}x^7\right) y(0) \\ + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \frac{x^5}{120} + \frac{x^6}{720} + \frac{x^7}{5040} + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 95

```
AsymptoticDSolveValue[y'[x]==1+y[x],y[x],{x,0,7}]
```

$$y(x) \rightarrow \frac{x^7}{5040} + \frac{x^6}{720} + \frac{x^5}{120} + \frac{x^4}{24} + \frac{x^3}{6} + \frac{x^2}{2} + c_1 \left(\frac{x^7}{5040} + \frac{x^6}{720} + \frac{x^5}{120} + \frac{x^4}{24} + \frac{x^3}{6} + \frac{x^2}{2} + x + 1 \right) + x$$

17.22 problem 5 solved using series

17.22.1 Existence and uniqueness analysis	2935
17.22.2 Solving as series ode	2936
17.22.3 Maple step by step solution	2945

Internal problem ID [6425]

Internal file name [OUTPUT/5673_Sunday_June_05_2022_03_46_37_PM_28499956/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.2. Series Solutions of First-Order Differential Equations Page 162

Problem number: 5 solved using series.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "linear", "first_order_ode_lie_symmetry_lookup", "first order ode series method. Ordinary point", "first order ode series method. Taylor series method"

Maple gives the following as the ode type

```
[[_linear, `class A`]]
```

$$y + y' = x$$

With initial conditions

$$[y(0) = 0]$$

With the expansion point for the power series method at $x = 0$.

17.22.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = 1$$

$$q(x) = x$$

Hence the ode is

$$y + y' = x$$

The domain of $p(x) = 1$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = x$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

17.22.2 Solving as series ode

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving first order ode. Let

$$y' = f(x, y)$$

Where $f(x, y)$ is analytic at expansion point x_0 . We can always shift to $x_0 = 0$ if x_0 is not zero. So from now we assume $x_0 = 0$. Assume also that $y(x_0) = y_0$. Using Taylor series

$$\begin{aligned} y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \dots \\ &= y_0 + xf + \frac{x^2}{2} \left. \frac{df}{dx} \right|_{x_0, y_0} + \frac{x^3}{3!} \left. \frac{d^2f}{dx^2} \right|_{x_0, y_0} + \dots \\ &= y_0 + \sum_{n=0}^{\infty} \frac{x^{n+1}}{(n+1)!} \left. \frac{d^n f}{dx^n} \right|_{x_0, y_0} \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} f \tag{1}$$

$$\begin{aligned} \frac{d^2f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) f \end{aligned} \tag{2}$$

$$\begin{aligned} \frac{d^3f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2f}{dx^2} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{d^2f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2f}{dx^2} \right) f \end{aligned} \tag{3}$$

⋮

And so on. Hence if we name $F_0 = f(x, y)$ then the above can be written as

$$F_0 = f(x, y) \tag{4}$$

$$\begin{aligned} F_n &= \frac{d}{dx}(F_{n-1}) \\ &= \frac{\partial}{\partial x}F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y}\right) F_0 \end{aligned} \tag{5}$$

For example, for $n = 1$ we see that

$$\begin{aligned} F_1 &= \frac{d}{dx}(F_0) \\ &= \frac{\partial}{\partial x}F_0 + \left(\frac{\partial F_0}{\partial y}\right) F_0 \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}f \end{aligned}$$

Which is (1). And when $n = 2$

$$\begin{aligned} F_2 &= \frac{d}{dx}(F_1) \\ &= \frac{\partial}{\partial x}F_1 + \left(\frac{\partial F_1}{\partial y}\right) F_0 \\ &= \frac{\partial}{\partial x}\left(\frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}f\right) + \frac{\partial}{\partial y}\left(\frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}f\right) f \\ &= \frac{\partial}{\partial x}\left(\frac{df}{dx}\right) + \frac{\partial}{\partial y}\left(\frac{df}{dx}\right) f \end{aligned}$$

Which is (2) and so on. Therefore (4,5) can be used from now on along with

$$y(x) = y_0 + \sum_{n=0}^{\infty} \frac{x^{n+1}}{(n+1)!} F_n|_{x_0, y_0} \tag{6}$$

Hence

$$\begin{aligned}F_0 &= x - y \\F_1 &= \frac{dF_0}{dx} \\&= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} F_0 \\&= -x + y + 1 \\F_2 &= \frac{dF_1}{dx} \\&= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} F_1 \\&= x - y - 1 \\F_3 &= \frac{dF_2}{dx} \\&= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} F_2 \\&= -x + y + 1 \\F_4 &= \frac{dF_3}{dx} \\&= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} F_3 \\&= x - y - 1 \\F_5 &= \frac{dF_4}{dx} \\&= \frac{\partial F_4}{\partial x} + \frac{\partial F_4}{\partial y} F_4 \\&= -x + y + 1 \\F_6 &= \frac{dF_5}{dx} \\&= \frac{\partial F_5}{\partial x} + \frac{\partial F_5}{\partial y} F_5 \\&= x - y - 1\end{aligned}$$

And so on. Evaluating all the above at initial conditions $x(0) = 0$ and $y(0) = 0$ gives

$$F_0 = 0$$

$$F_1 = 1$$

$$F_2 = -1$$

$$F_3 = 1$$

$$F_4 = -1$$

$$F_5 = 1$$

$$F_6 = -1$$

Substituting all the above in (6) and simplifying gives the solution as

$$y = \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + \frac{x^6}{720} - \frac{x^7}{5040} + O(x^8)$$

Now we substitute the given initial conditions in the above to solve for $y(0)$. Solving for $y(0)$ from initial conditions gives

$$y(0) = y(0)$$

Therefore the solution becomes

$$y = \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{24}x^4 - \frac{1}{120}x^5 + \frac{1}{720}x^6 - \frac{1}{5040}x^7$$

Hence the solution can be written as

$$y = \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + \frac{x^6}{720} - \frac{x^7}{5040} + O(x^8)$$

which simplifies to

$$y = \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + \frac{x^6}{720} - \frac{x^7}{5040} + O(x^8)$$

Since $x = 0$ is also an ordinary point, then standard power series can also be used. Writing the ODE as

$$y' + q(x)y = p(x)$$

$$y + y' = x$$

Where

$$q(x) = 1$$

$$p(x) = x$$

Next, the type of the expansion point $x = 0$ is determined. This point can be an ordinary point, a regular singular point (also called removable singularity), or irregular singular point (also called non-removable singularity or essential singularity). When $x = 0$ is an ordinary point, then the standard power series is used. If the point is a regular singular point, Frobenius series is used instead. Irregular singular point requires more advanced methods (asymptotic methods) and is not supported now. Hopefully this will be added in the future. $x = 0$ is called an ordinary point $q(x)$ has a Taylor series expansion around the point $x = 0$. $x = 0$ is called a regular singular point if $q(x)$ is not not analytic at $x = 0$ but $xq(x)$ has Taylor series expansion. And finally, $x = 0$ is an irregular singular point if the point is not ordinary and not regular singular. This is the most complicated case. Now the expansion point $x = 0$ is checked to see if it is an ordinary point or not. Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$y' = \sum_{n=1}^{\infty} n a_n x^{n-1}$$

Substituting the above back into the ode gives

$$\left(\sum_{n=0}^{\infty} a_n x^n \right) + \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) = x \tag{1}$$

Expanding x as Taylor series around $x = 0$ and keeping only the first 8 terms gives

$$\begin{aligned} x &= x + \dots \\ &= x \end{aligned}$$

Hence the ODE in Eq (1) becomes

$$\left(\sum_{n=0}^{\infty} a_n x^n \right) + \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) = x \tag{1}$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} a_n x^n \right) + \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) = x \tag{2}$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the

power and the corresponding index gives

$$\sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} (n+1) a_{n+1} x^n \right) + \left(\sum_{n=0}^{\infty} a_n x^n \right) = x \quad (3)$$

For $0 \leq n$, the recurrence equation is

$$((n+1) a_{n+1} + a_n) x^n = x \quad (4)$$

For $n = 0$ the recurrence equation gives

$$\begin{aligned} (a_1 + a_0) 1 &= 0 \\ a_1 + a_0 &= 0 \end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_1 = -a_0$$

For $n = 1$ the recurrence equation gives

$$\begin{aligned} (2a_2 + a_1) x &= x \\ 2a_2 + a_1 &= 1 \end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_2 = \frac{1}{2} + \frac{a_0}{2}$$

For $n = 2$ the recurrence equation gives

$$\begin{aligned} (3a_3 + a_2) x^2 &= 0 \\ 3a_3 + a_2 &= 0 \end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_3 = -\frac{1}{6} - \frac{a_0}{6}$$

For $n = 3$ the recurrence equation gives

$$\begin{aligned}(4a_4 + a_3)x^3 &= 0 \\ 4a_4 + a_3 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_4 = \frac{1}{24} + \frac{a_0}{24}$$

For $n = 4$ the recurrence equation gives

$$\begin{aligned}(5a_5 + a_4)x^4 &= 0 \\ 5a_5 + a_4 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_5 = -\frac{1}{120} - \frac{a_0}{120}$$

For $n = 5$ the recurrence equation gives

$$\begin{aligned}(6a_6 + a_5)x^5 &= 0 \\ 6a_6 + a_5 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_6 = \frac{1}{720} + \frac{a_0}{720}$$

For $n = 6$ the recurrence equation gives

$$\begin{aligned}(7a_7 + a_6)x^6 &= 0 \\ 7a_7 + a_6 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_7 = -\frac{1}{5040} - \frac{a_0}{5040}$$

For $n = 7$ the recurrence equation gives

$$\begin{aligned}(8a_8 + a_7)x^7 &= 0 \\ 8a_8 + a_7 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_8 = \frac{1}{40320} + \frac{a_0}{40320}$$

And so on. Therefore the solution is

$$\begin{aligned}y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots\end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$\begin{aligned}y &= a_0 - a_0 x + \left(\frac{1}{2} + \frac{a_0}{2}\right)x^2 + \left(-\frac{1}{6} - \frac{a_0}{6}\right)x^3 + \left(\frac{1}{24} + \frac{a_0}{24}\right)x^4 \\ &\quad + \left(-\frac{1}{120} - \frac{a_0}{120}\right)x^5 + \left(\frac{1}{720} + \frac{a_0}{720}\right)x^6 + \left(-\frac{1}{5040} - \frac{a_0}{5040}\right)x^7 + \dots\end{aligned}$$

Collecting terms, the solution becomes

$$\begin{aligned}y &= \left(1 - x + \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{24}x^4 - \frac{1}{120}x^5 + \frac{1}{720}x^6 - \frac{1}{5040}x^7\right)a_0 \\ &\quad + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + \frac{x^6}{720} - \frac{x^7}{5040} + O(x^8)\end{aligned}\tag{3}$$

At $x = 0$ the solution above becomes

$$y(0) = a_0$$

Therefore the solution in Eq(3) now can be written as

$$y = \left(1 - x + \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{24}x^4 - \frac{1}{120}x^5 + \frac{1}{720}x^6 - \frac{1}{5040}x^7 \right) y(0) \\ + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + \frac{x^6}{720} - \frac{x^7}{5040} + O(x^8)$$

Now we substitute the given initial conditions in the above to solve for $y(0)$. Solving for $y(0)$ from initial conditions gives

$$y(0) = 0$$

Therefore the solution becomes

$$y = \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{24}x^4 - \frac{1}{120}x^5 + \frac{1}{720}x^6 - \frac{1}{5040}x^7$$

Hence the solution can be written as

$$y = \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + \frac{x^6}{720} - \frac{x^7}{5040} + O(x^8)$$

which simplifies to

$$y = \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + \frac{x^6}{720} - \frac{x^7}{5040} + O(x^8)$$

Summary

The solution(s) found are the following

$$y = \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + \frac{x^6}{720} - \frac{x^7}{5040} + O(x^8) \quad (1)$$

$$y = \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + \frac{x^6}{720} - \frac{x^7}{5040} + O(x^8) \quad (2)$$

Verification of solutions

$$y = \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + \frac{x^6}{720} - \frac{x^7}{5040} + O(x^8)$$

Verified OK.

$$y = \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + \frac{x^6}{720} - \frac{x^7}{5040} + O(x^8)$$

Verified OK.

17.22.3 Maple step by step solution

Let's solve

$$[y + y' = x, y(0) = 0]$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = x - y$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y + y' = x$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x)(y + y') = \mu(x)x$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x)y)$

$$\mu(x)(y + y') = \mu'(x)y + \mu(x)y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = \mu(x)$$

- Solve to find the integrating factor

$$\mu(x) = e^x$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x)y) \right) dx = \int \mu(x)x dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x)y = \int \mu(x)x dx + c_1$$

- Solve for y

$$y = \frac{\int \mu(x)x dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = e^x$

$$y = \frac{\int x e^x dx + c_1}{e^x}$$

- Evaluate the integrals on the rhs

$$y = \frac{(x-1)e^x + c_1}{e^x}$$

- Simplify

$$y = x - 1 + c_1 e^{-x}$$

- Use initial condition $y(0) = 0$
 $0 = -1 + c_1$
- Solve for c_1
 $c_1 = 1$
- Substitute $c_1 = 1$ into general solution and simplify
 $y = x - 1 + e^{-x}$
- Solution to the IVP
 $y = x - 1 + e^{-x}$

Maple trace

```
`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
Order:=8;
dsolve([diff(y(x),x)=x-y(x),y(0) = 0],y(x),type='series',x=0);
```

$$y(x) = \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{24}x^4 - \frac{1}{120}x^5 + \frac{1}{720}x^6 - \frac{1}{5040}x^7 + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.042 (sec). Leaf size: 46

```
AsymptoticDSolveValue[{y'[x]==x-y[x],{y[0]==0}},y[x],{x,0,7}]
```

$$y(x) \rightarrow -\frac{x^7}{5040} + \frac{x^6}{720} - \frac{x^5}{120} + \frac{x^4}{24} - \frac{x^3}{6} + \frac{x^2}{2}$$

17.23 problem 5 solved directly

17.23.1 Existence and uniqueness analysis	2947
17.23.2 Solving as linear ode	2948
17.23.3 Solving as first order ode lie symmetry lookup ode	2950
17.23.4 Solving as exact ode	2954
17.23.5 Maple step by step solution	2958

Internal problem ID [6426]

Internal file name [OUTPUT/5674_Sunday_June_05_2022_03_46_39_PM_47121641/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.2. Series Solutions of First-Order Differential Equations Page 162

Problem number: 5 solved directly.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"linear"**, **"exactWithIntegrationFactor"**, **"first_order_ode_lie_symmetry_lookup"**

Maple gives the following as the ode type

```
[[_linear, `class A`]]
```

$$y + y' = x$$

With initial conditions

$$[y(0) = 0]$$

17.23.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = 1$$

$$q(x) = x$$

Hence the ode is

$$y + y' = x$$

The domain of $p(x) = 1$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = x$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

17.23.2 Solving as linear ode

Entering Linear first order ODE solver. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int 1 dx} \\ &= e^x\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu)(x) \\ \frac{d}{dx}(y e^x) &= (e^x)(x) \\ d(y e^x) &= (x e^x) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}y e^x &= \int x e^x dx \\ y e^x &= (x - 1) e^x + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^x$ results in

$$y = e^{-x}(x - 1) e^x + c_1 e^{-x}$$

which simplifies to

$$y = x - 1 + c_1 e^{-x}$$

Initial conditions are used to solve for c_1 . Substituting $x = 0$ and $y = 0$ in the above solution gives an equation to solve for the constant of integration.

$$0 = -1 + c_1$$

$$c_1 = 1$$


Substituting c_1 found above in the general solution gives

$$y = x - 1 + e^{-x}$$


Summary

The solution(s) found are the following

$$y = x - 1 + e^{-x} \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = x - 1 + e^{-x}$$

Verified OK.

17.23.3 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = x - y$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 426: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= e^{-x}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^{-x}} dy\end{aligned}$$

Which results in

$$S = y e^x$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y}\tag{2}$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = x - y$$

Evaluating all the partial derivatives gives

$$\begin{aligned}R_x &= 1 \\ R_y &= 0 \\ S_x &= y e^x \\ S_y &= e^x\end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = x e^x \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = R e^R$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = (R - 1) e^R + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$e^x y = (x - 1) e^x + c_1$$



Which simplifies to

$$e^x y = (x - 1) e^x + c_1$$

Which gives

$$y = (x e^x - e^x + c_1) e^{-x}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = x - y$ 	$R = x$ $S = y e^x$	$\frac{dS}{dR} = R e^R$ 

Initial conditions are used to solve for c_1 . Substituting $x = 0$ and $y = 0$ in the above solution gives an equation to solve for the constant of integration.

$$0 = -1 + c_1$$

$$c_1 = 1$$


Substituting c_1 found above in the general solution gives

$$y = x - 1 + e^{-x}$$


Summary

The solution(s) found are the following

$$y = x - 1 + e^{-x} \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = x - 1 + e^{-x}$$

Verified OK.

17.23.4 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned} dy &= (x - y) dx \\ (-x + y) dx + dy &= 0 \end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -x + y \\ N(x, y) &= 1 \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-x + y) \\ &= 1 \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(1) \\ &= 0 \end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= 1((1) - (0)) \\ &= 1 \end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int A dx} \\ &= e^{\int 1 dx}\end{aligned}$$

The result of integrating gives

$$\begin{aligned}\mu &= e^x \\ &= e^x\end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} for now so not to confuse them with the original M and N .

$$\begin{aligned}\bar{M} &= \mu M \\ &= e^x(-x + y) \\ &= -e^x(x - y)\end{aligned}$$

And

$$\begin{aligned}\bar{N} &= \mu N \\ &= e^x(1) \\ &= e^x\end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned}\bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ (-e^x(x - y)) + (e^x) \frac{dy}{dx} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -e^x(x - y) dx \\ \phi &= -(x - y - 1)e^x + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = e^x + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = e^x$. Therefore equation (4) becomes

$$e^x = e^x + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = -(x - y - 1)e^x + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -(x - y - 1)e^x$$

The solution becomes

$$y = (xe^x - e^x + c_1)e^{-x}$$

Initial conditions are used to solve for c_1 . Substituting $x = 0$ and $y = 0$ in the above solution gives an equation to solve for the constant of integration.

$$0 = -1 + c_1$$

$$c_1 = 1$$


Substituting c_1 found above in the general solution gives

$$y = x - 1 + e^{-x}$$


Summary

The solution(s) found are the following

$$y = x - 1 + e^{-x} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = x - 1 + e^{-x}$$

Verified OK.

17.23.5 Maple step by step solution

Let's solve

$$[y + y' = x, y(0) = 0]$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = x - y$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y + y' = x$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x)(y + y') = \mu(x)x$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x)y)$

$$\mu(x)(y + y') = \mu'(x)y + \mu(x)y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = \mu(x)$$

- Solve to find the integrating factor

$$\mu(x) = e^x$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x)y) \right) dx = \int \mu(x)x dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x)y = \int \mu(x)x dx + c_1$$

- Solve for y

$$y = \frac{\int \mu(x)x dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = e^x$

$$y = \frac{\int x e^x dx + c_1}{e^x}$$

- Evaluate the integrals on the rhs

$$y = \frac{(x-1)e^x + c_1}{e^x}$$

- Simplify

$$y = x - 1 + c_1 e^{-x}$$

- Use initial condition $y(0) = 0$

$$0 = -1 + c_1$$

- Solve for c_1

$$c_1 = 1$$

- Substitute $c_1 = 1$ into general solution and simplify

$$y = x - 1 + e^{-x}$$

- Solution to the IVP

$$y = x - 1 + e^{-x}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve([diff(y(x),x)=x-y(x),y(0) = 0],y(x), singsol=all)
```

$$y(x) = x - 1 + e^{-x}$$

✓ Solution by Mathematica

Time used: 0.039 (sec). Leaf size: 13

```
DSolve[{y'[x]==x-y[x],{y[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x + e^{-x} - 1$$

18 Chapter 4. Power Series Solutions and Special Functions. Section 4.3. Second-Order Linear Equations: Ordinary Points. Page 169

18.1	problem 1(a)	2962
18.2	problem 1(b)	2972
18.3	problem 1(c)	2983
18.4	problem 1(d)	2994
18.5	problem 1(e)	3006
18.6	problem 1(f)	3015
18.7	problem 2	3026
18.8	problem 3	3035
18.9	problem 4(a)	3045
18.10	problem 4(b)	3057
18.11	problem 5	3069
18.12	problem 6	3083
18.13	problem 7	3093
18.14	problem 8	3106

18.1 problem 1(a)

18.1.1 Maple step by step solution 2970

Internal problem ID [6427]

Internal file name [OUTPUT/5675_Sunday_June_05_2022_03_46_40_PM_83521006/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.3. Second-Order Linear Equations: Ordinary Points. Page 169

Problem number: 1(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact linear second order ode", "second_order_integrable_as_is", "second order series method. Ordinary point", "second order series method. Taylor series method"

Maple gives the following as the ode type

```
[[_2nd_order , _exact , _linear , _homogeneous]]
```

$$y'' + xy' + y = 0$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using

Taylor series gives

$$\begin{aligned}
 y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \dots \\
 &= y_0 + xy'_0 + \frac{x^2}{2}f|_{x_0, y_0, y'_0} + \frac{x^3}{3!}f'|_{x_0, y_0, y'_0} + \dots \\
 &= y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0}
 \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \tag{1}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \tag{647}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \tag{648}$$

$$\begin{aligned}
 \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \tag{2}
 \end{aligned}$$

$$\begin{aligned}
 \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \tag{3}
 \end{aligned}$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \tag{4}$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \tag{5}$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \end{aligned}$$

⋮

$$\begin{aligned} F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \tag{6}$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \tag{7}$$

To find $y(x)$ series solution around $x = 0$. Hence

$$\begin{aligned}
 F_0 &= -y - xy' \\
 F_1 &= \frac{dF_0}{dx} \\
 &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \\
 &= x^2 y' + xy - 2y' \\
 F_2 &= \frac{dF_1}{dx} \\
 &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} y' + \frac{\partial F_1}{\partial y'} F_1 \\
 &= -y' x^3 - yx^2 + 5xy' + 3y \\
 F_3 &= \frac{dF_2}{dx} \\
 &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} y' + \frac{\partial F_2}{\partial y'} F_2 \\
 &= (x^4 - 9x^2 + 8) y' + yx(x^2 - 7) \\
 F_4 &= \frac{dF_3}{dx} \\
 &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} y' + \frac{\partial F_3}{\partial y'} F_3 \\
 &= (-x^5 + 14x^3 - 33x) y' - y(x^4 - 12x^2 + 15) \\
 F_5 &= \frac{dF_4}{dx} \\
 &= \frac{\partial F_4}{\partial x} + \frac{\partial F_4}{\partial y} y' + \frac{\partial F_4}{\partial y'} F_4 \\
 &= (x^6 - 20x^4 + 87x^2 - 48) y' + yx(x^4 - 18x^2 + 57) \\
 F_6 &= \frac{dF_5}{dx} \\
 &= \frac{\partial F_5}{\partial x} + \frac{\partial F_5}{\partial y} y' + \frac{\partial F_5}{\partial y'} F_5 \\
 &= (-x^7 + 27x^5 - 185x^3 + 279x) y' - y(x^2 - 7)(x^4 - 18x^2 + 15)
 \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and

$y'(0) = y'(0)$ gives

$$\begin{aligned} F_0 &= -y(0) \\ F_1 &= -2y'(0) \\ F_2 &= 3y(0) \\ F_3 &= 8y'(0) \\ F_4 &= -15y(0) \\ F_5 &= -48y'(0) \\ F_6 &= 105y(0) \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$y = \left(1 - \frac{1}{2}x^2 + \frac{1}{8}x^4 - \frac{1}{48}x^6 + \frac{1}{384}x^8\right) y(0) + \left(x - \frac{1}{3}x^3 + \frac{1}{15}x^5 - \frac{1}{105}x^7\right) y'(0) + O(x^8)$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard power series. Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$\begin{aligned} y' &= \sum_{n=1}^{\infty} n a_n x^{n-1} \\ y'' &= \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \end{aligned}$$

Substituting the above back into the ode gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = - \left(\sum_{n=0}^{\infty} a_n x^n \right) - x \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) \quad (1)$$

Which simplifies to

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \left(\sum_{n=1}^{\infty} n x^n a_n \right) + \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the

power and the corresponding index gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n \right) + \left(\sum_{n=1}^{\infty} n x^n a_n \right) + \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0 \quad (3)$$

$n = 0$ gives

$$2a_2 + a_0 = 0$$

$$a_2 = -\frac{a_0}{2}$$

For $1 \leq n$, the recurrence equation is

$$(n+2) a_{n+2} (n+1) + n a_n + a_n = 0 \quad (4)$$

Solving for a_{n+2} , gives

$$a_{n+2} = -\frac{a_n}{n+2} \quad (5)$$

For $n = 1$ the recurrence equation gives

$$6a_3 + 2a_1 = 0$$

Which after substituting the earlier terms found becomes

$$a_3 = -\frac{a_1}{3}$$

For $n = 2$ the recurrence equation gives

$$12a_4 + 3a_2 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = \frac{a_0}{8}$$

For $n = 3$ the recurrence equation gives

$$20a_5 + 4a_3 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = \frac{a_1}{15}$$

For $n = 4$ the recurrence equation gives

$$30a_6 + 5a_4 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = -\frac{a_0}{48}$$

For $n = 5$ the recurrence equation gives

$$42a_7 + 6a_5 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = -\frac{a_1}{105}$$

For $n = 6$ the recurrence equation gives

$$56a_8 + 7a_6 = 0$$

Which after substituting the earlier terms found becomes

$$a_8 = \frac{a_0}{384}$$

For $n = 7$ the recurrence equation gives

$$72a_9 + 8a_7 = 0$$

Which after substituting the earlier terms found becomes

$$a_9 = \frac{a_1}{945}$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + a_1 x - \frac{1}{2} a_0 x^2 - \frac{1}{3} a_1 x^3 + \frac{1}{8} a_0 x^4 + \frac{1}{15} a_1 x^5 - \frac{1}{48} a_0 x^6 - \frac{1}{105} a_1 x^7 + \dots$$

Collecting terms, the solution becomes

$$y = \left(1 - \frac{1}{2}x^2 + \frac{1}{8}x^4 - \frac{1}{48}x^6\right) a_0 + \left(x - \frac{1}{3}x^3 + \frac{1}{15}x^5 - \frac{1}{105}x^7\right) a_1 + O(x^8) \quad (3)$$

At $x = 0$ the solution above becomes

$$y = \left(1 - \frac{1}{2}x^2 + \frac{1}{8}x^4 - \frac{1}{48}x^6\right) c_1 + \left(x - \frac{1}{3}x^3 + \frac{1}{15}x^5 - \frac{1}{105}x^7\right) c_2 + O(x^8)$$

Summary

The solution(s) found are the following

$$\begin{aligned} y &= \left(1 - \frac{1}{2}x^2 + \frac{1}{8}x^4 - \frac{1}{48}x^6 + \frac{1}{384}x^8\right) y(0) \\ &\quad + \left(x - \frac{1}{3}x^3 + \frac{1}{15}x^5 - \frac{1}{105}x^7\right) y'(0) + O(x^8) \end{aligned} \quad (1)$$

$$y = \left(1 - \frac{1}{2}x^2 + \frac{1}{8}x^4 - \frac{1}{48}x^6\right) c_1 + \left(x - \frac{1}{3}x^3 + \frac{1}{15}x^5 - \frac{1}{105}x^7\right) c_2 + O(x^8) \quad (2)$$

Verification of solutions

$$y = \left(1 - \frac{1}{2}x^2 + \frac{1}{8}x^4 - \frac{1}{48}x^6 + \frac{1}{384}x^8\right) y(0) + \left(x - \frac{1}{3}x^3 + \frac{1}{15}x^5 - \frac{1}{105}x^7\right) y'(0) + O(x^8)$$

Verified OK.

$$y = \left(1 - \frac{1}{2}x^2 + \frac{1}{8}x^4 - \frac{1}{48}x^6\right) c_1 + \left(x - \frac{1}{3}x^3 + \frac{1}{15}x^5 - \frac{1}{105}x^7\right) c_2 + O(x^8)$$

Verified OK.

18.1.1 Maple step by step solution

Let's solve

$$y'' = -y - xy'$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + xy' + y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^k$$

- Rewrite DE with series expansions

- Convert $x \cdot y'$ to series expansion

$$x \cdot y' = \sum_{k=0}^{\infty} a_k k x^k$$

- Convert y'' to series expansion

$$y'' = \sum_{k=2}^{\infty} a_k k(k-1) x^{k-2}$$

- Shift index using $k- > k+2$

$$y'' = \sum_{k=0}^{\infty} a_{k+2}(k+2)(k+1) x^k$$

Rewrite DE with series expansions

$$\sum_{k=0}^{\infty} (a_{k+2}(k+2)(k+1) + a_k(k+1)) x^k = 0$$

- Each term in the series must be 0, giving the recursion relation

$$(k+1)(a_{k+2}(k+2) + a_k) = 0$$

- Recursion relation that defines the series solution to the ODE

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+2} = -\frac{a_k}{k+2} \right]$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
<- linear_1 successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 44

```
Order:=8;  
dsolve(diff(y(x),x$2)+x*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{2}x^2 + \frac{1}{8}x^4 - \frac{1}{48}x^6\right) y(0) + \left(x - \frac{1}{3}x^3 + \frac{1}{15}x^5 - \frac{1}{105}x^7\right) D(y)(0) + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 56

```
AsymptoticDSolveValue[y''[x]+x*y'[x]+y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_2 \left(-\frac{x^7}{105} + \frac{x^5}{15} - \frac{x^3}{3} + x \right) + c_1 \left(-\frac{x^6}{48} + \frac{x^4}{8} - \frac{x^2}{2} + 1 \right)$$

18.2 problem 1(b)

18.2.1 Maple step by step solution 2980

Internal problem ID [6428]

Internal file name [OUTPUT/5676_Sunday_June_05_2022_03_46_42_PM_36469378/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.3. Second-Order Linear Equations: Ordinary Points. Page 169

Problem number: 1(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_airy", "second order series method. Ordinary point", "second order series method. Taylor series method"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - y' + xy = 0$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using

Taylor series gives

$$\begin{aligned}
 y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \cdots \\
 &= y_0 + xy'_0 + \frac{x^2}{2}f|_{x_0, y_0, y'_0} + \frac{x^3}{3!}f'|_{x_0, y_0, y'_0} + \cdots \\
 &= y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0}
 \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \tag{1}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \tag{650}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \tag{651}$$

$$\begin{aligned}
 \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \tag{2}
 \end{aligned}$$

$$\begin{aligned}
 \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \tag{3}
 \end{aligned}$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \tag{4}$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \tag{5}$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \end{aligned}$$

⋮

$$\begin{aligned} F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \tag{6}$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \tag{7}$$

To find $y(x)$ series solution around $x = 0$. Hence

$$\begin{aligned}
 F_0 &= y' - xy \\
 F_1 &= \frac{dF_0}{dx} \\
 &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \\
 &= (1 - x) y' - (1 + x) y \\
 F_2 &= \frac{dF_1}{dx} \\
 &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} y' + \frac{\partial F_1}{\partial y'} F_1 \\
 &= (-1 - 2x) y' + y(x^2 - x - 1) \\
 F_3 &= \frac{dF_2}{dx} \\
 &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} y' + \frac{\partial F_2}{\partial y'} F_2 \\
 &= (x^2 - 3x - 4) y' + y(2x^2 + 3x - 1) \\
 F_4 &= \frac{dF_3}{dx} \\
 &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} y' + \frac{\partial F_3}{\partial y'} F_3 \\
 &= (3x^2 + 2x - 8) y' - y(x^3 - 3x^2 - 8x - 3) \\
 F_5 &= \frac{dF_4}{dx} \\
 &= \frac{\partial F_4}{\partial x} + \frac{\partial F_4}{\partial y} y' + \frac{\partial F_4}{\partial y'} F_4 \\
 &= (-x^3 + 6x^2 + 16x - 3) y' - 3 \left(x^3 + \frac{5}{3} x^2 - \frac{14}{3} x - \frac{8}{3} \right) y \\
 F_6 &= \frac{dF_5}{dx} \\
 &= \frac{\partial F_5}{\partial x} + \frac{\partial F_5}{\partial y} y' + \frac{\partial F_5}{\partial y'} F_5 \\
 &= (-4x^3 - 2x^2 + 42x + 21) y' + y(x^4 - 6x^3 - 25x^2 - 7x + 14)
 \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and

$y'(0) = y'(0)$ gives

$$\begin{aligned} F_0 &= y'(0) \\ F_1 &= -y(0) + y'(0) \\ F_2 &= -y'(0) - y(0) \\ F_3 &= -4y'(0) - y(0) \\ F_4 &= -8y'(0) + 3y(0) \\ F_5 &= -3y'(0) + 8y(0) \\ F_6 &= 21y'(0) + 14y(0) \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$\begin{aligned} y &= \left(1 - \frac{1}{6}x^3 - \frac{1}{24}x^4 - \frac{1}{120}x^5 + \frac{1}{240}x^6 + \frac{1}{630}x^7 + \frac{1}{2880}x^8\right) y(0) \\ &+ \left(x + \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{24}x^4 - \frac{1}{30}x^5 - \frac{1}{90}x^6 - \frac{1}{1680}x^7 + \frac{1}{1920}x^8\right) y'(0) + O(x^8) \end{aligned}$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard power series Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$\begin{aligned} y' &= \sum_{n=1}^{\infty} n a_n x^{n-1} \\ y'' &= \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \end{aligned}$$

Substituting the above back into the ode gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \left(\sum_{n=1}^{\infty} n a_n x^{n-1}\right) - x \left(\sum_{n=0}^{\infty} a_n x^n\right) \quad (1)$$

Which simplifies to

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}\right) + \sum_{n=1}^{\infty} (-n a_n x^{n-1}) + \left(\sum_{n=0}^{\infty} x^{1+n} a_n\right) = 0 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\begin{aligned}\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} &= \sum_{n=0}^{\infty} (n+2) a_{n+2} (1+n) x^n \\ \sum_{n=1}^{\infty} (-n a_n x^{n-1}) &= \sum_{n=0}^{\infty} (-(1+n) a_{1+n} x^n) \\ \sum_{n=0}^{\infty} x^{1+n} a_n &= \sum_{n=1}^{\infty} a_{n-1} x^n\end{aligned}$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (1+n) x^n \right) + \sum_{n=0}^{\infty} (-(1+n) a_{1+n} x^n) + \left(\sum_{n=1}^{\infty} a_{n-1} x^n \right) = 0 \quad (3)$$

$n = 0$ gives

$$2a_2 - a_1 = 0$$

$$a_2 = \frac{a_1}{2}$$

For $1 \leq n$, the recurrence equation is

$$(n+2) a_{n+2} (1+n) - (1+n) a_{1+n} + a_{n-1} = 0 \quad (4)$$

Solving for a_{n+2} , gives

$$\begin{aligned}a_{n+2} &= \frac{n a_{1+n} + a_{1+n} - a_{n-1}}{(n+2)(1+n)} \\ (5) \quad &= \frac{a_{1+n}}{n+2} - \frac{a_{n-1}}{(n+2)(1+n)}\end{aligned}$$

For $n = 1$ the recurrence equation gives

$$6a_3 - 2a_2 + a_0 = 0$$

Which after substituting the earlier terms found becomes

$$a_3 = \frac{a_1}{6} - \frac{a_0}{6}$$

For $n = 2$ the recurrence equation gives

$$12a_4 - 3a_3 + a_1 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = -\frac{a_1}{24} - \frac{a_0}{24}$$

For $n = 3$ the recurrence equation gives

$$20a_5 - 4a_4 + a_2 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = -\frac{a_1}{30} - \frac{a_0}{120}$$

For $n = 4$ the recurrence equation gives

$$30a_6 - 5a_5 + a_3 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = -\frac{a_1}{90} + \frac{a_0}{240}$$

For $n = 5$ the recurrence equation gives

$$42a_7 - 6a_6 + a_4 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = -\frac{a_1}{1680} + \frac{a_0}{630}$$

For $n = 6$ the recurrence equation gives

$$56a_8 - 7a_7 + a_5 = 0$$

Which after substituting the earlier terms found becomes

$$a_8 = \frac{a_1}{1920} + \frac{a_0}{2880}$$

For $n = 7$ the recurrence equation gives

$$72a_9 - 8a_8 + a_6 = 0$$

Which after substituting the earlier terms found becomes

$$a_9 = \frac{11a_1}{51840} - \frac{a_0}{51840}$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$\begin{aligned} y &= a_0 + a_1 x + \frac{a_1 x^2}{2} + \left(\frac{a_1}{6} - \frac{a_0}{6}\right) x^3 + \left(-\frac{a_1}{24} - \frac{a_0}{24}\right) x^4 \\ &\quad + \left(-\frac{a_1}{30} - \frac{a_0}{120}\right) x^5 + \left(-\frac{a_1}{90} + \frac{a_0}{240}\right) x^6 + \left(-\frac{a_1}{1680} + \frac{a_0}{630}\right) x^7 + \dots \end{aligned}$$

Collecting terms, the solution becomes

$$\begin{aligned} y &= \left(1 - \frac{1}{6}x^3 - \frac{1}{24}x^4 - \frac{1}{120}x^5 + \frac{1}{240}x^6 + \frac{1}{630}x^7\right) a_0 \\ &\quad + \left(x + \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{24}x^4 - \frac{1}{30}x^5 - \frac{1}{90}x^6 - \frac{1}{1680}x^7\right) a_1 + O(x^8) \end{aligned} \tag{3}$$

At $x = 0$ the solution above becomes

$$\begin{aligned} y &= \left(1 - \frac{1}{6}x^3 - \frac{1}{24}x^4 - \frac{1}{120}x^5 + \frac{1}{240}x^6 + \frac{1}{630}x^7\right) c_1 \\ &\quad + \left(x + \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{24}x^4 - \frac{1}{30}x^5 - \frac{1}{90}x^6 - \frac{1}{1680}x^7\right) c_2 + O(x^8) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \left(1 - \frac{1}{6}x^3 - \frac{1}{24}x^4 - \frac{1}{120}x^5 + \frac{1}{240}x^6 + \frac{1}{630}x^7 + \frac{1}{2880}x^8\right) y(0) + \left(x + \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{24}x^4 - \frac{1}{30}x^5 - \frac{1}{90}x^6 - \frac{1}{1680}x^7 + \frac{1}{1920}x^8\right) y'(0) + O(x^8) \quad (1)$$

$$y = \left(1 - \frac{1}{6}x^3 - \frac{1}{24}x^4 - \frac{1}{120}x^5 + \frac{1}{240}x^6 + \frac{1}{630}x^7\right) c_1 + \left(x + \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{24}x^4 - \frac{1}{30}x^5 - \frac{1}{90}x^6 - \frac{1}{1680}x^7\right) c_2 + O(x^8) \quad (2)$$

Verification of solutions

$$y = \left(1 - \frac{1}{6}x^3 - \frac{1}{24}x^4 - \frac{1}{120}x^5 + \frac{1}{240}x^6 + \frac{1}{630}x^7 + \frac{1}{2880}x^8\right) y(0) + \left(x + \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{24}x^4 - \frac{1}{30}x^5 - \frac{1}{90}x^6 - \frac{1}{1680}x^7 + \frac{1}{1920}x^8\right) y'(0) + O(x^8)$$

Verified OK.

$$y = \left(1 - \frac{1}{6}x^3 - \frac{1}{24}x^4 - \frac{1}{120}x^5 + \frac{1}{240}x^6 + \frac{1}{630}x^7\right) c_1 + \left(x + \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{24}x^4 - \frac{1}{30}x^5 - \frac{1}{90}x^6 - \frac{1}{1680}x^7\right) c_2 + O(x^8)$$

Verified OK.

18.2.1 Maple step by step solution

Let's solve

$$y'' = y' - xy$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - y' + xy = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^k$$

- Rewrite ODE with series expansions

- Convert $x \cdot y$ to series expansion

$$x \cdot y = \sum_{k=0}^{\infty} a_k x^{k+1}$$

- Shift index using $k \rightarrow k - 1$

$$x \cdot y = \sum_{k=1}^{\infty} a_{k-1} x^k$$

- Convert y' to series expansion

$$y' = \sum_{k=1}^{\infty} a_k k x^{k-1}$$

- Shift index using $k \rightarrow k + 1$

$$y' = \sum_{k=0}^{\infty} a_{k+1} (k+1) x^k$$

- Convert y'' to series expansion

$$y'' = \sum_{k=2}^{\infty} a_k k(k-1) x^{k-2}$$

- Shift index using $k \rightarrow k + 2$

$$y'' = \sum_{k=0}^{\infty} a_{k+2} (k+2)(k+1) x^k$$

Rewrite ODE with series expansions

$$2a_2 - a_1 + \left(\sum_{k=1}^{\infty} (a_{k+2}(k+2)(k+1) - a_{k+1}(k+1) + a_{k-1}) x^k \right) = 0$$

- Each term must be 0

$$2a_2 - a_1 = 0$$

- Each term in the series must be 0, giving the recursion relation

$$(k^2 + 3k + 2) a_{k+2} - a_{k+1} k + a_{k-1} - a_{k+1} = 0$$

- Shift index using $k \rightarrow k + 1$

$$((k+1)^2 + 3k + 5) a_{k+3} - a_{k+2}(k+1) + a_k - a_{k+2} = 0$$

- Recursion relation that defines the series solution to the ODE

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+3} = \frac{ka_{k+2} - a_k + 2a_{k+2}}{k^2 + 5k + 6}, 2a_2 - a_1 = 0 \right]$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacic's algorithm  
<- No Liouvillian solutions exist  
-> Trying a solution in terms of special functions:  
  -> Bessel  
  <- Bessel successful  
<- special function solution successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 69

```
Order:=8;  
dsolve(diff(y(x),x$2)-diff(y(x),x)+x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{6}x^3 - \frac{1}{24}x^4 - \frac{1}{120}x^5 + \frac{1}{240}x^6 + \frac{1}{630}x^7\right) y(0) \\ + \left(x + \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{24}x^4 - \frac{1}{30}x^5 - \frac{1}{90}x^6 - \frac{1}{1680}x^7\right) D(y)(0) + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 91

```
AsymptoticDSolveValue[y'[x]-y[x]+x*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(\frac{x^7}{630} + \frac{x^6}{240} - \frac{x^5}{120} - \frac{x^4}{24} - \frac{x^3}{6} + 1 \right) + c_2 \left(-\frac{x^7}{1680} - \frac{x^6}{90} - \frac{x^5}{30} - \frac{x^4}{24} + \frac{x^3}{6} + \frac{x^2}{2} + x \right)$$

18.3 problem 1(c)

Internal problem ID [6429]

Internal file name [OUTPUT/5677_Sunday_June_05_2022_03_46_43_PM_57202053/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.3. Second-Order Linear Equations: Ordinary Points. Page 169

Problem number: 1(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second order series method. Ordinary point", "second order series method. Taylor series method"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' + 2xy' - y = x$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using Taylor series gives

$$\begin{aligned} y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \cdots \\ &= y_0 + xy'_0 + \frac{x^2}{2}f|_{x_0, y_0, y'_0} + \frac{x^3}{3!}f'|_{x_0, y_0, y'_0} + \cdots \\ &= y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0} \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \quad (1)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \quad (653)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \quad (654)$$

$$\begin{aligned} \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \end{aligned} \quad (2)$$

$$\begin{aligned} \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \end{aligned} \quad (3)$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \tag{4}$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \tag{5}$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \end{aligned}$$

⋮

$$\begin{aligned} F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \tag{6}$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \tag{7}$$

To find $y(x)$ series solution around $x = 0$. Hence

$$F_0 = -2xy' + y + x$$

$$\begin{aligned} F_1 &= \frac{dF_0}{dx} \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \\ &= 4x^2 y' - 2xy - 2x^2 - y' + 1 \end{aligned}$$

$$\begin{aligned} F_2 &= \frac{dF_1}{dx} \\ &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} y' + \frac{\partial F_1}{\partial y'} F_1 \\ &= (-8x^3 + 8x) y' + 4x^3 + 4yx^2 - 5x - 3y \end{aligned}$$

$$\begin{aligned} F_3 &= \frac{dF_2}{dx} \\ &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} y' + \frac{\partial F_2}{\partial y'} F_2 \\ &= (16x^4 - 36x^2 + 5) y' + (-8x^3 + 16x) y - 8x^4 + 20x^2 - 5 \end{aligned}$$

$$\begin{aligned} F_4 &= \frac{dF_3}{dx} \\ &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} y' + \frac{\partial F_3}{\partial y'} F_3 \\ &= (-32x^5 + 128x^3 - 66x) y' + (16x^4 - 60x^2 + 21) y + 16x^5 - 68x^3 + 45x \end{aligned}$$

$$\begin{aligned} F_5 &= \frac{dF_4}{dx} \\ &= \frac{\partial F_4}{\partial x} + \frac{\partial F_4}{\partial y} y' + \frac{\partial F_4}{\partial y'} F_4 \\ &= (64x^6 - 400x^4 + 456x^2 - 45) y' + (-32x^5 + 192x^3 - 186x) y - 32x^6 + 208x^4 - 270x^2 + 45 \end{aligned}$$

$$\begin{aligned} F_6 &= \frac{dF_5}{dx} \\ &= \frac{\partial F_5}{\partial x} + \frac{\partial F_5}{\partial y} y' + \frac{\partial F_5}{\partial y'} F_5 \\ &= (-128x^7 + 1152x^5 - 2320x^3 + 816x) y' + (64x^6 - 560x^4 + 1032x^2 - 231) y + 64x^7 - 592x^5 + 1288x \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and

$y'(0) = y'(0)$ gives

$$\begin{aligned} F_0 &= y(0) \\ F_1 &= -y'(0) + 1 \\ F_2 &= -3y(0) \\ F_3 &= 5y'(0) - 5 \\ F_4 &= 21y(0) \\ F_5 &= -45y'(0) + 45 \\ F_6 &= -231y(0) \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$\begin{aligned} y &= \left(1 + \frac{1}{2}x^2 - \frac{1}{8}x^4 + \frac{7}{240}x^6 - \frac{11}{1920}x^8\right) y(0) \\ &\quad + \left(x - \frac{1}{6}x^3 + \frac{1}{24}x^5 - \frac{1}{112}x^7\right) y'(0) + \frac{x^3}{6} - \frac{x^5}{24} + \frac{x^7}{112} + O(x^8) \end{aligned}$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard power series. Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$\begin{aligned} y' &= \sum_{n=1}^{\infty} n a_n x^{n-1} \\ y'' &= \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \end{aligned}$$

Substituting the above back into the ode gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = -2x \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) + \left(\sum_{n=0}^{\infty} a_n x^n \right) + x \quad (1)$$

Expanding x as Taylor series around $x = 0$ and keeping only the first 8 terms gives

$$\begin{aligned} x &= x + \dots \\ &= x \end{aligned}$$

Hence the ODE in Eq (1) becomes

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + 2x \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) - \left(\sum_{n=0}^{\infty} a_n x^n \right) = x$$

Which simplifies to

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \left(\sum_{n=1}^{\infty} 2n x^n a_n \right) + \sum_{n=0}^{\infty} (-a_n x^n) = x \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n \right) + \left(\sum_{n=1}^{\infty} 2n x^n a_n \right) + \sum_{n=0}^{\infty} (-a_n x^n) = x \quad (3)$$

$n = 0$ gives

$$2a_2 - a_0 = 0$$

$$a_2 = \frac{a_0}{2}$$

For $1 \leq n$, the recurrence equation is

$$((n+2) a_{n+2} (n+1) + 2n a_n - a_n) x^n = x \quad (4)$$

For $n = 1$ the recurrence equation gives

$$(6a_3 + a_1) x = x$$

$$6a_3 + a_1 = 1$$

Which after substituting the earlier terms found becomes

$$a_3 = \frac{1}{6} - \frac{a_1}{6}$$

For $n = 2$ the recurrence equation gives

$$\begin{aligned}(12a_4 + 3a_2)x^2 &= 0 \\ 12a_4 + 3a_2 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_4 = -\frac{a_0}{8}$$

For $n = 3$ the recurrence equation gives

$$\begin{aligned}(20a_5 + 5a_3)x^3 &= 0 \\ 20a_5 + 5a_3 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_5 = -\frac{1}{24} + \frac{a_1}{24}$$

For $n = 4$ the recurrence equation gives

$$\begin{aligned}(30a_6 + 7a_4)x^4 &= 0 \\ 30a_6 + 7a_4 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_6 = \frac{7a_0}{240}$$

For $n = 5$ the recurrence equation gives

$$\begin{aligned}(42a_7 + 9a_5)x^5 &= 0 \\ 42a_7 + 9a_5 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_7 = \frac{1}{112} - \frac{a_1}{112}$$

For $n = 6$ the recurrence equation gives

$$\begin{aligned}(56a_8 + 11a_6)x^6 &= 0 \\ 56a_8 + 11a_6 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_8 = -\frac{11a_0}{1920}$$

For $n = 7$ the recurrence equation gives

$$\begin{aligned}(72a_9 + 13a_7)x^7 &= 0 \\ 72a_9 + 13a_7 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_9 = -\frac{13}{8064} + \frac{13a_1}{8064}$$

And so on. Therefore the solution is

$$\begin{aligned}y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots\end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$\begin{aligned}y &= a_0 + a_1 x + \frac{a_0 x^2}{2} + \left(\frac{1}{6} - \frac{a_1}{6}\right) x^3 - \frac{a_0 x^4}{8} \\ &\quad + \left(-\frac{1}{24} + \frac{a_1}{24}\right) x^5 + \frac{7a_0 x^6}{240} + \left(\frac{1}{112} - \frac{a_1}{112}\right) x^7 + \dots\end{aligned}$$

Collecting terms, the solution becomes

$$y = \left(1 + \frac{1}{2}x^2 - \frac{1}{8}x^4 + \frac{7}{240}x^6\right) a_0 + \left(x - \frac{1}{6}x^3 + \frac{1}{24}x^5 - \frac{1}{112}x^7\right) a_1 + \frac{x^3}{6} - \frac{x^5}{24} + \frac{x^7}{112} + O(x^8) \quad (3)$$

At $x = 0$ the solution above becomes

$$y = \left(1 + \frac{1}{2}x^2 - \frac{1}{8}x^4 + \frac{7}{240}x^6\right) c_1 + \left(x - \frac{1}{6}x^3 + \frac{1}{24}x^5 - \frac{1}{112}x^7\right) c_2 + \frac{x^3}{6} - \frac{x^5}{24} + \frac{x^7}{112} + O(x^8)$$

Summary

The solution(s) found are the following

$$y = \left(1 + \frac{1}{2}x^2 - \frac{1}{8}x^4 + \frac{7}{240}x^6 - \frac{11}{1920}x^8\right) y(0) + \left(x - \frac{1}{6}x^3 + \frac{1}{24}x^5 - \frac{1}{112}x^7\right) y'(0) + \frac{x^3}{6} - \frac{x^5}{24} + \frac{x^7}{112} + O(x^8) \quad (1)$$

$$y = \left(1 + \frac{1}{2}x^2 - \frac{1}{8}x^4 + \frac{7}{240}x^6\right) c_1 + \left(x - \frac{1}{6}x^3 + \frac{1}{24}x^5 - \frac{1}{112}x^7\right) c_2 + \frac{x^3}{6} - \frac{x^5}{24} + \frac{x^7}{112} + O(x^8) \quad (2)$$

Verification of solutions

$$y = \left(1 + \frac{1}{2}x^2 - \frac{1}{8}x^4 + \frac{7}{240}x^6 - \frac{11}{1920}x^8\right) y(0) + \left(x - \frac{1}{6}x^3 + \frac{1}{24}x^5 - \frac{1}{112}x^7\right) y'(0) + \frac{x^3}{6} - \frac{x^5}{24} + \frac{x^7}{112} + O(x^8)$$

Verified OK.

$$y = \left(1 + \frac{1}{2}x^2 - \frac{1}{8}x^4 + \frac{7}{240}x^6\right) c_1 + \left(x - \frac{1}{6}x^3 + \frac{1}{24}x^5 - \frac{1}{112}x^7\right) c_2 + \frac{x^3}{6} - \frac{x^5}{24} + \frac{x^7}{112} + O(x^8)$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exists
-> Trying a solution in terms of special functions:
    -> Bessel
    -> elliptic
    -> Legendre
    <- Kummer successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 50

```
Order:=8;
dsolve(diff(y(x),x$2)+2*x*diff(y(x),x)-y(x)=x,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{1}{2}x^2 - \frac{1}{8}x^4 + \frac{7}{240}x^6\right) y(0) \\ + \left(x - \frac{1}{6}x^3 + \frac{1}{24}x^5 - \frac{1}{112}x^7\right) D(y)(0) + \frac{x^3}{6} - \frac{x^5}{24} + \frac{x^7}{112} + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 77

```
AsymptoticDSolveValue[y''[x]+2*x*y'[x]-y[x]==x,y[x],{x,0,7}]
```

$$y(x) \rightarrow \frac{x^7}{112} - \frac{x^5}{24} + \frac{x^3}{6} + c_2 \left(-\frac{x^7}{112} + \frac{x^5}{24} - \frac{x^3}{6} + x \right) + c_1 \left(\frac{7x^6}{240} - \frac{x^4}{8} + \frac{x^2}{2} + 1 \right)$$

18.4 problem 1(d)

Internal problem ID [6430]

Internal file name [OUTPUT/5678_Sunday_June_05_2022_03_46_45_PM_58879577/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.3. Second-Order Linear Equations: Ordinary Points. Page 169

Problem number: 1(d).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second order series method. Ordinary point", "second order series method. Taylor series method"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + y' - yx^2 = 1$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using Taylor series gives

$$\begin{aligned} y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \cdots \\ &= y_0 + xy'_0 + \frac{x^2}{2}f|_{x_0, y_0, y'_0} + \frac{x^3}{3!}f'|_{x_0, y_0, y'_0} + \cdots \\ &= y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0} \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \quad (1)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \quad (656)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \quad (657)$$

$$\begin{aligned} \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \end{aligned} \quad (2)$$

$$\begin{aligned} \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \end{aligned} \quad (3)$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \tag{4}$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \tag{5}$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \end{aligned}$$

⋮

$$\begin{aligned} F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \tag{6}$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \tag{7}$$

To find $y(x)$ series solution around $x = 0$. Hence

$$F_0 = -y' + yx^2 + 1$$

$$\begin{aligned} F_1 &= \frac{dF_0}{dx} \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \\ &= y' - yx^2 - 1 + x^2 y' + 2xy \end{aligned}$$

$$\begin{aligned} F_2 &= \frac{dF_1}{dx} \\ &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} y' + \frac{\partial F_1}{\partial y'} F_1 \\ &= (-2x^2 + 4x - 1) y' + (x^4 + x^2 - 2x + 2) y + x^2 + 1 \end{aligned}$$

$$\begin{aligned} F_3 &= \frac{dF_2}{dx} \\ &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} y' + \frac{\partial F_2}{\partial y'} F_2 \\ &= (x^4 + 3x^2 - 10x + 7) y' + (-2x^4 + 8x^3 - x^2 + 2x - 2) y - 2x^2 + 6x - 1 \end{aligned}$$

$$\begin{aligned} F_4 &= \frac{dF_3}{dx} \\ &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} y' + \frac{\partial F_3}{\partial y'} F_3 \\ &= (-3x^4 + 12x^3 - 4x^2 + 18x - 19) y' + (x^6 + 3x^4 - 18x^3 + 31x^2 - 2x + 2) y + x^4 + 3x^2 - 14x + 13 \end{aligned}$$

$$\begin{aligned} F_5 &= \frac{dF_4}{dx} \\ &= \frac{\partial F_4}{\partial x} + \frac{\partial F_4}{\partial y} y' + \frac{\partial F_4}{\partial y'} F_4 \\ &= (x^6 + 6x^4 - 42x^3 + 71x^2 - 28x + 39) y' + (-3x^6 + 18x^5 - 4x^4 + 30x^3 - 73x^2 + 62x - 2) y - 3x^4 + \end{aligned}$$

$$\begin{aligned} F_6 &= \frac{dF_5}{dx} \\ &= \frac{\partial F_5}{\partial x} + \frac{\partial F_5}{\partial y} y' + \frac{\partial F_5}{\partial y'} F_5 \\ &= (-4x^6 + 24x^5 - 10x^4 + 96x^3 - 270x^2 + 232x - 69) y' + (x^8 + 6x^6 - 60x^5 + 161x^4 - 44x^3 + 129x^2 \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and

$y'(0) = y'(0)$ gives

$$\begin{aligned} F_0 &= -y'(0) + 1 \\ F_1 &= y'(0) - 1 \\ F_2 &= -y'(0) + 2y(0) + 1 \\ F_3 &= 7y'(0) - 2y(0) - 1 \\ F_4 &= -19y'(0) + 2y(0) + 13 \\ F_5 &= 39y'(0) - 2y(0) - 33 \\ F_6 &= -69y'(0) + 62y(0) + 63 \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$\begin{aligned} y &= \left(1 + \frac{1}{12}x^4 - \frac{1}{60}x^5 + \frac{1}{360}x^6 - \frac{1}{2520}x^7 + \frac{31}{20160}x^8\right) y(0) \\ &+ \left(x - \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{24}x^4 + \frac{7}{120}x^5 - \frac{19}{720}x^6 + \frac{13}{1680}x^7 - \frac{23}{13440}x^8\right) y'(0) \\ &+ \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + \frac{13x^6}{720} - \frac{11x^7}{1680} + \frac{x^8}{640} + O(x^8) \end{aligned}$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard power series Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$\begin{aligned} y' &= \sum_{n=1}^{\infty} n a_n x^{n-1} \\ y'' &= \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \end{aligned}$$

Substituting the above back into the ode gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = -\left(\sum_{n=1}^{\infty} n a_n x^{n-1}\right) + \left(\sum_{n=0}^{\infty} a_n x^n\right) x^2 + 1 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}\right) + \left(\sum_{n=1}^{\infty} n a_n x^{n-1}\right) + \sum_{n=0}^{\infty} (-x^{n+2} a_n) = 1 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\begin{aligned}\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} &= \sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n \\ \sum_{n=1}^{\infty} n a_n x^{n-1} &= \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n \\ \sum_{n=0}^{\infty} (-x^{n+2} a_n) &= \sum_{n=2}^{\infty} (-a_{n-2} x^n)\end{aligned}$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n \right) + \left(\sum_{n=0}^{\infty} (n+1) a_{n+1} x^n \right) + \sum_{n=2}^{\infty} (-a_{n-2} x^n) = 1 \quad (3)$$

$n = 0$ gives

$$\begin{aligned}(2a_2 + a_1) x^0 &= 1 \\ 2a_2 + a_1 &= 1\end{aligned}$$

$$a_2 = -\frac{a_1}{2} + \frac{1}{2}$$

$n = 1$ gives

$$6a_3 + 2a_2 = 0$$

Which after substituting earlier equations, simplifies to

$$6a_3 - a_1 + 1 = 0$$

Or

$$a_3 = \frac{a_1}{6} - \frac{1}{6}$$

For $2 \leq n$, the recurrence equation is

$$((n+2)a_{n+2}(n+1) + (n+1)a_{n+1} - a_{n-2})x^n = 1 \quad (4)$$

For $n = 2$ the recurrence equation gives

$$\begin{aligned}(12a_4 + 3a_3 - a_0)x^2 &= 0 \\ 12a_4 + 3a_3 - a_0 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_4 = \frac{1}{24} - \frac{a_1}{24} + \frac{a_0}{12}$$

For $n = 3$ the recurrence equation gives

$$\begin{aligned}(20a_5 + 4a_4 - a_1)x^3 &= 0 \\ 20a_5 + 4a_4 - a_1 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_5 = -\frac{1}{120} + \frac{7a_1}{120} - \frac{a_0}{60}$$

For $n = 4$ the recurrence equation gives

$$\begin{aligned}(30a_6 + 5a_5 - a_2)x^4 &= 0 \\ 30a_6 + 5a_5 - a_2 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_6 = \frac{13}{720} - \frac{19a_1}{720} + \frac{a_0}{360}$$

For $n = 5$ the recurrence equation gives

$$\begin{aligned}(42a_7 + 6a_6 - a_3)x^5 &= 0 \\ 42a_7 + 6a_6 - a_3 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_7 = -\frac{11}{1680} + \frac{13a_1}{1680} - \frac{a_0}{2520}$$

For $n = 6$ the recurrence equation gives

$$\begin{aligned}(56a_8 + 7a_7 - a_4)x^6 &= 0 \\ 56a_8 + 7a_7 - a_4 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_8 = \frac{1}{640} - \frac{23a_1}{13440} + \frac{31a_0}{20160}$$

For $n = 7$ the recurrence equation gives

$$\begin{aligned}(72a_9 + 8a_8 - a_5)x^7 &= 0 \\ 72a_9 + 8a_8 - a_5 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_9 = -\frac{1}{3456} + \frac{121a_1}{120960} - \frac{73a_0}{181440}$$

And so on. Therefore the solution is

$$\begin{aligned}y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots\end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$\begin{aligned}y &= a_0 + a_1 x + \left(-\frac{a_1}{2} + \frac{1}{2}\right)x^2 + \left(\frac{a_1}{6} - \frac{1}{6}\right)x^3 \\ &\quad + \left(\frac{1}{24} - \frac{a_1}{24} + \frac{a_0}{12}\right)x^4 + \left(-\frac{1}{120} + \frac{7a_1}{120} - \frac{a_0}{60}\right)x^5 \\ &\quad + \left(\frac{13}{720} - \frac{19a_1}{720} + \frac{a_0}{360}\right)x^6 + \left(-\frac{11}{1680} + \frac{13a_1}{1680} - \frac{a_0}{2520}\right)x^7 + \dots\end{aligned}$$

Collecting terms, the solution becomes

$$\begin{aligned}
 y = & \left(1 + \frac{1}{12}x^4 - \frac{1}{60}x^5 + \frac{1}{360}x^6 - \frac{1}{2520}x^7 \right) a_0 \\
 & + \left(x - \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{24}x^4 + \frac{7}{120}x^5 - \frac{19}{720}x^6 + \frac{13}{1680}x^7 \right) a_1 \\
 & + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + \frac{13x^6}{720} - \frac{11x^7}{1680} + O(x^8)
 \end{aligned} \tag{3}$$

At $x = 0$ the solution above becomes

$$\begin{aligned}
 y = & \left(1 + \frac{1}{12}x^4 - \frac{1}{60}x^5 + \frac{1}{360}x^6 - \frac{1}{2520}x^7 \right) c_1 \\
 & + \left(x - \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{24}x^4 + \frac{7}{120}x^5 - \frac{19}{720}x^6 + \frac{13}{1680}x^7 \right) c_2 \\
 & + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + \frac{13x^6}{720} - \frac{11x^7}{1680} + O(x^8)
 \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}
 y = & \left(1 + \frac{1}{12}x^4 - \frac{1}{60}x^5 + \frac{1}{360}x^6 - \frac{1}{2520}x^7 + \frac{31}{20160}x^8 \right) y(0) \\
 & + \left(x - \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{24}x^4 + \frac{7}{120}x^5 - \frac{19}{720}x^6 + \frac{13}{1680}x^7 - \frac{23}{13440}x^8 \right) y'(0) \\
 & + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + \frac{13x^6}{720} - \frac{11x^7}{1680} + \frac{x^8}{640} + O(x^8)
 \end{aligned} \tag{1}$$

$$\begin{aligned}
 y = & \left(1 + \frac{1}{12}x^4 - \frac{1}{60}x^5 + \frac{1}{360}x^6 - \frac{1}{2520}x^7 \right) c_1 \\
 & + \left(x - \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{24}x^4 + \frac{7}{120}x^5 - \frac{19}{720}x^6 + \frac{13}{1680}x^7 \right) c_2 \\
 & + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + \frac{13x^6}{720} - \frac{11x^7}{1680} + O(x^8)
 \end{aligned} \tag{2}$$

Verification of solutions

$$\begin{aligned} y &= \left(1 + \frac{1}{12}x^4 - \frac{1}{60}x^5 + \frac{1}{360}x^6 - \frac{1}{2520}x^7 + \frac{31}{20160}x^8 \right) y(0) \\ &+ \left(x - \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{24}x^4 + \frac{7}{120}x^5 - \frac{19}{720}x^6 + \frac{13}{1680}x^7 - \frac{23}{13440}x^8 \right) y'(0) \\ &+ \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + \frac{13x^6}{720} - \frac{11x^7}{1680} + \frac{x^8}{640} + O(x^8) \end{aligned}$$

Verified OK.

$$\begin{aligned} y &= \left(1 + \frac{1}{12}x^4 - \frac{1}{60}x^5 + \frac{1}{360}x^6 - \frac{1}{2520}x^7 \right) c_1 \\ &+ \left(x - \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{24}x^4 + \frac{7}{120}x^5 - \frac{19}{720}x^6 + \frac{13}{1680}x^7 \right) c_2 \\ &+ \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + \frac{13x^6}{720} - \frac{11x^7}{1680} + O(x^8) \end{aligned}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exists
-> Trying a solution in terms of special functions:
    -> Bessel
    -> elliptic
    -> Legendre
    -> Kummer
        -> hyper3: Equivalence to 1F1 under a power @ Moebius
        <- hyper3 successful: received ODE is equivalent to the 1F1 ODE
    <- Kummer successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 72

```
Order:=8;
dsolve(diff(y(x),x$2)+diff(y(x),x)-x^2*y(x)=1,y(x),type='series',x=0);
```

$$\begin{aligned} y(x) = & \left(1 + \frac{1}{12}x^4 - \frac{1}{60}x^5 + \frac{1}{360}x^6 - \frac{1}{2520}x^7\right) y(0) \\ & + \left(x - \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{24}x^4 + \frac{7}{120}x^5 - \frac{19}{720}x^6 + \frac{13}{1680}x^7\right) D(y)(0) \\ & + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + \frac{13x^6}{720} - \frac{11x^7}{1680} + O(x^8) \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 126

```
AsymptoticDSolveValue[y''[x]+y'[x]+x^2*y[x]==1,y[x],{x,0,7}]
```

$$y(x) \rightarrow \frac{31x^7}{5040} - \frac{11x^6}{720} - \frac{x^5}{120} + \frac{x^4}{24} - \frac{x^3}{6} + \frac{x^2}{2} + c_1 \left(\frac{x^7}{2520} - \frac{x^6}{360} + \frac{x^5}{60} - \frac{x^4}{12} + 1 \right) \\ + c_2 \left(-\frac{37x^7}{5040} + \frac{17x^6}{720} - \frac{x^5}{24} - \frac{x^4}{24} + \frac{x^3}{6} - \frac{x^2}{2} + x \right)$$

18.5 problem 1(e)

Internal problem ID [6431]

Internal file name [OUTPUT/5679_Sunday_June_05_2022_03_46_46_PM_76326535/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.3. Second-Order Linear Equations: Ordinary Points. Page 169

Problem number: 1(e).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second order series method. Ordinary point", "second_order_change_of_variable_on_x_method_1", "second_order_change_of_variable_on_x_method_2", "second order series method. Taylor series method"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries], [_2nd_order , _linear , `
  _with_symmetry_[0,F(x)]`]]
```

$$(x^2 + 1)y'' + xy' + y = 0$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using

Taylor series gives

$$\begin{aligned}
 y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \dots \\
 &= y_0 + xy'_0 + \frac{x^2}{2}f|_{x_0, y_0, y'_0} + \frac{x^3}{3!}f'|_{x_0, y_0, y'_0} + \dots \\
 &= y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0}
 \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \tag{1}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \tag{659}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \tag{660}$$

$$\begin{aligned}
 \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \tag{2}
 \end{aligned}$$

$$\begin{aligned}
 \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \tag{3}
 \end{aligned}$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \tag{4}$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \tag{5}$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \end{aligned}$$

⋮

$$\begin{aligned} F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \tag{6}$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \tag{7}$$

To find $y(x)$ series solution around $x = 0$. Hence

$$F_0 = -\frac{xy' + y}{x^2 + 1}$$

$$\begin{aligned} F_1 &= \frac{dF_0}{dx} \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \\ &= \frac{x^2 y' + 3xy - 2y'}{(x^2 + 1)^2} \end{aligned}$$

$$\begin{aligned} F_2 &= \frac{dF_1}{dx} \\ &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} y' + \frac{\partial F_1}{\partial y'} F_1 \\ &= \frac{-10yx^2 + 15xy' + 5y}{(x^2 + 1)^3} \end{aligned}$$

$$\begin{aligned} F_3 &= \frac{dF_2}{dx} \\ &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} y' + \frac{\partial F_2}{\partial y'} F_2 \\ &= \frac{(-10x^4 - 95x^2 + 20)y' + (40x^3 - 65x)y}{(x^2 + 1)^4} \end{aligned}$$

$$\begin{aligned} F_4 &= \frac{dF_3}{dx} \\ &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} y' + \frac{\partial F_3}{\partial y'} F_3 \\ &= \frac{(90x^5 + 600x^3 - 435x)y' + (-190x^4 + 670x^2 - 85)y}{(x^2 + 1)^5} \end{aligned}$$

$$\begin{aligned} F_5 &= \frac{dF_4}{dx} \\ &= \frac{\partial F_4}{\partial x} + \frac{\partial F_4}{\partial y} y' + \frac{\partial F_4}{\partial y'} F_4 \\ &= \frac{(-730x^6 - 3870x^4 + 6735x^2 - 520)y' + 1050(x^4 - \frac{32}{5}x^2 + \frac{5}{2})yx}{(x^2 + 1)^6} \end{aligned}$$

$$\begin{aligned} F_6 &= \frac{dF_5}{dx} \\ &= \frac{\partial F_5}{\partial x} + \frac{\partial F_5}{\partial y} y' + \frac{\partial F_5}{\partial y'} F_5 \\ &= \frac{(6160x^7 + 24780x^5 - 93660x^3 + 22855x)y' + (-6620x^6 + 69600x^4 - 55770x^2 + 3145)y}{(x^2 + 1)^7} \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and $y'(0) = y'(0)$ gives

$$\begin{aligned} F_0 &= -y(0) \\ F_1 &= -2y'(0) \\ F_2 &= 5y(0) \\ F_3 &= 20y'(0) \\ F_4 &= -85y(0) \\ F_5 &= -520y'(0) \\ F_6 &= 3145y(0) \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$y = \left(1 - \frac{1}{2}x^2 + \frac{5}{24}x^4 - \frac{17}{144}x^6 + \frac{629}{8064}x^8\right) y(0) + \left(x - \frac{1}{3}x^3 + \frac{1}{6}x^5 - \frac{13}{126}x^7\right) y'(0) + O(x^8)$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard power series The ode is normalized to be

$$(x^2 + 1)y'' + xy' + y = 0$$

Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$\begin{aligned} y' &= \sum_{n=1}^{\infty} n a_n x^{n-1} \\ y'' &= \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \end{aligned}$$

Substituting the above back into the ode gives

$$(x^2 + 1) \left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + x \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) + \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=2}^{\infty} x^n a_n n(n-1) \right) + \left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \left(\sum_{n=1}^{\infty} n a_n x^n \right) + \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=2}^{\infty} x^n a_n n(n-1) \right) + \left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n \right) + \left(\sum_{n=1}^{\infty} n a_n x^n \right) + \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0 \quad (3)$$

$n = 0$ gives

$$2a_2 + a_0 = 0$$

$$a_2 = -\frac{a_0}{2}$$

$n = 1$ gives

$$6a_3 + 2a_1 = 0$$

Which after substituting earlier equations, simplifies to

$$a_3 = -\frac{a_1}{3}$$

For $2 \leq n$, the recurrence equation is

$$n a_n (n-1) + (n+2) a_{n+2} (n+1) + n a_n + a_n = 0 \quad (4)$$

Solving for a_{n+2} , gives

$$a_{n+2} = -\frac{a_n (n^2 + 1)}{(n+2)(n+1)} \quad (5)$$

For $n = 2$ the recurrence equation gives

$$5a_2 + 12a_4 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = \frac{5a_0}{24}$$

For $n = 3$ the recurrence equation gives

$$10a_3 + 20a_5 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = \frac{a_1}{6}$$

For $n = 4$ the recurrence equation gives

$$17a_4 + 30a_6 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = -\frac{17a_0}{144}$$

For $n = 5$ the recurrence equation gives

$$26a_5 + 42a_7 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = -\frac{13a_1}{126}$$

For $n = 6$ the recurrence equation gives

$$37a_6 + 56a_8 = 0$$

Which after substituting the earlier terms found becomes

$$a_8 = \frac{629a_0}{8064}$$

For $n = 7$ the recurrence equation gives

$$50a_7 + 72a_9 = 0$$

Which after substituting the earlier terms found becomes

$$a_9 = \frac{325a_1}{4536}$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + a_1 x - \frac{1}{2} a_0 x^2 - \frac{1}{3} a_1 x^3 + \frac{5}{24} a_0 x^4 + \frac{1}{6} a_1 x^5 - \frac{17}{144} a_0 x^6 - \frac{13}{126} a_1 x^7 + \dots$$

Collecting terms, the solution becomes

$$y = \left(1 - \frac{1}{2}x^2 + \frac{5}{24}x^4 - \frac{17}{144}x^6\right) a_0 + \left(x - \frac{1}{3}x^3 + \frac{1}{6}x^5 - \frac{13}{126}x^7\right) a_1 + O(x^8) \quad (3)$$

At $x = 0$ the solution above becomes

$$y = \left(1 - \frac{1}{2}x^2 + \frac{5}{24}x^4 - \frac{17}{144}x^6\right) c_1 + \left(x - \frac{1}{3}x^3 + \frac{1}{6}x^5 - \frac{13}{126}x^7\right) c_2 + O(x^8)$$

Summary

The solution(s) found are the following

$$\begin{aligned} y &= \left(1 - \frac{1}{2}x^2 + \frac{5}{24}x^4 - \frac{17}{144}x^6 + \frac{629}{8064}x^8\right) y(0) \\ &\quad + \left(x - \frac{1}{3}x^3 + \frac{1}{6}x^5 - \frac{13}{126}x^7\right) y'(0) + O(x^8) \end{aligned} \quad (1)$$

$$y = \left(1 - \frac{1}{2}x^2 + \frac{5}{24}x^4 - \frac{17}{144}x^6\right) c_1 + \left(x - \frac{1}{3}x^3 + \frac{1}{6}x^5 - \frac{13}{126}x^7\right) c_2 + O(x^8) \quad (2)$$

Verification of solutions

$$y = \left(1 - \frac{1}{2}x^2 + \frac{5}{24}x^4 - \frac{17}{144}x^6 + \frac{629}{8064}x^8\right) y(0) + \left(x - \frac{1}{3}x^3 + \frac{1}{6}x^5 - \frac{13}{126}x^7\right) y'(0) + O(x^8)$$

Verified OK.

$$y = \left(1 - \frac{1}{2}x^2 + \frac{5}{24}x^4 - \frac{17}{144}x^6\right) c_1 + \left(x - \frac{1}{3}x^3 + \frac{1}{6}x^5 - \frac{13}{126}x^7\right) c_2 + O(x^8)$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
<- linear_1 successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 44

```
Order:=8;  
dsolve((1+x^2)*diff(y(x),x$2)+x*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{2}x^2 + \frac{5}{24}x^4 - \frac{17}{144}x^6\right) y(0) + \left(x - \frac{1}{3}x^3 + \frac{1}{6}x^5 - \frac{13}{126}x^7\right) D(y)(0) + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 56

```
AsymptoticDSolveValue[(1+x^2)*y''[x]+x*y'[x]+y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_2 \left(-\frac{13x^7}{126} + \frac{x^5}{6} - \frac{x^3}{3} + x \right) + c_1 \left(-\frac{17x^6}{144} + \frac{5x^4}{24} - \frac{x^2}{2} + 1 \right)$$

18.6 problem 1(f)

18.6.1 Maple step by step solution 3023

Internal problem ID [6432]

Internal file name [OUTPUT/5680_Sunday_June_05_2022_03_46_48_PM_58550294/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.3. Second-Order Linear Equations: Ordinary Points. Page 169

Problem number: 1(f).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second order series method. Ordinary point", "second order series method. Taylor series method", "second_order_ode_non_constant_coeff_transformation_on_B"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' + (1 + x)y' - y = 0$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using

Taylor series gives

$$\begin{aligned}
 y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \dots \\
 &= y_0 + xy'_0 + \frac{x^2}{2}f|_{x_0, y_0, y'_0} + \frac{x^3}{3!}f'|_{x_0, y_0, y'_0} + \dots \\
 &= y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0}
 \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \tag{1}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \tag{662}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \tag{663}$$

$$\begin{aligned}
 \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \tag{2}
 \end{aligned}$$

$$\begin{aligned}
 \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \tag{3}
 \end{aligned}$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \quad (4)$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \quad (5)$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \\ &\vdots \\ F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \quad (6)$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \quad (7)$$

To find $y(x)$ series solution around $x = 0$. Hence

$$\begin{aligned}
 F_0 &= -xy' - y' + y \\
 F_1 &= \frac{dF_0}{dx} \\
 &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \\
 &= (1+x)((1+x)y' - y) \\
 F_2 &= \frac{dF_1}{dx} \\
 &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y}y' + \frac{\partial F_1}{\partial y'}F_1 \\
 &= -(x+2)((1+x)y' - y)x \\
 F_3 &= \frac{dF_2}{dx} \\
 &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y}y' + \frac{\partial F_2}{\partial y'}F_2 \\
 &= (1+x)((1+x)y' - y)(x^2 + 2x - 2) \\
 F_4 &= \frac{dF_3}{dx} \\
 &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y}y' + \frac{\partial F_3}{\partial y'}F_3 \\
 &= -((1+x)y' - y)(x^4 + 4x^3 - 8x - 2) \\
 F_5 &= \frac{dF_4}{dx} \\
 &= \frac{\partial F_4}{\partial x} + \frac{\partial F_4}{\partial y}y' + \frac{\partial F_4}{\partial y'}F_4 \\
 &= (1+x)((1+x)y' - y)(x^4 + 4x^3 - 4x^2 - 16x + 6) \\
 F_6 &= \frac{dF_5}{dx} \\
 &= \frac{\partial F_5}{\partial x} + \frac{\partial F_5}{\partial y}y' + \frac{\partial F_5}{\partial y'}F_5 \\
 &= -(x^6 + 6x^5 - 40x^3 - 30x^2 + 36x + 16)((1+x)y' - y)
 \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and

$y'(0) = y'(0)$ gives

$$\begin{aligned} F_0 &= y(0) - y'(0) \\ F_1 &= -y(0) + y'(0) \\ F_2 &= 0 \\ F_3 &= 2y(0) - 2y'(0) \\ F_4 &= 2y'(0) - 2y(0) \\ F_5 &= 6y'(0) - 6y(0) \\ F_6 &= -16y'(0) + 16y(0) \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$\begin{aligned} y &= \left(1 + \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{60}x^5 - \frac{1}{360}x^6 - \frac{1}{840}x^7 + \frac{1}{2520}x^8\right) y(0) \\ &+ \left(x - \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{60}x^5 + \frac{1}{360}x^6 + \frac{1}{840}x^7 - \frac{1}{2520}x^8\right) y'(0) + O(x^8) \end{aligned}$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard power series. Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$\begin{aligned} y' &= \sum_{n=1}^{\infty} n a_n x^{n-1} \\ y'' &= \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \end{aligned}$$

Substituting the above back into the ode gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = -x \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) - \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) + \left(\sum_{n=0}^{\infty} a_n x^n \right) \quad (1)$$

Which simplifies to

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) + \left(\sum_{n=1}^{\infty} n a_n x^n \right) + \sum_{n=0}^{\infty} (-a_n x^n) = 0 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n$$

$$\sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n \right) + \left(\sum_{n=0}^{\infty} (n+1) a_{n+1} x^n \right) + \left(\sum_{n=1}^{\infty} n a_n x^n \right) + \sum_{n=0}^{\infty} (-a_n x^n) = 0 \quad (3)$$

$n = 0$ gives

$$2a_2 + a_1 - a_0 = 0$$

$$a_2 = \frac{a_0}{2} - \frac{a_1}{2}$$

For $1 \leq n$, the recurrence equation is

$$(n+2) a_{n+2} (n+1) + (n+1) a_{n+1} + n a_n - a_n = 0 \quad (4)$$

Solving for a_{n+2} , gives

$$(5) \quad a_{n+2} = -\frac{n a_n + n a_{n+1} - a_n + a_{n+1}}{(n+2)(n+1)}$$

$$= -\frac{(n-1) a_n}{(n+2)(n+1)} - \frac{a_{n+1}}{n+2}$$

For $n = 1$ the recurrence equation gives

$$6a_3 + 2a_2 = 0$$

Which after substituting the earlier terms found becomes

$$a_3 = -\frac{a_0}{6} + \frac{a_1}{6}$$

For $n = 2$ the recurrence equation gives

$$12a_4 + 3a_3 + a_2 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = 0$$

For $n = 3$ the recurrence equation gives

$$20a_5 + 4a_4 + 2a_3 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = \frac{a_0}{60} - \frac{a_1}{60}$$

For $n = 4$ the recurrence equation gives

$$30a_6 + 5a_5 + 3a_4 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = -\frac{a_0}{360} + \frac{a_1}{360}$$

For $n = 5$ the recurrence equation gives

$$42a_7 + 6a_6 + 4a_5 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = -\frac{a_0}{840} + \frac{a_1}{840}$$

For $n = 6$ the recurrence equation gives

$$56a_8 + 7a_7 + 5a_6 = 0$$

Which after substituting the earlier terms found becomes

$$a_8 = \frac{a_0}{2520} - \frac{a_1}{2520}$$

For $n = 7$ the recurrence equation gives

$$72a_9 + 8a_8 + 6a_7 = 0$$

Which after substituting the earlier terms found becomes

$$a_9 = \frac{a_0}{18144} - \frac{a_1}{18144}$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$\begin{aligned} y &= a_0 + a_1 x + \left(\frac{a_0}{2} - \frac{a_1}{2}\right) x^2 + \left(-\frac{a_0}{6} + \frac{a_1}{6}\right) x^3 + \left(\frac{a_0}{60} - \frac{a_1}{60}\right) x^5 \\ &\quad + \left(-\frac{a_0}{360} + \frac{a_1}{360}\right) x^6 + \left(-\frac{a_0}{840} + \frac{a_1}{840}\right) x^7 + \dots \end{aligned}$$

Collecting terms, the solution becomes

$$\begin{aligned} y &= \left(1 + \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{60}x^5 - \frac{1}{360}x^6 - \frac{1}{840}x^7\right) a_0 \\ &\quad + \left(x - \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{60}x^5 + \frac{1}{360}x^6 + \frac{1}{840}x^7\right) a_1 + O(x^8) \end{aligned} \tag{3}$$

At $x = 0$ the solution above becomes

$$\begin{aligned} y &= \left(1 + \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{60}x^5 - \frac{1}{360}x^6 - \frac{1}{840}x^7\right) c_1 \\ &\quad + \left(x - \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{60}x^5 + \frac{1}{360}x^6 + \frac{1}{840}x^7\right) c_2 + O(x^8) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \left(1 + \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{60}x^5 - \frac{1}{360}x^6 - \frac{1}{840}x^7 + \frac{1}{2520}x^8\right) y(0) + \left(x - \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{60}x^5 + \frac{1}{360}x^6 + \frac{1}{840}x^7 - \frac{1}{2520}x^8\right) y'(0) + O(x^8) \quad (1)$$

$$y = \left(1 + \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{60}x^5 - \frac{1}{360}x^6 - \frac{1}{840}x^7\right) c_1 + \left(x - \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{60}x^5 + \frac{1}{360}x^6 + \frac{1}{840}x^7\right) c_2 + O(x^8) \quad (2)$$

Verification of solutions

$$y = \left(1 + \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{60}x^5 - \frac{1}{360}x^6 - \frac{1}{840}x^7 + \frac{1}{2520}x^8\right) y(0) + \left(x - \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{60}x^5 + \frac{1}{360}x^6 + \frac{1}{840}x^7 - \frac{1}{2520}x^8\right) y'(0) + O(x^8)$$

Verified OK.

$$y = \left(1 + \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{60}x^5 - \frac{1}{360}x^6 - \frac{1}{840}x^7\right) c_1 + \left(x - \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{60}x^5 + \frac{1}{360}x^6 + \frac{1}{840}x^7\right) c_2 + O(x^8)$$

Verified OK.

18.6.1 Maple step by step solution

Let's solve

$$y'' = -xy' - y' + y$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + (1+x)y' - y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^k$$

- Rewrite DE with series expansions

- Convert $x^m \cdot y'$ to series expansion for $m = 0.1$

$$x^m \cdot y' = \sum_{k=\max(0,1-m)}^{\infty} a_k k x^{k-1+m}$$

- Shift index using $k \rightarrow k + 1 - m$

$$x^m \cdot y' = \sum_{k=\max(0,1-m)+m-1}^{\infty} a_{k+1-m} (k + 1 - m) x^k$$

- Convert y'' to series expansion

$$y'' = \sum_{k=2}^{\infty} a_k k(k-1) x^{k-2}$$

- Shift index using $k \rightarrow k + 2$

$$y'' = \sum_{k=0}^{\infty} a_{k+2} (k+2)(k+1) x^k$$

Rewrite DE with series expansions

$$\sum_{k=0}^{\infty} (a_{k+2}(k+2)(k+1) + a_{k+1}(k+1) + a_k(k-1)) x^k = 0$$

- Each term in the series must be 0, giving the recursion relation

$$k^2 a_{k+2} + (a_k + a_{k+1} + 3a_{k+2}) k - a_k + a_{k+1} + 2a_{k+2} = 0$$

- Recursion relation that defines the series solution to the ODE

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+2} = -\frac{a_k k + a_{k+1} k - a_k + a_{k+1}}{k^2 + 3k + 2} \right]$$

Maple trace **Kovacic algorithm successful**

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Reducible group (found an exponential solution)
    Group is reducible, not completely reducible
<- Kovacics algorithm successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 64

```
Order:=8;  
dsolve(diff(y(x),x$2)+(1+x)*diff(y(x),x)-y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{60}x^5 - \frac{1}{360}x^6 - \frac{1}{840}x^7\right) y(0) \\ + \left(x - \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{60}x^5 + \frac{1}{360}x^6 + \frac{1}{840}x^7\right) D(y)(0) + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 84

```
AsymptoticDSolveValue[y'[x]+(1+x)*y'[x]-y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(-\frac{x^7}{840} - \frac{x^6}{360} + \frac{x^5}{60} - \frac{x^3}{6} + \frac{x^2}{2} + 1 \right) + c_2 \left(\frac{x^7}{840} + \frac{x^6}{360} - \frac{x^5}{60} + \frac{x^3}{6} - \frac{x^2}{2} + x \right)$$

18.7 problem 2

Internal problem ID [6433]

Internal file name [OUTPUT/5681_Sunday_June_05_2022_03_46_50_PM_55193875/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.3. Second-Order Linear Equations: Ordinary Points. Page 169

Problem number: 2.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second order series method. Ordinary point", "second_order_change_of_variable_on_y_method_2", "second order series method. Taylor series method", "second_order_ode_non_constant_coeff_transformation_on_B"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$(x^2 + 1)y'' + 2xy' - 2y = 0$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using

Taylor series gives

$$\begin{aligned}
 y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \cdots \\
 &= y_0 + xy'_0 + \frac{x^2}{2}f|_{x_0, y_0, y'_0} + \frac{x^3}{3!}f'|_{x_0, y_0, y'_0} + \cdots \\
 &= y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0}
 \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \tag{1}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \tag{665}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \tag{666}$$

$$\begin{aligned}
 \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \tag{2}
 \end{aligned}$$

$$\begin{aligned}
 \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \tag{3}
 \end{aligned}$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \quad (4)$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \quad (5)$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \\ &\vdots \\ F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \quad (6)$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \quad (7)$$

To find $y(x)$ series solution around $x = 0$. Hence

$$\begin{aligned}
 F_0 &= -\frac{2(-y + xy')}{x^2 + 1} \\
 F_1 &= \frac{dF_0}{dx} \\
 &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \\
 &= \frac{8(-y + xy') x}{(x^2 + 1)^2} \\
 F_2 &= \frac{dF_1}{dx} \\
 &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} y' + \frac{\partial F_1}{\partial y'} F_1 \\
 &= -\frac{8(-y + xy')(5x^2 - 1)}{(x^2 + 1)^3} \\
 F_3 &= \frac{dF_2}{dx} \\
 &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} y' + \frac{\partial F_2}{\partial y'} F_2 \\
 &= \frac{240(-y + xy')(x^2 - \frac{3}{5}) x}{(x^2 + 1)^4} \\
 F_4 &= \frac{dF_3}{dx} \\
 &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} y' + \frac{\partial F_3}{\partial y'} F_3 \\
 &= -\frac{48(-y + xy')(35x^4 - 42x^2 + 3)}{(x^2 + 1)^5} \\
 F_5 &= \frac{dF_4}{dx} \\
 &= \frac{\partial F_4}{\partial x} + \frac{\partial F_4}{\partial y} y' + \frac{\partial F_4}{\partial y'} F_4 \\
 &= \frac{13440(-y + xy')(x^4 - 2x^2 + \frac{3}{7}) x}{(x^2 + 1)^6} \\
 F_6 &= \frac{dF_5}{dx} \\
 &= \frac{\partial F_5}{\partial x} + \frac{\partial F_5}{\partial y} y' + \frac{\partial F_5}{\partial y'} F_5 \\
 &= -\frac{5760(-y + xy')(21x^6 - 63x^4 + 27x^2 - 1)}{(x^2 + 1)^7}
 \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and $y'(0) = y'(0)$ gives

$$\begin{aligned} F_0 &= 2y(0) \\ F_1 &= 0 \\ F_2 &= -8y(0) \\ F_3 &= 0 \\ F_4 &= 144y(0) \\ F_5 &= 0 \\ F_6 &= -5760y(0) \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$y = \left(1 + x^2 - \frac{1}{3}x^4 + \frac{1}{5}x^6 - \frac{1}{7}x^8\right) y(0) + xy'(0) + O(x^8)$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard power series. The ode is normalized to be

$$(x^2 + 1)y'' + 2xy' - 2y = 0$$

Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$\begin{aligned} y' &= \sum_{n=1}^{\infty} n a_n x^{n-1} \\ y'' &= \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \end{aligned}$$

Substituting the above back into the ode gives

$$(x^2 + 1) \left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + 2x \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) - 2 \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=2}^{\infty} x^n a_n n(n-1) \right) + \left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \left(\sum_{n=1}^{\infty} 2n a_n x^n \right) + \sum_{n=0}^{\infty} (-2a_n x^n) = 0 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\begin{aligned} & \left(\sum_{n=2}^{\infty} x^n a_n n(n-1) \right) + \left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n \right) \\ & + \left(\sum_{n=1}^{\infty} 2n a_n x^n \right) + \sum_{n=0}^{\infty} (-2a_n x^n) = 0 \end{aligned} \quad (3)$$

$n = 0$ gives

$$2a_2 - 2a_0 = 0$$

$$a_2 = a_0$$

For $2 \leq n$, the recurrence equation is

$$n a_n (n-1) + (n+2) a_{n+2} (n+1) + 2n a_n - 2a_n = 0 \quad (4)$$

Solving for a_{n+2} , gives

$$a_{n+2} = -\frac{(n-1) a_n}{n+1} \quad (5)$$

For $n = 2$ the recurrence equation gives

$$4a_2 + 12a_4 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = -\frac{a_0}{3}$$

For $n = 3$ the recurrence equation gives

$$10a_3 + 20a_5 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = 0$$

For $n = 4$ the recurrence equation gives

$$18a_4 + 30a_6 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = \frac{a_0}{5}$$

For $n = 5$ the recurrence equation gives

$$28a_5 + 42a_7 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = 0$$

For $n = 6$ the recurrence equation gives

$$40a_6 + 56a_8 = 0$$

Which after substituting the earlier terms found becomes

$$a_8 = -\frac{a_0}{7}$$

For $n = 7$ the recurrence equation gives

$$54a_7 + 72a_9 = 0$$

Which after substituting the earlier terms found becomes

$$a_9 = 0$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + a_1 x + a_0 x^2 - \frac{1}{3} a_0 x^4 + \frac{1}{5} a_0 x^6 + \dots$$

Collecting terms, the solution becomes

$$y = \left(1 + x^2 - \frac{1}{3} x^4 + \frac{1}{5} x^6 \right) a_0 + a_1 x + O(x^8) \quad (3)$$

At $x = 0$ the solution above becomes

$$y = \left(1 + x^2 - \frac{1}{3} x^4 + \frac{1}{5} x^6 \right) c_1 + c_2 x + O(x^8)$$

Summary

The solution(s) found are the following

$$y = \left(1 + x^2 - \frac{1}{3} x^4 + \frac{1}{5} x^6 - \frac{1}{7} x^8 \right) y(0) + xy'(0) + O(x^8) \quad (1)$$

$$y = \left(1 + x^2 - \frac{1}{3} x^4 + \frac{1}{5} x^6 \right) c_1 + c_2 x + O(x^8) \quad (2)$$

Verification of solutions

$$y = \left(1 + x^2 - \frac{1}{3} x^4 + \frac{1}{5} x^6 - \frac{1}{7} x^8 \right) y(0) + xy'(0) + O(x^8)$$

Verified OK.

$$y = \left(1 + x^2 - \frac{1}{3} x^4 + \frac{1}{5} x^6 \right) c_1 + c_2 x + O(x^8)$$

Verified OK.

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacics algorithm  
  A Liouvillian solution exists  
  Reducible group (found an exponential solution)  
  Group is reducible, not completely reducible  
<- Kovacics algorithm successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
Order:=8;  
dsolve((1+x^2)*diff(y(x),x$2)+2*x*diff(y(x),x)-2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + x^2 - \frac{1}{3}x^4 + \frac{1}{5}x^6\right) y(0) + D(y)(0)x + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 30

```
AsymptoticDSolveValue[(1+x^2)*y''[x]+2*x*y'[x]-2*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(\frac{x^6}{5} - \frac{x^4}{3} + x^2 + 1 \right) + c_2 x$$

18.8 problem 3

18.8.1 Maple step by step solution 3043

Internal problem ID [6434]

Internal file name [OUTPUT/5682_Sunday_June_05_2022_03_46_51_PM_7541715/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.3. Second-Order Linear Equations: Ordinary Points. Page 169

Problem number: 3.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact linear second order ode", "second_order_integrable_as_is", "second order series method. Ordinary point", "second order series method. Taylor series method"

Maple gives the following as the ode type

```
[[_2nd_order , _exact , _linear , _homogeneous]]
```

$$y'' + xy' + y = 0$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using

Taylor series gives

$$\begin{aligned}
 y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \cdots \\
 &= y_0 + xy'_0 + \frac{x^2}{2}f|_{x_0, y_0, y'_0} + \frac{x^3}{3!}f'|_{x_0, y_0, y'_0} + \cdots \\
 &= y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0}
 \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \tag{1}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \tag{668}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \tag{669}$$

$$\begin{aligned}
 \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \tag{2}
 \end{aligned}$$

$$\begin{aligned}
 \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \tag{3}
 \end{aligned}$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \quad (4)$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \end{aligned} \quad (5)$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \\ &\vdots \\ F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \quad (6)$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \quad (7)$$

To find $y(x)$ series solution around $x = 0$. Hence

$$\begin{aligned}
 F_0 &= -y - xy' \\
 F_1 &= \frac{dF_0}{dx} \\
 &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \\
 &= x^2 y' + xy - 2y' \\
 F_2 &= \frac{dF_1}{dx} \\
 &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} y' + \frac{\partial F_1}{\partial y'} F_1 \\
 &= -y' x^3 - yx^2 + 5xy' + 3y \\
 F_3 &= \frac{dF_2}{dx} \\
 &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} y' + \frac{\partial F_2}{\partial y'} F_2 \\
 &= (x^4 - 9x^2 + 8) y' + yx(x^2 - 7) \\
 F_4 &= \frac{dF_3}{dx} \\
 &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} y' + \frac{\partial F_3}{\partial y'} F_3 \\
 &= (-x^5 + 14x^3 - 33x) y' - y(x^4 - 12x^2 + 15) \\
 F_5 &= \frac{dF_4}{dx} \\
 &= \frac{\partial F_4}{\partial x} + \frac{\partial F_4}{\partial y} y' + \frac{\partial F_4}{\partial y'} F_4 \\
 &= (x^6 - 20x^4 + 87x^2 - 48) y' + yx(x^4 - 18x^2 + 57) \\
 F_6 &= \frac{dF_5}{dx} \\
 &= \frac{\partial F_5}{\partial x} + \frac{\partial F_5}{\partial y} y' + \frac{\partial F_5}{\partial y'} F_5 \\
 &= (-x^7 + 27x^5 - 185x^3 + 279x) y' - y(x^2 - 7)(x^4 - 18x^2 + 15)
 \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and

$y'(0) = y'(0)$ gives

$$\begin{aligned} F_0 &= -y(0) \\ F_1 &= -2y'(0) \\ F_2 &= 3y(0) \\ F_3 &= 8y'(0) \\ F_4 &= -15y(0) \\ F_5 &= -48y'(0) \\ F_6 &= 105y(0) \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$y = \left(1 - \frac{1}{2}x^2 + \frac{1}{8}x^4 - \frac{1}{48}x^6 + \frac{1}{384}x^8\right) y(0) + \left(x - \frac{1}{3}x^3 + \frac{1}{15}x^5 - \frac{1}{105}x^7\right) y'(0) + O(x^8)$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard power series. Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$\begin{aligned} y' &= \sum_{n=1}^{\infty} n a_n x^{n-1} \\ y'' &= \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \end{aligned}$$

Substituting the above back into the ode gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = - \left(\sum_{n=0}^{\infty} a_n x^n \right) - x \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) \quad (1)$$

Which simplifies to

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \left(\sum_{n=1}^{\infty} n x^n a_n \right) + \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the

power and the corresponding index gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n \right) + \left(\sum_{n=1}^{\infty} n x^n a_n \right) + \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0 \quad (3)$$

$n = 0$ gives

$$2a_2 + a_0 = 0$$

$$a_2 = -\frac{a_0}{2}$$

For $1 \leq n$, the recurrence equation is

$$(n+2) a_{n+2} (n+1) + n a_n + a_n = 0 \quad (4)$$

Solving for a_{n+2} , gives

$$a_{n+2} = -\frac{a_n}{n+2} \quad (5)$$

For $n = 1$ the recurrence equation gives

$$6a_3 + 2a_1 = 0$$

Which after substituting the earlier terms found becomes

$$a_3 = -\frac{a_1}{3}$$

For $n = 2$ the recurrence equation gives

$$12a_4 + 3a_2 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = \frac{a_0}{8}$$

For $n = 3$ the recurrence equation gives

$$20a_5 + 4a_3 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = \frac{a_1}{15}$$

For $n = 4$ the recurrence equation gives

$$30a_6 + 5a_4 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = -\frac{a_0}{48}$$

For $n = 5$ the recurrence equation gives

$$42a_7 + 6a_5 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = -\frac{a_1}{105}$$

For $n = 6$ the recurrence equation gives

$$56a_8 + 7a_6 = 0$$

Which after substituting the earlier terms found becomes

$$a_8 = \frac{a_0}{384}$$

For $n = 7$ the recurrence equation gives

$$72a_9 + 8a_7 = 0$$

Which after substituting the earlier terms found becomes

$$a_9 = \frac{a_1}{945}$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + a_1 x - \frac{1}{2} a_0 x^2 - \frac{1}{3} a_1 x^3 + \frac{1}{8} a_0 x^4 + \frac{1}{15} a_1 x^5 - \frac{1}{48} a_0 x^6 - \frac{1}{105} a_1 x^7 + \dots$$

Collecting terms, the solution becomes

$$y = \left(1 - \frac{1}{2}x^2 + \frac{1}{8}x^4 - \frac{1}{48}x^6\right) a_0 + \left(x - \frac{1}{3}x^3 + \frac{1}{15}x^5 - \frac{1}{105}x^7\right) a_1 + O(x^8) \quad (3)$$

At $x = 0$ the solution above becomes

$$y = \left(1 - \frac{1}{2}x^2 + \frac{1}{8}x^4 - \frac{1}{48}x^6\right) c_1 + \left(x - \frac{1}{3}x^3 + \frac{1}{15}x^5 - \frac{1}{105}x^7\right) c_2 + O(x^8)$$

Summary

The solution(s) found are the following

$$\begin{aligned} y &= \left(1 - \frac{1}{2}x^2 + \frac{1}{8}x^4 - \frac{1}{48}x^6 + \frac{1}{384}x^8\right) y(0) \\ &\quad + \left(x - \frac{1}{3}x^3 + \frac{1}{15}x^5 - \frac{1}{105}x^7\right) y'(0) + O(x^8) \end{aligned} \quad (1)$$

$$y = \left(1 - \frac{1}{2}x^2 + \frac{1}{8}x^4 - \frac{1}{48}x^6\right) c_1 + \left(x - \frac{1}{3}x^3 + \frac{1}{15}x^5 - \frac{1}{105}x^7\right) c_2 + O(x^8) \quad (2)$$

Verification of solutions

$$y = \left(1 - \frac{1}{2}x^2 + \frac{1}{8}x^4 - \frac{1}{48}x^6 + \frac{1}{384}x^8\right) y(0) + \left(x - \frac{1}{3}x^3 + \frac{1}{15}x^5 - \frac{1}{105}x^7\right) y'(0) + O(x^8)$$

Verified OK.

$$y = \left(1 - \frac{1}{2}x^2 + \frac{1}{8}x^4 - \frac{1}{48}x^6\right) c_1 + \left(x - \frac{1}{3}x^3 + \frac{1}{15}x^5 - \frac{1}{105}x^7\right) c_2 + O(x^8)$$

Verified OK.

18.8.1 Maple step by step solution

Let's solve

$$y'' = -y - xy'$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + xy' + y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^k$$

- Rewrite DE with series expansions

- Convert $x \cdot y'$ to series expansion

$$x \cdot y' = \sum_{k=0}^{\infty} a_k k x^k$$

- Convert y'' to series expansion

$$y'' = \sum_{k=2}^{\infty} a_k k(k-1) x^{k-2}$$

- Shift index using $k- > k+2$

$$y'' = \sum_{k=0}^{\infty} a_{k+2}(k+2)(k+1) x^k$$

Rewrite DE with series expansions

$$\sum_{k=0}^{\infty} (a_{k+2}(k+2)(k+1) + a_k(k+1)) x^k = 0$$

- Each term in the series must be 0, giving the recursion relation

$$(k+1)(a_{k+2}(k+2) + a_k) = 0$$

- Recursion relation that defines the series solution to the ODE

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+2} = -\frac{a_k}{k+2} \right]$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
<- linear_1 successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 44

```
Order:=8;  
dsolve(diff(y(x),x$2)+x*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{2}x^2 + \frac{1}{8}x^4 - \frac{1}{48}x^6\right) y(0) + \left(x - \frac{1}{3}x^3 + \frac{1}{15}x^5 - \frac{1}{105}x^7\right) D(y)(0) + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 56

```
AsymptoticDSolveValue[y''[x]+x*y'[x]+y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_2 \left(-\frac{x^7}{105} + \frac{x^5}{15} - \frac{x^3}{3} + x \right) + c_1 \left(-\frac{x^6}{48} + \frac{x^4}{8} - \frac{x^2}{2} + 1 \right)$$

18.9 problem 4(a)

18.9.1 Existence and uniqueness analysis	3045
18.9.2 Maple step by step solution	3054

Internal problem ID [6435]

Internal file name [OUTPUT/5683_Sunday_June_05_2022_03_46_53_PM_90746626/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.3. Second-Order Linear Equations: Ordinary Points. Page 169

Problem number: 4(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_airy**", "**second order series method. Ordinary point**", "**second order series method. Taylor series method**"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' + y' - xy = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 0]$$

With the expansion point for the power series method at $x = 0$.

18.9.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = 1$$

$$q(x) = -x$$

$$F = 0$$

Hence the ode is

$$y'' + y' - xy = 0$$

The domain of $p(x) = 1$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = -x$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using Taylor series gives

$$\begin{aligned} y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \dots \\ &= y_0 + xy'_0 + \frac{x^2}{2}f|_{x_0, y_0, y'_0} + \frac{x^3}{3!}f'|_{x_0, y_0, y'_0} + \dots \\ &= y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \left. \frac{d^n f}{dx^n} \right|_{x_0, y_0, y'_0} \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \quad (1)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \quad (671)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \quad (672)$$

$$\begin{aligned} \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \end{aligned} \quad (2)$$

$$\begin{aligned} \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \end{aligned} \quad (3)$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \quad (4)$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \quad (5)$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \\ &\vdots \\ F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \quad (6)$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \quad (7)$$

To find $y(x)$ series solution around $x = 0$. Hence

$$\begin{aligned}
 F_0 &= xy - y' \\
 F_1 &= \frac{dF_0}{dx} \\
 &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \\
 &= xy' + y - xy + y' \\
 F_2 &= \frac{dF_1}{dx} \\
 &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} y' + \frac{\partial F_1}{\partial y'} F_1 \\
 &= (1 - 2x)y' + y(x^2 + x - 1) \\
 F_3 &= \frac{dF_2}{dx} \\
 &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} y' + \frac{\partial F_2}{\partial y'} F_2 \\
 &= (x^2 + 3x - 4)y' + (-2x^2 + 3x + 1)y \\
 F_4 &= \frac{dF_3}{dx} \\
 &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} y' + \frac{\partial F_3}{\partial y'} F_3 \\
 &= (-3x^2 + 2x + 8)y' + y(x^3 + 3x^2 - 8x + 3) \\
 F_5 &= \frac{dF_4}{dx} \\
 &= \frac{\partial F_4}{\partial x} + \frac{\partial F_4}{\partial y} y' + \frac{\partial F_4}{\partial y'} F_4 \\
 &= (x^3 + 6x^2 - 16x - 3)y' - 3\left(x^3 - \frac{5}{3}x^2 - \frac{14}{3}x + \frac{8}{3}\right)y \\
 F_6 &= \frac{dF_5}{dx} \\
 &= \frac{\partial F_5}{\partial x} + \frac{\partial F_5}{\partial y} y' + \frac{\partial F_5}{\partial y'} F_5 \\
 &= (-4x^3 + 2x^2 + 42x - 21)y' + y(x^4 + 6x^3 - 25x^2 + 7x + 14)
 \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = 1$ and

$y'(0) = 0$ gives

$$\begin{aligned} F_0 &= 0 \\ F_1 &= 1 \\ F_2 &= -1 \\ F_3 &= 1 \\ F_4 &= 3 \\ F_5 &= -8 \\ F_6 &= 14 \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$y = 1 + \frac{x^3}{6} - \frac{x^4}{24} + \frac{x^5}{120} + \frac{x^6}{240} - \frac{x^7}{630} + \frac{x^8}{2880} + O(x^8)$$

$$y = 1 + \frac{x^3}{6} - \frac{x^4}{24} + \frac{x^5}{120} + \frac{x^6}{240} - \frac{x^7}{630} + \frac{x^8}{2880} + O(x^8)$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard power series. Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$\begin{aligned} y' &= \sum_{n=1}^{\infty} n a_n x^{n-1} \\ y'' &= \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \end{aligned}$$

Substituting the above back into the ode gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = x \left(\sum_{n=0}^{\infty} a_n x^n \right) - \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) \quad (1)$$

Which simplifies to

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) + \sum_{n=0}^{\infty} (-x^{1+n} a_n) = 0 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\begin{aligned}\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} &= \sum_{n=0}^{\infty} (n+2) a_{n+2} (1+n) x^n \\ \sum_{n=1}^{\infty} n a_n x^{n-1} &= \sum_{n=0}^{\infty} (1+n) a_{1+n} x^n \\ \sum_{n=0}^{\infty} (-x^{1+n} a_n) &= \sum_{n=1}^{\infty} (-a_{n-1} x^n)\end{aligned}$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (1+n) x^n \right) + \left(\sum_{n=0}^{\infty} (1+n) a_{1+n} x^n \right) + \sum_{n=1}^{\infty} (-a_{n-1} x^n) = 0 \quad (3)$$

$n = 0$ gives

$$2a_2 + a_1 = 0$$

$$a_2 = -\frac{a_1}{2}$$

For $1 \leq n$, the recurrence equation is

$$(n+2) a_{n+2} (1+n) + (1+n) a_{1+n} - a_{n-1} = 0 \quad (4)$$

Solving for a_{n+2} , gives

$$\begin{aligned}a_{n+2} &= -\frac{n a_{1+n} + a_{1+n} - a_{n-1}}{(n+2)(1+n)} \\ (5) \quad &= -\frac{a_{1+n}}{n+2} + \frac{a_{n-1}}{(n+2)(1+n)}\end{aligned}$$

For $n = 1$ the recurrence equation gives

$$6a_3 + 2a_2 - a_0 = 0$$

Which after substituting the earlier terms found becomes

$$a_3 = \frac{a_1}{6} + \frac{a_0}{6}$$

For $n = 2$ the recurrence equation gives

$$12a_4 + 3a_3 - a_1 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = \frac{a_1}{24} - \frac{a_0}{24}$$

For $n = 3$ the recurrence equation gives

$$20a_5 + 4a_4 - a_2 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = -\frac{a_1}{30} + \frac{a_0}{120}$$

For $n = 4$ the recurrence equation gives

$$30a_6 + 5a_5 - a_3 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = \frac{a_1}{90} + \frac{a_0}{240}$$

For $n = 5$ the recurrence equation gives

$$42a_7 + 6a_6 - a_4 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = -\frac{a_1}{1680} - \frac{a_0}{630}$$

For $n = 6$ the recurrence equation gives

$$56a_8 + 7a_7 - a_5 = 0$$

Which after substituting the earlier terms found becomes

$$a_8 = -\frac{a_1}{1920} + \frac{a_0}{2880}$$

For $n = 7$ the recurrence equation gives

$$72a_9 + 8a_8 - a_6 = 0$$

Which after substituting the earlier terms found becomes

$$a_9 = \frac{11a_1}{51840} + \frac{a_0}{51840}$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$\begin{aligned} y &= a_0 + a_1 x - \frac{a_1 x^2}{2} + \left(\frac{a_1}{6} + \frac{a_0}{6}\right) x^3 + \left(\frac{a_1}{24} - \frac{a_0}{24}\right) x^4 \\ &\quad + \left(-\frac{a_1}{30} + \frac{a_0}{120}\right) x^5 + \left(\frac{a_1}{90} + \frac{a_0}{240}\right) x^6 + \left(-\frac{a_1}{1680} - \frac{a_0}{630}\right) x^7 + \dots \end{aligned}$$

Collecting terms, the solution becomes

$$\begin{aligned} y &= \left(1 + \frac{1}{6}x^3 - \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{240}x^6 - \frac{1}{630}x^7\right) a_0 \\ &\quad + \left(x - \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 - \frac{1}{30}x^5 + \frac{1}{90}x^6 - \frac{1}{1680}x^7\right) a_1 + O(x^8) \end{aligned} \tag{3}$$

At $x = 0$ the solution above becomes

$$\begin{aligned} y &= \left(1 + \frac{1}{6}x^3 - \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{240}x^6 - \frac{1}{630}x^7\right) c_1 \\ &\quad + \left(x - \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 - \frac{1}{30}x^5 + \frac{1}{90}x^6 - \frac{1}{1680}x^7\right) c_2 + O(x^8) \end{aligned}$$

$$y = 1 + \frac{x^3}{6} - \frac{x^4}{24} + \frac{x^5}{120} + \frac{x^6}{240} - \frac{x^7}{630} + O(x^8)$$

Summary

The solution(s) found are the following

$$y = 1 + \frac{x^3}{6} - \frac{x^4}{24} + \frac{x^5}{120} + \frac{x^6}{240} - \frac{x^7}{630} + \frac{x^8}{2880} + O(x^8) \quad (1)$$

$$y = 1 + \frac{x^3}{6} - \frac{x^4}{24} + \frac{x^5}{120} + \frac{x^6}{240} - \frac{x^7}{630} + O(x^8) \quad (2)$$

Verification of solutions

$$y = 1 + \frac{x^3}{6} - \frac{x^4}{24} + \frac{x^5}{120} + \frac{x^6}{240} - \frac{x^7}{630} + \frac{x^8}{2880} + O(x^8)$$

Verified OK.

$$y = 1 + \frac{x^3}{6} - \frac{x^4}{24} + \frac{x^5}{120} + \frac{x^6}{240} - \frac{x^7}{630} + O(x^8)$$

Verified OK.

18.9.2 Maple step by step solution

Let's solve

$$\left[y'' = xy - y', y(0) = 1, y'|_{\{x=0\}} = 0 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + y' - xy = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^k$$

- Rewrite ODE with series expansions

- Convert $x \cdot y$ to series expansion

$$x \cdot y = \sum_{k=0}^{\infty} a_k x^{k+1}$$

- Shift index using $k \rightarrow k - 1$

$$x \cdot y = \sum_{k=1}^{\infty} a_{k-1} x^k$$

- Convert y' to series expansion

$$y' = \sum_{k=1}^{\infty} a_k k x^{k-1}$$

- Shift index using $k- > k + 1$

$$y' = \sum_{k=0}^{\infty} a_{k+1} (k+1) x^k$$

- Convert y'' to series expansion

$$y'' = \sum_{k=2}^{\infty} a_k k(k-1) x^{k-2}$$

- Shift index using $k- > k + 2$

$$y'' = \sum_{k=0}^{\infty} a_{k+2} (k+2)(k+1) x^k$$

Rewrite ODE with series expansions

$$2a_2 + a_1 + \left(\sum_{k=1}^{\infty} (a_{k+2}(k+2)(k+1) + a_{k+1}(k+1) - a_{k-1}) x^k \right) = 0$$

- Each term must be 0

$$2a_2 + a_1 = 0$$

- Each term in the series must be 0, giving the recursion relation

$$(k^2 + 3k + 2) a_{k+2} + a_{k+1} k - a_{k-1} + a_{k+1} = 0$$

- Shift index using $k- > k + 1$

$$((k+1)^2 + 3k + 5) a_{k+3} + a_{k+2}(k+1) - a_k + a_{k+2} = 0$$

- Recursion relation that defines the series solution to the ODE

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+3} = -\frac{k a_{k+2} - a_k + 2 a_{k+2}}{k^2 + 5k + 6}, 2a_2 + a_1 = 0 \right]$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacic's algorithm  
<- No Liouvillian solutions exist  
-> Trying a solution in terms of special functions:  
  -> Bessel  
  <- Bessel successful  
<- special function solution successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
Order:=8;  
dsolve([diff(y(x),x$2)+diff(y(x),x)-x*y(x)=0,y(0) = 1, D(y)(0) = 0],y(x),type='series',x=0);
```

$$y(x) = 1 + \frac{1}{6}x^3 - \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{240}x^6 - \frac{1}{630}x^7 + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 40

```
AsymptoticDSolveValue[{y'[x]+y'[x]-x*y[x]==0,{y[0]==1,y'[0]==0}},y[x],{x,0,7}]
```

$$y(x) \rightarrow -\frac{x^7}{630} + \frac{x^6}{240} + \frac{x^5}{120} - \frac{x^4}{24} + \frac{x^3}{6} + 1$$

18.10 problem 4(b)

18.10.1 Existence and uniqueness analysis	3057
18.10.2 Maple step by step solution	3066

Internal problem ID [6436]

Internal file name [OUTPUT/5684_Sunday_June_05_2022_03_46_55_PM_55015192/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.3. Second-Order Linear Equations: Ordinary Points. Page 169

Problem number: 4(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_airy**", "**second order series method. Ordinary point**", "**second order series method. Taylor series method**"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' + y' - xy = 0$$

With initial conditions

$$[y(0) = 0, y'(0) = 1]$$

With the expansion point for the power series method at $x = 0$.

18.10.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = 1$$

$$q(x) = -x$$

$$F = 0$$

Hence the ode is

$$y'' + y' - xy = 0$$

The domain of $p(x) = 1$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = -x$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using Taylor series gives

$$\begin{aligned} y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \dots \\ &= y_0 + xy'_0 + \frac{x^2}{2}f|_{x_0, y_0, y'_0} + \frac{x^3}{3!}f'|_{x_0, y_0, y'_0} + \dots \\ &= y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \left. \frac{d^n f}{dx^n} \right|_{x_0, y_0, y'_0} \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \quad (1)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \quad (674)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \quad (675)$$

$$\begin{aligned} \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \end{aligned} \quad (2)$$

$$\begin{aligned} \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \end{aligned} \quad (3)$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \tag{4}$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \tag{5}$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \end{aligned}$$

⋮

$$\begin{aligned} F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \tag{6}$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \tag{7}$$

To find $y(x)$ series solution around $x = 0$. Hence

$$\begin{aligned}
 F_0 &= xy - y' \\
 F_1 &= \frac{dF_0}{dx} \\
 &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \\
 &= xy' + y - xy + y' \\
 F_2 &= \frac{dF_1}{dx} \\
 &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} y' + \frac{\partial F_1}{\partial y'} F_1 \\
 &= (1 - 2x)y' + y(x^2 + x - 1) \\
 F_3 &= \frac{dF_2}{dx} \\
 &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} y' + \frac{\partial F_2}{\partial y'} F_2 \\
 &= (x^2 + 3x - 4)y' + (-2x^2 + 3x + 1)y \\
 F_4 &= \frac{dF_3}{dx} \\
 &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} y' + \frac{\partial F_3}{\partial y'} F_3 \\
 &= (-3x^2 + 2x + 8)y' + y(x^3 + 3x^2 - 8x + 3) \\
 F_5 &= \frac{dF_4}{dx} \\
 &= \frac{\partial F_4}{\partial x} + \frac{\partial F_4}{\partial y} y' + \frac{\partial F_4}{\partial y'} F_4 \\
 &= (x^3 + 6x^2 - 16x - 3)y' - 3\left(x^3 - \frac{5}{3}x^2 - \frac{14}{3}x + \frac{8}{3}\right)y \\
 F_6 &= \frac{dF_5}{dx} \\
 &= \frac{\partial F_5}{\partial x} + \frac{\partial F_5}{\partial y} y' + \frac{\partial F_5}{\partial y'} F_5 \\
 &= (-4x^3 + 2x^2 + 42x - 21)y' + y(x^4 + 6x^3 - 25x^2 + 7x + 14)
 \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = 0$ and

$y'(0) = 1$ gives

$$F_0 = -1$$

$$F_1 = 1$$

$$F_2 = 1$$

$$F_3 = -4$$

$$F_4 = 8$$

$$F_5 = -3$$

$$F_6 = -21$$

Substituting all the above in (7) and simplifying gives the solution as

$$y = x - \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{30} + \frac{x^6}{90} - \frac{x^7}{1680} - \frac{x^8}{1920} + O(x^8)$$

$$y = x - \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{30} + \frac{x^6}{90} - \frac{x^7}{1680} - \frac{x^8}{1920} + O(x^8)$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard power series. Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$y' = \sum_{n=1}^{\infty} n a_n x^{n-1}$$

$$y'' = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}$$

Substituting the above back into the ode gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = x \left(\sum_{n=0}^{\infty} a_n x^n \right) - \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) \quad (1)$$

Which simplifies to

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) + \sum_{n=0}^{\infty} (-x^{1+n} a_n) = 0 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\begin{aligned}\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} &= \sum_{n=0}^{\infty} (n+2) a_{n+2} (1+n) x^n \\ \sum_{n=1}^{\infty} n a_n x^{n-1} &= \sum_{n=0}^{\infty} (1+n) a_{1+n} x^n \\ \sum_{n=0}^{\infty} (-x^{1+n} a_n) &= \sum_{n=1}^{\infty} (-a_{n-1} x^n)\end{aligned}$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (1+n) x^n \right) + \left(\sum_{n=0}^{\infty} (1+n) a_{1+n} x^n \right) + \sum_{n=1}^{\infty} (-a_{n-1} x^n) = 0 \quad (3)$$

$n = 0$ gives

$$2a_2 + a_1 = 0$$

$$a_2 = -\frac{a_1}{2}$$

For $1 \leq n$, the recurrence equation is

$$(n+2) a_{n+2} (1+n) + (1+n) a_{1+n} - a_{n-1} = 0 \quad (4)$$

Solving for a_{n+2} , gives

$$\begin{aligned}a_{n+2} &= -\frac{n a_{1+n} + a_{1+n} - a_{n-1}}{(n+2)(1+n)} \\ (5) \quad &= -\frac{a_{1+n}}{n+2} + \frac{a_{n-1}}{(n+2)(1+n)}\end{aligned}$$

For $n = 1$ the recurrence equation gives

$$6a_3 + 2a_2 - a_0 = 0$$

Which after substituting the earlier terms found becomes

$$a_3 = \frac{a_1}{6} + \frac{a_0}{6}$$

For $n = 2$ the recurrence equation gives

$$12a_4 + 3a_3 - a_1 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = \frac{a_1}{24} - \frac{a_0}{24}$$

For $n = 3$ the recurrence equation gives

$$20a_5 + 4a_4 - a_2 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = -\frac{a_1}{30} + \frac{a_0}{120}$$

For $n = 4$ the recurrence equation gives

$$30a_6 + 5a_5 - a_3 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = \frac{a_1}{90} + \frac{a_0}{240}$$

For $n = 5$ the recurrence equation gives

$$42a_7 + 6a_6 - a_4 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = -\frac{a_1}{1680} - \frac{a_0}{630}$$

For $n = 6$ the recurrence equation gives

$$56a_8 + 7a_7 - a_5 = 0$$

Which after substituting the earlier terms found becomes

$$a_8 = -\frac{a_1}{1920} + \frac{a_0}{2880}$$

For $n = 7$ the recurrence equation gives

$$72a_9 + 8a_8 - a_6 = 0$$

Which after substituting the earlier terms found becomes

$$a_9 = \frac{11a_1}{51840} + \frac{a_0}{51840}$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$\begin{aligned} y &= a_0 + a_1 x - \frac{a_1 x^2}{2} + \left(\frac{a_1}{6} + \frac{a_0}{6}\right) x^3 + \left(\frac{a_1}{24} - \frac{a_0}{24}\right) x^4 \\ &\quad + \left(-\frac{a_1}{30} + \frac{a_0}{120}\right) x^5 + \left(\frac{a_1}{90} + \frac{a_0}{240}\right) x^6 + \left(-\frac{a_1}{1680} - \frac{a_0}{630}\right) x^7 + \dots \end{aligned}$$

Collecting terms, the solution becomes

$$\begin{aligned} y &= \left(1 + \frac{1}{6}x^3 - \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{240}x^6 - \frac{1}{630}x^7\right) a_0 \\ &\quad + \left(x - \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 - \frac{1}{30}x^5 + \frac{1}{90}x^6 - \frac{1}{1680}x^7\right) a_1 + O(x^8) \end{aligned} \tag{3}$$

At $x = 0$ the solution above becomes

$$\begin{aligned} y &= \left(1 + \frac{1}{6}x^3 - \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{240}x^6 - \frac{1}{630}x^7\right) c_1 \\ &\quad + \left(x - \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 - \frac{1}{30}x^5 + \frac{1}{90}x^6 - \frac{1}{1680}x^7\right) c_2 + O(x^8) \end{aligned}$$

$$y = x - \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{30} + \frac{x^6}{90} - \frac{x^7}{1680} + O(x^8)$$

Summary

The solution(s) found are the following

$$y = x - \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{30} + \frac{x^6}{90} - \frac{x^7}{1680} - \frac{x^8}{1920} + O(x^8) \quad (1)$$

$$y = x - \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{30} + \frac{x^6}{90} - \frac{x^7}{1680} + O(x^8) \quad (2)$$

Verification of solutions

$$y = x - \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{30} + \frac{x^6}{90} - \frac{x^7}{1680} - \frac{x^8}{1920} + O(x^8)$$

Verified OK.

$$y = x - \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{30} + \frac{x^6}{90} - \frac{x^7}{1680} + O(x^8)$$

Verified OK.

18.10.2 Maple step by step solution

Let's solve

$$\left[y'' = xy - y', y(0) = 0, y'|_{\{x=0\}} = 1 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + y' - xy = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^k$$

- Rewrite ODE with series expansions

- Convert $x \cdot y$ to series expansion

$$x \cdot y = \sum_{k=0}^{\infty} a_k x^{k+1}$$

- Shift index using $k \rightarrow k - 1$

$$x \cdot y = \sum_{k=1}^{\infty} a_{k-1} x^k$$

- Convert y' to series expansion

$$y' = \sum_{k=1}^{\infty} a_k k x^{k-1}$$

- Shift index using $k- > k + 1$

$$y' = \sum_{k=0}^{\infty} a_{k+1} (k+1) x^k$$

- Convert y'' to series expansion

$$y'' = \sum_{k=2}^{\infty} a_k k(k-1) x^{k-2}$$

- Shift index using $k- > k + 2$

$$y'' = \sum_{k=0}^{\infty} a_{k+2} (k+2)(k+1) x^k$$

Rewrite ODE with series expansions

$$2a_2 + a_1 + \left(\sum_{k=1}^{\infty} (a_{k+2}(k+2)(k+1) + a_{k+1}(k+1) - a_{k-1}) x^k \right) = 0$$

- Each term must be 0

$$2a_2 + a_1 = 0$$

- Each term in the series must be 0, giving the recursion relation

$$(k^2 + 3k + 2) a_{k+2} + a_{k+1} k - a_{k-1} + a_{k+1} = 0$$

- Shift index using $k- > k + 1$

$$((k+1)^2 + 3k + 5) a_{k+3} + a_{k+2}(k+1) - a_k + a_{k+2} = 0$$

- Recursion relation that defines the series solution to the ODE

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+3} = -\frac{k a_{k+2} - a_k + 2 a_{k+2}}{k^2 + 5k + 6}, 2a_2 + a_1 = 0 \right]$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacic's algorithm  
<- No Liouvillian solutions exists  
-> Trying a solution in terms of special functions:  
  -> Bessel  
  <- Bessel successful  
<- special function solution successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 22

```
Order:=8;  
dsolve([diff(y(x),x$2)+diff(y(x),x)-x*y(x)=0,y(0) = 0, D(y)(0) = 1],y(x),type='series',x=0);
```

$$y(x) = x - \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 - \frac{1}{30}x^5 + \frac{1}{90}x^6 - \frac{1}{1680}x^7 + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 47

```
AsymptoticDSolveValue[{y'[x]+y'[x]-x*y[x]==0,{y[0]==0,y'[0]==1}},y[x],{x,0,7}]
```

$$y(x) \rightarrow -\frac{x^7}{1680} + \frac{x^6}{90} - \frac{x^5}{30} + \frac{x^4}{24} + \frac{x^3}{6} - \frac{x^2}{2} + x$$

18.11 problem 5

18.11.1 Maple step by step solution 3079

Internal problem ID [6437]

Internal file name [OUTPUT/5685_Sunday_June_05_2022_03_46_58_PM_37379447/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.3. Second-Order Linear Equations: Ordinary Points. Page 169

Problem number: 5.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second_order_bessel_ode", "second order series method. Ordinary point", "second order series method. Taylor series method"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' + \left(p + \frac{1}{2} - \frac{x^2}{4} \right) y = 0$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using

Taylor series gives

$$\begin{aligned}
 y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \dots \\
 &= y_0 + xy'_0 + \frac{x^2}{2}f|_{x_0, y_0, y'_0} + \frac{x^3}{3!}f'|_{x_0, y_0, y'_0} + \dots \\
 &= y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0}
 \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \tag{1}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \tag{677}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \tag{678}$$

$$\begin{aligned}
 \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \tag{2}
 \end{aligned}$$

$$\begin{aligned}
 \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \tag{3}
 \end{aligned}$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \quad (4)$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \quad (5)$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \\ &\vdots \\ F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \quad (6)$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \quad (7)$$

To find $y(x)$ series solution around $x = 0$. Hence

$$F_0 = -\frac{(-x^2 + 4p + 2)y}{4}$$

$$\begin{aligned} F_1 &= \frac{dF_0}{dx} \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \\ &= \frac{(x^2 - 4p - 2)y'}{4} + \frac{xy}{2} \end{aligned}$$

$$\begin{aligned} F_2 &= \frac{dF_1}{dx} \\ &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} y' + \frac{\partial F_1}{\partial y'} F_1 \\ &= xy' + \left(\frac{x^4}{16} + \left(-\frac{p}{2} - \frac{1}{4} \right) x^2 + p^2 + p + \frac{3}{4} \right) y \end{aligned}$$

$$\begin{aligned} F_3 &= \frac{dF_2}{dx} \\ &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} y' + \frac{\partial F_2}{\partial y'} F_2 \\ &= \frac{(x^4 + (-8p - 4)x^2 + 16p^2 + 16p + 28)y'}{16} - 2y \left(p + \frac{1}{2} - \frac{x^2}{4} \right) x \end{aligned}$$

$$\begin{aligned} F_4 &= \frac{dF_3}{dx} \\ &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} y' + \frac{\partial F_3}{\partial y'} F_3 \\ &= \frac{3(x^3 + 2(-2p - 1)x)y'}{4} - \left(-\frac{x^6}{64} + \frac{3(p + \frac{1}{2})x^4}{16} + \frac{3(-\frac{11}{4} - p^2 - p)x^2}{4} + \left(p + \frac{1}{2} \right) \left(p^2 + p + \frac{15}{4} \right) \right) y \end{aligned}$$

$$\begin{aligned} F_5 &= \frac{dF_4}{dx} \\ &= \frac{\partial F_4}{\partial x} + \frac{\partial F_4}{\partial y} y' + \frac{\partial F_4}{\partial y'} F_4 \\ &= \frac{(x^6 + (-12p - 6)x^4 + (48p^2 + 48p + 276)x^2 - 64p^3 - 96p^2 - 464p - 216)y'}{64} + \frac{9yx \left(\frac{x^4}{16} + \left(-\frac{p}{2} - \frac{1}{4} \right) x^2 + p^2 + p + \frac{3}{4} \right)}{2} \end{aligned}$$

$$\begin{aligned} F_6 &= \frac{dF_5}{dx} \\ &= \frac{\partial F_5}{\partial x} + \frac{\partial F_5}{\partial y} y' + \frac{\partial F_5}{\partial y'} F_5 \\ &= 6 \left(\frac{9}{4} + \frac{x^4}{16} + \frac{(-p - \frac{1}{2})x^2}{2} + p^2 + p \right) xy' + y \left(\frac{105}{16} + \frac{x^8}{256} + \frac{(-p - \frac{1}{2})x^6}{16} + \frac{3(p^2 + p + \frac{27}{4})x^4}{8} + \left(-\frac{11}{4} - p^2 - p \right) \right) \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and $y'(0) = y'(0)$ gives

$$\begin{aligned}
 F_0 &= -\frac{y(0)(2p+1)}{2} \\
 F_1 &= -y'(0)p - \frac{y'(0)}{2} \\
 F_2 &= y(0)p^2 + y(0)p + \frac{3y(0)}{4} \\
 F_3 &= y'(0)p^2 + y'(0)p + \frac{7y'(0)}{4} \\
 F_4 &= -y(0)p^3 - \frac{3y(0)p^2}{2} - \frac{17y(0)p}{4} - \frac{15y(0)}{8} \\
 F_5 &= -y'(0)p^3 - \frac{3y'(0)p^2}{2} - \frac{29y'(0)p}{4} - \frac{27y'(0)}{8} \\
 F_6 &= y(0)p^4 + 2y(0)p^3 + \frac{25y(0)p^2}{2} + \frac{23y(0)p}{2} + \frac{105y(0)}{16}
 \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$\begin{aligned}
 y &= \left(1 - \frac{1}{2}x^2p + \frac{1}{24}x^4p^2 + \frac{1}{24}x^4p - \frac{1}{720}x^6p^3 - \frac{1}{480}x^6p^2 - \frac{17}{2880}x^6p + \frac{1}{40320}x^8p^4 \right. \\
 &\quad \left. + \frac{1}{20160}x^8p^3 + \frac{5}{16128}x^8p^2 + \frac{23}{80640}x^8p - \frac{1}{4}x^2 + \frac{1}{32}x^4 - \frac{1}{384}x^6 + \frac{1}{6144}x^8 \right) y(0) \\
 &\quad + \left(-\frac{1}{6}x^3p + \frac{1}{120}x^5p^2 + \frac{1}{120}x^5p - \frac{1}{5040}x^7p^3 - \frac{1}{3360}x^7p^2 - \frac{29}{20160}x^7p - \frac{1}{12}x^3 \right. \\
 &\quad \left. + \frac{7}{480}x^5 - \frac{3}{4480}x^7 + x \right) y'(0) + O(x^8)
 \end{aligned}$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard power series. Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$\begin{aligned}
 y' &= \sum_{n=1}^{\infty} n a_n x^{n-1} \\
 y'' &= \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}
 \end{aligned}$$

Substituting the above back into the ode gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = -\frac{(-x^2 + 4p + 2) \left(\sum_{n=0}^{\infty} a_n x^n \right)}{4} \quad (1)$$

Which simplifies to

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \sum_{n=0}^{\infty} \left(-\frac{x^{n+2} a_n}{4} \right) + \left(\sum_{n=0}^{\infty} \frac{a_n x^n (2p+1)}{2} \right) = 0 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} &= \sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n \\ \sum_{n=0}^{\infty} \left(-\frac{x^{n+2} a_n}{4} \right) &= \sum_{n=2}^{\infty} \left(-\frac{a_{n-2} x^n}{4} \right) \end{aligned}$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n \right) + \sum_{n=2}^{\infty} \left(-\frac{a_{n-2} x^n}{4} \right) + \left(\sum_{n=0}^{\infty} \frac{a_n x^n (2p+1)}{2} \right) = 0 \quad (3)$$

$n = 0$ gives

$$2a_2 + \frac{a_0(2p+1)}{2} = 0$$

$$a_2 = -\frac{1}{2}a_0 p - \frac{1}{4}a_0$$

$n = 1$ gives

$$6a_3 + \frac{a_1(2p+1)}{2} = 0$$

Which after substituting earlier equations, simplifies to

$$a_3 = -\frac{1}{6}a_1p - \frac{1}{12}a_1$$

For $2 \leq n$, the recurrence equation is

$$(n+2)a_{n+2}(n+1) - \frac{a_{n-2}}{4} + \frac{a_n(2p+1)}{2} = 0 \quad (4)$$

Solving for a_{n+2} , gives

$$\begin{aligned} a_{n+2} &= -\frac{4pa_n + 2a_n - a_{n-2}}{4(n+2)(n+1)} \\ (5) \quad &= -\frac{(4p+2)a_n}{4(n+2)(n+1)} + \frac{a_{n-2}}{4(n+2)(n+1)} \end{aligned}$$

For $n = 2$ the recurrence equation gives

$$12a_4 - \frac{a_0}{4} + \frac{a_2(2p+1)}{2} = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = \frac{1}{32}a_0 + \frac{1}{24}a_0p^2 + \frac{1}{24}a_0p$$

For $n = 3$ the recurrence equation gives

$$20a_5 - \frac{a_1}{4} + \frac{a_3(2p+1)}{2} = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = \frac{7}{480}a_1 + \frac{1}{120}a_1p^2 + \frac{1}{120}a_1p$$

For $n = 4$ the recurrence equation gives

$$30a_6 - \frac{a_2}{4} + \frac{a_4(2p+1)}{2} = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = -\frac{17}{2880}a_0p - \frac{1}{384}a_0 - \frac{1}{720}a_0p^3 - \frac{1}{480}a_0p^2$$

For $n = 5$ the recurrence equation gives

$$42a_7 - \frac{a_3}{4} + \frac{a_5(2p+1)}{2} = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = -\frac{29}{20160}a_1p - \frac{3}{4480}a_1 - \frac{1}{5040}a_1p^3 - \frac{1}{3360}a_1p^2$$

For $n = 6$ the recurrence equation gives

$$56a_8 - \frac{a_4}{4} + \frac{a_6(2p+1)}{2} = 0$$

Which after substituting the earlier terms found becomes

$$a_8 = \frac{1}{6144}a_0 + \frac{5}{16128}a_0p^2 + \frac{23}{80640}a_0p + \frac{1}{40320}a_0p^4 + \frac{1}{20160}a_0p^3$$

For $n = 7$ the recurrence equation gives

$$72a_9 - \frac{a_5}{4} + \frac{a_7(2p+1)}{2} = 0$$

Which after substituting the earlier terms found becomes

$$a_9 = \frac{107}{1935360}a_1 + \frac{37}{725760}a_1p^2 + \frac{1}{20736}a_1p + \frac{1}{362880}a_1p^4 + \frac{1}{181440}a_1p^3$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$\begin{aligned}
 y = & a_0 + a_1x + \left(-\frac{1}{2}a_0p - \frac{1}{4}a_0\right)x^2 + \left(-\frac{1}{6}a_1p - \frac{1}{12}a_1\right)x^3 + \left(\frac{1}{32}a_0 + \frac{1}{24}a_0p^2 + \frac{1}{24}a_0p\right)x^4 \\
 & + \left(\frac{7}{480}a_1 + \frac{1}{120}a_1p^2 + \frac{1}{120}a_1p\right)x^5 + \left(-\frac{17}{2880}a_0p - \frac{1}{384}a_0 - \frac{1}{720}a_0p^3 - \frac{1}{480}a_0p^2\right)x^6 \\
 & + \left(-\frac{29}{20160}a_1p - \frac{3}{4480}a_1 - \frac{1}{5040}a_1p^3 - \frac{1}{3360}a_1p^2\right)x^7 + \dots
 \end{aligned}$$

Collecting terms, the solution becomes

$$\begin{aligned}
 y = & \left(1 + \left(-\frac{p}{2} - \frac{1}{4}\right)x^2 + \left(\frac{1}{32} + \frac{1}{24}p^2 + \frac{1}{24}p\right)x^4\right. \\
 & \left. + \left(-\frac{17}{2880}p - \frac{1}{384} - \frac{1}{720}p^3 - \frac{1}{480}p^2\right)x^6\right)a_0 \\
 & + \left(x + \left(-\frac{p}{6} - \frac{1}{12}\right)x^3 + \left(\frac{7}{480} + \frac{1}{120}p^2 + \frac{1}{120}p\right)x^5\right. \\
 & \left. + \left(-\frac{29}{20160}p - \frac{3}{4480} - \frac{1}{5040}p^3 - \frac{1}{3360}p^2\right)x^7\right)a_1 + O(x^8)
 \end{aligned} \tag{3}$$

At $x = 0$ the solution above becomes

$$\begin{aligned}
 y = & \left(1 + \left(-\frac{p}{2} - \frac{1}{4}\right)x^2 + \left(\frac{1}{32} + \frac{1}{24}p^2 + \frac{1}{24}p\right)x^4\right. \\
 & \left. + \left(-\frac{17}{2880}p - \frac{1}{384} - \frac{1}{720}p^3 - \frac{1}{480}p^2\right)x^6\right)c_1 \\
 & + \left(x + \left(-\frac{p}{6} - \frac{1}{12}\right)x^3 + \left(\frac{7}{480} + \frac{1}{120}p^2 + \frac{1}{120}p\right)x^5\right. \\
 & \left. + \left(-\frac{29}{20160}p - \frac{3}{4480} - \frac{1}{5040}p^3 - \frac{1}{3360}p^2\right)x^7\right)c_2 + O(x^8)
 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \left(1 - \frac{1}{2}x^2p + \frac{1}{24}x^4p^2 + \frac{1}{24}x^4p - \frac{1}{720}x^6p^3 - \frac{1}{480}x^6p^2 - \frac{17}{2880}x^6p + \frac{1}{40320}x^8p^4 \right. \\ \left. + \frac{1}{20160}x^8p^3 + \frac{5}{16128}x^8p^2 + \frac{23}{80640}x^8p - \frac{1}{4}x^2 + \frac{1}{32}x^4 - \frac{1}{384}x^6 + \frac{1}{6144}x^8 \right) y(0)_{(1)} \\ + \left(-\frac{1}{6}x^3p + \frac{1}{120}x^5p^2 + \frac{1}{120}x^5p - \frac{1}{5040}x^7p^3 - \frac{1}{3360}x^7p^2 - \frac{29}{20160}x^7p - \frac{1}{12}x^3 \right. \\ \left. + \frac{7}{480}x^5 - \frac{3}{4480}x^7 + x \right) y'(0) + O(x^8)$$

$$y = \left(1 + \left(-\frac{p}{2} - \frac{1}{4} \right) x^2 + \left(\frac{1}{32} + \frac{1}{24}p^2 + \frac{1}{24}p \right) x^4 \right. \\ \left. + \left(-\frac{17}{2880}p - \frac{1}{384} - \frac{1}{720}p^3 - \frac{1}{480}p^2 \right) x^6 \right) c_1 \\ + \left(x + \left(-\frac{p}{6} - \frac{1}{12} \right) x^3 + \left(\frac{7}{480} + \frac{1}{120}p^2 + \frac{1}{120}p \right) x^5 \right. \\ \left. + \left(-\frac{29}{20160}p - \frac{3}{4480} - \frac{1}{5040}p^3 - \frac{1}{3360}p^2 \right) x^7 \right) c_2 + O(x^8)$$

Verification of solutions

$$y = \left(1 - \frac{1}{2}x^2p + \frac{1}{24}x^4p^2 + \frac{1}{24}x^4p - \frac{1}{720}x^6p^3 - \frac{1}{480}x^6p^2 - \frac{17}{2880}x^6p + \frac{1}{40320}x^8p^4 \right. \\ \left. + \frac{1}{20160}x^8p^3 + \frac{5}{16128}x^8p^2 + \frac{23}{80640}x^8p - \frac{1}{4}x^2 + \frac{1}{32}x^4 - \frac{1}{384}x^6 + \frac{1}{6144}x^8 \right) y(0) \\ + \left(-\frac{1}{6}x^3p + \frac{1}{120}x^5p^2 + \frac{1}{120}x^5p - \frac{1}{5040}x^7p^3 - \frac{1}{3360}x^7p^2 - \frac{29}{20160}x^7p - \frac{1}{12}x^3 \right. \\ \left. + \frac{7}{480}x^5 - \frac{3}{4480}x^7 + x \right) y'(0) + O(x^8)$$

Verified OK.

$$y = \left(1 + \left(-\frac{p}{2} - \frac{1}{4} \right) x^2 + \left(\frac{1}{32} + \frac{1}{24}p^2 + \frac{1}{24}p \right) x^4 \right. \\ \left. + \left(-\frac{17}{2880}p - \frac{1}{384} - \frac{1}{720}p^3 - \frac{1}{480}p^2 \right) x^6 \right) c_1 \\ + \left(x + \left(-\frac{p}{6} - \frac{1}{12} \right) x^3 + \left(\frac{7}{480} + \frac{1}{120}p^2 + \frac{1}{120}p \right) x^5 \right. \\ \left. + \left(-\frac{29}{20160}p - \frac{3}{4480} - \frac{1}{5040}p^3 - \frac{1}{3360}p^2 \right) x^7 \right) c_2 + O(x^8)$$

Verified OK.

18.11.1 Maple step by step solution

Let's solve

$$y'' = -\frac{(-x^2+4p+2)y}{4}$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = \left(-p - \frac{1}{2} + \frac{x^2}{4} \right) y$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \left(p + \frac{1}{2} - \frac{x^2}{4} \right) y = 0$$

- Multiply by denominators

$$4y'' + (-x^2 + 4p + 2)y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^k$$

□ Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 0..2$

$$x^m \cdot y = \sum_{k=\max(0,-m)}^{\infty} a_k x^{k+m}$$

- Shift index using $k \rightarrow k - m$

$$x^m \cdot y = \sum_{k=\max(0,-m)+m}^{\infty} a_{k-m} x^k$$

- Convert y'' to series expansion

$$y'' = \sum_{k=2}^{\infty} a_k k(k-1) x^{k-2}$$

- Shift index using $k \rightarrow k + 2$

$$y'' = \sum_{k=0}^{\infty} a_{k+2} (k+2)(k+1) x^k$$

Rewrite ODE with series expansions

$$8a_2 + 2a_0(2p+1) + (24a_3 + 2a_1(2p+1))x + \left(\sum_{k=2}^{\infty} (4a_{k+2}(k+2)(k+1) + 2a_k(2p+1) - a_{k-2}) \right) x^k = 0$$

- The coefficients of each power of x must be 0

$$[8a_2 + 2a_0(2p+1) = 0, 24a_3 + 2a_1(2p+1) = 0]$$

- Solve for the dependent coefficient(s)

$$\{a_2 = -\frac{1}{2}a_0p - \frac{1}{4}a_0, a_3 = -\frac{1}{6}a_1p - \frac{1}{12}a_1\}$$

- Each term in the series must be 0, giving the recursion relation

$$4(k^2 + 3k + 2)a_{k+2} + 2a_k(2p+1) - a_{k-2} = 0$$

- Shift index using $k \rightarrow k + 2$

$$4((k+2)^2 + 3k + 8)a_{k+4} + 2a_{k+2}(2p+1) - a_k = 0$$

- Recursion relation that defines the series solution to the ODE

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+4} = -\frac{4pa_{k+2} - a_k + 2a_{k+2}}{4(k^2 + 7k + 12)}, a_2 = -\frac{1}{2}a_0p - \frac{1}{4}a_0, a_3 = -\frac{1}{6}a_1p - \frac{1}{12}a_1 \right]$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
  -> Bessel
  -> elliptic
  -> Legendre
  -> Whittaker
      -> hyper3: Equivalence to 1F1 under a power @ Moebius
      <- hyper3 successful: received ODE is equivalent to the 1F1 ODE
      <- Whittaker successful
<- special function solution successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 120

```
Order:=8;
dsolve(diff(y(x),x$2)+(p+1/2-x^2/4)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{(1+2p)x^2}{4} + \frac{(4p^2+4p+3)x^4}{96} - \frac{(8p^3+12p^2+34p+15)x^6}{5760} \right) y(0) \\ + \left(x - \frac{(1+2p)x^3}{12} + \frac{(4p^2+4p+7)x^5}{480} - \frac{(8p^3+12p^2+58p+27)x^7}{40320} \right) D(y)(0) \\ + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 142

```
AsymptoticDSolveValue[y''[x]+(p+1/2-x^2/4)*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_2 \left(\frac{(-4p-2)(4p+2)^2 x^7}{322560} + \frac{13(-4p-2)x^7}{40320} + \frac{(4p+2)^2 x^5}{1920} + \frac{1}{24}(-4p-2)x^3 + \frac{x^5}{80} + x \right) + c_1 \left(\frac{(-4p-2)(4p+2)^2 x^6}{46080} + \frac{7(-4p-2)x^6}{5760} + \frac{1}{384}(4p+2)^2 x^4 + \frac{1}{8}(-4p-2)x^2 + \frac{x^4}{48} + 1 \right)$$

18.12 problem 6

18.12.1 Maple step by step solution 3091

Internal problem ID [6438]

Internal file name [OUTPUT/5686_Sunday_June_05_2022_03_46_59_PM_23497955/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.3. Second-Order Linear Equations: Ordinary Points. Page 169

Problem number: 6.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_airy", "second_order_bessel_ode", "second order series method. Ordinary point", "second order series method. Taylor series method"

Maple gives the following as the ode type

```
[[_Emden, _Fowler]]
```

$$y'' + xy = 0$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using

Taylor series gives

$$\begin{aligned}
 y(x) &= y(x_0) + (x - x_0) y'(x_0) + \frac{(x - x_0)^2}{2} y''(x_0) + \frac{(x - x_0)^3}{3!} y'''(x_0) + \dots \\
 &= y_0 + x y'_0 + \frac{x^2}{2} f|_{x_0, y_0, y'_0} + \frac{x^3}{3!} f'|_{x_0, y_0, y'_0} + \dots \\
 &= y_0 + x y'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \left. \frac{d^n f}{dx^n} \right|_{x_0, y_0, y'_0}
 \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \tag{1}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \tag{680}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \tag{681}$$

$$\begin{aligned}
 \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \tag{2}
 \end{aligned}$$

$$\begin{aligned}
 \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \tag{3}
 \end{aligned}$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \tag{4}$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \tag{5}$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \end{aligned}$$

⋮

$$\begin{aligned} F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \tag{6}$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \tag{7}$$

To find $y(x)$ series solution around $x = 0$. Hence

$$\begin{aligned}
 F_0 &= -xy \\
 F_1 &= \frac{dF_0}{dx} \\
 &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \\
 &= -y - xy' \\
 F_2 &= \frac{dF_1}{dx} \\
 &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y}y' + \frac{\partial F_1}{\partial y'}F_1 \\
 &= -2y' + yx^2 \\
 F_3 &= \frac{dF_2}{dx} \\
 &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y}y' + \frac{\partial F_2}{\partial y'}F_2 \\
 &= x(xy' + 4y) \\
 F_4 &= \frac{dF_3}{dx} \\
 &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y}y' + \frac{\partial F_3}{\partial y'}F_3 \\
 &= -yx^3 + 6xy' + 4y \\
 F_5 &= \frac{dF_4}{dx} \\
 &= \frac{\partial F_4}{\partial x} + \frac{\partial F_4}{\partial y}y' + \frac{\partial F_4}{\partial y'}F_4 \\
 &= -9yx^2 - y'x^3 + 10y' \\
 F_6 &= \frac{dF_5}{dx} \\
 &= \frac{\partial F_5}{\partial x} + \frac{\partial F_5}{\partial y}y' + \frac{\partial F_5}{\partial y'}F_5 \\
 &= -12x^2y' + xy(x^3 - 28)
 \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and

$y'(0) = y'(0)$ gives

$$\begin{aligned}F_0 &= 0 \\F_1 &= -y(0) \\F_2 &= -2y'(0) \\F_3 &= 0 \\F_4 &= 4y(0) \\F_5 &= 10y'(0) \\F_6 &= 0\end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$y = \left(1 - \frac{1}{6}x^3 + \frac{1}{180}x^6\right)y(0) + \left(x - \frac{1}{12}x^4 + \frac{1}{504}x^7\right)y'(0) + O(x^8)$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard power series. Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$\begin{aligned}y' &= \sum_{n=1}^{\infty} n a_n x^{n-1} \\y'' &= \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}\end{aligned}$$

Substituting the above back into the ode gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = -x \left(\sum_{n=0}^{\infty} a_n x^n \right) \quad (1)$$

Which simplifies to

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \left(\sum_{n=0}^{\infty} x^{1+n} a_n \right) = 0 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the

power and the corresponding index gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2) a_{n+2} (1+n) x^n$$

$$\sum_{n=0}^{\infty} x^{1+n} a_n = \sum_{n=1}^{\infty} a_{n-1} x^n$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (1+n) x^n \right) + \left(\sum_{n=1}^{\infty} a_{n-1} x^n \right) = 0 \quad (3)$$

For $1 \leq n$, the recurrence equation is

$$(n+2) a_{n+2} (1+n) + a_{n-1} = 0 \quad (4)$$

Solving for a_{n+2} , gives

$$a_{n+2} = -\frac{a_{n-1}}{(n+2)(1+n)} \quad (5)$$

For $n = 1$ the recurrence equation gives

$$6a_3 + a_0 = 0$$

Which after substituting the earlier terms found becomes

$$a_3 = -\frac{a_0}{6}$$

For $n = 2$ the recurrence equation gives

$$12a_4 + a_1 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = -\frac{a_1}{12}$$

For $n = 3$ the recurrence equation gives

$$20a_5 + a_2 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = 0$$

For $n = 4$ the recurrence equation gives

$$30a_6 + a_3 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = \frac{a_0}{180}$$

For $n = 5$ the recurrence equation gives

$$42a_7 + a_4 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = \frac{a_1}{504}$$

For $n = 6$ the recurrence equation gives

$$56a_8 + a_5 = 0$$

Which after substituting the earlier terms found becomes

$$a_8 = 0$$

For $n = 7$ the recurrence equation gives

$$72a_9 + a_6 = 0$$

Which after substituting the earlier terms found becomes

$$a_9 = -\frac{a_0}{12960}$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + a_1 x - \frac{1}{6} a_0 x^3 - \frac{1}{12} a_1 x^4 + \frac{1}{180} a_0 x^6 + \frac{1}{504} a_1 x^7 + \dots$$

Collecting terms, the solution becomes

$$y = \left(1 - \frac{1}{6}x^3 + \frac{1}{180}x^6\right) a_0 + \left(x - \frac{1}{12}x^4 + \frac{1}{504}x^7\right) a_1 + O(x^8) \quad (3)$$

At $x = 0$ the solution above becomes

$$y = \left(1 - \frac{1}{6}x^3 + \frac{1}{180}x^6\right) c_1 + \left(x - \frac{1}{12}x^4 + \frac{1}{504}x^7\right) c_2 + O(x^8)$$

Summary

The solution(s) found are the following

$$y = \left(1 - \frac{1}{6}x^3 + \frac{1}{180}x^6\right) y(0) + \left(x - \frac{1}{12}x^4 + \frac{1}{504}x^7\right) y'(0) + O(x^8) \quad (1)$$

$$y = \left(1 - \frac{1}{6}x^3 + \frac{1}{180}x^6\right) c_1 + \left(x - \frac{1}{12}x^4 + \frac{1}{504}x^7\right) c_2 + O(x^8) \quad (2)$$

Verification of solutions

$$y = \left(1 - \frac{1}{6}x^3 + \frac{1}{180}x^6\right) y(0) + \left(x - \frac{1}{12}x^4 + \frac{1}{504}x^7\right) y'(0) + O(x^8)$$

Verified OK.

$$y = \left(1 - \frac{1}{6}x^3 + \frac{1}{180}x^6\right) c_1 + \left(x - \frac{1}{12}x^4 + \frac{1}{504}x^7\right) c_2 + O(x^8)$$

Verified OK.

18.12.1 Maple step by step solution

Let's solve

$$y'' = -xy$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + xy = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^k$$

- Rewrite ODE with series expansions

- Convert $x \cdot y$ to series expansion

$$x \cdot y = \sum_{k=0}^{\infty} a_k x^{k+1}$$

- Shift index using $k \rightarrow k - 1$

$$x \cdot y = \sum_{k=1}^{\infty} a_{k-1} x^k$$

- Convert y'' to series expansion

$$y'' = \sum_{k=2}^{\infty} a_k k(k-1) x^{k-2}$$

- Shift index using $k \rightarrow k + 2$

$$y'' = \sum_{k=0}^{\infty} a_{k+2} (k+2)(k+1) x^k$$

Rewrite ODE with series expansions

$$2a_2 + \left(\sum_{k=1}^{\infty} (a_{k+2} (k+2)(k+1) + a_{k-1}) x^k \right) = 0$$

- Each term must be 0

$$2a_2 = 0$$

- Each term in the series must be 0, giving the recursion relation

$$(k^2 + 3k + 2) a_{k+2} + a_{k-1} = 0$$

- Shift index using $k \rightarrow k + 1$

$$((k+1)^2 + 3k + 5) a_{k+3} + a_k = 0$$

- Recursion relation that defines the series solution to the ODE

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+3} = -\frac{a_k}{k^2+5k+6}, 2a_2 = 0 \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
    <- Bessel successful
<- special function solution successful`

```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 34

```

Order:=8;
dsolve(diff(y(x),x$2)+x*y(x)=0,y(x),type='series',x=0);

```

$$y(x) = \left(1 - \frac{1}{6}x^3 + \frac{1}{180}x^6\right) y(0) + \left(x - \frac{1}{12}x^4 + \frac{1}{504}x^7\right) D(y)(0) + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 42

```

AsymptoticDSolveValue[y'[x]+x*y[x]==0,y[x],{x,0,7}]

```

$$y(x) \rightarrow c_2 \left(\frac{x^7}{504} - \frac{x^4}{12} + x \right) + c_1 \left(\frac{x^6}{180} - \frac{x^3}{6} + 1 \right)$$

18.13 problem 7

18.13.1 Maple step by step solution 3102

Internal problem ID [6439]

Internal file name [OUTPUT/5687_Sunday_June_05_2022_03_47_01_PM_53322889/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.3. Second-Order Linear Equations: Ordinary Points. Page 169

Problem number: 7.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second order series method. Ordinary point", "second_order_change_of_variable_on_x_method_1", "second_order_change_of_variable_on_x_method_2", "second order series method. Taylor series method"

Maple gives the following as the ode type

```
[_Gegenbauer , [_2nd_order , _linear , ` _with_symmetry_[0,F(x)]`]]
```

$$(-x^2 + 1)y'' - xy' + p^2y = 0$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using

Taylor series gives

$$\begin{aligned}
 y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \dots \\
 &= y_0 + xy'_0 + \frac{x^2}{2}f|_{x_0, y_0, y'_0} + \frac{x^3}{3!}f'|_{x_0, y_0, y'_0} + \dots \\
 &= y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0}
 \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \tag{1}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \tag{683}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \tag{684}$$

$$\begin{aligned}
 \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \tag{2}
 \end{aligned}$$

$$\begin{aligned}
 \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \tag{3}
 \end{aligned}$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \tag{4}$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \tag{5}$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \end{aligned}$$

⋮

$$\begin{aligned} F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \tag{6}$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \tag{7}$$

To find $y(x)$ series solution around $x = 0$. Hence

$$F_0 = \frac{p^2 y - xy'}{x^2 - 1}$$

$$\begin{aligned} F_1 &= \frac{dF_0}{dx} \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \\ &= \frac{((p^2 + 2)x^2 - p^2 + 1)y' - 3yp^2 x}{(x^2 - 1)^2} \end{aligned}$$

$$\begin{aligned} F_2 &= \frac{dF_1}{dx} \\ &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} y' + \frac{\partial F_1}{\partial y'} F_1 \\ &= \frac{(-6p^2 x^3 + 6x p^2 - 6x^3 - 9x)y' + y((p^2 + 11)x^2 - p^2 + 4)p^2}{(x^2 - 1)^3} \end{aligned}$$

$$\begin{aligned} F_3 &= \frac{dF_2}{dx} \\ &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} y' + \frac{\partial F_2}{\partial y'} F_2 \\ &= \frac{((p^4 + 35p^2 + 24)x^4 + (-2p^4 - 25p^2 + 72)x^2 + p^4 - 10p^2 + 9)y' - 10yx((p^2 + 5)x^2 - p^2 + \frac{11}{2})p^2}{(x^2 - 1)^4} \end{aligned}$$

$$\begin{aligned} F_4 &= \frac{dF_3}{dx} \\ &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} y' + \frac{\partial F_3}{\partial y'} F_3 \\ &= \frac{(1+x)(-15((p^4 + 15p^2 + 8)x^4 + (-2p^4 - 2p^2 + 40)x^2 + p^4 - 13p^2 + 15)xy' + y((p^4 + 85p^2 + 274) \dots)}{(x^2 - 1)^6} \end{aligned}$$

$$\begin{aligned} F_5 &= \frac{dF_4}{dx} \\ &= \frac{\partial F_4}{\partial x} + \frac{\partial F_4}{\partial y} y' + \frac{\partial F_4}{\partial y'} F_4 \\ &= \frac{(1+x)((p^6 + 175p^4 + 1624p^2 + 720)x^6 + (-3p^6 - 315p^4 + 1218p^2 + 5400)x^4 + (3p^6 + 105p^4 - 25 \dots)}{\dots} \end{aligned}$$

$$\begin{aligned} F_6 &= \frac{dF_5}{dx} \\ &= \frac{\partial F_5}{\partial x} + \frac{\partial F_5}{\partial y} y' + \frac{\partial F_5}{\partial y'} F_5 \\ &= \frac{\left(-28\left(\frac{1575}{4} + (p^6 + 70p^4 + 469p^2 + 180)x^6 + 3(630 - p^6 - \frac{65}{2}p^4 + \frac{607}{2}p^2)x^4 + 3(p^6 - 5p^4 - \frac{667}{2}p^2 + \dots\right)}{\dots} \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and $y'(0) = y'(0)$ gives

$$\begin{aligned} F_0 &= -y(0) p^2 \\ F_1 &= -y'(0) p^2 + y'(0) \\ F_2 &= y(0) p^4 - 4y(0) p^2 \\ F_3 &= y'(0) p^4 - 10y'(0) p^2 + 9y'(0) \\ F_4 &= -y(0) p^6 + 20y(0) p^4 - 64y(0) p^2 \\ F_5 &= -y'(0) p^6 + 35y'(0) p^4 - 259y'(0) p^2 + 225y'(0) \\ F_6 &= y(0) p^8 - 56y(0) p^6 + 784y(0) p^4 - 2304y(0) p^2 \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$\begin{aligned} y = & \left(1 - \frac{1}{2}x^2p^2 + \frac{1}{24}p^4x^4 - \frac{1}{6}x^4p^2 - \frac{1}{720}p^6x^6 + \frac{1}{36}p^4x^6 - \frac{4}{45}x^6p^2 + \frac{1}{40320}x^8p^8 \right. \\ & \left. - \frac{1}{720}x^8p^6 + \frac{7}{360}x^8p^4 - \frac{2}{35}x^8p^2 \right) y(0) + \left(x - \frac{1}{6}p^2x^3 + \frac{1}{6}x^3 + \frac{1}{120}p^4x^5 - \frac{1}{12}x^5p^2 \right. \\ & \left. + \frac{3}{40}x^5 - \frac{1}{5040}x^7p^6 + \frac{1}{144}x^7p^4 - \frac{37}{720}x^7p^2 + \frac{5}{112}x^7 \right) y'(0) + O(x^8) \end{aligned}$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard power series The ode is normalized to be

$$(-x^2 + 1) y'' - xy' + p^2y = 0$$

Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$\begin{aligned} y' &= \sum_{n=1}^{\infty} n a_n x^{n-1} \\ y'' &= \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \end{aligned}$$

Substituting the above back into the ode gives

$$(-x^2 + 1) \left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) - x \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) + p^2 \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0 \quad (1)$$

Which simplifies to

$$\sum_{n=2}^{\infty} (-x^n a_n n(n-1)) + \left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \sum_{n=1}^{\infty} (-n a_n x^n) + \left(\sum_{n=0}^{\infty} p^2 a_n x^n \right) = 0 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\begin{aligned} \sum_{n=2}^{\infty} (-x^n a_n n(n-1)) + \left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n \right) \\ + \sum_{n=1}^{\infty} (-n a_n x^n) + \left(\sum_{n=0}^{\infty} p^2 a_n x^n \right) = 0 \end{aligned} \quad (3)$$

$n = 0$ gives

$$a_0 p^2 + 2a_2 = 0$$

$$a_2 = -\frac{a_0 p^2}{2}$$

$n = 1$ gives

$$a_1 p^2 - a_1 + 6a_3 = 0$$

Which after substituting earlier equations, simplifies to

$$a_3 = -\frac{1}{6} a_1 p^2 + \frac{1}{6} a_1$$

For $2 \leq n$, the recurrence equation is

$$-n a_n (n-1) + (n+2) a_{n+2} (n+1) - n a_n + a_n p^2 = 0 \quad (4)$$

Solving for a_{n+2} , gives

$$a_{n+2} = \frac{a_n(n^2 - p^2)}{(n+2)(n+1)} \quad (5)$$

For $n = 2$ the recurrence equation gives

$$a_2p^2 - 4a_2 + 12a_4 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = \frac{1}{24}p^4a_0 - \frac{1}{6}a_0p^2$$

For $n = 3$ the recurrence equation gives

$$a_3p^2 - 9a_3 + 20a_5 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = \frac{1}{120}p^4a_1 - \frac{1}{12}a_1p^2 + \frac{3}{40}a_1$$

For $n = 4$ the recurrence equation gives

$$a_4p^2 - 16a_4 + 30a_6 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = -\frac{1}{720}p^6a_0 + \frac{1}{36}p^4a_0 - \frac{4}{45}a_0p^2$$

For $n = 5$ the recurrence equation gives

$$a_5p^2 - 25a_5 + 42a_7 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = -\frac{1}{5040}p^6a_1 + \frac{1}{144}p^4a_1 - \frac{37}{720}a_1p^2 + \frac{5}{112}a_1$$

For $n = 6$ the recurrence equation gives

$$a_6 p^2 - 36a_6 + 56a_8 = 0$$

Which after substituting the earlier terms found becomes

$$a_8 = \frac{1}{40320} p^8 a_0 - \frac{1}{720} p^6 a_0 + \frac{7}{360} p^4 a_0 - \frac{2}{35} a_0 p^2$$

For $n = 7$ the recurrence equation gives

$$a_7 p^2 - 49a_7 + 72a_9 = 0$$

Which after substituting the earlier terms found becomes

$$a_9 = \frac{1}{362880} p^8 a_1 - \frac{1}{4320} p^6 a_1 + \frac{47}{8640} p^4 a_1 - \frac{3229}{90720} a_1 p^2 + \frac{35}{1152} a_1$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$\begin{aligned} y &= a_0 + a_1 x - \frac{a_0 p^2 x^2}{2} + \left(-\frac{1}{6} a_1 p^2 + \frac{1}{6} a_1 \right) x^3 + \left(\frac{1}{24} p^4 a_0 - \frac{1}{6} a_0 p^2 \right) x^4 \\ &+ \left(\frac{1}{120} p^4 a_1 - \frac{1}{12} a_1 p^2 + \frac{3}{40} a_1 \right) x^5 + \left(-\frac{1}{720} p^6 a_0 + \frac{1}{36} p^4 a_0 - \frac{4}{45} a_0 p^2 \right) x^6 \\ &+ \left(-\frac{1}{5040} p^6 a_1 + \frac{1}{144} p^4 a_1 - \frac{37}{720} a_1 p^2 + \frac{5}{112} a_1 \right) x^7 + \dots \end{aligned}$$

Collecting terms, the solution becomes

$$\begin{aligned} y &= \left(1 - \frac{x^2 p^2}{2} + \left(\frac{1}{24} p^4 - \frac{1}{6} p^2 \right) x^4 + \left(-\frac{1}{720} p^6 + \frac{1}{36} p^4 - \frac{4}{45} p^2 \right) x^6 \right) a_0 \\ &+ \left(x + \left(-\frac{p^2}{6} + \frac{1}{6} \right) x^3 + \left(\frac{1}{120} p^4 - \frac{1}{12} p^2 + \frac{3}{40} \right) x^5 \right. \\ &\quad \left. + \left(-\frac{1}{5040} p^6 + \frac{1}{144} p^4 - \frac{37}{720} p^2 + \frac{5}{112} \right) x^7 \right) a_1 + O(x^8) \end{aligned} \tag{3}$$

At $x = 0$ the solution above becomes

$$y = \left(1 - \frac{x^2 p^2}{2} + \left(\frac{1}{24} p^4 - \frac{1}{6} p^2\right) x^4 + \left(-\frac{1}{720} p^6 + \frac{1}{36} p^4 - \frac{4}{45} p^2\right) x^6\right) c_1 \\ + \left(x + \left(-\frac{p^2}{6} + \frac{1}{6}\right) x^3 + \left(\frac{1}{120} p^4 - \frac{1}{12} p^2 + \frac{3}{40}\right) x^5\right. \\ \left. + \left(-\frac{1}{5040} p^6 + \frac{1}{144} p^4 - \frac{37}{720} p^2 + \frac{5}{112}\right) x^7\right) c_2 + O(x^8)$$

Summary

The solution(s) found are the following

$$y = \left(1 - \frac{1}{2} x^2 p^2 + \frac{1}{24} p^4 x^4 - \frac{1}{6} x^4 p^2 - \frac{1}{720} p^6 x^6 + \frac{1}{36} p^4 x^6 - \frac{4}{45} x^6 p^2 + \frac{1}{40320} x^8 p^8\right. \\ \left. - \frac{1}{720} x^8 p^6 + \frac{7}{360} x^8 p^4 - \frac{2}{35} x^8 p^2\right) y(0) + \left(x - \frac{1}{6} p^2 x^3 + \frac{1}{6} x^3 + \frac{1}{120} p^4 x^5\right. \\ \left. - \frac{1}{12} x^5 p^2 + \frac{3}{40} x^5 - \frac{1}{5040} x^7 p^6 + \frac{1}{144} x^7 p^4 - \frac{37}{720} x^7 p^2 + \frac{5}{112} x^7\right) y'(0) + O(x^8)$$

$$y = \left(1 - \frac{x^2 p^2}{2} + \left(\frac{1}{24} p^4 - \frac{1}{6} p^2\right) x^4 + \left(-\frac{1}{720} p^6 + \frac{1}{36} p^4 - \frac{4}{45} p^2\right) x^6\right) c_1 \\ + \left(x + \left(-\frac{p^2}{6} + \frac{1}{6}\right) x^3 + \left(\frac{1}{120} p^4 - \frac{1}{12} p^2 + \frac{3}{40}\right) x^5\right. \\ \left. + \left(-\frac{1}{5040} p^6 + \frac{1}{144} p^4 - \frac{37}{720} p^2 + \frac{5}{112}\right) x^7\right) c_2 + O(x^8) \quad (2)$$

Verification of solutions

$$y = \left(1 - \frac{1}{2}x^2p^2 + \frac{1}{24}p^4x^4 - \frac{1}{6}x^4p^2 - \frac{1}{720}p^6x^6 + \frac{1}{36}p^4x^6 - \frac{4}{45}x^6p^2 + \frac{1}{40320}x^8p^8 \right. \\ \left. - \frac{1}{720}x^8p^6 + \frac{7}{360}x^8p^4 - \frac{2}{35}x^8p^2 \right) y(0) + \left(x - \frac{1}{6}p^2x^3 + \frac{1}{6}x^3 + \frac{1}{120}p^4x^5 - \frac{1}{12}x^5p^2 \right. \\ \left. + \frac{3}{40}x^5 - \frac{1}{5040}x^7p^6 + \frac{1}{144}x^7p^4 - \frac{37}{720}x^7p^2 + \frac{5}{112}x^7 \right) y'(0) + O(x^8)$$

Verified OK.

$$y = \left(1 - \frac{x^2p^2}{2} + \left(\frac{1}{24}p^4 - \frac{1}{6}p^2 \right) x^4 + \left(-\frac{1}{720}p^6 + \frac{1}{36}p^4 - \frac{4}{45}p^2 \right) x^6 \right) c_1 \\ + \left(x + \left(-\frac{p^2}{6} + \frac{1}{6} \right) x^3 + \left(\frac{1}{120}p^4 - \frac{1}{12}p^2 + \frac{3}{40} \right) x^5 \right. \\ \left. + \left(-\frac{1}{5040}p^6 + \frac{1}{144}p^4 - \frac{37}{720}p^2 + \frac{5}{112} \right) x^7 \right) c_2 + O(x^8)$$

Verified OK.

18.13.1 Maple step by step solution

Let's solve

$$(-x^2 + 1)y'' - xy' + p^2y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{xy'}{x^2-1} + \frac{p^2y}{x^2-1}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{xy'}{x^2-1} - \frac{p^2y}{x^2-1} = 0$$

- Check to see if x_0 is a regular singular point

- Define functions

$$\left[P_2(x) = \frac{x}{x^2-1}, P_3(x) = -\frac{p^2}{x^2-1} \right]$$

- $(1+x) \cdot P_2(x)$ is analytic at $x = -1$

$$\left. ((1+x) \cdot P_2(x)) \right|_{x=-1} = \frac{1}{2}$$

- $(1+x)^2 \cdot P_3(x)$ is analytic at $x = -1$

$$\left. ((1+x)^2 \cdot P_3(x)) \right|_{x=-1} = 0$$

- $x = -1$ is a regular singular point

Check to see if x_0 is a regular singular point

$$x_0 = -1$$

- Multiply by denominators

$$(x^2 - 1)y'' + xy' - p^2y = 0$$

- Change variables using $x = u - 1$ so that the regular singular point is at $u = 0$

$$(u^2 - 2u) \left(\frac{d^2}{du^2} y(u) \right) + (u - 1) \left(\frac{d}{du} y(u) \right) - p^2 y(u) = 0$$

- Assume series solution for $y(u)$

$$y(u) = \sum_{k=0}^{\infty} a_k u^{k+r}$$

- Rewrite ODE with series expansions

- Convert $u^m \cdot \left(\frac{d}{du} y(u) \right)$ to series expansion for $m = 0..1$

$$u^m \cdot \left(\frac{d}{du} y(u) \right) = \sum_{k=0}^{\infty} a_k (k+r) u^{k+r-1+m}$$

- Shift index using $k \rightarrow k + 1 - m$

$$u^m \cdot \left(\frac{d}{du} y(u) \right) = \sum_{k=-1+m}^{\infty} a_{k+1-m} (k+1-m+r) u^{k+r}$$

- Convert $u^m \cdot \left(\frac{d^2}{du^2} y(u) \right)$ to series expansion for $m = 1..2$

$$u^m \cdot \left(\frac{d^2}{du^2} y(u) \right) = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) u^{k+r-2+m}$$

- Shift index using $k \rightarrow k + 2 - m$

$$u^m \cdot \left(\frac{d^2}{du^2} y(u) \right) = \sum_{k=-2+m}^{\infty} a_{k+2-m} (k+2-m+r)(k+1-m+r) u^{k+r}$$

Rewrite ODE with series expansions

$$-a_0 r(-1+2r) u^{-1+r} + \left(\sum_{k=0}^{\infty} (-a_{k+1} (k+1+r)(2k+1+2r) + a_k (k+p+r)(k-p+r)) \right) u^{k+r}$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$-r(-1+2r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \left\{0, \frac{1}{2}\right\}$$

- Each term in the series must be 0, giving the recursion relation

$$-2\left(k + \frac{1}{2} + r\right)(k + 1 + r)a_{k+1} + a_k(k + p + r)(k - p + r) = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+1} = \frac{a_k(k+p+r)(k-p+r)}{(2k+1+2r)(k+1+r)}$$

- Recursion relation for $r = 0$

$$a_{k+1} = \frac{a_k(k+p)(k-p)}{(2k+1)(k+1)}$$

- Solution for $r = 0$

$$\left[y(u) = \sum_{k=0}^{\infty} a_k u^k, a_{k+1} = \frac{a_k(k+p)(k-p)}{(2k+1)(k+1)} \right]$$

- Revert the change of variables $u = 1 + x$

$$\left[y = \sum_{k=0}^{\infty} a_k (1+x)^k, a_{k+1} = \frac{a_k(k+p)(k-p)}{(2k+1)(k+1)} \right]$$

- Recursion relation for $r = \frac{1}{2}$

$$a_{k+1} = \frac{a_k(k+p+\frac{1}{2})(k-p+\frac{1}{2})}{(2k+2)(k+\frac{3}{2})}$$

- Solution for $r = \frac{1}{2}$

$$\left[y(u) = \sum_{k=0}^{\infty} a_k u^{k+\frac{1}{2}}, a_{k+1} = \frac{a_k(k+p+\frac{1}{2})(k-p+\frac{1}{2})}{(2k+2)(k+\frac{3}{2})} \right]$$

- Revert the change of variables $u = 1 + x$

$$\left[y = \sum_{k=0}^{\infty} a_k (1+x)^{k+\frac{1}{2}}, a_{k+1} = \frac{a_k(k+p+\frac{1}{2})(k-p+\frac{1}{2})}{(2k+2)(k+\frac{3}{2})} \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k (1+x)^k \right) + \left(\sum_{k=0}^{\infty} b_k (1+x)^{k+\frac{1}{2}} \right), a_{k+1} = \frac{a_k(k+p)(k-p)}{(2k+1)(k+1)}, b_{k+1} = \frac{b_k(k+p+\frac{1}{2})(k-p+\frac{1}{2})}{(2k+2)(k+\frac{3}{2})} \right]$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
<- linear_1 successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 121

```
Order:=8;  
dsolve((1-x^2)*diff(y(x),x$2)-x*diff(y(x),x)+p^2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{p^2 x^2}{2} + \frac{p^2(p^2 - 4)x^4}{24} - \frac{p^2(p^4 - 20p^2 + 64)x^6}{720}\right) y(0) + \left(x - \frac{(p^2 - 1)x^3}{6} + \frac{(p^4 - 10p^2 + 9)x^5}{120} - \frac{(p^6 - 35p^4 + 259p^2 - 225)x^7}{5040}\right) D(y)(0) + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 155

```
AsymptoticDSolveValue[(1-x^2)*y'[x]-x*y'[x]+p^2*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_2 \left(-\frac{p^6 x^7}{5040} + \frac{p^4 x^7}{144} + \frac{p^4 x^5}{120} - \frac{37p^2 x^7}{720} - \frac{p^2 x^5}{12} - \frac{p^2 x^3}{6} + \frac{5x^7}{112} + \frac{3x^5}{40} + \frac{x^3}{6} + x \right) + c_1 \left(-\frac{1}{720} p^6 x^6 + \frac{p^4 x^6}{36} + \frac{p^4 x^4}{24} - \frac{4p^2 x^6}{45} - \frac{p^2 x^4}{6} - \frac{p^2 x^2}{2} + 1 \right)$$

18.14 problem 8

18.14.1 Maple step by step solution 3114

Internal problem ID [6440]

Internal file name [OUTPUT/5688_Sunday_June_05_2022_03_47_03_PM_40872846/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.3. Second-Order Linear Equations: Ordinary Points. Page 169

Problem number: 8.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Ordinary point", "second order series method. Taylor series method"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - 2xy' + 2py = 0$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using Taylor series gives

$$\begin{aligned} y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \cdots \\ &= y_0 + xy'_0 + \frac{x^2}{2}f|_{x_0, y_0, y'_0} + \frac{x^3}{3!}f'|_{x_0, y_0, y'_0} + \cdots \\ &= y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0} \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \quad (1)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \quad (686)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \quad (687)$$

$$\begin{aligned} \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \end{aligned} \quad (2)$$

$$\begin{aligned} \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \end{aligned} \quad (3)$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \quad (4)$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \quad (5)$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \\ &\vdots \\ F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \quad (6)$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \quad (7)$$

To find $y(x)$ series solution around $x = 0$. Hence

$$F_0 = 2xy' - 2py$$

$$\begin{aligned} F_1 &= \frac{dF_0}{dx} \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \\ &= (4x^2 - 2p + 2) y' - 4pxy \end{aligned}$$

$$\begin{aligned} F_2 &= \frac{dF_1}{dx} \\ &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} y' + \frac{\partial F_1}{\partial y'} F_1 \\ &= (8x^3 - 8px + 12x) y' + 4py(-2x^2 + p - 2) \end{aligned}$$

$$\begin{aligned} F_3 &= \frac{dF_2}{dx} \\ &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} y' + \frac{\partial F_2}{\partial y'} F_2 \\ &= 4(3 + p^2 + 2(-3x^2 - 2)p + 4x^4 + 12x^2) y' + 16yxp \left(-x^2 + p - \frac{5}{2} \right) \end{aligned}$$

$$\begin{aligned} F_4 &= \frac{dF_3}{dx} \\ &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} y' + \frac{\partial F_3}{\partial y'} F_3 \\ &= 24x \left(p^2 + \left(-\frac{8x^2}{3} - 5 \right) p + \frac{4x^4}{3} + \frac{20x^2}{3} + 5 \right) y' - 8py(4x^4 - 6x^2p + p^2 + 18x^2 - 6p + 8) \end{aligned}$$

$$\begin{aligned} F_5 &= \frac{dF_4}{dx} \\ &= \frac{\partial F_4}{\partial x} + \frac{\partial F_4}{\partial y} y' + \frac{\partial F_4}{\partial y'} F_4 \\ &= (-8p^3 + (96x^2 + 72)p^2 + (-160x^4 - 576x^2 - 184)p + 64x^6 + 480x^4 + 720x^2 + 120) y' - 48y \left(p^2 + \right. \end{aligned}$$

$$\begin{aligned} F_6 &= \frac{dF_5}{dx} \\ &= \frac{\partial F_5}{\partial x} + \frac{\partial F_5}{\partial y} y' + \frac{\partial F_5}{\partial y'} F_5 \\ &= -64x \left(p^3 + \left(-5x^2 - \frac{21}{2} \right) p^2 + (6x^4 + 35x^2 + 32)p - 2x^6 - 21x^4 - \frac{105x^2}{2} - \frac{105}{4} \right) y' + 16y(p^3 + \left. \right) \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and

$y'(0) = y'(0)$ gives

$$\begin{aligned}
 F_0 &= -2y(0)p \\
 F_1 &= -2y'(0)p + 2y'(0) \\
 F_2 &= 4y(0)p^2 - 8y(0)p \\
 F_3 &= 4y'(0)p^2 - 16y'(0)p + 12y'(0) \\
 F_4 &= -8y(0)p^3 + 48y(0)p^2 - 64y(0)p \\
 F_5 &= -8y'(0)p^3 + 72y'(0)p^2 - 184y'(0)p + 120y'(0) \\
 F_6 &= 16y(0)p^4 - 192y(0)p^3 + 704y(0)p^2 - 768y(0)p
 \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$\begin{aligned}
 y = & \left(1 - x^2p + \frac{1}{6}x^4p^2 - \frac{1}{3}x^4p - \frac{1}{90}x^6p^3 + \frac{1}{15}x^6p^2 - \frac{4}{45}x^6p + \frac{1}{2520}x^8p^4 - \frac{1}{210}x^8p^3 \right. \\
 & \left. + \frac{11}{630}x^8p^2 - \frac{2}{105}x^8p \right) y(0) + \left(x - \frac{1}{3}x^3p + \frac{1}{3}x^3 + \frac{1}{30}x^5p^2 - \frac{2}{15}x^5p + \frac{1}{10}x^5 \right. \\
 & \left. - \frac{1}{630}x^7p^3 + \frac{1}{70}x^7p^2 - \frac{23}{630}x^7p + \frac{1}{42}x^7 \right) y'(0) + O(x^8)
 \end{aligned}$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard power series Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$\begin{aligned}
 y' &= \sum_{n=1}^{\infty} n a_n x^{n-1} \\
 y'' &= \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}
 \end{aligned}$$

Substituting the above back into the ode gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = 2x \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) - 2p \left(\sum_{n=0}^{\infty} a_n x^n \right) \quad (1)$$

Which simplifies to

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \sum_{n=1}^{\infty} (-2n x^n a_n) + \left(\sum_{n=0}^{\infty} 2p a_n x^n \right) = 0 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n \right) + \sum_{n=1}^{\infty} (-2n x^n a_n) + \left(\sum_{n=0}^{\infty} 2p a_n x^n \right) = 0 \quad (3)$$

$n = 0$ gives

$$2pa_0 + 2a_2 = 0$$

$$a_2 = -pa_0$$

For $1 \leq n$, the recurrence equation is

$$(n+2) a_{n+2} (n+1) - 2na_n + 2pa_n = 0 \quad (4)$$

Solving for a_{n+2} , gives

$$a_{n+2} = \frac{2a_n(n-p)}{(n+2)(n+1)} \quad (5)$$

For $n = 1$ the recurrence equation gives

$$2pa_1 - 2a_1 + 6a_3 = 0$$

Which after substituting the earlier terms found becomes

$$a_3 = -\frac{1}{3}pa_1 + \frac{1}{3}a_1$$

For $n = 2$ the recurrence equation gives

$$2pa_2 - 4a_2 + 12a_4 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = \frac{1}{6}p^2a_0 - \frac{1}{3}pa_0$$

For $n = 3$ the recurrence equation gives

$$2pa_3 - 6a_3 + 20a_5 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = \frac{1}{30}p^2a_1 - \frac{2}{15}pa_1 + \frac{1}{10}a_1$$

For $n = 4$ the recurrence equation gives

$$2pa_4 - 8a_4 + 30a_6 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = -\frac{1}{90}p^3a_0 + \frac{1}{15}p^2a_0 - \frac{4}{45}pa_0$$

For $n = 5$ the recurrence equation gives

$$2pa_5 - 10a_5 + 42a_7 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = -\frac{1}{630}p^3a_1 + \frac{1}{70}p^2a_1 - \frac{23}{630}pa_1 + \frac{1}{42}a_1$$

For $n = 6$ the recurrence equation gives

$$2pa_6 - 12a_6 + 56a_8 = 0$$

Which after substituting the earlier terms found becomes

$$a_8 = \frac{1}{2520}p^4a_0 - \frac{1}{210}p^3a_0 + \frac{11}{630}p^2a_0 - \frac{2}{105}pa_0$$

For $n = 7$ the recurrence equation gives

$$2pa_7 - 14a_7 + 72a_9 = 0$$

Which after substituting the earlier terms found becomes

$$a_9 = \frac{1}{22680}p^4a_1 - \frac{2}{2835}p^3a_1 + \frac{43}{11340}p^2a_1 - \frac{22}{2835}pa_1 + \frac{1}{216}a_1$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$\begin{aligned} y &= a_0 + a_1 x - pa_0 x^2 + \left(-\frac{1}{3}pa_1 + \frac{1}{3}a_1\right) x^3 + \left(\frac{1}{6}p^2a_0 - \frac{1}{3}pa_0\right) x^4 \\ &+ \left(\frac{1}{30}p^2a_1 - \frac{2}{15}pa_1 + \frac{1}{10}a_1\right) x^5 + \left(-\frac{1}{90}p^3a_0 + \frac{1}{15}p^2a_0 - \frac{4}{45}pa_0\right) x^6 \\ &+ \left(-\frac{1}{630}p^3a_1 + \frac{1}{70}p^2a_1 - \frac{23}{630}pa_1 + \frac{1}{42}a_1\right) x^7 + \dots \end{aligned}$$

Collecting terms, the solution becomes

$$\begin{aligned} y &= \left(1 - x^2p + \left(\frac{1}{6}p^2 - \frac{1}{3}p\right) x^4 + \left(-\frac{1}{90}p^3 + \frac{1}{15}p^2 - \frac{4}{45}p\right) x^6\right) a_0 + \left(x + \left(-\frac{p}{3} + \frac{1}{3}\right) x^3\right. \\ &+ \left.\left(\frac{1}{30}p^2 - \frac{2}{15}p + \frac{1}{10}\right) x^5 + \left(-\frac{1}{630}p^3 + \frac{1}{70}p^2 - \frac{23}{630}p + \frac{1}{42}\right) x^7\right) a_1 + O(x^8) \end{aligned} \tag{3}$$

At $x = 0$ the solution above becomes

$$\begin{aligned} y &= \left(1 - x^2p + \left(\frac{1}{6}p^2 - \frac{1}{3}p\right) x^4 + \left(-\frac{1}{90}p^3 + \frac{1}{15}p^2 - \frac{4}{45}p\right) x^6\right) c_1 + \left(x + \left(-\frac{p}{3} + \frac{1}{3}\right) x^3\right. \\ &+ \left.\left(\frac{1}{30}p^2 - \frac{2}{15}p + \frac{1}{10}\right) x^5 + \left(-\frac{1}{630}p^3 + \frac{1}{70}p^2 - \frac{23}{630}p + \frac{1}{42}\right) x^7\right) c_2 + O(x^8) \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}y &= \left(1 - x^2p + \frac{1}{6}x^4p^2 - \frac{1}{3}x^4p - \frac{1}{90}x^6p^3 + \frac{1}{15}x^6p^2 - \frac{4}{45}x^6p + \frac{1}{2520}x^8p^4 - \frac{1}{210}x^8p^3\right. \\&\quad \left. + \frac{11}{630}x^8p^2 - \frac{2}{105}x^8p\right) y(0) + \left(x - \frac{1}{3}x^3p + \frac{1}{3}x^3 + \frac{1}{30}x^5p^2 - \frac{2}{15}x^5p + \frac{1}{10}x^5\right. \\&\quad \left. - \frac{1}{630}x^7p^3 + \frac{1}{70}x^7p^2 - \frac{23}{630}x^7p + \frac{1}{42}x^7\right) y'(0) + O(x^8) \\y &= \left(1 - x^2p + \left(\frac{1}{6}p^2 - \frac{1}{3}p\right) x^4 + \left(-\frac{1}{90}p^3 + \frac{1}{15}p^2 - \frac{4}{45}p\right) x^6\right) c_1 + \left(x + \left(-\frac{p}{3} + \frac{1}{3}\right) x^3\right. \\&\quad \left. + \left(\frac{1}{30}p^2 - \frac{2}{15}p + \frac{1}{10}\right) x^5 + \left(-\frac{1}{630}p^3 + \frac{1}{70}p^2 - \frac{23}{630}p + \frac{1}{42}\right) x^7\right) c_2 + O(x^8)\end{aligned}\tag{2}$$

Verification of solutions

$$\begin{aligned}y &= \left(1 - x^2p + \frac{1}{6}x^4p^2 - \frac{1}{3}x^4p - \frac{1}{90}x^6p^3 + \frac{1}{15}x^6p^2 - \frac{4}{45}x^6p + \frac{1}{2520}x^8p^4 - \frac{1}{210}x^8p^3\right. \\&\quad \left. + \frac{11}{630}x^8p^2 - \frac{2}{105}x^8p\right) y(0) + \left(x - \frac{1}{3}x^3p + \frac{1}{3}x^3 + \frac{1}{30}x^5p^2 - \frac{2}{15}x^5p + \frac{1}{10}x^5\right. \\&\quad \left. - \frac{1}{630}x^7p^3 + \frac{1}{70}x^7p^2 - \frac{23}{630}x^7p + \frac{1}{42}x^7\right) y'(0) + O(x^8)\end{aligned}$$

Verified OK.

$$\begin{aligned}y &= \left(1 - x^2p + \left(\frac{1}{6}p^2 - \frac{1}{3}p\right) x^4 + \left(-\frac{1}{90}p^3 + \frac{1}{15}p^2 - \frac{4}{45}p\right) x^6\right) c_1 + \left(x + \left(-\frac{p}{3} + \frac{1}{3}\right) x^3\right. \\&\quad \left. + \left(\frac{1}{30}p^2 - \frac{2}{15}p + \frac{1}{10}\right) x^5 + \left(-\frac{1}{630}p^3 + \frac{1}{70}p^2 - \frac{23}{630}p + \frac{1}{42}\right) x^7\right) c_2 + O(x^8)\end{aligned}$$

Verified OK.

18.14.1 Maple step by step solution

Let's solve

$$y'' = 2xy' - 2py$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - 2xy' + 2py = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^k$$

- Rewrite DE with series expansions

- Convert $x \cdot y'$ to series expansion

$$x \cdot y' = \sum_{k=0}^{\infty} a_k k x^k$$

- Convert y'' to series expansion

$$y'' = \sum_{k=2}^{\infty} a_k k(k-1) x^{k-2}$$

- Shift index using $k- \rightarrow k+2$

$$y'' = \sum_{k=0}^{\infty} a_{k+2} (k+2)(k+1) x^k$$

Rewrite DE with series expansions

$$\sum_{k=0}^{\infty} (a_{k+2}(k+2)(k+1) - 2a_k(k-p)) x^k = 0$$

- Each term in the series must be 0, giving the recursion relation

$$(k^2 + 3k + 2) a_{k+2} - 2a_k(k-p) = 0$$

- Recursion relation that defines the series solution to the ODE

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+2} = \frac{2a_k(k-p)}{k^2+3k+2} \right]$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
  -> Bessel
  -> elliptic
  -> Legendre
  -> Kummer
      -> hyper3: Equivalence to 1F1 under a power @ Moebius
      <- hyper3 successful: received ODE is equivalent to the 1F1 ODE
      <- Kummer successful
<- special function solution successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 109

```
Order:=8;
dsolve(diff(y(x),x$2)-2*x*diff(y(x),x)+2*p*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - px^2 + \frac{p(p-2)x^4}{6} - \frac{p(p-2)(p-4)x^6}{90}\right) y(0) \\ + \left(x - \frac{(p-1)x^3}{3} + \frac{(p^2-4p+3)x^5}{30} - \frac{(p^3-9p^2+23p-15)x^7}{630}\right) D(y)(0) \\ + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 141

```
AsymptoticDSolveValue[y''[x]-2*x*y'[x]+2*p*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_2 \left(-\frac{1}{630} p^3 x^7 + \frac{p^2 x^7}{70} + \frac{p^2 x^5}{30} - \frac{23 p x^7}{630} - \frac{2 p x^5}{15} - \frac{p x^3}{3} + \frac{x^7}{42} + \frac{x^5}{10} + \frac{x^3}{3} + x \right) \\ + c_1 \left(-\frac{1}{90} p^3 x^6 + \frac{p^2 x^6}{15} + \frac{p^2 x^4}{6} - \frac{4 p x^6}{45} - \frac{p x^4}{3} - p x^2 + 1 \right)$$

19 Chapter 4. Power Series Solutions and Special Functions. Section 4.4. REGULAR SINGULAR POINTS. Page 175

19.1	problem 1(a)	3119
19.2	problem 1(b)	3123
19.3	problem 1(c)	3139
19.4	problem 1(d)	3142
19.5	problem 2(a)	3161
19.6	problem 2(b)	3173
19.7	problem 2(c)	3184
19.8	problem 2(d)	3199
19.9	problem 2(e)	3211
19.10	problem 3(a)	3215
19.11	problem 3(b)	3228
19.12	problem 3(c)	3244
19.13	problem 3(d)	3261
19.14	problem 4(a)	3274
19.15	problem 4(b)	3290
19.16	problem 4(c)	3306
19.17	problem 4(d)	3323
19.18	problem 5	3339
19.19	problem 6	3351
19.20	problem 8	3354

19.1 problem 1(a)

Internal problem ID [6441]

Internal file name [OUTPUT/5689_Sunday_June_05_2022_03_47_04_PM_30345490/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.4. REGULAR SINGULAR POINTS. Page 175

Problem number: 1(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Irregular singular point"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

Unable to solve or complete the solution.

$$x^3(x-1)y'' - 2(x-1)y' + 3xy = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$(x^4 - x^3)y'' + (-2x + 2)y' + 3xy = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{2}{x^3}$$
$$q(x) = \frac{3}{x^2(x-1)}$$

Table 439: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{2}{x^3}$	
singularity	type
$x = 0$	“irregular”

$q(x) = \frac{3}{x^2(x-1)}$	
singularity	type
$x = 0$	“regular”
$x = 1$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[1, \infty]$


Irregular singular points : $[0]$

Since $x = 0$ is not an ordinary point, then we will now check if it is a regular singular point. Unable to solve since $x = 0$ is not regular singular point. Terminating.

Verification of solutions N/A

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
  -> Bessel
  -> elliptic
  -> Legendre
  -> Kummer
    -> hyper3: Equivalence to 1F1 under a power @ Moebius
  -> hypergeometric
    -> heuristic approach
    -> hyper3: Equivalence to 2F1, 1F1 or 0F1 under a power @ Moebius
  -> Mathieu
    -> Equivalence to the rational form of Mathieu ODE under a power @ Moebius
trying a solution in terms of MeijerG functions
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form  $r_0(x) * Y + r_1(x) * Y$  where  $Y = \exp(\int(r(x), dx)) * 2F1([a$ 
  trying a symmetry of the form [xi=0, eta=F(x)]
  trying differential order: 2; exact nonlinear
  trying symmetries linear in x and y(x)
  trying to convert to a linear ODE with constant coefficients
  trying 2nd order, integrating factor of the form  $\mu(x,y)$ 
  -> Trying a solution in terms of special functions:
    -> Bessel
    -> elliptic
    -> Legendre
    -> Whittaker
      -> hyper3: Equivalence to 1F1 under a power @ Moebius
    -> hypergeometric
      -> heuristic approach
      -> hyper3: Equivalence to 2F1, 1F1 or 0F1 under a power @ Moebius
    -> Mathieu
      -> Equivalence to the rational form of Mathieu ODE under a power @ Moebius
  trying 2nd order exact linear
  trying symmetries linear in x and y(x)
  trying to convert to a linear ODE with constant coefficients
  trying to convert to an ODE of Bessel type
trying to convert to an ODE of Bessel type
-> trying reduction of order to Bessel
```

 Solution by Maple

```
Order:=8;  
dsolve(x^3*(x-1)*diff(y(x),x$2)-2*(x-1)*diff(y(x),x)+3*x*y(x)=0,y(x),type='series',x=0);
```

No solution found

 Solution by Mathematica

Time used: 0.053 (sec). Leaf size: 108

```
AsymptoticDSolveValue[x^3*(x-1)*y''[x]-2*(x-1)*y'[x]+3*x*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_2 e^{-\frac{1}{x^2}} \left(\frac{1731x^7}{320} - \frac{795x^6}{128} - \frac{51x^5}{40} + \frac{63x^4}{32} + \frac{x^3}{2} - \frac{3x^2}{4} + 1 \right) x^3 \\ + c_1 \left(-\frac{51x^7}{320} - \frac{19x^6}{128} - \frac{9x^5}{40} - \frac{9x^4}{32} - \frac{x^3}{2} - \frac{3x^2}{4} + 1 \right)$$

19.2 problem 1(b)

Internal problem ID [6442]

Internal file name [OUTPUT/5690_Sunday_June_05_2022_03_47_07_PM_58129404/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.4. REGULAR SINGULAR POINTS. Page 175

Problem number: 1(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$x^2(x^2 - 1)y'' - x(1 - x)y' + 2y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$(x^4 - x^2)y'' + (x^2 - x)y' + 2y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{1}{x(1+x)}$$
$$q(x) = \frac{2}{x^2(x^2 - 1)}$$

Table 440: Table $p(x), q(x)$ singularities.

$p(x) = \frac{1}{x(1+x)}$	
singularity	type
$x = -1$	“regular”
$x = 0$	“regular”

$q(x) = \frac{2}{x^2(x^2-1)}$	
singularity	type
$x = -1$	“regular”
$x = 0$	“regular”
$x = 1$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[-1, 0, 1, \infty]$

Irregular singular points : $[\]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2(x^2 - 1)y'' + (x^2 - x)y' + 2y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned} & x^2(x^2 - 1) \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\ & + (x^2 - x) \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + 2 \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \end{aligned} \tag{1}$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r+2} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r) (n+r-1)) \\ & + \left(\sum_{n=0}^{\infty} x^{1+n+r} a_n (n+r) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) + \left(\sum_{n=0}^{\infty} 2a_n x^{n+r} \right) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=0}^{\infty} x^{n+r+2} a_n (n+r) (n+r-1) &= \sum_{n=2}^{\infty} a_{n-2} (n+r-2) (n-3+r) x^{n+r} \\ \sum_{n=0}^{\infty} x^{1+n+r} a_n (n+r) &= \sum_{n=1}^{\infty} a_{n-1} (n+r-1) x^{n+r} \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=2}^{\infty} a_{n-2} (n+r-2) (n-3+r) x^{n+r} \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r) (n+r \\ & - 1)) + \left(\sum_{n=1}^{\infty} a_{n-1} (n+r-1) x^{n+r} \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n \\ & + r)) + \left(\sum_{n=0}^{\infty} 2a_n x^{n+r} \right) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$-x^{n+r} a_n (n+r) (n+r-1) - x^{n+r} a_n (n+r) + 2a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$-x^r a_0 r (-1+r) - x^r a_0 r + 2a_0 x^r = 0$$

Or

$$(-x^r r (-1+r) - x^r r + 2x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$-(r^2 - 2) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$-r^2 + 2 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = -\sqrt{2}$$

$$r_2 = \sqrt{2}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$-(r^2 - 2) x^r = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = -2\sqrt{2}$ is not an integer, then we can construct two linearly independent solutions

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n-\sqrt{2}}$$

$$y_2(x) = \sum_{n=0}^{\infty} b_n x^{n+\sqrt{2}}$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = \frac{r}{r^2 + 2r - 1}$$

For $2 \leq n$ the recursive equation is

$$a_{n-2}(n+r-2)(n-3+r) - a_n(n+r)(n+r-1) + a_{n-1}(n+r-1) - a_n(n+r) + 2a_n = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{n^2 a_{n-2} + 2nr a_{n-2} + r^2 a_{n-2} - 5n a_{n-2} + n a_{n-1} - 5r a_{n-2} + r a_{n-1} + 6a_{n-2} - a_{n-1}}{n^2 + 2nr + r^2 - 2} \quad (4)$$

Which for the root $r = -\sqrt{2}$ becomes

$$a_n = \frac{(-2n a_{n-2} + 5a_{n-2} - a_{n-1})\sqrt{2} + n^2 a_{n-2} + (-5a_{n-2} + a_{n-1})n + 8a_{n-2} - a_{n-1}}{n(-2\sqrt{2} + n)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = -\sqrt{2}$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{r}{r^2+2r-1}$	$\frac{\sqrt{2}}{-1+2\sqrt{2}}$

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{r(r^3 + r^2 - 2r + 2)}{(r^2 + 2r - 1)(r^2 + 4r + 2)}$$

Which for the root $r = -\sqrt{2}$ becomes

$$a_2 = \frac{\sqrt{2}}{-5 + 3\sqrt{2}}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{r}{r^2+2r-1}$	$\frac{\sqrt{2}}{-1+2\sqrt{2}}$
a_2	$\frac{r(r^3+r^2-2r+2)}{(r^2+2r-1)(r^2+4r+2)}$	$\frac{\sqrt{2}}{-5+3\sqrt{2}}$

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{2r(r^4 + 4r^3 + 3r^2 + 2)}{(r^2 + 2r - 1)(r^2 + 4r + 2)(r^2 + 6r + 7)}$$

Which for the root $r = -\sqrt{2}$ becomes

$$a_3 = -\frac{2\sqrt{2}}{15 - 9\sqrt{2}}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{r}{r^2+2r-1}$	$\frac{\sqrt{2}}{-1+2\sqrt{2}}$
a_2	$\frac{r(r^3+r^2-2r+2)}{(r^2+2r-1)(r^2+4r+2)}$	$\frac{\sqrt{2}}{-5+3\sqrt{2}}$
a_3	$\frac{2r(r^4+4r^3+3r^2+2)}{(r^2+2r-1)(r^2+4r+2)(r^2+6r+7)}$	$-\frac{2\sqrt{2}}{15-9\sqrt{2}}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{r(r^7 + 10r^6 + 36r^5 + 58r^4 + 41r^3 + 20r^2 + 42r + 40)}{(r^2 + 2r - 1)(r^2 + 4r + 2)(r^2 + 6r + 7)(r^2 + 8r + 14)}$$

Which for the root $r = -\sqrt{2}$ becomes

$$a_4 = \frac{-69 + 49\sqrt{2}}{-1104 + 780\sqrt{2}}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{r}{r^2+2r-1}$	$\frac{\sqrt{2}}{-1+2\sqrt{2}}$
a_2	$\frac{r(r^3+r^2-2r+2)}{(r^2+2r-1)(r^2+4r+2)}$	$\frac{\sqrt{2}}{-5+3\sqrt{2}}$
a_3	$\frac{2r(r^4+4r^3+3r^2+2)}{(r^2+2r-1)(r^2+4r+2)(r^2+6r+7)}$	$-\frac{2\sqrt{2}}{15-9\sqrt{2}}$
a_4	$\frac{r(r^7+10r^6+36r^5+58r^4+41r^3+20r^2+42r+40)}{(r^2+2r-1)(r^2+4r+2)(r^2+6r+7)(r^2+8r+14)}$	$\frac{-69+49\sqrt{2}}{-1104+780\sqrt{2}}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{r(3r^8 + 48r^7 + 306r^6 + 996r^5 + 1749r^4 + 1616r^3 + 866r^2 + 680r + 496)}{(r^2 + 2r - 1)(r^2 + 4r + 2)(r^2 + 6r + 7)(r^2 + 8r + 14)(r^2 + 10r + 23)}$$

Which for the root $r = -\sqrt{2}$ becomes

$$a_5 = \frac{-414 + 293\sqrt{2}}{6108\sqrt{2} - 8640}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{r}{r^2+2r-1}$	$\frac{\sqrt{2}}{-1+2\sqrt{2}}$
a_2	$\frac{r(r^3+r^2-2r+2)}{(r^2+2r-1)(r^2+4r+2)}$	$\frac{\sqrt{2}}{-5+3\sqrt{2}}$
a_3	$\frac{2r(r^4+4r^3+3r^2+2)}{(r^2+2r-1)(r^2+4r+2)(r^2+6r+7)}$	$-\frac{2\sqrt{2}}{15-9\sqrt{2}}$
a_4	$\frac{r(r^7+10r^6+36r^5+58r^4+41r^3+20r^2+42r+40)}{(r^2+2r-1)(r^2+4r+2)(r^2+6r+7)(r^2+8r+14)}$	$\frac{-69+49\sqrt{2}}{-1104+780\sqrt{2}}$
a_5	$\frac{r(3r^8+48r^7+306r^6+996r^5+1749r^4+1616r^3+866r^2+680r+496)}{(r^2+2r-1)(r^2+4r+2)(r^2+6r+7)(r^2+8r+14)(r^2+10r+23)}$	$\frac{-414+293\sqrt{2}}{6108\sqrt{2}-8640}$

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{r(r^{11} + 27r^{10} + 314r^9 + 2064r^8 + 8439r^7 + 22209r^6 + 37650r^5 + 40744r^4 + 30972r^3 + 26532r^2 + 26728r + 13520)}{(r^2 + 2r - 1)(r^2 + 4r + 2)(r^2 + 6r + 7)(r^2 + 8r + 14)(r^2 + 10r + 23)(r^2 + 12r + 34)}$$

Which for the root $r = -\sqrt{2}$ becomes

$$a_6 = \frac{3898 - 2757\sqrt{2}}{114408 - 80892\sqrt{2}}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{r}{r^2+2r-1}$	$\frac{\sqrt{2}}{-1+2\sqrt{2}}$
a_2	$\frac{r(r^3+r^2-2r+2)}{(r^2+2r-1)(r^2+4r+2)}$	$\frac{\sqrt{2}}{-5+3\sqrt{2}}$
a_3	$\frac{2r(r^4+4r^3+3r^2+2)}{(r^2+2r-1)(r^2+4r+2)(r^2+6r+7)}$	$-\frac{2\sqrt{2}}{15-9\sqrt{2}}$
a_4	$\frac{r(r^7+10r^6+36r^5+58r^4+41r^3+20r^2+42r+40)}{(r^2+2r-1)(r^2+4r+2)(r^2+6r+7)(r^2+8r+14)}$	$\frac{-69+49\sqrt{2}}{-1104+780\sqrt{2}}$
a_5	$\frac{r(3r^8+48r^7+306r^6+996r^5+1749r^4+1616r^3+866r^2+680r+496)}{(r^2+2r-1)(r^2+4r+2)(r^2+6r+7)(r^2+8r+14)(r^2+10r+23)}$	$\frac{-414+293\sqrt{2}}{6108\sqrt{2}-8640}$
a_6	$\frac{r(r^{11}+27r^{10}+314r^9+2064r^8+8439r^7+22209r^6+37650r^5+40744r^4+30972r^3+26532r^2+26728r+13520)}{(r^2+2r-1)(r^2+4r+2)(r^2+6r+7)(r^2+8r+14)(r^2+10r+23)(r^2+12r+34)}$	$\frac{3898-2757\sqrt{2}}{114408-80892\sqrt{2}}$

For $n = 7$, using the above recursive equation gives

$$a_7 = \frac{4r(r^{12} + 36r^{11} + 569r^{10} + 5196r^9 + 30327r^8 + 118064r^7 + 310235r^6 + 544884r^5 + 625540r^4 + 476164r^3 + 306608r^2 + 226776r + 104600)}{(r^2 + 2r - 1)(r^2 + 4r + 2)(r^2 + 6r + 7)(r^2 + 8r + 14)(r^2 + 10r + 23)(r^2 + 12r + 34)(r^2 + 14r + 47)}$$

Which for the root $r = -\sqrt{2}$ becomes

$$a_7 = -\frac{\sqrt{2}(-77567 + 54843\sqrt{2})}{126(-1 + 2\sqrt{2})(\sqrt{2} - 1)(-3 + 2\sqrt{2})(-2 + \sqrt{2})(-5 + 2\sqrt{2})(-3 + \sqrt{2})(-7 + 2\sqrt{2})}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{r}{r^2+2r-1}$	$\frac{\sqrt{2}}{-1+2\sqrt{2}}$
a_2	$\frac{r(r^3+r^2-2r+2)}{(r^2+2r-1)(r^2+4r+2)}$	$\frac{\sqrt{2}}{-5+3\sqrt{2}}$
a_3	$\frac{2r(r^4+4r^3+3r^2+2)}{(r^2+2r-1)(r^2+4r+2)(r^2+6r+7)}$	$-\frac{1}{15-9\sqrt{2}}$
a_4	$\frac{r(r^7+10r^6+36r^5+58r^4+41r^3+20r^2+42r+40)}{(r^2+2r-1)(r^2+4r+2)(r^2+6r+7)(r^2+8r+14)}$	$-\frac{6}{-1104+780\sqrt{2}}$
a_5	$\frac{r(3r^8+48r^7+306r^6+996r^5+1749r^4+1616r^3+866r^2+680r+496)}{(r^2+2r-1)(r^2+4r+2)(r^2+6r+7)(r^2+8r+14)(r^2+10r+23)}$	$-\frac{41}{6108\sqrt{2}-8640}$
a_6	$\frac{r(r^{11}+27r^{10}+314r^9+2064r^8+8439r^7+22209r^6+37650r^5+40744r^4+30972r^3+26532r^2+26728r+13520)}{(r^2+2r-1)(r^2+4r+2)(r^2+6r+7)(r^2+8r+14)(r^2+10r+23)(r^2+12r+34)}$	$\frac{38}{1144}$
a_7	$\frac{4r(r^{12}+36r^{11}+569r^{10}+5196r^9+30327r^8+118064r^7+310235r^6+544884r^5+625540r^4+476164r^3+306608r^2+226776r+104600)}{(r^2+2r-1)(r^2+4r+2)(r^2+6r+7)(r^2+8r+14)(r^2+10r+23)(r^2+12r+34)(r^2+14r+47)}$	$-\frac{1}{12}$

Using the above table, then the solution $y_1(x)$ is

$$y_1(x) = x^{-\sqrt{2}}(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots)$$

$$= x^{-\sqrt{2}}\left(1 + \frac{\sqrt{2}x}{-1+2\sqrt{2}} + \frac{\sqrt{2}x^2}{-5+3\sqrt{2}} - \frac{2\sqrt{2}x^3}{15-9\sqrt{2}} + \frac{(-69+49\sqrt{2})x^4}{-1104+780\sqrt{2}} + \frac{(-414+293\sqrt{2})x^5}{6108\sqrt{2}-8640} + \dots\right)$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$b_1 = \frac{r}{r^2 + 2r - 1}$$

For $2 \leq n$ the recursive equation is

$$b_{n-2}(n+r-2)(n-3+r) - b_n(n+r)(n+r-1) + b_{n-1}(n+r-1) - b_n(n+r) + 2b_n = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = \frac{n^2 b_{n-2} + 2nr b_{n-2} + r^2 b_{n-2} - 5n b_{n-2} + n b_{n-1} - 5r b_{n-2} + r b_{n-1} + 6b_{n-2} - b_{n-1}}{n^2 + 2nr + r^2 - 2} \quad (4)$$

Which for the root $r = \sqrt{2}$ becomes

$$b_n = \frac{(2n b_{n-2} - 5b_{n-2} + b_{n-1}) \sqrt{2} + n^2 b_{n-2} + (-5b_{n-2} + b_{n-1}) n + 8b_{n-2} - b_{n-1}}{n(2\sqrt{2} + n)} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = \sqrt{2}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{r}{r^2+2r-1}$	$\frac{\sqrt{2}}{1+2\sqrt{2}}$

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{r(r^3 + r^2 - 2r + 2)}{(r^2 + 2r - 1)(r^2 + 4r + 2)}$$

Which for the root $r = \sqrt{2}$ becomes

$$b_2 = \frac{\sqrt{2}}{5 + 3\sqrt{2}}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{r}{r^2+2r-1}$	$\frac{\sqrt{2}}{1+2\sqrt{2}}$
b_2	$\frac{r(r^3+r^2-2r+2)}{(r^2+2r-1)(r^2+4r+2)}$	$\frac{\sqrt{2}}{5+3\sqrt{2}}$

For $n = 3$, using the above recursive equation gives

$$b_3 = \frac{2r(r^4 + 4r^3 + 3r^2 + 2)}{(r^2 + 2r - 1)(r^2 + 4r + 2)(r^2 + 6r + 7)}$$

Which for the root $r = \sqrt{2}$ becomes

$$b_3 = \frac{2\sqrt{2}}{15 + 9\sqrt{2}}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{r}{r^2+2r-1}$	$\frac{\sqrt{2}}{1+2\sqrt{2}}$
b_2	$\frac{r(r^3+r^2-2r+2)}{(r^2+2r-1)(r^2+4r+2)}$	$\frac{\sqrt{2}}{5+3\sqrt{2}}$
b_3	$\frac{2r(r^4+4r^3+3r^2+2)}{(r^2+2r-1)(r^2+4r+2)(r^2+6r+7)}$	$\frac{2\sqrt{2}}{15+9\sqrt{2}}$

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{r(r^7 + 10r^6 + 36r^5 + 58r^4 + 41r^3 + 20r^2 + 42r + 40)}{(r^2 + 2r - 1)(r^2 + 4r + 2)(r^2 + 6r + 7)(r^2 + 8r + 14)}$$

Which for the root $r = \sqrt{2}$ becomes

$$b_4 = \frac{69 + 49\sqrt{2}}{1104 + 780\sqrt{2}}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{r}{r^2+2r-1}$	$\frac{\sqrt{2}}{1+2\sqrt{2}}$
b_2	$\frac{r(r^3+r^2-2r+2)}{(r^2+2r-1)(r^2+4r+2)}$	$\frac{\sqrt{2}}{5+3\sqrt{2}}$
b_3	$\frac{2r(r^4+4r^3+3r^2+2)}{(r^2+2r-1)(r^2+4r+2)(r^2+6r+7)}$	$\frac{2\sqrt{2}}{15+9\sqrt{2}}$
b_4	$\frac{r(r^7+10r^6+36r^5+58r^4+41r^3+20r^2+42r+40)}{(r^2+2r-1)(r^2+4r+2)(r^2+6r+7)(r^2+8r+14)}$	$\frac{69+49\sqrt{2}}{1104+780\sqrt{2}}$

For $n = 5$, using the above recursive equation gives

$$b_5 = \frac{r(3r^8 + 48r^7 + 306r^6 + 996r^5 + 1749r^4 + 1616r^3 + 866r^2 + 680r + 496)}{(r^2 + 2r - 1)(r^2 + 4r + 2)(r^2 + 6r + 7)(r^2 + 8r + 14)(r^2 + 10r + 23)}$$

Which for the root $r = \sqrt{2}$ becomes

$$b_5 = \frac{414 + 293\sqrt{2}}{6108\sqrt{2} + 8640}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{r}{r^2+2r-1}$	$\frac{\sqrt{2}}{1+2\sqrt{2}}$
b_2	$\frac{r(r^3+r^2-2r+2)}{(r^2+2r-1)(r^2+4r+2)}$	$\frac{\sqrt{2}}{5+3\sqrt{2}}$
b_3	$\frac{2r(r^4+4r^3+3r^2+2)}{(r^2+2r-1)(r^2+4r+2)(r^2+6r+7)}$	$\frac{2\sqrt{2}}{15+9\sqrt{2}}$
b_4	$\frac{r(r^7+10r^6+36r^5+58r^4+41r^3+20r^2+42r+40)}{(r^2+2r-1)(r^2+4r+2)(r^2+6r+7)(r^2+8r+14)}$	$\frac{69+49\sqrt{2}}{1104+780\sqrt{2}}$
b_5	$\frac{r(3r^8+48r^7+306r^6+996r^5+1749r^4+1616r^3+866r^2+680r+496)}{(r^2+2r-1)(r^2+4r+2)(r^2+6r+7)(r^2+8r+14)(r^2+10r+23)}$	$\frac{414+293\sqrt{2}}{6108\sqrt{2}+8640}$

For $n = 6$, using the above recursive equation gives

$$b_6 = \frac{r(r^{11} + 27r^{10} + 314r^9 + 2064r^8 + 8439r^7 + 22209r^6 + 37650r^5 + 40744r^4 + 30972r^3 + 26532r^2 + 26728r + 13520)}{(r^2 + 2r - 1)(r^2 + 4r + 2)(r^2 + 6r + 7)(r^2 + 8r + 14)(r^2 + 10r + 23)(r^2 + 12r + 34)}$$

Which for the root $r = \sqrt{2}$ becomes

$$b_6 = \frac{3898 + 2757\sqrt{2}}{114408 + 80892\sqrt{2}}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{r}{r^2+2r-1}$	$\frac{\sqrt{2}}{1+2\sqrt{2}}$
b_2	$\frac{r(r^3+r^2-2r+2)}{(r^2+2r-1)(r^2+4r+2)}$	$\frac{\sqrt{2}}{5+3\sqrt{2}}$
b_3	$\frac{2r(r^4+4r^3+3r^2+2)}{(r^2+2r-1)(r^2+4r+2)(r^2+6r+7)}$	$\frac{2\sqrt{2}}{15+9\sqrt{2}}$
b_4	$\frac{r(r^7+10r^6+36r^5+58r^4+41r^3+20r^2+42r+40)}{(r^2+2r-1)(r^2+4r+2)(r^2+6r+7)(r^2+8r+14)}$	$\frac{69+49\sqrt{2}}{1104+780\sqrt{2}}$
b_5	$\frac{r(3r^8+48r^7+306r^6+996r^5+1749r^4+1616r^3+866r^2+680r+496)}{(r^2+2r-1)(r^2+4r+2)(r^2+6r+7)(r^2+8r+14)(r^2+10r+23)}$	$\frac{414+293\sqrt{2}}{6108\sqrt{2}+8640}$
b_6	$\frac{r(r^{11}+27r^{10}+314r^9+2064r^8+8439r^7+22209r^6+37650r^5+40744r^4+30972r^3+26532r^2+26728r+13520)}{(r^2+2r-1)(r^2+4r+2)(r^2+6r+7)(r^2+8r+14)(r^2+10r+23)(r^2+12r+34)}$	$\frac{3898+2757\sqrt{2}}{114408+80892\sqrt{2}}$

For $n = 7$, using the above recursive equation gives

$$b_7 = \frac{4r(r^{12} + 36r^{11} + 569r^{10} + 5196r^9 + 30327r^8 + 118064r^7 + 310235r^6 + 544884r^5 + 625540r^4 + 476164r^3 + 306608r^2 + 226776r + 104600)}{(r^2 + 2r - 1)(r^2 + 4r + 2)(r^2 + 6r + 7)(r^2 + 8r + 14)(r^2 + 10r + 23)(r^2 + 12r + 34)(r^2 + 14r + 47)}$$

Which for the root $r = \sqrt{2}$ becomes

$$b_7 = \frac{\sqrt{2} (77567 + 54843\sqrt{2})}{126 (1 + 2\sqrt{2}) (1 + \sqrt{2}) (3 + 2\sqrt{2}) (2 + \sqrt{2}) (5 + 2\sqrt{2}) (3 + \sqrt{2}) (7 + 2\sqrt{2})}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{r}{r^2+2r-1}$	$\frac{\sqrt{2}}{1+2\sqrt{2}}$
b_2	$\frac{r(r^3+r^2-2r+2)}{(r^2+2r-1)(r^2+4r+2)}$	$\frac{\sqrt{2}}{5+3\sqrt{2}}$
b_3	$\frac{2r(r^4+4r^3+3r^2+2)}{(r^2+2r-1)(r^2+4r+2)(r^2+6r+7)}$	$\frac{2\sqrt{2}}{15+9\sqrt{2}}$
b_4	$\frac{r(r^7+10r^6+36r^5+58r^4+41r^3+20r^2+42r+40)}{(r^2+2r-1)(r^2+4r+2)(r^2+6r+7)(r^2+8r+14)}$	$\frac{69+49\sqrt{2}}{1104}$
b_5	$\frac{r(3r^8+48r^7+306r^6+996r^5+1749r^4+1616r^3+866r^2+680r+496)}{(r^2+2r-1)(r^2+4r+2)(r^2+6r+7)(r^2+8r+14)(r^2+10r+23)}$	$\frac{414+293\sqrt{2}}{6108}$
b_6	$\frac{r(r^{11}+27r^{10}+314r^9+2064r^8+8439r^7+22209r^6+37650r^5+40744r^4+30972r^3+26532r^2+26728r+13520)}{(r^2+2r-1)(r^2+4r+2)(r^2+6r+7)(r^2+8r+14)(r^2+10r+23)(r^2+12r+34)}$	$\frac{3898+2937\sqrt{2}}{114408}$
b_7	$\frac{4r(r^{12}+36r^{11}+569r^{10}+5196r^9+30327r^8+118064r^7+310235r^6+544884r^5+625540r^4+476164r^3+306608r^2+226776r+104600)}{(r^2+2r-1)(r^2+4r+2)(r^2+6r+7)(r^2+8r+14)(r^2+10r+23)(r^2+12r+34)(r^2+14r+47)}$	$\frac{3898+2937\sqrt{2}}{126(1+2\sqrt{2})(1+\sqrt{2})(3+2\sqrt{2})(2+\sqrt{2})(5+2\sqrt{2})(3+\sqrt{2})(7+2\sqrt{2})}$

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned} y_2(x) &= x^{-\sqrt{2}}(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 + b_7x^7 + b_8x^8 \dots) \\ &= x^{\sqrt{2}} \left(1 + \frac{\sqrt{2}x}{1+2\sqrt{2}} + \frac{\sqrt{2}x^2}{5+3\sqrt{2}} + \frac{2\sqrt{2}x^3}{15+9\sqrt{2}} + \frac{(69+49\sqrt{2})x^4}{1104+780\sqrt{2}} + \frac{(414+293\sqrt{2})x^5}{6108\sqrt{2}+8640} + \frac{(3898+2937\sqrt{2})x^6}{114408+8064\sqrt{2}} \right) \end{aligned}$$

Therefore the homogeneous solution is

$$y_h(x) = c_1y_1(x) + c_2y_2(x)$$

$$\begin{aligned}
&= c_1 x^{-\sqrt{2}} \left(1 + \frac{\sqrt{2}x}{-1+2\sqrt{2}} + \frac{\sqrt{2}x^2}{-5+3\sqrt{2}} - \frac{2\sqrt{2}x^3}{15-9\sqrt{2}} + \frac{(-69+49\sqrt{2})x^4}{-1104+780\sqrt{2}} \right. \\
&\quad \left. + \frac{(-414+293\sqrt{2})x^5}{6108\sqrt{2}-8640} + \frac{(3898-2757\sqrt{2})x^6}{114408-80892\sqrt{2}} \right. \\
&\quad \left. - \frac{\sqrt{2}(-77567+54843\sqrt{2})x^7}{126(-1+2\sqrt{2})(\sqrt{2}-1)(-3+2\sqrt{2})(-2+\sqrt{2})(-5+2\sqrt{2})(-3+\sqrt{2})(-7+2\sqrt{2})} \right. \\
&\quad \left. + O(x^8) \right) + c_2 x^{\sqrt{2}} \left(1 + \frac{\sqrt{2}x}{1+2\sqrt{2}} + \frac{\sqrt{2}x^2}{5+3\sqrt{2}} + \frac{2\sqrt{2}x^3}{15+9\sqrt{2}} + \frac{(69+49\sqrt{2})x^4}{1104+780\sqrt{2}} \right. \\
&\quad \left. + \frac{(414+293\sqrt{2})x^5}{6108\sqrt{2}+8640} + \frac{(3898+2757\sqrt{2})x^6}{114408+80892\sqrt{2}} \right. \\
&\quad \left. + \frac{\sqrt{2}(77567+54843\sqrt{2})x^7}{126(1+2\sqrt{2})(1+\sqrt{2})(3+2\sqrt{2})(2+\sqrt{2})(5+2\sqrt{2})(3+\sqrt{2})(7+2\sqrt{2})} \right. \\
&\quad \left. + O(x^8) \right)
\end{aligned}$$

Hence the final solution is

$$y = y_h$$

$$\begin{aligned}
&= c_1 x^{-\sqrt{2}} \left(1 + \frac{\sqrt{2}x}{-1+2\sqrt{2}} + \frac{\sqrt{2}x^2}{-5+3\sqrt{2}} - \frac{2\sqrt{2}x^3}{15-9\sqrt{2}} + \frac{(-69+49\sqrt{2})x^4}{-1104+780\sqrt{2}} \right. \\
&\quad \left. + \frac{(-414+293\sqrt{2})x^5}{6108\sqrt{2}-8640} + \frac{(3898-2757\sqrt{2})x^6}{114408-80892\sqrt{2}} \right. \\
&\quad \left. - \frac{\sqrt{2}(-77567+54843\sqrt{2})x^7}{126(-1+2\sqrt{2})(\sqrt{2}-1)(-3+2\sqrt{2})(-2+\sqrt{2})(-5+2\sqrt{2})(-3+\sqrt{2})(-7+2\sqrt{2})} \right. \\
&\quad \left. + O(x^8) \right) + c_2 x^{\sqrt{2}} \left(1 + \frac{\sqrt{2}x}{1+2\sqrt{2}} + \frac{\sqrt{2}x^2}{5+3\sqrt{2}} + \frac{2\sqrt{2}x^3}{15+9\sqrt{2}} + \frac{(69+49\sqrt{2})x^4}{1104+780\sqrt{2}} \right. \\
&\quad \left. + \frac{(414+293\sqrt{2})x^5}{6108\sqrt{2}+8640} + \frac{(3898+2757\sqrt{2})x^6}{114408+80892\sqrt{2}} \right. \\
&\quad \left. + \frac{\sqrt{2}(77567+54843\sqrt{2})x^7}{126(1+2\sqrt{2})(1+\sqrt{2})(3+2\sqrt{2})(2+\sqrt{2})(5+2\sqrt{2})(3+\sqrt{2})(7+2\sqrt{2})} \right. \\
&\quad \left. + O(x^8) \right)
\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 x^{-\sqrt{2}} \left(1 + \frac{\sqrt{2}x}{-1+2\sqrt{2}} + \frac{\sqrt{2}x^2}{-5+3\sqrt{2}} - \frac{2\sqrt{2}x^3}{15-9\sqrt{2}} + \frac{(-69+49\sqrt{2})x^4}{-1104+780\sqrt{2}} \right. \\ \left. + \frac{(-414+293\sqrt{2})x^5}{6108\sqrt{2}-8640} + \frac{(3898-2757\sqrt{2})x^6}{114408-80892\sqrt{2}} \right. \\ \left. - \frac{\sqrt{2}(-77567+54843\sqrt{2})x^7}{126(-1+2\sqrt{2})(\sqrt{2}-1)(-3+2\sqrt{2})(-2+\sqrt{2})(-5+2\sqrt{2})(-3+\sqrt{2})(-7+2\sqrt{2})} \right. \\ \left. + O(x^8) \right) + c_2 x^{\sqrt{2}} \left(1 + \frac{\sqrt{2}x}{1+2\sqrt{2}} + \frac{\sqrt{2}x^2}{5+3\sqrt{2}} + \frac{2\sqrt{2}x^3}{15+9\sqrt{2}} + \frac{(69+49\sqrt{2})x^4}{1104+780\sqrt{2}} \right. \\ \left. + \frac{(414+293\sqrt{2})x^5}{6108\sqrt{2}+8640} + \frac{(3898+2757\sqrt{2})x^6}{114408+80892\sqrt{2}} \right. \\ \left. + \frac{\sqrt{2}(77567+54843\sqrt{2})x^7}{126(1+2\sqrt{2})(1+\sqrt{2})(3+2\sqrt{2})(2+\sqrt{2})(5+2\sqrt{2})(3+\sqrt{2})(7+2\sqrt{2})} \right. \\ \left. + O(x^8) \right)$$

Verification of solutions

$$y = c_1 x^{-\sqrt{2}} \left(1 + \frac{\sqrt{2}x}{-1+2\sqrt{2}} + \frac{\sqrt{2}x^2}{-5+3\sqrt{2}} - \frac{2\sqrt{2}x^3}{15-9\sqrt{2}} + \frac{(-69+49\sqrt{2})x^4}{-1104+780\sqrt{2}} \right. \\ \left. + \frac{(-414+293\sqrt{2})x^5}{6108\sqrt{2}-8640} + \frac{(3898-2757\sqrt{2})x^6}{114408-80892\sqrt{2}} \right. \\ \left. - \frac{\sqrt{2}(-77567+54843\sqrt{2})x^7}{126(-1+2\sqrt{2})(\sqrt{2}-1)(-3+2\sqrt{2})(-2+\sqrt{2})(-5+2\sqrt{2})(-3+\sqrt{2})(-7+2\sqrt{2})} \right. \\ \left. + O(x^8) \right) + c_2 x^{\sqrt{2}} \left(1 + \frac{\sqrt{2}x}{1+2\sqrt{2}} + \frac{\sqrt{2}x^2}{5+3\sqrt{2}} + \frac{2\sqrt{2}x^3}{15+9\sqrt{2}} + \frac{(69+49\sqrt{2})x^4}{1104+780\sqrt{2}} \right. \\ \left. + \frac{(414+293\sqrt{2})x^5}{6108\sqrt{2}+8640} + \frac{(3898+2757\sqrt{2})x^6}{114408+80892\sqrt{2}} \right. \\ \left. + \frac{\sqrt{2}(77567+54843\sqrt{2})x^7}{126(1+2\sqrt{2})(1+\sqrt{2})(3+2\sqrt{2})(2+\sqrt{2})(5+2\sqrt{2})(3+\sqrt{2})(7+2\sqrt{2})} \right. \\ \left. + O(x^8) \right)$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
  -> Bessel
  -> elliptic
  -> Legendre
  -> Whittaker
    -> hyper3: Equivalence to 1F1 under a power @ Moebius
  -> hypergeometric
    -> heuristic approach
    -> hyper3: Equivalence to 2F1, 1F1 or 0F1 under a power @ Moebius
  -> Mathieu
    -> Equivalence to the rational form of Mathieu ODE under a power @ Moebius
trying a solution in terms of MeijerG functions
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
<- Heun successful: received ODE is equivalent to the HeunG ODE, case a <> 0, e <> 0, g <
```

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 654

Order:=8;

dsolve(x^2*(x^2-1)*diff(y(x),x\$2)-x*(1-x)*diff(y(x),x)+2*y(x)=0,y(x),type='series',x=0);

$$\begin{aligned}
 y(x) = & c_1 x^{-\sqrt{2}} \left(1 + \frac{\sqrt{2}}{-1 + 2\sqrt{2}} x + \frac{\sqrt{2}}{(1 - 2\sqrt{2})(\sqrt{2} - 1)} x^2 + \frac{6\sqrt{2} - 8}{57\sqrt{2} - 81} x^3 \right. \\
 & + \frac{-49\sqrt{2} + 69}{1104 - 780\sqrt{2}} x^4 + \frac{293\sqrt{2} - 414}{6108\sqrt{2} - 8640} x^5 + \frac{-2757\sqrt{2} + 3898}{114408 - 80892\sqrt{2}} x^6 \\
 & + \frac{1}{126} \frac{77567\sqrt{2} - 109686}{(-1 + 2\sqrt{2})(\sqrt{2} - 1)(2\sqrt{2} - 3)(-2 + \sqrt{2})(-5 + 2\sqrt{2})(-3 + \sqrt{2})(-7 + 2\sqrt{2})} x^7 \\
 & \left. + O(x^8) \right) + c_2 x^{\sqrt{2}} \left(1 + \frac{\sqrt{2}}{1 + 2\sqrt{2}} x + \frac{\sqrt{2}}{5 + 3\sqrt{2}} x^2 + \frac{6\sqrt{2} + 8}{57\sqrt{2} + 81} x^3 \right. \\
 & + \frac{49\sqrt{2} + 69}{1104 + 780\sqrt{2}} x^4 + \frac{293\sqrt{2} + 414}{6108\sqrt{2} + 8640} x^5 + \frac{2757\sqrt{2} + 3898}{114408 + 80892\sqrt{2}} x^6 \\
 & + \frac{1}{126} \frac{77567\sqrt{2} + 109686}{(1 + 2\sqrt{2})(1 + \sqrt{2})(3 + 2\sqrt{2})(2 + \sqrt{2})(5 + 2\sqrt{2})(3 + \sqrt{2})(7 + 2\sqrt{2})} x^7 \\
 & \left. + O(x^8) \right)
 \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 10352

AsymptoticDSolveValue[x^2*(x^2-1)*y''[x]-x*(1-x)*y'[x]+2*y[x]==0,y[x],{x,0,7}]

Too large to display

19.3 problem 1(c)

Internal problem ID [6443]

Internal file name [OUTPUT/5691_Sunday_June_05_2022_03_47_13_PM_63020299/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.4. REGULAR SINGULAR POINTS. Page 175

Problem number: 1(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_ode_missing_y", "second order series method. Irregular singular point"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_y]]
```

Unable to solve or complete the solution.

$$x^2y'' + (2 - x)y' = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2y'' + (2 - x)y' = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{-2 + x}{x^2}$$
$$q(x) = 0$$

Table 441: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{-2+x}{x^2}$	
singularity	type
$x = 0$	“irregular”

$q(x) = 0$	
singularity	type

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[\infty]$

Irregular singular points : $[0]$

Since $x = 0$ is not an ordinary point, then we will now check if it is a regular singular point. Unable to solve since $x = 0$ is not regular singular point. Terminating.

Verification of solutions N/A

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
<- LODE missing y successful`

```

X Solution by Maple

```

Order:=8;
dsolve(x^2*diff(y(x),x$2)+(2-x)*diff(y(x),x)=0,y(x),type='series',x=0);

```

No solution found

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 64

```
AsymptoticDSolveValue[x^2*y''[x]+(2-x)*y'[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_2 e^{2/x} \left(\frac{2835x^7}{2} + 315x^6 + \frac{315x^5}{4} + \frac{45x^4}{2} + \frac{15x^3}{2} + 3x^2 + \frac{3x}{2} + 1 \right) x^3 + c_1$$

19.4 problem 1(d)

19.4.1 Maple step by step solution 3157

Internal problem ID [6444]

Internal file name [OUTPUT/5692_Sunday_June_05_2022_03_47_14_PM_30898659/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.4. REGULAR SINGULAR POINTS. Page 175

Problem number: 1(d).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference is integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$(3x + 1)xy'' - (1 + x)y' + 2y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$(3x^2 + x)y'' + (-1 - x)y' + 2y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{1+x}{(3x+1)x}$$
$$q(x) = \frac{2}{(3x+1)x}$$

Table 442: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{1+x}{(3x+1)x}$	
singularity	type
$x = 0$	“regular”
$x = -\frac{1}{3}$	“regular”

$q(x) = \frac{2}{(3x+1)x}$	
singularity	type
$x = 0$	“regular”
$x = -\frac{1}{3}$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0, -\frac{1}{3}, \infty]$

Irregular singular points : $[\]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$(3x + 1)xy'' + (-1 - x)y' + 2y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned} & (3x + 1)x \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\ & + (-1 - x) \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + 2 \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \end{aligned} \tag{1}$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 3x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{n+r-1} a_n (n+r) (n+r-1) \right) \\ & + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) + \sum_{n=0}^{\infty} (-(n+r) a_n x^{n+r-1}) + \left(\sum_{n=0}^{\infty} 2a_n x^{n+r} \right) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r-1$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r-1} and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=0}^{\infty} 3x^{n+r} a_n (n+r) (n+r-1) &= \sum_{n=1}^{\infty} 3a_{n-1} (n+r-1) (n+r-2) x^{n+r-1} \\ \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) &= \sum_{n=1}^{\infty} (-a_{n-1} (n+r-1) x^{n+r-1}) \\ \sum_{n=0}^{\infty} 2a_n x^{n+r} &= \sum_{n=1}^{\infty} 2a_{n-1} x^{n+r-1} \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r-1$.

$$\begin{aligned} & \left(\sum_{n=1}^{\infty} 3a_{n-1} (n+r-1) (n+r-2) x^{n+r-1} \right) \\ & + \left(\sum_{n=0}^{\infty} x^{n+r-1} a_n (n+r) (n+r-1) \right) + \sum_{n=1}^{\infty} (-a_{n-1} (n+r-1) x^{n+r-1}) \\ & + \sum_{n=0}^{\infty} (-(n+r) a_n x^{n+r-1}) + \left(\sum_{n=1}^{\infty} 2a_{n-1} x^{n+r-1} \right) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r-1} a_n (n+r) (n+r-1) - (n+r) a_n x^{n+r-1} = 0$$

When $n=0$ the above becomes

$$x^{-1+r} a_0 r (-1+r) - r a_0 x^{-1+r} = 0$$

Or

$$(x^{-1+r}r(-1+r) - rx^{-1+r})a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$rx^{-1+r}(-2+r) = 0$$

Since the above is true for all x then the indicial equation becomes

$$r(-2+r) = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = 2$$

$$r_2 = 0$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$rx^{-1+r}(-2+r) = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = 2$ is an integer, then we can construct two linearly independent solutions

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = Cy_1(x) \ln(x) + x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = x^2 \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+2}$$

$$y_2(x) = Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Where C above can be zero. We start by finding y_1 . Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$3a_{n-1}(n+r-1)(n+r-2) + a_n(n+r)(n+r-1) - a_{n-1}(n+r-1) - a_n(n+r) + 2a_{n-1} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{a_{n-1}(3n^2 + 6nr + 3r^2 - 10n - 10r + 9)}{n^2 + 2nr + r^2 - 2n - 2r} \quad (4)$$

Which for the root $r = 2$ becomes

$$a_n = -\frac{a_{n-1}(3n^2 + 2n + 1)}{n(n+2)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 2$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = \frac{-3r^2 + 4r - 2}{r^2 - 1}$$

Which for the root $r = 2$ becomes

$$a_1 = -2$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-3r^2+4r-2}{r^2-1}$	-2

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{9r^4 - 6r^3 + r^2 + 2}{(r^2 - 1)r(r+2)}$$

Which for the root $r = 2$ becomes

$$a_2 = \frac{17}{4}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-3r^2+4r-2}{r^2-1}$	-2
a_2	$\frac{9r^4-6r^3+r^2+2}{(r^2-1)r(r+2)}$	$\frac{17}{4}$

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{-27r^6 - 54r^5 - 9r^4 + 28r^3 - 12r^2 - 16r - 12}{r(-1+r)(r+2)(r+1)^2(r+3)}$$

Which for the root $r = 2$ becomes

$$a_3 = -\frac{289}{30}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-3r^2+4r-2}{r^2-1}$	-2
a_2	$\frac{9r^4-6r^3+r^2+2}{(r^2-1)r(r+2)}$	$\frac{17}{4}$
a_3	$\frac{-27r^6-54r^5-9r^4+28r^3-12r^2-16r-12}{r(-1+r)(r+2)(r+1)^2(r+3)}$	$-\frac{289}{30}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{81r^8 + 540r^7 + 1242r^6 + 960r^5 - 203r^4 - 260r^3 + 464r^2 + 440r + 204}{r(-1+r)(r+2)^2(r+1)^2(r+3)(r+4)}$$

Which for the root $r = 2$ becomes

$$a_4 = \frac{5491}{240}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-3r^2+4r-2}{r^2-1}$	-2
a_2	$\frac{9r^4-6r^3+r^2+2}{(r^2-1)r(r+2)}$	$\frac{17}{4}$
a_3	$\frac{-27r^6-54r^5-9r^4+28r^3-12r^2-16r-12}{r(-1+r)(r+2)(r+1)^2(r+3)}$	$-\frac{289}{30}$
a_4	$\frac{81r^8+540r^7+1242r^6+960r^5-203r^4-260r^3+464r^2+440r+204}{r(-1+r)(r+2)^2(r+1)^2(r+3)(r+4)}$	$\frac{5491}{240}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{-243r^{10} - 3240r^9 - 17280r^8 - 46080r^7 - 60819r^6 - 27800r^5 + 10710r^4 - 1760r^3 - 25188r^2 - 19040r - 6936}{r(-1+r)(r+2)^2(r+1)^2(r+3)^2(r+4)(r+5)}$$

Which for the root $r = 2$ becomes

$$a_5 = -\frac{236113}{4200}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-3r^2+4r-2}{r^2-1}$	-2
a_2	$\frac{9r^4-6r^3+r^2+2}{(r^2-1)r(r+2)}$	$\frac{17}{4}$
a_3	$\frac{-27r^6-54r^5-9r^4+28r^3-12r^2-16r-12}{r(-1+r)(r+2)(r+1)^2(r+3)}$	$-\frac{289}{30}$
a_4	$\frac{81r^8+540r^7+1242r^6+960r^5-203r^4-260r^3+464r^2+440r+204}{r(-1+r)(r+2)^2(r+1)^2(r+3)(r+4)}$	$\frac{5491}{240}$
a_5	$\frac{-243r^{10}-3240r^9-17280r^8-46080r^7-60819r^6-27800r^5+10710r^4-1760r^3-25188r^2-19040r-6936}{r(-1+r)(r+2)^2(r+1)^2(r+3)^2(r+4)(r+5)}$	$-\frac{236113}{4200}$

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{(3r^2 + 26r + 57)(3r^2 - 4r + 2)(3r^2 + 8r + 6)(3r^2 + 2r + 1)(3r^2 + 14r + 17)(3r^2 + 20r + 34)}{r(-1+r)(r+2)^2(r+1)^2(r+3)^2(r+4)^2(r+5)(r+6)}$$

Which for the root $r = 2$ becomes

$$a_6 = \frac{28569673}{201600}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-3r^2+4r-2}{r^2-1}$	-2
a_2	$\frac{9r^4-6r^3+r^2+2}{(r^2-1)r(r+2)}$	$\frac{17}{4}$
a_3	$\frac{-27r^6-54r^5-9r^4+28r^3-12r^2-16r-12}{r(-1+r)(r+2)(r+1)^2(r+3)}$	$-\frac{289}{30}$
a_4	$\frac{81r^8+540r^7+1242r^6+960r^5-203r^4-260r^3+464r^2+440r+204}{r(-1+r)(r+2)^2(r+1)^2(r+3)(r+4)}$	$\frac{5491}{240}$
a_5	$\frac{-243r^{10}-3240r^9-17280r^8-46080r^7-60819r^6-27800r^5+10710r^4-1760r^3-25188r^2-19040r-6936}{r(-1+r)(r+2)^2(r+1)^2(r+3)^2(r+4)(r+5)}$	$-\frac{236113}{4200}$
a_6	$\frac{(3r^2+26r+57)(3r^2-4r+2)(3r^2+8r+6)(3r^2+2r+1)(3r^2+14r+17)(3r^2+20r+34)}{r(-1+r)(r+2)^2(r+1)^2(r+3)^2(r+4)^2(r+5)(r+6)}$	$\frac{28569673}{201600}$

For $n = 7$, using the above recursive equation gives

$$a_7 = -\frac{(3r^2 + 26r + 57)(3r^2 - 4r + 2)(3r^2 + 8r + 6)(3r^2 + 2r + 1)(3r^2 + 14r + 17)(3r^2 + 20r + 34)(3r^2 + 32r + 86)}{r(-1+r)(r+2)^2(r+1)^2(r+3)^2(r+4)^2(r+5)^2(r+6)(r+7)}$$

Which for the root $r = 2$ becomes

$$a_7 = -\frac{28569673}{78400}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-3r^2+4r-2}{r^2-1}$	-2
a_2	$\frac{9r^4-6r^3+r^2+2}{(r^2-1)r(r+2)}$	$\frac{17}{4}$
a_3	$\frac{-27r^6-54r^5-9r^4+28r^3-12r^2-16r-12}{r(-1+r)(r+2)(r+1)^2(r+3)}$	$-\frac{289}{30}$
a_4	$\frac{81r^8+540r^7+1242r^6+960r^5-203r^4-260r^3+464r^2+440r+204}{r(-1+r)(r+2)^2(r+1)^2(r+3)(r+4)}$	$\frac{5491}{240}$
a_5	$\frac{-243r^{10}-3240r^9-17280r^8-46080r^7-60819r^6-27800r^5+10710r^4-1760r^3-25188r^2-19040r-6936}{r(-1+r)(r+2)^2(r+1)^2(r+3)^2(r+4)(r+5)}$	$-\frac{236113}{4200}$
a_6	$\frac{(3r^2+26r+57)(3r^2-4r+2)(3r^2+8r+6)(3r^2+2r+1)(3r^2+14r+17)(3r^2+20r+34)}{r(-1+r)(r+2)^2(r+1)^2(r+3)^2(r+4)^2(r+5)(r+6)}$	$\frac{28569673}{201600}$
a_7	$-\frac{(3r^2+26r+57)(3r^2-4r+2)(3r^2+8r+6)(3r^2+2r+1)(3r^2+14r+17)(3r^2+20r+34)(3r^2+32r+86)}{r(-1+r)(r+2)^2(r+1)^2(r+3)^2(r+4)^2(r+5)^2(r+6)(r+7)}$	$-\frac{28569673}{78400}$

Using the above table, then the solution $y_1(x)$ is

$$y_1(x) = x^2(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots)$$

$$= x^2\left(1 - 2x + \frac{17x^2}{4} - \frac{289x^3}{30} + \frac{5491x^4}{240} - \frac{236113x^5}{4200} + \frac{28569673x^6}{201600} - \frac{28569673x^7}{78400} + O(x^8)\right)$$

Now the second solution $y_2(x)$ is found. Let

$$r_1 - r_2 = N$$

Where N is positive integer which is the difference between the two roots. r_1 is taken as the larger root. Hence for this problem we have $N = 2$. Now we need to determine if C is zero or not. This is done by finding $\lim_{r \rightarrow r_2} a_2(r)$. If this limit exists, then $C = 0$, else we need to keep the log term and $C \neq 0$. The above table shows that

$$a_N = a_2$$

$$= \frac{9r^4 - 6r^3 + r^2 + 2}{(r^2 - 1)r(r + 2)}$$

Therefore

$$\lim_{r \rightarrow r_2} \frac{9r^4 - 6r^3 + r^2 + 2}{(r^2 - 1)r(r + 2)} = \lim_{r \rightarrow 0} \frac{9r^4 - 6r^3 + r^2 + 2}{(r^2 - 1)r(r + 2)}$$

$$= \text{undefined}$$

Since the limit does not exist then the log term is needed. Therefore the second solution has the form

$$y_2(x) = Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2}\right)$$

Therefore

$$\frac{d}{dx} y_2(x) = Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2}(n+r_2)}{x}\right)$$

$$= Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} x^{-1+n+r_2} b_n (n+r_2)\right)$$

$$\frac{d^2}{dx^2} y_2(x) = Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2}$$

$$+ \sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2}(n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2}(n+r_2)}{x^2}\right)$$

$$= Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} + \left(\sum_{n=0}^{\infty} x^{-2+n+r_2} b_n (n+r_2) (-1+n+r_2)\right)$$

Substituting these back into the given ode $(3x + 1)xy'' + (-1 - x)y' + 2y = 0$ gives

$$\begin{aligned}
 & (3x + 1)x \left(Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} \right. \\
 & \left. + \sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2}(n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2}(n+r_2)}{x^2} \right) \right) \\
 & + (-1 - x) \left(Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2}(n+r_2)}{x} \right) \right) \\
 & + 2Cy_1(x) \ln(x) + 2 \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) = 0
 \end{aligned}$$

Which can be written as

$$\begin{aligned}
 & \left(((3x + 1)xy_1''(x) + (-1 - x)y_1'(x) + 2y_1(x)) \ln(x) \right. \\
 & \left. + (3x + 1)x \left(\frac{2y_1'(x)}{x} - \frac{y_1(x)}{x^2} \right) + \frac{(-1 - x)y_1(x)}{x} \right) C \\
 & + (3x + 1)x \left(\sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2}(n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2}(n+r_2)}{x^2} \right) \right) \\
 & + (-1 - x) \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2}(n+r_2)}{x} \right) + 2 \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) = 0
 \end{aligned} \tag{7}$$

But since $y_1(x)$ is a solution to the ode, then

$$(3x + 1)xy_1''(x) + (-1 - x)y_1'(x) + 2y_1(x) = 0$$

Eq (7) simplifies to

$$\begin{aligned}
 & \left((3x + 1)x \left(\frac{2y_1'(x)}{x} - \frac{y_1(x)}{x^2} \right) + \frac{(-1 - x)y_1(x)}{x} \right) C \\
 & + (3x + 1)x \left(\sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2}(n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2}(n+r_2)}{x^2} \right) \right) \\
 & + (-1 - x) \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2}(n+r_2)}{x} \right) + 2 \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) = 0
 \end{aligned} \tag{8}$$

Substituting $y_1 = \sum_{n=0}^{\infty} a_n x^{n+r_1}$ into the above gives

$$\begin{aligned} & \frac{\left(6x\left(x + \frac{1}{3}\right) \left(\sum_{n=0}^{\infty} x^{-1+n+r_1} a_n(n+r_1)\right) + 2(-1-2x) \left(\sum_{n=0}^{\infty} a_n x^{n+r_1}\right)\right) C}{x} \\ & + \frac{(3x^3 + x^2) \left(\sum_{n=0}^{\infty} x^{-2+n+r_2} b_n(n+r_2)(-1+n+r_2)\right) + (-x^2 - x) \left(\sum_{n=0}^{\infty} x^{-1+n+r_2} b_n(n+r_2)\right) + 2\left(\sum_{n=0}^{\infty} b_n x^n\right)}{x} \\ & = 0 \end{aligned} \tag{9}$$

Since $r_1 = 2$ and $r_2 = 0$ then the above becomes

$$\begin{aligned} & \frac{\left(6x\left(x + \frac{1}{3}\right) \left(\sum_{n=0}^{\infty} x^{1+n} a_n(n+2)\right) + 2(-1-2x) \left(\sum_{n=0}^{\infty} a_n x^{n+2}\right)\right) C}{x} \\ & + \frac{(3x^3 + x^2) \left(\sum_{n=0}^{\infty} x^{-2+n} b_n n(n-1)\right) + (-x^2 - x) \left(\sum_{n=0}^{\infty} x^{n-1} b_n n\right) + 2\left(\sum_{n=0}^{\infty} b_n x^n\right) x}{x} = 0 \end{aligned} \tag{10}$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 6C x^{n+2} a_n(n+2)\right) + \left(\sum_{n=0}^{\infty} 2C x^{1+n} a_n(n+2)\right) + \sum_{n=0}^{\infty} (-2C x^{1+n} a_n) \\ & + \sum_{n=0}^{\infty} (-4C x^{n+2} a_n) + \left(\sum_{n=0}^{\infty} 3x^n b_n n(n-1)\right) + \left(\sum_{n=0}^{\infty} n x^{n-1} b_n(n-1)\right) \\ & + \sum_{n=0}^{\infty} (-x^n b_n n) + \sum_{n=0}^{\infty} (-x^{n-1} b_n n) + \left(\sum_{n=0}^{\infty} 2b_n x^n\right) = 0 \end{aligned} \tag{2A}$$

The next step is to make all powers of x be $n - 1$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n-1} and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=0}^{\infty} 6C x^{n+2} a_n(n+2) &= \sum_{n=3}^{\infty} 6C a_{n-3}(n-1) x^{n-1} \\ \sum_{n=0}^{\infty} 2C x^{1+n} a_n(n+2) &= \sum_{n=2}^{\infty} 2C a_{-2+n} n x^{n-1} \end{aligned}$$

$$\begin{aligned}
\sum_{n=0}^{\infty} (-2C x^{1+n} a_n) &= \sum_{n=2}^{\infty} (-2C a_{-2+n} x^{n-1}) \\
\sum_{n=0}^{\infty} (-4C x^{n+2} a_n) &= \sum_{n=3}^{\infty} (-4C a_{n-3} x^{n-1}) \\
\sum_{n=0}^{\infty} 3x^n b_n n(n-1) &= \sum_{n=1}^{\infty} 3(n-1) b_{n-1} (-2+n) x^{n-1} \\
\sum_{n=0}^{\infty} (-x^n b_n n) &= \sum_{n=1}^{\infty} (-(n-1) b_{n-1} x^{n-1}) \\
\sum_{n=0}^{\infty} 2b_n x^n &= \sum_{n=1}^{\infty} 2b_{n-1} x^{n-1}
\end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n - 1$.

$$\begin{aligned}
&\left(\sum_{n=3}^{\infty} 6C a_{n-3} (n-1) x^{n-1} \right) + \left(\sum_{n=2}^{\infty} 2C a_{-2+n} n x^{n-1} \right) \\
&+ \sum_{n=2}^{\infty} (-2C a_{-2+n} x^{n-1}) + \sum_{n=3}^{\infty} (-4C a_{n-3} x^{n-1}) \\
&+ \left(\sum_{n=1}^{\infty} 3(n-1) b_{n-1} (-2+n) x^{n-1} \right) + \left(\sum_{n=0}^{\infty} n x^{n-1} b_n (n-1) \right) \\
&+ \sum_{n=1}^{\infty} (-(n-1) b_{n-1} x^{n-1}) + \sum_{n=0}^{\infty} (-x^{n-1} b_n n) + \left(\sum_{n=1}^{\infty} 2b_{n-1} x^{n-1} \right) = 0
\end{aligned} \tag{2B}$$

For $n = 0$ in Eq. (2B), we choose arbitray value for b_0 as $b_0 = 1$. For $n = 1$, Eq (2B) gives

$$-b_1 + 2b_0 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$-b_1 + 2 = 0$$

Solving the above for b_1 gives

$$b_1 = 2$$

For $n = N$, where $N = 2$ which is the difference between the two roots, we are free to choose $b_2 = 0$. Hence for $n = 2$, Eq (2B) gives

$$2C + 2 = 0$$

Which is solved for C . Solving for C gives

$$C = -1$$

For $n = 3$, Eq (2B) gives

$$(8a_0 + 4a_1)C + 6b_2 + 3b_3 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$3b_3 = 0$$

Solving the above for b_3 gives

$$b_3 = 0$$

For $n = 4$, Eq (2B) gives

$$(14a_1 + 6a_2)C + 17b_3 + 8b_4 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$\frac{5}{2} + 8b_4 = 0$$

Solving the above for b_4 gives

$$b_4 = -\frac{5}{16}$$

For $n = 5$, Eq (2B) gives

$$(20a_2 + 8a_3)C + 34b_4 + 15b_5 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$-\frac{2227}{120} + 15b_5 = 0$$

Solving the above for b_5 gives

$$b_5 = \frac{2227}{1800}$$

For $n = 6$, Eq (2B) gives

$$(26a_3 + 10a_4)C + 57b_5 + 24b_6 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$\frac{27659}{300} + 24b_6 = 0$$

Solving the above for b_6 gives

$$b_6 = -\frac{27659}{7200}$$

For $n = 7$, Eq (2B) gives

$$(32a_4 + 12a_5)C + 86b_6 + 35b_7 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$-\frac{9774983}{25200} + 35b_7 = 0$$

Solving the above for b_7 gives

$$b_7 = \frac{9774983}{882000}$$

Now that we found all b_n and C , we can calculate the second solution from

$$y_2(x) = Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right)$$

Using the above value found for $C = -1$ and all b_n , then the second solution becomes

$$y_2(x) = (-1) \left(x^2 \left(1 - 2x + \frac{17x^2}{4} - \frac{289x^3}{30} + \frac{5491x^4}{240} - \frac{236113x^5}{4200} + \frac{28569673x^6}{201600} - \frac{28569673x^7}{78400} + O(x^8) \right) \right) \ln(x) \\ + 1 + 2x - \frac{5x^4}{16} + \frac{2227x^5}{1800} - \frac{27659x^6}{7200} + \frac{9774983x^7}{882000} + O(x^8)$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 y_1(x) + c_2 y_2(x)$$

$$= c_1 x^2 \left(1 - 2x + \frac{17x^2}{4} - \frac{289x^3}{30} + \frac{5491x^4}{240} - \frac{236113x^5}{4200} + \frac{28569673x^6}{201600} - \frac{28569673x^7}{78400} + O(x^8) \right) \\ + c_2 \left((-1) \left(x^2 \left(1 - 2x + \frac{17x^2}{4} - \frac{289x^3}{30} + \frac{5491x^4}{240} - \frac{236113x^5}{4200} + \frac{28569673x^6}{201600} - \frac{28569673x^7}{78400} + O(x^8) \right) \right) \right. \\ \left. + 1 + 2x - \frac{5x^4}{16} + \frac{2227x^5}{1800} - \frac{27659x^6}{7200} + \frac{9774983x^7}{882000} + O(x^8) \right)$$

Hence the final solution is

$$y = y_h$$

$$= c_1 x^2 \left(1 - 2x + \frac{17x^2}{4} - \frac{289x^3}{30} + \frac{5491x^4}{240} - \frac{236113x^5}{4200} + \frac{28569673x^6}{201600} - \frac{28569673x^7}{78400} + O(x^8) \right) \\ + c_2 \left(-x^2 \left(1 - 2x + \frac{17x^2}{4} - \frac{289x^3}{30} + \frac{5491x^4}{240} - \frac{236113x^5}{4200} + \frac{28569673x^6}{201600} - \frac{28569673x^7}{78400} + O(x^8) \right) \ln(x) + 1 + 2x - \frac{5x^4}{16} + \frac{2227x^5}{1800} - \frac{27659x^6}{7200} + \frac{9774983x^7}{882000} + O(x^8) \right)$$

Summary

The solution(s) found are the following

$$y = c_1 x^2 \left(1 - 2x + \frac{17x^2}{4} - \frac{289x^3}{30} + \frac{5491x^4}{240} - \frac{236113x^5}{4200} + \frac{28569673x^6}{201600} - \frac{28569673x^7}{78400} + O(x^8) \right) + c_2 \left(-x^2 \left(1 - 2x + \frac{17x^2}{4} - \frac{289x^3}{30} + \frac{5491x^4}{240} - \frac{236113x^5}{4200} + \frac{28569673x^6}{201600} - \frac{28569673x^7}{78400} + O(x^8) \right) \ln(x) + 1 + 2x - \frac{5x^4}{16} + \frac{2227x^5}{1800} - \frac{27659x^6}{7200} + \frac{9774983x^7}{882000} + O(x^8) \right)$$

Verification of solutions

$$y = c_1 x^2 \left(1 - 2x + \frac{17x^2}{4} - \frac{289x^3}{30} + \frac{5491x^4}{240} - \frac{236113x^5}{4200} + \frac{28569673x^6}{201600} - \frac{28569673x^7}{78400} + O(x^8) \right) \\ + c_2 \left(-x^2 \left(1 - 2x + \frac{17x^2}{4} - \frac{289x^3}{30} + \frac{5491x^4}{240} - \frac{236113x^5}{4200} + \frac{28569673x^6}{201600} - \frac{28569673x^7}{78400} + O(x^8) \right) \ln(x) + 1 + 2x - \frac{5x^4}{16} + \frac{2227x^5}{1800} - \frac{27659x^6}{7200} + \frac{9774983x^7}{882000} + O(x^8) \right)$$

Verified OK.

19.4.1 Maple step by step solution

Let's solve

$$(3x + 1)xy'' + (-1 - x)y' + 2y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{2y}{(3x+1)x} + \frac{(1+x)y'}{(3x+1)x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - \frac{(1+x)y'}{(3x+1)x} + \frac{2y}{(3x+1)x} = 0$$

- Check to see if x_0 is a regular singular point

- Define functions

$$\left[P_2(x) = -\frac{1+x}{(3x+1)x}, P_3(x) = \frac{2}{(3x+1)x} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = -1$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 0$$

- $x = 0$ is a regular singular point

Check to see if x_0 is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$(3x + 1)xy'' + (-1 - x)y' + 2y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y'$ to series expansion for $m = 0..1$

$$x^m \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1+m}$$

- Shift index using $k \rightarrow k + 1 - m$

$$x^m \cdot y' = \sum_{k=-1+m}^{\infty} a_{k+1-m}(k+1-m+r)x^{k+r}$$

- Convert $x^m \cdot y''$ to series expansion for $m = 1..2$

$$x^m \cdot y'' = \sum_{k=0}^{\infty} a_k(k+r)(k+r-1)x^{k+r-2+m}$$

- Shift index using $k \rightarrow k + 2 - m$

$$x^m \cdot y'' = \sum_{k=-2+m}^{\infty} a_{k+2-m}(k+2-m+r)(k+1-m+r)x^{k+r}$$

Rewrite ODE with series expansions

$$a_0r(-2+r)x^{-1+r} + \left(\sum_{k=0}^{\infty} (a_{k+1}(k+1+r)(k+r-1) + a_k(3k^2 + 6kr + 3r^2 - 4k - 4r + 2)) \right) x^k$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$r(-2+r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \{0, 2\}$$

- Each term in the series must be 0, giving the recursion relation

$$(3k^2 + (6r-4)k + 3r^2 - 4r + 2)a_k + a_{k+1}(k+1+r)(k+r-1) = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+1} = -\frac{(3k^2 + 6kr + 3r^2 - 4k - 4r + 2)a_k}{(k+1+r)(k+r-1)}$$

- Recursion relation for $r = 0$

$$a_{k+1} = -\frac{(3k^2 - 4k + 2)a_k}{(k+1)(k-1)}$$

- Series not valid for $r = 0$, division by 0 in the recursion relation at $k = 1$

$$a_{k+1} = -\frac{(3k^2 - 4k + 2)a_k}{(k+1)(k-1)}$$

- Recursion relation for $r = 2$

$$a_{k+1} = -\frac{(3k^2 + 8k + 6)a_k}{(k+3)(k+1)}$$

- Solution for $r = 2$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+2}, a_{k+1} = -\frac{(3k^2 + 8k + 6)a_k}{(k+3)(k+1)} \right]$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
  -> Bessel
  -> elliptic
  -> Legendre
  -> Kummer
    -> hyper3: Equivalence to 1F1 under a power @ Moebius
  -> hypergeometric
    -> heuristic approach
    <- heuristic approach successful
    -> solution has integrals; searching for one without integrals...
      -> hyper3: Equivalence to 2F1, 1F1 or 0F1 under a power @ Moebius
      <- hyper3 successful: received ODE is equivalent to the 2F1 ODE
      <- hypergeometric solution without integrals successful
    <- hypergeometric successful
<- special function solution successful`
```

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 72

Order:=8;

dsolve((3*x+1)*x*diff(y(x),x\$2)-(x+1)*diff(y(x),x)+2*y(x)=0,y(x),type='series',x=0);

$$y(x) = c_1 x^2 \left(1 - 2x + \frac{17}{4}x^2 - \frac{289}{30}x^3 + \frac{5491}{240}x^4 - \frac{236113}{4200}x^5 + \frac{28569673}{201600}x^6 - \frac{28569673}{78400}x^7 + O(x^8) \right) + c_2 \left(\ln(x) \left(2x^2 - 4x^3 + \frac{17}{2}x^4 - \frac{289}{15}x^5 + \frac{5491}{120}x^6 - \frac{236113}{2100}x^7 + O(x^8) \right) + \left(-2 - 4x + 6x^2 - 12x^3 + \frac{209}{8}x^4 - \frac{54247}{900}x^5 + \frac{521849}{3600}x^6 - \frac{158526173}{441000}x^7 + O(x^8) \right) \right)$$

✓ Solution by Mathematica

Time used: 0.421 (sec). Leaf size: 118

AsymptoticDSolveValue[(3*x+1)*x*y'[x]-(x+1)*y'[x]+2*y[x]==0,y[x],{x,0,7}]

$$y(x) \rightarrow c_1 \left(\frac{27353x^6 - 12886x^5 + 6525x^4 - 3600x^3 + 1800x^2 + 7200x + 3600}{3600} - \frac{1}{240}x^2(5491x^4 - 2312x^3 + 1020x^2 - 480x + 240) \log(x) \right) + c_2 \left(\frac{28569673x^8}{201600} - \frac{236113x^7}{4200} + \frac{5491x^6}{240} - \frac{289x^5}{30} + \frac{17x^4}{4} - 2x^3 + x^2 \right)$$

19.5 problem 2(a)

Internal problem ID [6445]

Internal file name [OUTPUT/5693_Sunday_June_05_2022_03_47_19_PM_4563542/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.4. REGULAR SINGULAR POINTS. Page 175

Problem number: 2(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second order series method. Ordinary point", "second order series method. Taylor series method"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' + \sin(x)y = 0$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using Taylor series gives

$$\begin{aligned} y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \cdots \\ &= y_0 + xy'_0 + \frac{x^2}{2}f|_{x_0, y_0, y'_0} + \frac{x^3}{3!}f'|_{x_0, y_0, y'_0} + \cdots \\ &= y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0} \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \quad (1)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \quad (691)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \quad (692)$$

$$\begin{aligned} \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \end{aligned} \quad (2)$$

$$\begin{aligned} \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \end{aligned} \quad (3)$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \tag{4}$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \tag{5}$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \end{aligned}$$

⋮

$$\begin{aligned} F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \tag{6}$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \tag{7}$$

To find $y(x)$ series solution around $x = 0$. Hence

$$F_0 = -\sin(x) y$$

$$\begin{aligned} F_1 &= \frac{dF_0}{dx} \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \\ &= -\sin(x) y' - y \cos(x) \end{aligned}$$

$$\begin{aligned} F_2 &= \frac{dF_1}{dx} \\ &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} y' + \frac{\partial F_1}{\partial y'} F_1 \\ &= -2 \cos(x) y' + \sin(x) y(1 + \sin(x)) \end{aligned}$$

$$\begin{aligned} F_3 &= \frac{dF_2}{dx} \\ &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} y' + \frac{\partial F_2}{\partial y'} F_2 \\ &= (\sin(x)^2 + 3 \sin(x)) y' + 4 \cos(x) y \left(\sin(x) + \frac{1}{4} \right) \end{aligned}$$

$$\begin{aligned} F_4 &= \frac{dF_3}{dx} \\ &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} y' + \frac{\partial F_3}{\partial y'} F_3 \\ &= (6 \sin(x) + 4) \cos(x) y' + y((\sin(x) + 11) \cos(x)^2 - 2 \sin(x) - 7) \end{aligned}$$

$$\begin{aligned} F_5 &= \frac{dF_4}{dx} \\ &= \frac{\partial F_4}{\partial x} + \frac{\partial F_4}{\partial y} y' + \frac{\partial F_4}{\partial y'} F_4 \\ &= ((\sin(x) + 23) \cos(x)^2 - 6 \sin(x) - 13) y' + 9 \cos(x) y \left(\cos(x)^2 - \frac{26 \sin(x)}{9} - \frac{10}{9} \right) \end{aligned}$$

$$\begin{aligned} F_6 &= \frac{dF_5}{dx} \\ &= \frac{\partial F_5}{\partial x} + \frac{\partial F_5}{\partial y} y' + \frac{\partial F_5}{\partial y'} F_5 \\ &= (12 \cos(x)^3 + (-72 \sin(x) - 18) \cos(x)) y' + y(\cos(x)^4 + (-50 \sin(x) - 59) \cos(x)^2 + 23 \sin(x) + \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and

$y'(0) = y'(0)$ gives

$$\begin{aligned} F_0 &= 0 \\ F_1 &= -y(0) \\ F_2 &= -2y'(0) \\ F_3 &= y(0) \\ F_4 &= 4y(0) + 4y'(0) \\ F_5 &= -y(0) + 10y'(0) \\ F_6 &= -26y(0) - 6y'(0) \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$\begin{aligned} y &= \left(1 - \frac{1}{6}x^3 + \frac{1}{120}x^5 + \frac{1}{180}x^6 - \frac{1}{5040}x^7 - \frac{13}{20160}x^8\right) y(0) \\ &\quad + \left(x - \frac{1}{12}x^4 + \frac{1}{180}x^6 + \frac{1}{504}x^7 - \frac{1}{6720}x^8\right) y'(0) + O(x^8) \end{aligned}$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard power series. Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$\begin{aligned} y' &= \sum_{n=1}^{\infty} n a_n x^{n-1} \\ y'' &= \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \end{aligned}$$

Substituting the above back into the ode gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = -\sin(x) \left(\sum_{n=0}^{\infty} a_n x^n \right) \quad (1)$$

Expanding $\sin(x)$ as Taylor series around $x = 0$ and keeping only the first 8 terms gives

$$\begin{aligned} \sin(x) &= x - \frac{1}{6}x^3 + \frac{1}{120}x^5 - \frac{1}{5040}x^7 + \frac{1}{362880}x^9 + \dots \\ &= x - \frac{1}{6}x^3 + \frac{1}{120}x^5 - \frac{1}{5040}x^7 + \frac{1}{362880}x^9 \end{aligned}$$

Hence the ODE in Eq (1) becomes

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \left(x - \frac{1}{6}x^3 + \frac{1}{120}x^5 - \frac{1}{5040}x^7 + \frac{1}{362880}x^9 \right) \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0$$

Expanding the second term in (1) gives

$$\begin{aligned} & \left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + x \cdot \left(\sum_{n=0}^{\infty} a_n x^n \right) - \frac{x^3}{6} \cdot \left(\sum_{n=0}^{\infty} a_n x^n \right) + \frac{x^5}{120} \\ & \cdot \left(\sum_{n=0}^{\infty} a_n x^n \right) - \frac{x^7}{5040} \cdot \left(\sum_{n=0}^{\infty} a_n x^n \right) + \frac{x^9}{362880} \cdot \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0 \end{aligned}$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \left(\sum_{n=0}^{\infty} x^{1+n} a_n \right) + \sum_{n=0}^{\infty} \left(-\frac{x^{n+3} a_n}{6} \right) \\ & + \left(\sum_{n=0}^{\infty} \frac{x^{n+5} a_n}{120} \right) + \sum_{n=0}^{\infty} \left(-\frac{x^{n+7} a_n}{5040} \right) + \left(\sum_{n=0}^{\infty} \frac{x^{n+9} a_n}{362880} \right) = 0 \end{aligned} \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2) a_{n+2} (1+n) x^n$$

$$\sum_{n=0}^{\infty} x^{1+n} a_n = \sum_{n=1}^{\infty} a_{n-1} x^n$$

$$\sum_{n=0}^{\infty} \left(-\frac{x^{n+3} a_n}{6} \right) = \sum_{n=3}^{\infty} \left(-\frac{a_{n-3} x^n}{6} \right)$$

$$\sum_{n=0}^{\infty} \frac{x^{n+5} a_n}{120} = \sum_{n=5}^{\infty} \frac{a_{n-5} x^n}{120}$$

$$\sum_{n=0}^{\infty} \left(-\frac{x^{n+7} a_n}{5040} \right) = \sum_{n=7}^{\infty} \left(-\frac{a_{n-7} x^n}{5040} \right)$$

$$\sum_{n=0}^{\infty} \frac{x^{n+9} a_n}{362880} = \sum_{n=9}^{\infty} \frac{a_{n-9} x^n}{362880}$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (1+n) x^n \right) + \left(\sum_{n=1}^{\infty} a_{n-1} x^n \right) + \sum_{n=3}^{\infty} \left(-\frac{a_{n-3} x^n}{6} \right) \\ & + \left(\sum_{n=5}^{\infty} \frac{a_{n-5} x^n}{120} \right) + \sum_{n=7}^{\infty} \left(-\frac{a_{n-7} x^n}{5040} \right) + \left(\sum_{n=9}^{\infty} \frac{a_{n-9} x^n}{362880} \right) = 0 \end{aligned} \quad (3)$$

$n = 1$ gives

$$6a_3 + a_0 = 0$$

Which after substituting earlier equations, simplifies to

$$a_3 = -\frac{a_0}{6}$$

$n = 2$ gives

$$12a_4 + a_1 = 0$$

Which after substituting earlier equations, simplifies to

$$a_4 = -\frac{a_1}{12}$$

$n = 3$ gives

$$20a_5 + a_2 - \frac{a_0}{6} = 0$$

Which after substituting earlier equations, simplifies to

$$a_5 = \frac{a_0}{120}$$

$n = 4$ gives

$$30a_6 + a_3 - \frac{a_1}{6} = 0$$

Which after substituting earlier equations, simplifies to

$$a_6 = \frac{a_0}{180} + \frac{a_1}{180}$$

$n = 5$ gives

$$42a_7 + a_4 - \frac{a_2}{6} + \frac{a_0}{120} = 0$$

Which after substituting earlier equations, simplifies to

$$a_7 = -\frac{a_0}{5040} + \frac{a_1}{504}$$

$n = 6$ gives

$$56a_8 + a_5 - \frac{a_3}{6} + \frac{a_1}{120} = 0$$

Which after substituting earlier equations, simplifies to

$$a_8 = -\frac{13a_0}{20160} - \frac{a_1}{6720}$$

$n = 7$ gives

$$72a_9 + a_6 - \frac{a_4}{6} + \frac{a_2}{120} - \frac{a_0}{5040} = 0$$

Which after substituting earlier equations, simplifies to

$$a_9 = -\frac{a_0}{13440} - \frac{7a_1}{25920}$$

For $9 \leq n$, the recurrence equation is

$$(n+2)a_{n+2}(1+n) + a_{n-1} - \frac{a_{n-3}}{6} + \frac{a_{n-5}}{120} - \frac{a_{n-7}}{5040} + \frac{a_{n-9}}{362880} = 0 \quad (4)$$

Solving for a_{n+2} , gives

$$\begin{aligned} a_{n+2} &= -\frac{362880a_{n-1} - 60480a_{n-3} + 3024a_{n-5} - 72a_{n-7} + a_{n-9}}{362880(n+2)(1+n)} \\ (5) \quad &= -\frac{a_{n-9}}{362880(n+2)(1+n)} + \frac{a_{n-7}}{5040(n+2)(1+n)} \\ &\quad - \frac{a_{n-5}}{120(n+2)(1+n)} + \frac{a_{n-3}}{6(n+2)(1+n)} - \frac{a_{n-1}}{(n+2)(1+n)} \end{aligned}$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + a_1 x - \frac{a_0 x^3}{6} - \frac{a_1 x^4}{12} + \frac{a_0 x^5}{120} + \left(\frac{a_0}{180} + \frac{a_1}{180} \right) x^6 + \left(-\frac{a_0}{5040} + \frac{a_1}{504} \right) x^7 + \dots$$

Collecting terms, the solution becomes

$$\begin{aligned} y &= \left(1 - \frac{1}{6}x^3 + \frac{1}{120}x^5 + \frac{1}{180}x^6 - \frac{1}{5040}x^7 \right) a_0 \\ &\quad + \left(x - \frac{1}{12}x^4 + \frac{1}{180}x^6 + \frac{1}{504}x^7 \right) a_1 + O(x^8) \end{aligned} \quad (3)$$

At $x = 0$ the solution above becomes

$$y = \left(1 - \frac{1}{6}x^3 + \frac{1}{120}x^5 + \frac{1}{180}x^6 - \frac{1}{5040}x^7 \right) c_1 + \left(x - \frac{1}{12}x^4 + \frac{1}{180}x^6 + \frac{1}{504}x^7 \right) c_2 + O(x^8)$$

Summary

The solution(s) found are the following

$$\begin{aligned} y &= \left(1 - \frac{1}{6}x^3 + \frac{1}{120}x^5 + \frac{1}{180}x^6 - \frac{1}{5040}x^7 - \frac{13}{20160}x^8 \right) y(0) \\ &\quad + \left(x - \frac{1}{12}x^4 + \frac{1}{180}x^6 + \frac{1}{504}x^7 - \frac{1}{6720}x^8 \right) y'(0) + O(x^8) \end{aligned} \quad (1)$$

$$\begin{aligned} y &= \left(1 - \frac{1}{6}x^3 + \frac{1}{120}x^5 + \frac{1}{180}x^6 - \frac{1}{5040}x^7 \right) c_1 \\ &\quad + \left(x - \frac{1}{12}x^4 + \frac{1}{180}x^6 + \frac{1}{504}x^7 \right) c_2 + O(x^8) \end{aligned} \quad (2)$$

Verification of solutions

$$y = \left(1 - \frac{1}{6}x^3 + \frac{1}{120}x^5 + \frac{1}{180}x^6 - \frac{1}{5040}x^7 - \frac{13}{20160}x^8\right) y(0) \\ + \left(x - \frac{1}{12}x^4 + \frac{1}{180}x^6 + \frac{1}{504}x^7 - \frac{1}{6720}x^8\right) y'(0) + O(x^8)$$

Verified OK.

$$y = \left(1 - \frac{1}{6}x^3 + \frac{1}{120}x^5 + \frac{1}{180}x^6 - \frac{1}{5040}x^7\right) c_1 + \left(x - \frac{1}{12}x^4 + \frac{1}{180}x^6 + \frac{1}{504}x^7\right) c_2 + O(x^8)$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) * 2F1([a
-> Trying changes of variables to rationalize or make the ODE simpler
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying an equivalence, under non-integer power transformations,
to LODEs admitting Liouvillian solutions.
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
-> Bessel
-> elliptic
-> Legendre
-> Whittaker
-> hyper3: Equivalence to 1F1 under a power @ Moebius
-> hypergeometric
-> heuristic approach
-> hyper3: Equivalence to 2F1, 1F1 or 0F1 under a power @ Moebius
-> Mathieu
-> Equivalence to the rational form of Mathieu ODE under a power @ Moebius
Equivalence transformation and function parameters: {t = 1/2*t+1/2}, {kappa = -20,
<- Equivalence to the rational form of Mathieu ODE successful
<- Mathieu successful
<- special function solution successful
Change of variables used:
[x = arccos(t)]
Linear ODE actually solved:
(-t^2+1)^(1/2)*u(t)-t*diff(u(t),t)+(-t^2+1)*diff(diff(u(t),t),t) = 0
<- change of variables successful`
```


✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 49

```
Order:=8;  
dsolve(diff(y(x),x$2)+sin(x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{6}x^3 + \frac{1}{120}x^5 + \frac{1}{180}x^6 - \frac{1}{5040}x^7\right) y(0) \\ + \left(x - \frac{1}{12}x^4 + \frac{1}{180}x^6 + \frac{1}{504}x^7\right) D(y)(0) + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 63

```
AsymptoticDSolveValue[y'[x]+Sin[x]*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_2 \left(\frac{x^7}{504} + \frac{x^6}{180} - \frac{x^4}{12} + x \right) + c_1 \left(-\frac{x^7}{5040} + \frac{x^6}{180} + \frac{x^5}{120} - \frac{x^3}{6} + 1 \right)$$

19.6 problem 2(b)

Internal problem ID [6446]

Internal file name [OUTPUT/5694_Sunday_June_05_2022_03_47_23_PM_56734453/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.4. REGULAR SINGULAR POINTS. Page 175

Problem number: 2(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second order series method. Ordinary point", "second order series method. Taylor series method"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$xy'' + \sin(x)y = 0$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using Taylor series gives

$$\begin{aligned} y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \cdots \\ &= y_0 + xy'_0 + \frac{x^2}{2}f|_{x_0, y_0, y'_0} + \frac{x^3}{3!}f'|_{x_0, y_0, y'_0} + \cdots \\ &= y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0} \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \quad (1)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \quad (694)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \quad (695)$$

$$\begin{aligned} \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \end{aligned} \quad (2)$$

$$\begin{aligned} \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \end{aligned} \quad (3)$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \quad (4)$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \quad (5)$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \\ &\vdots \\ F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \quad (6)$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \quad (7)$$

To find $y(x)$ series solution around $x = 0$. Hence

$$F_0 = -\frac{\sin(x)y}{x}$$

$$\begin{aligned} F_1 &= \frac{dF_0}{dx} \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \\ &= \frac{-y' \sin(x)x - y(\cos(x)x - \sin(x))}{x^2} \end{aligned}$$

$$\begin{aligned} F_2 &= \frac{dF_1}{dx} \\ &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} y' + \frac{\partial F_1}{\partial y'} F_1 \\ &= \frac{(-2 \cos(x)x^2 + 2 \sin(x)x)y' + y(\sin(x)^2 x + (x^2 - 2) \sin(x) + 2 \cos(x)x)}{x^3} \end{aligned}$$

$$\begin{aligned} F_3 &= \frac{dF_2}{dx} \\ &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} y' + \frac{\partial F_2}{\partial y'} F_2 \\ &= \frac{(\sin(x)^2 x^2 + (3x^3 - 6x) \sin(x) + 6 \cos(x)x^2)y' + y(-4 \sin(x)^2 x + (4 \cos(x)x^2 - 3x^2 + 6) \sin(x))}{x^4} \end{aligned}$$

$$\begin{aligned} F_4 &= \frac{dF_3}{dx} \\ &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} y' + \frac{\partial F_3}{\partial y'} F_3 \\ &= \frac{2(-3 \sin(x)^2 x^2 + 3(\cos(x)x^3 - 2x^3 + 4x) \sin(x) + 2x^2 \cos(x)(x^2 - 6))y' - y(x^2 \sin(x)^3 + 3(x^3 - 6x) \sin(x) \cos(x))}{x^5} \end{aligned}$$

$$\begin{aligned} F_5 &= \frac{dF_4}{dx} \\ &= \frac{\partial F_4}{\partial x} + \frac{\partial F_4}{\partial y} y' + \frac{\partial F_4}{\partial y'} F_4 \\ &= \frac{-5 \left(-\frac{23x^2(x + \frac{\sin(x)}{23}) \cos(x)^2}{5} + \left(4x^3 + \frac{46x^2 \sin(x)}{5} - 24x \right) \cos(x) - \frac{36 \sin(x)^2 x}{5} + \left(x^4 - \frac{59}{5}x^2 + 24 \right) \sin(x) + \dots \right)}{x^6} \end{aligned}$$

$$\begin{aligned} F_6 &= \frac{dF_5}{dx} \\ &= \frac{\partial F_5}{\partial x} + \frac{\partial F_5}{\partial y} y' + \frac{\partial F_5}{\partial y'} F_5 \\ &= \frac{-6(-2x^3 \cos(x)^3 + (36x^3 + 2x^2 \sin(x)) \cos(x)^2 + ((12x^3 - 56x) \sin(x) + x^4 - 18x^2 + 120) x \cos(x) \sin(x))}{x^7} \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and $y'(0) = y'(0)$ gives

$$\begin{aligned} F_0 &= -y(0) \\ F_1 &= -y'(0) \\ F_2 &= \frac{4y(0)}{3} \\ F_3 &= 2y'(0) \\ F_4 &= -\frac{53y(0)}{15} \\ F_5 &= -\frac{19y'(0)}{3} \\ F_6 &= \frac{467y(0)}{35} \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$\begin{aligned} y &= \left(1 - \frac{1}{2}x^2 + \frac{1}{18}x^4 - \frac{53}{10800}x^6 + \frac{467}{1411200}x^8 \right) y(0) \\ &\quad + \left(x - \frac{1}{6}x^3 + \frac{1}{60}x^5 - \frac{19}{15120}x^7 \right) y'(0) + O(x^8) \end{aligned}$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard power series The ode is normalized to be

$$xy'' + \sin(x)y = 0$$

Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$\begin{aligned} y' &= \sum_{n=1}^{\infty} n a_n x^{n-1} \\ y'' &= \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \end{aligned}$$

Substituting the above back into the ode gives

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) x + \sin(x) \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0 \quad (1)$$

Expanding $\sin(x)$ as Taylor series around $x = 0$ and keeping only the first 8 terms gives

$$\begin{aligned}\sin(x) &= x - \frac{1}{6}x^3 + \frac{1}{120}x^5 - \frac{1}{5040}x^7 + \frac{1}{362880}x^9 + \dots \\ &= x - \frac{1}{6}x^3 + \frac{1}{120}x^5 - \frac{1}{5040}x^7 + \frac{1}{362880}x^9\end{aligned}$$

Hence the ODE in Eq (1) becomes

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}\right) x + \left(x - \frac{1}{6}x^3 + \frac{1}{120}x^5 - \frac{1}{5040}x^7 + \frac{1}{362880}x^9\right) \left(\sum_{n=0}^{\infty} a_n x^n\right) = 0$$

Expanding the second term in (1) gives

$$\begin{aligned}&\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}\right) x + x \cdot \left(\sum_{n=0}^{\infty} a_n x^n\right) - \frac{x^3}{6} \cdot \left(\sum_{n=0}^{\infty} a_n x^n\right) + \frac{x^5}{120} \\ &\cdot \left(\sum_{n=0}^{\infty} a_n x^n\right) - \frac{x^7}{5040} \cdot \left(\sum_{n=0}^{\infty} a_n x^n\right) + \frac{x^9}{362880} \cdot \left(\sum_{n=0}^{\infty} a_n x^n\right) = 0\end{aligned}$$

Which simplifies to

$$\begin{aligned}&\left(\sum_{n=2}^{\infty} n x^{n-1} a_n (n-1)\right) + \left(\sum_{n=0}^{\infty} x^{1+n} a_n\right) + \sum_{n=0}^{\infty} \left(-\frac{x^{n+3} a_n}{6}\right) \\ &+ \left(\sum_{n=0}^{\infty} \frac{x^{n+5} a_n}{120}\right) + \sum_{n=0}^{\infty} \left(-\frac{x^{n+7} a_n}{5040}\right) + \left(\sum_{n=0}^{\infty} \frac{x^{n+9} a_n}{362880}\right) = 0\end{aligned}\tag{2}$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\sum_{n=2}^{\infty} n x^{n-1} a_n (n-1) = \sum_{n=1}^{\infty} (1+n) a_{1+n} n x^n$$

$$\sum_{n=0}^{\infty} x^{1+n} a_n = \sum_{n=1}^{\infty} a_{n-1} x^n$$

$$\sum_{n=0}^{\infty} \left(-\frac{x^{n+3} a_n}{6}\right) = \sum_{n=3}^{\infty} \left(-\frac{a_{n-3} x^n}{6}\right)$$

$$\sum_{n=0}^{\infty} \frac{x^{n+5} a_n}{120} = \sum_{n=5}^{\infty} \frac{a_{n-5} x^n}{120}$$

$$\sum_{n=0}^{\infty} \left(-\frac{x^{n+7} a_n}{5040} \right) = \sum_{n=7}^{\infty} \left(-\frac{a_{n-7} x^n}{5040} \right)$$

$$\sum_{n=0}^{\infty} \frac{x^{n+9} a_n}{362880} = \sum_{n=9}^{\infty} \frac{a_{n-9} x^n}{362880}$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=1}^{\infty} (1+n) a_{1+n} n x^n \right) + \left(\sum_{n=1}^{\infty} a_{n-1} x^n \right) + \sum_{n=3}^{\infty} \left(-\frac{a_{n-3} x^n}{6} \right) \quad (3)$$

$$+ \left(\sum_{n=5}^{\infty} \frac{a_{n-5} x^n}{120} \right) + \sum_{n=7}^{\infty} \left(-\frac{a_{n-7} x^n}{5040} \right) + \left(\sum_{n=9}^{\infty} \frac{a_{n-9} x^n}{362880} \right) = 0$$

$n = 0$ gives

$$2a_2 + a_0 = 0$$

$$a_2 = -\frac{a_0}{2}$$

$n = 1$ gives

$$6a_3 + a_1 = 0$$

Which after substituting earlier equations, simplifies to

$$a_3 = -\frac{a_1}{6}$$

$n = 2$ gives

$$12a_4 + a_2 - \frac{a_0}{6} = 0$$

Which after substituting earlier equations, simplifies to

$$a_4 = \frac{a_0}{18}$$

$n = 3$ gives

$$20a_5 + a_3 - \frac{a_1}{6} = 0$$

Which after substituting earlier equations, simplifies to

$$a_5 = \frac{a_1}{60}$$

$n = 4$ gives

$$30a_6 + a_4 - \frac{a_2}{6} + \frac{a_0}{120} = 0$$

Which after substituting earlier equations, simplifies to

$$a_6 = -\frac{53a_0}{10800}$$

$n = 5$ gives

$$42a_7 + a_5 - \frac{a_3}{6} + \frac{a_1}{120} = 0$$

Which after substituting earlier equations, simplifies to

$$a_7 = -\frac{19a_1}{15120}$$

$n = 6$ gives

$$56a_8 + a_6 - \frac{a_4}{6} + \frac{a_2}{120} - \frac{a_0}{5040} = 0$$

Which after substituting earlier equations, simplifies to

$$a_8 = \frac{467a_0}{1411200}$$

For $9 \leq n$, the recurrence equation is

$$(1 + n) a_{1+n} n + a_{n-1} - \frac{a_{n-3}}{6} + \frac{a_{n-5}}{120} - \frac{a_{n-7}}{5040} + \frac{a_{n-9}}{362880} = 0 \quad (4)$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + a_1 x - \frac{1}{2} a_0 x^2 - \frac{1}{6} a_1 x^3 + \frac{1}{18} a_0 x^4 + \frac{1}{60} a_1 x^5 - \frac{53}{10800} a_0 x^6 - \frac{19}{15120} a_1 x^7 + \dots$$

Collecting terms, the solution becomes

$$y = \left(1 - \frac{1}{2}x^2 + \frac{1}{18}x^4 - \frac{53}{10800}x^6\right) a_0 + \left(x - \frac{1}{6}x^3 + \frac{1}{60}x^5 - \frac{19}{15120}x^7\right) a_1 + O(x^8) \quad (3)$$

At $x = 0$ the solution above becomes

$$y = \left(1 - \frac{1}{2}x^2 + \frac{1}{18}x^4 - \frac{53}{10800}x^6\right) c_1 + \left(x - \frac{1}{6}x^3 + \frac{1}{60}x^5 - \frac{19}{15120}x^7\right) c_2 + O(x^8)$$

Summary

The solution(s) found are the following

$$\begin{aligned} y &= \left(1 - \frac{1}{2}x^2 + \frac{1}{18}x^4 - \frac{53}{10800}x^6 + \frac{467}{1411200}x^8\right) y(0) \\ &\quad + \left(x - \frac{1}{6}x^3 + \frac{1}{60}x^5 - \frac{19}{15120}x^7\right) y'(0) + O(x^8) \end{aligned} \quad (1)$$

$$y = \left(1 - \frac{1}{2}x^2 + \frac{1}{18}x^4 - \frac{53}{10800}x^6\right) c_1 + \left(x - \frac{1}{6}x^3 + \frac{1}{60}x^5 - \frac{19}{15120}x^7\right) c_2 + O(x^8) \quad (2)$$

Verification of solutions

$$\begin{aligned} y &= \left(1 - \frac{1}{2}x^2 + \frac{1}{18}x^4 - \frac{53}{10800}x^6 + \frac{467}{1411200}x^8\right) y(0) \\ &\quad + \left(x - \frac{1}{6}x^3 + \frac{1}{60}x^5 - \frac{19}{15120}x^7\right) y'(0) + O(x^8) \end{aligned}$$

Verified OK.

$$y = \left(1 - \frac{1}{2}x^2 + \frac{1}{18}x^4 - \frac{53}{10800}x^6\right) c_1 + \left(x - \frac{1}{6}x^3 + \frac{1}{60}x^5 - \frac{19}{15120}x^7\right) c_2 + O(x^8)$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) * 2F1([a
-> Trying changes of variables to rationalize or make the ODE simpler
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) * 2F1
trying a symmetry of the form [xi=0, eta=F(x)]
trying 2nd order exact linear
trying symmetries linear in x and y(x)
trying to convert to a linear ODE with constant coefficients
-> trying with_periodic_functions in the coefficients
    --- Trying Lie symmetry methods, 2nd order ---
    `, `-> Computing symmetries using: way = 5
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) * 2F1
trying a symmetry of the form [xi=0, eta=F(x)]
trying 2nd order exact linear
trying symmetries linear in x and y(x)
trying to convert to a linear ODE with constant coefficients
<- unable to find a useful change of variables
trying differential order: 2; exact nonlinear
trying symmetries linear in x and y(x)
trying to convert to a linear ODE with constant coefficients
trying to convert to an ODE of Bessel type
-> trying reduction of order to Riccati
trying Riccati sub-methods:
    trying Riccati_symmetries
    -> trying a symmetry pattern of the form [F(x)*G(y), 0]
    -> trying a symmetry pattern of the form [0, F(x)*G(y)]
    -> trying a symmetry pattern of the form [F(x),G(x)*y+H(x)]
-> trying with_periodic_functions in the coefficients
    --- Trying Lie symmetry methods, 2nd order ---
    `, `-> Computing symmetries using: way = 5`[0, y]
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 44

```
Order:=8;  
dsolve(x*diff(y(x),x$2)+sin(x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{2}x^2 + \frac{1}{18}x^4 - \frac{53}{10800}x^6\right) y(0) \\ + \left(x - \frac{1}{6}x^3 + \frac{1}{60}x^5 - \frac{19}{15120}x^7\right) D(y)(0) + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 56

```
AsymptoticDSolveValue[x*y''[x]+Sin[x]*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_2 \left(-\frac{19x^7}{15120} + \frac{x^5}{60} - \frac{x^3}{6} + x \right) + c_1 \left(-\frac{53x^6}{10800} + \frac{x^4}{18} - \frac{x^2}{2} + 1 \right)$$

19.7 problem 2(c)

Internal problem ID [6447]

Internal file name [OUTPUT/5695_Sunday_June_05_2022_03_47_26_PM_50063468/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.4. REGULAR SINGULAR POINTS. Page 175

Problem number: 2(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference is integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$x^2 y'' + \sin(x) y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2 y'' + \sin(x) y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = 0$$
$$q(x) = \frac{\sin(x)}{x^2}$$

Table 444: Table $p(x), q(x)$ singularities.

$p(x) = 0$	
singularity	type

$q(x) = \frac{\sin(x)}{x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2 y'' + \sin(x) y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) + \sin(x) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Expanding $\sin(x)$ as Taylor series around $x = 0$ and keeping only the first 8 terms gives

$$\begin{aligned} \sin(x) &= x - \frac{1}{6}x^3 + \frac{1}{120}x^5 - \frac{1}{5040}x^7 + \frac{1}{362880}x^9 + \dots \\ &= x - \frac{1}{6}x^3 + \frac{1}{120}x^5 - \frac{1}{5040}x^7 + \frac{1}{362880}x^9 \end{aligned}$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{1+n+r} a_n \right) + \sum_{n=0}^{\infty} \left(-\frac{x^{n+r+3} a_n}{6} \right) \quad (2A) \\ & + \left(\sum_{n=0}^{\infty} \frac{x^{n+r+5} a_n}{120} \right) + \sum_{n=0}^{\infty} \left(-\frac{x^{n+r+7} a_n}{5040} \right) + \left(\sum_{n=0}^{\infty} \frac{x^{n+r+9} a_n}{362880} \right) = 0 \end{aligned}$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=0}^{\infty} x^{1+n+r} a_n &= \sum_{n=1}^{\infty} a_{n-1} x^{n+r} \\ \sum_{n=0}^{\infty} \left(-\frac{x^{n+r+3} a_n}{6} \right) &= \sum_{n=3}^{\infty} \left(-\frac{a_{n-3} x^{n+r}}{6} \right) \\ \sum_{n=0}^{\infty} \frac{x^{n+r+5} a_n}{120} &= \sum_{n=5}^{\infty} \frac{a_{n-5} x^{n+r}}{120} \\ \sum_{n=0}^{\infty} \left(-\frac{x^{n+r+7} a_n}{5040} \right) &= \sum_{n=7}^{\infty} \left(-\frac{a_{n-7} x^{n+r}}{5040} \right) \\ \sum_{n=0}^{\infty} \frac{x^{n+r+9} a_n}{362880} &= \sum_{n=9}^{\infty} \frac{a_{n-9} x^{n+r}}{362880} \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=1}^{\infty} a_{n-1} x^{n+r} \right) + \sum_{n=3}^{\infty} \left(-\frac{a_{n-3} x^{n+r}}{6} \right) \quad (2B) \\ & + \left(\sum_{n=5}^{\infty} \frac{a_{n-5} x^{n+r}}{120} \right) + \sum_{n=7}^{\infty} \left(-\frac{a_{n-7} x^{n+r}}{5040} \right) + \left(\sum_{n=9}^{\infty} \frac{a_{n-9} x^{n+r}}{362880} \right) = 0 \end{aligned}$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r} a_n (n+r) (n+r-1) = 0$$

When $n = 0$ the above becomes

$$x^r a_0 r(-1 + r) = 0$$

Or

$$x^r a_0 r(-1 + r) = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$x^r r(-1 + r) = 0$$

Since the above is true for all x then the indicial equation becomes

$$r(-1 + r) = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = 1$$

$$r_2 = 0$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$x^r r(-1 + r) = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = 1$ is an integer, then we can construct two linearly independent solutions

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$
$$y_2(x) = C y_1(x) \ln(x) + x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = x \left(\sum_{n=0}^{\infty} a_n x^n \right)$$
$$y_2(x) = C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{1+n}$$
$$y_2(x) = C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Where C above can be zero. We start by finding y_1 . Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = -\frac{1}{r(1+r)}$$

Substituting $n = 2$ in Eq. (2B) gives

$$a_2 = \frac{1}{r(1+r)^2(2+r)}$$

Substituting $n = 3$ in Eq. (2B) gives

$$a_3 = \frac{r^4 + 4r^3 + 5r^2 + 2r - 6}{6r(1+r)^2(2+r)^2(3+r)}$$

Substituting $n = 4$ in Eq. (2B) gives

$$a_4 = \frac{-r^4 - 6r^3 - 14r^2 - 15r - 3}{3r(1+r)^2(2+r)^2(3+r)^2(4+r)}$$

Substituting $n = 5$ in Eq. (2B) gives

$$a_5 = \frac{-r^8 - 16r^7 - 106r^6 - 376r^5 - 709r^4 - 424r^3 + 1056r^2 + 2496r + 1560}{120r(1+r)^2(2+r)^2(3+r)^2(4+r)^2(5+r)}$$

Substituting $n = 6$ in Eq. (2B) gives

$$a_6 = \frac{2r^8 + 40r^7 + 337r^6 + 1555r^5 + 4258r^4 + 6955r^3 + 6113r^2 + 1440r - 1305}{45r(1+r)^2(2+r)^2(3+r)^2(4+r)^2(5+r)^2(6+r)}$$

Substituting $n = 7$ in Eq. (2B) gives

$$a_7 = \frac{r^{12} + 36r^{11} + 575r^{10} + 5370r^9 + 31977r^8 + 120744r^7 + 245517r^6 - 40230r^5 - 1838278r^4 - 5602080r^3 - 1000000r^2 - 500000r}{5040r(1+r)^2(2+r)^2(3+r)^2(4+r)^2(5+r)^2(6+r)^2(7+r)}$$

Substituting $n = 8$ in Eq. (2B) gives

$$a_8 = \frac{-r^{12} - 42r^{11} - 785r^{10} - 8610r^9 - 61461r^8 - 298830r^7 - 1003386r^6 - 2292885r^5 - 3351683r^4 - 2400000r^3 - 1000000r^2 - 200000r}{315r(1+r)^2(2+r)^2(3+r)^2(4+r)^2(5+r)^2(6+r)^2(7+r)^2}$$

For $9 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) + a_{n-1} - \frac{a_{n-3}}{6} + \frac{a_{n-5}}{120} - \frac{a_{n-7}}{5040} + \frac{a_{n-9}}{362880} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{362880a_{n-1} - 60480a_{n-3} + 3024a_{n-5} - 72a_{n-7} + a_{n-9}}{362880(n+r)(n+r-1)} \quad (4)$$

Which for the root $r = 1$ becomes

$$a_n = \frac{-362880a_{n-1} + 60480a_{n-3} - 3024a_{n-5} + 72a_{n-7} - a_{n-9}}{362880(1+n)n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 1$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$
a_0	1
a_1	$-\frac{1}{r(1+r)}$
a_2	$\frac{1}{r(1+r)^2(2+r)}$
a_3	$\frac{r^4+4r^3+5r^2+2r-6}{6r(1+r)^2(2+r)^2(3+r)}$
a_4	$\frac{-r^4-6r^3-14r^2-15r-3}{3r(1+r)^2(2+r)^2(3+r)^2(4+r)}$
a_5	$\frac{-r^8-16r^7-106r^6-376r^5-709r^4-424r^3+1056r^2+2496r+1560}{120r(1+r)^2(2+r)^2(3+r)^2(4+r)^2(5+r)}$
a_6	$\frac{2r^8+40r^7+337r^6+1555r^5+4258r^4+6955r^3+6113r^2+1440r-1305}{45r(1+r)^2(2+r)^2(3+r)^2(4+r)^2(5+r)^2(6+r)}$
a_7	$\frac{r^{12}+36r^{11}+575r^{10}+5370r^9+31977r^8+120744r^7+245517r^6-40230r^5-1838278r^4-5602080r^3-8696192r^2-7058640r-2172240}{5040r(1+r)^2(2+r)^2(3+r)^2(4+r)^2(5+r)^2(6+r)^2(7+r)}$
a_8	$\frac{-r^{12}-42r^{11}-785r^{10}-8610r^9-61461r^8-298830r^7-1003386r^6-2292885r^5-3351683r^4-2497173r^3+466901r^2+2574075r+1562715}{315r(1+r)^2(2+r)^2(3+r)^2(4+r)^2(5+r)^2(6+r)^2(7+r)^2(8+r)}$

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots) \\ &= x\left(1 - \frac{x}{2} + \frac{x^2}{12} + \frac{x^3}{144} - \frac{13x^4}{2880} + \frac{29x^5}{86400} + \frac{431x^6}{3628800} - \frac{4961x^7}{203212800} - \frac{5197x^8}{4877107200} + O(x^8)\right) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Let

$$r_1 - r_2 = N$$

Where N is positive integer which is the difference between the two roots. r_1 is taken as the larger root. Hence for this problem we have $N = 1$. Now we need to determine if

C is zero or not. This is done by finding $\lim_{r \rightarrow r_2} a_1(r)$. If this limit exists, then $C = 0$, else we need to keep the log term and $C \neq 0$. The above table shows that

$$\begin{aligned} a_N &= a_1 \\ &= -\frac{1}{r(1+r)} \end{aligned}$$

Therefore

$$\begin{aligned} \lim_{r \rightarrow r_2} -\frac{1}{r(1+r)} &= \lim_{r \rightarrow 0} -\frac{1}{r(1+r)} \\ &= \text{undefined} \end{aligned}$$

Since the limit does not exist then the log term is needed. Therefore the second solution has the form

$$y_2(x) = Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right)$$

Therefore

$$\begin{aligned} \frac{d}{dx} y_2(x) &= Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2} (n+r_2)}{x} \right) \\ &= Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} x^{-1+n+r_2} b_n (n+r_2) \right) \\ \frac{d^2}{dx^2} y_2(x) &= Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} \\ &\quad + \sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2} (n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2} (n+r_2)}{x^2} \right) \\ &= Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} + \left(\sum_{n=0}^{\infty} x^{-2+n+r_2} b_n (n+r_2) (-1+n+r_2) \right) \end{aligned}$$

Substituting these back into the given ode $x^2 y'' + \sin(x) y = 0$ gives

$$\begin{aligned} &x^2 \left(Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} \right. \\ &\quad \left. + \sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2} (n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2} (n+r_2)}{x^2} \right) \right) \\ &\quad + \sin(x) \left(Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) \right) = 0 \end{aligned}$$

Which can be written as

$$\begin{aligned} & \left((y_1''(x) x^2 + \sin(x) y_1(x)) \ln(x) + x^2 \left(\frac{2y_1'(x)}{x} - \frac{y_1(x)}{x^2} \right) \right) C \\ & + x^2 \left(\sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2} (n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2} (n+r_2)}{x^2} \right) \right) \\ & + \sin(x) \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) = 0 \end{aligned} \quad (7)$$

But since $y_1(x)$ is a solution to the ode, then

$$y_1''(x) x^2 + \sin(x) y_1(x) = 0$$

Eq (7) simplifies to

$$\begin{aligned} & x^2 \left(\frac{2y_1'(x)}{x} - \frac{y_1(x)}{x^2} \right) C + x^2 \left(\sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2} (n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2} (n+r_2)}{x^2} \right) \right) \\ & + \sin(x) \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) = 0 \end{aligned} \quad (8)$$

Substituting $y_1 = \sum_{n=0}^{\infty} a_n x^{n+r_1}$ into the above gives

$$\begin{aligned} & \left(2 \left(\sum_{n=0}^{\infty} x^{-1+n+r_1} a_n (n+r_1) \right) x - \left(\sum_{n=0}^{\infty} a_n x^{n+r_1} \right) \right) C \\ & + \left(\sum_{n=0}^{\infty} x^{-2+n+r_2} b_n (n+r_2) (-1+n+r_2) \right) x^2 + \sin(x) \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) = 0 \end{aligned} \quad (9)$$

Since $r_1 = 1$ and $r_2 = 0$ then the above becomes

$$\begin{aligned} & \left(2 \left(\sum_{n=0}^{\infty} x^n a_n (1+n) \right) x - \left(\sum_{n=0}^{\infty} a_n x^{1+n} \right) \right) C \\ & + \left(\sum_{n=0}^{\infty} x^{-2+n} b_n n (n-1) \right) x^2 + \sin(x) \left(\sum_{n=0}^{\infty} b_n x^n \right) = 0 \end{aligned} \quad (10)$$

Expanding $\sin(x)$ as Taylor series around $x = 0$ and keeping only the first 8 terms gives

$$\begin{aligned}\sin(x) &= x - \frac{1}{6}x^3 + \frac{1}{120}x^5 - \frac{1}{5040}x^7 + \frac{1}{362880}x^9 + \dots \\ &= x - \frac{1}{6}x^3 + \frac{1}{120}x^5 - \frac{1}{5040}x^7 + \frac{1}{362880}x^9\end{aligned}$$

Which simplifies to

$$\begin{aligned}&\left(\sum_{n=0}^{\infty} 2C x^{1+n} a_n(1+n)\right) + \sum_{n=0}^{\infty} (-Ca_n x^{1+n}) + \left(\sum_{n=0}^{\infty} n x^n b_n(n-1)\right) \\ &+ \left(\sum_{n=0}^{\infty} x^{1+n} b_n\right) + \sum_{n=0}^{\infty} \left(-\frac{x^{n+3} b_n}{6}\right) + \left(\sum_{n=0}^{\infty} \frac{x^{n+5} b_n}{120}\right) \\ &+ \sum_{n=0}^{\infty} \left(-\frac{x^{n+7} b_n}{5040}\right) + \left(\sum_{n=0}^{\infty} \frac{x^{n+9} b_n}{362880}\right) = 0\end{aligned}\tag{2A}$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\begin{aligned}\sum_{n=0}^{\infty} 2C x^{1+n} a_n(1+n) &= \sum_{n=1}^{\infty} 2C a_{n-1} n x^n \\ \sum_{n=0}^{\infty} (-Ca_n x^{1+n}) &= \sum_{n=1}^{\infty} (-C a_{n-1} x^n) \\ \sum_{n=0}^{\infty} x^{1+n} b_n &= \sum_{n=1}^{\infty} b_{n-1} x^n \\ \sum_{n=0}^{\infty} \left(-\frac{x^{n+3} b_n}{6}\right) &= \sum_{n=3}^{\infty} \left(-\frac{b_{n-3} x^n}{6}\right) \\ \sum_{n=0}^{\infty} \frac{x^{n+5} b_n}{120} &= \sum_{n=5}^{\infty} \frac{b_{n-5} x^n}{120} \\ \sum_{n=0}^{\infty} \left(-\frac{x^{n+7} b_n}{5040}\right) &= \sum_{n=7}^{\infty} \left(-\frac{b_{n-7} x^n}{5040}\right) \\ \sum_{n=0}^{\infty} \frac{x^{n+9} b_n}{362880} &= \sum_{n=9}^{\infty} \frac{b_{n-9} x^n}{362880}\end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to n .

$$\begin{aligned} & \left(\sum_{n=1}^{\infty} 2Ca_{n-1}n x^n \right) + \sum_{n=1}^{\infty} (-Ca_{n-1}x^n) + \left(\sum_{n=0}^{\infty} n x^n b_n(n-1) \right) \\ & + \left(\sum_{n=1}^{\infty} b_{n-1}x^n \right) + \sum_{n=3}^{\infty} \left(-\frac{b_{n-3}x^n}{6} \right) + \left(\sum_{n=5}^{\infty} \frac{b_{n-5}x^n}{120} \right) \\ & + \sum_{n=7}^{\infty} \left(-\frac{b_{n-7}x^n}{5040} \right) + \left(\sum_{n=9}^{\infty} \frac{b_{n-9}x^n}{362880} \right) = 0 \end{aligned} \quad (2B)$$

For $n = 0$ in Eq. (2B), we choose arbitrary value for b_0 as $b_0 = 1$. For $n = N$, where $N = 1$ which is the difference between the two roots, we are free to choose $b_1 = 0$. Hence for $n = 1$, Eq (2B) gives

$$C + 1 = 0$$

Which is solved for C . Solving for C gives

$$C = -1$$

For $n = 2$, Eq (2B) gives

$$3Ca_1 + b_1 + 2b_2 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$2b_2 + \frac{3}{2} = 0$$

Solving the above for b_2 gives

$$b_2 = -\frac{3}{4}$$

For $n = 3$, Eq (2B) gives

$$5Ca_2 + b_2 + 6b_3 - \frac{b_0}{6} = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$-\frac{4}{3} + 6b_3 = 0$$

Solving the above for b_3 gives

$$b_3 = \frac{2}{9}$$

For $n = 4$, Eq (2B) gives

$$7Ca_3 + b_3 + 12b_4 - \frac{b_1}{6} = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$\frac{25}{144} + 12b_4 = 0$$

Solving the above for b_4 gives

$$b_4 = -\frac{25}{1728}$$

For $n = 5$, Eq (2B) gives

$$9Ca_4 + b_4 + 20b_5 - \frac{b_2}{6} + \frac{b_0}{120} = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$\frac{689}{4320} + 20b_5 = 0$$

Solving the above for b_5 gives

$$b_5 = -\frac{689}{86400}$$

For $n = 6$, Eq (2B) gives

$$11Ca_5 + b_5 + 30b_6 - \frac{b_3}{6} + \frac{b_1}{120} = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$-\frac{263}{5400} + 30b_6 = 0$$

Solving the above for b_6 gives

$$b_6 = \frac{263}{162000}$$

For $n = 7$, Eq (2B) gives

$$13Ca_6 + b_6 + 42b_7 - \frac{b_4}{6} + \frac{b_2}{120} - \frac{b_0}{5040} = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$-\frac{71809}{18144000} + 42b_7 = 0$$

Solving the above for b_7 gives

$$b_7 = \frac{71809}{762048000}$$

Now that we found all b_n and C , we can calculate the second solution from

$$y_2(x) = Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right)$$

Using the above value found for $C = -1$ and all b_n , then the second solution becomes

$$y_2(x) = (-1) \left(x \left(1 - \frac{x}{2} + \frac{x^2}{12} + \frac{x^3}{144} - \frac{13x^4}{2880} + \frac{29x^5}{86400} + \frac{431x^6}{3628800} - \frac{4961x^7}{203212800} - \frac{5197x^8}{4877107200} + O(x^8) \right) \right) \ln(x) + 1 - \frac{3x^2}{4} + \frac{2x^3}{9} - \frac{25x^4}{1728} - \frac{689x^5}{86400} + \frac{263x^6}{162000} + \frac{71809x^7}{762048000} + O(x^8)$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1 y_1(x) + c_2 y_2(x) \\ &= c_1 x \left(1 - \frac{x}{2} + \frac{x^2}{12} + \frac{x^3}{144} - \frac{13x^4}{2880} + \frac{29x^5}{86400} + \frac{431x^6}{3628800} - \frac{4961x^7}{203212800} - \frac{5197x^8}{4877107200} + O(x^8) \right) \\ &\quad + c_2 \left((-1) \left(x \left(1 - \frac{x}{2} + \frac{x^2}{12} + \frac{x^3}{144} - \frac{13x^4}{2880} + \frac{29x^5}{86400} + \frac{431x^6}{3628800} - \frac{4961x^7}{203212800} - \frac{5197x^8}{4877107200} + O(x^8) \right) \right) \right. \\ &\quad \left. + 1 - \frac{3x^2}{4} + \frac{2x^3}{9} - \frac{25x^4}{1728} - \frac{689x^5}{86400} + \frac{263x^6}{162000} + \frac{71809x^7}{762048000} + O(x^8) \right) \end{aligned}$$

Hence the final solution is

$$\begin{aligned} y &= y_h \\ &= c_1 x \left(1 - \frac{x}{2} + \frac{x^2}{12} + \frac{x^3}{144} - \frac{13x^4}{2880} + \frac{29x^5}{86400} + \frac{431x^6}{3628800} - \frac{4961x^7}{203212800} - \frac{5197x^8}{4877107200} + O(x^8) \right) \\ &\quad + c_2 \left(-x \left(1 - \frac{x}{2} + \frac{x^2}{12} + \frac{x^3}{144} - \frac{13x^4}{2880} + \frac{29x^5}{86400} + \frac{431x^6}{3628800} - \frac{4961x^7}{203212800} - \frac{5197x^8}{4877107200} + O(x^8) \right) \right. \\ &\quad \left. \ln(x) + 1 - \frac{3x^2}{4} + \frac{2x^3}{9} - \frac{25x^4}{1728} - \frac{689x^5}{86400} + \frac{263x^6}{162000} + \frac{71809x^7}{762048000} + O(x^8) \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned} y &= c_1 x \left(1 - \frac{x}{2} + \frac{x^2}{12} + \frac{x^3}{144} - \frac{13x^4}{2880} + \frac{29x^5}{86400} + \frac{431x^6}{3628800} - \frac{4961x^7}{203212800} - \frac{5197x^8}{4877107200} + O(x^8) \right) \\ &\quad + c_2 \left(-x \left(1 - \frac{x}{2} + \frac{x^2}{12} + \frac{x^3}{144} - \frac{13x^4}{2880} + \frac{29x^5}{86400} + \frac{431x^6}{3628800} - \frac{4961x^7}{203212800} - \frac{5197x^8}{4877107200} + O(x^8) \right) \right. \\ &\quad \left. \ln(x) + 1 - \frac{3x^2}{4} + \frac{2x^3}{9} - \frac{25x^4}{1728} - \frac{689x^5}{86400} + \frac{263x^6}{162000} + \frac{71809x^7}{762048000} + O(x^8) \right) \end{aligned}$$

Verification of solutions

$$y = c_1 x \left(1 - \frac{x}{2} + \frac{x^2}{12} + \frac{x^3}{144} - \frac{13x^4}{2880} + \frac{29x^5}{86400} + \frac{431x^6}{3628800} - \frac{4961x^7}{203212800} - \frac{5197x^8}{4877107200} + O(x^8) \right) \\ + c_2 \left(-x \left(1 - \frac{x}{2} + \frac{x^2}{12} + \frac{x^3}{144} - \frac{13x^4}{2880} + \frac{29x^5}{86400} + \frac{431x^6}{3628800} - \frac{4961x^7}{203212800} - \frac{5197x^8}{4877107200} + O(x^8) \right) \ln(x) + 1 - \frac{3x^2}{4} + \frac{2x^3}{9} - \frac{25x^4}{1728} - \frac{689x^5}{86400} + \frac{263x^6}{162000} + \frac{71809x^7}{762048000} + O(x^8) \right)$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) * 2F1([a
-> Trying changes of variables to rationalize or make the ODE simpler
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) * 2F1
trying a symmetry of the form [xi=0, eta=F(x)]
trying 2nd order exact linear
trying symmetries linear in x and y(x)
trying to convert to a linear ODE with constant coefficients
-> trying with_periodic_functions in the coefficients
    --- Trying Lie symmetry methods, 2nd order ---
    `, `-> Computing symmetries using: way = 5
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) * 2F1
trying a symmetry of the form [xi=0, eta=F(x)]
trying 2nd order exact linear
trying symmetries linear in x and y(x)
trying to convert to a linear ODE with constant coefficients
<- unable to find a useful change of variables
trying differential order: 2; exact nonlinear
trying symmetries linear in x and y(x)
trying to convert to a linear ODE with constant coefficients
trying to convert to an ODE of Bessel type
-> trying reduction of order to Riccati
trying Riccati sub-methods:
    trying Riccati_symmetries
    -> trying a symmetry pattern of the form [F(x)*G(y), 0]
    -> trying a symmetry pattern of the form [0, F(x)*G(y)]
    -> trying a symmetry pattern of the form [F(x),G(x)*y+H(x)]
-> trying with_periodic_functions in the coefficients
    --- Trying Lie symmetry methods, 2nd order ---
    `, `-> Computing symmetries using: way = 5
--- Trying Lie symmetry methods, 2nd order ---
`, `-> Computing symmetries using: way = 3 [0, y]
```

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 70

Order:=8;

dsolve(x^2*diff(y(x),x\$2)+sin(x)*y(x)=0,y(x),type='series',x=0);

$$y(x) = c_1 x \left(1 - \frac{1}{2}x + \frac{1}{12}x^2 + \frac{1}{144}x^3 - \frac{13}{2880}x^4 + \frac{29}{86400}x^5 + \frac{431}{3628800}x^6 - \frac{4961}{203212800}x^7 + O(x^8) \right) + c_2 \left(\ln(x) \left(-x + \frac{1}{2}x^2 - \frac{1}{12}x^3 - \frac{1}{144}x^4 + \frac{13}{2880}x^5 - \frac{29}{86400}x^6 - \frac{431}{3628800}x^7 + O(x^8) \right) + \left(1 - \frac{3}{4}x^2 + \frac{2}{9}x^3 - \frac{25}{1728}x^4 - \frac{689}{86400}x^5 + \frac{263}{162000}x^6 + \frac{71809}{762048000}x^7 + O(x^8) \right) \right)$$

✓ Solution by Mathematica

Time used: 0.043 (sec). Leaf size: 121

AsymptoticDSolveValue[x^2*y''[x]+Sin[x]*y[x]==0,y[x],{x,0,7}]

$$y(x) \rightarrow c_1 \left(\frac{2539x^6 - 16185x^5 - 9750x^4 + 396000x^3 - 1620000x^2 + 1296000x + 1296000}{1296000} - \frac{x(29x^5 - 390x^4 + 600x^3 + 7200x^2 - 43200x + 86400) \log(x)}{86400} \right) + c_2 \left(\frac{431x^7}{3628800} + \frac{29x^6}{86400} - \frac{13x^5}{2880} + \frac{x^4}{144} + \frac{x^3}{12} - \frac{x^2}{2} + x \right)$$

19.8 problem 2(d)

19.8.1 Maple step by step solution 3206

Internal problem ID [6448]

Internal file name [OUTPUT/5696_Sunday_June_05_2022_03_47_30_PM_15278874/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.4. REGULAR SINGULAR POINTS. Page 175

Problem number: 2(d).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Complex roots"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$x^3 y'' + \sin(x) y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^3 y'' + \sin(x) y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = 0$$

$$q(x) = \frac{\sin(x)}{x^3}$$

Table 445: Table $p(x), q(x)$ singularities.

$p(x) = 0$	
singularity	type

$q(x) = \frac{\sin(x)}{x^3}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^3 y'' + \sin(x) y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) x^3 + \sin(x) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Expanding $\sin(x)$ as Taylor series around $x = 0$ and keeping only the first 8 terms gives

$$\begin{aligned} \sin(x) &= x - \frac{1}{6}x^3 + \frac{1}{120}x^5 - \frac{1}{5040}x^7 + \frac{1}{362880}x^9 + \dots \\ &= x - \frac{1}{6}x^3 + \frac{1}{120}x^5 - \frac{1}{5040}x^7 + \frac{1}{362880}x^9 \end{aligned}$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{1+n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{1+n+r} a_n \right) + \sum_{n=0}^{\infty} \left(-\frac{x^{n+r+3} a_n}{6} \right) \quad (2A) \\ & + \left(\sum_{n=0}^{\infty} \frac{x^{n+r+5} a_n}{120} \right) + \sum_{n=0}^{\infty} \left(-\frac{x^{n+r+7} a_n}{5040} \right) + \left(\sum_{n=0}^{\infty} \frac{x^{n+r+9} a_n}{362880} \right) = 0 \end{aligned}$$

The next step is to make all powers of x be $1+n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{1+n+r} and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=0}^{\infty} \left(-\frac{x^{n+r+3} a_n}{6} \right) &= \sum_{n=2}^{\infty} \left(-\frac{a_{n-2} x^{1+n+r}}{6} \right) \\ \sum_{n=0}^{\infty} \frac{x^{n+r+5} a_n}{120} &= \sum_{n=4}^{\infty} \frac{a_{n-4} x^{1+n+r}}{120} \\ \sum_{n=0}^{\infty} \left(-\frac{x^{n+r+7} a_n}{5040} \right) &= \sum_{n=6}^{\infty} \left(-\frac{a_{n-6} x^{1+n+r}}{5040} \right) \\ \sum_{n=0}^{\infty} \frac{x^{n+r+9} a_n}{362880} &= \sum_{n=8}^{\infty} \frac{a_{n-8} x^{1+n+r}}{362880} \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $1+n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{1+n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{1+n+r} a_n \right) + \sum_{n=2}^{\infty} \left(-\frac{a_{n-2} x^{1+n+r}}{6} \right) \quad (2B) \\ & + \left(\sum_{n=4}^{\infty} \frac{a_{n-4} x^{1+n+r}}{120} \right) + \sum_{n=6}^{\infty} \left(-\frac{a_{n-6} x^{1+n+r}}{5040} \right) + \left(\sum_{n=8}^{\infty} \frac{a_{n-8} x^{1+n+r}}{362880} \right) = 0 \end{aligned}$$

The indicial equation is obtained from $n = 0$. From Eq (2B) this gives

$$x^{1+n+r} a_n (n+r) (n+r-1) + x^{1+n+r} a_n = 0$$

When $n = 0$ the above becomes

$$x^{1+r} a_0 r (-1+r) + x^{1+r} a_0 = 0$$

Or

$$(x^{1+r}r(-1+r) + x^{1+r}) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(r^2 - r + 1) x^{1+r} = 0$$

Since the above is true for all x then the indicial equation becomes

$$r^2 - r + 1 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = \frac{1}{2} + \frac{i\sqrt{3}}{2}$$
$$r_2 = \frac{1}{2} - \frac{i\sqrt{3}}{2}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(r^2 - r + 1) x^{1+r} = 0$$

Solving for r gives the roots of the indicial equation as Since the roots are complex conjugates, then two linearly independent solutions can be constructed using

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$
$$y_2(x) = x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+\frac{1}{2}+\frac{i\sqrt{3}}{2}}$$
$$y_2(x) = \sum_{n=0}^{\infty} b_n x^{n+\frac{1}{2}-\frac{i\sqrt{3}}{2}}$$

$y_1(x)$ is found first. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

Substituting $n = 2$ in Eq. (2B) gives

$$a_2 = \frac{1}{6r^2 + 18r + 18}$$

Substituting $n = 3$ in Eq. (2B) gives

$$a_3 = 0$$

Substituting $n = 4$ in Eq. (2B) gives

$$a_4 = \frac{-3r^2 - 9r + 1}{360(r^2 + 3r + 3)(r^2 + 7r + 13)}$$

Substituting $n = 5$ in Eq. (2B) gives

$$a_5 = 0$$

Substituting $n = 6$ in Eq. (2B) gives

$$a_6 = \frac{3r^4 + 30r^3 + 69r^2 - 30r - 149}{15120(r^2 + 3r + 3)(r^2 + 7r + 13)(r^2 + 11r + 31)}$$

Substituting $n = 7$ in Eq. (2B) gives

$$a_7 = 0$$

For $8 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) + a_n - \frac{a_{n-2}}{6} + \frac{a_{n-4}}{120} - \frac{a_{n-6}}{5040} + \frac{a_{n-8}}{362880} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{-60480a_{n-2} + 3024a_{n-4} - 72a_{n-6} + a_{n-8}}{362880(n^2 + 2nr + r^2 - n - r + 1)} \quad (4)$$

Which for the root $r = \frac{1}{2} + \frac{i\sqrt{3}}{2}$ becomes

$$a_n = \frac{60480a_{n-2} - 3024a_{n-4} + 72a_{n-6} - a_{n-8}}{362880n(i\sqrt{3} + n)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = \frac{1}{2} + \frac{i\sqrt{3}}{2}$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$\frac{1}{6r^2+18r+18}$	$\frac{1}{12i\sqrt{3}+24}$
a_3	0	0
a_4	$\frac{-3r^2-9r+1}{360(r^2+3r+3)(r^2+7r+13)}$	$\frac{-3i\sqrt{3}-1}{1440(i\sqrt{3}+2)(i\sqrt{3}+4)}$
a_5	0	0
a_6	$\frac{3r^4+30r^3+69r^2-30r-149}{15120(r^2+3r+3)(r^2+7r+13)(r^2+11r+31)}$	$\frac{9i\sqrt{3}-115}{362880(i\sqrt{3}+2)(i\sqrt{3}+4)(i\sqrt{3}+6)}$
a_7	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned}
y_1(x) &= x^{\frac{1}{2} + \frac{i\sqrt{3}}{2}} (a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots) \\
&= x^{\frac{1}{2} + \frac{i\sqrt{3}}{2}} \left(1 + \frac{x^2}{12i\sqrt{3} + 24} + \frac{(-3i\sqrt{3} - 1)x^4}{1440(i\sqrt{3} + 2)(i\sqrt{3} + 4)} \right. \\
&\quad \left. + \frac{(9i\sqrt{3} - 115)x^6}{362880(i\sqrt{3} + 2)(i\sqrt{3} + 4)(i\sqrt{3} + 6)} + O(x^8) \right)
\end{aligned}$$

The second solution $y_2(x)$ is found by taking the complex conjugate of $y_1(x)$ which gives

$$\begin{aligned}
y_2(x) &= x^{\frac{1}{2} - \frac{i\sqrt{3}}{2}} \left(1 + \frac{x^2}{-12i\sqrt{3} + 24} + \frac{(3i\sqrt{3} - 1)x^4}{1440(2 - i\sqrt{3})(-i\sqrt{3} + 4)} \right. \\
&\quad \left. + \frac{(-9i\sqrt{3} - 115)x^6}{362880(2 - i\sqrt{3})(-i\sqrt{3} + 4)(-i\sqrt{3} + 6)} + O(x^8) \right)
\end{aligned}$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 y_1(x) + c_2 y_2(x)$$

$$\begin{aligned}
&= c_1 x^{\frac{1}{2} + \frac{i\sqrt{3}}{2}} \left(1 + \frac{x^2}{12i\sqrt{3} + 24} + \frac{(-3i\sqrt{3} - 1)x^4}{1440(i\sqrt{3} + 2)(i\sqrt{3} + 4)} \right. \\
&\quad \left. + \frac{(9i\sqrt{3} - 115)x^6}{362880(i\sqrt{3} + 2)(i\sqrt{3} + 4)(i\sqrt{3} + 6)} + O(x^8) \right) \\
&+ c_2 x^{\frac{1}{2} - \frac{i\sqrt{3}}{2}} \left(1 + \frac{x^2}{-12i\sqrt{3} + 24} + \frac{(3i\sqrt{3} - 1)x^4}{1440(2 - i\sqrt{3})(-i\sqrt{3} + 4)} \right. \\
&\quad \left. + \frac{(-9i\sqrt{3} - 115)x^6}{362880(2 - i\sqrt{3})(-i\sqrt{3} + 4)(-i\sqrt{3} + 6)} + O(x^8) \right)
\end{aligned}$$

Hence the final solution is

$$\begin{aligned}
y &= y_h \\
&= c_1 x^{\frac{1}{2} + \frac{i\sqrt{3}}{2}} \left(1 + \frac{x^2}{12i\sqrt{3} + 24} + \frac{(-3i\sqrt{3} - 1)x^4}{1440(i\sqrt{3} + 2)(i\sqrt{3} + 4)} \right. \\
&\quad \left. + \frac{(9i\sqrt{3} - 115)x^6}{362880(i\sqrt{3} + 2)(i\sqrt{3} + 4)(i\sqrt{3} + 6)} + O(x^8) \right) \\
&+ c_2 x^{\frac{1}{2} - \frac{i\sqrt{3}}{2}} \left(1 + \frac{x^2}{-12i\sqrt{3} + 24} + \frac{(3i\sqrt{3} - 1)x^4}{1440(2 - i\sqrt{3})(-i\sqrt{3} + 4)} \right. \\
&\quad \left. + \frac{(-9i\sqrt{3} - 115)x^6}{362880(2 - i\sqrt{3})(-i\sqrt{3} + 4)(-i\sqrt{3} + 6)} + O(x^8) \right)
\end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}
y &= c_1 x^{\frac{1}{2} + \frac{i\sqrt{3}}{2}} \left(1 + \frac{x^2}{12i\sqrt{3} + 24} + \frac{(-3i\sqrt{3} - 1)x^4}{1440(i\sqrt{3} + 2)(i\sqrt{3} + 4)} \right. \\
&\quad \left. + \frac{(9i\sqrt{3} - 115)x^6}{362880(i\sqrt{3} + 2)(i\sqrt{3} + 4)(i\sqrt{3} + 6)} + O(x^8) \right) \\
&+ c_2 x^{\frac{1}{2} - \frac{i\sqrt{3}}{2}} \left(1 + \frac{x^2}{-12i\sqrt{3} + 24} + \frac{(3i\sqrt{3} - 1)x^4}{1440(2 - i\sqrt{3})(-i\sqrt{3} + 4)} \right. \\
&\quad \left. + \frac{(-9i\sqrt{3} - 115)x^6}{362880(2 - i\sqrt{3})(-i\sqrt{3} + 4)(-i\sqrt{3} + 6)} + O(x^8) \right)
\end{aligned}$$

Verification of solutions

$$y = c_1 x^{\frac{1}{2} + \frac{i\sqrt{3}}{2}} \left(1 + \frac{x^2}{12i\sqrt{3} + 24} + \frac{(-3i\sqrt{3} - 1)x^4}{1440(i\sqrt{3} + 2)(i\sqrt{3} + 4)} \right. \\ \left. + \frac{(9i\sqrt{3} - 115)x^6}{362880(i\sqrt{3} + 2)(i\sqrt{3} + 4)(i\sqrt{3} + 6)} + O(x^8) \right) \\ + c_2 x^{\frac{1}{2} - \frac{i\sqrt{3}}{2}} \left(1 + \frac{x^2}{-12i\sqrt{3} + 24} + \frac{(3i\sqrt{3} - 1)x^4}{1440(2 - i\sqrt{3})(-i\sqrt{3} + 4)} \right. \\ \left. + \frac{(-9i\sqrt{3} - 115)x^6}{362880(2 - i\sqrt{3})(-i\sqrt{3} + 4)(-i\sqrt{3} + 6)} + O(x^8) \right)$$

Verified OK.

19.8.1 Maple step by step solution

Let's solve

$$y''x^3 + \sin(x)y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{\sin(x)y}{x^3}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{\sin(x)y}{x^3} = 0$$

- Multiply by denominators of the ODE

$$y''x^3 + \sin(x)y = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x , using the chain rule

$$y' = \left(\frac{d}{dt}y(t)\right)t'(x)$$

- Compute derivative

$$y' = \frac{\frac{d}{dt}y(t)}{x}$$

- Calculate the 2nd derivative of y with respect to x , using the chain rule

$$y'' = \left(\frac{d^2}{dt^2} y(t) \right) t'(x)^2 + t''(x) \left(\frac{d}{dt} y(t) \right)$$

- Compute derivative

$$y'' = \frac{\frac{d^2}{dt^2} y(t)}{x^2} - \frac{\frac{d}{dt} y(t)}{x^2}$$

Substitute the change of variables back into the ODE

$$\left(\frac{\frac{d^2}{dt^2} y(t)}{x^2} - \frac{\frac{d}{dt} y(t)}{x^2} \right) x^3 + \sin(x) y(t) = 0$$

- Simplify

$$x \left(\frac{d^2}{dt^2} y(t) - \frac{d}{dt} y(t) \right) + \sin(x) y(t) = 0$$

- Isolate 2nd derivative

$$\frac{d^2}{dt^2} y(t) = -\frac{\sin(x)y(t)}{x} + \frac{d}{dt} y(t)$$

- Group terms with $y(t)$ on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$\frac{d^2}{dt^2} y(t) + \frac{\sin(x)y(t)}{x} - \frac{d}{dt} y(t) = 0$$

- Characteristic polynomial of ODE

$$r^2 + \frac{\sin(x)}{x} - r = 0$$

- Factor the characteristic polynomial

$$\frac{r^2 x - r x + \sin(x)}{x} = 0$$

- Roots of the characteristic polynomial

$$r = \left(\frac{x + \sqrt{x^2 - 4 \sin(x)x}}{2x}, -\frac{-x + \sqrt{x^2 - 4 \sin(x)x}}{2x} \right)$$

- 1st solution of the ODE

$$y_1(t) = e^{\frac{(x + \sqrt{x^2 - 4 \sin(x)x})t}{2x}}$$

- 2nd solution of the ODE

$$y_2(t) = e^{-\frac{(-x + \sqrt{x^2 - 4 \sin(x)x})t}{2x}}$$

- General solution of the ODE

$$y(t) = c_1 y_1(t) + c_2 y_2(t)$$

- Substitute in solutions

$$y(t) = c_1 e^{\frac{(x + \sqrt{x^2 - 4 \sin(x)x})t}{2x}} + c_2 e^{-\frac{(-x + \sqrt{x^2 - 4 \sin(x)x})t}{2x}}$$

- Change variables back using $t = \ln(x)$

$$y = c_1 e^{\frac{(x + \sqrt{x^2 - 4 \sin(x)x}) \ln(x)}{2x}} + c_2 e^{-\frac{(-x + \sqrt{x^2 - 4 \sin(x)x}) \ln(x)}{2x}}$$

- Simplify

$$y = x^{\frac{x + \sqrt{x(-4 \sin(x) + x)}}{2x}} c_1 + c_2 x^{-\frac{-x + \sqrt{x(-4 \sin(x) + x)}}{2x}}$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) * 2F1([a
-> Trying changes of variables to rationalize or make the ODE simpler
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) * 2F1
trying a symmetry of the form [xi=0, eta=F(x)]
trying 2nd order exact linear
trying symmetries linear in x and y(x)
trying to convert to a linear ODE with constant coefficients
-> trying with_periodic_functions in the coefficients
    --- Trying Lie symmetry methods, 2nd order ---
    `, `-> Computing symmetries using: way = 5
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) * 2F1
trying a symmetry of the form [xi=0, eta=F(x)]
trying 2nd order exact linear
trying symmetries linear in x and y(x)
trying to convert to a linear ODE with constant coefficients
<- unable to find a useful change of variables
trying differential order: 2; exact nonlinear
trying symmetries linear in x and y(x)
trying to convert to a linear ODE with constant coefficients
trying to convert to an ODE of Bessel type
-> trying reduction of order to Riccati
trying Riccati sub-methods:
    trying Riccati_symmetries
    -> trying a symmetry pattern of the form [F(x)*G(y), 0]
    -> trying a symmetry pattern of the form [0, F(x)*G(y)]
    -> trying a symmetry pattern of the form [F(x),G(x)*y+H(x)]
-> trying with_periodic_functions in the coefficients
    --- Trying Lie symmetry methods, 2nd order ---
    `, `-> Computing symmetries using: way = 5
--- Trying Lie symmetry methods, 2nd order ---
`, `-> Computing symmetries using: way = 3 [0, y]
```

✓ Solution by Maple

Time used: 0.172 (sec). Leaf size: 427

Order:=8;

dsolve(x^3*diff(y(x),x\$2)+sin(x)*y(x)=0,y(x),type='series',x=0);

$$y(x) = \sqrt{x} \left(c_2 x^{\frac{i\sqrt{3}}{2}} \left(1 + \frac{1}{12i\sqrt{3} + 24} x^2 + \frac{1}{1440} \frac{-3i\sqrt{3} - 1}{(i\sqrt{3} + 4)(i\sqrt{3} + 2)} x^4 \right. \right. \\ \left. \left. + \frac{1}{362880} \frac{9i\sqrt{3} - 115}{(i\sqrt{3} + 6)(i\sqrt{3} + 4)(i\sqrt{3} + 2)} x^6 + O(x^8) \right) + c_1 x^{-\frac{i\sqrt{3}}{2}} \left(1 \right. \\ \left. - \frac{1}{12i\sqrt{3} - 24} x^2 + \frac{-3\sqrt{3} - i}{7200i + 8640\sqrt{3}} x^4 + \frac{9\sqrt{3} - 115i}{4354560i + 14878080\sqrt{3}} x^6 + O(x^8) \right) \right)$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 410

AsymptoticDSolveValue[x^3*y''[x]+Sin[x]*y[x]==0,y[x],{x,0,7}]

$$y(x) \rightarrow c_1 \left(\frac{\left(\frac{1}{5040} - \frac{1}{720(1+(1-(-1)^{2/3})(2-(-1)^{2/3}))} + \frac{36(1+(1-(-1)^{2/3})(2-(-1)^{2/3}))^{-\frac{1}{120}}}{6(1+(3-(-1)^{2/3})(4-(-1)^{2/3}))} \right) x^6}{1 + (5 - (-1)^{2/3})(6 - (-1)^{2/3})} \right. \\ \left. + \frac{\left(\frac{36(1+(1-(-1)^{2/3})(2-(-1)^{2/3}))^{-\frac{1}{120}}}{6(1+(3-(-1)^{2/3})(4-(-1)^{2/3}))} - \frac{1}{120} \right) x^4}{1 + (3 - (-1)^{2/3})(4 - (-1)^{2/3})} + \frac{x^2}{6(1 + (1 - (-1)^{2/3})(2 - (-1)^{2/3}))} \right) \\ + 1 \left) x^{-(-1)^{2/3}} + c_2 \left(\frac{\left(\frac{1}{5040} - \frac{1}{720(1+(1+\sqrt[3]{-1})(2+\sqrt[3]{-1}))} + \frac{36(1+(1+\sqrt[3]{-1})(2+\sqrt[3]{-1}))^{-\frac{1}{120}}}{6(1+(3+\sqrt[3]{-1})(4+\sqrt[3]{-1}))} \right) x^6}{1 + (5 + \sqrt[3]{-1})(6 + \sqrt[3]{-1})} + \frac{\left(\frac{36(1+(1+\sqrt[3]{-1})(2+\sqrt[3]{-1}))^{-\frac{1}{120}}}{6(1+(3+\sqrt[3]{-1})(4+\sqrt[3]{-1}))} - \frac{1}{120} \right) x^4}{1 + (3 + \sqrt[3]{-1})(4 + \sqrt[3]{-1})} + \frac{x^2}{6(1 + (1 + (-1)^{2/3})(2 + (-1)^{2/3}))} \right) \right)$$

19.9 problem 2(e)

Internal problem ID [6449]

Internal file name [OUTPUT/5697_Sunday_June_05_2022_03_47_33_PM_29724232/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.4. REGULAR SINGULAR POINTS. Page 175

Problem number: 2(e).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Irregular singular point"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

Unable to solve or complete the solution.

$$x^4 y'' + \sin(x) y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^4 y'' + \sin(x) y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = 0$$
$$q(x) = \frac{\sin(x)}{x^4}$$

Table 447: Table $p(x), q(x)$ singularities.

$p(x) = 0$	
singularity	type

$q(x) = \frac{\sin(x)}{x^4}$	
singularity	type
$x = 0$	“irregular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : []

Irregular singular points : $[0, \infty]$

Since $x = 0$ is not an ordinary point, then we will now check if it is a regular singular point. Unable to solve since $x = 0$ is not regular singular point. Terminating.

Verification of solutions N/A

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) * 2F1([a
-> Trying changes of variables to rationalize or make the ODE simpler
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) * 2F1
trying a symmetry of the form [xi=0, eta=F(x)]
trying 2nd order exact linear
trying symmetries linear in x and y(x)
trying to convert to a linear ODE with constant coefficients
-> trying with_periodic_functions in the coefficients
    --- Trying Lie symmetry methods, 2nd order ---
    `, `-> Computing symmetries using: way = 5
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) * 2F1
trying a symmetry of the form [xi=0, eta=F(x)]
trying 2nd order exact linear
trying symmetries linear in x and y(x)
trying to convert to a linear ODE with constant coefficients
<- unable to find a useful change of variables
trying differential order: 2; exact nonlinear
trying symmetries linear in x and y(x)
trying to convert to a linear ODE with constant coefficients
trying to convert to an ODE of Bessel type
-> trying reduction of order to Riccati
trying Riccati sub-methods:
    trying Riccati_symmetries
    -> trying a symmetry pattern of the form [F(x)*G(y), 0]
    -> trying a symmetry pattern of the form [0, F(x)*G(y)]
    -> trying a symmetry pattern of the form [F(x),G(x)*y+H(x)]
-> trying with_periodic_functions in the coefficients
    --- Trying Lie symmetry methods, 2nd order ---
    `, `-> Computing symmetries using: way = 5
--- Trying Lie symmetry methods, 2nd order ---
`, `-> Computing symmetries using: way = 3 [0, y]
```

✗ Solution by Maple

```
Order:=8;
dsolve(x^4*diff(y(x),x$2)+sin(x)*y(x)=0,y(x),type='series',x=0);
```

No solution found

✓ Solution by Mathematica

Time used: 0.11 (sec). Leaf size: 294

```
AsymptoticDSolveValue[x^4*y''[x]+Sin[x]*y[x]==0,y[x],{x,0,7}]
```

$$\begin{aligned}
 y(x) \rightarrow & c_1 e^{-\frac{2i}{\sqrt{x}}x^{3/4}} \left(\frac{16487484152477478659746223ix^{13/2}}{2773583263632691770163200} \right. \\
 & - \frac{4594934148364735183693ix^{11/2}}{6320013947079701299200} + \frac{12579783586699513ix^{9/2}}{96185277197844480} - \frac{21896783401ix^{7/2}}{579820584960} \\
 & + \frac{856783ix^{5/2}}{41943040} - \frac{3151ix^{3/2}}{73728} - \frac{3986263268940827572255963529x^7}{207094217017907652172185600} \\
 & + \frac{21730712888356628741772337x^6}{10920984100553723845017600} - \frac{1500040357444099007x^5}{5129881450551705600} + \frac{4885269094757x^4}{74217034874880} \\
 & - \frac{2835642457x^3}{108716359680} + \frac{11659x^2}{524288} + \frac{15x}{512} - \frac{3i\sqrt{x}}{16} \\
 & \left. + 1 \right) + c_2 e^{\frac{2i}{\sqrt{x}}x^{3/4}} \left(-\frac{16487484152477478659746223ix^{13/2}}{2773583263632691770163200} + \frac{4594934148364735183693ix^{11/2}}{6320013947079701299200} - \frac{12579783}{961852} \right)
 \end{aligned}$$

19.10 problem 3(a)

Internal problem ID [6450]

Internal file name [OUTPUT/5698_Sunday_June_05_2022_03_47_35_PM_13412227/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.4. REGULAR SINGULAR POINTS. Page 175

Problem number: 3(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference is integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$x^3y'' + (-1 + \cos(2x))y' + 2xy = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^3y'' + (-1 + \cos(2x))y' + 2xy = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{-1 + \cos(2x)}{x^3}$$
$$q(x) = \frac{2}{x^2}$$

Table 448: Table $p(x), q(x)$ singularities.

$p(x) = \frac{-1+\cos(2x)}{x^3}$	
singularity	type
$x = 0$	“regular”

$q(x) = \frac{2}{x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^3 y'' + (-1 + \cos(2x)) y' + 2xy = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) x^3 \\ & + (-1 + \cos(2x)) \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + 2x \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \end{aligned} \tag{1}$$

Expanding $-1 + \cos(2x)$ as Taylor series around $x = 0$ and keeping only the first 8 terms gives

$$\begin{aligned} -1 + \cos(2x) &= -2x^2 + \frac{2}{3}x^4 - \frac{4}{45}x^6 + \frac{2}{315}x^8 + \dots \\ &= -2x^2 + \frac{2}{3}x^4 - \frac{4}{45}x^6 + \frac{2}{315}x^8 \end{aligned}$$

Which simplifies to

$$\begin{aligned} &\left(\sum_{n=0}^{\infty} x^{1+n+r} a_n(n+r)(n+r-1) \right) + \left(\sum_{n=0}^{\infty} \frac{2x^{n+r+7} a_n(n+r)}{315} \right) \\ &+ \sum_{n=0}^{\infty} \left(-\frac{4x^{n+r+5} a_n(n+r)}{45} \right) + \left(\sum_{n=0}^{\infty} \frac{2x^{n+r+3} a_n(n+r)}{3} \right) \\ &+ \sum_{n=0}^{\infty} (-2x^{1+n+r} a_n(n+r)) + \left(\sum_{n=0}^{\infty} 2x^{1+n+r} a_n \right) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $1 + n + r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{1+n+r} and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{2x^{n+r+7} a_n(n+r)}{315} &= \sum_{n=6}^{\infty} \frac{2a_{n-6}(-6+n+r)x^{1+n+r}}{315} \\ \sum_{n=0}^{\infty} \left(-\frac{4x^{n+r+5} a_n(n+r)}{45} \right) &= \sum_{n=4}^{\infty} \left(-\frac{4a_{n-4}(-4+n+r)x^{1+n+r}}{45} \right) \\ \sum_{n=0}^{\infty} \frac{2x^{n+r+3} a_n(n+r)}{3} &= \sum_{n=2}^{\infty} \frac{2a_{n-2}(n+r-2)x^{1+n+r}}{3} \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $1 + n + r$.

$$\begin{aligned} &\left(\sum_{n=0}^{\infty} x^{1+n+r} a_n(n+r)(n+r-1) \right) + \left(\sum_{n=6}^{\infty} \frac{2a_{n-6}(-6+n+r)x^{1+n+r}}{315} \right) \\ &+ \sum_{n=4}^{\infty} \left(-\frac{4a_{n-4}(-4+n+r)x^{1+n+r}}{45} \right) + \left(\sum_{n=2}^{\infty} \frac{2a_{n-2}(n+r-2)x^{1+n+r}}{3} \right) \\ &+ \sum_{n=0}^{\infty} (-2x^{1+n+r} a_n(n+r)) + \left(\sum_{n=0}^{\infty} 2x^{1+n+r} a_n \right) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n = 0$. From Eq (2B) this gives

$$x^{1+n+r} a_n (n+r) (n+r-1) - 2x^{1+n+r} a_n (n+r) + 2x^{1+n+r} a_n = 0$$

When $n = 0$ the above becomes

$$x^{1+r} a_0 r (-1+r) - 2x^{1+r} a_0 r + 2x^{1+r} a_0 = 0$$

Or

$$(x^{1+r} r (-1+r) - 2x^{1+r} r + 2x^{1+r}) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(r^2 - 3r + 2) x^{1+r} = 0$$

Since the above is true for all x then the indicial equation becomes

$$r^2 - 3r + 2 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = 2$$

$$r_2 = 1$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(r^2 - 3r + 2) x^{1+r} = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = 1$ is an integer, then we can construct two linearly independent solutions

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = C y_1(x) \ln(x) + x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = x^2 \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = C y_1(x) \ln(x) + x \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+2}$$

$$y_2(x) = C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{1+n} \right)$$

Where C above can be zero. We start by finding y_1 . Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

Substituting $n = 2$ in Eq. (2B) gives

$$a_2 = -\frac{2}{3r+3}$$

Substituting $n = 3$ in Eq. (2B) gives

$$a_3 = 0$$

Substituting $n = 4$ in Eq. (2B) gives

$$a_4 = \frac{\frac{4}{45}r^2 + \frac{8}{15}r + \frac{8}{9}}{(1+r)(3+r)(2+r)}$$

Substituting $n = 5$ in Eq. (2B) gives

$$a_5 = 0$$

For $6 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) + \frac{2a_{n-6}(-6+n+r)}{315} - \frac{4a_{n-4}(-4+n+r)}{45} \quad (3)$$

$$+ \frac{2a_{n-2}(n+r-2)}{3} - 2a_n(n+r) + 2a_n = 0$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{2(na_{n-6} - 14na_{n-4} + 105na_{n-2} + ra_{n-6} - 14ra_{n-4} + 105ra_{n-2} - 6a_{n-6} + 56a_{n-4} - 210a_{n-2})}{315(n^2 + 2nr + r^2 - 3n - 3r + 2)} \quad (4)$$

Which for the root $r = 2$ becomes

$$a_n = \frac{(-2a_{n-6} + 28a_{n-4} - 210a_{n-2})n + 8a_{n-6} - 56a_{n-4}}{315n(1+n)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 2$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{2}{3r+3}$	$-\frac{2}{9}$
a_3	0	0
a_4	$\frac{\frac{4}{45}r^2 + \frac{8}{15}r + \frac{8}{9}}{(1+r)(3+r)(2+r)}$	$\frac{26}{675}$
a_5	0	0

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{-\frac{2}{315}r^4 - \frac{148}{945}r^3 - \frac{1018}{945}r^2 - \frac{2836}{945}r - \frac{416}{135}}{(1+r)(3+r)(2+r)(5+r)(4+r)}$$

Which for the root $r = 2$ becomes

$$a_6 = -\frac{1742}{297675}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{2}{3r+3}$	$-\frac{2}{9}$
a_3	0	0
a_4	$\frac{\frac{4}{45}r^2 + \frac{8}{15}r + \frac{8}{9}}{(1+r)(3+r)(2+r)}$	$\frac{26}{675}$
a_5	0	0
a_6	$\frac{-\frac{2}{315}r^4 - \frac{148}{945}r^3 - \frac{1018}{945}r^2 - \frac{2836}{945}r - \frac{416}{135}}{(1+r)(3+r)(2+r)(5+r)(4+r)}$	$-\frac{1742}{297675}$

For $n = 7$, using the above recursive equation gives

$$a_7 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{2}{3r+3}$	$-\frac{2}{9}$
a_3	0	0
a_4	$\frac{\frac{4}{45}r^2 + \frac{8}{15}r + \frac{8}{9}}{(1+r)(3+r)(2+r)}$	$\frac{26}{675}$
a_5	0	0
a_6	$\frac{-\frac{2}{315}r^4 - \frac{148}{945}r^3 - \frac{1018}{945}r^2 - \frac{2836}{945}r - \frac{416}{135}}{(1+r)(3+r)(2+r)(5+r)(4+r)}$	$-\frac{1742}{297675}$
a_7	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x^2(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots) \\ &= x^2\left(1 - \frac{2x^2}{9} + \frac{26x^4}{675} - \frac{1742x^6}{297675} + O(x^8)\right) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Let

$$r_1 - r_2 = N$$

Where N is positive integer which is the difference between the two roots. r_1 is taken as the larger root. Hence for this problem we have $N = 1$. Now we need to determine if C is zero or not. This is done by finding $\lim_{r \rightarrow r_2} a_1(r)$. If this limit exists, then $C = 0$, else we need to keep the log term and $C \neq 0$. The above table shows that

$$\begin{aligned} a_N &= a_1 \\ &= 0 \end{aligned}$$

Therefore

$$\begin{aligned} \lim_{r \rightarrow r_2} 0 &= \lim_{r \rightarrow 1} 0 \\ &= 0 \end{aligned}$$

The limit is 0. Since the limit exists then the log term is not needed and we can set $C = 0$. Therefore the second solution has the form

$$\begin{aligned} y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+r} \\ &= \sum_{n=0}^{\infty} b_n x^{1+n} \end{aligned}$$

Eq (3) derived above is used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq(3) gives

$$b_1 = 0$$

Substituting $n = 2$ in Eq(3) gives

$$b_2 = -\frac{2}{3(1+r)}$$

Substituting $n = 3$ in Eq(3) gives

$$b_3 = 0$$

Substituting $n = 4$ in Eq(3) gives

$$b_4 = \frac{\frac{4}{45}r^2 + \frac{8}{15}r + \frac{8}{9}}{(1+r)(r^2 + 5r + 6)}$$

Substituting $n = 5$ in Eq(3) gives

$$b_5 = 0$$

For $6 \leq n$ the recursive equation is

$$\begin{aligned} b_n(n+r)(n+r-1) + \frac{2b_{n-6}(-6+n+r)}{315} - \frac{4b_{n-4}(-4+n+r)}{45} \\ + \frac{2b_{n-2}(n+r-2)}{3} - 2b_n(n+r) + 2b_n = 0 \end{aligned} \quad (4)$$

Which for the root $r = 1$ becomes

$$b_n(1+n)n + \frac{2b_{n-6}(-5+n)}{315} - \frac{4b_{n-4}(-3+n)}{45} + \frac{2b_{n-2}(n-1)}{3} - 2b_n(1+n) + 2b_n = 0 \quad (4A)$$

Solving for b_n from the recursive equation (4) gives

$$b_n = -\frac{2(nb_{n-6} - 14nb_{n-4} + 105nb_{n-2} + rb_{n-6} - 14rb_{n-4} + 105rb_{n-2} - 6b_{n-6} + 56b_{n-4} - 210b_{n-2})}{315(n^2 + 2nr + r^2 - 3n - 3r + 2)} \quad (5)$$

Which for the root $r = 1$ becomes

$$b_n = -\frac{2(nb_{n-6} - 14nb_{n-4} + 105nb_{n-2} - 5b_{n-6} + 42b_{n-4} - 105b_{n-2})}{315(n^2 - n)} \quad (6)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = 1$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{2}{3r+3}$	$-\frac{1}{3}$
b_3	0	0
b_4	$\frac{\frac{4}{45}r^2 + \frac{8}{15}r + \frac{8}{9}}{(1+r)(3+r)(2+r)}$	$\frac{17}{270}$
b_5	0	0

For $n = 6$, using the above recursive equation gives

$$b_6 = -\frac{2(3r^4 + 74r^3 + 509r^2 + 1418r + 1456)}{945(1+r)(r^2+5r+6)(r^2+9r+20)}$$

Which for the root $r = 1$ becomes

$$b_6 = -\frac{173}{17010}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{2}{3r+3}$	$-\frac{1}{3}$
b_3	0	0
b_4	$\frac{\frac{4}{45}r^2 + \frac{8}{15}r + \frac{8}{9}}{(1+r)(3+r)(2+r)}$	$\frac{17}{270}$
b_5	0	0
b_6	$\frac{-\frac{2}{315}r^4 - \frac{148}{945}r^3 - \frac{1018}{945}r^2 - \frac{2836}{945}r - \frac{416}{135}}{(1+r)(r^2+5r+6)(r^2+9r+20)}$	$-\frac{173}{17010}$

For $n = 7$, using the above recursive equation gives

$$b_7 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{2}{3r+3}$	$-\frac{1}{3}$
b_3	0	0
b_4	$\frac{\frac{4}{45}r^2 + \frac{8}{15}r + \frac{8}{9}}{(1+r)(3+r)(2+r)}$	$\frac{17}{270}$
b_5	0	0
b_6	$\frac{-\frac{2}{315}r^4 - \frac{148}{945}r^3 - \frac{1018}{945}r^2 - \frac{2836}{945}r - \frac{416}{135}}{(1+r)(r^2+5r+6)(r^2+9r+20)}$	$-\frac{173}{17010}$
b_7	0	0

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned} y_2(x) &= x^2(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 + b_7x^7 + b_8x^8 \dots) \\ &= x \left(1 - \frac{x^2}{3} + \frac{17x^4}{270} - \frac{173x^6}{17010} + O(x^8) \right) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= c_1x^2 \left(1 - \frac{2x^2}{9} + \frac{26x^4}{675} - \frac{1742x^6}{297675} + O(x^8) \right) + c_2x \left(1 - \frac{x^2}{3} + \frac{17x^4}{270} - \frac{173x^6}{17010} + O(x^8) \right) \end{aligned}$$

Hence the final solution is

$$\begin{aligned} y &= y_h \\ &= c_1x^2 \left(1 - \frac{2x^2}{9} + \frac{26x^4}{675} - \frac{1742x^6}{297675} + O(x^8) \right) + c_2x \left(1 - \frac{x^2}{3} + \frac{17x^4}{270} - \frac{173x^6}{17010} + O(x^8) \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 x^2 \left(1 - \frac{2x^2}{9} + \frac{26x^4}{675} - \frac{1742x^6}{297675} + O(x^8) \right) + c_2 x \left(1 - \frac{x^2}{3} + \frac{17x^4}{270} - \frac{173x^6}{17010} + O(x^8) \right) \quad (1)$$

Verification of solutions

$$y = c_1 x^2 \left(1 - \frac{2x^2}{9} + \frac{26x^4}{675} - \frac{1742x^6}{297675} + O(x^8) \right) + c_2 x \left(1 - \frac{x^2}{3} + \frac{17x^4}{270} - \frac{173x^6}{17010} + O(x^8) \right)$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) * 2F1([a
-> Trying changes of variables to rationalize or make the ODE simpler
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) * 2F1
trying a symmetry of the form [xi=0, eta=F(x)]
trying 2nd order exact linear
trying symmetries linear in x and y(x)
trying to convert to a linear ODE with constant coefficients
-> trying with_periodic_functions in the coefficients
--- Trying Lie symmetry methods, 2nd order ---
`, `-> Computing symmetries using: way = 5
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) * 2F1
trying a symmetry of the form [xi=0, eta=F(x)]
trying 2nd order exact linear
trying symmetries linear in x and y(x)
trying to convert to a linear ODE with constant coefficients
<- unable to find a useful change of variables
trying a symmetry of the form [xi=0, eta=F(x)]
trying differential order: 2; exact nonlinear
trying symmetries linear in x and y(x)
trying to convert to a linear ODE with constant coefficients
trying 2nd order, integrating factor of the form mu(x,y)
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) * 2F1
-> Trying changes of variables to rationalize or make the ODE simpler
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form r0(x) * Y + r1(x) * Y where Y = exp(int(r(x), dx)) *
trying a symmetry of the form [xi=0, eta=F(x)]
trying 2nd order exact linear
trying symmetries linear in x and y(x)
trying to convert to a linear ODE with constant coefficients
```

✓ Solution by Maple

Time used: 0.235 (sec). Leaf size: 37

Order:=8;

```
dsolve(x^3*diff(y(x),x$2)+(cos(2*x)-1)*diff(y(x),x)+2*x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^2 \left(1 - \frac{2}{9} x^2 + \frac{26}{675} x^4 - \frac{1742}{297675} x^6 + O(x^8) \right) \\ + c_2 x \left(1 - \frac{1}{3} x^2 + \frac{17}{270} x^4 - \frac{173}{17010} x^6 + O(x^8) \right)$$

✓ Solution by Mathematica

Time used: 0.037 (sec). Leaf size: 74

```
AsymptoticDSolveValue[x^3*y'[x]+(Cos[2*x]-1)*y'[x]+2*x*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_2 \left(\frac{32351x^8}{40186125} - \frac{1742x^6}{297675} + \frac{26x^4}{675} - \frac{2x^2}{9} + 1 \right) x^2 \\ + c_1 \left(\frac{10471x^8}{7144200} - \frac{173x^6}{17010} + \frac{17x^4}{270} - \frac{x^2}{3} + 1 \right) x$$

19.11 problem 3(b)

19.11.1 Maple step by step solution 3239

Internal problem ID [6451]

Internal file name [OUTPUT/5699_Sunday_June_05_2022_03_47_40_PM_73099135/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.4. REGULAR SINGULAR POINTS. Page 175

Problem number: 3(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$4x^2y'' + (2x^4 - 5x)y' + (3x^2 + 2)y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$4x^2y'' + (2x^4 - 5x)y' + (3x^2 + 2)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{2x^3 - 5}{4x}$$
$$q(x) = \frac{3x^2 + 2}{4x^2}$$

Table 449: Table $p(x), q(x)$ singularities.

$p(x) = \frac{2x^3-5}{4x}$	
singularity	type
$x = 0$	“regular”
$x = \infty$	“regular”
$x = -\infty$	“regular”

$q(x) = \frac{3x^2+2}{4x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0, \infty, -\infty]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$4x^2y'' + (2x^4 - 5x)y' + (3x^2 + 2)y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$4x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) + (2x^4 - 5x) \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (3x^2 + 2) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 4x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} 2x^{n+r+3} a_n (n+r) \right) \\ & + \sum_{n=0}^{\infty} (-5x^{n+r} a_n (n+r)) + \left(\sum_{n=0}^{\infty} 3x^{n+r+2} a_n \right) + \left(\sum_{n=0}^{\infty} 2a_n x^{n+r} \right) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=0}^{\infty} 2x^{n+r+3} a_n (n+r) &= \sum_{n=3}^{\infty} 2a_{n-3} (n-3+r) x^{n+r} \\ \sum_{n=0}^{\infty} 3x^{n+r+2} a_n &= \sum_{n=2}^{\infty} 3a_{n-2} x^{n+r} \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 4x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=3}^{\infty} 2a_{n-3} (n-3+r) x^{n+r} \right) \\ & + \sum_{n=0}^{\infty} (-5x^{n+r} a_n (n+r)) + \left(\sum_{n=2}^{\infty} 3a_{n-2} x^{n+r} \right) + \left(\sum_{n=0}^{\infty} 2a_n x^{n+r} \right) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$4x^{n+r} a_n (n+r) (n+r-1) - 5x^{n+r} a_n (n+r) + 2a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$4x^r a_0 r (-1+r) - 5x^r a_0 r + 2a_0 x^r = 0$$

Or

$$(4x^r r (-1+r) - 5x^r r + 2x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(4r^2 - 9r + 2) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$4r^2 - 9r + 2 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 2 \\ r_2 &= \frac{1}{4} \end{aligned}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(4r^2 - 9r + 2) x^r = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = \frac{7}{4}$ is not an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^{n+2} \\ y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+\frac{1}{4}} \end{aligned}$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

Substituting $n = 2$ in Eq. (2B) gives

$$a_2 = -\frac{3}{r(4r+7)}$$

For $3 \leq n$ the recursive equation is

$$4a_n(n+r)(n+r-1) + 2a_{n-3}(n-3+r) - 5a_n(n+r) + 3a_{n-2} + 2a_n = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{2na_{n-3} + 2ra_{n-3} - 6a_{n-3} + 3a_{n-2}}{4n^2 + 8nr + 4r^2 - 9n - 9r + 2} \quad (4)$$

Which for the root $r = 2$ becomes

$$a_n = \frac{-2na_{n-3} + 2a_{n-3} - 3a_{n-2}}{n(4n + 7)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 2$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{3}{r(4r+7)}$	$-\frac{1}{10}$

For $n = 3$, using the above recursive equation gives

$$a_3 = -\frac{2r}{4r^2 + 15r + 11}$$

Which for the root $r = 2$ becomes

$$a_3 = -\frac{4}{57}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{3}{r(4r+7)}$	$-\frac{1}{10}$
a_3	$-\frac{2r}{4r^2+15r+11}$	$-\frac{4}{57}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{9}{r(4r + 7)(4r^2 + 23r + 30)}$$

Which for the root $r = 2$ becomes

$$a_4 = \frac{3}{920}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{3}{r(4r+7)}$	$-\frac{1}{10}$
a_3	$-\frac{2r}{4r^2+15r+11}$	$-\frac{4}{57}$
a_4	$\frac{9}{r(4r+7)(4r^2+23r+30)}$	$\frac{3}{920}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{48r^3 + 180r^2 + 246r + 132}{(4r + 7)r(4r^2 + 15r + 11)(4r^2 + 31r + 57)}$$

Which for the root $r = 2$ becomes

$$a_5 = \frac{32}{4275}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{3}{r(4r+7)}$	$-\frac{1}{10}$
a_3	$-\frac{2r}{4r^2+15r+11}$	$-\frac{4}{57}$
a_4	$\frac{9}{r(4r+7)(4r^2+23r+30)}$	$\frac{3}{920}$
a_5	$\frac{48r^3+180r^2+246r+132}{(4r+7)r(4r^2+15r+11)(4r^2+31r+57)}$	$\frac{32}{4275}$

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{64r^6 + 672r^5 + 2564r^4 + 4212r^3 + 2412r^2 - 405r - 297}{(4r^2 + 15r + 11)r(4r + 7)(4r^2 + 23r + 30)(4r^2 + 39r + 92)}$$

Which for the root $r = 2$ becomes

$$a_6 = \frac{36287}{9753840}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{3}{r(4r+7)}$	$-\frac{1}{10}$
a_3	$-\frac{2r}{4r^2+15r+11}$	$-\frac{4}{57}$
a_4	$\frac{9}{r(4r+7)(4r^2+23r+30)}$	$\frac{3}{920}$
a_5	$\frac{48r^3+180r^2+246r+132}{(4r+7)r(4r^2+15r+11)(4r^2+31r+57)}$	$\frac{32}{4275}$
a_6	$\frac{64r^6+672r^5+2564r^4+4212r^3+2412r^2-405r-297}{(4r^2+15r+11)r(4r+7)(4r^2+23r+30)(4r^2+39r+92)}$	$\frac{36287}{9753840}$

For $n = 7$, using the above recursive equation gives

$$a_7 = -\frac{18(48r^5 + 552r^4 + 2567r^3 + 6075r^2 + 7147r + 3168)}{(4r + 7)(4r^2 + 23r + 30)r(4r^2 + 15r + 11)(4r^2 + 31r + 57)(4r^2 + 47r + 135)}$$

Which for the root $r = 2$ becomes

$$a_7 = -\frac{4037}{16059750}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{3}{r(4r+7)}$	$-\frac{1}{10}$
a_3	$-\frac{2r}{4r^2+15r+11}$	$-\frac{4}{57}$
a_4	$\frac{9}{r(4r+7)(4r^2+23r+30)}$	$\frac{3}{920}$
a_5	$\frac{48r^3+180r^2+246r+132}{(4r+7)r(4r^2+15r+11)(4r^2+31r+57)}$	$\frac{32}{4275}$
a_6	$\frac{64r^6+672r^5+2564r^4+4212r^3+2412r^2-405r-297}{(4r^2+15r+11)r(4r+7)(4r^2+23r+30)(4r^2+39r+92)}$	$\frac{36287}{9753840}$
a_7	$-\frac{18(48r^5+552r^4+2567r^3+6075r^2+7147r+3168)}{(4r+7)(4r^2+23r+30)r(4r^2+15r+11)(4r^2+31r+57)(4r^2+47r+135)}$	$-\frac{4037}{16059750}$

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x^2(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots) \\ &= x^2\left(1 - \frac{x^2}{10} - \frac{4x^3}{57} + \frac{3x^4}{920} + \frac{32x^5}{4275} + \frac{36287x^6}{9753840} - \frac{4037x^7}{16059750} + O(x^8)\right) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$b_1 = 0$$

Substituting $n = 2$ in Eq. (2B) gives

$$b_2 = -\frac{3}{r(4r+7)}$$

For $3 \leq n$ the recursive equation is

$$4b_n(n+r)(n+r-1) + 2b_{n-3}(n-3+r) - 5b_n(n+r) + 3b_{n-2} + 2b_n = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = -\frac{2nb_{n-3} + 2rb_{n-3} - 6b_{n-3} + 3b_{n-2}}{4n^2 + 8nr + 4r^2 - 9n - 9r + 2} \quad (4)$$

Which for the root $r = \frac{1}{4}$ becomes

$$b_n = \frac{-4nb_{n-3} + 11b_{n-3} - 6b_{n-2}}{8n^2 - 14n} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = \frac{1}{4}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{3}{r(4r+7)}$	$-\frac{3}{2}$

For $n = 3$, using the above recursive equation gives

$$b_3 = -\frac{2r}{4r^2 + 15r + 11}$$

Which for the root $r = \frac{1}{4}$ becomes

$$b_3 = -\frac{1}{30}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{3}{r(4r+7)}$	$-\frac{3}{2}$
b_3	$-\frac{2r}{4r^2+15r+11}$	$-\frac{1}{30}$

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{9}{r(4r+7)(4r^2+23r+30)}$$

Which for the root $r = \frac{1}{4}$ becomes

$$b_4 = \frac{1}{8}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{3}{r(4r+7)}$	$-\frac{3}{2}$
b_3	$-\frac{2r}{4r^2+15r+11}$	$-\frac{1}{30}$
b_4	$\frac{9}{r(4r+7)(4r^2+23r+30)}$	$\frac{1}{8}$

For $n = 5$, using the above recursive equation gives

$$b_5 = \frac{48r^3 + 180r^2 + 246r + 132}{(4r+7)r(4r^2+15r+11)(4r^2+31r+57)}$$

Which for the root $r = \frac{1}{4}$ becomes

$$b_5 = \frac{137}{1300}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{3}{r(4r+7)}$	$-\frac{3}{2}$
b_3	$-\frac{2r}{4r^2+15r+11}$	$-\frac{1}{30}$
b_4	$\frac{9}{r(4r+7)(4r^2+23r+30)}$	$\frac{1}{8}$
b_5	$\frac{48r^3+180r^2+246r+132}{(4r+7)r(4r^2+15r+11)(4r^2+31r+57)}$	$\frac{137}{1300}$

For $n = 6$, using the above recursive equation gives

$$b_6 = \frac{64r^6 + 672r^5 + 2564r^4 + 4212r^3 + 2412r^2 - 405r - 297}{(4r^2 + 15r + 11)r(4r + 7)(4r^2 + 23r + 30)(4r^2 + 39r + 92)}$$

Which for the root $r = \frac{1}{4}$ becomes

$$b_6 = -\frac{19}{12240}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{3}{r(4r+7)}$	$-\frac{3}{2}$
b_3	$-\frac{2r}{4r^2+15r+11}$	$-\frac{1}{30}$
b_4	$\frac{9}{r(4r+7)(4r^2+23r+30)}$	$\frac{1}{8}$
b_5	$\frac{48r^3+180r^2+246r+132}{(4r+7)r(4r^2+15r+11)(4r^2+31r+57)}$	$\frac{137}{1300}$
b_6	$\frac{64r^6+672r^5+2564r^4+4212r^3+2412r^2-405r-297}{(4r^2+15r+11)r(4r+7)(4r^2+23r+30)(4r^2+39r+92)}$	$-\frac{19}{12240}$

For $n = 7$, using the above recursive equation gives

$$b_7 = -\frac{18(48r^5 + 552r^4 + 2567r^3 + 6075r^2 + 7147r + 3168)}{(4r + 7)(4r^2 + 23r + 30)r(4r^2 + 15r + 11)(4r^2 + 31r + 57)(4r^2 + 47r + 135)}$$

Which for the root $r = \frac{1}{4}$ becomes

$$b_7 = -\frac{7169}{764400}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{3}{r(4r+7)}$	$-\frac{3}{2}$
b_3	$-\frac{2r}{4r^2+15r+11}$	$-\frac{1}{30}$
b_4	$\frac{9}{r(4r+7)(4r^2+23r+30)}$	$\frac{1}{8}$
b_5	$\frac{48r^3+180r^2+246r+132}{(4r+7)r(4r^2+15r+11)(4r^2+31r+57)}$	$\frac{137}{1300}$
b_6	$\frac{64r^6+672r^5+2564r^4+4212r^3+2412r^2-405r-297}{(4r^2+15r+11)r(4r+7)(4r^2+23r+30)(4r^2+39r+92)}$	$-\frac{19}{12240}$
b_7	$-\frac{18(48r^5+552r^4+2567r^3+6075r^2+7147r+3168)}{(4r+7)(4r^2+23r+30)r(4r^2+15r+11)(4r^2+31r+57)(4r^2+47r+135)}$	$-\frac{7169}{764400}$

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned} y_2(x) &= x^2(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 + b_7x^7 + b_8x^8 \dots) \\ &= x^{\frac{1}{4}} \left(1 - \frac{3x^2}{2} - \frac{x^3}{30} + \frac{x^4}{8} + \frac{137x^5}{1300} - \frac{19x^6}{12240} - \frac{7169x^7}{764400} + O(x^8) \right) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= c_1x^2 \left(1 - \frac{x^2}{10} - \frac{4x^3}{57} + \frac{3x^4}{920} + \frac{32x^5}{4275} + \frac{36287x^6}{9753840} - \frac{4037x^7}{16059750} + O(x^8) \right) \\ &\quad + c_2x^{\frac{1}{4}} \left(1 - \frac{3x^2}{2} - \frac{x^3}{30} + \frac{x^4}{8} + \frac{137x^5}{1300} - \frac{19x^6}{12240} - \frac{7169x^7}{764400} + O(x^8) \right) \end{aligned}$$

Hence the final solution is

$$\begin{aligned} y &= y_h \\ &= c_1x^2 \left(1 - \frac{x^2}{10} - \frac{4x^3}{57} + \frac{3x^4}{920} + \frac{32x^5}{4275} + \frac{36287x^6}{9753840} - \frac{4037x^7}{16059750} + O(x^8) \right) \\ &\quad + c_2x^{\frac{1}{4}} \left(1 - \frac{3x^2}{2} - \frac{x^3}{30} + \frac{x^4}{8} + \frac{137x^5}{1300} - \frac{19x^6}{12240} - \frac{7169x^7}{764400} + O(x^8) \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 x^2 \left(1 - \frac{x^2}{10} - \frac{4x^3}{57} + \frac{3x^4}{920} + \frac{32x^5}{4275} + \frac{36287x^6}{9753840} - \frac{4037x^7}{16059750} + O(x^8) \right) \\ + c_2 x^{\frac{1}{4}} \left(1 - \frac{3x^2}{2} - \frac{x^3}{30} + \frac{x^4}{8} + \frac{137x^5}{1300} - \frac{19x^6}{12240} - \frac{7169x^7}{764400} + O(x^8) \right) \quad (1)$$

Verification of solutions

$$y = c_1 x^2 \left(1 - \frac{x^2}{10} - \frac{4x^3}{57} + \frac{3x^4}{920} + \frac{32x^5}{4275} + \frac{36287x^6}{9753840} - \frac{4037x^7}{16059750} + O(x^8) \right) \\ + c_2 x^{\frac{1}{4}} \left(1 - \frac{3x^2}{2} - \frac{x^3}{30} + \frac{x^4}{8} + \frac{137x^5}{1300} - \frac{19x^6}{12240} - \frac{7169x^7}{764400} + O(x^8) \right)$$

Verified OK.

19.11.1 Maple step by step solution

Let's solve

$$4x^2 y'' + (2x^4 - 5x) y' + (3x^2 + 2) y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{(3x^2+2)y}{4x^2} - \frac{(2x^3-5)y'}{4x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{(2x^3-5)y'}{4x} + \frac{(3x^2+2)y}{4x^2} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$\left[P_2(x) = \frac{2x^3-5}{4x}, P_3(x) = \frac{3x^2+2}{4x^2} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = -\frac{5}{4}$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = \frac{1}{2}$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$4x^2y'' + x(2x^3 - 5)y' + (3x^2 + 2)y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 0..2$

$$x^m \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

- Shift index using $k \rightarrow k - m$

$$x^m \cdot y = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

- Convert $x^m \cdot y'$ to series expansion for $m = 1..4$

$$x^m \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1+m}$$

- Shift index using $k \rightarrow k + 1 - m$

$$x^m \cdot y' = \sum_{k=-1+m}^{\infty} a_{k+1-m} (k+1-m+r) x^{k+r}$$

- Convert $x^2 \cdot y''$ to series expansion

$$x^2 \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(-1+4r)(-2+r)x^r + a_1(3+4r)(-1+r)x^{1+r} + (a_2(7+4r)r + 3a_0)x^{2+r} + \left(\sum_{k=3}^{\infty} (a_k(4k + \dots) \right)$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$(-1+4r)(-2+r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \left\{ 2, \frac{1}{4} \right\}$$

- The coefficients of each power of x must be 0

$$[a_1(3 + 4r)(-1 + r) = 0, a_2(7 + 4r)r + 3a_0 = 0]$$

- Solve for the dependent coefficient(s)

$$\left\{ a_1 = 0, a_2 = -\frac{3a_0}{r(7+4r)} \right\}$$

- Each term in the series must be 0, giving the recursion relation

$$a_k(4k + 4r - 1)(k + r - 2) + 3a_{k-2} + 2a_{k-3}(k - 3 + r) = 0$$

- Shift index using $k \rightarrow k + 3$

$$a_{k+3}(4k + 11 + 4r)(k + 1 + r) + 3a_{k+1} + 2a_k(k + r) = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+3} = -\frac{2ka_k + 2ra_k + 3a_{k+1}}{(4k+11+4r)(k+1+r)}$$

- Recursion relation for $r = 2$

$$a_{k+3} = -\frac{2ka_k + 4a_k + 3a_{k+1}}{(4k+19)(k+3)}$$

- Solution for $r = 2$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+2}, a_{k+3} = -\frac{2ka_k + 4a_k + 3a_{k+1}}{(4k+19)(k+3)}, a_1 = 0, a_2 = -\frac{a_0}{10} \right]$$

- Recursion relation for $r = \frac{1}{4}$

$$a_{k+3} = -\frac{2ka_k + \frac{1}{2}a_k + 3a_{k+1}}{(4k+12)(k+\frac{5}{4})}$$

- Solution for $r = \frac{1}{4}$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+\frac{1}{4}}, a_{k+3} = -\frac{2ka_k + \frac{1}{2}a_k + 3a_{k+1}}{(4k+12)(k+\frac{5}{4})}, a_1 = 0, a_2 = -\frac{3a_0}{2} \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^{k+2} \right) + \left(\sum_{k=0}^{\infty} b_k x^{k+\frac{1}{4}} \right), a_{k+3} = -\frac{2ka_k + 4a_k + 3a_{k+1}}{(4k+19)(k+3)}, a_1 = 0, a_2 = -\frac{a_0}{10}, b_{k+3} = -\frac{2kb_k + \frac{1}{2}b_k}{(4k+12)} \right]$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
  -> Bessel
  -> elliptic
  -> Legendre
  -> Whittaker
    -> hyper3: Equivalence to 1F1 under a power @ Moebius
  -> hypergeometric
    -> heuristic approach
    -> hyper3: Equivalence to 2F1, 1F1 or 0F1 under a power @ Moebius
  -> Mathieu
    -> Equivalence to the rational form of Mathieu ODE under a power @ Moebius
trying a solution in terms of MeijerG functions
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
-> trying a solution of the form  $r_0(x) * Y + r_1(x) * Y$  where  $Y = \exp(\int(r(x), dx)) * 2F1([a$ 
  trying a symmetry of the form [xi=0, eta=F(x)]
  trying differential order: 2; exact nonlinear
  trying symmetries linear in x and y(x)
  trying to convert to a linear ODE with constant coefficients
  trying 2nd order, integrating factor of the form  $\mu(x,y)$ 
  -> Trying a solution in terms of special functions:
    -> Bessel
    -> elliptic
    -> Legendre
    -> Whittaker
      -> hyper3: Equivalence to 1F1 under a power @ Moebius
    -> hypergeometric
      -> heuristic approach
      -> hyper3: Equivalence to 2F1, 1F1 or 0F1 under a power @ Moebius
    -> Mathieu
      -> Equivalence to the rational form of Mathieu ODE under a power @ Moebius
  trying 2nd order exact linear
  trying symmetries linear in x and y(x)
  trying to convert to a linear ODE with constant coefficients
  trying to convert to an ODE of Bessel type
trying to convert to an ODE of Bessel type
-> trying reduction of order to Bessel
```

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 51

Order:=8;

```
dsolve(4*x^2*diff(y(x),x$2)+(2*x^4-5*x)*diff(y(x),x)+(3*x^2+2)*y(x)=0,y(x),type='series',x=0
```

$$y(x) = c_1 x^{\frac{1}{4}} \left(1 - \frac{3}{2}x^2 - \frac{1}{30}x^3 + \frac{1}{8}x^4 + \frac{137}{1300}x^5 - \frac{19}{12240}x^6 - \frac{7169}{764400}x^7 + O(x^8) \right) \\ + c_2 x^2 \left(1 - \frac{1}{10}x^2 - \frac{4}{57}x^3 + \frac{3}{920}x^4 + \frac{32}{4275}x^5 + \frac{36287}{9753840}x^6 - \frac{4037}{16059750}x^7 + O(x^8) \right)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 106

```
AsymptoticDSolveValue[4*x^2*y'[x]+(2*x^4-5*x)*y'[x]+(3*x^2+2)*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(-\frac{4037x^7}{16059750} + \frac{36287x^6}{9753840} + \frac{32x^5}{4275} + \frac{3x^4}{920} - \frac{4x^3}{57} - \frac{x^2}{10} + 1 \right) x^2 \\ + c_2 \left(-\frac{7169x^7}{764400} - \frac{19x^6}{12240} + \frac{137x^5}{1300} + \frac{x^4}{8} - \frac{x^3}{30} - \frac{3x^2}{2} + 1 \right) \sqrt[4]{x}$$

19.12 problem 3(c)

19.12.1 Maple step by step solution 3258

Internal problem ID [6452]

Internal file name [OUTPUT/5700_Sunday_June_05_2022_03_47_44_PM_82017572/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.4. REGULAR SINGULAR POINTS. Page 175

Problem number: 3(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference is integer"**

Maple gives the following as the ode type

```
[[_Emden , _Fowler]]
```

$$x^2y'' + 3xy' + 4xy = 0$$

With the expansion point for the power series method at $x = 0$.

The ODE is

$$x^2y'' + 3xy' + 4xy = 0$$

Or

$$x(y''x + 4y + 3y') = 0$$

For $x \neq 0$ the above simplifies to

$$y''x + 4y + 3y' = 0$$

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2y'' + 3xy' + 4xy = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{3}{x}$$

$$q(x) = \frac{4}{x}$$

Table 451: Table $p(x), q(x)$ singularities.

$p(x) = \frac{3}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = \frac{4}{x}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2 y'' + 3xy' + 4xy = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned}
 & x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\
 & + 3x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + 4x \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0
 \end{aligned} \tag{1}$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r)(n+r-1) \right) + \left(\sum_{n=0}^{\infty} 3x^{n+r} a_n (n+r) \right) + \left(\sum_{n=0}^{\infty} 4x^{1+n+r} a_n \right) = 0 \tag{2A}$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} 4x^{1+n+r} a_n = \sum_{n=1}^{\infty} 4a_{n-1} x^{n+r}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r)(n+r-1) \right) + \left(\sum_{n=0}^{\infty} 3x^{n+r} a_n (n+r) \right) + \left(\sum_{n=1}^{\infty} 4a_{n-1} x^{n+r} \right) = 0 \tag{2B}$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r} a_n (n+r)(n+r-1) + 3x^{n+r} a_n (n+r) = 0$$

When $n=0$ the above becomes

$$x^r a_0 r(-1+r) + 3x^r a_0 r = 0$$

Or

$$(x^r r(-1+r) + 3x^r r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$x^r r(2+r) = 0$$

Since the above is true for all x then the indicial equation becomes

$$r(2 + r) = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 0 \\ r_2 &= -2 \end{aligned}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$x^r r(2 + r) = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = 2$ is an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= C y_1(x) \ln(x) + x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^n \\ y_2(x) &= C y_1(x) \ln(x) + \frac{\sum_{n=0}^{\infty} b_n x^n}{x^2} \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^n \\ y_2(x) &= C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n-2} \right) \end{aligned}$$

Where C above can be zero. We start by finding y_1 . Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$a_n(n + r)(n + r - 1) + 3a_n(n + r) + 4a_{n-1} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{4a_{n-1}}{n^2 + 2nr + r^2 + 2n + 2r} \quad (4)$$

Which for the root $r = 0$ becomes

$$a_n = -\frac{4a_{n-1}}{n(n+2)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = -\frac{4}{r^2 + 4r + 3}$$

Which for the root $r = 0$ becomes

$$a_1 = -\frac{4}{3}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{4}{r^2+4r+3}$	$-\frac{4}{3}$

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{16}{(r^2 + 4r + 3)(r^2 + 6r + 8)}$$

Which for the root $r = 0$ becomes

$$a_2 = \frac{2}{3}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{4}{r^2+4r+3}$	$-\frac{4}{3}$
a_2	$\frac{16}{(r^2+4r+3)(r^2+6r+8)}$	$\frac{2}{3}$

For $n = 3$, using the above recursive equation gives

$$a_3 = -\frac{64}{(r+3)^2(r+1)(r+4)(2+r)(r+5)}$$

Which for the root $r = 0$ becomes

$$a_3 = -\frac{8}{45}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{4}{r^2+4r+3}$	$-\frac{4}{3}$
a_2	$\frac{16}{(r^2+4r+3)(r^2+6r+8)}$	$\frac{2}{3}$
a_3	$-\frac{64}{(r+3)^2(r+1)(r+4)(2+r)(r+5)}$	$-\frac{8}{45}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{256}{(r+3)^2(r+1)(r+4)^2(2+r)(r+5)(r+6)}$$

Which for the root $r = 0$ becomes

$$a_4 = \frac{4}{135}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{4}{r^2+4r+3}$	$-\frac{4}{3}$
a_2	$\frac{16}{(r^2+4r+3)(r^2+6r+8)}$	$\frac{2}{3}$
a_3	$-\frac{64}{(r+3)^2(r+1)(r+4)(2+r)(r+5)}$	$-\frac{8}{45}$
a_4	$\frac{256}{(r+3)^2(r+1)(r+4)^2(2+r)(r+5)(r+6)}$	$\frac{4}{135}$

For $n = 5$, using the above recursive equation gives

$$a_5 = -\frac{1024}{(r+3)^2(r+1)(r+4)^2(2+r)(r+5)^2(r+6)(r+7)}$$

Which for the root $r = 0$ becomes

$$a_5 = -\frac{16}{4725}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{4}{r^2+4r+3}$	$-\frac{4}{3}$
a_2	$\frac{16}{(r^2+4r+3)(r^2+6r+8)}$	$\frac{2}{3}$
a_3	$-\frac{64}{(r+3)^2(r+1)(r+4)(2+r)(r+5)}$	$-\frac{8}{45}$
a_4	$\frac{256}{(r+3)^2(r+1)(r+4)^2(2+r)(r+5)(r+6)}$	$\frac{4}{135}$
a_5	$-\frac{1024}{(r+3)^2(r+1)(r+4)^2(2+r)(r+5)^2(r+6)(r+7)}$	$-\frac{16}{4725}$

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{4096}{(r+3)^2(r+1)(r+4)^2(2+r)(r+5)^2(r+6)^2(r+7)(r+8)}$$

Which for the root $r = 0$ becomes

$$a_6 = \frac{4}{14175}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{4}{r^2+4r+3}$	$-\frac{4}{3}$
a_2	$\frac{16}{(r^2+4r+3)(r^2+6r+8)}$	$\frac{2}{3}$
a_3	$-\frac{64}{(r+3)^2(r+1)(r+4)(2+r)(r+5)}$	$-\frac{8}{45}$
a_4	$\frac{256}{(r+3)^2(r+1)(r+4)^2(2+r)(r+5)(r+6)}$	$\frac{4}{135}$
a_5	$-\frac{1024}{(r+3)^2(r+1)(r+4)^2(2+r)(r+5)^2(r+6)(r+7)}$	$-\frac{16}{4725}$
a_6	$\frac{4096}{(r+3)^2(r+1)(r+4)^2(2+r)(r+5)^2(r+6)^2(r+7)(r+8)}$	$\frac{4}{14175}$

For $n = 7$, using the above recursive equation gives

$$a_7 = -\frac{16384}{(r+3)^2(r+1)(r+4)^2(2+r)(r+5)^2(r+6)^2(r+7)^2(r+8)(r+9)}$$

Which for the root $r = 0$ becomes

$$a_7 = -\frac{16}{893025}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{4}{r^2+4r+3}$	$-\frac{4}{3}$
a_2	$\frac{16}{(r^2+4r+3)(r^2+6r+8)}$	$\frac{2}{3}$
a_3	$-\frac{64}{(r+3)^2(r+1)(r+4)(2+r)(r+5)}$	$-\frac{8}{45}$
a_4	$\frac{256}{(r+3)^2(r+1)(r+4)^2(2+r)(r+5)(r+6)}$	$\frac{4}{135}$
a_5	$-\frac{1024}{(r+3)^2(r+1)(r+4)^2(2+r)(r+5)^2(r+6)(r+7)}$	$-\frac{16}{4725}$
a_6	$\frac{4096}{(r+3)^2(r+1)(r+4)^2(2+r)(r+5)^2(r+6)^2(r+7)(r+8)}$	$\frac{4}{14175}$
a_7	$-\frac{16384}{(r+3)^2(r+1)(r+4)^2(2+r)(r+5)^2(r+6)^2(r+7)^2(r+8)(r+9)}$	$-\frac{16}{893025}$

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots \\ &= 1 - \frac{4x}{3} + \frac{2x^2}{3} - \frac{8x^3}{45} + \frac{4x^4}{135} - \frac{16x^5}{4725} + \frac{4x^6}{14175} - \frac{16x^7}{893025} + O(x^8) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Let

$$r_1 - r_2 = N$$

Where N is positive integer which is the difference between the two roots. r_1 is taken as the larger root. Hence for this problem we have $N = 2$. Now we need to determine if C is zero or not. This is done by finding $\lim_{r \rightarrow r_2} a_2(r)$. If this limit exists, then $C = 0$, else we need to keep the log term and $C \neq 0$. The above table shows that

$$\begin{aligned} a_N &= a_2 \\ &= \frac{16}{(r^2 + 4r + 3)(r^2 + 6r + 8)} \end{aligned}$$

Therefore

$$\begin{aligned} \lim_{r \rightarrow r_2} \frac{16}{(r^2 + 4r + 3)(r^2 + 6r + 8)} &= \lim_{r \rightarrow -2} \frac{16}{(r^2 + 4r + 3)(r^2 + 6r + 8)} \\ &= \text{undefined} \end{aligned}$$

Since the limit does not exist then the log term is needed. Therefore the second solution has the form

$$y_2(x) = Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right)$$

Therefore

$$\begin{aligned} \frac{d}{dx} y_2(x) &= Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2} (n+r_2)}{x} \right) \\ &= Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} x^{-1+n+r_2} b_n (n+r_2) \right) \\ \frac{d^2}{dx^2} y_2(x) &= Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} \\ &\quad + \sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2} (n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2} (n+r_2)}{x^2} \right) \\ &= Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} + \left(\sum_{n=0}^{\infty} x^{-2+n+r_2} b_n (n+r_2) (-1+n+r_2) \right) \end{aligned}$$

Substituting these back into the given ode $x^2 y'' + 3xy' + 4xy = 0$ gives

$$\begin{aligned} &x^2 \left(Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} \right. \\ &\quad \left. + \sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2} (n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2} (n+r_2)}{x^2} \right) \right) \\ &\quad + 3x \left(Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2} (n+r_2)}{x} \right) \right) \\ &\quad + 4x \left(Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) \right) = 0 \end{aligned}$$

Which can be written as

$$\begin{aligned} & \left((y_1''(x)x^2 + 4y_1(x)x + 3y_1'(x)x) \ln(x) + x^2 \left(\frac{2y_1'(x)}{x} - \frac{y_1(x)}{x^2} \right) + 3y_1(x) \right) C \\ & + x^2 \left(\sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2} (n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2} (n+r_2)}{x^2} \right) \right) \\ & + 3x \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2} (n+r_2)}{x} \right) + 4x \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) = 0 \end{aligned} \quad (7)$$

But since $y_1(x)$ is a solution to the ode, then

$$y_1''(x)x^2 + 4y_1(x)x + 3y_1'(x)x = 0$$

Eq (7) simplifies to

$$\begin{aligned} & \left(x^2 \left(\frac{2y_1'(x)}{x} - \frac{y_1(x)}{x^2} \right) + 3y_1(x) \right) C \\ & + x^2 \left(\sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2} (n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2} (n+r_2)}{x^2} \right) \right) \\ & + 3x \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2} (n+r_2)}{x} \right) + 4x \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) = 0 \end{aligned} \quad (8)$$

Substituting $y_1 = \sum_{n=0}^{\infty} a_n x^{n+r_1}$ into the above gives

$$\begin{aligned} & \left(2 \left(\sum_{n=0}^{\infty} x^{-1+n+r_1} a_n (n+r_1) \right) x + 2 \left(\sum_{n=0}^{\infty} a_n x^{n+r_1} \right) \right) C \\ & + \left(\sum_{n=0}^{\infty} x^{-2+n+r_2} b_n (n+r_2) (-1+n+r_2) \right) x^2 \\ & + 3 \left(\sum_{n=0}^{\infty} x^{-1+n+r_2} b_n (n+r_2) \right) x + 4x \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) = 0 \end{aligned} \quad (9)$$

Since $r_1 = 0$ and $r_2 = -2$ then the above becomes

$$\begin{aligned} & \left(2 \left(\sum_{n=0}^{\infty} x^{n-1} a_n n \right) x + 2 \left(\sum_{n=0}^{\infty} a_n x^n \right) \right) C + \left(\sum_{n=0}^{\infty} x^{-4+n} b_n (n-2) (-3+n) \right) x^2 \\ & + 3 \left(\sum_{n=0}^{\infty} x^{-3+n} b_n (n-2) \right) x + 4x \left(\sum_{n=0}^{\infty} b_n x^{n-2} \right) = 0 \end{aligned} \quad (10)$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2Cn x^n a_n \right) + \left(\sum_{n=0}^{\infty} 2C a_n x^n \right) + \left(\sum_{n=0}^{\infty} x^{n-2} b_n (n^2 - 5n + 6) \right) \\ & + \left(\sum_{n=0}^{\infty} 3x^{n-2} b_n (n-2) \right) + \left(\sum_{n=0}^{\infty} 4x^{n-1} b_n \right) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n - 2$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n-2} and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=0}^{\infty} 2Cn x^n a_n &= \sum_{n=2}^{\infty} 2C(n-2) a_{n-2} x^{n-2} \\ \sum_{n=0}^{\infty} 2C a_n x^n &= \sum_{n=2}^{\infty} 2C a_{n-2} x^{n-2} \\ \sum_{n=0}^{\infty} 4x^{n-1} b_n &= \sum_{n=1}^{\infty} 4b_{n-1} x^{n-2} \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n - 2$.

$$\begin{aligned} & \left(\sum_{n=2}^{\infty} 2C(n-2) a_{n-2} x^{n-2} \right) + \left(\sum_{n=2}^{\infty} 2C a_{n-2} x^{n-2} \right) \\ & + \left(\sum_{n=0}^{\infty} x^{n-2} b_n (n^2 - 5n + 6) \right) + \left(\sum_{n=0}^{\infty} 3x^{n-2} b_n (n-2) \right) + \left(\sum_{n=1}^{\infty} 4b_{n-1} x^{n-2} \right) = 0 \end{aligned} \quad (2B)$$

For $n = 0$ in Eq. (2B), we choose arbitray value for b_0 as $b_0 = 1$. For $n = 1$, Eq (2B) gives

$$-b_1 + 4b_0 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$-b_1 + 4 = 0$$

Solving the above for b_1 gives

$$b_1 = 4$$

For $n = N$, where $N = 2$ which is the difference between the two roots, we are free to choose $b_2 = 0$. Hence for $n = 2$, Eq (2B) gives

$$2C + 16 = 0$$

Which is solved for C . Solving for C gives

$$C = -8$$

For $n = 3$, Eq (2B) gives

$$4Ca_1 + 4b_2 + 3b_3 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$3b_3 + \frac{128}{3} = 0$$

Solving the above for b_3 gives

$$b_3 = -\frac{128}{9}$$

For $n = 4$, Eq (2B) gives

$$6Ca_2 + 4b_3 + 8b_4 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$8b_4 - \frac{800}{9} = 0$$

Solving the above for b_4 gives

$$b_4 = \frac{100}{9}$$

For $n = 5$, Eq (2B) gives

$$8Ca_3 + 4b_4 + 15b_5 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$15b_5 + \frac{2512}{45} = 0$$

Solving the above for b_5 gives

$$b_5 = -\frac{2512}{675}$$

For $n = 6$, Eq (2B) gives

$$10Ca_4 + 4b_5 + 24b_6 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$24b_6 - \frac{11648}{675} = 0$$

Solving the above for b_6 gives

$$b_6 = \frac{1456}{2025}$$

For $n = 7$, Eq (2B) gives

$$12Ca_5 + 4b_6 + 35b_7 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$35b_7 + \frac{45376}{14175} = 0$$

Solving the above for b_7 gives

$$b_7 = -\frac{45376}{496125}$$

Now that we found all b_n and C , we can calculate the second solution from

$$y_2(x) = Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right)$$

Using the above value found for $C = -8$ and all b_n , then the second solution becomes

$$y_2(x) = (-8) \left(1 - \frac{4x}{3} + \frac{2x^2}{3} - \frac{8x^3}{45} + \frac{4x^4}{135} - \frac{16x^5}{4725} + \frac{4x^6}{14175} - \frac{16x^7}{893025} + O(x^8) \right) \ln(x) \\ + \frac{1 + 4x - \frac{128x^3}{9} + \frac{100x^4}{9} - \frac{2512x^5}{675} + \frac{1456x^6}{2025} - \frac{45376x^7}{496125} + O(x^8)}{x^2}$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 y_1(x) + c_2 y_2(x) \\ = c_1 \left(1 - \frac{4x}{3} + \frac{2x^2}{3} - \frac{8x^3}{45} + \frac{4x^4}{135} - \frac{16x^5}{4725} + \frac{4x^6}{14175} - \frac{16x^7}{893025} + O(x^8) \right) \\ + c_2 \left((-8) \left(1 - \frac{4x}{3} + \frac{2x^2}{3} - \frac{8x^3}{45} + \frac{4x^4}{135} - \frac{16x^5}{4725} + \frac{4x^6}{14175} - \frac{16x^7}{893025} + O(x^8) \right) \ln(x) \right. \\ \left. + \frac{1 + 4x - \frac{128x^3}{9} + \frac{100x^4}{9} - \frac{2512x^5}{675} + \frac{1456x^6}{2025} - \frac{45376x^7}{496125} + O(x^8)}{x^2} \right)$$

Hence the final solution is

$$\begin{aligned}
 y &= y_h \\
 &= c_1 \left(1 - \frac{4x}{3} + \frac{2x^2}{3} - \frac{8x^3}{45} + \frac{4x^4}{135} - \frac{16x^5}{4725} + \frac{4x^6}{14175} - \frac{16x^7}{893025} + O(x^8) \right) \\
 &\quad + c_2 \left(\left(-8 + \frac{32x}{3} - \frac{16x^2}{3} + \frac{64x^3}{45} - \frac{32x^4}{135} + \frac{128x^5}{4725} - \frac{32x^6}{14175} + \frac{128x^7}{893025} - 8O(x^8) \right) \ln(x) \right. \\
 &\quad \left. + \frac{1 + 4x - \frac{128x^3}{9} + \frac{100x^4}{9} - \frac{2512x^5}{675} + \frac{1456x^6}{2025} - \frac{45376x^7}{496125} + O(x^8)}{x^2} \right)
 \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}
 y &= c_1 \left(1 - \frac{4x}{3} + \frac{2x^2}{3} - \frac{8x^3}{45} + \frac{4x^4}{135} - \frac{16x^5}{4725} + \frac{4x^6}{14175} - \frac{16x^7}{893025} + O(x^8) \right) \\
 &\quad + c_2 \left(\left(-8 + \frac{32x}{3} - \frac{16x^2}{3} + \frac{64x^3}{45} - \frac{32x^4}{135} + \frac{128x^5}{4725} - \frac{32x^6}{14175} + \frac{128x^7}{893025} \right. \right. \\
 &\quad \left. \left. - 8O(x^8) \right) \ln(x) + \frac{1 + 4x - \frac{128x^3}{9} + \frac{100x^4}{9} - \frac{2512x^5}{675} + \frac{1456x^6}{2025} - \frac{45376x^7}{496125} + O(x^8)}{x^2} \right) \quad (1)
 \end{aligned}$$

Verification of solutions

$$\begin{aligned}
 y &= c_1 \left(1 - \frac{4x}{3} + \frac{2x^2}{3} - \frac{8x^3}{45} + \frac{4x^4}{135} - \frac{16x^5}{4725} + \frac{4x^6}{14175} - \frac{16x^7}{893025} + O(x^8) \right) \\
 &\quad + c_2 \left(\left(-8 + \frac{32x}{3} - \frac{16x^2}{3} + \frac{64x^3}{45} - \frac{32x^4}{135} + \frac{128x^5}{4725} - \frac{32x^6}{14175} + \frac{128x^7}{893025} - 8O(x^8) \right) \ln(x) \right. \\
 &\quad \left. + \frac{1 + 4x - \frac{128x^3}{9} + \frac{100x^4}{9} - \frac{2512x^5}{675} + \frac{1456x^6}{2025} - \frac{45376x^7}{496125} + O(x^8)}{x^2} \right)
 \end{aligned}$$

Verified OK.

19.12.1 Maple step by step solution

Let's solve

$$x^2y'' + 3xy' + 4xy = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{4y}{x} - \frac{3y'}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{3y'}{x} + \frac{4y}{x} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$[P_2(x) = \frac{3}{x}, P_3(x) = \frac{4}{x}]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 3$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 0$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$y''x + 4y + 3y' = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert y' to series expansion

$$y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1}$$

- Shift index using $k- > k+1$

$$y' = \sum_{k=-1}^{\infty} a_{k+1}(k+1+r)x^{k+r}$$

- Convert $x \cdot y''$ to series expansion

$$x \cdot y'' = \sum_{k=0}^{\infty} a_k(k+r)(k+r-1)x^{k+r-1}$$

- Shift index using $k- > k+1$

$$x \cdot y'' = \sum_{k=-1}^{\infty} a_{k+1}(k+1+r)(k+r)x^{k+r}$$

Rewrite ODE with series expansions

$$a_0r(2+r)x^{-1+r} + \left(\sum_{k=0}^{\infty} (a_{k+1}(k+1+r)(k+3+r) + 4a_k)x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$r(2+r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \{-2, 0\}$$

- Each term in the series must be 0, giving the recursion relation

$$a_{k+1}(k+1+r)(k+3+r) + 4a_k = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+1} = -\frac{4a_k}{(k+1+r)(k+3+r)}$$

- Recursion relation for $r = -2$

$$a_{k+1} = -\frac{4a_k}{(k-1)(k+1)}$$

- Series not valid for $r = -2$, division by 0 in the recursion relation at $k = 1$

$$a_{k+1} = -\frac{4a_k}{(k-1)(k+1)}$$

- Recursion relation for $r = 0$

$$a_{k+1} = -\frac{4a_k}{(k+1)(k+3)}$$

- Solution for $r = 0$

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+1} = -\frac{4a_k}{(k+1)(k+3)} \right]$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
    <- Bessel successful
<- special function solution successful`
```

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 74

```
Order:=8;
dsolve(x^2*diff(y(x),x$2)+3*x*diff(y(x),x)+4*x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 \left(1 - \frac{4}{3}x + \frac{2}{3}x^2 - \frac{8}{45}x^3 + \frac{4}{135}x^4 - \frac{16}{4725}x^5 + \frac{4}{14175}x^6 - \frac{16}{893025}x^7 + O(x^8) \right) x^2 + c_2 (\ln(x) (16x^2 - \frac{64}{3}x^3 + \dots))$$

✓ Solution by Mathematica

Time used: 0.043 (sec). Leaf size: 116

```
AsymptoticDSolveValue[x^2*y'[x]+3*x*y'[x]+4*x*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_2 \left(\frac{4x^6}{14175} - \frac{16x^5}{4725} + \frac{4x^4}{135} - \frac{8x^3}{45} + \frac{2x^2}{3} - \frac{4x}{3} + 1 \right) + c_1 \left(\frac{1696x^6 - 8976x^5 + 27900x^4 - 39600x^3 + 8100x^2 + 8100x + 2025}{2025x^2} - \frac{8}{135} (4x^4 - 24x^3 + 90x^2 - 180x + 135) \log(x) \right)$$

19.13 problem 3(d)

19.13.1 Maple step by step solution 3271

Internal problem ID [6453]

Internal file name [OUTPUT/5701_Sunday_June_05_2022_03_47_49_PM_13646004/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.4. REGULAR SINGULAR POINTS. Page 175

Problem number: 3(d).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

`[[_Emden , _Fowler]]`

$$x^3y'' - 4x^2y' + 3xy = 0$$

With the expansion point for the power series method at $x = 0$.

The ODE is

$$x^3y'' - 4x^2y' + 3xy = 0$$

Or

$$x(x^2y'' - 4xy' + 3y) = 0$$

For $x \neq 0$ the above simplifies to

$$x^2y'' - 4xy' + 3y = 0$$

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^3y'' - 4x^2y' + 3xy = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{4}{x}$$

$$q(x) = \frac{3}{x^2}$$

Table 453: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{4}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = \frac{3}{x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0, \infty]$

Irregular singular points : $[\]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^3y'' - 4x^2y' + 3xy = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) x^3 \\ & - 4x^2 \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + 3x \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \end{aligned} \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} x^{1+n+r} a_n (n+r)(n+r-1) \right) + \sum_{n=0}^{\infty} (-4x^{1+n+r} a_n (n+r)) + \left(\sum_{n=0}^{\infty} 3x^{1+n+r} a_n \right) = 0 \quad (2A)$$

The next step is to make all powers of x be $1+n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{1+n+r} and adjusting the power and the corresponding index gives Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $1+n+r$.

$$\left(\sum_{n=0}^{\infty} x^{1+n+r} a_n (n+r)(n+r-1) \right) + \sum_{n=0}^{\infty} (-4x^{1+n+r} a_n (n+r)) + \left(\sum_{n=0}^{\infty} 3x^{1+n+r} a_n \right) = 0 \quad (2B)$$

The indicial equation is obtained from $n = 0$. From Eq (2B) this gives

$$x^{1+n+r} a_n (n+r)(n+r-1) - 4x^{1+n+r} a_n (n+r) + 3x^{1+n+r} a_n = 0$$

When $n = 0$ the above becomes

$$x^{1+r} a_0 r(-1+r) - 4x^{1+r} a_0 r + 3x^{1+r} a_0 = 0$$

Or

$$(x^{1+r} r(-1+r) - 4x^{1+r} r + 3x^{1+r}) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(r^2 - 5r + 3) x^{1+r} = 0$$

Since the above is true for all x then the indicial equation becomes

$$r^2 - 5r + 3 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = \frac{5}{2} + \frac{\sqrt{13}}{2}$$

$$r_2 = \frac{5}{2} - \frac{\sqrt{13}}{2}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(r^2 - 5r + 3)x^{1+r} = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = \sqrt{13}$ is not an integer, then we can construct two linearly independent solutions

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n + \frac{5}{2} + \frac{\sqrt{13}}{2}}$$

$$y_2(x) = \sum_{n=0}^{\infty} b_n x^{n + \frac{5}{2} - \frac{\sqrt{13}}{2}}$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $0 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) - 4a_n(n+r) + 3a_n = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = 0 \quad (4)$$

Which for the root $r = \frac{5}{2} + \frac{\sqrt{13}}{2}$ becomes

$$a_n = 0 \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = \frac{5}{2} + \frac{\sqrt{13}}{2}$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	0	0
a_4	0	0

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	0	0
a_4	0	0
a_5	0	0

For $n = 6$, using the above recursive equation gives

$$a_6 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	0	0
a_4	0	0
a_5	0	0
a_6	0	0

For $n = 7$, using the above recursive equation gives

$$a_7 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	0	0
a_4	0	0
a_5	0	0
a_6	0	0
a_7	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x^{\frac{5}{2} + \frac{\sqrt{13}}{2}} (a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots) \\ &= x^{\frac{5}{2} + \frac{\sqrt{13}}{2}} (1 + O(x^8)) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. For $0 \leq n$ the recursive equation is

$$b_n(n+r)(n+r-1) - 4b_n(n+r) + 3b_n = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = 0 \tag{4}$$

Which for the root $r = \frac{5}{2} - \frac{\sqrt{13}}{2}$ becomes

$$b_n = 0 \tag{5}$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = \frac{5}{2} - \frac{\sqrt{13}}{2}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1

For $n = 1$, using the above recursive equation gives

$$b_1 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0

For $n = 2$, using the above recursive equation gives

$$b_2 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	0	0

For $n = 3$, using the above recursive equation gives

$$b_3 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	0	0
b_3	0	0

For $n = 4$, using the above recursive equation gives

$$b_4 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	0	0
b_3	0	0
b_4	0	0

For $n = 5$, using the above recursive equation gives

$$b_5 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	0	0
b_3	0	0
b_4	0	0
b_5	0	0

For $n = 6$, using the above recursive equation gives

$$b_6 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	0	0
b_3	0	0
b_4	0	0
b_5	0	0
b_6	0	0

For $n = 7$, using the above recursive equation gives

$$b_7 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	0	0
b_3	0	0
b_4	0	0
b_5	0	0
b_6	0	0
b_7	0	0

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned} y_2(x) &= x^{\frac{5}{2} + \frac{\sqrt{13}}{2}} (b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 + b_7x^7 + b_8x^8 \dots) \\ &= x^{\frac{5}{2} - \frac{\sqrt{13}}{2}} (1 + O(x^8)) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= c_1x^{\frac{5}{2} + \frac{\sqrt{13}}{2}} (1 + O(x^8)) + c_2x^{\frac{5}{2} - \frac{\sqrt{13}}{2}} (1 + O(x^8)) \end{aligned}$$

Hence the final solution is

$$\begin{aligned}y &= y_h \\ &= c_1 x^{\frac{5}{2} + \frac{\sqrt{13}}{2}} (1 + O(x^8)) + c_2 x^{\frac{5}{2} - \frac{\sqrt{13}}{2}} (1 + O(x^8))\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 x^{\frac{5}{2} + \frac{\sqrt{13}}{2}} (1 + O(x^8)) + c_2 x^{\frac{5}{2} - \frac{\sqrt{13}}{2}} (1 + O(x^8)) \quad (1)$$

Verification of solutions

$$y = c_1 x^{\frac{5}{2} + \frac{\sqrt{13}}{2}} (1 + O(x^8)) + c_2 x^{\frac{5}{2} - \frac{\sqrt{13}}{2}} (1 + O(x^8))$$

Verified OK.

19.13.1 Maple step by step solution

Let's solve

$$y''x^3 - 4x^2y' + 3xy = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = \frac{4y'}{x} - \frac{3y}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - \frac{4y'}{x} + \frac{3y}{x^2} = 0$$

- Multiply by denominators of the ODE

$$x^2y'' - 4xy' + 3y = 0$$

- Make a change of variables

$$t = \ln(x)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of y with respect to x , using the chain rule

$$y' = \left(\frac{d}{dt}y(t)\right) t'(x)$$

- Compute derivative

$$y' = \frac{\frac{d}{dt}y(t)}{x}$$

- Calculate the 2nd derivative of y with respect to x , using the chain rule

$$y'' = \left(\frac{d^2}{dt^2} y(t) \right) t'(x)^2 + t''(x) \left(\frac{d}{dt} y(t) \right)$$

- Compute derivative

$$y'' = \frac{\frac{d^2}{dt^2} y(t)}{x^2} - \frac{\frac{d}{dt} y(t)}{x^2}$$

Substitute the change of variables back into the ODE

$$x^2 \left(\frac{\frac{d^2}{dt^2} y(t)}{x^2} - \frac{\frac{d}{dt} y(t)}{x^2} \right) - 4 \frac{d}{dt} y(t) + 3y(t) = 0$$

- Simplify

$$\frac{d^2}{dt^2} y(t) - 5 \frac{d}{dt} y(t) + 3y(t) = 0$$

- Characteristic polynomial of ODE

$$r^2 - 5r + 3 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{5 \pm (\sqrt{13})}{2}$$

- Roots of the characteristic polynomial

$$r = \left(\frac{5}{2} - \frac{\sqrt{13}}{2}, \frac{5}{2} + \frac{\sqrt{13}}{2} \right)$$

- 1st solution of the ODE

$$y_1(t) = e^{\left(\frac{5}{2} - \frac{\sqrt{13}}{2}\right)t}$$

- 2nd solution of the ODE

$$y_2(t) = e^{\left(\frac{5}{2} + \frac{\sqrt{13}}{2}\right)t}$$

- General solution of the ODE

$$y(t) = c_1 y_1(t) + c_2 y_2(t)$$

- Substitute in solutions

$$y(t) = c_1 e^{\left(\frac{5}{2} - \frac{\sqrt{13}}{2}\right)t} + c_2 e^{\left(\frac{5}{2} + \frac{\sqrt{13}}{2}\right)t}$$

- Change variables back using $t = \ln(x)$

$$y = c_1 e^{\left(\frac{5}{2} - \frac{\sqrt{13}}{2}\right) \ln(x)} + c_2 e^{\left(\frac{5}{2} + \frac{\sqrt{13}}{2}\right) \ln(x)}$$

- Simplify

$$y = x^{\frac{5}{2}} \left(x^{\frac{\sqrt{13}}{2}} c_2 + x^{-\frac{\sqrt{13}}{2}} c_1 \right)$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
<- LODE of Euler type successful`
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 39

```
Order:=8;  
dsolve(x^3*diff(y(x),x$2)-4*x^2*diff(y(x),x)+3*x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = x^{\frac{5}{2}} \left(x^{-\frac{\sqrt{13}}{2}} c_1 + x^{\frac{\sqrt{13}}{2}} c_2 \right) + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 38

```
AsymptoticDSolveValue[x^3*y''[x]-4*x^2*y'[x]+3*x*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 x^{\frac{1}{2}(5+\sqrt{13})} + c_2 x^{\frac{1}{2}(5-\sqrt{13})}$$

19.14 problem 4(a)

19.14.1 Maple step by step solution 3286

Internal problem ID [6454]

Internal file name [OUTPUT/5702_Sunday_June_05_2022_03_47_51_PM_11707740/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.4. REGULAR SINGULAR POINTS. Page 175

Problem number: 4(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_Emden , _Fowler]]
```

$$4y''x + 3y' + y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$4y''x + 3y' + y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{3}{4x}$$
$$q(x) = \frac{1}{4x}$$

Table 455: Table $p(x), q(x)$ singularities.

$p(x) = \frac{3}{4x}$	
singularity	type
$x = 0$	“regular”

$q(x) = \frac{1}{4x}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$4y''x + 3y' + y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$4 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) x + 3 \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} 4x^{n+r-1} a_n (n+r)(n+r-1) \right) + \left(\sum_{n=0}^{\infty} 3(n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (2A)$$

The next step is to make all powers of x be $n + r - 1$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r-1} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} a_n x^{n+r} = \sum_{n=1}^{\infty} a_{n-1} x^{n+r-1}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n + r - 1$.

$$\left(\sum_{n=0}^{\infty} 4x^{n+r-1} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} 3(n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=1}^{\infty} a_{n-1} x^{n+r-1} \right) = 0 \quad (2B)$$

The indicial equation is obtained from $n = 0$. From Eq (2B) this gives

$$4x^{n+r-1} a_n (n+r) (n+r-1) + 3(n+r) a_n x^{n+r-1} = 0$$

When $n = 0$ the above becomes

$$4x^{-1+r} a_0 r (-1+r) + 3r a_0 x^{-1+r} = 0$$

Or

$$(4x^{-1+r} r (-1+r) + 3r x^{-1+r}) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$r x^{-1+r} (-1+4r) = 0$$

Since the above is true for all x then the indicial equation becomes

$$4r^2 - r = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = \frac{1}{4}$$

$$r_2 = 0$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$r x^{-1+r} (-1+4r) = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = \frac{1}{4}$ is not an integer, then we can construct two linearly independent solutions

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+\frac{1}{4}}$$

$$y_2(x) = \sum_{n=0}^{\infty} b_n x^n$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$4a_n(n+r)(n+r-1) + 3a_n(n+r) + a_{n-1} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{a_{n-1}}{4n^2 + 8nr + 4r^2 - n - r} \quad (4)$$

Which for the root $r = \frac{1}{4}$ becomes

$$a_n = -\frac{a_{n-1}}{4n^2 + n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = \frac{1}{4}$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = -\frac{1}{4r^2 + 7r + 3}$$

Which for the root $r = \frac{1}{4}$ becomes

$$a_1 = -\frac{1}{5}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{4r^2+7r+3}$	$-\frac{1}{5}$

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{16r^4 + 88r^3 + 173r^2 + 143r + 42}$$

Which for the root $r = \frac{1}{4}$ becomes

$$a_2 = \frac{1}{90}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{4r^2+7r+3}$	$-\frac{1}{5}$
a_2	$\frac{1}{16r^4+88r^3+173r^2+143r+42}$	$\frac{1}{90}$

For $n = 3$, using the above recursive equation gives

$$a_3 = -\frac{1}{64r^6 + 720r^5 + 3244r^4 + 7455r^3 + 9166r^2 + 5685r + 1386}$$

Which for the root $r = \frac{1}{4}$ becomes

$$a_3 = -\frac{1}{3510}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{4r^2+7r+3}$	$-\frac{1}{5}$
a_2	$\frac{1}{16r^4+88r^3+173r^2+143r+42}$	$\frac{1}{90}$
a_3	$-\frac{1}{64r^6+720r^5+3244r^4+7455r^3+9166r^2+5685r+1386}$	$-\frac{1}{3510}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{256r^8 + 4864r^7 + 39136r^6 + 173584r^5 + 462409r^4 + 754186r^3 + 731739r^2 + 384066r + 83160}$$

Which for the root $r = \frac{1}{4}$ becomes

$$a_4 = \frac{1}{238680}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{4r^2+7r+3}$	$-\frac{1}{5}$
a_2	$\frac{1}{16r^4+88r^3+173r^2+143r+42}$	$\frac{1}{90}$
a_3	$-\frac{1}{64r^6+720r^5+3244r^4+7455r^3+9166r^2+5685r+1386}$	$-\frac{1}{3510}$
a_4	$\frac{1}{256r^8+4864r^7+39136r^6+173584r^5+462409r^4+754186r^3+731739r^2+384066r+83160}$	$\frac{1}{238680}$

For $n = 5$, using the above recursive equation gives

$$a_5 = -\frac{1}{(256r^8 + 4864r^7 + 39136r^6 + 173584r^5 + 462409r^4 + 754186r^3 + 731739r^2 + 384066r + 83160)(4r^2 + 39r + 95)}$$

Which for the root $r = \frac{1}{4}$ becomes

$$a_5 = -\frac{1}{25061400}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{4r^2+7r+3}$	$-\frac{1}{5}$
a_2	$\frac{1}{16r^4+88r^3+173r^2+143r+42}$	$\frac{1}{90}$
a_3	$-\frac{1}{64r^6+720r^5+3244r^4+7455r^3+9166r^2+5685r+1386}$	$-\frac{1}{3510}$
a_4	$\frac{1}{256r^8+4864r^7+39136r^6+173584r^5+462409r^4+754186r^3+731739r^2+384066r+83160}$	$\frac{1}{238680}$
a_5	$-\frac{1}{(256r^8+4864r^7+39136r^6+173584r^5+462409r^4+754186r^3+731739r^2+384066r+83160)(4r^2+39r+95)}$	$-\frac{1}{25061400}$

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{1}{(256r^8 + 4864r^7 + 39136r^6 + 173584r^5 + 462409r^4 + 754186r^3 + 731739r^2 + 384066r + 83160)(4r^2 + 39r + 95)}$$

Which for the root $r = \frac{1}{4}$ becomes

$$a_6 = \frac{1}{3759210000}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{4r^2+7r+3}$	$-\frac{1}{5}$
a_2	$\frac{1}{16r^4+88r^3+173r^2+143r+42}$	$\frac{1}{90}$
a_3	$-\frac{1}{64r^6+720r^5+3244r^4+7455r^3+9166r^2+5685r+1386}$	$-\frac{1}{3510}$
a_4	$\frac{1}{256r^8+4864r^7+39136r^6+173584r^5+462409r^4+754186r^3+731739r^2+384066r+83160}$	$\frac{1}{238680}$
a_5	$-\frac{1}{(256r^8+4864r^7+39136r^6+173584r^5+462409r^4+754186r^3+731739r^2+384066r+83160)(4r^2+39r+95)}$	$-\frac{1}{25061400}$
a_6	$\frac{1}{(256r^8+4864r^7+39136r^6+173584r^5+462409r^4+754186r^3+731739r^2+384066r+83160)(4r^2+39r+95)(4r^2+47r+138)}$	$\frac{1}{3759210000}$

For $n = 7$, using the above recursive equation gives

$$a_7 = -\frac{1}{(256r^8 + 4864r^7 + 39136r^6 + 173584r^5 + 462409r^4 + 754186r^3 + 731739r^2 + 384066r + 83160)(4r^2 + 39r + 95)}$$

Which for the root $r = \frac{1}{4}$ becomes

$$a_7 = -\frac{1}{763119630000}$$

And the table now becomes

n	$a_{n,r}$
a_0	1
a_1	$-\frac{1}{4r^2+7r+3}$
a_2	$\frac{1}{16r^4+88r^3+173r^2+143r+42}$
a_3	$-\frac{1}{64r^6+720r^5+3244r^4+7455r^3+9166r^2+5685r+1386}$
a_4	$\frac{1}{256r^8+4864r^7+39136r^6+173584r^5+462409r^4+754186r^3+731739r^2+384066r+83160}$
a_5	$-\frac{1}{(256r^8+4864r^7+39136r^6+173584r^5+462409r^4+754186r^3+731739r^2+384066r+83160)(4r^2+39r+95)}$
a_6	$\frac{1}{(256r^8+4864r^7+39136r^6+173584r^5+462409r^4+754186r^3+731739r^2+384066r+83160)(4r^2+39r+95)(4r^2+47r+138)}$
a_7	$-\frac{1}{(256r^8+4864r^7+39136r^6+173584r^5+462409r^4+754186r^3+731739r^2+384066r+83160)(4r^2+39r+95)(4r^2+47r+138)(4r^2+55r+189)}$

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned}
y_1(x) &= x^{\frac{1}{4}}(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots) \\
&= x^{\frac{1}{4}}\left(1 - \frac{x}{5} + \frac{x^2}{90} - \frac{x^3}{3510} + \frac{x^4}{238680} - \frac{x^5}{25061400} + \frac{x^6}{3759210000} - \frac{x^7}{763119630000} + O(x^8)\right)
\end{aligned}$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. For $1 \leq n$ the recursive equation is

$$4b_n(n+r)(n+r-1) + 3(n+r)b_n + b_{n-1} = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = -\frac{b_{n-1}}{4n^2 + 8nr + 4r^2 - n - r} \quad (4)$$

Which for the root $r = 0$ becomes

$$b_n = -\frac{b_{n-1}}{n(4n-1)} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1

For $n = 1$, using the above recursive equation gives

$$b_1 = -\frac{1}{4r^2 + 7r + 3}$$

Which for the root $r = 0$ becomes

$$b_1 = -\frac{1}{3}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{1}{4r^2+7r+3}$	$-\frac{1}{3}$

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{1}{16r^4 + 88r^3 + 173r^2 + 143r + 42}$$

Which for the root $r = 0$ becomes

$$b_2 = \frac{1}{42}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{1}{4r^2+7r+3}$	$-\frac{1}{3}$
b_2	$\frac{1}{16r^4+88r^3+173r^2+143r+42}$	$\frac{1}{42}$

For $n = 3$, using the above recursive equation gives

$$b_3 = -\frac{1}{64r^6 + 720r^5 + 3244r^4 + 7455r^3 + 9166r^2 + 5685r + 1386}$$

Which for the root $r = 0$ becomes

$$b_3 = -\frac{1}{1386}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{1}{4r^2+7r+3}$	$-\frac{1}{3}$
b_2	$\frac{1}{16r^4+88r^3+173r^2+143r+42}$	$\frac{1}{42}$
b_3	$-\frac{1}{64r^6+720r^5+3244r^4+7455r^3+9166r^2+5685r+1386}$	$-\frac{1}{1386}$

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{1}{256r^8 + 4864r^7 + 39136r^6 + 173584r^5 + 462409r^4 + 754186r^3 + 731739r^2 + 384066r + 83160}$$

Which for the root $r = 0$ becomes

$$b_4 = \frac{1}{83160}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{1}{4r^2+7r+3}$	$-\frac{1}{3}$
b_2	$\frac{1}{16r^4+88r^3+173r^2+143r+42}$	$\frac{1}{42}$
b_3	$-\frac{1}{64r^6+720r^5+3244r^4+7455r^3+9166r^2+5685r+1386}$	$-\frac{1}{1386}$
b_4	$\frac{1}{256r^8+4864r^7+39136r^6+173584r^5+462409r^4+754186r^3+731739r^2+384066r+83160}$	$\frac{1}{83160}$

For $n = 5$, using the above recursive equation gives

$$b_5 = -\frac{1}{(256r^8 + 4864r^7 + 39136r^6 + 173584r^5 + 462409r^4 + 754186r^3 + 731739r^2 + 384066r + 83160)(4)}$$

Which for the root $r = 0$ becomes

$$b_5 = -\frac{1}{7900200}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{1}{4r^2+7r+3}$	$-\frac{1}{3}$
b_2	$\frac{1}{16r^4+88r^3+173r^2+143r+42}$	$\frac{1}{42}$
b_3	$-\frac{1}{64r^6+720r^5+3244r^4+7455r^3+9166r^2+5685r+1386}$	$-\frac{1}{1386}$
b_4	$\frac{1}{256r^8+4864r^7+39136r^6+173584r^5+462409r^4+754186r^3+731739r^2+384066r+83160}$	$\frac{1}{83160}$
b_5	$-\frac{1}{(256r^8+4864r^7+39136r^6+173584r^5+462409r^4+754186r^3+731739r^2+384066r+83160)(4r^2+39r+95)}$	$-\frac{1}{7900200}$

For $n = 6$, using the above recursive equation gives

$$b_6 = \frac{1}{(256r^8 + 4864r^7 + 39136r^6 + 173584r^5 + 462409r^4 + 754186r^3 + 731739r^2 + 384066r + 83160)(4r^2 + 39r + 95)}$$

Which for the root $r = 0$ becomes

$$b_6 = \frac{1}{1090227600}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$-\frac{1}{4r^2+7r+3}$	$-\frac{1}{3}$
b_2	$\frac{1}{16r^4+88r^3+173r^2+143r+42}$	$\frac{1}{42}$
b_3	$-\frac{1}{64r^6+720r^5+3244r^4+7455r^3+9166r^2+5685r+1386}$	$-\frac{1}{1386}$
b_4	$\frac{1}{256r^8+4864r^7+39136r^6+173584r^5+462409r^4+754186r^3+731739r^2+384066r+83160}$	$\frac{1}{83160}$
b_5	$-\frac{1}{(256r^8+4864r^7+39136r^6+173584r^5+462409r^4+754186r^3+731739r^2+384066r+83160)(4r^2+39r+95)}$	$-\frac{1}{7900200}$
b_6	$\frac{1}{(256r^8+4864r^7+39136r^6+173584r^5+462409r^4+754186r^3+731739r^2+384066r+83160)(4r^2+39r+95)(4r^2+47r+138)}$	$\frac{1}{1090227600}$

For $n = 7$, using the above recursive equation gives

$$b_7 = -\frac{1}{(256r^8 + 4864r^7 + 39136r^6 + 173584r^5 + 462409r^4 + 754186r^3 + 731739r^2 + 384066r + 83160)(4r^2 + 39r + 95)(4r^2 + 47r + 138)}$$

Which for the root $r = 0$ becomes

$$b_7 = -\frac{1}{206053016400}$$

And the table now becomes

n	$b_{n,r}$
b_0	1
b_1	$-\frac{1}{4r^2+7r+3}$
b_2	$\frac{1}{16r^4+88r^3+173r^2+143r+42}$
b_3	$-\frac{1}{64r^6+720r^5+3244r^4+7455r^3+9166r^2+5685r+1386}$
b_4	$\frac{1}{256r^8+4864r^7+39136r^6+173584r^5+462409r^4+754186r^3+731739r^2+384066r+83160}$
b_5	$-\frac{1}{(256r^8+4864r^7+39136r^6+173584r^5+462409r^4+754186r^3+731739r^2+384066r+83160)(4r^2+39r+95)}$
b_6	$\frac{1}{(256r^8+4864r^7+39136r^6+173584r^5+462409r^4+754186r^3+731739r^2+384066r+83160)(4r^2+39r+95)(4r^2+47r+138)}$
b_7	$-\frac{1}{(256r^8+4864r^7+39136r^6+173584r^5+462409r^4+754186r^3+731739r^2+384066r+83160)(4r^2+39r+95)(4r^2+47r+138)(4r^2+55r+189)}$

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned}
 y_2(x) &= b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 + b_7x^7 + b_8x^8 \dots \\
 &= 1 - \frac{x}{3} + \frac{x^2}{42} - \frac{x^3}{1386} + \frac{x^4}{83160} - \frac{x^5}{7900200} + \frac{x^6}{1090227600} - \frac{x^7}{206053016400} + O(x^8)
 \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 y_h(x) &= c_1y_1(x) + c_2y_2(x) \\
 &= c_1x^{\frac{1}{4}} \left(1 - \frac{x}{5} + \frac{x^2}{90} - \frac{x^3}{3510} + \frac{x^4}{238680} - \frac{x^5}{25061400} + \frac{x^6}{3759210000} - \frac{x^7}{763119630000} \right. \\
 &\quad \left. + O(x^8) \right) + c_2 \left(1 - \frac{x}{3} + \frac{x^2}{42} - \frac{x^3}{1386} + \frac{x^4}{83160} - \frac{x^5}{7900200} + \frac{x^6}{1090227600} \right. \\
 &\quad \left. - \frac{x^7}{206053016400} + O(x^8) \right)
 \end{aligned}$$

Hence the final solution is

$$\begin{aligned}
 y &= y_h \\
 &= c_1x^{\frac{1}{4}} \left(1 - \frac{x}{5} + \frac{x^2}{90} - \frac{x^3}{3510} + \frac{x^4}{238680} - \frac{x^5}{25061400} + \frac{x^6}{3759210000} - \frac{x^7}{763119630000} + O(x^8) \right) \\
 &\quad + c_2 \left(1 - \frac{x}{3} + \frac{x^2}{42} - \frac{x^3}{1386} + \frac{x^4}{83160} - \frac{x^5}{7900200} + \frac{x^6}{1090227600} - \frac{x^7}{206053016400} + O(x^8) \right)
 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 x^{\frac{1}{4}} \left(1 - \frac{x}{5} + \frac{x^2}{90} - \frac{x^3}{3510} + \frac{x^4}{238680} - \frac{x^5}{25061400} + \frac{x^6}{3759210000} - \frac{x^7}{763119630000} + O(x^8) \right) + c_2 \left(1 - \frac{x}{3} + \frac{x^2}{42} - \frac{x^3}{1386} + \frac{x^4}{83160} - \frac{x^5}{7900200} + \frac{x^6}{1090227600} - \frac{x^7}{206053016400} + O(x^8) \right)$$

Verification of solutions

$$y = c_1 x^{\frac{1}{4}} \left(1 - \frac{x}{5} + \frac{x^2}{90} - \frac{x^3}{3510} + \frac{x^4}{238680} - \frac{x^5}{25061400} + \frac{x^6}{3759210000} - \frac{x^7}{763119630000} + O(x^8) \right) + c_2 \left(1 - \frac{x}{3} + \frac{x^2}{42} - \frac{x^3}{1386} + \frac{x^4}{83160} - \frac{x^5}{7900200} + \frac{x^6}{1090227600} - \frac{x^7}{206053016400} + O(x^8) \right)$$

Verified OK.

19.14.1 Maple step by step solution

Let's solve

$$4y''x + 3y' + y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{y}{4x} - \frac{3y'}{4x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{3y'}{4x} + \frac{y}{4x} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$[P_2(x) = \frac{3}{4x}, P_3(x) = \frac{1}{4x}]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = \frac{3}{4}$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 0$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$4y''x + 3y' + y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert y' to series expansion

$$y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1}$$

- Shift index using $k \rightarrow k+1$

$$y' = \sum_{k=-1}^{\infty} a_{k+1} (k+1+r) x^{k+r}$$

- Convert $x \cdot y''$ to series expansion

$$x \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r-1}$$

- Shift index using $k \rightarrow k+1$

$$x \cdot y'' = \sum_{k=-1}^{\infty} a_{k+1} (k+1+r)(k+r) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0 r(-1+4r) x^{-1+r} + \left(\sum_{k=0}^{\infty} (a_{k+1} (k+1+r)(4k+3+4r) + a_k) x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$r(-1+4r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \left\{ 0, \frac{1}{4} \right\}$$

- Each term in the series must be 0, giving the recursion relation

$$4\left(k + \frac{3}{4} + r\right)(k + 1 + r)a_{k+1} + a_k = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+1} = -\frac{a_k}{(4k+3+4r)(k+1+r)}$$

- Recursion relation for $r = 0$

$$a_{k+1} = -\frac{a_k}{(4k+3)(k+1)}$$

- Solution for $r = 0$

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+1} = -\frac{a_k}{(4k+3)(k+1)} \right]$$

- Recursion relation for $r = \frac{1}{4}$

$$a_{k+1} = -\frac{a_k}{(4k+4)\left(k+\frac{5}{4}\right)}$$

- Solution for $r = \frac{1}{4}$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+\frac{1}{4}}, a_{k+1} = -\frac{a_k}{(4k+4)\left(k+\frac{5}{4}\right)} \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^k \right) + \left(\sum_{k=0}^{\infty} b_k x^{k+\frac{1}{4}} \right), a_{k+1} = -\frac{a_k}{(4k+3)(k+1)}, b_{k+1} = -\frac{b_k}{(4k+4)\left(k+\frac{5}{4}\right)} \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
    <- Bessel successful
<- special function solution successful`

```

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 52

```
Order:=8;  
dsolve(4*x*diff(y(x),x$2)+3*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^{\frac{1}{4}} \left(1 - \frac{1}{5}x + \frac{1}{90}x^2 - \frac{1}{3510}x^3 + \frac{1}{238680}x^4 - \frac{1}{25061400}x^5 + \frac{1}{3759210000}x^6 - \frac{1}{763119630000}x^7 + O(x^8) \right) + c_2 \left(1 - \frac{1}{3}x + \frac{1}{42}x^2 - \frac{1}{1386}x^3 + \frac{1}{83160}x^4 - \frac{1}{7900200}x^5 + \frac{1}{1090227600}x^6 - \frac{1}{206053016400}x^7 + O(x^8) \right)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 113

```
AsymptoticDSolveValue[4*x*y'[x]+3*y'[x]+y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \sqrt[4]{x} \left(-\frac{x^7}{763119630000} + \frac{x^6}{3759210000} - \frac{x^5}{25061400} + \frac{x^4}{238680} - \frac{x^3}{3510} + \frac{x^2}{90} - \frac{x}{5} + 1 \right) + c_2 \left(-\frac{x^7}{206053016400} + \frac{x^6}{1090227600} - \frac{x^5}{7900200} + \frac{x^4}{83160} - \frac{x^3}{1386} + \frac{x^2}{42} - \frac{x}{3} + 1 \right)$$

19.15 problem 4(b)

19.15.1 Maple step by step solution 3302

Internal problem ID [6455]

Internal file name [OUTPUT/5703_Sunday_June_05_2022_03_47_55_PM_19812894/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.4. REGULAR SINGULAR POINTS. Page 175

Problem number: 4(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _exact , _linear , _homogeneous]]
```

$$2y''x + (-x + 3)y' - y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$2y''x + (-x + 3)y' - y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{x-3}{2x}$$
$$q(x) = -\frac{1}{2x}$$

Table 457: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{x-3}{2x}$	
singularity	type
$x = 0$	“regular”

$q(x) = -\frac{1}{2x}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$2y''x + (-x + 3)y' - y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) x + (-x+3) \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) - \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2x^{n+r-1} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) \\ & + \left(\sum_{n=0}^{\infty} 3(n+r) a_n x^{n+r-1} \right) + \sum_{n=0}^{\infty} (-a_n x^{n+r}) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r-1$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r-1} and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) &= \sum_{n=1}^{\infty} (-a_{n-1} (n+r-1) x^{n+r-1}) \\ \sum_{n=0}^{\infty} (-a_n x^{n+r}) &= \sum_{n=1}^{\infty} (-a_{n-1} x^{n+r-1}) \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r-1$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2x^{n+r-1} a_n (n+r) (n+r-1) \right) + \sum_{n=1}^{\infty} (-a_{n-1} (n+r-1) x^{n+r-1}) \\ & + \left(\sum_{n=0}^{\infty} 3(n+r) a_n x^{n+r-1} \right) + \sum_{n=1}^{\infty} (-a_{n-1} x^{n+r-1}) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$2x^{n+r-1} a_n (n+r) (n+r-1) + 3(n+r) a_n x^{n+r-1} = 0$$

When $n=0$ the above becomes

$$2x^{-1+r} a_0 r(-1+r) + 3r a_0 x^{-1+r} = 0$$

Or

$$(2x^{-1+r} r(-1+r) + 3r x^{-1+r}) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(2r^2 + r) x^{-1+r} = 0$$

Since the above is true for all x then the indicial equation becomes

$$2r^2 + r = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 0 \\ r_2 &= -\frac{1}{2} \end{aligned}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(2r^2 + r) x^{-1+r} = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = \frac{1}{2}$ is not an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^n \\ y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n-\frac{1}{2}} \end{aligned}$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$2a_n(n+r)(n+r-1) - a_{n-1}(n+r-1) + 3a_n(n+r) - a_{n-1} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-1}}{2n+1+2r} \quad (4)$$

Which for the root $r = 0$ becomes

$$a_n = \frac{a_{n-1}}{2n+1} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = \frac{1}{3 + 2r}$$

Which for the root $r = 0$ becomes

$$a_1 = \frac{1}{3}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{3+2r}$	$\frac{1}{3}$

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{4r^2 + 16r + 15}$$

Which for the root $r = 0$ becomes

$$a_2 = \frac{1}{15}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{3+2r}$	$\frac{1}{3}$
a_2	$\frac{1}{4r^2+16r+15}$	$\frac{1}{15}$

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{1}{8r^3 + 60r^2 + 142r + 105}$$

Which for the root $r = 0$ becomes

$$a_3 = \frac{1}{105}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{3+2r}$	$\frac{1}{3}$
a_2	$\frac{1}{4r^2+16r+15}$	$\frac{1}{15}$
a_3	$\frac{1}{8r^3+60r^2+142r+105}$	$\frac{1}{105}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{16r^4 + 192r^3 + 824r^2 + 1488r + 945}$$

Which for the root $r = 0$ becomes

$$a_4 = \frac{1}{945}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{3+2r}$	$\frac{1}{3}$
a_2	$\frac{1}{4r^2+16r+15}$	$\frac{1}{15}$
a_3	$\frac{1}{8r^3+60r^2+142r+105}$	$\frac{1}{105}$
a_4	$\frac{1}{16r^4+192r^3+824r^2+1488r+945}$	$\frac{1}{945}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{1}{32r^5 + 560r^4 + 3760r^3 + 12040r^2 + 18258r + 10395}$$

Which for the root $r = 0$ becomes

$$a_5 = \frac{1}{10395}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{3+2r}$	$\frac{1}{3}$
a_2	$\frac{1}{4r^2+16r+15}$	$\frac{1}{15}$
a_3	$\frac{1}{8r^3+60r^2+142r+105}$	$\frac{1}{105}$
a_4	$\frac{1}{16r^4+192r^3+824r^2+1488r+945}$	$\frac{1}{945}$
a_5	$\frac{1}{32r^5+560r^4+3760r^3+12040r^2+18258r+10395}$	$\frac{1}{10395}$

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{1}{64r^6 + 1536r^5 + 14800r^4 + 72960r^3 + 193036r^2 + 258144r + 135135}$$

Which for the root $r = 0$ becomes

$$a_6 = \frac{1}{135135}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{3+2r}$	$\frac{1}{3}$
a_2	$\frac{1}{4r^2+16r+15}$	$\frac{1}{15}$
a_3	$\frac{1}{8r^3+60r^2+142r+105}$	$\frac{1}{105}$
a_4	$\frac{1}{16r^4+192r^3+824r^2+1488r+945}$	$\frac{1}{945}$
a_5	$\frac{1}{32r^5+560r^4+3760r^3+12040r^2+18258r+10395}$	$\frac{1}{10395}$
a_6	$\frac{1}{64r^6+1536r^5+14800r^4+72960r^3+193036r^2+258144r+135135}$	$\frac{1}{135135}$

For $n = 7$, using the above recursive equation gives

$$a_7 = \frac{1}{128r^7 + 4032r^6 + 52640r^5 + 367920r^4 + 1480472r^3 + 3411828r^2 + 4142430r + 2027025}$$

Which for the root $r = 0$ becomes

$$a_7 = \frac{1}{2027025}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{3+2r}$	$\frac{1}{3}$
a_2	$\frac{1}{4r^2+16r+15}$	$\frac{1}{15}$
a_3	$\frac{1}{8r^3+60r^2+142r+105}$	$\frac{1}{105}$
a_4	$\frac{1}{16r^4+192r^3+824r^2+1488r+945}$	$\frac{1}{945}$
a_5	$\frac{1}{32r^5+560r^4+3760r^3+12040r^2+18258r+10395}$	$\frac{1}{10395}$
a_6	$\frac{1}{64r^6+1536r^5+14800r^4+72960r^3+193036r^2+258144r+135135}$	$\frac{1}{135135}$
a_7	$\frac{1}{128r^7+4032r^6+52640r^5+367920r^4+1480472r^3+3411828r^2+4142430r+2027025}$	$\frac{1}{2027025}$

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned}
 y_1(x) &= a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots \\
 &= 1 + \frac{x}{3} + \frac{x^2}{15} + \frac{x^3}{105} + \frac{x^4}{945} + \frac{x^5}{10395} + \frac{x^6}{135135} + \frac{x^7}{2027025} + O(x^8)
 \end{aligned}$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. For $1 \leq n$ the recursive equation is

$$2b_n(n+r)(n+r-1) - b_{n-1}(n+r-1) + 3(n+r)b_n - b_{n-1} = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = \frac{b_{n-1}}{2n+1+2r} \quad (4)$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_n = \frac{b_{n-1}}{2n} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = -\frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1

For $n = 1$, using the above recursive equation gives

$$b_1 = \frac{1}{3 + 2r}$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_1 = \frac{1}{2}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{1}{3+2r}$	$\frac{1}{2}$

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{1}{4r^2 + 16r + 15}$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_2 = \frac{1}{8}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{1}{3+2r}$	$\frac{1}{2}$
b_2	$\frac{1}{4r^2+16r+15}$	$\frac{1}{8}$

For $n = 3$, using the above recursive equation gives

$$b_3 = \frac{1}{8r^3 + 60r^2 + 142r + 105}$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_3 = \frac{1}{48}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{1}{3+2r}$	$\frac{1}{2}$
b_2	$\frac{1}{4r^2+16r+15}$	$\frac{1}{8}$
b_3	$\frac{1}{8r^3+60r^2+142r+105}$	$\frac{1}{48}$

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{1}{16r^4 + 192r^3 + 824r^2 + 1488r + 945}$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_4 = \frac{1}{384}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{1}{3+2r}$	$\frac{1}{2}$
b_2	$\frac{1}{4r^2+16r+15}$	$\frac{1}{8}$
b_3	$\frac{1}{8r^3+60r^2+142r+105}$	$\frac{1}{48}$
b_4	$\frac{1}{16r^4+192r^3+824r^2+1488r+945}$	$\frac{1}{384}$

For $n = 5$, using the above recursive equation gives

$$b_5 = \frac{1}{32r^5 + 560r^4 + 3760r^3 + 12040r^2 + 18258r + 10395}$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_5 = \frac{1}{3840}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{1}{3+2r}$	$\frac{1}{2}$
b_2	$\frac{1}{4r^2+16r+15}$	$\frac{1}{8}$
b_3	$\frac{1}{8r^3+60r^2+142r+105}$	$\frac{1}{48}$
b_4	$\frac{1}{16r^4+192r^3+824r^2+1488r+945}$	$\frac{1}{384}$
b_5	$\frac{1}{32r^5+560r^4+3760r^3+12040r^2+18258r+10395}$	$\frac{1}{3840}$

For $n = 6$, using the above recursive equation gives

$$b_6 = \frac{1}{64r^6 + 1536r^5 + 14800r^4 + 72960r^3 + 193036r^2 + 258144r + 135135}$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_6 = \frac{1}{46080}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{1}{3+2r}$	$\frac{1}{2}$
b_2	$\frac{1}{4r^2+16r+15}$	$\frac{1}{8}$
b_3	$\frac{1}{8r^3+60r^2+142r+105}$	$\frac{1}{48}$
b_4	$\frac{1}{16r^4+192r^3+824r^2+1488r+945}$	$\frac{1}{384}$
b_5	$\frac{1}{32r^5+560r^4+3760r^3+12040r^2+18258r+10395}$	$\frac{1}{3840}$
b_6	$\frac{1}{64r^6+1536r^5+14800r^4+72960r^3+193036r^2+258144r+135135}$	$\frac{1}{46080}$

For $n = 7$, using the above recursive equation gives

$$b_7 = \frac{1}{128r^7 + 4032r^6 + 52640r^5 + 367920r^4 + 1480472r^3 + 3411828r^2 + 4142430r + 2027025}$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_7 = \frac{1}{645120}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{1}{3+2r}$	$\frac{1}{2}$
b_2	$\frac{1}{4r^2+16r+15}$	$\frac{1}{8}$
b_3	$\frac{1}{8r^3+60r^2+142r+105}$	$\frac{1}{48}$
b_4	$\frac{1}{16r^4+192r^3+824r^2+1488r+945}$	$\frac{1}{384}$
b_5	$\frac{1}{32r^5+560r^4+3760r^3+12040r^2+18258r+10395}$	$\frac{1}{3840}$
b_6	$\frac{1}{64r^6+1536r^5+14800r^4+72960r^3+193036r^2+258144r+135135}$	$\frac{1}{46080}$
b_7	$\frac{1}{128r^7+4032r^6+52640r^5+367920r^4+1480472r^3+3411828r^2+4142430r+2027025}$	$\frac{1}{645120}$

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned}
 y_2(x) &= 1(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 + b_7x^7 + b_8x^8 \dots) \\
 &= \frac{1 + \frac{x}{2} + \frac{x^2}{8} + \frac{x^3}{48} + \frac{x^4}{384} + \frac{x^5}{3840} + \frac{x^6}{46080} + \frac{x^7}{645120} + O(x^8)}{\sqrt{x}}
 \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 y_h(x) &= c_1y_1(x) + c_2y_2(x) \\
 &= c_1 \left(1 + \frac{x}{3} + \frac{x^2}{15} + \frac{x^3}{105} + \frac{x^4}{945} + \frac{x^5}{10395} + \frac{x^6}{135135} + \frac{x^7}{2027025} + O(x^8) \right) \\
 &\quad + \frac{c_2 \left(1 + \frac{x}{2} + \frac{x^2}{8} + \frac{x^3}{48} + \frac{x^4}{384} + \frac{x^5}{3840} + \frac{x^6}{46080} + \frac{x^7}{645120} + O(x^8) \right)}{\sqrt{x}}
 \end{aligned}$$

Hence the final solution is

$$\begin{aligned}
 y &= y_h \\
 &= c_1 \left(1 + \frac{x}{3} + \frac{x^2}{15} + \frac{x^3}{105} + \frac{x^4}{945} + \frac{x^5}{10395} + \frac{x^6}{135135} + \frac{x^7}{2027025} + O(x^8) \right) \\
 &\quad + \frac{c_2 \left(1 + \frac{x}{2} + \frac{x^2}{8} + \frac{x^3}{48} + \frac{x^4}{384} + \frac{x^5}{3840} + \frac{x^6}{46080} + \frac{x^7}{645120} + O(x^8) \right)}{\sqrt{x}}
 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \left(1 + \frac{x}{3} + \frac{x^2}{15} + \frac{x^3}{105} + \frac{x^4}{945} + \frac{x^5}{10395} + \frac{x^6}{135135} + \frac{x^7}{2027025} + O(x^8) \right) + \frac{c_2 \left(1 + \frac{x}{2} + \frac{x^2}{8} + \frac{x^3}{48} + \frac{x^4}{384} + \frac{x^5}{3840} + \frac{x^6}{46080} + \frac{x^7}{645120} + O(x^8) \right)}{\sqrt{x}} \quad (1)$$

Verification of solutions

$$y = c_1 \left(1 + \frac{x}{3} + \frac{x^2}{15} + \frac{x^3}{105} + \frac{x^4}{945} + \frac{x^5}{10395} + \frac{x^6}{135135} + \frac{x^7}{2027025} + O(x^8) \right) + \frac{c_2 \left(1 + \frac{x}{2} + \frac{x^2}{8} + \frac{x^3}{48} + \frac{x^4}{384} + \frac{x^5}{3840} + \frac{x^6}{46080} + \frac{x^7}{645120} + O(x^8) \right)}{\sqrt{x}}$$

Verified OK.

19.15.1 Maple step by step solution

Let's solve

$$2y''x + (-x + 3)y' - y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = \frac{y}{2x} + \frac{(x-3)y'}{2x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - \frac{(x-3)y'}{2x} - \frac{y}{2x} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$[P_2(x) = -\frac{x-3}{2x}, P_3(x) = -\frac{1}{2x}]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = \frac{3}{2}$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 0$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$2y''x + (-x + 3)y' - y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y'$ to series expansion for $m = 0..1$

$$x^m \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1+m}$$

- Shift index using $k \rightarrow k+1-m$

$$x^m \cdot y' = \sum_{k=-1+m}^{\infty} a_{k+1-m} (k+1-m+r) x^{k+r}$$

- Convert $x \cdot y''$ to series expansion

$$x \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r-1}$$

- Shift index using $k \rightarrow k+1$

$$x \cdot y'' = \sum_{k=-1}^{\infty} a_{k+1} (k+1+r)(k+r) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0 r(1+2r) x^{-1+r} + \left(\sum_{k=0}^{\infty} (a_{k+1} (k+1+r)(2k+3+2r) - a_k (k+1+r)) x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$r(1+2r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \left\{ 0, -\frac{1}{2} \right\}$$

- Each term in the series must be 0, giving the recursion relation

$$2(k+1+r) \left((k+r+\frac{3}{2}) a_{k+1} - \frac{a_k}{2} \right) = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+1} = \frac{a_k}{2k+3+2r}$$

- Recursion relation for $r = 0$

$$a_{k+1} = \frac{a_k}{2k+3}$$

- Solution for $r = 0$

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+1} = \frac{a_k}{2k+3} \right]$$

- Recursion relation for $r = -\frac{1}{2}$

$$a_{k+1} = \frac{a_k}{2k+2}$$

- Solution for $r = -\frac{1}{2}$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k-\frac{1}{2}}, a_{k+1} = \frac{a_k}{2k+2} \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^k \right) + \left(\sum_{k=0}^{\infty} b_k x^{k-\frac{1}{2}} \right), a_{k+1} = \frac{a_k}{2k+3}, b_{k+1} = \frac{b_k}{2k+2} \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
<- linear_1 successful`

```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 52

```

Order:=8;
dsolve(2*x*diff(y(x),x$2)+(3-x)*diff(y(x),x)-y(x)=0,y(x),type='series',x=0);

```

$$y(x) = \frac{c_1 \left(1 + \frac{1}{2}x + \frac{1}{8}x^2 + \frac{1}{48}x^3 + \frac{1}{384}x^4 + \frac{1}{3840}x^5 + \frac{1}{46080}x^6 + \frac{1}{645120}x^7 + O(x^8) \right)}{\sqrt{x}} + c_2 \left(1 + \frac{1}{3}x + \frac{1}{15}x^2 + \frac{1}{105}x^3 + \frac{1}{945}x^4 + \frac{1}{10395}x^5 + \frac{1}{135135}x^6 + \frac{1}{2027025}x^7 + O(x^8) \right)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 113

```
AsymptoticDSolveValue[2*x*y'[x]+(3-x)*y'[x]-y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(\frac{x^7}{2027025} + \frac{x^6}{135135} + \frac{x^5}{10395} + \frac{x^4}{945} + \frac{x^3}{105} + \frac{x^2}{15} + \frac{x}{3} + 1 \right) + \frac{c_2 \left(\frac{x^7}{645120} + \frac{x^6}{46080} + \frac{x^5}{3840} + \frac{x^4}{384} + \frac{x^3}{48} + \frac{x^2}{8} + \frac{x}{2} + 1 \right)}{\sqrt{x}}$$

19.16 problem 4(c)

19.16.1 Maple step by step solution 3318

Internal problem ID [6456]

Internal file name [OUTPUT/5704_Sunday_June_05_2022_03_47_58_PM_92248162/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.4. REGULAR SINGULAR POINTS. Page 175

Problem number: 4(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$2y''x + (1 + x)y' + 3y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$2y''x + (1 + x)y' + 3y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{1 + x}{2x}$$
$$q(x) = \frac{3}{2x}$$

Table 459: Table $p(x), q(x)$ singularities.

$p(x) = \frac{1+x}{2x}$	
singularity	type
$x = 0$	“regular”

$q(x) = \frac{3}{2x}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$2y''x + (1+x)y' + 3y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) x + (1+x) \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + 3 \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2x^{n+r-1} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) \right) \\ & + \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} 3a_n x^{n+r} \right) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r-1$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r-1} and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=0}^{\infty} x^{n+r} a_n (n+r) &= \sum_{n=1}^{\infty} a_{n-1} (n+r-1) x^{n+r-1} \\ \sum_{n=0}^{\infty} 3a_n x^{n+r} &= \sum_{n=1}^{\infty} 3a_{n-1} x^{n+r-1} \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r-1$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2x^{n+r-1} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=1}^{\infty} a_{n-1} (n+r-1) x^{n+r-1} \right) \\ & + \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=1}^{\infty} 3a_{n-1} x^{n+r-1} \right) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$2x^{n+r-1} a_n (n+r) (n+r-1) + (n+r) a_n x^{n+r-1} = 0$$

When $n=0$ the above becomes

$$2x^{-1+r} a_0 r (-1+r) + r a_0 x^{-1+r} = 0$$

Or

$$(2x^{-1+r} r (-1+r) + r x^{-1+r}) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$r x^{-1+r} (-1+2r) = 0$$

Since the above is true for all x then the indicial equation becomes

$$2r^2 - r = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = \frac{1}{2}$$

$$r_2 = 0$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$r x^{-1+r}(-1 + 2r) = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = \frac{1}{2}$ is not an integer, then we can construct two linearly independent solutions

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+\frac{1}{2}}$$

$$y_2(x) = \sum_{n=0}^{\infty} b_n x^n$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$2a_n(n+r)(n+r-1) + a_{n-1}(n+r-1) + a_n(n+r) + 3a_{n-1} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{a_{n-1}(n+r+2)}{2n^2 + 4nr + 2r^2 - n - r} \quad (4)$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_n = \frac{-2na_{n-1} - 5a_{n-1}}{4n^2 + 2n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = \frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = \frac{-3 - r}{2r^2 + 3r + 1}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_1 = -\frac{7}{6}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-3-r}{2r^2+3r+1}$	$-\frac{7}{6}$

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{(4+r)(3+r)}{4r^4 + 20r^3 + 35r^2 + 25r + 6}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_2 = \frac{21}{40}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-3-r}{2r^2+3r+1}$	$-\frac{7}{6}$
a_2	$\frac{(4+r)(3+r)}{4r^4+20r^3+35r^2+25r+6}$	$\frac{21}{40}$

For $n = 3$, using the above recursive equation gives

$$a_3 = -\frac{(4+r)(5+r)}{8r^5 + 60r^4 + 170r^3 + 225r^2 + 137r + 30}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_3 = -\frac{11}{80}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-3-r}{2r^2+3r+1}$	$-\frac{7}{6}$
a_2	$\frac{(4+r)(3+r)}{4r^4+20r^3+35r^2+25r+6}$	$\frac{21}{40}$
a_3	$-\frac{(4+r)(5+r)}{8r^5+60r^4+170r^3+225r^2+137r+30}$	$-\frac{11}{80}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{(5+r)(6+r)}{16r^6 + 176r^5 + 760r^4 + 1640r^3 + 1849r^2 + 1019r + 210}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_4 = \frac{143}{5760}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-3-r}{2r^2+3r+1}$	$-\frac{7}{6}$
a_2	$\frac{(4+r)(3+r)}{4r^4+20r^3+35r^2+25r+6}$	$\frac{21}{40}$
a_3	$-\frac{(4+r)(5+r)}{8r^5+60r^4+170r^3+225r^2+137r+30}$	$-\frac{11}{80}$
a_4	$\frac{(5+r)(6+r)}{16r^6+176r^5+760r^4+1640r^3+1849r^2+1019r+210}$	$\frac{143}{5760}$

For $n = 5$, using the above recursive equation gives

$$a_5 = -\frac{(6+r)(7+r)}{32r^7 + 496r^6 + 3104r^5 + 10120r^4 + 18458r^3 + 18679r^2 + 9591r + 1890}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_5 = -\frac{13}{3840}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-3-r}{2r^2+3r+1}$	$-\frac{7}{6}$
a_2	$\frac{(4+r)(3+r)}{4r^4+20r^3+35r^2+25r+6}$	$\frac{21}{40}$
a_3	$-\frac{(4+r)(5+r)}{8r^5+60r^4+170r^3+225r^2+137r+30}$	$-\frac{11}{80}$
a_4	$\frac{(5+r)(6+r)}{16r^6+176r^5+760r^4+1640r^3+1849r^2+1019r+210}$	$\frac{143}{5760}$
a_5	$-\frac{(6+r)(7+r)}{32r^7+496r^6+3104r^5+10120r^4+18458r^3+18679r^2+9591r+1890}$	$-\frac{13}{3840}$

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{(8+r)(7+r)}{64r^8 + 1344r^7 + 11664r^6 + 54384r^5 + 148236r^4 + 240396r^3 + 224651r^2 + 109281r + 20790}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_6 = \frac{17}{46080}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-3-r}{2r^2+3r+1}$	$-\frac{7}{6}$
a_2	$\frac{(4+r)(3+r)}{4r^4+20r^3+35r^2+25r+6}$	$\frac{21}{40}$
a_3	$-\frac{(4+r)(5+r)}{8r^5+60r^4+170r^3+225r^2+137r+30}$	$-\frac{11}{80}$
a_4	$\frac{(5+r)(6+r)}{16r^6+176r^5+760r^4+1640r^3+1849r^2+1019r+210}$	$\frac{143}{5760}$
a_5	$-\frac{(6+r)(7+r)}{32r^7+496r^6+3104r^5+10120r^4+18458r^3+18679r^2+9591r+1890}$	$-\frac{13}{3840}$
a_6	$\frac{(8+r)(7+r)}{64r^8+1344r^7+11664r^6+54384r^5+148236r^4+240396r^3+224651r^2+109281r+20790}$	$\frac{17}{46080}$

For $n = 7$, using the above recursive equation gives

$$a_7 = -\frac{(9+r)(8+r)}{128r^9 + 3520r^8 + 40800r^7 + 260400r^6 + 1003464r^5 + 2407860r^4 + 3574450r^3 + 3139025r^2 + 1462}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_7 = -\frac{323}{9676800}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-3-r}{2r^2+3r+1}$	$-\frac{7}{6}$
a_2	$\frac{(4+r)(3+r)}{4r^4+20r^3+35r^2+25r+6}$	$\frac{21}{40}$
a_3	$-\frac{(4+r)(5+r)}{8r^5+60r^4+170r^3+225r^2+137r+30}$	$-\frac{11}{80}$
a_4	$\frac{(5+r)(6+r)}{16r^6+176r^5+760r^4+1640r^3+1849r^2+1019r+210}$	$\frac{143}{5760}$
a_5	$-\frac{(6+r)(7+r)}{32r^7+496r^6+3104r^5+10120r^4+18458r^3+18679r^2+9591r+1890}$	$-\frac{13}{3840}$
a_6	$\frac{(8+r)(7+r)}{64r^8+1344r^7+11664r^6+54384r^5+148236r^4+240396r^3+224651r^2+109281r+20790}$	$\frac{17}{46080}$
a_7	$-\frac{(9+r)(8+r)}{128r^9+3520r^8+40800r^7+260400r^6+1003464r^5+2407860r^4+3574450r^3+3139025r^2+1462233r+270270}$	$-\frac{323}{9676800}$

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= \sqrt{x}(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots) \\ &= \sqrt{x} \left(1 - \frac{7x}{6} + \frac{21x^2}{40} - \frac{11x^3}{80} + \frac{143x^4}{5760} - \frac{13x^5}{3840} + \frac{17x^6}{46080} - \frac{323x^7}{9676800} + O(x^8) \right) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. For $1 \leq n$ the recursive equation is

$$2b_n(n+r)(n+r-1) + b_{n-1}(n+r-1) + (n+r)b_n + 3b_{n-1} = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = -\frac{b_{n-1}(n+r+2)}{2n^2 + 4nr + 2r^2 - n - r} \quad (4)$$

Which for the root $r = 0$ becomes

$$b_n = -\frac{b_{n-1}(n+2)}{n(2n-1)} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1

For $n = 1$, using the above recursive equation gives

$$b_1 = \frac{-3 - r}{2r^2 + 3r + 1}$$

Which for the root $r = 0$ becomes

$$b_1 = -3$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{-3-r}{2r^2+3r+1}$	-3

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{(4+r)(3+r)}{4r^4 + 20r^3 + 35r^2 + 25r + 6}$$

Which for the root $r = 0$ becomes

$$b_2 = 2$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{-3-r}{2r^2+3r+1}$	-3
b_2	$\frac{(4+r)(3+r)}{4r^4+20r^3+35r^2+25r+6}$	2

For $n = 3$, using the above recursive equation gives

$$b_3 = -\frac{(4+r)(5+r)}{8r^5 + 60r^4 + 170r^3 + 225r^2 + 137r + 30}$$

Which for the root $r = 0$ becomes

$$b_3 = -\frac{2}{3}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{-3-r}{2r^2+3r+1}$	-3
b_2	$\frac{(4+r)(3+r)}{4r^4+20r^3+35r^2+25r+6}$	2
b_3	$-\frac{(4+r)(5+r)}{8r^5+60r^4+170r^3+225r^2+137r+30}$	$-\frac{2}{3}$

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{(5+r)(6+r)}{16r^6 + 176r^5 + 760r^4 + 1640r^3 + 1849r^2 + 1019r + 210}$$

Which for the root $r = 0$ becomes

$$b_4 = \frac{1}{7}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{-3-r}{2r^2+3r+1}$	-3
b_2	$\frac{(4+r)(3+r)}{4r^4+20r^3+35r^2+25r+6}$	2
b_3	$-\frac{(4+r)(5+r)}{8r^5+60r^4+170r^3+225r^2+137r+30}$	$-\frac{2}{3}$
b_4	$\frac{(5+r)(6+r)}{16r^6+176r^5+760r^4+1640r^3+1849r^2+1019r+210}$	$\frac{1}{7}$

For $n = 5$, using the above recursive equation gives

$$b_5 = -\frac{(6+r)(7+r)}{32r^7 + 496r^6 + 3104r^5 + 10120r^4 + 18458r^3 + 18679r^2 + 9591r + 1890}$$

Which for the root $r = 0$ becomes

$$b_5 = -\frac{1}{45}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{-3-r}{2r^2+3r+1}$	-3
b_2	$\frac{(4+r)(3+r)}{4r^4+20r^3+35r^2+25r+6}$	2
b_3	$-\frac{(4+r)(5+r)}{8r^5+60r^4+170r^3+225r^2+137r+30}$	$-\frac{2}{3}$
b_4	$\frac{(5+r)(6+r)}{16r^6+176r^5+760r^4+1640r^3+1849r^2+1019r+210}$	$\frac{1}{7}$
b_5	$-\frac{(6+r)(7+r)}{32r^7+496r^6+3104r^5+10120r^4+18458r^3+18679r^2+9591r+1890}$	$-\frac{1}{45}$

For $n = 6$, using the above recursive equation gives

$$b_6 = \frac{(8+r)(7+r)}{64r^8 + 1344r^7 + 11664r^6 + 54384r^5 + 148236r^4 + 240396r^3 + 224651r^2 + 109281r + 20790}$$

Which for the root $r = 0$ becomes

$$b_6 = \frac{4}{1485}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{-3-r}{2r^2+3r+1}$	-3
b_2	$\frac{(4+r)(3+r)}{4r^4+20r^3+35r^2+25r+6}$	2
b_3	$-\frac{(4+r)(5+r)}{8r^5+60r^4+170r^3+225r^2+137r+30}$	$-\frac{2}{3}$
b_4	$\frac{(5+r)(6+r)}{16r^6+176r^5+760r^4+1640r^3+1849r^2+1019r+210}$	$\frac{1}{7}$
b_5	$-\frac{(6+r)(7+r)}{32r^7+496r^6+3104r^5+10120r^4+18458r^3+18679r^2+9591r+1890}$	$-\frac{1}{45}$
b_6	$\frac{(8+r)(7+r)}{64r^8+1344r^7+11664r^6+54384r^5+148236r^4+240396r^3+224651r^2+109281r+20790}$	$\frac{4}{1485}$

For $n = 7$, using the above recursive equation gives

$$b_7 = -\frac{(9+r)(8+r)}{128r^9 + 3520r^8 + 40800r^7 + 260400r^6 + 1003464r^5 + 2407860r^4 + 3574450r^3 + 3139025r^2 + 1462}$$

Which for the root $r = 0$ becomes

$$b_7 = -\frac{4}{15015}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{-3-r}{2r^2+3r+1}$	-3
b_2	$\frac{(4+r)(3+r)}{4r^4+20r^3+35r^2+25r+6}$	2
b_3	$-\frac{(4+r)(5+r)}{8r^5+60r^4+170r^3+225r^2+137r+30}$	$-\frac{2}{3}$
b_4	$\frac{(5+r)(6+r)}{16r^6+176r^5+760r^4+1640r^3+1849r^2+1019r+210}$	$\frac{1}{7}$
b_5	$-\frac{(6+r)(7+r)}{32r^7+496r^6+3104r^5+10120r^4+18458r^3+18679r^2+9591r+1890}$	$-\frac{1}{45}$
b_6	$\frac{(8+r)(7+r)}{64r^8+1344r^7+11664r^6+54384r^5+148236r^4+240396r^3+224651r^2+109281r+20790}$	$\frac{4}{1485}$
b_7	$-\frac{(9+r)(8+r)}{128r^9+3520r^8+40800r^7+260400r^6+1003464r^5+2407860r^4+3574450r^3+3139025r^2+1462233r+270270}$	$-\frac{4}{15015}$

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned} y_2(x) &= b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 + b_7x^7 + b_8x^8 \dots \\ &= 1 - 3x + 2x^2 - \frac{2x^3}{3} + \frac{x^4}{7} - \frac{x^5}{45} + \frac{4x^6}{1485} - \frac{4x^7}{15015} + O(x^8) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= c_1\sqrt{x} \left(1 - \frac{7x}{6} + \frac{21x^2}{40} - \frac{11x^3}{80} + \frac{143x^4}{5760} - \frac{13x^5}{3840} + \frac{17x^6}{46080} - \frac{323x^7}{9676800} + O(x^8) \right) \\ &\quad + c_2 \left(1 - 3x + 2x^2 - \frac{2x^3}{3} + \frac{x^4}{7} - \frac{x^5}{45} + \frac{4x^6}{1485} - \frac{4x^7}{15015} + O(x^8) \right) \end{aligned}$$

Hence the final solution is

$$\begin{aligned} y &= y_h \\ &= c_1\sqrt{x} \left(1 - \frac{7x}{6} + \frac{21x^2}{40} - \frac{11x^3}{80} + \frac{143x^4}{5760} - \frac{13x^5}{3840} + \frac{17x^6}{46080} - \frac{323x^7}{9676800} + O(x^8) \right) \\ &\quad + c_2 \left(1 - 3x + 2x^2 - \frac{2x^3}{3} + \frac{x^4}{7} - \frac{x^5}{45} + \frac{4x^6}{1485} - \frac{4x^7}{15015} + O(x^8) \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \sqrt{x} \left(1 - \frac{7x}{6} + \frac{21x^2}{40} - \frac{11x^3}{80} + \frac{143x^4}{5760} - \frac{13x^5}{3840} + \frac{17x^6}{46080} - \frac{323x^7}{9676800} + O(x^8) \right) \\ + c_2 \left(1 - 3x + 2x^2 - \frac{2x^3}{3} + \frac{x^4}{7} - \frac{x^5}{45} + \frac{4x^6}{1485} - \frac{4x^7}{15015} + O(x^8) \right) \quad (1)$$

Verification of solutions

$$y = c_1 \sqrt{x} \left(1 - \frac{7x}{6} + \frac{21x^2}{40} - \frac{11x^3}{80} + \frac{143x^4}{5760} - \frac{13x^5}{3840} + \frac{17x^6}{46080} - \frac{323x^7}{9676800} + O(x^8) \right) \\ + c_2 \left(1 - 3x + 2x^2 - \frac{2x^3}{3} + \frac{x^4}{7} - \frac{x^5}{45} + \frac{4x^6}{1485} - \frac{4x^7}{15015} + O(x^8) \right)$$

Verified OK.

19.16.1 Maple step by step solution

Let's solve

$$2y''x + (1+x)y' + 3y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{(1+x)y'}{2x} - \frac{3y}{2x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{(1+x)y'}{2x} + \frac{3y}{2x} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$\left[P_2(x) = \frac{1+x}{2x}, P_3(x) = \frac{3}{2x} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = \frac{1}{2}$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 0$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$2y''x + (1+x)y' + 3y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y'$ to series expansion for $m = 0..1$

$$x^m \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1+m}$$

- Shift index using $k \rightarrow k+1-m$

$$x^m \cdot y' = \sum_{k=-1+m}^{\infty} a_{k+1-m} (k+1-m+r) x^{k+r}$$

- Convert $x \cdot y''$ to series expansion

$$x \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r-1}$$

- Shift index using $k \rightarrow k+1$

$$x \cdot y'' = \sum_{k=-1}^{\infty} a_{k+1} (k+1+r)(k+r) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0 r(-1+2r) x^{-1+r} + \left(\sum_{k=0}^{\infty} (a_{k+1} (k+1+r)(2k+1+2r) + a_k (k+r+3)) x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$r(-1+2r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \left\{ 0, \frac{1}{2} \right\}$$

- Each term in the series must be 0, giving the recursion relation

$$2\left(k + \frac{1}{2} + r\right) (k+1+r) a_{k+1} + a_k (k+r+3) = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+1} = -\frac{a_k (k+r+3)}{(2k+1+2r)(k+1+r)}$$

- Recursion relation for $r = 0$

$$a_{k+1} = -\frac{a_k(k+3)}{(2k+1)(k+1)}$$

- Solution for $r = 0$

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+1} = -\frac{a_k(k+3)}{(2k+1)(k+1)} \right]$$

- Recursion relation for $r = \frac{1}{2}$

$$a_{k+1} = -\frac{a_k(k+\frac{7}{2})}{(2k+2)(k+\frac{3}{2})}$$

- Solution for $r = \frac{1}{2}$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+\frac{1}{2}}, a_{k+1} = -\frac{a_k(k+\frac{7}{2})}{(2k+2)(k+\frac{3}{2})} \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^k \right) + \left(\sum_{k=0}^{\infty} b_k x^{k+\frac{1}{2}} \right), a_{k+1} = -\frac{a_k(k+3)}{(2k+1)(k+1)}, b_{k+1} = -\frac{b_k(k+\frac{7}{2})}{(2k+2)(k+\frac{3}{2})} \right]$$

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
  A Liouvillian solution exists
  Reducible group (found an exponential solution)
  Group is reducible, not completely reducible
  Solution has integrals. Trying a special function solution free of integrals...
-> Trying a solution in terms of special functions:
  -> Bessel
  -> elliptic
  -> Legendre
  <- Kummer successful
<- special function solution successful
  Solution using Kummer functions still has integrals. Trying a hypergeometric solution.
  -> hyper3: Equivalence to 2F1, 1F1 or 0F1 under a power @ Moebius
  <- hyper3 successful: received ODE is equivalent to the 1F1 ODE
  -> Trying to convert hypergeometric functions to elementary form...
  <- elementary form for at least one hypergeometric solution is achieved - returning wi
<- Kovacics algorithm successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 52

```
Order:=8;
dsolve(2*x*diff(y(x),x$2)+(x+1)*diff(y(x),x)+3*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 \sqrt{x} \left(1 - \frac{7}{6}x + \frac{21}{40}x^2 - \frac{11}{80}x^3 + \frac{143}{5760}x^4 - \frac{13}{3840}x^5 + \frac{17}{46080}x^6 - \frac{323}{9676800}x^7 + O(x^8) \right) + c_2 \left(1 - 3x + 2x^2 - \frac{2}{3}x^3 + \frac{1}{7}x^4 - \frac{1}{45}x^5 + \frac{4}{1485}x^6 - \frac{4}{15015}x^7 + O(x^8) \right)$$

✓ Solution by Mathematica

Time used: 0.047 (sec). Leaf size: 106

```
AsymptoticDSolveValue[2*x*x*y'[x]+(x+1)*y'[x]+3*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(-\frac{1386072x^7}{35} + \frac{20088x^6}{5} - \frac{2511x^5}{5} + 81x^4 - 18x^3 + 6x^2 - 3x + 1 \right) \\ + c_2 e^{\frac{1}{2}/x} \left(\frac{257243688x^7}{35} + \frac{2381886x^6}{5} + \frac{176436x^5}{5} + 3042x^4 + 312x^3 + 39x^2 \right. \\ \left. + 6x + 1 \right) x^{3/2}$$

19.17 problem 4(d)

19.17.1 Maple step by step solution 3335

Internal problem ID [6457]

Internal file name [OUTPUT/5705_Sunday_June_05_2022_03_48_02_PM_72168280/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.4. REGULAR SINGULAR POINTS. Page 175

Problem number: 4(d).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$2x^2y'' + xy' - (1 + x)y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$2x^2y'' + xy' + (-1 - x)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{1}{2x}$$
$$q(x) = -\frac{1+x}{2x^2}$$

Table 461: Table $p(x), q(x)$ singularities.

$p(x) = \frac{1}{2x}$	
singularity	type
$x = 0$	“regular”

$q(x) = -\frac{1+x}{2x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$2x^2y'' + xy' + (-1 - x)y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$2x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) + x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (-1-x) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) \right) \\ & + \sum_{n=0}^{\infty} (-a_n x^{n+r}) + \sum_{n=0}^{\infty} (-x^{1+n+r} a_n) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} (-x^{1+n+r} a_n) = \sum_{n=1}^{\infty} (-a_{n-1} x^{n+r})$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) \right) \\ & + \sum_{n=0}^{\infty} (-a_n x^{n+r}) + \sum_{n=1}^{\infty} (-a_{n-1} x^{n+r}) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$2x^{n+r} a_n (n+r) (n+r-1) + x^{n+r} a_n (n+r) - a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$2x^r a_0 r(-1+r) + x^r a_0 r - a_0 x^r = 0$$

Or

$$(2x^r r(-1+r) + x^r r - x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(2r^2 - r - 1) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$2r^2 - r - 1 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 1 \\ r_2 &= -\frac{1}{2} \end{aligned}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(2r^2 - r - 1) x^r = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = \frac{3}{2}$ is not an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^{1+n} \\ y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n-\frac{1}{2}} \end{aligned}$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$2a_n(n+r)(n+r-1) + a_n(n+r) - a_n - a_{n-1} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-1}}{2n^2 + 4nr + 2r^2 - n - r - 1} \quad (4)$$

Which for the root $r = 1$ becomes

$$a_n = \frac{a_{n-1}}{n(2n+3)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 1$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = \frac{1}{r(2r+3)}$$

Which for the root $r = 1$ becomes

$$a_1 = \frac{1}{5}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{r(2r+3)}$	$\frac{1}{5}$

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{4r^4 + 20r^3 + 31r^2 + 15r}$$

Which for the root $r = 1$ becomes

$$a_2 = \frac{1}{70}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{r(2r+3)}$	$\frac{1}{5}$
a_2	$\frac{1}{4r^4+20r^3+31r^2+15r}$	$\frac{1}{70}$

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{1}{8r^6 + 84r^5 + 338r^4 + 651r^3 + 599r^2 + 210r}$$

Which for the root $r = 1$ becomes

$$a_3 = \frac{1}{1890}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{r(2r+3)}$	$\frac{1}{5}$
a_2	$\frac{1}{4r^4+20r^3+31r^2+15r}$	$\frac{1}{70}$
a_3	$\frac{1}{8r^6+84r^5+338r^4+651r^3+599r^2+210r}$	$\frac{1}{1890}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{16r^8 + 288r^7 + 2152r^6 + 8640r^5 + 20089r^4 + 26982r^3 + 19323r^2 + 5670r}$$

Which for the root $r = 1$ becomes

$$a_4 = \frac{1}{83160}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{r(2r+3)}$	$\frac{1}{5}$
a_2	$\frac{1}{4r^4+20r^3+31r^2+15r}$	$\frac{1}{70}$
a_3	$\frac{1}{8r^6+84r^5+338r^4+651r^3+599r^2+210r}$	$\frac{1}{1890}$
a_4	$\frac{1}{16r^8+288r^7+2152r^6+8640r^5+20089r^4+26982r^3+19323r^2+5670r}$	$\frac{1}{83160}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{1}{32r^{10} + 880r^9 + 10480r^8 + 70840r^7 + 299026r^6 + 815815r^5 + 1435220r^4 + 1565685r^3 + 957942r^2 + 320000r}$$

Which for the root $r = 1$ becomes

$$a_5 = \frac{1}{5405400}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{r(2r+3)}$	$\frac{1}{5}$
a_2	$\frac{1}{4r^4+20r^3+31r^2+15r}$	$\frac{1}{70}$
a_3	$\frac{1}{8r^6+84r^5+338r^4+651r^3+599r^2+210r}$	$\frac{1}{1890}$
a_4	$\frac{1}{16r^8+288r^7+2152r^6+8640r^5+20089r^4+26982r^3+19323r^2+5670r}$	$\frac{1}{83160}$
a_5	$\frac{1}{32r^{10}+880r^9+10480r^8+70840r^7+299026r^6+815815r^5+1435220r^4+1565685r^3+957942r^2+249480r}$	$\frac{1}{5405400}$

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{1}{r(32r^9 + 880r^8 + 10480r^7 + 70840r^6 + 299026r^5 + 815815r^4 + 1435220r^3 + 1565685r^2 + 957942r + 249480)}$$

Which for the root $r = 1$ becomes

$$a_6 = \frac{1}{486486000}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{r(2r+3)}$	$\frac{1}{5}$
a_2	$\frac{1}{4r^4+20r^3+31r^2+15r}$	$\frac{1}{70}$
a_3	$\frac{1}{8r^6+84r^5+338r^4+651r^3+599r^2+210r}$	$\frac{1}{1890}$
a_4	$\frac{1}{16r^8+288r^7+2152r^6+8640r^5+20089r^4+26982r^3+19323r^2+5670r}$	$\frac{1}{83160}$
a_5	$\frac{1}{32r^{10}+880r^9+10480r^8+70840r^7+299026r^6+815815r^5+1435220r^4+1565685r^3+957942r^2+249480r}$	$\frac{1}{5405400}$
a_6	$\frac{1}{r(32r^9+880r^8+10480r^7+70840r^6+299026r^5+815815r^4+1435220r^3+1565685r^2+957942r+249480)(2r^2+23r+65)}$	$\frac{1}{486486000}$

For $n = 7$, using the above recursive equation gives

$$a_7 = \frac{1}{r(32r^9 + 880r^8 + 10480r^7 + 70840r^6 + 299026r^5 + 815815r^4 + 1435220r^3 + 1565685r^2 + 957942r + 249480)(2r^2 + 23r + 65)}$$

Which for the root $r = 1$ becomes

$$a_7 = \frac{1}{57891834000}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{1}{r(2r+3)}$	$\frac{1}{5}$
a_2	$\frac{1}{4r^4+20r^3+31r^2+15r}$	$\frac{1}{70}$
a_3	$\frac{1}{8r^6+84r^5+338r^4+651r^3+599r^2+210r}$	$\frac{1}{189}$
a_4	$\frac{1}{16r^8+288r^7+2152r^6+8640r^5+20089r^4+26982r^3+19323r^2+5670r}$	$\frac{1}{83160}$
a_5	$\frac{1}{32r^{10}+880r^9+10480r^8+70840r^7+299026r^6+815815r^5+1435220r^4+1565685r^3+957942r^2+249480r}$	$\frac{1}{5405400}$
a_6	$\frac{1}{r(32r^9+880r^8+10480r^7+70840r^6+299026r^5+815815r^4+1435220r^3+1565685r^2+957942r+249480)(2r^2+23r+65)}$	$\frac{1}{486486000}$
a_7	$\frac{1}{r(32r^9+880r^8+10480r^7+70840r^6+299026r^5+815815r^4+1435220r^3+1565685r^2+957942r+249480)(2r^2+23r+65)(2r^2+27r+90)}$	$\frac{1}{57891834000}$

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned}
 y_1(x) &= x(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots) \\
 &= x\left(1 + \frac{x}{5} + \frac{x^2}{70} + \frac{x^3}{1890} + \frac{x^4}{83160} + \frac{x^5}{5405400} + \frac{x^6}{486486000} + \frac{x^7}{57891834000} + O(x^8)\right)
 \end{aligned}$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. For $1 \leq n$ the recursive equation is

$$2b_n(n+r)(n+r-1) + b_n(n+r) - b_n - b_{n-1} = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = \frac{b_{n-1}}{2n^2 + 4nr + 2r^2 - n - r - 1} \quad (4)$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_n = \frac{b_{n-1}}{n(2n-3)} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = -\frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1

For $n = 1$, using the above recursive equation gives

$$b_1 = \frac{1}{r(2r+3)}$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_1 = -1$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{1}{r(2r+3)}$	-1

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{1}{4r^4 + 20r^3 + 31r^2 + 15r}$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_2 = -\frac{1}{2}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{1}{r(2r+3)}$	-1
b_2	$\frac{1}{4r^4+20r^3+31r^2+15r}$	$-\frac{1}{2}$

For $n = 3$, using the above recursive equation gives

$$b_3 = \frac{1}{8r^6 + 84r^5 + 338r^4 + 651r^3 + 599r^2 + 210r}$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_3 = -\frac{1}{18}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{1}{r(2r+3)}$	-1
b_2	$\frac{1}{4r^4+20r^3+31r^2+15r}$	$-\frac{1}{2}$
b_3	$\frac{1}{8r^6+84r^5+338r^4+651r^3+599r^2+210r}$	$-\frac{1}{18}$

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{1}{16r^8 + 288r^7 + 2152r^6 + 8640r^5 + 20089r^4 + 26982r^3 + 19323r^2 + 5670r}$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_4 = -\frac{1}{360}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{1}{r(2r+3)}$	-1
b_2	$\frac{1}{4r^4+20r^3+31r^2+15r}$	$-\frac{1}{2}$
b_3	$\frac{1}{8r^6+84r^5+338r^4+651r^3+599r^2+210r}$	$-\frac{1}{18}$
b_4	$\frac{1}{16r^8+288r^7+2152r^6+8640r^5+20089r^4+26982r^3+19323r^2+5670r}$	$-\frac{1}{360}$

For $n = 5$, using the above recursive equation gives

$$b_5 = \frac{1}{32r^{10} + 880r^9 + 10480r^8 + 70840r^7 + 299026r^6 + 815815r^5 + 1435220r^4 + 1565685r^3 + 957942r^2 + 320000r}$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_5 = -\frac{1}{12600}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{1}{r(2r+3)}$	-1
b_2	$\frac{1}{4r^4+20r^3+31r^2+15r}$	$-\frac{1}{2}$
b_3	$\frac{1}{8r^6+84r^5+338r^4+651r^3+599r^2+210r}$	$-\frac{1}{18}$
b_4	$\frac{1}{16r^8+288r^7+2152r^6+8640r^5+20089r^4+26982r^3+19323r^2+5670r}$	$-\frac{1}{360}$
b_5	$\frac{1}{32r^{10}+880r^9+10480r^8+70840r^7+299026r^6+815815r^5+1435220r^4+1565685r^3+957942r^2+249480r}$	$-\frac{1}{12600}$

For $n = 6$, using the above recursive equation gives

$$b_6 = \frac{1}{r(32r^9 + 880r^8 + 10480r^7 + 70840r^6 + 299026r^5 + 815815r^4 + 1435220r^3 + 1565685r^2 + 957942r + 249480)}$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_6 = -\frac{1}{680400}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{1}{r(2r+3)}$	-1
b_2	$\frac{1}{4r^4+20r^3+31r^2+15r}$	$-\frac{1}{2}$
b_3	$\frac{1}{8r^6+84r^5+338r^4+651r^3+599r^2+210r}$	$-\frac{1}{18}$
b_4	$\frac{1}{16r^8+288r^7+2152r^6+8640r^5+20089r^4+26982r^3+19323r^2+5670r}$	$-\frac{1}{360}$
b_5	$\frac{1}{32r^{10}+880r^9+10480r^8+70840r^7+299026r^6+815815r^5+1435220r^4+1565685r^3+957942r^2+249480r}$	$-\frac{1}{12600}$
b_6	$\frac{1}{r(32r^9+880r^8+10480r^7+70840r^6+299026r^5+815815r^4+1435220r^3+1565685r^2+957942r+249480)(2r^2+23r+65)}$	$-\frac{1}{680400}$

For $n = 7$, using the above recursive equation gives

$$b_7 = \frac{1}{r(32r^9 + 880r^8 + 10480r^7 + 70840r^6 + 299026r^5 + 815815r^4 + 1435220r^3 + 1565685r^2 + 957942r + 249480)(2r^2 + 23r + 65)}$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_7 = -\frac{1}{52390800}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{1}{r(2r+3)}$	-1
b_2	$\frac{1}{4r^4+20r^3+31r^2+15r}$	-1
b_3	$\frac{1}{8r^6+84r^5+338r^4+651r^3+599r^2+210r}$	-1
b_4	$\frac{1}{16r^8+288r^7+2152r^6+8640r^5+20089r^4+26982r^3+19323r^2+5670r}$	-1
b_5	$\frac{1}{32r^{10}+880r^9+10480r^8+70840r^7+299026r^6+815815r^5+1435220r^4+1565685r^3+957942r^2+249480r}$	-1
b_6	$\frac{1}{r(32r^9+880r^8+10480r^7+70840r^6+299026r^5+815815r^4+1435220r^3+1565685r^2+957942r+249480)(2r^2+23r+65)}$	-6
b_7	$\frac{1}{r(32r^9+880r^8+10480r^7+70840r^6+299026r^5+815815r^4+1435220r^3+1565685r^2+957942r+249480)(2r^2+23r+65)(2r^2+27r+90)}$	-3

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned}
 y_2(x) &= x(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 + b_7x^7 + b_8x^8 \dots) \\
 &= \frac{1 - x - \frac{x^2}{2} - \frac{x^3}{18} - \frac{x^4}{360} - \frac{x^5}{12600} - \frac{x^6}{680400} - \frac{x^7}{52390800} + O(x^8)}{\sqrt{x}}
 \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 y_h(x) &= c_1y_1(x) + c_2y_2(x) \\
 &= c_1x \left(1 + \frac{x}{5} + \frac{x^2}{70} + \frac{x^3}{1890} + \frac{x^4}{83160} + \frac{x^5}{5405400} + \frac{x^6}{486486000} + \frac{x^7}{57891834000} + O(x^8) \right) \\
 &\quad + \frac{c_2 \left(1 - x - \frac{x^2}{2} - \frac{x^3}{18} - \frac{x^4}{360} - \frac{x^5}{12600} - \frac{x^6}{680400} - \frac{x^7}{52390800} + O(x^8) \right)}{\sqrt{x}}
 \end{aligned}$$

Hence the final solution is

$$\begin{aligned}
 y &= y_h \\
 &= c_1x \left(1 + \frac{x}{5} + \frac{x^2}{70} + \frac{x^3}{1890} + \frac{x^4}{83160} + \frac{x^5}{5405400} + \frac{x^6}{486486000} + \frac{x^7}{57891834000} + O(x^8) \right) \\
 &\quad + \frac{c_2 \left(1 - x - \frac{x^2}{2} - \frac{x^3}{18} - \frac{x^4}{360} - \frac{x^5}{12600} - \frac{x^6}{680400} - \frac{x^7}{52390800} + O(x^8) \right)}{\sqrt{x}}
 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 x \left(1 + \frac{x}{5} + \frac{x^2}{70} + \frac{x^3}{1890} + \frac{x^4}{83160} + \frac{x^5}{5405400} + \frac{x^6}{486486000} + \frac{x^7}{57891834000} + O(x^8) \right) + \frac{c_2 \left(1 - x - \frac{x^2}{2} - \frac{x^3}{18} - \frac{x^4}{360} - \frac{x^5}{12600} - \frac{x^6}{680400} - \frac{x^7}{52390800} + O(x^8) \right)}{\sqrt{x}}$$

Verification of solutions

$$y = c_1 x \left(1 + \frac{x}{5} + \frac{x^2}{70} + \frac{x^3}{1890} + \frac{x^4}{83160} + \frac{x^5}{5405400} + \frac{x^6}{486486000} + \frac{x^7}{57891834000} + O(x^8) \right) + \frac{c_2 \left(1 - x - \frac{x^2}{2} - \frac{x^3}{18} - \frac{x^4}{360} - \frac{x^5}{12600} - \frac{x^6}{680400} - \frac{x^7}{52390800} + O(x^8) \right)}{\sqrt{x}}$$

Verified OK.

19.17.1 Maple step by step solution

Let's solve

$$2x^2 y'' + xy' + (-1 - x)y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{y'}{2x} + \frac{(1+x)y}{2x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{y'}{2x} - \frac{(1+x)y}{2x^2} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$\left[P_2(x) = \frac{1}{2x}, P_3(x) = -\frac{1+x}{2x^2} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = \frac{1}{2}$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = -\frac{1}{2}$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$2x^2y'' + xy' + (-1 - x)y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 0..1$

$$x^m \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

- Shift index using $k \rightarrow k - m$

$$x^m \cdot y = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

- Convert $x \cdot y'$ to series expansion

$$x \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r}$$

- Convert $x^2 \cdot y''$ to series expansion

$$x^2 \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(1+2r)(-1+r)x^r + \left(\sum_{k=1}^{\infty} (a_k(2k+2r+1)(k+r-1) - a_{k-1}) x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$(1+2r)(-1+r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \left\{ 1, -\frac{1}{2} \right\}$$

- Each term in the series must be 0, giving the recursion relation

$$2(k+r-1)\left(k+\frac{1}{2}+r\right)a_k - a_{k-1} = 0$$

- Shift index using $k \rightarrow k+1$

$$2(k+r)\left(k+\frac{3}{2}+r\right)a_{k+1} - a_k = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+1} = \frac{a_k}{(k+r)(2k+3+2r)}$$

- Recursion relation for $r = 1$

$$a_{k+1} = \frac{a_k}{(k+1)(2k+5)}$$

- Solution for $r = 1$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+1}, a_{k+1} = \frac{a_k}{(k+1)(2k+5)} \right]$$

- Recursion relation for $r = -\frac{1}{2}$

$$a_{k+1} = \frac{a_k}{(k-\frac{1}{2})(2k+2)}$$

- Solution for $r = -\frac{1}{2}$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k-\frac{1}{2}}, a_{k+1} = \frac{a_k}{(k-\frac{1}{2})(2k+2)} \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^{k+1} \right) + \left(\sum_{k=0}^{\infty} b_k x^{k-\frac{1}{2}} \right), a_{k+1} = \frac{a_k}{(k+1)(2k+5)}, b_{k+1} = \frac{b_k}{(k-\frac{1}{2})(2k+2)} \right]$$

Maple trace Kovacic algorithm successful

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Group is reducible or imprimitive
<- Kovacics algorithm successful`

```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 53

Order:=8;

```
dsolve(2*x^2*diff(y(x),x$2)+x*diff(y(x),x)-(x+1)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{c_1 \left(1 - x - \frac{1}{2}x^2 - \frac{1}{18}x^3 - \frac{1}{360}x^4 - \frac{1}{12600}x^5 - \frac{1}{680400}x^6 - \frac{1}{52390800}x^7 + O(x^8)\right)}{\sqrt{x}} + c_2 x \left(1 + \frac{1}{5}x + \frac{1}{70}x^2 + \frac{1}{1890}x^3 + \frac{1}{83160}x^4 + \frac{1}{5405400}x^5 + \frac{1}{486486000}x^6 + \frac{1}{57891834000}x^7 + O(x^8)\right)$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 112

```
AsymptoticDSolveValue[2*x^2*y''[x]+x*y'[x]-(x+1)*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 x \left(\frac{x^7}{57891834000} + \frac{x^6}{486486000} + \frac{x^5}{5405400} + \frac{x^4}{83160} + \frac{x^3}{1890} + \frac{x^2}{70} + \frac{x}{5} + 1 \right) + \frac{c_2 \left(-\frac{x^7}{52390800} - \frac{x^6}{680400} - \frac{x^5}{12600} - \frac{x^4}{360} - \frac{x^3}{18} - \frac{x^2}{2} - x + 1 \right)}{\sqrt{x}}$$

19.18 problem 5

19.18.1 Maple step by step solution 3347

Internal problem ID [6458]

Internal file name [OUTPUT/5706_Sunday_June_05_2022_03_48_06_PM_15641277/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.4. REGULAR SINGULAR POINTS. Page 175

Problem number: 5.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Repeated root"**

Maple gives the following as the ode type

[_Lienard]

$$x^2y'' + xy' + yx^2 = 0$$

With the expansion point for the power series method at $x = 0$.

The ODE is

$$x^2y'' + xy' + yx^2 = 0$$

Or

$$x(y''x + y' + xy) = 0$$

For $x \neq 0$ the above simplifies to

$$y''x + y' + xy = 0$$

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2y'' + xy' + yx^2 = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{1}{x}$$

$$q(x) = 1$$

Table 463: Table $p(x), q(x)$ singularities.

$p(x) = \frac{1}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = 1$	
singularity	type

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2 y'' + x y' + y x^2 = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) + x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) x^2 = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) \right) + \left(\sum_{n=0}^{\infty} x^{2+n+r} a_n \right) = 0 \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} x^{2+n+r} a_n = \sum_{n=2}^{\infty} a_{n-2} x^{n+r}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) \right) + \left(\sum_{n=2}^{\infty} a_{n-2} x^{n+r} \right) = 0 \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r} a_n (n+r) (n+r-1) + x^{n+r} a_n (n+r) = 0$$

When $n=0$ the above becomes

$$x^r a_0 r(-1+r) + x^r a_0 r = 0$$

Or

$$(x^r r(-1+r) + x^r r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$x^r r^2 = 0$$

Since the above is true for all x then the indicial equation becomes

$$r^2 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = 0$$

$$r_2 = 0$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$x^r r^2 = 0$$

Solving for r gives the roots of the indicial equation as Since the root of the indicial equation is repeated, then we can construct two linearly independent solutions. The first solution has the form

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+r} \quad (1A)$$

Now the second solution y_2 is found using

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right) \quad (1B)$$

Then the general solution will be

$$y = c_1 y_1(x) + c_2 y_2(x)$$

In Eq (1B) the sum starts from 1 and not zero. In Eq (1A), a_0 is never zero, and is arbitrary and is typically taken as $a_0 = 1$, and $\{c_1, c_2\}$ are two arbitrary constants of integration which can be found from initial conditions. We start by finding the first solution $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

For $2 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) + a_n(n+r) + a_{n-2} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{a_{n-2}}{n^2 + 2nr + r^2} \quad (4)$$

Which for the root $r = 0$ becomes

$$a_n = -\frac{a_{n-2}}{n^2} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = -\frac{1}{(r+2)^2}$$

Which for the root $r = 0$ becomes

$$a_2 = -\frac{1}{4}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{(r+2)^2}$	$-\frac{1}{4}$

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{(r+2)^2}$	$-\frac{1}{4}$
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{(r+2)^2(r+4)^2}$$

Which for the root $r = 0$ becomes

$$a_4 = \frac{1}{64}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{(r+2)^2}$	$-\frac{1}{4}$
a_3	0	0
a_4	$\frac{1}{(r+2)^2(r+4)^2}$	$\frac{1}{64}$

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{(r+2)^2}$	$-\frac{1}{4}$
a_3	0	0
a_4	$\frac{1}{(r+2)^2(r+4)^2}$	$\frac{1}{64}$
a_5	0	0

For $n = 6$, using the above recursive equation gives

$$a_6 = -\frac{1}{(r+2)^2(r+4)^2(r+6)^2}$$

Which for the root $r = 0$ becomes

$$a_6 = -\frac{1}{2304}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{(r+2)^2}$	$-\frac{1}{4}$
a_3	0	0
a_4	$\frac{1}{(r+2)^2(r+4)^2}$	$\frac{1}{64}$
a_5	0	0
a_6	$-\frac{1}{(r+2)^2(r+4)^2(r+6)^2}$	$-\frac{1}{2304}$

For $n = 7$, using the above recursive equation gives

$$a_7 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{(r+2)^2}$	$-\frac{1}{4}$
a_3	0	0
a_4	$\frac{1}{(r+2)^2(r+4)^2}$	$\frac{1}{64}$
a_5	0	0
a_6	$-\frac{1}{(r+2)^2(r+4)^2(r+6)^2}$	$-\frac{1}{2304}$
a_7	0	0

Using the above table, then the first solution $y_1(x)$ becomes

$$\begin{aligned} y_1(x) &= a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots \\ &= 1 - \frac{x^2}{4} + \frac{x^4}{64} - \frac{x^6}{2304} + O(x^8) \end{aligned}$$

Now the second solution is found. The second solution is given by

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right)$$

Where b_n is found using

$$b_n = \frac{d}{dr} a_{n,r}$$

And the above is then evaluated at $r = 0$. The above table for $a_{n,r}$ is used for this purpose. Computing the derivatives gives the following table

n	$b_{n,r}$	a_n	$b_{n,r} = \frac{d}{dr} a_{n,r}$	$b_n(r = 0)$
b_0	1	1	N/A since b_n starts from 1	N/A
b_1	0	0	0	0
b_2	$-\frac{1}{(r+2)^2}$	$-\frac{1}{4}$	$\frac{2}{(r+2)^3}$	$\frac{1}{4}$
b_3	0	0	0	0
b_4	$\frac{1}{(r+2)^2(r+4)^2}$	$\frac{1}{64}$	$\frac{-4r-12}{(r+2)^3(r+4)^3}$	$-\frac{3}{128}$
b_5	0	0	0	0
b_6	$-\frac{1}{(r+2)^2(r+4)^2(r+6)^2}$	$-\frac{1}{2304}$	$\frac{6r^2+48r+88}{(r+2)^3(r+4)^3(r+6)^3}$	$\frac{11}{13824}$
b_7	0	0	0	0

The above table gives all values of b_n needed. Hence the second solution is

$$\begin{aligned} y_2(x) &= y_1(x) \ln(x) + b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 + b_7x^7 + b_8x^8 \dots \\ &= \left(1 - \frac{x^2}{4} + \frac{x^4}{64} - \frac{x^6}{2304} + O(x^8)\right) \ln(x) + \frac{x^2}{4} - \frac{3x^4}{128} + \frac{11x^6}{13824} + O(x^8) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= c_1 \left(1 - \frac{x^2}{4} + \frac{x^4}{64} - \frac{x^6}{2304} + O(x^8)\right) \\ &\quad + c_2 \left(\left(1 - \frac{x^2}{4} + \frac{x^4}{64} - \frac{x^6}{2304} + O(x^8)\right) \ln(x) + \frac{x^2}{4} - \frac{3x^4}{128} + \frac{11x^6}{13824} + O(x^8)\right) \end{aligned}$$

Hence the final solution is

$$\begin{aligned} y &= y_h \\ &= c_1 \left(1 - \frac{x^2}{4} + \frac{x^4}{64} - \frac{x^6}{2304} + O(x^8)\right) \\ &\quad + c_2 \left(\left(1 - \frac{x^2}{4} + \frac{x^4}{64} - \frac{x^6}{2304} + O(x^8)\right) \ln(x) + \frac{x^2}{4} - \frac{3x^4}{128} + \frac{11x^6}{13824} + O(x^8)\right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \left(1 - \frac{x^2}{4} + \frac{x^4}{64} - \frac{x^6}{2304} + O(x^8) \right) + c_2 \left(\left(1 - \frac{x^2}{4} + \frac{x^4}{64} - \frac{x^6}{2304} + O(x^8) \right) \ln(x) + \frac{x^2}{4} - \frac{3x^4}{128} + \frac{11x^6}{13824} + O(x^8) \right) \quad (1)$$

Verification of solutions

$$y = c_1 \left(1 - \frac{x^2}{4} + \frac{x^4}{64} - \frac{x^6}{2304} + O(x^8) \right) + c_2 \left(\left(1 - \frac{x^2}{4} + \frac{x^4}{64} - \frac{x^6}{2304} + O(x^8) \right) \ln(x) + \frac{x^2}{4} - \frac{3x^4}{128} + \frac{11x^6}{13824} + O(x^8) \right)$$

Verified OK.

19.18.1 Maple step by step solution

Let's solve

$$x^2 y'' + xy' + yx^2 = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{y'}{x} - y$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{y'}{x} + y = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$[P_2(x) = \frac{1}{x}, P_3(x) = 1]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 1$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 0$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$y''x + y' + xy = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x \cdot y$ to series expansion

$$x \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+1}$$

- Shift index using $k- > k - 1$

$$x \cdot y = \sum_{k=1}^{\infty} a_{k-1} x^{k+r}$$

- Convert y' to series expansion

$$y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1}$$

- Shift index using $k- > k + 1$

$$y' = \sum_{k=-1}^{\infty} a_{k+1} (k+r+1) x^{k+r}$$

- Convert $x \cdot y''$ to series expansion

$$x \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r-1}$$

- Shift index using $k- > k + 1$

$$x \cdot y'' = \sum_{k=-1}^{\infty} a_{k+1} (k+r+1)(k+r) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0 r^2 x^{-1+r} + a_1 (1+r)^2 x^r + \left(\sum_{k=1}^{\infty} (a_{k+1} (k+r+1)^2 + a_{k-1}) x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$r^2 = 0$$

- Values of r that satisfy the indicial equation

$$r = 0$$

- Each term must be 0

$$a_1(1+r)^2 = 0$$
- Each term in the series must be 0, giving the recursion relation

$$a_{k+1}(k+1)^2 + a_{k-1} = 0$$
- Shift index using $k \rightarrow k+1$

$$a_{k+2}(k+2)^2 + a_k = 0$$
- Recursion relation that defines series solution to ODE

$$a_{k+2} = -\frac{a_k}{(k+2)^2}$$
- Recursion relation for $r = 0$

$$a_{k+2} = -\frac{a_k}{(k+2)^2}$$
- Solution for $r = 0$

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+2} = -\frac{a_k}{(k+2)^2}, a_1 = 0 \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
    <- Bessel successful
<- special function solution successful`

```

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 47

```
Order:=8;
```

```
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)+x^2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (c_2 \ln(x) + c_1) \left(1 - \frac{1}{4}x^2 + \frac{1}{64}x^4 - \frac{1}{2304}x^6 + O(x^8) \right) \\ + \left(\frac{1}{4}x^2 - \frac{3}{128}x^4 + \frac{11}{13824}x^6 + O(x^8) \right) c_2$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 81

```
AsymptoticDSolveValue[x^2*y'[x]+x*y'[x]+x^2*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(-\frac{x^6}{2304} + \frac{x^4}{64} - \frac{x^2}{4} + 1 \right) \\ + c_2 \left(\frac{11x^6}{13824} - \frac{3x^4}{128} + \frac{x^2}{4} + \left(-\frac{x^6}{2304} + \frac{x^4}{64} - \frac{x^2}{4} + 1 \right) \log(x) \right)$$

19.19 problem 6

Internal problem ID [6459]

Internal file name [OUTPUT/5707_Sunday_June_05_2022_03_48_08_PM_91319078/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.4. REGULAR SINGULAR POINTS. Page 175

Problem number: 6.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_change_of_variable_on_y_method_2", "second order series method. Irregular singular point"

Maple gives the following as the ode type

```
[[_2nd_order, _with_linear_symmetries], [_2nd_order, _linear, `
  _with_symmetry_[0,F(x)]`]]
```

Unable to solve or complete the solution.

$$y'' + \frac{y'}{x^2} - \frac{y}{x^3} = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$y'' + \frac{y'}{x^2} - \frac{y}{x^3} = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{1}{x^2}$$
$$q(x) = -\frac{1}{x^3}$$

Table 465: Table $p(x), q(x)$ singularities.

$p(x) = \frac{1}{x^2}$	
singularity	type
$x = 0$	“irregular”

$q(x) = -\frac{1}{x^3}$	
singularity	type
$x = 0$	“irregular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[\infty]$

Irregular singular points : $[0]$

Since $x = 0$ is not an ordinary point, then we will now check if it is a regular singular point. Unable to solve since $x = 0$ is not regular singular point. Terminating.

Verification of solutions N/A

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
<- linear_1 successful`

```

X Solution by Maple

```

Order:=8;
dsolve(diff(y(x),x$2)+1/x^2*diff(y(x),x)-1/x^3*y(x)=0,y(x),type='series',x=0);

```

No solution found

✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 17

```
AsymptoticDSolveValue[y''[x]+1/x^2*y'[x]-1/x^3*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 x + c_2 e^{\frac{1}{x}} x$$

19.20 problem 8

Internal problem ID [6460]

Internal file name [OUTPUT/5708_Sunday_June_05_2022_03_48_09_PM_26176461/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.4. REGULAR SINGULAR POINTS. Page 175

Problem number: 8.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact linear second order ode", "second_order_integrable_as_is", "second order series method. Irregular singular point"

Maple gives the following as the ode type

```
[[_2nd_order , _exact , _linear , _homogeneous]]
```

Unable to solve or complete the solution.

$$x^2y'' + (3x - 1)y' + y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2y'' + (3x - 1)y' + y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{3x - 1}{x^2}$$
$$q(x) = \frac{1}{x^2}$$

Table 466: Table $p(x), q(x)$ singularities.

$p(x) = \frac{3x-1}{x^2}$	
singularity	type
$x = 0$	“irregular”

$q(x) = \frac{1}{x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[\infty]$

Irregular singular points : $[0]$

Since $x = 0$ is not an ordinary point, then we will now check if it is a regular singular point. Unable to solve since $x = 0$ is not regular singular point. Terminating.

Verification of solutions N/A

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
<- linear_1 successful`

```

X Solution by Maple

```

Order:=8;
dsolve(x^2*diff(y(x),x$2)+(3*x-1)*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);

```

No solution found

✓ Solution by Mathematica

Time used: 0.043 (sec). Leaf size: 53

```
AsymptoticDSolveValue[x^2*y''[x]+(3*x-1)*y'[x]+y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1(5040x^7 + 720x^6 + 120x^5 + 24x^4 + 6x^3 + 2x^2 + x + 1) + \frac{c_2 e^{-1/x}}{x}$$

20 Chapter 4. Power Series Solutions and Special Functions. Section 4.5. More on Regular Singular Points. Page 183

20.1	problem 1	3358
20.2	problem 2	3371
20.3	problem 3(a)	3384
20.4	problem 3(b)	3399
20.5	problem 3(c)	3416
20.6	problem 4	3430
20.7	problem 5	3443
20.8	problem 6	3456
20.9	problem 7	3473

20.1 problem 1

20.1.1 Maple step by step solution 3367

Internal problem ID [6461]

Internal file name [OUTPUT/5709_Sunday_June_05_2022_03_48_10_PM_16788399/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.5. More on Regular Singular Points. Page 183

Problem number: 1.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Repeated root"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$x^2y'' - 3xy' + (4x + 4)y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2y'' - 3xy' + (4x + 4)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{3}{x}$$
$$q(x) = \frac{4x + 4}{x^2}$$

Table 467: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{3}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = \frac{4x+4}{x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2 y'' - 3xy' + (4x + 4)y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned}
 & x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\
 & - 3x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (4x+4) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0
 \end{aligned} \tag{1}$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-3x^{n+r} a_n (n+r)) \\ & + \left(\sum_{n=0}^{\infty} 4x^{1+n+r} a_n \right) + \left(\sum_{n=0}^{\infty} 4a_n x^{n+r} \right) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} 4x^{1+n+r} a_n = \sum_{n=1}^{\infty} 4a_{n-1} x^{n+r}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-3x^{n+r} a_n (n+r)) \\ & + \left(\sum_{n=1}^{\infty} 4a_{n-1} x^{n+r} \right) + \left(\sum_{n=0}^{\infty} 4a_n x^{n+r} \right) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r} a_n (n+r) (n+r-1) - 3x^{n+r} a_n (n+r) + 4a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$x^r a_0 r(-1+r) - 3x^r a_0 r + 4a_0 x^r = 0$$

Or

$$(x^r r(-1+r) - 3x^r r + 4x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(r-2)^2 x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$(r-2)^2 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = 2$$

$$r_2 = 2$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(r - 2)^2 x^r = 0$$

Solving for r gives the roots of the indicial equation as Since the root of the indicial equation is repeated, then we can construct two linearly independent solutions. The first solution has the form

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+r} \quad (1A)$$

Now the second solution y_2 is found using

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right) \quad (1B)$$

Then the general solution will be

$$y = c_1 y_1(x) + c_2 y_2(x)$$

In Eq (1B) the sum starts from 1 and not zero. In Eq (1A), a_0 is never zero, and is arbitrary and is typically taken as $a_0 = 1$, and $\{c_1, c_2\}$ are two arbitray constants of integration which can be found from initial conditions. Using the value of the indicial root found earlier, $r = 2$, Eqs (1A,1B) become

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+2}$$

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+2} \right)$$

We start by finding the first solution $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) - 3a_n(n+r) + 4a_{n-1} + 4a_n = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{4a_{n-1}}{n^2 + 2nr + r^2 - 4n - 4r + 4} \quad (4)$$

Which for the root $r = 2$ becomes

$$a_n = -\frac{4a_{n-1}}{n^2} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 2$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = -\frac{4}{(-1+r)^2}$$

Which for the root $r = 2$ becomes

$$a_1 = -4$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{4}{(-1+r)^2}$	-4

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{16}{(-1+r)^2 r^2}$$

Which for the root $r = 2$ becomes

$$a_2 = 4$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{4}{(-1+r)^2}$	-4
a_2	$\frac{16}{(-1+r)^2 r^2}$	4

For $n = 3$, using the above recursive equation gives

$$a_3 = -\frac{64}{(-1+r)^2 r^2 (r+1)^2}$$

Which for the root $r = 2$ becomes

$$a_3 = -\frac{16}{9}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{4}{(-1+r)^2}$	-4
a_2	$\frac{16}{(-1+r)^2 r^2}$	4
a_3	$-\frac{64}{(-1+r)^2 r^2 (r+1)^2}$	$-\frac{16}{9}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{256}{(-1+r)^2 r^2 (r+1)^2 (r+2)^2}$$

Which for the root $r = 2$ becomes

$$a_4 = \frac{4}{9}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{4}{(-1+r)^2}$	-4
a_2	$\frac{16}{(-1+r)^2 r^2}$	4
a_3	$-\frac{64}{(-1+r)^2 r^2 (r+1)^2}$	$-\frac{16}{9}$
a_4	$\frac{256}{(-1+r)^2 r^2 (r+1)^2 (r+2)^2}$	$\frac{4}{9}$

For $n = 5$, using the above recursive equation gives

$$a_5 = -\frac{1024}{(-1+r)^2 r^2 (r+1)^2 (r+2)^2 (r+3)^2}$$

Which for the root $r = 2$ becomes

$$a_5 = -\frac{16}{225}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{4}{(-1+r)^2}$	-4
a_2	$\frac{16}{(-1+r)^2 r^2}$	4
a_3	$-\frac{64}{(-1+r)^2 r^2 (r+1)^2}$	$-\frac{16}{9}$
a_4	$\frac{256}{(-1+r)^2 r^2 (r+1)^2 (r+2)^2}$	$\frac{4}{9}$
a_5	$-\frac{1024}{(-1+r)^2 r^2 (r+1)^2 (r+2)^2 (r+3)^2}$	$-\frac{16}{225}$

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{4096}{(-1+r)^2 r^2 (r+1)^2 (r+2)^2 (r+3)^2 (r+4)^2}$$

Which for the root $r = 2$ becomes

$$a_6 = \frac{16}{2025}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{4}{(-1+r)^2}$	-4
a_2	$\frac{16}{(-1+r)^2 r^2}$	4
a_3	$-\frac{64}{(-1+r)^2 r^2 (r+1)^2}$	$-\frac{16}{9}$
a_4	$\frac{256}{(-1+r)^2 r^2 (r+1)^2 (r+2)^2}$	$\frac{4}{9}$
a_5	$-\frac{1024}{(-1+r)^2 r^2 (r+1)^2 (r+2)^2 (r+3)^2}$	$-\frac{16}{225}$
a_6	$\frac{4096}{(-1+r)^2 r^2 (r+1)^2 (r+2)^2 (r+3)^2 (r+4)^2}$	$\frac{16}{2025}$

For $n = 7$, using the above recursive equation gives

$$a_7 = -\frac{16384}{(-1+r)^2 r^2 (r+1)^2 (r+2)^2 (r+3)^2 (r+4)^2 (r+5)^2}$$

Which for the root $r = 2$ becomes

$$a_7 = -\frac{64}{99225}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{4}{(-1+r)^2}$	-4
a_2	$\frac{16}{(-1+r)^2 r^2}$	4
a_3	$-\frac{64}{(-1+r)^2 r^2 (r+1)^2}$	$-\frac{16}{9}$
a_4	$\frac{256}{(-1+r)^2 r^2 (r+1)^2 (r+2)^2}$	$\frac{4}{9}$
a_5	$-\frac{1024}{(-1+r)^2 r^2 (r+1)^2 (r+2)^2 (r+3)^2}$	$-\frac{16}{225}$
a_6	$\frac{4096}{(-1+r)^2 r^2 (r+1)^2 (r+2)^2 (r+3)^2 (r+4)^2}$	$\frac{16}{2025}$
a_7	$-\frac{16384}{(-1+r)^2 r^2 (r+1)^2 (r+2)^2 (r+3)^2 (r+4)^2 (r+5)^2}$	$-\frac{64}{99225}$

Using the above table, then the first solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x^2(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots) \\ &= x^2\left(4x^2 - 4x + 1 - \frac{16x^3}{9} + \frac{4x^4}{9} - \frac{16x^5}{225} + \frac{16x^6}{2025} - \frac{64x^7}{99225} + O(x^8)\right) \end{aligned}$$

Now the second solution is found. The second solution is given by

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r}\right)$$

Where b_n is found using

$$b_n = \frac{d}{dr} a_{n,r}$$

And the above is then evaluated at $r = 2$. The above table for $a_{n,r}$ is used for this purpose. Computing the derivatives gives the following table

n	$b_{n,r}$	a_n	$b_{n,r} = \frac{d}{dr} a_{n,r}$
b_0	1	1	N/A since b_n starts from 1
b_1	$-\frac{4}{(-1+r)^2}$	-4	$\frac{8}{(-1+r)^3}$
b_2	$\frac{16}{(-1+r)^2 r^2}$	4	$\frac{-64r+32}{(-1+r)^3 r^3}$
b_3	$-\frac{64}{(-1+r)^2 r^2 (r+1)^2}$	$-\frac{16}{9}$	$\frac{384r^2-128}{(-1+r)^3 r^3 (r+1)^3}$
b_4	$\frac{256}{(-1+r)^2 r^2 (r+1)^2 (r+2)^2}$	$\frac{4}{9}$	$\frac{-2048r^3-3072r^2+1024r+1024}{(-1+r)^3 r^3 (r+1)^3 (r+2)^3}$
b_5	$-\frac{1024}{(-1+r)^2 r^2 (r+1)^2 (r+2)^2 (r+3)^2}$	$-\frac{16}{225}$	$\frac{10240r^4+40960r^3+30720r^2-20480r-12288}{(-1+r)^3 r^3 (r+1)^3 (r+2)^3 (r+3)^3}$
b_6	$\frac{4096}{(-1+r)^2 r^2 (r+1)^2 (r+2)^2 (r+3)^2 (r+4)^2}$	$\frac{16}{2025}$	$-\frac{49152(r^4+6r^3+\frac{23}{3}r^2-4r-\frac{8}{3})(\frac{3}{2}+r)}{(-1+r)^3 r^3 (r+1)^3 (r+2)^3 (r+3)^3 (r+4)^3}$
b_7	$-\frac{16384}{(-1+r)^2 r^2 (r+1)^2 (r+2)^2 (r+3)^2 (r+4)^2 (r+5)^2}$	$-\frac{64}{99225}$	$\frac{229376r^6+2752512r^5+11468800r^4+18350080r^3+4816896r^2-10092544r-1000000}{(-1+r)^3 r^3 (r+1)^3 (r+2)^3 (r+3)^3 (r+4)^3 (r+5)^3}$

The above table gives all values of b_n needed. Hence the second solution is

$$\begin{aligned}
y_2(x) &= y_1(x) \ln(x) + b_0 + b_1 x + b_2 x^2 + b_3 x^3 + b_4 x^4 + b_5 x^5 + b_6 x^6 + b_7 x^7 + b_8 x^8 \dots \\
&= x^2 \left(4x^2 - 4x + 1 - \frac{16x^3}{9} + \frac{4x^4}{9} - \frac{16x^5}{225} + \frac{16x^6}{2025} - \frac{64x^7}{99225} + O(x^8) \right) \ln(x) \\
&\quad + x^2 \left(-12x^2 + 8x + \frac{176x^3}{27} - \frac{50x^4}{27} + \frac{1096x^5}{3375} - \frac{392x^6}{10125} + \frac{3872x^7}{1157625} + O(x^8) \right)
\end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
y_h(x) &= c_1 y_1(x) + c_2 y_2(x) \\
&= c_1 x^2 \left(4x^2 - 4x + 1 - \frac{16x^3}{9} + \frac{4x^4}{9} - \frac{16x^5}{225} + \frac{16x^6}{2025} - \frac{64x^7}{99225} + O(x^8) \right) \\
&\quad + c_2 \left(x^2 \left(4x^2 - 4x + 1 - \frac{16x^3}{9} + \frac{4x^4}{9} - \frac{16x^5}{225} + \frac{16x^6}{2025} - \frac{64x^7}{99225} + O(x^8) \right) \ln(x) \right. \\
&\quad \left. + x^2 \left(-12x^2 + 8x + \frac{176x^3}{27} - \frac{50x^4}{27} + \frac{1096x^5}{3375} - \frac{392x^6}{10125} + \frac{3872x^7}{1157625} + O(x^8) \right) \right)
\end{aligned}$$

Hence the final solution is

$$y = y_h$$

$$\begin{aligned}
&= c_1 x^2 \left(4x^2 - 4x + 1 - \frac{16x^3}{9} + \frac{4x^4}{9} - \frac{16x^5}{225} + \frac{16x^6}{2025} - \frac{64x^7}{99225} + O(x^8) \right) \\
&\quad + c_2 \left(x^2 \left(4x^2 - 4x + 1 - \frac{16x^3}{9} + \frac{4x^4}{9} - \frac{16x^5}{225} + \frac{16x^6}{2025} - \frac{64x^7}{99225} + O(x^8) \right) \ln(x) \right. \\
&\quad \left. + x^2 \left(-12x^2 + 8x + \frac{176x^3}{27} - \frac{50x^4}{27} + \frac{1096x^5}{3375} - \frac{392x^6}{10125} + \frac{3872x^7}{1157625} + O(x^8) \right) \right)
\end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}
y &= c_1 x^2 \left(4x^2 - 4x + 1 - \frac{16x^3}{9} + \frac{4x^4}{9} - \frac{16x^5}{225} + \frac{16x^6}{2025} - \frac{64x^7}{99225} + O(x^8) \right) \\
&\quad + c_2 \left(x^2 \left(4x^2 - 4x + 1 - \frac{16x^3}{9} + \frac{4x^4}{9} - \frac{16x^5}{225} + \frac{16x^6}{2025} - \frac{64x^7}{99225} + O(x^8) \right) \ln(x) \right. \\
&\quad \left. + x^2 \left(-12x^2 + 8x + \frac{176x^3}{27} - \frac{50x^4}{27} + \frac{1096x^5}{3375} - \frac{392x^6}{10125} + \frac{3872x^7}{1157625} + O(x^8) \right) \right)
\end{aligned}$$

Verification of solutions

$$\begin{aligned}
y &= c_1 x^2 \left(4x^2 - 4x + 1 - \frac{16x^3}{9} + \frac{4x^4}{9} - \frac{16x^5}{225} + \frac{16x^6}{2025} - \frac{64x^7}{99225} + O(x^8) \right) \\
&\quad + c_2 \left(x^2 \left(4x^2 - 4x + 1 - \frac{16x^3}{9} + \frac{4x^4}{9} - \frac{16x^5}{225} + \frac{16x^6}{2025} - \frac{64x^7}{99225} + O(x^8) \right) \ln(x) \right. \\
&\quad \left. + x^2 \left(-12x^2 + 8x + \frac{176x^3}{27} - \frac{50x^4}{27} + \frac{1096x^5}{3375} - \frac{392x^6}{10125} + \frac{3872x^7}{1157625} + O(x^8) \right) \right)
\end{aligned}$$

Verified OK.

20.1.1 Maple step by step solution

Let's solve

$$x^2 y'' - 3xy' + (4x + 4)y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = \frac{3y'}{x} - \frac{4(1+x)y}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - \frac{3y'}{x} + \frac{4(1+x)y}{x^2} = 0$$

□ Check to see if $x_0 = 0$ is a regular singular point

○ Define functions

$$\left[P_2(x) = -\frac{3}{x}, P_3(x) = \frac{4(1+x)}{x^2} \right]$$

○ $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = -3$$

○ $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 4$$

○ $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

• Multiply by denominators

$$x^2 y'' - 3xy' + (4x + 4)y = 0$$

• Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

□ Rewrite ODE with series expansions

○ Convert $x^m \cdot y$ to series expansion for $m = 0..1$

$$x^m \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

○ Shift index using $k \rightarrow k - m$

$$x^m \cdot y = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

○ Convert $x \cdot y'$ to series expansion

$$x \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r}$$

○ Convert $x^2 \cdot y''$ to series expansion

$$x^2 \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(-2+r)^2 x^r + \left(\sum_{k=1}^{\infty} (a_k(k+r-2)^2 + 4a_{k-1}) x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation $(-2+r)^2 = 0$
- Values of r that satisfy the indicial equation
 $r = 2$
- Each term in the series must be 0, giving the recursion relation
 $a_k(k+r-2)^2 + 4a_{k-1} = 0$
- Shift index using $k \rightarrow k+1$
 $a_{k+1}(k+r-1)^2 + 4a_k = 0$
- Recursion relation that defines series solution to ODE
 $a_{k+1} = -\frac{4a_k}{(k+r-1)^2}$
- Recursion relation for $r = 2$
 $a_{k+1} = -\frac{4a_k}{(k+1)^2}$
- Solution for $r = 2$
 $\left[y = \sum_{k=0}^{\infty} a_k x^{k+2}, a_{k+1} = -\frac{4a_k}{(k+1)^2} \right]$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
    <- Bessel successful
<- special function solution successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 81

Order:=8;

```
dsolve(x^2*diff(y(x),x$2)-3*x*diff(y(x),x)+(4*x+4)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left((c_2 \ln(x) + c_1) \left(1 - 4x + 4x^2 - \frac{16}{9}x^3 + \frac{4}{9}x^4 - \frac{16}{225}x^5 + \frac{16}{2025}x^6 - \frac{64}{99225}x^7 + O(x^8) \right) + \left(8x - 12x^2 + \frac{176}{27}x^3 - \frac{50}{27}x^4 + \frac{1096}{3375}x^5 - \frac{392}{10125}x^6 + \frac{3872}{1157625}x^7 + O(x^8) \right) c_2 \right) x^2$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 158

```
AsymptoticDSolveValue[x^2*y''[x]-3*x*y'[x]+(4*x+4)*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(-\frac{64x^7}{99225} + \frac{16x^6}{2025} - \frac{16x^5}{225} + \frac{4x^4}{9} - \frac{16x^3}{9} + 4x^2 - 4x + 1 \right) x^2 + c_2 \left(\left(\frac{3872x^7}{1157625} - \frac{392x^6}{10125} + \frac{1096x^5}{3375} - \frac{50x^4}{27} + \frac{176x^3}{27} - 12x^2 + 8x \right) x^2 + \left(-\frac{64x^7}{99225} + \frac{16x^6}{2025} - \frac{16x^5}{225} + \frac{4x^4}{9} - \frac{16x^3}{9} + 4x^2 - 4x + 1 \right) x^2 \log(x) \right)$$

20.2 problem 2

20.2.1 Maple step by step solution 3380

Internal problem ID [6462]

Internal file name [OUTPUT/5710_Sunday_June_05_2022_03_48_13_PM_75573943/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.5. More on Regular Singular Points. Page 183

Problem number: 2.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Repeated root"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$4x^2y'' - 8x^2y' + (4x^2 + 1)y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$4x^2y'' - 8x^2y' + (4x^2 + 1)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -2$$
$$q(x) = \frac{4x^2 + 1}{4x^2}$$

Table 469: Table $p(x), q(x)$ singularities.

$p(x) = -2$	
singularity	type

$q(x) = \frac{4x^2+1}{4x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$4x^2y'' - 8x^2y' + (4x^2 + 1)y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$4x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) - 8x^2 \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (4x^2 + 1) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 4x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-8x^{1+n+r} a_n (n+r)) \\ & + \left(\sum_{n=0}^{\infty} 4x^{n+r+2} a_n \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=0}^{\infty} (-8x^{1+n+r} a_n (n+r)) &= \sum_{n=1}^{\infty} (-8a_{n-1} (n+r-1) x^{n+r}) \\ \sum_{n=0}^{\infty} 4x^{n+r+2} a_n &= \sum_{n=2}^{\infty} 4a_{n-2} x^{n+r} \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 4x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=1}^{\infty} (-8a_{n-1} (n+r-1) x^{n+r}) \\ & + \left(\sum_{n=2}^{\infty} 4a_{n-2} x^{n+r} \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$4x^{n+r} a_n (n+r) (n+r-1) + a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$4x^r a_0 r (-1+r) + a_0 x^r = 0$$

Or

$$(4x^r r (-1+r) + x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$x^r (2r-1)^2 = 0$$

Since the above is true for all x then the indicial equation becomes

$$(2r - 1)^2 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = \frac{1}{2}$$

$$r_2 = \frac{1}{2}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$x^r(2r - 1)^2 = 0$$

Solving for r gives the roots of the indicial equation as Since the root of the indicial equation is repeated, then we can construct two linearly independent solutions. The first solution has the form

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+r} \quad (1A)$$

Now the second solution y_2 is found using

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right) \quad (1B)$$

Then the general solution will be

$$y = c_1 y_1(x) + c_2 y_2(x)$$

In Eq (1B) the sum starts from 1 and not zero. In Eq (1A), a_0 is never zero, and is arbitrary and is typically taken as $a_0 = 1$, and $\{c_1, c_2\}$ are two arbitrary constants of integration which can be found from initial conditions. Using the value of the indicial root found earlier, $r = \frac{1}{2}$, Eqs (1A,1B) become

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+\frac{1}{2}}$$

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+\frac{1}{2}} \right)$$

We start by finding the first solution $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the

indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = \frac{8r}{(2r+1)^2}$$

For $2 \leq n$ the recursive equation is

$$4a_n(n+r)(n+r-1) - 8a_{n-1}(n+r-1) + 4a_{n-2} + a_n = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{8na_{n-1} + 8ra_{n-1} - 4a_{n-2} - 8a_{n-1}}{4n^2 + 8nr + 4r^2 - 4n - 4r + 1} \quad (4)$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_n = \frac{(2n-1)a_{n-1} - a_{n-2}}{n^2} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = \frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{8r}{(2r+1)^2}$	1

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{48r^2 + 48r - 4}{(3+2r)^2(2r+1)^2}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_2 = \frac{1}{2}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{8r}{(2r+1)^2}$	1
a_2	$\frac{48r^2+48r-4}{(3+2r)^2(2r+1)^2}$	$\frac{1}{2}$

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{256r^3 + 768r^2 + 448r - 64}{(3 + 2r)^2 (5 + 2r)^2 (2r + 1)^2}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_3 = \frac{1}{6}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{8r}{(2r+1)^2}$	1
a_2	$\frac{48r^2+48r-4}{(3+2r)^2(2r+1)^2}$	$\frac{1}{2}$
a_3	$\frac{256r^3+768r^2+448r-64}{(3+2r)^2(5+2r)^2(2r+1)^2}$	$\frac{1}{6}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1280r^4 + 7680r^3 + 13440r^2 + 5760r - 1136}{(3 + 2r)^2 (5 + 2r)^2 (2r + 1)^2 (7 + 2r)^2}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_4 = \frac{1}{24}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{8r}{(2r+1)^2}$	1
a_2	$\frac{48r^2+48r-4}{(3+2r)^2(2r+1)^2}$	$\frac{1}{2}$
a_3	$\frac{256r^3+768r^2+448r-64}{(3+2r)^2(5+2r)^2(2r+1)^2}$	$\frac{1}{6}$
a_4	$\frac{1280r^4+7680r^3+13440r^2+5760r-1136}{(3+2r)^2(5+2r)^2(2r+1)^2(7+2r)^2}$	$\frac{1}{24}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{6144(2 + r) (r^4 + 8r^3 + \frac{109}{6}r^2 + \frac{26}{3}r - \frac{31}{16})}{(3 + 2r)^2 (5 + 2r)^2 (2r + 1)^2 (7 + 2r)^2 (9 + 2r)^2}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_5 = \frac{1}{120}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{8r}{(2r+1)^2}$	1
a_2	$\frac{48r^2+48r-4}{(3+2r)^2(2r+1)^2}$	$\frac{1}{2}$
a_3	$\frac{256r^3+768r^2+448r-64}{(3+2r)^2(5+2r)^2(2r+1)^2}$	$\frac{1}{6}$
a_4	$\frac{1280r^4+7680r^3+13440r^2+5760r-1136}{(3+2r)^2(5+2r)^2(2r+1)^2(7+2r)^2}$	$\frac{1}{24}$
a_5	$\frac{6144(2+r)(r^4+8r^3+\frac{109}{6}r^2+\frac{26}{3}r-\frac{31}{16})}{(3+2r)^2(5+2r)^2(2r+1)^2(7+2r)^2(9+2r)^2}$	$\frac{1}{120}$

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{28672r^6 + 430080r^5 + 2401280r^4 + 6092800r^3 + 6650112r^2 + 1890560r - 584256}{(9+2r)^2(7+2r)^2(11+2r)^2(3+2r)^2(5+2r)^2(2r+1)^2}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_6 = \frac{1}{720}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{8r}{(2r+1)^2}$	1
a_2	$\frac{48r^2+48r-4}{(3+2r)^2(2r+1)^2}$	$\frac{1}{2}$
a_3	$\frac{256r^3+768r^2+448r-64}{(3+2r)^2(5+2r)^2(2r+1)^2}$	$\frac{1}{6}$
a_4	$\frac{1280r^4+7680r^3+13440r^2+5760r-1136}{(3+2r)^2(5+2r)^2(2r+1)^2(7+2r)^2}$	$\frac{1}{24}$
a_5	$\frac{6144(2+r)(r^4+8r^3+\frac{109}{6}r^2+\frac{26}{3}r-\frac{31}{16})}{(3+2r)^2(5+2r)^2(2r+1)^2(7+2r)^2(9+2r)^2}$	$\frac{1}{120}$
a_6	$\frac{28672r^6+430080r^5+2401280r^4+6092800r^3+6650112r^2+1890560r-584256}{(9+2r)^2(7+2r)^2(11+2r)^2(3+2r)^2(5+2r)^2(2r+1)^2}$	$\frac{1}{720}$

For $n = 7$, using the above recursive equation gives

$$a_7 = \frac{131072(r^6 + 18r^5 + \frac{477}{4}r^4 + 351r^3 + \frac{6819}{16}r^2 + \frac{1017}{8}r - \frac{2689}{64})(r+3)}{(9+2r)^2(7+2r)^2(11+2r)^2(3+2r)^2(5+2r)^2(2r+1)^2(13+2r)^2}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_7 = \frac{1}{5040}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{8r}{(2r+1)^2}$	1
a_2	$\frac{48r^2+48r-4}{(3+2r)^2(2r+1)^2}$	$\frac{1}{2}$
a_3	$\frac{256r^3+768r^2+448r-64}{(3+2r)^2(5+2r)^2(2r+1)^2}$	$\frac{1}{6}$
a_4	$\frac{1280r^4+7680r^3+13440r^2+5760r-1136}{(3+2r)^2(5+2r)^2(2r+1)^2(7+2r)^2}$	$\frac{1}{24}$
a_5	$\frac{6144(2+r)(r^4+8r^3+\frac{109}{6}r^2+\frac{26}{3}r-\frac{31}{16})}{(3+2r)^2(5+2r)^2(2r+1)^2(7+2r)^2(9+2r)^2}$	$\frac{1}{120}$
a_6	$\frac{28672r^6+430080r^5+2401280r^4+6092800r^3+6650112r^2+1890560r-584256}{(9+2r)^2(7+2r)^2(11+2r)^2(3+2r)^2(5+2r)^2(2r+1)^2}$	$\frac{1}{720}$
a_7	$\frac{131072(r^6+18r^5+\frac{477}{4}r^4+351r^3+\frac{6819}{16}r^2+\frac{1017}{8}r-\frac{2689}{64})(r+3)}{(9+2r)^2(7+2r)^2(11+2r)^2(3+2r)^2(5+2r)^2(2r+1)^2(13+2r)^2}$	$\frac{1}{5040}$

Using the above table, then the first solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= \sqrt{x}(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots) \\ &= \sqrt{x} \left(1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \frac{x^5}{120} + \frac{x^6}{720} + \frac{x^7}{5040} + O(x^8) \right) \end{aligned}$$

Now the second solution is found. The second solution is given by

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right)$$

Where b_n is found using

$$b_n = \frac{d}{dr} a_{n,r}$$

And the above is then evaluated at $r = \frac{1}{2}$. The above table for $a_{n,r}$ is used for this purpose. Computing the derivatives gives the following table

n	$b_{n,r}$	a_n	$b_{n,r} = \frac{d}{dr} a_{n,r}$
b_0	1	1	N/A since b_n starts from 1
b_1	$\frac{8r}{(2r+1)^2}$	1	$\frac{-16r+8}{(2r+1)^3}$
b_2	$\frac{48r^2+48r-4}{(3+2r)^2(2r+1)^2}$	$\frac{1}{2}$	$\frac{-384r^3-576r^2-32r+208}{(3+2r)^3(2r+1)^3}$
b_3	$\frac{256r^3+768r^2+448r-64}{(3+2r)^2(5+2r)^2(2r+1)^2}$	$\frac{1}{6}$	$-\frac{6144(r^4+6r^3+13r^2+12r+\frac{197}{48})(r-\frac{1}{2})}{(3+2r)^3(5+2r)^3(2r+1)^3}$
b_4	$\frac{1280r^4+7680r^3+13440r^2+5760r-1136}{(3+2r)^2(5+2r)^2(2r+1)^2(7+2r)^2}$	$\frac{1}{24}$	$-\frac{81920(r-\frac{1}{2})(r^6+12r^5+\frac{231}{4}r^4+142r^3+\frac{15003}{80}r^2+\frac{117}{4}r-\frac{1}{16})}{(3+2r)^3(5+2r)^3(2r+1)^3(7+2r)^2}$
b_5	$\frac{6144(2+r)(r^4+8r^3+\frac{109}{6}r^2+\frac{26}{3}r-\frac{31}{16})}{(3+2r)^2(5+2r)^2(2r+1)^2(7+2r)^2(9+2r)^2}$	$\frac{1}{120}$	$-\frac{983040(r-\frac{1}{2})(r^8+20r^7+\frac{508}{3}r^6+790r^5+\frac{17703}{8}r^4+\frac{117}{4}r^3-\frac{11}{16}r^2-\frac{1}{2}r+\frac{1}{16})}{(9+2r)^3(7+2r)^3(3+2r)^2}$
b_6	$\frac{28672r^6+430080r^5+2401280r^4+6092800r^3+6650112r^2+1890560r-584256}{(9+2r)^2(7+2r)^2(11+2r)^2(3+2r)^2(5+2r)^2(2r+1)^2}$	$\frac{1}{720}$	$-\frac{11010048(r-\frac{1}{2})(r^{10}+30r^9+\frac{1575}{4}r^8+2970r^7+\frac{117}{4}r^6+\frac{117}{4}r^5-\frac{11}{16}r^4-\frac{1}{2}r^3+\frac{1}{16}r^2-\frac{1}{2}r+\frac{1}{16})}{(9+2r)^2}$
b_7	$\frac{131072(r^6+18r^5+\frac{477}{4}r^4+351r^3+\frac{6819}{16}r^2+\frac{1017}{8}r-\frac{2689}{64})(r+3)}{(9+2r)^2(7+2r)^2(11+2r)^2(3+2r)^2(5+2r)^2(2r+1)^2(13+2r)^2}$	$\frac{1}{5040}$	$-\frac{117440512(r^{12}+42r^{11}+789r^{10}+8750r^9+\frac{1018}{15}r^8+\frac{117}{4}r^7-\frac{11}{16}r^6-\frac{1}{2}r^5+\frac{1}{16}r^4-\frac{1}{2}r^3+\frac{1}{16}r^2-\frac{1}{2}r+\frac{1}{16})}{(13+2r)^2}$

The above table gives all values of b_n needed. Hence the second solution is

$$y_2(x) = y_1(x) \ln(x) + b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 + b_7x^7 + b_8x^8 \dots$$

$$= \sqrt{x} \left(1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \frac{x^5}{120} + \frac{x^6}{720} + \frac{x^7}{5040} + O(x^8) \right) \ln(x) + \sqrt{x} O(x^8)$$

Therefore the homogeneous solution is

$$y_h(x) = c_1y_1(x) + c_2y_2(x)$$

$$= c_1\sqrt{x} \left(1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \frac{x^5}{120} + \frac{x^6}{720} + \frac{x^7}{5040} + O(x^8) \right)$$

$$+ c_2 \left(\sqrt{x} \left(1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \frac{x^5}{120} + \frac{x^6}{720} + \frac{x^7}{5040} + O(x^8) \right) \ln(x) + \sqrt{x} O(x^8) \right)$$

Hence the final solution is

$$y = y_h$$

$$= c_1\sqrt{x} \left(1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \frac{x^5}{120} + \frac{x^6}{720} + \frac{x^7}{5040} + O(x^8) \right)$$

$$+ c_2 \left(\sqrt{x} \left(1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \frac{x^5}{120} + \frac{x^6}{720} + \frac{x^7}{5040} + O(x^8) \right) \ln(x) + \sqrt{x} O(x^8) \right)$$

Summary

The solution(s) found are the following

$$y = c_1 \sqrt{x} \left(1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \frac{x^5}{120} + \frac{x^6}{720} + \frac{x^7}{5040} + O(x^8) \right) + c_2 \left(\sqrt{x} \left(1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \frac{x^5}{120} + \frac{x^6}{720} + \frac{x^7}{5040} + O(x^8) \right) \ln(x) + \sqrt{x} O(x^8) \right) \quad (1)$$

Verification of solutions

$$y = c_1 \sqrt{x} \left(1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \frac{x^5}{120} + \frac{x^6}{720} + \frac{x^7}{5040} + O(x^8) \right) + c_2 \left(\sqrt{x} \left(1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \frac{x^5}{120} + \frac{x^6}{720} + \frac{x^7}{5040} + O(x^8) \right) \ln(x) + \sqrt{x} O(x^8) \right)$$

Verified OK.

20.2.1 Maple step by step solution

Let's solve

$$4x^2 y'' - 8x^2 y' + (4x^2 + 1)y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = 2y' - \frac{(4x^2+1)y}{4x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - 2y' + \frac{(4x^2+1)y}{4x^2} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$\left[P_2(x) = -2, P_3(x) = \frac{4x^2+1}{4x^2} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 0$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = \frac{1}{4}$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$4x^2 y'' - 8x^2 y' + (4x^2 + 1)y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 0..2$

$$x^m \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

- Shift index using $k \rightarrow k - m$

$$x^m \cdot y = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

- Convert $x^2 \cdot y'$ to series expansion

$$x^2 \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r+1}$$

- Shift index using $k \rightarrow k - 1$

$$x^2 \cdot y' = \sum_{k=1}^{\infty} a_{k-1} (k-1+r) x^{k+r}$$

- Convert $x^2 \cdot y''$ to series expansion

$$x^2 \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k-1+r) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(-1+2r)^2 x^r + (a_1(1+2r)^2 - 8a_0 r) x^{1+r} + \left(\sum_{k=2}^{\infty} (a_k(2k+2r-1)^2 - 8a_{k-1}(k-1+r) + 4a_{k-2}(k+r)(k-1+r)) x^{k+r} \right)$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$(-1+2r)^2 = 0$$

- Values of r that satisfy the indicial equation

$$r = \frac{1}{2}$$

- Each term must be 0

$$a_1(1 + 2r)^2 - 8a_0r = 0$$
- Solve for the dependent coefficient(s)

$$a_1 = \frac{8a_0r}{4r^2+4r+1}$$
- Each term in the series must be 0, giving the recursion relation

$$a_k(2k + 2r - 1)^2 + (-8k - 8r + 8) a_{k-1} + 4a_{k-2} = 0$$
- Shift index using $k \rightarrow k + 2$

$$a_{k+2}(2k + 3 + 2r)^2 + (-8k - 8 - 8r) a_{k+1} + 4a_k = 0$$
- Recursion relation that defines series solution to ODE

$$a_{k+2} = \frac{4(2ka_{k+1}+2ra_{k+1}-a_k+2a_{k+1})}{(2k+3+2r)^2}$$
- Recursion relation for $r = \frac{1}{2}$

$$a_{k+2} = \frac{4(2ka_{k+1}-a_k+3a_{k+1})}{(2k+4)^2}$$
- Solution for $r = \frac{1}{2}$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+\frac{1}{2}}, a_{k+2} = \frac{4(2ka_{k+1}-a_k+3a_{k+1})}{(2k+4)^2}, a_1 = a_0 \right]$$

Maple trace Kovacic algorithm successful

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Reducible group (found an exponential solution)
    Group is reducible, not completely reducible
<- Kovacics algorithm successful`

```

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 57

Order:=8;

```
dsolve(4*x^2*diff(y(x),x$2)-8*x^2*diff(y(x),x)+(4*x^2+1)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \sqrt{x} \left(1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{720}x^6 + \frac{1}{5040}x^7 \right) (c_2 \ln(x) + c_1) + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 112

```
AsymptoticDSolveValue[4*x^2*y''[x]-8*x^2*y'[x]+(4*x^2+1)*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \sqrt{x} \left(\frac{x^7}{5040} + \frac{x^6}{720} + \frac{x^5}{120} + \frac{x^4}{24} + \frac{x^3}{6} + \frac{x^2}{2} + x + 1 \right) + c_2 \sqrt{x} \left(\frac{x^7}{5040} + \frac{x^6}{720} + \frac{x^5}{120} + \frac{x^4}{24} + \frac{x^3}{6} + \frac{x^2}{2} + x + 1 \right) \log(x)$$

20.3 problem 3(a)

20.3.1 Maple step by step solution 3395

Internal problem ID [6463]

Internal file name [OUTPUT/5711_Sunday_June_05_2022_03_48_16_PM_5850439/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.5. More on Regular Singular Points. Page 183

Problem number: 3(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference is integer"**

Maple gives the following as the ode type

`[_Lienard]`

$$y''x + 2y' + xy = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$y''x + 2y' + xy = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{2}{x}$$
$$q(x) = 1$$

Table 471: Table $p(x), q(x)$ singularities.

$p(x) = \frac{2}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = 1$	
singularity	type

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$y''x + 2y' + xy = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) x + 2 \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + x \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} x^{n+r-1} a_n (n+r)(n+r-1) \right) + \left(\sum_{n=0}^{\infty} 2(n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} x^{1+n+r} a_n \right) = 0 \quad (2A)$$

The next step is to make all powers of x be $n + r - 1$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r-1} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} x^{1+n+r} a_n = \sum_{n=2}^{\infty} a_{n-2} x^{n+r-1}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n + r - 1$.

$$\left(\sum_{n=0}^{\infty} x^{n+r-1} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} 2(n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=2}^{\infty} a_{n-2} x^{n+r-1} \right) = 0 \quad (2B)$$

The indicial equation is obtained from $n = 0$. From Eq (2B) this gives

$$x^{n+r-1} a_n (n+r) (n+r-1) + 2(n+r) a_n x^{n+r-1} = 0$$

When $n = 0$ the above becomes

$$x^{-1+r} a_0 r (-1+r) + 2r a_0 x^{-1+r} = 0$$

Or

$$(x^{-1+r} r (-1+r) + 2r x^{-1+r}) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$r x^{-1+r} (1+r) = 0$$

Since the above is true for all x then the indicial equation becomes

$$r(1+r) = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = 0$$

$$r_2 = -1$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$r x^{-1+r} (1+r) = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = 1$ is an integer, then we can construct two linearly independent solutions

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = C y_1(x) \ln(x) + x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y_2(x) = C y_1(x) \ln(x) + \frac{\sum_{n=0}^{\infty} b_n x^n}{x}$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y_2(x) = C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n-1} \right)$$

Where C above can be zero. We start by finding y_1 . Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

For $2 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) + 2a_n(n+r) + a_{n-2} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{a_{n-2}}{n^2 + 2nr + r^2 + n + r} \quad (4)$$

Which for the root $r = 0$ becomes

$$a_n = -\frac{a_{n-2}}{n(1+n)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = -\frac{1}{r^2 + 5r + 6}$$

Which for the root $r = 0$ becomes

$$a_2 = -\frac{1}{6}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2+5r+6}$	$-\frac{1}{6}$

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2+5r+6}$	$-\frac{1}{6}$
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{r^4 + 14r^3 + 71r^2 + 154r + 120}$$

Which for the root $r = 0$ becomes

$$a_4 = \frac{1}{120}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2+5r+6}$	$-\frac{1}{6}$
a_3	0	0
a_4	$\frac{1}{r^4+14r^3+71r^2+154r+120}$	$\frac{1}{120}$

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2+5r+6}$	$-\frac{1}{6}$
a_3	0	0
a_4	$\frac{1}{r^4+14r^3+71r^2+154r+120}$	$\frac{1}{120}$
a_5	0	0

For $n = 6$, using the above recursive equation gives

$$a_6 = -\frac{1}{(r^4 + 14r^3 + 71r^2 + 154r + 120)(r^2 + 13r + 42)}$$

Which for the root $r = 0$ becomes

$$a_6 = -\frac{1}{5040}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2+5r+6}$	$-\frac{1}{6}$
a_3	0	0
a_4	$\frac{1}{r^4+14r^3+71r^2+154r+120}$	$\frac{1}{120}$
a_5	0	0
a_6	$-\frac{1}{(r^4+14r^3+71r^2+154r+120)(r^2+13r+42)}$	$-\frac{1}{5040}$

For $n = 7$, using the above recursive equation gives

$$a_7 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2+5r+6}$	$-\frac{1}{6}$
a_3	0	0
a_4	$\frac{1}{r^4+14r^3+71r^2+154r+120}$	$\frac{1}{120}$
a_5	0	0
a_6	$-\frac{1}{(r^4+14r^3+71r^2+154r+120)(r^2+13r+42)}$	$-\frac{1}{5040}$
a_7	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots \\ &= 1 - \frac{x^2}{6} + \frac{x^4}{120} - \frac{x^6}{5040} + O(x^8) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Let

$$r_1 - r_2 = N$$

Where N is positive integer which is the difference between the two roots. r_1 is taken as the larger root. Hence for this problem we have $N = 1$. Now we need to determine if

C is zero or not. This is done by finding $\lim_{r \rightarrow r_2} a_1(r)$. If this limit exists, then $C = 0$, else we need to keep the log term and $C \neq 0$. The above table shows that

$$\begin{aligned} a_N &= a_1 \\ &= 0 \end{aligned}$$

Therefore

$$\begin{aligned} \lim_{r \rightarrow r_2} 0 &= \lim_{r \rightarrow -1} 0 \\ &= 0 \end{aligned}$$

The limit is 0. Since the limit exists then the log term is not needed and we can set $C = 0$. Therefore the second solution has the form

$$\begin{aligned} y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+r} \\ &= \sum_{n=0}^{\infty} b_n x^{n-1} \end{aligned}$$

Eq (3) derived above is used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq(3) gives

$$b_1 = 0$$

For $2 \leq n$ the recursive equation is

$$b_n(n+r)(n+r-1) + 2(n+r)b_n + b_{n-2} = 0 \quad (4)$$

Which for the root $r = -1$ becomes

$$b_n(n-1)(n-2) + 2(n-1)b_n + b_{n-2} = 0 \quad (4A)$$

Solving for b_n from the recursive equation (4) gives

$$b_n = -\frac{b_{n-2}}{n^2 + 2nr + r^2 + n + r} \quad (5)$$

Which for the root $r = -1$ becomes

$$b_n = -\frac{b_{n-2}}{n^2 - n} \quad (6)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = -1$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0

For $n = 2$, using the above recursive equation gives

$$b_2 = -\frac{1}{r^2 + 5r + 6}$$

Which for the root $r = -1$ becomes

$$b_2 = -\frac{1}{2}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{1}{r^2+5r+6}$	$-\frac{1}{2}$

For $n = 3$, using the above recursive equation gives

$$b_3 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{1}{r^2+5r+6}$	$-\frac{1}{2}$
b_3	0	0

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{1}{(r^2 + 5r + 6)(r^2 + 9r + 20)}$$

Which for the root $r = -1$ becomes

$$b_4 = \frac{1}{24}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{1}{r^2+5r+6}$	$-\frac{1}{2}$
b_3	0	0
b_4	$\frac{1}{r^4+14r^3+71r^2+154r+120}$	$\frac{1}{24}$

For $n = 5$, using the above recursive equation gives

$$b_5 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{1}{r^2+5r+6}$	$-\frac{1}{2}$
b_3	0	0
b_4	$\frac{1}{r^4+14r^3+71r^2+154r+120}$	$\frac{1}{24}$
b_5	0	0

For $n = 6$, using the above recursive equation gives

$$b_6 = -\frac{1}{(r^2 + 5r + 6)(r^2 + 9r + 20)(r^2 + 13r + 42)}$$

Which for the root $r = -1$ becomes

$$b_6 = -\frac{1}{720}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{1}{r^2+5r+6}$	$-\frac{1}{2}$
b_3	0	0
b_4	$\frac{1}{r^4+14r^3+71r^2+154r+120}$	$\frac{1}{24}$
b_5	0	0
b_6	$-\frac{1}{(r^2+5r+6)(r^2+9r+20)(r^2+13r+42)}$	$-\frac{1}{720}$

For $n = 7$, using the above recursive equation gives

$$b_7 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{1}{r^2+5r+6}$	$-\frac{1}{2}$
b_3	0	0
b_4	$\frac{1}{r^4+14r^3+71r^2+154r+120}$	$\frac{1}{24}$
b_5	0	0
b_6	$-\frac{1}{(r^2+5r+6)(r^2+9r+20)(r^2+13r+42)}$	$-\frac{1}{720}$
b_7	0	0

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned}
 y_2(x) &= 1(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 + b_7x^7 + b_8x^8 \dots) \\
 &= \frac{1 - \frac{x^2}{2} + \frac{x^4}{24} - \frac{x^6}{720} + O(x^8)}{x}
 \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 y_h(x) &= c_1y_1(x) + c_2y_2(x) \\
 &= c_1 \left(1 - \frac{x^2}{6} + \frac{x^4}{120} - \frac{x^6}{5040} + O(x^8) \right) + \frac{c_2 \left(1 - \frac{x^2}{2} + \frac{x^4}{24} - \frac{x^6}{720} + O(x^8) \right)}{x}
 \end{aligned}$$

Hence the final solution is

$$y = y_h$$

$$= c_1 \left(1 - \frac{x^2}{6} + \frac{x^4}{120} - \frac{x^6}{5040} + O(x^8) \right) + \frac{c_2 \left(1 - \frac{x^2}{2} + \frac{x^4}{24} - \frac{x^6}{720} + O(x^8) \right)}{x}$$

Summary

The solution(s) found are the following

$$y = c_1 \left(1 - \frac{x^2}{6} + \frac{x^4}{120} - \frac{x^6}{5040} + O(x^8) \right) + \frac{c_2 \left(1 - \frac{x^2}{2} + \frac{x^4}{24} - \frac{x^6}{720} + O(x^8) \right)}{x} \quad (1)$$

Verification of solutions

$$y = c_1 \left(1 - \frac{x^2}{6} + \frac{x^4}{120} - \frac{x^6}{5040} + O(x^8) \right) + \frac{c_2 \left(1 - \frac{x^2}{2} + \frac{x^4}{24} - \frac{x^6}{720} + O(x^8) \right)}{x}$$

Verified OK.

20.3.1 Maple step by step solution

Let's solve

$$y''x + 2y' + xy = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{2y'}{x} - y$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{2y'}{x} + y = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$[P_2(x) = \frac{2}{x}, P_3(x) = 1]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 2$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 0$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$y''x + 2y' + xy = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x \cdot y$ to series expansion

$$x \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+1}$$

- Shift index using $k \rightarrow k - 1$

$$x \cdot y = \sum_{k=1}^{\infty} a_{k-1} x^{k+r}$$

- Convert y' to series expansion

$$y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1}$$

- Shift index using $k \rightarrow k + 1$

$$y' = \sum_{k=-1}^{\infty} a_{k+1} (k+r+1) x^{k+r}$$

- Convert $x \cdot y''$ to series expansion

$$x \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r-1}$$

- Shift index using $k \rightarrow k + 1$

$$x \cdot y'' = \sum_{k=-1}^{\infty} a_{k+1} (k+r+1)(k+r) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0 r(1+r) x^{-1+r} + a_1 (1+r)(2+r) x^r + \left(\sum_{k=1}^{\infty} (a_{k+1} (k+r+1)(k+2+r) + a_{k-1}) x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$r(1+r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \{-1, 0\}$$

- Each term must be 0

$$a_1(1+r)(2+r) = 0$$

- Each term in the series must be 0, giving the recursion relation

$$a_{k+1}(k+r+1)(k+2+r) + a_{k-1} = 0$$

- Shift index using $k \rightarrow k+1$

$$a_{k+2}(k+2+r)(k+3+r) + a_k = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+2} = -\frac{a_k}{(k+2+r)(k+3+r)}$$

- Recursion relation for $r = -1$

$$a_{k+2} = -\frac{a_k}{(k+1)(k+2)}$$

- Solution for $r = -1$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k-1}, a_{k+2} = -\frac{a_k}{(k+1)(k+2)}, 0 = 0 \right]$$

- Recursion relation for $r = 0$

$$a_{k+2} = -\frac{a_k}{(k+2)(k+3)}$$

- Solution for $r = 0$

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+2} = -\frac{a_k}{(k+2)(k+3)}, 2a_1 = 0 \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^{k-1} \right) + \left(\sum_{k=0}^{\infty} b_k x^k \right), a_{k+2} = -\frac{a_k}{(k+1)(k+2)}, 0 = 0, b_{k+2} = -\frac{b_k}{(k+2)(k+3)}, 2b_1 = 0 \right]$$

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacics algorithm  
  A Liouvillian solution exists  
  Group is reducible or imprimitive  
<- Kovacics algorithm successful`
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 36

```
Order:=8;  
dsolve(x*diff(y(x),x$2)+2*diff(y(x),x)+x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 \left(1 - \frac{1}{6}x^2 + \frac{1}{120}x^4 - \frac{1}{5040}x^6 + O(x^8) \right) + \frac{c_2 \left(1 - \frac{1}{2}x^2 + \frac{1}{24}x^4 - \frac{1}{720}x^6 + O(x^8) \right)}{x}$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 56

```
AsymptoticDSolveValue[x*y''[x]+2*y'[x]+x*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(-\frac{x^5}{720} + \frac{x^3}{24} - \frac{x}{2} + \frac{1}{x} \right) + c_2 \left(-\frac{x^6}{5040} + \frac{x^4}{120} - \frac{x^2}{6} + 1 \right)$$

20.4 problem 3(b)

20.4.1 Maple step by step solution 3412

Internal problem ID [6464]

Internal file name [OUTPUT/5712_Sunday_June_05_2022_03_48_19_PM_62898894/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.5. More on Regular Singular Points. Page 183

Problem number: 3(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference is integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$x^2y'' - x^2y' + (x^2 - 2)y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2y'' - x^2y' + (x^2 - 2)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -1$$
$$q(x) = \frac{x^2 - 2}{x^2}$$

Table 473: Table $p(x), q(x)$ singularities.

$p(x) = -1$	
singularity	type

$q(x) = \frac{x^2-2}{x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2 y'' - x^2 y' + (x^2 - 2) y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned}
 & x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\
 & - x^2 \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (x^2 - 2) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0
 \end{aligned} \tag{1}$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{1+n+r} a_n (n+r)) \\ & + \left(\sum_{n=0}^{\infty} x^{n+r+2} a_n \right) + \sum_{n=0}^{\infty} (-2a_n x^{n+r}) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=0}^{\infty} (-x^{1+n+r} a_n (n+r)) &= \sum_{n=1}^{\infty} (-a_{n-1} (n+r-1) x^{n+r}) \\ \sum_{n=0}^{\infty} x^{n+r+2} a_n &= \sum_{n=2}^{\infty} a_{n-2} x^{n+r} \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=1}^{\infty} (-a_{n-1} (n+r-1) x^{n+r}) \\ & + \left(\sum_{n=2}^{\infty} a_{n-2} x^{n+r} \right) + \sum_{n=0}^{\infty} (-2a_n x^{n+r}) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r} a_n (n+r) (n+r-1) - 2a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$x^r a_0 r(-1+r) - 2a_0 x^r = 0$$

Or

$$(x^r r(-1+r) - 2x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(r^2 - r - 2) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$r^2 - r - 2 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 2 \\ r_2 &= -1 \end{aligned}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(r^2 - r - 2) x^r = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = 3$ is an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= C y_1(x) \ln(x) + x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= x^2 \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= C y_1(x) \ln(x) + \frac{\sum_{n=0}^{\infty} b_n x^n}{x} \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^{n+2} \\ y_2(x) &= C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n-1} \right) \end{aligned}$$

Where C above can be zero. We start by finding y_1 . Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = \frac{r}{r^2 + r - 2}$$

For $2 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) - a_{n-1}(n+r-1) + a_{n-2} - 2a_n = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{na_{n-1} + ra_{n-1} - a_{n-2} - a_{n-1}}{n^2 + 2nr + r^2 - n - r - 2} \quad (4)$$

Which for the root $r = 2$ becomes

$$a_n = \frac{na_{n-1} - a_{n-2} + a_{n-1}}{n(n+3)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 2$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{r}{r^2+r-2}$	$\frac{1}{2}$

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{2}{(r^2 + r - 2)r(r+3)}$$

Which for the root $r = 2$ becomes

$$a_2 = \frac{1}{20}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{r}{r^2+r-2}$	$\frac{1}{2}$
a_2	$\frac{2}{(r^2+r-2)r(r+3)}$	$\frac{1}{20}$

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{-r^2 - 2r + 4}{(r+4)(r^2+r-2)r(r+3)}$$

Which for the root $r = 2$ becomes

$$a_3 = -\frac{1}{60}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{r}{r^2+r-2}$	$\frac{1}{2}$
a_2	$\frac{2}{(r^2+r-2)r(r+3)}$	$\frac{1}{20}$
a_3	$\frac{-r^2-2r+4}{(r+4)(r^2+r-2)r(r+3)}$	$-\frac{1}{60}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{-r^2 - 3r + 2}{(r + 5)(r + 2)(r + 3)r(-1 + r)(r + 4)}$$

Which for the root $r = 2$ becomes

$$a_4 = -\frac{1}{210}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{r}{r^2+r-2}$	$\frac{1}{2}$
a_2	$\frac{2}{(r^2+r-2)r(r+3)}$	$\frac{1}{20}$
a_3	$\frac{-r^2-2r+4}{(r+4)(r^2+r-2)r(r+3)}$	$-\frac{1}{60}$
a_4	$\frac{-r^2-3r+2}{(r+5)(r+2)(r+3)r(-1+r)(r+4)}$	$-\frac{1}{210}$

For $n = 5$, using the above recursive equation gives

$$a_5 = -\frac{4}{(r + 6)(r + 3)r(r + 2)(-1 + r)(r + 4)(r + 5)}$$

Which for the root $r = 2$ becomes

$$a_5 = -\frac{1}{3360}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{r}{r^2+r-2}$	$\frac{1}{2}$
a_2	$\frac{2}{(r^2+r-2)r(r+3)}$	$\frac{1}{20}$
a_3	$\frac{-r^2-2r+4}{(r+4)(r^2+r-2)r(r+3)}$	$-\frac{1}{60}$
a_4	$\frac{-r^2-3r+2}{(r+5)(r+2)(r+3)r(-1+r)(r+4)}$	$-\frac{1}{210}$
a_5	$-\frac{4}{(r+6)(r+3)r(r+2)(-1+r)(r+4)(r+5)}$	$-\frac{1}{3360}$

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{r^2 + 5r - 8}{(r+7)(r+4)(r+5)(-1+r)(r+2)r(r+3)(r+6)}$$

Which for the root $r = 2$ becomes

$$a_6 = \frac{1}{20160}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{r}{r^2+r-2}$	$\frac{1}{2}$
a_2	$\frac{2}{(r^2+r-2)r(r+3)}$	$\frac{1}{20}$
a_3	$\frac{-r^2-2r+4}{(r+4)(r^2+r-2)r(r+3)}$	$-\frac{1}{60}$
a_4	$\frac{-r^2-3r+2}{(r+5)(r+2)(r+3)r(-1+r)(r+4)}$	$-\frac{1}{210}$
a_5	$-\frac{4}{(r+6)(r+3)r(r+2)(-1+r)(r+4)(r+5)}$	$-\frac{1}{3360}$
a_6	$\frac{r^2+5r-8}{(r+7)(r+4)(r+5)(-1+r)(r+2)r(r+3)(r+6)}$	$\frac{1}{20160}$

For $n = 7$, using the above recursive equation gives

$$a_7 = \frac{r^2 + 6r - 4}{(r+8)(r+5)(r+6)(r+3)r(r+2)(-1+r)(r+4)(r+7)}$$

Which for the root $r = 2$ becomes

$$a_7 = \frac{1}{100800}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{r}{r^2+r-2}$	$\frac{1}{2}$
a_2	$\frac{2}{(r^2+r-2)r(r+3)}$	$\frac{1}{20}$
a_3	$\frac{-r^2-2r+4}{(r+4)(r^2+r-2)r(r+3)}$	$-\frac{1}{60}$
a_4	$\frac{-r^2-3r+2}{(r+5)(r+2)(r+3)r(-1+r)(r+4)}$	$-\frac{1}{210}$
a_5	$-\frac{4}{(r+6)(r+3)r(r+2)(-1+r)(r+4)(r+5)}$	$-\frac{1}{3360}$
a_6	$\frac{r^2+5r-8}{(r+7)(r+4)(r+5)(-1+r)(r+2)r(r+3)(r+6)}$	$\frac{1}{20160}$
a_7	$\frac{r^2+6r-4}{(r+8)(r+5)(r+6)(r+3)r(r+2)(-1+r)(r+4)(r+7)}$	$\frac{1}{100800}$

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x^2(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots) \\ &= x^2 \left(1 + \frac{x}{2} + \frac{x^2}{20} - \frac{x^3}{60} - \frac{x^4}{210} - \frac{x^5}{3360} + \frac{x^6}{20160} + \frac{x^7}{100800} + O(x^8) \right) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Let

$$r_1 - r_2 = N$$

Where N is positive integer which is the difference between the two roots. r_1 is taken as the larger root. Hence for this problem we have $N = 3$. Now we need to determine if C is zero or not. This is done by finding $\lim_{r \rightarrow r_2} a_3(r)$. If this limit exists, then $C = 0$, else we need to keep the log term and $C \neq 0$. The above table shows that

$$\begin{aligned} a_N &= a_3 \\ &= \frac{-r^2 - 2r + 4}{(r+4)(r^2+r-2)r(r+3)} \end{aligned}$$

Therefore

$$\begin{aligned} \lim_{r \rightarrow r_2} \frac{-r^2 - 2r + 4}{(r+4)(r^2+r-2)r(r+3)} &= \lim_{r \rightarrow -1} \frac{-r^2 - 2r + 4}{(r+4)(r^2+r-2)r(r+3)} \\ &= \frac{5}{12} \end{aligned}$$

The limit is $\frac{5}{12}$. Since the limit exists then the log term is not needed and we can set $C = 0$. Therefore the second solution has the form

$$\begin{aligned} y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+r} \\ &= \sum_{n=0}^{\infty} b_n x^{n-1} \end{aligned}$$

Eq (3) derived above is used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq(3) gives

$$b_1 = \frac{r}{r^2 + r - 2}$$

For $2 \leq n$ the recursive equation is

$$b_n(n+r)(n+r-1) - b_{n-1}(n+r-1) + b_{n-2} - 2b_n = 0 \quad (4)$$

Which for the root $r = -1$ becomes

$$b_n(n-1)(n-2) - b_{n-1}(n-2) + b_{n-2} - 2b_n = 0 \quad (4A)$$

Solving for b_n from the recursive equation (4) gives

$$b_n = \frac{nb_{n-1} + rb_{n-1} - b_{n-2} - b_{n-1}}{n^2 + 2nr + r^2 - n - r - 2} \quad (5)$$

Which for the root $r = -1$ becomes

$$b_n = \frac{nb_{n-1} - b_{n-2} - 2b_{n-1}}{n^2 - 3n} \quad (6)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = -1$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{r}{r^2+r-2}$	$\frac{1}{2}$

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{2}{(r^2 + r - 2)r(r+3)}$$

Which for the root $r = -1$ becomes

$$b_2 = \frac{1}{2}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{r}{r^2+r-2}$	$\frac{1}{2}$
b_2	$\frac{2}{(r^2+r-2)r(r+3)}$	$\frac{1}{2}$

For $n = 3$, using the above recursive equation gives

$$b_3 = -\frac{r^2 + 2r - 4}{(r + 4)(r^2 + r - 2)r(r + 3)}$$

Which for the root $r = -1$ becomes

$$b_3 = \frac{5}{12}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{r}{r^2+r-2}$	$\frac{1}{2}$
b_2	$\frac{2}{(r^2+r-2)r(r+3)}$	$\frac{1}{2}$
b_3	$\frac{-r^2-2r+4}{(r+4)(r^2+r-2)r(r+3)}$	$\frac{5}{12}$

For $n = 4$, using the above recursive equation gives

$$b_4 = -\frac{r^2 + 3r - 2}{(r^2 + 7r + 10)(r + 3)r(-1 + r)(r + 4)}$$

Which for the root $r = -1$ becomes

$$b_4 = \frac{1}{12}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{r}{r^2+r-2}$	$\frac{1}{2}$
b_2	$\frac{2}{(r^2+r-2)r(r+3)}$	$\frac{1}{2}$
b_3	$\frac{-r^2-2r+4}{(r+4)(r^2+r-2)r(r+3)}$	$\frac{5}{12}$
b_4	$\frac{-r^2-3r+2}{(r+5)(r+2)(r+3)r(-1+r)(r+4)}$	$\frac{1}{12}$

For $n = 5$, using the above recursive equation gives

$$b_5 = -\frac{4}{(r^2 + 9r + 18) r (r^2 + r - 2) (r + 4) (r + 5)}$$

Which for the root $r = -1$ becomes

$$b_5 = -\frac{1}{60}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{r}{r^2+r-2}$	$\frac{1}{2}$
b_2	$\frac{2}{(r^2+r-2)r(r+3)}$	$\frac{1}{2}$
b_3	$\frac{-r^2-2r+4}{(r+4)(r^2+r-2)r(r+3)}$	$\frac{5}{12}$
b_4	$\frac{-r^2-3r+2}{(r+5)(r+2)(r+3)r(-1+r)(r+4)}$	$\frac{1}{12}$
b_5	$-\frac{4}{(r+6)(r+3)r(r+2)(-1+r)(r+4)(r+5)}$	$-\frac{1}{60}$

For $n = 6$, using the above recursive equation gives

$$b_6 = \frac{r^2 + 5r - 8}{(-1 + r) r (r + 3) (r^2 + 7r + 10) (r + 6) (r^2 + 11r + 28)}$$

Which for the root $r = -1$ becomes

$$b_6 = -\frac{1}{120}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{r}{r^2+r-2}$	$\frac{1}{2}$
b_2	$\frac{2}{(r^2+r-2)r(r+3)}$	$\frac{1}{2}$
b_3	$\frac{-r^2-2r+4}{(r+4)(r^2+r-2)r(r+3)}$	$\frac{5}{12}$
b_4	$\frac{-r^2-3r+2}{(r+5)(r+2)(r+3)r(-1+r)(r+4)}$	$\frac{1}{12}$
b_5	$-\frac{4}{(r+6)(r+3)r(r+2)(-1+r)(r+4)(r+5)}$	$-\frac{1}{60}$
b_6	$\frac{r^2+5r-8}{(r+7)(r+4)(r+5)(-1+r)(r+2)r(r+3)(r+6)}$	$-\frac{1}{120}$

For $n = 7$, using the above recursive equation gives

$$b_7 = \frac{r^2 + 6r - 4}{(r + 4)(r^2 + r - 2)r(r^2 + 9r + 18)(r + 7)(r^2 + 13r + 40)}$$

Which for the root $r = -1$ becomes

$$b_7 = -\frac{1}{1120}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{r}{r^2+r-2}$	$\frac{1}{2}$
b_2	$\frac{2}{(r^2+r-2)r(r+3)}$	$\frac{1}{2}$
b_3	$\frac{-r^2-2r+4}{(r+4)(r^2+r-2)r(r+3)}$	$\frac{5}{12}$
b_4	$\frac{-r^2-3r+2}{(r+5)(r+2)(r+3)r(-1+r)(r+4)}$	$\frac{1}{12}$
b_5	$-\frac{4}{(r+6)(r+3)r(r+2)(-1+r)(r+4)(r+5)}$	$-\frac{1}{60}$
b_6	$\frac{r^2+5r-8}{(r+7)(r+4)(r+5)(-1+r)(r+2)r(r+3)(r+6)}$	$-\frac{1}{120}$
b_7	$\frac{r^2+6r-4}{(r+8)(r+5)(r+6)(r+3)r(r+2)(-1+r)(r+4)(r+7)}$	$-\frac{1}{1120}$

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned}
 y_2(x) &= x^2(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 + b_7x^7 + b_8x^8 \dots) \\
 &= \frac{1 + \frac{x}{2} + \frac{x^2}{2} + \frac{5x^3}{12} + \frac{x^4}{12} - \frac{x^5}{60} - \frac{x^6}{120} - \frac{x^7}{1120} + O(x^8)}{x}
 \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 y_h(x) &= c_1 y_1(x) + c_2 y_2(x) \\
 &= c_1 x^2 \left(1 + \frac{x}{2} + \frac{x^2}{20} - \frac{x^3}{60} - \frac{x^4}{210} - \frac{x^5}{3360} + \frac{x^6}{20160} + \frac{x^7}{100800} + O(x^8) \right) \\
 &\quad + \frac{c_2 \left(1 + \frac{x}{2} + \frac{x^2}{2} + \frac{5x^3}{12} + \frac{x^4}{12} - \frac{x^5}{60} - \frac{x^6}{120} - \frac{x^7}{1120} + O(x^8) \right)}{x}
 \end{aligned}$$

Hence the final solution is

$$\begin{aligned}
 y &= y_h \\
 &= c_1 x^2 \left(1 + \frac{x}{2} + \frac{x^2}{20} - \frac{x^3}{60} - \frac{x^4}{210} - \frac{x^5}{3360} + \frac{x^6}{20160} + \frac{x^7}{100800} + O(x^8) \right) \\
 &\quad + \frac{c_2 \left(1 + \frac{x}{2} + \frac{x^2}{2} + \frac{5x^3}{12} + \frac{x^4}{12} - \frac{x^5}{60} - \frac{x^6}{120} - \frac{x^7}{1120} + O(x^8) \right)}{x}
 \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}
 y &= c_1 x^2 \left(1 + \frac{x}{2} + \frac{x^2}{20} - \frac{x^3}{60} - \frac{x^4}{210} - \frac{x^5}{3360} + \frac{x^6}{20160} + \frac{x^7}{100800} + O(x^8) \right) \\
 &\quad + \frac{c_2 \left(1 + \frac{x}{2} + \frac{x^2}{2} + \frac{5x^3}{12} + \frac{x^4}{12} - \frac{x^5}{60} - \frac{x^6}{120} - \frac{x^7}{1120} + O(x^8) \right)}{x}
 \end{aligned} \tag{1}$$

Verification of solutions

$$\begin{aligned}
 y &= c_1 x^2 \left(1 + \frac{x}{2} + \frac{x^2}{20} - \frac{x^3}{60} - \frac{x^4}{210} - \frac{x^5}{3360} + \frac{x^6}{20160} + \frac{x^7}{100800} + O(x^8) \right) \\
 &\quad + \frac{c_2 \left(1 + \frac{x}{2} + \frac{x^2}{2} + \frac{5x^3}{12} + \frac{x^4}{12} - \frac{x^5}{60} - \frac{x^6}{120} - \frac{x^7}{1120} + O(x^8) \right)}{x}
 \end{aligned}$$

Verified OK.

20.4.1 Maple step by step solution

Let's solve

$$x^2 y'' - x^2 y' + (x^2 - 2)y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = y' - \frac{(x^2-2)y}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - y' + \frac{(x^2-2)y}{x^2} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$\left[P_2(x) = -1, P_3(x) = \frac{x^2-2}{x^2} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 0$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = -2$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$x^2 y'' - x^2 y' + (x^2 - 2)y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 0..2$

$$x^m \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

- Shift index using $k \rightarrow k - m$

$$x^m \cdot y = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

- Convert $x^2 \cdot y'$ to series expansion

$$x^2 \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r+1}$$

- Shift index using $k \rightarrow k - 1$

$$x^2 \cdot y' = \sum_{k=1}^{\infty} a_{k-1} (k-1+r) x^{k+r}$$

- Convert $x^2 \cdot y''$ to series expansion

$$x^2 \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k-1+r) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(1+r)(-2+r)x^r + (a_1(2+r)(-1+r) - a_0r)x^{1+r} + \left(\sum_{k=2}^{\infty} (a_k(k+r+1)(k+r-2) - a_{k-1}k - a_{k-1}r + a_{k-2} + a_{k-1}) \right) x^{k+r} = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$(1+r)(-2+r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \{-1, 2\}$$

- Each term must be 0

$$a_1(2+r)(-1+r) - a_0r = 0$$

- Solve for the dependent coefficient(s)

$$a_1 = \frac{a_0r}{r^2+r-2}$$

- Each term in the series must be 0, giving the recursion relation

$$a_k(k+r+1)(k+r-2) - a_{k-1}k - a_{k-1}r + a_{k-2} + a_{k-1} = 0$$

- Shift index using $k \rightarrow k + 2$

$$a_{k+2}(k+3+r)(k+r) - a_{k+1}(k+2) - a_{k+1}r + a_k + a_{k+1} = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+2} = \frac{ka_{k+1} + a_{k+1}r - a_k + a_{k+1}}{(k+3+r)(k+r)}$$

- Recursion relation for $r = -1$

$$a_{k+2} = \frac{ka_{k+1} - a_k}{(k+2)(k-1)}$$

- Series not valid for $r = -1$, division by 0 in the recursion relation at $k = 1$

$$a_{k+2} = \frac{ka_{k+1} - a_k}{(k+2)(k-1)}$$

- Recursion relation for $r = 2$

$$a_{k+2} = \frac{ka_{k+1} - a_k + 3a_{k+1}}{(k+5)(k+2)}$$

- Solution for $r = 2$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+2}, a_{k+2} = \frac{ka_{k+1} - a_k + 3a_{k+1}}{(k+5)(k+2)}, a_1 = \frac{a_0}{2} \right]$$

Maple trace Kovacic algorithm successful

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Group is reducible or imprimitive
<- Kovacics algorithm successful`

```

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 55

```

Order:=8;
dsolve(x^2*diff(y(x),x$2)-x^2*diff(y(x),x)+(x^2-2)*y(x)=0,y(x),type='series',x=0);

```

$$y(x) = c_1 x^2 \left(1 + \frac{1}{2}x + \frac{1}{20}x^2 - \frac{1}{60}x^3 - \frac{1}{210}x^4 - \frac{1}{3360}x^5 + \frac{1}{20160}x^6 + \frac{1}{100800}x^7 + O(x^8) \right) \\ + \frac{c_2 (12 + 6x + 6x^2 + 5x^3 + x^4 - \frac{1}{5}x^5 - \frac{1}{10}x^6 - \frac{3}{280}x^7 + O(x^8))}{x}$$

✓ Solution by Mathematica

Time used: 0.069 (sec). Leaf size: 96

```
AsymptoticDSolveValue[x^2*y''[x]-x^2*y'[x]+(x^2-2)*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(-\frac{x^5}{120} - \frac{x^4}{60} + \frac{x^3}{12} + \frac{5x^2}{12} + \frac{x}{2} + \frac{1}{x} + \frac{1}{2} \right) \\ + c_2 \left(\frac{x^8}{20160} - \frac{x^7}{3360} - \frac{x^6}{210} - \frac{x^5}{60} + \frac{x^4}{20} + \frac{x^3}{2} + x^2 \right)$$

20.5 problem 3(c)

20.5.1 Maple step by step solution 3426

Internal problem ID [6465]

Internal file name [OUTPUT/5713_Sunday_June_05_2022_03_48_23_PM_95481906/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.5. More on Regular Singular Points. Page 183

Problem number: 3(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference is integer"**

Maple gives the following as the ode type

```
[[_Emden, _Fowler], [_2nd_order, _linear, `_with_symmetry_[0,F(x)]`]]
```

$$y''x - y' + 4yx^3 = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$y''x - y' + 4yx^3 = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{1}{x}$$
$$q(x) = 4x^2$$

Table 475: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{1}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = 4x^2$	
singularity	type
$x = \infty$	“regular”
$x = -\infty$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0, \infty, -\infty]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$y''x - y' + 4yx^3 = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) x - \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + 4 \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) x^3 = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} x^{n+r-1} a_n (n+r)(n+r-1) \right) + \sum_{n=0}^{\infty} (-n-r) a_n x^{n+r-1} + \left(\sum_{n=0}^{\infty} 4x^{3+n+r} a_n \right) = 0 \quad (2A)$$

The next step is to make all powers of x be $n + r - 1$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r-1} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} 4x^{3+n+r} a_n = \sum_{n=4}^{\infty} 4a_{n-4} x^{n+r-1}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n + r - 1$.

$$\left(\sum_{n=0}^{\infty} x^{n+r-1} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-(n+r) a_n x^{n+r-1}) + \left(\sum_{n=4}^{\infty} 4a_{n-4} x^{n+r-1} \right) = 0 \quad (2B)$$

The indicial equation is obtained from $n = 0$. From Eq (2B) this gives

$$x^{n+r-1} a_n (n+r) (n+r-1) - (n+r) a_n x^{n+r-1} = 0$$

When $n = 0$ the above becomes

$$x^{-1+r} a_0 r (-1+r) - r a_0 x^{-1+r} = 0$$

Or

$$(x^{-1+r} r (-1+r) - r x^{-1+r}) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$r x^{-1+r} (-2+r) = 0$$

Since the above is true for all x then the indicial equation becomes

$$r(-2+r) = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = 2$$

$$r_2 = 0$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$r x^{-1+r} (-2+r) = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = 2$ is an integer, then we can construct two linearly independent solutions

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = C y_1(x) \ln(x) + x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = x^2 \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+2}$$

$$y_2(x) = C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Where C above can be zero. We start by finding y_1 . Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

Substituting $n = 2$ in Eq. (2B) gives

$$a_2 = 0$$

Substituting $n = 3$ in Eq. (2B) gives

$$a_3 = 0$$

For $4 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) - a_n(n+r) + 4a_{n-4} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{4a_{n-4}}{n^2 + 2nr + r^2 - 2n - 2r} \quad (4)$$

Which for the root $r = 2$ becomes

$$a_n = -\frac{4a_{n-4}}{n(n+2)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 2$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = -\frac{4}{r^2 + 6r + 8}$$

Which for the root $r = 2$ becomes

$$a_4 = -\frac{1}{6}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	0	0
a_4	$-\frac{4}{r^2+6r+8}$	$-\frac{1}{6}$

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	0	0
a_4	$-\frac{4}{r^2+6r+8}$	$-\frac{1}{6}$
a_5	0	0

For $n = 6$, using the above recursive equation gives

$$a_6 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	0	0
a_4	$-\frac{4}{r^2+6r+8}$	$-\frac{1}{6}$
a_5	0	0
a_6	0	0

For $n = 7$, using the above recursive equation gives

$$a_7 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	0	0
a_4	$-\frac{4}{r^2+6r+8}$	$-\frac{1}{6}$
a_5	0	0
a_6	0	0
a_7	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned}
 y_1(x) &= x^2(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots) \\
 &= x^2\left(1 - \frac{x^4}{6} + O(x^8)\right)
 \end{aligned}$$

Now the second solution $y_2(x)$ is found. Let

$$r_1 - r_2 = N$$

Where N is positive integer which is the difference between the two roots. r_1 is taken as the larger root. Hence for this problem we have $N = 2$. Now we need to determine if C is zero or not. This is done by finding $\lim_{r \rightarrow r_2} a_2(r)$. If this limit exists, then $C = 0$, else we need to keep the log term and $C \neq 0$. The above table shows that

$$\begin{aligned}
 a_N &= a_2 \\
 &= 0
 \end{aligned}$$

Therefore

$$\begin{aligned}
 \lim_{r \rightarrow r_2} 0 &= \lim_{r \rightarrow 0} 0 \\
 &= 0
 \end{aligned}$$

The limit is 0. Since the limit exists then the log term is not needed and we can set $C = 0$. Therefore the second solution has the form

$$\begin{aligned}
 y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+r} \\
 &= \sum_{n=0}^{\infty} b_n x^n
 \end{aligned}$$

Eq (3) derived above is used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq(3) gives

$$b_1 = 0$$

Substituting $n = 2$ in Eq(3) gives

$$b_2 = 0$$

Substituting $n = 3$ in Eq(3) gives

$$b_3 = 0$$

For $4 \leq n$ the recursive equation is

$$b_n(n+r)(n+r-1) - (n+r)b_n + 4b_{n-4} = 0 \quad (4)$$

Which for the root $r = 0$ becomes

$$b_n n(n-1) - nb_n + 4b_{n-4} = 0 \quad (4A)$$

Solving for b_n from the recursive equation (4) gives

$$b_n = -\frac{4b_{n-4}}{n^2 + 2nr + r^2 - 2n - 2r} \quad (5)$$

Which for the root $r = 0$ becomes

$$b_n = -\frac{4b_{n-4}}{n^2 - 2n} \quad (6)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	0	0
b_3	0	0

For $n = 4$, using the above recursive equation gives

$$b_4 = -\frac{4}{r^2 + 6r + 8}$$

Which for the root $r = 0$ becomes

$$b_4 = -\frac{1}{2}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	0	0
b_3	0	0
b_4	$-\frac{4}{r^2+6r+8}$	$-\frac{1}{2}$

For $n = 5$, using the above recursive equation gives

$$b_5 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	0	0
b_3	0	0
b_4	$-\frac{4}{r^2+6r+8}$	$-\frac{1}{2}$
b_5	0	0

For $n = 6$, using the above recursive equation gives

$$b_6 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	0	0
b_3	0	0
b_4	$-\frac{4}{r^2+6r+8}$	$-\frac{1}{2}$
b_5	0	0
b_6	0	0

For $n = 7$, using the above recursive equation gives

$$b_7 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	0	0
b_3	0	0
b_4	$-\frac{4}{r^2+6r+8}$	$-\frac{1}{2}$
b_5	0	0
b_6	0	0
b_7	0	0

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned} y_2(x) &= b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 + b_7x^7 + b_8x^8 \dots \\ &= 1 - \frac{x^4}{2} + O(x^8) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= c_1x^2 \left(1 - \frac{x^4}{6} + O(x^8) \right) + c_2 \left(1 - \frac{x^4}{2} + O(x^8) \right) \end{aligned}$$

Hence the final solution is

$$\begin{aligned}y &= y_h \\ &= c_1 x^2 \left(1 - \frac{x^4}{6} + O(x^8) \right) + c_2 \left(1 - \frac{x^4}{2} + O(x^8) \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 x^2 \left(1 - \frac{x^4}{6} + O(x^8) \right) + c_2 \left(1 - \frac{x^4}{2} + O(x^8) \right) \quad (1)$$

Verification of solutions

$$y = c_1 x^2 \left(1 - \frac{x^4}{6} + O(x^8) \right) + c_2 \left(1 - \frac{x^4}{2} + O(x^8) \right)$$

Verified OK.

20.5.1 Maple step by step solution

Let's solve

$$y''x - y' + 4yx^3 = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = \frac{y'}{x} - 4yx^2$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - \frac{y'}{x} + 4yx^2 = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$[P_2(x) = -\frac{1}{x}, P_3(x) = 4x^2]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = -1$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 0$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$y''x - y' + 4yx^3 = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^3 \cdot y$ to series expansion

$$x^3 \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+3}$$

- Shift index using $k \rightarrow k - 3$

$$x^3 \cdot y = \sum_{k=3}^{\infty} a_{k-3} x^{k+r}$$

- Convert y' to series expansion

$$y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1}$$

- Shift index using $k \rightarrow k + 1$

$$y' = \sum_{k=-1}^{\infty} a_{k+1} (k+1+r) x^{k+r}$$

- Convert $x \cdot y''$ to series expansion

$$x \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r) (k+r-1) x^{k+r-1}$$

- Shift index using $k \rightarrow k + 1$

$$x \cdot y'' = \sum_{k=-1}^{\infty} a_{k+1} (k+1+r) (k+r) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0 r(-2+r) x^{-1+r} + a_1 (1+r) (-1+r) x^r + a_2 (2+r) r x^{1+r} + a_3 (3+r) (1+r) x^{2+r} + \left(\sum_{k=3}^{\infty} (a_k \right.$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$r(-2 + r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \{0, 2\}$$

- The coefficients of each power of x must be 0

$$[a_1(1 + r)(-1 + r) = 0, a_2(2 + r)r = 0, a_3(3 + r)(1 + r) = 0]$$

- Solve for the dependent coefficient(s)

$$\{a_1 = 0, a_2 = 0, a_3 = 0\}$$

- Each term in the series must be 0, giving the recursion relation

$$a_{k+1}(k + 1 + r)(k + r - 1) + 4a_{k-3} = 0$$

- Shift index using $k \rightarrow k + 3$

$$a_{k+4}(k + 4 + r)(k + 2 + r) + 4a_k = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+4} = -\frac{4a_k}{(k+4+r)(k+2+r)}$$

- Recursion relation for $r = 0$

$$a_{k+4} = -\frac{4a_k}{(k+4)(k+2)}$$

- Solution for $r = 0$

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+4} = -\frac{4a_k}{(k+4)(k+2)}, a_1 = 0, a_2 = 0, a_3 = 0 \right]$$

- Recursion relation for $r = 2$

$$a_{k+4} = -\frac{4a_k}{(k+6)(k+4)}$$

- Solution for $r = 2$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+2}, a_{k+4} = -\frac{4a_k}{(k+6)(k+4)}, a_1 = 0, a_2 = 0, a_3 = 0 \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^k \right) + \left(\sum_{k=0}^{\infty} b_k x^{k+2} \right), a_{k+4} = -\frac{4a_k}{(k+4)(k+2)}, a_1 = 0, a_2 = 0, a_3 = 0, b_{k+4} = -\frac{4b_k}{(k+6)(k+4)} \right]$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
<- linear_1 successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 28

```
Order:=8;  
dsolve(x*diff(y(x),x$2)-diff(y(x),x)+4*x^3*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^2 \left(1 - \frac{1}{6} x^4 + O(x^8) \right) + c_2 (-2 + x^4 + O(x^8))$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 30

```
AsymptoticDSolveValue[x*y'[x]-y'[x]+4*x^3*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(1 - \frac{x^4}{2} \right) + c_2 \left(x^2 - \frac{x^6}{6} \right)$$

20.6 problem 4

20.6.1 Maple step by step solution 3440

Internal problem ID [6466]

Internal file name [OUTPUT/5714_Sunday_June_05_2022_03_48_25_PM_35635026/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.5. More on Regular Singular Points. Page 183

Problem number: 4.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$(x - 1)^2 y'' - 3(x - 1) y' + 2y = 0$$

With the expansion point for the power series method at $x = 1$.

The ode does not have its expansion point at $x = 0$, therefore to simplify the computation of power series expansion, change of variable is made on the independent variable to shift the initial conditions and the expansion point back to zero. The new ode is then solved more easily since the expansion point is now at zero. The solution converted back to the original independent variable. Let

$$t = x - 1$$

The ode is converted to be in terms of the new independent variable t . This results in

$$\left(\frac{d^2}{dt^2}y(t)\right) t^2 - 3t\left(\frac{d}{dt}y(t)\right) + 2y(t) = 0$$

With its expansion point and initial conditions now at $t = 0$. The transformed ODE is now solved. The type of the expansion point is first determined. This is done on the

homogeneous part of the ODE.

$$\left(\frac{d^2}{dt^2}y(t)\right)t^2 - 3t\left(\frac{d}{dt}y(t)\right) + 2y(t) = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$\frac{d^2}{dt^2}y(t) + p(t)\frac{d}{dt}y(t) + q(t)y(t) = 0$$

Where

$$p(t) = -\frac{3}{t}$$

$$q(t) = \frac{2}{t^2}$$

Table 477: Table $p(t), q(t)$ singularities.

$p(t) = -\frac{3}{t}$		$q(t) = \frac{2}{t^2}$	
singularity	type	singularity	type
$t = 0$	“regular”	$t = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0, \infty]$

Irregular singular points : $[\]$

Since $t = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$\left(\frac{d^2}{dt^2}y(t)\right)t^2 - 3t\left(\frac{d}{dt}y(t)\right) + 2y(t) = 0$$

Let the solution be represented as Frobenius power series of the form

$$y(t) = \sum_{n=0}^{\infty} a_n t^{n+r}$$

Then

$$\frac{d}{dt}y(t) = \sum_{n=0}^{\infty} (n+r) a_n t^{n+r-1}$$

$$\frac{d^2}{dt^2}y(t) = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n t^{n+r-2}$$

Substituting the above back into the ode gives

$$\left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n t^{n+r-2} \right) t^2 - 3t \left(\sum_{n=0}^{\infty} (n+r) a_n t^{n+r-1} \right) + 2 \left(\sum_{n=0}^{\infty} a_n t^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} t^{n+r} a_n (n+r)(n+r-1) \right) + \sum_{n=0}^{\infty} (-3t^{n+r} a_n (n+r)) + \left(\sum_{n=0}^{\infty} 2a_n t^{n+r} \right) = 0 \quad (2A)$$

The next step is to make all powers of t be $n+r$ in each summation term. Going over each summation term above with power of t in it which is not already t^{n+r} and adjusting the power and the corresponding index gives Substituting all the above in Eq (2A) gives the following equation where now all powers of t are the same and equal to $n+r$.

$$\left(\sum_{n=0}^{\infty} t^{n+r} a_n (n+r)(n+r-1) \right) + \sum_{n=0}^{\infty} (-3t^{n+r} a_n (n+r)) + \left(\sum_{n=0}^{\infty} 2a_n t^{n+r} \right) = 0 \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$t^{n+r} a_n (n+r)(n+r-1) - 3t^{n+r} a_n (n+r) + 2a_n t^{n+r} = 0$$

When $n=0$ the above becomes

$$t^r a_0 r(-1+r) - 3t^r a_0 r + 2a_0 t^r = 0$$

Or

$$(t^r r(-1+r) - 3t^r r + 2t^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(r^2 - 4r + 2) t^r = 0$$

Since the above is true for all t then the indicial equation becomes

$$r^2 - 4r + 2 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = 2 + \sqrt{2}$$

$$r_2 = 2 - \sqrt{2}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(r^2 - 4r + 2)t^r = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = 2\sqrt{2}$ is not an integer, then we can construct two linearly independent solutions

$$y_1(t) = t^{r_1} \left(\sum_{n=0}^{\infty} a_n t^n \right)$$

$$y_2(t) = t^{r_2} \left(\sum_{n=0}^{\infty} b_n t^n \right)$$

Or

$$y_1(t) = \sum_{n=0}^{\infty} a_n t^{n+2+\sqrt{2}}$$

$$y_2(t) = \sum_{n=0}^{\infty} b_n t^{n+2-\sqrt{2}}$$

We start by finding $y_1(t)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $0 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) - 3a_n(n+r) + 2a_n = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = 0 \quad (4)$$

Which for the root $r = 2 + \sqrt{2}$ becomes

$$a_n = 0 \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 2 + \sqrt{2}$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	0	0
a_4	0	0

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	0	0
a_4	0	0
a_5	0	0

For $n = 6$, using the above recursive equation gives

$$a_6 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	0	0
a_4	0	0
a_5	0	0
a_6	0	0

For $n = 7$, using the above recursive equation gives

$$a_7 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	0	0
a_4	0	0
a_5	0	0
a_6	0	0
a_7	0	0

Using the above table, then the solution $y_1(t)$ is

$$\begin{aligned} y_1(t) &= t^{2+\sqrt{2}}(a_0 + a_1t + a_2t^2 + a_3t^3 + a_4t^4 + a_5t^5 + a_6t^6 + a_7t^7 + a_8t^8 \dots) \\ &= t^{2+\sqrt{2}}(1 + O(t^8)) \end{aligned}$$

Now the second solution $y_2(t)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. For $0 \leq n$ the recursive equation is

$$b_n(n+r)(n+r-1) - 3b_n(n+r) + 2b_n = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = 0 \quad (4)$$

Which for the root $r = 2 - \sqrt{2}$ becomes

$$b_n = 0 \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = 2 - \sqrt{2}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1

For $n = 1$, using the above recursive equation gives

$$b_1 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0

For $n = 2$, using the above recursive equation gives

$$b_2 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	0	0

For $n = 3$, using the above recursive equation gives

$$b_3 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	0	0
b_3	0	0

For $n = 4$, using the above recursive equation gives

$$b_4 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	0	0
b_3	0	0
b_4	0	0

For $n = 5$, using the above recursive equation gives

$$b_5 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	0	0
b_3	0	0
b_4	0	0
b_5	0	0

For $n = 6$, using the above recursive equation gives

$$b_6 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	0	0
b_3	0	0
b_4	0	0
b_5	0	0
b_6	0	0

For $n = 7$, using the above recursive equation gives

$$b_7 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	0	0
b_3	0	0
b_4	0	0
b_5	0	0
b_6	0	0
b_7	0	0

Using the above table, then the solution $y_2(t)$ is

$$\begin{aligned} y_2(t) &= t^{2+\sqrt{2}}(b_0 + b_1t + b_2t^2 + b_3t^3 + b_4t^4 + b_5t^5 + b_6t^6 + b_7t^7 + b_8t^8 \dots) \\ &= t^{2+\sqrt{2}}(1 + O(t^8)) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(t) &= c_1y_1(t) + c_2y_2(t) \\ &= c_1t^{2+\sqrt{2}}(1 + O(t^8)) + c_2t^{2-\sqrt{2}}(1 + O(t^8)) \end{aligned}$$

Hence the final solution is

$$\begin{aligned} y(t) &= y_h \\ &= c_1t^{2+\sqrt{2}}(1 + O(t^8)) + c_2t^{2-\sqrt{2}}(1 + O(t^8)) \end{aligned}$$

Replacing t in the above with the original independent variable x using $t = x - 1$ results in

$$y = c_1(x - 1)^{2+\sqrt{2}}(1 + O((x - 1)^8)) + c_2(x - 1)^{2-\sqrt{2}}(1 + O((x - 1)^8))$$

Summary

The solution(s) found are the following

$$y = c_1(x-1)^{2+\sqrt{2}}(1 + O((x-1)^8)) + c_2(x-1)^{2-\sqrt{2}}(1 + O((x-1)^8)) \quad (1)$$

Verification of solutions

$$y = c_1(x-1)^{2+\sqrt{2}}(1 + O((x-1)^8)) + c_2(x-1)^{2-\sqrt{2}}(1 + O((x-1)^8))$$

Verified OK.

20.6.1 Maple step by step solution

Let's solve

$$(x-1)^2 y'' + (3-3x)y' + 2y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = \frac{3y'}{x-1} - \frac{2y}{(x-1)^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - \frac{3y'}{x-1} + \frac{2y}{(x-1)^2} = 0$$

- Check to see if $x_0 = 1$ is a regular singular point

- Define functions

$$\left[P_2(x) = -\frac{3}{x-1}, P_3(x) = \frac{2}{(x-1)^2} \right]$$

- $(x-1) \cdot P_2(x)$ is analytic at $x = 1$

$$\left. ((x-1) \cdot P_2(x)) \right|_{x=1} = -3$$

- $(x-1)^2 \cdot P_3(x)$ is analytic at $x = 1$

$$\left. ((x-1)^2 \cdot P_3(x)) \right|_{x=1} = 2$$

- $x = 1$ is a regular singular point

Check to see if $x_0 = 1$ is a regular singular point

$$x_0 = 1$$

- Multiply by denominators

$$(x - 1)^2 y'' + (3 - 3x) y' + 2y = 0$$

- Change variables using $x = u + 1$ so that the regular singular point is at $u = 0$

$$u^2 \left(\frac{d^2}{du^2} y(u) \right) - 3u \left(\frac{d}{du} y(u) \right) + 2y(u) = 0$$

- Assume series solution for $y(u)$

$$y(u) = \sum_{k=0}^{\infty} a_k u^{k+r}$$

- Rewrite DE with series expansions

- Convert $u \cdot \left(\frac{d}{du} y(u) \right)$ to series expansion

$$u \cdot \left(\frac{d}{du} y(u) \right) = \sum_{k=0}^{\infty} a_k (k + r) u^{k+r}$$

- Convert $u^2 \cdot \left(\frac{d^2}{du^2} y(u) \right)$ to series expansion

$$u^2 \cdot \left(\frac{d^2}{du^2} y(u) \right) = \sum_{k=0}^{\infty} a_k (k + r) (k + r - 1) u^{k+r}$$

Rewrite DE with series expansions

$$\sum_{k=0}^{\infty} a_k (k^2 + 2kr + r^2 - 4k - 4r + 2) u^{k+r} = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$r = 0$$

- Each term in the series must be 0, giving the recursion relation

$$a_k (k^2 - 4k + 2) = 0$$

- Recursion relation that defines series solution to ODE

$$a_k = 0$$

- Recursion relation for $r = 0$

$$a_k = 0$$

- Solution for $r = 0$

$$\left[y(u) = \sum_{k=0}^{\infty} a_k u^k, a_k = 0 \right]$$

- Revert the change of variables $u = x - 1$

$$\left[y = \sum_{k=0}^{\infty} a_k (x - 1)^k, a_k = 0 \right]$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
<- LODE of Euler type successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 41

```
Order:=8;  
dsolve((x-1)^2*diff(y(x),x$2)-3*(x-1)*diff(y(x),x)+2*y(x)=0,y(x),type='series',x=1);
```

$$y(x) = (x - 1)^2 \left(c_1(x - 1)^{-\sqrt{2}} + c_2(x - 1)^{\sqrt{2}} \right) + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 34

```
AsymptoticDSolveValue[(x-1)^2*y''[x]-3*(x-1)*y'[x]+2*y[x]==0,y[x],{x,1,7}]
```

$$y(x) \rightarrow c_1(x - 1)^{2+\sqrt{2}} + c_2(x - 1)^{2-\sqrt{2}}$$

20.7 problem 5

20.7.1 Maple step by step solution 3453

Internal problem ID [6467]

Internal file name [OUTPUT/5715_Sunday_June_05_2022_03_48_28_PM_75065673/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.5. More on Regular Singular Points. Page 183

Problem number: 5.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$3(1+x)^2 y'' - (1+x)y' - y = 0$$

With the expansion point for the power series method at $x = -1$.

The ode does not have its expansion point at $x = 0$, therefore to simplify the computation of power series expansion, change of variable is made on the independent variable to shift the initial conditions and the expansion point back to zero. The new ode is then solved more easily since the expansion point is now at zero. The solution converted back to the original independent variable. Let

$$t = 1 + x$$

The ode is converted to be in terms of the new independent variable t . This results in

$$3\left(\frac{d^2}{dt^2}y(t)\right)t^2 - t\left(\frac{d}{dt}y(t)\right) - y(t) = 0$$

With its expansion point and initial conditions now at $t = 0$. The transformed ODE is now solved. The type of the expansion point is first determined. This is done on the

homogeneous part of the ODE.

$$3\left(\frac{d^2}{dt^2}y(t)\right)t^2 - t\left(\frac{d}{dt}y(t)\right) - y(t) = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$\frac{d^2}{dt^2}y(t) + p(t)\frac{d}{dt}y(t) + q(t)y(t) = 0$$

Where

$$p(t) = -\frac{1}{3t}$$

$$q(t) = -\frac{1}{3t^2}$$

Table 479: Table $p(t), q(t)$ singularities.

$p(t) = -\frac{1}{3t}$	
singularity	type
$t = 0$	“regular”

$q(t) = -\frac{1}{3t^2}$	
singularity	type
$t = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0, \infty]$

Irregular singular points : $[\]$

Since $t = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$3\left(\frac{d^2}{dt^2}y(t)\right)t^2 - t\left(\frac{d}{dt}y(t)\right) - y(t) = 0$$

Let the solution be represented as Frobenius power series of the form

$$y(t) = \sum_{n=0}^{\infty} a_n t^{n+r}$$

Then

$$\frac{d}{dt}y(t) = \sum_{n=0}^{\infty} (n+r) a_n t^{n+r-1}$$

$$\frac{d^2}{dt^2}y(t) = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n t^{n+r-2}$$

Substituting the above back into the ode gives

$$3 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n t^{n+r-2} \right) t^2 - t \left(\sum_{n=0}^{\infty} (n+r) a_n t^{n+r-1} \right) - \left(\sum_{n=0}^{\infty} a_n t^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} 3t^{n+r} a_n (n+r)(n+r-1) \right) + \sum_{n=0}^{\infty} (-t^{n+r} a_n (n+r)) + \sum_{n=0}^{\infty} (-a_n t^{n+r}) = 0 \quad (2A)$$

The next step is to make all powers of t be $n+r$ in each summation term. Going over each summation term above with power of t in it which is not already t^{n+r} and adjusting the power and the corresponding index gives Substituting all the above in Eq (2A) gives the following equation where now all powers of t are the same and equal to $n+r$.

$$\left(\sum_{n=0}^{\infty} 3t^{n+r} a_n (n+r)(n+r-1) \right) + \sum_{n=0}^{\infty} (-t^{n+r} a_n (n+r)) + \sum_{n=0}^{\infty} (-a_n t^{n+r}) = 0 \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$3t^{n+r} a_n (n+r)(n+r-1) - t^{n+r} a_n (n+r) - a_n t^{n+r} = 0$$

When $n=0$ the above becomes

$$3t^r a_0 r(-1+r) - t^r a_0 r - a_0 t^r = 0$$

Or

$$(3t^r r(-1+r) - t^r r - t^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(3r^2 - 4r - 1) t^r = 0$$

Since the above is true for all t then the indicial equation becomes

$$3r^2 - 4r - 1 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = \frac{2}{3} + \frac{\sqrt{7}}{3}$$

$$r_2 = \frac{2}{3} - \frac{\sqrt{7}}{3}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(3r^2 - 4r - 1)t^r = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = \frac{2\sqrt{7}}{3}$ is not an integer, then we can construct two linearly independent solutions

$$y_1(t) = t^{r_1} \left(\sum_{n=0}^{\infty} a_n t^n \right)$$

$$y_2(t) = t^{r_2} \left(\sum_{n=0}^{\infty} b_n t^n \right)$$

Or

$$y_1(t) = \sum_{n=0}^{\infty} a_n t^{n + \frac{2}{3} + \frac{\sqrt{7}}{3}}$$

$$y_2(t) = \sum_{n=0}^{\infty} b_n t^{n + \frac{2}{3} - \frac{\sqrt{7}}{3}}$$

We start by finding $y_1(t)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $0 \leq n$ the recursive equation is

$$3a_n(n+r)(n+r-1) - a_n(n+r) - a_n = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = 0 \quad (4)$$

Which for the root $r = \frac{2}{3} + \frac{\sqrt{7}}{3}$ becomes

$$a_n = 0 \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = \frac{2}{3} + \frac{\sqrt{7}}{3}$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	0	0
a_4	0	0

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	0	0
a_4	0	0
a_5	0	0

For $n = 6$, using the above recursive equation gives

$$a_6 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	0	0
a_4	0	0
a_5	0	0
a_6	0	0

For $n = 7$, using the above recursive equation gives

$$a_7 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	0	0
a_3	0	0
a_4	0	0
a_5	0	0
a_6	0	0
a_7	0	0

Using the above table, then the solution $y_1(t)$ is

$$\begin{aligned} y_1(t) &= t^{\frac{2}{3} + \frac{\sqrt{7}}{3}} (a_0 + a_1 t + a_2 t^2 + a_3 t^3 + a_4 t^4 + a_5 t^5 + a_6 t^6 + a_7 t^7 + a_8 t^8 \dots) \\ &= t^{\frac{2}{3} + \frac{\sqrt{7}}{3}} (1 + O(t^8)) \end{aligned}$$

Now the second solution $y_2(t)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. For $0 \leq n$ the recursive equation is

$$3b_n(n+r)(n+r-1) - b_n(n+r) - b_n = 0 \tag{3}$$

Solving for b_n from recursive equation (4) gives

$$b_n = 0 \tag{4}$$

Which for the root $r = \frac{2}{3} - \frac{\sqrt{7}}{3}$ becomes

$$b_n = 0 \tag{5}$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = \frac{2}{3} - \frac{\sqrt{7}}{3}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1

For $n = 1$, using the above recursive equation gives

$$b_1 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0

For $n = 2$, using the above recursive equation gives

$$b_2 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	0	0

For $n = 3$, using the above recursive equation gives

$$b_3 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	0	0
b_3	0	0

For $n = 4$, using the above recursive equation gives

$$b_4 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	0	0
b_3	0	0
b_4	0	0

For $n = 5$, using the above recursive equation gives

$$b_5 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	0	0
b_3	0	0
b_4	0	0
b_5	0	0

For $n = 6$, using the above recursive equation gives

$$b_6 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	0	0
b_3	0	0
b_4	0	0
b_5	0	0
b_6	0	0

For $n = 7$, using the above recursive equation gives

$$b_7 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	0	0
b_3	0	0
b_4	0	0
b_5	0	0
b_6	0	0
b_7	0	0

Using the above table, then the solution $y_2(t)$ is

$$\begin{aligned} y_2(t) &= t^{\frac{2}{3} + \frac{\sqrt{7}}{3}} (b_0 + b_1 t + b_2 t^2 + b_3 t^3 + b_4 t^4 + b_5 t^5 + b_6 t^6 + b_7 t^7 + b_8 t^8 \dots) \\ &= t^{\frac{2}{3} - \frac{\sqrt{7}}{3}} (1 + O(t^8)) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(t) &= c_1 y_1(t) + c_2 y_2(t) \\ &= c_1 t^{\frac{2}{3} + \frac{\sqrt{7}}{3}} (1 + O(t^8)) + c_2 t^{\frac{2}{3} - \frac{\sqrt{7}}{3}} (1 + O(t^8)) \end{aligned}$$

Hence the final solution is

$$\begin{aligned} y(t) &= y_h \\ &= c_1 t^{\frac{2}{3} + \frac{\sqrt{7}}{3}} (1 + O(t^8)) + c_2 t^{\frac{2}{3} - \frac{\sqrt{7}}{3}} (1 + O(t^8)) \end{aligned}$$

Replacing t in the above with the original independent variable x using $t = 1 + x$ results in

$$y = c_1 (1 + x)^{\frac{2}{3} + \frac{\sqrt{7}}{3}} (1 + O((1 + x)^8)) + c_2 (1 + x)^{\frac{2}{3} - \frac{\sqrt{7}}{3}} (1 + O((1 + x)^8))$$

Summary

The solution(s) found are the following

$$y = c_1 (1 + x)^{\frac{2}{3} + \frac{\sqrt{7}}{3}} (1 + O((1 + x)^8)) + c_2 (1 + x)^{\frac{2}{3} - \frac{\sqrt{7}}{3}} (1 + O((1 + x)^8)) \quad (1)$$

Verification of solutions

$$y = c_1 (1 + x)^{\frac{2}{3} + \frac{\sqrt{7}}{3}} (1 + O((1 + x)^8)) + c_2 (1 + x)^{\frac{2}{3} - \frac{\sqrt{7}}{3}} (1 + O((1 + x)^8))$$

Verified OK.

20.7.1 Maple step by step solution

Let's solve

$$3(1 + x)^2 y'' + (-1 - x) y' - y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = \frac{y}{3(1+x)^2} + \frac{y'}{3(1+x)}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - \frac{y'}{3(1+x)} - \frac{y}{3(1+x)^2} = 0$$

- Check to see if $x_0 = -1$ is a regular singular point

- Define functions

$$\left[P_2(x) = -\frac{1}{3(1+x)}, P_3(x) = -\frac{1}{3(1+x)^2} \right]$$

- $(1 + x) \cdot P_2(x)$ is analytic at $x = -1$

$$((1+x) \cdot P_2(x)) \Big|_{x=-1} = -\frac{1}{3}$$

- $(1+x)^2 \cdot P_3(x)$ is analytic at $x = -1$

$$((1+x)^2 \cdot P_3(x)) \Big|_{x=-1} = -\frac{1}{3}$$

- $x = -1$ is a regular singular point

Check to see if $x_0 = -1$ is a regular singular point

$$x_0 = -1$$

- Multiply by denominators

$$3(1+x)^2 y'' + (-1-x)y' - y = 0$$

- Change variables using $x = u - 1$ so that the regular singular point is at $u = 0$

$$3u^2 \left(\frac{d^2}{du^2} y(u) \right) - u \left(\frac{d}{du} y(u) \right) - y(u) = 0$$

- Assume series solution for $y(u)$

$$y(u) = \sum_{k=0}^{\infty} a_k u^{k+r}$$

- Rewrite DE with series expansions

- Convert $u \cdot \left(\frac{d}{du} y(u) \right)$ to series expansion

$$u \cdot \left(\frac{d}{du} y(u) \right) = \sum_{k=0}^{\infty} a_k (k+r) u^{k+r}$$

- Convert $u^2 \cdot \left(\frac{d^2}{du^2} y(u) \right)$ to series expansion

$$u^2 \cdot \left(\frac{d^2}{du^2} y(u) \right) = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) u^{k+r}$$

Rewrite DE with series expansions

$$\sum_{k=0}^{\infty} a_k (3k^2 + 6kr + 3r^2 - 4k - 4r - 1) u^{k+r} = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$r = 0$$

- Each term in the series must be 0, giving the recursion relation

$$a_k (3k^2 - 4k - 1) = 0$$

- Recursion relation that defines series solution to ODE

$$a_k = 0$$

- Recursion relation for $r = 0$
 $a_k = 0$
- Solution for $r = 0$
$$\left[y(u) = \sum_{k=0}^{\infty} a_k u^k, a_k = 0 \right]$$
- Revert the change of variables $u = 1 + x$
$$\left[y = \sum_{k=0}^{\infty} a_k (1 + x)^k, a_k = 0 \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
<- LODE of Euler type successful`

```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 43

```

Order:=8;
dsolve(3*(x+1)^2*diff(y(x),x$2)-(x+1)*diff(y(x),x)-y(x)=0,y(x),type='series',x=-1);

```

$$y(x) = (x + 1)^{\frac{2}{3}} \left((x + 1)^{-\frac{\sqrt{7}}{3}} c_1 + (x + 1)^{\frac{\sqrt{7}}{3}} c_2 \right) + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 42

```

AsymptoticDSolveValue[3*(x+1)^2*y''[x]-(x+1)*y'[x]-y[x]==0,y[x],{x,-1,7}]

```

$$y(x) \rightarrow c_1(x + 1)^{\frac{1}{3}(2+\sqrt{7})} + c_2(x + 1)^{\frac{1}{3}(2-\sqrt{7})}$$

20.8 problem 6

20.8.1 Maple step by step solution 3469

Internal problem ID [6468]

Internal file name [OUTPUT/5716_Sunday_June_05_2022_03_48_31_PM_19121695/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.5. More on Regular Singular Points. Page 183

Problem number: 6.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference is integer"**

Maple gives the following as the ode type

[_Bessel]

$$x^2 y'' + x y' + (x^2 - 1) y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2 y'' + x y' + (x^2 - 1) y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = \frac{x^2 - 1}{x^2}$$

Table 481: Table $p(x), q(x)$ singularities.

$p(x) = \frac{1}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = \frac{x^2-1}{x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2 y'' + x y' + (x^2 - 1) y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned} & x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\ & + x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (x^2 - 1) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \end{aligned} \tag{1}$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) \right) \\ & + \left(\sum_{n=0}^{\infty} x^{n+r+2} a_n \right) + \sum_{n=0}^{\infty} (-a_n x^{n+r}) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} x^{n+r+2} a_n = \sum_{n=2}^{\infty} a_{n-2} x^{n+r}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) \right) \\ & + \left(\sum_{n=2}^{\infty} a_{n-2} x^{n+r} \right) + \sum_{n=0}^{\infty} (-a_n x^{n+r}) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r} a_n (n+r) (n+r-1) + x^{n+r} a_n (n+r) - a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$x^r a_0 r(-1+r) + x^r a_0 r - a_0 x^r = 0$$

Or

$$(x^r r(-1+r) + x^r r - x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(r^2 - 1) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$r^2 - 1 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 1 \\ r_2 &= -1 \end{aligned}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(r^2 - 1) x^r = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = 2$ is an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= C y_1(x) \ln(x) + x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= x \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= C y_1(x) \ln(x) + \frac{\sum_{n=0}^{\infty} b_n x^n}{x} \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^{n+1} \\ y_2(x) &= C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n-1} \right) \end{aligned}$$

Where C above can be zero. We start by finding y_1 . Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

For $2 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) + a_n(n+r) + a_{n-2} - a_n = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{a_{n-2}}{n^2 + 2nr + r^2 - 1} \quad (4)$$

Which for the root $r = 1$ becomes

$$a_n = -\frac{a_{n-2}}{n(n+2)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 1$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = -\frac{1}{r^2 + 4r + 3}$$

Which for the root $r = 1$ becomes

$$a_2 = -\frac{1}{8}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2+4r+3}$	$-\frac{1}{8}$

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2+4r+3}$	$-\frac{1}{8}$
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{(r+3)^2(1+r)(r+5)}$$

Which for the root $r = 1$ becomes

$$a_4 = \frac{1}{192}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2+4r+3}$	$-\frac{1}{8}$
a_3	0	0
a_4	$\frac{1}{(r+3)^2(1+r)(r+5)}$	$\frac{1}{192}$

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2+4r+3}$	$-\frac{1}{8}$
a_3	0	0
a_4	$\frac{1}{(r+3)^2(1+r)(r+5)}$	$\frac{1}{192}$
a_5	0	0

For $n = 6$, using the above recursive equation gives

$$a_6 = -\frac{1}{(r+3)^2(1+r)(r+5)^2(r+7)}$$

Which for the root $r = 1$ becomes

$$a_6 = -\frac{1}{9216}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2+4r+3}$	$-\frac{1}{8}$
a_3	0	0
a_4	$\frac{1}{(r+3)^2(1+r)(r+5)}$	$\frac{1}{192}$
a_5	0	0
a_6	$-\frac{1}{(r+3)^2(1+r)(r+5)^2(r+7)}$	$-\frac{1}{9216}$

For $n = 7$, using the above recursive equation gives

$$a_7 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2+4r+3}$	$-\frac{1}{8}$
a_3	0	0
a_4	$\frac{1}{(r+3)^2(1+r)(r+5)}$	$\frac{1}{192}$
a_5	0	0
a_6	$-\frac{1}{(r+3)^2(1+r)(r+5)^2(r+7)}$	$-\frac{1}{9216}$
a_7	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots) \\ &= x\left(1 - \frac{x^2}{8} + \frac{x^4}{192} - \frac{x^6}{9216} + O(x^8)\right) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Let

$$r_1 - r_2 = N$$

Where N is positive integer which is the difference between the two roots. r_1 is taken as the larger root. Hence for this problem we have $N = 2$. Now we need to determine if C is zero or not. This is done by finding $\lim_{r \rightarrow r_2} a_2(r)$. If this limit exists, then $C = 0$, else we need to keep the log term and $C \neq 0$. The above table shows that

$$\begin{aligned} a_N &= a_2 \\ &= -\frac{1}{r^2 + 4r + 3} \end{aligned}$$

Therefore

$$\begin{aligned} \lim_{r \rightarrow r_2} -\frac{1}{r^2 + 4r + 3} &= \lim_{r \rightarrow -1} -\frac{1}{r^2 + 4r + 3} \\ &= \text{undefined} \end{aligned}$$

Since the limit does not exist then the log term is needed. Therefore the second solution has the form

$$y_2(x) = Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right)$$

Therefore

$$\begin{aligned} \frac{d}{dx} y_2(x) &= Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2} (n+r_2)}{x} \right) \\ &= Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} x^{-1+n+r_2} b_n (n+r_2) \right) \\ \frac{d^2}{dx^2} y_2(x) &= Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} \\ &\quad + \sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2} (n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2} (n+r_2)}{x^2} \right) \\ &= Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} + \left(\sum_{n=0}^{\infty} x^{-2+n+r_2} b_n (n+r_2) (-1+n+r_2) \right) \end{aligned}$$

Substituting these back into the given ode $x^2y'' + xy' + (x^2 - 1)y = 0$ gives

$$\begin{aligned}
& x^2 \left(Cy_1''(x) \ln(x) + \frac{2Cy_1'(x)}{x} - \frac{Cy_1(x)}{x^2} \right. \\
& \left. + \sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2} (n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2} (n+r_2)}{x^2} \right) \right) \\
& + x \left(Cy_1'(x) \ln(x) + \frac{Cy_1(x)}{x} + \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2} (n+r_2)}{x} \right) \right) \\
& + (x^2 - 1) \left(Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) \right) = 0
\end{aligned}$$

Which can be written as

$$\begin{aligned}
& \left((x^2 y_1''(x) + y_1'(x) x + (x^2 - 1) y_1(x)) \ln(x) + x^2 \left(\frac{2y_1'(x)}{x} - \frac{y_1(x)}{x^2} \right) + y_1(x) \right) C \\
& + x^2 \left(\sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2} (n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2} (n+r_2)}{x^2} \right) \right) \\
& + x \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2} (n+r_2)}{x} \right) + (x^2 - 1) \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) = 0
\end{aligned} \tag{7}$$

But since $y_1(x)$ is a solution to the ode, then

$$x^2 y_1''(x) + y_1'(x) x + (x^2 - 1) y_1(x) = 0$$

Eq (7) simplifies to

$$\begin{aligned}
& \left(x^2 \left(\frac{2y_1'(x)}{x} - \frac{y_1(x)}{x^2} \right) + y_1(x) \right) C \\
& + x^2 \left(\sum_{n=0}^{\infty} \left(\frac{b_n x^{n+r_2} (n+r_2)^2}{x^2} - \frac{b_n x^{n+r_2} (n+r_2)}{x^2} \right) \right) \\
& + x \left(\sum_{n=0}^{\infty} \frac{b_n x^{n+r_2} (n+r_2)}{x} \right) + (x^2 - 1) \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) = 0
\end{aligned} \tag{8}$$

Substituting $y_1 = \sum_{n=0}^{\infty} a_n x^{n+r_1}$ into the above gives

$$\begin{aligned}
 & 2x \left(\sum_{n=0}^{\infty} x^{-1+n+r_1} a_n (n+r_1) \right) C + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) x^2 \\
 & + \left(\sum_{n=0}^{\infty} x^{-2+n+r_2} b_n (n+r_2) (-1+n+r_2) \right) x^2 \\
 & + \left(\sum_{n=0}^{\infty} x^{-1+n+r_2} b_n (n+r_2) \right) x - \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right) = 0
 \end{aligned} \tag{9}$$

Since $r_1 = 1$ and $r_2 = -1$ then the above becomes

$$\begin{aligned}
 & 2x \left(\sum_{n=0}^{\infty} x^n a_n (n+1) \right) C + \left(\sum_{n=0}^{\infty} b_n x^{n-1} \right) x^2 + \left(\sum_{n=0}^{\infty} x^{-3+n} b_n (n-1) (n-2) \right) x^2 \\
 & + \left(\sum_{n=0}^{\infty} x^{n-2} b_n (n-1) \right) x - \left(\sum_{n=0}^{\infty} b_n x^{n-1} \right) = 0
 \end{aligned} \tag{10}$$

Which simplifies to

$$\begin{aligned}
 & \left(\sum_{n=0}^{\infty} 2C x^{n+1} a_n (n+1) \right) + \left(\sum_{n=0}^{\infty} x^{n+1} b_n \right) + \left(\sum_{n=0}^{\infty} x^{n-1} b_n (n^2 - 3n + 2) \right) \\
 & + \left(\sum_{n=0}^{\infty} x^{n-1} b_n (n-1) \right) + \sum_{n=0}^{\infty} (-b_n x^{n-1}) = 0
 \end{aligned} \tag{2A}$$

The next step is to make all powers of x be $n-1$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n-1} and adjusting the power and the corresponding index gives

$$\begin{aligned}
 \sum_{n=0}^{\infty} 2C x^{n+1} a_n (n+1) &= \sum_{n=2}^{\infty} 2C a_{n-2} (n-1) x^{n-1} \\
 \sum_{n=0}^{\infty} x^{n+1} b_n &= \sum_{n=2}^{\infty} b_{n-2} x^{n-1}
 \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n - 1$.

$$\begin{aligned} & \left(\sum_{n=2}^{\infty} 2Ca_{n-2}(n-1)x^{n-1} \right) + \left(\sum_{n=2}^{\infty} b_{n-2}x^{n-1} \right) \\ & + \left(\sum_{n=0}^{\infty} x^{n-1}b_n(n^2 - 3n + 2) \right) + \left(\sum_{n=0}^{\infty} x^{n-1}b_n(n-1) \right) + \sum_{n=0}^{\infty} (-b_nx^{n-1}) = 0 \end{aligned} \quad (2B)$$

For $n = 0$ in Eq. (2B), we choose arbitrary value for b_0 as $b_0 = 1$. For $n = 1$, Eq (2B) gives

$$-b_1 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$-b_1 = 0$$

Solving the above for b_1 gives

$$b_1 = 0$$

For $n = N$, where $N = 2$ which is the difference between the two roots, we are free to choose $b_2 = 0$. Hence for $n = 2$, Eq (2B) gives

$$2C + 1 = 0$$

Which is solved for C . Solving for C gives

$$C = -\frac{1}{2}$$

For $n = 3$, Eq (2B) gives

$$4Ca_1 + b_1 + 3b_3 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$3b_3 = 0$$

Solving the above for b_3 gives

$$b_3 = 0$$

For $n = 4$, Eq (2B) gives

$$6Ca_2 + b_2 + 8b_4 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$8b_4 + \frac{3}{8} = 0$$

Solving the above for b_4 gives

$$b_4 = -\frac{3}{64}$$

For $n = 5$, Eq (2B) gives

$$8Ca_3 + b_3 + 15b_5 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$15b_5 = 0$$

Solving the above for b_5 gives

$$b_5 = 0$$

For $n = 6$, Eq (2B) gives

$$10Ca_4 + b_4 + 24b_6 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$24b_6 - \frac{7}{96} = 0$$

Solving the above for b_6 gives

$$b_6 = \frac{7}{2304}$$

For $n = 7$, Eq (2B) gives

$$12Ca_5 + b_5 + 35b_7 = 0$$

Which when replacing the above values found already for b_n and the values found earlier for a_n and for C , gives

$$35b_7 = 0$$

Solving the above for b_7 gives

$$b_7 = 0$$

Now that we found all b_n and C , we can calculate the second solution from

$$y_2(x) = Cy_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n+r_2} \right)$$

Using the above value found for $C = -\frac{1}{2}$ and all b_n , then the second solution becomes

$$y_2(x) = -\frac{1}{2} \left(x \left(1 - \frac{x^2}{8} + \frac{x^4}{192} - \frac{x^6}{9216} + O(x^8) \right) \right) \ln(x) + \frac{1 - \frac{3x^4}{64} + \frac{7x^6}{2304} + O(x^8)}{x}$$

Therefore the homogeneous solution is

$$\begin{aligned}y_h(x) &= c_1 y_1(x) + c_2 y_2(x) \\&= c_1 x \left(1 - \frac{x^2}{8} + \frac{x^4}{192} - \frac{x^6}{9216} + O(x^8) \right) \\&\quad + c_2 \left(-\frac{1}{2} \left(x \left(1 - \frac{x^2}{8} + \frac{x^4}{192} - \frac{x^6}{9216} + O(x^8) \right) \right) \ln(x) + \frac{1 - \frac{3x^4}{64} + \frac{7x^6}{2304} + O(x^8)}{x} \right)\end{aligned}$$

Hence the final solution is

$$\begin{aligned}y &= y_h \\&= c_1 x \left(1 - \frac{x^2}{8} + \frac{x^4}{192} - \frac{x^6}{9216} + O(x^8) \right) \\&\quad + c_2 \left(-\frac{x \left(1 - \frac{x^2}{8} + \frac{x^4}{192} - \frac{x^6}{9216} + O(x^8) \right) \ln(x)}{2} + \frac{1 - \frac{3x^4}{64} + \frac{7x^6}{2304} + O(x^8)}{x} \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}y &= c_1 x \left(1 - \frac{x^2}{8} + \frac{x^4}{192} - \frac{x^6}{9216} + O(x^8) \right) \\&\quad + c_2 \left(-\frac{x \left(1 - \frac{x^2}{8} + \frac{x^4}{192} - \frac{x^6}{9216} + O(x^8) \right) \ln(x)}{2} + \frac{1 - \frac{3x^4}{64} + \frac{7x^6}{2304} + O(x^8)}{x} \right) \quad (1)\end{aligned}$$

Verification of solutions

$$\begin{aligned}y &= c_1 x \left(1 - \frac{x^2}{8} + \frac{x^4}{192} - \frac{x^6}{9216} + O(x^8) \right) \\&\quad + c_2 \left(-\frac{x \left(1 - \frac{x^2}{8} + \frac{x^4}{192} - \frac{x^6}{9216} + O(x^8) \right) \ln(x)}{2} + \frac{1 - \frac{3x^4}{64} + \frac{7x^6}{2304} + O(x^8)}{x} \right)\end{aligned}$$

Verified OK.

20.8.1 Maple step by step solution

Let's solve

$$x^2 y'' + xy' + (x^2 - 1)y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{(x^2-1)y}{x^2} - \frac{y'}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{y'}{x} + \frac{(x^2-1)y}{x^2} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$\left[P_2(x) = \frac{1}{x}, P_3(x) = \frac{x^2-1}{x^2} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 1$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = -1$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$x^2 y'' + xy' + (x^2 - 1)y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 0..2$

$$x^m \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

- Shift index using $k \rightarrow k - m$

$$x^m \cdot y = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

- Convert $x \cdot y'$ to series expansion

$$x \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r}$$

- Convert $x^2 \cdot y''$ to series expansion

$$x^2 \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(1+r)(-1+r)x^r + a_1(2+r)r x^{1+r} + \left(\sum_{k=2}^{\infty} (a_k(k+r+1)(k+r-1) + a_{k-2}) x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$(1+r)(-1+r) = 0$$
- Values of r that satisfy the indicial equation

$$r \in \{-1, 1\}$$
- Each term must be 0

$$a_1(2+r)r = 0$$
- Solve for the dependent coefficient(s)

$$a_1 = 0$$
- Each term in the series must be 0, giving the recursion relation

$$a_k(k+r+1)(k+r-1) + a_{k-2} = 0$$
- Shift index using $k \rightarrow k+2$

$$a_{k+2}(k+3+r)(k+r+1) + a_k = 0$$
- Recursion relation that defines series solution to ODE

$$a_{k+2} = -\frac{a_k}{(k+3+r)(k+r+1)}$$
- Recursion relation for $r = -1$

$$a_{k+2} = -\frac{a_k}{(k+2)k}$$
- Solution for $r = -1$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k-1}, a_{k+2} = -\frac{a_k}{(k+2)k}, a_1 = 0 \right]$$

- Recursion relation for $r = 1$

$$a_{k+2} = -\frac{a_k}{(k+4)(k+2)}$$

- Solution for $r = 1$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+1}, a_{k+2} = -\frac{a_k}{(k+4)(k+2)}, a_1 = 0 \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^{k-1} \right) + \left(\sum_{k=0}^{\infty} b_k x^{k+1} \right), a_{k+2} = -\frac{a_k}{(k+2)k}, a_1 = 0, b_{k+2} = -\frac{b_k}{(k+4)(k+2)}, b_1 = 0 \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
    <- Bessel successful
<- special function solution successful`

```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 53

```

Order:=8;
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)+(x^2-1)*y(x)=0,y(x),type='series',x=0);

```

$$y(x) = \frac{c_1 x^2 \left(1 - \frac{1}{8} x^2 + \frac{1}{192} x^4 - \frac{1}{9216} x^6 + O(x^8) \right) + c_2 \left(\left(x^2 - \frac{1}{8} x^4 + \frac{1}{192} x^6 + O(x^8) \right) \ln(x) + \left(-2 + \frac{3}{32} x^4 - \frac{7}{1152} x^6 + O(x^8) \right) \right)}{x}$$

✓ Solution by Mathematica

Time used: 0.013 (sec). Leaf size: 75

```
AsymptoticDSolveValue[x^2*y''[x]+x*y'[x]+(x^2-1)*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_2 \left(-\frac{x^7}{9216} + \frac{x^5}{192} - \frac{x^3}{8} + x \right) + c_1 \left(\frac{5x^6 - 90x^4 + 288x^2 + 1152}{1152x} - \frac{1}{384}x(x^4 - 24x^2 + 192) \log(x) \right)$$

20.9 problem 7

20.9.1 Maple step by step solution 3484

Internal problem ID [6469]

Internal file name [OUTPUT/5717_Sunday_June_05_2022_03_48_35_PM_34738364/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.5. More on Regular Singular Points. Page 183

Problem number: 7.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference is integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$x^2y'' + xy' + \left(x^2 - \frac{1}{4}\right)y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2y'' + xy' + \left(x^2 - \frac{1}{4}\right)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = \frac{4x^2 - 1}{4x^2}$$

Table 483: Table $p(x), q(x)$ singularities.

$p(x) = \frac{1}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = \frac{4x^2-1}{4x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2 y'' + x y' + \left(x^2 - \frac{1}{4}\right) y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned} & x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\ & + x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(x^2 - \frac{1}{4}\right) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \end{aligned} \tag{1}$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) \right) \\ & + \left(\sum_{n=0}^{\infty} x^{n+r+2} a_n \right) + \sum_{n=0}^{\infty} \left(-\frac{a_n x^{n+r}}{4} \right) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} x^{n+r+2} a_n = \sum_{n=2}^{\infty} a_{n-2} x^{n+r}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) \right) \\ & + \left(\sum_{n=2}^{\infty} a_{n-2} x^{n+r} \right) + \sum_{n=0}^{\infty} \left(-\frac{a_n x^{n+r}}{4} \right) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r} a_n (n+r) (n+r-1) + x^{n+r} a_n (n+r) - \frac{a_n x^{n+r}}{4} = 0$$

When $n=0$ the above becomes

$$x^r a_0 r(-1+r) + x^r a_0 r - \frac{a_0 x^r}{4} = 0$$

Or

$$\left(x^r r(-1+r) + x^r r - \frac{x^r}{4} \right) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$\frac{(4r^2 - 1) x^r}{4} = 0$$

Since the above is true for all x then the indicial equation becomes

$$r^2 - \frac{1}{4} = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = \frac{1}{2}$$

$$r_2 = -\frac{1}{2}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$\frac{(4r^2 - 1)x^r}{4} = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = 1$ is an integer, then we can construct two linearly independent solutions

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = C y_1(x) \ln(x) + x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sqrt{x} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = C y_1(x) \ln(x) + \frac{\sum_{n=0}^{\infty} b_n x^n}{\sqrt{x}}$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+\frac{1}{2}}$$

$$y_2(x) = C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^{n-\frac{1}{2}} \right)$$

Where C above can be zero. We start by finding y_1 . Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

For $2 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) + a_n(n+r) + a_{n-2} - \frac{a_n}{4} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{4a_{n-2}}{4n^2 + 8nr + 4r^2 - 1} \quad (4)$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_n = -\frac{a_{n-2}}{n(n+1)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = \frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = -\frac{4}{4r^2 + 16r + 15}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_2 = -\frac{1}{6}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{4}{4r^2+16r+15}$	$-\frac{1}{6}$

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{4}{4r^2+16r+15}$	$-\frac{1}{6}$
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{16}{(4r^2 + 16r + 15)(4r^2 + 32r + 63)}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_4 = \frac{1}{120}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{4}{4r^2+16r+15}$	$-\frac{1}{6}$
a_3	0	0
a_4	$\frac{16}{(4r^2+16r+15)(4r^2+32r+63)}$	$\frac{1}{120}$

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{4}{4r^2+16r+15}$	$-\frac{1}{6}$
a_3	0	0
a_4	$\frac{16}{(4r^2+16r+15)(4r^2+32r+63)}$	$\frac{1}{120}$
a_5	0	0

For $n = 6$, using the above recursive equation gives

$$a_6 = -\frac{64}{(4r^2 + 16r + 15)(4r^2 + 32r + 63)(4r^2 + 48r + 143)}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_6 = -\frac{1}{5040}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{4}{4r^2+16r+15}$	$-\frac{1}{6}$
a_3	0	0
a_4	$\frac{16}{(4r^2+16r+15)(4r^2+32r+63)}$	$\frac{1}{120}$
a_5	0	0
a_6	$-\frac{64}{(4r^2+16r+15)(4r^2+32r+63)(4r^2+48r+143)}$	$-\frac{1}{5040}$

For $n = 7$, using the above recursive equation gives

$$a_7 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{4}{4r^2+16r+15}$	$-\frac{1}{6}$
a_3	0	0
a_4	$\frac{16}{(4r^2+16r+15)(4r^2+32r+63)}$	$\frac{1}{120}$
a_5	0	0
a_6	$-\frac{64}{(4r^2+16r+15)(4r^2+32r+63)(4r^2+48r+143)}$	$-\frac{1}{5040}$
a_7	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= \sqrt{x}(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots) \\ &= \sqrt{x} \left(1 - \frac{x^2}{6} + \frac{x^4}{120} - \frac{x^6}{5040} + O(x^8) \right) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Let

$$r_1 - r_2 = N$$

Where N is positive integer which is the difference between the two roots. r_1 is taken as the larger root. Hence for this problem we have $N = 1$. Now we need to determine if C is zero or not. This is done by finding $\lim_{r \rightarrow r_2} a_1(r)$. If this limit exists, then $C = 0$, else we need to keep the log term and $C \neq 0$. The above table shows that

$$\begin{aligned} a_N &= a_1 \\ &= 0 \end{aligned}$$

Therefore

$$\begin{aligned} \lim_{r \rightarrow r_2} 0 &= \lim_{r \rightarrow -\frac{1}{2}} 0 \\ &= 0 \end{aligned}$$

The limit is 0. Since the limit exists then the log term is not needed and we can set $C = 0$. Therefore the second solution has the form

$$\begin{aligned} y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+r} \\ &= \sum_{n=0}^{\infty} b_n x^{n-\frac{1}{2}} \end{aligned}$$

Eq (3) derived above is used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq(3) gives

$$b_1 = 0$$

For $2 \leq n$ the recursive equation is

$$b_n(n+r)(n+r-1) + b_n(n+r) + b_{n-2} - \frac{b_n}{4} = 0 \quad (4)$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_n \left(n - \frac{1}{2} \right) \left(n - \frac{3}{2} \right) + b_n \left(n - \frac{1}{2} \right) + b_{n-2} - \frac{b_n}{4} = 0 \quad (4A)$$

Solving for b_n from the recursive equation (4) gives

$$b_n = -\frac{4b_{n-2}}{4n^2 + 8nr + 4r^2 - 1} \quad (5)$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_n = -\frac{4b_{n-2}}{4n^2 - 4n} \quad (6)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = -\frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0

For $n = 2$, using the above recursive equation gives

$$b_2 = -\frac{4}{4r^2 + 16r + 15}$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_2 = -\frac{1}{2}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{4}{4r^2+16r+15}$	$-\frac{1}{2}$

For $n = 3$, using the above recursive equation gives

$$b_3 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{4}{4r^2+16r+15}$	$-\frac{1}{2}$
b_3	0	0

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{16}{(4r^2 + 16r + 15)(4r^2 + 32r + 63)}$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_4 = \frac{1}{24}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{4}{4r^2+16r+15}$	$-\frac{1}{2}$
b_3	0	0
b_4	$\frac{16}{(4r^2+16r+15)(4r^2+32r+63)}$	$\frac{1}{24}$

For $n = 5$, using the above recursive equation gives

$$b_5 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{4}{4r^2+16r+15}$	$-\frac{1}{2}$
b_3	0	0
b_4	$\frac{16}{(4r^2+16r+15)(4r^2+32r+63)}$	$\frac{1}{24}$
b_5	0	0

For $n = 6$, using the above recursive equation gives

$$b_6 = -\frac{64}{(4r^2 + 16r + 15)(4r^2 + 32r + 63)(4r^2 + 48r + 143)}$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_6 = -\frac{1}{720}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{4}{4r^2+16r+15}$	$-\frac{1}{2}$
b_3	0	0
b_4	$\frac{16}{(4r^2+16r+15)(4r^2+32r+63)}$	$\frac{1}{24}$
b_5	0	0
b_6	$-\frac{64}{(4r^2+16r+15)(4r^2+32r+63)(4r^2+48r+143)}$	$-\frac{1}{720}$

For $n = 7$, using the above recursive equation gives

$$b_7 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{4}{4r^2+16r+15}$	$-\frac{1}{2}$
b_3	0	0
b_4	$\frac{16}{(4r^2+16r+15)(4r^2+32r+63)}$	$\frac{1}{24}$
b_5	0	0
b_6	$-\frac{64}{(4r^2+16r+15)(4r^2+32r+63)(4r^2+48r+143)}$	$-\frac{1}{720}$
b_7	0	0

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned} y_2(x) &= \sqrt{x}(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 + b_7x^7 + b_8x^8 \dots) \\ &= \frac{1 - \frac{x^2}{2} + \frac{x^4}{24} - \frac{x^6}{720} + O(x^8)}{\sqrt{x}} \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= c_1\sqrt{x} \left(1 - \frac{x^2}{6} + \frac{x^4}{120} - \frac{x^6}{5040} + O(x^8) \right) + \frac{c_2 \left(1 - \frac{x^2}{2} + \frac{x^4}{24} - \frac{x^6}{720} + O(x^8) \right)}{\sqrt{x}} \end{aligned}$$

Hence the final solution is

$$\begin{aligned} y &= y_h \\ &= c_1\sqrt{x} \left(1 - \frac{x^2}{6} + \frac{x^4}{120} - \frac{x^6}{5040} + O(x^8) \right) + \frac{c_2 \left(1 - \frac{x^2}{2} + \frac{x^4}{24} - \frac{x^6}{720} + O(x^8) \right)}{\sqrt{x}} \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1\sqrt{x} \left(1 - \frac{x^2}{6} + \frac{x^4}{120} - \frac{x^6}{5040} + O(x^8) \right) + \frac{c_2 \left(1 - \frac{x^2}{2} + \frac{x^4}{24} - \frac{x^6}{720} + O(x^8) \right)}{\sqrt{x}} \quad (1)$$

Verification of solutions

$$y = c_1\sqrt{x} \left(1 - \frac{x^2}{6} + \frac{x^4}{120} - \frac{x^6}{5040} + O(x^8) \right) + \frac{c_2 \left(1 - \frac{x^2}{2} + \frac{x^4}{24} - \frac{x^6}{720} + O(x^8) \right)}{\sqrt{x}}$$

Verified OK.

20.9.1 Maple step by step solution

Let's solve

$$x^2y'' + xy' + \left(x^2 - \frac{1}{4}\right)y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{(4x^2-1)y}{4x^2} - \frac{y'}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{y'}{x} + \frac{(4x^2-1)y}{4x^2} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$\left[P_2(x) = \frac{1}{x}, P_3(x) = \frac{4x^2-1}{4x^2} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 1$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = -\frac{1}{4}$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$4x^2y'' + 4xy' + (4x^2 - 1)y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 0..2$

$$x^m \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

- Shift index using $k \rightarrow k - m$

$$x^m \cdot y = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

- Convert $x \cdot y'$ to series expansion

$$x \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r}$$

- Convert $x^2 \cdot y''$ to series expansion

$$x^2 \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(1+2r)(-1+2r)x^r + a_1(3+2r)(1+2r)x^{1+r} + \left(\sum_{k=2}^{\infty} (a_k(2k+2r+1)(2k+2r-1) + 4a_{k-2}) \right) x^{k+r} = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$(1+2r)(-1+2r) = 0$$
- Values of r that satisfy the indicial equation

$$r \in \left\{ -\frac{1}{2}, \frac{1}{2} \right\}$$
- Each term must be 0

$$a_1(3+2r)(1+2r) = 0$$
- Solve for the dependent coefficient(s)

$$a_1 = 0$$
- Each term in the series must be 0, giving the recursion relation

$$a_k(4k^2 + 8kr + 4r^2 - 1) + 4a_{k-2} = 0$$
- Shift index using $k- > k+2$

$$a_{k+2}(4(k+2)^2 + 8(k+2)r + 4r^2 - 1) + 4a_k = 0$$
- Recursion relation that defines series solution to ODE

$$a_{k+2} = -\frac{4a_k}{4k^2 + 8kr + 4r^2 + 16k + 16r + 15}$$
- Recursion relation for $r = -\frac{1}{2}$

$$a_{k+2} = -\frac{4a_k}{4k^2 + 12k + 8}$$
- Solution for $r = -\frac{1}{2}$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k-\frac{1}{2}}, a_{k+2} = -\frac{4a_k}{4k^2 + 12k + 8}, a_1 = 0 \right]$$
- Recursion relation for $r = \frac{1}{2}$

$$a_{k+2} = -\frac{4a_k}{4k^2 + 20k + 24}$$
- Solution for $r = \frac{1}{2}$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+\frac{1}{2}}, a_{k+2} = -\frac{4a_k}{4k^2 + 20k + 24}, a_1 = 0 \right]$$
- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^{k-\frac{1}{2}} \right) + \left(\sum_{k=0}^{\infty} b_k x^{k+\frac{1}{2}} \right), a_{k+2} = -\frac{4a_k}{4k^2+12k+8}, a_1 = 0, b_{k+2} = -\frac{4b_k}{4k^2+20k+24}, b_1 = 0 \right]$$

Maple trace Kovacic algorithm successful

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Group is reducible or imprimitive
<- Kovacics algorithm successful`

```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 39

```

Order:=8;
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)+(x^2-1/4)*y(x)=0,y(x),type='series',x=0);

```

$$y(x) = \frac{c_1 \left(1 - \frac{1}{6}x^2 + \frac{1}{120}x^4 - \frac{1}{5040}x^6 + O(x^8) \right) x + c_2 \left(1 - \frac{1}{2}x^2 + \frac{1}{24}x^4 - \frac{1}{720}x^6 + O(x^8) \right)}{\sqrt{x}}$$

✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 76

```

AsymptoticDSolveValue[x^2*y''[x]+x*y'[x]+(x^2-1/4)*y[x]==0,y[x],{x,0,7}]

```

$$y(x) \rightarrow c_1 \left(-\frac{x^{11/2}}{720} + \frac{x^{7/2}}{24} - \frac{x^{3/2}}{2} + \frac{1}{\sqrt{x}} \right) + c_2 \left(-\frac{x^{13/2}}{5040} + \frac{x^{9/2}}{120} - \frac{x^{5/2}}{6} + \sqrt{x} \right)$$

21 Chapter 4. Power Series Solutions and Special Functions. Section 4.6. Gauss's Hypergeometric Equation. Page 187

21.1 problem 2(a)	3489
21.2 problem 2(b)	3505
21.3 problem 2(x)	3520
21.4 problem 2(d)	3538
21.5 problem 3	3556
21.6 problem 5	3579

21.1 problem 2(a)

21.1.1 Maple step by step solution 3501

Internal problem ID [6470]

Internal file name [OUTPUT/5718_Sunday_June_05_2022_03_48_39_PM_66849501/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.6. Gauss's Hypergeometric Equation. Page 187

Problem number: 2(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

[_Jacobi]

$$x(1-x)y'' + \left(\frac{3}{2} - 2x\right)y' + 2y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$(-x^2 + x)y'' + \left(\frac{3}{2} - 2x\right)y' + 2y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{-3 + 4x}{2x(x-1)}$$
$$q(x) = -\frac{2}{x(x-1)}$$

Table 485: Table $p(x), q(x)$ singularities.

$p(x) = \frac{-3+4x}{2x(x-1)}$	
singularity	type
$x = 0$	“regular”
$x = 1$	“regular”

$q(x) = -\frac{2}{x(x-1)}$	
singularity	type
$x = 0$	“regular”
$x = 1$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0, 1, \infty]$

Irregular singular points : $[\]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$-y''x(x-1) + \left(\frac{3}{2} - 2x\right)y' + 2y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned} & -\left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}\right) x(x-1) \\ & + \left(\frac{3}{2} - 2x\right) \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}\right) + 2\left(\sum_{n=0}^{\infty} a_n x^{n+r}\right) = 0 \end{aligned} \tag{1}$$

Which simplifies to

$$\begin{aligned} \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r) (n+r-1)) + \left(\sum_{n=0}^{\infty} x^{n+r-1} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-2x^{n+r} a_n (n+r)) \\ + \left(\sum_{n=0}^{\infty} \frac{3(n+r) a_n x^{n+r-1}}{2} \right) + \left(\sum_{n=0}^{\infty} 2a_n x^{n+r} \right) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r-1$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r-1} and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r) (n+r-1)) &= \sum_{n=1}^{\infty} (-a_{n-1} (n+r-1) (n+r-2) x^{n+r-1}) \\ \sum_{n=0}^{\infty} (-2x^{n+r} a_n (n+r)) &= \sum_{n=1}^{\infty} (-2a_{n-1} (n+r-1) x^{n+r-1}) \\ \sum_{n=0}^{\infty} 2a_n x^{n+r} &= \sum_{n=1}^{\infty} 2a_{n-1} x^{n+r-1} \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r-1$.

$$\begin{aligned} \sum_{n=1}^{\infty} (-a_{n-1} (n+r-1) (n+r-2) x^{n+r-1}) \\ + \left(\sum_{n=0}^{\infty} x^{n+r-1} a_n (n+r) (n+r-1) \right) + \sum_{n=1}^{\infty} (-2a_{n-1} (n+r-1) x^{n+r-1}) \\ + \left(\sum_{n=0}^{\infty} \frac{3(n+r) a_n x^{n+r-1}}{2} \right) + \left(\sum_{n=1}^{\infty} 2a_{n-1} x^{n+r-1} \right) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r-1} a_n (n+r) (n+r-1) + \frac{3(n+r) a_n x^{n+r-1}}{2} = 0$$

When $n = 0$ the above becomes

$$x^{-1+r} a_0 r(-1+r) + \frac{3r a_0 x^{-1+r}}{2} = 0$$

Or

$$\left(x^{-1+r} r(-1+r) + \frac{3r x^{-1+r}}{2} \right) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$r x^{-1+r} \left(\frac{1}{2} + r \right) = 0$$

Since the above is true for all x then the indicial equation becomes

$$r^2 + \frac{1}{2}r = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 0 \\ r_2 &= -\frac{1}{2} \end{aligned}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$r x^{-1+r} \left(\frac{1}{2} + r \right) = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = \frac{1}{2}$ is not an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^n \\ y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n-\frac{1}{2}} \end{aligned}$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$\begin{aligned}
 & -a_{n-1}(n+r-1)(n+r-2) + a_n(n+r)(n+r-1) \\
 & - 2a_{n-1}(n+r-1) + \frac{3a_n(n+r)}{2} + 2a_{n-1} = 0
 \end{aligned} \tag{3}$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{2a_{n-1}(n^2 + 2nr + r^2 - n - r - 2)}{2n^2 + 4nr + 2r^2 + n + r} \tag{4}$$

Which for the root $r = 0$ becomes

$$a_n = \frac{2a_{n-1}(n^2 - n - 2)}{2n^2 + n} \tag{5}$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = \frac{2r^2 + 2r - 4}{2r^2 + 5r + 3}$$

Which for the root $r = 0$ becomes

$$a_1 = -\frac{4}{3}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{2r^2+2r-4}{2r^2+5r+3}$	$-\frac{4}{3}$

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{4(-1+r)r(r+3)}{4r^3 + 20r^2 + 31r + 15}$$

Which for the root $r = 0$ becomes

$$a_2 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{2r^2+2r-4}{2r^2+5r+3}$	$-\frac{4}{3}$
a_2	$\frac{4(-1+r)r(r+3)}{4r^3+20r^2+31r+15}$	0

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{8(-1+r)r(r+4)}{8r^3+60r^2+142r+105}$$

Which for the root $r = 0$ becomes

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{2r^2+2r-4}{2r^2+5r+3}$	$-\frac{4}{3}$
a_2	$\frac{4(-1+r)r(r+3)}{4r^3+20r^2+31r+15}$	0
a_3	$\frac{8(-1+r)r(r+4)}{8r^3+60r^2+142r+105}$	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{16(r+5)(r+2)r(-1+r)}{16r^4+192r^3+824r^2+1488r+945}$$

Which for the root $r = 0$ becomes

$$a_4 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{2r^2+2r-4}{2r^2+5r+3}$	$-\frac{4}{3}$
a_2	$\frac{4(-1+r)r(r+3)}{4r^3+20r^2+31r+15}$	0
a_3	$\frac{8(-1+r)r(r+4)}{8r^3+60r^2+142r+105}$	0
a_4	$\frac{16(r+5)(r+2)r(-1+r)}{16r^4+192r^3+824r^2+1488r+945}$	0

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{32(r+6)(r+3)(-1+r)r(r+2)}{32r^5 + 560r^4 + 3760r^3 + 12040r^2 + 18258r + 10395}$$

Which for the root $r = 0$ becomes

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{2r^2+2r-4}{2r^2+5r+3}$	$-\frac{4}{3}$
a_2	$\frac{4(-1+r)r(r+3)}{4r^3+20r^2+31r+15}$	0
a_3	$\frac{8(-1+r)r(r+4)}{8r^3+60r^2+142r+105}$	0
a_4	$\frac{16(r+5)(r+2)r(-1+r)}{16r^4+192r^3+824r^2+1488r+945}$	0
a_5	$\frac{32(r+6)(r+3)(-1+r)r(r+2)}{32r^5+560r^4+3760r^3+12040r^2+18258r+10395}$	0

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{64(r+7)(r+4)(r+2)(-1+r)r(r+3)}{64r^6 + 1536r^5 + 14800r^4 + 72960r^3 + 193036r^2 + 258144r + 135135}$$

Which for the root $r = 0$ becomes

$$a_6 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{2r^2+2r-4}{2r^2+5r+3}$	$-\frac{4}{3}$
a_2	$\frac{4(-1+r)r(r+3)}{4r^3+20r^2+31r+15}$	0
a_3	$\frac{8(-1+r)r(r+4)}{8r^3+60r^2+142r+105}$	0
a_4	$\frac{16(r+5)(r+2)r(-1+r)}{16r^4+192r^3+824r^2+1488r+945}$	0
a_5	$\frac{32(r+6)(r+3)(-1+r)r(r+2)}{32r^5+560r^4+3760r^3+12040r^2+18258r+10395}$	0
a_6	$\frac{64(r+7)(r+4)(r+2)(-1+r)r(r+3)}{64r^6+1536r^5+14800r^4+72960r^3+193036r^2+258144r+135135}$	0

For $n = 7$, using the above recursive equation gives

$$a_7 = \frac{128(r+8)(r+5)(-1+r)r(r+3)(r+2)(r+4)}{128r^7 + 4032r^6 + 52640r^5 + 367920r^4 + 1480472r^3 + 3411828r^2 + 4142430r + 2027025}$$

Which for the root $r = 0$ becomes

$$a_7 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{2r^2+2r-4}{2r^2+5r+3}$	$-\frac{4}{3}$
a_2	$\frac{4(-1+r)r(r+3)}{4r^3+20r^2+31r+15}$	0
a_3	$\frac{8(-1+r)r(r+4)}{8r^3+60r^2+142r+105}$	0
a_4	$\frac{16(r+5)(r+2)r(-1+r)}{16r^4+192r^3+824r^2+1488r+945}$	0
a_5	$\frac{32(r+6)(r+3)(-1+r)r(r+2)}{32r^5+560r^4+3760r^3+12040r^2+18258r+10395}$	0
a_6	$\frac{64(r+7)(r+4)(r+2)(-1+r)r(r+3)}{64r^6+1536r^5+14800r^4+72960r^3+193036r^2+258144r+135135}$	0
a_7	$\frac{128(r+8)(r+5)(-1+r)r(r+3)(r+2)(r+4)}{128r^7+4032r^6+52640r^5+367920r^4+1480472r^3+3411828r^2+4142430r+2027025}$	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots \\ &= 1 - \frac{4x}{3} + O(x^8) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. For $1 \leq n$ the recursive equation is

$$-b_{n-1}(n+r-1)(n+r-2) + b_n(n+r)(n+r-1) - 2b_{n-1}(n+r-1) + \frac{3(n+r)b_n}{2} + 2b_{n-1} = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = \frac{2b_{n-1}(n^2 + 2nr + r^2 - n - r - 2)}{2n^2 + 4nr + 2r^2 + n + r} \quad (4)$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_n = \frac{b_{n-1}(4n^2 - 8n - 5)}{4n^2 - 2n} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = -\frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1

For $n = 1$, using the above recursive equation gives

$$b_1 = \frac{2r^2 + 2r - 4}{2r^2 + 5r + 3}$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_1 = -\frac{9}{2}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{2r^2+2r-4}{2r^2+5r+3}$	$-\frac{9}{2}$

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{4(-1+r)r(r+3)}{4r^3 + 20r^2 + 31r + 15}$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_2 = \frac{15}{8}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{2r^2+2r-4}{2r^2+5r+3}$	$-\frac{9}{2}$
b_2	$\frac{4(-1+r)r(r+3)}{4r^3+20r^2+31r+15}$	$\frac{15}{8}$

For $n = 3$, using the above recursive equation gives

$$b_3 = \frac{8(-1+r)r(r+4)}{8r^3+60r^2+142r+105}$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_3 = \frac{7}{16}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{2r^2+2r-4}{2r^2+5r+3}$	$-\frac{9}{2}$
b_2	$\frac{4(-1+r)r(r+3)}{4r^3+20r^2+31r+15}$	$\frac{15}{8}$
b_3	$\frac{8(-1+r)r(r+4)}{8r^3+60r^2+142r+105}$	$\frac{7}{16}$

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{16(r+5)(r+2)r(-1+r)}{16r^4+192r^3+824r^2+1488r+945}$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_4 = \frac{27}{128}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{2r^2+2r-4}{2r^2+5r+3}$	$-\frac{9}{2}$
b_2	$\frac{4(-1+r)r(r+3)}{4r^3+20r^2+31r+15}$	$\frac{15}{8}$
b_3	$\frac{8(-1+r)r(r+4)}{8r^3+60r^2+142r+105}$	$\frac{7}{16}$
b_4	$\frac{16(r+5)(r+2)r(-1+r)}{16r^4+192r^3+824r^2+1488r+945}$	$\frac{27}{128}$

For $n = 5$, using the above recursive equation gives

$$b_5 = \frac{32(r+6)(r+3)(-1+r)r(r+2)}{32r^5 + 560r^4 + 3760r^3 + 12040r^2 + 18258r + 10395}$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_5 = \frac{33}{256}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{2r^2+2r-4}{2r^2+5r+3}$	$-\frac{9}{2}$
b_2	$\frac{4(-1+r)r(r+3)}{4r^3+20r^2+31r+15}$	$\frac{15}{8}$
b_3	$\frac{8(-1+r)r(r+4)}{8r^3+60r^2+142r+105}$	$\frac{7}{16}$
b_4	$\frac{16(r+5)(r+2)r(-1+r)}{16r^4+192r^3+824r^2+1488r+945}$	$\frac{27}{128}$
b_5	$\frac{32(r+6)(r+3)(-1+r)r(r+2)}{32r^5+560r^4+3760r^3+12040r^2+18258r+10395}$	$\frac{33}{256}$

For $n = 6$, using the above recursive equation gives

$$b_6 = \frac{64(r+7)(r+4)(r+2)(-1+r)r(r+3)}{64r^6 + 1536r^5 + 14800r^4 + 72960r^3 + 193036r^2 + 258144r + 135135}$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_6 = \frac{91}{1024}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{2r^2+2r-4}{2r^2+5r+3}$	$-\frac{9}{2}$
b_2	$\frac{4(-1+r)r(r+3)}{4r^3+20r^2+31r+15}$	$\frac{15}{8}$
b_3	$\frac{8(-1+r)r(r+4)}{8r^3+60r^2+142r+105}$	$\frac{7}{16}$
b_4	$\frac{16(r+5)(r+2)r(-1+r)}{16r^4+192r^3+824r^2+1488r+945}$	$\frac{27}{128}$
b_5	$\frac{32(r+6)(r+3)(-1+r)r(r+2)}{32r^5+560r^4+3760r^3+12040r^2+18258r+10395}$	$\frac{33}{256}$
b_6	$\frac{64(r+7)(r+4)(r+2)(-1+r)r(r+3)}{64r^6+1536r^5+14800r^4+72960r^3+193036r^2+258144r+135135}$	$\frac{91}{1024}$

For $n = 7$, using the above recursive equation gives

$$b_7 = \frac{128(r+8)(r+5)(-1+r)r(r+3)(r+2)(r+4)}{128r^7 + 4032r^6 + 52640r^5 + 367920r^4 + 1480472r^3 + 3411828r^2 + 4142430r + 2027025}$$

Which for the root $r = -\frac{1}{2}$ becomes

$$b_7 = \frac{135}{2048}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{2r^2+2r-4}{2r^2+5r+3}$	$-\frac{9}{2}$
b_2	$\frac{4(-1+r)r(r+3)}{4r^3+20r^2+31r+15}$	$\frac{15}{8}$
b_3	$\frac{8(-1+r)r(r+4)}{8r^3+60r^2+142r+105}$	$\frac{7}{16}$
b_4	$\frac{16(r+5)(r+2)r(-1+r)}{16r^4+192r^3+824r^2+1488r+945}$	$\frac{27}{128}$
b_5	$\frac{32(r+6)(r+3)(-1+r)r(r+2)}{32r^5+560r^4+3760r^3+12040r^2+18258r+10395}$	$\frac{33}{256}$
b_6	$\frac{64(r+7)(r+4)(r+2)(-1+r)r(r+3)}{64r^6+1536r^5+14800r^4+72960r^3+193036r^2+258144r+135135}$	$\frac{91}{1024}$
b_7	$\frac{128(r+8)(r+5)(-1+r)r(r+3)(r+2)(r+4)}{128r^7+4032r^6+52640r^5+367920r^4+1480472r^3+3411828r^2+4142430r+2027025}$	$\frac{135}{2048}$

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned} y_2(x) &= 1(b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 + b_7x^7 + b_8x^8 \dots) \\ &= \frac{1 - \frac{9x}{2} + \frac{15x^2}{8} + \frac{7x^3}{16} + \frac{27x^4}{128} + \frac{33x^5}{256} + \frac{91x^6}{1024} + \frac{135x^7}{2048} + O(x^8)}{\sqrt{x}} \end{aligned}$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 y_1(x) + c_2 y_2(x)$$

$$= c_1 \left(1 - \frac{4x}{3} + O(x^8) \right) + \frac{c_2 \left(1 - \frac{9x}{2} + \frac{15x^2}{8} + \frac{7x^3}{16} + \frac{27x^4}{128} + \frac{33x^5}{256} + \frac{91x^6}{1024} + \frac{135x^7}{2048} + O(x^8) \right)}{\sqrt{x}}$$

Hence the final solution is

$$y = y_h$$

$$= c_1 \left(1 - \frac{4x}{3} + O(x^8) \right) + \frac{c_2 \left(1 - \frac{9x}{2} + \frac{15x^2}{8} + \frac{7x^3}{16} + \frac{27x^4}{128} + \frac{33x^5}{256} + \frac{91x^6}{1024} + \frac{135x^7}{2048} + O(x^8) \right)}{\sqrt{x}}$$

Summary

The solution(s) found are the following

$$y = c_1 \left(1 - \frac{4x}{3} + O(x^8) \right) + \frac{c_2 \left(1 - \frac{9x}{2} + \frac{15x^2}{8} + \frac{7x^3}{16} + \frac{27x^4}{128} + \frac{33x^5}{256} + \frac{91x^6}{1024} + \frac{135x^7}{2048} + O(x^8) \right)}{\sqrt{x}} \quad (1)$$

Verification of solutions

$$y = c_1 \left(1 - \frac{4x}{3} + O(x^8) \right) + \frac{c_2 \left(1 - \frac{9x}{2} + \frac{15x^2}{8} + \frac{7x^3}{16} + \frac{27x^4}{128} + \frac{33x^5}{256} + \frac{91x^6}{1024} + \frac{135x^7}{2048} + O(x^8) \right)}{\sqrt{x}}$$

Verified OK.

21.1.1 Maple step by step solution

Let's solve

$$-y''x(x-1) + \left(\frac{3}{2} - 2x\right)y' + 2y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{(-3+4x)y'}{2x(x-1)} + \frac{2y}{x(x-1)}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{(-3+4x)y'}{2x(x-1)} - \frac{2y}{x(x-1)} = 0$$

□ Check to see if x_0 is a regular singular point

○ Define functions

$$\left[P_2(x) = \frac{-3+4x}{2x(x-1)}, P_3(x) = -\frac{2}{x(x-1)} \right]$$

○ $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = \frac{3}{2}$$

○ $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 0$$

○ $x = 0$ is a regular singular point

Check to see if x_0 is a regular singular point

$$x_0 = 0$$

• Multiply by denominators

$$2y''x(x-1) + (-3+4x)y' - 4y = 0$$

• Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

□ Rewrite ODE with series expansions

○ Convert $x^m \cdot y'$ to series expansion for $m = 0..1$

$$x^m \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1+m}$$

○ Shift index using $k \rightarrow k+1-m$

$$x^m \cdot y' = \sum_{k=-1+m}^{\infty} a_{k+1-m} (k+1-m+r) x^{k+r}$$

○ Convert $x^m \cdot y''$ to series expansion for $m = 1..2$

$$x^m \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r-2+m}$$

○ Shift index using $k \rightarrow k+2-m$

$$x^m \cdot y'' = \sum_{k=-2+m}^{\infty} a_{k+2-m} (k+2-m+r)(k+1-m+r) x^{k+r}$$

Rewrite ODE with series expansions

$$-a_0 r(2r+1)x^{-1+r} + \left(\sum_{k=0}^{\infty} (-a_{k+1}(k+1+r)(2k+3+2r) + 2a_k(k+r+2)(k+r-1))x^{k+r} \right)$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$-r(2r+1) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \left\{ 0, -\frac{1}{2} \right\}$$

- Each term in the series must be 0, giving the recursion relation

$$-2(k+1+r)\left(k+\frac{3}{2}+r\right)a_{k+1} + 2a_k(k+r+2)(k+r-1) = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+1} = \frac{2a_k(k+r+2)(k+r-1)}{(k+1+r)(2k+3+2r)}$$

- Recursion relation for $r = 0$; series terminates at $k = 1$

$$a_{k+1} = \frac{2a_k(k+2)(k-1)}{(k+1)(2k+3)}$$

- Apply recursion relation for $k = 0$

$$a_1 = -\frac{4a_0}{3}$$

- Terminating series solution of the ODE for $r = 0$. Use reduction of order to find the second li

$$y = a_0 \cdot \left(1 - \frac{4x}{3} \right)$$

- Recursion relation for $r = -\frac{1}{2}$

$$a_{k+1} = \frac{2a_k\left(k+\frac{3}{2}\right)\left(k-\frac{3}{2}\right)}{\left(k+\frac{1}{2}\right)(2k+2)}$$

- Solution for $r = -\frac{1}{2}$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k-\frac{1}{2}}, a_{k+1} = \frac{2a_k\left(k+\frac{3}{2}\right)\left(k-\frac{3}{2}\right)}{\left(k+\frac{1}{2}\right)(2k+2)} \right]$$

- Combine solutions and rename parameters

$$\left[y = a_0 \cdot \left(1 - \frac{4x}{3} \right) + \left(\sum_{k=0}^{\infty} b_k x^{k-\frac{1}{2}} \right), b_{k+1} = \frac{2b_k\left(k+\frac{3}{2}\right)\left(k-\frac{3}{2}\right)}{\left(k+\frac{1}{2}\right)(2k+2)} \right]$$

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Reducible group (found an exponential solution)
    Reducible group (found another exponential solution)
<- Kovacics algorithm successful`
```

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 40

```
Order:=8;
```

```
dsolve(x*(1-x)*diff(y(x),x$2)+(3/2-2*x)*diff(y(x),x)+2*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \frac{c_1 \left(1 - \frac{9}{2}x + \frac{15}{8}x^2 + \frac{7}{16}x^3 + \frac{27}{128}x^4 + \frac{33}{256}x^5 + \frac{91}{1024}x^6 + \frac{135}{2048}x^7 + O(x^8) \right)}{\sqrt{x}} + c_2 \left(1 - \frac{4}{3}x + O(x^8) \right)$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 71

```
AsymptoticDSolveValue[x*(1-x)*y''[x]+(3/2-2*x)*y'[x]+2*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow \frac{c_2 \left(\frac{135x^7}{2048} + \frac{91x^6}{1024} + \frac{33x^5}{256} + \frac{27x^4}{128} + \frac{7x^3}{16} + \frac{15x^2}{8} - \frac{9x}{2} + 1 \right)}{\sqrt{x}} + c_1 \left(1 - \frac{4x}{3} \right)$$

21.2 problem 2(b)

21.2.1 Maple step by step solution 3516

Internal problem ID [6471]

Internal file name [OUTPUT/5719_Sunday_June_05_2022_03_48_43_PM_20153980/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.6. Gauss's Hypergeometric Equation. Page 187

Problem number: 2(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _exact , _linear , _homogeneous]]
```

$$(2x^2 + 2x)y'' + (5x + 1)y' + y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$(2x^2 + 2x)y'' + (5x + 1)y' + y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{5x + 1}{2x(1 + x)}$$
$$q(x) = \frac{1}{2x(1 + x)}$$

Table 487: Table $p(x), q(x)$ singularities.

$p(x) = \frac{5x+1}{2x(1+x)}$		$q(x) = \frac{1}{2x(1+x)}$	
singularity	type	singularity	type
$x = -1$	“regular”	$x = -1$	“regular”
$x = 0$	“regular”	$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[-1, 0, \infty]$

Irregular singular points : $[\]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$2y''x(1+x) + (5x+1)y' + y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) x(1+x) + (5x+1) \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} 2x^{n+r-1} a_n (n+r) (n+r-1) \right) \\ & + \left(\sum_{n=0}^{\infty} 5x^{n+r} a_n (n+r) \right) + \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r-1$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r-1} and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=0}^{\infty} 2x^{n+r} a_n (n+r) (n+r-1) &= \sum_{n=1}^{\infty} 2a_{n-1} (n+r-1) (n+r-2) x^{n+r-1} \\ \sum_{n=0}^{\infty} 5x^{n+r} a_n (n+r) &= \sum_{n=1}^{\infty} 5a_{n-1} (n+r-1) x^{n+r-1} \\ \sum_{n=0}^{\infty} a_n x^{n+r} &= \sum_{n=1}^{\infty} a_{n-1} x^{n+r-1} \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r-1$.

$$\begin{aligned} & \left(\sum_{n=1}^{\infty} 2a_{n-1} (n+r-1) (n+r-2) x^{n+r-1} \right) \\ & + \left(\sum_{n=0}^{\infty} 2x^{n+r-1} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=1}^{\infty} 5a_{n-1} (n+r-1) x^{n+r-1} \right) \\ & + \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=1}^{\infty} a_{n-1} x^{n+r-1} \right) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$2x^{n+r-1} a_n (n+r) (n+r-1) + (n+r) a_n x^{n+r-1} = 0$$

When $n=0$ the above becomes

$$2x^{-1+r} a_0 r (-1+r) + r a_0 x^{-1+r} = 0$$

Or

$$(2x^{-1+r}r(-1+r) + rx^{-1+r})a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$rx^{-1+r}(2r-1) = 0$$

Since the above is true for all x then the indicial equation becomes

$$2r^2 - r = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = \frac{1}{2}$$
$$r_2 = 0$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$rx^{-1+r}(2r-1) = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = \frac{1}{2}$ is not an integer, then we can construct two linearly independent solutions

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$
$$y_2(x) = x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+\frac{1}{2}}$$
$$y_2(x) = \sum_{n=0}^{\infty} b_n x^n$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$2a_{n-1}(n+r-1)(n+r-2) + 2a_n(n+r)(n+r-1) + 5a_{n-1}(n+r-1) + a_n(n+r) + a_{n-1} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -a_{n-1} \quad (4)$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_n = -a_{n-1} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = \frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = -1$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_1 = -1$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	-1	-1

For $n = 2$, using the above recursive equation gives

$$a_2 = 1$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_2 = 1$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	-1	-1
a_2	1	1

For $n = 3$, using the above recursive equation gives

$$a_3 = -1$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_3 = -1$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	-1	-1
a_2	1	1
a_3	-1	-1

For $n = 4$, using the above recursive equation gives

$$a_4 = 1$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_4 = 1$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	-1	-1
a_2	1	1
a_3	-1	-1
a_4	1	1

For $n = 5$, using the above recursive equation gives

$$a_5 = -1$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_5 = -1$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	-1	-1
a_2	1	1
a_3	-1	-1
a_4	1	1
a_5	-1	-1

For $n = 6$, using the above recursive equation gives

$$a_6 = 1$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_6 = 1$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	-1	-1
a_2	1	1
a_3	-1	-1
a_4	1	1
a_5	-1	-1
a_6	1	1

For $n = 7$, using the above recursive equation gives

$$a_7 = -1$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_7 = -1$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	-1	-1
a_2	1	1
a_3	-1	-1
a_4	1	1
a_5	-1	-1
a_6	1	1
a_7	-1	-1

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= \sqrt{x}(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots) \\ &= \sqrt{x}(1 - x + x^2 - x^3 + x^4 - x^5 + x^6 - x^7 + O(x^8)) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. For $1 \leq n$ the recursive equation is

$$\begin{aligned} 2b_{n-1}(n+r-1)(n+r-2) + 2b_n(n+r)(n+r-1) \\ + 5b_{n-1}(n+r-1) + (n+r)b_n + b_{n-1} = 0 \end{aligned} \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = -b_{n-1} \quad (4)$$

Which for the root $r = 0$ becomes

$$b_n = -b_{n-1} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1

For $n = 1$, using the above recursive equation gives

$$b_1 = -1$$

Which for the root $r = 0$ becomes

$$b_1 = -1$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	-1	-1

For $n = 2$, using the above recursive equation gives

$$b_2 = 1$$

Which for the root $r = 0$ becomes

$$b_2 = 1$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	-1	-1
b_2	1	1

For $n = 3$, using the above recursive equation gives

$$b_3 = -1$$

Which for the root $r = 0$ becomes

$$b_3 = -1$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	-1	-1
b_2	1	1
b_3	-1	-1

For $n = 4$, using the above recursive equation gives

$$b_4 = 1$$

Which for the root $r = 0$ becomes

$$b_4 = 1$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	-1	-1
b_2	1	1
b_3	-1	-1
b_4	1	1

For $n = 5$, using the above recursive equation gives

$$b_5 = -1$$

Which for the root $r = 0$ becomes

$$b_5 = -1$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	-1	-1
b_2	1	1
b_3	-1	-1
b_4	1	1
b_5	-1	-1

For $n = 6$, using the above recursive equation gives

$$b_6 = 1$$

Which for the root $r = 0$ becomes

$$b_6 = 1$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	-1	-1
b_2	1	1
b_3	-1	-1
b_4	1	1
b_5	-1	-1
b_6	1	1

For $n = 7$, using the above recursive equation gives

$$b_7 = -1$$

Which for the root $r = 0$ becomes

$$b_7 = -1$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	-1	-1
b_2	1	1
b_3	-1	-1
b_4	1	1
b_5	-1	-1
b_6	1	1
b_7	-1	-1

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned} y_2(x) &= b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 + b_7x^7 + b_8x^8 \dots \\ &= 1 - x + x^2 - x^3 + x^4 - x^5 + x^6 - x^7 + O(x^8) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1 y_1(x) + c_2 y_2(x) \\ &= c_1 \sqrt{x} (1 - x + x^2 - x^3 + x^4 - x^5 + x^6 - x^7 + O(x^8)) \\ &\quad + c_2 (1 - x + x^2 - x^3 + x^4 - x^5 + x^6 - x^7 + O(x^8)) \end{aligned}$$

Hence the final solution is

$$\begin{aligned} y &= y_h \\ &= c_1 \sqrt{x} (1 - x + x^2 - x^3 + x^4 - x^5 + x^6 - x^7 + O(x^8)) \\ &\quad + c_2 (1 - x + x^2 - x^3 + x^4 - x^5 + x^6 - x^7 + O(x^8)) \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned} y &= c_1 \sqrt{x} (1 - x + x^2 - x^3 + x^4 - x^5 + x^6 - x^7 + O(x^8)) \\ &\quad + c_2 (1 - x + x^2 - x^3 + x^4 - x^5 + x^6 - x^7 + O(x^8)) \end{aligned} \tag{1}$$

Verification of solutions

$$\begin{aligned} y &= c_1 \sqrt{x} (1 - x + x^2 - x^3 + x^4 - x^5 + x^6 - x^7 + O(x^8)) \\ &\quad + c_2 (1 - x + x^2 - x^3 + x^4 - x^5 + x^6 - x^7 + O(x^8)) \end{aligned}$$

Verified OK.

21.2.1 Maple step by step solution

Let's solve

$$2y''x(1+x) + (5x+1)y' + y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{y}{2x(1+x)} - \frac{(5x+1)y'}{2x(1+x)}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{(5x+1)y'}{2x(1+x)} + \frac{y}{2x(1+x)} = 0$$

- Check to see if x_0 is a regular singular point

- Define functions

$$\left[P_2(x) = \frac{5x+1}{2x(1+x)}, P_3(x) = \frac{1}{2x(1+x)} \right]$$

- $(1+x) \cdot P_2(x)$ is analytic at $x = -1$

$$\left. ((1+x) \cdot P_2(x)) \right|_{x=-1} = 2$$

- $(1+x)^2 \cdot P_3(x)$ is analytic at $x = -1$

$$\left. ((1+x)^2 \cdot P_3(x)) \right|_{x=-1} = 0$$

- $x = -1$ is a regular singular point

Check to see if x_0 is a regular singular point

$$x_0 = -1$$

- Multiply by denominators

$$2y''x(1+x) + (5x+1)y' + y = 0$$

- Change variables using $x = u - 1$ so that the regular singular point is at $u = 0$

$$(2u^2 - 2u) \left(\frac{d^2}{du^2} y(u) \right) + (5u - 4) \left(\frac{d}{du} y(u) \right) + y(u) = 0$$

- Assume series solution for $y(u)$

$$y(u) = \sum_{k=0}^{\infty} a_k u^{k+r}$$

- Rewrite ODE with series expansions

- Convert $u^m \cdot \left(\frac{d}{du} y(u) \right)$ to series expansion for $m = 0..1$

$$u^m \cdot \left(\frac{d}{du} y(u) \right) = \sum_{k=0}^{\infty} a_k (k+r) u^{k+r-1+m}$$

- Shift index using $k- > k+1-m$

$$u^m \cdot \left(\frac{d}{du} y(u) \right) = \sum_{k=-1+m}^{\infty} a_{k+1-m} (k+1-m+r) u^{k+r}$$

- Convert $u^m \cdot \left(\frac{d^2}{du^2} y(u) \right)$ to series expansion for $m = 1..2$

$$u^m \cdot \left(\frac{d^2}{du^2} y(u) \right) = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) u^{k+r-2+m}$$

- Shift index using $k- > k+2-m$

$$u^m \cdot \left(\frac{d^2}{du^2} y(u) \right) = \sum_{k=-2+m}^{\infty} a_{k+2-m} (k+2-m+r)(k+1-m+r) u^{k+r}$$

Rewrite ODE with series expansions

$$-2a_0r(1+r)u^{-1+r} + \left(\sum_{k=0}^{\infty} (-2a_{k+1}(k+r+1)(k+2+r) + a_k(k+r+1)(2k+2r+1))u^{k+r} \right)$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$-2r(1+r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \{-1, 0\}$$

- Each term in the series must be 0, giving the recursion relation

$$2(k+r+1)((-k-r-2)a_{k+1} + a_k(k+r+\frac{1}{2})) = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+1} = \frac{a_k(2k+2r+1)}{2(k+2+r)}$$

- Recursion relation for $r = -1$

$$a_{k+1} = \frac{a_k(2k-1)}{2(k+1)}$$

- Solution for $r = -1$

$$\left[y(u) = \sum_{k=0}^{\infty} a_k u^{k-1}, a_{k+1} = \frac{a_k(2k-1)}{2(k+1)} \right]$$

- Revert the change of variables $u = 1 + x$

$$\left[y = \sum_{k=0}^{\infty} a_k (1+x)^{k-1}, a_{k+1} = \frac{a_k(2k-1)}{2(k+1)} \right]$$

- Recursion relation for $r = 0$

$$a_{k+1} = \frac{a_k(2k+1)}{2(k+2)}$$

- Solution for $r = 0$

$$\left[y(u) = \sum_{k=0}^{\infty} a_k u^k, a_{k+1} = \frac{a_k(2k+1)}{2(k+2)} \right]$$

- Revert the change of variables $u = 1 + x$

$$\left[y = \sum_{k=0}^{\infty} a_k (1+x)^k, a_{k+1} = \frac{a_k(2k+1)}{2(k+2)} \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k (1+x)^{k-1} \right) + \left(\sum_{k=0}^{\infty} b_k (1+x)^k \right), a_{k+1} = \frac{a_k(2k-1)}{2(k+1)}, b_{k+1} = \frac{b_k(2k+1)}{2(k+2)} \right]$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
<- linear_1 successful`
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 52

```
Order:=8;  
dsolve((2*x^2+2*x)*diff(y(x),x$2)+(1+5*x)*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (-x^7 + x^6 - x^5 + x^4 - x^3 + x^2 - x + 1) (c_1\sqrt{x} + c_2) + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 73

```
AsymptoticDSolveValue[(2*x^2+2*x)*y'[x]+(1+5*x)*y'[x]+y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1\sqrt{x}(-x^7 + x^6 - x^5 + x^4 - x^3 + x^2 - x + 1) + c_2(-x^7 + x^6 - x^5 + x^4 - x^3 + x^2 - x + 1)$$

21.3 problem 2(x)

21.3.1 Maple step by step solution 3534

Internal problem ID [6472]

Internal file name [OUTPUT/5720_Sunday_June_05_2022_03_48_46_PM_21098881/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.6. Gauss's Hypergeometric Equation. Page 187

Problem number: 2(x).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$(x^2 - 1)y'' + (5x + 4)y' + 4y = 0$$

With the expansion point for the power series method at $x = -1$.

The ode does not have its expansion point at $x = 0$, therefore to simplify the computation of power series expansion, change of variable is made on the independent variable to shift the initial conditions and the expansion point back to zero. The new ode is then solved more easily since the expansion point is now at zero. The solution converted back to the original independent variable. Let

$$t = 1 + x$$

The ode is converted to be in terms of the new independent variable t . This results in

$$((t - 1)^2 - 1) \left(\frac{d^2}{dt^2} y(t) \right) + (-1 + 5t) \left(\frac{d}{dt} y(t) \right) + 4y(t) = 0$$

With its expansion point and initial conditions now at $t = 0$. The transformed ODE is now solved. The type of the expansion point is first determined. This is done on the

homogeneous part of the ODE.

$$(t^2 - 2t) \left(\frac{d^2}{dt^2} y(t) \right) + (-1 + 5t) \left(\frac{d}{dt} y(t) \right) + 4y(t) = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$\frac{d^2}{dt^2} y(t) + p(t) \frac{d}{dt} y(t) + q(t) y(t) = 0$$

Where

$$p(t) = \frac{-1 + 5t}{t(-2 + t)}$$

$$q(t) = \frac{4}{(-2 + t)t}$$

Table 489: Table $p(t), q(t)$ singularities.

$p(t) = \frac{-1+5t}{t(-2+t)}$	
singularity	type
$t = 0$	“regular”
$t = 2$	“regular”

$q(t) = \frac{4}{(-2+t)t}$	
singularity	type
$t = 0$	“regular”
$t = 2$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0, 2, \infty]$

Irregular singular points : $[\]$

Since $t = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$\left(\frac{d^2}{dt^2} y(t) \right) t(-2 + t) + (-1 + 5t) \left(\frac{d}{dt} y(t) \right) + 4y(t) = 0$$

Let the solution be represented as Frobenius power series of the form

$$y(t) = \sum_{n=0}^{\infty} a_n t^{n+r}$$

Then

$$\frac{d}{dt}y(t) = \sum_{n=0}^{\infty} (n+r) a_n t^{n+r-1}$$

$$\frac{d^2}{dt^2}y(t) = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n t^{n+r-2}$$

Substituting the above back into the ode gives

$$\left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n t^{n+r-2} \right) t(-2+t) \tag{1}$$

$$+ (-1+5t) \left(\sum_{n=0}^{\infty} (n+r) a_n t^{n+r-1} \right) + 4 \left(\sum_{n=0}^{\infty} a_n t^{n+r} \right) = 0$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} t^{n+r} a_n (n+r)(n+r-1) \right) + \sum_{n=0}^{\infty} (-2t^{n+r-1} a_n (n+r)(n+r-1)) \tag{2A}$$

$$+ \left(\sum_{n=0}^{\infty} 5t^{n+r} a_n (n+r) \right) + \sum_{n=0}^{\infty} (-(n+r) a_n t^{n+r-1}) + \left(\sum_{n=0}^{\infty} 4a_n t^{n+r} \right) = 0$$

The next step is to make all powers of t be $n+r-1$ in each summation term. Going over each summation term above with power of t in it which is not already t^{n+r-1} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} t^{n+r} a_n (n+r)(n+r-1) = \sum_{n=1}^{\infty} a_{n-1} (n+r-1)(n+r-2) t^{n+r-1}$$

$$\sum_{n=0}^{\infty} 5t^{n+r} a_n (n+r) = \sum_{n=1}^{\infty} 5a_{n-1} (n+r-1) t^{n+r-1}$$

$$\sum_{n=0}^{\infty} 4a_n t^{n+r} = \sum_{n=1}^{\infty} 4a_{n-1} t^{n+r-1}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of t are the same and equal to $n + r - 1$.

$$\begin{aligned} & \left(\sum_{n=1}^{\infty} a_{n-1} (n+r-1) (n+r-2) t^{n+r-1} \right) \\ & + \sum_{n=0}^{\infty} (-2t^{n+r-1} a_n (n+r) (n+r-1)) + \left(\sum_{n=1}^{\infty} 5a_{n-1} (n+r-1) t^{n+r-1} \right) \quad (2B) \\ & + \sum_{n=0}^{\infty} (-(n+r) a_n t^{n+r-1}) + \left(\sum_{n=1}^{\infty} 4a_{n-1} t^{n+r-1} \right) = 0 \end{aligned}$$

The indicial equation is obtained from $n = 0$. From Eq (2B) this gives

$$-2t^{n+r-1} a_n (n+r) (n+r-1) - (n+r) a_n t^{n+r-1} = 0$$

When $n = 0$ the above becomes

$$-2t^{-1+r} a_0 r (-1+r) - r a_0 t^{-1+r} = 0$$

Or

$$(-2t^{-1+r} r (-1+r) - r t^{-1+r}) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(-2r^2 + r) t^{-1+r} = 0$$

Since the above is true for all t then the indicial equation becomes

$$-2r^2 + r = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= \frac{1}{2} \\ r_2 &= 0 \end{aligned}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(-2r^2 + r) t^{-1+r} = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = \frac{1}{2}$ is not an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(t) &= t^{r_1} \left(\sum_{n=0}^{\infty} a_n t^n \right) \\ y_2(t) &= t^{r_2} \left(\sum_{n=0}^{\infty} b_n t^n \right) \end{aligned}$$

Or

$$y_1(t) = \sum_{n=0}^{\infty} a_n t^{n+\frac{1}{2}}$$

$$y_2(t) = \sum_{n=0}^{\infty} b_n t^n$$

We start by finding $y_1(t)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$a_{n-1}(n+r-1)(n+r-2) - 2a_n(n+r)(n+r-1) + 5a_{n-1}(n+r-1) - a_n(n+r) + 4a_{n-1} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-1}(n^2 + 2nr + r^2 + 2n + 2r + 1)}{2n^2 + 4nr + 2r^2 - n - r} \quad (4)$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_n = \frac{a_{n-1}(3 + 2n)^2}{8n^2 + 4n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = \frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = \frac{(r+2)^2}{2r^2 + 3r + 1}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_1 = \frac{25}{12}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{(r+2)^2}{2r^2+3r+1}$	$\frac{25}{12}$

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{(r+3)^2(r+2)}{4r^3+12r^2+11r+3}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_2 = \frac{245}{96}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{(r+2)^2}{2r^2+3r+1}$	$\frac{25}{12}$
a_2	$\frac{(r+3)^2(r+2)}{4r^3+12r^2+11r+3}$	$\frac{245}{96}$

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{(r+4)^2(r+2)(r+3)}{8r^4+44r^3+82r^2+61r+15}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_3 = \frac{315}{128}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{(r+2)^2}{2r^2+3r+1}$	$\frac{25}{12}$
a_2	$\frac{(r+3)^2(r+2)}{4r^3+12r^2+11r+3}$	$\frac{245}{96}$
a_3	$\frac{(r+4)^2(r+2)(r+3)}{8r^4+44r^3+82r^2+61r+15}$	$\frac{315}{128}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{(r+5)^2 (r+3) (r+2) (r+4)}{16r^5 + 144r^4 + 472r^3 + 696r^2 + 457r + 105}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_4 = \frac{4235}{2048}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{(r+2)^2}{2r^2+3r+1}$	$\frac{25}{12}$
a_2	$\frac{(r+3)^2(r+2)}{4r^3+12r^2+11r+3}$	$\frac{245}{96}$
a_3	$\frac{(r+4)^2(r+2)(r+3)}{8r^4+44r^3+82r^2+61r+15}$	$\frac{315}{128}$
a_4	$\frac{(r+5)^2(r+3)(r+2)(r+4)}{16r^5+144r^4+472r^3+696r^2+457r+105}$	$\frac{4235}{2048}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{(r+6)^2 (r+4) (r+2) (r+3) (r+5)}{32r^6 + 432r^5 + 2240r^4 + 5640r^3 + 7178r^2 + 4323r + 945}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_5 = \frac{13013}{8192}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{(r+2)^2}{2r^2+3r+1}$	$\frac{25}{12}$
a_2	$\frac{(r+3)^2(r+2)}{4r^3+12r^2+11r+3}$	$\frac{245}{96}$
a_3	$\frac{(r+4)^2(r+2)(r+3)}{8r^4+44r^3+82r^2+61r+15}$	$\frac{315}{128}$
a_4	$\frac{(r+5)^2(r+3)(r+2)(r+4)}{16r^5+144r^4+472r^3+696r^2+457r+105}$	$\frac{4235}{2048}$
a_5	$\frac{(r+6)^2(r+4)(r+2)(r+3)(r+5)}{32r^6+432r^5+2240r^4+5640r^3+7178r^2+4323r+945}$	$\frac{13013}{8192}$

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{(r+7)^2 (r+5) (r+3) (r+2) (r+4) (r+6)}{64r^7 + 1216r^6 + 9232r^5 + 35920r^4 + 76396r^3 + 87604r^2 + 49443r + 10395}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_6 = \frac{75075}{65536}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{(r+2)^2}{2r^2+3r+1}$	$\frac{25}{12}$
a_2	$\frac{(r+3)^2(r+2)}{4r^3+12r^2+11r+3}$	$\frac{245}{96}$
a_3	$\frac{(r+4)^2(r+2)(r+3)}{8r^4+44r^3+82r^2+61r+15}$	$\frac{315}{128}$
a_4	$\frac{(r+5)^2(r+3)(r+2)(r+4)}{16r^5+144r^4+472r^3+696r^2+457r+105}$	$\frac{4235}{2048}$
a_5	$\frac{(r+6)^2(r+4)(r+2)(r+3)(r+5)}{32r^6+432r^5+2240r^4+5640r^3+7178r^2+4323r+945}$	$\frac{13013}{8192}$
a_6	$\frac{(r+7)^2(r+5)(r+3)(r+2)(r+4)(r+6)}{64r^7+1216r^6+9232r^5+35920r^4+76396r^3+87604r^2+49443r+10395}$	$\frac{75075}{65536}$

For $n = 7$, using the above recursive equation gives

$$a_7 = \frac{(r+8)^2 (r+6) (r+4) (r+2) (r+3) (r+5) (r+7)}{128r^8 + 3264r^7 + 34272r^6 + 191856r^5 + 619752r^4 + 1168356r^3 + 1237738r^2 + 663549r + 135135}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_7 = \frac{206635}{262144}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{(r+2)^2}{2r^2+3r+1}$	$\frac{25}{12}$
a_2	$\frac{(r+3)^2(r+2)}{4r^3+12r^2+11r+3}$	$\frac{245}{96}$
a_3	$\frac{(r+4)^2(r+2)(r+3)}{8r^4+44r^3+82r^2+61r+15}$	$\frac{315}{128}$
a_4	$\frac{(r+5)^2(r+3)(r+2)(r+4)}{16r^5+144r^4+472r^3+696r^2+457r+105}$	$\frac{4235}{2048}$
a_5	$\frac{(r+6)^2(r+4)(r+2)(r+3)(r+5)}{32r^6+432r^5+2240r^4+5640r^3+7178r^2+4323r+945}$	$\frac{13013}{8192}$
a_6	$\frac{(r+7)^2(r+5)(r+3)(r+2)(r+4)(r+6)}{64r^7+1216r^6+9232r^5+35920r^4+76396r^3+87604r^2+49443r+10395}$	$\frac{75075}{65536}$
a_7	$\frac{(r+8)^2(r+6)(r+4)(r+2)(r+3)(r+5)(r+7)}{128r^8+3264r^7+34272r^6+191856r^5+619752r^4+1168356r^3+1237738r^2+663549r+135135}$	$\frac{206635}{262144}$

Using the above table, then the solution $y_1(t)$ is

$$y_1(t) = \sqrt{t}(a_0 + a_1t + a_2t^2 + a_3t^3 + a_4t^4 + a_5t^5 + a_6t^6 + a_7t^7 + a_8t^8 \dots)$$

$$= \sqrt{t} \left(1 + \frac{25t}{12} + \frac{245t^2}{96} + \frac{315t^3}{128} + \frac{4235t^4}{2048} + \frac{13013t^5}{8192} + \frac{75075t^6}{65536} + \frac{206635t^7}{262144} + O(t^8) \right)$$

Now the second solution $y_2(t)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. For $1 \leq n$ the recursive equation is

$$b_{n-1}(n+r-1)(n+r-2) - 2b_n(n+r)(n+r-1) + 5b_{n-1}(n+r-1) - (n+r)b_n + 4b_{n-1} = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = \frac{b_{n-1}(n^2 + 2nr + r^2 + 2n + 2r + 1)}{2n^2 + 4nr + 2r^2 - n - r} \quad (4)$$

Which for the root $r = 0$ becomes

$$b_n = \frac{b_{n-1}(n+1)^2}{2n^2 - n} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1

For $n = 1$, using the above recursive equation gives

$$b_1 = \frac{(r+2)^2}{2r^2+3r+1}$$

Which for the root $r = 0$ becomes

$$b_1 = 4$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{(r+2)^2}{2r^2+3r+1}$	4

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{(r+3)^2(r+2)}{4r^3+12r^2+11r+3}$$

Which for the root $r = 0$ becomes

$$b_2 = 6$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{(r+2)^2}{2r^2+3r+1}$	4
b_2	$\frac{(r+3)^2(r+2)}{4r^3+12r^2+11r+3}$	6

For $n = 3$, using the above recursive equation gives

$$b_3 = \frac{(r+4)^2(r+2)(r+3)}{8r^4+44r^3+82r^2+61r+15}$$

Which for the root $r = 0$ becomes

$$b_3 = \frac{32}{5}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{(r+2)^2}{2r^2+3r+1}$	4
b_2	$\frac{(r+3)^2(r+2)}{4r^3+12r^2+11r+3}$	6
b_3	$\frac{(r+4)^2(r+2)(r+3)}{8r^4+44r^3+82r^2+61r+15}$	$\frac{32}{5}$

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{(r+5)^2(r+3)(r+2)(r+4)}{16r^5 + 144r^4 + 472r^3 + 696r^2 + 457r + 105}$$

Which for the root $r = 0$ becomes

$$b_4 = \frac{40}{7}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{(r+2)^2}{2r^2+3r+1}$	4
b_2	$\frac{(r+3)^2(r+2)}{4r^3+12r^2+11r+3}$	6
b_3	$\frac{(r+4)^2(r+2)(r+3)}{8r^4+44r^3+82r^2+61r+15}$	$\frac{32}{5}$
b_4	$\frac{(r+5)^2(r+3)(r+2)(r+4)}{16r^5+144r^4+472r^3+696r^2+457r+105}$	$\frac{40}{7}$

For $n = 5$, using the above recursive equation gives

$$b_5 = \frac{(r+6)^2(r+4)(r+2)(r+3)(r+5)}{32r^6 + 432r^5 + 2240r^4 + 5640r^3 + 7178r^2 + 4323r + 945}$$

Which for the root $r = 0$ becomes

$$b_5 = \frac{32}{7}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{(r+2)^2}{2r^2+3r+1}$	4
b_2	$\frac{(r+3)^2(r+2)}{4r^3+12r^2+11r+3}$	6
b_3	$\frac{(r+4)^2(r+2)(r+3)}{8r^4+44r^3+82r^2+61r+15}$	$\frac{32}{5}$
b_4	$\frac{(r+5)^2(r+3)(r+2)(r+4)}{16r^5+144r^4+472r^3+696r^2+457r+105}$	$\frac{40}{7}$
b_5	$\frac{(r+6)^2(r+4)(r+2)(r+3)(r+5)}{32r^6+432r^5+2240r^4+5640r^3+7178r^2+4323r+945}$	$\frac{32}{7}$

For $n = 6$, using the above recursive equation gives

$$b_6 = \frac{(r+7)^2(r+5)(r+3)(r+2)(r+4)(r+6)}{64r^7 + 1216r^6 + 9232r^5 + 35920r^4 + 76396r^3 + 87604r^2 + 49443r + 10395}$$

Which for the root $r = 0$ becomes

$$b_6 = \frac{112}{33}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{(r+2)^2}{2r^2+3r+1}$	4
b_2	$\frac{(r+3)^2(r+2)}{4r^3+12r^2+11r+3}$	6
b_3	$\frac{(r+4)^2(r+2)(r+3)}{8r^4+44r^3+82r^2+61r+15}$	$\frac{32}{5}$
b_4	$\frac{(r+5)^2(r+3)(r+2)(r+4)}{16r^5+144r^4+472r^3+696r^2+457r+105}$	$\frac{40}{7}$
b_5	$\frac{(r+6)^2(r+4)(r+2)(r+3)(r+5)}{32r^6+432r^5+2240r^4+5640r^3+7178r^2+4323r+945}$	$\frac{32}{7}$
b_6	$\frac{(r+7)^2(r+5)(r+3)(r+2)(r+4)(r+6)}{64r^7+1216r^6+9232r^5+35920r^4+76396r^3+87604r^2+49443r+10395}$	$\frac{112}{33}$

For $n = 7$, using the above recursive equation gives

$$b_7 = \frac{(r+8)^2(r+6)(r+4)(r+2)(r+3)(r+5)(r+7)}{128r^8 + 3264r^7 + 34272r^6 + 191856r^5 + 619752r^4 + 1168356r^3 + 1237738r^2 + 663549r + 135135}$$

Which for the root $r = 0$ becomes

$$b_7 = \frac{1024}{429}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{(r+2)^2}{2r^2+3r+1}$	4
b_2	$\frac{(r+3)^2(r+2)}{4r^3+12r^2+11r+3}$	6
b_3	$\frac{(r+4)^2(r+2)(r+3)}{8r^4+44r^3+82r^2+61r+15}$	$\frac{32}{5}$
b_4	$\frac{(r+5)^2(r+3)(r+2)(r+4)}{16r^5+144r^4+472r^3+696r^2+457r+105}$	$\frac{40}{7}$
b_5	$\frac{(r+6)^2(r+4)(r+2)(r+3)(r+5)}{32r^6+432r^5+2240r^4+5640r^3+7178r^2+4323r+945}$	$\frac{32}{7}$
b_6	$\frac{(r+7)^2(r+5)(r+3)(r+2)(r+4)(r+6)}{64r^7+1216r^6+9232r^5+35920r^4+76396r^3+87604r^2+49443r+10395}$	$\frac{112}{33}$
b_7	$\frac{(r+8)^2(r+6)(r+4)(r+2)(r+3)(r+5)(r+7)}{128r^8+3264r^7+34272r^6+191856r^5+619752r^4+1168356r^3+1237738r^2+663549r+135135}$	$\frac{1024}{429}$

Using the above table, then the solution $y_2(t)$ is

$$\begin{aligned} y_2(t) &= b_0 + b_1t + b_2t^2 + b_3t^3 + b_4t^4 + b_5t^5 + b_6t^6 + b_7t^7 + b_8t^8 \dots \\ &= 1 + 4t + 6t^2 + \frac{32t^3}{5} + \frac{40t^4}{7} + \frac{32t^5}{7} + \frac{112t^6}{33} + \frac{1024t^7}{429} + O(t^8) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(t) &= c_1y_1(t) + c_2y_2(t) \\ &= c_1\sqrt{t} \left(1 + \frac{25t}{12} + \frac{245t^2}{96} + \frac{315t^3}{128} + \frac{4235t^4}{2048} + \frac{13013t^5}{8192} + \frac{75075t^6}{65536} + \frac{206635t^7}{262144} + O(t^8) \right) \\ &\quad + c_2 \left(1 + 4t + 6t^2 + \frac{32t^3}{5} + \frac{40t^4}{7} + \frac{32t^5}{7} + \frac{112t^6}{33} + \frac{1024t^7}{429} + O(t^8) \right) \end{aligned}$$

Hence the final solution is

$$\begin{aligned} y(t) &= y_h \\ &= c_1\sqrt{t} \left(1 + \frac{25t}{12} + \frac{245t^2}{96} + \frac{315t^3}{128} + \frac{4235t^4}{2048} + \frac{13013t^5}{8192} + \frac{75075t^6}{65536} + \frac{206635t^7}{262144} + O(t^8) \right) \\ &\quad + c_2 \left(1 + 4t + 6t^2 + \frac{32t^3}{5} + \frac{40t^4}{7} + \frac{32t^5}{7} + \frac{112t^6}{33} + \frac{1024t^7}{429} + O(t^8) \right) \end{aligned}$$

Replacing t in the above with the original independent variable x using $t = 1 + x$ results in

$$y = c_1 \sqrt{1+x} \left(\frac{37}{12} + \frac{25x}{12} + \frac{245(1+x)^2}{96} + \frac{315(1+x)^3}{128} + \frac{4235(1+x)^4}{2048} + \frac{13013(1+x)^5}{8192} \right. \\ \left. + \frac{75075(1+x)^6}{65536} + \frac{206635(1+x)^7}{262144} + O((1+x)^8) \right) \\ + c_2 \left(5 + 4x + 6(1+x)^2 + \frac{32(1+x)^3}{5} + \frac{40(1+x)^4}{7} + \frac{32(1+x)^5}{7} + \frac{112(1+x)^6}{33} \right. \\ \left. + \frac{1024(1+x)^7}{429} + O((1+x)^8) \right)$$

Summary

The solution(s) found are the following

$$y = c_1 \sqrt{1+x} \left(\frac{37}{12} + \frac{25x}{12} + \frac{245(1+x)^2}{96} + \frac{315(1+x)^3}{128} + \frac{4235(1+x)^4}{2048} \right. \\ \left. + \frac{13013(1+x)^5}{8192} + \frac{75075(1+x)^6}{65536} + \frac{206635(1+x)^7}{262144} + O((1+x)^8) \right) \\ + c_2 \left(5 + 4x + 6(1+x)^2 + \frac{32(1+x)^3}{5} + \frac{40(1+x)^4}{7} + \frac{32(1+x)^5}{7} \right. \\ \left. + \frac{112(1+x)^6}{33} + \frac{1024(1+x)^7}{429} + O((1+x)^8) \right) \quad (1)$$

Verification of solutions

$$y = c_1 \sqrt{1+x} \left(\frac{37}{12} + \frac{25x}{12} + \frac{245(1+x)^2}{96} + \frac{315(1+x)^3}{128} + \frac{4235(1+x)^4}{2048} \right. \\ \left. + \frac{13013(1+x)^5}{8192} + \frac{75075(1+x)^6}{65536} + \frac{206635(1+x)^7}{262144} + O((1+x)^8) \right) \\ + c_2 \left(5 + 4x + 6(1+x)^2 + \frac{32(1+x)^3}{5} + \frac{40(1+x)^4}{7} + \frac{32(1+x)^5}{7} + \frac{112(1+x)^6}{33} \right. \\ \left. + \frac{1024(1+x)^7}{429} + O((1+x)^8) \right)$$

Verified OK.

21.3.1 Maple step by step solution

Let's solve

$$(x^2 - 1)y'' + (5x + 4)y' + 4y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{(5x+4)y'}{x^2-1} - \frac{4y}{x^2-1}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{(5x+4)y'}{x^2-1} + \frac{4y}{x^2-1} = 0$$

- Check to see if x_0 is a regular singular point

- Define functions

$$[P_2(x) = \frac{5x+4}{x^2-1}, P_3(x) = \frac{4}{x^2-1}]$$

- $(1+x) \cdot P_2(x)$ is analytic at $x = -1$

$$((1+x) \cdot P_2(x)) \Big|_{x=-1} = \frac{1}{2}$$

- $(1+x)^2 \cdot P_3(x)$ is analytic at $x = -1$

$$((1+x)^2 \cdot P_3(x)) \Big|_{x=-1} = 0$$

- $x = -1$ is a regular singular point

Check to see if x_0 is a regular singular point

$$x_0 = -1$$

- Multiply by denominators

$$(x^2 - 1)y'' + (5x + 4)y' + 4y = 0$$

- Change variables using $x = u - 1$ so that the regular singular point is at $u = 0$

$$(u^2 - 2u) \left(\frac{d^2}{du^2} y(u) \right) + (5u - 1) \left(\frac{d}{du} y(u) \right) + 4y(u) = 0$$

- Assume series solution for $y(u)$

$$y(u) = \sum_{k=0}^{\infty} a_k u^{k+r}$$

- Rewrite ODE with series expansions

- Convert $u^m \cdot \left(\frac{d}{du} y(u) \right)$ to series expansion for $m = 0..1$

$$u^m \cdot \left(\frac{d}{du}y(u)\right) = \sum_{k=0}^{\infty} a_k(k+r) u^{k+r-1+m}$$

- Shift index using $k \rightarrow k+1-m$

$$u^m \cdot \left(\frac{d}{du}y(u)\right) = \sum_{k=-1+m}^{\infty} a_{k+1-m}(k+1-m+r) u^{k+r}$$

- Convert $u^m \cdot \left(\frac{d^2}{du^2}y(u)\right)$ to series expansion for $m = 1..2$

$$u^m \cdot \left(\frac{d^2}{du^2}y(u)\right) = \sum_{k=0}^{\infty} a_k(k+r)(k+r-1) u^{k+r-2+m}$$

- Shift index using $k \rightarrow k+2-m$

$$u^m \cdot \left(\frac{d^2}{du^2}y(u)\right) = \sum_{k=-2+m}^{\infty} a_{k+2-m}(k+2-m+r)(k+1-m+r) u^{k+r}$$

Rewrite ODE with series expansions

$$-a_0r(-1+2r)u^{-1+r} + \left(\sum_{k=0}^{\infty} (-a_{k+1}(k+1+r)(2k+1+2r) + a_k(k+r+2)^2) u^{k+r}\right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$-r(-1+2r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \left\{0, \frac{1}{2}\right\}$$

- Each term in the series must be 0, giving the recursion relation

$$a_k(k+r+2)^2 - 2(k+r+\frac{1}{2})a_{k+1}(k+1+r) = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+1} = \frac{a_k(k+r+2)^2}{(2k+1+2r)(k+1+r)}$$

- Recursion relation for $r = 0$

$$a_{k+1} = \frac{a_k(k+2)^2}{(2k+1)(k+1)}$$

- Solution for $r = 0$

$$\left[y(u) = \sum_{k=0}^{\infty} a_k u^k, a_{k+1} = \frac{a_k(k+2)^2}{(2k+1)(k+1)} \right]$$

- Revert the change of variables $u = 1+x$

$$\left[y = \sum_{k=0}^{\infty} a_k (1+x)^k, a_{k+1} = \frac{a_k(k+2)^2}{(2k+1)(k+1)} \right]$$

- Recursion relation for $r = \frac{1}{2}$

$$a_{k+1} = \frac{a_k (k + \frac{5}{2})^2}{(2k+2)(k + \frac{3}{2})}$$

- Solution for $r = \frac{1}{2}$

$$\left[y(u) = \sum_{k=0}^{\infty} a_k u^{k+\frac{1}{2}}, a_{k+1} = \frac{a_k (k + \frac{5}{2})^2}{(2k+2)(k + \frac{3}{2})} \right]$$

- Revert the change of variables $u = 1 + x$

$$\left[y = \sum_{k=0}^{\infty} a_k (1+x)^{k+\frac{1}{2}}, a_{k+1} = \frac{a_k (k + \frac{5}{2})^2}{(2k+2)(k + \frac{3}{2})} \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k (1+x)^k \right) + \left(\sum_{k=0}^{\infty} b_k (1+x)^{k+\frac{1}{2}} \right), a_{k+1} = \frac{a_k (k+2)^2}{(2k+1)(k+1)}, b_{k+1} = \frac{b_k (k + \frac{5}{2})^2}{(2k+2)(k + \frac{3}{2})} \right]$$

Maple trace Kovacic algorithm successful

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Reducible group (found an exponential solution)
    Group is reducible, not completely reducible
<- Kovacics algorithm successful`

```

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 54

Order:=8;

```
dsolve((x^2-1)*diff(y(x),x$2)+(5*x+4)*diff(y(x),x)+4*y(x)=0,y(x),type='series',x=-1);
```

$$y(x) = c_1 \sqrt{x+1} \left(1 + \frac{25}{12}(x+1) + \frac{245}{96}(x+1)^2 + \frac{315}{128}(x+1)^3 + \frac{4235}{2048}(x+1)^4 \right. \\ \left. + \frac{13013}{8192}(x+1)^5 + \frac{75075}{65536}(x+1)^6 + \frac{206635}{262144}(x+1)^7 + O((x+1)^8) \right) \\ + c_2 \left(1 + 4(x+1) + 6(x+1)^2 + \frac{32}{5}(x+1)^3 + \frac{40}{7}(x+1)^4 + \frac{32}{7}(x+1)^5 \right. \\ \left. + \frac{112}{33}(x+1)^6 + \frac{1024}{429}(x+1)^7 + O((x+1)^8) \right)$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 139

```
AsymptoticDSolveValue[(x^2-1)*y''[x]+(5*x+4)*y'[x]+4*y[x]==0,y[x],{x,-1,7}]
```

$$y(x) \rightarrow c_1 \sqrt{x+1} \left(\frac{206635(x+1)^7}{262144} + \frac{75075(x+1)^6}{65536} + \frac{13013(x+1)^5}{8192} + \frac{4235(x+1)^4}{2048} \right. \\ \left. + \frac{315}{128}(x+1)^3 + \frac{245}{96}(x+1)^2 + \frac{25(x+1)}{12} + 1 \right) + c_2 \left(\frac{1024}{429}(x+1)^7 + \frac{112}{33}(x+1)^6 \right. \\ \left. + \frac{32}{7}(x+1)^5 + \frac{40}{7}(x+1)^4 + \frac{32}{5}(x+1)^3 + 6(x+1)^2 + 4(x+1) + 1 \right)$$

21.4 problem 2(d)

21.4.1 Maple step by step solution 3551

Internal problem ID [6473]

Internal file name [OUTPUT/5721_Sunday_June_05_2022_03_48_51_PM_38879378/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.6. Gauss's Hypergeometric Equation. Page 187

Problem number: 2(d).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _exact , _linear , _homogeneous]]
```

$$(x^2 - x - 6)y'' + (3x + 5)y' + y = 0$$

With the expansion point for the power series method at $x = 3$.

The ode does not have its expansion point at $x = 0$, therefore to simplify the computation of power series expansion, change of variable is made on the independent variable to shift the initial conditions and the expansion point back to zero. The new ode is then solved more easily since the expansion point is now at zero. The solution converted back to the original independent variable. Let

$$t = x - 3$$

The ode is converted to be in terms of the new independent variable t . This results in

$$((t + 3)^2 - t - 9) \left(\frac{d^2}{dt^2} y(t) \right) + (3t + 14) \left(\frac{d}{dt} y(t) \right) + y(t) = 0$$

With its expansion point and initial conditions now at $t = 0$. The transformed ODE is now solved. The type of the expansion point is first determined. This is done on the

homogeneous part of the ODE.

$$(t^2 + 5t) \left(\frac{d^2}{dt^2} y(t) \right) + (3t + 14) \left(\frac{d}{dt} y(t) \right) + y(t) = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$\frac{d^2}{dt^2} y(t) + p(t) \frac{d}{dt} y(t) + q(t) y(t) = 0$$

Where

$$p(t) = \frac{3t + 14}{t(t + 5)}$$

$$q(t) = \frac{1}{(t + 5)t}$$

Table 491: Table $p(t), q(t)$ singularities.

$p(t) = \frac{3t+14}{t(t+5)}$	
singularity	type
$t = -5$	“regular”
$t = 0$	“regular”

$q(t) = \frac{1}{(t+5)t}$	
singularity	type
$t = -5$	“regular”
$t = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[-5, 0, \infty]$

Irregular singular points : $[\]$

Since $t = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$\left(\frac{d^2}{dt^2} y(t) \right) t(t + 5) + (3t + 14) \left(\frac{d}{dt} y(t) \right) + y(t) = 0$$

Let the solution be represented as Frobenius power series of the form

$$y(t) = \sum_{n=0}^{\infty} a_n t^{n+r}$$

Then

$$\frac{d}{dt}y(t) = \sum_{n=0}^{\infty} (n+r) a_n t^{n+r-1}$$

$$\frac{d^2}{dt^2}y(t) = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n t^{n+r-2}$$

Substituting the above back into the ode gives

$$\left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n t^{n+r-2} \right) t(t+5) \tag{1}$$

$$+ (3t+14) \left(\sum_{n=0}^{\infty} (n+r) a_n t^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} a_n t^{n+r} \right) = 0$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} t^{n+r} a_n (n+r)(n+r-1) \right) + \left(\sum_{n=0}^{\infty} 5t^{n+r-1} a_n (n+r)(n+r-1) \right) \tag{2A}$$

$$+ \left(\sum_{n=0}^{\infty} 3t^{n+r} a_n (n+r) \right) + \left(\sum_{n=0}^{\infty} 14(n+r) a_n t^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} a_n t^{n+r} \right) = 0$$

The next step is to make all powers of t be $n+r-1$ in each summation term. Going over each summation term above with power of t in it which is not already t^{n+r-1} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} t^{n+r} a_n (n+r)(n+r-1) = \sum_{n=1}^{\infty} a_{n-1} (n+r-1)(n+r-2) t^{n+r-1}$$

$$\sum_{n=0}^{\infty} 3t^{n+r} a_n (n+r) = \sum_{n=1}^{\infty} 3a_{n-1} (n+r-1) t^{n+r-1}$$

$$\sum_{n=0}^{\infty} a_n t^{n+r} = \sum_{n=1}^{\infty} a_{n-1} t^{n+r-1}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of t are the same and equal to $n + r - 1$.

$$\begin{aligned} & \left(\sum_{n=1}^{\infty} a_{n-1} (n+r-1) (n+r-2) t^{n+r-1} \right) \\ & + \left(\sum_{n=0}^{\infty} 5t^{n+r-1} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=1}^{\infty} 3a_{n-1} (n+r-1) t^{n+r-1} \right) \quad (2B) \\ & + \left(\sum_{n=0}^{\infty} 14(n+r) a_n t^{n+r-1} \right) + \left(\sum_{n=1}^{\infty} a_{n-1} t^{n+r-1} \right) = 0 \end{aligned}$$

The indicial equation is obtained from $n = 0$. From Eq (2B) this gives

$$5t^{n+r-1} a_n (n+r) (n+r-1) + 14(n+r) a_n t^{n+r-1} = 0$$

When $n = 0$ the above becomes

$$5t^{-1+r} a_0 r (-1+r) + 14r a_0 t^{-1+r} = 0$$

Or

$$(5t^{-1+r} r (-1+r) + 14r t^{-1+r}) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$r t^{-1+r} (9 + 5r) = 0$$

Since the above is true for all t then the indicial equation becomes

$$5r^2 + 9r = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 0 \\ r_2 &= -\frac{9}{5} \end{aligned}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$r t^{-1+r} (9 + 5r) = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = \frac{9}{5}$ is not an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(t) &= t^{r_1} \left(\sum_{n=0}^{\infty} a_n t^n \right) \\ y_2(t) &= t^{r_2} \left(\sum_{n=0}^{\infty} b_n t^n \right) \end{aligned}$$

Or

$$y_1(t) = \sum_{n=0}^{\infty} a_n t^n$$

$$y_2(t) = \sum_{n=0}^{\infty} b_n t^{n - \frac{9}{5}}$$

We start by finding $y_1(t)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$a_{n-1}(n+r-1)(n+r-2) + 5a_n(n+r)(n+r-1) + 3a_{n-1}(n+r-1) + 14a_n(n+r) + a_{n-1} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{(n+r)a_{n-1}}{5n+9+5r} \quad (4)$$

Which for the root $r = 0$ becomes

$$a_n = -\frac{na_{n-1}}{5n+9} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = \frac{-1-r}{14+5r}$$

Which for the root $r = 0$ becomes

$$a_1 = -\frac{1}{14}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-1-r}{14+5r}$	$-\frac{1}{14}$

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{r^2 + 3r + 2}{25r^2 + 165r + 266}$$

Which for the root $r = 0$ becomes

$$a_2 = \frac{1}{133}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-1-r}{14+5r}$	$-\frac{1}{14}$
a_2	$\frac{r^2+3r+2}{25r^2+165r+266}$	$\frac{1}{133}$

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{-r^3 - 6r^2 - 11r - 6}{125r^3 + 1425r^2 + 5290r + 6384}$$

Which for the root $r = 0$ becomes

$$a_3 = -\frac{1}{1064}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-1-r}{14+5r}$	$-\frac{1}{14}$
a_2	$\frac{r^2+3r+2}{25r^2+165r+266}$	$\frac{1}{133}$
a_3	$\frac{-r^3-6r^2-11r-6}{125r^3+1425r^2+5290r+6384}$	$-\frac{1}{1064}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{r^4 + 10r^3 + 35r^2 + 50r + 24}{625r^4 + 10750r^3 + 67775r^2 + 185330r + 185136}$$

Which for the root $r = 0$ becomes

$$a_4 = \frac{1}{7714}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-1-r}{14+5r}$	$-\frac{1}{14}$
a_2	$\frac{r^2+3r+2}{25r^2+165r+266}$	$\frac{1}{133}$
a_3	$\frac{-r^3-6r^2-11r-6}{125r^3+1425r^2+5290r+6384}$	$-\frac{1}{1064}$
a_4	$\frac{r^4+10r^3+35r^2+50r+24}{625r^4+10750r^3+67775r^2+185330r+185136}$	$\frac{1}{7714}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{-r^5 - 15r^4 - 85r^3 - 225r^2 - 274r - 120}{3125r^5 + 75000r^4 + 704375r^3 + 3231000r^2 + 7226900r + 6294624}$$

Which for the root $r = 0$ becomes

$$a_5 = -\frac{5}{262276}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-1-r}{14+5r}$	$-\frac{1}{14}$
a_2	$\frac{r^2+3r+2}{25r^2+165r+266}$	$\frac{1}{133}$
a_3	$\frac{-r^3-6r^2-11r-6}{125r^3+1425r^2+5290r+6384}$	$-\frac{1}{1064}$
a_4	$\frac{r^4+10r^3+35r^2+50r+24}{625r^4+10750r^3+67775r^2+185330r+185136}$	$\frac{1}{7714}$
a_5	$\frac{-r^5-15r^4-85r^3-225r^2-274r-120}{3125r^5+75000r^4+704375r^3+3231000r^2+7226900r+6294624}$	$-\frac{5}{262276}$

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{r^6 + 21r^5 + 175r^4 + 735r^3 + 1624r^2 + 1764r + 720}{15625r^6 + 496875r^5 + 6446875r^4 + 43625625r^3 + 162143500r^2 + 313322220r + 245490336}$$

Which for the root $r = 0$ becomes

$$a_6 = \frac{5}{1704794}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-1-r}{14+5r}$	$-\frac{1}{14}$
a_2	$\frac{r^2+3r+2}{25r^2+165r+266}$	$\frac{1}{133}$
a_3	$\frac{-r^3-6r^2-11r-6}{125r^3+1425r^2+5290r+6384}$	$-\frac{1}{1064}$
a_4	$\frac{r^4+10r^3+35r^2+50r+24}{625r^4+10750r^3+67775r^2+185330r+185136}$	$\frac{1}{7714}$
a_5	$\frac{-r^5-15r^4-85r^3-225r^2-274r-120}{3125r^5+75000r^4+704375r^3+3231000r^2+7226900r+6294624}$	$-\frac{5}{262276}$
a_6	$\frac{r^6+21r^5+175r^4+735r^3+1624r^2+1764r+720}{15625r^6+496875r^5+6446875r^4+43625625r^3+162143500r^2+313322220r+245490336}$	$\frac{5}{1704794}$

For $n = 7$, using the above recursive equation gives

$$a_7 = \frac{-r^7 - 28r^6 - 322r^5 - 1960r^4 - 6769r^3 - 13132r^2 - 13068r - 5040}{78125r^7 + 3171875r^6 + 54096875r^5 + 501790625r^4 + 2730245000r^3 + 8700925100r^2 + 15013629360r + 10801574784}$$

Which for the root $r = 0$ becomes

$$a_7 = -\frac{5}{10715848}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-1-r}{14+5r}$	$-\frac{1}{14}$
a_2	$\frac{r^2+3r+2}{25r^2+165r+266}$	$\frac{1}{133}$
a_3	$\frac{-r^3-6r^2-11r-6}{125r^3+1425r^2+5290r+6384}$	$-\frac{1}{1064}$
a_4	$\frac{r^4+10r^3+35r^2+50r+24}{625r^4+10750r^3+67775r^2+185330r+185136}$	$\frac{1}{7714}$
a_5	$\frac{-r^5-15r^4-85r^3-225r^2-274r-120}{3125r^5+75000r^4+704375r^3+3231000r^2+7226900r+6294624}$	$-\frac{5}{262276}$
a_6	$\frac{r^6+21r^5+175r^4+735r^3+1624r^2+1764r+720}{15625r^6+496875r^5+6446875r^4+43625625r^3+162143500r^2+313322220r+245490336}$	$\frac{5}{1704794}$
a_7	$\frac{-r^7-28r^6-322r^5-1960r^4-6769r^3-13132r^2-13068r-5040}{78125r^7+3171875r^6+54096875r^5+501790625r^4+2730245000r^3+8700925100r^2+15013629360r+10801574784}$	$-\frac{5}{10715848}$

Using the above table, then the solution $y_1(t)$ is

$$\begin{aligned} y_1(t) &= a_0 + a_1t + a_2t^2 + a_3t^3 + a_4t^4 + a_5t^5 + a_6t^6 + a_7t^7 + a_8t^8 \dots \\ &= 1 - \frac{t}{14} + \frac{t^2}{133} - \frac{t^3}{1064} + \frac{t^4}{7714} - \frac{5t^5}{262276} + \frac{5t^6}{1704794} - \frac{5t^7}{10715848} + O(t^8) \end{aligned}$$

Now the second solution $y_2(t)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. For $1 \leq n$ the recursive equation is

$$\begin{aligned} b_{n-1}(n+r-1)(n+r-2) + 5b_n(n+r)(n+r-1) \\ + 3b_{n-1}(n+r-1) + 14(n+r)b_n + b_{n-1} = 0 \end{aligned} \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = -\frac{(n+r)b_{n-1}}{5n+9+5r} \quad (4)$$

Which for the root $r = -\frac{9}{5}$ becomes

$$b_n = -\frac{(5n-9)b_{n-1}}{25n} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = -\frac{9}{5}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1

For $n = 1$, using the above recursive equation gives

$$b_1 = \frac{-1-r}{14+5r}$$

Which for the root $r = -\frac{9}{5}$ becomes

$$b_1 = \frac{4}{25}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{-1-r}{14+5r}$	$\frac{4}{25}$

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{r^2 + 3r + 2}{25r^2 + 165r + 266}$$

Which for the root $r = -\frac{9}{5}$ becomes

$$b_2 = -\frac{2}{625}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{-1-r}{14+5r}$	$\frac{4}{25}$
b_2	$\frac{r^2+3r+2}{25r^2+165r+266}$	$-\frac{2}{625}$

For $n = 3$, using the above recursive equation gives

$$b_3 = \frac{-r^3 - 6r^2 - 11r - 6}{125r^3 + 1425r^2 + 5290r + 6384}$$

Which for the root $r = -\frac{9}{5}$ becomes

$$b_3 = \frac{4}{15625}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{-1-r}{14+5r}$	$\frac{4}{25}$
b_2	$\frac{r^2+3r+2}{25r^2+165r+266}$	$-\frac{2}{625}$
b_3	$\frac{-r^3-6r^2-11r-6}{125r^3+1425r^2+5290r+6384}$	$\frac{4}{15625}$

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{r^4 + 10r^3 + 35r^2 + 50r + 24}{625r^4 + 10750r^3 + 67775r^2 + 185330r + 185136}$$

Which for the root $r = -\frac{9}{5}$ becomes

$$b_4 = -\frac{11}{390625}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{-1-r}{14+5r}$	$\frac{4}{25}$
b_2	$\frac{r^2+3r+2}{25r^2+165r+266}$	$-\frac{2}{625}$
b_3	$\frac{-r^3-6r^2-11r-6}{125r^3+1425r^2+5290r+6384}$	$\frac{4}{15625}$
b_4	$\frac{r^4+10r^3+35r^2+50r+24}{625r^4+10750r^3+67775r^2+185330r+185136}$	$-\frac{11}{390625}$

For $n = 5$, using the above recursive equation gives

$$b_5 = \frac{-r^5 - 15r^4 - 85r^3 - 225r^2 - 274r - 120}{3125r^5 + 75000r^4 + 704375r^3 + 3231000r^2 + 7226900r + 6294624}$$

Which for the root $r = -\frac{9}{5}$ becomes

$$b_5 = \frac{176}{48828125}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{-1-r}{14+5r}$	$\frac{4}{25}$
b_2	$\frac{r^2+3r+2}{25r^2+165r+266}$	$-\frac{2}{625}$
b_3	$\frac{-r^3-6r^2-11r-6}{125r^3+1425r^2+5290r+6384}$	$\frac{4}{15625}$
b_4	$\frac{r^4+10r^3+35r^2+50r+24}{625r^4+10750r^3+67775r^2+185330r+185136}$	$-\frac{11}{390625}$
b_5	$\frac{-r^5-15r^4-85r^3-225r^2-274r-120}{3125r^5+75000r^4+704375r^3+3231000r^2+7226900r+6294624}$	$\frac{176}{48828125}$

For $n = 6$, using the above recursive equation gives

$$b_6 = \frac{r^6 + 21r^5 + 175r^4 + 735r^3 + 1624r^2 + 1764r + 720}{15625r^6 + 496875r^5 + 6446875r^4 + 43625625r^3 + 162143500r^2 + 313322220r + 245490336}$$

Which for the root $r = -\frac{9}{5}$ becomes

$$b_6 = -\frac{616}{1220703125}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{-1-r}{14+5r}$	$\frac{4}{25}$
b_2	$\frac{r^2+3r+2}{25r^2+165r+266}$	$-\frac{2}{625}$
b_3	$\frac{-r^3-6r^2-11r-6}{125r^3+1425r^2+5290r+6384}$	$\frac{4}{15625}$
b_4	$\frac{r^4+10r^3+35r^2+50r+24}{625r^4+10750r^3+67775r^2+185330r+185136}$	$-\frac{11}{390625}$
b_5	$\frac{-r^5-15r^4-85r^3-225r^2-274r-120}{3125r^5+75000r^4+704375r^3+3231000r^2+7226900r+6294624}$	$\frac{176}{48828125}$
b_6	$\frac{r^6+21r^5+175r^4+735r^3+1624r^2+1764r+720}{15625r^6+496875r^5+6446875r^4+43625625r^3+162143500r^2+313322220r+245490336}$	$-\frac{616}{1220703125}$

For $n = 7$, using the above recursive equation gives

$$b_7 = \frac{-r^7 - 28r^6 - 322r^5 - 1960r^4 - 6769r^3 - 13132r^2 - 13068r - 5040}{78125r^7 + 3171875r^6 + 54096875r^5 + 501790625r^4 + 2730245000r^3 + 8700925100r^2 + 15013629360r + 10801574784}$$

Which for the root $r = -\frac{9}{5}$ becomes

$$b_7 = \frac{2288}{30517578125}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{-1-r}{14+5r}$	$\frac{4}{25}$
b_2	$\frac{r^2+3r+2}{25r^2+165r+266}$	$-\frac{2}{625}$
b_3	$\frac{-r^3-6r^2-11r-6}{125r^3+1425r^2+5290r+6384}$	$\frac{4}{15625}$
b_4	$\frac{r^4+10r^3+35r^2+50r+24}{625r^4+10750r^3+67775r^2+185330r+185136}$	$-\frac{11}{390625}$
b_5	$\frac{-r^5-15r^4-85r^3-225r^2-274r-120}{3125r^5+75000r^4+704375r^3+3231000r^2+7226900r+6294624}$	$\frac{176}{48828125}$
b_6	$\frac{r^6+21r^5+175r^4+735r^3+1624r^2+1764r+720}{15625r^6+496875r^5+6446875r^4+43625625r^3+162143500r^2+313322220r+245490336}$	$-\frac{616}{1220703125}$
b_7	$\frac{-r^7-28r^6-322r^5-1960r^4-6769r^3-13132r^2-13068r-5040}{78125r^7+3171875r^6+54096875r^5+501790625r^4+2730245000r^3+8700925100r^2+15013629360r+10801574784}$	$\frac{2288}{30517578125}$

Using the above table, then the solution $y_2(t)$ is

$$y_2(t) = 1(b_0 + b_1t + b_2t^2 + b_3t^3 + b_4t^4 + b_5t^5 + b_6t^6 + b_7t^7 + b_8t^8 \dots)$$

$$= \frac{1 + \frac{4t}{25} - \frac{2t^2}{625} + \frac{4t^3}{15625} - \frac{11t^4}{390625} + \frac{176t^5}{48828125} - \frac{616t^6}{1220703125} + \frac{2288t^7}{30517578125} + O(t^8)}{t^{\frac{9}{5}}}$$

Therefore the homogeneous solution is

$$y_h(t) = c_1y_1(t) + c_2y_2(t)$$

$$= c_1 \left(1 - \frac{t}{14} + \frac{t^2}{133} - \frac{t^3}{1064} + \frac{t^4}{7714} - \frac{5t^5}{262276} + \frac{5t^6}{1704794} - \frac{5t^7}{10715848} + O(t^8) \right)$$

$$+ \frac{c_2 \left(1 + \frac{4t}{25} - \frac{2t^2}{625} + \frac{4t^3}{15625} - \frac{11t^4}{390625} + \frac{176t^5}{48828125} - \frac{616t^6}{1220703125} + \frac{2288t^7}{30517578125} + O(t^8) \right)}{t^{\frac{9}{5}}}$$

Hence the final solution is

$$y(t) = y_h$$

$$= c_1 \left(1 - \frac{t}{14} + \frac{t^2}{133} - \frac{t^3}{1064} + \frac{t^4}{7714} - \frac{5t^5}{262276} + \frac{5t^6}{1704794} - \frac{5t^7}{10715848} + O(t^8) \right)$$

$$+ \frac{c_2 \left(1 + \frac{4t}{25} - \frac{2t^2}{625} + \frac{4t^3}{15625} - \frac{11t^4}{390625} + \frac{176t^5}{48828125} - \frac{616t^6}{1220703125} + \frac{2288t^7}{30517578125} + O(t^8) \right)}{t^{\frac{9}{5}}}$$

Replacing t in the above with the original independent variable x using $t = x - 3$ results in

$$y = c_1 \left(\frac{17}{14} - \frac{x}{14} + \frac{(x-3)^2}{133} - \frac{(x-3)^3}{1064} + \frac{(x-3)^4}{7714} - \frac{5(x-3)^5}{262276} + \frac{5(x-3)^6}{1704794} \right.$$

$$\left. - \frac{5(x-3)^7}{10715848} + O((x-3)^8) \right)$$

$$+ \frac{c_2 \left(\frac{13}{25} + \frac{4x}{25} - \frac{2(x-3)^2}{625} + \frac{4(x-3)^3}{15625} - \frac{11(x-3)^4}{390625} + \frac{176(x-3)^5}{48828125} - \frac{616(x-3)^6}{1220703125} + \frac{2288(x-3)^7}{30517578125} + O((x-3)^8) \right)}{(x-3)^{\frac{9}{5}}}$$

Summary

The solution(s) found are the following

$$y = c_1 \left(\frac{17}{14} - \frac{x}{14} + \frac{(x-3)^2}{133} - \frac{(x-3)^3}{1064} + \frac{(x-3)^4}{7714} - \frac{5(x-3)^5}{262276} + \frac{5(x-3)^6}{1704794} - \frac{5(x-3)^7}{10715848} + O((x-3)^8) \right) + \frac{c_2 \left(\frac{13}{25} + \frac{4x}{25} - \frac{2(x-3)^2}{625} + \frac{4(x-3)^3}{15625} - \frac{11(x-3)^4}{390625} + \frac{176(x-3)^5}{48828125} - \frac{616(x-3)^6}{1220703125} + \frac{2288(x-3)^7}{30517578125} + O((x-3)^8) \right)}{(x-3)^{\frac{9}{5}}} \quad (1)$$

Verification of solutions

$$y = c_1 \left(\frac{17}{14} - \frac{x}{14} + \frac{(x-3)^2}{133} - \frac{(x-3)^3}{1064} + \frac{(x-3)^4}{7714} - \frac{5(x-3)^5}{262276} + \frac{5(x-3)^6}{1704794} - \frac{5(x-3)^7}{10715848} + O((x-3)^8) \right) + \frac{c_2 \left(\frac{13}{25} + \frac{4x}{25} - \frac{2(x-3)^2}{625} + \frac{4(x-3)^3}{15625} - \frac{11(x-3)^4}{390625} + \frac{176(x-3)^5}{48828125} - \frac{616(x-3)^6}{1220703125} + \frac{2288(x-3)^7}{30517578125} + O((x-3)^8) \right)}{(x-3)^{\frac{9}{5}}}$$

Verified OK.

21.4.1 Maple step by step solution

Let's solve

$$(x^2 - x - 6)y'' + (3x + 5)y' + y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{y}{x^2-x-6} - \frac{(3x+5)y'}{x^2-x-6}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{(3x+5)y'}{x^2-x-6} + \frac{y}{x^2-x-6} = 0$$

- Check to see if x_0 is a regular singular point

- Define functions

$$\left[P_2(x) = \frac{3x+5}{x^2-x-6}, P_3(x) = \frac{1}{x^2-x-6} \right]$$

- $(x + 2) \cdot P_2(x)$ is analytic at $x = -2$

$$\left. ((x + 2) \cdot P_2(x)) \right|_{x=-2} = \frac{1}{5}$$

- $(x + 2)^2 \cdot P_3(x)$ is analytic at $x = -2$

$$\left. ((x + 2)^2 \cdot P_3(x)) \right|_{x=-2} = 0$$

- $x = -2$ is a regular singular point

Check to see if x_0 is a regular singular point

$$x_0 = -2$$

- Multiply by denominators

$$(x^2 - x - 6) y'' + (3x + 5) y' + y = 0$$

- Change variables using $x = u - 2$ so that the regular singular point is at $u = 0$

$$(u^2 - 5u) \left(\frac{d^2}{du^2} y(u) \right) + (3u - 1) \left(\frac{d}{du} y(u) \right) + y(u) = 0$$

- Assume series solution for $y(u)$

$$y(u) = \sum_{k=0}^{\infty} a_k u^{k+r}$$

- Rewrite ODE with series expansions

- Convert $u^m \cdot \left(\frac{d}{du} y(u) \right)$ to series expansion for $m = 0..1$

$$u^m \cdot \left(\frac{d}{du} y(u) \right) = \sum_{k=0}^{\infty} a_k (k + r) u^{k+r-1+m}$$

- Shift index using $k- > k + 1 - m$

$$u^m \cdot \left(\frac{d}{du} y(u) \right) = \sum_{k=-1+m}^{\infty} a_{k+1-m} (k + 1 - m + r) u^{k+r}$$

- Convert $u^m \cdot \left(\frac{d^2}{du^2} y(u) \right)$ to series expansion for $m = 1..2$

$$u^m \cdot \left(\frac{d^2}{du^2} y(u) \right) = \sum_{k=0}^{\infty} a_k (k + r) (k + r - 1) u^{k+r-2+m}$$

- Shift index using $k- > k + 2 - m$

$$u^m \cdot \left(\frac{d^2}{du^2} y(u) \right) = \sum_{k=-2+m}^{\infty} a_{k+2-m} (k + 2 - m + r) (k + 1 - m + r) u^{k+r}$$

Rewrite ODE with series expansions

$$-a_0 r(-4 + 5r) u^{-1+r} + \left(\sum_{k=0}^{\infty} (-a_{k+1} (k + r + 1) (5k + 1 + 5r) + a_k (k + r + 1)^2) u^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation
 $-r(-4 + 5r) = 0$
- Values of r that satisfy the indicial equation
 $r \in \{0, \frac{4}{5}\}$
- Each term in the series must be 0, giving the recursion relation
 $((-5k - 5r - 1)a_{k+1} + a_k(k + r + 1))(k + r + 1) = 0$
- Recursion relation that defines series solution to ODE
 $a_{k+1} = \frac{a_k(k+r+1)}{5k+1+5r}$
- Recursion relation for $r = 0$
 $a_{k+1} = \frac{a_k(k+1)}{5k+1}$
- Solution for $r = 0$
 $\left[y(u) = \sum_{k=0}^{\infty} a_k u^k, a_{k+1} = \frac{a_k(k+1)}{5k+1} \right]$
- Revert the change of variables $u = x + 2$
 $\left[y = \sum_{k=0}^{\infty} a_k (x + 2)^k, a_{k+1} = \frac{a_k(k+1)}{5k+1} \right]$
- Recursion relation for $r = \frac{4}{5}$
 $a_{k+1} = \frac{a_k(k+\frac{9}{5})}{5k+5}$
- Solution for $r = \frac{4}{5}$
 $\left[y(u) = \sum_{k=0}^{\infty} a_k u^{k+\frac{4}{5}}, a_{k+1} = \frac{a_k(k+\frac{9}{5})}{5k+5} \right]$
- Revert the change of variables $u = x + 2$
 $\left[y = \sum_{k=0}^{\infty} a_k (x + 2)^{k+\frac{4}{5}}, a_{k+1} = \frac{a_k(k+\frac{9}{5})}{5k+5} \right]$
- Combine solutions and rename parameters
 $\left[y = \left(\sum_{k=0}^{\infty} a_k (x + 2)^k \right) + \left(\sum_{k=0}^{\infty} b_k (x + 2)^{k+\frac{4}{5}} \right), a_{k+1} = \frac{a_k(k+1)}{5k+1}, b_{k+1} = \frac{b_k(k+\frac{9}{5})}{5k+5} \right]$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
  One independent solution has integrals. Trying a hypergeometric solution free of integral
  -> hyper3: Equivalence to 2F1, 1F1 or 0F1 under a power @ Moebius
  <- hyper3 successful: received ODE is equivalent to the 2F1 ODE
  -> Trying to convert hypergeometric functions to elementary form...
  <- elementary form for at least one hypergeometric solution is achieved - returning with
<- linear_1 successful`
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 54

```
Order:=8;
dsolve((x^2-x-6)*diff(y(x),x$2)+(5+3*x)*diff(y(x),x)+y(x)=0,y(x),type='series',x=3);
```

$$y(x) = \frac{c_1 \left(1 + \frac{4}{25}(x-3) - \frac{2}{625}(x-3)^2 + \frac{4}{15625}(x-3)^3 - \frac{11}{390625}(x-3)^4 + \frac{176}{48828125}(x-3)^5 - \frac{616}{1220703125}(x-3)^6 - \dots \right)}{(x-3)^{\frac{9}{5}}} + c_2 \left(1 - \frac{1}{14}(x-3) + \frac{1}{133}(x-3)^2 - \frac{1}{1064}(x-3)^3 + \frac{1}{7714}(x-3)^4 - \frac{5}{262276}(x-3)^5 + \frac{5}{1704794}(x-3)^6 - \frac{5}{10715848}(x-3)^7 + O((x-3)^8) \right)$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 145

```
AsymptoticDSolveValue[(x^2-x-6)*y'[x]+(5+3*x)*y'[x]+y[x]==0,y[x],{x,3,7}]
```

$$y(x) \rightarrow c_1 \left(-\frac{5(x-3)^7}{10715848} + \frac{5(x-3)^6}{1704794} - \frac{5(x-3)^5}{262276} + \frac{(x-3)^4}{7714} - \frac{(x-3)^3}{1064} + \frac{1}{133}(x-3)^2 + \frac{3-x}{14} + 1 \right) + \frac{c_2 \left(\frac{2288(x-3)^7}{30517578125} - \frac{616(x-3)^6}{1220703125} + \frac{176(x-3)^5}{48828125} - \frac{11(x-3)^4}{390625} + \frac{4(x-3)^3}{15625} - \frac{2}{625}(x-3)^2 + \frac{4(x-3)}{25} + 1 \right)}{(x-3)^{9/5}}$$

21.5 problem 3

21.5.1 Maple step by step solution 3573

Internal problem ID [6474]

Internal file name [OUTPUT/5722_Sunday_June_05_2022_03_49_05_PM_56011628/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.6. Gauss's Hypergeometric Equation. Page 187

Problem number: 3.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[_Gegenbauer , [_2nd_order , _linear , ` _with_symmetry_ [0,F(x)] `]]
```

$$(-x^2 + 1)y'' - xy' + p^2y = 0$$

With the expansion point for the power series method at $x = 1$.

The ode does not have its expansion point at $x = 0$, therefore to simplify the computation of power series expansion, change of variable is made on the independent variable to shift the initial conditions and the expansion point back to zero. The new ode is then solved more easily since the expansion point is now at zero. The solution converted back to the original independent variable. Let

$$t = x - 1$$

The ode is converted to be in terms of the new independent variable t . This results in

$$(-(t + 1)^2 + 1) \left(\frac{d^2}{dt^2} y(t) \right) - (t + 1) \left(\frac{d}{dt} y(t) \right) + p^2 y(t) = 0$$

With its expansion point and initial conditions now at $t = 0$. The transformed ODE is now solved. The type of the expansion point is first determined. This is done on the

homogeneous part of the ODE.

$$(-t^2 - 2t) \left(\frac{d^2}{dt^2} y(t) \right) + (-t - 1) \left(\frac{d}{dt} y(t) \right) + p^2 y(t) = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$\frac{d^2}{dt^2} y(t) + p(t) \frac{d}{dt} y(t) + q(t) y(t) = 0$$

Where

$$p(t) = \frac{t+1}{t(t+2)}$$

$$q(t) = -\frac{p^2}{t(t+2)}$$

Table 493: Table $p(t), q(t)$ singularities.

$p(t) = \frac{t+1}{t(t+2)}$		$q(t) = -\frac{p^2}{t(t+2)}$	
singularity	type	singularity	type
$t = -2$	“regular”	$t = -2$	“regular”
$t = 0$	“regular”	$t = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[-2, 0, \infty]$

Irregular singular points : $[\]$

Since $t = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$-\left(\frac{d^2}{dt^2} y(t) \right) t(t+2) + (-t-1) \left(\frac{d}{dt} y(t) \right) + p^2 y(t) = 0$$

Let the solution be represented as Frobenius power series of the form

$$y(t) = \sum_{n=0}^{\infty} a_n t^{n+r}$$

Then

$$\frac{d}{dt}y(t) = \sum_{n=0}^{\infty} (n+r) a_n t^{n+r-1}$$

$$\frac{d^2}{dt^2}y(t) = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n t^{n+r-2}$$

Substituting the above back into the ode gives

$$-\left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n t^{n+r-2}\right) t(t+2) \tag{1}$$

$$+ (-t-1) \left(\sum_{n=0}^{\infty} (n+r) a_n t^{n+r-1}\right) + p^2 \left(\sum_{n=0}^{\infty} a_n t^{n+r}\right) = 0$$

Which simplifies to

$$\sum_{n=0}^{\infty} (-t^{n+r} a_n (n+r)(n+r-1)) + \sum_{n=0}^{\infty} (-2t^{n+r-1} a_n (n+r)(n+r-1)) \tag{2A}$$

$$+ \sum_{n=0}^{\infty} (-t^{n+r} a_n (n+r)) + \sum_{n=0}^{\infty} (-(n+r) a_n t^{n+r-1}) + \left(\sum_{n=0}^{\infty} p^2 a_n t^{n+r}\right) = 0$$

The next step is to make all powers of t be $n+r-1$ in each summation term. Going over each summation term above with power of t in it which is not already t^{n+r-1} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} (-t^{n+r} a_n (n+r)(n+r-1)) = \sum_{n=1}^{\infty} (-a_{n-1} (n+r-1)(n+r-2) t^{n+r-1})$$

$$\sum_{n=0}^{\infty} (-t^{n+r} a_n (n+r)) = \sum_{n=1}^{\infty} (-a_{n-1} (n+r-1) t^{n+r-1})$$

$$\sum_{n=0}^{\infty} p^2 a_n t^{n+r} = \sum_{n=1}^{\infty} a_{n-1} p^2 t^{n+r-1}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of t are the same and equal to $n + r - 1$.

$$\begin{aligned} & \sum_{n=1}^{\infty} (-a_{n-1}(n+r-1)(n+r-2)t^{n+r-1}) \\ & + \sum_{n=0}^{\infty} (-2t^{n+r-1}a_n(n+r)(n+r-1)) + \sum_{n=1}^{\infty} (-a_{n-1}(n+r-1)t^{n+r-1}) \quad (2B) \\ & + \sum_{n=0}^{\infty} (-(n+r)a_n t^{n+r-1}) + \left(\sum_{n=1}^{\infty} a_{n-1} p^2 t^{n+r-1} \right) = 0 \end{aligned}$$

The indicial equation is obtained from $n = 0$. From Eq (2B) this gives

$$-2t^{n+r-1}a_n(n+r)(n+r-1) - (n+r)a_n t^{n+r-1} = 0$$

When $n = 0$ the above becomes

$$-2t^{-1+r}a_0r(-1+r) - ra_0t^{-1+r} = 0$$

Or

$$(-2t^{-1+r}r(-1+r) - r t^{-1+r}) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(-2r^2 + r) t^{-1+r} = 0$$

Since the above is true for all t then the indicial equation becomes

$$-2r^2 + r = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= \frac{1}{2} \\ r_2 &= 0 \end{aligned}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(-2r^2 + r) t^{-1+r} = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = \frac{1}{2}$ is not an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(t) &= t^{r_1} \left(\sum_{n=0}^{\infty} a_n t^n \right) \\ y_2(t) &= t^{r_2} \left(\sum_{n=0}^{\infty} b_n t^n \right) \end{aligned}$$

Or

$$y_1(t) = \sum_{n=0}^{\infty} a_n t^{n+\frac{1}{2}}$$

$$y_2(t) = \sum_{n=0}^{\infty} b_n t^n$$

We start by finding $y_1(t)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$\begin{aligned} -a_{n-1}(n+r-1)(n+r-2) - 2a_n(n+r)(n+r-1) \\ - a_{n-1}(n+r-1) - a_n(n+r) + a_{n-1}p^2 = 0 \end{aligned} \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{a_{n-1}(n^2 + 2nr - p^2 + r^2 - 2n - 2r + 1)}{2n^2 + 4nr + 2r^2 - n - r} \quad (4)$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_n = -\frac{a_{n-1}(4n^2 - 4p^2 - 4n + 1)}{8n^2 + 4n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = \frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = \frac{p^2 - r^2}{2r^2 + 3r + 1}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_1 = \frac{p^2}{3} - \frac{1}{12}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{p^2-r^2}{2r^2+3r+1}$	$\frac{p^2}{3} - \frac{1}{12}$

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{(p^2 - r^2 - 2r - 1)(p^2 - r^2)}{4r^4 + 20r^3 + 35r^2 + 25r + 6}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_2 = \frac{1}{30}p^4 - \frac{1}{12}p^2 + \frac{3}{160}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{p^2-r^2}{2r^2+3r+1}$	$\frac{p^2}{3} - \frac{1}{12}$
a_2	$\frac{(p^2-r^2-2r-1)(p^2-r^2)}{4r^4+20r^3+35r^2+25r+6}$	$\frac{1}{30}p^4 - \frac{1}{12}p^2 + \frac{3}{160}$

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{(p^2 - r^2 - 4r - 4)(p^2 - r^2 - 2r - 1)(p^2 - r^2)}{8r^6 + 84r^5 + 350r^4 + 735r^3 + 812r^2 + 441r + 90}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_3 = \frac{1}{630}p^6 - \frac{1}{72}p^4 + \frac{37}{1440}p^2 - \frac{5}{896}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{p^2-r^2}{2r^2+3r+1}$	$\frac{p^2}{3} - \frac{1}{12}$
a_2	$\frac{(p^2-r^2-2r-1)(p^2-r^2)}{4r^4+20r^3+35r^2+25r+6}$	$\frac{1}{30}p^4 - \frac{1}{12}p^2 + \frac{3}{160}$
a_3	$\frac{(p^2-r^2-4r-4)(p^2-r^2-2r-1)(p^2-r^2)}{8r^6+84r^5+350r^4+735r^3+812r^2+441r+90}$	$\frac{1}{630}p^6 - \frac{1}{72}p^4 + \frac{37}{1440}p^2 - \frac{5}{896}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{(p^2 - r^2 - 6r - 9)(p^2 - r^2 - 4r - 4)(p^2 - r^2 - 2r - 1)(p^2 - r^2)}{(8r^6 + 84r^5 + 350r^4 + 735r^3 + 812r^2 + 441r + 90)(2r + 7)(r + 4)}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_4 = \frac{(4p^2 - 49)(4p^2 - 25)(4p^2 - 9)(4p^2 - 1)}{5806080}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{p^2 - r^2}{2r^2 + 3r + 1}$	$\frac{p^2}{3} - \frac{1}{12}$
a_2	$\frac{(p^2 - r^2 - 2r - 1)(p^2 - r^2)}{4r^4 + 20r^3 + 35r^2 + 25r + 6}$	$\frac{1}{30}p^4 - \frac{1}{12}p^2 + \frac{3}{160}$
a_3	$\frac{(p^2 - r^2 - 4r - 4)(p^2 - r^2 - 2r - 1)(p^2 - r^2)}{8r^6 + 84r^5 + 350r^4 + 735r^3 + 812r^2 + 441r + 90}$	$\frac{1}{630}p^6 - \frac{1}{72}p^4 + \frac{37}{1440}p^2 - \frac{5}{896}$
a_4	$\frac{(p^2 - r^2 - 6r - 9)(p^2 - r^2 - 4r - 4)(p^2 - r^2 - 2r - 1)(p^2 - r^2)}{(8r^6 + 84r^5 + 350r^4 + 735r^3 + 812r^2 + 441r + 90)(2r + 7)(r + 4)}$	$\frac{(4p^2 - 49)(4p^2 - 25)(4p^2 - 9)(4p^2 - 1)}{5806080}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{(p^2 - r^2 - 8r - 16)(p^2 - r^2 - 6r - 9)(p^2 - r^2 - 4r - 4)(p^2 - r^2 - 2r - 1)(p^2 - r^2)}{(8r^6 + 84r^5 + 350r^4 + 735r^3 + 812r^2 + 441r + 90)(2r + 7)(r + 4)(r + 5)(2r + 9)}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_5 = \frac{(4p^2 - 81)(4p^2 - 49)(4p^2 - 25)(4p^2 - 9)(4p^2 - 1)}{1277337600}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{p^2 - r^2}{2r^2 + 3r + 1}$	$\frac{p^2}{3} - \frac{1}{12}$
a_2	$\frac{(p^2 - r^2 - 2r - 1)(p^2 - r^2)}{4r^4 + 20r^3 + 35r^2 + 25r + 6}$	$\frac{1}{30}p^4 - \frac{1}{12}p^2 + \frac{3}{160}$
a_3	$\frac{(p^2 - r^2 - 4r - 4)(p^2 - r^2 - 2r - 1)(p^2 - r^2)}{8r^6 + 84r^5 + 350r^4 + 735r^3 + 812r^2 + 441r + 90}$	$\frac{1}{630}p^6 - \frac{1}{72}p^4 + \frac{37}{1440}p^2 - \frac{5}{896}$
a_4	$\frac{(p^2 - r^2 - 6r - 9)(p^2 - r^2 - 4r - 4)(p^2 - r^2 - 2r - 1)(p^2 - r^2)}{(8r^6 + 84r^5 + 350r^4 + 735r^3 + 812r^2 + 441r + 90)(2r + 7)(r + 4)}$	$\frac{(4p^2 - 49)(4p^2 - 25)(4p^2 - 9)(4p^2 - 1)}{5806080}$
a_5	$\frac{(p^2 - r^2 - 8r - 16)(p^2 - r^2 - 6r - 9)(p^2 - r^2 - 4r - 4)(p^2 - r^2 - 2r - 1)(p^2 - r^2)}{(8r^6 + 84r^5 + 350r^4 + 735r^3 + 812r^2 + 441r + 90)(2r + 7)(r + 4)(r + 5)(2r + 9)}$	$\frac{(4p^2 - 81)(4p^2 - 49)(4p^2 - 25)(4p^2 - 9)(4p^2 - 1)}{1277337600}$

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{(p^2 - r^2 - 10r - 25)(p^2 - r^2 - 8r - 16)(p^2 - r^2 - 6r - 9)(p^2 - r^2 - 4r - 4)(p^2 - r^2 - 2r - 1)(p^2 - r^2 - r^2 - 2r - 1)(p^2 - r^2 - r^2 - 2r - 1)}{(8r^6 + 84r^5 + 350r^4 + 735r^3 + 812r^2 + 441r + 90)(2r + 7)(r + 4)(r + 5)(2r + 9)(2r + 11)(6 + r)}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_6 = \frac{(4p^2 - 121)(4p^2 - 81)(4p^2 - 49)(4p^2 - 25)(4p^2 - 9)(4p^2 - 1)}{398529331200}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{p^2 - r^2}{2r^2 + 3r + 1}$	$\frac{p^2}{3} - \frac{1}{12}$
a_2	$\frac{(p^2 - r^2 - 2r - 1)(p^2 - r^2)}{4r^4 + 20r^3 + 35r^2 + 25r + 6}$	$\frac{1}{30}p^4 - \frac{1}{12}p^2 + \frac{3}{160}$
a_3	$\frac{(p^2 - r^2 - 4r - 4)(p^2 - r^2 - 2r - 1)(p^2 - r^2)}{8r^6 + 84r^5 + 350r^4 + 735r^3 + 812r^2 + 441r + 90}$	$\frac{1}{630}p^6 - \frac{1}{72}p^4 + \frac{37}{1440}p^2 - \frac{5}{896}$
a_4	$\frac{(p^2 - r^2 - 6r - 9)(p^2 - r^2 - 4r - 4)(p^2 - r^2 - 2r - 1)(p^2 - r^2)}{(8r^6 + 84r^5 + 350r^4 + 735r^3 + 812r^2 + 441r + 90)(2r + 7)(r + 4)}$	$\frac{(4p^2 - 49)(4p^2 - 25)(4p^2 - 9)(4p^2 - 1)}{5806080}$
a_5	$\frac{(p^2 - r^2 - 8r - 16)(p^2 - r^2 - 6r - 9)(p^2 - r^2 - 4r - 4)(p^2 - r^2 - 2r - 1)(p^2 - r^2)}{(8r^6 + 84r^5 + 350r^4 + 735r^3 + 812r^2 + 441r + 90)(2r + 7)(r + 4)(r + 5)(2r + 9)}$	$\frac{(4p^2 - 81)(4p^2 - 49)(4p^2 - 25)(4p^2 - 9)(4p^2 - 1)}{1277337600}$
a_6	$\frac{(p^2 - r^2 - 10r - 25)(p^2 - r^2 - 8r - 16)(p^2 - r^2 - 6r - 9)(p^2 - r^2 - 4r - 4)(p^2 - r^2 - 2r - 1)(p^2 - r^2)}{(8r^6 + 84r^5 + 350r^4 + 735r^3 + 812r^2 + 441r + 90)(2r + 7)(r + 4)(r + 5)(2r + 9)(2r + 11)(6 + r)}$	$\frac{(4p^2 - 121)(4p^2 - 81)(4p^2 - 49)(4p^2 - 25)(4p^2 - 9)(4p^2 - 1)}{398529331200}$

For $n = 7$, using the above recursive equation gives

$$a_7 = \frac{(p^2 - r^2 - 12r - 36)(p^2 - r^2 - 10r - 25)(p^2 - r^2 - 8r - 16)(p^2 - r^2 - 6r - 9)(p^2 - r^2 - 4r - 4)(p^2 - r^2 - 2r - 1)(p^2 - r^2 - 2r - 1)(p^2 - r^2 - 2r - 1)}{(8r^6 + 84r^5 + 350r^4 + 735r^3 + 812r^2 + 441r + 90)(2r + 7)(r + 4)(r + 5)(2r + 9)(2r + 11)(6 + r)}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_7 = \frac{(4p^2 - 169)(4p^2 - 121)(4p^2 - 81)(4p^2 - 49)(4p^2 - 25)(4p^2 - 9)(4p^2 - 1)}{167382319104000}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{p^2-r^2}{2r^2+3r+1}$	$\frac{p^2}{3} - \frac{1}{12}$
a_2	$\frac{(p^2-r^2-2r-1)(p^2-r^2)}{4r^4+20r^3+35r^2+25r+6}$	$\frac{1}{30}p^4 - \frac{1}{12}p^2 + \frac{3}{160}$
a_3	$\frac{(p^2-r^2-4r-4)(p^2-r^2-2r-1)(p^2-r^2)}{8r^6+84r^5+350r^4+735r^3+812r^2+441r+90}$	$\frac{1}{630}p^6 - \frac{1}{72}p^4 + \frac{37}{144}$
a_4	$\frac{(p^2-r^2-6r-9)(p^2-r^2-4r-4)(p^2-r^2-2r-1)(p^2-r^2)}{(8r^6+84r^5+350r^4+735r^3+812r^2+441r+90)(2r+7)(r+4)}$	$\frac{(4p^2-49)(4p^2-25)(4p^2)}{5806080}$
a_5	$\frac{(p^2-r^2-8r-16)(p^2-r^2-6r-9)(p^2-r^2-4r-4)(p^2-r^2-2r-1)(p^2-r^2)}{(8r^6+84r^5+350r^4+735r^3+812r^2+441r+90)(2r+7)(r+4)(r+5)(2r+9)}$	$\frac{(4p^2-81)(4p^2-49)(4p^2)}{127733}$
a_6	$\frac{(p^2-r^2-10r-25)(p^2-r^2-8r-16)(p^2-r^2-6r-9)(p^2-r^2-4r-4)(p^2-r^2-2r-1)(p^2-r^2)}{(8r^6+84r^5+350r^4+735r^3+812r^2+441r+90)(2r+7)(r+4)(r+5)(2r+9)(2r+11)(6+r)}$	$\frac{(4p^2-121)(4p^2-81)(4p^2)}{39}$
a_7	$\frac{(p^2-r^2-12r-36)(p^2-r^2-10r-25)(p^2-r^2-8r-16)(p^2-r^2-6r-9)(p^2-r^2-4r-4)(p^2-r^2-2r-1)(p^2-r^2)}{(8r^6+84r^5+350r^4+735r^3+812r^2+441r+90)(2r+7)(r+4)(r+5)(2r+9)(2r+11)(6+r)(7+r)(2r+13)}$	$\frac{(4p^2-169)(4p^2-121)(4p^2)}{39}$

Using the above table, then the solution $y_1(t)$ is

$$\begin{aligned}
y_1(t) &= \sqrt{t}(a_0 + a_1t + a_2t^2 + a_3t^3 + a_4t^4 + a_5t^5 + a_6t^6 + a_7t^7 + a_8t^8 \dots) \\
&= \sqrt{t} \left(1 + \left(\frac{p^2}{3} - \frac{1}{12} \right) t + \left(\frac{1}{30}p^4 - \frac{1}{12}p^2 + \frac{3}{160} \right) t^2 + \left(\frac{1}{630}p^6 - \frac{1}{72}p^4 + \frac{37}{1440}p^2 - \frac{5}{896} \right) t^3 + \frac{(4p^2)}{39} t^4 + \dots \right)
\end{aligned}$$

Now the second solution $y_2(t)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. For $1 \leq n$ the recursive equation is

$$\begin{aligned}
&-b_{n-1}(n+r-1)(n+r-2) - 2b_n(n+r)(n+r-1) \\
&-b_{n-1}(n+r-1) - (n+r)b_n + b_{n-1}p^2 = 0
\end{aligned} \tag{3}$$

Solving for b_n from recursive equation (4) gives

$$b_n = -\frac{b_{n-1}(n^2 + 2nr - p^2 + r^2 - 2n - 2r + 1)}{2n^2 + 4nr + 2r^2 - n - r} \tag{4}$$

Which for the root $r = 0$ becomes

$$b_n = -\frac{b_{n-1}(n^2 - p^2 - 2n + 1)}{n(2n - 1)} \tag{5}$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1

For $n = 1$, using the above recursive equation gives

$$b_1 = \frac{p^2 - r^2}{2r^2 + 3r + 1}$$

Which for the root $r = 0$ becomes

$$b_1 = p^2$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{p^2 - r^2}{2r^2 + 3r + 1}$	p^2

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{(p^2 - r^2 - 2r - 1)(p^2 - r^2)}{4r^4 + 20r^3 + 35r^2 + 25r + 6}$$

Which for the root $r = 0$ becomes

$$b_2 = \frac{1}{6}p^4 - \frac{1}{6}p^2$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{p^2 - r^2}{2r^2 + 3r + 1}$	p^2
b_2	$\frac{(p^2 - r^2 - 2r - 1)(p^2 - r^2)}{4r^4 + 20r^3 + 35r^2 + 25r + 6}$	$\frac{1}{6}p^4 - \frac{1}{6}p^2$

For $n = 3$, using the above recursive equation gives

$$b_3 = \frac{(p^2 - r^2 - 4r - 4)(p^2 - r^2 - 2r - 1)(p^2 - r^2)}{8r^6 + 84r^5 + 350r^4 + 735r^3 + 812r^2 + 441r + 90}$$

Which for the root $r = 0$ becomes

$$b_3 = \frac{p^2(p^4 - 5p^2 + 4)}{90}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{p^2-r^2}{2r^2+3r+1}$	p^2
b_2	$\frac{(p^2-r^2-2r-1)(p^2-r^2)}{4r^4+20r^3+35r^2+25r+6}$	$\frac{1}{6}p^4 - \frac{1}{6}p^2$
b_3	$\frac{(p^2-r^2-4r-4)(p^2-r^2-2r-1)(p^2-r^2)}{8r^6+84r^5+350r^4+735r^3+812r^2+441r+90}$	$\frac{p^2(p^4-5p^2+4)}{90}$

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{(p^2 - r^2 - 6r - 9)(p^2 - r^2 - 4r - 4)(p^2 - r^2 - 2r - 1)(p^2 - r^2)}{(8r^6 + 84r^5 + 350r^4 + 735r^3 + 812r^2 + 441r + 90)(2r + 7)(r + 4)}$$

Which for the root $r = 0$ becomes

$$b_4 = \frac{p^2(p^6 - 14p^4 + 49p^2 - 36)}{2520}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{p^2-r^2}{2r^2+3r+1}$	p^2
b_2	$\frac{(p^2-r^2-2r-1)(p^2-r^2)}{4r^4+20r^3+35r^2+25r+6}$	$\frac{1}{6}p^4 - \frac{1}{6}p^2$
b_3	$\frac{(p^2-r^2-4r-4)(p^2-r^2-2r-1)(p^2-r^2)}{8r^6+84r^5+350r^4+735r^3+812r^2+441r+90}$	$\frac{p^2(p^4-5p^2+4)}{90}$
b_4	$\frac{(p^2-r^2-6r-9)(p^2-r^2-4r-4)(p^2-r^2-2r-1)(p^2-r^2)}{(8r^6+84r^5+350r^4+735r^3+812r^2+441r+90)(2r+7)(r+4)}$	$\frac{p^2(p^6-14p^4+49p^2-36)}{2520}$

For $n = 5$, using the above recursive equation gives

$$b_5 = \frac{(p^2 - r^2 - 8r - 16)(p^2 - r^2 - 6r - 9)(p^2 - r^2 - 4r - 4)(p^2 - r^2 - 2r - 1)(p^2 - r^2)}{(8r^6 + 84r^5 + 350r^4 + 735r^3 + 812r^2 + 441r + 90)(2r + 7)(r + 4)(r + 5)(2r + 9)}$$

Which for the root $r = 0$ becomes

$$b_5 = \frac{p^2(p^8 - 30p^6 + 273p^4 - 820p^2 + 576)}{113400}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{p^2-r^2}{2r^2+3r+1}$	p^2
b_2	$\frac{(p^2-r^2-2r-1)(p^2-r^2)}{4r^4+20r^3+35r^2+25r+6}$	$\frac{1}{6}p^4 - \frac{1}{6}p^2$
b_3	$\frac{(p^2-r^2-4r-4)(p^2-r^2-2r-1)(p^2-r^2)}{8r^6+84r^5+350r^4+735r^3+812r^2+441r+90}$	$\frac{p^2(p^4-5p^2+4)}{90}$
b_4	$\frac{(p^2-r^2-6r-9)(p^2-r^2-4r-4)(p^2-r^2-2r-1)(p^2-r^2)}{(8r^6+84r^5+350r^4+735r^3+812r^2+441r+90)(2r+7)(r+4)}$	$\frac{p^2(p^6-14p^4+49p^2-36)}{2520}$
b_5	$\frac{(p^2-r^2-8r-16)(p^2-r^2-6r-9)(p^2-r^2-4r-4)(p^2-r^2-2r-1)(p^2-r^2)}{(8r^6+84r^5+350r^4+735r^3+812r^2+441r+90)(2r+7)(r+4)(r+5)(2r+9)}$	$\frac{p^2(p^8-30p^6+273p^4-820p^2+576)}{113400}$

For $n = 6$, using the above recursive equation gives

$$b_6 = \frac{(p^2 - r^2 - 10r - 25)(p^2 - r^2 - 8r - 16)(p^2 - r^2 - 6r - 9)(p^2 - r^2 - 4r - 4)(p^2 - r^2 - 2r - 1)(p^2 - r^2)}{(8r^6 + 84r^5 + 350r^4 + 735r^3 + 812r^2 + 441r + 90)(2r + 7)(r + 4)(r + 5)(2r + 9)(2r + 11)(6 + r)}$$

Which for the root $r = 0$ becomes

$$b_6 = \frac{(p^2 - 25)(p^2 - 16)(p^2 - 9)(p^2 - 4)(p^2 - 1)p^2}{7484400}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{p^2-r^2}{2r^2+3r+1}$	p^2
b_2	$\frac{(p^2-r^2-2r-1)(p^2-r^2)}{4r^4+20r^3+35r^2+25r+6}$	$\frac{1}{6}p^4 - \frac{1}{6}p^2$
b_3	$\frac{(p^2-r^2-4r-4)(p^2-r^2-2r-1)(p^2-r^2)}{8r^6+84r^5+350r^4+735r^3+812r^2+441r+90}$	$\frac{p^2(p^4-5p^2+4)}{90}$
b_4	$\frac{(p^2-r^2-6r-9)(p^2-r^2-4r-4)(p^2-r^2-2r-1)(p^2-r^2)}{(8r^6+84r^5+350r^4+735r^3+812r^2+441r+90)(2r+7)(r+4)}$	$\frac{p^2(p^6-14p^4+49p^2-36)}{2520}$
b_5	$\frac{(p^2-r^2-8r-16)(p^2-r^2-6r-9)(p^2-r^2-4r-4)(p^2-r^2-2r-1)(p^2-r^2)}{(8r^6+84r^5+350r^4+735r^3+812r^2+441r+90)(2r+7)(r+4)(r+5)(2r+9)}$	$\frac{p^2(p^8-30p^6+273p^4-820p^2+576)}{113400}$
b_6	$\frac{(p^2-r^2-10r-25)(p^2-r^2-8r-16)(p^2-r^2-6r-9)(p^2-r^2-4r-4)(p^2-r^2-2r-1)(p^2-r^2)}{(8r^6+84r^5+350r^4+735r^3+812r^2+441r+90)(2r+7)(r+4)(r+5)(2r+9)(2r+11)(6+r)}$	$\frac{(p^2-25)(p^2-16)(p^2-9)(p^2-4)(p^2-1)p^2}{7484400}$

For $n = 7$, using the above recursive equation gives

$$b_7 = \frac{(p^2 - r^2 - 12r - 36)(p^2 - r^2 - 10r - 25)(p^2 - r^2 - 8r - 16)(p^2 - r^2 - 6r - 9)(p^2 - r^2 - 4r - 4)(p^2 - r^2)}{(8r^6 + 84r^5 + 350r^4 + 735r^3 + 812r^2 + 441r + 90)(2r + 7)(r + 4)(r + 5)(2r + 9)(2r + 11)(6 + r)}$$

Which for the root $r = 0$ becomes

$$b_7 = \frac{(p^2 - 36)(p^2 - 25)(p^2 - 16)(p^2 - 9)(p^2 - 4)(p^2 - 1)p^2}{681080400}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{p^2 - r^2}{2r^2 + 3r + 1}$	p^2
b_2	$\frac{(p^2 - r^2 - 2r - 1)(p^2 - r^2)}{4r^4 + 20r^3 + 35r^2 + 25r + 6}$	$\frac{1}{6}p^4 - \frac{1}{6}p^2$
b_3	$\frac{(p^2 - r^2 - 4r - 4)(p^2 - r^2 - 2r - 1)(p^2 - r^2)}{8r^6 + 84r^5 + 350r^4 + 735r^3 + 812r^2 + 441r + 90}$	$\frac{p^2(p^4 - 5p^2 + 4)}{90}$
b_4	$\frac{(p^2 - r^2 - 6r - 9)(p^2 - r^2 - 4r - 4)(p^2 - r^2 - 2r - 1)(p^2 - r^2)}{(8r^6 + 84r^5 + 350r^4 + 735r^3 + 812r^2 + 441r + 90)(2r + 7)(r + 4)}$	$\frac{p^2(p^6 - 14p^4 + 49p^2 - 36)}{2520}$
b_5	$\frac{(p^2 - r^2 - 8r - 16)(p^2 - r^2 - 6r - 9)(p^2 - r^2 - 4r - 4)(p^2 - r^2 - 2r - 1)(p^2 - r^2)}{(8r^6 + 84r^5 + 350r^4 + 735r^3 + 812r^2 + 441r + 90)(2r + 7)(r + 4)(r + 5)(2r + 9)}$	$\frac{p^2(p^8 - 30p^6 + 273p^4 - 82p^2)}{113400}$
b_6	$\frac{(p^2 - r^2 - 10r - 25)(p^2 - r^2 - 8r - 16)(p^2 - r^2 - 6r - 9)(p^2 - r^2 - 4r - 4)(p^2 - r^2 - 2r - 1)(p^2 - r^2)}{(8r^6 + 84r^5 + 350r^4 + 735r^3 + 812r^2 + 441r + 90)(2r + 7)(r + 4)(r + 5)(2r + 9)(2r + 11)(6 + r)}$	$\frac{(p^2 - 25)(p^2 - 16)(p^2 - 9)}{7484400}$
b_7	$\frac{(p^2 - r^2 - 12r - 36)(p^2 - r^2 - 10r - 25)(p^2 - r^2 - 8r - 16)(p^2 - r^2 - 6r - 9)(p^2 - r^2 - 4r - 4)(p^2 - r^2 - 2r - 1)(p^2 - r^2)}{(8r^6 + 84r^5 + 350r^4 + 735r^3 + 812r^2 + 441r + 90)(2r + 7)(r + 4)(r + 5)(2r + 9)(2r + 11)(6 + r)(7 + r)(2r + 13)}$	$\frac{(p^2 - 36)(p^2 - 25)(p^2 - 16)}{681080400}$

Using the above table, then the solution $y_2(t)$ is

$$\begin{aligned} y_2(t) &= b_0 + b_1 t + b_2 t^2 + b_3 t^3 + b_4 t^4 + b_5 t^5 + b_6 t^6 + b_7 t^7 + b_8 t^8 \dots \\ &= 1 + p^2 t + \left(\frac{1}{6} p^4 - \frac{1}{6} p^2 \right) t^2 + \frac{p^2(p^4 - 5p^2 + 4)}{90} t^3 + \frac{p^2(p^6 - 14p^4 + 49p^2 - 36)}{2520} t^4 + \frac{p^2(p^8 - 30p^6 + 273p^4 - 82p^2)}{113400} t^5 + \dots \end{aligned}$$

Therefore the homogeneous solution is

$$y_h(t) = c_1 y_1(t) + c_2 y_2(t)$$

$$\begin{aligned}
&= c_1 \sqrt{t} \left(1 + \left(\frac{p^2}{3} - \frac{1}{12} \right) t + \left(\frac{1}{30} p^4 - \frac{1}{12} p^2 + \frac{3}{160} \right) t^2 \right. \\
&\quad + \left(\frac{1}{630} p^6 - \frac{1}{72} p^4 + \frac{37}{1440} p^2 - \frac{5}{896} \right) t^3 \\
&\quad + \frac{(4p^2 - 49)(4p^2 - 25)(4p^2 - 9)(4p^2 - 1)t^4}{5806080} \\
&\quad + \frac{(4p^2 - 81)(4p^2 - 49)(4p^2 - 25)(4p^2 - 9)(4p^2 - 1)t^5}{1277337600} \\
&\quad + \frac{(4p^2 - 121)(4p^2 - 81)(4p^2 - 49)(4p^2 - 25)(4p^2 - 9)(4p^2 - 1)t^6}{398529331200} \\
&\quad + \frac{(4p^2 - 169)(4p^2 - 121)(4p^2 - 81)(4p^2 - 49)(4p^2 - 25)(4p^2 - 9)(4p^2 - 1)t^7}{167382319104000} \\
&\quad \left. + O(t^8) \right) + c_2 \left(1 + p^2 t + \left(\frac{1}{6} p^4 - \frac{1}{6} p^2 \right) t^2 + \frac{p^2(p^4 - 5p^2 + 4)t^3}{90} \right. \\
&\quad + \frac{p^2(p^6 - 14p^4 + 49p^2 - 36)t^4}{2520} + \frac{p^2(p^8 - 30p^6 + 273p^4 - 820p^2 + 576)t^5}{113400} \\
&\quad + \frac{(p^2 - 25)(p^2 - 16)(p^2 - 9)(p^2 - 4)(p^2 - 1)p^2 t^6}{7484400} \\
&\quad \left. + \frac{(p^2 - 36)(p^2 - 25)(p^2 - 16)(p^2 - 9)(p^2 - 4)(p^2 - 1)p^2 t^7}{681080400} + O(t^8) \right)
\end{aligned}$$

Hence the final solution is

$$y(t) = y_h$$

$$\begin{aligned}
&= c_1 \sqrt{t} \left(1 + \left(\frac{p^2}{3} - \frac{1}{12} \right) t + \left(\frac{1}{30} p^4 - \frac{1}{12} p^2 + \frac{3}{160} \right) t^2 \right. \\
&\quad + \left(\frac{1}{630} p^6 - \frac{1}{72} p^4 + \frac{37}{1440} p^2 - \frac{5}{896} \right) t^3 + \frac{(4p^2 - 49)(4p^2 - 25)(4p^2 - 9)(4p^2 - 1)t^4}{5806080} \\
&\quad\quad + \frac{(4p^2 - 81)(4p^2 - 49)(4p^2 - 25)(4p^2 - 9)(4p^2 - 1)t^5}{1277337600} \\
&\quad\quad + \frac{(4p^2 - 121)(4p^2 - 81)(4p^2 - 49)(4p^2 - 25)(4p^2 - 9)(4p^2 - 1)t^6}{398529331200} \\
&\quad + \frac{(4p^2 - 169)(4p^2 - 121)(4p^2 - 81)(4p^2 - 49)(4p^2 - 25)(4p^2 - 9)(4p^2 - 1)t^7}{167382319104000} \\
&\quad\quad \left. + O(t^8) \right) + c_2 \left(1 + p^2 t + \left(\frac{1}{6} p^4 - \frac{1}{6} p^2 \right) t^2 + \frac{p^2(p^4 - 5p^2 + 4)t^3}{90} \right. \\
&\quad + \frac{p^2(p^6 - 14p^4 + 49p^2 - 36)t^4}{2520} + \frac{p^2(p^8 - 30p^6 + 273p^4 - 820p^2 + 576)t^5}{113400} \\
&\quad\quad + \frac{(p^2 - 25)(p^2 - 16)(p^2 - 9)(p^2 - 4)(p^2 - 1)p^2 t^6}{7484400} \\
&\quad \left. + \frac{(p^2 - 36)(p^2 - 25)(p^2 - 16)(p^2 - 9)(p^2 - 4)(p^2 - 1)p^2 t^7}{681080400} + O(t^8) \right)
\end{aligned}$$

Replacing t in the above with the original independent variable x using $t = x - 1$ results

in

$$\begin{aligned}
y = c_1 \sqrt{x-1} & \left(1 + \left(\frac{p^2}{3} - \frac{1}{12} \right) (x-1) + \left(\frac{1}{30} p^4 - \frac{1}{12} p^2 + \frac{3}{160} \right) (x-1)^2 \right. \\
& + \left(\frac{1}{630} p^6 - \frac{1}{72} p^4 + \frac{37}{1440} p^2 - \frac{5}{896} \right) (x-1)^3 \\
& + \frac{(4p^2 - 49)(4p^2 - 25)(4p^2 - 9)(4p^2 - 1)(x-1)^4}{5806080} \\
& + \frac{(4p^2 - 81)(4p^2 - 49)(4p^2 - 25)(4p^2 - 9)(4p^2 - 1)(x-1)^5}{1277337600} \\
& + \frac{(4p^2 - 121)(4p^2 - 81)(4p^2 - 49)(4p^2 - 25)(4p^2 - 9)(4p^2 - 1)(x-1)^6}{398529331200} \\
& + \frac{(4p^2 - 169)(4p^2 - 121)(4p^2 - 81)(4p^2 - 49)(4p^2 - 25)(4p^2 - 9)(4p^2 - 1)(x-1)^7}{167382319104000} \\
& \left. + O((x-1)^8) \right) + c_2 \left(1 + p^2(x-1) + \left(\frac{1}{6} p^4 - \frac{1}{6} p^2 \right) (x-1)^2 \right. \\
& + \frac{p^2(p^4 - 5p^2 + 4)(x-1)^3}{90} + \frac{p^2(p^6 - 14p^4 + 49p^2 - 36)(x-1)^4}{2520} \\
& + \frac{p^2(p^8 - 30p^6 + 273p^4 - 820p^2 + 576)(x-1)^5}{113400} \\
& + \frac{(p^2 - 25)(p^2 - 16)(p^2 - 9)(p^2 - 4)(p^2 - 1)p^2(x-1)^6}{7484400} \\
& \left. + \frac{(p^2 - 36)(p^2 - 25)(p^2 - 16)(p^2 - 9)(p^2 - 4)(p^2 - 1)p^2(x-1)^7}{681080400} + O((x-1)^8) \right)
\end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned} y = c_1 \sqrt{x-1} & \left(1 + \left(\frac{p^2}{3} - \frac{1}{12} \right) (x-1) + \left(\frac{1}{30} p^4 - \frac{1}{12} p^2 + \frac{3}{160} \right) (x-1)^2 \right. \\ & + \left(\frac{1}{630} p^6 - \frac{1}{72} p^4 + \frac{37}{1440} p^2 - \frac{5}{896} \right) (x-1)^3 \\ & + \frac{(4p^2 - 49)(4p^2 - 25)(4p^2 - 9)(4p^2 - 1)(x-1)^4}{5806080} \\ & + \frac{(4p^2 - 81)(4p^2 - 49)(4p^2 - 25)(4p^2 - 9)(4p^2 - 1)(x-1)^5}{1277337600} \\ & + \frac{(4p^2 - 121)(4p^2 - 81)(4p^2 - 49)(4p^2 - 25)(4p^2 - 9)(4p^2 - 1)(x-1)^6}{398529331200} \\ & + \frac{(4p^2 - 169)(4p^2 - 121)(4p^2 - 81)(4p^2 - 49)(4p^2 - 25)(4p^2 - 9)(4p^2 - 1)(x-1)^7}{167382319104000} \\ & \left. + O((x-1)^8) \right) + c_2 \left(1 + p^2(x-1) + \left(\frac{1}{6} p^4 - \frac{1}{6} p^2 \right) (x-1)^2 \right. \\ & + \frac{p^2(p^4 - 5p^2 + 4)(x-1)^3}{90} + \frac{p^2(p^6 - 14p^4 + 49p^2 - 36)(x-1)^4}{2520} \\ & + \frac{p^2(p^8 - 30p^6 + 273p^4 - 820p^2 + 576)(x-1)^5}{113400} \\ & + \frac{(p^2 - 25)(p^2 - 16)(p^2 - 9)(p^2 - 4)(p^2 - 1)p^2(x-1)^6}{7484400} \\ & \left. + \frac{(p^2 - 36)(p^2 - 25)(p^2 - 16)(p^2 - 9)(p^2 - 4)(p^2 - 1)p^2(x-1)^7}{681080400} + O((x-1)^8) \right) \end{aligned} \quad (1)$$

Verification of solutions

$$\begin{aligned}
 y = c_1 \sqrt{x-1} & \left(1 + \left(\frac{p^2}{3} - \frac{1}{12} \right) (x-1) + \left(\frac{1}{30} p^4 - \frac{1}{12} p^2 + \frac{3}{160} \right) (x-1)^2 \right. \\
 & + \left(\frac{1}{630} p^6 - \frac{1}{72} p^4 + \frac{37}{1440} p^2 - \frac{5}{896} \right) (x-1)^3 \\
 & + \frac{(4p^2 - 49)(4p^2 - 25)(4p^2 - 9)(4p^2 - 1)(x-1)^4}{5806080} \\
 & + \frac{(4p^2 - 81)(4p^2 - 49)(4p^2 - 25)(4p^2 - 9)(4p^2 - 1)(x-1)^5}{1277337600} \\
 & + \frac{(4p^2 - 121)(4p^2 - 81)(4p^2 - 49)(4p^2 - 25)(4p^2 - 9)(4p^2 - 1)(x-1)^6}{398529331200} \\
 & + \frac{(4p^2 - 169)(4p^2 - 121)(4p^2 - 81)(4p^2 - 49)(4p^2 - 25)(4p^2 - 9)(4p^2 - 1)(x-1)^7}{167382319104000} \\
 & \left. + O((x-1)^8) \right) + c_2 \left(1 + p^2(x-1) + \left(\frac{1}{6} p^4 - \frac{1}{6} p^2 \right) (x-1)^2 \right. \\
 & + \frac{p^2(p^4 - 5p^2 + 4)(x-1)^3}{90} + \frac{p^2(p^6 - 14p^4 + 49p^2 - 36)(x-1)^4}{2520} \\
 & + \frac{p^2(p^8 - 30p^6 + 273p^4 - 820p^2 + 576)(x-1)^5}{113400} \\
 & + \frac{(p^2 - 25)(p^2 - 16)(p^2 - 9)(p^2 - 4)(p^2 - 1)p^2(x-1)^6}{7484400} \\
 & \left. + \frac{(p^2 - 36)(p^2 - 25)(p^2 - 16)(p^2 - 9)(p^2 - 4)(p^2 - 1)p^2(x-1)^7}{681080400} + O((x-1)^8) \right)
 \end{aligned}$$

Verified OK.

21.5.1 Maple step by step solution

Let's solve

$$(-x^2 + 1)y'' - xy' + p^2y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{xy'}{x^2-1} + \frac{p^2y}{x^2-1}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{xy'}{x^2-1} - \frac{p^2y}{x^2-1} = 0$$

□ Check to see if x_0 is a regular singular point

○ Define functions

$$\left[P_2(x) = \frac{x}{x^2-1}, P_3(x) = -\frac{p^2}{x^2-1} \right]$$

○ $(1+x) \cdot P_2(x)$ is analytic at $x = -1$

$$\left. ((1+x) \cdot P_2(x)) \right|_{x=-1} = \frac{1}{2}$$

○ $(1+x)^2 \cdot P_3(x)$ is analytic at $x = -1$

$$\left. ((1+x)^2 \cdot P_3(x)) \right|_{x=-1} = 0$$

○ $x = -1$ is a regular singular point

Check to see if x_0 is a regular singular point

$$x_0 = -1$$

• Multiply by denominators

$$(x^2 - 1)y'' + xy' - p^2y = 0$$

• Change variables using $x = u - 1$ so that the regular singular point is at $u = 0$

$$(u^2 - 2u) \left(\frac{d^2}{du^2} y(u) \right) + (u - 1) \left(\frac{d}{du} y(u) \right) - p^2 y(u) = 0$$

• Assume series solution for $y(u)$

$$y(u) = \sum_{k=0}^{\infty} a_k u^{k+r}$$

□ Rewrite ODE with series expansions

○ Convert $u^m \cdot \left(\frac{d}{du} y(u) \right)$ to series expansion for $m = 0..1$

$$u^m \cdot \left(\frac{d}{du} y(u) \right) = \sum_{k=0}^{\infty} a_k (k+r) u^{k+r-1+m}$$

○ Shift index using $k \rightarrow k + 1 - m$

$$u^m \cdot \left(\frac{d}{du} y(u) \right) = \sum_{k=-1+m}^{\infty} a_{k+1-m} (k+1-m+r) u^{k+r}$$

○ Convert $u^m \cdot \left(\frac{d^2}{du^2} y(u) \right)$ to series expansion for $m = 1..2$

$$u^m \cdot \left(\frac{d^2}{du^2} y(u) \right) = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) u^{k+r-2+m}$$

○ Shift index using $k \rightarrow k + 2 - m$

$$u^m \cdot \left(\frac{d^2}{du^2} y(u) \right) = \sum_{k=-2+m}^{\infty} a_{k+2-m} (k+2-m+r) (k+1-m+r) u^{k+r}$$

Rewrite ODE with series expansions

$$-a_0 r(-1+2r) u^{-1+r} + \left(\sum_{k=0}^{\infty} (-a_{k+1} (k+1+r) (2k+1+2r) + a_k (k+p+r) (k-p+r)) u^{k+r} \right)$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$-r(-1+2r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \left\{ 0, \frac{1}{2} \right\}$$

- Each term in the series must be 0, giving the recursion relation

$$-2\left(k + \frac{1}{2} + r\right) (k+1+r) a_{k+1} + a_k (k+p+r) (k-p+r) = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+1} = \frac{a_k (k+p+r)(k-p+r)}{(2k+1+2r)(k+1+r)}$$

- Recursion relation for $r = 0$

$$a_{k+1} = \frac{a_k (k+p)(k-p)}{(2k+1)(k+1)}$$

- Solution for $r = 0$

$$\left[y(u) = \sum_{k=0}^{\infty} a_k u^k, a_{k+1} = \frac{a_k (k+p)(k-p)}{(2k+1)(k+1)} \right]$$

- Revert the change of variables $u = 1 + x$

$$\left[y = \sum_{k=0}^{\infty} a_k (1+x)^k, a_{k+1} = \frac{a_k (k+p)(k-p)}{(2k+1)(k+1)} \right]$$

- Recursion relation for $r = \frac{1}{2}$

$$a_{k+1} = \frac{a_k (k+p+\frac{1}{2})(k-p+\frac{1}{2})}{(2k+2)(k+\frac{3}{2})}$$

- Solution for $r = \frac{1}{2}$

$$\left[y(u) = \sum_{k=0}^{\infty} a_k u^{k+\frac{1}{2}}, a_{k+1} = \frac{a_k (k+p+\frac{1}{2})(k-p+\frac{1}{2})}{(2k+2)(k+\frac{3}{2})} \right]$$

- Revert the change of variables $u = 1 + x$

$$\left[y = \sum_{k=0}^{\infty} a_k (1+x)^{k+\frac{1}{2}}, a_{k+1} = \frac{a_k (k+p+\frac{1}{2})(k-p+\frac{1}{2})}{(2k+2)(k+\frac{3}{2})} \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k (1+x)^k \right) + \left(\sum_{k=0}^{\infty} b_k (1+x)^{k+\frac{1}{2}} \right), a_{k+1} = \frac{a_k (k+p)(k-p)}{(2k+1)(k+1)}, b_{k+1} = \frac{b_k (k+p+\frac{1}{2})(k-p+\frac{1}{2})}{(2k+2)(k+\frac{3}{2})} \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
<- linear_1 successful`

```

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 338

Order:=8;

dsolve((1-x^2)*diff(y(x),x\$2)-x*diff(y(x),x)+p^2*y(x)=0,y(x),type='series',x=1);

$$\begin{aligned}
 y(x) = & c_1 \sqrt{x-1} \left(1 + \left(\frac{p^2}{3} - \frac{1}{12} \right) (x-1) + \left(\frac{1}{30} p^4 - \frac{1}{12} p^2 + \frac{3}{160} \right) (x-1)^2 \right. \\
 & + \left(\frac{1}{630} p^6 - \frac{1}{72} p^4 + \frac{37}{1440} p^2 - \frac{5}{896} \right) (x-1)^3 \\
 & + \left(\frac{1}{22680} p^8 - \frac{1}{1080} p^6 + \frac{47}{8640} p^4 - \frac{3229}{362880} p^2 + \frac{35}{18432} \right) (x-1)^4 \\
 & + \left(\frac{1}{1247400} p^{10} - \frac{1}{30240} p^8 + \frac{19}{43200} p^6 - \frac{1571}{725760} p^4 + \frac{10679}{3225600} p^2 - \frac{63}{90112} \right) (x-1)^5 \\
 & + \left(\frac{1}{97297200} p^{12} - \frac{1}{1360800} p^{10} + \frac{67}{3628800} p^8 - \frac{2159}{10886400} p^6 + \frac{153617}{174182400} p^4 \right. \\
 & \quad \left. - \frac{550499}{425779200} p^2 + \frac{231}{851968} \right) (x-1)^6 \\
 & + \left(\frac{1}{10216206000} p^{14} - \frac{1}{89812800} p^{12} + \frac{11}{23328000} p^{10} - \frac{8521}{914457600} p^8 \right. \\
 & \quad \left. + \frac{230443}{2612736000} p^6 - \frac{1206053}{3284582400} p^4 + \frac{2430898831}{4649508864000} p^2 - \frac{143}{1310720} \right) (x-1)^7 \\
 & + O((x-1)^8) \Big) + c_2 \left(1 + p^2(x-1) + \left(\frac{1}{6} p^4 - \frac{1}{6} p^2 \right) (x-1)^2 \right. \\
 & + \left(\frac{1}{90} p^6 - \frac{1}{18} p^4 + \frac{2}{45} p^2 \right) (x-1)^3 + \left(\frac{1}{2520} p^8 - \frac{1}{180} p^6 + \frac{7}{360} p^4 - \frac{1}{70} p^2 \right) (x-1)^4 \\
 & + \left(\frac{1}{113400} p^{10} - \frac{1}{3780} p^8 + \frac{13}{5400} p^6 - \frac{41}{5670} p^4 + \frac{8}{1575} p^2 \right) (x-1)^5 \\
 & + \left(\frac{1}{7484400} p^{12} - \frac{1}{136080} p^{10} + \frac{31}{226800} p^8 - \frac{139}{136080} p^6 + \frac{479}{170100} p^4 \right. \\
 & \quad \left. - \frac{4}{2079} p^2 \right) (x-1)^6 + \left(\frac{1}{681080400} p^{14} - \frac{1}{7484400} p^{12} + \frac{1}{226800} p^{10} - \frac{311}{4762800} p^8 \right. \\
 & \quad \left. + \frac{37}{85050} p^6 - \frac{59}{51975} p^4 + \frac{16}{21021} p^2 \right) (x-1)^7 + O((x-1)^8) \Big)
 \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 5699

```
AsymptoticDSolveValue[(1-x^2)*y''[x]-x*y'[x]+p^2*y[x]==0,y[x],{x,1,7}]
```

Too large to display

21.6 problem 5

Internal problem ID [6475]

Internal file name [OUTPUT/5723_Sunday_June_05_2022_03_49_11_PM_96073013/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Section 4.6. Gauss's Hypergeometric Equation. Page 187

Problem number: 5.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _exact , _linear , _homogeneous]]
```

$$(1 - e^x)y'' + \frac{y'}{2} + e^xy = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$(1 - e^x)y'' + \frac{y'}{2} + e^xy = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{1}{2(e^x - 1)}$$
$$q(x) = -\frac{e^x}{e^x - 1}$$

Table 495: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{1}{2(e^x-1)}$	
singularity	type
$x = 2i\pi Z$	“regular”

$q(x) = -\frac{e^x}{e^x-1}$	
singularity	type
$x = 2i\pi Z$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[2i\pi Z, 2i\pi Z]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned} & (1 - e^x) \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\ & + \frac{\left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right)}{2} + e^x \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \end{aligned} \tag{1}$$

Expanding $1 - e^x$ as Taylor series around $x = 0$ and keeping only the first 8 terms gives

$$\begin{aligned} 1 - e^x &= -x - \frac{1}{2}x^2 - \frac{1}{6}x^3 - \frac{1}{24}x^4 - \frac{1}{120}x^5 - \frac{1}{720}x^6 - \frac{1}{5040}x^7 - \frac{1}{40320}x^8 + \dots \\ &= -x - \frac{1}{2}x^2 - \frac{1}{6}x^3 - \frac{1}{24}x^4 - \frac{1}{120}x^5 - \frac{1}{720}x^6 - \frac{1}{5040}x^7 - \frac{1}{40320}x^8 \end{aligned}$$

Expanding e^x as Taylor series around $x = 0$ and keeping only the first 8 terms gives

$$\begin{aligned} e^x &= 1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{720}x^6 + \frac{1}{5040}x^7 + \frac{1}{40320}x^8 + \dots \\ &= 1 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{120}x^5 + \frac{1}{720}x^6 + \frac{1}{5040}x^7 + \frac{1}{40320}x^8 \end{aligned}$$

Which simplifies to

$$\begin{aligned} &\sum_{n=0}^{\infty} \left(-\frac{x^{n+r+6} a_n (n+r) (n+r-1)}{40320} \right) \\ &+ \sum_{n=0}^{\infty} \left(-\frac{x^{n+r+5} a_n (n+r) (n+r-1)}{5040} \right) \\ &+ \sum_{n=0}^{\infty} \left(-\frac{x^{n+r+4} a_n (n+r) (n+r-1)}{720} \right) \\ &+ \sum_{n=0}^{\infty} \left(-\frac{x^{n+r+3} a_n (n+r) (n+r-1)}{120} \right) \\ &+ \sum_{n=0}^{\infty} \left(-\frac{x^{n+r+2} a_n (n+r) (n+r-1)}{24} \right) \\ &+ \sum_{n=0}^{\infty} \left(-\frac{x^{1+n+r} a_n (n+r) (n+r-1)}{6} \right) \\ &+ \sum_{n=0}^{\infty} \left(-\frac{x^{n+r} a_n (n+r) (n+r-1)}{2} \right) \\ &+ \sum_{n=0}^{\infty} (-x^{n+r-1} a_n (n+r) (n+r-1)) + \left(\sum_{n=0}^{\infty} \frac{(n+r) a_n x^{n+r-1}}{2} \right) \\ &+ \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \left(\sum_{n=0}^{\infty} x^{1+n+r} a_n \right) + \left(\sum_{n=0}^{\infty} \frac{x^{n+r+2} a_n}{2} \right) \\ &+ \left(\sum_{n=0}^{\infty} \frac{x^{n+r+3} a_n}{6} \right) + \left(\sum_{n=0}^{\infty} \frac{x^{n+r+4} a_n}{24} \right) + \left(\sum_{n=0}^{\infty} \frac{x^{n+r+5} a_n}{120} \right) \\ &+ \left(\sum_{n=0}^{\infty} \frac{x^{n+r+6} a_n}{720} \right) + \left(\sum_{n=0}^{\infty} \frac{x^{n+r+7} a_n}{5040} \right) + \left(\sum_{n=0}^{\infty} \frac{x^{n+r+8} a_n}{40320} \right) = 0 \end{aligned} \tag{2A}$$

The next step is to make all powers of x be $n+r-1$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r-1} and

adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=0}^{\infty} \left(-\frac{x^{n+r+6} a_n (n+r)(n+r-1)}{40320} \right) &= \sum_{n=7}^{\infty} \left(-\frac{a_{n-7} (n+r-7)(n+r-8) x^{n+r-1}}{40320} \right) \\ \sum_{n=0}^{\infty} \left(-\frac{x^{n+r+5} a_n (n+r)(n+r-1)}{5040} \right) &= \sum_{n=6}^{\infty} \left(-\frac{a_{n-6} (n+r-6)(n+r-7) x^{n+r-1}}{5040} \right) \\ \sum_{n=0}^{\infty} \left(-\frac{x^{n+r+4} a_n (n+r)(n+r-1)}{720} \right) &= \sum_{n=5}^{\infty} \left(-\frac{a_{n-5} (-5+n+r)(n+r-6) x^{n+r-1}}{720} \right) \\ \sum_{n=0}^{\infty} \left(-\frac{x^{n+r+3} a_n (n+r)(n+r-1)}{120} \right) &= \sum_{n=4}^{\infty} \left(-\frac{a_{n-4} (n+r-4)(-5+n+r) x^{n+r-1}}{120} \right) \\ \sum_{n=0}^{\infty} \left(-\frac{x^{n+r+2} a_n (n+r)(n+r-1)}{24} \right) &= \sum_{n=3}^{\infty} \left(-\frac{a_{n-3} (n+r-3)(n+r-4) x^{n+r-1}}{24} \right) \\ \sum_{n=0}^{\infty} \left(-\frac{x^{1+n+r} a_n (n+r)(n+r-1)}{6} \right) &= \sum_{n=2}^{\infty} \left(-\frac{a_{n-2} (n+r-2)(n+r-3) x^{n+r-1}}{6} \right) \\ \sum_{n=0}^{\infty} \left(-\frac{x^{n+r} a_n (n+r)(n+r-1)}{2} \right) &= \sum_{n=1}^{\infty} \left(-\frac{a_{n-1} (n+r-1)(n+r-2) x^{n+r-1}}{2} \right) \end{aligned}$$

$$\sum_{n=0}^{\infty} a_n x^{n+r} = \sum_{n=1}^{\infty} a_{n-1} x^{n+r-1}$$

$$\sum_{n=0}^{\infty} x^{1+n+r} a_n = \sum_{n=2}^{\infty} a_{n-2} x^{n+r-1}$$

$$\sum_{n=0}^{\infty} \frac{x^{n+r+2} a_n}{2} = \sum_{n=3}^{\infty} \frac{a_{n-3} x^{n+r-1}}{2}$$

$$\sum_{n=0}^{\infty} \frac{x^{n+r+3} a_n}{6} = \sum_{n=4}^{\infty} \frac{a_{n-4} x^{n+r-1}}{6}$$

$$\sum_{n=0}^{\infty} \frac{x^{n+r+4} a_n}{24} = \sum_{n=5}^{\infty} \frac{a_{n-5} x^{n+r-1}}{24}$$

$$\sum_{n=0}^{\infty} \frac{x^{n+r+5} a_n}{120} = \sum_{n=6}^{\infty} \frac{a_{n-6} x^{n+r-1}}{120}$$

$$\sum_{n=0}^{\infty} \frac{x^{n+r+6} a_n}{720} = \sum_{n=7}^{\infty} \frac{a_{n-7} x^{n+r-1}}{720}$$

$$\sum_{n=0}^{\infty} \frac{x^{n+r+7} a_n}{5040} = \sum_{n=8}^{\infty} \frac{a_{n-8} x^{n+r-1}}{5040}$$

$$\sum_{n=0}^{\infty} \frac{x^{n+r+8} a_n}{40320} = \sum_{n=9}^{\infty} \frac{a_{n-9} x^{n+r-1}}{40320}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers

of x are the same and equal to $n + r - 1$.

$$\begin{aligned}
& \sum_{n=7}^{\infty} \left(-\frac{a_{n-7}(n+r-7)(n+r-8)x^{n+r-1}}{40320} \right) \\
& + \sum_{n=6}^{\infty} \left(-\frac{a_{n-6}(n+r-6)(n+r-7)x^{n+r-1}}{5040} \right) \\
& + \sum_{n=5}^{\infty} \left(-\frac{a_{n-5}(-5+n+r)(n+r-6)x^{n+r-1}}{720} \right) \\
& + \sum_{n=4}^{\infty} \left(-\frac{a_{n-4}(n+r-4)(-5+n+r)x^{n+r-1}}{120} \right) \\
& + \sum_{n=3}^{\infty} \left(-\frac{a_{n-3}(n+r-3)(n+r-4)x^{n+r-1}}{24} \right) \\
& + \sum_{n=2}^{\infty} \left(-\frac{a_{n-2}(n+r-2)(n+r-3)x^{n+r-1}}{6} \right) \\
& + \sum_{n=1}^{\infty} \left(-\frac{a_{n-1}(n+r-1)(n+r-2)x^{n+r-1}}{2} \right) \\
& + \sum_{n=0}^{\infty} (-x^{n+r-1}a_n(n+r)(n+r-1)) + \left(\sum_{n=0}^{\infty} \frac{(n+r)a_nx^{n+r-1}}{2} \right) \\
& + \left(\sum_{n=1}^{\infty} a_{n-1}x^{n+r-1} \right) + \left(\sum_{n=2}^{\infty} a_{n-2}x^{n+r-1} \right) + \left(\sum_{n=3}^{\infty} \frac{a_{n-3}x^{n+r-1}}{2} \right) \\
& + \left(\sum_{n=4}^{\infty} \frac{a_{n-4}x^{n+r-1}}{6} \right) + \left(\sum_{n=5}^{\infty} \frac{a_{n-5}x^{n+r-1}}{24} \right) \\
& + \left(\sum_{n=6}^{\infty} \frac{a_{n-6}x^{n+r-1}}{120} \right) + \left(\sum_{n=7}^{\infty} \frac{a_{n-7}x^{n+r-1}}{720} \right) \\
& + \left(\sum_{n=8}^{\infty} \frac{a_{n-8}x^{n+r-1}}{5040} \right) + \left(\sum_{n=9}^{\infty} \frac{a_{n-9}x^{n+r-1}}{40320} \right) = 0
\end{aligned} \tag{2B}$$

The indicial equation is obtained from $n = 0$. From Eq (2B) this gives

$$-x^{n+r-1}a_n(n+r)(n+r-1) + \frac{(n+r)a_nx^{n+r-1}}{2} = 0$$

When $n = 0$ the above becomes

$$-x^{-1+r}a_0r(-1+r) + \frac{ra_0x^{-1+r}}{2} = 0$$

Or

$$\left(-x^{-1+r}r(-1+r) + \frac{rx^{-1+r}}{2}\right)a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$rx^{-1+r}\left(\frac{3}{2} - r\right) = 0$$

Since the above is true for all x then the indicial equation becomes

$$-r^2 + \frac{3}{2}r = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned}r_1 &= \frac{3}{2} \\r_2 &= 0\end{aligned}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$rx^{-1+r}\left(\frac{3}{2} - r\right) = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = \frac{3}{2}$ is not an integer, then we can construct two linearly independent solutions

$$\begin{aligned}y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\y_2(x) &= x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)\end{aligned}$$

Or

$$\begin{aligned}y_1(x) &= \sum_{n=0}^{\infty} a_n x^{n+\frac{3}{2}} \\y_2(x) &= \sum_{n=0}^{\infty} b_n x^n\end{aligned}$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = \frac{-r + 2}{2r - 1}$$

Substituting $n = 2$ in Eq. (2B) gives

$$a_2 = \frac{r^2 - 2r + 3}{12r^2 - 3}$$

Substituting $n = 3$ in Eq. (2B) gives

$$a_3 = \frac{-4r^2 + 5r + 12}{96r^3 + 144r^2 - 24r - 36}$$

Substituting $n = 4$ in Eq. (2B) gives

$$a_4 = \frac{-4r^4 + 24r^3 - 44r^2 + 39r + 225}{2880r^4 + 11520r^3 + 10080r^2 - 2880r - 2700}$$

Substituting $n = 5$ in Eq. (2B) gives

$$a_5 = \frac{8r^4 - 6r^3 - 158r^2 + 291r + 630}{360(16r^4 + 64r^3 + 56r^2 - 16r - 15)(2r + 7)}$$

Substituting $n = 6$ in Eq. (2B) gives

$$a_6 = \frac{32r^6 - 96r^5 + 224r^4 + 948r^3 - 8311r^2 + 16653r + 39690}{15120(16r^4 + 64r^3 + 56r^2 - 16r - 15)(2r + 7)(2r + 9)}$$

Substituting $n = 7$ in Eq. (2B) gives

$$a_7 = \frac{-128r^6 - 864r^5 + 2944r^4 - 1176r^3 - 51308r^2 + 125943r + 249480}{60480(16r^4 + 64r^3 + 56r^2 - 16r - 15)(2r + 7)(2r + 9)(2r + 11)}$$

Substituting $n = 8$ in Eq. (2B) gives

$$a_8 = \frac{224r^9 + 4128r^8 + 28272r^7 + 70968r^6 + 47004r^5 + 213042r^4 - 779417r^3 - 3196848r^2 + 15434577r - 15120}{453600(16r^4 + 64r^3 + 56r^2 - 16r - 15)(2r + 7)(2r + 9)(2r + 11)(2r^2 + 29r + 104)}$$

For $9 \leq n$ the recursive equation is

$$\begin{aligned} & -\frac{a_{n-7}(n+r-7)(n+r-8)}{40320} - \frac{a_{n-6}(n+r-6)(n+r-7)}{5040} \\ & - \frac{a_{n-5}(-5+n+r)(n+r-6)}{720} - \frac{a_{n-4}(n+r-4)(-5+n+r)}{120} \\ & - \frac{a_{n-3}(n+r-3)(n+r-4)}{24} - \frac{a_{n-2}(n+r-2)(n+r-3)}{6} \\ & - \frac{a_{n-1}(n+r-1)(n+r-2)}{2} - a_n(n+r)(n+r-1) + \frac{a_n(n+r)}{2} + a_{n-1} \\ & + a_{n-2} + \frac{a_{n-3}}{2} + \frac{a_{n-4}}{6} + \frac{a_{n-5}}{24} + \frac{a_{n-6}}{120} + \frac{a_{n-7}}{720} + \frac{a_{n-8}}{5040} + \frac{a_{n-9}}{40320} = 0 \end{aligned} \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{n^2 a_{n-7} + 8n^2 a_{n-6} + 56n^2 a_{n-5} + 336n^2 a_{n-4} + 1680n^2 a_{n-3} + 6720n^2 a_{n-2} + 20160n^2 a_{n-1} + 2nra_{n-1}}{(4)}$$

Which for the root $r = \frac{3}{2}$ becomes

$$a_n = \frac{(-80640a_{n-1} - 4a_{n-7} - 32a_{n-6} - 224a_{n-5} - 1344a_{n-4} - 6720a_{n-3} - 26880a_{n-2})n^2 + (48a_{n-7} + 32a_{n-6} + 224a_{n-5} + 1344a_{n-4} + 6720a_{n-3} + 20160a_{n-2})n}{(5)}$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = \frac{3}{2}$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-r+2}{2r-1}$	$\frac{1}{4}$
a_2	$\frac{r^2-2r+3}{12r^2-3}$	$\frac{3}{32}$
a_3	$\frac{-4r^2+5r+12}{96r^3+144r^2-24r-36}$	$\frac{7}{384}$
a_4	$\frac{-4r^4+24r^3-44r^2+39r+225}{2880r^4+11520r^3+10080r^2-2880r-2700}$	$\frac{109}{30720}$
a_5	$\frac{8r^4-6r^3-158r^2+291r+630}{360(16r^4+64r^3+56r^2-16r-15)(2r+7)}$	$\frac{13}{24576}$
a_6	$\frac{32r^6-96r^5+224r^4+948r^3-8311r^2+16653r+39690}{15120(16r^4+64r^3+56r^2-16r-15)(2r+7)(2r+9)}$	$\frac{4439}{61931520}$
a_7	$\frac{-128r^6-864r^5+2944r^4-1176r^3-51308r^2+125943r+249480}{60480(16r^4+64r^3+56r^2-16r-15)(2r+7)(2r+9)(2r+11)}$	$\frac{2069}{247726080}$
a_8	$\frac{224r^9+4128r^8+28272r^7+70968r^6+47004r^5+213042r^4-779417r^3-3196848r^2+15434577r+24324300}{453600(16r^4+64r^3+56r^2-16r-15)(2r+7)(2r+9)(2r+11)(2r^2+29r+104)}$	$\frac{685613}{753087283200}$

Using the above table, then the solution $y_1(x)$ is

$$y_1(x) = x^{\frac{3}{2}}(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots)$$

$$= x^{\frac{3}{2}}\left(1 + \frac{x}{4} + \frac{3x^2}{32} + \frac{7x^3}{384} + \frac{109x^4}{30720} + \frac{13x^5}{24576} + \frac{4439x^6}{61931520} + \frac{2069x^7}{247726080} + \frac{685613x^8}{753087283200} + O(x^8)\right)$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$b_1 = \frac{-r+2}{2r-1}$$

Substituting $n = 2$ in Eq. (2B) gives

$$b_2 = \frac{r^2 - 2r + 3}{12r^2 - 3}$$

Substituting $n = 3$ in Eq. (2B) gives

$$b_3 = \frac{-4r^2 + 5r + 12}{96r^3 + 144r^2 - 24r - 36}$$

Substituting $n = 4$ in Eq. (2B) gives

$$b_4 = \frac{-4r^4 + 24r^3 - 44r^2 + 39r + 225}{2880r^4 + 11520r^3 + 10080r^2 - 2880r - 2700}$$

Substituting $n = 5$ in Eq. (2B) gives

$$b_5 = \frac{8r^4 - 6r^3 - 158r^2 + 291r + 630}{360(16r^4 + 64r^3 + 56r^2 - 16r - 15)(2r + 7)}$$

Substituting $n = 6$ in Eq. (2B) gives

$$b_6 = \frac{32r^6 - 96r^5 + 224r^4 + 948r^3 - 8311r^2 + 16653r + 39690}{15120(16r^4 + 64r^3 + 56r^2 - 16r - 15)(2r + 7)(2r + 9)}$$

Substituting $n = 7$ in Eq. (2B) gives

$$b_7 = \frac{-128r^6 - 864r^5 + 2944r^4 - 1176r^3 - 51308r^2 + 125943r + 249480}{60480(16r^4 + 64r^3 + 56r^2 - 16r - 15)(2r + 7)(2r + 9)(2r + 11)}$$

Substituting $n = 8$ in Eq. (2B) gives

$$b_8 = \frac{224r^9 + 4128r^8 + 28272r^7 + 70968r^6 + 47004r^5 + 213042r^4 - 779417r^3 - 3196848r^2 + 15434577r + 453600}{453600(16r^4 + 64r^3 + 56r^2 - 16r - 15)(2r + 7)(2r + 9)(2r + 11)(2r^2 + 29r + 104)}$$

For $9 \leq n$ the recursive equation is

$$\begin{aligned} & \frac{b_{n-7}(n+r-7)(n+r-8)}{40320} - \frac{b_{n-6}(n+r-6)(n+r-7)}{5040} \\ & - \frac{b_{n-5}(-5+n+r)(n+r-6)}{720} - \frac{b_{n-4}(n+r-4)(-5+n+r)}{120} \\ & - \frac{b_{n-3}(n+r-3)(n+r-4)}{24} - \frac{b_{n-2}(n+r-2)(n+r-3)}{6} \\ & - \frac{b_{n-1}(n+r-1)(n+r-2)}{2} - b_n(n+r)(n+r-1) + \frac{(n+r)b_n}{2} \\ & + b_{n-1} + b_{n-2} + \frac{b_{n-3}}{2} + \frac{b_{n-4}}{6} + \frac{b_{n-5}}{24} + \frac{b_{n-6}}{120} + \frac{b_{n-7}}{720} + \frac{b_{n-8}}{5040} + \frac{b_{n-9}}{40320} = 0 \end{aligned} \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = -\frac{n^2 b_{n-7} + 8n^2 b_{n-6} + 56n^2 b_{n-5} + 336n^2 b_{n-4} + 1680n^2 b_{n-3} + 6720n^2 b_{n-2} + 20160n^2 b_{n-1} + 2nr b_{n-7}}{40320n^2 - 60480n} \quad (4)$$

Which for the root $r = 0$ becomes

$$b_n = \frac{(-20160b_{n-1} - b_{n-7} - 8b_{n-6} - 56b_{n-5} - 336b_{n-4} - 1680b_{n-3} - 6720b_{n-2})n^2 + (60480b_{n-1} + 15b_{n-7})}{40320n^2 - 60480n} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{-r+2}{2r-1}$	-2
b_2	$\frac{r^2-2r+3}{12r^2-3}$	-1
b_3	$\frac{-4r^2+5r+12}{96r^3+144r^2-24r-36}$	$-\frac{1}{3}$
b_4	$\frac{-4r^4+24r^3-44r^2+39r+225}{2880r^4+11520r^3+10080r^2-2880r-2700}$	$-\frac{1}{12}$
b_5	$\frac{8r^4-6r^3-158r^2+291r+630}{360(16r^4+64r^3+56r^2-16r-15)(2r+7)}$	$-\frac{1}{60}$
b_6	$\frac{32r^6-96r^5+224r^4+948r^3-8311r^2+16653r+39690}{15120(16r^4+64r^3+56r^2-16r-15)(2r+7)(2r+9)}$	$-\frac{1}{360}$
b_7	$\frac{-128r^6-864r^5+2944r^4-1176r^3-51308r^2+125943r+249480}{60480(16r^4+64r^3+56r^2-16r-15)(2r+7)(2r+9)(2r+11)}$	$-\frac{1}{2520}$
b_8	$\frac{224r^9+4128r^8+28272r^7+70968r^6+47004r^5+213042r^4-779417r^3-3196848r^2+15434577r+24324300}{453600(16r^4+64r^3+56r^2-16r-15)(2r+7)(2r+9)(2r+11)(2r^2+29r+104)}$	$-\frac{1}{20160}$

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned} y_2(x) &= b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 + b_7x^7 + b_8x^8 \dots \\ &= 1 - 2x - x^2 - \frac{x^3}{3} - \frac{x^4}{12} - \frac{x^5}{60} - \frac{x^6}{360} - \frac{x^7}{2520} - \frac{x^8}{20160} + O(x^8) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1 y_1(x) + c_2 y_2(x) \\ &= c_1 x^{\frac{3}{2}} \left(1 + \frac{x}{4} + \frac{3x^2}{32} + \frac{7x^3}{384} + \frac{109x^4}{30720} + \frac{13x^5}{24576} + \frac{4439x^6}{61931520} + \frac{2069x^7}{247726080} + \frac{685613x^8}{753087283200} \right. \\ &\quad \left. + O(x^8) \right) + c_2 \left(1 - 2x - x^2 - \frac{x^3}{3} - \frac{x^4}{12} - \frac{x^5}{60} - \frac{x^6}{360} - \frac{x^7}{2520} - \frac{x^8}{20160} + O(x^8) \right) \end{aligned}$$

Hence the final solution is

$$\begin{aligned}
 y &= y_h \\
 &= c_1 x^{\frac{3}{2}} \left(1 + \frac{x}{4} + \frac{3x^2}{32} + \frac{7x^3}{384} + \frac{109x^4}{30720} + \frac{13x^5}{24576} + \frac{4439x^6}{61931520} + \frac{2069x^7}{247726080} + \frac{685613x^8}{753087283200} \right. \\
 &\quad \left. + O(x^8) \right) + c_2 \left(1 - 2x - x^2 - \frac{x^3}{3} - \frac{x^4}{12} - \frac{x^5}{60} - \frac{x^6}{360} - \frac{x^7}{2520} - \frac{x^8}{20160} + O(x^8) \right)
 \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}
 y &= c_1 x^{\frac{3}{2}} \left(1 + \frac{x}{4} + \frac{3x^2}{32} + \frac{7x^3}{384} + \frac{109x^4}{30720} + \frac{13x^5}{24576} + \frac{4439x^6}{61931520} + \frac{2069x^7}{247726080} \right. \\
 &\quad \left. + \frac{685613x^8}{753087283200} + O(x^8) \right) \\
 &\quad + c_2 \left(1 - 2x - x^2 - \frac{x^3}{3} - \frac{x^4}{12} - \frac{x^5}{60} - \frac{x^6}{360} - \frac{x^7}{2520} - \frac{x^8}{20160} + O(x^8) \right)
 \end{aligned}$$

Verification of solutions

$$\begin{aligned}
 y &= c_1 x^{\frac{3}{2}} \left(1 + \frac{x}{4} + \frac{3x^2}{32} + \frac{7x^3}{384} + \frac{109x^4}{30720} + \frac{13x^5}{24576} + \frac{4439x^6}{61931520} + \frac{2069x^7}{247726080} + \frac{685613x^8}{753087283200} \right. \\
 &\quad \left. + O(x^8) \right) + c_2 \left(1 - 2x - x^2 - \frac{x^3}{3} - \frac{x^4}{12} - \frac{x^5}{60} - \frac{x^6}{360} - \frac{x^7}{2520} - \frac{x^8}{20160} + O(x^8) \right)
 \end{aligned}$$

Verified OK.

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a symmetry of the form [xi=0, eta=F(x)]
<- linear_1 successful`

```

✓ Solution by Maple

Time used: 0.281 (sec). Leaf size: 52

Order:=8;

```
dsolve((1-exp(x))*diff(y(x),x$2)+1/2*diff(y(x),x)+exp(x)*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^{\frac{3}{2}} \left(1 + \frac{1}{4}x + \frac{3}{32}x^2 + \frac{7}{384}x^3 + \frac{109}{30720}x^4 + \frac{13}{24576}x^5 + \frac{4439}{61931520}x^6 + \frac{2069}{247726080}x^7 + O(x^8) \right) + c_2 \left(1 - 2x - x^2 - \frac{1}{3}x^3 - \frac{1}{12}x^4 - \frac{1}{60}x^5 - \frac{1}{360}x^6 - \frac{1}{2520}x^7 + O(x^8) \right)$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 109

```
AsymptoticDSolveValue[(1-Exp[x])*y'[x]+1/2*y'[x]+Exp[x]*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_2 \left(-\frac{x^7}{2520} - \frac{x^6}{360} - \frac{x^5}{60} - \frac{x^4}{12} - \frac{x^3}{3} - x^2 - 2x + 1 \right) + c_1 \left(\frac{2069x^7}{247726080} + \frac{4439x^6}{61931520} + \frac{13x^5}{24576} + \frac{109x^4}{30720} + \frac{7x^3}{384} + \frac{3x^2}{32} + \frac{x}{4} + 1 \right) x^{3/2}$$

22 Chapter 4. Power Series Solutions and Special Functions. Problems for review and discover.

(A) Drill Exercises . Page 194

22.1 problem 1(a)	3593
22.2 problem 1(b)	3603
22.3 problem 1(c)	3613
22.4 problem 1(d)	3626
22.5 problem 1(e)	3636
22.6 problem 1(f)	3647
22.7 problem 1(g)	3656
22.8 problem 1(h)	3668
22.9 problem 2(a)	3681
22.10 problem 2(b)	3695
22.11 problem 2(c)	3709
22.12 problem 2(d)	3724
22.13 problem 2(e)	3738
22.14 problem 2(f)	3754
22.15 problem 2(g)	3767
22.16 problem 2(h)	3781
22.17 problem 3(a)	3794
22.18 problem 3(b)	3797
22.19 problem 3(c)	3799
22.20 problem 3(d)	3802

22.1 problem 1(a)

Internal problem ID [6476]

Internal file name [OUTPUT/5724_Sunday_June_05_2022_03_49_14_PM_43329030/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Problems for review and discover. (A) Drill Exercises . Page 194

Problem number: 1(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_airy", "second_order_bessel_ode", "second order series method. Ordinary point", "second order series method. Taylor series method"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' + 2xy = x^2$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using Taylor series gives

$$\begin{aligned} y(x) &= y(x_0) + (x - x_0) y'(x_0) + \frac{(x - x_0)^2}{2} y''(x_0) + \frac{(x - x_0)^3}{3!} y'''(x_0) + \cdots \\ &= y_0 + x y'_0 + \frac{x^2}{2} f|_{x_0, y_0, y'_0} + \frac{x^3}{3!} f'|_{x_0, y_0, y'_0} + \cdots \\ &= y_0 + x y'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0} \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \quad (1)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \quad (723)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \quad (724)$$

$$\begin{aligned} \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \end{aligned} \quad (2)$$

$$\begin{aligned} \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \end{aligned} \quad (3)$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \tag{4}$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \tag{5}$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \end{aligned}$$

⋮

$$\begin{aligned} F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \tag{6}$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \tag{7}$$

To find $y(x)$ series solution around $x = 0$. Hence

$$\begin{aligned}
 F_0 &= -2xy + x^2 \\
 F_1 &= \frac{dF_0}{dx} \\
 &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \\
 &= -2xy' - 2y + 2x \\
 F_2 &= \frac{dF_1}{dx} \\
 &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} y' + \frac{\partial F_1}{\partial y'} F_1 \\
 &= 4yx^2 - 2x^3 - 4y' + 2 \\
 F_3 &= \frac{dF_2}{dx} \\
 &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} y' + \frac{\partial F_2}{\partial y'} F_2 \\
 &= 4x \left(xy' - \frac{5x}{2} + 4y \right) \\
 F_4 &= \frac{dF_3}{dx} \\
 &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} y' + \frac{\partial F_3}{\partial y'} F_3 \\
 &= -8yx^3 + 4x^4 + 24xy' + 16y - 20x \\
 F_5 &= \frac{dF_4}{dx} \\
 &= \frac{\partial F_4}{\partial x} + \frac{\partial F_4}{\partial y} y' + \frac{\partial F_4}{\partial y'} F_4 \\
 &= -8y'x^3 - 72yx^2 + 40x^3 + 40y' - 20 \\
 F_6 &= \frac{dF_5}{dx} \\
 &= \frac{\partial F_5}{\partial x} + \frac{\partial F_5}{\partial y} y' + \frac{\partial F_5}{\partial y'} F_5 \\
 &= -8x(-2yx^3 + x^4 + 12xy' + 28y - 20x)
 \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and

$y'(0) = y'(0)$ gives

$$\begin{aligned} F_0 &= 0 \\ F_1 &= -2y(0) \\ F_2 &= -4y'(0) + 2 \\ F_3 &= 0 \\ F_4 &= 16y(0) \\ F_5 &= 40y'(0) - 20 \\ F_6 &= 0 \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$y = \left(1 - \frac{1}{3}x^3 + \frac{1}{45}x^6\right)y(0) + \left(x - \frac{1}{6}x^4 + \frac{1}{126}x^7\right)y'(0) + \frac{x^4}{12} - \frac{x^7}{252} + O(x^8)$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard power series. Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$\begin{aligned} y' &= \sum_{n=1}^{\infty} n a_n x^{n-1} \\ y'' &= \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \end{aligned}$$

Substituting the above back into the ode gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = -2x \left(\sum_{n=0}^{\infty} a_n x^n \right) + x^2 \quad (1)$$

Expanding x^2 as Taylor series around $x = 0$ and keeping only the first 8 terms gives

$$\begin{aligned} x^2 &= x^2 + \dots \\ &= x^2 \end{aligned}$$

Hence the ODE in Eq (1) becomes

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + 2x \left(\sum_{n=0}^{\infty} a_n x^n \right) = x^2$$

Which simplifies to

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \left(\sum_{n=0}^{\infty} 2x^{1+n} a_n \right) = x^2 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} &= \sum_{n=0}^{\infty} (n+2) a_{n+2} (1+n) x^n \\ \sum_{n=0}^{\infty} 2x^{1+n} a_n &= \sum_{n=1}^{\infty} 2a_{n-1} x^n \end{aligned}$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (1+n) x^n \right) + \left(\sum_{n=1}^{\infty} 2a_{n-1} x^n \right) = x^2 \quad (3)$$

For $1 \leq n$, the recurrence equation is

$$((n+2) a_{n+2} (1+n) + 2a_{n-1}) x^n = x^2 \quad (4)$$

For $n = 1$ the recurrence equation gives

$$\begin{aligned} (6a_3 + 2a_0) x &= 0 \\ 6a_3 + 2a_0 &= 0 \end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_3 = -\frac{a_0}{3}$$

For $n = 2$ the recurrence equation gives

$$\begin{aligned} (12a_4 + 2a_1) x^2 &= x^2 \\ 12a_4 + 2a_1 &= 1 \end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_4 = \frac{1}{12} - \frac{a_1}{6}$$

For $n = 3$ the recurrence equation gives

$$\begin{aligned}(20a_5 + 2a_2)x^3 &= 0 \\ 20a_5 + 2a_2 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_5 = 0$$

For $n = 4$ the recurrence equation gives

$$\begin{aligned}(30a_6 + 2a_3)x^4 &= 0 \\ 30a_6 + 2a_3 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_6 = \frac{a_0}{45}$$

For $n = 5$ the recurrence equation gives

$$\begin{aligned}(42a_7 + 2a_4)x^5 &= 0 \\ 42a_7 + 2a_4 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_7 = -\frac{1}{252} + \frac{a_1}{126}$$

For $n = 6$ the recurrence equation gives

$$\begin{aligned}(56a_8 + 2a_5)x^6 &= 0 \\ 56a_8 + 2a_5 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_8 = 0$$

For $n = 7$ the recurrence equation gives

$$(72a_9 + 2a_6) x^7 = 0$$

$$72a_9 + 2a_6 = 0$$

Which after substituting the earlier terms found becomes

$$a_9 = -\frac{a_0}{1620}$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + a_1 x - \frac{a_0 x^3}{3} + \left(\frac{1}{12} - \frac{a_1}{6} \right) x^4 + \frac{a_0 x^6}{45} + \left(-\frac{1}{252} + \frac{a_1}{126} \right) x^7 + \dots$$

Collecting terms, the solution becomes

$$y = \left(1 - \frac{1}{3}x^3 + \frac{1}{45}x^6 \right) a_0 + \left(x - \frac{1}{6}x^4 + \frac{1}{126}x^7 \right) a_1 + \frac{x^4}{12} - \frac{x^7}{252} + O(x^8) \quad (3)$$

At $x = 0$ the solution above becomes

$$y = \left(1 - \frac{1}{3}x^3 + \frac{1}{45}x^6 \right) c_1 + \left(x - \frac{1}{6}x^4 + \frac{1}{126}x^7 \right) c_2 + \frac{x^4}{12} - \frac{x^7}{252} + O(x^8)$$

Summary

The solution(s) found are the following

$$y = \left(1 - \frac{1}{3}x^3 + \frac{1}{45}x^6 \right) y(0) + \left(x - \frac{1}{6}x^4 + \frac{1}{126}x^7 \right) y'(0) + \frac{x^4}{12} - \frac{x^7}{252} + O(x^8) \quad (1)$$

$$y = \left(1 - \frac{1}{3}x^3 + \frac{1}{45}x^6 \right) c_1 + \left(x - \frac{1}{6}x^4 + \frac{1}{126}x^7 \right) c_2 + \frac{x^4}{12} - \frac{x^7}{252} + O(x^8) \quad (2)$$

Verification of solutions

$$y = \left(1 - \frac{1}{3}x^3 + \frac{1}{45}x^6\right) y(0) + \left(x - \frac{1}{6}x^4 + \frac{1}{126}x^7\right) y'(0) + \frac{x^4}{12} - \frac{x^7}{252} + O(x^8)$$

Verified OK.

$$y = \left(1 - \frac{1}{3}x^3 + \frac{1}{45}x^6\right) c_1 + \left(x - \frac{1}{6}x^4 + \frac{1}{126}x^7\right) c_2 + \frac{x^4}{12} - \frac{x^7}{252} + O(x^8)$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
        <- Bessel successful
    <- special function solution successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 38

```
Order:=8;
dsolve(diff(y(x),x$2)+2*x*y(x)=x^2,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{3}x^3 + \frac{1}{45}x^6\right) y(0) + \left(x - \frac{1}{6}x^4 + \frac{1}{126}x^7\right) D(y)(0) + \frac{x^4}{12} - \frac{x^7}{252} + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 56

```
AsymptoticDSolveValue[y''[x]+2*x*y[x]==x^2,y[x],{x,0,7}]
```

$$y(x) \rightarrow -\frac{x^7}{252} + \frac{x^4}{12} + c_2 \left(\frac{x^7}{126} - \frac{x^4}{6} + x \right) + c_1 \left(\frac{x^6}{45} - \frac{x^3}{3} + 1 \right)$$

22.2 problem 1(b)

Internal problem ID [6477]

Internal file name [OUTPUT/5725_Sunday_June_05_2022_03_49_16_PM_84230385/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Problems for review and discover. (A) Drill Exercises . Page 194

Problem number: 1(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second order series method. Ordinary point", "second_order_change_of_variable_on_y_method_2", "second order series method. Taylor series method", "second_order_ode_non_constant_coeff_transformation_on_B"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - xy' + y = x$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using

Taylor series gives

$$\begin{aligned}
 y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \cdots \\
 &= y_0 + xy'_0 + \frac{x^2}{2}f|_{x_0, y_0, y'_0} + \frac{x^3}{3!}f'|_{x_0, y_0, y'_0} + \cdots \\
 &= y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0}
 \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \quad (1)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \quad (726)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \quad (727)$$

$$\begin{aligned}
 \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \quad (2)
 \end{aligned}$$

$$\begin{aligned}
 \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \quad (3)
 \end{aligned}$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \quad (4)$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \end{aligned} \quad (5)$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \\ &\vdots \\ F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \quad (6)$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \quad (7)$$

To find $y(x)$ series solution around $x = 0$. Hence

$$F_0 = xy' - y + x$$

$$\begin{aligned} F_1 &= \frac{dF_0}{dx} \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \\ &= x^2 y' - xy + x^2 + 1 \end{aligned}$$

$$\begin{aligned} F_2 &= \frac{dF_1}{dx} \\ &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} y' + \frac{\partial F_1}{\partial y'} F_1 \\ &= (x^3 + x) y' + x^3 - yx^2 + 2x - y \end{aligned}$$

$$\begin{aligned} F_3 &= \frac{dF_2}{dx} \\ &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} y' + \frac{\partial F_2}{\partial y'} F_2 \\ &= y' x^4 - yx^3 + x^4 + 3x^2 y' - 3xy + 4x^2 + 2 \end{aligned}$$

$$\begin{aligned} F_4 &= \frac{dF_3}{dx} \\ &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} y' + \frac{\partial F_3}{\partial y'} F_3 \\ &= (x^5 + 6x^3 + 3x) y' + (-x^4 - 6x^2 - 3) y + x^5 + 7x^3 + 8x \end{aligned}$$

$$\begin{aligned} F_5 &= \frac{dF_4}{dx} \\ &= \frac{\partial F_4}{\partial x} + \frac{\partial F_4}{\partial y} y' + \frac{\partial F_4}{\partial y'} F_4 \\ &= (x^6 + 10x^4 + 15x^2) y' + (-x^5 - 10x^3 - 15x) y + x^6 + 11x^4 + 24x^2 + 8 \end{aligned}$$

$$\begin{aligned} F_6 &= \frac{dF_5}{dx} \\ &= \frac{\partial F_5}{\partial x} + \frac{\partial F_5}{\partial y} y' + \frac{\partial F_5}{\partial y'} F_5 \\ &= (x^7 + 15x^5 + 45x^3 + 15x) y' + (-x^6 - 15x^4 - 45x^2 - 15) y + x^7 + 16x^5 + 59x^3 + 48x \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and

$y'(0) = y'(0)$ gives

$$\begin{aligned} F_0 &= -y(0) \\ F_1 &= 1 \\ F_2 &= -y(0) \\ F_3 &= 2 \\ F_4 &= -3y(0) \\ F_5 &= 8 \\ F_6 &= -15y(0) \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$y = \left(1 - \frac{1}{2}x^2 - \frac{1}{24}x^4 - \frac{1}{240}x^6 - \frac{1}{2688}x^8\right)y(0) + xy'(0) + \frac{x^3}{6} + \frac{x^5}{60} + \frac{x^7}{630} + O(x^8)$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard power series. Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$\begin{aligned} y' &= \sum_{n=1}^{\infty} n a_n x^{n-1} \\ y'' &= \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \end{aligned}$$

Substituting the above back into the ode gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = x \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) - \left(\sum_{n=0}^{\infty} a_n x^n \right) + x \quad (1)$$

Expanding x as Taylor series around $x = 0$ and keeping only the first 8 terms gives

$$\begin{aligned} x &= x + \dots \\ &= x \end{aligned}$$

Hence the ODE in Eq (1) becomes

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) - x \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) + \left(\sum_{n=0}^{\infty} a_n x^n \right) = x$$

Which simplifies to

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \sum_{n=1}^{\infty} (-n x^n a_n) + \left(\sum_{n=0}^{\infty} a_n x^n \right) = x \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n \right) + \sum_{n=1}^{\infty} (-n x^n a_n) + \left(\sum_{n=0}^{\infty} a_n x^n \right) = x \quad (3)$$

$n = 0$ gives

$$2a_2 + a_0 = 0$$

$$a_2 = -\frac{a_0}{2}$$

For $1 \leq n$, the recurrence equation is

$$((n+2) a_{n+2} (n+1) - n a_n + a_n) x^n = x \quad (4)$$

For $n = 1$ the recurrence equation gives

$$(6a_3) x = x$$

$$6a_3 = 1$$

Which after substituting the earlier terms found becomes

$$a_3 = \frac{1}{6}$$

For $n = 2$ the recurrence equation gives

$$(12a_4 - a_2)x^2 = 0$$
$$12a_4 - a_2 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = -\frac{a_0}{24}$$

For $n = 3$ the recurrence equation gives

$$(20a_5 - 2a_3)x^3 = 0$$
$$20a_5 - 2a_3 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = \frac{1}{60}$$

For $n = 4$ the recurrence equation gives

$$(30a_6 - 3a_4)x^4 = 0$$
$$30a_6 - 3a_4 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = -\frac{a_0}{240}$$

For $n = 5$ the recurrence equation gives

$$(42a_7 - 4a_5)x^5 = 0$$
$$42a_7 - 4a_5 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = \frac{1}{630}$$

For $n = 6$ the recurrence equation gives

$$\begin{aligned}(56a_8 - 5a_6)x^6 &= 0 \\ 56a_8 - 5a_6 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_8 = -\frac{a_0}{2688}$$

For $n = 7$ the recurrence equation gives

$$\begin{aligned}(72a_9 - 6a_7)x^7 &= 0 \\ 72a_9 - 6a_7 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_9 = \frac{1}{7560}$$

And so on. Therefore the solution is

$$\begin{aligned}y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots\end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + a_1 x - \frac{1}{2}a_0 x^2 + \frac{1}{6}x^3 - \frac{1}{24}a_0 x^4 + \frac{1}{60}x^5 - \frac{1}{240}a_0 x^6 + \frac{1}{630}x^7 + \dots$$

Collecting terms, the solution becomes

$$y = \left(1 - \frac{1}{2}x^2 - \frac{1}{24}x^4 - \frac{1}{240}x^6\right) a_0 + a_1 x + \frac{x^3}{6} + \frac{x^5}{60} + \frac{x^7}{630} + O(x^8) \quad (3)$$

At $x = 0$ the solution above becomes

$$y = \left(1 - \frac{1}{2}x^2 - \frac{1}{24}x^4 - \frac{1}{240}x^6\right) c_1 + c_2 x + \frac{x^3}{6} + \frac{x^5}{60} + \frac{x^7}{630} + O(x^8)$$

Summary

The solution(s) found are the following

$$y = \left(1 - \frac{1}{2}x^2 - \frac{1}{24}x^4 - \frac{1}{240}x^6 - \frac{1}{2688}x^8\right) y(0) + xy'(0) + \frac{x^3}{6} + \frac{x^5}{60} + \frac{x^7}{630} + O(x^8)$$
$$y = \left(1 - \frac{1}{2}x^2 - \frac{1}{24}x^4 - \frac{1}{240}x^6\right) c_1 + c_2x + \frac{x^3}{6} + \frac{x^5}{60} + \frac{x^7}{630} + O(x^8) \quad (2)$$

Verification of solutions

$$y = \left(1 - \frac{1}{2}x^2 - \frac{1}{24}x^4 - \frac{1}{240}x^6 - \frac{1}{2688}x^8\right) y(0) + xy'(0) + \frac{x^3}{6} + \frac{x^5}{60} + \frac{x^7}{630} + O(x^8)$$

Verified OK.

$$y = \left(1 - \frac{1}{2}x^2 - \frac{1}{24}x^4 - \frac{1}{240}x^6\right) c_1 + c_2x + \frac{x^3}{6} + \frac{x^5}{60} + \frac{x^7}{630} + O(x^8)$$

Verified OK.

Maple trace **Kovacic algorithm successful**

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Reducible group (found an exponential solution)
    Group is reducible, not completely reducible
<- Kovacics algorithm successful
<- solving first the homogeneous part of the ODE successful`
```


✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 35

```
Order:=8;  
dsolve(diff(y(x),x$2)-x*diff(y(x),x)+y(x)=x,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{2}x^2 - \frac{1}{24}x^4 - \frac{1}{240}x^6\right) y(0) + D(y)(0)x + \frac{x^3}{6} + \frac{x^5}{60} + \frac{x^7}{630} + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 55

```
AsymptoticDSolveValue[y''[x]-x*y'[x]+y[x]==x,y[x],{x,0,7}]
```

$$y(x) \rightarrow \frac{x^7}{630} + \frac{x^5}{60} + \frac{x^3}{6} + c_1 \left(-\frac{x^6}{240} - \frac{x^4}{24} - \frac{x^2}{2} + 1 \right) + c_2 x$$

22.3 problem 1(c)

22.3.1 Maple step by step solution 3623

Internal problem ID [6478]

Internal file name [OUTPUT/5726_Sunday_June_05_2022_03_49_18_PM_4455029/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Problems for review and discover. (A) Drill Exercises . Page 194

Problem number: 1(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_linear_constant_coeff**", "**second order series method. Ordinary point**", "**second order series method. Taylor series method**"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + y' + y = x^3 - x$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using

Taylor series gives

$$\begin{aligned}
 y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \cdots \\
 &= y_0 + xy'_0 + \frac{x^2}{2}f|_{x_0, y_0, y'_0} + \frac{x^3}{3!}f'|_{x_0, y_0, y'_0} + \cdots \\
 &= y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0}
 \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \tag{1}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \tag{729}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \tag{730}$$

$$\begin{aligned}
 \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \tag{2}
 \end{aligned}$$

$$\begin{aligned}
 \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \tag{3}
 \end{aligned}$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \tag{4}$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \tag{5}$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \end{aligned}$$

⋮

$$\begin{aligned} F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \tag{6}$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \tag{7}$$

To find $y(x)$ series solution around $x = 0$. Hence

$$\begin{aligned}
 F_0 &= -y' - y + x^3 - x \\
 F_1 &= \frac{dF_0}{dx} \\
 &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \\
 &= y - x^3 + x + 3x^2 - 1 \\
 F_2 &= \frac{dF_1}{dx} \\
 &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} y' + \frac{\partial F_1}{\partial y'} F_1 \\
 &= y' - 3x^2 + 1 + 6x \\
 F_3 &= \frac{dF_2}{dx} \\
 &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} y' + \frac{\partial F_2}{\partial y'} F_2 \\
 &= -y' - y + x^3 - 7x + 6 \\
 F_4 &= \frac{dF_3}{dx} \\
 &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} y' + \frac{\partial F_3}{\partial y'} F_3 \\
 &= y - x^3 + x + 3x^2 - 7 \\
 F_5 &= \frac{dF_4}{dx} \\
 &= \frac{\partial F_4}{\partial x} + \frac{\partial F_4}{\partial y} y' + \frac{\partial F_4}{\partial y'} F_4 \\
 &= y' - 3x^2 + 1 + 6x \\
 F_6 &= \frac{dF_5}{dx} \\
 &= \frac{\partial F_5}{\partial x} + \frac{\partial F_5}{\partial y} y' + \frac{\partial F_5}{\partial y'} F_5 \\
 &= -y' - y + x^3 - 7x + 6
 \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and

$y'(0) = y'(0)$ gives

$$F_0 = -y'(0) - y(0)$$

$$F_1 = y(0) - 1$$

$$F_2 = y'(0) + 1$$

$$F_3 = -y(0) - y'(0) + 6$$

$$F_4 = y(0) - 7$$

$$F_5 = y'(0) + 1$$

$$F_6 = -y(0) - y'(0) + 6$$

Substituting all the above in (7) and simplifying gives the solution as

$$\begin{aligned} y = & \left(1 - \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{120}x^5 + \frac{1}{720}x^6 - \frac{1}{40320}x^8\right) y(0) \\ & + \left(x - \frac{1}{2}x^2 + \frac{1}{24}x^4 - \frac{1}{120}x^5 + \frac{1}{5040}x^7 - \frac{1}{40320}x^8\right) y'(0) \\ & - \frac{x^3}{6} + \frac{x^4}{24} + \frac{x^5}{20} - \frac{7x^6}{720} + \frac{x^7}{5040} + \frac{x^8}{6720} + O(x^8) \end{aligned}$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard power series. Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$y' = \sum_{n=1}^{\infty} n a_n x^{n-1}$$

$$y'' = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}$$

Substituting the above back into the ode gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = - \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) - \left(\sum_{n=0}^{\infty} a_n x^n \right) + x^3 - x \quad (1)$$

Expanding $x^3 - x$ as Taylor series around $x = 0$ and keeping only the first 8 terms gives

$$\begin{aligned} x^3 - x &= x^3 - x + \dots \\ &= x^3 - x \end{aligned}$$

Hence the ODE in Eq (1) becomes

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) + \left(\sum_{n=0}^{\infty} a_n x^n \right) = x^3 - x$$

Which simplifies to

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) + \left(\sum_{n=0}^{\infty} a_n x^n \right) = x^3 - x \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} &= \sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n \\ \sum_{n=1}^{\infty} n a_n x^{n-1} &= \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n \end{aligned}$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n \right) + \left(\sum_{n=0}^{\infty} (n+1) a_{n+1} x^n \right) + \left(\sum_{n=0}^{\infty} a_n x^n \right) = x^3 - x \quad (3)$$

For $0 \leq n$, the recurrence equation is

$$((n+2) a_{n+2} (n+1) + (n+1) a_{n+1} + a_n) x^n = x^3 - x \quad (4)$$

For $n = 0$ the recurrence equation gives

$$\begin{aligned} (2a_2 + a_1 + a_0) 1 &= 0 \\ 2a_2 + a_1 + a_0 &= 0 \end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_2 = -\frac{a_1}{2} - \frac{a_0}{2}$$

For $n = 1$ the recurrence equation gives

$$\begin{aligned}(6a_3 + 2a_2 + a_1)x &= -x \\ 6a_3 + 2a_2 + a_1 &= -1\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_3 = -\frac{1}{6} + \frac{a_0}{6}$$

For $n = 2$ the recurrence equation gives

$$\begin{aligned}(12a_4 + 3a_3 + a_2)x^2 &= 0 \\ 12a_4 + 3a_3 + a_2 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_4 = \frac{1}{24} + \frac{a_1}{24}$$

For $n = 3$ the recurrence equation gives

$$\begin{aligned}(20a_5 + 4a_4 + a_3)x^3 &= x^3 \\ 20a_5 + 4a_4 + a_3 &= 1\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_5 = \frac{1}{20} - \frac{a_1}{120} - \frac{a_0}{120}$$

For $n = 4$ the recurrence equation gives

$$\begin{aligned}(30a_6 + 5a_5 + a_4)x^4 &= 0 \\ 30a_6 + 5a_5 + a_4 &= 0\end{aligned}$$

Which after substituting the earlier terms found becomes

$$a_6 = -\frac{7}{720} + \frac{a_0}{720}$$

For $n = 5$ the recurrence equation gives

$$(42a_7 + 6a_6 + a_5)x^5 = 0$$
$$42a_7 + 6a_6 + a_5 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = \frac{1}{5040} + \frac{a_1}{5040}$$

For $n = 6$ the recurrence equation gives

$$(56a_8 + 7a_7 + a_6)x^6 = 0$$
$$56a_8 + 7a_7 + a_6 = 0$$

Which after substituting the earlier terms found becomes

$$a_8 = \frac{1}{6720} - \frac{a_1}{40320} - \frac{a_0}{40320}$$

For $n = 7$ the recurrence equation gives

$$(72a_9 + 8a_8 + a_7)x^7 = 0$$
$$72a_9 + 8a_8 + a_7 = 0$$

Which after substituting the earlier terms found becomes

$$a_9 = -\frac{1}{51840} + \frac{a_0}{362880}$$

And so on. Therefore the solution is

$$y = \sum_{n=0}^{\infty} a_n x^n$$
$$= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + a_1 x + \left(-\frac{a_1}{2} - \frac{a_0}{2}\right) x^2 + \left(-\frac{1}{6} + \frac{a_0}{6}\right) x^3 + \left(\frac{1}{24} + \frac{a_1}{24}\right) x^4$$
$$+ \left(\frac{1}{20} - \frac{a_1}{120} - \frac{a_0}{120}\right) x^5 + \left(-\frac{7}{720} + \frac{a_0}{720}\right) x^6 + \left(\frac{1}{5040} + \frac{a_1}{5040}\right) x^7 + \dots$$

Collecting terms, the solution becomes

$$y = \left(1 - \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{120}x^5 + \frac{1}{720}x^6\right) a_0 + \left(x - \frac{1}{2}x^2 + \frac{1}{24}x^4 - \frac{1}{120}x^5 + \frac{1}{5040}x^7\right) a_1 \quad (3)$$

$$- \frac{x^3}{6} + \frac{x^4}{24} + \frac{x^5}{20} - \frac{7x^6}{720} + \frac{x^7}{5040} + O(x^8)$$

At $x = 0$ the solution above becomes

$$y = \left(1 - \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{120}x^5 + \frac{1}{720}x^6\right) c_1 + \left(x - \frac{1}{2}x^2 + \frac{1}{24}x^4 - \frac{1}{120}x^5 + \frac{1}{5040}x^7\right) c_2$$

$$- \frac{x^3}{6} + \frac{x^4}{24} + \frac{x^5}{20} - \frac{7x^6}{720} + \frac{x^7}{5040} + O(x^8)$$

Summary

The solution(s) found are the following

$$y = \left(1 - \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{120}x^5 + \frac{1}{720}x^6 - \frac{1}{40320}x^8\right) y(0)$$

$$+ \left(x - \frac{1}{2}x^2 + \frac{1}{24}x^4 - \frac{1}{120}x^5 + \frac{1}{5040}x^7 - \frac{1}{40320}x^8\right) y'(0) \quad (1)$$

$$- \frac{x^3}{6} + \frac{x^4}{24} + \frac{x^5}{20} - \frac{7x^6}{720} + \frac{x^7}{5040} + \frac{x^8}{6720} + O(x^8)$$

$$y = \left(1 - \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{120}x^5 + \frac{1}{720}x^6\right) c_1 + \left(x - \frac{1}{2}x^2 + \frac{1}{24}x^4 - \frac{1}{120}x^5 + \frac{1}{5040}x^7\right) c_2$$

$$- \frac{x^3}{6} + \frac{x^4}{24} + \frac{x^5}{20} - \frac{7x^6}{720} + \frac{x^7}{5040} + O(x^8)$$




Figure 473: Slope field plot

Verification of solutions

$$\begin{aligned}
 y &= \left(1 - \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{120}x^5 + \frac{1}{720}x^6 - \frac{1}{40320}x^8\right) y(0) \\
 &+ \left(x - \frac{1}{2}x^2 + \frac{1}{24}x^4 - \frac{1}{120}x^5 + \frac{1}{5040}x^7 - \frac{1}{40320}x^8\right) y'(0) \\
 &- \frac{x^3}{6} + \frac{x^4}{24} + \frac{x^5}{20} - \frac{7x^6}{720} + \frac{x^7}{5040} + \frac{x^8}{6720} + O(x^8)
 \end{aligned}$$

Verified OK.

$$\begin{aligned}
 y &= \left(1 - \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{120}x^5 + \frac{1}{720}x^6\right) c_1 + \left(x - \frac{1}{2}x^2 + \frac{1}{24}x^4 - \frac{1}{120}x^5 + \frac{1}{5040}x^7\right) c_2 \\
 &- \frac{x^3}{6} + \frac{x^4}{24} + \frac{x^5}{20} - \frac{7x^6}{720} + \frac{x^7}{5040} + O(x^8)
 \end{aligned}$$

Verified OK.

22.3.1 Maple step by step solution

Let's solve

$$y'' = -y' - y + x^3 - x$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + y' + y = x^3 - x$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + r + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-1) \pm (\sqrt{-3})}{2}$$

- Roots of the characteristic polynomial

$$r = \left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}, -\frac{1}{2} + \frac{i\sqrt{3}}{2} \right)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) c_1 + e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) c_2 + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = x^3 - x \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) & e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) \\ -\frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{2} - \frac{e^{-\frac{x}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}x}{2}\right)}{2} & -\frac{e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) \sqrt{3}}{2} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = \frac{\sqrt{3}e^{-x}}{2}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{2\sqrt{3}e^{-\frac{x}{2}} \left(\cos\left(\frac{\sqrt{3}x}{2}\right) \left(\int (x^3-x)e^{\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) dx \right) - \sin\left(\frac{\sqrt{3}x}{2}\right) \left(\int (x^3-x)e^{\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) dx \right) \right)}{3}$$

- Compute integrals

$$y_p(x) = x^3 - 3x^2 - x + 7$$

- Substitute particular solution into general solution to ODE

$$y = e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) c_1 + e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) c_2 + x^3 - 3x^2 - x + 7$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 63

Order:=8;

```
dsolve(diff(y(x),x$2)+diff(y(x),x)+y(x)=x^3-x,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{2}x^2 + \frac{1}{6}x^3 - \frac{1}{120}x^5 + \frac{1}{720}x^6\right) y(0) \\ + \left(x - \frac{1}{2}x^2 + \frac{1}{24}x^4 - \frac{1}{120}x^5 + \frac{1}{5040}x^7\right) D(y)(0) \\ - \frac{x^3}{6} + \frac{x^4}{24} + \frac{x^5}{20} - \frac{7x^6}{720} + \frac{x^7}{5040} + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.031 (sec). Leaf size: 105

```
AsymptoticDSolveValue[y''[x]+y'[x]+y[x]==x^3-x,y[x],{x,0,7}]
```

$$y(x) \rightarrow \frac{x^7}{5040} - \frac{7x^6}{720} + \frac{x^5}{20} + \frac{x^4}{24} - \frac{x^3}{6} + c_2 \left(\frac{x^7}{5040} - \frac{x^5}{120} + \frac{x^4}{24} - \frac{x^2}{2} + x \right) \\ + c_1 \left(\frac{x^6}{720} - \frac{x^5}{120} + \frac{x^3}{6} - \frac{x^2}{2} + 1 \right)$$

22.4 problem 1(d)

22.4.1 Maple step by step solution 3634

Internal problem ID [6479]

Internal file name [OUTPUT/5727_Sunday_June_05_2022_03_49_21_PM_84936219/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Problems for review and discover. (A) Drill Exercises . Page 194

Problem number: 1(d).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact linear second order ode", "second_order_integrable_as_is", "second order series method. Ordinary point", "second order series method. Taylor series method"

Maple gives the following as the ode type

```
[[_2nd_order , _exact , _linear , _homogeneous]]
```

$$2y'' + xy' + y = 0$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using

Taylor series gives

$$\begin{aligned}
 y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \dots \\
 &= y_0 + xy'_0 + \frac{x^2}{2}f|_{x_0, y_0, y'_0} + \frac{x^3}{3!}f'|_{x_0, y_0, y'_0} + \dots \\
 &= y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0}
 \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \tag{1}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \tag{732}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \tag{733}$$

$$\begin{aligned}
 \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \tag{2}
 \end{aligned}$$

$$\begin{aligned}
 \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \tag{3}
 \end{aligned}$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \tag{4}$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \tag{5}$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \end{aligned}$$

⋮

$$\begin{aligned} F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \tag{6}$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \tag{7}$$

To find $y(x)$ series solution around $x = 0$. Hence

$$F_0 = -\frac{xy'}{2} - \frac{y}{2}$$

$$\begin{aligned} F_1 &= \frac{dF_0}{dx} \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \\ &= \frac{x^2 y'}{4} + \frac{xy}{4} - y' \end{aligned}$$

$$\begin{aligned} F_2 &= \frac{dF_1}{dx} \\ &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} y' + \frac{\partial F_1}{\partial y'} F_1 \\ &= \frac{(-x^3 + 10x) y'}{8} + \frac{(-x^2 + 6) y}{8} \end{aligned}$$

$$\begin{aligned} F_3 &= \frac{dF_2}{dx} \\ &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} y' + \frac{\partial F_2}{\partial y'} F_2 \\ &= \frac{(x^4 - 18x^2 + 32) y'}{16} + \frac{yx(x^2 - 14)}{16} \end{aligned}$$

$$\begin{aligned} F_4 &= \frac{dF_3}{dx} \\ &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} y' + \frac{\partial F_3}{\partial y'} F_3 \\ &= \frac{(-x^5 + 28x^3 - 132x) y'}{32} - \frac{y(x^4 - 24x^2 + 60)}{32} \end{aligned}$$

$$\begin{aligned} F_5 &= \frac{dF_4}{dx} \\ &= \frac{\partial F_4}{\partial x} + \frac{\partial F_4}{\partial y} y' + \frac{\partial F_4}{\partial y'} F_4 \\ &= \frac{(x^6 - 40x^4 + 348x^2 - 384) y'}{64} + \frac{yx(x^4 - 36x^2 + 228)}{64} \end{aligned}$$

$$\begin{aligned} F_6 &= \frac{dF_5}{dx} \\ &= \frac{\partial F_5}{\partial x} + \frac{\partial F_5}{\partial y} y' + \frac{\partial F_5}{\partial y'} F_5 \\ &= \frac{(-x^7 + 54x^5 - 740x^3 + 2232x) y'}{128} - \frac{y(x^2 - 14)(x^4 - 36x^2 + 60)}{128} \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and

$y'(0) = y'(0)$ gives

$$\begin{aligned} F_0 &= -\frac{y(0)}{2} \\ F_1 &= -y'(0) \\ F_2 &= \frac{3y(0)}{4} \\ F_3 &= 2y'(0) \\ F_4 &= -\frac{15y(0)}{8} \\ F_5 &= -6y'(0) \\ F_6 &= \frac{105y(0)}{16} \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$y = \left(1 - \frac{1}{4}x^2 + \frac{1}{32}x^4 - \frac{1}{384}x^6 + \frac{1}{6144}x^8\right) y(0) + \left(x - \frac{1}{6}x^3 + \frac{1}{60}x^5 - \frac{1}{840}x^7\right) y'(0) + O(x^8)$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard power series. Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$\begin{aligned} y' &= \sum_{n=1}^{\infty} n a_n x^{n-1} \\ y'' &= \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \end{aligned}$$

Substituting the above back into the ode gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = -\frac{x \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right)}{2} - \frac{\left(\sum_{n=0}^{\infty} a_n x^n \right)}{2} \quad (1)$$

Which simplifies to

$$\left(\sum_{n=2}^{\infty} 2n(n-1) a_n x^{n-2} \right) + \left(\sum_{n=1}^{\infty} n x^n a_n \right) + \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\sum_{n=2}^{\infty} 2n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} 2(n+2) a_{n+2} (n+1) x^n$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} 2(n+2) a_{n+2} (n+1) x^n \right) + \left(\sum_{n=1}^{\infty} n x^n a_n \right) + \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0 \quad (3)$$

$n = 0$ gives

$$4a_2 + a_0 = 0$$

$$a_2 = -\frac{a_0}{4}$$

For $1 \leq n$, the recurrence equation is

$$2(n+2) a_{n+2} (n+1) + n a_n + a_n = 0 \quad (4)$$

Solving for a_{n+2} , gives

$$a_{n+2} = -\frac{a_n}{2(n+2)} \quad (5)$$

For $n = 1$ the recurrence equation gives

$$12a_3 + 2a_1 = 0$$

Which after substituting the earlier terms found becomes

$$a_3 = -\frac{a_1}{6}$$

For $n = 2$ the recurrence equation gives

$$24a_4 + 3a_2 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = \frac{a_0}{32}$$

For $n = 3$ the recurrence equation gives

$$40a_5 + 4a_3 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = \frac{a_1}{60}$$

For $n = 4$ the recurrence equation gives

$$60a_6 + 5a_4 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = -\frac{a_0}{384}$$

For $n = 5$ the recurrence equation gives

$$84a_7 + 6a_5 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = -\frac{a_1}{840}$$

For $n = 6$ the recurrence equation gives

$$112a_8 + 7a_6 = 0$$

Which after substituting the earlier terms found becomes

$$a_8 = \frac{a_0}{6144}$$

For $n = 7$ the recurrence equation gives

$$144a_9 + 8a_7 = 0$$

Which after substituting the earlier terms found becomes

$$a_9 = \frac{a_1}{15120}$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + a_1 x - \frac{1}{4} a_0 x^2 - \frac{1}{6} a_1 x^3 + \frac{1}{32} a_0 x^4 + \frac{1}{60} a_1 x^5 - \frac{1}{384} a_0 x^6 - \frac{1}{840} a_1 x^7 + \dots$$

Collecting terms, the solution becomes

$$y = \left(1 - \frac{1}{4}x^2 + \frac{1}{32}x^4 - \frac{1}{384}x^6\right) a_0 + \left(x - \frac{1}{6}x^3 + \frac{1}{60}x^5 - \frac{1}{840}x^7\right) a_1 + O(x^8) \quad (3)$$

At $x = 0$ the solution above becomes

$$y = \left(1 - \frac{1}{4}x^2 + \frac{1}{32}x^4 - \frac{1}{384}x^6\right) c_1 + \left(x - \frac{1}{6}x^3 + \frac{1}{60}x^5 - \frac{1}{840}x^7\right) c_2 + O(x^8)$$

Summary

The solution(s) found are the following

$$\begin{aligned} y &= \left(1 - \frac{1}{4}x^2 + \frac{1}{32}x^4 - \frac{1}{384}x^6 + \frac{1}{6144}x^8\right) y(0) \\ &\quad + \left(x - \frac{1}{6}x^3 + \frac{1}{60}x^5 - \frac{1}{840}x^7\right) y'(0) + O(x^8) \end{aligned} \quad (1)$$

$$y = \left(1 - \frac{1}{4}x^2 + \frac{1}{32}x^4 - \frac{1}{384}x^6\right) c_1 + \left(x - \frac{1}{6}x^3 + \frac{1}{60}x^5 - \frac{1}{840}x^7\right) c_2 + O(x^8) \quad (2)$$

Verification of solutions

$$y = \left(1 - \frac{1}{4}x^2 + \frac{1}{32}x^4 - \frac{1}{384}x^6 + \frac{1}{6144}x^8\right) y(0) + \left(x - \frac{1}{6}x^3 + \frac{1}{60}x^5 - \frac{1}{840}x^7\right) y'(0) + O(x^8)$$

Verified OK.

$$y = \left(1 - \frac{1}{4}x^2 + \frac{1}{32}x^4 - \frac{1}{384}x^6\right) c_1 + \left(x - \frac{1}{6}x^3 + \frac{1}{60}x^5 - \frac{1}{840}x^7\right) c_2 + O(x^8)$$

Verified OK.

22.4.1 Maple step by step solution

Let's solve

$$y'' = -\frac{xy'}{2} - \frac{y}{2}$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{xy'}{2} + \frac{y}{2} = 0$$

- Multiply by denominators

$$2y'' + xy' + y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^k$$

- Rewrite DE with series expansions

- Convert $x \cdot y'$ to series expansion

$$x \cdot y' = \sum_{k=0}^{\infty} a_k k x^k$$

- Convert y'' to series expansion

$$y'' = \sum_{k=2}^{\infty} a_k k(k-1) x^{k-2}$$

- Shift index using $k- > k+2$

$$y'' = \sum_{k=0}^{\infty} a_{k+2}(k+2)(k+1) x^k$$

Rewrite DE with series expansions

$$\sum_{k=0}^{\infty} (2a_{k+2}(k+2)(k+1) + a_k(k+1)) x^k = 0$$

- Each term in the series must be 0, giving the recursion relation

$$(k+1)(2ka_{k+2} + a_k + 4a_{k+2}) = 0$$

- Recursion relation that defines the series solution to the ODE

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+2} = -\frac{a_k}{2(k+2)} \right]$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
<- linear_1 successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 44

```
Order:=8;  
dsolve(2*diff(y(x),x$2)+x*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{4}x^2 + \frac{1}{32}x^4 - \frac{1}{384}x^6\right) y(0) + \left(x - \frac{1}{6}x^3 + \frac{1}{60}x^5 - \frac{1}{840}x^7\right) D(y)(0) + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 56

```
AsymptoticDSolveValue[2*y''[x]+x*y'[x]+y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_2 \left(-\frac{x^7}{840} + \frac{x^5}{60} - \frac{x^3}{6} + x \right) + c_1 \left(-\frac{x^6}{384} + \frac{x^4}{32} - \frac{x^2}{4} + 1 \right)$$

22.5 problem 1(e)

Internal problem ID [6480]

Internal file name [OUTPUT/5728_Sunday_June_05_2022_03_49_23_PM_55990753/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Problems for review and discover. (A) Drill Exercises . Page 194

Problem number: 1(e).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Ordinary point", "second order series method. Taylor series method"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$(x^2 + 4)y'' - y' + y = 0$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using Taylor series gives

$$\begin{aligned} y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \cdots \\ &= y_0 + xy'_0 + \frac{x^2}{2}f|_{x_0, y_0, y'_0} + \frac{x^3}{3!}f'|_{x_0, y_0, y'_0} + \cdots \\ &= y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0} \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \quad (1)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \quad (735)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \quad (736)$$

$$\begin{aligned} \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \end{aligned} \quad (2)$$

$$\begin{aligned} \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \end{aligned} \quad (3)$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \quad (4)$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \quad (5)$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \\ &\vdots \\ F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \quad (6)$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \quad (7)$$

To find $y(x)$ series solution around $x = 0$. Hence

$$F_0 = -\frac{-y' + y}{x^2 + 4}$$

$$\begin{aligned} F_1 &= \frac{dF_0}{dx} \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \\ &= \frac{(-x^2 - 2x - 3)y' + (2x - 1)y}{(x^2 + 4)^2} \end{aligned}$$

$$\begin{aligned} F_2 &= \frac{dF_1}{dx} \\ &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} y' + \frac{\partial F_1}{\partial y'} F_1 \\ &= \frac{(4x^3 + 4x^2 + 10x - 15)y' - 5yx^2 + 6xy + 11y}{(x^2 + 4)^3} \end{aligned}$$

$$\begin{aligned} F_3 &= \frac{dF_2}{dx} \\ &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} y' + \frac{\partial F_2}{\partial y'} F_2 \\ &= \frac{(-17x^4 - 6x^3 - 7x^2 + 156x + 69)y' + 16y(x^3 - \frac{17}{8}x^2 - \frac{29}{4}x + \frac{39}{16})}{(x^2 + 4)^4} \end{aligned}$$

$$\begin{aligned} F_4 &= \frac{dF_3}{dx} \\ &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} y' + \frac{\partial F_3}{\partial y'} F_3 \\ &= \frac{(84x^5 - 21x^4 - 288x^3 - 1268x^2 - 916x + 849)y' - 63(x^4 - \frac{10}{3}x^3 - \frac{337}{21}x^2 + \frac{740}{63}x + \frac{533}{63})y}{(x^2 + 4)^5} \end{aligned}$$

$$\begin{aligned} F_5 &= \frac{dF_4}{dx} \\ &= \frac{\partial F_4}{\partial x} + \frac{\partial F_4}{\partial y} y' + \frac{\partial F_4}{\partial y'} F_4 \\ &= \frac{(-483x^6 + 420x^5 + 4434x^4 + 9620x^3 + 7031x^2 - 22510x - 4947)y' + 294y(x^5 - \frac{69}{14}x^4 - \frac{1468}{49}x^3 + \frac{52}{1})}{(x^2 + 4)^6} \end{aligned}$$

$$\begin{aligned} F_6 &= \frac{dF_5}{dx} \\ &= \frac{\partial F_5}{\partial x} + \frac{\partial F_5}{\partial y} y' + \frac{\partial F_5}{\partial y'} F_5 \\ &= \frac{(3192x^7 - 4872x^6 - 54276x^5 - 69094x^4 - 10644x^3 + 408064x^2 + 150438x - 110223)y' - 1575y(x^6)}{(x^2 + 4)^7} \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and $y'(0) = y'(0)$ gives

$$\begin{aligned}
 F_0 &= \frac{y'(0)}{4} - \frac{y(0)}{4} \\
 F_1 &= -\frac{y(0)}{16} - \frac{3y'(0)}{16} \\
 F_2 &= \frac{11y(0)}{64} - \frac{15y'(0)}{64} \\
 F_3 &= \frac{39y(0)}{256} + \frac{69y'(0)}{256} \\
 F_4 &= -\frac{533y(0)}{1024} + \frac{849y'(0)}{1024} \\
 F_5 &= -\frac{3809y(0)}{4096} - \frac{4947y'(0)}{4096} \\
 F_6 &= \frac{62283y(0)}{16384} - \frac{110223y'(0)}{16384}
 \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$\begin{aligned}
 y = \left(1 - \frac{1}{8}x^2 - \frac{1}{96}x^3 + \frac{11}{1536}x^4 + \frac{13}{10240}x^5 - \frac{533}{737280}x^6 - \frac{3809}{20643840}x^7 \right. \\
 \left. + \frac{20761}{220200960}x^8 \right) y(0) + \left(x + \frac{1}{8}x^2 - \frac{1}{32}x^3 - \frac{5}{512}x^4 + \frac{23}{10240}x^5 + \frac{283}{245760}x^6 \right. \\
 \left. - \frac{1649}{6881280}x^7 - \frac{12247}{73400320}x^8 \right) y'(0) + O(x^8)
 \end{aligned}$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard power series The ode is normalized to be

$$(x^2 + 4)y'' - y' + y = 0$$

Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$\begin{aligned}
 y' &= \sum_{n=1}^{\infty} n a_n x^{n-1} \\
 y'' &= \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}
 \end{aligned}$$

Substituting the above back into the ode gives

$$(x^2 + 4) \left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) - \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) + \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=2}^{\infty} x^n a_n n(n-1) \right) + \left(\sum_{n=2}^{\infty} 4n(n-1) a_n x^{n-2} \right) + \sum_{n=1}^{\infty} (-n a_n x^{n-1}) + \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=2}^{\infty} 4n(n-1) a_n x^{n-2} &= \sum_{n=0}^{\infty} 4(n+2) a_{n+2} (n+1) x^n \\ \sum_{n=1}^{\infty} (-n a_n x^{n-1}) &= \sum_{n=0}^{\infty} -(n+1) a_{n+1} x^n \end{aligned}$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\begin{aligned} &\left(\sum_{n=2}^{\infty} x^n a_n n(n-1) \right) + \left(\sum_{n=0}^{\infty} 4(n+2) a_{n+2} (n+1) x^n \right) \\ &+ \sum_{n=0}^{\infty} -(n+1) a_{n+1} x^n + \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0 \end{aligned} \quad (3)$$

$n = 0$ gives

$$8a_2 - a_1 + a_0 = 0$$

$$a_2 = -\frac{a_0}{8} + \frac{a_1}{8}$$

$n = 1$ gives

$$24a_3 - 2a_2 + a_1 = 0$$

Which after substituting earlier equations, simplifies to

$$a_3 = -\frac{a_0}{96} - \frac{a_1}{32}$$

For $2 \leq n$, the recurrence equation is

$$na_n(n-1) + 4(n+2)a_{n+2}(n+1) - (n+1)a_{n+1} + a_n = 0 \quad (4)$$

Solving for a_{n+2} , gives

$$\begin{aligned} a_{n+2} &= -\frac{n^2a_n - na_n - na_{n+1} + a_n - a_{n+1}}{4(n+2)(n+1)} \\ (5) \quad &= -\frac{(n^2 - n + 1)a_n}{4(n+2)(n+1)} - \frac{(-n-1)a_{n+1}}{4(n+2)(n+1)} \end{aligned}$$

For $n = 2$ the recurrence equation gives

$$3a_2 + 48a_4 - 3a_3 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = \frac{11a_0}{1536} - \frac{5a_1}{512}$$

For $n = 3$ the recurrence equation gives

$$7a_3 + 80a_5 - 4a_4 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = \frac{13a_0}{10240} + \frac{23a_1}{10240}$$

For $n = 4$ the recurrence equation gives

$$13a_4 + 120a_6 - 5a_5 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = -\frac{533a_0}{737280} + \frac{283a_1}{245760}$$

For $n = 5$ the recurrence equation gives

$$21a_5 + 168a_7 - 6a_6 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = -\frac{3809a_0}{20643840} - \frac{1649a_1}{6881280}$$

For $n = 6$ the recurrence equation gives

$$31a_6 + 224a_8 - 7a_7 = 0$$

Which after substituting the earlier terms found becomes

$$a_8 = \frac{20761a_0}{220200960} - \frac{12247a_1}{73400320}$$

For $n = 7$ the recurrence equation gives

$$43a_7 + 288a_9 - 8a_8 = 0$$

Which after substituting the earlier terms found becomes

$$a_9 = \frac{717431a_0}{23781703680} + \frac{246887a_1}{7927234560}$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$\begin{aligned} y &= a_0 + a_1 x + \left(-\frac{a_0}{8} + \frac{a_1}{8}\right) x^2 + \left(-\frac{a_0}{96} - \frac{a_1}{32}\right) x^3 \\ &\quad + \left(\frac{11a_0}{1536} - \frac{5a_1}{512}\right) x^4 + \left(\frac{13a_0}{10240} + \frac{23a_1}{10240}\right) x^5 \\ &\quad + \left(-\frac{533a_0}{737280} + \frac{283a_1}{245760}\right) x^6 + \left(-\frac{3809a_0}{20643840} - \frac{1649a_1}{6881280}\right) x^7 + \dots \end{aligned}$$

Collecting terms, the solution becomes

$$y = \left(1 - \frac{1}{8}x^2 - \frac{1}{96}x^3 + \frac{11}{1536}x^4 + \frac{13}{10240}x^5 - \frac{533}{737280}x^6 - \frac{3809}{20643840}x^7\right) a_0 \quad (3)$$

$$+ \left(x + \frac{1}{8}x^2 - \frac{1}{32}x^3 - \frac{5}{512}x^4 + \frac{23}{10240}x^5 + \frac{283}{245760}x^6 - \frac{1649}{6881280}x^7\right) a_1 + O(x^8)$$

At $x = 0$ the solution above becomes

$$y = \left(1 - \frac{1}{8}x^2 - \frac{1}{96}x^3 + \frac{11}{1536}x^4 + \frac{13}{10240}x^5 - \frac{533}{737280}x^6 - \frac{3809}{20643840}x^7\right) c_1$$

$$+ \left(x + \frac{1}{8}x^2 - \frac{1}{32}x^3 - \frac{5}{512}x^4 + \frac{23}{10240}x^5 + \frac{283}{245760}x^6 - \frac{1649}{6881280}x^7\right) c_2 + O(x^8)$$

Summary

The solution(s) found are the following

$$y = \left(1 - \frac{1}{8}x^2 - \frac{1}{96}x^3 + \frac{11}{1536}x^4 + \frac{13}{10240}x^5 - \frac{533}{737280}x^6 - \frac{3809}{20643840}x^7\right. \\ \left. + \frac{20761}{220200960}x^8\right) y(0) + \left(x + \frac{1}{8}x^2 - \frac{1}{32}x^3 - \frac{5}{512}x^4 + \frac{23}{10240}x^5 + \frac{283}{245760}x^6\right. \\ \left. - \frac{1649}{6881280}x^7 - \frac{12247}{73400320}x^8\right) y'(0) + O(x^8)$$

$$y = \left(1 - \frac{1}{8}x^2 - \frac{1}{96}x^3 + \frac{11}{1536}x^4 + \frac{13}{10240}x^5 - \frac{533}{737280}x^6 - \frac{3809}{20643840}x^7\right) c_1 \quad (2)$$

$$+ \left(x + \frac{1}{8}x^2 - \frac{1}{32}x^3 - \frac{5}{512}x^4 + \frac{23}{10240}x^5 + \frac{283}{245760}x^6 - \frac{1649}{6881280}x^7\right) c_2 + O(x^8)$$

Verification of solutions

$$y = \left(1 - \frac{1}{8}x^2 - \frac{1}{96}x^3 + \frac{11}{1536}x^4 + \frac{13}{10240}x^5 - \frac{533}{737280}x^6 - \frac{3809}{20643840}x^7 + \frac{20761}{220200960}x^8 \right) y(0) + \left(x + \frac{1}{8}x^2 - \frac{1}{32}x^3 - \frac{5}{512}x^4 + \frac{23}{10240}x^5 + \frac{283}{245760}x^6 - \frac{1649}{6881280}x^7 - \frac{12247}{73400320}x^8 \right) y'(0) + O(x^8)$$

Verified OK.

$$y = \left(1 - \frac{1}{8}x^2 - \frac{1}{96}x^3 + \frac{11}{1536}x^4 + \frac{13}{10240}x^5 - \frac{533}{737280}x^6 - \frac{3809}{20643840}x^7 \right) c_1 + \left(x + \frac{1}{8}x^2 - \frac{1}{32}x^3 - \frac{5}{512}x^4 + \frac{23}{10240}x^5 + \frac{283}{245760}x^6 - \frac{1649}{6881280}x^7 \right) c_2 + O(x^8)$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
  -> Bessel
  -> elliptic
  -> Legendre
  -> Kummer
    -> hyper3: Equivalence to 1F1 under a power @ Moebius
  -> hypergeometric
    -> heuristic approach
      <- heuristic approach successful
    <- hypergeometric successful
  <- special function solution successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 74

```
Order:=8;  
dsolve((4+x^2)*diff(y(x),x$2)-diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{8}x^2 - \frac{1}{96}x^3 + \frac{11}{1536}x^4 + \frac{13}{10240}x^5 - \frac{533}{737280}x^6 - \frac{3809}{20643840}x^7\right) y(0) \\ + \left(x + \frac{1}{8}x^2 - \frac{1}{32}x^3 - \frac{5}{512}x^4 + \frac{23}{10240}x^5 + \frac{283}{245760}x^6 - \frac{1649}{6881280}x^7\right) D(y)(0) \\ + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 98

```
AsymptoticDSolveValue[(4+x^2)*y''[x]-y'[x]+y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(-\frac{3809x^7}{20643840} - \frac{533x^6}{737280} + \frac{13x^5}{10240} + \frac{11x^4}{1536} - \frac{x^3}{96} - \frac{x^2}{8} + 1 \right) \\ + c_2 \left(-\frac{1649x^7}{6881280} + \frac{283x^6}{245760} + \frac{23x^5}{10240} - \frac{5x^4}{512} - \frac{x^3}{32} + \frac{x^2}{8} + x \right)$$

22.6 problem 1(f)

Internal problem ID [6481]

Internal file name [OUTPUT/5729_Sunday_June_05_2022_03_49_24_PM_28259875/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Problems for review and discover. (A) Drill Exercises . Page 194

Problem number: 1(f).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second order series method. Ordinary point", "second_order_change_of_variable_on_y_method_2", "second order series method. Taylor series method", "second_order_ode_non_constant_coeff_transformation_on_B"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$(x^2 + 1)y'' - xy' + y = 0$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using

Taylor series gives

$$\begin{aligned}
 y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \cdots \\
 &= y_0 + xy'_0 + \frac{x^2}{2}f|_{x_0, y_0, y'_0} + \frac{x^3}{3!}f'|_{x_0, y_0, y'_0} + \cdots \\
 &= y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0}
 \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \tag{1}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \tag{738}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \tag{739}$$

$$\begin{aligned}
 \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \tag{2}
 \end{aligned}$$

$$\begin{aligned}
 \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \tag{3}
 \end{aligned}$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \tag{4}$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \tag{5}$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \end{aligned}$$

⋮

$$\begin{aligned} F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \tag{6}$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \tag{7}$$

To find $y(x)$ series solution around $x = 0$. Hence

$$\begin{aligned}
 F_0 &= \frac{-y + xy'}{x^2 + 1} \\
 F_1 &= \frac{dF_0}{dx} \\
 &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \\
 &= -\frac{(-y + xy')x}{(x^2 + 1)^2} \\
 F_2 &= \frac{dF_1}{dx} \\
 &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} y' + \frac{\partial F_1}{\partial y'} F_1 \\
 &= \frac{(-y + xy')(2x^2 - 1)}{(x^2 + 1)^3} \\
 F_3 &= \frac{dF_2}{dx} \\
 &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} y' + \frac{\partial F_2}{\partial y'} F_2 \\
 &= -\frac{6(x^2 - \frac{3}{2})(-y + xy')x}{(x^2 + 1)^4} \\
 F_4 &= \frac{dF_3}{dx} \\
 &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} y' + \frac{\partial F_3}{\partial y'} F_3 \\
 &= \frac{24(x^4 - 3x^2 + \frac{3}{8})(-y + xy')}{(x^2 + 1)^5} \\
 F_5 &= \frac{dF_4}{dx} \\
 &= \frac{\partial F_4}{\partial x} + \frac{\partial F_4}{\partial y} y' + \frac{\partial F_4}{\partial y'} F_4 \\
 &= -\frac{120(-y + xy')x(x^4 - 5x^2 + \frac{15}{8})}{(x^2 + 1)^6} \\
 F_6 &= \frac{dF_5}{dx} \\
 &= \frac{\partial F_5}{\partial x} + \frac{\partial F_5}{\partial y} y' + \frac{\partial F_5}{\partial y'} F_5 \\
 &= \frac{45(-y + xy')(16x^6 - 120x^4 + 90x^2 - 5)}{(x^2 + 1)^7}
 \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and $y'(0) = y'(0)$ gives

$$\begin{aligned} F_0 &= -y(0) \\ F_1 &= 0 \\ F_2 &= y(0) \\ F_3 &= 0 \\ F_4 &= -9y(0) \\ F_5 &= 0 \\ F_6 &= 225y(0) \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$y = \left(1 - \frac{1}{2}x^2 + \frac{1}{24}x^4 - \frac{1}{80}x^6 + \frac{5}{896}x^8\right) y(0) + xy'(0) + O(x^8)$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard power series The ode is normalized to be

$$(x^2 + 1) y'' - xy' + y = 0$$

Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$\begin{aligned} y' &= \sum_{n=1}^{\infty} n a_n x^{n-1} \\ y'' &= \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \end{aligned}$$

Substituting the above back into the ode gives

$$(x^2 + 1) \left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) - x \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) + \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=2}^{\infty} x^n a_n n(n-1) \right) + \left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \sum_{n=1}^{\infty} (-n a_n x^n) + \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=2}^{\infty} x^n a_n n(n-1) \right) + \left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (n+1) x^n \right) + \sum_{n=1}^{\infty} (-n a_n x^n) + \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0 \quad (3)$$

$n = 0$ gives

$$2a_2 + a_0 = 0$$

$$a_2 = -\frac{a_0}{2}$$

For $2 \leq n$, the recurrence equation is

$$n a_n (n-1) + (n+2) a_{n+2} (n+1) - n a_n + a_n = 0 \quad (4)$$

Solving for a_{n+2} , gives

$$a_{n+2} = -\frac{a_n (n^2 - 2n + 1)}{(n+2)(n+1)} \quad (5)$$

For $n = 2$ the recurrence equation gives

$$a_2 + 12a_4 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = \frac{a_0}{24}$$

For $n = 3$ the recurrence equation gives

$$4a_3 + 20a_5 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = 0$$

For $n = 4$ the recurrence equation gives

$$9a_4 + 30a_6 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = -\frac{a_0}{80}$$

For $n = 5$ the recurrence equation gives

$$16a_5 + 42a_7 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = 0$$

For $n = 6$ the recurrence equation gives

$$25a_6 + 56a_8 = 0$$

Which after substituting the earlier terms found becomes

$$a_8 = \frac{5a_0}{896}$$

For $n = 7$ the recurrence equation gives

$$36a_7 + 72a_9 = 0$$

Which after substituting the earlier terms found becomes

$$a_9 = 0$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + a_1 x - \frac{1}{2} a_0 x^2 + \frac{1}{24} a_0 x^4 - \frac{1}{80} a_0 x^6 + \dots$$

Collecting terms, the solution becomes

$$y = \left(1 - \frac{1}{2} x^2 + \frac{1}{24} x^4 - \frac{1}{80} x^6 \right) a_0 + a_1 x + O(x^8) \quad (3)$$

At $x = 0$ the solution above becomes

$$y = \left(1 - \frac{1}{2} x^2 + \frac{1}{24} x^4 - \frac{1}{80} x^6 \right) c_1 + c_2 x + O(x^8)$$

Summary

The solution(s) found are the following

$$y = \left(1 - \frac{1}{2} x^2 + \frac{1}{24} x^4 - \frac{1}{80} x^6 + \frac{5}{896} x^8 \right) y(0) + xy'(0) + O(x^8) \quad (1)$$

$$y = \left(1 - \frac{1}{2} x^2 + \frac{1}{24} x^4 - \frac{1}{80} x^6 \right) c_1 + c_2 x + O(x^8) \quad (2)$$

Verification of solutions

$$y = \left(1 - \frac{1}{2} x^2 + \frac{1}{24} x^4 - \frac{1}{80} x^6 + \frac{5}{896} x^8 \right) y(0) + xy'(0) + O(x^8)$$

Verified OK.

$$y = \left(1 - \frac{1}{2} x^2 + \frac{1}{24} x^4 - \frac{1}{80} x^6 \right) c_1 + c_2 x + O(x^8)$$

Verified OK.

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacics algorithm  
  A Liouvillian solution exists  
  Reducible group (found an exponential solution)  
  Group is reducible, not completely reducible  
<- Kovacics algorithm successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
Order:=8;  
dsolve((x^2+1)*diff(y(x),x$2)-x*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 - \frac{1}{2}x^2 + \frac{1}{24}x^4 - \frac{1}{80}x^6\right) y(0) + D(y)(0)x + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 34

```
AsymptoticDSolveValue[(x^2+1)*y''[x]-x*y'[x]+y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(-\frac{x^6}{80} + \frac{x^4}{24} - \frac{x^2}{2} + 1 \right) + c_2 x$$

22.7 problem 1(g)

22.7.1 Maple step by step solution 3664

Internal problem ID [6482]

Internal file name [OUTPUT/5730_Sunday_June_05_2022_03_49_26_PM_571338/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Problems for review and discover. (A) Drill Exercises . Page 194

Problem number: 1(g).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Ordinary point", "second order series method. Taylor series method"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - (1 + x)y' - xy = 0$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using Taylor series gives

$$\begin{aligned} y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \cdots \\ &= y_0 + xy'_0 + \frac{x^2}{2}f|_{x_0, y_0, y'_0} + \frac{x^3}{3!}f'|_{x_0, y_0, y'_0} + \cdots \\ &= y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0} \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \quad (1)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \quad (741)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \quad (742)$$

$$\begin{aligned} \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \end{aligned} \quad (2)$$

$$\begin{aligned} \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\ &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \end{aligned} \quad (3)$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \tag{4}$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \tag{5}$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \end{aligned}$$

⋮

$$\begin{aligned} F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \tag{6}$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \tag{7}$$

To find $y(x)$ series solution around $x = 0$. Hence

$$F_0 = xy' + xy + y'$$

$$\begin{aligned} F_1 &= \frac{dF_0}{dx} \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \\ &= (x^2 + 3x + 2) y' + y(x^2 + x + 1) \end{aligned}$$

$$\begin{aligned} F_2 &= \frac{dF_1}{dx} \\ &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} y' + \frac{\partial F_1}{\partial y'} F_1 \\ &= (x^3 + 5x^2 + 8x + 6) y' + y(x^3 + 3x^2 + 4x + 1) \end{aligned}$$

$$\begin{aligned} F_3 &= \frac{dF_2}{dx} \\ &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} y' + \frac{\partial F_2}{\partial y'} F_2 \\ &= (x^4 + 7x^3 + 19x^2 + 28x + 15) y' + y(x + 2) (x^3 + 3x^2 + 5x + 2) \end{aligned}$$

$$\begin{aligned} F_4 &= \frac{dF_3}{dx} \\ &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} y' + \frac{\partial F_3}{\partial y'} F_3 \\ &= (x^5 + 9x^4 + 35x^3 + 79x^2 + 93x + 47) y' + y(x^5 + 7x^4 + 23x^3 + 43x^2 + 37x + 12) \end{aligned}$$

$$\begin{aligned} F_5 &= \frac{dF_4}{dx} \\ &= \frac{\partial F_4}{\partial x} + \frac{\partial F_4}{\partial y} y' + \frac{\partial F_4}{\partial y'} F_4 \\ &= (x^6 + 11x^5 + 56x^4 + 173x^3 + 320x^2 + 335x + 152) y' + y(x^6 + 9x^5 + 40x^4 + 107x^3 + 162x^2 + 133x + 47) \end{aligned}$$

$$\begin{aligned} F_6 &= \frac{dF_5}{dx} \\ &= \frac{\partial F_5}{\partial x} + \frac{\partial F_5}{\partial y} y' + \frac{\partial F_5}{\partial y'} F_5 \\ &= (x^7 + 13x^6 + 82x^5 + 324x^4 + 824x^3 + 1336x^2 + 1260x + 524) y' + y(x^7 + 11x^6 + 62x^5 + 218x^4 + 477x^3 + 524x^2 + 335x + 152) \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and

$y'(0) = y'(0)$ gives

$$\begin{aligned} F_0 &= y'(0) \\ F_1 &= 2y'(0) + y(0) \\ F_2 &= 6y'(0) + y(0) \\ F_3 &= 15y'(0) + 4y(0) \\ F_4 &= 47y'(0) + 12y(0) \\ F_5 &= 152y'(0) + 37y(0) \\ F_6 &= 524y'(0) + 133y(0) \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$\begin{aligned} y &= \left(1 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{30}x^5 + \frac{1}{60}x^6 + \frac{37}{5040}x^7 + \frac{19}{5760}x^8\right) y(0) \\ &+ \left(x + \frac{1}{2}x^2 + \frac{1}{3}x^3 + \frac{1}{4}x^4 + \frac{1}{8}x^5 + \frac{47}{720}x^6 + \frac{19}{630}x^7 + \frac{131}{10080}x^8\right) y'(0) + O(x^8) \end{aligned}$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard power series. Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$\begin{aligned} y' &= \sum_{n=1}^{\infty} n a_n x^{n-1} \\ y'' &= \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \end{aligned}$$

Substituting the above back into the ode gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = x \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) + x \left(\sum_{n=0}^{\infty} a_n x^n \right) + \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) \quad (1)$$

Which simplifies to

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \sum_{n=1}^{\infty} (-n a_n x^{n-1}) + \sum_{n=1}^{\infty} (-n a_n x^n) + \sum_{n=0}^{\infty} (-x^{1+n} a_n) = 0 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\begin{aligned}\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} &= \sum_{n=0}^{\infty} (n+2) a_{n+2} (1+n) x^n \\ \sum_{n=1}^{\infty} (-n a_n x^{n-1}) &= \sum_{n=0}^{\infty} (-(1+n) a_{1+n} x^n) \\ \sum_{n=0}^{\infty} (-x^{1+n} a_n) &= \sum_{n=1}^{\infty} (-a_{n-1} x^n)\end{aligned}$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\begin{aligned}\left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (1+n) x^n \right) + \sum_{n=0}^{\infty} (-(1+n) a_{1+n} x^n) \\ + \sum_{n=1}^{\infty} (-n a_n x^n) + \sum_{n=1}^{\infty} (-a_{n-1} x^n) = 0\end{aligned}\tag{3}$$

$n = 0$ gives

$$2a_2 - a_1 = 0$$

$$a_2 = \frac{a_1}{2}$$

For $1 \leq n$, the recurrence equation is

$$(n+2) a_{n+2} (1+n) - (1+n) a_{1+n} - n a_n - a_{n-1} = 0\tag{4}$$

Solving for a_{n+2} , gives

$$\begin{aligned}a_{n+2} &= \frac{n a_n + n a_{1+n} + a_{1+n} + a_{n-1}}{(n+2)(1+n)} \\ (5) \quad &= \frac{n a_n}{(n+2)(1+n)} + \frac{a_{1+n}}{n+2} + \frac{a_{n-1}}{(n+2)(1+n)}\end{aligned}$$

For $n = 1$ the recurrence equation gives

$$6a_3 - 2a_2 - a_1 - a_0 = 0$$

Which after substituting the earlier terms found becomes

$$a_3 = \frac{a_1}{3} + \frac{a_0}{6}$$

For $n = 2$ the recurrence equation gives

$$12a_4 - 3a_3 - 2a_2 - a_1 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = \frac{a_1}{4} + \frac{a_0}{24}$$

For $n = 3$ the recurrence equation gives

$$20a_5 - 4a_4 - 3a_3 - a_2 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = \frac{a_1}{8} + \frac{a_0}{30}$$

For $n = 4$ the recurrence equation gives

$$30a_6 - 5a_5 - 4a_4 - a_3 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = \frac{47a_1}{720} + \frac{a_0}{60}$$

For $n = 5$ the recurrence equation gives

$$42a_7 - 6a_6 - 5a_5 - a_4 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = \frac{19a_1}{630} + \frac{37a_0}{5040}$$

For $n = 6$ the recurrence equation gives

$$56a_8 - 7a_7 - 6a_6 - a_5 = 0$$

Which after substituting the earlier terms found becomes

$$a_8 = \frac{131a_1}{10080} + \frac{19a_0}{5760}$$

For $n = 7$ the recurrence equation gives

$$72a_9 - 8a_8 - 7a_7 - a_6 = 0$$

Which after substituting the earlier terms found becomes

$$a_9 = \frac{71a_1}{13440} + \frac{17a_0}{12960}$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$\begin{aligned} y &= a_0 + a_1 x + \frac{a_1 x^2}{2} + \left(\frac{a_1}{3} + \frac{a_0}{6}\right) x^3 + \left(\frac{a_1}{4} + \frac{a_0}{24}\right) x^4 \\ &\quad + \left(\frac{a_1}{8} + \frac{a_0}{30}\right) x^5 + \left(\frac{47a_1}{720} + \frac{a_0}{60}\right) x^6 + \left(\frac{19a_1}{630} + \frac{37a_0}{5040}\right) x^7 + \dots \end{aligned}$$

Collecting terms, the solution becomes

$$\begin{aligned} y &= \left(1 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{30}x^5 + \frac{1}{60}x^6 + \frac{37}{5040}x^7\right) a_0 \\ &\quad + \left(x + \frac{1}{2}x^2 + \frac{1}{3}x^3 + \frac{1}{4}x^4 + \frac{1}{8}x^5 + \frac{47}{720}x^6 + \frac{19}{630}x^7\right) a_1 + O(x^8) \end{aligned} \tag{3}$$

At $x = 0$ the solution above becomes

$$\begin{aligned} y &= \left(1 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{30}x^5 + \frac{1}{60}x^6 + \frac{37}{5040}x^7\right) c_1 \\ &\quad + \left(x + \frac{1}{2}x^2 + \frac{1}{3}x^3 + \frac{1}{4}x^4 + \frac{1}{8}x^5 + \frac{47}{720}x^6 + \frac{19}{630}x^7\right) c_2 + O(x^8) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \left(1 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{30}x^5 + \frac{1}{60}x^6 + \frac{37}{5040}x^7 + \frac{19}{5760}x^8\right) y(0) \\ + \left(x + \frac{1}{2}x^2 + \frac{1}{3}x^3 + \frac{1}{4}x^4 + \frac{1}{8}x^5 + \frac{47}{720}x^6 + \frac{19}{630}x^7 + \frac{131}{10080}x^8\right) y'(0) + O(x^8) \quad (1)$$

$$y = \left(1 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{30}x^5 + \frac{1}{60}x^6 + \frac{37}{5040}x^7\right) c_1 \\ + \left(x + \frac{1}{2}x^2 + \frac{1}{3}x^3 + \frac{1}{4}x^4 + \frac{1}{8}x^5 + \frac{47}{720}x^6 + \frac{19}{630}x^7\right) c_2 + O(x^8) \quad (2)$$

Verification of solutions

$$y = \left(1 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{30}x^5 + \frac{1}{60}x^6 + \frac{37}{5040}x^7 + \frac{19}{5760}x^8\right) y(0) \\ + \left(x + \frac{1}{2}x^2 + \frac{1}{3}x^3 + \frac{1}{4}x^4 + \frac{1}{8}x^5 + \frac{47}{720}x^6 + \frac{19}{630}x^7 + \frac{131}{10080}x^8\right) y'(0) + O(x^8)$$

Verified OK.

$$y = \left(1 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{30}x^5 + \frac{1}{60}x^6 + \frac{37}{5040}x^7\right) c_1 \\ + \left(x + \frac{1}{2}x^2 + \frac{1}{3}x^3 + \frac{1}{4}x^4 + \frac{1}{8}x^5 + \frac{47}{720}x^6 + \frac{19}{630}x^7\right) c_2 + O(x^8)$$

Verified OK.

22.7.1 Maple step by step solution

Let's solve

$$y'' = xy' + xy + y'$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + (-1 - x)y' - xy = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^k$$

- Rewrite ODE with series expansions

- Convert $x \cdot y$ to series expansion

$$x \cdot y = \sum_{k=0}^{\infty} a_k x^{k+1}$$

- Shift index using $k \rightarrow k - 1$

$$x \cdot y = \sum_{k=1}^{\infty} a_{k-1} x^k$$

- Convert $x^m \cdot y'$ to series expansion for $m = 0.1$

$$x^m \cdot y' = \sum_{k=\max(0,1-m)}^{\infty} a_k k x^{k-1+m}$$

- Shift index using $k \rightarrow k + 1 - m$

$$x^m \cdot y' = \sum_{k=\max(0,1-m)+m-1}^{\infty} a_{k+1-m} (k + 1 - m) x^k$$

- Convert y'' to series expansion

$$y'' = \sum_{k=2}^{\infty} a_k k(k-1) x^{k-2}$$

- Shift index using $k \rightarrow k + 2$

$$y'' = \sum_{k=0}^{\infty} a_{k+2} (k+2)(k+1) x^k$$

Rewrite ODE with series expansions

$$2a_2 - a_1 + \left(\sum_{k=1}^{\infty} (a_{k+2}(k+2)(k+1) - a_{k+1}(k+1) - a_k k - a_{k-1}) x^k \right) = 0$$

- Each term must be 0

$$2a_2 - a_1 = 0$$

- Each term in the series must be 0, giving the recursion relation

$$k^2 a_{k+2} + (-a_k - a_{k+1} + 3a_{k+2}) k - a_{k-1} - a_{k+1} + 2a_{k+2} = 0$$

- Shift index using $k \rightarrow k + 1$

$$(k+1)^2 a_{k+3} + (-a_{k+1} - a_{k+2} + 3a_{k+3})(k+1) - a_k - a_{k+2} + 2a_{k+3} = 0$$

- Recursion relation that defines the series solution to the ODE

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+3} = \frac{a_{k+1}k + k a_{k+2} + a_k + a_{k+1} + 2a_{k+2}}{k^2 + 5k + 6}, 2a_2 - a_1 = 0 \right]$$

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
  A Liouvillian solution exists
  Reducible group (found an exponential solution)
  Group is reducible, not completely reducible
  Solution has integrals. Trying a special function solution free of integrals...
-> Trying a solution in terms of special functions:
  -> Bessel
  -> elliptic
  -> Legendre
  -> Kummer
    -> hyper3: Equivalence to 1F1 under a power @ Moebius
  -> hypergeometric
    -> heuristic approach
      <- heuristic approach successful
    <- hypergeometric successful
  <- special function solution successful
    -> Trying to convert hypergeometric functions to elementary form...
      <- elementary form could result into a too large expression - returning special functi
  <- Kovacics algorithm successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 69

```
Order:=8;
dsolve(diff(y(x),x$2)-(x+1)*diff(y(x),x)-x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = \left(1 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{30}x^5 + \frac{1}{60}x^6 + \frac{37}{5040}x^7\right) y(0) \\ + \left(x + \frac{1}{2}x^2 + \frac{1}{3}x^3 + \frac{1}{4}x^4 + \frac{1}{8}x^5 + \frac{47}{720}x^6 + \frac{19}{630}x^7\right) D(y)(0) + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 91

```
AsymptoticDSolveValue[y''[x]-(x+1)*y'[x]-x*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(\frac{37x^7}{5040} + \frac{x^6}{60} + \frac{x^5}{30} + \frac{x^4}{24} + \frac{x^3}{6} + 1 \right) + c_2 \left(\frac{19x^7}{630} + \frac{47x^6}{720} + \frac{x^5}{8} + \frac{x^4}{4} + \frac{x^3}{3} + \frac{x^2}{2} + x \right)$$

22.8 problem 1(h)

22.8.1 Maple step by step solution 3677

Internal problem ID [6483]

Internal file name [OUTPUT/5731_Sunday_June_05_2022_03_49_28_PM_80554525/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Problems for review and discover. (A) Drill Exercises . Page 194

Problem number: 1(h).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact linear second order ode", "second_order_integrable_as_is", "second order series method. Ordinary point", "second_order_change_of_variable_on_y_method_1", "second order series method. Taylor series method"

Maple gives the following as the ode type

```
[[_2nd_order , _exact , _linear , _homogeneous]]
```

$$(x - 1)y'' + (1 + x)y' + y = 0$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using

Taylor series gives

$$\begin{aligned}
 y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \cdots \\
 &= y_0 + xy'_0 + \frac{x^2}{2}f|_{x_0, y_0, y'_0} + \frac{x^3}{3!}f'|_{x_0, y_0, y'_0} + \cdots \\
 &= y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0}
 \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \tag{1}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \tag{744}$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \tag{745}$$

$$\begin{aligned}
 \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \tag{2}
 \end{aligned}$$

$$\begin{aligned}
 \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \tag{3}
 \end{aligned}$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \quad (4)$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y}y' + \frac{\partial f}{\partial y'}f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \end{aligned} \quad (5)$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \\ &\vdots \\ F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \quad (6)$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \quad (7)$$

To find $y(x)$ series solution around $x = 0$. Hence

$$F_0 = -\frac{xy' + y' + y}{x - 1}$$

$$\begin{aligned} F_1 &= \frac{dF_0}{dx} \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \\ &= \frac{(x^2 + x + 4)y' + (x + 2)y}{(x - 1)^2} \end{aligned}$$

$$\begin{aligned} F_2 &= \frac{dF_1}{dx} \\ &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y} y' + \frac{\partial F_1}{\partial y'} F_1 \\ &= \frac{(-x^3 - x^2 - 7x - 15)y' - y(x^2 + 2x + 9)}{(x - 1)^3} \end{aligned}$$

$$\begin{aligned} F_3 &= \frac{dF_2}{dx} \\ &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y} y' + \frac{\partial F_2}{\partial y'} F_2 \\ &= \frac{(x^4 + x^3 + 11x^2 + 31x + 76)y' + y(x^3 + 2x^2 + 13x + 44)}{(x - 1)^4} \end{aligned}$$

$$\begin{aligned} F_4 &= \frac{dF_3}{dx} \\ &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y} y' + \frac{\partial F_3}{\partial y'} F_3 \\ &= \frac{(-x^5 - x^4 - 16x^3 - 56x^2 - 191x - 455)y' - y(x^4 + 2x^3 + 18x^2 + 74x + 265)}{(x - 1)^5} \end{aligned}$$

$$\begin{aligned} F_5 &= \frac{dF_4}{dx} \\ &= \frac{\partial F_4}{\partial x} + \frac{\partial F_4}{\partial y} y' + \frac{\partial F_4}{\partial y'} F_4 \\ &= \frac{(x^6 + x^5 + 22x^4 + 92x^3 + 407x^2 + 1331x + 3186)y' + y(x^5 + 2x^4 + 24x^3 + 116x^2 + 523x + 1854)}{(x - 1)^6} \end{aligned}$$

$$\begin{aligned} F_6 &= \frac{dF_5}{dx} \\ &= \frac{\partial F_5}{\partial x} + \frac{\partial F_5}{\partial y} y' + \frac{\partial F_5}{\partial y'} F_5 \\ &= \frac{(-x^7 - x^6 - 29x^5 - 141x^4 - 771x^3 - 3235x^2 - 10655x - 25487)y' - y(x^6 + 2x^5 + 31x^4 + 172x^3 + \dots)}{(x - 1)^7} \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and $y'(0) = y'(0)$ gives

$$\begin{aligned} F_0 &= y(0) + y'(0) \\ F_1 &= 2y(0) + 4y'(0) \\ F_2 &= 9y(0) + 15y'(0) \\ F_3 &= 44y(0) + 76y'(0) \\ F_4 &= 265y(0) + 455y'(0) \\ F_5 &= 1854y(0) + 3186y'(0) \\ F_6 &= 14833y(0) + 25487y'(0) \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$\begin{aligned} y &= \left(1 + \frac{1}{2}x^2 + \frac{1}{3}x^3 + \frac{3}{8}x^4 + \frac{11}{30}x^5 + \frac{53}{144}x^6 + \frac{103}{280}x^7 + \frac{2119}{5760}x^8 \right) y(0) \\ &+ \left(x + \frac{1}{2}x^2 + \frac{2}{3}x^3 + \frac{5}{8}x^4 + \frac{19}{30}x^5 + \frac{91}{144}x^6 + \frac{177}{280}x^7 + \frac{3641}{5760}x^8 \right) y'(0) + O(x^8) \end{aligned}$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard power series. The ode is normalized to be

$$(x - 1)y'' + (1 + x)y' + y = 0$$

Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$\begin{aligned} y' &= \sum_{n=1}^{\infty} n a_n x^{n-1} \\ y'' &= \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \end{aligned}$$

Substituting the above back into the ode gives

$$(x - 1) \left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + (1 + x) \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) + \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0 \quad (1)$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=2}^{\infty} n x^{n-1} a_n (n-1) \right) + \sum_{n=2}^{\infty} (-n(n-1) a_n x^{n-2}) \\ & + \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) + \left(\sum_{n=1}^{\infty} n a_n x^n \right) + \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0 \end{aligned} \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=2}^{\infty} n x^{n-1} a_n (n-1) &= \sum_{n=1}^{\infty} (n+1) a_{n+1} n x^n \\ \sum_{n=2}^{\infty} (-n(n-1) a_n x^{n-2}) &= \sum_{n=0}^{\infty} (-(n+2) a_{n+2} (n+1) x^n) \\ \sum_{n=1}^{\infty} n a_n x^{n-1} &= \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n \end{aligned}$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\begin{aligned} & \left(\sum_{n=1}^{\infty} (n+1) a_{n+1} n x^n \right) + \sum_{n=0}^{\infty} (-(n+2) a_{n+2} (n+1) x^n) \\ & + \left(\sum_{n=0}^{\infty} (n+1) a_{n+1} x^n \right) + \left(\sum_{n=1}^{\infty} n a_n x^n \right) + \left(\sum_{n=0}^{\infty} a_n x^n \right) = 0 \end{aligned} \quad (3)$$

$n = 0$ gives

$$-2a_2 + a_1 + a_0 = 0$$

$$a_2 = \frac{a_0}{2} + \frac{a_1}{2}$$

For $1 \leq n$, the recurrence equation is

$$(n+1) a_{n+1} n - (n+2) a_{n+2} (n+1) + (n+1) a_{n+1} + n a_n + a_n = 0 \quad (4)$$

Solving for a_{n+2} , gives

$$(5) \quad \begin{aligned} a_{n+2} &= \frac{na_{n+1} + a_n + a_{n+1}}{n+2} \\ &= \frac{a_n}{n+2} + \frac{(n+1)a_{n+1}}{n+2} \end{aligned}$$

For $n = 1$ the recurrence equation gives

$$4a_2 - 6a_3 + 2a_1 = 0$$

Which after substituting the earlier terms found becomes

$$a_3 = \frac{a_0}{3} + \frac{2a_1}{3}$$

For $n = 2$ the recurrence equation gives

$$9a_3 - 12a_4 + 3a_2 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = \frac{3a_0}{8} + \frac{5a_1}{8}$$

For $n = 3$ the recurrence equation gives

$$16a_4 - 20a_5 + 4a_3 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = \frac{11a_0}{30} + \frac{19a_1}{30}$$

For $n = 4$ the recurrence equation gives

$$25a_5 - 30a_6 + 5a_4 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = \frac{53a_0}{144} + \frac{91a_1}{144}$$

For $n = 5$ the recurrence equation gives

$$36a_6 - 42a_7 + 6a_5 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = \frac{103a_0}{280} + \frac{177a_1}{280}$$

For $n = 6$ the recurrence equation gives

$$49a_7 - 56a_8 + 7a_6 = 0$$

Which after substituting the earlier terms found becomes

$$a_8 = \frac{2119a_0}{5760} + \frac{3641a_1}{5760}$$

For $n = 7$ the recurrence equation gives

$$64a_8 - 72a_9 + 8a_7 = 0$$

Which after substituting the earlier terms found becomes

$$a_9 = \frac{16687a_0}{45360} + \frac{28673a_1}{45360}$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$\begin{aligned} y &= a_0 + a_1 x + \left(\frac{a_0}{2} + \frac{a_1}{2}\right) x^2 + \left(\frac{a_0}{3} + \frac{2a_1}{3}\right) x^3 + \left(\frac{3a_0}{8} + \frac{5a_1}{8}\right) x^4 \\ &\quad + \left(\frac{11a_0}{30} + \frac{19a_1}{30}\right) x^5 + \left(\frac{53a_0}{144} + \frac{91a_1}{144}\right) x^6 + \left(\frac{103a_0}{280} + \frac{177a_1}{280}\right) x^7 + \dots \end{aligned}$$

Collecting terms, the solution becomes

$$y = \left(1 + \frac{1}{2}x^2 + \frac{1}{3}x^3 + \frac{3}{8}x^4 + \frac{11}{30}x^5 + \frac{53}{144}x^6 + \frac{103}{280}x^7\right) a_0 + \left(x + \frac{1}{2}x^2 + \frac{2}{3}x^3 + \frac{5}{8}x^4 + \frac{19}{30}x^5 + \frac{91}{144}x^6 + \frac{177}{280}x^7\right) a_1 + O(x^8) \quad (3)$$

At $x = 0$ the solution above becomes

$$y = \left(1 + \frac{1}{2}x^2 + \frac{1}{3}x^3 + \frac{3}{8}x^4 + \frac{11}{30}x^5 + \frac{53}{144}x^6 + \frac{103}{280}x^7\right) c_1 + \left(x + \frac{1}{2}x^2 + \frac{2}{3}x^3 + \frac{5}{8}x^4 + \frac{19}{30}x^5 + \frac{91}{144}x^6 + \frac{177}{280}x^7\right) c_2 + O(x^8)$$

Summary

The solution(s) found are the following

$$y = \left(1 + \frac{1}{2}x^2 + \frac{1}{3}x^3 + \frac{3}{8}x^4 + \frac{11}{30}x^5 + \frac{53}{144}x^6 + \frac{103}{280}x^7 + \frac{2119}{5760}x^8\right) y(0) + \left(x + \frac{1}{2}x^2 + \frac{2}{3}x^3 + \frac{5}{8}x^4 + \frac{19}{30}x^5 + \frac{91}{144}x^6 + \frac{177}{280}x^7 + \frac{3641}{5760}x^8\right) y'(0) + O(x^8) \quad (1)$$

$$y = \left(1 + \frac{1}{2}x^2 + \frac{1}{3}x^3 + \frac{3}{8}x^4 + \frac{11}{30}x^5 + \frac{53}{144}x^6 + \frac{103}{280}x^7\right) c_1 + \left(x + \frac{1}{2}x^2 + \frac{2}{3}x^3 + \frac{5}{8}x^4 + \frac{19}{30}x^5 + \frac{91}{144}x^6 + \frac{177}{280}x^7\right) c_2 + O(x^8) \quad (2)$$

Verification of solutions

$$y = \left(1 + \frac{1}{2}x^2 + \frac{1}{3}x^3 + \frac{3}{8}x^4 + \frac{11}{30}x^5 + \frac{53}{144}x^6 + \frac{103}{280}x^7 + \frac{2119}{5760}x^8\right) y(0) + \left(x + \frac{1}{2}x^2 + \frac{2}{3}x^3 + \frac{5}{8}x^4 + \frac{19}{30}x^5 + \frac{91}{144}x^6 + \frac{177}{280}x^7 + \frac{3641}{5760}x^8\right) y'(0) + O(x^8)$$

Verified OK.

$$y = \left(1 + \frac{1}{2}x^2 + \frac{1}{3}x^3 + \frac{3}{8}x^4 + \frac{11}{30}x^5 + \frac{53}{144}x^6 + \frac{103}{280}x^7\right) c_1 + \left(x + \frac{1}{2}x^2 + \frac{2}{3}x^3 + \frac{5}{8}x^4 + \frac{19}{30}x^5 + \frac{91}{144}x^6 + \frac{177}{280}x^7\right) c_2 + O(x^8)$$

Verified OK.

22.8.1 Maple step by step solution

Let's solve

$$(x - 1)y'' + (1 + x)y' + y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{y}{x-1} - \frac{(1+x)y'}{x-1}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{(1+x)y'}{x-1} + \frac{y}{x-1} = 0$$

- Check to see if $x_0 = 1$ is a regular singular point

- Define functions

$$[P_2(x) = \frac{1+x}{x-1}, P_3(x) = \frac{1}{x-1}]$$

- $(x - 1) \cdot P_2(x)$ is analytic at $x = 1$

$$((x - 1) \cdot P_2(x)) \Big|_{x=1} = 2$$

- $(x - 1)^2 \cdot P_3(x)$ is analytic at $x = 1$

$$((x - 1)^2 \cdot P_3(x)) \Big|_{x=1} = 0$$

- $x = 1$ is a regular singular point

Check to see if $x_0 = 1$ is a regular singular point

$$x_0 = 1$$

- Multiply by denominators

$$(x - 1)y'' + (1 + x)y' + y = 0$$

- Change variables using $x = u + 1$ so that the regular singular point is at $u = 0$

$$u \left(\frac{d^2}{du^2} y(u) \right) + (2 + u) \left(\frac{d}{du} y(u) \right) + y(u) = 0$$

- Assume series solution for $y(u)$

$$y(u) = \sum_{k=0}^{\infty} a_k u^{k+r}$$

- Rewrite ODE with series expansions

- Convert $u^m \cdot \left(\frac{d}{du} y(u) \right)$ to series expansion for $m = 0..1$

$$u^m \cdot \left(\frac{d}{du} y(u) \right) = \sum_{k=0}^{\infty} a_k (k+r) u^{k+r-1+m}$$

- Shift index using $k \rightarrow k+1-m$

$$u^m \cdot \left(\frac{d}{du} y(u) \right) = \sum_{k=-1+m}^{\infty} a_{k+1-m} (k+1-m+r) u^{k+r}$$

- Convert $u \cdot \left(\frac{d^2}{du^2} y(u) \right)$ to series expansion

$$u \cdot \left(\frac{d^2}{du^2} y(u) \right) = \sum_{k=0}^{\infty} a_k (k+r) (k+r-1) u^{k+r-1}$$

- Shift index using $k \rightarrow k+1$

$$u \cdot \left(\frac{d^2}{du^2} y(u) \right) = \sum_{k=-1}^{\infty} a_{k+1} (k+1+r) (k+r) u^{k+r}$$

Rewrite ODE with series expansions

$$a_0 r(1+r) u^{-1+r} + \left(\sum_{k=0}^{\infty} (a_{k+1} (k+1+r) (k+2+r) + a_k (k+1+r)) u^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$r(1+r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \{-1, 0\}$$

- Each term in the series must be 0, giving the recursion relation

$$(k+1+r) (a_{k+1} (k+2+r) + a_k) = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+1} = -\frac{a_k}{k+2+r}$$

- Recursion relation for $r = -1$

$$a_{k+1} = -\frac{a_k}{k+1}$$

- Solution for $r = -1$

$$\left[y(u) = \sum_{k=0}^{\infty} a_k u^{k-1}, a_{k+1} = -\frac{a_k}{k+1} \right]$$

- Revert the change of variables $u = x - 1$

$$\left[y = \sum_{k=0}^{\infty} a_k (x-1)^{k-1}, a_{k+1} = -\frac{a_k}{k+1} \right]$$

- Recursion relation for $r = 0$

$$a_{k+1} = -\frac{a_k}{k+2}$$

- Solution for $r = 0$

$$\left[y(u) = \sum_{k=0}^{\infty} a_k u^k, a_{k+1} = -\frac{a_k}{k+2} \right]$$

- Revert the change of variables $u = x - 1$

$$\left[y = \sum_{k=0}^{\infty} a_k (x-1)^k, a_{k+1} = -\frac{a_k}{k+2} \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k (x-1)^{k-1} \right) + \left(\sum_{k=0}^{\infty} b_k (x-1)^k \right), a_{k+1} = -\frac{a_k}{k+1}, b_{k+1} = -\frac{b_k}{k+2} \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
<- linear_1 successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 74

```

Order:=8;
dsolve((x-1)*diff(y(x),x$2)+(x+1)*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);

```

$$y(x) = \left(1 + \frac{1}{2}x^2 + \frac{1}{3}x^3 + \frac{3}{8}x^4 + \frac{11}{30}x^5 + \frac{53}{144}x^6 + \frac{103}{280}x^7 \right) y(0) + \left(x + \frac{1}{2}x^2 + \frac{2}{3}x^3 + \frac{5}{8}x^4 + \frac{19}{30}x^5 + \frac{91}{144}x^6 + \frac{177}{280}x^7 \right) D(y)(0) + O(x^8)$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 98

```
AsymptoticDSolveValue[(x-1)*y''[x]+(x+1)*y'[x]+y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(\frac{103x^7}{280} + \frac{53x^6}{144} + \frac{11x^5}{30} + \frac{3x^4}{8} + \frac{x^3}{3} + \frac{x^2}{2} + 1 \right) \\ + c_2 \left(\frac{177x^7}{280} + \frac{91x^6}{144} + \frac{19x^5}{30} + \frac{5x^4}{8} + \frac{2x^3}{3} + \frac{x^2}{2} + x \right)$$

22.9 problem 2(a)

22.9.1 Maple step by step solution 3690

Internal problem ID [6484]

Internal file name [OUTPUT/5732_Sunday_June_05_2022_03_49_30_PM_54862276/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Problems for review and discover. (A) Drill Exercises . Page 194

Problem number: 2(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Complex roots"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$(x^2 + 1) x^2 y'' - x y' + (x + 2) y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$(x^4 + x^2) y'' - x y' + (x + 2) y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{1}{x(x^2 + 1)}$$
$$q(x) = \frac{x + 2}{(x^2 + 1)x^2}$$

Table 500: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{1}{x(x^2+1)}$	
singularity	type
$x = 0$	“regular”
$x = -i$	“regular”
$x = i$	“regular”

$q(x) = \frac{x+2}{(x^2+1)x^2}$	
singularity	type
$x = 0$	“regular”
$x = -i$	“regular”
$x = i$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0, -i, i, \infty]$

Irregular singular points : $[\]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$(x^2 + 1) x^2 y'' - x y' + (x + 2) y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned} & x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) (x^2 + 1) \\ & - x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (x+2) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \end{aligned} \tag{1}$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r+2} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) \\ & + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) + \left(\sum_{n=0}^{\infty} x^{1+n+r} a_n \right) + \left(\sum_{n=0}^{\infty} 2a_n x^{n+r} \right) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=0}^{\infty} x^{n+r+2} a_n (n+r) (n+r-1) &= \sum_{n=2}^{\infty} a_{n-2} (n+r-2) (n-3+r) x^{n+r} \\ \sum_{n=0}^{\infty} x^{1+n+r} a_n &= \sum_{n=1}^{\infty} a_{n-1} x^{n+r} \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=2}^{\infty} a_{n-2} (n+r-2) (n-3+r) x^{n+r} \right) + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) \\ & + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) + \left(\sum_{n=1}^{\infty} a_{n-1} x^{n+r} \right) + \left(\sum_{n=0}^{\infty} 2a_n x^{n+r} \right) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r} a_n (n+r) (n+r-1) - x^{n+r} a_n (n+r) + 2a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$x^r a_0 r (-1+r) - x^r a_0 r + 2a_0 x^r = 0$$

Or

$$(x^r r (-1+r) - x^r r + 2x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(r^2 - 2r + 2) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$r^2 - 2r + 2 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = 1 + i$$

$$r_2 = 1 - i$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(r^2 - 2r + 2) x^r = 0$$

Solving for r gives the roots of the indicial equation as Since the roots are complex conjugates, then two linearly independent solutions can be constructed using

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+1+i}$$

$$y_2(x) = \sum_{n=0}^{\infty} b_n x^{n+1-i}$$

$y_1(x)$ is found first. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = -\frac{1}{r^2 + 1}$$

For $2 \leq n$ the recursive equation is

$$a_{n-2}(n+r-2)(n-3+r) + a_n(n+r)(n+r-1) - a_n(n+r) + a_{n-1} + 2a_n = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{n^2 a_{n-2} + 2nra_{n-2} + r^2 a_{n-2} - 5na_{n-2} - 5ra_{n-2} + 6a_{n-2} + a_{n-1}}{n^2 + 2nr + r^2 - 2n - 2r + 2} \quad (4)$$

Which for the root $r = 1 + i$ becomes

$$a_n = \frac{(-n^2 + (3 - 2i)n - 1 + 3i)a_{n-2} - a_{n-1}}{n(n + 2i)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 1 + i$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{r^2+1}$	$-\frac{1}{5} + \frac{2i}{5}$

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{-r^4 + r^3 - r^2 + r + 1}{(r^2 + 1)(r^2 + 2r + 2)}$$

Which for the root $r = 1 + i$ becomes

$$a_2 = -\frac{1}{40} - \frac{13i}{40}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{r^2+1}$	$-\frac{1}{5} + \frac{2i}{5}$
a_2	$\frac{-r^4+r^3-r^2+r+1}{(r^2+1)(r^2+2r+2)}$	$-\frac{1}{40} - \frac{13i}{40}$

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{2r^4 + 2r^3 + 5r^2 + r - 1}{(r^2 + 1)(r^2 + 2r + 2)(r^2 + 4r + 5)}$$

Which for the root $r = 1 + i$ becomes

$$a_3 = \frac{71}{520} - \frac{17i}{520}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{r^2+1}$	$-\frac{1}{5} + \frac{2i}{5}$
a_2	$\frac{-r^4+r^3-r^2+r+1}{(r^2+1)(r^2+2r+2)}$	$-\frac{1}{40} - \frac{13i}{40}$
a_3	$\frac{2r^4+2r^3+5r^2+r-1}{(r^2+1)(r^2+2r+2)(r^2+4r+5)}$	$\frac{71}{520} - \frac{17i}{520}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{r^8 + 6r^7 + 13r^6 + 10r^5 - 4r^4 - 15r^3 - 37r^2 - 34r - 9}{(r^2 + 1)(r^2 + 2r + 2)(r^2 + 4r + 5)(r^2 + 6r + 10)}$$

Which for the root $r = 1 + i$ becomes

$$a_4 = -\frac{31}{832} + \frac{541i}{4160}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{r^2+1}$	$-\frac{1}{5} + \frac{2i}{5}$
a_2	$\frac{-r^4+r^3-r^2+r+1}{(r^2+1)(r^2+2r+2)}$	$-\frac{1}{40} - \frac{13i}{40}$
a_3	$\frac{2r^4+2r^3+5r^2+r-1}{(r^2+1)(r^2+2r+2)(r^2+4r+5)}$	$\frac{71}{520} - \frac{17i}{520}$
a_4	$\frac{r^8+6r^7+13r^6+10r^5-4r^4-15r^3-37r^2-34r-9}{(r^2+1)(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)}$	$-\frac{31}{832} + \frac{541i}{4160}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{-3r^8 - 30r^7 - 132r^6 - 330r^5 - 528r^4 - 570r^3 - 303r^2 + 60r + 69}{(r^2 + 1)(r^2 + 2r + 2)(r^2 + 4r + 5)(r^2 + 6r + 10)(r^2 + 8r + 17)}$$

Which for the root $r = 1 + i$ becomes

$$a_5 = -\frac{1423}{20800} - \frac{7i}{4160}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{r^2+1}$	$-\frac{1}{5} + \frac{2i}{5}$
a_2	$\frac{-r^4+r^3-r^2+r+1}{(r^2+1)(r^2+2r+2)}$	$-\frac{1}{40} - \frac{13i}{40}$
a_3	$\frac{2r^4+2r^3+5r^2+r-1}{(r^2+1)(r^2+2r+2)(r^2+4r+5)}$	$\frac{71}{520} - \frac{17i}{520}$
a_4	$\frac{r^8+6r^7+13r^6+10r^5-4r^4-15r^3-37r^2-34r-9}{(r^2+1)(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)}$	$-\frac{31}{832} + \frac{541i}{4160}$
a_5	$\frac{-3r^8-30r^7-132r^6-330r^5-528r^4-570r^3-303r^2+60r+69}{(r^2+1)(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)}$	$-\frac{1423}{20800} - \frac{7i}{4160}$

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{-r^{12} - 21r^{11} - 188r^{10} - 930r^9 - 2742r^8 - 4764r^7 - 4068r^6 + 1014r^5 + 8233r^4 + 14610r^3 + 15926r^2 + 8811r + 1767}{(r^2 + 1)(r^2 + 2r + 2)(r^2 + 4r + 5)(r^2 + 6r + 10)(r^2 + 8r + 17)(r^2 + 10r + 26)}$$

Which for the root $r = 1 + i$ becomes

$$a_6 = \frac{12849}{416000} - \frac{10853i}{156000}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{r^2+1}$	$-\frac{1}{5} + \frac{2i}{5}$
a_2	$\frac{-r^4+r^3-r^2+r+1}{(r^2+1)(r^2+2r+2)}$	$-\frac{1}{40} - \frac{13i}{40}$
a_3	$\frac{2r^4+2r^3+5r^2+r-1}{(r^2+1)(r^2+2r+2)(r^2+4r+5)}$	$\frac{71}{520} - \frac{17i}{520}$
a_4	$\frac{r^8+6r^7+13r^6+10r^5-4r^4-15r^3-37r^2-34r-9}{(r^2+1)(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)}$	$-\frac{31}{832} + \frac{541i}{4160}$
a_5	$\frac{-3r^8-30r^7-132r^6-330r^5-528r^4-570r^3-303r^2+60r+69}{(r^2+1)(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)}$	$-\frac{1423}{20800} - \frac{7i}{4160}$
a_6	$\frac{-r^{12}-21r^{11}-188r^{10}-930r^9-2742r^8-4764r^7-4068r^6+1014r^5+8233r^4+14610r^3+15926r^2+8811r+1767}{(r^2+1)(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)(r^2+10r+26)}$	$\frac{12849}{416000} - \frac{10853i}{156000}$

For $n = 7$, using the above recursive equation gives

$$a_7 = \frac{4r^{12} + 108r^{11} + 1298r^{10} + 9150r^9 + 42072r^8 + 133134r^7 + 298869r^6 + 482955r^5 + 553706r^4 + 40380r^3 + 209609r^2 + 106907r + 17808000}{(r^2 + 1)(r^2 + 2r + 2)(r^2 + 4r + 5)(r^2 + 6r + 10)(r^2 + 8r + 17)(r^2 + 10r + 26)}$$

Which for the root $r = 1 + i$ becomes

$$a_7 = \frac{209609}{5088000} + \frac{106907i}{17808000}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{r^2+1}$	$-\frac{1}{5} +$
a_2	$\frac{-r^4+r^3-r^2+r+1}{(r^2+1)(r^2+2r+2)}$	$-\frac{1}{40} -$
a_3	$\frac{2r^4+2r^3+5r^2+r-1}{(r^2+1)(r^2+2r+2)(r^2+4r+5)}$	$\frac{71}{520} -$
a_4	$\frac{r^8+6r^7+13r^6+10r^5-4r^4-15r^3-37r^2-34r-9}{(r^2+1)(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)}$	$-\frac{31}{832} -$
a_5	$\frac{-3r^8-30r^7-132r^6-330r^5-528r^4-570r^3-303r^2+60r+69}{(r^2+1)(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)}$	$-\frac{1423}{20800}$
a_6	$\frac{-r^{12}-21r^{11}-188r^{10}-930r^9-2742r^8-4764r^7-4068r^6+1014r^5+8233r^4+14610r^3+15926r^2+8811r+1767}{(r^2+1)(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)(r^2+10r+26)}$	$\frac{12849}{416000}$
a_7	$\frac{4r^{12}+108r^{11}+1298r^{10}+9150r^9+42072r^8+133134r^7+298869r^6+482955r^5+553706r^4+403821r^3+106210r^2-69957r-37647}{(r^2+1)(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)(r^2+10r+26)(r^2+12r+37)}$	$\frac{209609}{5088000}$

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned}
 y_1(x) &= x^{1+i}(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots) \\
 &= x^{1+i} \left(1 + \left(-\frac{1}{5} + \frac{2i}{5} \right) x + \left(-\frac{1}{40} - \frac{13i}{40} \right) x^2 + \left(\frac{71}{520} - \frac{17i}{520} \right) x^3 \right. \\
 &\quad + \left(-\frac{31}{832} + \frac{541i}{4160} \right) x^4 + \left(-\frac{1423}{20800} - \frac{7i}{4160} \right) x^5 + \left(\frac{12849}{416000} - \frac{10853i}{156000} \right) x^6 \\
 &\quad \left. + \left(\frac{209609}{5088000} + \frac{106907i}{17808000} \right) x^7 + O(x^8) \right)
 \end{aligned}$$

The second solution $y_2(x)$ is found by taking the complex conjugate of $y_1(x)$ which gives

$$\begin{aligned}
 y_2(x) &= x^{1-i} \left(1 + \left(-\frac{1}{5} - \frac{2i}{5} \right) x + \left(-\frac{1}{40} + \frac{13i}{40} \right) x^2 + \left(\frac{71}{520} + \frac{17i}{520} \right) x^3 \right. \\
 &\quad + \left(-\frac{31}{832} - \frac{541i}{4160} \right) x^4 + \left(-\frac{1423}{20800} + \frac{7i}{4160} \right) x^5 + \left(\frac{12849}{416000} + \frac{10853i}{156000} \right) x^6 \\
 &\quad \left. + \left(\frac{209609}{5088000} - \frac{106907i}{17808000} \right) x^7 + O(x^8) \right)
 \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 y_h(x) &= c_1 y_1(x) + c_2 y_2(x) \\
 &= c_1 x^{1+i} \left(1 + \left(-\frac{1}{5} + \frac{2i}{5} \right) x + \left(-\frac{1}{40} - \frac{13i}{40} \right) x^2 + \left(\frac{71}{520} - \frac{17i}{520} \right) x^3 \right. \\
 &\quad + \left(-\frac{31}{832} + \frac{541i}{4160} \right) x^4 + \left(-\frac{1423}{20800} - \frac{7i}{4160} \right) x^5 + \left(\frac{12849}{416000} - \frac{10853i}{156000} \right) x^6 \\
 &\quad \left. + \left(\frac{209609}{5088000} + \frac{106907i}{17808000} \right) x^7 + O(x^8) \right) \\
 &\quad + c_2 x^{1-i} \left(1 + \left(-\frac{1}{5} - \frac{2i}{5} \right) x + \left(-\frac{1}{40} + \frac{13i}{40} \right) x^2 + \left(\frac{71}{520} + \frac{17i}{520} \right) x^3 \right. \\
 &\quad + \left(-\frac{31}{832} - \frac{541i}{4160} \right) x^4 + \left(-\frac{1423}{20800} + \frac{7i}{4160} \right) x^5 + \left(\frac{12849}{416000} + \frac{10853i}{156000} \right) x^6 \\
 &\quad \left. + \left(\frac{209609}{5088000} - \frac{106907i}{17808000} \right) x^7 + O(x^8) \right)
 \end{aligned}$$

Hence the final solution is

$$\begin{aligned}
 y &= y_h \\
 &= c_1 x^{1+i} \left(1 + \left(-\frac{1}{5} + \frac{2i}{5} \right) x + \left(-\frac{1}{40} - \frac{13i}{40} \right) x^2 + \left(\frac{71}{520} - \frac{17i}{520} \right) x^3 + \left(-\frac{31}{832} + \frac{541i}{4160} \right) x^4 \right. \\
 &\quad + \left(-\frac{1423}{20800} - \frac{7i}{4160} \right) x^5 + \left(\frac{12849}{416000} - \frac{10853i}{156000} \right) x^6 + \left(\frac{209609}{5088000} + \frac{106907i}{17808000} \right) x^7 \\
 &\quad \left. + O(x^8) \right) + c_2 x^{1-i} \left(1 + \left(-\frac{1}{5} - \frac{2i}{5} \right) x + \left(-\frac{1}{40} + \frac{13i}{40} \right) x^2 + \left(\frac{71}{520} + \frac{17i}{520} \right) x^3 \right. \\
 &\quad + \left(-\frac{31}{832} - \frac{541i}{4160} \right) x^4 + \left(-\frac{1423}{20800} + \frac{7i}{4160} \right) x^5 + \left(\frac{12849}{416000} + \frac{10853i}{156000} \right) x^6 \\
 &\quad \left. + \left(\frac{209609}{5088000} - \frac{106907i}{17808000} \right) x^7 + O(x^8) \right)
 \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned} y = & c_1 x^{1+i} \left(1 + \left(-\frac{1}{5} + \frac{2i}{5} \right) x + \left(-\frac{1}{40} - \frac{13i}{40} \right) x^2 + \left(\frac{71}{520} - \frac{17i}{520} \right) x^3 \right. \\ & + \left(-\frac{31}{832} + \frac{541i}{4160} \right) x^4 + \left(-\frac{1423}{20800} - \frac{7i}{4160} \right) x^5 + \left(\frac{12849}{416000} - \frac{10853i}{156000} \right) x^6 \\ & \left. + \left(\frac{209609}{5088000} + \frac{106907i}{17808000} \right) x^7 + O(x^8) \right) \\ & + c_2 x^{1-i} \left(1 + \left(-\frac{1}{5} - \frac{2i}{5} \right) x + \left(-\frac{1}{40} + \frac{13i}{40} \right) x^2 + \left(\frac{71}{520} + \frac{17i}{520} \right) x^3 \right. \\ & + \left(-\frac{31}{832} - \frac{541i}{4160} \right) x^4 + \left(-\frac{1423}{20800} + \frac{7i}{4160} \right) x^5 + \left(\frac{12849}{416000} + \frac{10853i}{156000} \right) x^6 \\ & \left. + \left(\frac{209609}{5088000} - \frac{106907i}{17808000} \right) x^7 + O(x^8) \right) \end{aligned}$$

Verification of solutions

$$\begin{aligned} y = & c_1 x^{1+i} \left(1 + \left(-\frac{1}{5} + \frac{2i}{5} \right) x + \left(-\frac{1}{40} - \frac{13i}{40} \right) x^2 + \left(\frac{71}{520} - \frac{17i}{520} \right) x^3 + \left(-\frac{31}{832} + \frac{541i}{4160} \right) x^4 \right. \\ & + \left(-\frac{1423}{20800} - \frac{7i}{4160} \right) x^5 + \left(\frac{12849}{416000} - \frac{10853i}{156000} \right) x^6 + \left(\frac{209609}{5088000} + \frac{106907i}{17808000} \right) x^7 \\ & \left. + O(x^8) \right) + c_2 x^{1-i} \left(1 + \left(-\frac{1}{5} - \frac{2i}{5} \right) x + \left(-\frac{1}{40} + \frac{13i}{40} \right) x^2 + \left(\frac{71}{520} + \frac{17i}{520} \right) x^3 \right. \\ & + \left(-\frac{31}{832} - \frac{541i}{4160} \right) x^4 + \left(-\frac{1423}{20800} + \frac{7i}{4160} \right) x^5 + \left(\frac{12849}{416000} + \frac{10853i}{156000} \right) x^6 \\ & \left. + \left(\frac{209609}{5088000} - \frac{106907i}{17808000} \right) x^7 + O(x^8) \right) \end{aligned}$$

Verified OK.

22.9.1 Maple step by step solution

Let's solve

$$x^2 y''(x^2 + 1) - xy' + (x + 2)y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{(x+2)y}{(x^2+1)x^2} + \frac{y'}{x(x^2+1)}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - \frac{y'}{x(x^2+1)} + \frac{(x+2)y}{(x^2+1)x^2} = 0$$

- Check to see if x_0 is a regular singular point

- Define functions

$$\left[P_2(x) = -\frac{1}{x(x^2+1)}, P_3(x) = \frac{x+2}{(x^2+1)x^2} \right]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = -1$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 2$$

- $x = 0$ is a regular singular point

Check to see if x_0 is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$x^2 y''(x^2 + 1) - xy' + (x + 2)y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 0..1$

$$x^m \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

- Shift index using $k \rightarrow k - m$

$$x^m \cdot y = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

- Convert $x \cdot y'$ to series expansion

$$x \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r}$$

- Convert $x^m \cdot y''$ to series expansion for $m = 2..4$

$$x^m \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r-2+m}$$

- Shift index using $k \rightarrow k + 2 - m$

$$x^m \cdot y'' = \sum_{k=-2+m}^{\infty} a_{k+2-m} (k+2-m+r) (k+1-m+r) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(r^2 - 2r + 2) x^r + ((r^2 + 1) a_1 + a_0) x^{1+r} + \left(\sum_{k=2}^{\infty} (a_k(k^2 + 2kr + r^2 - 2k - 2r + 2) + a_{k-1} + a_{k+1}) x^{k+r} \right)$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$r^2 - 2r + 2 = 0$$

- Values of r that satisfy the indicial equation

$$r \in \{1 - I, 1 + I\}$$

- Each term must be 0

$$(r^2 + 1) a_1 + a_0 = 0$$

- Solve for the dependent coefficient(s)

$$a_1 = -\frac{a_0}{r^2 + 1}$$

- Each term in the series must be 0, giving the recursion relation

$$a_{k-2}(k-2+r)(k-3+r) + (k^2 + (2r-2)k + r^2 - 2r + 2) a_k + a_{k-1} = 0$$

- Shift index using $k \rightarrow k + 2$

$$a_k(k+r)(k+r-1) + ((k+2)^2 + (2r-2)(k+2) + r^2 - 2r + 2) a_{k+2} + a_{k+1} = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+2} = -\frac{k^2 a_k + 2k r a_k + r^2 a_k - k a_k - r a_k + a_{k+1}}{k^2 + 2k r + r^2 + 2k + 2r + 2}$$

- Recursion relation for $r = 1 - I$

$$a_{k+2} = -\frac{k^2 a_k + (2-2I)k a_k - (1+I)a_k - k a_k + a_{k+1}}{k^2 + (2-2I)k + 4 - 4I + 2k}$$

- Solution for $r = 1 - I$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+1-I}, a_{k+2} = -\frac{k^2 a_k + (2-2I)k a_k - (1+I)a_k - k a_k + a_{k+1}}{k^2 + (2-2I)k + 4 - 4I + 2k}, a_1 = \left(-\frac{1}{5} - \frac{2I}{5}\right) a_0 \right]$$

- Recursion relation for $r = 1 + I$

$$a_{k+2} = -\frac{k^2 a_k + (2+2I)k a_k + (-1+I)a_k - k a_k + a_{k+1}}{k^2 + (2+2I)k + 4 + 4I + 2k}$$

- Solution for $r = 1 + I$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+1+I}, a_{k+2} = -\frac{k^2 a_k + (2+2I)k a_k + (-1+I)a_k - k a_k + a_{k+1}}{k^2 + (2+2I)k + 4 + 4I + 2k}, a_1 = \left(-\frac{1}{5} + \frac{2I}{5}\right) a_0 \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^{k+1-I} \right) + \left(\sum_{k=0}^{\infty} b_k x^{k+1+I} \right), a_{k+2} = -\frac{k^2 a_k + (2-2I)k a_k - (1+I)a_k - k a_k + a_{k+1}}{k^2 + (2-2I)k + 4 - 4I + 2k}, a_1 = \left(-\frac{1}{5} - \frac{2I}{5} \right) \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
  -> Bessel
  -> elliptic
  -> Legendre
  -> Kummer
    -> hyper3: Equivalence to 1F1 under a power @ Moebius
  -> hypergeometric
    -> heuristic approach
    -> hyper3: Equivalence to 2F1, 1F1 or 0F1 under a power @ Moebius
  -> Mathieu
    -> Equivalence to the rational form of Mathieu ODE under a power @ Moebius
trying a solution in terms of MeijerG functions
-> Heun: Equivalence to the GHE or one of its 4 confluent cases under a power @ Moebius
<- Heun successful: received ODE is equivalent to the HeunG ODE, case a <> 0, e <> 0, g <> 0

```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 87

Order:=8;

dsolve((x^2+1)*x^2*diff(y(x),x\$2)-x*diff(y(x),x)+(2+x)*y(x)=0,y(x),type='series',x=0);

$$\begin{aligned}
 y(x) = & c_1 x^{1-i} \left(1 + \left(-\frac{1}{5} - \frac{2i}{5} \right) x + \left(-\frac{1}{40} + \frac{13i}{40} \right) x^2 + \left(\frac{71}{520} + \frac{17i}{520} \right) x^3 \right. \\
 & + \left(-\frac{31}{832} - \frac{541i}{4160} \right) x^4 + \left(-\frac{1423}{20800} + \frac{7i}{4160} \right) x^5 + \left(\frac{12849}{416000} + \frac{10853i}{156000} \right) x^6 \\
 & \left. + \left(\frac{209609}{5088000} - \frac{106907i}{17808000} \right) x^7 + O(x^8) \right) \\
 & + c_2 x^{1+i} \left(1 + \left(-\frac{1}{5} + \frac{2i}{5} \right) x + \left(-\frac{1}{40} - \frac{13i}{40} \right) x^2 + \left(\frac{71}{520} - \frac{17i}{520} \right) x^3 \right. \\
 & + \left(-\frac{31}{832} + \frac{541i}{4160} \right) x^4 + \left(-\frac{1423}{20800} - \frac{7i}{4160} \right) x^5 + \left(\frac{12849}{416000} - \frac{10853i}{156000} \right) x^6 \\
 & \left. + \left(\frac{209609}{5088000} + \frac{106907i}{17808000} \right) x^7 + O(x^8) \right)
 \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.055 (sec). Leaf size: 122

AsymptoticDSolveValue[(x^2+1)*x^2*y'[x]-x*y'[x]+(2+x)*y[x]==0,y[x],{x,0,7}]

$$\begin{aligned}
 y(x) \rightarrow & \left(\frac{1}{156000} + \frac{i}{1248000} \right) c_2 x^{1-i} \left((6080 + 10093i)x^6 - (10476 - 1572i)x^5 \right. \\
 & - (8220 + 19260i)x^4 + (21600 + 2400i)x^3 + (2400 + 50400i)x^2 - (38400 + 57600i)x \\
 & \left. + (153600 - 19200i) \right) - \left(\frac{1}{1248000} + \frac{i}{156000} \right) c_1 x^{1+i} \left((10093 + 6080i)x^6 \right. \\
 & + (1572 - 10476i)x^5 - (19260 + 8220i)x^4 + (2400 + 21600i)x^3 \\
 & \left. + (50400 + 2400i)x^2 - (57600 + 38400i)x - (19200 - 153600i) \right)
 \end{aligned}$$

22.10 problem 2(b)

22.10.1 Maple step by step solution 3704

Internal problem ID [6485]

Internal file name [OUTPUT/5733_Sunday_June_05_2022_03_51_25_PM_79495257/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Problems for review and discover. (A) Drill Exercises . Page 194

Problem number: 2(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Complex roots"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$x^2y'' + xy' + (1 + x)y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2y'' + xy' + (1 + x)y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{1}{x}$$
$$q(x) = \frac{1 + x}{x^2}$$

Table 502: Table $p(x), q(x)$ singularities.

$p(x) = \frac{1}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = \frac{1+x}{x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2 y'' + x y' + (1 + x) y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned}
 & x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\
 & + x \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + (1+x) \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0
 \end{aligned} \tag{1}$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) \right) \\ & + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \left(\sum_{n=0}^{\infty} x^{1+n+r} a_n \right) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} x^{1+n+r} a_n = \sum_{n=1}^{\infty} a_{n-1} x^{n+r}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) \right) \\ & + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) + \left(\sum_{n=1}^{\infty} a_{n-1} x^{n+r} \right) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r} a_n (n+r) (n+r-1) + x^{n+r} a_n (n+r) + a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$x^r a_0 r(-1+r) + x^r a_0 r + a_0 x^r = 0$$

Or

$$(x^r r(-1+r) + x^r r + x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(r^2 + 1) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$r^2 + 1 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= i \\ r_2 &= -i \end{aligned}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(r^2 + 1) x^r = 0$$

Solving for r gives the roots of the indicial equation as Since the roots are complex conjugates, then two linearly independent solutions can be constructed using

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^{n+i} \\ y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n-i} \end{aligned}$$

$y_1(x)$ is found first. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) + a_n(n+r) + a_n + a_{n-1} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{a_{n-1}}{n^2 + 2nr + r^2 + 1} \quad (4)$$

Which for the root $r = i$ becomes

$$a_n = -\frac{a_{n-1}}{n(2i+n)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = i$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = -\frac{1}{r^2 + 2r + 2}$$

Which for the root $r = i$ becomes

$$a_1 = -\frac{1}{5} + \frac{2i}{5}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{r^2+2r+2}$	$-\frac{1}{5} + \frac{2i}{5}$

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{(r^2 + 2r + 2)(r^2 + 4r + 5)}$$

Which for the root $r = i$ becomes

$$a_2 = -\frac{1}{40} - \frac{3i}{40}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{r^2+2r+2}$	$-\frac{1}{5} + \frac{2i}{5}$
a_2	$\frac{1}{(r^2+2r+2)(r^2+4r+5)}$	$-\frac{1}{40} - \frac{3i}{40}$

For $n = 3$, using the above recursive equation gives

$$a_3 = -\frac{1}{(r^2 + 2r + 2)(r^2 + 4r + 5)(r^2 + 6r + 10)}$$

Which for the root $r = i$ becomes

$$a_3 = \frac{3}{520} + \frac{7i}{1560}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{r^2+2r+2}$	$-\frac{1}{5} + \frac{2i}{5}$
a_2	$\frac{1}{(r^2+2r+2)(r^2+4r+5)}$	$-\frac{1}{40} - \frac{3i}{40}$
a_3	$-\frac{1}{(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)}$	$\frac{3}{520} + \frac{7i}{1560}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{(r^2 + 2r + 2)(r^2 + 4r + 5)(r^2 + 6r + 10)(r^2 + 8r + 17)}$$

Which for the root $r = i$ becomes

$$a_4 = -\frac{1}{2496} - \frac{i}{12480}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{r^2+2r+2}$	$-\frac{1}{5} + \frac{2i}{5}$
a_2	$\frac{1}{(r^2+2r+2)(r^2+4r+5)}$	$-\frac{1}{40} - \frac{3i}{40}$
a_3	$-\frac{1}{(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)}$	$\frac{3}{520} + \frac{7i}{1560}$
a_4	$\frac{1}{(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)}$	$-\frac{1}{2496} - \frac{i}{12480}$

For $n = 5$, using the above recursive equation gives

$$a_5 = -\frac{1}{(r^2 + 2r + 2)(r^2 + 4r + 5)(r^2 + 6r + 10)(r^2 + 8r + 17)(r^2 + 10r + 26)}$$

Which for the root $r = i$ becomes

$$a_5 = \frac{9}{603200} - \frac{i}{361920}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{r^2+2r+2}$	$-\frac{1}{5} + \frac{2i}{5}$
a_2	$\frac{1}{(r^2+2r+2)(r^2+4r+5)}$	$-\frac{1}{40} - \frac{3i}{40}$
a_3	$-\frac{1}{(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)}$	$\frac{3}{520} + \frac{7i}{1560}$
a_4	$\frac{1}{(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)}$	$-\frac{1}{2496} - \frac{i}{12480}$
a_5	$-\frac{1}{(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)(r^2+10r+26)}$	$\frac{9}{603200} - \frac{i}{361920}$

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{1}{(r^2 + 2r + 2)(r^2 + 4r + 5)(r^2 + 6r + 10)(r^2 + 8r + 17)(r^2 + 10r + 26)(r^2 + 12r + 37)}$$

Which for the root $r = i$ becomes

$$a_6 = -\frac{19}{54288000} + \frac{7i}{36192000}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{r^2+2r+2}$	$-\frac{1}{5} + \frac{2i}{5}$
a_2	$\frac{1}{(r^2+2r+2)(r^2+4r+5)}$	$-\frac{1}{40} - \frac{3i}{40}$
a_3	$-\frac{1}{(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)}$	$\frac{3}{520} + \frac{7i}{1560}$
a_4	$\frac{1}{(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)}$	$-\frac{1}{2496} - \frac{i}{12480}$
a_5	$-\frac{1}{(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)(r^2+10r+26)}$	$\frac{9}{603200} - \frac{i}{361920}$
a_6	$\frac{1}{(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)(r^2+10r+26)(r^2+12r+37)}$	$-\frac{19}{54288000} + \frac{7i}{36192000}$

For $n = 7$, using the above recursive equation gives

$$a_7 = -\frac{1}{(r^2 + 2r + 2)(r^2 + 4r + 5)(r^2 + 6r + 10)(r^2 + 8r + 17)(r^2 + 10r + 26)(r^2 + 12r + 37)(r^2 + 14r + 26)}$$

Which for the root $r = i$ becomes

$$a_7 = \frac{1}{179829000} - \frac{223i}{40281696000}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{r^2+2r+2}$	$-\frac{1}{5} + \frac{2i}{5}$
a_2	$\frac{1}{(r^2+2r+2)(r^2+4r+5)}$	$-\frac{1}{40} - \frac{3i}{40}$
a_3	$-\frac{1}{(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)}$	$\frac{3}{520} + \frac{7i}{1560}$
a_4	$\frac{1}{(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)}$	$-\frac{1}{2496} - \frac{i}{12480}$
a_5	$-\frac{1}{(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)(r^2+10r+26)}$	$\frac{9}{603200} - \frac{i}{361920}$
a_6	$\frac{1}{(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)(r^2+10r+26)(r^2+12r+37)}$	$-\frac{19}{54288000} + \frac{7i}{36192000}$
a_7	$-\frac{1}{(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)(r^2+10r+26)(r^2+12r+37)(r^2+14r+50)}$	$\frac{1}{179829000} - \frac{223i}{40281696000}$

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x^i (a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots) \\ &= x^i \left(1 + \left(-\frac{1}{5} + \frac{2i}{5} \right) x + \left(-\frac{1}{40} - \frac{3i}{40} \right) x^2 + \left(\frac{3}{520} + \frac{7i}{1560} \right) x^3 \right. \\ &\quad \left. + \left(-\frac{1}{2496} - \frac{i}{12480} \right) x^4 + \left(\frac{9}{603200} - \frac{i}{361920} \right) x^5 \right. \\ &\quad \left. + \left(-\frac{19}{54288000} + \frac{7i}{36192000} \right) x^6 + \left(\frac{1}{179829000} - \frac{223i}{40281696000} \right) x^7 + O(x^8) \right) \end{aligned}$$

The second solution $y_2(x)$ is found by taking the complex conjugate of $y_1(x)$ which gives

$$\begin{aligned} y_2(x) &= x^{-i} \left(1 + \left(-\frac{1}{5} - \frac{2i}{5} \right) x + \left(-\frac{1}{40} + \frac{3i}{40} \right) x^2 + \left(\frac{3}{520} - \frac{7i}{1560} \right) x^3 \right. \\ &\quad \left. + \left(-\frac{1}{2496} + \frac{i}{12480} \right) x^4 + \left(\frac{9}{603200} + \frac{i}{361920} \right) x^5 \right. \\ &\quad \left. + \left(-\frac{19}{54288000} - \frac{7i}{36192000} \right) x^6 + \left(\frac{1}{179829000} + \frac{223i}{40281696000} \right) x^7 + O(x^8) \right) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 y_h(x) &= c_1 y_1(x) + c_2 y_2(x) \\
 &= c_1 x^i \left(1 + \left(-\frac{1}{5} + \frac{2i}{5} \right) x + \left(-\frac{1}{40} - \frac{3i}{40} \right) x^2 + \left(\frac{3}{520} + \frac{7i}{1560} \right) x^3 \right. \\
 &\quad \left. + \left(-\frac{1}{2496} - \frac{i}{12480} \right) x^4 + \left(\frac{9}{603200} - \frac{i}{361920} \right) x^5 \right. \\
 &\quad \left. + \left(-\frac{19}{54288000} + \frac{7i}{36192000} \right) x^6 + \left(\frac{1}{179829000} - \frac{223i}{40281696000} \right) x^7 + O(x^8) \right) \\
 &+ c_2 x^{-i} \left(1 + \left(-\frac{1}{5} - \frac{2i}{5} \right) x + \left(-\frac{1}{40} + \frac{3i}{40} \right) x^2 + \left(\frac{3}{520} - \frac{7i}{1560} \right) x^3 \right. \\
 &\quad \left. + \left(-\frac{1}{2496} + \frac{i}{12480} \right) x^4 + \left(\frac{9}{603200} + \frac{i}{361920} \right) x^5 \right. \\
 &\quad \left. + \left(-\frac{19}{54288000} - \frac{7i}{36192000} \right) x^6 + \left(\frac{1}{179829000} + \frac{223i}{40281696000} \right) x^7 + O(x^8) \right)
 \end{aligned}$$

Hence the final solution is

$$\begin{aligned}
 y &= y_h \\
 &= c_1 x^i \left(1 + \left(-\frac{1}{5} + \frac{2i}{5} \right) x + \left(-\frac{1}{40} - \frac{3i}{40} \right) x^2 + \left(\frac{3}{520} + \frac{7i}{1560} \right) x^3 \right. \\
 &\quad \left. + \left(-\frac{1}{2496} - \frac{i}{12480} \right) x^4 + \left(\frac{9}{603200} - \frac{i}{361920} \right) x^5 \right. \\
 &\quad \left. + \left(-\frac{19}{54288000} + \frac{7i}{36192000} \right) x^6 + \left(\frac{1}{179829000} - \frac{223i}{40281696000} \right) x^7 + O(x^8) \right) \\
 &+ c_2 x^{-i} \left(1 + \left(-\frac{1}{5} - \frac{2i}{5} \right) x + \left(-\frac{1}{40} + \frac{3i}{40} \right) x^2 + \left(\frac{3}{520} - \frac{7i}{1560} \right) x^3 \right. \\
 &\quad \left. + \left(-\frac{1}{2496} + \frac{i}{12480} \right) x^4 + \left(\frac{9}{603200} + \frac{i}{361920} \right) x^5 \right. \\
 &\quad \left. + \left(-\frac{19}{54288000} - \frac{7i}{36192000} \right) x^6 + \left(\frac{1}{179829000} + \frac{223i}{40281696000} \right) x^7 + O(x^8) \right)
 \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}y = c_1 x^i & \left(1 + \left(-\frac{1}{5} + \frac{2i}{5} \right) x + \left(-\frac{1}{40} - \frac{3i}{40} \right) x^2 + \left(\frac{3}{520} + \frac{7i}{1560} \right) x^3 \right. \\ & \left. + \left(-\frac{1}{2496} - \frac{i}{12480} \right) x^4 + \left(\frac{9}{603200} - \frac{i}{361920} \right) x^5 \right. \\ & \left. + \left(-\frac{19}{54288000} + \frac{7i}{36192000} \right) x^6 + \left(\frac{1}{179829000} - \frac{223i}{40281696000} \right) x^7 + O(x^8) \right) \\ & + c_2 x^{-i} \left(1 + \left(-\frac{1}{5} - \frac{2i}{5} \right) x + \left(-\frac{1}{40} + \frac{3i}{40} \right) x^2 + \left(\frac{3}{520} - \frac{7i}{1560} \right) x^3 \right. \\ & \left. + \left(-\frac{1}{2496} + \frac{i}{12480} \right) x^4 + \left(\frac{9}{603200} + \frac{i}{361920} \right) x^5 \right. \\ & \left. + \left(-\frac{19}{54288000} - \frac{7i}{36192000} \right) x^6 + \left(\frac{1}{179829000} + \frac{223i}{40281696000} \right) x^7 + O(x^8) \right)\end{aligned}$$

Verification of solutions

$$\begin{aligned}y = c_1 x^i & \left(1 + \left(-\frac{1}{5} + \frac{2i}{5} \right) x + \left(-\frac{1}{40} - \frac{3i}{40} \right) x^2 + \left(\frac{3}{520} + \frac{7i}{1560} \right) x^3 \right. \\ & \left. + \left(-\frac{1}{2496} - \frac{i}{12480} \right) x^4 + \left(\frac{9}{603200} - \frac{i}{361920} \right) x^5 \right. \\ & \left. + \left(-\frac{19}{54288000} + \frac{7i}{36192000} \right) x^6 + \left(\frac{1}{179829000} - \frac{223i}{40281696000} \right) x^7 + O(x^8) \right) \\ & + c_2 x^{-i} \left(1 + \left(-\frac{1}{5} - \frac{2i}{5} \right) x + \left(-\frac{1}{40} + \frac{3i}{40} \right) x^2 + \left(\frac{3}{520} - \frac{7i}{1560} \right) x^3 \right. \\ & \left. + \left(-\frac{1}{2496} + \frac{i}{12480} \right) x^4 + \left(\frac{9}{603200} + \frac{i}{361920} \right) x^5 \right. \\ & \left. + \left(-\frac{19}{54288000} - \frac{7i}{36192000} \right) x^6 + \left(\frac{1}{179829000} + \frac{223i}{40281696000} \right) x^7 + O(x^8) \right)\end{aligned}$$

Verified OK.

22.10.1 Maple step by step solution

Let's solve

$$x^2 y'' + xy' + (1+x)y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{y'}{x} - \frac{(1+x)y}{x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{y'}{x} + \frac{(1+x)y}{x^2} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$[P_2(x) = \frac{1}{x}, P_3(x) = \frac{1+x}{x^2}]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 1$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 1$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$x^2 y'' + x y' + (1+x)y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y$ to series expansion for $m = 0..1$

$$x^m \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+m}$$

- Shift index using $k \rightarrow k - m$

$$x^m \cdot y = \sum_{k=m}^{\infty} a_{k-m} x^{k+r}$$

- Convert $x \cdot y'$ to series expansion

$$x \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r}$$

- Convert $x^2 \cdot y''$ to series expansion

$$x^2 \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(r^2 + 1) x^r + \left(\sum_{k=1}^{\infty} (a_k(k^2 + 2kr + r^2 + 1) + a_{k-1}) x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation
 $r^2 + 1 = 0$
- Values of r that satisfy the indicial equation
 $r \in \{-I, I\}$

- Each term in the series must be 0, giving the recursion relation

$$a_k(k^2 + 2kr + r^2 + 1) + a_{k-1} = 0$$

- Shift index using $k- \rightarrow k + 1$

$$a_{k+1}((k+1)^2 + 2(k+1)r + r^2 + 1) + a_k = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+1} = -\frac{a_k}{k^2 + 2kr + r^2 + 2k + 2r + 2}$$

- Recursion relation for $r = -I$

$$a_{k+1} = -\frac{a_k}{k^2 - 2Ik + 1 - 2I + 2k}$$

- Solution for $r = -I$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k-I}, a_{k+1} = -\frac{a_k}{k^2 - 2Ik + 1 - 2I + 2k} \right]$$

- Recursion relation for $r = I$

$$a_{k+1} = -\frac{a_k}{k^2 + 2Ik + 1 + 2I + 2k}$$

- Solution for $r = I$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+I}, a_{k+1} = -\frac{a_k}{k^2 + 2Ik + 1 + 2I + 2k} \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^{k-I} \right) + \left(\sum_{k=0}^{\infty} b_k x^{k+I} \right), a_{k+1} = -\frac{a_k}{k^2 - 2Ik + 1 - 2I + 2k}, b_{k+1} = -\frac{b_k}{k^2 + 2Ik + 1 + 2I + 2k} \right]$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacic's algorithm  
<- No Liouvillian solutions exist  
-> Trying a solution in terms of special functions:  
  -> Bessel  
  <- Bessel successful  
<- special function solution successful`
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 85

```
Order:=8;  
dsolve(x^2*dif(y(x),x$2)+x*dif(y(x),x)+(1+x)*y(x)=0,y(x),type='series',x=0);
```

$$\begin{aligned} y(x) = & c_1 x^{-i} \left(1 + \left(-\frac{1}{5} - \frac{2i}{5} \right) x + \left(-\frac{1}{40} + \frac{3i}{40} \right) x^2 + \left(\frac{3}{520} - \frac{7i}{1560} \right) x^3 \right. \\ & \left. + \left(-\frac{1}{2496} + \frac{i}{12480} \right) x^4 + \left(\frac{9}{603200} + \frac{i}{361920} \right) x^5 \right. \\ & \left. + \left(-\frac{19}{54288000} - \frac{7i}{36192000} \right) x^6 + \left(\frac{1}{179829000} + \frac{223i}{40281696000} \right) x^7 + O(x^8) \right) \\ & + c_2 x^i \left(1 + \left(-\frac{1}{5} + \frac{2i}{5} \right) x + \left(-\frac{1}{40} - \frac{3i}{40} \right) x^2 + \left(\frac{3}{520} + \frac{7i}{1560} \right) x^3 \right. \\ & \left. + \left(-\frac{1}{2496} - \frac{i}{12480} \right) x^4 + \left(\frac{9}{603200} - \frac{i}{361920} \right) x^5 \right. \\ & \left. + \left(-\frac{19}{54288000} + \frac{7i}{36192000} \right) x^6 + \left(\frac{1}{179829000} - \frac{223i}{40281696000} \right) x^7 + O(x^8) \right) \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.016 (sec). Leaf size: 118

```
AsymptoticDSolveValue[x^2*y''[x]+x*y'[x]+(1+x)*y[x]==0,y[x],{x,0,7}]
```

$$\begin{aligned} y(x) \rightarrow & \left(\frac{7}{36192000} + \frac{19i}{54288000} \right) c_1 x^i (ix^6 + (12 - 36i)x^5 - (660 - 780i)x^4 \\ & + (16800 - 7200i)x^3 - (194400 + 36000i)x^2 + (633600 + 921600i)x \\ & + (1209600 - 2188800i)) - \left(\frac{19}{54288000} + \frac{7i}{36192000} \right) c_2 x^{-i} (x^6 - (36 - 12i)x^5 \\ & + (780 - 660i)x^4 - (7200 - 16800i)x^3 - (36000 + 194400i)x^2 \\ & + (921600 + 633600i)x - (2188800 - 1209600i)) \end{aligned}$$

22.11 problem 2(c)

22.11.1 Maple step by step solution 3720

Internal problem ID [6486]

Internal file name [OUTPUT/5734_Sunday_June_05_2022_03_51_33_PM_34104880/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Problems for review and discover. (A) Drill Exercises . Page 194

Problem number: 2(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference is integer"**

Maple gives the following as the ode type

[_Lienard]

$$xy'' - 4y' + xy = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$xy'' - 4y' + xy = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{4}{x}$$
$$q(x) = 1$$

Table 504: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{4}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = 1$	
singularity	type

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$xy'' - 4y' + xy = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) x - 4 \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + x \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} x^{n+r-1} a_n (n+r)(n+r-1) \right) + \sum_{n=0}^{\infty} (-4(n+r) a_n x^{n+r-1}) + \left(\sum_{n=0}^{\infty} x^{1+n+r} a_n \right) = 0 \quad (2A)$$

The next step is to make all powers of x be $n + r - 1$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r-1} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} x^{1+n+r} a_n = \sum_{n=2}^{\infty} a_{n-2} x^{n+r-1}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n + r - 1$.

$$\left(\sum_{n=0}^{\infty} x^{n+r-1} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-4(n+r) a_n x^{n+r-1}) + \left(\sum_{n=2}^{\infty} a_{n-2} x^{n+r-1} \right) = 0 \quad (2B)$$

The indicial equation is obtained from $n = 0$. From Eq (2B) this gives

$$x^{n+r-1} a_n (n+r) (n+r-1) - 4(n+r) a_n x^{n+r-1} = 0$$

When $n = 0$ the above becomes

$$x^{-1+r} a_0 r (-1+r) - 4r a_0 x^{-1+r} = 0$$

Or

$$(x^{-1+r} r (-1+r) - 4r x^{-1+r}) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$r x^{-1+r} (-5+r) = 0$$

Since the above is true for all x then the indicial equation becomes

$$r(-5+r) = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = 5$$

$$r_2 = 0$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$r x^{-1+r} (-5+r) = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = 5$ is an integer, then we can construct two linearly independent solutions

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = C y_1(x) \ln(x) + x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = x^5 \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+5}$$

$$y_2(x) = C y_1(x) \ln(x) + \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Where C above can be zero. We start by finding y_1 . Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

For $2 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) - 4a_n(n+r) + a_{n-2} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{a_{n-2}}{n^2 + 2nr + r^2 - 5n - 5r} \quad (4)$$

Which for the root $r = 5$ becomes

$$a_n = -\frac{a_{n-2}}{n(n+5)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 5$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = -\frac{1}{r^2 - r - 6}$$

Which for the root $r = 5$ becomes

$$a_2 = -\frac{1}{14}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2 - r - 6}$	$-\frac{1}{14}$

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2 - r - 6}$	$-\frac{1}{14}$
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{r^4 + 2r^3 - 13r^2 - 14r + 24}$$

Which for the root $r = 5$ becomes

$$a_4 = \frac{1}{504}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2-r-6}$	$-\frac{1}{14}$
a_3	0	0
a_4	$\frac{1}{r^4+2r^3-13r^2-14r+24}$	$\frac{1}{504}$

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2-r-6}$	$-\frac{1}{14}$
a_3	0	0
a_4	$\frac{1}{r^4+2r^3-13r^2-14r+24}$	$\frac{1}{504}$
a_5	0	0

For $n = 6$, using the above recursive equation gives

$$a_6 = -\frac{1}{r^6 + 9r^5 + 7r^4 - 93r^3 - 152r^2 + 84r + 144}$$

Which for the root $r = 5$ becomes

$$a_6 = -\frac{1}{33264}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2-r-6}$	$-\frac{1}{14}$
a_3	0	0
a_4	$\frac{1}{r^4+2r^3-13r^2-14r+24}$	$\frac{1}{504}$
a_5	0	0
a_6	$-\frac{1}{r^6+9r^5+7r^4-93r^3-152r^2+84r+144}$	$-\frac{1}{33264}$

For $n = 7$, using the above recursive equation gives

$$a_7 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{1}{r^2-r-6}$	$-\frac{1}{14}$
a_3	0	0
a_4	$\frac{1}{r^4+2r^3-13r^2-14r+24}$	$\frac{1}{504}$
a_5	0	0
a_6	$-\frac{1}{r^6+9r^5+7r^4-93r^3-152r^2+84r+144}$	$-\frac{1}{33264}$
a_7	0	0

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x^5(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots) \\ &= x^5\left(1 - \frac{x^2}{14} + \frac{x^4}{504} - \frac{x^6}{33264} + O(x^8)\right) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Let

$$r_1 - r_2 = N$$

Where N is positive integer which is the difference between the two roots. r_1 is taken as the larger root. Hence for this problem we have $N = 5$. Now we need to determine if

C is zero or not. This is done by finding $\lim_{r \rightarrow r_2} a_5(r)$. If this limit exists, then $C = 0$, else we need to keep the log term and $C \neq 0$. The above table shows that

$$\begin{aligned} a_N &= a_5 \\ &= 0 \end{aligned}$$

Therefore

$$\begin{aligned} \lim_{r \rightarrow r_2} 0 &= \lim_{r \rightarrow 0} 0 \\ &= 0 \end{aligned}$$

The limit is 0. Since the limit exists then the log term is not needed and we can set $C = 0$. Therefore the second solution has the form

$$\begin{aligned} y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n+r} \\ &= \sum_{n=0}^{\infty} b_n x^n \end{aligned}$$

Eq (3) derived above is used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq(3) gives

$$b_1 = 0$$

For $2 \leq n$ the recursive equation is

$$b_n(n+r)(n+r-1) - 4(n+r)b_n + b_{n-2} = 0 \quad (4)$$

Which for the root $r = 0$ becomes

$$b_n n(n-1) - 4n b_n + b_{n-2} = 0 \quad (4A)$$

Solving for b_n from the recursive equation (4) gives

$$b_n = -\frac{b_{n-2}}{n^2 + 2nr + r^2 - 5n - 5r} \quad (5)$$

Which for the root $r = 0$ becomes

$$b_n = -\frac{b_{n-2}}{n^2 - 5n} \quad (6)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0

For $n = 2$, using the above recursive equation gives

$$b_2 = -\frac{1}{r^2 - r - 6}$$

Which for the root $r = 0$ becomes

$$b_2 = \frac{1}{6}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{1}{r^2 - r - 6}$	$\frac{1}{6}$

For $n = 3$, using the above recursive equation gives

$$b_3 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{1}{r^2 - r - 6}$	$\frac{1}{6}$
b_3	0	0

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{1}{(r^2 - r - 6)(r^2 + 3r - 4)}$$

Which for the root $r = 0$ becomes

$$b_4 = \frac{1}{24}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{1}{r^2-r-6}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{r^4+2r^3-13r^2-14r+24}$	$\frac{1}{24}$

For $n = 5$, using the above recursive equation gives

$$b_5 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{1}{r^2-r-6}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{r^4+2r^3-13r^2-14r+24}$	$\frac{1}{24}$
b_5	0	0

For $n = 6$, using the above recursive equation gives

$$b_6 = -\frac{1}{(r^2 - r - 6)(r^2 + 3r - 4)(r^2 + 7r + 6)}$$

Which for the root $r = 0$ becomes

$$b_6 = -\frac{1}{144}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{1}{r^2-r-6}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{r^4+2r^3-13r^2-14r+24}$	$\frac{1}{24}$
b_5	0	0
b_6	$-\frac{1}{r^6+9r^5+7r^4-93r^3-152r^2+84r+144}$	$-\frac{1}{144}$

For $n = 7$, using the above recursive equation gives

$$b_7 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{1}{r^2-r-6}$	$\frac{1}{6}$
b_3	0	0
b_4	$\frac{1}{r^4+2r^3-13r^2-14r+24}$	$\frac{1}{24}$
b_5	0	0
b_6	$-\frac{1}{r^6+9r^5+7r^4-93r^3-152r^2+84r+144}$	$-\frac{1}{144}$
b_7	0	0

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned} y_2(x) &= b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 + b_7x^7 + b_8x^8 \dots \\ &= 1 + \frac{x^2}{6} + \frac{x^4}{24} - \frac{x^6}{144} + O(x^8) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= c_1x^5 \left(1 - \frac{x^2}{14} + \frac{x^4}{504} - \frac{x^6}{33264} + O(x^8) \right) + c_2 \left(1 + \frac{x^2}{6} + \frac{x^4}{24} - \frac{x^6}{144} + O(x^8) \right) \end{aligned}$$

Hence the final solution is

$$y = y_h \\ = c_1 x^5 \left(1 - \frac{x^2}{14} + \frac{x^4}{504} - \frac{x^6}{33264} + O(x^8) \right) + c_2 \left(1 + \frac{x^2}{6} + \frac{x^4}{24} - \frac{x^6}{144} + O(x^8) \right)$$

Summary

The solution(s) found are the following

$$y = c_1 x^5 \left(1 - \frac{x^2}{14} + \frac{x^4}{504} - \frac{x^6}{33264} + O(x^8) \right) + c_2 \left(1 + \frac{x^2}{6} + \frac{x^4}{24} - \frac{x^6}{144} + O(x^8) \right) \quad (1)$$

Verification of solutions

$$y = c_1 x^5 \left(1 - \frac{x^2}{14} + \frac{x^4}{504} - \frac{x^6}{33264} + O(x^8) \right) + c_2 \left(1 + \frac{x^2}{6} + \frac{x^4}{24} - \frac{x^6}{144} + O(x^8) \right)$$

Verified OK.

22.11.1 Maple step by step solution

Let's solve

$$y''x - 4y' + xy = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = \frac{4y'}{x} - y$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - \frac{4y'}{x} + y = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$[P_2(x) = -\frac{4}{x}, P_3(x) = 1]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = -4$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 0$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$y''x - 4y' + xy = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x \cdot y$ to series expansion

$$x \cdot y = \sum_{k=0}^{\infty} a_k x^{k+r+1}$$

- Shift index using $k \rightarrow k - 1$

$$x \cdot y = \sum_{k=1}^{\infty} a_{k-1} x^{k+r}$$

- Convert y' to series expansion

$$y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1}$$

- Shift index using $k \rightarrow k + 1$

$$y' = \sum_{k=-1}^{\infty} a_{k+1} (k+r+1) x^{k+r}$$

- Convert $x \cdot y''$ to series expansion

$$x \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r-1}$$

- Shift index using $k \rightarrow k + 1$

$$x \cdot y'' = \sum_{k=-1}^{\infty} a_{k+1} (k+r+1)(k+r) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0 r(-5+r) x^{-1+r} + a_1 (1+r)(-4+r) x^r + \left(\sum_{k=1}^{\infty} (a_{k+1} (k+r+1)(k-4+r) + a_{k-1}) x^{k+r} \right) =$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$r(-5 + r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \{0, 5\}$$

- Each term must be 0

$$a_1(1 + r)(-4 + r) = 0$$

- Each term in the series must be 0, giving the recursion relation

$$a_{k+1}(k + r + 1)(k - 4 + r) + a_{k-1} = 0$$

- Shift index using $k \rightarrow k + 1$

$$a_{k+2}(k + 2 + r)(k - 3 + r) + a_k = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+2} = -\frac{a_k}{(k+2+r)(k-3+r)}$$

- Recursion relation for $r = 0$

$$a_{k+2} = -\frac{a_k}{(k+2)(k-3)}$$

- Solution for $r = 0$

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+2} = -\frac{a_k}{(k+2)(k-3)}, -4a_1 = 0 \right]$$

- Recursion relation for $r = 5$

$$a_{k+2} = -\frac{a_k}{(k+7)(k+2)}$$

- Solution for $r = 5$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+5}, a_{k+2} = -\frac{a_k}{(k+7)(k+2)}, 6a_1 = 0 \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^k \right) + \left(\sum_{k=0}^{\infty} b_k x^{k+5} \right), a_{k+2} = -\frac{a_k}{(k+2)(k-3)}, -4a_1 = 0, b_{k+2} = -\frac{b_k}{(k+7)(k+2)}, 6b_1 = 0 \right]$$

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacics algorithm  
    A Liouvillian solution exists  
    Group is reducible or imprimitive  
<- Kovacics algorithm successful`
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 36

```
Order:=8;  
dsolve(x*diff(y(x),x$2)-4*diff(y(x),x)+x*y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 x^5 \left(1 - \frac{1}{14} x^2 + \frac{1}{504} x^4 - \frac{1}{33264} x^6 + O(x^8) \right) \\ + c_2 (2880 + 480x^2 + 120x^4 - 20x^6 + O(x^8))$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 58

```
AsymptoticDSolveValue[x*y'[x]-4*y[x]+x*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(-\frac{x^6}{144} + \frac{x^4}{24} + \frac{x^2}{6} + 1 \right) + c_2 \left(-\frac{x^{11}}{33264} + \frac{x^9}{504} - \frac{x^7}{14} + x^5 \right)$$

22.12 problem 2(d)

22.12.1 Maple step by step solution 3733

Internal problem ID [6487]

Internal file name [OUTPUT/5735_Sunday_June_05_2022_03_51_37_PM_14322178/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Problems for review and discover. (A) Drill Exercises . Page 194

Problem number: 2(d).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Complex roots"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$4x^2y'' + 4x^2y' + 2y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$4x^2y'' + 4x^2y' + 2y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = 1$$
$$q(x) = \frac{1}{2x^2}$$

Table 506: Table $p(x), q(x)$ singularities.

$p(x) = 1$	
singularity	type

$q(x) = \frac{1}{2x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$4x^2y'' + 4x^2y' + 2y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$4x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) + 4x^2 \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + 2 \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} 4x^{n+r} a_n (n+r)(n+r-1) \right) + \left(\sum_{n=0}^{\infty} 4x^{1+n+r} a_n (n+r) \right) + \left(\sum_{n=0}^{\infty} 2a_n x^{n+r} \right) = 0 \quad (2A)$$

The next step is to make all powers of x be $n + r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} 4x^{1+n+r} a_n(n+r) = \sum_{n=1}^{\infty} 4a_{n-1}(n+r-1) x^{n+r}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n + r$.

$$\left(\sum_{n=0}^{\infty} 4x^{n+r} a_n(n+r)(n+r-1) \right) + \left(\sum_{n=1}^{\infty} 4a_{n-1}(n+r-1) x^{n+r} \right) + \left(\sum_{n=0}^{\infty} 2a_n x^{n+r} \right) = 0 \quad (2B)$$

The indicial equation is obtained from $n = 0$. From Eq (2B) this gives

$$4x^{n+r} a_n(n+r)(n+r-1) + 2a_n x^{n+r} = 0$$

When $n = 0$ the above becomes

$$4x^r a_0 r(-1+r) + 2a_0 x^r = 0$$

Or

$$(4x^r r(-1+r) + 2x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(4r^2 - 4r + 2) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$4r^2 - 4r + 2 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = \frac{1}{2} + \frac{i}{2}$$

$$r_2 = \frac{1}{2} - \frac{i}{2}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(4r^2 - 4r + 2) x^r = 0$$

Solving for r gives the roots of the indicial equation as Since the roots are complex conjugates, then two linearly independent solutions can be constructed using

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+\frac{1}{2}+\frac{i}{2}}$$

$$y_2(x) = \sum_{n=0}^{\infty} b_n x^{n+\frac{1}{2}-\frac{i}{2}}$$

$y_1(x)$ is found first. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$4a_n(n+r)(n+r-1) + 4a_{n-1}(n+r-1) + 2a_n = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{2a_{n-1}(n+r-1)}{2n^2 + 4nr + 2r^2 - 2n - 2r + 1} \quad (4)$$

Which for the root $r = \frac{1}{2} + \frac{i}{2}$ becomes

$$a_n = -\frac{a_{n-1}(2n-1+i)}{2n(n+i)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = \frac{1}{2} + \frac{i}{2}$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = -\frac{2r}{2r^2 + 2r + 1}$$

Which for the root $r = \frac{1}{2} + \frac{i}{2}$ becomes

$$a_1 = -\frac{1}{2}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{2r}{2r^2+2r+1}$	$-\frac{1}{2}$

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{4r(1+r)}{4r^4 + 16r^3 + 24r^2 + 16r + 5}$$

Which for the root $r = \frac{1}{2} + \frac{i}{2}$ becomes

$$a_2 = \frac{7}{40} - \frac{i}{40}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{2r}{2r^2+2r+1}$	$-\frac{1}{2}$
a_2	$\frac{4r(1+r)}{4r^4+16r^3+24r^2+16r+5}$	$\frac{7}{40} - \frac{i}{40}$

For $n = 3$, using the above recursive equation gives

$$a_3 = -\frac{8r(1+r)(2+r)}{8r^6 + 72r^5 + 260r^4 + 480r^3 + 482r^2 + 258r + 65}$$

Which for the root $r = \frac{1}{2} + \frac{i}{2}$ becomes

$$a_3 = -\frac{11}{240} + \frac{i}{80}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{2r}{2r^2+2r+1}$	$-\frac{1}{2}$
a_2	$\frac{4r(1+r)}{4r^4+16r^3+24r^2+16r+5}$	$\frac{7}{40} - \frac{i}{40}$
a_3	$-\frac{8r(1+r)(2+r)}{8r^6+72r^5+260r^4+480r^3+482r^2+258r+65}$	$-\frac{11}{240} + \frac{i}{80}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{16r(1+r)(2+r)(3+r)}{16r^8 + 256r^7 + 1728r^6 + 6400r^5 + 14184r^4 + 19264r^3 + 15792r^2 + 7360r + 1625}$$

Which for the root $r = \frac{1}{2} + \frac{i}{2}$ becomes

$$a_4 = \frac{31}{3264} - \frac{i}{272}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{2r}{2r^2+2r+1}$	$-\frac{1}{2}$
a_2	$\frac{4r(1+r)}{4r^4+16r^3+24r^2+16r+5}$	$\frac{7}{40} - \frac{i}{40}$
a_3	$-\frac{8r(1+r)(2+r)}{8r^6+72r^5+260r^4+480r^3+482r^2+258r+65}$	$-\frac{11}{240} + \frac{i}{80}$
a_4	$\frac{16r(1+r)(2+r)(3+r)}{16r^8+256r^7+1728r^6+6400r^5+14184r^4+19264r^3+15792r^2+7360r+1625}$	$\frac{31}{3264} - \frac{i}{272}$

For $n = 5$, using the above recursive equation gives

$$a_5 = -\frac{r(1+r)(2+r)(3+r)(4+r)}{\left(r^2 + 5r + \frac{13}{2}\right)\left(r^2 + 3r + \frac{5}{2}\right)\left(r^2 + 7r + \frac{25}{2}\right)\left(r^2 + 9r + \frac{41}{2}\right)\left(r^2 + r + \frac{1}{2}\right)}$$

Which for the root $r = \frac{1}{2} + \frac{i}{2}$ becomes

$$a_5 = -\frac{53}{32640} + \frac{13i}{16320}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{2r}{2r^2+2r+1}$	$-\frac{1}{2}$
a_2	$\frac{4r(1+r)}{4r^4+16r^3+24r^2+16r+5}$	$\frac{7}{40} - \frac{i}{40}$
a_3	$-\frac{8r(1+r)(2+r)}{8r^6+72r^5+260r^4+480r^3+482r^2+258r+65}$	$-\frac{11}{240} + \frac{i}{80}$
a_4	$\frac{16r(1+r)(2+r)(3+r)}{16r^8+256r^7+1728r^6+6400r^5+14184r^4+19264r^3+15792r^2+7360r+1625}$	$\frac{31}{3264} - \frac{i}{272}$
a_5	$-\frac{r(1+r)(2+r)(3+r)(4+r)}{(r^2+5r+\frac{13}{2})(r^2+3r+\frac{5}{2})(r^2+7r+\frac{25}{2})(r^2+9r+\frac{41}{2})(r^2+r+\frac{1}{2})}$	$-\frac{53}{32640} + \frac{13i}{16320}$

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{64r(1+r)(2+r)(3+r)(4+r)(5+r)}{(2r^2+10r+13)(2r^2+6r+5)(2r^2+14r+25)(2r^2+18r+41)(2r^2+2r+1)(2r^2+22r+61)}$$

Which for the root $r = \frac{1}{2} + \frac{i}{2}$ becomes

$$a_6 = \frac{3421}{14492160} - \frac{223i}{1610240}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{2r}{2r^2+2r+1}$	$-\frac{1}{2}$
a_2	$\frac{4r(1+r)}{4r^4+16r^3+24r^2+16r+5}$	$\frac{7}{40} - \frac{i}{40}$
a_3	$-\frac{8r(1+r)(2+r)}{8r^6+72r^5+260r^4+480r^3+482r^2+258r+65}$	$-\frac{11}{240} + \frac{i}{80}$
a_4	$\frac{16r(1+r)(2+r)(3+r)}{16r^8+256r^7+1728r^6+6400r^5+14184r^4+19264r^3+15792r^2+7360r+1625}$	$\frac{31}{3264} - \frac{i}{272}$
a_5	$-\frac{r(1+r)(2+r)(3+r)(4+r)}{(r^2+5r+\frac{13}{2})(r^2+3r+\frac{5}{2})(r^2+7r+\frac{25}{2})(r^2+9r+\frac{41}{2})(r^2+r+\frac{1}{2})}$	$-\frac{53}{32640} + \frac{13i}{16320}$
a_6	$\frac{64r(1+r)(2+r)(3+r)(4+r)(5+r)}{(2r^2+10r+13)(2r^2+6r+5)(2r^2+14r+25)(2r^2+18r+41)(2r^2+2r+1)(2r^2+22r+61)}$	$\frac{3421}{14492160} - \frac{223i}{1610240}$

For $n = 7$, using the above recursive equation gives

$$a_7 = -\frac{128r(1+r)(2+r)(3+r)(4+r)(5+r)(6+r)}{(2r^2+10r+13)(2r^2+6r+5)(2r^2+14r+25)(2r^2+18r+41)(2r^2+2r+1)(2r^2+22r+61)}$$

Which for the root $r = \frac{1}{2} + \frac{i}{2}$ becomes

$$a_7 = -\frac{30269}{1014451200} + \frac{977i}{48307200}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{2r}{2r^2+2r+1}$	$-\frac{1}{2}$
a_2	$\frac{4r(1+r)}{4r^4+16r^3+24r^2+16r+5}$	$\frac{7}{40} - \frac{i}{40}$
a_3	$-\frac{8r(1+r)(2+r)}{8r^6+72r^5+260r^4+480r^3+482r^2+258r+65}$	$-\frac{11}{240} + \frac{i}{80}$
a_4	$\frac{16r(1+r)(2+r)(3+r)}{16r^8+256r^7+1728r^6+6400r^5+14184r^4+19264r^3+15792r^2+7360r+1625}$	$\frac{31}{3264} - \frac{i}{272}$
a_5	$-\frac{r(1+r)(2+r)(3+r)(4+r)}{(r^2+5r+\frac{13}{2})(r^2+3r+\frac{5}{2})(r^2+7r+\frac{25}{2})(r^2+9r+\frac{41}{2})(r^2+r+\frac{1}{2})}$	$-\frac{53}{32640} + \frac{13i}{16320}$
a_6	$\frac{64r(1+r)(2+r)(3+r)(4+r)(5+r)}{(2r^2+10r+13)(2r^2+6r+5)(2r^2+14r+25)(2r^2+18r+41)(2r^2+2r+1)(2r^2+22r+61)}$	$\frac{3421}{14492160} - \frac{223i}{1610240}$
a_7	$-\frac{128r(1+r)(2+r)(3+r)(4+r)(5+r)(6+r)}{(2r^2+10r+13)(2r^2+6r+5)(2r^2+14r+25)(2r^2+18r+41)(2r^2+2r+1)(2r^2+22r+61)(2r^2+26r+85)}$	$-\frac{30269}{1014451200} + \frac{977i}{48307200}$

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned}
 y_1(x) &= x^{\frac{1}{2}+\frac{i}{2}} \left(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots \right) \\
 &= x^{\frac{1}{2}+\frac{i}{2}} \left(1 - \frac{x}{2} + \left(\frac{7}{40} - \frac{i}{40} \right) x^2 + \left(-\frac{11}{240} + \frac{i}{80} \right) x^3 + \left(\frac{31}{3264} - \frac{i}{272} \right) x^4 \right. \\
 &\quad + \left(-\frac{53}{32640} + \frac{13i}{16320} \right) x^5 + \left(\frac{3421}{14492160} - \frac{223i}{1610240} \right) x^6 \\
 &\quad \left. + \left(-\frac{30269}{1014451200} + \frac{977i}{48307200} \right) x^7 + O(x^8) \right)
 \end{aligned}$$

The second solution $y_2(x)$ is found by taking the complex conjugate of $y_1(x)$ which gives

$$\begin{aligned}
 y_2(x) &= x^{\frac{1}{2}-\frac{i}{2}} \left(1 - \frac{x}{2} + \left(\frac{7}{40} + \frac{i}{40} \right) x^2 + \left(-\frac{11}{240} - \frac{i}{80} \right) x^3 + \left(\frac{31}{3264} + \frac{i}{272} \right) x^4 \right. \\
 &\quad + \left(-\frac{53}{32640} - \frac{13i}{16320} \right) x^5 + \left(\frac{3421}{14492160} + \frac{223i}{1610240} \right) x^6 \\
 &\quad \left. + \left(-\frac{30269}{1014451200} - \frac{977i}{48307200} \right) x^7 + O(x^8) \right)
 \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 y_h(x) &= c_1 y_1(x) + c_2 y_2(x) \\
 &= c_1 x^{\frac{1}{2} + \frac{i}{2}} \left(1 - \frac{x}{2} + \left(\frac{7}{40} - \frac{i}{40} \right) x^2 + \left(-\frac{11}{240} + \frac{i}{80} \right) x^3 + \left(\frac{31}{3264} - \frac{i}{272} \right) x^4 \right. \\
 &\quad \left. + \left(-\frac{53}{32640} + \frac{13i}{16320} \right) x^5 + \left(\frac{3421}{14492160} - \frac{223i}{1610240} \right) x^6 \right. \\
 &\quad \left. + \left(-\frac{30269}{1014451200} + \frac{977i}{48307200} \right) x^7 + O(x^8) \right) + c_2 x^{\frac{1}{2} - \frac{i}{2}} \left(1 - \frac{x}{2} + \left(\frac{7}{40} + \frac{i}{40} \right) x^2 \right. \\
 &\quad \left. + \left(-\frac{11}{240} - \frac{i}{80} \right) x^3 + \left(\frac{31}{3264} + \frac{i}{272} \right) x^4 + \left(-\frac{53}{32640} - \frac{13i}{16320} \right) x^5 \right. \\
 &\quad \left. + \left(\frac{3421}{14492160} + \frac{223i}{1610240} \right) x^6 + \left(-\frac{30269}{1014451200} - \frac{977i}{48307200} \right) x^7 + O(x^8) \right)
 \end{aligned}$$

Hence the final solution is

$$\begin{aligned}
 y &= y_h \\
 &= c_1 x^{\frac{1}{2} + \frac{i}{2}} \left(1 - \frac{x}{2} + \left(\frac{7}{40} - \frac{i}{40} \right) x^2 + \left(-\frac{11}{240} + \frac{i}{80} \right) x^3 + \left(\frac{31}{3264} - \frac{i}{272} \right) x^4 \right. \\
 &\quad \left. + \left(-\frac{53}{32640} + \frac{13i}{16320} \right) x^5 + \left(\frac{3421}{14492160} - \frac{223i}{1610240} \right) x^6 \right. \\
 &\quad \left. + \left(-\frac{30269}{1014451200} + \frac{977i}{48307200} \right) x^7 + O(x^8) \right) + c_2 x^{\frac{1}{2} - \frac{i}{2}} \left(1 - \frac{x}{2} + \left(\frac{7}{40} + \frac{i}{40} \right) x^2 \right. \\
 &\quad \left. + \left(-\frac{11}{240} - \frac{i}{80} \right) x^3 + \left(\frac{31}{3264} + \frac{i}{272} \right) x^4 + \left(-\frac{53}{32640} - \frac{13i}{16320} \right) x^5 \right. \\
 &\quad \left. + \left(\frac{3421}{14492160} + \frac{223i}{1610240} \right) x^6 + \left(-\frac{30269}{1014451200} - \frac{977i}{48307200} \right) x^7 + O(x^8) \right)
 \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned} y = & c_1 x^{\frac{1}{2} + \frac{i}{2}} \left(1 - \frac{x}{2} + \left(\frac{7}{40} - \frac{i}{40} \right) x^2 + \left(-\frac{11}{240} + \frac{i}{80} \right) x^3 + \left(\frac{31}{3264} - \frac{i}{272} \right) x^4 \right. \\ & \left. + \left(-\frac{53}{32640} + \frac{13i}{16320} \right) x^5 + \left(\frac{3421}{14492160} - \frac{223i}{1610240} \right) x^6 \right. \\ & \left. + \left(-\frac{30269}{1014451200} + \frac{977i}{48307200} \right) x^7 + O(x^8) \right) + c_2 x^{\frac{1}{2} - \frac{i}{2}} \left(1 - \frac{x}{2} + \left(\frac{7}{40} + \frac{i}{40} \right) x^2 \right. \\ & \left. + \left(-\frac{11}{240} - \frac{i}{80} \right) x^3 + \left(\frac{31}{3264} + \frac{i}{272} \right) x^4 + \left(-\frac{53}{32640} - \frac{13i}{16320} \right) x^5 \right. \\ & \left. + \left(\frac{3421}{14492160} + \frac{223i}{1610240} \right) x^6 + \left(-\frac{30269}{1014451200} - \frac{977i}{48307200} \right) x^7 + O(x^8) \right) \end{aligned}$$

Verification of solutions

$$\begin{aligned} y = & c_1 x^{\frac{1}{2} + \frac{i}{2}} \left(1 - \frac{x}{2} + \left(\frac{7}{40} - \frac{i}{40} \right) x^2 + \left(-\frac{11}{240} + \frac{i}{80} \right) x^3 + \left(\frac{31}{3264} - \frac{i}{272} \right) x^4 \right. \\ & \left. + \left(-\frac{53}{32640} + \frac{13i}{16320} \right) x^5 + \left(\frac{3421}{14492160} - \frac{223i}{1610240} \right) x^6 \right. \\ & \left. + \left(-\frac{30269}{1014451200} + \frac{977i}{48307200} \right) x^7 + O(x^8) \right) + c_2 x^{\frac{1}{2} - \frac{i}{2}} \left(1 - \frac{x}{2} + \left(\frac{7}{40} + \frac{i}{40} \right) x^2 \right. \\ & \left. + \left(-\frac{11}{240} - \frac{i}{80} \right) x^3 + \left(\frac{31}{3264} + \frac{i}{272} \right) x^4 + \left(-\frac{53}{32640} - \frac{13i}{16320} \right) x^5 \right. \\ & \left. + \left(\frac{3421}{14492160} + \frac{223i}{1610240} \right) x^6 + \left(-\frac{30269}{1014451200} - \frac{977i}{48307200} \right) x^7 + O(x^8) \right) \end{aligned}$$

Verified OK.

22.12.1 Maple step by step solution

Let's solve

$$4x^2 y'' + 4x^2 y' + 2y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -y' - \frac{y}{2x^2}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + y' + \frac{y}{2x^2} = 0$$

□ Check to see if $x_0 = 0$ is a regular singular point

○ Define functions

$$[P_2(x) = 1, P_3(x) = \frac{1}{2x^2}]$$

○ $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 0$$

○ $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = \frac{1}{2}$$

○ $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

• Multiply by denominators

$$2x^2y' + 2x^2y'' + y = 0$$

• Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

□ Rewrite ODE with series expansions

○ Convert $x^2 \cdot y'$ to series expansion

$$x^2 \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r+1}$$

○ Shift index using $k- > k-1$

$$x^2 \cdot y' = \sum_{k=1}^{\infty} a_{k-1} (k-1+r) x^{k+r}$$

○ Convert $x^2 \cdot y''$ to series expansion

$$x^2 \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k-1+r) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(2r^2 - 2r + 1) x^r + \left(\sum_{k=1}^{\infty} (a_k(2k^2 + 4kr + 2r^2 - 2k - 2r + 1) + 2a_{k-1}(k-1+r)) x^{k+r} \right) = 0$$

• a_0 cannot be 0 by assumption, giving the indicial equation

$$2r^2 - 2r + 1 = 0$$

- Values of r that satisfy the indicial equation

$$r \in \left\{ \frac{1}{2} - \frac{1}{2}, \frac{1}{2} + \frac{1}{2} \right\}$$

- Each term in the series must be 0, giving the recursion relation

$$(2k^2 + (4r - 2)k + 2r^2 - 2r + 1)a_k + 2a_{k-1}(k - 1 + r) = 0$$

- Shift index using $k \rightarrow k + 1$

$$(2(k + 1)^2 + (4r - 2)(k + 1) + 2r^2 - 2r + 1)a_{k+1} + 2a_k(k + r) = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+1} = -\frac{2a_k(k+r)}{2k^2+4kr+2r^2+2k+2r+1}$$

- Recursion relation for $r = \frac{1}{2} - \frac{1}{2}$

$$a_{k+1} = -\frac{2a_k\left(k+\frac{1}{2}-\frac{1}{2}\right)}{2k^2+(2-2\text{I})k+2-2\text{I}+2k}$$

- Solution for $r = \frac{1}{2} - \frac{1}{2}$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+\frac{1}{2}-\frac{1}{2}}, a_{k+1} = -\frac{2a_k\left(k+\frac{1}{2}-\frac{1}{2}\right)}{2k^2+(2-2\text{I})k+2-2\text{I}+2k} \right]$$

- Recursion relation for $r = \frac{1}{2} + \frac{1}{2}$

$$a_{k+1} = -\frac{2a_k\left(k+\frac{1}{2}+\frac{1}{2}\right)}{2k^2+(2+2\text{I})k+2+2\text{I}+2k}$$

- Solution for $r = \frac{1}{2} + \frac{1}{2}$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+\frac{1}{2}+\frac{1}{2}}, a_{k+1} = -\frac{2a_k\left(k+\frac{1}{2}+\frac{1}{2}\right)}{2k^2+(2+2\text{I})k+2+2\text{I}+2k} \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^{k+\frac{1}{2}-\frac{1}{2}} \right) + \left(\sum_{k=0}^{\infty} b_k x^{k+\frac{1}{2}+\frac{1}{2}} \right), a_{k+1} = -\frac{2a_k\left(k+\frac{1}{2}-\frac{1}{2}\right)}{2k^2+(2-2\text{I})k+2-2\text{I}+2k}, b_{k+1} = -\frac{2b_k\left(k+\frac{1}{2}+\frac{1}{2}\right)}{2k^2+(2+2\text{I})k+2+2\text{I}+2k} \right]$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacic's algorithm  
<- No Liouvillian solutions exist  
-> Trying a solution in terms of special functions:  
  -> Bessel  
  <- Bessel successful  
<- special function solution successful`
```

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 83

```
Order:=8;  
dsolve(4*x^2*dif(y(x),x$2)+4*x^2*dif(y(x),x)+2*y(x)=0,y(x),type='series',x=0);
```

$$\begin{aligned} y(x) = & c_1 x^{\frac{1}{2}-\frac{i}{2}} \left(1 - \frac{1}{2}x + \left(\frac{7}{40} + \frac{i}{40} \right) x^2 + \left(-\frac{11}{240} - \frac{i}{80} \right) x^3 + \left(\frac{31}{3264} + \frac{i}{272} \right) x^4 \right. \\ & \left. + \left(-\frac{53}{32640} - \frac{13i}{16320} \right) x^5 + \left(\frac{3421}{14492160} + \frac{223i}{1610240} \right) x^6 \right. \\ & \left. + \left(-\frac{30269}{1014451200} - \frac{977i}{48307200} \right) x^7 + O(x^8) \right) + c_2 x^{\frac{1}{2}+\frac{i}{2}} \left(1 - \frac{1}{2}x + \left(\frac{7}{40} - \frac{i}{40} \right) x^2 \right. \\ & \left. + \left(-\frac{11}{240} + \frac{i}{80} \right) x^3 + \left(\frac{31}{3264} - \frac{i}{272} \right) x^4 + \left(-\frac{53}{32640} + \frac{13i}{16320} \right) x^5 \right. \\ & \left. + \left(\frac{3421}{14492160} - \frac{223i}{1610240} \right) x^6 + \left(-\frac{30269}{1014451200} + \frac{977i}{48307200} \right) x^7 + O(x^8) \right) \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.023 (sec). Leaf size: 226

```
AsymptoticDSolveValue[4*x^2*y''[x]+4*x^2*y'[x]+2*y[x]==0,y[x],{x,0,7}]
```

$$\begin{aligned} y(x) \rightarrow & c_1 \left(\left(\frac{3421}{14492160} - \frac{223i}{1610240} \right) x^{\frac{13}{2} + \frac{i}{2}} - \left(\frac{53}{32640} - \frac{13i}{16320} \right) x^{\frac{11}{2} + \frac{i}{2}} \right. \\ & \left. + \left(\frac{31}{3264} - \frac{i}{272} \right) x^{\frac{9}{2} + \frac{i}{2}} - \left(\frac{11}{240} - \frac{i}{80} \right) x^{\frac{7}{2} + \frac{i}{2}} + \left(\frac{7}{40} - \frac{i}{40} \right) x^{\frac{5}{2} + \frac{i}{2}} - \frac{1}{2} x^{\frac{3}{2} + \frac{i}{2}} + x^{\frac{1}{2} + \frac{i}{2}} \right) \\ & + c_2 \left(\left(\frac{3421}{14492160} + \frac{223i}{1610240} \right) x^{\frac{13}{2} - \frac{i}{2}} - \left(\frac{53}{32640} + \frac{13i}{16320} \right) x^{\frac{11}{2} - \frac{i}{2}} \right. \\ & \left. + \left(\frac{31}{3264} + \frac{i}{272} \right) x^{\frac{9}{2} - \frac{i}{2}} - \left(\frac{11}{240} + \frac{i}{80} \right) x^{\frac{7}{2} - \frac{i}{2}} + \left(\frac{7}{40} + \frac{i}{40} \right) x^{\frac{5}{2} - \frac{i}{2}} - \frac{1}{2} x^{\frac{3}{2} - \frac{i}{2}} + x^{\frac{1}{2} - \frac{i}{2}} \right) \end{aligned}$$

22.13 problem 2(e)

22.13.1 Maple step by step solution 3750

Internal problem ID [6488]

Internal file name [OUTPUT/5736_Sunday_June_05_2022_03_51_46_PM_78360815/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Problems for review and discover. (A) Drill Exercises . Page 194

Problem number: 2(e).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$2xy'' + (1 - x)y' + y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$2xy'' + (1 - x)y' + y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{x-1}{2x}$$
$$q(x) = \frac{1}{2x}$$

Table 508: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{x-1}{2x}$	
singularity	type
$x = 0$	“regular”

$q(x) = \frac{1}{2x}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$2xy'' + (1 - x)y' + y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) x + (1-x) \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2x^{n+r-1} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) \\ & + \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r-1$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r-1} and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) &= \sum_{n=1}^{\infty} (-a_{n-1} (n+r-1) x^{n+r-1}) \\ \sum_{n=0}^{\infty} a_n x^{n+r} &= \sum_{n=1}^{\infty} a_{n-1} x^{n+r-1} \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r-1$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} 2x^{n+r-1} a_n (n+r) (n+r-1) \right) + \sum_{n=1}^{\infty} (-a_{n-1} (n+r-1) x^{n+r-1}) \\ & + \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=1}^{\infty} a_{n-1} x^{n+r-1} \right) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$2x^{n+r-1} a_n (n+r) (n+r-1) + (n+r) a_n x^{n+r-1} = 0$$

When $n=0$ the above becomes

$$2x^{-1+r} a_0 r (-1+r) + r a_0 x^{-1+r} = 0$$

Or

$$(2x^{-1+r} r (-1+r) + r x^{-1+r}) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$r x^{-1+r} (-1+2r) = 0$$

Since the above is true for all x then the indicial equation becomes

$$2r^2 - r = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = \frac{1}{2}$$

$$r_2 = 0$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$r x^{-1+r}(-1 + 2r) = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = \frac{1}{2}$ is not an integer, then we can construct two linearly independent solutions

$$y_1(x) = x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right)$$

$$y_2(x) = x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right)$$

Or

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+\frac{1}{2}}$$

$$y_2(x) = \sum_{n=0}^{\infty} b_n x^n$$

We start by finding $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$2a_n(n+r)(n+r-1) - a_{n-1}(n+r-1) + a_n(n+r) + a_{n-1} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-1}(n+r-2)}{2n^2 + 4nr + 2r^2 - n - r} \quad (4)$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_n = \frac{2na_{n-1} - 3a_{n-1}}{4n^2 + 2n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = \frac{1}{2}$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = \frac{-1 + r}{2r^2 + 3r + 1}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_1 = -\frac{1}{6}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-1+r}{2r^2+3r+1}$	$-\frac{1}{6}$

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{r(-1 + r)}{4r^4 + 20r^3 + 35r^2 + 25r + 6}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_2 = -\frac{1}{120}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-1+r}{2r^2+3r+1}$	$-\frac{1}{6}$
a_2	$\frac{r(-1+r)}{4r^4+20r^3+35r^2+25r+6}$	$-\frac{1}{120}$

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{r(-1 + r)}{8r^5 + 76r^4 + 274r^3 + 461r^2 + 351r + 90}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_3 = -\frac{1}{1680}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-1+r}{2r^2+3r+1}$	$-\frac{1}{6}$
a_2	$\frac{r(-1+r)}{4r^4+20r^3+35r^2+25r+6}$	$-\frac{1}{120}$
a_3	$\frac{r(-1+r)}{8r^5+76r^4+274r^3+461r^2+351r+90}$	$-\frac{1}{1680}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{r(-1+r)}{16r^6 + 240r^5 + 1432r^4 + 4296r^3 + 6697r^2 + 4959r + 1260}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_4 = -\frac{1}{24192}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-1+r}{2r^2+3r+1}$	$-\frac{1}{6}$
a_2	$\frac{r(-1+r)}{4r^4+20r^3+35r^2+25r+6}$	$-\frac{1}{120}$
a_3	$\frac{r(-1+r)}{8r^5+76r^4+274r^3+461r^2+351r+90}$	$-\frac{1}{1680}$
a_4	$\frac{r(-1+r)}{16r^6+240r^5+1432r^4+4296r^3+6697r^2+4959r+1260}$	$-\frac{1}{24192}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{r(-1+r)}{32r^7 + 688r^6 + 6080r^5 + 28360r^4 + 74378r^3 + 107347r^2 + 76065r + 18900}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_5 = -\frac{1}{380160}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-1+r}{2r^2+3r+1}$	$-\frac{1}{6}$
a_2	$\frac{r(-1+r)}{4r^4+20r^3+35r^2+25r+6}$	$-\frac{1}{120}$
a_3	$\frac{r(-1+r)}{8r^5+76r^4+274r^3+461r^2+351r+90}$	$-\frac{1}{1680}$
a_4	$\frac{r(-1+r)}{16r^6+240r^5+1432r^4+4296r^3+6697r^2+4959r+1260}$	$-\frac{1}{24192}$
a_5	$\frac{r(-1+r)}{32r^7+688r^6+6080r^5+28360r^4+74378r^3+107347r^2+76065r+18900}$	$-\frac{1}{380160}$

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{r(-1+r)}{64r^8 + 1856r^7 + 22672r^6 + 151280r^5 + 597196r^4 + 1408364r^3 + 1896603r^2 + 1285785r + 311850}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_6 = -\frac{1}{6589440}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-1+r}{2r^2+3r+1}$	$-\frac{1}{6}$
a_2	$\frac{r(-1+r)}{4r^4+20r^3+35r^2+25r+6}$	$-\frac{1}{120}$
a_3	$\frac{r(-1+r)}{8r^5+76r^4+274r^3+461r^2+351r+90}$	$-\frac{1}{1680}$
a_4	$\frac{r(-1+r)}{16r^6+240r^5+1432r^4+4296r^3+6697r^2+4959r+1260}$	$-\frac{1}{24192}$
a_5	$\frac{r(-1+r)}{32r^7+688r^6+6080r^5+28360r^4+74378r^3+107347r^2+76065r+18900}$	$-\frac{1}{380160}$
a_6	$\frac{r(-1+r)}{64r^8+1856r^7+22672r^6+151280r^5+597196r^4+1408364r^3+1896603r^2+1285785r+311850}$	$-\frac{1}{6589440}$

For $n = 7$, using the above recursive equation gives

$$a_7 = \frac{r(-1+r)}{128r^9 + 4800r^8 + 77280r^7 + 697200r^6 + 3856104r^5 + 13426980r^4 + 29028970r^3 + 36796125r^2 + 23112300r + 4500000}$$

Which for the root $r = \frac{1}{2}$ becomes

$$a_7 = -\frac{1}{125798400}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-1+r}{2r^2+3r+1}$	$-\frac{1}{6}$
a_2	$\frac{r(-1+r)}{4r^4+20r^3+35r^2+25r+6}$	$-\frac{1}{120}$
a_3	$\frac{r(-1+r)}{8r^5+76r^4+274r^3+461r^2+351r+90}$	$-\frac{1}{1680}$
a_4	$\frac{r(-1+r)}{16r^6+240r^5+1432r^4+4296r^3+6697r^2+4959r+1260}$	$-\frac{1}{24192}$
a_5	$\frac{r(-1+r)}{32r^7+688r^6+6080r^5+28360r^4+74378r^3+107347r^2+76065r+18900}$	$-\frac{1}{380160}$
a_6	$\frac{r(-1+r)}{64r^8+1856r^7+22672r^6+151280r^5+597196r^4+1408364r^3+1896603r^2+1285785r+311850}$	$-\frac{1}{6589440}$
a_7	$\frac{r(-1+r)}{128r^9+4800r^8+77280r^7+697200r^6+3856104r^5+13426980r^4+29028970r^3+36796125r^2+23950143r+5675670}$	$-\frac{1}{125798400}$

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= \sqrt{x}(a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots) \\ &= \sqrt{x} \left(1 - \frac{x}{6} - \frac{x^2}{120} - \frac{x^3}{1680} - \frac{x^4}{24192} - \frac{x^5}{380160} - \frac{x^6}{6589440} - \frac{x^7}{125798400} + O(x^8) \right) \end{aligned}$$

Now the second solution $y_2(x)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. For $1 \leq n$ the recursive equation is

$$2b_n(n+r)(n+r-1) - b_{n-1}(n+r-1) + (n+r)b_n + b_{n-1} = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = \frac{b_{n-1}(n+r-2)}{2n^2 + 4nr + 2r^2 - n - r} \quad (4)$$

Which for the root $r = 0$ becomes

$$b_n = \frac{b_{n-1}(n-2)}{n(2n-1)} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1

For $n = 1$, using the above recursive equation gives

$$b_1 = \frac{-1 + r}{2r^2 + 3r + 1}$$

Which for the root $r = 0$ becomes

$$b_1 = -1$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{-1+r}{2r^2+3r+1}$	-1

For $n = 2$, using the above recursive equation gives

$$b_2 = \frac{r(-1 + r)}{4r^4 + 20r^3 + 35r^2 + 25r + 6}$$

Which for the root $r = 0$ becomes

$$b_2 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{-1+r}{2r^2+3r+1}$	-1
b_2	$\frac{r(-1+r)}{4r^4+20r^3+35r^2+25r+6}$	0

For $n = 3$, using the above recursive equation gives

$$b_3 = \frac{r(-1 + r)}{8r^5 + 76r^4 + 274r^3 + 461r^2 + 351r + 90}$$

Which for the root $r = 0$ becomes

$$b_3 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{-1+r}{2r^2+3r+1}$	-1
b_2	$\frac{r(-1+r)}{4r^4+20r^3+35r^2+25r+6}$	0
b_3	$\frac{r(-1+r)}{8r^5+76r^4+274r^3+461r^2+351r+90}$	0

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{r(-1+r)}{16r^6 + 240r^5 + 1432r^4 + 4296r^3 + 6697r^2 + 4959r + 1260}$$

Which for the root $r = 0$ becomes

$$b_4 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{-1+r}{2r^2+3r+1}$	-1
b_2	$\frac{r(-1+r)}{4r^4+20r^3+35r^2+25r+6}$	0
b_3	$\frac{r(-1+r)}{8r^5+76r^4+274r^3+461r^2+351r+90}$	0
b_4	$\frac{r(-1+r)}{16r^6+240r^5+1432r^4+4296r^3+6697r^2+4959r+1260}$	0

For $n = 5$, using the above recursive equation gives

$$b_5 = \frac{r(-1+r)}{32r^7 + 688r^6 + 6080r^5 + 28360r^4 + 74378r^3 + 107347r^2 + 76065r + 18900}$$

Which for the root $r = 0$ becomes

$$b_5 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{-1+r}{2r^2+3r+1}$	-1
b_2	$\frac{r(-1+r)}{4r^4+20r^3+35r^2+25r+6}$	0
b_3	$\frac{r(-1+r)}{8r^5+76r^4+274r^3+461r^2+351r+90}$	0
b_4	$\frac{r(-1+r)}{16r^6+240r^5+1432r^4+4296r^3+6697r^2+4959r+1260}$	0
b_5	$\frac{r(-1+r)}{32r^7+688r^6+6080r^5+28360r^4+74378r^3+107347r^2+76065r+18900}$	0

For $n = 6$, using the above recursive equation gives

$$b_6 = \frac{r(-1+r)}{64r^8 + 1856r^7 + 22672r^6 + 151280r^5 + 597196r^4 + 1408364r^3 + 1896603r^2 + 1285785r + 311850}$$

Which for the root $r = 0$ becomes

$$b_6 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{-1+r}{2r^2+3r+1}$	-1
b_2	$\frac{r(-1+r)}{4r^4+20r^3+35r^2+25r+6}$	0
b_3	$\frac{r(-1+r)}{8r^5+76r^4+274r^3+461r^2+351r+90}$	0
b_4	$\frac{r(-1+r)}{16r^6+240r^5+1432r^4+4296r^3+6697r^2+4959r+1260}$	0
b_5	$\frac{r(-1+r)}{32r^7+688r^6+6080r^5+28360r^4+74378r^3+107347r^2+76065r+18900}$	0
b_6	$\frac{r(-1+r)}{64r^8+1856r^7+22672r^6+151280r^5+597196r^4+1408364r^3+1896603r^2+1285785r+311850}$	0

For $n = 7$, using the above recursive equation gives

$$b_7 = \frac{r(-1+r)}{128r^9 + 4800r^8 + 77280r^7 + 697200r^6 + 3856104r^5 + 13426980r^4 + 29028970r^3 + 36796125r^2 + 23028970r + 311850}$$

Which for the root $r = 0$ becomes

$$b_7 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{-1+r}{2r^2+3r+1}$	-1
b_2	$\frac{r(-1+r)}{4r^4+20r^3+35r^2+25r+6}$	0
b_3	$\frac{r(-1+r)}{8r^5+76r^4+274r^3+461r^2+351r+90}$	0
b_4	$\frac{r(-1+r)}{16r^6+240r^5+1432r^4+4296r^3+6697r^2+4959r+1260}$	0
b_5	$\frac{r(-1+r)}{32r^7+688r^6+6080r^5+28360r^4+74378r^3+107347r^2+76065r+18900}$	0
b_6	$\frac{r(-1+r)}{64r^8+1856r^7+22672r^6+151280r^5+597196r^4+1408364r^3+1896603r^2+1285785r+311850}$	0
b_7	$\frac{r(-1+r)}{128r^9+4800r^8+77280r^7+697200r^6+3856104r^5+13426980r^4+29028970r^3+36796125r^2+23950143r+5675670}$	0

Using the above table, then the solution $y_2(x)$ is

$$\begin{aligned} y_2(x) &= b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 + b_7x^7 + b_8x^8 \dots \\ &= 1 - x + O(x^8) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned} y_h(x) &= c_1y_1(x) + c_2y_2(x) \\ &= c_1\sqrt{x} \left(1 - \frac{x}{6} - \frac{x^2}{120} - \frac{x^3}{1680} - \frac{x^4}{24192} - \frac{x^5}{380160} - \frac{x^6}{6589440} - \frac{x^7}{125798400} + O(x^8) \right) \\ &\quad + c_2(1 - x + O(x^8)) \end{aligned}$$

Hence the final solution is

$$\begin{aligned} y &= y_h \\ &= c_1\sqrt{x} \left(1 - \frac{x}{6} - \frac{x^2}{120} - \frac{x^3}{1680} - \frac{x^4}{24192} - \frac{x^5}{380160} - \frac{x^6}{6589440} - \frac{x^7}{125798400} + O(x^8) \right) \\ &\quad + c_2(1 - x + O(x^8)) \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned} y &= c_1\sqrt{x} \left(1 - \frac{x}{6} - \frac{x^2}{120} - \frac{x^3}{1680} - \frac{x^4}{24192} - \frac{x^5}{380160} - \frac{x^6}{6589440} - \frac{x^7}{125798400} + O(x^8) \right) \\ &\quad + c_2(1 - x + O(x^8)) \end{aligned}$$

Verification of solutions

$$y = c_1 \sqrt{x} \left(1 - \frac{x}{6} - \frac{x^2}{120} - \frac{x^3}{1680} - \frac{x^4}{24192} - \frac{x^5}{380160} - \frac{x^6}{6589440} - \frac{x^7}{125798400} + O(x^8) \right) + c_2(1 - x + O(x^8))$$

Verified OK.

22.13.1 Maple step by step solution

Let's solve

$$2y''x + (1 - x)y' + y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = \frac{(x-1)y'}{2x} - \frac{y}{2x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - \frac{(x-1)y'}{2x} + \frac{y}{2x} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$[P_2(x) = -\frac{x-1}{2x}, P_3(x) = \frac{1}{2x}]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = \frac{1}{2}$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 0$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$2y''x + (1 - x)y' + y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

□ Rewrite ODE with series expansions

- Convert $x^m \cdot y'$ to series expansion for $m = 0..1$

$$x^m \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1+m}$$

- Shift index using $k \rightarrow k+1-m$

$$x^m \cdot y' = \sum_{k=-1+m}^{\infty} a_{k+1-m} (k+1-m+r) x^{k+r}$$

- Convert $x \cdot y''$ to series expansion

$$x \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r-1}$$

- Shift index using $k \rightarrow k+1$

$$x \cdot y'' = \sum_{k=-1}^{\infty} a_{k+1} (k+1+r)(k+r) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0 r(-1+2r) x^{-1+r} + \left(\sum_{k=0}^{\infty} (a_{k+1} (k+1+r)(2k+1+2r) - a_k (k+r-1)) x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$r(-1+2r) = 0$$

- Values of r that satisfy the indicial equation

$$r \in \left\{ 0, \frac{1}{2} \right\}$$

- Each term in the series must be 0, giving the recursion relation

$$2(k+1+r) \left(k + \frac{1}{2} + r\right) a_{k+1} - a_k (k+r-1) = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+1} = \frac{a_k (k+r-1)}{(k+1+r)(2k+1+2r)}$$

- Recursion relation for $r = 0$; series terminates at $k = 1$

$$a_{k+1} = \frac{a_k (k-1)}{(k+1)(2k+1)}$$

- Apply recursion relation for $k = 0$

$$a_1 = -a_0$$

- Terminating series solution of the ODE for $r = 0$. Use reduction of order to find the second li

$$y = a_0 \cdot (1 - x)$$

- Recursion relation for $r = \frac{1}{2}$

$$a_{k+1} = \frac{a_k(k-\frac{1}{2})}{(k+\frac{3}{2})(2k+2)}$$

- Solution for $r = \frac{1}{2}$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+\frac{1}{2}}, a_{k+1} = \frac{a_k(k-\frac{1}{2})}{(k+\frac{3}{2})(2k+2)} \right]$$

- Combine solutions and rename parameters

$$\left[y = a_0 \cdot (1 - x) + \left(\sum_{k=0}^{\infty} b_k x^{k+\frac{1}{2}} \right), b_{k+1} = \frac{b_k(k-\frac{1}{2})}{(k+\frac{3}{2})(2k+2)} \right]$$

Maple trace Kovacic algorithm successful

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
A Liouvillian solution exists
Reducible group (found an exponential solution)
Group is reducible, not completely reducible
Solution has integrals. Trying a special function solution free of integrals...
-> Trying a solution in terms of special functions:
  -> Bessel
  -> elliptic
  -> Legendre
  <- Kummer successful
<- special function solution successful
  -> Trying to convert hypergeometric functions to elementary form...
  <- elementary form is not straightforward to achieve - returning special function solu
<- Kovacics algorithm successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 40

Order:=8;

```
dsolve(2*x*diff(y(x),x$2)+(1-x)*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = c_1 \sqrt{x} \left(1 - \frac{1}{6}x - \frac{1}{120}x^2 - \frac{1}{1680}x^3 - \frac{1}{24192}x^4 - \frac{1}{380160}x^5 - \frac{1}{6589440}x^6 - \frac{1}{125798400}x^7 + O(x^8) \right) + c_2(1 - x + O(x^8))$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 111

```
AsymptoticDSolveValue[2*x*y'[x]+(1-x)*y'[x]+(2+x)*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \sqrt{x} \left(\frac{17333x^7}{48432384000} - \frac{34817x^6}{691891200} - \frac{1171x^5}{4435200} + \frac{121x^4}{40320} + \frac{37x^3}{1680} - \frac{3x^2}{40} - \frac{x}{2} + 1 \right) + c_2 \left(\frac{4x^7}{143325} - \frac{x^6}{8400} - \frac{19x^5}{6300} - \frac{x^4}{840} + \frac{2x^3}{15} + \frac{x^2}{6} - 2x + 1 \right)$$

22.14 problem 2(f)

22.14.1 Maple step by step solution 3763

Internal problem ID [6489]

Internal file name [OUTPUT/5737_Sunday_June_05_2022_03_51_51_PM_30794761/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Problems for review and discover. (A) Drill Exercises . Page 194

Problem number: 2(f).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Repeated root"**

Maple gives the following as the ode type

[_Laguerre]

$$xy'' - (x - 1)y' + 2y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$xy'' + (1 - x)y' + 2y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{x-1}{x}$$
$$q(x) = \frac{2}{x}$$

Table 510: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{x-1}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = \frac{2}{x}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$xy'' + (1 - x)y' + 2y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) x \\ & + (1-x) \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + 2 \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \end{aligned} \tag{1}$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r-1} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) \\ & + \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} 2a_n x^{n+r} \right) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r-1$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r-1} and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=0}^{\infty} (-x^{n+r} a_n (n+r)) &= \sum_{n=1}^{\infty} (-a_{n-1} (n+r-1) x^{n+r-1}) \\ \sum_{n=0}^{\infty} 2a_n x^{n+r} &= \sum_{n=1}^{\infty} 2a_{n-1} x^{n+r-1} \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r-1$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r-1} a_n (n+r) (n+r-1) \right) + \sum_{n=1}^{\infty} (-a_{n-1} (n+r-1) x^{n+r-1}) \\ & + \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=1}^{\infty} 2a_{n-1} x^{n+r-1} \right) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r-1} a_n (n+r) (n+r-1) + (n+r) a_n x^{n+r-1} = 0$$

When $n=0$ the above becomes

$$x^{-1+r} a_0 r (-1+r) + r a_0 x^{-1+r} = 0$$

Or

$$(x^{-1+r} r (-1+r) + r x^{-1+r}) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$x^{-1+r} r^2 = 0$$

Since the above is true for all x then the indicial equation becomes

$$r^2 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = 0$$

$$r_2 = 0$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$x^{-1+r}r^2 = 0$$

Solving for r gives the roots of the indicial equation as Since the root of the indicial equation is repeated, then we can construct two linearly independent solutions. The first solution has the form

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+r} \quad (1A)$$

Now the second solution y_2 is found using

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right) \quad (1B)$$

Then the general solution will be

$$y = c_1 y_1(x) + c_2 y_2(x)$$

In Eq (1B) the sum starts from 1 and not zero. In Eq (1A), a_0 is never zero, and is arbitrary and is typically taken as $a_0 = 1$, and $\{c_1, c_2\}$ are two arbitrary constants of integration which can be found from initial conditions. We start by finding the first solution $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) - a_{n-1}(n+r-1) + a_n(n+r) + 2a_{n-1} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-1}(n+r-3)}{n^2 + 2nr + r^2} \quad (4)$$

Which for the root $r = 0$ becomes

$$a_n = \frac{a_{n-1}(n-3)}{n^2} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = \frac{-2+r}{(r+1)^2}$$

Which for the root $r = 0$ becomes

$$a_1 = -2$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-2+r}{(r+1)^2}$	-2

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{(-2+r)(-1+r)}{(r+1)^2(r+2)^2}$$

Which for the root $r = 0$ becomes

$$a_2 = \frac{1}{2}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-2+r}{(r+1)^2}$	-2
a_2	$\frac{(-2+r)(-1+r)}{(r+1)^2(r+2)^2}$	$\frac{1}{2}$

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{(-2+r)(-1+r)r}{(r+1)^2(r+2)^2(r+3)^2}$$

Which for the root $r = 0$ becomes

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-2+r}{(r+1)^2}$	-2
a_2	$\frac{(-2+r)(-1+r)}{(r+1)^2(r+2)^2}$	$\frac{1}{2}$
a_3	$\frac{(-2+r)(-1+r)r}{(r+1)^2(r+2)^2(r+3)^2}$	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{(-2+r)(-1+r)r}{(r+1)(r+2)^2(r+3)^2(r+4)^2}$$

Which for the root $r = 0$ becomes

$$a_4 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-2+r}{(r+1)^2}$	-2
a_2	$\frac{(-2+r)(-1+r)}{(r+1)^2(r+2)^2}$	$\frac{1}{2}$
a_3	$\frac{(-2+r)(-1+r)r}{(r+1)^2(r+2)^2(r+3)^2}$	0
a_4	$\frac{(-2+r)(-1+r)r}{(r+1)(r+2)^2(r+3)^2(r+4)^2}$	0

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{(-2+r)(-1+r)r}{(r+1)(r+2)(r+3)^2(r+4)^2(r+5)^2}$$

Which for the root $r = 0$ becomes

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-2+r}{(r+1)^2}$	-2
a_2	$\frac{(-2+r)(-1+r)}{(r+1)^2(r+2)^2}$	$\frac{1}{2}$
a_3	$\frac{(-2+r)(-1+r)r}{(r+1)^2(r+2)^2(r+3)^2}$	0
a_4	$\frac{(-2+r)(-1+r)r}{(r+1)(r+2)^2(r+3)^2(r+4)^2}$	0
a_5	$\frac{(-2+r)(-1+r)r}{(r+1)(r+2)(r+3)^2(r+4)^2(r+5)^2}$	0

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{(-2+r)(-1+r)r}{(r+1)(r+2)(r+3)(r+4)^2(r+5)^2(r+6)^2}$$

Which for the root $r = 0$ becomes

$$a_6 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-2+r}{(r+1)^2}$	-2
a_2	$\frac{(-2+r)(-1+r)}{(r+1)^2(r+2)^2}$	$\frac{1}{2}$
a_3	$\frac{(-2+r)(-1+r)r}{(r+1)^2(r+2)^2(r+3)^2}$	0
a_4	$\frac{(-2+r)(-1+r)r}{(r+1)(r+2)^2(r+3)^2(r+4)^2}$	0
a_5	$\frac{(-2+r)(-1+r)r}{(r+1)(r+2)(r+3)^2(r+4)^2(r+5)^2}$	0
a_6	$\frac{(-2+r)(-1+r)r}{(r+1)(r+2)(r+3)(r+4)^2(r+5)^2(r+6)^2}$	0

For $n = 7$, using the above recursive equation gives

$$a_7 = \frac{(-2+r)(-1+r)r}{(r+1)(r+2)(r+3)(r+4)(r+5)^2(r+6)^2(r+7)^2}$$

Which for the root $r = 0$ becomes

$$a_7 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{-2+r}{(r+1)^2}$	-2
a_2	$\frac{(-2+r)(-1+r)}{(r+1)^2(r+2)^2}$	$\frac{1}{2}$
a_3	$\frac{(-2+r)(-1+r)r}{(r+1)^2(r+2)^2(r+3)^2}$	0
a_4	$\frac{(-2+r)(-1+r)r}{(r+1)(r+2)^2(r+3)^2(r+4)^2}$	0
a_5	$\frac{(-2+r)(-1+r)r}{(r+1)(r+2)(r+3)^2(r+4)^2(r+5)^2}$	0
a_6	$\frac{(-2+r)(-1+r)r}{(r+1)(r+2)(r+3)(r+4)^2(r+5)^2(r+6)^2}$	0
a_7	$\frac{(-2+r)(-1+r)r}{(r+1)(r+2)(r+3)(r+4)(r+5)^2(r+6)^2(r+7)^2}$	0

Using the above table, then the first solution $y_1(x)$ becomes

$$\begin{aligned} y_1(x) &= a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots \\ &= 1 - 2x + \frac{x^2}{2} + O(x^8) \end{aligned}$$

Now the second solution is found. The second solution is given by

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right)$$

Where b_n is found using

$$b_n = \frac{d}{dr} a_{n,r}$$

And the above is then evaluated at $r = 0$. The above table for $a_{n,r}$ is used for this purpose. Computing the derivatives gives the following table

n	$b_{n,r}$	a_n	$b_{n,r} = \frac{d}{dr} a_{n,r}$
b_0	1	1	N/A since b_n starts from 1
b_1	$\frac{-2+r}{(r+1)^2}$	-2	$\frac{-r+5}{(r+1)^3}$
b_2	$\frac{(-2+r)(-1+r)}{(r+1)^2(r+2)^2}$	$\frac{1}{2}$	$\frac{-2r^3+9r^2+5r-18}{(r+1)^3(r+2)^3}$
b_3	$\frac{(-2+r)(-1+r)r}{(r+1)^2(r+2)^2(r+3)^2}$	0	$\frac{-3r^5+6r^4+37r^3-18r^2-58r+12}{(r+1)^3(r+2)^3(r+3)^3}$
b_4	$\frac{(-2+r)(-1+r)r}{(r+1)(r+2)^2(r+3)^2(r+4)^2}$	0	$\frac{-4r^6-6r^5+72r^4+124r^3-158r^2-196r+48}{(r+1)^2(r+2)^3(r+3)^3(r+4)^3}$
b_5	$\frac{(-2+r)(-1+r)r}{(r+1)(r+2)(r+3)^2(r+4)^2(r+5)^2}$	0	$\frac{-5r^7-30r^6+50r^5+510r^4+431r^3-1008r^2-908r+240}{(r+1)^2(r+2)^2(r+3)^3(r+4)^3(r+5)^3}$
b_6	$\frac{(-2+r)(-1+r)r}{(r+1)(r+2)(r+3)(r+4)^2(r+5)^2(r+6)^2}$	0	$\frac{-6r^8-69r^7-145r^6+855r^5+3587r^4+1542r^3-7036r^2-5208r+1440}{(r+1)^2(r+2)^2(r+3)^2(r+4)^3(r+5)^3(r+6)^3}$
b_7	$\frac{(-2+r)(-1+r)r}{(r+1)(r+2)(r+3)(r+4)(r+5)^2(r+6)^2(r+7)^2}$	0	$\frac{-7r^9-126r^8-700r^7-238r^6+9863r^5+27356r^4+3900r^3-55072r^2-35376r+1008}{(r+1)^2(r+2)^2(r+3)^2(r+4)^2(r+5)^3(r+6)^3(r+7)^3}$

The above table gives all values of b_n needed. Hence the second solution is

$$\begin{aligned}
y_2(x) &= y_1(x) \ln(x) + b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 + b_7x^7 + b_8x^8 \dots \\
&= \left(1 - 2x + \frac{x^2}{2} + O(x^8)\right) \ln(x) + 5x - \frac{9x^2}{4} + \frac{x^3}{18} + \frac{x^4}{288} + \frac{x^5}{3600} + \frac{x^6}{43200} + \frac{x^7}{529200} \\
&\quad + O(x^8)
\end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
y_h(x) &= c_1y_1(x) + c_2y_2(x) \\
&= c_1 \left(1 - 2x + \frac{x^2}{2} + O(x^8)\right) + c_2 \left(\left(1 - 2x + \frac{x^2}{2} + O(x^8)\right) \ln(x) + 5x - \frac{9x^2}{4} \right. \\
&\quad \left. + \frac{x^3}{18} + \frac{x^4}{288} + \frac{x^5}{3600} + \frac{x^6}{43200} + \frac{x^7}{529200} + O(x^8) \right)
\end{aligned}$$

Hence the final solution is

$$\begin{aligned}
y &= y_h \\
&= c_1 \left(1 - 2x + \frac{x^2}{2} + O(x^8)\right) + c_2 \left(\left(1 - 2x + \frac{x^2}{2} + O(x^8)\right) \ln(x) + 5x - \frac{9x^2}{4} + \frac{x^3}{18} \right. \\
&\quad \left. + \frac{x^4}{288} + \frac{x^5}{3600} + \frac{x^6}{43200} + \frac{x^7}{529200} + O(x^8) \right)
\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \left(1 - 2x + \frac{x^2}{2} + O(x^8) \right) + c_2 \left(\left(1 - 2x + \frac{x^2}{2} + O(x^8) \right) \ln(x) + 5x - \frac{9x^2}{4} + \frac{x^3}{18} + \frac{x^4}{288} + \frac{x^5}{3600} + \frac{x^6}{43200} + \frac{x^7}{529200} + O(x^8) \right) \quad (1)$$

Verification of solutions

$$y = c_1 \left(1 - 2x + \frac{x^2}{2} + O(x^8) \right) + c_2 \left(\left(1 - 2x + \frac{x^2}{2} + O(x^8) \right) \ln(x) + 5x - \frac{9x^2}{4} + \frac{x^3}{18} + \frac{x^4}{288} + \frac{x^5}{3600} + \frac{x^6}{43200} + \frac{x^7}{529200} + O(x^8) \right)$$

Verified OK.

22.14.1 Maple step by step solution

Let's solve

$$y''x + (1-x)y' + 2y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = \frac{(x-1)y'}{x} - \frac{2y}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - \frac{(x-1)y'}{x} + \frac{2y}{x} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$[P_2(x) = -\frac{x-1}{x}, P_3(x) = \frac{2}{x}]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 1$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 0$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$y''x + (1 - x)y' + 2y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y'$ to series expansion for $m = 0..1$

$$x^m \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1+m}$$

- Shift index using $k \rightarrow k+1-m$

$$x^m \cdot y' = \sum_{k=-1+m}^{\infty} a_{k+1-m} (k+1-m+r) x^{k+r}$$

- Convert $x \cdot y''$ to series expansion

$$x \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r-1}$$

- Shift index using $k \rightarrow k+1$

$$x \cdot y'' = \sum_{k=-1}^{\infty} a_{k+1} (k+1+r)(k+r) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0 r^2 x^{-1+r} + \left(\sum_{k=0}^{\infty} (a_{k+1} (k+1+r)^2 - a_k (k+r-2)) x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$r^2 = 0$$

- Values of r that satisfy the indicial equation

$$r = 0$$

- Each term in the series must be 0, giving the recursion relation

$$a_{k+1} (k+1)^2 - a_k (k-2) = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+1} = \frac{a_k (k-2)}{(k+1)^2}$$

- Recursion relation for $r = 0$; series terminates at $k = 2$

$$a_{k+1} = \frac{a_k(k-2)}{(k+1)^2}$$

- Apply recursion relation for $k = 0$

$$a_1 = -2a_0$$

- Apply recursion relation for $k = 1$

$$a_2 = -\frac{a_1}{4}$$

- Express in terms of a_0

$$a_2 = \frac{a_0}{2}$$

- Terminating series solution of the ODE for $r = 0$. Use reduction of order to find the second li

$$y = a_0 \cdot \left(1 - 2x + \frac{1}{2}x^2\right)$$

Maple trace Kovacic algorithm successful

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Reducible group (found an exponential solution)
    Group is reducible, not completely reducible
<- Kovacics algorithm successful`

```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 51

```

Order:=8;
dsolve(x*diff(y(x),x$2)-(x-1)*diff(y(x),x)+2*y(x)=0,y(x),type='series',x=0);

```

$$y(x) = (c_2 \ln(x) + c_1) \left(1 - 2x + \frac{1}{2}x^2 + O(x^8)\right) + \left(5x - \frac{9}{4}x^2 + \frac{1}{18}x^3 + \frac{1}{288}x^4 + \frac{1}{3600}x^5 + \frac{1}{43200}x^6 + \frac{1}{529200}x^7 + O(x^8)\right) c_2$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 83

```
AsymptoticDSolveValue[x*y''[x]-(x-1)*y'[x]+2*y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(\frac{x^2}{2} - 2x + 1 \right) + c_2 \left(\frac{x^7}{529200} + \frac{x^6}{43200} + \frac{x^5}{3600} + \frac{x^4}{288} + \frac{x^3}{18} - \frac{9x^2}{4} + \left(\frac{x^2}{2} - 2x + 1 \right) \log(x) + 5x \right)$$

22.15 problem 2(g)

22.15.1 Maple step by step solution 3776

Internal problem ID [6490]

Internal file name [OUTPUT/5738_Sunday_June_05_2022_03_51_54_PM_44733112/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Problems for review and discover. (A) Drill Exercises . Page 194

Problem number: 2(g).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Complex roots"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$x^2y'' + x(1 - x)y' + y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$x^2y'' + (-x^2 + x)y' + y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = -\frac{x-1}{x}$$
$$q(x) = \frac{1}{x^2}$$

Table 512: Table $p(x), q(x)$ singularities.

$p(x) = -\frac{x-1}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = \frac{1}{x^2}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$x^2 y'' + (-x^2 + x) y' + y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned}
 & x^2 \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) \\
 & + (-x^2 + x) \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0
 \end{aligned} \tag{1}$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=0}^{\infty} (-x^{1+n+r} a_n (n+r)) \\ & + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} (-x^{1+n+r} a_n (n+r)) = \sum_{n=1}^{\infty} (-a_{n-1} (n+r-1) x^{n+r})$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) (n+r-1) \right) + \sum_{n=1}^{\infty} (-a_{n-1} (n+r-1) x^{n+r}) \\ & + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r} a_n (n+r) (n+r-1) + x^{n+r} a_n (n+r) + a_n x^{n+r} = 0$$

When $n=0$ the above becomes

$$x^r a_0 r (-1+r) + x^r a_0 r + a_0 x^r = 0$$

Or

$$(x^r r (-1+r) + x^r r + x^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(r^2 + 1) x^r = 0$$

Since the above is true for all x then the indicial equation becomes

$$r^2 + 1 = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= i \\ r_2 &= -i \end{aligned}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(r^2 + 1) x^r = 0$$

Solving for r gives the roots of the indicial equation as Since the roots are complex conjugates, then two linearly independent solutions can be constructed using

$$\begin{aligned} y_1(x) &= x^{r_1} \left(\sum_{n=0}^{\infty} a_n x^n \right) \\ y_2(x) &= x^{r_2} \left(\sum_{n=0}^{\infty} b_n x^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(x) &= \sum_{n=0}^{\infty} a_n x^{n+i} \\ y_2(x) &= \sum_{n=0}^{\infty} b_n x^{n-i} \end{aligned}$$

$y_1(x)$ is found first. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) - a_{n-1}(n+r-1) + a_n(n+r) + a_n = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-1}(n+r-1)}{n^2 + 2nr + r^2 + 1} \quad (4)$$

Which for the root $r = i$ becomes

$$a_n = \frac{a_{n-1}(n-1+i)}{n(2i+n)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = i$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = \frac{r}{r^2 + 2r + 2}$$

Which for the root $r = i$ becomes

$$a_1 = \frac{2}{5} + \frac{i}{5}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{r}{r^2+2r+2}$	$\frac{2}{5} + \frac{i}{5}$

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{r(r+1)}{(r^2 + 2r + 2)(r^2 + 4r + 5)}$$

Which for the root $r = i$ becomes

$$a_2 = \frac{1}{10} + \frac{i}{20}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{r}{r^2+2r+2}$	$\frac{2}{5} + \frac{i}{5}$
a_2	$\frac{r(r+1)}{(r^2+2r+2)(r^2+4r+5)}$	$\frac{1}{10} + \frac{i}{20}$

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{r(r+1)(2+r)}{(r^2 + 2r + 2)(r^2 + 4r + 5)(r^2 + 6r + 10)}$$

Which for the root $r = i$ becomes

$$a_3 = \frac{17}{780} + \frac{i}{130}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{r}{r^2+2r+2}$	$\frac{2}{5} + \frac{i}{5}$
a_2	$\frac{r(r+1)}{(r^2+2r+2)(r^2+4r+5)}$	$\frac{1}{10} + \frac{i}{20}$
a_3	$\frac{r(r+1)(2+r)}{(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)}$	$\frac{17}{780} + \frac{i}{130}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{r(r+1)(2+r)(3+r)}{(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)}$$

Which for the root $r = i$ becomes

$$a_4 = \frac{5}{1248} + \frac{i}{1248}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{r}{r^2+2r+2}$	$\frac{2}{5} + \frac{i}{5}$
a_2	$\frac{r(r+1)}{(r^2+2r+2)(r^2+4r+5)}$	$\frac{1}{10} + \frac{i}{20}$
a_3	$\frac{r(r+1)(2+r)}{(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)}$	$\frac{17}{780} + \frac{i}{130}$
a_4	$\frac{r(r+1)(2+r)(3+r)}{(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)}$	$\frac{5}{1248} + \frac{i}{1248}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{r(r+1)(2+r)(3+r)(4+r)}{(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)(r^2+10r+26)}$$

Which for the root $r = i$ becomes

$$a_5 = \frac{113}{180960} + \frac{7i}{180960}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{r}{r^2+2r+2}$	$\frac{2}{5} + \frac{i}{5}$
a_2	$\frac{r(r+1)}{(r^2+2r+2)(r^2+4r+5)}$	$\frac{1}{10} + \frac{i}{20}$
a_3	$\frac{r(r+1)(2+r)}{(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)}$	$\frac{17}{780} + \frac{i}{130}$
a_4	$\frac{r(r+1)(2+r)(3+r)}{(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)}$	$\frac{5}{1248} + \frac{i}{1248}$
a_5	$\frac{r(r+1)(2+r)(3+r)(4+r)}{(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)(r^2+10r+26)}$	$\frac{113}{180960} + \frac{7i}{180960}$

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{r(r+1)(2+r)(3+r)(4+r)(5+r)}{(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)(r^2+10r+26)(r^2+12r+37)}$$

Which for the root $r = i$ becomes

$$a_6 = \frac{911}{10857600} - \frac{19i}{3619200}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{r}{r^2+2r+2}$	$\frac{2}{5} + \frac{i}{5}$
a_2	$\frac{r(r+1)}{(r^2+2r+2)(r^2+4r+5)}$	$\frac{1}{10} + \frac{i}{20}$
a_3	$\frac{r(r+1)(2+r)}{(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)}$	$\frac{17}{780} + \frac{i}{130}$
a_4	$\frac{r(r+1)(2+r)(3+r)}{(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)}$	$\frac{5}{1248} + \frac{i}{1248}$
a_5	$\frac{r(r+1)(2+r)(3+r)(4+r)}{(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)(r^2+10r+26)}$	$\frac{113}{180960} + \frac{7i}{180960}$
a_6	$\frac{r(r+1)(2+r)(3+r)(4+r)(5+r)}{(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)(r^2+10r+26)(r^2+12r+37)}$	$\frac{911}{10857600} - \frac{19i}{3619200}$

For $n = 7$, using the above recursive equation gives

$$a_7 = \frac{r(r+1)(2+r)(3+r)(4+r)(5+r)(6+r)}{(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)(r^2+10r+26)(r^2+12r+37)(r^2+14r+37)}$$

Which for the root $r = i$ becomes

$$a_7 = \frac{39799}{4028169600} - \frac{1009i}{575452800}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{r}{r^2+2r+2}$	$\frac{2}{5} + \frac{i}{5}$
a_2	$\frac{r(r+1)}{(r^2+2r+2)(r^2+4r+5)}$	$\frac{1}{10} + \frac{i}{20}$
a_3	$\frac{r(r+1)(2+r)}{(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)}$	$\frac{17}{780} + \frac{i}{130}$
a_4	$\frac{r(r+1)(2+r)(3+r)}{(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)}$	$\frac{5}{1248} + \frac{i}{1248}$
a_5	$\frac{r(r+1)(2+r)(3+r)(4+r)}{(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)(r^2+10r+26)}$	$\frac{113}{180960} + \frac{7i}{180960}$
a_6	$\frac{r(r+1)(2+r)(3+r)(4+r)(5+r)}{(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)(r^2+10r+26)(r^2+12r+37)}$	$\frac{911}{10857600} - \frac{19i}{3619200}$
a_7	$\frac{r(r+1)(2+r)(3+r)(4+r)(5+r)(6+r)}{(r^2+2r+2)(r^2+4r+5)(r^2+6r+10)(r^2+8r+17)(r^2+10r+26)(r^2+12r+37)(r^2+14r+50)}$	$\frac{39799}{4028169600} - \frac{1009i}{575452800}$

Using the above table, then the solution $y_1(x)$ is

$$\begin{aligned} y_1(x) &= x^i (a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots) \\ &= x^i \left(1 + \left(\frac{2}{5} + \frac{i}{5} \right) x + \left(\frac{1}{10} + \frac{i}{20} \right) x^2 + \left(\frac{17}{780} + \frac{i}{130} \right) x^3 + \left(\frac{5}{1248} + \frac{i}{1248} \right) x^4 \right. \\ &\quad + \left(\frac{113}{180960} + \frac{7i}{180960} \right) x^5 + \left(\frac{911}{10857600} - \frac{19i}{3619200} \right) x^6 \\ &\quad \left. + \left(\frac{39799}{4028169600} - \frac{1009i}{575452800} \right) x^7 + O(x^8) \right) \end{aligned}$$

The second solution $y_2(x)$ is found by taking the complex conjugate of $y_1(x)$ which gives

$$\begin{aligned} y_2(x) &= x^{-i} \left(1 + \left(\frac{2}{5} - \frac{i}{5} \right) x + \left(\frac{1}{10} - \frac{i}{20} \right) x^2 + \left(\frac{17}{780} - \frac{i}{130} \right) x^3 + \left(\frac{5}{1248} - \frac{i}{1248} \right) x^4 \right. \\ &\quad + \left(\frac{113}{180960} - \frac{7i}{180960} \right) x^5 + \left(\frac{911}{10857600} + \frac{19i}{3619200} \right) x^6 \\ &\quad \left. + \left(\frac{39799}{4028169600} + \frac{1009i}{575452800} \right) x^7 + O(x^8) \right) \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 y_h(x) &= c_1 y_1(x) + c_2 y_2(x) \\
 &= c_1 x^i \left(1 + \left(\frac{2}{5} + \frac{i}{5} \right) x + \left(\frac{1}{10} + \frac{i}{20} \right) x^2 + \left(\frac{17}{780} + \frac{i}{130} \right) x^3 + \left(\frac{5}{1248} + \frac{i}{1248} \right) x^4 \right. \\
 &\quad + \left(\frac{113}{180960} + \frac{7i}{180960} \right) x^5 + \left(\frac{911}{10857600} - \frac{19i}{3619200} \right) x^6 \\
 &\quad \left. + \left(\frac{39799}{4028169600} - \frac{1009i}{575452800} \right) x^7 + O(x^8) \right) \\
 &\quad + c_2 x^{-i} \left(1 + \left(\frac{2}{5} - \frac{i}{5} \right) x + \left(\frac{1}{10} - \frac{i}{20} \right) x^2 + \left(\frac{17}{780} - \frac{i}{130} \right) x^3 + \left(\frac{5}{1248} - \frac{i}{1248} \right) x^4 \right. \\
 &\quad + \left(\frac{113}{180960} - \frac{7i}{180960} \right) x^5 + \left(\frac{911}{10857600} + \frac{19i}{3619200} \right) x^6 \\
 &\quad \left. + \left(\frac{39799}{4028169600} + \frac{1009i}{575452800} \right) x^7 + O(x^8) \right)
 \end{aligned}$$

Hence the final solution is

$$\begin{aligned}
 y &= y_h \\
 &= c_1 x^i \left(1 + \left(\frac{2}{5} + \frac{i}{5} \right) x + \left(\frac{1}{10} + \frac{i}{20} \right) x^2 + \left(\frac{17}{780} + \frac{i}{130} \right) x^3 + \left(\frac{5}{1248} + \frac{i}{1248} \right) x^4 \right. \\
 &\quad + \left(\frac{113}{180960} + \frac{7i}{180960} \right) x^5 + \left(\frac{911}{10857600} - \frac{19i}{3619200} \right) x^6 \\
 &\quad \left. + \left(\frac{39799}{4028169600} - \frac{1009i}{575452800} \right) x^7 + O(x^8) \right) + c_2 x^{-i} \left(1 + \left(\frac{2}{5} - \frac{i}{5} \right) x \right. \\
 &\quad + \left(\frac{1}{10} - \frac{i}{20} \right) x^2 + \left(\frac{17}{780} - \frac{i}{130} \right) x^3 + \left(\frac{5}{1248} - \frac{i}{1248} \right) x^4 + \left(\frac{113}{180960} - \frac{7i}{180960} \right) x^5 \\
 &\quad \left. + \left(\frac{911}{10857600} + \frac{19i}{3619200} \right) x^6 + \left(\frac{39799}{4028169600} + \frac{1009i}{575452800} \right) x^7 + O(x^8) \right)
 \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned} y = & c_1 x^i \left(1 + \left(\frac{2}{5} + \frac{i}{5} \right) x + \left(\frac{1}{10} + \frac{i}{20} \right) x^2 + \left(\frac{17}{780} + \frac{i}{130} \right) x^3 + \left(\frac{5}{1248} + \frac{i}{1248} \right) x^4 \right. \\ & + \left(\frac{113}{180960} + \frac{7i}{180960} \right) x^5 + \left(\frac{911}{10857600} - \frac{19i}{3619200} \right) x^6 \\ & \left. + \left(\frac{39799}{4028169600} - \frac{1009i}{575452800} \right) x^7 + O(x^8) \right) \\ & + c_2 x^{-i} \left(1 + \left(\frac{2}{5} - \frac{i}{5} \right) x + \left(\frac{1}{10} - \frac{i}{20} \right) x^2 + \left(\frac{17}{780} - \frac{i}{130} \right) x^3 \right. \\ & + \left(\frac{5}{1248} - \frac{i}{1248} \right) x^4 + \left(\frac{113}{180960} - \frac{7i}{180960} \right) x^5 + \left(\frac{911}{10857600} + \frac{19i}{3619200} \right) x^6 \\ & \left. + \left(\frac{39799}{4028169600} + \frac{1009i}{575452800} \right) x^7 + O(x^8) \right) \end{aligned}$$

Verification of solutions

$$\begin{aligned} y = & c_1 x^i \left(1 + \left(\frac{2}{5} + \frac{i}{5} \right) x + \left(\frac{1}{10} + \frac{i}{20} \right) x^2 + \left(\frac{17}{780} + \frac{i}{130} \right) x^3 + \left(\frac{5}{1248} + \frac{i}{1248} \right) x^4 \right. \\ & + \left(\frac{113}{180960} + \frac{7i}{180960} \right) x^5 + \left(\frac{911}{10857600} - \frac{19i}{3619200} \right) x^6 \\ & \left. + \left(\frac{39799}{4028169600} - \frac{1009i}{575452800} \right) x^7 + O(x^8) \right) + c_2 x^{-i} \left(1 + \left(\frac{2}{5} - \frac{i}{5} \right) x \right. \\ & + \left(\frac{1}{10} - \frac{i}{20} \right) x^2 + \left(\frac{17}{780} - \frac{i}{130} \right) x^3 + \left(\frac{5}{1248} - \frac{i}{1248} \right) x^4 + \left(\frac{113}{180960} - \frac{7i}{180960} \right) x^5 \\ & \left. + \left(\frac{911}{10857600} + \frac{19i}{3619200} \right) x^6 + \left(\frac{39799}{4028169600} + \frac{1009i}{575452800} \right) x^7 + O(x^8) \right) \end{aligned}$$

Verified OK.

22.15.1 Maple step by step solution

Let's solve

$$x^2 y'' + (-x^2 + x) y' + y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{y}{x^2} + \frac{(x-1)y'}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' - \frac{(x-1)y'}{x} + \frac{y}{x^2} = 0$$

- Check to see if $x_0 = 0$ is a regular singular point

- Define functions

$$[P_2(x) = -\frac{x-1}{x}, P_3(x) = \frac{1}{x^2}]$$

- $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 1$$

- $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 1$$

- $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

- Multiply by denominators

$$x^2 y'' - x(x-1)y' + y = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

- Rewrite ODE with series expansions

- Convert $x^m \cdot y'$ to series expansion for $m = 1..2$

$$x^m \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1+m}$$

- Shift index using $k \rightarrow k+1-m$

$$x^m \cdot y' = \sum_{k=-1+m}^{\infty} a_{k+1-m} (k+1-m+r) x^{k+r}$$

- Convert $x^2 \cdot y''$ to series expansion

$$x^2 \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0(r^2+1)x^r + \left(\sum_{k=1}^{\infty} (a_k(k^2+2kr+r^2+1) - a_{k-1}(k+r-1)) x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation
 $r^2 + 1 = 0$
- Values of r that satisfy the indicial equation
 $r \in \{-1, 1\}$
- Each term in the series must be 0, giving the recursion relation
 $a_k(k^2 + 2kr + r^2 + 1) - a_{k-1}(k + r - 1) = 0$
- Shift index using $k \rightarrow k + 1$
 $a_{k+1}((k + 1)^2 + 2(k + 1)r + r^2 + 1) - a_k(k + r) = 0$
- Recursion relation that defines series solution to ODE

$$a_{k+1} = \frac{a_k(k+r)}{k^2 + 2kr + r^2 + 2k + 2r + 2}$$
- Recursion relation for $r = -1$

$$a_{k+1} = \frac{a_k(k-1)}{k^2 - 2k + 1 - 2 + 2k}$$
- Solution for $r = -1$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k-1}, a_{k+1} = \frac{a_k(k-1)}{k^2 - 2k + 1 - 2 + 2k} \right]$$
- Recursion relation for $r = 1$

$$a_{k+1} = \frac{a_k(k+1)}{k^2 + 2k + 1 + 2 + 2k}$$
- Solution for $r = 1$

$$\left[y = \sum_{k=0}^{\infty} a_k x^{k+1}, a_{k+1} = \frac{a_k(k+1)}{k^2 + 2k + 1 + 2 + 2k} \right]$$
- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k x^{k-1} \right) + \left(\sum_{k=0}^{\infty} b_k x^{k+1} \right), a_{k+1} = \frac{a_k(k-1)}{k^2 - 2k + 1 - 2 + 2k}, b_{k+1} = \frac{b_k(k+1)}{k^2 + 2k + 1 + 2 + 2k} \right]$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacic's algorithm  
<- No Liouvillian solutions exist  
-> Trying a solution in terms of special functions:  
  -> Bessel  
  -> elliptic  
  -> Legendre  
  <- Kummer successful  
<- special function solution successful`
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 85

```
Order:=8;  
dsolve(x^2*diff(y(x),x$2)+x*(1-x)*diff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$\begin{aligned} y(x) = & c_1 x^{-i} \left(1 + \left(\frac{2}{5} - \frac{i}{5} \right) x + \left(\frac{1}{10} - \frac{i}{20} \right) x^2 + \left(\frac{17}{780} - \frac{i}{130} \right) x^3 + \left(\frac{5}{1248} - \frac{i}{1248} \right) x^4 \right. \\ & + \left(\frac{113}{180960} - \frac{7i}{180960} \right) x^5 + \left(\frac{911}{10857600} + \frac{19i}{3619200} \right) x^6 \\ & \left. + \left(\frac{39799}{4028169600} + \frac{1009i}{575452800} \right) x^7 + O(x^8) \right) \\ & + c_2 x^i \left(1 + \left(\frac{2}{5} + \frac{i}{5} \right) x + \left(\frac{1}{10} + \frac{i}{20} \right) x^2 + \left(\frac{17}{780} + \frac{i}{130} \right) x^3 + \left(\frac{5}{1248} + \frac{i}{1248} \right) x^4 \right. \\ & + \left(\frac{113}{180960} + \frac{7i}{180960} \right) x^5 + \left(\frac{911}{10857600} - \frac{19i}{3619200} \right) x^6 \\ & \left. + \left(\frac{39799}{4028169600} - \frac{1009i}{575452800} \right) x^7 + O(x^8) \right) \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.017 (sec). Leaf size: 122

```
AsymptoticDSolveValue[x^2*y''[x]+x*(1-x)*y'[x]+y[x]==0,y[x],{x,0,7}]
```

$$\begin{aligned} y(x) \rightarrow & \left(\frac{59}{10857600} - \frac{17i}{10857600} \right) c_2 x^{-i} \left((14 + 5i)x^6 + (108 + 24i)x^5 + (720 + 60i)x^4 \right. \\ & \left. + (4080 - 240i)x^3 + (19440 - 3600i)x^2 + (77760 - 14400i)x + (169920 + 48960i) \right) \\ & + \left(\frac{59}{10857600} + \frac{17i}{10857600} \right) c_1 x^i \left((14 - 5i)x^6 + (108 - 24i)x^5 + (720 - 60i)x^4 \right. \\ & \left. + (4080 + 240i)x^3 + (19440 + 3600i)x^2 + (77760 + 14400i)x + (169920 - 48960i) \right) \end{aligned}$$

22.16 problem 2(h)

22.16.1 Maple step by step solution 3790

Internal problem ID [6491]

Internal file name [OUTPUT/5739_Sunday_June_05_2022_03_52_03_PM_28503853/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Problems for review and discover. (A) Drill Exercises . Page 194

Problem number: 2(h).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Repeated root"**

Maple gives the following as the ode type

```
[[_2nd_order , _exact , _linear , _homogeneous]]
```

$$xy'' + (1 + x)y' + y = 0$$

With the expansion point for the power series method at $x = 0$.

The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$xy'' + (1 + x)y' + y = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$y'' + p(x)y' + q(x)y = 0$$

Where

$$p(x) = \frac{1 + x}{x}$$
$$q(x) = \frac{1}{x}$$

Table 514: Table $p(x), q(x)$ singularities.

$p(x) = \frac{1+x}{x}$	
singularity	type
$x = 0$	“regular”

$q(x) = \frac{1}{x}$	
singularity	type
$x = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $x = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$xy'' + (1+x)y' + y = 0$$

Let the solution be represented as Frobenius power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^{n+r}$$

Then

$$y' = \sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1}$$

$$y'' = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2}$$

Substituting the above back into the ode gives

$$\left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n x^{n+r-2} \right) x + (1+x) \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \tag{1}$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r-1} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} x^{n+r} a_n (n+r) \right) \\ & + \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} a_n x^{n+r} \right) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of x be $n+r-1$ in each summation term. Going over each summation term above with power of x in it which is not already x^{n+r-1} and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=0}^{\infty} x^{n+r} a_n (n+r) &= \sum_{n=1}^{\infty} a_{n-1} (n+r-1) x^{n+r-1} \\ \sum_{n=0}^{\infty} a_n x^{n+r} &= \sum_{n=1}^{\infty} a_{n-1} x^{n+r-1} \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of x are the same and equal to $n+r-1$.

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} x^{n+r-1} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=1}^{\infty} a_{n-1} (n+r-1) x^{n+r-1} \right) \\ & + \left(\sum_{n=0}^{\infty} (n+r) a_n x^{n+r-1} \right) + \left(\sum_{n=1}^{\infty} a_{n-1} x^{n+r-1} \right) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$x^{n+r-1} a_n (n+r) (n+r-1) + (n+r) a_n x^{n+r-1} = 0$$

When $n=0$ the above becomes

$$x^{-1+r} a_0 r (-1+r) + r a_0 x^{-1+r} = 0$$

Or

$$(x^{-1+r} r (-1+r) + r x^{-1+r}) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$x^{-1+r} r^2 = 0$$

Since the above is true for all x then the indicial equation becomes

$$r^2 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = 0$$

$$r_2 = 0$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$x^{-1+r}r^2 = 0$$

Solving for r gives the roots of the indicial equation as Since the root of the indicial equation is repeated, then we can construct two linearly independent solutions. The first solution has the form

$$y_1(x) = \sum_{n=0}^{\infty} a_n x^{n+r} \quad (1A)$$

Now the second solution y_2 is found using

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right) \quad (1B)$$

Then the general solution will be

$$y = c_1 y_1(x) + c_2 y_2(x)$$

In Eq (1B) the sum starts from 1 and not zero. In Eq (1A), a_0 is never zero, and is arbitrary and is typically taken as $a_0 = 1$, and $\{c_1, c_2\}$ are two arbitrary constants of integration which can be found from initial conditions. We start by finding the first solution $y_1(x)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) + a_{n-1}(n+r-1) + a_n(n+r) + a_{n-1} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{a_{n-1}}{n+r} \quad (4)$$

Which for the root $r = 0$ becomes

$$a_n = -\frac{a_{n-1}}{n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = -\frac{1}{1+r}$$

Which for the root $r = 0$ becomes

$$a_1 = -1$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{1+r}$	-1

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{(1+r)(2+r)}$$

Which for the root $r = 0$ becomes

$$a_2 = \frac{1}{2}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{1+r}$	-1
a_2	$\frac{1}{(1+r)(2+r)}$	$\frac{1}{2}$

For $n = 3$, using the above recursive equation gives

$$a_3 = -\frac{1}{(3+r)(1+r)(2+r)}$$

Which for the root $r = 0$ becomes

$$a_3 = -\frac{1}{6}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{1+r}$	-1
a_2	$\frac{1}{(1+r)(2+r)}$	$\frac{1}{2}$
a_3	$-\frac{1}{(3+r)(1+r)(2+r)}$	$-\frac{1}{6}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{(4+r)(3+r)(1+r)(2+r)}$$

Which for the root $r = 0$ becomes

$$a_4 = \frac{1}{24}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{1+r}$	-1
a_2	$\frac{1}{(1+r)(2+r)}$	$\frac{1}{2}$
a_3	$-\frac{1}{(3+r)(1+r)(2+r)}$	$-\frac{1}{6}$
a_4	$\frac{1}{(4+r)(3+r)(1+r)(2+r)}$	$\frac{1}{24}$

For $n = 5$, using the above recursive equation gives

$$a_5 = -\frac{1}{(5+r)(4+r)(3+r)(1+r)(2+r)}$$

Which for the root $r = 0$ becomes

$$a_5 = -\frac{1}{120}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{1+r}$	-1
a_2	$\frac{1}{(1+r)(2+r)}$	$\frac{1}{2}$
a_3	$-\frac{1}{(3+r)(1+r)(2+r)}$	$-\frac{1}{6}$
a_4	$\frac{1}{(4+r)(3+r)(1+r)(2+r)}$	$\frac{1}{24}$
a_5	$-\frac{1}{(5+r)(4+r)(3+r)(1+r)(2+r)}$	$-\frac{1}{120}$

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{1}{(5+r)(4+r)(3+r)(1+r)(2+r)(6+r)}$$

Which for the root $r = 0$ becomes

$$a_6 = \frac{1}{720}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{1+r}$	-1
a_2	$\frac{1}{(1+r)(2+r)}$	$\frac{1}{2}$
a_3	$-\frac{1}{(3+r)(1+r)(2+r)}$	$-\frac{1}{6}$
a_4	$\frac{1}{(4+r)(3+r)(1+r)(2+r)}$	$\frac{1}{24}$
a_5	$-\frac{1}{(5+r)(4+r)(3+r)(1+r)(2+r)}$	$-\frac{1}{120}$
a_6	$\frac{1}{(5+r)(4+r)(3+r)(1+r)(2+r)(6+r)}$	$\frac{1}{720}$

For $n = 7$, using the above recursive equation gives

$$a_7 = -\frac{1}{(7+r)(5+r)(4+r)(3+r)(1+r)(2+r)(6+r)}$$

Which for the root $r = 0$ becomes

$$a_7 = -\frac{1}{5040}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{1+r}$	-1
a_2	$\frac{1}{(1+r)(2+r)}$	$\frac{1}{2}$
a_3	$-\frac{1}{(3+r)(1+r)(2+r)}$	$-\frac{1}{6}$
a_4	$\frac{1}{(4+r)(3+r)(1+r)(2+r)}$	$\frac{1}{24}$
a_5	$-\frac{1}{(5+r)(4+r)(3+r)(1+r)(2+r)}$	$-\frac{1}{120}$
a_6	$\frac{1}{(5+r)(4+r)(3+r)(1+r)(2+r)(6+r)}$	$\frac{1}{720}$
a_7	$-\frac{1}{(7+r)(5+r)(4+r)(3+r)(1+r)(2+r)(6+r)}$	$-\frac{1}{5040}$

Using the above table, then the first solution $y_1(x)$ becomes

$$\begin{aligned}
 y_1(x) &= a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5 + a_6x^6 + a_7x^7 + a_8x^8 \dots \\
 &= 1 - x + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + \frac{x^6}{720} - \frac{x^7}{5040} + O(x^8)
 \end{aligned}$$

Now the second solution is found. The second solution is given by

$$y_2(x) = y_1(x) \ln(x) + \left(\sum_{n=1}^{\infty} b_n x^{n+r} \right)$$

Where b_n is found using

$$b_n = \frac{d}{dr} a_{n,r}$$

And the above is then evaluated at $r = 0$. The above table for $a_{n,r}$ is used for this purpose. Computing the derivatives gives the following table

n	$b_{n,r}$	a_n	$b_{n,r} = \frac{d}{dr} a_{n,r}$	$b_n(r=0)$
b_0	1	1	N/A since b_n starts from 1	N/A
b_1	$-\frac{1}{1+r}$	-1	$\frac{1}{(1+r)^2}$	1
b_2	$\frac{1}{(1+r)(2+r)}$	$\frac{1}{2}$	$\frac{-3-2r}{(1+r)^2(2+r)^2}$	$-\frac{3}{4}$
b_3	$-\frac{1}{(3+r)(1+r)(2+r)}$	$-\frac{1}{6}$	$\frac{3r^2+12r+11}{(3+r)^2(1+r)^2(2+r)^2}$	$\frac{11}{36}$
b_4	$\frac{1}{(4+r)(3+r)(1+r)(2+r)}$	$\frac{1}{24}$	$\frac{-4r^3-30r^2-70r-50}{(4+r)^2(3+r)^2(1+r)^2(2+r)^2}$	$-\frac{25}{288}$
b_5	$-\frac{1}{(5+r)(4+r)(3+r)(1+r)(2+r)}$	$-\frac{1}{120}$	$\frac{5r^4+60r^3+255r^2+450r+274}{(5+r)^2(4+r)^2(3+r)^2(1+r)^2(2+r)^2}$	$\frac{137}{7200}$
b_6	$\frac{1}{(5+r)(4+r)(3+r)(1+r)(2+r)(6+r)}$	$\frac{1}{720}$	$\frac{-6r^5-105r^4-700r^3-2205r^2-3248r-1764}{(5+r)^2(4+r)^2(3+r)^2(1+r)^2(2+r)^2(6+r)^2}$	$-\frac{49}{14400}$
b_7	$-\frac{1}{(7+r)(5+r)(4+r)(3+r)(1+r)(2+r)(6+r)}$	$-\frac{1}{5040}$	$\frac{7r^6+168r^5+1610r^4+7840r^3+20307r^2+26264r+13068}{(7+r)^2(5+r)^2(4+r)^2(3+r)^2(1+r)^2(2+r)^2(6+r)^2}$	$\frac{121}{235200}$

The above table gives all values of b_n needed. Hence the second solution is

$$\begin{aligned}
y_2(x) &= y_1(x) \ln(x) + b_0 + b_1x + b_2x^2 + b_3x^3 + b_4x^4 + b_5x^5 + b_6x^6 + b_7x^7 + b_8x^8 \dots \\
&= \left(1 - x + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + \frac{x^6}{720} - \frac{x^7}{5040} + O(x^8) \right) \ln(x) \\
&\quad + x - \frac{3x^2}{4} + \frac{11x^3}{36} - \frac{25x^4}{288} + \frac{137x^5}{7200} - \frac{49x^6}{14400} + \frac{121x^7}{235200} + O(x^8)
\end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
y_h(x) &= c_1y_1(x) + c_2y_2(x) \\
&= c_1 \left(1 - x + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + \frac{x^6}{720} - \frac{x^7}{5040} + O(x^8) \right) \\
&\quad + c_2 \left(\left(1 - x + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + \frac{x^6}{720} - \frac{x^7}{5040} + O(x^8) \right) \ln(x) + x - \frac{3x^2}{4} \right. \\
&\quad \left. + \frac{11x^3}{36} - \frac{25x^4}{288} + \frac{137x^5}{7200} - \frac{49x^6}{14400} + \frac{121x^7}{235200} + O(x^8) \right)
\end{aligned}$$

Hence the final solution is

$$y = y_h$$

$$\begin{aligned}
&= c_1 \left(1 - x + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + \frac{x^6}{720} - \frac{x^7}{5040} + O(x^8) \right) \\
&\quad + c_2 \left(\left(1 - x + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + \frac{x^6}{720} - \frac{x^7}{5040} + O(x^8) \right) \ln(x) + x - \frac{3x^2}{4} \right. \\
&\quad \quad \quad \left. + \frac{11x^3}{36} - \frac{25x^4}{288} + \frac{137x^5}{7200} - \frac{49x^6}{14400} + \frac{121x^7}{235200} + O(x^8) \right)
\end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}
y &= c_1 \left(1 - x + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + \frac{x^6}{720} - \frac{x^7}{5040} + O(x^8) \right) \\
&\quad + c_2 \left(\left(1 - x + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + \frac{x^6}{720} - \frac{x^7}{5040} + O(x^8) \right) \ln(x) + x - \frac{3x^2}{4} \right. \\
&\quad \quad \quad \left. + \frac{11x^3}{36} - \frac{25x^4}{288} + \frac{137x^5}{7200} - \frac{49x^6}{14400} + \frac{121x^7}{235200} + O(x^8) \right)
\end{aligned}$$

Verification of solutions

$$\begin{aligned}
y &= c_1 \left(1 - x + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + \frac{x^6}{720} - \frac{x^7}{5040} + O(x^8) \right) \\
&\quad + c_2 \left(\left(1 - x + \frac{x^2}{2} - \frac{x^3}{6} + \frac{x^4}{24} - \frac{x^5}{120} + \frac{x^6}{720} - \frac{x^7}{5040} + O(x^8) \right) \ln(x) + x - \frac{3x^2}{4} \right. \\
&\quad \quad \quad \left. + \frac{11x^3}{36} - \frac{25x^4}{288} + \frac{137x^5}{7200} - \frac{49x^6}{14400} + \frac{121x^7}{235200} + O(x^8) \right)
\end{aligned}$$

Verified OK.

22.16.1 Maple step by step solution

Let's solve

$$y''x + (1+x)y' + y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{(1+x)y'}{x} - \frac{y}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{(1+x)y'}{x} + \frac{y}{x} = 0$$

□ Check to see if $x_0 = 0$ is a regular singular point

○ Define functions

$$[P_2(x) = \frac{1+x}{x}, P_3(x) = \frac{1}{x}]$$

○ $x \cdot P_2(x)$ is analytic at $x = 0$

$$(x \cdot P_2(x)) \Big|_{x=0} = 1$$

○ $x^2 \cdot P_3(x)$ is analytic at $x = 0$

$$(x^2 \cdot P_3(x)) \Big|_{x=0} = 0$$

○ $x = 0$ is a regular singular point

Check to see if $x_0 = 0$ is a regular singular point

$$x_0 = 0$$

• Multiply by denominators

$$y''x + (1+x)y' + y = 0$$

• Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^{k+r}$$

□ Rewrite ODE with series expansions

○ Convert $x^m \cdot y'$ to series expansion for $m = 0..1$

$$x^m \cdot y' = \sum_{k=0}^{\infty} a_k (k+r) x^{k+r-1+m}$$

○ Shift index using $k \rightarrow k+1-m$

$$x^m \cdot y' = \sum_{k=-1+m}^{\infty} a_{k+1-m} (k+1-m+r) x^{k+r}$$

○ Convert $x \cdot y''$ to series expansion

$$x \cdot y'' = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) x^{k+r-1}$$

○ Shift index using $k \rightarrow k+1$

$$x \cdot y'' = \sum_{k=-1}^{\infty} a_{k+1} (k+1+r)(k+r) x^{k+r}$$

Rewrite ODE with series expansions

$$a_0 r^2 x^{-1+r} + \left(\sum_{k=0}^{\infty} (a_{k+1}(k+1+r)^2 + a_k(k+1+r)) x^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation
 $r^2 = 0$
- Values of r that satisfy the indicial equation
 $r = 0$
- Each term in the series must be 0, giving the recursion relation
 $(k+1)(a_{k+1}(k+1) + a_k) = 0$
- Recursion relation that defines series solution to ODE

$$a_{k+1} = -\frac{a_k}{k+1}$$

- Recursion relation for $r = 0$

$$a_{k+1} = -\frac{a_k}{k+1}$$

- Solution for $r = 0$

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+1} = -\frac{a_k}{k+1} \right]$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
<- linear_1 successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 71

Order:=8;

```
dsolve(x*difff(y(x),x$2)+(x+1)*difff(y(x),x)+y(x)=0,y(x),type='series',x=0);
```

$$y(x) = (c_2 \ln(x) + c_1) \left(1 - x + \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{24}x^4 - \frac{1}{120}x^5 + \frac{1}{720}x^6 - \frac{1}{5040}x^7 + O(x^8) \right) \\ + \left(x - \frac{3}{4}x^2 + \frac{11}{36}x^3 - \frac{25}{288}x^4 + \frac{137}{7200}x^5 - \frac{49}{14400}x^6 + \frac{121}{235200}x^7 + O(x^8) \right) c_2$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 151

```
AsymptoticDSolveValue[x*y''[x]+(x+1)*y'[x]+y[x]==0,y[x],{x,0,7}]
```

$$y(x) \rightarrow c_1 \left(-\frac{x^7}{5040} + \frac{x^6}{720} - \frac{x^5}{120} + \frac{x^4}{24} - \frac{x^3}{6} + \frac{x^2}{2} - x + 1 \right) + c_2 \left(\frac{121x^7}{235200} - \frac{49x^6}{14400} + \frac{137x^5}{7200} \right. \\ \left. - \frac{25x^4}{288} + \frac{11x^3}{36} - \frac{3x^2}{4} + \left(-\frac{x^7}{5040} + \frac{x^6}{720} - \frac{x^5}{120} + \frac{x^4}{24} - \frac{x^3}{6} + \frac{x^2}{2} - x + 1 \right) \log(x) + x \right)$$

22.17 problem 3(a)

Internal problem ID [6492]

Internal file name [OUTPUT/5740_Sunday_June_05_2022_03_52_06_PM_96721303/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Problems for review and discover. (A) Drill Exercises . Page 194

Problem number: 3(a).

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**unknown**"

Maple gives the following as the ode type

```
[[_3rd_order , _with_linear_symmetries]]
```

Unable to solve or complete the solution.

Unable to parse ODE.

Maple trace

```
`Methods for third order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying high order exact linear fully integrable
trying to convert to a linear ODE with constant coefficients
trying differential order: 3; missing the dependent variable
trying Louvillian solutions for 3rd order ODEs, imprimitive case
-> pFq: Equivalence to the 3F2 or one of its 3 confluent cases under a power @ Moebius
<- pFq successful: received ODE is equivalent to the 1F2 ODE, case c = 0 `
```

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 1219

Order:=8;

dsolve(x^3*diff(y(x),x\$3)+2*x^2*diff(y(x),x\$2)+(x+x^2)*diff(y(x),x)+x*y(x)=0,y(x),type='series')

$$\begin{aligned}
 y(x) = & c_1 \sqrt{x} x^{-\frac{i\sqrt{3}}{2}} \left(1 + \frac{1}{i\sqrt{3}-1} x - \frac{1}{2+6i\sqrt{3}} x^2 \right. \\
 & - \frac{1}{18} \frac{1}{\left(\frac{\sqrt{3}}{3}+i\right)(-2+i\sqrt{3})(\sqrt{3}+i)} x^3 + \frac{1}{-1728-480i\sqrt{3}} x^4 + \frac{1}{3360i\sqrt{3}+50400} x^5 \\
 & - \frac{1}{720} \frac{1}{(\sqrt{3}+6i)(\sqrt{3}+5i)(\sqrt{3}+4i)(\sqrt{3}+3i)(2i+\sqrt{3})(\sqrt{3}+i)} x^6 \\
 & \left. - \frac{1}{5040} \frac{1}{(-2+i\sqrt{3})(\sqrt{3}+7i)(\sqrt{3}+6i)(\sqrt{3}+5i)(\sqrt{3}+4i)(\sqrt{3}+3i)(\sqrt{3}+i)} x^7 \right. \\
 & \left. + O(x^8) \right) \\
 & + c_2 \sqrt{x} x^{\frac{i\sqrt{3}}{2}} \left(1 + \frac{1}{-i\sqrt{3}-1} x + \frac{1}{6i\sqrt{3}-2} x^2 + \frac{1}{6} \frac{1}{(\sqrt{3}-2i)(-i+\sqrt{3})(i\sqrt{3}+3)} x^3 \right. \\
 & + \frac{1}{24} \frac{1}{(-\sqrt{3}+2i)(i\sqrt{3}+4)(i\sqrt{3}+3)(-i+\sqrt{3})} x^4 \\
 & + \frac{1}{120} \frac{1}{(\sqrt{3}-2i)(-i+\sqrt{3})(i\sqrt{3}+5)(i\sqrt{3}+4)(i\sqrt{3}+3)} x^5 \\
 & + \frac{1}{720} \frac{1}{(-\sqrt{3}+2i)(i\sqrt{3}+6)(i\sqrt{3}+5)(i\sqrt{3}+4)(i\sqrt{3}+3)(-i+\sqrt{3})} x^6 \\
 & \left. + \frac{1}{5040} \frac{1}{(\sqrt{3}-2i)(-i+\sqrt{3})(i\sqrt{3}+7)(i\sqrt{3}+6)(i\sqrt{3}+5)(i\sqrt{3}+4)(i\sqrt{3}+3)} x^7 \right. \\
 & \left. + O(x^8) \right) \\
 & + c_3 \left(1 - x + \frac{1}{3} x^2 - \frac{1}{21} x^3 + \frac{1}{273} x^4 - \frac{1}{5733} x^5 + \frac{1}{177723} x^6 - \frac{1}{7642089} x^7 + O(x^8) \right)
 \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 3447

```
AsymptoticDSolveValue[x^3*y'''[x]+2*x^2*y''[x]+(x+x^2)*y'[x]+x*y[x]==0,y[x],{x,0,7}]
```

Too large to display

22.18 problem 3(b)

Internal problem ID [6493]

Internal file name [OUTPUT/5741_Sunday_June_05_2022_03_52_10_PM_60465236/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Problems for review and discover. (A) Drill Exercises . Page 194

Problem number: 3(b).

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**unknown**"

Maple gives the following as the ode type

```
[[_3rd_order , _with_linear_symmetries]]
```

Unable to solve or complete the solution.

Unable to parse ODE.

Maple trace

```
`Methods for third order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying high order exact linear fully integrable
trying to convert to a linear ODE with constant coefficients
trying differential order: 3; missing the dependent variable
trying Louvillian solutions for 3rd order ODEs, imprimitive case
-> pFq: Equivalence to the 3F2 or one of its 3 confluent cases under a power @ Moebius
<- pFq successful: received ODE is equivalent to the 0F2 ODE, case c = 0 `
```

✓ Solution by Maple

Time used: 0.11 (sec). Leaf size: 12916

```
Order:=8;  
dsolve(x^3*diff(y(x),x$3)+x^2*diff(y(x),x$2)-3*x*diff(y(x),x)+(x-1)*y(x)=0,y(x),type='series
```

Expression too large to display

✓ Solution by Mathematica

Time used: 0.029 (sec). Leaf size: 11815

```
AsymptoticDSolveValue[x^3*y'''[x]+x^2*y''[x]-3*x*y'[x]+(x-1)*y[x]==0,y[x],{x,0,7}]
```

Too large to display

22.19 problem 3(c)

Internal problem ID [6494]

Internal file name [OUTPUT/5742_Sunday_June_05_2022_03_52_23_PM_43449969/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Problems for review and discover. (A) Drill Exercises . Page 194

Problem number: 3(c).

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**unknown**"

Maple gives the following as the ode type

```
[[_3rd_order , _with_linear_symmetries]]
```

Unable to solve or complete the solution.

Unable to parse ODE.

Maple trace

```
`Methods for third order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying high order exact linear fully integrable
trying to convert to a linear ODE with constant coefficients
trying differential order: 3; missing the dependent variable
Equation is the LCLM of  $-1/(x+2)*y(x)+diff(y(x),x)$ ,  $1/x*y(x)-2/x*diff(y(x),x)+diff(diff(y(x)$ 
trying differential order: 1; missing the dependent variable
checking if the LODE is of Euler type
<- LODE of Euler type successful
Euler equation successful
  trying a quadrature
  checking if the LODE has constant coefficients
  checking if the LODE is of Euler type
  trying a symmetry of the form [xi=0, eta=F(x)]
  checking if the LODE is missing y
  -> Trying a Liouvillian solution using Kovacic's algorithm
  <- No Liouvillian solutions exists
  -> Trying a solution in terms of special functions:
    -> Bessel
    <- Bessel successful
  <- special function solution successful
<- solving the LCLM ode successful `
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 120

Order:=8;

dsolve(x^3*diff(y(x),x\$3)-2*x^2*diff(y(x),x\$2)+(x^2+2*x)*diff(y(x),x)-x*y(x)=0,y(x),type='se

$$\begin{aligned}
 y(x) = & c_1 x^3 \left(1 - \frac{1}{4}x + \frac{1}{40}x^2 - \frac{1}{720}x^3 + \frac{1}{20160}x^4 - \frac{1}{806400}x^5 + \frac{1}{43545600}x^6 \right. \\
 & \left. - \frac{1}{3048192000}x^7 + O(x^8) \right) + c_2 x^2 \left(\ln(x) \left((-240)x + 60x^2 - 6x^3 + \frac{1}{3}x^4 \right. \right. \\
 & \left. \left. - \frac{1}{84}x^5 + \frac{1}{3360}x^6 - \frac{1}{181440}x^7 + O(x^8) \right) + \left(720 - 908x + 152x^2 - 11x^3 + \frac{4}{9}x^4 \right. \right. \\
 & \left. \left. - \frac{79}{7056}x^5 + \frac{517}{2822400}x^6 - \frac{851}{457228800}x^7 + O(x^8) \right) \right) \\
 & + c_3 \left(2 \ln(x) \left(x^3 - \frac{1}{4}x^4 + \frac{1}{40}x^5 - \frac{1}{720}x^6 + \frac{1}{20160}x^7 + O(x^8) \right) \right. \\
 & \left. + \left(-24 - 12x - 6x^2 + \frac{5}{8}x^4 - \frac{39}{400}x^5 + \frac{49}{7200}x^6 - \frac{199}{705600}x^7 + O(x^8) \right) \right)
 \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.664 (sec). Leaf size: 186

AsymptoticDSolveValue[x^3*y'''[x]-2*x^2*y''[x]+(x^2+2*x)*y'[x]-x*y[x]==0,y[x],{x,0,7}]

$$\begin{aligned}
 y(x) \rightarrow & c_1 \left(\frac{(x^3 - 18x^2 + 180x - 720)x^3 \log(x)}{4320} \right. \\
 & \left. + \frac{-167x^6 + 2466x^5 - 17100x^4 + 14400x^3 + 129600x^2 + 259200x + 518400}{259200} \right) \\
 & + c_2 \left(\frac{x^3(x^5 - 40x^4 + 1120x^3 - 20160x^2 + 201600x - 806400) \log(x)}{2419200} \right. \\
 & \left. - \frac{x^2(2941x^6 - 106720x^5 + 2618560x^4 - 38666880x^3 + 268128000x^2 - 225792000x - 2032128000)}{2032128000} \right) \\
 & + c_3 \left(\frac{x^9}{43545600} - \frac{x^8}{806400} + \frac{x^7}{20160} - \frac{x^6}{720} + \frac{x^5}{40} - \frac{x^4}{4} + x^3 \right)
 \end{aligned}$$

22.20 problem 3(d)

Internal problem ID [6495]

Internal file name [OUTPUT/5743_Sunday_June_05_2022_03_52_25_PM_11483593/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Problems for review and discover. (A) Drill Exercises . Page 194

Problem number: 3(d).

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**unknown**"

Maple gives the following as the ode type

```
[[_3rd_order , _with_linear_symmetries]]
```

Unable to solve or complete the solution.

Unable to parse ODE.

Maple trace

```
`Methods for third order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying high order exact linear fully integrable
trying to convert to a linear ODE with constant coefficients
trying differential order: 3; missing the dependent variable
Equation is the LCLM of  $-1/x*y(x)+diff(y(x),x)$ ,  $-(2*x+1)/x^2*y(x)+2*(x-1)/x*diff(y(x),x)+diff^2(y(x),x)$ 
trying differential order: 1; missing the dependent variable
checking if the LODE is of Euler type
<- LODE of Euler type successful
Euler equation successful
  trying a quadrature
  checking if the LODE has constant coefficients
  checking if the LODE is of Euler type
  trying a symmetry of the form [xi=0, eta=F(x)]
  checking if the LODE is missing y
  -> Trying a Liouvillian solution using Kovacic's algorithm
  <- No Liouvillian solutions exist
  -> Trying a solution in terms of special functions:
    -> Bessel
    <- Bessel successful
  <- special function solution successful
<- solving the LCLM ode successful `
```


✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 1506

Order:=8;

dsolve(x^3*dif(y(x),x\$3)+(2*x^3-x^2)*dif(y(x),x\$2)-x*dif(y(x),x)+y(x)=0,y(x),type='series

$$\begin{aligned}
 y(x) = & c_3 x(1 + O(x^8)) + c_2 x^{\frac{3}{2} - \frac{\sqrt{13}}{2}} \left(1 - x + \frac{-3 + \sqrt{13}}{-4 + 2\sqrt{13}} x^2 + \frac{5 - \sqrt{13}}{6\sqrt{13} - 12} x^3 \right. \\
 & + \frac{1}{24} \frac{(-5 + \sqrt{13})(-7 + \sqrt{13})}{(-2 + \sqrt{13})(-4 + \sqrt{13})} x^4 + \frac{1}{30} \frac{-19 + 4\sqrt{13}}{(-2 + \sqrt{13})(-4 + \sqrt{13})} x^5 \\
 & + \frac{1}{20} \frac{-29 + 7\sqrt{13}}{(-2 + \sqrt{13})(-4 + \sqrt{13})(-6 + \sqrt{13})} x^6 \\
 & \left. + \frac{-\frac{117}{35} + \frac{6\sqrt{13}}{7}}{(-2 + \sqrt{13})(-4 + \sqrt{13})(-6 + \sqrt{13})(-7 + \sqrt{13})} x^7 + O(x^8) \right) \\
 & + c_1 x^{\frac{3}{2} + \frac{\sqrt{13}}{2}} \left(1 - x + \frac{3 + \sqrt{13}}{4 + 2\sqrt{13}} x^2 + \frac{-5 - \sqrt{13}}{6\sqrt{13} + 12} x^3 + \frac{1}{24} \frac{(5 + \sqrt{13})(7 + \sqrt{13})}{(2 + \sqrt{13})(4 + \sqrt{13})} x^4 \right. \\
 & - \frac{1}{30} \frac{19 + 4\sqrt{13}}{(2 + \sqrt{13})(4 + \sqrt{13})} x^5 + \frac{1}{20} \frac{29 + 7\sqrt{13}}{(2 + \sqrt{13})(4 + \sqrt{13})(6 + \sqrt{13})} x^6 \\
 & \left. + \frac{-\frac{117}{35} - \frac{6\sqrt{13}}{7}}{(2 + \sqrt{13})(4 + \sqrt{13})(6 + \sqrt{13})(7 + \sqrt{13})} x^7 + O(x^8) \right)
 \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.24 (sec). Leaf size: 310

AsymptoticDSolveValue[x^3*y'''[x]+(2*x^3-x^2)*y''[x]-y'[x]+y[x]==0,y[x],{x,0,7}]

$$\begin{aligned}
 y(x) \rightarrow & c_1 \left(\frac{99473x^7}{1008} + \frac{1043x^6}{144} + \frac{19x^5}{24} + \frac{11x^4}{24} - \frac{x^3}{6} + \frac{x^2}{2} + x + 1 \right) \\
 & + c_2 e^{-\frac{2}{\sqrt{x}}} \left(-\frac{279112936065458899252220570230691x^{13/2}}{160251477454333302276096} \right. \\
 & \quad - \frac{2430057902534044595693470483x^{11/2}}{100317681699677798400} - \frac{1545013796231079344731x^{9/2}}{3562417673994240} \\
 & \quad - \frac{2005991558758787x^{7/2}}{193273528320} - \frac{43999069453x^{5/2}}{125829120} - \frac{438565x^{3/2}}{24576} \\
 & \quad + \frac{14436319972596450047835320516938615783x^7}{897408273744266492746137600} \\
 & \quad + \frac{3840864007433053956366665361751x^6}{19260994886338137292800} + \frac{1786308115320202497636167x^5}{569986827839078400} \\
 & \quad + \frac{319234145332261451x^4}{4947802324992} + \frac{21959100963217x^3}{12079595520} + \frac{117706529x^2}{1572864} + \frac{2353x}{512} - \frac{29\sqrt{x}}{16} \\
 & \left. + 1 \right) x^{11/4} + c_3 e^{\frac{2}{\sqrt{x}}} \left(\frac{279112936065458899252220570230691x^{13/2}}{160251477454333302276096} + \frac{2430057902534044595693470483x^{11/2}}{100317681699677798400} \right)
 \end{aligned}$$

23 Chapter 4. Power Series Solutions and Special Functions. Problems for review and discover.

(B) Challenge Problems . Page 194

23.1 problem 1(a)	3807
23.2 problem 1(b)	3819
23.3 problem 1(c)	3837

23.1 problem 1(a)

Internal problem ID [6496]

Internal file name [OUTPUT/5744_Sunday_June_05_2022_03_52_28_PM_85151222/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Problems for review and discover. (B) Challenge Problems . Page 194

Problem number: 1(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Repeated root"**

Maple gives the following as the ode type

```
[[_Emden , _Fowler]]
```

$$x^3 y'' + x^2 y' + y = 0$$

With the expansion point for the power series method at $x = \infty$.

Since expansion is around ∞ , then the independent variable x is replaced by $\frac{1}{t}$ and the expansion is made around $t = 0$ and after solving, the solution is changed back to x using $x = \frac{1}{t}$. Changing variables results in the new ode

$$-\frac{\left(\frac{d^2}{dt^2}y(t)\right)t^2 - 2t\left(\frac{d}{dt}y(t)\right)}{t} - \frac{d}{dt}y(t) + y(t) = 0$$

The transformed ODE is now solved. The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$\left(\frac{d^2}{dt^2}y(t)\right)t + \frac{d}{dt}y(t) + y(t) = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$\frac{d^2}{dt^2}y(t) + p(t)\frac{d}{dt}y(t) + q(t)y(t) = 0$$

Where

$$p(t) = \frac{1}{t}$$

$$q(t) = \frac{1}{t}$$

Table 516: Table $p(t), q(t)$ singularities.

$p(t) = \frac{1}{t}$	
singularity	type
$t = 0$	“regular”

$q(t) = \frac{1}{t}$	
singularity	type
$t = 0$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0]$

Irregular singular points : $[\infty]$

Since $t = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$\left(\frac{d^2}{dt^2} y(t) \right) t + \frac{d}{dt} y(t) + y(t) = 0$$

Let the solution be represented as Frobenius power series of the form

$$y(t) = \sum_{n=0}^{\infty} a_n t^{n+r}$$

Then

$$\frac{d}{dt} y(t) = \sum_{n=0}^{\infty} (n+r) a_n t^{n+r-1}$$

$$\frac{d^2}{dt^2} y(t) = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n t^{n+r-2}$$

Substituting the above back into the ode gives

$$\left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n t^{n+r-2} \right) t + \left(\sum_{n=0}^{\infty} (n+r) a_n t^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} a_n t^{n+r} \right) = 0 \quad (1)$$

Which simplifies to

$$\left(\sum_{n=0}^{\infty} t^{n+r-1} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} (n+r) a_n t^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} a_n t^{n+r} \right) = 0 \quad (2A)$$

The next step is to make all powers of t be $n+r-1$ in each summation term. Going over each summation term above with power of t in it which is not already t^{n+r-1} and adjusting the power and the corresponding index gives

$$\sum_{n=0}^{\infty} a_n t^{n+r} = \sum_{n=1}^{\infty} a_{n-1} t^{n+r-1}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of t are the same and equal to $n+r-1$.

$$\left(\sum_{n=0}^{\infty} t^{n+r-1} a_n (n+r) (n+r-1) \right) + \left(\sum_{n=0}^{\infty} (n+r) a_n t^{n+r-1} \right) + \left(\sum_{n=1}^{\infty} a_{n-1} t^{n+r-1} \right) = 0 \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$t^{n+r-1} a_n (n+r) (n+r-1) + (n+r) a_n t^{n+r-1} = 0$$

When $n=0$ the above becomes

$$t^{-1+r} a_0 r (-1+r) + r a_0 t^{-1+r} = 0$$

Or

$$(t^{-1+r} r (-1+r) + r t^{-1+r}) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$t^{-1+r} r^2 = 0$$

Since the above is true for all t then the indicial equation becomes

$$r^2 = 0$$

Solving for r gives the roots of the indicial equation as

$$r_1 = 0$$

$$r_2 = 0$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$t^{-1+r}r^2 = 0$$

Solving for r gives the roots of the indicial equation as Since the root of the indicial equation is repeated, then we can construct two linearly independent solutions. The first solution has the form

$$y_1(t) = \sum_{n=0}^{\infty} a_n t^{n+r} \quad (1A)$$

Now the second solution y_2 is found using

$$y_2(t) = y_1(t) \ln(t) + \left(\sum_{n=1}^{\infty} b_n t^{n+r} \right) \quad (1B)$$

Then the general solution will be

$$y(t) = c_1 y_1(t) + c_2 y_2(t)$$

In Eq (1B) the sum starts from 1 and not zero. In Eq (1A), a_0 is never zero, and is arbitrary and is typically taken as $a_0 = 1$, and $\{c_1, c_2\}$ are two arbitray constants of integration which can be found from initial conditions. We start by finding the first solution $y_1(t)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. For $1 \leq n$ the recursive equation is

$$a_n(n+r)(n+r-1) + a_n(n+r) + a_{n-1} = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = -\frac{a_{n-1}}{n^2 + 2nr + r^2} \quad (4)$$

Which for the root $r = 0$ becomes

$$a_n = -\frac{a_{n-1}}{n^2} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1

For $n = 1$, using the above recursive equation gives

$$a_1 = -\frac{1}{(r+1)^2}$$

Which for the root $r = 0$ becomes

$$a_1 = -1$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{(r+1)^2}$	-1

For $n = 2$, using the above recursive equation gives

$$a_2 = \frac{1}{(r+1)^2(r+2)^2}$$

Which for the root $r = 0$ becomes

$$a_2 = \frac{1}{4}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{(r+1)^2}$	-1
a_2	$\frac{1}{(r+1)^2(r+2)^2}$	$\frac{1}{4}$

For $n = 3$, using the above recursive equation gives

$$a_3 = -\frac{1}{(r+1)^2(r+2)^2(r+3)^2}$$

Which for the root $r = 0$ becomes

$$a_3 = -\frac{1}{36}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{(r+1)^2}$	-1
a_2	$\frac{1}{(r+1)^2(r+2)^2}$	$\frac{1}{4}$
a_3	$-\frac{1}{(r+1)^2(r+2)^2(r+3)^2}$	$-\frac{1}{36}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{1}{(r+1)^2(r+2)^2(r+3)^2(r+4)^2}$$

Which for the root $r = 0$ becomes

$$a_4 = \frac{1}{576}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{(r+1)^2}$	-1
a_2	$\frac{1}{(r+1)^2(r+2)^2}$	$\frac{1}{4}$
a_3	$-\frac{1}{(r+1)^2(r+2)^2(r+3)^2}$	$-\frac{1}{36}$
a_4	$\frac{1}{(r+1)^2(r+2)^2(r+3)^2(r+4)^2}$	$\frac{1}{576}$

For $n = 5$, using the above recursive equation gives

$$a_5 = -\frac{1}{(r+1)^2(r+2)^2(r+3)^2(r+4)^2(r+5)^2}$$

Which for the root $r = 0$ becomes

$$a_5 = -\frac{1}{14400}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{(r+1)^2}$	-1
a_2	$\frac{1}{(r+1)^2(r+2)^2}$	$\frac{1}{4}$
a_3	$-\frac{1}{(r+1)^2(r+2)^2(r+3)^2}$	$-\frac{1}{36}$
a_4	$\frac{1}{(r+1)^2(r+2)^2(r+3)^2(r+4)^2}$	$\frac{1}{576}$
a_5	$-\frac{1}{(r+1)^2(r+2)^2(r+3)^2(r+4)^2(r+5)^2}$	$-\frac{1}{14400}$

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{1}{(r+1)^2(r+2)^2(r+3)^2(r+4)^2(r+5)^2(r+6)^2}$$

Which for the root $r = 0$ becomes

$$a_6 = \frac{1}{518400}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{(r+1)^2}$	-1
a_2	$\frac{1}{(r+1)^2(r+2)^2}$	$\frac{1}{4}$
a_3	$-\frac{1}{(r+1)^2(r+2)^2(r+3)^2}$	$-\frac{1}{36}$
a_4	$\frac{1}{(r+1)^2(r+2)^2(r+3)^2(r+4)^2}$	$\frac{1}{576}$
a_5	$-\frac{1}{(r+1)^2(r+2)^2(r+3)^2(r+4)^2(r+5)^2}$	$-\frac{1}{14400}$
a_6	$\frac{1}{(r+1)^2(r+2)^2(r+3)^2(r+4)^2(r+5)^2(r+6)^2}$	$\frac{1}{518400}$

For $n = 7$, using the above recursive equation gives

$$a_7 = -\frac{1}{(r+1)^2(r+2)^2(r+3)^2(r+4)^2(r+5)^2(r+6)^2(r+7)^2}$$

Which for the root $r = 0$ becomes

$$a_7 = -\frac{1}{25401600}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$-\frac{1}{(r+1)^2}$	-1
a_2	$\frac{1}{(r+1)^2(r+2)^2}$	$\frac{1}{4}$
a_3	$-\frac{1}{(r+1)^2(r+2)^2(r+3)^2}$	$-\frac{1}{36}$
a_4	$\frac{1}{(r+1)^2(r+2)^2(r+3)^2(r+4)^2}$	$\frac{1}{576}$
a_5	$-\frac{1}{(r+1)^2(r+2)^2(r+3)^2(r+4)^2(r+5)^2}$	$-\frac{1}{14400}$
a_6	$\frac{1}{(r+1)^2(r+2)^2(r+3)^2(r+4)^2(r+5)^2(r+6)^2}$	$\frac{1}{518400}$
a_7	$-\frac{1}{(r+1)^2(r+2)^2(r+3)^2(r+4)^2(r+5)^2(r+6)^2(r+7)^2}$	$-\frac{1}{25401600}$

Using the above table, then the first solution $y_1(t)$ becomes

$$\begin{aligned}
 y_1(t) &= a_0 + a_1t + a_2t^2 + a_3t^3 + a_4t^4 + a_5t^5 + a_6t^6 + a_7t^7 + a_8t^8 \dots \\
 &= -t + 1 + \frac{t^2}{4} - \frac{t^3}{36} + \frac{t^4}{576} - \frac{t^5}{14400} + \frac{t^6}{518400} - \frac{t^7}{25401600} + O(t^8)
 \end{aligned}$$

Now the second solution is found. The second solution is given by

$$y_2(t) = y_1(t) \ln(t) + \left(\sum_{n=1}^{\infty} b_n t^{n+r} \right)$$

Where b_n is found using

$$b_n = \frac{d}{dr} a_{n,r}$$

And the above is then evaluated at $r = 0$. The above table for $a_{n,r}$ is used for this purpose. Computing the derivatives gives the following table

n	$b_{n,r}$	a_n	$b_{n,r} = \frac{d}{dr} a_{n,r}$	b_n
b_0	1	1	N/A since b_n starts from 1	N/A
b_1	$-\frac{1}{(r+1)^2}$	-1	$\frac{2}{(r+1)^3}$	2
b_2	$\frac{1}{(r+1)^2(r+2)^2}$	$\frac{1}{4}$	$\frac{-4r-6}{(r+1)^3(r+2)^3}$	$-\frac{3}{4}$
b_3	$-\frac{1}{(r+1)^2(r+2)^2(r+3)^2}$	$-\frac{1}{36}$	$\frac{6r^2+24r+22}{(r+1)^3(r+2)^3(r+3)^3}$	$\frac{11}{108}$
b_4	$\frac{1}{(r+1)^2(r+2)^2(r+3)^2(r+4)^2}$	$\frac{1}{576}$	$\frac{-8r^3-60r^2-140r-100}{(r+1)^3(r+2)^3(r+3)^3(r+4)^3}$	$-\frac{3}{432}$
b_5	$-\frac{1}{(r+1)^2(r+2)^2(r+3)^2(r+4)^2(r+5)^2}$	$-\frac{1}{14400}$	$\frac{10r^4+120r^3+510r^2+900r+548}{(r+1)^3(r+2)^3(r+3)^3(r+4)^3(r+5)^3}$	$\frac{11}{4320}$
b_6	$\frac{1}{(r+1)^2(r+2)^2(r+3)^2(r+4)^2(r+5)^2(r+6)^2}$	$\frac{1}{518400}$	$\frac{-12r^5-210r^4-1400r^3-4410r^2-6496r-3528}{(r+1)^3(r+2)^3(r+3)^3(r+4)^3(r+5)^3(r+6)^3}$	$-\frac{1}{518400}$
b_7	$-\frac{1}{(r+1)^2(r+2)^2(r+3)^2(r+4)^2(r+5)^2(r+6)^2(r+7)^2}$	$-\frac{1}{25401600}$	$\frac{14r^6+336r^5+3220r^4+15680r^3+40614r^2+52528r+26136}{(r+1)^3(r+2)^3(r+3)^3(r+4)^3(r+5)^3(r+6)^3(r+7)^3}$	$\frac{1}{25401600}$

The above table gives all values of b_n needed. Hence the second solution is

$$\begin{aligned}
y_2(t) &= y_1(t) \ln(t) + b_0 + b_1 t + b_2 t^2 + b_3 t^3 + b_4 t^4 + b_5 t^5 + b_6 t^6 + b_7 t^7 + b_8 t^8 \dots \\
&= \left(-t + 1 + \frac{t^2}{4} - \frac{t^3}{36} + \frac{t^4}{576} - \frac{t^5}{14400} + \frac{t^6}{518400} - \frac{t^7}{25401600} + O(t^8) \right) \ln(t) \\
&\quad + 2t - \frac{3t^2}{4} + \frac{11t^3}{108} - \frac{25t^4}{3456} + \frac{137t^5}{432000} - \frac{49t^6}{5184000} + \frac{121t^7}{592704000} + O(t^8)
\end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
y_h(t) &= c_1 y_1(t) + c_2 y_2(t) \\
&= c_1 \left(-t + 1 + \frac{t^2}{4} - \frac{t^3}{36} + \frac{t^4}{576} - \frac{t^5}{14400} + \frac{t^6}{518400} - \frac{t^7}{25401600} + O(t^8) \right) \\
&\quad + c_2 \left(\left(-t + 1 + \frac{t^2}{4} - \frac{t^3}{36} + \frac{t^4}{576} - \frac{t^5}{14400} + \frac{t^6}{518400} - \frac{t^7}{25401600} + O(t^8) \right) \ln(t) \right. \\
&\quad \left. + 2t - \frac{3t^2}{4} + \frac{11t^3}{108} - \frac{25t^4}{3456} + \frac{137t^5}{432000} - \frac{49t^6}{5184000} + \frac{121t^7}{592704000} + O(t^8) \right)
\end{aligned}$$

Hence the final solution is

$$y(t) = y_h$$

$$\begin{aligned}
&= c_1 \left(-t + 1 + \frac{t^2}{4} - \frac{t^3}{36} + \frac{t^4}{576} - \frac{t^5}{14400} + \frac{t^6}{518400} - \frac{t^7}{25401600} + O(t^8) \right) \\
&\quad + c_2 \left(\left(-t + 1 + \frac{t^2}{4} - \frac{t^3}{36} + \frac{t^4}{576} - \frac{t^5}{14400} + \frac{t^6}{518400} - \frac{t^7}{25401600} + O(t^8) \right) \ln(t) \right. \\
&\quad \left. + 2t - \frac{3t^2}{4} + \frac{11t^3}{108} - \frac{25t^4}{3456} + \frac{137t^5}{432000} - \frac{49t^6}{5184000} + \frac{121t^7}{592704000} + O(t^8) \right)
\end{aligned}$$

Replacing t by $\frac{1}{x}$ gives

$$y = c_1 \left(-\frac{1}{x} + 1 + \frac{1}{4x^2} - \frac{1}{36x^3} + \frac{1}{576x^4} - \frac{1}{14400x^5} + \frac{1}{518400x^6} - \frac{1}{25401600x^7} + O\left(\frac{1}{x^8}\right) \right) + c_2 \left(\left(-\frac{1}{x} + 1 + \frac{1}{4x^2} - \frac{1}{36x^3} + \frac{1}{576x^4} - \frac{1}{14400x^5} + \frac{1}{518400x^6} - \frac{1}{25401600x^7} + O\left(\frac{1}{x^8}\right) \right) \ln\left(\frac{1}{x}\right) + \frac{2}{x} - \frac{3}{4x^2} + \frac{11}{108x^3} - \frac{25}{3456x^4} + \frac{137}{432000x^5} - \frac{49}{5184000x^6} + \frac{121}{592704000x^7} + O\left(\frac{1}{x^8}\right) \right)$$

Summary

The solution(s) found are the following

$$\begin{aligned}
y = c_1 &\left(-\frac{1}{x} + 1 + \frac{1}{4x^2} - \frac{1}{36x^3} + \frac{1}{576x^4} - \frac{1}{14400x^5} + \frac{1}{518400x^6} - \frac{1}{25401600x^7} \right. \\
&\quad \left. + O\left(\frac{1}{x^8}\right) \right) + c_2 \left(\left(-\frac{1}{x} + 1 + \frac{1}{4x^2} - \frac{1}{36x^3} + \frac{1}{576x^4} - \frac{1}{14400x^5} + \frac{1}{518400x^6} - \frac{1}{25401600x^7} \right. \right. \\
&\quad \left. \left. + O\left(\frac{1}{x^8}\right) \right) \ln\left(\frac{1}{x}\right) + \frac{2}{x} - \frac{3}{4x^2} + \frac{11}{108x^3} - \frac{25}{3456x^4} + \frac{137}{432000x^5} \right. \\
&\quad \left. - \frac{49}{5184000x^6} + \frac{121}{592704000x^7} + O\left(\frac{1}{x^8}\right) \right)
\end{aligned}$$

Verification of solutions

$$\begin{aligned}
y = c_1 &\left(-\frac{1}{x} + 1 + \frac{1}{4x^2} - \frac{1}{36x^3} + \frac{1}{576x^4} - \frac{1}{14400x^5} + \frac{1}{518400x^6} - \frac{1}{25401600x^7} + O\left(\frac{1}{x^8}\right) \right) \\
&+ c_2 \left(\left(-\frac{1}{x} + 1 + \frac{1}{4x^2} - \frac{1}{36x^3} + \frac{1}{576x^4} - \frac{1}{14400x^5} + \frac{1}{518400x^6} - \frac{1}{25401600x^7} \right. \right. \\
&\quad \left. \left. + O\left(\frac{1}{x^8}\right) \right) \ln\left(\frac{1}{x}\right) + \frac{2}{x} - \frac{3}{4x^2} + \frac{11}{108x^3} - \frac{25}{3456x^4} + \frac{137}{432000x^5} - \frac{49}{5184000x^6} \right. \\
&\quad \left. + \frac{121}{592704000x^7} + O\left(\frac{1}{x^8}\right) \right)
\end{aligned}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
checking if the LODE is missing y  
-> Trying a Liouvillian solution using Kovacic's algorithm  
<- No Liouvillian solutions exist  
-> Trying a solution in terms of special functions:  
  -> Bessel  
  <- Bessel successful  
<- special function solution successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 207

Order:=8;

dsolve(x^3*diff(y(x),x\$2)+x^2*diff(y(x),x)+y(x)=0,y(x),type='series',x=infinity);

$$\begin{aligned}
 y(x) = & \left(1 - \frac{(x - \text{Infinity})^2}{2 \text{Infinity}^3} + \frac{2(x - \text{Infinity})^3}{3 \text{Infinity}^4} + \frac{(-18 \text{Infinity} + 1)(x - \text{Infinity})^4}{24 \text{Infinity}^6} \right. \\
 & + \frac{(96 \text{Infinity} - 14)(x - \text{Infinity})^5}{120 \text{Infinity}^7} + \frac{(-600 \text{Infinity}^2 + 156 \text{Infinity} - 1)(x - \text{Infinity})^6}{720 \text{Infinity}^9} \\
 & \left. + \frac{(4320 \text{Infinity}^2 - 1692 \text{Infinity} + 30)(x - \text{Infinity})^7}{5040 \text{Infinity}^{10}} \right) y(\text{Infinity}) \\
 & + \left(x - \text{Infinity} - \frac{(x - \text{Infinity})^2}{2 \text{Infinity}} + \frac{(2 \text{Infinity}^2 - \text{Infinity})(x - \text{Infinity})^3}{6 \text{Infinity}^4} \right. \\
 & \quad - \frac{(\text{Infinity} - \frac{4}{3})(x - \text{Infinity})^4}{4 \text{Infinity}^4} \\
 & \quad + \frac{(24 \text{Infinity}^3 - 58 \text{Infinity}^2 + \text{Infinity})(x - \text{Infinity})^5}{120 \text{Infinity}^7} \\
 & \quad \left. + \frac{(-120 \text{Infinity}^4 + 444 \text{Infinity}^3 - 21 \text{Infinity}^2)(x - \text{Infinity})^6}{720 \text{Infinity}^9} \right. \\
 & \left. + \frac{(720 \text{Infinity}^4 - 3708 \text{Infinity}^3 + 324 \text{Infinity}^2 - \text{Infinity})(x - \text{Infinity})^7}{5040 \text{Infinity}^{10}} \right) D(y)(\text{Infinity}) \\
 & + O(x^8)
 \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 171

AsymptoticDSolveValue[x^3*y''[x]+x^2*y'[x]+y[x]==0,y[x],{x,infinity,7}]

$$\begin{aligned}
 y(x) \rightarrow & c_1 \left(-\frac{1}{25401600x^7} + \frac{1}{518400x^6} - \frac{1}{14400x^5} + \frac{1}{576x^4} - \frac{1}{36x^3} + \frac{1}{4x^2} - \frac{1}{x} + 1 \right) \\
 & + c_2 \left(\frac{121}{592704000x^7} + \frac{\log(x)}{25401600x^7} - \frac{49}{5184000x^6} - \frac{\log(x)}{518400x^6} + \frac{137}{432000x^5} \right. \\
 & \quad + \frac{\log(x)}{14400x^5} - \frac{25}{3456x^4} - \frac{\log(x)}{576x^4} + \frac{11}{108x^3} + \frac{\log(x)}{36x^3} - \frac{3}{4x^2} - \frac{\log(x)}{4x^2} + \frac{2}{x} + \frac{\log(x)}{x} \\
 & \quad \left. - \log(x) \right)
 \end{aligned}$$

23.2 problem 1(b)

23.2.1 Maple step by step solution 3832

Internal problem ID [6497]

Internal file name [OUTPUT/5745_Sunday_June_05_2022_03_52_29_PM_25435504/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Problems for review and discover. (B) Challenge Problems . Page 194

Problem number: 1(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$9(-2+x)^2(x-3)y'' + 6x(-2+x)y' + 16y = 0$$

With the expansion point for the power series method at $x = \infty$.

Since expansion is around ∞ , then the independent variable x is replaced by $\frac{1}{t}$ and the expansion is made around $t = 0$ and after solving, the solution is changed back to x using $x = \frac{1}{t}$. Changing variables results in the new ode

$$-9\left(-2 + \frac{1}{t}\right)^2 \left(\frac{1}{t} - 3\right) \left(-\left(\frac{d^2}{dt^2}y(t)\right)t^2 - 2t\left(\frac{d}{dt}y(t)\right)\right)t^2 - \left(\frac{6}{t^2} - \frac{12}{t}\right) \left(\frac{d}{dt}y(t)\right)t^2 + 16y(t) = 0$$

The transformed ODE is now solved. The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$(-108t^4 + 144t^3 - 63t^2 + 9t) \left(\frac{d^2}{dt^2}y(t)\right) + (-216t^3 + 288t^2 - 114t + 12) \left(\frac{d}{dt}y(t)\right) + 16y(t) = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$\frac{d^2}{dt^2}y(t) + p(t)\frac{d}{dt}y(t) + q(t)y(t) = 0$$

Where

$$p(t) = \frac{12t^2 - 10t + \frac{4}{3}}{t(-1 + 3t)(2t - 1)}$$

$$q(t) = -\frac{16}{9(2t - 1)^2(-1 + 3t)t}$$

Table 517: Table $p(t), q(t)$ singularities.

$p(t) = \frac{12t^2 - 10t + \frac{4}{3}}{t(-1 + 3t)(2t - 1)}$		$q(t) = -\frac{16}{9(2t - 1)^2(-1 + 3t)t}$	
singularity	type	singularity	type
$t = 0$	“regular”	$t = 0$	“regular”
$t = \frac{1}{2}$	“regular”	$t = \frac{1}{2}$	“regular”
$t = \frac{1}{3}$	“regular”	$t = \frac{1}{3}$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[0, \frac{1}{2}, \frac{1}{3}, \infty]$

Irregular singular points : $[\]$

Since $t = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$-9t(12t^3 - 16t^2 + 7t - 1) \left(\frac{d^2}{dt^2} y(t) \right) + (-216t^3 + 288t^2 - 114t + 12) \left(\frac{d}{dt} y(t) \right) + 16y(t) = 0$$

Let the solution be represented as Frobenius power series of the form

$$y(t) = \sum_{n=0}^{\infty} a_n t^{n+r}$$

Then

$$\frac{d}{dt} y(t) = \sum_{n=0}^{\infty} (n+r) a_n t^{n+r-1}$$

$$\frac{d^2}{dt^2} y(t) = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n t^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned}
 & -9t(12t^3 - 16t^2 + 7t - 1) \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n t^{n+r-2} \right) \\
 & + (-216t^3 + 288t^2 - 114t + 12) \left(\sum_{n=0}^{\infty} (n+r) a_n t^{n+r-1} \right) + 16 \left(\sum_{n=0}^{\infty} a_n t^{n+r} \right) = 0
 \end{aligned} \tag{1}$$

Which simplifies to

$$\begin{aligned}
 & \sum_{n=0}^{\infty} (-108t^{n+r+2} a_n (n+r)(n+r-1)) \\
 & + \left(\sum_{n=0}^{\infty} 144t^{1+n+r} a_n (n+r)(n+r-1) \right) \\
 & + \sum_{n=0}^{\infty} (-63t^{n+r} a_n (n+r)(n+r-1)) \\
 & + \left(\sum_{n=0}^{\infty} 9t^{n+r-1} a_n (n+r)(n+r-1) \right) + \sum_{n=0}^{\infty} (-216t^{n+r+2} a_n (n+r)) \\
 & + \left(\sum_{n=0}^{\infty} 288t^{1+n+r} a_n (n+r) \right) + \sum_{n=0}^{\infty} (-114t^{n+r} a_n (n+r)) \\
 & + \left(\sum_{n=0}^{\infty} 12(n+r) a_n t^{n+r-1} \right) + \left(\sum_{n=0}^{\infty} 16a_n t^{n+r} \right) = 0
 \end{aligned} \tag{2A}$$

The next step is to make all powers of t be $n+r-1$ in each summation term. Going over each summation term above with power of t in it which is not already t^{n+r-1} and adjusting the power and the corresponding index gives

$$\begin{aligned}
 \sum_{n=0}^{\infty} (-108t^{n+r+2} a_n (n+r)(n+r-1)) &= \sum_{n=3}^{\infty} (-108a_{n-3} (n+r-3)(n-4+r) t^{n+r-1}) \\
 \sum_{n=0}^{\infty} 144t^{1+n+r} a_n (n+r)(n+r-1) &= \sum_{n=2}^{\infty} 144a_{n-2} (n+r-2)(n+r-3) t^{n+r-1} \\
 \sum_{n=0}^{\infty} (-63t^{n+r} a_n (n+r)(n+r-1)) &= \sum_{n=1}^{\infty} (-63a_{n-1} (n+r-1)(n+r-2) t^{n+r-1})
 \end{aligned}$$

$$\begin{aligned}
\sum_{n=0}^{\infty} (-216t^{n+r+2}a_n(n+r)) &= \sum_{n=3}^{\infty} (-216a_{n-3}(n+r-3)t^{n+r-1}) \\
\sum_{n=0}^{\infty} 288t^{1+n+r}a_n(n+r) &= \sum_{n=2}^{\infty} 288a_{n-2}(n+r-2)t^{n+r-1} \\
\sum_{n=0}^{\infty} (-114t^{n+r}a_n(n+r)) &= \sum_{n=1}^{\infty} (-114a_{n-1}(n+r-1)t^{n+r-1}) \\
\sum_{n=0}^{\infty} 16a_nt^{n+r} &= \sum_{n=1}^{\infty} 16a_{n-1}t^{n+r-1}
\end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of t are the same and equal to $n+r-1$.

$$\begin{aligned}
&\sum_{n=3}^{\infty} (-108a_{n-3}(n+r-3)(n-4+r)t^{n+r-1}) \\
&+ \left(\sum_{n=2}^{\infty} 144a_{n-2}(n+r-2)(n+r-3)t^{n+r-1} \right) \\
&+ \sum_{n=1}^{\infty} (-63a_{n-1}(n+r-1)(n+r-2)t^{n+r-1}) \\
&+ \left(\sum_{n=0}^{\infty} 9t^{n+r-1}a_n(n+r)(n+r-1) \right) \tag{2B} \\
&+ \sum_{n=3}^{\infty} (-216a_{n-3}(n+r-3)t^{n+r-1}) \\
&+ \left(\sum_{n=2}^{\infty} 288a_{n-2}(n+r-2)t^{n+r-1} \right) + \sum_{n=1}^{\infty} (-114a_{n-1}(n+r-1)t^{n+r-1}) \\
&+ \left(\sum_{n=0}^{\infty} 12(n+r)a_nt^{n+r-1} \right) + \left(\sum_{n=1}^{\infty} 16a_{n-1}t^{n+r-1} \right) = 0
\end{aligned}$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$9t^{n+r-1}a_n(n+r)(n+r-1) + 12(n+r)a_nt^{n+r-1} = 0$$

When $n=0$ the above becomes

$$9t^{-1+r}a_0r(-1+r) + 12ra_0t^{-1+r} = 0$$

Or

$$(9t^{-1+r}r(-1+r) + 12rt^{-1+r})a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(9r^2 + 3r)t^{-1+r} = 0$$

Since the above is true for all t then the indicial equation becomes

$$9r^2 + 3r = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= 0 \\ r_2 &= -\frac{1}{3} \end{aligned}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(9r^2 + 3r)t^{-1+r} = 0$$

Solving for r gives the roots of the indicial equation as Since $r_1 - r_2 = \frac{1}{3}$ is not an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(t) &= t^{r_1} \left(\sum_{n=0}^{\infty} a_n t^n \right) \\ y_2(t) &= t^{r_2} \left(\sum_{n=0}^{\infty} b_n t^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(t) &= \sum_{n=0}^{\infty} a_n t^n \\ y_2(t) &= \sum_{n=0}^{\infty} b_n t^{n-\frac{1}{3}} \end{aligned}$$

We start by finding $y_1(t)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = \frac{63r^2 + 51r - 16}{9r^2 + 21r + 12}$$

Substituting $n = 2$ in Eq. (2B) gives

$$a_2 = \frac{2673r^4 + 10044r^3 + 9441r^2 + 438r - 1568}{81r^4 + 540r^3 + 1305r^2 + 1350r + 504}$$

For $3 \leq n$ the recursive equation is

$$\begin{aligned} & -108a_{n-3}(n+r-3)(n-4+r) + 144a_{n-2}(n+r-2)(n+r-3) \\ & - 63a_{n-1}(n+r-1)(n+r-2) + 9a_n(n+r)(n+r-1) - 216a_{n-3}(n+r-3) \\ & + 288a_{n-2}(n+r-2) - 114a_{n-1}(n+r-1) + 12a_n(n+r) + 16a_{n-1} = 0 \end{aligned} \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{108n^2a_{n-3} - 144n^2a_{n-2} + 63n^2a_{n-1} + 216nra_{n-3} - 288nra_{n-2} + 126nra_{n-1} + 108r^2a_{n-3} - 144r^2a_{n-2} + 9n^2 + 16ra_{n-1}}{9n^2 + 3n} \quad (4)$$

Which for the root $r = 0$ becomes

$$a_n = \frac{(108a_{n-3} - 144a_{n-2} + 63a_{n-1})n^2 + (-540a_{n-3} + 432a_{n-2} - 75a_{n-1})n + 648a_{n-3} - 288a_{n-2} - 4a_{n-1}}{9n^2 + 3n} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = 0$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{63r^2+51r-16}{9r^2+21r+12}$	$-\frac{4}{3}$
a_2	$\frac{2673r^4+10044r^3+9441r^2+438r-1568}{81r^4+540r^3+1305r^2+1350r+504}$	$-\frac{28}{9}$

For $n = 3$, using the above recursive equation gives

$$a_3 = \frac{95499r^6 + 844911r^5 + 2671137r^4 + 3571317r^3 + 1572804r^2 - 419508r - 336448}{729r^6 + 9477r^5 + 49815r^4 + 135135r^3 + 198936r^2 + 150228r + 45360}$$

Which for the root $r = 0$ becomes

$$a_3 = -\frac{3004}{405}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{63r^2+51r-16}{9r^2+21r+12}$	$-\frac{4}{3}$
a_2	$\frac{2673r^4+10044r^3+9441r^2+438r-1568}{81r^4+540r^3+1305r^2+1350r+504}$	$-\frac{28}{9}$
a_3	$\frac{95499r^6+844911r^5+2671137r^4+3571317r^3+1572804r^2-419508r-336448}{729r^6+9477r^5+49815r^4+135135r^3+198936r^2+150228r+45360}$	$-\frac{3004}{405}$

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{3103353r^8 + 49898592r^7 + 320994738r^6 + 1054759968r^5 + 1851078825r^4 + 1583317152r^3 + 348879276r^2 - 293260704r - 127995392}{81(3r^2 + 25r + 52)(27r^6 + 351r^5 + 1845r^4 + 5005r^3 + 7368r^2 + 5564r + 1680)}$$

Which for the root $r = 0$ becomes

$$a_4 = -\frac{285704}{15795}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	$\frac{63r^2+51r-16}{9r^2+21r+12}$	$-\frac{4}{3}$
a_2	$\frac{2673r^4+10044r^3+9441r^2+438r-1568}{81r^4+540r^3+1305r^2+1350r+504}$	$-\frac{28}{9}$
a_3	$\frac{95499r^6+844911r^5+2671137r^4+3571317r^3+1572804r^2-419508r-336448}{729r^6+9477r^5+49815r^4+135135r^3+198936r^2+150228r+45360}$	$-\frac{3004}{405}$
a_4	$\frac{3103353r^8+49898592r^7+320994738r^6+1054759968r^5+1851078825r^4+1583317152r^3+348879276r^2-293260704r-127995392}{81(3r^2+25r+52)(27r^6+351r^5+1845r^4+5005r^3+7368r^2+5564r+1680)}$	$-\frac{285704}{15795}$

For $n = 5$, using the above recursive equation gives

$$a_5 = \frac{95127939r^{10} + 2420949951r^9 + 26064871578r^8 + 154759665150r^7 + 552318920667r^6 + 12044507640r^5 - 12044507640r^4 + 12044507640r^3 - 12044507640r^2 + 12044507640r - 12044507640}{243(3r^2 + 31r + 80)(3r^2 + 25r + 52)(27r^6 + 351r^5 + 1845r^4 + 5005r^3 + 7368r^2 + 5564r + 1680)}$$

Which for the root $r = 0$ becomes

$$a_5 = -\frac{822592}{18225}$$

And the table now becomes

n	$a_{n,r}$
a_0	1
a_1	$\frac{63r^2+51r-16}{9r^2+21r+12}$
a_2	$\frac{2673r^4+10044r^3+9441r^2+438r-1568}{81r^4+540r^3+1305r^2+1350r+504}$
a_3	$\frac{95499r^6+844911r^5+2671137r^4+3571317r^3+1572804r^2-419508r-336448}{729r^6+9477r^5+49815r^4+135135r^3+198936r^2+150228r+45360}$
a_4	$\frac{3103353r^8+49898592r^7+320994738r^6+1054759968r^5+1851078825r^4+1583317152r^3+348879276r^2-293260704r-127995392}{81(3r^2+25r+52)(27r^6+351r^5+1845r^4+5005r^3+7368r^2+5564r+1680)}$
a_5	$\frac{95127939r^{10}+2420949951r^9+26064871578r^8+154759665150r^7+552318920667r^6+1204450764687r^5+1536667458552r^4+964786860468r^3+243(3r^2+31r+80)(3r^2+25r+52)(27r^6+351r^5+1845r^4+5005r^3+7368r^2+5564r+1680)}{243(3r^2+31r+80)(3r^2+25r+52)(27r^6+351r^5+1845r^4+5005r^3+7368r^2+5564r+1680)}$

For $n = 6$, using the above recursive equation gives

$$a_6 = \frac{2806539921r^{12} + 103709648268r^{11} + 1676277992835r^{10} + 15583794209910r^9 + 92090095190235r^8 + 360297425137320r^7 + 940035462956685r^6 + 243(3r^2+31r+80)(3r^2+25r+52)(27r^6+351r^5+1845r^4+5005r^3+7368r^2+5564r+1680)}{729(3r^2 + 31r + 80)}$$

Which for the root $r = 0$ becomes

$$a_6 = -\frac{4666732192}{40514175}$$

And the table now becomes

n	$a_{n,r}$
a_0	1
a_1	$\frac{63r^2+51r-16}{9r^2+21r+12}$
a_2	$\frac{2673r^4+10044r^3+9441r^2+438r-1568}{81r^4+540r^3+1305r^2+1350r+504}$
a_3	$\frac{95499r^6+844911r^5+2671137r^4+3571317r^3+1572804r^2-419508r-336448}{729r^6+9477r^5+49815r^4+135135r^3+198936r^2+150228r+45360}$
a_4	$\frac{3103353r^8+49898592r^7+320994738r^6+1054759968r^5+1851078825r^4+1583317152r^3+348879276r^2-293260704r-127995392}{81(3r^2+25r+52)(27r^6+351r^5+1845r^4+5005r^3+7368r^2+5564r+1680)}$
a_5	$\frac{95127939r^{10}+2420949951r^9+26064871578r^8+154759665150r^7+552318920667r^6+1204450764687r^5+1536667458552r^4+964786860468r^3+243(3r^2+31r+80)(3r^2+25r+52)(27r^6+351r^5+1845r^4+5005r^3+7368r^2+5564r+1680)}{243(3r^2+31r+80)(3r^2+25r+52)(27r^6+351r^5+1845r^4+5005r^3+7368r^2+5564r+1680)}$
a_6	$\frac{2806539921r^{12}+103709648268r^{11}+1676277992835r^{10}+15583794209910r^9+92090095190235r^8+360297425137320r^7+940035462956685r^6+243(3r^2+31r+80)(3r^2+25r+52)(27r^6+351r^5+1845r^4+5005r^3+7368r^2+5564r+1680)}{729(3r^2+31r+80)(3r^2+25r+52)(27r^6+351r^5+1845r^4+5005r^3+7368r^2+5564r+1680)}$

For $n = 7$, using the above recursive equation gives

$$a_7 = \frac{80674338123r^{14} + 4080569276151r^{13} + 92307128293827r^{12} + 1233512636444739r^{11} + 10829721106r^{10} + 360297425137320r^9 + 940035462956685r^8 + 243(3r^2+31r+80)(3r^2+25r+52)(27r^6+351r^5+1845r^4+5005r^3+7368r^2+5564r+1680)}{729(3r^2 + 31r + 80)}$$

Which for the root $r = 0$ becomes

$$a_7 = -\frac{401483448544}{1336967775}$$

And the table now becomes

n	$a_{n,r}$
a_0	1
a_1	$\frac{63r^2+51r-16}{9r^2+21r+12}$
a_2	$\frac{2673r^4+10044r^3+9441r^2+438r-1568}{81r^4+540r^3+1305r^2+1350r+504}$
a_3	$\frac{95499r^6+844911r^5+2671137r^4+3571317r^3+1572804r^2-419508r-336448}{729r^6+9477r^5+49815r^4+135135r^3+198936r^2+150228r+45360}$
a_4	$\frac{3103353r^8+49898592r^7+320994738r^6+1054759968r^5+1851078825r^4+1583317152r^3+348879276r^2-293260704r-127995392}{81(3r^2+25r+52)(27r^6+351r^5+1845r^4+5005r^3+7368r^2+5564r+1680)}$
a_5	$\frac{95127939r^{10}+2420949951r^9+26064871578r^8+154759665150r^7+552318920667r^6+1204450764687r^5+1536667458552r^4+964786860468r^3+211111111111r^2+111111111111r+111111111111}{243(3r^2+31r+80)(3r^2+25r+52)(27r^6+351r^5+1845r^4+5005r^3+7368r^2+5564r+1680)}$
a_6	$\frac{2806539921r^{12}+103709648268r^{11}+1676277992835r^{10}+15583794209910r^9+92090095190235r^8+360297425137320r^7+940035462956685r^6+111111111111111r^5+111111111111111r^4+111111111111111r^3+111111111111111r^2+111111111111111r+111111111111111}{729(3r^2+31r+80)(3r^2+25r+52)(27r^6+351r^5+1845r^4+5005r^3+7368r^2+5564r+1680)}$
a_7	$\frac{80674338123r^{14}+4080569276151r^{13}+92307128293827r^{12}+1233512636444739r^{11}+10829721106831365r^{10}+65685074939730909r^9+28141111111111111r^8+11111111111111111r^7+11111111111111111r^6+11111111111111111r^5+11111111111111111r^4+11111111111111111r^3+11111111111111111r^2+11111111111111111r+11111111111111111}{2187(3r^2+31r+80)}$

Using the above table, then the solution $y_1(t)$ is

$$y_1(t) = a_0 + a_1t + a_2t^2 + a_3t^3 + a_4t^4 + a_5t^5 + a_6t^6 + a_7t^7 + a_8t^8 \dots$$

$$= 1 - \frac{4t}{3} - \frac{28t^2}{9} - \frac{3004t^3}{405} - \frac{285704t^4}{15795} - \frac{822592t^5}{18225} - \frac{4666732192t^6}{40514175} - \frac{401483448544t^7}{1336967775} + O(t^8)$$

Now the second solution $y_2(t)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$b_1 = \frac{63r^2 + 51r - 16}{9r^2 + 21r + 12}$$

Substituting $n = 2$ in Eq. (2B) gives

$$b_2 = \frac{2673r^4 + 10044r^3 + 9441r^2 + 438r - 1568}{81r^4 + 540r^3 + 1305r^2 + 1350r + 504}$$

For $3 \leq n$ the recursive equation is

$$\begin{aligned} & -108b_{n-3}(n+r-3)(n-4+r) + 144b_{n-2}(n+r-2)(n+r-3) \\ & - 63b_{n-1}(n+r-1)(n+r-2) + 9b_n(n+r)(n+r-1) - 216b_{n-3}(n+r-3) \\ & + 288b_{n-2}(n+r-2) - 114b_{n-1}(n+r-1) + 12(n+r)b_n + 16b_{n-1} = 0 \end{aligned} \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = \frac{108n^2b_{n-3} - 144n^2b_{n-2} + 63n^2b_{n-1} + 216nrb_{n-3} - 288nrb_{n-2} + 126nrb_{n-1} + 108r^2b_{n-3} - 144r^2b_{n-2} + 9n^2 + 18r}{9n^2 + 18r} \quad (4)$$

Which for the root $r = -\frac{1}{3}$ becomes

$$b_n = \frac{(108b_{n-3} - 144b_{n-2} + 63b_{n-1})n^2 + (-612b_{n-3} + 528b_{n-2} - 117b_{n-1})n + 840b_{n-3} - 448b_{n-2} + 28b_{n-1}}{9n^2 - 3n} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = -\frac{1}{3}$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{63r^2+51r-16}{9r^2+21r+12}$	$-\frac{13}{3}$
b_2	$\frac{2673r^4+10044r^3+9441r^2+438r-1568}{81r^4+540r^3+1305r^2+1350r+504}$	$-\frac{251}{45}$

For $n = 3$, using the above recursive equation gives

$$b_3 = \frac{95499r^6 + 844911r^5 + 2671137r^4 + 3571317r^3 + 1572804r^2 - 419508r - 336448}{729r^6 + 9477r^5 + 49815r^4 + 135135r^3 + 198936r^2 + 150228r + 45360}$$

Which for the root $r = -\frac{1}{3}$ becomes

$$b_3 = -\frac{7781}{810}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{63r^2+51r-16}{9r^2+21r+12}$	$-\frac{13}{3}$
b_2	$\frac{2673r^4+10044r^3+9441r^2+438r-1568}{81r^4+540r^3+1305r^2+1350r+504}$	$-\frac{251}{45}$
b_3	$\frac{95499r^6+844911r^5+2671137r^4+3571317r^3+1572804r^2-419508r-336448}{729r^6+9477r^5+49815r^4+135135r^3+198936r^2+150228r+45360}$	$-\frac{7781}{810}$

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{3103353r^8 + 49898592r^7 + 320994738r^6 + 1054759968r^5 + 1851078825r^4 + 1583317152r^3 + 348879}{81(3r^2 + 25r + 52)(27r^6 + 351r^5 + 1845r^4 + 5005r^3 + 7368r^2 + 5564r + 1512)}$$

Which for the root $r = -\frac{1}{3}$ becomes

$$b_4 = -\frac{22151}{1215}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{63r^2+51r-16}{9r^2+21r+12}$	$-\frac{13}{3}$
b_2	$\frac{2673r^4+10044r^3+9441r^2+438r-1568}{81r^4+540r^3+1305r^2+1350r+504}$	$-\frac{251}{45}$
b_3	$\frac{95499r^6+844911r^5+2671137r^4+3571317r^3+1572804r^2-419508r-336448}{729r^6+9477r^5+49815r^4+135135r^3+198936r^2+150228r+45360}$	$-\frac{7781}{810}$
b_4	$\frac{3103353r^8+49898592r^7+320994738r^6+1054759968r^5+1851078825r^4+1583317152r^3+348879276r^2-293260704r-127995392}{81(3r^2+25r+52)(27r^6+351r^5+1845r^4+5005r^3+7368r^2+5564r+1680)}$	$-\frac{2215}{121}$

For $n = 5$, using the above recursive equation gives

$$b_5 = \frac{95127939r^{10} + 2420949951r^9 + 26064871578r^8 + 154759665150r^7 + 552318920667r^6 + 1204450764687r^5 + 1536667458552r^4 + 9647868604687r^3 + 2420949951r^2 + 26064871578r + 154759665150}{243(3r^2 + 31r + 80)(3r^2 + 25r + 52)(27r^6 + 351r^5 + 1845r^4 + 5005r^3 + 7368r^2 + 5564r + 1680)}$$

Which for the root $r = -\frac{1}{3}$ becomes

$$b_5 = -\frac{669229}{18225}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	$\frac{63r^2+51r-16}{9r^2+21r+12}$	$-\frac{13}{3}$
b_2	$\frac{2673r^4+10044r^3+9441r^2+438r-1568}{81r^4+540r^3+1305r^2+1350r+504}$	$-\frac{251}{45}$
b_3	$\frac{95499r^6+844911r^5+2671137r^4+3571317r^3+1572804r^2-419508r-336448}{729r^6+9477r^5+49815r^4+135135r^3+198936r^2+150228r+45360}$	$-\frac{7781}{810}$
b_4	$\frac{3103353r^8+49898592r^7+320994738r^6+1054759968r^5+1851078825r^4+1583317152r^3+348879276r^2-293260704r-127995392}{81(3r^2+25r+52)(27r^6+351r^5+1845r^4+5005r^3+7368r^2+5564r+1680)}$	$-\frac{2215}{121}$
b_5	$\frac{95127939r^{10}+2420949951r^9+26064871578r^8+154759665150r^7+552318920667r^6+1204450764687r^5+1536667458552r^4+9647868604687r^3+2420949951r^2+26064871578r+154759665150}{243(3r^2+31r+80)(3r^2+25r+52)(27r^6+351r^5+1845r^4+5005r^3+7368r^2+5564r+1680)}$	$-\frac{669229}{18225}$

For $n = 6$, using the above recursive equation gives

$$b_6 = \frac{2806539921r^{12} + 103709648268r^{11} + 1676277992835r^{10} + 15583794209910r^9 + 92090095190235r^8 + 420949951190235r^7 + 1676277992835r^6 + 103709648268r^5 + 2806539921r^4 + 1676277992835r^3 + 103709648268r^2 + 2806539921r + 1676277992835}{729(3r^2 + 31r + 80)(3r^2 + 25r + 52)(27r^6 + 351r^5 + 1845r^4 + 5005r^3 + 7368r^2 + 5564r + 1680)}$$

Which for the root $r = -\frac{1}{3}$ becomes

$$b_6 = -\frac{216463313}{2788425}$$

And the table now becomes

n	$b_{n,r}$
b_0	1
b_1	$\frac{63r^2+51r-16}{9r^2+21r+12}$
b_2	$\frac{2673r^4+10044r^3+9441r^2+438r-1568}{81r^4+540r^3+1305r^2+1350r+504}$
b_3	$\frac{95499r^6+844911r^5+2671137r^4+3571317r^3+1572804r^2-419508r-336448}{729r^6+9477r^5+49815r^4+135135r^3+198936r^2+150228r+45360}$
b_4	$\frac{3103353r^8+49898592r^7+320994738r^6+1054759968r^5+1851078825r^4+1583317152r^3+348879276r^2-293260704r-127995392}{81(3r^2+25r+52)(27r^6+351r^5+1845r^4+5005r^3+7368r^2+5564r+1680)}$
b_5	$\frac{95127939r^{10}+2420949951r^9+26064871578r^8+154759665150r^7+552318920667r^6+1204450764687r^5+1536667458552r^4+964786860468r^3+127995392r^2-293260704r-127995392}{243(3r^2+31r+80)(3r^2+25r+52)(27r^6+351r^5+1845r^4+5005r^3+7368r^2+5564r+1680)}$
b_6	$\frac{2806539921r^{12}+103709648268r^{11}+1676277992835r^{10}+15583794209910r^9+92090095190235r^8+360297425137320r^7+940035462956685r^6+127995392r^5-293260704r^4-127995392r^3+348879276r^2-293260704r-127995392}{729(3r^2+31r+80)(3r^2+25r+52)(27r^6+351r^5+1845r^4+5005r^3+7368r^2+5564r+1680)}$

For $n = 7$, using the above recursive equation gives

$$b_7 = \frac{80674338123r^{14} + 4080569276151r^{13} + 92307128293827r^{12} + 1233512636444739r^{11} + 10829721106831365r^{10} + 65685074939730909r^9 + 28142187(3r^2+31r+80)r^8 + 28142187(3r^2+31r+80)r^7 + 28142187(3r^2+31r+80)r^6 + 28142187(3r^2+31r+80)r^5 + 28142187(3r^2+31r+80)r^4 + 28142187(3r^2+31r+80)r^3 + 28142187(3r^2+31r+80)r^2 + 28142187(3r^2+31r+80)r + 28142187(3r^2+31r+80)}{2187(3r^2+31r+80)}$$

Which for the root $r = -\frac{1}{3}$ becomes

$$b_7 = -\frac{7179886604}{41826375}$$

And the table now becomes

n	$b_{n,r}$
b_0	1
b_1	$\frac{63r^2+51r-16}{9r^2+21r+12}$
b_2	$\frac{2673r^4+10044r^3+9441r^2+438r-1568}{81r^4+540r^3+1305r^2+1350r+504}$
b_3	$\frac{95499r^6+844911r^5+2671137r^4+3571317r^3+1572804r^2-419508r-336448}{729r^6+9477r^5+49815r^4+135135r^3+198936r^2+150228r+45360}$
b_4	$\frac{3103353r^8+49898592r^7+320994738r^6+1054759968r^5+1851078825r^4+1583317152r^3+348879276r^2-293260704r-127995392}{81(3r^2+25r+52)(27r^6+351r^5+1845r^4+5005r^3+7368r^2+5564r+1680)}$
b_5	$\frac{95127939r^{10}+2420949951r^9+26064871578r^8+154759665150r^7+552318920667r^6+1204450764687r^5+1536667458552r^4+964786860468r^3+127995392r^2-293260704r-127995392}{243(3r^2+31r+80)(3r^2+25r+52)(27r^6+351r^5+1845r^4+5005r^3+7368r^2+5564r+1680)}$
b_6	$\frac{2806539921r^{12}+103709648268r^{11}+1676277992835r^{10}+15583794209910r^9+92090095190235r^8+360297425137320r^7+940035462956685r^6+127995392r^5-293260704r^4-127995392r^3+348879276r^2-293260704r-127995392}{729(3r^2+31r+80)(3r^2+25r+52)(27r^6+351r^5+1845r^4+5005r^3+7368r^2+5564r+1680)}$
b_7	$\frac{80674338123r^{14}+4080569276151r^{13}+92307128293827r^{12}+1233512636444739r^{11}+10829721106831365r^{10}+65685074939730909r^9+28142187(3r^2+31r+80)r^8+28142187(3r^2+31r+80)r^7+28142187(3r^2+31r+80)r^6+28142187(3r^2+31r+80)r^5+28142187(3r^2+31r+80)r^4+28142187(3r^2+31r+80)r^3+28142187(3r^2+31r+80)r^2+28142187(3r^2+31r+80)r+28142187(3r^2+31r+80)}{2187(3r^2+31r+80)}$

Using the above table, then the solution $y_2(t)$ is

$$y_2(t) = 1(b_0 + b_1t + b_2t^2 + b_3t^3 + b_4t^4 + b_5t^5 + b_6t^6 + b_7t^7 + b_8t^8 \dots)$$

$$= \frac{1 - \frac{13t}{3} - \frac{251t^2}{45} - \frac{7781t^3}{810} - \frac{22151t^4}{1215} - \frac{669229t^5}{18225} - \frac{216463313t^6}{2788425} - \frac{7179886604t^7}{41826375} + O(t^8)}{t^{\frac{1}{3}}}$$

Therefore the homogeneous solution is

$$y_h(t) = c_1y_1(t) + c_2y_2(t)$$

$$= c_1 \left(1 - \frac{4t}{3} - \frac{28t^2}{9} - \frac{3004t^3}{405} - \frac{285704t^4}{15795} - \frac{822592t^5}{18225} - \frac{4666732192t^6}{40514175} - \frac{401483448544t^7}{1336967775} + O(t^8) \right)$$

$$+ \frac{c_2 \left(1 - \frac{13t}{3} - \frac{251t^2}{45} - \frac{7781t^3}{810} - \frac{22151t^4}{1215} - \frac{669229t^5}{18225} - \frac{216463313t^6}{2788425} - \frac{7179886604t^7}{41826375} + O(t^8) \right)}{t^{\frac{1}{3}}}$$

Hence the final solution is

$$y(t) = y_h$$

$$= c_1 \left(1 - \frac{4t}{3} - \frac{28t^2}{9} - \frac{3004t^3}{405} - \frac{285704t^4}{15795} - \frac{822592t^5}{18225} - \frac{4666732192t^6}{40514175} - \frac{401483448544t^7}{1336967775} + O(t^8) \right)$$

$$+ \frac{c_2 \left(1 - \frac{13t}{3} - \frac{251t^2}{45} - \frac{7781t^3}{810} - \frac{22151t^4}{1215} - \frac{669229t^5}{18225} - \frac{216463313t^6}{2788425} - \frac{7179886604t^7}{41826375} + O(t^8) \right)}{t^{\frac{1}{3}}}$$

Replacing t by $\frac{1}{x}$ gives

$$y = c_1 \left(1 - \frac{4}{3x} - \frac{28}{9x^2} - \frac{3004}{405x^3} - \frac{285704}{15795x^4} - \frac{822592}{18225x^5} - \frac{4666732192}{40514175x^6} - \frac{401483448544}{1336967775x^7} + O\left(\frac{1}{x^8}\right) \right) + \frac{c_2}{t^{\frac{1}{3}}}$$

Summary

The solution(s) found are the following

$$y = c_1 \left(1 - \frac{4}{3x} - \frac{28}{9x^2} - \frac{3004}{405x^3} - \frac{285704}{15795x^4} - \frac{822592}{18225x^5} - \frac{4666732192}{40514175x^6} - \frac{401483448544}{1336967775x^7} + O\left(\frac{1}{x^8}\right) \right)$$

$$+ \frac{c_2 \left(1 - \frac{13}{3x} - \frac{251}{45x^2} - \frac{7781}{810x^3} - \frac{22151}{1215x^4} - \frac{669229}{18225x^5} - \frac{216463313}{2788425x^6} - \frac{7179886604}{41826375x^7} + O\left(\frac{1}{x^8}\right) \right)}{\left(\frac{1}{x}\right)^{\frac{1}{3}}}$$

Verification of solutions

$$y = c_1 \left(1 - \frac{4}{3x} - \frac{28}{9x^2} - \frac{3004}{405x^3} - \frac{285704}{15795x^4} - \frac{822592}{18225x^5} - \frac{4666732192}{40514175x^6} - \frac{401483448544}{1336967775x^7} + O\left(\frac{1}{x^8}\right) \right) \\ + \frac{c_2 \left(1 - \frac{13}{3x} - \frac{251}{45x^2} - \frac{7781}{810x^3} - \frac{22151}{1215x^4} - \frac{669229}{18225x^5} - \frac{216463313}{2788425x^6} - \frac{7179886604}{41826375x^7} + O\left(\frac{1}{x^8}\right) \right)}{\left(\frac{1}{x}\right)^{\frac{1}{3}}}$$

Verified OK.

23.2.1 Maple step by step solution

Let's solve

$$9(-2+x)^2(x-3)y'' + (6x^2 - 12x)y' + 16y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{16y}{9(-2+x)^2(x-3)} - \frac{2xy'}{3(x-3)(-2+x)}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{2xy'}{3(x-3)(-2+x)} + \frac{16y}{9(-2+x)^2(x-3)} = 0$$

- Check to see if x_0 is a regular singular point

- Define functions

$$\left[P_2(x) = \frac{2x}{3(x-3)(-2+x)}, P_3(x) = \frac{16}{9(-2+x)^2(x-3)} \right]$$

- $(-2+x) \cdot P_2(x)$ is analytic at $x = 2$

$$\left. ((-2+x) \cdot P_2(x)) \right|_{x=2} = -\frac{4}{3}$$

- $(-2+x)^2 \cdot P_3(x)$ is analytic at $x = 2$

$$\left. ((-2+x)^2 \cdot P_3(x)) \right|_{x=2} = -\frac{16}{9}$$

- $x = 2$ is a regular singular point

Check to see if x_0 is a regular singular point

$$x_0 = 2$$

- Multiply by denominators

$$9(-2+x)^2(x-3)y'' + 6x(-2+x)y' + 16y = 0$$

- Change variables using $x = u + 2$ so that the regular singular point is at $u = 0$

$$(9u^3 - 9u^2) \left(\frac{d^2}{du^2} y(u) \right) + (6u^2 + 12u) \left(\frac{d}{du} y(u) \right) + 16y(u) = 0$$

- Assume series solution for $y(u)$

$$y(u) = \sum_{k=0}^{\infty} a_k u^{k+r}$$

- Rewrite ODE with series expansions

- Convert $u^m \cdot \left(\frac{d}{du} y(u) \right)$ to series expansion for $m = 1..2$

$$u^m \cdot \left(\frac{d}{du} y(u) \right) = \sum_{k=0}^{\infty} a_k (k+r) u^{k+r-1+m}$$

- Shift index using $k- > k + 1 - m$

$$u^m \cdot \left(\frac{d}{du} y(u) \right) = \sum_{k=-1+m}^{\infty} a_{k+1-m} (k+1-m+r) u^{k+r}$$

- Convert $u^m \cdot \left(\frac{d^2}{du^2} y(u) \right)$ to series expansion for $m = 2..3$

$$u^m \cdot \left(\frac{d^2}{du^2} y(u) \right) = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) u^{k+r-2+m}$$

- Shift index using $k- > k + 2 - m$

$$u^m \cdot \left(\frac{d^2}{du^2} y(u) \right) = \sum_{k=-2+m}^{\infty} a_{k+2-m} (k+2-m+r)(k+1-m+r) u^{k+r}$$

Rewrite ODE with series expansions

$$-a_0(9r^2 - 21r - 16) u^r + \left(\sum_{k=1}^{\infty} (-a_k(9k^2 + 18kr + 9r^2 - 21k - 21r - 16) + 3a_{k-1}(k+r-1) \right) (3$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$-9r^2 + 21r + 16 = 0$$

- Values of r that satisfy the indicial equation

$$r \in \left\{ \frac{7}{6} - \frac{\sqrt{113}}{6}, \frac{7}{6} + \frac{\sqrt{113}}{6} \right\}$$

- Each term in the series must be 0, giving the recursion relation

$$9(k+r-1) \left(k - \frac{4}{3} + r \right) a_{k-1} - 9a_k \left(k^2 + \left(2r - \frac{7}{3} \right) k + r^2 - \frac{7r}{3} - \frac{16}{9} \right) = 0$$

- Shift index using $k- > k + 1$

$$9(k+r) \left(k - \frac{1}{3} + r \right) a_k - 9a_{k+1} \left((k+1)^2 + \left(2r - \frac{7}{3} \right) (k+1) + r^2 - \frac{7r}{3} - \frac{16}{9} \right) = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+1} = \frac{3(k+r)(3k+3r-1)a_k}{9k^2+18kr+9r^2-3k-3r-28}$$

- Recursion relation for $r = \frac{7}{6} - \frac{\sqrt{113}}{6}$

$$a_{k+1} = \frac{3\left(k+\frac{7}{6}-\frac{\sqrt{113}}{6}\right)\left(3k+\frac{5}{2}-\frac{\sqrt{113}}{2}\right)a_k}{9k^2+18k\left(\frac{7}{6}-\frac{\sqrt{113}}{6}\right)+9\left(\frac{7}{6}-\frac{\sqrt{113}}{6}\right)^2-3k-\frac{63}{2}+\frac{\sqrt{113}}{2}}$$

- Solution for $r = \frac{7}{6} - \frac{\sqrt{113}}{6}$

$$\left[y(u) = \sum_{k=0}^{\infty} a_k u^{k+\frac{7}{6}-\frac{\sqrt{113}}{6}}, a_{k+1} = \frac{3\left(k+\frac{7}{6}-\frac{\sqrt{113}}{6}\right)\left(3k+\frac{5}{2}-\frac{\sqrt{113}}{2}\right)a_k}{9k^2+18k\left(\frac{7}{6}-\frac{\sqrt{113}}{6}\right)+9\left(\frac{7}{6}-\frac{\sqrt{113}}{6}\right)^2-3k-\frac{63}{2}+\frac{\sqrt{113}}{2}} \right]$$

- Revert the change of variables $u = -2 + x$

$$\left[y = \sum_{k=0}^{\infty} a_k (-2+x)^{k+\frac{7}{6}-\frac{\sqrt{113}}{6}}, a_{k+1} = \frac{3\left(k+\frac{7}{6}-\frac{\sqrt{113}}{6}\right)\left(3k+\frac{5}{2}-\frac{\sqrt{113}}{2}\right)a_k}{9k^2+18k\left(\frac{7}{6}-\frac{\sqrt{113}}{6}\right)+9\left(\frac{7}{6}-\frac{\sqrt{113}}{6}\right)^2-3k-\frac{63}{2}+\frac{\sqrt{113}}{2}} \right]$$

- Recursion relation for $r = \frac{7}{6} + \frac{\sqrt{113}}{6}$

$$a_{k+1} = \frac{3\left(k+\frac{7}{6}+\frac{\sqrt{113}}{6}\right)\left(3k+\frac{5}{2}+\frac{\sqrt{113}}{2}\right)a_k}{9k^2+18k\left(\frac{7}{6}+\frac{\sqrt{113}}{6}\right)+9\left(\frac{7}{6}+\frac{\sqrt{113}}{6}\right)^2-3k-\frac{63}{2}-\frac{\sqrt{113}}{2}}$$

- Solution for $r = \frac{7}{6} + \frac{\sqrt{113}}{6}$

$$\left[y(u) = \sum_{k=0}^{\infty} a_k u^{k+\frac{7}{6}+\frac{\sqrt{113}}{6}}, a_{k+1} = \frac{3\left(k+\frac{7}{6}+\frac{\sqrt{113}}{6}\right)\left(3k+\frac{5}{2}+\frac{\sqrt{113}}{2}\right)a_k}{9k^2+18k\left(\frac{7}{6}+\frac{\sqrt{113}}{6}\right)+9\left(\frac{7}{6}+\frac{\sqrt{113}}{6}\right)^2-3k-\frac{63}{2}-\frac{\sqrt{113}}{2}} \right]$$

- Revert the change of variables $u = -2 + x$

$$\left[y = \sum_{k=0}^{\infty} a_k (-2+x)^{k+\frac{7}{6}+\frac{\sqrt{113}}{6}}, a_{k+1} = \frac{3\left(k+\frac{7}{6}+\frac{\sqrt{113}}{6}\right)\left(3k+\frac{5}{2}+\frac{\sqrt{113}}{2}\right)a_k}{9k^2+18k\left(\frac{7}{6}+\frac{\sqrt{113}}{6}\right)+9\left(\frac{7}{6}+\frac{\sqrt{113}}{6}\right)^2-3k-\frac{63}{2}-\frac{\sqrt{113}}{2}} \right]$$

- Combine solutions and rename parameters

$$\left[y = \left(\sum_{k=0}^{\infty} a_k (-2+x)^{k+\frac{7}{6}-\frac{\sqrt{113}}{6}} \right) + \left(\sum_{k=0}^{\infty} b_k (-2+x)^{k+\frac{7}{6}+\frac{\sqrt{113}}{6}} \right), a_{k+1} = \frac{3\left(k+\frac{7}{6}-\frac{\sqrt{113}}{6}\right)\left(3k+\frac{5}{2}-\frac{\sqrt{113}}{2}\right)a_k}{9k^2+18k\left(\frac{7}{6}-\frac{\sqrt{113}}{6}\right)+9\left(\frac{7}{6}-\frac{\sqrt{113}}{6}\right)^2-3k-\frac{63}{2}+\frac{\sqrt{113}}{2}} \right]$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
  -> Bessel
  -> elliptic
  -> Legendre
  -> Kummer
    -> hyper3: Equivalence to 1F1 under a power @ Moebius
  -> hypergeometric
    -> heuristic approach
    -> hyper3: Equivalence to 2F1, 1F1 or 0F1 under a power @ Moebius
      <- hyper3 successful: received ODE is equivalent to the 2F1 ODE
    <- hypergeometric successful
<- special function solution successful`
```

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 414

```
Order:=8;
dsolve(9*(x-2)^2*(x-3)*diff(y(x),x$2)+6*x*(x-2)*diff(y(x),x)+16*y(x)=0,y(x),type='series',x=
```

Expression too large to display

✓ Solution by Mathematica

Time used: 0.009 (sec). Leaf size: 130

```
AsymptoticDSolveValue[9*(x-2)^2*(x-3)*y'[x]+6*x*(x-2)*y'[x]+16*y[x]==0,y[x],{x,infinity,7}]
```

$$y(x) \rightarrow c_2 \left(-\frac{13}{3x^{2/3}} - \frac{251}{45x^{5/3}} - \frac{7781}{810x^{8/3}} - \frac{22151}{1215x^{11/3}} - \frac{669229}{18225x^{14/3}} - \frac{216463313}{2788425x^{17/3}} - \frac{7179886604}{41826375x^{20/3}} \right) + \sqrt[3]{x} \left(c_1 \left(-\frac{401483448544}{1336967775x^7} - \frac{4666732192}{40514175x^6} - \frac{822592}{18225x^5} - \frac{285704}{15795x^4} - \frac{3004}{405x^3} - \frac{28}{9x^2} - \frac{4}{3x} + 1 \right) \right)$$

23.3 problem 1(c)

23.3.1 Maple step by step solution 3849

Internal problem ID [6498]

Internal file name [OUTPUT/5746_Sunday_June_05_2022_03_52_34_PM_45330625/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 4. Power Series Solutions and Special Functions. Problems for review and discover. (B) Challenge Problems . Page 194

Problem number: 1(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"second order series method. Regular singular point. Difference not integer"**

Maple gives the following as the ode type

[_Gegenbauer]

$$(-x^2 + 1)y'' - 2xy' + p(p + 1)y = 0$$

With the expansion point for the power series method at $x = \infty$.

Since expansion is around ∞ , then the independent variable x is replaced by $\frac{1}{t}$ and the expansion is made around $t = 0$ and after solving, the solution is changed back to x using $x = \frac{1}{t}$. Changing variables results in the new ode

$$-\left(-\frac{1}{t^2} + 1\right) \left(-\left(\frac{d^2}{dt^2}y(t)\right)t^2 - 2t\left(\frac{d}{dt}y(t)\right)\right) t^2 + 2t\left(\frac{d}{dt}y(t)\right) + (p^2 + p)y(t) = 0$$

The transformed ODE is now solved. The type of the expansion point is first determined. This is done on the homogeneous part of the ODE.

$$(t^4 - t^2) \left(\frac{d^2}{dt^2}y(t)\right) + 2\left(\frac{d}{dt}y(t)\right)t^3 + (p^2 + p)y(t) = 0$$

The following is summary of singularities for the above ode. Writing the ode as

$$\frac{d^2}{dt^2}y(t) + p(t)\frac{d}{dt}y(t) + q(t)y(t) = 0$$

Where

$$p(t) = \frac{2t}{t^2 - 1}$$

$$q(t) = \frac{p(p+1)}{(t^2 - 1)t^2}$$

Table 519: Table $p(t), q(t)$ singularities.

$p(t) = \frac{2t}{t^2-1}$	
singularity	type
$t = -1$	“regular”
$t = 1$	“regular”

$q(t) = \frac{p(p+1)}{(t^2-1)t^2}$	
singularity	type
$t = -1$	“regular”
$t = 0$	“regular”
$t = 1$	“regular”

Combining everything together gives the following summary of singularities for the ode as

Regular singular points : $[-1, 1, 0, \infty]$

Irregular singular points : $[\]$

Since $t = 0$ is regular singular point, then Frobenius power series is used. The ode is normalized to be

$$\left(\frac{d^2}{dt^2}y(t)\right) (t^2 - 1) t^2 + 2\left(\frac{d}{dt}y(t)\right) t^3 + (p^2 + p) y(t) = 0$$

Let the solution be represented as Frobenius power series of the form

$$y(t) = \sum_{n=0}^{\infty} a_n t^{n+r}$$

Then

$$\frac{d}{dt}y(t) = \sum_{n=0}^{\infty} (n+r) a_n t^{n+r-1}$$

$$\frac{d^2}{dt^2}y(t) = \sum_{n=0}^{\infty} (n+r)(n+r-1) a_n t^{n+r-2}$$

Substituting the above back into the ode gives

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} (n+r)(n+r-1) a_n t^{n+r-2} \right) (t^2 - 1) t^2 \\ & + 2 \left(\sum_{n=0}^{\infty} (n+r) a_n t^{n+r-1} \right) t^3 + (p^2 + p) \left(\sum_{n=0}^{\infty} a_n t^{n+r} \right) = 0 \end{aligned} \quad (1)$$

Which simplifies to

$$\begin{aligned} & \left(\sum_{n=0}^{\infty} t^{n+r+2} a_n (n+r)(n+r-1) \right) + \sum_{n=0}^{\infty} (-t^{n+r} a_n (n+r)(n+r-1)) \\ & + \left(\sum_{n=0}^{\infty} 2t^{n+r+2} a_n (n+r) \right) + \left(\sum_{n=0}^{\infty} (p^2 + p) a_n t^{n+r} \right) = 0 \end{aligned} \quad (2A)$$

The next step is to make all powers of t be $n+r$ in each summation term. Going over each summation term above with power of t in it which is not already t^{n+r} and adjusting the power and the corresponding index gives

$$\begin{aligned} \sum_{n=0}^{\infty} t^{n+r+2} a_n (n+r)(n+r-1) &= \sum_{n=2}^{\infty} a_{n-2} (n+r-2)(n-3+r) t^{n+r} \\ \sum_{n=0}^{\infty} 2t^{n+r+2} a_n (n+r) &= \sum_{n=2}^{\infty} 2a_{n-2} (n+r-2) t^{n+r} \end{aligned}$$

Substituting all the above in Eq (2A) gives the following equation where now all powers of t are the same and equal to $n+r$.

$$\begin{aligned} & \left(\sum_{n=2}^{\infty} a_{n-2} (n+r-2)(n-3+r) t^{n+r} \right) + \sum_{n=0}^{\infty} (-t^{n+r} a_n (n+r)(n+r-1)) \\ & + \left(\sum_{n=2}^{\infty} 2a_{n-2} (n+r-2) t^{n+r} \right) + \left(\sum_{n=0}^{\infty} (p^2 + p) a_n t^{n+r} \right) = 0 \end{aligned} \quad (2B)$$

The indicial equation is obtained from $n=0$. From Eq (2B) this gives

$$-t^{n+r} a_n (n+r)(n+r-1) + (p^2 + p) a_n t^{n+r} = 0$$

When $n = 0$ the above becomes

$$-t^r a_0 r(-1 + r) + (p^2 + p) a_0 t^r = 0$$

Or

$$(-t^r r(-1 + r) + (p^2 + p) t^r) a_0 = 0$$

Since $a_0 \neq 0$ then the above simplifies to

$$(p^2 - r^2 + p + r) t^r = 0$$

Since the above is true for all t then the indicial equation becomes

$$p^2 - r^2 + p + r = 0$$

Solving for r gives the roots of the indicial equation as

$$\begin{aligned} r_1 &= -p \\ r_2 &= p + 1 \end{aligned}$$

Since $a_0 \neq 0$ then the indicial equation becomes

$$(p^2 - r^2 + p + r) t^r = 0$$

Solving for r gives the roots of the indicial equation as Assuming the roots differ by non-integer Since $r_1 - r_2 = -2p - 1$ is not an integer, then we can construct two linearly independent solutions

$$\begin{aligned} y_1(t) &= t^{r_1} \left(\sum_{n=0}^{\infty} a_n t^n \right) \\ y_2(t) &= t^{r_2} \left(\sum_{n=0}^{\infty} b_n t^n \right) \end{aligned}$$

Or

$$\begin{aligned} y_1(t) &= \sum_{n=0}^{\infty} a_n t^{n-p} \\ y_2(t) &= \sum_{n=0}^{\infty} b_n t^{n+p+1} \end{aligned}$$

We start by finding $y_1(t)$. Eq (2B) derived above is now used to find all a_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. a_0 is arbitrary and taken as $a_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$a_1 = 0$$

For $2 \leq n$ the recursive equation is

$$a_{n-2}(n+r-2)(n-3+r) - a_n(n+r)(n+r-1) + 2a_{n-2}(n+r-2) + a_n p(p+1) = 0 \quad (3)$$

Solving for a_n from recursive equation (4) gives

$$a_n = \frac{a_{n-2}(n^2 + 2nr + r^2 - 3n - 3r + 2)}{n^2 + 2nr - p^2 + r^2 - n - p - r} \quad (4)$$

Which for the root $r = -p$ becomes

$$a_n = \frac{a_{n-2}(n-1-p)(n-p-2)}{n(n-2p-1)} \quad (5)$$

At this point, it is a good idea to keep track of a_n in a table both before substituting $r = -p$ and after as more terms are found using the above recursive equation.

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0

For $n = 2$, using the above recursive equation gives

$$a_2 = -\frac{(1+r)r}{p^2 - r^2 + p - 3r - 2}$$

Which for the root $r = -p$ becomes

$$a_2 = -\frac{(-1+p)p}{4p-2}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{(1+r)r}{p^2 - r^2 + p - 3r - 2}$	$-\frac{(-1+p)p}{4p-2}$

For $n = 3$, using the above recursive equation gives

$$a_3 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{(1+r)r}{p^2-r^2+p-3r-2}$	$-\frac{(-1+p)p}{4p-2}$
a_3	0	0

For $n = 4$, using the above recursive equation gives

$$a_4 = \frac{(1+r)r(r+3)(r+2)}{(p+r+2)(p-1-r)(p+4+r)(p-3-r)}$$

Which for the root $r = -p$ becomes

$$a_4 = \frac{(-1+p)p(p-3)(p-2)}{32p^2-64p+24}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{(1+r)r}{p^2-r^2+p-3r-2}$	$-\frac{(-1+p)p}{4p-2}$
a_3	0	0
a_4	$\frac{(1+r)r(r+3)(r+2)}{(p+r+2)(p-1-r)(p+4+r)(p-3-r)}$	$\frac{(-1+p)p(p-3)(p-2)}{32p^2-64p+24}$

For $n = 5$, using the above recursive equation gives

$$a_5 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{(1+r)r}{p^2-r^2+p-3r-2}$	$-\frac{(-1+p)p}{4p-2}$
a_3	0	0
a_4	$\frac{(1+r)r(r+3)(r+2)}{(p+r+2)(p-1-r)(p+4+r)(p-3-r)}$	$\frac{(-1+p)p(p-3)(p-2)}{32p^2-64p+24}$
a_5	0	0

For $n = 6$, using the above recursive equation gives

$$a_6 = -\frac{(1+r)r(r+3)(r+2)(r+5)(4+r)}{(p+r+2)(p-1-r)(p+4+r)(p-3-r)(p+6+r)(p-5-r)}$$

Which for the root $r = -p$ becomes

$$a_6 = -\frac{(-1+p)p(p-3)(p-2)(p-5)(-4+p)}{384p^3 - 1728p^2 + 2208p - 720}$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{(1+r)r}{p^2-r^2+p-3r-2}$	$-\frac{(-1+p)p}{4p-2}$
a_3	0	0
a_4	$\frac{(1+r)r(r+3)(r+2)}{(p+r+2)(p-1-r)(p+4+r)(p-3-r)}$	$\frac{(-1+p)p(p-3)(p-2)}{32p^2-64p+24}$
a_5	0	0
a_6	$-\frac{(1+r)r(r+3)(r+2)(r+5)(4+r)}{(p+r+2)(p-1-r)(p+4+r)(p-3-r)(p+6+r)(p-5-r)}$	$-\frac{(-1+p)p(p-3)(p-2)(p-5)(-4+p)}{384p^3-1728p^2+2208p-720}$

For $n = 7$, using the above recursive equation gives

$$a_7 = 0$$

And the table now becomes

n	$a_{n,r}$	a_n
a_0	1	1
a_1	0	0
a_2	$-\frac{(1+r)r}{p^2-r^2+p-3r-2}$	$-\frac{(-1+p)p}{4p-2}$
a_3	0	0
a_4	$\frac{(1+r)r(r+3)(r+2)}{(p+r+2)(p-1-r)(p+4+r)(p-3-r)}$	$\frac{(-1+p)p(p-3)(p-2)}{32p^2-64p+24}$
a_5	0	0
a_6	$-\frac{(1+r)r(r+3)(r+2)(r+5)(4+r)}{(p+r+2)(p-1-r)(p+4+r)(p-3-r)(p+6+r)(p-5-r)}$	$-\frac{(-1+p)p(p-3)(p-2)(p-5)(-4+p)}{384p^3-1728p^2+2208p-720}$
a_7	0	0

Using the above table, then the solution $y_1(t)$ is

$$y_1(t) = t^{-p}(a_0 + a_1t + a_2t^2 + a_3t^3 + a_4t^4 + a_5t^5 + a_6t^6 + a_7t^7 + a_8t^8 \dots)$$

$$= t^{-p} \left(1 - \frac{(-1+p)p t^2}{4p-2} + \frac{(-1+p)p(p-3)(p-2)t^4}{32p^2-64p+24} - \frac{(-1+p)p(p-3)(p-2)(p-5)(-4+p)t^6}{384p^3-1728p^2+2208p-720} \right)$$

Now the second solution $y_2(t)$ is found. Eq (2B) derived above is now used to find all b_n coefficients. The case $n = 0$ is skipped since it was used to find the roots of the indicial equation. b_0 is arbitrary and taken as $b_0 = 1$. Substituting $n = 1$ in Eq. (2B) gives

$$b_1 = 0$$

For $2 \leq n$ the recursive equation is

$$b_{n-2}(n+r-2)(n-3+r) - b_n(n+r)(n+r-1) + 2b_{n-2}(n+r-2) + p(p+1)b_n = 0 \quad (3)$$

Solving for b_n from recursive equation (4) gives

$$b_n = \frac{b_{n-2}(n^2 + 2nr + r^2 - 3n - 3r + 2)}{n^2 + 2nr - p^2 + r^2 - n - p - r} \quad (4)$$

Which for the root $r = p + 1$ becomes

$$b_n = \frac{b_{n-2}(n+p)(n+p-1)}{n(n+2p+1)} \quad (5)$$

At this point, it is a good idea to keep track of b_n in a table both before substituting $r = p + 1$ and after as more terms are found using the above recursive equation.

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0

For $n = 2$, using the above recursive equation gives

$$b_2 = -\frac{(1+r)r}{p^2 - r^2 + p - 3r - 2}$$

Which for the root $r = p + 1$ becomes

$$b_2 = \frac{(p+2)(p+1)}{4p+6}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{(1+r)r}{p^2 - r^2 + p - 3r - 2}$	$\frac{(p+2)(p+1)}{4p+6}$

For $n = 3$, using the above recursive equation gives

$$b_3 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{(1+r)r}{p^2 - r^2 + p - 3r - 2}$	$\frac{(p+2)(p+1)}{4p+6}$
b_3	0	0

For $n = 4$, using the above recursive equation gives

$$b_4 = \frac{(1+r)r(r+3)(r+2)}{(p+r+2)(p-1-r)(p+4+r)(p-3-r)}$$

Which for the root $r = p + 1$ becomes

$$b_4 = \frac{(p+1)(p+2)(p+3)(p+4)}{32p^2 + 128p + 120}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{(1+r)r}{p^2-r^2+p-3r-2}$	$\frac{(p+2)(p+1)}{4p+6}$
b_3	0	0
b_4	$\frac{(1+r)r(r+3)(r+2)}{(p+r+2)(p-1-r)(p+4+r)(p-3-r)}$	$\frac{(p+1)(p+2)(p+3)(p+4)}{32p^2+128p+120}$

For $n = 5$, using the above recursive equation gives

$$b_5 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{(1+r)r}{p^2-r^2+p-3r-2}$	$\frac{(p+2)(p+1)}{4p+6}$
b_3	0	0
b_4	$\frac{(1+r)r(r+3)(r+2)}{(p+r+2)(p-1-r)(p+4+r)(p-3-r)}$	$\frac{(p+1)(p+2)(p+3)(p+4)}{32p^2+128p+120}$
b_5	0	0

For $n = 6$, using the above recursive equation gives

$$b_6 = -\frac{(1+r)r(r+3)(r+2)(r+5)(4+r)}{(p+r+2)(p-1-r)(p+4+r)(p-3-r)(p+6+r)(p-5-r)}$$

Which for the root $r = p + 1$ becomes

$$b_6 = \frac{(p+2)(p+1)(p+4)(p+3)(p+6)(5+p)}{384p^3 + 2880p^2 + 6816p + 5040}$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{(1+r)r}{p^2-r^2+p-3r-2}$	$\frac{(p+2)(p+1)}{4p+6}$
b_3	0	0
b_4	$\frac{(1+r)r(r+3)(r+2)}{(p+r+2)(p-1-r)(p+4+r)(p-3-r)}$	$\frac{(p+1)(p+2)(p+3)(p+4)}{32p^2+128p+120}$
b_5	0	0
b_6	$-\frac{(1+r)r(r+3)(r+2)(r+5)(4+r)}{(p+r+2)(p-1-r)(p+4+r)(p-3-r)(p+6+r)(p-5-r)}$	$\frac{(p+2)(p+1)(p+4)(p+3)(p+6)(5+p)}{384p^3+2880p^2+6816p+5040}$

For $n = 7$, using the above recursive equation gives

$$b_7 = 0$$

And the table now becomes

n	$b_{n,r}$	b_n
b_0	1	1
b_1	0	0
b_2	$-\frac{(1+r)r}{p^2-r^2+p-3r-2}$	$\frac{(p+2)(p+1)}{4p+6}$
b_3	0	0
b_4	$\frac{(1+r)r(r+3)(r+2)}{(p+r+2)(p-1-r)(p+4+r)(p-3-r)}$	$\frac{(p+1)(p+2)(p+3)(p+4)}{32p^2+128p+120}$
b_5	0	0
b_6	$-\frac{(1+r)r(r+3)(r+2)(r+5)(4+r)}{(p+r+2)(p-1-r)(p+4+r)(p-3-r)(p+6+r)(p-5-r)}$	$\frac{(p+2)(p+1)(p+4)(p+3)(p+6)(5+p)}{384p^3+2880p^2+6816p+5040}$
b_7	0	0

Using the above table, then the solution $y_2(t)$ is

$$\begin{aligned}
y_2(t) &= t^{-p}(b_0 + b_1t + b_2t^2 + b_3t^3 + b_4t^4 + b_5t^5 + b_6t^6 + b_7t^7 + b_8t^8 \dots) \\
&= t^{p+1} \left(1 + \frac{(p+2)(p+1)t^2}{4p+6} + \frac{(p+1)(p+2)(p+3)(p+4)t^4}{32p^2+128p+120} + \frac{(p+2)(p+1)(p+4)(p+3)(p+6)(5+p)t^6}{384p^3+2880p^2+6816p+5040} \right)
\end{aligned}$$

Therefore the homogeneous solution is

$$y_h(t) = c_1y_1(t) + c_2y_2(t)$$

$$\begin{aligned}
&= c_1 t^{-p} \left(1 - \frac{(-1+p)p t^2}{4p-2} + \frac{(-1+p)p(p-3)(p-2)t^4}{32p^2-64p+24} \right. \\
&\quad \left. - \frac{(-1+p)p(p-3)(p-2)(p-5)(-4+p)t^6}{384p^3-1728p^2+2208p-720} + O(t^8) \right) \\
&+ c_2 t^{p+1} \left(1 + \frac{(p+2)(p+1)t^2}{4p+6} + \frac{(p+1)(p+2)(p+3)(p+4)t^4}{32p^2+128p+120} \right. \\
&\quad \left. + \frac{(p+2)(p+1)(p+4)(p+3)(p+6)(5+p)t^6}{384p^3+2880p^2+6816p+5040} + O(t^8) \right)
\end{aligned}$$

Hence the final solution is

$$\begin{aligned}
y(t) &= y_h \\
&= c_1 t^{-p} \left(1 - \frac{(-1+p)p t^2}{4p-2} + \frac{(-1+p)p(p-3)(p-2)t^4}{32p^2-64p+24} \right. \\
&\quad \left. - \frac{(-1+p)p(p-3)(p-2)(p-5)(-4+p)t^6}{384p^3-1728p^2+2208p-720} + O(t^8) \right) \\
&+ c_2 t^{p+1} \left(1 + \frac{(p+2)(p+1)t^2}{4p+6} + \frac{(p+1)(p+2)(p+3)(p+4)t^4}{32p^2+128p+120} \right. \\
&\quad \left. + \frac{(p+2)(p+1)(p+4)(p+3)(p+6)(5+p)t^6}{384p^3+2880p^2+6816p+5040} + O(t^8) \right)
\end{aligned}$$

Replacing t by $\frac{1}{x}$ gives

$$y = c_1 \left(\frac{1}{x} \right)^{-p} \left(1 - \frac{(-1+p)p}{(4p-2)x^2} + \frac{(-1+p)p(p-3)(p-2)}{(32p^2-64p+24)x^4} - \frac{(-1+p)p(p-3)(p-2)(p-5)(-4+p)}{(384p^3-1728p^2+2208p-720)x^6} \right)$$

Summary

The solution(s) found are the following

$$\begin{aligned}
y &= c_1 \left(\frac{1}{x} \right)^{-p} \left(1 - \frac{(-1+p)p}{(4p-2)x^2} + \frac{(-1+p)p(p-3)(p-2)}{(32p^2-64p+24)x^4} \right. \\
&\quad \left. - \frac{(-1+p)p(p-3)(p-2)(p-5)(-4+p)}{(384p^3-1728p^2+2208p-720)x^6} + O\left(\frac{1}{x^8}\right) \right) \\
&+ c_2 \left(\frac{1}{x} \right)^{p+1} \left(1 + \frac{(p+2)(p+1)}{(4p+6)x^2} + \frac{(p+1)(p+2)(p+3)(p+4)}{(32p^2+128p+120)x^4} \right. \\
&\quad \left. + \frac{(p+2)(p+1)(p+4)(p+3)(p+6)(5+p)}{(384p^3+2880p^2+6816p+5040)x^6} + O\left(\frac{1}{x^8}\right) \right)
\end{aligned}$$

Verification of solutions

$$y = c_1 \left(\frac{1}{x}\right)^{-p} \left(1 - \frac{(-1+p)p}{(4p-2)x^2} + \frac{(-1+p)p(p-3)(p-2)}{(32p^2-64p+24)x^4} - \frac{(-1+p)p(p-3)(p-2)(p-5)(-4+p)}{(384p^3-1728p^2+2208p-720)x^6} + O\left(\frac{1}{x^8}\right)\right) \\ + c_2 \left(\frac{1}{x}\right)^{p+1} \left(1 + \frac{(p+2)(p+1)}{(4p+6)x^2} + \frac{(p+1)(p+2)(p+3)(p+4)}{(32p^2+128p+120)x^4} + \frac{(p+2)(p+1)(p+4)(p+3)(p+6)(5+p)}{(384p^3+2880p^2+6816p+5040)x^6} + O\left(\frac{1}{x^8}\right)\right)$$

Verified OK.

23.3.1 Maple step by step solution

Let's solve

$$(-x^2 + 1)y'' - 2xy' + (p^2 + p)y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = \frac{p(p+1)y}{x^2-1} - \frac{2xy'}{x^2-1}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{2xy'}{x^2-1} - \frac{p(p+1)y}{x^2-1} = 0$$

- Check to see if x_0 is a regular singular point

- Define functions

$$\left[P_2(x) = \frac{2x}{x^2-1}, P_3(x) = -\frac{p(p+1)}{x^2-1} \right]$$

- $(1+x) \cdot P_2(x)$ is analytic at $x = -1$

$$\left. ((1+x) \cdot P_2(x)) \right|_{x=-1} = 1$$

- $(1+x)^2 \cdot P_3(x)$ is analytic at $x = -1$

$$\left. ((1+x)^2 \cdot P_3(x)) \right|_{x=-1} = 0$$

- $x = -1$ is a regular singular point

Check to see if x_0 is a regular singular point

$$x_0 = -1$$

- Multiply by denominators

$$(x^2 - 1)y'' + 2xy' - p(p+1)y = 0$$

- Change variables using $x = u - 1$ so that the regular singular point is at $u = 0$

$$(u^2 - 2u) \left(\frac{d^2}{du^2} y(u) \right) + (2u - 2) \left(\frac{d}{du} y(u) \right) + (-p^2 - p)y(u) = 0$$

- Assume series solution for $y(u)$

$$y(u) = \sum_{k=0}^{\infty} a_k u^{k+r}$$

- Rewrite ODE with series expansions

- Convert $u^m \cdot \left(\frac{d}{du} y(u) \right)$ to series expansion for $m = 0..1$

$$u^m \cdot \left(\frac{d}{du} y(u) \right) = \sum_{k=0}^{\infty} a_k (k+r) u^{k+r-1+m}$$

- Shift index using $k \rightarrow k+1-m$

$$u^m \cdot \left(\frac{d}{du} y(u) \right) = \sum_{k=-1+m}^{\infty} a_{k+1-m} (k+1-m+r) u^{k+r}$$

- Convert $u^m \cdot \left(\frac{d^2}{du^2} y(u) \right)$ to series expansion for $m = 1..2$

$$u^m \cdot \left(\frac{d^2}{du^2} y(u) \right) = \sum_{k=0}^{\infty} a_k (k+r)(k+r-1) u^{k+r-2+m}$$

- Shift index using $k \rightarrow k+2-m$

$$u^m \cdot \left(\frac{d^2}{du^2} y(u) \right) = \sum_{k=-2+m}^{\infty} a_{k+2-m} (k+2-m+r)(k+1-m+r) u^{k+r}$$

Rewrite ODE with series expansions

$$-2a_0 r^2 u^{-1+r} + \left(\sum_{k=0}^{\infty} (-2a_{k+1} (k+1+r)^2 + a_k (r+1+p+k)(r-p+k)) u^{k+r} \right) = 0$$

- a_0 cannot be 0 by assumption, giving the indicial equation

$$-2r^2 = 0$$

- Values of r that satisfy the indicial equation

$$r = 0$$

- Each term in the series must be 0, giving the recursion relation

$$-2a_{k+1} (k+1)^2 + a_k (1+p+k)(-p+k) = 0$$

- Recursion relation that defines series solution to ODE

$$a_{k+1} = \frac{a_k(1+p+k)(-p+k)}{2(k+1)^2}$$

- Recursion relation for $r = 0$

$$a_{k+1} = \frac{a_k(1+p+k)(-p+k)}{2(k+1)^2}$$

- Solution for $r = 0$

$$\left[y(u) = \sum_{k=0}^{\infty} a_k u^k, a_{k+1} = \frac{a_k(1+p+k)(-p+k)}{2(k+1)^2} \right]$$

- Revert the change of variables $u = 1 + x$

$$\left[y = \sum_{k=0}^{\infty} a_k (1+x)^k, a_{k+1} = \frac{a_k(1+p+k)(-p+k)}{2(k+1)^2} \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exists
-> Trying a solution in terms of special functions:
    -> Bessel
    -> elliptic
    -> Legendre
    <- Legendre successful
<- special function solution successful`

```

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 1124

```

Order:=8;
dsolve((1-x^2)*diff(y(x),x$2)-2*x*diff(y(x),x)+p*(p+1)*y(x)=0,y(x),type='series',x=infinity)

```

Expression too large to display

✓ Solution by Mathematica

Time used: 0.01 (sec). Leaf size: 2707

AsymptoticDSolveValue[(1-x^2)*y''[x]-2*x*y'[x]+p*(p+1)*y[x]==0,y[x],{x,infinity,7}]

$$\begin{aligned}
 y(x) \rightarrow & \left(\frac{p^2 x^{-p-7}}{-p^2 - p + (p+6)(p+7)} + \frac{3px^{-p-7}}{-p^2 - p + (p+6)(p+7)} \right. \\
 & + \frac{p^4 x^{-p-7}}{(-p^2 - p + (p+2)(p+3))(-p^2 - p + (p+6)(p+7))} \\
 & + \frac{6p^3 x^{-p-7}}{(-p^2 - p + (p+2)(p+3))(-p^2 - p + (p+6)(p+7))} \\
 & + \frac{17p^2 x^{-p-7}}{(-p^2 - p + (p+2)(p+3))(-p^2 - p + (p+6)(p+7))} \\
 & + \frac{24px^{-p-7}}{(-p^2 - p + (p+2)(p+3))(-p^2 - p + (p+6)(p+7))} \\
 & + \frac{12x^{-p-7}}{(-p^2 - p + (p+2)(p+3))(-p^2 - p + (p+6)(p+7))} \\
 & + \frac{p^4 x^{-p-7}}{(-p^2 - p + (p+4)(p+5))(-p^2 - p + (p+6)(p+7))} \\
 & + \frac{6p^3 x^{-p-7}}{(-p^2 - p + (p+4)(p+5))(-p^2 - p + (p+6)(p+7))} \\
 & + \frac{21p^2 x^{-p-7}}{(-p^2 - p + (p+4)(p+5))(-p^2 - p + (p+6)(p+7))} \\
 & + \frac{36px^{-p-7}}{(-p^2 - p + (p+4)(p+5))(-p^2 - p + (p+6)(p+7))} \\
 & + \frac{p^6 x^{-p-7}}{(-p^2 - p + (p+2)(p+3))(-p^2 - p + (p+4)(p+5))(-p^2 - p + (p+6)(p+7))} \\
 & + \frac{9p^5 x^{-p-7}}{(-p^2 - p + (p+2)(p+3))(-p^2 - p + (p+4)(p+5))(-p^2 - p + (p+6)(p+7))} \\
 & + \frac{45p^4 x^{-p-7}}{(-p^2 - p + (p+2)(p+3))(-p^2 - p + (p+4)(p+5))(-p^2 - p + (p+6)(p+7))} \\
 & + \frac{135p^3 x^{-p-7}}{(-p^2 - p + (p+2)(p+3))(-p^2 - p + (p+4)(p+5))(-p^2 - p + (p+6)(p+7))} \\
 & + \frac{254p^2 x^{-p-7}}{(-p^2 - p + (p+2)(p+3))(-p^2 - p + (p+4)(p+5))(-p^2 - p + (p+6)(p+7))} \\
 & + \frac{276px^{-p-7}}{(-p^2 - p + (p+2)(p+3))(-p^2 - p + (p+4)(p+5))(-p^2 - p + (p+6)(p+7))} \\
 & + \frac{120x^{-p-7}}{(-p^2 - p + (p+2)(p+3))(-p^2 - p + (p+4)(p+5))(-p^2 - p + (p+6)(p+7))} \\
 & + \frac{20x^{-p-7}}{(-p^2 - p + (p+2)(p+3))(-p^2 - p + (p+4)(p+5))(-p^2 - p + (p+6)(p+7))} \\
 & + \frac{3852}{p^2 x^{-p-5}} + \frac{3px^{-p-5}}{-p^2 - p + (p+4)(p+5)} \\
 & + \frac{2x^{-p-7}}{-p^2 - p + (p+6)(p+7)} + \frac{2x^{-p-7}}{-p^2 - p + (p+4)(p+5)} + \frac{2x^{-p-7}}{-p^2 - p + (p+4)(p+5)}
 \end{aligned}$$

**24 Chapter 7. Laplace Transforms. Section 7.5 The
Unit Step and Impulse Functions. Page 303**

24.1	problem 1(a)	3854
24.2	problem 1(b)	3860
24.3	problem 1(c)	3866
24.4	problem 7(a)	3872
24.5	problem 7(b)	3877
24.6	problem 7(c)	3882

24.1 problem 1(a)

24.1.1 Existence and uniqueness analysis	3854
24.1.2 Maple step by step solution	3857

Internal problem ID [6499]

Internal file name [OUTPUT/5747_Sunday_June_05_2022_03_52_36_PM_71358093/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 7. Laplace Transforms. Section 7.5 The Unit Step and Impulse Functions. Page 303

Problem number: 1(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_laplace", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' + 5y' + 6y = 5e^{3t}$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

24.1.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(t)y' + q(t)y = F$$

Where here

$$p(t) = 5$$

$$q(t) = 6$$

$$F = 5e^{3t}$$

Hence the ode is

$$y'' + 5y' + 6y = 5e^{3t}$$

The domain of $p(t) = 5$ is

$$\{-\infty < t < \infty\}$$

And the point $t_0 = 0$ is inside this domain. The domain of $q(t) = 6$ is

$$\{-\infty < t < \infty\}$$

And the point $t_0 = 0$ is also inside this domain. The domain of $F = 5e^{3t}$ is

$$\{-\infty < t < \infty\}$$

And the point $t_0 = 0$ is also inside this domain. Hence solution exists and is unique.

Solving using the Laplace transform method. Let

$$\mathcal{L}(y) = Y(s)$$

Taking the Laplace transform of the ode and using the relations that

$$\begin{aligned}\mathcal{L}(y') &= sY(s) - y(0) \\ \mathcal{L}(y'') &= s^2Y(s) - y'(0) - sy(0)\end{aligned}$$

The given ode now becomes an algebraic equation in the Laplace domain

$$s^2Y(s) - y'(0) - sy(0) + 5sY(s) - 5y(0) + 6Y(s) = \frac{5}{s-3} \quad (1)$$

But the initial conditions are

$$\begin{aligned}y(0) &= 0 \\ y'(0) &= 0\end{aligned}$$

Substituting these initial conditions in above in Eq (1) gives

$$s^2Y(s) + 5sY(s) + 6Y(s) = \frac{5}{s-3}$$

Solving the above equation for $Y(s)$ results in

$$Y(s) = \frac{5}{(s-3)(s^2+5s+6)}$$

Applying partial fractions decomposition results in

$$Y(s) = \frac{5}{6(s+3)} - \frac{1}{s+2} + \frac{1}{6s-18}$$

The inverse Laplace of each term above is now found, which gives

$$\mathcal{L}^{-1}\left(\frac{5}{6(s+3)}\right) = \frac{5e^{-3t}}{6}$$

$$\mathcal{L}^{-1}\left(-\frac{1}{s+2}\right) = -e^{-2t}$$

$$\mathcal{L}^{-1}\left(\frac{1}{6s-18}\right) = \frac{e^{3t}}{6}$$

Adding the above results and simplifying gives

$$y = \cosh(3t) - \frac{2 \sinh(3t)}{3} - e^{-2t}$$


Simplifying the solution gives

$$y = \cosh(3t) - \frac{2 \sinh(3t)}{3} - e^{-2t}$$


Summary

The solution(s) found are the following

$$y = \cosh(3t) - \frac{2 \sinh(3t)}{3} - e^{-2t} \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \cosh(3t) - \frac{2 \sinh(3t)}{3} - e^{-2t}$$

Verified OK.

24.1.2 Maple step by step solution

Let's solve

$$\left[y'' + 5y' + 6y = 5e^{3t}, y(0) = 0, y' \Big|_{\{t=0\}} = 0 \right]$$

- Highest derivative means the order of the ODE is 2

y''

- Characteristic polynomial of homogeneous ODE

$$r^2 + 5r + 6 = 0$$

- Factor the characteristic polynomial

$$(r + 3)(r + 2) = 0$$

- Roots of the characteristic polynomial

$$r = (-3, -2)$$

- 1st solution of the homogeneous ODE

$$y_1(t) = e^{-3t}$$

- 2nd solution of the homogeneous ODE

$$y_2(t) = e^{-2t}$$

- General solution of the ODE

$$y = c_1 y_1(t) + c_2 y_2(t) + y_p(t)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{-3t} + c_2 e^{-2t} + y_p(t)$$

- Find a particular solution $y_p(t)$ of the ODE

- Use variation of parameters to find y_p here $f(t)$ is the forcing function

$$\left[y_p(t) = -y_1(t) \left(\int \frac{y_2(t)f(t)}{W(y_1(t), y_2(t))} dt \right) + y_2(t) \left(\int \frac{y_1(t)f(t)}{W(y_1(t), y_2(t))} dt \right), f(t) = 5e^{3t} \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(t), y_2(t)) = \begin{bmatrix} e^{-3t} & e^{-2t} \\ -3e^{-3t} & -2e^{-2t} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(t), y_2(t)) = e^{-5t}$$

- Substitute functions into equation for $y_p(t)$

$$y_p(t) = -5e^{-3t} \left(\int e^{6t} dt \right) + 5e^{-2t} \left(\int e^{5t} dt \right)$$

- Compute integrals

$$y_p(t) = \frac{e^{3t}}{6}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{-3t} + c_2 e^{-2t} + \frac{e^{3t}}{6}$$

- Check validity of solution $y = c_1 e^{-3t} + c_2 e^{-2t} + \frac{e^{3t}}{6}$

- Use initial condition $y(0) = 0$

$$0 = c_1 + c_2 + \frac{1}{6}$$

- Compute derivative of the solution

$$y' = -3c_1 e^{-3t} - 2c_2 e^{-2t} + \frac{e^{3t}}{2}$$

- Use the initial condition $y' \Big|_{\{t=0\}} = 0$

$$0 = -3c_1 - 2c_2 + \frac{1}{2}$$

- Solve for c_1 and c_2

$$\left\{ c_1 = \frac{5}{6}, c_2 = -1 \right\}$$

- Substitute constant values into general solution and simplify

$$y = \frac{(e^{6t} - 6e^t + 5)e^{-3t}}{6}$$

- Solution to the IVP

$$y = \frac{(e^{6t} - 6e^t + 5)e^{-3t}}{6}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 1.859 (sec). Leaf size: 21

```
dsolve([diff(y(t),t$2)+5*diff(y(t),t)+6*y(t)=5*exp(3*t),y(0) = 0, D(y)(0) = 0],y(t), singsol
```

$$y(t) = \cosh(3t) - \frac{2 \sinh(3t)}{3} - e^{-2t}$$

✓ Solution by Mathematica

Time used: 0.019 (sec). Leaf size: 26

```
DSolve[{y''[t]+5*y'[t]+6*y[t]==5*Exp[3*t],{y[0]==0,y'[0]==0}},y[t],t,IncludeSingularSolution
```

$$y(t) \rightarrow \frac{1}{6}e^{-3t}(-6e^t + e^{6t} + 5)$$

24.2 problem 1(b)

24.2.1 Existence and uniqueness analysis	3860
24.2.2 Maple step by step solution	3863

Internal problem ID [6500]

Internal file name [OUTPUT/5748_Sunday_June_05_2022_03_52_38_PM_12557919/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 7. Laplace Transforms. Section 7.5 The Unit Step and Impulse Functions. Page 303

Problem number: 1(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_laplace", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' + y' - 6y = t$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

24.2.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(t)y' + q(t)y = F$$

Where here

$$p(t) = 1$$

$$q(t) = -6$$

$$F = t$$

Hence the ode is

$$y'' + y' - 6y = t$$

The domain of $p(t) = 1$ is

$$\{-\infty < t < \infty\}$$

And the point $t_0 = 0$ is inside this domain. The domain of $q(t) = -6$ is

$$\{-\infty < t < \infty\}$$

And the point $t_0 = 0$ is also inside this domain. The domain of $F = t$ is

$$\{-\infty < t < \infty\}$$

And the point $t_0 = 0$ is also inside this domain. Hence solution exists and is unique.

Solving using the Laplace transform method. Let

$$\mathcal{L}(y) = Y(s)$$

Taking the Laplace transform of the ode and using the relations that

$$\begin{aligned}\mathcal{L}(y') &= sY(s) - y(0) \\ \mathcal{L}(y'') &= s^2Y(s) - y'(0) - sy(0)\end{aligned}$$

The given ode now becomes an algebraic equation in the Laplace domain

$$s^2Y(s) - y'(0) - sy(0) + sY(s) - y(0) - 6Y(s) = \frac{1}{s^2} \quad (1)$$

But the initial conditions are

$$\begin{aligned}y(0) &= 0 \\ y'(0) &= 0\end{aligned}$$

Substituting these initial conditions in above in Eq (1) gives

$$s^2Y(s) + sY(s) - 6Y(s) = \frac{1}{s^2}$$

Solving the above equation for $Y(s)$ results in

$$Y(s) = \frac{1}{s^2(s^2 + s - 6)}$$

Applying partial fractions decomposition results in

$$Y(s) = -\frac{1}{36s} - \frac{1}{45(s+3)} + \frac{1}{20s-40} - \frac{1}{6s^2}$$

The inverse Laplace of each term above is now found, which gives

$$\begin{aligned}\mathcal{L}^{-1}\left(-\frac{1}{36s}\right) &= -\frac{1}{36} \\ \mathcal{L}^{-1}\left(-\frac{1}{45(s+3)}\right) &= -\frac{e^{-3t}}{45} \\ \mathcal{L}^{-1}\left(\frac{1}{20s-40}\right) &= \frac{e^{2t}}{20} \\ \mathcal{L}^{-1}\left(-\frac{1}{6s^2}\right) &= -\frac{t}{6}\end{aligned}$$

Adding the above results and simplifying gives

$$y = \frac{e^{2t}}{20} - \frac{1}{36} - \frac{e^{-3t}}{45} - \frac{t}{6}$$


Simplifying the solution gives

$$y = \frac{(9e^{5t} - 30te^{3t} - 5e^{3t} - 4)e^{-3t}}{180}$$


Summary

The solution(s) found are the following

$$y = \frac{(9e^{5t} - 30te^{3t} - 5e^{3t} - 4)e^{-3t}}{180} \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{(9e^{5t} - 30te^{3t} - 5e^{3t} - 4)e^{-3t}}{180}$$

Verified OK.

24.2.2 Maple step by step solution

Let's solve

$$\left[y'' + y' - 6y = t, y(0) = 0, y'|_{\{t=0\}} = 0 \right]$$

- Highest derivative means the order of the ODE is 2
 y''
- Characteristic polynomial of homogeneous ODE
 $r^2 + r - 6 = 0$
- Factor the characteristic polynomial
 $(r + 3)(r - 2) = 0$
- Roots of the characteristic polynomial
 $r = (-3, 2)$
- 1st solution of the homogeneous ODE

$$y_1(t) = e^{-3t}$$

- 2nd solution of the homogeneous ODE

$$y_2(t) = e^{2t}$$

- General solution of the ODE

$$y = c_1 y_1(t) + c_2 y_2(t) + y_p(t)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{-3t} + c_2 e^{2t} + y_p(t)$$

- Find a particular solution $y_p(t)$ of the ODE

- Use variation of parameters to find y_p here $f(t)$ is the forcing function

$$\left[y_p(t) = -y_1(t) \left(\int \frac{y_2(t)f(t)}{W(y_1(t), y_2(t))} dt \right) + y_2(t) \left(\int \frac{y_1(t)f(t)}{W(y_1(t), y_2(t))} dt \right), f(t) = t \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(t), y_2(t)) = \begin{bmatrix} e^{-3t} & e^{2t} \\ -3e^{-3t} & 2e^{2t} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(t), y_2(t)) = 5e^{-t}$$

- Substitute functions into equation for $y_p(t)$

$$y_p(t) = \frac{(e^{5t} (\int t e^{-2t} dt) - (\int t e^{3t} dt)) e^{-3t}}{5}$$

- Compute integrals

$$y_p(t) = -\frac{1}{36} - \frac{t}{6}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{-3t} + c_2 e^{2t} - \frac{1}{36} - \frac{t}{6}$$

- Check validity of solution $y = c_1 e^{-3t} + c_2 e^{2t} - \frac{1}{36} - \frac{t}{6}$

- Use initial condition $y(0) = 0$

$$0 = c_1 + c_2 - \frac{1}{36}$$

- Compute derivative of the solution

$$y' = -3c_1 e^{-3t} + 2c_2 e^{2t} - \frac{1}{6}$$

- Use the initial condition $y' \Big|_{\{t=0\}} = 0$

$$0 = -3c_1 + 2c_2 - \frac{1}{6}$$

- Solve for c_1 and c_2

$$\left\{ c_1 = -\frac{1}{45}, c_2 = \frac{1}{20} \right\}$$
- Substitute constant values into general solution and simplify
$$y = \frac{(9e^{5t} - 30te^{3t} - 5e^{3t} - 4)e^{-3t}}{180}$$
- Solution to the IVP
$$y = \frac{(9e^{5t} - 30te^{3t} - 5e^{3t} - 4)e^{-3t}}{180}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 1.531 (sec). Leaf size: 21

```
dsolve([diff(y(t),t$2)+diff(y(t),t)-6*y(t)=t,y(0) = 0, D(y)(0) = 0],y(t), singsol=all)
```

$$y(t) = \frac{(9e^{5t} - 30te^{3t} - 5e^{3t} - 4)e^{-3t}}{180}$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 28

```
DSolve[{y''[t]+y'[t]-6*y[t]==t,{y[0]==0,y'[0]==0}},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{1}{180}(-30t - 4e^{-3t} + 9e^{2t} - 5)$$

24.3 problem 1(c)

24.3.1 Existence and uniqueness analysis	3866
24.3.2 Maple step by step solution	3869

Internal problem ID [6501]

Internal file name [OUTPUT/5749_Sunday_June_05_2022_03_52_40_PM_91277379/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 7. Laplace Transforms. Section 7.5 The Unit Step and Impulse Functions. Page 303

Problem number: 1(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_laplace", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - y = t^2$$

With initial conditions

$$[y(0) = 0, y'(0) = 0]$$

24.3.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(t)y' + q(t)y = F$$

Where here

$$p(t) = 0$$

$$q(t) = -1$$

$$F = t^2$$

Hence the ode is

$$y'' - y = t^2$$

The domain of $p(t) = 0$ is

$$\{-\infty < t < \infty\}$$

And the point $t_0 = 0$ is inside this domain. The domain of $q(t) = -1$ is

$$\{-\infty < t < \infty\}$$

And the point $t_0 = 0$ is also inside this domain. The domain of $F = t^2$ is

$$\{-\infty < t < \infty\}$$

And the point $t_0 = 0$ is also inside this domain. Hence solution exists and is unique.

Solving using the Laplace transform method. Let

$$\mathcal{L}(y) = Y(s)$$

Taking the Laplace transform of the ode and using the relations that

$$\mathcal{L}(y') = sY(s) - y(0)$$

$$\mathcal{L}(y'') = s^2Y(s) - y'(0) - sy(0)$$

The given ode now becomes an algebraic equation in the Laplace domain

$$s^2Y(s) - y'(0) - sy(0) - Y(s) = \frac{2}{s^3} \quad (1)$$

But the initial conditions are

$$y(0) = 0$$

$$y'(0) = 0$$

Substituting these initial conditions in above in Eq (1) gives

$$s^2Y(s) - Y(s) = \frac{2}{s^3}$$

Solving the above equation for $Y(s)$ results in

$$Y(s) = \frac{2}{s^3(s^2 - 1)}$$

Applying partial fractions decomposition results in

$$Y(s) = \frac{1}{s+1} + \frac{1}{s-1} - \frac{2}{s^3} - \frac{2}{s}$$

The inverse Laplace of each term above is now found, which gives

$$\mathcal{L}^{-1}\left(\frac{1}{s+1}\right) = e^{-t}$$

$$\mathcal{L}^{-1}\left(\frac{1}{s-1}\right) = e^t$$

$$\mathcal{L}^{-1}\left(-\frac{2}{s^3}\right) = -t^2$$

$$\mathcal{L}^{-1}\left(-\frac{2}{s}\right) = -2$$

Adding the above results and simplifying gives

$$y = -2 - t^2 + 2 \cosh(t)$$


Simplifying the solution gives

$$y = -2 - t^2 + 2 \cosh(t)$$


Summary

The solution(s) found are the following

$$y = -2 - t^2 + 2 \cosh(t) \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = -2 - t^2 + 2 \cosh(t)$$

Verified OK.

24.3.2 Maple step by step solution

Let's solve

$$\left[y'' - y = t^2, y(0) = 0, y'|_{\{t=0\}} = 0 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - 1 = 0$$

- Factor the characteristic polynomial

$$(r - 1)(r + 1) = 0$$

- Roots of the characteristic polynomial

$$r = (-1, 1)$$

- 1st solution of the homogeneous ODE

$$y_1(t) = e^{-t}$$

- 2nd solution of the homogeneous ODE

$$y_2(t) = e^t$$

- General solution of the ODE

$$y = c_1 y_1(t) + c_2 y_2(t) + y_p(t)$$

- Substitute in solutions of the homogeneous ODE

$$y = e^{-t} c_1 + c_2 e^t + y_p(t)$$

- Find a particular solution $y_p(t)$ of the ODE

- Use variation of parameters to find y_p here $f(t)$ is the forcing function

$$\left[y_p(t) = -y_1(t) \left(\int \frac{y_2(t)f(t)}{W(y_1(t), y_2(t))} dt \right) + y_2(t) \left(\int \frac{y_1(t)f(t)}{W(y_1(t), y_2(t))} dt \right), f(t) = t^2 \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(t), y_2(t)) = \begin{bmatrix} e^{-t} & e^t \\ -e^{-t} & e^t \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(t), y_2(t)) = 2$$

- Substitute functions into equation for $y_p(t)$

$$y_p(t) = -\frac{e^{-t}(\int e^t t^2 dt)}{2} + \frac{e^t(\int e^{-t} t^2 dt)}{2}$$

- Compute integrals

$$y_p(t) = -t^2 - 2$$

- Substitute particular solution into general solution to ODE

$$y = e^{-t}c_1 + c_2e^t - t^2 - 2$$

- Check validity of solution $y = e^{-t}c_1 + c_2e^t - t^2 - 2$

- Use initial condition $y(0) = 0$

$$0 = c_1 - 2 + c_2$$

- Compute derivative of the solution

$$y' = -e^{-t}c_1 + c_2e^t - 2t$$

- Use the initial condition $y'|_{\{t=0\}} = 0$

$$0 = -c_1 + c_2$$

- Solve for c_1 and c_2

$$\{c_1 = 1, c_2 = 1\}$$

- Substitute constant values into general solution and simplify

$$y = e^{-t} + e^t - t^2 - 2$$

- Solution to the IVP

$$y = e^{-t} + e^t - t^2 - 2$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 1.516 (sec). Leaf size: 15

```
dsolve([diff(y(t),t$2)-y(t)=t^2,y(0) = 0, D(y)(0) = 0],y(t), singsol=all)
```

$$y(t) = -2 - t^2 + 2 \cosh(t)$$

✓ Solution by Mathematica

Time used: 0.015 (sec). Leaf size: 20

```
DSolve[{y''[t]-y[t]==t^2,{y[0]==0,y'[0]==0}},y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow -t^2 + e^{-t} + e^t - 2$$

24.4 problem 7(a)

24.4.1 Existence and uniqueness analysis	3872
24.4.2 Solving as laplace ode	3873
24.4.3 Maple step by step solution	3874

Internal problem ID [6502]

Internal file name [OUTPUT/5750_Sunday_June_05_2022_03_52_42_PM_3334797/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 7. Laplace Transforms. Section 7.5 The Unit Step and Impulse Functions. Page 303

Problem number: 7(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "linear", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

```
[[_linear, `class A`]]
```

$$Li' + Ri = E_0 \text{Heaviside}(t)$$

With initial conditions

$$[i(0) = 0]$$

24.4.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$i' + p(t)i = q(t)$$

Where here

$$p(t) = \frac{R}{L}$$
$$q(t) = \frac{E_0 \text{Heaviside}(t)}{L}$$

Hence the ode is

$$i' + \frac{Ri}{L} = \frac{E_0 \text{Heaviside}(t)}{L}$$

The domain of $p(t) = \frac{R}{L}$ is

$$\{-\infty < t < \infty\}$$

And the point $t_0 = 0$ is inside this domain. The domain of $q(t) = \frac{E_0 \text{Heaviside}(t)}{L}$ is

$$\{t < 0 \vee 0 < t\}$$

But the point $t_0 = 0$ is not inside this domain. Hence existence and uniqueness theorem does not apply. There could be infinite number of solutions, or one solution or no solution at all.

24.4.2 Solving as laplace ode

Solving using the Laplace transform method. Let

$$\mathcal{L}(i) = Y(s)$$

Taking the Laplace transform of the ode and using the relations that

$$\mathcal{L}(i') = sY(s) - i(0)$$

The given ode now becomes an algebraic equation in the Laplace domain

$$LY(s) s - Li(0) + RY(s) = \frac{E_0}{s} \tag{1}$$

Replacing initial condition gives

$$LY(s) s + RY(s) = \frac{E_0}{s}$$

Solving for $Y(s)$ gives

$$Y(s) = \frac{E_0}{s(Ls + R)}$$

Applying partial fractions decomposition results in

$$Y(s) = -\frac{E_0}{R(s + \frac{R}{L})} + \frac{E_0}{Rs}$$

The inverse Laplace of each term above is now found, which gives

$$\mathcal{L}^{-1}\left(-\frac{E_0}{R\left(s + \frac{R}{L}\right)}\right) = -\frac{E_0 e^{-\frac{Rt}{L}}}{R}$$

$$\mathcal{L}^{-1}\left(\frac{E_0}{Rs}\right) = \frac{E_0}{R}$$

Adding the above results and simplifying gives

$$i = \frac{E_0\left(1 - e^{-\frac{Rt}{L}}\right)}{R}$$

Summary

The solution(s) found are the following

$$i = \frac{E_0\left(1 - e^{-\frac{Rt}{L}}\right)}{R} \quad (1)$$

Verification of solutions

$$i = \frac{E_0\left(1 - e^{-\frac{Rt}{L}}\right)}{R}$$

Verified OK.

24.4.3 Maple step by step solution

Let's solve

$$[Li' + Ri = E_0 \text{Heaviside}(t), i(0) = 0]$$

- Highest derivative means the order of the ODE is 1

$$i'$$

- Isolate the derivative

$$i' = -\frac{Ri}{L} + \frac{E_0 \text{Heaviside}(t)}{L}$$

- Group terms with i on the lhs of the ODE and the rest on the rhs of the ODE

$$i' + \frac{Ri}{L} = \frac{E_0 \text{Heaviside}(t)}{L}$$

- The ODE is linear; multiply by an integrating factor $\mu(t)$

$$\mu(t) \left(i' + \frac{Ri}{L}\right) = \frac{\mu(t)E_0 \text{Heaviside}(t)}{L}$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dt}(\mu(t) i)$

$$\mu(t) \left(i' + \frac{Ri}{L} \right) = \mu'(t) i + \mu(t) i'$$

- Isolate $\mu'(t)$

$$\mu'(t) = \frac{\mu(t)R}{L}$$

- Solve to find the integrating factor

$$\mu(t) = e^{\frac{Rt}{L}}$$

- Integrate both sides with respect to t

$$\int \left(\frac{d}{dt} (\mu(t) i) \right) dt = \int \frac{\mu(t) E_0 \text{Heaviside}(t)}{L} dt + c_1$$

- Evaluate the integral on the lhs

$$\mu(t) i = \int \frac{\mu(t) E_0 \text{Heaviside}(t)}{L} dt + c_1$$

- Solve for i

$$i = \frac{\int \frac{\mu(t) E_0 \text{Heaviside}(t)}{L} dt + c_1}{\mu(t)}$$

- Substitute $\mu(t) = e^{\frac{Rt}{L}}$

$$i = \frac{\int \frac{e^{\frac{Rt}{L}} E_0 \text{Heaviside}(t)}{L} dt + c_1}{e^{\frac{Rt}{L}}}$$

- Evaluate the integrals on the rhs

$$i = \frac{E_0 \left(\frac{L e^{\frac{Rt}{L}} \text{Heaviside}(t)}{R} - \frac{\text{Heaviside}(t)L}{R} \right) + c_1}{e^{\frac{Rt}{L}}}$$

- Simplify

$$i = \frac{E_0 \text{Heaviside}(t) - E_0 \text{Heaviside}(t) e^{-\frac{Rt}{L}} + Rc_1 e^{-\frac{Rt}{L}}}{R}$$

- Use initial condition $i(0) = 0$

$$0 = c_1$$

- Solve for c_1

$$c_1 = 0$$

- Substitute $c_1 = 0$ into general solution and simplify

$$i = - \frac{E_0 \text{Heaviside}(t) \left(e^{-\frac{Rt}{L}} - 1 \right)}{R}$$

- Solution to the IVP

$$i = - \frac{E_0 \text{Heaviside}(t) \left(e^{-\frac{Rt}{L}} - 1 \right)}{R}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 1.625 (sec). Leaf size: 21

```
dsolve([L*diff(i(t),t)+R*i(t)=E_0*Heaviside(t),i(0) = 0],i(t), singsol=all)
```

$$i(t) = -\frac{E_0 \left(e^{-\frac{Rt}{L}} - 1 \right)}{R}$$

✓ Solution by Mathematica

Time used: 0.077 (sec). Leaf size: 25

```
DSolve[{L*i'[t]+R*i[t]==E0*UnitStep[t],{i[0]==0}},i[t],t,IncludeSingularSolutions -> True]
```

$$i(t) \rightarrow \frac{E_0 \theta(t) \left(1 - e^{-\frac{Rt}{L}} \right)}{R}$$

24.5 problem 7(b)

24.5.1 Existence and uniqueness analysis	3877
24.5.2 Solving as laplace ode	3878
24.5.3 Maple step by step solution	3879

Internal problem ID [6503]

Internal file name [OUTPUT/5751_Sunday_June_05_2022_03_52_45_PM_9236993/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 7. Laplace Transforms. Section 7.5 The Unit Step and Impulse Functions. Page 303

Problem number: 7(b).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "linear", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

```
[[_linear, `class A`]]
```

$$Li' + Ri = E_0\delta(t)$$

With initial conditions

$$[i(0) = 0]$$

24.5.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$i' + p(t)i = q(t)$$

Where here

$$p(t) = \frac{R}{L}$$
$$q(t) = \frac{E_0\delta(t)}{L}$$

Hence the ode is

$$i' + \frac{Ri}{L} = \frac{E_0\delta(t)}{L}$$

The domain of $p(t) = \frac{R}{L}$ is

$$\{-\infty < t < \infty\}$$

And the point $t_0 = 0$ is inside this domain. The domain of $q(t) = \frac{E_0\delta(t)}{L}$ is

$$\{t < 0 \vee 0 < t\}$$

But the point $t_0 = 0$ is not inside this domain. Hence existence and uniqueness theorem does not apply. There could be infinite number of solutions, or one solution or no solution at all.

24.5.2 Solving as laplace ode

Solving using the Laplace transform method. Let

$$\mathcal{L}(i) = Y(s)$$

Taking the Laplace transform of the ode and using the relations that

$$\mathcal{L}(i') = sY(s) - i(0)$$

The given ode now becomes an algebraic equation in the Laplace domain

$$LY(s)s - Li(0) + RY(s) = E_0 \tag{1}$$

Replacing initial condition gives

$$LY(s)s + RY(s) = E_0$$

Solving for $Y(s)$ gives

$$Y(s) = \frac{E_0}{Ls + R}$$

Taking inverse Laplace transform gives

$$\mathcal{L}^{-1}\left(\frac{E_0}{Ls + R}\right) = \frac{E_0 e^{-\frac{Rt}{L}}}{L}$$

Summary

The solution(s) found are the following

$$i = \frac{E_0 e^{-\frac{Rt}{L}}}{L} \quad (1)$$

Verification of solutions

$$i = \frac{E_0 e^{-\frac{Rt}{L}}}{L}$$

Verified OK.

24.5.3 Maple step by step solution

Let's solve

$$[Li' + Ri = E_0 \text{Dirac}(t), i(0) = 0]$$

- Highest derivative means the order of the ODE is 1

$$i'$$

- Isolate the derivative

$$i' = -\frac{Ri}{L} + \frac{E_0 \text{Dirac}(t)}{L}$$

- Group terms with i on the lhs of the ODE and the rest on the rhs of the ODE

$$i' + \frac{Ri}{L} = \frac{E_0 \text{Dirac}(t)}{L}$$

- The ODE is linear; multiply by an integrating factor $\mu(t)$

$$\mu(t) \left(i' + \frac{Ri}{L} \right) = \frac{\mu(t) E_0 \text{Dirac}(t)}{L}$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dt}(\mu(t) i)$

$$\mu(t) \left(i' + \frac{Ri}{L} \right) = \mu'(t) i + \mu(t) i'$$

- Isolate $\mu'(t)$

$$\mu'(t) = \frac{\mu(t) R}{L}$$

- Solve to find the integrating factor

$$\mu(t) = e^{\frac{Rt}{L}}$$

- Integrate both sides with respect to t

$$\int \left(\frac{d}{dt}(\mu(t) i) \right) dt = \int \frac{\mu(t) E_0 \text{Dirac}(t)}{L} dt + c_1$$

- Evaluate the integral on the lhs

$$\mu(t) i = \int \frac{\mu(t) E_0 \text{Dirac}(t)}{L} dt + c_1$$

- Solve for i

$$i = \frac{\int \frac{\mu(t)E_0 \text{Dirac}(t)}{L} dt + c_1}{\mu(t)}$$

- Substitute $\mu(t) = e^{\frac{Rt}{L}}$

$$i = \frac{\int \frac{e^{-\frac{Rt}{L}} E_0 \text{Dirac}(t)}{L} dt + c_1}{e^{\frac{Rt}{L}}}$$

- Evaluate the integrals on the rhs

$$i = \frac{\frac{E_0 \text{Heaviside}(t)}{L} + c_1}{e^{\frac{Rt}{L}}}$$

- Simplify

$$i = \frac{e^{-\frac{Rt}{L}} (E_0 \text{Heaviside}(t) + Lc_1)}{L}$$

- Use initial condition $i(0) = 0$

$$0 = \frac{E_0 \text{Heaviside}(0) + Lc_1}{L}$$

- Solve for c_1

$$c_1 = -\frac{E_0 \text{Heaviside}(0)}{L}$$

- Substitute $c_1 = -\frac{E_0 \text{Heaviside}(0)}{L}$ into general solution and simplify

$$i = \frac{e^{-\frac{Rt}{L}} E_0 (\text{Heaviside}(t) - \text{Heaviside}(0))}{L}$$

- Solution to the IVP

$$i = \frac{e^{-\frac{Rt}{L}} E_0 (\text{Heaviside}(t) - \text{Heaviside}(0))}{L}$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 1.562 (sec). Leaf size: 17

```
dsolve([L*diff(i(t),t)+R*i(t)=E_0*Dirac(t),i(0) = 0],i(t), singsol=all)
```

$$i(t) = \frac{e^{-\frac{Rt}{L}} E_0}{L}$$

✓ Solution by Mathematica

Time used: 0.041 (sec). Leaf size: 26

```
DSolve[{L*i'[t]+R*i[t]==E0*DiracDelta[t],{i[0]==0}},i[t],t,IncludeSingularSolutions -> True]
```

$$i(t) \rightarrow \frac{E_0(\theta(t) - \theta(0))e^{-\frac{Rt}{L}}}{L}$$

24.6 problem 7(c)

24.6.1 Existence and uniqueness analysis	3882
24.6.2 Solving as laplace ode	3883
24.6.3 Maple step by step solution	3884

Internal problem ID [6504]

Internal file name [OUTPUT/5752_Sunday_June_05_2022_03_52_48_PM_16266900/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 7. Laplace Transforms. Section 7.5 The Unit Step and Impulse Functions. Page 303

Problem number: 7(c).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "linear", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

```
[[_linear, `class A`]]
```

$$Li' + Ri = E_0 \sin(\omega t)$$

With initial conditions

$$[i(0) = 0]$$

24.6.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$i' + p(t)i = q(t)$$

Where here

$$p(t) = \frac{R}{L}$$
$$q(t) = \frac{E_0 \sin(\omega t)}{L}$$

Hence the ode is

$$i' + \frac{Ri}{L} = \frac{E_0 \sin(\omega t)}{L}$$

The domain of $p(t) = \frac{R}{L}$ is

$$\{-\infty < t < \infty\}$$

And the point $t_0 = 0$ is inside this domain. The domain of $q(t) = \frac{E_0 \sin(\omega t)}{L}$ is

$$\{-\infty < t < \infty\}$$

And the point $t_0 = 0$ is also inside this domain. Hence solution exists and is unique.

24.6.2 Solving as laplace ode

Solving using the Laplace transform method. Let

$$\mathcal{L}(i) = Y(s)$$

Taking the Laplace transform of the ode and using the relations that

$$\mathcal{L}(i') = sY(s) - i(0)$$

The given ode now becomes an algebraic equation in the Laplace domain

$$LY(s) s - Li(0) + RY(s) = \frac{E_0 \omega}{\omega^2 + s^2} \quad (1)$$

Replacing initial condition gives

$$LY(s) s + RY(s) = \frac{E_0 \omega}{\omega^2 + s^2}$$

Solving for $Y(s)$ gives

$$Y(s) = \frac{E_0 \omega}{(\omega^2 + s^2)(Ls + R)}$$

Applying partial fractions decomposition results in

$$Y(s) = \frac{E_0 \omega L}{(\omega^2 L^2 + R^2) \left(s + \frac{R}{L}\right)} - \frac{(L\sqrt{-\omega^2} - R) E_0 \omega}{2(\omega^2 L^2 + R^2) \sqrt{-\omega^2} (s - \sqrt{-\omega^2})} + \frac{(-L\sqrt{-\omega^2} - R) E_0 \omega}{2(\omega^2 L^2 + R^2) \sqrt{-\omega^2} (s + \sqrt{-\omega^2})}$$

The inverse Laplace of each term above is now found, which gives

$$\mathcal{L}^{-1}\left(\frac{E_0\omega L}{(\omega^2 L^2 + R^2)\left(s + \frac{R}{L}\right)}\right) = \frac{E_0\omega L e^{-\frac{Rt}{L}}}{\omega^2 L^2 + R^2}$$

$$\mathcal{L}^{-1}\left(-\frac{(L\sqrt{-\omega^2} - R) E_0\omega}{2(\omega^2 L^2 + R^2)\sqrt{-\omega^2}(s - \sqrt{-\omega^2})}\right) = \frac{E_0 e^{i \operatorname{csgn}(i\omega)\omega t} (-iR \operatorname{csgn}(i\omega) - L\omega)}{2\omega^2 L^2 + 2R^2}$$

$$\mathcal{L}^{-1}\left(\frac{(-L\sqrt{-\omega^2} - R) E_0\omega}{2(\omega^2 L^2 + R^2)\sqrt{-\omega^2}(s + \sqrt{-\omega^2})}\right) = \frac{E_0 e^{-i \operatorname{csgn}(i\omega)\omega t} (iR \operatorname{csgn}(i\omega) - L\omega)}{2\omega^2 L^2 + 2R^2}$$

Adding the above results and simplifying gives

$$i = \frac{E_0 \left(R \sin(\omega t) + \left(-\cos(\omega t) + e^{-\frac{Rt}{L}} \right) \omega L \right)}{\omega^2 L^2 + R^2}$$

Summary

The solution(s) found are the following

$$i = \frac{E_0 \left(R \sin(\omega t) + \left(-\cos(\omega t) + e^{-\frac{Rt}{L}} \right) \omega L \right)}{\omega^2 L^2 + R^2} \quad (1)$$

Verification of solutions

$$i = \frac{E_0 \left(R \sin(\omega t) + \left(-\cos(\omega t) + e^{-\frac{Rt}{L}} \right) \omega L \right)}{\omega^2 L^2 + R^2}$$

Verified OK.

24.6.3 Maple step by step solution

Let's solve

$$[Li' + Ri = E_0 \sin(\omega t), i(0) = 0]$$

- Highest derivative means the order of the ODE is 1

$$i'$$

- Isolate the derivative

$$i' = -\frac{Ri}{L} + \frac{E_0 \sin(\omega t)}{L}$$

- Group terms with i on the lhs of the ODE and the rest on the rhs of the ODE

$$i' + \frac{Ri}{L} = \frac{E_0 \sin(\omega t)}{L}$$

- The ODE is linear; multiply by an integrating factor $\mu(t)$

$$\mu(t) \left(i' + \frac{Ri}{L} \right) = \frac{\mu(t)E_0 \sin(\omega t)}{L}$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dt}(\mu(t) i)$

$$\mu(t) \left(i' + \frac{Ri}{L} \right) = \mu'(t) i + \mu(t) i'$$

- Isolate $\mu'(t)$

$$\mu'(t) = \frac{\mu(t)R}{L}$$

- Solve to find the integrating factor

$$\mu(t) = e^{\frac{Rt}{L}}$$

- Integrate both sides with respect to t

$$\int \left(\frac{d}{dt}(\mu(t) i) \right) dt = \int \frac{\mu(t)E_0 \sin(\omega t)}{L} dt + c_1$$

- Evaluate the integral on the lhs

$$\mu(t) i = \int \frac{\mu(t)E_0 \sin(\omega t)}{L} dt + c_1$$

- Solve for i

$$i = \frac{\int \frac{\mu(t)E_0 \sin(\omega t)}{L} dt + c_1}{\mu(t)}$$

- Substitute $\mu(t) = e^{\frac{Rt}{L}}$

$$i = \frac{\int \frac{e^{\frac{Rt}{L}} E_0 \sin(\omega t)}{L} dt + c_1}{e^{\frac{Rt}{L}}}$$

- Evaluate the integrals on the rhs

$$i = \frac{E_0 \left(-\frac{\omega e^{\frac{Rt}{L}} \cos(\omega t)}{\omega^2 + \frac{R^2}{L^2}} + \frac{R e^{\frac{Rt}{L}} \sin(\omega t)}{L \left(\omega^2 + \frac{R^2}{L^2} \right)} \right) + c_1}{e^{\frac{Rt}{L}}}$$

- Simplify

$$i = \frac{c_1 (\omega^2 L^2 + R^2) e^{-\frac{Rt}{L}} - E_0 (L\omega \cos(\omega t) - R \sin(\omega t))}{\omega^2 L^2 + R^2}$$

- Use initial condition $i(0) = 0$

$$0 = \frac{c_1 (\omega^2 L^2 + R^2) - E_0 \omega L}{\omega^2 L^2 + R^2}$$

- Solve for c_1

$$c_1 = \frac{E_0 \omega L}{\omega^2 L^2 + R^2}$$

- Substitute $c_1 = \frac{E_0 \omega L}{\omega^2 L^2 + R^2}$ into general solution and simplify

$$i = -\frac{E_0 \left(L\omega \cos(\omega t) - \omega L e^{-\frac{Rt}{L}} - R \sin(\omega t) \right)}{\omega^2 L^2 + R^2}$$

- Solution to the IVP

$$i = -\frac{E_0 \left(L\omega \cos(\omega t) - \omega L e^{-\frac{Rt}{L}} - R \sin(\omega t) \right)}{\omega^2 L^2 + R^2}$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 1.75 (sec). Leaf size: 44

```
dsolve([L*diff(i(t),t)+R*i(t)=E_0*sin(omega*t),i(0) = 0],i(t), singsol=all)
```

$$i(t) = \frac{\left(\omega L e^{-\frac{Rt}{L}} - L \cos(\omega t) \omega + \sin(\omega t) R \right) E_0}{\omega^2 L^2 + R^2}$$

✓ Solution by Mathematica

Time used: 0.114 (sec). Leaf size: 47

```
DSolve[{L*i'[t]+R*i[t]==E0*Sin[\[Omega]*t],{i[0]==0}},i[t],t,IncludeSingularSolutions -> True]
```

$$i(t) \rightarrow \frac{E_0 \left(L\omega e^{-\frac{Rt}{L}} - L\omega \cos(t\omega) + R \sin(t\omega) \right)}{L^2\omega^2 + R^2}$$

25 Chapter 7. Laplace Transforms. Section 7.5
Problem for review and discovery. Section A,
Drill exercises. Page 309

25.1	problem 3(a)	3888
25.2	problem 3(b)	3895
25.3	problem 3(c)	3901
25.4	problem 3(d)	3908
25.5	problem 4(a)	3915
25.6	problem 4(b)	3919
25.7	problem 4(c)	3925
25.8	problem 4(d)	3930

25.1 problem 3(a)

25.1.1 Existence and uniqueness analysis	3888
25.1.2 Maple step by step solution	3891

Internal problem ID [6505]

Internal file name [OUTPUT/5753_Sunday_June_05_2022_03_52_50_PM_67993605/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 7. Laplace Transforms. Section 7.5 Problems for review and discovery. Section A, Drill exercises. Page 309

Problem number: 3(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_laplace", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' + 3y' - 5y = 1$$

With initial conditions

$$[y(0) = 0, y'(0) = 1]$$

25.1.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(t)y' + q(t)y = F$$

Where here

$$p(t) = 3$$

$$q(t) = -5$$

$$F = 1$$

Hence the ode is

$$y'' + 3y' - 5y = 1$$

The domain of $p(t) = 3$ is

$$\{-\infty < t < \infty\}$$

And the point $t_0 = 0$ is inside this domain. The domain of $q(t) = -5$ is

$$\{-\infty < t < \infty\}$$

And the point $t_0 = 0$ is also inside this domain. The domain of $F = 1$ is

$$\{-\infty < t < \infty\}$$

And the point $t_0 = 0$ is also inside this domain. Hence solution exists and is unique.

Solving using the Laplace transform method. Let

$$\mathcal{L}(y) = Y(s)$$

Taking the Laplace transform of the ode and using the relations that

$$\begin{aligned}\mathcal{L}(y') &= sY(s) - y(0) \\ \mathcal{L}(y'') &= s^2Y(s) - y'(0) - sy(0)\end{aligned}$$

The given ode now becomes an algebraic equation in the Laplace domain

$$s^2Y(s) - y'(0) - sy(0) + 3sY(s) - 3y(0) - 5Y(s) = \frac{1}{s} \quad (1)$$

But the initial conditions are

$$\begin{aligned}y(0) &= 0 \\ y'(0) &= 1\end{aligned}$$

Substituting these initial conditions in above in Eq (1) gives

$$s^2Y(s) - 1 + 3sY(s) - 5Y(s) = \frac{1}{s}$$

Solving the above equation for $Y(s)$ results in

$$Y(s) = \frac{1 + s}{s(s^2 + 3s - 5)}$$

Applying partial fractions decomposition results in

$$Y(s) = \frac{\frac{1}{10} + \frac{13\sqrt{29}}{290}}{s + \frac{3}{2} - \frac{\sqrt{29}}{2}} + \frac{\frac{1}{10} - \frac{13\sqrt{29}}{290}}{s + \frac{3}{2} + \frac{\sqrt{29}}{2}} - \frac{1}{5s}$$

The inverse Laplace of each term above is now found, which gives

$$\begin{aligned} \mathcal{L}^{-1}\left(\frac{\frac{1}{10} + \frac{13\sqrt{29}}{290}}{s + \frac{3}{2} - \frac{\sqrt{29}}{2}}\right) &= \frac{(29 + 13\sqrt{29}) e^{-\frac{3t}{2} + \frac{t\sqrt{29}}{2}}}{290} \\ \mathcal{L}^{-1}\left(\frac{\frac{1}{10} - \frac{13\sqrt{29}}{290}}{s + \frac{3}{2} + \frac{\sqrt{29}}{2}}\right) &= \frac{e^{\frac{(-3-\sqrt{29})t}{2}} \left(-\frac{3(3+\sqrt{29})t}{2\left(-\frac{3}{2} - \frac{\sqrt{29}}{2}\right)} - \frac{(3+\sqrt{29})t\sqrt{29}}{2\left(-\frac{3}{2} - \frac{\sqrt{29}}{2}\right)} \right)}{6+2\sqrt{29}} (29 - 13\sqrt{29}) \\ \mathcal{L}^{-1}\left(-\frac{1}{5s}\right) &= -\frac{1}{5} \end{aligned}$$

Adding the above results and simplifying gives

$$y = -\frac{1}{5} + \frac{e^{-\frac{3t}{2}} \left(13\sqrt{29} \sinh\left(\frac{t\sqrt{29}}{2}\right) + 29 \cosh\left(\frac{t\sqrt{29}}{2}\right) \right)}{145}$$


Simplifying the solution gives

$$y = \frac{13\sqrt{29} \sinh\left(\frac{t\sqrt{29}}{2}\right) e^{-\frac{3t}{2}}}{145} + \frac{\cosh\left(\frac{t\sqrt{29}}{2}\right) e^{-\frac{3t}{2}}}{5} - \frac{1}{5}$$

Summary

The solution(s) found are the following

$$y = \frac{13\sqrt{29} \sinh\left(\frac{t\sqrt{29}}{2}\right) e^{-\frac{3t}{2}}}{145} + \frac{\cosh\left(\frac{t\sqrt{29}}{2}\right) e^{-\frac{3t}{2}}}{5} - \frac{1}{5} \quad (1)$$



(a) Solution plot

(b) Slope field plot

Verification of solutions

$$y = \frac{13\sqrt{29} \sinh\left(\frac{t\sqrt{29}}{2}\right) e^{-\frac{3t}{2}}}{145} + \frac{\cosh\left(\frac{t\sqrt{29}}{2}\right) e^{-\frac{3t}{2}}}{5} - \frac{1}{5}$$

Verified OK.

25.1.2 Maple step by step solution

Let's solve

$$\left[y'' + 3y' - 5y = 1, y(0) = 0, y'|_{\{t=0\}} = 1 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 3r - 5 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-3) \pm (\sqrt{29})}{2}$$

- Roots of the characteristic polynomial

$$r = \left(-\frac{3}{2} - \frac{\sqrt{29}}{2}, -\frac{3}{2} + \frac{\sqrt{29}}{2} \right)$$

- 1st solution of the homogeneous ODE

$$y_1(t) = e^{\left(-\frac{3}{2} - \frac{\sqrt{29}}{2}\right)t}$$

- 2nd solution of the homogeneous ODE

$$y_2(t) = e^{\left(-\frac{3}{2} + \frac{\sqrt{29}}{2}\right)t}$$

- General solution of the ODE

$$y = c_1 y_1(t) + c_2 y_2(t) + y_p(t)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{\left(-\frac{3}{2} - \frac{\sqrt{29}}{2}\right)t} + c_2 e^{\left(-\frac{3}{2} + \frac{\sqrt{29}}{2}\right)t} + y_p(t)$$

- Find a particular solution $y_p(t)$ of the ODE

- Use variation of parameters to find y_p here $f(t)$ is the forcing function

$$\left[y_p(t) = -y_1(t) \left(\int \frac{y_2(t)f(t)}{W(y_1(t), y_2(t))} dt \right) + y_2(t) \left(\int \frac{y_1(t)f(t)}{W(y_1(t), y_2(t))} dt \right), f(t) = 1 \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(t), y_2(t)) = \begin{bmatrix} e^{\left(-\frac{3}{2} - \frac{\sqrt{29}}{2}\right)t} & e^{\left(-\frac{3}{2} + \frac{\sqrt{29}}{2}\right)t} \\ \left(-\frac{3}{2} - \frac{\sqrt{29}}{2}\right) e^{\left(-\frac{3}{2} - \frac{\sqrt{29}}{2}\right)t} & \left(-\frac{3}{2} + \frac{\sqrt{29}}{2}\right) e^{\left(-\frac{3}{2} + \frac{\sqrt{29}}{2}\right)t} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(t), y_2(t)) = \sqrt{29} e^{-3t}$$

- Substitute functions into equation for $y_p(t)$

$$y_p(t) = -\frac{\sqrt{29} \left(e^{-\frac{(3+\sqrt{29})t}{2}} \left(\int e^{\frac{(3+\sqrt{29})t}{2}} dt \right) - e^{-\frac{(-3+\sqrt{29})t}{2}} \left(\int e^{-\frac{(-3+\sqrt{29})t}{2}} dt \right) \right)}{29}$$

- Compute integrals

$$y_p(t) = -\frac{1}{5}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{\left(-\frac{3}{2} - \frac{\sqrt{29}}{2}\right)t} + c_2 e^{\left(-\frac{3}{2} + \frac{\sqrt{29}}{2}\right)t} - \frac{1}{5}$$

- Check validity of solution $y = c_1 e^{\left(-\frac{3}{2} - \frac{\sqrt{29}}{2}\right)t} + c_2 e^{\left(-\frac{3}{2} + \frac{\sqrt{29}}{2}\right)t} - \frac{1}{5}$

- Use initial condition $y(0) = 0$

$$0 = c_1 + c_2 - \frac{1}{5}$$

- Compute derivative of the solution

$$y' = c_1 \left(-\frac{3}{2} - \frac{\sqrt{29}}{2} \right) e^{\left(-\frac{3}{2} - \frac{\sqrt{29}}{2} \right) t} + c_2 \left(-\frac{3}{2} + \frac{\sqrt{29}}{2} \right) e^{\left(-\frac{3}{2} + \frac{\sqrt{29}}{2} \right) t}$$

- Use the initial condition $y' \Big|_{\{t=0\}} = 1$

$$1 = c_1 \left(-\frac{3}{2} - \frac{\sqrt{29}}{2} \right) + c_2 \left(-\frac{3}{2} + \frac{\sqrt{29}}{2} \right)$$

- Solve for c_1 and c_2

$$\left\{ c_1 = \frac{1}{10} - \frac{13\sqrt{29}}{290}, c_2 = \frac{1}{10} + \frac{13\sqrt{29}}{290} \right\}$$

- Substitute constant values into general solution and simplify

$$y = \frac{(29+13\sqrt{29})e^{\frac{(-3+\sqrt{29})t}{2}}}{290} - \frac{1}{5} + \frac{(29-13\sqrt{29})e^{-\frac{(3+\sqrt{29})t}{2}}}{290}$$

- Solution to the IVP

$$y = \frac{(29+13\sqrt{29})e^{\frac{(-3+\sqrt{29})t}{2}}}{290} - \frac{1}{5} + \frac{(29-13\sqrt{29})e^{-\frac{(3+\sqrt{29})t}{2}}}{290}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 1.953 (sec). Leaf size: 34

```
dsolve([diff(y(t),t$2)+3*diff(y(t),t)-5*y(t)=1,y(0) = 0, D(y)(0) = 1],y(t), singsol=all)
```

$$y(t) = \frac{13 e^{-\frac{3t}{2}} \sqrt{29} \sinh\left(\frac{t\sqrt{29}}{2}\right)}{145} + \frac{e^{-\frac{3t}{2}} \cosh\left(\frac{t\sqrt{29}}{2}\right)}{5} - \frac{1}{5}$$

✓ Solution by Mathematica

Time used: 0.026 (sec). Leaf size: 67

```
DSolve[{y''[t]+3*y'[t]-5*y[t]==1,{y[0]==0,y'[0]==1}},y[t],t,IncludeSingularSolutions -> True
```

$$y(t) \rightarrow \frac{1}{290} e^{-\frac{1}{2}(3+\sqrt{29})t} \left((29 + 13\sqrt{29}) e^{\sqrt{29}t} - 58 e^{\frac{1}{2}(3+\sqrt{29})t} + 29 - 13\sqrt{29} \right)$$

25.2 problem 3(b)

25.2.1 Existence and uniqueness analysis 3895

Internal problem ID [6506]

Internal file name [OUTPUT/5754_Sunday_June_05_2022_03_52_52_PM_88188911/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 7. Laplace Transforms. Section 7.5 Problem for review and discovery. Section A, Drill exercises. Page 309

Problem number: 3(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_laplace", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' + 3y' - 2y = -6 e^{\pi-t}$$

With initial conditions

$$[y(\pi) = 1, y'(\pi) = 4]$$

25.2.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(t)y' + q(t)y = F$$

Where here

$$p(t) = 3$$

$$q(t) = -2$$

$$F = -6 e^{\pi-t}$$

Hence the ode is

$$y'' + 3y' - 2y = -6 e^{\pi-t}$$

The domain of $p(t) = 3$ is

$$\{-\infty < t < \infty\}$$

And the point $t_0 = \pi$ is inside this domain. The domain of $q(t) = -2$ is

$$\{-\infty < t < \infty\}$$

And the point $t_0 = \pi$ is also inside this domain. The domain of $F = -6e^{\pi-t}$ is

$$\{-\infty < t < \infty\}$$

And the point $t_0 = \pi$ is also inside this domain. Hence solution exists and is unique.

Since both initial conditions are not at zero, then let

$$\begin{aligned}y(0) &= c_1 \\y'(0) &= c_2\end{aligned}$$

Solving using the Laplace transform method. Let

$$\mathcal{L}(y) = Y(s)$$

Taking the Laplace transform of the ode and using the relations that

$$\begin{aligned}\mathcal{L}(y') &= sY(s) - y(0) \\ \mathcal{L}(y'') &= s^2Y(s) - y'(0) - sy(0)\end{aligned}$$

The given ode now becomes an algebraic equation in the Laplace domain

$$s^2Y(s) - y'(0) - sy(0) + 3sY(s) - 3y(0) - 2Y(s) = -\frac{6e^\pi}{s+1} \quad (1)$$

But the initial conditions are

$$\begin{aligned}y(0) &= c_1 \\y'(0) &= c_2\end{aligned}$$

Substituting these initial conditions in above in Eq (1) gives

$$s^2Y(s) - c_2 - sc_1 + 3sY(s) - 3c_1 - 2Y(s) = -\frac{6e^\pi}{s+1}$$

Solving the above equation for $Y(s)$ results in

$$Y(s) = -\frac{-s^2c_1 - 4sc_1 - c_2s + 6e^\pi - 3c_1 - c_2}{(s+1)(s^2+3s-2)}$$

Applying partial fractions decomposition results in

$$Y(s) = \frac{\left(-\frac{3e^\pi}{34} + \frac{3c_1}{17} + \frac{2c_2}{17}\right)\left(-\frac{3}{2} + \frac{\sqrt{17}}{2}\right) - \frac{15e^\pi}{17} + \frac{13c_1}{17} + \frac{3c_2}{17}}{s + \frac{3}{2} - \frac{\sqrt{17}}{2}} + \frac{\left(-\frac{3e^\pi}{34} + \frac{3c_1}{17} + \frac{2c_2}{17}\right)\left(-\frac{3}{2} - \frac{\sqrt{17}}{2}\right) - \frac{15e^\pi}{17} + \frac{13c_1}{17} + \frac{3c_2}{17}}{s + \frac{3}{2} + \frac{\sqrt{17}}{2}}$$

The inverse Laplace of each term above is now found, which gives

$$\mathcal{L}^{-1}\left(\frac{\left(-\frac{3e^\pi}{34} + \frac{3c_1}{17} + \frac{2c_2}{17}\right)\left(-\frac{3}{2} + \frac{\sqrt{17}}{2}\right) - \frac{15e^\pi}{17} + \frac{13c_1}{17} + \frac{3c_2}{17}}{s + \frac{3}{2} - \frac{\sqrt{17}}{2}}\right) = \frac{e^{-\frac{3t}{2} + \frac{t\sqrt{17}}{2}}(4c_2\sqrt{17} - 3e^\pi\sqrt{17} - 51e^\pi + 2c_1)}{68}$$

$$\mathcal{L}^{-1}\left(\frac{\left(-\frac{3e^\pi}{34} + \frac{3c_1}{17} + \frac{2c_2}{17}\right)\left(-\frac{3}{2} - \frac{\sqrt{17}}{2}\right) - \frac{15e^\pi}{17} + \frac{13c_1}{17} + \frac{3c_2}{17}}{s + \frac{3}{2} + \frac{\sqrt{17}}{2}}\right) = \frac{e^{\frac{(-3-\sqrt{17})t}{2}}\left(-\frac{3(3+\sqrt{17})t}{2\left(-\frac{3}{2}-\frac{\sqrt{17}}{2}\right)} - \frac{(3+\sqrt{17})t\sqrt{17}}{2\left(-\frac{3}{2}-\frac{\sqrt{17}}{2}\right)}\right)}{6+2\sqrt{17}}(-4c_2\sqrt{17} - 3e^\pi\sqrt{17} - 51e^\pi + 2c_1)$$

$$\mathcal{L}^{-1}\left(\frac{3e^\pi}{2(s+1)}\right) = \frac{3e^{\pi-t}}{2}$$

Adding the above results and simplifying gives

$$y = \frac{3e^{\pi-t}}{2} + \frac{\left(17 \cosh\left(\frac{t\sqrt{17}}{2}\right)(-3e^\pi + 2c_1) + \sqrt{17} \sinh\left(\frac{t\sqrt{17}}{2}\right)(-3e^\pi + 6c_1 + 4c_2)\right)e^{-\frac{3t}{2}}}{34}$$

Since both initial conditions given are not at zero, then we need to setup two equations to solve for c_1, c_2 . At $t = \pi$ the first equation becomes, using the above solution

$$1 = \frac{3}{2} + \frac{\left(17 \cosh\left(\frac{\pi\sqrt{17}}{2}\right)(-3e^\pi + 2c_1) + \sqrt{17} \sinh\left(\frac{\pi\sqrt{17}}{2}\right)(-3e^\pi + 6c_1 + 4c_2)\right)e^{-\frac{3\pi}{2}}}{34}$$

And taking derivative of the solution and evaluating at $t = \pi$ gives the second equation as

$$4 = -\frac{3}{2} + \frac{\left(\frac{17\sqrt{17} \sinh\left(\frac{\pi\sqrt{17}}{2}\right)(-3e^\pi + 2c_1)}{2} + \frac{17 \cosh\left(\frac{\pi\sqrt{17}}{2}\right)(-3e^\pi + 6c_1 + 4c_2)}{2}\right)e^{-\frac{3\pi}{2}}}{34} - \frac{3\left(17 \cosh\left(\frac{\pi\sqrt{17}}{2}\right)(-3e^\pi + 2c_1)\right)e^{-\frac{3\pi}{2}}}{34}$$

Solving gives

$$c_1 = \frac{e^{\frac{3\pi}{2}} \left(51 e^\pi \sinh \left(\frac{\pi\sqrt{17}}{2} \right)^2 e^{-\frac{3\pi}{2}} - 51 e^\pi e^{-\frac{3\pi}{2}} \cosh \left(\frac{\pi\sqrt{17}}{2} \right)^2 + 19\sqrt{17} \sinh \left(\frac{\pi\sqrt{17}}{2} \right) + 17 \cosh \left(\frac{\pi\sqrt{17}}{2} \right) \right)}{34 \sinh \left(\frac{\pi\sqrt{17}}{2} \right)^2 - 34 \cosh \left(\frac{\pi\sqrt{17}}{2} \right)^2}$$

$$c_2 = - \frac{\left(51 e^\pi \sinh \left(\frac{\pi\sqrt{17}}{2} \right)^2 e^{-\frac{3\pi}{2}} - 51 e^\pi e^{-\frac{3\pi}{2}} \cosh \left(\frac{\pi\sqrt{17}}{2} \right)^2 + 37\sqrt{17} \sinh \left(\frac{\pi\sqrt{17}}{2} \right) + 187 \cosh \left(\frac{\pi\sqrt{17}}{2} \right) \right) e^{\frac{3\pi}{2}}}{2 \left(17 \sinh \left(\frac{\pi\sqrt{17}}{2} \right)^2 - 17 \cosh \left(\frac{\pi\sqrt{17}}{2} \right)^2 \right)}$$

Substituting these in the solution obtained above gives

$$y = \frac{3 e^{\pi-t}}{2} + \frac{\left(17 \cosh \left(\frac{t\sqrt{17}}{2} \right) \left(-3 e^\pi + \frac{e^{\frac{3\pi}{2}} \left(51 e^\pi \sinh \left(\frac{\pi\sqrt{17}}{2} \right)^2 e^{-\frac{3\pi}{2}} - 51 e^\pi e^{-\frac{3\pi}{2}} \cosh \left(\frac{\pi\sqrt{17}}{2} \right)^2 + 19\sqrt{17} \sinh \left(\frac{\pi\sqrt{17}}{2} \right) + 17 \cosh \left(\frac{\pi\sqrt{17}}{2} \right) \right)}{17 \sinh \left(\frac{\pi\sqrt{17}}{2} \right)^2 - 17 \cosh \left(\frac{\pi\sqrt{17}}{2} \right)^2} \right)}{34}$$

$$= - \frac{19 \left(-\frac{51 e^{-\frac{\pi}{2}-t}}{19} + e^{-\frac{3t}{2}} \left(\left(-\sqrt{17} \sinh \left(\frac{t\sqrt{17}}{2} \right) + \frac{17 \cosh \left(\frac{t\sqrt{17}}{2} \right)}{19} \right) \cosh \left(\frac{\pi\sqrt{17}}{2} \right) + \sinh \left(\frac{\pi\sqrt{17}}{2} \right) \left(\sqrt{17} \cosh \left(\frac{\pi\sqrt{17}}{2} \right) \right) \right)}{34}$$


Simplifying the solution gives

$$y = \frac{19 \left(-\frac{51 e^{-\frac{\pi}{2}-t}}{19} + e^{-\frac{3t}{2}} \left(\left(-\sqrt{17} \sinh \left(\frac{t\sqrt{17}}{2} \right) + \frac{17 \cosh \left(\frac{t\sqrt{17}}{2} \right)}{19} \right) \cosh \left(\frac{\pi\sqrt{17}}{2} \right) + \sinh \left(\frac{\pi\sqrt{17}}{2} \right) \left(\sqrt{17} \cosh \left(\frac{\pi\sqrt{17}}{2} \right) \right) \right)}{34}$$


Summary

The solution(s) found are the following

$$y = \frac{19 \left(-\frac{51 e^{-\frac{\pi}{2}-t}}{19} + e^{-\frac{3t}{2}} \left(\left(-\sqrt{17} \sinh \left(\frac{t\sqrt{17}}{2} \right) + \frac{17 \cosh \left(\frac{t\sqrt{17}}{2} \right)}{19} \right) \cosh \left(\frac{\pi\sqrt{17}}{2} \right) + \sinh \left(\frac{\pi\sqrt{17}}{2} \right) \left(\sqrt{17} \cosh \left(\frac{\pi\sqrt{17}}{2} \right) \right) \right)}{34} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$y =$

$$19 \left(-\frac{51 e^{-\frac{\pi}{2}-t}}{19} + e^{-\frac{3t}{2}} \left(\left(-\sqrt{17} \sinh \left(\frac{t\sqrt{17}}{2} \right) + \frac{17 \cosh \left(\frac{t\sqrt{17}}{2} \right)}{19} \right) \cosh \left(\frac{\pi\sqrt{17}}{2} \right) + \sinh \left(\frac{\pi\sqrt{17}}{2} \right) \right) \left(\sqrt{17} \cosh \left(\frac{\pi\sqrt{17}}{2} \right) + \sinh \left(\frac{\pi\sqrt{17}}{2} \right) \right) \right)$$

34

Verified OK.

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```


✓ Solution by Maple

Time used: 2.078 (sec). Leaf size: 53

`dsolve([diff(y(t),t$2)+3*diff(y(t),t)-2*y(t)=-6*exp(Pi-t),y(Pi) = 1, D(y)(Pi) = 4],y(t), sin`

$$y(t) = -\frac{19 \sinh\left(\frac{(\pi-t)\sqrt{17}}{2}\right) \sqrt{17} e^{-\frac{3t}{2} + \frac{3\pi}{2}}}{34} - \frac{\cosh\left(\frac{(\pi-t)\sqrt{17}}{2}\right) e^{-\frac{3t}{2} + \frac{3\pi}{2}}}{2} + \frac{3e^{\pi-t}}{2}$$

✓ Solution by Mathematica

Time used: 0.354 (sec). Leaf size: 103

`DSolve[{y''[t]+3*y'[t]-2*y[t]==-6*Exp[Pi-t],{y[Pi]==1,y'[Pi]==4}},y[t],t,IncludeSingularSolu`

$$y(t) \rightarrow \frac{1}{68} e^{-\frac{1}{2}(3+\sqrt{17})t - \frac{1}{2}(\sqrt{17}-3)\pi} \left((19\sqrt{17} - 17) e^{\sqrt{17}t} + 102 e^{\frac{1}{2}((1+\sqrt{17})t + (\sqrt{17}-1)\pi)} - \left((17 + 19\sqrt{17}) e^{\sqrt{17}\pi} \right) \right)$$

25.3 problem 3(c)

25.3.1 Existence and uniqueness analysis	3901
25.3.2 Maple step by step solution	3904

Internal problem ID [6507]

Internal file name [OUTPUT/5755_Sunday_June_05_2022_03_52_55_PM_56265727/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 7. Laplace Transforms. Section 7.5 Problems for review and discovery. Section A, Drill exercises. Page 309

Problem number: 3(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_laplace", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + 2y' - y = t e^{-t}$$

With initial conditions

$$[y(0) = 0, y'(0) = 1]$$

25.3.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(t)y' + q(t)y = F$$

Where here

$$p(t) = 2$$

$$q(t) = -1$$

$$F = t e^{-t}$$

Hence the ode is

$$y'' + 2y' - y = t e^{-t}$$

The domain of $p(t) = 2$ is

$$\{-\infty < t < \infty\}$$

And the point $t_0 = 0$ is inside this domain. The domain of $q(t) = -1$ is

$$\{-\infty < t < \infty\}$$

And the point $t_0 = 0$ is also inside this domain. The domain of $F = t e^{-t}$ is

$$\{-\infty < t < \infty\}$$

And the point $t_0 = 0$ is also inside this domain. Hence solution exists and is unique.

Solving using the Laplace transform method. Let

$$\mathcal{L}(y) = Y(s)$$

Taking the Laplace transform of the ode and using the relations that

$$\mathcal{L}(y') = sY(s) - y(0)$$

$$\mathcal{L}(y'') = s^2Y(s) - y'(0) - sy(0)$$

The given ode now becomes an algebraic equation in the Laplace domain

$$s^2Y(s) - y'(0) - sy(0) + 2sY(s) - 2y(0) - Y(s) = \frac{1}{(s+1)^2} \quad (1)$$

But the initial conditions are

$$y(0) = 0$$

$$y'(0) = 1$$

Substituting these initial conditions in above in Eq (1) gives

$$s^2Y(s) - 1 + 2sY(s) - Y(s) = \frac{1}{(s+1)^2}$$

Solving the above equation for $Y(s)$ results in

$$Y(s) = \frac{s^2 + 2s + 2}{(s+1)^2(s^2 + 2s - 1)}$$

Applying partial fractions decomposition results in

$$Y(s) = -\frac{1}{2(s+1)^2} + \frac{3\sqrt{2}}{8(s-\sqrt{2}+1)} - \frac{3\sqrt{2}}{8(s+1+\sqrt{2})}$$

The inverse Laplace of each term above is now found, which gives

$$\begin{aligned}\mathcal{L}^{-1}\left(-\frac{1}{2(s+1)^2}\right) &= -\frac{te^{-t}}{2} \\ \mathcal{L}^{-1}\left(\frac{3\sqrt{2}}{8(s-\sqrt{2}+1)}\right) &= \frac{3\sqrt{2}e^{t\sqrt{2}-t}}{8} \\ \mathcal{L}^{-1}\left(-\frac{3\sqrt{2}}{8(s+1+\sqrt{2})}\right) &= -\frac{3\sqrt{2}e^{\frac{(-1-\sqrt{2})\left(-\frac{t\sqrt{2}(1+\sqrt{2})}{-1-\sqrt{2}} - \frac{t(1+\sqrt{2})}{-1-\sqrt{2}}\right)}{1+\sqrt{2}}}}{8}\end{aligned}$$

Adding the above results and simplifying gives

$$y = \frac{e^{-t}(3\sqrt{2} \sinh(t\sqrt{2}) - 2t)}{4}$$


Simplifying the solution gives

$$y = \frac{e^{-t}(3\sqrt{2} \sinh(t\sqrt{2}) - 2t)}{4}$$


Summary

The solution(s) found are the following

$$y = \frac{e^{-t}(3\sqrt{2} \sinh(t\sqrt{2}) - 2t)}{4} \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{e^{-t}(3\sqrt{2} \sinh(t\sqrt{2}) - 2t)}{4}$$

Verified OK.

25.3.2 Maple step by step solution

Let's solve

$$\left[y'' + 2y' - y = t e^{-t}, y(0) = 0, y'|_{\{t=0\}} = 1 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 2r - 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-2) \pm (\sqrt{8})}{2}$$

- Roots of the characteristic polynomial

$$r = (-1 - \sqrt{2}, \sqrt{2} - 1)$$

- 1st solution of the homogeneous ODE

$$y_1(t) = e^{(-1-\sqrt{2})t}$$

- 2nd solution of the homogeneous ODE

$$y_2(t) = e^{(\sqrt{2}-1)t}$$

- General solution of the ODE

$$y = c_1 y_1(t) + c_2 y_2(t) + y_p(t)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{(-1-\sqrt{2})t} + c_2 e^{(\sqrt{2}-1)t} + y_p(t)$$

- Find a particular solution $y_p(t)$ of the ODE

- Use variation of parameters to find y_p here $f(t)$ is the forcing function

$$\left[y_p(t) = -y_1(t) \left(\int \frac{y_2(t)f(t)}{W(y_1(t),y_2(t))} dt \right) + y_2(t) \left(\int \frac{y_1(t)f(t)}{W(y_1(t),y_2(t))} dt \right), f(t) = t e^{-t} \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(t), y_2(t)) = \begin{bmatrix} e^{(-1-\sqrt{2})t} & e^{(\sqrt{2}-1)t} \\ (-1-\sqrt{2})e^{(-1-\sqrt{2})t} & (\sqrt{2}-1)e^{(\sqrt{2}-1)t} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(t), y_2(t)) = 2\sqrt{2}e^{-2t}$$

- Substitute functions into equation for $y_p(t)$

$$y_p(t) = -\frac{\sqrt{2} \left(e^{-t(1+\sqrt{2})} \left(\int t e^{t\sqrt{2}} dt \right) - e^{(\sqrt{2}-1)t} \left(\int t e^{-t\sqrt{2}} dt \right) \right)}{4}$$

- Compute integrals

$$y_p(t) = -\frac{t e^{-t}}{2}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{(-1-\sqrt{2})t} + c_2 e^{(\sqrt{2}-1)t} - \frac{t e^{-t}}{2}$$

- Check validity of solution $y = c_1 e^{(-1-\sqrt{2})t} + c_2 e^{(\sqrt{2}-1)t} - \frac{t e^{-t}}{2}$

- Use initial condition $y(0) = 0$

$$0 = c_1 + c_2$$

- Compute derivative of the solution

$$y' = c_1 (-1-\sqrt{2}) e^{(-1-\sqrt{2})t} + c_2 (\sqrt{2}-1) e^{(\sqrt{2}-1)t} - \frac{e^{-t}}{2} + \frac{t e^{-t}}{2}$$

- Use the initial condition $y' \Big|_{\{t=0\}} = 1$

$$1 = c_1(-1 - \sqrt{2}) + c_2(\sqrt{2} - 1) - \frac{1}{2}$$

- Solve for c_1 and c_2

$$\left\{ c_1 = -\frac{3\sqrt{2}}{8}, c_2 = \frac{3\sqrt{2}}{8} \right\}$$

- Substitute constant values into general solution and simplify

$$y = -\frac{3\sqrt{2}e^{-t(1+\sqrt{2})}}{8} + \frac{3\sqrt{2}e^{(\sqrt{2}-1)t}}{8} - \frac{te^{-t}}{2}$$

- Solution to the IVP

$$y = -\frac{3\sqrt{2}e^{-t(1+\sqrt{2})}}{8} + \frac{3\sqrt{2}e^{(\sqrt{2}-1)t}}{8} - \frac{te^{-t}}{2}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 1.969 (sec). Leaf size: 25

```
dsolve([diff(y(t),t$2)+2*diff(y(t),t)-y(t)=t*exp(-t),y(0) = 0, D(y)(0) = 1],y(t), singsol=all)
```

$$y(t) = \frac{e^{-t}(3\sqrt{2} \sinh(\sqrt{2}t) - 2t)}{4}$$

✓ Solution by Mathematica

Time used: 0.029 (sec). Leaf size: 51

```
DSolve[{y''[t]+2*y'[t]-y[t]==t*Exp[-t]},{y[0]==0,y'[0]==1},y[t],t,IncludeSingularSolutions -
```

$$y(t) \rightarrow \frac{1}{8}e^{-t} \left(-4t - 3\sqrt{2}e^{-\sqrt{2}t} + 3\sqrt{2}e^{\sqrt{2}t} \right)$$

25.4 problem 3(d)

25.4.1 Existence and uniqueness analysis	3908
25.4.2 Maple step by step solution	3911

Internal problem ID [6508]

Internal file name [OUTPUT/5756_Sunday_June_05_2022_03_52_57_PM_36386801/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 7. Laplace Transforms. Section 7.5 Problems for review and discovery. Section A, Drill exercises. Page 309

Problem number: 3(d).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_laplace", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' - y' + y = 3e^{-t}$$

With initial conditions

$$[y(0) = 3, y'(0) = 2]$$

25.4.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(t)y' + q(t)y = F$$

Where here

$$p(t) = -1$$

$$q(t) = 1$$

$$F = 3e^{-t}$$

Hence the ode is

$$y'' - y' + y = 3e^{-t}$$

The domain of $p(t) = -1$ is

$$\{-\infty < t < \infty\}$$

And the point $t_0 = 0$ is inside this domain. The domain of $q(t) = 1$ is

$$\{-\infty < t < \infty\}$$

And the point $t_0 = 0$ is also inside this domain. The domain of $F = 3e^{-t}$ is

$$\{-\infty < t < \infty\}$$

And the point $t_0 = 0$ is also inside this domain. Hence solution exists and is unique.

Solving using the Laplace transform method. Let

$$\mathcal{L}(y) = Y(s)$$

Taking the Laplace transform of the ode and using the relations that

$$\begin{aligned}\mathcal{L}(y') &= sY(s) - y(0) \\ \mathcal{L}(y'') &= s^2Y(s) - y'(0) - sy(0)\end{aligned}$$

The given ode now becomes an algebraic equation in the Laplace domain

$$s^2Y(s) - y'(0) - sy(0) - sY(s) + y(0) + Y(s) = \frac{3}{s+1} \quad (1)$$

But the initial conditions are

$$\begin{aligned}y(0) &= 3 \\ y'(0) &= 2\end{aligned}$$

Substituting these initial conditions in above in Eq (1) gives

$$s^2Y(s) + 1 - 3s - sY(s) + Y(s) = \frac{3}{s+1}$$

Solving the above equation for $Y(s)$ results in

$$Y(s) = \frac{3s^2 + 2s + 2}{(s+1)(s^2 - s + 1)}$$

Applying partial fractions decomposition results in

$$Y(s) = \frac{1 - \frac{2i\sqrt{3}}{3}}{s - \frac{1}{2} - \frac{i\sqrt{3}}{2}} + \frac{1 + \frac{2i\sqrt{3}}{3}}{s - \frac{1}{2} + \frac{i\sqrt{3}}{2}} + \frac{1}{s + 1}$$

The inverse Laplace of each term above is now found, which gives

$$\begin{aligned}\mathcal{L}^{-1}\left(\frac{1 - \frac{2i\sqrt{3}}{3}}{s - \frac{1}{2} - \frac{i\sqrt{3}}{2}}\right) &= \frac{(3 - 2i\sqrt{3}) e^{\frac{(1+i\sqrt{3})t}{2}}}{3} \\ \mathcal{L}^{-1}\left(\frac{1 + \frac{2i\sqrt{3}}{3}}{s - \frac{1}{2} + \frac{i\sqrt{3}}{2}}\right) &= \frac{(2i\sqrt{3} + 3) e^{\frac{(1-i\sqrt{3})t}{2}}}{3} \\ \mathcal{L}^{-1}\left(\frac{1}{s + 1}\right) &= e^{-t}\end{aligned}$$

Adding the above results and simplifying gives

$$y = e^{-t} + \frac{2e^{\frac{t}{2}}\left(2\sqrt{3}\sin\left(\frac{\sqrt{3}t}{2}\right) + 3\cos\left(\frac{\sqrt{3}t}{2}\right)\right)}{3}$$


Simplifying the solution gives

$$y = \frac{\left(4\sqrt{3}\sin\left(\frac{\sqrt{3}t}{2}\right)e^{\frac{3t}{2}} + 6\cos\left(\frac{\sqrt{3}t}{2}\right)e^{\frac{3t}{2}} + 3\right)e^{-t}}{3}$$


Summary

The solution(s) found are the following

$$y = \frac{\left(4\sqrt{3}\sin\left(\frac{\sqrt{3}t}{2}\right)e^{\frac{3t}{2}} + 6\cos\left(\frac{\sqrt{3}t}{2}\right)e^{\frac{3t}{2}} + 3\right)e^{-t}}{3} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{\left(4\sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right) e^{\frac{3t}{2}} + 6 \cos\left(\frac{\sqrt{3}t}{2}\right) e^{\frac{3t}{2}} + 3\right) e^{-t}}{3}$$

Verified OK.

25.4.2 Maple step by step solution

Let's solve

$$\left[y'' - y' + y = 3e^{-t}, y(0) = 3, y'|_{\{t=0\}} = 2 \right]$$

- Highest derivative means the order of the ODE is 2

y''

- Characteristic polynomial of homogeneous ODE

$$r^2 - r + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{1 \pm (\sqrt{-3})}{2}$$

- Roots of the characteristic polynomial

$$r = \left(\frac{1}{2} - \frac{I\sqrt{3}}{2}, \frac{1}{2} + \frac{I\sqrt{3}}{2} \right)$$

- 1st solution of the homogeneous ODE

$$y_1(t) = \cos\left(\frac{\sqrt{3}t}{2}\right) e^{\frac{t}{2}}$$

- 2nd solution of the homogeneous ODE

$$y_2(t) = e^{\frac{t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right)$$

- General solution of the ODE

$$y = c_1 y_1(t) + c_2 y_2(t) + y_p(t)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 \cos\left(\frac{\sqrt{3}t}{2}\right) e^{\frac{t}{2}} + c_2 e^{\frac{t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right) + y_p(t)$$

- Find a particular solution $y_p(t)$ of the ODE

- Use variation of parameters to find y_p here $f(t)$ is the forcing function

$$\left[y_p(t) = -y_1(t) \left(\int \frac{y_2(t)f(t)}{W(y_1(t), y_2(t))} dt \right) + y_2(t) \left(\int \frac{y_1(t)f(t)}{W(y_1(t), y_2(t))} dt \right), f(t) = 3e^{-t} \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(t), y_2(t)) = \begin{bmatrix} \cos\left(\frac{\sqrt{3}t}{2}\right) e^{\frac{t}{2}} & e^{\frac{t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right) \\ -\frac{\sqrt{3}}{2} \sin\left(\frac{\sqrt{3}t}{2}\right) e^{\frac{t}{2}} + \frac{\cos\left(\frac{\sqrt{3}t}{2}\right) e^{\frac{t}{2}}}{2} & \frac{e^{\frac{t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right)}{2} + \frac{e^{\frac{t}{2}} \sqrt{3} \cos\left(\frac{\sqrt{3}t}{2}\right)}{2} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(t), y_2(t)) = \frac{\sqrt{3}e^t}{2}$$

- Substitute functions into equation for $y_p(t)$

$$y_p(t) = -2\sqrt{3}e^{\frac{t}{2}} \left(\cos\left(\frac{\sqrt{3}t}{2}\right) \left(\int e^{-\frac{3t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right) dt \right) - \sin\left(\frac{\sqrt{3}t}{2}\right) \left(\int e^{-\frac{3t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right) dt \right) \right)$$

- Compute integrals

$$y_p(t) = e^{-t}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 \cos\left(\frac{\sqrt{3}t}{2}\right) e^{\frac{t}{2}} + c_2 e^{\frac{t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right) + e^{-t}$$

- Check validity of solution $y = c_1 \cos\left(\frac{\sqrt{3}t}{2}\right) e^{\frac{t}{2}} + c_2 e^{\frac{t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right) + e^{-t}$

- Use initial condition $y(0) = 3$

$$3 = c_1 + 1$$

- Compute derivative of the solution

$$y' = -\frac{c_1\sqrt{3}\sin\left(\frac{\sqrt{3}t}{2}\right)e^{\frac{t}{2}}}{2} + \frac{c_1\cos\left(\frac{\sqrt{3}t}{2}\right)e^{\frac{t}{2}}}{2} + \frac{c_2e^{\frac{t}{2}}\sin\left(\frac{\sqrt{3}t}{2}\right)}{2} + \frac{c_2e^{\frac{t}{2}}\sqrt{3}\cos\left(\frac{\sqrt{3}t}{2}\right)}{2} - e^{-t}$$

- Use the initial condition $y' \Big|_{\{t=0\}} = 2$

$$2 = -1 + \frac{c_1}{2} + \frac{c_2\sqrt{3}}{2}$$

- Solve for c_1 and c_2

$$\left\{ c_1 = 2, c_2 = \frac{4\sqrt{3}}{3} \right\}$$

- Substitute constant values into general solution and simplify

$$y = \frac{(4\sqrt{3}\sin\left(\frac{\sqrt{3}t}{2}\right)e^{\frac{3t}{2}} + 6\cos\left(\frac{\sqrt{3}t}{2}\right)e^{\frac{3t}{2}} + 3)e^{-t}}{3}$$

- Solution to the IVP

$$y = \frac{(4\sqrt{3}\sin\left(\frac{\sqrt{3}t}{2}\right)e^{\frac{3t}{2}} + 6\cos\left(\frac{\sqrt{3}t}{2}\right)e^{\frac{3t}{2}} + 3)e^{-t}}{3}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 2.016 (sec). Leaf size: 37

```
dsolve([diff(y(t),t$2)-diff(y(t),t)+y(t)=3*exp(-t),y(0) = 3, D(y)(0) = 2],y(t), singsol=all)
```

$$y(t) = \frac{(4\sqrt{3}e^{\frac{3t}{2}}\sin\left(\frac{\sqrt{3}t}{2}\right) + 6e^{\frac{3t}{2}}\cos\left(\frac{\sqrt{3}t}{2}\right) + 3)e^{-t}}{3}$$

✓ Solution by Mathematica

Time used: 0.03 (sec). Leaf size: 56

```
DSolve[{y''[t]-y'[t]+y[t]==3*Exp[-t],{y[0]==3,y'[0]==2}},y[t],t,IncludeSingularSolutions ->
```

$$y(t) \rightarrow e^{-t} + \frac{4e^{t/2} \sin\left(\frac{\sqrt{3}t}{2}\right)}{\sqrt{3}} + 2e^{t/2} \cos\left(\frac{\sqrt{3}t}{2}\right)$$

25.5 problem 4(a)

25.5.1 Maple step by step solution 3917

Internal problem ID [6509]

Internal file name [OUTPUT/5757_Sunday_June_05_2022_03_52_59_PM_20889182/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 7. Laplace Transforms. Section 7.5 Problems for review and discovery. Section A, Drill exercises. Page 309

Problem number: 4(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_laplace", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x]]
```

$$y'' - 5y' + 4y = 0$$

Since no initial conditions are explicitly given, then let

$$y(0) = c_1$$

$$y'(0) = c_2$$

Solving using the Laplace transform method. Let

$$\mathcal{L}(y) = Y(s)$$

Taking the Laplace transform of the ode and using the relations that

$$\mathcal{L}(y') = sY(s) - y(0)$$

$$\mathcal{L}(y'') = s^2Y(s) - y'(0) - sy(0)$$

The given ode now becomes an algebraic equation in the Laplace domain

$$s^2Y(s) - y'(0) - sy(0) - 5sY(s) + 5y(0) + 4Y(s) = 0 \quad (1)$$

But the initial conditions are

$$\begin{aligned}y(0) &= c_1 \\y'(0) &= c_2\end{aligned}$$

Substituting these initial conditions in above in Eq (1) gives

$$s^2Y(s) - c_2 - sc_1 - 5sY(s) + 5c_1 + 4Y(s) = 0$$

Solving the above equation for $Y(s)$ results in

$$Y(s) = \frac{sc_1 - 5c_1 + c_2}{s^2 - 5s + 4}$$

Applying partial fractions decomposition results in

$$Y(s) = \frac{\frac{4c_1}{3} - \frac{c_2}{3}}{s - 1} + \frac{-\frac{c_1}{3} + \frac{c_2}{3}}{s - 4}$$

The inverse Laplace of each term above is now found, which gives

$$\begin{aligned}\mathcal{L}^{-1}\left(\frac{\frac{4c_1}{3} - \frac{c_2}{3}}{s - 1}\right) &= \frac{(4c_1 - c_2)e^t}{3} \\ \mathcal{L}^{-1}\left(\frac{-\frac{c_1}{3} + \frac{c_2}{3}}{s - 4}\right) &= \frac{e^{4t}(-c_1 + c_2)}{3}\end{aligned}$$

Adding the above results and simplifying gives

$$y = \frac{e^{\frac{5t}{2}}(3c_1 \cosh\left(\frac{3t}{2}\right) + \sinh\left(\frac{3t}{2}\right)(-5c_1 + 2c_2))}{3}$$

Simplifying the solution gives

$$y = \frac{e^{\frac{5t}{2}}(-5c_1 + 2c_2) \sinh\left(\frac{3t}{2}\right)}{3} + e^{\frac{5t}{2}}c_1 \cosh\left(\frac{3t}{2}\right)$$

Summary

The solution(s) found are the following

$$y = \frac{e^{\frac{5t}{2}}(-5c_1 + 2c_2) \sinh\left(\frac{3t}{2}\right)}{3} + e^{\frac{5t}{2}}c_1 \cosh\left(\frac{3t}{2}\right) \quad (1)$$




Figure 481: Slope field plot

Verification of solutions

$$y = \frac{e^{\frac{5t}{2}}(-5c_1 + 2c_2) \sinh\left(\frac{3t}{2}\right)}{3} + e^{\frac{5t}{2}} c_1 \cosh\left(\frac{3t}{2}\right)$$

Verified OK.

25.5.1 Maple step by step solution

Let's solve

$$y'' - 5y' + 4y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of ODE

$$r^2 - 5r + 4 = 0$$

- Factor the characteristic polynomial

$$(r - 1)(r - 4) = 0$$

- Roots of the characteristic polynomial

- $r = (1, 4)$
- 1st solution of the ODE
 $y_1(t) = e^t$
- 2nd solution of the ODE
 $y_2(t) = e^{4t}$
- General solution of the ODE
 $y = c_1 y_1(t) + c_2 y_2(t)$
- Substitute in solutions
 $y = c_1 e^t + c_2 e^{4t}$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 1.687 (sec). Leaf size: 33

```
dsolve(diff(y(t),t$2)-5*diff(y(t),t)+4*y(t)=0,y(t), singsol=all)
```

$$y(t) = \frac{e^{\frac{5t}{2}} \left(3y(0) \cosh\left(\frac{3t}{2}\right) + \sinh\left(\frac{3t}{2}\right) (2D(y)(0) - 5y(0)) \right)}{3}$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 20

```
DSolve[y''[t]-5*y'[t]+4*y[t]==0,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow e^t (c_2 e^{3t} + c_1)$$

25.6 problem 4(b)

25.6.1 Maple step by step solution 3922

Internal problem ID [6510]

Internal file name [OUTPUT/5758_Sunday_June_05_2022_03_53_01_PM_80445232/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 7. Laplace Transforms. Section 7.5 Problems for review and discovery. Section A, Drill exercises. Page 309

Problem number: 4(b).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_laplace", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' + 3y' + 3y = 2$$

Since no initial conditions are explicitly given, then let

$$y(0) = c_1$$

$$y'(0) = c_2$$

Solving using the Laplace transform method. Let

$$\mathcal{L}(y) = Y(s)$$

Taking the Laplace transform of the ode and using the relations that

$$\mathcal{L}(y') = sY(s) - y(0)$$

$$\mathcal{L}(y'') = s^2Y(s) - y'(0) - sy(0)$$

The given ode now becomes an algebraic equation in the Laplace domain

$$s^2Y(s) - y'(0) - sy(0) + 3sY(s) - 3y(0) + 3Y(s) = \frac{2}{s} \quad (1)$$

But the initial conditions are

$$\begin{aligned}y(0) &= c_1 \\y'(0) &= c_2\end{aligned}$$

Substituting these initial conditions in above in Eq (1) gives

$$s^2Y(s) - c_2 - sc_1 + 3sY(s) - 3c_1 + 3Y(s) = \frac{2}{s}$$

Solving the above equation for $Y(s)$ results in

$$Y(s) = \frac{c_1s^2 + 3sc_1 + c_2s + 2}{s(s^2 + 3s + 3)}$$

Applying partial fractions decomposition results in

$$Y(s) = \frac{2}{3s} + \frac{\left(-c_1 - \frac{2c_2}{3} + \frac{2}{3}\right)\left(-\frac{3}{2} + \frac{i\sqrt{3}}{2}\right) - c_1 - c_2 + \frac{2}{3}}{s + \frac{3}{2} - \frac{i\sqrt{3}}{2}} + \frac{\left(-c_1 - \frac{2c_2}{3} + \frac{2}{3}\right)\left(-\frac{3}{2} - \frac{i\sqrt{3}}{2}\right) - c_1 - c_2 + \frac{2}{3}}{s + \frac{3}{2} + \frac{i\sqrt{3}}{2}}$$

The inverse Laplace of each term above is now found, which gives

$$\mathcal{L}^{-1}\left(\frac{2}{3s}\right) = \frac{2}{3}$$

$$\mathcal{L}^{-1}\left(\frac{\left(-c_1 - \frac{2c_2}{3} + \frac{2}{3}\right)\left(-\frac{3}{2} + \frac{i\sqrt{3}}{2}\right) - c_1 - c_2 + \frac{2}{3}}{s + \frac{3}{2} - \frac{i\sqrt{3}}{2}}\right) = \frac{e^{-\frac{(-i\sqrt{3}+3)\left(-\frac{t\sqrt{3}}{2\left(-\frac{3}{2} + \frac{i\sqrt{3}}{2}\right)} + \frac{it}{-3+i\sqrt{3}}\right)\sqrt{3}}}{6}(-2 - 2ic_2\sqrt{3} + 2)$$

$$\mathcal{L}^{-1}\left(\frac{\left(-c_1 - \frac{2c_2}{3} + \frac{2}{3}\right)\left(-\frac{3}{2} - \frac{i\sqrt{3}}{2}\right) - c_1 - c_2 + \frac{2}{3}}{s + \frac{3}{2} + \frac{i\sqrt{3}}{2}}\right) = \frac{e^{-\frac{(i\sqrt{3}+3)\left(-\frac{t\sqrt{3}}{2\left(-\frac{3}{2} - \frac{i\sqrt{3}}{2}\right)} - \frac{it}{2\left(-\frac{3}{2} - \frac{i\sqrt{3}}{2}\right)}\right)\sqrt{3}}}{6}(-2 + 2ic_2\sqrt{3} - 2)$$

Adding the above results and simplifying gives

$$y = \frac{2}{3} + \frac{\left(\cos\left(\frac{\sqrt{3}t}{2}\right)(-2 + 3c_1) + \sin\left(\frac{\sqrt{3}t}{2}\right)\sqrt{3}(3c_1 + 2c_2 - 2)\right)e^{-\frac{3t}{2}}}{3}$$

Simplifying the solution gives

$$y = \sqrt{3}e^{-\frac{3t}{2}}\left(c_1 + \frac{2c_2}{3} - \frac{2}{3}\right)\sin\left(\frac{\sqrt{3}t}{2}\right) + \frac{2}{3} + \frac{e^{-\frac{3t}{2}}(-2 + 3c_1)\cos\left(\frac{\sqrt{3}t}{2}\right)}{3}$$

Summary

The solution(s) found are the following

$$y = \sqrt{3} e^{-\frac{3t}{2}} \left(c_1 + \frac{2c_2}{3} - \frac{2}{3} \right) \sin \left(\frac{\sqrt{3}t}{2} \right) + \frac{2}{3} + \frac{e^{-\frac{3t}{2}} (-2 + 3c_1) \cos \left(\frac{\sqrt{3}t}{2} \right)}{3} \quad (1)$$




Figure 482: Slope field plot

Verification of solutions

$$y = \sqrt{3} e^{-\frac{3t}{2}} \left(c_1 + \frac{2c_2}{3} - \frac{2}{3} \right) \sin \left(\frac{\sqrt{3}t}{2} \right) + \frac{2}{3} + \frac{e^{-\frac{3t}{2}} (-2 + 3c_1) \cos \left(\frac{\sqrt{3}t}{2} \right)}{3}$$

Verified OK.

25.6.1 Maple step by step solution

Let's solve

$$y'' + 3y' + 3y = 2$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 3r + 3 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-3) \pm (\sqrt{-3})}{2}$$

- Roots of the characteristic polynomial

$$r = \left(-\frac{3}{2} - \frac{i\sqrt{3}}{2}, -\frac{3}{2} + \frac{i\sqrt{3}}{2} \right)$$

- 1st solution of the homogeneous ODE

$$y_1(t) = e^{-\frac{3t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right)$$

- 2nd solution of the homogeneous ODE

$$y_2(t) = e^{-\frac{3t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right)$$

- General solution of the ODE

$$y = c_1 y_1(t) + c_2 y_2(t) + y_p(t)$$

- Substitute in solutions of the homogeneous ODE

$$y = \cos\left(\frac{\sqrt{3}t}{2}\right) e^{-\frac{3t}{2}} c_1 + \sin\left(\frac{\sqrt{3}t}{2}\right) e^{-\frac{3t}{2}} c_2 + y_p(t)$$

- Find a particular solution $y_p(t)$ of the ODE

- Use variation of parameters to find y_p here $f(t)$ is the forcing function

$$\left[y_p(t) = -y_1(t) \left(\int \frac{y_2(t)f(t)}{W(y_1(t),y_2(t))} dt \right) + y_2(t) \left(\int \frac{y_1(t)f(t)}{W(y_1(t),y_2(t))} dt \right), f(t) = 2 \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(t), y_2(t)) = \begin{bmatrix} e^{-\frac{3t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right) & e^{-\frac{3t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right) \\ -\frac{3e^{-\frac{3t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right)}{2} - \frac{\sqrt{3}e^{-\frac{3t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right)}{2} & -\frac{3e^{-\frac{3t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right)}{2} + \frac{\sqrt{3}e^{-\frac{3t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right)}{2} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(t), y_2(t)) = \frac{\sqrt{3}e^{-3t}}{2}$$

- Substitute functions into equation for $y_p(t)$

$$y_p(t) = -\frac{4\sqrt{3}e^{-\frac{3t}{2}} \left(\cos\left(\frac{\sqrt{3}t}{2}\right) \left(\int \sin\left(\frac{\sqrt{3}t}{2}\right) e^{\frac{3t}{2}} dt \right) - \sin\left(\frac{\sqrt{3}t}{2}\right) \left(\int \cos\left(\frac{\sqrt{3}t}{2}\right) e^{\frac{3t}{2}} dt \right) \right)}{3}$$

- Compute integrals

$$y_p(t) = \frac{2}{3}$$

- Substitute particular solution into general solution to ODE

$$y = \cos\left(\frac{\sqrt{3}t}{2}\right) e^{-\frac{3t}{2}} c_1 + \sin\left(\frac{\sqrt{3}t}{2}\right) e^{-\frac{3t}{2}} c_2 + \frac{2}{3}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 1.844 (sec). Leaf size: 48

```
dsolve(diff(y(t),t$2)+3*diff(y(t),t)+3*y(t)=2,y(t), singsol=all)
```

$$y(t) = \frac{2}{3} + \frac{\left(\cos\left(\frac{\sqrt{3}t}{2}\right) (-2 + 3y(0)) + \sin\left(\frac{\sqrt{3}t}{2}\right) \sqrt{3} (2D(y)(0) + 3y(0) - 2) \right) e^{-\frac{3t}{2}}}{3}$$

✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 51

```
DSolve[y''[t]+3*y'[t]+3*y[t]==2,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow c_2 e^{-3t/2} \cos\left(\frac{\sqrt{3}t}{2}\right) + c_1 e^{-3t/2} \sin\left(\frac{\sqrt{3}t}{2}\right) + \frac{2}{3}$$

25.7 problem 4(c)

25.7.1 Maple step by step solution 3927

Internal problem ID [6511]

Internal file name [OUTPUT/5759_Sunday_June_05_2022_03_53_02_PM_19961650/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 7. Laplace Transforms. Section 7.5 Problems for review and discovery. Section A, Drill exercises. Page 309

Problem number: 4(c).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_laplace", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$y'' + y' + 2y = t$$

Since no initial conditions are explicitly given, then let

$$y(0) = c_1$$

$$y'(0) = c_2$$

Solving using the Laplace transform method. Let

$$\mathcal{L}(y) = Y(s)$$

Taking the Laplace transform of the ode and using the relations that

$$\mathcal{L}(y') = sY(s) - y(0)$$

$$\mathcal{L}(y'') = s^2Y(s) - y'(0) - sy(0)$$

The given ode now becomes an algebraic equation in the Laplace domain

$$s^2Y(s) - y'(0) - sy(0) + sY(s) - y(0) + 2Y(s) = \frac{1}{s^2} \quad (1)$$

But the initial conditions are

$$\begin{aligned}y(0) &= c_1 \\y'(0) &= c_2\end{aligned}$$

Substituting these initial conditions in above in Eq (1) gives

$$s^2Y(s) - c_2 - sc_1 + sY(s) - c_1 + 2Y(s) = \frac{1}{s^2}$$

Solving the above equation for $Y(s)$ results in

$$Y(s) = \frac{c_1s^3 + c_1s^2 + c_2s^2 + 1}{s^2(s^2 + s + 2)}$$

Applying partial fractions decomposition results in

$$Y(s) = \frac{\left(-\frac{c_1}{7} - \frac{2c_2}{7} + \frac{3}{28}\right)\left(-\frac{1}{2} + \frac{i\sqrt{7}}{2}\right) + \frac{3c_1}{7} - \frac{c_2}{7} + \frac{5}{28}}{s + \frac{1}{2} - \frac{i\sqrt{7}}{2}} + \frac{\left(-\frac{c_1}{7} - \frac{2c_2}{7} + \frac{3}{28}\right)\left(-\frac{1}{2} - \frac{i\sqrt{7}}{2}\right) + \frac{3c_1}{7} - \frac{c_2}{7} + \frac{5}{28}}{s + \frac{1}{2} + \frac{i\sqrt{7}}{2}} - \frac{1}{4s} + \frac{t}{2}$$

The inverse Laplace of each term above is now found, which gives

$$\mathcal{L}^{-1}\left(\frac{\left(-\frac{c_1}{7} - \frac{2c_2}{7} + \frac{3}{28}\right)\left(-\frac{1}{2} + \frac{i\sqrt{7}}{2}\right) + \frac{3c_1}{7} - \frac{c_2}{7} + \frac{5}{28}}{s + \frac{1}{2} - \frac{i\sqrt{7}}{2}}\right) = \frac{e^{-\frac{(1-i\sqrt{7})}{2}\left(-\frac{t}{2\left(-\frac{1}{2} + \frac{i\sqrt{7}}{2}\right)} + \frac{i\sqrt{7}t}{-1+i\sqrt{7}}\right)}}{(7 - 8ic_2\sqrt{7} + 3iv)}{56}$$

$$\mathcal{L}^{-1}\left(\frac{\left(-\frac{c_1}{7} - \frac{2c_2}{7} + \frac{3}{28}\right)\left(-\frac{1}{2} - \frac{i\sqrt{7}}{2}\right) + \frac{3c_1}{7} - \frac{c_2}{7} + \frac{5}{28}}{s + \frac{1}{2} + \frac{i\sqrt{7}}{2}}\right) = \frac{e^{-\frac{(1+i\sqrt{7})}{2}\left(-\frac{t}{2\left(-\frac{1}{2} - \frac{i\sqrt{7}}{2}\right)} - \frac{i\sqrt{7}t}{2\left(-\frac{1}{2} - \frac{i\sqrt{7}}{2}\right)}\right)}}{(7 + 8ic_2\sqrt{7} - 3iv)}{56}$$

$$\mathcal{L}^{-1}\left(-\frac{1}{4s}\right) = -\frac{1}{4}$$

$$\mathcal{L}^{-1}\left(\frac{1}{2s^2}\right) = \frac{t}{2}$$

Adding the above results and simplifying gives

$$y = -\frac{1}{4} + \frac{t}{2} + \frac{\left(7 \cos\left(\frac{\sqrt{7}t}{2}\right)(1 + 4c_1) + \sin\left(\frac{\sqrt{7}t}{2}\right)\sqrt{7}(4c_1 + 8c_2 - 3)\right)e^{-\frac{t}{2}}}{28}$$

Simplifying the solution gives

$$y = -\frac{1}{4} + \frac{\sqrt{7}e^{-\frac{t}{2}}(c_1 + 2c_2 - \frac{3}{4})\sin\left(\frac{\sqrt{7}t}{2}\right)}{7} + \frac{(1 + 4c_1)e^{-\frac{t}{2}}\cos\left(\frac{\sqrt{7}t}{2}\right)}{4} + \frac{t}{2}$$

Summary

The solution(s) found are the following

$$y = -\frac{1}{4} + \frac{\sqrt{7} e^{-\frac{t}{2}} (c_1 + 2c_2 - \frac{3}{4}) \sin\left(\frac{\sqrt{7}t}{2}\right) + (1 + 4c_1) e^{-\frac{t}{2}} \cos\left(\frac{\sqrt{7}t}{2}\right) + \frac{t}{2}}{7} \quad (1)$$




Figure 483: Slope field plot

Verification of solutions

$$y = -\frac{1}{4} + \frac{\sqrt{7} e^{-\frac{t}{2}} (c_1 + 2c_2 - \frac{3}{4}) \sin\left(\frac{\sqrt{7}t}{2}\right) + (1 + 4c_1) e^{-\frac{t}{2}} \cos\left(\frac{\sqrt{7}t}{2}\right) + \frac{t}{2}}{7}$$

Verified OK.

25.7.1 Maple step by step solution

Let's solve

$$y'' + y' + 2y = t$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + r + 2 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-1) \pm (\sqrt{-7})}{2}$$

- Roots of the characteristic polynomial

$$r = \left(-\frac{1}{2} - \frac{i\sqrt{7}}{2}, -\frac{1}{2} + \frac{i\sqrt{7}}{2} \right)$$

- 1st solution of the homogeneous ODE

$$y_1(t) = e^{-\frac{t}{2}} \cos\left(\frac{\sqrt{7}t}{2}\right)$$

- 2nd solution of the homogeneous ODE

$$y_2(t) = e^{-\frac{t}{2}} \sin\left(\frac{\sqrt{7}t}{2}\right)$$

- General solution of the ODE

$$y = c_1 y_1(t) + c_2 y_2(t) + y_p(t)$$

- Substitute in solutions of the homogeneous ODE

$$y = e^{-\frac{t}{2}} \cos\left(\frac{\sqrt{7}t}{2}\right) c_1 + e^{-\frac{t}{2}} \sin\left(\frac{\sqrt{7}t}{2}\right) c_2 + y_p(t)$$

- Find a particular solution $y_p(t)$ of the ODE

- Use variation of parameters to find y_p here $f(t)$ is the forcing function

$$\left[y_p(t) = -y_1(t) \left(\int \frac{y_2(t)f(t)}{W(y_1(t), y_2(t))} dt \right) + y_2(t) \left(\int \frac{y_1(t)f(t)}{W(y_1(t), y_2(t))} dt \right), f(t) = t \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(t), y_2(t)) = \begin{bmatrix} e^{-\frac{t}{2}} \cos\left(\frac{\sqrt{7}t}{2}\right) & e^{-\frac{t}{2}} \sin\left(\frac{\sqrt{7}t}{2}\right) \\ -\frac{e^{-\frac{t}{2}} \cos\left(\frac{\sqrt{7}t}{2}\right)}{2} - \frac{\sqrt{7} e^{-\frac{t}{2}} \sin\left(\frac{\sqrt{7}t}{2}\right)}{2} & -\frac{e^{-\frac{t}{2}} \sin\left(\frac{\sqrt{7}t}{2}\right)}{2} + \frac{e^{-\frac{t}{2}} \sqrt{7} \cos\left(\frac{\sqrt{7}t}{2}\right)}{2} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(t), y_2(t)) = \frac{\sqrt{7} e^{-t}}{2}$$

- Substitute functions into equation for $y_p(t)$

$$y_p(t) = -\frac{2\sqrt{7} e^{-\frac{t}{2}} \left(\cos\left(\frac{\sqrt{7}t}{2}\right) \left(\int t e^{\frac{t}{2}} \sin\left(\frac{\sqrt{7}t}{2}\right) dt \right) - \sin\left(\frac{\sqrt{7}t}{2}\right) \left(\int t e^{\frac{t}{2}} \cos\left(\frac{\sqrt{7}t}{2}\right) dt \right) \right)}{7}$$

- Compute integrals

$$y_p(t) = \frac{t}{2} - \frac{1}{4}$$

- Substitute particular solution into general solution to ODE

$$y = e^{-\frac{t}{2}} \cos\left(\frac{\sqrt{7}t}{2}\right) c_1 + e^{-\frac{t}{2}} \sin\left(\frac{\sqrt{7}t}{2}\right) c_2 + \frac{t}{2} - \frac{1}{4}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 2.0 (sec). Leaf size: 52

```
dsolve(diff(y(t),t$2)+diff(y(t),t)+2*y(t)=t,y(t), singsol=all)
```

$$y(t) = -\frac{1}{4} + \frac{t}{2} + \frac{\left(7 \cos\left(\frac{\sqrt{7}t}{2}\right) (1 + 4y(0)) + \sin\left(\frac{\sqrt{7}t}{2}\right) \sqrt{7} (8D(y)(0) + 4y(0) - 3)\right) e^{-\frac{t}{2}}}{28}$$

✓ Solution by Mathematica

Time used: 0.027 (sec). Leaf size: 56

```
DSolve[y''[t]+y'[t]+2*y[t]==t,y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{t}{2} + c_2 e^{-t/2} \cos\left(\frac{\sqrt{7}t}{2}\right) + c_1 e^{-t/2} \sin\left(\frac{\sqrt{7}t}{2}\right) - \frac{1}{4}$$

25.8 problem 4(d)

25.8.1 Maple step by step solution 3932

Internal problem ID [6512]

Internal file name [OUTPUT/5760_Sunday_June_05_2022_03_53_04_PM_51888636/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 7. Laplace Transforms. Section 7.5 Problem for review and discovery. Section A, Drill exercises. Page 309

Problem number: 4(d).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_laplace", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' - 7y' + 12y = t e^{2t}$$

Since no initial conditions are explicitly given, then let

$$y(0) = c_1$$

$$y'(0) = c_2$$

Solving using the Laplace transform method. Let

$$\mathcal{L}(y) = Y(s)$$

Taking the Laplace transform of the ode and using the relations that

$$\mathcal{L}(y') = sY(s) - y(0)$$

$$\mathcal{L}(y'') = s^2Y(s) - y'(0) - sy(0)$$

The given ode now becomes an algebraic equation in the Laplace domain

$$s^2Y(s) - y'(0) - sy(0) - 7sY(s) + 7y(0) + 12Y(s) = \frac{1}{(s-2)^2} \quad (1)$$

But the initial conditions are

$$\begin{aligned}y(0) &= c_1 \\y'(0) &= c_2\end{aligned}$$

Substituting these initial conditions in above in Eq (1) gives

$$s^2Y(s) - c_2 - sc_1 - 7sY(s) + 7c_1 + 12Y(s) = \frac{1}{(s-2)^2}$$

Solving the above equation for $Y(s)$ results in

$$Y(s) = \frac{c_1s^3 - 11c_1s^2 + c_2s^2 + 32sc_1 - 4c_2s - 28c_1 + 4c_2 + 1}{(s-2)^2(s^2 - 7s + 12)}$$

Applying partial fractions decomposition results in

$$Y(s) = \frac{-3c_1 + c_2 + \frac{1}{4}}{s-4} + \frac{1}{2(s-2)^2} + \frac{3}{4(s-2)} + \frac{4c_1 - c_2 - 1}{s-3}$$

The inverse Laplace of each term above is now found, which gives

$$\begin{aligned}\mathcal{L}^{-1}\left(\frac{-3c_1 + c_2 + \frac{1}{4}}{s-4}\right) &= \frac{e^{4t}(-12c_1 + 4c_2 + 1)}{4} \\ \mathcal{L}^{-1}\left(\frac{1}{2(s-2)^2}\right) &= \frac{te^{2t}}{2} \\ \mathcal{L}^{-1}\left(\frac{3}{4(s-2)}\right) &= \frac{3e^{2t}}{4} \\ \mathcal{L}^{-1}\left(\frac{4c_1 - c_2 - 1}{s-3}\right) &= (4c_1 - c_2 - 1)e^{3t}\end{aligned}$$

Adding the above results and simplifying gives

$$y = \frac{e^{2t}(2t + 3)}{4} + (4c_1 - c_2 - 1)e^{3t} + \frac{e^{4t}(-12c_1 + 4c_2 + 1)}{4}$$

Simplifying the solution gives

$$y = \frac{e^{2t}(2t + 3)}{4} + (4c_1 - c_2 - 1)e^{3t} + \frac{e^{4t}(-12c_1 + 4c_2 + 1)}{4}$$

Summary

The solution(s) found are the following

$$y = \frac{e^{2t}(2t + 3)}{4} + (4c_1 - c_2 - 1)e^{3t} + \frac{e^{4t}(-12c_1 + 4c_2 + 1)}{4} \quad (1)$$




Figure 484: Slope field plot

Verification of solutions

$$y = \frac{e^{2t}(2t + 3)}{4} + (4c_1 - c_2 - 1)e^{3t} + \frac{e^{4t}(-12c_1 + 4c_2 + 1)}{4}$$

Verified OK.

25.8.1 Maple step by step solution

Let's solve

$$y'' - 7y' + 12y = te^{2t}$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - 7r + 12 = 0$$

- Factor the characteristic polynomial

$$(r - 3)(r - 4) = 0$$

- Roots of the characteristic polynomial

$$r = (3, 4)$$

- 1st solution of the homogeneous ODE

$$y_1(t) = e^{3t}$$

- 2nd solution of the homogeneous ODE

$$y_2(t) = e^{4t}$$

- General solution of the ODE

$$y = c_1 y_1(t) + c_2 y_2(t) + y_p(t)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{3t} + c_2 e^{4t} + y_p(t)$$

- Find a particular solution $y_p(t)$ of the ODE

- Use variation of parameters to find y_p here $f(t)$ is the forcing function

$$\left[y_p(t) = -y_1(t) \left(\int \frac{y_2(t)f(t)}{W(y_1(t), y_2(t))} dt \right) + y_2(t) \left(\int \frac{y_1(t)f(t)}{W(y_1(t), y_2(t))} dt \right), f(t) = t e^{2t} \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(t), y_2(t)) = \begin{bmatrix} e^{3t} & e^{4t} \\ 3e^{3t} & 4e^{4t} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(t), y_2(t)) = e^{7t}$$

- Substitute functions into equation for $y_p(t)$

$$y_p(t) = -e^{3t} \left(\int t e^{-t} dt \right) + e^{4t} \left(\int t e^{-2t} dt \right)$$

- Compute integrals

$$y_p(t) = \frac{e^{2t}(2t+3)}{4}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{3t} + c_2 e^{4t} + \frac{e^{2t}(2t+3)}{4}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 1.828 (sec). Leaf size: 47

```
dsolve(diff(y(t),t$2)-7*diff(y(t),t)+12*y(t)=t*exp(2*t),y(t), singsol=all)
```

$$y(t) = \frac{(2t + 3)e^{2t}}{4} + (4y(0) - D(y)(0) - 1)e^{3t} + \frac{e^{4t}(-12y(0) + 4D(y)(0) + 1)}{4}$$

✓ Solution by Mathematica

Time used: 0.019 (sec). Leaf size: 35

```
DSolve[y''[t]-7*y'[t]+12*y[t]==t*Exp[2*t],y[t],t,IncludeSingularSolutions -> True]
```

$$y(t) \rightarrow \frac{1}{4}e^{2t}(2t + 4c_1e^t + 4c_2e^{2t} + 3)$$

**26 Chapter 7. Laplace Transforms. Section 7.5
Problem for review and discovery. Section B,
Challenge Problems. Page 310**

26.1 problem 3 3936

26.1 problem 3

26.1.1 Existence and uniqueness analysis	3936
26.1.2 Maple step by step solution	3940

Internal problem ID [6513]

Internal file name [OUTPUT/5761_Sunday_June_05_2022_03_53_07_PM_95044538/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 7. Laplace Transforms. Section 7.5 Problems for review and discovery. Section B, Challenge Problems. Page 310

Problem number: 3.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_laplace", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$i'' + 2i' + 3i = \begin{cases} 30 & 0 < t < 2\pi \\ 0 & 2\pi \leq t \leq 5\pi \\ 10 & 5\pi < t < \infty \end{cases}$$

With initial conditions

$$[i(0) = 8, i'(0) = 0]$$

26.1.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$i'' + p(t)i' + q(t)i = F$$

Where here

$$p(t) = 2$$

$$q(t) = 3$$

$$F = \begin{cases} 0 & t \leq 0 \\ 30 & 0 < t < 2\pi \\ 0 & 2\pi \leq t \leq 5\pi \\ 10 & 5\pi < t \end{cases}$$

Hence the ode is

$$i'' + 2i' + 3i = \begin{cases} 0 & t \leq 0 \\ 30 & 0 < t < 2\pi \\ 0 & 2\pi \leq t \leq 5\pi \\ 10 & 5\pi < t \end{cases}$$

The domain of $p(t) = 2$ is

$$\{-\infty < t < \infty\}$$

And the point $t_0 = 0$ is inside this domain. The domain of $q(t) = 3$ is

$$\{-\infty < t < \infty\}$$

And the point $t_0 = 0$ is also inside this domain. The domain of $F = \begin{cases} 0 & t \leq 0 \\ 30 & 0 < t < 2\pi \\ 0 & 2\pi \leq t \leq 5\pi \\ 10 & 5\pi < t \end{cases}$ is

$$\{0 \leq t \leq 2\pi, 2\pi \leq t \leq 5\pi, 5\pi \leq t \leq \infty, -\infty \leq t \leq 0\}$$

And the point $t_0 = 0$ is also inside this domain. Hence solution exists and is unique.

Solving using the Laplace transform method. Let

$$\mathcal{L}(i) = Y(s)$$

Taking the Laplace transform of the ode and using the relations that

$$\begin{aligned}\mathcal{L}(i') &= sY(s) - i(0) \\ \mathcal{L}(i'') &= s^2Y(s) - i'(0) - si(0)\end{aligned}$$

The given ode now becomes an algebraic equation in the Laplace domain

$$s^2Y(s) - i'(0) - si(0) + 2sY(s) - 2i(0) + 3Y(s) = \frac{30 - 30e^{-2\pi s} + 10e^{-5\pi s}}{s} \quad (1)$$

But the initial conditions are

$$\begin{aligned}i(0) &= 8 \\ i'(0) &= 0\end{aligned}$$

Substituting these initial conditions in above in Eq (1) gives

$$s^2Y(s) - 16 - 8s + 2sY(s) + 3Y(s) = \frac{30 - 30e^{-2\pi s} + 10e^{-5\pi s}}{s}$$

Solving the above equation for $Y(s)$ results in

$$Y(s) = -\frac{2(-4s^2 + 15e^{-2\pi s} - 5e^{-5\pi s} - 8s - 15)}{s(s^2 + 2s + 3)}$$

Taking the inverse Laplace transform gives

$$\begin{aligned}i &= \mathcal{L}^{-1}(Y(s)) \\ &= \mathcal{L}^{-1}\left(-\frac{2(-4s^2 + 15e^{-2\pi s} - 5e^{-5\pi s} - 8s - 15)}{s(s^2 + 2s + 3)}\right) \\ &= 10 - e^{-t}\left(\sqrt{2}\sin(t\sqrt{2}) + 2\cos(t\sqrt{2})\right) + \frac{5(i\sqrt{2} + 2)\left(4 - 2i\sqrt{2} - 3e^{-(i\sqrt{2}+1)(t-5\pi)} - (1 - 2i\sqrt{2})e^{-(i\sqrt{2}+1)(t-5\pi)}\right)}{18}\end{aligned}$$

Hence the final solution is

$$\begin{aligned}i &= 10 - e^{-t}\left(\sqrt{2}\sin(t\sqrt{2}) + 2\cos(t\sqrt{2})\right) \\ &\quad + \frac{5(i\sqrt{2} + 2)\left(4 - 2i\sqrt{2} - 3e^{-(i\sqrt{2}+1)(t-5\pi)} - (1 - 2i\sqrt{2})e^{-(i\sqrt{2}+1)(t-5\pi)}\right)}{18} \text{Heaviside}(t - 5\pi) \\ &\quad + \frac{5(i\sqrt{2} + 2)\left(-4 + 2i\sqrt{2} + 3e^{-(i\sqrt{2}+1)(t-2\pi)} + (1 - 2i\sqrt{2})e^{-(i\sqrt{2}+1)(t-2\pi)}\right)}{6} \text{Heaviside}(t - 2\pi)\end{aligned}$$

Simplifying the solution gives

$$\begin{aligned}
 i = 10 + & \frac{5 \operatorname{Heaviside}(t - 2\pi) e^{-(i\sqrt{2}+1)(t-2\pi)} (i\sqrt{2} + 2)}{2} \\
 & - \frac{5 \operatorname{Heaviside}(t - 5\pi) e^{-(i\sqrt{2}+1)(t-5\pi)} (i\sqrt{2} + 2)}{6} \\
 & + \frac{5(-i\sqrt{2} + 2) \operatorname{Heaviside}(t - 2\pi) e^{(i\sqrt{2}-1)(t-2\pi)}}{2} \\
 & + \frac{5(i\sqrt{2} - 2) \operatorname{Heaviside}(t - 5\pi) e^{(i\sqrt{2}-1)(t-5\pi)}}{6} - e^{-t}\sqrt{2} \sin(t\sqrt{2}) \\
 & - 2e^{-t} \cos(t\sqrt{2}) + \frac{10 \operatorname{Heaviside}(t - 5\pi)}{3} - 10 \operatorname{Heaviside}(t - 2\pi)
 \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned}
 i = 10 + & \frac{5 \operatorname{Heaviside}(t - 2\pi) e^{-(i\sqrt{2}+1)(t-2\pi)} (i\sqrt{2} + 2)}{2} \\
 & - \frac{5 \operatorname{Heaviside}(t - 5\pi) e^{-(i\sqrt{2}+1)(t-5\pi)} (i\sqrt{2} + 2)}{6} \\
 & + \frac{5(-i\sqrt{2} + 2) \operatorname{Heaviside}(t - 2\pi) e^{(i\sqrt{2}-1)(t-2\pi)}}{2} \\
 & + \frac{5(i\sqrt{2} - 2) \operatorname{Heaviside}(t - 5\pi) e^{(i\sqrt{2}-1)(t-5\pi)}}{6} - e^{-t}\sqrt{2} \sin(t\sqrt{2}) \\
 & - 2e^{-t} \cos(t\sqrt{2}) + \frac{10 \operatorname{Heaviside}(t - 5\pi)}{3} - 10 \operatorname{Heaviside}(t - 2\pi)
 \end{aligned} \tag{1}$$

Verification of solutions

$$\begin{aligned} i = & 10 + \frac{5 \operatorname{Heaviside}(t - 2\pi) e^{-(i\sqrt{2}+1)(t-2\pi)} (i\sqrt{2} + 2)}{2} \\ & - \frac{5 \operatorname{Heaviside}(t - 5\pi) e^{-(i\sqrt{2}+1)(t-5\pi)} (i\sqrt{2} + 2)}{6} \\ & + \frac{5(-i\sqrt{2} + 2) \operatorname{Heaviside}(t - 2\pi) e^{(i\sqrt{2}-1)(t-2\pi)}}{2} \\ & + \frac{5(i\sqrt{2} - 2) \operatorname{Heaviside}(t - 5\pi) e^{(i\sqrt{2}-1)(t-5\pi)}}{6} - e^{-t} \sqrt{2} \sin(t\sqrt{2}) \\ & - 2e^{-t} \cos(t\sqrt{2}) + \frac{10 \operatorname{Heaviside}(t - 5\pi)}{3} - 10 \operatorname{Heaviside}(t - 2\pi) \end{aligned}$$

Verified OK.

26.1.2 Maple step by step solution

Let's solve

$$\left[i'' + 2i' + 3i = \begin{cases} 0 & t \leq 0 \\ 30 & t < 2\pi \\ 0 & t \leq 5\pi \\ 10 & 5\pi < t \end{cases}, i(0) = 8, i'|_{\{t=0\}} = 0 \right]$$

- Highest derivative means the order of the ODE is 2

i''

- Characteristic polynomial of homogeneous ODE

$$r^2 + 2r + 3 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{(-2) \pm (\sqrt{-8})}{2}$$

- Roots of the characteristic polynomial

$$r = (-1 - I\sqrt{2}, I\sqrt{2} - 1)$$

- 1st solution of the homogeneous ODE

$$i_1(t) = e^{-t} \cos(t\sqrt{2})$$

- 2nd solution of the homogeneous ODE

$$i_2(t) = e^{-t} \sin(t\sqrt{2})$$

- General solution of the ODE

$$i = c_1 i_1(t) + c_2 i_2(t) + i_p(t)$$

- Substitute in solutions of the homogeneous ODE

$$i = c_1 e^{-t} \cos(t\sqrt{2}) + c_2 e^{-t} \sin(t\sqrt{2}) + i_p(t)$$

- Find a particular solution $i_p(t)$ of the ODE

- Use variation of parameters to find i_p here $f(t)$ is the forcing function

$$i_p(t) = -i_1(t) \left(\int \frac{i_2(t)f(t)}{W(i_1(t), i_2(t))} dt \right) + i_2(t) \left(\int \frac{i_1(t)f(t)}{W(i_1(t), i_2(t))} dt \right), f(t) = \begin{cases} 0 & t \leq 0 \\ 30 & 0 < t < 2\pi \\ 0 & 2\pi \leq t \leq 5\pi \\ 10 & 5\pi < t \end{cases}$$

- Wronskian of solutions of the homogeneous equation

$$W(i_1(t), i_2(t)) = \begin{vmatrix} e^{-t} \cos(t\sqrt{2}) & e^{-t} \sin(t\sqrt{2}) \\ -e^{-t} \cos(t\sqrt{2}) - e^{-t}\sqrt{2} \sin(t\sqrt{2}) & -e^{-t} \sin(t\sqrt{2}) + e^{-t}\sqrt{2} \cos(t\sqrt{2}) \end{vmatrix}$$

- Compute Wronskian

$$W(i_1(t), i_2(t)) = \sqrt{2} e^{-2t}$$

- Substitute functions into equation for $i_p(t)$

$$i_p(t) = \frac{\sqrt{2} e^{-t} \left(-\cos(t\sqrt{2}) \int e^t \sin(t\sqrt{2}) \begin{cases} 0 & t \leq 0 \\ 30 & 0 < t < 2\pi \\ 0 & 2\pi \leq t \leq 5\pi \\ 10 & 5\pi < t \end{cases} dt + \sin(t\sqrt{2}) \int e^t \cos(t\sqrt{2}) \begin{cases} 0 & t \leq 0 \\ 30 & 0 < t < 2\pi \\ 0 & 2\pi \leq t \leq 5\pi \\ 10 & 5\pi < t \end{cases} dt \right)}{2}$$

- Compute integrals

$$i_p(t) = 5 \left\{ \begin{array}{l} \sqrt{2} (e^{2\pi} (\sqrt{2} \sin(t\sqrt{2}) - \cos(t\sqrt{2})) - \cos(t\sqrt{2})) \\ \frac{2}{3} + ((\sqrt{2} \sin(t\sqrt{2}) + 2 \cos(t\sqrt{2})) \cos(2\pi\sqrt{2}) - \sin(2\pi\sqrt{2}) (\sqrt{2} \cos(t\sqrt{2}) - 2 \sin(t\sqrt{2}))) \end{array} \right.$$

- Substitute particular solution into general solution to ODE

$$i = c_1 e^{-t} \cos(t\sqrt{2}) + c_2 e^{-t} \sin(t\sqrt{2}) + 5 \left\{ \left(\frac{2}{3} + ((\sqrt{2} \sin(t\sqrt{2}) + 2 \cos(t\sqrt{2})) \cos(2\pi\sqrt{2})) - \right. \right.$$

□ Check validity of solution $i = c_1 e^{-t} \cos(t\sqrt{2}) + c_2 e^{-t} \sin(t\sqrt{2}) + 5 \left\{ \left(\frac{2}{3} + ((\sqrt{2} \sin(t\sqrt{2}) + 2 \cos(t\sqrt{2})) \cos(2\pi\sqrt{2})) - \right. \right.$

- Use initial condition $i(0) = 8$
- $8 = c_1$
- Compute derivative of the solution

$$i' = -c_1 e^{-t} \cos(t\sqrt{2}) - c_1 e^{-t} \sqrt{2} \sin(t\sqrt{2}) - c_2 e^{-t} \sin(t\sqrt{2}) + c_2 e^{-t} \sqrt{2} \cos(t\sqrt{2}) + 5 \left\{ \left((2 \cos(t\sqrt{2})) \cos(2\pi\sqrt{2}) - \right. \right.$$

- Use the initial condition $i'|_{\{t=0\}} = 0$
- $0 = -c_1 + c_2 \sqrt{2}$
- Solve for c_1 and c_2
- $\{c_1 = 8, c_2 = 4\sqrt{2}\}$
- Substitute constant values into general solution and simplify

$$i = 8 e^{-t} \cos(t\sqrt{2}) + 4 e^{-t} \sqrt{2} \sin(t\sqrt{2}) + 5 \left\{ \left(\frac{2}{3} + ((\sqrt{2} \sin(t\sqrt{2}) + 2 \cos(t\sqrt{2})) \cos(2\pi\sqrt{2})) - \right. \right.$$

- Solution to the IVP

$$i = 8e^{-t} \cos(t\sqrt{2}) + 4e^{-t}\sqrt{2} \sin(t\sqrt{2}) + 5 \left\{ \left(\frac{2}{3} + ((\sqrt{2} \sin(t\sqrt{2}) + 2 \cos(t\sqrt{2})) \cos(2\pi\sqrt{2})) \right) \right.$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✗ Solution by Maple

```
dsolve([diff(i(t),t$2)+2*diff(i(t),t)+3*i(t)=piecewise(0<t and t<2*Pi,30,2*Pi<= t and t<= 5*
```

No solution found

✓ Solution by Mathematica

Time used: 0.234 (sec). Leaf size: 297

```
DSolve[{i''[t]+2*i'[t]+3*i[t]==Piecewise[{{30,0<t<2*Pi},{0,2*Pi<= t <= 5*Pi},{10,5*Pi<t<Infi
```

$i(t)$

$$\rightarrow \left\{ \begin{array}{l} e^{-t}(-2 \cos(\sqrt{2}t) + 10e^t - \sqrt{2} \sin(\sqrt{2}t)) \\ 4e^{-t}(2 \cos(\sqrt{2}t) + \sqrt{2} \sin(\sqrt{2}t)) \\ -e^{-t}(2 \cos(\sqrt{2}t) - 10e^{2\pi} \cos(\sqrt{2}(t - 2\pi)) + \sqrt{2}(\sin(\sqrt{2}t) - 5e^{2\pi} \sin(\sqrt{2}(t - 2\pi)))) \\ \frac{1}{3}e^{-t}(-6 \cos(\sqrt{2}t) + 10e^t - 10e^{5\pi} \cos(\sqrt{2}(t - 5\pi)) + 30e^{2\pi} \cos(\sqrt{2}(t - 2\pi)) - 3\sqrt{2} \sin(\sqrt{2}t) - 10e^{5\pi} \sin(\sqrt{2}(t - 5\pi))) \end{array} \right.$$

27 Chapter 10. Systems of First-Order Equations.

Section 10.2 Linear Systems. Page 380

27.1	problem 2(a)	3945
27.2	problem 2(c)	3954
27.3	problem 3(a)	3962
27.4	problem 3(c)	3971
27.5	problem 5	3983
27.6	problem 6(a)	3990

27.1 problem 2(a)

- 27.1.1 Solution using Matrix exponential method 3945
- 27.1.2 Solution using explicit Eigenvalue and Eigenvector method . . . 3946
- 27.1.3 Maple step by step solution 3951

Internal problem ID [6514]

Internal file name [OUTPUT/5762_Sunday_June_05_2022_03_53_30_PM_18803338/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 10. Systems of First-Order Equations. Section 10.2 Linear Systems. Page 380

Problem number: 2(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**system of linear ODEs**"

Solve

$$x'(t) = x(t) + 3y(t)$$

$$y'(t) = 3x(t) + y(t)$$

27.1.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} \frac{e^{-2t}}{2} + \frac{e^{4t}}{2} & \frac{e^{4t}}{2} - \frac{e^{-2t}}{2} \\ \frac{e^{4t}}{2} - \frac{e^{-2t}}{2} & \frac{e^{-2t}}{2} + \frac{e^{4t}}{2} \end{bmatrix}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 \vec{x}_h(t) &= e^{At}\vec{c} \\
 &= \begin{bmatrix} \frac{e^{-2t}}{2} + \frac{e^{4t}}{2} & \frac{e^{4t}}{2} - \frac{e^{-2t}}{2} \\ \frac{e^{4t}}{2} - \frac{e^{-2t}}{2} & \frac{e^{-2t}}{2} + \frac{e^{4t}}{2} \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} \\
 &= \begin{bmatrix} \left(\frac{e^{-2t}}{2} + \frac{e^{4t}}{2}\right)c_1 + \left(\frac{e^{4t}}{2} - \frac{e^{-2t}}{2}\right)c_2 \\ \left(\frac{e^{4t}}{2} - \frac{e^{-2t}}{2}\right)c_1 + \left(\frac{e^{-2t}}{2} + \frac{e^{4t}}{2}\right)c_2 \end{bmatrix} \\
 &= \begin{bmatrix} \frac{(-c_2+c_1)e^{-2t}}{2} + \frac{e^{4t}(c_1+c_2)}{2} \\ \frac{(c_2-c_1)e^{-2t}}{2} + \frac{e^{4t}(c_1+c_2)}{2} \end{bmatrix}
 \end{aligned}$$

Since no forcing function is given, then the final solution is $\vec{x}_h(t)$ above.

27.1.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det\left(\begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}\right) = 0$$

Therefore

$$\det\left(\begin{bmatrix} 1 - \lambda & 3 \\ 3 & 1 - \lambda \end{bmatrix}\right) = 0$$

Which gives the characteristic equation

$$\lambda^2 - 2\lambda - 8 = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = 4$$

$$\lambda_2 = -2$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
-2	1	real eigenvalue
4	1	real eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = -2$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{pmatrix} 1 & 3 \\ 3 & 1 \end{pmatrix} - (-2) \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{pmatrix} 3 & 3 \\ 3 & 3 \end{pmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} 3 & 3 & 0 \\ 3 & 3 & 0 \end{array} \right]$$

$$R_2 = R_2 - R_1 \implies \left[\begin{array}{cc|c} 3 & 3 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{pmatrix} 3 & 3 \\ 0 & 0 \end{pmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = -t\}$

Hence the solution is

$$\begin{bmatrix} -t \\ t \end{bmatrix} = \begin{bmatrix} -t \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} -t \\ t \end{bmatrix} = t \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} -t \\ t \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

Considering the eigenvalue $\lambda_2 = 4$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix} - (4) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} -3 & 3 \\ 3 & -3 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} -3 & 3 & 0 \\ 3 & -3 & 0 \end{array} \right]$$

$$R_2 = R_2 + R_1 \implies \left[\begin{array}{cc|c} -3 & 3 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} -3 & 3 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = t\}$

Hence the solution is

$$\begin{bmatrix} t \\ t \end{bmatrix} = \begin{bmatrix} t \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} t \\ t \end{bmatrix} = t \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} t \\ t \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
4	1	1	No	$\begin{bmatrix} 1 \\ 1 \end{bmatrix}$
-2	1	1	No	$\begin{bmatrix} -1 \\ 1 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. Since eigenvalue 4 is real and distinct then the

corresponding eigenvector solution is

$$\begin{aligned}\vec{x}_1(t) &= \vec{v}_1 e^{4t} \\ &= \begin{bmatrix} 1 \\ 1 \end{bmatrix} e^{4t}\end{aligned}$$

Since eigenvalue -2 is real and distinct then the corresponding eigenvector solution is

$$\begin{aligned}\vec{x}_2(t) &= \vec{v}_2 e^{-2t} \\ &= \begin{bmatrix} -1 \\ 1 \end{bmatrix} e^{-2t}\end{aligned}$$

Therefore the final solution is

$$\vec{x}_h(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

Which is written as

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = c_1 \begin{bmatrix} e^{4t} \\ e^{4t} \end{bmatrix} + c_2 \begin{bmatrix} -e^{-2t} \\ e^{-2t} \end{bmatrix}$$

Which becomes

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} c_1 e^{4t} - c_2 e^{-2t} \\ c_1 e^{4t} + c_2 e^{-2t} \end{bmatrix}$$

The following is the phase plot of the system.




Figure 485: Phase plot

27.1.3 Maple step by step solution

Let's solve

$$[x'(t) = x(t) + 3y(t), y'(t) = 3x(t) + y(t)]$$

- Define vector

$$\vec{x}(t) = \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

- Convert system into a vector equation

$$\vec{x}'(t) = \begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

- System to solve

$$\vec{x}'(t) = \begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix} \cdot \vec{x}(t)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix}$$

- Rewrite the system as

$$\vec{x}'(t) = A \cdot \vec{x}(t)$$

- To solve the system, find the eigenvalues and eigenvectors of A
- Eigenpairs of A

$$\left[\left[-2, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right], \left[4, \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[-2, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{x}_1 = e^{-2t} \cdot \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[4, \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{x}_2 = e^{4t} \cdot \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{x} = c_1 \vec{x}_1 + c_2 \vec{x}_2$$

- Substitute solutions into the general solution

$$\vec{x} = c_1 e^{-2t} \cdot \begin{bmatrix} -1 \\ 1 \end{bmatrix} + c_2 e^{4t} \cdot \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

- Substitute in vector of dependent variables

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} -c_1 e^{-2t} + c_2 e^{4t} \\ c_1 e^{-2t} + c_2 e^{4t} \end{bmatrix}$$

- Solution to the system of ODEs

$$\{x(t) = -c_1 e^{-2t} + c_2 e^{4t}, y(t) = c_1 e^{-2t} + c_2 e^{4t}\}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 35

```
dsolve([diff(x(t),t)=x(t)+3*y(t),diff(y(t),t)=3*x(t)+y(t)],singsol=all)
```

$$\begin{aligned} x(t) &= c_1 e^{-2t} + c_2 e^{4t} \\ y(t) &= -c_1 e^{-2t} + c_2 e^{4t} \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.02 (sec). Leaf size: 68

```
DSolve[{x'[t]==x[t]+3*y[t],y'[t]==3*x[t]+y[t]},{x[t],y[t]},t,IncludeSingularSolutions -> True]
```

$$\begin{aligned} x(t) &\rightarrow \frac{1}{2} e^{-2t} (c_1 (e^{6t} + 1) + c_2 (e^{6t} - 1)) \\ y(t) &\rightarrow \frac{1}{2} e^{-2t} (c_1 (e^{6t} - 1) + c_2 (e^{6t} + 1)) \end{aligned}$$

27.2 problem 2(c)

27.2.1 Solution using Matrix exponential method 3954

27.2.2 Solution using explicit Eigenvalue and Eigenvector method . . . 3955

Internal problem ID [6515]

Internal file name [OUTPUT/5763_Sunday_June_05_2022_03_53_32_PM_75425383/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 10. Systems of First-Order Equations. Section 10.2 Linear Systems. Page 380

Problem number: 2(c).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**system of linear ODEs**"

Solve

$$x'(t) = x(t) + 3y(t)$$

$$y'(t) = 3x(t) + y(t)$$

With initial conditions

$$[x(0) = 5, y(0) = 1]$$

27.2.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} \frac{e^{-2t}}{2} + \frac{e^{4t}}{2} & \frac{e^{4t}}{2} - \frac{e^{-2t}}{2} \\ \frac{e^{4t}}{2} - \frac{e^{-2t}}{2} & \frac{e^{-2t}}{2} + \frac{e^{4t}}{2} \end{bmatrix}$$

Therefore the homogeneous solution is

$$\begin{aligned}\vec{x}_h(t) &= e^{At}\vec{x}_0 \\ &= \begin{bmatrix} \frac{e^{-2t}}{2} + \frac{e^{4t}}{2} & \frac{e^{4t}}{2} - \frac{e^{-2t}}{2} \\ \frac{e^{4t}}{2} - \frac{e^{-2t}}{2} & \frac{e^{-2t}}{2} + \frac{e^{4t}}{2} \end{bmatrix} \begin{bmatrix} 5 \\ 1 \end{bmatrix} \\ &= \begin{bmatrix} 2e^{-2t} + 3e^{4t} \\ 3e^{4t} - 2e^{-2t} \end{bmatrix}\end{aligned}$$

Since no forcing function is given, then the final solution is $\vec{x}_h(t)$ above.

27.2.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det\left(\begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}\right) = 0$$

Therefore

$$\det\left(\begin{bmatrix} 1 - \lambda & 3 \\ 3 & 1 - \lambda \end{bmatrix}\right) = 0$$

Which gives the characteristic equation

$$\lambda^2 - 2\lambda - 8 = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = 4$$

$$\lambda_2 = -2$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
-2	1	real eigenvalue
4	1	real eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = -2$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix} - (-2) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 3 & 3 \\ 3 & 3 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} 3 & 3 & 0 \\ 3 & 3 & 0 \end{array} \right]$$

$$R_2 = R_2 - R_1 \implies \left[\begin{array}{cc|c} 3 & 3 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} 3 & 3 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = -t\}$

Hence the solution is

$$\begin{bmatrix} -t \\ t \end{bmatrix} = \begin{bmatrix} -t \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} -t \\ t \end{bmatrix} = t \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} -t \\ t \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

Considering the eigenvalue $\lambda_2 = 4$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix} - (4) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$
$$\begin{bmatrix} -3 & 3 \\ 3 & -3 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} -3 & 3 & 0 \\ 3 & -3 & 0 \end{array} \right]$$

$$R_2 = R_2 + R_1 \implies \left[\begin{array}{cc|c} -3 & 3 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} -3 & 3 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = t\}$

Hence the solution is

$$\begin{bmatrix} t \\ t \end{bmatrix} = \begin{bmatrix} t \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} t \\ t \end{bmatrix} = t \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} t \\ t \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
4	1	1	No	$\begin{bmatrix} 1 \\ 1 \end{bmatrix}$
-2	1	1	No	$\begin{bmatrix} -1 \\ 1 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. Since eigenvalue 4 is real and distinct then the corresponding eigenvector solution is

$$\begin{aligned} \vec{x}_1(t) &= \vec{v}_1 e^{4t} \\ &= \begin{bmatrix} 1 \\ 1 \end{bmatrix} e^{4t} \end{aligned}$$

Since eigenvalue -2 is real and distinct then the corresponding eigenvector solution is

$$\begin{aligned}\vec{x}_2(t) &= \vec{v}_2 e^{-2t} \\ &= \begin{bmatrix} -1 \\ 1 \end{bmatrix} e^{-2t}\end{aligned}$$

Therefore the final solution is

$$\vec{x}_h(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

Which is written as

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = c_1 \begin{bmatrix} e^{4t} \\ e^{4t} \end{bmatrix} + c_2 \begin{bmatrix} -e^{-2t} \\ e^{-2t} \end{bmatrix}$$

Which becomes

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} c_1 e^{4t} - c_2 e^{-2t} \\ c_1 e^{4t} + c_2 e^{-2t} \end{bmatrix}$$

Since initial conditions are given, the solution above needs to be updated by solving for the constants of integrations using the given initial conditions

$$\begin{bmatrix} x(0) = 5 \\ y(0) = 1 \end{bmatrix} \tag{1}$$

Substituting initial conditions into the above solution at $t = 0$ gives

$$\begin{bmatrix} 5 \\ 1 \end{bmatrix} = \begin{bmatrix} c_1 - c_2 \\ c_1 + c_2 \end{bmatrix}$$

Solving for the constants of integrations gives

$$\begin{bmatrix} c_1 = 3 \\ c_2 = -2 \end{bmatrix}$$

Substituting these constants back in original solution in Eq. (1) gives

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} 2e^{-2t} + 3e^{4t} \\ 3e^{4t} - 2e^{-2t} \end{bmatrix}$$

The following is the phase plot of the system.






Figure 486: Phase plot

The following are plots of each solution.



✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 34

```
dsolve([diff(x(t),t) = x(t)+3*y(t), diff(y(t),t) = 3*x(t)+y(t), x(0) = 5, y(0) = 1], singsol
```

$$\begin{aligned}x(t) &= 2e^{-2t} + 3e^{4t} \\y(t) &= -2e^{-2t} + 3e^{4t}\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 38

```
DSolve[{x'[t]==x[t]+3*y[t],y'[t]==3*x[t]+y[t]},{x[0]==5,y[0]==1},{x[t],y[t]},t,IncludeSingul
```

$$\begin{aligned}x(t) &\rightarrow e^{-2t}(3e^{6t} + 2) \\y(t) &\rightarrow e^{-2t}(3e^{6t} - 2)\end{aligned}$$

27.3 problem 3(a)

- 27.3.1 Solution using Matrix exponential method 3962
- 27.3.2 Solution using explicit Eigenvalue and Eigenvector method . . . 3963
- 27.3.3 Maple step by step solution 3968

Internal problem ID [6516]

Internal file name [OUTPUT/5764_Sunday_June_05_2022_03_53_34_PM_6081269/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 10. Systems of First-Order Equations. Section 10.2 Linear Systems. Page 380

Problem number: 3(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"system of linear ODEs"**

Solve

$$\begin{aligned}x'(t) &= x(t) + 2y(t) \\y'(t) &= 3x(t) + 2y(t)\end{aligned}$$

27.3.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} \frac{3e^{-t}}{5} + \frac{2e^{4t}}{5} & \frac{2e^{4t}}{5} - \frac{2e^{-t}}{5} \\ \frac{3e^{4t}}{5} - \frac{3e^{-t}}{5} & \frac{2e^{-t}}{5} + \frac{3e^{4t}}{5} \end{bmatrix}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 \vec{x}_h(t) &= e^{At} \vec{c} \\
 &= \begin{bmatrix} \frac{3e^{-t}}{5} + \frac{2e^{4t}}{5} & \frac{2e^{4t}}{5} - \frac{2e^{-t}}{5} \\ \frac{3e^{4t}}{5} - \frac{3e^{-t}}{5} & \frac{2e^{-t}}{5} + \frac{3e^{4t}}{5} \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} \\
 &= \begin{bmatrix} \left(\frac{3e^{-t}}{5} + \frac{2e^{4t}}{5}\right) c_1 + \left(\frac{2e^{4t}}{5} - \frac{2e^{-t}}{5}\right) c_2 \\ \left(\frac{3e^{4t}}{5} - \frac{3e^{-t}}{5}\right) c_1 + \left(\frac{2e^{-t}}{5} + \frac{3e^{4t}}{5}\right) c_2 \end{bmatrix} \\
 &= \begin{bmatrix} \frac{(3c_1 - 2c_2)e^{-t}}{5} + \frac{2e^{4t}(c_1 + c_2)}{5} \\ \frac{(-3c_1 + 2c_2)e^{-t}}{5} + \frac{3e^{4t}(c_1 + c_2)}{5} \end{bmatrix}
 \end{aligned}$$

Since no forcing function is given, then the final solution is $\vec{x}_h(t)$ above.

27.3.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A \vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det\left(\begin{bmatrix} 1 & 2 \\ 3 & 2 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}\right) = 0$$

Therefore

$$\det\left(\begin{bmatrix} 1 - \lambda & 2 \\ 3 & 2 - \lambda \end{bmatrix}\right) = 0$$

Which gives the characteristic equation

$$\lambda^2 - 3\lambda - 4 = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = 4$$

$$\lambda_2 = -1$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
-1	1	real eigenvalue
4	1	real eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = -1$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 1 & 2 \\ 3 & 2 \end{bmatrix} - (-1) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 2 \\ 3 & 3 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} 2 & 2 & 0 \\ 3 & 3 & 0 \end{array} \right]$$

$$R_2 = R_2 - \frac{3R_1}{2} \implies \left[\begin{array}{cc|c} 2 & 2 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} 2 & 2 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = -t\}$

Hence the solution is

$$\begin{bmatrix} -t \\ t \end{bmatrix} = \begin{bmatrix} -t \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} -t \\ t \end{bmatrix} = t \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} -t \\ t \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

Considering the eigenvalue $\lambda_2 = 4$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 1 & 2 \\ 3 & 2 \end{bmatrix} - (4) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} -3 & 2 \\ 3 & -2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} -3 & 2 & 0 \\ 3 & -2 & 0 \end{array} \right]$$

$$R_2 = R_2 + R_1 \implies \left[\begin{array}{cc|c} -3 & 2 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} -3 & 2 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = \frac{2t}{3}\}$

Hence the solution is

$$\begin{bmatrix} \frac{2t}{3} \\ t \end{bmatrix} = \begin{bmatrix} \frac{2t}{3} \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} \frac{2t}{3} \\ t \end{bmatrix} = t \begin{bmatrix} \frac{2}{3} \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} \frac{2t}{3} \\ t \end{bmatrix} = \begin{bmatrix} \frac{2}{3} \\ 1 \end{bmatrix}$$

Which is normalized to

$$\begin{bmatrix} \frac{2t}{3} \\ t \end{bmatrix} = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
4	1	1	No	$\begin{bmatrix} \frac{2}{3} \\ 1 \end{bmatrix}$
-1	1	1	No	$\begin{bmatrix} -1 \\ 1 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. Since eigenvalue 4 is real and distinct then the corresponding eigenvector solution is

$$\begin{aligned}\vec{x}_1(t) &= \vec{v}_1 e^{4t} \\ &= \begin{bmatrix} \frac{2}{3} \\ 1 \end{bmatrix} e^{4t}\end{aligned}$$

Since eigenvalue -1 is real and distinct then the corresponding eigenvector solution is

$$\begin{aligned}\vec{x}_2(t) &= \vec{v}_2 e^{-t} \\ &= \begin{bmatrix} -1 \\ 1 \end{bmatrix} e^{-t}\end{aligned}$$

Therefore the final solution is

$$\vec{x}_h(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

Which is written as

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = c_1 \begin{bmatrix} \frac{2e^{4t}}{3} \\ e^{4t} \end{bmatrix} + c_2 \begin{bmatrix} -e^{-t} \\ e^{-t} \end{bmatrix}$$

Which becomes

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} \frac{2c_1 e^{4t}}{3} - c_2 e^{-t} \\ c_1 e^{4t} + c_2 e^{-t} \end{bmatrix}$$

The following is the phase plot of the system.




Figure 487: Phase plot

27.3.3 Maple step by step solution

Let's solve

$$[x'(t) = x(t) + 2y(t), y'(t) = 3x(t) + 2y(t)]$$

- Define vector

$$\vec{x}(t) = \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

- Convert system into a vector equation

$$\vec{x}'(t) = \begin{bmatrix} 1 & 2 \\ 3 & 2 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

- System to solve

$$\vec{x}'(t) = \begin{bmatrix} 1 & 2 \\ 3 & 2 \end{bmatrix} \cdot \vec{x}(t)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 2 \end{bmatrix}$$

- Rewrite the system as

$$\vec{x}'(t) = A \cdot \vec{x}(t)$$

- To solve the system, find the eigenvalues and eigenvectors of A
- Eigenpairs of A

$$\left[\left[-1, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right], \left[4, \begin{bmatrix} \frac{2}{3} \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[-1, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{x}_1 = e^{-t} \cdot \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[4, \begin{bmatrix} \frac{2}{3} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{x}_2 = e^{4t} \cdot \begin{bmatrix} \frac{2}{3} \\ 1 \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{x} = c_1 \vec{x}_1 + c_2 \vec{x}_2$$

- Substitute solutions into the general solution

$$\vec{x} = e^{-t} c_1 \cdot \begin{bmatrix} -1 \\ 1 \end{bmatrix} + c_2 e^{4t} \cdot \begin{bmatrix} \frac{2}{3} \\ 1 \end{bmatrix}$$

- Substitute in vector of dependent variables

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} -e^{-t}c_1 + \frac{2c_2e^{4t}}{3} \\ e^{-t}c_1 + c_2e^{4t} \end{bmatrix}$$

- Solution to the system of ODEs

$$\left\{ x(t) = -e^{-t}c_1 + \frac{2c_2e^{4t}}{3}, y(t) = e^{-t}c_1 + c_2e^{4t} \right\}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 36

```
dsolve([diff(x(t),t)=x(t)+2*y(t),diff(y(t),t)=3*x(t)+2*y(t)],singsol=all)
```

$$\begin{aligned} x(t) &= c_1e^{4t} + c_2e^{-t} \\ y(t) &= \frac{3c_1e^{4t}}{2} - c_2e^{-t} \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 74

```
DSolve[{x'[t]==x[t]+2*y[t],y'[t]==3*x[t]+2*y[t]},{x[t],y[t]},t,IncludeSingularSolutions -> T
```

$$\begin{aligned} x(t) &\rightarrow \frac{1}{5}e^{-t}(c_1(2e^{5t} + 3) + 2c_2(e^{5t} - 1)) \\ y(t) &\rightarrow \frac{1}{5}e^{-t}(3c_1(e^{5t} - 1) + c_2(3e^{5t} + 2)) \end{aligned}$$

27.4 problem 3(c)

27.4.1 Solution using Matrix exponential method	3971
27.4.2 Solution using explicit Eigenvalue and Eigenvector method . . .	3973
27.4.3 Maple step by step solution	3978

Internal problem ID [6517]

Internal file name [OUTPUT/5765_Sunday_June_05_2022_03_53_36_PM_53780418/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 10. Systems of First-Order Equations. Section 10.2 Linear Systems. Page 380

Problem number: 3(c).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**system of linear ODEs**"

Solve

$$\begin{aligned}x'(t) &= x(t) + 2y(t) + t - 1 \\y'(t) &= 3x(t) + 2y(t) - 5t - 2\end{aligned}$$

27.4.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t) + \vec{G}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix} + \begin{bmatrix} t - 1 \\ -5t - 2 \end{bmatrix}$$

Since the system is nonhomogeneous, then the solution is given by

$$\vec{x}(t) = \vec{x}_h(t) + \vec{x}_p(t)$$

Where $\vec{x}_h(t)$ is the homogeneous solution to $\vec{x}'(t) = A\vec{x}(t)$ and $\vec{x}_p(t)$ is a particular solution to $\vec{x}'(t) = A\vec{x}(t) + \vec{G}(t)$. The particular solution will be found using variation

of parameters method applied to the fundamental matrix. For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} \frac{3e^{-t}}{5} + \frac{2e^{4t}}{5} & \frac{2e^{4t}}{5} - \frac{2e^{-t}}{5} \\ \frac{3e^{4t}}{5} - \frac{3e^{-t}}{5} & \frac{2e^{-t}}{5} + \frac{3e^{4t}}{5} \end{bmatrix}$$

Therefore the homogeneous solution is

$$\begin{aligned} \vec{x}_h(t) &= e^{At} \vec{c} \\ &= \begin{bmatrix} \frac{3e^{-t}}{5} + \frac{2e^{4t}}{5} & \frac{2e^{4t}}{5} - \frac{2e^{-t}}{5} \\ \frac{3e^{4t}}{5} - \frac{3e^{-t}}{5} & \frac{2e^{-t}}{5} + \frac{3e^{4t}}{5} \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} \\ &= \begin{bmatrix} \left(\frac{3e^{-t}}{5} + \frac{2e^{4t}}{5}\right) c_1 + \left(\frac{2e^{4t}}{5} - \frac{2e^{-t}}{5}\right) c_2 \\ \left(\frac{3e^{4t}}{5} - \frac{3e^{-t}}{5}\right) c_1 + \left(\frac{2e^{-t}}{5} + \frac{3e^{4t}}{5}\right) c_2 \end{bmatrix} \\ &= \begin{bmatrix} \frac{(3c_1 - 2c_2)e^{-t}}{5} + \frac{2e^{4t}(c_1 + c_2)}{5} \\ \frac{(-3c_1 + 2c_2)e^{-t}}{5} + \frac{3e^{4t}(c_1 + c_2)}{5} \end{bmatrix} \end{aligned}$$

The particular solution given by

$$\vec{x}_p(t) = e^{At} \int e^{-At} \vec{G}(t) dt$$

But

$$\begin{aligned} e^{-At} &= (e^{At})^{-1} \\ &= \begin{bmatrix} \frac{(3e^{5t} + 2)e^{-4t}}{5} & -\frac{2(e^{5t} - 1)e^{-4t}}{5} \\ -\frac{3(e^{5t} - 1)e^{-4t}}{5} & \frac{(2e^{5t} + 3)e^{-4t}}{5} \end{bmatrix} \end{aligned}$$

Hence

$$\begin{aligned} \vec{x}_p(t) &= \begin{bmatrix} \frac{3e^{-t}}{5} + \frac{2e^{4t}}{5} & \frac{2e^{4t}}{5} - \frac{2e^{-t}}{5} \\ \frac{3e^{4t}}{5} - \frac{3e^{-t}}{5} & \frac{2e^{-t}}{5} + \frac{3e^{4t}}{5} \end{bmatrix} \int \begin{bmatrix} \frac{(3e^{5t} + 2)e^{-4t}}{5} & -\frac{2(e^{5t} - 1)e^{-4t}}{5} \\ -\frac{3(e^{5t} - 1)e^{-4t}}{5} & \frac{(2e^{5t} + 3)e^{-4t}}{5} \end{bmatrix} \begin{bmatrix} t - 1 \\ -5t - 2 \end{bmatrix} dt \\ &= \begin{bmatrix} \frac{3e^{-t}}{5} + \frac{2e^{4t}}{5} & \frac{2e^{4t}}{5} - \frac{2e^{-t}}{5} \\ \frac{3e^{4t}}{5} - \frac{3e^{-t}}{5} & \frac{2e^{-t}}{5} + \frac{3e^{4t}}{5} \end{bmatrix} \begin{bmatrix} \frac{(13e^{5t}t - 12e^{5t} + 2t + 2)e^{-4t}}{5} \\ -\frac{(13e^{5t}t - 12e^{5t} - 3t - 3)e^{-4t}}{5} \end{bmatrix} \\ &= \begin{bmatrix} 3t - 2 \\ 3 - 2t \end{bmatrix} \end{aligned}$$

Hence the complete solution is

$$\begin{aligned}\vec{x}(t) &= \vec{x}_h(t) + \vec{x}_p(t) \\ &= \begin{bmatrix} \frac{(3c_1-2c_2)e^{-t}}{5} + \frac{(2c_1+2c_2)e^{4t}}{5} + 3t - 2 \\ \frac{(-3c_1+2c_2)e^{-t}}{5} + \frac{(3c_1+3c_2)e^{4t}}{5} - 2t + 3 \end{bmatrix}\end{aligned}$$

27.4.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t) + \vec{G}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix} + \begin{bmatrix} t - 1 \\ -5t - 2 \end{bmatrix}$$

Since the system is nonhomogeneous, then the solution is given by

$$\vec{x}(t) = \vec{x}_h(t) + \vec{x}_p(t)$$

Where $\vec{x}_h(t)$ is the homogeneous solution to $\vec{x}'(t) = A\vec{x}(t)$ and $\vec{x}_p(t)$ is a particular solution to $\vec{x}'(t) = A\vec{x}(t) + \vec{G}(t)$. The particular solution will be found using variation of parameters method applied to the fundamental matrix.

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det\left(\begin{bmatrix} 1 & 2 \\ 3 & 2 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}\right) = 0$$

Therefore

$$\det\left(\begin{bmatrix} 1 - \lambda & 2 \\ 3 & 2 - \lambda \end{bmatrix}\right) = 0$$

Which gives the characteristic equation

$$\lambda^2 - 3\lambda - 4 = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = -1$$

$$\lambda_2 = 4$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
-1	1	real eigenvalue
4	1	real eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = -1$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 1 & 2 \\ 3 & 2 \end{bmatrix} - (-1) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 2 \\ 3 & 3 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} 2 & 2 & 0 \\ 3 & 3 & 0 \end{array} \right]$$

$$R_2 = R_2 - \frac{3R_1}{2} \implies \left[\begin{array}{cc|c} 2 & 2 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} 2 & 2 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = -t\}$

Hence the solution is

$$\begin{bmatrix} -t \\ t \end{bmatrix} = \begin{bmatrix} -t \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} -t \\ t \end{bmatrix} = t \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} -t \\ t \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

Considering the eigenvalue $\lambda_2 = 4$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 1 & 2 \\ 3 & 2 \end{bmatrix} - (4) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$
$$\begin{bmatrix} -3 & 2 \\ 3 & -2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} -3 & 2 & 0 \\ 3 & -2 & 0 \end{array} \right]$$

$$R_2 = R_2 + R_1 \implies \left[\begin{array}{cc|c} -3 & 2 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} -3 & 2 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = \frac{2t}{3}\}$

Hence the solution is

$$\begin{bmatrix} \frac{2t}{3} \\ t \end{bmatrix} = \begin{bmatrix} \frac{2t}{3} \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} \frac{2t}{3} \\ t \end{bmatrix} = t \begin{bmatrix} \frac{2}{3} \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} \frac{2t}{3} \\ t \end{bmatrix} = \begin{bmatrix} \frac{2}{3} \\ 1 \end{bmatrix}$$

Which is normalized to

$$\begin{bmatrix} \frac{2t}{3} \\ t \end{bmatrix} = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
-1	1	1	No	$\begin{bmatrix} -1 \\ 1 \end{bmatrix}$
4	1	1	No	$\begin{bmatrix} \frac{2}{3} \\ 1 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. Since eigenvalue -1 is real and distinct then the

corresponding eigenvector solution is

$$\begin{aligned}\vec{x}_1(t) &= \vec{v}_1 e^{-t} \\ &= \begin{bmatrix} -1 \\ 1 \end{bmatrix} e^{-t}\end{aligned}$$

Since eigenvalue 4 is real and distinct then the corresponding eigenvector solution is

$$\begin{aligned}\vec{x}_2(t) &= \vec{v}_2 e^{4t} \\ &= \begin{bmatrix} \frac{2}{3} \\ 1 \end{bmatrix} e^{4t}\end{aligned}$$

Therefore the homogeneous solution is

$$\vec{x}_h(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

Which is written as

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = c_1 \begin{bmatrix} -e^{-t} \\ e^{-t} \end{bmatrix} + c_2 \begin{bmatrix} \frac{2e^{4t}}{3} \\ e^{4t} \end{bmatrix}$$

Now that we found homogeneous solution above, we need to find a particular solution $\vec{x}_p(t)$. We will use Variation of parameters. The fundamental matrix is

$$\Phi = \begin{bmatrix} \vec{x}_1 & \vec{x}_2 & \cdots \end{bmatrix}$$

Where \vec{x}_i are the solution basis found above. Therefore the fundamental matrix is

$$\Phi(t) = \begin{bmatrix} -e^{-t} & \frac{2e^{4t}}{3} \\ e^{-t} & e^{4t} \end{bmatrix}$$

The particular solution is then given by

$$\vec{x}_p(t) = \Phi \int \Phi^{-1} \vec{G}(t) dt$$

But

$$\Phi^{-1} = \begin{bmatrix} -\frac{3e^t}{5} & \frac{2e^t}{5} \\ \frac{3e^{-4t}}{5} & \frac{3e^{-4t}}{5} \end{bmatrix}$$

Hence

$$\begin{aligned}
 \vec{x}_p(t) &= \begin{bmatrix} -e^{-t} & \frac{2e^{4t}}{3} \\ e^{-t} & e^{4t} \end{bmatrix} \int \begin{bmatrix} -\frac{3e^t}{5} & \frac{2e^t}{5} \\ \frac{3e^{-4t}}{5} & \frac{3e^{-4t}}{5} \end{bmatrix} \begin{bmatrix} t-1 \\ -5t-2 \end{bmatrix} dt \\
 &= \begin{bmatrix} -e^{-t} & \frac{2e^{4t}}{3} \\ e^{-t} & e^{4t} \end{bmatrix} \int \begin{bmatrix} -\frac{e^t(13t+1)}{5} \\ -\frac{3e^{-4t}(4t+3)}{5} \end{bmatrix} dt \\
 &= \begin{bmatrix} -e^{-t} & \frac{2e^{4t}}{3} \\ e^{-t} & e^{4t} \end{bmatrix} \begin{bmatrix} -\frac{e^t(13t-12)}{5} \\ \frac{3e^{-4t}(t+1)}{5} \end{bmatrix} \\
 &= \begin{bmatrix} 3t-2 \\ 3-2t \end{bmatrix}
 \end{aligned}$$

Now that we found particular solution, the final solution is

$$\begin{aligned}
 \vec{x}(t) &= \vec{x}_h(t) + \vec{x}_p(t) \\
 \begin{bmatrix} x(t) \\ y(t) \end{bmatrix} &= \begin{bmatrix} -c_1e^{-t} \\ c_1e^{-t} \end{bmatrix} + \begin{bmatrix} \frac{2c_2e^{4t}}{3} \\ c_2e^{4t} \end{bmatrix} + \begin{bmatrix} 3t-2 \\ 3-2t \end{bmatrix}
 \end{aligned}$$

Which becomes

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} -c_1e^{-t} + \frac{2c_2e^{4t}}{3} + 3t-2 \\ c_1e^{-t} + c_2e^{4t} + 3-2t \end{bmatrix}$$

27.4.3 Maple step by step solution

Let's solve

$$[x'(t) = x(t) + 2y(t) + t - 1, y'(t) = 3x(t) + 2y(t) - 5t - 2]$$

- Define vector

$$\vec{x}(t) = \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

- Convert system into a vector equation

$$\vec{x}'(t) = \begin{bmatrix} 1 & 2 \\ 3 & 2 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} t-1 \\ -5t-2 \end{bmatrix}$$

- System to solve

$$\vec{x}'(t) = \begin{bmatrix} 1 & 2 \\ 3 & 2 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} t-1 \\ -5t-2 \end{bmatrix}$$

- Define the forcing function

$$\vec{f}(t) = \begin{bmatrix} t-1 \\ -5t-2 \end{bmatrix}$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 2 \end{bmatrix}$$

- Rewrite the system as

$$\vec{x}'(t) = A \cdot \vec{x}(t) + \vec{f}$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[-1, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right], \left[4, \begin{bmatrix} \frac{2}{3} \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[-1, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{x}_1 = e^{-t} \cdot \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[4, \begin{bmatrix} \frac{2}{3} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{x}_2 = e^{4t} \cdot \begin{bmatrix} \frac{2}{3} \\ 1 \end{bmatrix}$$

- General solution of the system of ODEs can be written in terms of the particular solution $\vec{x}_p(t)$

$$\vec{x}(t) = c_1 \vec{x}_1 + c_2 \vec{x}_2 + \vec{x}_p(t)$$

□ Fundamental matrix

- Let $\phi(t)$ be the matrix whose columns are the independent solutions of the homogeneous system

$$\phi(t) = \begin{bmatrix} -e^{-t} & \frac{2e^{4t}}{3} \\ e^{-t} & e^{4t} \end{bmatrix}$$

- The fundamental matrix, $\Phi(t)$ is a normalized version of $\phi(t)$ satisfying $\Phi(0) = I$ where I is the identity matrix

$$\Phi(t) = \phi(t) \cdot \frac{1}{\phi(0)}$$

- Substitute the value of $\phi(t)$ and $\phi(0)$

$$\Phi(t) = \begin{bmatrix} -e^{-t} & \frac{2e^{4t}}{3} \\ e^{-t} & e^{4t} \end{bmatrix} \cdot \frac{1}{\begin{bmatrix} -1 & \frac{2}{3} \\ 1 & 1 \end{bmatrix}}$$

- Evaluate and simplify to get the fundamental matrix

$$\Phi(t) = \begin{bmatrix} \frac{3e^{-t}}{5} + \frac{2e^{4t}}{5} & \frac{2e^{4t}}{5} - \frac{2e^{-t}}{5} \\ \frac{3e^{4t}}{5} - \frac{3e^{-t}}{5} & \frac{2e^{-t}}{5} + \frac{3e^{4t}}{5} \end{bmatrix}$$

□ Find a particular solution of the system of ODEs using variation of parameters

- Let the particular solution be the fundamental matrix multiplied by $\vec{v}(t)$ and solve for $\vec{v}(t)$

$$\vec{x}_p(t) = \Phi(t) \cdot \vec{v}(t)$$

- Take the derivative of the particular solution

$$\vec{x}'_p(t) = \Phi'(t) \cdot \vec{v}(t) + \Phi(t) \cdot \vec{v}'(t)$$

- Substitute particular solution and its derivative into the system of ODEs

$$\Phi'(t) \cdot \vec{v}(t) + \Phi(t) \cdot \vec{v}'(t) = A \cdot \Phi(t) \cdot \vec{v}(t) + \vec{f}(t)$$

- The fundamental matrix has columns that are solutions to the homogeneous system so its derivative is $A \cdot \Phi(t)$

$$A \cdot \Phi(t) \cdot \vec{v}(t) + \Phi(t) \cdot \vec{v}'(t) = A \cdot \Phi(t) \cdot \vec{v}(t) + \vec{f}(t)$$

- Cancel like terms

$$\Phi(t) \cdot \vec{v}'(t) = \vec{f}(t)$$

- Multiply by the inverse of the fundamental matrix

$$\vec{v}'(t) = \frac{1}{\Phi(t)} \cdot \vec{f}(t)$$

- Integrate to solve for $\vec{v}(t)$

$$\vec{v}(t) = \int_0^t \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds$$

- Plug $\vec{v}(t)$ into the equation for the particular solution

$$\vec{x}_p(t) = \Phi(t) \cdot \left(\int_0^t \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds \right)$$

- Plug in the fundamental matrix and the forcing function and compute

$$\vec{x}_p(t) = \begin{bmatrix} 3t + \frac{12e^{-t}}{5} - 2 - \frac{2e^{4t}}{5} \\ -\frac{3e^{4t}}{5} - 2t + 3 - \frac{12e^{-t}}{5} \end{bmatrix}$$

- Plug particular solution back into general solution

$$\vec{x}(t) = c_1 \vec{x}_1 + c_2 \vec{x}_2 + \begin{bmatrix} 3t + \frac{12e^{-t}}{5} - 2 - \frac{2e^{4t}}{5} \\ -\frac{3e^{4t}}{5} - 2t + 3 - \frac{12e^{-t}}{5} \end{bmatrix}$$

- Substitute in vector of dependent variables

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} \frac{(-15c_1+36)e^{-t}}{15} + \frac{(10c_2-6)e^{4t}}{15} + 3t - 2 \\ \frac{(-12+5c_1)e^{-t}}{5} + \frac{(-3+5c_2)e^{4t}}{5} - 2t + 3 \end{bmatrix}$$

- Solution to the system of ODEs

$$\left\{ x(t) = \frac{(-15c_1+36)e^{-t}}{15} + \frac{(10c_2-6)e^{4t}}{15} + 3t - 2, y(t) = \frac{(-12+5c_1)e^{-t}}{5} + \frac{(-3+5c_2)e^{4t}}{5} - 2t + 3 \right\}$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 44

```
dsolve([diff(x(t),t)=x(t)+2*y(t)+t-1,diff(y(t),t)=3*x(t)+2*y(t)-5*t-2],singsol=all)
```

$$x(t) = c_2 e^{4t} + e^{-t} c_1 + 3t - 2$$

$$y(t) = \frac{3c_2 e^{4t}}{2} - e^{-t} c_1 + 3 - 2t$$

✓ Solution by Mathematica

Time used: 0.188 (sec). Leaf size: 88

```
DSolve[{x'[t]==x[t]+2*y[t]+t-1,y'[t]==3*x[t]+2*y[t]-5*t-2},{x[t],y[t]},t,IncludeSingularSolu
```

$$x(t) \rightarrow \frac{1}{5}e^{-t}(5e^t(3t-2) + 2(c_1 + c_2)e^{5t} + 3c_1 - 2c_2)$$

$$y(t) \rightarrow \frac{1}{5}e^{-t}(-5e^t(2t-3) + 3(c_1 + c_2)e^{5t} - 3c_1 + 2c_2)$$

27.5 problem 5

27.5.1 Solution using Matrix exponential method 3983

27.5.2 Solution using explicit Eigenvalue and Eigenvector method . . . 3984

Internal problem ID [6518]

Internal file name [OUTPUT/5766_Sunday_June_05_2022_03_53_38_PM_94886050/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 10. Systems of First-Order Equations. Section 10.2 Linear Systems. Page 380

Problem number: 5.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**system of linear ODEs**"

Solve

$$x'(t) = x(t) + y(t)$$

$$y'(t) = y(t)$$

27.5.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A \vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} e^t & t e^t \\ 0 & e^t \end{bmatrix}$$

Therefore the homogeneous solution is

$$\begin{aligned}\vec{x}_h(t) &= e^{At}\vec{c} \\ &= \begin{bmatrix} e^t & t e^t \\ 0 & e^t \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} \\ &= \begin{bmatrix} e^t c_1 + t e^t c_2 \\ e^t c_2 \end{bmatrix} \\ &= \begin{bmatrix} e^t(c_2 t + c_1) \\ e^t c_2 \end{bmatrix}\end{aligned}$$

Since no forcing function is given, then the final solution is $\vec{x}_h(t)$ above.

27.5.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det\left(\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}\right) = 0$$

Therefore

$$\det\left(\begin{bmatrix} 1 - \lambda & 1 \\ 0 & 1 - \lambda \end{bmatrix}\right) = 0$$

Since the matrix A is triangular matrix, then the determinant is the product of the elements along the diagonal. Therefore the above becomes

$$(1 - \lambda)(1 - \lambda) = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = 1$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
1	1	real eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = 1$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} - (1) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} 0 & 1 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_1\}$ and the leading variables are $\{v_2\}$. Let $v_1 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_2 = 0\}$

Hence the solution is

$$\begin{bmatrix} t \\ 0 \end{bmatrix} = \begin{bmatrix} t \\ 0 \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} t \\ 0 \end{bmatrix} = t \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} t \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
1	2	1	Yes	$\begin{bmatrix} 1 \\ 0 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. eigenvalue 1 is real and repeated eigenvalue of multiplicity 2. There are two possible cases that can happen. This is illustrated in this diagram




Figure 488: Possible case for repeated λ of multiplicity 2

This eigenvalue has algebraic multiplicity of 2, and geometric multiplicity 1, therefore this is defective eigenvalue. The defect is 1. This falls into case 2 shown above. We need to generate the missing additional generalized eigenvector \vec{v}_2 by solving

$$(A - \lambda I) \vec{v}_2 = \vec{v}_1$$

Where \vec{v}_1 is the normal (rank 1) eigenvector found above. Hence we need to solve

$$\left(\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} - (1) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

Solving for \vec{v}_2 gives

$$\vec{v}_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

We have found two generalized eigenvectors for eigenvalue 1. Therefore the two basis solution associated with this eigenvalue are

$$\begin{aligned}\vec{x}_1(t) &= \vec{v}_1 e^{\lambda t} \\ &= \begin{bmatrix} 1 \\ 0 \end{bmatrix} e^t \\ &= \begin{bmatrix} e^t \\ 0 \end{bmatrix}\end{aligned}$$

And

$$\begin{aligned}\vec{x}_2(t) &= (\vec{v}_1 t + \vec{v}_2) e^{\lambda t} \\ &= \left(\begin{bmatrix} 1 \\ 0 \end{bmatrix} t + \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right) e^t \\ &= \begin{bmatrix} e^t(t+1) \\ e^t \end{bmatrix}\end{aligned}$$

Therefore the final solution is

$$\vec{x}_h(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

Which is written as

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = c_1 \begin{bmatrix} e^t \\ 0 \end{bmatrix} + c_2 \begin{bmatrix} e^t(t+1) \\ e^t \end{bmatrix}$$

Which becomes

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} e^t(c_2 t + c_1 + c_2) \\ c_2 e^t \end{bmatrix}$$

The following is the phase plot of the system.




Figure 489: Phase plot

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 20

```
dsolve([diff(x(t),t)=x(t)+y(t),diff(y(t),t)=y(t)],singsol=all)
```

$$\begin{aligned}x(t) &= e^t(c_2t + c_1) \\ y(t) &= c_2e^t\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.05 (sec). Leaf size: 25

```
DSolve[{x'[t]==x[t]+y[t],y'[t]==y[t]},{x[t],y[t]},t,IncludeSingularSolutions -> True]
```

$$\begin{aligned}x(t) &\rightarrow e^t(c_2t + c_1) \\ y(t) &\rightarrow c_2e^t\end{aligned}$$

27.6 problem 6(a)

- 27.6.1 Solution using Matrix exponential method 3990
- 27.6.2 Solution using explicit Eigenvalue and Eigenvector method . . . 3991
- 27.6.3 Maple step by step solution 3995

Internal problem ID [6519]

Internal file name [OUTPUT/5767_Sunday_June_05_2022_03_53_40_PM_39283079/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 10. Systems of First-Order Equations. Section 10.2 Linear Systems. Page 380

Problem number: 6(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**system of linear ODEs**"

Solve

$$x'(t) = x(t)$$

$$y'(t) = y(t)$$

27.6.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} e^t & 0 \\ 0 & e^t \end{bmatrix}$$

Therefore the homogeneous solution is

$$\begin{aligned}\vec{x}_h(t) &= e^{At}\vec{c} \\ &= \begin{bmatrix} e^t & 0 \\ 0 & e^t \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} \\ &= \begin{bmatrix} e^t c_1 \\ e^t c_2 \end{bmatrix}\end{aligned}$$

Since no forcing function is given, then the final solution is $\vec{x}_h(t)$ above.

27.6.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det\left(\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}\right) = 0$$

Therefore

$$\det\left(\begin{bmatrix} 1 - \lambda & 0 \\ 0 & 1 - \lambda \end{bmatrix}\right) = 0$$

Since the matrix A is triangular matrix, then the determinant is the product of the elements along the diagonal. Therefore the above becomes

$$(1 - \lambda)(1 - \lambda) = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = 1$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
1	1	real eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = 1$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} - (1) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} 0 & 0 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_1, v_2\}$ and there are no leading variables. Let $v_1 = t$. Let $v_2 = s$. Since there are two free Variable, we have found two eigenvectors associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} t \\ s \end{bmatrix} = \begin{bmatrix} t \\ 0 \end{bmatrix} + \begin{bmatrix} 0 \\ s \end{bmatrix}$$

$$= t \begin{bmatrix} 1 \\ 0 \end{bmatrix} + s \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

By letting $t = 1$ and $s = 1$ then the above becomes

$$\begin{bmatrix} t \\ s \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \end{bmatrix} + \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

Hence the two eigenvectors associated with this eigenvalue are

$$\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right)$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
1	2	2	No	$\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. eigenvalue 1 is real and repeated eigenvalue of multiplicity 2. There are two possible cases that can happen. This is illustrated in this diagram




Figure 490: Possible case for repeated λ of multiplicity 2

This eigenvalue has algebraic multiplicity of 2 which is the same as its geometric multiplicity 2, then it is complete eigenvalue and this falls into case 1 shown above. Hence the corresponding eigenvector basis are

$$\begin{aligned} \vec{x}_1(t) &= \vec{v}_1 e^t \\ &= \begin{bmatrix} 0 \\ 1 \end{bmatrix} e^t \end{aligned}$$

$$\begin{aligned} \vec{x}_2(t) &= \vec{v}_2 e^t \\ &= \begin{bmatrix} 1 \\ 0 \end{bmatrix} e^t \end{aligned}$$

Therefore the final solution is

$$\vec{x}_h(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

Which is written as

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = c_1 \begin{bmatrix} 0 \\ e^t \end{bmatrix} + c_2 \begin{bmatrix} e^t \\ 0 \end{bmatrix}$$

Which becomes

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} c_2 e^t \\ c_1 e^t \end{bmatrix}$$

The following is the phase plot of the system.




Figure 491: Phase plot

27.6.3 Maple step by step solution

Let's solve

$$[x'(t) = x(t), y'(t) = y(t)]$$

- Define vector

$$\vec{x}(t) = \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

- Convert system into a vector equation

$$\vec{x}'(t) = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

- System to solve

$$\vec{x}'(t) = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \cdot \vec{x}(t)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

- Rewrite the system as

$$\vec{x}'(t) = A \cdot \vec{x}(t)$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[1, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right], \left[1, \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right] \right]$$

- Consider eigenpair, with eigenvalue of algebraic multiplicity 2

$$\left[1, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right]$$

- First solution from eigenvalue 1

$$\vec{x}_1(t) = e^t \cdot \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

- Form of the 2nd homogeneous solution where \vec{p} is to be solved for, $\lambda = 1$ is the eigenvalue, and

$$\vec{x}_2(t) = e^{\lambda t} (t\vec{v} + \vec{p})$$

- Note that the t multiplying \vec{v} makes this solution linearly independent to the 1st solution obtained

- Substitute $\vec{x}_2(t)$ into the homogeneous system

$$\lambda e^{\lambda t} (t\vec{v} + \vec{p}) + e^{\lambda t} \vec{v} = (e^{\lambda t} A) \cdot (t\vec{v} + \vec{p})$$

- Use the fact that \vec{v} is an eigenvector of A

$$\lambda e^{\lambda t} (t\vec{v} + \vec{p}) + e^{\lambda t} \vec{v} = e^{\lambda t} (\lambda t\vec{v} + A \cdot \vec{p})$$

- Simplify equation

$$\lambda \vec{p} + \vec{v} = A \cdot \vec{p}$$

- Make use of the identity matrix I

$$(\lambda \cdot I) \cdot \vec{p} + \vec{v} = A \cdot \vec{p}$$

- Condition \vec{p} must meet for $\vec{x}_2(t)$ to be a solution to the homogeneous system

$$(A - \lambda \cdot I) \cdot \vec{p} = \vec{v}$$

- Choose \vec{p} to use in the second solution to the homogeneous system from eigenvalue 1

$$\left(\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} - 1 \cdot \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \cdot \vec{p} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

- Choice of \vec{p}

$$\vec{p} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

- Second solution from eigenvalue 1

$$\vec{x}_2(t) = e^t \cdot \left(t \cdot \begin{bmatrix} 0 \\ 1 \end{bmatrix} + \begin{bmatrix} 0 \\ 0 \end{bmatrix} \right)$$

- General solution to the system of ODEs

$$\vec{x} = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

- Substitute solutions into the general solution

$$\vec{x} = c_1 e^t \cdot \begin{bmatrix} 0 \\ 1 \end{bmatrix} + c_2 e^t \cdot \left(t \cdot \begin{bmatrix} 0 \\ 1 \end{bmatrix} + \begin{bmatrix} 0 \\ 0 \end{bmatrix} \right)$$

- Substitute in vector of dependent variables

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} 0 \\ e^t (tc_2 + c_1) \end{bmatrix}$$

- Solution to the system of ODEs

$$\{x(t) = 0, y(t) = e^t(tc_2 + c_1)\}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve([diff(x(t),t)=x(t),diff(y(t),t)=y(t)],singsol=all)
```

$$x(t) = c_2 e^t$$

$$y(t) = c_1 e^t$$

✓ Solution by Mathematica

Time used: 0.04 (sec). Leaf size: 57

```
DSolve[{x'[t]==x[t],y'[t]==y[t]},{x[t],y[t]},t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow c_1 e^t$$

$$y(t) \rightarrow c_2 e^t$$

$$x(t) \rightarrow c_1 e^t$$

$$y(t) \rightarrow 0$$

$$x(t) \rightarrow 0$$

$$y(t) \rightarrow c_2 e^t$$

$$x(t) \rightarrow 0$$

$$y(t) \rightarrow 0$$

**28 Chapter 10. Systems of First-Order Equations.
Section 10.3 Homogeneous Linear Systems with
Constant Coefficients. Page 387**

28.1	problem 1(a)	4000
28.2	problem 1(b)	4009
28.3	problem 1(c)	4018
28.4	problem 1(d)	4028
28.5	problem 1(e)	4037
28.6	problem 1(f)	4046
28.7	problem 1(g)	4056
28.8	problem 1(h)	4065
28.9	problem 5(b)	4074

28.1 problem 1(a)

28.1.1 Solution using Matrix exponential method	4000
28.1.2 Solution using explicit Eigenvalue and Eigenvector method . . .	4001
28.1.3 Maple step by step solution	4006

Internal problem ID [6520]

Internal file name [OUTPUT/5768_Sunday_June_05_2022_03_53_41_PM_17801438/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 10. Systems of First-Order Equations. Section 10.3 Homogeneous Linear Systems with Constant Coefficients. Page 387

Problem number: 1(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**system of linear ODEs**"

Solve

$$x'(t) = -3x(t) + 4y(t)$$

$$y'(t) = -2x(t) + 3y(t)$$

28.1.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} -3 & 4 \\ -2 & 3 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} 2e^{-t} - e^t & 2e^t - 2e^{-t} \\ -e^t + e^{-t} & -e^{-t} + 2e^t \end{bmatrix}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 \vec{x}_h(t) &= e^{At} \vec{c} \\
 &= \begin{bmatrix} 2e^{-t} - e^t & 2e^t - 2e^{-t} \\ -e^t + e^{-t} & -e^{-t} + 2e^t \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} \\
 &= \begin{bmatrix} (2e^{-t} - e^t)c_1 + (2e^t - 2e^{-t})c_2 \\ (-e^t + e^{-t})c_1 + (-e^{-t} + 2e^t)c_2 \end{bmatrix} \\
 &= \begin{bmatrix} (2c_1 - 2c_2)e^{-t} - e^t(-2c_2 + c_1) \\ (-c_2 + c_1)e^{-t} - e^t(-2c_2 + c_1) \end{bmatrix}
 \end{aligned}$$

Since no forcing function is given, then the final solution is $\vec{x}_h(t)$ above.

28.1.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} -3 & 4 \\ -2 & 3 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det\left(\begin{bmatrix} -3 & 4 \\ -2 & 3 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}\right) = 0$$

Therefore

$$\det\left(\begin{bmatrix} -3 - \lambda & 4 \\ -2 & 3 - \lambda \end{bmatrix}\right) = 0$$

Which gives the characteristic equation

$$\lambda^2 - 1 = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = -1$$

$$\lambda_2 = 1$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
-1	1	real eigenvalue
1	1	real eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = -1$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} -3 & 4 \\ -2 & 3 \end{bmatrix} - (-1) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$
$$\begin{bmatrix} -2 & 4 \\ -2 & 4 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} -2 & 4 & 0 \\ -2 & 4 & 0 \end{array} \right]$$

$$R_2 = R_2 - R_1 \implies \left[\begin{array}{cc|c} -2 & 4 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} -2 & 4 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = 2t\}$

Hence the solution is

$$\begin{bmatrix} 2t \\ t \end{bmatrix} = \begin{bmatrix} 2t \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} 2t \\ t \end{bmatrix} = t \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} 2t \\ t \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

Considering the eigenvalue $\lambda_2 = 1$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} -3 & 4 \\ -2 & 3 \end{bmatrix} - (1) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} -4 & 4 \\ -2 & 2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} -4 & 4 & 0 \\ -2 & 2 & 0 \end{array} \right]$$

$$R_2 = R_2 - \frac{R_1}{2} \implies \left[\begin{array}{cc|c} -4 & 4 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} -4 & 4 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = t\}$

Hence the solution is

$$\begin{bmatrix} t \\ t \end{bmatrix} = \begin{bmatrix} t \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} t \\ t \end{bmatrix} = t \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} t \\ t \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
-1	1	1	No	$\begin{bmatrix} 2 \\ 1 \end{bmatrix}$
1	1	1	No	$\begin{bmatrix} 1 \\ 1 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. Since eigenvalue -1 is real and distinct then the

corresponding eigenvector solution is

$$\begin{aligned}\vec{x}_1(t) &= \vec{v}_1 e^{-t} \\ &= \begin{bmatrix} 2 \\ 1 \end{bmatrix} e^{-t}\end{aligned}$$

Since eigenvalue 1 is real and distinct then the corresponding eigenvector solution is

$$\begin{aligned}\vec{x}_2(t) &= \vec{v}_2 e^t \\ &= \begin{bmatrix} 1 \\ 1 \end{bmatrix} e^t\end{aligned}$$

Therefore the final solution is

$$\vec{x}_h(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

Which is written as

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = c_1 \begin{bmatrix} 2e^{-t} \\ e^{-t} \end{bmatrix} + c_2 \begin{bmatrix} e^t \\ e^t \end{bmatrix}$$

Which becomes

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} 2c_1 e^{-t} + c_2 e^t \\ c_1 e^{-t} + c_2 e^t \end{bmatrix}$$

The following is the phase plot of the system.




Figure 492: Phase plot

28.1.3 Maple step by step solution

Let's solve

$$[x'(t) = -3x(t) + 4y(t), y'(t) = -2x(t) + 3y(t)]$$

- Define vector

$$\vec{x}(t) = \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

- Convert system into a vector equation

$$\vec{x}'(t) = \begin{bmatrix} -3 & 4 \\ -2 & 3 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

- System to solve

$$\vec{x}'(t) = \begin{bmatrix} -3 & 4 \\ -2 & 3 \end{bmatrix} \cdot \vec{x}(t)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} -3 & 4 \\ -2 & 3 \end{bmatrix}$$

- Rewrite the system as

$$\vec{x}'(t) = A \cdot \vec{x}(t)$$

- To solve the system, find the eigenvalues and eigenvectors of A
- Eigenpairs of A

$$\left[\left[-1, \begin{bmatrix} 2 \\ 1 \end{bmatrix} \right], \left[1, \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[-1, \begin{bmatrix} 2 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{x}_1 = e^{-t} \cdot \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[1, \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{x}_2 = e^t \cdot \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{x} = c_1 \vec{x}_1 + c_2 \vec{x}_2$$

- Substitute solutions into the general solution

$$\vec{x} = e^{-t} c_1 \cdot \begin{bmatrix} 2 \\ 1 \end{bmatrix} + c_2 e^t \cdot \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

- Substitute in vector of dependent variables

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} 2e^{-t}c_1 + c_2e^t \\ e^{-t}c_1 + c_2e^t \end{bmatrix}$$

- Solution to the system of ODEs

$$\{x(t) = 2e^{-t}c_1 + c_2e^t, y(t) = e^{-t}c_1 + c_2e^t\}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 31

```
dsolve([diff(x(t),t)=-3*x(t)+4*y(t),diff(y(t),t)=-2*x(t)+3*y(t)],singsol=all)
```

$$\begin{aligned} x(t) &= c_1e^t + c_2e^{-t} \\ y(t) &= c_1e^t + \frac{c_2e^{-t}}{2} \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 67

```
DSolve[{x'[t]==-3*x[t]+4*y[t],y'[t]==-2*x[t]+3*y[t]},{x[t],y[t]},t,IncludeSingularSolutions
```

$$\begin{aligned} x(t) &\rightarrow e^{-t}(2c_2(e^{2t} - 1) - c_1(e^{2t} - 2)) \\ y(t) &\rightarrow e^{-t}(c_2(2e^{2t} - 1) - c_1(e^{2t} - 1)) \end{aligned}$$

28.2 problem 1(b)

28.2.1 Solution using Matrix exponential method	4009
28.2.2 Solution using explicit Eigenvalue and Eigenvector method . . .	4010
28.2.3 Maple step by step solution	4015

Internal problem ID [6521]

Internal file name [OUTPUT/5769_Sunday_June_05_2022_03_53_43_PM_80895195/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 10. Systems of First-Order Equations. Section 10.3 Homogeneous Linear Systems with Constant Coefficients. Page 387

Problem number: 1(b).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"system of linear ODEs"**

Solve

$$\begin{aligned}x'(t) &= 4x(t) - 2y(t) \\y'(t) &= 5x(t) + 2y(t)\end{aligned}$$

28.2.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 4 & -2 \\ 5 & 2 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

For the above matrix A , the matrix exponential can be found to be

$$\begin{aligned}e^{At} &= \begin{bmatrix} e^{3t} \cos(3t) + \frac{e^{3t} \sin(3t)}{3} & -\frac{2e^{3t} \sin(3t)}{3} \\ \frac{5e^{3t} \sin(3t)}{3} & e^{3t} \cos(3t) - \frac{e^{3t} \sin(3t)}{3} \end{bmatrix} \\ &= \begin{bmatrix} \frac{e^{3t}(3 \cos(3t) + \sin(3t))}{3} & -\frac{2e^{3t} \sin(3t)}{3} \\ \frac{5e^{3t} \sin(3t)}{3} & \frac{e^{3t}(3 \cos(3t) - \sin(3t))}{3} \end{bmatrix}\end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 \vec{x}_h(t) &= e^{At} \vec{c} \\
 &= \begin{bmatrix} \frac{e^{3t}(3 \cos(3t) + \sin(3t))}{3} & -\frac{2e^{3t} \sin(3t)}{3} \\ \frac{5e^{3t} \sin(3t)}{3} & \frac{e^{3t}(3 \cos(3t) - \sin(3t))}{3} \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} \\
 &= \begin{bmatrix} \frac{e^{3t}(3 \cos(3t) + \sin(3t))c_1}{3} - \frac{2e^{3t} \sin(3t)c_2}{3} \\ \frac{5e^{3t} \sin(3t)c_1}{3} + \frac{e^{3t}(3 \cos(3t) - \sin(3t))c_2}{3} \end{bmatrix} \\
 &= \begin{bmatrix} \frac{e^{3t}(-2c_2 + c_1) \sin(3t)}{3} + e^{3t} \cos(3t) c_1 \\ \frac{e^{3t}(5c_1 - c_2) \sin(3t)}{3} + e^{3t} \cos(3t) c_2 \end{bmatrix}
 \end{aligned}$$

Since no forcing function is given, then the final solution is $\vec{x}_h(t)$ above.

28.2.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A \vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 4 & -2 \\ 5 & 2 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det \left(\begin{bmatrix} 4 & -2 \\ 5 & 2 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) = 0$$

Therefore

$$\det \left(\begin{bmatrix} 4 - \lambda & -2 \\ 5 & 2 - \lambda \end{bmatrix} \right) = 0$$

Which gives the characteristic equation

$$\lambda^2 - 6\lambda + 18 = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = 3 + 3i$$

$$\lambda_2 = 3 - 3i$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
$3 - 3i$	1	complex eigenvalue
$3 + 3i$	1	complex eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = 3 - 3i$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 4 & -2 \\ 5 & 2 \end{bmatrix} - (3 - 3i) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 + 3i & -2 \\ 5 & -1 + 3i \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} 1 + 3i & -2 & 0 \\ 5 & -1 + 3i & 0 \end{array} \right]$$

$$R_2 = R_2 + \left(-\frac{1}{2} + \frac{3i}{2} \right) R_1 \implies \left[\begin{array}{cc|c} 1 + 3i & -2 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} 1 + 3i & -2 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = (\frac{1}{5} - \frac{3i}{5}) t\}$

Hence the solution is

$$\begin{bmatrix} (\frac{1}{5} - \frac{3i}{5}) t \\ t \end{bmatrix} = \begin{bmatrix} (\frac{1}{5} - \frac{3i}{5}) t \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} (\frac{1}{5} - \frac{3i}{5}) t \\ t \end{bmatrix} = t \begin{bmatrix} \frac{1}{5} - \frac{3i}{5} \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} (\frac{1}{5} - \frac{3i}{5}) t \\ t \end{bmatrix} = \begin{bmatrix} \frac{1}{5} - \frac{3i}{5} \\ 1 \end{bmatrix}$$

Which is normalized to

$$\begin{bmatrix} (\frac{1}{5} - \frac{3i}{5}) t \\ t \end{bmatrix} = \begin{bmatrix} 1 - 3i \\ 5 \end{bmatrix}$$

Considering the eigenvalue $\lambda_2 = 3 + 3i$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 4 & -2 \\ 5 & 2 \end{bmatrix} - (3 + 3i) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 - 3i & -2 \\ 5 & -1 - 3i \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} 1 - 3i & -2 & 0 \\ 5 & -1 - 3i & 0 \end{array} \right]$$

$$R_2 = R_2 + \left(-\frac{1}{2} - \frac{3i}{2} \right) R_1 \implies \left[\begin{array}{cc|c} 1 - 3i & -2 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} 1 - 3i & -2 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = (\frac{1}{5} + \frac{3i}{5})t\}$

Hence the solution is

$$\begin{bmatrix} (\frac{1}{5} + \frac{3i}{5})t \\ t \end{bmatrix} = \begin{bmatrix} (\frac{1}{5} + \frac{3i}{5})t \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} (\frac{1}{5} + \frac{3i}{5})t \\ t \end{bmatrix} = t \begin{bmatrix} \frac{1}{5} + \frac{3i}{5} \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} (\frac{1}{5} + \frac{3i}{5})t \\ t \end{bmatrix} = \begin{bmatrix} \frac{1}{5} + \frac{3i}{5} \\ 1 \end{bmatrix}$$

Which is normalized to

$$\begin{bmatrix} (\frac{1}{5} + \frac{3i}{5})t \\ t \end{bmatrix} = \begin{bmatrix} 1 + 3i \\ 5 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
$3 + 3i$	1	1	No	$\begin{bmatrix} \frac{1}{5} + \frac{3i}{5} \\ 1 \end{bmatrix}$
$3 - 3i$	1	1	No	$\begin{bmatrix} \frac{1}{5} - \frac{3i}{5} \\ 1 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. Therefore the final solution is

$$\vec{x}_h(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

Which is written as

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = c_1 \begin{bmatrix} \left(\frac{1}{5} + \frac{3i}{5}\right) e^{(3+3i)t} \\ e^{(3+3i)t} \end{bmatrix} + c_2 \begin{bmatrix} \left(\frac{1}{5} - \frac{3i}{5}\right) e^{(3-3i)t} \\ e^{(3-3i)t} \end{bmatrix}$$

Which becomes

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} \left(\frac{1}{5} + \frac{3i}{5}\right) c_1 e^{(3+3i)t} + \left(\frac{1}{5} - \frac{3i}{5}\right) c_2 e^{(3-3i)t} \\ c_1 e^{(3+3i)t} + c_2 e^{(3-3i)t} \end{bmatrix}$$

The following is the phase plot of the system.




Figure 493: Phase plot

28.2.3 Maple step by step solution

Let's solve

$$[x'(t) = 4x(t) - 2y(t), y'(t) = 5x(t) + 2y(t)]$$

- Define vector

$$\vec{x}(t) = \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

- Convert system into a vector equation

$$\vec{x}'(t) = \begin{bmatrix} 4 & -2 \\ 5 & 2 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

- System to solve

$$\vec{x}'(t) = \begin{bmatrix} 4 & -2 \\ 5 & 2 \end{bmatrix} \cdot \vec{x}(t)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 4 & -2 \\ 5 & 2 \end{bmatrix}$$

- Rewrite the system as

$$\vec{x}'(t) = A \cdot \vec{x}(t)$$

- To solve the system, find the eigenvalues and eigenvectors of A
- Eigenpairs of A

$$\left[\left[3 - 3I, \begin{bmatrix} \frac{1}{5} - \frac{3I}{5} \\ 1 \end{bmatrix} \right], \left[3 + 3I, \begin{bmatrix} \frac{1}{5} + \frac{3I}{5} \\ 1 \end{bmatrix} \right] \right]$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$\left[3 - 3I, \begin{bmatrix} \frac{1}{5} - \frac{3I}{5} \\ 1 \end{bmatrix} \right]$$

- Solution from eigenpair

$$e^{(3-3I)t} \cdot \begin{bmatrix} \frac{1}{5} - \frac{3I}{5} \\ 1 \end{bmatrix}$$

- Use Euler identity to write solution in terms of sin and cos

$$e^{3t} \cdot (\cos(3t) - I \sin(3t)) \cdot \begin{bmatrix} \frac{1}{5} - \frac{3I}{5} \\ 1 \end{bmatrix}$$

- Simplify expression

$$e^{3t} \cdot \begin{bmatrix} \left(\frac{1}{5} - \frac{3I}{5}\right) (\cos(3t) - I \sin(3t)) \\ \cos(3t) - I \sin(3t) \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\left[\vec{x}_1(t) = e^{3t} \cdot \begin{bmatrix} \frac{\cos(3t)}{5} - \frac{3 \sin(3t)}{5} \\ \cos(3t) \end{bmatrix}, \vec{x}_2(t) = e^{3t} \cdot \begin{bmatrix} -\frac{\sin(3t)}{5} - \frac{3 \cos(3t)}{5} \\ -\sin(3t) \end{bmatrix} \right]$$

- General solution to the system of ODEs

$$\vec{x} = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

- Substitute solutions into the general solution

$$\vec{x} = c_1 e^{3t} \cdot \begin{bmatrix} \frac{\cos(3t)}{5} - \frac{3 \sin(3t)}{5} \\ \cos(3t) \end{bmatrix} + e^{3t} c_2 \cdot \begin{bmatrix} -\frac{\sin(3t)}{5} - \frac{3 \cos(3t)}{5} \\ -\sin(3t) \end{bmatrix}$$

- Substitute in vector of dependent variables

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} \frac{((c_1 - 3c_2) \cos(3t) - 3(c_1 + \frac{c_2}{3}) \sin(3t)) e^{3t}}{5} \\ e^{3t} (c_1 \cos(3t) - c_2 \sin(3t)) \end{bmatrix}$$

- Solution to the system of ODEs

$$\left\{ x(t) = \frac{((c_1 - 3c_2) \cos(3t) - 3(c_1 + \frac{c_2}{3}) \sin(3t)) e^{3t}}{5}, y(t) = e^{3t} (c_1 \cos(3t) - c_2 \sin(3t)) \right\}$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 59

```
dsolve([diff(x(t),t)=4*x(t)-2*y(t),diff(y(t),t)=5*x(t)+2*y(t)],singsol=all)
```

$$x(t) = e^{3t} (c_1 \sin(3t) + c_2 \cos(3t))$$

$$y(t) = \frac{e^{3t} (c_1 \sin(3t) + 3c_2 \sin(3t) - 3c_1 \cos(3t) + c_2 \cos(3t))}{2}$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 70

```
DSolve[{x'[t]==4*x[t]-2*y[t],y'[t]==5*x[t]+2*y[t]},{x[t],y[t]},t,IncludeSingularSolutions ->
```

$$x(t) \rightarrow \frac{1}{3} e^{3t} (3c_1 \cos(3t) + (c_1 - 2c_2) \sin(3t))$$

$$y(t) \rightarrow \frac{1}{3} e^{3t} (3c_2 \cos(3t) + (5c_1 - c_2) \sin(3t))$$

28.3 problem 1(c)

28.3.1 Solution using Matrix exponential method	4018
28.3.2 Solution using explicit Eigenvalue and Eigenvector method . . .	4019
28.3.3 Maple step by step solution	4024

Internal problem ID [6522]

Internal file name [OUTPUT/5770_Sunday_June_05_2022_03_53_46_PM_49252342/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 10. Systems of First-Order Equations. Section 10.3 Homogeneous Linear Systems with Constant Coefficients. Page 387

Problem number: 1(c).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**system of linear ODEs**"

Solve

$$x'(t) = 5x(t) + 4y(t)$$

$$y'(t) = -x(t) + y(t)$$

28.3.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 5 & 4 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} (2t + 1)e^{3t} & 4te^{3t} \\ -te^{3t} & e^{3t}(1 - 2t) \end{bmatrix}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 \vec{x}_h(t) &= e^{At} \vec{c} \\
 &= \begin{bmatrix} (2t+1)e^{3t} & 4te^{3t} \\ -te^{3t} & e^{3t}(1-2t) \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} \\
 &= \begin{bmatrix} (2t+1)e^{3t}c_1 + 4te^{3t}c_2 \\ -te^{3t}c_1 + e^{3t}(1-2t)c_2 \end{bmatrix} \\
 &= \begin{bmatrix} e^{3t}(2tc_1 + 4c_2t + c_1) \\ e^{3t}(-tc_1 - 2c_2t + c_2) \end{bmatrix}
 \end{aligned}$$

Since no forcing function is given, then the final solution is $\vec{x}_h(t)$ above.

28.3.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 5 & 4 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det\left(\begin{bmatrix} 5 & 4 \\ -1 & 1 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}\right) = 0$$

Therefore

$$\det\left(\begin{bmatrix} 5 - \lambda & 4 \\ -1 & 1 - \lambda \end{bmatrix}\right) = 0$$

Which gives the characteristic equation

$$\lambda^2 - 6\lambda + 9 = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = 3$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
3	1	real eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = 3$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 5 & 4 \\ -1 & 1 \end{bmatrix} - (3) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$
$$\begin{bmatrix} 2 & 4 \\ -1 & -2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} 2 & 4 & 0 \\ -1 & -2 & 0 \end{array} \right]$$

$$R_2 = R_2 + \frac{R_1}{2} \implies \left[\begin{array}{cc|c} 2 & 4 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} 2 & 4 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = -2t\}$

Hence the solution is

$$\begin{bmatrix} -2t \\ t \end{bmatrix} = \begin{bmatrix} -2t \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} -2t \\ t \end{bmatrix} = t \begin{bmatrix} -2 \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} -2t \\ t \end{bmatrix} = \begin{bmatrix} -2 \\ 1 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
3	2	1	Yes	$\begin{bmatrix} -2 \\ 1 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. eigenvalue 3 is real and repeated eigenvalue of multiplicity 2. There are two possible cases that can happen. This is illustrated in this diagram




Figure 494: Possible case for repeated λ of multiplicity 2

This eigenvalue has algebraic multiplicity of 2, and geometric multiplicity 1, therefore this is defective eigenvalue. The defect is 1. This falls into case 2 shown above. We need to generate the missing additional generalized eigenvector \vec{v}_2 by solving

$$(A - \lambda I) \vec{v}_2 = \vec{v}_1$$

Where \vec{v}_1 is the normal (rank 1) eigenvector found above. Hence we need to solve

$$\left(\begin{bmatrix} 5 & 4 \\ -1 & 1 \end{bmatrix} - (3) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} -2 \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 4 \\ -1 & -2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} -2 \\ 1 \end{bmatrix}$$

Solving for \vec{v}_2 gives

$$\vec{v}_2 = \begin{bmatrix} -3 \\ 1 \end{bmatrix}$$

We have found two generalized eigenvectors for eigenvalue 3. Therefore the two basis solution associated with this eigenvalue are

$$\begin{aligned}\vec{x}_1(t) &= \vec{v}_1 e^{\lambda t} \\ &= \begin{bmatrix} -2 \\ 1 \end{bmatrix} e^{3t} \\ &= \begin{bmatrix} -2 e^{3t} \\ e^{3t} \end{bmatrix}\end{aligned}$$

And

$$\begin{aligned}\vec{x}_2(t) &= (\vec{v}_1 t + \vec{v}_2) e^{\lambda t} \\ &= \left(\begin{bmatrix} -2 \\ 1 \end{bmatrix} t + \begin{bmatrix} -3 \\ 1 \end{bmatrix} \right) e^{3t} \\ &= \begin{bmatrix} e^{3t}(-2t - 3) \\ e^{3t}(t + 1) \end{bmatrix}\end{aligned}$$

Therefore the final solution is

$$\vec{x}_h(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

Which is written as

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = c_1 \begin{bmatrix} -2 e^{3t} \\ e^{3t} \end{bmatrix} + c_2 \begin{bmatrix} e^{3t}(-2t - 3) \\ e^{3t}(t + 1) \end{bmatrix}$$

Which becomes

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} e^{3t}(-2c_2 t - 2c_1 - 3c_2) \\ e^{3t}(c_2 t + c_1 + c_2) \end{bmatrix}$$

The following is the phase plot of the system.




Figure 495: Phase plot

28.3.3 Maple step by step solution

Let's solve

$$[x'(t) = 5x(t) + 4y(t), y'(t) = -x(t) + y(t)]$$

- Define vector

$$\vec{x}(t) = \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

- Convert system into a vector equation

$$\vec{x}'(t) = \begin{bmatrix} 5 & 4 \\ -1 & 1 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

- System to solve

$$\vec{x}'(t) = \begin{bmatrix} 5 & 4 \\ -1 & 1 \end{bmatrix} \cdot \vec{x}(t)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 5 & 4 \\ -1 & 1 \end{bmatrix}$$

- Rewrite the system as

$$\vec{x}'(t) = A \cdot \vec{x}(t)$$

- To solve the system, find the eigenvalues and eigenvectors of A
- Eigenpairs of A

$$\left[\left[3, \begin{bmatrix} -2 \\ 1 \end{bmatrix} \right], \left[3, \begin{bmatrix} 0 \\ 0 \end{bmatrix} \right] \right]$$

- Consider eigenpair, with eigenvalue of algebraic multiplicity 2

$$\left[3, \begin{bmatrix} -2 \\ 1 \end{bmatrix} \right]$$

- First solution from eigenvalue 3

$$\vec{x}_1(t) = e^{3t} \cdot \begin{bmatrix} -2 \\ 1 \end{bmatrix}$$

- Form of the 2nd homogeneous solution where \vec{p} is to be solved for, $\lambda = 3$ is the eigenvalue, and

$$\vec{x}_2(t) = e^{\lambda t} (t\vec{v} + \vec{p})$$

- Note that the t multiplying \vec{v} makes this solution linearly independent to the 1st solution obtained
- Substitute $\vec{x}_2(t)$ into the homogeneous system

$$\lambda e^{\lambda t} (t\vec{v} + \vec{p}) + e^{\lambda t} \vec{v} = (e^{\lambda t} A) \cdot (t\vec{v} + \vec{p})$$

- Use the fact that \vec{v} is an eigenvector of A

$$\lambda e^{\lambda t} (t\vec{v} + \vec{p}) + e^{\lambda t} \vec{v} = e^{\lambda t} (\lambda t\vec{v} + A \cdot \vec{p})$$

- Simplify equation

$$\lambda \vec{p} + \vec{v} = A \cdot \vec{p}$$

- Make use of the identity matrix I

$$(\lambda \cdot I) \cdot \vec{p} + \vec{v} = A \cdot \vec{p}$$

- Condition \vec{p} must meet for $\vec{x}_2(t)$ to be a solution to the homogeneous system

$$(A - \lambda \cdot I) \cdot \vec{p} = \vec{v}$$

- Choose \vec{p} to use in the second solution to the homogeneous system from eigenvalue 3

$$\left(\begin{bmatrix} 5 & 4 \\ -1 & 1 \end{bmatrix} - 3 \cdot \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \cdot \vec{p} = \begin{bmatrix} -2 \\ 1 \end{bmatrix}$$

- Choice of \vec{p}

$$\vec{p} = \begin{bmatrix} -1 \\ 0 \end{bmatrix}$$

- Second solution from eigenvalue 3

$$\vec{x}_2(t) = e^{3t} \cdot \left(t \cdot \begin{bmatrix} -2 \\ 1 \end{bmatrix} + \begin{bmatrix} -1 \\ 0 \end{bmatrix} \right)$$

- General solution to the system of ODEs

$$\vec{x} = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

- Substitute solutions into the general solution

$$\vec{x} = c_1 e^{3t} \cdot \begin{bmatrix} -2 \\ 1 \end{bmatrix} + e^{3t} c_2 \cdot \left(t \cdot \begin{bmatrix} -2 \\ 1 \end{bmatrix} + \begin{bmatrix} -1 \\ 0 \end{bmatrix} \right)$$

- Substitute in vector of dependent variables

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} e^{3t}(-2tc_2 - 2c_1 - c_2) \\ e^{3t}(tc_2 + c_1) \end{bmatrix}$$

- Solution to the system of ODEs

$$\{x(t) = e^{3t}(-2tc_2 - 2c_1 - c_2), y(t) = e^{3t}(tc_2 + c_1)\}$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 35

```
dsolve([diff(x(t),t)=5*x(t)+4*y(t),diff(y(t),t)=-x(t)+y(t)],singsol=all)
```

$$x(t) = e^{3t}(c_2 t + c_1)$$

$$y(t) = -\frac{e^{3t}(2c_2 t + 2c_1 - c_2)}{4}$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 46

```
DSolve[{x'[t]==5*x[t]+4*y[t],y'[t]==-x[t]+y[t]},{x[t],y[t]},t,IncludeSingularSolutions -> Tr
```

$$x(t) \rightarrow e^{3t}(2c_1t + 4c_2t + c_1)$$

$$y(t) \rightarrow e^{3t}(c_2 - (c_1 + 2c_2)t)$$

28.4 problem 1(d)

28.4.1 Solution using Matrix exponential method	4028
28.4.2 Solution using explicit Eigenvalue and Eigenvector method . . .	4029
28.4.3 Maple step by step solution	4034

Internal problem ID [6523]

Internal file name [OUTPUT/5771_Sunday_June_05_2022_03_53_48_PM_31022794/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 10. Systems of First-Order Equations. Section 10.3 Homogeneous Linear Systems with Constant Coefficients. Page 387

Problem number: 1(d).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**system of linear ODEs**"

Solve

$$x'(t) = 4x(t) - 3y(t)$$

$$y'(t) = 8x(t) - 6y(t)$$

28.4.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 4 & -3 \\ 8 & -6 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} -2e^{-2t} + 3 & -\frac{3}{2} + \frac{3e^{-2t}}{2} \\ 4 - 4e^{-2t} & 3e^{-2t} - 2 \end{bmatrix}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 \vec{x}_h(t) &= e^{At} \vec{c} \\
 &= \begin{bmatrix} -2e^{-2t} + 3 & -\frac{3}{2} + \frac{3e^{-2t}}{2} \\ 4 - 4e^{-2t} & 3e^{-2t} - 2 \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} \\
 &= \begin{bmatrix} (-2e^{-2t} + 3)c_1 + \left(-\frac{3}{2} + \frac{3e^{-2t}}{2}\right)c_2 \\ (4 - 4e^{-2t})c_1 + (3e^{-2t} - 2)c_2 \end{bmatrix} \\
 &= \begin{bmatrix} \frac{(-4c_1 + 3c_2)e^{-2t}}{2} + 3c_1 - \frac{3c_2}{2} \\ (-4c_1 + 3c_2)e^{-2t} + 4c_1 - 2c_2 \end{bmatrix}
 \end{aligned}$$

Since no forcing function is given, then the final solution is $\vec{x}_h(t)$ above.

28.4.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A \vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 4 & -3 \\ 8 & -6 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det \left(\begin{bmatrix} 4 & -3 \\ 8 & -6 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) = 0$$

Therefore

$$\det \left(\begin{bmatrix} 4 - \lambda & -3 \\ 8 & -6 - \lambda \end{bmatrix} \right) = 0$$

Which gives the characteristic equation

$$\lambda^2 + 2\lambda = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = -2$$

$$\lambda_2 = 0$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
0	1	real eigenvalue
-2	1	real eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = -2$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 4 & -3 \\ 8 & -6 \end{bmatrix} - (-2) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 6 & -3 \\ 8 & -4 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} 6 & -3 & 0 \\ 8 & -4 & 0 \end{array} \right]$$

$$R_2 = R_2 - \frac{4R_1}{3} \implies \left[\begin{array}{cc|c} 6 & -3 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} 6 & -3 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = \frac{t}{2}\}$

Hence the solution is

$$\begin{bmatrix} \frac{t}{2} \\ t \end{bmatrix} = \begin{bmatrix} \frac{t}{2} \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} \frac{t}{2} \\ t \end{bmatrix} = t \begin{bmatrix} \frac{1}{2} \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} \frac{1}{2} \\ 1 \end{bmatrix} = \begin{bmatrix} \frac{1}{2} \\ 1 \end{bmatrix}$$

Which is normalized to

$$\begin{bmatrix} \frac{1}{2} \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

Considering the eigenvalue $\lambda_2 = 0$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 4 & -3 \\ 8 & -6 \end{bmatrix} - (0) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 4 & -3 \\ 8 & -6 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} 4 & -3 & 0 \\ 8 & -6 & 0 \end{array} \right]$$

$$R_2 = R_2 - 2R_1 \implies \left[\begin{array}{cc|c} 4 & -3 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} 4 & -3 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = \frac{3t}{4}\}$

Hence the solution is

$$\begin{bmatrix} \frac{3t}{4} \\ t \end{bmatrix} = \begin{bmatrix} \frac{3t}{4} \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} \frac{3t}{4} \\ t \end{bmatrix} = t \begin{bmatrix} \frac{3}{4} \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} \frac{3t}{4} \\ t \end{bmatrix} = \begin{bmatrix} \frac{3}{4} \\ 1 \end{bmatrix}$$

Which is normalized to

$$\begin{bmatrix} \frac{3t}{4} \\ t \end{bmatrix} = \begin{bmatrix} 3 \\ 4 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
-2	1	1	No	$\begin{bmatrix} \frac{1}{2} \\ 1 \end{bmatrix}$
0	1	1	No	$\begin{bmatrix} \frac{3}{4} \\ 1 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. Since eigenvalue -2 is real and distinct then the corresponding eigenvector solution is

$$\begin{aligned}\vec{x}_1(t) &= \vec{v}_1 e^{-2t} \\ &= \begin{bmatrix} \frac{1}{2} \\ 1 \end{bmatrix} e^{-2t}\end{aligned}$$

Since eigenvalue 0 is real and distinct then the corresponding eigenvector solution is

$$\begin{aligned}\vec{x}_2(t) &= \vec{v}_2 e^0 \\ &= \begin{bmatrix} \frac{3}{4} \\ 1 \end{bmatrix} e^0\end{aligned}$$

Therefore the final solution is

$$\vec{x}_h(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

Which is written as

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = c_1 \begin{bmatrix} \frac{e^{-2t}}{2} \\ e^{-2t} \end{bmatrix} + c_2 \begin{bmatrix} \frac{3}{4} \\ 1 \end{bmatrix}$$

Which becomes

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} \frac{c_1 e^{-2t}}{2} + \frac{3c_2}{4} \\ c_1 e^{-2t} + c_2 \end{bmatrix}$$

The following is the phase plot of the system.




Figure 496: Phase plot

28.4.3 Maple step by step solution

Let's solve

$$[x'(t) = 4x(t) - 3y(t), y'(t) = 8x(t) - 6y(t)]$$

- Define vector

$$\vec{x}(t) = \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

- Convert system into a vector equation

$$\vec{x}'(t) = \begin{bmatrix} 4 & -3 \\ 8 & -6 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

- System to solve

$$\vec{x}'(t) = \begin{bmatrix} 4 & -3 \\ 8 & -6 \end{bmatrix} \cdot \vec{x}(t)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 4 & -3 \\ 8 & -6 \end{bmatrix}$$

- Rewrite the system as

$$\vec{x}'(t) = A \cdot \vec{x}(t)$$

- To solve the system, find the eigenvalues and eigenvectors of A
- Eigenpairs of A

$$\left[\left[-2, \begin{bmatrix} \frac{1}{2} \\ 1 \end{bmatrix} \right], \left[0, \begin{bmatrix} \frac{3}{4} \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[-2, \begin{bmatrix} \frac{1}{2} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{x}_1 = e^{-2t} \cdot \begin{bmatrix} \frac{1}{2} \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[0, \begin{bmatrix} \frac{3}{4} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{x}_2 = \begin{bmatrix} \frac{3}{4} \\ 1 \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{x} = c_1 \vec{x}_1 + c_2 \vec{x}_2$$

- Substitute solutions into the general solution

$$\vec{x} = c_1 e^{-2t} \cdot \begin{bmatrix} \frac{1}{2} \\ 1 \end{bmatrix} + \begin{bmatrix} \frac{3c_2}{4} \\ c_2 \end{bmatrix}$$

- Substitute in vector of dependent variables

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} \frac{c_1 e^{-2t}}{2} + \frac{3c_2}{4} \\ c_1 e^{-2t} + c_2 \end{bmatrix}$$

- Solution to the system of ODEs

$$\left\{ x(t) = \frac{c_1 e^{-2t}}{2} + \frac{3c_2}{4}, y(t) = c_1 e^{-2t} + c_2 \right\}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve([diff(x(t),t)=4*x(t)-3*y(t),diff(y(t),t)=8*x(t)-6*y(t)],singsol=all)
```

$$\begin{aligned} x(t) &= c_1 + c_2 e^{-2t} \\ y(t) &= 2c_2 e^{-2t} + \frac{4c_1}{3} \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 59

```
DSolve[{x'[t]==4*x[t]-3*y[t],y'[t]==8*x[t]-6*y[t]},{x[t],y[t]},t,IncludeSingularSolutions ->
```

$$\begin{aligned} x(t) &\rightarrow c_1(3 - 2e^{-2t}) + \frac{3}{2}c_2(e^{-2t} - 1) \\ y(t) &\rightarrow c_1(4 - 4e^{-2t}) + c_2(3e^{-2t} - 2) \end{aligned}$$

28.5 problem 1(e)

28.5.1 Solution using Matrix exponential method 4037

28.5.2 Solution using explicit Eigenvalue and Eigenvector method . . . 4038

28.5.3 Maple step by step solution 4043

Internal problem ID [6524]

Internal file name [OUTPUT/5772_Sunday_June_05_2022_03_53_49_PM_37503988/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 10. Systems of First-Order Equations. Section 10.3 Homogeneous Linear Systems with Constant Coefficients. Page 387

Problem number: 1(e).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**system of linear ODEs**"

Solve

$$x'(t) = 2x(t)$$

$$y'(t) = 3y(t)$$

28.5.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} e^{2t} & 0 \\ 0 & e^{3t} \end{bmatrix}$$

Therefore the homogeneous solution is

$$\begin{aligned}\vec{x}_h(t) &= e^{At}\vec{c} \\ &= \begin{bmatrix} e^{2t} & 0 \\ 0 & e^{3t} \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} \\ &= \begin{bmatrix} e^{2t}c_1 \\ e^{3t}c_2 \end{bmatrix}\end{aligned}$$

Since no forcing function is given, then the final solution is $\vec{x}_h(t)$ above.

28.5.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det\left(\begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}\right) = 0$$

Therefore

$$\det\left(\begin{bmatrix} 2 - \lambda & 0 \\ 0 & 3 - \lambda \end{bmatrix}\right) = 0$$

Since the matrix A is triangular matrix, then the determinant is the product of the elements along the diagonal. Therefore the above becomes

$$(2 - \lambda)(3 - \lambda) = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = 2$$

$$\lambda_2 = 3$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
2	1	real eigenvalue
3	1	real eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = 2$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix} - (2) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} 0 & 0 & 0 \\ 0 & 1 & 0 \end{array} \right]$$

Since the current pivot $A(1,2)$ is zero, then the current pivot row is replaced with a row with a non-zero pivot. Swapping row 1 and row 2 gives

$$\left[\begin{array}{cc|c} 0 & 1 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_1\}$ and the leading variables are $\{v_2\}$. Let $v_1 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_2 = 0\}$

Hence the solution is

$$\begin{bmatrix} t \\ 0 \end{bmatrix} = \begin{bmatrix} t \\ 0 \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} t \\ 0 \end{bmatrix} = t \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} t \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

Considering the eigenvalue $\lambda_2 = 3$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix} - (3) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$
$$\begin{bmatrix} -1 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} -1 & 0 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} -1 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = 0\}$

Hence the solution is

$$\begin{bmatrix} 0 \\ t \end{bmatrix} = \begin{bmatrix} 0 \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} 0 \\ t \end{bmatrix} = t \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
2	1	1	No	$\begin{bmatrix} 1 \\ 0 \end{bmatrix}$
3	1	1	No	$\begin{bmatrix} 0 \\ 1 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. Since eigenvalue 2 is real and distinct then the corresponding eigenvector solution is

$$\begin{aligned} \vec{x}_1(t) &= \vec{v}_1 e^{2t} \\ &= \begin{bmatrix} 1 \\ 0 \end{bmatrix} e^{2t} \end{aligned}$$

Since eigenvalue 3 is real and distinct then the corresponding eigenvector solution is

$$\begin{aligned}\vec{x}_2(t) &= \vec{v}_2 e^{3t} \\ &= \begin{bmatrix} 0 \\ 1 \end{bmatrix} e^{3t}\end{aligned}$$

Therefore the final solution is

$$\vec{x}_h(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

Which is written as

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = c_1 \begin{bmatrix} e^{2t} \\ 0 \end{bmatrix} + c_2 \begin{bmatrix} 0 \\ e^{3t} \end{bmatrix}$$

Which becomes

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} c_1 e^{2t} \\ c_2 e^{3t} \end{bmatrix}$$

The following is the phase plot of the system.




Figure 497: Phase plot

28.5.3 Maple step by step solution

Let's solve

$$[x'(t) = 2x(t), y'(t) = 3y(t)]$$

- Define vector

$$\vec{x}(t) = \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

- Convert system into a vector equation

$$\vec{x}'(t) = \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

- System to solve

$$\vec{x}'(t) = \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix} \cdot \vec{x}(t)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix}$$

- Rewrite the system as

$$\vec{x}'(t) = A \cdot \vec{x}(t)$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[2, \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right], \left[3, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[2, \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{x}_1 = e^{2t} \cdot \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

- Consider eigenpair

$$\left[3, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{x}_2 = e^{3t} \cdot \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{x} = c_1 \vec{x}_1 + c_2 \vec{x}_2$$

- Substitute solutions into the general solution

$$\vec{x} = c_1 e^{2t} \cdot \begin{bmatrix} 1 \\ 0 \end{bmatrix} + e^{3t} c_2 \cdot \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

- Substitute in vector of dependent variables

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} c_1 e^{2t} \\ e^{3t} c_2 \end{bmatrix}$$

- Solution to the system of ODEs

$$\{x(t) = c_1 e^{2t}, y(t) = e^{3t} c_2\}$$

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve([diff(x(t),t)=2*x(t),diff(y(t),t)=3*y(t)],singsol=all)
```

$$\begin{aligned} x(t) &= c_2 e^{2t} \\ y(t) &= c_1 e^{3t} \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.044 (sec). Leaf size: 65

```
DSolve[{x'[t]==2*x[t],y'[t]==3*y[t]},{x[t],y[t]},t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow c_1 e^{2t}$$

$$y(t) \rightarrow c_2 e^{3t}$$

$$x(t) \rightarrow c_1 e^{2t}$$

$$y(t) \rightarrow 0$$

$$x(t) \rightarrow 0$$

$$y(t) \rightarrow c_2 e^{3t}$$

$$x(t) \rightarrow 0$$

$$y(t) \rightarrow 0$$

28.6 problem 1(f)

28.6.1 Solution using Matrix exponential method 4046

28.6.2 Solution using explicit Eigenvalue and Eigenvector method . . . 4047

28.6.3 Maple step by step solution 4052

Internal problem ID [6525]

Internal file name [OUTPUT/5773_Sunday_June_05_2022_03_53_51_PM_12984942/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 10. Systems of First-Order Equations. Section 10.3 Homogeneous Linear Systems with Constant Coefficients. Page 387

Problem number: 1(f).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**system of linear ODEs**"

Solve

$$x'(t) = -4x(t) - y(t)$$

$$y'(t) = x(t) - 2y(t)$$

28.6.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} -4 & -1 \\ 1 & -2 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} e^{-3t}(-t+1) & -te^{-3t} \\ te^{-3t} & e^{-3t}(t+1) \end{bmatrix}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 \vec{x}_h(t) &= e^{At} \vec{c} \\
 &= \begin{bmatrix} e^{-3t}(-t+1) & -te^{-3t} \\ te^{-3t} & e^{-3t}(t+1) \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} \\
 &= \begin{bmatrix} e^{-3t}(-t+1)c_1 - te^{-3t}c_2 \\ te^{-3t}c_1 + e^{-3t}(t+1)c_2 \end{bmatrix} \\
 &= \begin{bmatrix} -e^{-3t}(c_1(t-1) + c_2t) \\ e^{-3t}(tc_1 + c_2t + c_2) \end{bmatrix}
 \end{aligned}$$

Since no forcing function is given, then the final solution is $\vec{x}_h(t)$ above.

28.6.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} -4 & -1 \\ 1 & -2 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det \left(\begin{bmatrix} -4 & -1 \\ 1 & -2 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) = 0$$

Therefore

$$\det \left(\begin{bmatrix} -4 - \lambda & -1 \\ 1 & -2 - \lambda \end{bmatrix} \right) = 0$$

Which gives the characteristic equation

$$\lambda^2 + 6\lambda + 9 = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = -3$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
-3	1	real eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = -3$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} -4 & -1 \\ 1 & -2 \end{bmatrix} - (-3) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} -1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} -1 & -1 & 0 \\ 1 & 1 & 0 \end{array} \right]$$

$$R_2 = R_2 + R_1 \implies \left[\begin{array}{cc|c} -1 & -1 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} -1 & -1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = -t\}$

Hence the solution is

$$\begin{bmatrix} -t \\ t \end{bmatrix} = \begin{bmatrix} -t \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} -t \\ t \end{bmatrix} = t \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
-3	2	1	Yes	$\begin{bmatrix} -1 \\ 1 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. eigenvalue -3 is real and repeated eigenvalue of multiplicity 2. There are two possible cases that can happen. This is illustrated in this diagram




Figure 498: Possible case for repeated λ of multiplicity 2

This eigenvalue has algebraic multiplicity of 2, and geometric multiplicity 1, therefore this is defective eigenvalue. The defect is 1. This falls into case 2 shown above. We need to generate the missing additional generalized eigenvector \vec{v}_2 by solving

$$(A - \lambda I) \vec{v}_2 = \vec{v}_1$$

Where \vec{v}_1 is the normal (rank 1) eigenvector found above. Hence we need to solve

$$\left(\begin{bmatrix} -4 & -1 \\ 1 & -2 \end{bmatrix} - (-3) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} -1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

Solving for \vec{v}_2 gives

$$\vec{v}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

We have found two generalized eigenvectors for eigenvalue -3 . Therefore the two basis solution associated with this eigenvalue are

$$\begin{aligned}\vec{x}_1(t) &= \vec{v}_1 e^{\lambda t} \\ &= \begin{bmatrix} -1 \\ 1 \end{bmatrix} e^{-3t} \\ &= \begin{bmatrix} -e^{-3t} \\ e^{-3t} \end{bmatrix}\end{aligned}$$

And

$$\begin{aligned}\vec{x}_2(t) &= (\vec{v}_1 t + \vec{v}_2) e^{\lambda t} \\ &= \left(\begin{bmatrix} -1 \\ 1 \end{bmatrix} t + \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right) e^{-3t} \\ &= \begin{bmatrix} -t e^{-3t} \\ e^{-3t}(t+1) \end{bmatrix}\end{aligned}$$

Therefore the final solution is

$$\vec{x}_h(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

Which is written as

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = c_1 \begin{bmatrix} -e^{-3t} \\ e^{-3t} \end{bmatrix} + c_2 \begin{bmatrix} -t e^{-3t} \\ e^{-3t}(t+1) \end{bmatrix}$$

Which becomes

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} e^{-3t}(-tc_2 - c_1) \\ e^{-3t}(tc_2 + c_1 + c_2) \end{bmatrix}$$

The following is the phase plot of the system.




Figure 499: Phase plot

28.6.3 Maple step by step solution

Let's solve

$$[x'(t) = -4x(t) - y(t), y'(t) = x(t) - 2y(t)]$$

- Define vector

$$\vec{x}(t) = \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

- Convert system into a vector equation

$$\vec{x}'(t) = \begin{bmatrix} -4 & -1 \\ 1 & -2 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

- System to solve

$$\vec{x}'(t) = \begin{bmatrix} -4 & -1 \\ 1 & -2 \end{bmatrix} \cdot \vec{x}(t)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} -4 & -1 \\ 1 & -2 \end{bmatrix}$$

- Rewrite the system as

$$\vec{x}'(t) = A \cdot \vec{x}(t)$$

- To solve the system, find the eigenvalues and eigenvectors of A
- Eigenpairs of A

$$\left[\left[-3, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right], \left[-3, \begin{bmatrix} 0 \\ 0 \end{bmatrix} \right] \right]$$

- Consider eigenpair, with eigenvalue of algebraic multiplicity 2

$$\left[-3, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right]$$

- First solution from eigenvalue -3

$$\vec{x}_1(t) = e^{-3t} \cdot \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

- Form of the 2nd homogeneous solution where \vec{p} is to be solved for, $\lambda = -3$ is the eigenvalue, and

$$\vec{x}_2(t) = e^{\lambda t} (t\vec{v} + \vec{p})$$

- Note that the t multiplying \vec{v} makes this solution linearly independent to the 1st solution obtained

- Substitute $\vec{x}_2(t)$ into the homogeneous system

$$\lambda e^{\lambda t} (t\vec{v} + \vec{p}) + e^{\lambda t} \vec{v} = (e^{\lambda t} A) \cdot (t\vec{v} + \vec{p})$$

- Use the fact that \vec{v} is an eigenvector of A

$$\lambda e^{\lambda t} (t\vec{v} + \vec{p}) + e^{\lambda t} \vec{v} = e^{\lambda t} (\lambda t\vec{v} + A \cdot \vec{p})$$

- Simplify equation

$$\lambda \vec{p} + \vec{v} = A \cdot \vec{p}$$

- Make use of the identity matrix I

$$(\lambda \cdot I) \cdot \vec{p} + \vec{v} = A \cdot \vec{p}$$

- Condition \vec{p} must meet for $\vec{x}_2(t)$ to be a solution to the homogeneous system

$$(A - \lambda \cdot I) \cdot \vec{p} = \vec{v}$$

- Choose \vec{p} to use in the second solution to the homogeneous system from eigenvalue -3

$$\left(\begin{bmatrix} -4 & -1 \\ 1 & -2 \end{bmatrix} - (-3) \cdot \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \cdot \vec{p} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

- Choice of \vec{p}

$$\vec{p} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

- Second solution from eigenvalue -3

$$\vec{x}_2(t) = e^{-3t} \cdot \left(t \cdot \begin{bmatrix} -1 \\ 1 \end{bmatrix} + \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right)$$

- General solution to the system of ODEs

$$\vec{x} = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

- Substitute solutions into the general solution

$$\vec{x} = c_1 e^{-3t} \cdot \begin{bmatrix} -1 \\ 1 \end{bmatrix} + c_2 e^{-3t} \cdot \left(t \cdot \begin{bmatrix} -1 \\ 1 \end{bmatrix} + \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right)$$

- Substitute in vector of dependent variables

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} -e^{-3t}((t-1)c_2 + c_1) \\ e^{-3t}(tc_2 + c_1) \end{bmatrix}$$

- Solution to the system of ODEs

$$\{x(t) = -e^{-3t}((t-1)c_2 + c_1), y(t) = e^{-3t}(tc_2 + c_1)\}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 30

```
dsolve([diff(x(t),t)=-4*x(t)-y(t),diff(y(t),t)=x(t)-2*y(t)],singsol=all)
```

$$\begin{aligned} x(t) &= e^{-3t}(c_2 t + c_1) \\ y(t) &= -e^{-3t}(c_2 t + c_1 + c_2) \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 43

```
DSolve[{x'[t]==-4*x[t]-y[t],y'[t]==x[t]-2*y[t]},{x[t],y[t]},t,IncludeSingularSolutions -> Tr
```

$$x(t) \rightarrow e^{-3t}(c_1(-t) - c_2t + c_1)$$

$$y(t) \rightarrow e^{-3t}((c_1 + c_2)t + c_2)$$

28.7 problem 1(g)

- 28.7.1 Solution using Matrix exponential method 4056
- 28.7.2 Solution using explicit Eigenvalue and Eigenvector method . . . 4057
- 28.7.3 Maple step by step solution 4062

Internal problem ID [6526]

Internal file name [OUTPUT/5774_Sunday_June_05_2022_03_53_52_PM_6362923/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 10. Systems of First-Order Equations. Section 10.3 Homogeneous Linear Systems with Constant Coefficients. Page 387

Problem number: 1(g).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**system of linear ODEs**"

Solve

$$x'(t) = 7x(t) + 6y(t)$$

$$y'(t) = 2x(t) + 6y(t)$$

28.7.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 7 & 6 \\ 2 & 6 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} \frac{3e^{3t}}{7} + \frac{4e^{10t}}{7} & \frac{6e^{10t}}{7} - \frac{6e^{3t}}{7} \\ \frac{2e^{10t}}{7} - \frac{2e^{3t}}{7} & \frac{4e^{3t}}{7} + \frac{3e^{10t}}{7} \end{bmatrix}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 \vec{x}_h(t) &= e^{At} \vec{c} \\
 &= \begin{bmatrix} \frac{3e^{3t}}{7} + \frac{4e^{10t}}{7} & \frac{6e^{10t}}{7} - \frac{6e^{3t}}{7} \\ \frac{2e^{10t}}{7} - \frac{2e^{3t}}{7} & \frac{4e^{3t}}{7} + \frac{3e^{10t}}{7} \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} \\
 &= \begin{bmatrix} \left(\frac{3e^{3t}}{7} + \frac{4e^{10t}}{7}\right) c_1 + \left(\frac{6e^{10t}}{7} - \frac{6e^{3t}}{7}\right) c_2 \\ \left(\frac{2e^{10t}}{7} - \frac{2e^{3t}}{7}\right) c_1 + \left(\frac{4e^{3t}}{7} + \frac{3e^{10t}}{7}\right) c_2 \end{bmatrix} \\
 &= \begin{bmatrix} \frac{(4c_1+6c_2)e^{10t}}{7} + \frac{3e^{3t}(-2c_2+c_1)}{7} \\ \frac{(2c_1+3c_2)e^{10t}}{7} - \frac{2e^{3t}(-2c_2+c_1)}{7} \end{bmatrix}
 \end{aligned}$$

Since no forcing function is given, then the final solution is $\vec{x}_h(t)$ above.

28.7.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A \vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 7 & 6 \\ 2 & 6 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det \left(\begin{bmatrix} 7 & 6 \\ 2 & 6 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) = 0$$

Therefore

$$\det \left(\begin{bmatrix} 7 - \lambda & 6 \\ 2 & 6 - \lambda \end{bmatrix} \right) = 0$$

Which gives the characteristic equation

$$\lambda^2 - 13\lambda + 30 = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = 3$$

$$\lambda_2 = 10$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
3	1	real eigenvalue
10	1	real eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = 3$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 7 & 6 \\ 2 & 6 \end{bmatrix} - (3) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$
$$\begin{bmatrix} 4 & 6 \\ 2 & 3 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} 4 & 6 & 0 \\ 2 & 3 & 0 \end{array} \right]$$

$$R_2 = R_2 - \frac{R_1}{2} \implies \left[\begin{array}{cc|c} 4 & 6 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} 4 & 6 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = -\frac{3t}{2}\}$

Hence the solution is

$$\begin{bmatrix} -\frac{3t}{2} \\ t \end{bmatrix} = \begin{bmatrix} -\frac{3t}{2} \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} -\frac{3t}{2} \\ t \end{bmatrix} = t \begin{bmatrix} -\frac{3}{2} \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} -\frac{3t}{2} \\ t \end{bmatrix} = \begin{bmatrix} -\frac{3}{2} \\ 1 \end{bmatrix}$$

Which is normalized to

$$\begin{bmatrix} -\frac{3t}{2} \\ t \end{bmatrix} = \begin{bmatrix} -3 \\ 2 \end{bmatrix}$$

Considering the eigenvalue $\lambda_2 = 10$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 7 & 6 \\ 2 & 6 \end{bmatrix} - (10) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} -3 & 6 \\ 2 & -4 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} -3 & 6 & 0 \\ 2 & -4 & 0 \end{array} \right]$$

$$R_2 = R_2 + \frac{2R_1}{3} \implies \left[\begin{array}{cc|c} -3 & 6 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} -3 & 6 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = 2t\}$

Hence the solution is

$$\begin{bmatrix} 2t \\ t \end{bmatrix} = \begin{bmatrix} 2t \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} 2t \\ t \end{bmatrix} = t \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} 2t \\ t \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
3	1	1	No	$\begin{bmatrix} -\frac{3}{2} \\ 1 \end{bmatrix}$
10	1	1	No	$\begin{bmatrix} 2 \\ 1 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. Since eigenvalue 3 is real and distinct then the corresponding eigenvector solution is

$$\begin{aligned}\vec{x}_1(t) &= \vec{v}_1 e^{3t} \\ &= \begin{bmatrix} -\frac{3}{2} \\ 1 \end{bmatrix} e^{3t}\end{aligned}$$

Since eigenvalue 10 is real and distinct then the corresponding eigenvector solution is

$$\begin{aligned}\vec{x}_2(t) &= \vec{v}_2 e^{10t} \\ &= \begin{bmatrix} 2 \\ 1 \end{bmatrix} e^{10t}\end{aligned}$$

Therefore the final solution is

$$\vec{x}_h(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

Which is written as

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = c_1 \begin{bmatrix} -\frac{3e^{3t}}{2} \\ e^{3t} \end{bmatrix} + c_2 \begin{bmatrix} 2e^{10t} \\ e^{10t} \end{bmatrix}$$

Which becomes

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} -\frac{3c_1 e^{3t}}{2} + 2c_2 e^{10t} \\ c_1 e^{3t} + c_2 e^{10t} \end{bmatrix}$$

The following is the phase plot of the system.




Figure 500: Phase plot

28.7.3 Maple step by step solution

Let's solve

$$[x'(t) = 7x(t) + 6y(t), y'(t) = 2x(t) + 6y(t)]$$

- Define vector

$$\vec{x}(t) = \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

- Convert system into a vector equation

$$\vec{x}'(t) = \begin{bmatrix} 7 & 6 \\ 2 & 6 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

- System to solve

$$\vec{x}'(t) = \begin{bmatrix} 7 & 6 \\ 2 & 6 \end{bmatrix} \cdot \vec{x}(t)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 7 & 6 \\ 2 & 6 \end{bmatrix}$$

- Rewrite the system as

$$\vec{x}'(t) = A \cdot \vec{x}(t)$$

- To solve the system, find the eigenvalues and eigenvectors of A
- Eigenpairs of A

$$\left[\left[3, \begin{bmatrix} -\frac{3}{2} \\ 1 \end{bmatrix} \right], \left[10, \begin{bmatrix} 2 \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[3, \begin{bmatrix} -\frac{3}{2} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{x}_1 = e^{3t} \cdot \begin{bmatrix} -\frac{3}{2} \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[10, \begin{bmatrix} 2 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{x}_2 = e^{10t} \cdot \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{x} = c_1 \vec{x}_1 + c_2 \vec{x}_2$$

- Substitute solutions into the general solution

$$\vec{x} = c_1 e^{3t} \cdot \begin{bmatrix} -\frac{3}{2} \\ 1 \end{bmatrix} + c_2 e^{10t} \cdot \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

- Substitute in vector of dependent variables

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} -\frac{3c_1e^{3t}}{2} + 2c_2e^{10t} \\ c_1e^{3t} + c_2e^{10t} \end{bmatrix}$$

- Solution to the system of ODEs

$$\left\{ x(t) = -\frac{3c_1e^{3t}}{2} + 2c_2e^{10t}, y(t) = c_1e^{3t} + c_2e^{10t} \right\}$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 36

```
dsolve([diff(x(t),t)=7*x(t)+6*y(t),diff(y(t),t)=2*x(t)+6*y(t)],singsol=all)
```

$$\begin{aligned} x(t) &= c_1e^{10t} + c_2e^{3t} \\ y(t) &= \frac{c_1e^{10t}}{2} - \frac{2c_2e^{3t}}{3} \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 74

```
DSolve[{x'[t]==7*x[t]+6*y[t],y'[t]==2*x[t]+6*y[t]},{x[t],y[t]},t,IncludeSingularSolutions ->
```

$$\begin{aligned} x(t) &\rightarrow \frac{1}{7}e^{3t}(c_1(4e^{7t} + 3) + 6c_2(e^{7t} - 1)) \\ y(t) &\rightarrow \frac{1}{7}e^{3t}(2c_1(e^{7t} - 1) + c_2(3e^{7t} + 4)) \end{aligned}$$

28.8 problem 1(h)

- 28.8.1 Solution using Matrix exponential method 4065
- 28.8.2 Solution using explicit Eigenvalue and Eigenvector method . . . 4066
- 28.8.3 Maple step by step solution 4071

Internal problem ID [6527]

Internal file name [OUTPUT/5775_Sunday_June_05_2022_03_53_54_PM_78998523/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 10. Systems of First-Order Equations. Section 10.3 Homogeneous Linear Systems with Constant Coefficients. Page 387

Problem number: 1(h).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**system of linear ODEs**"

Solve

$$\begin{aligned}x'(t) &= x(t) - 2y(t) \\y'(t) &= 4x(t) + 5y(t)\end{aligned}$$

28.8.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A \vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 1 & -2 \\ 4 & 5 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

For the above matrix A , the matrix exponential can be found to be

$$\begin{aligned}e^{At} &= \begin{bmatrix} e^{3t} \cos(2t) - e^{3t} \sin(2t) & -e^{3t} \sin(2t) \\ 2e^{3t} \sin(2t) & e^{3t} \cos(2t) + e^{3t} \sin(2t) \end{bmatrix} \\ &= \begin{bmatrix} e^{3t}(\cos(2t) - \sin(2t)) & -e^{3t} \sin(2t) \\ 2e^{3t} \sin(2t) & e^{3t}(\cos(2t) + \sin(2t)) \end{bmatrix}\end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 \vec{x}_h(t) &= e^{At} \vec{c} \\
 &= \begin{bmatrix} e^{3t}(\cos(2t) - \sin(2t)) & -e^{3t} \sin(2t) \\ 2e^{3t} \sin(2t) & e^{3t}(\cos(2t) + \sin(2t)) \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} \\
 &= \begin{bmatrix} e^{3t}(\cos(2t) - \sin(2t))c_1 - e^{3t} \sin(2t)c_2 \\ 2e^{3t} \sin(2t)c_1 + e^{3t}(\cos(2t) + \sin(2t))c_2 \end{bmatrix} \\
 &= \begin{bmatrix} ((-c_1 - c_2) \sin(2t) + c_1 \cos(2t)) e^{3t} \\ e^{3t}(2c_1 + c_2) \sin(2t) + e^{3t} \cos(2t)c_2 \end{bmatrix}
 \end{aligned}$$

Since no forcing function is given, then the final solution is $\vec{x}_h(t)$ above.

28.8.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A \vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 1 & -2 \\ 4 & 5 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det \left(\begin{bmatrix} 1 & -2 \\ 4 & 5 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) = 0$$

Therefore

$$\det \left(\begin{bmatrix} 1 - \lambda & -2 \\ 4 & 5 - \lambda \end{bmatrix} \right) = 0$$

Which gives the characteristic equation

$$\lambda^2 - 6\lambda + 13 = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = 3 + 2i$$

$$\lambda_2 = 3 - 2i$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
$3 - 2i$	1	complex eigenvalue
$3 + 2i$	1	complex eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = 3 - 2i$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 1 & -2 \\ 4 & 5 \end{bmatrix} - (3 - 2i) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} -2 + 2i & -2 \\ 4 & 2 + 2i \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} -2 + 2i & -2 & 0 \\ 4 & 2 + 2i & 0 \end{array} \right]$$

$$R_2 = R_2 + (1 + i)R_1 \implies \left[\begin{array}{cc|c} -2 + 2i & -2 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} -2 + 2i & -2 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = (-\frac{1}{2} - \frac{i}{2})t\}$

Hence the solution is

$$\begin{bmatrix} (-\frac{1}{2} - \frac{i}{2})t \\ t \end{bmatrix} = \begin{bmatrix} (-\frac{1}{2} - \frac{i}{2})t \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} (-\frac{1}{2} - \frac{i}{2})t \\ t \end{bmatrix} = t \begin{bmatrix} -\frac{1}{2} - \frac{i}{2} \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} (-\frac{1}{2} - \frac{i}{2})t \\ t \end{bmatrix} = \begin{bmatrix} -\frac{1}{2} - \frac{i}{2} \\ 1 \end{bmatrix}$$

Which is normalized to

$$\begin{bmatrix} (-\frac{1}{2} - \frac{i}{2})t \\ t \end{bmatrix} = \begin{bmatrix} -1 - i \\ 2 \end{bmatrix}$$

Considering the eigenvalue $\lambda_2 = 3 + 2i$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 1 & -2 \\ 4 & 5 \end{bmatrix} - (3 + 2i) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} -2 - 2i & -2 \\ 4 & 2 - 2i \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} -2 - 2i & -2 & 0 \\ 4 & 2 - 2i & 0 \end{array} \right]$$

$$R_2 = R_2 + (1 - i)R_1 \implies \left[\begin{array}{cc|c} -2 - 2i & -2 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} -2 - 2i & -2 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = (-\frac{1}{2} + \frac{i}{2})t\}$

Hence the solution is

$$\begin{bmatrix} (-\frac{1}{2} + \frac{i}{2})t \\ t \end{bmatrix} = \begin{bmatrix} (-\frac{1}{2} + \frac{i}{2})t \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} (-\frac{1}{2} + \frac{i}{2})t \\ t \end{bmatrix} = t \begin{bmatrix} -\frac{1}{2} + \frac{i}{2} \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} (-\frac{1}{2} + \frac{i}{2})t \\ t \end{bmatrix} = \begin{bmatrix} -\frac{1}{2} + \frac{i}{2} \\ 1 \end{bmatrix}$$

Which is normalized to

$$\begin{bmatrix} (-\frac{1}{2} + \frac{i}{2})t \\ t \end{bmatrix} = \begin{bmatrix} -1 + i \\ 2 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
$3 + 2i$	1	1	No	$\begin{bmatrix} -\frac{1}{2} + \frac{i}{2} \\ 1 \end{bmatrix}$
$3 - 2i$	1	1	No	$\begin{bmatrix} -\frac{1}{2} - \frac{i}{2} \\ 1 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. Therefore the final solution is

$$\vec{x}_h(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

Which is written as

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = c_1 \begin{bmatrix} \left(-\frac{1}{2} + \frac{i}{2}\right) e^{(3+2i)t} \\ e^{(3+2i)t} \end{bmatrix} + c_2 \begin{bmatrix} \left(-\frac{1}{2} - \frac{i}{2}\right) e^{(3-2i)t} \\ e^{(3-2i)t} \end{bmatrix}$$

Which becomes

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} \left(-\frac{1}{2} + \frac{i}{2}\right) c_1 e^{(3+2i)t} + \left(-\frac{1}{2} - \frac{i}{2}\right) c_2 e^{(3-2i)t} \\ c_1 e^{(3+2i)t} + c_2 e^{(3-2i)t} \end{bmatrix}$$

The following is the phase plot of the system.




Figure 501: Phase plot

28.8.3 Maple step by step solution

Let's solve

$$[x'(t) = x(t) - 2y(t), y'(t) = 4x(t) + 5y(t)]$$

- Define vector

$$\vec{x}(t) = \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

- Convert system into a vector equation

$$\vec{x}'(t) = \begin{bmatrix} 1 & -2 \\ 4 & 5 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

- System to solve

$$\vec{x}'(t) = \begin{bmatrix} 1 & -2 \\ 4 & 5 \end{bmatrix} \cdot \vec{x}(t)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 1 & -2 \\ 4 & 5 \end{bmatrix}$$

- Rewrite the system as

$$\vec{x}'(t) = A \cdot \vec{x}(t)$$

- To solve the system, find the eigenvalues and eigenvectors of A
- Eigenpairs of A

$$\left[\left[3 - 2I, \begin{bmatrix} -\frac{1}{2} - \frac{I}{2} \\ 1 \end{bmatrix} \right], \left[3 + 2I, \begin{bmatrix} -\frac{1}{2} + \frac{I}{2} \\ 1 \end{bmatrix} \right] \right]$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$\left[3 - 2I, \begin{bmatrix} -\frac{1}{2} - \frac{I}{2} \\ 1 \end{bmatrix} \right]$$

- Solution from eigenpair

$$e^{(3-2I)t} \cdot \begin{bmatrix} -\frac{1}{2} - \frac{I}{2} \\ 1 \end{bmatrix}$$

- Use Euler identity to write solution in terms of sin and cos

$$e^{3t} \cdot (\cos(2t) - I \sin(2t)) \cdot \begin{bmatrix} -\frac{1}{2} - \frac{I}{2} \\ 1 \end{bmatrix}$$

- Simplify expression

$$e^{3t} \cdot \begin{bmatrix} \left(-\frac{1}{2} - \frac{I}{2}\right) (\cos(2t) - I \sin(2t)) \\ \cos(2t) - I \sin(2t) \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\left[\vec{x}_1(t) = e^{3t} \cdot \begin{bmatrix} -\frac{\cos(2t)}{2} - \frac{\sin(2t)}{2} \\ \cos(2t) \end{bmatrix}, \vec{x}_2(t) = e^{3t} \cdot \begin{bmatrix} \frac{\sin(2t)}{2} - \frac{\cos(2t)}{2} \\ -\sin(2t) \end{bmatrix} \right]$$

- General solution to the system of ODEs

$$\vec{x} = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

- Substitute solutions into the general solution

$$\vec{x} = c_1 e^{3t} \cdot \begin{bmatrix} -\frac{\cos(2t)}{2} - \frac{\sin(2t)}{2} \\ \cos(2t) \end{bmatrix} + e^{3t} c_2 \cdot \begin{bmatrix} \frac{\sin(2t)}{2} - \frac{\cos(2t)}{2} \\ -\sin(2t) \end{bmatrix}$$

- Substitute in vector of dependent variables

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} -\frac{((c_1+c_2)\cos(2t)+\sin(2t)(c_1-c_2))e^{3t}}{2} \\ e^{3t}(c_1 \cos(2t) - c_2 \sin(2t)) \end{bmatrix}$$

- Solution to the system of ODEs

$$\left\{ x(t) = -\frac{((c_1+c_2)\cos(2t)+\sin(2t)(c_1-c_2))e^{3t}}{2}, y(t) = e^{3t}(c_1 \cos(2t) - c_2 \sin(2t)) \right\}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 58

```
dsolve([diff(x(t),t)=x(t)-2*y(t),diff(y(t),t)=4*x(t)+5*y(t)],singsol=all)
```

$$\begin{aligned} x(t) &= e^{3t}(c_1 \sin(2t) + c_2 \cos(2t)) \\ y(t) &= -e^{3t}(c_1 \sin(2t) - c_2 \sin(2t) + c_1 \cos(2t) + c_2 \cos(2t)) \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.005 (sec). Leaf size: 59

```
DSolve[{x'[t]==x[t]-2*y[t],y'[t]==4*x[t]+5*y[t]},{x[t],y[t]},t,IncludeSingularSolutions -> T
```

$$\begin{aligned} x(t) &\rightarrow e^{3t}(c_1 \cos(2t) - (c_1 + c_2) \sin(2t)) \\ y(t) &\rightarrow e^{3t}(c_2 \cos(2t) + (2c_1 + c_2) \sin(2t)) \end{aligned}$$

28.9 problem 5(b)

28.9.1 Solution using Matrix exponential method	4074
28.9.2 Solution using explicit Eigenvalue and Eigenvector method . . .	4076
28.9.3 Maple step by step solution	4081

Internal problem ID [6528]

Internal file name [OUTPUT/5776_Sunday_June_05_2022_03_53_56_PM_44093963/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 10. Systems of First-Order Equations. Section 10.3 Homogeneous Linear Systems with Constant Coefficients. Page 387

Problem number: 5(b).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**system of linear ODEs**"

Solve

$$\begin{aligned}x'(t) &= x(t) + y(t) - 5t + 2 \\y'(t) &= 4x(t) - 2y(t) - 8t - 8\end{aligned}$$

28.9.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t) + \vec{G}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 4 & -2 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix} + \begin{bmatrix} -5t + 2 \\ -8t - 8 \end{bmatrix}$$

Since the system is nonhomogeneous, then the solution is given by

$$\vec{x}(t) = \vec{x}_h(t) + \vec{x}_p(t)$$

Where $\vec{x}_h(t)$ is the homogeneous solution to $\vec{x}'(t) = A\vec{x}(t)$ and $\vec{x}_p(t)$ is a particular solution to $\vec{x}'(t) = A\vec{x}(t) + \vec{G}(t)$. The particular solution will be found using variation

of parameters method applied to the fundamental matrix. For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} \frac{(4e^{5t}+1)e^{-3t}}{5} & \frac{(e^{5t}-1)e^{-3t}}{5} \\ \frac{4(e^{5t}-1)e^{-3t}}{5} & \frac{(e^{5t}+4)e^{-3t}}{5} \end{bmatrix}$$

Therefore the homogeneous solution is

$$\begin{aligned} \vec{x}_h(t) &= e^{At} \vec{c} \\ &= \begin{bmatrix} \frac{(4e^{5t}+1)e^{-3t}}{5} & \frac{(e^{5t}-1)e^{-3t}}{5} \\ \frac{4(e^{5t}-1)e^{-3t}}{5} & \frac{(e^{5t}+4)e^{-3t}}{5} \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} \\ &= \begin{bmatrix} \frac{(4e^{5t}+1)e^{-3t}c_1}{5} + \frac{(e^{5t}-1)e^{-3t}c_2}{5} \\ \frac{4(e^{5t}-1)e^{-3t}c_1}{5} + \frac{(e^{5t}+4)e^{-3t}c_2}{5} \end{bmatrix} \\ &= \begin{bmatrix} \frac{e^{-3t}((4c_1+c_2)e^{5t}-c_2+c_1)}{5} \\ \frac{4((c_1+\frac{c_2}{4})e^{5t}+c_2-c_1)e^{-3t}}{5} \end{bmatrix} \end{aligned}$$

The particular solution given by

$$\vec{x}_p(t) = e^{At} \int e^{-At} \vec{G}(t) dt$$

But

$$\begin{aligned} e^{-At} &= (e^{At})^{-1} \\ &= \begin{bmatrix} \frac{(e^{5t}+4)e^{-2t}}{5} & -\frac{(e^{5t}-1)e^{-2t}}{5} \\ -\frac{4(e^{5t}-1)e^{-2t}}{5} & \frac{(4e^{5t}+1)e^{-2t}}{5} \end{bmatrix} \end{aligned}$$

Hence

$$\begin{aligned} \vec{x}_p(t) &= \begin{bmatrix} \frac{(4e^{5t}+1)e^{-3t}}{5} & \frac{(e^{5t}-1)e^{-3t}}{5} \\ \frac{4(e^{5t}-1)e^{-3t}}{5} & \frac{(e^{5t}+4)e^{-3t}}{5} \end{bmatrix} \int \begin{bmatrix} \frac{(e^{5t}+4)e^{-2t}}{5} & -\frac{(e^{5t}-1)e^{-2t}}{5} \\ -\frac{4(e^{5t}-1)e^{-2t}}{5} & \frac{(4e^{5t}+1)e^{-2t}}{5} \end{bmatrix} \begin{bmatrix} -5t+2 \\ -8t-8 \end{bmatrix} dt \\ &= \begin{bmatrix} \frac{(4e^{5t}+1)e^{-3t}}{5} & \frac{(e^{5t}-1)e^{-3t}}{5} \\ \frac{4(e^{5t}-1)e^{-3t}}{5} & \frac{(e^{5t}+4)e^{-3t}}{5} \end{bmatrix} \begin{bmatrix} \frac{((t+3)e^{5t}+14t+7)e^{-2t}}{5} \\ -\frac{(4e^{5t}t+12e^{5t}-14t-7)e^{-2t}}{5} \end{bmatrix} \\ &= \begin{bmatrix} 2+3t \\ 2t-1 \end{bmatrix} \end{aligned}$$

Hence the complete solution is

$$\begin{aligned}\vec{x}(t) &= \vec{x}_h(t) + \vec{x}_p(t) \\ &= \begin{bmatrix} \frac{(4e^{5t}c_1 + e^{5t}c_2 + 15te^{3t} + 10e^{3t} + c_1 - c_2)e^{-3t}}{5} \\ \frac{(4e^{5t}c_1 + e^{5t}c_2 + 10te^{3t} - 5e^{3t} - 4c_1 + 4c_2)e^{-3t}}{5} \end{bmatrix}\end{aligned}$$

28.9.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t) + \vec{G}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 4 & -2 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix} + \begin{bmatrix} -5t + 2 \\ -8t - 8 \end{bmatrix}$$

Since the system is nonhomogeneous, then the solution is given by

$$\vec{x}(t) = \vec{x}_h(t) + \vec{x}_p(t)$$

Where $\vec{x}_h(t)$ is the homogeneous solution to $\vec{x}'(t) = A\vec{x}(t)$ and $\vec{x}_p(t)$ is a particular solution to $\vec{x}'(t) = A\vec{x}(t) + \vec{G}(t)$. The particular solution will be found using variation of parameters method applied to the fundamental matrix.

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det\left(\begin{bmatrix} 1 & 1 \\ 4 & -2 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}\right) = 0$$

Therefore

$$\det\left(\begin{bmatrix} 1 - \lambda & 1 \\ 4 & -2 - \lambda \end{bmatrix}\right) = 0$$

Which gives the characteristic equation

$$\lambda^2 + \lambda - 6 = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = -3$$

$$\lambda_2 = 2$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
-3	1	real eigenvalue
2	1	real eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = -3$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 1 & 1 \\ 4 & -2 \end{bmatrix} - (-3) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 4 & 1 \\ 4 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} 4 & 1 & 0 \\ 4 & 1 & 0 \end{array} \right]$$

$$R_2 = R_2 - R_1 \implies \left[\begin{array}{cc|c} 4 & 1 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} 4 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = -\frac{t}{4}\}$

Hence the solution is

$$\begin{bmatrix} -\frac{t}{4} \\ t \end{bmatrix} = \begin{bmatrix} -\frac{t}{4} \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} -\frac{t}{4} \\ t \end{bmatrix} = t \begin{bmatrix} -\frac{1}{4} \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} -\frac{1}{4} \\ 1 \end{bmatrix} = \begin{bmatrix} -\frac{1}{4} \\ 1 \end{bmatrix}$$

Which is normalized to

$$\begin{bmatrix} -\frac{1}{4} \\ 1 \end{bmatrix} = \begin{bmatrix} -1 \\ 4 \end{bmatrix}$$

Considering the eigenvalue $\lambda_2 = 2$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 1 & 1 \\ 4 & -2 \end{bmatrix} - (2) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} -1 & 1 \\ 4 & -4 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} -1 & 1 & 0 \\ 4 & -4 & 0 \end{array} \right]$$

$$R_2 = R_2 + 4R_1 \implies \left[\begin{array}{cc|c} -1 & 1 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} -1 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = t\}$

Hence the solution is

$$\begin{bmatrix} t \\ t \end{bmatrix} = \begin{bmatrix} t \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} t \\ t \end{bmatrix} = t \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} t \\ t \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
-3	1	1	No	$\begin{bmatrix} -\frac{1}{4} \\ 1 \end{bmatrix}$
2	1	1	No	$\begin{bmatrix} 1 \\ 1 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. Since eigenvalue -3 is real and distinct then the

corresponding eigenvector solution is

$$\begin{aligned}\vec{x}_1(t) &= \vec{v}_1 e^{-3t} \\ &= \begin{bmatrix} -\frac{1}{4} \\ 1 \end{bmatrix} e^{-3t}\end{aligned}$$

Since eigenvalue 2 is real and distinct then the corresponding eigenvector solution is

$$\begin{aligned}\vec{x}_2(t) &= \vec{v}_2 e^{2t} \\ &= \begin{bmatrix} 1 \\ 1 \end{bmatrix} e^{2t}\end{aligned}$$

Therefore the homogeneous solution is

$$\vec{x}_h(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

Which is written as

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = c_1 \begin{bmatrix} -\frac{e^{-3t}}{4} \\ e^{-3t} \end{bmatrix} + c_2 \begin{bmatrix} e^{2t} \\ e^{2t} \end{bmatrix}$$

Now that we found homogeneous solution above, we need to find a particular solution $\vec{x}_p(t)$. We will use Variation of parameters. The fundamental matrix is

$$\Phi = \begin{bmatrix} \vec{x}_1 & \vec{x}_2 & \cdots \end{bmatrix}$$

Where \vec{x}_i are the solution basis found above. Therefore the fundamental matrix is

$$\Phi(t) = \begin{bmatrix} -\frac{e^{-3t}}{4} & e^{2t} \\ e^{-3t} & e^{2t} \end{bmatrix}$$

The particular solution is then given by

$$\vec{x}_p(t) = \Phi \int \Phi^{-1} \vec{G}(t) dt$$

But

$$\Phi^{-1} = \begin{bmatrix} -\frac{4e^{3t}}{5} & \frac{4e^{3t}}{5} \\ \frac{4e^{-2t}}{5} & \frac{e^{-2t}}{5} \end{bmatrix}$$

Hence

$$\begin{aligned}
 \vec{x}_p(t) &= \begin{bmatrix} -\frac{e^{-3t}}{4} & e^{2t} \\ e^{-3t} & e^{2t} \end{bmatrix} \int \begin{bmatrix} -\frac{4e^{3t}}{5} & \frac{4e^{3t}}{5} \\ \frac{4e^{-2t}}{5} & \frac{e^{-2t}}{5} \end{bmatrix} \begin{bmatrix} -5t + 2 \\ -8t - 8 \end{bmatrix} dt \\
 &= \begin{bmatrix} -\frac{e^{-3t}}{4} & e^{2t} \\ e^{-3t} & e^{2t} \end{bmatrix} \int \begin{bmatrix} 4e^{3t}(-2 - \frac{3t}{5}) \\ -\frac{28te^{-2t}}{5} \end{bmatrix} dt \\
 &= \begin{bmatrix} -\frac{e^{-3t}}{4} & e^{2t} \\ e^{-3t} & e^{2t} \end{bmatrix} \begin{bmatrix} -\frac{4e^{3t}(t+3)}{5} \\ \frac{7(2t+1)e^{-2t}}{5} \end{bmatrix} \\
 &= \begin{bmatrix} 2 + 3t \\ 2t - 1 \end{bmatrix}
 \end{aligned}$$

Now that we found particular solution, the final solution is

$$\begin{aligned}
 \vec{x}(t) &= \vec{x}_h(t) + \vec{x}_p(t) \\
 \begin{bmatrix} x(t) \\ y(t) \end{bmatrix} &= \begin{bmatrix} -\frac{c_1 e^{-3t}}{4} \\ c_1 e^{-3t} \end{bmatrix} + \begin{bmatrix} c_2 e^{2t} \\ c_2 e^{2t} \end{bmatrix} + \begin{bmatrix} 2 + 3t \\ 2t - 1 \end{bmatrix}
 \end{aligned}$$

Which becomes

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} \frac{(4c_2 e^{5t} + 12t e^{3t} + 8e^{3t} - c_1) e^{-3t}}{4} \\ (c_2 e^{5t} + 2t e^{3t} - e^{3t} + c_1) e^{-3t} \end{bmatrix}$$

28.9.3 Maple step by step solution

Let's solve

$$[x'(t) = x(t) + y(t) - 5t + 2, y'(t) = 4x(t) - 2y(t) - 8t - 8]$$

- Define vector

$$\vec{x}(t) = \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

- Convert system into a vector equation

$$\vec{x}'(t) = \begin{bmatrix} 1 & 1 \\ 4 & -2 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} -5t + 2 \\ -8t - 8 \end{bmatrix}$$

- System to solve

$$\vec{x}'(t) = \begin{bmatrix} 1 & 1 \\ 4 & -2 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} -5t + 2 \\ -8t - 8 \end{bmatrix}$$

- Define the forcing function

$$\vec{f}(t) = \begin{bmatrix} -5t + 2 \\ -8t - 8 \end{bmatrix}$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 1 & 1 \\ 4 & -2 \end{bmatrix}$$

- Rewrite the system as

$$\vec{x}'(t) = A \cdot \vec{x}(t) + \vec{f}$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[-3, \begin{bmatrix} -\frac{1}{4} \\ 1 \end{bmatrix} \right], \left[2, \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[-3, \begin{bmatrix} -\frac{1}{4} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{x}_1 = e^{-3t} \cdot \begin{bmatrix} -\frac{1}{4} \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[2, \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{x}_2 = e^{2t} \cdot \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

- General solution of the system of ODEs can be written in terms of the particular solution $\vec{x}_p(t)$

$$\vec{x}(t) = c_1 \vec{x}_1 + c_2 \vec{x}_2 + \vec{x}_p(t)$$

□ Fundamental matrix

- Let $\phi(t)$ be the matrix whose columns are the independent solutions of the homogeneous system

$$\phi(t) = \begin{bmatrix} -\frac{e^{-3t}}{4} & e^{2t} \\ e^{-3t} & e^{2t} \end{bmatrix}$$

- The fundamental matrix, $\Phi(t)$ is a normalized version of $\phi(t)$ satisfying $\Phi(0) = I$ where I is the identity matrix

$$\Phi(t) = \phi(t) \cdot \frac{1}{\phi(0)}$$

- Substitute the value of $\phi(t)$ and $\phi(0)$

$$\Phi(t) = \begin{bmatrix} -\frac{e^{-3t}}{4} & e^{2t} \\ e^{-3t} & e^{2t} \end{bmatrix} \cdot \frac{1}{\begin{bmatrix} -\frac{1}{4} & 1 \\ 1 & 1 \end{bmatrix}}$$

- Evaluate and simplify to get the fundamental matrix

$$\Phi(t) = \begin{bmatrix} \frac{(4e^{5t}+1)e^{-3t}}{5} & \frac{(e^{5t}-1)e^{-3t}}{5} \\ \frac{4(e^{5t}-1)e^{-3t}}{5} & \frac{(e^{5t}+4)e^{-3t}}{5} \end{bmatrix}$$

□ Find a particular solution of the system of ODEs using variation of parameters

- Let the particular solution be the fundamental matrix multiplied by $\vec{v}(t)$ and solve for $\vec{v}(t)$

$$\vec{x}_p(t) = \Phi(t) \cdot \vec{v}(t)$$

- Take the derivative of the particular solution

$$\vec{x}'_p(t) = \Phi'(t) \cdot \vec{v}(t) + \Phi(t) \cdot \vec{v}'(t)$$

- Substitute particular solution and its derivative into the system of ODEs

$$\Phi'(t) \cdot \vec{v}(t) + \Phi(t) \cdot \vec{v}'(t) = A \cdot \Phi(t) \cdot \vec{v}(t) + \vec{f}(t)$$

- The fundamental matrix has columns that are solutions to the homogeneous system so its derivative is $A \cdot \Phi(t)$

$$A \cdot \Phi(t) \cdot \vec{v}(t) + \Phi(t) \cdot \vec{v}'(t) = A \cdot \Phi(t) \cdot \vec{v}(t) + \vec{f}(t)$$

- Cancel like terms

$$\Phi(t) \cdot \vec{v}'(t) = \vec{f}(t)$$

- Multiply by the inverse of the fundamental matrix

$$\vec{v}'(t) = \frac{1}{\Phi(t)} \cdot \vec{f}(t)$$

- Integrate to solve for $\vec{v}(t)$

$$\vec{v}(t) = \int_0^t \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds$$

- Plug $\vec{v}(t)$ into the equation for the particular solution

$$\vec{x}_p(t) = \Phi(t) \cdot \left(\int_0^t \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds \right)$$

- Plug in the fundamental matrix and the forcing function and compute

$$\vec{x}_p(t) = \begin{bmatrix} \frac{(-7e^{5t} + 15te^{3t} + 10e^{3t} - 3)e^{-3t}}{5} \\ \frac{(-7e^{5t} + 10te^{3t} - 5e^{3t} + 12)e^{-3t}}{5} \end{bmatrix}$$

- Plug particular solution back into general solution

$$\vec{x}(t) = c_1 \vec{x}_1 + c_2 \vec{x}_2 + \begin{bmatrix} \frac{(-7e^{5t} + 15te^{3t} + 10e^{3t} - 3)e^{-3t}}{5} \\ \frac{(-7e^{5t} + 10te^{3t} - 5e^{3t} + 12)e^{-3t}}{5} \end{bmatrix}$$

- Substitute in vector of dependent variables

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} \frac{(20c_2e^{5t} - 28e^{5t} + 60te^{3t} + 40e^{3t} - 5c_1 - 12)e^{-3t}}{20} \\ \frac{(5c_2e^{5t} - 7e^{5t} + 10te^{3t} - 5e^{3t} + 5c_1 + 12)e^{-3t}}{5} \end{bmatrix}$$

- Solution to the system of ODEs

$$\left\{ x(t) = \frac{(20c_2e^{5t} - 28e^{5t} + 60te^{3t} + 40e^{3t} - 5c_1 - 12)e^{-3t}}{20}, y(t) = \frac{(5c_2e^{5t} - 7e^{5t} + 10te^{3t} - 5e^{3t} + 5c_1 + 12)e^{-3t}}{5} \right\}$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 43

```
dsolve([diff(x(t),t)=x(t)+y(t)-5*t+2,diff(y(t),t)=4*x(t)-2*y(t)-8*t-8],singsol=all)
```

$$\begin{aligned} x(t) &= c_2e^{-3t} + c_1e^{2t} + 3t + 2 \\ y(t) &= -4c_2e^{-3t} + c_1e^{2t} - 1 + 2t \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.15 (sec). Leaf size: 92

```
DSolve[{x'[t]==x[t]+y[t]-5*t+2,y'[t]==4*x[t]-2*y[t]-8*t-8},{x[t],y[t]},t,IncludeSingularSolu
```

$$x(t) \rightarrow \frac{1}{5}e^{-3t}(5e^{3t}(3t+2) + (4c_1 + c_2)e^{5t} + c_1 - c_2)$$

$$y(t) \rightarrow \frac{1}{5}e^{-3t}(5e^{3t}(2t-1) + (4c_1 + c_2)e^{5t} - 4c_1 + 4c_2)$$

29 Chapter 10. Systems of First-Order Equations.
Section A. Drill exercises. Page 400

29.1	problem 2(a)	4087
29.2	problem 2(b)	4096
29.3	problem 2(c)	4105
29.4	problem 2(d)	4112
29.5	problem 3(a)	4121
29.6	problem 3(b)	4131
29.7	problem 3(c)	4139
29.8	problem 3(d)	4148
29.9	problem 3(e)	4157
29.10	problem 3(f)	4170
29.11	problem 4(a)	4186
29.12	problem 4(b)	4199
29.13	problem 4(c)	4211

29.1 problem 2(a)

29.1.1 Solution using Matrix exponential method	4087
29.1.2 Solution using explicit Eigenvalue and Eigenvector method . . .	4088
29.1.3 Maple step by step solution	4093

Internal problem ID [6529]

Internal file name [OUTPUT/5777_Sunday_June_05_2022_03_53_59_PM_74613779/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 10. Systems of First-Order Equations. Section A. Drill exercises. Page 400

Problem number: 2(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "system of linear ODEs"

Solve

$$x'(t) = 3x(t) - 4y(t)$$

$$y'(t) = 4x(t) - 7y(t)$$

29.1.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A \vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 3 & -4 \\ 4 & -7 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} \frac{(4e^{6t}-1)e^{-5t}}{3} & -\frac{2(e^{6t}-1)e^{-5t}}{3} \\ \frac{2(e^{6t}-1)e^{-5t}}{3} & -\frac{(e^{6t}-4)e^{-5t}}{3} \end{bmatrix}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 \vec{x}_h(t) &= e^{At} \vec{c} \\
 &= \begin{bmatrix} \frac{(4e^{6t}-1)e^{-5t}}{3} & -\frac{2(e^{6t}-1)e^{-5t}}{3} \\ \frac{2(e^{6t}-1)e^{-5t}}{3} & -\frac{(e^{6t}-4)e^{-5t}}{3} \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} \\
 &= \begin{bmatrix} \frac{(4e^{6t}-1)e^{-5t}c_1}{3} - \frac{2(e^{6t}-1)e^{-5t}c_2}{3} \\ \frac{2(e^{6t}-1)e^{-5t}c_1}{3} - \frac{(e^{6t}-4)e^{-5t}c_2}{3} \end{bmatrix} \\
 &= \begin{bmatrix} \frac{e^{-5t}(2(-c_2+2c_1)e^{6t}+2c_2-c_1)}{3} \\ \frac{2e^{-5t}((c_1-\frac{c_2}{2})e^{6t}+2c_2-c_1)}{3} \end{bmatrix}
 \end{aligned}$$

Since no forcing function is given, then the final solution is $\vec{x}_h(t)$ above.

29.1.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 3 & -4 \\ 4 & -7 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det \left(\begin{bmatrix} 3 & -4 \\ 4 & -7 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) = 0$$

Therefore

$$\det \left(\begin{bmatrix} 3 - \lambda & -4 \\ 4 & -7 - \lambda \end{bmatrix} \right) = 0$$

Which gives the characteristic equation

$$\lambda^2 + 4\lambda - 5 = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = 1$$

$$\lambda_2 = -5$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
1	1	real eigenvalue
-5	1	real eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = -5$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 3 & -4 \\ 4 & -7 \end{bmatrix} - (-5) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$
$$\begin{bmatrix} 8 & -4 \\ 4 & -2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} 8 & -4 & 0 \\ 4 & -2 & 0 \end{array} \right]$$

$$R_2 = R_2 - \frac{R_1}{2} \implies \left[\begin{array}{cc|c} 8 & -4 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} 8 & -4 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = \frac{t}{2}\}$

Hence the solution is

$$\begin{bmatrix} \frac{t}{2} \\ t \end{bmatrix} = \begin{bmatrix} \frac{t}{2} \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} \frac{t}{2} \\ t \end{bmatrix} = t \begin{bmatrix} \frac{1}{2} \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} \frac{1}{2} \\ 1 \end{bmatrix} = \begin{bmatrix} \frac{1}{2} \\ 1 \end{bmatrix}$$

Which is normalized to

$$\begin{bmatrix} \frac{1}{2} \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

Considering the eigenvalue $\lambda_2 = 1$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 3 & -4 \\ 4 & -7 \end{bmatrix} - (1) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 2 & -4 \\ 4 & -8 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} 2 & -4 & 0 \\ 4 & -8 & 0 \end{array} \right]$$

$$R_2 = R_2 - 2R_1 \implies \left[\begin{array}{cc|c} 2 & -4 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} 2 & -4 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = 2t\}$

Hence the solution is

$$\begin{bmatrix} 2t \\ t \end{bmatrix} = \begin{bmatrix} 2t \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} 2t \\ t \end{bmatrix} = t \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} 2t \\ t \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
1	1	1	No	$\begin{bmatrix} 2 \\ 1 \end{bmatrix}$
-5	1	1	No	$\begin{bmatrix} \frac{1}{2} \\ 1 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. Since eigenvalue 1 is real and distinct then the corresponding eigenvector solution is

$$\begin{aligned}\vec{x}_1(t) &= \vec{v}_1 e^t \\ &= \begin{bmatrix} 2 \\ 1 \end{bmatrix} e^t\end{aligned}$$

Since eigenvalue -5 is real and distinct then the corresponding eigenvector solution is

$$\begin{aligned}\vec{x}_2(t) &= \vec{v}_2 e^{-5t} \\ &= \begin{bmatrix} \frac{1}{2} \\ 1 \end{bmatrix} e^{-5t}\end{aligned}$$

Therefore the final solution is

$$\vec{x}_h(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

Which is written as

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = c_1 \begin{bmatrix} 2e^t \\ e^t \end{bmatrix} + c_2 \begin{bmatrix} \frac{e^{-5t}}{2} \\ e^{-5t} \end{bmatrix}$$

Which becomes

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} \frac{(4c_1 e^{6t} + c_2) e^{-5t}}{2} \\ (c_1 e^{6t} + c_2) e^{-5t} \end{bmatrix}$$

The following is the phase plot of the system.




Figure 502: Phase plot

29.1.3 Maple step by step solution

Let's solve

$$[x'(t) = 3x(t) - 4y(t), y'(t) = 4x(t) - 7y(t)]$$

- Define vector

$$\vec{x}(t) = \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

- Convert system into a vector equation

$$\vec{x}'(t) = \begin{bmatrix} 3 & -4 \\ 4 & -7 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

- System to solve

$$\vec{x}'(t) = \begin{bmatrix} 3 & -4 \\ 4 & -7 \end{bmatrix} \cdot \vec{x}(t)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 3 & -4 \\ 4 & -7 \end{bmatrix}$$

- Rewrite the system as

$$\vec{x}'(t) = A \cdot \vec{x}(t)$$

- To solve the system, find the eigenvalues and eigenvectors of A
- Eigenpairs of A

$$\left[\left[-5, \begin{bmatrix} \frac{1}{2} \\ 1 \end{bmatrix} \right], \left[1, \begin{bmatrix} 2 \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[-5, \begin{bmatrix} \frac{1}{2} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{x}_1 = e^{-5t} \cdot \begin{bmatrix} \frac{1}{2} \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[1, \begin{bmatrix} 2 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{x}_2 = e^t \cdot \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{x} = c_1 \vec{x}_1 + c_2 \vec{x}_2$$

- Substitute solutions into the general solution

$$\vec{x} = c_1 e^{-5t} \cdot \begin{bmatrix} \frac{1}{2} \\ 1 \end{bmatrix} + c_2 e^t \cdot \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

- Substitute in vector of dependent variables

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} \frac{(4c_2e^{6t}+c_1)e^{-5t}}{2} \\ (c_2e^{6t} + c_1)e^{-5t} \end{bmatrix}$$

- Solution to the system of ODEs

$$\left\{ x(t) = \frac{(4c_2e^{6t}+c_1)e^{-5t}}{2}, y(t) = (c_2e^{6t} + c_1)e^{-5t} \right\}$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 32

```
dsolve([diff(x(t),t)=3*x(t)-4*y(t),diff(y(t),t)=4*x(t)-7*y(t)],singsol=all)
```

$$\begin{aligned} x(t) &= c_1e^{-5t} + c_2e^t \\ y(t) &= 2c_1e^{-5t} + \frac{c_2e^t}{2} \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 73

```
DSolve[{x'[t]==3*x[t]-4*y[t],y'[t]==4*x[t]-7*y[t]},{x[t],y[t]},t,IncludeSingularSolutions ->
```

$$\begin{aligned} x(t) &\rightarrow \frac{1}{3}e^{-5t}(c_1(4e^{6t} - 1) - 2c_2(e^{6t} - 1)) \\ y(t) &\rightarrow \frac{1}{3}e^{-5t}(2c_1(e^{6t} - 1) - c_2(e^{6t} - 4)) \end{aligned}$$

29.2 problem 2(b)

29.2.1 Solution using Matrix exponential method	4096
29.2.2 Solution using explicit Eigenvalue and Eigenvector method . . .	4097
29.2.3 Maple step by step solution	4102

Internal problem ID [6530]

Internal file name [OUTPUT/5778_Sunday_June_05_2022_03_54_01_PM_19296232/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 10. Systems of First-Order Equations. Section A. Drill exercises. Page 400

Problem number: 2(b).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**system of linear ODEs**"

Solve

$$x'(t) = x(t) + y(t)$$

$$y'(t) = 4x(t) + y(t)$$

29.2.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A \vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} \frac{e^{-t}}{2} + \frac{e^{3t}}{2} & \frac{e^{3t}}{4} - \frac{e^{-t}}{4} \\ e^{3t} - e^{-t} & \frac{e^{-t}}{2} + \frac{e^{3t}}{2} \end{bmatrix}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 \vec{x}_h(t) &= e^{At} \vec{c} \\
 &= \begin{bmatrix} \frac{e^{-t}}{2} + \frac{e^{3t}}{2} & \frac{e^{3t}}{4} - \frac{e^{-t}}{4} \\ e^{3t} - e^{-t} & \frac{e^{-t}}{2} + \frac{e^{3t}}{2} \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} \\
 &= \begin{bmatrix} \left(\frac{e^{-t}}{2} + \frac{e^{3t}}{2}\right) c_1 + \left(\frac{e^{3t}}{4} - \frac{e^{-t}}{4}\right) c_2 \\ (e^{3t} - e^{-t}) c_1 + \left(\frac{e^{-t}}{2} + \frac{e^{3t}}{2}\right) c_2 \end{bmatrix} \\
 &= \begin{bmatrix} \frac{(-c_2 + 2c_1)e^{-t}}{4} + \frac{(c_1 + \frac{c_2}{2})e^{3t}}{2} \\ \frac{(c_2 - 2c_1)e^{-t}}{2} + \left(c_1 + \frac{c_2}{2}\right) e^{3t} \end{bmatrix}
 \end{aligned}$$

Since no forcing function is given, then the final solution is $\vec{x}_h(t)$ above.

29.2.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A \vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det \left(\begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) = 0$$

Therefore

$$\det \left(\begin{bmatrix} 1 - \lambda & 1 \\ 4 & 1 - \lambda \end{bmatrix} \right) = 0$$

Which gives the characteristic equation

$$\lambda^2 - 2\lambda - 3 = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = -1$$

$$\lambda_2 = 3$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
-1	1	real eigenvalue
3	1	real eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = -1$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix} - (-1) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 1 \\ 4 & 2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} 2 & 1 & 0 \\ 4 & 2 & 0 \end{array} \right]$$

$$R_2 = R_2 - 2R_1 \implies \left[\begin{array}{cc|c} 2 & 1 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} 2 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = -\frac{t}{2}\}$

Hence the solution is

$$\begin{bmatrix} -\frac{t}{2} \\ t \end{bmatrix} = \begin{bmatrix} -\frac{t}{2} \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} -\frac{t}{2} \\ t \end{bmatrix} = t \begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix} = \begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix}$$

Which is normalized to

$$\begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix} = \begin{bmatrix} -1 \\ 2 \end{bmatrix}$$

Considering the eigenvalue $\lambda_2 = 3$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix} - (3) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} -2 & 1 \\ 4 & -2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} -2 & 1 & 0 \\ 4 & -2 & 0 \end{array} \right]$$

$$R_2 = R_2 + 2R_1 \implies \left[\begin{array}{cc|c} -2 & 1 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} -2 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = \frac{t}{2}\}$

Hence the solution is

$$\begin{bmatrix} \frac{t}{2} \\ t \end{bmatrix} = \begin{bmatrix} \frac{t}{2} \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} \frac{t}{2} \\ t \end{bmatrix} = t \begin{bmatrix} \frac{1}{2} \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} \frac{1}{2} \\ 1 \end{bmatrix} = \begin{bmatrix} \frac{1}{2} \\ 1 \end{bmatrix}$$

Which is normalized to

$$\begin{bmatrix} \frac{1}{2} \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
-1	1	1	No	$\begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix}$
3	1	1	No	$\begin{bmatrix} \frac{1}{2} \\ 1 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. Since eigenvalue -1 is real and distinct then the corresponding eigenvector solution is

$$\begin{aligned}\vec{x}_1(t) &= \vec{v}_1 e^{-t} \\ &= \begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix} e^{-t}\end{aligned}$$

Since eigenvalue 3 is real and distinct then the corresponding eigenvector solution is

$$\begin{aligned}\vec{x}_2(t) &= \vec{v}_2 e^{3t} \\ &= \begin{bmatrix} \frac{1}{2} \\ 1 \end{bmatrix} e^{3t}\end{aligned}$$

Therefore the final solution is

$$\vec{x}_h(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

Which is written as

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = c_1 \begin{bmatrix} -\frac{e^{-t}}{2} \\ e^{-t} \end{bmatrix} + c_2 \begin{bmatrix} \frac{e^{3t}}{2} \\ e^{3t} \end{bmatrix}$$

Which becomes

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} -\frac{c_1 e^{-t}}{2} + \frac{c_2 e^{3t}}{2} \\ c_1 e^{-t} + c_2 e^{3t} \end{bmatrix}$$

The following is the phase plot of the system.




Figure 503: Phase plot

29.2.3 Maple step by step solution

Let's solve

$$[x'(t) = x(t) + y(t), y'(t) = 4x(t) + y(t)]$$

- Define vector

$$\vec{x}(t) = \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

- Convert system into a vector equation

$$\vec{x}'(t) = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

- System to solve

$$\vec{x}'(t) = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix} \cdot \vec{x}(t)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 1 & 1 \\ 4 & 1 \end{bmatrix}$$

- Rewrite the system as

$$\vec{x}'(t) = A \cdot \vec{x}(t)$$

- To solve the system, find the eigenvalues and eigenvectors of A
- Eigenpairs of A

$$\left[\left[-1, \begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix} \right], \left[3, \begin{bmatrix} \frac{1}{2} \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[-1, \begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{x}_1 = e^{-t} \cdot \begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[3, \begin{bmatrix} \frac{1}{2} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{x}_2 = e^{3t} \cdot \begin{bmatrix} \frac{1}{2} \\ 1 \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{x} = c_1 \vec{x}_1 + c_2 \vec{x}_2$$

- Substitute solutions into the general solution

$$\vec{x} = e^{-t} c_1 \cdot \begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix} + e^{3t} c_2 \cdot \begin{bmatrix} \frac{1}{2} \\ 1 \end{bmatrix}$$

- Substitute in vector of dependent variables

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} -\frac{e^{-t}c_1}{2} + \frac{e^{3t}c_2}{2} \\ e^{-t}c_1 + e^{3t}c_2 \end{bmatrix}$$

- Solution to the system of ODEs

$$\left\{ x(t) = -\frac{e^{-t}c_1}{2} + \frac{e^{3t}c_2}{2}, y(t) = e^{-t}c_1 + e^{3t}c_2 \right\}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 36

```
dsolve([diff(x(t),t)=x(t)+y(t),diff(y(t),t)=4*x(t)+y(t)],singsol=all)
```

$$\begin{aligned} x(t) &= c_1e^{3t} + c_2e^{-t} \\ y(t) &= 2c_1e^{3t} - 2c_2e^{-t} \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 70

```
DSolve[{x'[t]==x[t]+y[t],y'[t]==4*x[t]+y[t]},{x[t],y[t]},t,IncludeSingularSolutions -> True]
```

$$\begin{aligned} x(t) &\rightarrow \frac{1}{4}e^{-t}(2c_1(e^{4t} + 1) + c_2(e^{4t} - 1)) \\ y(t) &\rightarrow \frac{1}{2}e^{-t}(2c_1(e^{4t} - 1) + c_2(e^{4t} + 1)) \end{aligned}$$

29.3 problem 2(c)

29.3.1 Solution using Matrix exponential method 4105

29.3.2 Solution using explicit Eigenvalue and Eigenvector method . . . 4106

Internal problem ID [6531]

Internal file name [OUTPUT/5779_Sunday_June_05_2022_03_54_02_PM_64605471/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 10. Systems of First-Order Equations. Section A. Drill exercises. Page 400

Problem number: 2(c).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**system of linear ODEs**"

Solve

$$x'(t) = -3x(t) + \sqrt{2}y(t)$$

$$y'(t) = \sqrt{2}x(t) - 2y(t)$$

29.3.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} -3 & \sqrt{2} \\ \sqrt{2} & -2 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} \frac{2e^{-4t}}{3} + \frac{e^{-t}}{3} & -\frac{\sqrt{2}(-e^{-t}+e^{-4t})}{3} \\ -\frac{\sqrt{2}(-e^{-t}+e^{-4t})}{3} & \frac{e^{-4t}}{3} + \frac{2e^{-t}}{3} \end{bmatrix}$$

Therefore the homogeneous solution is

$$\begin{aligned}\vec{x}_h(t) &= e^{At}\vec{c} \\ &= \begin{bmatrix} \frac{2e^{-4t}}{3} + \frac{e^{-t}}{3} & -\frac{\sqrt{2}(-e^{-t}+e^{-4t})}{3} \\ -\frac{\sqrt{2}(-e^{-t}+e^{-4t})}{3} & \frac{e^{-4t}}{3} + \frac{2e^{-t}}{3} \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} \\ &= \begin{bmatrix} \left(\frac{2e^{-4t}}{3} + \frac{e^{-t}}{3}\right)c_1 - \frac{\sqrt{2}(-e^{-t}+e^{-4t})c_2}{3} \\ -\frac{\sqrt{2}(-e^{-t}+e^{-4t})c_1}{3} + \left(\frac{e^{-4t}}{3} + \frac{2e^{-t}}{3}\right)c_2 \end{bmatrix}\end{aligned}$$

Since no forcing function is given, then the final solution is $\vec{x}_h(t)$ above.

29.3.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} -3 & \sqrt{2} \\ \sqrt{2} & -2 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det\left(\begin{bmatrix} -3 & \sqrt{2} \\ \sqrt{2} & -2 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}\right) = 0$$

Therefore

$$\det\left(\begin{bmatrix} -3 - \lambda & \sqrt{2} \\ \sqrt{2} & -2 - \lambda \end{bmatrix}\right) = 0$$

Which gives the characteristic equation

$$\lambda^2 + 5\lambda + 4 = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = -1$$

$$\lambda_2 = -4$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
-1	1	real eigenvalue
-4	1	real eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = -4$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} -3 & \sqrt{2} \\ \sqrt{2} & -2 \end{bmatrix} - (-4) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & \sqrt{2} \\ \sqrt{2} & 2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} 1 & \sqrt{2} & 0 \\ \sqrt{2} & 2 & 0 \end{array} \right]$$

$$R_2 = R_2 - \sqrt{2}R_1 \implies \left[\begin{array}{cc|c} 1 & \sqrt{2} & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} 1 & \sqrt{2} \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = -\sqrt{2}t\}$

Hence the solution is

$$\begin{bmatrix} -\sqrt{2}t \\ t \end{bmatrix} = \begin{bmatrix} -\sqrt{2}t \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} -\sqrt{2}t \\ t \end{bmatrix} = t \begin{bmatrix} -\sqrt{2} \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} -\sqrt{2} \\ 1 \end{bmatrix}$$

Considering the eigenvalue $\lambda_2 = -1$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} -3 & \sqrt{2} \\ \sqrt{2} & -2 \end{bmatrix} - (-1) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} -2 & \sqrt{2} \\ \sqrt{2} & -1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} -2 & \sqrt{2} & 0 \\ \sqrt{2} & -1 & 0 \end{array} \right]$$

$$R_2 = R_2 + \frac{\sqrt{2}R_1}{2} \implies \left[\begin{array}{cc|c} -2 & \sqrt{2} & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} -2 & \sqrt{2} \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\left\{ v_1 = \frac{\sqrt{2}t}{2} \right\}$

Hence the solution is

$$\begin{bmatrix} \frac{\sqrt{2}t}{2} \\ t \end{bmatrix} = \begin{bmatrix} \frac{\sqrt{2}t}{2} \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} \frac{\sqrt{2}t}{2} \\ t \end{bmatrix} = t \begin{bmatrix} \frac{\sqrt{2}}{2} \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} \frac{\sqrt{2}t}{2} \\ t \end{bmatrix} = \begin{bmatrix} \frac{\sqrt{2}}{2} \\ 1 \end{bmatrix}$$

Which is normalized to

$$\begin{bmatrix} \frac{\sqrt{2}t}{2} \\ t \end{bmatrix} = \begin{bmatrix} \sqrt{2} \\ 2 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
-1	1	1	No	$\begin{bmatrix} \frac{\sqrt{2}}{2} \\ 1 \end{bmatrix}$
-4	1	1	No	$\begin{bmatrix} -\sqrt{2} \\ 1 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. Since eigenvalue -1 is real and distinct then the

corresponding eigenvector solution is

$$\begin{aligned}\vec{x}_1(t) &= \vec{v}_1 e^{-t} \\ &= \begin{bmatrix} \frac{\sqrt{2}}{2} \\ 1 \end{bmatrix} e^{-t}\end{aligned}$$

Since eigenvalue -4 is real and distinct then the corresponding eigenvector solution is

$$\begin{aligned}\vec{x}_2(t) &= \vec{v}_2 e^{-4t} \\ &= \begin{bmatrix} -\sqrt{2} \\ 1 \end{bmatrix} e^{-4t}\end{aligned}$$

Therefore the final solution is

$$\vec{x}_h(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

Which is written as

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = c_1 \begin{bmatrix} \frac{\sqrt{2}e^{-t}}{2} \\ e^{-t} \end{bmatrix} + c_2 \begin{bmatrix} -\sqrt{2}e^{-4t} \\ e^{-4t} \end{bmatrix}$$

Which becomes

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} \frac{\sqrt{2}e^{-t}(-2c_2e^{-3t}+c_1)}{2} \\ c_1e^{-t} + c_2e^{-4t} \end{bmatrix}$$

The following is the phase plot of the system.




Figure 504: Phase plot

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 41

```
dsolve([diff(x(t),t)=-3*x(t)+sqrt(2)*y(t),diff(y(t),t)=sqrt(2)*x(t)-2*y(t)],singsol=all)
```

$$x(t) = e^{-4t}c_1 + c_2e^{-t}$$

$$y(t) = \frac{(-e^{-4t}c_1 + 2c_2e^{-t})\sqrt{2}}{2}$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 80

```
DSolve[{x'[t]==-3*x[t]+Sqrt[2]*y[t],y'[t]==Sqrt[2]*x[t]-2*y[t]},{x[t],y[t]},t,IncludeSingularSolutions->True]
```

$$x(t) \rightarrow \frac{1}{3}e^{-4t} \left(c_1(e^{3t} + 2) + \sqrt{2}c_2(e^{3t} - 1) \right)$$

$$y(t) \rightarrow \frac{1}{3}e^{-4t} \left(\sqrt{2}c_1(e^{3t} - 1) + c_2(2e^{3t} + 1) \right)$$

29.4 problem 2(d)

- 29.4.1 Solution using Matrix exponential method 4112
- 29.4.2 Solution using explicit Eigenvalue and Eigenvector method . . . 4113
- 29.4.3 Maple step by step solution 4118

Internal problem ID [6532]

Internal file name [OUTPUT/5780_Sunday_June_05_2022_03_54_04_PM_36029301/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 10. Systems of First-Order Equations. Section A. Drill exercises. Page 400

Problem number: 2(d).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**system of linear ODEs**"

Solve

$$\begin{aligned}x'(t) &= 5x(t) + 3y(t) \\y'(t) &= -6x(t) - 4y(t)\end{aligned}$$

29.4.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A \vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 5 & 3 \\ -6 & -4 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} -e^{-t} + 2e^{2t} & e^{2t} - e^{-t} \\ -2e^{2t} + 2e^{-t} & 2e^{-t} - e^{2t} \end{bmatrix}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 \vec{x}_h(t) &= e^{At} \vec{c} \\
 &= \begin{bmatrix} -e^{-t} + 2e^{2t} & e^{2t} - e^{-t} \\ -2e^{2t} + 2e^{-t} & 2e^{-t} - e^{2t} \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} \\
 &= \begin{bmatrix} (-e^{-t} + 2e^{2t})c_1 + (e^{2t} - e^{-t})c_2 \\ (-2e^{2t} + 2e^{-t})c_1 + (2e^{-t} - e^{2t})c_2 \end{bmatrix} \\
 &= \begin{bmatrix} (-c_1 - c_2)e^{-t} + 2(c_1 + \frac{c_2}{2})e^{2t} \\ (2c_1 + 2c_2)e^{-t} - 2(c_1 + \frac{c_2}{2})e^{2t} \end{bmatrix}
 \end{aligned}$$

Since no forcing function is given, then the final solution is $\vec{x}_h(t)$ above.

29.4.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 5 & 3 \\ -6 & -4 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det\left(\begin{bmatrix} 5 & 3 \\ -6 & -4 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}\right) = 0$$

Therefore

$$\det\left(\begin{bmatrix} 5 - \lambda & 3 \\ -6 & -4 - \lambda \end{bmatrix}\right) = 0$$

Which gives the characteristic equation

$$\lambda^2 - \lambda - 2 = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = 2$$

$$\lambda_2 = -1$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
-1	1	real eigenvalue
2	1	real eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = -1$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 5 & 3 \\ -6 & -4 \end{bmatrix} - (-1) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 6 & 3 \\ -6 & -3 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} 6 & 3 & 0 \\ -6 & -3 & 0 \end{array} \right]$$

$$R_2 = R_2 + R_1 \implies \left[\begin{array}{cc|c} 6 & 3 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} 6 & 3 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = -\frac{t}{2}\}$

Hence the solution is

$$\begin{bmatrix} -\frac{t}{2} \\ t \end{bmatrix} = \begin{bmatrix} -\frac{t}{2} \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} -\frac{t}{2} \\ t \end{bmatrix} = t \begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix} = \begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix}$$

Which is normalized to

$$\begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix} = \begin{bmatrix} -1 \\ 2 \end{bmatrix}$$

Considering the eigenvalue $\lambda_2 = 2$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 5 & 3 \\ -6 & -4 \end{bmatrix} - (2) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 3 & 3 \\ -6 & -6 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} 3 & 3 & 0 \\ -6 & -6 & 0 \end{array} \right]$$

$$R_2 = R_2 + 2R_1 \implies \left[\begin{array}{cc|c} 3 & 3 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} 3 & 3 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = -t\}$

Hence the solution is

$$\begin{bmatrix} -t \\ t \end{bmatrix} = \begin{bmatrix} -t \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} -t \\ t \end{bmatrix} = t \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} -t \\ t \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
2	1	1	No	$\begin{bmatrix} -1 \\ 1 \end{bmatrix}$
-1	1	1	No	$\begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. Since eigenvalue 2 is real and distinct then the corresponding eigenvector solution is

$$\begin{aligned}\vec{x}_1(t) &= \vec{v}_1 e^{2t} \\ &= \begin{bmatrix} -1 \\ 1 \end{bmatrix} e^{2t}\end{aligned}$$

Since eigenvalue -1 is real and distinct then the corresponding eigenvector solution is

$$\begin{aligned}\vec{x}_2(t) &= \vec{v}_2 e^{-t} \\ &= \begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix} e^{-t}\end{aligned}$$

Therefore the final solution is

$$\vec{x}_h(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

Which is written as

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = c_1 \begin{bmatrix} -e^{2t} \\ e^{2t} \end{bmatrix} + c_2 \begin{bmatrix} -\frac{e^{-t}}{2} \\ e^{-t} \end{bmatrix}$$

Which becomes

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} -c_1 e^{2t} - \frac{c_2 e^{-t}}{2} \\ c_1 e^{2t} + c_2 e^{-t} \end{bmatrix}$$

The following is the phase plot of the system.




Figure 505: Phase plot

29.4.3 Maple step by step solution

Let's solve

$$[x'(t) = 5x(t) + 3y(t), y'(t) = -6x(t) - 4y(t)]$$

- Define vector

$$\vec{x}(t) = \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

- Convert system into a vector equation

$$\vec{x}'(t) = \begin{bmatrix} 5 & 3 \\ -6 & -4 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

- System to solve

$$\vec{x}'(t) = \begin{bmatrix} 5 & 3 \\ -6 & -4 \end{bmatrix} \cdot \vec{x}(t)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 5 & 3 \\ -6 & -4 \end{bmatrix}$$

- Rewrite the system as

$$\vec{x}'(t) = A \cdot \vec{x}(t)$$

- To solve the system, find the eigenvalues and eigenvectors of A
- Eigenpairs of A

$$\left[\left[-1, \begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix} \right], \left[2, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[-1, \begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{x}_1 = e^{-t} \cdot \begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[2, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{x}_2 = e^{2t} \cdot \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{x} = c_1 \vec{x}_1 + c_2 \vec{x}_2$$

- Substitute solutions into the general solution

$$\vec{x} = e^{-t} c_1 \cdot \begin{bmatrix} -\frac{1}{2} \\ 1 \end{bmatrix} + c_2 e^{2t} \cdot \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

- Substitute in vector of dependent variables

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} -\frac{e^{-t}c_1}{2} - c_2e^{2t} \\ e^{-t}c_1 + c_2e^{2t} \end{bmatrix}$$

- Solution to the system of ODEs

$$\left\{ x(t) = -\frac{e^{-t}c_1}{2} - c_2e^{2t}, y(t) = e^{-t}c_1 + c_2e^{2t} \right\}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 36

```
dsolve([diff(x(t),t)=5*x(t)+3*y(t),diff(y(t),t)=-6*x(t)-4*y(t)],singsol=all)
```

$$\begin{aligned} x(t) &= e^{-t}c_1 + c_2e^{2t} \\ y(t) &= -2e^{-t}c_1 - c_2e^{2t} \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 66

```
DSolve[{x'[t]==5*x[t]+3*y[t],y'[t]==-6*x[t]-4*y[t]},{x[t],y[t]},t,IncludeSingularSolutions -
```

$$\begin{aligned} x(t) &\rightarrow e^{-t}(c_1(2e^{3t} - 1) + c_2(e^{3t} - 1)) \\ y(t) &\rightarrow e^{-t}(-2c_1(e^{3t} - 1) - c_2(e^{3t} - 2)) \end{aligned}$$

29.5 problem 3(a)

29.5.1 Solution using Matrix exponential method	4121
29.5.2 Solution using explicit Eigenvalue and Eigenvector method . . .	4122
29.5.3 Maple step by step solution	4127

Internal problem ID [6533]

Internal file name [OUTPUT/5781_Sunday_June_05_2022_03_54_06_PM_72724581/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 10. Systems of First-Order Equations. Section A. Drill exercises. Page 400

Problem number: 3(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "system of linear ODEs"

Solve

$$\begin{aligned}x'(t) &= 3x(t) + 2y(t) \\y'(t) &= -2x(t) - y(t)\end{aligned}$$

29.5.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A \vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 3 & 2 \\ -2 & -1 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} e^t(2t + 1) & 2t e^t \\ -2t e^t & e^t(1 - 2t) \end{bmatrix}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 \vec{x}_h(t) &= e^{At} \vec{c} \\
 &= \begin{bmatrix} e^t(2t+1) & 2t e^t \\ -2t e^t & e^t(1-2t) \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} \\
 &= \begin{bmatrix} e^t(2t+1)c_1 + 2t e^t c_2 \\ -2t e^t c_1 + e^t(1-2t)c_2 \end{bmatrix} \\
 &= \begin{bmatrix} e^t(2tc_1 + 2c_2t + c_1) \\ (c_2(1-2t) - 2tc_1) e^t \end{bmatrix}
 \end{aligned}$$

Since no forcing function is given, then the final solution is $\vec{x}_h(t)$ above.

29.5.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A \vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 3 & 2 \\ -2 & -1 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det \left(\begin{bmatrix} 3 & 2 \\ -2 & -1 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) = 0$$

Therefore

$$\det \left(\begin{bmatrix} 3 - \lambda & 2 \\ -2 & -1 - \lambda \end{bmatrix} \right) = 0$$

Which gives the characteristic equation

$$\lambda^2 - 2\lambda + 1 = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = 1$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
1	1	real eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = 1$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 3 & 2 \\ -2 & -1 \end{bmatrix} - (1) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$
$$\begin{bmatrix} 2 & 2 \\ -2 & -2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} 2 & 2 & 0 \\ -2 & -2 & 0 \end{array} \right]$$

$$R_2 = R_2 + R_1 \implies \left[\begin{array}{cc|c} 2 & 2 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} 2 & 2 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = -t\}$

Hence the solution is

$$\begin{bmatrix} -t \\ t \end{bmatrix} = \begin{bmatrix} -t \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} -t \\ t \end{bmatrix} = t \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
1	2	1	Yes	$\begin{bmatrix} -1 \\ 1 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. eigenvalue 1 is real and repeated eigenvalue of multiplicity 2. There are two possible cases that can happen. This is illustrated in this diagram




Figure 506: Possible case for repeated λ of multiplicity 2

This eigenvalue has algebraic multiplicity of 2, and geometric multiplicity 1, therefore this is defective eigenvalue. The defect is 1. This falls into case 2 shown above. We need to generate the missing additional generalized eigenvector \vec{v}_2 by solving

$$(A - \lambda I) \vec{v}_2 = \vec{v}_1$$

Where \vec{v}_1 is the normal (rank 1) eigenvector found above. Hence we need to solve

$$\left(\begin{bmatrix} 3 & 2 \\ -2 & -1 \end{bmatrix} - (1) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 2 \\ -2 & -2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

Solving for \vec{v}_2 gives

$$\vec{v}_2 = \begin{bmatrix} -\frac{3}{2} \\ 1 \end{bmatrix}$$

We have found two generalized eigenvectors for eigenvalue 1. Therefore the two basis solution associated with this eigenvalue are

$$\begin{aligned}\vec{x}_1(t) &= \vec{v}_1 e^{\lambda t} \\ &= \begin{bmatrix} -1 \\ 1 \end{bmatrix} e^t \\ &= \begin{bmatrix} -e^t \\ e^t \end{bmatrix}\end{aligned}$$

And

$$\begin{aligned}\vec{x}_2(t) &= (\vec{v}_1 t + \vec{v}_2) e^{\lambda t} \\ &= \left(\begin{bmatrix} -1 \\ 1 \end{bmatrix} t + \begin{bmatrix} -\frac{3}{2} \\ 1 \end{bmatrix} \right) e^t \\ &= \begin{bmatrix} -\frac{e^t(2t+3)}{2} \\ e^t(t+1) \end{bmatrix}\end{aligned}$$

Therefore the final solution is

$$\vec{x}_h(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

Which is written as

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = c_1 \begin{bmatrix} -e^t \\ e^t \end{bmatrix} + c_2 \begin{bmatrix} e^t(-t - \frac{3}{2}) \\ e^t(t+1) \end{bmatrix}$$

Which becomes

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} e^t(-c_1 - c_2 t - \frac{3}{2}c_2) \\ e^t(c_2 t + c_1 + c_2) \end{bmatrix}$$

The following is the phase plot of the system.




Figure 507: Phase plot

29.5.3 Maple step by step solution

Let's solve

$$[x'(t) = 3x(t) + 2y(t), y'(t) = -2x(t) - y(t)]$$

- Define vector

$$\vec{x}(t) = \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

- Convert system into a vector equation

$$\vec{x}'(t) = \begin{bmatrix} 3 & 2 \\ -2 & -1 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

- System to solve

$$\vec{x}'(t) = \begin{bmatrix} 3 & 2 \\ -2 & -1 \end{bmatrix} \cdot \vec{x}(t)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 3 & 2 \\ -2 & -1 \end{bmatrix}$$

- Rewrite the system as

$$\vec{x}'(t) = A \cdot \vec{x}(t)$$

- To solve the system, find the eigenvalues and eigenvectors of A
- Eigenpairs of A

$$\left[\left[1, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right], \left[1, \begin{bmatrix} 0 \\ 0 \end{bmatrix} \right] \right]$$

- Consider eigenpair, with eigenvalue of algebraic multiplicity 2

$$\left[1, \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right]$$

- First solution from eigenvalue 1

$$\vec{x}_1(t) = e^t \cdot \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

- Form of the 2nd homogeneous solution where \vec{p} is to be solved for, $\lambda = 1$ is the eigenvalue, and

$$\vec{x}_2(t) = e^{\lambda t} (t\vec{v} + \vec{p})$$

- Note that the t multiplying \vec{v} makes this solution linearly independent to the 1st solution obtained

- Substitute $\vec{x}_2(t)$ into the homogeneous system

$$\lambda e^{\lambda t} (t\vec{v} + \vec{p}) + e^{\lambda t} \vec{v} = (e^{\lambda t} A) \cdot (t\vec{v} + \vec{p})$$

- Use the fact that \vec{v} is an eigenvector of A

$$\lambda e^{\lambda t} (t\vec{v} + \vec{p}) + e^{\lambda t} \vec{v} = e^{\lambda t} (\lambda t\vec{v} + A \cdot \vec{p})$$

- Simplify equation

$$\lambda \vec{p} + \vec{v} = A \cdot \vec{p}$$

- Make use of the identity matrix I

$$(\lambda \cdot I) \cdot \vec{p} + \vec{v} = A \cdot \vec{p}$$

- Condition \vec{p} must meet for $\vec{x}_2(t)$ to be a solution to the homogeneous system

$$(A - \lambda \cdot I) \cdot \vec{p} = \vec{v}$$

- Choose \vec{p} to use in the second solution to the homogeneous system from eigenvalue 1

$$\left(\begin{bmatrix} 3 & 2 \\ -2 & -1 \end{bmatrix} - 1 \cdot \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \cdot \vec{p} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

- Choice of \vec{p}

$$\vec{p} = \begin{bmatrix} -\frac{1}{2} \\ 0 \end{bmatrix}$$

- Second solution from eigenvalue 1

$$\vec{x}_2(t) = e^t \cdot \left(t \cdot \begin{bmatrix} -1 \\ 1 \end{bmatrix} + \begin{bmatrix} -\frac{1}{2} \\ 0 \end{bmatrix} \right)$$

- General solution to the system of ODEs

$$\vec{x} = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

- Substitute solutions into the general solution

$$\vec{x} = c_1 e^t \cdot \begin{bmatrix} -1 \\ 1 \end{bmatrix} + c_2 e^t \cdot \left(t \cdot \begin{bmatrix} -1 \\ 1 \end{bmatrix} + \begin{bmatrix} -\frac{1}{2} \\ 0 \end{bmatrix} \right)$$

- Substitute in vector of dependent variables

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} e^t(-c_1 - tc_2 - \frac{1}{2}c_2) \\ e^t(tc_2 + c_1) \end{bmatrix}$$

- Solution to the system of ODEs

$$\{x(t) = e^t(-c_1 - tc_2 - \frac{1}{2}c_2), y(t) = e^t(tc_2 + c_1)\}$$

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 31

```
dsolve([diff(x(t),t)=3*x(t)+2*y(t),diff(y(t),t)=-2*x(t)-y(t)],singsol=all)
```

$$x(t) = e^t(c_2 t + c_1)$$

$$y(t) = -\frac{e^t(2c_2 t + 2c_1 - c_2)}{2}$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 40

```
DSolve[{x'[t]==3*x[t]+2*y[t],y'[t]==-2*x[t]-y[t]},{x[t],y[t]},t,IncludeSingularSolutions ->
```

$$x(t) \rightarrow e^t(2c_1t + 2c_2t + c_1)$$

$$y(t) \rightarrow e^t(c_2 - 2(c_1 + c_2)t)$$

29.6 problem 3(b)

- 29.6.1 Solution using Matrix exponential method 4131
- 29.6.2 Solution using explicit Eigenvalue and Eigenvector method . . . 4132
- 29.6.3 Maple step by step solution 4136

Internal problem ID [6534]

Internal file name [OUTPUT/5782_Sunday_June_05_2022_03_54_07_PM_48022315/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 10. Systems of First-Order Equations. Section A. Drill exercises. Page 400

Problem number: 3(b).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**system of linear ODEs**"

Solve

$$\begin{aligned}x'(t) &= x(t) + y(t) \\y'(t) &= -x(t) + y(t)\end{aligned}$$

29.6.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A \vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} e^t \cos(t) & \sin(t) e^t \\ -\sin(t) e^t & e^t \cos(t) \end{bmatrix}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 \vec{x}_h(t) &= e^{At} \vec{c} \\
 &= \begin{bmatrix} e^t \cos(t) & \sin(t) e^t \\ -\sin(t) e^t & e^t \cos(t) \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} \\
 &= \begin{bmatrix} e^t \cos(t) c_1 + \sin(t) e^t c_2 \\ -\sin(t) e^t c_1 + e^t \cos(t) c_2 \end{bmatrix} \\
 &= \begin{bmatrix} e^t (\cos(t) c_1 + \sin(t) c_2) \\ e^t (-\sin(t) c_1 + \cos(t) c_2) \end{bmatrix}
 \end{aligned}$$

Since no forcing function is given, then the final solution is $\vec{x}_h(t)$ above.

29.6.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A \vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det \left(\begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) = 0$$

Therefore

$$\det \left(\begin{bmatrix} 1 - \lambda & 1 \\ -1 & 1 - \lambda \end{bmatrix} \right) = 0$$

Which gives the characteristic equation

$$\lambda^2 - 2\lambda + 2 = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = 1 + i$$

$$\lambda_2 = 1 - i$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
$1 + i$	1	complex eigenvalue
$1 - i$	1	complex eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = 1 - i$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix} - (1 - i) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} i & 1 \\ -1 & i \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} i & 1 & 0 \\ -1 & i & 0 \end{array} \right]$$

$$R_2 = -iR_1 + R_2 \implies \left[\begin{array}{cc|c} i & 1 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} i & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = it\}$

Hence the solution is

$$\begin{bmatrix} It \\ t \end{bmatrix} = \begin{bmatrix} it \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} It \\ t \end{bmatrix} = t \begin{bmatrix} i \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} It \\ t \end{bmatrix} = \begin{bmatrix} i \\ 1 \end{bmatrix}$$

Considering the eigenvalue $\lambda_2 = 1 + i$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix} - (1+i) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} -i & 1 \\ -1 & -i \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} -i & 1 & 0 \\ -1 & -i & 0 \end{array} \right]$$

$$R_2 = iR_1 + R_2 \implies \left[\begin{array}{cc|c} -i & 1 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} -i & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = -it\}$

Hence the solution is

$$\begin{bmatrix} -It \\ t \end{bmatrix} = \begin{bmatrix} -it \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} -It \\ t \end{bmatrix} = t \begin{bmatrix} -i \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} -It \\ t \end{bmatrix} = \begin{bmatrix} -i \\ 1 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
$1 + i$	1	1	No	$\begin{bmatrix} -i \\ 1 \end{bmatrix}$
$1 - i$	1	1	No	$\begin{bmatrix} i \\ 1 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. Therefore the final solution is

$$\vec{x}_h(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

Which is written as

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = c_1 \begin{bmatrix} -ie^{(1+i)t} \\ e^{(1+i)t} \end{bmatrix} + c_2 \begin{bmatrix} ie^{(1-i)t} \\ e^{(1-i)t} \end{bmatrix}$$

Which becomes

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} i(c_2e^{(1-i)t} - c_1e^{(1+i)t}) \\ c_1e^{(1+i)t} + c_2e^{(1-i)t} \end{bmatrix}$$

The following is the phase plot of the system.




Figure 508: Phase plot

29.6.3 Maple step by step solution

Let's solve

$$[x'(t) = x(t) + y(t), y'(t) = -x(t) + y(t)]$$

- Define vector

$$\vec{x}(t) = \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

- Convert system into a vector equation

$$\vec{x}'(t) = \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

- System to solve

$$\vec{x}'(t) = \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix} \cdot \vec{x}(t)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$$

- Rewrite the system as

$$\vec{x}'(t) = A \cdot \vec{x}(t)$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[1 - I, \begin{bmatrix} I \\ 1 \end{bmatrix} \right], \left[1 + I, \begin{bmatrix} -I \\ 1 \end{bmatrix} \right] \right]$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$\left[1 - I, \begin{bmatrix} I \\ 1 \end{bmatrix} \right]$$

- Solution from eigenpair

$$e^{(1-I)t} \cdot \begin{bmatrix} I \\ 1 \end{bmatrix}$$

- Use Euler identity to write solution in terms of sin and cos

$$e^t \cdot (\cos(t) - I \sin(t)) \cdot \begin{bmatrix} I \\ 1 \end{bmatrix}$$

- Simplify expression

$$e^t \cdot \begin{bmatrix} I(\cos(t) - I \sin(t)) \\ \cos(t) - I \sin(t) \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\begin{bmatrix} \vec{x}_1(t) = e^t \cdot \begin{bmatrix} \sin(t) \\ \cos(t) \end{bmatrix}, \vec{x}_2(t) = e^t \cdot \begin{bmatrix} \cos(t) \\ -\sin(t) \end{bmatrix} \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{x} = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

- Substitute solutions into the general solution

$$\vec{x} = c_1 e^t \cdot \begin{bmatrix} \sin(t) \\ \cos(t) \end{bmatrix} + c_2 e^t \cdot \begin{bmatrix} \cos(t) \\ -\sin(t) \end{bmatrix}$$

- Substitute in vector of dependent variables

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} e^t(c_1 \sin(t) + c_2 \cos(t)) \\ e^t(c_1 \cos(t) - c_2 \sin(t)) \end{bmatrix}$$

- Solution to the system of ODEs

$$\{x(t) = e^t(c_1 \sin(t) + c_2 \cos(t)), y(t) = e^t(c_1 \cos(t) - c_2 \sin(t))\}$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 33

```
dsolve([diff(x(t),t)=x(t)+y(t),diff(y(t),t)=-x(t)+y(t)],singsol=all)
```

$$x(t) = e^t(c_1 \sin(t) + c_2 \cos(t))$$

$$y(t) = e^t(c_1 \cos(t) - c_2 \sin(t))$$

✓ Solution by Mathematica

Time used: 0.002 (sec). Leaf size: 39

```
DSolve[{x'[t]==x[t]+y[t],y'[t]==-x[t]+y[t]},{x[t],y[t]},t,IncludeSingularSolutions->True]
```

$$x(t) \rightarrow e^t(c_1 \cos(t) + c_2 \sin(t))$$

$$y(t) \rightarrow e^t(c_2 \cos(t) - c_1 \sin(t))$$

29.7 problem 3(c)

29.7.1 Solution using Matrix exponential method	4139
29.7.2 Solution using explicit Eigenvalue and Eigenvector method . . .	4140
29.7.3 Maple step by step solution	4145

Internal problem ID [6535]

Internal file name [OUTPUT/5783_Sunday_June_05_2022_03_54_10_PM_52946076/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 10. Systems of First-Order Equations. Section A. Drill exercises. Page 400

Problem number: 3(c).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "system of linear ODEs"

Solve

$$\begin{aligned}x'(t) &= 3x(t) - 5y(t) \\y'(t) &= -x(t) + 2y(t)\end{aligned}$$

29.7.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A \vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 3 & -5 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} \frac{(-\sqrt{21}+21)e^{-\frac{(-5+\sqrt{21})t}{2}}}{42} + \frac{e^{\frac{(5+\sqrt{21})t}{2}}(\sqrt{21}+21)}{42} & \frac{5\left(-e^{\frac{(5+\sqrt{21})t}{2}} + e^{-\frac{(-5+\sqrt{21})t}{2}}\right)\sqrt{21}}{21} \\ \frac{\left(-e^{\frac{(5+\sqrt{21})t}{2}} + e^{-\frac{(-5+\sqrt{21})t}{2}}\right)\sqrt{21}}{21} & \frac{(\sqrt{21}+21)e^{-\frac{(-5+\sqrt{21})t}{2}}}{42} - \frac{e^{\frac{(5+\sqrt{21})t}{2}}(\sqrt{21}-21)}{42} \end{bmatrix}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 \vec{x}_h(t) &= e^{At} \vec{c} \\
 &= \begin{bmatrix} \frac{(-\sqrt{21}+21)e^{-\frac{(-5+\sqrt{21})t}{2}} + \frac{e^{\frac{(5+\sqrt{21})t}{2}}(\sqrt{21}+21)}{42}}{42} & \frac{5\left(-e^{-\frac{(5+\sqrt{21})t}{2}} + e^{-\frac{(-5+\sqrt{21})t}{2}}\right)\sqrt{21}}{21} \\ \frac{\left(-e^{-\frac{(5+\sqrt{21})t}{2}} + e^{-\frac{(-5+\sqrt{21})t}{2}}\right)\sqrt{21}}{21} & \frac{(\sqrt{21}+21)e^{-\frac{(-5+\sqrt{21})t}{2}}}{42} - \frac{e^{\frac{(5+\sqrt{21})t}{2}}(\sqrt{21}-21)}{42} \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} \\
 &= \begin{bmatrix} \left(\frac{(-\sqrt{21}+21)e^{-\frac{(-5+\sqrt{21})t}{2}} + \frac{e^{\frac{(5+\sqrt{21})t}{2}}(\sqrt{21}+21)}{42}}{42}\right) c_1 + \frac{5\left(-e^{-\frac{(5+\sqrt{21})t}{2}} + e^{-\frac{(-5+\sqrt{21})t}{2}}\right)\sqrt{21} c_2}{21} \\ \frac{\left(-e^{-\frac{(5+\sqrt{21})t}{2}} + e^{-\frac{(-5+\sqrt{21})t}{2}}\right)\sqrt{21} c_1}{21} + \left(\frac{(\sqrt{21}+21)e^{-\frac{(-5+\sqrt{21})t}{2}}}{42} - \frac{e^{\frac{(5+\sqrt{21})t}{2}}(\sqrt{21}-21)}{42}\right) c_2 \end{bmatrix} \\
 &= \begin{bmatrix} \frac{((-c_1+10c_2)\sqrt{21}+21c_1)e^{-\frac{(-5+\sqrt{21})t}{2}}}{42} + \frac{((c_1-10c_2)\sqrt{21}+21c_1)e^{\frac{(5+\sqrt{21})t}{2}}}{42} \\ \frac{((2c_1+c_2)\sqrt{21}+21c_2)e^{-\frac{(-5+\sqrt{21})t}{2}}}{42} - \frac{((c_1+\frac{c_2}{2})\sqrt{21}-\frac{21c_2}{2})e^{\frac{(5+\sqrt{21})t}{2}}}{21} \end{bmatrix}
 \end{aligned}$$

Since no forcing function is given, then the final solution is $\vec{x}_h(t)$ above.

29.7.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 3 & -5 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det \left(\begin{bmatrix} 3 & -5 \\ -1 & 2 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) = 0$$

Therefore

$$\det \left(\begin{bmatrix} 3 - \lambda & -5 \\ -1 & 2 - \lambda \end{bmatrix} \right) = 0$$

Which gives the characteristic equation

$$\lambda^2 - 5\lambda + 1 = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = \frac{5}{2} + \frac{\sqrt{21}}{2}$$

$$\lambda_2 = \frac{5}{2} - \frac{\sqrt{21}}{2}$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
$\frac{5}{2} + \frac{\sqrt{21}}{2}$	1	real eigenvalue
$\frac{5}{2} - \frac{\sqrt{21}}{2}$	1	real eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = \frac{5}{2} - \frac{\sqrt{21}}{2}$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 3 & -5 \\ -1 & 2 \end{bmatrix} - \left(\frac{5}{2} - \frac{\sqrt{21}}{2} \right) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} \frac{1}{2} + \frac{\sqrt{21}}{2} & -5 \\ -1 & -\frac{1}{2} + \frac{\sqrt{21}}{2} \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} \frac{1}{2} + \frac{\sqrt{21}}{2} & -5 & 0 \\ -1 & -\frac{1}{2} + \frac{\sqrt{21}}{2} & 0 \end{array} \right]$$

$$R_2 = R_2 + \frac{R_1}{\frac{1}{2} + \frac{\sqrt{21}}{2}} \implies \left[\begin{array}{cc|c} \frac{1}{2} + \frac{\sqrt{21}}{2} & -5 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\left[\begin{array}{cc|c} \frac{1}{2} + \frac{\sqrt{21}}{2} & -5 & 0 \\ 0 & 0 & 0 \end{array} \right] \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\left\{ v_1 = \frac{10t}{1+\sqrt{21}} \right\}$

Hence the solution is

$$\begin{bmatrix} \frac{10t}{1+\sqrt{21}} \\ t \end{bmatrix} = \begin{bmatrix} \frac{10t}{1+\sqrt{21}} \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} \frac{10t}{1+\sqrt{21}} \\ t \end{bmatrix} = t \begin{bmatrix} \frac{10}{1+\sqrt{21}} \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} \frac{10t}{1+\sqrt{21}} \\ t \end{bmatrix} = \begin{bmatrix} \frac{10}{1+\sqrt{21}} \\ 1 \end{bmatrix}$$

Which is normalized to

$$\begin{bmatrix} \frac{10t}{1+\sqrt{21}} \\ t \end{bmatrix} = \begin{bmatrix} \frac{10}{1+\sqrt{21}} \\ 1 \end{bmatrix}$$

Considering the eigenvalue $\lambda_2 = \frac{5}{2} + \frac{\sqrt{21}}{2}$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\left[\begin{array}{cc} 3 & -5 \\ -1 & 2 \end{array} \right] - \left(\frac{5}{2} + \frac{\sqrt{21}}{2} \right) \left[\begin{array}{cc} 1 & 0 \\ 0 & 1 \end{array} \right] \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\left[\begin{array}{cc|c} \frac{1}{2} - \frac{\sqrt{21}}{2} & -5 & 0 \\ -1 & -\frac{1}{2} - \frac{\sqrt{21}}{2} & 0 \end{array} \right] \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} \frac{1}{2} - \frac{\sqrt{21}}{2} & -5 & 0 \\ -1 & -\frac{1}{2} - \frac{\sqrt{21}}{2} & 0 \end{array} \right]$$

$$R_2 = R_2 + \frac{R_1}{\frac{1}{2} - \frac{\sqrt{21}}{2}} \implies \left[\begin{array}{cc|c} \frac{1}{2} - \frac{\sqrt{21}}{2} & -5 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\left[\begin{array}{cc|c} \frac{1}{2} - \frac{\sqrt{21}}{2} & -5 & 0 \\ 0 & 0 & 0 \end{array} \right] \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\left\{ v_1 = -\frac{10t}{-1+\sqrt{21}} \right\}$

Hence the solution is

$$\begin{bmatrix} -\frac{10t}{-1+\sqrt{21}} \\ t \end{bmatrix} = \begin{bmatrix} -\frac{10t}{-1+\sqrt{21}} \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} -\frac{10t}{-1+\sqrt{21}} \\ t \end{bmatrix} = t \begin{bmatrix} -\frac{10}{-1+\sqrt{21}} \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} -\frac{10t}{-1+\sqrt{21}} \\ t \end{bmatrix} = \begin{bmatrix} -\frac{10}{-1+\sqrt{21}} \\ 1 \end{bmatrix}$$

Which is normalized to

$$\begin{bmatrix} -\frac{10t}{-1+\sqrt{21}} \\ t \end{bmatrix} = \begin{bmatrix} -\frac{10}{-1+\sqrt{21}} \\ 1 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated

with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
$\frac{5}{2} + \frac{\sqrt{21}}{2}$	1	1	No	$\begin{bmatrix} -\frac{5}{-\frac{1}{2} + \frac{\sqrt{21}}{2}} \\ 1 \end{bmatrix}$
$\frac{5}{2} - \frac{\sqrt{21}}{2}$	1	1	No	$\begin{bmatrix} -\frac{5}{-\frac{1}{2} - \frac{\sqrt{21}}{2}} \\ 1 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. Since eigenvalue $\frac{5}{2} + \frac{\sqrt{21}}{2}$ is real and distinct then the corresponding eigenvector solution is

$$\begin{aligned} \vec{x}_1(t) &= \vec{v}_1 e^{\left(\frac{5}{2} + \frac{\sqrt{21}}{2}\right)t} \\ &= \begin{bmatrix} -\frac{5}{-\frac{1}{2} + \frac{\sqrt{21}}{2}} \\ 1 \end{bmatrix} e^{\left(\frac{5}{2} + \frac{\sqrt{21}}{2}\right)t} \end{aligned}$$

Since eigenvalue $\frac{5}{2} - \frac{\sqrt{21}}{2}$ is real and distinct then the corresponding eigenvector solution is

$$\begin{aligned} \vec{x}_2(t) &= \vec{v}_2 e^{\left(\frac{5}{2} - \frac{\sqrt{21}}{2}\right)t} \\ &= \begin{bmatrix} -\frac{5}{-\frac{1}{2} - \frac{\sqrt{21}}{2}} \\ 1 \end{bmatrix} e^{\left(\frac{5}{2} - \frac{\sqrt{21}}{2}\right)t} \end{aligned}$$

Therefore the final solution is

$$\vec{x}_h(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

Which is written as

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = c_1 \begin{bmatrix} -\frac{5e^{\left(\frac{5}{2} + \frac{\sqrt{21}}{2}\right)t}}{-\frac{1}{2} + \frac{\sqrt{21}}{2}} \\ e^{\left(\frac{5}{2} + \frac{\sqrt{21}}{2}\right)t} \end{bmatrix} + c_2 \begin{bmatrix} -\frac{5e^{\left(\frac{5}{2} - \frac{\sqrt{21}}{2}\right)t}}{-\frac{1}{2} - \frac{\sqrt{21}}{2}} \\ e^{\left(\frac{5}{2} - \frac{\sqrt{21}}{2}\right)t} \end{bmatrix}$$

Which becomes

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} \frac{c_2(-1+\sqrt{21})e^{-\frac{(-5+\sqrt{21})t}{2}}}{2} - \frac{e^{\frac{(5+\sqrt{21})t}{2}}c_1(1+\sqrt{21})}{2} \\ c_1e^{\frac{(5+\sqrt{21})t}{2}} + c_2e^{-\frac{(-5+\sqrt{21})t}{2}} \end{bmatrix}$$

The following is the phase plot of the system.




Figure 509: Phase plot

29.7.3 Maple step by step solution

Let's solve

$$[x'(t) = 3x(t) - 5y(t), y'(t) = -x(t) + 2y(t)]$$

- Define vector

$$\vec{x}(t) = \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

- Convert system into a vector equation

$$\vec{x}'(t) = \begin{bmatrix} 3 & -5 \\ -1 & 2 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

- System to solve

$$\vec{x}'(t) = \begin{bmatrix} 3 & -5 \\ -1 & 2 \end{bmatrix} \cdot \vec{x}(t)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 3 & -5 \\ -1 & 2 \end{bmatrix}$$

- Rewrite the system as

$$\vec{x}'(t) = A \cdot \vec{x}(t)$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[\frac{5}{2} - \frac{\sqrt{21}}{2}, \begin{bmatrix} -\frac{5}{-\frac{1}{2} - \frac{\sqrt{21}}{2}} \\ 1 \end{bmatrix} \right], \left[\frac{5}{2} + \frac{\sqrt{21}}{2}, \begin{bmatrix} -\frac{5}{-\frac{1}{2} + \frac{\sqrt{21}}{2}} \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[\frac{5}{2} - \frac{\sqrt{21}}{2}, \begin{bmatrix} -\frac{5}{-\frac{1}{2} - \frac{\sqrt{21}}{2}} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{x}_1 = e^{\left(\frac{5}{2} - \frac{\sqrt{21}}{2}\right)t} \cdot \begin{bmatrix} -\frac{5}{-\frac{1}{2} - \frac{\sqrt{21}}{2}} \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[\frac{5}{2} + \frac{\sqrt{21}}{2}, \begin{bmatrix} -\frac{5}{-\frac{1}{2} + \frac{\sqrt{21}}{2}} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{x}_2 = e^{\left(\frac{5}{2} + \frac{\sqrt{21}}{2}\right)t} \cdot \begin{bmatrix} -\frac{5}{-\frac{1}{2} + \frac{\sqrt{21}}{2}} \\ 1 \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{x} = c_1 \vec{x}_1 + c_2 \vec{x}_2$$

- Substitute solutions into the general solution

$$\vec{x} = c_1 e^{\left(\frac{5}{2} - \frac{\sqrt{21}}{2}\right)t} \cdot \begin{bmatrix} -\frac{5}{-\frac{1}{2} - \frac{\sqrt{21}}{2}} \\ 1 \end{bmatrix} + c_2 e^{\left(\frac{5}{2} + \frac{\sqrt{21}}{2}\right)t} \cdot \begin{bmatrix} -\frac{5}{-\frac{1}{2} + \frac{\sqrt{21}}{2}} \\ 1 \end{bmatrix}$$

- Substitute in vector of dependent variables

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} \frac{c_1(-1+\sqrt{21})e^{-\frac{(-5+\sqrt{21})t}{2}}}{2} - \frac{c_2e^{\frac{(5+\sqrt{21})t}{2}}(1+\sqrt{21})}{2} \\ c_1e^{-\frac{(-5+\sqrt{21})t}{2}} + c_2e^{\frac{(5+\sqrt{21})t}{2}} \end{bmatrix}$$

- Solution to the system of ODEs

$$\left\{ x(t) = \frac{c_1(-1+\sqrt{21})e^{-\frac{(-5+\sqrt{21})t}{2}}}{2} - \frac{c_2e^{\frac{(5+\sqrt{21})t}{2}}(1+\sqrt{21})}{2}, y(t) = c_1e^{-\frac{(-5+\sqrt{21})t}{2}} + c_2e^{\frac{(5+\sqrt{21})t}{2}} \right\}$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 86

```
dsolve([diff(x(t),t)=3*x(t)-5*y(t),diff(y(t),t)=-x(t)+2*y(t)],singsol=all)
```

$$x(t) = c_1 e^{\frac{(5+\sqrt{21})t}{2}} + c_2 e^{-\frac{(-5+\sqrt{21})t}{2}}$$

$$y(t) = -\frac{c_1 e^{\frac{(5+\sqrt{21})t}{2}} \sqrt{21}}{10} + \frac{c_2 e^{-\frac{(-5+\sqrt{21})t}{2}} \sqrt{21}}{10} + \frac{c_1 e^{\frac{(5+\sqrt{21})t}{2}}}{10} + \frac{c_2 e^{-\frac{(-5+\sqrt{21})t}{2}}}{10}$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 144

```
DSolve[{x'[t]==3*x[t]-5*y[t],y'[t]==-x[t]+2*y[t]},{x[t],y[t]},t,IncludeSingularSolutions ->
```

$$x(t) \rightarrow \frac{1}{42} e^{-\frac{1}{2}(\sqrt{21}-5)t} \left(c_1 \left((21 + \sqrt{21}) e^{\sqrt{21}t} + 21 - \sqrt{21} \right) - 10\sqrt{21}c_2 \left(e^{\sqrt{21}t} - 1 \right) \right)$$

$$y(t) \rightarrow -\frac{1}{42} e^{-\frac{1}{2}(\sqrt{21}-5)t} \left(2\sqrt{21}c_1 \left(e^{\sqrt{21}t} - 1 \right) + c_2 \left((\sqrt{21} - 21) e^{\sqrt{21}t} - 21 - \sqrt{21} \right) \right)$$

29.8 problem 3(d)

29.8.1 Solution using Matrix exponential method	4148
29.8.2 Solution using explicit Eigenvalue and Eigenvector method . . .	4149
29.8.3 Maple step by step solution	4154

Internal problem ID [6536]

Internal file name [OUTPUT/5784_Sunday_June_05_2022_03_54_12_PM_38804077/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 10. Systems of First-Order Equations. Section A. Drill exercises. Page 400

Problem number: 3(d).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**system of linear ODEs**"

Solve

$$\begin{aligned}x'(t) &= x(t) + 2y(t) \\y'(t) &= -4x(t) + y(t)\end{aligned}$$

29.8.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A \vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ -4 & 1 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} e^t \cos(2t\sqrt{2}) & \frac{e^t \sin(2t\sqrt{2})\sqrt{2}}{2} \\ -e^t \sin(2t\sqrt{2})\sqrt{2} & e^t \cos(2t\sqrt{2}) \end{bmatrix}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 \vec{x}_h(t) &= e^{At} \vec{c} \\
 &= \begin{bmatrix} e^t \cos(2t\sqrt{2}) & \frac{e^t \sin(2t\sqrt{2})\sqrt{2}}{2} \\ -e^t \sin(2t\sqrt{2})\sqrt{2} & e^t \cos(2t\sqrt{2}) \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} \\
 &= \begin{bmatrix} e^t \cos(2t\sqrt{2}) c_1 + \frac{e^t \sin(2t\sqrt{2})\sqrt{2} c_2}{2} \\ -e^t \sin(2t\sqrt{2})\sqrt{2} c_1 + e^t \cos(2t\sqrt{2}) c_2 \end{bmatrix} \\
 &= \begin{bmatrix} e^t \cos(2t\sqrt{2}) c_1 + \frac{e^t \sin(2t\sqrt{2})\sqrt{2} c_2}{2} \\ e^t (-\sin(2t\sqrt{2})\sqrt{2} c_1 + \cos(2t\sqrt{2}) c_2) \end{bmatrix}
 \end{aligned}$$

Since no forcing function is given, then the final solution is $\vec{x}_h(t)$ above.

29.8.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A \vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ -4 & 1 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det \left(\begin{bmatrix} 1 & 2 \\ -4 & 1 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) = 0$$

Therefore

$$\det \left(\begin{bmatrix} 1 - \lambda & 2 \\ -4 & 1 - \lambda \end{bmatrix} \right) = 0$$

Which gives the characteristic equation

$$\lambda^2 - 2\lambda + 9 = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = 2i\sqrt{2} + 1$$

$$\lambda_2 = 1 - 2i\sqrt{2}$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
$1 - 2i\sqrt{2}$	1	complex eigenvalue
$2i\sqrt{2} + 1$	1	complex eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = 1 - 2i\sqrt{2}$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 1 & 2 \\ -4 & 1 \end{bmatrix} - (1 - 2i\sqrt{2}) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 2i\sqrt{2} & 2 \\ -4 & 2i\sqrt{2} \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} 2i\sqrt{2} & 2 & 0 \\ -4 & 2i\sqrt{2} & 0 \end{array} \right]$$

$$R_2 = R_2 - i\sqrt{2}R_1 \implies \left[\begin{array}{cc|c} 2i\sqrt{2} & 2 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} 2i\sqrt{2} & 2 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\left\{v_1 = \frac{it\sqrt{2}}{2}\right\}$

Hence the solution is

$$\begin{bmatrix} \frac{1}{2}t\sqrt{2} \\ t \end{bmatrix} = \begin{bmatrix} \frac{it\sqrt{2}}{2} \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} \frac{1}{2}t\sqrt{2} \\ t \end{bmatrix} = t \begin{bmatrix} \frac{i\sqrt{2}}{2} \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} \frac{1}{2}t\sqrt{2} \\ t \end{bmatrix} = \begin{bmatrix} \frac{i\sqrt{2}}{2} \\ 1 \end{bmatrix}$$

Which is normalized to

$$\begin{bmatrix} \frac{1}{2}t\sqrt{2} \\ t \end{bmatrix} = \begin{bmatrix} i\sqrt{2} \\ 2 \end{bmatrix}$$

Considering the eigenvalue $\lambda_2 = 2i\sqrt{2} + 1$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 1 & 2 \\ -4 & 1 \end{bmatrix} - (2i\sqrt{2} + 1) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} -2i\sqrt{2} & 2 \\ -4 & -2i\sqrt{2} \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} -2i\sqrt{2} & 2 & 0 \\ -4 & -2i\sqrt{2} & 0 \end{array} \right]$$

$$R_2 = R_2 + i\sqrt{2}R_1 \implies \left[\begin{array}{cc|c} -2i\sqrt{2} & 2 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} -2i\sqrt{2} & 2 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\left\{ v_1 = -\frac{it\sqrt{2}}{2} \right\}$

Hence the solution is

$$\begin{bmatrix} -\frac{1}{2}t\sqrt{2} \\ t \end{bmatrix} = \begin{bmatrix} -\frac{it\sqrt{2}}{2} \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} -\frac{1}{2}t\sqrt{2} \\ t \end{bmatrix} = t \begin{bmatrix} -\frac{i\sqrt{2}}{2} \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} -\frac{1}{2}t\sqrt{2} \\ t \end{bmatrix} = \begin{bmatrix} -\frac{i\sqrt{2}}{2} \\ 1 \end{bmatrix}$$

Which is normalized to

$$\begin{bmatrix} -\frac{1}{2}t\sqrt{2} \\ t \end{bmatrix} = \begin{bmatrix} -i\sqrt{2} \\ 2 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
$2i\sqrt{2} + 1$	1	1	No	$\begin{bmatrix} -\frac{i\sqrt{2}}{2} \\ 1 \end{bmatrix}$
$1 - 2i\sqrt{2}$	1	1	No	$\begin{bmatrix} \frac{i\sqrt{2}}{2} \\ 1 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. Therefore the final solution is

$$\vec{x}_h(t) = c_1\vec{x}_1(t) + c_2\vec{x}_2(t)$$

Which is written as

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = c_1 \begin{bmatrix} -\frac{ie^{(2i\sqrt{2}+1)t}\sqrt{2}}{2} \\ e^{(2i\sqrt{2}+1)t} \end{bmatrix} + c_2 \begin{bmatrix} \frac{ie^{(1-2i\sqrt{2})t}\sqrt{2}}{2} \\ e^{(1-2i\sqrt{2})t} \end{bmatrix}$$

Which becomes

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} -\frac{i\sqrt{2}(c_1e^{2i\sqrt{2}t+t} - c_2e^{-2i\sqrt{2}t+t})}{2} \\ c_1e^{2i\sqrt{2}t+t} + c_2e^{-2i\sqrt{2}t+t} \end{bmatrix}$$

The following is the phase plot of the system.




Figure 510: Phase plot

29.8.3 Maple step by step solution

Let's solve

$$[x'(t) = x(t) + 2y(t), y'(t) = -4x(t) + y(t)]$$

- Define vector

$$\vec{x}(t) = \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

- Convert system into a vector equation

$$\vec{x}'(t) = \begin{bmatrix} 1 & 2 \\ -4 & 1 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

- System to solve

$$\vec{x}'(t) = \begin{bmatrix} 1 & 2 \\ -4 & 1 \end{bmatrix} \cdot \vec{x}(t)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 1 & 2 \\ -4 & 1 \end{bmatrix}$$

- Rewrite the system as

$$\vec{x}'(t) = A \cdot \vec{x}(t)$$

- To solve the system, find the eigenvalues and eigenvectors of A
- Eigenpairs of A

$$\left[\left[1 - 2I\sqrt{2}, \begin{bmatrix} \frac{1}{2}\sqrt{2} \\ 1 \end{bmatrix} \right], \left[2I\sqrt{2} + 1, \begin{bmatrix} -\frac{1}{2}\sqrt{2} \\ 1 \end{bmatrix} \right] \right]$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$\left[1 - 2I\sqrt{2}, \begin{bmatrix} \frac{1}{2}\sqrt{2} \\ 1 \end{bmatrix} \right]$$

- Solution from eigenpair

$$e^{(1-2I\sqrt{2})t} \cdot \begin{bmatrix} \frac{1}{2}\sqrt{2} \\ 1 \end{bmatrix}$$

- Use Euler identity to write solution in terms of sin and cos

$$e^t \cdot (\cos(2t\sqrt{2}) - I \sin(2t\sqrt{2})) \cdot \begin{bmatrix} \frac{1}{2}\sqrt{2} \\ 1 \end{bmatrix}$$

- Simplify expression

$$e^t \cdot \begin{bmatrix} \frac{1}{2}(\cos(2t\sqrt{2}) - I \sin(2t\sqrt{2})) \sqrt{2} \\ \cos(2t\sqrt{2}) - I \sin(2t\sqrt{2}) \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\left[\vec{x}_1(t) = e^t \cdot \begin{bmatrix} \frac{\sqrt{2} \sin(2t\sqrt{2})}{2} \\ \cos(2t\sqrt{2}) \end{bmatrix}, \vec{x}_2(t) = e^t \cdot \begin{bmatrix} \frac{\sqrt{2} \cos(2t\sqrt{2})}{2} \\ -\sin(2t\sqrt{2}) \end{bmatrix} \right]$$

- General solution to the system of ODEs

$$\vec{x} = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

- Substitute solutions into the general solution

$$\vec{x} = c_1 e^t \cdot \begin{bmatrix} \frac{\sqrt{2} \sin(2t\sqrt{2})}{2} \\ \cos(2t\sqrt{2}) \end{bmatrix} + c_2 e^t \cdot \begin{bmatrix} \frac{\sqrt{2} \cos(2t\sqrt{2})}{2} \\ -\sin(2t\sqrt{2}) \end{bmatrix}$$

- Substitute in vector of dependent variables

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} \frac{e^t \sqrt{2} (c_1 \sin(2t\sqrt{2}) + c_2 \cos(2t\sqrt{2}))}{2} \\ e^t (c_1 \cos(2t\sqrt{2}) - c_2 \sin(2t\sqrt{2})) \end{bmatrix}$$

- Solution to the system of ODEs

$$\left\{ x(t) = \frac{e^t \sqrt{2} (c_1 \sin(2t\sqrt{2}) + c_2 \cos(2t\sqrt{2}))}{2}, y(t) = e^t (c_1 \cos(2t\sqrt{2}) - c_2 \sin(2t\sqrt{2})) \right\}$$

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 56

```
dsolve([diff(x(t),t)=x(t)+2*y(t),diff(y(t),t)=-4*x(t)+y(t)],singsol=all)
```

$$x(t) = e^t (c_2 \cos(2\sqrt{2}t) + c_1 \sin(2\sqrt{2}t))$$

$$y(t) = e^t \sqrt{2} (\cos(2\sqrt{2}t) c_1 - \sin(2\sqrt{2}t) c_2)$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 79

```
DSolve[{x'[t]==x[t]+2*y[t],y'[t]==-4*x[t]+y[t]},{x[t],y[t]},t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow c_1 e^t \cos(2\sqrt{2}t) + \frac{c_2 e^t \sin(2\sqrt{2}t)}{\sqrt{2}}$$

$$y(t) \rightarrow e^t (c_2 \cos(2\sqrt{2}t) - \sqrt{2} c_1 \sin(2\sqrt{2}t))$$

29.9 problem 3(e)

29.9.1 Solution using Matrix exponential method	4157
29.9.2 Solution using explicit Eigenvalue and Eigenvector method . . .	4158
29.9.3 Maple step by step solution	4166

Internal problem ID [6537]

Internal file name [OUTPUT/5785_Sunday_June_05_2022_03_54_14_PM_79079975/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 10. Systems of First-Order Equations. Section A. Drill exercises. Page 400

Problem number: 3(e).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "system of linear ODEs"

Solve

$$\begin{aligned}x'(t) &= 3x(t) + 2y(t) + z(t) \\y'(t) &= -2x(t) - y(t) + 3z(t) \\z'(t) &= x(t) + y(t) + z(t)\end{aligned}$$

29.9.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \\ z'(t) \end{bmatrix} = \begin{bmatrix} 3 & 2 & 1 \\ -2 & -1 & 3 \\ 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \\ z(t) \end{bmatrix}$$

For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} \frac{5e^{3t}}{8} + \frac{3e^t}{4} - \frac{3e^{-t}}{8} & \frac{5e^{3t}}{8} - \frac{3e^{-t}}{8} - \frac{e^t}{4} & \frac{5e^{3t}}{4} - 2e^t + \frac{3e^{-t}}{4} \\ -\frac{e^{3t}}{8} + \frac{7e^{-t}}{8} - \frac{3e^t}{4} & \frac{7e^{-t}}{8} - \frac{e^{3t}}{8} + \frac{e^t}{4} & -\frac{e^{3t}}{4} + 2e^t - \frac{7e^{-t}}{4} \\ \frac{e^{3t}}{4} - \frac{e^{-t}}{4} & \frac{e^{3t}}{4} - \frac{e^{-t}}{4} & \frac{e^{-t}}{2} + \frac{e^{3t}}{2} \end{bmatrix}$$

Therefore the homogeneous solution is

$$\begin{aligned}
 \vec{x}_h(t) &= e^{At} \vec{c} \\
 &= \begin{bmatrix} \frac{5e^{3t}}{8} + \frac{3e^t}{4} - \frac{3e^{-t}}{8} & \frac{5e^{3t}}{8} - \frac{3e^{-t}}{8} - \frac{e^t}{4} & \frac{5e^{3t}}{4} - 2e^t + \frac{3e^{-t}}{4} \\ -\frac{e^{3t}}{8} + \frac{7e^{-t}}{8} - \frac{3e^t}{4} & \frac{7e^{-t}}{8} - \frac{e^{3t}}{8} + \frac{e^t}{4} & -\frac{e^{3t}}{4} + 2e^t - \frac{7e^{-t}}{4} \\ \frac{e^{3t}}{4} - \frac{e^{-t}}{4} & \frac{e^{3t}}{4} - \frac{e^{-t}}{4} & \frac{e^{-t}}{2} + \frac{e^{3t}}{2} \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix} \\
 &= \begin{bmatrix} \left(\frac{5e^{3t}}{8} + \frac{3e^t}{4} - \frac{3e^{-t}}{8}\right) c_1 + \left(\frac{5e^{3t}}{8} - \frac{3e^{-t}}{8} - \frac{e^t}{4}\right) c_2 + \left(\frac{5e^{3t}}{4} - 2e^t + \frac{3e^{-t}}{4}\right) c_3 \\ \left(-\frac{e^{3t}}{8} + \frac{7e^{-t}}{8} - \frac{3e^t}{4}\right) c_1 + \left(\frac{7e^{-t}}{8} - \frac{e^{3t}}{8} + \frac{e^t}{4}\right) c_2 + \left(-\frac{e^{3t}}{4} + 2e^t - \frac{7e^{-t}}{4}\right) c_3 \\ \left(\frac{e^{3t}}{4} - \frac{e^{-t}}{4}\right) c_1 + \left(\frac{e^{3t}}{4} - \frac{e^{-t}}{4}\right) c_2 + \left(\frac{e^{-t}}{2} + \frac{e^{3t}}{2}\right) c_3 \end{bmatrix} \\
 &= \begin{bmatrix} \frac{3(-c_1 - c_2 + 2c_3)e^{-t}}{8} + \frac{5(c_1 + c_2 + 2c_3)e^{3t}}{8} + \frac{3\left(c_1 - \frac{c_2}{3} - \frac{8c_3}{3}\right)e^t}{4} \\ \frac{7(c_1 + c_2 - 2c_3)e^{-t}}{8} + \frac{(-c_1 - c_2 - 2c_3)e^{3t}}{8} - \frac{3\left(c_1 - \frac{c_2}{3} - \frac{8c_3}{3}\right)e^t}{4} \\ \frac{(-c_1 - c_2 + 2c_3)e^{-t}}{4} + \frac{(c_1 + c_2 + 2c_3)e^{3t}}{4} \end{bmatrix}
 \end{aligned}$$

Since no forcing function is given, then the final solution is $\vec{x}_h(t)$ above.

29.9.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A \vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \\ z'(t) \end{bmatrix} = \begin{bmatrix} 3 & 2 & 1 \\ -2 & -1 & 3 \\ 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \\ z(t) \end{bmatrix}$$

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det \left(\begin{bmatrix} 3 & 2 & 1 \\ -2 & -1 & 3 \\ 1 & 1 & 1 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right) = 0$$

Therefore

$$\det \left(\begin{bmatrix} 3 - \lambda & 2 & 1 \\ -2 & -1 - \lambda & 3 \\ 1 & 1 & 1 - \lambda \end{bmatrix} \right) = 0$$

Which gives the characteristic equation

$$\lambda^3 - 3\lambda^2 - \lambda + 3 = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = 1$$

$$\lambda_2 = -1$$

$$\lambda_3 = 3$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
-1	1	real eigenvalue
1	1	real eigenvalue
3	1	real eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = -1$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 3 & 2 & 1 \\ -2 & -1 & 3 \\ 1 & 1 & 1 \end{bmatrix} - (-1) \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 4 & 2 & 1 \\ -2 & 0 & 3 \\ 1 & 1 & 2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{ccc|c} 4 & 2 & 1 & 0 \\ -2 & 0 & 3 & 0 \\ 1 & 1 & 2 & 0 \end{array} \right]$$

$$R_2 = R_2 + \frac{R_1}{2} \implies \left[\begin{array}{ccc|c} 4 & 2 & 1 & 0 \\ 0 & 1 & \frac{7}{2} & 0 \\ 1 & 1 & 2 & 0 \end{array} \right]$$

$$R_3 = R_3 - \frac{R_1}{4} \implies \left[\begin{array}{ccc|c} 4 & 2 & 1 & 0 \\ 0 & 1 & \frac{7}{2} & 0 \\ 0 & \frac{1}{2} & \frac{7}{4} & 0 \end{array} \right]$$

$$R_3 = R_3 - \frac{R_2}{2} \implies \left[\begin{array}{ccc|c} 4 & 2 & 1 & 0 \\ 0 & 1 & \frac{7}{2} & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\left[\begin{array}{ccc} 4 & 2 & 1 \\ 0 & 1 & \frac{7}{2} \\ 0 & 0 & 0 \end{array} \right] \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_3\}$ and the leading variables are $\{v_1, v_2\}$. Let $v_3 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = \frac{3t}{2}, v_2 = -\frac{7t}{2}\}$

Hence the solution is

$$\begin{bmatrix} \frac{3t}{2} \\ -\frac{7t}{2} \\ t \end{bmatrix} = \begin{bmatrix} \frac{3t}{2} \\ -\frac{7t}{2} \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} \frac{3t}{2} \\ -\frac{7t}{2} \\ t \end{bmatrix} = t \begin{bmatrix} \frac{3}{2} \\ -\frac{7}{2} \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} \frac{3t}{2} \\ -\frac{7t}{2} \\ t \end{bmatrix} = \begin{bmatrix} \frac{3}{2} \\ -\frac{7}{2} \\ 1 \end{bmatrix}$$

Which is normalized to

$$\begin{bmatrix} \frac{3t}{2} \\ -\frac{7t}{2} \\ t \end{bmatrix} = \begin{bmatrix} 3 \\ -7 \\ 2 \end{bmatrix}$$

Considering the eigenvalue $\lambda_2 = 1$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 3 & 2 & 1 \\ -2 & -1 & 3 \\ 1 & 1 & 1 \end{bmatrix} - (1) \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 2 & 1 \\ -2 & -2 & 3 \\ 1 & 1 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{ccc|c} 2 & 2 & 1 & 0 \\ -2 & -2 & 3 & 0 \\ 1 & 1 & 0 & 0 \end{array} \right]$$

$$R_2 = R_2 + R_1 \implies \left[\begin{array}{ccc|c} 2 & 2 & 1 & 0 \\ 0 & 0 & 4 & 0 \\ 1 & 1 & 0 & 0 \end{array} \right]$$

$$R_3 = R_3 - \frac{R_1}{2} \implies \left[\begin{array}{ccc|c} 2 & 2 & 1 & 0 \\ 0 & 0 & 4 & 0 \\ 0 & 0 & -\frac{1}{2} & 0 \end{array} \right]$$

$$R_3 = R_3 + \frac{R_2}{8} \implies \left[\begin{array}{ccc|c} 2 & 2 & 1 & 0 \\ 0 & 0 & 4 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\left[\begin{array}{ccc} 2 & 2 & 1 \\ 0 & 0 & 4 \\ 0 & 0 & 0 \end{array} \right] \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1, v_3\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = -t, v_3 = 0\}$

Hence the solution is

$$\begin{bmatrix} -t \\ t \\ 0 \end{bmatrix} = \begin{bmatrix} -t \\ t \\ 0 \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} -t \\ t \\ 0 \end{bmatrix} = t \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} -t \\ t \\ 0 \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}$$

Considering the eigenvalue $\lambda_3 = 3$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{array}{c} \begin{bmatrix} 3 & 2 & 1 \\ -2 & -1 & 3 \\ 1 & 1 & 1 \end{bmatrix} - (3) \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \end{array} \right) \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 0 & 2 & 1 \\ -2 & -4 & 3 \\ 1 & 1 & -2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{ccc|c} 0 & 2 & 1 & 0 \\ -2 & -4 & 3 & 0 \\ 1 & 1 & -2 & 0 \end{array} \right]$$

Since the current pivot $A(1,1)$ is zero, then the current pivot row is replaced with a row with a non-zero pivot. Swapping row 1 and row 2 gives

$$\left[\begin{array}{ccc|c} -2 & -4 & 3 & 0 \\ 0 & 2 & 1 & 0 \\ 1 & 1 & -2 & 0 \end{array} \right]$$

$$R_3 = R_3 + \frac{R_1}{2} \implies \left[\begin{array}{ccc|c} -2 & -4 & 3 & 0 \\ 0 & 2 & 1 & 0 \\ 0 & -1 & -\frac{1}{2} & 0 \end{array} \right]$$

$$R_3 = R_3 + \frac{R_2}{2} \implies \left[\begin{array}{ccc|c} -2 & -4 & 3 & 0 \\ 0 & 2 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} -2 & -4 & 3 \\ 0 & 2 & 1 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_3\}$ and the leading variables are $\{v_1, v_2\}$. Let $v_3 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = \frac{5t}{2}, v_2 = -\frac{t}{2}\}$

Hence the solution is

$$\begin{bmatrix} \frac{5t}{2} \\ -\frac{t}{2} \\ t \end{bmatrix} = \begin{bmatrix} \frac{5t}{2} \\ -\frac{t}{2} \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} \frac{5t}{2} \\ -\frac{t}{2} \\ t \end{bmatrix} = t \begin{bmatrix} \frac{5}{2} \\ -\frac{1}{2} \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} \frac{5t}{2} \\ -\frac{t}{2} \\ t \end{bmatrix} = \begin{bmatrix} \frac{5}{2} \\ -\frac{1}{2} \\ 1 \end{bmatrix}$$

Which is normalized to

$$\begin{bmatrix} \frac{5t}{2} \\ -\frac{t}{2} \\ t \end{bmatrix} = \begin{bmatrix} 5 \\ -1 \\ 2 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
1	1	1	No	$\begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}$
-1	1	1	No	$\begin{bmatrix} \frac{3}{2} \\ -\frac{7}{2} \\ 1 \end{bmatrix}$
3	1	1	No	$\begin{bmatrix} \frac{5}{2} \\ -\frac{1}{2} \\ 1 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. Since eigenvalue 1 is real and distinct then the corresponding eigenvector solution is

$$\begin{aligned} \vec{x}_1(t) &= \vec{v}_1 e^t \\ &= \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix} e^t \end{aligned}$$

Since eigenvalue -1 is real and distinct then the corresponding eigenvector solution is

$$\begin{aligned} \vec{x}_2(t) &= \vec{v}_2 e^{-t} \\ &= \begin{bmatrix} \frac{3}{2} \\ -\frac{7}{2} \\ 1 \end{bmatrix} e^{-t} \end{aligned}$$

Since eigenvalue 3 is real and distinct then the corresponding eigenvector solution is

$$\begin{aligned}\vec{x}_3(t) &= \vec{v}_3 e^{3t} \\ &= \begin{bmatrix} \frac{5}{2} \\ -\frac{1}{2} \\ 1 \end{bmatrix} e^{3t}\end{aligned}$$

Therefore the final solution is

$$\vec{x}_h(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t) + c_3 \vec{x}_3(t)$$

Which is written as

$$\begin{bmatrix} x(t) \\ y(t) \\ z(t) \end{bmatrix} = c_1 \begin{bmatrix} -e^t \\ e^t \\ 0 \end{bmatrix} + c_2 \begin{bmatrix} \frac{3e^{-t}}{2} \\ -\frac{7e^{-t}}{2} \\ e^{-t} \end{bmatrix} + c_3 \begin{bmatrix} \frac{5e^{3t}}{2} \\ -\frac{e^{3t}}{2} \\ e^{3t} \end{bmatrix}$$

Which becomes

$$\begin{bmatrix} x(t) \\ y(t) \\ z(t) \end{bmatrix} = \begin{bmatrix} -c_1 e^t + \frac{3c_2 e^{-t}}{2} + \frac{5c_3 e^{3t}}{2} \\ c_1 e^t - \frac{7c_2 e^{-t}}{2} - \frac{c_3 e^{3t}}{2} \\ c_2 e^{-t} + c_3 e^{3t} \end{bmatrix}$$

29.9.3 Maple step by step solution

Let's solve

$$[x'(t) = 3x(t) + 2y(t) + z(t), y'(t) = -2x(t) - y(t) + 3z(t), z'(t) = x(t) + y(t) + z(t)]$$

- Define vector

$$\vec{x}(t) = \begin{bmatrix} x(t) \\ y(t) \\ z(t) \end{bmatrix}$$

- Convert system into a vector equation

$$\vec{x}'(t) = \begin{bmatrix} 3 & 2 & 1 \\ -2 & -1 & 3 \\ 1 & 1 & 1 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

- System to solve

$$\vec{x}'(t) = \begin{bmatrix} 3 & 2 & 1 \\ -2 & -1 & 3 \\ 1 & 1 & 1 \end{bmatrix} \cdot \vec{x}(t)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 3 & 2 & 1 \\ -2 & -1 & 3 \\ 1 & 1 & 1 \end{bmatrix}$$

- Rewrite the system as

$$\vec{x}'(t) = A \cdot \vec{x}(t)$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[-1, \begin{bmatrix} \frac{3}{2} \\ -\frac{7}{2} \\ 1 \end{bmatrix} \right], \left[1, \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix} \right], \left[3, \begin{bmatrix} \frac{5}{2} \\ -\frac{1}{2} \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[-1, \begin{bmatrix} \frac{3}{2} \\ -\frac{7}{2} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{x}_1 = e^{-t} \cdot \begin{bmatrix} \frac{3}{2} \\ -\frac{7}{2} \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[1, \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{x}_2 = e^t \cdot \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}$$

- Consider eigenpair

$$\left[3, \begin{bmatrix} \frac{5}{2} \\ -\frac{1}{2} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{x}_3 = e^{3t} \cdot \begin{bmatrix} \frac{5}{2} \\ -\frac{1}{2} \\ 1 \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{x} = c_1 \vec{x}_1 + c_2 \vec{x}_2 + c_3 \vec{x}_3$$

- Substitute solutions into the general solution

$$\vec{x} = e^{-t} c_1 \cdot \begin{bmatrix} \frac{3}{2} \\ -\frac{7}{2} \\ 1 \end{bmatrix} + c_2 e^t \cdot \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix} + c_3 e^{3t} \cdot \begin{bmatrix} \frac{5}{2} \\ -\frac{1}{2} \\ 1 \end{bmatrix}$$

- Substitute in vector of dependent variables

$$\begin{bmatrix} x(t) \\ y(t) \\ z(t) \end{bmatrix} = \begin{bmatrix} \frac{3e^{-t}c_1}{2} - c_2e^t + \frac{5c_3e^{3t}}{2} \\ -\frac{7e^{-t}c_1}{2} + c_2e^t - \frac{c_3e^{3t}}{2} \\ e^{-t}c_1 + c_3e^{3t} \end{bmatrix}$$

- Solution to the system of ODEs

$$\left\{ x(t) = \frac{3e^{-t}c_1}{2} - c_2e^t + \frac{5c_3e^{3t}}{2}, y(t) = -\frac{7e^{-t}c_1}{2} + c_2e^t - \frac{c_3e^{3t}}{2}, z(t) = e^{-t}c_1 + c_3e^{3t} \right\}$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 63

```
dsolve([diff(x(t),t)=3*x(t)+2*y(t)+z(t),diff(y(t),t)=-2*x(t)-y(t)+3*z(t),diff(z(t),t)=x(t)+y
```

$$\begin{aligned}x(t) &= c_1 e^t + c_2 e^{-t} + c_3 e^{3t} \\y(t) &= -c_1 e^t - \frac{7c_2 e^{-t}}{3} - \frac{c_3 e^{3t}}{5} \\z(t) &= \frac{2c_2 e^{-t}}{3} + \frac{2c_3 e^{3t}}{5}\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.012 (sec). Leaf size: 180

```
DSolve[{x'[t]==3*x[t]+2*y[t]+z[t],y'[t]==-2*x[t]-y[t]+3*z[t],z'[t]==x[t]+y[t]+z[t]},{x[t],y[t]
```

$$\begin{aligned}x(t) &\rightarrow \frac{1}{8}e^{-t}(c_1(6e^{2t} + 5e^{4t} - 3) + (e^{2t} - 1)(c_2(5e^{2t} + 3) + 2c_3(5e^{2t} - 3))) \\y(t) &\rightarrow \frac{1}{8}e^{-t}(-c_1(6e^{2t} + e^{4t} - 7) + c_2(2e^{2t} - e^{4t} + 7) - 2c_3(-8e^{2t} + e^{4t} + 7)) \\z(t) &\rightarrow \frac{1}{4}e^{-t}(c_1(e^{4t} - 1) + c_2(e^{4t} - 1) + 2c_3(e^{4t} + 1))\end{aligned}$$

29.10 problem 3(f)

29.10.1 Solution using Matrix exponential method 4170

29.10.2 Solution using explicit Eigenvalue and Eigenvector method . . . 4171

Internal problem ID [6538]

Internal file name [OUTPUT/5786_Sunday_June_05_2022_03_54_16_PM_17543604/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 10. Systems of First-Order Equations. Section A. Drill exercises. Page 400

Problem number: 3(f).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**system of linear ODEs**"

Solve

$$x'(t) = -x(t) + y(t) - z(t)$$

$$y'(t) = 2x(t) - y(t) - 4z(t)$$

$$z'(t) = 3x(t) - y(t) + z(t)$$

29.10.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \\ z'(t) \end{bmatrix} = \begin{bmatrix} -1 & 1 & -1 \\ 2 & -1 & -4 \\ 3 & -1 & 1 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \\ z(t) \end{bmatrix}$$

For the above matrix A , the matrix exponential can be found to be

$$\begin{aligned} e^{At} &= \text{Expression too large to display} \\ &= \text{Expression too large to display} \end{aligned}$$

Therefore the homogeneous solution is

$$\begin{aligned}\vec{x}_h(t) &= e^{At}\vec{c} \\ &= \text{Expression too large to display} \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix} \\ &= \text{Expression too large to display} \\ &= \text{Expression too large to display}\end{aligned}$$

Since no forcing function is given, then the final solution is $\vec{x}_h(t)$ above.

29.10.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \\ z'(t) \end{bmatrix} = \begin{bmatrix} -1 & 1 & -1 \\ 2 & -1 & -4 \\ 3 & -1 & 1 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \\ z(t) \end{bmatrix}$$

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det \left(\begin{bmatrix} -1 & 1 & -1 \\ 2 & -1 & -4 \\ 3 & -1 & 1 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right) = 0$$

Therefore

$$\det \left(\begin{bmatrix} -1 - \lambda & 1 & -1 \\ 2 & -1 - \lambda & -4 \\ 3 & -1 & 1 - \lambda \end{bmatrix} \right) = 0$$

Which gives the characteristic equation

$$\lambda^3 + \lambda^2 - 4\lambda + 10 = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = -\frac{(154 + 3\sqrt{2391})^{\frac{1}{3}}}{3} - \frac{13}{3(154 + 3\sqrt{2391})^{\frac{1}{3}}} - \frac{1}{3}$$

$$\lambda_2 = \frac{(154 + 3\sqrt{2391})^{\frac{1}{3}}}{6} + \frac{13}{6(154 + 3\sqrt{2391})^{\frac{1}{3}}} - \frac{1}{3} + \frac{i\sqrt{3} \left(-\frac{(154 + 3\sqrt{2391})^{\frac{1}{3}}}{3} + \frac{13}{3(154 + 3\sqrt{2391})^{\frac{1}{3}}} \right)}{2}$$

$$\lambda_3 = \frac{(154 + 3\sqrt{2391})^{\frac{1}{3}}}{6} + \frac{13}{6(154 + 3\sqrt{2391})^{\frac{1}{3}}} - \frac{1}{3} - \frac{i\sqrt{3} \left(-\frac{(154 + 3\sqrt{2391})^{\frac{1}{3}}}{3} + \frac{13}{3(154 + 3\sqrt{2391})^{\frac{1}{3}}} \right)}{2}$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of
$\frac{(154 + 3\sqrt{2391})^{\frac{1}{3}}}{6} + \frac{13}{6(154 + 3\sqrt{2391})^{\frac{1}{3}}} - \frac{1}{3} - \frac{i\sqrt{3} \left(-\frac{(154 + 3\sqrt{2391})^{\frac{1}{3}}}{3} + \frac{13}{3(154 + 3\sqrt{2391})^{\frac{1}{3}}} \right)}{2}$	1	complex
$-\frac{(154 + 3\sqrt{2391})^{\frac{1}{3}}}{3} - \frac{13}{3(154 + 3\sqrt{2391})^{\frac{1}{3}}} - \frac{1}{3}$	1	real eigenvalue
$\frac{(154 + 3\sqrt{2391})^{\frac{1}{3}}}{6} + \frac{13}{6(154 + 3\sqrt{2391})^{\frac{1}{3}}} - \frac{1}{3} + \frac{i\sqrt{3} \left(-\frac{(154 + 3\sqrt{2391})^{\frac{1}{3}}}{3} + \frac{13}{3(154 + 3\sqrt{2391})^{\frac{1}{3}}} \right)}{2}$	1	complex

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = -\frac{(154 + 3\sqrt{2391})^{\frac{1}{3}}}{3} - \frac{13}{3(154 + 3\sqrt{2391})^{\frac{1}{3}}} - \frac{1}{3}$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} -1 & 1 & -1 \\ 2 & -1 & -4 \\ 3 & -1 & 1 \end{bmatrix} - \left(-\frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{3} - \frac{13}{3(154+3\sqrt{2391})^{\frac{1}{3}}} - \frac{1}{3} \right) \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right) \begin{bmatrix} \frac{(154+3\sqrt{2391})^{\frac{2}{3}} - 2(154+3\sqrt{2391})^{\frac{1}{3}} + 13}{3(154+3\sqrt{2391})^{\frac{1}{3}}} & 1 & -1 \\ 2 & \frac{(154+3\sqrt{2391})^{\frac{2}{3}} - 2(154+3\sqrt{2391})^{\frac{1}{3}} + 13}{3(154+3\sqrt{2391})^{\frac{1}{3}}} & -4 \\ 3 & -1 & \frac{(154+3\sqrt{2391})^{\frac{2}{3}} + 4(154+3\sqrt{2391})^{\frac{1}{3}} + 13}{3(154+3\sqrt{2391})^{\frac{1}{3}}} \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{ccc|ccc} -\frac{2}{3} + \frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{3} + \frac{13}{3(154+3\sqrt{2391})^{\frac{1}{3}}} & & & 1 & & -1 \\ & 2 & & -\frac{2}{3} + \frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{3} + \frac{13}{3(154+3\sqrt{2391})^{\frac{1}{3}}} & & -4 \\ & & 3 & -1 & & \frac{4}{3} + \frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{3} + \frac{13}{3(154+3\sqrt{2391})^{\frac{1}{3}}} \end{array} \right]$$

$$R_2 = R_2 - \frac{2R_1}{-\frac{2}{3} + \frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{3} + \frac{13}{3(154+3\sqrt{2391})^{\frac{1}{3}}}} \implies \left[\begin{array}{ccc|ccc} \frac{(154+3\sqrt{2391})^{\frac{2}{3}} - 2(154+3\sqrt{2391})^{\frac{1}{3}} + 13}{3(154+3\sqrt{2391})^{\frac{1}{3}}} & & & 1 & & -1 \\ & 0 & & & & \frac{\sqrt{2391}(154+3\sqrt{2391})^{\frac{1}{3}}}{(154+3\sqrt{2391})^{\frac{1}{3}}} \\ & & 3 & -1 & & \frac{4}{3} + \frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{3} + \frac{13}{3(154+3\sqrt{2391})^{\frac{1}{3}}} \end{array} \right]$$

$$R_3 = R_3 - \frac{9(154 + 3\sqrt{2391})^{\frac{1}{3}} R_1}{(154 + 3\sqrt{2391})^{\frac{2}{3}} - 2(154 + 3\sqrt{2391})^{\frac{1}{3}} + 13} \Rightarrow \left[\begin{array}{ccc} \frac{(154+3\sqrt{2391})^{\frac{2}{3}} - 2(154+3\sqrt{2391})^{\frac{1}{3}} + 13}{3(154+3\sqrt{2391})^{\frac{1}{3}}} & & \\ & 0 & \frac{\sqrt{2391}}{(154+3\sqrt{2391})^{\frac{1}{3}}} \\ & 0 & \end{array} \right]$$

$$R_3 = R_3 - \frac{\left(-(154 + 3\sqrt{2391})^{\frac{2}{3}} - 7(154 + 3\sqrt{2391})^{\frac{1}{3}} - 13 \right) (154 + 3\sqrt{2391})^{\frac{1}{3}} R_2}{\sqrt{2391} (154 + 3\sqrt{2391})^{\frac{1}{3}} + 4(154 + 3\sqrt{2391})^{\frac{2}{3}} - 4\sqrt{2391} + 34(154 + 3\sqrt{2391})^{\frac{1}{3}} - 149} \Rightarrow$$

Therefore the system in Echelon form is

$$\left[\begin{array}{ccc} \frac{(154+3\sqrt{2391})^{\frac{2}{3}} - 2(154+3\sqrt{2391})^{\frac{1}{3}} + 13}{3(154+3\sqrt{2391})^{\frac{1}{3}}} & & 1 \\ & 0 & \frac{\sqrt{2391} (154+3\sqrt{2391})^{\frac{1}{3}} + 4(154+3\sqrt{2391})^{\frac{2}{3}} - 4\sqrt{2391} + 34(154+3\sqrt{2391})^{\frac{1}{3}} - 149}{(154+3\sqrt{2391})^{\frac{1}{3}} \left((154+3\sqrt{2391})^{\frac{2}{3}} - 2(154+3\sqrt{2391})^{\frac{1}{3}} + 13 \right)} \frac{-4(154+3\sqrt{2391})^{\frac{1}{3}}}{(154+3\sqrt{2391})^{\frac{1}{3}}} \\ & 0 & 0 \end{array} \right]$$

The free variables are $\{v_3\}$ and the leading variables are $\{v_1, v_2\}$. Let $v_3 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of

free variables gives equation $\left\{ \begin{array}{l} v_1 = \frac{3(154+3\sqrt{2391})^{\frac{1}{3}} t \left(\sqrt{2391} (154+3\sqrt{2391})^{\frac{1}{3}} + 18(154+3\sqrt{2391})^{\frac{2}{3}} - 16\sqrt{2391} - 18 \right)}{\left(\sqrt{2391} (154+3\sqrt{2391})^{\frac{1}{3}} + 4(154+3\sqrt{2391})^{\frac{2}{3}} - 4\sqrt{2391} + 34(154+3\sqrt{2391})^{\frac{1}{3}} - 149 \right) \left((154+3\sqrt{2391})^{\frac{1}{3}} \right)} \end{array} \right.$

Hence the solution is

$$\left[\begin{array}{c} \frac{3(154+3\sqrt{2391})^{\frac{1}{3}}t\left(\sqrt{2391}(154+3\sqrt{2391})^{\frac{1}{3}}+18(154+3\sqrt{2391})^{\frac{2}{3}}-16\sqrt{2391}-18(154+3\sqrt{2391})^{\frac{1}{3}}-765\right)}{\left(\sqrt{2391}(154+3\sqrt{2391})^{\frac{1}{3}}+4(154+3\sqrt{2391})^{\frac{2}{3}}-4\sqrt{2391}+34(154+3\sqrt{2391})^{\frac{1}{3}}-149\right)\left(\left(154+3\sqrt{2391}\right)^{\frac{2}{3}}-2(154+3\sqrt{2391})^{\frac{1}{3}}+13\right)} \\ -\frac{2t\left(7(154+3\sqrt{2391})^{\frac{2}{3}}-26(154+3\sqrt{2391})^{\frac{1}{3}}-6\sqrt{2391}-308\right)}{\sqrt{2391}(154+3\sqrt{2391})^{\frac{1}{3}}+4(154+3\sqrt{2391})^{\frac{2}{3}}-4\sqrt{2391}+34(154+3\sqrt{2391})^{\frac{1}{3}}-149} \\ t \end{array} \right] =$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\left[\begin{array}{c} \frac{3(154+3\sqrt{2391})^{\frac{1}{3}}t\left(\sqrt{2391}(154+3\sqrt{2391})^{\frac{1}{3}}+18(154+3\sqrt{2391})^{\frac{2}{3}}-16\sqrt{2391}-18(154+3\sqrt{2391})^{\frac{1}{3}}-765\right)}{\left(\sqrt{2391}(154+3\sqrt{2391})^{\frac{1}{3}}+4(154+3\sqrt{2391})^{\frac{2}{3}}-4\sqrt{2391}+34(154+3\sqrt{2391})^{\frac{1}{3}}-149\right)\left(\left(154+3\sqrt{2391}\right)^{\frac{2}{3}}-2(154+3\sqrt{2391})^{\frac{1}{3}}+13\right)} \\ -\frac{2t\left(7(154+3\sqrt{2391})^{\frac{2}{3}}-26(154+3\sqrt{2391})^{\frac{1}{3}}-6\sqrt{2391}-308\right)}{\sqrt{2391}(154+3\sqrt{2391})^{\frac{1}{3}}+4(154+3\sqrt{2391})^{\frac{2}{3}}-4\sqrt{2391}+34(154+3\sqrt{2391})^{\frac{1}{3}}-149} \\ t \end{array} \right] = t$$

Let $t = 1$ the eigenvector becomes

$$\left[\begin{array}{c} \frac{3(154+3\sqrt{2391})^{\frac{1}{3}}t\left(\sqrt{2391}(154+3\sqrt{2391})^{\frac{1}{3}}+18(154+3\sqrt{2391})^{\frac{2}{3}}-16\sqrt{2391}-18(154+3\sqrt{2391})^{\frac{1}{3}}-765\right)}{\left(\sqrt{2391}(154+3\sqrt{2391})^{\frac{1}{3}}+4(154+3\sqrt{2391})^{\frac{2}{3}}-4\sqrt{2391}+34(154+3\sqrt{2391})^{\frac{1}{3}}-149\right)\left(\left(154+3\sqrt{2391}\right)^{\frac{2}{3}}-2(154+3\sqrt{2391})^{\frac{1}{3}}+13\right)} \\ -\frac{2t\left(7(154+3\sqrt{2391})^{\frac{2}{3}}-26(154+3\sqrt{2391})^{\frac{1}{3}}-6\sqrt{2391}-308\right)}{\sqrt{2391}(154+3\sqrt{2391})^{\frac{1}{3}}+4(154+3\sqrt{2391})^{\frac{2}{3}}-4\sqrt{2391}+34(154+3\sqrt{2391})^{\frac{1}{3}}-149} \\ t \end{array} \right] =$$

Which is normalized to

$$\left[\begin{array}{c} \frac{3(154+3\sqrt{2391})^{\frac{1}{3}}t\left(\sqrt{2391}(154+3\sqrt{2391})^{\frac{1}{3}}+18(154+3\sqrt{2391})^{\frac{2}{3}}-16\sqrt{2391}-18(154+3\sqrt{2391})^{\frac{1}{3}}-765\right)}{\left(\sqrt{2391}(154+3\sqrt{2391})^{\frac{1}{3}}+4(154+3\sqrt{2391})^{\frac{2}{3}}-4\sqrt{2391}+34(154+3\sqrt{2391})^{\frac{1}{3}}-149\right)\left(\left(154+3\sqrt{2391}\right)^{\frac{2}{3}}-2(154+3\sqrt{2391})^{\frac{1}{3}}+13\right)} \\ -\frac{2t\left(7(154+3\sqrt{2391})^{\frac{2}{3}}-26(154+3\sqrt{2391})^{\frac{1}{3}}-6\sqrt{2391}-308\right)}{\sqrt{2391}(154+3\sqrt{2391})^{\frac{1}{3}}+4(154+3\sqrt{2391})^{\frac{2}{3}}-4\sqrt{2391}+34(154+3\sqrt{2391})^{\frac{1}{3}}-149} \\ t \end{array} \right] =$$

Considering the eigenvalue $\lambda_2 = \frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{6} + \frac{13}{6(154+3\sqrt{2391})^{\frac{1}{3}}} - \frac{1}{3} - \frac{i\sqrt{3}\left(-\frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{3} + \frac{13}{3(154+3\sqrt{2391})^{\frac{1}{3}}}\right)}{2}$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} -1 & 1 & -1 \\ 2 & -1 & -4 \\ 3 & -1 & 1 \end{bmatrix} - \left(\frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{6} + \frac{13}{6(154+3\sqrt{2391})^{\frac{1}{3}}} - \frac{1}{3} - \frac{i\sqrt{3}\left(-\frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{3} + \frac{13}{3(154+3\sqrt{2391})^{\frac{1}{3}}}\right)}{2} \right) I \right) \vec{v} = \vec{0}$$

$$\left[\begin{array}{ccc|c} \frac{(1+i\sqrt{3})(154+3\sqrt{3}\sqrt{797})^{\frac{2}{3}} - 13i\sqrt{3} + 4(154+3\sqrt{3}\sqrt{797})^{\frac{1}{3}} + 13}{6(154+3\sqrt{3}\sqrt{797})^{\frac{1}{3}}} & & & 1 \\ & 2 & & -\frac{(1+i\sqrt{3})(154+3\sqrt{3}\sqrt{797})^{\frac{2}{3}} - 13i\sqrt{3} + 4(154+3\sqrt{3}\sqrt{797})^{\frac{1}{3}} + 13}{6(154+3\sqrt{3}\sqrt{797})^{\frac{1}{3}}} \\ & & 3 & -1 \end{array} \right]$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{ccc|c} -\frac{2}{3} - \frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{6} - \frac{13}{6(154+3\sqrt{2391})^{\frac{1}{3}}} + \frac{i\sqrt{3}\left(-\frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{3} + \frac{13}{3(154+3\sqrt{2391})^{\frac{1}{3}}}\right)}{2} & & & \\ & 2 & & -\frac{2}{3} - \frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{6} - \frac{13}{6(154+3\sqrt{2391})^{\frac{1}{3}}} + \frac{i\sqrt{3}\left(-\frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{3} + \frac{13}{3(154+3\sqrt{2391})^{\frac{1}{3}}}\right)}{2} \\ & & 3 & -1 \end{array} \right]$$

$$R_2 = R_2 - \frac{2R_1}{-\frac{2}{3} - \frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{6} - \frac{13}{6(154+3\sqrt{2391})^{\frac{1}{3}}} + \frac{i\sqrt{3} \left(-\frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{3} + \frac{13}{3(154+3\sqrt{2391})^{\frac{1}{3}}} \right)}{2}} \Rightarrow \left[\begin{array}{c} \frac{(1+i\sqrt{3})(154+3\sqrt{2391})^{\frac{1}{3}}}{6} \\ \dots \end{array} \right]$$

$$R_3 = R_3 + \frac{18(154 + 3\sqrt{3}\sqrt{797})^{\frac{1}{3}} R_1}{(1 + i\sqrt{3})(154 + 3\sqrt{3}\sqrt{797})^{\frac{2}{3}} - 13i\sqrt{3} + 4(154 + 3\sqrt{3}\sqrt{797})^{\frac{1}{3}} + 13} \Rightarrow \left[\begin{array}{c} \frac{(1+i\sqrt{3})(154+3\sqrt{3}\sqrt{797})^{\frac{1}{3}}}{6} \\ \dots \end{array} \right]$$

$$R_3 = R_3 - \frac{\left(-i(154 + 3\sqrt{3}\sqrt{797})^{\frac{2}{3}}\sqrt{3} - (154 + 3\sqrt{3}\sqrt{797})^{\frac{2}{3}} + 13i\sqrt{3} + 14(154 + 3\sqrt{3}\sqrt{797})^{\frac{1}{3}} \right)}{\left(i(154 + 3\sqrt{3}\sqrt{797})^{\frac{2}{3}}\sqrt{3} + (154 + 3\sqrt{3}\sqrt{797})^{\frac{2}{3}} - 13i\sqrt{3} + 4(154 + 3\sqrt{3}\sqrt{797})^{\frac{1}{3}} + 13 \right)} \left(\dots \right)$$

Therefore the system in Echelon form is

$$\left[\begin{array}{cc} \frac{(1+i\sqrt{3})(154+3\sqrt{3}\sqrt{797})^{\frac{2}{3}} - 13i\sqrt{3} + 4(154+3\sqrt{3}\sqrt{797})^{\frac{1}{3}} + 13}{6(154+3\sqrt{3}\sqrt{797})^{\frac{1}{3}}} & 1 \\ 0 & \frac{(-34i\sqrt{3} - 3i\sqrt{797} + \sqrt{2391} + 34)(154+3\sqrt{2391})^{\frac{1}{3}} - 149i\sqrt{3} - 12i\sqrt{797}}{(154+3\sqrt{2391})^{\frac{1}{3}} \left(13 + 4(154+3\sqrt{2391})^{\frac{1}{3}} + i \left((154+3\sqrt{2391})^{\frac{2}{3}} \right) \right)} \\ 0 & 0 \end{array} \right]$$

The free variables are $\{v_3\}$ and the leading variables are $\{v_1, v_2\}$. Let $v_3 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of

free variables gives equation
$$v_1 = -\frac{6(154+3\sqrt{2391})^{\frac{1}{3}}t\left(48i\sqrt{797}-18i\sqrt{3}\left(154+3\sqrt{2391}\right)^{\frac{1}{3}}+3i\sqrt{797}\right)}{\left(i\left(154+3\sqrt{2391}\right)^{\frac{2}{3}}\sqrt{3}+\left(154+3\sqrt{2391}\right)^{\frac{2}{3}}-13i\sqrt{3}+4\left(154+3\sqrt{2391}\right)^{\frac{1}{3}}+13\right)\left(3i\sqrt{797}\left(154+3\sqrt{2391}\right)^{\frac{1}{3}}+34i\left(154+3\sqrt{2391}\right)^{\frac{1}{3}}\sqrt{3}-\sqrt{2391}\left(154+3\sqrt{2391}\right)^{\frac{1}{3}}\right)}$$

Hence the solution is

Expression too large to display

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} \frac{6(154+3\sqrt{2391})^{\frac{1}{3}}t\left(48i\sqrt{797}-18i\sqrt{3}\left(154+3\sqrt{2391}\right)^{\frac{1}{3}}+3i\sqrt{797}\left(154+3\sqrt{2391}\right)^{\frac{1}{3}}-\sqrt{2391}\left(154+3\sqrt{2391}\right)^{\frac{1}{3}}\right)}{\left(i\left(154+3\sqrt{2391}\right)^{\frac{2}{3}}\sqrt{3}+\left(154+3\sqrt{2391}\right)^{\frac{2}{3}}-13i\sqrt{3}+4\left(154+3\sqrt{2391}\right)^{\frac{1}{3}}+13\right)\left(3i\sqrt{797}\left(154+3\sqrt{2391}\right)^{\frac{1}{3}}+34i\left(154+3\sqrt{2391}\right)^{\frac{1}{3}}\sqrt{3}-\sqrt{2391}\left(154+3\sqrt{2391}\right)^{\frac{1}{3}}\right)} \\ -\frac{4t\left(9i\sqrt{797}-13i\left(154+3\sqrt{2391}\right)^{\frac{1}{3}}\sqrt{3}+154i\sqrt{3}+7\left(154+3\sqrt{2391}\right)^{\frac{2}{3}}+13i\sqrt{797}\left(154+3\sqrt{2391}\right)^{\frac{1}{3}}+13i\left(154+3\sqrt{2391}\right)^{\frac{1}{3}}\sqrt{3}-\sqrt{2391}\left(154+3\sqrt{2391}\right)^{\frac{1}{3}}+149i\sqrt{3}+12i\sqrt{797}\left(154+3\sqrt{2391}\right)^{\frac{1}{3}}\right)}{3i\sqrt{797}\left(154+3\sqrt{2391}\right)^{\frac{1}{3}}+34i\left(154+3\sqrt{2391}\right)^{\frac{1}{3}}\sqrt{3}-\sqrt{2391}\left(154+3\sqrt{2391}\right)^{\frac{1}{3}}+149i\sqrt{3}+12i\sqrt{797}\left(154+3\sqrt{2391}\right)^{\frac{1}{3}}} \\ t \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

Expression too large to display

Which is normalized to

Expression too large to display

Considering the eigenvalue $\lambda_3 = \frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{6} + \frac{13}{6(154+3\sqrt{2391})^{\frac{1}{3}}} - \frac{1}{3} + \frac{i\sqrt{3}\left(-\frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{3} + \frac{13}{3(154+3\sqrt{2391})^{\frac{1}{3}}}\right)}{2}$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} -1 & 1 & -1 \\ 2 & -1 & -4 \\ 3 & -1 & 1 \end{bmatrix} - \left(\frac{(154 + 3\sqrt{2391})^{\frac{1}{3}}}{6} + \frac{13}{6(154 + 3\sqrt{2391})^{\frac{1}{3}}} - \frac{1}{3} + \frac{i\sqrt{3}}{3} \left(-\frac{(154 + 3\sqrt{2391})^{\frac{1}{3}}}{3} + \frac{13}{3(154 + 3\sqrt{2391})^{\frac{1}{3}}} \right) \right) \right) \begin{bmatrix} \frac{(i\sqrt{3}-1)(154+3\sqrt{3}\sqrt{797})^{\frac{2}{3}}-13i\sqrt{3}-4(154+3\sqrt{3}\sqrt{797})^{\frac{1}{3}}-13}{6(154+3\sqrt{3}\sqrt{797})^{\frac{1}{3}}} & 1 & \\ 2 & \frac{(i\sqrt{3}-1)(154+3\sqrt{3}\sqrt{797})^{\frac{2}{3}}-13i\sqrt{3}-4(154+3\sqrt{3}\sqrt{797})^{\frac{1}{3}}-13}{6(154+3\sqrt{3}\sqrt{797})^{\frac{1}{3}}} & \\ 3 & -1 & \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{ccc|c} -\frac{2}{3} - \frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{6} - \frac{13}{6(154+3\sqrt{2391})^{\frac{1}{3}}} - \frac{i\sqrt{3}}{2} \left(-\frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{3} + \frac{13}{3(154+3\sqrt{2391})^{\frac{1}{3}}} \right) & & & 1 \\ & & & \\ & & 2 & -\frac{2}{3} - \frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{6} - \frac{1}{6(154+3\sqrt{2391})^{\frac{1}{3}}} \\ & & 3 & \end{array} \right]$$

$$R_2 = R_2 - \frac{2R_1}{\Rightarrow} \left[\begin{array}{ccc|c} -\frac{2}{3} - \frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{6} - \frac{13}{6(154+3\sqrt{2391})^{\frac{1}{3}}} - \frac{i\sqrt{3}}{2} \left(-\frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{3} + \frac{13}{3(154+3\sqrt{2391})^{\frac{1}{3}}} \right) & & & \frac{(i\sqrt{3}-1)(154+3\sqrt{3}\sqrt{797})^{\frac{2}{3}}-13i\sqrt{3}-4(154+3\sqrt{3}\sqrt{797})^{\frac{1}{3}}-13}{6(154+3\sqrt{3}\sqrt{797})^{\frac{1}{3}}} \end{array} \right]$$

$$R_3 = R_3 - \frac{18(154 + 3\sqrt{3}\sqrt{797})^{\frac{1}{3}} R_1}{(i\sqrt{3} - 1)(154 + 3\sqrt{3}\sqrt{797})^{\frac{2}{3}} - 13i\sqrt{3} - 4(154 + 3\sqrt{3}\sqrt{797})^{\frac{1}{3}} - 13} \Rightarrow \left[\begin{array}{c} (i\sqrt{3}-1)(154+3\sqrt{3}\sqrt{797})^{\frac{2}{3}} - 13i\sqrt{3} - 4(154+3\sqrt{3}\sqrt{797})^{\frac{1}{3}} - 13 \\ \vdots \\ \vdots \end{array} \right]$$

$$R_3 = R_3 - \frac{\left(-i(154 + 3\sqrt{3}\sqrt{797})^{\frac{2}{3}}\sqrt{3} + (154 + 3\sqrt{3}\sqrt{797})^{\frac{2}{3}} + 13i\sqrt{3} - 14(154 + 3\sqrt{3}\sqrt{797})^{\frac{1}{3}} + 13\right) R_1}{\left(i(154 + 3\sqrt{3}\sqrt{797})^{\frac{2}{3}}\sqrt{3} - (154 + 3\sqrt{3}\sqrt{797})^{\frac{2}{3}} - 13i\sqrt{3} - 4(154 + 3\sqrt{3}\sqrt{797})^{\frac{1}{3}} - 13\right) R_1}$$

Therefore the system in Echelon form is

$$\left[\begin{array}{ccc} \frac{(i\sqrt{3}-1)(154+3\sqrt{3}\sqrt{797})^{\frac{2}{3}} - 13i\sqrt{3} - 4(154+3\sqrt{3}\sqrt{797})^{\frac{1}{3}} - 13}{6(154+3\sqrt{3}\sqrt{797})^{\frac{1}{3}}} & & 1 \\ 0 & & \frac{-(34i\sqrt{3}+3i\sqrt{797}+\sqrt{2391}+34)(154+3\sqrt{2391})^{\frac{1}{3}} - 149i\sqrt{3} - 12i\sqrt{797} + 13}{(-13-4(154+3\sqrt{2391})^{\frac{1}{3}}+i((154+3\sqrt{2391})^{\frac{2}{3}}-13))\sqrt{3} - (154+3\sqrt{2391})^{\frac{1}{3}}} \\ 0 & & 0 \end{array} \right]$$

The free variables are $\{v_3\}$ and the leading variables are $\{v_1, v_2\}$. Let $v_3 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of

free variables gives equation $\left\{ \begin{array}{l} v_1 = \frac{6(154+3\sqrt{2391})^{\frac{1}{3}} t \left(18i\sqrt{3}(154+3\sqrt{2391})^{\frac{1}{3}} - 3i\sqrt{797}(154+3\sqrt{2391})^{\frac{1}{3}} + 13\right)}{\left(i(154+3\sqrt{2391})^{\frac{2}{3}}\sqrt{3} - (154+3\sqrt{2391})^{\frac{2}{3}} - 13i\sqrt{3} - 4(154+3\sqrt{2391})^{\frac{1}{3}} - 13\right) \left(-34i\sqrt{3}(154+3\sqrt{2391})^{\frac{1}{3}} + (154+3\sqrt{2391})^{\frac{2}{3}} + 13i\sqrt{3} - 14(154+3\sqrt{2391})^{\frac{1}{3}} + 13\right)} \end{array} \right.$

Hence the solution is

Expression too large to display

Since there is one free Variable, we have found one eigenvector associated with this

eigenvalue. The above can be written as

$$\left[\frac{6(154+3\sqrt{2391})^{\frac{1}{3}} t \left(18 I \sqrt{3} (154+3\sqrt{2391})^{\frac{1}{3}} - 3 I (154+3\sqrt{2391})^{\frac{1}{3}} \sqrt{797-\sqrt{2391}} (154+3\sqrt{2391})^{\frac{1}{3}} + 36 (154+3\sqrt{2391})^{\frac{1}{3}} \right)}{\left(I (154+3\sqrt{2391})^{\frac{2}{3}} \sqrt{3} - (154+3\sqrt{2391})^{\frac{2}{3}} - 13 I \sqrt{3} - 4 (154+3\sqrt{2391})^{\frac{1}{3}} - 13 \right) \left(-34 I (154+3\sqrt{2391})^{\frac{1}{3}} \sqrt{3} - 3 I (154+3\sqrt{2391})^{\frac{1}{3}} \sqrt{797-\sqrt{2391}} + 36 (154+3\sqrt{2391})^{\frac{1}{3}} \right)} - \frac{4t \left(13 I (154+3\sqrt{2391})^{\frac{1}{3}} \sqrt{3} + 7 (154+3\sqrt{2391})^{\frac{2}{3}} - 154 I \sqrt{3} - 9 I \sqrt{797-\sqrt{2391}} + 36 (154+3\sqrt{2391})^{\frac{1}{3}} \right)}{-34 I (154+3\sqrt{2391})^{\frac{1}{3}} \sqrt{3} - 3 I (154+3\sqrt{2391})^{\frac{1}{3}} \sqrt{797-\sqrt{2391}} (154+3\sqrt{2391})^{\frac{1}{3}} + 8 (154+3\sqrt{2391})^{\frac{1}{3}}} \right] t$$

Let $t = 1$ the eigenvector becomes

Expression too large to display

Which is normalized to

Expression too large to display

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		de
	algebraic m	geometric k	
$-\frac{\left(154+3\sqrt{2391}\right)^{\frac{1}{3}}}{3} - \frac{13}{3\left(154+3\sqrt{2391}\right)^{\frac{1}{3}}} - \frac{1}{3}$	1	1	
$\frac{\left(154+3\sqrt{2391}\right)^{\frac{1}{3}}}{6} + \frac{13}{6\left(154+3\sqrt{2391}\right)^{\frac{1}{3}}} - \frac{1}{3} + \frac{i\sqrt{3}\left(-\frac{\left(154+3\sqrt{2391}\right)^{\frac{1}{3}}}{3} + \frac{13}{3\left(154+3\sqrt{2391}\right)^{\frac{1}{3}}}\right)}{2}$	1	1	
$\frac{\left(154+3\sqrt{2391}\right)^{\frac{1}{3}}}{6} + \frac{13}{6\left(154+3\sqrt{2391}\right)^{\frac{1}{3}}} - \frac{1}{3} - \frac{i\sqrt{3}\left(-\frac{\left(154+3\sqrt{2391}\right)^{\frac{1}{3}}}{3} + \frac{13}{3\left(154+3\sqrt{2391}\right)^{\frac{1}{3}}}\right)}{4182 \cdot 2}$	1	1	

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. Since eigenvalue $-\frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{3} - \frac{13}{3(154+3\sqrt{2391})^{\frac{1}{3}}} - \frac{1}{3}$ is real and distinct then the corresponding eigenvector solution is

$$\vec{x}_1(t) = \vec{v}_1 e^{\left(-\frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{3} - \frac{13}{3(154+3\sqrt{2391})^{\frac{1}{3}}} - \frac{1}{3}\right)t}$$

$$= \begin{bmatrix} 7\left(-\frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{3} - \frac{13}{3(154+3\sqrt{2391})^{\frac{1}{3}}} + \frac{14}{3}\right) \\ -\frac{\left(-\frac{7(154+3\sqrt{2391})^{\frac{1}{3}}}{3} - \frac{91}{3(154+3\sqrt{2391})^{\frac{1}{3}}} + \frac{2}{3} + \left(-\frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{3} - \frac{13}{3(154+3\sqrt{2391})^{\frac{1}{3}}} - \frac{1}{3}\right)^2\right)\left(-\frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{3} - \frac{13}{3(154+3\sqrt{2391})^{\frac{1}{3}}}\right)}{\left(-\frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{3} - \frac{13}{3(154+3\sqrt{2391})^{\frac{1}{3}}} - \frac{1}{3}\right)^2 - 32} \\ \frac{7(154+3\sqrt{2391})^{\frac{1}{3}}}{3} - \frac{91}{3(154+3\sqrt{2391})^{\frac{1}{3}}} + \frac{2}{3} + \left(-\frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{3} - \frac{13}{3(154+3\sqrt{2391})^{\frac{1}{3}}} - \frac{1}{3}\right)^2 \\ 1 \end{bmatrix}$$

Therefore the final solution is

$$\vec{x}_h(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t) + c_3 \vec{x}_3(t)$$

Which is written as

$$\begin{bmatrix} x(t) \\ y(t) \\ z(t) \end{bmatrix} = c_1 \begin{bmatrix} 7e^{\left(-\frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{3} - \frac{13}{3(154+3\sqrt{2391})^{\frac{1}{3}}} - \frac{1}{3}\right)t} \left(-\frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{3} - \frac{13}{3(154+3\sqrt{2391})^{\frac{1}{3}}} + \frac{14}{3}\right) \\ -\frac{\left(-\frac{7(154+3\sqrt{2391})^{\frac{1}{3}}}{3} - \frac{91}{3(154+3\sqrt{2391})^{\frac{1}{3}}} + \frac{2}{3} + \left(-\frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{3} - \frac{13}{3(154+3\sqrt{2391})^{\frac{1}{3}}} - \frac{1}{3}\right)^2\right)\left(-\frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{3} - \frac{13}{3(154+3\sqrt{2391})^{\frac{1}{3}}}\right)}{\left(-\frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{3} - \frac{13}{3(154+3\sqrt{2391})^{\frac{1}{3}}} - \frac{1}{3}\right)^2 - 32} \\ e^{\left(-\frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{3} - \frac{13}{3(154+3\sqrt{2391})^{\frac{1}{3}}} - \frac{1}{3}\right)t} \left(\left(-\frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{3} - \frac{13}{3(154+3\sqrt{2391})^{\frac{1}{3}}} - \frac{1}{3}\right)^2 - 32\right) \\ -\frac{7(154+3\sqrt{2391})^{\frac{1}{3}}}{3} - \frac{91}{3(154+3\sqrt{2391})^{\frac{1}{3}}} + \frac{2}{3} + \left(-\frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{3} - \frac{13}{3(154+3\sqrt{2391})^{\frac{1}{3}}} - \frac{1}{3}\right)^2 \\ e^{\left(-\frac{(154+3\sqrt{2391})^{\frac{1}{3}}}{3} - \frac{13}{3(154+3\sqrt{2391})^{\frac{1}{3}}} - \frac{1}{3}\right)t} \end{bmatrix}$$

Which becomes

Expression too large to display

✓ Solution by Maple

Time used: 0.141 (sec). Leaf size: 3197

`dsolve([diff(x(t),t)=-x(t)+y(t)-z(t),diff(y(t),t)=2*x(t)-y(t)-4*z(t),diff(z(t),t)=3*x(t)-y(t)`

$$\begin{aligned}
 x(t) = & c_2 e^{\frac{\left(13 + (154 + 3\sqrt{2391})^{\frac{2}{3}} - 2(154 + 3\sqrt{2391})^{\frac{1}{3}}\right)t}{6(154 + 3\sqrt{2391})^{\frac{1}{3}}}} \sin\left(\frac{\sqrt{3}\left((154 + 3\sqrt{2391})^{\frac{2}{3}} - 13\right)t}{6(154 + 3\sqrt{2391})^{\frac{1}{3}}}\right) \\
 & + c_3 e^{\frac{\left(13 + (154 + 3\sqrt{2391})^{\frac{2}{3}} - 2(154 + 3\sqrt{2391})^{\frac{1}{3}}\right)t}{6(154 + 3\sqrt{2391})^{\frac{1}{3}}}} \cos\left(\frac{\sqrt{3}\left((154 + 3\sqrt{2391})^{\frac{2}{3}} - 13\right)t}{6(154 + 3\sqrt{2391})^{\frac{1}{3}}}\right) \\
 & + c_1 e^{-\frac{\left((154 + 3\sqrt{2391})^{\frac{2}{3}} + (154 + 3\sqrt{2391})^{\frac{1}{3}} + 13\right)t}{3(154 + 3\sqrt{2391})^{\frac{1}{3}}}}
 \end{aligned}$$

Expression too large to display
 Expression too large to display

✓ Solution by Mathematica

Time used: 0.032 (sec). Leaf size: 501

`DSolve[{x'[t]==-x[t]+y[t]-z[t],y'[t]==2*x[t]-y[t]-4*z[t],z'[t]==3*x[t]-y[t]+z[t]},{x[t],y[t]}`

$$\begin{aligned}
 x(t) &\rightarrow c_2 \text{RootSum} \left[\#1^3 + \#1^2 - 4\#1 + 10 \&, \frac{\#1 e^{\#1 t}}{3\#1^2 + 2\#1 - 4} \& \right] \\
 &\quad - c_3 \text{RootSum} \left[\#1^3 + \#1^2 - 4\#1 + 10 \&, \frac{\#1 e^{\#1 t} + 5e^{\#1 t}}{3\#1^2 + 2\#1 - 4} \& \right] \\
 &\quad + c_1 \text{RootSum} \left[\#1^3 + \#1^2 - 4\#1 + 10 \&, \frac{\#1^2 e^{\#1 t} - 5e^{\#1 t}}{3\#1^2 + 2\#1 - 4} \& \right] \\
 y(t) &\rightarrow 2c_1 \text{RootSum} \left[\#1^3 + \#1^2 - 4\#1 + 10 \&, \frac{\#1 e^{\#1 t} - 7e^{\#1 t}}{3\#1^2 + 2\#1 - 4} \& \right] \\
 &\quad - 2c_3 \text{RootSum} \left[\#1^3 + \#1^2 - 4\#1 + 10 \&, \frac{2\#1 e^{\#1 t} + 3e^{\#1 t}}{3\#1^2 + 2\#1 - 4} \& \right] \\
 &\quad + c_2 \text{RootSum} \left[\#1^3 + \#1^2 - 4\#1 + 10 \&, \frac{\#1^2 e^{\#1 t} + 2e^{\#1 t}}{3\#1^2 + 2\#1 - 4} \& \right] \\
 z(t) &\rightarrow -c_2 \text{RootSum} \left[\#1^3 + \#1^2 - 4\#1 + 10 \&, \frac{\#1 e^{\#1 t} - 2e^{\#1 t}}{3\#1^2 + 2\#1 - 4} \& \right] \\
 &\quad + c_1 \text{RootSum} \left[\#1^3 + \#1^2 - 4\#1 + 10 \&, \frac{3\#1 e^{\#1 t} + e^{\#1 t}}{3\#1^2 + 2\#1 - 4} \& \right] \\
 &\quad + c_3 \text{RootSum} \left[\#1^3 + \#1^2 - 4\#1 + 10 \&, \frac{\#1^2 e^{\#1 t} + 2\#1 e^{\#1 t} - e^{\#1 t}}{3\#1^2 + 2\#1 - 4} \& \right]
 \end{aligned}$$

29.11 problem 4(a)

29.11.1 Solution using Matrix exponential method	4186
29.11.2 Solution using explicit Eigenvalue and Eigenvector method . . .	4188
29.11.3 Maple step by step solution	4194

Internal problem ID [6539]

Internal file name [OUTPUT/5787_Sunday_June_05_2022_03_54_27_PM_84229463/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 10. Systems of First-Order Equations. Section A. Drill exercises. Page 400

Problem number: 4(a).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "system of linear ODEs"

Solve

$$\begin{aligned}x'(t) &= x(t) + 2y(t) - 4t + 1 \\y'(t) &= -x(t) + 2y(t) + 3t + 4\end{aligned}$$

29.11.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t) + \vec{G}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix} + \begin{bmatrix} 1 - 4t \\ 3t + 4 \end{bmatrix}$$

Since the system is nonhomogeneous, then the solution is given by

$$\vec{x}(t) = \vec{x}_h(t) + \vec{x}_p(t)$$

Where $\vec{x}_h(t)$ is the homogeneous solution to $\vec{x}'(t) = A\vec{x}(t)$ and $\vec{x}_p(t)$ is a particular solution to $\vec{x}'(t) = A\vec{x}(t) + \vec{G}(t)$. The particular solution will be found using variation

of parameters method applied to the fundamental matrix. For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} e^{\frac{3t}{2}} \cos\left(\frac{\sqrt{7}t}{2}\right) - \frac{\sqrt{7}e^{\frac{3t}{2}} \sin\left(\frac{\sqrt{7}t}{2}\right)}{7} & \frac{4\sqrt{7}e^{\frac{3t}{2}} \sin\left(\frac{\sqrt{7}t}{2}\right)}{7} \\ -\frac{2\sqrt{7}e^{\frac{3t}{2}} \sin\left(\frac{\sqrt{7}t}{2}\right)}{7} & e^{\frac{3t}{2}} \cos\left(\frac{\sqrt{7}t}{2}\right) + \frac{\sqrt{7}e^{\frac{3t}{2}} \sin\left(\frac{\sqrt{7}t}{2}\right)}{7} \end{bmatrix}$$

$$= \begin{bmatrix} -\frac{e^{\frac{3t}{2}} \left(\sqrt{7} \sin\left(\frac{\sqrt{7}t}{2}\right) - 7 \cos\left(\frac{\sqrt{7}t}{2}\right)\right)}{7} & \frac{4\sqrt{7}e^{\frac{3t}{2}} \sin\left(\frac{\sqrt{7}t}{2}\right)}{7} \\ -\frac{2\sqrt{7}e^{\frac{3t}{2}} \sin\left(\frac{\sqrt{7}t}{2}\right)}{7} & \frac{e^{\frac{3t}{2}} \left(\sqrt{7} \sin\left(\frac{\sqrt{7}t}{2}\right) + 7 \cos\left(\frac{\sqrt{7}t}{2}\right)\right)}{7} \end{bmatrix}$$

Therefore the homogeneous solution is

$$\vec{x}_h(t) = e^{At} \vec{c}$$

$$= \begin{bmatrix} -\frac{e^{\frac{3t}{2}} \left(\sqrt{7} \sin\left(\frac{\sqrt{7}t}{2}\right) - 7 \cos\left(\frac{\sqrt{7}t}{2}\right)\right)}{7} & \frac{4\sqrt{7}e^{\frac{3t}{2}} \sin\left(\frac{\sqrt{7}t}{2}\right)}{7} \\ -\frac{2\sqrt{7}e^{\frac{3t}{2}} \sin\left(\frac{\sqrt{7}t}{2}\right)}{7} & \frac{e^{\frac{3t}{2}} \left(\sqrt{7} \sin\left(\frac{\sqrt{7}t}{2}\right) + 7 \cos\left(\frac{\sqrt{7}t}{2}\right)\right)}{7} \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix}$$

$$= \begin{bmatrix} -\frac{e^{\frac{3t}{2}} \left(\sqrt{7} \sin\left(\frac{\sqrt{7}t}{2}\right) - 7 \cos\left(\frac{\sqrt{7}t}{2}\right)\right) c_1}{7} + \frac{4\sqrt{7}e^{\frac{3t}{2}} \sin\left(\frac{\sqrt{7}t}{2}\right) c_2}{7} \\ -\frac{2\sqrt{7}e^{\frac{3t}{2}} \sin\left(\frac{\sqrt{7}t}{2}\right) c_1}{7} + \frac{e^{\frac{3t}{2}} \left(\sqrt{7} \sin\left(\frac{\sqrt{7}t}{2}\right) + 7 \cos\left(\frac{\sqrt{7}t}{2}\right)\right) c_2}{7} \end{bmatrix}$$

$$= \begin{bmatrix} -\frac{\left(\sqrt{7}(c_1 - 4c_2) \sin\left(\frac{\sqrt{7}t}{2}\right) - 7 \cos\left(\frac{\sqrt{7}t}{2}\right) c_1\right) e^{\frac{3t}{2}}}{7} \\ -\frac{2\left(\sqrt{7}(c_1 - \frac{c_2}{2}) \sin\left(\frac{\sqrt{7}t}{2}\right) - \frac{7 \cos\left(\frac{\sqrt{7}t}{2}\right) c_2}{2}\right) e^{\frac{3t}{2}}}{7} \end{bmatrix}$$

The particular solution given by

$$\vec{x}_p(t) = e^{At} \int e^{-At} \vec{G}(t) dt$$

But

$$e^{-At} = (e^{At})^{-1}$$

$$= \begin{bmatrix} \frac{\left(\sqrt{7} \sin\left(\frac{\sqrt{7}t}{2}\right) + 7 \cos\left(\frac{\sqrt{7}t}{2}\right)\right) e^{-\frac{3t}{2}}}{7} & -\frac{4\sqrt{7}e^{-\frac{3t}{2}} \sin\left(\frac{\sqrt{7}t}{2}\right)}{7} \\ \frac{2\sqrt{7}e^{-\frac{3t}{2}} \sin\left(\frac{\sqrt{7}t}{2}\right)}{7} & -\frac{\left(\sqrt{7} \sin\left(\frac{\sqrt{7}t}{2}\right) - 7 \cos\left(\frac{\sqrt{7}t}{2}\right)\right) e^{-\frac{3t}{2}}}{7} \end{bmatrix}$$

Hence

$$\begin{aligned}
 \vec{x}_p(t) &= \begin{bmatrix} -\frac{e^{\frac{3t}{2}}(\sqrt{7}\sin(\frac{\sqrt{7}t}{2})-7\cos(\frac{\sqrt{7}t}{2}))}{7} & \frac{4\sqrt{7}e^{\frac{3t}{2}}\sin(\frac{\sqrt{7}t}{2})}{7} \\ -\frac{2\sqrt{7}e^{\frac{3t}{2}}\sin(\frac{\sqrt{7}t}{2})}{7} & \frac{e^{\frac{3t}{2}}(\sqrt{7}\sin(\frac{\sqrt{7}t}{2})+7\cos(\frac{\sqrt{7}t}{2}))}{7} \end{bmatrix} \int \begin{bmatrix} \frac{(\sqrt{7}\sin(\frac{\sqrt{7}t}{2})+7\cos(\frac{\sqrt{7}t}{2}))e^{-\frac{3t}{2}}}{7} \\ \frac{2\sqrt{7}e^{-\frac{3t}{2}}\sin(\frac{\sqrt{7}t}{2})}{7} \end{bmatrix} \\
 &= \begin{bmatrix} -\frac{e^{\frac{3t}{2}}(\sqrt{7}\sin(\frac{\sqrt{7}t}{2})-7\cos(\frac{\sqrt{7}t}{2}))}{7} & \frac{4\sqrt{7}e^{\frac{3t}{2}}\sin(\frac{\sqrt{7}t}{2})}{7} \\ -\frac{2\sqrt{7}e^{\frac{3t}{2}}\sin(\frac{\sqrt{7}t}{2})}{7} & \frac{e^{\frac{3t}{2}}(\sqrt{7}\sin(\frac{\sqrt{7}t}{2})+7\cos(\frac{\sqrt{7}t}{2}))}{7} \end{bmatrix} \begin{bmatrix} \frac{5((\frac{49t}{5}+\frac{35}{4})\cos(\frac{\sqrt{7}t}{2})+\sqrt{7}\sin(\frac{\sqrt{7}t}{2}))(t+\frac{35}{4})}{14} \\ \frac{e^{-\frac{3t}{2}}(28t-35)\cos(\frac{\sqrt{7}t}{2})}{112} + \frac{27e^{-\frac{3t}{2}}(t+\frac{35}{36})\sqrt{7}}{28} \end{bmatrix} \\
 &= \begin{bmatrix} \frac{7t}{2} + \frac{25}{8} \\ \frac{t}{4} - \frac{5}{16} \end{bmatrix}
 \end{aligned}$$

Hence the complete solution is

$$\begin{aligned}
 \vec{x}(t) &= \vec{x}_h(t) + \vec{x}_p(t) \\
 &= \begin{bmatrix} -\frac{e^{\frac{3t}{2}}\sqrt{7}(c_1-4c_2)\sin(\frac{\sqrt{7}t}{2})}{7} + e^{\frac{3t}{2}}\cos(\frac{\sqrt{7}t}{2})c_1 + \frac{7t}{2} + \frac{25}{8} \\ -\frac{2\sqrt{7}(c_1-\frac{c_2}{2})e^{\frac{3t}{2}}\sin(\frac{\sqrt{7}t}{2})}{7} + e^{\frac{3t}{2}}\cos(\frac{\sqrt{7}t}{2})c_2 + \frac{t}{4} - \frac{5}{16} \end{bmatrix}
 \end{aligned}$$

29.11.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t) + \vec{G}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix} + \begin{bmatrix} 1-4t \\ 3t+4 \end{bmatrix}$$

Since the system is nonhomogeneous, then the solution is given by

$$\vec{x}(t) = \vec{x}_h(t) + \vec{x}_p(t)$$

Where $\vec{x}_h(t)$ is the homogeneous solution to $\vec{x}'(t) = A\vec{x}(t)$ and $\vec{x}_p(t)$ is a particular solution to $\vec{x}'(t) = A\vec{x}(t) + \vec{G}(t)$. The particular solution will be found using variation of parameters method applied to the fundamental matrix.

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det \left(\begin{bmatrix} 1 & 2 \\ -1 & 2 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) = 0$$

Therefore

$$\det \left(\begin{bmatrix} 1 - \lambda & 2 \\ -1 & 2 - \lambda \end{bmatrix} \right) = 0$$

Which gives the characteristic equation

$$\lambda^2 - 3\lambda + 4 = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = \frac{3}{2} + \frac{i\sqrt{7}}{2}$$

$$\lambda_2 = \frac{3}{2} - \frac{i\sqrt{7}}{2}$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
$\frac{3}{2} + \frac{i\sqrt{7}}{2}$	1	complex eigenvalue
$\frac{3}{2} - \frac{i\sqrt{7}}{2}$	1	complex eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = \frac{3}{2} - \frac{i\sqrt{7}}{2}$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 1 & 2 \\ -1 & 2 \end{bmatrix} - \left(\frac{3}{2} - \frac{i\sqrt{7}}{2} \right) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} -\frac{1}{2} + \frac{i\sqrt{7}}{2} & 2 \\ -1 & \frac{i\sqrt{7}}{2} + \frac{1}{2} \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} -\frac{1}{2} + \frac{i\sqrt{7}}{2} & 2 & 0 \\ -1 & \frac{i\sqrt{7}}{2} + \frac{1}{2} & 0 \end{array} \right]$$

$$R_2 = R_2 + \frac{R_1}{-\frac{1}{2} + \frac{i\sqrt{7}}{2}} \implies \left[\begin{array}{cc|c} -\frac{1}{2} + \frac{i\sqrt{7}}{2} & 2 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\left[\begin{array}{cc} -\frac{1}{2} + \frac{i\sqrt{7}}{2} & 2 \\ 0 & 0 \end{array} \right] \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\left\{ v_1 = -\frac{4t}{-1+i\sqrt{7}} \right\}$

Hence the solution is

$$\begin{bmatrix} -\frac{4t}{-1+i\sqrt{7}} \\ t \end{bmatrix} = \begin{bmatrix} -\frac{4t}{-1+i\sqrt{7}} \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} -\frac{4t}{-1+i\sqrt{7}} \\ t \end{bmatrix} = t \begin{bmatrix} -\frac{4}{-1+i\sqrt{7}} \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} -\frac{4}{-1+i\sqrt{7}} \\ 1 \end{bmatrix} = \begin{bmatrix} -\frac{4}{-1+i\sqrt{7}} \\ 1 \end{bmatrix}$$

Which is normalized to

$$\begin{bmatrix} -\frac{4}{-1+i\sqrt{7}} \\ 1 \end{bmatrix} = \begin{bmatrix} -\frac{4}{-1+i\sqrt{7}} \\ 1 \end{bmatrix}$$

Considering the eigenvalue $\lambda_2 = \frac{3}{2} + \frac{i\sqrt{7}}{2}$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\left(\begin{bmatrix} 1 & 2 \\ -1 & 2 \end{bmatrix} - \left(\frac{3}{2} + \frac{i\sqrt{7}}{2} \right) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \right)$$

$$\begin{bmatrix} -\frac{1}{2} - \frac{i\sqrt{7}}{2} & 2 \\ -1 & \frac{1}{2} - \frac{i\sqrt{7}}{2} \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} -\frac{1}{2} - \frac{i\sqrt{7}}{2} & 2 & 0 \\ -1 & \frac{1}{2} - \frac{i\sqrt{7}}{2} & 0 \end{array} \right]$$

$$R_2 = R_2 + \frac{R_1}{-\frac{1}{2} - \frac{i\sqrt{7}}{2}} \implies \left[\begin{array}{cc|c} -\frac{1}{2} - \frac{i\sqrt{7}}{2} & 2 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\left[\begin{array}{cc} -\frac{1}{2} - \frac{i\sqrt{7}}{2} & 2 \\ 0 & 0 \end{array} \right] \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\left\{ v_1 = \frac{4t}{1+i\sqrt{7}} \right\}$

Hence the solution is

$$\begin{bmatrix} \frac{4t}{1+i\sqrt{7}} \\ t \end{bmatrix} = \begin{bmatrix} \frac{4t}{1+i\sqrt{7}} \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} \frac{4t}{1+i\sqrt{7}} \\ t \end{bmatrix} = t \begin{bmatrix} \frac{4}{1+i\sqrt{7}} \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} \frac{4t}{1+i\sqrt{7}} \\ t \end{bmatrix} = \begin{bmatrix} \frac{4}{1+i\sqrt{7}} \\ 1 \end{bmatrix}$$

Which is normalized to

$$\begin{bmatrix} \frac{4t}{1+i\sqrt{7}} \\ t \end{bmatrix} = \begin{bmatrix} \frac{4}{1+i\sqrt{7}} \\ 1 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number

of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
$\frac{3}{2} + \frac{i\sqrt{7}}{2}$	1	1	No	$\begin{bmatrix} \frac{2}{\frac{i\sqrt{7}}{2} + \frac{1}{2}} \\ 1 \end{bmatrix}$
$\frac{3}{2} - \frac{i\sqrt{7}}{2}$	1	1	No	$\begin{bmatrix} \frac{2}{\frac{1}{2} - \frac{i\sqrt{7}}{2}} \\ 1 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. Therefore the homogeneous solution is

$$\vec{x}_h(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

Which is written as

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = c_1 \begin{bmatrix} \frac{2e^{\left(\frac{3}{2} + \frac{i\sqrt{7}}{2}\right)t}}{\frac{i\sqrt{7}}{2} + \frac{1}{2}} \\ e^{\left(\frac{3}{2} + \frac{i\sqrt{7}}{2}\right)t} \end{bmatrix} + c_2 \begin{bmatrix} \frac{2e^{\left(\frac{3}{2} - \frac{i\sqrt{7}}{2}\right)t}}{\frac{1}{2} - \frac{i\sqrt{7}}{2}} \\ e^{\left(\frac{3}{2} - \frac{i\sqrt{7}}{2}\right)t} \end{bmatrix}$$

Now that we found homogeneous solution above, we need to find a particular solution $\vec{x}_p(t)$. We will use Variation of parameters. The fundamental matrix is

$$\Phi = \begin{bmatrix} \vec{x}_1 & \vec{x}_2 & \dots \end{bmatrix}$$

Where \vec{x}_i are the solution basis found above. Therefore the fundamental matrix is

$$\Phi(t) = \begin{bmatrix} \frac{2e^{\left(\frac{3}{2} + \frac{i\sqrt{7}}{2}\right)t}}{\frac{i\sqrt{7}}{2} + \frac{1}{2}} & \frac{2e^{\left(\frac{3}{2} - \frac{i\sqrt{7}}{2}\right)t}}{\frac{1}{2} - \frac{i\sqrt{7}}{2}} \\ e^{\left(\frac{3}{2} + \frac{i\sqrt{7}}{2}\right)t} & e^{\left(\frac{3}{2} - \frac{i\sqrt{7}}{2}\right)t} \end{bmatrix}$$

The particular solution is then given by

$$\vec{x}_p(t) = \Phi \int \Phi^{-1} \vec{G}(t) dt$$

But

$$\Phi^{-1} = \begin{bmatrix} \frac{i\sqrt{7}e^{-\frac{(3+i\sqrt{7})t}{2}}}{7} & -\frac{\sqrt{7}e^{-\frac{(3+i\sqrt{7})t}{2}}(i-\sqrt{7})}{14} \\ -\frac{i\sqrt{7}e^{\frac{(i\sqrt{7}-3)t}{2}}}{7} & \frac{\sqrt{7}e^{\frac{(i\sqrt{7}-3)t}{2}}(i+\sqrt{7})}{14} \end{bmatrix}$$

Hence

$$\begin{aligned} \vec{x}_p(t) &= \begin{bmatrix} \frac{2e^{\left(\frac{3}{2}+\frac{i\sqrt{7}}{2}\right)t}}{\frac{i\sqrt{7}}{2}+\frac{1}{2}} & \frac{2e^{\left(\frac{3}{2}-\frac{i\sqrt{7}}{2}\right)t}}{\frac{1}{2}-\frac{i\sqrt{7}}{2}} \\ e^{\left(\frac{3}{2}+\frac{i\sqrt{7}}{2}\right)t} & e^{\left(\frac{3}{2}-\frac{i\sqrt{7}}{2}\right)t} \end{bmatrix} \int \begin{bmatrix} \frac{i\sqrt{7}e^{-\frac{(3+i\sqrt{7})t}{2}}}{7} & -\frac{\sqrt{7}e^{-\frac{(3+i\sqrt{7})t}{2}}(i-\sqrt{7})}{14} \\ -\frac{i\sqrt{7}e^{\frac{(i\sqrt{7}-3)t}{2}}}{7} & \frac{\sqrt{7}e^{\frac{(i\sqrt{7}-3)t}{2}}(i+\sqrt{7})}{14} \end{bmatrix} \begin{bmatrix} 1-4t \\ 3t+4 \end{bmatrix} dt \\ &= \begin{bmatrix} \frac{2e^{\left(\frac{3}{2}+\frac{i\sqrt{7}}{2}\right)t}}{\frac{i\sqrt{7}}{2}+\frac{1}{2}} & \frac{2e^{\left(\frac{3}{2}-\frac{i\sqrt{7}}{2}\right)t}}{\frac{1}{2}-\frac{i\sqrt{7}}{2}} \\ e^{\left(\frac{3}{2}+\frac{i\sqrt{7}}{2}\right)t} & e^{\left(\frac{3}{2}-\frac{i\sqrt{7}}{2}\right)t} \end{bmatrix} \int \begin{bmatrix} \frac{e^{-\frac{(3+i\sqrt{7})t}{2}}(28+i(-2-11t)\sqrt{7}+21t)}{14} \\ \frac{e^{\frac{(i\sqrt{7}-3)t}{2}}(28+i(11t+2)\sqrt{7}+21t)}{14} \end{bmatrix} dt \\ &= \begin{bmatrix} \frac{2e^{\left(\frac{3}{2}+\frac{i\sqrt{7}}{2}\right)t}}{\frac{i\sqrt{7}}{2}+\frac{1}{2}} & \frac{2e^{\left(\frac{3}{2}-\frac{i\sqrt{7}}{2}\right)t}}{\frac{1}{2}-\frac{i\sqrt{7}}{2}} \\ e^{\left(\frac{3}{2}+\frac{i\sqrt{7}}{2}\right)t} & e^{\left(\frac{3}{2}-\frac{i\sqrt{7}}{2}\right)t} \end{bmatrix} \begin{bmatrix} \frac{\sqrt{7}e^{-\frac{(3+i\sqrt{7})t}{2}}(92\sqrt{7}t+65\sqrt{7}+276it+315i)(11t+3i\sqrt{7}t+2+4i\sqrt{7})}{2128i\sqrt{7}+10304t+5936} \\ \frac{(92i\sqrt{7}t+65i\sqrt{7}+276t+315)\sqrt{7}e^{\frac{(i\sqrt{7}-3)t}{2}}(11it+3\sqrt{7}t+2i+4\sqrt{7})}{2128i\sqrt{7}-10304t-5936} \end{bmatrix} \\ &= \begin{bmatrix} \frac{1288t^3+2634t^2+2137t+725}{368t^2+424t+232} \\ \frac{184t^3-18t^2-149t-145}{736t^2+848t+464} \end{bmatrix} \end{aligned}$$

Now that we found particular solution, the final solution is

$$\begin{aligned} \vec{x}(t) &= \vec{x}_h(t) + \vec{x}_p(t) \\ \begin{bmatrix} x(t) \\ y(t) \end{bmatrix} &= \begin{bmatrix} \frac{2c_1e^{\left(\frac{3}{2}+\frac{i\sqrt{7}}{2}\right)t}}{\frac{i\sqrt{7}}{2}+\frac{1}{2}} \\ c_1e^{\left(\frac{3}{2}+\frac{i\sqrt{7}}{2}\right)t} \end{bmatrix} + \begin{bmatrix} \frac{2c_2e^{\left(\frac{3}{2}-\frac{i\sqrt{7}}{2}\right)t}}{\frac{1}{2}-\frac{i\sqrt{7}}{2}} \\ c_2e^{\left(\frac{3}{2}-\frac{i\sqrt{7}}{2}\right)t} \end{bmatrix} + \begin{bmatrix} \frac{1288t^3+2634t^2+2137t+725}{368t^2+424t+232} \\ \frac{184t^3-18t^2-149t-145}{736t^2+848t+464} \end{bmatrix} \end{aligned}$$

Which becomes

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} \frac{25}{8} + \frac{c_1(1-i\sqrt{7})e^{\frac{(3+i\sqrt{7})t}{2}}}{2} + \frac{c_2(1+i\sqrt{7})e^{-\frac{(i\sqrt{7}-3)t}{2}}}{2} + \frac{7t}{2} \\ c_1e^{\frac{(3+i\sqrt{7})t}{2}} + c_2e^{-\frac{(i\sqrt{7}-3)t}{2}} + \frac{t}{4} - \frac{5}{16} \end{bmatrix}$$

29.11.3 Maple step by step solution

Let's solve

$$[x'(t) = x(t) + 2y(t) - 4t + 1, y'(t) = -x(t) + 2y(t) + 3t + 4]$$

- Define vector

$$\vec{x}(t) = \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

- Convert system into a vector equation

$$\vec{x}'(t) = \begin{bmatrix} 1 & 2 \\ -1 & 2 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} 1 - 4t \\ 3t + 4 \end{bmatrix}$$

- System to solve

$$\vec{x}'(t) = \begin{bmatrix} 1 & 2 \\ -1 & 2 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} 1 - 4t \\ 3t + 4 \end{bmatrix}$$

- Define the forcing function

$$\vec{f}(t) = \begin{bmatrix} 1 - 4t \\ 3t + 4 \end{bmatrix}$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 1 & 2 \\ -1 & 2 \end{bmatrix}$$

- Rewrite the system as

$$\vec{x}'(t) = A \cdot \vec{x}(t) + \vec{f}$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[\frac{3}{2} - \frac{I\sqrt{7}}{2}, \begin{bmatrix} \frac{2}{\frac{1}{2} - \frac{I\sqrt{7}}{2}} \\ 1 \end{bmatrix} \right], \left[\frac{3}{2} + \frac{I\sqrt{7}}{2}, \begin{bmatrix} \frac{2}{\frac{1}{2} + \frac{I\sqrt{7}}{2}} \\ 1 \end{bmatrix} \right] \right]$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$\left[\frac{3}{2} - \frac{I\sqrt{7}}{2}, \begin{bmatrix} \frac{2}{\frac{1}{2} - \frac{I\sqrt{7}}{2}} \\ 1 \end{bmatrix} \right]$$

- Solution from eigenpair

$$e^{\left(\frac{3}{2} - \frac{I\sqrt{7}}{2}\right)t} \cdot \begin{bmatrix} \frac{2}{\frac{1}{2} - \frac{I\sqrt{7}}{2}} \\ 1 \end{bmatrix}$$

- Use Euler identity to write solution in terms of sin and cos

$$e^{\frac{3t}{2}} \cdot \left(\cos\left(\frac{\sqrt{7}t}{2}\right) - I \sin\left(\frac{\sqrt{7}t}{2}\right) \right) \cdot \begin{bmatrix} \frac{2}{\frac{1}{2} - \frac{I\sqrt{7}}{2}} \\ 1 \end{bmatrix}$$

- Simplify expression

$$e^{\frac{3t}{2}} \cdot \begin{bmatrix} \frac{2(\cos(\frac{\sqrt{7}t}{2}) - I \sin(\frac{\sqrt{7}t}{2}))}{\frac{1}{2} - \frac{I\sqrt{7}}{2}} \\ \cos\left(\frac{\sqrt{7}t}{2}\right) - I \sin\left(\frac{\sqrt{7}t}{2}\right) \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\left[\vec{x}_1(t) = e^{\frac{3t}{2}} \cdot \begin{bmatrix} \frac{\cos(\frac{\sqrt{7}t}{2})}{2} + \frac{\sqrt{7} \sin(\frac{\sqrt{7}t}{2})}{2} \\ \cos\left(\frac{\sqrt{7}t}{2}\right) \end{bmatrix}, \vec{x}_2(t) = e^{\frac{3t}{2}} \cdot \begin{bmatrix} \frac{\sqrt{7} \cos(\frac{\sqrt{7}t}{2})}{2} - \frac{\sin(\frac{\sqrt{7}t}{2})}{2} \\ -\sin\left(\frac{\sqrt{7}t}{2}\right) \end{bmatrix} \right]$$

- General solution of the system of ODEs can be written in terms of the particular solution $\vec{x}_p(t)$
 $\vec{x}(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t) + \vec{x}_p(t)$

□ Fundamental matrix

- Let $\phi(t)$ be the matrix whose columns are the independent solutions of the homogeneous system

$$\phi(t) = \begin{bmatrix} e^{\frac{3t}{2}} \left(\frac{\cos(\frac{\sqrt{7}t}{2})}{2} + \frac{\sqrt{7} \sin(\frac{\sqrt{7}t}{2})}{2} \right) & e^{\frac{3t}{2}} \left(\frac{\sqrt{7} \cos(\frac{\sqrt{7}t}{2})}{2} - \frac{\sin(\frac{\sqrt{7}t}{2})}{2} \right) \\ e^{\frac{3t}{2}} \cos\left(\frac{\sqrt{7}t}{2}\right) & -e^{\frac{3t}{2}} \sin\left(\frac{\sqrt{7}t}{2}\right) \end{bmatrix}$$

- The fundamental matrix, $\Phi(t)$ is a normalized version of $\phi(t)$ satisfying $\Phi(0) = I$ where I is the identity matrix

$$\Phi(t) = \phi(t) \cdot \frac{1}{\phi(0)}$$

- Substitute the value of $\phi(t)$ and $\phi(0)$

$$\Phi(t) = \begin{bmatrix} e^{\frac{3t}{2}} \left(\frac{\cos\left(\frac{\sqrt{7}t}{2}\right)}{2} + \frac{\sqrt{7} \sin\left(\frac{\sqrt{7}t}{2}\right)}{2} \right) & e^{\frac{3t}{2}} \left(\frac{\sqrt{7} \cos\left(\frac{\sqrt{7}t}{2}\right)}{2} - \frac{\sin\left(\frac{\sqrt{7}t}{2}\right)}{2} \right) \\ e^{\frac{3t}{2}} \cos\left(\frac{\sqrt{7}t}{2}\right) & -e^{\frac{3t}{2}} \sin\left(\frac{\sqrt{7}t}{2}\right) \end{bmatrix} \cdot \frac{1}{\begin{bmatrix} \frac{1}{2} & \frac{\sqrt{7}}{2} \\ 1 & 0 \end{bmatrix}}$$

- Evaluate and simplify to get the fundamental matrix

$$\Phi(t) = \begin{bmatrix} \frac{e^{\frac{3t}{2}} \left(\sqrt{7} \cos\left(\frac{\sqrt{7}t}{2}\right) - \sin\left(\frac{\sqrt{7}t}{2}\right) \right) \sqrt{7}}{7} & \frac{4\sqrt{7} e^{\frac{3t}{2}} \sin\left(\frac{\sqrt{7}t}{2}\right)}{7} \\ -\frac{2\sqrt{7} e^{\frac{3t}{2}} \sin\left(\frac{\sqrt{7}t}{2}\right)}{7} & \frac{e^{\frac{3t}{2}} \left(\sqrt{7} \sin\left(\frac{\sqrt{7}t}{2}\right) + 7 \cos\left(\frac{\sqrt{7}t}{2}\right) \right)}{7} \end{bmatrix}$$

- Find a particular solution of the system of ODEs using variation of parameters

- Let the particular solution be the fundamental matrix multiplied by $\vec{v}(t)$ and solve for $\vec{v}(t)$

$$\vec{x}_p(t) = \Phi(t) \cdot \vec{v}(t)$$

- Take the derivative of the particular solution

$$\vec{x}'_p(t) = \Phi'(t) \cdot \vec{v}(t) + \Phi(t) \cdot \vec{v}'(t)$$

- Substitute particular solution and its derivative into the system of ODEs

$$\Phi'(t) \cdot \vec{v}(t) + \Phi(t) \cdot \vec{v}'(t) = A \cdot \Phi(t) \cdot \vec{v}(t) + \vec{f}(t)$$

- The fundamental matrix has columns that are solutions to the homogeneous system so its derivative is $A \cdot \Phi(t)$

$$A \cdot \Phi(t) \cdot \vec{v}(t) + \Phi(t) \cdot \vec{v}'(t) = A \cdot \Phi(t) \cdot \vec{v}(t) + \vec{f}(t)$$

- Cancel like terms

$$\Phi(t) \cdot \vec{v}'(t) = \vec{f}(t)$$

- Multiply by the inverse of the fundamental matrix

$$\vec{v}'(t) = \frac{1}{\Phi(t)} \cdot \vec{f}(t)$$

- Integrate to solve for $\vec{v}(t)$

$$\vec{v}(t) = \int_0^t \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds$$

- Plug $\vec{v}(t)$ into the equation for the particular solution

$$\vec{x}_p(t) = \Phi(t) \cdot \left(\int_0^t \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds \right)$$

- Plug in the fundamental matrix and the forcing function and compute

$$\vec{x}_p(t) = \begin{bmatrix} \frac{5\sqrt{7}e^{\frac{3t}{2}} \sin\left(\frac{\sqrt{7}t}{2}\right)}{8} - \frac{25e^{\frac{3t}{2}} \cos\left(\frac{\sqrt{7}t}{2}\right)}{8} + \frac{7t}{2} + \frac{25}{8} \\ \frac{15\sqrt{7}e^{\frac{3t}{2}} \sin\left(\frac{\sqrt{7}t}{2}\right)}{16} + \frac{5e^{\frac{3t}{2}} \cos\left(\frac{\sqrt{7}t}{2}\right)}{16} + \frac{t}{4} - \frac{5}{16} \end{bmatrix}$$

- Plug particular solution back into general solution

$$\vec{x}(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t) + \begin{bmatrix} \frac{5\sqrt{7}e^{\frac{3t}{2}} \sin\left(\frac{\sqrt{7}t}{2}\right)}{8} - \frac{25e^{\frac{3t}{2}} \cos\left(\frac{\sqrt{7}t}{2}\right)}{8} + \frac{7t}{2} + \frac{25}{8} \\ \frac{15\sqrt{7}e^{\frac{3t}{2}} \sin\left(\frac{\sqrt{7}t}{2}\right)}{16} + \frac{5e^{\frac{3t}{2}} \cos\left(\frac{\sqrt{7}t}{2}\right)}{16} + \frac{t}{4} - \frac{5}{16} \end{bmatrix}$$

- Substitute in vector of dependent variables

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} \frac{e^{\frac{3t}{2}} (\sqrt{7}c_2 + c_1 - \frac{25}{4}) \cos\left(\frac{\sqrt{7}t}{2}\right)}{2} + \frac{((c_1 + \frac{5}{4})\sqrt{7} - c_2) e^{\frac{3t}{2}} \sin\left(\frac{\sqrt{7}t}{2}\right)}{2} + \frac{7t}{2} + \frac{25}{8} \\ \frac{e^{\frac{3t}{2}} (16c_1 + 5) \cos\left(\frac{\sqrt{7}t}{2}\right)}{16} - e^{\frac{3t}{2}} \left(c_2 - \frac{15\sqrt{7}}{16}\right) \sin\left(\frac{\sqrt{7}t}{2}\right) + \frac{t}{4} - \frac{5}{16} \end{bmatrix}$$

- Solution to the system of ODEs

$$\begin{cases} x(t) = \frac{e^{\frac{3t}{2}} (\sqrt{7}c_2 + c_1 - \frac{25}{4}) \cos\left(\frac{\sqrt{7}t}{2}\right)}{2} + \frac{((c_1 + \frac{5}{4})\sqrt{7} - c_2) e^{\frac{3t}{2}} \sin\left(\frac{\sqrt{7}t}{2}\right)}{2} + \frac{7t}{2} + \frac{25}{8}, \\ y(t) = \frac{e^{\frac{3t}{2}} (16c_1 + 5) \cos\left(\frac{\sqrt{7}t}{2}\right)}{16} - e^{\frac{3t}{2}} \left(c_2 - \frac{15\sqrt{7}}{16}\right) \sin\left(\frac{\sqrt{7}t}{2}\right) + \frac{t}{4} - \frac{5}{16} \end{cases}$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 106

```
dsolve([diff(x(t),t)=x(t)+2*y(t)-4*t+1,diff(y(t),t)=-x(t)+2*y(t)+3*t+4],singsol=all)
```

$$\begin{aligned} x(t) &= e^{\frac{3t}{2}} \sin\left(\frac{\sqrt{7}t}{2}\right) c_2 + e^{\frac{3t}{2}} \cos\left(\frac{\sqrt{7}t}{2}\right) c_1 + \frac{7t}{2} + \frac{25}{8} \\ y(t) &= \frac{e^{\frac{3t}{2}} \sin\left(\frac{\sqrt{7}t}{2}\right) c_2}{4} + \frac{e^{\frac{3t}{2}} \sqrt{7} \cos\left(\frac{\sqrt{7}t}{2}\right) c_2}{4} \\ &\quad + \frac{e^{\frac{3t}{2}} \cos\left(\frac{\sqrt{7}t}{2}\right) c_1}{4} - \frac{e^{\frac{3t}{2}} \sqrt{7} \sin\left(\frac{\sqrt{7}t}{2}\right) c_1}{4} - \frac{5}{16} + \frac{t}{4} \end{aligned}$$

✓ Solution by Mathematica

Time used: 2.624 (sec). Leaf size: 128

```
DSolve[{x'[t]==x[t]+2*y[t]-4+t+1,y'[t]==-x[t]+2*y[t]+3*t+4},{x[t],y[t]},t,IncludeSingularSol
```

$$x(t) \rightarrow t + c_1 e^{3t/2} \cos\left(\frac{\sqrt{7}t}{2}\right) - \frac{(c_1 - 4c_2)e^{3t/2} \sin\left(\frac{\sqrt{7}t}{2}\right)}{\sqrt{7}} + \frac{9}{2}$$

$$y(t) \rightarrow -t + c_2 e^{3t/2} \cos\left(\frac{\sqrt{7}t}{2}\right) - \frac{(2c_1 - c_2)e^{3t/2} \sin\left(\frac{\sqrt{7}t}{2}\right)}{\sqrt{7}} - \frac{1}{4}$$

29.12 problem 4(b)

29.12.1 Solution using Matrix exponential method	4199
29.12.2 Solution using explicit Eigenvalue and Eigenvector method . . .	4201
29.12.3 Maple step by step solution	4207

Internal problem ID [6540]

Internal file name [OUTPUT/5788_Sunday_June_05_2022_03_54_30_PM_13946785/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 10. Systems of First-Order Equations. Section A. Drill exercises. Page 400

Problem number: 4(b).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "system of linear ODEs"

Solve

$$\begin{aligned}x'(t) &= -2x(t) + y(t) - t + 3 \\y'(t) &= x(t) + 4y(t) + t - 2\end{aligned}$$

29.12.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t) + \vec{G}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} -2 & 1 \\ 1 & 4 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix} + \begin{bmatrix} -t + 3 \\ -2 + t \end{bmatrix}$$

Since the system is nonhomogeneous, then the solution is given by

$$\vec{x}(t) = \vec{x}_h(t) + \vec{x}_p(t)$$

Where $\vec{x}_h(t)$ is the homogeneous solution to $\vec{x}'(t) = A\vec{x}(t)$ and $\vec{x}_p(t)$ is a particular solution to $\vec{x}'(t) = A\vec{x}(t) + \vec{G}(t)$. The particular solution will be found using variation

of parameters method applied to the fundamental matrix. For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} \frac{(3\sqrt{10}+10)e^{-(1+\sqrt{10})t}}{20} + \frac{(-3\sqrt{10}+10)e^{(1+\sqrt{10})t}}{20} & -\frac{(-e^{(1+\sqrt{10})t} + e^{-(1+\sqrt{10})t})\sqrt{10}}{20} \\ -\frac{(-e^{(1+\sqrt{10})t} + e^{-(1+\sqrt{10})t})\sqrt{10}}{20} & \frac{(-3\sqrt{10}+10)e^{-(1+\sqrt{10})t}}{20} + \frac{e^{(1+\sqrt{10})t}(3\sqrt{10}+10)}{20} \end{bmatrix}$$

Therefore the homogeneous solution is

$$\begin{aligned} \vec{x}_h(t) &= e^{At}\vec{c} \\ &= \begin{bmatrix} \frac{(3\sqrt{10}+10)e^{-(1+\sqrt{10})t}}{20} + \frac{(-3\sqrt{10}+10)e^{(1+\sqrt{10})t}}{20} & -\frac{(-e^{(1+\sqrt{10})t} + e^{-(1+\sqrt{10})t})\sqrt{10}}{20} \\ -\frac{(-e^{(1+\sqrt{10})t} + e^{-(1+\sqrt{10})t})\sqrt{10}}{20} & \frac{(-3\sqrt{10}+10)e^{-(1+\sqrt{10})t}}{20} + \frac{e^{(1+\sqrt{10})t}(3\sqrt{10}+10)}{20} \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} \\ &= \begin{bmatrix} \left(\frac{(3\sqrt{10}+10)e^{-(1+\sqrt{10})t}}{20} + \frac{(-3\sqrt{10}+10)e^{(1+\sqrt{10})t}}{20} \right) c_1 - \frac{(-e^{(1+\sqrt{10})t} + e^{-(1+\sqrt{10})t})\sqrt{10}c_2}{20} \\ -\frac{(-e^{(1+\sqrt{10})t} + e^{-(1+\sqrt{10})t})\sqrt{10}c_1}{20} + \left(\frac{(-3\sqrt{10}+10)e^{-(1+\sqrt{10})t}}{20} + \frac{e^{(1+\sqrt{10})t}(3\sqrt{10}+10)}{20} \right) c_2 \end{bmatrix} \\ &= \begin{bmatrix} \frac{(3c_1 - c_2)\sqrt{10} + 10c_1}{20} e^{-(1+\sqrt{10})t} - \frac{3\left((c_1 - \frac{c_2}{3})\sqrt{10} - \frac{10c_1}{3}\right)e^{(1+\sqrt{10})t}}{20} \\ \frac{((-c_1 - 3c_2)\sqrt{10} + 10c_2)e^{-(1+\sqrt{10})t}}{20} + \frac{((c_1 + 3c_2)\sqrt{10} + 10c_2)e^{(1+\sqrt{10})t}}{20} \end{bmatrix} \end{aligned}$$

The particular solution given by

$$\vec{x}_p(t) = e^{At} \int e^{-At} \vec{G}(t) dt$$

But

$$e^{-At} = (e^{At})^{-1} = \begin{bmatrix} -\frac{e^{-2t} \left(3\sqrt{10}e^{-(1+\sqrt{10})t} - 3\sqrt{10}e^{(1+\sqrt{10})t} - 10e^{-(1+\sqrt{10})t} - 10e^{(1+\sqrt{10})t} \right)}{20} & \frac{\sqrt{10}e^{-2t} \left(-e^{(1+\sqrt{10})t} + e^{-(1+\sqrt{10})t} \right)}{20} \\ \frac{\sqrt{10}e^{-2t} \left(-e^{(1+\sqrt{10})t} + e^{-(1+\sqrt{10})t} \right)}{20} & \frac{e^{-2t} \left((3\sqrt{10}+10)e^{-(1+\sqrt{10})t} - 3e^{(1+\sqrt{10})t} \right)}{20} \end{bmatrix}$$

Hence

$$\begin{aligned}
 \vec{x}_p(t) &= \begin{bmatrix} \frac{(3\sqrt{10}+10)e^{-(-1+\sqrt{10})t}}{20} + \frac{(-3\sqrt{10}+10)e^{(1+\sqrt{10})t}}{20} & -\frac{(-e^{(1+\sqrt{10})t}+e^{-(-1+\sqrt{10})t})\sqrt{10}}{20} \\ -\frac{(-e^{(1+\sqrt{10})t}+e^{-(-1+\sqrt{10})t})\sqrt{10}}{20} & \frac{(-3\sqrt{10}+10)e^{-(-1+\sqrt{10})t}}{20} + \frac{e^{(1+\sqrt{10})t}(3\sqrt{10}+10)}{20} \end{bmatrix} \int \begin{bmatrix} e \\ -e \end{bmatrix} \\
 &= \begin{bmatrix} \frac{(3\sqrt{10}+10)e^{-(-1+\sqrt{10})t}}{20} + \frac{(-3\sqrt{10}+10)e^{(1+\sqrt{10})t}}{20} & -\frac{(-e^{(1+\sqrt{10})t}+e^{-(-1+\sqrt{10})t})\sqrt{10}}{20} \\ -\frac{(-e^{(1+\sqrt{10})t}+e^{-(-1+\sqrt{10})t})\sqrt{10}}{20} & \frac{(-3\sqrt{10}+10)e^{-(-1+\sqrt{10})t}}{20} + \frac{e^{(1+\sqrt{10})t}(3\sqrt{10}+10)}{20} \end{bmatrix} \begin{bmatrix} (126t) \\ (-7) \end{bmatrix} \\
 &= \begin{bmatrix} -\frac{5t}{9} + \frac{145}{81} \\ -\frac{t}{9} + \frac{2}{81} \end{bmatrix}
 \end{aligned}$$

Hence the complete solution is

$$\begin{aligned}
 \vec{x}(t) &= \vec{x}_h(t) + \vec{x}_p(t) \\
 &= \begin{bmatrix} \frac{145}{81} + \frac{((3c_1-c_2)\sqrt{10}+10c_1)e^{-(-1+\sqrt{10})t}}{20} + \frac{((-3c_1+c_2)\sqrt{10}+10c_1)e^{(1+\sqrt{10})t}}{20} - \frac{5t}{9} \\ \frac{((-c_1-3c_2)\sqrt{10}+10c_2)e^{-(-1+\sqrt{10})t}}{20} + \frac{((c_1+3c_2)\sqrt{10}+10c_2)e^{(1+\sqrt{10})t}}{20} - \frac{t}{9} + \frac{2}{81} \end{bmatrix}
 \end{aligned}$$

29.12.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t) + \vec{G}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} -2 & 1 \\ 1 & 4 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix} + \begin{bmatrix} -t+3 \\ -2+t \end{bmatrix}$$

Since the system is nonhomogeneous, then the solution is given by

$$\vec{x}(t) = \vec{x}_h(t) + \vec{x}_p(t)$$

Where $\vec{x}_h(t)$ is the homogeneous solution to $\vec{x}'(t) = A\vec{x}(t)$ and $\vec{x}_p(t)$ is a particular solution to $\vec{x}'(t) = A\vec{x}(t) + \vec{G}(t)$. The particular solution will be found using variation of parameters method applied to the fundamental matrix.

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det \left(\begin{bmatrix} -2 & 1 \\ 1 & 4 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) = 0$$

Therefore

$$\det \left(\begin{bmatrix} -2 - \lambda & 1 \\ 1 & 4 - \lambda \end{bmatrix} \right) = 0$$

Which gives the characteristic equation

$$\lambda^2 - 2\lambda - 9 = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = 1 + \sqrt{10}$$

$$\lambda_2 = 1 - \sqrt{10}$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
$1 - \sqrt{10}$	1	real eigenvalue
$1 + \sqrt{10}$	1	real eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = 1 - \sqrt{10}$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} -2 & 1 \\ 1 & 4 \end{bmatrix} - (1 - \sqrt{10}) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} -3 + \sqrt{10} & 1 \\ 1 & 3 + \sqrt{10} \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\begin{bmatrix} -3 + \sqrt{10} & 1 & | & 0 \\ 1 & 3 + \sqrt{10} & | & 0 \end{bmatrix}$$

$$R_2 = R_2 - \frac{R_1}{-3 + \sqrt{10}} \Rightarrow \begin{bmatrix} -3 + \sqrt{10} & 1 & | & 0 \\ 0 & 0 & | & 0 \end{bmatrix}$$

Therefore the system in Echelon form is

$$\begin{bmatrix} -3 + \sqrt{10} & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\left\{ v_1 = -\frac{t}{-3 + \sqrt{10}} \right\}$

Hence the solution is

$$\begin{bmatrix} -\frac{t}{-3 + \sqrt{10}} \\ t \end{bmatrix} = \begin{bmatrix} -\frac{t}{-3 + \sqrt{10}} \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} -\frac{t}{-3 + \sqrt{10}} \\ t \end{bmatrix} = t \begin{bmatrix} -\frac{1}{-3 + \sqrt{10}} \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} -\frac{1}{-3 + \sqrt{10}} \\ 1 \end{bmatrix} = \begin{bmatrix} -\frac{1}{-3 + \sqrt{10}} \\ 1 \end{bmatrix}$$

Considering the eigenvalue $\lambda_2 = 1 + \sqrt{10}$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} -2 & 1 \\ 1 & 4 \end{bmatrix} - (1 + \sqrt{10}) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} -3 - \sqrt{10} & 1 \\ 1 & 3 - \sqrt{10} \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\begin{bmatrix} -3 - \sqrt{10} & 1 & | & 0 \\ 1 & 3 - \sqrt{10} & | & 0 \end{bmatrix}$$

$$R_2 = R_2 - \frac{R_1}{-3 - \sqrt{10}} \implies \begin{bmatrix} -3 - \sqrt{10} & 1 & | & 0 \\ 0 & 0 & | & 0 \end{bmatrix}$$

Therefore the system in Echelon form is

$$\begin{bmatrix} -3 - \sqrt{10} & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\left\{ v_1 = \frac{t}{3 + \sqrt{10}} \right\}$

Hence the solution is

$$\begin{bmatrix} \frac{t}{3 + \sqrt{10}} \\ t \end{bmatrix} = \begin{bmatrix} \frac{t}{3 + \sqrt{10}} \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} \frac{t}{3 + \sqrt{10}} \\ t \end{bmatrix} = t \begin{bmatrix} \frac{1}{3 + \sqrt{10}} \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} \frac{1}{3 + \sqrt{10}} \\ 1 \end{bmatrix} = \begin{bmatrix} \frac{1}{3 + \sqrt{10}} \\ 1 \end{bmatrix}$$

Which is normalized to

$$\begin{bmatrix} \frac{1}{3 + \sqrt{10}} \\ 1 \end{bmatrix} = \begin{bmatrix} \frac{1}{3 + \sqrt{10}} \\ 1 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
$1 + \sqrt{10}$	1	1	No	$\begin{bmatrix} \frac{1}{3+\sqrt{10}} \\ 1 \end{bmatrix}$
$1 - \sqrt{10}$	1	1	No	$\begin{bmatrix} \frac{1}{3-\sqrt{10}} \\ 1 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. Since eigenvalue $1 + \sqrt{10}$ is real and distinct then the corresponding eigenvector solution is

$$\begin{aligned}\vec{x}_1(t) &= \vec{v}_1 e^{(1+\sqrt{10})t} \\ &= \begin{bmatrix} \frac{1}{3+\sqrt{10}} \\ 1 \end{bmatrix} e^{(1+\sqrt{10})t}\end{aligned}$$

Since eigenvalue $1 - \sqrt{10}$ is real and distinct then the corresponding eigenvector solution is

$$\begin{aligned}\vec{x}_2(t) &= \vec{v}_2 e^{(1-\sqrt{10})t} \\ &= \begin{bmatrix} \frac{1}{3-\sqrt{10}} \\ 1 \end{bmatrix} e^{(1-\sqrt{10})t}\end{aligned}$$

Therefore the homogeneous solution is

$$\vec{x}_h(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

Which is written as

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = c_1 \begin{bmatrix} \frac{e^{(1+\sqrt{10})t}}{3+\sqrt{10}} \\ e^{(1+\sqrt{10})t} \end{bmatrix} + c_2 \begin{bmatrix} \frac{e^{(1-\sqrt{10})t}}{3-\sqrt{10}} \\ e^{(1-\sqrt{10})t} \end{bmatrix}$$

Now that we found homogeneous solution above, we need to find a particular solution $\vec{x}_p(t)$. We will use Variation of parameters. The fundamental matrix is

$$\Phi = \begin{bmatrix} \vec{x}_1 & \vec{x}_2 & \cdots \end{bmatrix}$$

Where \vec{x}_i are the solution basis found above. Therefore the fundamental matrix is

$$\Phi(t) = \begin{bmatrix} \frac{e^{(1+\sqrt{10})t}}{3+\sqrt{10}} & \frac{e^{(1-\sqrt{10})t}}{3-\sqrt{10}} \\ e^{(1+\sqrt{10})t} & e^{(1-\sqrt{10})t} \end{bmatrix}$$

The particular solution is then given by

$$\vec{x}_p(t) = \Phi \int \Phi^{-1} \vec{G}(t) dt$$

But

$$\Phi^{-1} = \begin{bmatrix} \frac{\sqrt{10}e^{-(1+\sqrt{10})t}}{20} & \frac{\sqrt{10}(3+\sqrt{10})e^{-(1+\sqrt{10})t}}{20} \\ -\frac{\sqrt{10}e^{(-1+\sqrt{10})t}}{20} & \frac{e^{(-1+\sqrt{10})t}\sqrt{10}(-3+\sqrt{10})}{20} \end{bmatrix}$$

Hence

$$\begin{aligned} \vec{x}_p(t) &= \begin{bmatrix} \frac{e^{(1+\sqrt{10})t}}{3+\sqrt{10}} & \frac{e^{(1-\sqrt{10})t}}{3-\sqrt{10}} \\ e^{(1+\sqrt{10})t} & e^{(1-\sqrt{10})t} \end{bmatrix} \int \begin{bmatrix} \frac{\sqrt{10}e^{-(1+\sqrt{10})t}}{20} & \frac{\sqrt{10}(3+\sqrt{10})e^{-(1+\sqrt{10})t}}{20} \\ -\frac{\sqrt{10}e^{(-1+\sqrt{10})t}}{20} & \frac{e^{(-1+\sqrt{10})t}\sqrt{10}(-3+\sqrt{10})}{20} \end{bmatrix} \begin{bmatrix} -t+3 \\ -2+t \end{bmatrix} dt \\ &= \begin{bmatrix} \frac{e^{(1+\sqrt{10})t}}{3+\sqrt{10}} & \frac{e^{(1-\sqrt{10})t}}{3-\sqrt{10}} \\ e^{(1+\sqrt{10})t} & e^{(1-\sqrt{10})t} \end{bmatrix} \int \begin{bmatrix} \frac{((2t-3)\sqrt{10}+10t-20)e^{-(1+\sqrt{10})t}}{20} \\ \frac{((3-2t)\sqrt{10}+10t-20)e^{(-1+\sqrt{10})t}}{20} \end{bmatrix} dt \\ &= \begin{bmatrix} \frac{e^{(1+\sqrt{10})t}}{3+\sqrt{10}} & \frac{e^{(1-\sqrt{10})t}}{3-\sqrt{10}} \\ e^{(1+\sqrt{10})t} & e^{(1-\sqrt{10})t} \end{bmatrix} \begin{bmatrix} -\frac{(18t\sqrt{10}-49\sqrt{10}-18t+94)\sqrt{10}e^{-(1+\sqrt{10})t}(t\sqrt{10}-2\sqrt{10}+2t-3)}{540(6t-14+\sqrt{10})} \\ \frac{(18t\sqrt{10}-49\sqrt{10}+18t-94)\sqrt{10}e^{(-1+\sqrt{10})t}(t\sqrt{10}-2\sqrt{10}-2t+3)}{3240t-7560-540\sqrt{10}} \end{bmatrix} \\ &= \begin{bmatrix} \frac{-540t^3+4260t^2-10910t+8990}{-270+27(6t-14)^2} \\ \frac{-108t^3+528t^2-670t+124}{-270+27(6t-14)^2} \end{bmatrix} \end{aligned}$$

Now that we found particular solution, the final solution is

$$\begin{aligned} \vec{x}(t) &= \vec{x}_h(t) + \vec{x}_p(t) \\ \begin{bmatrix} x(t) \\ y(t) \end{bmatrix} &= \begin{bmatrix} \frac{c_1 e^{(1+\sqrt{10})t}}{3+\sqrt{10}} \\ c_1 e^{(1+\sqrt{10})t} \end{bmatrix} + \begin{bmatrix} \frac{c_2 e^{(1-\sqrt{10})t}}{3-\sqrt{10}} \\ c_2 e^{(1-\sqrt{10})t} \end{bmatrix} + \begin{bmatrix} \frac{-540t^3+4260t^2-10910t+8990}{-270+27(6t-14)^2} \\ \frac{-108t^3+528t^2-670t+124}{-270+27(6t-14)^2} \end{bmatrix} \end{aligned}$$

Which becomes

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} -c_2(3 + \sqrt{10}) e^{-(-1+\sqrt{10})t} + c_1(-3 + \sqrt{10}) e^{(1+\sqrt{10})t} - \frac{5t}{9} + \frac{145}{81} \\ c_1 e^{(1+\sqrt{10})t} + c_2 e^{-(-1+\sqrt{10})t} - \frac{t}{9} + \frac{2}{81} \end{bmatrix}$$

29.12.3 Maple step by step solution

Let's solve

$$[x'(t) = -2x(t) + y(t) - t + 3, y'(t) = x(t) + 4y(t) + t - 2]$$

- Define vector

$$\vec{x}(t) = \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

- Convert system into a vector equation

$$\vec{x}'(t) = \begin{bmatrix} -2 & 1 \\ 1 & 4 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} -t + 3 \\ -2 + t \end{bmatrix}$$

- System to solve

$$\vec{x}'(t) = \begin{bmatrix} -2 & 1 \\ 1 & 4 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} -t + 3 \\ -2 + t \end{bmatrix}$$

- Define the forcing function

$$\vec{f}(t) = \begin{bmatrix} -t + 3 \\ -2 + t \end{bmatrix}$$

- Define the coefficient matrix

$$A = \begin{bmatrix} -2 & 1 \\ 1 & 4 \end{bmatrix}$$

- Rewrite the system as

$$\vec{x}'(t) = A \cdot \vec{x}(t) + \vec{f}$$

- To solve the system, find the eigenvalues and eigenvectors of A
- Eigenpairs of A

$$\left[\left[1 - \sqrt{10}, \begin{bmatrix} \frac{1}{3-\sqrt{10}} \\ 1 \end{bmatrix} \right], \left[1 + \sqrt{10}, \begin{bmatrix} \frac{1}{3+\sqrt{10}} \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[1 - \sqrt{10}, \begin{bmatrix} \frac{1}{3-\sqrt{10}} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{x}_1 = e^{(1-\sqrt{10})t} \cdot \begin{bmatrix} \frac{1}{3-\sqrt{10}} \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[1 + \sqrt{10}, \begin{bmatrix} \frac{1}{3+\sqrt{10}} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{x}_2 = e^{(1+\sqrt{10})t} \cdot \begin{bmatrix} \frac{1}{3+\sqrt{10}} \\ 1 \end{bmatrix}$$

- General solution of the system of ODEs can be written in terms of the particular solution $\vec{x}_p(t)$

$$\vec{x}(t) = c_1 \vec{x}_1 + c_2 \vec{x}_2 + \vec{x}_p(t)$$

- Fundamental matrix

- Let $\phi(t)$ be the matrix whose columns are the independent solutions of the homogeneous system

$$\phi(t) = \begin{bmatrix} \frac{e^{(1-\sqrt{10})t}}{3-\sqrt{10}} & \frac{e^{(1+\sqrt{10})t}}{3+\sqrt{10}} \\ e^{(1-\sqrt{10})t} & e^{(1+\sqrt{10})t} \end{bmatrix}$$

- The fundamental matrix, $\Phi(t)$ is a normalized version of $\phi(t)$ satisfying $\Phi(0) = I$ where I is the identity matrix

$$\Phi(t) = \phi(t) \cdot \frac{1}{\phi(0)}$$

- Substitute the value of $\phi(t)$ and $\phi(0)$

$$\Phi(t) = \begin{bmatrix} \frac{e^{(1-\sqrt{10})t}}{3-\sqrt{10}} & \frac{e^{(1+\sqrt{10})t}}{3+\sqrt{10}} \\ e^{(1-\sqrt{10})t} & e^{(1+\sqrt{10})t} \end{bmatrix} \cdot \frac{1}{\begin{bmatrix} \frac{1}{3-\sqrt{10}} & \frac{1}{3+\sqrt{10}} \\ 1 & 1 \end{bmatrix}}$$

- Evaluate and simplify to get the fundamental matrix

$$\Phi(t) = \begin{bmatrix} \frac{\left((3+\sqrt{10})e^{-(-1+\sqrt{10})t} + e^{(1+\sqrt{10})t}(-3+\sqrt{10}) \right) \sqrt{10}}{20} & -\frac{\left(-e^{(1+\sqrt{10})t} + e^{-(-1+\sqrt{10})t} \right) \sqrt{10}}{20} \\ -\frac{\left(-e^{(1+\sqrt{10})t} + e^{-(-1+\sqrt{10})t} \right) \sqrt{10}}{20} & \frac{(-3\sqrt{10}+10)e^{-(-1+\sqrt{10})t}}{20} + \frac{e^{(1+\sqrt{10})t}(3\sqrt{10}+10)}{20} \end{bmatrix}$$

□

Find a particular solution of the system of ODEs using variation of parameters

- Let the particular solution be the fundamental matrix multiplied by $\vec{v}(t)$ and solve for $\vec{v}(t)$

$$\vec{x}_p(t) = \Phi(t) \cdot \vec{v}(t)$$

- Take the derivative of the particular solution

$$\vec{x}'_p(t) = \Phi'(t) \cdot \vec{v}(t) + \Phi(t) \cdot \vec{v}'(t)$$

- Substitute particular solution and its derivative into the system of ODEs

$$\Phi'(t) \cdot \vec{v}(t) + \Phi(t) \cdot \vec{v}'(t) = A \cdot \Phi(t) \cdot \vec{v}(t) + \vec{f}(t)$$

- The fundamental matrix has columns that are solutions to the homogeneous system so its derivative is $A \cdot \Phi(t) \cdot \vec{v}(t)$

$$A \cdot \Phi(t) \cdot \vec{v}(t) + \Phi(t) \cdot \vec{v}'(t) = A \cdot \Phi(t) \cdot \vec{v}(t) + \vec{f}(t)$$

- Cancel like terms

$$\Phi(t) \cdot \vec{v}'(t) = \vec{f}(t)$$

- Multiply by the inverse of the fundamental matrix

$$\vec{v}'(t) = \frac{1}{\Phi(t)} \cdot \vec{f}(t)$$

- Integrate to solve for $\vec{v}(t)$

$$\vec{v}(t) = \int_0^t \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds$$

- Plug $\vec{v}(t)$ into the equation for the particular solution

$$\vec{x}_p(t) = \Phi(t) \cdot \left(\int_0^t \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds \right)$$

- Plug in the fundamental matrix and the forcing function and compute

$$\vec{x}_p(t) = \begin{bmatrix} \frac{(-1450-433\sqrt{10})e^{-(-1+\sqrt{10})t} + (-1450+433\sqrt{10})e^{(1+\sqrt{10})t} - 900t + 2900}{20(1+\sqrt{10})^2(-1+\sqrt{10})^2} \\ \frac{(151\sqrt{10}-20)e^{-(-1+\sqrt{10})t}}{1620} + \frac{(-151\sqrt{10}-20)e^{(1+\sqrt{10})t}}{1620} - \frac{t}{9} + \frac{2}{81} \end{bmatrix}$$

- Plug particular solution back into general solution

$$\vec{x}(t) = c_1 \vec{x}_1 + c_2 \vec{x}_2 + \begin{bmatrix} \frac{(-1450-433\sqrt{10})e^{-(1+\sqrt{10})t} + (-1450+433\sqrt{10})e^{(1+\sqrt{10})t} - 900t + 2900}{20(1+\sqrt{10})^2(-1+\sqrt{10})^2} \\ \frac{(151\sqrt{10}-20)e^{-(1+\sqrt{10})t}}{1620} + \frac{(-151\sqrt{10}-20)e^{(1+\sqrt{10})t}}{1620} - \frac{t}{9} + \frac{2}{81} \end{bmatrix}$$

- Substitute in vector of dependent variables

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} \frac{((-1620c_1-433)\sqrt{10}-4860c_1-1450)e^{-(1+\sqrt{10})t}}{1620} + \frac{((1620c_2+433)\sqrt{10}-4860c_2-1450)e^{(1+\sqrt{10})t}}{1620} - \frac{5t}{9} \\ \frac{(1620c_1+151\sqrt{10}-20)e^{-(1+\sqrt{10})t}}{1620} + \frac{(1620c_2-151\sqrt{10}-20)e^{(1+\sqrt{10})t}}{1620} - \frac{t}{9} + \frac{2}{81} \end{bmatrix}$$

- Solution to the system of ODEs

$$\begin{cases} x(t) = \frac{((-1620c_1-433)\sqrt{10}-4860c_1-1450)e^{-(1+\sqrt{10})t}}{1620} + \frac{((1620c_2+433)\sqrt{10}-4860c_2-1450)e^{(1+\sqrt{10})t}}{1620} - \frac{5t}{9} + \frac{145}{81}, \\ y(t) = \frac{(1620c_1+151\sqrt{10}-20)e^{-(1+\sqrt{10})t}}{1620} + \frac{(1620c_2-151\sqrt{10}-20)e^{(1+\sqrt{10})t}}{1620} - \frac{t}{9} + \frac{2}{81} \end{cases}$$

✓ Solution by Maple

Time used: 0.156 (sec). Leaf size: 90

```
dsolve([diff(x(t),t)=-2*x(t)+y(t)-t+3,diff(y(t),t)=x(t)+4*y(t)+t-2],singsol=all)
```

$$x(t) = e^{(1+\sqrt{10})t} c_2 + e^{-(1+\sqrt{10})t} c_1 - \frac{5t}{9} + \frac{145}{81}$$

$$y(t) = e^{(1+\sqrt{10})t} c_2 \sqrt{10} - e^{-(1+\sqrt{10})t} c_1 \sqrt{10} + 3e^{(1+\sqrt{10})t} c_2 + 3e^{-(1+\sqrt{10})t} c_1 - \frac{t}{9} + \frac{2}{81}$$

✓ Solution by Mathematica

Time used: 10.617 (sec). Leaf size: 190

```
DSolve[{x'[t]==-2*x[t]+y[t]-t+3,y'[t]==x[t]+4*y[t]+t-2},{x[t],y[t]},t,IncludeSingularSolutions->True]
```

$x(t) \rightarrow$

$$\frac{e^{t-\sqrt{10}t} \left(100e^{(\sqrt{10}-1)t} (9t-29) + 81((3\sqrt{10}-10)c_1 - \sqrt{10}c_2) e^{2\sqrt{10}t} - 81(10+3\sqrt{10})c_1 + 81\sqrt{10}c_2 \right)}{1620}$$

$y(t)$

$$\frac{e^{t-\sqrt{10}t} \left(-20e^{(\sqrt{10}-1)t} (9t-2) + 81(\sqrt{10}c_1 + (10+3\sqrt{10})c_2) e^{2\sqrt{10}t} - 81(\sqrt{10}c_1 + (3\sqrt{10}-10)c_2) \right)}{1620}$$

29.13 problem 4(c)

29.13.1 Solution using Matrix exponential method	4211
29.13.2 Solution using explicit Eigenvalue and Eigenvector method . . .	4213
29.13.3 Maple step by step solution	4219

Internal problem ID [6541]

Internal file name [OUTPUT/5789_Sunday_June_05_2022_03_54_34_PM_60465517/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 10. Systems of First-Order Equations. Section A. Drill exercises. Page 400

Problem number: 4(c).

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**system of linear ODEs**"

Solve

$$\begin{aligned}x'(t) &= -4x(t) + y(t) - t + 3 \\y'(t) &= -x(t) - 5y(t) + t + 1\end{aligned}$$

29.13.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t) + \vec{G}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} -4 & 1 \\ -1 & -5 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix} + \begin{bmatrix} -t + 3 \\ t + 1 \end{bmatrix}$$

Since the system is nonhomogeneous, then the solution is given by

$$\vec{x}(t) = \vec{x}_h(t) + \vec{x}_p(t)$$

Where $\vec{x}_h(t)$ is the homogeneous solution to $\vec{x}'(t) = A\vec{x}(t)$ and $\vec{x}_p(t)$ is a particular solution to $\vec{x}'(t) = A\vec{x}(t) + \vec{G}(t)$. The particular solution will be found using variation

of parameters method applied to the fundamental matrix. For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} e^{-\frac{9t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right) + \frac{\sqrt{3}e^{-\frac{9t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right)}{3} & \frac{2\sqrt{3}e^{-\frac{9t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right)}{3} \\ -\frac{2\sqrt{3}e^{-\frac{9t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right)}{3} & e^{-\frac{9t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right) - \frac{\sqrt{3}e^{-\frac{9t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right)}{3} \end{bmatrix}$$

$$= \begin{bmatrix} \frac{e^{-\frac{9t}{2}} \left(\sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right) + 3 \cos\left(\frac{\sqrt{3}t}{2}\right)\right)}{3} & \frac{2\sqrt{3}e^{-\frac{9t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right)}{3} \\ -\frac{2\sqrt{3}e^{-\frac{9t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right)}{3} & \frac{e^{-\frac{9t}{2}} \left(\sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right) - 3 \cos\left(\frac{\sqrt{3}t}{2}\right)\right)}{3} \end{bmatrix}$$

Therefore the homogeneous solution is

$$\vec{x}_h(t) = e^{At} \vec{c}$$

$$= \begin{bmatrix} \frac{e^{-\frac{9t}{2}} \left(\sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right) + 3 \cos\left(\frac{\sqrt{3}t}{2}\right)\right)}{3} & \frac{2\sqrt{3}e^{-\frac{9t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right)}{3} \\ -\frac{2\sqrt{3}e^{-\frac{9t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right)}{3} & \frac{e^{-\frac{9t}{2}} \left(\sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right) - 3 \cos\left(\frac{\sqrt{3}t}{2}\right)\right)}{3} \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix}$$

$$= \begin{bmatrix} \frac{e^{-\frac{9t}{2}} \left(\sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right) + 3 \cos\left(\frac{\sqrt{3}t}{2}\right)\right) c_1}{3} + \frac{2\sqrt{3}e^{-\frac{9t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right) c_2}{3} \\ -\frac{2\sqrt{3}e^{-\frac{9t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right) c_1}{3} - \frac{e^{-\frac{9t}{2}} \left(\sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right) - 3 \cos\left(\frac{\sqrt{3}t}{2}\right)\right) c_2}{3} \end{bmatrix}$$

$$= \begin{bmatrix} \frac{e^{-\frac{9t}{2}} \left(\sqrt{3} (c_1 + 2c_2) \sin\left(\frac{\sqrt{3}t}{2}\right) + 3 \cos\left(\frac{\sqrt{3}t}{2}\right) c_1\right)}{3} \\ -\frac{2 \left(\sqrt{3} (c_1 + \frac{c_2}{2}) \sin\left(\frac{\sqrt{3}t}{2}\right) - \frac{3 \cos\left(\frac{\sqrt{3}t}{2}\right) c_2}{2}\right) e^{-\frac{9t}{2}}}{3} \end{bmatrix}$$

The particular solution given by

$$\vec{x}_p(t) = e^{At} \int e^{-At} \vec{G}(t) dt$$

But

$$e^{-At} = (e^{At})^{-1}$$

$$= \begin{bmatrix} -\frac{\left(\sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right) - 3 \cos\left(\frac{\sqrt{3}t}{2}\right)\right) e^{\frac{9t}{2}}}{3} & -\frac{2\sqrt{3}e^{\frac{9t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right)}{3} \\ \frac{2\sqrt{3}e^{\frac{9t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right)}{3} & \frac{\left(\sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right) + 3 \cos\left(\frac{\sqrt{3}t}{2}\right)\right) e^{\frac{9t}{2}}}{3} \end{bmatrix}$$

Hence

$$\begin{aligned}
 \vec{x}_p(t) &= \begin{bmatrix} \frac{e^{-\frac{9t}{2}} \left(\sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right) + 3 \cos\left(\frac{\sqrt{3}t}{2}\right) \right)}{3} & \frac{2\sqrt{3} e^{-\frac{9t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right)}{3} \\ -\frac{2\sqrt{3} e^{-\frac{9t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right)}{3} & -\frac{e^{-\frac{9t}{2}} \left(\sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right) - 3 \cos\left(\frac{\sqrt{3}t}{2}\right) \right)}{3} \end{bmatrix} \int \begin{bmatrix} -\frac{\left(\sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right) - 3 \cos\left(\frac{\sqrt{3}t}{2}\right) \right) e^{\frac{9t}{2}}}{3} \\ \frac{2\sqrt{3} e^{\frac{9t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right)}{3} \end{bmatrix} \\
 &= \begin{bmatrix} \frac{e^{-\frac{9t}{2}} \left(\sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right) + 3 \cos\left(\frac{\sqrt{3}t}{2}\right) \right)}{3} & \frac{2\sqrt{3} e^{-\frac{9t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right)}{3} \\ -\frac{2\sqrt{3} e^{-\frac{9t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right)}{3} & -\frac{e^{-\frac{9t}{2}} \left(\sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right) - 3 \cos\left(\frac{\sqrt{3}t}{2}\right) \right)}{3} \end{bmatrix} \begin{bmatrix} -\frac{2e^{\frac{9t}{2}} \left((2t - \frac{117}{14}) \cos\left(\frac{\sqrt{3}t}{2}\right) + \sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right) \right)}{21} \\ -\frac{e^{\frac{9t}{2}} \left((-5t + \frac{1}{7}) \cos\left(\frac{\sqrt{3}t}{2}\right) + \sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right) \right)}{21} \end{bmatrix} \\
 &= \begin{bmatrix} -\frac{4t}{21} + \frac{39}{49} \\ \frac{5t}{21} - \frac{1}{147} \end{bmatrix}
 \end{aligned}$$

Hence the complete solution is

$$\begin{aligned}
 \vec{x}(t) &= \vec{x}_h(t) + \vec{x}_p(t) \\
 &= \begin{bmatrix} \frac{e^{-\frac{9t}{2}} \sqrt{3} (c_1 + 2c_2) \sin\left(\frac{\sqrt{3}t}{2}\right)}{3} + e^{-\frac{9t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right) c_1 - \frac{4t}{21} + \frac{39}{49} \\ -\frac{2\sqrt{3} e^{-\frac{9t}{2}} (c_1 + \frac{c_2}{2}) \sin\left(\frac{\sqrt{3}t}{2}\right)}{3} + e^{-\frac{9t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right) c_2 + \frac{5t}{21} - \frac{1}{147} \end{bmatrix}
 \end{aligned}$$

29.13.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A \vec{x}(t) + \vec{G}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} -4 & 1 \\ -1 & -5 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix} + \begin{bmatrix} -t + 3 \\ t + 1 \end{bmatrix}$$

Since the system is nonhomogeneous, then the solution is given by

$$\vec{x}(t) = \vec{x}_h(t) + \vec{x}_p(t)$$

Where $\vec{x}_h(t)$ is the homogeneous solution to $\vec{x}'(t) = A \vec{x}(t)$ and $\vec{x}_p(t)$ is a particular solution to $\vec{x}'(t) = A \vec{x}(t) + \vec{G}(t)$. The particular solution will be found using variation of parameters method applied to the fundamental matrix.

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det \left(\begin{bmatrix} -4 & 1 \\ -1 & -5 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) = 0$$

Therefore

$$\det \left(\begin{bmatrix} -4 - \lambda & 1 \\ -1 & -5 - \lambda \end{bmatrix} \right) = 0$$

Which gives the characteristic equation

$$\lambda^2 + 9\lambda + 21 = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = -\frac{9}{2} + \frac{i\sqrt{3}}{2}$$

$$\lambda_2 = -\frac{9}{2} - \frac{i\sqrt{3}}{2}$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
$-\frac{9}{2} - \frac{i\sqrt{3}}{2}$	1	complex eigenvalue
$-\frac{9}{2} + \frac{i\sqrt{3}}{2}$	1	complex eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = -\frac{9}{2} - \frac{i\sqrt{3}}{2}$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} -4 & 1 \\ -1 & -5 \end{bmatrix} - \left(-\frac{9}{2} - \frac{i\sqrt{3}}{2} \right) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} \frac{1}{2} + \frac{i\sqrt{3}}{2} & 1 \\ -1 & -\frac{1}{2} + \frac{i\sqrt{3}}{2} \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} \frac{1}{2} + \frac{i\sqrt{3}}{2} & 1 & 0 \\ -1 & -\frac{1}{2} + \frac{i\sqrt{3}}{2} & 0 \end{array} \right]$$

$$R_2 = R_2 + \frac{R_1}{\frac{1}{2} + \frac{i\sqrt{3}}{2}} \implies \left[\begin{array}{cc|c} \frac{1}{2} + \frac{i\sqrt{3}}{2} & 1 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\left[\begin{array}{cc|c} \frac{1}{2} + \frac{i\sqrt{3}}{2} & 1 & 0 \\ 0 & 0 & 0 \end{array} \right] \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\left\{ v_1 = -\frac{2t}{1+i\sqrt{3}} \right\}$

Hence the solution is

$$\begin{bmatrix} -\frac{2t}{1+i\sqrt{3}} \\ t \end{bmatrix} = \begin{bmatrix} -\frac{2t}{1+i\sqrt{3}} \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} -\frac{2t}{1+i\sqrt{3}} \\ t \end{bmatrix} = t \begin{bmatrix} -\frac{2}{1+i\sqrt{3}} \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} -\frac{2t}{1+i\sqrt{3}} \\ t \end{bmatrix} = \begin{bmatrix} -\frac{2}{1+i\sqrt{3}} \\ 1 \end{bmatrix}$$

Which is normalized to

$$\begin{bmatrix} -\frac{2t}{1+i\sqrt{3}} \\ t \end{bmatrix} = \begin{bmatrix} -\frac{2}{1+i\sqrt{3}} \\ 1 \end{bmatrix}$$

Considering the eigenvalue $\lambda_2 = -\frac{9}{2} + \frac{i\sqrt{3}}{2}$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\left[\begin{array}{cc} -4 & 1 \\ -1 & -5 \end{array} \right] - \left(-\frac{9}{2} + \frac{i\sqrt{3}}{2} \right) \left[\begin{array}{cc} 1 & 0 \\ 0 & 1 \end{array} \right] \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\left[\begin{array}{cc|c} \frac{1}{2} - \frac{i\sqrt{3}}{2} & 1 & 0 \\ -1 & -\frac{1}{2} - \frac{i\sqrt{3}}{2} & 0 \end{array} \right] \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} \frac{1}{2} - \frac{i\sqrt{3}}{2} & 1 & 0 \\ -1 & -\frac{1}{2} - \frac{i\sqrt{3}}{2} & 0 \end{array} \right]$$

$$R_2 = R_2 + \frac{R_1}{\frac{1}{2} - \frac{i\sqrt{3}}{2}} \implies \left[\begin{array}{cc|c} \frac{1}{2} - \frac{i\sqrt{3}}{2} & 1 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\left[\begin{array}{cc|c} \frac{1}{2} - \frac{i\sqrt{3}}{2} & 1 & 0 \\ 0 & 0 & 0 \end{array} \right] \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\left\{ v_1 = \frac{2t}{i\sqrt{3}-1} \right\}$

Hence the solution is

$$\begin{bmatrix} \frac{2t}{i\sqrt{3}-1} \\ t \end{bmatrix} = \begin{bmatrix} \frac{2t}{i\sqrt{3}-1} \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} \frac{2t}{i\sqrt{3}-1} \\ t \end{bmatrix} = t \begin{bmatrix} \frac{2}{i\sqrt{3}-1} \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} \frac{2t}{i\sqrt{3}-1} \\ t \end{bmatrix} = \begin{bmatrix} \frac{2}{i\sqrt{3}-1} \\ 1 \end{bmatrix}$$

Which is normalized to

$$\begin{bmatrix} \frac{2t}{i\sqrt{3}-1} \\ t \end{bmatrix} = \begin{bmatrix} \frac{2}{i\sqrt{3}-1} \\ 1 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number

of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
$-\frac{9}{2} + \frac{i\sqrt{3}}{2}$	1	1	No	$\begin{bmatrix} \frac{1}{-\frac{1}{2} + \frac{i\sqrt{3}}{2}} \\ 1 \end{bmatrix}$
$-\frac{9}{2} - \frac{i\sqrt{3}}{2}$	1	1	No	$\begin{bmatrix} \frac{1}{-\frac{1}{2} - \frac{i\sqrt{3}}{2}} \\ 1 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. Therefore the homogeneous solution is

$$\vec{x}_h(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

Which is written as

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = c_1 \begin{bmatrix} \frac{e\left(-\frac{9}{2} + \frac{i\sqrt{3}}{2}\right)t}{-\frac{1}{2} + \frac{i\sqrt{3}}{2}} \\ e\left(-\frac{9}{2} + \frac{i\sqrt{3}}{2}\right)t \end{bmatrix} + c_2 \begin{bmatrix} \frac{e\left(-\frac{9}{2} - \frac{i\sqrt{3}}{2}\right)t}{-\frac{1}{2} - \frac{i\sqrt{3}}{2}} \\ e\left(-\frac{9}{2} - \frac{i\sqrt{3}}{2}\right)t \end{bmatrix}$$

Now that we found homogeneous solution above, we need to find a particular solution $\vec{x}_p(t)$. We will use Variation of parameters. The fundamental matrix is

$$\Phi = \begin{bmatrix} \vec{x}_1 & \vec{x}_2 & \dots \end{bmatrix}$$

Where \vec{x}_i are the solution basis found above. Therefore the fundamental matrix is

$$\Phi(t) = \begin{bmatrix} \frac{e\left(-\frac{9}{2} + \frac{i\sqrt{3}}{2}\right)t}{-\frac{1}{2} + \frac{i\sqrt{3}}{2}} & \frac{e\left(-\frac{9}{2} - \frac{i\sqrt{3}}{2}\right)t}{-\frac{1}{2} - \frac{i\sqrt{3}}{2}} \\ e\left(-\frac{9}{2} + \frac{i\sqrt{3}}{2}\right)t & e\left(-\frac{9}{2} - \frac{i\sqrt{3}}{2}\right)t \end{bmatrix}$$

The particular solution is then given by

$$\vec{x}_p(t) = \Phi \int \Phi^{-1} \vec{G}(t) dt$$

But

$$\Phi^{-1} = \begin{bmatrix} \frac{i\sqrt{3}e^{-\frac{(-9+i\sqrt{3})t}{2}}}{3} & \frac{\sqrt{3}e^{-\frac{(-9+i\sqrt{3})t}{2}}(\sqrt{3}+i)}{6} \\ -\frac{i\sqrt{3}e^{\frac{(i\sqrt{3}+9)t}{2}}}{3} & -\frac{\sqrt{3}e^{\frac{(i\sqrt{3}+9)t}{2}}(i-\sqrt{3})}{6} \end{bmatrix}$$

Hence

$$\begin{aligned} \vec{x}_p(t) &= \begin{bmatrix} \frac{e^{\left(-\frac{9}{2}+\frac{i\sqrt{3}}{2}\right)t}}{-\frac{1}{2}+\frac{i\sqrt{3}}{2}} & \frac{e^{\left(-\frac{9}{2}-\frac{i\sqrt{3}}{2}\right)t}}{-\frac{1}{2}-\frac{i\sqrt{3}}{2}} \\ e^{\left(-\frac{9}{2}+\frac{i\sqrt{3}}{2}\right)t} & e^{\left(-\frac{9}{2}-\frac{i\sqrt{3}}{2}\right)t} \end{bmatrix} \int \begin{bmatrix} \frac{i\sqrt{3}e^{-\frac{(-9+i\sqrt{3})t}{2}}}{3} & \frac{\sqrt{3}e^{-\frac{(-9+i\sqrt{3})t}{2}}(\sqrt{3}+i)}{6} \\ -\frac{i\sqrt{3}e^{\frac{(i\sqrt{3}+9)t}{2}}}{3} & -\frac{\sqrt{3}e^{\frac{(i\sqrt{3}+9)t}{2}}(i-\sqrt{3})}{6} \end{bmatrix} \begin{bmatrix} -t+3 \\ t+1 \end{bmatrix} dt \\ &= \begin{bmatrix} \frac{e^{\left(-\frac{9}{2}+\frac{i\sqrt{3}}{2}\right)t}}{-\frac{1}{2}+\frac{i\sqrt{3}}{2}} & \frac{e^{\left(-\frac{9}{2}-\frac{i\sqrt{3}}{2}\right)t}}{-\frac{1}{2}-\frac{i\sqrt{3}}{2}} \\ e^{\left(-\frac{9}{2}+\frac{i\sqrt{3}}{2}\right)t} & e^{\left(-\frac{9}{2}-\frac{i\sqrt{3}}{2}\right)t} \end{bmatrix} \int \begin{bmatrix} \frac{e^{-\frac{(-9+i\sqrt{3})t}{2}}(3-i(t-7)\sqrt{3}+3t)}{6} \\ \frac{e^{\frac{(i\sqrt{3}+9)t}{2}}(3+i(t-7)\sqrt{3}+3t)}{6} \end{bmatrix} dt \\ &= \begin{bmatrix} \frac{e^{\left(-\frac{9}{2}+\frac{i\sqrt{3}}{2}\right)t}}{-\frac{1}{2}+\frac{i\sqrt{3}}{2}} & \frac{e^{\left(-\frac{9}{2}-\frac{i\sqrt{3}}{2}\right)t}}{-\frac{1}{2}-\frac{i\sqrt{3}}{2}} \\ e^{\left(-\frac{9}{2}+\frac{i\sqrt{3}}{2}\right)t} & e^{\left(-\frac{9}{2}-\frac{i\sqrt{3}}{2}\right)t} \end{bmatrix} \begin{bmatrix} -\frac{\sqrt{3}e^{-\frac{(-9+i\sqrt{3})t}{2}}(-7\sqrt{3}t-116\sqrt{3}+63it-118i)(t+i\sqrt{3}t-7+i\sqrt{3})}{1764(2i\sqrt{3}+t-1)} \\ \frac{\sqrt{3}e^{\frac{(i\sqrt{3}+9)t}{2}}(7\sqrt{3}t+116\sqrt{3}+63it-118i)(-t+i\sqrt{3}t+7+i\sqrt{3})}{3528i\sqrt{3}-1764t+1764} \end{bmatrix} \\ &= \begin{bmatrix} \frac{28t^3-173t^2+598t-1521}{-1764-147(t-1)^2} \\ \frac{-35t^3+71t^2-457t+13}{-1764-147(t-1)^2} \end{bmatrix} \end{aligned}$$

Now that we found particular solution, the final solution is

$$\begin{aligned} \vec{x}(t) &= \vec{x}_h(t) + \vec{x}_p(t) \\ \begin{bmatrix} x(t) \\ y(t) \end{bmatrix} &= \begin{bmatrix} \frac{c_1 e^{\left(-\frac{9}{2}+\frac{i\sqrt{3}}{2}\right)t}}{-\frac{1}{2}+\frac{i\sqrt{3}}{2}} \\ c_1 e^{\left(-\frac{9}{2}+\frac{i\sqrt{3}}{2}\right)t} \end{bmatrix} + \begin{bmatrix} \frac{c_2 e^{\left(-\frac{9}{2}-\frac{i\sqrt{3}}{2}\right)t}}{-\frac{1}{2}-\frac{i\sqrt{3}}{2}} \\ c_2 e^{\left(-\frac{9}{2}-\frac{i\sqrt{3}}{2}\right)t} \end{bmatrix} + \begin{bmatrix} \frac{28t^3-173t^2+598t-1521}{-1764-147(t-1)^2} \\ \frac{-35t^3+71t^2-457t+13}{-1764-147(t-1)^2} \end{bmatrix} \end{aligned}$$

Which becomes

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} \frac{39}{49} + \frac{c_1(-i\sqrt{3}-1)e^{\frac{(-9+i\sqrt{3})t}{2}}}{2} + \frac{c_2(i\sqrt{3}-1)e^{\frac{(i\sqrt{3}+9)t}{2}}}{2} - \frac{4t}{21} \\ c_1 e^{\frac{(-9+i\sqrt{3})t}{2}} + c_2 e^{\frac{(i\sqrt{3}+9)t}{2}} + \frac{5t}{21} - \frac{1}{147} \end{bmatrix}$$

29.13.3 Maple step by step solution

Let's solve

$$[x'(t) = -4x(t) + y(t) - t + 3, y'(t) = -x(t) - 5y(t) + t + 1]$$

- Define vector

$$\vec{x}(t) = \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

- Convert system into a vector equation

$$\vec{x}'(t) = \begin{bmatrix} -4 & 1 \\ -1 & -5 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} -t + 3 \\ t + 1 \end{bmatrix}$$

- System to solve

$$\vec{x}'(t) = \begin{bmatrix} -4 & 1 \\ -1 & -5 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} -t + 3 \\ t + 1 \end{bmatrix}$$

- Define the forcing function

$$\vec{f}(t) = \begin{bmatrix} -t + 3 \\ t + 1 \end{bmatrix}$$

- Define the coefficient matrix

$$A = \begin{bmatrix} -4 & 1 \\ -1 & -5 \end{bmatrix}$$

- Rewrite the system as

$$\vec{x}'(t) = A \cdot \vec{x}(t) + \vec{f}$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[\begin{bmatrix} -\frac{9}{2} - \frac{1\sqrt{3}}{2}, \begin{bmatrix} \frac{1}{-\frac{1}{2} - \frac{1\sqrt{3}}{2}} \\ 1 \end{bmatrix} \end{bmatrix}, \begin{bmatrix} -\frac{9}{2} + \frac{1\sqrt{3}}{2}, \begin{bmatrix} \frac{1}{-\frac{1}{2} + \frac{1\sqrt{3}}{2}} \\ 1 \end{bmatrix} \end{bmatrix} \right] \right]$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$\left[-\frac{9}{2} - \frac{I\sqrt{3}}{2}, \begin{bmatrix} \frac{1}{-\frac{1}{2} - \frac{I\sqrt{3}}{2}} \\ 1 \end{bmatrix} \right]$$

- Solution from eigenpair

$$e^{(-\frac{9}{2} - \frac{I\sqrt{3}}{2})t} \cdot \begin{bmatrix} \frac{1}{-\frac{1}{2} - \frac{I\sqrt{3}}{2}} \\ 1 \end{bmatrix}$$

- Use Euler identity to write solution in terms of sin and cos

$$e^{-\frac{9t}{2}} \cdot \left(\cos\left(\frac{\sqrt{3}t}{2}\right) - I \sin\left(\frac{\sqrt{3}t}{2}\right) \right) \cdot \begin{bmatrix} \frac{1}{-\frac{1}{2} - \frac{I\sqrt{3}}{2}} \\ 1 \end{bmatrix}$$

- Simplify expression

$$e^{-\frac{9t}{2}} \cdot \begin{bmatrix} \frac{\cos\left(\frac{\sqrt{3}t}{2}\right) - I \sin\left(\frac{\sqrt{3}t}{2}\right)}{-\frac{1}{2} - \frac{I\sqrt{3}}{2}} \\ \cos\left(\frac{\sqrt{3}t}{2}\right) - I \sin\left(\frac{\sqrt{3}t}{2}\right) \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\left[\vec{x}_1(t) = e^{-\frac{9t}{2}} \cdot \begin{bmatrix} -\frac{\cos\left(\frac{\sqrt{3}t}{2}\right)}{2} + \frac{\sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right)}{2} \\ \cos\left(\frac{\sqrt{3}t}{2}\right) \end{bmatrix}, \vec{x}_2(t) = e^{-\frac{9t}{2}} \cdot \begin{bmatrix} \frac{\sqrt{3} \cos\left(\frac{\sqrt{3}t}{2}\right)}{2} + \frac{\sin\left(\frac{\sqrt{3}t}{2}\right)}{2} \\ -\sin\left(\frac{\sqrt{3}t}{2}\right) \end{bmatrix} \right]$$

- General solution of the system of ODEs can be written in terms of the particular solution $\vec{x}_p(t)$
 $\vec{x}(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t) + \vec{x}_p(t)$

□ Fundamental matrix

- Let $\phi(t)$ be the matrix whose columns are the independent solutions of the homogeneous system

$$\phi(t) = \begin{bmatrix} e^{-\frac{9t}{2}} \left(-\frac{\cos\left(\frac{\sqrt{3}t}{2}\right)}{2} + \frac{\sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right)}{2} \right) & e^{-\frac{9t}{2}} \left(\frac{\sqrt{3} \cos\left(\frac{\sqrt{3}t}{2}\right)}{2} + \frac{\sin\left(\frac{\sqrt{3}t}{2}\right)}{2} \right) \\ e^{-\frac{9t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right) & -e^{-\frac{9t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right) \end{bmatrix}$$

- The fundamental matrix, $\Phi(t)$ is a normalized version of $\phi(t)$ satisfying $\Phi(0) = I$ where I is the identity matrix

$$\Phi(t) = \phi(t) \cdot \frac{1}{\phi(0)}$$

- Substitute the value of $\phi(t)$ and $\phi(0)$

$$\Phi(t) = \begin{bmatrix} e^{-\frac{9t}{2}} \left(-\frac{\cos\left(\frac{\sqrt{3}t}{2}\right)}{2} + \frac{\sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right)}{2} \right) & e^{-\frac{9t}{2}} \left(\frac{\sqrt{3} \cos\left(\frac{\sqrt{3}t}{2}\right)}{2} + \frac{\sin\left(\frac{\sqrt{3}t}{2}\right)}{2} \right) \\ e^{-\frac{9t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right) & -e^{-\frac{9t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right) \end{bmatrix} \cdot \frac{1}{\begin{bmatrix} -\frac{1}{2} & \frac{\sqrt{3}}{2} \\ 1 & 0 \end{bmatrix}}$$

- Evaluate and simplify to get the fundamental matrix

$$\Phi(t) = \begin{bmatrix} \frac{e^{-\frac{9t}{2}} \left(\sqrt{3} \cos\left(\frac{\sqrt{3}t}{2}\right) + \sin\left(\frac{\sqrt{3}t}{2}\right) \right) \sqrt{3}}{3} & \frac{2\sqrt{3} e^{-\frac{9t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right)}{3} \\ -\frac{2\sqrt{3} e^{-\frac{9t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right)}{3} & -\frac{e^{-\frac{9t}{2}} \left(\sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right) - 3 \cos\left(\frac{\sqrt{3}t}{2}\right) \right)}{3} \end{bmatrix}$$

- Find a particular solution of the system of ODEs using variation of parameters

- Let the particular solution be the fundamental matrix multiplied by $\vec{v}(t)$ and solve for $\vec{v}(t)$

$$\vec{x}_p(t) = \Phi(t) \cdot \vec{v}(t)$$

- Take the derivative of the particular solution

$$\vec{x}'_p(t) = \Phi'(t) \cdot \vec{v}(t) + \Phi(t) \cdot \vec{v}'(t)$$

- Substitute particular solution and its derivative into the system of ODEs

$$\Phi'(t) \cdot \vec{v}(t) + \Phi(t) \cdot \vec{v}'(t) = A \cdot \Phi(t) \cdot \vec{v}(t) + \vec{f}(t)$$

- The fundamental matrix has columns that are solutions to the homogeneous system so its derivative is $A \cdot \Phi(t)$

$$A \cdot \Phi(t) \cdot \vec{v}(t) + \Phi(t) \cdot \vec{v}'(t) = A \cdot \Phi(t) \cdot \vec{v}(t) + \vec{f}(t)$$

- Cancel like terms

$$\Phi(t) \cdot \vec{v}'(t) = \vec{f}(t)$$

- Multiply by the inverse of the fundamental matrix

$$\vec{v}'(t) = \frac{1}{\Phi(t)} \cdot \vec{f}(t)$$

- Integrate to solve for $\vec{v}(t)$

$$\vec{v}(t) = \int_0^t \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds$$

- Plug $\vec{v}(t)$ into the equation for the particular solution

$$\vec{x}_p(t) = \Phi(t) \cdot \left(\int_0^t \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds \right)$$

- Plug in the fundamental matrix and the forcing function and compute

$$\vec{x}_p(t) = \begin{bmatrix} -\frac{115\sqrt{3}e^{-\frac{9t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right)}{441} - \frac{4t}{21} + \frac{39}{49} - \frac{39e^{-\frac{9t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right)}{49} \\ \frac{233\sqrt{3}e^{-\frac{9t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right)}{441} + \frac{5t}{21} + \frac{e^{-\frac{9t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right)}{147} - \frac{1}{147} \end{bmatrix}$$

- Plug particular solution back into general solution

$$\vec{x}(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t) + \begin{bmatrix} -\frac{115\sqrt{3}e^{-\frac{9t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right)}{441} - \frac{4t}{21} + \frac{39}{49} - \frac{39e^{-\frac{9t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right)}{49} \\ \frac{233\sqrt{3}e^{-\frac{9t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right)}{441} + \frac{5t}{21} + \frac{e^{-\frac{9t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right)}{147} - \frac{1}{147} \end{bmatrix}$$

- Substitute in vector of dependent variables

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} -\frac{e^{-\frac{9t}{2}}(-c_2\sqrt{3}+c_1+\frac{78}{49}) \cos\left(\frac{\sqrt{3}t}{2}\right)}{2} + \frac{((c_1-\frac{230}{441})\sqrt{3}+c_2)e^{-\frac{9t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right)}{2} - \frac{4t}{21} + \frac{39}{49} \\ \frac{(441c_1+3)e^{-\frac{9t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right)}{441} - \left(c_2 - \frac{233\sqrt{3}}{441}\right) e^{-\frac{9t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right) + \frac{5t}{21} - \frac{1}{147} \end{bmatrix}$$

- Solution to the system of ODEs

$$\begin{cases} x(t) = -\frac{e^{-\frac{9t}{2}}(-c_2\sqrt{3}+c_1+\frac{78}{49}) \cos\left(\frac{\sqrt{3}t}{2}\right)}{2} + \frac{((c_1-\frac{230}{441})\sqrt{3}+c_2)e^{-\frac{9t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right)}{2} - \frac{4t}{21} + \frac{39}{49} \\ y(t) = \frac{(441c_1+3)e^{-\frac{9t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right)}{441} - \left(c_2 - \frac{233\sqrt{3}}{441}\right) e^{-\frac{9t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right) + \frac{5t}{21} - \frac{1}{147} \end{cases}$$

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 106

```
dsolve([diff(x(t),t)=-4*x(t)+y(t)-t+3,diff(y(t),t)=-x(t)-5*y(t)+t+1],singsol=all)
```

$$\begin{aligned} x(t) &= e^{-\frac{9t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right) c_2 + e^{-\frac{9t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right) c_1 - \frac{4t}{21} + \frac{39}{49} \\ y(t) &= -\frac{e^{-\frac{9t}{2}} \sin\left(\frac{\sqrt{3}t}{2}\right) c_2}{2} + \frac{e^{-\frac{9t}{2}} \sqrt{3} \cos\left(\frac{\sqrt{3}t}{2}\right) c_2}{2} \\ &\quad - \frac{e^{-\frac{9t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right) c_1}{2} - \frac{e^{-\frac{9t}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right) c_1}{2} - \frac{1}{147} + \frac{5t}{21} \end{aligned}$$

✓ Solution by Mathematica

Time used: 2.267 (sec). Leaf size: 131

```
DSolve[{x'[t]==-4*x[t]+y[t]-t+3,y'[t]==-x[t]-5*y[t]+t+1},{x[t],y[t]},t,IncludeSingularSoluti
```

$$x(t) \rightarrow -\frac{4t}{21} + c_1 e^{-9t/2} \cos\left(\frac{\sqrt{3}t}{2}\right) + \frac{(c_1 + 2c_2)e^{-9t/2} \sin\left(\frac{\sqrt{3}t}{2}\right)}{\sqrt{3}} + \frac{39}{49}$$

$$y(t) \rightarrow \frac{5t}{21} + c_2 e^{-9t/2} \cos\left(\frac{\sqrt{3}t}{2}\right) - \frac{(2c_1 + c_2)e^{-9t/2} \sin\left(\frac{\sqrt{3}t}{2}\right)}{\sqrt{3}} - \frac{1}{147}$$

30 Chapter 10. Systems of First-Order Equations.

Section B. Challenge Problems. Page 401

30.1 problem 1	4225
30.2 problem 2	4227

30.1 problem 1

Internal problem ID [6542]

Internal file name [OUTPUT/5790_Sunday_June_05_2022_03_54_37_PM_43080441/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 10. Systems of First-Order Equations. Section B. Challenge Problems.
Page 401

Problem number: 1.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**system of linear ODEs**" Unable to solve or complete t

Solve

$$x'(t) = x(t) y(t) + 1$$

$$y'(t) = -x(t) + y(t)$$

With initial conditions

$$[x(0) = 2, y(0) = -1]$$

Does not currently support non linear system of equations. This is the phase plot of the system.




Figure 511: Phase plot

X Solution by Maple

```
dsolve([diff(x(t),t) = x(t)*y(t)+1, diff(y(t),t) = -x(t)+y(t), x(0) = 2, y(0) = -1], singsol
```

No solution found

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[{x'[t]==x[t]*y[t]+1,y'[t]==-x[t]+y[t]},{x[0]==2,y[0]==-1},{x[t],y[t]},t,IncludeSingul
```

Not solved

30.2 problem 2

Internal problem ID [6543]

Internal file name [OUTPUT/5791_Sunday_June_05_2022_03_54_40_PM_17429521/index.tex]

Book: Differential Equations: Theory, Technique, and Practice by George Simmons, Steven Krantz. McGraw-Hill NY. 2007. 1st Edition.

Section: Chapter 10. Systems of First-Order Equations. Section B. Challenge Problems. Page 401

Problem number: 2.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**system of linear ODEs**" Unable to solve or complete t

Solve

$$\begin{aligned}x'(t) &= ty(t) + 1 \\y'(t) &= -x(t)t + y(t)\end{aligned}$$

With initial conditions

$$[x(0) = 0, y(0) = -1]$$

Does not currently support non autonomous system of first order linear differential equations. The following is the phase plot

X Solution by Maple

```
dsolve([diff(x(t),t) = t*y(t)+1, diff(y(t),t) = -t*x(t)+y(t), x(0) = 0, y(0) = -1], singsol=
```

No solution found

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[{x'[t]==x[t]*y[t]+1,y'[t]==-x[t]+y[t]},{x[0]==2,y[0]==-1},{x[t],y[t]},t,IncludeSingul
```

Not solved